

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 November 2006 (02.11.2006)

PCT

(10) International Publication Number
WO 2006/116583 A2

(51) International Patent Classification:
A61K 31/787 (2006.01)

(21) International Application Number:
PCT/US2006/015983

(22) International Filing Date: 25 April 2006 (25.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/675,260 26 April 2005 (26.04.2005) US

(71) Applicant (for all designated States except US): **THE CURATORS OF THE UNIVERSITY OF MISSOURI** [US/US]; Office of Technology and Special Projects, 475 McReynolds Hall, Columbia, Missouri 65211 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **PHILLIPS, Jeffrey** [US/US]; 1250 E. Nashville Church Road, Ashland, Missouri 65010 (US).

(74) Agent: **FOURNIER, David, B.**; Mayer, Brown, Rowe & Maw LLP, P.o. Box 2828, Chicago, Illinois 60690-2828 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS COMPRISING A POLYMERIZED BENZIMIDAZOLE AND A BUFFERING AGENT AND METHODS OF USING SAME

(57) Abstract: The present invention provides pharmaceutical compositions comprising at least one polymerized benzimidazole, at least one buffering agent and/or a protein component, and one or more optional pharmaceutical excipients. Compositions of the invention are useful in treating, inter alia, gastrointestinal diseases and disorders.



WO 2006/116583 A2

COMPOSITIONS COMPRISING A POLYMERIZED BENZIMIDAZOLE AND A BUFFERING AGENT AND METHODS OF USING SAME

FIELD OF THE INVENTION

5 The present invention relates to pharmaceutical compositions comprising a polymerized benzimidazole and a buffering agent, to methods for manufacture of such compositions, and to use of such compositions in treating various diseases and/or disorders.

BACKGROUND OF THE INVENTION

10 Gastrointestinal disorders such as active duodenal ulcers, gastric ulcers, gastroesophageal reflux disease (GERD), severe erosive esophagitis, poorly responsive symptomatic GERD, and pathological hypersecretory conditions such as Zollinger Ellison syndrome represent a major global health concern impacting millions of people globally. In fact, it is estimated that as many as 60 million
15 Americans experience reflux at least once a month, while approximately 19 million Americans suffer from GERD.

 In the past, the above-described (and other related) gastrointestinal disorders and their associated symptoms have been treated with H₂ histamine antagonists and antacids. Unfortunately, many such available treatments are not
20 completely effective in ameliorating the disorders themselves or their symptoms; additionally, many produce adverse side effects including, among others, constipation, diarrhea, and thrombocytopenia. Moreover, H₂ antagonists such as ranitidine and cimetidine are relatively costly modes of therapy generally requiring multiple daily doses to produce some control of acid secretion. In addition,
25 tolerance to H₂ antagonists increases with continued use thus prohibiting clinical utility in chronic dosing settings.

 More recently, at least some of the above-described gastrointestinal disorders have been treated with proton pump inhibitors (also called PPIs). PPIs are believed to reduce gastric acid production by inhibiting H⁺, K⁺-ATPase of the
30 parietal cell—the final common pathway for gastric acid secretion. One particular class of PPIs includes substituted benzimidazole compounds that contain a sulfinyl group bridging substituted benzimidazole and pyridine rings.

At neutral pH, these PPIs are chemically stable, lipid-soluble compounds that have little or no inhibitory activity. It is believed that the neutral PPIs reach parietal cells from the blood and diffuse into the secretory canaliculi where they become protonated and thereby trapped. The protonated agent is then believed to
5 rearrange to form a sulfenic acid and a sulfenamide. The sulfenamide, in turn, is thought to interact covalently with sulfhydryl groups at critical sites in the extracellular (luminal) domain of the membrane-spanning H^+ , K^+ -ATPase. *See e.g. Hardman et al., Goodman & Gilman's The Pharmacological Basis of Therapeutics*, p. 907, 9th ed. (1996).

10 Unfortunately, commercially available substituted benzimidazole compounds are unstable at neutral or acidic pH and undergo decomposition in gastrointestinal fluid upon oral administration, thereby resulting in loss of therapeutic activity. To overcome this acid instability, such compounds are typically formulated for oral delivery as enteric coated solid dosage forms, for
15 example enteric coated tablets, which coating protects the drug from contact with acidic stomach secretions. An undesirable consequence of such enteric coating is that therapeutic onset time is significantly delayed by comparison with non-enteric coated dosage forms. Such prolonged time to therapeutic onset is particularly undesirable for patients in need of rapid relief from one or more of the above
20 described disorders or symptoms.

Specific examples of enteric coated PPIs include omeprazole (Prilosec®), lansoprazole (Prevacid®), perprazole (also referred to as esomeprazole; Nexium®), rabeprazole (Aciphex®), and pantoprazole (Protonix®). Omeprazole, a substituted benzimidazole, has the following chemical name: 5-methoxy-2-[(4-
25 methoxy-3,5-dimethyl-2-pyridinyl) methyl] sulfinyl]-1H-benzimidazole.

U.S. Patent No. 4,786,505 to Lovgren *et al.* discloses that a pharmaceutical oral solid dosage form of omeprazole must be protected from contact with acidic gastric juice by an enteric coating to maintain its pharmaceutical activity. That patent describes an enteric coated omeprazole preparation containing an alkaline
30 core comprising omeprazole, a subcoating over the core, and an enteric coating over the subcoating.

One recent attempt to overcome acid instability associated with PPIs is disclosed in U.S. Patent No. 6,617,338 to Mali *et al.* The Mali patent discloses use of polymerized benzimidazoles for the treatment or prevention of peptic ulcers,

gastrointestinal inflammatory disease or gastrointestinal disorder. Polymerized benzimidazoles disclosed therein are said to be more acid stable than respective parent benzimidazole moieties. In fact, Mali *et al.*, state that the polymeric benzimidazoles of that invention are found to be acid stable due to the polymeric N-substitution and therefore do not disintegrate in the gastrointestinal fluid and are suitable for administration without enteric coating.

However, it is presently believed that polymeric benzimidazole compositions disclosed by Mali *et al.*, will not adequately overcome the *in vivo* acid degradation problems described above. Therefore, a continuing unmet need exists for orally deliverable compositions comprising acid-labile drugs (for example PPIs) which formulations are capable of providing rapid onset of therapeutic effect while improving the acid degradation problems associated therewith. A significant advance in the art would result if compositions overcoming these and other problems could be prepared.

SUMMARY OF THE INVENTION

In one embodiment, the present invention provides pharmaceutical compositions comprising at least one polymerized benzimidazole, for example a polymerized proton pump inhibitor, one or more buffering agents and one or more optional pharmaceutical excipients. Such compositions can further comprise a protein component.

In another embodiment, the present invention provides pharmaceutical compositions comprising at least one polymerized benzimidazole, for example a PPI, a protein component and one or more optional pharmaceutical excipients. Such compositions can further comprise a buffering agent.

In another embodiment, the present invention provides an orally deliverable pharmaceutical composition comprising a polymerized PPI and one or more buffering agents (and/or a protein component) wherein upon oral administration of the composition to a plurality of fasted adult human subjects, the subjects exhibit an average plasma concentration of the PPI moiety of at least about 0.1 mg/ml at any time within about 30 minutes after administration.

Also provided by the present invention are methods of treating acid related gastrointestinal disorders by administering to a subject one or more compositions described herein.

DETAILED DESCRIPTION OF THE INVENTION

While the present invention is capable of being embodied in various forms, the description below of several embodiments is made with the understanding that the present disclosure is to be considered as an exemplification of the invention, and is not intended to limit the invention to the specific embodiments illustrated. Headings are provided for convenience only and are not to be construed to limit the invention in any way. Embodiments illustrated under any heading may be combined with embodiments illustrated under any other heading.

The use of numerical values in the various ranges specified in this application, unless expressly indicated otherwise, are stated as approximations as though the minimum and maximum values within the stated ranges were both preceded by the word "about." In this manner, slight variations above and below the stated ranges can be used to achieve substantially the same results as values within the ranges. As used herein, the terms "about" and "approximately" when referring to a numerical value shall have their plain and ordinary meanings to one skilled in the art of pharmaceutical sciences or the art relevant to the range or element at issue. The amount of broadening from the strict numerical boundary depends upon many factors. For example, some of the factors to be considered may include the criticality of the element and/or the effect a given amount of variation will have on the performance of the claimed subject matter, as well as other considerations known to those of skill in the art. Thus, as a general matter, "about" or "approximately" broaden the numerical value. For example, in some cases, "about" or "approximately" may mean $\pm 5\%$, or $\pm 10\%$, or $\pm 20\%$, or $\pm 30\%$ depending on the relevant technology. Also, the disclosure of ranges is intended as a continuous range including every value between the minimum and maximum values recited as well as any ranges that can be formed thereby.

It is also to be understood that any ranges, ratios and ranges of ratios that can be formed by any of the numbers or data present herein represent further embodiments of the present invention. This includes ranges that can be formed that do or do not include a finite upper and/or lower boundary. For example, by way of illustration and not limitation, in one embodiment, a benzimidazole moiety is present in a composition of the invention in an amount of about 1 to about 2000 mg; in another embodiment, a buffering agent is present in a composition of the

invention in an amount of about 200 mg to about 3500 mg. One of skill in the art will therefore recognize, for example, that additional embodiments include situations where a composition has a benzimidazole:buffering agent weight ratio of less than or greater than 1:200, less than or greater than 1:3500, less than or greater than 10:1 (2000/200), less than or greater than 1:3500, less than or greater than about 200:3500, or in ranges of about 1:200 to about 1:3500, about 1:3500 to about 10:1, 1:200 to about to about 200:3500, *etc.* Accordingly, the skilled person will appreciate that many such ratios, ranges, and ranges of ratios can be unambiguously derived from the data and numbers presented herein and all represent embodiments of the present invention.

Polymeric Benzimidazoles

Compositions of the present invention comprise a polymeric benzimidazole, also referred to herein as “polymer-based benzimidazoles”, or “polymerized benzimidazoles”. The term “polymeric benzimidazole” and the interchangeable terms listed immediately above refer to a compound comprising a benzimidazole moiety, wherein the compound is bonded or linked to a least one pharmaceutically acceptable polymer. The term “benzimidazole moiety” refers to the benzimidazole moiety of formulas (II) and (III) as shown herein below, not including any pharmaceutically acceptable polymer substituted at the N position of the benzimidazole ring.

Benzimidazole moiety-containing compounds or drugs useful in the present invention include any pharmacologically active benzimidazole compounds that are susceptible to acid catalyzed degradation and that can be substituted at the N position of the benzimidazole ring. The term “pharmacologically active benzimidazole compound” includes any such compound or drug which is capable of producing a pharmacological response in a subject, irrespective of whether therapeutic, diagnostic, or prophylactic in nature.


While not wishing to be bound by any one theory, the acidic decomposition of an acid-labile benzimidazole compound is believed to be due to an acid catalyzed reaction as described by, for example, G. Rackur *et al.*, in *Biochem. Biophys. Res. Commun.* 1985: 128(1). p. 477 - 484. Thus, pharmacologically

active agents useful in the present invention are those which are degraded by acids, even organic acids, or are degraded in acid catalyzed reactions.

A particularly preferred class of benzimidazole moiety-containing acid-labile compounds suitable for polymeric N-substitution and preparation of polymeric benzimidazoles useful in the present invention include substituted benzimidazole compounds possessing pharmacological activity as an inhibitor of H^+ , K^+ -ATPase (*i.e.* proton pump inhibitors). Such proton pump inhibitors can, if desired, be in the form of a free base, a free acid, a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a prodrug, a polymorph, a derivative or the like, and can be in racemic or enantiomeric form.




Polymeric benzimidazoles suitable for use in the present invention can be prepared by any process known in the art. For example, otherwise acid-labile benzimidazole compounds can be used as starting agents for formation of polymeric benzimidazoles by polymeric N-substitution as described in U.S. Patent No. 6,617,338, which is hereby incorporated by reference in its entirety. Without being bound by theory, it is believed that upon oral administration of these polymeric N-substituted benzimidazoles with a buffering agent or protein component as described herein, enzymes and chemicals in the gastrointestinal fluid cleave at the hydrolysable group (*e.g.* (E) of formula (I)) to release an N-substituted benzimidazole derivative that may or may not be the original starting benzimidazole moiety (*e.g.* moiety B below) having a portion of the polymer remaining attached thereto. Without being held to a particular theory, it is also presently believed that the N-substituted benzimidazole derivative, in the presence of the buffering agent or protein component, maintains its pharmacological activity as an inhibitor of H^+ , K^+ -ATPase, while the cleaved portion of the polymer becomes an inert, non-toxic and/or non-absorbable metabolite that is excreted from the body.

In one embodiment, suitable polymeric benzimidazoles are those compounds of formula (I):



wherein R_7 , R_8 , R_9 , and R_{10} , are independently H or a lower alkyl (preferably methyl or ethyl), for example a C_{1-4} alkyl;

U is



5


R_{11} is H, CH₃, C₂H₅, or CONH₂;

Y and V are independently OH or NH₂;

E is —COO—; and

10

B is a benzimidazole moiety of formula (II):



wherein R^1 is hydrogen, alkyl, halogen, cyano, carboxy, carboalkoxy, carboalkoxyalkyl, carbamoyl, carbamoylalkyl, hydroxy, alkoxy which may optionally be fluorinated, hydroxyalkyl, trifluoromethyl, acyl, carbamoyloxy, nitro, acyloxy, aryl, aryloxy, alkylthio, or alkylsulfinyl;


5 Q is nitrogen, CH, or CR^1 ;

W is nitrogen, CH, or CR^1 ; and

y is an integer of 0 through 4;

or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a polymorph, a polymorph, a prodrug, or a derivative thereof.

10 In another embodiment, (B) is a benzimidazole moiety of formula (III):



wherein:

R^1 is hydrogen, alkyl, halogen, cyano, carboxy, carboalkoxy, carboalkoxyalkyl, carbamoyl, carbamoylalkyl, hydroxy, alkoxy which may optionally be fluorinated, hydroxyalkyl, trifluoromethyl, acyl, carbamoyloxy, nitro, acyloxy, aryl, aryloxy, alkylthio, or alkylsulfinyl;

R^3 and R^5 are independently hydrogen, alkyl, alkoxy, amino, or alkoxyalkoxy;

20 R^4 is hydrogen, alkyl, alkoxy which may optionally be fluorinated, or alkoxyalkoxy;

Q is nitrogen, CH, or CR^1 ;

W is nitrogen, CH, or CR^1 ;

y is an integer of 0 through 4; and

25 Z is nitrogen, CH, or CR^1 ;

or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a polymorph, a prodrug, or a derivative thereof.

Specific examples of suitable PPIs for use as the benzimidazole moiety (B) include esomeprazole (also referred to as S-omeprazole), ilaprazole (U.S. Pat. No. 5,703,097), lansoprazole, omeprazole, pantoprazole, pariprazole, rabeprazole, tenatoprazole, and nepaprazole or a free base, a free acid, or a salt, hydrate, ester, amide, enantiomer, isomer, tautomer, polymorph, prodrug, or derivative of such compounds.

Proton pump inhibitors as well as their salts, hydrates, esters, amides, enantiomers, isomers, tautomers, polymorphs, prodrugs, and derivatives may be prepared using standard procedures known to those skilled in the art of synthetic organic chemistry. See, *e.g.*, March, *Advanced Organic Chemistry: Reactions, Mechanisms and Structure*, 4th Ed. (New York: Wiley-Interscience, 1992); Leonard *et al.*, *Advanced Practical Organic Chemistry* (1992); Howarth *et al.*, *Core Organic Chemistry* (1998); and Weisermel *et al.*, *Industrial Organic Chemistry* (2002).

“Pharmaceutically acceptable salts,” or “salts,” include the salt of a proton pump inhibitor prepared from formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, stearic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic, methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, cyclohexylaminosulfonic, algenic, beta.-hydroxybutyric, galactaric and galacturonic acids.

Acid addition salts can be prepared from free base forms using conventional methodology involving reaction of the free base with a suitable acid. Suitable acids for preparing acid addition salts include both organic acids, *e.g.*, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like, as well as inorganic acids, *e.g.*, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like.

Acid addition salts can be reconverted to the free base by treatment with a suitable base. In a further embodiment, the acid addition salts of the proton pump inhibitors are halide salts, which are prepared using hydrochloric or hydrobromic

acids. In still other embodiments, the basic salts are alkali metal salts, *e.g.*, sodium salt.

Salt forms of proton pump inhibitors include, but are not limited to: a sodium salt form such as esomeprazole sodium, omeprazole sodium, rabeprazole sodium, pantoprazole sodium; or a magnesium salt form such as esomeprazole magnesium or omeprazole magnesium, described in U.S. Pat. No. 5,900,424; a calcium salt form; or a potassium salt form such as the potassium salt of esomeprazole, described in U.S. Pat. No. 6,511,996. Other salts of esomeprazole are described in U.S. Pat. Nos.: 4,738,974 and 6,369,085. Salt forms of pantoprazole and lansoprazole are discussed in U.S. Pat. Nos. 4,758,579 and 4,628,098, respectively.

In one embodiment, preparation of esters involves functionalizing hydroxyl and/or carboxyl groups that may be present within the molecular structure of the drug. In another embodiment, the esters are acyl-substituted derivatives of free alcohol groups, *e.g.*, moieties derived from carboxylic acids of the formula RCOOR₁ where R₁ is a lower alkyl group. Esters can be reconverted to the free acids, if desired, by using conventional procedures such as hydrogenolysis or hydrolysis.

“Amides” may be prepared using techniques known to those skilled in the art or described in the pertinent literature. For example, amides may be prepared from esters, using suitable amine reactants, or they may be prepared from an anhydride or an acid chloride by reaction with an amine group such as ammonia or a lower alkyl amine.

“Tautomers” of substituted bicyclic aryl-imidazoles include, *e.g.*, tautomers of omeprazole such as those described in U.S. Pat. Nos. 6,262,085; 6,262,086; 6,268,385; 6,312,723; 6,316,020; 6,326,384; 6,369,087; and 6,444,689.

An exemplary “isomer” of a substituted bicyclic aryl-imidazole is the isomer of omeprazole including but not limited to isomers described in: Oishi *et al.*, Acta Cryst. (1989), C45, 1921-1923; U.S. Pat. No. 6,150,380; U.S. Patent Publication No. 02/0156284; and PCT Publication No. WO 02/085889.

Exemplary “polymorphs” include, but are not limited to, those described in PCT Publication No. WO 92/08716, and U.S. Pat. Nos. 4,045,563; 4,182,766; 4,508,905; 4,628,098; 4,636,499; 4,689,333; 4,758,579; 4,783,974; 4,786,505; 4,808,596; 4,853,230; 5,026,560; 5,013,743; 5,035,899; 5,045,321; 5,045,552; 5,093,132; 5,093,342; 5,433,959; 5,464,632; 5,536,735; 5,576,025; 5,599,794;

5,629,305; 5,639,478; 5,690,960; 5,703,110; 5,705,517; 5,714,504; 5,731,006;
5,879,708; 5,900,424; 5,948,773; 5,997,903; 6,017,560; 6,123,962; 6,147,103;
6,150,380; 6,166,213; 6,191,148; 5,187,340; 6,268,385; 6,262,086; 6,262,085;
6,296,875; 6,316,020; 6,328,994; 6,326,384; 6,369,085; 6,369,087; 6,380,234;
5 6,428,810; 6,444,689; and 6,462,0577.

Compositions of the invention comprise at least one polymeric benzimidazole. In one embodiment, at least one polymeric benzimidazole is present in a composition of the invention in an amount corresponding to a benzimidazole moiety weight of about 0.0005% to about 80%, about 0.005% to
10 about 70%, about 0.05% to about 60%, or about 0.5% to about 50%, by weight of the composition.

In another embodiment, the benzimidazole moiety itself (without regard for the weight of the polymer portion) is present in a composition of the invention in a total amount of about 1 to about 2000 mg, about 2.5 to about 1700 mg, about 5 to
15 about 1500 mg, about 7.5 to about 1250 mg, about 7.5 to about 1000 mg, about 7.5 to about 750 mg, about 7.5 to about 500 mg, or about 7.5 mg to about 250 mg, for example about 5 mg, about 6 mg, about 7 mg, about 8 mg, about 9 mg, about 10 mg, about 11 mg, about 12 mg, about 13 mg, about 14 mg, about 15 mg, about 16 mg, about 17 mg, about 18 mg, about 19 mg, about 20 mg, about 21 mg, about 22
20 mg, about 23 mg, about 24 mg, about 25 mg, about 26 mg, about 27 mg, about 28 mg, about 29 mg, about 30 mg, about 40 mg, about 41 mg, about 42 mg, about 43 mg, about 44 mg, about 45 mg, about 46 mg, about 47 mg, about 48 mg, about 49 mg, about 50 mg, about 51 mg, about 52 mg, about 53 mg, about 54 mg, about 55 mg about 56 mg, about 57 mg, about 58 mg, about 59 mg, about 60 mg, about 61
25 mg, about 62 mg, about 63 mg, about 64 mg, about 65 mg, about 66 mg, about 67 mg, about 68 mg, about 69 mg, about 70 mg, about 71 mg, about 72 mg, about 73 mg, about 74 mg, about 75 mg, about 77 mg, about 78 mg, about 79 mg, about 80 mg, about 81 mg, about 82 mg, about 83 mg, about 84 mg, about 85 mg, about 86 mg, about 87 mg, about 88 mg, about 89 mg, about 90 mg, about 91 mg, about 92
30 mg, about 93 mg, about 94 mg, about 95 mg, about 96 mg, about 97 mg, about 98 mg, about 99 mg, about 100 mg, about 100 mg, about 110 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about

210 mg, about 220 mg, about 230 mg, about 240 mg, about 150 mg, about 175 mg, about 200 mg, about 250 mg, about 260 mg, about 270 mg, about 280 mg, about 290 mg, about 300 mg, about 350 mg, about 400 mg, 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 750 mg, about 800 mg, 5 about 850 mg, about 900 mg, about 950 mg, about 1,000 mg, about 1,100 mg, about 1,200 mg, about 1,300 mg, about 1,400 mg, about 1,500 mg, about 1,600 mg, about 1,700 mg, about 1,800 mg, about 1,900 mg, or about 2,000 mg of benzimidazole moiety.

In another embodiment, compositions of the invention comprise at least one 10 polymeric benzimidazole in an amount of about 0.0005% to about 80%, about 0.005% to about 70%, about 0.05% to about 60%, or about 0.5% to about 50%, by weight of the composition.


In yet another embodiment, at least one polymeric benzimidazole is present in the composition in a total amount of about 5 to about 2500 mg, about 7.5 to 15 about 2000 mg, about 10 to about 1500 mg, or about 15 to about 1000 mg, for example about 5 mg, about 6 mg, about 7 mg, about 8 mg, about 9 mg, about 10 mg, about 11 mg, about 12 mg, about 13 mg, about 14 mg, about 15 mg, about 16 mg, about 17 mg, about 18 mg, about 19 mg, about 20 mg, about 21 mg, about 22 mg, about 23 mg, about 24 mg, about 25 mg, about 26 mg, about 27 mg, about 28 20 mg, about 29 mg, about 30 mg, about 40 mg, about 41 mg, about 42 mg, about 43 mg, about 44 mg, about 45 mg, about 46 mg, about 47 mg, about 48 mg, about 49 mg, about 50 mg, about 51 mg, about 52 mg, about 53 mg, about 54 mg, about 55 mg about 56 mg, about 57 mg, about 58 mg, about 59 mg, about 60 mg, about 61 mg, about 62 mg, about 63 mg, about 64 mg, about 65 mg, about 66 mg, about 67 25 mg, about 68 mg, about 69 mg, about 70 mg, about 71 mg, about 72 mg, about 73 mg, about 74 mg, about 75 mg, about 77 mg, about 78 mg, about 79 mg, about 80 mg, about 81 mg, about 82 mg, about 83 mg, about 84 mg, about 85 mg, about 86 mg, about 87 mg, about 88 mg, about 89 mg, about 90 mg, about 91 mg, about 92 mg, about 93 mg, about 94 mg, about 95 mg, about 96 mg, about 97 mg, about 98 30 mg, about 99 mg, about 100 mg, about 100 mg, about 110 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 210 mg, about 220 mg, about 230 mg, about 240 mg, about 150 mg, about 175 mg,

about 200 mg, about 250 mg, about 260 mg, about 270 mg, about 280 mg, about 290 mg, about 300 mg, about 350 mg, about 400 mg, 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 750 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, about 1,000 mg, about 1,100 mg,
5 about 1,200 mg, about 1,300 mg, about 1,400 mg, about 1,500 mg, about 1,600 mg, about 1,700 mg, about 1,800 mg, about 1,900 mg, or about 2,000 mg

In one embodiment, no portion of the polymeric benzimidazole is enteric coated. In another embodiment, at least a portion of the polymeric benzimidazole is not enteric coated. In another embodiment, at least a therapeutically effective
10 portion of the polymeric benzimidazole is not enteric coated. In another embodiment, at least about 5%, about 15%, about 20%, about 30%, about 40%, about 50% or about 60% of the polymeric benzimidazole is not enteric coated.

It will be understood that the amount of benzimidazole moiety that is administered to a subject is dependent on, *inter alia*, the type of subject, the sex,
15 age, general health, diet, and/or body weight of the subject. Illustratively, where a composition of the invention comprises a benzimidazole moiety such as omeprazole, tenatoprazole, lansoprazole, pantoprazole, rabeprazole, esomeprazole or pariprazole, and the subject is a child or a small animal (*e.g.* a dog), a relatively low amount of the benzimidazole moiety in the dose range provided herein is
20 likely to provide blood serum concentrations consistent with therapeutic effectiveness. For even smaller mammals, such as a guinea pig, even smaller amounts of benzimidazole moiety in the dosage ranges provided herein are required. Where the subject is a large animal, for example a horse, a relatively high amount of the compound in the dose range provided herein is likely to provide blood serum
25 concentrations consistent with therapeutic effectiveness.

In addition to polymeric benzimidazoles, compositions of the invention can also optionally comprise one or more non-polymeric benzimidazoles. Illustrative non-polymeric benzimidazoles are those of formula (IV):



wherein

R^1 is hydrogen, alkyl, halogen, cyano, carboxy, carboalkoxy, carboalkoxyalkyl, carbamoyl, carbamoylalkyl, hydroxy, alkoxy which is optionally
 5 fluorinated, hydroxyalkyl, trifluoromethyl, acyl, carbamoyloxy, nitro, acyloxy, aryl, aryloxy, alkylthio, or alkylsulfinyl;

R^2 is hydrogen, alkyl, acyl, acyloxy, alkoxy, amino, aralkyl, carboalkoxy, carbamoyl, alkylcarbamoyl, dialkylcarbamoyl, alkylcarbonylmethyl, alkoxy carbonylmethyl, or alkylsulfonyl;

10 R^3 and R^5 are the same or different and each is hydrogen, alkyl, alkoxy, amino, or alkoxyalkoxy;

R^4 is hydrogen, alkyl, alkoxy which may optionally be fluorinated, or alkoxyalkoxy;

Q is nitrogen, CH, or CR^1 ;

15 W is nitrogen, CH, or CR^1 ;

y is an integer of 0 through 4; and

Z is nitrogen, CH, or CR^1 ;

or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a prodrug, a polymorph, or a derivative thereof.

20 Illustrative optional non-polymeric benzimidazoles include omeprazole, tenatoprazole, lansoprazole, rabeprazole, esomeprazole (also referred to as S-omeprazole), pantoprazole, pariprazole, and nepaprazole or a free base, a free acid, or a salt, hydrate, ester, amide, enantiomer, isomer, tautomer, polymorph, prodrug, or derivative of these compounds, as well as all those substituted benzimidazoles
 25 described herein above.

Buffering Agent

Compositions of the invention comprise one or more pharmaceutically acceptable antacids or "buffering agents". Buffering agents useful in the present invention include agents possessing pharmacological activity as a weak or strong
5 base. In one embodiment, the buffering agent, when formulated with or administered substantially simultaneously with a PPI, functions to raise the pH of gastrointestinal fluid and thereby to substantially prevent or inhibit acid degradation of the PPI by gastrointestinal fluid for a period of time. In one
10 embodiment, the period of time is a period of time sufficient for allow for absorption of a therapeutically effective amount of the proton pump inhibitor.

In another embodiment, buffering agents useful in accordance with the present invention comprise a salt of a Group IA metal including, for example, a bicarbonate salt of a Group IA metal, a carbonate salt of a Group IA metal, an
15 alkaline earth metal buffering agent, an amino acid, an alkaline salt of an amino acid, an aluminum buffering agent, a calcium buffering agent, a sodium buffering agent, or a magnesium buffering agent. Other suitable buffering agents include alkali (sodium and potassium) or alkaline earth (calcium and magnesium)
carbonates, phosphates, bicarbonates, citrates, borates, acetates, phthalates, tartrates, succinates and the like, such as sodium or potassium phosphate, citrate,
20 borate, acetate, bicarbonate and carbonate.

Non-limiting examples of suitable buffering agents include aluminum, magnesium hydroxide, aluminum hydroxide/magnesium hydroxide co-precipitate, aluminum hydroxide/sodium bicarbonate co-precipitate, aluminum glycinate, calcium acetate, calcium bicarbonate, calcium borate, calcium carbonate, calcium
25 citrate, calcium gluconate, calcium glycerophosphate, calcium hydroxide, calcium lactate, calcium phthalate, calcium phosphate, calcium succinate, calcium tartrate, dibasic sodium phosphate, dipotassium hydrogen phosphate, dipotassium phosphate, disodium hydrogen phosphate, disodium succinate, dry aluminum hydroxide gel, L-arginine, magnesium acetate, magnesium aluminate, magnesium
30 borate, magnesium bicarbonate, magnesium carbonate, magnesium citrate, magnesium gluconate, magnesium hydroxide, magnesium lactate, magnesium metasilicate aluminate, magnesium oxide, magnesium phthalate, magnesium phosphate, magnesium silicate, magnesium succinate, magnesium tartrate,

potassium acetate, potassium carbonate, potassium bicarbonate, potassium borate, potassium citrate, potassium metaphosphate, potassium phthalate, potassium phosphate, potassium polyphosphate, potassium pyrophosphate, potassium succinate, potassium tartrate, sodium acetate, sodium bicarbonate, sodium borate, sodium carbonate, sodium citrate, sodium gluconate, sodium hydrogen phosphate, sodium hydroxide, sodium lactate, sodium phthalate, sodium phosphate, sodium polyphosphate, sodium pyrophosphate, sodium sesquicarbonate, sodium succinate, sodium tartrate, sodium tripolyphosphate, synthetic hydrotalcite, tetrapotassium pyrophosphate, tetrasodium pyrophosphate, tripotassium phosphate, trisodium phosphate, and trometamol. (Based in part upon the list provided in The Merck Index, Merck & Co. Rahway, N.J. (2001)). In addition, due to the ability of proteins or protein hydrolysates to react with stomach acids, they too can serve as buffering agents in the present invention. Furthermore, combinations or mixtures of the above mentioned buffering agents can be used in the pharmaceutical formulations described herein.

Buffering agents useful in the present invention also include buffering agents or combinations of buffering agents that interact with HCl (or other acids in the environment of interest) faster than the proton pump inhibitor interacts with the same acids. When placed in a liquid phase such as water, these buffering agents produce and maintain a pH greater than the pKa of the proton pump inhibitor.

In various other embodiments of the present invention, the buffering agent is present in a total amount of about 0.1 mEq/mg to about 5 mEq/mg of the benzimidazole moiety, about 0.5 mEq/mg to about 3 mEq/mg of the benzimidazole moiety, about 0.6 mEq/mg to about 2.5 mEq/mg of the benzimidazole moiety, about 0.7 mEq/mg to about 2.0 mEq/mg of the benzimidazole moiety, about 0.8 mEq/mg to about 1.8 mEq/mg of the benzimidazole moiety, about 1.0 mEq/mg to about 1.5 mEq/mg of the benzimidazole moiety. In another embodiment, the buffering agent is present in an amount of at least about 0.5 mEq/mg of the benzimidazole moiety, at least about 0.75 mEq/mg of the benzimidazole moiety, or at least about 1 mEq/mg of the benzimidazole moiety on a dry weight basis.

In another embodiment, one or more buffering agents are present in a total amount of about 0.5 mEq to about 160 mEq, about 1 mEq to about 150 mEq, about 10 mEq to about 150 mEq, about 10 mEq to about 75 mEq, about 10 mEq to about 60 mEq, or about 10 mEq to about 50 mEq. Illustratively, a composition of the

invention can comprise about 1 mEq, or about 5 mEq, or about 10 mEq, or about 15 mEq, or about 20 mEq, or about 25 mEq, or about 30 mEq, or about 35 mEq, or about 40 mEq, or about 45 mEq, or about 50 mEq, or about 60 mEq, or about 70 mEq, or about 80 mEq, or about 90 mEq, or about 100 mEq, or about 110 mEq, or about 120 mEq, or about 130 mEq, or about 140 mEq, or about 150 mEq, or about 160 mEq of buffering agent.

In yet another embodiment, one or more buffering agents are present in a total amount of at least about 10 mEq, at least about 11 mEq, at least about 12 mEq, at least about 13 mEq, at least about 14 mEq, or at least about 15 mEq.

In still another embodiment, one or more buffering agents and the benzimidazole moiety are present in a weight ratio of at least about 5:1, at least about 7:1, at least about 10:1, at least about 20:1, greater than 20:1, at least about 21:1, at least about 22:1, at least about 23:1, at least about 25:1, at least about 30:1, at least about 35:1, at least about 40:1, greater than 40:1, or at least about 45:1.

In another embodiment, the amount of buffering agent present in a composition of the invention ranges from about 200 to about 3500 mg, about 300 to about 3000 mg, about 400 to about 2500 mg, or about 500 to about 2200 mg. In other embodiments, the amount of buffering agent present in a composition of the invention is about 200 mgs, or about 300 mgs, or about 400 mgs, or about 500 mgs, or about 600 mgs, or about 700 mgs, or about 800 mgs, or about 900 mgs, or about 1000 mgs, or about 1100 mgs, or about 1200 mgs, or about 1300 mgs, or about 1400 mgs, or about 1500 mgs, or about 1600 mgs, or about 1700 mgs, or about 1800 mgs, or about 1900 mgs, or about 2000 mgs, or about 2100 mgs, or about 2200 mgs, or about 2300 mgs, or about 2400 mgs, or about 2500 mgs, or about 2600 mgs, or about 2700 mgs, or about 2800 mgs, or about 2900 mgs, or about 3000 mgs, or about 3200 mgs, or about 3500 mgs.

In another embodiment, one or more buffering agents are present in a composition of the invention in a total amount that is greater than 800 mg, for example at least about 920 mg or at least about 1000 mg.

In still another embodiment, particularly where the composition is other than a dosage form selected from the group consisting of a suspension tablet, a chewable tablet, an effervescent powder, an effervescent tablet, lozenge and/or a troche, the buffering agent and PPI are present in a weight ratio greater than 20:1, not less than about 21:1, not less than about 22:1, not less than about 23:1, not less

than about 24:1, not less than about 25:1, not less than about 26:1, not less than about 27:1, not less than about 28:1, not less than about 29:1, not less than about 30:1, not less than about 31:1, not less than about 32:1, not less than about 33:1, not less than about 34:1, not less than about 35:1, not less than about 36:1, not less than about 37:1, not less than about 38:1, not less than about 39:1, not less than about 40:1, not less than about 41:1, not less than about 42:1, not less than about 43:1, not less than about 44:1, not less than about 45:1, not less than about 46:1, not less than about 47:1, not less than about 48:1, not less than about 49:1, or not less than about 50:1.

10 In another embodiment, a composition is provided that comprises a combination of at least two non-amino acid buffering agents, wherein the combination of at least two non-amino acid buffering agents comprises substantially no aluminum hydroxide-sodium bicarbonate co-precipitate. In a related embodiment, if such a composition comprises a poly[phosphoryl/sulfon]-
15 ated carbohydrate, the weight ratio of poly[phosphoryl/sulfon]-ated carbohydrate to buffering agent is less than 1:5 (0.2), less than 1:10 (0.1) or less than 1:20 (0.05). Alternatively, the poly[phosphoryl/sulfon]-ated carbohydrate is present in the composition, if at all, in an amount less than 50 mg, less than 25 mg, less than 10 mg or less than 5 mg.

20 In other embodiments, if the pharmaceutical composition comprises an amino acid buffering agent, the total amount of amino acid buffering agent present in the pharmaceutical composition is less than about 5 mEq, or less than about 4 mEq, or less than about 3 mEq.

The phrase "amino acid buffering agent" as used herein includes amino
25 acids, amino acid salts, and amino acid alkali salts including: glycine, alanine, threonine, isoleucine, valine, phenylalanine, glutamic acid, asparaginic acid, lysine, aluminum glycinate and/or lysine glutamic acid salt, glycine hydrochloride, L-alanine, DL-alanine, L-threonine, DL-threonine, L-isoleucine, L-valine, L-phenylalanine, L-glutamic acid, L-glutamic acid hydrochloride, L-glutamic acid
30 sodium salt, L-asparaginic acid, L-asparaginic acid sodium salt, L-lysine and L-lysine-L-glutamic acid salt. The term "non-amino acid buffering agent" herein includes buffering agents as defined hereinabove but does not include amino acid buffering agents.

In another embodiment, a composition of the invention comprises at least one non-amino acid buffering agent wherein the non-amino acid buffering agent is present in the composition in a total amount greater than 800 mg. In a related embodiment, if such a composition comprises a poly[phosphoryl/sulfon]-ated carbohydrate, the weight ratio of poly[phosphoryl/sulfon]-ated carbohydrate to buffering agent is less than 1:5 (0.2), less than 1:10 (0.1) or less than 1:20 (0.05). Alternatively, the poly[phosphoryl/sulfon]-ated carbohydrate is present in the composition, if at all, in an amount less than 50 mg, less than 25 mg, less than 10 mg or less than 5 mg.

In still another embodiment, a composition is provided that comprises at least one buffering agent in a total amount of at least about 10 mEq. In a related embodiment, if an amino acid buffering agent is present in the composition, at least one of the following conditions is met: (1) the weight ratio of amino acid buffering agent:proton pump inhibitor is greater than 20:1; (2) the composition comprises at least two non-amino acid buffering agents; (3) the composition comprises at least one non-amino acid buffering agent wherein the weight ratio of the at least one non-amino acid buffering agent:proton pump inhibitor is greater than 20:1; and/or (4) the weight ratio of total buffering agent:proton pump inhibitor is greater than 40:1.

In other embodiments, where two or more buffering agents are present, the two or more buffering agents comprise at least two non-amino acid buffering agents, wherein the combination of at least two non-amino acid buffering agents comprises substantially no aluminum hydroxide-sodium bicarbonate co-precipitate.

In still another embodiment, the buffering agent comprises a mixture of sodium bicarbonate, calcium carbonate, and magnesium hydroxide, wherein the sodium bicarbonate, calcium carbonate, and magnesium hydroxide are each present in an amount of about 0.1 mEq/mg proton pump inhibitor to about 5 mEq/mg of the proton pump inhibitor.

Also provided herein are pharmaceutical compositions comprising at least one soluble buffering agent. The term "soluble buffering agent" as used herein refers to an antacid that has a solubility of at least about 500 mg/mL, or at least about 300 mg/mL, or at least about 200 mg/mL, or at least about 100 mL/mL in gastrointestinal fluid or simulated gastrointestinal fluid.

In some embodiments of the present invention, the buffering agent has a defined particle size distribution. For example, in various embodiments, the D₅₀, D₇₀, D₈₅, or D₉₀ particle size of the buffering agent, by weight or by number, is no greater than about 20 µm, no greater than about 30 µm, no greater than about 40 µm, no greater than about 50 µm, no greater than about 60 µm, no greater than about 70 µm, no greater than about 80 µm, no greater than about 90 µm, no greater than about 100 µm in diameter, no greater than about 200 µm in diameter, no greater than about 300 µm in diameter, no greater than about 400 µm in diameter, or no greater than about 100 µm in diameter.

10 Protein Component

Compositions of the invention optionally comprise a protein component in addition to or instead of the buffering agent. The term “protein component” as used herein includes protein isolates, hydrolyzed proteins (protein hydrolysates) as well as protein concentrates. Also included within the definition of a protein component are peptone, tryptone, and peptides. The term “protein component” does not embrace amino acids. A protein component can replace a buffering agent in any embodiment of the invention described herein.

Compositions of the invention can comprise one or more of a protein isolate, a protein hydrolysate, a protein concentrate, peptone, tryptone, and/or peptides. A suitable protein component can be derived from any origin including plants, animals, or a combination thereof. Non-limiting examples of suitable sources of protein component include soy, corn, whey, egg, casein, fish, meat, poultry *etc.*

Protein isolate typically comprises at least about 85%, for example about 85 – 95% protein on a dry basis. Suitable protein isolates can be prepared using any suitable procedure, for example by using an alcohol wash, water wash or ionization concentration techniques that separate at least a portion of carbohydrates and fats from the protein itself.

Protein concentrate typically comprises about 50% to about 85% protein on a dry basis, for example about 60 to about 85%. Protein concentrate can be prepared using any suitable process, for example by concentrating the desired protein through high heat drying (dehydration), acid extraction or filtration to reduce the original source to a more concentrated protein.

Protein hydrolysates are protein molecules that have been lysed, typically but not exclusively with water, into smaller peptides. Protein isolates suitable for the instant invention include substantially pure protein isolate or protein isolate formulations, for example liquid or powder formulations. Non-limiting examples
5 of powder protein hydrolysate formulations include Alimentum, Nutramigen, and Pregestimil.

In one embodiment, compositions of the invention optionally comprise a protein component in a total amount of about 1% to about 95%, about 5% to about 90%, or about 10% to about 85% on a dry weight basis in the composition.

10 In another embodiment, compositions of the invention optionally comprise a protein component in a total amount of about 1 mg to about 100 g, about 1 mg to about 20 g, about 1 mg to about 10 g, about 5 mg to about 5 g, about 10 mg to about 2.5 g, about 10 mg to about 1.0 g, or about 10 mg to about 0.5 g on a dry weight basis.

15 In another embodiment, the weight ratio of protein component (if desired) to benzimidazole moiety, on a dry basis, is about 0.001 to about 1, about 0.0025 to about 0.5, or about 0.1 to about 0.05.

In another embodiment of the invention, the optional protein component has a Protein Digestibility-Corrected Amino Acid Score (PDCAAS) of at least
20 about 0.68, at least about 0.75, at least about 0.80 at least about 0.85, at least about 0.90, at least about 0.92, at least about 0.95, at least about 0.98, or about 1.

In another embodiment, the optional protein component has a Protein Digestibility-Corrected Amino Acid Score (PDCAAS) of about 0.68 to about 1,
25 about 0.80 to about 1, about 0.90 to about 1, about 0.92 to about 1 or about 0.95 to about 1.

Pharmaceutical Excipients

Compositions of the invention can, if desired, comprise one or more pharmaceutically acceptable excipients. The term "excipient" herein means any substance, not itself a therapeutic agent, used as a carrier or vehicle for delivery of
30 a therapeutic agent to a subject or added to a pharmaceutical composition to improve its handling or storage properties or to permit or facilitate formation of a unit dose of the composition. Excipients include, by way of illustration and not limitation, diluents, disintegrants, binding agents, adhesives, wetting agents,

lubricants, glidants, surface modifying agents, substances added to mask or counteract a disagreeable taste or odor, flavors, dyes, fragrances, and substances added to improve appearance of the composition.

Excipients optionally employed in compositions of the invention can be
5 solids, semi-solids, liquids or combinations thereof. Compositions of the invention containing excipients can be prepared by any known technique of pharmacy that comprises admixing an excipient with a drug or therapeutic agent.

Compositions of the invention optionally comprise one or more pharmaceutically acceptable diluents as excipients. Suitable diluents illustratively
10 include, either individually or in combination, lactose, including anhydrous lactose and lactose monohydrate; starches, including directly compressible starch and hydrolyzed starches (*e.g.*, Celutab™ and Emdex™); mannitol; sorbitol; xylitol; dextrose (*e.g.*, Cerelease™ 2000) and dextrose monohydrate; dibasic calcium phosphate dihydrate; sucrose-based diluents; confectioner's sugar; monobasic
15 calcium sulfate monohydrate; calcium sulfate dihydrate; granular calcium lactate trihydrate; dextrates; inositol; hydrolyzed cereal solids; amylose; celluloses including microcrystalline cellulose, food grade sources of α - and amorphous cellulose (*e.g.*, Rexcel™) and powdered cellulose; calcium carbonate; glycine; bentonite; polyvinylpyrrolidone; and the like. Such diluents, if present, constitute
20 in total about 5% to about 99%, about 10% to about 85%, or about 20% to about 80%, of the total weight of the composition. The diluent or diluents selected preferably exhibit suitable flow properties and, where tablets are desired, compressibility.

Lactose and microcrystalline cellulose, either individually or in
25 combination, are preferred diluents. The use of extragranular microcrystalline cellulose (that is, microcrystalline cellulose added to a wet granulated composition after a drying step) can be used to improve hardness (for tablets) and/or disintegration time.

Compositions of the invention optionally comprise one or more
30 pharmaceutically acceptable disintegrants as excipients, particularly for tablet formulations. Suitable disintegrants include, either individually or in combination, starches, including sodium starch glycolate (*e.g.*, Explotab™ of PenWest) and pregelatinized corn starches (*e.g.*, National™ 1551, National™ 1550, and Colocorn™ 1500), clays (*e.g.*, Veegum™ HV), celluloses such as purified

cellulose, microcrystalline cellulose, methylcellulose, carboxymethylcellulose and sodium carboxymethylcellulose, croscarmellose sodium (*e.g.*, Ac-Di-Sol™ of FMC), alginates, crospovidone, and gums such as agar, guar, xanthan, locust bean, karaya, pectin and tragacanth gums.

5 Disintegrants may be added at any suitable step during the preparation of the composition, particularly prior to a granulation step or during a lubrication step prior to compression. Such disintegrants, if present, constitute in total about 0.2% to about 30%, about 0.2% to about 10%, or about 0.2% to about 5%, of the total weight of the composition.

10 Croscarmellose sodium is a preferred disintegrant for tablet or capsule disintegration, and, if present, preferably constitutes about 0.2% to about 10%, about 0.2% to about 7%, or about 0.2% to about 5%, of the total weight of the composition.

Compositions of the invention optionally comprise one or more
15 pharmaceutically acceptable binding agents or adhesives as excipients, particularly for tablet formulations. Such binding agents and adhesives preferably impart sufficient cohesion to the powder being tableted to allow for normal processing operations such as sizing, lubrication, compression and packaging, but still allow the tablet to disintegrate and the composition to be absorbed upon ingestion.

20 Suitable binding agents and adhesives include, either individually or in combination, acacia; tragacanth; sucrose; gelatin; glucose; starches such as, but not limited to, pregelatinized starches (*e.g.*, National™ 1511 and National™ 1500); celluloses such as, but not limited to, methylcellulose and carmellose sodium (*e.g.*, Tylose™); alginic acid and salts of alginic acid; magnesium aluminum silicate;
25 PEG; guar gum; polysaccharide acids; bentonites; povidone, for example povidone K-15, K-30 and K-29/32; polymethacrylates; HPMC; hydroxypropylcellulose (*e.g.*, Klucel™); and ethylcellulose (*e.g.*, Ethocel™). Such binding agents and/or adhesives, if present, constitute in total about 0.5% to about 25%, about 0.75% to about 15%, or about 1% to about 10%, of the total weight of the composition.

30 Compositions of the invention optionally comprise one or more pharmaceutically acceptable wetting agents as excipients. Non-limiting examples of surfactants that can be used as wetting agents in compositions of the invention include quaternary ammonium compounds, for example benzalkonium chloride, benzethonium chloride and cetylpyridinium chloride, dioctyl sodium

sulfosuccinate, polyoxyethylene alkylphenyl ethers, for example nonoxynol 9, nonoxynol 10, and octoxynol 9, poloxamers (polyoxyethylene and polyoxypropylene block copolymers), polyoxyethylene fatty acid glycerides and oils, for example polyoxyethylene (8) caprylic/capric mono- and diglycerides (*e.g.*,
5 Labrasol™ of Gattefossé), polyoxyethylene (35) castor oil and polyoxyethylene (40) hydrogenated castor oil; polyoxyethylene alkyl ethers, for example polyoxyethylene (20) cetostearyl ether, polyoxyethylene fatty acid esters, for example polyoxyethylene (40) stearate, polyoxyethylene sorbitan esters, for example polysorbate 20 and polysorbate 80 (*e.g.*, Tween™ 80 of ICI), propylene
10 glycol fatty acid esters, for example propylene glycol laurate (*e.g.*, Lauroglycol™ of Gattefossé), sodium lauryl sulfate, fatty acids and salts thereof, for example oleic acid, sodium oleate and triethanolamine oleate, glyceryl fatty acid esters, for example glyceryl monostearate, sorbitan esters, for example sorbitan monolaurate, sorbitan monooleate, sorbitan monopalmitate and sorbitan monostearate, tyloxapol,
15 and mixtures thereof. Such wetting agents, if present, constitute in total about 0.25% to about 15%, about 0.4% to about 10%, or about 0.5% to about 5%, of the total weight of the composition.

Compositions of the invention optionally comprise one or more pharmaceutically acceptable lubricants (including anti-adherents and/or glidants)
20 as excipients. Suitable lubricants include, either individually or in combination, glyceryl behenate (*e.g.*, Compritol™ 888); stearic acid and salts thereof, including magnesium, calcium and sodium stearates; hydrogenated vegetable oils (*e.g.*, Sterotex™); colloidal silica; talc; waxes; boric acid; sodium benzoate; sodium acetate; sodium fumarate; sodium chloride; DL-leucine; PEG (*e.g.*, Carbowax™
25 4000 and Carbowax™ 6000); sodium oleate; sodium lauryl sulfate; and magnesium lauryl sulfate. Such lubricants, if present, constitute in total about 0.1% to about 10%, about 0.2% to about 8%, or about 0.25% to about 5%, of the total weight of the composition.

Magnesium stearate is a preferred lubricant used, for example, to reduce
30 friction between the equipment and granulated mixture during compression of tablet formulations.

Suitable anti-adherents include talc, cornstarch, DL-leucine, sodium lauryl sulfate and metallic stearates. Talc is a preferred anti-adherent or glidant used, for example, to reduce formulation sticking to equipment surfaces and also to reduce

static in the blend. Talc, if present, constitutes about 0.1% to about 10%, about 0.25% to about 5%, or about 0.5% to about 2%, of the total weight of the composition.

Glidants can be used to promote powder flow of a solid formulation.

- 5 Suitable glidants include colloidal silicon dioxide, starch, talc, tribasic calcium phosphate, powdered cellulose and magnesium trisilicate. Colloidal silicon dioxide is particularly preferred.

Compositions of the present invention can comprise one or more anti-foaming agents. Dimethicone is a preferred anti-foaming agent.

- 10 Compositions of the present invention can comprise one or more flavoring agents, sweetening agents, and/or colorants. Flavoring agents useful in the present invention include, without limitation, acacia syrup, alitame, anise, apple, aspartame, banana, Bavarian cream, berry, black currant, butter, butter pecan, butterscotch, calcium citrate, camphor, caramel, cherry, cherry cream, chocolate,
15 cinnamon, citrus, citrus punch, citrus cream, cocoa, coffee, cola, cool cherry, cool citrus, cyclamate, cyclamate, dextrose, eucalyptus, eugenol, fructose, fruit punch, ginger, glycyrrhetinate, glycyrrhiza (licorice) syrup, grape, grapefruit, honey, isomalt, lemon, lime, lemon cream, MagnaSweet®, maltol, mannitol, maple, menthol, mint, mint cream, mixed berry, nut, orange, peanut butter, pear,
20 peppermint, peppermint cream, Prosweet® Powder, raspberry, root beer, rum, saccharin, saffron, sorbitol, spearmint, spearmint cream, strawberry, strawberry cream, stevia, sucralose, sucrose, Swiss cream, tagatose, tangerine, thaumatin, tutti frutti, vanilla, walnut, watermelon, wild cherry, wintergreen, xylitol, and combinations thereof, for example, anise-menthol, cherry-anise, cinnamon-orange,
25 cherry-cinnamon, chocolate-mint, honey-lemon, lemon-lime, lemon-mint, menthol-eucalyptus, orange-cream, vanilla-mint, *etc.*

- Sweetening agents that can be used in the present invention include, for example, acesulfame potassium (acesulfame K), alitame, aspartame, cyclamate, cyclamate, dextrose, isomalt, MagnaSweet®, maltitol, mannitol, neohesperidine
30 DC, neotame, Prosweet® Powder, saccharin, sorbitol, stevia, sucralose, sucrose, tagatose, thaumatin, xylitol, and the like.

The foregoing excipients can have multiple roles as is known in the art. For example, starch can serve as a filler as well as a disintegrant. The classification of excipients above is not to be construed as limiting in any manner.

Pharmaceutical Dosage Forms

Compositions of the present invention can be formulated as solid, liquid or semi-solid dosage forms. In one embodiment, such compositions are in the form of discrete dose units or dosage units. The terms “dose unit” and/or “dosage unit”
5 herein refer to a portion of a pharmaceutical composition that contains an amount of a therapeutic agent suitable for a single administration to provide a therapeutic effect. Such dosage units may be administered one to a small plurality (*i.e.* 1 to about 4) of times per day, or as many times as needed to elicit a therapeutic response. A particular dosage form can be selected to accommodate any desired
10 frequency of administration to achieve a specified daily dose. Typically one dose unit, or a small plurality (*i.e.* up to about 4) of dose units, provides a sufficient amount of the active drug (*e.g.* benzimidazole moiety) to result in the desired response or effect.

Alternatively, compositions of the invention can also be formulated for
15 rectal, topical, or parenteral (*e.g.* subcutaneous, intramuscular, intravenous and intradermal or infusion) delivery.

In one embodiment, compositions of the invention are suitable for rapid onset of therapeutic effect, particularly with respect to the PPI component. In another embodiment, upon oral administration of a composition of the invention to
20 a subject, at least a therapeutically effective amount of the PPI is available for absorption in the stomach of the subject. As discussed above, most commercially available PPIs require enteric coating to prevent exposure of the PPI to gastrointestinal fluids (and consequent drug degradation) by way of pH dependent coatings. Such coating, in turn, prevents rapid PPI absorption and therapeutic
25 onset of action. Compositions of the present invention, by contrast, do not require enteric coating to maintain drug stability in gastrointestinal fluids and thereby provide for rapid absorption and onset of therapeutic effect.

In one embodiment, a single dosage unit, be it solid or liquid, comprises a therapeutically effective amount or a therapeutically and/or prophylactically
30 effective amount of PPI. The term “therapeutically effective amount” or “therapeutically and/or prophylactically effective amount” as used herein refers to an amount of compound or agent that is sufficient to elicit the required or desired therapeutic and/or prophylactic response, as the particular treatment context may

require.

It will be understood that a therapeutically and/or prophylactically effective amount of a drug for a subject is dependent *inter alia* on the body weight of the subject. A “subject” herein to which a therapeutic agent or composition thereof
5 can be administered includes a human subject of either sex and of any age, and also includes any nonhuman animal, particularly a domestic or companion animal, illustratively a cat, dog or a horse.

Solid Dosage Forms

In some embodiments, compositions of the invention are in the form of
10 solid dosage forms or units. Non-limiting examples of suitable solid dosage forms include tablets (*e.g.* suspension tablets, bite suspension tablets, rapid dispersion tablets, chewable tablets, effervescent tablets, bilayer tablets, *etc.*), caplets, capsules (*e.g.* a soft or a hard gelatin capsule), powder (*e.g.* a packaged powder, a dispensable powder or an effervescent powder), lozenges, sachets, cachets, troches,
15 pellets, granules, microgranules, encapsulated microgranules, powder aerosol formulations, or any other solid dosage form reasonably adapted for oral administration.

Tablets are an illustrative dosage form for compositions of the invention. Tablets can be prepared according to any of the many relevant, well known
20 pharmacy techniques. In one embodiment, tablets or other solid dosage forms can be prepared by processes that employ one or a combination of methods including, without limitation, (1) dry mixing, (2) direct compression, (3) milling, (4) dry or non-aqueous granulation, (5) wet granulation, or (6) fusion.

The individual steps in the wet granulation process of tablet preparation
25 typically include milling and sieving of the ingredients, dry powder mixing, wet massing, granulation and final grinding. Dry granulation involves compressing a powder mixture into a rough tablet or “slug” on a heavy-duty rotary tablet press. The slugs are then broken up into granular particles by a grinding operation, usually by passage through an oscillation granulator. The individual steps include
30 mixing of the powders, compressing (slugging) and grinding (slug reduction or granulation). Typically, no wet binder or moisture is involved in any of the steps.

In another embodiment, solid dosage forms such as tablets can be prepared by mixing a PPI with at least one buffering agent as described herein above and, if

desired, with one or more optional pharmaceutical excipient to form a substantially homogeneous preformulation blend. The preformulation blend can then be subdivided and optionally further processed (*e.g.* compressed, encapsulated, packaged, dispersed, *etc.*) into any desired dosage forms.

5 Compressed tablets can be prepared by compacting a powder or granulation composition of the invention. The term “compressed tablet” generally refers to a plain, uncoated tablet suitable for oral ingestion, prepared by a single compression or by pre-compaction tapping followed by a final compression. Tablets of the present invention may be coated or otherwise compounded to provide a dosage
10 form affording the advantage of improved handling or storage characteristics. Preferably, however, any such coating will be selected so as to not substantially delay onset of therapeutic effect of a composition of the invention upon administration to a subject. The term “suspension tablet” as used herein refers to a compressed tablet that rapidly disintegrates after placement in water.

15 In one embodiment, a composition of the invention comprises a multi-layer tablet having a core comprising a proton pump inhibitor; the core is substantially or completely surrounded by the buffering agent. In one such embodiment, the buffering agent layer completely surrounds the core. In another embodiment, the buffering agent layer partially surrounds the core. In yet another embodiment, the
20 buffering agent layer is in contact with a portion of or with all of the surface area of the core.

 In another embodiment, one or more intermediate layers exists in between the core and the buffering agent. The intermediate layers can comprise any pharmaceutically acceptable material, preferably inert and non-pH sensitive
25 coating materials such as polymer based coatings.

 In still another embodiment, compositions of the invention can be microencapsulated, for example as is described in U.S. Patent Publication No. 2005/0037070, hereby incorporated by reference herein in its entirety.

 In another embodiment, a composition of the invention comprises a proton
30 pump inhibitor and a buffering agent mixed together in powder form and optionally filled into a capsule, for example a hard or soft gelatin or HPMC capsule.

Liquid Dosage Forms

In another embodiment of the invention, compositions can be in the form of liquid dosage forms or units. Non-limiting examples of suitable liquid dosage forms include solutions, suspension, elixirs, syrups, liquid aerosol formulations, *etc.*

In one embodiment, a liquid composition comprising water, PPI and a buffering agent can be prepared. In another embodiment, compositions of the invention are in the form of a powder for suspension that can be suspended in a liquid vehicle prior to administration to a subject. While the powder for suspension itself, can be a solid dosage form of the present invention, the powder dispersed in liquid also comprises a liquid embodiment of the invention.

Generally, a liquid composition of PPI (without a buffering agent) would exhibit a very short period of stability, even when maintained under refrigerated conditions. This is particularly inconvenient in the hospital setting as fresh batches of suspension are continually required.

Suspension compositions of the invention comprise at least one PPI, a buffering agent, a liquid media (*e.g.* water, de-ionized water, *etc.*) and one or more optional pharmaceutical excipients. Such compositions, upon storage in a closed container maintained at either room temperature, refrigerated (*e.g.* about 5 -10 °C) temperature, or freezing temperature for a period of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months, exhibit at least about 90%, at least about 92.5%, at least about 95%, or at least about 97.5% of the original PPI present therein.

Storage Stability

In one embodiment, compositions of the invention are in the form of a powder for suspension that is ultimately to be suspended in a liquid vehicle prior to administration to a subject. Liquid compositions comprising an acid labile PPI suspended in a liquid vehicle, without more, would typically exhibit short periods of stability, even when maintained under refrigerated conditions. This is particularly inconvenient in the hospital setting as fresh batches of suspension are continually required. Suspension compositions of the invention are believed to exhibit improved storage stability.

Illustrative suspension compositions of the invention comprise at least one PPI, at least one buffering agent, vitamin B₁₂, water, and one or more optional

pharmaceutical excipients. Such compositions, upon storage in a closed container maintained at room temperature, refrigerated (*e.g.* about 5 to about -10 °C) temperature, or frozen for a period of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months, preferably exhibit at least about 90%, more preferably at least about 92.5%, still more preferably at least about 95%, and yet more preferably at least about 97.5% of the original benzimidazole moiety present therein.

Parietal Cell Activators

In one embodiment, a composition of the present invention can further include one or more parietal cell activators. Parietal cell activators are particularly preferred where the benzimidazole moiety is a PPI. Parietal cell activators such as chocolate, calcium and sodium bicarbonate and other alkaline substances stimulate the parietal cells and enhance the pharmacologic activity of the PPI administered. For the purposes of this application, “parietal cell activator” or “activator” shall mean any compound or mixture of compounds possessing such stimulatory effect including, but not limited to, chocolate, sodium bicarbonate, calcium (for example, calcium carbonate, calcium gluconate, calcium hydroxide, calcium acetate and calcium glycerophosphate), peppermint oil, spearmint oil, coffee, tea and colas (even if decaffeinated), caffeine, theophylline, theobromine, and amino acids (particularly aromatic amino acids such as phenylalanine and tryptophan) and combinations thereof.

Parietal cell activators, if desired, are typically present in a composition of the invention in an amount sufficient to produce the desired stimulatory effect without causing untoward side effects to patients. For example, chocolate, as raw cocoa, is administered in an amount of about 5 mg to 2.5 g per 20 mg dose of omeprazole (or equivalent pharmacologic dose of another proton pump inhibiting agent). The dose of activator administered to a subject, for example, a human, in the context of the present invention should be sufficient to result in enhanced effect of a PPI over a desired time frame.

Illustratively, the approximate effective ranges for various parietal cell activators per 20 mg dose of omeprazole (or equivalent dose of other PPI) include, Chocolate (raw cocoa) – 5 mg to 2.5 g; Sodium bicarbonate – 7 mEq to 25 mEq; Calcium carbonate – 1 mg to 1.5 g; Calcium gluconate – 1 mg to 1.5 g; Calcium lactate – 1 mg to 1.5 g; Calcium hydroxide – 1 mg to 1.5 g; Calcium acetate – 0.5

- mg to 1.5 g; Calcium glycerophosphate – 0.5 mg to 1.5 g; Peppermint oil –
(powdered form) 1 mg to 1 g; Spearmint oil – (powdered form) 1 mg to 1 g; Coffee
– 20 ml to 240 ml; Tea – 20 ml to 240 ml; Cola – 20 ml to 240 ml; Caffeine – 0.5
mg to 1.5 g; Theophylline – 0.5 mg to 1.5 g; Theobromine – 0.5 mg to 1.5g;
5 Phenylalanine – 0.5 mg to 1.5 g; and Tryptophan – 0.5 mg to 1.5 g.

Administration

- Compositions of the present invention are useful for treating and/or
preventing, *inter alia*, gastrointestinal disorders and, in particular, acid related
gastrointestinal disorders. The phrase “acid related gastrointestinal disorder” or
10 “acid related gastrointestinal disease” refers generally to a disease or disorder that
occurs due to an imbalance between acid and pepsin production on the one hand,
so-called aggressive factors, and mucous, bicarbonate, and prostaglandin
production on the other hand, so-called defensive factors.

- The term “treat” or “treatment” as used herein refers to any treatment of a
15 disorder or disease associated with a gastrointestinal disorder, and includes, but is
not limited to, preventing the disorder or disease from occurring in a subject that
may be predisposed to the disorder or disease but has not yet been diagnosed as
having the disorder or disease; inhibiting the disorder or disease, for example,
arresting the development of the disorder or disease; relieving the disorder or
20 disease, for example, causing regression of the disorder or disease; or relieving the
condition caused by the disease or disorder, for example, stopping the symptoms of
the disease or disorder.

- The term “prevent” or “prevention,” in relation to a gastrointestinal
disorder or disease, means preventing the onset of gastrointestinal disorder or
25 disease development if none had occurred, or preventing further gastrointestinal
disorder or disease development if the gastrointestinal disorder or disease was
already present.

- In mammals gastrointestinal disorders include, but are not limited to,
duodenal ulcer, gastric ulcer, acid dyspepsia, gastroesophageal reflux disease
30 (GERD), severe erosive esophagitis, poorly responsive symptomatic
gastroesophageal reflux disease, (acid reflux), heartburn, nighttime heartburn
symptoms, nocturnal acid breakthrough (NAB), and gastrointestinal pathological
hypersecretory conditions such as Zollinger Ellison Syndrome. Illustrative acid-

related gastrointestinal disorders including duodenal ulcer disease, gastric ulcer disease, gastroesophageal reflux disease (GERD), erosive esophagitis, poorly responsive symptomatic gastroesophageal reflux disease (acid reflux), pathological gastrointestinal hypersecretory disease, Zollinger Ellison Syndrome, acid
5 dyspepsia, heartburn, and/or NSAID induced ulcer.

Where the disorder is heartburn, the heartburn can be meal-related or induced, sleep-related or induced, and/or nighttime-related or induced heartburn. Sleep-related heartburn and/or nighttime-related heartburn can be caused, for example, by breakthrough gastritis between conventional doses of a therapeutic
10 agent, such as while sleeping or in the early morning hours after a night's sleep. Treatment of these conditions is accomplished by administering to a subject a gastrointestinal-disorder-effective amount (or a therapeutically-effective amount) of a pharmaceutical composition according to the present invention. A subject may be experiencing one or more of the above conditions or disorders or related
15 symptoms.

Compositions of the invention can be administered to a subject at any suitable time, for example upon waking, prior to a meal, during the day, or at night time (*e.g.* before bed). In one embodiment, a composition of the invention is useful for treating and/or preventing nighttime heartburn or nighttime heartburn
20 symptoms, nocturnal acid breakthrough (NAB), and/or for providing nighttime pH control. NAB herein refers to intragastric pH less than 4 for more than 1 hour in the overnight period.

In another embodiment, a composition of the invention is administered to a subject between about 6:00 pm and about 1:00 am, about 7:00 pm and about 1:00
25 am, about 8:00 pm and about 12:00 am, about 8:00 pm and about 11:pm, about 8:00 pm and about 10:30 pm, or about 9:00 pm and about 10:30 pm, for example at about 9:00 pm, 9:15 pm, 9:30 pm, 9:45 pm, 10:00 pm, 10:15 pm or 10:30 pm.

In another embodiment, a composition of the invention is administered to a subject within about 3 hours before or after the subject has eaten dinner, within
30 about 2 hours before or after the subject has eaten dinner, within about 1 hour before or after the subject has eaten dinner or within about 30 minutes before or after the subject has eaten dinner.

In another embodiment, a composition of the invention is administered to a subject at such a time to result in a blood serum concentration of the proton pump

inhibitor of at least about 0.1 micromolar, at least about 0.2 micromolar, at least about 0.3 micromolar, at least about 0.4 micromolar, at least about 0.5 micromolar, at least about 0.6 micromolar, at least about 0.7 micromolar, at least about 0.8 micromolar, at least about 0.9 micromolar that occurs at any time during the
5 subject's typical period of nocturnal acid breakthrough.

The term "the subject's typical period of nocturnal acid breakthrough" refers to a period of time at night during which a subject tends to experiences intragastric pH not greater than 4 for a continuous period of about one hour. This can be determined, for example, by measuring a subject's intragastric pH for 1 to 3
10 nights and calculating the beginning and ending times (of pH less than 4), or the average beginning and ending times if pH is measured during more than 1 night. For example, if a subject experiences nocturnal acid breakthrough on a first measured day starting at 1:00 am and ending at 4:00 am, and on a second measured day that subject experiences nocturnal acid breakthrough starting at 2:00 am and
15 ending at 5:00 am, the subject's typical period of nocturnal acid breakthrough could be characterized as 1:30 am to 4:30 am. Thus, a composition of the invention could be administered to the subject at such a time so as to provide a blood serum concentration of the proton pump inhibitor of at least about 0.1 micromolar, at least about 0.2 micromolar, at least about 0.3 micromolar, at least
20 about 0.4 micromolar, at least about 0.5 micromolar, at least about 0.6 micromolar, at least about 0.7 micromolar, at least about 0.8 micromolar, or at least about 0.9 micromolar at any time during the period of about 12:00 am to about 6:00 am, about 12:30 am to about 5:30 am, about 1:00 am to about 5:00 am, or about 2:00 am to about 4:00 am.

25 In another embodiment, a composition of the invention is administered to a subject at such a time to result in a blood serum concentration of the proton pump inhibitor of at least about 0.1 micromolar, at least about 0.2 micromolar, at least about 0.3 micromolar, at least about 0.4 micromolar, at least about 0.5 micromolar, at least about 0.6 micromolar, at least about 0.7 micromolar, at least about 0.8
30 micromolar, at least about 0.9 micromolar or at least about 1 micromolar continuously throughout the subject's typical period of nocturnal acid breakthrough.

In another embodiment, a composition of the invention is administered to a subject at such a time to result in a blood serum concentration of the proton pump

inhibitor of at least about 0.1 micromolar, at least about 0.2 micromolar, at least about 0.3 micromolar, at least about 0.4 micromolar, at least about 0.5 micromolar, at least about 0.6 micromolar, at least about 0.7 micromolar, at least about 0.8 micromolar, at least about 0.9 micromolar, or at least about 1 micromolar at any
5 time point during the period of about 12:00 am to about 6:00 am, about 12:30 am to about 5:30 am, about 1:00 am to about 5:00 am, or about 2:00 am to about 4:00 am.

In another embodiment, a composition of the invention is administered to a subject at such a time to result in a blood serum concentration of the proton pump
10 inhibitor of at least about 0.1 micromolar, at least about 0.2 micromolar, at least about 0.3 micromolar, at least about 0.4 micromolar, at least about 0.5 micromolar, at least about 0.6 micromolar, at least about 0.7 micromolar, at least about 0.8 micromolar, at least about 0.9 micromolar or at least about 1 micromolar continuously from about 12:00 am to about 6:00 am, from about 12:30 am to about
15 5:30 am, from about 1:00 am to about 5:00 am, or from about 2:00 am to about 4:00 am.

In another embodiment, a composition of the invention is administered to a subject at such a time to result in a blood serum concentration of the proton pump inhibitor of about 0.1 micromolar to about 5 micromolar, about 0.2 micromolar to
20 about 2.5 micromolar, or about 0.3 micromolar to about 2 micromolar continuously from about 12:00 am to about 6:00 am, from about 12:30 am to about 5:30 am, from about 1:00 am to about 5:00 am, or from about 2:00 am to about 4:00 am.

In another embodiment, a composition of the invention is administered to a
25 subject at such a time to result in a blood serum concentration of the proton pump inhibitor of about 0.1 micromolar to about 5 micromolar, about 0.2 micromolar to about 2.5 micromolar, or about 0.3 micromolar to about 2 micromolar from about 12:00 am to about 6:00 am, from about 12:30 am to about 5:30 am, at any time during the period of about 1:00 am to about 5:00 am, or from about 2:00 am to
30 about 4:00 am.

In still another embodiment, a composition of the invention is administered to a subject at such a time to result in a $C_{\max}$ of the proton pump inhibitor that occurs during the subject's typical period of nocturnal acid breakthrough.

In another embodiment, a composition of the invention is administered to a subject at such a time to result in a $C_{\max}$ of the proton pump inhibitor that occurs about 0.1 to about 5 hours, about 0.1 to about 4 hours, about 0.1 to about 3 hours, or about 0.1 to about 2 hours before the subject's typical period of nocturnal acid
5 breakthrough.

Compositions of the invention are to be administered and dosed in accordance with good medical practice, taking into account the clinical condition of the individual patient, the site and method of administration, scheduling of administration and other factors known to medical practitioners. In human
10 therapy, it is important to provide a dosage form that delivers the required therapeutic amount of the drug *in vivo*, and renders the drug bioavailable in a rapid manner. In addition to the dosage forms described herein, the dosage forms described by Phillips *et al.* in U.S. Pat. No. 6,489,346 are incorporated herein by reference.

15 The percent of intact drug that is absorbed into the bloodstream is not narrowly critical, as long as a therapeutic-disorder-effective amount, for example a gastrointestinal-disorder-effective amount of a proton pump inhibiting agent, is absorbed following administration of the pharmaceutical composition to a subject. It will be understood that the amount of proton pump inhibiting agent and/or
20 antacid that is administered to a subject is dependent on various factors including the sex, general health, diet, and/or body weight of the subject.

Illustratively, when administering a PPI to a young child or a small animal, such as a dog, a relatively low amount of the proton pump inhibitor, *e.g.*, about 1 mg to about 30 mg, will often provide blood serum concentrations consistent with
25 therapeutic effectiveness. Where the subject is an adult human or a large animal, such as a horse, achievement of a therapeutically effective blood serum concentration may require larger dosage units, for example about 10 mg, about 15 mg, about 20 mg, about 30 mg, about 40 mg, about 80 mg, or about 120 mg dose for an adult human, or about 150 mg, or about 200 mg, or about 400 mg, or about
30 800 mg, or about 1000 mg dose, or about 1500 mg dose, or about 2000 mg dose, or about 2500 mg dose, or about 3000 mg dose, or about 3200 mg dose, or about 3500 mg dose for an adult horse.

In various other embodiments of the present invention, the amount of proton pump inhibitor administered to a subject is about 1-2 mg/Kg of body

weight, illustratively about 0.5 mg/Kg of body weight, about 1 mg/Kg of body weight, about 1.5 mg/Kg of body weight, or about 2 mg/Kg of body weight.

Treatment dosages generally may be titrated to optimize safety and efficacy. Typically, dosage-effect relationships from *in vitro* and/or *in vivo* tests
5 initially can provide useful guidance on the proper doses for subject administration. Studies in animal models generally may be used for guidance regarding effective dosages for treatment of gastrointestinal disorders or diseases in accordance with the present invention. In terms of treatment protocols, it should be appreciated that the dosage to be administered will depend on several factors,
10 including the particular agent that is administered, the route chosen for administration, the condition of the particular subject.

In another embodiment of the present invention, the composition is administered to a subject in an amount sufficient to achieve a measurable serum concentration of a non-acid degraded or non-acid reacted proton pump inhibitor
15 greater than about 100 ng/ml within about 30 minutes or about 15 minutes or about 10 minutes after administration of the composition.

In another embodiment of the present invention, the composition is administered to a subject in an amount sufficient to achieve a measurable serum concentration of the proton pump inhibitor greater than about 150 ng/ml within
20 about 15 minutes and to maintain a serum concentration of the proton pump inhibitor of greater than about 150 ng/ml from about 15 minutes to about 1 hour after administration of the composition. In yet another embodiment of the present invention, the composition is administered to the subject in an amount sufficient to achieve a measurable serum concentration of the proton pump inhibitor greater
25 than about 250 ng/ml within about 15 minutes and to maintain a serum concentration of the proton pump inhibiting agent of greater than about 150 ng/ml from about 15 minutes to about 1 hour after administration of the composition.

In another embodiment of the present invention, the composition is administered to the subject in an amount sufficient to achieve a measurable serum
30 concentration of the proton pump inhibitor greater than about 350 ng/ml within about 15 minutes and to maintain a serum concentration of the proton pump inhibitor of greater than about 150 ng/ml from about 15 minutes to about 1 hour after administration of the composition. In another embodiment, the composition is administered to the subject in an amount sufficient to achieve a measurable

serum concentration of the proton pump inhibiting agent greater than about 450 ng/ml within about 15 minutes and to maintain a serum concentration of the proton pump inhibiting agent of greater than about 150 ng/ml from about 15 minutes to about 1 hour after administration of the composition.

- 5 In another embodiment, the composition is administered to the subject in an amount sufficient to achieve a measurable serum concentration of the proton pump inhibitor greater than about 150 ng/ml within about 30 minutes and to maintain a serum concentration of the proton pump inhibitor of greater than about 150 ng/ml from about 30 minutes to about 1 hour after administration of the composition. In
- 10 yet another embodiment of the present invention, the composition is administered to the subject in an amount to achieve a measurable serum concentration of the proton pump inhibitor greater than about 250 ng/ml within about 30 minutes and to maintain a serum concentration of the proton pump inhibitor of greater than about 150 ng/ml from about 30 minutes to about 1 hour after administration of the
- 15 composition. In another embodiment of the present invention, the composition is administered to the subject in an amount sufficient to achieve a measurable serum concentration of the proton pump inhibitor greater than about 350 ng/ml within about 30 minutes and to maintain a serum concentration of the proton pump inhibitor of greater than about 150 ng/ml from about 30 minutes to about 1 hour
- 20 after administration of the composition. In another embodiment of the present invention, the composition is administered to the subject in an amount sufficient to achieve a measurable serum concentration of the proton pump inhibitor greater than about 450 ng/ml within about 30 minutes and to maintain a serum concentration of the proton pump inhibiting agent of greater than about 150 ng/ml
- 25 from about 30 minutes to about 1 hour after administration of the composition.

- In still another embodiment of the present invention, the composition is administered to the subject in an amount sufficient to achieve a measurable serum concentration of a non-acid degraded or non-acid reacted proton pump inhibitor greater than about 500 ng/ml within about 1 hour after administration of the
- 30 composition. In yet another embodiment of the present invention, the composition is administered to the subject in an amount to achieve a measurable serum concentration of a non-acid degraded or non-acid reacted proton pump inhibitor greater than about 300 ng/ml within about 45 minutes after administration of the composition.

Contemplated compositions of the present invention provide a therapeutic effect as proton pump inhibiting agent medications over an interval of about 5 minutes to about 24 hours after administration, enabling, for example, once-a-day, twice-a-day, or three times a day administration if desired.

5 In another embodiment, upon oral administration of a composition of the invention to a plurality of fasted human subjects, the subjects exhibit an average $T_{\max}$ of PPI within about 30 seconds to about 90 minutes, within about 1 minute to about 80 minutes, within about 5 minutes to about 60 minutes, within about 10 minutes to about 50 minutes, or within about 15 minutes to about 45 minutes.

10 In still another embodiment, upon administration of a composition of the invention to a plurality of fasted human subjects, the subjects exhibit an average plasma concentration of the PPI of at least about 0.1 $\mu\text{g/ml}$, at least about 0.15 $\mu\text{g/ml}$, at least about 0.2 $\mu\text{g/ml}$, at least about 0.3 $\mu\text{g/ml}$, at least about 0.4 $\mu\text{g/ml}$, at least about 0.5 $\mu\text{g/ml}$, at least about 0.6 $\mu\text{g/ml}$, at least about 0.7 $\mu\text{g/ml}$, at least
15 about 0.8 $\mu\text{g/ml}$, at least about 0.9 $\mu\text{g/ml}$, at least about 1 $\mu\text{g/ml}$, at least about 1.5 $\mu\text{g/ml}$, or at least about 2.0 $\mu\text{g/ml}$ at any time within about 90 minutes, within about 75 minutes, within about 60 minutes, within about 55 minutes, within about 50 minutes, within about 45 minutes, within about 40 minutes, within about 35 minutes, within about 30 minutes, within about 25 minutes, within about 20
20 minutes, within about 17 minutes, within about 15 minutes, within about 12 minutes, or within about 10 minutes after administration.

 In yet another embodiment, upon administration of a composition of the invention to a plurality of fasted human subjects, the subjects exhibit a plasma concentration of PPI of at least about 0.1 $\mu\text{g/ml}$, at least about 0.15 $\mu\text{g/ml}$, at least
25 about 0.2 $\mu\text{g/ml}$, at least about 0.3 $\mu\text{g/ml}$, at least about 0.4 $\mu\text{g/ml}$, at least about 0.5 $\mu\text{g/ml}$, at least about 0.6 $\mu\text{g/ml}$, at least about 0.7 $\mu\text{g/ml}$, at least about 0.8 $\mu\text{g/ml}$, at least about 0.9 $\mu\text{g/ml}$, at least about 1.0 $\mu\text{g/ml}$, at least about 1.5 $\mu\text{g/ml}$ or at least about 2.0 $\mu\text{g/ml}$, maintained from at latest about 15 minutes to at earliest about 60 minutes after administration, preferably at latest about 15 minutes after
30 administration to at earliest about 90 minutes after administration, more preferably at latest about 15 minutes to at earliest about 120 minutes after administration, and still more preferably at latest about 15 minutes to at earliest about 180 minutes after administration.

In another embodiment, upon administration of a composition of the invention to a plurality of fasted human subjects, the subjects exhibit at least one of: a mean $C_{\max}$ of PPI of about 500 $\mu\text{g/ml}$ to about 2000 $\mu\text{g/ml}$, about 600 $\mu\text{g/ml}$ to about 1900 $\mu\text{g/ml}$, or about 700 ng/ml to about 1800 $\mu\text{g/ml}$; a mean $T_{\max}$ of PPI of about 0.15 to about 2 hours, about 0.25 to about 1.75 hours, or about 0.3 hours to about 1 hour; and/or a mean $\text{AUC}_{(0-\infty)}$ of PPI of about 1000 to about 3000 $\mu\text{g}\cdot\text{hr/ml}$, about 1500 to about 2700 $\mu\text{g}\cdot\text{hr/ml}$, or about 1700 to about 2500 $\mu\text{g}\cdot\text{hr/ml}$.

In another embodiment, upon administration of a composition of the invention to a plurality of fasted adult human subjects, the subjects exhibit: a mean $C_{\max}$ of PPI of about 500 $\mu\text{g/ml}$ to about 2000 $\mu\text{g/ml}$, about 600 $\mu\text{g/ml}$ to about 1900 $\mu\text{g/ml}$, or about 700 $\mu\text{g/ml}$ to about 1800 $\mu\text{g/ml}$; a mean $T_{\max}$ of PPI of about 0.15 to about 2 hours, about 0.25 to about 1.75 hours, or about 0.3 hours to about 1 hour; and a mean $\text{AUC}_{(0-\infty)}$ of PPI of about 1000 to about 3000 $\text{ng}\cdot\text{hr/ml}$, about 1500 to about 2700 $\text{ng}\cdot\text{hr/ml}$, or about 1700 to about 2500 $\text{ng}\cdot\text{hr/ml}$.

Gastrointestinal Disorders

Compositions of the present invention are particularly useful for treating and/or preventing gastrointestinal disorders. Illustratively gastrointestinal disorders are acid-caused gastrointestinal disorders including duodenal ulcer disease, gastric ulcer disease, gastroesophageal reflux disease (GERD), erosive esophagitis, poorly responsive symptomatic gastroesophageal reflux disease, pathological gastrointestinal hypersecretory disease, Zollinger Ellison Syndrome, acid dyspepsia, heartburn, nocturnal acid breakthrough (NAB), and/or NSAID induced ulcer.

Where the disorder is heartburn, the heartburn can be meal related or induced, sleep related or induced, and/or nighttime related or induced. Sleep related or induced heartburn and/or nighttime related or induced heartburn can be caused, for example, by breakthrough gastritis between conventional doses of a therapeutic agent, such as while sleeping or in the early morning hours after a night's sleep. Treatment of these conditions is accomplished by administering to a subject a gastrointestinal-disorder-effective amount (or a therapeutically-effective amount) of a pharmaceutical composition according to the present invention. A subject may be experiencing one or more of these conditions or disorders.

The term "treat" or "treatment" as used herein refers to any treatment of a disorder or disease associated with a gastrointestinal disorder, and includes, but is not limited to, preventing the disorder or disease from occurring in a subject that may be predisposed to the disorder or disease but has not yet been diagnosed as

5 having the disorder or disease; inhibiting the disorder or disease, for example, arresting the development of the disorder or disease; relieving the disorder or disease, for example, causing regression of the disorder or disease; or relieving the condition caused by the disease or disorder, for example, stopping the symptoms of the disease or disorder.

10 The term "prevent" or "prevention," in relation to a gastrointestinal disorder or disease, means preventing the onset of gastrointestinal disorder or disease development if none had occurred, or preventing further gastrointestinal disorder or disease development if the gastrointestinal disorder or disease was already present.

15 Those skilled in the art will readily appreciate that numerous other embodiments, modifications and equivalents are contemplated and encompassed by the disclosure of the present invention.

All U.S. Patents and published U.S. patent applications listed herein are hereby incorporated by reference in their entirety. All patents, patent applications

20 and publications referenced herein are hereby incorporated by reference herein to the fullest extent allowed under the law.

EXAMPLES

Example 1

The following formulations, shown in Table 1, can be prepared as tablets, capsules, or other solid dosage forms..

5

Table 1


	Composition No.							
Component	1	2	3	4	5	6	7	8
Polymerized Omeprazole (mg)	10	20	40	60				
Polymerized Lansoprazole (mg)					10	15	30	45
Sodium Bicarbonate (mg)	900	800	600	1600	1200	800	600	1400
Calcium Carbonate (mg)				100		200		
Sodium Carbonate (mg)			600				300	
Magnesium Hydroxide (mg)	300	400				100		

CLAIMS

What is claimed is:

1. A pharmaceutical composition comprising:




(a) at least one polymeric benzimidazole compound of formula (II):



wherein,

R_7 , R_8 , R_9 , and R_{10} , are independently H or CH_3 ;


U is




R_{11} is H, CH_3 , C_2H_5 , or CONH_2 ;

Y and V are independently OH or NH_2 ,

E is



B is a benzimidazole moiety of formula (III):



wherein,

R^1 is hydrogen, alkyl, halogen, cyano, carboxy, carboalkoxy, carboalkoxyalkyl, carbamoyl, carbamoylalkyl, hydroxy, alkoxy which may optionally be fluorinated, hydroxyalkyl, trifluoromethyl, acyl, carbamoyloxy, nitro, acyloxy, aryl, aryloxy, alkylthio, or alkylsulfinyl;

R^3 and R^5 are the same or different and each is hydrogen, alkyl, alkoxy, amino, or alkoxyalkoxy;

R^4 is hydrogen, alkyl, alkoxy which may optionally be fluorinated, or alkoxyalkoxy;

Q is nitrogen, CH, or CR^1 ;

W is nitrogen, CH, or CR^1 ;


y is an integer of 0 through 4;

Z is nitrogen, CH, or CR^1 ;

or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a polymorph, a prodrug, or a derivative of a compound of formula (III); and

(b) at least one buffering agent in an amount of about 0.1 mEq to about 10 mEq per mg of the benzimidazole moiety (B).

2. The composition of claim 1, wherein B is a benzimidazole moiety of formula (III):



wherein,

R^1 is hydrogen or alkoxy which may optionally be fluorinated;

R^3 and R^5 are the same or different and each is hydrogen, alkyl, alkoxy, amino, or alkoxyalkoxy;

R^4 is hydrogen, alkyl, alkoxy which may optionally be fluorinated, or alkoxyalkoxy;

Q is CH;

W is CH;

y is an integer of 0 through 4;

Z is nitrogen;


or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a polymorph, a prodrug, or a derivative of a compound of formula (III).

3. The composition of claim 1, wherein the at least one buffering agent is selected from the group consisting of a bicarbonate salt of a Group IA metal, a carbonate salt of a Group IA metal, an alkaline earth metal buffering agent, an aluminum buffering agent, a calcium buffering agent, a magnesium buffering agent, an amino acid, an alkaline salt of an amino acid, and mixtures thereof.
4. The composition of claim 3, wherein the at least one buffering agent is selected from the group consisting of aluminum hydroxide, aluminum hydroxide/magnesium carbonate, aluminum hydroxide/magnesium carbonate/calcium carbonate co-precipitate, aluminum magnesium hydroxide, aluminum hydroxide/magnesium hydroxide co-precipitate, aluminum hydroxide/sodium bicarbonate coprecipitate, aluminum glycinate, calcium acetate, calcium bicarbonate, calcium borate, calcium

- carbonate, calcium citrate, calcium chloride, calcium gluconate, calcium glycerophosphate, calcium hydroxide, calcium lactate, calcium phthalate, calcium phosphate, calcium succinate, calcium tartrate, dibasic sodium phosphate, dipotassium hydrogen phosphate, dipotassium phosphate, disodium hydrogen phosphate, disodium succinate, dry aluminum hydroxide gel, L-arginine, magnesium acetate, magnesium aluminate, magnesium borate, magnesium bicarbonate, magnesium carbonate, magnesium citrate, magnesium gluconate, magnesium hydroxide, magnesium lactate, magnesium metasilicate aluminate, magnesium oxide, magnesium phthalate, magnesium phosphate, magnesium silicate, magnesium succinate, magnesium tartrate, potassium acetate, potassium carbonate, potassium bicarbonate, potassium borate, potassium citrate, potassium metaphosphate, potassium phthalate, potassium phosphate, potassium polyphosphate, potassium pyrophosphate, potassium succinate, potassium tartrate, sodium acetate, sodium bicarbonate, sodium borate, sodium carbonate, sodium citrate, sodium gluconate, sodium dihydrogen phosphate, sodium hydrogen phosphate, sodium hydroxide, sodium lactate, sodium phthalate, sodium phosphate, sodium polyphosphate, sodium pyrophosphate, sodium sesquicarbonate, sodium succinate, sodium tartrate, sodium tripolyphosphate, synthetic hydrotalcite, tetrapotassium pyrophosphate, tetrasodium pyrophosphate, trihydroxymethylaminomethane, tripotassium phosphate, trisodium phosphate, and trometamol, and mixtures thereof.
5. The composition of claim 1 wherein the at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 0.0005% to about 80% of benzimidazole moiety, by total weight of the composition
6. The composition of claim 1 wherein the at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 0.5% to about 50% of benzimidazole moiety, by total weight of the composition.

7. The composition of claim 1 wherein at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 1 to about 2000 mg of benzimidazole moiety.
8. The composition of claim 1 wherein at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 7.5 to about 250 mg of benzimidazole moiety.
9. The composition of claim 1 wherein the at least one buffering agent is present in a composition of the invention in a total amount of about 0.5 mEq to about 150 mEq.
10. The composition of claim 1 wherein the at least one buffering agent is present in a composition of the invention in a total amount of about 1 mEq to about 110 mEq.
11. The composition of claim 1 wherein the at least one buffering agent is present in a composition of the invention in a total amount of about 0.05 mEq to about 10 mEq per mg of benzimidazole moiety.
12. The composition of claim 1 wherein the at least one buffering agent is present in a composition of the invention in a total amount of about 0.2 mEq to about 2.5 mEq per mg of benzimidazole moiety.
13. The composition of claim 1 wherein the at least one buffering agent and the benzimidazole moiety are present in a weight ratio of at least about 5:1.
14. The composition of claim 1 wherein the at least one buffering agent and the benzimidazole moiety are present in a weight ratio of at least about 20:1.
15. The composition of claim 1, wherein the composition is a dosage form selected from the group consisting of a tablet, a suspension tablet, a bite suspension tablet, a rapid dispersion tablet, a chewable tablet, an effervescent tablet, a bilayer tablet, and a tablet-in-a-tablet a powder, a filled capsule, a caplet, a lozenge, a sachet, a cachet, a troche, a pellet, a granule, a solution and a suspension.
16. The composition of claim 15, wherein the capsule is selected from the group consisting of a soft gelatin capsule and a hard gelatin capsule.




17. The composition of claim 1, wherein the composition further comprises a pharmaceutically acceptable excipient selected from the group consisting of a binder, a filling agent, a suspending agent, a flavoring agent, a sweetening agent, a disintegrant, a flow aid, a lubricant, an adjuvant, a colorant, a diluent, a solubilizer, a moistening agent, a stabilizer, a wetting agent, an anti-adherent, a glidant, a preservative, a parietal cell activator, an anti-foaming agent, an antioxidant, a chelating agent, an antifungal agent, an antibacterial agent, an isotonic agent, and mixtures thereof.
18. A pharmaceutical composition comprising:
- (a) at least one polymeric benzimidazole compound of formula (II):



wherein,

R_7 , R_8 , R_9 , and R_{10} , are independently H or CH_3 ;


U is



R_{11} is H, CH_3 , C_2H_5 , or $CONH_2$;


Y and V are independently OH or NH_2 ,

E is



5

B is a benzimidazole moiety of formula (III):



wherein,

R^1 is hydrogen, alkyl, halogen, cyano, carboxy, carboalkoxy, carboalkoxyalkyl, carbamoyl, carbamoylalkyl, hydroxy, alkoxy which may optionally be fluorinated, hydroxyalkyl, trifluoromethyl, acyl, carbamoyloxy, nitro, acyloxy, aryl, aryloxy, alkylthio, or alkylsulfinyl;

10

R^3 and R^5 are the same or different and each is hydrogen, alkyl, alkoxy, amino, or alkoxyalkoxy;

R^4 is hydrogen, alkyl, alkoxy which may optionally be fluorinated, or alkoxyalkoxy;

15

Q is nitrogen, CH, or CR^1 ;

W is nitrogen, CH, or CR^1 ;

y is an integer of 0 through 4;

Z is nitrogen, CH, or CR^1 ;


20

or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a polymorph, a prodrug, or a derivative of a compound of formula (III); and

(b) a protein component.

19. The composition of claim 18, wherein B is a benzimidazole moiety of formula (III):

25



wherein,

R^1 is hydrogen or alkoxy which may optionally be fluorinated;

R^3 and R^5 are the same or different and each is hydrogen, alkyl, alkoxy, amino, or alkoxyalkoxy;

R^4 is hydrogen, alkyl, alkoxy which may optionally be fluorinated, or alkoxyalkoxy;

Q is CH;

W is CH;

y is an integer of 0 through 4;

Z is nitrogen;

or a salt, an ester, a hydrate, an amide, an enantiomer, an isomer, a tautomer, a polymorph, a prodrug, or a derivative of a compound of formula (III).

20. The composition of claim 18, wherein the at least one buffering agent is selected from the group consisting of a bicarbonate salt of a Group IA metal, a carbonate salt of a Group IA metal, an alkaline earth metal buffering agent, an aluminum buffering agent, a calcium buffering agent, a magnesium buffering agent, an amino acid, an alkaline salt of an amino acid, and mixtures thereof.
21. The composition of claim 18, wherein the protein component is selected from protein isolates, hydrolyzed proteins (protein hydrolysates) as well as protein concentrates and mixtures thereof.
22. The composition of claim 18 wherein the at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 0.0005% to about 80% of benzimidazole moiety, by total weight of the composition

23. The composition of claim 18 wherein the at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 0.5% to about 50% of benzimidazole moiety, by total weight of the composition.
- 5 24. The composition of claim 18 wherein at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to about 1 to about 2000 mg of benzimidazole moiety.
25. The composition of claim 18 wherein at least one polymeric benzimidazole compound is present in the composition in a total amount corresponding to
10 about 7.5 to about 250 mg of benzimidazole moiety.
26. The composition of claim 18 wherein the protein component is present in a composition of the invention in a total amount of about 1 % to about 95% on a dry weight basis.
27. The composition of claim 18 wherein the at least one buffering agent is
15 present in a composition of the invention in a total amount of about 110% to about 85% on a dry weight basis.
28. The composition of claim 18, wherein the composition is a dosage form selected from the group consisting of a tablet, a suspension tablet, a bite suspension tablet, a rapid dispersion tablet, a chewable tablet, an
20 effervescent tablet, a bilayer tablet, and a tablet-in-a-tablet a powder, a filled capsule, a caplet, a lozenge, a sachet, a cachet, a troche, a pellet, a granule, a solution and a suspension.
29. The composition of claim 18, wherein the capsule is selected from the group consisting of a soft gelatin capsule and a hard gelatin capsule.
- 25 30. The composition of claim 18, wherein the composition further comprises a pharmaceutically acceptable excipient selected from the group consisting of a binder, a filling agent, a suspending agent, a flavoring agent, a sweetening agent, a disintegrant, a flow aid, a lubricant, an adjuvant, a colorant, a diluent, a solubilizer, a moistening agent, a stabilizer, a wetting agent, an
30 anti-adherent, a glidant, a preservative, a parietal cell activator, an anti-foaming agent, an antioxidant, a chelating agent, an antifungal agent, an antibacterial agent, an isotonic agent, and mixtures thereof.

31. A method of treating a gastrointestinal disorder in a subject in need thereof, the method comprising administering to the subject a gastrointestinal disorder-effective amount of a composition of claim 1 or claim 18.
32. The method of claim 31, wherein the gastrointestinal disorder is a duodenal ulcer disease, a gastric ulcer disease, a gastroesophageal reflux disease, an erosive esophagitis, a poorly responsive symptomatic gastroesophageal reflux disease, a pathological gastrointestinal hypersecretory disease, Zollinger Ellison Syndrome, acid dyspepsia, heartburn, nocturnal acid breakthrough, an esophageal disorder, a non-erosive reflux disorder, or an NSAID induced ulcer.
33. The method of claim 32, wherein the gastrointestinal disorder is heartburn.
34. Use of a pharmaceutical composition of any one of claims 1 – 30 in the manufacture of a medicament for use for the treatment of a gastric acid related disorder in a patient in need thereof.
35. Use as claimed in claim 36 wherein the gastric acid related disorder is selected from duodenal ulcer disease, gastric ulcer disease, gastroesophageal reflux disease (GERD), erosive esophagitis, poorly responsive symptomatic gastroesophageal reflux disease, pathological gastrointestinal hypersecretory disease, Zollinger Ellison Syndrome, acid dyspepsia, heartburn, nocturnal acid breakthrough (NAB), and/or NSAID induced ulcer.