

(19) **DANMARK**



Patent- og  
Varemærkestyrelsen

(10) **DK/EP 4073098 T5**

(12) **Rettet oversættelse af  
europæisk patentskrift**

- 
- (51) Int.Cl.: **C 07 K 14/62 (2006.01)** **A 61 P 3/08 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-07-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-09-27**
- (86) Europæisk ansøgning nr.: **20830444.4**
- (86) Europæisk indleveringsdag: **2020-12-07**
- (87) Den europæiske ansøgnings publiceringsdag: **2022-10-19**
- (86) International ansøgning nr.: **US2020063673**
- (87) Internationalt publikationsnr.: **WO2021126584**
- (30) Prioritet: **2019-12-19 US 201962950803 P** **2020-03-12 US 202062988441 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Akston Biosciences Corporation, 100 Cummings Center, Ste 454C, Beverly, Massachusetts 01915, USA**
- (72) Opfinder: **LANCASTER, Thomas, M., 166 Larch Row, Wenham, MA 01984, USA**  
**ZION, Todd, C., 72 Nanepashemet Street, Marblehead, MA 01945, USA**
- (74) Fuldmægtig i Danmark: **Murgitroyd & Company, Mannerheimvägen 12 B, 5:e våningen, FI-00100 Helsingfors, Finland**
- (54) Benævnelse:  
**Ultralangtidsvirkende insulin-Fc-fusionsproteiner og fremgangsmåder til anvendelse**
- (56) Fremdragne publikationer:  
**WO-A1-2016/178905**  
**WO-A1-2018/107117**  
**WO-A1-2020/006529**  
**XINHUA WANG ET AL: "IgG Fc engineering to modulate antibody effector functions", PROTEIN & CELL, vol. 9, no. 1, 6 October 2017 (2017-10-06), pages 63-73, XP055457296, Beijing, CN ISSN: 1674-800X, DOI: 10.1007/s13238-017-0473-8**  
**Chunpeng Yang ET AL: "Engineering of Fc Fragments with Optimized Physicochemical Properties Implying Improvement of Clinical Potentials for Fc-Based Therapeutics", Frontiers in Immunology, vol. 8, 8 January 2018 (2018-01-08), pages 1-14, XP055671009, DOI: 10.3389/fimmu.2017.01860**  
**SAZINSKY STEPHEN L ET AL: "Aglycosylated immunoglobulin G1 variants productively engage activating Fc receptors", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, NATIONAL ACADEMY OF SCIENCES, vol. 105, no. 51, 23 December 2008 (2008-12-23), pages 20167-20172, XP009143223, ISSN: 0027-8424, DOI: 10.1073/PNAS.0809257105**



**Description****TECHNICAL FIELD**

5     **[0001]** The present technology relates to compositions of insulin-Fc fusion proteins and their use to treat diabetes in humans.

**BACKGROUND**

10    **[0002]** The following description of the background of the present technology is provided simply as an aid in understanding the present technology and is not admitted to describe or constitute prior art to the present technology.

15    **[0003]** Diabetes is a chronic condition characterized by an insulin deficiency and/or ineffective use of insulin. Diabetics that have an absolute deficiency of insulin are categorized as having type 1 or insulin-dependent diabetes mellitus (IDDM). Type 1 diabetics are thought to have a genetic predisposition combined with immunologic destruction of the insulin-producing  $\beta$ -cells of the pancreas. In comparison, diabetics that can still produce some insulin but have a relative deficiency due to insulin resistance or other dysfunction, are classified as having type 2 or non-insulin-dependent diabetes mellitus (NIDDM). Type 2 diabetes is linked to genetic predisposition, obesity, and certain medications. Women can also develop temporary insulin resistance during pregnancy in what is called gestational diabetes. Some adults are diagnosed with latent autoimmune diabetes in adults (LADA), a slow-progressing form of autoimmune diabetes. Similar to type 1 diabetes, in LADA the insulin-producing  $\beta$ -cells of the pancreas are destroyed but at a slower rate. A small percentage of people are diagnosed with maturity onset diabetes of the young (MODY), which refers to any of several hereditary forms of diabetes mellitus caused by mutations in an autosomal dominant gene disrupting insulin production.

20    **[0004]** When a type 1 diabetes, LADA, or MODY patient's pancreas does not produce enough insulin, the patient generally exhibits an atypical glycemia phenotype marked by hyperglycemia. In these cases, the patients are treated with chronic insulin injection therapy. In type 2 and gestational diabetes, patients also often exhibit hyperglycemia as they are unable to properly utilize the insulin that is being produced by the pancreas. In these cases, the patients can be treated with oral medication with or without changes in diet and exercise; however, many subjects eventually progress to resemble a type 1 diabetes condition (inflammatory disease in pancreas with significant loss of beta cell mass) and become dependent on exogenous insulin. Left untreated, diabetes can lead to weight loss, loss of appetite, vomiting, dehydration, problems with motor function, coma, and even death.

25    **[0005]** Approximately 30 million people, or 9.4% of the United States population, have diabetes. Type 1 diabetes accounts for about 5% of all diagnosed cases of diabetes, affecting approximately 1.5 million people. Current diabetes therapies include a variety of short-acting (e.g. Humalog® (Eli Lilly, Indianapolis, IN) and NovoLog® (Novo Nordisk, Bagsværd, Denmark)) and long-acting insulin products (e.g. Lantus® (Sanofi, Paris, France) and Levemir® (Novo Nordisk, Bagsværd, Denmark)), which are administered via subcutaneous injection multiple times a day or through a wearable subcutaneous infusion pump. The burden of frequent injections results in a lack of treatment regimen compliance and under-dosing, leading to poor long-term health outcomes. In fact, each year over 7 million hospital discharges related to diabetes are reported among United States adults due to cardiovascular events, amputations, and ketoacidosis. Furthermore, each year over 14 million emergency department visits related to diabetes are reported among United States adults due to hypoglycemia and hyperglycemia crises among other conditions. Among U.S. adults aged 20 years or older with diagnosed diabetes, the estimated prevalence of kidney disease is over 36%. Diabetes is the seventh leading cause of death in the United States with a total estimated annual cost of over \$245 billion. Other research includes, WO 2018/107117 which is directed to insulin-fc fusions proteins and their use to treat autoimmune disease. The disclosed insulin-fc proteins comprise an insulin polypeptide fused to a Fc domain, wherein the insulin polypeptide comprises a B-chain peptide, a C-chain peptide, and an A-chain peptide, wherein the amino acid sequence of the C-chain peptide is AAK. Wang et al, "IgG Fc engineering to modulate antibody effector functions", PROTEIN & CELL, vol. 9, no. 1, pages 63 - 73, is another review article relating to various antibody engineering approaches to improve safety and efficacy of IgG. However, there remains a need for cost effective and less burdensome treatment options for this disease.

**SUMMARY OF THE PRESENT TECHNOLOGY**

30    **[0006]** It will be appreciated that the scope is in accordance with the claims. Accordingly, there is provided a fusion protein, as defined in claim 1. Further features are provided in accordance with the dependant claims. The specification may also include description of arrangements outside the scope of the invention provided as background and to assist in understanding the invention.

In an aspect defined in claim 2, the present disclosure provides a fusion protein comprising an insulin polypeptide and an Fc fragment, wherein the insulin polypeptide and the Fc fragment are connected by a linker, wherein the fusion protein

comprises the following sequence:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGA  
 5 GGGGAGGGGAGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
 PEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 10 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 87).

**[0007]** As defined in claim 1, the present disclosure provides a fusion protein comprising an insulin polypeptide and an Fc fragment, wherein the insulin polypeptide and the Fc fragment are connected by a linker, comprising domains in the following orientation from N- to C- terminus: (N-terminus)-insulin polypeptide-linker-Fc fragment-(C-terminus), wherein the Fc fragment is of human origin and comprises the following sequence:

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH  
 NAKTKPREEQYX<sub>1</sub>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY  
 TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKS  
 20 SVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 77) wherein Xi is Sor R, and wherein the insulin polypeptide comprises the following sequence:  
 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYC (SEQ ID NO: 10) and wherein the linker comprises the following sequence:

GGGGGAGGGGAGGGGAGGGGG (SEQ ID NO: 67); GGGGGQGGGGQGGGGQGGGGGQGGGG (SEQ ID NO: 99); or  
 GGGGGQGGGGQGGGGQGGGGG (SEQ ID NO: 13).

**[0008]** In some embodiments, the fusion protein comprises the following sequence:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
 GGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
 35 PEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 78).

**[0009]** In some embodiments, the fusion protein comprises the following sequence:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
 GGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
 45 PEVKFNWYVDGVEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 84).

**[0010]** In some embodiments, the fusion protein is a homodimer. In examples, the percentage homodimer of the fusion protein is greater than 90%. In some examples, the fusion protein is made using HEK293 cells or CHO cells, and the resulting homodimer titer after purification using Protein A beads or a Protein A column is greater than 150 mg/L. In embodiments, the insulin receptor IC<sub>50</sub> for the fusion protein is less than or equal to 5000 nM. In some embodiments, the insulin receptor IC<sub>50</sub> for the fusion protein is less than or equal to 2400 nM.

**[0011]** In some embodiments, the human FcRn receptor EC<sub>50</sub> for the fusion protein is less than or equal to 1000 ng/mL. In examples, the human Fc(γ)RI receptor assay OD<sub>450</sub> Ratio at a biotinylated-Fc(γ)RI concentration of 3000 ng/mL of the fusion protein is less than or equal to 0.50. In embodiments, the human C1q assay OD<sub>450</sub> Ratio

at a biotinylated-C1q concentration of 1000 ng/mL is less than or equal to 0.35. In embodiments, the fusion protein is formulated as a pharmaceutical composition. In some examples, the concentration of the fusion protein in the pharmaceutical composition is about 3 mg/mL or greater. In some embodiments, the pharmaceutical composition is suitable for subcutaneous administration.

**[0012]** In embodiments, a physiologically effective amount of the fusion protein or a pharmaceutical composition thereof may be used as a medicament for lowering the blood glucose level of the patient. In examples, the patient is diagnosed with diabetes. In some embodiments, the fusion protein is administered subcutaneously. The fusion protein may be administered daily, twice weekly, or once weekly to the patient. In some examples, the fusion protein is administered once weekly to the patient at a dose between 0.025 and 0.500 mg/kg/week.

**[0013]** In embodiments, a cell may be engineered to express the fusion protein. The cell may be transfected with a nucleic acid encoding the fusion protein. In some examples, the cell is a HEK293 cell or a CHO cell.

**[0014]** In embodiments, the nucleic acid (cDNA) encoding the fusion protein of SEQ ID NO: 87 comprises the following nucleic acid sequence:

```

atggaatggagctgggtctttcttcttctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgcggctccacctggtggaagctctggc
actcgtgtgcggcgagcggggctccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaacagtgtcgcacctccacctgct
ccctggaccagctggaaaactactgcggtggcggaggtggtgcaggaggcgggtggaggtggggctggaggaggcgggggagacaa
aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaaccaaggacacctcatgatctccgg
acccttgaggtcacatgcgtggtgggtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag
acaaagccgcgaggaggagcagtagcagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt
caaggtctccaacaaagccctcccagccccatcgaaaaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacacctgccccat
ccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatccagcgacatcgccgtggagtgggagagcaatgggcag
ccggagaacaactacaagaccagcctccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag
gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 88).

```

**[0015]** In embodiments, the nucleic acid (cDNA) encoding the fusion protein of SEQ ID NO: 78 comprises the following nucleic acid sequence:

```

atggaatggagctgggtctttcttcttctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgcggctccacctggtggaagctctggc
actcgtgtgcggcgagcggggctccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaacagtgtcgcacctccacctgct
ccctggaccagctggaaaactactgcggtggcggaggtggtcaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa
aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaaccaaggacacctcatgatctccgg
acccttgaggtcacatgcgtggtgggtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag
acaaagccgcgaggaggagcagtagcagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt
caaggtctccaacaaagccctcccagccccatcgaaaaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacacctgccccat
ccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatccagcgacatcgccgtggagtgggagagcaatgggcag
ccggagaacaactacaagaccagcctccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag
gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 79).

```

**[0016]** In embodiments, the nucleic acid (cDNA) encoding the fusion protein of SEQ ID NO: 84 comprises the following nucleic acid sequence:

atggaatggagctgggtctttcttcttctcgtcagtaacgactgggtgtccactccttcgtgaaccagcacctgtgcggctcccacctggggaagctctggc  
 actcgtgtgcggcgagcggggtccactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtgcacctccacctgct  
 5 ccctggaccagctggaaaactactgcgggtggcggaggtgtcaaggaggcgggtggacaggggtggaggtgggcaggaggaggcgggggagacaa  
 aactcacacatgcccacgtgcccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacacctcatgatctcccg  
 acccctgaggtcacatgcgtgggtgggtggacgtgagccacgaagacctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaag  
 10 acaaagccgcgggaggagcagtagacaagcacgtaccgtgtgtgtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt  
 caaggtctcaacaaagccctcccagccccatcgagaaaaccatctcaaagccaaaggcagccccgagaaccacaggtgtacacctgccccat  
 cccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatccagcgacatcgccgtggagtgaggagcaatgggcag  
 ccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
 15 gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacgcagaagagcctctcctgtctccgggttag (SEQ ID NO: 85).

## BRIEF DESCRIPTION OF THE DRAWINGS

[0017]

FIG. 1 shows a schematic representation of an exemplary insulin-Fc fusion protein homodimer.

FIG. 2 shows average % fasting blood glucose levels from Day 0 to Day 3 for N=3 dogs dosed intravenously on Day 0 at 0.2 mg/kg with the homodimer of SEQ ID NO: 31 (not claimed).

FIG. 3 illustrates a side-by-side sequence comparison of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, and SEQ ID NO: 35 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 4 illustrates a side-by-side sequence comparison of SEQ ID NO: 31, SEQ ID NO: 36, SEQ ID NO: 37, and SEQ ID NO: 38 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 5 shows average % fasting blood glucose levels from Day 0 to Day 7 for N=3 dogs dosed intravenously on Day 0 at 0.2 mg/kg with the homodimer of SEQ ID NO: 36 (not claimed).

FIG. 6 shows average % fasting blood glucose levels from Day 0 to Day 7 for N=6 dogs dosed subcutaneously on Day 0 at 0.33 mg/kg with the homodimer of SEQ ID NO: 36 (not claimed).

FIG. 7 shows the average anti-drug antibody titer ( $\mu\text{g/mL}$ ) for N=3 dogs dosed subcutaneously on Day 0 (0.30 mg/kg), Day 28 (0.33 mg/kg), Day 35 (0.33 mg/kg), Day 42 (0.50 mg/kg), Day 49 (1.00 mg/kg) and Day 56 (1.00 mg/kg) with the homodimer of SEQ ID NO: 36 (not claimed).

FIG. 8 illustrates a side-by-side sequence comparison of SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 41, and SEQ ID NO: 42 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 9 shows the average anti-drug antibody titer ( $\mu\text{g/mL}$ ) for N=1 dog dosed subcutaneously on Day 0 (0.33 mg/kg), Day 7 (0.50 mg/kg), Day 14 (0.50 mg/kg), and Day 21 (0.50 mg/kg) with the homodimer of SEQ ID NO: 42 (not claimed).

FIG. 10 shows the average anti-drug antibody titer ( $\mu\text{g/mL}$ ) for N=1 dogs dosed subcutaneously on Day 0 (0.33 mg/kg) and Day 14 (0.16 mg/kg) with the homodimer of SEQ ID NO: 43 (not claimed).

FIG. 11 shows average % fasting blood glucose levels from Day 0 to Day 7 for N=2 dogs dosed subcutaneously on Day 0 at 0.33 mg/kg with the homodimer of SEQ ID NO: 43 (not claimed).

FIG. 12 illustrates a side-by-side sequence comparison of SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47 and SEQ ID NO: 48 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 13 illustrates a side-by-side sequence comparison of SEQ ID NO: 43, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, and SEQ ID NO: 52 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 14 illustrates a side-by-side sequence comparison of SEQ ID NO: 43, SEQ ID NO: 48 and SEQ ID NO: 53 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 15 illustrates a side-by-side sequence comparison of SEQ ID NO: 43, SEQ ID NO: 51, SEQ ID NO: 52 and SEQ ID NO: 54 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 16 illustrates a side-by-side sequence comparison of SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 43, SEQ ID NO: 55, SEQ ID NO: 56 and SEQ ID NO: 57 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 17 shows % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously on Day 0 at 0.16 mg/kg with the homodimer of SEQ ID NO: 28 (not claimed).

FIG. 18 shows the anti-drug antibody titer ( $\mu\text{g/mL}$ ) for N=1 dog dosed subcutaneously on Day 0 (0.16 mg/kg), Day 14 (0.16 mg/kg), Day 28 (0.16 mg/kg), and Day 42 (0.16 mg/kg) with the homodimer of SEQ ID NO: 28 (not claimed).

FIG. 19 shows % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously on Day 0 at 0.33 mg/kg with the homodimer of SEQ ID NO: 26 (not claimed).

FIG. 20 shows % fasting blood glucose levels from Day 0 to Day 60 for N=1 dog dosed subcutaneously on Day 0 (0.33 mg/kg), Day 15 (0.16 mg/kg), Day 31 (0.16 mg/kg) and Day 45 (0.15 mg/kg) with the homodimer of SEQ ID NO: 26 (not claimed).

FIG. 21 shows the anti-drug antibody titer ( $\mu\text{g/mL}$ ) for N=1 dogs dosed subcutaneously on Day 0 (0.33 mg/kg), Day 15 (0.16 mg/kg), Day 31 (0.16 mg/kg) and Day 45 (0.15 mg/kg) with the homodimer of SEQ ID NO: 26 (not claimed).

FIG. 22 shows % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously on Day 0 at 0.16 mg/kg with the homodimer of SEQ ID NO: 58 (not claimed).

FIG. 23 shows % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously on Day 0 at 0.16 mg/kg with the homodimer of SEQ ID NO: 59 (not claimed).

FIG. 24 illustrates a side-by-side sequence comparison of SEQ ID NO: 61 and SEQ ID NO: 62 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 25 shows % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously on Day 0 at 0.16 mg/kg with the homodimer of SEQ ID NO: 61, and % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously on Day 0 at 0.16 mg/kg with the homodimer of SEQ ID NO: 62 (not claimed).

FIG. 26 shows % fasting blood glucose levels from Day 0 to Day 7 for N=1 dog dosed subcutaneously with the homodimer of SEQ ID NO: 30 (not claimed) in addition to the times that the dog was given food.

FIG. 27 illustrates a side-by-side sequence comparison of SEQ ID NO: 76 (not claimed), SEQ ID NO: 91 (not claimed), and SEQ ID NO: 78. "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 28 illustrates a side-by-side sequence comparison of SEQ ID NO: 75, SEQ ID NO: 76, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, and SEQ ID NO: 95 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 29 illustrates a side-by-side sequence comparison of SEQ ID NO: 76 (not claimed), SEQ ID NO: 78, SEQ ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), SEQ ID NO: 84, and SEQ ID NO: 86 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 30 illustrates a side-by-side sequence comparison of SEQ ID NO: 87, SEQ ID NO: 96, SEQ ID NO: 78, SEQ ID NO: 97 (not claimed), SEQ ID NO: 89 (not claimed), and SEQ ID NO: 98 (not claimed). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FIG. 31 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 78) and its corresponding nucleic acid sequence (SEQ ID NO: 79).

FIG. 32 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 80 (not claimed)) and its corresponding nucleic acid sequence (SEQ ID NO: 81 (not claimed)).

FIG. 33 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 82 (not claimed)) and its corresponding nucleic acid sequence (SEQ ID NO: 83 (not claimed)).

FIG. 34 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 84) and its corresponding nucleic acid sequence (SEQ ID NO: 85).

FIG. 35 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 87) and its corresponding nucleic acid sequence (SEQ ID NO: 88).

FIG. 36 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 89 (not claimed)) and its corresponding nucleic acid sequence (SEQ ID NO: 90 (not claimed)).

FIG. 37 illustrates the full amino acid sequence including the leader sequence of a fusion protein (SEQ ID NO: 86 (not claimed)) and its corresponding nucleic acid sequence (SEQ ID NO: 100 (not claimed)).

FIG. 38 shows average % fasting blood glucose levels from Hour 0 to Hour 10 for N=12 Balb/c mice fasted 1 hour before subcutaneous injection at a dose of 300  $\mu$ g/kg with the homodimers of SEQ ID NO: 87 and SEQ ID NO: 89(not claimed).

## DETAILED DESCRIPTION

**[0018]** An insulin treatment that requires less frequent dosing (e.g., once-weekly injections) would be less burdensome on patients, leading to better compliance, better glucose control, and ultimately better long-term health outcomes. As disclosed herein, proposed ultra-long acting insulin treatments for human clinical use comprise an insulin-Fc fusion protein making use of a human Fc fragment to prolong their action in vivo. An insulin-Fc fusion protein suitable for an ultra-long acting treatment for diabetes should meet various design goals. An insulin-Fc fusion protein suitable for an ultra-long acting treatment for diabetes should be manufacturable in mammalian cells, for example human embryonic kidney (HEK, e.g. HEK293) cells, with an acceptable titer of the desired homodimer product (e.g., greater than 50 mg/L homodimer titer from transiently transfected HEK cells, greater than 75 mg/L from transiently transfected HEK cells, greater than 100 mg/L from transiently transfected HEK cells, greater than 150 mg/L from transiently transfected HEK cells, etc.). Only human insulin-Fc fusion protein configurations with a homodimer titer of greater than 150 mg/L are considered useful in the present invention, because experience has demonstrated that homodimer titers less than this level in transiently transfected HEK cells will not likely result in commercial production homodimer titers in stably transfected Chinese hamster ovary (CHO) cells that meet the low manufacturing cost requirements for the relatively commoditized human insulin market.

**[0019]** In addition, the insulin-Fc fusion protein must bind the IR with an appreciable affinity (e.g., IC<sub>50</sub> less than 5000 nM, IC<sub>50</sub> less than 4000 nM, IC<sub>50</sub> less than 3000 nM, IC<sub>50</sub> less than 2400 nM, IC<sub>50</sub> more preferably less than 2000 nM, etc.) as measured in the 4°C IM-9 IR binding assay. Based on experience, only molecules exhibiting IR activity IC<sub>50</sub> values less than 5000 nM are deemed likely to exhibit the requisite bioactivity. In preferred embodiments, the insulin-Fc fusion protein exhibits an IR activity IC<sub>50</sub> value less than 2400 nM, and more preferably less than 2000 nM. The insulin-Fc fusion protein configuration must also exhibit sustained bioactivity in vivo (e.g., demonstrate glucose lowering activity greater than about 2 hours, 6 hours, 9 hours, 12 hours, 18 hours, 1 day, 1.5 days, 2 days, 2.5 days, 3 days, 4 days, 5 days, 6 days, 7 days, or longer) to justify less frequent dosing. The insulin-Fc fusion protein configuration must also demonstrate prolonged system residence time in vivo (e.g., the serum half-life must be greater than 3 days, or longer). The sustained bioactivity and prolonged residence time of a given insulin-Fc fusion protein configuration may be predicted by its ability to bind the FcRn receptor, which is responsible for the prolonged in vivo elimination half-lives of antibodies and Fc-fusion proteins. FcRn receptor activity is typically measured by the concentration of an insulin-Fc fusion protein that causes the insulin-Fc fusion protein to reach half of its maximum binding (i.e., EC<sub>50</sub> value) as measured on an assay (e.g., an enzyme-linked immunosorbent assay (ELISA) assay) using OD 450 nm values as measured on a microplate reader. Based on experience, insulin-Fc fusion protein configurations exhibiting human FcRn receptor EC<sub>50</sub> values less than or equal to 1500 ng/mL (and more preferably less than 1000 ng/mL) are the most likely to exhibit long enough half-lives to justify once-a-week dosing.

**[0020]** Lastly, to be useful for treating a chronic disease such as diabetes, the insulin-Fc fusion protein configuration must not induce the production of anti-drug antibodies, especially antibodies that neutralize the bioactivity of the molecule after repeated dosing. The propensity of a given insulin-Fc fusion protein configuration to induce an adverse immunogenic response may be predicted first by its ability to bind the Fc(gamma)RI receptor, which plays a key role in many immune system effector functions, including phagocytosis of opsonized molecules, release of inflammatory mediators, and antibody-dependent cellular cytotoxicity. Fc(gamma)RI receptor activity is typically measured by the absorbance at a wavelength at 450 nm (OD<sub>450</sub>) value obtained on a microplate reader in an enzyme-linked immunosorbent assay (ELISA) assay at a given concentration of the insulin-Fc fusion protein. Based on experience, insulin-Fc fusion protein configurations exhibiting human Fc(gamma)RI receptor assay OD<sub>450</sub> Ratios (where the reference insulin-Fc fusion protein configuration for the ratio is SEQ ID NO: 76) at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL less than or equal to 0.50 are likely to exhibit sufficiently low immunogenicity to justify repeated once-a-week dosing. The propensity of a given insulin-Fc fusion protein configuration to induce an adverse immunogenic response may also be predicted by its ability to bind the complement component 1q (C1q), which activates the complement cascade causing

phagocytes to clear the bound molecule, inflammation to attract additional phagocytes, and activation of the cell-killing membrane attack complex. C1q activity is typically measured by the OD450 value obtained on a microplate reader in an enzyme-linked immunosorbent assay (ELISA) assay at a given concentration of the insulin-Fc fusion protein configuration that is coated onto a microplate. Based on experience, insulin-Fc fusion protein configurations exhibiting human C1q receptor assay OD450 Ratios (where the reference insulin-Fc fusion protein configuration for the ratio is SEQ ID NO: 76) at a biotinylated-C1q concentration of 1000 ng/mL less than or equal to 0.35 are likely to exhibit sufficiently low immunogenicity to justify repeated once-a-week dosing.

**[0021]** Proposed ultra-long acting insulin treatments for human clinical use comprise an insulin-Fc fusion protein making use of a human Fc fragment to prolong their action in vivo. To understand the behavior of insulin-Fc fusion protein configurations of various designs, insulin-Fc fusion protein configurations useful as ultra-long acting insulins for dogs were considered first. As a human Fc fragment is expected to be immunogenic and therefore capable of inducing the production of anti-drug antibodies in dogs, the human Fc fragment was replaced with a canine Fc fragment.

**[0022]** However, it was unexpectedly found that a simple exchange between the human Fc fragment and any canine Fc fragment in the insulin-Fc fusion protein configuration did not necessarily yield a product with an acceptable homodimer titer (e.g., a homodimer titer greater than 50 mg/L) or a sufficiently high NAOC value (e.g., a NAOC greater than 150 %FBGL·days·kg/mg). For example, in some cases only a specific isotype (e.g., canine IgGB) for the Fc fragment resulted in an insulin-Fc fusion protein configuration with a high enough homodimer titer to meet the design goals (e.g., a homodimer titer of canine insulin-Fc fusion protein of greater than 50 mg/L) and an acceptably high NAOC value (e.g., a NAOC greater than 150 %FBGL·days·kg/mg). In other cases, specific amino acids of the insulin polypeptide of the insulin-Fc fusion protein configuration were found to be immunogenic in the target species thereby requiring site-directed mutations to find the relatively small number of insulin-Fc fusion protein configurations that were both non-immunogenic and bioactive in the target species with acceptably high NAOC values (e.g., NAOC values greater than 150 %FBGL·days·kg/mg) and NAOCR values after the third weekly subcutaneous dose that were greater than 0.50.

**[0023]** In further cases, when the Fc fragments were mutated to prevent glycosylation and thereby further reduce the immunogenicity of the insulin-Fc fusion protein configuration, it was discovered unexpectedly that only specific amino acid mutations in the Fc fragment led to the desired homodimer titers (e.g., homodimer titers for canine insulin-Fc fusion protein configurations of greater than 50 mg/L) and NAOC values (e.g., NAOC greater values than 150 %FBGL·days·kg/mg). Furthermore, it was discovered that an additional mutation in the insulin component of the insulin-Fc fusion protein configuration was required to produce these Fc-mutated, non-glycosylated insulin Fc-fusion protein configurations with the desired homodimer titers (e.g., homodimer titers for canine insulin-Fc fusion protein configurations of greater than 50 mg/L) and NAOC values (e.g., NAOC greater values than 150 %FBGL·days·kg/mg), while also achieving NAOCR values after the third weekly subcutaneous dose that were greater than 0.50.

**[0024]** For human ultra-long acting insulins, to maximize homodimer titers and reduce manufacturing costs, insulin-Fc fusion protein configurations were produced comprising Fc fragments based on different isotypes of the human IgG molecule (e.g., IgG1 and IgG2). It was discovered rather unexpectedly that switching from an Fc fragment based on the IgG2 molecule to an Fc fragment based on the IgG1 molecule in the insulin-Fc fusion protein configuration increased the average homodimer titer by over 50% without significantly compromising the IR or FcRn binding activity. However, the resulting IgG1-derived insulin-Fc fusion protein configuration was deemed much more likely to interact adversely with the immune system and develop neutralizing antibodies given that its human Fc(gamma)RI receptor assay OD450 Ratio (where the reference insulin-Fc fusion protein configuration for the ratio is SEQ ID NO: 76) at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL was much greater than 0.50 and its human C1q receptor assay OD450 Ratio (where the reference insulin-Fc fusion protein configuration for the ratio is SEQ ID NO: 76) at biotinylated-C1q concentration of 1000 ng/mL was much greater than 0.35.

**[0025]** In an attempt to reduce unwanted immunogenicity, the insulin-Fc fusion protein configuration was mutated to prevent the glycosylation of the Fc fragment during synthesis in the host cell. Specifically, the conserved asparagine (N)-glycosylation site in the CH2 domain of the heavy chain of the IgG1 Fc region was mutated to different amino acids (e.g., S, D, A, R, and Q) in an attempt to maintain the improved insulin-Fc fusion protein homodimer yields while reducing Fc(gamma)RI and C1q interactions. Interestingly, the S, D (not claimed), A (not claimed), and R mutated non-glycosylated insulin-Fc fusion protein configurations gave improved homodimer titers relative to the glycosylated parent configurations; however, the homodimer titer of the insulin-Fc fusion protein configuration with the Q mutation at the conserved asparagine (N)-glycosylation site was too low (i.e., less than the 150 mg/L design goal for human insulin-Fc fusion protein configurations) to support the low cost of manufacturing requirements. Additionally, the Fc(gamma)RI and C1q binding decreased significantly for the S, D (not claimed), A (not claimed), and R mutated non-glycosylated insulin-Fc fusion protein configurations. However, in these insulin-Fc fusion protein configurations the improvements in yield and immunogenicity were offset by unexpectedly lower FcRn binding affinity and significantly lower IR binding relative to the glycosylated parent configurations, indicating an unacceptable reduction in bioactivity and residence time in vivo.

**[0026]** As was unexpectedly discovered with the insulin-Fc fusion proteins configurations comprising an Fc fragment of canine origin, it was discovered that by changing a single amino acid in the insulin sequence in human insulin-Fc

fusion protein configurations, the improvements in yield and immunogenicity of the S, D (not claimed), A (not claimed), and R mutated non-glycosylated insulin-Fc fusion protein configurations were maintained or improved upon, while the IR and FcRn binding affinities increased (e.g. lower IR assay IC50 value), rather than decreased relative to the original glycosylated parent configuration. The resulting non-glycosylated, mutated insulin-Fc fusion protein configurations are expected, therefore, to exhibit acceptable in vivo glucose lowering potencies and prolonged residence times. Unexpectedly, changing the same amino acid in the non-glycosylated insulin-Fc fusion protein configuration with the Q mutation at the conserved asparagine (N)-glycosylation site resulted in a significant decrease in the already poor homodimer titer.

**[0027]** To understand how the properties of the S, D (not claimed), A (not claimed), and R non-glycosylated mutant insulin-Fc fusion proteins configurations could be further manipulated, modifications were made to the linker region of the insulin-Fc fusion protein configuration, connecting the insulin polypeptide to the Fc fragment. Test results indicated that without a linker, the homodimer titers of the resulting canine insulin-Fc fusion protein configurations were unacceptably low (i.e., less than 50 mg/L). For insulin-Fc fusion protein configurations comprising linkers of the same length, it was discovered that certain amino acid sequences were preferred over others in terms of homodimer titers, IR and FcRn receptor binding, and Fc(gamma)RI and C1q binding of the insulin-Fc fusion protein configuration.

**[0028]** Provided herein, therefore, are specific manufacturable, high purity, long-acting, bioactive, non-immunogenic insulin-Fc fusion protein configurations, each of which comprises a mutated insulin polypeptide, a non-glycosylated Fc fragment, and a linker between the mutated insulin polypeptide and the non-glycosylated Fc fragment, and wherein the insulin-Fc fusion protein configurations meet the design goals of acceptably high homodimer titers (e.g., homodimer titers greater than 150 mg/L), IR assay IC50 values (e.g., IC50 less than 5000 nM, less than 2400 nM, and more preferably less than 2000 nM), human FcRn receptor EC50 values (e.g., EC50 less than or equal to 1500 ng/L, and more preferably less than 1000 ng/mL), human Fc(gamma)RI receptor OD 450 Ratios (e.g. OD450 Ratios at a biotinylated-Fc(gamma)RI receptor concentration of 3000 ng/mL less than or equal to 0.50 where the reference insulin-Fc fusion protein configuration for the ratio is SEQ ID NO: 76), and human C1q receptor OD 450 Ratios (e.g. OD450 Ratios at a biotinylated-C1q concentration of 1000 ng/mL less than or equal to 0.35 where the reference insulin-Fc fusion protein configuration for the ratio is SEQ ID NO: 76). These exemplary insulin-Fc fusion protein configurations are expected to exhibit sufficiently low immunogenicity long enough half-lives to justify repeated once-a-week dosing, rendering them suitable for the treatment of diabetes.

### Definitions

**[0029]** As used herein, the articles "a" and "an" refer to one or more than one, e.g., to at least one, of the grammatical object of the article. The use of the words "a" or "an" when used in conjunction with the term "comprising" herein may mean "one," but it is also consistent with the meaning of "one or more," "at least one," and "one or more than one."

**[0030]** As used herein, "about" and "approximately" generally mean an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Exemplary degrees of error are within 20 percent (%), typically, within 10%, and more typically, within 5% of a given range of values.

**[0031]** As used herein, an amount of a molecule, compound, conjugate, or substance effective to treat a disorder (e.g., a disorder described herein), "therapeutically effective amount," or "effective amount" refers to an amount of the molecule, compound, conjugate, or substance which is effective, upon single or multiple dose administration(s) to a subject, in treating a subject, or in curing, alleviating, relieving or improving a subject with a disorder (e.g., a disorder described herein) beyond that expected in the absence of such treatment.

**[0032]** As used herein, the term "analog" refers to a compound or conjugate (e.g., a compound or conjugate as described herein, e.g., insulin) having a chemical structure similar to that of another compound or conjugate but differing from it in at least one aspect.

**[0033]** As used herein, the term "antibody" or "antibody molecule" refers to an immunoglobulin molecule (Ig), immunologically active portions of an immunoglobulin (Ig) molecule, i.e., a molecule that contains an antigen binding site that specifically binds, e.g., immunoreacts with, an antigen. As used herein, the term "antibody domain" refers to a variable or constant region of an immunoglobulin. As used herein, the term "antibody domain" refers to a variable or constant region of an immunoglobulin. It is documented in the art that antibodies comprise several classes, for example IgA, IgM, or IgG in the case of mammals (e.g., humans). Classes of immunoglobulins can be further classified into different isotypes, such as IgGA, IgGB, IgGC, and IgGD for canines, and IgG1, IgG2, IgG3, and IgG4 for humans. Those skilled in the art will recognize that immunoglobulin isotypes of a given immunoglobulin class will comprise different amino acid sequences, structures, and functional properties from one another (e.g., different binding affinities to Fc(gamma) receptors). "Specifically binds" or "immunoreacts with" means that the antibody reacts with one or more antigenic determinants of the desired antigen and has a lower affinity for other polypeptides, e.g., does not react with other polypeptides.

**[0034]** As used herein, the term "area-under-the-curve" or "AUC" refers to the integrated area under the %FBGL (Fasting Blood Glucose Level) vs. time curve for a subject after a given dose of an insulin-Fc fusion protein is administered. As used herein, the term "area-over-the curve" or "AOC" is used as a measure of the biological potency of an insulin-

Fc fusion protein such that the AOC equals the difference between the total possible area under the %FBGL vs. time curve and the AUC value. As used herein, the "normalized area-over-the curve," "normalized AOC," or "NAOC" is the AOC value divided by the actual dose of insulin-Fc fusion protein administered. As used herein, the term "normalized AOC ratio" or "NAOCR" is the ratio of the NAOC resulting from a particular administration of an insulin-Fc fusion protein to the NAOC resulting from the first administration of an insulin-Fc fusion protein in a series of administrations. The NAOCR thus provides a measure of the change in biological activity of an insulin-Fc fusion protein after repeated administrations.

**[0035]** As used herein, the term "bioactivity," "activity," "biological activity," "potency," "bioactive potency," or "biological potency" refers to the extent to which an insulin-Fc fusion protein activates the IR and/or exerts a reduction in blood glucose levels in a target subject. As used herein, "in vitro activity" or "IR activity" refers to the affinity with which an insulin-Fc fusion protein binds to the IR and is typically measured by the concentration at which an insulin-Fc fusion protein displaces half of an insulin reference standard from the IR in a competitive binding assay (i.e., IC50). As used herein, "in vivo activity" refers to the extent and duration of reduction in a target subject's fasting blood glucose level after administration of an insulin-Fc fusion protein.

**[0036]** As used herein, the term "biosynthesis," "recombinant synthesis," or "recombinantly made" refers to the process by which an insulin-Fc fusion protein is expressed within a host cell by transfecting the cell with a nucleic acid molecule (e.g., vector) encoding the insulin-Fc fusion protein (e.g., where the entire insulin-Fc fusion protein is encoded by a single nucleic acid molecule). Exemplary host cells include mammalian cells, e.g., HEK293 cells or CHO cells. The cells can be cultured using standard methods in the art and the expressed insulin-Fc fusion protein may be harvested and purified from the cell culture using standard methods in the art.

**[0037]** As used herein, the term "cell surface receptor" refers to a molecule such as a protein, generally found on the external surface of the membrane of a cell and which interacts with soluble molecules, e.g., molecules that circulate in the blood supply. In some embodiments, a cell surface receptor may include a hormone receptor (e.g., an insulin hormone receptor or insulin receptor (IR)) or an Fc receptor which binds to an Fc fragment or the Fc region of an antibody (e.g., an Fc(gamma) receptor, for example Fc(gamma)RI, or an Fc neonatal receptor, for example FcRn). As used herein, "in vitro activity" or "Fc(gamma) receptor activity" or "Fc(gamma) receptor binding" or "FcRn receptor activity" or "FcRn binding" refers to the affinity with which an insulin-Fc fusion protein binds to the Fc receptor (e.g. Fc(gamma) receptor or FcRn receptor) and is typically measured by the concentration of an insulin-Fc fusion protein that causes the insulin-Fc fusion protein to reach half of its maximum binding (i.e., EC50 value) as measured on an assay (e.g., an enzyme-linked immunosorbent assay (ELISA) assay) using OD 450 nm values as measured on a microplate reader. Alternatively, the affinity with which an insulin-Fc fusion protein binds to the Fc receptor (e.g. Fc(gamma) receptor or FcRn receptor) is measured by the OD 450 nm value obtained on a microplate reader in an enzyme-linked immunosorbent assay (ELISA) assay at a given concentration of the insulin-Fc fusion protein.

**[0038]** As used herein, the term "C1q" or "complement component 1q" means a protein complex involved in the complement system, which is part of the innate immune system. C1q together with C1r and C1s form the C1 complex. C1q plays a role in involved in specific antigen presentation by dendritic cells to T cells and B cells.

**[0039]** As used herein, the term "fasting blood glucose level" or "FBGL" refers to the average blood glucose level in a target subject at the end of a period during which no food is administered and just prior to the time at which an insulin-Fc fusion protein is administered. As used herein, the term "percent fasting blood glucose level," "% fasting blood glucose level," or "%FBGL" refers to the ratio of a given blood glucose level to the fasting blood glucose level multiplied by 100.

**[0040]** As used herein, the term "immunogenic" or "immunogenicity" refers to the capacity for a given molecule (e.g., an insulin-Fc fusion protein of the present invention) to provoke the immune system of a target subject such that after repeated administrations of the molecule, the subject develops antibodies capable of specifically binding the molecule (i.e., anti-drug antibodies). As used herein, the terms "neutralizing," "neutralizing antibodies", or "neutralizing anti-drug antibodies" refer to the capacity for antibodies to interfere with the compound's biological activity in the target subject. As used herein, the term "immunogenic epitopes," "immunogenic hot spots," or "hot spots" refers to the mutations or epitopes of a given molecule (e.g., an insulin-Fc fusion protein of the present invention) that are responsible for moderate or strong binding of the anti-drug antibodies.

**[0041]** As used herein, the term "insulin reference standard" is any one of: (i) a naturally occurring insulin from a mammal (e.g., a dog, or a human); (ii) an insulin polypeptide that does not comprise an Fc fragment; or (iii) a standard of care insulin (e.g., a commercially available insulin).

**[0042]** As used herein, the term "monomer" refers to a protein or a fusion protein comprising a single polypeptide. In embodiments, the "monomer" is a protein or a fusion protein, e.g., a single polypeptide, comprising an insulin polypeptide and an Fc fragment polypeptide, wherein the insulin and Fc fragment polypeptides are joined by peptide bonds to form the single polypeptide. In embodiments, the monomer is encoded by a single nucleic acid molecule.

**[0043]** As used herein, "N-terminus" refers to the start of a protein or polypeptide that is initiated by an amino acid containing a free amine group that is the alpha-amino group of the amino acid (e.g. the free amino that is covalently linked to one carbon atom that is located adjacent to a second carbon atom, wherein the second carbon atom is part of

the carbonyl group of the amino acid). As used herein, "C-terminus" refers to the end of a protein or polypeptide that is terminated by an amino acid containing a carboxylic acid group, wherein the carbon atom of the carboxylic acid group is located adjacent to the alpha-amino group of the amino acid.

**[0044]** As used herein, "OD450", "optical density at 450 nm", and "absorbance at 450 nm" may be used interchangeably and refer to the use of a microplate reader to read the absorbance of light at 450 nm that is passed through a sample in an assay, e.g. a microplate-based assay, e.g. an enzyme-linked immunosorbent assay, e.g. an ELISA assay.

**[0045]** As used herein, "OD450 Ratio" for a particular assay refers to a way to compare an OD450 value obtained for a first test insulin-Fc fusion protein run at a particular time against an OD450 value obtained for a second test insulin-Fc fusion protein run at another time. The OD450 Ratio is obtained by dividing the OD450 value for a first test article by the OD450 value of a reference insulin-Fc fusion protein. Similarly, a second OD450 Ratio can be obtained for a second test article by dividing the OD450 value for a second test article by the OD450 value of the same reference insulin-Fc fusion protein as used for calculating the OD450 Ratio of the first test article. As a result, the assay properties of a first test insulin-Fc fusion protein and a second test insulin-Fc fusion protein can be compared. The reference insulin-Fc fusion protein configuration used for calculating OD450 Ratios is SEQ ID NO: 76.

**[0046]** As used herein, "pharmacodynamics" or "PD" generally refers to the biological effects of an insulin-Fc fusion protein in a subject. Specifically, herein the PD refers to the measure of the reduction in fasting blood glucose level over time in a subject after the administration of an insulin-Fc fusion protein.

**[0047]** As used herein, "pharmacokinetics" or "PK" generally refers to the characteristic interactions of an insulin-Fc fusion protein and the body of the subject in terms of its absorption, distribution, metabolism, and excretion. Specifically, herein the PK refers to the concentration of an insulin-Fc fusion protein in the blood or serum of a subject at a given time after the administration of the insulin-Fc fusion protein. As used herein, "half-life" refers to the time taken for the concentration of insulin-Fc fusion protein in the blood or serum of a subject to reach half of its original value as calculated from a first order exponential decay model for drug elimination. Insulin-Fc fusion proteins with greater "half-life" values demonstrate greater duration of action in the target subject.

**[0048]** The terms "sequence identity" "sequence homology" "homology" or "identical" in amino acid or nucleotide sequences as used herein describes that the same nucleotides or amino acid residues are found within the variant and reference sequences when a specified, contiguous segment of the nucleotide sequence or amino acid sequence of the variant is aligned and compared to the nucleotide sequence or amino acid sequence of the reference sequence. Methods for sequence alignment and for determining identity between sequences are known in the art, including the use of Clustal Omega, which organizes, aligns, and compares sequences for similarity, wherein the software highlights each sequence position and compares across all sequences at that position and assigns one of the following scores: an "\*" (asterisk) for sequence positions which have a single, fully conserved residue, a ":" (colon) indicates conservation between groups of strongly similar properties with scoring greater than 0.5 in the Gonnet PAM 250 matrix, and a "." (period) indicates conservation between groups of weakly similar properties with scoring less than or equal to 0.5 in the Gonnet PAM 250 matrix, a "-" (dash) indicates a sequence gap, meaning that no local homology exists within a particular set of comparisons within a certain range of the sequences, and an empty space " " indicates little or no sequence homology for that particular position across the compared sequences. See, for example Ausubel et al., eds. (1995) Current Protocols in Molecular Biology, Chapter 19 (Greene Publishing and Wiley- Interscience, New York); and the ALIGN program Dayhoff (1978) in Atlas of Polypeptide Sequence and Structure 5: Suppl. 3 (National Biomedical Research Foundation, Washington, D.C.). With respect to optimal alignment of two nucleotide sequences, the contiguous segment of the variant nucleotide sequence may have additional nucleotides or deleted nucleotides with respect to the reference nucleotide sequence. Likewise, for purposes of optimal alignment of two amino acid sequences, the contiguous segment of the variant amino acid sequence may have additional amino acid residues or deleted amino acid residues with respect to the reference amino acid sequence. In some embodiments, the contiguous segment used for comparison to the reference nucleotide sequence or reference amino acid sequence will comprise at least 6, 10, 15, or 20 contiguous nucleotides, or amino acid residues, and may be 30, 40, 50, 100, or more nucleotides or amino acid residues. Corrections for increased sequence identity associated with inclusion of gaps in the variant's nucleotide sequence or amino acid sequence can be made by assigning gap penalties. Methods of sequence alignment are known in the art.

**[0049]** In embodiments, the determination of percent identity or "homology" between two sequences is accomplished using a mathematical algorithm. For example, the percent identity of an amino acid sequence is determined using the Smith-Waterman homology search algorithm using an affine 6 gap search with a gap open penalty of 12 and a gap extension penalty of 2, BLOSUM matrix 62. The Smith-Waterman homology search algorithm is described in Smith and Waterman (1981) Adv. Appl. Math 2:482-489. In embodiments, the percent identity of a nucleotide sequence is determined using the Smith-Waterman homology search algorithm using a gap open penalty of 25 and a gap extension penalty of 5. Such a determination of sequence identity can be performed using, for example, the DeCypher Hardware Accelerator from TimeLogic.

**[0050]** As used herein, the term "homology" is used to compare two or more proteins by locating common structural characteristics and common spatial distribution of, for instance, beta strands, helices, and folds. Accordingly, homologous

protein structures are defined by spatial analyses. Measuring structural homology involves computing the geometric-topological features of a space. One approach used to generate and analyze three-dimensional (3D) protein structures is homology modeling (also called comparative modeling or knowledge-based modeling) which works by finding similar sequences on the basis of the fact that 3D similarity reflects 2D similarity. Homologous structures do not imply sequence similarity as a necessary condition.

[0051] As used herein, the terms "subject" and "patient" are intended to include canine animals and humans having a disease or a disorder, e.g., diabetes or another disease or disorder described herein, or normal subjects.

[0052] As used herein, the term "titer" or "yield" refers to the amount of a fusion protein product (e.g., an insulin-Fc fusion protein described herein) resulting from the biosynthesis (e.g., in a mammalian cell, e.g., in a HEK293 cell or CHO cell) per volume of the cell culture. The amount of product may be determined at any step of the production process (e.g., before or after purification), but the yield or titer is always stated per volume of the original cell culture. As used herein, the term "product yield" or "total protein yield" refers to the total amount of insulin-Fc fusion protein expressed by cells and purified via at least one affinity chromatography step (e.g. Protein A or Protein G) and includes monomers of insulin-Fc fusion protein, homodimers of insulin-Fc fusion protein, and higher-order molecular aggregates of homodimers of insulin-Fc fusion protein. As used herein, the term "percent homodimer" or "%homodimer" refers to the proportion of a fusion protein product (e.g., an insulin-Fc fusion protein described herein) that is the desired homodimer. As used herein, the term "homodimer titer" refers to the product of the %homodimer and the total protein yield after Protein A purification step reported per volume of the cell culture.

[0053] As used herein, the terms "treat" or "treating" a subject having a disease or a disorder refer to subjecting the subject to a regimen, for example the administration of a fusion protein such as a fusion protein described herein, such that at least one symptom of the disease or disorder is cured, healed, alleviated, relieved, altered, remedied, ameliorated, or improved. Treating includes administering an amount effective to alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease or disorder, or the symptoms of the disease or disorder. The treatment may inhibit deterioration or worsening of a symptom of a disease or disorder.

#### ***Insulin-Fc Fusion Protein Components and Structure***

[0054] The present disclosure relates to a composition of a fusion protein (i.e., an insulin-Fc fusion protein) comprising an insulin polypeptide linked via a peptide linker to a species-specific Fc fragment, and its use to treat diabetes (e.g., in humans and/or dogs) as defined in claim 1. As used herein, the terms "fusion protein" and "insulin-Fc fusion protein" refer to a protein comprising more than one part, for example from different sources (different proteins, polypeptides, cells, etc.), that are covalently linked through peptide bonds. The insulin-Fc fusion proteins are covalently linked by (i) connecting the genes that encode for each part into a single nucleic acid molecule and (ii) expressing in a host cell (e.g., HEK or CHO) the protein for which the nucleic acid molecule encodes as follows: (N-terminus)--insulin polypeptide--linker--Fc fragment--(C-terminus). The fully recombinant synthesis approach is preferred over methods in which the insulin polypeptide and Fc fragments are synthesized separately and then chemically conjugated. The chemical conjugation step and subsequent purification process increase the manufacturing complexity, reduce product yield, and increase cost.

[0055] As used herein, the term "dimer" refers to a protein or a fusion protein comprising two polypeptides linked covalently. In embodiments, two identical polypeptides are linked covalently (e.g., via disulfide bonds) forming a "homodimer" (diagrammatically represented in FIG. 1). Disulfide bonds are shown in FIG. 1; the total number of disulfide bonds in actuality may be greater or less than the number shown in FIG. 1. In embodiments, the homodimer is encoded by a single nucleic acid molecule, wherein the homodimer is made recombinantly inside a cell by first forming insulin-Fc fusion protein monomers and by then assembling two identical insulin-Fc fusion protein monomers into the homodimer upon further processing inside the cell.

[0056] As used herein, the terms "multimer," "multimeric," or "multimeric state" refer to noncovalent, associated forms of Fc fusion protein dimers that may be in equilibrium with Fc fusion protein dimers or may act as permanently aggregated versions of Fc fusion protein dimers (e.g., dimers of Fc fusion protein homodimers, trimers of Fc fusion protein homodimers, tetramers of Fc fusion protein homodimers, or higher order aggregates containing five or more Fc fusion protein homodimers). It may be expected that multimeric forms of Fc fusion proteins may have different physical, stability, or pharmacologic activities from that of the insulin-Fc fusion protein homodimers.

#### ***Insulin Polypeptide***

[0057] The insulin polypeptide of the invention is defined in claim 1. An insulin polypeptide may be, for example, an insulin or insulin analog produced by  $\beta$ -cells in the islets of Langerhans within the pancreas. Insulin functions by regulating the absorption of glucose from the blood. Upon a stimulus, such as increased protein and glucose levels, insulin is released from  $\beta$ -cells and binds to the IR, initiating a signal cascade that affects many aspects of mammalian (e.g.,

human) metabolism. Disruption of this process is directly related to several diseases, notably diabetes, insulinoma, insulin resistance, metabolic syndromes, and polycystic ovary syndrome. Insulin analogs of the present disclosure may be related to the structure of insulin yet contain one or more modifications. In some embodiments, the insulin analog comprises at least one amino acid substitution, deletion, addition or chemical modification relative to insulin, which may impact a particular feature or characteristic of the insulin-Fc fusion protein configuration. For example, the modifications or alterations described herein may impact the structure, stability, pH sensitivity, bioactivity, or binding affinity of the insulin-Fc fusion protein configuration to a cell surface receptor (e.g. an insulin hormone receptor) relative to a reference standard.

**[0058]** The amino acid sequence of insulin is strongly conserved throughout evolution, particularly in vertebrates. For example, native canine and porcine insulins differ by only one amino acid from human insulin, native bovine insulin differs by only three amino acids from human insulin, and native feline insulin differs by just four amino acids from human insulin. As used herein, the terms "B-chain or B-chain analog", "C-peptide" or "C-chain", and "A-chain or A-chain analog" refer to the peptide segments of an insulin polypeptide as illustrated in FIG. 1. Insulin is a 51 amino acid hormone containing two peptide chains (i.e., a B-chain and an A-chain) connected via disulfide bonds (e.g., disulfide bonds formed by one or more B-chain cysteine side chain thiols and one or more A-chain cysteine side chain thiols). The A-chain of insulin is 21 amino acids in length and the B-chain of insulin is 30 amino acids in length. In the native form of insulin, the A-chain contains one intrachain disulfide bond formed by two A-chain cysteine side chain thiols. For reference purposes, the sequences for the human insulin B-chain of SEQ ID NO: 1 and the human insulin A-chain of SEQ ID NO: 2 are shown below:

FVNQHLCGSHLVEALYLVCGERGFFYTPKT (SEQ ID NO: 1) GIVEQCCTSICSLYQLENYCN (SEQ ID NO: 2).

**[0059]** As used herein, the term "insulin" or "insulin polypeptide" encompasses mature insulin, proinsulin, and naturally occurring insulin, or analogs thereof. In embodiments, an insulin polypeptide can be a full-length insulin polypeptide or a fragment thereof. In embodiments, an insulin polypeptide can comprise one or more fragments from mature insulin, proinsulin, or naturally occurring insulin.

**[0060]** Insulin is normally constructed as a N-terminus--B-chain:C-chain:A-chain--C-terminus polypeptide, wherein the C-chain is cleaved in order to make it bioactive. For reference purposes, the sequence of the entire human insulin molecule including the C-chain (i.e., human proinsulin) is shown below with the C-chain underlined:

FVNQHLCGSHLVEALYLVCGERGFFYTPKTRREAEDLQVGQVELGGPGAGSLQPLALEGSLQK  
RGIVEQCCTSICSLYQLENYCN (SEQ ID NO: 3).

The transformation of the single-chain insulin polypeptide into a bioactive two-chain polypeptide is normally accomplished within the  $\beta$ -cells of the islets of Langerhans prior to glucose-stimulated insulin secretion by two endoproteases, Type I endoproteases, PC1 and PC3, that disrupt the C peptide-B chain connection and PC2, and a Type II endoprotease, that cleaves the C peptide-A chain bond at exactly the right sites. However, cell systems used for the biosynthesis of therapeutic molecules such as insulin (e.g. bacteria, yeast, and mammalian (e.g. HEK and CHO) cell systems) do not possess this pathway, and therefore the transformation must take place after expression and harvesting of the single chain polypeptide using chemical or enzymatic methods. All the known techniques for cleaving the C-chain after expression and harvesting rely on first modifying the C-chain such that it terminates in a lysine just before the N-terminus of the A-chain. Then, using an enzyme selected from the trypsin or Lys-C families, which clips peptide bonds specifically at the C-termini of lysine residues, the single chain-insulin polypeptide is cleaved at the C-terminal lysine of the C-chain and at the C-terminal lysine at the 29<sup>th</sup> position from the N-terminus of the B-chain. In some cases, the resulting bioactive two-chain insulin is used without reattaching the clipped amino acid at the 30<sup>th</sup> position from the N-terminus of the B-chain, and in some cases the clipped amino acid at the 30<sup>th</sup> position from the N-terminus of the B-chain is added back to the molecule using an additional enzymatic method. Such a process works well with insulin because it contains only one lysine in its entire two chain polypeptide form. However, this process cannot be used on the insulin-Fc fusion proteins contained herein, because all known Fc fragments contain multiple lysine residues. The enzymatic cleavage process would, therefore, digest the Fc fragment into non-functional parts, thereby eliminating the ability of the Fc fragment to prolong the action of the insulin polypeptide in vivo. Therefore, an insulin-Fc fusion protein of the present invention must comprise an insulin polypeptide that does not require C-chain cleavage and is therefore bioactive in its single chain form.

**[0061]** A number of bioactive single chain insulin polypeptides have been described in the art. In all cases, the single chain insulin polypeptides contain C-chains of specific length and composition as well as A-chains and B-chains mutated at specific amino acid sites in order to achieve electrostatic balance, prevent aggregation, and enhance IR binding and/or downstream signaling to achieve bioactivity at levels comparable to that of the native two-chain insulin. Herein, the location of mutations on peptide segments are notated using the name of the segment (e.g., B-chain, C-chain, A-chain) and the number of the amino acid counting from the N-terminus of the segment. For example, the notation "B16" refers to the 16<sup>th</sup> amino acid from the N-terminus of the amino acid sequence of the B-chain. The notation "A8" refers to the

8<sup>th</sup> amino acid from the N-terminus of the A-chain. Furthermore, if an amino acid is mutated from its native form to a new amino acid at a particular location, the location is appended with the one letter amino acid code for the new amino acid. For example, B16A refers to an alanine mutation at the 16<sup>th</sup> amino acid from the N-terminus of the amino acid sequence of the B-chain and A8H refers to a histidine mutation at the 8<sup>th</sup> amino acid from the N-terminus of the amino acid sequence of the A-chain.

**[0062]** US9855318B2 describes a single chain insulin analog with a C-chain ("first linker") of the sequence GGS<sup>GGGG</sup> (SEQ ID NO: 72), substitutions in the A-chain, and substitutions and deletions in the B-chain (non-native amino acids are underlined and deleted native amino acids represented with an underlined Z):

FVNQHLCGSHLVEALELVCGERGFHY<sup>GGGGGGSGGGGG</sup>GIVEQCCTSTC<sup>SLD</sup>QLENYC<sup>Z</sup> (SEQ ID NO: 7\_NULL).

**[0063]** The following is a restatement of the sequence shown above but with the absent amino acids of symbol Z removed from the notation of the insulin polypeptide sequence. Again, as before, the non-native amino acids are underlined. Despite the two separate notations, the paired sequences refer to exactly the same insulin polypeptide.

FVNQHLCGSHLVEALELVCGERGFHY<sup>GGGGGGSGGGGG</sup>GIVEQCCTSTC<sup>SLD</sup>QLENYC (SEQ ID NO: 7)

**[0064]** In some embodiments, it was discovered unexpectedly that insulin-Fc fusion protein configurations wherein alanine is substituted for glutamic acid at position 16 from the N-terminus of the B-chain (i.e., B16) in SEQ ID NO: 7 to produce SEQ ID NO: 10, led to improved homodimer titers, IR binding affinity, and FcRn binding of the insulin-Fc fusion protein while preserving reduced immunogenicity as measured by low Fc(gamma)RI and C1q binding affinities. This particular amino acid substitution at B16 was originally motivated by the fact that an alanine in this position is known to be less capable of activating insulin-specific T cells (Alleva, D.G., Gaur, A., Jin, L., Wegmann, D., Gottlieb, P.A., Pahuja, A., Johnson, E.B., Motheral, T., Putnam, A., Crowe, P.D., Ling, N., Boehme, S.A., Conlon, P.J., (2002) Diabetes Vol. 51, No. 7 pp 2126-2134). SEQ ID NO: 10 is listed below with each of the non-native amino acids underlined:

FVNQHLCGSHLVEALALVCGERGFHY<sup>GGGGGGSGGGGG</sup>GIVEQCCTSTC<sup>SLD</sup>QLENYC (SEQ ID NO: 10).

#### Linker

**[0065]** The successful construction of a recombinantly made insulin-Fc fusion protein requires a linker connecting the insulin polypeptide to the Fc fragment. In embodiments, insulin-Fc fusion protein configurations described herein comprise a peptide linker between the insulin polypeptide and the Fc fragment comprising amino acids (e.g., natural or unnatural amino acids). In embodiments, the peptide linker can be encoded by a nucleic acid molecule, for example such that a single nucleic acid molecule can encode the various peptides within an insulin polypeptide as well as the peptide linker and the Fc fragment. The choice of peptide linker (for example, the length, composition, hydrophobicity, and secondary structure) could impact the manufacturability of the insulin-Fc fusion protein configuration (i.e., the homodimer titer), the chemical and enzymatic stability, the bioactivity (i.e., the NAOC value), parameters that correlate with bioactivity (i.e., the FcRn assay EC50 value), and the immunogenicity of the insulin-Fc fusion protein (Chen, X., Zaro, J., Shen, W.C., Adv Drug Deliv Rev. 2013 October 15; 65(10): 1357-1369). Table 1 lists several linkers used in the design of insulin-Fc fusion protein configurations with the goal of improving the homodimer titer and the bioactivity.

| Table 1: Peptide Linker Between A-chain and Fc Fragment in an Insulin-Fc Fusion Protein |                                                 |
|-----------------------------------------------------------------------------------------|-------------------------------------------------|
|                                                                                         | GGGGAGGGG (SEQ ID NO: 11) (not claimed)         |
|                                                                                         | GGGGSGGGG (SEQ ID NO: 12) (not claimed)         |
|                                                                                         | GGGGAGGGG (SEQ ID NO: 64) (not claimed)         |
|                                                                                         | GGGGSGGGSGGGSGGGG (SEQ ID NO: 65) (not claimed) |
|                                                                                         | GGGGKGGGKGGGKGGG (SEQ ID NO: 66) (not claimed)  |
|                                                                                         | GGGGAGGGAGGGAGGGG (SEQ ID NO: 67)               |
|                                                                                         | GGGGQGGGQGGGQGGG (SEQ ID NO: 13)                |
|                                                                                         | SGGGQGGGQGGGQGGG (SEQ ID NO: 68) (not claimed)  |
|                                                                                         | HGGGQGGGQGGGQGGG (SEQ ID NO: 69) (not claimed)  |
|                                                                                         | PGGGQGGGQGGGQGGG (SEQ ID NO: 70) (not claimed)  |
|                                                                                         | GGGGQGGGQGGGQGGG (SEQ ID NO: 99)                |

**[0066]** As defined in claim 1, the peptide linker comprises the sequence: GGGGQGGGQGGGQGGG (SEQ ID NO: 13); GGGGAGGGAGGGAGGGG (SEQ ID NO: 67) or the sequence: GGGGQGGGQGGGQGGG-

GGGGQGGGG (SEQ ID NO: 99) .

**[0067]** In constructing a recombinantly made insulin-Fc fusion protein configuration with a peptide linker like the one of SEQ ID NO: 13, attention must be paid to the possibility of unwanted enzymatic cleavage between the C-terminus of the insulin A-chain and the N-terminus of the peptide linker. Cleavage of the linker and Fc-fragment from the insulin polypeptide would render the insulin-Fc fusion protein configuration incapable of providing an extended duration of bioactivity. A known enzymatic cleavage site exists between asparagine-glycine bonds (Vlasak, J., Ionescu, R., (2011) MAbs Vol. 3, No. 3 pp 253-263). In many peptide linker embodiments, including the preferred peptide linker of SEQ ID NO: 13, the N-terminal amino acid is a glycine. Furthermore, the C-terminus of the insulin A-chain (i.e. the 21st amino acid from the N-terminus of the A-chain (i.e., A21)) is an asparagine. Therefore, the A21 asparagine is omitted in the insulin polypeptides of SEQ ID NO: 7, SEQ ID NO: 9, and SEQ ID NO: 10 to eliminate the potentially enzymatically cleavable asparagine-glycine bond that would form between the A-chain and the peptide linker in insulin-Fc fusion protein configurations. Unexpectedly, an insulin-Fc fusion protein configuration constructed from the insulin polypeptide of SEQ ID NO: 8 (not claimed), which retains the asparagine at the C-terminus of the A-chain, demonstrates manufacturability in mammalian cells with an acceptable homodimer titer (i.e., a homodimer titer for the canine insulin-Fc fusion protein of greater than 50 mg/L), an acceptable bioactivity in vivo in dogs (i.e., a NAOC greater than 150 %FBGL·days·kg/mg), and sustained levels of bioactivity after multiple doses (i.e., a NAOCR value after the third injection in the dog of greater than 0.5). The results indicate that, contrary to expectations based on prior teachings, there is no risk of enzymatic cleavage or deactivation of insulin-Fc fusion protein configurations containing the asparagine-glycine link between the insulin polypeptide and peptide linker, at least for insulin-Fc fusion protein configurations comprising the Fc fragment sequences disclosed herein.

**[0068]** It was discovered that for the same insulin polypeptide and Fc fragment compositions, mutating the Glutamine (Q) in SEQ ID NO: 13 to Alanine (A), generating the peptide linker of GGGGAGGGGAGGGGAGGGGG (SEQ ID NO: 67), resulted in an insulin-Fc fusion protein configuration with higher homodimer titer, increased binding affinity to the IR, and increased binding affinity to the FcRn receptor.

**[0069]** It was discovered that for the same insulin polypeptide and Fc fragment compositions, the peptide linker of SEQ ID NO: 67 could be shortened without significantly affecting the homodimer titer or the binding affinity to the FcRn receptor of the insulin-Fc fusion protein, but with a 60% increase in the IR assay IC50 value. The shortened peptide linker comprises the sequence: GGGGAGGGG (SEQ ID NO: 11) (not claimed).

#### *Fc Fragment*

**[0070]** The terms "Fc fragment," "Fc region," "Fc domain," or "Fc polypeptide," are used herein to define a C-terminal region of an immunoglobulin heavy chain. The Fc fragment, region, domain or polypeptide may be a native sequence Fc region or a variant/mutant Fc region. Although the boundaries of the Fc region of an immunoglobulin heavy chain may vary, they generally comprise some or all of the hinge region of the heavy chain, the CH2 region of the heavy chain, and the CH3 region of the heavy chain. The hinge region of a canine or human Fc fragment comprise amino acid sequences that connect the CH1 domain of the heavy chain to the CH2 region of the heavy chain and which contain one or more cysteines that form one or more interheavy chain disulfide bridges to form a homodimer of an Fc fusion protein from two identical but separate monomers of the Fc fusion protein. The hinge region may comprise all or part of a naturally occurring amino acid sequence or a non-naturally occurring amino acid sequence.

**[0071]** An Fc receptor (FcR) refers to a receptor that binds to an Fc fragment or to the Fc region of an antibody. As disclosed, the FcR is a native sequence of the canine or human FcR. In embodiments, the FcR is one which binds an Fc fragment or the Fc region of an IgG antibody (a gamma receptor) and includes without limitation, receptors of the Fc(gamma)RI, Fc(gamma)RIIa, Fc(gamma)RIIb, and Fc(gamma)RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. "FcR" also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgG molecules to the fetus (Guyer et al., 1976 J. Immunol., 117:587; and Kim et al., 1994, J. Immunol., 24:249) and is also responsible for the prolonged in vivo elimination half-lives of antibodies and Fc-fusion proteins in vivo. Those skilled in the art will understand that mammalian FcR from one species (e.g., FcR of human origin) are capable of in vitro binding of insulin-Fc fusion proteins comprising Fc fragments of the same species (e.g., of human origin) and also are sometimes capable of in vitro binding of Fc fragments from another mammalian species (e.g. of canine origin). FcR of human origin are used in vitro (e.g., in an assay) to measure the binding properties of insulin-Fc fusion protein configurations comprising Fc fragments of human or canine origin so as to assess their FcR binding properties. FcR of canine origin are used in vitro (e.g., in an assay) to measure the binding of insulin-Fc fusion protein configurations comprising Fc fragments of canine origin.

**[0072]** The C-terminal lysine that is often found in native canine and human IgG isotype Fc fragment amino acid sequences (i.e., the lysine that represents the last amino acid of the Fc fragment sequence) is omitted to prevent the accidental production of unwanted amino acid sequence variants during manufacturing (e.g., Fc fragments containing the C-terminal lysine becoming mixed with Fc fragments where the C-terminal lysine is omitted, which can occur during

production of the desired protein within cells (Dick, LW., (2008) Biotechnol Bioeng. Aug 15;100(6) pp1132-43).

**[0073]** In canine insulin-Fc fusion protein embodiments, the Fc fragment comprises the Fc region (e.g., hinge region, CH2 domain, and CH3 domain) of a canine IgGA Fc fragment (SEQ ID NO: 14), a canine IgGB Fc fragment (SEQ ID NO: 15), a canine IgGC Fc fragment (SEQ ID NO: 16), or a canine IgGD Fc fragment (SEQ ID NO: 17). Therefore, in canine insulin-Fc fusion protein configurations, the canine Fc fragment sequences lacking a C-terminal lysine are:

RCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPEVTCVVLDLGREDPEVQISWFVDGKEVHTAKT  
 QSREEQFNGTYRVVSVLPIEHQDWLTGKEFKCRVNHIDLPSPIERTISKARGRAHKPSVYVLPPSPK  
 ELSSSDTVSITCLIKDFYPPDIDVEWQSNQQEPERKHRMTPPQLDEDGSYFLYSKLSVDKSRWQQ  
 GDPFTCAVMHETLQNHYTDLSLSHSPG (SEQ ID NO: 14) (not claimed)

DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQMQTAKTQ  
 PREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
 ELSKNTVSLTCLIKDFFPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
 DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 15) (not claimed)

CNNCPCPGCGLLGGPSVFIFPPKPKDILVTARTPTVTCVVVDLDPENPEVQISWFVDSKQVQTANT  
 QPREEQSNGTYRVVSVLPIGHQDWLSGKQFKCKVNNKALPSPIEEIISKTPGQAHQPNVYVLPPSR  
 DEMSKNTVTLTCLVKDFFPPEIDVEWQSNQQEPESKYRMTTPPQLDEDGSYFLYSKLSVDKSRWQ  
 RGDTFICAVMHEALHNHYTQISLSHSPG (SEQ ID NO: 16) (not claimed)

CISPCPVPESLGGPSVFIFPPKPKDILRITRTPEITCVVLDLGREDPEVQISWFVDGKEVHTAKTQPRE  
 QQFNSTYRVVSVLPIEHQDWLTGKEFKCRVNHIGLPSPIERTISKARGQAHQPSVYVLPPSPKELSS  
 SDTVTLTCLIKDFFPPEIDVEWQSNQQEPESKYHTTAPQLDEDGSYFLYSKLSVDKSRWQQGDTF  
 TCAVMHEALQNHYTDLSLSHSPG (SEQ ID NO: 17) (not claimed).

In human insulin-Fc fusion protein configurations, the human Fc fragment sequences lacking a C-terminal lysine are:

DKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH  
 NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY  
 TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSR  
 RWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 73) (not claimed)

ECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTK  
 PREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSRE  
 EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQG  
 NVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 74) (not claimed).

**[0074]** For dogs, the canine IgGA is preferable to minimize any unwanted immunogenicity, due to the IgGA isotype's lack of Fc(gamma) effector function in dogs (much like the human IgG2 isotype in humans). However, in an insulin-Fc

fusion protein embodiment (not claimed) containing the insulin polypeptide of SEQ ID NO: 4 and the peptide linker of SEQ ID NO: 11, it was unexpectedly discovered that the insulin-Fc fusion protein configuration comprising the canine IgGA fragment (SEQ ID NO: 14) was highly aggregated with low titers of the desired homodimer (i.e., homodimer titers of the canine insulin-Fc fusion protein of less than 50 mg/L). Furthermore, the insulin-Fc fusion protein configuration was non-bioactive in dogs (i.e., the NAOC value was less than 150 %FBGL-days·kg/mg), presumably due to its high level of aggregation (e.g. low %homodimer). Despite mutating the insulin polypeptide of SEQ ID NO: 4, the canine IgGA Fc fragment (SEQ ID NO: 14), and/or the linker, there was no insulin-Fc fusion protein embodiment comprising the canine IgGA Fc fragment that exhibited a low enough degree of aggregation and a high enough titer of the desired homodimer. However, replacing of the canine IgGA Fc fragment (SEQ ID NO: 14) with the canine IgGB Fc fragment (SEQ ID NO: 15) in insulin-Fc fusion protein configurations yielded significantly less aggregated compounds with a comparatively high titer of the desired homodimer. Furthermore, the insulin-Fc fusion protein configuration containing the insulin polypeptide of SEQ ID NO: 4 and the canine IgGB Fc fragment (SEQ ID NO: 15) was bioactive in dogs, exhibiting glucose lowering bioactivity over multiple days (i.e., the NAOC value was greater than 150 %FBGL-days·kg/mg).

**[0075]** The preference for the canine IgGB Fc fragment over the canine IgGA Fc fragment was confirmed in insulin-Fc fusion protein configurations (not claimed) containing the insulin polypeptide of SEQ ID NO: 7 and the peptide linker of SEQ ID NO: 13, both of which vary considerably from the insulin polypeptide of SEQ ID NO: 4 and the peptide linker of SEQ ID NO: 11. Insulin-Fc fusion protein configurations containing the insulin polypeptide of SEQ ID NO: 7 and the peptide linker of SEQ ID NO: 13 were synthesized using Fc fragments from the canine IgGA (SEQ ID NO: 14), canine IgGB (SEQ ID NO: 15), canine IgGC (SEQ ID NO: 16), or canine IgGD (SEQ ID NO: 17) immunoglobulins. Using a conventional purification method, only the insulin-Fc fusion protein configurations comprising the canine IgGA and the canine IgGB showed any appreciable protein yields. However as previously seen, the canine IgGA configuration of the insulin-Fc fusion protein was highly aggregated with low levels of bioactivity, whereas the canine IgGB configuration of the insulin-Fc fusion protein exhibited a low degree of aggregation (i.e. high %homodimer), a high titer of the desired homodimer (i.e., a homodimer titer of the canine insulin-Fc fusion protein configurations of greater than 50 mg/L), and appreciable levels of long-duration glucose lowering bioactivity in dogs (i.e., the NAOC value was greater than 150 %FBGL-days·kg/mg). Using an alternative purification method, the canine IgGC configuration of the insulin-Fc fusion protein was recovered with low degrees of aggregation, but it was minimally bioactive in dogs (i.e., the NAOC value was less than 150 %FBGL-days·kg/mg), presumably due to its low affinity for the FcRn receptor. Therefore, with respect to a dog-specific product, the canine IgGB (SEQ ID NO: 15) is the preferred Fc fragment for all insulin-Fc fusion protein configurations used in dogs, regardless of the choice of insulin polypeptide.

**[0076]** Given that the canine IgGB isotype interacts with the canine Fc(gamma) receptor with higher affinities than the canine IgGA isotype, there could be a risk of unwanted immunogenicity after repeated injections of an insulin-Fc fusion protein configuration comprising canine IgGB. Therefore, various mutations to the canine IgGB Fc fragment were investigated in an effort to preserve greater insulin-Fc fusion protein homodimer titers while decreasing the affinity for the canine Fc(gamma)RI receptor.

**[0077]** One method for reducing the Fc(gamma)RI interaction involves deglycosylating or preventing the glycosylation of the Fc fragment during synthesis of the insulin-Fc fusion protein in the host cell. Each IgG fragment contains a conserved asparagine (N)-glycosylation site in the CH2 domain of each heavy chain of the Fc region. Herein, the notation used to refer to the conserved N-glycosylation site is "cNg". One way to remove the attached glycan from a synthesized insulin-Fc fusion protein is to mutate the cNg site to prevent the attachment of glycans altogether during production in the host cell. Herein, the notation used to describe a cNg mutation is cNg-(substituted amino acid). For example, if the asparagine at the cNg site is mutated to serine, this mutation is notated as "cNg-S".

**[0078]** The absolute position of the cNg site from the N-terminus of the B-chain of the insulin-Fc fusion protein configuration varies depending on the length of the insulin polypeptide, the length of the linker, and any omitted amino acids in the Fc fragment prior to the cNg site. Herein, the notation used to refer to the absolute position of the cNg site in a given insulin-Fc fusion protein sequence (as measured counting from the N-terminus of the B-chain of the insulin-Fc fusion protein) is "NB(number)". For example, if the cNg site is found at the 155<sup>th</sup> amino acid position as counted from the N-terminus of the B-chain, the absolute position of this site is referred to as "cNg-NB155". As a further example, if the cNg site is found at the 155<sup>th</sup> amino acid position as counted from the N-terminus of the B-chain, and the asparagine at this site is mutated to serine, this mutation is noted as "cNg-NB155-S".

**[0079]** In insulin-Fc fusion protein embodiments (not claimed) containing the insulin polypeptide of SEQ ID NO: 4 and the canine IgGB Fc fragment with the cNg-Q, cNg-S, cNg-D, and cNg-K mutations, it was unexpectedly discovered that only the compounds containing the cNg-K and cNg-S mutations exhibited the requisite homodimer titer of the canine insulin-Fc fusion protein of greater than 50 mg/L and lowest Fc(gamma)RI binding affinities. On the other hand, in an insulin-Fc fusion protein embodiment containing the insulin polypeptide of SEQ ID NO: 7 and the canine IgGB Fc fragment with the cNg-S mutation, it was unexpectedly discovered that the resulting insulin-Fc fusion protein was significantly less bioactive in dogs compared to the counterpart insulin-Fc fusion protein configuration but comprising the native

canine IgGB Fc (i.e., the NAOC value was significantly lower for the insulin-Fc fusion protein counterpart containing the native glycosylation site amino acid, e.g., cNg-N). The bioactivity of the insulin-Fc fusion protein was unexpectedly restored in the cNg-S mutant (i.e., the NAOC value increased significantly) when the B 16 amino acid was mutated to alanine as described above for insulin polypeptide SEQ ID NO: 10. Taken together, there is an unexpected and significant interaction between the choice of cNg mutation on the Fc fragment of the insulin-Fc fusion protein configuration and the composition of the insulin polypeptide such that experimentation is required to identify the preferred insulin-Fc fusion protein embodiments. In specific insulin-Fc fusion protein embodiments, the canine IgGB Fc mutant containing the cNg-S mutation is preferred and the sequence with underlined cNg-S is shown as:

DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKTQ  
 PREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
 ELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
 DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 18) (not claimed).

**[0080]** Generally, the human Fc IgG2 isotype is preferred over the other isotypes due to its lack of Fc(gamma) effector function and therefore its lower propensity to induce unwanted immunogenicity. As an illustration, in one execution of the Fc(gamma)RI binding and C1q binding ELISAs, the insulin-Fc fusion protein embodiment of SEQ ID NO: 76 comprising the human IgG1 fragment exhibited a human Fc(gamma)RI binding assay OD450 at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL of **2.078**, and exhibited a human C1q binding assay OD450 at a biotinylated-C1q concentration of 1000 ng/mL of **3.006**, such high values indicating a likelihood of SEQ ID NO: 76 exhibiting immunogenicity in patients. In contrast, in the same execution of the Fc(gamma)RI binding and C1q binding ELISAs, the insulin-Fc fusion protein configuration (SEQ ID NO: 75) comprising the insulin polypeptide of SEQ ID NO: 7, the peptide linker of SEQ ID NO: 13, and the human IgG2 Fc fragment (SEQ ID NO: 74) exhibited a human Fc(gamma)RI binding assay OD450 at a biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL of **0.093** and a human C1q binding assay OD450 at a biotinylated-C1q concentration of 1,000 ng/mL of **0.928**, indicating a significantly reduced likelihood of SEQ ID NO: 75 exhibiting immunogenicity in patients.

**[0081]** The absolute value of OD450 measurements for an insulin-Fc fusion protein configuration can vary from one execution of the ELISA to the next, for example due to small variations in execution of the assay. This makes comparisons of absolute value measurements of OD450 for different insulin-Fc fusion protein configurations across different ELISA executions less reliable, even when the concentration of the insulin-Fc fusion configuration under test is held constant. In contrast, the ratio of the OD450 measurement of one insulin-Fc fusion protein configuration in an ELISA execution with the OD450 measurement of a second insulin-Fc fusion protein configuration from the same ELISA execution (again holding the insulin-Fc fusion protein concentrations constant) will be relatively stable for the same two insulin-Fc fusion protein configurations across different ELISA executions. Accordingly, insulin-Fc fusion protein design goals for the OD450 for a human Fc(gamma)RI binding assay and C1q binding assay have been expressed as OD 450 Ratios, where the ratio is made of the absolute OD450 value of the insulin-Fc fusion protein under analysis to the absolute OD 450 value of a reference insulin-Fc fusion protein configuration, where both measurements are made within the same ELISA experiment. The insulin-Fc fusion protein configuration sample biotinylated-Fc(gamma)RI concentration in the Fc(gamma)RI binding ELISA OD450 is set to 3000 ng/mL and the insulin-Fc fusion protein configuration sample biotinylated-C1q concentration in the C1q binding ELISA OD450 is set to 1000 ng/mL for all samples tested.

**[0082]** The insulin-Fc fusion protein configuration of SEQ ID NO: 76 (comprising the human IgG1 Fc fragment) is used as the reference insulin-Fc fusion protein for the Fc(gamma)RI binding ELISA and C1q binding ELISA OD450 Ratio calculations. The Fc(gamma)RI binding ELISA OD450 Ratio for the insulin-Fc fusion protein configuration of SEQ ID NO: 75 (comprising the human IgG2 Fc fragment) using the measured OD450 values given above is:

$$Fc(\gamma)RI \text{ binding ELISA OD450 Ratio}_{\left(\frac{SEQ ID NO:75}{SEQ ID NO:76}\right)} = \frac{0.093}{2.078} = 0.045$$

The C1q binding ELISA OD450 Ratio for the insulin-Fc fusion protein configuration of SEQ ID NO: 75 using the measured OD450 values given above is:

$$C1q \text{ binding ELISA OD450 Ratio}_{\left(\frac{SEQ ID NO:75}{SEQ ID NO:76}\right)} = \frac{0.928}{3.06} = 0.309$$

**[0083]** The Fc(gamma)RI binding ELISA OD450 Ratio and the C1q binding ELISA OD450 Ratio of the insulin-Fc fusion protein configuration of SEQ ID NO: 75 (hIgG2) with respect to the insulin-Fc fusion protein configuration of SEQ ID NO: 76 (hIgG 1) represents an aspirational benchmark that was used in creating the design goals for evaluating the Fc(gamma)RI binding and C1q binding for different insulin-Fc fusion configurations in order to reduce the propensity for unwanted immunogenicity. Accordingly, the design goal established for human Fc(gamma)RI binding (where the biotinylated-Fc(gamma)RI concentration of the insulin-Fc fusion protein under test is 3000 ng/mL) is an OD450 Ratio < 0.50 and the design goal established for human C1q binding (where the biotinylated-C1q concentration of the insulin-Fc fusion protein under test is 1000 ng/mL) is an OD450 ratio <0.35.

**[0084]** The resulting average homodimer titer from two separate syntheses of this insulin-Fc fusion protein configuration (SEQ ID NO: 75) comprising the human IgG2 Fc fragment was 117 mg/L.

**[0085]** By comparison, in an insulin-Fc fusion protein embodiment (SEQ ID NO: 76) (not claimed) comprising the same insulin polypeptide of SEQ ID NO: 7, the same peptide linker of SEQ ID NO: 13, and the human IgG1 fragment (SEQ ID NO: 73) it was found that the insulin-Fc fusion protein exhibited an average homodimer titer from two separate syntheses of 180 mg/L, which was over 50% greater than that obtained for the analogous insulin-Fc fusion protein configuration comprising the human IgG2 fragment. Given that the human IgG1 isotype interacts with the human Fc(gamma)RI receptor with higher affinities than the human IgG2 isotype, there could be a risk of unwanted immunogenicity after repeated injections of an insulin-Fc fusion protein configuration comprising human IgG1.

**[0086]** Therefore, various mutations to the human IgG1 Fc fragment were investigated in an effort to preserve greater insulin-Fc fusion protein homodimer titers while decreasing the affinity for the human Fc(gamma)RI receptor and C1q.

**[0087]** As discussed previously, one method for reducing the Fc(gamma)RI interaction involves deglycosylating or preventing the glycosylation of the Fc fragment during synthesis of the insulin-Fc fusion protein in the host cell. One way to remove the attached glycan from a synthesized insulin-Fc fusion protein is to mutate the cNg site to prevent the attachment of glycans altogether during production in the host cell.

**[0088]** The mutated human IgG1 Fc fragment generalized configuration with cNg site underlined is as follows:

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH  
NAKTKPREEQYX<sub>1</sub>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY  
TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKS RWQQGNVFSC-  
SVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 77), where in examples Xi is S, D (not claimed), A (not claimed), R or Q (not claimed).

**[0089]** The human IgG1 Fc fragment comprising the insulin polypeptide of SEQ ID NO: 7, the peptide linker of SEQ ID NO: 13, and the human IgG1 Fc fragment of SEQ ID NO: 77 where Xi is S (i.e., with the cNg-NB155-S mutation) is shown below:

FVNQHLCGSHLVEALELVCGERGEFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
EVKFNWYVDGVEVHNAKTKPREEQY<sub>2</sub>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD  
DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 91) (not  
claimed).

**[0090]** It was unexpectedly discovered that the non-glycosylated insulin-Fc fusion protein embodiment of SEQ ID NO: 91 gave improved homodimer titers relative to the glycosylated parent material (SEQ ID NO: 76). Additionally, the human Fc(gamma)RI assay OD450 Ratio at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL for the insulin-Fc fusion protein embodiment of SEQ ID NO: 91 against the reference insulin-Fc fusion protein of SEQ ID NO: 76 was less than 0.50 and the human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1,000 ng/mL for the insulin-Fc fusion protein configuration of SEQ ID NO: 91 against the reference insulin-Fc fusion protein of SEQ ID NO: 76 was less than 0.35. However, the resulting non-glycosylated insulin-Fc fusion protein of SEQ ID NO: 91 with respect to its parent glycosylated insulin-Fc fusion protein of SEQ ID NO: 76 exhibited decreased IR binding (increased IR assay IC50 value) and decreased FcRn binding affinity (increased EC50 value), indicating a high likelihood of an unacceptable reduction in bioactivity with residence time in vivo respectively.

#### *Insulin-Fc Fusion Proteins*

**[0091]** Provided herein are insulin-Fc fusion protein configurations comprising an insulin polypeptide, an Fc fragment,

and a linker between the insulin polypeptide and the Fc fragment. The insulin polypeptide comprises domains in the following orientation from N- to C-termini: (N-terminus)--B-chain--C-chain--A-chain--(C-terminus). In embodiments, the insulin polypeptide is located on the N-terminal side of the Fc fragment. In embodiments, the fusion protein comprises domains in the following orientation from N- to C-termini: (N-terminus)--insulin polypeptide--linker--Fc fragment--(C-terminus) (e.g., (N-terminus)--B-chain--C-chain--A-chain--linker--Fc fragment--(C-terminus)) as illustrated in FIG. 1.

### Canine Insulin-Fc Fusion Proteins

**[0092]** In embodiments (not claimed), the non-immunogenic, bioactive insulin polypeptide of SEQ ID NO: 5 was combined with the preferred canine IgGB Fc fragment of SEQ ID NO: 15 using the preferred linker of SEQ ID NO: 13 to produce a family of high homodimer titer-yielding, non-aggregated, bioactive, non-immunogenic canine insulin-Fc fusion configurations of SEQ ID NO: 20 that exhibit homodimer titers greater than 50 mg/L, NAOC values greater than 150 %FBGL·days·kg/mg in dogs, and NAOCR values greater than 0.5 after the third injection in a series of repeated injections in dogs. The following shows SEQ ID NO: 20 with non-native amino acids underlined:

FVNQHLCGSX<sub>1</sub>LVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCX<sub>2</sub>STCSLDQLENYCX<sub>3</sub>GGGG  
 QGGGGGGGGGGGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPE  
 VQISWFDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTIS  
 KARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDED  
 GSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 20), where X<sub>1</sub> is not D, X<sub>2</sub> is not H,  
 and X<sub>3</sub> is absent or N (not claimed).

**[0093]** In canine insulin-Fc fusion protein embodiments comprising SEQ ID NO: 20, X<sub>1</sub> is H, X<sub>2</sub> is T, and X<sub>3</sub> is absent or N. The selections produce the high homodimer titer-yielding, non-aggregated, bioactive, non-immunogenic canine insulin-Fc fusion protein configurations of SEQ ID NO: 21 that exhibit homodimer titers greater than 50 mg/L, NAOC values greater than 150 %FBGL·days·kg/mg in dogs, and NAOCR values greater than 0.5 after the third injection in a series of repeated injections in dogs. The following shows SEQ ID NO: 21 with non-native amino acids underlined:

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCX<sub>3</sub>GGGGG  
 QGGGGGGGGGGGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEV  
 QISWFDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISK  
 ARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDG  
 SYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 21), where X<sub>3</sub> is absent or N (not  
 claimed).

**[0094]** In embodiments (not claimed), X<sub>3</sub> is absent in SEQ ID NO: 21 to produce the high homodimer-yielding, non-aggregated, bioactive, non-immunogenic canine insulin-Fc fusion protein of SEQ ID NO: 26 that exhibits a homodimer titer greater than 50 mg/L, a NAOC value greater than 150 %FBGL·days·kg/mg in dogs, and a NAOCR value greater than 0.5 after the third injection in a series of repeated injections in dogs. The following shows SEQ ID NO: 26 with non-native amino acids underlined:

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGGGG  
 GGGGGGGGGGGGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQIS  
 WFDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
 GQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYF  
 LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 26) (not claimed).

**[0095]** In embodiments (not claimed), X<sub>3</sub> is N in SEQ ID NO: 21 to produce the high homodimer titer-yielding, non-aggregated, bioactive, non-immunogenic canine insulin-Fc fusion protein of SEQ ID NO: 28 that exhibits a homodimer titer greater than 50 mg/L, a NAOC value greater than 150 %FBGL·days·kg/mg in dogs, and a NAOCR value greater than 0.5 after the third injection in a series of repeated injections in dogs. The following shows SEQ ID NO: 28 with non-native amino acids underlined:

FVNQBLCGSBLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCNGGGGGGGG  
 GGGGGGGGGGGGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQ  
 ISWFDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKA  
 RGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGS  
 YFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 28) (not claimed).

**[0096]** In embodiments (not claimed), the preferred non-glycosylated, cNg-S mutated canine IgGB Fc fragment of SEQ ID NO: 18 is combined with the preferred B 16A mutated insulin polypeptide sequence of SEQ ID NO: 9 using the

preferred linker of SEQ ID NO: 13 to produce a family of high homodimer titer-yielding, non-aggregated, bioactive, non-immunogenic canine insulin-Fc fusion proteins of SEQ ID NO: 22 that exhibit homodimer titers greater than 50 mg/L, NAOC values greater than 150 %FBGL·days·kg/mg in dogs, and NAOCR values greater than 0.5 after the third injection in a series of repeated injections in dogs. The following shows SEQ ID NO: 22 with non-native amino acids underlined:

5 FVNQHLCGSX<sub>1</sub>LVEALALVCGERGFHYGGGGGGSGGGGGIVEQCX<sub>2</sub>STCSLDQLENYCGGGGG  
 QGGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEV  
 QISWFVDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISK  
 ARGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDG  
 10 SYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 22), where X<sub>1</sub> is not D and X<sub>2</sub> is not H (not claimed).

[0097] In embodiments (not claimed), X<sub>1</sub> is H and X<sub>2</sub> is T in SEQ ID NO: 22 to produce the high homodimer titer-yielding, non-aggregated, bioactive, non-immunogenic canine insulin-Fc fusion protein of SEQ ID NO: 30 that exhibits a homodimer titer greater than 50 mg/L, a NAOC value greater than 150 %FBGL·days·kg/mg in dogs, and a NAOCR value greater than 0.5 after the third injection in a series of repeated injections in dogs. The following shows SEQ ID  
 15 NO: 30 with non-native amino acids underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
 GGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQ  
 20 ISWFVDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKA  
 RGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDGS  
 YFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 30) (not claimed).

[0098] The unexpected discovery of the highly manufacturable and effective ultra-long acting canine insulin-Fc fusion protein embodiment of SEQ ID NO: 30, led the inventors to attempt to produce similarly manufacturable and effective ultra-long acting insulin-Fc fusion protein configurations for human patients.

#### 30 Human Insulin-Fc Fusion Proteins

[0099] Further experiments were done to determine if other non-glycosylated insulin-Fc fusion protein configurations exhibited the same behavior. The human IgG1 Fc fragment generalized configuration with cNg site underlined is as follows:

35 DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH  
 NAKTKPREEQYX<sub>1</sub>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVY  
 TLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKS RWQQGNVFSC-  
 SVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 77), where X<sub>1</sub> is S, D (not claimed), A (not claimed), R or Q (not claimed).

[0100] Insulin-Fc fusion protein embodiments (not claimed) comprising the insulin polypeptide of SEQ ID NO: 7 (no B16A mutation) with the linker of SEQ ID NO: 13 and with the human IgG1 Fc fragment configuration with cNg mutation options to prevent glycosylation (SEQ ID NO: 77 where X<sub>1</sub> is S, D, A, R or Q) are shown below:

45 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
 EVKFNWYVDGVEVHNAKTKPREEQY<sub>2</sub>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD  
 50 DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 91) (not  
 claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP  
 5 EVKFNWYVDGVEVHNAKTKPREEQYDSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLY  
 10 SKLTVDKSRWQQGNVFSVMSHEALHNHYTQKSLSLSPG (SEQ ID NO: 92) (not  
 claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP  
 15 EVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLY  
 20 SKLTVDKSRWQQGNVFSVMSHEALHNHYTQKSLSLSPG (SEQ ID NO: 93) (not  
 claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP  
 25 EVKFNWYVDGVEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLY  
 30 SKLTVDKSRWQQGNVFSVMSHEALHNHYTQKSLSLSPG (SEQ ID NO: 94) (not  
 claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP  
 EVKFNWYVDGVEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 40 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLY  
 SKLTVDKSRWQQGNVFSVMSHEALHNHYTQKSLSLSPG (SEQ ID NO: 95) (not  
 claimed)

45 **[0101]** In embodiments (not claimed) containing the insulin polypeptide of SEQ ID NO: 7 using the preferred linker of  
 SEQ ID NO: 13 and the human IgG1 Fc fragment of SEQ ID NO: 77 wherein Xi is S (cNg-NB155-S - SEQ ID NO: 91),  
 D (cNg-NB155-D - SEQ ID NO: 92), A (cNg-NB155-A - SEQ ID NO: 93), R (cNg-NB155-R - SEQ ID NO: 94), or Q (cNg-  
 NB155-Q - SEQ ID NO: 95), it was unexpectedly discovered that each of these non-glycosylated insulin-Fc fusion  
 50 proteins gave improved homodimer titers relative to the glycosylated parent material (SEQ ID NO: 76). Additionally, for  
 each of the deglycosylated insulin-Fc fusion proteins of SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO:  
 94 and SEQ ID NO: 95, the human Fc(gamma)RI assay OD450 Ratios at a biotinylated-Fc(gamma)RI concentration of  
 3000 ng/mL (where the reference insulin-Fc fusion protein used in OD450 Ratio calculations is the insulin-Fc fusion  
 protein of SEQ ID NO: 76) were all less than 0.50 and the human C1q binding assay OD450 Ratios at a biotinylated-  
 55 C1q concentration of 1,000 ng/mL (where the reference insulin-Fc fusion protein used in OD450 Ratio calculations is  
 the insulin-Fc fusion protein of SEQ ID NO: 76) were also less than 0.35. However, the deglycosylated insulin-Fc fusion  
 proteins of SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94 and SEQ ID NO: 95 all demonstrated lower  
 IR binding affinities (higher IR assay IC50 values) and lower FcRn binding affinities (higher EC50 values) relative to the

glycosylated parent material (SEQ ID NO: 76), indicating a high likelihood of these compounds displaying an unacceptable reduction in bioactivity and residence time in vivo. Unexpectedly, in the embodiment (not claimed) comprising the insulin polypeptide of SEQ ID NO: 7 using the preferred linker of SEQ ID NO: 13 and the human IgG1 Fc fragment of SEQ ID NO: 77 wherein Xi is Q (SEQ ID NO: 95), the resulting non-glycosylated insulin-Fc fusion protein gave a homodimer titer of 136 mg/L, such that this insulin-Fc fusion protein configuration did not meet the design goal for homodimer titer of 150 mg/L.

**[0102]** FIG. 28 illustrates a side-by-side sequence comparison of SEQ ID NO: 75 (comprising native human IgG2) and SEQ ID NO: 76 (comprising native human IgG1), with sequences comprising the variants of SEQ ID NO: 77, that is SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, and SEQ ID NO: 95. "\*" represents complete homology across all sequences at a given sequence position, while ":", "." or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

**[0103]** Applying the learning from the unexpected results with respect to the canine insulin-Fc fusion protein configurations discussed previously, the 16<sup>th</sup> amino acid on the B-chain (B 16) of the insulin polypeptide in the insulin-Fc fusion protein embodiments of SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93 and SEQ ID NO: 94 was mutated to alanine, resulting in SEQ ID NO: 78 (with cNg-NB155-S), SEQ ID NO: 80 (with cNg-NB155-D) (not claimed), SEQ ID NO: 82 (with cNg-NB155-A) (not claimed), SEQ ID NO: 84 (with cNg-NB155-R). Acceptable insulin receptor binding was unexpectedly restored in the insulin-Fc fusion protein configurations of SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82 and SEQ ID NO: 84 (i.e., the IR binding and FcRn receptor binding affinities increased significantly) when the B16 amino acid was mutated to alanine as described above resulting in the insulin polypeptide of SEQ ID NO: 10, without compromising the increased homodimer titers or the reduction in Fc(gamma)RI and C1q binding affinities.

**[0104]** As defined in claim 2, the human insulin-Fc fusion protein configuration of SEQ ID NO: 78 comprises the insulin polypeptide of SEQ ID NO: 10, the linker of SEQ ID NO: 13, and the cNg-NB155-S mutant of the human IgG1 Fc fragment of SEQ ID NO: 77. The insulin-Fc fusion protein of SEQ ID NO: 78 exhibits a homodimer titer greater than 150 mg/L. The following shows the insulin-Fc fusion protein of SEQ ID NO: 78 with the B16A and cNg-NB155-S mutations underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
GGGGQGGGGQGGGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHED  
PEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 78).

**[0105]** Similar to the unexpected discovery with canine insulin-Fc fusion protein configurations, the bioactivity of the non-glycosylated insulin-Fc fusion protein embodiment of SEQ ID NO: 78 comprising the B 16A mutation and a human IgG1 Fc fragment with the cNg-NB 155-S mutation was unexpectedly restored (i.e., the IR binding and FcRn receptor binding affinities increased significantly) as compared to the bioactivity of the non-glycosylated insulin-Fc fusion protein embodiment of SEQ ID NO: 91 which lacked the B 16A mutation. The insulin-Fc fusion protein embodiment of SEQ ID NO: 78 demonstrated an IR assay IC50 value less than 2400 nM and more preferably less than 2000 nM, a human FcRn assay EC50 value less than 1500 ng/L and more preferably less than 1000 ng/mL, a human Fc(gamma)RI assay OD450 Ratio at a biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL less than 0.50 (where the reference insulin-Fc fusion protein used in OD450 Ratio calculations is the insulin-Fc fusion protein of SEQ ID NO: 76), and a human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1,000 ng/mL (where the reference insulin-Fc fusion protein used in OD450 Ratio calculations is the insulin-Fc fusion protein of SEQ ID NO: 76) less than 0.35.

**[0106]** The insulin-Fc fusion protein of SEQ ID NO: 80 (not claimed) comprises the insulin polypeptide of SEQ ID NO: 10, the linker of SEQ ID NO: 13, and the cNg-NB 155-D mutant of the Fc fragment of SEQ ID NO: 77 (Xi is D). The insulin-Fc fusion protein of SEQ ID NO: 80 exhibits a homodimer titer greater than 150 mg/L, an IR assay IC50 value less than 2400 nM, a human FcRn assay EC50 value less than 1000 ng/mL, a human Fc(gamma)RI assay OD450 Ratio at a biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL less than 0.50, and a human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1,000 ng/mL less than 0.35. The following shows SEQ ID NO: 80 with the B 16A and cNg-NB 155-D mutations underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
 GGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
 5 PEVKFNWYVDGVEVHNAKTKPREEQYDSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 80) (not  
 10 claimed).

**[0107]** The insulin-Fc fusion protein of SEQ ID NO: 82 comprises the insulin polypeptide of SEQ ID NO: 10, the linker of SEQ ID NO: 13, and the cNg-NB155-A mutant of the Fc fragment of SEQ ID NO: 77 (Xi is A). The insulin-Fc fusion protein of SEQ ID NO: 82 exhibits a homodimer titer greater than 150 mg/L, an IR assay IC50 value less than 2400 nM,  
 15 a human FcRn assay EC50 value less than 1000 ng/mL, a human Fc(gamma)RI assay OD450 Ratio at a biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL less than 0.50, and a human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1,000 ng/mL less than 0.35. The following shows SEQ ID NO: 82 with the B16A and cNg-NB155-A mutations underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
 GGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
 PEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 25 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 82) (not  
 claimed).

**[0108]** As defined in claim 4, the insulin-Fc fusion protein of SEQ ID NO: 84 comprises the insulin polypeptide of SEQ ID NO: 10, the linker of SEQ ID NO: 13, and the cNg-NB 155-R mutant of the Fc fragment of SEQ ID NO: 77 (Xi is R). The insulin-Fc fusion protein of SEQ ID NO: 84 exhibits a homodimer titer greater than 150 mg/L, an IR assay IC50 value less than 2400 nM, a human FcRn assay EC50 value less than 1000 ng/mL, a human Fc(gamma)RI assay OD450  
 35 Ratio at a biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL less than 0.50, and a human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1,000 ng/mL less than 0.35. The following shows SEQ ID NO: 84 with the B16A and cNg-NB155-R mutations underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
 GGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
 PEVKFNWYVDGVEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 45 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 84).

**[0109]** As was the case with SEQ ID NO: 78, the bioactivity of the non-glycosylated insulin-Fc fusion protein configurations comprising human IgG1 FC fragments with the cNg-NB155-D, cNg-NB155-A, cNg-NB155-R and cNg-NB155-Q mutations (SEQ ID NO: 92 (not claimed), SEQ ID NO: 93 (not claimed), SEQ ID NO: 94 (not claimed) and SEQ ID NO: 95 (not claimed)) was restored (i.e., the IR binding and FcRn receptor binding affinities increased significantly) when the 16<sup>th</sup> amino acid on the B-chain (B16) of the insulin polypeptide was mutated to alanine (resulting in the insulin polypeptide of SEQ ID NO: 10), without compromising the increased homodimer titers or the reduction in Fc(gamma)RI and C1q binding affinities of the non-glycosylated insulin-Fc fusion protein configurations of SEQ ID NO: 92, SEQ ID  
 55 NO: 93, and SEQ ID NO: 94 (as previously discussed, the homodimer titer of the insulin-Fc fusion protein configuration of SEQ ID NO: 95 did not meet the design goal of 150 mg/L).

**[0110]** However, the insulin-Fc fusion protein of SEQ ID NO: 86 comprising the insulin polypeptide of SEQ ID NO: 10, the linker of SEQ ID NO: 13, and the cNg-NB155-Q mutant of the Fc fragment of SEQ ID NO: 77 (Xi is Q) did not exhibit

the reduction in Fc(gamma)RI and C1q binding affinities of the non-glycosylated insulin-Fc fusion protein configuration of SEQ ID NO: 95, however the homodimer titer of the insulin-Fc fusion protein configuration of SEQ ID NO: 86 dropped further from that of SEQ ID NO: 95 to 111 mg/L, which does not meet the design goal of an insulin-Fc fusion protein homodimer titer greater than 150 mg/L. The following shows SEQ ID NO: 86 with the B16A and cNg-NB155-Q mutations underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
GGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
PEVKFNWYVDGVEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 86) (not  
claimed).

**[0111]** These results indicate that there is an unexpected and significant interaction between the choice of cNg mutation on the human IgG1 Fc fragment and the composition of the insulin polypeptide such that experimentation is required to identify the preferred configurations with respect to IR and FcRn binding affinities of the resulting insulin-Fc fusion proteins.

**[0112]** In some configurations, insulin-Fc fusion protein configurations described herein do not include a leader amino acid sequence at the N-terminus. In other configurations, insulin-Fc fusion protein configurations described herein includes a leader sequence, e.g., at the N-terminus. In some embodiments, an exemplary leader sequence includes the amino acid sequence: MEWSWVFLFFLSVTTGVHS (SEQ ID NO: 24). In some embodiments, insulin-Fc fusion protein configurations described herein are encoded by a nucleic acid molecule comprising a leader sequence, e.g., for expression (e.g., recombinant expression) in cells (e.g., eukaryotic, e.g., mammalian cells). In certain embodiments, the leader sequence is cleaved off, e.g., in the cell culture, during expression. An exemplary nucleic acid sequence encoding a leader sequence includes the nucleic acid sequence:

atggaatggagctgggtctttcttcttctcgtcagtaacgactgggtccactccttcgtgaaccagcacctgtgcggctcccacctgggtggaagctctggc  
actcgtgtgcggcgagcggggctccactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtcgcacctccacctgct  
ccctggaccagctggaaaactactgcggtggcggaggtggtcaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa  
aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttccccccaaaaccaaaggacacctcatgatctccggg  
accctgaggtcacatgcgtggtggtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag  
acaaagccgcgggaggagcagtacagcagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtg  
caaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccat  
cccggtatgagctgaccaagaaccaggtcagcctgacctgcctggtaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcag  
ccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 79).

**[0113]** In a preferred embodiment comprising the insulin-Fc fusion protein configuration of SEQ ID NO: 78, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtctttcttcttctcgtcagtaacgactgggtccactccttcgtgaaccagcacctgtgcggctcccacctgggtggaagctctggc  
actcgtgtgcggcgagcggggctccactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtcgcacctccacctgct  
ccctggaccagctggaaaactactgcggtggcggaggtggtcaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa  
aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttccccccaaaaccaaaggacacctcatgatctccggg  
accctgaggtcacatgcgtggtggtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag  
acaaagccgcgggaggagcagtacagcagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtg  
caaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccat  
cccggtatgagctgaccaagaaccaggtcagcctgacctgcctggtaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcag  
ccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 79).

**[0114]** In a preferred embodiment comprising the insulin-Fc fusion protein configuration of SEQ ID NO: 80, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtctttcttcttctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgcggctccacctgggtggaagctctggc  
actcgtgtgcggcgagcggggcttcactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtgcacctccacctgct  
 5 ccctggaccagctggaaaactactgcggtggcggaggtggtaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa  
aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacacccctcatgatctccgg  
accctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag  
 10 acaaagccgcgggaggagcagtagcacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt  
caaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccat  
ccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatccagcgacatcgccgtggagtgaggagagcaatgggcag  
 15 ccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 81)  
 (not claimed).

[0115] In an embodiment comprising the insulin-Fc fusion protein configuration of SEQ ID NO: 82, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtctttcttcttctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgcggctccacctgggtggaagctctggc  
actcgtgtgcggcgagcggggcttcactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtgcacctccacctgct  
 25 ccctggaccagctggaaaactactgcggtggcggaggtggtaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa  
aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacacccctcatgatctccgg  
 30 accctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag  
acaaagccgcgggaggagcagtagccagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt  
caaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccat  
 35 ccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatccagcgacatcgccgtggagtgaggagagcaatgggcag  
ccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 83)  
 40 (not claimed).

[0116] In a preferred embodiment comprising the insulin-Fc fusion protein configuration of SEQ ID NO: 84, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtcttctcttctcgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgcggctcccacctgggtggaagctctggc  
 actcgtgtgcggcgagcggggcttccactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtgcacctccacctgct  
 ccctggaccagctggaaaactactgcggtggcggagggtgtcaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa  
 aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaaccaaggacacccctcatgatctccggg  
 acccctgaggtcacatgcgtgggtggacgtgagccacgaagaccctgaggtcaagtcaactgggtacgtggacggcgtggaggtgcataatgccaa  
 acaaagccgaggaggagcagtagaagcacgtaccgtgtgtgtagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt  
 caaggtctccaacaaagccctcccagccccatcgagaaaaccatctcaaagccaaaggcagccccgagaaccacaggtgtacacctgccccat  
 cccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcag  
 ccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
 gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 85).

**[0117]** In a preferred embodiment comprising the insulin-Fc fusion protein configuration of SEQ ID NO: 86, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtcttcttcttctcgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgcggctcccacctgggtggaagctctggc  
 actcgtgtgcggcgagcggggcttccactacgggggtggcggaggaggttctgggtggcggcgaggcatcgtggaacagtgtgcacctccacctgct  
 ccctggaccagctggaaaactactgcggtggcggagggtgtcaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaa  
 aactcacacatgccaccgtgccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaaccaaggacacccctcatgatctccggg  
 acccctgaggtcacatgcgtgggtggacgtgagccacgaagaccctgaggtcaagtcaactggtacgtggacggcgtggaggtgcataatgccaa  
 acaaagccgaggaggagcagtagcaaaagcacgtaccgtgtgtgtagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagt  
 caaggtctccaacaaagccctcccagccccatcgagaaaaccatctcaaagccaaaggcagccccgagaaccacaggtgtacacctgccccat  
 cccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcag  
 ccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcag  
 gggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO:  
 100) (not claimed).

**[0118]** In some applications, it may be desirable to covalently modify the insulin-Fc fusion protein to further modify its properties. For example, the insulin-Fc fusion protein may be conjugated to various lengths of polyethylene glycol (PEG) molecules to further extend its circulating half-life as described in Dozier, J.; Distefano, M. Site-Specific PEGylation of Therapeutic Proteins. Int. J. Mol. Sci. 2015, 16(10), 25831-25864). One such method of conjugation relies on a class of enzymes called transglutaminases (TGases) which create a covalent bond between a primary amine (e.g. at the terminus of a PEG molecule) and the carboxamide group of glutamine (e.g. Q on a target protein, e.g. on a target insulin-Fc fusion protein) via an acyl transfer reaction (Dozier, J.; Distefano, M., page 25853). The Q at position NB153 of the insulin-Fc fusion proteins of SEQ ID NO: 86, and SEQ ID NO: 95 is a preferred TGase conjugation site, but because each of the insulin-Fc fusion proteins comprises a GGGGGQGGGGQGGGGQGGGGG (SEQ ID NO: 13) linker, the linker will also be conjugated at the Q residues. Therefore, to minimize unwanted TGase conjugation at the linker Q sites, alanine (A) for glutamine (Q) mutations in the linker were tested. Accordingly, in a preferred embodiment, the insulin-Fc fusion protein of SEQ ID NO: 87 comprises the insulin polypeptide of SEQ ID NO: 10, the linker of GGGGGAGGGGAGGGGAGGGG (SEQ ID NO: 67), and the cNg-NB 155-S mutant of the Fc fragment of SEQ ID NO: 77 (Xi is S). The insulin-Fc fusion protein of SEQ ID NO: 87 exhibits a homodimer titer greater than 150 mg/L, an IR assay IC50 value less than 2400 nM, a human FcRn assay EC50 value less than 1000 ng/mL, a human Fc(gamma)RI assay OD450 Ratio at biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL less than 0.50, and a human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1,000 ng/mL less than 0.35. The following shows SEQ ID NO: 87 with the B16A and cNg-NB155-S mutations as well as the linker sequence underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGA  
GGGGAGGGGAGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHED  
 5 PEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
 KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD  
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 87).

10 **[0119]** In an embodiment comprising the insulin-Fc fusion protein of SEQ ID NO: 87, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtcttctcttctctctgtcagtaacgactgggtccactcctcgtgaaccagcacctgtgcggctcccactgggtggaagctctggc  
 15 actcgtgtgcggcgagcggggtccactacgggggtggcggaggaggttctggtggcgggaggagcatcgtggaacagtgtgcacctccacctgct  
ccctggaccagctggaaaactactgcggtggcgagggtgtgcaggaggcggtggagctgggggtggaggaggcgggggagacaa  
 20 aactcacacatgcccacgtgcccagcacctgaactcctggggggaccgtcagttcttcttcccccaaaacccaaggacacctcatgatctccgg  
acccctgagggtcacatgcgtgggtgggacgtgagccacgaagacctgagggtcaagttcaactggtacgtggacggcgtggagggtgcataatgccaag  
acaaagccgcgaggagcagtagcagcagcagcgtaccgtgtggtcagcgtcctcaccgtcctgaccaggactggctgaatggcaaggagtacaagt  
 25 caaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccccat  
ccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttctatccagcgacatcgccgtggagtgggagagcaatgggcag  
ccggagaacaactacaagaccacgcctcccgctgctggactccgacggctccttctctctacagcaagctcaccgtggacaagagcaggtggcagcag  
 30 gggaactcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccgggttag (SEQ ID NO: 88).

35 **[0120]** Linker length in Fc-fusion protein conjugates has been known to affect characteristics of the fusion protein ranging from yield to bioactivity. Accordingly, an insulin-Fc fusion protein configuration with a shorter linker was tested to determine any gain or loss of yield or function due to the length of the linker. In another preferred embodiment, the insulin-Fc fusion protein of SEQ ID NO: 89 (not claimed) comprises the insulin polypeptide of SEQ ID NO: 10, the linker of GGGGAGGGG (SEQ ID NO: 11), and the cNg-NB155-S mutant of the Fc fragment of SEQ ID NO: 77 (Xi is S). The insulin-Fc fusion protein of SEQ ID NO: 89 exhibits a homodimer titer greater than 150 mg/L, an IR assay IC<sub>50</sub> value less than 2400 nM, a human FcRn assay EC<sub>50</sub> value less than 1000 ng/mL, a human Fc(gamma)RI assay OD<sub>450</sub> Ratio at a biotinylated-Fc(gamma)RI concentration of 3,000 ng/mL less than 0.50, and a human C1q binding assay OD<sub>450</sub> Ratio at a biotinylated-C1q concentration of 1,000 ng/mL less than 0.35.

40 **[0121]** The following shows SEQ ID NO: 89 with the B16A and cNg-NB155-S mutations as well as the linker sequence underlined:

FVNQHLCGSHLVEALALVCGERGFHYGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGAG  
 45 GGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV  
 EVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ  
 VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD  
 50 KSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 89) (not claimed).

**[0122]** In an embodiment comprising the insulin-Fc fusion protein of SEQ ID NO: 89, the nucleic acid sequence (leader sequence underlined) is:

atggaatggagctgggtctttcttcttctcgtcagtaacgactgggtgccactcctcgtgaaccagcacctgtgcggctcccacgtggtggaagctctggc  
 actcgtgtgcggcgagcggggcttccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaacagtgtgcacctccacctgct  
 5 ccttgaccagctggaaaactactgcggtggcggagggtgccggaggcgggggagacaaaactcacacatgccaccgtgccagcacctgaactcct  
 ggggggaccgtcagttcttcttcccccaaaacccaaggacaccctcatgatctccggaccctgaggtcacatgcgtggtggtggacgtgagccac  
 gaagaccctgaggtcaagttcaactggtacgtggacggcgtggagggtcataatgccaagacaaagccggggaggagcagtacagcagcacgtacc  
 10 gtgtggtcagcgtcctcaccgtctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaacaagccctcccagccccatcgaga  
 aaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacaccctgcccccatccgggatgagctgaccaagaaccaggtcagcctgac  
 ctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagcaatgggcagccggagaaactacaagaccacgcctccgtgctgg  
 15 actccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgttctctcatgctccgtgatgcatgaggctctg  
 cacaaccactacagcagaagagcctctccctgtctccgggttag (SEQ ID NO: 90) (not claimed).

[0123] In general preferred configurations as claimed, the less-immunogenic, non-glycosylated, cNg-mutated human  
 20 IgG1 Fc fragments of SEQ ID NO: 77 (Xi is S, D (not claimed), A(not claimed), or R) combined with the preferred B16A-  
 mutated insulin polypeptide of SEQ ID NO: 10 and with various linkers (SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO:  
 67) produce a family of high homodimer titer-yielding, bioactive, non-immunogenic human insulin-Fc fusion protein  
 configurations that are expected to exhibit sufficient in vivo potency and duration of action to enable chronic, once-a-  
 week dosing in diabetic patients.

#### ***Insulin-Fc fusion Protein Production***

[0124] In embodiments, a fusion protein can be expressed by a cell as described in more detail in the Examples section.

#### ***Expression and Purification***

[0125] In embodiments, an insulin-Fc fusion protein can be expressed recombinantly, e.g., in a eukaryotic cell, e.g.,  
 mammalian cell or non-mammalian cell. Exemplary mammalian cells used for expression include HEK cells (e.g., HEK293  
 cells) or CHO cells. CHO cells can be subdivided into various strains or subclasses, (e.g. CHO DG44, CHO-M, and  
 35 CHO-K1), and some of these cell strains may be genetically engineered for optimal use with a particular type of nucleic  
 acid molecule (e.g., a vector comprising DNA) or a particular cell growth media composition as described in the Examples  
 section. In embodiments, cells are transfected with a nucleic acid molecule (e.g., vector) encoding the insulin-Fc fusion  
 protein (e.g., where the entire insulin-Fc fusion protein is encoded by a single nucleic acid molecule). In embodiments,  
 40 HEK293 cells are transfected with a vector that encodes for the insulin-Fc fusion protein, but only results in temporary  
 expression of the insulin-Fc fusion protein for a period of time (e.g., 3 days, 4 days, 5, days, 7 days, 10 days, 12 days,  
 14 days, or more) before the host cell stops expressing appreciable levels of the insulin-Fc fusion protein (i.e., transient  
 transfection). HEK293 cells that are transiently transfected with nucleic acid sequences encoding for insulin-Fc fusion  
 proteins often allow for more rapid production of recombinant proteins which facilitates making and screening multiple  
 45 insulin-Fc fusion protein candidates. In embodiments, CHO cells are transfected with a vector that is permanently  
 incorporated into the host cell DNA and leads to consistent and permanent expression (i.e., stable transfection) of the  
 insulin-Fc fusion protein as long as the cells are cultured appropriately. CHO cells and CHO cell lines that are stably  
 transfected with nucleic acids encoding for insulin-Fc fusion proteins often take longer to develop, but they often produce  
 higher total protein yields and are more amenable to manufacturing low cost products (e.g., products for the relatively  
 commoditized human insulin market). Cells and cell lines can be cultured using standard methods in the art.

[0126] In preferred embodiments, HEK cells comprising any one of the cDNA sequences with SEQ ID NO: 79 (corre-  
 sponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 78), SEQ ID NO: 81 (corresponding to the insulin-  
 Fc fusion protein embodiment of SEQ ID NO: 80) (not claimed), SEQ ID NO: 83 (corresponding to the insulin-Fc fusion  
 protein embodiment of SEQ ID NO: 82) (not claimed), SEQ ID NO: 85 (corresponding to the insulin-Fc fusion protein  
 embodiment of SEQ ID NO: 84), SEQ ID NO: 88 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID  
 55 NO: 87), and SEQ ID NO: 90 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 89) (not claimed)  
 are used to express insulin-Fc fusion proteins. In preferred embodiments, CHO cells comprising any one of the cDNA  
 sequences with SEQ ID NO: 79 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 78), SEQ  
 ID NO: 81 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 80) (not claimed), SEQ ID NO:

83 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 82) (not claimed), SEQ ID NO: 85 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 84), SEQ ID NO: 88 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 87), and SEQ ID NO: 90 (corresponding to the insulin-Fc fusion protein embodiment of SEQ ID NO: 89) (not claimed) are used to express insulin-Fc fusion proteins.

**[0127]** In some embodiments, the insulin-Fc fusion protein is purified or isolated from the cells (e.g., by lysis of the cells). In other embodiments, the insulin-Fc fusion protein is secreted by the cells and purified or isolated from the cell culture media in which the cells were grown. Purification of the insulin-Fc fusion protein can include using column chromatography (e.g., affinity chromatography) or using other separation methods based on differences in size, charge, and/or affinity for certain molecules. In embodiments, purification of the insulin-Fc fusion protein involves selecting or enriching for proteins containing an Fc fragment, e.g., by using Protein A beads or a Protein A column that cause proteins containing an Fc fragment to become bound with high affinity at neutral solution pH to the Protein A covalently conjugated to the Protein A beads. The bound insulin-Fc fusion protein may then be eluted from the Protein A beads by a change in a solution variable (e.g. a decrease in the solution pH). Other separation methods such as ion exchange chromatography and/or gel filtration chromatography can also be employed alternatively or additionally. In embodiments, purification of the insulin-Fc fusion protein further comprises filtering or centrifuging the protein preparation. In embodiments, further purification of the insulin-Fc fusion protein comprises diafiltration, ultrafiltration, and filtration through porous membranes of various sizes, as well as final formulation with excipients.

**[0128]** The purified insulin-Fc fusion protein can be characterized, e.g., for purity, total protein yield, structure, and/or activity, using a variety of methods, e.g., absorbance at 280 nm (e.g., to determine total protein yield), size exclusion or capillary electrophoresis (e.g., to determine the molecular weight, percent aggregation, and/or purity), mass spectrometry (MS) and/or liquid chromatography (LC-MS) (e.g., to determine purity and/or glycosylation), and/or ELISA (e.g., to determine extent of binding, e.g., affinity, to an anti-insulin antibody). Exemplary methods of characterization are also described in the Examples section.

**[0129]** In embodiments, the total protein yield of an insulin-Fc fusion protein after production in transiently transfected HEK cells and protein A purification is greater than 5 mg/L, 10 mg/L, or 20 mg/L. In preferred embodiments, the total protein yield of a human insulin-Fc fusion protein after production in transiently transfected HEK cells and protein A purification is greater than 100 mg/L (e.g., greater than 150 mg/L). In embodiments, the %homodimer of an insulin-Fc fusion protein after production in transiently transfected HEK cells and protein A purification is greater than 70% (e.g., greater than 80%, greater than 85%, greater than 90%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, greater than 99%). In preferred embodiments, the homodimer titer of a human insulin-Fc fusion protein after production in transiently transfected HEK cells and protein A purification, calculated as the product between the insulin-Fc fusion total protein yield and the %homodimer is greater than 100 mg/L (e.g., greater than 150 mg/L). Only human insulin-Fc fusion protein configurations with a homodimer titer of greater than 150 mg/L were considered useful in the present invention, because experience has demonstrated that homodimer titers less than this level will not likely result in commercial production titers in CHO cells that meet the stringently low manufacturing cost requirements for the relatively commoditized human insulin market.

**[0130]** In embodiments, the total protein yield of an insulin-Fc fusion protein after production in stably transfected CHO cells (e.g., CHO cell lines or CHO cell clones) and protein A purification is greater than 100 mg of insulin-Fc fusion protein per L (e.g. mg/L of culture media). In preferred embodiments, the total protein yield of an insulin-Fc fusion protein after production in stably transfected CHO cells (e.g. CHO cell lines or CHO cell clones) and protein A purification is greater than 150 mg insulin-Fc fusion protein/L of culture media (e.g., greater than 200 mg/L, greater than 300 mg/L, greater than 400 mg/L, greater than 500 mg/L, greater than 600 mg/L or more). In embodiments, the %homodimer of an insulin-Fc fusion protein after production in stably transfected CHO cells (e.g. CHO cell lines or CHO cell clones) and protein A purification is greater than 70% (e.g., greater than 80%, greater than 85%, greater than 90%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, greater than 99%). In embodiments, the homodimer titer of an insulin-Fc fusion protein after production in stably transfected CHO cells (e.g. CHO cell lines or CHO cell clones) and protein A purification, calculated as the product between the insulin-Fc fusion total protein yield and the %homodimer is greater than 150 mg/L (e.g., greater than 200 mg/L, greater than 300 mg/L, greater than 400 mg/L, greater than 500 mg/L, greater than 600 mg/L or more).

### ***Functional Features of Insulin-Fc Fusion Proteins***

**[0131]** Described herein are methods and uses for interacting with the insulin receptors to lower blood glucose in the target subject (e.g., dogs or humans), wherein the methods and uses comprise administering to the subject an insulin-Fc fusion protein, e.g., a fusion protein described herein. In some embodiments, the subject has been diagnosed with diabetes (e.g., canine diabetes for dogs, or type 1 diabetes or type 2 diabetes for humans).

**[0132]** In embodiments, an insulin-Fc fusion protein described herein binds to the IR with an appreciable affinity as measured by the IC<sub>50</sub> in the 4°C IM-9 IR binding assay described in Example 12 (e.g. IC<sub>50</sub> less than 5000 nM, IC<sub>50</sub>

less than 4000 nM, IC<sub>50</sub> less than 3000 nM, IC less than 2400 nM, IC<sub>50</sub> less than 2000 nM). Based on experience, only compounds exhibiting IR activity IC<sub>50</sub> values less than 5000 nM, preferably less than 2400 nM and more preferably less than 2000 nM, were deemed likely to exhibit bioactivity in the target subject. Generally, higher affinity IR binding (i.e., lower IC<sub>50</sub> values) is preferred. However, it is well-known that the clearance of insulin and insulin analogs (e.g., insulin polypeptides described herein) is governed primarily through binding to the IR followed by IR internalization and degradation within the cell. Therefore, insulin-Fc fusion protein configurations with too high of an IR binding affinity (i.e., too low of an IC<sub>50</sub>, e.g. an IC<sub>50</sub> less than 500 nM) may be cleared too quickly from circulation resulting in a lower than desired duration of glucose-lowering bioactivity in the target subject.

**[0133]** In embodiments, the insulin-Fc fusion protein configurations described herein bind to the FcRn receptor with an affinity that is higher than that of an insulin-Fc fusion protein reference standard as measured according to Example 19. In some embodiments, the FcRn receptor affinity of the insulin-Fc fusion protein configurations as described herein and as measured by the human FcRn receptor assay EC<sub>50</sub> value is less than or equal to 1500 ng/mL, and more preferably less than or equal to 1000 ng/mL.

**[0134]** In embodiments, the insulin-Fc fusion proteins described herein are capable of lowering glucose levels (e.g., blood glucose levels) after administration in a subject. In embodiments, the glucose lowering activity of the insulin-Fc fusion proteins is greater than that of an insulin reference standard. In some embodiments, the duration of activity of the insulin-Fc fusion proteins can be measured by a decrease, e.g., a statistically significant decrease, in fasting blood glucose relative to a pre-dose fasting blood glucose level. In embodiments, the duration of activity of the insulin-Fc fusion proteins (e.g., the time during which there is a statistically significant decrease in fasting blood glucose level in a subject relative to a pre-dose level) is longer than about 2 hours. In embodiments, the duration of activity of the insulin-Fc fusion proteins (e.g., the time during which there is a statistically significant decrease in blood glucose level in a subject relative to a pre-dose level) is longer than about 6 hours, 9 hours, 12 hours, 18 hours, 1 day, 1.5 days, 2 days, 2.5 days, 3 days, 4 days, 5 days, 6 days, 7 days, or longer. In embodiments, the insulin-Fc fusion proteins are long-acting (e.g., have a long half-life, e.g., in serum).

**[0135]** In embodiments, the serum half-life of the insulin-Fc fusion proteins described herein in the target subject is longer than that of an insulin reference standard or control formulation. In embodiments, the serum half-life of the insulin-Fc fusion proteins (e.g., in the blood of a subject upon administration) in the target subject is longer than about 2 hours. In embodiments, the serum half-life of the insulin-Fc fusion protein in the target subject is about 0.5 days, 1 day, 2 days, or 2.5 days. In preferred embodiments, the serum half-life of the insulin-Fc fusion protein in the target subject is about 3 days or longer.

**[0136]** In embodiments, the combination of potency and duration of bioactivity of the insulin-Fc fusion proteins described herein may be quantified by calculating the area over the percent fasting blood glucose (%FBGL) curve normalized to a given dose in mg/kg (NAOC) with units of %FBGL·days·kg/mg. In embodiments, the NAOC of the insulin-Fc fusion proteins described herein is greater than 150 %FBGL·days·kg/mg (e.g. greater than 200 %FBGL·days·kg/mg, greater than 250 %FBGL·days·kg/mg or more). Again, based on experience, at NAOC values greater than 150 %FBGL·days·kg/mg, the dose requirements in the target subject will be sufficiently low so as to achieve an acceptable treatment cost. In embodiments, the NAOC of the insulin-Fc fusion proteins must be maintained after repeated dosing in the target subject (i.e., the ratio of the NAOC after the third dose to the NAOC after the first dose of the insulin-Fc fusion protein is greater than 0.5 (e.g., greater than 0.6, greater than 0.7, greater than 0.8, greater than 0.9, or more).

**[0137]** In some embodiments, the insulin-Fc fusion protein configurations described herein bind to the Fc(γ) receptor with an affinity that is lower than that of an insulin-Fc fusion protein reference standard as measured according to Example 15. In some embodiments, the ratio of the Fc(γ) receptor affinity of the insulin-Fc fusion proteins to that of an insulin-Fc fusion protein reference standard (i.e., the insulin-Fc fusion protein of SEQ ID NO: 76) is less than 0.50 (e.g. less than 0.40, less than 0.30, less than 0.20). In some embodiments, the Fc(γ) receptor affinity of the insulin-Fc fusion protein as measured by the human Fc(γ)RI receptor assay OD450 Ratio at 3000 ng/mL biotinylated-Fc(γ)RI (with respect to the reference insulin-Fc fusion protein of SEQ ID NO: 76) is less than or equal to 0.50.

**[0138]** In some embodiments, the insulin-Fc fusion protein configurations described herein bind to C1q with an affinity that is lower than that of an insulin-Fc fusion protein reference standard as measured according to Example 16. In some embodiments, the ratio of the C1q binding affinity of the insulin-Fc fusion proteins to that of an insulin-Fc fusion protein reference standard (i.e., the insulin-Fc fusion protein of SEQ ID NO: 76) is less than 0.50 (e.g. less than 0.40, less than 0.30, less than 0.20). In some embodiments, the C1q binding affinity of the insulin-Fc fusion proteins as measured by the human C1q binding assay OD450 Ratio at a biotinylated-C1q concentration of 1000 ng/mL (with respect to the reference insulin-Fc fusion protein of SEQ ID NO: 76) is less than or equal to 0.35.

#### ***Methods of Treatment (not claimed) and Characteristics of Subject Selection***

**[0139]** Described herein are methods for treating diabetes (e.g., type 1 diabetes or type 2 diabetes in humans), the

methods comprising the administration of an insulin-Fc fusion protein (e.g., an insulin-Fc fusion protein configuration as described herein) to a target subject (not claimed). Also described herein are fusion proteins as described herein or a pharmaceutical composition thereof for use as a medicament in lowering the blood glucose level of a patient.

**[0140]** In embodiments, a reference standard used in any method or use described herein comprises a reference treatment or reference therapy. In some embodiments, the reference comprises a standard of care agent for diabetes treatment (e.g., a standard of care agent for canine diabetes or a standard of care agent for type 1 diabetes in humans or a standard of care agent for type 2 diabetes in humans). In some embodiments, the reference standard is a commercially available insulin or insulin analog. In some embodiments, the reference standard comprises a long-lasting insulin, intermediate-lasting insulin, shortlasting insulin, rapid-acting insulin, short-acting, intermediate-acting, long-acting insulin. In some embodiments, the reference standard for canine insulin comprises Vetsulin®, Prozinc®, insulin NPH, insulin glargine (Lantus®) or recombinant human insulin. In some embodiments, the reference standard for human insulin comprises Humalog®, NovoLog®, Novolin®R (Novo Nordisk, Bagsværd, Denmark), Novolin®N (Novo Nordisk, Bagsværd, Denmark), Humulin®R (Eli Lilly, Indianapolis, IN), Humulin®N (Eli Lilly, Indianapolis, IN), Lantus®, and Levemir®, or generic recombinant human insulin.

**[0141]** In embodiments, a reference standard used in any method described herein includes an outcome, e.g., outcome described herein, of a diabetes therapy (e.g., a canine diabetes therapy or a human diabetes therapy).

**[0142]** In embodiments, a reference standard is a level of a marker (e.g., blood glucose or HbA1c) in the target subject prior to initiation of a therapy, e.g., an insulin-Fc fusion protein therapy described herein; where the target subject has diabetes. In embodiments, the blood glucose level in a target subject is greater than 200 mg/dL (e.g. greater than 250 mg/dL, 300 mg/dL, 350 mg/dL, 400 mg/dL or more) prior to initiation of therapy. In embodiments, the fructosamine level in a dog is greater than 250 micromol/L, 350 micromol/L (e.g. greater than 400 micromol/L, 450 micromol/L, 500 micromol/L, 550 micromol/L, 600 micromol/L, 650 micromol/L, 700 micromol/L, 750 micromol/L or more) prior to initiation of therapy. In embodiments, the HbA1c level in a human target subject is greater than 7 mmol/L (e.g. greater than 7.5 mmol/L, 8 mmol/L, 9 mmol/L, 10 mmol/L, 11 mmol/L, 12 mmol/L, or more) prior to initiation of therapy. In embodiments, a reference standard is a measure of the presence of, the progression of, or the severity of the disease. In embodiments, a reference standard is a measure of the presence of or the severity of the disease symptoms prior to initiation of a therapy, e.g., an insulin-Fc fusion protein therapy described herein, e.g., where the target subject has diabetes.

### ***Pharmaceutical Compositions and Routes of Administration***

**[0143]** Provided herein are pharmaceutical compositions containing an insulin-Fc fusion protein configuration as described herein that can be used to lower blood glucose in target subjects. The amount and concentration of the insulin-Fc fusion protein in the pharmaceutical compositions, as well as the quantity of the pharmaceutical composition administered to a target subject, can be selected based on clinically relevant factors, such as medically relevant characteristics of the subject (e.g., age, weight, gender, other medical conditions, and the like), the solubility of compounds in the pharmaceutical compositions, the potency and activity of the compounds, and the manner of administration of the pharmaceutical compositions. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

**[0144]** Formulations of the present disclosure include those suitable for parenteral administration. The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by intravenous or subcutaneous injection.

**[0145]** Examples of suitable aqueous and non-aqueous carriers that may be employed in the pharmaceutical compositions of the disclosure include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants, e.g., Tween-like surfactants. In some embodiments, the pharmaceutical composition (e.g., as described herein) comprises a Tween-like surfactant, e.g., polysorbate-20, Tween-20 or Tween-80. In some embodiments, the pharmaceutical composition (e.g., as described herein) comprises a Tween-like surfactant, e.g., Tween-80, at a concentration between about 0.001% and about 2%, or between about 0.005% and about 0.1%, or between about 0.01% and about 0.5%.

**[0146]** In some embodiments, the concentration of the insulin-Fc fusion protein in the aqueous carrier is about 3 mg/mL. In some embodiments, the concentration of the insulin-Fc fusion protein in the aqueous carrier is about 6 mg/mL. In some embodiments, the concentration of the insulin-Fc fusion protein in the aqueous carrier is about 8 mg/mL, 9 mg/mL, 10 mg/mL, 12 mg/mL, 15 mg/mL or more.

**[0147]** In some embodiments, the insulin-Fc fusion protein is administered as a bolus, infusion, or an intravenous push. In some embodiments, the fusion protein is administered through syringe injection, pump, pen, needle, or indwelling catheter. In some embodiments, the insulin-Fc fusion protein is administered by a subcutaneous bolus injection. Methods

of introduction may also be provided by rechargeable or biodegradable devices. Various slow release polymeric devices have been developed and tested in vivo in recent years for the controlled delivery of drugs, including proteinaceous biopharmaceuticals. A variety of biocompatible polymers (including hydrogels), including both biodegradable and non-degradable polymers, can be used to form an implant for the sustained release of a compound at a particular target site.

## Dosages

[0148] Actual dosage levels of an insulin-Fc fusion protein of the configurations described herein can be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular target subject (e.g., dog or human). The selected dosage level will depend upon a variety of factors including the activity of the particular fusion protein employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular fusion protein employed, the age, sex, weight, condition, general health and prior medical history of the subject being treated, and like factors well known in the medical arts.

[0149] In general, a suitable dose of an insulin-Fc fusion protein will be the amount that is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Generally, intravenous and subcutaneous doses of the insulin-Fc fusion protein for a target subject will range from about 0.001 to about 1 mg per kilogram (e.g. mg/kg) of body weight per day, e.g., about 0.001 to 1 mg/kg/day, about 0.01 to 0.1 mg/kg/day, about 0.1 to 1 mg/kg/day, or about 0.01 to 1 mg/kg/day. In still other embodiments, the fusion protein is administered at a dose between 0.025 and 4 mg per kilogram of body weight per week, e.g., between 0.025 and 1.0 mg/kg/week.

[0150] The present disclosure contemplates formulation of the insulin-Fc fusion protein in any of the aforementioned pharmaceutical compositions and preparations. Furthermore, the present disclosure contemplates administration via any of the foregoing routes of administration. One of skill in the art can select the appropriate formulation and route of administration based on the condition being treated and the overall health, age, and size of the subject being treated.

## EXAMPLES

[0151] The present technology is further illustrated by the following Examples, which should not be construed as limiting in any way.

## GENERAL METHODS, ASSAYS, AND MATERIALS

### Example 1: Synthesis and Methods of Making an Insulin-Fc Fusion Protein in HEK293 Cells.

[0152] Insulin-Fc fusion proteins were synthesized as follows. A gene sequence of interest was constructed using proprietary software (LakePharma, Belmont, CA) and was cloned into a high expression mammalian vector. HEK293 cells were seeded in a shake flask 24 hours before transfection and were grown using serum-free chemically defined media. A DNA expression construct that encodes the insulin-Fc fusion protein of interest was transiently transfected into a suspension of HEK293 cells using the (LakePharma, Belmont, CA) standard operating procedure for transient transfection. After 20 hours, the cells were counted to determine the viability and viable cell count, and the titer was measured by FortéBio® Octet® (Pall FortéBio LLC, Fremont, CA). Additional readings were taken throughout the transient transfection production run. The culture was harvested on or after day 5.

### Example 2: Synthesis and Methods of Making an Insulin-Fc Fusion Protein in CHO Cells.

[0153] A CHO cell line was originally derived from CHO-K1 (LakePharma, Belmont, CA), and the endogenous glutamine synthetase (GS) genes were knocked out by recombinant technology using methods known in the art. Stable expression DNA vectors were designed and optimized for CHO expression and GS selection and incorporated into a high expression mammalian vector (LakePharma, Belmont, CA). The sequence of each completed construct was confirmed prior to initiating scale up experiments. The suspension-adapted CHO cells were cultured in a humidified 5% CO<sub>2</sub> incubator at 37°C in a chemically defined media (CD OptiCHO; Invitrogen, Carlsbad, CA). No serum or other animal-derived products were used in culturing the CHO cells.

[0154] Approximately 80 million suspension-adapted CHO cells, growing in CD OptiCHO media during the exponential growth phase, were transfected by electroporation using MaxCyte® STX® system (MaxCyte, Inc., Gaithersburg, MD) with 80 µg DNA to create a stable CHO cell line for each insulin-Fc fusion protein (DNA construct contains the full-length sequence of the insulin-Fc fusion protein). After twenty-four hours, the transfected cells were counted and placed

under selection for stable integration of the insulin-Fc fusion genes. The transfected cells were seeded into CD OptiCHO selection media containing between 0-100  $\mu$ M methionine sulfoximine (MSX) at a cell density of  $0.5 \times 10^6$  cells/mL in a shaker flask and were incubated at 37°C with 5% CO<sub>2</sub>. During a selection process, the cells were spun down and resuspended in fresh selection media every 2-3 days until the CHO stable pool recovered its growth rate and viability.

The cell culture was monitored for growth and titer.

**[0155]** The cells were grown to  $2.5 \times 10^6$  cells per mL. At the time of harvest for cell banking, the viability was above 95%. The cells were then centrifuged, and the cell pellet was resuspended in the CD OptiCHO media with 7.5% dimethyl sulfoxide (DMSO) to a cell count of  $15 \times 10^6$  cells per mL per vial. Vials were cryopreserved for storage in liquid nitrogen.

**[0156]** A small-scale-up production was performed using the CHO cells as follows. The cells were scaled up for production in CD OptiCHO growth medium containing 100  $\mu$ M MSX at 37°C and fed every 2-4 days as needed, with CD OptiCHO growth medium supplemented with glucose and additional amino acids as necessary for approximately 14-21 days. The conditioned media supernatant harvested from the stable pool production run was clarified by centrifuge spinning. The protein was run over a Protein A (MabSelect, GE Healthcare, Little Chalfont, United Kingdom) column pre-equilibrated with binding buffer. Washing buffer was then passed through the column until the OD280 value (NanoDrop, Thermo Scientific) was measured to be at or near background levels. The insulin-Fc fusion protein was eluted using a low pH buffer, elution fractions were collected, and the OD280 value of each fraction was recorded. Fractions containing the target insulin-Fc fusion protein were pooled and optionally further filtered using a 0.2  $\mu$ M membrane filter.

**[0157]** The cell line was optionally further subcloned to monoclonality and optionally further selected for high titer insulin-Fc-fusion protein-expressing clones using the method of limiting dilution, a method known to those skilled in the art. After obtaining a high titer, monoclonal insulin-Fc fusion protein-expressing cell line, production of the insulin-Fc fusion protein was accomplished as described above in growth medium without MSX, or optionally in growth medium containing MSX, to obtain a cell culture supernatant containing the recombinant, CHO-made, insulin-Fc fusion protein. The MSX concentration was optionally increased over time to exert additional selectivity for clones capable of yielding higher product titers.

### Example 3: Synthesis and Methods of Making an Insulin-Fc Fusion Protein in CHO Cells.

**[0158]** A CHO cell line is originally derived from CHO-K1 (LakePharma, Belmont, CA), and the endogenous glutamine synthetase (GS) genes are knocked out by recombinant technology using methods known in the art. Stable expression DNA vectors are designed and optimized for CHO expression and GS selection and incorporated into a high expression mammalian vector (LakePharma, Belmont, CA). The sequence of each completed construct is confirmed prior to initiating scale up experiments. The suspension-adapted CHO cells are cultured in a humidified 5% CO<sub>2</sub> incubator at 37°C in a chemically defined media (CD OptiCHO; Invitrogen, Carlsbad, CA). No serum or other animal-derived products are used in culturing the CHO cells.

**[0159]** Approximately 80 million suspension-adapted CHO cells, growing in CD OptiCHO media during the exponential growth phase, are transfected by electroporation using MaxCyte® STX® system (MaxCyte, Inc., Gaithersburg, MD) with 80  $\mu$ g DNA to create a stable CHO cell line for each insulin-Fc fusion protein (DNA construct contains the full-length sequence of the insulin-Fc fusion protein). After twenty-four hours, the transfected cells are counted and placed under selection for stable integration of the insulin-Fc fusion genes. The transfected cells are seeded into CD OptiCHO selection media containing between 0-100  $\mu$ M methionine sulfoximine (MSX) at a cell density of  $0.5 \times 10^6$  cells/mL in a shaker flask and are incubated at 37°C with 5% CO<sub>2</sub>. During a selection process, the cells are spun down and resuspended in fresh selection media every 2-3 days until the CHO stable pool recovered its growth rate and viability. The cell culture is monitored for growth and titer.

**[0160]** The cells are grown to  $2.5 \times 10^6$  cells per mL. At the time of harvest for cell banking, the viability is to remain above 95%. The cells are then centrifuged, and the cell pellet resuspended in the CD OptiCHO media with 7.5% dimethyl sulfoxide (DMSO) to a cell count of  $15 \times 10^6$  cells per mL per vial. Vials are cryopreserved for storage in liquid nitrogen.

**[0161]** A small-scale-up production is performed using the CHO cells as follows. The cells are scaled up for production in CD OptiCHO growth medium containing 100  $\mu$ M MSX at 37°C and fed every 2-4 days as needed, with CD OptiCHO growth medium supplemented with glucose and additional amino acids as necessary for approximately 14-21 days. The conditioned media supernatant harvested from the stable pool production run is clarified by centrifuge spinning. The protein is run over a Protein A (MabSelect, GE Healthcare, Little Chalfont, United Kingdom) column pre-equilibrated with binding buffer. Washing buffer is then passed through the column until the OD280 value (NanoDrop, Thermo Scientific) is measured to be at or near background levels. The insulin-Fc fusion protein is eluted using a low pH buffer, elution fractions are collected, and the OD280 value of each fraction is recorded. Fractions containing the target insulin-Fc fusion protein are pooled and optionally further filtered using a 0.2  $\mu$ M membrane filter.

**[0162]** The cell line is optionally further subcloned to monoclonality and optionally further selected for high titer insulin-Fc-fusion protein-expressing clones using the method of limiting dilution, a method known to those skilled in the art. After obtaining a high titer, monoclonal insulin-Fc fusion protein-expressing cell line, production of the insulin-Fc fusion

protein is accomplished as described above in growth medium without MSX, or optionally in growth medium containing MSX, to obtain a cell culture supernatant containing the recombinant, CHO-made, insulin-Fc fusion protein. The MSX concentration is optionally increased over time to exert additional selectivity for clones capable of yielding higher product titers.

Example 4: Purification of an Insulin-Fc Fusion Protein.

**[0163]** Purification of an insulin-Fc fusion protein was performed as follows. Conditioned media supernatants containing the secreted insulin-Fc fusion protein were harvested from the transiently transfected HEK, stably transfected HEK, or stably transfected CHO production runs and were clarified by centrifugation. The supernatant containing the desired insulin-Fc fusion protein was run over a Protein A column, washed with various wash buffers including 0.15-0.50M sodium chloride, and then eluted using a low pH solution. Afterwards, the eluted fractions containing the desired protein were pooled and buffer exchanged into 200 mM HEPES, 100 mM NaCl, 50 mM NaOAc, pH 7.0 buffer. A final filtration step was performed using a 0.2  $\mu$ m membrane filter. The final protein concentration was calculated from the solution optical density at 280 nm. Further optional purification by ion-exchange chromatography (e.g. using an anion exchange bead resin or a cation exchange bead resin), gel filtration chromatography, or other methods was performed as necessary.

Example 5: Purification of an Insulin-Fc Fusion Protein.

**[0164]** Purification of an insulin-Fc fusion protein is performed as follows. Conditioned media supernatants containing the secreted insulin-Fc fusion protein are harvested from the transiently transfected HEK, stably transfected HEK, or stably transfected CHO production runs and are clarified by centrifugation. The supernatant containing the desired insulin-Fc fusion protein is run over a Protein A column, washed with various wash buffers including 0.15-0.50M sodium chloride, and then eluted using a low pH solution. Afterwards, the eluted fractions containing the desired protein are pooled and buffer exchanged into 200 mM HEPES, 100 mM NaCl, 50 mM NaOAc, pH 7.0 buffer. A final filtration step is performed using a 0.2  $\mu$ m membrane filter. The final protein concentration is calculated from the solution optical density at 280 nm. Further optional purification by ion-exchange chromatography (e.g. using an anion exchange bead resin or a cation exchange bead resin), gel filtration chromatography, or other methods is performed as necessary.

Example 6: Structure Confirmation by Non-reducing and Reducing CE-SDS.

**[0165]** Capillary electrophoresis sodium dodecyl sulfate (CE-SDS) analysis was performed in a LabChip® GXII (Perkin Elmer, Waltham, MA) on a solution of a purified insulin-Fc fusion protein dissolved in 200 mM HEPES, 100 mM NaCl, 50 mM NaOAc, pH 7.0 buffer, and the electropherogram was plotted. Under non-reducing conditions, the sample was run against known molecular weight (MW) protein standards, and the eluting peak represented the 'apparent' MW of the insulin-Fc fusion protein homodimer.

**[0166]** Under reducing conditions (e.g. using beta-mercaptoethanol to break disulfide bonds of the insulin-Fc fusion homodimer), the apparent MW of the resulting insulin-Fc fusion protein monomer is compared against half the molecular weight of the insulin-Fc fusion protein homodimer as a way of determining that the structural purity of the insulin-Fc fusion protein is likely to be correct.

Example 7: Structure Confirmation by Non-reducing and Reducing CE-SDS.

**[0167]** Capillary electrophoresis sodium dodecyl sulfate (CE-SDS) analysis is performed in a LabChip® GXII (Perkin Elmer, Waltham, MA) on a solution of a purified insulin-Fc fusion protein dissolved in 200 mM HEPES, 100 mM NaCl, 50 mM NaOAc, pH 7.0 buffer, and the electropherogram is plotted. Under non-reducing conditions, the sample is run against known molecular weight (MW) protein standards, and the eluting peak represents the 'apparent' MW of the insulin-Fc fusion protein homodimer.

**[0168]** Under reducing conditions (e.g. using beta-mercaptoethanol to break disulfide bonds of the insulin-Fc fusion homodimer), the apparent MW of the resulting insulin-Fc fusion protein monomer is compared against half the molecular weight of the insulin-Fc fusion protein homodimer as a way of determining that the structural purity of the insulin-Fc fusion protein is likely to be correct.

Example 8: Sequence Identification by LC-MS with Glycan Removal for Compounds Containing Glycan.

**[0169]** To obtain an accurate estimate of the insulin-Fc fusion protein mass via mass spectroscopy (MS) in cases where the insulin-Fc fusion protein is naturally glycosylated, the sample was first treated to remove naturally occurring glycan that might interfere with the MS analysis. 100  $\mu$ L of a 2.5 mg/mL insulin-Fc fusion protein dissolved in 200 mM

HEPES, 100 mM NaCl, 50 mM NaOAc, pH 7.0 buffer solution was first buffer exchanged into 0.1 M Tris, pH 8.0 buffer containing 5 mM EDTA using a Zeba desalting column (Pierce, ThermoFisher Scientific, Waltham, MA). 1.67  $\mu$ L of PNGase F enzyme (Prozyme N-glycanase) was added to this solution in order to remove N-linked glycan present in the fusion protein (e.g., glycan linked to the side chain of the asparagine located at the cNg-N site), and the mixture was incubated at 37°C overnight in an incubator. The sample was then analyzed via LC-MS (NovaBioassays, Woburn, MA) resulting in a molecular mass of the molecule which corresponds to the desired homodimer without the glycan. This mass was then further corrected since the enzymatic process used to cleave the glycan from the cNg-asparagine also deaminates the asparagine side chain to form an aspartic acid, and in doing so the enzymatically treated homodimer gains 2 Da overall, corresponding to a mass of 1 Da for each chain present in the homodimer. Therefore, the actual molecular mass was the measured mass minus 2 Da to correct for the enzymatic modification of the insulin-Fc fusion protein structure in the analytical sample.

**[0170]** In cases where the insulin-Fc fusion protein amino acid composition prevents natural glycosylation from occurring at the cNg site, LC-MS (NovaBioassays, Woburn, MA) was used to directly obtain an accurate estimate of the insulin-Fc fusion protein mass without pretreatment of the molecule with PNGase enzyme.

Example 9: Sequence Identification by LC-MS with Glycan Removal for Compounds Containing Glycan.

**[0171]** To obtain an accurate estimate of the insulin-Fc fusion protein mass via mass spectroscopy (MS) in cases where the insulin-Fc fusion protein is naturally glycosylated, the sample is first treated to remove naturally occurring glycan that might interfere with the MS analysis. 100  $\mu$ L of a 2.5 mg/mL insulin-Fc fusion protein dissolved in 200 mM HEPES, 100 mM NaCl, 50 mM NaOAc, pH 7.0 buffer solution is first buffer exchanged into 0.1 M Tris, pH 8.0 buffer containing 5 mM EDTA using a Zeba desalting column (Pierce, ThermoFisher Scientific, Waltham, MA). 1.67  $\mu$ L of PNGase F enzyme (Prozyme N-glycanase) is added to this solution in order to remove N-linked glycan present in the fusion protein (e.g., glycan linked to the side chain of the asparagine located at the cNg-N site), and the mixture is incubated at 37°C overnight in an incubator. The sample is then analyzed via LC-MS (NovaBioassays, Woburn, MA) resulting in a molecular mass of the molecule which corresponds to the desired homodimer without the glycan. This mass is then further corrected since the enzymatic process used to cleave the glycan from the cNg-asparagine also deaminates the asparagine side chain to form an aspartic acid, and in doing so the enzymatically treated homodimer gains 2 Da overall, corresponding to a mass of 1 Da for each chain present in the homodimer. Therefore, the actual molecular mass is the measured mass minus 2 Da to correct for the enzymatic modification of the insulin-Fc fusion protein structure in the analytical sample.

**[0172]** In cases where the insulin-Fc fusion protein amino acid composition prevents natural glycosylation from occurring at the cNg site, LC-MS (NovaBioassays, Woburn, MA) can be used to directly obtain an accurate estimate of the insulin-Fc fusion protein mass without pretreatment of the molecule with PNGase enzyme.

Example 10: %Homodimer by Size-Exclusion Chromatography.

**[0173]** Size-exclusion chromatography (SEC-HPLC) of insulin-Fc fusion proteins was carried out using a Waters 2795HT HPLC (Waters Corporation, Milford, MA) connected to a 2998 Photodiode array at a wavelength of 280 nm. 100  $\mu$ L or less of a sample containing an insulin-Fc fusion protein of interest was injected into a MAbPac SEC-1, 5  $\mu$ m, 4  $\times$  300 mm column (ThermoFisher Scientific, Waltham, MA) operating at a flow rate of 0.2 mL/min and with a mobile phase comprising 50 mM sodium phosphate, 300 mM NaCl, and 0.05% w/v sodium azide, pH 6.2. The MAbPac SEC-1 column operates on the principle of molecular size separation. Therefore, larger soluble insulin-Fc aggregates (e.g. multimers of insulin-Fc fusion protein homodimers) eluted at earlier retention times, and the non-aggregated homodimers eluted at later retention times. In separating the mixture of homodimers from aggregated multimeric homodimers via analytical SEC-HPLC, the purity of the insulin-Fc fusion protein solution in terms of the percentage of non-aggregated homodimer was ascertained.

Example 11: %Homodimer by Size-Exclusion Chromatography.

**[0174]** Size-exclusion chromatography (SEC-HPLC) of insulin-Fc fusion proteins is carried out using a Waters 2795HT HPLC (Waters Corporation, Milford, MA) connected to a 2998 Photodiode array at a wavelength of 280 nm. 100  $\mu$ L or less of a sample containing an insulin-Fc fusion protein of interest is injected into a MAbPac SEC-1, 5  $\mu$ m, 4  $\times$  300 mm column (Thermo Fisher Scientific, Waltham, MA) operating at a flow rate of 0.2 mL/min and with a mobile phase comprising 50 mM sodium phosphate, 300 mM NaCl, and 0.05% w/v sodium azide, pH 6.2. The MAbPac SEC-1 column operates on the principle of molecular size separation. Therefore, larger soluble insulin-Fc aggregates (e.g. multimers of insulin-Fc fusion protein homodimers) will elute at earlier retention times, and the non-aggregated homodimers will elute at later retention times. In separating the mixture of homodimers from aggregated multimeric homodimers via analytical SEC-

HPLC, the purity of the insulin-Fc fusion protein solution in terms of the percentage of non-aggregated homodimer is ascertained.

*Example 12: In vitro IM-9 Insulin Receptor (IR) Binding of an Exemplary Insulin-Fc Fusion Protein at 4°C.*

**[0175]** Human IM-9 cells (ATTC# CCL-159) that express human IR were cultured and maintained in complete RPMI 5% FBS medium at 70-80% confluency. Cultures of IM-9 cells were centrifuged at 250×g (~1000 rpm) for 10 min to pellet the cells. Cells were washed once with HBSS or PBS buffer, resuspended in cold FACS staining medium (HBSS/2mM EDTA/0.1% Na-azide + 4% horse serum) to a concentration of 8×10<sup>6</sup> cells/mL and kept on ice or at 4°C until test solutions were made. The insulin-Fc protein was diluted in FACS buffer in 1:3 serial dilutions as 2× concentrations in 1.2 mL tubes (approx. 60 µL volume of each dilution), and the solutions were kept cold on ice until ready for pipetting.

**[0176]** Biotinylated-RHI was diluted in FACS staining medium to a concentration of 1.25 µg/mL. 40 µL of the serially diluted test compound and 8 µL of 1.25 µg/mL Biotin-RHI were added into each well of a V bottom microtiter plate, mixed by slow vortexing, and placed on ice. 40 µL of an IM-9 cell suspension (8×10<sup>6</sup> cells/mL) was then added to each well by multichannel pipette, mixed again gently and incubated on ice for 30 min to allow competitive binding on the IR on IM-9 cells. Cells were then washed twice with 275µL of ice-cold FACS wash buffer (HBSS/2mM EDTA/0.1% Na-azide + 0.5% horse serum) by centrifuging the V-bottom plate at 3000 rpm for 3 min and aspirating the supernatant. Cells were then resuspended in 40µL of FACS staining medium containing 1:100 diluted Streptavidin-PE (Life Technologies) for 20 min on ice. Cells were then washed once with 275 µL of ice-cold FACS buffer and finally fixed with 3% paraformaldehyde for 10 min at room temp. Cells were then washed once with 275 µL of ice-cold FACS buffer and resuspended in 250µL of FACS buffer for analysis.

**[0177]** The V-bottom plates containing cells were then analyzed on a Guava 8-HT flow cytometer (Millipore). Biotinylated-RHI binding to IR was quantitated by the median fluorescence intensity (MFI) of the cells on the FACS FL-2 channel for each concentration of the test compound. Control wells were labeled only with biotinylated-RHI and were used to calculate the percent (%) inhibition resulting from each test compound concentration. The % inhibition by test compounds of biotinylated-RHI binding on IM-9 cells was plotted against log concentrations of the test compound, and the resulting IC<sub>50</sub> values were calculated using GraphPad Prism (GraphPad Software, La Jolla, CA) for the test compounds. Lower IC<sub>50</sub> values of the test compound therefore indicate greater levels of biotinylated-RHI inhibition at lower concentrations indicating stronger binding of the insulin-Fc fusion protein to the IR. A control compound, such as unlabeled recombinant human insulin (RHI) was also used as an internal standard to generate an RHI IC<sub>50</sub> against which a given compound IC<sub>50</sub> could be ratioed (IC<sub>50</sub>(compound)/IC<sub>50</sub>(RHI)). Lower IC<sub>50</sub> ratios have more similar binding to RHI (stronger binding to IR), while higher IC<sub>50</sub> ratios have weaker binding to the IR relative to RHI.

*Example 13: In vitro IM-9 Insulin Receptor (IR) Binding of an Exemplary Insulin-Fc Fusion Protein at 4°C.*

**[0178]** Human IM-9 cells (ATTC# CCL-159) that express human IR are cultured and maintained in complete RPMI 5% FBS medium at 70-80% confluency. Cultures of IM-9 cells are centrifuged at 250×g (~1000 rpm) for 10 min to pellet the cells. Cells are washed once with HBSS or PBS buffer, resuspended in cold FACS staining medium (HBSS/2mM EDTA/0.1% Na-azide + 4% horse serum) to a concentration of 8×10<sup>6</sup> cells/mL and kept on ice or at 4°C until test solutions are made. The insulin-Fc protein is diluted in FACS buffer in 1:3 serial dilutions as 2× concentrations in 1.2 mL tubes (approx. 60 µL volume of each dilution), and the solutions are kept cold on ice until ready for pipetting.

**[0179]** Biotinylated-RHI is diluted in FACS staining medium to a concentration of 1.25 µg/mL. 40 µL of the serially diluted test compound and 8 µL of 1.25 µg/mL Biotin-RHI are added into each well of a V bottom microtiter plate, mixed by slow vortexing, and placed on ice. 40 µL of an IM-9 cell suspension (8×10<sup>6</sup> cells/mL) is then added to each well by multichannel pipette, mixed again gently and incubated on ice for 30 min to allow competitive binding on the IR on IM-9 cells. Cells are then washed twice with 275µL of ice-cold FACS wash buffer (HBSS/2mM EDTA/0.1% Na-azide + 0.5% horse serum) by centrifuging the V-bottom plate at 3000 rpm for 3 min and aspirating the supernatant. Cells are then resuspended in 40µL of FACS staining medium containing 1:100 diluted Streptavidin-PE (Life Technologies) for 20 min on ice. Cells are then washed once with 275 µL of ice-cold FACS buffer and finally fixed with 3% paraformaldehyde for 10 min at room temp. Cells are then washed once with 275 µL of ice-cold FACS buffer and resuspended in 250µL of FACS buffer for analysis.

**[0180]** The V-bottom plates containing cells are then analyzed on a Guava 8-HT flow cytometer (Millipore). Biotinylated-RHI binding to IR is quantitated by the median fluorescence intensity (MFI) of the cells on the FACS FL-2 channel for each concentration of the test compound. Control wells are labeled only with biotinylated-RHI and are used to calculate the percent (%) inhibition resulting from each test compound concentration. The % inhibition by test compounds of biotinylated-RHI binding on IM-9 cells is plotted against log concentrations of the test compound, and the resulting IC<sub>50</sub> values are calculated using GraphPad Prism (GraphPad Software, La Jolla, CA) for the test compounds. Lower IC<sub>50</sub> values of the test compound therefore indicate greater levels of biotinylated-RHI inhibition at lower concentrations indi-

cating stronger binding of the insulin-Fc fusion protein to the IR. A control compound, such as unlabeled recombinant human insulin (RHI) is also used as an internal standard to generate an RHI IC<sub>50</sub> against which a given compound IC<sub>50</sub> could be ratioed (IC<sub>50</sub>(compound)/IC<sub>50</sub>(RHI)). Lower IC<sub>50</sub> ratios have more similar binding to RHI (stronger binding to IR), while higher IC<sub>50</sub> ratios have weaker binding to the IR relative to RHI.

*Example 14: In vitro Human Fc(gamma)RI Binding Affinity Assay.*

**[0181]** The binding of insulin-Fc fusion proteins to the Fc(gamma)RI at pH 7.4 was conducted using an ELISA assay as follows using human Fc(gamma)RI (i.e., rhFc(gamma)RI). Insulin-Fc fusion proteins were diluted to 10 µg/mL in sodium bicarbonate buffer at pH 9.6 and coated on Maxisorp (Nunc) microtiter plates overnight at 4°C, after which the microplate strips were washed 5 times with PBST (PBS/0.05% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Serial dilutions of biotinylated rhFc(gamma)RI (recombinant human Fc(gamma)RI; R&D Systems) were prepared in PBST/10% Superblock buffer from 3000 ng/mL to 4.1 ng/mL and loaded at 100 µL/well onto the microplate strips coated with insulin-Fc fusion protein. The microtiter plate was incubated for 1 hour at room temperature after which time the microplate strips were washed 5 times with PBST and then loaded with 100 µL/well of streptavidin-HRP diluted 1:10000 in PBST/10% Superblock buffer. After incubating for 45 min, the microplate strips were washed again 5 times with PBST. TMB was added to reveal the bound Fc(gamma)RI proteins and stopped with ELISA stop reagent (Boston Bioproducts). The plate was read in an ELISA plate reader at 450 nm, and the OD450 values (proportional to the binding of rhFc(gamma)RI to insulin-Fc protein) were plotted against log concentrations of rhFc(gamma)RI added to each well to generate binding curves using GraphPad Prism software. For groupings of compounds with somewhat similar curves, the OD450 at one of the higher concentrations, for instance at rhFc(gamma)RI concentrations of 3000 ng/mL, can be used to identify differences between coated insulin-Fc fusion protein compounds. In order to compare differences across multiple insulin-Fc fusion proteins run at different times, the human Fc(gamma)RI assay OD450 Ratio was calculated as the OD450 value of a test insulin-Fc fusion protein compound obtained at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL, divided by the OD450 value of a reference insulin-Fc fusion protein of SEQ ID NO: 76 obtained at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL.

*Example 15: In vitro Human Fc(gama)RI Binding Affinity Assay.*

**[0182]** The binding of insulin-Fc fusion proteins to the Fc(gamma)RI at pH 7.4 is conducted using an ELISA assay as follows using human Fc(gamma)RI (i.e., rhFc(gamma)RI). Insulin-Fc fusion proteins are diluted to 10 µg/mL in sodium bicarbonate buffer at pH 9.6 and coated on Maxisorp (Nunc) microtiter plates overnight at 4°C, after which the microplate strips are washed 5 times with PBST (PBS/0.05% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Serial dilutions of biotinylated rhFc(gamma)RI (recombinant human Fc(gamma)RI; R&D Systems) are prepared in PBST/10% Superblock buffer from 3000 ng/mL to 4.1 ng/mL and loaded at 100 µL/well onto the microplate strips coated with insulin-Fc fusion protein. The microtiter plate is incubated for 1 hour at room temperature after which time the microplate strips are washed 5 times with PBST and then loaded with 100 µL/well of streptavidin-HRP diluted 1:10000 in PBST/10% Superblock buffer. After incubating for 45 min, the microplate strips are washed again 5 times with PBST. TMB is added to reveal the bound Fc(gamma)RI proteins and is stopped with ELISA stop reagent (Boston Bioproducts). The plate is read in an ELISA plate reader at 450 nm, and the OD450 values (proportional to the binding of rhFc(gamma)RI to insulin-Fc protein) are plotted against log concentrations of rhFc(gamma)RI added to each well to generate binding curves using GraphPad Prism software. For groupings of compounds with somewhat similar curves, the OD450 at one of the higher concentrations, for instance at rhFc(gamma)RI concentrations of 3000 ng/mL, can be used to identify differences between coated insulin-Fc fusion protein compounds. In order to compare differences across multiple insulin-Fc fusion proteins run at different times, the human Fc(gamma)RI assay OD450 Ratio is calculated as the OD450 value of a test insulin-Fc fusion protein compound obtained at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL, divided by the OD450 value of a reference insulin-Fc fusion protein of SEQ ID NO: 76 obtained at a biotinylated-Fc(gamma)RI concentration of 3000 ng/mL.

*Example 16: In Vitro C1q Binding Affinity Assay.*

**[0183]** The binding of insulin-Fc fusion proteins to complement component C1q at pH 7.4 was conducted using an ELISA assay as follows using human complement component C1q. Insulin-Fc fusion proteins were diluted to 10 µg/mL in sodium bicarbonate buffer at pH 9.6 and coated on Maxisorp (Nunc) microtiter plates overnight at 4°C, after which the microplate strips were washed 5 times with PBST (PBS/0.05% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Serial dilutions of biotinylated complement component C1q (human complement component C1q; Sigma-Aldrich) were prepared in PBST/10% Superblock buffer from 1000 ng/mL to 1.4 ng/mL and loaded at 100 µL/well onto the microplate strips coated with insulin-Fc fusion protein. The microtiter plate was incubated for 1

hour at room temperature after which the microplate strips were washed 5 times with PBST and then loaded with 100  $\mu$ L/well of streptavidin-HRP diluted 1:12000 in PBST/10% Superblock buffer. After incubating for 45 min, the microplate strips were washed again 5 times with PBST. TMB was added to reveal the bound complement C1q proteins and stopped with ELISA stop reagent (Boston Bioproducts). The plate absorbance was read in an ELISA plate reader at 450 nm (OD450), and the OD450 values (proportional to the binding of complement component C1q to insulin-Fc protein) were plotted against log concentrations of complement component C1q added to each well to generate binding curves using GraphPad Prism software. For groupings of compounds with somewhat similar curves, the OD450 at one of the higher concentrations, for instance at complement component C1q concentrations of 1000 ng/mL, was used to identify differences between coated insulin-Fc fusion protein compounds. In order to further compare differences across multiple insulin-Fc fusion proteins run at different times, the human C1q assay OD450 Ratio was calculated as the OD450 of a test insulin-Fc fusion protein compound obtained at a biotinylated-C1q concentration of 1000 ng/mL, divided by the OD450 of a reference insulin-Fc fusion protein of SEQ ID NO: 76 obtained at a biotinylated-C1q concentration of 1000 ng/mL.

*Example 17: In Vitro C1q Binding Affinity Assay.*

**[0184]** The binding of insulin-Fc fusion proteins to complement component C1q at pH 7.4 is conducted using an ELISA assay as follows using human complement component C1q. Insulin-Fc compounds are diluted to 10  $\mu$ g/mL in sodium bicarbonate buffer at pH 9.6 and coated on Maxisorp (Nunc) microtiter plates overnight at 4°C, after which the microplate strips are washed 5 times with PBST (PBS/0.05% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Serial dilutions of biotinylated complement component C1q (human complement component C1q; Sigma-Aldrich) are prepared in PBST/10% Superblock buffer from 1000 ng/mL to 1.4 ng/mL and loaded at 100  $\mu$ L/well onto the microplate strips coated with insulin-Fc fusion protein. The microtiter plate is incubated for 1 hour at room temperature after which the microplate strips are washed 5 times with PBST and then loaded with 100  $\mu$ L/well of streptavidin-HRP diluted 1:12000 in PBST/10% Superblock buffer. After incubating for 45 min, the microplate strips are washed again 5 times with PBST. TMB is added to reveal the bound complement C1q proteins and stopped with ELISA stop reagent (Boston Bioproducts). The plate absorbance is read in an ELISA plate reader at 450 nm (OD450), and the OD450 values (proportional to the binding of complement component C1q to insulin-Fc protein) are plotted against log concentrations of complement component C1q added to each well to generate binding curves using GraphPad Prism software. For groupings of compounds with somewhat similar curves, the OD450 at one of the higher concentrations, for instance at complement component C1q concentrations of 1000 ng/mL, can be used to identify differences between coated insulin-Fc fusion protein compounds. In order to further compare differences across multiple insulin-Fc fusion proteins run at different times, the human C1q assay OD450 Ratio is calculated as the OD450 of a test insulin-Fc fusion protein compound obtained at a biotinylated-C1q concentration of 1000 ng/mL, divided by the OD450 of a reference insulin-Fc fusion protein of SEQ ID NO: 76 obtained at a biotinylated-C1q concentration of 1000 ng/mL.

*Example 18: In vitro Measurement of Insulin-Fc Fusion Protein Affinity for the Canine FcRn receptor.*

**[0185]** In vitro binding affinity of insulin-Fc fusion proteins containing Fc fragments of canine IgG origin to the canine FcRn receptor was measured via an ELISA technique conducted at a solution pH of 5.5. The slightly acidic pH is the preferred binding environment for Fc fragment-containing molecules to bind to the FcRn receptor. In vivo, cells express FcRn on their surfaces and internally in the endosomes. As molecules containing Fc fragments are brought into the cell through natural processes (e.g. pinocytosis or endocytosis), the pH changes to a lower pH in the endosomes, where the FcRn receptor binds to Fc fragment-containing molecules that would otherwise be degraded in the endosomal-lysosomal compartments, thereby allowing these molecules to recycle back to the cellular surface where the pH is closer to neutral (e.g., pH 7.0-7.4). Neutral pH disfavors binding to the FcRn receptor and allows release of the Fc-fragment containing molecules back into circulation. This is a primary mechanism by which Fc fragment-containing molecules exhibit prolonged circulatory pharmacokinetic half-lives in vivo.

**[0186]** Insulin-Fc fusion proteins comprising Fc fragments of canine origin were diluted to 10  $\mu$ g/ml in sodium bicarbonate pH 9.6 buffer and coated in duplicate on Maxisorb ELISA plate strips for 1-2 hours at RT. The strips were then washed 4 times with PBST (PBS/0.1% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Strips for FcRn binding were then washed again twice with pH 5.5 MES/NaCl/Tween (50mM MES/150mM NaCl/0.1% Tween-20) buffer before addition of the FcRn reagent (biotinylated canine FcRn; Immunitrack). Serial dilutions (1:3X dilutions) of biotinylated FcRn reagent were prepared in pH 5.5 MES/NaCl/Tween/10%Superblock buffer at concentrations from 1000 ng/ml to 0.45 ng/mL and loaded at 100  $\mu$ L/well using a multichannel pipettor onto the strips coated with the insulin-Fc fusion protein compounds. The assay plate was then incubated for 1 hour at room temperature. FcRn binding strips were washed 4 times with pH 5.5 MES/NaCl/Tween buffer and then loaded with 100  $\mu$ L/well streptavidin-HRP diluted 1:10000 in pH 5.5 MES/NaCl /10%Superblock buffer. After incubating for 45 minutes, strips were washed

again 4 times with pH 5.5 MES/NaCl/Tween buffer. TMB was finally added to reveal the bound biotinylated-canine FcRn reagent, and the color development was stopped with the ELISA stop reagent. The plate was read in an ELISA plate reader at a wavelength of 450 nm. The OD values (proportional to the binding of canine-FcRn to the insulin-Fc fusion protein test compounds) were plotted against log concentrations of FcRn added to each well to generate binding curves using GraphPad Prism software. EC50 values for each binding curve were calculated to compare between different compounds.

*Example 19: In vitro Measurement of Insulin-Fc Fusion Protein Affinity for the Human FcRn receptor.*

**[0187]** In vitro binding affinity of insulin-Fc fusion proteins containing Fc fragments of human IgG origin to the human FcRn receptor was measured via an ELISA technique conducted at a solution pH of 5.5. The slightly acidic pH is the preferred binding environment for Fc fragment-containing molecules to bind to the FcRn receptor. In vivo, cells express FcRn on their surfaces and internally in the endosomes. As molecules containing Fc fragments are brought into the cell through natural processes (e.g. pinocytosis or endocytosis), the pH changes to a lower pH in the endosomes, where the FcRn receptor binds to Fc fragment-containing molecules that would otherwise be degraded in the endosomal-lysosomal compartments, thereby allowing these molecules to recycle back to the cellular surface where the pH is closer to neutral (e.g., pH 7.0-7.4). Neutral pH disfavors binding to the FcRn receptor and allows release of the Fc-fragment containing molecules back into circulation. This is a primary mechanism by which Fc fragment-containing molecules exhibit prolonged circulatory pharmacokinetic half-lives in vivo.

**[0188]** Insulin-Fc fusion proteins comprising Fc fragments were diluted to 10  $\mu$ g/mL in sodium bicarbonate pH 9.6 buffer and coated in duplicate on Maxisorb ELISA plate strips for 1-2 hours at RT. The strips were then washed 5 times with PBST (PBS/0.1% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Strips for FcRn binding were then washed again 3 times with pH 5.5 MES/NaCl/Tween (50mM MES/150mM NaCl/0.1% Tween-20) buffer before addition of the FcRn reagent (biotinylated human FcRn; Immunitrack). Serial dilutions (1:3X dilutions) of biotinylated FcRn reagent were prepared in pH 5.5 MES/NaCl/Tween/5%Superblock buffer at concentrations from 6000 ng/mL to 8.23 ng/mL and loaded at 100  $\mu$ L/well using a multichannel pipettor onto the strips coated with the insulin-Fc fusion protein compounds. The assay plate was then incubated for 1.5 hours at room temperature. FcRn binding strips were washed 4 times with pH 5.5 MES/NaCl/Tween buffer and then loaded with 100  $\mu$ L/well streptavidin-HRP diluted 1:10000 in pH 5.5 MES/NaCl /5%Superblock buffer. After incubating for 45 minutes, strips were washed again 5 times with pH 5.5 MES/NaCl/Tween buffer. TMB was finally added to reveal the bound biotinylated-FcRn reagent, and the color development was stopped with the ELISA stop reagent. The plate was read in an ELISA plate reader at a wavelength of 450 nm. The OD450 values (proportional to the binding of human-FcRn to the insulin-Fc fusion protein test compounds) were plotted against log concentrations of FcRn added to each well to generate binding curves using GraphPad Prism software. EC50 values for each binding curve were calculated to compare between different compounds.

*Example 20: In vitro Measurement of Insulin-Fc Fusion Protein Affinity for the Human FcRn receptor.*

**[0189]** In vitro binding affinity of insulin-Fc fusion proteins containing Fc fragments of human IgG origin to the human FcRn receptor is measured via an ELISA technique conducted at a solution pH of 5.5. The slightly acidic pH is the preferred binding environment for Fc fragment-containing molecules to bind to the FcRn receptor. In vivo, cells express FcRn on their surfaces and internally in the endosomes. As molecules containing Fc fragments are brought into the cell through natural processes (e.g. pinocytosis or endocytosis), the pH changes to a lower pH in the endosomes, where the FcRn receptor binds to Fc fragment-containing molecules that would otherwise be degraded in the endosomal-lysosomal compartments, thereby allowing these molecules to recycle back to the cellular surface where the pH is closer to neutral (e.g., pH 7.0-7.4). Neutral pH disfavors binding to the FcRn receptor and allows release of the Fc-fragment containing molecules back into circulation. This is a primary mechanism by which Fc fragment-containing molecules exhibit prolonged circulatory pharmacokinetic half-lives in vivo.

**[0190]** Insulin-Fc fusion proteins comprising Fc fragments are diluted to 10  $\mu$ g/mL in sodium bicarbonate pH 9.6 buffer and coated in duplicate on Maxisorb ELISA plate strips for 1-2 hours at RT. The strips are then washed 5 times with PBST (PBS/0.1% Tween-20) buffer and blocked with Superblock blocking reagent (ThermoFisher). Strips for FcRn binding are then washed again 3 times with pH 5.5 MES/NaCl/Tween (50mM MES/150mM NaCl/0.1% Tween-20) buffer before addition of the FcRn reagent (biotinylated human FcRn; Immunitrack). Serial dilutions (1:3X dilutions) of biotinylated FcRn reagent are prepared in pH 5.5 MES/NaCl/Tween/5%Superblock buffer at concentrations from 6000 ng/mL to 8.23 ng/mL and loaded at 100  $\mu$ L/well using a multichannel pipettor onto the strips coated with the insulin-Fc fusion protein compounds. The assay plate is then incubated for 1.5 hours at room temperature. FcRn binding strips are washed 4 times with pH 5.5 MES/NaCl/Tween buffer and then loaded with 100  $\mu$ L/well streptavidin-HRP diluted 1:10000 in pH 5.5 MES/NaCl /5%Superblock buffer. After incubating for 45 minutes, strips are washed again 5 times with pH 5.5 MES/NaCl/Tween buffer. TMB is finally added to reveal the bound biotinylated-FcRn reagent, and the color development

is stopped with the ELISA stop reagent. The plate is read in an ELISA plate reader at a wavelength of 450 nm. The OD450 values (proportional to the binding of human-FcRn to the insulin-Fc fusion protein test compounds) are plotted against log concentrations of FcRn added to each well to generate binding curves using GraphPad Prism software. EC50 values for each binding curve are calculated to compare between different compounds.

**Example 21: Generalized Procedure for Determination of In vivo Pharmacodynamics (PD) After Single Administration of Canine Insulin Fc-Fusion Proteins in Dogs.**

**[0191]** Insulin-Fc fusion proteins were assessed for their effects on fasting blood glucose levels in dogs as follows. N=1, 2, 3 or more healthy, antibody-naïve, dogs weighing approximately 10-15 kg were used, one for each insulin-Fc fusion protein. Animals were also observed twice daily for signs of anaphylaxis, lethargy, distress, pain, etc., and, optionally for some compounds, treatment was continued for an additional three weekly subcutaneous injections or more to observe if the glucose lowering capability of the compounds lessened over time, a key sign of potential induction of neutralizing anti-drug antibodies. On day 0, the animals received a single injection either via intravenous or subcutaneous administration of a pharmaceutical composition containing an insulin Fc-fusion protein homodimer at a concentration between 1 and 10 mg/mL in a solution of between 10-50 mM sodium hydrogen phosphate, 50-150 mM sodium chloride, 0.005-0.05% v/v Tween-80, and optionally a bacteriostat (e.g. phenol, m-cresol, or methylparaben) at a concentration of between 0.02-1.00 mg/mL, at a solution pH of between 7.0-8.0, at a dose of 0.08-0.80 mg insulin-Fc fusion protein/kg (or approximately equivalent to 1.2-12.3 nmol/kg or approximately equivalent to 0.4-4.0 U/kg insulin equivalent on molar basis). On day 0, blood was collected from a suitable vein immediately prior to injection and at 15, 30, 45, 60, 120, 240, 360, and 480 min and at 1, 2, 3, 4, 5, 6, and 7 days post injection.

**[0192]** For each time point, a minimum of 1 mL of whole blood was collected. A glucose level reading was immediately determined using a glucose meter (ACCU-CHEK® Aviva Plus), which required approximately one drop of blood. Average % fasting blood glucose levels (% FBGL) from day 0 to day 7 were plotted to assess the bioactivity of a given insulin-Fc fusion protein.

**Example 22: Generalized Procedure for Determination of In vivo Pharmacodynamics (PD) After Repeated Administration of Canine Insulin-Fc Fusion Proteins in Dogs.**

**[0193]** Insulin-Fc fusion proteins were assessed for their effects on blood glucose levels over repeated injections as follows. Healthy, antibody-naïve, dogs weighing approximately between 10kg and 20 kg were used, and each animal was administered doses of an insulin-Fc fusion protein. Animals were observed twice daily for signs of anaphylaxis, lethargy, distress, pain, and other negative side effects, and optionally for some compounds, treatment was continued for up to an additional two to five subcutaneous injections to observe if the glucose lowering capability of the compounds decreased over time, indicating the possible presence of neutralizing anti-drug antibodies in vivo. On day 0, the animals received a single subcutaneous injection of a pharmaceutical composition containing an insulin Fc-fusion protein in a solution of 10-50 mM sodium hydrogen phosphate, 50-150 mM sodium chloride, 0.005-0.05% v/v Tween-80, and optionally a bacteriostat (e.g. phenol, m-cresol, or methylparaben) at a concentration of between 0.02-1.00 mg/mL, at a solution pH of between 7.0-8.0, at a dose of 0.08-0.80 mg insulin-Fc fusion protein/kg (or approximately equivalent to 1.2-12.3 nmol/kg or approximately equivalent to 0.4-4.0 U/kg insulin equivalent on molar basis). On day 0, blood was collected from a suitable vein immediately prior to injection and at 15, 30, 45, 60, 120, 240, 360, and 480 min and at 1, 2, 3, 4, 5, 6, and 7 days post injection.

**[0194]** Subsequent subcutaneous injections were given no more frequently than once-weekly, and in some cases the injections were given at different intervals based on the pharmacodynamics of a given insulin-Fc fusion protein formulation. Subsequent injections for each insulin-Fc fusion protein were adjusted to higher or lower doses, depending on the demonstrated pharmacodynamics of the insulin-Fc fusion protein. For instance, if the dose of a first injection on day 0 was found to be ineffective at lowering blood glucose, the subsequent dose levels of injected insulin-Fc fusion protein were adjusted upward. In a similar manner, if the dose of a first injection on day 0 was found to lower glucose in too strong a manner, then subsequent dose levels of injected insulin-Fc fusion protein were adjusted downward. It was also found that interim doses or final doses could be adjusted in a similar manner as needed. For each dose, blood was collected from a suitable vein just immediately prior to injection and at 15, 30, 45, 60, 120, 240, 360, and 480 min and at 1, 2, 3, 4, 5, 6, 7 days (and optionally 14 days) post injection. For each time point, a minimum of 1 mL of whole blood was collected. A glucose level reading was immediately determined using a glucose meter (ACCU-CHEK® Aviva Plus), which required approximately one drop of blood. Average % fasting blood glucose levels (% FBGL) from throughout the study were plotted against time which allows the bioactivity of a fusion protein to be determined.

**[0195]** To determine the bioactivity of each dose, an area-over-the-curve (AOC) analysis was conducted as follows. After constructing the %FBGL versus time data, the data were then entered into data analysis software (GraphPad Prism, GraphPad Software, San Diego CA). The software was used to first conduct an area-under-the curve analysis

(AUC) to integrate the area under the %FBGL vs. time curve for each dose. To convert the AUC data into the desired AOC data, the following equation was used:  $AOC = TPA - AUC$ ; where TPA is the total possible area obtained by multiplying each dose lifetime (e.g., 7 days, 14 days, etc.) by 100% (where 100% represents the  $y = 100\%$  of the %FBGL vs. time curve). For example, given a dose lifetime of 7 days and a calculated AUC of 500 %FBGL days, a calculation gave the following for AOC:  $AOC = (100\% \text{ FBGL} \times 7 \text{ days}) - (500\% \text{ FBGL days}) = 200\% \text{ FBGL days}$ . The analysis was performed for each injected dose in a series of injected doses to obtain the AOC values for injection 1, injection 2, injection 3, etc.

**[0196]** As the doses of insulin-Fc fusion protein may vary as previously discussed, it is often more convenient to normalize all calculated AOC values for a given insulin-Fc fusion protein to a particular dose of that insulin-Fc fusion protein. Doing so allows for convenient comparison of the glucose-lowering potency of an insulin-Fc fusion protein across multiple injections, even if the dose levels change across the injections of a given study. Normalized AOC (NAOC) for a given dose was calculated as follows:  $NAOC = AOC / D$  with units of %FBGL-days/kg/mg; where D is the actual dose injected into the animal in mg/kg. NAOC values were calculated for each injection in a series of injections for a given animal and were averaged across a group of animals receiving the same insulin-Fc fusion protein formulation.

**[0197]** The NAOC ratio (NAOCR) was calculated for each injection in a series of injections for a given animal by taking the NAOC values for each injection (e.g. injections 1, 2, 3,...N) and dividing each NAOC for a given injection by the NAOC from injection 1 as follows:  $NAOCR = (NAOC(N^{\text{th}} \text{ injection}) / NAOC(\text{injection } 1))$ . By evaluating the NAOCR of a given insulin-Fc fusion protein formulation for the  $N^{\text{th}}$  injection in a series of injections, it was possible to determine whether the in vivo glucose lowering activity of a given insulin-Fc fusion protein had substantially retained its bioactivity over a series of N doses (e.g., NAOCR for the Nth dose of greater than 0.5) or whether the in vivo glucose lowering activity of a given insulin-Fc fusion protein had lost a substantial portion of its potency (e.g., NAOCR of the  $N^{\text{th}}$  dose is less than 0.5) over a course of N doses, indicating the potential formation of neutralizing anti-drug antibodies in vivo. In preferred embodiments, the ratio of NAOC following the third subcutaneous injection to the NAOC following the first subcutaneous injection was greater than 0.5 (i.e., the NAOCR of the third subcutaneous injection was greater than 0.5).

*Example 23: Generalized Procedure for the Determination of In vivo Pharmacokinetics (PK) of Canine Insulin-Fc Fusion Proteins in Canine Serum.*

**[0198]** An assay was constructed for measuring the concentrations of insulin-Fc fusion proteins comprising Fc fragments of a canine isotype in canine serum as follows. The assay comprised a sandwich ELISA format in which therapeutic compounds in serum samples were captured by an anti-insulin/proinsulin monoclonal antibody (mAb) coated on the ELISA plates and were then detected by an HRP-conjugated anti-canine IgG Fc specific antibody followed by use of a TMB substrate system for color development. Maxisorp ELISA Plates (Nunc) were coated with the anti-insulin mAb clone D6C4 (Biorad) in coating buffer (pH=9.6 sodium carbonate-sodium biocarbonate buffer) at 5  $\mu\text{g/ml}$  overnight at 4°C. Plates were then washed 5 times with PBST (PBS + 0.05% Tween 20) and blocked for a minimum of one hour at room temperature (or overnight at 4C) with SuperBlock blocking solution (ThermoFisher). Test serum samples were diluted to 1:20 in PBST/SB/20%HS sample dilution buffer (PBS+0.1%Tween 20+10% SuperBlock+20% horse serum). For making a standard curve, the insulin-Fc fusion protein of interest was diluted in sample dilution buffer (PBST/SB/20%HS) + 5% of pooled beagle serum (BioIVT) from a concentration range of 200 ng/ml to 0.82 ng/ml in 1:2.5 serial dilutions. Standards and diluted serum samples were added to the blocked plates at 100  $\mu\text{l/well}$  in duplicate and were incubated for 1 hour at room temperature. Following incubation, samples and standards were washed 5 times with PBST. HRP-conjugated goat anti-canine IgG Fc (Sigma) detection antibody was diluted to about 1:15,000 in PBST/SB/20%HS buffer and 100  $\mu\text{l}$  was added to all the wells and incubated for 45 minutes at room temperature in the dark. Plates were washed 5 times with PBST and once with deionized water and were developed by the addition of 100  $\mu\text{l/well}$  TMB (Invitrogen) for 8-10 minutes at room temperature. Color development was then stopped by the addition of 100  $\mu\text{l/well}$  ELISA Stop Solution (Boston Bioproducts) and the absorbance was read at 450 nm using a SpectraMax plate reader (Molecular Devices) within 30 minutes. Concentrations of insulin-Fc fusion protein compounds in the samples were calculated by interpolation on a 4-PL curve using SoftMaxPro software.

*Example 24: Assay Protocol for Measuring Anti-Drug Antibodies in Canine Serum.*

**[0199]** Maxisorp ELISA Plates (Nunc) were coated with the insulin-Fc fusion protein of interest diluted in coating buffer (pH=9.6 Carbonate-Biocarbonate buffer) at 10  $\mu\text{g/mL}$  overnight at 4°C for measuring ADAs against the test compound. For measuring ADAs against the insulin portion of the insulin-Fc fusion protein containing an Fc fragment of canine IgG origin, plates were coated with purified insulin at 30  $\mu\text{g/mL}$  in coating buffer. Plates were then washed 5 times with PBST (PBS + 0.05% Tween 20) and blocked for at least 1 hour (or overnight) with SuperBlock blocking solution (ThermoFisher, Waltham MA). For calculating the ADAs in canine IgG units, strips were directly coated with 1:2 serial dilutions of canine IgG (Jackson ImmunoResearch Laboratories, West Grove PA) in pH=9.6 Carb-Biocarb coating buffer at concentrations

between 300-4.69ng/ml overnight at 4°C and used to create a 7-point pseudo-standard curve. The standards strip plates were also washed and blocked with SuperBlock blocking solution for at least 1 hour (or overnight).

[0200] Test serum samples were diluted to greater than or equal to 1: 100 (typically tested as 1:200) in PBST/SB/20%HS sample dilution buffer (PBS+0.1% Tween 20+10% SuperBlock+20% horse serum) and added to the insulin-Fc fusion protein coated (or RHI coated) strips at 100 µL/well in duplicate. Duplicate strips of canine IgG coated standard strips were also added to each plate and filled with PBST/SB (PBS+0.1% Tween 20+10% SuperBlock) buffer at 100 µL/well. Plates were incubated for 1 hour at room temperature and then washed 5 times with PBST. For detection of ADAs, HRP-conjugated Goat anti-feline IgG F(ab')<sub>2</sub> (anti-feline IgG F(ab')<sub>2</sub> reagent was cross-reacted to canine antibodies; Jackson ImmunoResearch Laboratories, West Grove PA), which was diluted in PBST/SB to 1:10000 and added at 100 µL/well to both sample and standard wells and incubated for 45 minutes at room temperature in the dark. Plates were washed 5 times with PBST and then one time with deionized water and then developed by adding 100 µL/well TMB substrate (Invitrogen, ThermoFisher Scientific, Waltham MA) for 15-20 minutes at room temperature in the dark. Color development was then stopped by addition of 100 µL/well of ELISA Stop Solution (Boston Bioproducts) and the absorbance was read at 450 nm using a SpectraMax plate reader within 30 minutes. The anti-drug antibody concentration was determined by interpolating the OD values in the 4-PL pseudo-standard curve using SoftMax Pro Software (Molecular Devices, San Jose CA).

[0201] To demonstrate the specificity of the detected ADAs, an "inhibition" assay was carried out. In the drug inhibition ADA assay, serum samples were diluted 1:100 in PBST/SB/20%HS buffer and mixed with an equal volume of 300 µg/mL of the relevant therapeutic compound (final sample dilution at 1:200 and final inhibitory compound at 150 µg/mL) and were incubated for 30-40 minutes at room temperature to allow anti-drug antibodies to bind the free inhibitor (i.e., the therapeutic compound). After preincubation, the samples were added to insulin-Fc fusion protein coated (or RHI coated) strips at 100 µL/well in duplicate. Samples diluted 1:200 in PBST/SB/20%HS buffer without the inhibitory compound were also tested in the sample plates along with duplicate strips of canine IgG coated standards. Remaining steps of the assay procedure were carried out as described above. The ADAs measured in the drug-inhibited wells were matched with the non-inhibited ADA concentrations to assess the specificity of the ADAs. If significant inhibition of ADA signals was observed in the drug-inhibited wells, this meant the ADAs were specific to the therapeutic compound.

Example 25: Generalized Procedure for Determination of In vivo Pharmacodynamics (PD) After a Single Administration of Human Insulin Fc-Fusion Proteins in Mice.

[0202] Insulin-Fc fusion proteins were assessed for their effects on fasting blood glucose levels as follows. Data was collected from N=3 balb/c mice or diabetic wt NOD mice (Jackson Laboratories) per group. The animals were fasted one hour before the experiment and then at time = 0 hours, the mice received a single subcutaneous administration of a pharmaceutical composition containing an insulin Fc-fusion protein homodimer at a concentration of 300 µg/kg of insulin-Fc fusion protein in a solution of between 10-50 mM sodium hydrogen phosphate, 50-150 mM sodium chloride, 0.005-0.05% v/v Tween-80, and optionally a bacteriostat (e.g. phenol, m-cresol, or methylparaben) at a concentration of between 0.02-1.00 mg/mL, at a final solution pH of between 7.0-8.0 adjusted using sodium hydroxide and/or hydrochloric acid.

[0203] For each time point, a sample of blood was collected, and a glucose level reading was immediately determined using a glucose meter (AlphaTRAK® 2 pet glucometer), which requires approximately one drop of blood. Average % fasting blood glucose levels (% FBGL) from 0 to 9 hours were plotted to assess the bioactivity of a given insulin-Fc fusion protein.

Example 26: Generalized Procedure for Determination of In vivo Pharmacodynamics (PD) After a Single Administration of Human Insulin Fc-Fusion Proteins in Mice.

[0204] Insulin-Fc fusion proteins are assessed for their effects on fasting blood glucose levels as follows. Data is collected from N=3 balb/c mice or diabetic wt NOD mice (Jackson Laboratories) per group. The animals are fasted one hour before the experiment and then at time = 0 hours, the mice receive a single subcutaneous administration of a pharmaceutical composition containing an insulin Fc-fusion protein homodimer at a concentration of 300 µg/kg of insulin-Fc fusion protein in a solution of between 10-50 mM sodium hydrogen phosphate, 50-150 mM sodium chloride, 0.005-0.05% v/v Tween-80, and optionally a bacteriostat (e.g. phenol, m-cresol, or methylparaben) at a concentration of between 0.02-1.00 mg/mL, at a final solution pH of between 7.0-8.0 adjusted using sodium hydroxide and/or hydrochloric acid.

[0205] For each time point, a sample of blood is collected, and a glucose level reading is immediately determined using a glucose meter (AlphaTRAK® 2 pet glucometer), which requires approximately one drop of blood. Average % fasting blood glucose levels (% FBGL) from 0 to 9 hours are plotted to assess the bioactivity of a given insulin-Fc fusion protein.

Example 27: Assay Procedure for Immunogenic Epitope Identification.

[0206] Maxisorp ELISA microplates (Nunc) are coated with a library of insulin-Fc fusion protein homodimer compounds with known amino acid sequences, and the coated plates are blocked in a similar manner as described in the anti-drug antibody ELISA assay of Example 25, except that each compound in the library is coated on a separate individual strip of ELISA microplate wells. The compounds in the library comprise a range of insulin-Fc fusion proteins with different insulin polypeptide amino acid compositions, including various B-chain, C-chain, and A-chain amino acid mutations, different linker compositions, and different Fc fragment compositions, including some of human origin. Separately, some plate strip wells are directly coated with 1:2 serial dilutions of canine IgG (Jackson ImmunoResearch Laboratories, West Grove PA) for calculating the anti-drug antibodies (ADA) in canine IgG units, respectively, as described in Example 24.

[0207] Serum obtained from individual dogs receiving repeated doses of an insulin-Fc fusion protein is first screened on the anti-drug antibody ELISA assay of Example 24. Serum samples demonstrating moderate or high positivity (e.g. moderate or high titers of antibodies) on the assay of Example 24 are serially diluted (1:200 to 1:8000) in PBST/SB/20%HS sample dilution buffer (PBS+0.1% Tween 20+10% SuperBlock+20% horse serum) and added to the plates coated with the library of insulin-Fc fusion protein compounds for 1 hour at RT. Following incubation, the plates are washed 5 times with PBST. For detection of canine antibodies capable of cross-reacting to the coated compound library, HRP conjugated goat anti-feline IgG F(ab')<sub>2</sub> (Jackson ImmunoResearch Laboratories, West Grove PA), which is cross-reactive to canine IgGs, is diluted in PBST/SB to 1:10000 and added at 100  $\mu$ L/well to both sample and standard wells and incubated for 45 min at RT in the dark. Plates are washed 5 times with PBST, once with deionized water, and developed by the adding 100  $\mu$ L/well TMB substrate (Invitrogen, ThermoFisher Scientific, Waltham MA) for 15-20 min at RT in the dark. Color development is then stopped by addition of 100  $\mu$ L/well of ELISA Stop Solution (Boston Bioproducts, Ashland MA) and absorbance is read at 450 nm using a SpectraMax plate reader within 30 min. Anti-compound cross-reactive antibody concentrations present in the serum samples are determined by interpolating the OD values in the 4-PL pseudo-standard curve against the directly coated canine IgG antibody controls using SoftMax Pro Software (Molecular Devices, San Jose CA).

[0208] By correlating the resulting antibody concentrations from the assay with the known amino acid compositions of the coated insulin-Fc fusion protein library, one can determine whether particular amino acid mutations or epitopes are responsible for causing none, some, most, or all of the total antibody signal on the assay, indicating no binding, weak binding, or strong binding to various insulin-Fc fusion protein homodimers. The mutations or epitopes responsible for moderate or strong binding are herein referred to as immunogenic "hot spots".

Example 28: Design Process for Obtaining Canine Insulin-Fc fusion Proteins with High Homodimer Titers and Acceptable Levels of Acute and Repeated Dose Bioactivity.

[0209] The process for meeting the design goals described in the Detailed Description of the Invention comprised the following steps. First, the insulin polypeptide of SEQ ID NO: 4 or SEQ ID NO: 5 was combined with a species-specific Fc fragment of a particular IgG isotype and a linker such that the resulting insulin-Fc fusion protein was most likely to yield a long acting bioactivity product with minimal immunogenicity (e.g., a species-specific IgG isotype was chosen with minimal Fc(gamma)receptor I binding). The DNA sequence coding for the desired fusion protein was prepared, cloned into a vector (LakePharma, San Carlos, CA), and the vector was then used to transiently transfect HEK cells according to the procedure described in Example 1. The insulin-Fc fusion protein was then purified according to Example 4 and the overall protein yield and %homodimer measured according to Example 10. Only candidates with a homodimer titer of greater than 50 mg/L were considered acceptable, because titers less than this level are not likely to result in commercial production titers that meet the stringently low manufacturing cost requirements for veterinary products. Selected insulin-Fc fusion proteins were then screened for indicators of bioactivity through in vitro insulin receptor binding studies as described in Example 12. Based on experience, only compounds that exhibited IR activity IC<sub>50</sub> values less than 5000 nM were deemed likely to exhibit bioactivity in the target species. Although the in vitro IR IC<sub>50</sub> value is a useful qualitative screening tool, it utilizes human IM-9 cells which express the human insulin receptor and therefore it may not capture some of the small differences in affinity between the canine IR and the human IR. Furthermore, factors other than insulin receptor binding may influence a compound's bioactivity in vivo (e.g., affinity for canine FcRn to allow for extended pharmacokinetic elimination half-lives in vivo). Therefore, selected insulin-Fc fusion proteins that were acceptable from a manufacturing and IR activity IC<sub>50</sub> value standpoint were further screened for bioactivity dogs to screen out any materials with less than the desired potency and/or duration of bioactivity (e.g., NAOC of less than 150 %FB-GL.days.kg/mg). Again, based on experience, at NAOC values of greater than 150 %FBGL-days-kg/mg, the dose requirements in the target species will be sufficiently low so as to reach an acceptable treatment cost. Lastly, an additional evaluation criterion was added which is mentioned rarely if ever in the art. As discussed in more detail in the Examples below, many insulin-Fc fusion protein embodiments that exhibit acceptable NAOC levels in the target species after the first dose, unexpectedly fail to maintain that level of bioactivity after repeated doses. Furthermore, in most cases the

reduction in repeated dose bioactivity in the target species is correlated with the development of neutralizing anti-drug antibodies. This propensity to generate anti-drug antibodies and the failure to maintain activity render such insulin-Fc fusion proteins impractical for use in treating a chronic disease such as canine diabetes. Therefore, only the insulin-Fc fusions proteins exhibiting acceptable levels of repeated dose bioactivity (e.g., NAOCR values greater than 0.5 for the third dose relative to the first dose) with minimal levels of anti-drug antibodies were deemed acceptable for use in the present invention.

Example 29: Design Process for Obtaining Human Insulin-Fc fusion Proteins with High Homodimer Titers and Acceptable Levels of Acute and Repeated Dose Bioactivity.

**[0210]** The process for meeting the design goals described in the Detailed Description of the invention comprised the following steps. First, the insulin polypeptide of SEQ ID NO: 7 or SEQ ID NO: 10 was combined with a human Fc fragment of a particular IgG isotype (IgG2 or IgG1) and a linker such that the resulting insulin-Fc fusion protein was most likely to yield a long acting bioactivity product. The DNA sequence coding for the desired fusion protein was prepared, cloned into a vector (LakePharma, San Carlos, CA), and the vector was then used to transiently transfect HEK cells according to the procedure described in Example 1. The insulin-Fc fusion protein was then purified according to Example 4 and the total protein yield and %homodimer was measured according to Example 10 and the homodimer titer calculated. Only candidates with a homodimer titer of greater than 150 mg/L were considered acceptable, because homodimer titers less than this level are not likely to translate to high homodimer titer CHO stably infected cell lines, and therefore not likely to result in commercial production titers that meet the low manufacturing cost requirements for the relatively commoditized human insulin market. Selected insulin-Fc fusion protein configurations were then screened for indicators of bioactivity through in vitro IR binding studies as described in Example 12. Based on experience, only compounds that exhibited IR activity IC<sub>50</sub> values less than 2400 nM and more preferably less than 2000 nM were deemed likely to exhibit bioactivity in vivo. The in vitro IR IC<sub>50</sub> value is a useful qualitative screening tool, as the assay utilizes human IM-9 cells which express the human IR, and the greater binding (lower IR IC<sub>50</sub> values) would be expected to give greater in vivo potency for a given dose compared to compounds with decreased binding (higher IR IC<sub>50</sub> values). Furthermore, factors other than IR binding may influence a compound's bioactivity in vivo. For example, the affinity of Fc fusion proteins for the human FcRn receptor is related to a compound's in vivo pharmacokinetic elimination half-life. Extended in vivo half-life of insulin-Fc fusion proteins of several days or more correlates well with EC<sub>50</sub> values less than 1500 ng/mL and more preferably less than 1000 ng/mL, as measured by an in vitro human FcRn binding assay (Example 19).

**[0211]** Selected insulin-Fc fusion protein configurations that were acceptable in the areas of manufacturing, IR activity IC<sub>50</sub> value, and human FcRn activity EC<sub>50</sub> value were further screened for their potential for in vivo immunogenicity by testing the human Fc(gamma)RI receptor and human C1q binding affinities. Molecules that bind more strongly to these immune system components are more likely to go through increased antigen cell presentation (APC) and exhibit a greater immunogenicity profile as measured by anti-drug antibodies. Anti-drug antibodies are unwanted and may hinder the in vivo pharmacokinetic half-life of the molecule, may neutralize the activity of the drug, or may perform both functions. The impact of potential anti-drug antibodies on insulin-Fc in vivo performance could be quite problematic. Therefore, work was performed to screen candidate insulin-Fc fusion proteins against Fc(gamma)RI (Example 14) and C1q (Example 16) to mitigate the risk of unwanted antigen presentation and potential anti-drug antibody-inducing immunogenicity (Guilliams, Martin & Bruhns, Pierre & Saeys, Yvan & Hammad, Hamida & Lambrecht, Bart. (2014). The function of Fc gamma receptors in dendritic cells and macrophages. Nature reviews. Immunology. 14. 10.1038/nri3582). The design goal established for human Fc(gamma)RI binding (where the biotinylated-Fc(gamma)RI concentration of the insulin-Fc fusion protein under test is 3000 ng/mL) is an OD<sub>450</sub> Ratio < 0.50 (with respect to the insulin-Fc fusion protein configuration of SEQ ID NO: 76) and the design goal established for C1q binding (where the biotinylated-C1q concentration of the insulin-Fc fusion protein under test is 1000 ng/mL) is an OD<sub>450</sub> ratio < 0.35 (with respect to the insulin-Fc fusion protein configuration of SEQ ID NO: 76).

**[0212]** As described elsewhere, human IgG1 Fc containing insulin-Fc fusion protein configurations were unexpectedly found to be much higher yielding than the human IgG2 Fc containing configurations and were therefore preferred with respect to manufacturability. However, human IgG1 Fc containing insulin-Fc fusion protein configurations, when containing natural N-linked glycosylation at the cNg site often display very high binding to both Fc(gamma)RI and C1q, and strong binding to one or both of these moieties has been correlated to a high potential for enhanced antigen presentation and immunogenicity in vivo (Kouser, L., Madhukaran, S. P., Shastri, A., Saraon, A., Ferluga, J., Al-Mozaini, M., & Kishore, U. (2015). Emerging and Novel Functions of Complement Protein C1q. Frontiers in Immunology, 6, 317. doi:10.3389/fimmu.2015.00317). Therefore, several variants of hIgG1 isotype, Fc-containing insulin-Fc fusion protein configurations were tested by first screening compounds whose native glycosylation site (cNg) was mutated to prevent natural glycosylation during biosynthesis. These molecules were then further mutated to improve their properties by manipulating the insulin polypeptide composition. Lastly, several linker variants of different composition and length were examined to identify whether various design properties could be further optimized. After screening through multiple rounds of

optimization, only the insulin-Fc fusions protein configurations exhibiting acceptable levels of homodimer titer, IR binding, FcRn binding, Fc(gamma)RI binding, and C1q binding were deemed acceptable for use in the present invention.

#### **RESULTS - INSULIN-FC FUSION PROTEINS COMPRISING A CANINE Fc FRAGMENT** *Comparative Example 30:*

##### Canine Insulin-Fc Fusion Protein Comprising the Canine Fc IgGA Isotype.

**[0213]** An attempt was made to produce an insulin-Fc fusion protein comprising the insulin polypeptide sequence of SEQ ID NO: 4 and the Fc fragment of the canine IgGA isotype (SEQ ID NO: 14) using the peptide linker of SEQ ID NO: 11. The full amino acid sequence for the resulting insulin-Fc fusion protein is as follows:

FVNQHLCGSDLEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
GRCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPEVTCVVLDLGREDPEVQISWFDGKEVHTA  
KTQSREQQFNGTYRVVSVLPIDHQDWLTGKEFKCRVNHIDLPSPERTISKARGRAHKPSVYVLPPS  
PKELSSSDTVSITCLIKDFYPPDIDVEWQSNGQQEPPERKHRMTPPQLDEDGSYFLYSKLSVDKSRW  
QQGDPFTCAVMHETLQNHYTDLSSLHSPG (SEQ ID NO: 31).

**[0214]** The insulin-Fc fusion protein of SEQ ID NO: 31 was synthesized in HEK cells according to Example 1 and purified according to Example 4. The protein yield was 22 mg/L after the Protein A purification step. The structure of the insulin-Fc fusion protein was confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequence was further identified by LC-MS with glycan removal according to Example 8. The %homodimer was measured by size-exclusion chromatography according to Example 10 and determined to be 24%, indicating a high degree of homodimer aggregates. The resulting homodimer titer was therefore only 5 mg/L. In summary, manufacturing of the insulin-Fc fusion protein of SEQ ID NO: 31 in HEK cells resulted in a high level of aggregates and a low homodimer titer (5 mg/L), which did not meet the design goal of a homodimer titer of greater than 50 mg/L.

**[0215]** Nevertheless, the insulin-Fc fusion protein of SEQ ID NO: 31 as evaluated for bioactivity. First, the insulin receptor binding of the insulin-Fc fusion protein of SEQ ID NO: 31 was measured according to Example 12, resulting in an IC<sub>50</sub> value of 2,733 nM indicating that the compound is likely to be bioactive in vivo (i.e. IC<sub>50</sub> less than 5000 nM).

**[0216]** Next, the in vivo pharmacodynamics (PD) of the insulin-Fc fusion protein of SEQ ID NO: 31 was measured after a single intravenous administration of the compound to N=3 canines, according to Example 21. FIG. 2 shows the percent fasting blood glucose level of SEQ NO: 31 as a function of time. The NAOC for SEQ ID NO: 31 was calculated to be 105 %FBGL-days-kg/mg according to the procedure of Example 22. The in vivo half-life of SEQ ID NO: 31 was calculated to be less than one day using the method of Example 23. The relatively low NAOC was likely the result of the high amount of aggregates in the sample (i.e., low %homodimer), but what soluble homodimer remained in circulation still only had a pharmacokinetic elimination half-life of less than one day which was deemed unlikely support of once-weekly administration.

##### Comparative Example 31: Mutations of the Fc Fragment Region of Insulin-Fc Fusion Proteins Comprising the Canine IgGA Isotype.

**[0217]** In an attempt to increase the %homodimer content, improve the bioactivity, and increase the half-life of the insulin-Fc fusion protein of SEQ ID NO: 31, mutations were inserted into the Fc fragment CH3 region to try to prevent intermolecular association (e.g., Fc fragment-Fc fragment interactions between molecules) and encourage stronger binding to the FcRn receptor (e.g., higher affinity for the FcRn) to increase recycling and systemic circulation time. The following insulin-Fc fusion proteins were synthesized in HEK cells according to Example 1, purified according to Example 4, and tested according to Example 6, Example 8 and Example 10, the test results of which are shown below in Table 2. The sequence alignment of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, and SEQ ID NO: 35 against SEQ ID NO: 31 and the differences in amino acid sequences are shown in Fig. 3.

FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
 GRCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPEVTCVVLDLGREDPEVQISWFVDGKEVHTA  
 5 KTQSREQQFNGTYRVVSVLPIEHQDWLTGKEFKCRVNHIDLPSPIERTISKARGRAHKPSVYVLPPS  
 PKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQEPPERKHRMTPPQLDEDGSYFLYSKLSVDKSRW  
 QQGDPFTCAVLHEALHSHYTQKSLSLSPG (SEQ ID NO: 32)

10 FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
 GRCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPEVTCVVLDLGREDPEVQISWFVDGKEVHTA  
 KTQSREQQFNGTYRVVSVLPIEHQDWLTGKEFKCRVNHIDLPSPIERTISKARGRAHKPSVYVLPPS  
 15 PKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQEPPERKHRMTPPQLDEDGSYFLYSKLSVDKSRW  
 QQGDPFTCAVLHETLQSHYTDLSLHSPG (SEQ ID NO: 33)

20 FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
 GRCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPEVTCVVLDLGREDPEVQISWFVDGKEVHTA  
 KTQSREQQFNGTYRVVSVLPIEHQDWLTGKEFKCRVNHIDLPSPIERTISKARGRAHKPSVYVLPPS  
 25 PKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQEPPERKHRMTPPQLDEDGSYFLYSKLSVDKSRW  
 QQGDPFTCAVMHETLQSHYTDLSLHSPG (SEQ ID NO: 34)

30 FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
 GRCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPEVTCVVLDLGREDPEVQISWFVDGKEVHTA  
 KTQSREQQFNGTYRVVSVLPIEHQDWLTGKEFKCRVNHIDLPSPIERTISKARGRAHKPSVYVLPPS  
 PKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQEPPERKHRMTPPQLDEDGSYFLYSKLSVDKSRW  
 35 QQGDPFTCAVLHETLQNHYTDLSLHSPG (SEQ ID NO: 35)

[0218] The insulin-Fc fusion proteins based on canine IgGA variants are listed in Table 2 along with the corresponding protein yields, %homodimer, and homodimer titers. The results show that the various mutations to the IgGA Fc fragment, instead of improving the %homodimer and homodimer titer, gave rise to highly aggregated proteins with extremely low homodimer titers of less than 5 mg/L. As such, the in vivo bioactivity and pharmacokinetics of the compounds could not be evaluated.

**Table 2: Homodimer titers for sequences utilizing a native or mutated canine IgGA Fc fragment CH3 region**

| SEQ ID NO:    | Protein Yield (mg/L) | %Homodimer | Homodimer Titer (mg/L) |
|---------------|----------------------|------------|------------------------|
| SEQ ID NO: 31 | 22                   | 24%        | 5                      |
| SEQ ID NO: 32 | 33                   | 0%         | 0                      |
| SEQ ID NO: 33 | 57                   | 0%         | 0                      |
| SEQ ID NO: 34 | 67                   | 0%         | 0                      |
| SEQ ID NO: 35 | 80                   | 0%         | 0                      |

*Comparative Example 32: Canine Insulin-Fc Fusion Protein Using Other Canine Fc Fragment Isotypes.*

[0219] As described above, canine IgGA is thought to be the preferred isotype for the Fc fragment to produce non-immunogenic insulin-Fc fusion protein for dogs due to its lack of Fc(gamma) I effector function in canines (much like the

human IgG2 isotype in humans). However, insulin-Fc fusion proteins manufactured with a canine IgGA Fc fragment were highly aggregated with an unacceptably low homodimer titer and unacceptably low levels of bioactivity and duration of action. Therefore, Fc fragments from the other canine IgG isotypes (canine IgGB of SEQ ID NO: 15), canine IgGC of SEQ ID NO: 16, and canine IgGD of SEQ ID NO: 17) were evaluated as replacements for the canine IgGA Fc fragment of the insulin-Fc fusion of SEQ ID NO: 31. The three insulin-Fc fusion proteins containing Fc fragments based on the canine IgGB, IgGC, and IgGD isotypes were synthesized using the same insulin polypeptide of SEQ ID NO: 4 and peptide linker of SEQ ID NO: 11 as were used to make the insulin-Fc fusion protein of SEQ ID NO: 31. The proteins were manufactured in HEK293 cells according to Example 1. The insulin-Fc fusion proteins were then purified using a Protein A column according to Example 4. The structures of the insulin-Fc fusion proteins were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. The %homodimer was measured by size-exclusion chromatography according to Example 10. Their sequences are shown below and their sequence alignment comparison against SEQ ID NO: 31 is shown in FIG. 4:

FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
GDCPKCPAPEMLGGPSVFIFPPKPKDILLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQMQTAKT  
QPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPSPR  
EELSKNTVSLTCLIKDFFPPDIDVEWQSNQGQEPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQR  
GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 36)

FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
GCNNCPGCGLLGGPSVFIFPPKPKDILVTARTPTVTCVVVDLDPENPEVQISWFVDSKQVQTAN  
TQPREEQSNGTYRVVSVLPIGHQDWLSGKQFKCKVNNKALPSPIEEIISKTPGQAHQPNVYVLPSPS  
RDEMSKNTVTLTCLVKDFFPPEIDVEWQSNQGQEPESKYRMTTPQLDEDGSYFLYSKLSVDKSRW  
QRGDTFICAVMHEALHNHYTQISLSHSPG (SEQ ID NO: 37)

FVNQHLCGSDLVEALALVCGERGGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
GCISPCPVPESLGGPSVFIFPPKPKDILRITRTEITCVVLDLGREDPEVQISWFVDGKEVHTAKTQPR  
EQQFNSTYRVVSVLPIDHQLDWTGKEFKCRVNHGLPSPIERTISKARGQAHQPSVYVLPSPKELS  
SSDVTVTLTCLIKDFFPPEIDVEWQSNQGQEPESKYHTTAPQLDEDGSYFLYSKLSVDKSRWQQGDT  
FTCAVMHEALQNHYTDLSLSHSPG (SEQ ID NO: 38)

**[0220]** The resulting protein yields, %homodimer, and homodimer titers are given in Table 3. Unexpectedly, only the insulin-Fc fusion protein of SEQ ID NO: 36 comprising an Fc fragment based on the canine IgGB isotype demonstrated a homodimer titer which met the design criteria of greater than 50 mg/L. The insulin-Fc fusion protein of SEQ ID NO: 37 comprising an Fc fragment based on the canine IgGC isotype did not yield any compound at all, and the insulin-Fc fusion protein of SEQ ID NO: 38 comprising an Fc fragment based on the canine IgGD isotype demonstrated an appreciable protein yield but with a high degree of aggregation and therefore an unacceptably low homodimer titer.

**[0221]** In vitro insulin receptor binding for the insulin-Fc fusion proteins of SEQ ID NO: 36 and SEQ ID NO: 38 was tested according to the procedure of Example 12. The insulin-Fc fusion protein of SEQ ID NO: 38 demonstrated an IC<sub>50</sub> of greater than 5000 nM, indicating that the compound was highly unlikely to show bioactivity in vivo. However, the insulin-Fc fusion protein of SEQ ID NO: 36 demonstrated an IC<sub>50</sub> of 28 nM indicating that this sequence was likely to be bioactive in vivo.

**Table 3: Homodimer titers for sequences utilizing native canine IgGB, IgGC, and IgGD Fc fragments**

| SEQ ID NO:             | IgG Fragment | Protein Yield (mg/L) | %Homodimer | Homodimer Titer (mg/L) | IR Binding, IC50 (nM) |
|------------------------|--------------|----------------------|------------|------------------------|-----------------------|
| SEQ ID NO: 31          | IgGA         | 21                   | 24%        | 5                      | 2,733                 |
| SEQ ID NO: 36          | IgGB         | 80                   | 93%        | 74                     | 28                    |
| SEQ ID NO: 37          | IgGC         | 0                    | 0%         | 0                      | DNM*                  |
| SEQ ID NO: 38          | IgGD         | 134                  | 12%        | 16                     | >5000                 |
| *DNM = Did Not Measure |              |                      |            |                        |                       |

**Comparative Example 33: In vivo Efficacy of an Insulin-Fc fusion Protein Comprising the Insulin Polypeptide of SEQ ID NO: 4 with a Canine IgGB Isotype Fc Fragment.**

[0222] Given the promising homodimer titer and insulin receptor activity results in Example 32, the insulin-Fc fusion protein of SEQ ID NO: 36 was tested for in vivo bioactivity according to Example 21 following an intravenous injection in each of N=3 healthy, antibody-naïve, beagle dogs weighing approximately 10 kg. In a separate experiment, the compound was administered subcutaneously to N=3 naïve beagle dogs. FIG. 5 shows the %FBGL versus time for a single intravenous administration of the insulin-Fc fusion protein of SEQ ID NO: 36, and FIG. 6 shows the %FBGL vs. time for a single subcutaneous administration of the insulin-Fc fusion protein of SEQ ID NO: 36, both of which demonstrate that the insulin-Fc fusion protein of SEQ ID NO: 36 is significantly bioactive in dogs.

[0223] The NAOC was calculated according to the procedure of Example 22 to determine the relative bioactivity and duration of action of the insulin-Fc fusion protein. The NAOC of the insulin-Fc fusion protein of SEQ ID NO: 36 injected intravenously was 399 %FBGL·days·kg/mg which was 3.8 times the NAOC of the insulin-Fc fusion protein of SEQ ID NO: 31 injected intravenously, illustrating significantly increased bioactivity for the insulin-Fc fusion protein comprising the canine IgGB Fc fragment versus the insulin-Fc fusion protein comprising the canine IgGA Fc fragment. The NAOC of the insulin-Fc fusion protein of SEQ ID NO: 36 injected subcutaneously was 366 %FBGL·days·kg/mg, demonstrating a level of bioactivity via subcutaneous administration that is similar to that obtained via intravenous administration.

**Example 34: In Vivo Immunogenicity Screening After Repeated Subcutaneous Doses of the Insulin-Fc Fusion Protein Comprising the Insulin Polypeptide of SEQ ID NO: 4 with a Canine IgGB Isotype Fc Fragment.**

[0224] The repeated dose subcutaneous bioactivity of the insulin-Fc fusion protein of SEQ ID NO: 36 was tested in dogs as per the method described in Example 22. N=3 animals were dosed subcutaneously at day 0, at day 35, and at day 42, and the %FBGL was measured for the 7-day window after each dose according to Example 22. The NAOC and NAOCR were calculated according to the procedure of Example 22 for each repeated subcutaneous injection. As illustrated in Table 4, repeated subcutaneous dosing in dogs unexpectedly revealed a significant decay in bioactivity by the third dose as measured by a significant decrease in the NAOCR (i.e., the NAOC for the third injection was only 0.40, or 40%, of the NAOC for the first injection).

**Table 4: NAOC per dose and NAOCR for repeated doses of SEQ ID NO: 36**

| Injection Number of SEQ ID NO: 36 | NAOC (%FBGL·days·kg/mg) | NAOCR (ratioed to Week 1) |
|-----------------------------------|-------------------------|---------------------------|
| 1                                 | 330                     | 1.0                       |
| 2                                 | 339                     | 1.1                       |
| 3                                 | 115                     | 0.4                       |

[0225] Without being bound to any particular explanation it was postulated that the cause of the significant reduction in bioactivity of the insulin-Fc fusion protein of SEQ ID NO: 36 after the third repeated subcutaneous dose in dogs was due to the development of anti-drug antibodies that neutralized its biological activity. Anti-drug antibodies may be directed

against the insulin polypeptide, linker, or Fc-fragment portions of an insulin-Fc fusion protein. The immunogenic response manifests as interactions between antigen presenting cells, T-helper cells, B-cells, and their associated cytokines, which may lead to the production of endogenous antibodies against the drug (e.g. anti-drug antibodies). Binding antibodies are all isotypes capable of binding the insulin-Fc fusion protein, and these may be detected in an immunoassay as described in Example 24. Neutralizing antibodies that inhibit functional activity of the insulin-Fc fusion protein are generally directed against an epitope that is required for bioactivity. To assess whether this was the case, serum that was collected prior to the administration of each dose and at the end of the experiment described in Example 11 and Example 12 was tested to quantify the levels of anti-drug antibodies according to Example 24. As shown in FIG. 7, levels of anti-drug antibodies did indeed increase with multiple subcutaneous administrations of the compound, indicating that the generation of neutralizing anti-drug antibodies were the likely cause for the reduction in the NAOCR after the third injection of the insulin Fc-fusion protein of SEQ ID NO: 36.

Comparative Example 35: Non-Glycosylated Insulin-Fc Fusion Protein Comprising the Insulin Polypeptide of SEQ ID NO: 4 with Canine IgGB Isotype Fc Fragments to Reduce the Potential Risk of Immunogenicity.

[0226] As shown in Example 32 and Example 33, the insulin-Fc fusion protein of SEQ ID NO: 36 showed acceptable %homodimer content, homodimer titer, and bioactivity in dogs; however, its use for a chronic disease such as diabetes is compromised by the reduction in bioactivity (Example 34) and generation of anti-drug antibodies (Example 34) with repeated subcutaneous dosing. Without being bound to any particular theory, one possible cause of the generation of anti-drug antibodies and the reduction in bioactivity is the increased interaction of the canine IgGB Fc fragment with various receptors of the canine immune system (e.g. Fc(gamma) receptors, e.g. Fc(gamma)RI). Nevertheless, the canine IgGB isotype was the only one of the four canine IgG isotypes that, when used for the Fc fragment, resulted in an insulin-Fc fusion protein meeting the manufacturability and single-dose bioactivity design goals (Example 28). As described in the Detailed Description of the Invention, one method for reducing the Fc(gamma) interaction involves mutating the Fc fragment cNg site to prevent glycosylation during synthesis in the host cell. Therefore, cNg site mutations were made to the Fc fragment region of SEQ ID NO: 36 to reduce the binding affinity of the Fc fragment for Fc(gamma) receptors in vivo, as measured by binding in an in vitro human Fc(gamma)RI assay described in Example 14. Verification of the lack of glycan were performed using the LC-MS method of Example 8, but with omission of the PNGase F treatment step. The position of the cNg site in the insulin-Fc fusion protein of SEQ ID NO: 36 is cNg-NB139. Mutations to SEQ ID NO: 36 included SEQ ID NO: 39 comprising a mutation of cNg-NB139-Q, SEQ ID NO: 40 comprising a mutation of cNg-NB139-S, SEQ ID NO: 41 comprising a mutation of cNg-NB139-D, and SEQ ID NO: 42 comprising a mutation of cNg-NB139-K. The full amino acid sequences of the cNg-mutated insulin-Fc fusion proteins are listed below (with the NB139 position underlined) and the resulting sequence alignments are shown in FIG. 8 (Clustal Omega):

FVNQHLCGSDLEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
QPREEQFQGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSR  
EELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQR  
GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 39)

FVNQHLCGSDLEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
QPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSR  
EELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQR  
GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 40)

FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
 GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
 QPREEQFDGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPSSR  
 EELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQR  
 GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 41)

FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGGGAGGG  
 GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
 QPREEQFKGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPSSR  
 EELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQR  
 GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 42).

**[0227]** The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. The structures of the insulin-Fc fusion proteins were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. The %homodimer was measured by size-exclusion chromatography according to Example 10. As shown in Table 5, the homodimer titers of the insulin-Fc fusion proteins of SEQ ID NO: 40, SEQ ID NO: 41, and SEQ ID NO: 42 meet the design goal, while unexpectedly the insulin-Fc fusion protein of SEQ ID NO: 39 containing the cNg- NB 139-Q mutation did not meet the design goal for homodimer titer.

**Table 5: Homodimer titers for cNg variations of SEQ ID NO: 36**

| SEQ ID NO:    | cNg Mutation | Protein Yield (mg/L) | %Homodimer | Homodimer Titer (mg/L) |
|---------------|--------------|----------------------|------------|------------------------|
| SEQ ID NO: 39 | cNg-Q        | 37                   | 98%        | 36                     |
| SEQ ID NO: 40 | cNg-S        | 77                   | 98%        | 75                     |
| SEQ ID NO: 41 | cNg-D        | 88                   | 98%        | 86                     |
| SEQ ID NO: 42 | cNg-K        | 68                   | 98%        | 67                     |

**[0228]** To determine which of the remaining three compounds was most likely to exhibit reduced immunogenicity, the Fc(gamma) receptor binding was measured according to the procedure of Example 15. Low Fc(gamma) receptor binding is most likely to correlate with minimum immunogenicity. Table 6 compares the Fc(gamma) receptor binding of these insulin-Fc fusion proteins with the Fc(gamma) receptor binding of the insulin-Fc fusion protein of SEQ ID NO: 36 demonstrating unexpectedly that the insulin-Fc fusion protein of SEQ ID NO: 41, containing the cNg-D mutation, exhibits an Fc (gamma) receptor binding activity that is approximately twice that of the insulin-Fc fusion proteins of SEQ ID NO: 40, containing the cNg-S mutation and SEQ ID NO: 42 containing the cNg-K mutation. Therefore, only the insulin-Fc fusion proteins comprising the latter two compounds containing the cNg-S mutation and the cNg-K mutations were deemed suitable for repeated dose bioactivity testing in dogs.

**Table 6: Fc(gamma) receptor binding for cNg variations of SEQ ID NO: 36**

| SEQ ID NO:    | cNg Mutation | OD450nm Log[Fc (gamma) RI] (ng/mL) | OD450nm Minus Assay Background | Ratio to SEQ ID NO: 36 |
|---------------|--------------|------------------------------------|--------------------------------|------------------------|
| SEQ ID NO: 36 | Native cNg   | 0.386                              | 0.323                          | 1.00                   |
| SEQ ID NO: 40 | cNg-S        | 0.140                              | 0.077                          | 0.24                   |
| SEQ ID NO: 41 | cNg-D        | 0.204                              | 0.141                          | 0.44                   |
| SEQ ID NO: 42 | cNg-K        | 0.126                              | 0.063                          | 0.20                   |

(continued)

**Table 6: Fc(gamma) receptor binding for cNg variations of SEQ ID NO: 36**

| SEQ ID NO:                     | cNg Mutation | OD450nm Log[Fc (gamma) RI] (ng/mL) | OD450nm Minus Assay Background | Ratio to SEQ ID NO: 36 |
|--------------------------------|--------------|------------------------------------|--------------------------------|------------------------|
| Assay background (no compound) | N/A          | 0.063                              | 0.000                          | N/A                    |

*Comparative Example 36: Evaluation of In Vivo Bioactivity and Immunogenicity of an Insulin Polypeptide of SEQ ID NO: 4 with the Non-Glycosylated cNg-K and cNg-S Canine IgGB Isotype Fc Fragments.*

**[0229]** To determine if the insulin-Fc fusion protein of SEQ ID NO: 40, containing the cNg-S mutation, improved the repeated dose bioactivity performance in dogs, the compound was administered subcutaneously to N=1 dog on day 0, day 7, day 14, and on day 28 according to the procedure of Example 22. When the dog's %FBGL dropped too low, the dog was given food to raise the blood glucose to a safe level. The NAOC for the first injection was 191 %FBGL·days·kg/mg, showing that the insulin-Fc fusion protein of SEQ ID NO: 40 was satisfactorily bioactive in vivo. The NAOC and NAOCR were also measured for each subsequent dose according to the general procedure of Example 22, calculated from the time the dose was administered until just before the next dose was administered. The NAOC and the NAOCR shown in Table 7 illustrate that the insulin-Fc fusion protein of SEQ ID NO: 40 exhibited an NAOCR that decreased significantly on doses 3 and 4 of a four dose regimen. Therefore, the insulin-Fc fusion protein of SEQ ID NO: 40, containing the cNg-S mutation, was unable to demonstrate repeated dose bioactivity in dogs despite having a low Fc(gamma)RI binding four times lower than that of the insulin-Fc fusion protein of SEQ ID NO: 36.

**Table 7: NAOC per dose for repeated doses of SEQ ID NO: 40**

| Injection Number of SEQ ID NO: 40 | NAOC (%FBGL·days·kg/mg) | NAOCR |
|-----------------------------------|-------------------------|-------|
| 1                                 | 191                     | 1.0   |
| 2                                 | 240                     | 1.3   |
| 3                                 | 0                       | 0.0   |
| 4                                 | 39                      | 0.2   |

**[0230]** To determine if the insulin-Fc fusion protein of SEQ ID NO: 42, containing the cNg-K mutation, improved the repeated dose bioactivity performance in dogs, the compound was administered subcutaneously to N=1 dog on day 0, day 7, day 14, and on day 28 according to the procedure of Example 22. When the dog's %FBGL dropped too low, the dog was given food to raise the blood glucose to a safe level. The NAOC for the first injection was 449 %FBGL·days·kg/mg, showing that the insulin-Fc fusion protein of SEQ ID NO: 42 was satisfactorily bioactive in vivo. The pharmacokinetic profile of the compound was also measured by the method of Example 23 using ELISA, and a two-compartment model was fit to the data to determine its elimination half-life which was about 0.9 days. The NAOC and NAOCR were also measured for each subsequent dose according to the general procedure of Example 22, calculated from the time the dose was administered until just before the next dose was administered. The NAOC and the NAOCR shown in Table 8 illustrate that the insulin-Fc fusion protein of SEQ ID NO: 42 maintains an NAOCR greater than 0.6 throughout the four doses. Therefore, unexpectedly, the insulin-Fc fusion protein of SEQ ID NO: 42, containing the cNg-K mutation, was the only non-glycosylated mutant of the insulin-Fc fusion protein of SEQ ID NO: 36 resulting in significantly improved repeated dose bioactivity in dogs.

**Table 8: NAOC per dose for repeated doses of SEQ ID NO: 42**

| Injection Number of SEQ ID NO: 42 | NAOC (%FBGL·days·kg/mg) | NAOCR |
|-----------------------------------|-------------------------|-------|
| 1                                 | 449                     | 1.0   |
| 2                                 | 361                     | 0.8   |
| 3                                 | 259                     | 0.6   |
| 4                                 | 638                     | 1.4   |

**[0231]** The levels of anti-drug and anti-insulin antibodies were also measured throughout the course of treatment (28 days) and for an additional two weeks according to Example 24. FIG. 9 illustrates that the insulin-Fc fusion protein of SEQ ID NO: 42 still generated anti-drug antibodies with repeated subcutaneous dosing in dogs, but the anti-drug antibody titers were much lower than those generated by the insulin-Fc fusion protein of SEQ ID NO: 36 (Example 32).

*Comparative Example 37: Screening of Canine Serum Containing Anti-Drug Antibodies and Identification of Potential Immunogenic Epitopes at the B10D and A8H Positions of the Insulin Polypeptide.*

**[0232]** Mutating the cNg site of the canine IgGB Fc fragment to a Lys (i.e., cNg-K) did improve the repeated dose bioactivity of the insulin-fusion protein comprising the insulin polypeptide of SEQ ID NO: 4 and the peptide linker of SEQ ID NO: 11 (Example 36), but the resulting insulin-Fc fusion protein of SEQ ID NO: 42 still gave rise to anti-drug antibodies (Example 36). It was hypothesized, therefore, that the insulin polypeptide of SEQ ID NO: 4 may unexpectedly contain specific epitopes (i.e., immunogenic "hot spots") against which the dog's immune system is directed. Therefore, the binding specificity of the antibodies present in the serum samples described in Example 24 were evaluated according to the general procedure of Example 27. The analysis of the antibody-containing serum samples from the repeated dosing of the insulin-Fc fusion protein of SEQ ID NO: 36 (Example 32) against the coated insulin-Fc fusion protein library demonstrated that there were unexpectedly two primary "hot spots" present within the insulin polypeptide sequence of SEQ ID NO: 5: the aspartic acid mutation at the 10th position from the N-terminus of the B-chain (i.e., B 10), and, separately, the histidine mutation at the 8th position from the N-terminal end of the A-chain (i.e., A8). The results suggest that insulin-Fc fusion proteins comprising insulin polypeptide amino acid compositions containing these two particular amino acid mutations are likely to be immunogenic in dogs and therefore likely to give rise to anti-drug antibodies that neutralize the bioactivity after repeated injections. Therefore, it was determined that insulin polypeptides that do not contain the B 10 aspartic acid and A8 histidine are preferred for insulin-Fc fusion proteins that need to be repeatedly dosed in dogs over long periods long-term (e.g., to treat canine diabetes).

*Comparative Example 38: An Insulin-Fc Fusion Protein Comprising the Insulin Polypeptide of SEQ ID NO: 4 and a Non-Glycosylated Canine IgGB Isotype Fc Fragment in Which the B10D and A8H Mutations of the Insulin Polypeptide are Restored to Native Compositions to Reduce the Potential Risk of Immunogenicity.*

**[0233]** To evaluate whether replacing the "hot spot" mutations would improve the immunogenicity and repeated dose bioactivity of insulin-Fc fusion proteins comprising the insulin polypeptide of SEQ ID NO: 4 and the canine IgGB isotype fragment, an exemplary insulin-Fc fusion protein (SEQ ID NO: 43) was synthesized in which the B10 and A8 amino acids of the insulin polypeptide were restored to their native histidine and threonine compositions, respectively (SEQ ID NO: 63) listed below with non-native amino acids underlined.

FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSI<sup>1</sup>CSLYQLENYCN (SEQ ID NO: 63).

**[0234]** Furthermore, given the additional potential benefits of the non-glycosylated cNg mutants, the insulin-Fc fusion protein of SEQ ID NO: 43 contains the cNg-Q mutation. The entire amino acid sequence of the insulin-Fc fusion protein of SEQ ID NO: 43 is given below:

FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSI<sup>1</sup>CSLYQLENYCNGGGGAGGG  
GDCPKCPAPEMLGGPSVFIFPPKPKD<sup>2</sup>TL<sup>3</sup>LIARTPEVTCVVDLDPEDPEVQISW<sup>4</sup>FDGKQM<sup>5</sup>QTAKT  
QPREEQFQGT<sup>6</sup>YRVVSVLP<sup>7</sup>IGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQA<sup>8</sup>HQPSVYVLPPSR  
EELSKNTVSLTCLIKDFP<sup>9</sup>PDIDVEWQSN<sup>10</sup>GQEPESKYRTTP<sup>11</sup>QLDEDGSYFLYSKLSVDKSRWQR  
GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 43).

**[0235]** The insulin-Fc fusion protein of SEQ ID NO: 43 was manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. The resulting protein yield was only 21 mg/L. The structure was confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequence was further identified by LC-MS with glycan removal according to Example 8. The %homodimer as measured by size-exclusion chromatography according to Example 10, was 98.0% indicating that the protein was relatively free of aggregates.

**[0236]** Despite the relatively low homodimer titer of 21 mg/L, the insulin-Fc fusion protein of SEQ ID NO: 43 was evaluated in dogs for in vivo bioactivity and immunogenicity according to the procedures of Example 22, Example 23, and Example 24, respectively. FIG. 10 demonstrates that restoration of the B10D and A8H mutations to their native amino acids (i.e., B10H and A8T) in the insulin-Fc fusion protein of SEQ ID NO: 43 did significantly reduce the immunogenicity of the parent compound (SEQ ID NO: 36).

**[0237]** However, as shown in FIG. 11, the insulin-Fc fusion protein of SEQ ID NO: 43 containing the native B10 and A8 amino acids was not bioactive (i.e., the NAOC was essentially zero).

*Comparative Example 39: Attempts to Incorporate Additional B-chain and A-chain Mutations into the Insulin Polypeptide of SEQ ID NO: 63 to Improve the Bioactivity of the Associated Insulin-Fc Fusion Proteins Containing the Canine IgGB Fc Fragment.*

**[0238]** The fact that the insulin-Fc fusion protein of SEQ ID NO: 43 did not generate anti-drug antibodies (Example 38) compared to the insulin-Fc fusion protein of SEQ ID NO: 36 (Example 33) provides strong evidence for the theory that the B10D and A8H mutations in the insulin polypeptide of SEQ ID NO: 4 are likely the immunogenic epitopes responsible for the production of anti-drug antibodies. However, the lack of in vivo potency of the insulin-Fc fusion protein of SEQ ID NO: 43 compared to that of SEQ ID NO: 36 indicates that these two amino acid mutations are also responsible for achieving acceptable levels of bioactivity. The lack of in vivo potency for the insulin-Fc fusion protein of SEQ ID NO: 43 correlates with its high IC<sub>50</sub> (shown in Table 9 below) as measured by the insulin receptor binding assay according to the method of Example 12. Therefore, further efforts were required to increase the insulin-Fc fusion protein bioactivity (i.e., decrease the insulin receptor binding assay IC<sub>50</sub> value to less than 5000 nM, or more preferably less than 4000 nM, or even more preferably less than 3000 nM) while maintaining a low degree of immunogenicity by keeping the native B10 and A8 amino acids in the insulin polypeptide.

**[0239]** It is known that various portions of the insulin B-chain and A-chain are required for strong binding to the IR (Hubbard S.R., "Structural biology: Insulin meets its receptor", Nature. 2013; 493(7431):171-172). Therefore, portions of the B-chain or A-chain were modified while keeping the B10 and A8 the same as in native insulin and the C-chain and peptide linker constant. Several of these insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10, and their insulin receptor binding affinities were measured according to Example 12. Their sequences are shown below, and the resulting sequence alignments against SEQ ID NO: 43 are shown in FIG. 12 (Clustal Omega).

FVNQHLCGSHLVQALYLCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGAGGGG  
DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKTQ  
PREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
ELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 44)

FVNQHLCGSELVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGAGGGG  
DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKTQ  
PREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
ELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 45)

FVNQHLCGSHLVEALALVCGEAGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGAGGGG  
DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKTQ  
PREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
ELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 46)

FVNQHLCGSHLVEALALVCGERGFYYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGAGGG  
 GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
 5 QPREEQFSGTyrVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSR  
 EELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQR  
 GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 47)

10 FVNQHLCGSHLVEALALVCGERGFYYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGAGGGG  
 DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKTQ  
 PREEQFSGTyrVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
 15 ELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
 DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 48)

20

| Table 9: %homodimer, homodimer titers, and IR IC50 values for various SEQ ID NOs. |            |                            |              |
|-----------------------------------------------------------------------------------|------------|----------------------------|--------------|
| SEQ ID NO:                                                                        | %Homodimer | HEK homodimer titer (mg/L) | IR IC50 (nM) |
| SEQ ID NO: 43                                                                     | 98.0%      | 21                         | >5000        |
| 25 SEQ ID NO: 44                                                                  | 97.6%      | 9                          | 2624         |
| SEQ ID NO: 45                                                                     | 81.4%      | 17                         | 633          |
| SEQ ID NO: 46                                                                     | 99.1%      | 22                         | >5000        |
| SEQ ID NO: 47                                                                     | 96.6%      | 25                         | 2402         |
| 30 SEQ ID NO: 48                                                                  | 98.0%      | 6                          | >5000        |

[0240] In only three cases (SEQ ID NO: 44, SEQ ID NO: 45, and SEQ ID NO: 47 did the proposed mutations improve the IR binding (i.e., lower the IC50 value) as compared to SEQ ID NO: 43. However, none of the mutations resulted in compounds that meet the manufacturing design goal of a homodimer titer greater than 50 mg/L, and in some cases, the mutations lead to significantly reduced manufacturability (e.g., homodimer titers less than 20 mg/L).

Comparative Example 40: Attempts to Incorporate C-chain Mutations into the Insulin Polypeptide of SEQ ID NO: 63 to Improve the Bioactivity of the Associated Insulin-Fc Fusion Proteins Containing the Canine IgGB Fc Fragment.

40 [0241] The results obtained in Example 39 showed that all attempts to mutate the A-chain and B-chain of the insulin polypeptide of SEQ ID NO: 63 resulted in unacceptably low HEK homodimer titers of the associated insulin-Fc fusion (i.e., homodimer titers less than or equal to 25 mg/L). Therefore, there was a need for further experimentation. In the present example, the C-chain composition of the insulin polypeptide of SEQ ID NO: 63 was mutated by making it longer or by increasing its flexibility. Native insulin (e.g. human insulin) has been shown to undergo a significant conformational change that involves movement of both the B-chain and A-chain folding as it binds to the insulin receptor (e.g., as described by Menting, et al., Nature, 2013; 493(7431): pp241-245). Native insulin, unlike the insulin polypeptides of the present invention, is freely able to undergo this conformational change at the insulin receptor, because it is a two-chain polypeptide in its native form, connected only through two disulfide bonds with no C-chain constraining the mobility of the A- and B-chains. Without being bound by any particular theory, it was hypothesized that the C-chain contained within the insulin polypeptide of SEQ ID NO: 63 was too inflexible (e.g. an amino acid composition and sequence that does not permit facile movement between the B-chain and A-chain) and/or too short (e.g. not enough amino acids between the C-terminus of the B-chain and the N-terminus of the A-chain) thus preventing the insulin polypeptide from undergoing the necessary change in molecular shape required for strong binding to the insulin receptor. Therefore, several insulin-Fc fusion proteins were synthesized based on the insulin-Fc fusion protein of SEQ ID NO: 43 with variations in the insulin polypeptide C-chain as shown below with the resulting sequence alignments against SEQ ID NO: 43 shown in FIG. 13 (Clustal Omega).

FVNQHLCGSHLVQALYLCGERGFFYTDPTQRGGGGGGQRGIVEQCCTSICSLYQLENYCGGGGA  
 GGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQMOT  
 5 AKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVL  
 PPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDGSYFLYSKLSVDKSR  
 WQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 49)

10 FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGGGGSGGGGGIVEQCCTSICSLYQLENYCGGGG  
 AGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQM  
 QTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVY  
 15 VLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDGSYFLYSKLSVDK  
 SRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 50)

20 FVNQHLCGSHLVEALALVCGERGFFYTDPGGGGGGGGGIVEQCCTSICSLYQLENYCGGGGAGG  
 GGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQMOTAK  
 TQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPS  
 25 REELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQ  
 RGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 51)

30 FVNQHLCGSHLVEALALVCGERGFFYTPGGGGGGGGGGIVEQCCTSICSLYQLENYCGGGGAGGG  
 GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQMOTAKT  
 QPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSR  
 35 EELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQR  
 GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 52)

**Table 10: %homodimer, homodimer titers, and IR IC50 values for various SEQ ID NOs.**

| SEQ ID NO:    | %Homodimer | HEK homodimer titer (mg/L) | IR IC50 (nM) |
|---------------|------------|----------------------------|--------------|
| SEQ ID NO: 43 | 98.0%      | 21                         | >5000        |
| SEQ ID NO: 49 | 94.0%      | 8                          | 4176         |
| SEQ ID NO: 50 | 99.6%      | 37                         | 1609         |
| SEQ ID NO: 51 | 98.3%      | 42                         | >5000        |
| SEQ ID NO: 52 | 98.6%      | 33                         | 4720         |

**[0242]** The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10, and their insulin receptor binding affinities were measured according to Example 12. The results are shown in Table 10. In only one case, (SEQ ID NO: 50) which comprises the longest C-chain (GGGGGGSGGGG - SEQ ID NO: 71), did a C-chain mutation significantly improve the insulin receptor binding affinity (IC50 less than 3000 nM) compared to that of the insulin-Fc fusion protein of SEQ ID NO: 43. However, none of these C-chain-mutated insulin-Fc fusion proteins exhibited a homodimer titer greater than the manufacturing design goal of 50 mg/L. In fact, in one case (SEQ ID NO: 49) the C-

chain mutation unexpectedly led to significantly lower homodimer titers.

*Comparative Example 41: Attempts to Incorporate Peptide Linker Mutations into Insulin-Fc Fusion Proteins Containing the Insulin Polypeptide of SEQ ID NO: 63 and the Canine IgGB Fc Fragment to Improve Bioactivity.*

**[0243]** Without being bound by any particular theory, another possible reason for the poor insulin receptor binding of the insulin-Fc fusion protein of SEQ ID NO: 43 was thought to involve the steric hindrance between the insulin polypeptide and the insulin receptor resulting from the close proximity of the much larger Fc fragment molecule attached to the insulin polypeptide through the peptide linker. Shorter peptide linkers or more tightly folded peptide linkers were thought to potentially exacerbate this issue, while longer peptide linkers or peptide linkers that are resistant to folding back on themselves (e.g., linkers with more molecular stiffness) may alleviate this issue by creating more space between the insulin polypeptide and the Fc fragment. The increased space between the insulin polypeptide and the Fc fragment would also increase the distance between the insulin receptor and the Fc fragment leading to less interference during insulin receptor binding. The peptide linker of SEQ ID NO: 11 (i.e., GGGGAGGGG) used to construct the insulin-Fc fusion protein of SEQ ID NO: 43 was hypothesized to be potentially too short and/or too flexible, because the amino acids that comprise the linker contain no side chains (i.e., it contains only glycine and alanine amino acids). Therefore, to test this hypothesis, two other insulin-Fc fusion protein variants of the insulin-Fc fusion protein of SEQ ID NO: 43 were synthesized. The insulin-Fc fusion protein of SEQ ID NO: 48 contained the same peptide linker as was used to construct the insulin-Fc fusion protein of SEQ ID NO: 43 but with an insulin polypeptide in which the asparagine at the 21<sup>st</sup> position from the N-terminus of the A chain (i.e., A21) was absent (i.e., des-A21). This particular mutation was incorporated to see whether the junction between the A-chain and the peptide linker affects the protein yield and/or bioactivity of the molecule. The other insulin-Fc fusion protein of SEQ ID NO: 53 contains this des-A21N A-chain mutation and a peptide linker that is more than twice the length of that used to construct the insulin-Fc fusion protein of SEQ ID NO: 43. In this longer peptide linker, alanine is disfavored and instead is replaced with a glutamine, which contains a polar amide side chain. The glutamine substitutions were expected to increase the hydrophilic nature of the peptide linker and potentially prevent the linker from folding back against itself. The sequences are shown below with the resulting sequence alignments against SEQ ID NO: 43 shown in FIG. 14 (Clustal Omega).

FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGGQGGG  
GQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF  
VDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQA  
HQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQGQEPESKYRTTPPQLDEDGSYFLYS  
KLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 53)

FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCGGGGAGGGG  
DCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFVDGKQMQTAKTQ  
PREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSRE  
ELSKNTVSLTCLIKDFFPPDIDVEWQSNQGQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRG  
DTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 48)

**Table 11: %homodimer, homodimer titers, and IR IC50 values for various SEQ ID NOs.**

| SEQ ID NO:    | %Homodimer | HEK Homodimer titer (mg/L) | IR IC50 (nM) |
|---------------|------------|----------------------------|--------------|
| SEQ ID NO: 43 | 98.0%      | 21                         | >5000        |
| SEQ ID NO: 48 | 98.0%      | 6                          | >5000        |
| SEQ ID NO: 53 | 99.6%      | 11                         | 1281         |

**[0244]** The two insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-

reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10, and their insulin receptor binding affinities were measured according to Example 12. The results are given in Table 11. The incorporation of a longer peptide linker of different composition (GGGGGQGGGGQGGGGQGGGGG (SEQ ID NO: 13) for SEQ ID NO: 53 vs. GGGGAGGGG (SEQ ID NO: 11) for SEQ ID NO: 43) did improve the insulin receptor binding as measured by a significant reduction in the IC50 value, indicating that longer linkers may be a strategy for increasing insulin receptor binding for other insulin-Fc fusion proteins. However, the incorporation of a longer linker still did not improve the homodimer titers to above the manufacturing design goal of greater than 50 mg/L.

*Comparative Example 42: Attempts to Delete Portions of the B-chain of the Insulin Polypeptide of SEQ ID NO: 63 to Improve the Homodimer Titer of the Associated Insulin-Fc Fusion Proteins Containing the Canine IgGB Fc Fragment.*

**[0245]** The results from Example 41 demonstrate that the peptide linker can be modified to increase the insulin receptor binding affinity of the insulin-Fc fusion protein of SEQ ID NO: 43, which contains the native B10 and A8 amino acids. However, the peptide linker mutation failed to increase the homodimer titer enough to meet the manufacturing design goal. Because the homodimer titer is a function of several properties, including the intracellular synthesis and processing within cells, it was hypothesized that perhaps the insulin-Fc molecule was self-associating (i.e., aggregating) during and after synthesis either intramolecularly between the two monomers of the homodimer or intermolecularly between two or more separate homodimers. This aggregation would lead to unacceptably low homodimer titers obtained from the cell culture supernatants during the production process described in Example 1, Example 4, and Example 10. This potential interaction between the insulin-Fc fusion protein molecules could be due, in part, to insulin's well-known propensity to self-associate and form aggregates. One method known in the art to reduce the propensity for insulin to self-associate involves mutating the amino acids near the C-terminus of the B-chain. For example, insulin lispro (B28K; B29P mutations) and insulin aspart (B28D mutation) are well-known commercial two-chain insulins with non-native B-chain mutations that prevent association and aggregation thus giving rise to a predominantly monomeric form of insulin in solution. Another approach to prevent aggregation involves amino acid structural deletions. For example, a two-chain insulin known as despentapeptide insulin (DPPI; see Brange J., Dodson G.G., Edwards J., Holden P.H., Whittingham J.L. 1997b. "A model of insulin fibrils derived from the x-ray crystal structure of a monomeric insulin (despentapeptide insulin)" Proteins 27 507-516), is identical to native two-chain human insulin except that the five C-terminal amino acids of the B-chain (YTPKT) are removed. DPPI has a lower binding affinity to the insulin receptor as compared to the native two-chain human insulin, but it is completely monomeric in solution, meaning that there is no significant association or aggregation between DPPI molecules. Therefore, in an attempt to decrease the potential for intramolecular and intermolecular self-association and improve the insulin-Fc fusion protein homodimer titer, several variants of the insulin-Fc fusion protein of SEQ ID NO: 43 were constructed using partial B-chain amino acid truncation and B-chain amino acid mutations as described above for DPPI, insulin lispro, and insulin aspart. The sequences are shown below with the resulting sequence alignments against SEQ ID NO: 43 shown in FIG. 15 (Clustal Omega).

FVNQHLCGSHLVEALALVCGERGFFYTDPGGGGGGGGGIVEQCCTSICSLYQLENYCGGGGAGG  
GGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAK  
TQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPS  
REELSKNTVSLTCLIKDFFPPDIDVEWQSNQGQEPESKYRTPPQLDEDGSYFLYSKLSVDKSRWQ  
RGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 51)

FVNQHLCGSHLVEALALVCGERGFFYTDPGGGGGGGGGIVEQCCTSICSLYQLENYCGGGGAGGG  
GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
QPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPPSR  
EELSKNTVSLTCLIKDFFPPDIDVEWQSNQGQEPESKYRTPPQLDEDGSYFLYSKLSVDKSRWQR  
GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 52)

FVNQHLCGSHLVEALALVCGERGFYFYTQGGGGGGGGGIVEQCCTSICSLYQLENYCGGGGAGGG  
 GDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWFDGKQMQTAKT  
 5 QPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKARGQAHQPSVYVLPSSR  
 EELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQR  
 GDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 54)

10

| Table 12: %homodimer, homodimer titers, and IR IC50 values for various SEQ ID NOs. |            |                            |              |
|------------------------------------------------------------------------------------|------------|----------------------------|--------------|
| SEQ ID NO:                                                                         | %Homodimer | HEK Homodimer titer (mg/L) | IR IC50 (nM) |
| 15 SEQ ID NO: 43                                                                   | 98.0%      | 21                         | >5000        |
| SEQ ID NO: 51                                                                      | 98.3%      | 42                         | 1915         |
| SEQ ID NO: 54                                                                      | 99.4%      | 22                         | 2195         |
| SEQ ID NO: 52                                                                      | 98.6%      | 33                         | 1930         |

20 [0246] The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10, and their  
 25 insulin receptor binding affinities were measured according to Example 12. The results are given in Table 12. The homodimer titer of the resulting compounds was only significantly increased in one case (SEQ ID NO: 51), but unexpectedly, the insulin receptor affinity was improved for all of the mutated compounds (SEQ ID NO: 51, SEQ ID NO: 54, and SEQ ID NO: 52).

30 Example 43: Attempts to Combine B-chain, C-chain, and A-chain Mutations, B-chain Truncation, and Linker Mutations to the Insulin-Fc Fusion Protein of SEQ ID NO: 43 to Further Improve Homodimer Titer and Bioactivity.

[0247] As shown in Example 39, Example 40, Example 41 and Example 42, no single strategy successfully incorporated  
 35 an insulin polypeptide comprising the non-immunogenic native B10 and A8 amino acids with the canine IgGB Fc fragment to form an insulin-Fc fusion protein with acceptable insulin receptor activity and homodimer titer. Therefore, the concepts of a longer C-chain, a longer peptide linker, and truncation of the C-terminal amino acids of the B-chain were combined. In addition, to potentially further decrease the propensity for self-association and aggregation, additional point mutations were introduced to the native insulin hydrophobic amino acid residue sites using less hydrophobic amino acids, including  
 40 those with side groups that are negatively or positively charged at physiological pH. Example mutations included tyrosine to alanine, tyrosine to glutamic acid, isoleucine to threonine, and phenylalanine to histidine. Furthermore, to simplify the analysis, in all cases the cNg site of the canine IgGB Fc fragment was restored to its native asparagine. The sequences for these insulin-Fc fusion protein variants are shown below with the resulting sequence alignments against SEQ ID NO: 43 shown in FIG. 16 (Clustal Omega).

45 FVNQHLCGSHLVEALELVCGERGFYTPKTGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQIS  
 WFVDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
 50 GQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPPQLDEDGSYF  
 LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 55)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCNHGGGGQ  
 GGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQ  
 5 ISWFVDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKA  
 RGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPQLDEDGS  
 YFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 56)

10 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCNGGGGGQ  
 GGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQ  
 15 ISWFVDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKA  
 RGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPQLDEDGS  
 YFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 28)

20 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
 GGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQIS  
 WFVDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
 25 GQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPQLDEDGSYF  
 LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 26)

30 FVNQHLCGSHLVEALELVCGERGFFYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
 GGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQIS  
 WFVDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
 35 GQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNGQQEPESKYRTTPQLDEDGSYF  
 LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 57)

| Table 13: %homodimer, homodimer titers, and IR IC50 values for various SEQ ID NOs. |            |                            |              |
|------------------------------------------------------------------------------------|------------|----------------------------|--------------|
| SEQ ID NO:                                                                         | %Homodimer | HEK homodimer titer (mg/L) | IR IC50 (nM) |
| SEQ ID NO: 43                                                                      | 98.0%      | 21                         | >5000        |
| SEQ ID NO: 55                                                                      | 97.9%      | 69                         | 3869         |
| SEQ ID NO: 56                                                                      | 99.5%      | 101                        | 554          |
| SEQ ID NO: 28                                                                      | 99.7%      | 107                        | 1247         |
| SEQ ID NO: 57                                                                      | 99.7%      | 128                        | 2043         |
| SEQ ID NO: 26                                                                      | 99.4%      | 187                        | 2339         |

[0248] The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10, and their insulin receptor binding affinities were measured according to Example 12. The results are given in Table 13. The results show that a combination of decreasing the hydrophobicity of certain B-chain and A-chain amino acids, using longer and more flexible C-peptide sequences, truncating several C-terminal B-chain amino acids, and using a longer peptide linker

resulted in several useful insulin-Fc fusion proteins that meet the minimum homodimer titer and insulin receptor binding activity design criteria. SEQ ID NO: 56, SEQ ID NO: 28, SEQ ID NO: 26, and SEQ ID NO: 57 showed more preferable insulin receptor IC50 values (less than 3000 nM) and more preferable HEK homodimer titer values (greater than 100 mg/L) than either SEQ ID NO: 43 or SEQ ID NO: 55. Surprisingly, changing just a few amino acids leads to a multifold improvement in insulin receptor affinity, and, in the case of the insulin-Fc fusion protein of SEQ ID NO: 26 a dramatic increase in homodimer titer over the original insulin-Fc fusion protein of SEQ ID NO: 43.

*Comparative Example 44: In Vivo Bioactivity, Repeated Dose Bioactivity, and Immunogenicity of Insulin-Fc Fusion Proteins Constructed from the Insulin Polypeptide of SEQ ID NO: 7, the Peptide Linker of SEQ ID NO: 13, and the Canine IgGB Fc Fragment of SEQ ID NO: 15.*

**[0249]** Given the positive homodimer titer and insulin receptor binding activity results from Example 43, two of the most promising insulin-Fc fusion proteins (SEQ ID NO: 26 and SEQ ID NO: 28) were tested in dogs to evaluate the repeated dose bioactivity and immunogenicity. Each compound comprises the longer, more hydrophilic peptide linker of SEQ ID NO: 13 and the more manufacturable, less aggregated canine IgGB Fc fragment of SEQ ID NO: 15. Most importantly, both insulin-Fc fusion proteins comprise insulin polypeptides with the putatively less immunogenic native B10 and A8 amino acids (i.e. general SEQ ID NO: 6). In the case of the insulin-Fc fusion protein of SEQ ID NO: 28, the asparagine at position A21 is present (i.e. the insulin polypeptide comprises SEQ ID NO: 8). In the case of the insulin-Fc fusion protein of SEQ ID NO: 26, the asparagine at position A21 is absent (i.e. the insulin polypeptide comprises SEQ ID NO: 7).

**[0250]** The in vivo bioactivity of the insulin-Fc fusion protein of SEQ ID NO: 28 was tested in N=1 dog according to the procedure of Example 21. The results shown in FIG. 17 for a single subcutaneous dose demonstrate that the insulin-Fc fusion protein of SEQ ID NO: 28 is indeed bioactive in vivo with an NAOC of 1076 %FBGL·days·kg/mg calculated according to the procedure in Example 22. The insulin-Fc fusion protein of SEQ ID NO: 28 pharmacokinetic profile was measured by the method of Example 23 using ELISA, and a two-compartment model was fit to the data to determine its elimination half-life which was 3.5 days.

**[0251]** The repeated dose bioactivity was then evaluated by continuing to subcutaneously administer the insulin-Fc fusion protein of SEQ ID NO: 28 to N=1 dog on day 14, day 28, and day 42 after the initial injection according to the procedure of Example 15. When the dog's %FBGL dropped too low, the dog was given food to raise the blood glucose to a safe level. The NAOC and NAOCR were measured for each subsequent dose according to the general procedure of Example 22, calculated from the time the dose was administered until just before the next dose was administered. The NAOC and the NAOCR shown in Table 14 illustrate that the insulin-Fc fusion protein of SEQ ID NO: 28 maintains an NAOCR greater than 0.8 throughout the four doses thus meeting the repeated dose bioactivity design goal.

| Table 14: NAOC per dose for repeated doses of SEQ ID NO: 28 |     |                         |       |
|-------------------------------------------------------------|-----|-------------------------|-------|
| Injection#                                                  | Day | NAOC (%FBGL·days·kg/mg) | NAOCR |
| 1                                                           | 0   | 1076                    | 1.0   |
| 2                                                           | 14  | 1005                    | 0.9   |
| 3                                                           | 28  | 900                     | 0.8   |
| 4                                                           | 42  | 838                     | 0.8   |

**[0252]** The immunogenicity of the insulin-Fc fusion protein of SEQ ID NO: 28 was tested according to the procedure of Example 24. FIG. 18 demonstrates that the insulin-Fc fusion protein of SEQ ID NO: 28 exhibits no apparent immunogenicity in vivo in agreement with the maintenance of in vivo bioactivity throughout the repeated dose experiment.

**[0253]** The insulin-Fc fusion protein of SEQ ID NO: 26, with the asparagine at A21 of the insulin polypeptide chain absent, was also evaluated for repeated dose bioactivity performance in dogs. The compound was administered subcutaneously to N=1 dog on day 0, day 14, day 28, and on day 42 according to the procedure of Example 22. When the dog's %FBGL dropped too low, the dog was given food to raise the blood glucose to a safe level. The NAOC for the first injection was an impressive 2278 %FBGL·days·kg/mg, showing that the insulin-Fc fusion protein of SEQ ID NO: 26 was satisfactorily bioactive in vivo at almost twice the potency of the insulin-Fc fusion protein of SEQ ID NO: 28. The pharmacokinetic profile of the insulin-Fc fusion protein was measured by the method of Example 23 using ELISA, and a two-compartment model was fit to the data to determine its elimination half-life which was  $4.1 \pm 0.7$  days. FIG. 19 and FIG. 20 show the single dose blood glucose control and the multidose, multiweek blood glucose control for animals receiving the homodimer of SEQ ID NO: 26. The NAOC and NAOCR were also measured for each subsequent dose according to the general procedure of Example 22, calculated from the time the dose was administered until just before the next

dose was administered. The NAOC and the NAOCR shown in Table 15 illustrate that the insulin-Fc fusion protein of SEQ ID NO: 26 maintains an NAOCR greater than or equal to 1.0 throughout the four doses thus meeting the repeated dose bioactivity design goal described in Example 29.

[0254] The immunogenicity of the insulin-Fc fusion protein of SEQ ID NO: 26 was tested according to the procedure of Example 24. FIG. 21 demonstrates that the insulin-Fc fusion protein of SEQ ID NO: 26 exhibits no apparent immunogenicity in vivo in agreement with the maintenance of in vivo bioactivity throughout the repeated dose experiment.

| Table 15: NAOC per dose for repeated doses of SEQ ID NO: 26 |     |                         |       |
|-------------------------------------------------------------|-----|-------------------------|-------|
| Injection#                                                  | Day | NAOC (%FBGL-days-kg/mg) | NAOCR |
| 1                                                           | 0   | 2278                    | 1.0   |
| 2                                                           | 14  | 4029                    | 1.8   |
| 3                                                           | 28  | 3450                    | 1.5   |
| 4                                                           | 42  | 3257                    | 1.4   |

[0255] As discussed in the Detailed Description of the invention, a known enzymatic cleavage site exists between asparagine-glycine bonds (Vlasak, J., Ionescu, R., (2011) MAb Vol. 3, No. 3 pp 253-263). Omitting the asparagine at the 21st amino acid in the A chain (i.e., A21) in the insulin polypeptide of SEQ ID NO: 7 contained in the insulin-Fc fusion protein of SEQ ID NO: 26 with the peptide linker of SEQ ID NO: 13, eliminates the possibility of enzymatic cleavage of the asparagine-glycine bond between the C-terminus of the A-chain and the N-terminus of the peptide linker. However, the insulin-Fc fusion protein of SEQ ID NO: 28 comprises the peptide linker of SEQ ID NO: 13 and the insulin polypeptide of SEQ ID NO: 8, which keeps the asparagine at A21. Therefore, it would have been expected that the insulin-Fc fusion protein of SEQ ID NO: 28 would have been enzymatically digested during synthesis or in vivo following subcutaneous administration. However, rather unexpectedly the insulin-Fc fusion protein of SEQ ID NO: 28 was manufacturable in HEK cells with an acceptable homodimer titer and demonstrated acceptable bioactivity in vivo with no signs of enzymatic digestion compromising its bioactivity.

Comparative Example 45: Confirmation of the Canine IgGB isotype Fc Fragment for Optimal Manufacturability and In Vivo Efficacy of Insulin-Fc Fusion Proteins Comprising the Preferred Insulin Polypeptide of SEQ ID NO: 7 and the Preferred Peptide Linker of SEQ ID NO: 13.

[0256] Having discovered a new insulin polypeptide and peptide linker combination resulting in non-immunogenic, high yielding, high purity, and highly bioactive insulin-Fc fusion proteins as described in Example 43 and Example 44, a question remained as to whether the canine IgGB Fc fragment was still the preferred isotype with respect to homodimer titer and bioactivity as was the case for the insulin-Fc fusion proteins in Example 32 and Example 33. Therefore, additional insulin-Fc fusion proteins were designed wherein the insulin polypeptide (SEQ ID NO: 7) and peptide linker (SEQ ID NO: 13) of the insulin-Fc fusion protein of SEQ ID NO: 26 were kept constant, and the canine IgGB Fc fragment of SEQ ID NO: 15 was replaced by the canine IgGA Fc fragment of SEQ ID NO: 14, the canine IgGC Fc fragment of SEQ ID NO: 16, or the canine IgGD Fc fragment of SEQ ID NO: 17. The sequences for these resulting insulin-Fc fusion protein variants are shown below:

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQIS  
WFVDGKQMQTAKTQPREEQFNGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
GGAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYF  
LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 26)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGRCCTDTPPCVPPEPLGGPSVLIFPPKPKDILRITRTPVTCVVLDLGREDPEV  
 5 QISWFVDGKEVHTAKTQSREQQFNQTYRVS SVLPPIEHQDWLTGKEFKCRVNHIDLPSPERTISKA  
 RGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNGQQEPERKHRMTPPQLDEDGS  
 YFLYSKLSVDKSRWQQGDPFTCAVMHETLQNHYTDLSLSHSPG (SEQ ID NO: 58)

10 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGCNNCPGCGLLGGPSVFIFPPKPKDILVTARTPTVTCVVVDLDPENPEVQ  
 ISWFVDSKQVQTANTQPREEQSNGTYRVS SVLPPIGHQDWLSGKQFKCKVNNKALPSPIEEIISKTP  
 15 GQAHQPNVYVLPSPKELSSSDTVTLTCLIKDFFPPEIDVEWQSNGQQEPESKYRMTTPPQLDEDGS  
 YFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQISLSHSPG (SEQ ID NO: 59)

20 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGCISPCVPPELGGPSVFIFPPKPKDILRITRTPVTCVVLDLGREDPEVQISWF  
 VDGKEVHTAKTQPREQQFNSTYRVS SVLPPIEHQDWLTGKEFKCRVNHIGLPSPIERTISKARGQAH  
 25 QPSVYVLPSPKELSSSDTVTLTCLIKDFFPPEIDVEWQSNGQQEPESKYHTTAPQLDEDGSYFLYS  
 KLSVDKSRWQQGDTFTCAVMHEALQNHYTDLSLSHSPG (SEQ ID NO: 60).

[0257] The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using  
 a Protein A or Protein G columns according to Example 4. Their structures were confirmed according to Example 6 by  
 30 non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according  
 to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10,  
 and their insulin receptor binding affinities were measured according to Example 12. Additionally, the insulin-Fc fusion  
 protein affinities for the canine FcRn receptor were measured according to Example 15. As is shown in Table 16, the  
 35 insulin-Fc fusion protein of SEQ ID NO: 26 comprising the canine IgGB Fc fragment demonstrated the highest homodimer  
 titer of these sequences. The insulin-Fc fusion protein of SEQ ID NO: 58 comprising the canine IgGA Fc fragment  
 exhibited poor homodimer titer when purified using a Protein A column; however, when it purified using a Protein G  
 column, the homodimer titer was significantly improved, exceeding the design goal of greater than 50 mg/L. The same  
 was true for the insulin-Fc fusion protein of SEQ ID NO: 59 comprising the canine IgGC Fc fragment. The insulin-Fc  
 40 fusion protein of SEQ ID NO: 60 comprising the canine IgGD Fc fragment did not yield any compound when purified  
 with either a Protein A or a Protein G column. Therefore, as was demonstrated with the insulin-Fc fusion protein of SEQ  
 ID NO: 36 containing a different insulin polypeptide (SEQ ID NO: 4) and peptide linker (SEQ ID NO: 11), the canine  
 IgGB was the preferred Fc fragment with respect to homodimer titer (see Example 32).

**Table 16: Homodimer titers, IR binding, and FcRn binding for sequences utilizing native canine IgGA, IgGB, IgGC, and IgGD Fc fragments**

| SEQ ID NO:    | Fc Fragment IgG Isotype | Protein Yield Protein A / (Protein G) (mg/L) | %Homo-dimer Protein A / (Protein G) | Homo-dimer Titer (mg/L) | IR Binding, IC50 (nM) | FcRn Binding, EC50 (ng/mL) | First dose NAOC (%FBGL· days·kg/mg) |
|---------------|-------------------------|----------------------------------------------|-------------------------------------|-------------------------|-----------------------|----------------------------|-------------------------------------|
| SEQ ID NO: 26 | IgGB                    | 187 / (DNM)                                  | 99% / (DNM)                         | 185                     | 2339                  | 599                        | 2278                                |

(continued)

**Table 16: Homodimer titers, IR binding, and FcRn binding for sequences utilizing native canine IgGA, IgGB, IgGC, and IgGD Fc fragments**

| SEQ ID NO:    | Fc Fragment IgG Isotype | Protein Yield Protein A / (Protein G) (mg/L) | %Homo-dimer Protein A / (Protein G) | Homo-dimer Titer (mg/L) | IR Binding, IC50 (nM) | FcRn Binding, EC50 (ng/mL) | First dose NAOC (%FBGL·days·kg/mg) |
|---------------|-------------------------|----------------------------------------------|-------------------------------------|-------------------------|-----------------------|----------------------------|------------------------------------|
| SEQ ID NO: 58 | IgGA                    | 10 / (69)                                    | 45% / (91%)                         | 62‡                     | 2586 <sup>#</sup>     | 1610                       | 174                                |
| SEQ ID NO: 59 | IgGC                    | 0/(86)                                       | 0% / (94%)                          | 81‡                     | 2084‡                 | >200000                    | 39                                 |
| SEQ ID NO: 60 | IgGD                    | 0 / (0)                                      | (DNM)/ (DNM)                        | 0                       | DNM                   | DNM                        | DNM                                |

DNM = did not measure; <sup>#</sup> = purified via Protein A; ‡ = purified by Protein G.

**[0258]** The in vivo bioactivity of the insulin-Fc fusion protein of SEQ ID NO: 58 comprising the canine IgGA Fc fragment that was purified via Protein G was tested according to the procedure of Example 21. The results illustrated in FIG. 22 show that the insulin-Fc fusion protein of SEQ ID NO: 58 is only somewhat bioactive in vivo with a NAOC of only 174 %FBGL·days·kg/mg calculated according to Example 22.

**[0259]** The in vivo bioactivity of the insulin-Fc fusion protein of SEQ ID NO: 59 comprising the canine IgGC Fc fragment was purified via Protein G tested according to the procedure of Example 21. The results illustrated in FIG. 23 show that the insulin-Fc fusion protein of SEQ ID NO: 59 is only somewhat bioactive in vivo with a NAOC of only 39 %FBGL·days·kg/mg calculated according to Example 22.

**[0260]** Therefore, as was demonstrated with the insulin-Fc fusion protein of SEQ ID NO: 36 containing a different insulin polypeptide (SEQ ID NO: 4 and peptide linker (SEQ ID NO: 11), the canine IgGB was the preferred Fc fragment with respect to bioactivity (see Example 32 and Example 33 and Table 16 above).

Comparative Example 46: Non-Glycosylated Insulin-Fc fusion Proteins Comprising the Insulin Polypeptide of SEQ ID NO: 8, the Peptide Linker of SEQ ID NO: 13, and the Canine IgGB Fc Fragment to Reduce the Potential Risk of Immunogenicity.

**[0261]** While the insulin-Fc fusion protein of SEQ ID NO: 26 meets all of the design goals (Example 28), there may or may not be a risk of immunogenicity over extended periods of treatment (e.g., 6 months, 1 year, 2 years or more) which could compromise the use of this insulin-Fc fusion protein for treating diabetes should this occur. As described in the Detailed Description of the Invention and in Example 34 and Example 35, one possible cause of a reduction in bioactivity after repeated doses is the unwanted interaction of the canine IgGB Fc fragment with the dog's immune system resulting in the production of neutralizing anti-drug antibodies. However, the results shown in Example 45 demonstrate that unexpectedly, the canine IgGB isotype was the only option of the four canine IgG isotypes that yielded the desired manufacturability and bioactivity. Therefore, further Fc mutations were explored to achieve non-glycosylated insulin-Fc fusion proteins with low Fc(gamma)RI receptor binding, which should reduce the long-term, chronic immunogenicity risk.

**[0262]** As described in the Detailed Description of the Invention, one method for reducing the Fc(gamma)RI interaction involves mutating the Fc fragment cNg site to prevent glycosylation during synthesis in the host cell. Therefore, cNg site mutations were made to the Fc fragment region of SEQ ID NO: 26 to reduce the binding affinity of the Fc fragment for Fc(gamma) receptors in vivo, as measured by binding in an in vitro human Fc(gamma)RI assay described in Example 15. The position of the cNg site in the insulin-Fc fusion protein of SEQ ID NO: 26 is cNg-NB151. Mutations to SEQ ID NO: 26 included SEQ ID NO: 62 comprising a cNg-NB151-S mutation and SEQ ID NO: 61 comprising the same cNg-NB151-S mutation as well as a NB119-A mutation. The NB119-A was incorporated in a further attempt to reduce the interaction with Fc(gamma)RI as has been described only for use in mouse antibodies by Lo, M. et al. "Effector attenuating

substitutions that maintain antibody stability and reduce toxicity in mice", J. Biol. Chem. (2017), pp. 1-20. The full amino acid sequences of the resulting insulin-Fc fusion proteins are listed below (NB119 and NB151 sites underlined for clarity) along with their sequence alignments (Clustal Omega) which are shown in FIG. 24:

5 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVDLDPEDPEVQIS  
WFVDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
10 GQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYF  
LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 61)

15 FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVDLDPEDPEVQIS  
WFVDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKAR  
20 GQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPPQLDEDGSYF  
LYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 62).

[0263] The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10, and their insulin receptor binding affinities were measured according to Example 12. As shown in Table 17, incorporating the cNg-NB151-S mutations on the Fc fragment decreased the %homodimer, indicating an unacceptably high level of aggregation (i.e., the %homodimer dropped to just above 70%).

**Table 17: Homodimer titers for non-glycosylated insulin-Fc fusion proteins of SEQ ID NO: 61 and 104**

| SEQ ID NO:    | IgG Fragment | Relevant Mutations    | Protein Yield (mg/L) | %Homodimer | Homodimer Titer (mg/L) | IR Binding, IC50 (nM) |
|---------------|--------------|-----------------------|----------------------|------------|------------------------|-----------------------|
| SEQ ID NO: 26 | IgGB         | cNg-NB-151-N          | 187                  | 99%        | 185                    | 2339                  |
| SEQ ID NO: 61 | IgGB         | cNg-NB-151-S, NB119-A | 78                   | 73%        | 57                     | 3093                  |
| SEQ ID NO: 62 | IgGB         | cNg-NB151-S           | 130                  | 71%        | 93                     | 2302                  |

[0264] The in vivo bioactivity of the insulin-Fc fusion proteins of SEQ ID NO: 61 and SEQ ID NO: 62 was tested in N=1 dog each according to the procedure of Example 21. The results shown in FIG. 25 for a single subcutaneous dose demonstrate that both compounds were significantly less bioactive in vivo than the insulin-Fc fusion protein of SEQ ID NO: 26 (NAOC for SEQ ID NO: 62 = 574 %FBGL·days·kg/mg; NAOC for SEQ ID NO: 61 = 921 %FBGL·days·kg/mg). The results indicate that incorporating cNg-NB151-S mutations on the Fc fragment to produce non-glycosylated versions of the insulin-Fc fusion protein of SEQ ID NO: 26 unexpectedly decreased the in vivo bioactivity of the resulting compounds.

[0265] In an attempt to lessen the degree of aggregation and improve the bioactivity of the insulin-Fc fusion protein of SEQ ID NO: 62 containing the cNg-NB 151-S site mutation, various insulin-polypeptide B-chain variants were investigated with mutations in the region thought to be responsible for aggregation. The insulin-Fc fusion proteins were manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. Their structures were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. Their %homodimer content was measured by size-exclusion chromatography according to Example 10. Among the B-chain variants tested, one insulin Fc-fusion protein (SEQ ID NO: 30) containing a tyrosine to alanine substitution at the 16<sup>th</sup> amino acid from the N-terminus of the

B-chain (i.e., B16) was unexpectedly found to have high homodimer titers (105 mg/L) with low aggregation (99% homodimer), resulting in a homodimer titer of 104 mg/L. The insulin receptor binding measured according to Example 12 was acceptable with an IC<sub>50</sub> of 2040 nM. The FcRn receptor binding affinity EC<sub>50</sub> value measured according to Example 16 was 1194 ng/mL. The pharmacokinetic profile of the insulin-Fc fusion protein of SEQ ID NO: 30 was measured by the method of Example 23 using ELISA, and a two-compartment model was fit to the data to determine its elimination half-life which was  $4.1 \pm 0.7$  days. The sequence of SEQ ID NO: 30 is shown below (B16A and cNg-NB151-S mutations underlined for clarity).

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
GGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQ  
ISWFDVGKQMQTAKTQPRREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISKA  
RGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTPQLDEDGS  
YFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG (SEQ ID NO: 30).

[0266] The insulin-Fc fusion protein of SEQ ID NO: 30 was then evaluated for repeated dose bioactivity performance in dogs. The compound was administered subcutaneously to N=1 dog on day 0, day 7, day 14, and on day 28 according to the procedure of Example 22. When the dog's %FBGL dropped too low, the dog was given food to raise the blood glucose to a safe level. Unexpectedly, compared to the insulin-Fc fusion protein of SEQ ID NO: 62, the NAO for the first injection of the insulin-Fc fusion protein of SEQ ID NO: 30 containing the B16A mutation, was significantly higher (1185 %FBGL·days·kg/mg). The first dose in vivo bioactivity plot is shown in FIG. 26. The pharmacokinetic profile of the compound was also measured by the method of Example 23 using ELISA, and a two-compartment model was fit to the data to determine its elimination half-life which was 3.5 days. The NAO and NAOCR were also measured for each subsequent dose according to the general procedure of Example 22, calculated from the time the dose was administered until just before the next dose was administered. The NAO and the NAOCR shown in Table 18 illustrate that the insulin-Fc fusion protein of SEQ ID NO: 30 maintains an NAOCR greater than or equal to 0.6 throughout the four doses thus meeting the repeated dose bioactivity design goal. Taken together, the results indicate that it was necessary to mutate the insulin B-chain sequence to obtain a suitable, non-glycosylated cNg-S variant of SEQ ID NO: 26. Therefore, the insulin polypeptide of SEQ ID NO: 10 was preferred for non-glycosylated insulin-Fc fusion proteins comprising cNg-mutated canine IgGB Fc fragments.

**Table 18: NAO per dose for repeated doses of SEQ ID NO: 30**

| Injection# | Day | NAOC (%FBGL·days·kg/mg) | NAOCR |
|------------|-----|-------------------------|-------|
| 1          | 0   | 1185                    | 1.0   |
| 2          | 7   | 954                     | 0.8   |
| 3          | 14  | 764                     | 0.6   |
| 4          | 28  | 991                     | 0.8   |

[0267] Finally, select compounds were tested for their likelihood to interact with the immune system by measuring their Fc(gamma) receptor binding activity according to the procedure of Example 15. Table 19 compares the Fc(gamma) receptor binding of these insulin-Fc fusion proteins with the Fc(gamma) receptor binding of the insulin-Fc fusion protein of SEQ ID NO: 36. It can be seen that the non-glycosylated insulin-Fc fusion proteins (achieved through a cNg-S mutation) exhibited the lowest Fc(gamma) receptor binding ratio to SEQ ID NO: 36.

**Table 19: Fc(gamma) receptor binding for cNg variations of SEQ ID NO: 36**

| SEQ ID NO:    | Species / Fc Isotype | Glycosylation Mutation | OD450nm at a Fc(gamma)RI concentration of 3000 (ng/mL) | OD450nm Minus Assay Background | Ratio to SEQ ID NO: 36 |
|---------------|----------------------|------------------------|--------------------------------------------------------|--------------------------------|------------------------|
| SEQ ID NO: 36 | Canine / IgGB        | Native cNg             | 0.428                                                  | 0.371                          | 1.00                   |

(continued)

**Table 19: Fc(gamma) receptor binding for cNg variations of SEQ ID NO: 36**

| SEQ ID NO:    | Species / Fc Isotype | Glycosylation Mutation | OD450nm at a Fc(gamma)RI concentration of 3000 (ng/mL) | OD450nm Minus Assay Background | Ratio to SEQ ID NO: 36 |
|---------------|----------------------|------------------------|--------------------------------------------------------|--------------------------------|------------------------|
| SEQ ID NO: 26 | Canine / IgGB        | Native cNg             | 0.368                                                  | 0.311                          | 0.84                   |
| SEQ ID NO: 58 | Canine / IgGA        | Native cNg             | 0.253                                                  | 0.196                          | 0.53                   |
| SEQ ID NO: 62 | Canine / IgGB        | cNg-S                  | 0.175                                                  | 0.118                          | 0.32                   |
| SEQ ID NO: 61 | Canine / IgGB        | cNg-S and NB119-A      | 0.166                                                  | 0.109                          | 0.29                   |
| SEQ ID NO: 30 | Canine / IgGB        | cNg-S and B16A         | 0.177                                                  | 0.120                          | 0.32                   |

*Comparative Example 47: Exemplary CHO-Based Production Runs Using Preferred Insulin-Fc Fusion Proteins Comprising Fc Fragments of Canine IgGB Origin Made via Stably Transfected CHO Cell Lines.*

**[0268]** Separate CHO cell lines stably transfected with vectors encoding for SEQ ID NO: 26, or SEQ ID NO: 30 were constructed as described in Example 2. Fed-batch shake flask 14-day production runs (0.5-2.0 L media scale) were seeded at 0.5 million cells/mL in an incubator-shaker set at 37°C and 5% carbon dioxide, and the runs were conducted as described in Example 2 above, except that CD OptiCHO was substituted for Dynamis as the growth media (ThermoFisher) and Efficient Feed C (ThermoFisher) was used as the feed. Feed was added at 3% v/v starting on production run day 3, and on day 4, the shake-flask temperature was adjusted to 32°C and the incubator-shaker carbon dioxide concentration was lowered from 5% to 2%. During the run, the cells increased to between 8-14 million cells/mL, and on Day 14 the production run was harvested to remove the cells and the culture supernatant was purified and tested to obtain the insulin-Fc fusion protein as described in Example 4, Example 6, Example 8, and Example 10. Table 20 describes the manufacturing data obtained from the production runs with stably transfected CHO cell lines.

**Table 20: Homodimer titers for non-glycosylated insulin-Fc fusion proteins of SEQ ID NO: 26 and SEQ ID NO: 30**

| SEQ ID NO:    | Protein Yield (mg/L) | %Homodimer | Homodimer Titer (mg/L) |
|---------------|----------------------|------------|------------------------|
| SEQ ID NO: 26 | 485                  | 99.3%      | 482                    |
| SEQ ID NO: 30 | 260                  | 99.0%      | 257                    |

*Comparative Example 48: Exemplary CHO-Based Production Runs Using Preferred Insulin-Fc Fusion Proteins Comprising Fc Fragments of Canine IgGB Origin Made via Stably Transfected CHO Cell Lines.*

**[0269]** A CHO cell line stably transfected with vectors encoding for SEQ ID NO: 28 is constructed as described in Example 2. Fed-batch shake flask 14-day production runs (0.5-2.0 L media scale) is seeded at 0.5 million cells/mL in an incubator-shaker set at 37°C and 5% carbon dioxide, and the run is conducted as described in Example 2, except that CD OptiCHO is substituted for Dynamis as the growth media (ThermoFisher) and Efficient Feed C (ThermoFisher) is used as the feed. Feed is added at 3% v/v starting on production run day 3, and on day 4, the shake-flask temperature is adjusted to 32°C and the incubator-shaker carbon dioxide concentration is lowered from 5% to 2%. On Day 14, the production run is harvested to remove the cells, and the culture supernatant is purified and tested to obtain the insulin-Fc fusion protein as described in Example 4, Example 6, Example 8, and Example 10. The resulting production run is expected to give a protein yield of greater than 200 mg/L, greater than 95% homodimer, and greater than 190 mg/L homodimer titer of SEQ ID NO: 28.

**RESULTS - INSULIN-Fc FUSION PROTEINS COMPRISING A HUMAN Fc FRAGMENT** *Comparative Example 49:*  
*Insulin-Fc Fusion Proteins Comprising Fc Fragment of the Human Fc IgG1 and IgG2 Isotypes.*

**[0270]** An attempt was made to produce an insulin-Fc fusion protein comprising the insulin polypeptide sequence of SEQ ID NO: 7 and the Fc fragment of the human IgG2 isotype (SEQ ID NO: 74) using the peptide linker of SEQ ID NO: 13. The full amino acid sequence for the resulting insulin-Fc fusion protein is as follows:

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFN  
 WYVDGVEVHNAKTKPREEQFNSTFRVVSFLTIVHQQDWLNGKEYKCKVSNKGLPAPIEKTISKTK  
 GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFF  
 LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 75).

**[0271]** A further attempt was made to produce an insulin-Fc fusion protein comprising the insulin polypeptide sequence of SEQ ID NO: 7 and the Fc fragment of the human IgG1 isotype (SEQ ID NO: 73) using the peptide linker of SEQ ID NO: 13. The full amino acid sequence for the resulting insulin-Fc fusion protein is as follows:

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
 EVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSFLTIVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD  
 SGGFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 76).

**[0272]** The insulin-Fc fusion protein configurations of SEQ ID NO: 75 and SEQ ID NO: 76 were synthesized in HEK cells according to Example 1 and purified according to Example 4. The total protein yields gave average homodimer titers as shown in Table 21 after two separate synthesis and Protein A purification steps for each insulin-Fc fusion protein configuration. The structures of the insulin-Fc fusion protein configurations were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS with glycan removal according to Example 8. The %homodimer was measured by size-exclusion chromatography according to Example 10 and determined as shown in Table 21 below. In summary, manufacturing of the insulin-Fc fusion protein of SEQ ID NO: 76 in HEK cells resulted in acceptable aggregate and homodimer titer levels, which meet the design goal of a homodimer titer of greater than 150 mg/L, while the manufacturing of the insulin-Fc fusion protein of SEQ ID NO: 75 in HEK cells did not meet the homodimer titer design goal.

**[0273]** Next, the insulin-Fc fusion protein of SEQ ID NO: 76 was evaluated for the potential for in vivo bioactivity. First, the IR binding of the insulin-Fc fusion protein of SEQ ID NO: 76 was measured according to Example 12, resulting in an IC<sub>50</sub> value of 1799 nM indicating that the compound is likely to be bioactive in vivo (i.e., IC<sub>50</sub> less than 2400 nM). The FcRn assay EC<sub>50</sub> value was 578 ng/mL, indicating a strong likelihood of having a prolonged in vivo bioactivity profile when measured in a canine screening study (e.g., as described in Example 28) or in a human diabetic patient.

**[0274]** Next, the propensity for unwanted immunogenic potential was screened by measuring each compound's in vitro binding to the Fc(γ)RI and C1q by the methods of Example 14 and Example 16. The insulin-Fc fusion protein of SEQ ID NO: 75 demonstrated Fc(γ)RI and C1q assay OD<sub>450</sub> Ratios (where the reference insulin-Fc fusion protein in the OD<sub>450</sub> Ratio is SEQ ID NO: 76) of 0.045 and 0.309 respectively, indicating that the homodimer of SEQ ID NO: 75 did meet the design goals (Example 29) for Fc(γ)RI binding (assay OD<sub>450</sub> Ratio less than 0.50) and the C1q binding (assay OD<sub>450</sub> Ratio less than 0.35). However, as described above, the yield of the insulin-Fc fusion protein of SEQ ID NO: 75 in HEK cells did not meet the homodimer titer design goal.

| Table 21: Homodimer titers and receptor binding data for sequences utilizing an Fc fragment of IgG1 or IgG2 isotype                                                                                     |                    |                                             |             |                        |                    |                                |                                                                             |                                                             |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|---------------------------------------------|-------------|------------------------|--------------------|--------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------|
| Candidate ID                                                                                                                                                                                            | Species/Fe Isotype | Transiently Transfected HEK293 Yield (mg/L) | %Homo-dimer | Homodimer Titer (mg/L) | IR Assay IC50 (nM) | Human FeRn Assay EC50, (ng/mL) | Human Fc(gamma)RI Assay OD450 Ratio at 3,000 ng/mL biotinylated-Fc(gamma)RI | Human C1q Assay OD450 Ratio at 1,000 ng/mL biotinylated-C1q |
| SEQ ID NO: 75                                                                                                                                                                                           | Human/IgG2         | 118 <sup>#</sup>                            | 99.5        | 117 <sup>#</sup>       | 1726               | 462                            | 0.045                                                                       | 0.309                                                       |
| SEQ ID NO: 76                                                                                                                                                                                           | Human/IgG 1        | 181 <sup>#</sup>                            | 99.5        | 180 <sup>#</sup>       | 1799               | 578                            | 1.000 <sup>*</sup>                                                          | 1.000 <sup>*</sup>                                          |
| <sup>#</sup> = averaged from two independent experiments.<br><sup>^</sup> = averaged from three independent experiments.<br><sup>*</sup> = ratio of reference insulin-Fc fusion protein against itself. |                    |                                             |             |                        |                    |                                |                                                                             |                                                             |

Comparative Example 50: cNg Mutation of the Fc Fragment Region of an Insulin-Fc Fusion Protein Comprising the Human IgG1 Isotype.

**[0275]** In an attempt to maintain %homodimer content, IR binding affinity, and acceptable binding affinity to the FcRn receptor of the insulin-Fc fusion protein of SEQ ID NO: 76, a cNg-S mutation was inserted into the Fc fragment CH3 region to try to decrease binding to Fc(gamma)RI and C1q, resulting in the insulin-Fc fusion protein configuration of SEQ ID NO: 91 as follows:

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
EVKFNWYVDGVEVHNAKTKPREEQY<sup>S</sup>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFS<sup>S</sup>CSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 91).

**[0276]** The non-glycosylated insulin-Fc fusion protein configuration of SEQ ID NO: 91 was synthesized in HEK cells according to Example 1, purified according to Example 4, and tested according to Example 6, Example 8, Example 10, Example 12, Example 14 and Example 16 with results as shown in Table 22, together with the results of the parent glycosylated insulin-Fc fusion protein configuration of SEQ ID NO: 76 showing total protein yields, %homodimer, and homodimer titers. The results show that the cNg-S mutation within the human IgG1 Fc fragment of SEQ ID NO: 76 resulting in the insulin-Fc fusion protein configuration of SEQ ID NO: 91 did result in desirable decreases in Fc(gamma)RI affinity and C1q affinity. The homodimer titer of the insulin-Fc fusion protein configuration of SEQ ID NO: 91 increased significantly relative to the homodimer titer of the insulin-Fc fusion protein configuration of SEQ ID NO: 76. Despite the improvement in homodimer titer, the insulin-Fc fusion protein configuration of SEQ ID NO: 91 gave IR assay IC50 values that were greater than 2400 nM (outside of the design criteria of less than 2400 nM) and substantially higher (i.e., lower IR affinity) than that of SEQ ID NO: 76, indicating that this insulin-Fc fusion protein configuration was not likely to exhibit acceptable potency or meet the design goals (Example 29).

| Table 22: Homodimer titers and receptor binding data for sequences comprising an Fc fragment of IgG1 and cNg mutations |                       |                                             |             |                        |                    |                                |                                                                             |                                                             |  |
|------------------------------------------------------------------------------------------------------------------------|-----------------------|---------------------------------------------|-------------|------------------------|--------------------|--------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------|--|
| Candidate ID                                                                                                           | Fc Isotype / Mutation | Transiently Transfected HEK293 Yield (mg/L) | %Homo-dimer | Homodimer Titer (mg/L) | IR Assay IC50 (nM) | Human FeRn Assay EC50, (ng/mL) | Human Fc(gamma)RI Assay OD450 Ratio at 3,000 ng/mL biotinylated-Fc(gamma)RI | Human C1q Assay OD450 Ratio at 1,000 ng/mL biotinylated-C1q |  |
| SEQ ID NO: 76                                                                                                          | hIgG1 / cNg-NB 155    | 181 <sup>#</sup>                            | 99.5        | 180                    | 1799               | 578                            | 1.000                                                                       | 1.000                                                       |  |
| SEQ ID NO: 91                                                                                                          | hIgG1 / cNg-NB155-S   | 193                                         | 98.1        | 189                    | 2937               | 943                            | 0.199                                                                       | 0.102                                                       |  |
| # = averaged from two independent experiments.<br>DNT = did not test.                                                  |                       |                                             |             |                        |                    |                                |                                                                             |                                                             |  |

Example 51: Human Insulin-Fc Fusion Protein Using Mutated Insulin Polypeptide (SEQ ID NO: 10), Linker (SEQ ID NO: 13), and Fc IgG1 Isotype with cNg-S Mutation.

**[0277]** As described above, a cNg-S, IgG1 isotype Fc fragment mutation produced an insulin-Fc fusion protein configuration with significantly reduced affinity for the Fc(gamma)RI and C1q, but this insulin-Fc fusion protein configuration has IR receptor affinity that is unacceptably low (i.e., IC50 value greater than 2400 nM). Therefore, a mutation to the insulin polypeptide was made and the resulting materials was screened against the IR IC50 assay in an effort to see if the mutation led to improved IR affinity. A switch from tyrosine to alanine was made at the 16<sup>th</sup> amino acid from the N-terminus of the B-chain (i.e., B16A) to give the insulin polypeptide of SEQ ID NO: 10. The mutated insulin polypeptide of SEQ ID NO: 10 and peptide linker SEQ ID NO: 13 were used with the cNg mutated Fc fragment of SEQ ID NO: 77 with X<sub>1</sub> is S, resulting in the insulin-Fc fusion protein configuration of SEQ ID NO: 78. The cNg-S mutation in the insulin-Fc fusion protein configuration was made to the Fc fragment region to reduce the binding affinity of the Fc fragment for Fc(gamma) receptors in vivo, as measured by binding in an in vitro human Fc(gamma)RI assay described in Example 15. The position of the cNg-S site in the insulin-Fc fusion protein of SEQ ID NO: 78 is cNg-NB155, comprising a mutation at the cNg site from "N" to "S".

**[0278]** The insulin-Fc fusion protein configuration of SEQ ID NO: 78 was manufactured in HEK293 cells according to Example 1 and purified using a Protein A column according to Example 4. The structure of the insulin-Fc fusion protein of SEQ ID NO: 78 was confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequence was further identified by LC-MS without glycan removal according to Example 8. The compound MW of SEQ ID NO: 78 was assessed by the LC-MS method of Example 8 without enzymatic deglycosylation. The target mass of SEQ ID NO: 78 is 65046.9 Da and the found mass was 65046.7 Da, which confirmed the compound amino acid composition and homodimer structure of SEQ ID NO: 78 were correct.

**[0279]** The %homodimer was measured by size-exclusion chromatography according to Example 10. The full amino acid sequence of the cNg-mutated insulin-Fc fusion proteins is listed below (with the NB155 position underlined):

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
GGGGQGGGGQGGGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHED  
PEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 78).

**[0280]** The resulting total protein yield, %homodimer, and homodimer titer of the insulin-Fc fusion protein configuration of SEQ ID NO: 78 is given in Table 23 and compared to that of SEQ ID NO: 76 and SEQ ID NO: 91. Including the B 16A mutation in the insulin polypeptide sequence with the non-glycosylated cNg-S mutated insulin-Fc fusion protein improved the homodimer titer relative to SEQ ID NO: 76. In vitro IR binding for the insulin-Fc fusion protein of SEQ ID NO: 78 was tested according to the procedure of Example 12. The inclusion of the B16A mutation into the insulin polypeptide of SEQ ID NO: 10 in the insulin-Fc fusion protein of SEQ ID NO: 78 led to marked improvements in IR affinity (i.e., lower IR IC50 assay values) compared to SEQ ID NO: 91 in which the insulin polypeptide contained the B16E mutation (SEQ ID NO: 7). This is especially surprising, given that this B16A mutation in a different insulin-Fc fusion protein (SEQ ID NO: 3 in US10,597,435) gave IR IC50 assay values of greater than 5000 nM and showed no glucose-lowering bioactivity in vivo. The sequence alignment of SEQ ID NO: 76, SEQ ID NO: 78 and SEQ ID NO: 91 highlighting the differences in amino acid sequences is shown in FIG. 27. "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

**Table 23: Homodimer titers and receptor binding data for sequences comprising an Fc fragment of IgG1, cNg mutations and B16A mutation.**

| Candidate ID                | Fc Isotype / Mutation       | Transiently Transfected HEK293 Yield (mg/L) | %Homo-dimer | Homodimer Titer (mg/L) | IR Assay IC50 (nM) | Human FeRn Assay EC50, (ng/mL) | Human Fc(gamma)RI Assay OD450 Ratio at 3,000 ng/mL biotinylated-Fc(gamma)RI | Human C1q Assay OD450 Ratio at 1,000 ng/mL biotinylated-C1q |
|-----------------------------|-----------------------------|---------------------------------------------|-------------|------------------------|--------------------|--------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------|
| SEQ ID NO: 76 (Not claimed) | hIgG1 / cNg-NB 155          | 181 <sup>#</sup>                            | 99.5        | 180                    | 1799               | 578                            | 1.000                                                                       | 1.000                                                       |
| SEQ ID NO: 91 (not claimed) | hIgG1 / cNg-NB155-S         | 193                                         | 98.1        | 189                    | 2937               | 943                            | 0.199                                                                       | 0.102                                                       |
| SEQ ID NO: 78               | hIgG1 / B16A / cNg-NB 155-S | 189                                         | 93.2        | 176                    | 1162               | 825                            | 0.197                                                                       | 0.169                                                       |

**[0281]** Furthermore, inclusion of the B16A mutation in the insulin polypeptide (resulting in SEQ ID NO: 10) of the insulin-Fc fusion protein of SEQ ID NO: 78 led to acceptable FcRn binding as measured by the procedure in Example 19, indicating that this insulin-Fc fusion protein will likely exhibit prolonged in vivo pharmacokinetic half-life and demonstrate a prolonged glucose-lowering bioactivity profile in vivo. It was noted that the FcRn binding affinity for SEQ ID NO: 78 is greater (as demonstrated by a lower EC50 measurement in the FcRn binding assay) than that of the similar insulin-Fc fusion protein not containing the B 16A mutation (SEQ ID NO: 91), further highlighting that the inclusion of the B16A mutation unexpectedly impacts FcRn receptor affinity in addition to IR receptor affinity. It is thought that this incremental increases in FcRn receptor affinity may be desirable, since higher FcRn affinities often correlate with longer therapeutic pharmacokinetic half-lives in vivo.

**[0282]** The Fc(gamma)RI and C1q binding properties were measured by the methods of Example 14 and Example 16 for the insulin-Fc fusion protein of SEQ ID NO: 78 and were shown to be acceptable. The C1q binding affinity was also much lower for the insulin-Fc fusion protein of SEQ ID NO: 78, showing greater than 80% reduction in C1q binding relative to the insulin-Fc fusion protein of SEQ ID NO: 76. Thus, the insulin-Fc fusion protein of SEQ ID NO: 78 met the design goals (Example 29) for manufacturability, in vivo potency, prolonged in vivo bioactivity, and a low potential for immunogenicity.

Comparative Example 52: Alternate cNg Mutations of the Fc Fragment Region of Insulin-Fc Fusion Proteins Comprising the Human IgG1 Isotype.

**[0283]** In further attempt to create insulin-Fc fusion protein configurations with prolonged in vivo bioactivity and a low potential for immunogenicity while maintaining similar %homodimer content and IR binding affinity to the insulin-Fc fusion protein of SEQ ID NO: 76, and acceptable binding affinity to the FcRn receptor, alternate cNg mutations were inserted into the Fc fragment CH3 region to try to decrease binding to Fc(gamma)RI and C1q, resulting in SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, and SEQ ID NO: 95. The insulin-Fc fusion protein embodiments were synthesized in HEK cells according to Example 1, purified according to Example 4, and tested according to Example 6, Example 8, Example 10, Example 12, Example 14 and Example 16 with results as shown in Table 24. The sequence alignment of SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, and SEQ ID NO: 95 against SEQ ID NO: 75, SEQ ID NO: 76 and SEQ ID NO: 91 highlighting the differences in amino acid sequences are shown in FIG. 28 (Clustal Omega). "\*" represents complete homology across all sequences at a given sequence position, while ":", "." or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

**[0284]** The full amino acid sequences for the insulin-Fc fusion protein embodiments are shown below, with the cNg mutations underlined.

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
EVKFNWYVDGVEVHNAKTKPREEQYDSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV  
FSCSMHEALHNHYTQKSLSLSPG (SEQ ID NO: 92) (not claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP  
EVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNV  
FSCSMHEALHNHYTQKSLSLSPG (SEQ ID NO: 93) (not claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP  
 5 EVKFNWYVDGVEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDS  
 DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 94) (not  
 10 claimed)

FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQG  
 GGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP  
 15 EVKFNWYVDGVEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK  
 TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDS  
 DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 95) (not  
 20 claimed).

**[0285]** The insulin-Fc fusion protein embodiments above comprising the human IgG1 Fc fragment of SEQ ID NO: 77  
 where X1 is D, A, R, and Q, are listed in Table 24 along with SEQ ID NO: 76, illustrating the corresponding total protein  
 25 yields, %homodimer, and homodimer titers. The results show that the various mutations to the cNg site contained within  
 the human IgG1 Fc fragment for SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94 and SEQ ID NO: 95 did result in  
 desirable decreases in Fc(gamma)RI affinity and C1q affinity. The homodimer titer of SEQ ID NO: 92, SEQ ID NO: 93,  
 and SEQ ID NO: 94 increased significantly relative to SEQ ID NO: 76. The homodimer titer of SEQ ID NO: 95 was too  
 30 low to meet design goals (Example 29) and so insulin receptor binding was not assessed. Despite the improvements in  
 homodimer titer, SEQ ID NO: 92, SEQ ID NO: 93, and SEQ ID NO: 94 gave IR assay IC50 values that were greater  
 than 2400 nM in all cases and substantially higher (i.e., lower IR affinity) than that of SEQ ID NO: 76. The IR assay IC50  
 values in all cases were greater than the design criterion of less than 2400 nM, indicating that these insulin-Fc fusion  
 protein embodiments were not likely to exhibit acceptable potency and cost of goods. None of SEQ ID NO: 91, SEQ ID  
 35 NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, and SEQ ID NO: 95 meet the design goals (Example 29).

| Table 24: Homodimer titers and receptor binding data for sequences comprising an Fc fragment of IgG1 and cNg mutations |                       |                                             |             |                        |                    |                                |                                                                             |                                                             |  |
|------------------------------------------------------------------------------------------------------------------------|-----------------------|---------------------------------------------|-------------|------------------------|--------------------|--------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------|--|
| Candidate ID                                                                                                           | Fc Isotype / Mutation | Transiently Transfected HEK293 Yield (mg/L) | %Homo-dimer | Homodimer Titer (mg/L) | IR Assay IC50 (nM) | Human FeRn Assay EC50, (ng/mL) | Human Fc(gamma)RI Assay OD450 Ratio at 3,000 ng/mL biotinylated-Fc(gamma)RI | Human C1q Assay OD450 Ratio at 1,000 ng/mL biotinylated-C1q |  |
| SEQ ID NO: 76 (not claimed)                                                                                            | hIgG1 / cNg-NB 155    | 181#                                        | 99.5        | 180                    | 1799               | 578                            | 1.000                                                                       | 1.000                                                       |  |
| SEQ ID NO: 92 (not claimed)                                                                                            | hIgG1 / cNg-NB 155-D  | 243                                         | 97.8        | 238                    | 2438               | 883                            | 0.379                                                                       | 0.116                                                       |  |
| SEQ ID NO: 93 (not claimed)                                                                                            | hIgG1 / cNg-NB 155-A  | 228                                         | 96.7        | 220                    | 2852               | 900                            | 0.245                                                                       | 0.151                                                       |  |
| SEQ ID NO: 94 (not claimed)                                                                                            | hIgG1 / cNg-NB 155-R  | 219                                         | 94.7        | 207                    | 3767               | 913                            | 0.150                                                                       | 0.127                                                       |  |
| SEQ ID NO: 95 (not claimed)                                                                                            | hIgG1 / cNg-NB 155-Q  | 139                                         | 97.6        | 136                    | DNT                | 775                            | 0.192                                                                       | 0.222                                                       |  |
| # = averaged from two independent experiments.<br>DNT = did not test.                                                  |                       |                                             |             |                        |                    |                                |                                                                             |                                                             |  |

*Example 53: Human Insulin-Fc Fusion Proteins Using Insulin Polypeptide (SEQ ID NO: 10), Linker (SEQ ID NO: 13), and Fc IgG1 Isotypes Comprising Various cNg Mutations.*

**[0286]** As described above, cNg-site, IgG1 isotype Fc fragment mutations produce insulin-Fc fusion protein with significantly reduced affinities for the Fc(gamma)RI and C1q, but these compounds have IR receptor affinities that are unacceptably low (i.e., IC50 values greater than 2400 nM). Therefore, the same mutation that was made to the insulin polypeptide in SEQ ID NO: 78 (a switch from tyrosine to alanine was made at the 16<sup>th</sup> amino acid from the N-terminus of the B-chain (i.e., B16A) to give the insulin polypeptide of SEQ ID NO: 10) was made and the resulting insulin-Fc fusion protein configurations were screened against the IR IC50 assay in an effort to see if any of the mutated insulin polypeptides led to improved IR affinity. The mutated insulin polypeptide of SEQ ID NO: 10 and peptide linker SEQ ID NO: 13 were used to make the insulin-Fc fusion protein configurations of SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, and SEQ ID NO: 86 as shown below. Each of these insulin-Fc fusion protein configurations comprise cNg site mutations that were made to the Fc fragment region to reduce the binding affinity of the Fc fragment for Fc(gamma) receptors in vivo, as measured by binding in an in vitro human Fc(gamma)RI assay described in Example 15. The position of the cNg site in the insulin-Fc fusion protein configurations of SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, and SEQ ID NO: 86 is cNg-NB155. The insulin-Fc fusion protein of SEQ ID NO: 80 comprises a mutation at the cNg site from "N" to "D". The insulin-Fc fusion protein of SEQ ID NO: 82 comprises a mutation at the cNg site from "N" to "A". The insulin-Fc fusion protein of SEQ ID NO: 84 comprises a mutation at the cNg site from "N" to "R". The insulin-Fc fusion protein of SEQ ID NO: 86 comprises a mutation at the cNg site from "N" to "Q".

**[0287]** The proteins were manufactured in HEK293 cells according to Example 1. Surprisingly, the homodimer titer for the insulin-Fc fusion protein configuration of SEQ ID NO: 86 comprising the cNg-NB155-Q mutation together with the B16A mutation was significantly lower than required to meet the design goal for homodimer titer. The insulin-Fc fusion protein configurations of SEQ ID NO: 80, SEQ ID NO: 82, and SEQ ID NO: 84 were then purified using a Protein A column according to Example 4. The structures of the insulin-Fc fusion protein configurations were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS without glycan removal according to Example 8. The compound MW of the insulin-Fc fusion protein of SEQ ID NO: 80 was assessed by the LC-MS method of Example 8 without enzymatic deglycosylation. The target mass of SEQ ID NO: 80 is 65102.9 Da and the found mass was 65102.8 Da, which confirmed the compound amino acid composition and homodimer structure of SEQ ID NO: 80 were correct. The compound MW of the insulin-Fc fusion protein of SEQ ID NO: 82 was assessed by the LC-MS method of Example 8 without enzymatic deglycosylation. The target mass of SEQ ID NO: 82 is 65014.9 Da and the found mass was 65014.9 Da, which confirmed the compound amino acid composition and homodimer structure of SEQ ID NO: 82 were correct. The compound MW of the insulin-Fc fusion protein of SEQ ID NO: 84 was assessed by the LC-MS method of Example 8 without enzymatic deglycosylation. The target mass of SEQ ID NO: 84 is 65185.1 Da and the found mass was 65184.8 Da, which confirmed the compound amino acid composition and homodimer structure of SEQ ID NO: 84 were correct.

**[0288]** The %homodimer was measured by size-exclusion chromatography according to Example 10. The full amino acid sequences of the cNg-mutated insulin-Fc fusion protein configurations are listed below (SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, and SEQ ID NO: 86 with the NB155 position underlined) and the resulting sequence alignments are shown against SEQ ID NO: 76 and SEQ ID NO: 78 in FIG. 29 (Clustal Omega). "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FVNQHLCGSHLVEALALVCGERGFHYGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQGGGGQG  
GGGQGGGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDG  
VEVHNAKTKPREEQYDSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL  
PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV  
FSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 80) (not claimed)

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQGGGGQG  
 GGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDG  
 5 VEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL  
 PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV  
 FSCSVMEALHNHYTQKSLSLSPG (SEQ ID NO: 82) (not claimed)

10 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQGGGGQG  
 GGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDG  
 VEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL  
 15 PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV  
 FSCSVMEALHNHYTQKSLSLSPG (SEQ ID NO: 84)

20 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGGGGQGGGGQG  
 GGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDG  
 VEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL  
 25 PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV  
 FSCSVMEALHNHYTQKSLSLSPG (SEQ ID NO: 86) (not claimed).

[0289] The resulting total protein yields, %homodimer, and homodimer titers are given in Table 25. With the exception  
 of SEQ ID NO: 86 as described above, including the B16A mutation in the insulin polypeptide sequence with the non-  
 glycosylated cNg mutated insulin-Fc fusion protein embodiments improved the homodimer titers relative to SEQ ID NO:  
 76. The cNg-NB155-D composition exhibited the highest homodimer titer at 222 mg/L. In vitro IR binding for the insulin-  
 Fc fusion proteins was tested according to the procedure of Example 12. As was the case with SEQ ID NO: 78, the  
 inclusion of the B16A mutation into the insulin polypeptides of the insulin-Fc fusion protein embodiments of SEQ ID NO:  
 80 (not claimed), SEQ ID NO: 82 (not claimed), and SEQ ID NO: 84 led to marked improvements in IR affinity (i.e., lower  
 IR IC50 assay values) compared to the insulin-Fc fusion protein configurations of SEQ ID NO: 92, SEQ ID NO: 93, and  
 SEQ ID NO: 94 (not claimed) in which the insulin polypeptide contained the B16E mutation.

[0290] Furthermore, inclusion of the B 16A mutation in the insulin polypeptides of the insulin-Fc fusion proteins of SEQ  
 ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), and SEQ ID NO: 84 led to acceptable FcRn binding for these  
 insulin-Fc fusion protein embodiments as measured by the procedure in Example 19, indicating that these compounds  
 will likely exhibit prolonged in vivo pharmacokinetic half-lives and demonstrate prolonged glucose-lowering bioactivity  
 profiles in vivo. It was noted that the FcRn binding affinities for the insulin-Fc fusion protein configurations containing  
 the B16A mutation (SEQ ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), and SEQ ID NO: 84) were significantly  
 greater (e.g. lower EC50's in the FcRn binding assay) than those of the similar insulin-Fc fusion protein configurations  
 not containing the B16A mutation (SEQ ID NO: 92, SEQ ID NO: 93, and SEQ ID NO: 94 (not claimed)), further highlighting  
 that the inclusion of the B16A mutation unexpectedly impacts FcRn receptor affinity in addition to IR receptor affinity. It  
 is thought that these incremental increases FcRn receptor affinity may be desirable, since higher FcRn affinities often  
 correlate with longer therapeutic pharmacokinetic half-lives in vivo.

[0291] The Fc(gamma)RI and C1q binding properties were measured by the methods of Example 14 and Example  
 16 for the insulin-Fc fusion protein configurations of SEQ ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), and  
 SEQ ID NO: 84, and were shown to be acceptable. Additionally, the Fc(gamma)RI binding affinities decreased very  
 significantly with the cNg-NB155-R (SEQ ID NO: 84) mutation, demonstrating greater than 80% reduction in Fc(gamma)RI  
 binding relative to the insulin-Fc fusion protein of SEQ ID NO: 76. The C1q binding affinities were also much lower for  
 the insulin-Fc fusion proteins of SEQ ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), and SEQ ID NO: 84, with  
 all of these insulin-Fc fusion protein embodiments showing greater than 80% reduction in C1q binding relative to the  
 insulin-Fc fusion protein of SEQ ID NO: 76. Thus, the insulin-Fc fusion proteins of SEQ ID NO: 80 (not claimed), SEQ  
 ID NO: 82 (not claimed), and SEQ ID NO: 84 met the design goals (Example 29) for manufacturability, in vivo potency,  
 prolonged in vivo bioactivity, and a low potential for immunogenicity.

| Table 25: Homodimer titers and receptor binding data for sequences comprising an Fc fragment of IgG1 |                             |                                             |            |                        |                    |                                |                                                                             |                                                             |  |
|------------------------------------------------------------------------------------------------------|-----------------------------|---------------------------------------------|------------|------------------------|--------------------|--------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------|--|
| Candidate ID                                                                                         | Fc Isotype / Mutation       | Transiently Transfected HEK293 Yield (mg/L) | %Homodimer | Homodimer Titer (mg/L) | IR Assay IC50 (nM) | Human FeRn Assay EC50, (ng/mL) | Human Fc(gamma)RI Assay OD450 Ratio at 3,000 ng/mL biotinylated-Fc(gamma)RI | Human C1q Assay OD450 Ratio at 1,000 ng/mL biotinylated-C1q |  |
| SEQ ID NO: 76 (not claimed)                                                                          | hIgG1 / cNg-NB 155          | 181#                                        | 99.5       | 180                    | 1799               | 578                            | 1.000                                                                       | 1.000                                                       |  |
| SEQ ID NO: 80 (not claimed)                                                                          | hIgG1 / B16A / cNg-NB155-D  | 234                                         | 94.9       | 222                    | 881                | 788                            | 0.346                                                                       | 0.109                                                       |  |
| SEQ ID NO: 82 (not claimed)                                                                          | hIgG1 / B16A / cNg-NB 155-A | 227                                         | 93.1       | 211                    | 1125               | 690                            | 0.267                                                                       | 0.181                                                       |  |
| SEQ ID NO: 84                                                                                        | hIgG1 / B16A / cNg-NB 155-R | 193                                         | 96.1       | 185                    | 911                | 696                            | 0.167                                                                       | 0.173                                                       |  |
| SEQ ID NO: 86 (not claimed)                                                                          | hIgG1 / B16A / cNg-NB 155-Q | 114                                         | 97.1       | 111                    | DNT                | 750                            | 0.192                                                                       | 0.311                                                       |  |
| # = averaged from two independent experiments.<br>DNT = did not test.                                |                             |                                             |            |                        |                    |                                |                                                                             |                                                             |  |

Example 54: Human Insulin-Fc Fusion Proteins with an Insulin Polypeptide SEQ ID NO: 10, the Fc IgG1 Isotype Comprising the cNg-S Mutation, and Various Linkers.

[0292] As described above, the IgG1 isotype Fc fragment mutations at the cNg site and mutations on the insulin polypeptide led to unexpected changes in the material properties, including for the IR, FcRn, Fc(gamma)RI receptor, and C1q binding. To better understand how the linker composition impacted these properties, a particular insulin-Fc fusion protein configuration (SEQ ID NO: 78) containing the B 16A-mutated insulin polypeptide of SEQ ID NO: 10, the non-glycosylated hlgG1 Fc fragment with the cNg-S mutation (SEQ ID NO: 77 where X<sub>1</sub> is S), and the linker sequence of SEQ ID NO: 13 was chosen as the starting point from which the linker length and composition were modified as shown in Table 26 below.

| Table 26: Sequence ID NOs and linker amino acid compositions, and position of cNg-S site |                                            |                          |
|------------------------------------------------------------------------------------------|--------------------------------------------|--------------------------|
| Candidate ID                                                                             | Linker Amino Acid Sequence                 | Position of the cNg Site |
| SEQ ID NO: 96                                                                            | GGGGGQGGGGQGGGGQGGGGGQGGGG (SEQ ID NO: 99) | cNg-NB 160-S             |
| SEQ ID NO: 78                                                                            | GGGGGQGGGGQGGGGQGGGGG (SEQ ID NO: 13)      | cNg-NB155-S              |
| SEQ ID NO: 97 (not claimed)                                                              | GGGGQGGGG (SEQ ID NO: 19)                  | cNg-NB143-S              |
| SEQ ID NO: 98 (not claimed)                                                              | <Linker Absent>                            | cNg-NB 136-S             |
| SEQ ID NO: 87                                                                            | GGGGGAGGGGAGGGGAGGGGG (SEQ ID NO: 67)      | cNg-NB155-S              |
| SEQ ID NO: 89 (not claimed)                                                              | GGGGAGGGG (SEQ ID NO: 11)                  | cNg-NB143-S              |

[0293] The insulin-Fc fusion protein embodiments of SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 87, and SEQ ID NO: 89 were manufactured in HEK293 cells according to Example 1. The insulin-Fc fusion protein configurations were then purified using a Protein A column according to Example 4. The structures of the insulin-Fc fusion protein configurations were confirmed according to Example 6 by non-reducing and reducing CE-SDS, and the sequences were further identified by LC-MS without glycan removal according to Example 8. Verification of the lack of glycan was performed using the LC-MS method of Example 8, but with omission of the PNGase F treatment step. The compound MW of the insulin-Fc fusion protein configuration of SEQ ID NO: 87 was assessed by the LC-MS method of Example 8 without enzymatic deglycosylation. The target mass is 64704.6 Da and the found mass was 64704.4 Da, which confirmed the compound amino acid composition and homodimer structure of SEQ ID NO: 87 were correct. The compound MW of the insulin-Fc fusion protein configuration of SEQ ID NO: 89 was assessed by the LC-MS method of Example 8 without enzymatic deglycosylation. The target mass is 63279.2 Da and the found mass was 63278.8 Da, which confirmed the compound amino acid composition and homodimer structure of SEQ ID NO: 89 were correct.

[0294] The %homodimer was measured by size-exclusion chromatography according to Example 10. In vitro IM-9 IR binding IC<sub>50</sub> was measured as described in Example 12, Fc(gamma) receptor binding affinity was measured using an in vitro human Fc(gamma)RI assay as described in Example 15, C1q binding affinity was measured as described in Example 16, and affinity for the human FcRn receptor was measured as described in Example 19. The results are given in Table 27.

| Table 27: Homodimer titers and receptor binding data for sequences comprising a B16A, Fc fragment of IgG1, cNg-S mutations, and various linker compositions |                                             |            |                        |                    |                                |                                                                             |                                                             |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------|------------|------------------------|--------------------|--------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------|
| Candidate ID                                                                                                                                                | Transiently Transfected HEK293 Titer (mg/L) | %Homodimer | Homodimer Titer (mg/L) | IR Assay IC50 (nM) | Human FeRn Assay EC50, (ng/mL) | Human Fc(gamma)RI Assay OD450 Ratio at 3,000 ng/mL biotinylated-Fc(gamma)RI | Human C1q Assay OD450 Ratio at 1,000 ng/mL biotinylated-C1q |
| SEQ ID NO: 96                                                                                                                                               | 227                                         | 94.2       | 214                    | 760                | 672                            | 0.216                                                                       | 0.166                                                       |
| SEQ ID NO: 78                                                                                                                                               | 189                                         | 93.2       | 176                    | 1162               | 825                            | 0.197                                                                       | 0.169                                                       |
| SEQ ID NO: 97 (not claimed)                                                                                                                                 | 187                                         | 95.5       | 179                    | 954                | 679                            | 0.208                                                                       | 0.314                                                       |
| SEQ ID NO: 98 (not claimed)                                                                                                                                 | 40                                          | 94.8       | 38                     | 526                | 401                            | 0.146                                                                       | 0.311                                                       |
| SEQ ID NO: 87                                                                                                                                               | 224                                         | 95.2       | 213                    | 789                | 701                            | 0.193                                                                       | 0.204                                                       |
| SEQ ID NO: 89 (not claimed)                                                                                                                                 | 193                                         | 94.5       | 182                    | 1265               | 657                            | 0.192                                                                       | 0.305                                                       |

**[0295]** The full amino acid sequences of the resulting insulin-Fc fusion proteins are listed below (with the cNg-S position underlined). FIG. 30 (Clustal Omega) illustrates a side-by-side sequence comparison of SEQ ID NO: 87, SEQ ID NO: 96, SEQ ID NO: 78, SEQ ID NO: 97, SEQ ID NO: 89, and SEQ ID NO: 98. "\*" represents complete homology across all sequences at a given sequence position, while ":", ".", or spaces refer to conservative, moderate, or very different amino acid mutations across the sequences at a given sequence position respectively.

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGA  
GGGGAGGGGAGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED  
PEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD  
SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 87);

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGGQ  
GGGGQGGGGQGGGGGGQGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV  
DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNK  
ALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK  
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO:  
96);

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGGQ  
GGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV  
EVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ  
VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD  
KSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 97) (not claimed);

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCGGGGAG  
GGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV  
EVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ  
VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD  
KSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 89) (not claimed);

FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENYCDKTHTCP  
PCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPR  
EEQYSSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDE  
LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV  
FSCFSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 98) (not claimed).

**[0296]** The results indicate that in the insulin-Fc fusion protein configurations without a linker, wherein the C-terminal end of the insulin polypeptide is directly linked to the N-terminal end of the Fc fragment (SEQ ID NO: 98), the homodimer titer is unacceptably low (i.e., 38 mg/L). In the insulin-Fc fusion protein configurations with the glycine-glutamine linker composition (SEQ ID NO: 97 (not claimed), SEQ ID NO: 78, and SEQ ID NO: 96) comprising at least nine amino acids

in length, there did not appear to be any significant trends in physiochemical properties that varied with linker length. A comparison of the insulin-Fc fusion protein configuration of SEQ ID NO: 87, containing a linker of glycine-alanine composition, with the insulin-Fc fusion protein configuration of SEQ ID NO: 78 containing a linker of glycine-glutamine composition, suggested that for an identical linker length of 21 amino acids, the insulin-Fc fusion protein configuration of SEQ ID NO: 87 demonstrated improved homodimer titer, stronger IR binding, and stronger FcRn binding, while having nearly identical Fc(gamma)RI and C1q binding properties. Therefore, while the insulin-Fc fusion protein configurations of SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 87, and SEQ ID NO: 89 all met the design goals (Example 29), the homodimer of SEQ ID NO: 87 is a preferred embodiment.

**Example 55: *In vivo Efficacy of Insulin-Fc fusion Proteins Comprising the B16A Insulin Polypeptide Mutant, Human IgG1 Isotype Fc Fragment with cNg-NB155-X Site Mutations, and Linker (SEQ ID NO: 13).***

**[0297]** Given the promising homodimer titer, IR activity, and FcRn activity results in Example 53, the insulin-Fc fusion proteins of SEQ ID NO: 78, SEQ ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), and SEQ ID NO: 84 will be tested for in vivo bioactivity according to Example 22 following a subcutaneous injection in each of N=3 healthy, antibody-naive, beagle dogs weighing approximately 10 kg. A plot of %FBGL versus time for a single subcutaneous administration of each insulin-Fc fusion protein of SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, and SEQ ID NO: 84 will be constructed, and the %FBGL vs. time data is expected to demonstrate that the insulin-Fc fusion proteins of SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, and SEQ ID NO: 84 are significantly bioactive in dogs.

**Comparative Example 56: *In vivo Efficacy of Insulin-Fc fusion Proteins Comprising the B16A Insulin Polypeptide Mutant, Human IgG1 Isotype Fc Fragment with cNg-NB155-S Mutation, and Linker (SEQ ID NO: 11).***

**[0298]** Given the promising homodimer titer, IR activity, and FcRn activity results in Example 54, the insulin-Fc fusion protein of SEQ ID NO: 89 (not claimed) will be tested for in vivo bioactivity according to Example 22 following a subcutaneous injection in each of N=3 healthy, antibody-naive, beagle dogs weighing approximately 10 kg. A plot of %FBGL versus time for a single subcutaneous administration of the insulin-Fc fusion protein of SEQ ID NO: 89 will be constructed, and the %FBGL vs. time data is expected to demonstrate that the insulin-Fc fusion protein of SEQ ID NO: 89 is significantly bioactive in dogs.

**Example 57: *In vivo Efficacy of Insulin-Fc fusion Proteins Comprising the B16A Insulin Polypeptide Mutant, Human IgG1 Isotype Fc Fragment with cNg-NB155-S Mutation, and Linker (SEQ ID NO: 67).***

**[0299]** Given the promising homodimer titer, IR activity, and FcRn activity results in Example 54, the insulin-Fc fusion protein of SEQ ID NO: 87 will be tested for in vivo bioactivity according to Example 22 following a subcutaneous injection in each of N=3 healthy, antibody-naive, beagle dogs weighing approximately 10 kg. A plot of %FBGL versus time for a single subcutaneous administration of the insulin-Fc fusion protein of SEQ ID NO: 87 will be constructed, and the %FBGL vs. time data is expected to demonstrate that the insulin-Fc fusion protein of SEQ ID NO: 87 is significantly bioactive in dogs.

**Example 58: *Exemplary CHO-Based Production Runs Using Preferred Insulin-Fc Fusion Proteins Comprising Fc Fragments of Human IgG1 Origin Made via Stably Transfected CHO Cell Lines.***

**[0300]** Separate CHO cell lines stably transfected with vectors encoding for the insulin-Fc fusion protein configurations of SEQ ID NO: 78, SEQ ID NO: 80 (not claimed), SEQ ID NO: 82 (not claimed), SEQ ID NO: 84, SEQ ID NO: 87, and SEQ ID NO: 89 (not claimed) will be constructed as described in Example 2. Fed-batch shake flask 14-day production runs (0.5-2.0 L media scale) will be seeded at 0.5 million cells/mL in an incubator-shaker set at 37°C and 5% carbon dioxide, and the runs will be conducted as described in Example 2, except that CD OptiCHO will be substituted for Dynamis as the growth media (ThermoFisher) and Efficient Feed C (ThermoFisher) will be used as the feed. Feed will be added at 3% v/v starting on production run day 3, and on day 4, the shake-flask temperature will be adjusted to 32°C and the incubator-shaker carbon dioxide concentration will be lowered from 5% to 2%. During the run, the cell densities will increase to between 8-14 million cells/mL, and on Day 14 the production run will be harvested to remove the cells and the culture supernatant purified and tested to obtain the insulin-Fc fusion proteins as described in Example 4, Example 6, Example 8, and Example 10. The manufacturing and in vitro testing data from the separate production runs of the insulin-Fc fusion protein configurations of SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 87 and SEQ ID NO: 89 with stably transfected CHO cell lines are expected to exhibit homodimer titers greater than those obtained in the transiently transfected HEK293 cells, with IR, FcRn receptor, Fc(gamma)RI receptor binding and C1q binding properties that meet the design goals (Example 29).

Example 59: In vivo Pharmacodynamics (PD) After a Single Administration of Human Insulin Fc-Fusion Proteins in Mice.

[0301] Insulin-Fc fusion proteins were assessed for their effects on fasting blood glucose levels as follows. Data was collected from N=3 balb/c mice or diabetic wt NOD mice (Jackson Laboratories) per group. The animals were fasted one hour before the experiment and then at time = 0 hours, the mice received a single subcutaneous administration of a pharmaceutical composition containing an insulin Fc-fusion protein homodimer at a concentration of 300  $\mu$ g/kg of insulin-Fc fusion protein in a solution of between 10-50 mM sodium hydrogen phosphate, 50-150 mM sodium chloride, 0.005-0.05% v/v Tween-80, and optionally a bacteriostat (e.g. phenol, m-cresol, or methylparaben) at a concentration of between 0.02-1.00 mg/mL, at a final solution pH of between 7.0-8.0 adjusted using sodium hydroxide and/or hydrochloric acid.

[0302] For each time point, a sample of blood was collected, and a glucose level reading was immediately determined using a glucose meter (AlphaTRAK<sup>®</sup> 2 pet glucometer), which requires approximately one drop of blood. Average % fasting blood glucose levels (% FBGL) from 0 to 9 hours were plotted to assess the bioactivity of a given insulin-Fc fusion protein configuration. The results shown in FIG. 38 for a single subcutaneous dose of SEQ ID NO: 87 and SEQ ID NO: 89 (not claimed) demonstrate that both compounds showed significant bioactivity in mice.

Example 60: In vivo Pharmacodynamics (PD) After a Single Administration of Human Insulin Fc-Fusion Proteins in Mice.

[0303] Insulin-Fc fusion proteins are assessed for their effects on fasting blood glucose levels as follows. Data is collected from N=3 balb/c mice or diabetic wt NOD mice (Jackson Laboratories) per group. The animals are fasted one hour before the experiment and then at time = 0 hours, the mice receive a single subcutaneous administration of a pharmaceutical composition containing an insulin Fc-fusion protein homodimer at a concentration of 300  $\mu$ g/kg of insulin-Fc fusion protein in a solution of between 10-50 mM sodium hydrogen phosphate, 50-150 mM sodium chloride, 0.005-0.05% v/v Tween-80, and optionally a bacteriostat (e.g. phenol, m-cresol, or methylparaben) at a concentration of between 0.02-1.00 mg/mL, at a final solution pH of between 7.0-8.0 adjusted using sodium hydroxide and/or hydrochloric acid.

[0304] For each time point, a sample of blood is collected, and a glucose level reading is immediately determined using a glucose meter (AlphaTRAK<sup>®</sup> 2 pet glucometer), which requires approximately one drop of blood. Average % fasting blood glucose levels (% FBGL) from 0 to 9 hours are plotted to assess the bioactivity of a given insulin-Fc fusion protein configuration. It is expected that a single subcutaneous dose of SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, or SEQ ID NO: 84 will demonstrate that these insulin-Fc fusion proteins show significant bioactivity in mice.

Example 61: Exemplary Insulin-Fc Fusion Protein Domains and Sequences.

[0305] Exemplary insulin-Fc fusion protein amino acid sequences and corresponding DNA sequences used in the above Examples are shown in FIG. 31, FIG. 32, FIG. 33, FIG. 34, FIG. 35, FIG. 36 and FIG.

**EQUIVALENTS**

[0306] In the claims, articles such as "a," "an," and "the" may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context.

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- WO 2018107117 A [0005]
- US 9855318 B2 [0062]
- US 10597435 B [0280]

## Non-patent literature cited in the description

- **AAK. WANG et al.** IgG Fc engineering to modulate antibody effector functions. *PROTEIN & CELL*, vol. 9 (1), 63-73 [0005]
- Current Protocols in Molecular Biology. Greene Publishing and Wiley- Interscience, 1995 [0048]
- **DAYHOFF.** Atlas of Polypeptide Sequence and Structure. National Biomedical Research Foundation, 1978, vol. 5 [0048]
- **SMITH ; WATERMAN.** *Adv. Appl. Math.*, 1981, vol. 2, 482-489 [0049]
- **ALLEVA, D.G. ; GAUR, A. ; JIN, L. ; WEGMANN, D. ; GOTTLIEB, P.A. ; PAHUJA, A. ; JOHNSON, E.B. ; MOTHERAL, T. ; PUTNAM, A. ; CROWE, P.D.** *Diabetes*, 2002, vol. 51 (7), 2126-2134 [0064]
- **CHEN, X. ; ZARO, J. ; SHEN, W.C.** *Adv Drug Deliv Rev.*, 15 October 2013, vol. 65 (10), 1357-1369 [0065]
- **VLASAK, J. ; IONESCU, R.** *MAbs*, 2011, vol. 3 (3), 253-263 [0067] [0255]
- **GUYER et al.** *J. Immunol.*, 1976, vol. 117, 587 [0071]
- **KIM et al.** *J. Immunol.*, 1994, vol. 24, 249 [0071]
- **DICK, LW.** *Biotechnol Bioeng.*, 15 August 2008, vol. 100 (6), 1132-43 [0072]
- **DOZIER, J. ; DISTEFANO, M.** Site-Specific PEGylation of Therapeutic Proteins. *Int. J. Mol. Sci.*, 2015, vol. 16 (10), 25831-25864 [0118]
- **CORWIN HANSCH.** Comprehensive Medicinal Chemistry. Pergamon Press, 1990, vol. 5 [0143]
- **GUILLIAMS, MARTIN ; BRUHNS, PIERRE ; SAEYS, YVAN ; HAMMAD, HAMIDA ; LAMBRECHT, BART.** The function of Fc gamma receptors in dendritic cells and macrophages. *Nature reviews. Immunology.*, 2014, vol. 14 [0211]
- **KOUSER, L. ; MADHUKARAN, S. P. ; SHASTRI, A. ; SARAON, A. ; FERLUGA, J. ; AL-MOZAINI, M. ; KISHORE, U.** Emerging and Novel Functions of Complement Protein C1q. *Frontiers in Immunology*, 2015, vol. 6, 317 [0212]
- **HUBBARD S.R.** Structural biology: Insulin meets its receptor. *Nature*, 2013, vol. 493 (7431), 171-172 [0239]
- **MENTING et al.** *Nature*, 2013, vol. 493 (7431), 241-245 [0241]
- **BRANGE J. ; DODSON G.G. ; EDWARDS J. ; HOLDEN P.H. ; WHITTINGHAM J.L.** A model of insulin fibrils derived from the x-ray crystal structure of a monomeric insulin (despentapeptide insulin). *Proteins*, 1997, vol. 27, 507-516 [0245]
- **LO, M. et al.** Effector attenuating substitutions that maintain antibody stability and reduce toxicity in mice. *J. Biol. Chem.*, 2017, 1-20 [0262]

**PATENTKRAV:**

1. Fusionprotein, som omfatter et insulinpolypeptid og et Fc-fragment, hvor insulinpolypeptidet og Fc-fragmentet er forbundet via en linker, der omfatter domæner i følgende orientering fra N- til C-terminalen: (N-terminal)—insulinpolypeptid—linker—Fc-fragment—(C-terminal),  
 hvor Fc-fragmentet er af human oprindelse og omfatter følgende sekvens:  
 DKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY  
 VDGVEVHNAKTKPREEQYX<sub>1</sub>STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT  
 ISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT  
 TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG (SEQ  
 ID NO: 77), hvor X<sub>1</sub> er S eller R;  
 hvor insulinpolypeptidet omfatter følgende sekvens:  
 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENY  
 C (SEQ ID NO: 10); og  
 hvor linkeren omfatter følgende sekvens:  
 GGGGGAGGGGAGGGGAGGGGG (SEQ ID NO: 67);  
 GGGGGQGGGGQGGGGQGGGGQGGGG (SEQ ID NO: 99); eller  
 GGGGGQGGGGQGGGGQGGGG (SEQ ID NO: 13).
2. Fusionsprotein ifølge krav 1, som omfatter et insulinpolypeptid og et Fc-fragment, hvor insulinpolypeptidet og Fc-fragmentet er forbundet via en linker, hvor fusionsproteinet omfatter følgende sekvens:  
 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENY  
 CGGGGGAGGGGAGGGGAGGGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMI  
 SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLH  
 QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL  
 VKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFC  
 SVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 87).
3. Fusionsprotein ifølge krav 1, hvor fusionsproteinet omfatter følgende sekvens:  
 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENY  
 CGGGGGQGGGGQGGGGQGGGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMI  
 SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLH  
 QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL  
 VKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFC  
 SVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 78).

4. Fusionsprotein ifølge krav 1, hvor fusionsproteinet omfatter følgende sekvens:  
 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENY  
 CGGGGGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI  
 SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYRSTYRVSVLTVLH  
 5 QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL  
 VKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSC  
 SVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 84).
5. Fusionsprotein ifølge krav 1, hvor fusionsproteinet omfatter følgende sekvens:  
 10 FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGGGIVEQCCTSTCSLDQLENY  
 CGGGGGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPK  
 KDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVS  
 VLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQ  
 VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQ  
 15 GNVFSCSVMHEALHNHYTQKSLSLSPG (SEQ ID NO: 96).
6. Fusionsprotein ifølge et hvilket som helst af ovennævnte krav, hvor fusionsproteinet er en homodimer, og hvor procentdelen af homodimer i fusionsproteinet er over 90 %.
- 20 7. Fusionsprotein ifølge krav 6, hvor fusionsproteinet er produceret ved anvendelse af én af HEK293-celler eller CHO-celler, og den resulterende homodimertiter efter oprensning ved anvendelse af Protein A-perler eller en Protein-A-søjle er over 150 mg/l.
8. Fusionsprotein ifølge krav 6, hvor insulinreceptor-IC<sub>50</sub> for fusionsproteinet er mindre end eller lig med 5000 nM.  
 25
9. Fusionsprotein ifølge krav 6, hvor human FcRn-receptor-EC<sub>50</sub> for fusionsproteinet er mindre end eller lig med 1000 ng/ml.
- 30 10. Fusionsprotein ifølge krav 6, hvor human Fc(gamma)RI-receptor assay-OD<sub>450</sub>-ratio ved en biotinyleret Fc(gamma)RI-koncentration på 3000 ng/ml af fusionsproteinet er mindre end eller lig med 0,50.
11. Fusionsprotein ifølge krav 6, hvor human C1q-assay-OD<sub>450</sub>-ratio ved en biotinyleret  
 35 C1q-koncentration på 1000 ng/ml er mindre end eller lig med 0,35.

12. Fusionsprotein ifølge krav 6, hvor fusionsproteinet er formuleret som en farmaceutisk sammensætning, og hvor fusionsproteinet er til stede i den farmaceutiske sammensætning ved en koncentration på ca. 3 mg/ml eller derover.
- 5 13. Farmaceutisk sammensætning ifølge krav 12, hvor sammensætningen er egnet til subkutan administration.
14. Fusionsprotein ifølge krav 6 eller en farmaceutisk sammensætning deraf til anvendelse som et medikament til sænkning af blodglucoseniveauet hos en patient.
- 10 15. Celle, der er modificeret til ekspression af fusionsproteinet ifølge et hvilket som helst af ovennævnte krav.
16. cDNA, der koder for fusionsproteinet ifølge SEQ ID NO: 87, som omfatter følgende nukleinsyresekvens:
- 15 atggaatggagctgggtcttctcttctcctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgctggctccc  
acctgggtgaagctctggcactcgtgtgctggcgagcggggctccactacgggggtggcgaggagggttctggtggcg  
cggaggcatcgtggaacagtgtctgcacctccacctgctccctggaccagctggaaaactactgctgggtggcgagggtgt  
gcaggaggcggtggagccggtggagggtggggctggaggaggcgggggagacaaaactcacacatgccaccgtgc  
20 ccagcacctgaactcctggggggaccgtcagcttctccttccccccaaaacccaaggacacctcatgatctccggac  
ccctgagggtcacatgctggtgggtggacgtgagccacgaagaccctgagggtcaagtcaactggtagctggacggcggtg  
gagggtgcataatgccaagacaaagccgctggaggagcagtagcagcagcagtagctggtgagcgtcctcaccgt  
cctgcaccaggactggctgaatggcaaggagtacaagtgaagggtcctcaacaaagccctccagcccccatcgaga  
aaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacaccctgcccccatccgggatgagctgac  
25 caagaaccagggtcagcctgacctgctggtcaaaggcttctatccagcgacatcgccgtggagtgaggagcaatgg  
gcagccgggagaacaactacaagaccacgcctcccgctgctggactccgacgggtccttctcctctacagcaagctcacc  
gtggacaagagcagggtggcagcaggggaacgtcttctcatgctccgtgatgatgaggctctgcacaaccactacacgc  
agaagagcctctccctgtctccgggttag (SEQ ID NO: 88).
- 30 17. cDNA, der koder for fusionsproteinet ifølge SEQ ID NO: 78, som omfatter følgende nukleinsyresekvens:
- atggaatggagctgggtcttctcttctcctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgctggctccc  
acctgggtgaagctctggcactcgtgtgctggcgagcggggctccactacgggggtggcgaggagggttctggtggcg  
cggaggcatcgtggaacagtgtctgcacctccacctgctccctggaccagctggaaaactactgctgggtggcgagggtgt  
35 caaggaggcggtggacaggggtggagggtgggcagggaggaggcgggggagacaaaactcacacatgccaccgtg  
cccagcacctgaactcctggggggaccgtcagcttctccttccccccaaaacccaaggacacctcatgatctccggga  
cccctgagggtcacatgctggtgggtggacgtgagccacgaagaccctgagggtcaagtcaactggtagctggacggcgt  
ggagggtgcataatgccaagacaaagccgctggaggagcagtagcagcagcagtagctggtgagcgtcctcaccg

5      tcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaacaaagccctcccagccccatcgaga  
aaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgac  
caagaaccaggctcagcctgacctgctggtaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgg  
gcagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcacc  
gtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgc  
agaagagcctctccctgtctccgggttag (SEQ ID NO: 79).

18. cDNA, der koder for fusionsproteinet ifølge SEQ ID NO: 84, som omfatter følgende nukleinsyresekvens:
- 10      atggaatggagctgggtctttcttcttctgtcagtaacgactgggtgccactccttcgtgaaccagcacctgtgaggctccc  
acctgggtgaagctctggcactcgtgtgcccgcagcggggctccactacgggggtggcggaggagggttctgtggcgg  
cggaggcatcgtggaacagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcggagggtgt  
caaggaggcgggtggacaggggtggagggtgggcagggaggaggcgggggagacaaaactcacacatgccaccgtg  
cccagcacctgaactcctggggggaccgtcagtcttcttctcccccaaaacccaaggacaccctcatgatctccgga  
15      cccctgaggtcacatgctggtggtggacgtgagccacgaagaccctgaggtcaagtcaactggtacgtggacggcgt  
ggagggtgcataatgccaagacaaagccgcgggaggagcagtacagaagcacgtaccgtgtggtcagcgtcctcacc  
gtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaacaaagccctcccagccccatcgag  
aaaaccatctcaaagccaaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctga  
ccaagaaccaggctcagcctgacctgcctggtaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatg  
20      ggcagccgggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctctacagcaagctcac  
cgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacg  
cagaagagcctctccctgtctccgggttag (SEQ ID NO: 85).

# DRAWINGS




FIG. 1




FIG. 2

```

SEQ ID NO: 32      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 33      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 34      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 31      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 35      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
                    *****

SEQ ID NO: 32      GAGGGGRC TDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTEVTCVVLDLGREDPEVQI 120
SEQ ID NO: 33      GAGGGGRC TDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTEVTCVVLDLGREDPEVQI 120
SEQ ID NO: 34      GAGGGGRC TDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTEVTCVVLDLGREDPEVQI 120
SEQ ID NO: 31      GAGGGGRC TDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTEVTCVVLDLGREDPEVQI 120
SEQ ID NO: 35      GAGGGGRC TDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTEVTCVVLDLGREDPEVQI 120
                    *****

SEQ ID NO: 32      SWFVDGKEVHTAKTQSREQQFNGTYRVVSVLP IEHQDWLTGKEFKCRVNHIDLPSP IERT 180
SEQ ID NO: 33      SWFVDGKEVHTAKTQSREQQFNGTYRVVSVLP IEHQDWLTGKEFKCRVNHIDLPSP IERT 180
SEQ ID NO: 34      SWFVDGKEVHTAKTQSREQQFNGTYRVVSVLP IEHQDWLTGKEFKCRVNHIDLPSP IERT 180
SEQ ID NO: 31      SWFVDGKEVHTAKTQSREQQFNGTYRVVSVLP IEHQDWLTGKEFKCRVNHIDLPSP IERT 180
SEQ ID NO: 35      SWFVDGKEVHTAKTQSREQQFNGTYRVVSVLP IEHQDWLTGKEFKCRVNHIDLPSP IERT 180
                    *****

SEQ ID NO: 32      ISKARGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQQEPEKHR 240
SEQ ID NO: 33      ISKARGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQQEPEKHR 240
SEQ ID NO: 34      ISKARGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQQEPEKHR 240
SEQ ID NO: 31      ISKARGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQQEPEKHR 240
SEQ ID NO: 35      ISKARGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQQEPEKHR 240
                    *****

SEQ ID NO: 32      MTPPQLDEDGSYFLYSKLSVDKSRWQQGDPFTCAVLHEALHSHYTQKSLSLSPG 294
SEQ ID NO: 33      MTPPQLDEDGSYFLYSKLSVDKSRWQQGDPFTCAVLHETLQSHYTDLSLSHSPG 294
SEQ ID NO: 34      MTPPQLDEDGSYFLYSKLSVDKSRWQQGDPFTCAVMHETLQSHYTDLSLSHSPG 294
SEQ ID NO: 31      MTPPQLDEDGSYFLYSKLSVDKSRWQQGDPFTCAVMHETLQNHYTDLNHYTDLSLSHSPG 294
SEQ ID NO: 35      MTPPQLDEDGSYFLYSKLSVDKSRWQQGDPFTCAVLHETLQNHYTDLNHYTDLSLSHSPG 294
                    *****:.*:.*:.*:.*:.*:

```

FIG. 3

```

SEQ ID NO: 31      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 38      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 36      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
SEQ ID NO: 37      FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG 60
                    *****

SEQ ID NO: 31      GAGGGGRCTDTPPCPVPEPLGGPSVLIFPPKPKDILRITRTPETCVVLDLGRDPEVQI 120
SEQ ID NO: 38      GAGGGGC---ISPCVPPESLGGPSVFIFFPKPKDILRITRTPETCVVLDLGRDPEVQI 117
SEQ ID NO: 36      GAGGGGDCPK---CPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLDPEDPEVQI 117
SEQ ID NO: 37      GAGGGG-CNN-CPCPGCGLLGGPSVFIFFPKPKDILVTARTPTVTCVVVDLDPENPEVQI 118
                    *****      **      *****:***** *      :*** :****:*. *:*****

SEQ ID NO: 31      SWFVDGKEVHTAKTQSRQQFNQNGTYRVVSVLPPIEQDWLTGKEFKCRVNHIDLPSPERT 180
SEQ ID NO: 38      SWFVDGKEVHTAKTQPREQQFNSTYRVVSVLPPIEQDWLTGKEFKCRVNHIGLPSPIERT 177
SEQ ID NO: 36      SWFVDGKQMQTAKTQPREQQFNQNGTYRVVSVLPPIEQDWLKGKQFTCKVNNKALPSPIERT 177
SEQ ID NO: 37      SWFVDSKQVQTANTQPREQQSNGTYRVVSVLPPIEQDWLSGKQFKCKVNNKALPSPIEEI 178
                    *****:*.::*:** *: * *.***** *****:*.*:*.*: *****.

SEQ ID NO: 31      ISKARGRAHKPSVYVLPSPKELSSSDTVSITCLIKDFYPPDIDVEWQSNQQEPEPKHR 240
SEQ ID NO: 38      ISKARGQAHQPSVYVLPSPKELSSSDTVTLTCLIKDFFPPEIDVEWQSNQQEPEPESKYH 237
SEQ ID NO: 36      ISKARGQAHQPSVYVLPSPREELS-KNTVSLTCLIKDFFPPEIDVEWQSNQQEPEPESKYR 236
SEQ ID NO: 37      ISKTPGQAHQPNVYVLPSPRDEMS-KNTVTLTCLVKDFFPPEIDVEWQSNQQEPEPESKYR 237
                    ***: *:***:*.***** .*: * .:***:***:***:***:***** *** *:;

SEQ ID NO: 31      MTPPQLDEDEGSYFLYSKLSVDKSRWQQGDPFTCAVMHETLQNHYTDLSLSHSPG 294
SEQ ID NO: 38      TTAPQLDEDEGSYFLYSKLSVDKSRWQQGDTFTCAVMHEALQNHYTDLSLSHSPG 291
SEQ ID NO: 36      TTPPQLDEDEGSYFLYSKLSVDKSRWQRGDTFTCAVMHEALHNHYTQESLSHSPG 290
SEQ ID NO: 37      MTPPQLDEDEGSYFLYSKLSVDKSRWQRGDTFTCAVMHEALHNHYTQISLSHSPG 291
                    * *****:*****:*** * *****:*.*****: *****

```

FIG. 4




FIG. 5




FIG. 6




FIG. 7

|               |                                                              |     |
|---------------|--------------------------------------------------------------|-----|
| SEQ ID NO: 39 | FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG | 60  |
| SEQ ID NO: 40 | FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG | 60  |
| SEQ ID NO: 41 | FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG | 60  |
| SEQ ID NO: 42 | FVNQHLCGSDLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCHSICSLYQLENYCNGGG | 60  |
| *****         |                                                              |     |
| SEQ ID NO: 39 | GAGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF | 120 |
| SEQ ID NO: 40 | GAGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF | 120 |
| SEQ ID NO: 41 | GAGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF | 120 |
| SEQ ID NO: 42 | GAGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF | 120 |
| *****         |                                                              |     |
| SEQ ID NO: 39 | VDGKQMQTAKTQPREEQFGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISK  | 180 |
| SEQ ID NO: 40 | VDGKQMQTAKTQPREEQFGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISK  | 180 |
| SEQ ID NO: 41 | VDGKQMQTAKTQPREEQFGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISK  | 180 |
| SEQ ID NO: 42 | VDGKQMQTAKTQPREEQFGTYRVVSVLPIGHQDWLKGKQFTCKVNNKALPSPIERTISK  | 180 |
| *****         |                                                              |     |
| SEQ ID NO: 39 | ARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP  | 240 |
| SEQ ID NO: 40 | ARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP  | 240 |
| SEQ ID NO: 41 | ARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP  | 240 |
| SEQ ID NO: 42 | ARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP  | 240 |
| *****         |                                                              |     |
| SEQ ID NO: 39 | QLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG           | 290 |
| SEQ ID NO: 40 | QLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG           | 290 |
| SEQ ID NO: 41 | QLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG           | 290 |
| SEQ ID NO: 42 | QLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG           | 290 |
| *****         |                                                              |     |

FIG. 8




FIG. 9




FIG. 10




FIG. 11

```

SEQ ID NO: 43      FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCNGGG 60
SEQ ID NO: 44      FVNQHLCGSHLVQALYLVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCNGG-G 59
SEQ ID NO: 45      FVNQHLCGSELVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCNGG-G 59
SEQ ID NO: 46      FVNQHLCGSHLVEALALVCGEAGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCNGG-G 59
SEQ ID NO: 47      FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCNGG-G 59
SEQ ID NO: 48      FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSICSLYQLENYCNGG-G 59
                    ***** . ** : * ***** ** : ***** ***** . * *

SEQ ID NO: 43      GAGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF 120
SEQ ID NO: 44      GAGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF 119
SEQ ID NO: 45      GAGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF 119
SEQ ID NO: 46      GAGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF 119
SEQ ID NO: 47      GAGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF 119
SEQ ID NO: 48      GAGGGGDCPKCPAPEMLGGPSVFIFPPKPKDTLLIARTPEVTCVVVDLDPEDPEVQISWF 119
                    *****

SEQ ID NO: 43      VDGKQMQTAKTQPREEQFGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSP IERTISK 180
SEQ ID NO: 44      VDGKQMQTAKTQPREEQFGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSP IERTISK 179
SEQ ID NO: 45      VDGKQMQTAKTQPREEQFGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSP IERTISK 179
SEQ ID NO: 46      VDGKQMQTAKTQPREEQFGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSP IERTISK 179
SEQ ID NO: 47      VDGKQMQTAKTQPREEQFGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSP IERTISK 179
SEQ ID NO: 48      VDGKQMQTAKTQPREEQFGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSP IERTISK 179
                    *****

SEQ ID NO: 43      ARGQAHQPSVYVLPSPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP 240
SEQ ID NO: 44      ARGQAHQPSVYVLPSPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP 239
SEQ ID NO: 45      ARGQAHQPSVYVLPSPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP 239
SEQ ID NO: 46      ARGQAHQPSVYVLPSPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP 239
SEQ ID NO: 47      ARGQAHQPSVYVLPSPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP 239
SEQ ID NO: 48      ARGQAHQPSVYVLPSPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKYRTTTP 239
                    *****

SEQ ID NO: 43      QLDEGGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG      290
SEQ ID NO: 44      QLDEGGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG      289
SEQ ID NO: 45      QLDEGGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG      289
SEQ ID NO: 46      QLDEGGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG      289
SEQ ID NO: 47      QLDEGGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG      289
SEQ ID NO: 48      QLDEGGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG      289
                    *****

```

FIG. 12

|               |                                                               |     |
|---------------|---------------------------------------------------------------|-----|
| SEQ ID NO: 43 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGG-----GPRRGIVEQCCTSI CSLYQLENY | 55  |
| SEQ ID NO: 49 | FVNQHLCGSHLVQALYLVCGERGFFYTDPTQRGGG--GGQRGIVEQCCTSI CSLYQLENY | 58  |
| SEQ ID NO: 50 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGGGGSGGGGGIVEQCCTSI CSLYQLENY | 60  |
| SEQ ID NO: 51 | FVNQHLCGSHLVEALALVCGERGFFYTDPPGGG---GGGGGIVEQCCTSI CSLYQLENY  | 56  |
| SEQ ID NO: 52 | FVNQHLCGSHLVEALALVCGERGFFYT-PGGGG---GGGGGIVEQCCTSI CSLYQLENY  | 55  |
|               | *****. ** ***** *                                             |     |
|               |                                                               |     |
| SEQ ID NO: 43 | CNGGGGAGGGGDCPKCPAPEMLGGPSVFI FPPKPKDTLLIARTPEVTCVVVDLDPEDPEV | 115 |
| SEQ ID NO: 49 | CGG-GGAGGGGDCPKCPAPEMLGGPSVFI FPPKPKDTLLIARTPEVTCVVVDLDPEDPEV | 117 |
| SEQ ID NO: 50 | CGG-GGAGGGGDCPKCPAPEMLGGPSVFI FPPKPKDTLLIARTPEVTCVVVDLDPEDPEV | 119 |
| SEQ ID NO: 51 | CGG-GGAGGGGDCPKCPAPEMLGGPSVFI FPPKPKDTLLIARTPEVTCVVVDLDPEDPEV | 115 |
| SEQ ID NO: 52 | CGG-GGAGGGGDCPKCPAPEMLGGPSVFI FPPKPKDTLLIARTPEVTCVVVDLDPEDPEV | 114 |
|               | * , *****                                                     |     |
|               |                                                               |     |
| SEQ ID NO: 43 | QISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSPIE | 175 |
| SEQ ID NO: 49 | QISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSPIE | 177 |
| SEQ ID NO: 50 | QISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSPIE | 179 |
| SEQ ID NO: 51 | QISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSPIE | 175 |
| SEQ ID NO: 52 | QISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLP IGHQDWLKGKQFTCKVNNKALPSPIE | 174 |
|               | *****. *****                                                  |     |
|               |                                                               |     |
| SEQ ID NO: 43 | RTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKY   | 235 |
| SEQ ID NO: 49 | RTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKY   | 237 |
| SEQ ID NO: 50 | RTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKY   | 239 |
| SEQ ID NO: 51 | RTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKY   | 235 |
| SEQ ID NO: 52 | RTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQEPESKY   | 234 |
|               | *****                                                         |     |
|               |                                                               |     |
| SEQ ID NO: 43 | RTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG       | 290 |
| SEQ ID NO: 49 | RTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG       | 292 |
| SEQ ID NO: 50 | RTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG       | 294 |
| SEQ ID NO: 51 | RTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG       | 290 |
| SEQ ID NO: 52 | RTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG       | 289 |
|               | *****                                                         |     |

FIG. 13

|               |                                                                        |     |
|---------------|------------------------------------------------------------------------|-----|
| SEQ ID NO: 43 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSIICSLYQLENYCNGGG          | 60  |
| SEQ ID NO: 53 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSIICSLYQLENYCGGGG          | 60  |
| SEQ ID NO: 48 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGGGPRRGIVEQCCTSIICSLYQLENYCGGGG<br>***** | 60  |
| SEQ ID NO: 43 | GA-----GGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLD                  | 109 |
| SEQ ID NO: 53 | GQGGGQGGGQGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLD               | 120 |
| SEQ ID NO: 48 | A-----GGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDLD<br>.<br>*****     | 108 |
| SEQ ID NO: 43 | PEDPEVQISWFVDGKQMQTAKTQPREEQFGTYRVVSVLPIGHQDWLKGKQFTCKVNNKA            | 169 |
| SEQ ID NO: 53 | PEDPEVQISWFVDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKA           | 180 |
| SEQ ID NO: 48 | PEDPEVQISWFVDGKQMQTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNKA<br>*****  | 168 |
| SEQ ID NO: 43 | LPSPiERTISKARGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQ            | 229 |
| SEQ ID NO: 53 | LPSPiERTISKARGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQ            | 240 |
| SEQ ID NO: 48 | LPSPiERTISKARGQAHQPSVYVLPPSREELSKNTVSLTCLIKDFFPPDIDVEWQSNQQ<br>*****   | 228 |
| SEQ ID NO: 43 | EPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSP            | 289 |
| SEQ ID NO: 53 | EPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSP            | 300 |
| SEQ ID NO: 48 | EPESKYRTTPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSP<br>*****   | 288 |
| SEQ ID NO: 43 | G 290                                                                  |     |
| SEQ ID NO: 53 | G 301                                                                  |     |
| SEQ ID NO: 48 | G 289<br>*                                                             |     |

FIG. 14

|               |                                                      |     |
|---------------|------------------------------------------------------|-----|
| SEQ ID NO: 43 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGGG-PRRGIVEQCCTSI      | 59  |
| SEQ ID NO: 51 | FVNQHLCGSHLVEALALVCGERGFFYTDPGGGGGGGGIVEQCCTSI       | 59  |
| SEQ ID NO: 54 | FVNQHLCGSHLVEALALVCGERGFFYTDG-<br>GGGGGGGGGIVEQCCTSI | 58  |
| SEQ ID NO: 52 | FVNQHLCGSHLVEALALVCGERGFFYTPG-<br>GGGGGGGGGIVEQCCTSI | 58  |
|               | *****                                                |     |
| SEQ ID NO: 43 | GGAGGGGDCPKCPAPEMLGGPSVFI                            | 119 |
| SEQ ID NO: 51 | GGAGGGGDCPKCPAPEMLGGPSVFI                            | 119 |
| SEQ ID NO: 54 | GGAGGGGDCPKCPAPEMLGGPSVFI                            | 118 |
| SEQ ID NO: 52 | GGAGGGGDCPKCPAPEMLGGPSVFI                            | 118 |
|               | *****                                                |     |
| SEQ ID NO: 43 | FVDGKQMQTAKTQPREEQFGTYRVVSVLP                        | 179 |
| SEQ ID NO: 51 | FVDGKQMQTAKTQPREEQFGTYRVVSVLP                        | 179 |
| SEQ ID NO: 54 | FVDGKQMQTAKTQPREEQFGTYRVVSVLP                        | 178 |
| SEQ ID NO: 52 | FVDGKQMQTAKTQPREEQFGTYRVVSVLP                        | 178 |
|               | *****                                                |     |
| SEQ ID NO: 43 | KARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQ        | 239 |
| SEQ ID NO: 51 | KARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQ        | 239 |
| SEQ ID NO: 54 | KARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQ        | 238 |
| SEQ ID NO: 52 | KARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQ        | 238 |
|               | *****                                                |     |
| SEQ ID NO: 43 | PQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG  | 290 |
| SEQ ID NO: 51 | PQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG  | 290 |
| SEQ ID NO: 54 | PQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG  | 289 |
| SEQ ID NO: 52 | PQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHSPG  | 289 |
|               | *****                                                |     |

FIG. 15

|               |                                                                                             |     |
|---------------|---------------------------------------------------------------------------------------------|-----|
| SEQ ID NO: 43 | FVNQHLCGSHLVEALALVCGERGFFYTDPTGG-GPRRGIVEQCCTSTCSLDQLENYCNGG                                | 59  |
| SEQ ID NO: 55 | FVNQHLCGSHLVEALELVCGERGFFYTPKTGGSGGGGGIVEQCCTSTCSLDQLENYCGG-                                | 59  |
| SEQ ID NO: 56 | FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCNHG                                | 60  |
| SEQ ID NO: 28 | FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCNGG                                | 60  |
| SEQ ID NO: 26 | FVNQHLCGSHLVEALELVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGG-                                | 59  |
| SEQ ID NO: 57 | FVNQHLCGSHLVEALELVCGERGFFYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGG-<br>*****.*****.* ** * ***** ** | 59  |
| SEQ ID NO: 43 | GG-----AGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVD                                       | 107 |
| SEQ ID NO: 55 | GGGQGGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVD                                | 119 |
| SEQ ID NO: 56 | GGGQGGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVD                                | 120 |
| SEQ ID NO: 28 | GGGQGGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVD                                | 120 |
| SEQ ID NO: 26 | GGGQGGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVD                                | 119 |
| SEQ ID NO: 57 | GGGQGGGGQGGGGQGGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVD<br>** .*****                   | 119 |
| SEQ ID NO: 43 | LDPEDPEVQISWFDGKQMOTAKTQPREEQFNGTYRVVSVLP IGHQDWLKGKQFTCKVNN                                | 167 |
| SEQ ID NO: 55 | LDPEDPEVQISWFDGKQMOTAKTQPREEQFNGTYRVVSVLP IGHQDWLKGKQFTCKVNN                                | 179 |
| SEQ ID NO: 56 | LDPEDPEVQISWFDGKQMOTAKTQPREEQFNGTYRVVSVLP IGHQDWLKGKQFTCKVNN                                | 180 |
| SEQ ID NO: 28 | LDPEDPEVQISWFDGKQMOTAKTQPREEQFNGTYRVVSVLP IGHQDWLKGKQFTCKVNN                                | 180 |
| SEQ ID NO: 26 | LDPEDPEVQISWFDGKQMOTAKTQPREEQFNGTYRVVSVLP IGHQDWLKGKQFTCKVNN                                | 179 |
| SEQ ID NO: 57 | LDPEDPEVQISWFDGKQMOTAKTQPREEQFNGTYRVVSVLP IGHQDWLKGKQFTCKVNN<br>*****:*****                 | 179 |
| SEQ ID NO: 43 | KALPSPIERTISKARGQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNG                                | 227 |
| SEQ ID NO: 55 | KALPSPIERTISKARGQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNG                                | 239 |
| SEQ ID NO: 56 | KALPSPIERTISKARGQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNG                                | 240 |
| SEQ ID NO: 28 | KALPSPIERTISKARGQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNG                                | 240 |
| SEQ ID NO: 26 | KALPSPIERTISKARGQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNG                                | 239 |
| SEQ ID NO: 57 | KALPSPIERTISKARGQAHQPSVYVLPSSREELSKNTVSLTCLIKDFFPPDIDVEWQSNG<br>*****                       | 239 |
| SEQ ID NO: 43 | QQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSH                                | 287 |
| SEQ ID NO: 55 | QQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSH                                | 299 |
| SEQ ID NO: 56 | QQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSH                                | 300 |
| SEQ ID NO: 28 | QQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSH                                | 300 |
| SEQ ID NO: 26 | QQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSH                                | 299 |
| SEQ ID NO: 57 | QQEPESKYRTTPPQLDEDGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSH<br>*****                       | 299 |
| SEQ ID NO: 43 | SPG 290                                                                                     |     |
| SEQ ID NO: 55 | SPG 302                                                                                     |     |
| SEQ ID NO: 56 | SPG 303                                                                                     |     |
| SEQ ID NO: 28 | SPG 303                                                                                     |     |
| SEQ ID NO: 26 | SPG 302                                                                                     |     |
| SEQ ID NO: 57 | SPG 302<br>***                                                                              |     |

FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23

|               |                                                              |     |
|---------------|--------------------------------------------------------------|-----|
| SEQ ID NO: 61 | FVNQHLCGSHLVEALELVCGERGFHYGGGGGGGGGGGIVEQCCTSTCSLDQLENYCGGG  | 60  |
| SEQ ID NO: 62 | FVNQHLCGSHLVEALELVCGERGFHYGGGGGGGGGGGIVEQCCTSTCSLDQLENYCGGG  | 60  |
| *****         |                                                              |     |
| SEQ ID NO: 61 | GGQGGGGQGGGGQGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVAL  | 120 |
| SEQ ID NO: 62 | GGQGGGGQGGGGQGGGGDCPKCPAPEMLGGPSVFIFFPKPKDTLLIARTPEVTCVVVDL  | 120 |
| *****         |                                                              |     |
| SEQ ID NO: 61 | DPEDPEVQISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNK | 180 |
| SEQ ID NO: 62 | DPEDPEVQISWFVDGKQMOTAKTQPREEQFSGTYRVVSVLPIGHQDWLKGKQFTCKVNNK | 180 |
| *****         |                                                              |     |
| SEQ ID NO: 61 | ALPSPIERTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQG | 240 |
| SEQ ID NO: 62 | ALPSPIERTISKARGQAHQPSVYVLPSPREELSKNTVSLTCLIKDFFPPDIDVEWQSNQG | 240 |
| *****         |                                                              |     |
| SEQ ID NO: 61 | QEPESKYRTTPQLDEDEGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHS | 300 |
| SEQ ID NO: 62 | QEPESKYRTTPQLDEDEGSYFLYSKLSVDKSRWQRGDTFICAVMHEALHNHYTQESLSHS | 300 |
| *****         |                                                              |     |
| SEQ ID NO: 61 | PG 302                                                       |     |
| SEQ ID NO: 62 | PG 302                                                       |     |
| **            |                                                              |     |

FIG. 24




FIG. 25




FIG. 26

|               |                                                              |     |
|---------------|--------------------------------------------------------------|-----|
| SEQ ID NO: 76 | FVNQHLCGSHLVEALELVCGERGFIHYGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| SEQ ID NO: 91 | FVNQHLCGSHLVEALELVCGERGFIHYGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| SEQ ID NO: 78 | FVNQHLCGSHLVEALALVCGERGFIHYGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
|               | *****                                                        |     |
| SEQ ID NO: 76 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV | 120 |
| SEQ ID NO: 91 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV | 120 |
| SEQ ID NO: 78 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV | 120 |
|               | *****                                                        |     |
| SEQ ID NO: 76 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
| SEQ ID NO: 91 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
| SEQ ID NO: 78 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
|               | *****                                                        |     |
| SEQ ID NO: 76 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 91 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 78 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
|               | *****                                                        |     |
| SEQ ID NO: 76 | SNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSMHEALHNYTQKSLS    | 300 |
| SEQ ID NO: 91 | SNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSMHEALHNYTQKSLS    | 300 |
| SEQ ID NO: 78 | SNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSMHEALHNYTQKSLS    | 300 |
|               | *****                                                        |     |
| SEQ ID NO: 76 | LSPG                                                         | 304 |
| SEQ ID NO: 91 | LSPG                                                         | 304 |
| SEQ ID NO: 78 | LSPG                                                         | 304 |
|               | ****                                                         |     |

FIG. 27

|                             |                                                              |     |
|-----------------------------|--------------------------------------------------------------|-----|
| SEQ ID NO: 75               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| SEQ ID NO: 76               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| SEQ ID NO: 91               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| SEQ ID NO: 92               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| SEQ ID NO: 93               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| SEQ ID NO: 94               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| SEQ ID NO: 95               | FVNQHLCGSHLVEALELVCGERGFHYGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG   | 60  |
| *****                       |                                                              |     |
| SEQ ID NO: 75               | GGQGGGGQGGGGQGGGGG---ECPPCPAPPV-AGPSVFLFPPKPKDTLMISRTPEVTCV  | 115 |
| SEQ ID NO: 76               | GGQGGGGQGGGGQGGGGGDKHTCPCPAPELLLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 91               | GGQGGGGQGGGGQGGGGGDKHTCPCPAPELLLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 92               | GGQGGGGQGGGGQGGGGGDKHTCPCPAPELLLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 93               | GGQGGGGQGGGGQGGGGGDKHTCPCPAPELLLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 94               | GGQGGGGQGGGGQGGGGGDKHTCPCPAPELLLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 95               | GGQGGGGQGGGGQGGGGGDKHTCPCPAPELLLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| ***** : .*****              |                                                              |     |
| SEQ ID NO: 75               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCK | 175 |
| SEQ ID NO: 76               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
| SEQ ID NO: 91               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| SEQ ID NO: 92               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYDSTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
| SEQ ID NO: 93               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
| SEQ ID NO: 94               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCK | 180 |
| SEQ ID NO: 95               | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| *****:*****: **:*****:***** |                                                              |     |
| SEQ ID NO: 75               | VSNKGLPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 235 |
| SEQ ID NO: 76               | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 91               | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 92               | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 93               | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 94               | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| SEQ ID NO: 95               | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE | 240 |
| ****.*****:*****:*.*****    |                                                              |     |
| SEQ ID NO: 75               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 295 |
| SEQ ID NO: 76               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 300 |
| SEQ ID NO: 91               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 300 |
| SEQ ID NO: 92               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 300 |
| SEQ ID NO: 93               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 300 |
| SEQ ID NO: 94               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 300 |
| SEQ ID NO: 95               | SNGQPENNYKTTTPVLDSGSGFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS  | 300 |
| *****:*****                 |                                                              |     |
| SEQ ID NO: 75               | LSPG 299                                                     |     |
| SEQ ID NO: 76               | LSPG 304                                                     |     |
| SEQ ID NO: 91               | LSPG 304                                                     |     |
| SEQ ID NO: 92               | LSPG 304                                                     |     |
| SEQ ID NO: 93               | LSPG 304                                                     |     |
| SEQ ID NO: 94               | LSPG 304                                                     |     |
| SEQ ID NO: 95               | LSPG 304                                                     |     |
| ****                        |                                                              |     |

FIG. 28

|               |                                                               |     |
|---------------|---------------------------------------------------------------|-----|
| SEQ ID NO: 76 | FVNQHLCGSHLVEALVLVCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG  | 60  |
| SEQ ID NO: 78 | FVNQHLCGSHLVEALALVLCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| SEQ ID NO: 80 | FVNQHLCGSHLVEALALVLCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| SEQ ID NO: 82 | FVNQHLCGSHLVEALALVLCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| SEQ ID NO: 84 | FVNQHLCGSHLVEALALVLCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| SEQ ID NO: 86 | FVNQHLCGSHLVEALALVLCGERGFHYGGGGGGSGGGGGIVEQCCTSTCSLDQLENYCGGG | 60  |
| *****         |                                                               |     |
| SEQ ID NO: 76 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 78 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 80 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 82 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 84 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| SEQ ID NO: 86 | GGQGGGGQGGGGQGGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  | 120 |
| *****         |                                                               |     |
| SEQ ID NO: 76 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| SEQ ID NO: 78 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| SEQ ID NO: 80 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYDSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| SEQ ID NO: 82 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| SEQ ID NO: 84 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYRSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| SEQ ID NO: 86 | VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCK  | 180 |
| *****         |                                                               |     |
| SEQ ID NO: 76 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE  | 240 |
| SEQ ID NO: 78 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE  | 240 |
| SEQ ID NO: 80 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE  | 240 |
| SEQ ID NO: 82 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE  | 240 |
| SEQ ID NO: 84 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE  | 240 |
| SEQ ID NO: 86 | VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE  | 240 |
| *****         |                                                               |     |
| SEQ ID NO: 76 | SNGQFENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS    | 300 |
| SEQ ID NO: 78 | SNGQFENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS    | 300 |
| SEQ ID NO: 80 | SNGQFENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS    | 300 |
| SEQ ID NO: 82 | SNGQFENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS    | 300 |
| SEQ ID NO: 84 | SNGQFENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS    | 300 |
| SEQ ID NO: 86 | SNGQFENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLS    | 300 |
| *****         |                                                               |     |
| SEQ ID NO: 76 | LSPG 304                                                      |     |
| SEQ ID NO: 78 | LSPG 304                                                      |     |
| SEQ ID NO: 80 | LSPG 304                                                      |     |
| SEQ ID NO: 82 | LSPG 304                                                      |     |
| SEQ ID NO: 84 | LSPG 304                                                      |     |
| SEQ ID NO: 86 | LSPG 304                                                      |     |
| ****          |                                                               |     |

FIG. 29

|               |                                                                       |     |
|---------------|-----------------------------------------------------------------------|-----|
| SEQ ID NO: 87 | FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG           | 60  |
| SEQ ID NO: 96 | FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG           | 60  |
| SEQ ID NO: 78 | FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG           | 60  |
| SEQ ID NO: 97 | FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG           | 60  |
| SEQ ID NO: 89 | FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGIVEQCCTSTCSLDQLENYCGGG           | 60  |
| SEQ ID NO: 98 | FVNQHLCGSHLVEALALVCGERGFHYGGGGGGSGGGGIVEQCCTSTCSLDQLENYCD--<br>*****  | 58  |
| SEQ ID NO: 87 | GGAGGGGAGGGGA-----GGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT            | 115 |
| SEQ ID NO: 96 | GGQGGGGQGGGGQGGGGQGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT            | 120 |
| SEQ ID NO: 78 | GGQGGGGQGGGGQGG-----GGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT           | 115 |
| SEQ ID NO: 97 | G-----QGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT                       | 103 |
| SEQ ID NO: 89 | G-----AGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT                       | 103 |
| SEQ ID NO: 98 | -----KTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT<br>*****                     | 94  |
| SEQ ID NO: 87 | EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGK          | 175 |
| SEQ ID NO: 96 | EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGK          | 180 |
| SEQ ID NO: 78 | EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGK          | 175 |
| SEQ ID NO: 97 | EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGK          | 163 |
| SEQ ID NO: 89 | EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGK          | 163 |
| SEQ ID NO: 98 | EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYSSTYRVVSVLTVLHQDWLNGK<br>***** | 154 |
| SEQ ID NO: 87 | EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI          | 235 |
| SEQ ID NO: 96 | EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI          | 240 |
| SEQ ID NO: 78 | EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI          | 235 |
| SEQ ID NO: 97 | EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI          | 223 |
| SEQ ID NO: 89 | EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI          | 223 |
| SEQ ID NO: 98 | EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI<br>***** | 214 |
| SEQ ID NO: 87 | AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOGNVFSCSVMEALHNHYT           | 295 |
| SEQ ID NO: 96 | AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOGNVFSCSVMEALHNHYT           | 300 |
| SEQ ID NO: 78 | AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOGNVFSCSVMEALHNHYT           | 295 |
| SEQ ID NO: 97 | AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOGNVFSCSVMEALHNHYT           | 283 |
| SEQ ID NO: 89 | AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOGNVFSCSVMEALHNHYT           | 283 |
| SEQ ID NO: 98 | AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOGNVFSCSVMEALHNHYT<br>*****  | 274 |
| SEQ ID NO: 87 | QKSLSLSPG                                                             | 304 |
| SEQ ID NO: 96 | QKSLSLSPG                                                             | 309 |
| SEQ ID NO: 78 | QKSLSLSPG                                                             | 304 |
| SEQ ID NO: 97 | QKSLSLSPG                                                             | 292 |
| SEQ ID NO: 89 | QKSLSLSPG                                                             | 292 |
| SEQ ID NO: 98 | QKSLSLSPG<br>*****                                                    | 283 |

FIG. 30

SEQ ID NO: 79 atggaatggagctgggtctttctcttcttctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 78 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggtcccacctggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcggaggaggttctggtggcggcggaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcggaggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 ggtcaaggaggcgggtggacaggggtggaggtgggcaggaggaggcgggggagacaaaact  
 G Q G G G G Q G G G G Q G G G G G D K T  
 cacacatgcccacgtgcccagcacctgaactcctggggggaccgtcagtcttctcttc  
 H T C P P C P A P E L L G G P S V F L F  
 cccccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtgggtg  
 P P K P K D T L M I S R T P E V T C V V  
 gtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggag  
 V D V S H E D P E V K F N W Y V D G V E  
 gtgcataatgccaagacaaagccgcgggaggagcagtacagcagcacgtaccgtgtggtc  
 V H N A K T K P R E E Q Y S S T Y R V V  
 agcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtc  
 S V L T V L H Q D W L N G K E Y K C K V  
 tccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccc  
 S N K A L P A P I E K T I S K A K G Q P  
 cgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtc  
 R E P Q V Y T L P P S R D E L T K N Q V  
 agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagagc  
 S L T C L V K G F Y P S D I A V E W E S  
 aatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgacggctcc  
 N G Q P E N N Y K T T P P V L D S D G S  
 ttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttc  
 F F L Y S K L T V D K S R W Q Q G N V F  
 tcatgtccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctg  
 S C S V M H E A L H N H Y T Q K S L S L  
 tctccgggttag  
 S P G -

FIG. 31

SEQ ID NO: 81 atggaatggagctgggtctttctcttcttcctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 80 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggctcccacctgggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgtcacctccacctgctccctggaccagctggaaaactactgcggtggcggaggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 ggtcaaggaggcgggtggacagggtggaggtgggcaggaggaggcgggggagacaaaact  
 G Q G G G G Q G G G G Q G G G G G D K T  
 cacacatgcccacgtgcccagcacctgaactcctggggggaccgtcagtcttcctcttc  
 H T C P P C P A P E L L G G P S V F L F  
 cccccaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtg  
 P P K P K D T L M I S R T P E V T C V V  
 gtggcgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggag  
 V D V S H E D P E V K F N W Y V D G V E  
 gtgcataatgccaagacaaagccgcgaggaggagcagtagcacagcacgtaccgtgtggtc  
 V H N A K T K P R E E Q Y D S T Y R V V  
 agcgtcctcacgtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtc  
 S V L T V L H Q D W L N G K E Y K C K V  
 tccaacaaagccctcccagcccccacgagaaaaccatctccaaagccaaaggcagccc  
 S N K A L P A P I E K T I S K A K G Q P  
 cgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtc  
 R E P Q V Y T L P P S R D E L T K N Q V  
 agcctgacctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagc  
 S L T C L V K G F Y P S D I A V E W E S  
 aatgggcagccggagaaactacaagaccacgcctcccgtgctggactccgacgggtcc  
 N G Q P E N N Y K T T P P V L D S D G S  
 ttcttcctctacagcaagctcacctggacaagagcaggtggcagcaggggaacgtcttc  
 F F L Y S K L T V D K S R W Q Q G N V F  
 tcatgtccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctg  
 S C S V M H E A L H N H Y T Q K S L S L  
 tctccgggttag  
 S P G -

FIG. 32

SEQ ID NO: 83 atggaatggagctgggtctttctcttcttctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 82 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggctcccacctggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcggaggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 ggtcaaggaggcgggtggacaggggtggaggtgggcaggaggaggcgggggagacaaaact  
 G Q G G G G Q G G G G Q G G G G G D K T  
 cacacatgcccacgtgccacgacactgaactcctggggggaccgtcagtcttctcttc  
 H T C P P C P A P E L L G G P S V F L F  
 cccccaaaacccaaggacacctcatgatctcccgaccctgaggtcacatgcgtgggtg  
 P P K P K D T L M I S R T P E V T C V V  
 gtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggag  
 V D V S H E D P E V K F N W Y V D G V E  
 gtgcataatgccaaagacaaagccgcgaggagcagtagccagcacgtaccgtgtggtc  
 V H N A K T K P R E E Q Y A S T Y R V V  
 agcgtcctcacgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtc  
 S V L T V L H Q D W L N G K E Y K C K V  
 tccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaaggcagccc  
 S N K A L P A P I E K T I S K A K G Q P  
 cgagaaccacaggtgtacacctgcccccatcccgggatgagctgaccaagaaccaggtc  
 R E P Q V Y T L P P S R D E L T K N Q V  
 agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagc  
 S L T C L V K G F Y P S D I A V E W E S  
 aatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctcc  
 N G Q P E N N Y K T T P P V L D S D G S  
 ttcttctctacagcaagctcacctggacaagagcaggtggcagcaggggaacgtcttc  
 F F L Y S K L T V D K S R W Q Q G N V F  
 tcatgtcctgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctg  
 S C S V M H E A L H N H Y T Q K S L S L  
 tctccgggttag  
 S P G -

FIG. 33

SEQ ID NO: 85 atggaatggagctgggtctttctcttcttctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 84 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggtcccacctggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcgaggaggttctggtggcgggcgaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcgagggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 ggtcaaggaggcggtggacaggggtggaggtgggcagggaggaggcgggggagacaaaact  
 G Q G G G G Q G G G G Q G G G G G D K T  
 cacacatgcccaccgtgcccagcacctgaactcctggggggaccgtcagtcttctcttc  
 H T C P P C P A P E L L G G P S V F L F  
 cccccaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtg  
 P P K P K D T L M I S R T P E V T C V V  
 gtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcggtggag  
 V D V S H E D P E V K F N W Y V D G V E  
 gtgcataatgccaagacaaagccgcgaggagcagtagagaagcacgtaccgtgtggtc  
 V H N A K T K P R E E Q Y R S T Y R V V  
 agcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggctc  
 S V L T V L H Q D W L N G K E Y K C K V  
 tccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccc  
 S N K A L P A P I E K T I S K A K G Q P  
 cgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtc  
 R E P Q V Y T L P P S R D E L T K N Q V  
 agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagc  
 S L T C L V K G F Y P S D I A V E W E S  
 aatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctcc  
 N G Q P E N N Y K T T P P V L D S D G S  
 ttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttc  
 F F L Y S K L T V D K S R W Q Q G N V F  
 tcatgtcctgcatgcatgaggctctgcacaaccactacacgcagaagagcctctccctg  
 S C S V M H E A L H N H Y T Q K S L S L  
 tctccgggttag  
 S P G -

FIG. 34

SEQ ID NO: 88 atggaatggagctgggtctttctcttcttctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 87 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggtcccacctgggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcggaggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 ggtgcaggaggcgggtggagccggtggaggtggggctggaggaggcgggggagacaaaact  
 G A G G G G A G G G G A G G G G G D K T  
 cacacatgcccaccgtgcccagcacctgaactcctggggggaccgtcagtcttctcttc  
 H T C P P C P A P E L L G G P S V F L F  
 cccccaaaacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtgggtg  
 P P K P K D T L M I S R T P E V T C V V  
 gtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggag  
 V D V S H E D P E V K F N W Y V D G V E  
 gtgcataatgccaagacaaagccgcgaggagcagtagcagcagcacgtaccgtgtggtc  
 V H N A K T K P R E E Q Y S S T Y R V V  
 agcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggctc  
 S V L T V L H Q D W L N G K E Y K C K V  
 tccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccc  
 S N K A L P A P I E K T I S K A K G Q P  
 cgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtc  
 R E P Q V Y T L P P S R D E L T K N Q V  
 agcctgacctgcctgggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagc  
 S L T C L V K G F Y P S D I A V E W E S  
 aatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctcc  
 N G Q P E N N Y K T T P P V L D S D G S  
 ttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttc  
 F F L Y S K L T V D K S R W Q Q G N V F  
 tcatgtccgtgatgcatgaggtctgcacaaccactacacgcagaagagcctctccctg  
 S C S V M H E A L H N H Y T Q K S L S L  
 tctccgggttag  
 S P G -

FIG. 35

SEQ ID NO: 90 atggaatggagctgggtctttctcttcttctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 89 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggctcccacctggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcggaggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 gccggaggcgggggagacaaaactcacacatgccaccgtgccagcacctgaactcctg  
 A G G G G D K T H T C P P C P A P E L L  
 gggggaccgtcagtcttctcttccccccaaaacccaaggacaccctcatgatctcccg  
 G G P S V F L F P P K P K D T L M I S R  
 acccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttc  
 T P E V T C V V V D V S H E D P E V K F  
 aactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgaggagcag  
 N W Y V D G V E V H N A K T K P R E E Q  
 tacagcagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggtgaat  
 Y S S T Y R V V S V L T V L H Q D W L N  
 ggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaacc  
 G K E Y K C K V S N K A L P A P I E K T  
 atctccaaagccaaaggcagccccgagaaccacaggtgtacaccctgcccccatcccgg  
 I S K A K G Q P R E P Q V Y T L P P S R  
 gatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagc  
 D E L T K N Q V S L T C L V K G F Y P S  
 gacatcgccgtggagtgggagagcaatgggcagccggagaaactacaagaccacgcct  
 D I A V E W E S N G Q P E N N Y K T T P  
 cccgtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagc  
 P V L D S D G S F F L Y S K L T V D K S  
 aggtggcagcaggggaacgtcttctcatgctccgtgatgcaggtctgcacaaccac  
 R W Q Q G N V F S C S V M H E A L H N H  
 tacacgcagaagagcctctccctgtctccgggttag  
 Y T Q K S L S L S P G -

FIG. 36

SEQ ID NO: 100 atggaatggagctgggtctttctcttcttctgtcagtaacgactgggtgtccactccttc  
 SEQ ID NO: 86 M E W S W V F L F F L S V T T G V H S F  
 gtgaaccagcacctgtgcggtcccacctgggtggaagctctggcactcgtgtgcggcgag  
 V N Q H L C G S H L V E A L A L V C G E  
 cggggcttccactacgggggtggcggaggaggttctggtggcggcgaggcatcgtggaa  
 R G F H Y G G G G G S G G G G G I V E  
 cagtgtgcacctccacctgctccctggaccagctggaaaactactgcggtggcggaggt  
 Q C C T S T C S L D Q L E N Y C G G G G  
 ggtcaaggaggcgggtggacagggtggaggtggcaggaggaggcggggagacaaaact  
 G Q G G G G Q G G G G Q G G G G G D K T  
 cacacatgcccaccgtgccagcacctgaactcctggggggaccgtcagttcttctcttc  
 H T C P P C P A P E L L G G P S V F L F  
 ccccaaaacccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtg  
 P P K P K D T L M I S R T P E V T C V V  
 gtggacgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggcgtggag  
 V D V S H E D P E V K F N W Y V D G V E  
 gtgcataatgccaaagacaaagccgcgggaggagcagtaacaaagcacgtaccgtgtggtc  
 V H N A K T K P R E E Q Y Q S T Y R V V  
 agcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtaacagtgcaaggtc  
 S V L T V L H Q D W L N G K E Y K C K V  
 tccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccc  
 S N K A L P A P I E K T I S K A K G Q P  
 cgagaaccacaggtgtacacctgccccatcccgggatgagctgaccaagaaccaggtc  
 R E P Q V Y T L P P S R D E L T K N Q V  
 agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagc  
 S L T C L V K G F Y P S D I A V E W E S  
 aatgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctcc  
 N G Q P E N N Y K T T P P V L D S D G S  
 ttcttctctacagcaagctcacctggacaagagcaggtggcagcaggggaacgtcttc  
 F F L Y S K L T V D K S R W Q Q G N V F  
 tcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctg  
 S C S V M H E A L H N H Y T Q K S L S L  
 tctccgggttag  
 S P G -

FIG. 37




FIG. 38

## SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

