

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2014205114 B2**

(54) Title  
**Methods and compositions for affecting the flavor and aroma profile of consumables**

(51) International Patent Classification(s)  
**A23D 7/005 (2006.01) A23J 1/06 (2006.01)**  
**A23D 9/013 (2006.01) A23J 1/12 (2006.01)**  
**A23J 1/00 (2006.01) A23J 1/14 (2006.01)**

(21) Application No: **2014205114** (22) Date of Filing: **2014.01.13**

(87) WIPO No: **WO14/110532**

(30) Priority Data

(31) Number (32) Date (33) Country  
**13/941,211 2013.07.12 US**  
**61/908,634 2013.11.25 US**  
**61/751,816 2013.01.11 US**

(43) Publication Date: **2014.07.17**  
(44) Accepted Journal Date: **2017.11.16**

(71) Applicant(s)  
**Impossible Foods Inc.**

(72) Inventor(s)  
**Fraser, Rachel;Brown, Patrick O'Reilly;Karr, Jessica;Holz-Schietinger, Celeste;Cohn, Elysia**

(74) Agent / Attorney  
**Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000, AU**

(56) Related Art  
**US 2008268112 A1**  
**US 2009/0274817 A1**  
**US 3829582 A**  
**US 5443852 A**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/110532 A3

(43) International Publication Date

17 July 2014 (17.07.2014)

(51) International Patent Classification:

A23D 7/005 (2006.01) A23L 1/30 (2006.01)  
A23D 9/013 (2006.01)

(21) International Application Number:

PCT/US2014/011347

(22) International Filing Date:

13 January 2014 (13.01.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/751,816	11 January 2013 (11.01.2013)	US
13/941,211	12 July 2013 (12.07.2013)	US
61/908,634	25 November 2013 (25.11.2013)	US

(71) Applicant: MARAXI, INC. [US/US]; 525 Chesapeake Drive, Redwood City, California 94063 (US).

(72) Inventors; and

(71) Applicants : FRASER, Rachel [US/US]; 3653 24th Street, Apt. 2, San Francisco, California 94110 (US). BROWN, Patrick O'Reilly [US/US]; 76 Peter Coutts Circle, Stanford, California 94305 (US). KARR, Jessica [US/US]; 456 Kansas Street, San Francisco, California 94107 (US). HOLZ-SCHIETINGER, Celeste [US/US]; 1159 Sage Street, East Palo Alto, California 94303 (US). COHN, Elysia [US/US]; 1885 California Street, Apt. 49, Mountain View, California 94041 (US).

(74) Agents: LAVOIE, Teresa et al.; Fish & Richardson P.C., P.O. Box 1022, Minneapolis, Minnesota 55440 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

4 September 2014

WO 2014/110532 A3

(54) Title: METHODS AND COMPOSITIONS FOR AFFECTING THE FLAVOR AND AROMA PROFILE OF CONSUMABLES

(57) Abstract: This document relates to food products containing highly conjugated heterocyclic rings complexed to an iron ion and one or more flavor precursors, and using such food products to modulate the flavor and/or aroma profile of other foods. The food products described herein can be prepared in various ways and can be formulated to be free of animal products.

## METHODS AND COMPOSITIONS FOR AFFECTING THE FLAVOR AND AROMA PROFILE OF CONSUMABLES

### **CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims priority to U.S. Application Serial No. 13/941,211, filed July 12, 2013, U.S. Application Serial No. 61/908,634, filed November 25, 2013, and to U.S. Application Serial No. 61/751,816, filed January 11, 2013, and is related to the following 5 co-pending patent applications: Application Serial No. PCT/US12/46560; Application Serial No PCT/US12/46552; Application Serial No. 61,876,676, filed September 11, 2013; and Application Serial No. 61/751,818, filed January 11, 2013, all of which are incorporated herein by reference.

10

### **TECHNICAL FIELD**

This invention relates to food products and more particularly, to food products that include a highly conjugated heterocyclic ring complexed to iron such as a heme-cofactor and one or more flavor precursor molecules.

15

### **BACKGROUND**

20

Food is any substance that is either eaten or drunk by any animal, including humans, for nutrition or pleasure. It is usually of plant or animal origin, and can contain essential nutrients, such as carbohydrates, fats, proteins, vitamins, or minerals. The substance is ingested by an organism and assimilated by the organism's cells in an effort to produce energy, maintain life, or stimulate growth.

Food typically has its origin in a photosynthetic organism, such as a plant. Some food is obtained directly from plants, but even animals that are used as food sources are raised by feeding them food which is typically derived from plants.

25

In most cases, the plant or animal food source is fractionated into a variety of different portions, depending upon the purpose of the food. Often, certain portions of the plant, such as the seeds or fruits, are more highly prized by humans than others and these

are selected for human consumption, while other less desirable portions, such as the stalks of grasses, are typically used for feeding animals.

Current plant-based meat substitutes have largely failed to cause a shift to a vegetarian diet. Meat substitute compositions are typically extruded soy/grain mixtures which largely fail to replicate the experience of cooking and eating meat. Common limitations of plant-based meat substitute products are a texture and mouth-feel that are more homogenous than that of equivalent meat products. Furthermore, as these products must largely be sold pre-cooked, with artificial flavors and aromas pre-incorporated, they fail to replicate the aromas, flavors, and other key features, such as texture and mouth-feel, associated with cooking or cooked meat. As a result, these products appeal largely to a limited consumer base that is already committed to vegetarianism/veganism, but have failed to appeal to the larger consumer segment accustomed to eating meat. It would be useful to have improved plant-based meat substitutes which better replicate the aromas and flavors of meat, particularly during and/or after cooking.

## SUMMARY

In a first aspect, the present invention provides a ground beef-like food product, comprising: a) 0.1%-5% by weight of a heme-containing protein; b) a compound selected from glucose, ribose, fructose, lactose, xylose, arabinose, glucose-6-phosphate, maltose, and galactose, and mixtures of two or more thereof; c) at least 1.5 mM of a compound selected from cysteine, cysteine, thiamine, methionine, and mixtures of two or more thereof; and d) one or more plant proteins, wherein the ground beef-like food product contains no animal product; wherein cooking the ground beef-like food product results in the production of at least two volatile compounds which have a beef-associate aroma.

Provided herein are methods and compositions for modulating the flavor and/or aroma profile of consumable food products, including animal- or non-animal (e.g., plant) based food products, or mixtures of animal- and non-animal-based food products. In some embodiments, the methods and compositions are useful for modulating the flavor and/or aroma profile of a consumable food product during and/or after the cooking process. In some embodiments, the methods and compositions are used to generate one or more chemical compounds that modulate the flavor and/or aroma profile of the consumable food product during and/or after the cooking process.

As provided herein, and without being bound by theory, certain characteristic meaty flavors and/or aromas (e.g., beefy, bacony, umami, savory, bloody, brothy, gravy, metallic, bouillon-like; see Tables 2, 7, and 11), including one or more specific chemical compounds

2a

associated with the same (see Tables 3, 8, 9, 12, 14, 16, or 17), are believed to be produced during the cooking process of a consumable food product by chemical reaction of one or more flavor precursor molecules or compositions catalyzed by the

presence of a highly conjugated heterocyclic ring complexed to an iron ion (e.g., a heme moiety; or a porphyrin; a porphyrinogen; a corrin; a corrinoid; a chlorin; a bacteriochlorophyll; a corphin; a chlorophyllin; a bacteriochlorin; or an isobacteriochlorin moiety complexed to an iron ion). Such highly conjugated heterocyclic moieties 5 include heterocyclic aromatic rings composed of one or more (2, 3, or 4 more) pyrrole, pyrrole-like, and/or pyrroline subunits. The highly conjugated heterocyclic ring complexed to an iron ion is referred to herein as an iron complex. In some embodiments, the heme moiety can be a heme cofactor such as a heme moiety bound to a protein; a heme moiety bound to a non-proteinaceous polymer; a heme moiety bound to a solid 10 support; or a heme moiety encapsulated in a liposome. In some embodiments, the flavors and/or aromas are not generated in the absence of the iron complex (e.g., in the absence of a ferrous chlorin) or are not generated in the absence of a heme-cofactor (e.g., in the absence of a heme-containing protein). Accordingly, as described herein, the iron complexes such as isolated chlorin-iron complexes or heme-cofactors (e.g., heme-containing proteins) can be used to generate meaty flavors and/or aromas in a variety of 15 food products, such as during the cooking process.

Combining one or more iron complexes such as a heme-cofactor (e.g., a heme-containing protein, including, for example a plant-derived heme protein such as a plant leghemoglobin (legH)), with one or more flavor precursor molecules or compositions 20 (see, e.g., Table 1 or Table 13) can generate or provide a range of savory and meaty aromas and tastes (see, e.g., Tables 2, 7, and/or 11) in a cooked consumable food product. Flavor precursor molecules or compositions can be added to the uncooked food product in purified form and/or can be derived from ingredients in the uncooked consumable food product that contain and/or are enriched with one or more of the particular flavor 25 precursors or compositions, including, for example, yeast extract, vegetable oil, corn oil, soybean oil, palm fruit oil, palm kernel oil, safflower oil, flaxseed oil, rice bran oil, cottonseed oil, olive oil, canola oil, sunflower oil, coconut oil, mango oil, or an algal oil. The resultant flavor and/or aroma profile can be modulated by the type and concentration 30 of the flavor precursors, the pH of the reaction, the length of cooking, the type and amount of iron complex (e.g., a heme cofactor such as a heme-containing protein), the

temperature of the reaction, and the amount of water activity in the product, among other factors.

One or more flavor precursor molecules or compositions can be added along with a iron complex (e.g., ferrous chlorophyllin or a heme cofactor such as a heme-containing protein), to an uncooked food product, before and/or during the cooking process, to give the cooked consumable food product a particular meaty taste and smell, for example, the taste and smell of beef, bacon, pork, lamb, or chicken. Consumable food products can be animal or non-animal based (e.g., plant) food products, or combinations of an animal and non-animal based food product. For example, a plant based veggie burger or an animal-based burger, such as a chicken burger, can be modified with the compositions and methods of the present disclosure to result in a burger having a cooked flavor and/or aroma profile that is more meat like, e.g., beef-like, lamb-like, pork-like, turkey-like, duck-like, deer-like, yak-like, bison-like or other desirable meat flavor.

Food products for use in the present disclosure include those that have an iron-complex (e.g., a heme cofactor such as a heme-containing protein), and one or more flavor precursor molecules included therein. The iron-complex such as a heme cofactor (e.g., a heme-containing protein) and the one or more flavor precursor molecules can be homogenously or heterogeneously included in the food products. A heme protein can be isolated and purified prior to inclusion in the food product. Non-limiting examples of consumable food products which can include an iron complex such as a heme-cofactor (e.g., a heme-containing protein) and one or more flavor precursor molecules include animal-based or non-animal (e.g., plant-based), or combinations of animal-based and non-animal-based, food products in the form of hot dogs, burgers, ground meat, sausages, steaks, filets, roasts, breasts, thighs, wings, meatballs, meatloaf, bacon, strips, fingers, nuggets, cutlets, or cubes.

Consumable food products for use in the present disclosure can be flavor additive compositions, e.g., for addition to another consumable food product before, during, or after its cooking process. A flavor additive composition can include an iron complex such as a heme-cofactor (e.g., a heme-containing protein), and one or more flavor precursors.

A flavor additive composition can include a heme protein, e.g., an isolated and purified heme protein; such a flavor additive composition can be used to modulate the flavor and/or aroma profile of a consumable food product that comprises one or more flavor precursor molecules or compositions. A flavor additive composition can include 5 one or more flavor precursor molecules or compositions; such a flavor additive composition can be used to modulate the flavor and/or aroma profile of a consumable food product that comprises the heme protein, e.g., an isolated and purified heme protein.

A flavor additive composition can be in the form, of but not limited to, soup or stew bases, bouillon, e.g., powder or cubes, flavor packets, or seasoning packets or 10 shakers. Such flavor additive compositions can be used to modulate the flavor and/or aroma profile for a variety of consumable food products, and can be added to a consumable food product before, during, or after cooking of the consumable food product.

In some embodiments, a flavor additive composition such as one including an 15 iron complex (e.g., ferrous chlorin or a heme protein) and one or more flavor precursors can be reacted (e.g., *in vitro*) with heating to generate a particular flavor and/or aroma profile of interest and the resultant product mixture can be added to the consumable food product of interest, which can then be eaten as-is or can be additionally modified, e.g., by additional cooking. In some embodiments, the iron complex can be removed from the 20 resultant product mixture before adding the product mixture to the consumable food product of interest. For example, the iron complex can be removed from the product mixture using chromatographic techniques such as column chromatography, e.g., a column containing heme or iron-chlorin.

In some embodiments, the iron complex such as a heme-cofactor, e.g., a heme- 25 protein, and the one or more flavor precursor flavor additive compositions can be soy-free, wheat-free, yeast-free, MSG-free, and free of protein hydrolysis products, and can taste meaty, highly savory, and without off odors or flavors.

In one aspect, this document features a food product that includes an iron complex such as a heme moiety, or a porphyrin, a porphyrinogen, a corrin, a corrinoid, a chlorin, a 30 bacteriochlorophyll, a corphin, a chlorophyllin, a bacteriochlorin, or an isobacteriochlorin

moiety complexed to an iron ion and one or more flavor precursor molecules selected from the group consisting of glucose, fructose, ribose, arabinose, glucose-6-phosphate, fructose 6-phosphate, fructose 1,6-diphosphate, inositol, maltose, sucrose, maltodextrin, glycogen, nucleotide-bound sugars, molasses, a phospholipid, a lecithin, inosine, inosine monophosphate (IMP), guanosine monophosphate (GMP), pyrazine, adenosine monophosphate (AMP), lactic acid, succinic acid, glycolic acid, thiamine, creatine, pyrophosphate, vegetable oil, algal oil, corn oil, soybean oil, palm fruit oil, palm kernel oil, safflower oil, flaxseed oil, rice bran oil, cottonseed oil, sunflower oil, canola oil, olive oil, a free fatty acid, cysteine, methionine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophan, valine, arginine, histidine, alanine, asparagine, aspartate, glutamate, glutamine, glycine, proline, serine, tyrosine, glutathione, an amino acid derivative, a protein hydrolysate, a malt extract, a yeast extract, and a peptone. The heme moiety can be a heme-containing protein, a heme moiety bound to a non-peptidic polymer; or a heme moiety bound to a solid support. The heme-containing protein can be a plant, mammalian, a yeast or filamentous fungi, or bacterial heme-containing protein.

The food product can include two to one hundred, two to fifty flavor precursors, two to forty flavor precursors, two to thirty-five flavor precursors, two to ten flavor precursors, or two to six flavor precursors. In some embodiments, the one or more flavor precursor molecules are selected from the group consisting of glucose, ribose, cysteine, a cysteine derivative, thiamine, alanine, methionine, lysine, a lysine derivative, glutamic acid, a glutamic acid derivative, IMP, GMP, lactic acid, maltodextrin, creatine, alanine, arginine, asparagine, aspartate, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, methionine, phenylalanine, proline, threonine, tryptophan, tyrosine, valine, linoleic acid, and mixtures thereof. The heme-containing protein can be a non-symbiotic hemoglobin or a leghemoglobin (e.g., a plant leghemoglobin such as one from soybean, alfalfa, lupin, pea, cow pea, or lupin). The heme-containing protein can include an amino acid sequence having at least 80% sequence identity to a polypeptide set forth in SEQ ID NOs:1-26.

The heme-containing protein can be isolated and purified. The food product further can include a food-grade oil, a seasoning agent, a flavoring agent, a protein, a protein concentrate, an emulsifier, a gelling agent, or a fiber. The food product can be a meat

substitute, a soup base, stew base, snack food, bouillon powder, bouillon cube, a flavor packet, or a frozen food product. Any of the food products can be free of animal products. The food product can be sealed within a packet or shaker.

This document also features a method for producing a flavor compound. The method can include combining an iron complex (e.g., a heme moiety, a porphyrin, a porphyrinogen, a corrin, a corrinoid, a chlorin, a bacteriochlorophyll, a corphin, a chlorophyllin, a bacteriochlorin, or an isobacteriochlorin complexed to an iron) and one or more flavor precursor molecules to form a mixture, the one or more flavor precursor molecules selected from the group consisting of glucose, fructose, arabinose, ribose 5  
glucose-6-phosphate, fructose 6-phosphate, fructose 1,6-diphosphate, inositol, maltose, sucrose, maltodextrin, glycogen, nucleotide-bound sugars, molasses, a phospholipid, a lecithin, inosine, inosine monophosphate (IMP), guanosine monophosphate (GMP), pyrazine, adenosine monophosphate (AMP), lactic acid, succinic acid, glycolic acid, thiamine, creatine, pyrophosphate, vegetable oil, algal oil, corn oil, soybean oil, palm 10  
fruit oil, palm kernel oil, safflower oil, flaxseed oil, rice bran oil, cottonseed oil, canola oil, olive oil, sunflower oil, flaxseed oil, coconut oil, mango oil, a free fatty acid, cysteine, methionine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophan, valine, arginine, histidine, alanine, asparagine, aspartate, glutamate, glutamine, glycine, 15  
proline, serine, tyrosine, glutathione, an amino acid derivative, a protein hydrolysate, a malt extract, a yeast extract, and a peptone; and heating the mixture to form one or more flavor compounds selected from the group consisting of phenylacetaldehyde, 1-octen-3-one, 2-n-heptylfuran, 2-thiophenecarboxaldehyde, 3-thiophenecarboxaldehyde, butyrolactone, 2-undecenal, pyrazine, methyl-, furfural, 2-decanone, pyrrole, 1-octen-3-ol, 2-acetylthiazole, (E)-2-octenal, decanal, benzaldehyde, (E)-2-nonenal, pyrazine, 1- 20  
hexanol, 1-heptanol, dimethyl trisulfide, 2-nonenone, 2-pentanone, 2-heptanone, 2,3-butanedione, heptanal, nonanal, 2-octanone, 1-octanol, 3-ethylcyclopentanone, 3-octen-2-one, (E,E)-2,4-heptadienal, (Z)-2-heptenal, 2-heptanone, 6-methyl-, (Z)-4-heptenal, (E,Z)-2,6-nonadienal, 3-methyl-2-butenal, 2-pentyl-furan, thiazole, (E, E)-2,4-decadienal, hexanoic acid, 1-ethyl-5-methylcyclopentene, (E,E)-2,4-nonadienal, (Z)-2-decenal, dihydro-5-pentyl-2(3H)-furanone, trans-3-nonen-2-one, (E,E)-3,5-octadien-2- 25  
30

one, (Z)-2-octen-1-ol, 5-ethyldihydro-2(3H)-furanone, 2-butenal, 1-penten-3-ol, (E)-2-hexenal, formic acid, heptyl ester, 2-pentyl-thiophene, (Z)-2-nonenal, 2-hexyl-thiophene, (E)-2-decenal, 2-ethyl-5-methyl-pyrazine, 3-ethyl-2,5-dimethyl-pyrazine, 2-ethyl-1-hexanol, thiophene, 2-methyl-furan, pyridine, butanal, 2-ethyl-furan, 3-methyl-butanal, 5 trichloromethane, 2-methyl-butanal, methacrolein, 2-methyl-propanal, propanal, acetaldehyde, 2-propyl-furan, dihydro-5-propyl-2(3H)-furanone, 1,3-hexadiene, 4-decyne, pentanal, 1-propanol, heptanoic acid, trimethyl-ethanethiol, , 1-butanol, 1-penten-3-one, dimethyl sulfide, 2-ethyl furan, 2-pentyl-thiophene, 2-propenal, 2-tridecen-1-ol, 4-octene, 2-methyl thiazole,, methyl-pyrazine, 2-butanone, 2-pentyl-furan, 2-methyl-propanal, butyrolactone,, 3-methyl-butanal, methyl-thiirane, 2-hexyl-furan, 10 butanal,, 2-methyl-butanal, 2-methyl-furan, furan, octanal, 2-heptenal, 1-octene, formic acid heptyl ester, 3-pentyl-furan, and 4-penten-2-one. The heme moiety can be a heme-containing protein, a heme moiety bound to a non-peptidic polymer; or a heme moiety bound to a solid support. The method can include combining cysteine, ribose, lactic acid, 15 lysine, and/or thiamine with the heme-containing protein.

In another aspect, this document features a method for producing a flavor compound. The method includes combining an iron complex, such as a heme-containing protein, and one or more flavor precursor molecules to form a mixture, the one or more flavor precursor molecules selected from the group consisting of glucose, fructose, 20 ribose, arabinose, glucose-6-phosphate, fructose 6-phosphate, fructose 1,6-diphosphate, inositol, maltose, sucrose, maltodextrin, glycogen, nucleotide-bound sugars, molasses, a phospholipid, a lecithin, inosine, IMP, GMP, pyrazine, AMP, lactic acid, succinic acid, glycolic acid, thiamine, creatine, pyrophosphate, vegetable oil, algal oil, corn oil, soybean oil, palm fruit oil, palm kernel oil, safflower oil, flaxseed oil, rice bran oil, cottonseed oil, 25 olive oil, sunflower oil, canola oil, flaxseed oil, coconut oil, mango oil, a free fatty acid, methionine, cysteine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophan, valine, arginine, histidine, alanine, asparagine, aspartate, glutamate, glutamine, glycine, proline, serine, tyrosine, glutathione, an amino acid derivative, a protein hydrolysate, a malt extract, a yeast extract, and a peptone; and heating the mixture to form one or more

flavor compounds set forth in Tables 3, 8, or 9. For example, the flavor precursors can include cysteine, a sugar, and one or more other precursors.

This document also features a method for imparting a meat like flavor (e.g., beef-like, chicken like, pork-like, lamb-like, turkey-like, duck-like, deer-like, or bison-like) to a food product. The method includes contacting the food product with a flavoring composition, the flavoring composition comprising i) an iron complex, such as a heme moiety (e.g., a heme-containing protein); and ii) one or more flavor precursor molecules selected from the group consisting of glucose, fructose, ribose, arabinose, glucose-6-phosphate, fructose 6-phosphate, fructose 1,6-diphosphate, inositol, maltose, sucrose, maltodextrin, glycogen, nucleotide-bound sugars, molasses, a phospholipid, a lecithin, inosine, IMP, GMP, pyrazine, AMP, lactic acid, succinic acid, glycolic acid, thiamine, creatine, pyrophosphate, vegetable oil, algal oil, corn oil, soybean oil, palm fruit oil, palm kernel oil, safflower oil, flaxseed oil, rice bran oil, cottonseed oil, olive oil, sunflower oil, canola oil, flaxseed oil, coconut oil, mango oil, a free fatty acid, cysteine, methionine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophan, valine, arginine, histidine, alanine, asparagine, aspartate, glutamate, glutamine, glycine, proline, serine, tyrosine, glutathione, an amino acid derivative, a protein hydrolysate, a malt extract, a yeast extract, and a peptone; wherein after heating the food product and the flavoring composition together, a meat like flavor (e.g., beef-like, chicken like, pork-like, lamb-like, turkey-like, duck-like, deer-like, or bison-like) is imparted to the food product. In some embodiments, the iron complex is removed from the food product. The flavoring composition further can include a seasoning agent, a flavoring agent, a protein, a protein concentrate, or an emulsifier. The flavoring composition can be sealed within a packet or shaker.

In another aspect, this document features a method of making a food product. The method includes combining an isolated heme-containing protein and one or more flavor precursor molecules to form a mixture, the one or more flavor precursor molecules selected from the group consisting of glucose, fructose, ribose, arabinose, glucose-6-phosphate, fructose 6-phosphate, fructose 1,6-diphosphate, inositol, maltose, sucrose, maltodextrin, glycogen, nucleotide-bound sugars, molasses, a phospholipid, a lecithin,

5 inosine, IMP, GMP, pyrazine, AMP, lactic acid, succinic acid, glycolic acid, thiamine, creatine, pyrophosphate, sunflower oil, coconut oil, canola oil, flaxseed oil, mango oil, a free fatty acid, cysteine, methionine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophan, valine, arginine, histidine, alanine, asparagine, aspartate, glutamate, glutamine, glycine, proline, serine, tyrosine, glutathione, an amino acid derivative, a protein hydrolysate, a malt extract, a yeast extract, and a peptone; and heating the mixture.

10 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used to practice the invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, 15 methods, and examples are illustrative only and not intended to be limiting.

20 The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims. The word “comprising” in the claims may be replaced by “consisting essentially of” or with “consisting of,” according to standard practice in patent law.

## **DESCRIPTION OF THE DRAWINGS**

FIG. 1 contains amino acid sequences of exemplary heme-containing proteins.

25 FIG. 2 is a bar graph of the beefiness rating of the meat replica with or without the Magic Mix, both samples in triplicate with 1% w/v LegH protein. Tasters rated beefiness on a scale from 1-7, with 1 being not beefy at all and 7 being exactly like ground beef.

## **DETAILED DESCRIPTION**

This document is based on methods and materials for modulating the taste and/or aroma profile of food products. As described herein, compositions containing one or

more flavor precursors and one or more highly conjugated heterocyclic rings complexed to an iron (referred to herein as an iron complex) can be used to modulate the taste and/or aroma profile of food products. Such iron complexes include heme moieties or other highly conjugated heterocyclic rings complexed to an iron ion (referred to as an iron complex). “Heme” refers to a prosthetic group bound to iron ( $Fe^{2+}$  or  $Fe^{3+}$ ) in the center of a porphyrin ring. Thus, an iron complex can be a heme moiety, or a porphyrin, porphyrinogen, corrin, corrinoid, chlorin, bacteriochlorophyll, corphin, chlorophyllin, bacteriochlorin, or isobacteriochlorin moiety complexed to iron ion. The heme moiety that can be used to modulate the taste and/or aroma profile of food products can be a heme cofactor such as a heme-containing protein; a heme moiety bound to a non-peptidic polymer or other macromolecule such as a liposome, a polyethylene glycol, a carbohydrate, a polysaccharide, a cyclodextrin, a polyethylenimine, a polyacrylate, or derivatives thereof; a siderophore (i.e., an iron chelating compound); or a heme moiety bound to a solid support (e.g., beads) composed of a chromatography resin, cellulose, graphite, charcoal, or diatomaceous earth.

In some embodiments, the iron complexes catalyze some reactions and produce flavor precursors without heating or cooking. In some embodiments, the iron complex destabilizes upon heating or cooking and releases the iron, e.g., the protein is denatured, so flavor precursors can be generated.

Suitable flavor precursors include sugars, sugar alcohols, sugar derivatives, oils (e.g., vegetable oils), free fatty acids, alpha-hydroxy acids, dicarboxylic acids, amino acids and derivatives thereof, nucleosides, nucleotides, vitamins, peptides, protein hydrolysates, extracts, phospholipids, lecithin, and organic molecules. Non-limiting examples of such flavor precursors are provided in Table 1.

25

**TABLE 1****Flavor Precursor Molecules**

Sugars, sugar alcohols, sugar acids, and sugar derivatives: glucose, fructose, ribose, sucrose, arabinose, glucose-6-phosphate, fructose-6-phosphate, fructose 1,6-diphosphate, inositol, maltose, molasses, maltodextrin, glycogen, galactose,

lactose, ribitol, gluconic acid and glucuronic acid, amylose, amylopectin, or xylose
Oils: coconut oil, mango oil, sunflower oil, cottonseed oil, safflower oil, rice bran oil, cocoa butter, palm fruit oil, palm oil, soybean oil, canola oil, corn oil, sesame oil, walnut oil, flaxseed, jojoba oil, castor, grapeseed oil, peanut oil, olive oil, algal oil, oil from bacteria or fungi
Free fatty acids: caprylic acid, capric acid, lauric acid, myristic acid, palmititic acid, palmitoleic acid, stearic, oleic acid, linoleic acid, alpha linolenic acid, gamma linolenic acid, arachidic acid, arachidonic acid, behenic acid, or erucic acid
Amino acids and derivatives thereof: cysteine, cystine, a cysteine sulfoxide, allicin, selenocysteine, methionine, isoleucine, leucine, lysine, phenylalanine, threonine, tryptophan, 5-hydroxytryptophan, valine, arginine, histidine, alanine, asparagine, aspartate, glutamate, glutamine, glycine, proline, serine, or tyrosine
Nucleosides and Nucleotides: inosine, inosine monophosphate (IMP), guanosine, guanoside monophosphate (GMP), adenosine, adenosine monophosphate (AMP)
Vitamins: thiamine, vitamin C, Vitamin D, Vitamin B6, or Vitamin E
Misc: phospholipid, lecithin, pyrazine, creatine, pyrophosphate
Acids: acetic acid, alpha hydroxy acids such as lactic acid or glycolic acid, tricarboxylic acids such as citric acid, dicarboxylic acids such as succinic acid or tartaric acid
Peptides and protein hydrolysates: glutathione, vegetable protein hydrolysates, soy protein hydrolysates, yeast protein hydrolysates, algal protein hydrolysates, meat protein hydrolysates
Extracts: a malt extract, a yeast extract, and a peptone

In some embodiments, one flavor precursor or combinations of two to one hundred flavor precursors, two to ninety, two to eighty, two to seventy, two to sixty, or two to fifty flavor precursors are used. For example, combinations of two to forty flavor

precursors, two to thirty-five flavor precursors, two to ten flavor precursors, or two to six flavor precursors can be used with the one or more iron complexes (e.g., heme co-factors such as a heme-containing proteins). For example, the one or more flavor precursors can be glucose, ribose, cysteine, a cysteine derivative, thiamine, lysine, a lysine derivative, 5 glutamic acid, a glutamic acid derivative, alanine, methionine, IMP, GMP, lactic acid, and mixtures thereof (e.g., glucose and cysteine; cysteine and ribose; cysteine, glucose or ribose, and thiamine; cysteine, glucose or ribose, IMP, and GMP; cysteine, glucose or ribose, and lactic acid). For example, the one or more flavor precursors can be alanine, 10 arginine, asparagine, aspartate, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, threonine, tryptophan, tyrosine, valine, glucose, ribose, maltodextrin, thiamine, IMP, GMP, lactic acid, and creatine.

As used herein, the term “heme containing protein” can be used interchangeably with “heme containing polypeptide” or “heme protein” or “heme polypeptide” and 15 includes any polypeptide that can covalently or noncovalently bind a heme moiety. In some embodiments, the heme-containing polypeptide is a globin and can include a globin fold, which comprises a series of seven to nine alpha helices. Globin type proteins can be of any class (e.g., class I, class II, or class III), and in some embodiments, can transport or store oxygen. For example, a heme-containing protein can be a non-symbiotic type of 20 hemoglobin or a leghemoglobin. A heme-containing polypeptide can be a monomer, i.e., a single polypeptide chain, or can be a dimer, a trimer, tetramer, and/or higher order oligomers. The life-time of the oxygenated  $Fe^{2+}$  state of a heme-containing protein can be similar to that of myoglobin or can exceed it by 10%, 20%, 30% 50%, 100% or more under conditions in which the heme-protein-containing consumable is manufactured, 25 stored, handled or prepared for consumption. The life-time of the unoxygenated  $Fe^{2+}$  state of a heme-containing protein can be similar to that of myoglobin or can exceed it by 10%, 20%, 30% 50%, 100% or more under conditions in which the heme-protein-containing consumable is manufactured, stored, handled or prepared for consumption

30 Non-limiting examples of heme-containing polypeptides can include an androglobin, a cytoglobin, a globin E, a globin X, a globin Y, a hemoglobin, a

myoglobin, an erythrocytins, a beta hemoglobin, an alpha hemoglobin, a protoglobin, a cyanoglobin, a cytoglobin, a histoglobin, a neuroglobins, a chlorocytins, a truncated hemoglobin (e.g., HbN or HbO), a truncated 2/2 globin, a hemoglobin 3 (e.g., Glb3), a cytochrome, or a peroxidase.

5 Heme-containing proteins that can be used in the compositions and food products described herein can be from mammals (e.g., farm animals such as cows, goats, sheep, pigs, ox, or rabbits), birds, plants, algae, fungi (e.g., yeast or filamentous fungi), ciliates, or bacteria. For example, a heme-containing protein can be from a mammal such as a farm animal (e.g., a cow, goat, sheep, pig, ox, or rabbit) or a bird such as a turkey or chicken. Heme-containing proteins can be from a plant such as *Nicotiana tabacum* or *Nicotiana sylvestris* (tobacco); *Zea mays* (corn), *Arabidopsis thaliana*, a legume such as *Glycine max* (soybean), *Cicer arietinum* (garbanzo or chick pea), *Pisum sativum* (pea) varieties such as garden peas or sugar snap peas, *Phaseolus vulgaris* varieties of common beans such as green beans, black beans, navy beans, northern beans, or pinto beans, *Vigna unguiculata* varieties (cow peas), *Vigna radiata* (Mung beans), *Lupinus albus* (lupin), or *Medicago sativa* (alfalfa); *Brassica napus* (canola); *Triticum* sps. (wheat, including wheat berries, and spelt); *Gossypium hirsutum* (cotton); *Oryza sativa* (rice); *Zizania* sps. (wild rice); *Helianthus annuus* (sunflower); *Beta vulgaris* (sugarbeet); *Pennisetum glaucum* (pearl millet); *Chenopodium* sp. (quinoa); *Sesamum* sp. (sesame); *Linum usitatissimum* (flax); or *Hordeum vulgare* (barley). Heme-containing proteins can be isolated from fungi such as *Saccharomyces cerevisiae*, *Pichia pastoris*, *Magnaporthe oryzae*, *Fusarium graminearum*, *Aspergillus oryzae*, *Trichoderma reesei*, *Mycelopthora thermophile*, *Kluyvera lactis*, or *Fusarium oxysporum*. Heme-containing proteins can be isolated from bacteria such as *Escherichia coli*, *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus megaterium*, *Synechocystis* sp., *Aquifex aeolicus*, *Methylacidiphilum infernorum*, or thermophilic bacteria such as *Thermophilus*. The sequences and structure of numerous heme-containing proteins are known. See for example, Reedy, *et al.*, *Nucleic Acids Research*, 2008, Vol. 36, Database issue D307–D313 and the Heme Protein Database available on the world wide web at <http://hemeprotein.info/heme.php>.

For example, a non-symbiotic hemoglobin can be from a plant selected from the group consisting of soybean, sprouted soybean, alfalfa, golden flax, black bean, black eyed pea, northern, garbanzo, moong bean, cowpeas, pinto beans, pod peas, quinoa, sesame, sunflower, wheat berries, spelt, barley, wild rice, or rice.

5 Any of the heme-containing proteins described herein that can be used for producing food products can have at least 70% (e.g., at least 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100%) sequence identity to the amino acid sequence of the corresponding wild-type heme-containing protein or fragments thereof that contain a heme-binding motif. For example, a heme-containing protein can have at least 70% 10 sequence identity to an amino acid sequence set forth in FIG. 1, including a non-symbiotic hemoglobin such as that from *Vigna radiata* (SEQ ID NO:1), *Hordeum vulgare* (SEQ ID NO:5), *Zea mays* (SEQ ID NO:13), *Oryza sativa* subsp. *japonica* (rice) (SEQ ID NO:14), or *Arabidopsis thaliana* (SEQ ID NO:15), a Hell's gate globin I such as that from *Methylacidiphilum infernorum* (SEQ ID NO:2), a flavohemoprotein such as that from *Aquifex aeolicus* (SEQ ID NO:3), a leghemoglobin such as that from *Glycine max* (SEQ ID NO:4), *Pisum sativum* (SEQ ID NO:16), or *Vigna unguiculata* (SEQ ID NO:17), a heme-dependent peroxidase such as from *Magnaporthe oryzae*, (SEQ ID NO:6) or *Fusarium oxysporum* (SEQ ID NO:7), a cytochrome c peroxidase from *Fusarium graminearum* (SEQ ID NO:8), a truncated hemoglobin from *Chlamydomonas moewusii* (SEQ ID NO:9), *Tetrahymena pyriformis* (SEQ ID NO:10, group I truncated), *Paramecium caudatum* (SEQ ID NO:11, group I truncated), a hemoglobin from *Aspergillus niger* (SEQ ID NO:12), or a mammalian myoglobin protein such as the *Bos taurus* (SEQ ID NO:18) myoglobin, *Sus scrofa* (SEQ ID NO:19) myoglobin, *Equus caballus* (SEQ ID NO:20) myoglobin, a heme-protein from *Nicotiana benthamiana* (SEQ ID NO:21), *Bacillus subtilis* (SEQ ID NO:22), *Corynebacterium glutamicum* (SEQ ID NO:23), *Synechocystis* PCC6803 (SEQ ID NO:24), *Synechococcus* sp. PCC 7335 (SEQ ID NO:25), or *Nostoc commune* (SEQ ID NO:26).

30 The percent identity between two amino acid sequences can be determined as follows. First, the amino acid sequences are aligned using the BLAST 2 Sequences (Bl2seq) program from the stand-alone version of BLASTZ containing BLASTP version

2.0.14. This stand-alone version of BLASTZ can be obtained from Fish & Richardson's web site (e.g., [www.ncbi.nlm.nih.gov/blast/](http://www.ncbi.nlm.nih.gov/blast/)) or the U.S. government's National Center for Biotechnology Information web site ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). Instructions explaining how to use the Bl2seq program can be found in the readme file accompanying BLASTZ.

5 Bl2seq performs a comparison between two amino acid sequences using the BLASTP algorithm. To compare two amino acid sequences, the options of Bl2seq are set as follows: -i is set to a file containing the first amino acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second amino acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastp; -o is set to any desired file name (e.g., C:\output.txt); and all other options are left at their default setting. For example, the following 10 command can be used to generate an output file containing a comparison between two amino acid sequences: C:\Bl2seq -i c:\seq1.txt -j c:\seq2.txt -p blastp -o c:\output.txt. If the two compared sequences share homology, then the designated output file will present those regions of homology as aligned sequences. If the two compared sequences do not 15 share homology, then the designated output file will not present aligned sequences.

Similar procedures can be followed for nucleic acid sequences except that blastn is used.

Once aligned, the number of matches is determined by counting the number of positions where an identical amino acid residue is presented in both sequences. The percent identity is determined by dividing the number of matches by the length of the 20 full-length polypeptide amino acid sequence followed by multiplying the resulting value by 100. It is noted that the percent identity value is rounded to the nearest tenth. For example, 78.11, 78.12, 78.13, and 78.14 is rounded down to 78.1, while 78.15, 78.16, 78.17, 78.18, and 78.19 is rounded up to 78.2. It also is noted that the length value will always be an integer.

25 It will be appreciated that a number of nucleic acids can encode a polypeptide having a particular amino acid sequence. The degeneracy of the genetic code is well known to the art; i.e., for many amino acids, there is more than one nucleotide triplet that serves as the codon for the amino acid. For example, codons in the coding sequence for a given enzyme can be modified such that optimal expression in a particular species (e.g., 30 bacteria or fungus) is obtained, using appropriate codon bias tables for that species.

Heme-containing proteins can be extracted from the source material (e.g., extracted from animal tissue, or plant, fungal, algal, or bacterial biomass, or from the culture supernatant for secreted proteins) or from a combination of source materials (e.g., multiple plant species). Leghemoglobin is readily available as an unused by-product of commodity legume crops (e.g., soybean, alfalfa, or pea). The amount of leghemoglobin in the roots of these crops in the United States exceeds the myoglobin content of all the red meat consumed in the United States.

In some embodiments, extracts of heme-containing proteins include one or more non-heme-containing proteins from the source material (e.g., other animal, plant, fungal, algal, or bacterial proteins) or from a combination of source materials (e.g., different animal, plant, fungi, algae, or bacteria).

In some embodiments, heme-containing proteins are isolated and purified from other components of the source material (e.g., other animal, plant, fungal, algal, or bacterial proteins). As used herein, the term “isolated and purified” indicates that the preparation of heme-containing protein is at least 60% pure, e.g., greater than 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% pure. Without being bound by theory, isolating and purifying proteins can allow the food products to be made with greater consistency and greater control over the properties of the food product as unwanted material is eliminated. Proteins can be separated on the basis of their molecular weight, for example, by size exclusion chromatography, ultrafiltration through membranes, or density centrifugation. In some embodiments, the proteins can be separated based on their surface charge, for example, by isoelectric precipitation, anion exchange chromatography, or cation exchange chromatography. Proteins also can be separated on the basis of their solubility, for example, by ammonium sulfate precipitation, isoelectric precipitation, surfactants, detergents or solvent extraction. Proteins also can be separated by their affinity to another molecule, using, for example, hydrophobic interaction chromatography, reactive dyes, or hydroxyapatite. Affinity chromatography also can include using antibodies having specific binding affinity for the heme-containing protein, nickel NTA for His-tagged recombinant proteins, lectins to bind to sugar moieties on a glycoprotein, or other molecules which specifically binds the protein.

Heme-containing proteins also can be recombinantly produced using polypeptide expression techniques (e.g., heterologous expression techniques using bacterial cells, insect cells, fungal cells such as yeast, plant cells such as tobacco, soybean, or *Arabidopsis*, or mammalian cells). In some cases, standard polypeptide synthesis techniques (e.g., liquid-phase polypeptide synthesis techniques or solid-phase polypeptide synthesis techniques) can be used to produce heme-containing proteins synthetically. In some cases, *in vitro* transcription-translation techniques can be used to produce heme-containing proteins.

The protein used in the consumable may be soluble in a solution. In some 10 embodiments, the isolated and purified proteins are soluble in solution at greater than 5, 10, 15, 20, 25, 50, 100, 150, 200, or 250g/L.

In some embodiments, the isolated and purified protein is substantially in its native fold and water soluble. In some embodiments, the isolated and purified protein is more than 50, 60, 70, 80, or 90% in its native fold. In some embodiments, the isolated 15 and purified protein is more than 50, 60, 70, 80, or 90% water soluble.

#### *Modulating Flavor and/or Aroma Profiles*

As described herein, different combinations of flavor precursors can be used with one or more iron complexes (e.g., a ferrous chlorin, a chlorin-iron complex, or a heme-cofactor such as a heme-containing protein or heme bound to a non-peptidic polymer such as polyethylene glycol or to a solid support) to produce different flavor and aroma profiles when the flavor precursors and iron complexes are heated together (e.g., during cooking). The resultant flavor and/or aroma profile can be modulated by the type and concentration of the flavor precursors, the pH of the reaction, the length of cooking, the 20 type and amount of iron complex (e.g., a heme-cofactor such as heme-containing protein, heme bound to non-peptidic polymer or macromolecule, or heme bound to a solid support), the temperature of the reaction, and the amount of water activity in the product, among other factors. In embodiments in which a heme moiety is bound to a solid support 25 such as cellulose or a chromatography resin, graphite, charcoal, or diatomaceous earth, the solid support (e.g., beads) can be incubated with sugars and/or one or more other 30

flavor precursors to generate flavors, and then the solid support with attached heme moiety can be re-used, i.e., incubated again with sugars and/or one or more other flavor precursors to generate flavors.

Table 2 provides non-limiting examples of flavor types that can be generated by combining one or more flavor precursors and one or more heme co-factors (e.g., heme-containing proteins). See also Tables 7 and/or 11.

**TABLE 2**

<b>Flavor Types</b>	
beef	beef broth
beef dripping	cheesy
cold-cut deli meat	squash
bacon	sharp
meaty	fruity
brothy	floral
ramen	musty
egg	fried food
malty	caramel
bready	barbeque
sulfur	chocolate
fried chicken	sweet
browned	potato
pretzel	french toast
grassy	breadcrust
bloody	mushroom
broccoli	chicken
brothy	cumin
buttery	umami
metallic	raisin

yeasty	goaty
vegetable broth	

Flavor and aroma profiles are created by different chemical compounds formed by chemical reactions between the heme co-factor (e.g., heme-containing protein) and flavor precursors. Gas chromatography-mass spectrometry (GCMS) can be used to separate and identify the different chemical compounds within a test sample. For example, volatile chemicals can be isolated from the head space after heating a heme-containing protein and one or more flavor precursors.

Table 3 provides non-limiting examples of compounds that can be produced. See also Tables 8, 9, 12, and/or 14.

10

**TABLE 3****Compounds Produced**

phenylacetaldehyde	2-butenal, 2-ethyl-	1,3-hexadiene
1-octen-3-one	acetonitrile	4-decyne
2-n-heptylfuran		pentanal
2-thiophenecarboxaldehyde	(E)-2-Hexenal	1-propanol
3-thiophenecarboxaldehyde	4-ethyl-phenol,	heptanoic acid
1-octene	3-octanone	ethanethiol
butyrolactone	styrene	2-methyl-1-heptene
2-undecenal	furan, 3-pentyl-	(E)-4-octene
propyl-cyclopropane	formic acid, heptyl ester	2-methyl-2-heptene
methyl-pyrazine	(E)-2-Heptenal	pentanoic acid
1-hydroxy-propanone	6-methyl-5-hepten-2-one	nonanoic acid
acetic acid	n-caproic acid vinyl ester	1,3-dimethyl-benzene
furfural	2-ethyl-2-hexenal	
2-decanone	1-hepten-3-ol	toluene
pyrrole	1-ethyl-1-methyl-cyclopentane	1-butanol
1-octen-3-ol	3-ethyl-2-methyl-1,3-hexadiene	2,3,3-trimethyl-pentane
2-acetylthiazole	2-pentyl-thiophene	isopropyl alcohol
(E)-2-octenal	(Z)-2-nonenal	2,2,4,6,6-pentamethyl-heptane
decanal	2-n-octylfuran	phenol

benzaldehyde	2-hexyl-thiophene	1-penten-3-one
(E)-2-Nonenal	4-cyclopentene-1,3-dione	dimethyl sulfide
pyrazine	1-nonanol	thiirane
1-pentanol	(E)-2-decenal	(E)-2-octen-1-ol
trans-2-(2-pentenyl)furan	4-ethyl-benzaldehyde	2,4-dimethyl-1-heptene
1-hexanol	1,7-octadien-3-ol	1,3-bis(1,1-dimethylethyl)-benzene
1-heptanol	octanoic acid	heptane
dimethyl trisulfide	2-ethyl-5-methyl-pyrazine	4,7-dimethyl-undecane
2-nonanone	3-ethyl-2,5-dimethyl-pyrazine	acetophenone
2-pentanone	1,3,5-cycloheptatriene	tridecane
2-heptanone	2-ethyl-1-hexanol	thiophosphoramide, s-methyl ester
2,3-butanedione	4-methyl-octanoic acid	2-methyl-thiazole
heptanal	m-aminophenylacetylene	3-(1-methylethoxy)-propanenitrile,
nonanal	benzene	2,4-bis(1,1-dimethylethyl)-phenol
2-octanone	thiophene	3-ethyl-2,2-dimethyl-pentane
2-butanone	2-methyl-furan	3-ethyl-pentane
octanal	pyridine	2,3,4-trimethyl-pentane
1-octanol	furan	2,4,6-trimethyl-octane
3-ethylcyclopentanone	butanal	2,6-dimethyl-nonane
8-methyl-1-undecene	2-ethyl-furan	2-hexyl-furan
3-octen-2-one	carbon disulfide	4-methyl-5-thiazoleethanol
2,4-Heptadienal, (E,E)-	Furan, 2-hexyl-:2	4-penten-2-one
(Z)-2-heptenal	3-methyl-butanal	4-methylthiazole
6-methyl-2-heptanone	2-methyl-butanal	2-methyl-3-pentanone
(Z)-4-heptenal	methacrolein	2,3-pentanedione
(E,Z)-2,6-nonadienal	octane	(E)-2-tridecen-1-ol
3-methyl-2-butenal	ethanol	2-thiophenemethanamine
2-pentyl-furan	2-methyl-propanal	(Z)-2-nonenal,
thiazole	acetone	methyl thiolacetate
(E,E)-2,4-decadienal	propanal	methyl ethanoate
hexanoic acid	methyl-thiirane	isothiazole
1-ethyl-5-methylcyclopentene	acetaldehyde	3,3-dimethyl-hexane

(E,E)-2,4-nonadienal	2-propenal	4-methyl-heptane
(Z)-2-decenal	2-propyl-furan	2,4-dimethyl-heptane
dihydro-5-pentyl-2(3h)-furanone	dihydro-5-propyl-2(3H)-furanone	2,3,4-trimethyl-heptane
trans-3-nonen-2-one	dihydro-3-(2H)-thiophenone	2-methyl-heptane
(E,E)-3,5-octadien-2-one	2,2,6-trimethyl-decane	2-methyl-3-furanthiol
(Z)-2-octen-1-ol	3,3'-dithiobis[2-methyl-furan	4-amino-1,2,5-oxadiazole-3-carbonitrile
5-ethyldihydro-2(3h)-furanone	1-heptene	1,2-benzisothiazol-3(2H)-one
2-butenal	1,3-octadiene	2-acetyl-propen-2-ol,
1-penten-3-ol	1-nonene	1-decen-3-one
1-(ethylthio)-2-(methylthio)-buta-1,3-diene		

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein is heated in the presence of ground chicken, to increase specific volatile flavor and odorant components typically elevated in beef. For example, propanal, butanal, 2-ethyl-furan, heptanal, octanal, trans-2-(2-pentenyl)furan, (Z)-2-heptenal, (E)-2-octenal, pyrrole, 2,4-dodecadienal, 1-octanal, (Z)-2-decenal, or 2-undecenal can be increased in the presence of the heme-containing protein, which can impart a more beefy flavor to the chicken.

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein is heated in the presence of cysteine and glucose or other combinations of flavor precursors to provide a different profile of volatile odorants than when any subset of the three components are used individually. Volatile flavor components that are increased under these conditions include but are not limited to furan, acetone, thiazole, furfural, benzaldehyde, 2-pyridinecarboxaldehyde, 5-methyl-2-thiophenecarboxaldehyde, 3-methyl-2-thiophenecarboxaldehyde, 3-thiophenemethanol and decanol. See, e.g., Tables 8 and 9. Under these conditions, cysteine and glucose alone or in the presence of iron salts such as ferrous glucanate produced a sulfurous, odor, but addition of heme-containing proteins reduced the

sulfurous odor and replaced it with flavors including but not limited to chicken broth, burnt mushroom, molasses, and bread.

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein is heated in the presence of cysteine and ribose to provide a different profile of volatile odorants. Heating in the presence of ribose created some additional compounds as compared to when a heme-containing protein and glucose were heated together. See Tables 8 and 9.

In some embodiments, an iron complex (e.g., a ferrous chlorophillin or a heme-cofactor such as a heme-containing protein) described herein can be heated in the presence of thiamine and a sugar to affect the formation of 5-Thiazoleethanol, 4-methylfuran, 3,3'-dithiobis[2-methyl-furan, and/or 4-Methylthiazole. These compounds are known to be present in meat and have beefy, meaty taste notes.

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein can be heated in the presence of a nucleotide such as inosine monophosphate and/or guanosine monophosphate to control the formation of flavor compounds such as (E)-4-octene, 2-ethyl-furan, 2-pentanone, 2,3-butanedione, 2-methyl-thiazole, methyl-pyrazine, tridecane, (E)-2-octenal, 2-thiopenecarboxaldehyde, and/or 3-thiopenecarboxaldehyde. These compounds are known to be present in meat and have a beefy, meaty, buttery, and or savory flavor notes.

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein can be heated in the presence of lysine, a sugar such as ribose, and cysteine to control the formation of flavor compounds such as dimethyl trisulfide, nonanal, 2-pentyl thiophene, 2-nonenal furfural, 1-octanol, 2-nonenal, thiazole, 2-acetylthiazole, phenylacetaldehyde, and/or 2-acetylthiazole. These compounds are known to be present in meat and some have a beefy, meaty, and or savory flavor.

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein can be heated in the presence of lactic acid, a sugar such as ribose, and cysteine to control the formation of the flavor compounds nonanal, thiazole, 2-acetylthiazole, and/or 8-methyl 1-undecene. These

compounds are known to be present in meat and have beefy, savory, browned, bready, and malty notes.

In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein can be heated in the presence of 5 amino acids, sugars such as glucose, ribose, and maltodextrin, lactic acid, thiamine, IMP, GMP, creatine, and salts such as potassium chloride and sodium chloride, to control the formation of flavor compounds such as 1,3-bis(1,1-dimethylethyl)-benzene, 2-methyl 3-furanthiol, and/or bis(2-methyl-4,5-dihydro-3-furyl) disulfide. These compounds are known to be present in meat and have beefy notes. See also Table 14.

10 In some embodiments, a particular type of heme-containing protein is chosen to control the formation of flavor compounds. See, for example, the results of Table 9, which shows that the addition of different types of heme-proteins (LegH, Barley, *B. myoglobin*, or *A. aeolicus*) in flavor reaction mixtures containing one or more flavor precursor compounds results in many of the same key meat flavors, including but not 15 limited to pentanone, 3-methyl butanal, 2-methyl butanal, 2-heptenal, 1-octene, nonanal, 2-propenal, 2-decenal, 2-nonenone, 2-octanone, 2-tridecen-1-ol, 2-octanone, 2-octenal, 4-methyl-2-heptanone, octanal, 2-undecenal, butyrolactone, 1-octen-3-one, 3-methylheptyl acetate, and 2-pentyl-thiophene. These differences in flavor compounds can change the overall taste profile.

20 In some embodiments, an iron complex (e.g., a ferrous chlorin or a heme-cofactor such as a heme-containing protein) described herein and one or more flavor precursors can be reacted (e.g., *in vitro*) with heating to generate a particular flavor and/or aroma profile of interest and the resultant flavor additive composition can be added to the consumable food product of interest, which can then be eaten as-is or can be additionally 25 modified, e.g., by additional cooking.

In some embodiments, any undesirable flavors can be minimized by deodorizing with activated charcoal or by removing enzymes such as lipoxygenases (LOX), which can be present in trace amounts when using preparations of plant proteins, and which can convert unsaturated triacylglycerides (such as linoleic acid or linolenic acid) into smaller 30 and more volatile molecules. LOX are naturally present in legumes such as peas,

soybeans, and peanuts, as well as rice, potatoes, and olives. When legume flours are fractionated into separate protein fractions, LOX can act as undesirable “time-bombs” that can cause undesirable flavors on aging or storage. Compositions containing plant proteins (e.g., from ground plant seeds) can be subjected to purification to remove LOX using, for example, an affinity resin that binds to LOX and removes it from the protein sample. The affinity resin can be linoleic acid, linolenic acid, stearic acid, oleic acid, propyl gallate, or epigallocatechin gallate attached to a solid support such as a bead or resin. See, e.g., WO2013138793. In addition, depending on the protein component of the food product, certain combinations of antioxidants and/or LOX inhibitors can be used as effective agents to minimize off-flavor or off-odor generation especially in the presence of fats and oils. Such compounds can include, for example, one or more of  $\beta$ -carotene,  $\alpha$ -tocopherol, caffeic acid, propyl gallate, or epigallocatechin gallate.

In some embodiments, specific flavor compounds, such as those described in Tables 3, 8, 9, 12, 14, 16, or 17 can be isolated and purified from the flavor additive composition. These isolated and purified compounds can be used as an ingredient to create flavors useful to the food and fragrance industry.

A flavor additive composition can be in the form, of but not limited to, soup or stew bases, bouillon, e.g., powder or cubes, flavor packets, or seasoning packets or shakers. Such flavor additive compositions can be used to modulate the flavor and/or aroma profile for a variety of food products, and can be added to a consumable food product before, during, or after cooking of the food product.

### *Food Products*

Food products containing one or more flavor precursors and one or more heme-containing proteins can be used as a base for formulating a variety of additional food products, including meat substitutes, soup bases, stew bases, snack foods, bouillon powders, bouillon cubes, flavor packets, or frozen food products. Meat substitutes can be formulated, for example, as hot dogs, burgers, ground meat, sausages, steaks, filets, roasts, breasts, thighs, wings, meatballs, meatloaf, bacon, strips, fingers, nuggets, cutlets, or cubes.

5 In addition, food products described herein can be used to modulate the taste and/or aroma profile of other food products (e.g., meat replicas, meat substitutes, tofu, mock duck or other gluten based vegetable product, textured vegetable protein such as textured soy protein, pork, fish, lamb, or poultry products such as chicken or turkey products) and can be applied to the other food product before or during cooking. Using the food products described herein can provide a particular meaty taste and smell, for example, the taste and smell of beef or bacon, to a non-meat product or to a poultry product.

10 Food products described herein can be packaged in various ways, including being sealed within individual packets or shakers, such that the composition can be sprinkled or spread on top of a food product before or during cooking.

15 Food products described herein can include additional ingredients including food-grade oils such as canola, corn, sunflower, soybean, olive or coconut oil, seasoning agents such as edible salts (e.g., sodium or potassium chloride) or herbs (e.g., rosemary, thyme, basil, sage, or mint), flavoring agents, proteins (e.g., soy protein isolate, wheat gluten, pea vicilin, and/or pea legumin), protein concentrates (e.g., soy protein concentrate), emulsifiers (e.g., lecithin), gelling agents (e.g., k-carrageenan or gelatin), fibers (e.g., bamboo fiber or inulin), or minerals (e.g., iodine, zinc, and/or calcium).

20 Food products described herein also can include a natural coloring agent such as turmeric or beet juice, or an artificial coloring agent such as azo dyes, triphenylmethanes, xanthenes, quinines, indigoids, titanium dioxide, red #3, red #40, blue #1, or yellow #5.

Food products described herein also can include meat shelf life extenders such as carbon monoxide, nitrites, sodium metabisulfite, Bombal, vitamin E, rosemary extract, green tea extract, catechins and other anti-oxidants.

25 Food products described herein can be free of animal products (e.g., animal heme-containing proteins or other animal products).

In some embodiments, the food products can be soy-free, wheat-free, yeast-free, MSG-free, and/or free of protein hydrolysis products, and can taste meaty, highly savory, and without off odors or flavors.

*Assessment of Food Products*

Food products described herein can be assessed using trained human panelists.

The evaluations can involve eyeing, feeling, chewing, and tasting of the product to judge product appearance, color, integrity, texture, flavor, and mouth feel, etc. Panelists can be served samples under red or under white light. Samples can be assigned random three-digit numbers and rotated in ballot position to prevent bias. Sensory judgments can be scaled for “acceptance” or “likeability” or use special terminology. For example, letter scales (A for excellent, B for good, C for poor) or number scales may be used (1 = dislike, 2 = fair, 3 = good; 4 = very good; 5=excellent). A scale can be used to rate the overall acceptability or quality of the food product or specific quality attributes such beefiness, texture, and flavor. Panelists can be encouraged to rinse their mouths with water between samples, and given opportunity to comment on each sample.

In some embodiments, a food product described herein can be compared to another food product (e.g., meat or meat substitute) based upon olfactometer readings. In various embodiments, the olfactometer can be used to assess odor concentration and odor thresholds, odor suprathresholds with comparison to a reference gas, hedonic scale scores to determine the degree of appreciation, or relative intensity of odors.

In some embodiments, an olfactometer allows the training and automatic evaluation of expert panels. In some embodiments, a food product described herein causes similar or identical olfactometer readings. In some embodiments, the differences between flavors generated using the methods of the invention and meat are sufficiently small to be below the detection threshold of human perception.

In some embodiments, volatile chemicals identified using GCMS can be evaluated. For example, a human can rate the experience of smelling the chemical responsible for a certain peak. This information could be used to further refine the profile of flavor and aroma compounds produced using a heme-containing protein and one or more flavor precursors.

Characteristic flavor and fragrance components are mostly produced during the cooking process by chemical reactions molecules including amino acids, fats and sugars which are found in plants as well as meat. Therefore, in some embodiments, a food

product is tested for similarity to meat during or after cooking. In some embodiments human ratings, human evaluation, olfactometer readings, or GCMS measurements, or combinations thereof, are used to create an olfactory map of the food product. Similarly, an olfactory map of the food product, for example, a meat replica, can be created. These 5 maps can be compared to assess how similar the cooked food product is to meat.

In some embodiments, the olfactory map of the food product during or after cooking is similar to or indistinguishable from that of cooked or cooking meat. In some 10 embodiments the similarity is sufficient to be beyond the detection threshold of human perception. The food product can be created so its characteristics are similar to a food product after cooking, but the uncooked food product may have properties that are different from the predicate food product prior to cooking.

These results will demonstrate that the compositions of the invention are judged 15 as acceptably equivalent to real meat products. Additionally, these results can demonstrate that compositions of the invention are preferred by panelist over other commercially available meat substitutes. So, in some embodiments the present invention provides for consumables that are significantly similar to traditional meats and are more meat like than previously known meat alternatives.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

## EXAMPLES

### Example 1: Addition of heme-protein increases beefy qualities of replica burgers

Replica burgers containing the ingredients in Table 4 and the flavor precursors  
 5 cysteine (10 mM), glutamic acid (10 mM), glucose (10 mM), and thiamine (1 mM) were prepared. Water was added to make up the balance. See, for example, U.S. Provisional Application No. 61/751,816, filed January 11, 2013. Control burgers were prepared as in Table 4 with precursors cysteine (10 mM), glutamic acid (10 mM), glucose (10 mM), and thiamine (1 mM) except LegH was omitted.

10 After cooking for 5 minutes at 150 C, the replica burgers were evaluated by a trained sensory panel. Panelists were served samples under red lights and each panelist individually evaluated the samples. Samples were assigned a random three-digit number and rotated in ballot position to prevent bias. Panelists were asked to evaluate cooked replica burger samples on multiple flavor, aroma, taste, texture and appearance attributes  
 15 including but not limited to: beefiness, bloody quality, savory quality, and overall acceptability using a 7-point scale from 1=dislike extremely, to 7=like extremely. Panelists were encouraged to rinse their mouths with water between samples, and to fill out a survey to record their evaluation of each sample.

When replica burgers containing the LegH were compared to the control replica  
 20 burgers without LegH, the samples containing LegH were rated significantly beefier, bloodier, more savory, and overall preferred compared to those that did not include LegH. See Table 5.

TABLE 4  
 25 Replica Burger Ingredients

Replica burger	% precooked w/w
Pea vicilin	3.86
Soy protein concentrate (SPC)	2.52
Bamboo fiber	0.34

NaCl	0.54
Pea legumin	2
Soy Protein Isolate (SPI) (Solae, St. Louis, MO)	4.68
Wheat gluten	4.68
Coconut oil	15
Soy lecithin	0.1
k-carrageenan	1
LegH	1

**TABLE 5**  
**Sensory evaluation of replica burger with Heme**

Attribute	Beef 20/80	No Heme	1% Heme
Beefyness			
mean	5.33	1.30	3.20
STDEV	1.58	0.67	0.79
Bloody			
mean	4.00	1.10	2.78
STDEV	1.32	0.32	1.64
Savory			
mean	4.67	3.00	5.10
STDEV	1.22	1.63	0.57

5                   **Example 2: Replica burgers with a flavor precursor mixture taste beefy and  
bloody**

Replica burgers containing a flavor precursor mixture of glucose, cysteine, thiamine, and glutamic acid and 1% LegH pre-cooked w/w (see Table 4) were prepared as described in Example 1, and evaluated by a trained sensory panel after the burgers 10 were cooked for 5 minutes at 150 C. Control burgers included LegH and all other ingredients except for the flavor precursor mixture.

Panelists were asked to evaluate the samples overall improvement in taste and descriptively analyze each sample using a 5-point scale from 1=dislike extremely, to 5=like extremely. Panelists were encouraged to rinse their mouths with water between samples, and to fill out a survey to record their evaluation of each sample. The replicate burgers which included LegH and the flavor precursor mixture were described as having bouillon, gravy, meaty, bloody, savory, and beefy notes on taste, and were preferred to the same replica burger with LegH but no added flavor precursor mixture. See, Table 6

**TABLE 6**

**Improvement of overall taste with precursors added to LegH burgers**

	with precursors	without precursors
Average	3.5	1.8
STDV	0.6	0.5

10

**Example 3: Replica burgers with flavor precursor mixture resulting in a bacon taste**

Replica burgers (see Table 4) were cooked with different precursor mixes (see Table 7) and 1% LegH and evaluated by a trained sensory panel after the burgers were cooked for 5 minutes at 150 C. Control burgers contained LegH and all of the other ingredients except for the flavor precursors. Panelists were asked to evaluate each sample and descriptively analyze of each sample. 5-point scale from 1=dislike extremely, to 5=like extremely. Panelists were encouraged to rinse their mouths with water between samples, and to fill out a survey to record their evaluation of each sample. A replica burger with a precursor mixture of 10 mM glucose, 10 mM ribose, 10 mM cysteine, 1 mM thiamine, 1 mM glutamic acid, 1 mM GMP, and LegH was described as having a bacon aroma and taste, and overall meatiness, savory quality, a very umami quality, a brothy quality, and slight beefy notes. See Table 7 for a summary of the flavor description for the various combinations of flavor precursors and heme-containing protein.

TABLE 7  
Flavors generated by addition of precursors to LegH (1 %)

Precursor (concentration)			Flavor Description
ribose (10 mM)	cysteine (10 mM)		some kind of cold-cut/sliced deli meat
ribose (10 mM)	cysteine (10 mM)		bread crust with beef drippings, sweet, grassy, umami
ribose (10 mM)	cysteine (10 mM)		lactic acid (1 mM)
ribose (10 mM)	cysteine (10 mM)		bread-y, malty, browned, breadcrust
ribose (10 mM)	cysteine (10 mM)		lysine (5 mM)
ribose (10 mM)	cysteine (10 mM)		alanine (5 mM)
ribose (10 mM)	cysteine (10 mM)		savory, weak beefy, brothy, little metallic
ribose (10 mM)	cysteine (10 mM)		I+G (2 mM)
ribose (10 mM)	cysteine (10 mM)		savory, weak beefy, brothy, sweet
ribose (10 mM)	cysteine (10 mM)		methionine
ribose (10 mM)	cysteine (10 mM)		cooked potato
ribose (10 mM)	cysteine (10 mM)		little meaty, pretzel, brothy, savory, sweet, chocolate
glucose (10 mM)	cysteine (10 mM)		glutamic acid (5 mM)
glucose (10 mM)	cysteine (10 mM)		glutamic acid (5 mM)
glucose (10 mM)	cysteine (10 mM)		slight beefy, browned, grassy,
glucose (10 mM)	cysteine (10 mM)		IMP (2 mM)
glucose (10 mM)	cysteine (10 mM)		bacon, very umami, savory, brothy, slight beef
			beef jerky, bloody, meaty, brothy
			savory, beefy, bloody, meaty, savory, gravy

glucose (10 mM)	cysteine (10 mM)	thiamine (2 mM)	glutamic acid (5 mM)	lysine (5 mM)	roast beef
glucose (10 mM)	cysteine (10 mM)	thiamine (2 mM)	glutamic acid (5 mM)	alanine (5 mM)	boiled beef, sweet
glucose (10 mM)	cysteine (10 mM)	thiamine (2 mM)	glutamic acid (5 mM)	I+G (2 mM)	beefy with a sulfury note
glucose (10 mM)	cysteine (10 mM)			I+G (2 mM)	sweet, malty, umami, meaty
glucose (10 mM)				I+G (2 mM)	savory, roast beef, grassy
glucose (10 mM)			glutamic acid (5 mM)		umami, savory, meaty, sweaty, fermented

**Example 4: Type of sugar modulates flavor compounds created in the presence of Hemeprotein**

The addition of different sugars to flavor reaction mixtures containing a hemeprotein and one or more flavor precursor compounds resulted in distinct differences 5 in the flavor compounds generated and the overall flavor profile. LegH heme protein at 1% pre-cooked w/w was mixed with cysteine (10 mM) and glucose (20 mM) at pH 6 in phosphate buffer to form a flavor reaction mixture and heated to 150 C for 3 minutes; this reaction created flavor compounds known to be present in meat; *see* Table 8. Similarly, a flavor reaction mixture made when LegH heme protein at 1% was mixed with cysteine 10 (10 mM) and ribose (20 mM) at pH 6 and heated to 150 C for 3 minutes created flavor compounds known to be in meat; *see* Table 8.

The characteristic flavor and fragrance components were mostly produced during the cooking process when the flavor precursor molecules reacted with the heme-protein. Gas chromatography–mass spectrometry (GCMS) is a method that combines the features 15 of gas-liquid chromatography and mass spectrometry to separate and identify different substances within a test sample. Samples were evaluated by GCMS to identify the flavor compounds generated after heating and also evaluated for their sensory profiles. Volatile chemicals were isolated from the head space around the flavor reactions. The profile of the volatile chemicals in the headspace around the flavor reaction mixtures is shown in 20 Table 8. In particular, the use of ribose created some additional compounds as compared to glucose, as shown in Table 8.

Notably, the control mixtures of cysteine with ribose or glucose heated in the absence of the LegH heme-protein did not generate the same set of flavor compounds. The flavor reaction mixtures containing LegH also were evaluated by a blinded trained 25 sensory panel, which described the samples with ribose as having beefy, savory, brothy, and gravy-like notes, and the samples with glucose as savory, bloody, metallic, raw meat, and bouillon-like.

**TABLE 8**  
**Flavor compounds generated with cysteine, LegH, and either glucose or ribose in**  
**the flavor reaction mixture.**

LegH 1%		
Compounds created	cysteine (10 mM), glucose (20 mM)	cysteine (10 mM), ribose (20 mM)
benzaldehyde	X	X
2-butanone	X	X
dimethyl trisulfide	X	X
2-pentyl-furan	X	X
2-methyl-propanal	X	X
thiazole	X	X
butyrolactone	X	X
2-acetylthiazole	X	X
pentanal	X	X
3-methyl-butanal	X	X
methyl-thiirane	X	X
nonanal	X	X
heptanal	X	X
2,3-butanedione	X	X
1,3,5-cycloheptatriene	X	X
propyl-cyclopropane	X	X
2-hexyl-furan	X	X
butanal	X	X
2-methyl-butanal	X	X
2-ethyl-furan		X
2-octanone	X	X
propanal	X	X

trichloromethane	X	
2-methyl-furan	X	X
furan	X	X
pyrazine	X	X
thiophene	X	X
1,3-dimethyl-benzene	X	X
octane		X
octanal	X	X
thiazole	X	X
2-pentanone		X
furfural	X	X
2-nonenone	X	X
(Z)-2-heptenal	X	X
(E)-2-heptenal	X	X
1-octene	X	X
formic acid, heptyl ester	X	X
2-pentyl-thiophene		X
1-octen-3-one	X	X
3-pentyl-furan	X	X
2-propenal		X
(E)-2-tridecen-1-ol		X
benzene		X
(E)-4-octene		X
1-penten-3-one		X
4-penten-2-one	X	X
2-methyl-thiazole		X
methyl-pyrazine		X
trans-2-(2-pentenyl)furan		X
3-ethylcyclopentanone		X

pyrrole	X	X
2-thiophenecarboxaldehyde		X
3-thiophenecarboxaldehyde		X

**Example 5: Heme-protein in the presence of thiamine affects the production of certain flavor compounds**

The addition of thiamine in a flavor reaction mixtures with a heme protein and other flavor precursors affected the formation of 5-Thiazoleethanol, 4-methyl-furan, 3,3'-dithiobis[2-methyl-thiazole, and 4-methylthiazole. These compounds are known to be present in meat and have beefy, meaty taste notes.

Flavor reaction mixtures at pH 6 containing LegH (1%), cysteine (10 mM), thiamine (1 mM), either glucose or ribose (20 mM), and with or without glutamic acid (10 mM) were prepared and subsequently heated to 150 C for 3 minutes. These flavor reaction samples then were evaluated by GCMS for the flavor compounds generated and evaluated by a trained panel for their sensory profiles. Volatile chemicals were isolated from the head space around the flavor reactions. GCMS showed 4-methyl-5-thiazoleethanol, 3,3'-dithiobis[2-methyl]-furan, and 4-methylthiazole compounds were created by a mixture of LegH with thiamine, a sugar (either glucose or ribose), and cysteine. The same flavor reaction mixtures without thiamine did not generate these compounds; additionally these compounds were not generated when heme-proteins were not present in the flavor reaction mixtures.

The flavor reaction samples also were evaluated by a blinded trained sensory panel, which described the samples with the addition of thiamine as more complex in taste and more beefy, meaty, and savory.

**Example 6: Heme-proteins with nucleotides controls particular flavor compound production.**

The addition of inosine monophosphate and guanosine monophosphate in mixes with heme protein and other precursors controlled the formation of flavor compounds (E)-4-octene, 2-ethyl-furan, 2-pentanone, 2,3-butanedione, 2-methyl-thiazole, , methyl-

pyrazine, tridecane, (E)-2-octenal, 2-thiophenecarboxaldehyde, and 3-thiophenecarboxaldehyde. These compounds are known to be present in meat and have a beefy, meaty, buttery, and or savory flavor notes.

Reactions containing heme protein at 1% (LegH) with cysteine (10 mM), and glucose (20 mM), 1 mM IMP and 1 mM GMP, at pH 6.0 were prepared and heated to 150C for 3 minutes. Characteristic flavor and fragrance components were mostly produced during the cooking process where precursors reacting heme-protein. These samples were evaluated by GCMS for the flavor compounds generated and evaluated for the sensory experience. Volatile chemicals were isolated from the head space around the flavor reaction and identified using GCMS, creating a profile of the volatile chemicals in the headspace around the flavor reaction mixture. GCMS showed 4-octene, 2-ethyl furan, 2-pentanone, 2,3-butanedione, 2-methyl-thiazole, methyl-pyrazine, tridecane, 2-octenal, 2-thiophenecarboxaldehyde, 3-thiophenecarboxaldehyde compounds were created by a mixture of hemeprotein LegH with IMP, GMP, glucose, and cysteine. The same samples without IMP and GMP did not generate these compounds, additionally these compounds were also not created when heme-proteins were not present, just precursor molecules. Sensory evaluation by blinded trained panelist found the samples with the addition of inosine and guanosine as described as having more complexity in taste and more beefy, meaty, brothy and savory. Figure 2 shows the abundance of the novel flavor compounds created with heme protein at 1% was mixed in a reaction at pH 6, with cysteine (10 mM), and glucose (20 mM), IMP (1 mM) and GMP (1 mM), and detected by solid phase microextraction (SPME) and then detected by GCMS.

#### **Example 7: Flavor generation with the addition of a particular organic acid**

The addition of lactic acid in mixes with heme protein, ribose, and cysteine controlled the formation of the flavor compounds nonanal, thiazole, 2-acetylthiazole, and 8-methyl-1-undecene. These compounds are known to be present in meat.

Reactions containing heme protein at 1%, cysteine (10 mM), and ribose (20 mM), and lactic acid (1mM), pH 6.0, were prepared and heated to 150C for 3 minutes. Characteristic flavor and fragrance components were mostly produced during the cooking

process where precursors reacting heme-protein. These samples were evaluated by GCMS for the flavor compounds generated and evaluated for the sensory experience. Volatile chemicals were isolated from the head space around the flavor reaction and identified using GCMS, creating a profile of the generated compounds. Nonanal, 5 thiazole, 2-acetylthiazole, and 8-methyl-1-undecene compounds were created by a mixture of LegH with lactic acid, ribose, and cysteine. The same samples without lactic acid did not generate these compounds, additionally these compounds were not created in the absence of heme-proteins.

10 Sensory evaluation by blinded trained panelist found the samples with the addition of lactic acid as described as beefy, savory, browned, bready, and having malty notes. The sample with everything but lactic acid rated lower in browned, bready and malty notes.

**Example 8: Flavor generated with the addition of a particular amino acid.**

15 The addition of lysine in mixes with heme protein ribose, and cysteine controlled the formation of flavor compounds dimethyl trisulfide, nonanal, 2-pentyl-thiophene, furfural, 2-nonenal, 1-octanol, 2-nonenal, thiazole, 2-acetylthiazole, phenylacetaldehyde, 2-acetylthiazole. These compounds are known to be present in meat and some have a beefy, meaty, and or savory flavor.

20 Reactions containing heme protein at 1%, cysteine (10 mM), and ribose (20 mM), and lysine (1mM), at pH 6.0, were prepared and heated to 150C for 3 minutes. These samples were evaluated by GCMS for the flavor compounds generated and evaluated for the sensory experience. Characteristic flavor and fragrance components were mostly produced during the cooking process where precursors could react with the heme-protein. 25 These samples were evaluated by GCMS for the flavor compounds generated and evaluated for the sensory experience. Volatile chemicals were isolated from the head space around the flavor reaction. Dimethyl trisulfide, nonanal, 2-pentyl-thiophene, furfural, 2-nonenal, 1-octanol, 2-nonenal, thiazole, 2-acetylthiazole, phenylacetaldehyde, 2-acetylthiazole compounds were created by a mixture of LegH with lactic acid, ribose, 30 and cysteine. The same samples without lactic acid did not generate these compounds,

additionally these compounds were not created when heme-proteins were not present, just precursor molecules. Sensory evaluation by blinded trained panelist found the samples with the addition of lysine as described as roast beefy, savory, and browned. The addition of lysine increased the roasted browned notes.

5

**Example 9 - Flavor compound production by different Heme-proteins**

The addition of different types of heme-proteins (LegH, Barley, *B. myoglobin*, or *A. aeolicus*) in flavor reaction mixtures containing one or more flavor precursor compounds results in many of the same key meat flavors, including but not limited to 2-pentyl-furan, 2,3-Butanedione, Thiophene, 2-methyl-thiazole, Pyrazine, Furan, Pyrrole, 2-methyl-furan and distinct differences in the flavor compounds, including but not limited to 2-pentyl-thiophene, Nonanal, 2-Nonanone, and 1-Octen-3-one. These differences in flavor compounds can change the overall taste profile. The different types of heme-protein were LegH, Barley, *B. myoglobin*, or *A. aeolicus* used at 1% w/w in a reaction mixed with cysteine (10 mM) and ribose (10 mM) at pH 6. The pre- reaction mixture was heated to 150 C for 3 minutes; this reaction created flavor compounds known to be present in meat; see Table 9. The characteristic flavor and fragrance components are mostly produced during the cooking process where the flavor precursor molecules react with the heme-protein. Samples were evaluated by GCMS to identify the flavor compounds generated after heating and also evaluated for their sensory profiles. Volatile chemicals were isolated from the head space around the flavor reactions. Table 9 shows the similarity and differences in volatile flavor compounds created by the different types of heme-proteins.

10

15

20

25

**TABLE 9**  
**Flavor compounds created by different heme-protein**  
**when heated with ribose and cysteine.**

Name	LegH	Barley	<i>B. myoglobin</i>	<i>A. aeolicus</i>
Furan	x	x	x	x
Thiazole	x	x	x	x
benzaldehyde	x	x	x	x
2-acetylthiazole	x	x	x	x

2-methyl-propanal	x	x	x	x
furfural	x	x	x	x
2,3-butanedione	x	x	x	x
2-pentyl-furan	x	x	x	x
2-pentanone	x	x		
pyrazine	x	x	x	x
dimethyl trisulfide	x	x	x	x
3-methyl-butanal	x	x		x
2-methyl-thiazole	x	x	x	x
pentanal	x	x	x	x
1,3,5-cycloheptatriene	x	x	x	x
methacrolein	x	x	x	x
heptanal	x	x	x	x
2-methyl-butanal	x	x		x
isothiazole	x	x	x	x
thiophene	x	x	x	x
propanal	x	x	x	x
2-heptenal	x		x	x
methyl-pyrazine	x	x	x	x
1-octene	x		x	x
butanal	x	x	x	x
2-acetyl-propen-2-ol	x	x	x	x
pyrrole	x	x	x	x
2-methyl-furan	x	x	x	x
nonanal		x	x	x
2-propenal		x	x	x
2-decenal		x	x	x
2-nonenone		x		x
2-octanone		x	x	x
2-tridecen-1-ol,			x	x
2-octanone			x	
2-octenal			x	x
4-methyl-2-heptanone			x	x
octanal			x	x
2-undecenal				x
butyrolactone				x
1-octen-3-one				x
3-methylheptyl acetate				x

2-pentyl-thiophene				x
--------------------	--	--	--	---

### Example 10 - Generation of Meat flavors from different lipids

Several different samples including oils (canola oil or coconut oil), free fatty acids (FFA) (linoleic acid (C18:2), oleic acid (C18:1), stearic acid (C18:0), or myristic acid (C14:0)) and phospholipids (PL) (beef heart polar lipids extract, Biolipon95 (from Perimond), or NatCholinePC40 (from Perimond)) were tested for their ability to produce beefy flavor in the absence and in the presents of other precursors. Oils, FFAs, and PLs were added to 50 mM potassium phosphate buffer (PPB) pH 6.0 or a Maillard reaction mix (MRM) containing 50 mM potassium phosphate pH 6.0, 5 mM Cysteine, 10 mM Glucose, 0.1 mM Thiamine, and 0.1% (w/v) LegHemoglobin. Lipids in combination with MRM were designed to capture the cross reactions of lipid degradation and Maillard reaction productions while lipids in phosphate buffer functioned as a lipid control. The oils were added at 3% of the total 1 mL volume of solution while FFAs and PLs were added at 1% of the total 1 mL volumes. All samples were cooked at 150°C for 3 mins, cooled to 50°C and then analyzed using GCMS (SPME fiber sampling of headspace). After all samples were analyzed by GCMS the caps were removed and samples were smelled by a trained flavor scientist and aromas recorded.

**Table 10**  
**Legend showing components of each sample**

Sample Name	Solution	Additives
MRM None	Maillard Reaction Mix	None
MRM_Linoelic Acid	Maillard Reaction Mix	1% linoleic acid
MRM_Oleic Acid	Maillard Reaction Mix	1% oleic acid
MRM_C14	Maillard Reaction Mix	1% C14:0 free fatty acid
MRM_C18	Maillard Reaction Mix	1% C18:0 free fatty acid
MRM_Canola	Maillard Reaction Mix	3% Canola Oil
MRM_Coconut	Maillard Reaction Mix	3% Coconut Oil
MRM_BeefHeart	Maillard Reaction Mix	1% Beef Heart Polar Lipids Extract
MRM_Biolipon95	Maillard Reaction Mix	1% Biolipon95 (emulsifier)

MRM_NatCholinePC40	Maillard Reaction Mix	1% NatCholinePC40 (emulsifier)
KPhos6_Linoelic Acid	PPB, pH 6	1% linoelic acid
KPhos6_Oleic Acid	PPB, pH 6	1% oleic acid
KPhos6_C14	PPB, pH 6	1% C14:0 free fatty acid
KPhos6_C18	PPB, pH 6	1% C18:0 free fatty acid
KPhos6_Canola	PPB pH 6	3% Canola Oil
KPhos6_Coconut	PPB, pH 6	3% Coconut Oil
KPhos6_BeefHeart	PPB, pH 6	1% Beef Heart Polar Lipids Extract
KPhos6_Biolipon95	PPB, pH 6	1% Biolipon95 (emulsifier)
KPhos6_NatCholinePC40	PPB, pH 6	1% NatCholinePC40 (emulsifier)

Table 11 contains the aroma descriptions and Table 12 contains the GCMS data from the most interesting samples analyzed. Many of the lipids introduced a “fatty” aroma to MRM that was otherwise absent. The combinations of Linoleic Acid or 5 NatCholinePC40 in MRM produced the greatest abundance of fatty compounds suggesting that these lipids may improve the flavor perception of beef tallow. Linoleic Acid and NatCholinePC40 also showed high abundance of earthy-mushroom aromas. The addition of lipids to MRM significantly increased the abundance of “nutty & roasted” aromas. Less desirable “green” aroma compounds were most prominent in 10 samples with unsaturated free fatty acids (linoleic acid or oleic acid) or phospholipids. In general, the addition of lipids significantly increased the number of target beef compounds made.

15 **TABLE 11**  
**Aroma descriptions of each sample after it was cooked.**

Sample Names	Aroma Descriptions
MRM_Only	brothy, malty, beef stew
KPhos6_BeefHeart	fatty, creamy, beef tallow, slight sweet, slight roasted nutty
MRM_BeefHeart	fatty, beef tallow, old meat, mushroom
KPhos6_Biolipon95	fatty, fresh
MRM_Biolipon95	fatty, brothy, hay, malty green

KPhos6_NatCholinePC40	light fatty, fresh
MRM_NatCholinePC40	fatty, beef tallow, brothy
K-Phos6_C14	light/faint plastic/waxy
MRM_C14	brothy, beefy, minty, fresh
K-Phos6_C18	light/faint plastic/waxy
MRM_C18	beefy with cucumber &/or pepper aroma
K-Phos6_Canola	fresh, cucumber
MRM_Canola	fatty, brothy, oil, roasted nuts
K-Phos6_Coconut	nothing
MRM_Coconut	brothy, beefy, slight fatty, crackers
K-Phos6_Oleic Acid	fresh, cucumber, camphorous/minty-like
MRM_Oleic Acid	herbal, plastic, slight cheesy, brothy
K-Phos6_Linoelic Acid	light plastic
MRM_Linoelic Acid	fatty, light waxy, brothy, herbal

**Table 12**

**List of aromatic compounds found in Beef by GCMS and a chart showing which were detected in each lipid plus MRM sample.**

Compounds in Beef	MRM only	MRM_BeefHeart	MRM_NatCholinePC40	MRM_Linoelic acid
(s)-isopropyl lactate	N	N	N	N
1-ethyl-5-methylcyclopentene	Y	Y	Y	Y
1-heptanol	N	Y	N	N
1-hepten-3-ol	N	Y	Y	Y
1-heptene	N	Y	Y	Y
2-methyl-1-heptene	N	N	N	N
1-hexanol	N	Y	Y	Y
2-ethyl-1-hexanol	N	N	N	N
1-nonanol	N	N	Y	N
1-nonene	N	Y	Y	N
1-octanol	N	Y	Y	N
1-octen-3-ol	N	Y	Y	Y

1-octen-3-one	Y	Y	Y	Y
1-octene	N	N	N	N
1-pentanol	N	Y	Y	Y
1-penten-3-ol	N	Y	Y	N
1-propanol	N	N	N	N
8-methyl-1-undecene	N	Y	Y	Y
1,3-hexadiene	N	N	N	Y
3-ethyl-2-methyl-1,3-hexadiene	N	Y	Y	Y
1,3-octadiene	Y	N	N	Y
1,3,5-cycloheptatriene	N	N	N	N
2,3-dihydro-5,6-dimethyl-1,4-dioxin	N	N	N	N
1,7-octadien-3-ol	N	Y	N	N
1h-pyrrole-2-carboxaldehyde	N	N	N	N
2-methyl-1H-pyrrole	N	N	N	N
2-acetyl-2-thiazoline	Y	N	N	N
2-acetylthiazole	Y	Y	Y	Y
2-butanone	N	Y	Y	Y
2-butenal	N	Y	Y	Y
2-ethyl-2-butenal	N	N	N	Y
3-methyl-2-butenal	N	N	Y	Y
3-methyl-2-cyclohexen-1-one	N	N	N	N
2-decanone	Y	Y	Y	N
(E)-2-decenal	N	N	N	N
(Z)-2-decenal	Y	Y	Y	Y
2-furanmethanol	N	N	N	N
2-heptanone	Y	Y	Y	Y
6-methyl-2-heptanone	N	N	Y	N
(E)-2-heptenal	N	Y	Y	Y
(Z)-2-heptenal	N	N	N	Y
(E)-2-hexenal	N	Y	Y	Y
2-ethyl-2-hexenal	N	N	N	N
2-methyl-2-heptene	Y	N	N	N
2-n-heptylfuran	Y	N	N	N
2-n-octylfuran	Y	Y	Y	N

2-nonanone	N	Y	Y	N
(E)-2-nonenal	Y	Y	Y	Y
(Z)-2-nonenal	N	N	N	Y
2-octanone	Y	Y	Y	Y
(Z)-2-octen-1-ol	Y	Y	Y	Y
(E)-2-octenal	N	Y	Y	Y
2-pentanone	N	Y	Y	N
1-propoxy-2-propanol	N	N	N	N
1-(acetyloxy)-2-propanone	Y	N	N	N
1-hydroxy-2-propanone	Y	N	N	N
2-propenal	N	N	N	Y
2-thiophenecarboxaldehyde	Y	Y	Y	Y
2-undecenal	N	Y	Y	Y
2,3-butanedione	N	N	N	Y
2,3-pentanedione	N	N	N	N
(E,E)-2,4-decadienal	N	Y	Y	Y
2,4-decadienal	N	N	N	Y
(E,E)-2,4-heptadienal	N	Y	Y	Y
(E,E)-2,4-nonadienal	N	Y	Y	Y
2,6-dimethylpyrazine	N	N	N	N
(E,Z)-2,6-nonadienal	N	N	Y	N
5-ethylhydro-2(3H)-furanone	N	Y	Y	Y
5-methyl-2(3H)-furanone	N	N	N	N
dihydro-5-pentyl-2(3H)-furanone	N	N	Y	Y
dihydro-5-propyl-2(3H)-furanone	N	N	N	N
2(5H)-furanone	N	N	N	N
tetrahydro-6-methyl-2H-pyran-2-one	N	N	N	N
3-ethylcyclopentanone	N	Y	Y	Y
3-hexanone	N	N	N	N
3-methyl-2-thiophenecarboxaldehyde	N	N	N	N
3-octanone	Y	Y	N	Y
3-octen-2-one	N	Y	Y	Y
3-thiophenecarboxaldehyde	N	Y	Y	Y

	N	N	Y	Y
(E,E)-3,5-octadien-2-one				
dihydro-2-methyl-3(2H)-furanone	N	N	N	N
4-cyanocyclohexene	N	N	N	N
4-cyclopentene-1,3-dione	N	N	Y	N
4-decyne	N	Y	N	N
(Z)-4-heptenal	N	Y	Y	Y
4-methyloctanoic acid	N	N	N	N
(E)-4-octene	N	N	N	N
2,3-dihydro-3,5-dihydroxy-6-methyl-4(H)-pyran-4-one	Y	N	N	N
6-methyl-5-hepten-2-one	Y	N	N	N
acetaldehyde	N	N	N	Y
acetic acid	N	N	N	N
acetic acid ethenyl ester	Y	N	N	N
acetoin	Y	N	N	N
acetone	Y	N	N	Y
acetonitrile	N	N	N	Y
benzaldehyde	Y	Y	Y	Y
4-ethyl-benzaldehyde	N	Y	Y	N
benzene	Y	N	N	N
benzoic acid, hydrazide	Y	N	N	N
butanal	Y	N	N	Y
2-methyl-butanal	N	N	N	N
3-methyl-butanal	Y	N	N	N
butanoic acid	N	N	N	N
butyrolactone	Y	Y	N	Y
caprolactam	N	N	N	N
carbon disulfide	N	N	N	Y
1-ethyl-1-methyl-cyclopentane	Y	Y	Y	Y
propyl-cyclopropane	N	N	Y	Y
decanal	N	Y	Y	N
dihydro-3-(2H)-thiophenone	N	N	N	N
Dimethyl sulfide	Y	N	N	N
dimethyl sulfone	N	N	N	N
dimethyl trisulfide	Y	Y	N	N
ethanethiol	N	N	N	N

	N	N	N	Y
ethanol				
1-(1(H)-pyrrol-2-yl)-ethanone	N	N	N	N
1-(2-furanyl)-ethanone	N	N	N	N
ethosuximide	Y	N	N	N
formic acid, heptyl ester	Y	Y	N	N
furan	Y	N	N	Y
2-ethyl-furan	Y	N	N	N
2-hexyl-furan	Y	N	N	Y
2-methyl-furan	N	N	N	Y
2-pentyl-furan	N	Y	Y	Y
2-propyl-furan	N	N	Y	Y
3-methyl-furan	Y	N	N	N
3-pentyl-furan	Y	Y	Y	Y
furfural	N	Y	Y	Y
heptanal	N	Y	Y	Y
heptanoic acid	N	N	N	Y
2-methyl-hex-2-yn-4-one	N	N	N	N
hexanoic acid	N	N	N	Y
hydrogen sulfide	N	N	N	N
m-aminophenylacetylene	N	N	N	N
maleic anhydride	N	N	N	N
methacrolein	N	N	N	N
methanethiol	N	N	N	N
methyl ethanoate	N	N	N	N
methyl isobutyl ketone	Y	N	N	N
n-caproic acid vinyl ester	N	Y	Y	N
nonanal	N	Y	Y	Y
3-methyl-nonane	Y	N	N	N
nonanoic acid	Y	N	N	N
octanal	N	Y	Y	Y
octane	N	N	N	Y
octanoic acid	N	N	N	Y
oxalic acid, isobutyl pentyl ester	Y	N	N	N
p-cresol	N	N	N	N
pentanal	N	N	N	Y
pentanoic acid	Y	N	N	Y

4-ethyl-phenol	N	Y	Y	N
phenylacetaldehyde	Y	Y	Y	Y
(p-hydroxyphenyl)-phosphonic acid	Y	N	N	N
propanal	N	N	N	Y
2-methyl-propanal	N	N	N	N
propanoic acid	N	N	N	N
2-methyl-propanoic acid	Y	N	N	N
propanoic acid, ethenyl ester	N	N	N	N
pyrazine	N	Y	N	Y
2-ethyl-5-methyl-pyrazine	N	N	N	N
2-ethyl-6-methyl-pyrazine	N	N	N	N
2,3-dimethyl-pyrazine	N	N	N	N
2,5-dimethyl-pyrazine	N	N	N	N
3-ethyl-2,5-dimethyl-pyrazine	Y	N	N	N
ethyl-pyrazine	N	N	N	N
methyl-pyrazine	N	N	N	N
trimethyl-pyrazine	Y	N	N	N
pyridine	Y	N	Y	N
pyrrole	Y	Y	Y	Y
styrene	Y	N	Y	N
thiazole	Y	Y	Y	Y
methyl-thiirane	N	N	N	N
thiophene	N	N	N	Y
2-hexyl-thiophene	Y	N	Y	N
2-pentyl-thiophene	N	Y	N	N
trans-2-(2-pentenyl)furan	N	Y	Y	N
trans-3-nonen-2-one	N	Y	Y	Y
undecanoic acid	N	N	N	N
Total # of Compounds				
Detected:	54	63	66	76

In samples having fatty or creamy aromas, 2,4-decadienal, (E,E)-2,4-nonadienal, (E,E)-2,4-heptadienal, and/or (E,E)-2,4-decadienal were detected in the KPhos6\_BeefHeart, MRM\_BeefHeart, MRM\_BioLipon95, MRM\_NatCholinePC40,

Kphos6\_Canola, MRM\_Canola, KPhos6\_Oleic Acid, KPhos6\_Linoleic acid and MRM\_Linoleic acid samples. For (E,E)-2,4-decadienal, the strongest signal intensity was in the MRM\_NatCholinePC40 sample, followed by the MRM\_Linoleic acid, KPhos6\_Linoleic acid, MRM\_BeefHeart, MRM\_BioLipon95, KPhos6\_BeefHeart, 5 MRM\_Oleic Acid, and KPhos6\_Oleic Acid samples. For (E,E)-2,4-heptadienal, the strongest signal intensity was in the MRM\_NatCholinePC40 sample followed by the MRM\_Canola sample. (E,E)-2,4-heptadienal also was detected in the MRM\_BioLipon95, MRM\_BeefHeart, and MRM\_Linoleic acid samples. For (E,E)-2,4- 10 nonadienal, the strongest signal intensity was in the MRM\_Canola and MRM\_Linoleic acid samples. (E,E)-2,4-nonadienal also was detected in the Kphos6\_Canola, MRM\_NatCholinePC40, MRM\_BioLipon95, MRM\_BeefHeart, and KPhos6\_Linoleic acid samples. For 2,4-decadienal, the strongest signal intensity was in the MRM\_Linoleic acid sample. 2,4-decadienal also was detected in KPhos6\_Linoleic acid, MRM\_Canola, and KPhos6\_Oleic Acid samples.

15 In samples having earthy or mushroom aromas, 3-octen-2-one, 1-octen-3-one, 3-octanone, and/or 1-octen-3-ol were detected in the KPhos6\_BeefHeart, MRM\_BeefHeart, Kphos\_BioLipon95, MRM\_BioLipon95, Kphos\_NatCholinePC40, MRM\_NatCholinePC40, MRM\_Canola, KPhos6\_Oleic Acid, MRM\_Oleic Acid, KPhos6\_Linoleic acid, and MRM\_Linoleic acid samples. For 1-octen-3-ol, the strongest 20 signal intensity was in the MRM\_Linoleic acid sample, followed by MRM\_NatCholinePC40, KPhos6\_Linoleic acid, MRM\_BeefHeart, KPhos6\_BeefHeart, MRM\_Canola, MRM\_BioLipon95, KPhos6\_Oleic Acid, and MRM\_Oleic Acid samples. 3-octanone was detected in the MRM\_Oleic Acid, KPhos6\_Linoleic acid, and MRM\_Linoleic acid samples. For 1-octen-3-one, the strongest signal intensity was in the 25 MRM\_Linoleic acid and MRM\_BeefHeart samples, followed by KPhos6\_Linoleic acid, MRM\_NatCholinePC40, KPhos6\_BeefHeart, MRM\_BioLipon95, MRM\_Oleic Acid, and KPhos6\_Oleic Acid samples. For 3-octen-2-one, the strongest signal intensity was in the KPhos6\_Linoleic acid sample, followed by MRM\_Linoleic acid, MRM\_NatCholinePC40, KPhos6\_BeefHeart, KPhos6\_Oleic Acid, MRM\_Oleic Acid, 30 MRM\_BeefHeart, MRM\_BioLipon95, MRM\_Canola, Kphos\_BioLipon95, and

Kphos\_NatCholinePC40. Pyrazine was detected in the MRM\_Coconut, MRM\_C18, MRM\_C14, and MRM\_BioLipon95 samples.

In samples having a nutty and roasted aroma, thiazole and 2-acetylthiazole were the most abundant compounds detected, along with pyrazine, methyl pyrazine, trimethyl pyrazine, and 3-ethyl-2,5-dimethylpyrazine. 2-acetylthiazole was detected in all samples with MRM and most abundant in samples with MRM\_Beefheat, MRM\_biolipon95, MRM\_Canola, and MRM\_coconut. Thiazole was created in samples with MRM-Coconut, MRM\_BeefHeat, MRM\_Biolipon95, MRM\_C14, MRM\_C18, MRM\_Canola, MRM\_Oleic acid and MRM\_Linoleic acid and MRM\_NatCholinePC40. Pyrazine was present in the largest amount in samples with MRM-Coconut, followed by samples MRM\_BeefHeat, MRM\_Biolipon95, MRM\_C14, MRM\_C18, MRM\_Canola having roughly equal amount, MRM\_Oleic acid and MRM\_Linoleic acid sample had even less. Methyl-pyrazine was present in MRM\_Biolipon95 and MRM\_Coconut. 3-ethyl-2,5-dimethyl- pyrazine and trimethyl- pyrazine, were present only without phospholipids in the MRM.

In samples having green, vegetable, or grass aromas, 1-heptanol, 1-hepten-3-ol, 1-hexanol, (E)-2-heptenal, (Z)-2-heptenal, (E)-2-hexenal, 2-pentyl-furan, and/or heptanal were detected in the KPhos6\_BeefHeart, MRM\_BeefHeart, Kphos\_BioLipon95, MRM\_BioLipon95, Kphos\_NatCholinePC40, MRM\_NatCholinePC40, Kphos\_C14, MRM\_C14, Kphos\_C18, MRM\_C18, MRM\_Canola, MRM\_Coconut, KPhos6\_Oleic Acid, MRM\_Oleic Acid, KPhos6\_Linoleic acid, and MRM\_Linoleic acid samples. For 2-pentyl-furan, the strongest signal intensity was in the KPhos6\_BeefHeart sample, followed by the KPhos6\_Linoleic acid, MRM\_BioLipon95, MRM\_Linoleic acid, MRM\_BeefHeart, MRM\_Oleic Acid, MRM\_NatCholinePC40, MRM\_Canola, KPhos6\_Oleic Acid, and Kphos\_NatCholinePC40 samples. For (E)-2-heptenal, the strongest signal intensity was in the MRM\_BeefHeart, MRM\_Canola, MRM\_Oleic Acid, and KPhos6\_Linoleic acid samples, followed by the KPhos6\_Oleic Acid, MRM\_BioLipon95, KPhos6\_BeefHeart, MRM\_Linoleic acid, MRM\_NatCholinePC40, Kphos\_BioLipon95, and Kphos\_NatCholinePC40 samples. For (Z)-2-heptenal, the strongest signal intensity was in the MRM\_Linoleic acid sample. MRM\_Linoleic acid

also was detected in the KPhos6\_Linoleic acid sample. For heptanal, the strongest signal intensity was in the MRM\_Oleic Acid sample, followed by the KPhos6\_Oleic Acid, MRM\_C14, MRM\_C18, MRM\_Canola, MRM\_BeefHeart, MRM\_NatCholinePC40, MRM\_Linoleic acid, and KPhos6\_BeefHeart samples. For , (E)-2-hexenal, the strongest signal intensity was in the MRM\_Linoleic acid sample, followed by the MRM\_NatCholinePC40, KPhos6\_Linoleic acid, and MRM\_Oleic Acid samples.

#### **Example 11 - Creation of beefy flavors using complex precursor mixtures**

A formulation was prepared (the “magic mix,” see Table 13 containing the estimated concentrations of amino acids, sugars, and other small molecules in beef based on their values reported in literature. The magic mix was tested for its ability to produce beefy flavors in the presence of LegHemoglobin (LegH). The magic mix and 1% w/v LegH were added to the meat replica, pH 6.0 (see Table 4) and baked in a convection oven for 7 minutes at 160°C. A control sample was prepared by adding 1% w/v LegH to the meat replica, pH 6.0 and baking in a convection oven for 7 minutes at 160°C.

The meat replica sample containing only LegH, was compared to the meat replica sample containing the magic mix and LegH by a sensory panel and GCMS analysis. Five tasters rated the flavored meat replicas for beefiness, bitterness, and levels of savory flavors, and off flavors. Each property was rated on a 7 point scale in which 7 was the highest amount of the specified property (e.g., a standard 80:20 ground beef would be rated 7 on the beefy scale). The Magic Mix flavor was rated one point higher in beefy character than the LegH only sample (FIG. 1).

To determine which chemical products were produced upon heating, a solution of Magic Mix was prepared with 1% w/v LegH at pH 6.0. The samples were cooked with shaking at 150°C for three minutes, then Solid Phase Micro Extraction (SPME) was performed for twelve minutes at 50°C to extract the volatile compounds above the headspace of the reaction. A search algorithm was used to analyze the retention time and mass fingerprint information of the volatile compounds and assign chemical names to peaks. Table 14 shows the compounds identified in both the Magic Mix + LegH (MM, average of two samples) and in the LegH alone in buffer (LegH, average of five samples)

samples. The compounds in Table 14 are listed in order of the retention time (R.T., in seconds), and are designated as having a zero peak area (0), or a small (S), medium (M), or large (L) average peak area. Hundreds of compounds were identified between the samples, many of which are characteristic of beefy aroma, including but not limited to 5 1,3-bis(1,1-dimethylethyl)-benzene, 2-methyl 3-furanthiol, and Bis(2-methyl-4,5-dihydro-3-furyl) disulfide, which increased in the samples containing the Magic Mix and LegH.

10 **Table 13**  
**Chemical entities added to the Magic Mix**

<u>Chemical entity</u>	<u>mM</u>
Alanine	5.6
Arginine	0.6
Asparagine	0.8
Aspartate	0.8
Cysteine	0.8
Glutamic acid	3.4
Glutamine	0.7
Glycine	1.3
Histidine	0.6
Isoleucine	0.8
Leucine	0.8
Lysine	0.7
Methionine	0.7
Phenylalanine	0.6
Proline	0.9
Threonine	0.8
Tryptophan	0.5
Tyrosine	0.6

Valine	0.9
glucose	5.6
Ribose	6.7
Maltodextrin	5.0
Thiamine	0.5
GMP	0.24
IMP	0.6
Lactic acid	1.0
creatine	1.0
NaCl	10
KCl	10
Kphos pH 6.0	10

**TABLE 14**  
**Compounds identified with GC-MS analysis in samples with MM and LegH, or**  
**LegH alone (average of five samples)**

R.T.(s)	Name	MM with LegH	LegH alone
248	acetaldehyde	L	S
256.3	carbon disulfide	L	S
264.3	dimethyl sulfide	S	0
265	oxalic acid, isobutyl pentyl ester	M	0
268.1	2,3,4-trimethyl-pentane	M	0
269.2	methanethiol	S	0
283.4	propanal	M	0
285.4	octane	M	0
287.1	furan	M	0
295.3	2-methyl-propanal	L	S
297.6	acetone	L	S
319.3	2-propenal	M	S
338.1	2-methyl-furan	M	S
342.1	butanal	L	S

344.2	2,4-dimethyl-1-heptene	M	0
346.3	methacrolein	M	0
357.4	methyl-thiirane	L	0
360.2	3-methyl-furan	S	0
363.7	butanone	L	S
368.9	2,3-dihydro-5-methyl-furan	M	S
376.4	2-methyl-butanal	L	M
381.1	3-methyl-butanal	L	M
390.6	isopropyl alcohol	0	S
399.6	ethanol	L	M
406.2	2-propenoic acid, methyl ester	M	0
408.2	benzene	S	0
414.4	methyl vinyl ketone	M	0
416.4	2,2,4,6,6-pentamethyl-heptane	M	0
422.6	2-ethyl-furan	S	0
438.4	2-ethylacrolein	M	0
449.9	2-pantanone	S	0
453.2	pentanal/2,3-butanedione	L	0
453.8	2,3-butanedione	L	M
472.8	4,7-dimethyl-undecane	M	S
485.9	2-methyl-pentanal	M	0
492.6	2-methyl-1-penten-1-one	S	0
496.6	(E)-3-penten-2-one	M	0
508.6	1-penten-3-one	M	0
510.6	trichloromethane	M	M
520.4	p-dithiane-2,5-diol	M	0
525.5	3-methyl-pantanone	M	0
535.1	(E)-5-decene	M	0
536.5	toluene	M	S
537.9	2-butenal	M	S
543.8	4-penten-2-one	M	0
550.8	methyl thiolacetate	M	0
683.7	p-xylene	S	0
727.4	dimethyl selenone	M	0
738.3	methyl isopropyl disulphide	M	0
755	2-heptanone	M	0
758.7	heptanal	L	0
781.9	1,3-diisopropoxy-1,3-dimethyl-1,3-	S	M

	disilacyclobutane		
789.4	3-methyl-2-butenal	M	0
793.4	4-methyl-2-heptanone	M	0
810.4	pyrazine	M	0
818.8	isothiazole	S	0
827.1	acetyl valeryl	M	0
831.8	2-pentyl-furan	L	0
851	2-methyl-thiazole	S	0
853.3	isothiocyanato-methane	S	0
870.9	thiazole	L	0
879.2	styrene	M	0
890.7	1-(methylthio)-propane	M	0
895.6	methyl-pyrazine	M	0
910.5	thiocyanic acid, methyl ester	S	0
918.6	4-methylthiazole	M	0
921.4	2-octanone	M	0
923.9	2-methyl-cyclopentanone	M	0
927.9	octanal	L	S
934.3	tridecane	M	0
948.8	trans-2-(2-pentenyl)furan	S	0
961.9	1-hydroxy-2-propanone	M	0
974.5	(E)-2-heptenal	M	0
987.4	5-methyl-1-undecene	M	0
993.8	2-hexyl-furan	M	0
1007.8	7-methyl-(E)-5-undecene	M	0
1024.1	2-methyl-5-(methylthio)-furan,	S	0
1058.6	2-butyl-1-decene	M	0
1079.3	dimethyl trisulfide	L	S
1085.3	2-nonenone	M	0
1093.2	nonanal	L	M
1142.3	1,3-bis(1,1-dimethylethyl)-benzene	M	0
1149.6	(E)-2-octenal	M	0
1164.5	1-heptanol	M	0
1193.5	methional	L	0
1198.8	acetic acid	M	S
1207.2	furfural	M	0
1242.1	2-decanone	M	0
1250.8	decanal	M	0

1265.2	1-decen-3-one	M	0
1283.3	pyrrole	M	0
1292.6	5-ethenyl-4-methyl-thiazole	M	0
1294.3	benzaldehyde	L	M
1303.7	2-n-octylfuran	M	0
1305.6	(E)-2-nonenal	M	0
1341.4	1-octanol	M	0
1361.1	2-methyl-1(H)-pyrrole	S	0
1391.7	2-undecanone	M	0
1401.2	(E)-2-octen-1-ol	M	0
1448	butyrolactone	S	S
1456.3	(E)-2-decenal	M	0
1462.4	phenylacetaldehyde	L	S
1466.3	2-acetylthiazole	L	0
1471.3	acetophenone	M	S
1475.4	1-nonanol	M	0
1487	methyl (methylthio)methyl disulfide	M	0
1497.1	5-(2-chloroethyl)-4-methylthiazole	L	0
1497.5	1-(ethylthio)-2-(methylthio)-buta-1,3-diene	L	S
1512	3-thiophenecarboxaldehyde	M	0
1518.8	2-non-en-4-one	M	0
1531.7	2-thiophenecarboxaldehyde	S	0
1543.9	dodecanal	M	0
1551.6	4-ethyl-2-methyl-pyrrole	S	0
1558.2	3-(methylthio)-propanenitrile	S	0
1561.2	3-decen-2-one	M	0
1613.1	bis(2-methyl-4,5-dihydro-3-furyl) disulfide	M	0
1615.6	1,10-undecadiene	M	0
1619.5	2-undecenal	S	0
1668.9	2-phenylpropenal	M	0
1692.3	(Z)-3-decen-1-ol, acetate	M	0
1733.1	3-phenyl-furan	S	0
1739.7	4-nitrophenyl 2-thiophenecarboxylic acid ester	S	0
1741.2	5-formyl-4-methylthiazole	M	0
1749.7	pentanoic acid, 2,2,4-trimethyl-3-hydroxy-, isobutyl ester	M	0
1765.5	benzyl alcohol	S	0

1774.2	pentanoic acid, 2,2,4-trimethyl-3-hydroxy-, isobutyl ester	S	0
1796.9	dodecanal	M	0
1806.1	(1-ethyl-1-propenyl)-benzene	S	0
1825.6	1-undecanol	M	S
1827.9	2-methyl-3-furanthiol	M	0
1828.3	2-methyl-3-(methylthio) furan	M	0
1836.1	4-chloro-2,6-bis(1,1-dimethylethyl)-phenol	S	0
1844.1	phenol	S	S
1845.3	[(methylsulfonyl)methyl]-benzene	S	0
1850.3	(e)-2-tridecen-1-ol	M	0
1859.9	1-heptyl-1,2,3,4-tetrahydro-4-methyl-naphthalene	S	0
1863.2	2,4-decadienal	S	0
1905.1	3,3'-dithiobis[2-methyl]-furan	M	0
1906.9	3,5-di-tert-butylbenzoic acid	S	0
1909.6	4-ethoxy-benzoic acid, ethyl ester	S	0
1921.5	3-(phenylmethyl)-2,5-piperazinedione	S	0
1944.5	9-octadecenal	M	0
1959.7	3,5-bis(1,1-dimethylethyl)-phenol	M	S
1968.4	4-methyl-5-thiazoleethanol	M	S
2007.8	1,1'-(1,2-cyclobutanediyl)bis-cis-benzene	S	0
2019.8	benzoic acid	S	S
2026.4	4-quinolinecarboxaldehyde	S	0
2027.8	m-aminophenylacetylene	M	0

**Example 12 – Ferrous Chlorin Catalyzes Production of Meat-like flavor compounds**

5 Fresh green spinach (10 lb) was added to 500 mL water and finely ground in a Vitamix blender to yield 2 L of green suspension. Acetone (8L) was added with mixing and the material was allowed to extract for 1 hour. The material was filtered through Whatman filter paper and the acetone was removed on a rotary evaporator (Buchi). To the residual green suspension (500 mL) was added 2 mL of 10 M HCl, causing the suspension to turn brown. To this was added 1g of FeCl<sub>2</sub>.4H<sub>2</sub>O in 10 mL H<sub>2</sub>O. The

10

5 solution was shaken then left at 4°C for 16 hours. This suspension was extracted with diethyl ether (3x50mL) to give a bright green organic phase, the combined organics were washed with saturated sodium chloride solution, dried over sodium sulfate, filtered and evaporated to leave a black paste (1.1g). The pellet was dissolved in chloroform for fractionation.

10 Chlorophyll and Ferrous chlorin crude fractions were stored at -20°C. Crude extracts were fractionated by reverse-phase high-pressure liquid chromatography (RP-HPLC). HPLC conditions are outlined in Table 15. Both chlorophyll and ferrous chlorophyll were eluted from the column with a peak retention time of 7.6 minutes. Eluted material was collected from 7.3 – 8.0 minutes. Collected fractions were pooled and stored on ice. Collected fractions were re-chromatographed and showed a single peak with retention time 7.6 minutes. The desired fractions were pooled, then 10% sunflower oil was added, methanol was removed on a rotary evaporator (Buchi).

15

**TABLE 15**

**HPLC conditions for purification of chlorophyll and ferrous chlorin from crude extract.**

Sample:	Chlorophyll or Fe-chlorin (~2mg/mL in CHCl <sub>3</sub> )
System:	Agilent 1100 with Chemstation
Column:	Zorbax Bonus-RP (4.6 x 250 mm, 5uM)
Mobile phase:	acetonitrile, methanol, ethyl acetate (60:20:20) isocratic flow
Temperature:	30°C
Flow Rate:	1.0 mL per minute
Injection volume:	0.05 mL

**Preparation of Flavor Reaction Containing Ferrous Chlorin or Leghemoglobin**

20 A solution of ferrous chlorophyll was mixed with the Magic Mix (Table 13) to a final concentration of 0.35% ferrous chlorin, 1% glycerol, 0.005% tween-20, 5% sunflower oil, 100 mM NaCl, 20 mM phosphate at pH 6. Leghemoglobin (0.35%) at pH 6 in phosphate buffer (20 mM), 100 mM NaCl, was mixed with the Magic Mix (Table

13), 1% glycerol, and 0.005% tween-20. The flavor reaction mixtures were heated to 150°C for 3 minutes; this reaction created flavor compounds known to be present in meat, created by hemoglobin and also created by ferrous chlorin; *see* Table 16.

The characteristic flavor and fragrance components were mostly produced during the cooking process when the flavor precursor molecules reacted with the heme-protein or the ferrous chlorophyll. Samples were evaluated by GCMS to identify the flavor compounds generated after heating. Volatile chemicals were isolated from the headspace around the flavor reactions. The profile of the volatile chemicals in the headspace around the flavor reaction mixtures that were similar between heme-protein and ferrous chlorin are shown in Table 16. Notably, many of the compounds created by the ferrous chlorin are important in the flavor of meat.

**TABLE 16**

**Flavor Compounds created by both Ferrous Chlorin and LegH with Magic Mix.**

1-heptanol	acetone
1-hexanol	acetonitrile
1-octanol	benzaldehyde
1-octen-3-ol	butanal
1-octen-3-one	2-methyl-butanal
1-pentanol	dimethyl trisulfide
2-acetylthiazole	ethyl acetate
2-butenal	furan
3-methyl-2-butenal,	2-ethyl-furan
(Z)-2-decenal	2-hexyl furan
6-methyl-2-heptanone	2-pentyl-furan
(E)-2-heptenal	furfural
(E)-2-hexenal	heptanal
2-methyl-3-furanthiol	aminophenylacetylene
(E)-2-nonenal	methacrolein
(E)-2-octenal	methional
2-pentanone	octanal
1-hydroxy-2-propanone	octane
2-thiophenecarboxaldehyde	oxalic acid, diallyl ester
2-undecenal	2,3-butanedione

3-methyl-3-buten-2-one	2-methyl-propanal
3-thiophenecarboxaldehyde	pyrazine
(E)-4-octene,	2,3-dimethyl-pyrazine
methyl-pyrazine	2,5-dimethyl-pyrazine
thiazole	

### Example 13 - Flavor creation by immobilized Hemin

#### Preparation of hemin linked CM Sepharose.

200 mg of bovine hemin (Sigma Aldrich) was loaded into a scintillation vial. A small magnetic stir bar, 800  $\mu$ L acetonitrile, 64  $\mu$ L 4-methylmorpholine, and 71 mg of N-hydroxysuccinimide were added in that order. The vial was placed in an ice bath and chilled then 118 mg of N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide hydrochloride was added with stirring, followed by 845  $\mu$ L of Jeffamine ED900. This was stirred while allowing the black mixture to warm to ambient temperature. Chloroform (10 mL) was added to the mixture followed by water (4 mL). A flashlight was used to distinguish between organic and aqueous layers since both were black and the organic layer was pipetted off and concentrated to a dark black oil. The oil was dissolved in a 4:1 mixture of acetonitrile and ethanol to make an approximately 10% strength solution that was inky black in color.

2 mL of water swelled and equilibrated CM Sepharose was equilibrated in a BioRad minicolumn with 3 volumes of acetonitrile. The resin was resuspended in 1 mL acetonitrile and pipetted into a scintillation vial. This was followed with 44 microliters 4-methylmorpholine, 23 mg N-hydroxysuccinimide, and 39 mg of solid N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide hydrochloride. The mixture was vortexed vigorously and then shaken for three hours. To this white solid was added 570 microliters of inky black 20% strength hemin coupled diamine. The black solid was vortexed and shaken for an hour. The slurry strongly resembled Turkish coffee. The mixture was poured into a BioRad minicolumn and filtered, washed with acetonitrile until what came out no longer resembled espresso, then switched to deionized water, and finally 20 mM pH 9 sodium carbonate buffer. The black solid was washed until the effluent ran clear and then resuspended in 2 mL of buffer for storage until use.

### Flavor reaction

The flavor reaction was created with heme protein (equine myoglobin -Sigma) at 0.35% in a phosphate buffer (20 mM) at pH 6.0 with 100 mM NaCl, this was mixed with 5 Magic Mix (Table 13). Another flavor reaction was created with Immobilized Hemin at 0.35% in a phosphate buffer (20 mM) at pH 6.0 with 100 mM NaCl, this was mixed with Magic Mix (Table 13). The flavor reaction mixtures were heated to 150°C for 3 minutes; this reaction created flavor compounds known to be present in meat.

10 The characteristic flavor and fragrance components were mostly produced during the cooking process when the flavor precursor molecules reacted with the Heme-protein or the immobilized Hemin. Samples were evaluated by GCMS to identify the flavor compounds generated after heating. Volatile chemicals were isolated from the headspace around the flavor reactions. As can be seen in Table 17, immobilized hemin catalyzed 15 production of compounds similar to those whose production was catalyzed by myoglobin free in solution. Notably, the profiles of flavor compounds, measured by GCMS, produced by cooking mixtures containing the immobilized hemin and the heme-protein, respectively, were very similar.

20

**TABLE 17**

**Flavor compounds produced by cooking mixtures containing either myoglobin free  
in solution or hemin coupled to a solid support**

Flavor compound	myoglobin	hemin-linker-resin
2-methyl-5-(methylthio)-thiophene	Low	
dihydro-5-propyl-2(3H)-furanone	Low	
octane	Low	
pyrrole	Low	Low
methanethiol	Low	Low
2-thiophenecarboxaldehyde	Low	Low
methyl-pyrazine	Low	Low
1-hydroxy-2-propanone	Low	Low
propanal	Low	Low

thiophene	Low	medium
pyridine	Low	Low
2-methyl-furan	Low	medium
oxalic acid, butyl propyl ester	Low	Low
pyrazine	medium	Low
oxalic acid, diallyl ester	medium	medium
2-butenal	medium	large
furfural	medium	medium
nonanal	medium	medium
2-ethyl-furan	medium	Low
ethanol	medium	very large
tert-butanol	medium	
3,3'-dithiobis[2-methyl]-furan	medium	medium
m-aminophenylacetylene	medium	medium
2,5-dihydro-3,4-dimethyl-furan	medium	medium
2-acetylthiazole	medium	medium
cyclohexane	medium	
ethyl tert-butyl ether	medium	
carbon disulfide	medium	medium
thiazole	medium	medium
acetonitrile	medium	large
2-pentyl-furan	medium	Low
3-thiophenecarboxaldehyde	medium	medium
2-methyl-butanal	medium	medium
thiazole	medium	large
2-methyl-3-furanthiol	large	large
2-propenal	large	large
3-methyl-2-butenal	large	medium
2-methyl-3-(methylthio) furan	large	large
ethyl acetate	large	medium
methacrolein	large	medium
methyl-thiirane	large	large
methional	large	large
methyl alcohol	large	medium
2-butanone	large	Low
2,3-butanedione	large	medium
acetone	large	large
furan	large	medium

benzaldehyde	large	medium
methyl thiolacetate	large	medium
acetaldehyde	very large	very large
2-methyl-propanal	very large	very large
dimethyl trisulfide	very large	very large
3-methyl-butanal	very large	very large
propyl-cyclopropane		medium
(E)-2-octenal		medium
2-n-propylaziridine		medium
thiirane		medium
ethyl formate		medium
methyl vinyl ketone		medium
2-propenoic acid, ethyl ester		medium
1-nonal		large
1-octene		large
1-heptanol		large
1-dodecene		large
phorone		very large

**Example 14. The combination of precursors with Heme protein drives flavor reactions.**

Three samples were compared: precursor mix alone, 1% heme protein alone, and precursor mix with 1% heme. The precursor mix was made of glucose (20 mM), ribose (20 mM), cysteine (10 mM), thiamine (1 mM), and glutamic acid (1 mM). Reactions were all at pH 6.0, prepared and heated to 150°C for 3 minutes. These three samples were run in duplicate. These samples were evaluated by GCMS for the flavor compounds generated. Characteristic flavor and fragrance components were mostly produced during the cooking process where precursors could react with the heme-protein. These samples were evaluated by GCMS for the flavor compounds generated and evaluated for the sensory experience. Volatile chemicals were isolated from the head space around the flavor reaction. The flavor compounds created in each sample is indicated in Table 18. As shown most of the flavor molecules were created on when the precursors are combined with the heme protein.

**Table 18**  
**Flavor molecules created by the combination of LegH and precursor mix.**

<b>Compound</b>	<b>Precursor mix</b>	<b>LegH</b>	<b>Precursor mix + Leg H</b>
carbon disulfide	medium	medium	high
isopropyl alcohol	medium	medium	low
2-methyl-furan	low		low
butanal	low		medium
thiophene	low		low
2,3-butanedione	low	low	high
furan	low		medium
2,4-dimethyl-1-heptene		high	high
acetone		high	high
dimethyl trisulfide		medium	medium
2-methyl-heptane		medium	medium
2-pentanone		medium	
pentanal		medium	medium
2-pentyl-furan		medium	medium
2-methyl-propanal		low	high
2-acetyl-1-propene		low	low
2-methyl-butanal		low	medium
1,3-dimethyl-benzene		low	low
octane		low	low
benzene		low	low
benzaldehyde			very high
2-butanone			very high
furfural			very high
thiazole			high
nonanal			high
thiazole			high
2-acetylthiazole			medium
3-methyl-butanal			medium
(Z)-2-heptenal			medium
heptanal			medium
methyl-thiirane			medium
3-ethyl-pentane			medium
phenylacetaldehyde			medium
2-hexyl-furan			medium

2-nonanone			medium
propanal			medium
pyrazine			medium
(Z)-2-heptenal			medium
2-methyl-1-heptene			medium
2-ethyl-furan			medium
octanal			medium
(E)-4-octene			low
(E)-2-octenal			low
2-methyl-thiazole			low
2-propenal			low
1-octen-3-one			low
1-octene			low
2-octanone			low
dimethyl sulfide			low
3-pentyl-furan			low
2-n-octylfuran			low
2-pentyl-thiophene			low

## OTHER EMBODIMENTS

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

## CLAIMS

1. A ground beef-like food product, comprising:
  - a) 0.1%-5% by weight of a heme-containing protein;
  - b) a compound selected from glucose, ribose, fructose, lactose, xylose, arabinose, glucose-6-phosphate, maltose, and galactose, and mixtures of two or more thereof;
  - c) at least 1.5 mM of a compound selected from cysteine, cysteine, thiamine, methionine, and mixtures of two or more thereof; and
  - d) one or more plant proteins,  
wherein the ground beef-like food product contains no animal product;  
wherein cooking the ground beef-like food product results in the production of at least two volatile compounds which have a beef-associate aroma.
2. The ground beef-like food product of claim 1, comprising at least 10 mM of a compound selected from cysteine, cysteine, thiamine, methionine, and mixtures of two or more thereof.
3. The ground beef-like food product of claim 1 or claim 2, wherein the heme-containing protein is globin.
4. The ground beef-like food product of claim 1 or claim 2, wherein the heme-containing protein comprises an amino acid sequence having at least 80% sequence identity to a polypeptide set forth in SEQ ID Nos. 1-26.
5. The ground beef-like food product of any one of claims 1 to 4, further comprising one or more of inosine, inosine monophosphate (IMP), guanosine, guanosine monophosphate (GMP), and adenosine monophosphate (AMP).
6. The ground beef-like food product of any one of claims 1 to 4, further comprising one or more of beta-carotene, alpha-tocopherol, caffeic acid, propyl gallate, and epigallocatechin gallate.
7. The ground beef-like food product of any one of claims 1 to 4, further comprising one or more of a vegetable oil, an algal oil, sunflower oil, corn oil, soybean oil, palm fruit oil, palm kernel oil, safflower oil, flaxseed oil, rice bran oil, cottonseed oil, olive oil, canola oil, flaxseed oil, coconut oil, and mango oil.

8. The ground beef-like food product of any one of claims 1 to 4, comprising a textured vegetable protein.
9. The ground beef-like food product of any one of claims 1 to 8, wherein the ground beef-like food product has a pink to red color before cooking to indicate a raw or uncooked state.
10. The ground beef-like food product of any one of claims 1 to 9, wherein at least a portion of the ground beef-like food product, upon cooking, transitions in color from a pink to red color in a raw or uncooked state to a lighter pink to brown color in a partially cooked or fully cooked state.
11. The ground beef-like food product of any one of claims 1 to 10, comprising about 5.6 to about 20 mM of the compound selected from glucose, ribose, fructose, lactose, xylose, arabinose, glucose-6-phosphate, maltose, and galactose, and mixtures of two or more thereof.
12. The ground beef-like food product of any one of claims 1 to 11, comprising about 0.8 mM to about 10 mM cysteine.
13. The ground beef-like food product of any one of claims 1 to 12, comprising about 0.1 mM to about 2 mM thiamine.
14. The ground beef-like food product of any one of claims 1 to 13, wherein the at least two volatile compounds are selected from 2-methyl-furan, bis(2-methyl-3-furyl)disulfide, 2-pentyl-furan, 3,3'-dithiobis-2-methyl-furan, 2,5-dimethyl-pyrazine, 2-methyl-3-furanthiol, dihydro-3-(2H)-thiophenone, 5-methyl-2-thiophenecarboxaldehyde, 3-methyl-2-thiophenecarboxaldehyde, 2-methyl-thiazole, dimethyl sulfide, decanal, 5-ethyldihydro-2(3H)-furanone, dihydro-5-pentyl-2(3H)-furanone, 2-octanone, 3,5-octadien-2-one, p-Cresol, and hexanoic acid.
15. The ground beef-like food product of any one of claims 1 to 14, wherein cooking comprises heating the ground beef-like food product at 150°C for about 3 to about 5 minutes.
16. The ground beef-like food product of any one of claims 1 to 15, further comprising one or more of acetic acid, lactic acid, glycolic acid, citric acid, succinic acid, tartaric acid, caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid, linoleic acid, alpha linolenic acid, gamma linolenic acid, arachidic acid, arachidonic acid, behenic acid, and erucic acid.

17. The ground beef-like food product of any one of claims 1 to 16, wherein cooking the ground beef-like food product results in the production of at least five volatile compounds which have a beef-associated aroma.
18. The ground beef-like food product of any one of claims 1 to 17, wherein cooking the ground beef-like food product results in the production of at least ten volatile compounds which have a beef-associated aroma.
19. The ground beef-like food product of any one of claims 1 to 18, wherein cooking the ground beef-like food product results in the production of at least twenty volatile compounds which have a beef-associated aroma.
20. The ground beef-like food product of any one of claims 1 to 19, wherein the at least two volatile compounds are 2-methyl-furan and bis(2-methyl-3-furyl)disulfide.

**Impossible Foods Inc.**  
**Patent Attorneys for the Applicant/Nominated Person**  
**SPRUSON & FERGUSON**

## FIGURE 1

SEQ ID NO:1 *Vigna radiata*

MTTTLERGFTEEQEALVVKSWNVMKKNSGELGLKFFLKIFEIAPSAQKLFSFLRDSTVP  
LEQNPKLKPHAVSVFVMTCDASVQLRKAGKVTVRESNLKKGATHFRTGVANEHFEVTK  
FALLETIKEAVPEMWSPAMKNAWGEAYDQLVDAIKYEMKPPSS

SEQ ID NO:2 *Methylacidiphilum infernorum*

MIDQKEKELIKESWKRIEPNKNEIGLLFYANLFKEEPTVSVLFQNPISQSRSRKLMQVLG  
ILVQGIDNLEGLIPTLQDLGRRHKQYGVVDSHYPLVGDCLLKSIQEYLGQGFTEEAKAA  
WTKVYGINAAQVMTAE

SEQ ID NO:3 *Aquifex aeolicus*

MLSEETIRVIKSTVPLLKEHGTEITARMYELLFSKYPKTKELFAGASEEQPKKLANAI  
AYATYIDRLEELDNAISTIARSHVRRNVKPEHYPLVKECLLQAIIEEVLPGEEVLKAW  
EAYDFLAKTLITLEKKLYSQP

SEQ ID NO:4 *Glycine max*

MGAFTEKQEALVSSSFEAFKANI PQYSVVFYTSILEKAPAAKDLFSFLSNGVDPSNPKL  
TGHAEKLFGLVRDSDAGQLKANGTVVADAALGSIHAQKAITDPQFVVVKEALLKTIKEAV  
GDKWSDELSSAWEVAYDELAAAIKKAF

SEQ ID NO:5 *Hordeum vulgare*

MSAAEGAVVFSEEKEALVLKSWAIMKKDSANLGLRFFLKIFEIAPSARQMFPLRDSDV  
PLETNPKLKTHAVSVFVMTCEAAAQLRKAGKITVRETTLKRLGGTHLKYGVADGHFEVT  
RFALLETIKEALPADMWGPEMRNAWGEAYDQLVAAIKQEMKPAE

*Magnaporthe oryzae*, (SEQ ID NO:6)

1 mdgavrlldwt gldltgheih dgvpriasrvq vmvspfplfkd qhiimsskes psrksstigg  
61 strngscqad tqkqqlppvg ekpkpvkenp mkklikemsqr plptqhgdt yptekkltg  
121 gedlkhirgy dvktllamvk sklkgek1kd dktm1mervm qlvarlptes kkraeltdsl  
181 inelwesldh pp1n1ylgpeh syrtpdgsyn hpfnpqlgaa gsryarsvip tvtppgalpd  
241 pglifdsimg rtpnsyrkhp nnvssilwyw atiiihd1fw tdprrdintnk sssyldlapl  
301 ygnsgemqds irtfkdgrmk pdcyadkrla gmppgvs11 imfnrfhnhv aenlalineg  
361 grfnkpsd11 egeareaawk kydndlqva r1vtsglyin it1vdyvrni vnlrvdttw  
421 tlaprqdaga hvgtadgaer gtgnavsaef nlcyrwhsci sekdskfvea qfqnifgkpa  
481 sevrpdemwk gfakmeqntp adpgqrtfgg fkrgpdgkfd dddlvrcise avedvagafg  
541 arnvpqamkv vetcgiiqgr kwnvaglnef rkhfh1kpy5 tfedinsdpg vaealrrlyd

601 hpdnvelypg lvaeedkqpm vpgvgiaptv tisrvvlsda vclvrgdrfy ttdftprnlt  
 661 nwgykevdyd lsvnhgcvfy klfirafpnh fkqnsvyahy pmvvpsenkr ilealgradl  
 721 fdfeapkyip prvnitsygg aeyiletqek ykvttwheglg flmgeggglkf mlsqddplha  
 781 qqrkcmmaql ykdgwteavk aftyagmmeel lvsksyflgn nkhrhvdair dvgnmvvhvf  
 841 asqvfqplk taknptgvft eqemygila ifttiffdld psksfplrtk trevcqklak  
 901 lveanvklin kipwsrgmfv gkpakdepls iygkmtikgl kahglsdydi awshvvptsg  
 961 amvpnqaqvf aqavdyylsp agmhyipeih mvalqpstpe tdalllgyam egirlagtfq  
 1021 syreavddv vkedngrqvp vkagdrvfts fvdaardpkh fpdpesvnpn rpakkyihyg  
 1081 vgphaclgrd asqiaitemf rclfrnnvr rvpqpgelk kvprpggfyv ymredwgglf  
 1141 pfpvtrmrvmw dde

*Fusarium oxysporum* (SEQ ID NO:7)

1 mkgsatlafa lvqfsaasql vwpskwdeve dlymqggfn krgfadalrt cefgsnvpqt  
 61 qntaewlra fhdaitdhak agtggldasi ywessrpenp gkafnntfgf fsgfhnpnrat  
 121 asdltalgtv lavgacngpr ipfragrida ykagpagvpe pstnlkdtfa aftkaagftke  
 181 emtamvacgh aiggvhsvdf peivgikadp nndtnvpfqqk dvssfhngiv teylagtskn  
 241 plvasknatf hsdkrifdnd katmkklstf agfnsmcadl ltrmidtvpk svqltpvlea  
 301 ydvrpyitel slnnknkihf tgsvrvritn nirdnndlai nliyvgrdgk kvtvptqqvt  
 361 fqqgtsfgag evfanfefdt tmdakngitk ffiqevkpst katvthdnqk tggkvddtv  
 421 lyqlqqscav leklnaplv vtamvrdrara kdaltlrvah kkpvkgsivp rfqtaitnfk  
 481 atgkksssgyt gfqaktmfee qstyfdiulg gspasgvqfl tsqampsqcs

*Fusarium graminearum* (SEQ ID NO:8)

1 masatrqfar aatratrnf aiaprqvirq qgrryysssep aqkssssawiw ltgaavagga  
 61 gyyfygnsas satakvfnps kedyqkvne iaarleekdd yddgsygpvl vrlawhasgt  
 121 ydketgtggs ngatmrfape sdhganagla aardflqpvk ekfpwitysd lwilagvcai  
 181 qemlgpaipy rpgrsdrdvs gctpdgrlrd askrqdhlrg ifgrmgfndq eivalsgaha  
 241 lgrchtdrsg ysgpwtfspt vltndyfrll veekwqwkww ngpaqyedks tkslmmlpsd  
 301 ialiedkkfk pwvekyakdn daffkdfsnv vrlrfelgvp faqgtenqrw tfkpthqe

SEQ ID NO: 9 *Chlamydomonas eugametos*

MSLFAKLGGREAVEAAVDKFYNKIVADPTVSTYFSNTDMKVQRSKQFAFLAYALGGASE  
 WKGKDMRTAHKDLVPHLSVDVHFQAVARHLSDTLTELGVPPEDITDAMAVVASTRTEVLN  
 MPQQ

SEQ ID NO:10 *Tetrahymena pyriformis*

MNKPQTIYEKLGGENAMKAAVPLFYKKVLADERVKHFFKNTDMDHQTKQQTDFLTMLLG  
 GPNHYKGKNMTEAHKGMLQNLHFDAAI IENLAATLKEGLGVTDAVINEAAKVIEHTRKDM  
 LGK

SEQ ID NO:11 *Paramecium caudatum*

MSLFEQLGGQAAVQAVTAQFYANIQADATVATFFNGIDMPNQTNKTA AFLCAALGGPNA  
WTGRNLKEVHANMGVSNAQFTVIGHRLSALTGAGVAAALVEQTVAVAETVRGDVVTV

SEQ ID NO:12 *Aspergillus niger*

MPLTPEQIKIIKATVPVLQEYGTKITTAFYMNSTMPELNAVFTANQVKGHQARALA  
GALFAYASHIDDLGALGPAVELICNKHASLYIQADEYKIVGKYLLEAMKEVLGDACTDD  
ILDAGAAYWALADIMINREAALYKQSQG

SEQ ID NO:13 *Zea mays*

MALAEADDGAVVFGEEQEALVLKSWAVMKDKAANLGLRFFLKVFIA  
PSAEQMFSFLRD SDVPLEKNPKLKTHAMSVFVMTCEAAAQLRKAGKVT  
VRETTLKRLGATHFKYGVADGHF EVTG  
FALLETIKEALPADMWSEM  
KKAWAEAYSQ  
LVA  
AIKRE  
MKP  
DA

SEQ ID NO:14 *Oryza sativa subsp. japonica*

MALVEGNNGVSGGAVSFSEEQEALVLKSWAIMKKDSANIGLRF  
FLKIFEVAPSASQMFS  
FLRNSDVP  
LEKNPKLKTHAMSVFVMTCEAAAQLRKAGKVT  
VRETTLKRLGATHFKYGVG  
DAHFEV  
TRF  
ALLETIKEA  
VPV  
DMW  
SPAM  
KSA  
WSE  
AYN  
Q  
LVA  
AIK  
QEM  
KPAE

SEQ ID NO:15 *Arabidopsis thaliana*

MESEGKIVFTEEQEALVVKSWSVMKKNSAELGLKLF  
IKIFEIAP  
TTKKMFSFLRD  
S  
PIP  
AEQNP  
KLKPHAM  
SVF  
VMCC  
ESAV  
QLRKT  
GKV  
TRE  
TTLKRLG  
ASH  
SKYGV  
VDEH  
FEV  
AK  
Y  
ALLET  
IKEA  
VP  
PEM  
W  
PEM  
KVA  
WGQAY  
DHL  
VAA  
AIKA  
EMN  
LSN

SEQ ID NO:16 *Pisum sativum*

MGFTDKQEALVNSSWESFKQNLSGNSILFYTIILEKAPAAKGLFSFLKDTAGVEDSPKL  
QAHAEQVFGVRDSSAQLRTKGEVVLGNATLGAIHVQRGVTDPHFVVVKEALLQTIKKA  
SGNNWSEELNTAWEVAYDGLATAIKKAMT

SEQ ID NO:17 *Vigna unguiculata*

MVAFSDKQEALVNGAYEAFKANIPKYSVVFYTTILEKAPAAKNLFSFLANGVADATNPKL  
TGHAEKLFGLVRDSSAQLRASGGVVADAALGAVHSQKAVNDAQFVVVKEALVKTLEAV  
GDKWSDELGTAVELAYDELAAAIKKAY

SEQ ID NO:18 *Bos taurus*

MGLSDGEWQLVLNAWGKVEADVAGHGQEVLIRLFKGPETLEKFDKFHKHLKTEAEMKAS  
EDLKKHGNTVLTALGGILKKKGHEAEVHKLAESHANKHKIPVKYLEFISDAIIHVLHA  
KHPDSFGADAQAMSKALELFRNDMAAQYKVLGFHG

SEQ ID NO:19 *Sus scrofa*

MGLSDGEWQLVNVWGKVEADVAGHGQEVLIRLFKGPETLEKFDKFHKHLKSEDEMKA  
EDLKKHGNTVLTALGGILKKKGHEAEELPLAQSHATHKHPIVKYLEFISEAIIQVLQS  
KH PGDFGADAQGAMSKALELFRNDMAAKYKELGFQG

SEQ ID NO: 20 *Equus caballus*

MGLSDGEWQQVLNVWGKVEADIAGHGQEVLIRLFKGPETLEKFDKFHKHLKTEAEMKAS  
EDLKKHGTVVLTAALGGILKKKGHEAEELPLAQSHATHKHPIVKYLEFISDAIIHVLHS  
KH PGDFGADAQGAMTKALELFRNDIAAKYKELGFQG

SEQ ID NO: 21 *Nicotiana benthamiana*

MSSFTEEQEAJVVKSWDSMKNAGEWGLKLFKIFEIAPSACKLFSFLKDSNVPL  
EQNAKLKPHSKSVFVMTCEAAVQLRKAGKVVVRDSTLKKLGATHFKYGVADE  
HFEVTKFALLETIKEAVPEMWSVDMKNAWGEAFDQLVNAIKTEMK

SEQ ID NO: 22 *Bacillus subtilis*

MGQSFNAPYEAIGEELLSQLVDTFYERVASHPLKPIFPSDLTETARKQQQLTQY  
LGGPPLYTEEHGHPMLRARHLPFPTNERADAWLSCMKDAMDHVGLEGEIREFL  
FGRLELTARHMQVNQTEAEDRSS

SEQ ID NO: 23 *Corynebacterium glutamicum*

MTTSENFYDSVGEETFSLIVHRYEQVPNDDILGPMYPPDDFEGAEQRLKMF  
QYWGGPKDYQEQRGHPRLRMHVNVPIGVTAAERWLQLMSNALDGVDLTAEQ  
REAIWEHMVRAADMLINSNPDPHA

SEQ ID NO: 24 *Synechocystis PCC6803*

MSTLYEKLGGTTAVDLAVDKFYERVLQDDRIKHFFADVDMAKQRAHQKAFLTY  
AFGGTDKYDGRYMREAHKELVENHGLNGEHFDAVAEDLLATLKGVPEDLIA  
EVAAVAGAPAHKRDVLNQ

SEQ ID NO: 25 *Synechococcus* sp. PCC 7335

MDVALLEKSFEQISPRAIEFSASFYQNLFHHPPELKPLFAETSQTIQEKKLIFSLAAI  
IENLRNPDLQPALKSLGARHAEVGTIKSHYPLVGQALIETFAEYLAADWTEQLA  
TAWVEAYDVIASTMIEGADNPAAAYLEPELTFYEWLDLYGEESPKVRNAIATLTH  
FHYGEDPQDVQRDSRG


SEQ ID NO: 26 *Nostoc commune*

MSTLYDNIGGQPAIEQVVDELHKRIATDSLLAPVFAGTDMVKQRNHLVAFLAQIF  
EGPKQYGGRPMKDTHAGLNLQQPHFDAIAKHLGERMAVRGVSSENTKAALDR  
VTNMKGAILNK

SEQ ID NO: 27 *Bacillus megaterium*

MREKIHSPYELLGGEHTISKLVDAFYTRVGQHPELAPIFPDNLTETARKQQQLTQ  
YLGGPSLYTEEHGHPMLRARHLPFEITPSRAKAWLTCMHEAMDEINLEGP  
DELYHRLILTAQHMINSPQTDEKGFSH

FIGURE 2



60900957.doc