

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 July 2010 (08.07.2010)

(10) International Publication Number
WO 2010/077882 A2

(51) International Patent Classification:
C07D 413/10 (2006.01) *A61P 11/06* (2006.01)
C07D 261/06 (2006.01) *A61K 31/553* (2006.01)
C07D 413/06 (2006.01)

(21) International Application Number:
PCT/US2009/068105

(22) International Filing Date:
15 December 2009 (15.12.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/122,568 15 December 2008 (15.12.2008) US

(71) Applicant (for all designated States except US): **AMIRA PHARMACEUTICALS, INC.** [US/US]; 9535 Waples Street, Suite 100, San Diego, California 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HUTCHINSON, John, Howard** [US/US]; 5948 Via Zurita, La Jolla, California 92037 (US). **SEIDERS, Thomas, Jon** [US/US]; 5257 Soledad Mountain Road, San Diego, California 92109 (US). **WANG, Bowei** [US/US]; 10 Kirkview Circle, Westfield, New Jersey 07090 (US). **ARRUDA, Jeannie, M.** [US/US]; 3611 Florida St. #4, San Diego, California 92104 (US). **ROPPE, Jeffrey, Roger** [US/US]; 32182 Camino Guarda, Temecula, California 92592 (US). **PARR, Timothy** [US/US]; 8504 Lemon Avenue, La Mesa, California 91941 (US). **ZHAO, Lucy** [US/US]; 11254 Caminito Corriente, San Diego, California 92128 (US).

(74) Agents: **AKHAVAN, Ramin** et al.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, California 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))



WO 2010/077882 A2

(54) Title: ANTAGONISTS OF LYSOPHOSPHATIDIC ACID RECEPTORS

(57) Abstract: Described herein are compounds that are antagonists of lysophosphatidic receptor(s). Also described are pharmaceutical compositions and medicaments that include the compounds described herein, as well as methods of using such antagonists, alone and in combination with other compounds, for treating LPA-dependent or LPA-mediated conditions or diseases.

ANTAGONISTS OF LYSOPHOSPHATIDIC ACID RECEPTORS

RELATED APPLICATIONS

[0001] This application claims benefit of U.S. Provisional Application No. 61/122,568, entitled “ANTAGONISTS OF LYSOPHOSPHATIDIC ACID RECEPTORS” filed on December 15, 5 2008, which is herein incorporated by reference.

FIELD OF THE INVENTION

[0002] Described herein are compounds, methods of making such compounds, pharmaceutical compositions and medicaments comprising such compounds, and methods of using such compounds to treat, prevent or diagnose diseases, disorders or conditions associated with one or 10 more of the lysophosphatidic acid (LPA) receptors.

BACKGROUND OF THE INVENTION

[0003] Lysophospholipids are membrane-derived bioactive lipid mediators. Lysophospholipids affect fundamental cellular functions that include proliferation, differentiation, survival, migration, adhesion, invasion, and morphogenesis. These functions influence many biological 15 processes that include, but are not limited to, neurogenesis, angiogenesis, wound healing, fibrosis, immunity, and carcinogenesis.

[0004] Lysophosphatidic acid (LPA) is a lysophospholipid that has been shown to act through sets of specific G protein-coupled receptors (GPCRs) in an autocrine and paracrine fashion. LPA binding to its cognate GPCRs (LPA₁, LPA₂, LPA₃, LPA₄, LPA₅, LPA₆) activates intracellular 20 signaling pathways to produce a variety of biological responses. Antagonists of the LPA receptors find use in the treatment of diseases, disorders or conditions in which LPA plays a role.

SUMMARY OF THE INVENTION

[0005] In one aspect, presented herein are compounds of Formula (I), Formula (II) and Formula (III) that inhibit the physiological activity of lysophosphatidic acid (LPA), and therefore, are 25 useful as agents for the treatment or prevention of diseases in which inhibition of the physiological activity of LPA is useful, such as diseases in which an LPA receptor participates, is involved in the etiology or pathology of the disease, or is otherwise associated with at least one symptom of the disease.


[0006] In one aspect, the compounds of Formula (I), Formula (II) and Formula (III) are useful 30 for the treatment of fibrosis of organs (liver, kidney, lung, heart and the like), liver diseases (acute hepatitis, chronic hepatitis, liver fibrosis, liver cirrhosis, portal hypertension, regenerative failure, non-alcoholic steatohepatitis (NASH), liver hypofunction, hepatic blood flow disorder,

and the like), cell proliferative disease (cancer (solid tumor, solid tumor metastasis, vascular fibroma, myeloma, multiple myeloma, Kaposi's sarcoma, leukemia, chronic lymphocytic leukemia (CLL) and the like) and invasive metastasis of cancer cell, and the like), inflammatory disease (psoriasis, nephropathy, pneumonia and the like), gastrointestinal tract disease (irritable bowel syndrome (IBS), inflammatory bowel disease (IBD), abnormal pancreatic secretion, and the like), renal disease, urinary tract-associated disease (benign prostatic hyperplasia or symptoms associated with neuropathic bladder disease, spinal cord tumor, hernia of intervertebral disk, spinal canal stenosis, symptoms derived from diabetes, lower urinary tract disease (obstruction of lower urinary tract, and the like), inflammatory disease of lower urinary tract, dysuria, frequent urination, and the like), pancreas disease, abnormal angiogenesis-associated disease (arterial obstruction and the like), scleroderma, brain-associated disease (cerebral infarction, cerebral hemorrhage, and the like), neuropathic pain, peripheral neuropathy, and the like, ocular disease (age-related macular degeneration (AMD), diabetic retinopathy, proliferative vitreoretinopathy (PVR), cicatricial pemphigoid, glaucoma filtration surgery scarring, and the like). In one aspect, the compounds of Formula (I), Formula (II) and Formula (III) are used in the treatment of fibrotic diseases or conditions.

[0007] In one aspect, described herein are compounds of Formula (I), Formula (II) and Formula (III), pharmaceutically acceptable salts, pharmaceutically acceptable solvates, and prodrugs thereof. Compounds of Formula (I), Formula (II) and Formula (III) are antagonists of at least one of the LPA receptors selected from LPA₁, LPA₂, LPA₃, LPA₄, LPA₅ and LPA₆. In one embodiment, compounds of Formula (I), Formula (II) and Formula (III) are antagonists of LPA₁. In one embodiment, compounds of Formula (I), Formula (II) and Formula (III) are antagonists of LPA₁ and/or LPA₃. In some embodiments, compounds of Formula (I), Formula (II) and Formula (III) are antagonists of LPA₁ and/or LPA₂. In some embodiments, compounds of Formula (I), Formula (II) and Formula (III) are selective antagonists for one of the LPA receptors relative to the other LPA receptors. In some embodiments, such a selective antagonist is selective for the LPA₁ receptor. In some embodiments, such a selective antagonist is selective for the LPA₂ receptor. In some embodiments, such a selective antagonist is selective for the LPA₃ receptor.

[0008] Compounds of Formula (I), Formula (II) and Formula (III) are used in the treatment of diseases, disorders, or conditions in which activation of at least one LPA receptor by LPA contributes to the symptomology or progression of the disease, disorder or condition. These diseases, disorders, or conditions may arise from one or more of a genetic, iatrogenic, immunological, infectious, metabolic, oncological, toxic, surgical, and/or traumatic etiology. In one aspect, the methods, compounds, pharmaceutical compositions, and medicaments described

herein comprise antagonists of LPA receptors. In one aspect, the methods, compounds, pharmaceutical compositions, and medicaments described herein comprise antagonists of LPA₁. [0009] In one aspect, provided herein is a compound of Formula (I), pharmaceutically acceptable salt, pharmaceutically acceptable solvate, or prodrug thereof:



Formula (I)

wherein

R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -C(=O)NH-OH, -C(=O)NH-CN, -CN, -C(=NH)-NH₂, -C(=NH)-NHC(=O)R^D, -C(=O)NHCH₂CH₂SO₃H, 10 tetrazolyl or carboxylic acid bioisostere;

R^D is H or C₁-C₆alkyl;

ring A is a substituted or unsubstituted C₃-C₁₀cycloalkylene, a substituted or unsubstituted C₂-C₁₀heterocycloalkylene, a substituted or unsubstituted arylene, or a substituted or unsubstituted heteroarylene, where if ring A is substituted then ring A is substituted with 1, 2, or 3 R^A; 15

ring B is a substituted or unsubstituted C₃-C₁₀cycloalkylene, a substituted or unsubstituted C₂-C₁₀heterocycloalkylene, a substituted or unsubstituted arylene, or a substituted or unsubstituted heteroarylene, where if ring B is substituted then ring B is substituted with 1, 2, or 3 R^B;

20 L¹ is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, or C₁-C₆heteroalkylene;

L² is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, C₁-C₆heteroalkylene, -O-, -S-, -SO-, -SO₂-, -NR²-, or -C(=O)-;

R² is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

R³ is H, C₁-C₄alkyl, C₃-C₆cycloalkyl, or C₁-C₄fluoroalkyl;

25 R⁴ is -NR⁷C(=O)OCH(R⁸)-CY, -NR⁷C(=O)OC(R⁸)₂-CY, or -NR⁷C(=O)O-CY;

R⁷ is H or C₁-C₄alkyl;

R⁸ is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

CY is C₁-C₆alkyl, a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl, wherein if CY is substituted then CY is substituted with 1, 2, or 3 R^C; or 30

R⁸ and CY are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted carbocycle or a substituted or unsubstituted heterocycle;



each of R^A , R^B , and R^C are independently selected from H, halogen, -CN, -NO₂, -OH, -OR¹⁰, -SR¹⁰, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R⁹)S(=O)R¹⁰, -S(=O)N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR⁹C(=O)N(R⁹)₂, -NR⁹C(=O)R¹⁰, -NR⁹C(=O)OR¹⁰, C₁-C₄alkyl, C₁-C₄fluoroalkyl, 5 C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl; each R⁹ is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted cycloalkyl), 10 -C₁-C₄alkylene-(substituted or unsubstituted heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted heterocycle; 15 R¹⁰ is selected from C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene- substituted or unsubstituted cycloalkyl, -C₁-C₄alkylene-(substituted or unsubstituted heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl). 20



[0010] For any and all of the embodiments, substituents are selected from among from a subset of the listed alternatives. For example, in some embodiments, R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -CN, -C(=NH)-NH₂, -C(=NH)-NHC(=O)R^D, or tetrazolyl. In some embodiments, R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -CN, or 25 tetrazolyl. In some embodiments, R¹ is -CO₂H, -CO₂R^D, or -C(=O)NHSO₂R¹⁰. In some embodiments, R¹ is -CO₂H or -CO₂R^D. In some embodiments, R¹ is -CO₂H. In some embodiments, R¹ is -C(=O)NHSO₂R¹⁰.



[0011] In some embodiments, R⁴ is -NR⁷C(=O)OCH(R⁸)-CY, or -NR⁷C(=O)O-CY. In some embodiments, R⁴ is -NR⁷C(=O)OCH(R⁸)-CY.

[0012] In some embodiments, CY is a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted C₂-C₁₀heterocycloalkyl, a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl, wherein if CY is substituted then CY is substituted with 1, 2, or 3 R^C; or R⁸ and CY are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted carbocycle or a substituted or unsubstituted heterocycle. 30

[0013] In some embodiments, R^1 is $-CO_2H$, $-CO_2R^D$, $-C(=O)NHSO_2R^{10}$, $-C(=O)N(R^9)_2$, $-CN$, $-C(=NH)-NH_2$, $-C(=NH)-NHC(=O)R^D$, or tetrazolyl; R^3 is C_1-C_4 alkyl; R^4 is $-NHC(=O)OCH(R^8)$; ring A is a substituted or unsubstituted phenylene, a substituted or unsubstituted naphthylene, or a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring A is substituted then ring A is substituted with

5 1 or 2 R^A ; where the groups  and  are on non-adjacent ring atoms of ring A; ring B is a substituted or unsubstituted monocyclic cycloalkylene, a substituted or unsubstituted bicyclic cycloalkylene, a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted bicyclic heterocycloalkylene, a substituted or 10 unsubstituted phenylene, a substituted or unsubstituted naphthylene, a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring B is substituted then ring B is substituted with 1 or 2 R^B ; CY is a substituted or unsubstituted C_3-C_{10} cycloalkyl, a substituted or unsubstituted phenyl, a substituted or unsubstituted napthyl, a substituted or unsubstituted monocyclic heteroaryl containing 0-4 N atoms, or a substituted or unsubstituted bicyclic heteroaryl containing 0-4 N atoms, wherein if CY is substituted then CY is substituted with 1 or 2 R^C .

15 [0014] In some embodiments, ring A is a monocyclic ring wherein the groups  and  are in a 1,3-relationship on ring A (i.e. an meta relationship).


20 [0015] In some embodiments, ring A is a monocyclic ring wherein the groups  and  are in a 1,4-relationship on ring A (i.e. an para relationship).

[0016] In some embodiments, ring A is a substituted or unsubstituted phenylene, or a substituted or unsubstituted monocyclic heteroarylene containing 0 or 1 O atoms, 0 or 1 S atoms, 0-3 N atoms, and at least 2 carbon atoms in the ring, where if ring A is substituted then ring A is substituted with 1 or 2 R^A ; ring B is a substituted or unsubstituted monocyclic cycloalkylene, a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted phenylene, or a substituted or unsubstituted monocyclic heteroarylene, where if ring B is substituted then ring B is substituted with 1 or 2 R^B ; CY is a substituted or unsubstituted C_3-C_6 cycloalkyl, a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic

heteroaryl containing 0-4 N atoms, wherein if CY is substituted then CY is substituted with 1 or 2 R^C; each of R^A, R^B, and R^C are independently selected from H, halogen, -CN, -NO₂, -OH, -OR¹⁰, -N(R⁹)₂, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

5 [0017] In some embodiments, each of R^A, R^B, and R^C are independently selected from H, halogen, -CN, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[0018] In some embodiments, the compound of Formula (I) has the structure of Formula (II):



10

Formula (II)

wherein,

each of A¹, A², and A³ is independently selected from CR^A and N;


each R^A is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃;

each R^B is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃.

15 [0019] In some embodiments, each of A¹, A², and A³ is CR^A; or one of A¹, A², and A³ is N. In some embodiments, each of A¹, A², and A³ is CR^A. In some embodiments, one of A¹, A², and A³ is N.

[0020] In some embodiments, ring A is a substituted or unsubstituted phen-1,3-ylene, or a substituted or unsubstituted phen-1,4-ylene, where if ring A is substituted then ring A is

20 substituted with R^A. In some embodiments, ring A is a substituted or unsubstituted phen-1,4-ylene, where if ring A is substituted then ring A is substituted with R^A.



[0021] In some embodiments, the group

embodiments, the group

embodiments, the group

25 [0022] In some embodiments, R¹ is -CO₂H, -CO₂R^D, or -C(=O)NHSO₂R¹⁰; R^D is H, -CH₃, or -CH₂CH₃; R⁸ is H, -CH₃, -CH₂CH₃ or -CF₃. In some embodiments, R⁸ is H or -CH₃. In some embodiments, R⁸ is -CH₃.

[0023] In some embodiments, R¹ is -CO₂H; R³ is -CH₃ or -CH₂CH₃; R⁸ is H, or -CH₃.

[0024] In some embodiments, R³ is -CH₃; L¹ is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, or - NR¹¹-C₁-C₄alkylene -; R¹¹ is H or C₁-C₂alkyl; L² is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₆alkylene-O-, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₆alkylene-S-, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₆alkylene-SO-, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₆alkylene-SO₂-, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₄alkylene-NR¹¹-, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, - NR¹¹-C₁-C₄alkylene -, -O-, -S-, -SO-, -SO₂- or -NR²-; R¹¹ is H or C₁-C₂alkyl; CY is a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl containing 1-3 N atoms, wherein if CY is substituted then CY is substituted with 1 or 2 R^C; each R^C is independently selected from H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[0025] In some embodiments, L¹ is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, or - NR¹¹-C₁-C₄alkylene -; R¹¹ is H or C₁-C₂alkyl. In some embodiments, L¹ is absent or C₁-C₆alkylene. In some embodiments, L¹ is absent or C₁-C₄alkylene. In some embodiments, L¹ is C₁-C₆alkylene. In some embodiments, L¹ is absent, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, -CH₂CH(CH₃)-, or -CH₂C(CH₃)₂-. In some embodiments, L¹ is absent, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH(CH₃)-, or -C(CH₃)₂-. In some embodiments, L¹ is -CH₂-.

[0026] In some embodiments, L² is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₆alkylene-O-, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₆alkylene-S-, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₆alkylene-SO-, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₆alkylene-SO₂-, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₄alkylene-NR¹¹-, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, - NR¹¹-C₁-C₄alkylene -, -O-, -S-, -SO-, -SO₂- or -NR²-; R¹¹ is H or C₁-C₂alkyl.


[0027] In some embodiments, L² is absent, C₁-C₄alkylene, C₁-C₄fluoroalkylene, -C₁-C₄alkylene-O-, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₄alkylene -, -C₁-C₄alkylene-S-, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₄alkylene -, -C₁-C₄alkylene-SO-, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₄alkylene -, -C₁-C₄alkylene-SO₂-, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₄alkylene -, -O-, -S-, -SO-, or -SO₂-.

[0028] In some embodiments, L^2 is absent, C_1 - C_4 alkylene, $-C_1$ - C_2 alkylene-O-, -O-C $_1$ -C $_2$ alkylene -, -C $_1$ -C $_2$ alkylene-S-, -S-C $_1$ -C $_2$ alkylene -, -C $_1$ -C $_2$ alkylene-SO-, -SO-C $_1$ -C $_2$ alkylene -, -C $_1$ -C $_2$ alkylene-SO $_2$ -, -SO $_2$ -C $_1$ -C $_2$ alkylene -, -O-, or -S-.

[0029] In some embodiments, L^2 is absent, -CH $_2$ -, -CH $_2$ O-, -OCH $_2$ -, -CH $_2$ S-, -SCH $_2$ -, -CH $_2$ NH-, -NHCH $_2$ -, -O-, -S-, or -NH-.


[0030] In some embodiments, L^2 is absent, -CH $_2$ -, -CH $_2$ O-, -OCH $_2$ -, -CH $_2$ S-, -SCH $_2$ -, -O-, or -S-. In some embodiments, L^2 is absent.

[0031] In some embodiments, L^1 is absent, -CH $_2$ -, -CH $_2$ CH $_2$ -, -CH $_2$ CH $_2$ CH $_2$ -, -CH(CH $_3$)-, -C(CH $_3$) $_2$ -, -CH(CH $_2$ CH $_3$)-, -C(CH $_2$ CH $_3$) $_2$ -, -CH $_2$ CH(CH $_3$)-, or -CH $_2$ C(CH $_3$) $_2$ -, L^2 is absent, -CH $_2$ -, -CH $_2$ O-, -OCH $_2$ -, -CH $_2$ S-, -SCH $_2$ -, -CH $_2$ NH-, -NHCH $_2$ -, -O-, -S-, or -NH-.




[0032] In some embodiments, R^4 is ; ring B is substituted or unsubstituted phenylene, where if ring B is substituted then ring B is substituted with 1 or 2 R^B ; CY is a substituted or unsubstituted phenyl, wherein if CY is substituted then CY is substituted with 1 or 2 R^C . In some embodiments, CY is a substituted or unsubstituted phenyl, wherein if CY is substituted then CY is substituted with R^C .


[0033] In some embodiments, ring B is a substituted or unsubstituted phen-1,2-ylene, a substituted or unsubstituted phen-1,3-ylene, or a substituted or unsubstituted phen-1,4-ylene, where if ring B is substituted then ring B is substituted with R^B . In some embodiments, ring B is a substituted or unsubstituted phen-1,3-ylene, or a substituted or unsubstituted phen-1,4-ylene, where if ring B is substituted then ring B is substituted with R^B . In some embodiments, ring B is a substituted or unsubstituted phen-1,4-ylene, where if ring B is substituted then ring B is substituted with R^B .




[0034] In some embodiments, R^4 is . In some embodiments, R^4 is




[0035] In some embodiments, L^1 is absent, -CH $_2$ -, -CH(CH $_3$)-, -C(CH $_3$) $_2$ -, -CH $_2$ CH $_2$ -, or -CH $_2$ CH $_2$ CH $_2$ -, L^2 is absent; ring A is ; ring B is ; or ; R^8 is -CH $_3$; each R^C is independently selected from H, F, Cl, Br, I, -CH $_3$, -CF $_3$, -OH, -OCF $_3$, and -OCH $_3$.





[0036] In some embodiments, ring A is or . In some embodiments, ring


A is . In some embodiments, ring A is .



[0037] In some embodiments, ring B is or . In some embodiments, ring


B is . In some embodiments, ring B is .

5 [0038] In some embodiments, L¹ is absent or -CH₂-; each R^A is independently selected from H, F, Cl, -CH₃, and -OH; each R^B is independently selected from H, F, Cl, -CH₃, and -OH; each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃.



[0039] In some embodiments, L¹ is -CH₂-; L² is absent; ring B is ; CY is phenyl, 2-fluorophenyl or 2-chloro-phenyl.

10 [0040] In some embodiments, the compound of Formula (I) has the structure of Formula (III):



Formula (III)

wherein

R¹ is -CO₂H, -CO₂R^D, -CN, tetrazolyl, -C(=O)N(R⁹)₂, -C(=O)NHCH₂CH₂SO₃H, or -

15 C(=O)NHSO₂R¹⁰; R^D is H or C₁-C₄alkyl;

L¹ is absent or C₁-C₆alkylene;

R³ is H, C₁-C₄alkyl, C₃-C₆cycloalkyl, or C₁-C₄fluoroalkyl;

R⁷ is H or C₁-C₄alkyl;

R⁸ is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

R⁹ is a H, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₃-C₆cycloalkyl, or a substituted or unsubstituted phenyl;

R¹⁰ is a C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₃-C₆cycloalkyl, or a substituted or unsubstituted phenyl;

5 each of R^A, R^B, and R^C are independently selected from H, F, Cl, Br, I, -CN, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl;

m is 0, 1, or 2; n is 0, 1, or 2; p is 0, 1, or 2.

[0041] In some embodiments, R¹ is -CO₂H, -CO₂R^D or -C(=O)NHSO₂R¹⁰; R³ is C₁-C₄alkyl; R⁷

10 is H; R¹⁰ is a C₁-C₆alkyl or a substituted or unsubstituted phenyl; each R^A is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; each R^B is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; each R^C is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; m is 0 or 1; p is 0 or 1.

[0042] In some embodiments, R¹ is -CO₂H or -CO₂R^D; R³ is -CH₃ or -CH₂CH₃; R⁸ is H, -CH₃


15 or -CF₃; R^D is H, -CH₃, or -CH₂CH₃.

[0043] In some embodiments, R¹ is -CO₂H; R³ is -CH₃; R⁸ is -CH₃; L¹ is absent, -CH₂-, -

CH₂CH₂-, -CH₂CH₂CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, -

CH₂CH(CH₃)-, or -CH₂C(CH₃)₂-.

[0044] In some embodiments, the compound of Formula (III) has the following structure:




20

[0045] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, or -C(CH₂CH₃)₂-, each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; m is 0; n is 0, 1, or 2; p is 0.

[0046] In some embodiments, L¹ is absent or -CH₂-, each R^C is independently selected from H,

25 F, Cl, -CH₃, -CF₃, and -OH; n is 0 or 1.


[0047] In some embodiments, the compound of Formula (III) has the following structure:




[0048] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, or -C(CH₂CH₃)₂-, each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; m is 0; n is 0, 1, or 2; p is 0.

5 [0049] In some embodiments, L¹ is absent or -CH₂-; each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, and -OH; n is 0 or 1.


[0050] In some embodiments, R¹ is -C(=O)NHSO₂R¹⁰, R³ is -CH₃ or -CH₂CH₃; R⁸ is H, -CH₃ or -CF₃; R¹⁰ is -CH₃, or -CH₂CH₃.




[0051] In some embodiments,



10 [0052] In some embodiments,



[0053] In some embodiments,



[0054] In some embodiments,

[0055] Any combination of the groups described above for the various variables is contemplated herein. Throughout the specification, groups and substituents thereof are chosen by one skilled in the field to provide stable moieties and compounds.

[0056] In one aspect, provided is a compound presented in Table 1, Table 2, Table 3, Table 4, Table 5, Table 6, Table 7.

[0057] The compound of Formula (I), Formula (II) or Formula (III) is an antagonist of a LPA receptor. In some embodiments, the compound of Formula (I), Formula (II) or Formula (III) is an antagonist of LPA₁. In some embodiments, the compound of Formula (I), Formula (II) or

Formula (III) is an antagonist of LPA₂. In some embodiments, the compound of Formula (I), Formula (II) or Formula (III) is an antagonist of LPA₃.

[0058] In some embodiments, presented herein are compounds selected from active metabolites, tautomers, pharmaceutically acceptable solvates, pharmaceutically acceptable salts or prodrugs of a compound of Formula (I), Formula (II) or Formula (III).

[0059] In some embodiments, provided is a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III). In some embodiments, the pharmaceutical composition also contains at least one pharmaceutically acceptable inactive ingredient.

[0060] In some embodiments, provided is a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable inactive ingredient. In one aspect, the pharmaceutical composition is formulated for intravenous injection, subcutaneous injection, oral administration, inhalation, nasal administration, topical administration, ophthalmic administration or otic administration. In some embodiments, the pharmaceutical composition is a tablet, a pill, a capsule, a liquid, an inhalant, a nasal spray solution, a suppository, a suspension, a gel, a colloid, a dispersion, a suspension, a solution, an emulsion, an ointment, a lotion, an eye drop or an ear drop.

[0061] In some embodiments, the pharmaceutical composition further comprises one or more additional therapeutically active agents selected from: corticosteroids, immunosuppressants, analgesics, anti-cancer agent, anti-inflammatories, chemokine receptor antagonists, bronchodilators, leukotriene receptor antagonists, leukotriene formation inhibitors, monoacylglycerol kinase inhibitors, phospholipase A₁ inhibitors, phospholipase A₂ inhibitors, and lysophospholipase D (lysoPLD) inhibitors, autotaxin inhibitors, decongestants, antihistamines, mucolytics, anticholinergics, antitussives, expectorants, and β-2 agonists.

[0062] In some embodiments, provided is a method comprising administering a compound of Formula (I), Formula (II) or Formula (III) to a human with a LPA-dependent or LPA-mediated disease or condition. In some embodiments, the human is already being administered one or more additional therapeutically active agents other than a compound of Formula (I), Formula (II) or Formula (III). In some embodiments, the method further comprises administering one or more additional therapeutically active agents other than a compound of Formula (I), Formula (II) or Formula (III).

[0063] In some embodiments, the one or more additional therapeutically active agents other than a compound of Formula (I), Formula (II) or Formula (III) are selected from: corticosteroids, immunosuppressants, analgesics, anti-cancer agent, anti-inflammatories, chemokine receptor

antagonists, bronchodilators, leukotriene receptor antagonists, leukotriene formation inhibitors, monoacylglycerol kinase inhibitors, phospholipase A₁ inhibitors, phospholipase A₂ inhibitors, and lysophospholipase D (lysoPLD) inhibitors, autotaxin inhibitors, decongestants, antihistamines, mucolytics, anticholinergics, antitussives, expectorants, and β-2 agonists.

5 [0064] Pharmaceutical formulations described herein are administerable to a subject in a variety of ways by multiple administration routes, including but not limited to, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical or transdermal administration routes. The pharmaceutical formulations described herein include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal dispersions, aerosols, solid dosage forms, powders, immediate release formulations, controlled 10 release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate and controlled release formulations.

15 [0065] In some embodiments, the compounds of Formula (I), Formula (II) and Formula (III) are administered orally.

20 [0066] In some embodiments, the compounds of Formula (I), Formula (II) and Formula (III) are administered topically. In such embodiments, the compound of Formula (I), Formula (II) or Formula (III) is formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, shampoos, scrubs, rubs, smears, medicated sticks, medicated bandages, balms, creams or ointments. Such pharmaceutical compounds can contain 25 solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives. In one aspect, the compounds of Formula (I), Formula (II) and Formula (III) are administered topically to the skin.

[0067] In another aspect, the compounds of Formula (I), Formula (II) and Formula (III) are administered by inhalation. In one embodiment, the compounds of Formula (I), Formula (II) and Formula (III) are administered by inhalation that directly targets the pulmonary system.

[0068] In another aspect, the compounds of Formula (I), Formula (II) and Formula (III) are formulated for intranasal administration. Such formulations include nasal sprays, nasal mists, and the like.

30 [0069] In another aspect, the compounds of Formula (I), Formula (II) and Formula (III) are formulated as eye drops.

[0070] In another aspect is the use of a compound of Formula (I), Formula (II) or Formula (III) in the manufacture of a medicament for treating a disease, disorder or conditions in which the activity of at least one LPA receptor contributes to the pathology and/or symptoms of the disease or condition. In one embodiment of this aspect, the LPA is selected from LPA₁, LPA₂, LPA₃,

LPA₄, LPA₅ and LPA₆. In one aspect, the LPA receptor is LPA₁. In one aspect, the disease or condition is any of the diseases or conditions specified herein.

[0071] In any of the aforementioned aspects are further embodiments in which: (a) the effective amount of the compound of Formula (I), Formula (II) or Formula (III) is systemically

5 administered to the mammal; and/or (b) the effective amount of the compound is administered orally to the mammal; and/or (c) the effective amount of the compound is intravenously administered to the mammal; and/or (d) the effective amount of the compound is administered by inhalation; and/or (e) the effective amount of the compound is administered by nasal administration; or and/or (f) the effective amount of the compound is administered by injection to 10 the mammal; and/or (g) the effective amount of the compound is administered topically to the mammal; and/or (h) the effective amount of the compound is administered by ophthalmic administration; and/or (i) the effective amount of the compound is administered rectally to the mammal; and/or (j) the effective amount is administered non-systemically or locally to the mammal.

15 **[0072]** In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in which (i) the compound is administered once; (ii) the compound is administered to the mammal multiple times over the span of one day; (iii) continually; or (iv) continuously.

[0073] In any of the aforementioned aspects are further embodiments comprising multiple 20 administrations of the effective amount of the compound, including further embodiments in which (i) the compound is administered continuously or intermittently: as in a single dose; (ii) the time between multiple administrations is every 6 hours; (iii) the compound is administered to the mammal every 8 hours; (iv) the compound is administered to the mammal every 12 hours; (v) the compound is administered to the mammal every 24 hours. In further or alternative 25 embodiments, the method comprises a drug holiday, wherein the administration of the compound is temporarily suspended or the dose of the compound being administered is temporarily reduced; at the end of the drug holiday, dosing of the compound is resumed. In one embodiment, the length of the drug holiday varies from 2 days to 1 year.

[0074] Also provided is a method of inhibiting the physiological activity of LPA in a mammal 30 comprising administering a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III) or a pharmaceutically acceptable salt thereof to the mammal in need thereof.

[0075] In one aspect, provided is a medicament for treating a LPA-dependent or LPA-mediated disease or condition in a mammal comprising a therapeutically effective amount of a compound 35 of Formula (I), Formula (II) or Formula (III).

[0076] In some cases disclosed herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the manufacture of a medicament for the treatment of a LPA-dependent or LPA-mediated disease or condition.

5 [0077] In some cases disclosed herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the treatment or prevention of a LPA-dependent or LPA-mediated disease or condition.

[0078] In one aspect, is a method for treating or preventing a LPA-dependent or LPA-mediated disease or condition in a mammal comprising administering a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III).

10 [0079] In one aspect, LPA-dependent or LPA-mediated diseases or conditions include, but are not limited to, fibrosis of organs or tissues, scarring, liver diseases, dermatological conditions, cancer, cardiovascular disease, respiratory diseases or conditions, inflammatory disease, gastrointestinal tract disease, renal disease, urinary tract-associated disease, inflammatory disease of lower urinary tract, dysuria, frequent urination, pancreas disease, arterial obstruction, cerebral infarction, cerebral hemorrhage, pain, peripheral neuropathy, and fibromyalgia.

15 [0080] In one aspect, the LPA-dependent or LPA-mediated disease or condition is a respiratory disease or condition. In some embodiments, the respiratory disease or condition is asthma, chronic obstructive pulmonary disease (COPD), pulmonary fibrosis, pulmonary arterial hypertension or acute respiratory distress syndrome.

20 [0081] In some embodiments, the LPA-dependent or LPA-mediated disease or condition is selected from idiopathic pulmonary fibrosis; other diffuse parenchymal lung diseases of different etiologies including iatrogenic drug-induced fibrosis, occupational and/or environmental induced fibrosis, granulomatous diseases (sarcoidosis, hypersensitivity pneumonia), collagen vascular disease, alveolar proteinosis, langerhans cell granulomatosis, lymphangioleiomyomatosis, 25 inherited diseases (Hermansky-Pudlak Syndrome, tuberous sclerosis, neurofibromatosis, metabolic storage disorders, familial interstitial lung disease); radiation induced fibrosis; chronic obstructive pulmonary disease (COPD); scleroderma; bleomycin induced pulmonary fibrosis; chronic asthma; silicosis; asbestos induced pulmonary fibrosis; acute respiratory distress syndrome (ARDS); kidney fibrosis; tubulointerstitium fibrosis; glomerular nephritis; focal segmental glomerular sclerosis; IgA nephropathy; hypertension; Alport; gut fibrosis; liver 30 fibrosis; cirrhosis; alcohol induced liver fibrosis; toxic/drug induced liver fibrosis; hemochromatosis; nonalcoholic steatohepatitis (NASH); biliary duct injury; primary biliary cirrhosis; infection induced liver fibrosis; viral induced liver fibrosis; and autoimmune hepatitis; corneal scarring; hypertrophic scarring; Duputren disease, keloids, cutaneous fibrosis; cutaneous scleroderma; spinal cord injury/fibrosis; myelofibrosis; vascular restenosis; atherosclerosis;

arteriosclerosis; Wegener's granulomatosis; Peyronie's disease, chronic lymphocytic leukemia, tumor metastasis, transplant organ rejection, endometreosis, neonatal respiratory distress syndrome and neuropathic pain.

[0082] In one aspect, the LPA-dependent or LPA-mediated disease or condition is described
5 herein.

[0083] In one aspect, provided is a method for the treatment or prevention of organ fibrosis in a mammal comprising administering a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III) or a pharmaceutically acceptable salt thereof to a mammal in need thereof.

10 [0084] In one aspect, the organ fibrosis comprises lung fibrosis, renal fibrosis, or hepatic fibrosis.

[0085] In one aspect, provided is a method of improving lung function in a mammal comprising administering a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III) or a pharmaceutically acceptable salt thereof to the mammal in need thereof. In one aspect, the mammal has been diagnosed as having lung fibrosis.

15 [0086] In one aspect, compounds disclosed herein are used to treat idiopathic pulmonary fibrosis (usual interstitial pneumonia) in a mammal.

[0087] In some embodiments, compounds disclosed herein are used to treat diffuse parenchymal interstitial lung diseases in mammal: iatrogenic drug induced, occupational/environmental (Farmer lung), granulomatous diseases (sarcoidosis, hypersensitivity pneumonia), collagen 20 vascular disease (scleroderma and others), alveolar proteinosis, langerhans cell granulomatosis, lymphangioleiomyomatosis, Hermansky-Pudlak Syndrome, Tuberous sclerosis, neurofibromatosis, metabolic storage disorders, familial interstitial lung disease.

[0088] In some embodiments, compounds disclosed herein are used to treat post-transplant fibrosis associated with chronic rejection in a mammal: Bronchiolitis obliterans for lung transplant.

25 [0089] In some embodiments, compounds disclosed herein are used to treat cutaneous fibrosis in a mammal: cutaneous scleroderma, Dupuytren disease, keloids.

[0090] In one aspect, compounds disclosed herein are used to treat hepatic fibrosis with or without cirrhosis in a mammal: toxic/drug induced (hemochromatosis), alcoholic liver disease, 30 viral hepatitis (hepatitis B virus, hepatitis C virus, HCV), nonalcoholic liver disease (NASH), metabolic and auto-immune.

[0091] In one aspect, compounds disclosed herein are used to treat renal fibrosis in a mammal: tubulointerstitium fibrosis, glomerular sclerosis.

[0092] In any of the aforementioned aspects involving the treatment of LPA dependent diseases 35 or conditions are further embodiments comprising administering at least one additional agent in

addition to the administration of a compound having the structure of Formula (I), Formula (II) or Formula (III). In various embodiments, each agent is administered in any order, including simultaneously.

[0093] In any of the embodiments disclosed herein, the mammal is a human.

5 [0094] In some embodiments, compounds provided herein are administered to a human.

[0095] In some embodiments, compounds provided herein are orally administered.

[0096] In some embodiments, compounds provided herein are used as antagonists of at least one LPA receptor. In some embodiments, compounds provided herein are used for inhibiting the activity of at least one LPA receptor or for the treatment of a disease or condition that would 10 benefit from inhibition of the activity of at least one LPA receptor. In one aspect, the LPA receptor is LPA₁.

[0097] In other embodiments, compounds provided herein are used for the formulation of a medicament for the inhibition of LPA₁ activity.

[0098] Articles of manufacture, which include packaging material, a compound of Formula (I),

15 Formula (II) or Formula (III), within the packaging material, and a label that indicates that the compound or composition, or pharmaceutically acceptable salt, tautomers, pharmaceutically acceptable N-oxide, pharmaceutically active metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, is used for inhibiting the activity of at least one LPA receptor, or for the treatment, prevention or amelioration of one or more symptoms of a disease 20 or condition that would benefit from inhibition of the activity of at least one LPA receptor, are provided.

[0099] Other objects, features and advantages of the compounds, methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating 25 specific embodiments, are given by way of illustration only, since various changes and modifications within the spirit and scope of the instant disclosure will become apparent to those skilled in the art from this detailed description

DETAILED DESCRIPTION OF THE INVENTION

30 [00100] Lysophospholipids are membrane-derived bioactive lipid mediators. Lysophospholipids include, but are not limited to, lysophosphatidic acid (1-acyl-2-hydroxy-*sn*-glycero-3-phosphate; LPA), sphingosine 1-phosphate (S1P), lysophosphatidylcholine (LPC), and sphingosylphosphorylcholine (SPC). Lysophospholipids affect fundamental cellular functions that include cellular proliferation, differentiation, survival, migration, adhesion, invasion, and

morphogenesis. These functions influence many biological processes that include neurogenesis, angiogenesis, wound healing, immunity, and carcinogenesis.

[00101] LPA acts through sets of specific G protein-coupled receptors (GPCRs) in an autocrine and paracrine fashion. LPA binding to its cognate GPCRs (LPA₁, LPA₂, LPA₃, LPA₄, LPA₅,

5 LPA₆) activates intracellular signaling pathways to produce a variety of biological responses.

[00102] Lysophospholipids, such as LPA, are quantitatively minor lipid species compared to their major phospholipid counterparts (e.g., phosphatidylcholine, phosphatidylethanolamine, and sphingomyelin). LPA has a role as a biological effector molecule, and has a diverse range of physiological actions such as, but not limited to, effects on blood pressure, platelet activation, 10 and smooth muscle contraction, and a variety of cellular effects, which include cell growth, cell rounding, neurite retraction, and actin stress fiber formation and cell migration. The effects of LPA are predominantly receptor mediated.

[00103] Activation of the LPA receptors (LPA₁, LPA₂, LPA₃, LPA₄, LPA₅, LPA₆) with LPA

mediates a range of downstream signaling cascades. These include, but are not limited to,

15 mitogen-activated protein kinase (MAPK) activation, adenylyl cyclase (AC)

inhibition/activation, phospholipase C (PLC) activation/Ca²⁺ mobilization, arachidonic acid release, Akt/PKB activation, and the activation of small GTPases, Rho, ROCK, Rac, and Ras.

Other pathways that are affected by LPA receptor activation include, but are not limited to, cyclic 20 adenosine monophosphate (cAMP), cell division cycle 42/GTP-binding protein (Cdc42) , proto-

oncogene serine/threonine-protein kinase Raf (c-RAF), proto-oncogene tyrosine-protein kinase Src (c-src), extracellular signal-regulated kinase (ERK), focal adhesion kinase (FAK), guanine

nucleotide exchange factor (GEF), glycogen synthase kinase 3b (GSK3b), c-jun amino-terminal kinase (JNK), MEK, myosin light chain II (MLC II), nuclear factor kB (NF-kB), N-methyl-D-aspartate (NMDA) receptor activation, phosphatidylinositol 3-kinase (PI3K), protein kinase A

25 (PKA), protein kinase C (PKC), ras-related C3 botulinum toxin substrate 1 (RAC1). The actual

pathway and realized end point are dependent on a range of variables that include receptor usage, cell type, expression level of a receptor or signaling protein, and LPA concentration. Nearly all

mammalian cells, tissues and organs co-express several LPA-receptor subtypes, which indicates 30 that LPA receptors signal in a cooperative manner. LPA₁, LPA₂, and LPA₃ share high amino acid sequence similarity.

[00104] LPA is produced from activated platelets, activated adipocytes, neuronal cells, and other cell types. Serum LPA is produced by multiple enzymatic pathways that involve monoacylglycerol kinase, phospholipase A₁, secretory phospholipase A₂, and lysophospholipase D (lysoPLD), including autotaxin. Several enzymes are involved in LPA degradation:

35 lysophospholipase, lipid phosphate phosphatase, and LPA acyl transferase such as endophilin.

LPA concentrations in human serum are estimated to be 1–5 μ M. Serum LPA is bound to albumin, low-density lipoproteins, or other proteins, which possibly protect LPA from rapid degradation. LPA molecular species with different acyl chain lengths and saturation are naturally occurring, including 1-palmitoyl (16:0), 1-palmitoleoyl (16:1), 1-stearoyl (18:0), 1-oleoyl (18:1), 5 1-linoleoyl (18:2), and 1-arachidonyl (20:4) LPA. Quantitatively minor alkyl LPA has biological activities similar to acyl LPA, and different LPA species activate LPA receptor subtypes with varied efficacies.

LPA RECEPTORS

[00105] LPA₁ (previously called VZG-1/EDG-2/mrec1.3) couples with three types of G proteins,

10 G_{i/o}, G_q, and G_{12/13}. Through activation of these G proteins, LPA induces a range of cellular responses through LPA₁ including but not limited to: cell proliferation, serum-response element (SRE) activation, mitogen-activated protein kinase (MAPK) activation, adenylyl cyclase (AC) inhibition, phospholipase C (PLC) activation, Ca²⁺ mobilization, Akt activation, and Rho activation.

15 [00106] Wide expression of LPA₁ is observed in adult mice, with clear presence in testis, brain, heart, lung, small intestine, stomach, spleen, thymus, and skeletal muscle. Similarly, human tissues also express LPA₁; it is present in brain, heart, lung, placenta, colon, small intestine, prostate, testis, ovary, pancreas, spleen, kidney, skeletal muscle, and thymus.

[00107] LPA₂ (EDG-4) also couples with three types of G proteins, G_{i/o}, G_q, and G_{12/13}, to mediate

20 LPA-induced cellular signaling. Expression of LPA₂ is observed in the testis, kidney, lung, thymus, spleen, and stomach of adult mice and in the human testis, pancreas, prostate, thymus, spleen, and peripheral blood leukocytes. Expression of LPA₂ is upregulated in various cancer cell lines, and several human LPA₂ transcriptional variants with mutations in the 3'-untranslated region have been observed. Targeted deletion of LPA₂ in mice has not shown any obvious 25 phenotypic abnormalities, but has demonstrated a significant loss of normal LPA signaling (e.g., PLC activation, Ca²⁺ mobilization, and stress fiber formation) in primary cultures of mouse embryonic fibroblasts (MEFs). Creation of *lpa1*(-/-) *lpa2* (-/-) double-null mice has revealed that many LPA-induced responses, which include cell proliferation, AC inhibition, PLC activation, Ca²⁺ mobilization, JNK and Akt activation, and stress fiber formation, are absent or severely 30 reduced in double-null MEFs. All these responses, except for AC inhibition (AC inhibition is nearly abolished in LPA₁ (-/-) MEFs), are only partially affected in either LPA₁ (-/-) or LPA₂ (-/-) MEFs. LPA₂ contributes to normal LPA-mediated signaling responses in at least some cell types (Choi *et al*, *Biochimica et Biophysica Acta* 2008, 1781, p531-539).

[00108] LPA₃ (EDG-7) is distinct from LPA₁ and LPA₂ in its ability to couple with G_{i/o} and G_q

35 but not G_{12/13} and is much less responsive to LPA species with saturated acyl chains. LPA₃ can

mediate pleiotropic LPA-induced signaling that includes PLC activation, Ca^{2+} mobilization, AC inhibition/activation, and MAPK activation. Overexpression of LPA₃ in neuroblastoma cells leads to neurite elongation, whereas that of LPA₁ or LPA₂ results in neurite retraction and cell rounding when stimulated with LPA. Expression of LPA₃ is observed in adult mouse testis,

5 kidney, lung, small intestine, heart, thymus, and brain. In humans, it is found in the heart, pancreas, prostate, testis, lung, ovary, and brain (frontal cortex, hippocampus, and amygdala).

[00109] LPA₄ (p2y₉/GPR23) is of divergent sequence compared to LPA₁, LPA₂, and LPA₃ with closer similarity to the platelet-activating factor (PAF) receptor. LPA₄ mediates LPA induced Ca^{2+} mobilization and cAMP accumulation, and functional coupling to the G protein Gs for AC

10 activation, as well as coupling to other G proteins. The LPA₄ gene is expressed in the ovary, pancreas, thymus, kidney and skeletal muscle.

[00110] LPA₅ (GPR92) is a member of the purinocluster of GPCRs and is structurally most closely related to LPA₄. LPA₅ is expressed in human heart, placenta, spleen, brain, lung and gut. LPA₅ also shows very high expression in the CD8+ lymphocyte compartment of the

15 gastrointestinal tract.

[00111] LPA₆ (p2y5) is a member of the purinocluster of GPCRs and is structurally most closely related to LPA₄. LPA₆ is an LPA receptor coupled to the G12/13-Rho signaling pathways and is expressed in the inner root sheaths of human hair follicles.

Illustrative Biological Activity

20 Wound Healing

[00112] Normal wound healing occurs by a highly coordinated sequence of events in which cellular, soluble factors and matrix components act in concert to repair the injury. The healing response can be described as taking place in four broad, overlapping phases—hemostasis, inflammation, proliferation, and remodeling. Many growth factors and cytokines are released into

25 a wound site to initiate and perpetuate wound healing processes.

[00113] When wounded, damaged blood vessels activate platelets. The activated platelets play pivotal roles in subsequent repair processes by releasing bioactive mediators to induce cell proliferation, cell migration, blood coagulation, and angiogenesis. LPA is one such mediator that is released from activated platelets; this induces platelet aggregation along with

30 mitogenic/migration effects on the surrounding cells, such as endothelial cells, smooth muscle cells, fibroblasts, and keratinocytes.

[00114] Topical application of LPA to cutaneous wounds in mice promotes repair processes (wound closure and increased neoepithelial thickness) by increasing cell proliferation/ migration without affecting secondary inflammation.

[00115] Activation of dermal fibroblasts by growth factors and cytokines leads to their subsequent migration from the edges of the wound into the provisional matrix formed by the fibrin clot whereupon the fibroblasts proliferate and start to restore the dermis by secreting and organizing the characteristic dermal extracellular matrix (ECM). The increasing number of fibroblasts within

5 the wound and continuous precipitation of ECM enhances matrix rigidity by applying small tractional forces to the newly formed granulation tissue. The increase in mechanical stress, in conjunction with transforming growth factor β (TGF β), induces α -smooth muscle actin (α -SMA) expression and the subsequent transformation of fibroblasts into myofibroblasts. Myofibroblasts facilitate granulation tissue remodeling via myofibroblast contraction and through the production

10 of ECM components.

[00116] LPA regulates many important functions of fibroblasts in wound healing, including proliferation, migration, differentiation and contraction. Fibroblast proliferation is required in wound healing in order to fill an open wound. In contrast, fibrosis is characterized by intense proliferation and accumulation of myofibroblasts that actively synthesize ECM and

15 proinflammatory cytokines. LPA can either increase or suppress the proliferation of cell types important in wound healing, such as epithelial and endothelial cells (EC), macrophages, keratinocytes, and fibroblasts. A role for LPA₁ in LPA-induced proliferation was provided by the observation that LPA-stimulated proliferation of fibroblasts isolated from LPA₁ receptor null mice was attenuated (Mills *et al*, *Nat Rev. Cancer* 2003; 3: 582-591). LPA induces cytoskeletal changes that are integral to fibroblast adhesion, migration, differentiation and contraction.

Fibrosis

[00117] Tissue injury initiates a complex series of host wound-healing responses; if successful, these responses restore normal tissue structure and function. If not, these responses can lead to tissue fibrosis and loss of function.

25 [00118] For the majority of organs and tissues the development of fibrosis involves a multitude of events and factors. Molecules involved in the development of fibrosis include proteins or peptides (profibrotic cytokines, chemokines, metalloproteinases etc.) and phospholipids. Phospholipids involved in the development of fibrosis include platelet activating factor (PAF), phosphatidyl choline, sphingosine-1 phosphate (S1P) and lysophosphatidic acid (LPA).

30 [00119] A number of muscular dystrophies are characterized by a progressive weakness and wasting of musculature, and by extensive fibrosis. It has been shown that LPA treatment of cultured myoblasts induced significant expression of connective tissue growth factor (CTGF). CTGF subsequently induces collagen, fibronectin and integrin expression and induces dedifferentiation of these myoblasts. Treatment of a variety of cell types with LPA induces reproducible and high level induction of CTGF (J.P. Pradere, *et al.*, LPA₁ receptor activation

promotes renal interstitial fibrosis, *J. Am. Soc. Nephrol.* 18 (2007) 3110–3118; N. Wiedmaier, *et al.*, *Int J Med Microbiol*; 298(3-4):231-43, 2008). CTGF is a profibrotic cytokine, signaling down-stream and in parallel with TGF β .

5 [00120] CTGF expression by gingival epithelial cells, which are involved in the development of gingival fibromatosis, was found to be exacerbated by LPA treatment (A. Kantarci, *et al.*, *J. Pathol.* 210 (2006) 59–66).

10 [00121] LPA is associated with the progression of liver fibrosis. *In vitro*, LPA induces stellate cell and hepatocyte proliferation. These activated cells are the main cell type responsible for the accumulation of ECM in the liver. Furthermore, LPA plasma levels rise during CCl₄-induced liver fibrosis in rodents, or in hepatitis C virus-induced liver fibrosis in human (N. Watanabe, *et al.*, Plasma lysophosphatidic acid level and serum autotaxin activity are increased in liver injury in rats in relation to its severity, *Life Sci.* 81 (2007) 1009–1015; N. Watanabe, *et al.*, *J. Clin. Gastroenterol.* 41 (2007) 616–623).

15 [00122] An increase of phospholipid concentrations in the bronchoalveolar lavage fluid in rabbits and rodents injected with bleomycin has been reported (K. Kuroda, *et al.*, Phospholipid concentration in lung lavage fluid as biomarker for pulmonary fibrosis, *Inhal. Toxicol.* 18 (2006) 389–393; K. Yasuda, *et al.*, *Lung* 172 (1994) 91–102).

20 [00123] LPA is associated with heart disease and myocardial remodeling. Serum LPA levels are increased after myocardial infarction in patients and LPA stimulates rat cardiac fibroblast proliferation and collagen production (*Chen et al. FEBS Lett.* 2006 Aug 21;580(19):4737-45).

Pulmonary Fibrosis

[00124] In the lung, aberrant wound healing responses to injury contribute to the pathogenesis of fibrotic lung diseases. Fibrotic lung diseases, such as idiopathic pulmonary fibrosis (IPF), are associated with high morbidity and mortality.

25 [00125] LPA is an important mediator of fibroblast recruitment in pulmonary fibrosis. LPA and LPA₁ play key pathogenic roles in pulmonary fibrosis. Fibroblast chemoattractant activity plays an important role in the lungs in patients with pulmonary fibrosis. Profibrotic effects of LPA₁-receptor stimulation is explained by LPA₁-receptor-mediated vascular leakage and increased fibroblast recruitment, both profibrotic events. The LPA-LPA₁ pathway has a role in mediating 30 fibroblast migration and vascular leakage in IPF. The end result is the aberrant healing process that characterises this fibrotic condition.

[00126] The LPA₁ receptor is the LPA receptor most highly expressed on fibroblasts obtained from patients with IPF. Furthermore, BAL obtained from IPF patients induced chemotaxis of human foetal lung fibroblasts that was blocked by the dual LPA₁- LPA₃ receptor antagonist Ki16425. In an experimental bleomycin-induced lung injury mouse model, it was shown that

LPA levels were high in bronchoalveolar lavage samples compared with unexposed controls. LPA₁ knockout mice are protected from fibrosis after bleomycin challenge with reduced fibroblast accumulation and vascular leakage. In human subjects with IPF, high LPA levels were observed in bronchoalveolar lavage samples compared with healthy controls. Increased fibroblast 5 chemotactic activity in these samples was inhibited by the Ki16425 indicating that fibroblast migration is mediated by the LPA-LPA receptor(s) pathway (Tager *et al. Nature Medicine*, vol. 14, no. 1, 45-54, 2008).

[00127] The LPA-LPA₁ pathway is crucial in fibroblast recruitment and vascular leakage in pulmonary fibrosis.

10 [00128] Activation of latent TGF- β by the av β 6 integrin plays a critical role in the development of lung injury and fibrosis (Munger *et al. Cell*, vol. 96, 319-328, 1999). LPA induces av β 6-mediated TGF- β activation on human lung epithelial cells (Xu *et al. Am. J. Pathology*, vol. 174, no. 2, 1264-1279, 2009). The LPA-induced av β 6-mediated TGF- β activation is mediated by the LPA2 receptor. Expression of the LPA2 receptor is increased in epithelial cells and mesenchymal 15 cells in areas of lung fibrosis from IPF patients compared to normal human lung tissue. The LPA-LPA2 pathway contributes to the activation of the TGF- β pathway in pulmonary fibrosis. In some embodiments, compounds that inhibit LPA2 show efficacy in the treatment of lung fibrosis. In some embodiments, compounds that inhibit both LPA1 and LPA2 show improved efficacy in the treatment of lung fibrosis compared to compounds which inhibit only LPA1 or LPA2.

20 Renal Fibrosis

[00129] LPA and LPA₁ are involved in the etiology of kidney fibrosis. LPA has effects on both proliferation and contraction of glomerular mesangial cells and thus has been implicated in proliferative glomerulonephritis (C.N. Inoue, *et al., Clin. Sci. (Colch.)* 96, 431– 436, 1999). In an animal model of renal fibrosis, unilateral ureteral obstruction (UUO), it was found that renal LPA 25 receptors are expressed under basal conditions with an expression order of LPA₂>LPA₃=LPA₁>>LPA₄. This model mimics in an accelerated manner the development of renal fibrosis including renal inflammation, fibroblast activation and accumulation of extracellular matrix in the tubulointerstitium. UUO significantly induced LPA₁-receptor expression. This was paralleled by renal LPA production (3.3 fold increase) in conditioned media 30 from kidney explants. Contra-lateral kidneys exhibited no significant changes in LPA release and LPA-receptors expression. This shows that a prerequisite for an action of LPA in fibrosis is met: production of a ligand (LPA) and induction of one of its receptors (the LPA₁ receptor) (J.P. Pradere *et al., Biochim Biophys Acta*. 2008 Sep;1781(9):582-7).

[00130] In mice invalidated for the LPA₁ receptor (LPA₁ (-/-)), the development of renal fibrosis was significantly attenuated. UUO mice treated with the LPA receptor antagonist Ki16425 closely resembled the LPA₁ (-/-) mice.

5 [00131] LPA can participate in intraperitoneal accumulation of monocyte/macrophages and that LPA can induce expression of the profibrotic cytokine CTGF in primary cultures of human fibroblasts (J.S. Koh,*et al.*, *J. Clin. Invest.* 102 (1998) 716–727).

10 [00132] LPA treatment of a mouse epithelial renal cell line, MCT, induced a rapid increase in the expression of the profibrotic cytokine CTGF. CTGF plays a crucial role in UUO-induced tubulointerstitial fibrosis (TIF), and is involved in the profibrotic activity of TGF β . This induction was almost completely suppressed by co-treatment with the LPA-receptor antagonist Ki16425. In one aspect, the profibrotic activity of LPA in kidney results from a direct action of LPA on kidney cells involving induction of CTGF.

Hepatic fibrosis

15 [00133] LPA is implicated in liver disease and fibrosis. Plasma LPA levels and serum autotoxin (enzyme responsible for LPA production) are elevated in hepatitis patients and animal models of liver injury in correlation with increased fibrosis. LPA also regulates liver cell function. LPA₁ and LPA₂ receptors are expressed by mouse hepatic stellate cells and LPA stimulates migration of hepatic myofibroblasts.

Ocular Fibrosis

20 [00134] LPA is involved in wound healing in the eye. LPA₁ and LPA₃ receptors are detectable in the normal rabbit corneal epithelial cells, keratocytes and endothelial cells and LPA₁ and LPA₃ expression are increased in corneal epithelial cells following injury.

[00135] LPA and its homologues are present in the aqueous humor and the lacrimal gland fluid of the rabbit eye and these levels are increased in a rabbit corneal injury model.

25 [00136] LPA induces actin stress fiber formation in rabbit corneal endothelial and epithelial cells and promotes contraction corneal fibroblasts. LPA also stimulates proliferation of human retinal pigmented epithelial cells

Cardiac fibrosis

30 [00137] LPA is implicated in myocardial infarction and cardiac fibrosis. Serum LPA levels are increased in patients following myocardial infarction (MI) and LPA stimulates proliferation and collagen production (fibrosis) by rat cardiac fibroblasts. Both LPA1 and LPA3 receptors are highly expressed in human heart tissue.

Treatment of Fibrosis

35 [00138] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are used to treat or prevent fibrosis in a mammal. In one aspect, compounds of Formula (I), Formula (II) and

Formula (III) are used to treat fibrosis of an organ or tissue in a mammal. In one aspect is a method for preventing a fibrosis condition in a mammal, the method comprising administering to the mammal at risk of developing one or more fibrosis conditions a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III). In one aspect, the mammal

5 has been exposed to one or more environmental conditions that are known to increase the risk of fibrosis of an organ or tissue. In one aspect, the mammal has been exposed to one or more environmental conditions that are known to increase the risk of lung, liver or kidney fibrosis. In one aspect, the mammal has a genetic predisposition of developing fibrosis of an organ or tissue. In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is administered to a

10 mammal to prevent or minimize scarring following injury. In one aspect, injury includes surgery.

[00139] The terms “fibrosis” or “fibrosing disorder,” as used herein, refers to conditions that are associated with the abnormal accumulation of cells and/or fibronectin and/or collagen and/or increased fibroblast recruitment and include but are not limited to fibrosis of individual organs or tissues such as the heart, kidney, liver, joints, lung, pleural tissue, peritoneal tissue, skin, cornea, retina, musculoskeletal and digestive tract.

[00140] Exemplary diseases, disorders, or conditions that involve fibrosis include, but are not limited to: Lung diseases associated with fibrosis, e.g., idiopathic pulmonary fibrosis, pulmonary fibrosis secondary to systemic inflammatory disease such as rheumatoid arthritis, scleroderma, lupus, cryptogenic fibrosing alveolitis, radiation induced fibrosis, chronic obstructive pulmonary

20 disease (COPD), scleroderma, chronic asthma, silicosis, asbestos induced pulmonary or pleural fibrosis, acute lung injury and acute respiratory distress (including bacterial pneumonia induced, trauma induced, viral pneumonia induced, ventilator induced, non-pulmonary sepsis induced, and aspiration induced); Chronic nephropathies associated with injury/fibrosis (kidney fibrosis), e.g., glomerulonephritis secondary to systemic inflammatory diseases such as lupus and scleroderma,

25 diabetes, glomerular nephritis, focal segmental glomerular sclerosis, IgA nephropathy, hypertension, allograft and Alport; Gut fibrosis, e.g., scleroderma, and radiation induced gut fibrosis; Liver fibrosis, e.g., cirrhosis, alcohol induced liver fibrosis, nonalcoholic steatohepatitis (NASH), biliary duct injury, primary biliary cirrhosis, infection or viral induced liver fibrosis (e.g., chronic HCV infection), and autoimmune hepatitis; Head and neck fibrosis, e.g., radiation

30 induced; Corneal scarring, e.g., LASIK (laser-assisted in situ keratomileusis), corneal transplant, and trabeculectomy; Hypertrophic scarring and keloids, e.g., burn induced or surgical; and Other fibrotic diseases, e.g., sarcoidosis, scleroderma, spinal cord injury/fibrosis, myelofibrosis, vascular restenosis, atherosclerosis, arteriosclerosis, Wegener's granulomatosis, mixed connective tissue disease, and Peyronie's disease.

[00141] In one aspect, a mammal suffering from one of the following non-limiting exemplary diseases, disorders, or conditions will benefit from therapy with a compound of Formula (I), Formula (II) or Formula (III): atherosclerosis, thrombosis, heart disease, vasculitis, formation of scar tissue, restenosis, phlobitis, COPD (chronic obstructive pulmonary disease), pulmonary hypertension, pulmonary fibrosis, pulmonary inflammation, bowel adhesions, bladder fibrosis and cystitis, fibrosis of the nasal passages, sinusitis, inflammation mediated by neutrophils, and fibrosis mediated by fibroblasts.

[00142] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is administered to a mammal with fibrosis of an organ or tissue or with a predisposition of developing fibrosis of an organ or tissue with one or more other agents that are used to treat fibrosis. In one aspect, the one or more agents include corticosteroids. In one aspect, the one or more agents include immunosuppresants. In one aspect, the one or more agents include B-cell antagonists. In one aspect, the one or more agents include uteroglobin.

[00143] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used to treat a dermatological disorders in a mammal. The term “dermatological disorder,” as used herein refers to a skin disorder. Such dermatological disorders include, but are not limited to, proliferative or inflammatory disorders of the skin such as, atopic dermatitis, bullous disorders, collagenoses, psoriasis, psoriatic lesions, dermatitis, contact dermatitis, eczema, urticaria, rosacea, wound healing, scarring, hypertrophic scarring, keloids, Kawasaki Disease, rosacea, Sjogren-Larsso Syndrome, urticaria.

[00144] In some embodiments, provided is a method of reducing lung injury, vascular leakage, inflammation and/or fibrosis in a mammal comprising administering to the mammal a selective LPA1 receptor antagonist. In some embodiments, provided is a method of reducing lung injury, vascular leakage, inflammation and fibrosis in a mammal comprising administering to the mammal a selective LPA1 receptor antagonist. In some embodiments, provided is a method of attenuating fibrosis in a mammal comprising administering a selective LPA1 receptor antagonist. In some embodiments, provided is a method of attenuating tissue remodeling and fibrosis in a mammal comprising administering a selective LPA1 receptor antagonist.

[00145] In some embodiments, provided is a method of decreasing cytokine production in a mammal comprising administering a selective LPA1 receptor antagonist. In some embodiments, the method of decreasing cytokine production in a mammal comprising administering a selective LPA1 receptor antagonist results in a reduction of tissue damage and fibrosis in a mammal.

[00146] In some embodiments, provided is a method of treating fibrosis in a mammal comprising administering to the mammal a selective LPA1 receptor antagonist. In some embodiments, provided is a method of treating fibrosis in a mammal while maintaining body weight in the

mammal comprising administering to the mammal a selective LPA₁ receptor antagonist. In some embodiments, provided is a method of treating respiratory disease in a mammal comprising administering to the mammal a selective LPA₁ receptor antagonist.

[00147] In some embodiments, provided is a method of treating fibrosis in a mammal with a

5 selective LPA₁ receptor antagonist, wherein the fibrosis in the mammal is not responsive to treatment with pirfenidone. In some embodiments, the LPA₁ receptor antagonist is a compound of Formula (I), Formula (II) or Formula (III).

[00148] As shown in the Examples, a selective LPA₁ receptor antagonist reduced lung fibrosis, kidney fibrosis and liver fibrosis in various animal models of fibrosis.

10 [00149] In a mouse bleomycin lung fibrosis model, a selective LPA₁ receptor antagonist (e.g. Compound 1-88), reduced lung injury, vascular leakage, inflammation and fibrosis at multiple timepoints following intratracheal bleomycin instillation. In the acute setting (3 day), a selective LPA₁ receptor antagonist (e.g. Compound 1-88) significantly reduced total protein, lactate dehydrogenase activity (LDH; tissue injury marker) and metalloproteinase-1 (TIMP-1; pro-fibrotic marker), TGF β 1, hyaluronan and MMP7 concentrations in bronchoalveolar lavage fluid (BALF). In the chronic setting (14 and 28 day) model, a selective LPA₁ receptor antagonist (e.g. Compound 1-88) maintained mouse body weight and decreased inflammatory cell influx and fibrosis after a single bleomycin (1.5 units/kg) instillation and decreased pulmonary resistance and lung fibrosis following repeated bleomycin (3.0-5.0 units/kg/week) instillations. A selective 15 LPA₁ receptor antagonist (e.g. Compound 1-88) also demonstrated superior efficacy to pirfenidone at reducing lung tissue damage and fibrosis in a 14 day mouse bleomycin model.

Pain

[00150] Since LPA is released following tissue injury, LPA₁ plays an important role in the initiation of neuropathic pain. LPA₁, unlike LPA₂ or LPA₃, is expressed in both dorsal root 25 ganglion (DRG) and dorsal root neurons. Using the antisense oligodeoxynucleotide (AS-ODN) for LPA₁ and LPA₁-null mice, it was found that LPA-induced mechanical allodynia and hyperalgesia is mediated in an LPA₁-dependent manner. LPA₁ and downstream Rho-ROCK activation play a role in the initiation of neuropathic pain signaling. Pretreatment with Clostridium botulinum C3 exoenzyme (BoTXC3, Rho inhibitor) or Y-27632 (ROCK inhibitor) 30 completely abolished the allodynia and hyperalgesia in nerve-injured mice. LPA also induced demyelination of the dorsal root, which was prevented by BoTXC3. The dorsal root demyelination by injury was not observed in LPA₁-null mice or AS-ODN injected wild-type mice. LPA signaling appears to induce important neuropathic pain markers such as protein kinase C γ (PKC γ) and a voltage-gated calcium channel α 2 δ 1 subunit (Ca α 2 δ 1) in an LPA₁ and Rho-

dependent manner (M. Inoue, *et al.*, Initiation of neuropathic pain requires lysophosphatidic acid receptor signaling, *Nat. Med.* 10 (2004) 712–718).

[00151] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of pain in a mammal. In one aspect, the pain is acute pain or chronic pain. In another aspect, the pain is neuropathic pain. In another aspect, the pain is cancer pain.

[00152] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of fibromyalgia. In one aspect, fibromyalgia stems from the formation of fibrous scar tissue in contractile (voluntary) muscles. Fibrosis binds the tissue and inhibits blood flow, resulting in pain.

10 **Cancer**

[00153] Lysophospholipid receptor signaling plays a role in the etiology of cancer.

Lysophosphatidic acid (LPA) and its G protein-coupled receptors (GPCRs) LPA₁, LPA₂, and/or LPA₃ play a role in the development of several types of cancers. The initiation, progression and metastasis of cancer involve several concurrent and sequential processes including cell proliferation and growth, survival and anti-apoptosis, migration of cells, penetration of foreign cells into defined cellular layers and/or organs, and promotion of angiogenesis. The control of each of these processes by LPA signaling in physiological and pathophysiological conditions underscores the potential therapeutic usefulness of modulating LPA signaling pathways for the treatment of cancer, especially at the level of the LPA receptors or ATX/lysoPLD. Autotaxin (ATX) is a prometastatic enzyme initially isolated from the conditioned medium of human melanoma cells that stimulates a myriad of biological activities, including angiogenesis and the promotion of cell growth, migration, survival, and differentiation through the production of LPA (*Mol Cancer Ther* 2008;7(10):3352–62).

[00154] LPA signals through its own GPCRs leading to activation of multiple downstream effector pathways. Such downstream effector pathways play a role in cancer. LPA and its GPCRs are linked to cancer through major oncogenic signaling pathways.

[00155] LPA contributes to tumorigenesis by increasing motility and invasiveness of cells. LPA has been implicated in the initiation or progression of ovarian cancer. LPA is present at significant concentrations (2–80 μ M) in the ascitic fluid of ovarian cancer patients. Ovarian cancer cells constitutively produce increased amounts of LPA as compared to normal ovarian surface epithelial cells, the precursor of ovarian epithelial cancer. Elevated LPA levels are also detected in plasma from patients with early-stage ovarian cancers compared with controls. LPA receptors (LPA₂ and LPA₃) are also overexpressed in ovarian cancer cells as compared to normal ovarian surface epithelial cells. LPA stimulates Cox-2 expression through transcriptional activation and post-transcriptional enhancement of Cox-2 mRNA in ovarian cancer cells.

Prostaglandins produced by Cox-2 have been implicated in a number of human cancers and pharmacological inhibition of Cox-2 activity reduces colon cancer development and decreases the size and number of adenomas in patients with familial adenomatous polyposis. LPA has also been implicated in the initiation or progression of prostate cancer, breast cancer, melanoma, head and neck cancer, bowel cancer (colorectal cancer), thyroid cancer, glioblastoma, and other cancers (Gardell *et al*, *Trends in Molecular Medicine*, vol. 12, no. 2, p 65-75, 2006; Ishii *et al*, *Annu. Rev. Biochem.*, 73, 321-354, 2004; Mills *et al*., *Nat. Rev. Cancer*, 3, 582-591, 2003; Murph *et al*., *Biochimica et Biophysica Acta*, 1781, 547-557, 2008; Kishi *et al*., *J. Biol. Chem.*, 281, 17492-17500, 2006;).

[00156] The cellular responses to LPA are mediated through the lysophosphatidic acid receptors. For example, LPA receptors mediate both migration of and invasion by pancreatic cancer cell lines: an antagonist of LPA₁ and LPA₃ (Ki16425) and LPA₁-specific siRNA effectively blocked in vitro migration in response to LPA and peritoneal fluid (ascites) from pancreatic cancer patients; in addition, Ki16425 blocked the LPA-induced and ascites-induced invasion activity of a highly peritoneal metastatic pancreatic cancer cell line (Yamada *et al*, *J. Biol. Chem.*, 279, 6595-6605, 2004).

[00157] Colorectal carcinoma cell lines show significant expression of LPA₁ mRNA and respond to LPA by cell migration and production of angiogenic factors. Overexpression of LPA receptors has a role in the pathogenesis of thyroid cancer. LPA₃ was originally cloned from prostate cancer cells, concordant with the ability of LPA to induce autocrine proliferation of prostate cancer cells.

[00158] LPA has stimulatory roles in cancer progression in many types of cancer. LPA is produced from and induces proliferation of prostate cancer cell lines. LPA induces human colon carcinoma DLD1 cell proliferation, migration, adhesion, and secretion of angiogenic factors through LPA₁ signalling. In other human colon carcinoma cells lines (HT29 and WiDR), LPA enhances cell proliferation and secretion of angiogenic factors. In other colon cancer cell lines, LPA₂ and LPA₃ receptor activation results in proliferation of the cells. LPA₁ is implicated in bone metastasis and the LPA₁/LPA₃ dual antagonist Ki16425 has been shown to inhibit metastasis to bone in vivo (Boucharaba *et al*., *Proc. Natl. Acad. Sci USA*, 103, 9643-9648, 2006). The genetic or pharmacological manipulation of LPA metabolism, specific blockade of receptor signaling, and/or inhibition of downstream signal transduction pathways, represent approaches for cancer therapies.

[00159] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of cancer. In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of malignant and benign proliferative disease. In one aspect, a compound of

Formula (I), Formula (II) or Formula (III) is used to prevent or reduce proliferation of tumor cells, invasion and metastasis of carcinomas, pleural mesothelioma (Yamada, *Cancer Sci.*, 2008, 99(8), 1603-1610) or peritoneal mesothelioma, cancer pain, bone metastases (Boucharaba *et al.*, *J. Clin. Invest.*, 2004, 114(12), 1714-1725; Boucharaba *et al.*, *Proc. Natl. Acad. Sci.*, 2006,

5 103(25) 9643-9648). In one aspect is a method of treating cancer in a mammal, the method comprising administering to the mammal a compound of Formula (I), Formula (II) or Formula (III) and a second therapeutic agent, wherein the second therapeutic agent is an anti-cancer agent.

[00160] The term “cancer,” as used herein refers to an abnormal growth of cells which tend to proliferate in an uncontrolled way and, in some cases, to metastasize (spread). The types of 10 cancer include, but is not limited to, solid tumors (such as those of the bladder, bowel, brain, breast, endometrium, heart, kidney, lung, lymphatic tissue (lymphoma), ovary, pancreas or other endocrine organ (thyroid), prostate, skin (melanoma or basal cell cancer) or hematological tumors (such as the leukemias) at any stage of the disease with or without metastases.

[00161] The increased concentrations of LPA and vesicles in ascites from ovarian cancer patients 15 and breast cancer effusions indicate that it could be an early diagnostic marker, a prognostic indicator or an indicator of response to therapy (Mills *et al.*, *Nat. Rev. Cancer.*, 3, 582-591, 2003; Sutphen *et al.*, *Cancer Epidemiol. Biomarkers Prev.* 13, 1185-1191, 2004). LPA concentrations are consistently higher in ascites samples than in matched plasma samples.

Respiratory and Allergic Disorders

20 [00162] In one aspect, LPA is a contributor to the pathogenesis of respiratory diseases. In one aspect the respiratory disease is asthma. Proinflammatory effects of LPA include degranulation of mast cells, contraction of smooth-muscle cells and release of cytokines from dendritic cells. Airway smooth muscle cells, epithelial cells and lung fibroblasts all show responses to LPA.

25 LPA induces the secretion of IL-8 from human bronchial epithelial cells. IL-8 is found in increased concentrations in BAL fluids from patients with asthma, chronic obstructive lung disease, pulmonary sarcoidosis and acute respiratory distress syndrome and IL-8 has been shown to exacerbate airway inflammation and airway remodeling of asthmatics. LPA1, LPA2 and LPA3 receptors have all been shown to contribute to the LPA-induced IL-8 production. Studies cloning multiple GPCRs that are activated by LPA allowed the demonstration of the presence of mRNA 30 for the LPA₁, LPA₂ and LPA₃ in the lung (J.J.A. Contos, *et al.*, *Mol. Pharmacol.* 58, 1188-1196, 2000).

35 [00163] The release of LPA from platelets activated at a site of injury and its ability to promote fibroblast proliferation and contraction are features of LPA as a mediator of wound repair. In the context of airway disease, asthma is an inflammatory disease where inappropriate airway “repair” processes lead to structural “remodeling” of the airway. In asthma, the cells of the

airway are subject to ongoing injury due to a variety of insults, including allergens, pollutants, other inhaled environmental agents, bacteria and viruses, leading to the chronic inflammation that characterizes asthma.

[00164] In one aspect, in the asthmatic individual, the release of normal repair mediators, including LPA, is exaggerated or the actions of the repair mediators are inappropriately prolonged leading to inappropriate airway remodeling. Major structural features of the remodeled airway observed in asthma include a thickened lamina reticularis (the basement membrane-like structure just beneath the airway epithelial cells), increased numbers and activation of myofibroblasts, thickening of the smooth muscle layer, increased numbers of mucus glands and mucus secretions, and alterations in the connective tissue and capillary bed throughout the airway wall. In one aspect, LPA contributes to these structural changes in the airway. In one aspect, LPA is involved in acute airway hyperresponsiveness in asthma. The lumen of the remodeled asthmatic airway is narrower due to the thickening of the airway wall, thus decreasing airflow. In one aspect, LPA contributes to the long-term structural remodeling and the acute hyperresponsiveness of the asthmatic airway. In one aspect, LPA contributes to the hyper-responsiveness that is a primary feature of acute exacerbations of asthma.

[00165] In addition to the cellular responses mediated by LPA, several of the LPA signaling pathway components leading to these responses are relevant to asthma. EGF receptor upregulation is induced by LPA and is also seen in asthmatic airways (M. Amishima, *et al.*, *Am. J. Respir. Crit. Care Med.* 157, 1907– 1912, 1998). Chronic inflammation is a contributor to asthma, and several of the transcription factors that are activated by LPA are known to be involved in inflammation (Ediger *et al.*, *Eur Respir J* 21:759-769, 2003).

[00166] In one aspect, the fibroblast proliferation and contraction and extracellular matrix secretion stimulated by LPA contributes to the fibroproliferative features of other airway diseases, such as the peribronchiolar fibrosis present in chronic bronchitis, emphysema, and interstitial lung disease. Emphysema is also associated with a mild fibrosis of the alveolar wall, a feature which is believed to represent an attempt to repair alveolar damage. In another aspect, LPA plays a role in the fibrotic interstitial lung diseases and obliterative bronchiolitis, where both collagen and myofibroblasts are increased. In another aspect, LPA is involved in several of the various syndromes that constitute chronic obstructive pulmonary disease.

[00167] Administration of LPA *in vivo* induces airway hyper-responsiveness, itch-scratch responses, infiltration and activation of eosinophils and neutrophils, vascular remodeling, and nociceptive flexor responses. LPA also induces histamine release from mouse and rat mast cells. In an acute allergic reaction, histamine induces various responses, such as contraction of smooth muscle, plasma exudation, and mucus production. Plasma exudation is important in the airway,

because the leakage and subsequent airway-wall edema contribute to the development of airway hyperresponsiveness. Plasma exudation progresses to conjunctival swelling in ocular allergic disorder and nasal blockage in allergic rhinitis (Hashimoto *et al.*, *J Pharmacol Sci* 100, 82 – 87, 2006). In one aspect, plasma exudation induced by LPA is mediated by histamine release from mast cells via one or more LPA receptors. In one aspect, the LPA receptor(s) include LPA₁ and/or LPA₃. In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of various allergic disorders in a mammal. In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of respiratory diseases, disorders or conditions in a mammal. In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of asthma in a mammal. In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is used in the treatment of chronic asthma in a mammal.

[00168] The term “respiratory disease,” as used herein, refers to diseases affecting the organs that are involved in breathing, such as the nose, throat, larynx, eustachian tubes, trachea, bronchi, lungs, related muscles (e.g., diaphragm and intercostals), and nerves. Respiratory diseases include, but are not limited to, asthma, adult respiratory distress syndrome and allergic (extrinsic) asthma, non-allergic (intrinsic) asthma, acute severe asthma, chronic asthma, clinical asthma, nocturnal asthma, allergen-induced asthma, aspirin-sensitive asthma, exercise-induced asthma, isocapnic hyperventilation, child-onset asthma, adult-onset asthma, cough-variant asthma, occupational asthma, steroid-resistant asthma, seasonal asthma, seasonal allergic rhinitis, perennial allergic rhinitis, chronic obstructive pulmonary disease, including chronic bronchitis or emphysema, pulmonary hypertension, interstitial lung fibrosis and/or airway inflammation and cystic fibrosis, and hypoxia.

[00169] The term “asthma” as used herein refers to any disorder of the lungs characterized by variations in pulmonary gas flow associated with airway constriction of whatever cause (intrinsic, extrinsic, or both; allergic or non-allergic). The term asthma may be used with one or more adjectives to indicate cause.

[00170] In one aspect, presented herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the treatment or prevention of chronic obstructive pulmonary disease in a mammal comprising administering to the mammal at least once an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III). In addition, chronic obstructive pulmonary disease includes, but is not limited to, chronic bronchitis or emphysema, pulmonary hypertension, interstitial lung fibrosis and/or airway inflammation, and cystic fibrosis.

Nervous System

[00171] The nervous system is a major locus for LPA₁ expression; there it is spatially and temporally regulated throughout brain development. Oligodendrocytes, the myelinating cells in

the central nervous system (CNS), express LPA₁ in mammals. In addition, Schwann cells, the myelinating cells of the peripheral nervous system, also express LPA₁, which is involved in regulating Schwann cell survival and morphology. These observations identify important functions for receptor-mediated LPA signaling in neurogenesis, cell survival, and myelination.

5 [00172] Exposure of peripheral nervous system cell lines to LPA produces a rapid retraction of their processes resulting in cell rounding, which was, in part, mediated by polymerization of the actin cytoskeleton. In one aspect, LPA causes neuronal degeneration under pathological conditions when the blood-brain barrier is damaged and serum components leak into the brain (Moolenaar, *Curr. Opin. Cell Biol.* 7:203-10, 1995). Immortalized CNS neuroblast cell lines 10 from the cerebral cortex also display retraction responses to LPA exposure through Rho activation and actomyosin interactions. In one aspect, LPA is associated with post-ischemic neural damage (*J. Neurochem.* 61, 340, 1993; *J. Neurochem.*, 70:66, 1998).

15 [00173] In one aspect, provided is a compound of Formula (I), Formula (II) or Formula (III) for use in the treatment or prevention of a nervous system disorder in a mammal. The term “nervous system disorder,” as used herein, refers to conditions that alter the structure or function of the brain, spinal cord or peripheral nervous system, including but not limited to Alzheimer’s Disease, cerebral edema, cerebral ischemia, stroke, multiple sclerosis, neuropathies, Parkinson’s Disease, those found after blunt or surgical trauma (including post-surgical cognitive dysfunction and spinal cord or brain stem injury), as well as the neurological aspects of disorders such as 20 degenerative disk disease and sciatica.

25 [00174] In one aspect, provided is a compound of Formula (I), Formula (II) or Formula (III) for use in the treatment or prevention of a CNS disorder in a mammal. CNS disorders include, but are not limited to, multiple sclerosis, Parkinson’s disease, Alzheimer’s disease, stroke, cerebral ischemia, retinal ischemia, post-surgical cognitive dysfunction, migraine, peripheral neuropathy/neuropathic pain, spinal cord injury, cerebral edema and head injury.

Cardiovascular Disorders

30 [00175] Cardiovascular phenotypes observed after targeted deletion of lysophospholipid receptors reveal important roles for lysophospholipid signaling in the development and maturation of blood vessels, formation of atherosclerotic plaques and maintenance of heart rate (Ishii, I. *et al. Annu. Rev. Biochem.* 73, 321-354, 2004). Angiogenesis, the formation of new capillary networks from pre-existing vasculature, is normally invoked in wound healing, tissue growth and myocardial angiogenesis after ischemic injury. Peptide growth factors (e.g. vascular endothelial growth factor (VEGF)) and lysophospholipids control coordinated proliferation, migration, adhesion, differentiation and assembly of vascular endothelial cells (VECs) and surrounding vascular 35 smooth-muscle cells (VSMCs). In one aspect, dysregulation of the processes mediating

angiogenesis leads to atherosclerosis, hypertension, tumor growth, rheumatoid arthritis and diabetic retinopathy (Osborne, N. and Stainier, D.Y. *Annu. Rev. Physiol.* 65, 23–43, 2003).

[00176] Downstream signaling pathways evoked by lysophospholipid receptors include Rac-dependent lamellipodia formation (e.g. LPA₁) and Rho-dependent stress-fiber formation (e.g.

5 LPA₁), which is important in cell migration and adhesion. Dysfunction of the vascular endothelium can shift the balance from vasodilatation to vasoconstriction and lead to hypertension and vascular remodeling, which are risk factors for atherosclerosis (Maguire, J.J. et al., *Trends Pharmacol. Sci.* 26, 448–454, 2005).

[00177] LPA contributes to both the early phase (barrier dysfunction and monocyte adhesion of 10 the endothelium) and the late phase (platelet activation and intra-arterial thrombus formation) of atherosclerosis, in addition to its overall progression. In the early phase, LPA from numerous sources accumulates in lesions and activates its cognate GPCRs (LPA₁ and LPA₃) expressed on platelets (Siess, W. *Biochim. Biophys. Acta* 1582, 204–215, 2002; Rother, E. et al. *Circulation* 108, 741–747, 2003). This triggers platelet shape change and aggregation, leading to intra-arterial 15 thrombus formation and, potentially, myocardial infarction and stroke. In support of its atherogenic activity, LPA can also be a mitogen and motogen to VSMCs and an activator of endothelial cells and macrophages. LPA has been shown to be involved in ischemia-reperfusion injury. Blockade of the LPA₃ receptor in a murine model of renal ischemia-reperfusion injury reduced the severity of injury. This effect was reversed in the presence of the selective LPA₃ 20 receptor agonist OMPT (Okusa et al., *Am. J. Physiol. Renal Physiol.*, 285, F565-F574, 2003). In one aspect, mammals with cardiovascular disease benefit from LPA receptor antagonists that prevent thrombus and neointima plaque formation.

[00178] The specific effects of LPA are receptor-mediated.

[00179] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are used to treat 25 or prevent cardiovascular disease in mammal.

[00180] The term “cardiovascular disease,” as used herein refers to diseases affecting the heart or blood vessels or both, including but not limited to: arrhythmia (atrial or ventricular or both); atherosclerosis and its sequelae; angina; cardiac rhythm disturbances; myocardial ischemia; myocardial infarction; cardiac or vascular aneurysm; vasculitis, stroke; peripheral obstructive 30 arteriopathy of a limb, an organ, or a tissue; reperfusion injury following ischemia of the brain, heart, kidney or other organ or tissue; endotoxic, surgical, or traumatic shock; hypertension, valvular heart disease, heart failure, abnormal blood pressure; shock; vasoconstriction (including that associated with migraines); vascular abnormality, inflammation, insufficiency limited to a single organ or tissue..

[00181] In one aspect, provided herein are methods for preventing or treating vasoconstriction, atherosclerosis and its sequelae myocardial ischemia, myocardial infarction, aortic aneurysm, vasculitis and stroke comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III) or pharmaceutical

5 composition or medicament which includes a compound of Formula (I), Formula (II) or Formula (III).

[00182] In one aspect, provided herein are methods for reducing cardiac reperfusion injury following myocardial ischemia and/or endotoxic shock comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or

10 Formula (III).

[00183] In one aspect, provided herein are methods for reducing the constriction of blood vessels in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

[00184] In one aspect, provided herein are methods for lowering or preventing an increase in

15 blood pressure of a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

Inflammation

[00185] LPA has been shown to regulate immunological responses by modulating activities/functions of immune cells such as T-/B-lymphocytes and macrophages. In activated T

20 cells, LPA activates IL-2 production/cell proliferation through LPA₁ (Gardell et al, *TRENDS in Molecular Medicine* Vol.12 No.2 February 2006). Expression of LPA-induced inflammatory

response genes is mediated by LPA₁ and LPA₃ (*Biochem Biophys Res Commun.* 363(4):1001-8, 2007). In addition, LPA modulates the chemotaxis of inflammatory cells (*Biochem Biophys Res Commun.*, 1993, 15;193(2), 497). The proliferation and cytokine-secreting activity in response to

25 LPA of immune cells (*J. Immunol.* 1999, 162, 2049), platelet aggregation activity in response to LPA, acceleration of migration activity in monocytes, activation of NF-κB in fibroblast, enhancement of fibronectin-binding to the cell surface, and the like are known. Thus, LPA is associated with various inflammatory/immune diseases .

[00186] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are used to treat

30 or prevent inflammation in a mammal. In one aspect, antagonists of LPA₁ and/or LPA₃ find use in the treatment or prevention of inflammatory/immune disorders in a mammal. In one aspect, the antagonist of LPA₁ is a compound of Formula (I), Formula (II) or Formula (III).

[00187] Examples of inflammatory/immune disorders include psoriasis, rheumatoid arthritis, vasculitis, inflammatory bowel disease, dermatitis, osteoarthritis, asthma, inflammatory muscle disease, allergic rhinitis, vaginitis, interstitial cystitis, scleroderma, eczema, allogeneic or

xenogeneic transplantation (organ, bone marrow, stem cells and other cells and tissues) graft rejection, graft-versus-host disease, lupus erythematosus, inflammatory disease, type I diabetes, pulmonary fibrosis, dermatomyositis, Sjogren's syndrome, thyroiditis (e.g., Hashimoto's and autoimmune thyroiditis), myasthenia gravis, autoimmune hemolytic anemia, multiple sclerosis, 5 cystic fibrosis, chronic relapsing hepatitis, primary biliary cirrhosis, allergic conjunctivitis and atopic dermatitis.

Other Diseases, Disorders or Conditions

[00188] In accordance with one aspect, are methods for treating, preventing, reversing, halting or slowing the progression of LPA-dependent or LPA-mediated diseases or conditions once it

10 becomes clinically evident, or treating the symptoms associated with or related to LPA-dependent or LPA-mediated diseases or conditions, by administering to the mammal a compound of Formula (I), Formula (II) or Formula (III). In certain embodiments, the subject already has a LPA-dependent or LPA-mediated disease or condition at the time of administration, or is at risk of developing a LPA-dependent or LPA-mediated disease or condition.

15 [00189] In certain aspects, the activity of LPA₁ in a mammal is directly or indirectly modulated by the administration of (at least once) a therapeutically effective amount of at least one compound of Formula (I), Formula (II) or Formula (III). Such modulation includes, but is not limited to, reducing and/or inhibiting the activity of LPA₁. In additional aspects, the activity of LPA in a mammal is directly or indirectly modulated, including reducing and/or inhibiting, by the 20 administration of (at least once) a therapeutically effective amount of at least one compound of Formula (I), Formula (II) or Formula (III). Such modulation includes, but is not limited to, reducing and/or inhibiting the amount and/or activity of a LPA receptor. In one aspect, the LPA receptor is LPA₁.

25 [00190] In one aspect, LPA has a contracting action on bladder smooth muscle cell isolated from bladder, and promotes growth of prostate-derived epithelial cell (*The Journal of Urology* , 1999, vol. 162, pp. 1779-1784; *The Journal of Urology* , 2000, vol. 163, pp. 1027-1032). In another aspect, LPA contracts the urinary tract and prostate *in vitro* and increases intraurethral pressure *in vivo* (WO 02/062389).

30 [00191] In certain aspects, are methods for preventing or treating eosinophil and/or basophil and/or dendritic cell and/or neutrophil and/or monocyte and/or T-cell recruitment comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

35 [00192] In certain aspects, are methods for the treatment of cystitis, including, e.g., interstitial cystitis, comprising administering at least once to the mammal a therapeutically effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

[00193] In accordance with one aspect, methods described herein include the diagnosis or determination of whether or not a patient is suffering from a LPA-dependent or LPA-mediated disease or condition by administering to the subject a therapeutically effective amount of a compound of Formula (I), Formula (II) or Formula (III) and determining whether or not the patient responds to the treatment.

[00194] In one aspect provided herein are compounds of Formula (I), Formula (II) and Formula (III), pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, which are antagonists of LPA₁, and are used to treat patients suffering from one or more LPA-dependent or LPA-mediated conditions or diseases, including, but not limited to, lung fibrosis, kidney fibrosis, liver fibrosis, scarring, asthma, rhinitis, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, pain, proliferative disorders and inflammatory conditions. In some embodiments, LPA-dependent conditions or diseases include those wherein an absolute or relative excess of LPA is present and/or observed.

[00195] In any of the aforementioned aspects the LPA-dependent or LPA-mediated diseases or conditions include, but are not limited to, organ fibrosis, asthma, allergic disorders, chronic obstructive pulmonary disease, pulmonary hypertension, lung or pleural fibrosis, peritoneal fibrosis, arthritis, allergy, cancer, cardiovascular disease, ult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, and cancer.

[00196] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are used to improve the corneal sensitivity decrease caused by corneal operations such as laser-assisted in situ keratomileusis (LASIK) or cataract operation, corneal sensitivity decrease caused by corneal degeneration, and dry eye symptom caused thereby.

[00197] In one aspect, presented herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the treatment or prevention of ocular inflammation and allergic conjunctivitis, vernal keratoconjunctivitis, and papillary conjunctivitis in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

[00198] In one aspect, presented herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the treatment or prevention of Sjogren disease or inflammatory disease with dry eyes in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

[00199] In one aspect, LPA and LPA receptors (e.g. LPA₁) are involved in the pathogenesis of osteoarthritis (Kotani *et al*, *Hum. Mol. Genet.*, 2008, 17, 1790-1797). In one aspect, presented

herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the treatment or prevention of osteoarthritis in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).


[00200] In one aspect, LPA receptors (e.g. LPA₁, LPA₃) contribute to the pathogenesis of rheumatoid arthritis (Zhao *et al*, *Mol. Pharmacol.*, 2008, 73(2), 587-600). In one aspect, presented herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the treatment or prevention of rheumatoid arthritis in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

[00201] In one aspect, LPA receptors (e.g. LPA₁) contribute to adipogenesis. (Simon *et al*, *J.Biol. Chem.*, 2005, vol. 280, no. 15, p.14656). In one aspect, presented herein is the use of a compound of Formula (I), Formula (II) or Formula (III) in the promotion of adipose tissue formation in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of Formula (I), Formula (II) or Formula (III).

15 Compounds

[00202] Compounds of Formula (I), Formula (II) and Formula (III), including pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, antagonize or modulate LPA₁ and are used to treat patients suffering from LPA₁-dependent or LPA₁-mediated conditions or diseases.

[00203] In one aspect, provided herein is a compound of Formula (I), pharmaceutically acceptable salt, solvate, metabolite, or prodrug thereof:



Formula (I)

wherein

25 R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -C(=O)NH-OH, -C(=O)NH-CN, -CN, tetrazolyl or carboxylic acid bioisostere;

R^D is H or C₁-C₆alkyl;

ring A is a substituted or unsubstituted cycloalkylene, a substituted or unsubstituted heterocycloalkylene, a substituted or unsubstituted arylene, or a substituted or unsubstituted heteroarylene, where if ring A is substituted then each substituent on ring A is H or R^A;

ring B is a substituted or unsubstituted cycloalkylene, a substituted or unsubstituted heterocycloalkylene, a substituted or unsubstituted arylene, or a substituted or

unsubstituted heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B;

5 L¹ is absent, a substituted or unsubstituted alkylene, a substituted or unsubstituted fluoroalkylene, or a substituted or unsubstituted heteroalkylene;

L² is absent, a substituted or unsubstituted alkylene, a substituted or unsubstituted fluoroalkylene, a substituted or unsubstituted heteroalkylene, -O-, -S-, -SO-, -SO₂-, -NR²-, or -C(=O)-;

R² is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

R³ is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

10 R⁴ is -NR⁷C(=O)X¹CH(R⁸)-CY, or -NR⁷C(=O)X¹-CY;

R⁷ is H or C₁-C₄alkyl;

R⁸ is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

X¹ is -O-, NR⁷, or -C(R⁸)₂-;

CY is a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl, wherein if CY is substituted then each substituent on CY is H or R^C; or

15 R⁸ and CY are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted carbocycle or a substituted or unsubstituted heterocycle;

20 n is 0, 1, or 2;

each of R^A, R^B, and R^C are independently selected from H, halogen, -CN, -NO₂, -OH, -OR¹⁰, -S(=O)₂R¹⁰, -SR¹⁰, -S(=O)R¹⁰, -N(R⁹)S(=O)₂R¹⁰, -S(=O)₂N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR⁹C(=O)N(R⁹)₂, -NR⁹C(=O)R¹⁰, -NR⁹C(=O)OR¹⁰, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl;

25 each R⁹ is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted -C₁-C₄alkylene-cycloalkyl, a substituted or unsubstituted -C₁-C₄alkylene-heterocycloalkyl, a substituted or unsubstituted -C₁-C₄alkylene-aryl, or a substituted or unsubstituted -C₁-C₄alkylene-heteroaryl; or

30 two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted heterocycle;

R¹⁰ is selected from C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted -C₁-C₄alkylene-cycloalkyl, a substituted or unsubstituted -C₁-C₄alkylene-heterocycloalkyl, a substituted or unsubstituted -C₁-C₄alkylene-aryl, and a substituted or unsubstituted -C₁-C₄alkylene-heteroaryl.

[00204] For any and all of the embodiments, substituents are selected from among from a subset of the listed alternatives. For example, in some embodiments, R³ is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl. In other embodiments, R³ is C₁-C₄alkyl, C₃-C₆cycloalkyl, or C₁-C₄fluoroalkyl. In

10 some embodiments, R³ is H, C₁-C₄alkyl, cyclopropyl, or C₁-C₄fluoroalkyl. In some embodiments, R³ is C₁-C₄alkyl, cyclopropyl, or C₁-C₄fluoroalkyl. In other embodiments, R³ is C₁-C₄alkyl, or C₁-C₄fluoroalkyl. In yet other embodiments, R³ is C₁-C₄alkyl. In some embodiments, R³ is -CH₃ or -CH₂CH₃. In some embodiments, R³ is -CH₃, -CF₃, -CH₂CH₃, -CHCH₂, or cyclopropyl. In some embodiments, R³ is -CH₃, -CF₃, -CH₂CH₃, or -CHCH₂. In one aspect, R³ is -CH₃.

[00205] In some embodiments, ring A is a substituted or unsubstituted arylene, or a substituted or unsubstituted heteroarylene, where if ring A is substituted then each substituent on ring A is H or R^A.

20 [00206] In some embodiments, ring A is a substituted or unsubstituted phenylene, a substituted or unsubstituted napthylene, or a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring A is substituted then each substituent on ring A is H or R^A.



[00207] In some embodiments, ring A is a substituted or unsubstituted phenylene, or a substituted or unsubstituted monocyclic heteroarylene containing 0 or 1 O atoms, 0 or 1 S atoms, 0-3 N atoms, and at least 2 carbon atoms in the ring, where if ring A is substituted then each substituent on ring A is H or R^A.

[00208] In some embodiments, ring A is a substituted or unsubstituted phenylene, or a substituted or unsubstituted pyridinylene, where if ring A is substituted then each substituent on ring A is H or R^A.


30 [00209] In some embodiments, ring B is a substituted or unsubstituted monocyclic cycloalkylene, a substituted or unsubstituted bicyclic cycloalkylene, a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted bicyclic heterocycloalkylene, a substituted or unsubstituted phenylene, a substituted or unsubstituted naphthylene, a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B.

[00210] In some embodiments, each of R^A , R^B , and R^C are independently selected from H, halogen, -CN, -NO₂, -OH, -OR¹⁰, -N(R⁹)₂, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[00211] In some embodiments, each of R^A , R^B , and R^C are independently selected from H, halogen, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[00212] In one aspect, the groups  and  are on non-adjacent ring atoms of ring A.

[00213] In one aspect, the compound of Formula (I) has the structure of Formula (II):



10 Formula (II)

wherein

each of A¹, A², and A³ is independently selected from CR^A and N.

[00214] In some embodiments, each R^A is independently selected from H, halogen, -CH₃, -CF₃, -OCF₃, and -OCH₃.

15 [00215] In some embodiments, each of A¹, A², and A³ is CR^A.

[00216] In some embodiments, one of A¹, A², and A³ is N.

[00217] In some embodiments, A¹ is N. In some embodiments, A² is N. In some embodiments, A³ is N.

[00218] In some embodiments, R⁴ is -NR⁷C(=O)OCH(R⁸)-CY, -NR⁷C(=O)NR⁷CH(R⁸)-CY, -NR⁷C(=O)CH₂CH(R⁸)-CY, or -NR⁷C(=O)O-CY.

20 [00219] In one aspect, R⁴ is -NR⁷C(=O)OCH(R⁸)-CY.

[00220] In one aspect, R⁷ is H. In some embodiments, R⁸ is H, C₁-C₄alkyl or C₁-C₄fluoroalkyl. In some embodiments, R⁸ is C₁-C₄alkyl or C₁-C₄fluoroalkyl. In some other embodiments, R⁸ is H, -CH₃ or -CF₃. In yet other embodiments, R⁸ is -CH₃ or -CF₃. In one aspect, R⁸ is -CH₃.

25 [00221] In some embodiments, R¹ is -CO₂H or -CO₂R^D. In some embodiments, R^D is H, -CH₃, or -CH₂CH₃.

[00222] In one aspect, R¹ is -CO₂H.

[00223] In some embodiments, CY is a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl, wherein if CY is substituted then each substituent on CY is H or R^C.

[00224] In some embodiments, CY is a substituted or unsubstituted monocyclic cycloalkyl, a substituted or unsubstituted bicyclic cycloalkyl, a substituted or unsubstituted phenyl, a

substituted or unsubstituted napthyl, a substituted or unsubstituted monocyclic heteroaryl containing 0-4 N atoms, or a substituted or unsubstituted bicyclic heteroaryl containing 0-4 N atoms, wherein if CY is substituted then each substituent on CY is H or R^C.

[00225] In some embodiments, CY is a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl containing 1-3 N atoms, wherein if CY is substituted then each substituent on CY is H or R^C.

[00226] In some embodiments, each R^C is independently selected from H, halogen, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[00227] In one aspect, in any of the embodiments described herein, A is unsubstituted or monosubstituted with R^A. In one aspect, in any of the embodiments described herein, A is unsubstituted.

[00228] In one aspect, in any of the embodiments described herein, B is unsubstituted or monosubstituted with R^B. In one aspect, in any of the embodiments described herein, B is unsubstituted.

[00229] In one aspect, in any of the embodiments described herein, CY is unsubstituted or monosubstituted with R^C.

[00230] In some embodiments, L¹ is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, or C₁-C₆heteroalkylene.

[00231] In alternative embodiments, L² is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, C₁-C₆heteroalkylene, -O-, -S-, -SO-, -SO₂- or -NR²-.

[00232] In some other embodiments, L² is absent, C₁-C₆alkylene, -O-, -S-, -SO-, -SO₂- or -NR²-.

In yet other embodiments, L² is absent, -CH₂-, -O-, -S-, -SO-, -SO₂- or -NR²-.

In yet other embodiments, L² is absent, -CH₂-, -O-, -S-, -SO-, or -SO₂-.

In some embodiments, L² is absent.

In yet other embodiments, L² is -CH₂-.

In some embodiments, L² is -O-, -S-, -SO-, or -SO₂-.

[00233] In one aspect, ring B is a substituted or unsubstituted monocyclic cycloalkylene, a substituted or unsubstituted bicyclic cycloalkylene, a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted bicyclic heterocycloalkylene, a substituted or unsubstituted phenylene, a substituted or unsubstituted naphthylene, a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B.

[00234] In some embodiments, each R^B is independently selected from H, halogen, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[00235] In some embodiments, L² is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₆alkylene-O-, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₆alkylene-S-, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₆alkylene-SO-, -C₁-C₃alkylene-SO-C₁-

C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₆alkylene-SO₂-, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₄alkylene-NR¹¹-, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, -NR¹¹-C₁-C₄alkylene -, -O-, -S-, -SO-, -SO₂- or -NR²-; R¹¹ is H or C₁-C₂alkyl.

[00236] In some embodiments, L² is absent, -CH₂-, -CH₂O-, -OCH₂-, -CH₂S-, -SCH₂-, -CH₂NH-, -NHCH₂-, -O-, -S-, or -NH-. In some other embodiments, L² is -CH₂O-, -OCH₂-, -CH₂S-, -SCH₂-, -CH₂NH-, or -NHCH₂-. In some embodiments, L² is absent, -CH₂-, or -CH₂O-. In some embodiments, L² is absent.

[00237] In alternative embodiments, L¹ is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, or -NR¹¹-C₁-C₄alkylene -; R¹¹ is H or C₁-C₂alkyl.

[00238] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -OCH₂-, -OCH(CH₃)-, -OC(CH₃)₂-, -OCH₂CH₂-, -CH₂OCH₂-, -CH₂OCH₂CH₂-, -CH₂CH₂OCH₂-, -SCH₂-, -SCH(CH₃)-, -SC(CH₃)₂-, -SCH₂CH₂-, -CH₂SCH₂-, -CH₂SCH₂CH₂-, -CH₂CH₂SCH₂-, -SO₂CH₂-, -SO₂CH(CH₃)-, -SO₂C(CH₃)₂-, -SO₂CH₂CH₂-, -CH₂SO₂CH₂-, -CH₂SO₂CH₂CH₂-, -CH₂CH₂SO₂CH₂-, -NHCH₂-, -NHCH(CH₃)-, -NHC(CH₃)₂-, -NHCH₂CH₂-, -CH₂NHCH₂-, -CH₂NHCH₂CH₂-, or -CH₂CH₂NHCH₂-.

[00239] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -OCH₂-, -OCH(CH₃)-, -OC(CH₃)₂-, -OCH₂CH₂-, -CH₂OCH₂-, -CH₂OCH₂CH₂-, -CH₂CH₂OCH₂-, -SCH₂-, -SCH(CH₃)-, -SC(CH₃)₂-, -SCH₂CH₂-, -CH₂SCH₂-, -CH₂SCH₂CH₂-, or -CH₂CH₂SCH₂-.

[00240] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂- or -CH₂CH₂CH₂-. In some embodiments, L¹ is -CH₂-, -CH(CH₃)-, or -C(CH₃)₂-. In some embodiments, L¹ is -CH₂-.

[00241] In some embodiments, ring B is a substituted or unsubstituted monocyclic cycloalkylene, or a substituted or unsubstituted phenylene, where if ring B is substituted then each substituent on ring B is H or R^B.

[00242] In some other embodiments, ring B is a substituted or unsubstituted cyclopentylene, a substituted or unsubstituted cyclopentenylene, a substituted or unsubstituted cyclohexylene, a substituted or unsubstituted cyclohexenylene, or a substituted or unsubstituted phenylene, where if ring B is substituted then each substituent on ring B is H or R^B.

[00243] In other embodiments, ring B is a substituted or unsubstituted phenylene, where if ring B is substituted then each substituent on ring B is H or R^B. In other embodiments, ring B is an unsubstituted phenylene.

[00244] In some embodiments, ring B is a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted bicyclic heterocycloalkylene, a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B.

5 [00245] In some embodiments, ring B is a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B.


[00246] In some embodiments, ring B is a substituted or unsubstituted monocyclic heteroarylene containing 0 or 1 O atoms, 0 or 1 S atoms, 0-3 N atoms, and at least 2 carbon atoms in the ring, 10 where if ring B is substituted then each substituent on ring B is H or R^B.

[00247] In some embodiments, ring B is a substituted or unsubstituted phenylene, or a substituted or unsubstituted monocyclic heteroarylene containing 0 or 1 O atoms, 0 or 1 S atoms, 0-3 N atoms, and at least 2 carbon atoms in the ring, where if ring B is substituted then each substituent on ring B is H or R^B.


15 [00248] In one aspect, ring B is a unsubstituted or substituted divalent ring selected from phenylene, naphthylene, furanylene, pyrrolylene, oxazolylene, thiazolylene, imidazolylene, pyrazolylene, triazolylene, tetrazolylene, isoxazolylene, isothiazolylene, oxadiazolylene, thiadiazolylene, pyridinylene, pyrimidinylene, pyrazinylene, pyridazinylene, triazinylene, quinolinylene, isoquinolinylene, quinazolinylene, quinoxalinylene, naphthyridinylene, 20 indolylene, indazolylene, benzoxazolylene, benzisoxazolylene, benzofuranylene, benzothienylene, benzothiazolylene, benzimidazolylene, purinylene, cinnolinylene, phthalazinylene, pteridinylene, pyridopyrimidinylene, pyrazolopyrimidinylene, azaindolylene, cyclopentylene, cyclopentenylene, cycloalkylene, and cyclalkenylene.

[00249] In one aspect, ring B is a unsubstituted or substituted divalent ring selected from phenylene, furanylene, pyrrolylene, oxazolylene, thiazolylene, imidazolylene, pyrazolylene, triazolylene, tetrazolylene, isoxazolylene, isothiazolylene, oxadiazolylene, thiadiazolylene, pyridinylene, pyrimidinylene, pyrazinylene, pyridazinylene, triazinylene, cyclopentylene, cyclopentenylene, cycloalkylene, and cyclalkenylene.


[00250] In one aspect, ring B is a unsubstituted or substituted divalent ring selected from pyrrolylene, oxazolylene, thiazolylene, imidazolylene, pyrazolylene, triazolylene, tetrazolylene, isoxazolylene, isothiazolylene, oxadiazolylene, thiadiazolylene, pyridinylene, pyrimidinylene, pyrazinylene, pyridazinylene, triazinylene, cyclopentylene, cyclopentenylene, cycloalkylene, and cyclalkenylene.



[00252] In some embodiments, R⁸ is -CH₃ or -CF₃. In other embodiments, R⁸ is -CH₃.



[00253] In some embodiments, R⁴ is . In other embodiments, R⁴ is



[00254] In some embodiments, CY is a substituted or unsubstituted phenyl, wherein if CY is

5 substituted then each substituent on CY is H or R^C; each R^C is independently selected from H, halogen, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[00255] In some embodiments, each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OCF₃, -OCH₃.

[00256] In some embodiments, CY is phenyl, 2-fluorophenyl or 2-chloro-phenyl. In some 10 embodiments, CY is phenyl.



[00257] In one aspect, provided is a 3-substituted-{4-[1-(2-R^C-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzene compound, wherein 3-substituent is -L²-B-L¹-R¹, or pharmaceutically acceptable salts, pharmaceutically acceptable solvates, or prodrugs thereof; R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -C(=O)NH-OH, -C(=O)NH-CN, 15 tetrazolyl or carboxylic acid bioisostere; R^D is H or C₁-C₆alkyl; B is a divalent group selected from a substituted or unsubstituted cycloalkylene, a substituted or unsubstituted heterocycloalkylene, a substituted or unsubstituted arylene, and a substituted or unsubstituted heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B; each R^B is selected from H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-20 C₄alkoxy, and C₁-C₄heteroalkyl; L¹ is absent, a substituted or unsubstituted alkylene, a substituted or unsubstituted fluoroalkylene, or a substituted or unsubstituted heteroalkylene; L² is absent, a substituted or unsubstituted alkylene, a substituted or unsubstituted fluoroalkylene, a substituted or unsubstituted heteroalkylene, -O-, -S-, -SO-, -SO₂-, -NR²-, or -C(=O)-; R² is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl; R^C is selected from H, halogen, -OH, C₁-C₄alkyl, C₁-25 C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.



[00258] In one aspect, provided is a 4-substituted-{4-[1-(2-R^C-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzene compound, wherein 4-substituent is -L²-B-L¹-R¹, or pharmaceutically acceptable salts, pharmaceutically acceptable solvates, or prodrugs thereof; R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -C(=O)NH-OH, -C(=O)NH-CN, 30 tetrazolyl or carboxylic acid bioisostere; R^D is H or C₁-C₆alkyl; B is a divalent group selected from a substituted or unsubstituted cycloalkylene, a substituted or unsubstituted heterocycloalkylene, a substituted or unsubstituted arylene, and a substituted or unsubstituted heteroarylene, a substituted or unsubstituted



heteroarylene, where if ring B is substituted then each substituent on ring B is H or R^B; each R^B is selected from H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl; L¹ is absent, a substituted or unsubstituted alkylene, a substituted or unsubstituted fluoroalkylene, or a substituted or unsubstituted heteroalkylene; L² is absent, a substituted or unsubstituted alkylene, a substituted or unsubstituted fluoroalkylene, a substituted or unsubstituted heteroalkylene, -O-, -S-, -SO-, -SO₂-, -NR²-, or -C(=O)-; R² is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl; R^C is selected from H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

[00259] In some embodiments, R^C is H, F, Cl, -CH₃, or CF₃. In some other embodiments, R^C is F or Cl. In other embodiments, R^C is Cl. In yet other embodiments, R^C is F.


[00260] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂- or -

CH₂CH₂CH₂-; L² is absent; ring B is  or ; each R^C is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃.

[00261] In some embodiments, ring B is  or . In some embodiments, ring

15 B is . In some embodiments, ring B is .

[00262] In some embodiments, the compound of Formula (I) has the structure of Formula (III):



Formula (III)

wherein

20 R¹ is -CO₂H, -CO₂R^D, -CN, tetrazolyl, -C(=O)N(R⁹)₂, -C(=O)NHCH₂CH₂SO₃H, or -C(=O)NHSO₂R¹⁰; R^D is H or C₁-C₄alkyl; L¹ is absent or C₁-C₆alkylene;

R³ is H, C₁-C₄alkyl, C₃-C₆cycloalkyl, or C₁-C₄fluoroalkyl;

R⁷ is H or C₁-C₄alkyl;

R⁸ is H, C₁-C₄alkyl, or C₁-C₄fluoroalkyl;

R⁹ is a H, C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₃-C₆cycloalkyl, or a substituted or unsubstituted phenyl;

R¹⁰ is a C₁-C₆alkyl, C₁-C₆fluoroalkyl, C₃-C₆cycloalkyl, or a substituted or unsubstituted phenyl;

each of R^A, R^B, and R^C are independently selected from H, F, Cl, Br, I, -CN, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl;


m is 0, 1, or 2; n is 0, 1, or 2; p is 0, 1, or 2.

[00263] In some embodiments, R¹ is -CO₂H, -CO₂R^D or -C(=O)NHSO₂R¹⁰; R³ is C₁-C₄alkyl; R⁷ is H; R¹⁰ is a C₁-C₆alkyl or a substituted or unsubstituted phenyl; each R^A is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; each R^B is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; each R^C is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; m is 0 or 1; p is 0 or 1.

[00264] In some embodiments, R¹ is -CO₂H or -CO₂R^D; R³ is -CH₃ or -CH₂CH₃; R⁸ is H, -CH₃ or -CF₃; R^D is H, -CH₃, or -CH₂CH₃.

[00265] In some embodiments, R¹ is -CO₂H; R³ is -CH₃; R⁸ is -CH₃; L¹ is absent, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, -CH₂CH(CH₃)-, or -CH₂C(CH₃)₂-.


[00266] In some embodiments, the compound of Formula (III) has the following structure:



[00267] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, or -C(CH₂CH₃)₂-, each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; m is 0; n is 0, 1, or 2; p is 0.

[00268] In some embodiments, L¹ is absent or -CH₂-, each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, and -OH; n is 0 or 1.


[00269] In some embodiments, the compound of Formula (III) has the following structure:




[00270] In some embodiments, L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, or -C(CH₂CH₃)₂-, each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃; m is 0; n is 0, 1, or 2; p is 0.

5 [00271] In some embodiments, L¹ is absent or -CH₂-; each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, and -OH; n is 0 or 1.


[00272] In some embodiments, R¹ is -C(=O)NHSO₂R¹⁰, R³ is -CH₃ or -CH₂CH₃; R⁸ is H, -CH₃ or -CF₃; R¹⁰ is -CH₃, or -CH₂CH₃.




[00273] In some embodiments,



10 [00274] In some embodiments,




[00275] In some embodiments,



[00276] In some embodiments,

[00277] In one aspect, the compound of Formula (I) has a structure selected from:




[00278] In one aspect, B is as described in any one of Tables 1 through 6. In one aspect, CY is as described in any one of Tables 1 through 6. In one aspect, L^2 is as described in any one of Tables 1 through 6. In one aspect, R^8 is as described in any one of Tables 1 through 6. In one aspect, R^1 is as described in any one of Tables 1 through 7. In one aspect, R^D is as described in Table 1. In one aspect, R^C is as described in any one of Tables 3 through 7.

[00279] Any combination of the groups described above for the various variables is contemplated herein. Throughout the specification, groups and substituents thereof are chosen by one skilled in the field to provide stable moieties and compounds.

[00280] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) include, but are not limited to, those described in Tables 1 through 6:

Table 1:



Cmpd #	A ¹	A ²	Position of L ²	-L ² -	B	-L ¹ -	R ^D	CY	R ⁸
1-1	CH	CH	4	-CH ₂ -	1 <i>H</i> -[1,2,4]Triazol-1,3-ylene	-	H	2-Chlorophenyl	CH ₃
1-2	CH	CH	4	-CH ₂ -	1 <i>H</i> -Imidazol-1,4-ylene	-	H	2-Chlorophenyl	CH ₃
1-3	CH	CH	4	-CH ₂ -	1 <i>H</i> -Imidazol-1,5-ylene	-	H	2-Chlorophenyl	CH ₃
1-4	CH	CH	4	-CH ₂ -	5-Trifluoromethyl-1-1 <i>H</i> -pyrazol-1,4-ylene	-	H	2-Chlorophenyl	CH ₃
1-5	CH	CH	4	-CH ₂ -	Phen-1,3-ylene	-	H	2-Chlorophenyl	CH ₃
1-6	CH	CH	4	-CH ₂ -	Phen-1,3-ylene	-	H	2-Bromo-phenyl	H
1-7	CH	CH	4	-CH ₂ -	2-Methoxyphen-1,5-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-9	CH	CH	4	CH ₂	1 <i>H</i> -Pyrazol-1,4-ylene	-	H	2-Chlorophenyl	CH ₃
1-11	CH	CH	4	-	Phen-1,3-ylene	-	H	Pyridin-2-yl	(R)-CH ₃
1-12	CH	CH	4	CH ₂ O-	Phen-1,3-ylene	-	H	2-Chlorophenyl	CH ₃
1-13	CH	CH	4	-	Phen-1,3-ylene	-	H	6-Methylpyridin-2-yl	H
1-14	CH	CH	4	-CH ₂ -	2-Chlorophen-1,5-ylene	-	H	2-Chlorophenyl	CH ₃
1-15	CH	CH	4	-CH ₂ -	Thiophen-2,5-yl	-	H	2-Chlorophenyl	CH ₃
1-16	CH	CH	4	-	Phen-1,3-ylene	-	H	Indan-1-yl	
1-17	CH	CH	4	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-18	CH	CH	4	-CH ₂ S-	Phen-1,3-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-19	CH	CH	4	-CH ₂ S-	Phen-1,2-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-20	CH	CH	4	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-21	CH	CH	4	-CH ₂ -	Phen-1,3-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-22	CH	CH	4	-CH ₂ -	Phen-1,2-ylene	-CH ₂ -	H	2-Chlorophenyl	CH ₃
1-23	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Trifluoromethyl-phenyl	CH ₃
1-24	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	3-Trifluoromethyl-phenyl	CH ₃
1-25	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2,4-Dichlorophenyl	CH ₃
1-26	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluorophenyl	CH ₃
1-27	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	3-Bromo-	CH ₃

Cmpd #	A ¹	A ²	Position of L ²	-L ² -	B	-L ¹ -	R ^D	CY	R ⁸
								phenyl	
1-28	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Methoxy-phenyl	CH ₃
1-29	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	Benzyl	CH ₃
1-30	CH	CH	3	-	2-Methoxyphen-1,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-31	CH	CH	3	-	Phen-1,3-ylene	-	H	2-Chloro-phenyl	CH ₃
1-32	CH	CH	3	-	Phen-1,2-ylene	-	H	2-Chloro-phenyl	CH ₃
1-34	CH	CH	3	-	Pyridin-3,6-ylene	-	H	2-Chloro-phenyl	CH ₃
1-35	CH	CH	3	-	Pyridin-2,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-36	CH	CH	3	-	Pyridin-2,6-ylene	-	H	2-Chloro-phenyl	CH ₃
1-37	CH	CH	3	-	Pyridin-3,5-ylene	-	H	2-Chloro-phenyl	CH ₃
1-38	CH	CH	3	-	2-Methoxypyridin-3,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-39	CH	CH	3	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-40	CH	CH	3	-	Indol-5,1-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-41	CH	CH	3	-	Pyridin-3,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-42	CH	CH	3	-	2-Fluorophen-1,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-43	CH	CH	3	-	4-Fluorophen-1,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-44	CH	CH	3	-	Phen-1,2-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-45	CH	CH	3	-	Cyclohex-1-en-1,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-46	CH	CH	3	-	Cyclohex-1,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-47	N	CH	4	-	Phen-1,4-ylene	-CH ₂ -	CH ₂ CH ₃	2-Fluoro-phenyl	CH ₃
1-48	N	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-phenyl	CH ₃
1-49	CH	CH	4	-	2-Methoxyphen-1,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-50	CH	CH	4	-	Pyridin-2,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-51	CH	CH	4	-	Phen-1,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-52	CH	CH	4	-	Phen-1,2-ylene	-	H	2-Chloro-phenyl	CH ₃
1-53	CH	CH	4	-	Pyridin-2,6-ylene	-	H	2-Chloro-phenyl	CH ₃
1-54	CH	CH	4	-	Pyridin-2,5-	-	H	2-Chloro-	CH ₃

Cmpd #	A ¹	A ²	Position of L ²	-L ² -	B	-L ¹ -	R ^D	CY	R ⁸
					ylene			phenyl	
1-55	CH	CH	4	-	Pyridin-3,5-ylene	-	H	2-Chloro-phenyl	CH ₃
1-56	CH	CH	4	-	Pyridin-3,6-ylene	-	H	2-Chloro-phenyl	CH ₃
1-57	CH	CH	4	-	Phen-1,2-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-58	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-59	CH	CH	4	-	Phen-1,3-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-60	CH	CH	3	-	Phen-1,3-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-61	CH	CH	4	-	Pyridin-2,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-62	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-63	CH	CH	4	-	Pyridin-3,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-64	CH	CH	4	-	2-Fluorophen-1,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-65	CH	CH	4	-	4-Fluorophen-1,5-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-66	CH	CH	4	-	Cyclohex-1-en-1,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-67	CH	CH	4	-	Cyclohex-1,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-69	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	CH ₃	2-Chloro-phenyl	CH ₃
1-70	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	CH ₂ CH ₃	2-Chloro-phenyl	CH ₃
1-71	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	H	2-Chloro-phenyl	CH ₃
1-72	CH	CH	4	-	Phen-1,4-ylene	-C(CH ₃) ₂ -	H	2-Chloro-phenyl	CH ₃
1-73	CH	N	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-74	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃) ₂ -	H	2-Fluoro-phenyl	CH ₃
1-75	CH	CH	4	-	Phen-1,4-ylene	CH ₂ CH ₂ C _{H₂} -	H	2-Chloro-phenyl	CH ₃
1-76	CH	CH	4	-	Phen-1,3-ylene	-	H	2-Chloro-phenyl	CH ₃
1-77	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	4-chloro-2-Fluoro-phenyl	CH ₃
1-78	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-79	CH	CH	4	-CH ₂ O-	Phen-1,2-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-80	CH	CH	4	-CH ₂ -	Phen-1,3-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-81	CH	CH	4	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-	(R)-CH ₃

Cmpd #	A ¹	A ²	Position of L ²	-L ² -	B	-L ¹ -	R ^D	CY	R ⁸
								phenyl	
1-82	CH	CH	4	-CH ₂ -	2-Fluorophen-1,5-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-83	CH	CH	4	-O-	Phen-1,4-ylene	-CH ₂ -	CH ₂ CH ₃	2-Fluoro-phenyl	(R)-CH ₃
1-84	CH	CH	4	-O-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-85	C(CH ₃)	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-86	CH	CH	4	-	Phen-1,4-ylene	-C(CH ₃) ₂ -	H	2-Fluoro-phenyl	CH ₃
1-87	CH	CH	4	-	Pyridin-2,5-ylene	-C(CH ₃) ₂ -	H	2-Fluoro-phenyl	CH ₃
1-88	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-89	CH	CH	4	-	Phen-1,4-ylene	-C(CH ₃) ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-90	CH	CH	4	-	Phen-1,4-ylene	-C(CH ₃) ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-91	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	H	2-Fluoro-phenyl	(R)-CH ₃
1-92	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	H	2-Chloro-phenyl	(R)-CH ₃
1-93	CH	CH	4	-	Thiazol-2,4-ylene	-	H	2-Chloro-phenyl	CH ₃
1-94	CH	CH	4	-	Thiazol-2,5-ylene	-	H	2-Chloro-phenyl	CH ₃
1-95	CH	CH	4	-O-	Phen-1,3-ylene	-CH ₂ -	CH ₃	2-Fluoro-phenyl	CH ₃
1-96	CH	CH	4	-O-	Phen-1,3-ylene	-CH ₂ -	H	2-Fluoro-phenyl	CH ₃
1-97	CH	CH	4	-CH ₂ SO ₂ -	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-98	CH	CH	4	-CH ₂ SO ₂ -	Phen-1,3-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-99	CH	CH	4	-CH ₂ SO ₂ -	Phen-1,2-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-100	CH	CH	4	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	CH ₃
1-101	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2,6-Dichloro-phenyl	CH ₃
1-102	CH	CH	4	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	-CH ₃	2-Chloro-phenyl	(R)-CH ₃
1-103	CH	CH	4	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-104	CH	CH	4	-OCH ₂ -	Phen-1,4-ylene	-	H	2-Fluoro-phenyl	(R)-CH ₃
1-105	CH	CH	4	-S-	Phen-1,4-ylene	-CH ₂ -	CH ₂ CH ₃	2-Fluoro-phenyl	(R)-CH ₃
1-106	CH	CH	4	-S-	Phen-1,3-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-107	CH	CH	4	-S-	Phen-1,2-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃

Cmpd #	A ¹	A ²	Position of L ²	-L ² -	B	-L ¹ -	R ^D	CY	R ⁸
1-108	CH	CH	4	-S-	Pyridin-2,5-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-109	CH	CH	4	-S-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-110	C(CH ₃)	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	H	2-Fluoro-phenyl	(R)-CH ₃
1-111	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(S)-CH ₃
1-112	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(S)-CH ₃
1-113	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	H
1-114	CH	CH	4	-CH ₂ O-	Phen-1,2-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-115	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	Phenyl	(R)-CH ₃
1-116	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2,3-Difluoro-phenyl	CH ₃
1-117	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2,4-Difluoro-phenyl	CH ₃
1-118	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Fluoro-4-methoxy-phenyl	CH ₃
1-119	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2,5-Difluoro-phenyl	CH ₃
1-120	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2,6-Difluoro-phenyl	CH ₃
1-121	CH	CH	4	-	Phen-1,3-ylene	-CH ₂ -	H	Phenyl	(R)-CH ₃
1-122	CH	CH	4	-	Phen-1,4-ylene	-	H	Phenyl	(R)-CH ₃
1-123	CH	CH	4	-	Phen-1,2-ylene	-CH ₂ -	H	Phenyl	(R)-CH ₃
1-124	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Methyl-phenyl	(R)-CH ₃
1-125	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	-CH ₃	-CH ₃
1-126	CH	N	4	-	Pyridin-2,5-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-127	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	H	2-Chloro-phenyl	(R)-CH ₃
1-128	CH	CH	4	-	Phen-1,4-ylene	-CH(CH ₃)-	H	2-Chloro-phenyl	(R)-CH ₃
1-129	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-130	CH	CH	4	-	Phen-1,4-ylene	-CH-(CH ₂ CH ₃)-	H	2-Chloro-phenyl	(R)-CH ₃
1-131	C(Cl)	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-132	C(F)	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-133	CH	CH	4	-	Phen-1,4-ylene	-	H	2-Chloro-phenyl	(R)-CH ₃
1-134	CH	CH	4	-	Tetrazol-2,5-ylene	-CH ₂ -	CH ₃	2-Fluoro-phenyl	(R)-CH ₃

Cmpd #	A ¹	A ²	Position of L ²	-L ² -	B	-L ¹ -	R ^D	CY	R ⁸
1-135	CH	CH	4	-	Tetrazol-2,5-ylene	-CH ₂ -	H	2-Fluoro-phenyl	(R)-CH ₃
1-136	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	t-butyl	
1-137	CH	CH	4	-O-	Phen-1,3-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-138	CH	CH	4	-O-	Phen-1,4-ylene	-CH ₂ -	H	2-Chloro-phenyl	(R)-CH ₃
1-139	CH	CH	4	-	Phen-1,4-ylene	-CH(OH)-	H	2-Fluoro-phenyl	(R)-CH ₃
1-140	CH	CH	4	-CH ₂ -	Azetidin-1,3-ylene	-	H	Phenyl	(R)-CH ₃
1-141	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	3,5-Dibromo-phenyl	(R)-CH ₃
1-142	N	CH	4	-	Phen-1,4-ylene	-CH-(CH ₂ CH ₃)-	H	Phenyl	(R)-CH ₃
1-143	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	Phenyl	(S)-CH ₃
1-144	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	3-Hydroxy-phenyl	(R)-CH ₃
1-145	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	Phenyl	CH ₃
1-146	CH	CH	4	-	Phen-1,4-ylene	-CH ₂ -	H	Phenyl-d5	CD ₃

[00281] Compounds in Table 1 are named:

1-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-1H-[1,2,4]triazole-3-carboxylic acid* (Compound 1-1); 1-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-1H-imidazole-4-carboxylic acid* (Compound 1-2); 3-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-3H-imidazole-4-carboxylic acid* (Compound 1-3); 1-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-5-trifluoromethyl-1H-pyrazole-4-carboxylic acid* (Compound 1-4); 3-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-benzoic acid* (Compound 1-5); 3-*{4-[4-(2-Bromo-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl]-benzyl}-benzoic acid* (Compound 1-6); [3-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-4-methoxy-phenyl]-acetic acid (Compound 1-7); 1-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-1H-pyrazole-4-carboxylic acid* (Compound 1-9); 4'-*[3-Methyl-4-((R)-1-pyridin-2-yl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid* (Compound 1-11); 3-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-benzoic acid* (Compound 1-12); 4'-*[3-Methyl-4-(6-methyl-pyridin-2-ylmethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid* (Compound 1-13); 4-Chloro-3-*(4-[4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-benzoic acid* (Compound 1-14); 5-*(4-[4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-benzyl)-thiophene-2-carboxylic acid* (Compound 1-15); 4'-*[4-(Indan-1-**

ylloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-16); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-17); [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-18); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-19); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-20); [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-21); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-22); (4'-{3-Methyl-4-[1-(2-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-23); (4'-{3-Methyl-4-[1-(3-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-24); (4'-{4-[1-(2,4-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-25); (4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-26); (4'-{4-[1-(3-Bromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-27); (4'-{4-[1-(2-Methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-28); {4'-[3-Methyl-4-(1-methyl-2-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-29); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid (Compound 1-30); 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid (Compound 1-31); 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid (Compound 1-32); 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-34); 2-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid (Compound 1-35); 6-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-36); 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-37); [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid (Compound 1-38); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-39); [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-indol-1-yl]-acetic acid (Compound 1-40); [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-41); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-


3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-42); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-4-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-43); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-yl)-acetic acid (Compound 1-44); 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid (Compound 1-45); 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid (Compound 1-46); [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid ethyl ester (Compound 1-47); [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid (Compound 1-48); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-methoxy-biphenyl-3-yl)-acetic acid (Compound 1-49); 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid (Compound 1-50); 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid (Compound 1-51); 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid (Compound 1-52); 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-53); 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-54); 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-55); 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-56); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-yl)-acetic acid (Compound 1-57); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-58); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-yl)-acetic acid (Compound 1-59); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-yl)-acetic acid (Compound 1-60); [6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-61); 3-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-62); [5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-63); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-64); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-4-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-65); 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid (Compound 1-66); 4-(4-{4-[1-(2-Chloro-phenyl)-

ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid (Compound 1-67); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid methyl ester (Compound 1-69); 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester (Compound 1-70); 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-71); 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-72); [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-3-yl)-phenyl]-acetic acid (Compound 1-73); 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-74); 4-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid (Compound 1-75); 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid (Compound 1-76); (4'-{4-[1-(4-Chloro-2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-77); (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-78); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-79); [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-80); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-81); [4-Fluoro-3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-82); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid ethyl ester (Compound 1-83); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-84); (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-acetic acid (Compound 1-85); 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-86); 2-[6-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-2-methyl-propionic acid (Compound 1-87); (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-88); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-89); 2-(4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-90); 2-(4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-91); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-

ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-92); 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-4-carboxylic acid (Compound 1-93); 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-5-carboxylic acid (Compound 1-94); [3-(4-{4-[1-(2-5 Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester (Compound 1-95); [3-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-96); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-97); [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-10 yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-98); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-99); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-100); (4'-{4-[1-(2,6-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-101); [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester (Compound 1-102); [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-103); 4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxyethyl)-benzoic acid (Compound 1-104); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester (Compound 1-105); [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-106); [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-107); [6-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-pyridin-3-yl]-acetic acid (Compound 1-108); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-109); 2-(4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-propionic acid (Compound 1-110); (4'-{4-[(S)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-111); (4'-{4-[(S)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-112); {4'-[4-(2-Chloro-benzyloxy carbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-113); [2-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-114); {4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-115); (4'-{4-[1-(2,3-Difluoro-phenyl)-

ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-116); (4'-{4-[1-(2,4-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-117); (4'-{4-[1-(2-Fluoro-4-methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-118); (4'-{4-[1-(2,5-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-119); (4'-{4-[1-(2,6-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-120); {4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-yl}-acetic acid (Compound 1-121); 4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-carboxylic acid (Compound 1-122); {4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-2-yl}-acetic acid (Compound 1-123); {4'-[3-Methyl-4-((R)-1-o-tolyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-124); 4'-{(4-Isopropoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid (Compound 1-125); (6'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-[2,3']bipyridinyl-5-yl)-acetic acid (Compound 1-126); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-127); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-128); (3'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-129); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid (Compound 1-130); (2'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-131); (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-fluorobiphenyl-4-yl)-acetic acid (Compound 1-132); 4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid (Compound 1-133); [5-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid methyl ester (Compound 1-134); [5-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid (Compound 1-135); [4'-(4-*tert*-Butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid (Compound 1-136); [3-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-137); [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-138); (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-hydroxy-acetic acid (Compound 1-139); 1-{4-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-benzyl}-azetidine-3-carboxylic acid (Compound 1-140);

(4'-{4-[*(R*)-1-(3,5-Dibromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-141); 2-(4-{5-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-pyridin-2-yl}-phenyl)-butyric acid (Compound 1-142); {4'-[3-Methyl-4-((*S*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-143); (4'-{4-[*(R*)-1-(3-Hydroxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-144); {4'-[3-Methyl-4-(1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-145); {4'-[3-Methyl-4-(1-phenyl-ethoxy-d9-carbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-146).


10 **Table 2:**

Cmpd #	L ²	B	L ¹	R ¹	CY	R ⁸
2-1	-CH ₂ -	Phen-1,4-ylene	-	-C(=O)NH-(cyclopropyl)	2-Chloro-phenyl	CH ₃
2-2	-	Phen-1,4-ylene	-CH ₂ -	-CN	2-Chloro-phenyl	(<i>R</i>)-CH ₃
2-3	-	Phen-1,4-ylene	-CH ₂ -	-CN	2-Fluoro-phenyl	(<i>R</i>)-CH ₃
2-4	-	Phen-1,4-ylene	-CH ₂ -	2H-Tetrazol-5-yl	2-Fluoro-phenyl	(<i>R</i>)-CH ₃
2-5	-	Phen-1,4-ylene	-CH ₂ -	2H-Tetrazol-5-yl	2-Chloro-phenyl	(<i>R</i>)-CH ₃
2-6	-	Phen-1,4-ylene	-CH ₂ -	-C(=NH)-NH ₂	2-Fluoro-phenyl	(<i>R</i>)-CH ₃
2-7	-	Phen-1,4-ylene	-CH ₂ -	-C(=NH)-NH-Ac	2-Fluoro-phenyl	(<i>R</i>)-CH ₃
2-8	-	Phen-1,4-ylene	-CH ₂ -	-C(=O)NH-CH ₂ CH ₂ SO ₃ H	phenyl	(<i>R</i>)-CH ₃

[00282] Compounds in Table 2 are named:

15 {5-[4-(4-Cyclopropylcarbamoyl-benzyl)-phenyl]-3-methyl-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (Compound 2-1); [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (*R*)-1-(2-chloro-phenyl)-ethyl ester (Compound 2-2); [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (*R*)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-3); {3-Methyl-5-[4'-(2*H*-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (*R*)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-4); {3-Methyl-5-[4'-(2*H*-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (*R*)-1-(2-chloro-phenyl)-ethyl ester (Compound 2-5); [5-(4'-Carbamimidoylmethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (*R*)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-6); {5-[4'-(2-Acetylamino-2-imino-ethyl)-biphenyl-4-yl]-3-methyl-isoxazol-4-yl}-carbamic acid (*R*)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-7); 2-(2-{4'-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetylamino)-ethanesulfonic acid (Compound 2-8).

[00283] Table 3:




Cmpd #	A	-L ² -	B	-L ¹ -	R ^D	R ^C	R ⁸
3-1	Naphthalen-2,6-ylene	-	Phen-1,4-ylene	-CH ₂ -	H	Cl	CH ₃
3-2	Naphthalen-2,6-ylene	-	Phen-1,4-ylene	-CH ₂ SCH ₂ CH ₂ -	H	Cl	CH ₃
3-3	Thiazol-4,2-ylene	-	Phen-1,4-ylene	-CH ₂ -	H	F	(R)-CH ₃
3-4	Thiazol-5,2-ylene	-	Phen-1,4-ylene	-CH ₂ -	H	F	(R)-CH ₃
3-5	Thiazol-4,2-ylene	-	Phen-1,2-ylene	-CH ₂ -	H	F	(R)-CH ₃
3-6	Thiazol-4,2-ylene	-	Phen-1,3-ylene	-CH ₂ -	H	F	(R)-CH ₃

[00284] Compounds in Table 3 are named:

5 [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-phenyl]-acetic acid (Compound 3-1); 3-[4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-benzylsulfanyl]-propionic acid (Compound 3-2); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-3); [4-(5-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-4); [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-5); and [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-6).


Table 4:



15


Cmpd #	Position of L ²	-L ² -	B (see note a)	-L ¹ -	R ^C
4-1	3	-	Phen-1,4-ylene	-CH ₂ -	Cl
4-2	4	-	Phen-1,4-ylene	-CH ₂ -	Cl
4-3	3	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
4-4	4	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
4-5	3	-O-	Phen-1,4-ylene	-CH ₂ -	Cl
4-6	4	-O-	Phen-1,4-ylene	-CH ₂ -	Cl

4-7	3	-S-	Phen-1,4-ylene	-CH ₂ -	Cl
4-8	4	-S-	Phen-1,4-ylene	-CH ₂ -	Cl
4-9	3	-NH-	Phen-1,4-ylene	-CH ₂ -	Cl
4-10	4	-NH-	Phen-1,4-ylene	-CH ₂ -	Cl
4-11	3	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	Cl
4-12	4	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	Cl
4-13	3	-OCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
4-14	4	-OCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
4-15	3	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	Cl
4-16	4	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	Cl
4-17	3	-CH ₂ NH-	Phen-1,4-ylene	-CH ₂ -	Cl
4-18	4	-CH ₂ NH-	Phen-1,4-ylene	-CH ₂ -	Cl
4-19	3	-NHCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
4-20	4	-NHCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl

Table 5:

Cmpd #	Position of L ²	-L ² -	B (see note a)	-L ¹ -	R ^C
5-1	3	-	Phen-1,4-ylene	-CH ₂ -	Cl
5-2	4	-	Phen-1,4-ylene	-CH ₂ -	Cl
5-3	3	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
5-4	4	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
5-5	3	-O-	Phen-1,4-ylene	-CH ₂ -	Cl
5-6	4	-O-	Phen-1,4-ylene	-CH ₂ -	Cl
5-7	3	-S-	Phen-1,4-ylene	-CH ₂ -	Cl
5-8	4	-S-	Phen-1,4-ylene	-CH ₂ -	Cl
5-9	3	-NH-	Phen-1,4-ylene	-CH ₂ -	Cl
5-10	4	-NH-	Phen-1,4-ylene	-CH ₂ -	Cl
5-11	3	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	Cl
5-12	4	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	Cl
5-13	3	-OCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
5-14	4	-OCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
5-15	3	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	Cl
5-16	4	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	Cl
5-17	3	-CH ₂ NH-	Phen-1,4-ylene	-CH ₂ -	Cl
5-18	4	-CH ₂ NH-	Phen-1,4-ylene	-CH ₂ -	Cl
5-19	3	-NHCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
5-20	4	-NHCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl

Table 6:



Cmpd #	Position of L ²	-L ² -	B (see note a)	-L ¹ -	R ^c
6-1	3	-	Phen-1,4-ylene	-CH ₂ -	Cl
6-2	4	-	Phen-1,4-ylene	-CH ₂ -	Cl
6-3	3	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
6-4	4	-CH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
6-5	3	-O-	Phen-1,4-ylene	-CH ₂ -	Cl
6-6	4	-O-	Phen-1,4-ylene	-CH ₂ -	Cl
6-7	3	-S-	Phen-1,4-ylene	-CH ₂ -	Cl
6-8	4	-S-	Phen-1,4-ylene	-CH ₂ -	Cl
6-9	3	-NH-	Phen-1,4-ylene	-CH ₂ -	Cl
6-10	4	-NH-	Phen-1,4-ylene	-CH ₂ -	Cl
6-11	3	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	Cl
6-12	4	-CH ₂ O-	Phen-1,4-ylene	-CH ₂ -	Cl
6-13	3	-OCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
6-14	4	-OCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
6-15	3	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	Cl
6-16	4	-CH ₂ S-	Phen-1,4-ylene	-CH ₂ -	Cl
6-17	3	-CH ₂ NH-	Phen-1,4-ylene	-CH ₂ -	Cl
6-18	4	-CH ₂ NH-	Phen-1,4-ylene	-CH ₂ -	Cl
6-19	3	-NHCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl
6-20	4	-NHCH ₂ -	Phen-1,4-ylene	-CH ₂ -	Cl

[00285] In one aspect, Phen-1,4-ylene (i.e. ring B) in Tables 4 through 6 is replaced with a group selected from Phen-1,2-ylene, Phen-1,3-ylene, Pyridin-2,3-ylene, Pyridin-2,4-ylene, Pyridin-2,5-

5 ylene, Pyridin-2,6-ylene, Pyridin-3,4-ylene, Pyridin-3,5-ylene, Pyridin-3,6-ylene, Pyridin-4,5-ylene, Pyridin-4,6-ylene, Pyridin-5,6-ylene, Pyrimidin-2,4-ylene, Pyrimidin-2,5-ylene, Pyrimidin-4,5-ylene, Pyrimidin-4,6-ylene, Pyrazin-2,3-ylene, Pyrazin-2,5-ylene, Pyrazin-2,6-ylene, Pyridazin-3,4-ylene, Pyridazin-3,5-ylene, Pyridazin-3,6-ylene, Pyridazin-4,5-ylene, 1H-Imidazol-1,2-ylene, 1H-Imidazol-1,4-ylene, 1H-Imidazol-1,5-ylene, 1H-Pyrazol-1,3-ylene, 1H-
10 Pyrazol-1,4-ylene, 1H-Pyrazol-1,5-ylene, 1H-[1,2,3]Triazol-1,4-ylene, 1H-[1,2,3]Triazol-1,5-ylene, 1H-[1,2,4]Triazol-1,3-ylene, and 1H-[1,2,4]Triazol-1,5-ylene.

[00286] In one aspect, Phen-1,4-ylene (i.e. ring B) in Tables 4 through 6 is replaced with a group selected from Phen-1,2-ylene, Phen-1,3-ylene, Pyridin-2,3-ylene, Pyridin-2,4-ylene, Pyridin-2,5-

15 ylene, Pyridin-2,6-ylene, Pyridin-3,4-ylene, Pyridin-3,5-ylene, Pyridin-3,6-ylene, Pyridin-4,5-ylene, Pyridin-4,6-ylene, Pyridin-5,6-ylene, Pyrimidin-2,4-ylene, Pyrimidin-2,5-ylene, Pyrimidin-4,5-ylene, Pyrimidin-4,6-ylene, Pyrazin-2,3-ylene, Pyrazin-2,5-ylene, Pyrazin-2,6-

ylene, Pyridazin-3,4-ylene, Pyridazin-3,5-ylene, Pyridazin-3,6-ylene, Pyridazin-4,5-ylene, 1*H*-Imidazol-1,2-ylene, 1*H*-Imidazol-1,4-ylene, 1*H*-Imidazol-1,5-ylene, 1*H*-Pyrazol-1,3-ylene, 1*H*-Pyrazol-1,4-ylene, 1*H*-Pyrazol-1,5-ylene, 1*H*-[1,2,3]Triazol-1,4-ylene, 1*H*-[1,2,3]Triazol-1,5-ylene, 1*H*-[1,2,4]Triazol-1,3-ylene, and 1*H*-[1,2,4]Triazol-1,5-ylene; and Cl (i.e. R^C) is replaced with F.

[00287] In one aspect, Cl (i.e. R^C) in Tables 4 through 6 is replaced with F.

Table 7:

Cmpd #	R ¹	R ^C	R ⁸
7-1	-C(=O)NHSO ₂ Me	H	(R)-CH ₃
7-2	-C(=O)NHSO ₂ Me	H	H
7-3	-C(=O)NHSO ₂ Me	2-F	(R)-CH ₃
7-4	-C(=O)NHSO ₂ Me	2-F	H
7-5	-C(=O)NHSO ₂ Me	2-Cl	(R)-CH ₃
7-6	-C(=O)NHSO ₂ Me	2-Cl	H
7-7	-C(=O)NHSO ₂ Me	2-CH ₃	(R)-CH ₃
7-8	-C(=O)NHSO ₂ Me	2-CH ₃	H
7-9	-C(=O)NHSO ₂ Me	2-CF ₃	(R)-CH ₃
7-10	-C(=O)NHSO ₂ Me	2-CF ₃	H
7-11	-C(=O)NHSO ₂ Me	3-F	(R)-CH ₃
7-12	-C(=O)NHSO ₂ Me	3-F	H
7-13	-C(=O)NHSO ₂ Me	3-Cl	(R)-CH ₃
7-14	-C(=O)NHSO ₂ Me	3-Cl	H
7-15	-C(=O)NHSO ₂ Me	3-CH ₃	(R)-CH ₃
7-16	-C(=O)NHSO ₂ Me	3-CH ₃	H
7-17	-C(=O)NHSO ₂ Me	3-CF ₃	(R)-CH ₃
7-18	-C(=O)NHSO ₂ Me	3-CF ₃	H
7-19	-C(=O)NHSO ₂ Ph	H	(R)-CH ₃
7-20	-C(=O)NHSO ₂ Ph	H	H
7-21	-C(=O)NHSO ₂ Ph	2-F	(R)-CH ₃
7-22	-C(=O)NHSO ₂ Ph	2-F	H
7-23	-C(=O)NHSO ₂ Ph	2-Cl	(R)-CH ₃
7-24	-C(=O)NHSO ₂ Ph	2-Cl	H
7-25	-C(=O)NHSO ₂ Ph	2-CH ₃	(R)-CH ₃
7-26	-C(=O)NHSO ₂ Ph	2-CH ₃	H
7-27	-C(=O)NHSO ₂ Ph	2-CF ₃	(R)-CH ₃
7-28	-C(=O)NHSO ₂ Ph	2-CF ₃	H
7-29	-C(=O)NHSO ₂ Ph	3-F	(R)-CH ₃
7-30	-C(=O)NHSO ₂ Ph	3-F	H

Cmpd #	R ¹	R ^C	R ⁸
7-31	-C(=O)NHSO ₂ Ph	3-Cl	(R)-CH ₃
7-32	-C(=O)NHSO ₂ Ph	3-Cl	H
7-33	-C(=O)NHSO ₂ Ph	3-CH ₃	(R)-CH ₃
7-34	-C(=O)NHSO ₂ Ph	3-CH ₃	H
7-35	-C(=O)NHSO ₂ Ph	3-CF ₃	(R)-CH ₃
7-36	-C(=O)NHSO ₂ Ph	3-CF ₃	H

Synthesis of Compounds

[00288] Compounds of Formula (I), Formula (II) and Formula (III) described herein are synthesized using standard synthetic techniques or using methods known in the art in

5 combination with methods described herein. In additions, solvents, temperatures and other reaction conditions presented herein may vary.

[00289] The starting material used for the synthesis of the compounds of Formula (I), Formula (II) and Formula (III) are either synthesized or obtained from commercial sources, such as, but not limited to, Sigma-Aldrich, Fluka, Acros Organics, Alfa Aesar, and the like. The compounds


10 described herein, and other related compounds having different substituents are synthesized using techniques and materials described herein or otherwise known, including those found in March, ADVANCED ORGANIC CHEMISTRY 4th Ed., (Wiley 1992); Carey and Sundberg, ADVANCED ORGANIC CHEMISTRY 4th Ed., Vols. A and B (Plenum 2000, 2001), and Green and Wuts,

PROTECTIVE GROUPS IN ORGANIC SYNTHESIS 3rd Ed., (Wiley 1999). General methods for the

15 preparation of compounds can be modified by the use of appropriate reagents and conditions for the introduction of the various moieties found in the formulae as provided herein.


[00290] In one aspect, the compounds of Formula (I), Formula (II) and Formula (III) are prepared as outlined in the following Scheme.

Scheme 1:



[00291] In one aspect, the synthesis of compounds of Formula (I), Formula (II) and Formula (III) described herein begins with the reaction of an alkyl acetoacetate with methylamine to provide a compound of structure **1-II**. Compounds of structure **1-II** are reacted with a substituted or

5 unsubstituted 4-halo-benzoyl chloride (structure **1-III**, where A^1 and A^2 are both CH) to provide compounds of structure **1-IV**. In one aspect, n is 0, 1, or 2. In one aspect, substituted napthylene carbonyl chlorides or substituted heteroaryl carbonyl chlorides are used instead of compounds of

10 structure **1-III**, resulting in the synthesis of compounds of structure **1-VIII**, wherein  is replaced with other monocyclic or bicyclic rings. In some embodiments, n is 0. In other


embodiments, n is 1.

[00292] Treatment of compounds of structure **1-IV** with hydroxyl amine and acetic acid provides isoxazoles of structure **1-V**. Hydrolysis of the ester group of isoxazoles of structure **1-V** provides carboxylic acids of structure **1-VI**. A Curtius rearrangement of carboxylic acids of structure **1-VI** in the presence of hydroxy compounds of structure **1-VII** provides carbamate compounds of

15 structure **1-VIII**.

[00293] In one aspect, ureas of structure **XIV** are prepared as outlined in Scheme 2.

Scheme 2




[00294] A Curtius rearrangement of carboxylic acids of structure **2-I** in the presence of tert-butanol provides carbamate compounds of structure **2-II**. Removal of the tert-butyl group with acid provides amines of structure **2-III**. Amines of structure **2-III** are reacted with isocyanates of structure **2-IV** to provide ureas of structure **2-V**. Ureas of structure **2-V** are used in the synthesis of compounds of Formula (I), Formula (II) and Formula (III) as described herein.

[00295] Common methods for the synthesis of isocyanates include the Curtius rearrangement of acyl azides and the Lossen rearrangement of hydroxamic acids. The synthesis of ureas include the following procedures: C. Han, J. A. Porco, Jr, *Org. Lett.*, 2007, 9, 1517-1520; H. Lebel, O. Leogane, *Org. Lett.*, 2006, 8, 5717-5720; M. B. Bertrand, J. P. Wolfe, *Tetrahedron*, 2005, 61, 6447-6459; M. McLaughlin, M. Palucki, I. W. Davies, *Org. Lett.*, 2006, 8, 3311-3314; J. A. Fritz, et al., *Org. Lett.*, 2006, 8, 2531-2534; L. Marinescu, et al., *J. Org. Chem.*, 2003, 68, 9453-9455; S.-H. Lee, et al., *Tetrahedron*, 2004, 60, 3439-3443.

[00296] In one aspect, amides of structure **3-III** are prepared as outlined in Scheme 3.

Scheme 3




[00297] Amines of structure **2-III** are reacted with acid chlorides of structure **3-II** to provide amides of structure **3-III**. Amides of structure **3-III** are used in the synthesis of compounds of Formula (I), Formula (II) and Formula (III) as described herein.

[00298] In some embodiments, amines of structure **2-III** are reacted with, but not limited to, a carboxylic acid ($CY-(CH(R^8))_nCO_2H$, where n is 0, 1, or 2) and coupling reagent such as EDC EDC, DCC, BOP, HATU or the like, or a carboxylic acid activated ester or an acid chloride (i.e. compound of structure **3-II**), in the presence of a suitable base and a suitable solvent to afford compounds of structure **3-III**.

[00299] In one aspect, metal mediated couplings are used to provide compounds of Formula (I), Formula (II) and Formula (III) as described in Scheme 4, Scheme 5, Scheme 6, and Scheme 7.

Scheme 4




5 [00300] In one aspect, a Suzuki reaction between compounds of structure **4-III** and compounds of structure **4-I** or **4-II** is used to provide compounds of structure **4-IV** or **4-V**. In one aspect, the phenyl ring of compounds of structure **4-III** is replaced with other cyclic rings (e.g. napthyl, cycloalkyl, monocyclic heteroaryl, bicyclic heteroaryl). Other metal mediated coupling reactions contemplated for the preparation of compounds of structure **4-IV** and **4-V** include, but are not limited to Suzuki reactions, Stille cross couplings, Negishi couplings, Kumada couplings, Ullmann reactions, Hiyama Coupling, and variants thereof (Metal-Catalyzed Cross-Coupling Reactions, Armin de Meijere (Editor), François Diederich (Editor), John Wiley & Sons; 2nd edition, 2004; Özdemir, *et al.*, *Tetrahedron*, 2005, 61, 9791-9798; Ackermann, *et al.*, *Org. Lett.*, 2006, 8, 3457-3460; Blakey, *et al.*, *J. Am. Chem. Soc.*, 2003, 125, 6046-6047; Dai, *et al.*, *Org. Lett.*, 2004, 6, 221-224; Yoshikai, *et al.*, *J. Am. Chem. Soc.*, 2005, 127, 17978-17979; Tang, *et al.*, *J. Org. Chem.*, 2006, 71, 2167-2169; Murata, *et al.*, *Synthesis*, 2001, 2231-2233).

10 [00301] In one aspect, the coupling partners in the metal mediated reaction condition includes those outlined in Scheme 5.

15


Scheme 5



[00302] In one aspect, halides of structure **4-I** are reacted with a borylating agent using transition metal mediated reaction conditions to form boronate compounds of structure **5-II**. Boronate compounds of structure **5-II** are reacted with compounds of structure **5-III** under palladium mediated coupling conditions to form biaryl aldehydes of structure **5-IV**.

[00303] In one aspect, compounds of Formula (I) where L^2 is $-O-$ are prepared as outlined in Scheme 6.

Scheme 6




[00304] In one aspect, boronate esters of structure **5-II** are oxidized to provide hydroxy compounds of structure **6-II**. An Ullman ether reaction between hydroxy compounds of structure **6-II** and boronic acid compounds of structure **6-III** provides compounds of structure **6-IV**. Methods of forming diaryl ethers, such as metal mediated reactions, include but are not limited to the Ulman Ether synthesis, Chan-Lam coupling, and Buchwald-Hartwig synthesis (D. Ma, Q. Cai, *Org. Lett.*, 2003, 5, 3799-3802; C. G. Bates, *et al.*, *Org. Lett.*, 2002, 4, 2803-2806; C. H. Burgos, *et al.*, *Angew. Chem. Int. Ed.*, 2006, 45, 4321-4326; C. H. Burgos, *et al.*, *Angew. Chem. Int. Ed.*, 2006, 45, 4321-4326; D. M. T. Chan, *et al.*, *Tetrahedron Lett.*, 1998, 39, 2933-2936; Z. Liu, R. C. Larock, *J. Org. Chem.*, 2006, 71, 3198-3209; Y.-J. Chen, H.-H. Chen, *Org. Lett.*,

2006, 8, 5609-5612; F. Li, Q. et al., *Org. Lett.*, 2003, 5, 2169-2171; D. A. Evans, et al., *Tetrahedron Letters*, 1998, 39, 2937-2940; C.-E. Yeom, et al., *Synlett*, 2007, 146-150).

[00305] In one aspect, compounds of Formula (I) where L^2 is $-O-$, $-S-$, or $-NH-$ are prepared as outlined Scheme 7.


5 **Scheme 7**



[00306] An Ullman ether reaction is used to couple boronic esters, or boronoc acids, or structure **5-II** with compounds of structure **7-II** to provide compounds of structure **7-III**. As described for Scheme 6, other coupling conditions exist.

10 [00307] In one aspect, benzyl halides of structure **8-II** are reacted with hydroxy compounds of structure **6-II** to provide ethers of structure **8-III**.


Scheme 8



[00308] Treatment of hydroxy compounds of structure **6-II** with alkyl halides of structure **8-II** in the presence of a suitable base and a suitable solvent provide ethers of structure **8-III**. In one aspect, the suitable base is selected from potassium carbonate, sodium carbonate, triethylamine and the like.

[00309] In one aspect, compounds of Formula (I) where L^2 is a heteroalkyl are prepared as outlined in Scheme 9.


Scheme 9



[00310] Treatment of alkyl halides of structure **9-I** with compounds of **9-II** in the presence of a suitable base and a suitable solvent provides compounds of structure **9-III**. In one aspect, L^2 is $-CH_2-O-$, $-CH_2-S-$, $-CH_2-NH-$, and the like.

[00311] In yet another aspect, compounds of Formula (I) are prepared as outlined in Scheme 10.



Scheme 10



[00312] Following the same procedures as described in Scheme 1, compounds of structure **10-I** are used to prepare compounds of Formula (I). In one aspect, compounds of structure **10-I** are prepared using the methodology described herein or known in the art.

[00313] A non-limiting example of the synthesis of compounds of structure **10-I** is shown in Scheme 11.

Scheme 11



[00314] In any of the schemes,  can be replaced with other cyclic groups as described herein.

[00315] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are synthesized as outlined in the Examples.

[00316] Throughout the specification, groups and substituents thereof are chosen by one skilled in the field to provide stable moieties and compounds.

[00317] A detailed description of techniques applicable to the creation of protecting groups and their removal are described in Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, and Kocienski, *Protective Groups*, Thieme Verlag, New York, NY, 1994, which are incorporated herein by reference for such disclosure.

Further Forms of Compounds

[00318] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) possess one or more stereocenters and each stereocenter exists independently in either the R or S configuration. The compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. The compounds and methods provided herein include all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof. In certain embodiments, compounds of Formula (I), Formula (II) and Formula (III) are prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds/salts, separating the diastereomers and recovering the optically pure enantiomers. In some embodiments, resolution of enantiomers is carried out using covalent diastereomeric derivatives of the compounds described herein. In another embodiment, diastereomers are separated by separation/resolution techniques based upon differences in solubility. In other embodiments, separation of stereoisomers is performed by chromatography or by the forming diastereomeric salts and separation by recrystallization, or chromatography, or any combination

thereof. Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981. In one aspect, stereoisomers are obtained by stereoselective synthesis.

[00319] The methods and compositions described herein include the use of amorphous forms as well as crystalline forms (also known as polymorphs). In one aspect, compounds described herein are in the form of pharmaceutically acceptable salts. As well, active metabolites of these compounds having the same type of activity are included in the scope of the present disclosure. In addition, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00320] In some embodiments, compounds described herein are prepared as prodrugs. A "prodrug" refers to an agent that is converted into the parent drug *in vivo*. Prodrugs are often useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. In some embodiments, the design of a prodrug increases the effective water solubility. An example, without limitation, of a prodrug is a compound described herein, which is administered as an ester (the "prodrug") to facilitate transmittal across a cell membrane where water solubility is detrimental to mobility but which then is metabolically hydrolyzed to the carboxylic acid, the active entity, once inside the cell where water-solubility is beneficial. A further example of a prodrug might be a short peptide (polyaminoacid) bonded to an acid group where the peptide is metabolized to reveal the active moiety. In certain embodiments, upon *in vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically active form of the compound. In certain embodiments, a prodrug is enzymatically metabolized by one or more steps or processes to the biologically, pharmaceutically or therapeutically active form of the compound.

[00321] In one aspect, prodrugs are designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacokinetic, pharmacodynamic processes and drug metabolism *in vivo*, once a pharmaceutically active compound is known, the design prodrugs of the compound is possible. (see, for example, Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New York, pages 388-392; Silverman (1992), *The Organic Chemistry of Drug Design and Drug Action*, Academic Press, Inc., San Diego, pages 352-401, Rooseboom *et al.*, *Pharmacological Reviews*, 56:53-102, 2004; Aesop Cho, "Recent Advances in Oral Prodrug Discovery", *Annual*

Reports in Medicinal Chemistry, Vol. 41, 395-407, 2006; T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series).

[00322] Prodrug forms of the herein described compounds, wherein the prodrug is metabolized *in vivo* to produce a compound of Formula (I), Formula (II) or Formula (III) as set forth herein are

5 included within the scope of the claims. In some cases, some of the herein-described compounds may be a prodrug for another derivative or active compound.

[00323] In some embodiments, sites on the aromatic ring portion of compounds of Formula (I),

Formula (II) and Formula (III) are susceptible to various metabolic reactions Therefore

10 incorporation of appropriate substituents on the aromatic ring structures will reduce, minimize or eliminate this metabolic pathway. In specific embodiments, the appropriate substituent to decrease or eliminate the susceptibility of the aromatic ring to metabolic reactions is, by way of example only, a halogen, or an alkyl group.

[00324] In another embodiment, the compounds described herein are labeled isotopically (e.g.

with a radioisotope) or by another other means, including, but not limited to, the use of

15 chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

[00325] Compounds described herein include isotopically-labeled compounds, which are identical to those recited in the various formulae and structures presented herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be

20 incorporated into the present compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, fluorine and chlorine, such as, for example, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{35}S , ^{18}F , ^{36}Cl . In one aspect, isotopically-labeled compounds described herein, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. In one aspect, substitution with isotopes such as deuterium affords

25 certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements.

[00326] In additional or further embodiments, the compounds described herein are metabolized upon administration to an organism in need to produce a metabolite that is then used to produce a desired effect, including a desired therapeutic effect.

30 [00327] "Pharmaceutically acceptable," as used herein, refers a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is relatively nontoxic, i.e., the material may be administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

[00328] The term “pharmaceutically acceptable salt” refers to a formulation of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. In some embodiments, pharmaceutically acceptable salts are obtained by reacting a compound of Formula (I), Formula 5 (II) or Formula (III) with acids. Pharmaceutically acceptable salts are also obtained by reacting a compound of Formula (I) with a base to form a salt.

[00329] Compounds described herein may be formed as, and/or used as, pharmaceutically acceptable salts. The type of pharmaceutical acceptable salts, include, but are not limited to: (1) acid addition salts, formed by reacting the free base form of the compound with a 10 pharmaceutically acceptable: inorganic acid, such as, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, metaphosphoric acid, and the like; or with an organic acid, such as, for example, acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, trifluoroacetic acid, tartaric acid, citric acid, benzoic acid, 15 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, toluenesulfonic acid, 2-naphthalenesulfonic acid, 4-methylbicyclo-[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, butyric acid, phenylacetic acid, phenylbutyric acid, valproic acid, and the like; (2) salts formed when an acidic proton present in the parent compound is replaced by a metal ion, e.g., an alkali metal ion (e.g. lithium, sodium, potassium), an alkaline earth ion (e.g. magnesium, or calcium), or an aluminum ion. In some embodiments, where the compound of Formula (I), Formula (II) or 20 Formula (III) has an acidic proton, a sodium salt of the compound of Formula (I), Formula (II) or Formula (III) is formed. In some cases, compounds described herein may coordinate with an organic base, such as, but not limited to, ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, dicyclohexylamine, tris(hydroxymethyl)methylamine. In other cases, compounds described herein may form salts with amino acids such as, but not 25 limited to, arginine, lysine, and the like. Acceptable inorganic bases used to form salts with compounds that include an acidic proton, include, but are not limited to, aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, sodium hydroxide, and the like.

[00330] It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs. Solvates 30 contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed

5 during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds described herein can be conveniently prepared or formed during the processes described herein. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

10 [00331] Compounds described herein, such as compounds of Formula (I), Formula (II) and Formula (III), may be in various forms, including but not limited to, amorphous forms, milled forms and nano-particulate forms. In addition, compounds described herein include crystalline forms, also known as polymorphs. Polymorphs include the different crystal packing arrangements of the same elemental composition of a compound. Polymorphs usually have different X-ray diffraction patterns, melting points, density, hardness, crystal shape, optical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of 15 crystallization, and storage temperature may cause a single crystal form to dominate.

15 [00332] Throughout the specification, groups and substituents thereof can be chosen by one skilled in the field to provide stable moieties and compounds.

Certain Terminology

20 [00333] Unless otherwise stated, the following terms used in this application, including the specification and claims, have the definitions given below. It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Unless otherwise indicated, conventional methods of mass spectroscopy, NMR, HPLC, protein chemistry, biochemistry, recombinant 25 DNA techniques and pharmacology are employed. In this application, the use of "or" or "and" means "and/or" unless stated otherwise. Furthermore, use of the term "including" as well as other forms, such as "include", "includes," and "included," is not limiting. The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

30 [00334] An "alkyl" group refers to an aliphatic hydrocarbon group. The alkyl group may be a saturated alkyl group (which means that it does not contain any carbon-carbon double bonds or carbon-carbon triple bonds) or the alkyl group may be an unsaturated alkyl group (which means that it contains at least one carbon-carbon double bonds or carbon-carbon triple bond). The alkyl moiety, whether saturated or unsaturated, may be branched or straight chain.

35 [00335] The "alkyl" group may have 1 to 10 carbon atoms (whenever it appears herein, a numerical range such as "1 to 10" refers to each integer in the given range; *e.g.*, "1 to 10 carbon

atoms" means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms, although the present definition also covers the occurrence of the term "alkyl" where no numerical range is designated). The alkyl group of the compounds described herein may be designated as "C₁-C₆ alkyl" or similar designations. By way of example only, "C₁-C₆ alkyl" indicates that there are one, two, three, four, five, or six carbon atoms in the alkyl chain. In one aspect the alkyl is selected from the group consisting of methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tertiary butyl, pentyl, neopentyl, hexyl, allyl, but-2-enyl, but-3-enyl, and the like. In one aspect, an alkyl is a C₁-C₆alkyl.

[00336] The term "alkylene" refers to a divalent alkyl radical. Any of the above mentioned monovalent alkyl groups may be an alkylene by abstraction of a second hydrogen atom from the alkyl. In one aspect, an alkylene is a C₁-C₆alkylene. In another aspect, an alkylene is a C₁-C₄alkylene. Typical alkylene groups include, but are not limited to, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH(CH₃)-, -CH₂C(CH₃)₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, and the like.

[00337] An "alkoxy" group refers to a (alkyl)O- group, where alkyl is as defined herein.

[00338] The term "alkylamine" refers to the -N(alkyl)_xH_y group, where x and y are selected from the group x=1, y=1 and x=2, y=0. In some embodiments, when x=2 and y=0, the alkyl groups taken together with the nitrogen atom to which they are attached form a cyclic ring system.

[00339] The term "aromatic" refers to a planar ring having a delocalized π -electron system containing 4n+2 π electrons, where n is an integer. Aromatic rings can be formed from five, six, seven, eight, nine, ten, or more than ten atoms. Aromatics are optionally substituted. The term "aromatic" includes both carbocyclic aryl ("aryl", *e.g.*, phenyl) and heterocyclic aryl (or "heteroaryl" or "heteroaromatic") groups (*e.g.*, pyridine). The term includes monocyclic or fused-ring polycyclic (*i.e.*, rings which share adjacent pairs of carbon atoms) groups.

[00340] The term "carbocyclic" or "carbocycle" refers to a ring or ring system where the atoms forming the backbone of the ring are all carbon atoms. The term thus distinguishes carbocyclic from heterocyclic rings in which the ring backbone contains at least one atom which is different from carbon.

[00341] As used herein, the term "aryl" refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. Aryl rings are formed by five, six, seven, eight, nine, or more than nine carbon atoms. Aryl groups are optionally substituted. In one aspect, an aryl is a phenyl or a naphthalenyl. In one aspect, an aryl is a phenyl. In one aspect, an aryl is a C₆-C₁₀aryl. Depending on the structure, an aryl group can be a monoradical or a diradical (*i.e.*, an arylene

group). Exemplary arylenes include, but are not limited to, phenyl-1,2-ene, phenyl-1,3-ene, and phenyl-1,4-ene.

[00342] The term “cycloalkyl” refers to a monocyclic or polycyclic aliphatic, non-aromatic radical, wherein each of the atoms forming the ring (i.e. skeletal atoms) is a carbon atom.

5 Cycloalkyls may be saturated, or partially unsaturated. Cycloalkyls may be fused with an aromatic ring, and the point of attachment is at a carbon that is not an aromatic ring carbon atom. Cycloalkyl groups include groups having from 3 to 10 ring atoms. In some embodiments, cycloalkyl groups are selected from among cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cyclooctyl. Cycloalkyl groups may be substituted or 10 unsubstituted. Depending on the structure, a cycloalkyl group can be a monoradical or a diradical (i.e., an cycloalkylene group, such as, but not limited to, cyclopropan-1,1-diyl, cyclobutan-1,1-diyl, cyclopentan-1,1-diyl, cyclohexan-1,1-diyl, cyclohexan-1,4-diyl, cycloheptan-1,1-diyl, and the like). In one aspect, a cycloalkyl is a C₃-C₆cycloalkyl.

15 [00343] The term “halo” or, alternatively, “halogen” or “halide” means fluoro, chloro, bromo or iodo.

[00344] The term “haloalkyl” refers to an alkyl group in which one or more hydrogen atoms are replaced by one or more halide atoms. In one aspect, a haloalkyl is a C₁-C₄haloalkyl.

20 [00345] The term “haloalkylene” refers to an alkylene group in which one or more hydrogen atoms are replaced by one or more halide atoms. In one aspect, a haloalkylene is a C₁-C₆haloalkylene. In another aspect, a haloalkylene is a C₁-C₄haloalkylene.

[00346] The term “fluoroalkyl” refers to an alkyl in which one or more hydrogen atoms are replaced by a fluorine atom. In one aspect, a fluoralkyl is a C₁-C₄fluoroalkyl.

25 [00347] The term “fluoroalkylene” refers to an alkylene in which one or more hydrogen atoms are replaced by a fluorine atom. In one aspect, a fluoralkylene is a C₁-C₆fluoroalkylene. In another aspect, a fluoralkylene is a C₁-C₄fluoroalkylene.

[00348] The term “heteroalkyl” refers to an alkyl group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorus or combinations thereof. In one aspect, a heteroalkyl is a C₁-C₆heteroalkyl.

30 [00349] The term “heteroalkylene” refers to an alkylene group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorus or combinations thereof. In one aspect, a heteroalkylene is a C₁-C₆heteroalkylene. In another aspect, a heteroalkylene is a C₁-C₄heteroalkylene. Exemplary heteroalkylenes include, but are not limited to, -OCH₂-, -OCH(CH₃)-, -OC(CH₃)₂-, -OCH₂CH₂-, -CH₂O-, -CH(CH₃)O-, -C(CH₃)₂O-, -CH₂CH₂O-, -CH₂OCH₂-, -CH₂OCH₂CH₂-, -CH₂CH₂OCH₂-, -SCH₂-, -SCH(CH₃)-, -SC(CH₃)₂-, -SCH₂CH₂-, -CH₂S-, -CH(CH₃)S-, -C(CH₃)₂S-, -CH₂CH₂S-, -CH₂SCH₂-, -

CH₂SCH₂CH₂-, -CH₂CH₂SCH₂-, -SO₂CH₂-, -SO₂CH(CH₃)-, -SO₂C(CH₃)₂-, -SO₂CH₂CH₂-, -CH₂SO₂-, -CH(CH₃)SO₂-, -C(CH₃)₂SO₂-, -CH₂CH₂SO₂-, -CH₂SO₂CH₂-, -CH₂SO₂CH₂CH₂-, -CH₂CH₂SO₂CH₂-, -NHCH₂-, -NHCH(CH₃)-, -NHC(CH₃)₂-, -NHCH₂CH₂-, -CH₂NH-, -CH(CH₃)NH-, -C(CH₃)₂NH-, -CH₂CH₂NH-, -CH₂NHCH₂-, -CH₂NHCH₂CH₂-, -

5 CH₂CH₂NHCH₂-, and the like.

[00350] The term "heterocycle" or "heterocyclic" refers to heteroaromatic rings (also known as heteroaryls) and heterocycloalkyl rings (also known as heteroalicyclic groups) containing one to four heteroatoms in the ring(s), where each heteroatom in the ring(s) is selected from O, S and N, wherein each heterocyclic group has from 4 to 10 atoms in its ring system, and with the proviso

10 that the any ring does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups (also known as heterocycloalkyls) include groups having only 3 atoms in their ring system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems. An example of a 3-membered heterocyclic group is aziridinyl. An example of a 4-membered heterocyclic group is azetidinyl. An example of a 5-

15 membered heterocyclic group is thiazolyl. An example of a 6-membered heterocyclic group is pyridyl, and an example of a 10-membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl,

tetrahydrothienyl, oxazolidinonyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, thioxanyl, piperazinyl, aziridinyl, azetidinyl, oxetanyl,


20 thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 1,2,3,6-tetrahydropyridinyl, pyrrolin-2-yl, pyrrolin-3-yl, indolinyl, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazolanyl, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl,

dihydrofuranyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl and quinolizinyl. Examples of aromatic heterocyclic

25 groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indolizinyl, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, and

30 furopyridinyl. The foregoing groups may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl or imidazol-3-yl (both N-attached) or imidazol-2-yl, imidazol-4-yl or imidazol-5-yl (all C-attached). The heterocyclic groups include benzo-fused ring systems. Non-aromatic heterocycles may be substituted with one or two oxo (=O) moieties, such as pyrrolidin-2-one.


[00351] The terms “heteroaryl” or, alternatively, “heteroaromatic” refers to an aryl group that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur. Illustrative examples of heteroaryl groups include the following moieties:



include pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl,

10 thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, triazinyl, oxadiazolyl, thiadiazolyl, and furazanyl. In one aspect, a heteroaryl contains 0-3 N atoms. In another aspect, a heteroaryl contains 1-3 N atoms. In another aspect, a heteroaryl contains 0-3 N atoms, 0-1 O atoms, and 0-1 S atoms. In another aspect, a heteroaryl is a monocyclic or bicyclic heteroaryl. In one aspect, heteroaryl is a C₁-C₉heteroaryl. In one aspect, monocyclic heteroaryl is a C₁-C₅heteroaryl. In one aspect, monocyclic heteroaryl is a 5-membered or 6-membered heteroaryl. In one aspect, bicyclic heteroaryl is a C₆-C₉heteroaryl. Depending on the structure, a heteroaryl group can be a monoradical or a diradical (i.e., a heteroarylene group).

[00352] A “heterocycloalkyl” or “heteroalicyclic” group refers to a cycloalkyl group that includes at least one heteroatom selected from nitrogen, oxygen and sulfur. The radicals may be fused with an aryl or heteroaryl. Illustrative examples of heterocycloalkyl groups, also referred to as non-aromatic heterocycles, include:



and the like. In some embodiments, the heterocycloalkyl is selected from oxazolidinonyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, and indolinyl. The term heteroalicyclic also includes all ring forms of the carbohydrates, including but not limited to the monosaccharides, the disaccharides and the oligosaccharides. In one aspect, a heterocycloalkyl is a C₂-C₁₀heterocycloalkyl. In another aspect, a heterocycloalkyl is a C₄-C₁₀heterocycloalkyl. In one aspect, a heterocycloalkyl contains 0-2 N atoms. In another aspect, a heterocycloalkyl contains 0-2 N atoms, 0-2 O atoms or 0-1 S atoms.


[00353] The term “bond” or “single bond” refers to a chemical bond between two atoms, or two 10 moieties when the atoms joined by the bond are considered to be part of larger substructure. In one aspect, when a group described herein is a bond, the referenced group is absent thereby allowing a bond to be formed between the remaining identified groups.

[00354] The term “membered ring” includes any cyclic structure. The term “membered” is meant to denote the number of skeletal atoms that constitute the ring. Thus, for example, cyclohexyl, 15 pyridinyl, pyranyl and thiopyranyl are 6-membered rings and cyclopentyl, pyrrolyl, furanyl, and thieryl are 5-membered rings.

[00355] The term “moiety” refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

20 [00356] As used herein, “carboxylic acid bioisostere” refers to a functional group or moiety that exhibits similar physical, biological and/or chemical properties as a carboxylic acid moiety.

Examples of carboxylic acid bioisosteres include, but are not limited to,



and the like.

25 [00357] The term “optionally substituted” or “substituted” means that the referenced group may be substituted with one or more additional group(s) individually and independently selected from alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, arylsulfone, cyano, halo, nitro, haloalkyl, fluoroalkyl, fluoroalkoxy, and amino, including mono- and di-substituted amino groups, and the protected 30 derivatives thereof. By way of example an optional substituents may be halide, -CN, -NO₂, or L_sR_s, wherein each L_s is independently selected from a bond, -O-, -C(=O)-, -C(=O)O-, -S-, -

S(=O)-, -S(=O)₂-, -NH-, -NHC(=O)-, -C(=O)NH-, S(=O)₂NH-, -NHS(=O)₂, -OC(=O)NH-, -NHC(=O)O-, or -(C₁-C₆ alkylene)-; and each R_s is selected from H, alkyl, fluoroalkyl, heteroalkyl, cycloalkyl, aryl, heteroaryl, or heterocycloalkyl. The protecting groups that may form the protective derivatives of the above substituents may be found in sources such as Greene and Wuts, above. In some embodiments, optional substituents are selected from halogen, -CN, -NH₂, -OH, -N(CH₃)₂, alkyl, fluoroalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, alkoxy, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, and arylsulfone. In some embodiments, an optional substituent is halogen, -CN, -NH₂, -OH, -NH(CH₃), -N(CH₃)₂, -CO₂H, -CO₂alkyl, -C(=O)NH₂, -C(=O)NHalkyl, -C(=O)N(alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(alkyl), -S(=O)₂N(alkyl)₂, alkyl, cycloalkyl, fluoroalkyl, heteroalkyl, alkoxy, fluoroalkoxy, -S-alkyl, or -S(=O)₂alkyl. In some embodiments, an optional substituent is selected from halogen, -CN, -NH₂, -OH, -NH(CH₃), -N(CH₃)₂, -CH₃, -CH₂CH₃, -CF₃, -OCH₃, and -OCF₃. In some embodiments, substituted groups are substituted with one or two of the preceding groups. In some embodiments, substituted groups are substituted with one of the preceding groups. In some embodiments, an optional substituent on an aliphatic carbon atom (acyclic or cyclic, saturated or unsaturated carbon atoms, excluding aromatic carbon atoms) includes oxo (=O).

[00358] In certain embodiments, the compounds presented herein possess one or more stereocenters and each center independently exists in either the R or S configuration. The compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. Stereoisomers are obtained, if desired, by methods such as, stereoselective synthesis and/or the separation of stereoisomers by chiral chromatographic columns.

[00359] The methods and formulations described herein include the use of *N*-oxides (if appropriate), crystalline forms (also known as polymorphs), or pharmaceutically acceptable salts of compounds having the structure of Formula (I), Formula (II) or Formula (III), as well as active metabolites of these compounds having the same type of activity. In some situations, compounds may exist as tautomers. All tautomers are included within the scope of the compounds presented herein. In specific embodiments, the compounds described herein exist in solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In other embodiments, the compounds described herein exist in unsolvated form.

Certain Terminology

[00360] The term “acceptable” with respect to a formulation, composition or ingredient, as used herein, means having no persistent detrimental effect on the general health of the subject being treated.

[00361] The term “modulate,” as used herein, means to interact with a target either directly or indirectly so as to alter the activity of the target, including, by way of example only, to enhance the activity of the target, to inhibit the activity of the target, to limit the activity of the target, or to extend the activity of the target.

5 [00362] The term “modulator,” as used herein, refers to a molecule that interacts with a target either directly or indirectly. The interactions include, but are not limited to, the interactions of an agonist, partial agonist, an inverse agonist and antagonist. In one embodiment, a modulator is an antagonist.

10 [00363] The term “agonist,” as used herein, refers to a molecule such as a compound, a drug, an enzyme activator or a hormone modulator that binds to a specific receptor and triggers a response in the cell. An agonist mimics the action of an endogenous ligand (such as LPA, prostaglandin, hormone or neurotransmitter) that binds to the same receptor.

[00364] The term “antagonist,” as used herein, refers to a molecule such as a compound, which diminishes, inhibits, or prevents the action of another molecule or the activity of a receptor site.

15 Antagonists include, but are not limited to, competitive antagonists, non-competitive antagonists, uncompetitive antagonists, partial agonists and inverse agonists.

[00365] The term “LPA-dependent”, as used herein, refers to conditions or disorders that would not occur, or would not occur to the same extent, in the absence of LPA.

20 [00366] The term “LPA-mediated”, as used herein, refers to conditions or disorders that might occur in the absence of LPA but can occur in the presence of LPA.

[00367] “Selectivity” for one LPA receptor versus other LPA receptors means that the compound has an IC₅₀ (Ca Flux assay) for the indicated LPA receptor that is at least 10-fold less than the IC₅₀ for other LPA receptors. In some embodiments, selectivity for one LPA receptor versus other LPA receptor means that the compound has an IC₅₀ for the indicated LPA receptor that is at least 10-fold, at least 20-fold, at least 40-fold, at least 50-fold, at least 100-fold, at least 200-fold, at least 500-fold, or at least 1000-fold, less than the IC₅₀ for other LPA receptors. For example, a selective LPA₁ receptor antagonist has an IC₅₀ that is at least 10-fold, at least 20-fold, at least 40-fold, at least 50-fold, at least 100-fold, at least 200-fold, at least 500-fold, or at least 1000-fold, less than the IC₅₀ for other LPA receptors (e.g. LPA₂, LPA₃).

25 [00368] The terms “co-administration” or the like, as used herein, are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different time.

30 [00369] The terms “effective amount” or “therapeutically effective amount,” as used herein, refer to a sufficient amount of an agent or a compound being administered which will relieve to some

extent one or more of the symptoms of the disease or condition being treated. The result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic uses is the amount of the composition comprising a compound as disclosed herein required to provide a 5 clinically significant decrease in disease symptoms. An appropriate “effective” amount in any individual case may be determined using techniques, such as a dose escalation study.

[00370] The terms “enhance” or “enhancing,” as used herein, means to increase or prolong either in potency or duration a desired effect. Thus, in regard to enhancing the effect of therapeutic agents, the term “enhancing” refers to the ability to increase or prolong, either in potency or 10 duration, the effect of other therapeutic agents on a system. An “enhancing-effective amount,” as used herein, refers to an amount adequate to enhance the effect of another therapeutic agent in a desired system.

[00371] The terms “kit” and “article of manufacture” are used as synonyms.

[00372] A “metabolite” of a compound disclosed herein is a derivative of that compound that is 15 formed when the compound is metabolized. The term “active metabolite” refers to a biologically active derivative of a compound that is formed when the compound is metabolized. The term “metabolized,” as used herein, refers to the sum of the processes (including, but not limited to, hydrolysis reactions and reactions catalyzed by enzymes) by which a particular substance is changed by an organism. Thus, enzymes may produce specific structural alterations to a 20 compound. For example, cytochrome P450 catalyzes a variety of oxidative and reductive reactions while uridine diphosphate glucuronyltransferases catalyze the transfer of an activated glucuronic-acid molecule to aromatic alcohols, aliphatic alcohols, carboxylic acids, amines and free sulphhydryl groups. In some embodiments, carboxylic acid containing compounds form taurine conjugates *in vivo*. Metabolites of the compounds disclosed herein are optionally 25 identified either by administration of compounds to a host and analysis of tissue samples from the host, or by incubation of compounds with hepatic cells *in vitro* and analysis of the resulting compounds.

[00373] The term “pharmaceutical combination” as used herein, means a product that results from 30 the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term “fixed combination” means that the active ingredients, e.g. a compound of Formula (I), Formula (II) or Formula (III) and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage. The term “non-fixed combination” means that the active ingredients, e.g. a compound of Formula (I), Formula (II) or Formula (III) and a co-agent, are administered to a patient as separate entities 35 either simultaneously, concurrently or sequentially with no specific intervening time limits,

wherein such administration provides effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[00374] The term “subject” or “patient” encompasses mammals and non-mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. In one embodiment, the mammal is a human.

[00375] The terms “treat,” “treating” or “treatment,” as used herein, include alleviating, abating or ameliorating at least one symptom of a disease disease or condition, preventing additional symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

Routes of Administration

[00376] Suitable routes of administration include, but are not limited to, oral, intravenous, rectal, aerosol, parenteral, ophthalmic, pulmonary, transmucosal, transdermal, vaginal, otic, nasal, and topical administration. In addition, by way of example only, parenteral delivery includes intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intralymphatic, and intranasal injections.

[00377] In certain embodiments, a compound as described herein is administered in a local rather than systemic manner, for example, via injection of the compound directly into an organ, often in a depot preparation or sustained release formulation. In specific embodiments, long acting formulations are administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Furthermore, in other embodiments, the drug is delivered in a targeted drug delivery system, for example, in a liposome coated with organ-specific antibody. In such embodiments, the liposomes are targeted to and taken up selectively by the organ. In yet other embodiments, the compound as described herein is provided in the form of a rapid release formulation, in the form of an extended release formulation, or in the form of an intermediate release formulation. In yet other embodiments, the compound described herein is administered topically.

Pharmaceutical Compositions/Formulations

[00378] In some embodiments, the compounds described herein are formulated into pharmaceutical compositions. Pharmaceutical compositions are formulated in a conventional

manner using one or more pharmaceutically acceptable inactive ingredients that facilitate processing of the active compounds into preparations that can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein can be found, for example, in Remington: The Science and

5 Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference for such disclosure.

10 [00379] Provided herein are pharmaceutical compositions that include a compound of Formula (I), Formula (II) or Formula (III) and at least one pharmaceutically acceptable inactive ingredient. In some embodiments, the compounds described herein are administered as pharmaceutical compositions in which a compound of Formula (I), Formula (II) or Formula (III) is mixed with other active ingredients, as in combination therapy. In other embodiments, the pharmaceutical 15 compositions include other medicinal or pharmaceutical agents, carriers, adjuvants, preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure, and/or buffers. In yet other embodiments, the pharmaceutical compositions include other therapeutically valuable substances.

[00380] A pharmaceutical composition, as used herein, refers to a mixture of a compound of 20 Formula (I), Formula (II) or Formula (III) with other chemical components (i.e. pharmaceutically acceptable inactive ingredients), such as carriers, excipients, binders, filling agents, suspending agents, flavoring agents, sweetening agents, disintegrating agents, dispersing agents, surfactants, lubricants, colorants, diluents, solubilizers, moistening agents, plasticizers, stabilizers, penetration enhancers, wetting agents, anti-foaming agents, antioxidants, preservatives, or one or 25 more combination thereof. The pharmaceutical composition facilitates administration of the compound to an organism. In practicing the methods of treatment or use provided herein, therapeutically effective amounts of compounds described herein are administered in a pharmaceutical composition to a mammal having a disease, disorder, or condition to be treated. In some embodiments, the mammal is a human. A therapeutically effective amount can vary 30 widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors. The compounds can be used singly or in combination with one or more therapeutic agents as components of mixtures.

[00381] The pharmaceutical formulations described herein are administered to a subject by appropriate administration routes, including but not limited to, oral, parenteral (e.g., intravenous, 35 subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration

routes. The pharmaceutical formulations described herein include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal dispersions, aerosols, solid dosage forms, powders, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release 5 formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate and controlled release formulations.

[00382] Pharmaceutical compositions including a compound of Formula (I), Formula (II) or Formula (III) are manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, 10 encapsulating, entrapping or compression processes.

[00383] The pharmaceutical compositions will include at least one compound of Formula (I), Formula (II) or Formula (III) as an active ingredient in free-acid or free-base form, or in a pharmaceutically acceptable salt form. In addition, the methods and pharmaceutical compositions described herein include the use of *N*-oxides (if appropriate), crystalline forms, amorphous 15 phases, as well as active metabolites of these compounds having the same type of activity. In some embodiments, compounds described herein exist in unsolvated form or in solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00384] In certain embodiments, compositions provided herein include one or more preservatives 20 to inhibit microbial activity. Suitable preservatives include mercury-containing substances such as merfen and thiomersal; stabilized chlorine dioxide; and quaternary ammonium compounds such as benzalkonium chloride, cetyltrimethylammonium bromide and cetylpyridinium chloride.

[00385] In some embodiments, formulations described herein benefit from antioxidants, metal 25 chelating agents, thiol containing compounds and other general stabilizing agents. Examples of such stabilizing agents, include, but are not limited to: (a) about 0.5% to about 2% w/v glycerol, (b) about 0.1% to about 1% w/v methionine, (c) about 0.1% to about 2% w/v monothioglycerol, (d) about 1 mM to about 10 mM EDTA, (e) about 0.01% to about 2% w/v ascorbic acid, (f) 0.003% to about 0.02% w/v polysorbate 80, (g) 0.001% to about 0.05% w/v. polysorbate 20, (h) arginine, (i) heparin, (j) dextran sulfate, (k) cyclodextrins, (l) pentosan polysulfate and other 30 heparinoids, (m) divalent cations such as magnesium and zinc; or (n) combinations thereof.

[00386] The pharmaceutical compositions described herein, which include a compound of Formula (I), Formula (II) or Formula (III) are formulated into any suitable dosage form, including but not limited to, aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions, solid oral dosage forms, aerosols, controlled release formulations, fast melt 35 formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills,

dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[00387] Pharmaceutical preparations for oral use are obtained by mixing one or more solid excipient with one or more of the compounds described herein, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methylcellulose, 10 microcrystalline cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose; or others such as: polyvinylpyrrolidone (PVP or povidone) or calcium phosphate. If desired, disintegrating agents are added, such as the cross-linked croscarmellose sodium, polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. In some embodiments, dyestuffs or pigments are added to the tablets or dragee coatings for identification 15 or to characterize different combinations of active compound doses.

[00388] Pharmaceutical preparations that are administered orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, 20 optionally, stabilizers. In soft capsules, the active compounds are dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In some embodiments, stabilizers are added.

[00389] All formulations for oral administration are in dosages suitable for such administration.

[00390] In one aspect, solid oral dosage forms are prepared by mixing a compound of Formula (I), 25 Formula (II) or Formula (III) with one or more of the following: antioxidants, flavoring agents, and carrier materials such as binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, and diluents.

[00391] In some embodiments, the solid dosage forms disclosed herein are in the form of a tablet, (including a suspension tablet, a fast-melt tablet, a bite-disintegration tablet, a rapid- 30 disintegration tablet, an effervescent tablet, or a caplet), a pill, a powder, a capsule, solid dispersion, solid solution, bioerodible dosage form, controlled release formulations, pulsatile release dosage forms, multiparticulate dosage forms, beads, pellets, granules. In other embodiments, the pharmaceutical formulation is in the form of a powder. In still other embodiments, the pharmaceutical formulation is in the form of a tablet. In other embodiments,

pharmaceutical formulations of the compound of Formula (I), Formula (II) or Formula (III) is in the form of a capsule.

[00392] In some embodiments, solid dosage forms, e.g., tablets, effervescent tablets, and capsules, are prepared by mixing particles of a compound of Formula (I), Formula (II) or Formula (III)

5 with one or more pharmaceutical excipients to form a bulk blend composition. The bulk blend is readily subdivided into equally effective unit dosage forms, such as tablets, pills, and capsules. In some embodiments, the individual unit dosages include film coatings. These formulations are manufactured by conventional formulation techniques.

[00393] Conventional formulation techniques include, e.g., one or a combination of methods: (1)

10 dry mixing, (2) direct compression, (3) milling, (4) dry or non-aqueous granulation, (5) wet granulation, or (6) fusion. Other methods include, e.g., spray drying, pan coating, melt granulation, granulation, fluidized bed spray drying or coating (e.g., wurster coating), tangential coating, top spraying, tabletting, extruding and the like.

[00394] Suitable carriers for use in the solid dosage forms described herein include, but are not

15 limited to, acacia, gelatin, colloidal silicon dioxide, calcium glycerophosphate, calcium lactate, maltodextrin, glycerine, magnesium silicate, sodium caseinate, soy lecithin, sodium chloride, tricalcium phosphate, dipotassium phosphate, sodium stearoyl lactylate, carrageenan, monoglyceride, diglyceride, pregelatinized starch, hydroxypropylmethylcellulose, hydroxypropylmethylcellulose acetate stearate, sucrose, microcrystalline cellulose, lactose, 20 mannitol and the like.

[00395] Suitable filling agents for use in the solid dosage forms described herein include, but are not limited to, lactose, calcium carbonate, calcium phosphate, dibasic calcium phosphate, calcium sulfate, microcrystalline cellulose, cellulose powder, dextrose, dextrose, dextrates, dextran, starches, pregelatinized starch, hydroxypropylmethylcellulose (HPMC),

25 hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate stearate (HPMCAS), sucrose, xylitol, lactitol, mannitol, sorbitol, sodium chloride, polyethylene glycol, and the like.

[00396] Suitable disintegrants for use in the solid dosage forms described herein include, but are not limited to, natural starch such as corn starch or potato starch, a pregelatinized starch, or

30 sodium starch glycolate, a cellulose such as methylcrystalline cellulose, methylcellulose, microcrystalline cellulose, croscarmellose, or a cross-linked cellulose, such as cross-linked sodium carboxymethylcellulose, cross-linked carboxymethylcellulose, or cross-linked croscarmellose, a cross-linked starch such as sodium starch glycolate, a cross-linked polymer such as crospovidone, a cross-linked polyvinylpyrrolidone, alginate such as alginic acid or a salt 35 of alginic acid such as sodium alginate, a gum such as agar, guar, locust bean, Karaya, pectin, or

tragacanth, sodium starch glycolate, bentonite, sodium lauryl sulfate, sodium lauryl sulfate in combination starch, and the like.

[00397] Binders impart cohesiveness to solid oral dosage form formulations: for powder filled capsule formulation, they aid in plug formation that can be filled into soft or hard shell capsules and for tablet formulation, they ensure the tablet remaining intact after compression and help assure blend uniformity prior to a compression or fill step. Materials suitable for use as binders in the solid dosage forms described herein include, but are not limited to, carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, hydroxypropylmethylcellulose acetate stearate, hydroxyethylcellulose, hydroxypropylcellulose, ethylcellulose, and microcrystalline cellulose, microcrystalline dextrose, amylose, magnesium aluminum silicate, polysaccharide acids, bentonites, gelatin, polyvinylpyrrolidone/vinyl acetate copolymer, crospovidone, povidone, starch, pregelatinized starch, tragacanth, dextrin, a sugar, such as sucrose, glucose, dextrose, molasses, mannitol, sorbitol, xylitol, lactose, a natural or synthetic gum such as acacia, tragacanth, ghatti gum, mucilage of isapol husks, starch, polyvinylpyrrolidone, larch arabogalactan, polyethylene glycol, waxes, sodium alginate, and the like.

[00398] In general, binder levels of 20-70% are used in powder-filled gelatin capsule formulations. Binder usage level in tablet formulations varies whether direct compression, wet granulation, roller compaction, or usage of other excipients such as fillers which itself can act as moderate binder. Binder levels of up to 70% in tablet formulations is common.

[00399] Suitable lubricants or glidants for use in the solid dosage forms described herein include, but are not limited to, stearic acid, calcium hydroxide, talc, corn starch, sodium stearyl fumerate, alkali-metal and alkaline earth metal salts, such as aluminum, calcium, magnesium, zinc, stearic acid, sodium stearates, magnesium stearate, zinc stearate, waxes, Stearowet[®], boric acid, sodium benzoate, sodium acetate, sodium chloride, leucine, a polyethylene glycol or a methoxypolyethylene glycol such as CarbowaxTM, PEG 4000, PEG 5000, PEG 6000, propylene glycol, sodium oleate, glyceryl behenate, glyceryl palmitostearate, glyceryl benzoate, magnesium or sodium lauryl sulfate, and the like.

[00400] Suitable diluents for use in the solid dosage forms described herein include, but are not limited to, sugars (including lactose, sucrose, and dextrose), polysaccharides (including dextrates and maltodextrin), polyols (including mannitol, xylitol, and sorbitol), cyclodextrins and the like.

[00401] Suitable wetting agents for use in the solid dosage forms described herein include, for example, oleic acid, glyceryl monostearate, sorbitan monooleate, sorbitan monolaurate, triethanolamine oleate, polyoxyethylene sorbitan monooleate, polyoxyethylene sorbitan monolaurate, quaternary ammonium compounds (e.g., Polyquat 10[®]), sodium oleate, sodium lauryl sulfate, magnesium stearate, sodium docusate, triacetin, vitamin E TPGS and the like.

[00402] Suitable surfactants for use in the solid dosage forms described herein include, for example, sodium lauryl sulfate, sorbitan monooleate, polyoxyethylene sorbitan monooleate, polysorbates, polaxomers, bile salts, glyceryl monostearate, copolymers of ethylene oxide and propylene oxide, e.g., Pluronic® (BASF), and the like.

5 [00403] Suitable suspending agents for use in the solid dosage forms described here include, but are not limited to, polyvinylpyrrolidone, e.g., polyvinylpyrrolidone K12, polyvinylpyrrolidone K17, polyvinylpyrrolidone K25, or polyvinylpyrrolidone K30, polyethylene glycol, e.g., the polyethylene glycol can have a molecular weight of about 300 to about 6000, or about 3350 to about 4000, or about 7000 to about 5400, vinyl pyrrolidone/vinyl acetate copolymer (S630),
10 sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, polysorbate-80, hydroxyethylcellulose, sodium alginate, gums, such as, e.g., gum tragacanth and gum acacia, guar gum, xanthans, including xanthan gum, sugars, cellulosics, such as, e.g., sodium carboxymethylcellulose, methylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose, polysorbate-80, sodium alginate,
15 polyethoxylated sorbitan monolaurate, polyethoxylated sorbitan monolaurate, povidone and the like.

[00404] Suitable antioxidants for use in the solid dosage forms described herein include, for example, e.g., butylated hydroxytoluene (BHT), sodium ascorbate, and tocopherol.

20 [00405] It should be appreciated that there is considerable overlap between additives used in the solid dosage forms described herein. Thus, the above-listed additives should be taken as merely exemplary, and not limiting, of the types of additives that can be included in solid dosage forms of the pharmaceutical compositions described herein. The amounts of such additives can be readily determined by one skilled in the art, according to the particular properties desired.

25 [00406] Compressed tablets are solid dosage forms prepared by compacting the bulk blend of the formulations described above.

[00407] In various embodiments, tablets will include one or more flavoring agents.

30 [00408] In other embodiments, the tablets will include a film surrounding the final compressed tablet. In some embodiments, the film coating can provide a delayed release of the compound of Formula (I), Formula (II) or Formula (III) from the formulation. In other embodiments, the film coating aids in patient compliance (e.g., Opadry® coatings or sugar coating). Film coatings including Opadry® typically range from about 1% to about 3% of the tablet weight.

35 [00409] A capsule may be prepared, for example, by placing the bulk blend of the formulation of the compound described above, inside of a capsule. In some embodiments, the formulations (non-aqueous suspensions and solutions) are placed in a soft gelatin capsule. In other embodiments, the formulations are placed in standard gelatin capsules or non-gelatin capsules

such as capsules comprising HPMC. In other embodiments, the formulation is placed in a sprinkle capsule, wherein the capsule is swallowed whole or the capsule is opened and the contents sprinkled on food prior to eating.

[00410] In various embodiments, the particles of the compound of Formula (I), Formula (II) or

5 Formula (III) and one or more excipients are dry blended and compressed into a mass, such as a tablet, having a hardness sufficient to provide a pharmaceutical composition that substantially disintegrates within less than about 30 minutes, less than about 35 minutes, less than about 40 minutes, less than about 45 minutes, less than about 50 minutes, less than about 55 minutes, or less than about 60 minutes, after oral administration, thereby releasing the formulation into the

10 gastrointestinal fluid.

[00411] In other embodiments, a powder including a compound of Formula (I), Formula (II) or Formula (III) is formulated to include one or more pharmaceutical excipients and flavors. Such a powder is prepared, for example, by mixing the the compound of Formula (I), Formula (II) or Formula (III) and optional pharmaceutical excipients to form a bulk blend composition.

15 Additional embodiments also include a suspending agent and/or a wetting agent. This bulk blend is uniformly subdivided into unit dosage packaging or multi-dosage packaging units.

[00412] In still other embodiments, effervescent powders are also prepared. Effervescent salts have been used to disperse medicines in water for oral administration.

[00413] In some embodiments, the pharmaceutical solid oral dosage forms are formulated to 20 provide a controlled release of the compound of Formula (I), Formula (II) or Formula (III).

Controlled release refers to the release of the compound of Formula (I), Formula (II) or Formula (III) from a dosage form in which it is incorporated according to a desired profile over an extended period of time. Controlled release profiles include, for example, sustained release, prolonged release, pulsatile release, and delayed release profiles. In contrast to immediate release 25 compositions, controlled release compositions allow delivery of an agent to a subject over an extended period of time according to a predetermined profile. Such release rates can provide therapeutically effective levels of agent for an extended period of time and thereby provide a longer period of pharmacologic response while minimizing side effects as compared to conventional rapid release dosage forms. Such longer periods of response provide for many 30 inherent benefits that are not achieved with the corresponding short acting, immediate release preparations.

[00414] In some embodiments, the solid dosage forms described herein are formulated as enteric coated delayed release oral dosage forms, i.e., as an oral dosage form of a pharmaceutical composition as described herein which utilizes an enteric coating to affect release in the small intestine or large intestine. In one aspect, the enteric coated dosage form is a compressed or

molded or extruded tablet/mold (coated or uncoated) containing granules, powder, pellets, beads or particles of the active ingredient and/or other composition components, which are themselves coated or uncoated. In one aspect, the enteric coated oral dosage form is in the form of a capsule containing pellets, beads or granules, which include a compound of Formula (I), Formula (II) or 5 Formula (III), that are coated or uncoated.

[00415] Any coatings should be applied to a sufficient thickness such that the entire coating does not dissolve in the gastrointestinal fluids at pH below about 5, but does dissolve at pH about 5 and above. Coatings are typically selected from any of the following:

[00416] Shellac - this coating dissolves in media of pH >7; Acrylic polymers - examples of 10 suitable acrylic polymers include methacrylic acid copolymers and ammonium methacrylate copolymers. The Eudragit series E, L, S, RL, RS and NE (Rohm Pharma) are available as solubilized in organic solvent, aqueous dispersion, or dry powders. The Eudragit series RL, NE, and RS are insoluble in the gastrointestinal tract but are permeable and are used primarily for colonic targeting. The Eudragit series E dissolve in the stomach. The Eudragit series L, L-30D 15 and S are insoluble in stomach and dissolve in the intestine; Poly Vinyl Acetate Phthalate (PVAP) - PVAP dissolves in pH >5, and it is much less permeable to water vapor and gastric fluids.

[00417] Conventional coating techniques such as spray or pan coating are employed to apply 20 coatings. The coating thickness must be sufficient to ensure that the oral dosage form remains intact until the desired site of topical delivery in the intestinal tract is reached.

[00418] In other embodiments, the formulations described herein are delivered using a pulsatile dosage form. A pulsatile dosage form is capable of providing one or more immediate release pulses at predetermined time points after a controlled lag time or at specific sites. Exemplary pulsatile dosage forms and methods of their manufacture are disclosed in U.S. Pat. Nos. 25 5,011,692, 5,017,381, 5,229,135, 5,840,329 and 5,837,284. In one embodiment, the pulsatile dosage form includes at least two groups of particles, (i.e. multiparticulate) each containing the formulation described herein. The first group of particles provides a substantially immediate dose of the compound of Formula (I), Formula (II) or Formula (III) upon ingestion by a mammal. The first group of particles can be either uncoated or include a coating and/or sealant. In one aspect, 30 the second group of particles comprises coated particles. The coating on the second group of particles provides a delay of from about 2 hours to about 7 hours following ingestion before release of the second dose. Suitable coatings for pharmaceutical compositions are described herein or known in the art.

[00419] In some embodiments, pharmaceutical formulations are provided that include particles of 35 a compound of Formula (I), Formula (II) or Formula (III) and at least one dispersing agent or

suspending agent for oral administration to a subject. The formulations may be a powder and/or granules for suspension, and upon admixture with water, a substantially uniform suspension is obtained.

[00420] In one aspect, liquid formulation dosage forms for oral administration are in the form of aqueous suspensions selected from the group including, but not limited to, pharmaceutically acceptable aqueous oral dispersions, emulsions, solutions, elixirs, gels, and syrups. See, e.g., Singh *et al.*, Encyclopedia of Pharmaceutical Technology, 2nd Ed., pp. 754-757 (2002). In addition to the particles of the compound of Formula (I), Formula (II) or Formula (III), the liquid dosage forms include additives, such as: (a) disintegrating agents; (b) dispersing agents; (c) wetting agents; (d) at least one preservative, (e) viscosity enhancing agents, (f) at least one sweetening agent, and (g) at least one flavoring agent. In some embodiments, the aqueous dispersions can further include a crystalline inhibitor.

[00421] Furthermore, pharmaceutical compositions optionally include one or more pH adjusting agents or buffering agents, including acids such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, sodium bicarbonate and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in an acceptable range.

[00422] Additionally, pharmaceutical compositions optionally include one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate.

[00423] Other pharmaceutical compositions optionally include one or more preservatives to inhibit microbial activity. Suitable preservatives include mercury-containing substances such as merfen and thiomersal; stabilized chlorine dioxide; and quaternary ammonium compounds such as benzalkonium chloride, cetyltrimethylammonium bromide and cetylpyridinium chloride.

[00424] In one embodiment, the aqueous suspensions and dispersions described herein remain in a homogenous state, as defined in The USP Pharmacists' Pharmacopeia (2005 edition, chapter 905), for at least 4 hours. In one embodiment, an aqueous suspension is re-suspended into a homogenous suspension by physical agitation lasting less than 1 minute. In still another embodiment, no agitation is necessary to maintain a homogeneous aqueous dispersion.

[00425] Examples of disintegrating agents for use in the aqueous suspensions and dispersions include, but are not limited to, a starch, e.g., a natural starch such as corn starch or potato starch, a pregelatinized starch, or sodium starch glycolate; a cellulose such as methylcrystalline

cellulose, methylcellulose, croscarmellose, or a cross-linked cellulose, such as cross-linked sodium carboxymethylcellulose, cross-linked carboxymethylcellulose, or cross-linked croscarmellose; a cross-linked starch such as sodium starch glycolate; a cross-linked polymer such as cespovidone; a cross-linked polyvinylpyrrolidone; alginate such as alginic acid or a salt 5 of alginic acid such as sodium alginate; a gum such as agar, guar, locust bean, Karaya, pectin, or tragacanth; sodium starch glycolate; bentonite; a natural sponge; a surfactant; a resin such as a cation-exchange resin; citrus pulp; sodium lauryl sulfate; sodium lauryl sulfate in combination starch; and the like.

[00426] In some embodiments, the dispersing agents suitable for the aqueous suspensions and 10 dispersions described herein include, for example, hydrophilic polymers, Tween[®] 60 or 80, PEG, polyvinylpyrrolidone, and the carbohydrate-based dispersing agents such as, for example, hydroxypropylcellulose and hydroxypropyl cellulose ethers, hydroxypropyl methylcellulose and hydroxypropyl methylcellulose ethers, carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethyl-cellulose phthalate, 15 hydroxypropylmethyl-cellulose acetate stearate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol (PVA), polyvinylpyrrolidone/vinyl acetate copolymer, 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde (also known as tyloxapol), poloxamers; and poloxamines. In other embodiments, the dispersing agent is selected from a group not comprising one of the following agents: hydrophilic polymers; 20 electrolytes; Tween[®] 60 or 80; PEG; polyvinylpyrrolidone (PVP); hydroxypropylcellulose and hydroxypropyl cellulose ethers; hydroxypropyl methylcellulose and hydroxypropyl methylcellulose ethers; carboxymethylcellulose sodium; methylcellulose; hydroxyethylcellulose; hydroxypropylmethyl-cellulose phthalate; hydroxypropylmethyl-cellulose acetate stearate; non-crystalline cellulose; magnesium aluminum silicate; triethanolamine; polyvinyl alcohol (PVA); 25 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde; poloxamers; or poloxamines.

[00427] Wetting agents suitable for the aqueous suspensions and dispersions described herein include, but are not limited to, cetyl alcohol, glycerol monostearate, polyoxyethylene sorbitan fatty acid esters (e.g., the commercially available Tweens[®] such as e.g., Tween 20[®] and Tween 30 80[®], and polyethylene glycols, oleic acid, glyceryl monostearate, sorbitan monooleate, sorbitan monolaurate, triethanolamine oleate, polyoxyethylene sorbitan monooleate, polyoxyethylene sorbitan monolaurate, sodium oleate, sodium lauryl sulfate, sodium docusate, triacetin, vitamin E TPGS, sodium taurocholate, simethicone, phosphotidylcholine and the like

[00428] Suitable preservatives for the aqueous suspensions or dispersions described herein 35 include, for example, potassium sorbate, parabens (e.g., methylparaben and propylparaben),

benzoic acid and its salts, other esters of parahydroxybenzoic acid such as butylparaben, alcohols such as ethyl alcohol or benzyl alcohol, phenolic compounds such as phenol, or quaternary compounds such as benzalkonium chloride. Preservatives, as used herein, are incorporated into the dosage form at a concentration sufficient to inhibit microbial growth.

5 [00429] Suitable viscosity enhancing agents for the aqueous suspensions or dispersions described herein include, but are not limited to, methyl cellulose, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, Plasdon[®] S-630, carbomer, polyvinyl alcohol, alginates, acacia, chitosans and combinations thereof. The concentration of the viscosity enhancing agent will depend upon the agent selected and the viscosity desired.

10 [00430] Examples of sweetening agents suitable for the aqueous suspensions or dispersions described herein include, for example, acacia syrup, acesulfame K, alitame, aspartame, chocolate, cinnamon, citrus, cocoa, cyclamate, dextrose, fructose, ginger, glycyrrhetinate, glycyrrhiza (licorice) syrup, monoammonium glyrrhizinate (MagnaSweet[®]), maltol, mannitol, menthol, neohesperidine DC, neotame, Prosweet[®] Powder, saccharin, sorbitol, stevia, sucralose, sucrose, 15 sodium saccharin, saccharin, aspartame, acesulfame potassium, mannitol, sucralose, tagatose, thaumatin, vanilla, xylitol, or any combination thereof.

[00431] In some embodiments, the liquid formulations also include inert diluents commonly used in the art, such as water or other solvents, solubilizing agents, and emulsifiers. Exemplary emulsifiers are ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, 20 benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, dimethylformamide, sodium lauryl sulfate, sodium doccuscate, cholesterol, cholesterol esters, taurocholic acid, phosphotidylcholine, oils, such as cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil, and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols, fatty acid esters of sorbitan, or mixtures of these substances, and the like.

25 [00432] Representative intranasal formulations are described in, for example, U.S. Pat. Nos. 4,476,116, 5,116,817 and 6,391,452. Formulations that include a compound of Formula (I), Formula (II) or Formula (III) are prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, fluorocarbons, and/or other solubilizing or dispersing agents known in the art. See, for example, Ansel, H. C. *et al.*, *Pharmaceutical Dosage Forms and Drug Delivery* 30 Systems, Sixth Ed. (1995). Preferably these compositions and formulations are prepared with suitable nontoxic pharmaceutically acceptable ingredients. These ingredients are known to those skilled in the preparation of nasal dosage forms and some of these can be found in REMINGTON: THE SCIENCE AND PRACTICE OF PHARMACY, 21st edition, 2005. The choice of suitable carriers is dependent upon the exact nature of the nasal dosage form desired, e.g., solutions, suspensions, ointments, or gels. Nasal dosage forms generally contain large 35

amounts of water in addition to the active ingredient. Minor amounts of other ingredients such as pH adjusters, emulsifiers or dispersing agents, preservatives, surfactants, gelling agents, or buffering and other stabilizing and solubilizing agents are optionally present. Preferably, the nasal dosage form should be isotonic with nasal secretions.

5 [00433] For administration by inhalation, a compound of Formula (I), Formula (II) or Formula (III) is formulated for use as an aerosol, a mist or a powder. Pharmaceutical compositions described herein are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or 10 other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, such as, by way of example only, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound described herein and a suitable powder base such as lactose or starch.

15 [00434] Buccal formulations that include a compound of Formula (I), Formula (II) or Formula (III) are administered using a variety of formulations known in the art. For example, such formulations include, but are not limited to, U.S. Pat. Nos. 4,229,447, 4,596,795, 4,755,386, and 5,739,136. In addition, the buccal dosage forms described herein can further include a 20 bioerodible (hydrolysable) polymeric carrier that also serves to adhere the dosage form to the buccal mucosa. For buccal or sublingual administration, the compositions may take the form of tablets, lozenges, or gels formulated in a conventional manner.

25 [00435] In some embodiments, compounds of Formula (I), Formula (II) and Formula (III) are prepared as transdermal dosage forms. In one embodiments, the transdermal formulations described herein include at least three components: (1) a formulation of a compound of Formula (I), Formula (II) or Formula (III); (2) a penetration enhancer; and (3) an aqueous adjuvant. In some embodiments the transdermal formulations include additional components such as, but not limited to, gelling agents, creams and ointment bases, and the like. In some embodiments, the transdermal formulation further include a woven or non-woven backing material to enhance absorption and prevent the removal of the transdermal formulation from the skin. In other embodiments, the transdermal formulations described herein can maintain a saturated or 30 supersaturated state to promote diffusion into the skin.

35 [00436] In one aspect, formulations suitable for transdermal administration of compounds described herein employ transdermal delivery devices and transdermal delivery patches and can be lipophilic emulsions or buffered, aqueous solutions, dissolved and/or dispersed in a polymer or an adhesive. In one aspect, such patches are constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents. Still further, transdermal delivery of the compounds

described herein can be accomplished by means of iontophoretic patches and the like. In one aspect, transdermal patches provide controlled delivery of the compound of Formula (I), Formula (II) or Formula (III). In one aspect, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a 5 rate controlling barrier to deliver the compound to the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

[00437] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is formulated into a pharmaceutical composition suitable for intramuscular, subcutaneous, or intravenous injection. In one aspect, formulations suitable for intramuscular, subcutaneous, or intravenous 10 injection include physiologically acceptable sterile aqueous or non-aqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and non-aqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (propyleneglycol, polyethylene-glycol, glycerol, cremophor and the like), suitable mixtures thereof, vegetable oils (such as olive 15 oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. In some embodiments, formulations suitable for subcutaneous injection also contain additives such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the growth of microorganisms can be ensured 20 by various antibacterial and antifungal agents, such as parabens, chlorobutanol, phenol, sorbic acid, and the like. In some cases it is desirable to include isotonic agents, such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, such as aluminum monostearate and gelatin.

[00438] For intravenous injections, compounds described herein are formulated in aqueous 25 solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. For other parenteral injections, appropriate formulations include aqueous or nonaqueous solutions, preferably with physiologically compatible buffers or excipients. Such 30 excipients are known.

[00439] Parenteral injections may involve bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The pharmaceutical composition described herein may be in a form suitable for parenteral injection as a sterile suspensions, solutions or emulsions in oily or aqueous 35 vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing

agents. In one aspect, the active ingredient is in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[00440] In certain embodiments, delivery systems for pharmaceutical compounds may be employed, such as, for example, liposomes and emulsions. In certain embodiments, compositions 5 provided herein can also include an mucoadhesive polymer, selected from among, for example, carboxymethylcellulose, carbomer (acrylic acid polymer), poly(methylmethacrylate), polyacrylamide, polycarbophil, acrylic acid/butyl acrylate copolymer, sodium alginate and dextran.

[00441] In some embodiments, the compounds described herein may be administered topically 10 and can be formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, medicated sticks, balms, creams or ointments. Such pharmaceutical compounds can contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

[00442] In some embodiments, the compounds of Formula (I), Formula (II) and Formula (III) are 15 formulated in rectal compositions such as enemas, rectal gels, rectal foams, rectal aerosols, suppositories, jelly suppositories, or retention enemas, containing conventional suppository bases such as cocoa butter or other glycerides, as well as synthetic polymers such as polyvinylpyrrolidone, PEG, and the like. In suppository forms of the compositions, a low-melting wax such as, but not limited to, a mixture of fatty acid glycerides, optionally in combination with 20 cocoa butter is first melted.

Methods of Dosing and Treatment Regimens

[00443] In one embodiment, the compounds of Formula (I), Formula (II) and Formula (III) are used in the preparation of medicaments for the treatment of LPA-dependent or LPA-mediated diseases or conditions. In addition, a method for treating any of the diseases or conditions 25 described herein in a subject in need of such treatment, involves administration of pharmaceutical compositions that include at least one compound of Formula (I), Formula (II) or Formula (III) or a pharmaceutically acceptable salt, pharmaceutically active metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said subject.

[00444] In certain embodiments, the compositions containing the compound(s) described herein 30 are administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest at least one of the symptoms of the disease or condition. Amounts effective for this use depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the 35

drugs, and the judgment of the treating physician. Therapeutically effective amounts are optionally determined by methods including, but not limited to, a dose escalation clinical trial.

[00445] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder or

5 condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. When used in a patient, effective amounts for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician. In one aspect, prophylactic treatments include
10 administering to a mammal, who previously experienced at least one symptom of the disease being treated and is currently in remission, a pharmaceutical composition comprising a compound of Formula (I), Formula (II) or Formula (III) in order to prevent a return of the symptoms of the disease or condition.

[00446] In certain embodiments wherein the patient's condition does not improve, upon the
15 doctor's discretion the administration of the compounds are administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[00447] In certain embodiments wherein a patient's status does improve, the dose of drug being administered may be temporarily reduced or temporarily suspended for a certain length of time
20 (*i.e.*, a "drug holiday"). In specific embodiments, the length of the drug holiday is between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, or more than 28 days. The dose reduction during a drug holiday is, by way of example only, by 10%-100%, including by way of example only 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%,
25 95%, and 100%.

[00448] In certain embodiments in which the patient is presented with a situation in which the activity of LPA needs to be enhanced, for example, to assist with wound healing, the dose of drug being administered may be temporarily reduced or temporarily suspended for a certain length of time (*i.e.*, a "drug diversion"). In specific embodiments, the length of the drug diversion
30 is between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, or more than 28 days. The dose reduction during a drug diversion is, by way of example only, by 10%-100%, including by way of example only 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, and 100%. Once the situation requiring enhanced activity of LPA is
35 alleviated, the normal dosing schedule is optionally reinstated.

[00449] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, in specific embodiments, the dosage or the frequency of administration, or both, is reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In certain embodiments, however, the patient 5 requires intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00450] The amount of a given agent that corresponds to such an amount varies depending upon factors such as the particular compound, disease condition and its severity, the identity (e.g., weight, sex) of the subject or host in need of treatment, but can nevertheless be determined according to the particular circumstances surrounding the case, including, e.g., the specific agent 10 being administered, the route of administration, the condition being treated, and the subject or host being treated. In general, however, doses employed for adult human treatment are typically in the range of 0.01mg-5000 mg per day. In one aspect, doses employed for adult human treatment are from about 1mg to about 1000 mg per day. In one embodiment, the desired dose is conveniently presented in a single dose or in divided doses administered simultaneously (or over 15 a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[00451] In one embodiment, the daily dosages appropriate for the compound of Formula (I), Formula (II) or Formula (III) described herein are from about 0.01 to about 10 mg/kg per body weight. In specific embodiments, an indicated daily dosage in a large mammal, including, but not 20 limited to, humans, is in the range from about 0.5 mg to about 1000 mg, conveniently administered in divided doses, including, but not limited to, up to four times a day. In one embodiment, the daily dosage is administered in extended release form. In certain embodiments, suitable unit dosage forms for oral administration comprise from about 1 to 500 mg active ingredient. In other embodiments, the daily dosage or the amount of active in the dosage form are 25 lower or higher than the ranges indicated herein, based on a number of variables in regard to an individual treatment regime. In various embodiments, the daily and unit dosages are altered depending on a number of variables including, but not limited to, the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the 30 practitioner.

[00452] Toxicity and therapeutic efficacy of such therapeutic regimens are determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ and the ED₅₀. The dose ratio between the toxic and therapeutic effects is the therapeutic index and it is expressed as the ratio between LD₅₀ and 35 ED₅₀. In certain embodiments, the data obtained from cell culture assays and animal studies are

used in formulating the therapeutically effective daily dosage range and/or the therapeutically effective unit dosage amount for use in mammals, including humans. In some embodiments, the daily dosage amount of the compounds described herein lies within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. In certain embodiments, the daily dosage range and/or the unit dosage amount varies within this range depending upon the dosage form employed and the route of administration utilized.

5 **Patient Selection**

[00453] In any of the aforementioned aspects involving the prevention or treatment of LPA-mediated diseases or conditions are further embodiments comprising identifying patients by screening for LPA receptor gene SNPs. A SNP located in the promoter region of LPA₁ showed significant association with knee osteoarthritis in two independent populations (Mototani *et al.* *Hum. Mol. Genetics*, vol. 17, no. 12, 2008). Patients can be further selected based on increased LPA receptor expression in the tissue of interest. For example, chronic lymphocytic leukemia (CLL) is characterized by the accumulation of CD19+/CD5+ B-lymphocytes in the peripheral blood, bone marrow and lymphoid organs which occurs as a result of a block in B-lymphocyte apoptosis. LPA can protect some CLL cells from apoptosis and the cells that are protected by LPA have high levels of LPA₁ mRNA. In some embodiments, CLL patients are selected based on the expression of the LPA1R. LPA receptor expression are determined by methods including, but not limited to, northern blotting, western blotting, quantitative PCR (qPCR), flow cytometry, autoradiography (using a small molecule radioligand or PET ligand). In some embodiments, patients are selected based on the concentration of serum or tissue LPA measured by mass spectrometry. LPA concentrations are high in ovarian cancer ascites and in some breast cancer effusions. In some embodiments, patients are selected based on a combination of the above markers (increased LPA concentrations and increased LPA receptor expression).

25 **Combination Treatments**

[00454] In certain instances, it is appropriate to administer at least one compound of Formula (I), Formula (II) or Formula (III) in combination with another therapeutic agent. By way of example only, if one of the side effects experienced by a patient upon receiving one of the compounds herein is inflammation, then it may be appropriate to administer an anti-inflammatory agent in combination with the initial therapeutic agent. Or in another example, a patient is presented with a situation in which antagonism of LPA receptors provides potential harm, for example, if the patient is wounded, antagonism of LPA receptors may lead to a delay in wound healing. In such an event, in certain embodiments, the patient benefits by administration of a local wound-healing agent (at the site of the wound) in combination with the co-existing administration of a compound of Formula (I), Formula (II) or Formula (III).

[00455] Or, in one embodiment, the therapeutic effectiveness of one of the compounds described herein is enhanced by administration of an adjuvant (*i.e.*, by itself the adjuvant may have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, in some embodiments, the benefit experienced by a patient

5 is increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit.

[00456] In one specific embodiment, a compound of Formula (I), Formula (II) or Formula (III) is co-administered with a second therapeutic agent, wherein the compound of Formula (I), Formula (II) or Formula (III) and the second therapeutic agent modulate different aspects of the disease,

10 disorder or condition being treated, thereby providing a greater overall benefit than administration of either therapeutic agent alone.

[00457] In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

15 [00458] In certain embodiments, different therapeutically-effective dosages of the compounds disclosed herein will be utilized in formulating pharmaceutical composition and/or in treatment regimens when the compounds disclosed herein are administered in combination with one or more additional agent, such as an additional therapeutically effective drug, an adjuvant or the like. Therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens can be determined by means similar to those set forth hereinabove for the actives themselves. Furthermore, the methods of prevention/treatment described herein encompass the use of metronomic dosing, *i.e.*, providing more frequent, lower doses in order to minimize toxic side effects. In some embodiments, a combination treatment regimen encompasses treatment regimens in which administration of a compound of Formula (I), Formula (II) or Formula (III) is initiated prior to, during, or after treatment with a second agent described herein, and continues until any time during treatment with the second agent or after termination of treatment with the second agent. It also includes treatments in which a compound of Formula (I), Formula (II) or Formula (III) and the second agent being used in combination are administered simultaneously or at different times and/or at decreasing or increasing intervals

20 during the treatment period. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient.

25

[00459] Compositions and methods for combination therapy are provided herein. In accordance with one aspect, the pharmaceutical compositions disclosed herein are used to treat LPA-dependent or LPA-mediated conditions.

[00460] It is understood that the dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, is modified in accordance with a variety of factors. These factors include the disease, disorder or condition from which the subject suffers, as well as the age, weight, sex, diet, and medical condition of the subject. Thus, in some instances, the dosage regimen actually employed varies and, in some embodiments, deviates from the dosage regimens set forth herein.

[00461] For combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In additional embodiments, when co-administered with one or more other therapeutic agents, the compound provided herein is administered either simultaneously with the one or more other therapeutic agents, or sequentially.

[00462] In combination therapies, the multiple therapeutic agents (one of which is one of the compounds described herein) are administered in any order or even simultaneously. If administration is simultaneous, the multiple therapeutic agents are, by way of example only, provided in a single, unified form, or in multiple forms (e.g., as a single pill or as two separate pills). In one embodiment, one of the therapeutic agents is given in multiple doses, and in another, two (or more if present) are given as multiple doses. In some embodiments of non-simultaneous administration, the timing between the multiple doses vary from more than zero weeks to less than four weeks. In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents; the use of multiple therapeutic combinations is also envisioned.

[00463] The compounds of Formula (I), Formula (II) and Formula (III) and combination therapies are administered before, during or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound varies. Thus, in one embodiment, the compounds described herein are used as a prophylactic and are administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. In another embodiment, the compounds and compositions are administered to a subject during or as soon as possible after the onset of the symptoms. In specific embodiments, a compound described herein is administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease. In some embodiments, the length required for treatment varies, and the treatment length is adjusted to suit the specific needs of each subject. For example, in specific embodiments, a compound described herein or a formulation containing the compound is administered for at least 2 weeks, about 1 month to about 5 years.

[00464] By way of example, therapies which combine a compound of Formula (I), Formula (II) or Formula (III) with inhibitors of LPA synthesis or LPA receptor antagonists, either acting at the same or other points in the LPA synthesis or signalling pathway, are encompassed herein for treating LPA-dependent or LPA-mediated diseases or conditions.

5 **Exemplary Agent for use in Combination with Compounds of Formula (I), Formula (II) and Formula (III)**

[00465] In another embodiment described herein, methods for treatment of LPA-dependent or LPA-mediated conditions or diseases, such as proliferative disorders, including cancer, comprises administration to a mammal a compound of Formula (I), Formula (II) or Formula (III)

10 in combination with at least one additional agent selected, by way of example only, alemtuzumab, arsenic trioxide, asparaginase (pegylated or non-), bevacizumab, cetuximab, platinum-based compounds such as cisplatin, cladribine, daunorubicin/doxorubicin/idarubicin, irinotecan, fludarabine, 5-fluorouracil, gemtuzumab, methotrexate, PaclitaxelTM, taxol, temozolamide, thioguanine, or classes of drugs including hormones (an antiestrogen, an 15 antiandrogen, or gonadotropin releasing hormone analogues, interferons such as alpha interferon, nitrogen mustards such as busulfan or melphalan or mechlorethamine, retinoids such as tretinoin, topoisomerase inhibitors such as irinotecan or topotecan, tyrosine kinase inhibitors such as gefinitinib or imatinib, or agents to treat signs or symptoms induced by such therapy including allopurinol, filgrastim, granisetron/ondansetron/palonosetron, dronabinol).

20 [00466] In one aspect, the compound of Formula (I), Formula (II) or Formula (III) is administered or formulated in combination with one or more anti-cancer agents. In some embodiments, one or more of the anti-cancer agents are proapoptotic agents. Examples of anti-cancer agents include, but are not limited to, any of the following: gossypol, genasense, polyphenol E, Chlorofusin, all trans-retinoic acid (ATRA), bryostatin, tumor necrosis factor-related apoptosis-inducing ligand (TRAIL), 5-aza-2'-deoxycytidine, all trans retinoic acid, doxorubicin, vincristine, etoposide, gemcitabine, imatinib, geldanamycin, 17-N-Allylamino-17-Demethoxygeldanamycin (17-AAG), flavopiridol, LY294002, bortezomib, trastuzumab, BAY 11-7082, PKC412, or PD184352, TaxolTM (paclitaxel), and analogs of TaxolTM, such as TaxotereTM. Compounds that have the basic taxane skeleton as a common structure feature, have also been shown to have the ability to 25 arrest cells in the G2-M phases due to stabilized microtubules and may be useful for treating cancer in combination with the compounds described herein.

30 [00467] Further examples of anti-cancer agents for use in combination with the compounds of Formula (I), Formula (II) and Formula (III) include inhibitors of mitogen-activated protein kinase signaling, e.g., U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063,

SP600125, BAY 43-9006, wortmannin, or LY294002; Syk inhibitors; mTOR inhibitors; and antibodies (e.g., rituxan).

[00468] Other anti-cancer agents for use in combination with the compounds of Formula (I), Formula (II) and Formula (III) include one or more of the following: Adriamycin, Dactinomycin,

5 Bleomycin, Vinblastine, Cisplatin, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone;

10 caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitucin;

15 enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; iimofosine; interleukin II (including recombinant interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1; interferon alfa-n3; interferon beta-1 a; interferon gamma-1 b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprolol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril;

20 mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazoic; nogalamycin; ormaplatin; oxisuran; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin;

25 prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprime; thioguanine; thioteplatin; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate;

trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride.

5 **[00469]** Yet other anticancer agents for use in combination with the compounds of Formula (I), Formula (II) and Formula (III) include alkylating agents, antimetabolites, natural products, or hormones, e.g., nitrogen mustards (e.g., mechloroethamine, cyclophosphamide, chlorambucil, etc.), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomusitne, etc.), or triazenes (decarbazine, etc.). Examples of antimetabolites include but are not limited to folic acid analog 10 (e.g., methotrexate), or pyrimidine analogs (e.g., Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin).

15 **[00470]** Examples of natural products for use in combination with the compounds of Formula (I), Formula (II) and Formula (III) include but are not limited to vinca alkaloids (e.g., vinblastin, vincristine), epipodophyllotoxins (e.g., etoposide), antibiotics (e.g., daunorubicin, doxorubicin, bleomycin), enzymes (e.g., L-asparaginase), or biological response modifiers (e.g., interferon alpha).

20 **[00471]** Examples of alkylating agents for use in combination with the compounds of Formula (I), Formula (II) and Formula (III) include, but are not limited to, nitrogen mustards (e.g., mechloroethamine, cyclophosphamide, chlorambucil, meiphalan, etc.), ethylenimine and methylmelamines (e.g., hexamethylmelamine, thiotepa), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomusitne, semustine, streptozocin, etc.), or triazenes (decarbazine, etc.). Examples of antimetabolites include, but are not limited to folic acid analog 25 (e.g., methotrexate), or pyrimidine analogs (e.g., fluorouracil, floxouridine, Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin).

25 **[00472]** Examples of hormones and antagonists for use in combination with the compounds of Formula (I), Formula (II) and Formula (III) include, but are not limited to, adrenocorticosteroids (e.g., prednisone), progestins (e.g., hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate), estrogens (e.g., diethylstilbestrol, ethinyl estradiol), antiestrogen (e.g., tamoxifen), androgens (e.g., testosterone propionate, fluoxymesterone), antiandrogen (e.g., 30 flutamide), gonadotropin releasing hormone analog (e.g., leuprolide). Other agents that can be used in the methods and compositions described herein for the treatment or prevention of cancer include platinum coordination complexes (e.g., cisplatin, carboblatin), anthracenedione (e.g., mitoxantrone), substituted urea (e.g., hydroxyurea), methyl hydrazine derivative (e.g., procarbazine), adrenocortical suppressant (e.g., mitotane, aminoglutethimide).

[00473] Examples of anti-cancer agents which act by arresting cells in the G2-M phases due to stabilized microtubules include without limitation the following marketed drugs and drugs in development: Erbulozole, Dolastatin 10, Mivobulin isethionate, Vincristine, NSC-639829, Discodermolide, ABT-751, Altorhyrtins (such as Altorhyrtin A and Altorhyrtin C), Spongistatins (such as Spongistatin 1, Spongistatin 2, Spongistatin 3, Spongistatin 4, Spongistatin 5, Spongistatin 6, Spongistatin 7, Spongistatin 8, and Spongistatin 9), Cemadotin hydrochloride, Epothilones (such as Epothilone A, Epothilone B, Epothilone C, Epothilone D, Epothilone E, Epothilone F, Epothilone B N-oxide, Epothilone A N-oxide, 16-aza-epothilone B, 21-aminoepothilone B, 21-hydroxyepothilone D, 26-fluoroepothilone, Auristatin PE, Soblidotin, Vincristine sulfate, Cryptophycin 52, Vitilevuamide, Tubulysin A, Canadensol, Centaureidin, Oncocidin A1 Fijianolide B, Laulimalide, Narcosine, Nascapine, Hemiasterlin, Vanadocene acetylacetone, Indanocine Eleutherobins (such as Desmethylleleutherobin, Desaetyleleutherobin, Isoeleutherobin A, and Z-Eleutherobin), Caribaeoside, Caribaeolin, Halichondrin B, Diazonamide A, Taccalonolide A, Diozostatin, (-)-Phenylahistin, Myoseverin B, Resverastatin phosphate sodium.

[00474] In one aspect, a compound of Formula (I), Formula (II) or Formula (III) is co-administered with thrombolytic agents (e.g., alteplase anistreplase, streptokinase, urokinase, or tissue plasminogen activator), heparin, tinzaparin, warfarin, dabigatran (e.g., dabigatran etexilate), factor Xa inhibitors (e.g., fondaparinux, draparinux, rivaroxaban, DX-9065a, otamixaban, LY517717, or YM150), ticlopidine, clopidogrel, CS-747 (prasugrel, LY640315), ximelagatran, or BIBR 1048.

[00475] In some embodiments, a compound of Formula (I), Formula (II) or Formula (III) is used in combination with anti-emetic agents to treat nausea or emesis, which may result from the use of a compound of Formula (I), Formula (II) or Formula (III), anti-cancer agent(s) and/or radiation therapy.

[00476] Anti-emetic agents include, but are not limited to: neurokinin-1 receptor antagonists, 5HT3 receptor antagonists (such as ondansetron, granisetron, tropisetron, Palonosetron, and zatisetron), GABA_B receptor agonists (such as baclofen), corticosteroids (such as dexamethasone, prednisone, prednisolone, or others), dopamine antagonists (such as, but not limited to, domperidone, droperidol, haloperidol, chlorpromazine, promethazine, prochlorperazine, metoclopramide), antihistamines (H1 histamine receptor antagonists, such as but not limited to, cyclizine, diphenhydramine, dimenhydrinate, meclizine, promethazine, hydroxyzine), cannabinoids (such as but not limited to, cannabis, marinol, dronabinol), and others (such as, but not limited to, trimethobenzamide; ginger, emetrol, propofol).

[00477] In some embodiments, a compound of Formula (I), Formula (II) or Formula (III) is used in combination with an agent useful in the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis receptor activator (such as epoetin- α).

[00478] In some embodiments, a compound of Formula (I), Formula (II) or Formula (III) is used in combination with an agent useful in the treatment of neutropenia. Examples of agents useful in the treatment of neutropenia include, but are not limited to, a hematopoietic growth factor which regulates the production and function of neutrophils such as a human granulocyte colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

[00479] In some embodiments, a compound of Formula (I), Formula (II) or Formula (III) is used in combination with radiation therapy (or radiotherapy). Radiation therapy is the treatment of cancer and other diseases with ionizing radiation. Radiation therapy can be used to treat localized solid tumors, such as cancers of the skin, tongue, larynx, brain, breast, prostate, colon, uterus and/or cervix. It can also be used to treat leukemia and lymphoma (cancers of the blood-forming cells and lymphatic system, respectively).

[00480] A technique for delivering radiation to cancer cells is to place radioactive implants directly in a tumor or body cavity. This is called internal radiotherapy (brachytherapy, interstitial irradiation, and intracavitary irradiation are types of internal radiotherapy.) Using internal radiotherapy, the radiation dose is concentrated in a small area, and the patient stays in the hospital for a few days. Internal radiotherapy is frequently used for cancers of the tongue, uterus, prostate, colon, and cervix.

[00481] The term “radiotherapy” or “ionizing radiation” include all forms of radiation, including but not limited to α , β , and γ radiation and ultraviolet light.

Immunosuppressants

[00482] In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are to treat or reduce fibrosis in a mammal. In one aspect, compounds of Formula (I), Formula (II) and Formula (III) are administered in combination with one or more immunosuppressants. Immunosuppressive therapy is clinically used to treat or prevent the rejection of transplanted organs and tissues (e.g. bone marrow, heart, kidney, liver); treatment of autoimmune diseases or diseases that are most likely of autoimmune origin (e.g. rheumatoid arthritis, myasthenia gravis, systemic lupus erythematosus, Crohn's disease, and ulcerative colitis); and treatment of some other non-autoimmune inflammatory diseases (e.g. long term allergic asthma control), and in the treatment of fibrotic conditions.

[00483] In some embodiments, a compound of Formula (I), Formula (II) or Formula (III) is administered with corticosteroids. In some embodiments, a compound of Formula (I), Formula (II) or Formula (III) is administered with an a therapeutic agent selected from among:

Calcineurin inhibitors (such as, but not limited to, cyclosporin, tacrolimus); mTOR inhibitors (such as, but not limited to, sirolimus, everolimus); anti-proliferatives (such as, but not limited to, azathioprine, mycophenolic acid); corticosteroids (such as, but not limited to, prednisone, cortisone acetate, prednisolone, methylprednisolone, dexamethasone, betamethasone, 5 triamcinolone, beclometasone, fludrocortisone acetate, deoxycorticosterone acetate, aldosterone, hydrocortisone); antibodies (such as, but not limited to, monoclonal anti-IL-2R α receptor antibodies (basiliximab, daclizumab), polyclonal anti-T-cell antibodies (anti-thymocyte globulin (ATG), anti-lymphocyte globulin (ALG)), B-cell antagonists, rituximab, natalizumab).

[00484] Other therapeutic agents include, but are not limited to: cyclophosphamide, penicillamine, 10 cyclosporine, nitrosoureas, cisplatin, carboplatin, oxaliplatin, methotrexate, azathioprine, mercaptopurine, pyrimidine analogues, protein synthesis inhibitors, dactinomycin, anthracyclines, mitomycin C, bleomycin, mithramycin, Atgam^(R), Thymoglobuline^(R), OKT3^(R), basiliximab, daclizumab, cyclosporin, tacrolimus, sirolimus, Interferons (IFN- β , IFN- γ), opioids, TNF binding proteins (infliximab, etanercept, adalimumab, golimumab), leflunomide, gold 15 thioglucose, gold thiomalate, aurofin, sulfasalazine, hydroxychloroquine, minocycline, rapamicin, mycophenolic acid, mycophenolate mofetil, FTY720, as well as those listed in US 7,060,697.

[00485] In one embodiment, a compound of Formula (I), Formula (II) or Formula (III) is administered in combination with Cyclosporin A (CsA) or tacrolimus (FK506). In one 20 embodiment, a compound of Formula (I), Formula (II) or Formula (III) is administered to a mammal in combination with an anti-inflammatory agent including, but not limited to, non-steroidal anti-inflammatory drugs (NSAIDs) and corticosteroids (glucocorticoids).

[00486] NSAIDs include, but are not limited to: aspirin, salicylic acid, gentisic acid, choline magnesium salicylate, choline salicylate, choline magnesium salicylate, choline salicylate, 25 magnesium salicylate, sodium salicylate, diflunisal, carprofen, fenoprofen, fenoprofen calcium, flurobiprofen, ibuprofen, ketoprofen, nabutone, ketorolac, ketorolac tromethamine, naproxen, oxaprozin, diclofenac, etodolac, indomethacin, sulindac, tolmetin, meclofenamate, meclofenamate sodium, mefenamic acid, piroxicam, meloxicam, COX-2 specific inhibitors (such as, but not limited to, celecoxib, rofecoxib, valdecoxib, parecoxib, etoricoxib, lumiracoxib, CS- 30 502, JTE-522, L-745,337 and NS398).

[00487] Corticosteroids, include, but are not limited to: betamethasone, prednisone, alclometasone, aldosterone, amcinonide, beclometasone, betamethasone, budesonide, ciclesonide, clobetasol, clobetasone, clocortolone, cloprednol, cortisone, cortivazol, deflazacort, deoxycorticosterone, desonide, desoximetasone, desoxycortone, dexamethasone, diflorasone, 35 diflucortolone, difluprednate, fluclorolone, fludrocortisone, fludroxycoertide, flumetasone,

flunisolide, fluocinolone acetonide, fluocinonide, fluocortin, fluocortolone, fluorometholone, fluperolone, fluprednidene, fluticasone, formocortal, halcinonide, halometasone, hydrocortisone/cortisol, hydrocortisone aceponate, hydrocortisone buteprate, hydrocortisone butyrate, loteprednol, medrysone, meprednisone, methylprednisolone, methylprednisolone aceponate, mometasone furoate, paramethasone, prednicarbate, prednisone/prednisolone, rimexolone, tixocortol, triamcinolone, and ulobetasol.

[00488] In one embodiment, a compound of Formula (I), Formula (II) or Formula (III) is administered in combination with leukotriene receptor antagonists including, but are not limited to, BAY u9773 (see EP 00791576; published 27 Aug 1997), DUO-LT (Tsuji *et al*, *Org. Biomol. Chem.*, 1, 3139-3141, 2003), zafirlukast, montelukast, prankulast, and derivatives or analogs thereof.

Other Combination Therapies

[00489] In another embodiment described herein, methods for treatment of LPA-dependent or LPA-mediated conditions or diseases, such as atherosclerosis, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected, by way of example only, HMG-CoA reductase inhibitors (e.g., statins in their lactonized or dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof, including but not limited to lovastatin; simvastatin; dihydroxy open-acid simvastatin, particularly the ammonium or calcium salts thereof; pravastatin, particularly the sodium salt thereof; fluvastatin, particularly the sodium salt thereof; atorvastatin, particularly the calcium salt thereof; nisvastatin, also referred to as NK-104; rosuvastatin); agents that have both lipid-altering effects and other pharmaceutical activities; HMG-CoA synthase inhibitors; cholesterol absorption inhibitors such as ezetimibe; cholesterol ester transfer protein (CETP) inhibitors, for example JTT-705 and CP529, 414; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors); acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of ACAT-1 or ACAT-2 as well as dual inhibitors of ACAT-1 and-2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; platelet aggregation inhibitors, for example glycoprotein IIb/IIIa fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR γ) agonists, including the compounds commonly referred to as glitazones, for example troglitazone, pioglitazone and rosiglitazone and including those compounds included within the structural class known as thiazolidinediones as well as those PPAR γ agonists outside the thiazolidinedione structural class; PPAR α agonists such as clofibrate, fenofibrate including micronized fenofibrate, and gemfibrozil ; PPAR dual α/γ agonists such as 5-[(2, 4-dioxo-5-

thiazolidinyl)methyl]-2-methoxy-N-[[4-(trifluoromethyl)phenyl]methyl]-benzamide, known as KRP-297; vitamin B6 (also known as pyridoxine) and the pharmaceutically acceptable salts thereof such as the HCl salt; vitamin B12 (also known as cyanocobalamin); folic acid or a pharmaceutically acceptable salt or ester thereof such as the sodium salt and the

5 methylglucamine salt; anti-oxidant vitamins such as vitamin C and E and beta carotene; beta-blockers; angiotensin II antagonists such as losartan; angiotensin converting enzyme inhibitors such as enalapril and captopril; calcium channel blockers such as nifedipine and diltiazam; endothelial antagonists; agents that enhance ABC1 gene expression; FXR and LXR ligands including both inhibitors and agonists; bisphosphonate compounds such as alendronate sodium; 10 and cyclooxygenase-2 inhibitors such as rofecoxib and celecoxib.

[00490] In another embodiment described herein, methods for treatment of LPA-dependent or LPA-mediated conditions or diseases, such as the therapy of stroke, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from, by way of example only, COX-2 15 inhibitors; nitric oxide synthase inhibitors, such as N-(3-(aminomethyl)benzyl) acetamidine; Rho kinase inhibitors, such as fasudil; angiotension II type-1 receptor antagonists, including candesartan, losartan, irbesartan, eprosartan, telmisartan and valsartan; glycogen synthase kinase 3 inhibitors; sodium or calcium channel blockers, including crobenetine; p38 MAP kinase inhibitors, including SKB 239063; thromboxane AX- synthetase inhibitors, including isbogrel, 20 ozagrel, ridogrel and dazoxiben; statins (HMG CoA reductase inhibitors), including lovastatin, simvastatin, dihydroxy open-acid simvastatin, pravastatin, fluvastatin, atorvastatin, nisvastatin, and rosuvastatin; neuroprotectants, including free radical scavengers, calcium channel blockers, excitatory amino acid antagonists, growth factors, antioxidants, such as edaravone, vitamin C, TROLOX™, citicoline and minicycline, and reactive astrocyte inhibitors, such as (2R)-2- 25 propyloctanoic acid; beta andrenergic blockers, such as propranolol, nadolol, timolol, pindolol, labetalol, metoprolol, atenolol, esmolol and acebutolol; NMDA receptor antagonists, including memantine; NR2B antagonists, such as traxoprodil; 5-HT1A agonists; receptor platelet fibrinogen receptor antagonists, including tirofiban and lamifiban; thrombin inhibitors; antithrombotics, such as argatroban; antihypertensive agents, such as enalapril; vasodilators, such as cyclandelate; 30 nociceptin antagonists; DPIV antagonists; GABA 5 inverse agonists; and selective androgen receptor modulators.

[00491] In another embodiment described herein, methods for treatment of LPA-dependent or LPA-mediated conditions or diseases, such as the therapy of interstitial cystitis, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described

herein in combination with at least one additional agent selected from, by way of example only, dimethylsulfoxide, omalizumab, and pentosan polysulfate.

[00492] In yet another embodiment described herein, methods for treating LPA-dependent or LPA-mediated conditions or diseases, such as the therapy of respiratory disorders (e.g., asthma,

5 COPD and rhinitis), comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one agent used in the treatment of respiratory conditions. Agents used in the treatment of respiratory conditions include, but are not limited to, bronchodilators (e.g., sympathomimetic agents and xanthine derivatives), leukotriene receptor antagonists, leukotriene formation inhibitors, leukotriene modulators, nasal decongestants, respiratory enzymes, lung surfactants, antihistamines (e.g., mepyramine (pyrilamine), antazoline, diphenhydramine, carboxamine, doxylamine, clemastine, dimenhydrinate, pheniramine, chlorphenamine (chlorpheniramine), dexchlorpheniramine, brompheniramine, triprolidine, cetirizine, cyclizine, chlorcyclizine, hydroxyzine, meclizine, loratadine, desloratadine, promethazine, alimemazine (trimeprazine), cyproheptadine, azatadine, 10 ketotifen, acrivastine, astemizole, cetirizine, mizolastine, terfenadine, azelastine, levocabastine, olopatadine, levocetirizine, fexofenadine), mucolytics, corticosteroids, anticholinergics, antitussives, analgesics, expectorants, albuterol, ephedrine, epinephrine, fomoterol, metaproterenol, terbutaline, budesonide, ciclesonide, dexamethasone, flunisolide, fluticasone propionate, triamcinolone acetonide, ipratropium bromide, pseudoephedrine, theophylline, 15 montelukast, zafirlukast, ambrisentan, bosentan, enrasentan, sitaxsentan, tezosentan, iloprost, treprostinil, pirfenidone, 5-lipoxygenase-activating protein (FLAP) inhibitors, FLAP modulators and 5-LO inhibitors.

[00493] In a specific embodiment described herein, methods for treating LPA-dependent or LPA-mediated conditions or diseases, such as the therapy of asthma and/or COPD, comprises

25 administration to a patient anti-inflammatory agents. In certain embodiments, methods for treating LPA-dependent or LPA-mediated conditions or diseases, such as the therapy of asthma and/or COPD, comprise administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from, but not limited to, epinephrine, isoproterenol, orciprenaline, bronchodilators, glucocorticoids, 30 leukotriene modifiers, mast-cell stabilizers, xanthines, anticholinergics, β -2 agonists, FLAP inhibitors, FLAP modulators or 5-LO inhibitors. β -2 agonists include, but are not limited to, short-acting β -2 agonists (e.g., salbutamol (albuterol), levalbuterol, terbutaline, pirbuterol, procaterol, metaproterenol, fenoterol and bitolterol mesylate) and long-acting β -2 agonists (e.g., salmeterol, formoterol, bambuterol and clenbuterol). FLAP inhibitors and/or FLAP modulators 35 include, but are not limited to, 3-[3-tert-butylsulfanyl-1-[4-(6-methoxy-pyridin-3-yl)-benzyl]-5-

(pyridin-2-ylmethoxy)-1H-indol-2-yl]-2,2-dimethyl-propionic acid, 3-[3-*tert*-butylsulfanyl-1-[4-(6-ethoxy-pyridin-3-yl)-benzyl]-5-(5-methyl-pyridin-2-ylmethoxy)-1H-indol-2-yl]-2,2-dimethyl-propionic acid, MK-886, MK-0591, BAY-x1005 and compounds found in US 2007/0225285, US 2007/0219206, US 2007/0173508, US 2007/0123522 and US 2007/0105866 (each of which are

5 hereby incorporated by reference). Glucocorticoids include, but are not limited to, beclometasone, budesonide, ciclesonide, fluticasone and mometasone. Anticholinergics include, but are not limited to, ipratropium and tiotropium. Mast cell stabilizers include, but are not limited to, cromoglicate and nedocromil. Xanthines include, but are not limited to, amminophylline, theobromine and theophylline. Leukotriene antagonists include, but are not 10 limited to, montelukast, tomelukast, pranlukast and zafirlukast. 5-LO inhibitors include, but are not limited to, zileuton, VIA-2291 (ABT761), AZ-4407 and ZD-2138 and compounds found in US 2007/0149579, WO2007/016784.

[00494] In another specific embodiment described herein, methods for treating LPA-dependent or LPA-mediated conditions or diseases, such as the therapy of allergic diseases or conditions, 15 comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from, by way of example only, antihistamines, leukotriene antagonists, corticosteroids and decongestants.

Leukotriene antagonists include, but are not limited to, montelukast, tomelukast, pranlukast and zafirlukast.

[00495] In one aspect, LPA receptor antagonists described herein are administered in combination with one or more agents used to treat used to treat asthma, including, but not limited to: combination inhalers (fluticasone and salmeterol oral inhalation (e.g. Advair)); inhaled Beta-2 agonists (albuterol inhaler; albuterol nebulizer solution; formoterol; isoproterenol oral inhalation; levalbuterol; metaproterenol inhalation; pirbuterol acetate oral inhalation; salmeterol aerosol inhalation; salmeterol powder inhalation; terbutaline inhaler); inhaled corticosteroids (beclomethasone oral inhalation; budesonide inhalation solution; budesonide inhaler; flunisolide oral inhalation; fluticasone inhalation aerosol; fluticasone powder for oral inhalation; mometasone inhalation powder; triamcinolone oral inhalation); leukotriene modifiers (montelukast; zafirlukast; zileuton); mast cell stabilizers (cromolyn inhaler; nedocromil oral inhalation); monoclonal antibodies (omalizumab); oral Beta-2 agonists (albuterol oral syrup; albuterol oral tablets; metaproterenol; terbutaline); bronchodilator (aminophylline; oxtriphylline; theophylline).

[00496] In one aspect, LPA receptor antagonists described herein are administered in combination with one or more agents used to treat allergy, including, but not limited to: antihistamine and decongestant combinations (cetirizine and pseudoephedrine; desloratadine and pseudoephedrine

ER; fexofenadine and pseudoephedrine; loratadine and pseudoephedrine); antihistamines (azelastine nasal spray; brompheniramine; brompheniramine oral suspension; carbinoxamine; cetirizine; chlorpheniramine; clemastine; desloratadine; dexchlorpheniramine ER; dexchlorpheniramine oral syrup; diphenhydramine oral; fexofenadine; loratadine; promethazine);
5 decongestants (pseudoephedrine); leukotriene modifiers (montelukast; montelukast granules); nasal anticholinergics (ipratropium); nasal corticosteroids (beclomethasone nasal inhalation; budesonide nasal inhaler; flunisolide nasal inhalation; fluticasone nasal inhalation; mometasone nasal spray; triamcinolone nasal inhalation; triamcinolone nasal spray); nasal decongestants (phenylephrine); nasal mast cell stabilizers (cromolyn nasal spray).

10 [00497] In one aspect, LPA receptor antagonists described herein are administered in combination with one or more agents used to treat chronic obstructive pulmonary disease (COPD), including, but not limited to: anticholinergics - ipratropium bromide oral inhalation); combination Inhalers (albuterol and ipratropium (e.g. Combivent, DuoNeb); fluticasone and salmeterol oral inhalation (e.g. Advair)); corticosteroids (dexamethasone tablets; fludrocortisone acetate; hydrocortisone
15 tablets; methylprednisolone; prednisolone liquid; prednisone oral; triamcinolone oral); inhaled Beta-2 Agonists (albuterol inhaler; albuterol nebulizer solution; formoterol; isoproterenol oral inhalation; levalbuterol; metaproterenol inhalation; pirbuterol acetate oral inhalation; salmeterol aerosol inhalation; salmeterol powder inhalation; terbutaline inhaler); inhaled Corticosteroids (beclomethasone oral inhalation; budesonide inhalation solution; budesonide inhaler; flunisolide
20 oral inhalation; fluticasone inhalation aerosol; fluticasone powder for oral inhalation; triamcinolone oral inhalation); mukolitics (guaifenesin); oral Beta-2 agonists (albuterol oral syrup; albuterol oral tablets; metaproterenol; terbutaline); bronchodilator (aminophylline; oxtriphylline; theophylline).

25 [00498] In one embodiment, compounds of Formula (I), Formula (II) and Formula (III) described herein are administered to a patient in combination with inhaled corticosteroids.

[00499] In one embodiment, compounds of Formula (I), Formula (II) and Formula (III) described herein are administered to a patient in combination with beta2-adrenergic receptor agonists. In one embodiment, compounds of Formula (I), Formula (II) and Formula (III) are administered to a patient in combination with short acting beta2-adrenergic receptor agonists. In one embodiment, 30 compounds of Formula (I), Formula (II) and Formula (III) are administered to a patient in combination with long-acting beta2-adrenergic receptor agonists.

Kits/Articles of Manufacture

[00500] For use in the therapeutic applications described herein, kits and articles of manufacture are also described herein. Such kits can comprise a carrier, package, or container that is 35 compartmentalized to receive one or more containers such as vials, tubes, and the like, each of

the container(s) comprising one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers are formed from any acceptable material including, e.g., glass or plastic.

[00501] For example, the container(s) can comprise one or more compounds described herein,

5 optionally in a composition or in combination with another agent as disclosed herein. The container(s) optionally have a sterile access port (for example the container can be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits optionally comprising a compound with an identifying description or label or instructions relating to its use in the methods described herein.

10 [00502] A kit will typically comprise one or more additional containers, each with one or more of various materials (such as reagents, optionally in concentrated form, and/or devices) desirable from a commercial and user standpoint for use of a compound described herein. Non-limiting examples of such materials include, but not limited to, buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and 15 package inserts with instructions for use. A set of instructions will also typically be included.

[00503] A label can be on or associated with the container. A label can be on a container when letters, numbers or other characters forming the label are attached, molded or etched into the container itself; a label can be associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. A label can be used to indicate 20 that the contents are to be used for a specific therapeutic application. The label can also indicate directions for use of the contents, such as in the methods described herein.


EXAMPLES

[00504] These examples are provided for illustrative purposes only and not to limit the scope of

25 the claims provided herein.


Synthesis of Compounds

(R)-2'-chloro-alpha-methylbenzyl alcohol




[00505] Using the procedure of Meier *et al* (Tetrahedron, 1996, 52, 589; Method 3), 2'-


30 chloroacetophenone (Aldrich) was reduced to give (R)-2'-chloro-alpha-methylbenzylalcohol in 92 %e.e. (HPLC analysis of the acetate derivative (made by reacting the benzyl alcohol with acetyl chloride and triethylamine in methylene chloride) using Chiralcel OD eluted with 99:1 Hexane:Ethanol. R isomer retention time 4.3 minutes)

(S)-2'-chloro-alpha-methylbenzyl alcohol


[00506] Using the procedure of Meier *et al* (Tetrahedron, 1996, 52, 589; Method 3), 2'-chloroacetophenone (Aldrich) was reduced to give (R)-2'-chloro-alpha-methylbenzylalcohol in 80 %e.e. (HPLC analysis of the acetate derivative (made by reacting the benzyl alcohol with acetyl chloride and triethylamine in methylene chloride) using Chiralcel OD eluted with 99:1 Hexane:Ethanol. S isomer retention time 5.3 minutes)

(R)-2'-fluoro-alpha-methylbenzyl alcohol


[00507] Using the procedure of Meier *et al* (Tetrahedron, 1996, 52, 589; Method 3), 2'-fluoroacetophenone (Aldrich) was reduced to give (R)-2'-fluoro-alpha-methylbenzylalcohol in 88.6 %e.e. (HPLC analysis of the acetate derivative (made by reacting the benzyl alcohol with acetyl chloride and triethylamine in methylene chloride) using Chiralcel OD eluted with 99.8:0.2 Hexane:Ethanol. R isomer retention time 5.9 minutes)

15 (S)-2'-fluoro-alpha-methylbenzyl alcohol

[00508] Using the procedure of Meier *et al* (Tetrahedron, 1996, 52, 589; Method 3), 2'-fluoroacetophenone (Aldrich) was reduced to give (S)-2'-fluoro-alpha-methylbenzylalcohol in 88.6 %e.e. (HPLC analysis of the acetate derivative (made by reacting the benzyl alcohol with acetyl chloride and triethylamine in methylene chloride) using Chiralcel OD eluted with 99.8:0.2 Hexane:Ethanol. S isomer retention time 6.7 minutes)


Example 1: Synthesis of 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1H-[1,2,4]triazole-3-carboxylic acid (Compound 1-1)**25 Step 1: 3-Methylamino-but-2-enoic acid methyl ester**

[00509] To a solution of methyl acetoacetate (29.4g, 253mmol) in MeOH (30mL) was added methylamine (33 wt% in EtOH; 48mL, 385mmol) dropwise at room temperature. The reaction was stirred for 1 hour, and then concentrated and dried to give the title compound as a white crystalline solid.




Step 2: 2-(4-Chloromethyl-benzoyl)-3-oxo-butyric acid methyl ester

[00510] To 3-methylamino-but-2-enoic acid methyl ester (18.5g, 143mmol) in THF (350mL) was added pyridine (17mL) dropwise. The mixture was cooled to 0°C, and 4-chloromethylbenzoyl chloride (27.5g, 146mmol) in THF (100mL) was added dropwise. The reaction was stirred at room temperature overnight, and then H₂O was added. The mixture was extracted with EtOAc, and the combined organic layers were washed with H₂O, dried over Na₂SO₄, filtered, and concentrated to give the title compound.



10 **Step 3: 5-(4-Chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester**


[00511] To a mixture of 2-(4-chloromethyl-benzoyl)-3-oxo-butyric acid methyl ester (35g, 131mmol) in acetic acid (110mL) was added hydroxylamine hydrochloride (8.9g, 127mmol), and the reaction was stirred at 115°C for 2 hours. After cooling, saturated aqueous NaHCO₃ was added to the mixture to adjust to pH 8. The solution was extracted with EtOAc, and the combined organic layers were washed three times with brine, dried over Na₂SO₄, filtered, and concentrated. The crude material was purified by silica gel chromatography (10-12% EtOAc in hexanes) to give the title compound.



Step 4: 5-(4-Chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid


20 [00512] Lithium hydroxide (17.8g, 424mmol) was added to a solution of 5-(4-chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester (22.5g, 84.9mmol) in 2:1 THF:H₂O (300mL), and the reaction was stirred for 2 days at room temperature. 1N Aqueous HCl was

added to adjust to pH 3, and the mixture was extracted with CHCl_3 . The combined organic layers were washed three times with brine, dried over Na_2SO_4 , filtered, and concentrated to give the title compound.




5 **Step 5: [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chlorophenyl)-ethyl ester**

[00513] 5-(4-Chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid (2.0g, 8.0mmol), 2-chloro-alpha-methylbenzyl alcohol (1.5g, 9.5mol), diphenylphosphoryl azide (3.3g, 12.0mmol), and triethylamine (2.2mL, 15.8mmol) were combined in toluene in a sealed container and stirred 10 at 120°C for 2 hours. After cooling, the mixture was concentrated, and the residue was purified by silica gel chromatography to give the title compound.



Step 6: 1-(4-{4-[1-(2-Chlorophenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1H-[1,2,4]triazole-3-carboxylic acid methyl ester

15 [00514] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chlorophenyl)-ethyl ester (0.340g, 0.84mmol), methyl 1,2,4-triazole-3-carboxylate (0.112g, 0.88mmol), and potassium carbonate (0.232g, 0.84mmol) were combined in DMF, and the reaction was stirred at 70°C for 3 hours. After aqueous worked-up, the crude material was used directly in the hydrolysis step.




Step 7: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1H-[1,2,4]triazole-3-carboxylic acid

[00515] 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-


5 1H-[1,2,4]triazole-3-carboxylic acid methyl ester (0.84mmol) was treated with lithium hydroxide (0.088g, 2.10mmol) in 2:1 THF:H₂O (6mL) overnight. The mixture was acidified to pH 3 and purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 481.

Example 2: Synthesis of 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1H-imidazole-4-carboxylic acid (Compound 1-2) and 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-3H-imidazole-4-carboxylic acid (Compound 1-3)



Step 1: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1H-imidazole-4-carboxylic acid methyl ester and 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-3H-imidazole-4-carboxylic acid methyl ester


[00516] Prepared according to the procedure described in Example 1, Step 6, using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and methyl 4-imidazolecarboxylate.



Step 2: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1*H*-imidazole-4-carboxylic acid and 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-3*H*-imidazole-4-carboxylic acid


5 [00517] Prepared according to the procedure described in Example 1, Step 7, using the following starting material: a mixture of 1-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1*H*-imidazole-4-carboxylic acid methyl ester and 3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-3*H*-imidazole-4-carboxylic acid methyl ester; the compounds were separated during preparative HPLC purification. Mass spec. 10 data (M+H) = 481.

Example 3: Synthesis of 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-5-trifluoromethyl-1*H*-pyrazole-4-carboxylic acid (Compound 1-4)



Step 1: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-5-trifluoromethyl-1*H*-pyrazole-4-carboxylic acid ethyl ester


15 [00518] Prepared according to the procedure described in Example 1, Step 6, using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, ethyl 3-(trifluoromethyl)pyrazole-4-carboxylate, and cesium carbonate.



Step 2: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-5-trifluoromethyl-1H-pyrazole-4-carboxylic acid

[00519] Prepared according to the procedure described in Example 1, Step 7, using the following starting material: 1-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-5-trifluoromethyl-1H-pyrazole-4-carboxylic acid ethyl ester. Mass spec. data ($M+H$) = 549.

Example 4: Synthesis of 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-benzoic acid (Compound 1-5)




10

Step 1: 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-benzoic acid


[00520] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.160g, 0.40mmol), 3-carboxyphenylboronic acid (0.079g, 0.48mmol), 15 tetrakis(triphenylphosphine)palladium(0) (0.046g, 0.04mmol), and potassium carbonate (0.110g, 0.80mmol) were combined in 3:1 DME:H₂O, and the mixture was purged with N₂. The reaction was heated for 4 days, and then purified by silica gel chromatography, followed by preparative HPLC to give the title compound. Mass spec. data ($M+H$) = 491.

Example 5: Synthesis of 3-{4-[4-(2-Bromo-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl]-benzyl}-benzoic acid (Compound 1-6)



Step 1: [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 2-bromo-benzyl ester


[00521] Prepared according to the procedure described in Example 1, Step 5, using the following starting materials: 5-(4-chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 2-bromobenzyl alcohol.



Step 2: 3-{4-[4-(2-Bromo-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl]-benzyl}-benzoic acid

[00522] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 2-bromo-benzyl ester (0.220g, 0.50mmol), 3-carboxyphenylboronic acid (0.100g, 0.60mmol), tetrakis(triphenylphosphine)palladium(0) (0.058g, 0.05mmol), and potassium carbonate (0.138g, 1.00mmol) were combined in 3:1 DME:H₂O, and the mixture was purged with N₂ and stirred overnight at 68°C. After aqueous work-up, the crude material was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 522.


Example 6: Synthesis of [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-4-methoxy-phenyl]-acetic acid (Compound 1-7)



Step 1: (3-Bromo-4-methoxy-phenyl)-acetic acid ethyl ester

[00523] To 3-bromo-4-methoxyphenylacetic acid (24g, 97.9mmol) in EtOH (240mL) was added thionyl chloride (7.8mL, 107.7mmol), and the reaction was stirred at room temperature for 1 hour. Once no starting material was seen by analytical LCMS, the mixture was basified with


saturated aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined organic layers were dried over MgSO₄, filtered, and concentrated to give the desired product.



Step 2: [4-Methoxy-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester


5

[00524] (3-Bromo-4-methoxy-phenyl)-acetic acid ethyl ester (27.4g, 100.3mmol), bis(pinacolato)diboron (25.47g, 100.3mmol), and potassium acetate (24.6g, 250.8mmol) were combined in 1,4-dioxane (250mL) under N₂. The solution was purged with N₂, and then (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II) (4.10g, 5.02mmol) was added and the 10 reaction was heated to 110°C overnight. The mixture was filtered through Celite and partitioned between EtOAc and brine. The aqueous layer was separated and extracted twice with EtOAc, and the combined organic layers were dried and concentrated. The residue was purified by silica gel chromatography (20-60% EtOAc in hexanes) to give the title compound.



15 **Step 3: [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-4-methoxy-phenyl]-acetic acid ethyl ester**


[00525] Prepared according to the procedure described in Example 5, Step 2, using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and [4-methoxy-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-4-methoxy-phenyl]-acetic acid


[00526] Prepared according to the procedure described in Example 1, Step 7, using the following starting material: [3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-4-methoxy-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 535.

Example 7: Synthesis of 4'-[3-Methyl-4-(3-phenyl-propionylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-8)




Step 1: 2-(4-Bromo-benzoyl)-3-oxo-butyric acid methyl ester

[00527] Prepared according to the procedure described in Example 1, Step 2, using the following starting materials: 3-methylamino-but-2-enoic acid methyl ester and 4-bromobenzoyl chloride.




Step 2: 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00528] Prepared according to the procedure described in Example 1, Step 3, using the following starting materials: 2-(4-bromo-benzoyl)-3-oxo-butyric acid methyl ester and hydroxylamine hydrochloride.




Step 3: 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid

[00529] Prepared according to the procedure described in Example 1, Step 4, using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




Step 4: 5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-ylamine

[00530] 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (1.1g, 4.1mmol), diphenylphosphoryl azide (1.2mL, 4.3mmol), and triethylamine (1.14mL, 8.2mmol) were combined in toluene and stirred at 120°C for 25 minutes. After cooling to room temperature, H₂O (2mL) in THF (2mL) was added, and the reaction was stirred at 88°C for 2 hours. The mixture was cooled to room temperature and extracted with EtOAc. The combined organic layers were concentrated to give the title compound.



10 **Step 5: N-[5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-3-phenyl-propionamide**


[00531] Hydrocinnamoyl chloride (0.73mL, 4.9mmol) was added to a mixture of 5-(4-bromo-phenyl)-3-methyl-isoxazol-4-ylamine (4.1mmol), triethylamine (1.14mL, 8.2mmol), and a few drops of DMF in CH₂Cl₂, and the reaction was stirred at room temperature for 2 hours. The mixture was diluted with EtOAc and H₂O, and the organic layer was separated, washed with brine, and concentrated. The residue was purified by dry-load silica gel chromatography (28-48% EtOAc in hexanes) to give the title compound.



Step 6: 4'-(3-Methyl-4-(3-phenyl-propionylamino)-isoxazol-5-yl)-biphenyl-3-carboxylic acid

[00532] Prepared according to the procedure described in Example 5, Step 2, using the following 20 starting materials: N-[5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-3-phenyl-propionamide and 3-carboxyphenylboronic acid.

Example 8: Synthesis of 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1*H*-pyrazole-4-carboxylic acid (Compound 1-9)




Step 1: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-

5 **benzyl)-1*H*-pyrazole-4-carboxylic acid ethyl ester**

[00533] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.156g, 0.39mmol), ethyl 4-pyrazolecarboxylate (0.054g, 0.39mmol), and cesium carbonate (0.254g, 0.78mmol) were combined in DMF, and the reaction was stirred at room temperature for 4 hours. After aqueous worked-up, the crude material was used directly in the

10 hydrolysis step.




Step 2: 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1*H*-pyrazole-4-carboxylic acid

[00534] 1-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-1*H*-pyrazole-4-carboxylic acid ethyl ester (0.090g, 0.19mmol) was treated with lithium hydroxide (0.040g, 0.94mmol) in 3:1 THF:H₂O (3mL) at 60°C for 4 hours. The mixture was acidified to pH 3 and purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 481.

Example 9: Synthesis of 4'-[4-(3-Benzyl-ureido)-3-methyl-isoxazol-5-yl]-biphenyl-3-

20 **carboxylic acid (Compound 1-10)**




Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid *tert*-butyl ester

[00535] 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (1.6g, 6.0mmol),

diphenylphosphoryl azide (1.6g, 6.0mmol), and triethylamine (1.3mL, 9.3mmol) were combined

5 in t-BuOH and refluxed overnight. After cooling, the mixture was concentrated, and the residue was partitioned between EtOAc and H₂O. The organic layer was separated and concentrated, and the residue was purified by silica gel chromatography (16-18% EtOAc in hexanes) to give the title compound.




10 **Step 2: 5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-ylamine**

[00536] [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid *tert*-butyl ester (0.38g,

1.07mmol) was treated with 50% trifluoroacetic acid in CH₂Cl₂ and stirred for 3 hours. Once no

starting material was seen by analytical LCMS, the mixture was concentrated to give the title compound.




15

Step 3: 1-Benzyl-3-[5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-urea

[00537] Sodium hydride (60% in mineral oil; 0.040g, 1.00mmol) was added to a solution of 5-(4-bromo-phenyl)-3-methyl-isoxazol-4-ylamine (0.170g, 0.67mmol) in DMF, and the mixture was stirred for 5 minutes at room temperature. Benzyl isocyanate (0.08mL, 0.67mmol) was added,


20 and the reaction was stirred at 80°C for 2 hours. After aqueous work-up, the crude material was used directly in the coupling step.



Step 4: 4'-(4-(3-Benzyl-ureido)-3-methyl-isoxazol-5-yl)-biphenyl-3-carboxylic acid


[00538] Prepared according to the procedure described in Example 5, Step 2, using the following starting materials: 1-benzyl-3-[5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-urea, 3-carboxyphenylboronic acid.

Example 10: Synthesis of 4'-(3-Methyl-4-((R)-1-pyridin-2-yl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-3-carboxylic acid (Compound 1-11)




Step 1: 2-(4-Bromo-benzoyl)-3-oxo-butyric acid methyl ester

[00539] Prepared according to the procedure described in Example 1, Step 2, using the following starting materials: 3-methylamino-but-2-enoic acid methyl ester and 4-bromobenzoyl chloride.




Step 2: 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00540] Prepared according to the procedure described in Example 1, Step 3, using the following starting materials: 2-(4-bromo-benzoyl)-3-oxo-butyric acid methyl ester and hydroxylamine hydrochloride.



Step 3: 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid

[00541] Prepared according to the procedure described in Example 1, Step 4, using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




5

Step 4: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-pyridin-2-yl-ethyl ester

[00542] Prepared according to the procedure described in Example 1, Step 5, using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-2-(1-


10 hydroxyethyl)pyridine.

**Step 5: 4'-[3-Methyl-4-((R)-1-pyridin-2-yl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid**

[00543] Prepared according to the procedure described in Example 5, Step 2, using the following


15 starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-pyridin-2-yl-ethyl ester and 3-carboxyphenylboronic acid. Mass spec. data (M+H) = 444.

Example 11: Synthesis of 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-benzoic acid (Compound 1-12)



Step 1: 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-benzoic acid methyl ester


[00544] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.160g, 0.40mmol), methyl 3-hydroxybenzoate (0.073g, 0.48mmol), and potassium carbonate (0.110g, 0.8mmol) were combined in acetone and stirred overnight at 60°C. The mixture was dissolved in DMSO and purified by preparative HPLC to give the title compound.



Step 2: 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-benzoic acid


[00545] Prepared according to the procedure described in Example 1, Step 7, using the following starting material: 3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-benzoic acid methyl ester. Mass spec. data (M+H) = 507.

Example 12: Synthesis of 4'-[3-Methyl-4-(6-methyl-pyridin-2-ylmethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-13)



Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 6-methyl-pyridin-2-ylmethyl ester


[00546] Prepared according to the procedure described in Example 1, Step 5, using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 6-methyl-2-pyridinemethanol.



Step 2: 4'-[3-Methyl-4-(6-methyl-pyridin-2-ylmethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid

[00547] Prepared according to the procedure described in Example 5, Step 2, using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 6-methyl-pyridin-2-ylmethyl ester and 3-carboxyphenylboronic acid. Mass spec. data ($M+H$) = 444.


Example 13: Synthesis of 4-Chloro-3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-benzoic acid (Compound 1-14)



15 Step 1: 4-Chloro-3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-benzoic acid

[00548] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 2-chloro-5-carboxyphenylboronic acid. Mass spec. data ($M+H$) = 525.


Example 14: Synthesis of 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-thiophene-2-carboxylic acid (Compound 1-15)



Step 1: 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-thiophene-2-carboxylic acid

[00549] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 2-carboxythiophene-5-boronic acid. Mass spec. data ($M+H$) = 497.


Example 15: Synthesis of 4'-[4-(Indan-1-yloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-16)



10

Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid indan-1-yl ester


[00550] Prepared according to the procedure described in Example 1, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-indanol.



15 Step 2: 4'-[4-(Indan-1-yloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-3-carboxylic acid

[00551] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid indan-1-yl ester and 3-carboxyphenylboronic acid.


Example 16: Synthesis of [4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-17)



Step 1: [4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid

[00552] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.100g, 0.25mmol), 4-mercaptophenylacetic acid (0.050g, 0.30mmol), and potassium carbonate (0.086g, 0.62mmol) were combined in DMF, and the reaction was stirred overnight at 46°C. The crude material was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 537.


Example 17: Synthesis of [3-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-18)



Step 1: [3-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid

[00553] Prepared according to the procedure described in Example 16, Step 1 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 3-mercaptophenylacetic acid. Mass spec. data (M+H) = 537.


Example 18: Synthesis of [2-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-19)



Step 1: [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid


[00554] Prepared according to the procedure described in Example 16, Step 1 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 2-mercaptophenylacetic acid. Mass spec. data ($M+H$) = 537.

Example 19: Synthesis of [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-20)



10 **Step 1: [4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester**


[00555] Ethyl 4-bromophenylacetate (12g, 49mmol), bis(pinacolato)diboron (12.4g, 59mmol), 1,1'-bis(diphenylphosphino)ferrocene-dichloropalladium(II) (2g, 2.4mmol), and potassium acetate (9.6g, 98mmol) were combined in 1,4-dioxane (100mL), and the reaction was heated to 80°C for 6 hours. The mixture was worked-up to give the title compound.



15


Step 2: 4-(Ethoxycarbonyl)methylphenylboronic acid

[00556] [4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl (0.460g, 1.58mmol), sodium periodate (0.746g, 3.49mmol), and ammonium acetate (0.309g, 4.01mmol) were combined in 1:1 acetone:H₂O and stirred overnight at room temperature. Aqueous work-up provided the title compound.



Step 3: [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid ethyl ester


[00557] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 4-((ethoxycarbonyl)methyl)phenylboronic acid.



Step 4: [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid


[00558] Prepared according to the procedure described in Example 1, Step 7 using the following starting materials: [4-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid ethyl ester. Mass spec. data ($M+H$) = 505.

Example 20: Synthesis of [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-21)




Step 1: [3-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid methyl ester

[00559] Prepared according to the procedure described in Example 19, Step 1 using the following starting materials: methyl 2-(3-bromophenyl)acetate and bis(pinacolato)diboron.




Step 2: 3-((Methoxycarbonyl)methyl)phenylboronic acid

[00560] Prepared according to the procedure described in Example 19, Step 2 using the following starting material: [3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid methyl ester.



Step 3: [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid methyl ester


[00561] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 3-((methoxycarbonyl)methyl)phenylboronic acid.



Step 4: [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid


[00562] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid methyl ester. Mass spec. data (M+H) = 505.

Example 21: Synthesis of [2-(4-{1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-22)




Step 1: (2-Bromo-phenyl)-acetic acid ethyl ester

5 [00563] 2-Bromophenylacetic acid (1.6g, 7.44mmol) was treated with sulfuric acid (1mL) in EtOH (10mL) at 80°C for 1 hour. The mixture was concentrated and purified by silica gel chromatography to give the title compound.




Step 2: [2-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester

10 [00564] Prepared according to the procedure described in Example 19, Step 1 using the following starting materials: (2-bromo-phenyl)-acetic acid ethyl ester and bis(pinacolato)diboron.




Step 3: 2-((Ethoxycarbonyl)methyl)phenylboronic acid

15 [00565] Prepared according to the procedure described in Example 19, Step 2 using the following starting material: [2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: [2-(4-{1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid ethyl ester


[00566] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 2-((ethoxycarbonyl)methyl)phenylboronic acid.



5 **Step 5: [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid**


[00567] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [2-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 505.

10 **Example 22: Synthesis of (4'-{3-Methyl-4-[1-(2-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-23)**



Step 1: 5-(4'-Ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid


[00568] 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (2.0g, 7.07mmol), [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester (2.46g, 8.5mmol), tetrakis(triphenylphosphine)palladium(0) (0.80g, 0.70mmol), and sodium bicarbonate (4.3g, 52mmol) were combined in 1,4-dioxane (50mL) and H₂O (10mL), and the reaction was stirred overnight at 80°C to give the title compound.



Step 2: (4'-(3-Methyl-4-[1-(2-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester

[00569] 5-(4'-Ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid


5 (0.150g, 0.41mmol), alpha-methyl-2-trifluoromethylbenzyl alcohol (0.094g, 0.49mmol), diphenylphosphoryl azide (0.11mL, 0.49mmol), and triethylamine (0.11mL, 0.49mmol) were combined in toluene (3mL) and stirred at 80°C for 2 hours. After cooling, the mixture was worked-up to give the title compound.



10 **Step 3: (4'-(3-Methyl-4-[1-(2-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid**


[00570] (4'-(3-Methyl-4-[1-(2-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester (0.41mmol) was treated with aqueous lithium hydroxide in MeOH at 60°C for 30 minutes. The mixture was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 525.

Example 23: Synthesis of (4'-(3-Methyl-4-[1-(3-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-24)



Step 1: (4'-(3-Methyl-4-[1-(3-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00571] Prepared according to the procedure described in Example 22, Step 2 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and alpha-methyl-3-trifluoromethylbenzyl alcohol.



Step 2: (4'-(3-Methyl-4-[1-(3-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00572] Prepared according to the procedure described in Example 22, Step 3 using the following starting material: (4'-(3-methyl-4-[1-(3-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 525.

Example 24: Synthesis of (4'-(4-[1-(2,4-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-25)



Step 1: (4'-(4-[1-(2,4-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00573] Prepared according to the procedure described in Example 22, Step 2 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 2,4-dichloro-alpha-methylbenzyl alcohol



Step 2: (4'-(4-[1-(2,4-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00574] Prepared according to the procedure described in Example 22, Step 3 using the following starting material: (4'-(4-[1-(2,4-dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 525.

Example 25: Synthesis of (4'-(4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-26)



Step 1: (4'-(4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00575] Prepared according to the procedure described in Example 22, Step 2 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 2-fluoro-alpha-methylbenzyl alcohol.



Step 2: (4'-(4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00576] Prepared according to the procedure described in Example 22, Step 3 using the following starting material: (4'-(4-[1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 475.

Example 26: Synthesis of (4'-(4-[1-(3-Bromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-27)



Step 1: (4'-(4-[(3-Bromo-phenyl)-ethoxycarbonylaminobiphenyl-4-yl)-3-methylisoxazol-5-yl]-acetic acid ethyl ester


[00577] Prepared according to the procedure described in Example 22, Step 2 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 3-bromo-alpha-methylbenzyl alcohol.



Step 2: (4'-(4-[(3-Bromo-phenyl)-ethoxycarbonylaminobiphenyl-4-yl)-3-methylisoxazol-5-yl]-acetic acid


[00578] Prepared according to the procedure described in Example 22, Step 3 using the following starting material: (4'-(4-[(3-bromo-phenyl)-ethoxycarbonylaminobiphenyl-4-yl)-3-methylisoxazol-5-yl]-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 535.

Example 27: Synthesis of (4'-(4-[(2-Methoxy-phenyl)-ethoxycarbonylaminobiphenyl-4-yl)-3-methylisoxazol-5-yl]-biphenyl-4-yl)-acetic acid (Compound 1-28)



Step 1: (4'-(4-[(1-ethoxy-1-methylethoxy)carbonyl]amino)-3-methylisoxazol-5-yl)biphenyl-4-yl acetic acid ethyl ester


[00579] Prepared according to the procedure described in Example 22, Step 2 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2-methoxyphenyl)ethanol.



Step 2: (4'-(4-[(1-hydroxy-1-methylethoxy)carbonyl]amino)-3-methylisoxazol-5-yl)biphenyl-4-yl acetic acid


[00580] Prepared according to the procedure described in Example 22, Step 3 using the following starting material: (4'-(4-[(1-hydroxy-1-methylethoxy)carbonyl]amino)-3-methylisoxazol-5-yl)biphenyl-4-yl acetic acid. Mass spec. data ($M+H$) = 487.

Example 28: Synthesis of {4'-(3-Methyl-4-(1-methyl-2-phenyl-ethoxycarbonyl)amino)-isoxazol-5-yl}-biphenyl-4-yl acetic acid (Compound 1-29)



Step 1: {4'-(3-Methyl-4-(1-methyl-2-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid ethyl ester

[00581] Prepared according to the procedure described in Example 22, Step 2 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 1-phenyl-2-propanol.



Step 2: {4'-(3-Methyl-4-(1-methyl-2-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid

[00582] Prepared according to the procedure described in Example 22, Step 3 using the following starting material: {4'-(3-methyl-4-(1-methyl-2-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid ethyl ester. Mass spec. data (M+H) = 471.


Example 29: Synthesis of (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-methoxy-biphenyl-3-yl)-acetic acid (Compound 1-30)



15

Step 1: 3-Bromo-benzoyl chloride


[00583] To 3-bromobenzoic acid (9.42g, 46.9mmol) in CH_2Cl_2 (500mL) was added DMF (0.4mL), followed by oxalyl chloride (5.3mL, 60.9mmol) slowly over 10 minutes by syringe. The reaction was stirred for 1 hour at room temperature, and then concentrated to give the title compound.



5

Step 2: 2-(3-Bromo-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester

[00584] To 3-bromo-benzoyl chloride (46.9mmol) in THF (400mL) was added 3-methylamino-but-2-enoic acid methyl ester (6.0g, 46.9mmol), followed by pyridine (4.2mL, 51.6mmol), and the reaction was stirred overnight at room temperature. Solid pyridine hydrochloride coated the flask, so the mixture was decanted, and the solid was washed three times with EtOAc. The combined solutions were concentrated and diluted with EtOAc (500mL), and the mixture was washed three times with H_2O . The aqueous layer was separated and back-extracted with EtOAc, and the combined organic layers were dried over MgSO_4 , filtered, and concentrated to give the title compound.



15


Step 3: 5-(3-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00585] 2-(3-Bromo-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester (15g, 47mmol) and hydroxylamine hydrochloride (3.26g, 47mmol) were combined in acetic acid (300mL) and stirred at 60°C overnight. Once no starting material was seen by analytical LCMS, the mixture was concentrated, and the residue was partitioned between EtOAc and H_2O . Saturated aqueous NaHCO_3 was added to neutralize residual acetic acid, and the organic layer was separated and washed twice with H_2O . The combined aqueous layers were back-extracted with EtOAc, and the combined organic layers were dried, filtered, and concentrated. The crude material was filtered through a plug of silica gel, eluting with 30% EtOAc in hexanes (400mL), and the filtrate was concentrated to give the title compound.




Step 4: 5-(3-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid

[00586] 5-(3-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester (3.64g, 12.3mmol) was hydrolyzed with 1N aqueous LiOH (30mL, 30mmol) for 6 hours to give the title compound.



Step 5: [5-(3-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester


[00587] 5-(3-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (1.128g, 4.0mmol) was dissolved in toluene (25mL). Triethylamine (0.61mL, 4.4mmol) was added, followed by diphenylphosphoryl azide (0.86mL, 4.0mmol), and the mixture was stirred for 5 minutes. 2-Chloro-alpha-methylbenzyl alcohol (0.691g, 4.4mmol) was added, and the reaction was stirred at 80°C for 2 hours. The mixture was cooled to room temperature and diluted with EtOAc (150mL). The organic layer was washed twice with H₂O and once with brine, and the combined aqueous layers were back-extracted with EtOAc. The combined organic layers were dried, filtered, and concentrated, and the residue was purified by silica gel chromatography to give the title compound.



Step 6: (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-methoxy-biphenyl-3-yl)-acetic acid ethyl ester

[00588] [5-(3-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.085g, 0.20mmol) and [4-methoxy-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester (0.077g, 0.24mmol) were combined in 2:1 DME:H₂O (3mL), and


the solution was purged with N₂ for 5 minutes. Potassium carbonate (0.083g, 0.6mmol) was added, followed by tetrakis(triphenylphosphine)palladium(0) (0.014g, 0.02mmol), and the reaction was stirred at 70°C for 30 minutes, until no starting material was seen by analytical LCMS and tlc. After aqueous work-up, the crude material was purified by silica gel chromatography to give the title compound.



Step 7: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-methoxy-biphenyl-3-yl)-acetic acid


[00589] (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-methoxy-biphenyl-3-yl)-acetic acid ethyl ester was hydrolyzed with lithium hydroxide in MeOH and H₂O to give the title compound. Mass spec. data (M+H) = 521.

Example 30: Synthesis of 3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-carboxylic acid (Compound 1-31)




15 **Step 1: {3-Methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester**

[00590] Prepared according to the procedure described in Example 6, Step 2 using the following starting materials: [5-(3-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and bis(pinacolato)diboron.



Step 2: 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid methyl ester


[00591] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and methyl 3-bromobenzoate.



Step 3: 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid


[00592] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 3'-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid methyl ester. Mass spec. data ($M+H$) = 477.

Example 31: Synthesis of 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid (Compound 1-32)



Step 1: 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid methyl ester


[00593] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and methyl 2-bromobenzoate.



Step 2: 3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-carboxylic acid


[00594] Prepared according to the procedure described in Example 29, Step 7 using the following 5 starting material: 3'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-carboxylic acid methyl ester. Mass spec. data ($M+H$) = 477.

Example 33: Synthesis of 5-(3-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-pyridine-2-carboxylic acid (Compound 1-34)



10 **Step 1: 5-(3-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-pyridine-2-carboxylic acid methyl ester**


[00595] {3-Methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.100g, 0.21mmol), methyl 5-bromopicolinate (0.054g, 0.25mmol), and sodium bicarbonate (0.053g, 0.63mmol) were combined in 1,4-dioxane (3mL) and H_2O (1.5mL). The solution was purged with N_2 for 10 minutes, and then (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II) (0.009g, 0.01mmol) was added, and the reaction was stirred at 50°C for 1.5 hours, until no starting material was seen by analytical LMCS. The mixture was worked-up and purified by silica gel chromatography to give the title compound.



Step 2: 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid


[00596] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 5-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester. Mass spec. data (M+H) = 478.

Example 34: Synthesis of 2-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid (Compound 1-35)



10 **Step 1: 2-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid methyl ester**

[00597] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and methyl 2-bromoisonicotinate.




15

Step 2: 2-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid


[00598] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 2-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid methyl ester. Mass spec. data (M+H) = 478.

5 **Example 35: Synthesis of 6-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-36)**



Step 1: 6-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid ethyl ester


10 [00599] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and ethyl 6-bromopicolinate.



Step 2: 6-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid


15 [00600] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 6-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid ethyl ester. Mass spec. data (M+H) = 478.

Example 36: Synthesis of 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-37)



Step 1: 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid methyl ester


[00601] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and methyl 5-bromopyridine-3-carboxylate.



Step 2: 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid


[00602] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 5-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid methyl ester. Mass spec. data ($M+H$) = 478.

Example 37: Synthesis of [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid (Compound 1-38)



Step 1: 3-Bromo-2-methoxy-5-methyl-pyridine


[00603] 3-Bromo-2-hydroxy-5-methylpyridine (5.012g, 26.7mmol), silver carbonate (9.58g, 34.7mmol), and iodomethane (5mL, 80.0mmol) were combined in $CHCl_3$ (75mL) and stirred at 20 $55^\circ C$ for 3 hours. Once no starting material was seen by analytical LCMS, the mixture was cooled and filtered through Celite, and the filtrate was concentrated and purified by silica gel chromatography (0-100% EtOAc in hexanes) to give the title compound.



Step 2: 3-Bromo-5-bromomethyl-2-methoxy-pyridine


[00604] 3-Bromo-2-methoxy-5-methyl-pyridine (4.0g, 19.8mmol), N-bromosuccinimide (4.45g, 23.8mmol), and azobisisobutyronitrile (0.163g, 0.99mmol) were combined in benzene (40mL) and stirred overnight at 80°C. Analytical LCMS indicated the starting material was still present, so the reaction was stirred for another 24 hours at 80°C, and then additional N-bromosuccinimide (1.8g, 10.1mmol), and azobisisobutyronitrile (catalytic) were added. After stirring for another 2 hours at 80°C, the mixture was diluted with H₂O and extracted with CH₂Cl₂. The residue was purified by silica gel chromatography to give the title compound.

10




Step 3: 3-Bromo-5-dibromomethyl-2-methoxy-pyridine

[00605] 3-Bromo-5-bromomethyl-2-methoxy-pyridine (6.03g, 21.5mmol), N-bromosuccinimide (4.09g, 23.0mmol), and azobisisobutyronitrile (one scoop) were combined in benzene (40mL) and stirred at 80°C for 4 hours. After cooling, the mixture was diluted with H₂O and extracted with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and concentrated to give the title compound.




Step 4: 5-Bromo-6-methoxy-pyridine-3-carbaldehyde and 5-Bromo-6-methoxy-nicotinic acid ethyl ester

[00606] To 3-bromo-5-dibromomethyl-2-methoxy-pyridine (7.1g, 19.7mmol) in EtOH (200mL) was added silver nitrate (8.38g, 49.3mmol) in H₂O (40mL) slowly, and the reaction was stirred at 75°C overnight. Analytical LCMS indicated that the diethyl acetal product may have formed, so the mixture was cooled, diluted with brine, filtered, and concentrated. 1,4-Dioxane (70mL) and 1N aqueous HCl (70mL) was added, and the reaction was stirred overnight. Analytical LCMS showed that two products were present. The mixture was neutralized and extracted with EtOAc, and the combined organic layers were dried, filtered, and concentrated. The residue was purified by silica gel chromatography to give the title compounds, which were combined and reduced together in the next step.




Step 5: (5-Bromo-6-methoxy-pyridin-3-yl)-methanol


[00607] A mixture of 5-bromo-6-methoxy-pyridine-3-carbaldehyde and 5-bromo-6-methoxy-nicotinic acid ethyl ester (3g, 13.9mmol) was dissolved in THF (30mL) and cooled to 0°C, and lithium triethylborohydride (1M in THF; 42mL, 42mmol) was added. The reaction was stirred 5 for 5 minutes, and then quenched with EtOH, acidified with 1N aqueous HCl, and extracted with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and concentrated to give the title compound.

**Step 6: 3-Bromo-5-bromomethyl-2-methoxy-pyridine**


10 [00608] To (5-bromo-6-methoxy-pyridin-3-yl)-methanol (2.86g, 13.1mmol) in DME (30mL) was added phosphorus tribromide (1.9mL, 19.7mmol), and the reaction was stirred for 15 minutes. The mixture was worked up with EtOAc and saturated aqueous NaHCO₃, and the combined organic layers were dried over MgSO₄, filtered, and concentrated to give the title compound.

**15 Step 7: (5-Bromo-6-methoxy-pyridin-3-yl)-acetonitrile**

[00609] To 3-bromo-5-bromomethyl-2-methoxy-pyridine (1.42g, 5.05mmol) in EtOH (14mL) and H₂O (1.5mL) was added sodium cyanide (0.25g, 5.05mmol), and the reaction was stirred at 50°C overnight. The mixture was cooled to room temperature and worked up with CH₂Cl₂ and saturated aqueous NaHCO₃. The combined organic layers were dried, filtered, and concentrated 20 to give the title compound.


**Step 8: (5-Bromo-6-methoxy-pyridin-3-yl)-acetic acid ethyl ester**

[00610] To (5-bromo-6-methoxy-pyridin-3-yl)-acetonitrile (1.139g, 5.02mmol) in EtOH (11ml) was added acetyl chloride (3.6mL, 50.54mmol), and the reaction was stirred at 50°C for 24 hours, 25 and at room temperature for 2 days. The reaction was worked up with CH₂Cl₂ and saturated aqueous NaHCO₃, and then reacidified and extracted with CH₂Cl₂. The combined organic layers were dried, filtered, and concentrated to give the title compound.



Step 9: [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid ethyl ester


[00611] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and (5-bromo-6-methoxy-pyridin-3-yl)-acetic acid ethyl ester.



Step 10: [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid


[00612] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: [5-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid ethyl ester. Mass spec. data (M+H) = 522.

Example 38: Synthesis of (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-39)



Step 1: (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester

[00613] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and ethyl 4-bromophenylacetate.




Step 2: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid

[00614] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (3'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester; further purification by preparative HPLC was required. Mass spec. data (M+H) = 491.


Example 39: Synthesis of [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-indol-1-yl]-acetic acid (Compound 1-40)

10



Step 1: (5-Bromo-indol-1-yl)-acetic acid ethyl ester


[00615] To 5-bromoindole (0.150g, 0.77mmol) in DMF (10mL) was added sodium hydride (60% in mineral oil; 0.028g, 1.15mmol), and the mixture was stirred for 1 hour. Ethyl bromoacetate (0.13mL, 1.15mmol) was added, and the reaction was stirred for 1 hour at room temperature. The mixture was quenched with H₂O and 1N aqueous HCl (5mL), and extracted with EtOAc. The combined organic layers were washed twice with H₂O and once with brine, and then dried and concentrated to give the title compound.



Step 2: [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-indol-1-yl]-acetic acid ethyl ester

[00616] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-


isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and (5-bromo-indol-1-yl)-acetic acid ethyl ester; purification by preparative HPLC was utilized.



Step 3: [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-indol-1-yl]-acetic acid


5 [00617] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: [5-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-indol-1-yl]-acetic acid ethyl ester. Mass spec. data (M+H) = 530.

10 **Example 40: Synthesis of [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-41)**



Step 1: (5-Bromo-pyridin-3-yl)-acetic acid ethyl ester


[00618] 5-Bromo-3-pyridylacetic acid was treated with 5% concentrated sulfuric acid in EtOH at reflux overnight to give the title compound.



15

Step 2: [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid ethyl ester


[00619] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and (5-bromo-pyridin-3-yl)-acetic acid ethyl ester.



Step 3: [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid


[00620] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: [5-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid ethyl ester. Mass spec. data (M+H) = 492.

Example 41: Synthesis of (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-42)




10 **Step 1: (3-Bromo-4-fluoro-phenyl)-acetic acid ethyl ester**

[00621] 3-Bromo-4-fluorophenylacetic acid (2.35g, 9.0mmol) was treated with concentrated sulfuric acid (2mL) in EtOH (100mL) at reflux for 2 hours. Purification by silica gel chromatography provided the title compound.



15 **Step 2: (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid ethyl ester**


[00622] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and (3-bromo-4-fluoro-phenyl)-acetic acid ethyl ester.



Step 3: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-fluoro-biphenyl-3-yl)-acetic acid


[00623] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (3'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-fluoro-biphenyl-3-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 509.

Example 42: Synthesis of (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-43)




10 **Step 1: (5-Bromo-2-fluoro-phenyl)-acetic acid ethyl ester**

[00624] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 5-bromo-2-fluorophenylacetic acid.



15 **Step 2: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid ethyl ester**


[00625] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and (5-bromo-2-fluoro-phenyl)-acetic acid ethyl ester.



Step 3: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid

[00626] Prepared according to the procedure described in Example 29, Step 7 using the following 5 starting material: (3'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 509.


Example 43: Synthesis of (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-yl)-acetic acid (Compound 1-44)



10 **Step 1: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-yl)-acetic acid ethyl ester**

[00627] {3-Methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.241g, 0.5mmol) and ethyl 2-bromophenylacetate (0.146g, 0.6mmol) were combined in 1,4-dioxane (6mL) and H_2O (3mL).


15 The solution was purged with N_2 for 10 minutes, and then sodium bicarbonate (0.126g, 1.5mmol) was added, followed by bis(triphenylphosphine)palladium(II) dichloride (0.018g, 0.03mmol), and the reaction was stirred at 80°C for 1 hour. After work-up, the crude material was purified by silica gel chromatography to give the title compound.



Step 2: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-yl)-acetic acid

[00628] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (3'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-yl)-acetic acid ethyl ester; further purification by preparative HPLC was required. Mass spec. data ($M+H$) = 491.


Example 44: Synthesis of 4-(3-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-cyclohex-3-enecarboxylic acid (Compound 1-45)



10

Step 1: 4-Trifluoromethanesulfonyloxy-cyclohex-3-enecarboxylic acid ethyl ester

[00629] Ethyl-4-oxocyclohexane carboxylate (0.5mL, 3.14mmol) was dissolved in dry THF (20mL) and cooled to -78°C. Lithium bis(trimethylsilyl)amide (1M in hexane; 4mL, 4.0mmol) was added over 3 minutes, and the solution was stirred at -78°C for 2 hours. *N,N*-Bis(trifluoromethylsulfonyl)-5-chloro-2-pyridylamine (1.36g, 3.45mmol) was added, and the reaction was stirred at room temperature for 3 hours. After aqueous work-up, the crude material was purified by silica gel chromatography (0-30% EtOAc in hexanes) to give the title compound.




20

Step 2: 4-(3-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-cyclohex-3-enecarboxylic acid ethyl ester

[00630] {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.400g, 0.83mmol) and 4-


trifluoromethanesulfonyloxy-cyclohex-3-enecarboxylic acid ethyl ester (0.277g, 0.91mmol) were combined in DME (6mL) and H₂O (3mL). The solution was purged with N₂ for 5 minutes, and then sodium carbonate (0.265g, 2.49mmol) was added, followed by bis(triphenylphosphine)palladium(II) dichloride (0.030g, 0.04mmol), and the reaction was stirred 5 at 75°C for 1 hour. After work-up, the crude material was purified by silica gel chromatography to give the title compound.



Step 3: 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid

10 [00631] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 4-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid ethyl ester. Mass spec. data (M+H) = 481.


Example 45: Synthesis of 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid (Compound 1-46)



15

Step 1: 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid ethyl ester


[00632] 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid ethyl ester (0.225g, 0.44mmol) was dissolved in 10% MeOH in 20 EtOAc (20mL), and the solution was purged with N₂ for 6 minutes. 10% Palladium on carbon (50% wet; 0.100g, 0.044mmol) was added, and the reaction was stirred under a balloon of H₂ for 2 hours at room temperature. The mixture was worked-up, and the crude material was purified by silica gel chromatography to give the title compound.



Step 2: 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid


[00633] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 4-(3-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid ethyl ester.

Example 46: Synthesis of [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid ethyl ester (Compound 1-47)




10 **Step 1: 6-Bromo-nicotinoyl chloride**

[00634] Prepared according to the procedure described in Example 29, Step 1 using the following starting material: 6-bromopyridine-3-carboxylic acid.




Step 2: 2-(6-Chloro-pyridine-3-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester

15 [00635] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 6-bromo-nicotinoyl chloride and 3-methylamino-but-2-enoic acid methyl ester; the product was purified by silica gel chromatoagrophy. An analytical LCMS indicated that the bromide was displaced by a chloride during the reaction conditions.




Step 3: 5-(6-Chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00636] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(6-chloro-pyridine-3-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.



Step 3: 5-(6-Chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid


[00637] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(6-chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.



10

Step 4: 5-[6-(4-Ethoxycarbonylmethyl-phenyl)-pyridin-3-yl]-3-methyl-isoxazole-4-carboxylic acid

[00638] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: 5-(6-chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.




Step 5: [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid ethyl ester

[00639] Prepared according to the procedure described in Example 29, Step 5 using the following

5 starting materials: 5-[6-(4-ethoxycarbonylmethyl-phenyl)-pyridin-3-yl]-3-methyl-isoxazole-4-carboxylic acid and 2-fluoro-alpha-methylbenzyl alcohol. Mass spec. data (M+H) = 504.


Example 47: Synthesis of [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid (Compound 1-48)



10 **Step 1: [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid**

[00640] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: [4-(5-{4-[1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 476.


15 **Example 48: Synthesis of (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-methoxy-biphenyl-3-yl)-acetic acid (Compound 1-49)**



Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester

[00641] To 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (5.764g, 20.4mmol) in


5 toluene (200mL) was added triethylamine (3.13mL, 22.4mmol) and diphenylphosphoryl azide (4.42mL, 20.4mmol), and the mixture was stirred for 5 minutes at room temperature. 2-Chloro-alpha-methylbenzyl alcohol (3.2g, 22.4mol) was added, and the reaction was stirred at 80°C for 2 hours. The mixture was diluted with EtOAc (200mL) and extracted with EtOAc. The combined organic layers were washed twice with H₂O and once with brine, and the combined aqueous 10 layers were back-extracted with EtOAc. The combined organic layers were concentrated, and the residue was purified by silica gel chromatography to give the title compound.



Step 2: (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-methoxy-biphenyl-3-yl)-acetic acid ethyl ester

[00642] To [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.100g, 0.23mmol) in DME (5mL) was added [4-methoxy-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester (0.074g, 0.23mmol), followed by potassium carbonate (0.079g, 0.58mmol) and H₂O (3mL). The solution was purged with N₂ for 10 minutes, and then tetrakis(triphenylphosphine)palladium(0) (0.027g, 0.02mmol) was added.

20 The reaction was stirred at 100°C until no starting material was seen by analytical LCMS and tlc, and then cooled to room temperature and diluted with EtOAc. The mixture was extracted with EtOAc, and the combined organic layers were washed with H₂O and brine. The combined aqueous layers were back-extracted with EtOAc, and the combined organic layers were dried and concentrated. The residue was purified by silica gel chromatography to give the title compound.




Step 3: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-methoxy-biphenyl-3-yl)-acetic acid

[00643] To (4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-6-

5 methoxy-biphenyl-3-yl)-acetic acid ethyl ester (0.105g, 0.20mmol) in MeOH (20mL) was added 1N aqueous LiOH (5mL, 5mmol), and the reaction was stirred for 2 hours, until no starting material was seen by analytical LCMS. The mixture was diluted with H₂O and extracted with EtOAc. The combined organic layers were dried and concentrated to give the title compound. Mass spec. data (M+H) = 521.


10 **Example 49: Synthesis of 2-(4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-isonicotinic acid (Compound 1-50)**



Step 1: {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester

15 [00644] [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (3.48g, 8.0mmol), bis(pinacolato)diboron (2.24g, 8.8mmol), and potassium acetate (2.88g, 32.0mmol) were combined in 1,4-dioxane (200mL) under N₂. The solution was purged with N₂ for 10 minutes, and then (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II) (0.652g, 0.8mmol) was added and the reaction was heated to 80°C for 6 hours, and then cooled to 20 room temperature and stirred overnight. The mixture was diluted with H₂O and extracted with EtOAc. The combined organic layers were washed twice with H₂O and once with brine, and the combined aqueous layers were back-extracted with EtOAc. The combined organic layers were


concentrated and the residue was purified by silica gel chromatography (0-100% EtOAc in hexanes) to give the title compound.



Step 2: 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid

[00645] {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.100g, 0.21mmol), methyl 2-bromoisonicotinate (0.045g, 0.21mmol), and potassium carbonate (0.073g, 0.53mmol) were combined in DME (10mL) and H₂O (5mL). The solution was purged with N₂ for 10 minutes, and then (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II) (0.017g, 0.02mmol) was added. The reaction was stirred at 70°C for 4 hours, until no starting material was seen by analytical LCMS and the hydrolyzed product was observed. The mixture was cooled to room temperature and neutralized with 1N aqueous HCl (3mL). The mixture was extracted with EtOAc, and the combined organic layers were dried and concentrated. The residue was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 478.

Example 50: Synthesis of 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid (Compound 1-51)




Step 1: 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid

[00646] Prepared according to the procedure described in Example 49, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-

isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 4-bromobenzoic acid. Mass spec. data ($M+H$) = 477.


Example 51: Synthesis of 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid (Compound 1-52)



5

Step 1: 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid methyl ester


[00647] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, methyl 2-bromobenzoate, and bis(triphenylphosphine)palladium(II) dichloride.



Step 2: 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid


[00648] To 4'-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid methyl ester (0.21mmol) in MeOH was added 1N aqueous LiOH (0.5mL), and the reaction was stirred overnight at room temperature. Analytical LCMS indicated that only starting material was present, so the mixture was concentrated, redissolved in MeOH, and treated with 1N aqueous LiOH (2mL) and stirred overnight at room temperature. No reaction was observed, so the solution was stirred at 40°C for 3 days. Analytical LCMS showed that no starting material remained, so the mixture was purified by preparative HPLC to give the title compound. Mass spec. data ($M+H$) = 477.

Example 52: Synthesis of 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-53)



Step 1: 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-2-carboxylic acid ethyl ester


[00649] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, ethyl 6-bromopicolinate, and bis(triphenylphosphine)palladium(II) dichloride; purification by preparative HPLC was utilized.



Step 2: 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-2-carboxylic acid

[00650] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 6-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-2-carboxylic acid ethyl ester. Mass spec. data (M+H) = 478.


Example 53: Synthesis of 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-54)



Step 1: 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid

[00651] Prepared according to the procedure described in Example 49, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 6-bromopyridine-3-carboxylic acid. Mass spec. data ($M+H$) = 478.

Example 54: Synthesis of 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-55)




10

Step 1: 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid


[00652] Prepared according to the procedure described in Example 49, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and methyl 5-bromopyridine-3-carboxylate. Mass spec. data ($M+H$) = 478.

Example 55: Synthesis of 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-56)



Step 1: 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester


[00653] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, methyl 5-bromopicolinate, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).



Step 2: 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid

[00654] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 5-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid methyl ester. Mass spec. data (M+H) = 478.


Example 56: Synthesis of (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-yl)-acetic acid (Compound 1-57)



Step 1: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-2-yl)-acetic acid

[00655] Prepared according to the procedure described in Example 49, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 2-bromophenylacetic acid. Mass spec. data ($M+H$) = 491.


Example 57: Synthesis of (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-58)



10

Step 1: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00656] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, ethyl 4-bromophenylacetate, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).



Step 2: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00657] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: (4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 491.

Example 58: Synthesis of (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid (Compound 1-59)




10 **Step 1: (3-Bromo-phenyl)-acetic acid ethyl ester**

[00658] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 3-bromophenylacetic acid.



Step 2: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid ethyl ester


[00659] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, (3-bromo-phenyl)-acetic acid ethyl ester, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).



Step 3: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid


[00660] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 491.

Example 59: Synthesis of (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid (Compound 1-60)




10 **Step 1: (3-Bromo-phenyl)-acetic acid ethyl ester**

[00661] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 3-bromophenylacetic acid.



15 **Step 2: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid ethyl ester**


[00662] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, (3-bromo-phenyl)-acetic acid ethyl ester, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).



Step 3: (3'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid


[00663] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: (3'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 491.

Example 60: Synthesis of [6-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl]-pyridin-3-yl]-acetic acid (Compound 1-61)



10 **Step 1: (6-Chloro-pyridin-3-yl)-acetic acid methyl ester**


[00664] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 2-chloropyridine-5-acetic acid.



Step 2: [6-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl]-pyridin-3-yl]-acetic acid


[00665] Prepared according to the procedure described in Example 49, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, (6-chloro-pyridin-3-yl)-acetic acid methyl ester, and tetrakis(triphenylphosphine)palladium(0). Mass spec. data ($M+H$) = 492.

Example 61: Synthesis of 3-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-62)




Step 1: 3-(4-Bromo-phenyl)-propionic acid ethyl ester

5 [00666] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 3-(4-bromophenyl)propionic acid.



Step 2: 3-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester


10 [00667] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, 3-(4-bromo-phenyl)-propionic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride.



Step 3: 3-(4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid


[00668] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 3-(4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid ethyl ester. Mass spec. data ($M+H$) = 505.

Example 62: Synthesis of [5-(4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-63)



10 **Step 1: (5-Bromo-pyridin-3-yl)-acetic acid ethyl ester**


[00669] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 5-bromo-3-pyridylacetic acid.



15 **Step 2: [5-(4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-pyridin-3-yl]-acetic acid**

[00670] Prepared according to the procedure described in Example 49, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, (5-bromo-pyridin-3-yl)-acetic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride. Mass spec. data ($M+H$) = 492.

Example 63: Synthesis of (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-64)




Step 1: (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid ethyl ester

5

[00671] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, (3-bromo-4-fluoro-phenyl)-acetic acid ethyl ester, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).

10




Step 2: (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid

15


[00672] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: (4'-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 509.

Example 64: Synthesis of (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-4-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-65)



Step 1: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid ethyl ester


[00673] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, (5-bromo-2-fluoro-phenyl)-acetic acid ethyl ester, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).



Step 2: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid


[00674] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: (4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-4-fluoro-biphenyl-3-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 509.

Example 65: Synthesis of 4-(4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-phenyl)-cyclohex-3-enecarboxylic acid (Compound 1-66)



Step 1: 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid ethyl ester


[00675] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, 4-trifluoromethanesulfonyloxy-cyclohex-3-enecarboxylic acid ethyl ester, and (1,1'-bis(diphenylphosphino)ferrocene)-dichloropalladium(II).



10 **Step 2: 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid**


[00676] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 4-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid ethyl ester. Mass spec. data (M+H) = 481.

15 **Example 66: Synthesis of 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid (Compound 1-67)**



Step 1: 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid ethyl ester


[00677] Prepared according to the procedure described in Example 45, Step 1 using the following starting material: 4-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid ethyl ester.



Step 2: 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid

[00678] 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid ethyl ester (0.089g, 0.17mmol) was dissolved in MeOH (10mL) and THF (2mL), and treated with 1N aqueous LiOH (2mL). The reaction was stirred overnight at room temperature. Analytical LCMS indicated that some starting material was still present, so additional 1N aqueous LiOH (3mL) was added, and the reaction was stirred for 1.5 hours, until no starting material was seen by analytical LCMS. The mixture was quenched with 1N aqueous HCl (6mL) and extracted with EtOAc. The combined organic layers were concentrated, and the residue was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 483.

Example 67: Synthesis of (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid methyl ester (Compound 1-69)




Step 1: (4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid methyl ester

[00679] To (4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-

4-yl)-acetic acid (0.050g, 0.1mmol) in benzene (17mL) and MeOH (2mL) was added (trimethylsilyl)diazomethane (2M in hexanes; 0.1mL, 0.2mmol), and the reaction was stirred at room temperature for 5 minutes. Analytical LCMS showed the starting material was still present, so additional (trimethylsilyl)diazomethane (2M in hexanes; 0.25mL, 0.5mmol) was added, and the reaction was stirred until no starting material was seen by analytical LCMS. The mixture was quenched with H₂O and extracted with EtOAc. The combined organic layers were washed twice with H₂O and once with brine, and then dried and concentrated. The residue was purified by silica gel chromatography (0-80% EtOAc in hexanes) to give the title compound. Mass spec. data (M+H) = 505.

Example 68: Synthesis of 2-(4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-

isoxazol-5-yl)-biphenyl-4-yl)-propionic acid ethyl ester (Compound 1-70)




Step 1: 2-(4-Bromo-phenyl)-propionic acid ethyl ester

[00680] To ethyl 4-bromophenylacetate (2.137g, 8.8mmol) in THF (20mL) at -78°C was added

lithium bis(trimethylsilyl)amide (1M in hexane; 11.4mL, 11.4mmol), and the solution was stirred for 1 hour at -78°C. Iodomethane (0.71mL, 11.4mmol) in THF (5mL) was added, and the reaction was warmed to room temperature. Analytical LCMS indicated that the reaction was complete, so the mixture was quenched with H₂O and concentrated. The residue was dissolved in EtOAc and washed three times with H₂O and once with brine. The organic layer was dried and


concentrated, and the crude material was purified by silica gel chromatography (0-60% EtOAc in hexanes) to give the title compound.



Step 2: 2-(4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid ethyl ester

[00681] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, 2-(4-bromo-phenyl)-propionic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride; further purification by preparative HPLC was required. Mass spec. data ($M+H$) = 533.


Example 69: Synthesis of 2-(4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid (Compound 1-71)



Step 1: 2-(4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid


[00682] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-(4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid ethyl ester. Mass spec. data ($M+H$) = 505.

Example 70: Synthesis of 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-72)




Step 1: 2-(4-Bromo-phenyl)-2-methyl-propionic acid ethyl ester

5 [00683] To sodium hydride (0.859g, 35.8mmol) and iodomethane (2.23mL, 35.8mmol) in DMF (80mL) at 0°C was added ethyl 4-bromophenylacetate (2.17g, 9.0mmol) dropwise, and the reaction was warmed to room temperature and stirred for 4 hours while being monitored by analytical LCMS. Once the reaction was complete, the mixture was quenched with H₂O and 1N aqueous HCl (20mL), and then extracted with 1:1 EtOAc:hexanes. The combined organic layers 10 were washed five times with H₂O and once with brine, and the combined aqueous layers were back-extracted with EtOAc. The combined organic layers were dried and concentrated, and the residue was purified by silica gel chromatography (0-20% EtOAc in hexanes). Further purification by preparative HPLC gave the title compound.



15 **Step 2: 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester**


[00684] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, 2-(4-bromo-phenyl)-2-methyl-propionic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride; further purification by preparative HPLC was required.



Step 3: 2-(4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid

[00685] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-(4'-(4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester; purification by preparative HPLC was utilized. Mass spec. data ($M+H$) = 519.


Example 71: Synthesis of [4-(6-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-pyridin-3-yl)-phenyl]-acetic acid (Compound 1-73)



10


Step 1: 5-Bromo-pyridine-2-carbonyl chloride

[00686] Prepared according to the procedure described in Example 29, Step 1 using the following starting material: 5-bromopyridine-2-carboxylic acid.




15 **Step 2: 2-(5-Bromo-pyridine-2-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester**

[00687] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 5-bromo-pyridine-2-carbonyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




Step 3: 5-(5-Bromo-pyridin-2-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00688] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(5-bromo-pyridine-2-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.



Step 4: 5-(5-Bromo-pyridin-2-yl)-3-methyl-isoxazole-4-carboxylic acid

[00689] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(5-bromo-pyridin-2-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




10

Step 5: [5-(5-Bromo-pyridin-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chlorophenyl)-ethyl ester


[00690] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(5-bromo-pyridin-2-yl)-3-methyl-isoxazole-4-carboxylic acid and 2-chloro-alpha-methylbenzyl alcohol.

15



Step 6: [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-3-yl)-phenyl]-acetic acid ethyl ester


[00691] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: [5-(5-bromo-pyridin-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chlorophenyl)-ethyl ester, [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride; purification by preparative HPLC was utilized.



10 **Step 7: [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-3-yl)-phenyl]-acetic acid**


[00692] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: [4-(6-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-3-yl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 492.

15 **Example 72: Synthesis of 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-74)**




Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-fluoro-phenyl)-ethyl ester

[00693] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 2-fluoro-alpha-methylbenzyl alcohol.



Step 2: {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-fluoro-phenyl)-ethyl ester


[00694] Prepared according to the procedure described in Example 49, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-fluoro-phenyl)-ethyl ester and bis(pinacolato)diboron.



Step 3: 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester

[00695] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-


isoxazol-4-yl}-carbamic acid 1-(2-fluoro-phenyl)-ethyl ester, 2-(4-bromo-phenyl)-propionic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride; purification by preparative HPLC, followed by silica gel chromatography, was utilized.



5 **Step 4: 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid**


[00696] Prepared according to the procedure described in Example 48, Step 3 using the following starting materials: 2-(4'-{4-[1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester. Mass spec. data (M+H) = 489.

10 **Example 73: Synthesis of 4-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid (Compound 1-75)**




Step 1: 4-(4-Bromo-phenyl)-butyric acid ethyl ester

[00697] Prepared according to the procedure described in Example 41, Step 1 using the following starting material: 4-(4-bromophenyl)butanoic acid.



Step 2: 4-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid ethyl ester


[00698] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester, 4-(4-bromo-phenyl)-butyric acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride.



Step 3: 4-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid

[00699] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 4-(4'-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid ethyl ester. Mass spec. data (M+H) = 519.

Example 74: Synthesis of 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid (Compound 1-76)




Step 1: 4'-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-3-carboxylic acid

[00700] [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl

5 ester (0.200g, 0.46mmol), 3-carboxyphenylboronic acid (0.092g, 0.55mmol), and potassium carbonate (0.190g, 1.37mmol) were combined in 2:1 DME:H₂O (5mL), and the solution was purged with N₂ for 10 minutes. Tetrakis(triphenylphosphine)palladium(0) (0.053g, 0.05mmol) was added, and the reaction was purged with N₂ for an additional 10 minutes, and then sealed and stirred at 80°C overnight. The mixture was partitioned between EtOAc and H₂O, and the 10 aqueous layer was extracted with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and concentrated, and the residue was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 477.

Example 75: Synthesis of (4'-(4-[1-(4-Chloro-2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-77)




15

Step 1: (4'-(4-[1-(4-Chloro-2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester

[00701] Prepared according to the procedure described in Example 29, Step 5 using the following


20 starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 1-(4-chloro-2-fluorophenyl)ethanol.



Step 2: (4'-(4-Chloro-2-fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid


[00702] Excess lithium hydroxide was added to a solution of (4'-(4-chloro-2-fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester (0.100g, 0.19mmol) in MeOH and H₂O, and the reaction was stirred at room temperature overnight. The mixture was acidified with 1N aqueous HCl and extracted with EtOAc, and the combined organic layers were dried, filtered, and concentrated. The residue was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 509.

10 **Example 76: Synthesis of (4'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-78)**




Step 1: (R)-1-(2-Fluoro-phenyl)-ethanol synthesis described on page 118

[00703] 2'-Fluoroacetophenone (25g, 180mmol) and (S)-(-)-2-methyl-CBS-oxazaborolidine (6.02g, 22mmol) were combined in THF (200mL). Borane methyl sulfide complex (2M in THF; 59.4mL, 119mmol) was added over 20 minutes, and the reaction was stirred at room temperature for 30 minutes. MeOH was added, and the mixture was worked-up with CH₂Cl₂ and H₂O. The organic layer was concentrated to give the title compound in 90.6% e.e.



Step 2: (4'-(4'-(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


5 [00704] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



Step 3: (4'-(4'-(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


10 [00705] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4'-(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 475.

Example 77: Synthesis of [2-(4-(4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-benzyloxy)-phenyl]-acetic acid (Compound 1-79)



Step 1: [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid ethyl ester


[00706] [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-5 ethyl ester (0.200g, 0.50mmol), ethyl 2-hydroxyphenylacetate (0.095g, 0.53mmol), and potassium carbonate (0.138g, 1.0mmol) were combined in DMSO and stirred at room temperature for 1.5 days to give the title compound.



Step 2: [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid


[00707] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [2-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 521.

Example 78: Synthesis of [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-80)




Step 1: [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00708] Prepared according to the procedure described in Example 1, Step 5 using the following starting materials: 5-(4-chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



Step 2: [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid methyl ester


[00709] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 3-((methoxycarbonyl)methyl)phenylboronic acid.



Step 3: [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid


[00710] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid methyl ester. Mass spec. data (M+H) = 489.

Example 79: Synthesis of [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-81)



Step 1: [4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-benzyl]-phenyl]-acetic acid ethyl ester


[00711] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 4-((ethoxycarbonyl)methyl)phenylboronic acid.



Step 2: [4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-benzyl]-phenyl]-acetic acid

[00712] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [4-(4-{(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-benzyl]-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 489.


Example 80: Synthesis of [4-Fluoro-3-(4-{(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-benzyl]-phenyl]-acetic acid (Compound 1-82)



15

Step 1: [4-Fluoro-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester


[00713] Prepared according to the procedure described in Example 19, Step 1 using the following starting materials: (3-bromo-4-fluoro-phenyl)-acetic acid ethyl ester and bis(pinacolato)diboron.



Step 2: 5-((Ethoxycarbonyl)methyl)-2-fluorophenylboronic acid

Prepared according to the procedure described in Example 19, Step 2 using the following starting material: [4-fluoro-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.


5



Step 3: [4-Fluoro-3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid ethyl ester

[00714] Prepared according to the procedure described in Example 5, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 5-((ethoxycarbonyl)methyl)-2-fluorophenylboronic acid.


10



Step 4: [4-Fluoro-3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid


[00715] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [4-fluoro-3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 507.

Example 81: Synthesis of [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid ethyl ester (Compound 1-83)




Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00716] Prepared according to the procedure described in Example 1, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



Step 2: {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester


[00717] Prepared according to the procedure described in Example 6, Step 2 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and bis(pinacolato)diboron.



Step 3: [5-(4-Hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00718] To {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (1.2g, 2.58mmol) in MeOH was added 1N aqueous NaOH (12.8mL, 12.8mmol) and hydrogen peroxide (30% in H2O; 1.45g, 12.8mmol), and the reaction was stirred at room temperature for 30 minutes. The mixture was acidified to pH


3, which resulted in the formation of a precipitate. The precipitate was collected and washed five times with H₂O, and then dried to give the title compound.



Step 4: [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid ethyl ester

[00719] [5-(4-Hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (0.100g, 0.28mmol), 4-((ethoxycarbonyl)methyl)phenylboronic acid (0.065g, 0.31mmol), copper(II) acetate (0.102g, 0.56mmol), and pyridine (0.070g, 0.89mmol) were combined in CH₂Cl₂ and stirred at room temperature for 18 hours. After aqueous work-up, the crude material was purified by silica gel chromatography (10-30% EtOAc in hexanes) to give the title compound. Mass spec. data (M+H) = 519.


Example 82: Synthesis of [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-84)



Step 1: [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid


[00720] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [4-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 491.

Example 83: Synthesis of (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-acetic acid (Compound 1-85)



Step 1: 4-Bromo-3-methyl-benzoyl chloride


[00721] Prepared according to the procedure described in Example 29, Step 1 using the following starting material: 4-bromo-3-methylbenzoic acid.



5

Step 2: 2-(4-Bromo-3-methyl-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester


[00722] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 4-bromo-3-methyl-benzoyl chloride and 3-methylamino-but-2-enoic acid methyl ester.



10

Step 3: 5-(4-Bromo-3-methyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester


[00723] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(4-bromo-3-methyl-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.



15


Step 4: 5-(4-Bromo-3-methyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid

[00724] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(4-bromo-3-methyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




Step 5: [5-(4-Bromo-3-methyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00725] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-3-methyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



Step 6: (4'-{4-[{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-acetic acid ethyl ester


[00726] Prepared according to the procedure described in Example 43, Step 1 using the following starting materials: [5-(4-bromo-3-methyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-acetic acid


[00727] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (4'-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 489.

Example 84: Synthesis of 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-86)



Step 1: 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester

[00728] Prepared according to the procedure described in Example 43, Step 1 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-fluoro-phenyl)-ethyl ester and 2-(4-bromo-phenyl)-2-methyl-propionic acid ethyl ester; purification by preparative HPLC was utilized.




Step 2: 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid

[00729] 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester (0.51mmol) in MeOH (50mL) was treated with lithium hydroxide (excess) as well as 1N aqueous LiOH (excess, and the reaction was stirred at room


temperature. Analytical LCMS indicated that starting material was still present, so the reaction was stirred at 80°C overnight. Once no starting material was seen by analytical LCMS, the mixture was neutralized with 1N HCl (20mL) and extracted with EtOAc. The combined organic layers were washed with H₂O, dried, and concentrated, and the residue was purified by 5 preparative HPLC to give the title compound. Mass spec. data (M+H) = 503.

Example 85: Synthesis of 2-[6-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-2-methyl-propionic acid (Compound 1-87)




Step 1: 2-(6-Chloro-pyridin-3-yl)-2-methyl-propionic acid methyl ester

10 [00730] Potassium bis(trimethylsilyl)amide (0.5M in toluene; 13.68mL, 6.84mmol) was cooled to -20°C. (6-Chloro-pyridin-3-yl)-acetic acid methyl ester (0.425g, 2.28mmol) in THF (20mL) was added dropwise via syringe over 5 minutes. The solution was stirred for 30 minutes, and then iodomethane (0.33mL, 5.24mmol) was added dropwise. The reaction was warmed to room temperature and stirred overnight. The mixture was quenched with H₂O and NH₄Cl, and 15 extracted with EtOAc. The combined organic layers were washed with H₂O, dried, and concentrated, and the residue was purified by preparative HPLC to give the title compound.



Step 2: 2-[6-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-2-methyl-propionic acid methyl ester


20 [00731] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-fluoro-phenyl)-ethyl ester and 2-(6-chloro-pyridin-3-yl)-2-methyl-propionic acid methyl ester; further purification by preparative HPLC was required.



Step 3: 2-[6-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-2-methyl-propionic acid


[00732] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-[6-(4-{4-[1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-2-methyl-propionic acid methyl ester. Mass spec. data ($M+H$) = 504.

Example 86: Synthesis of (4'-{4-[{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-88)



10 **Step 1: 3-Methylamino-but-2-enoic acid methyl ester**


[00733] To a solution of methyl acetoacetate (29.4g, 253mmol) in MeOH (30mL) was added methylamine (33 wt% in EtOH; 48mL, 385mmol) dropwise at room temperature. The reaction was stirred for 1 hour, and then concentrated and dried to give the title compound as a white crystalline solid.



15


Step 2: 2-(4-Bromo-benzoyl)-3-oxo-butyric acid methyl ester

[00734] To 3-methylamino-but-2-enoic acid methyl ester (5.0g, 39.1mmol) in THF (70mL) was added pyridine (3.7mL, 47mmol) dropwise. The mixture was cooled to 0°C, and 4-bromobenzoyl chloride (8.55g, 39.1mmol) in THF (30mL) was added dropwise. The reaction was stirred at room temperature overnight, and then H₂O was added. The mixture was extracted with EtOAc, and the combined organic layers were washed with H₂O, dried, filtered, and concentrated to give the title compound.




Step 3: 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00735] To a mixture of 2-(4-bromo-benzoyl)-3-oxo-butyric acid methyl ester (11g, 39mmol) in acetic acid (50mL) was added hydroxylamine hydrochloride (2.66g, 39mmol), and the reaction 5 was stirred at 115°C for 1 hours. After cooling, saturated aqueous NaHCO₃ was added to the mixture to adjust to pH 8. The solution was extracted with EtOAc, and the combined organic layers were washed with brine, dried, filtered, and concentrated to give the title compound.




Step 4: 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid

10 [00736] Lithium hydroxide (2g, 48mmol) was added to a solution of 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester (39mmol) in MeOH (50mL) and H₂O (10mL), and the reaction was stirred at 60°C for 1 hour. Acidic work-up gave the title compound.



15 **Step 5: (R)-1-(2-Chloro-phenyl)-ethanol**


[00737] 2'-Chloroacetophenone (8.4mL, 65mmol) and (S)-(-)-2-methyl-CBS-oxazaborolidine (0.90g, 0.32mmol) were combined in THF (75mL). Borane methyl sulfide complex (2M in THF; 21.5mL, 43mmol) was added over 20 minutes, and the reaction was stirred at room temperature for 30 minutes. MeOH was added, and the mixture was worked-up with CH₂Cl₂ and H₂O. The 20 organic layer was concentrated to give the title compound.



Step 6: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester

[00738] 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (10g, 35mmol), (R)-1-(2-chloro-phenyl)-ethanol (6.6g, 42mmol), triethylamine (10mL, 70mmol), and diphenylphosphoryl

5 azide (11.5g, 42mmol) were combined in toluene (100mL) and stirred at 90°C for 1 hour. The mixture was concentrated, and the residue was purified by silica gel chromatography (0-30% EtOAc in hexanes). The isolated product was recrystallized in 5:1 hexanes:acetone to give the title compound in >99% e.e (by chiral HPLC. Chiracel OD 98.4% hexanes/1.6% Ethanol. Major isomer 27.9 min minor isomer 32.7 min).




10

Step 7: [4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester

[00739] Ethyl 4-bromophenylacetate (9.72g, 40mmol), bis(pinacolato)diboron (12.2g, 48mmol), and potassium acetate (18.8g, 1690mmol) were combined in 1,4-dioxane, and the mixture was purged with N₂ for 10 minutes. 1,1'-Bis(diphenylphosphino)ferrocene)-dichloropalladium(II)

15 (3.26g, 4.0mmol) was added, and the reaction was heated to 80°C for 24 hours, and at room temperature for 2 days. The mixture was diluted with H₂O and extracted with EtOAc. The combined organic layers were washed with 4 times with H₂O, once with brine, and then dried, filtered, and concentrated. The residue was dissolved in CH₂Cl₂ and filtered through Celite to remove solids. The filtrate was concentrated, and the crude material was purified by silica gel

20 chromatography (0-30% EtOAc in hexanes) to give the title compound.




Step 8: (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester

[00740] [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-

25 ethyl ester (3.0g, 6.9mmol), [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid

ethyl ester (2.2g, 7.6mmol), dichloro-bis(triphenylphosphine)palladium(II) (0.25g, 0.35mmol), and sodium bicarbonate (1.7g, 20.7mmol) were combined in DME (30mL) and H₂O (10mL), and the reaction was stirred at 80°C for 2 hours. After aqueous work-up, the crude material was purified by silica gel chromatography (0-40% EtOAc in hexanes) to give the title compound.




5

Step 9: (4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00741] 4'-{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester (3g, 5.9mmol) in MeOH and H₂O was treated with lithium hydroxide (1g, 23.8mmol), and the reaction was stirred at 60°C. Acidic work-up provided the title compound. Mass spec. data (M+H) = 491.

Example 87: Synthesis of 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-89)




15 **Step 1: 2-Methyl-2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester**

[00742] Prepared according to the procedure described in Example 49, Step 1 using the following starting materials: 2-(4-Bromo-phenyl)-2-methyl-propionic acid ethyl ester and bis(pinacolato)diboron.



Step 2: 2-(4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester


[00743] Prepared according to the procedure described in Example 43, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and 2-methyl-2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester.



Step 3: 2-(4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid


[00744] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-(4'-(4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester. Mass spec. data (M+H) = 519.

Example 88: Synthesis of 2-(4'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-90)



Step 1: 2-(4'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester


[00745] Prepared according to the procedure described in Example 43, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 2-methyl-2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester.



Step 2: 2-(4'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid


[00746] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-(4'-(4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-2-methyl-propionic acid ethyl ester. Mass spec. data (M+H) = 503.

Example 89: Synthesis of 2-(4'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid (Compound 1-91)




Step 1: 2-[4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester

[00747] Prepared according to the procedure described in Example 49, Step 1 using the following 5 starting materials: 2-(4-bromo-phenyl)-propionic acid ethyl ester and bis(pinacolato)diboron.



Step 2: 2-(4'-(4-[R]-1-(2-fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-biphenyl-4-yl-propionic acid ethyl ester


[00748] Prepared according to the procedure described in Example 43, Step 1 using the following 10 starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 2-[4-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester.



Step 3: 2-(4'-(4-[R]-1-(2-fluoro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-biphenyl-4-yl-propionic acid

[00749] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-(4'-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester. Mass spec. data (M+H) = 489.


Example 90: Synthesis of 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-92)



Step 1: 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester

[00750] Prepared according to the procedure described in Example 43, Step 1 using the following


10 starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and 2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester.



Step 2: 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid

[00751] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 2-(4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester. Mass spec. data (M+H) = 505.

Example 91: Synthesis of 2-(4-{4-[(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-4-carboxylic acid (Compound 1-93)




Step 1: 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-4-carboxylic acid ethyl ester

[00752] {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-

5 carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.100g, 0.21mmol), ethyl 2-bromothiazole-4-carboxylate (0.076g, 0.32mmol), potassium carbonate (0.073g, 0.53mmol), and tetrakis(triphenylphosphine)palladium(0) (0.023g, 0.02mmol) were combined in 2:1 DME:H₂O, and the mixture was purged with N₂ for 5 minutes. The reaction was stirred at 90°C for 3 hours, until no starting material was seen by analytical LCMS. The solution was used directly in the

10 hydrolysis step.




Step 2: 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-4-carboxylic acid

[00753] 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-


15 thiazole-4-carboxylic acid ethyl ester (0.21mmol) was treated with 1N aqueous LiOH (excess), and the reaction was stirred for 2 hours at room temperature. Once no starting material was seen by analytical LCMS, the mixture was acidified with 1N aqueous HCl and extracted with EtOAc. The combined organic layers were concentrated, and the residue was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 484.

20 **Example 92: Synthesis of 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-5-carboxylic acid (Compound 1-94)**



Step 1: 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-5-carboxylic acid ethyl ester


[00754] Prepared according to the procedure described in Example 91, Step 1 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and ethyl 2-bromothiazole-5-carboxylate.



Step 2: 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-5-carboxylic acid

[00755] Prepared according to the procedure described in Example 91, Step 2 using the following starting material: 2-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-5-carboxylic acid ethyl ester. Mass spec. data (M+H) = 484.


Example 93: Synthesis of {5-[4-(4-Cyclopropylcarbamoyl-benzyl)-phenyl]-3-methyl-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (Compound 2-1)



Step 1: {5-[4-(4-Cyclopropylcarbamoyl-benzyl)-phenyl]-3-methyl-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester

[00756] Prepared according to the procedure described in Example 5, Step 2, using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 4-(cyclopropylaminocarbonyl)phenylboronic acid. Mass spec. data ($M+H$) = 530.

Example 94: Synthesis of [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Compound 2-2)




10

Step 1: [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester

[00757] Prepared according to the procedure described in Example 43, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and 4-cyanomethylphenylboronic acid. Mass spec. data ($M+H$) = 472.


Example 95: Synthesis of [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-3)



Step 1: [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00758] Prepared according to the procedure described in Example 43, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 4-cyanomethylphenylboronic acid. Mass spec. data ($M+H$) = 456.


Example 96: Synthesis of {3-Methyl-5-[4'-(2*H*-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-4)



10 **Step 1: {3-Methyl-5-[4'-(2*H*-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester**

[00759] To a solution of [5-(4'-cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (0.200g, 0.44mmol) in toluene (5mL) was added dibutyltin oxide (0.011g, 0.04mmol), followed by azidotrimethylsilane (0.07mL, 0.53mmol), and the reaction was stirred at 100°C for 5 hours, and then stirred at room temperature overnight. The mixture was concentrated, and the residue was dissolved in EtOAc and washed with H₂O and brine. The organic layer was dried and concentrated, and the crude material was purified by preparative HPLC to give the title compound. Mass spec. data ($M+H$) = 499.


20 **Example 97: Synthesis of {3-Methyl-5-[4'-(2*H*-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Compound 2-5)**



Step 1: {3-Methyl-5-[4'-(2H-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester

[00760] Prepared according to the procedure described in Example 100, Step 1 using the following starting materials: [5-(4'-cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and azidotrimethylsilane. Mass spec. data ($M+H$) = 515.


Example 98: Synthesis of [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-phenyl]-acetic acid (Compound 3-1)



10


Step 1: 6-Bromo-naphthalene-2-carbonyl chloride

[00761] 6-Bromo-2-naphthoic acid (5.0g, 19.9mmol) was treated with thionyl chloride (20mL) and stirred at reflux for 2 hours. The mixture was concentrated to give the title compound.



15 **Step 2: 2-(6-Bromo-naphthalene-2-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester**


[00762] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 6-bromo-naphthalene-2-carbonyl chloride and 3-methylamino-but-2-enoic acid methyl ester.



Step 3: 5-(6-Bromo-naphthalen-2-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00763] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(6-bromo-naphthalene-2-carbonyl)-3-[(E)-methylimino]-butyric acid methyl


5 ester and hydroxylamine hydrochloride.



Step 4: 5-(6-Bromo-naphthalen-2-yl)-3-methyl-isoxazole-4-carboxylic acid

[00764] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(6-bromo-naphthalen-2-yl)-3-methyl-isoxazole-4-carboxylic acid methyl


10 ester.



Step 5: [5-(6-Bromo-naphthalen-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chlorophenyl)-ethyl ester

[00765] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(6-bromo-naphthalen-2-yl)-3-methyl-isoxazole-4-carboxylic acid and 2-


15 chloro-alpha-methylbenzyl alcohol.



Step 6: [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-phenyl]-acetic acid ethyl ester

[00766] Prepared according to the procedure described in Example 91, Step 1 using the following


5 starting materials: [5-(6-bromo-naphthalen-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-phenyl]-acetic acid

[00767] Prepared according to the procedure described in Example 91, Step 2 using the following starting material: [4-(6-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 541.


Example 99: Synthesis of 3-[4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-benzylsulfanyl]-propionic acid (Compound 3-2)



Step 1: 3-(4-Bromo-benzylsulfanyl)-propionic acid methyl ester

[00768] To a solution of 4-bromobenzyl bromide (6.0g, 24.0mmol) in THF (20mL) was added triethylamine (6.7mL, 48.0mmol), followed by methyl 3-mercaptopropionate (2.9mL,


5 26.4mmol), and the reaction was stirred at room temperature for 2 hours. Once no starting material was seen by analytical tlc, the mixture was worked-up with EtOAc and H₂O, and the residue was purified by silica gel chromatography to give the title compound.



Step 2: 3-[4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzylsulfanyl]-propionic acid

10 **methyl ester**


[00769] Prepared according to the procedure described in Example 49, Step 1 using the following starting materials: 3-(4-bromobenzylsulfanyl)-propionic acid methyl ester and bis(pinacolato)diboron.



15 **Step 3: 3-[4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-benzylsulfanyl]-propionic acid methyl ester**

[00770] Prepared according to the procedure described in Example 91, Step 1 using the following starting materials: [5-(6-bromo-naphthalen-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and 3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-

20 benzylsulfanyl]-propionic acid methyl ester.




Step 4: 3-[4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-benzylsulfanyl]-propionic acid

[00771] Prepared according to the procedure described in Example 91, Step 2 using the following


5 starting material: 3-[4-(6-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-benzylsulfanyl]-propionic acid methyl ester. Mass spec. data (M+H) = 601.

Example 100: Synthesis of [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-3)




10 **Step 1: 2-Bromo-thiazole-4-carbonyl chloride**

[00772] Prepared according to the procedure described in Example 29, Step 1 using the following starting material: 2-bromo-4-thiazolecarboxylic acid.



Step 2: 2-(2-Bromo-thiazole-4-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester

15 [00773] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 2-bromo-thiazole-4-carbonyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




Step 3: 5-(2-Bromo-thiazol-4-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00774] Prepared according to the procedure described in Example 29, Step 3 using the following

starting materials: 2-(2-bromo-thiazole-4-carbonyl)-3-[(E)-methylimino]-butyric acid methyl


5 ester and hydroxylamine hydrochloride.



Step 4: 5-(2-Bromo-thiazol-4-yl)-3-methyl-isoxazole-4-carboxylic acid

[00775] Prepared according to the procedure described in Example 29, Step 4 using the following

starting material: 5-(2-bromo-thiazol-4-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




10

Step 5: [5-(2-Bromo-thiazol-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00776] Prepared according to the procedure described in Example 29, Step 5 using the following

starting materials: 5-(2-bromo-thiazol-4-yl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-


15 fluoro-phenyl)-ethanol.



Step 6: [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid ethyl ester

[00777] Prepared according to the procedure described in Example 91, Step 1 using the following starting materials: [5-(2-bromo-thiazol-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-


5 fluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid

10 [00778] Prepared according to the procedure described in Example 91, Step 2 using the following starting material: [4-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 482.


Example 101: Synthesis of [4-(5-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-4)



15


Step 1: 2-Bromo-thiazole-5-carbonyl chloride

[00779] Prepared according to the procedure described in Example 29, Step 1 using the following starting material: 2-bromo-5-thiazolecarboxylic acid.




20 **Step 2: 2-(2-Bromo-thiazole-5-carbonyl)-3-[(E)-methylimino]butyric acid methyl ester**

[00780] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 2-bromo-thiazole-5-carbonyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




5 **Step 3: 5-(2-Bromo-thiazol-5-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester**

[00781] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(2-bromo-thiazole-5-carbonyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.




10 **Step 4: 5-(2-Bromo-thiazol-5-yl)-3-methyl-isoxazole-4-carboxylic acid**

[00782] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(2-bromo-thiazol-5-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




Step 5: [5-(2-Bromo-thiazol-5-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00783] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(2-bromo-thiazol-5-yl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



Step 6: [4-(5-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid ethyl ester


[00784] Prepared according to the procedure described in Example 91, Step 1 using the following starting materials: [5-(2-bromo-thiazol-5-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: [4-(5-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid


[00785] Prepared according to the procedure described in Example 91, Step 2 using the following starting material: [4-(5-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 482.

Example 102: Synthesis of {4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ((Compound 1-115)



Step 1: (R)-1-Phenyl-ethanol

[00786] To acetophenone (19.47mL, 166.6mmol) in THF (100mL) was added (S)-(-)-2-methyl-CBS-oxazaborolidine (4.62g, 16.6mmol), and the reaction was cooled to 0°C. Borane methyl sulfide complex (2M in THF; 50mL, 100mmol) was added over 15 minutes, and the reaction was stirred at room temperature. Aqueous work-up gave the title compound.




Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester

[00787] 5-(4-Bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid (25g, 88.7mmol) in toluene

(500mL) was added triethylamine (18.5mL, 133mmol), followed by diphenylphosphoryl azide

5 (22.1mL, 101.9mmol). (R)-(+)-1-Phenylethyl alcohol (11.9mL, 97.5mmol) was added, and the reaction was stirred at 75°C for 2 hours. The mixture was partitioned between EtOAc and H₂O and filtered through Celite. The aqueous layer was extracted with EtOAc, and the combined organic layers were dried over MgSO₄, filtered, and concentrated to give the title compound.




10 **Step 3: {4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester**

[00788] [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester

(39g, 97.2mmol), [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester (31g, 107mmol), and sodium bicarbonate (32.6g, 389mmol) were combined in 3:1

15 DME:H₂O (500mL), and the mixture was purged with N₂ for 15 minutes. (1,1'-Bis(diphenylphosphino)ferrocene)-dichloropalladium(II) (2.13g, 2.91mmol) was added, and the reaction was purged with N₂ for an additional 10 minutes and then stirred at 90°C overnight. The mixture was partitioned between EtOAc and H₂O, and the aqueous layer was extracted with EtOAc. The combined organic layers were washed with H₂O, dried over MgSO₄, filtered, and 20 concentrated, and the residue was purified by silica gel chromatography (EtOAc/hexane gradient) to give the title compound.




Step 4: {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid

[00789] To a suspension of {4'-(3-methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-

5 biphenyl-4-yl}-acetic acid ethyl ester (24g, 49mmol) in 3:1 MeOH:H₂O (300mL) was added lithium hydroxide (8.3g, 198mmol), and the reaction was stirred at room temperature overnight. The mixture was acidified and partitioned between EtOAc and H₂O. The aqueous layer was extracted with EtOAc, and the combined organic layers were washed with H₂O, dried over MgSO₄, filtered, and concentrated to give the title compound. Mass spec. data (M+H) = 457.


10 **Example 104: Synthesis of {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-3-yl}-acetic acid (Compound 1-121)**



Step 1: (3-Bromo-phenyl)-acetic acid ethyl ester


[00790] Prepared according to the procedure described in Example 21, Step 1 using the following

15 starting materials: 3-bromophenylacetic acid and ethanol.




Step 1: [3-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester

[00791] Prepared according to the procedure described in Example 49, Step 1 using the following starting materials: (3-Bromo-phenyl)-acetic acid ethyl ester and bis(pinacolato)diboron.



Step 2: {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-3-yl}-acetic acid ethyl ester


[00792] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester, [3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester, and bis(triphenylphosphine)palladium(II) dichloride.



Step 3: {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-3-yl}-acetic acid


[00793] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: {4'-(3-methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-3-yl}-acetic acid ethyl ester. Mass spec. data (M+H) = 457.

Example 105: Synthesis of 4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-carboxylic acid (Compound 1-122)



Step 1: 4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-carboxylic acid ethyl ester


[00794] Prepared according to the procedure described in Example 48, Step 2 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-5-ethyl ester and 4-ethoxycarbonylphenylboronic acid.



Step 2: 4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-carboxylic acid

[00795] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: 4'-[3-methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-carboxylic acid ethyl ester. Mass spec. data (M+H) = 443.


Example 106: Synthesis of [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-109)



15 **Step 1: 3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-propionic acid 2-ethyl-hexyl ester**


[00796] [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (2g, 4.8mmol), 3-mercaptopropionic acid 2-ethylhexyl ester (1.04g, 4.8mmol), diisopropylethylamine (1.7mL, 9.5mmol), (dibenzylideneacetone)dipalladium(0) (0.109g, 0.12mmol), and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.123g, 0.24mmol) were combined in 1,4-dioxane (10mL), and the reaction was purged with N₂ for 10 minutes and then stirred at 110°C for 16 hours. Once no starting material was seen by analytical LCMS, the

mixture was filtered, the filtrate was concentrated, and the residue was purified by silica gel chromatography to give the title compound.




Step 2: [5-(4-Mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00797] A solution of 3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-propionic acid 2-ethyl-hexyl ester (0.5g, 0.90mmol) in 1,4-dioxane (5mL) was purged with N₂ for 5 minutes, and then treated with potassium *tert*-butoxide (0.111g, 0.94mmol). The reaction was purged with N₂ for 3 minutes, and then stirred overnight at room temperature to give the title compound, which was used as a solution in 1,4-dioxane in the next step.



Step 3: [4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester


[00798] A solution of [5-(4-mercaptophenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (0.9mmol) in 1,4-dioxane was purged with N₂ for 3 minutes. Ethyl 4-bromophenylacetate (0.218g, 0.90mmol), diisopropylethylamine (0.3mL, 1.8mmol), (dibenzylideneacetone)dipalladium(0) (0.021g, 0.02mmol), and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.026g, 0.04mmol) were added, and the reaction was purged with N₂ for another 3 minutes and then stirred at 110°C for 2 hours. Once no starting material was seen by analytical LCMS, the mixture was filtered, the filtrate was concentrated, and the residue was purified by silica gel chromatography to give the title compound.



Step 4: [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid


[00799] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: [4-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 507.

Example 107: Synthesis of [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-103)



10 **Step 1: [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester**


[00800] Prepared according to the procedure described in Example 1, Step 5, using the following starting materials: 5-(4-chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-chloro-phenyl)-ethanol.



Step 2: [4-(4-{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester

[00801] Methyl 4-hydroxyphenylacetate (0.145g, 0.88mmol) in MeCN was treated with cesium carbonate (0.098g, 0.30mmol), and the mixture was stirred at 60°C for 30 minutes and then


5 cooled to room temperature. [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (0.100g, 0.25mmol) was added, and the reaction was stirred at room temperature for 4 hours. Once no starting material was seen by analytical LCMS, the mixture was purified by preparative HPLC to give the title compound.



10 **Step 3: [4-(4-{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid**


[00802] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [4-(4-{(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester. Mass spec. data (M+H) = 521.

15 **Example 108: Synthesis of [4-(4-{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-138)**




Step 1: {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester

[00803] Prepared according to the procedure described in Example 6, Step 2 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and bis(pinacolato)diboron.



5 **Step 2: [5-(4-Hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester**


[00804] Prepared according to the procedure described in Example 81, Step 3 using the following starting material: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester.



10

Step 3: [4-(4-{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-phenoxy]-phenyl]-acetic acid ethyl ester


[00805] Prepared according to the procedure described in Example 81, Step 4 using the following starting materials: [5-(4-hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and 4-ethoxycarbonylmethylphenylboronic acid.



Step 4: [4-(4-{(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-phenoxy]-phenyl]-acetic acid

[00806] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [4-(4-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 507.


5 **Example 109: Synthesis of [3-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester (Compound 1-95)**



Step 1: [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester

10 [00807] Prepared according to the procedure described in Example 81, Step 4 using the following starting materials: [5-(4-hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 3-((methoxycarbonyl)methyl)phenylboronic acid. Mass spec. data (M+H) = 505.

Example 110 : Synthesis of [3-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-96)




15

Step 1: [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid

20 [00808] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester. Mass spec. data (M+H) = 491.

Example 111 : Synthesis of [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-97)




Step 1: [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid

[00809] [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-

5 benzylsulfonyl)-phenyl]-acetic acid (0.050g, 0.09mmol) in CH_2Cl_2 (2mL) and EtOH (1mL) was treated with meta-chloroperoxybenzoic acid (0.104g, 0.46mmol), and the reaction was stirred at room temperature for 1 hour. The crude material was purified by preparative HPLC to give the title compound. Mass spec. data ($\text{M}+\text{H}$) = 569.

Example 112 : Synthesis of [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-

10 **isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-98)**




Step 1: [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid

[00810] Prepared according to the procedure described in Example 111, Step 1 using the

15 following starting material: [3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfonyl)-phenyl]-acetic acid. Mass spec. data ($\text{M}+\text{H}$) = 569.


Example 113: Synthesis of [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-99)



Step 1: [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid


[00811] Prepared according to the procedure described in Example 111, Step 1 using the following starting material: [2-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid. Mass spec. data (M+H) = 569.

Example 114: Synthesis of [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-100)



10 **Step 1: {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-ylmethyl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester**


[00812] Prepared according to the procedure described in Example 49, Step 1 using the following starting materials: [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester and bis(pinacolato)diboron.



15


Step 2: {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-ylmethyl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester

[00813] Prepared according to the procedure described in Example 81, Step 3 using the following starting material: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-ylmethyl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester.




5 **Step 3: [5-(4-Bromomethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester**

[00814] {3-Methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-ylmethyl)-phenyl]-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.390g, 1.0mmol) and phosphorus tribromide (0.142mL, 1.5mmol) were combined in DME, and the reaction was stirred at room temperature 10 for 1.5 hours. The mixture was partitioned between H₂O and EtOAc, and the organic layer was separated, concentrated, and purified by silica gel chromatography to give the title compound.



Step 4: [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester


15 [00815] [5-(4-Bromomethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (0.080g, 0.19mmol), methyl 4-hydroxyphenylacetate (0.040g, 0.23mmol), and cesium carbonate (0.093g, 0.29mmol) were combined in MeCN and stirred at room temperature for 30 minutes. The mixture was partitioned between H₂O and EtOAc, and the organic layer was separated and concentrated to give the title compound.



Step 5: [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid


[00816] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [4-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester. Mass spec. data ($M+H$) = 521.

Example 115: Synthesis of (4'-{4-[1-(2,6-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-101)



10 **Step 1: (4'-{4-[1-(2,6-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester**


[00817] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2,6-dichlorophenyl)ethanol.



Step 2: (4'-(4-[1-(2,6-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid

[00818] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[1-(2,6-dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 525.


Example 116: Synthesis of [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester (Compound 1-102)



10 **Step 1: [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester**


[00819] Prepared according to the procedure described in Example 107, Step 2, using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and methyl 4-hydroxyphenylacetate. Mass spec. data ($M+H$) = 535.

Example 117 : Synthesis of 4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)methyl)-benzoic acid (Compound 1-104)



Step 1: 4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-phenoxy-methyl-benzoic acid methyl ester


[00820] Prepared according to the procedure described in Example 114, Step 4 using the following starting materials: [5-(4-hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and methyl 4-(bromomethyl)benzoate.



Step 2: 4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-phenoxy-methyl-benzoic acid

[00821] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: 4-(4-{(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-phenoxy-methyl-benzoic acid methyl ester. Mass spec. data (M+H) = 491.


Example 118: Synthesis of [4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-phenylsulfanyl]-phenyl]-acetic acid ethyl ester (Compound 1-105)



Step 1: [5-(4-Mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00822] Prepared according to the procedure described in Example 106, Step 2 using the


5 following starting material: 3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-propionic acid 2-ethyl-hexyl ester.



Step 2: [4-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester

10 [00823] Prepared according to the procedure described in Example 106, Step 3 using the following starting materials: [5-(4-mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and ethyl 4-bromophenylacetate. Mass spec. data (M+H) = 535.

Example 119: Synthesis of [3-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-106)




15

Step 1: [5-(4-Mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00824] Prepared according to the procedure described in Example 106, Step 2 using the


20 following starting material: 3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-propionic acid 2-ethyl-hexyl ester.



Step 2: [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester

[00825] Prepared according to the procedure described in Example 106, Step 3 using the


5 following starting materials: [5-(4-mercaptop-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and ethyl 2-(3-bromophenyl)acetate.



Step 3: [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid

10 [00826] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: [3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 507.


Example 120: Synthesis of [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-107)



15


Step 1: [5-(4-Mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00827] Prepared according to the procedure described in Example 106, Step 2 using the following starting material: 3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-propionic acid 2-ethyl-hexyl ester.



5 **Step 2: [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester**

[00828] Prepared according to the procedure described in Example 106, Step 3 using the following starting materials: [5-(4-mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and ethyl 2-bromophenylacetate.




10

Step 3: [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid


[00829] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: [2-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 507.

Example 121: Synthesis of [6-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-pyridin-3-yl]-acetic acid (Compound 1-108)




Step 1: [5-(4-Mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00830] Prepared according to the procedure described in Example 106, Step 2 using the following starting material: 3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-propionic acid 2-ethyl-hexyl ester.



Step 2: [6-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-pyridin-3-yl]-acetic acid methyl ester


[00831] Prepared according to the procedure described in Example 106, Step 3 using the following starting materials: [5-(4-mercapto-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and (6-chloro-pyridin-3-yl)-acetic acid methyl ester.



Step 3: [6-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-pyridin-3-yl]-acetic acid


[00832] Prepared according to the procedure described in Example 48, Step 3 using the following starting material: [6-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-pyridin-3-yl]-acetic acid methyl ester. Mass spec. data (M+H) = 508.

Example 122: Synthesis of 2-(4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-propionic acid (Compound 1-110)



Step 1: 2-(4'-(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-2'-methyl-biphenyl-4-yl)-propionic acid ethyl ester


[00833] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-3-methyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionic acid ethyl ester.



Step 2: 2-(4'-(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-2'-methyl-biphenyl-4-yl)-propionic acid


[00834] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 2-(4'-(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-2'-methyl-biphenyl-4-yl)-propionic acid ethyl ester. Mass spec. data (M+H) = 503.

Example 123: Synthesis of (4'-(S)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-111)




Step 1: (S)-1-(2-Fluoro-phenyl)-ethanol

[00835] Prepared according to the procedure described in Example 76, Step 1 using the following starting materials: 2'-fluoroacetophenone and (R)-(+)-2-methyl-CBS-oxazaborolidine.



Step 2: (4'-(4-[(S)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00836] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and (S)-1-(2-fluoro-phenyl)-ethanol.



10 Step 3: (4'-(4-[(S)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00837] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[(S)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 475.

15 Example 124: Synthesis of (4'-(4-[(S)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-112)



Step 1: (S)-1-(2-Chloro-phenyl)-ethanol


[00838] Prepared according to the procedure described in Example 76, Step 1 using the following starting materials: 2'-chloroacetophenone and (R)-(+)-2-methyl-CBS-oxazaborolidine.



Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (S)-1-(2-chloro-phenyl)-ethyl ester

5


[00839] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and (S)-1-(2-chloro-phenyl)-ethanol.



10 Step 3: (4'-(S)-1-(2-Chloro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester

[00840] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (S)-1-(2-chloro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.


15



Step 4: (4'-(4-[(S)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00841] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[(S)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 491.

Example 125: Synthesis of {4'-(4-Chlorobenzyloxycarbonylamino)-3-methyl-isoxazol-5-yl}-biphenyl-4-yl}-acetic acid (Compound 1-113)




10 **Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 2-chloro-benzyl ester**

[00842] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials :5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and chlorobenzyl alcohol.



Step 2: {4'-[4-(2-Chloro-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester


[00843] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 2-chloro-benzyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 2: {4'-[4-(2-Chloro-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid


[00844] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: {4'-[4-(2-chloro-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester. Mass spec. data ($M+H$) = 477.

Example 126: Synthesis of [2-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-114)



15 Step 1: [2-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid ethyl ester


[00845] Prepared according to the procedure described in Example 107, Step 2 using the following starting materials: [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and ethyl 2-hydroxyphenylacetate.



Step 2: [2-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid


[00846] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: [2-(4-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 521.

Example 127: Synthesis of (4'-{4-[1-(2,3-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-116)



10 **Step 1: 1-(2,3-Difluoro-phenyl)-ethanol**


[00847] 2',3'-Difluoroacetophenone (0.337g, 2.16mmol) in MeOH (15mL) was treated with sodium borohydride (0.090g, 2.37mmol), and the reaction was stirred at room temperature for 30 minutes. The mixture was partitioned between CH₂Cl₂ and H₂O, and the organic layer was separated, dried over MgSO₄, filtered, and concentrated to give the title compound.



15


Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,3-difluoro-phenyl)-ethyl ester

[00848] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2,3-difluoro-phenyl)-ethanol.



Step 3: (4'-(4-[1-(2,3-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00849] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,3-difluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: (4'-(4-[1-(2,3-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00850] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[1-(2,3-difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 493.

Example 128: Synthesis of (4'-(4-[1-(2,4-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-117)




Step 1: 1-(2,4-Difluorophenyl)ethanol

[00851] Prepared according to the procedure described in Example 127, Step 1 using the following starting material: 2',4'-difluoroacetophenone.




5 **Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,4-difluoro-phenyl)-ethyl ester**

[00852] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2,4-difluorophenyl)ethanol.



10 **Step 3: (4'-{4-[1-(2,4-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester**

[00853] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,4-difluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: (4'-(4-[1-(2,4-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00854] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[1-(2,4-difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 493.

Example 129: Synthesis of (4'-(4-[1-(2-Fluoro-4-methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-118)




10 **Step 1: 1-(2-Fluoro-4-methoxyphenyl)ethan-1-ol**

[00855] Prepared according to the procedure described in Example 127, Step 1 using the following starting material: 2'-fluoro-4'-methoxyacetophenone.




15 **Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-fluoro-4-methoxy-phenyl)-ethyl ester**

[00856] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2-fluoro-4-methoxyphenyl)ethan-1-ol.



Step 3: (4'-(4-[1-(2-Fluoro-4-methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester

[00857] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2-fluoro-4-methoxy-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: (4'-(4-[1-(2-Fluoro-4-methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00858] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[1-(2-fluoro-4-methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 505.

Example 130: Synthesis of (4'-(4-[1-(2,5-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-119)




Step 1: 1-(2,5-Difluorophenyl)ethanol

[00859] Prepared according to the procedure described in Example 127, Step 1 using the following starting material: 2',5'-difluoroacetophenone.




Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,5-difluoro-phenyl)-5-ethyl ester

[00860] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2,5-difluorophenyl)ethanol.



10 Step 3: (4'-{4-[1-(2,5-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester


[00861] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,5-difluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: (4'-(4-[1-(2,5-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00862] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[1-(2,5-difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 493.

Example 131: Synthesis of (4'-(4-[1-(2,6-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-120)




10 **Step 1: 1-(2,6-Difluorophenyl)ethan-1-ol**

[00863] Prepared according to the procedure described in Example 127, Step 1 using the following starting material: 2',6'-difluoroacetophenone.




15 **Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,6-difluoro-phenyl)-ethyl ester**

[00864] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-(2,6-difluorophenyl)ethan-1-ol.



Step 3: (4'-(4-[1-(2,6-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester


[00865] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-(2,6-difluoro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: (4'-(4-[1-(2,6-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid


[00866] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-(4-[1-(2,6-difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 493.

Example 132: Synthesis of {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-2-yl}-acetic acid (Compound 1-123)



Step 1: {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-2-yl}-acetic acid ethyl ester


[00867] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester and [2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 2: {4'-(3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-2-yl}-acetic acid

[00868] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: {4'-(3-methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-2-yl}-acetic acid ethyl ester. Mass spec. data (M+H) = 457.


Example 133: Synthesis of {4'-(3-Methyl-4-((R)-1-*o*-tolyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid (Compound 1-124)



15


Step 1: (R)-1-*o*-Tolyl-ethanol

[00869] Prepared according to the procedure described in Example 76, Step 1 using the following starting materials: 2'-methylacetophenone and (S)-(-)-2-methyl-CBS-oxazaborolidine.



Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-*o*-tolyl-ethyl ester


[00870] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-*o*-tolyl-ethanol.



5


Step 3: {4'-[3-Methyl-4-((R)-1-*o*-tolyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester

[00871] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-*o*-tolyl-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.

**Step 4: {4'-[3-Methyl-4-((R)-1-*o*-tolyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid**

[00872] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: {4'-[3-methyl-4-((R)-1-*o*-tolyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester. Mass spec. data (M+H) = 471.

Example 134: Synthesis of 4'-(4-Isopropoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid (Compound 1-125)




Step 1: 5-(4'-Ethoxycarbonyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid

[00873] Prepared according to the procedure described in Example 33, Step 1 using the following

starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 4-


5 ethoxycarbonylphenylboronic acid.



Step 2: 4'-(4-Isopropoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid ethyl ester

[00874] Prepared according to the procedure described in Example 29, Step 5 using the following

10 starting materials: 5-(4'-ethoxycarbonyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and isopropanol.




Step 3: 4'-(4-Isopropoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid

[00875] Prepared according to the procedure described in Example 75, Step 3 using the following

15 starting material: 4'-(4-isopropoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid ethyl ester. Mass spec. data (M+H) = 381.


Example 135: Synthesis of (6'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-[2,3']bipyridinyl-5-yl)-acetic acid (Compound 1-126)



Step 1: [5-(5-Bromo-pyridin-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-

5 phenyl)-ethyl ester


[00876] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(5-bromo-pyridin-2-yl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



10 Step 2: (6-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-3-yl)-boronic acid


and {3-Methyl-5-[5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-pyridin-2-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00877] Prepared according to the procedure described in Example 6, Step 2 using the following starting materials: [5-(5-Bromo-pyridin-2-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and bis(pinacolato)diboron; a 3:1 mixture of the boronic acid to the boronate was obtained.



Step 3: (6'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-[2,3'bipyridinyl-5-yl)-acetic acid methyl ester


[00878] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: a 3:1 mixture of (6-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-pyridin-3-yl)-boronic acid and {3-methyl-5-[5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-pyridin-2-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester, and (6-chloro-pyridin-3-yl)-acetic acid methyl ester.



10 **Step 4: (6'-(4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-[2,3'bipyridinyl-5-yl)-acetic acid**


[00879] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (6'-(4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-[2,3'bipyridinyl-5-yl)-acetic acid methyl ester. Mass spec. data (M+H) = 477.

15 **Example 136: Synthesis of 2-(4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid (Diastereomer A) (Compound 1-127)**



Step 1: 2-(4-Bromo-phenyl)-propionic acid ethyl ester


[00880] 2-(4-Bromo-phenyl)-propionic acid ethyl ester (10.0g, 41mmol) was dissolved in THF (400mL) and cooled to -78°C. Sodium bis(trimethylsilyl)amide (2M in THF; 53.5mL, 107mmol) was added, and the mixture was stirred for 1 hour at -78°C. Iodomethane (2.814mL, 45.1mmol) was added, and the reaction was slowly warmed to room temperature and stirred overnight. The mixture was concentrated, and the residue was extracted with EtOAc. The combined organic layers were washed with H₂O and brine, and then dried and concentrated. The crude material was purified by silica gel chromatography (0-15% EtOAc in hexanes) to give the title compound.



10


Step 1: 2-(4-Bromo-phenyl)-propionic acid

[00881] Prepared according to the procedure described in Example 1, Step 7 using the following starting material: 2-(4-bromo-phenyl)-propionic acid ethyl ester.



15 **Step 2: 2-(4-Bromo-phenyl)-propionyl chloride**

[00882] 2-(4-Bromo-phenyl)-propionic acid (0.500g, 2.18mmol) in CH₂Cl₂ (20mL) was treated with DMF (0.03mL, 0.44mmol), followed by oxalyl chloride (0.248mL, 2.8mmol) dropwise over 5 minutes, and the reaction was stirred at room temperature for 1 hour. The mixture was then concentrated and dried under vacuum to give the title compound.




Step 3: (4R,5S)-3-[2-(4-Bromo-phenyl)-propionyl]-4-methyl-5-phenyl-oxazolidin-2-one

[00883] (4R,5S)-(+)-4-Methyl-5-phenyl-2-oxazolidinone (0.348g, 1.96mmol) was dissolved in

THF (20mL) and cooled to -78°C. n-Butyllithium (1.6M in hexanes; 1.64mL, 2.62mmol) was


5 added, and the reaction was stirred for 1 hour at -78°C. 2-(4-Bromo-phenyl)-propionyl chloride (0.540g, 2.18mmol) in THF (10mL) was added, and the reaction was stirred for 30 minutes. The mixture was concentrated, and the residue was partitioned between EtOAc and H₂O. The organic layer was separated, dried, and concentrated, and the crude material was purified by silica gel chromatography (0-25% EtOAc in hexanes) to give two diastereomers as separated

10 products. Diastereomer A was the first product to come off the column, while Diastereomer B was the second product to come off the column.




Step 4: (4R,5S)-4-Methyl-5-phenyl-3-{2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionyl}-oxazolidin-2-one (Diastereomer A)

15 [00884] Prepared according to the procedure described in Example 6, Step 2 using the following starting materials: (4R,5S)-3-[2-(4-bromo-phenyl)-propionyl]-4-methyl-5-phenyl-oxazolidin-2-one (Diastereomer A) and bis(pinacolato)diboron.



Step 5: (3-Methyl-5-{4'-[1-methyl-2-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidin-3-yl)-2-oxo-ethyl]-biphenyl-4-yl}-isoxazol-4-yl)-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Diastereomer A)


5 [00885] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: (4R,5S)-4-methyl-5-phenyl-3-{2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionyl}-oxazolidin-2-one (Diastereomer A) and [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester.



10 **Step 6: 2-(4'-(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl-biphenyl-4-yl)-propionic acid (Diastereomer A)**


[00886] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (3-methyl-5-{4'-[1-methyl-2-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidin-3-yl)-2-oxo-ethyl]-biphenyl-4-yl}-isoxazol-4-yl)-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Diastereomer A). Mass spec. data (M+H) = 505.

Example 137: Synthesis of 2-(4'-(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl-biphenyl-4-yl)-propionic acid (Diastereomer B) (Compound 1-128)




Step 1: (4R,5S)-4-Methyl-5-phenyl-3-{2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionyl}-oxazolidin-2-one (Diastereomer B)

[00887] Prepared according to the procedure described in Example 6, Step 2 using the following starting materials: (4R,5S)-3-[2-(4-bromo-phenyl)-propionyl]-4-methyl-5-phenyl-oxazolidin-2-one (Diastereomer B) and bis(pinacolato)diboron.



Step 2: (3-Methyl-5-{4'-[1-methyl-2-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidin-3-yl)-2-oxo-ethyl]-biphenyl-4-yl}-isoxazol-4-yl)-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Diastereomer B)

[00888] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: (4R,5S)-4-methyl-5-phenyl-3-{2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propionyl}-oxazolidin-2-one (Diastereomer B) and [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester.




Step 3: 2-(4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-propionic acid (Diastereomer B)

[00889] Prepared according to the procedure described in Example 75, Step 2 using the following


5 starting materials: (3-methyl-5-{4'-(1-methyl-2-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidin-3-yl)-2-oxo-ethyl)-biphenyl-4-yl}-isoxazol-4-yl)-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Diastereomer B). Mass spec. data ($M+H$) = 505.

Example 138: Synthesis of (3'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-10 129)



Step 1: 4-Bromo-2-chloro-benzoyl chloride


[00890] Prepared according to the procedure described in Example 29, Step 1 using the following starting materials: 4-bromo-2-chlorobenzoic acid and oxalyl chloride.



15


Step 2: 2-(4-Bromo-2-chloro-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester

[00891] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 4-bromo-2-chloro-benzoyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




Step 3: 5-(4-Bromo-2-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00892] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(4-bromo-2-chloro-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.




Step 4: 5-(4-Bromo-2-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid

[00893] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(4-bromo-2-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




Step 5: [5-(4-Bromo-2-chloro-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester

[00894] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-2-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-chloro-phenyl)-ethanol.



Step 6: (3'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester


[00895] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(4-bromo-2-chloro-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: (3'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid

[00896] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (3'-chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data ($M+H$) = 525.

Example 139: Synthesis of 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid (Compound 1-130)




Step 1: 2-(4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-butyric acid

[00897] Prepared according to the procedure described in Example 33, Step 1 using the following


5 starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and 2-(4-bromophenyl)butanoic acid. Mass spec. data ($M+H$) = 519.

Example 140: Synthesis of (2'-Chloro-4'-(4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-yl)-acetic acid (Compound 1-131)



Step 1: 4-Bromo-3-chloro-benzoyl chloride


[00898] Prepared according to the procedure described in Example 29, Step 1 using the following starting materials: 4-bromo-3-chlorobenzoic acid and oxalyl chloride.



15

Step 2: 2-(4-Bromo-3-chloro-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester

[00899] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 4-bromo-3-chloro-benzoyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




Step 3: 5-(4-Bromo-3-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

[00900] Prepared according to the procedure described in Example 29, Step 3 using the following

starting materials: 2-(4-bromo-3-chloro-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester

5 and hydroxylamine hydrochloride.




Step 4: 5-(4-Bromo-3-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid

[00901] Prepared according to the procedure described in Example 29, Step 4 using the following

starting material: 5-(4-bromo-3-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl


10 ester.



Step 5: [5-(4-Bromo-3-chloro-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester


[00902] Prepared according to the procedure described in Example 29, Step 5 using the following

15 starting material: 5-(4-bromo-3-chloro-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-chloro-phenyl)-ethanol.



Step 6: (2'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester


[00903] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(4-bromo-3-chloro-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: (2'-Chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid


[00904] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (2'-chloro-4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 526.

Example 141: Synthesis of (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-fluoro-biphenyl-4-yl)-acetic acid (Compound 1-132)




Step 1: 4-Bromo-3-fluoro-benzoyl chloride

[00905] Prepared according to the procedure described in Example 29, Step 1 using the following starting materials: 4-bromo-3-fluorobenzoyl chloride and oxalyl chloride.




Step 2: 2-(4-Bromo-3-fluoro-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester

5 [00906] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 4-bromo-3-fluoro-benzoyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




Step 3: 5-(4-Bromo-3-fluoro-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

10 [00907] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(4-bromo-3-fluoro-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.




Step 4: 5-(4-Bromo-3-fluoro-phenyl)-3-methyl-isoxazole-4-carboxylic acid

15 [00908] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(4-bromo-3-fluoro-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




Step 5: [5-(4-Bromo-3-fluoro-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester

[00909] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-3-fluoro-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-chloro-phenyl)-ethanol.



Step 6: (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-fluoro-biphenyl-4-yl)-acetic acid ethyl ester


[00910] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(4-bromo-3-fluoro-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 7: (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-fluoro-biphenyl-4-yl)-acetic acid


[00911] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: (4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-fluoro-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 509.

Example 142: Synthesis of 4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid (Compound 1-133)



Step 1: 4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid ethyl ester


[00912] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-chlorophenyl)-ethyl ester and 4-ethoxycarbonylphenylboronic acid.



Step 2: 4'-(4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid

[00913] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 4'-(4-[(R)-1-(2-chlorophenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid ethyl ester. Mass spec. data (M+H) = 477.


Example 143: Synthesis of [5-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid methyl ester (Compound 1-134)



15


Step 1: 2-(4-Cyano-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester

[00914] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 4-cyanobenzoyl chloride and 3-methylamino-but-2-enoic acid methyl ester.




Step 2: 5-(4-Cyano-phenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester

5 [00915] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(4-cyano-benzoyl)-3-[(E)-methylimino]-butyric acid methyl ester and hydroxylamine hydrochloride.




Step 3: 5-(4-Cyano-phenyl)-3-methyl-isoxazole-4-carboxylic acid

10 [00916] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(4-cyanophenyl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.



Step 4: [5-(4-Cyano-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester


15 [00917] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-cyanophenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(2-fluoro-phenyl)-ethanol.



Step 5: {3-Methyl-5-[4-(2H-tetrazol-5-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00918] [5-(4-Cyano-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-

5 ethyl ester (0.310g, 0.85mmol), trimethylsilyl azide (0.146mL, 1.10mmol), and dibutyltin(IV) oxide (0.021g, 0.085mmol) were combined in toluene, and the reaction was subjected to microwave at 120°C for 5 minutes. The mixture was partitioned between EtOAc and H₂O, and the organic layer was extracted 3 times with saturated aqueous NaHCO₃. The aqueous layer was acidified with 1N aqueous HCl and extracted twice with EtOAc. The combined organic layers 10 were dried over MgSO₄, filtered, and concentrated. The crude material was resubjected to the same reaction conditions, treated with trimethylsilyl azide and dibutyltin(IV) oxide in toluene at 120°C and microwaved, and the mixture was worked-up in the same manner to give the title compound.



15 **Step 6: [5-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid methyl ester**

[00919] {3-Methyl-5-[4-(2H-tetrazol-5-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-

phenyl)-ethyl ester (0.055g, 0.135mmol), methyl bromoacetate (0.015mL, 0.162mmol), and


potassium carbonate (0.037g, 0.27mmol) were combined in MeCN and stirred overnight at room

20 temperature. The mixture was concentrated, and the residue was diluted with CH₂Cl₂ and

filtered. The crude material was purified by silica gel chromatography (0-100% EtOAc in

hexanes) to give the title compound. Mass spec. data (M+H) = 480.


Example 144: Synthesis of [5-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid (Compound 1-135)



Step 1: [5-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid


5 [00920] Prepared according to the procedure described in Example 75, Step 2 using the following starting materials: [5-(4-{(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid methyl ester. Mass spec. data ($M+H$) = 467.

10 **Example 147: Synthesis of [4'-(4-*tert*-Butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid (Compound 1-136)**



Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid *tert*-butyl ester


15 [00921] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and *tert*-butanol.



15

Step 2: [4'-(4-*tert*-Butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid ethyl ester


[00922] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid *tert*-butyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



5 **Step 3: [4'-(4-*tert*-Butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid**


[00923] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: [4'-(4-*tert*-butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid ethyl ester. Mass spec. data ($M+H$) = 409.

10 **Example 148: Synthesis of [3-(4-{4-[*(R*)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-137)**



Step 1: [3-(4-{4-[*(R*)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester


[00924] Prepared according to the procedure described in Example 81, Step 4 using the following starting materials: [5-(4-hydroxy-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (*R*)-1-(2-chlorophenyl)-ethyl ester and 3-(2-methoxy-2-oxoethyl)phenylboronic acid.



Step 2: [3-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid

[00925] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: [3-(4-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester. Mass spec. data (M+H) = 507.


Example 149: Synthesis of (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-hydroxy-acetic acid (Compound 1-139)



Step 1: (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-hydroxy-acetic acid


[00926] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: {3-methyl-5-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 4-bromomandelic acid. Mass spec. data (M+H) = 491.

Example 150: Synthesis of 1-{4-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-benzyl}-azetidine-3-carboxylic acid (Compound 1-140)




Step 1: [5-(4-Chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester

[00927] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-chloromethyl-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-phenyl-ethanol.



Step 2: 1-[4-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-benzyl]-azetidine-3-carboxylic acid methyl ester


[00928] To [5-(4-chloromethyl-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester (0.060g, 0.16mmol) in MeCN (2mL) was added diisopropylethylamine (0.07mL, 0.41mmol), followed by methyl azetidine-3-carboxylate hydrochloride (0.030g, 0.19mmol), and the reaction was stirred at room temperature overnight. Analytical LCMS indicated that no starting material remained, and the solution was used directly in the hydrolysis step.



10 **Step 3: 1-[4-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-benzyl]-azetidine-3-carboxylic acid**

[00929] A solution of 1-[4-[3-methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-benzyl]-azetidine-3-carboxylic acid methyl ester (0.16mmol) in MeCN (2mL) was treated with 3N aqueous NaOH (1mL) and stirred at room temperature for 5 minutes. The mixture was 1N aqueous NaOH and washed with Et₂O. The aqueous layer was acidified with concentrated HCl and extracted with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and concentrated to give the title compound. Mass spec. data (M+H) = 436.

Example 151: Synthesis of (4'-{4-[(R)-1-(3,5-Dibromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-141)




20

Step 1: 1-(3,5-Dibromo-phenyl)-ethanone


[00930] To a solution of 3,5-dibromobenzoic acid (2.5g, 8.9mmol) in Et₂O (30mL) at 0°C was added methyl lithium (1.6M in Et₂O; 12.3mL, 19.6mmol) was added dropwise. The reaction was warmed to room temperature and stirred for 2 hours. Acidic work-up, followed by silica gel chromatography, gave the title compound.

5



Step 2: (R)-1-(3,5-Dibromo-phenyl)-ethanol


[00931] Prepared according to the procedure described in Example 76, Step 1 using the following starting materials: 1-(3,5-dibromo-phenyl)-ethanone and (S)-(-)-2-methyl-CBS-oxazaborolidine.



10 **Step 3: (4'-{4-[(R)-1-(3,5-Dibromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester**

[00932] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4'-ethoxycarbonylmethyl-biphenyl-4-yl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-(3,5-dibromo-phenyl)-ethanol.


15



Step 4: (4'-{4-[(R)-1-(3,5-Dibromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid


[00933] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-{4-[(R)-1-(3,5-dibromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 615.

5 **Example 152: Synthesis of 2-(4-{5-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-pyridin-2-yl}-phenyl)-butyric acid (Compound 1-142)**



Step 1: 2-(6-Chloro-pyridine-3-carbonyl)-3-oxo-butyric acid methyl ester


[00934] Prepared according to the procedure described in Example 29, Step 2 using the following starting materials: 6-chloronicotinoyl chloride and 3-methylamino-but-2-enoic acid methyl ester.



10

Step 2: 5-(6-Chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester


[00935] Prepared according to the procedure described in Example 29, Step 3 using the following starting materials: 2-(6-chloro-pyridine-3-carbonyl)-3-oxo-butyric acid methyl ester and hydroxylamine hydrochloride.



15


Step 3: 5-(6-Chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid

[00936] Prepared according to the procedure described in Example 29, Step 4 using the following starting material: 5-(6-chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid methyl ester.




Step 4: [5-(6-Chloro-pyridin-3-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-phenyl-ethyl ester

[00937] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(6-chloro-pyridin-3-yl)-3-methyl-isoxazole-4-carboxylic acid and (R)-1-phenyl-ethanol.



Step 5: 2-(4-Bromo-phenyl)-butyric acid ethyl ester


[00938] Prepared according to the procedure described in Example 21, Step 1 using the following starting materials: 2-(4-bromophenyl)butanoic acid and ethanol.



10

Step 6: 2-[4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-butyric acid ethyl ester


[00939] Prepared according to the procedure described in Example 49, Step 1 using the following starting materials: 2-(4-bromo-phenyl)-butyric acid ethyl ester and bis(pinacolato)diboron.



15 **Step 7: 2-(4-{5-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-pyridin-2-yl}-phenyl)-butyric acid ethyl ester**

[00940] Prepared according to the procedure described in Example 29, Step 6 using the following starting materials: [5-(6-chloro-pyridin-3-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-


phenyl-ethyl ester and 2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-butyric acid ethyl ester.



Step 8: 2-(4-{5-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-pyridin-2-yl}-phenyl)-butyric acid


[00941] Prepared according to the procedure described in Example 29, Step 7 using the following starting material: 2-(4-{5-[3-methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-pyridin-2-yl}-phenyl)-butyric acid ethyl ester. Mass spec. data (M+H) = 486.

Example 153: Synthesis of {4'-[3-Methyl-4-((S)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-143)




Step 1: (S)-1-Phenyl-ethanol

[00942] Prepared according to the procedure described in Example 76, Step 1 using the following starting materials: acetophenone and (R)-(+)-2-methyl-CBS-oxazaborolidine.




Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (S)-1-phenyl-ethyl ester

[00943] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and (S)-1-phenyl-ethanol.



Step 3: {4'-(3-Methyl-4-((S)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid ethyl ester


[00944] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (S)-1-phenyl-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 4: {4'-(3-Methyl-4-((S)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid

[00945] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: {4'-(3-methyl-4-((S)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid ethyl ester. Mass spec. data (M+H) = 457.


Example 154: Synthesis of (4'-{4-[(R)-1-(3-Hydroxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-144)



15


Step 1: 3-((R)-1-Hydroxy-ethyl)-phenol

[00946] Prepared according to the procedure described in Example 76, Step 1 using the following starting materials: 3'-hydroxyacetophenone and (S)-(-)-2-methyl-CBS-oxazaborolidine.




Step 2: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(3-hydroxy-phenyl)-ethyl ester

[00947] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 3-((R)-1-hydroxy-ethyl)-phenol.



Step 3: (4'-{4-[(R)-1-(3-Hydroxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester


[00948] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(3-hydroxy-phenyl)-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



[00949] **Step 4: (4'-{4-[(R)-1-(3-Hydroxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid**

[00949] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: (4'-{4-[{(R)-1-(3-hydroxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid ethyl ester. Mass spec. data (M+H) = 473.


5 **Example 155: Synthesis of {4'-[3-Methyl-4-(1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-145)**



Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-phenyl-ethyl ester


[00950] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-

10 phenylethanol.



Step 2: {4'-[3-Methyl-4-(1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester


[00951] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-phenyl-ethyl ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 3: {4'-[3-Methyl-4-(1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid


[00952] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: {4'-[3-Methyl-4-(1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester. Mass spec. data ($M+H$) = 457.

Example 156: Synthesis of {4'-[3-Methyl-4-(1-phenyl-ethoxy-d9-carbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-146)




Step 1: [5-(4-Bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-phenyl-ethyl-d9 ester

[00953] Prepared according to the procedure described in Example 29, Step 5 using the following starting materials: 5-(4-bromo-phenyl)-3-methyl-isoxazole-4-carboxylic acid and 1-phenylethanol-d9 (obtained from Carbocore).



Step 2: {4'-[3-Methyl-4-(1-phenyl-ethoxy-d9-carbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid ethyl ester


[00954] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid 1-phenyl-ethyl-d9 ester and [4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl.



Step 3: {4'-(3-Methyl-4-(1-phenyl-ethoxy-d9-carbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid


[00955] Prepared according to the procedure described in Example 75, Step 2 using the following starting material: {4'-(3-Methyl-4-(1-phenyl-ethoxy-d9-carbonylamino)-isoxazol-5-yl)-biphenyl-4-yl}-acetic acid ethyl ester. Mass spec. data ($M+H$) = 466.

Example 157: Synthesis of [5-(4'-Carbamimidoylmethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-6)



10 **Step 1: [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester**

[00956] Prepared according to the procedure described in Example 33, Step 1 using the following starting materials: [5-(4-bromo-phenyl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and 4-cyanomethylphenylboronic acid.




Step 2: [5-(4'-Carbamimidoylmethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00957] [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-

5 fluoro-phenyl)-ethyl ester (0.400g, 0.88mmol) in EtOH (5mL) was treated with 4N HCl in 1,4-dioxane (5mL), and the reaction was stirred overnight at room temperature. The mixture was concentrated to dryness, and then dissolved in 2M NH₃ in MeOH. The reaction was stirred overnight at room temperature, and then additional 2M NH₃ in MeOH was added and the reaction was stirred at 40°C for 1.5 hours. The mixture was concentrated, and the residue was

10 purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 473.

Example 158: Synthesis of {5-[4'-(2-Acetyl-amino-2-imino-ethyl)-biphenyl-4-yl]-3-methyl-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-7)



Step 1: {5-[4'-(2-Acetyl-amino-2-imino-ethyl)-biphenyl-4-yl]-3-methyl-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester

[00958] To [5-(4'-carbamimidoylmethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid

(R)-1-(2-fluoro-phenyl)-ethyl ester (0.055g, 0.12mmol) in CH₂Cl₂ (5mL) was added


diisopropylethylamine (0.052mL, 0.3mmol), followed by acetyl chloride (0.009mL, 0.126mmol),

and the reaction was stirred at room temperature. Additional acetyl chloride (0.009mL,

20 0.126mmol) was added, and the reaction was stirred at room temperature. The mixture was


quenched with H₂O and extracted with CH₂Cl₂. The combined organic layers were dried, filtered, and concentrated, and the residue was purified by preparative HPLC to give the title compound. Mass spec. data (M+H) = 515.

5 **Example 159: Synthesis of [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-5)**



Step 1: [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid ethyl ester


[00959] Prepared according to the procedure described in Example 91, Step 1 using the following starting materials: [5-(2-bromo-thiazol-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and [2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



Step 2: [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid


[00960] Prepared according to the procedure described in Example 91, Step 2 using the following starting material: [2-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid ethyl ester. Mass spec. data (M+H) = 482.

20 **Example 160: Synthesis of [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-6)**



Step 1: [3-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-thiazol-2-yl]-acetic acid ethyl ester


[00961] Prepared according to the procedure described in Example 91, Step 1 using the following starting materials: [5-(2-bromo-thiazol-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester and [3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetic acid ethyl ester.



10 **Step 2: [3-(4-{(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-thiazol-2-yl]-acetic acid**


[00962] Prepared according to the procedure described in Example 91, Step 2 using the following starting material: [3-(4-{(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino}-3-methyl-isoxazol-5-yl)-thiazol-2-yl]-acetic acid ethyl ester. Mass spec. data (M+H) = 482.

Example 161: Synthesis of [4'-(4-Amino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid



[00963] [4'-(4-*tert*-Butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid (1.0mmol) was treated with trifluoroacetic acid (5mL) for 1 hour. Work-up provided the title compound. Mass spec. data (M+H) = 309.

Example 162: Synthesis of 2-(2-{4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetylamino)-ethanesulfonic acid (Compound 2-8)



[00964] Rats were dosed with {4'-[3-ethyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (30 mg/kg) and the bile was collected over 6 hours. The title compound was purified from bile by reverse phase HPLC. Mass spec. data (M+H) = 564.

10 [00965] In some embodiments, Mass spectrometric data (mass spec. data) is obtained on with a Shimadzu LCMS 2010A.

***In vitro* Assay(s)**

Example 163. Establishment of a CHO Cell Line Stably Expressing Human LPA₁

[00966] A 1.1 kb cDNA encoding the human LPA₁ receptor was cloned from human lung. Human 15 lung RNA (Clontech Laboratories, Inc. USA) was reverse transcribed using the RETROscript kit (Ambion, Inc.) and the full-length cDNA for human LPA₁ was obtained by PCR of the reverse transcription reaction. The nucleotide sequence of the cloned human LPA₁ was determined by sequencing and confirmed to be identical to the published human LPA₁ sequence (An *et al.* *Biochem. Biophys. Res. Commun.* 231:619 (1997)). The cDNA was cloned into the pCDNA5/FRT expression plasmid and transfected in CHO cells using lipofectamine 2000 (Invitrogen Corp., USA). Clones stably expressing human LPA₁ were selected using hygromycin and identified as cells that show Ca-influx in response to LPA.

Example 164. Generation of Cells Transiently Expressing Human LPA₂

[00967] A vector containing the human LPA₂ receptor cDNA was obtained from the Missouri 25 S&T cDNA Resource Center (www.cdna.org). The full-length cDNA fragment for human LPA₂ was obtained by PCR from the vector. The nucleotide sequence of the cloned human LPA₂ was determined by sequencing and confirmed to be identical to the published human LPA₂ sequence (NCBI accession number NM_004720). The cDNA was cloned into the pCDNA3.1 expression

plasmid and transfected into B103 cells (Invitrogen Corp., USA) by seeding cells in a 96-well poly-D-lysine coated plate at 30,000-35,000 cells per well together with 0.2 μ l lipofectamine 2000 and 0.2 μ g of the LPA₂ expression vector. Cells were cultured overnight in complete media before being assayed for LPA-induced Ca-influx.

5 **Example 165. Establishment of a CHO Cell Line Stably Expressing Human LPA₃**

[00968] A vector containing the human LPA₃ receptor cDNA was obtained from the Missouri S&T cDNA Resource Center (www.cdna.org). The full-length cDNA fragment for human LPA₃ was obtained by PCR from the vector. The nucleotide sequence of the cloned human LPA₃ was determined by sequencing and confirmed to be identical to the published human LPA₃ sequence (NCBI accession number NM_012152). The cDNA was cloned into the pCDNA5/FRT expression plasmid and transfected in CHO cells using lipofectamine 2000 (Invitrogen Corp., USA). Clones stably expressing human LPA₃ were selected using hygromycin and identified as cells that show Ca-influx in response to LPA.

Example 166. LPA₁ and LPA₃ Calcium Flux Assays.

[00969] Human LPA₁ or LPA₃ expressing CHO cells are seeded at 20,000-45,000 cells per well in a 96-well poly-D-lysine coated plate one or two days before the assay. Prior to the assay, the cells are washed once with PBS and then cultured in serum-free media overnight. On the day of the assay, a calcium indicator dye (Calcium 4, Molecular Devices) in assay buffer (HBSS with Ca²⁺ and Mg²⁺ and containing 20 mM Hepes and 0.3% fatty-acid free human serum albumin) is added to each well and incubation continued for 1 hour at 37°C. 10 μ l of test compounds in 2.5% DMSO are added to the cells and incubation continued at room temperature for 30 minutes. Cells are stimulated by the addition of 10 nM LPA and intracellular Ca²⁺ measured using the Flexstation 3 (Molecular Devices). IC₅₀s are determined using Graphpad prism analysis of drug titration curves.

25 **Example 167. LPA₂ Calcium Flux Assay.**

[00970] Following an overnight culture with lipofectamine 2000 and the LPA₂ expression vector, the B103 cells are washed once with PBS then serum starved for 4 hours. A calcium indicator dye (Calcium 4, Molecular Devices) in assay buffer (HBSS with Ca²⁺ and Mg²⁺ and containing 20 mM Hepes and 0.3% fatty-acid free human serum albumin) is added to each well and incubation continued for 1 hour at 37°C. 10 μ l of test compounds in 2.5% DMSO are added to the cells and incubation continued at room temperature for 30 minutes. Cells are stimulated by the addition of 10 nM LPA and intracellular Ca²⁺ measured using the Flexstation 3 (Molecular Devices). IC₅₀s are determined using Graphpad prism analysis of drug titration curves.

Example 168. GTP γ S Binding Assay

[00971] The ability of a compound to inhibit binding of GTP to LPA₁ is assessed via a membrane GTP γ S assay. CHO cells stably expressing the recombinant human LPA₁ receptor are

resuspended in 10 mM Hepes, 7.4 containing 1 mM DTT, lysed and centrifuged at 75,000 xg to pellet the membranes. The membranes are resuspended in 10 mM Hepes, 7.4 containing 1 mM

5 DTT and 10% glycerol. Membranes (~25 μ g per well) are incubated in 96-well plates with 0.1 nM [³⁵S]-GTP γ S, 900 nM LPA, 5 μ M GDP, and test compound in Assay Buffer (50 mM Hepes, pH 7.4, 100 mM NaCl, 10 mM MgCl₂, 50 μ g/ml saponin and 0.2% fatty-acid free human serum albumin) for 30 minutes at 30°C. The reactions are terminated by rapid filtration through

Whatman GF/B glass fibre filter plates. The filter plates are washed 3 times with 1 ml cold Wash

10 Buffer (50 mM Hepes, 7.5, 100 mM NaCl and 10 mM MgCl₂) and dried. Scintillant is then added to the plates and the radioactivity retained on the filters is determined on a Packard

TopCount (Perkin Elmer). Specific binding is determined as total radioactive binding minus non-specific binding in the absence of the ligand (900 nM LPA). IC₅₀s were determined using Graphpad prism analysis of drug titration curves.

15 [00972] Illustrative *in vitro* biological data for representative compounds of Formula (I), Formula (II) and Formula (III) is presented in the Table below. Unless otherwise noted, compounds that were tested had an IC₅₀ of less than 50 μ M in the HLPA1 Ca Flux assay.

Compound Number	HLPA1 Ca Flux IC ₅₀ (μ M)	HLPA3 Ca Flux IC ₅₀ (μ M)
Compound 1-1	A	C
Compound 1-2	ND	ND
Compound 1-3	A	C
Compound 1-4	A	C
Compound 1-5	A	C
Compound 1-6	C	ND
Compound 1-7	B	C
Compound 1-9	A	C
Compound 1-11	C	ND
Compound 1-12	A	C
Compound 1-13	C	ND
Compound 1-14	B	ND
Compound 1-15	A	C
Compound 1-16	D	ND
Compound 1-17	A	C
Compound 1-18	A	ND
Compound 1-19	A	C
Compound 1-20	A	C
Compound 1-21	A	C
Compound 1-22	A	ND
Compound 1-23	A	C
Compound 1-24	B	C
Compound 1-25	A	C
Compound 1-26	A	C
Compound 1-27	A	C
Compound 1-28	A	D

Compound 1-29	B	ND
Compound 1-30	B	C
Compound 1-31	C	C
Compound 1-32	B	ND
Compound 1-34	B	ND
Compound 1-35	B	ND
Compound 1-36	B	ND
Compound 1-37	C	ND
Compound 1-38	C	ND
Compound 1-39	C	ND
Compound 1-40	D	ND
Compound 1-41	C	ND
Compound 1-42	C	ND
Compound 1-43	C	ND
Compound 1-44	C	ND
Compound 1-45	A	D
Compound 1-46	A	ND
Compound 1-47	B	ND
Compound 1-48	A	D
Compound 1-49	B	C
Compound 1-50	A	C
Compound 1-51	A	C
Compound 1-52	C	ND
Compound 1-53	A	C
Compound 1-54	A	C
Compound 1-55	A	ND
Compound 1-56	A	C
Compound 1-57	A	C
Compound 1-58	A	C
Compound 1-59	A	C
Compound 1-60	C	C
Compound 1-61	A	C
Compound 1-62	A	C
Compound 1-63	A	C
Compound 1-64	A	C
Compound 1-65	A	C
Compound 1-66	A	C
Compound 1-67	A	C
Compound 1-69	A	C
Compound 1-70	A	ND
Compound 1-71	A	C
Compound 1-72	A	ND
Compound 1-73	A	ND
Compound 1-74	A	C
Compound 1-75	A	ND
Compound 1-76	A	C
Compound 1-77	A	C
Compound 1-78	A	C
Compound 1-79	A	C
Compound 1-80	A	C
Compound 1-81	A	C
Compound 1-82	B	ND
Compound 1-83	B	ND
Compound 1-84	A	C
Compound 1-85	A	C

Compound 1-86	A	C
Compound 1-87	A	C
Compound 1-88	A	C
Compound 1-89	A	B
Compound 1-90	A	C
Compound 1-91	A	C
Compound 1-92	A	B
Compound 1-93	A	C
Compound 1-94	A	C
Compound 1-95	A	C
Compound 1-96	A	C
Compound 1-97	A	C
Compound 1-98	A	D
Compound 1-99	A	C
Compound 1-100	A	C
Compound 1-101	C	ND
Compound 1-102	A	C
Compound 1-103	A	C
Compound 1-104	C	ND
Compound 1-105	C	ND
Compound 1-106	B	ND
Compound 1-107	C	ND
Compound 1-108	A	ND
Compound 1-109	A	C
Compound 1-110	A	
Compound 1-111	A	C
Compound 1-112	B	C
Compound 1-113	B	ND
Compound 1-114	A	C
Compound 1-115	A	C
Compound 1-116	A	C
Compound 1-117	A	C
Compound 1-118	B	ND
Compound 1-119	A	C
Compound 1-120	A	D
Compound 1-121	A	D
Compound 1-122	A	C
Compound 1-123	C	ND
Compound 1-124	A	C
Compound 1-125	C	ND
Compound 1-126	A	C
Compound 1-127	A	B
Compound 1-128	A	C
Compound 1-129	A	C
Compound 1-130	A	B
Compound 1-131	A	C
Compound 1-132	A	C
Compound 1-133	A	B
Compound 1-134	C	ND
Compound 1-135	C	ND
Compound 1-136	B	ND
Compound 1-137	A	C
Compound 1-138	A	B
Compound 1-139	A	C
Compound 1-140	C	ND

Compound 1-141	C	ND
Compound 1-142	A	C
Compound 1-143	A	C
Compound 1-144	C	D
Compound 1-145	ND	ND
Compound 1-146	ND	ND
Compound 2-1	A	ND
Compound 2-2	A	C
Compound 2-3	A	D
Compound 2-4	A	C
Compound 2-5	A	C
Compound 2-6	C	ND
Compound 2-7	A	ND
Compound 2-7	A	ND
Compound 3-1	B	ND
Compound 3-2	B	ND
Compound 3-3	C	ND
Compound 3-4	A	D
Compound 3-5	C	ND
Compound 3-6	B	ND

A = less than 0.3 μ M; B = greater than 0.3 μ M and less than 1 μ M;

C= greater than 1 μ M and less than 50 μ M; D= greater than 50 μ M.

ND = not determined

Example 169. LPA1 Chemotaxis Assay.

5 [00973] Chemotaxis of the A2058 human melanoma cells was measured using the Neuroprobe ChemoTx® System plates (8 μ m pore size, 5.7 mm diameter sites). The filter sites were coated with 0.001% fibronectin (Sigma) in 20 mM Hepes, pH 7.4 and allowed to dry. A2058 cells were serum-starved for 24 hours , then harvested with Cell Stripper and resuspended in DMEM containing 0.1% fatty-acid-free bovine serum albumin (BSA) to a concentration of 1 x 10⁶/ml.

10 Cells were mixed with an equal volume of test compound (2X) in DMEM containing 0.1% fatty-acid-free BSA and incubated at 37°C for 15 minutes. LPA (100 nM in DMEM containing 0.1% fatty-acid-free BSA) or vehicle was added to each well of the lower chamber and 50 μ l of the cell suspension/test compound mix was applied to the upper portion of the ChemoTx plate. Plates were incubated at 37°C for three hours and then the cells removed from the upper portion by

15 rinsing with PBS and scraping. The filter was dried then stained with HEMA 3 Staining System (Fisher Scientific). The absorbance of the filter was read at 590 nM and IC₅₀s were determined using Symyx Assay Explorer.

[00974] Compound 1-88 and Compound 1-115 inhibited LPA-driven chemotaxis (IC₅₀ less than 300 nM) of human A2058 melanoma cells.

20 In vivo Assay(s)

Example 170: Bleomycin-induced Lung fibrosis model in mice

[00975] Female CD-1 mice (Harlan, 25-30g) are housed 4 per cage, given free access to food and water and allowed to acclimate for at least 7 days prior to test initiation. After the habituation

phase, mice are lightly anesthetized with isoflurane (5% in 100% O₂) and administered with bleomycin sulfate (0.01-5 U/kg, Henry Schein) via intratracheal instillation (Cuzzocrea S *et al.* *Am J Physiol Lung Cell Mol Physiol.* 2007 May;292(5):L1095-104. Epub 2007 Jan 12.). Mice are returned to their cages and monitored daily for the duration of the experiment. Test

5 compound or vehicle is delivered po, ip or sc daily. The route and frequency of dosing is based on previously determined pharmacokinetic properties. All animals are sacrificed using inhaled isoflurane 3, 7, 14, 21 or 28 days after bleomycin instillation. Following sacrifice, mice are intubated with a 20 gauge angiocatheter attached to a 1 ml syringe. Lungs are lavaged with saline to obtain bronchoalveolar lavage fluid (BALF) and then removed and fixed in 10% neutral 10 buffered formalin for subsequent histopathological analysis. BALF is centrifuged for 10 min at 800 x g to pellet the cells and the cell supernatant removed and frozen at -80 °C for subsequent protein analysis using the DC protein assay kit (Biorad, Hercules, CA.) and soluble collagen analysis using Sircol (Biocolor Ltd, UK). BALF is analyzed for concentrations of inflammatory, 15 pro-fibrotic and tissue injury biomarkers including transforming growth factor β1, hyaluronic acid, tissue inhibitor of metalloproteinase-1, matrix metalloproteinase-7, connective tissue growth factor and lactate dehydrogenase activity, using commercially available ELISA. The cell pellet is re-suspended in PBS. Total cell counts are then obtained using a Hemavet hematology system (Drew Scientific, Wayne, PA.) and differential cells counts are determined using Shandon cytocentrifuge (Thermo Scientific, Waltham, MA.). Lung tissue is stained using hematoxylin and eosin (H&E) and trichrome and lung fibrosis is determined by semiquantitative histopathological 20 scoring (Ashcroft T. *et al. J. Clin. Path.* 1988;41;4, 467-470.) using light microscopy (10x magnification) and quantitative, computer-assisted densitometry of collagen in lung tissue sections using light microscopy. The data are plotted using Graphpad prism and statistical differences between groups determined.

25 [00976] In the acute setting (3 day), Compound 1-88 significantly reduced total protein, lactate dehydrogenase activity (LDH; tissue injury marker) and tissue inhibitor of metalloproteinase-1 (TIMP-1; pro-fibrotic marker) concentrations in bronchoalveolar lavage fluid (BALF). In the chronic setting (14 and 28 day) model, Compound 1-88 maintained mouse body weight and decreased inflammatory cell influx and fibrosis after a single bleomycin (1.5 units/kg) instillation 30 and decreased pulmonary resistance and lung fibrosis following repeated bleomycin (3.0-5.0 units/kg/week) instillations.

[00977] Compound 1-115 reduced total protein, lactate and TIMP-1 in the BALF in the acute setting (3-day). Compound 1-115 decreased inflammatory cell influx and fibrosis after a single bleomycin instillation (3.0 units) in the chronic setting (14-days only).

35 **Example 171: Mouse carbon tetrachloride (CCl₄)-induced liver fibrosis model**

[00978] Female C57BL/6 mice (Harlan, 20-25g) housed 4/cage are given free access to food and water and allowed to acclimate for at least 7 days prior to test initiation. After the habituation phase, mice receive CCl₄ (0.5-1.0 ml/kg body weight) diluted in corn oil vehicle (100 µL volume) via i.p. injection twice a week for 4-6 weeks. (Higazi, A. A. *et al.*, *Clin Exp Immunol.*

5 2008 Apr;152(1):163-73. Epub 2008 Feb 14.). Control mice receive an equivalent volume of corn oil vehicle only. Test compound or vehicle is delivered po, ip or sc daily. At the end of the study (4-6 weeks after first i.p. injection of CCl₄), mice are sacrificed using inhaled isoflurane and blood is drawn via cardiac puncture for subsequent analysis of ALT/AST levels. The liver is harvested, and one half of the liver is frozen at -80°C and the other half is fixed in 10% neutral
10 buffered formalin for histological assessment of liver fibrosis using light microscopy (10x magnification). Liver tissue homogenates are analyzed for collagen levels using Sircol (Bicolor Ltd, UK). Fixed Liver tissue is stained using hematoxylin and eosin (H&E) and trichrome and liver fibrosis is determined by quantitative, computer-assisted densitometry of collagen in liver tissue sections using light microscopy. Plasma and liver tissue lysates are also analyzed for
15 concentrations of inflammatory, pro-fibrotic and tissue injury biomarkers including transforming growth factor β1, hyaluronic acid, tissue inhibitor of metalloproteinase-1, matrix metalloproteinase-7, connective tissue growth factor and lactate dehydrogenase activity, using commercially available ELISA. The resulting data are plotted using Graphpad prism and statistical differences between groups determined.

20 [00979] In this experiment, Compound 1-88 significantly reduced collagen deposition in the liver as compared to the untreated group. Compound 1-88 (30 mg/kg, po, qd) had the same effect on collagen deposition as pirfenidone. Compound 1-115 significantly reduced collagen deposition in the liver as compared to untreated control group.

Example 172: Mouse intravenous LPA-induced histamine release

25 [00980] A mouse intravenous LPA-induced histamine release model is utilized to determine the in vivo potency of LPA₁ and LPA₃ receptor antagonists. Female CD-1 mice (weighing 25 – 35 grams) are administered compound (i.p., s.c. or p.o.) in a volume of 10ml/kg 30 minutes to 24 hours prior to intravenous LPA challenge (300 µg/mouse in 0.1% FAF BSA). Immediately following LPA challenge mice are placed into an enclosed Plexiglas chamber and exposed to an
30 isoflurane for a period of 2 minutes. They are removed, decapitated and trunk blood collected into tubes containing EDTA. Blood is then centrifuged at 10,000 X g for 10 minutes at 4°C. Histamine concentrations in the plasma are determined by EIA. Drug concentrations in plasma are determined by mass spectrometry. The dose to achieve 50% inhibition of blood histamine release is calculated by nonlinear regression (Graphpad Prism) and plotted as the ED₅₀. The
35 plasma concentration associated with this dose is plotted as the EC₅₀.

Example 173: Mouse unilateral ureteral obstruction kidney fibrosis model

[00981] Female C57BL/6 mice (Harlan, 20-25g) housed 4/cage will be given free access to food and water and allowed to acclimate for at least 7 days prior to test initiation. After the habituation phase, mice undergo unilateral ureteral obstruction (UUO) surgery or sham to left kidney.

5 Briefly, a longitudinal, upper left incision is performed to expose the left kidney. The renal artery is located and 6/0 silk thread is passed between the artery and the ureter. The thread is looped around the ureter and knotted 3 times insuring full ligation of ureter. The kidney is returned to abdomen, the abdominal muscle is sutured and the skin is stapled closed. Mice are returned to their cages and monitored daily for the duration of the experiment. Test compound or
10 vehicle is delivered po, ip or sc daily. The route and frequency of dosing is based on previously determined pharmacokinetic properties. All animals are sacrificed using inhaled isoflurane 4, 8 or 14 days after UUO surgery. Following sacrifice blood is drawn via cardiac puncture, the kidneys are harvested and one half of the kidney is frozen at -80°C and the other half is fixed in 10% neutral buffered formalin for histological assessment of kidney fibrosis using light microscopy
15 (10x magnification). Kidney tissue homogenates are analyzed for collagen levels using Sircol (Biocolor Ltd, UK). Fixed kidney tissue is also stained using hematoxylin and eosin (H&E) and trichrome and kidney fibrosis is determined by quantitative, computer-assisted densitometry of collagen in liver tissue sections using light microscopy. Plasma and kidney tissue lysates are also analyzed for concentrations of inflammatory, pro-fibrotic and tissue injury biomarkers including
20 transforming growth factor β 1, hyaluronic acid, tissue inhibitor of metalloproteinase-1, matrix metalloproteinase-7, connective tissue growth factor and lactate dehydrogenase activity, using commercially available ELISA. The resulting data are plotted using Graphpad prism and statistical differences between groups determined.

[00982] In this experiment, Compound 1-88 reduced kidney fibrosis by at least 20% as compared
25 to the untreated group. Compound 1-115 reduced kidney fibrosis by 55% as compared to untreated group.

Example 174: Clinical Trial in Humans with Idiopathic Pulmonary Fibrosis (IPF)**Purpose**

[00983] The purposes of this study is to assess the efficacy of treatment with a compound of
30 Formula (I), Formula (II) or Formula (III) compared with placebo in patients with idiopathic pulmonary fibrosis (IPF) and to assess the safety of treatment with a compound of Formula (I), Formula (II) or Formula (III) compared with placebo in patients with IPF.

[00984] The primary outcome variable is the absolute change in percent predicted forced vital capacity (FVC) from baseline to Week 72.

[00985] Secondary outcome measures include: composite outcomes of important IPF-related events; progression-free survival; categorical assessment of absolute change in percent predicted FVC from baseline to Week 72; change in Shortness-of-Breath from baseline to Week 72; change in percent predicted hemoglobin (Hb)-corrected carbon monoxide diffusing capacity (DLco) of the lungs from baseline to Week 72; change in oxygen saturation during the 6 minute walk test (6MWT) from baseline to Week 72; change in high-resolution computed tomography (HRCT) assessment from baseline to Week 72; change in distance walked in the 6MWT from baseline to Week 72.

Criteria

[00986] Patients eligible for this study include those patients that satisfy the following inclusion criteria: diagnosis of IPF; 40 to 80 years of age; FVC \geq 50% predicted value; DLco \geq 35% predicted value; either FVC or DLco \leq 90% predicted value; no improvement in past year; able to walk 150 meters in 6 minutes and maintain saturation \geq 83% while on no more than 6 L/min supplemental oxygen.

[00987] Patients are excluded from this study if they satisfy any of the following criteria: unable to undergo pulmonary function testing; evidence of significant obstructive lung disease or airway hyper-responsiveness; in the clinical opinion of the investigator, the patient is expected to need and be eligible for a lung transplant within 72 weeks of randomization; active infection; liver disease; cancer or other medical condition likely to result in death within 2 years; diabetes; pregnancy or lactation; substance abuse; personal or family history of long QT syndrome; other IPF treatment; unable to take study medication; withdrawal from other IPF trials.

[00988] Patients are orally dosed with either placebo or an amount of compound of Formula (I), Formula (II) or Formula (III) (1 mg/day – 1000 mg/day). The primary outcome variable will be the absolute change in percent predicted FVC from Baseline to Week 72. Patients will receive blinded study treatment from the time of randomization until the last patient randomized has been treated for 72 weeks. A Data Monitoring Committee (DMC) will periodically review safety and efficacy data to ensure patient safety.

[00989] After week 72, patients who meet the Progression of Disease (POD) definition, which is a \geq 10% absolute decrease in percent predicted FVC or a \geq 15% absolute decrease in percent predicted DLco, will be eligible to receive permitted IPF therapies in addition to their blinded study drug. Permitted IPF therapies include corticosteroids, azathioprine, cyclophosphamide and N-acetyl-cysteine.

Pharmaceutical Compositions

Example 175: Parenteral Composition

[00990] To prepare a parenteral pharmaceutical composition suitable for administration by injection (subcutaneous, intravenous, and the like), 100 mg of a water-soluble salt of a compound of Formula (I), Formula (II) or Formula (III) is dissolved in DMSO and then mixed with 10 mL of 0.9% sterile saline. The mixture is incorporated into a dosage unit form suitable for

5 administration by injection

[00991] In another embodiment, the following ingredients are mixed to form an injectable formulation: 1.2 g of a compound of Formulas (I), 2.0 mL of sodium acetate buffer solution (0.4 M), HCl (1 N) or NaOH (1 M) (q.s. to suitable pH), water (distilled, sterile) (q.s.to 20 mL). All of the above ingredients, except water, are combined and stirred and if necessary, with slight

10 heating if necessary. A sufficient quantity of water is then added.

Example 176: Oral Composition

[00992] To prepare a pharmaceutical composition for oral delivery, 100 mg of a compound of Formula (I), Formula (II) or Formula (III) is mixed with 750 mg of starch. The mixture is incorporated into an oral dosage unit for, such as a hard gelatin capsule, which is suitable for oral

15 administration.

Example 177: Sublingual (Hard Lozenge) Composition

[00993] To prepare a pharmaceutical composition for buccal delivery, such as a hard lozenge, mix 100 mg of a compound of Formula (I), Formula (II) or Formula (III) with 420 mg of powdered sugar mixed, with 1.6 mL of light corn syrup, 2.4 mL distilled water, and 0.42 mL mint extract.

20 The mixture is gently blended and poured into a mold to form a lozenge suitable for buccal administration.

Example 178: Fast-Disintegrating Sublingual Tablet

[00994] A fast-disintegrating sublingual tablet is prepared by mixing 48.5% by weigh of a compound of Formula (I), Formula (II) or Formula (III), 44.5% by weight of microcrystalline cellulose (KG-802), 5% by weight of low-substituted hydroxypropyl cellulose (50 μ m), and 2% by weight of magnesium stearate. Tablets are prepared by direct compression (*AAPS PharmSciTech*. 2006;7(2):E41). The total weight of the compressed tablets is maintained at 150 mg. The formulation is prepared by mixing the amount of compound of Formula (I), Formula (II) or Formula (III) with the total quantity of microcrystalline cellulose (MCC) and two-thirds of the

30 quantity of low-substituted hydroxypropyl cellulose (L-HPC) by using a three dimensional manual mixer (Inversina®, Bioengineering AG, Switzerland) for 4.5 minutes. All of the magnesium stearate (MS) and the remaining one-third of the quantity of L-HPC are added 30 seconds before the end of mixing.

Example 179: Inhalation Composition

[00995] To prepare a pharmaceutical composition for inhalation delivery, 20 mg of a compound of Formula (I), Formula (II) or Formula (III) is mixed with 50 mg of anhydrous citric acid and 100 mL of 0.9% sodium chloride solution. The mixture is incorporated into an inhalation delivery unit, such as a nebulizer, which is suitable for inhalation administration.

5 [00996] In another embodiment, compound of Formula (I), Formula (II) or Formula (III) (500 mg) is suspended in sterile water (100 mL), Span 85 (1 g) is added followed by addition of dextrose (5.5 g) and ascorbic acid (10 mg). Benzalkonium chloride (3 mL of a 1:750 aqueous solution) is added and the pH is adjusted to 7 with phosphate buffer. The suspension is packaged in sterile

10 nebulizers.

Example 180: Rectal Gel Composition

[00997] To prepare a pharmaceutical composition for rectal delivery, 100 mg of a compound of Formula (I), Formula (II) or Formula (III) is mixed with 2.5 g of methylcellulose (1500 mPa), 100 mg of methylparaben, 5 g of glycerin and 100 mL of purified water. The resulting gel mixture is then incorporated into rectal delivery units, such as syringes, which are suitable for rectal administration.

Example 1810: Topical Gel Composition

[00998] To prepare a pharmaceutical topical gel composition, 100 mg of a compound of Formula (I), Formula (II) or Formula (III) is mixed with 1.75 g of hydroxypropyl cellulose, 10 mL of propylene glycol, 10 mL of isopropyl myristate and 100 mL of purified alcohol USP. The resulting gel mixture is then incorporated into containers, such as tubes, which are suitable for topical administration.

Example 182: Ophthalmic Solution Composition

[00999] To prepare a pharmaceutical ophthalmic solution composition, 100 mg of a compound of Formula (I), Formula (II) or Formula (III) is mixed with 0.9 g of NaCl in 100 mL of purified water and filtered using a 0.2 micron filter. The resulting isotonic solution is then incorporated into ophthalmic delivery units, such as eye drop containers, which are suitable for ophthalmic administration.


Example 183: Nasal spray solution

30 [001000] To prepare a pharmaceutical nasal spray solution, 10 g of a compound of Formula (I), Formula (II) or Formula (III) is mixed with 30 mL of a 0.05M phosphate buffer solution (pH 4.4). The solution is placed in a nasal administrator designed to deliver 100 μ l of spray for each application.

[001001] The examples and embodiments described herein are for illustrative purposes only and various modifications or changes suggested to persons skilled in the art are to be included within the spirit and purview of this application and scope of the appended claims.

WHAT IS CLAIMED IS:

1. A compound having the structure of Formula (I), pharmaceutically acceptable salt, pharmaceutically acceptable solvate, or prodrug thereof:



Formula (I)

wherein

R^1 is $-CO_2H$, $-CO_2R^D$, $-C(=O)NHSO_2R^{10}$, $-C(=O)N(R^9)_2$, $-C(=O)NH-OH$, $-C(=O)NH-CN$, $-CN$, $-C(=NH)-NH_2$, $-C(=NH)-NHC(=O)R^D$, $-C(=O)NHCH_2CH_2SO_3H$,
10 tetrazolyl or carboxylic acid bioisostere;

R^D is H or C_1-C_6 alkyl;

ring A is a substituted or unsubstituted C_3-C_{10} cycloalkylene, a substituted or unsubstituted C_2-C_{10} heterocycloalkylene, a substituted or unsubstituted arylene, or a substituted or unsubstituted heteroarylene, where if ring A is substituted then ring A is substituted with 1, 2, or 3 R^A ;
15

ring B is a substituted or unsubstituted C_3-C_{10} cycloalkylene, a substituted or unsubstituted C_2-C_{10} heterocycloalkylene, a substituted or unsubstituted arylene, or a substituted or unsubstituted heteroarylene, where if ring B is substituted then ring B is substituted with 1, 2, or 3 R^B ;

20 L^1 is absent, C_1-C_6 alkylene, C_1-C_6 fluoroalkylene, or C_1-C_6 heteroalkylene;

L^2 is absent, C_1-C_6 alkylene, C_1-C_6 fluoroalkylene, C_1-C_6 heteroalkylene, $-O-$, $-S-$, $-SO-$, $-SO_2-$, $-NR^2-$, or $-C(=O)-$;

R^2 is H, C_1-C_4 alkyl, or C_1-C_4 fluoroalkyl;

R^3 is H, C_1-C_4 alkyl, C_3-C_6 cycloalkyl, or C_1-C_4 fluoroalkyl;

25 R^4 is $-NR^7C(=O)OCH(R^8)-CY$, $-NR^7C(=O)OC(R^8)_2-CY$, or $-NR^7C(=O)O-CY$;

R^7 is H or C_1-C_4 alkyl;

R^8 is H, C_1-C_4 alkyl, or C_1-C_4 fluoroalkyl;

CY is C_1-C_6 alkyl, a substituted or unsubstituted C_3-C_{10} cycloalkyl, a substituted or unsubstituted C_2-C_{10} heterocycloalkyl, a substituted or unsubstituted aryl, or a substituted or unsubstituted heteroaryl, wherein if CY is substituted then CY is substituted with 1, 2, or 3 R^C ; or
30

R^8 and CY are taken together with the carbon atom to which they are attached to form a substituted or unsubstituted carbocycle or a substituted or unsubstituted heterocycle;

each of R^A , R^B , and R^C are independently selected from H, halogen, -CN, -NO₂, -OH, -OR¹⁰, -SR¹⁰, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -N(R⁹)S(=O)R¹⁰, -S(=O)N(R⁹)₂, -C(=O)R¹⁰, -OC(=O)R¹⁰, -CO₂R⁹, -OCO₂R¹⁰, -N(R⁹)₂, -C(=O)N(R⁹)₂, -OC(=O)N(R⁹)₂, -NR⁹C(=O)N(R⁹)₂, -NR⁹C(=O)R¹⁰, -NR⁹C(=O)OR¹⁰, C₁-C₄alkyl, C₁-C₄fluoroalkyl, 5 C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl; each R⁹ is independently selected from H, C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene-(substituted or unsubstituted cycloalkyl), 10 -C₁-C₄alkylene-(substituted or unsubstituted heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), or -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl); or two R⁹ groups attached to the same N atom are taken together with the N atom to which they are attached to form a substituted or unsubstituted heterocycle; 15 R¹⁰ is selected from C₁-C₆alkyl, C₁-C₆heteroalkyl, C₁-C₆fluoroalkyl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, -C₁-C₄alkylene- substituted or unsubstituted cycloalkyl, -C₁-C₄alkylene-(substituted or unsubstituted heterocycloalkyl), -C₁-C₄alkylene-(substituted or unsubstituted aryl), and -C₁-C₄alkylene-(substituted or unsubstituted heteroaryl). 20



2. The compound of claim 1, wherein:

R¹ is -CO₂H, -CO₂R^D, -C(=O)NHSO₂R¹⁰, -C(=O)N(R⁹)₂, -CN, -C(=NH)-NH₂, -C(=NH)-NHC(=O)R^D, or tetrazolyl;

R³ is C₁-C₄alkyl;

25 R⁴ is -NHC(=O)OCH(R⁸)-CY;

ring A is a substituted or unsubstituted phenylene, a substituted or unsubstituted naphthylene, or a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring A is substituted then ring A is

substituted with 1 or 2 R^A; where the groups  and  are

30 on non-adjacent ring atoms of ring A;

ring B is a substituted or unsubstituted monocyclic cycloalkylene, a substituted or unsubstituted bicyclic cycloalkylene, a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted bicyclic heterocycloalkylene, a

substituted or unsubstituted phenylene, a substituted or unsubstituted naphthylene, a substituted or unsubstituted monocyclic heteroarylene, or a substituted or unsubstituted bicyclic heteroarylene, where if ring B is substituted then ring B is substituted with 1 or 2 R^B;

5 CY is a substituted or unsubstituted C₃-C₁₀cycloalkyl, a substituted or unsubstituted phenyl, a substituted or unsubstituted napthyl, a substituted or unsubstituted monocyclic heteroaryl containing 0-4 N atoms, or a substituted or unsubstituted bicyclic heteroaryl containing 0-4 N atoms, wherein if CY is substituted then CY is substituted with 1 or 2 R^C.

10 3. The compound of claim 2, wherein:


ring A is a substituted or unsubstituted phenylene, or a substituted or unsubstituted monocyclic heteroarylene containing 0 or 1 O atoms, 0 or 1 S atoms, 0-3 N atoms, and at least 2 carbon atoms in the ring, where if ring A is substituted then ring A is substituted with 1 or 2 R^A;

15 ring B is a substituted or unsubstituted monocyclic cycloalkylene, a substituted or unsubstituted monocyclic heterocycloalkylene, a substituted or unsubstituted phenylene, or a substituted or unsubstituted monocyclic heteroarylene, where if ring B is substituted then ring B is substituted with 1 or 2 R^B;

CY is a substituted or unsubstituted C₃-C₆cycloalkyl, a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl containing 0-4 N atoms, wherein if CY is substituted then CY is substituted with 1 or 2 R^C;

each of R^A, R^B, and R^C are independently selected from H, halogen, -CN, -NO₂, -OH, -OR¹⁰, -N(R⁹)₂, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

25 4. The compound of claim 3, wherein the compound of Formula (I) has the structure of Formula (II):



Formula (II)

wherein,

30 each of A¹, A², and A³ is independently selected from CR^A and N;

each R^A is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃;

each R^B is independently selected from H, F, Cl, Br, I, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃.

5. The compound of claim 4, wherein:

each of A¹, A², and A³ is CR^A; or

5 one of A¹, A², and A³ is N.

6. The compound of claim 5, wherein:

R¹ is -CO₂H, -CO₂R^D, or -C(=O)NHSO₂R¹⁰;

R^D is H, -CH₃, or -CH₂CH₃;

R⁸ is H, -CH₃, -CH₂CH₃ or -CF₃.

10 7. The compound of claim 6, wherein:

R¹ is -CO₂H;

R³ is -CH₃ or -CH₂CH₃;

R⁸ is H, or -CH₃.

8. The compound of claim 7, wherein:

15 R³ is -CH₃;

L¹ is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, or - NR¹¹-C₁-C₄alkylene -; R¹¹ is H or C₁-C₂alkyl;

20 L² is absent, C₁-C₆alkylene, C₁-C₆fluoroalkylene, -C₁-C₆alkylene-O-, -C₁-C₃alkylene-O-C₁-C₃alkylene -, -O-C₁-C₆alkylene -, -C₁-C₆alkylene-S-, -C₁-C₃alkylene-S-C₁-C₃alkylene -, -S-C₁-C₆alkylene -, -C₁-C₆alkylene-SO-, -C₁-C₃alkylene-SO-C₁-C₃alkylene -, -SO-C₁-C₆alkylene -, -C₁-C₆alkylene-SO₂-, -C₁-C₃alkylene-SO₂-C₁-C₃alkylene -, -SO₂-C₁-C₆alkylene -, -C₁-C₄alkylene-NR¹¹-, -C₁-C₂alkylene-NR¹¹-C₁-C₂alkylene -, - NR¹¹-C₁-C₄alkylene -, -O-, -S-, -SO-, -SO₂- or -NR²-;

25 R¹¹ is H or C₁-C₂alkyl;

CY is a substituted or unsubstituted phenyl, or a substituted or unsubstituted monocyclic heteroaryl containing 1-3 N atoms, wherein if CY is substituted then CY is substituted with 1 or 2 R^C;


30 each R^C is independently selected from H, halogen, -OH, C₁-C₄alkyl, C₁-C₄fluoroalkyl, C₁-C₄fluoroalkoxy, C₁-C₄alkoxy, and C₁-C₄heteroalkyl.

9. The compound of claim 8, wherein:

L¹ is absent, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, -CH₂CH(CH₃)-, or -CH₂C(CH₃)₂;

L^2 is absent, $-\text{CH}_2-$, $-\text{CH}_2\text{O}-$, $-\text{OCH}_2-$, $-\text{CH}_2\text{S}-$, $-\text{SCH}_2-$, $-\text{CH}_2\text{NH}-$, $-\text{NHCH}_2-$, $-\text{O}-$, $-\text{S}-$, or $-\text{NH}-$.

10. The compound of claim 9, wherein:





5 ring B is substituted or unsubstituted phenylene, where if ring B is substituted then ring B is substituted with 1 or 2 R^B ;

CY is a substituted or unsubstituted phenyl, wherein if CY is substituted then CY is substituted with 1 or 2 R^C .

11. The compound of claim 10, wherein:

10 L^1 is absent, $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{CH}_2\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2\text{CH}_2-$;

L^2 is absent;



R^8 is $-\text{CH}_3$;

15 each R^C is independently selected from H, F, Cl, Br, I, $-\text{CH}_3$, $-\text{CF}_3$, $-\text{OH}$, $-\text{OCF}_3$, and $-\text{OCH}_3$.

12. The compound of claim 11, wherein:

L^1 is absent or $-\text{CH}_2-$;

each R^A is independently selected from H, F, Cl, $-\text{CH}_3$, and $-\text{OH}$;


20 each R^B is independently selected from H, F, Cl, $-\text{CH}_3$, and $-\text{OH}$;

each R^C is independently selected from H, F, Cl, $-\text{CH}_3$, $-\text{CF}_3$, $-\text{OH}$, $-\text{OCF}_3$, and $-\text{OCH}_3$.

13. The compound of claim 12, wherein:


L^1 is $-\text{CH}_2-$;

L^2 is absent;



CY is phenyl, 2-fluorophenyl or 2-chloro-phenyl.

14. The compound of claim 1, wherein the compound of Formula (I) has the structure of Formula (III):



Formula (III)

wherein

5 R^1 is $-CO_2H$, $-CO_2R^D$, $-CN$, tetrazolyl, $-C(=O)N(R^9)_2$, $-C(=O)NHCH_2CH_2SO_3H$, or $-C(=O)NHSO_2R^{10}$; R^D is H or C_1 - C_4 alkyl;

L^1 is absent or C_1 - C_6 alkylene;

R^3 is H, C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, or C_1 - C_4 fluoroalkyl;

R^7 is H or C_1 - C_4 alkyl;

R^8 is H, C_1 - C_4 alkyl, or C_1 - C_4 fluoroalkyl;

10 R^9 is a H, C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_3 - C_6 cycloalkyl, or a substituted or unsubstituted phenyl;

R^{10} is a C_1 - C_6 alkyl, C_1 - C_6 fluoroalkyl, C_3 - C_6 cycloalkyl, or a substituted or unsubstituted phenyl; each of R^A , R^B , and R^C are independently selected from H, F, Cl, Br, I, $-CN$, $-OH$, C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, C_1 - C_4 fluoroalkoxy, C_1 - C_4 alkoxy, and C_1 - C_4 heteroalkyl;

15 m is 0, 1, or 2;

n is 0, 1, or 2;

p is 0, 1, or 2.

15. The compound of claim 14, wherein:

20 R^1 is $-CO_2H$, $-CO_2R^D$ or $-C(=O)NHSO_2R^{10}$;

R^3 is C_1 - C_4 alkyl;

R^7 is H;

R^{10} is a C_1 - C_6 alkyl or a substituted or unsubstituted phenyl;

each R^A is independently selected from H, F, Cl, Br, I, $-CH_3$, $-CF_3$, $-OH$, $-OCF_3$, and $-OCH_3$;

each R^B is independently selected from H, F, Cl, Br, I, $-CH_3$, $-CF_3$, $-OH$, $-OCF_3$, and $-OCH_3$;

each R^C is independently selected from H, F, Cl, Br, I, $-CH_3$, $-CF_3$, $-OH$, $-OCF_3$, and $-OCH_3$;

m is 0 or 1;

p is 0 or 1.

16. The compound of claim 15, wherein:

R¹ is -CO₂H or -CO₂R^D;

5 R³ is -CH₃ or -CH₂CH₃;

R⁸ is H, -CH₃ or -CF₃;

R^D is H, -CH₃, or -CH₂CH₃.

17. The compound of claim 16, wherein:


R¹ is -CO₂H;

10 R³ is -CH₃;

R⁸ is -CH₃;

L¹ is absent, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, -CH₂CH(CH₃)-, or -CH₂C(CH₃)₂-.

18. The compound of claim 17, wherein the compound of Formula (III) has the following structure:



19. The compound of claim 18, wherein:

L¹ is absent, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, or -C(CH₂CH₃)₂-,

each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, -OH, -OCF₃, and -OCH₃;

20 m is 0;

n is 0, 1, or 2;

p is 0.


20. The compound of claim 19, wherein:

L¹ is absent or -CH₂-;

25 each R^C is independently selected from H, F, Cl, -CH₃, -CF₃, and -OH;

n is 0 or 1.

21. The compound of claim 16, wherein wherein the compound of Formula (III) has the following structure:



22. The compound of claim 21, wherein:

L^1 is absent, $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{CH}(\text{CH}_2\text{CH}_3)-$, or $-\text{C}(\text{CH}_2\text{CH}_3)_2-$;
 each R^C is independently selected from H, F, Cl, $-\text{CH}_3$, $-\text{CF}_3$, $-\text{OH}$, $-\text{OCF}_3$, and $-\text{OCH}_3$;
 5 m is 0;
 n is 0, 1, or 2;
 p is 0.

23. The compound of claim 22, wherein:

L^1 is absent or $-\text{CH}_2-$;
 10 each R^C is independently selected from H, F, Cl, $-\text{CH}_3$, $-\text{CF}_3$, and $-\text{OH}$;
 n is 0 or 1.

24. The compound of claim 15, wherein:

R^1 is $-\text{C}(=\text{O})\text{NHSO}_2\text{R}^{10}$;
 R^3 is $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$;
 15 R^8 is H, $-\text{CH}_3$ or $-\text{CF}_3$;
 R^{10} is $-\text{CH}_3$, or $-\text{CH}_2\text{CH}_3$.

25. The compound of claim 1 selected from:

1-(4-((1-(2-Chloro-phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-benzyl)-
 1*H*-[1,2,4]triazole-3-carboxylic acid (Compound 1-1); 1-(4-((1-(2-Chloro-phenyl)-
 ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-benzyl)-1*H*-imidazole-4-carboxylic acid
 20 (Compound 1-2); 3-(4-((1-(2-Chloro-phenyl)-ethoxycarbonylamino)-3-methyl-
 isoxazol-5-yl)-benzyl)-3*H*-imidazole-4-carboxylic acid (Compound 1-3); 1-(4-((1-(2-Chloro-
 phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-benzyl)-5-
 trifluoromethyl-1*H*-pyrazole-4-carboxylic acid (Compound 1-4); 3-(4-((1-(2-Chloro-
 phenyl)-ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-benzyl)-benzoic acid
 25 (Compound 1-5); 3-((4-(2-Bromo-benzyloxycarbonylamino)-3-methyl-isoxazol-5-yl)-
 benzyl)-benzoic acid (Compound 1-6); [3-(4-((1-(2-Chloro-phenyl)-
 ethoxycarbonylamino)-3-methyl-isoxazol-5-yl)-benzyl)-4-methoxy-phenyl]-acetic acid
 (Compound 1-7); 1-(4-((1-(2-Chloro-phenyl)-ethoxycarbonylamino)-3-methyl-

5 isoxazol-5-yl}-benzyl)-1*H*-pyrazole-4-carboxylic acid (Compound 1-9); 4'-[3-Methyl-4-((R)-1-pyridin-2-yl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-11); 3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-benzoic acid (Compound 1-12); 4'-[3-Methyl-4-(6-methyl-pyridin-2-ylmethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-13); 4-Chloro-3-(4-{4-[1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-benzoic acid (Compound 1-14); 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-thiophene-2-carboxylic acid (Compound 1-15); 4'-[4-(Indan-1-yloxycarbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-3-carboxylic acid (Compound 1-16); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-17); [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-18); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzylsulfanyl)-phenyl]-acetic acid (Compound 1-19); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-20); [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-21); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-22); (4'-{3-Methyl-4-[1-(2-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-23); (4'-{3-Methyl-4-[1-(3-trifluoromethyl-phenyl)-ethoxycarbonylamino]-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-24); (4'-{4-[1-(2,4-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-25); (4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-26); (4'-{4-[1-(3-Bromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-27); (4'-{4-[1-(2-Methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-28); {4'-[3-Methyl-4-(1-methyl-2-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-29); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-methoxy-biphenyl-3-yl)-acetic acid (Compound 1-30); 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid (Compound 1-31); 3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid (Compound 1-32); 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid

(Compound 1-34); 2-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid (Compound 1-35); 6-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-36); 5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid (Compound 1-38); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-39); [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-6-methoxy-pyridin-3-yl]-acetic acid (Compound 1-40); [5-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-41); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-42); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-4-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-43); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-yl)-acetic acid (Compound 1-44); 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid (Compound 1-45); 4-(3-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid (Compound 1-46); [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid ethyl ester (Compound 1-47); [4-(5-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-2-yl)-phenyl]-acetic acid (Compound 1-48); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-methoxy-biphenyl-3-yl)-acetic acid (Compound 1-49); 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-isonicotinic acid (Compound 1-50); 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid (Compound 1-51); 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-carboxylic acid (Compound 1-52); 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-53); 6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-54); 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-nicotinic acid (Compound 1-55); 5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridine-2-carboxylic acid (Compound 1-56); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-2-yl)-acetic acid (Compound 1-35)

57); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-58); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-yl)-acetic acid (Compound 1-59); (3'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-yl)-acetic acid (Compound 1-60); [6-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-61); 3-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-62); [5-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-acetic acid (Compound 1-63); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-6-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-64); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-4-fluoro-biphenyl-3-yl)-acetic acid (Compound 1-65); 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohex-3-enecarboxylic acid (Compound 1-66); 4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-cyclohexanecarboxylic acid (Compound 1-67); (4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid methyl ester (Compound 1-69); 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid ethyl ester (Compound 1-70); 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-71); 2-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-72); [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-pyridin-3-yl)-phenyl]-acetic acid (Compound 1-73); 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-74); 4-(4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid (Compound 1-75); 4'-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-3-carboxylic acid (Compound 1-76); (4'-{4-[1-(4-Chloro-2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-77); (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-78); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-79); [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-80); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-

yl}-benzyl)-phenyl]-acetic acid (Compound 1-81); [4-Fluoro-3-(4-{4-[(R)-1-(2-fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyl)-phenyl]-acetic acid (Compound 1-82); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid ethyl ester (Compound 1-83); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-84); (4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-acetic acid (Compound 1-85); 2-(4'-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-86); 2-[6-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-pyridin-3-yl]-2-methyl-propionic acid (Compound 1-87); (4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-88); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-89); 2-(4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-2-methyl-propionic acid (Compound 1-90); 2-(4'-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-91); 2-(4'-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-92); 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-4-carboxylic acid (Compound 1-93); 2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-thiazole-5-carboxylic acid (Compound 1-94); [3-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid methyl ester (Compound 1-95); [3-(4-{4-[1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-96); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-97); [3-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-98); [2-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylmethanesulfonyl)-phenyl]-acetic acid (Compound 1-99); [4-(4-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-100); (4'-{4-[1-(2,6-Dichloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-101); [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid methyl ester (Compound 1-102); [4-(4-{4-[(R)-1-(2-Chloro-phenyl)-

5

10

15

20

25

30

35

ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid
(Compound 1-103); 4-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy-methyl)-benzoic acid (Compound 1-104); [4-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid ethyl ester (Compound 1-105); [3-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-106); [2-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-107); [6-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-pyridin-3-yl]-acetic acid (Compound 1-108); [4-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenylsulfanyl)-phenyl]-acetic acid (Compound 1-109); 2-(4'-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-methyl-biphenyl-4-yl)-propionic acid (Compound 1-110); (4'-{4-[*(S)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-111); (4'-{4-[*(S)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-112); {4'-[4-(2-Chloro-benzyloxy-carbonylamino)-3-methyl-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-113); [2-(4-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-benzyloxy)-phenyl]-acetic acid (Compound 1-114); {4'-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-115); (4'-{4-[1-(2,3-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-116); (4'-{4-[1-(2,4-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-117); (4'-{4-[1-(2-Fluoro-4-methoxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-118); (4'-{4-[1-(2,5-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-119); (4'-{4-[1-(2,6-Difluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-120); {4'-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-3-yl}-acetic acid (Compound 1-121); 4'-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-carboxylic acid (Compound 1-122); {4'-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-2-yl}-acetic acid (Compound 1-123); {4'-[3-Methyl-4-((*R*)-1-*o*-tolyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-124); 4'-{(4-Isopropoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-carboxylic acid (Compound 1-125); (6'-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-

isoxazol-5-yl}-[2,3']bipyridinyl-5-yl)-acetic acid (Compound 1-126); 2-(4'-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-127); 2-(4'-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-propionic acid (Compound 1-128); (3'-Chloro-4'-{4-[*(R)*-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-129); 2-(4'-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-butyric acid (Compound 1-130); (2'-Chloro-4'-{4-[*(R)*-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-131); (4'-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-2'-fluoro-biphenyl-4-yl)-acetic acid (Compound 1-132); 4'-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-carboxylic acid (Compound 1-133); [5-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid methyl ester (Compound 1-134); [5-(4-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenyl)-tetrazol-2-yl]-acetic acid (Compound 1-135); [4'-(4-*tert*-Butoxycarbonylamino-3-methyl-isoxazol-5-yl)-biphenyl-4-yl]-acetic acid (Compound 1-136); [3-(4-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-137); [4-(4-{4-[*(R)*-1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-phenoxy)-phenyl]-acetic acid (Compound 1-138); (4'-{4-[*(R)*-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-hydroxy-acetic acid (Compound 1-139); 1-{4-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-benzyl}-azetidine-3-carboxylic acid (Compound 1-140); (4'-{4-[*(R)*-1-(3,5-Dibromo-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-141); 2-(4-{5-[3-Methyl-4-((*R*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-pyridin-2-yl}-phenyl)-butyric acid (Compound 1-142); {4'-[3-Methyl-4-((*S*)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-143); (4'-{4-[*(R)*-1-(3-Hydroxy-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-biphenyl-4-yl)-acetic acid (Compound 1-144); {4'-[3-Methyl-4-(1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-145); {4'-[3-Methyl-4-(1-phenyl-ethoxy-d9-carbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetic acid (Compound 1-146); {5-[4-(4-Cyclopropylcarbamoyl-benzyl)-phenyl]-3-methyl-isoxazol-4-yl}-carbamic acid 1-(2-chloro-phenyl)-ethyl ester (Compound 2-1); [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (*R*)-1-(2-chloro-phenyl)-ethyl ester (Compound 2-35)

2); [5-(4'-Cyanomethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-3); {3-Methyl-5-[4'-(2H-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-4); {3-Methyl-5-[4'-(2H-tetrazol-5-ylmethyl)-biphenyl-4-yl]-isoxazol-4-yl}-carbamic acid (R)-1-(2-chloro-phenyl)-ethyl ester (Compound 2-5); [5-(4'-Carbamimidoylmethyl-biphenyl-4-yl)-3-methyl-isoxazol-4-yl]-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-6); {5-[4'-(2-Acetylamino-2-imino-ethyl)-biphenyl-4-yl]-3-methyl-isoxazol-4-yl}-carbamic acid (R)-1-(2-fluoro-phenyl)-ethyl ester (Compound 2-7); 2-(2-{4'-[3-Methyl-4-((R)-1-phenyl-ethoxycarbonylamino)-isoxazol-5-yl]-biphenyl-4-yl}-acetylamino)-ethanesulfonic acid (Compound 2-8); [4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-phenyl]-acetic acid (Compound 3-1); 3-[4-(6-{4-[1-(2-Chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-naphthalen-2-yl)-benzylsulfanyl]-propionic acid (Compound 3-2); [4-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-3); [4-(5-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-4); [2-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-5); and [3-(4-{4-[(R)-1-(2-Fluoro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl}-thiazol-2-yl)-phenyl]-acetic acid (Compound 3-6).

26. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of claims 1-25, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable inactive ingredient.

25 27. The pharmaceutical composition of claim 26, wherein the pharmaceutical composition is:
(i) formulated for intravenous injection, subcutaneous injection, oral administration, inhalation, nasal administration, topical administration, ophthalmic administration or otic administration; or
(ii) a tablet, a pill, a capsule, a liquid, an inhalant, a nasal spray solution, a suppository, a suspension, a gel, a colloid, a dispersion, a suspension, a solution, an emulsion, an ointment, a lotion, an eye drop or an ear drop.

30 28. The pharmaceutical composition of claim 26 or claim 27 further comprising one or more additional therapeutically active agents selected from: corticosteroids, immunosuppresant, analgesics, anti-cancer agent, anti-inflammatories, chemokine receptor antagonists, bronchodilators, leukotriene receptor antagonists, leukotriene formation inhibitors,

monoacylglycerol kinase inhibitors, phospholipase A₁ inhibitors, phospholipase A₂ inhibitors, and lysophospholipase D (lysoPLD) inhibitors, autotoxin inhibitors, decongestants, antihistamines, mucolytics, anticholinergics, antitussives, expectorants, and β-2 agonists.

5 29. A method of inhibiting the physiological activity of LPA in a mammal comprising administering a therapeutically effective amount of a compound according to any one of claims 1-25 or a pharmaceutically acceptable salt thereof to the mammal in need thereof.

10 30. A method for treating or preventing a LPA-dependent or LPA-mediated disease or condition in a mammal comprising administering a therapeutically effective amount of a compound according to any one of claims 1-25 or a pharmaceutically acceptable salt thereof to the mammal in need thereof.

15 31. The method of claim 30, wherein the LPA-dependent or LPA-mediated disease or condition is selected from fibrosis of organs or tissues, scarring, liver diseases, dermatological conditions, cancer, cardiovascular diseases, respiratory diseases or conditions, inflammatory diseases, gastrointestinal tract diseases, renal diseases, urinary tract-associated diseases, inflammatory diseases of lower urinary tract, dysuria, frequent urination, pancreas disease, arterial obstruction, cerebral infarction, cerebral hemorrhage, pain, peripheral neuropathy, and fibromyalgia.

20 32. The method of claim 30, wherein the LPA-dependent or LPA-mediated disease or condition is a respiratory disease or condition.

33. The method of claim 32, wherein the respiratory disease or condition is asthma, chronic obstructive pulmonary disease (COPD), pulmonary fibrosis, pulmonary arterial hypertension, or acute respiratory distress syndrome.

25 34. The method of claim 30, wherein the LPA-dependent or LPA-mediated disease or condition is selected from idiopathic pulmonary fibrosis; other diffuse parenchymal lung diseases of different etiologies including iatrogenic drug-induced fibrosis, occupational and/or environmental induced fibrosis, granulomatous diseases (sarcoidosis, hypersensitivity pneumonia), collagen vascular disease, alveolar proteinosis, langerhans cell granulomatosis, lymphangioleiomyomatosis, inherited diseases (Hermansky-Pudlak Syndrome, tuberous sclerosis, neurofibromatosis, metabolic storage disorders, familial interstitial lung disease); radiation induced fibrosis; chronic obstructive pulmonary disease (COPD); scleroderma; systemic sclerosis, bleomycin induced pulmonary fibrosis; chronic asthma; silicosis; asbestos induced pulmonary fibrosis; acute respiratory distress syndrome (ARDS); kidney fibrosis; tubulointerstitium fibrosis; glomerular nephritis; 30 focal segmental glomerular sclerosis; lupus nephritis, IgA nephropathy; hypertension;

5 Alport; gut fibrosis; liver fibrosis; cirrhosis; alcohol induced liver fibrosis; toxic/drug induced liver fibrosis; hemochromatosis; nonalcoholic steatohepatitis (NASH); biliary duct injury; primary biliary cirrhosis; infection induced liver fibrosis; viral induced liver fibrosis; and autoimmune hepatitis; corneal scarring; hypertrophic scarring; Dupuytren's disease, keloids, cutaneous fibrosis; cutaneous scleroderma; spinal cord injury/fibrosis; myelofibrosis; vascular restenosis; atherosclerosis; arteriosclerosis; Wegener's granulomatosis; Peyronie's disease, chronic lymphocytic leukemia, tumor metastasis, transplant organ rejection, endometreosis, neonatal respiratory distress syndrome and neuropathic pain.

10 35. A method of controlling the activation of LPA receptors in a tissue in a mammal comprising administering a compound of any one of claims 1-25 to mammal.

36. The method of claim 43, wherein the activation of LPA receptors in a tissue in a mammal results in fibrosis.

37. A method for the treatment or prevention of organ fibrosis in a mammal comprising 15 administering a therapeutically effective amount of a selective LPA1 receptor antagonist to the mammal in need thereof.

38. The method of claim 37, wherein the organ fibrosis comprises lung fibrosis, renal fibrosis, hepatic fibrosis or cutaneous fibrosis.

39. A method of improving lung function in a mammal comprising administering a 20 therapeutically effective amount of a selective LPA1 receptor antagonist to the mammal in need thereof.

40. The method of claim 39, wherein the mammal has been diagnosed as having lung fibrosis.

41. A method of treating idiopathic pulmonary fibrosis in a mammal comprising 25 administering a therapeutically effective amount of a LPA1 receptor antagonist to the mammal in need thereof.

42. A method of controlling an abnormal accumulation or activation of cells, fibronectin, collagen or increased fibroblast recruitment in a tissue of a mammal comprising administering a selective LPA1 receptor antagonist to mammal.

30 43. The method of claim 42, wherein the abnormal accumulation or activation of cells, fibronectin, collagen or increased fibroblast recruitment in the tissue results in fibrosis.

44. A method of controlling an abnormal healing process in a tissue of a mammal comprising administering a selective LPA1 receptor antagonist to mammal.

45. The method of claim 44, wherein the abnormal healing process in the tissue of the 35 mammal results in fibrosis.

46. The method of any one of claims 37-45, wherein the selective LPA1 receptor antagonist is a compound according to any one of claims 1-25 or a pharmaceutically acceptable salt.