



US 20130104259A1

(19) **United States**(12) **Patent Application Publication**
Sampson et al.(10) **Pub. No.: US 2013/0104259 A1**(43) **Pub. Date: Apr. 25, 2013**(54) **AXMI-115, AXMI-113, AXMI-005, AXMI-163
AND AXMI-184: INSECTICIDAL PROTEINS
AND METHODS FOR THEIR USE**(71) Applicant: **ATHENIX CORPORATION,**
Morrisville, NC (US)(72) Inventors: **Kimberly S. Sampson,** Durham, NC
(US); **Daniel J. Tomso,** Bahama, NC
(US); **Shruti Agarwal,** Durham, NC
(US); **Brian McNulty,** Raleigh, NC
(US); **Chris Campbell,** Cary, NC (US)(73) Assignee: **ATHENIX CORPORATION,**
Morrisville, NC (US)(21) Appl. No.: **13/675,572**(22) Filed: **Nov. 13, 2012****Related U.S. Application Data**(63) Continuation of application No. 12/497,221, filed on
Jul. 2, 2009, now Pat. No. 8,334,431.(60) Provisional application No. 61/077,812, filed on Jul. 2,
2008, provisional application No. 61/158,953, filed on
Mar. 10, 2009.**Publication Classification**(51) **Int. Cl.****C07K 14/325** (2006.01)**A01N 37/18** (2006.01)**C12N 15/82** (2006.01)**C07K 16/12** (2006.01)(52) **U.S. Cl.**CPC **C07K 14/325** (2013.01); **C07K 16/1278**
(2013.01); **A01N 37/18** (2013.01); **C12N**
15/8286 (2013.01)USPC **800/279**; 536/23.71; 530/350; 530/387.9;
514/4.5; 435/320.1; 800/302; 435/419; 435/412;
435/252.33; 435/252.31

(57)

ABSTRACT

Compositions and methods for conferring insecticidal activity to host cells are provided. Compositions comprising a coding sequence for a delta-endotoxin polypeptide are provided. The coding sequences can be used in DNA constructs or expression cassettes for transformation and expression in host cells. Compositions also comprise transformed host cells. In particular, isolated delta-endotoxin nucleic acid molecules are provided. Additionally, amino acid sequences corresponding to the polynucleotides are encompassed, and antibodies specifically binding to those amino acid sequences. In particular, the present invention provides for isolated nucleic acid molecules comprising nucleotide sequences encoding the amino acid sequence shown in SEQ ID NO:4, 5, 6, 13, or 14, or the nucleotide sequence set forth in SEQ ID NO:1, 2, 3, 11, or 12, as well as variants and fragments thereof.

Optaxmi113	MMNMNTKLSTRALPSFIDYFNGLIGFATGKIDYNNMIFKFTTGGGLFLDLILKMQQLNE	60
Optaxmi005	MMNMNTKLNAARALPSFIDYFNGLIGFATGKIDYNNMIFKFTTGGGLFLDLILKMQQLNE	60
Optaxmi115	MMNMNTKLNAARALPSFIDYFNGLIGFATGKIDYNNMIFKFTTGGGLFLDLILKMQQLNE *****:*****:*****:*****:*****:*****:*****:*****	60
Optaxmi113	ISGKLDGVNGSINDLIAQGNLNTLSKEILKIANEQNVQVLDVNMKLDIAINTMLHITYLPK	120
Optaxmi005	ISGKLDGVNGSINDLIAQGNLNTLSKEILKIANEQNVQVLDVNMKLDIAINTMLHITYLPK	120
Optaxmi115	ISGKLDGVNGSINDLIAQGNLNTLSKEILKIANEQNVQVLDVNMKLDIAINTMLHITYLPK *****:*****:*****:*****:*****:*****:*****:*****	120
Optaxmi113	ITSMLSGVMMQNYALSLQIEYLSKQIQEISDKLDIIVNVNLINSTLTETTPAYQRKYVN	180
Optaxmi005	ITSMLSGVMMQNYALSLQIEYLSKQIQEISDKLDIIVNVNLINSTLTETTPAYQRKYVN	180
Optaxmi115	ITSMLSGVMMQNYALSLQIEYLSKQIQEISDKLDIIVNVNLINSTLTETTPAYQRKYVN *****:*****:*****:*****:*****:*****:*****:*****	180
Optaxmi113	EKFEELTFATETNLKVKK----DGSPADILCELTELTELAKSVTNQDVGGFEFYINTFH	235
Optaxmi005	EKFEELTFATETNLKVKK----DGSPADILCELTELTELAKSVTNQDVGGFEFYINTFH	235
Optaxmi115	EKFEELTFATETNLKVKK----DGSPADILCELTELTELAKSVTNQDVGGFEFYINTFH *****:*****:*****:*****:*****:*****:*****:*****	240
Optaxmi113	DVMYGNMLFGSALKTASELITRENKVTSGSEVNVYNFLVIVLTALQAKAFLTLTTCRKL	295
Optaxmi005	DVMYGNMLFGSALKTASELITRENKVTSGSEVNVYNFLVIVLTALQAKAFLTLTTCRKL	295
Optaxmi115	DVLIGNMLFGSALKTASELITRENKVTSGSEVNVYNFLVIVLTALQAKAFLTLTTCRKL *:*****:*****:*****:*****:*****:*****:*****:*****	300
Optaxmi113	LGLADIDYTSIMNNEHLNKEKEEFERNVILEFTLSNFTSNPNXINTKGSDEDAEVLIQAEPGH	355
Optaxmi005	LGLADIDYTSIMNNEHLNKEKEEFERNVILEFTLSNFTSNPNXINTKGSDEDAEVLIQAEPGH	355
Optaxmi115	LGLSDIDYTSIMNNEHLNKEKEEFERNVILEFTLSNFTSNPNXINTKGSDEDAEVLIQAEPGH *****:*****:*****:*****:*****:*****:*****:*****	360
Optaxmi113	ALVGFEMINHPSPALKVYQAKLTNYQVQDKSLSETVYGDMDKILQPKSKQMYLIHNIT	415
Optaxmi005	ALVGFEMINHPSPALKVYQAKLTNYQVQDKSLSETVYGDMDKILQPKSKQMYLIHNIT	415
Optaxmi115	ALVGFEMINHPSPALKVYQAKLTNYQVQDKSLSETVYGDMDKILQPKSKQMYLIHNIT *****:*****:*****:*****:*****:*****:*****:*****	420
Optaxmi113	FPNEYVITETIETKKKNSIRYEVIANYYEFSGGDIDLNKKVKS--SEAEVSTLSVSDND	472
Optaxmi005	FPNEYVITETIETKKKNSIRYEVIANYYEFSGGDIDLNKKVKS--SEAEVSTLSVSDND	472
Optaxmi115	FPDGYVITKITFEKKLNNLIYEATANFYDPSTGSDIDLNKKQVESTFPQTEYITMDIGDD *:*****:*****:*****:*****:*****:*****:*****:*****	480
Optaxmi113	AIYMPGLVISETFLTPFKGFLTPDESSRLVTLTKCSYLREIILATDLSNKKATKLIAPPN	532
Optaxmi005	GIYMPGLVISETFLTPFKGFLTPDESSRLVTLTKCSYLREIILATDLSNKKATKLIAPPN	532
Optaxmi115	GIYMPGLVISETFLTPFKGFLTPDESSRLVTLTKCSYLREIILATDLSNKKATKLIAPPN :*****:*****:*****:*****:*****:*****:*****:*****	540
Optaxmi113	GFISNLVENGDEADNIEPKNKNNKRAYVQHTGVNGIKALYTDGGEFSQFLQDKIKSK	592
Optaxmi005	GFISNLVENGDEADNIEPKNKNNKRAYVQHTGVNGIKALYTDGGEFSQFLQDKIKSK	592
Optaxmi115	VFISNVKNWDIEEDSLFVPWANKNRAYVQHTGVNGIKALYTDGGEFSQFLQDKIKSK *****:*****:*****:*****:*****:*****:*****:*****	600
Optaxmi113	TEYIIQYTVKKGESIHLNKNENTGYIHYEDTNNMLDYQTIKRAFTTGDLKGVYLLIKSQ	652
Optaxmi005	TEYIIQYTVKKGESIHLNKNENTGYIHYEDTNNMLDYQTIKRAFTTGDLKGVYLLIKSQ	652
Optaxmi115	TEYIIQYTVKKGESIHLNKNENTGYIHYEDTNNMLDYQTIKRAFTTGDLKGVYLLIKSQ *:*****:*****:*****:*****:*****:*****:*****:*****	660

FIG. 1A

```
Optaxmi113      NCYKAWGDNFLITEIRPRE-VVSPELIKVENWICMOGSNHVNPDSLLLFTCCRSILKQNL 711
Optaxmi005      NGDEAWGDNFTILEISFSEKLLSPELINVNNWIRTG-STHISGNLTILYQGGGGLKQNL 711
Optaxmi115      NGYEAWGDNFIILEAKLFETPESPELIKENDWERFG-TTYLTGNELRIDHSRGGYFROSL 719
                ** :***** * *      * *****:.* * :.:. : * : . . :.*.*

Optaxmi113      QLDSYSTYRVRFSLMVIGKAKVIIRNSS-EVLFEKSYVNDSEGVLEGVSEFTTKSIQDN 770
Optaxmi005      QLDSYSTYRVNFS--VTGDANVRIIRNSR-EVLFEKKYMSG---AKDVSEIFTTKLGKDN 764
Optaxmi115      NIDSYSTYDLSFS-FSGLWAKVIVKNSRGVLFEEKVKNNGSS--YEDISESPTASNKDG 776
                :.*:*** ; **      *:* :.* ***** ..      :.* ** * :*,

Optaxmi113      FYVELSNEGTFGSKDVAYFYNFSIR---- 795
Optaxmi005      FYIELSQGNLYGGPLVKFNDVSIK---- 789
Optaxmi115      FFIELTAER--TSSTFHSFRDISIKERIE 803
                *.:** : . . * :.*: ~~~~~
```

FIG. 1B




FIG. 2

AXMI-115, AXMI-113, AXMI-005, AXMI-163 AND AXMI-184: INSECTICIDAL PROTEINS AND METHODS FOR THEIR USE

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application is a continuation of U.S. patent application Ser. No. 12/497,221, filed Jul. 2, 2009, which claims the benefit of U.S. Provisional Application Ser. No. 61/077,812, filed Jul. 2, 2008, and U.S. Provisional Application Ser. No. 61/158,953, filed Mar. 10, 2009, the contents of which are hereby incorporated in their entirety by reference herein.

REFERENCE TO SEQUENCE LISTING SUBMITTED AS A TEXT FILE VIA EFS-WEB

[0002] The official copy of the sequence listing is submitted concurrently with the specification as an ASCII formatted text file via EFS-Web, with a file name of "APA058US01NSEQLIST.txt", a creation date of Nov. 9, 2012, and a size of 88 kilobytes. The sequence listing filed via EFS-Web is part of the specification and is hereby incorporated in its entirety by reference herein.

FIELD OF THE INVENTION

[0003] This invention relates to the field of molecular biology. Provided are novel genes that encode insecticidal proteins. These proteins and the nucleic acid sequences that encode them are useful in preparing insecticidal formulations and in the production of transgenic insect-resistant plants.

BACKGROUND OF THE INVENTION

[0004] *Bacillus thuringiensis* is a Gram-positive spore forming soil bacterium characterized by its ability to produce crystalline inclusions that are specifically toxic to certain orders and species of insects, but are harmless to plants and other non-targeted organisms. For this reason, compositions including *Bacillus thuringiensis* strains or their insecticidal proteins can be used as environmentally-acceptable insecticides to control agricultural insect pests or insect vectors for a variety of human or animal diseases.

[0005] Crystal (Cry) proteins (delta-endotoxins) from *Bacillus thuringiensis* have potent insecticidal activity against predominantly Lepidopteran, Dipteran, and Coleopteran larvae. These proteins also have shown activity against *Hymenoptera*, *Homoptera*, *Phthiraptera*, *Mallophaga*, and *Acari* pest orders, as well as other invertebrate orders such as Nematelminthes, Platyhelminthes, and Sarcostomastigophora (Feitelson (1993) The *Bacillus Thuringiensis* family tree. In *Advanced Engineered Pesticides*, Marcel Dekker, Inc., New York, N.Y.) These proteins were originally classified as CryI to CryV based primarily on their insecticidal activity. The major classes were Lepidoptera-specific (I), Lepidoptera- and Diptera-specific (II), Coleoptera-specific (III), Diptera-specific (IV), and nematode-specific (V) and (VI). The proteins were further classified into subfamilies; more highly related proteins within each family were assigned divisional letters such as Cry1A, Cry1B, Cry1C, etc. Even more closely related proteins within each division were given names such as Cry1C1, C1C2, etc.

[0006] A new nomenclature was recently described for the Cry genes based upon amino acid sequence homology rather than insect target specificity (Crickmore et al. (1998) *Micro-*

biol. Mol. Biol. Rev. 62:807-813). In the new classification, each toxin is assigned a unique name incorporating a primary rank (an Arabic number), a secondary rank (an uppercase letter), a tertiary rank (a lowercase letter), and a quaternary rank (another Arabic number). In the new classification, Roman numerals have been exchanged for Arabic numerals in the primary rank. Proteins with less than 45% sequence identity have different primary ranks, and the criteria for secondary and tertiary ranks are 78% and 95%, respectively.

[0007] The crystal protein does not exhibit insecticidal activity until it has been ingested and solubilized in the insect midgut. The ingested protoxin is hydrolyzed by proteases in the insect digestive tract to an active toxic molecule. (Höfte and Whiteley (1989) *Microbiol. Rev.* 53:242-255). This toxin binds to apical brush border receptors in the midgut of the target larvae and inserts into the apical membrane creating ion channels or pores, resulting in larval death.

[0008] Delta-endotoxins generally have five conserved sequence domains, and three conserved structural domains (see, for example, de Maagd et al. (2001) *Trends Genetics* 17:193-199). The first conserved structural domain consists of seven alpha helices and is involved in membrane insertion and pore formation. Domain II consists of three beta-sheets arranged in a "Greek key" configuration, and domain III consists of two antiparallel beta-sheets in "jelly-roll" formation (de Maagd et al., 2001, supra). Domains II and III are involved in receptor recognition and binding, and are therefore considered determinants of toxin specificity.

[0009] Aside from delta-endotoxins, there are several other known classes of pesticidal protein toxins. The VIP1/VIP2 toxins (see, for example, U.S. Pat. No. 5,770,696) are binary pesticidal toxins that exhibit strong activity on insects by a mechanism believed to involve receptor-mediated endocytosis followed by cellular toxification, similar to the mode of action of other binary ("A/B") toxins. A/B toxins such as VIP, C2, CDT, CST, or the *B. anthracis* edema and lethal toxins initially interact with target cells via a specific, receptor-mediated binding of "B" components as monomers. These monomers then form homoheptamers. The "B" heptamer-receptor complex then acts as a docking platform that subsequently binds and allows the translocation of an enzymatic "A" component(s) into the cytosol via receptor-mediated endocytosis. Once inside the cell's cytosol, "A" components inhibit normal cell function by, for example, ADP-ribosylation of G-actin, or increasing intracellular levels of cyclic AMP (cAMP). See Barth et al. (2004) *Microbiol Mol Biol Rev* 68:373-402.

[0010] The intensive use of *B. thuringiensis*-based insecticides has already given rise to resistance in field populations of the diamondback moth, *Plutella xylostella* (Ferré and Van Rie (2002) *Annu. Rev. Entomol.* 47:501-533). The most common mechanism of resistance is the reduction of binding of the toxin to its specific midgut receptor(s). This may also confer cross-resistance to other toxins that share the same receptor (Ferré and Van Rie (2002)).

SUMMARY OF INVENTION

[0011] Compositions and methods for conferring insect resistance to bacteria, plants, plant cells, tissues and seeds are provided. Compositions include nucleic acid molecules encoding sequences for delta-endotoxin polypeptides, vectors comprising those nucleic acid molecules, and host cells comprising the vectors. Compositions also include the polypeptide sequences of the endotoxin, and antibodies to

those polypeptides. The nucleotide sequences can be used in DNA constructs or expression cassettes for transformation and expression in organisms, including microorganisms and plants. The nucleotide or amino acid sequences may be synthetic sequences that have been designed for expression in an organism including, but not limited to, a microorganism or a plant. Compositions also comprise transformed bacteria, plants, plant cells, tissues, and seeds.

[0012] In particular, isolated nucleic acid molecules corresponding to delta-endotoxin nucleic acid sequences are provided. Additionally, amino acid sequences corresponding to the polynucleotides are encompassed. In particular, the present invention provides for an isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence shown in any of SEQ ID NO:4, 5, 6, 13, or 14, or a nucleotide sequence set forth in any of SEQ ID NO:1, 2, 3, 11, or 12, as well as variants and fragments thereof. Nucleotide sequences that are complementary to a nucleotide sequence of the invention, or that hybridize to a sequence of the invention are also encompassed.

[0013] The compositions and methods of the invention are useful for the production of organisms with insecticide resistance, specifically bacteria and plants. These organisms and compositions derived from them are desirable for agricultural purposes. The compositions of the invention are also useful for generating altered or improved delta-endotoxin proteins that have insecticidal activity, or for detecting the presence of delta-endotoxin proteins or nucleic acids in products or organisms.

[0014] The following embodiments are encompassed by the present invention:

[0015] 1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

[0016] a) the nucleotide sequence of any of SEQ ID NO:1, 2, 3, 11, or 12, or a complement thereof;

[0017] b) a nucleotide sequence that encodes a polypeptide comprising the amino acid sequence of any of SEQ ID NO:4, 5, 6, 13, or 14;

[0018] c) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 96% sequence identity to the amino acid sequence of SEQ ID NO:4 or 14, wherein said amino acid sequence has insecticidal activity;

[0019] d) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO:6, wherein said amino acid sequence has insecticidal activity; and

[0020] e) a nucleotide sequence encoding an insecticidal polypeptide that is a variant of SEQ ID NO:4, 5, 6, 13, or 14, wherein said variant is the result of one or more domain(s) of SEQ ID NO:4, 5, 6, 13, or 14 being exchanged with the corresponding domain(s) of SEQ ID NO:4, 5, 6, 13, or 14, wherein said nucleotide sequence encodes an insecticidal polypeptide.

[0021] 2. The isolated nucleic acid molecule of embodiment 1, wherein said nucleotide sequence is a synthetic sequence that has been designed for expression in a plant.

[0022] 3. The isolated nucleic acid molecule of embodiment 2, wherein said nucleotide sequence is selected from the group consisting of SEQ ID NO:15, 16, 17, and 18.

[0023] 4. An expression cassette comprising the nucleic acid molecule of embodiment 1.

[0024] 5. The expression cassette of embodiment 4, further comprising a nucleic acid molecule encoding a heterologous polypeptide.

[0025] 6. A host cell that contains the expression cassette of embodiment 4.

[0026] 7. The host cell of embodiment 6 that is a bacterial host cell.

[0027] 8. The host cell of embodiment 6 that is a plant cell.

[0028] 9. A transgenic plant comprising the host cell of embodiment 8.

[0029] 10. The transgenic plant of embodiment 9, wherein said plant is selected from the group consisting of maize, sorghum, wheat, cabbage, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape.

[0030] 11. An isolated polypeptide with insecticidal activity, selected from the group consisting of:

[0031] a) a polypeptide comprising the amino acid sequence of any of SEQ ID NO:4, 5, 6, 13, or 14;

[0032] b) a polypeptide comprising an amino acid sequence having at least 96% sequence identity to the amino acid sequence of SEQ ID NO:4 or 14, wherein said amino acid sequence has insecticidal activity;

[0033] c) a polypeptide comprising an amino acid sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO:6, wherein said amino acid sequence has insecticidal activity;

[0034] d) a polypeptide that is encoded by the nucleotide sequence of any of SEQ ID NO:1, 2, 3, 11, or 12; and

[0035] e) a polypeptide that is a variant of SEQ ID NO:4, 5, 6, 13, or 14, wherein said variant is the result of one or more domain(s) of SEQ ID NO:4, 5, 6, 13, or 14 being exchanged with the corresponding domain(s) of SEQ ID NO:4, 5, 6, 13, or 14, wherein said polypeptide has insecticidal activity.

[0036] 12. The polypeptide of embodiment 11 further comprising heterologous amino acid sequences.

[0037] 13. An antibody that selectively binds to the polypeptide of embodiment 11.

[0038] 14. A composition comprising the polypeptide of embodiment 11.

[0039] 15. The composition of embodiment 14, wherein said composition is selected from the group consisting of a powder, dust, pellet, granule, spray, emulsion, colloid, and solution.

[0040] 16. The composition of embodiment 14, wherein said composition is prepared by desiccation, lyophilization, homogenization, extraction, filtration, centrifugation, sedimentation, or concentration of a culture of *Bacillus thuringiensis* cells.

[0041] 17. The composition of embodiment 14, comprising from about 1% to about 99% by weight of said polypeptide.

[0042] 18. A method for controlling a lepidopteran or coleopteran pest population comprising contacting said population with an insecticidally-effective amount of the polypeptide of embodiment 11.

[0043] 19. A method for killing a lepidopteran or coleopteran pest, comprising contacting said pest with, or feeding to said pest, an insecticidally-effective amount of the polypeptide of embodiment 11.

[0044] 20. A method for producing a polypeptide with insecticidal activity, comprising culturing the host cell of embodiment 6 under conditions in which the nucleic acid molecule encoding the polypeptide is expressed.

[0045] 21. A plant having stably incorporated into its genome a DNA construct comprising a nucleotide sequence that encodes a protein having insecticidal activity, wherein said nucleotide sequence is selected from the group consisting of:

[0046] a) the nucleotide sequence of any of SEQ ID NO:1, 2, 3, 11, or 12;

[0047] b) a nucleotide sequence that encodes a polypeptide comprising the amino acid sequence of any of SEQ ID NO:4, 5, 6, 13, or 14;

[0048] c) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 96% sequence identity to the amino acid sequence of SEQ ID NO:4 or 14, wherein said amino acid sequence has insecticidal activity;

[0049] d) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO:6, wherein said amino acid sequence has insecticidal activity; and

[0050] e) a nucleotide sequence encoding an insecticidal polypeptide that is a variant of SEQ ID NO:4, 5, 6, 13, or 14, wherein said variant is the result of one or more domain(s) of SEQ ID NO:4, 5, 6, 13, or 14 being exchanged with the corresponding domain(s) of SEQ ID NO:4, 5, 6, 13, or 14, wherein said nucleotide sequence encodes an insecticidal polypeptide;

wherein said nucleotide sequence is operably linked to a promoter that drives expression of a coding sequence in a plant cell.

[0051] 22. The plant of embodiment 21, wherein said plant is a plant cell.

[0052] 23. A transgenic seed of the plant of embodiment 21, wherein said seed comprises a nucleotide sequence selected from the group consisting of:

[0053] a) the nucleotide sequence of any of SEQ ID NO:1, 2, 3, 11, or 12;

[0054] b) a nucleotide sequence that encodes a polypeptide comprising the amino acid sequence of any of SEQ ID NO:4, 5, 6, 13, or 14;

[0055] c) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 96% sequence identity to the amino acid sequence of SEQ ID NO:4 or 14, wherein said amino acid sequence has insecticidal activity;

[0056] d) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO:6, wherein said amino acid sequence has insecticidal activity; and

[0057] e) a nucleotide sequence encoding an insecticidal polypeptide that is a variant of SEQ ID NO:4, 5, 6, 13, or 14, wherein said variant is the result of one or more domain(s) of SEQ ID NO:4, 5, 6, 13, or 14 being exchanged with the corresponding domain(s) of SEQ ID NO:4, 5, 6, 13, or 14, wherein said nucleotide sequence encodes an insecticidal polypeptide.

[0058] 24. A method for protecting a plant from an insect pest, comprising introducing into said plant or cell thereof at least one expression vector comprising a nucleotide sequence that encodes an insecticidal polypeptide, wherein said nucleotide sequence is selected from the group consisting of:

[0059] a) the nucleotide sequence of any of SEQ ID NO:1, 2, 3, 11, or 12;

[0060] b) a nucleotide sequence that encodes a polypeptide comprising the amino acid sequence of any of SEQ ID NO:4, 5, 6, 13, or 14;

[0061] c) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 96% sequence identity to the amino acid sequence of SEQ ID NO:4 or 14, wherein said amino acid sequence has insecticidal activity;

[0062] d) a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO:6, wherein said amino acid sequence has insecticidal activity; and

[0063] e) a nucleotide sequence encoding an insecticidal polypeptide that is a variant of SEQ ID NO:4, 5, 6, 13, or 14, wherein said variant is the result of one or more domain(s) of SEQ ID NO:4, 5, 6, 13, or 14 being exchanged with the corresponding domain(s) of SEQ ID NO:4, 5, 6, 13, or 14, wherein said nucleotide sequence encodes an insecticidal polypeptide.

[0064] 25. The method of embodiment 24, wherein said plant produces an insecticidal polypeptide having insecticidal activity against a lepidopteran or coleopteran pest.

BRIEF DESCRIPTION OF THE FIGURES

[0065] FIGS. 1A and 1B show an alignment of AXMI-113 (SEQ ID NO:5), AXMI-005 (SEQ ID NO:4), and AXMI-115 (SEQ ID NO:6). The left and right arrows mark the boundaries of the C-terminal $\frac{1}{3}$ rd region of the proteins.

[0066] FIG. 2 depicts domains within AXMI-005 and AXMI-115 that are swapped to generate new toxins.

DETAILED DESCRIPTION

[0067] The present invention is drawn to compositions and methods for regulating insect resistance in organisms, particularly plants or plant cells. The methods involve transforming organisms with a nucleotide sequence encoding a delta-endotoxin protein of the invention. In particular, the nucleotide sequences of the invention are useful for preparing plants and microorganisms that possess insecticidal activity. Thus, transformed bacteria, plants, plant cells, plant tissues and seeds are provided. Compositions are delta-endotoxin nucleic acids and proteins of *Bacillus thuringiensis*. The sequences find use in the construction of expression vectors for subsequent transformation into organisms of interest, as probes for the isolation of other delta-endotoxin genes, and for the generation of altered insecticidal proteins by methods known in the art, such as domain swapping or DNA shuffling. See, for example, Table 2 and FIG. 2. The proteins find use in controlling or killing lepidopteran, coleopteran, and other insect populations, and for producing compositions with insecticidal activity.

[0068] By "delta-endotoxin" is intended a toxin from *Bacillus thuringiensis* that has toxic activity against one or more pests, including, but not limited to, members of the Lepidoptera, Diptera, and Coleoptera orders, or a protein that has homology to such a protein. In some cases, delta-endotoxin proteins have been isolated from other organisms, including *Clostridium bifermentans* and *Paenibacillus popilliae*. Delta-endotoxin proteins include amino acid sequences deduced from the full-length nucleotide sequences disclosed herein, and amino acid sequences that are shorter than the full-length sequences, either due to the use of an

alternate downstream start site, or due to processing that produces a shorter protein having insecticidal activity. Processing may occur in the organism the protein is expressed in, or in the pest after ingestion of the protein.

[0069] Delta-endotoxins include proteins identified as cry1 through cry43, cyt1 and cyt2, and Cyt-like toxin. There are currently over 250 known species of delta-endotoxins with a wide range of specificities and toxicities. For an expansive list see Crickmore et al. (1998), *Microbiol. Mol. Biol. Rev.* 62:807-813, and for regular updates see Crickmore et al. (2003) "Bacillus thuringiensis toxin nomenclature," at www.biols.susx.ac.uk/Home/Neil_Crickmore/Bt/index.

[0070] Provided herein are novel isolated nucleotide sequences that confer insecticidal activity. Also provided are the amino acid sequences of the delta-endotoxin proteins. The protein resulting from translation of this gene allows cells to control or kill insects that ingest it.

Isolated Nucleic Acid Molecules, and Variants and Fragments Thereof

[0071] One aspect of the invention pertains to isolated or recombinant nucleic acid molecules comprising nucleotide sequences encoding delta-endotoxin proteins and polypeptides or biologically active portions thereof, as well as nucleic acid molecules sufficient for use as hybridization probes to identify delta-endotoxin encoding nucleic acids. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., recombinant DNA, cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

[0072] An "isolated" or "purified" nucleic acid molecule or protein, or biologically active portion thereof, is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For purposes of the invention, "isolated" when used to refer to nucleic acid molecules excludes isolated chromosomes. For example, in various embodiments, the isolated delta-endotoxin encoding nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. A delta-endotoxin protein that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, or 5% (by dry weight) of non-delta-endotoxin protein (also referred to herein as a "contaminating protein").

[0073] Nucleotide sequences encoding the proteins of the present invention include the sequence set forth in SEQ ID NO:1, 2, 3, 11, or 12, and variants, fragments, and complements thereof. By "complement" is intended a nucleotide sequence that is sufficiently complementary to a given nucleotide sequence such that it can hybridize to the given nucleotide sequence to thereby form a stable duplex. The corresponding amino acid sequence for the delta-endotoxin protein encoded by this nucleotide sequence are set forth in SEQ ID NO:4, 5, 6, 13, or 14.

[0074] Nucleic acid molecules that are fragments of these delta-endotoxin encoding nucleotide sequences are also encompassed by the present invention. By "fragment" is intended a portion of the nucleotide sequence encoding a delta-endotoxin protein. A fragment of a nucleotide sequence may encode a biologically active portion of a delta-endotoxin protein, or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. Nucleic acid molecules that are fragments of a delta-endotoxin nucleotide sequence comprise at least about 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, 2000, 2050, 2100, 2150, 2200, 2250, 2300, 2350, 2400, 2450, 2500, 2550, 2600, 2650, 2700, 2750, 2800, 2850, 2900, 2950, 3000, 3050, 3100, 3150, 3200, 3250, 3300, 3350 contiguous nucleotides, or up to the number of nucleotides present in a full-length delta-endotoxin encoding nucleotide sequence disclosed herein depending upon the intended use. By "contiguous" nucleotides is intended nucleotide residues that are immediately adjacent to one another. Fragments of the nucleotide sequences of the present invention will encode protein fragments that retain the biological activity of the delta-endotoxin protein and, hence, retain insecticidal activity. By "retains activity" is intended that the fragment will have at least about 30%, at least about 50%, at least about 70%, 80%, 90%, 95% or higher of the insecticidal activity of the delta-endotoxin protein. Methods for measuring insecticidal activity are well known in the art. See, for example, Czaplak and Lang (1990) *J. Econ. Entomol.* 83:2480-2485; Andrews et al. (1988) *Biochem. J.* 252:199-206; Marrone et al. (1985) *J. of Economic Entomology* 78:290-293; and U.S. Pat. No. 5,743,477, all of which are herein incorporated by reference in their entirety.

[0075] A fragment of a delta-endotoxin encoding nucleotide sequence that encodes a biologically active portion of a protein of the invention will encode at least about 15, 25, 30, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100 contiguous amino acids, or up to the total number of amino acids present in a full-length delta-endotoxin protein of the invention.

[0076] Preferred delta-endotoxin proteins of the present invention are encoded by a nucleotide sequence sufficiently identical to the nucleotide sequence of SEQ ID NO:1, 2, 3, 11, or 12. By "sufficiently identical" is intended an amino acid or nucleotide sequence that has at least about 60% or 65% sequence identity, about 70% or 75% sequence identity, about 80% or 85% sequence identity, about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater sequence identity compared to a reference sequence using one of the alignment programs described herein using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like.

[0077] To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., percent identity=number of identical positions/total number of positions (e.g., overlapping positions)×100). In one embodiment, the two sequences are the same length. In another embodi-

ment, the comparison is across the entirety of the reference sequence (e.g., across the entirety of one of SEQ ID NO:1, 2, 3, 11, or 12, or across the entirety of one of SEQ ID NO:4, 5, 6, 13, or 14). The percent identity between two sequences can be determined using techniques similar to those described below, with or without allowing gaps. In calculating percent identity, typically exact matches are counted.

[0078] The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A nonlimiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the BLASTN and BLASTX programs of Altschul et al. (1990) *J. Mol. Biol.* 215:403. BLAST nucleotide searches can be performed with the BLASTN program, score=100, wordlength=12, to obtain nucleotide sequences homologous to delta-endotoxin-like nucleic acid molecules of the invention. BLAST protein searches can be performed with the BLASTX program, score=50, wordlength=3, to obtain amino acid sequences homologous to delta-endotoxin protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-Blast can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) *supra*. When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., BLASTX and BLASTN) can be used. Alignment may also be performed manually by inspection.

[0079] Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the ClustalW algorithm (Higgins et al. (1994) *Nucleic Acids Res.* 22:4673-4680). ClustalW compares sequences and aligns the entirety of the amino acid or DNA sequence, and thus can provide data about the sequence conservation of the entire amino acid sequence. The ClustalW algorithm is used in several commercially available DNA/amino acid analysis software packages, such as the ALIGNX module of the Vector NTI Program Suite (Invitrogen Corporation, Carlsbad, Calif.). After alignment of amino acid sequences with ClustalW, the percent amino acid identity can be assessed. A non-limiting example of a software program useful for analysis of ClustalW alignments is GENEDOC™. GENEDOC™ (Karl Nicholas) allows assessment of amino acid (or DNA) similarity and identity between multiple proteins. Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN program (version 2.0), which is part of the GCG Wisconsin Genetics Software Package, Version 10 (available from Accelrys, Inc., 9685 Scranton Rd., San Diego, Calif., USA). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

[0080] Unless otherwise stated, GAP Version 10, which uses the algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48(3):443-453, will be used to determine sequence identity or similarity using the following parameters: % identity and % similarity for a nucleotide sequence using GAP Weight of 50 and Length Weight of 3, and the nws gapdna.cmp scor-

ing matrix; % identity or % similarity for an amino acid sequence using GAP weight of 8 and length weight of 2, and the BLOSUM62 scoring program. Equivalent programs may also be used. By “equivalent program” is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by GAP Version 10.

[0081] The invention also encompasses variant nucleic acid molecules. “Variants” of the delta-endotoxin encoding nucleotide sequences include those sequences that encode the delta-endotoxin proteins disclosed herein but that differ conservatively because of the degeneracy of the genetic code as well as those that are sufficiently identical as discussed above. Naturally occurring allelic variants can be identified with the use of well-known molecular biology techniques, such as polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences that have been generated, for example, by using site-directed mutagenesis but which still encode the delta-endotoxin proteins disclosed in the present invention as discussed below. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the native protein, that is, retaining insecticidal activity. By “retains activity” is intended that the variant will have at least about 30%, at least about 50%, at least about 70%, or at least about 80% of the insecticidal activity of the native protein. Methods for measuring insecticidal activity are well known in the art. See, for example, Czaplá and Lang (1990) *J. Econ. Entomol.* 83: 2480-2485; Andrews et al. (1988) *Biochem. J.* 252:199-206; Marrone et al. (1985) *J. of Economic Entomology* 78:290-293; and U.S. Pat. No. 5,743,477, all of which are herein incorporated by reference in their entirety.

[0082] The skilled artisan will further appreciate that changes can be introduced by mutation of the nucleotide sequences of the invention thereby leading to changes in the amino acid sequence of the encoded delta-endotoxin proteins, without altering the biological activity of the proteins. Thus, variant isolated nucleic acid molecules can be created by introducing one or more nucleotide substitutions, additions, or deletions into the corresponding nucleotide sequence disclosed herein, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Such variant nucleotide sequences are also encompassed by the present invention.

[0083] For example, conservative amino acid substitutions may be made at one or more predicted, nonessential amino acid residues. A “nonessential” amino acid residue is a residue that can be altered from the wild-type sequence of a delta-endotoxin protein without altering the biological activity, whereas an “essential” amino acid residue is required for biological activity. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine,

threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine).

[0084] Delta-endotoxins generally have five conserved sequence domains, and three conserved structural domains (see, for example, de Maagd et al. (2001) *Trends Genetics* 17:193-199). The first conserved structural domain consists of seven alpha helices and is involved in membrane insertion and pore formation. Domain II consists of three beta-sheets arranged in a Greek key configuration, and domain III consists of two antiparallel beta-sheets in "jelly-roll" formation (de Maagd et al., 2001, supra). Domains II and III are involved in receptor recognition and binding, and are therefore considered determinants of toxin specificity.

[0085] Amino acid substitutions may be made in nonconserved regions that retain function. In general, such substitutions would not be made for conserved amino acid residues, or for amino acid residues residing within a conserved motif, where such residues are essential for protein activity. Examples of residues that are conserved and that may be essential for protein activity include, for example, residues that are identical between all proteins contained in an alignment of the amino acid sequences of the present invention and known delta-endotoxin sequences. Examples of residues that are conserved but that may allow conservative amino acid substitutions and still retain activity include, for example, residues that have only conservative substitutions between all proteins contained in an alignment of the amino acid sequences of the present invention and known delta-endotoxin sequences. However, one of skill in the art would understand that functional variants may have minor conserved or nonconserved alterations in the conserved residues.

[0086] Alternatively, variant nucleotide sequences can be made by introducing mutations randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for ability to confer delta-endotoxin activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed recombinantly, and the activity of the protein can be determined using standard assay techniques.

[0087] Using methods such as PCR, hybridization, and the like corresponding delta-endotoxin sequences can be identified, such sequences having substantial identity to the sequences of the invention. See, for example, Sambrook and Russell (2001) *Molecular Cloning: A Laboratory Manual*. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.) and Innis, et al. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, NY).

[0088] In a hybridization method, all or part of the delta-endotoxin nucleotide sequence can be used to screen cDNA or genomic libraries. Methods for construction of such cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook and Russell, 2001, supra. The so-called hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments, or other oligonucleotides, and may be labeled with a detectable group such as ^{32}P , or any other detectable marker, such as other radioisotopes, a fluorescent compound, an enzyme, or an enzyme co-factor. Probes for hybridization can be made by labeling synthetic oligonucleotides based on the known delta-endotoxin-encoding nucleotide sequence disclosed herein. Degenerate primers designed on the basis of conserved nucleotides or amino

acid residues in the nucleotide sequence or encoded amino acid sequence can additionally be used. The probe typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, at least about 25, at least about 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, or 400 consecutive nucleotides of delta-endotoxin encoding nucleotide sequence of the invention or a fragment or variant thereof. Methods for the preparation of probes for hybridization are generally known in the art and are disclosed in Sambrook and Russell, 2001, supra herein incorporated by reference.

[0089] For example, an entire delta-endotoxin sequence disclosed herein, or one or more portions thereof, may be used as a probe capable of specifically hybridizing to corresponding delta-endotoxin-like sequences and messenger RNAs. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique and are preferably at least about 10 nucleotides in length, or at least about 20 nucleotides in length. Such probes may be used to amplify corresponding delta-endotoxin sequences from a chosen organism by PCR. This technique may be used to isolate additional coding sequences from a desired organism or as a diagnostic assay to determine the presence of coding sequences in an organism. Hybridization techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, for example, Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.).

[0090] Hybridization of such sequences may be carried out under stringent conditions. By "stringent conditions" or "stringent hybridization conditions" is intended conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in depending upon circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

[0091] Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55° C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37° C., and a wash in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60 to 65° C. Optionally, wash buffers may comprise about 0.1% to about 1% SDS. Duration of hybridization is generally less than about 24 hours, usually about 4 to about 12 hours.

[0092] Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284: $T_m = 81.5^\circ \text{C} + 16.6 (\log M) + 0.41 (\% \text{ GC}) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1°C . for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $\geq 90\%$ identity are sought, the T_m can be decreased 10°C . Generally, stringent conditions are selected to be about 5°C . lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at $1, 2, 3$, or 4°C . lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at $6, 7, 8, 9$, or 10°C . lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at $11, 12, 13, 14, 15$, or 20°C . lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C . (aqueous solution) or 32°C . (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel et al., eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.).

Isolated Proteins and Variants and Fragments Thereof

[0093] Delta-endotoxin proteins are also encompassed within the present invention. By "delta-endotoxin protein" is intended a protein having the amino acid sequence set forth in SEQ ID NO:4, 5, 6, 13, or 14. Fragments, biologically active portions, and variants thereof are also provided, and may be used to practice the methods of the present invention.

[0094] "Fragments" or "biologically active portions" include polypeptide fragments comprising amino acid sequences sufficiently identical to the amino acid sequence set forth in any of SEQ ID NO:4, 5, 6, 13, or 14 and that exhibit insecticidal activity. A biologically active portion of a delta-endotoxin protein can be a polypeptide that is, for example, 10, 25, 50, 100 or more amino acids in length. Such biologically active portions can be prepared by recombinant techniques and evaluated for insecticidal activity. Methods for measuring insecticidal activity are well known in the art. See, for example, Czapla and Lang (1990) *J. Econ. Entomol.* 83:2480-2485; Andrews et al. (1988) *Biochem. J.* 252:199-206; Marrone et al. (1985) *J. of Economic Entomology*

78:290-293; and U.S. Pat. No. 5,743,477, all of which are herein incorporated by reference in their entirety. As used here, a fragment comprises at least 8 contiguous amino acids of SEQ ID NO:4, 5, 6, 13, or 14. The invention encompasses other fragments, however, such as any fragment in the protein greater than about 10, 20, 30, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, or 1300 amino acids.

[0095] By "variants" is intended proteins or polypeptides having an amino acid sequence that is at least about 60%, 65%, about 70%, 75%, about 80%, 85%, about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of any of SEQ ID NO:4, 5, 6, 13, or 14. Variants also include polypeptides encoded by a nucleic acid molecule that hybridizes to the nucleic acid molecule of SEQ ID NO:1, 2, 3, 11, or 12, or a complement thereof, under stringent conditions. Variants include polypeptides that differ in amino acid sequence due to mutagenesis. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the native protein, that is, retaining insecticidal activity. Methods for measuring insecticidal activity are well known in the art. See, for example, Czapla and Lang (1990) *J. Econ. Entomol.* 83:2480-2485; Andrews et al. (1988) *Biochem. J.* 252:199-206; Marrone et al. (1985) *J. of Economic Entomology* 78:290-293; and U.S. Pat. No. 5,743,477, all of which are herein incorporated by reference in their entirety.

[0096] Bacterial genes, such as the axmi genes of this invention, quite often possess multiple methionine initiation codons in proximity to the start of the open reading frame. Often, translation initiation at one or more of these start codons will lead to generation of a functional protein. These start codons can include ATG codons. However, bacteria such as *Bacillus* sp. also recognize the codon GTG as a start codon, and proteins that initiate translation at GTG codons contain a methionine at the first amino acid. Furthermore, it is not often determined a priori which of these codons are used naturally in the bacterium. Thus, it is understood that use of one of the alternate methionine codons may also lead to generation of delta-endotoxin proteins that encode insecticidal activity. These delta-endotoxin proteins are encompassed in the present invention and may be used in the methods of the present invention.

[0097] Antibodies to the polypeptides of the present invention, or to variants or fragments thereof, are also encompassed. Methods for producing antibodies are well known in the art (see, for example, Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.; U.S. Pat. No. 4,196,265).

Altered or Improved Variants

[0098] It is recognized that DNA sequences of a delta-endotoxin may be altered by various methods, and that these alterations may result in DNA sequences encoding proteins with amino acid sequences different than that encoded by a delta-endotoxin of the present invention. This protein may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions of one or more amino acids of SEQ ID NO:4, 5, 6, 13, or 14, including up to about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 15, about 20, about 25, about 30, about 35, about 40, about 45, about 50, about 55, about 60, about 65, about 70, about 75, about 80, about 85, about 90, about 100,

about 105, about 110, about 115, about 120, about 125, about 130 or more amino acid substitutions, deletions or insertions.

[0099] Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of a delta-endotoxin protein can be prepared by mutations in the DNA. This may also be accomplished by one of several forms of mutagenesis and/or in directed evolution. In some aspects, the changes encoded in the amino acid sequence will not substantially affect the function of the protein. Such variants will possess the desired insecticidal activity. However, it is understood that the ability of a delta-endotoxin to confer insecticidal activity may be improved by the use of such techniques upon the compositions of this invention. For example, one may express a delta-endotoxin in host cells that exhibit high rates of base misincorporation during DNA replication, such as XL-1 Red (Stratagene). After propagation in such strains, one can isolate the delta-endotoxin DNA (for example by preparing plasmid DNA, or by amplifying by PCR and cloning the resulting PCR fragment into a vector), culture the delta-endotoxin mutations in a non-mutagenic strain, and identify mutated delta-endotoxin genes with insecticidal activity, for example by performing an assay to test for insecticidal activity. Generally, the protein is mixed and used in feeding assays. See, for example Marrone et al. (1985) *J. of Economic Entomology* 78:290-293. Such assays can include contacting plants with one or more insects and determining the plant's ability to survive and/or cause the death of the insects. Examples of mutations that result in increased toxicity are found in Schnepf et al. (1998) *Microbiol. Mol. Biol. Rev.* 62:775-806.

[0100] Alternatively, alterations may be made to the protein sequence of many proteins at the amino or carboxy terminus without substantially affecting activity. This can include insertions, deletions, or alterations introduced by modern molecular methods, such as PCR, including PCR amplifications that alter or extend the protein coding sequence by virtue of inclusion of amino acid encoding sequences in the oligonucleotides utilized in the PCR amplification. Alternatively, the protein sequences added can include entire protein-coding sequences, such as those used commonly in the art to generate protein fusions. Such fusion proteins are often used to (1) increase expression of a protein of interest (2) introduce a binding domain, enzymatic activity, or epitope to facilitate either protein purification, protein detection, or other experimental uses known in the art (3) target secretion or translation of a protein to a subcellular organelle, such as the periplasmic space of Gram-negative bacteria, or the endoplasmic reticulum of eukaryotic cells, the latter of which often results in glycosylation of the protein.

[0101] Variant nucleotide and amino acid sequences of the present invention also encompass sequences derived from mutagenic and recombinogenic procedures such as DNA shuffling. With such a procedure, one or more different delta-endotoxin protein coding regions can be used to create a new delta-endotoxin protein possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined in vitro or in vivo. For example, using this approach, sequence motifs encoding a domain of interest may be shuffled between a delta-endotoxin gene of the invention and other known delta-endotoxin genes to obtain a new gene coding for a protein with an improved property of interest, such as an

increased insecticidal activity. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Cramer et al. (1997) *Nature Biotech.* 15:436-438; Moore et al. (1997) *J. Mol. Biol.* 272:336-347; Zhang et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Cramer et al. (1998) *Nature* 391:288-291; and U.S. Pat. Nos. 5,605,793 and 5,837,458.

[0102] Domain swapping or shuffling is another mechanism for generating altered delta-endotoxin proteins. Domains II and III may be swapped between delta-endotoxin proteins, resulting in hybrid or chimeric toxins with improved insecticidal activity or target spectrum. Methods for generating recombinant proteins and testing them for insecticidal activity are well known in the art (see, for example, Naimov et al. (2001) *Appl. Environ. Microbiol.* 67:5328-5330; de Maagd et al. (1996) *Appl. Environ. Microbiol.* 62:1537-1543; Ge et al. (1991) *J. Biol. Chem.* 266:17954-17958; Schnepf et al. (1990) *J. Biol. Chem.* 265:20923-20930; Rang et al. (1999) *Appl. Environ. Microbiol.* 65:2918-2925).

Vectors

[0103] A delta-endotoxin sequence of the invention may be provided in an expression cassette for expression in a plant of interest. By "plant expression cassette" is intended a DNA construct that is capable of resulting in the expression of a protein from an open reading frame in a plant cell. Typically these contain a promoter and a coding sequence. Often, such constructs will also contain a 3' untranslated region. Such constructs may contain a "signal sequence" or "leader sequence" to facilitate co-translational or post-translational transport of the peptide to certain intracellular structures such as the chloroplast (or other plastid), endoplasmic reticulum, or Golgi apparatus.

[0104] By "signal sequence" is intended a sequence that is known or suspected to result in cotranslational or post-translational peptide transport across the cell membrane. In eukaryotes, this typically involves secretion into the Golgi apparatus, with some resulting glycosylation. By "leader sequence" is intended any sequence that when translated, results in an amino acid sequence sufficient to trigger cotranslational transport of the peptide chain to a sub-cellular organelle. Thus, this includes leader sequences targeting transport and/or glycosylation by passage into the endoplasmic reticulum, passage to vacuoles, plastids including chloroplasts, mitochondria, and the like.

[0105] By "plant transformation vector" is intended a DNA molecule that is necessary for efficient transformation of a plant cell. Such a molecule may consist of one or more plant expression cassettes, and may be organized into more than one "vector" DNA molecule. For example, binary vectors are plant transformation vectors that utilize two non-contiguous DNA vectors to encode all requisite cis- and trans-acting functions for transformation of plant cells (Hellens and Mulineaux (2000) *Trends in Plant Science* 5:446-451). "Vector" refers to a nucleic acid construct designed for transfer between different host cells. "Expression vector" refers to a vector that has the ability to incorporate, integrate and express heterologous DNA sequences or fragments in a foreign cell. The cassette will include 5' and 3' regulatory sequences operably linked to a sequence of the invention. By "operably linked" is intended a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence cor-

responding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes.

[0106] “Promoter” refers to a nucleic acid sequence that functions to direct transcription of a downstream coding sequence. The promoter together with other transcriptional and translational regulatory nucleic acid sequences (also termed “control sequences”) are necessary for the expression of a DNA sequence of interest.

[0107] Such an expression cassette is provided with a plurality of restriction sites for insertion of the delta-endotoxin sequence to be under the transcriptional regulation of the regulatory regions.

[0108] The expression cassette will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region (i.e., a promoter), a DNA sequence of the invention, and a translational and transcriptional termination region (i.e., termination region) functional in plants. The promoter may be native, or analogous, or foreign or heterologous, to the plant host and/or to the DNA sequence of the invention. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. Where the promoter is “native” or “homologous” to the plant host, it is intended that the promoter is found in the native plant into which the promoter is introduced. Where the promoter is “foreign” or “heterologous” to the DNA sequence of the invention, it is intended that the promoter is not the native or naturally occurring promoter for the operably linked DNA sequence of the invention.

[0109] The termination region may be native with the transcriptional initiation region, may be native with the operably linked DNA sequence of interest, may be native with the plant host, or may be derived from another source (i.e., foreign or heterologous to the promoter, the DNA sequence of interest, the plant host, or any combination thereof). Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also Guerineau et al. (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon et al. (1991) *Genes Dev.* 5:141-149; Mogen et al. (1990) *Plant Cell* 2:1261-1272; Munroe et al. (1990) *Gene* 91:151-158; Ballas et al. (1989) *Nucleic Acids Res.* 17:7891-7903; and Joshi et al. (1987) *Nucleic Acid Res.* 15:9627-9639.

[0110] Where appropriate, the gene(s) may be optimized for increased expression in the transformed host cell. That is, the genes can be synthesized using host cell-preferred codons for improved expression, or may be synthesized using codons at a host-preferred codon usage frequency. Generally, the GC content of the gene will be increased. See, for example, Campbell and Gowri (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Pat. Nos. 5,380,831, and 5,436,391, and Murray et al. (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

[0111] In one embodiment, the delta-endotoxin is targeted to the chloroplast for expression. In this manner, where the delta-endotoxin is not directly inserted into the chloroplast,

the expression cassette will additionally contain a nucleic acid encoding a transit peptide to direct the delta-endotoxin to the chloroplasts. Such transit peptides are known in the art. See, for example, Von Heijne et al. (1991) *Plant Mol. Biol. Rep.* 9:104-126; Clark et al. (1989) *J. Biol. Chem.* 264:17544-17550; Della-Cioppa et al. (1987) *Plant Physiol.* 84:965-968; Romer et al. (1993) *Biochem. Biophys. Res. Commun.* 196:1414-1421; and Shah et al. (1986) *Science* 233:478-481.

[0112] The delta-endotoxin gene to be targeted to the chloroplast may be optimized for expression in the chloroplast to account for differences in codon usage between the plant nucleus and this organelle. In this manner, the nucleic acids of interest may be synthesized using chloroplast-preferred codons. See, for example, U.S. Pat. No. 5,380,831, herein incorporated by reference.

Plant Transformation

[0113] Methods of the invention involve introducing a nucleotide construct into a plant. By “introducing” is intended to present to the plant the nucleotide construct in such a manner that the construct gains access to the interior of a cell of the plant. The methods of the invention do not require that a particular method for introducing a nucleotide construct to a plant is used, only that the nucleotide construct gains access to the interior of at least one cell of the plant. Methods for introducing nucleotide constructs into plants are known in the art including, but not limited to, stable transformation methods, transient transformation methods, and virus-mediated methods.

[0114] By “plant” is intended whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, propagules, embryos and progeny of the same. Plant cells can be differentiated or undifferentiated (e.g. callus, suspension culture cells, protoplasts, leaf cells, root cells, phloem cells, pollen).

[0115] “Transgenic plants” or “transformed plants” or “stably transformed” plants or cells or tissues refers to plants that have incorporated or integrated exogenous nucleic acid sequences or DNA fragments into the plant cell. These nucleic acid sequences include those that are exogenous, or not present in the untransformed plant cell, as well as those that may be endogenous, or present in the untransformed plant cell. “Heterologous” generally refers to the nucleic acid sequences that are not endogenous to the cell or part of the native genome in which they are present, and have been added to the cell by infection, transfection, microinjection, electroporation, microprojection, or the like.

[0116] Transformation of plant cells can be accomplished by one of several techniques known in the art. The delta-endotoxin gene of the invention may be modified to obtain or enhance expression in plant cells. Typically a construct that expresses such a protein would contain a promoter to drive transcription of the gene, as well as a 3' untranslated region to allow transcription termination and polyadenylation. The organization of such constructs is well known in the art. In some instances, it may be useful to engineer the gene such that the resulting peptide is secreted, or otherwise targeted within the plant cell. For example, the gene can be engineered to contain a signal peptide to facilitate transfer of the peptide to the endoplasmic reticulum. It may also be preferable to engineer the plant expression cassette to contain an intron, such that mRNA processing of the intron is required for expression.

[0117] Typically this “plant expression cassette” will be inserted into a “plant transformation vector”. This plant trans-

formation vector may be comprised of one or more DNA vectors needed for achieving plant transformation. For example, it is a common practice in the art to utilize plant transformation vectors that are comprised of more than one contiguous DNA segment. These vectors are often referred to in the art as “binary vectors”. Binary vectors as well as vectors with helper plasmids are most often used for *Agrobacterium*-mediated transformation, where the size and complexity of DNA segments needed to achieve efficient transformation is quite large, and it is advantageous to separate functions onto separate DNA molecules. Binary vectors typically contain a plasmid vector that contains the cis-acting sequences required for T-DNA transfer (such as left border and right border), a selectable marker that is engineered to be capable of expression in a plant cell, and a “gene of interest” (a gene engineered to be capable of expression in a plant cell for which generation of transgenic plants is desired). Also present on this plasmid vector are sequences required for bacterial replication. The cis-acting sequences are arranged in a fashion to allow efficient transfer into plant cells and expression therein. For example, the selectable marker gene and the delta-endotoxin are located between the left and right borders. Often a second plasmid vector contains the trans-acting factors that mediate T-DNA transfer from *Agrobacterium* to plant cells. This plasmid often contains the virulence functions (Vir genes) that allow infection of plant cells by *Agrobacterium*, and transfer of DNA by cleavage at border sequences and vir-mediated DNA transfer, as is understood in the art (Hellens and Mullineaux (2000) *Trends in Plant Science* 5:446-451). Several types of *Agrobacterium* strains (e.g. LBA4404, GV3101, EHA101, EHA105, etc.) can be used for plant transformation. The second plasmid vector is not necessary for transforming the plants by other methods such as microprojection, microinjection, electroporation, polyethylene glycol, etc.

[0118] In general, plant transformation methods involve transferring heterologous DNA into target plant cells (e.g. immature or mature embryos, suspension cultures, undifferentiated callus, protoplasts, etc.), followed by applying a maximum threshold level of appropriate selection (depending on the selectable marker gene) to recover the transformed plant cells from a group of untransformed cell mass. Explants are typically transferred to a fresh supply of the same medium and cultured routinely. Subsequently, the transformed cells are differentiated into shoots after placing on regeneration medium supplemented with a maximum threshold level of selecting agent. The shoots are then transferred to a selective rooting medium for recovering rooted shoot or plantlet. The transgenic plantlet then grows into a mature plant and produces fertile seeds (e.g. Hiei et al. (1994) *The Plant Journal* 6:271-282; Ishida et al. (1996) *Nature Biotechnology* 14:745-750). Explants are typically transferred to a fresh supply of the same medium and cultured routinely. A general description of the techniques and methods for generating transgenic plants are found in Ayres and Park (1994) *Critical Reviews in Plant Science* 13:219-239 and Bommineni and Jauhar (1997) *Maydica* 42:107-120. Since the transformed material contains many cells; both transformed and non-transformed cells are present in any piece of subjected target callus or tissue or group of cells. The ability to kill non-transformed cells and allow transformed cells to proliferate results in transformed plant cultures. Often, the ability to remove non-transformed cells is a limitation to rapid recovery of transformed plant cells and successful generation of transgenic plants.

[0119] Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. Generation of transgenic plants may be performed by one of several methods, including, but not limited to, microinjection, electroporation, direct gene transfer, introduction of heterologous DNA by *Agrobacterium* into plant cells (*Agrobacterium*-mediated transformation), bombardment of plant cells with heterologous foreign DNA adhered to particles, ballistic particle acceleration, aerosol beam transformation (U.S. Published Application No. 20010026941; U.S. Pat. No. 4,945,050; International Publication No. WO 91/00915; U.S. Published Application No. 2002015066), Lec1 transformation, and various other non-particle direct-mediated methods to transfer DNA.

[0120] Methods for transformation of chloroplasts are known in the art. See, for example, Svab et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530; Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917; Svab and Maliga (1993) *EMBO J.* 12:601-606. The method relies on particle gun delivery of DNA containing a selectable marker and targeting of the DNA to the plastid genome through homologous recombination. Additionally, plastid transformation can be accomplished by transactivation of a silent plastid-borne transgene by tissue-preferred expression of a nuclear-encoded and plastid-directed RNA polymerase. Such a system has been reported in McBride et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:7301-7305.

[0121] Following integration of heterologous foreign DNA into plant cells, one then applies a maximum threshold level of appropriate selection in the medium to kill the untransformed cells and separate and proliferate the putatively transformed cells that survive from this selection treatment by transferring regularly to a fresh medium. By continuous passage and challenge with appropriate selection, one identifies and proliferates the cells that are transformed with the plasmid vector. Molecular and biochemical methods can then be used to confirm the presence of the integrated heterologous gene of interest into the genome of the transgenic plant.

[0122] The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick et al. (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the desired phenotypic characteristic has been achieved. In this manner, the present invention provides transformed seed (also referred to as “transgenic seed”) having a nucleotide construct of the invention, for example, an expression cassette of the invention, stably incorporated into their genome.

Evaluation of Plant Transformation

[0123] Following introduction of heterologous foreign DNA into plant cells, the transformation or integration of heterologous gene in the plant genome is confirmed by various methods such as analysis of nucleic acids, proteins and metabolites associated with the integrated gene.

[0124] PCR analysis is a rapid method to screen transformed cells, tissue or shoots for the presence of incorporated gene at the earlier stage before transplanting into the soil

(Sambrook and Russell (2001) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). PCR is carried out using oligonucleotide primers specific to the gene of interest or *Agrobacterium* vector background, etc.

[0125] Plant transformation may be confirmed by Southern blot analysis of genomic DNA (Sambrook and Russell, 2001, supra). In general, total DNA is extracted from the transformant, digested with appropriate restriction enzymes, resolved in an agarose gel and transferred to a nitrocellulose or nylon membrane. The membrane or "blot" is then probed with, for example, radiolabeled ³²P target DNA fragment to confirm the integration of introduced gene into the plant genome according to standard techniques (Sambrook and Russell, 2001, supra).

[0126] In Northern blot analysis, RNA is isolated from specific tissues of transformant, fractionated in a formaldehyde agarose gel, and blotted onto a nylon filter according to standard procedures that are routinely used in the art (Sambrook and Russell, 2001, supra). Expression of RNA encoded by the delta-endotoxin is then tested by hybridizing the filter to a radioactive probe derived from a delta-endotoxin, by methods known in the art (Sambrook and Russell, 2001, supra).

[0127] Western blot, biochemical assays and the like may be carried out on the transgenic plants to confirm the presence of protein encoded by the delta-endotoxin gene by standard procedures (Sambrook and Russell, 2001, supra) using antibodies that bind to one or more epitopes present on the delta-endotoxin protein.

Insecticidal Activity in Plants

[0128] In another aspect of the invention, one may generate transgenic plants expressing a delta-endotoxin that has insecticidal activity. Methods described above by way of example may be utilized to generate transgenic plants, but the manner in which the transgenic plant cells are generated is not critical to this invention. Methods known or described in the art such as *Agrobacterium*-mediated transformation, biolistic transformation, and non-particle-mediated methods may be used at the discretion of the experimenter. Plants expressing a delta-endotoxin may be isolated by common methods described in the art, for example by transformation of callus, selection of transformed callus, and regeneration of fertile plants from such transgenic callus. In such process, one may use any gene as a selectable marker so long as its expression in plant cells confers ability to identify or select for transformed cells.

[0129] A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, the aminoglycoside G418, hygromycin, or the like. Other genes that encode a product involved in chloroplast metabolism may also be used as selectable markers. For example, genes that provide resistance to plant herbicides such as glyphosate, bromoxynil, or imidazolinone may find particular use. Such genes have been reported (Stalker et al. (1985) *J. Biol. Chem.* 263:6310-6314 (bromoxynil resistance nitrilase gene); and Sathasivan et al. (1990) *Nucl. Acids Res.* 18:2188 (AHAS imidazolinone resistance gene). Additionally, the genes disclosed herein are useful as markers to assess transformation of bacterial or plant cells. Methods for detecting the presence of a transgene in a plant, plant organ (e.g., leaves, stems, roots, etc.), seed, plant cell, propagule, embryo or progeny of the

same are well known in the art. In one embodiment, the presence of the transgene is detected by testing for insecticidal activity.

[0130] Fertile plants expressing a delta-endotoxin may be tested for insecticidal activity, and the plants showing optimal activity selected for further breeding. Methods are available in the art to assay for insect activity. Generally, the protein is mixed and used in feeding assays. See, for example Marrone et al. (1985) *J. of Economic Entomology* 78:290-293.

[0131] The present invention may be used for transformation of any plant species, including, but not limited to, monocots and dicots. Examples of plants of interest include, but are not limited to, corn (maize), sorghum, wheat, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape, *Brassica* sp., alfalfa, rye, millet, safflower, peanuts, sweet potato, cassava, coffee, coconut, pineapple, citrus trees, cocoa, tea, banana, avocado, fig, guava, mango, olive, papaya, cashew, macadamia, almond, oats, vegetables, ornamentals, and conifers. Vegetables include, but are not limited to, tomatoes, lettuce, green beans, lima beans, peas, and members of the genus *Curcumis* such as cucumber, cantaloupe, and musk melon. Ornamentals include, but are not limited to, azalea, hydrangea, hibiscus, roses, tulips, daffodils, petunias, carnation, poinsettia, and chrysanthemum. Preferably, plants of the present invention are crop plants (for example, maize, sorghum, wheat, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, oilseed rape, etc.).

Use in Insect Control

[0132] General methods for employing strains comprising a nucleotide sequence of the present invention, or a variant thereof, in insect control or in engineering other organisms as insecticidal agents are known in the art. See, for example U.S. Pat. No. 5,039,523 and EP 0480762A2.

[0133] The *Bacillus* strains containing a nucleotide sequence of the present invention, or a variant thereof, or the microorganisms that have been genetically altered to contain an insecticidal gene and protein may be used for protecting agricultural crops and products from insects. In one aspect of the invention, whole, i.e., unlysed, cells of a toxin (insecticide)-producing organism are treated with reagents that prolong the activity of the toxin produced in the cell when the cell is applied to the environment of target insect(s).

[0134] Alternatively, the insecticide is produced by introducing a delta-endotoxin gene into a cellular host. Expression of the delta-endotoxin gene results, directly or indirectly, in the intracellular production and maintenance of the insecticide. In one aspect of this invention, these cells are then treated under conditions that prolong the activity of the toxin produced in the cell when the cell is applied to the environment of target insect(s). The resulting product retains the toxicity of the toxin. These naturally encapsulated insecticides may then be formulated in accordance with conventional techniques for application to the environment hosting a target insect, e.g., soil, water, and foliage of plants. See, for example EPA 0192319, and the references cited therein. Alternatively, one may formulate the cells expressing a gene of this invention such as to allow application of the resulting material as an insecticide.

[0135] Insecticidal Compositions

[0136] The active ingredients of the present invention are normally applied in the form of compositions and can be

applied to the crop area or plant to be treated, simultaneously or in succession, with other compounds. These compounds can be fertilizers, weed killers, cryoprotectants, surfactants, detergents, insecticidal soaps, dormant oils, polymers, and/or time-release or biodegradable carrier formulations that permit long-term dosing of a target area following a single application of the formulation. They can also be selective herbicides, chemical insecticides, virucides, microbicides, amoebicides, pesticides, fungicides, bacteriocides, nematocides, molluscicides or mixtures of several of these preparations, if desired, together with further agriculturally acceptable carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. Suitable carriers and adjuvants can be solid or liquid and correspond to the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, binders or fertilizers. Likewise the formulations may be prepared into edible "baits" or fashioned into pest "traps" to permit feeding or ingestion by a target pest of the insecticidal formulation.

[0137] Methods of applying an active ingredient of the present invention or an agrochemical composition of the present invention that contains at least one of the insecticidal proteins produced by the bacterial strains of the present invention include leaf application, seed coating and soil application. The number of applications and the rate of application depend on the intensity of infestation by the corresponding insect.

[0138] The composition may be formulated as a powder, dust, pellet, granule, spray, emulsion, colloid, solution, or such like, and may be prepared by such conventional means as desiccation, lyophilization, homogenation, extraction, filtration, centrifugation, sedimentation, or concentration of a culture of cells comprising the polypeptide. In all such compositions that contain at least one such insecticidal polypeptide, the polypeptide may be present in a concentration of from about 1% to about 99% by weight.

[0139] Lepidopteran, coleopteran, or other insects may be killed or reduced in numbers in a given area by the methods of the invention, or may be prophylactically applied to an environmental area to prevent infestation by a susceptible insect. Preferably the insect ingests, or is contacted with, an insecticidally-effective amount of the polypeptide. By "insecticidally-effective amount" is intended an amount of the insecticide that is able to bring about death to at least one insect, or to noticeably reduce insect growth, feeding, or normal physiological development. This amount will vary depending on such factors as, for example, the specific target insects to be controlled, the specific environment, location, plant, crop, or agricultural site to be treated, the environmental conditions, and the method, rate, concentration, stability, and quantity of application of the insecticidally-effective polypeptide composition. The formulations may also vary with respect to climatic conditions, environmental considerations, and/or frequency of application and/or severity of insect infestation.

[0140] The insecticide compositions described may be made by formulating either the bacterial cell, crystal and/or spore suspension, or isolated protein component with the desired agriculturally-acceptable carrier. The compositions may be formulated prior to administration in an appropriate means such as lyophilized, freeze-dried, desiccated, or in an aqueous carrier, medium or suitable diluent, such as saline or other buffer. The formulated compositions may be in the form of a dust or granular material, or a suspension in oil (vegetable

or mineral), or water or oil/water emulsions, or as a wettable powder, or in combination with any other carrier material suitable for agricultural application. Suitable agricultural carriers can be solid or liquid and are well known in the art. The term "agriculturally-acceptable carrier" covers all adjuvants, inert components, dispersants, surfactants, tackifiers, binders, etc. that are ordinarily used in insecticide formulation technology; these are well known to those skilled in insecticide formulation. The formulations may be mixed with one or more solid or liquid adjuvants and prepared by various means, e.g., by homogeneously mixing, blending and/or grinding the insecticidal composition with suitable adjuvants using conventional formulation techniques. Suitable formulations and application methods are described in U.S. Pat. No. 6,468,523, herein incorporated by reference.

[0141] "Pest" includes but is not limited to, insects, fungi, bacteria, nematodes, mites, ticks, and the like. Insect pests include insects selected from the orders Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc., particularly Coleoptera, Lepidoptera, and Diptera.

[0142] The order Coleoptera includes the suborders Adephaga and Polyphaga. Suborder Adephaga includes the superfamilies Caraboidea and Gyrinoidea, while suborder Polyphaga includes the superfamilies Hydrophiloidea, Staphylinoidea, Cantharoidea, Cleroidea, Elateroidea, Dascilloidea, Dryopoidea, Byrrhoidea, Cucujoidea, Meloidea, Mordelloidea, Tenebrionoidea, Bostrichioidea, Scarabaeoidea, Cerambycoidea, Chrysomeloidea, and Curculionoidea. Superfamily Caraboidea includes the families Cicindelidae, Carabidae, and Dytiscidae. Superfamily Gyrinoidea includes the family Gyrinidae. Superfamily Hydrophiloidea includes the family Hydrophilidae. Superfamily Staphylinoidea includes the families Silphidae and Staphylinidae. Superfamily Cantharoidea includes the families Cantharidae and Lampyridae. Superfamily Cleroidea includes the families Cleridae and Dermestidae. Superfamily Elateroidea includes the families Elateridae and Buprestidae. Superfamily Cucujoidea includes the family Coccinellidae. Superfamily Meloidea includes the family Meloidea. Superfamily Tenebrionoidea includes the family Tenebrionidae. Superfamily Scarabaeoidea includes the families Passalidae and Scarabaeidae. Superfamily Cerambycoidea includes the family Cerambycidae. Superfamily Chrysomeloidea includes the family Chrysomelidae. Superfamily Curculionoidea includes the families Curculionidae and Scolytidae.

[0143] The order Diptera includes the Suborders Nemato-cera, Brachycera, and Cyclorrhapha. Suborder Nemato-cera includes the families Tipulidae, Psychodidae, Culicidae, Ceratopogonidae, Chironomidae, Simuliidae, Bibionidae, and Cecidomyiidae. Suborder Brachycera includes the families Stratiomyidae, Tabanidae, Therevidae, Asilidae, Mydidae, Bombyliidae, and Dolichopodidae. Suborder Cyclorrhapha includes the Divisions Aschiza and Schizura. Division Aschiza includes the families Phoridae, Syrphidae, and Conopidae. Division Schizura includes the Sections Acalyptratae and Calyptratae. Section Acalyptratae includes the families Otitidae, Tephritidae, Agromyzidae, and Drosophilidae. Section Calyptratae includes the families Hippoboscidae, Oestridae, Tachinidae, Anthomyiidae, Muscidae, Caliphoridae, and Sarcophagidae.

[0144] The order Lepidoptera includes the families Papilionidae, Pieridae, Lycaenidae, Nymphalidae, Danaidae,

Satyridae, Hesperidae, Sphingidae, Saturniidae, Geometridae, Arctiidae, Noctuidae, Lymantriidae, Sesiidae, Crambidae, and Tineidae.

[0145] Nematodes include parasitic nematodes such as root-knot, cyst, and lesion nematodes, including *Heterodera* spp., *Meloidogyne* spp., and *Globodera* spp.; particularly members of the cyst nematodes, including, but not limited to, *Heterodera glycines* (soybean cyst nematode); *Heterodera schachtii* (beet cyst nematode); *Heterodera avenae* (cereal cyst nematode); and *Globodera rostochiensis* and *Globodera pailida* (potato cyst nematodes). Lesion nematodes include *Pratylenchus* spp.

[0146] Insect pests of the invention for the major crops include: Maize: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Helicoverpa zea*, corn earworm; *Spodoptera frugiperda*, fall armyworm; *Diatraea grandiosella*, southwestern corn borer; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Diatraea saccharalis*, sugarcane borer; *Diabrotica virgifera*, western corn rootworm; *Diabrotica longicornis barberi*, northern corn rootworm; *Diabrotica undecimpunctata howardi*, southern corn rootworm; *Melanotus* spp., wireworms; *Cyclocephala borealis*, northern masked chafer (white grub); *Cyclocephala immaculata*, southern masked chafer (white grub); *Popillia japonica*, Japanese beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Anuraphis maidiradicis*, corn root aphid; *Blissus leucopterus leucopterus*, chinch bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Hylemya platura*, seedcorn maggot; *Agromyza parvicornis*, corn blot leafminer; *Anaphothrips obscurus*, grass thrips; *Solenopsis milesta*, thief ant; *Tetranychus urticae*, twospotted spider mite; *Sorghum*: *Chilo partellus*, sorghum borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Feltia subterranea*, granulate cutworm; *Phyllophaga crinita*, white grub; *Eleodes*, *Conoderus*, and *Aeolus* spp., wireworms; *Oulema melanopus*, cereal leaf beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Sipha flava*, yellow sugarcane aphid; *Blissus leucopterus leucopterus*, chinch bug; *Contarinia sorghicola*, sorghum midge; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Wheat: *Pseudaletia unipunctata*, army worm; *Spodoptera frugiperda*, fall armyworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Agrotis orthogonia*, western cutworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Oulema melanopus*, cereal leaf beetle; *Hypera punctata*, clover leaf weevil; *Diabrotica undecimpunctata howardi*, southern corn rootworm; Russian wheat aphid; *Schizaphis graminum*, greenbug; *Macrosiphum avenae*, English grain aphid; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Mayetiola destructor*, Hessian fly; *Sitodiplosis mosellana*, wheat midge; *Meromyza americana*, wheat stem maggot; *Hylemya coarctata*, wheat bulb fly; *Frankliniella fusca*, tobacco thrips; *Cephus cinctus*, wheat stem sawfly; *Aceria tulipae*, wheat curl mite; Sunflower: *Suleima helianthana*, sunflower bud moth; *Homoeosoma electellum*, sunflower moth; *zygogramma exclamationis*, sunflower beetle; *Bothyrus gibbosus*, carrot beetle; *Neolasioptera murtfeldtiana*, sunflower seed midge; Cotton: *Heliothis virescens*, cotton budworm; *Heli-*

coverpa zea, cotton bollworm; *Spodoptera exigua*, beet armyworm; *Pectinophora gossypiella*, pink bollworm; *Anthonomus grandis*, boll weevil; *Aphis gossypii*, cotton aphid; *Pseudatomoscelis seriatus*, cotton fleahopper; *Trialeurodes abutilonea*, bandedwinged whitefly; *Lygus lineolaris*, tarnished plant bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Thrips tabaci*, onion thrips; *Frankliniella fusca*, tobacco thrips; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Rice: *Diatraea saccharalis*, sugarcane borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Colaspis brunnea*, grape colaspis; *Lissorhoptrus oryzophilus*, rice water weevil; *Sitophilus oryzae*, rice weevil; *Nephotettix nigropictus*, rice leafhopper; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; Soybean: *Pseudoplusia includens*, soybean looper; *Anticarsia gemmatilis*, velvetbean caterpillar; *Plathypena scabra*, green cloverworm; *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Spodoptera exigua*, beet armyworm; *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Epilachna varivestis*, Mexican bean beetle; *Myzus persicae*, green peach aphid; *Empoasca fabae*, potato leafhopper; *Acrosternum hilare*, green stink bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Hylemya platura*, seedcorn maggot; *Sericothrips variabilis*, soybean thrips; *Thrips tabaci*, onion thrips; *Tetranychus turkestanii*, strawberry spider mite; *Tetranychus urticae*, twospotted spider mite; Barley: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Schizaphis graminum*, greenbug; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; *Euschistus servus*, brown stink bug; *Delia platura*, seedcorn maggot; *Mayetiola destructor*, Hessian fly; *Petrobia latens*, brown wheat mite; Oil Seed Rape: *Brevicoryne brassicae*, cabbage aphid; *Phyllotreta cruciferae*, Flea beetle; *Mamestra configurata*, Bertha armyworm; *Plutella xylostella*, Diamond-back moth; *Delia* spp., Root maggots.

Methods for Increasing Plant Yield

[0147] Methods for increasing plant yield are provided. The methods comprise introducing into a plant or plant cell a polynucleotide comprising an insecticidal sequence disclosed herein. As defined herein, the “yield” of the plant refers to the quality and/or quantity of biomass produced by the plant. By “biomass” is intended any measured plant product. An increase in biomass production is any improvement in the yield of the measured plant product. Increasing plant yield has several commercial applications. For example, increasing plant leaf biomass may increase the yield of leafy vegetables for human or animal consumption. Additionally, increasing leaf biomass can be used to increase production of plant-derived pharmaceutical or industrial products. An increase in yield can comprise any statistically significant increase including, but not limited to, at least a 1% increase, at least a 3% increase, at least a 5% increase, at least a 10% increase, at least a 20% increase, at least a 30%, at least a 50%, at least a 70%, at least a 100% or a greater increase in yield compared to a plant not expressing the insecticidal sequence.

[0148] In specific methods, plant yield is increased as a result of improved insect resistance of a plant expressing an insecticidal protein disclosed herein. Expression of the insect-

ticidal protein results in a reduced ability of an insect to infest or feed on the plant, thus improving plant yield.

[0149] The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

Example 1

Discovery of a Novel Toxin Gene Axmi115 from the *Bacillus thuringiensis* Strain ATX12983

[0150] The complete gene sequence was identified from the selected strain via the MiDAS genomics approach as follows:

[0151] Preparation of extrachromosomal DNA from the strain. Extrachromosomal DNA contains a mixture of some or all of the following: plasmids of various size; phage chromosomes; genomic DNA fragments not separated by the purification protocol; other uncharacterized extrachromosomal molecules.

[0152] Mechanical or enzymatic shearing of the extrachromosomal DNA to generate size-distributed fragments.

[0153] Sequencing of the fragmented DNA

[0154] Identification of putative toxin genes via homology and/or other computational analyses.

[0155] When required, sequence finishing of the gene of interest by one of several PCR or cloning strategies (e.g. TAIL-PCR).

The novel gene is referred to herein as axmi-115 (SEQ ID NO:3), and the encoded amino acid referred to as AXMI-115 (SEQ ID NO:6). Synthetic nucleotide sequences encoding AXMI-115 is set forth in SEQ ID NO:15 and 16.

Gene and Protein Characteristics

[0156] Gene length, DNA base pairs: 2409

Protein length, amino acid residues: 803

Estimated protein molecular weight, Da: 90877

Known homologs and approximate percent identity:

[0157] Vip3Afl—70.7%

[0158] Axmi005—70.4%

[0159] Axmi026—70.4%

[0160] Vip3Aa7—70.1

Example 2

Novel Insecticidal Protein AXMI-005 from the *Bacillus thuringiensis* Strain ATX13002

[0161] The AXMI-005 insecticidal gene was identified from the strain ATX13002 using the MiDAS approach as described in U.S. Patent Publication No. 20040014091, which is herein incorporated by reference in its entirety, using the following steps:

Steps taken in the current strategy to gene discovery:

Step 1: Culture from the strain was grown in large quantities. The plasmid DNA was then separated from the chromosomal DNA by a cesium chloride gradient spun in an ultracentrifuge. The purified plasmid DNA was then nebulized to a 5-10 kb size range appropriate for coverage of an average sized coding region. The fragment ends were polished then ligated overnight into a vector cut with a restriction enzyme producing blunt ends.

Step 3: Once the shotgun library quality was checked and confirmed, colonies were grown, prepped and sequenced in a

96-well format. The library plates were end sequenced off of the vector backbone for initial screening.

Step 5: All of the reads were compiled into an assembly project and aligned together to form contigs. These contigs, along with any individual read that may not have been added to a contig, were analyzed using BLAST, using a batch format, against an internal database made up of all classes of known delta-endotoxin genes. Any contigs or individual reads that pulled up any homology to a known gene were analyzed further by selecting a single clone from the library that covered the entire hypothesized coding region.

Step 6: The individual clone covering the area of interest was then walked over read by read by designing primers to extend the sequence. This was done until both end reads of the clone were joined and the coverage was at least 2x. The completed contig of the single clone was then analyzed using BLAST (both blastn and blastx) against a public database of all known insecticidal genes. Hits from both searches were then pulled from an internal database of all the genes (clipped to coding sequence only) and aligned with the completed library clone sequence to determine the percentage of divergence from the known gene.

[0162] A novel gene, referred to herein as axmi-005 (SEQ ID NO:1), and the encoded amino acid referred to as AXMI-005 (SEQ ID NO:4) were identified by this approach. Searching of public sequence databases, including the GENBANK® databases, showed that AXMI-005 is a unique protein that has highest homology (94.9%) to the vip3Aa insecticidal protein (GenePept ID L48841).

[0163] A synthetic sequence encoding the AXMI-005 protein was designed and termed optaxmi-005. The nucleotide sequence is set forth in SEQ ID NO:7 and encodes the amino acid sequence set forth in SEQ ID NO:9 (with the addition of a C-terminal histidine tag). The optaxmi-005 gene disclosed herein can be used with or without the C-terminal histidine tag.

Example 3

Discovery of a Novel Toxin Gene Axmi-113 from the *Bacillus thuringiensis* Strain ATX12987

[0164] The complete gene sequence was identified from the selected strain via the MiDAS genomics approach as follows:

[0165] Preparation of extrachromosomal DNA from the strain. Extrachromosomal DNA contains a mixture of some or all of the following: plasmids of various size; phage chromosomes; genomic DNA fragments not separated by the purification protocol; other uncharacterized extrachromosomal molecules.

[0166] Mechanical or enzymatic shearing of the extrachromosomal DNA to generate size-distributed fragments.

[0167] Sequencing of the fragmented DNA

[0168] Identification of putative toxin genes via homology and/or other computational analyses.

[0169] When required, sequence finishing of the gene of interest by one of several PCR or cloning strategies (e.g. TAIL-PCR).

The novel gene is referred to herein as axmi-113 (SEQ ID NO:2), and the encoded amino acid referred to as AXMI-113 (SEQ ID NO:5).

Gene and Protein Characteristics

[0170] Gene length, DNA base pairs: 2385
 Protein length, amino acid residues: 795
 Estimated protein molecular weight, Da: 89475
 Known homologs and approximate percent identity:

[0171] Vip3Ah—99%

[0172] Vip3Aa18—79.8%

[0173] Axmi005—79%

[0174] A synthetic sequence encoding the AXMI-113 protein was designed and termed optaxmi-113. The nucleotide sequence is set forth in SEQ ID NO:8 and encodes the amino acid sequence set forth in SEQ ID NO:5 or 10 (with the addition of a C-terminal histidine tag). The optaxmi-113 gene disclosed herein can be used with or without the C-terminal histidine tag.

Example 4

Discovery of Novel Toxin Genes Axmi-163 and Axmi-184 from the *Bacillus thuringiensis* Strain ATX14775

[0175] The complete gene sequence for each was identified from the selected strain via the MiDAS genomics approach as follows:

[0176] Preparation of extrachromosomal DNA from the strain. Extrachromosomal DNA contains a mixture of some or all of the following: plasmids of various size; phage chromosomes; genomic DNA fragments not separated by the purification protocol; other uncharacterized extrachromosomal molecules.

[0177] Mechanical or enzymatic shearing of the extrachromosomal DNA to generate size-distributed fragments.

[0178] Sequencing of the fragmented DNA

[0179] Identification of putative toxin genes via homology and/or other computational analyses.

[0180] When required, sequence finishing of the gene of interest by one of several PCR or cloning strategies (e.g. TAIL-PCR).

The novel gene referred to herein as axmi-163 is set forth in SEQ ID NO:11, and the encoded amino acid referred to as AXMI-163 is set forth in SEQ ID NO:13.

Gene and Protein Characteristics

[0181] Gene length, DNA base pairs: 2370
 Protein length, amino acid residues: 790
 Estimated protein molecular weight, Da: 88,700
 Known homologs and approximate percent identity:

[0182] SEQ ID NO:17 from U.S. Pat. No. 7,129,212—98%

[0183] Axmi005—78%

The novel gene referred to herein as axmi-184 is set forth in SEQ ID NO:12, and the encoded amino acid referred to as AXMI-184 is set forth in SEQ ID NO:14. Synthetic nucleotide sequences encoding AXMI-184 are set forth in SEQ ID NO:17 and 18.

Gene and Protein Characteristics

[0184] Gene length, DNA base pairs: 2370
 Protein length, amino acid residues: 790
 Estimated protein molecular weight, Da: 88,300
 Known homologs and approximate percent identity:

[0185] Vip3Afl—93%

[0186] Axmi005—86%

Example 5

Construction of Synthetic Sequences

[0187] In one aspect of the invention, synthetic axmi sequences are generated. These synthetic sequences have an altered DNA sequence relative to the parent axmi sequence, and encode a protein that is collinear with the parent AXMI protein to which it corresponds, but lacks the C-terminal “crystal domain” present in many delta-endotoxin proteins.

[0188] In another aspect of the invention, modified versions of synthetic genes are designed such that the resulting peptide is targeted to a plant organelle, such as the endoplasmic reticulum or the apoplast. Peptide sequences known to result in targeting of fusion proteins to plant organelles are known in the art. For example, the N-terminal region of the acid phosphatase gene from the White Lupin *Lupinus albus* (Genebank ID GI:14276838; Miller et al. (2001) Plant Physiology 127: 594-606) is known in the art to result in endoplasmic reticulum targeting of heterologous proteins. If the resulting fusion protein also contains an endoplasmic retention sequence comprising the peptide N-terminus-lysine-aspartic acid-glutamic acid-leucine (i.e. the “KDEL” motif (SEQ ID NO:19) at the C-terminus, the fusion protein will be targeted to the endoplasmic reticulum. If the fusion protein lacks an endoplasmic reticulum targeting sequence at the C-terminus, the protein will be targeted to the endoplasmic reticulum, but will ultimately be sequestered in the apoplast.

Example 6

Expression in *Bacillus*

[0189] As an example of the expression of the genes and proteins of the invention in *Bacillus* species, the insecticidal gene disclosed herein is amplified by PCR, and the PCR product is cloned into the *Bacillus* expression vector pAX916, or another suitable vector, by methods well known in the art. The resulting *Bacillus* strain, containing the vector with axmi gene is cultured on a conventional growth media, such as CYS media (10 g/l Bacto-casitone; 3 g/l yeast extract; 6 g/l KH_2PO_4 ; 14 g/l K_2HPO_4 ; 0.5 mM MgSO_4 ; 0.05 mM MnCl_2 ; 0.05 mM FeSO_4), until sporulation is evident by microscopic examination. Samples are prepared and tested for activity in bioassays.

Example 7

Expression in *E. coli*

[0190] As an example of a method of expression of the genes and proteins of the invention in *E. coli* based systems, the complete ORF of each axmi gene is cloned into an *E. coli* expression vector based on pRSF1b. The resulting clones are confirmed by restriction analysis and finally, by complete sequencing of the cloned gene.

[0191] For expression in *E. coli*, BL21*DE3 is transformed with the vector expressing the axmi gene. Single colonies are inoculated in LB supplemented with kanamycin and grown overnight at 37° C. The following day, fresh medium is inoculated in duplicate with 1% of overnight culture and grown at 37° C. to logarithmic phase. Subsequently, cultures are induced with 1 mM isopropyl β -D-1-thiogalactopyranoside (IPTG) for 3 hours at 37° C. or overnight at 20° C. Each cell pellet is suspended in 50 mM sodium carbonate buffer, pH

10.5 supplemented with 1 mM DTT dithiothreitol and sonicated. Samples are prepared and tested for activity in bioassays.

Example 8

Expression of AXMI-115, AXMI-113, and AXMI-005 in *E. coli*

[0192] *E. coli* clones were generated that contained DNA segments containing the complete open reading frame as well as a portion of the DNA region naturally occurring upstream and adjacent to each gene. This DNA segment for each of axmi-113 (SEQ ID:2), axmi-115 (SEQ ID NO:3) or axmi-005 (SEQ ID NO:1) was amplified and cloned into the vector pAX916, to yield the clones pAX5463, pAX5464 and pAX5465 respectively. The resulting clones were confirmed by restriction analysis and by complete sequencing of the cloned fragments.

[0193] *E. coli* cells were transformed with each of the clones pAX5463, pAX5464 and pAX5465.

[0194] axmi005, axmi113 and axmi115 genes that had their codons optimized for expression in corn, and had a C-terminal his6-tag added, were also expressed from *E. coli* expression vector utilizing T7 promoter. In addition, constructs were also generated that expressed N-terminal his6-tagged or untagged versions of optaxmi005 (pAX5475, pAX5478) and optaxmi115 (pAX5476, pAX5477).

[0195] Single *E. coli* colonies of the axmi-115, axmi-113, and axmi-005 expressing clones were then grown overnight at 37° C. in LB medium. The following day, fresh medium was inoculated in duplicate with 1% of overnight culture and grown at 37° C. to logarithmic phase. Subsequently, cultures were induced with 1 mM IPTG overnight at 20° C. The resulting cells were collected by centrifugation, and suspended in either 50 mM sodium carbonate buffer, pH 10.5 supplemented with 1 mM DTT or 50 mM Tris Cl buffer, pH 8 with 1 mM DTT prior to sonication. SDS-PAGE analysis showed expression of a ~90 kD protein in all samples.

Example 9

Insect Bioassays of *E. coli* Expressed Proteins

[0196] Soluble extracts containing AXMI-005, AXMI-113, or AXMI-115 were tested in insect assays with appropriate controls. Twenty four well tissue culture plates (Corning) were filled with 1 ml of multi-species diet (Bio-Serv) and allowed to solidify. Once solidified, 40 µl of protein sample was placed on the diet surface of each well and allowed to soak in/dry at room temperature. Depending upon the experiment, either egg masses or neonate larvae were placed in each well. Plates were sealed with gas-permeable membranes (Research Products International) and incubated at 25° C. and 90% relative humidity. After five or seven days, samples were scored visually compared to a buffer only or non-transformed extract control.

[0197] Strong activity of AXMI-005 extracts was observed on *Helicoverpa zea* (HZ), *Heliothis virescens* (HV), Fall Armyworm (FAW), Black cutworm (BCW), Sugarcane borer (SCB), and Velvet Bean caterpillar (VBC). AXMI-005 also showed activity on Southwestern Corn Borer (SWCB).

[0198] Strong activity of AXMI-115 extracts was observed on *Heliothis virescens*, Fall Armyworm, Black cutworm, and Velvet Bean caterpillar. AXMI-115 also exhibited activity on the European Corn Borer (ECB), SCB, SWCB, and Dia-

mondback moth (DBM). Activity of AXMI-115 on *Helicoverpa zea* (HZ) was less pronounced than for the other insects tested, but was still significant.

[0199] Activity of each of the AXMI-005 and AXMI-115 extracts was scored, and assigned a number from 1 to 5 based on relative activity in the assays. A summary of the scores in a particular assay is shown in Table 1.

[0200] AXMI-005 showed some activity on SWCB (score of 2) and high levels of activity on HZ, Hv, FAW, BCW and VBC (scores of 4 to 5). AXMI-115 showed high levels of activity on SWCB (80% mortality), ECB, FAW and VBC (scores of 4 to 5) and lesser activity on HZ and Hv. AXMI-113 also showed high activity on SWCB (score of 4 with 20% mortality) and on SCB. No activity was seen on the other insects tested.

TABLE 1

Insecticidal activity of AXMI-115, AXMI-113, and AXMI-005*					
	Axmi115 (pH 10.5)	Axmi115 (pH 8)	Axmi005 (pH10.5)	Axmi005 (pH 8)	Axmi113 (pH 10.5)
HZ	0	0	4	4	0
ECB	2	4	0	0	0
live infest					
Hv	0	0	4	4/5	0
FAW	4	2	4/5	4/5	0
BCW	0	0	3/4	4	0
VBC	4	3	4	4/5	0
SWCB	4; 80% mortality	3; 50% mortality	2	ND	4; 20% mortality
SCB	ND	3; 25% mortality	ND	4; 100% mortality	3; 50% mortality
DBM	ND	2/3	0	0	0

*= represented as stunt and mortality percent where stunting is scored according to the following scale:

Score	Definition
0	No Activity
1	Slight, non-uniform stunt
2	Non-uniform stunt
3	Uniform stunt
4	Uniform stunt with mortality (expressed as a percentage)
5	Uniform stunt with 100% mortality

Example 10

Bioassay of Axmi184

Gene Expression and Purification

[0201] The DNA region encoding the toxin domain of Axmi184 was cloned into an *E. coli* expression vector pMAL-C4x behind the maltE gene coding for Maltose binding protein (MBP). This in-frame fusion resulted in MBP-Axmi084 fusion protein expression in *E. coli*.

[0202] For expression in *E. coli*, BL21*DE3 was transformed with individual plasmids. Single colony was inoculated in LB supplemented with carbenicillin and glucose, and grown overnight at 37° C. The following day, fresh medium was inoculated with 1% of overnight culture and grown at 37° C. to logarithmic phase. Subsequently, cultures were induced with 0.3 mM IPTG for overnight at 20° C. Each cell pellet was suspended in 20 mM Tris-Cl buffer, pH 7.4+200 mM NaCl+1 mM DTT+protease inhibitors and sonicated. Analysis by SDS-PAGE confirmed expression of fusion proteins.

[0203] Total cell free extracts were run over amylose column attached to FPLC for affinity purification of MBP-AXMI184 fusion proteins. Bound fusion protein was eluted from the resin with 10 mM maltose solution. Purified fusion proteins were then cleaved with either Factor Xa or trypsin to remove the amino terminal MBP tag from the AXMI184 protein. Cleavage and solubility of the proteins was determined by SDS-PAGE.

[0204] Cleaved proteins were tested in insect assays with appropriate controls. A 5-day read of the plates showed following activities of AXMI-184 against Diamondback moth.

Example 11

Domain Swapping

[0205] axmi005, axmi113 and axmi115 genes that had their codons optimized for expression in corn were used in this example. Plasmids expressing untagged versions of optaxmi005 (pAX5478), optaxmi113 (pAX5493) and optaxmi115 (pAX5477) were used to design DNA swap constructs as describe here.

[0206] AXMI-005, AXMI-113 and AXMI-115 have significant sequence identity/similarity in their N-terminal $\frac{2}{3}^{rd}$ region. The remaining $\frac{1}{3}^{rd}$ region in their C-termini (CT) shows substantial sequence divergence as seen in the protein sequence alignment provided as FIGS. 1A and 1B.

[0207] The protein region of AXMI-113 between the forward and reverse arrows shown in FIG. 1 was replaced with the corresponding fragment of either AXMI-005 (to give pAX5492) or AXMI-115 (pAX5494).

[0208] For expression in *E. coli*, BL21*DE3 was transformed with individual constructs. A single colony was inoculated in LB supplemented with kanamycin and grown overnight at 37° C. The following day, fresh medium was inoculated in duplicate with 1% of overnight culture and grown at 37° C. to logarithmic phase. Subsequently, cultures

were induced with 1 mM IPTG overnight at 20° C. Cell pellet was suspended in 50 mM sodium carbonate buffer, pH 10.5 supplemented with 1 mM DTT, and sonicated. Analysis by SDS-PAGE showed extremely good soluble expression of all proteins.

[0209] Filter sterilized, soluble extracts expressing OptAxmi005, 113, 115, Optaxmi113+CT of Optaxmi005 and Optaxmi113+CT of Optaxmi115 were tested in insect assays with appropriate controls. As shown in Example 9, AXMI-113 showed high activity on SWCB (25% mortality). It showed an additional activity on SCB (50% mortality). Also as shown in Example 9, AXMI-005 showed activity on SWCB, Hz, Hv, FAW, BCW and VBC. It showed an additional activity on SCB (25% mortality). AXMI-115 was also found to have some activity on SCB.

[0210] The fusion of AXMI-113+CT of AXMI-005 showed all of the insect activities seen with AXMI-005. In other words, replacement of the C-terminal fragment of AXMI-113 with that of AXMI-005 bestowed upon it the insect activities that were otherwise missing in its naturally occurring form.

[0211] Additional toxin protein sequences can be generated by swapping domains from one protein into another. For example, one or more of the AXMI-005 domains shown in FIG. 2 are introduced into AXMI-115. The domains are introduced using the sense ("s") and antisense ("a") oligonucleotides shown in Table 2. The portion of the axmi-005 sequence that is being introduced into the axmi-115 sequence is shown in bold print. The flanking sequences in each oligonucleotide are axmi-115 sequences that are used for annealing the oligonucleotides to the axmi-115 template. The number following the term "sub" in each primer name corresponds to the numbered boxes in FIG. 2. Similar oligonucleotides can be designed to swap domains between multiple sequences, for example, between the AXMI-005, AXMI-113, AXMI-115, AXMI-163 and AXMI-184 sequences described herein.

TABLE 2

Oligonucleotide primer	Sequence	SEQ ID NO:
axmi115sub1 s	AAC ACC GGC GGC GTC AAT GGA ACA AGG GCG CTC TTC ACC CA	20
axmi115sub1 a	TGG GTG AAG AGC GCC CTT GTT CCA TTG ACG CCG CCG GTG TT	21
axmi115sub10 s	GCC CGG AGC TCA TCA ATG TCA ACA ACT GGA TCA GAA CTG GCA CCA CCT ACA TCA C	22
axmi115sub10 a	GTG ATG TAG GTG GTG CCA GTT CTG ATC CAG TTG TTG ACA TTG ATG AGC TCC GGG C	23
axmi115sub11 s	ATG ATT GGG AGA GGT TCG GAA GCA CCC ACA TCA GCG GCA ATG AGC TGA GG	24
axmi115sub11 a	CCT CAG CTC ATT GCC GCT GAT GTG GGT GCT TCC GAA CCT CTC CCA ATC AT	25
axmi115sub12 s	CTA CAT CAC CGG CAA TAC CTT GAC GCT CTA CCA AGG AGG AGG AGG CTA CTT CCG C	26
axmi115sub12 a	GCG GAA GTA GCC TCC TCC TCC TTG GTA GAG CGT CAA GGT ATT GCC GGT GAT GTA G	27
axmi115sub14 s	CGA CAG CTA CAG CAC CTA CAG GGT GAA CTT CTC CGT CAC CGG CTG GGC CAA GGT GAT	28

TABLE 2 -continued

Oligonucleotide primer	Sequence	SEQ ID NO:
axmi115sub14 a	ATC ACC TTG GCC CAG CCG GTG ACG GAG AAG TTC ACC CTG TAG GTG CTG TAG CTG TCG	29
axmi115sub15 s	GCT TCA GCG GCC TCG ACG CCA ATG TGA GGA TCA GAA ACA GCC GCG GC	30
axmi115sub15 a	GCC GCG GCT GTT TCT GAT CCT CAC ATT GGC GTC GAG GCC GCT GAA GC	31
axmi115sub16 s	GTG AAG AAC AGC CGC GAG GTG CTC TTC GAG AAG AGA TAC ATG AAT GGA AGC AGC TAT GA	32
axmi115sub16 a	TCA TAG CTG CTT CCA TTC ATG TAT CTC TTC TCG AAG AGC ACC TCG CGG CTG TTC TTC AC	33
axmi115sub17 s	TTC GAG AAG GTG AAG AAC AGC GGC GCC AAG GAT GTT TCA GAG AGC TTC ACC ACC	34
axmi115sub17 a	GGT GGT GAA GCT CTC TGA AAC ATC CTT GGC GCC GCT GTT CTT CAC CTT CTC GAA	35
axmi115sub19 s	GCT TCT TCA TCG AGC TCA GCC AAG GCA ACA ACC TCT ATA GCA GCA CCT TCC AC	36
axmi115sub19 a	GTG GAA GGT GCT GCT ATA GAG GTT GTT GCC TTG GCT GAG CTC GAT GAA GAA GC	37
axmi115sub2 s	GAA GCA AGG CGC TCT ATG TTC ACA AGG ATG GAG GCT TCA GCC AGT TCA TCG	38
axmi115sub2 a	CGA TGA ACT GGC TGA AGC CTC CAT CCT TGT GAA CAT AGA GCG CCT TGC TTC	39
axmi115sub20 s	CCG CCG AGA GGA CAG GAG GGC CGC TGG TGA AGT TCA GAG ACA TCA GCA TC	40
axmi115sub20 a	GAT GCT GAT GTC TCT GAA CTT CAC CAG CGG CCC TCC TGT CCT CTC GGC GG	41
axmi115sub21 s	AGC ACC TTC CAC AGC TTC AAT GAT GTG AGC ATC AAG TAA GGC GCG CCG	42
axmi115sub21 a	CGG CGC GCC TTA CTT GAT GCT CAC ATC ATT GAA GCT GTG GAA GGT GCT	43
axmi115sub3 s	CGA CAA GCT AAA GCC CAA GAC AGA ATA TGT CAT CCA GTA CAC CGT CAA G	44
axmi115sub3 a	CTT GAC GGT GTA CTG GAT GAC ATA TTC TGT CTT GGG CTT TAG CTT GTC G	45
axmi115sub5 s	CCT ACG AGG ACA CCA ATA ACA ACA ACC TGG AGG ACT ACC AAA CAA TTG CTG TGA AG	46
axmi115sub5 a	CTT CAC AGC AAT TGT TTG GTA GTC CTC CAG GTT GTT GTT ATT GGT GTC CTC GTA GG	47
axmi115sub6 s	GAG GAG TTC CAA ACA ATT ACC AAG AGG TTC ACC ACC GGC ACA GAT TTG AGC CAG ACC	48
axmi115sub6 a	GGT CTG GCT CAA ATC TGT GCC GGT GGT GAA CCT CTT GGT AAT TGT TTG GAA CTC CTC	49
axmi115sub7 s	CAC CTC AGA AAC AGA TTT GAA GGG CGT CTA CCT CAT CTT GAA GAG CCA AAA TGG ATA T	50

TABLE 2 -continued

Oligonucleotide primer	Sequence	SEQ ID NO:
axmi115sub7 a	ATA TCC ATT TTG GCT CTT CAA GAT GAG GTA GAC GCC CTT CAA ATC TGT TTC TGA GGT G	51
axmi115sub9 s	TCC TGG AGG CCA AGC CAT CAG AGA AGC TGC TCA GCC CGG AGC TCA	52
axmi115sub9 a	TGA GCT CCG GGC TGA GCA GCT TCT CTG ATG GCT TGG CCT CCA GGA	53
axmi115sub13 s	ATC ATT CAA GAG GAG GCA ACC TCA AGC AGA ACC TCC AGC TTG ACA GCT TCA GCA CCT ACG ACC TCA G	54
axmi115sub13 a	CTG AGG TCG TAG GTG CTG AAG CTG TCA AGC TGG AGG TTC TGC TTG AGG TTG CCT CCT CTT GAA TGA T	55
axmi115sub18 s	GCT ATG AGG ACA TCT CAG AGA TCT TCA CCA CCA AGC TGG GCA AGG ACA ACT TC TAC A TCG AGC TCA CCG C	56
axmi115sub18 a	GCG GTG AGC TCG AT GTA G AAG TTG TCC TTG CCC AGC TTG GTG GTG AAG ATC TCT GAG ATG TCC TCA TAG C	57
axmi115sub4 s	CAA GGG CAA GCC GTC AAT CCA CCT CAA GAA TGA GAA CAC CGG CTA CAT CCA CTAC GA GGA CAC CAA TGG	58
axmi115sub4 a	CCA TTG GTG TCC TCG TAG TGG ATG TAG CCG GTG TTC TCA TTC TTG AGG TGG ATT GAC GGC TTG CCC TTG	59
axmi115sub8 s	CAA GAG CCA AAA TGG AGA TGA AGC ATG GGG AGA CAA CTT CAC CAT CCT GGA GAT CTC GCT CTT CGA GAC ACC AGA A	60
axmi115sub8 a	TTC TGG TGT CTC GAA GAG CGA GAT CTC CAG GAT GGT GAA GTT GTC TCC CCA TGC TTC ATC TCC ATT TTG GCT CTT G	61

Example 12

Additional Assays for Pesticidal Activity

[0212] The ability of an insecticidal protein to act as a pesticide upon a pest is often assessed in a number of ways. One way well known in the art is to perform a feeding assay. In such a feeding assay, one exposes the pest to a sample containing either compounds to be tested, or control samples. Often this is performed by placing the material to be tested, or a suitable dilution of such material, onto a material that the pest will ingest, such as an artificial diet. The material to be tested may be in a liquid, solid, or slurry form. The material to be tested may be placed upon the surface and then allowed to dry or incorporate into the diet. Alternatively, the material to be tested may be mixed with a molten artificial diet, then dispensed into the assay chamber. The assay chamber may be, for example, a cup, a dish, or a well of a microtiter plate.

[0213] Assays for sucking pests (for example aphids) may involve separating the test material from the insect by a partition, ideally a portion that can be pierced by the sucking mouth parts of the sucking insect, to allow ingestion of the test material. Often the test material is mixed with a feeding stimulant, such as sucrose, to promote ingestion of the test compound.

[0214] Other types of assays can include microinjection of the test material into the mouth, or gut of the pest, as well as development of transgenic plants, followed by test of the ability of the pest to feed upon the transgenic plant. Plant testing may involve isolation of the plant parts normally consumed, for example, small cages attached to a leaf, or isolation of entire plants in cages containing insects.

[0215] Other methods and approaches to assay pests are known in the art, and can be found, for example in Robertson, J. L. & H. K. Preisler. 1992. *Pesticide bioassays with arthropods*. CRC, Boca Raton, Fla. Alternatively, assays are commonly described in the journals "Arthropod Management Tests" and "Journal of Economic Entomology" or by discussion with members of the Entomological Society of America (ESA).

Example 13

Vectoring of the Insecticidal Genes of the Invention for Plant Expression

[0216] Each of the coding regions of the genes of the invention is connected independently with appropriate promoter and terminator sequences for expression in plants. Such sequences are well known in the art and may include the rice

actin promoter or maize ubiquitin promoter for expression in monocots, the *Arabidopsis* UBQ3 promoter or CaMV 35S promoter for expression in dicots, and the nos or PinII terminators. Techniques for producing and confirming promoter—gene—terminator constructs also are well known in the art.

Example 14

Transformation of the Genes of the Invention into Plant Cells by *Agrobacterium*-Mediated Transformation

[0217] Ears are collected 8-12 days after pollination. Embryos are isolated from the ears, and those embryos 0.8-1.5 mm in size are used for transformation. Embryos are plated scutellum side-up on a suitable incubation media, and incubated overnight at 25° C. in the dark. However, it is not necessary per se to incubate the embryos overnight. Embryos are contacted with an *Agrobacterium* strain containing the appropriate vectors for Ti plasmid mediated transfer for 5-10 min, and then plated onto co-cultivation media for 3 days (25° C. in the dark). After co-cultivation, explants are transferred to recovery period media for five days (at 25° C. in the dark). Explants are incubated in selection media for up to eight weeks, depending on the nature and characteristics of the particular selection utilized. After the selection period, the resulting callus is transferred to embryo maturation media, until the formation of mature somatic embryos is observed. The resulting mature somatic embryos are then placed under low light, and the process of regeneration is initiated as known in the art. The resulting shoots are allowed to root on rooting media, and the resulting plants are transferred to nursery pots and propagated as transgenic plants.

Example 15

Transformation of Maize Cells with the Insecticidal Genes of the Invention

[0218] Maize ears are collected 8-12 days after pollination. Embryos are isolated from the ears, and those embryos 0.8-1.5 mm in size are used for transformation. Embryos are plated scutellum side-up on a suitable incubation media, such as DN62A5S media (3.98 g/L N6 Salts; 1 mL/L (of 1000× Stock) N6 Vitamins; 800 mg/L L-Asparagine; 100 mg/L Myo-inositol; 1.4 g/L L-Proline; 100 mg/L Casaminoacids; 50 g/L sucrose; 1 mL/L (of 1 mg/mL Stock) 2,4-D), and incubated overnight at 25° C. in the dark.

[0219] The resulting explants are transferred to mesh squares (30-40 per plate), transferred onto osmotic media for 30-45 minutes, then transferred to a beaming plate (see, for example, PCT Publication No. WO/0138514 and U.S. Pat. No. 5,240,842).

[0220] DNA constructs designed to express the genes of the invention in plant cells are accelerated into plant tissue using

an aerosol beam accelerator, using conditions essentially as described in PCT Publication No. WO/0138514. After beaming, embryos are incubated for 30 min on osmotic media, then placed onto incubation media overnight at 25° C. in the dark. To avoid unduly damaging beamed explants, they are incubated for at least 24 hours prior to transfer to recovery media. Embryos are then spread onto recovery period media, for 5 days, 25° C. in the dark, then transferred to a selection media. Explants are incubated in selection media for up to eight weeks, depending on the nature and characteristics of the particular selection utilized. After the selection period, the resulting callus is transferred to embryo maturation media, until the formation of mature somatic embryos is observed. The resulting mature somatic embryos are then placed under low light, and the process of regeneration is initiated by methods known in the art. The resulting shoots are allowed to root on rooting media, and the resulting plants are transferred to nursery pots and propagated as transgenic plants.

Materials

[0221]

DN62A5S Media		
Components	per Liter	Source
Chu's N6 Basal Salt Mixture (Prod. No. C 416)	3.98 g/L	Phytotechnology Labs
Chu's N6 Vitamin Solution (Prod. No. C 149)	1 mL/L (of 1000x Stock)	Phytotechnology Labs
L-Asparagine	800 mg/L	Phytotechnology Labs
Myo-inositol	100 mg/L	Sigma
L-Proline	1.4 g/L	Phytotechnology Labs
Casaminoacids	100 mg/L	Fisher Scientific
Sucrose	50 g/L	Phytotechnology Labs
2,4-D (Prod. No. D-7299)	1 mL/L (of 1 mg/mL Stock)	Sigma

[0222] Adjust the pH of the solution to pH to 5.8 with 1N KOH/1N KCl, add Gelrite (Sigma) to 3 g/L, and autoclave. After cooling to 50° C., add 2 ml/l of a 5 mg/ml stock solution of Silver Nitrate (Phytotechnology Labs). This recipe yields about 20 plates.

[0223] All publications and patent applications mentioned in the specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[0224] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 61

<210> SEQ ID NO 1

<211> LENGTH: 2370

<212> TYPE: DNA

-continued

<213> ORGANISM: *Bacillus thuringiensis*

<400> SEQUENCE: 1

```

atgaatatga ataatactaa attaaacgca agggccctac cgagttttat tgattatttt    60
aatggcattt atggatttgc cactggtatc aaagacatta tgaatatgat ttttaaacg    120
gatacagggtg gtaatctaac cttagacgaa atcctaaaga atcagcagtt actaaatgag    180
atttctggta aattggatgg ggtaaatggg agcttaaatg atcttatcgc acagggaaac    240
ttaaatcacg aattatctaa ggaaatctta aaaattgcaa atgaacagaa tcaagtctta    300
aatgatgtta ataacaaact cgatgcgata aatagcatgc ttcatatata tctacctaaa    360
attacatcta tgtaagtga tgtaatgaag caaaattatg cgctaagtct gcaaatagaa    420
tacttaagta agcaattgca agaaatttct gataaattag atattattaa cgtaaatgtt    480
cttattaact ctacacttac tgaaattaca cctgcataac aacggattaa atatgtgaat    540
gaaaattttg aagaattaac ttttgcatac gaaaccactt taaaagtaaa aaaggatagc    600
tcgcctgctg atattcttga tgagttaact gaattaactg aactagcgaa aagtgttaca    660
aaaaatgacg ttgatggttt tgaattttac cttaatacat tccacgatgt aatggtagga    720
aataatttat tcgggcgctc agcttttaaa actgcttcag aattaattgc taaagaaaat    780
gtgaaaacaa gtggcagtga agtaggaat gtttataatt tcttaattgt attaacagct    840
ctacaagcaa aagcttttct tactttaaca acatgccgaa aattattagg cttagcagat    900
attgattata cttctattat gaatgaacat ttaataaagg aaaagagga atttagagta    960
aacatccttc ctacacttct taatactttt tctaattccta attatgcaa agttaaagga   1020
agtgatgaag atgcaaatg gattgtggaa gctaaaccag gacatgcatt ggttgggttt   1080
gaaatgagca atgattcaat cacagtatta aaagtatatg aggctaagct aaaacaaaat   1140
tatcaagttg ataaggatcc cttatcggag gttatttatg gtgatatgga taaattattg   1200
tgtccagatc aatctgaaca aatttattat acaataata tagtatttcc aatgaatat   1260
gtaattacta aaattgattt tactaagaaa atgaagactt taagatatga ggtaacagct   1320
aattcttatg attctctac aggagaaatt gacttaata agaagaaagt agaatcaagt   1380
gaagcggagt ataggacgtt aagtgcataa gatgatggag tgtatatgcc gttaggtgtc   1440
atcagtgaac catttttgac tccgattaat gggtttgccc tccaagctga tgaatttca   1500
agattaatta ctttaacatg taaatcatat ttaagagaac tactgttagc aacagactta   1560
agcaataaag aaactaaatt gatcgtcccg ccaagtgggt ttattagtaa tattgtagaa   1620
aatgggaact tagagggaga aaacttagag ccgtggatag caaataataa gaatgcgtat   1680
gtagatcata caggcggagt gaatggtagt agagctttat atgttcataa ggacggagga   1740
ttttcacaat ttattggaga taagttaaaa ccgaaaactg agtatgtaat ccaatatact   1800
gttaaggaaa aaccttctat tcatttaaaa aatgaaaata ctggatatat tcattatgaa   1860
gatacaataa acaatttaga agattatcaa actattacta aacgttttac tacaggaact   1920
gatttaaagg gagtgtattt aattttaaaa agtcaaatg gagatgaagc ctggggagat   1980
aactttacaa ttttgaaaat tagtccttct gaaaagtatt taagtccaga attaatcaat   2040
gtaataaatt ggattcgcac gggatcaact catattagcg gtaatacact cactctctat   2100
caggaggagg gaggaatct aaaacaaaac cttcaattag acagtttttc aacttataga   2160
gtgaattttt ctgtgaccgg agatgctaata gtaaggatta gaaattctag ggaagtgtta   2220

```

-continued

tttgaaaaac gatatatgag cgggtgctaaa gatgtttctg aaattttcac tacaaaattg	2280
gggaaagata acttttatat agagctttct caagggaata atttatatgg tggtcctctt	2340
gtaaagttta acgatgtctc aattaagtaa	2370

<210> SEQ ID NO 2
 <211> LENGTH: 2385
 <212> TYPE: DNA
 <213> ORGANISM: *Bacillus thuringiensis*

<400> SEQUENCE: 2

atgaacatga ataatactaa attaagcaca agagccttac caagttttat tgattat	60
aatggcattt atggatttgc cactgggtatc aaagacatta tgaacatgat ttttaaaacg	120
gatacaggtg gtgatctaac cctagacgaa attttaaaga atcagcagtt actaaatgaa	180
atttctggta aattggatgg ggtaaatgga agcttaaag atcttatcgc acagggaaac	240
ttaaatacag aattatctaa ggaaatatta aaaattgcaa atgaacaaaa tcaagtttta	300
aatgatgtta ataacaaact tgatgcgata aatacgtatc tgcataata tctacctaaa	360
attacctcta tgttaagtga tgtaaatgaa caaaattatg cgctaagtct gcaaatagaa	420
tacctaatga aacagttgca agaaatttct gataaattgg atattattaa cgtaaatgta	480
cttattaact ctacacttac tgaaattaca cctgcgtatc aacggattaa atatgtgaac	540
gaaaaatttg aggaattaac ttttgctaca gaaactaatt taaaagttaa aaaggatggc	600
tctctgcag atattcttga tgagttaact gagttaactg aactagcgaa aagtgtaca	660
aaaaatgatg tggatgggtt tgaattttat cttaatacat tccacgatgt aatggtaggg	720
aataatttat tggggcggtc agctttaaaa actgcatcgg aattaattac taaagaaaat	780
gtgaaaacaa gtggcagtga ggtcggaat gtttataact tcttaattgt attaacagct	840
ctgcaagcaa aagcttttct tactttaaca acatgccgaa aattattagg cttagcagat	900
attgattata cttctattat gaatgaacat ttaaataagg aaaaagagga atttagagta	960
aacatccttc ctacacttcc taatactttt tctaactcta attatataaa aactaaagga	1020
agcgatgagg atgcggaagt gattattcaa gctgaaccag gacatgcttt ggtgggattt	1080
gaaatgatta atgatccaag tccagcgtta aaagtgtatc aggctaaact aacaacaaat	1140
tatcaagttg ataagcagtc attatctgaa actgtgtatg gtgatatgga taaaatatta	1200
tgccagata aatctcaaca aatgtattat ctgcataata taacattccc aaatgaatat	1260
gtgattacag aaattatttt tacaaagaaa aagaacagtt taaggatga ggtgatagca	1320
aattattatg agttctcttc aggagatata gacttaaata agaagttagt aaaatcaagt	1380
gaagcagagt atagtacgtt aagtgttagt aacgatgcta tttatagcc gctaggtgtt	1440
atcagtgaat cttttttgac cccaattaaa ggattcggac taacagttga tgaagttca	1500
agactggtaa cttaacgtg taaatcatat ttaagggaaa ttctgttagc aacagattta	1560
agcaataaag caacaaaatt aattgtccca cctaattggt ttattagtaa tttggtggaa	1620
aatggggata ttgagcgtga caacatagaa ccatggaagg gaaataataa aaatgcgtat	1680
gttgatcata caggtggggg gaatggaact aaagctctgt aactcaaga tgatggggaa	1740
ttttcacaat ttattggaga taagtgaaa tcgaaaactg agtatataat tcaatatact	1800
gtaaaaggaa atacttctat ttatttgaaa gataaaaaaa atgaaaatgt tatttatgaa	1860

-continued

gataaaaata ataatttaga ggcttttcaa actattacta aaagggtttac tacagaattg	1920
gattcttcag atgtttactt agtgtttaaa tgcaaaaatg gctataaagc ttggggagac	1980
aactttctaa ttacagaaat taggcctaag gaagtggtaa gccagaatt gataaaagta	2040
gaaaattgga ttggaatggg tggtagtaat catgtaaacc ctgattcact tttgcttttt	2100
acagggtgga ggtcaatttt aaaacaaaat ctccaattag atagttattc aacctataga	2160
gtaagatttt ctttaatggt aattggtaag gctaaggta ttataaggaa ttcaagtga	2220
gtactgtttg aaaaaagta tgtgaatgat tctgaagggtg ttttagaagg tgtttctgaa	2280
acttttacta caaaatcgat tcaagataac ttctatgtag aactttctaa tgagggcact	2340
tttggaaagta aagatgttgc atacttttat aatttttcta ttaga	2385

<210> SEQ ID NO 3

<211> LENGTH: 2409

<212> TYPE: DNA

<213> ORGANISM: *Bacillus thuringiensis*

<400> SEQUENCE: 3

atgaacatga ataactactaa attaaacgca agggccttac caagttttat tgattatttt	60
aatggcattt atggatttgc cactggatc aaagacatta tgaacatgat ttttaaacg	120
gatacagggtg gtgatctaac cctagacgaa attttaaaga atcagcagtt actaaatgaa	180
atttctggta aattggatgg ggtaaatggg agcttaaatg atcttatcgc acagggaac	240
ttaaatacag aattatctaa ggaaatctta aaaattgcaa atgaacaaaa tcaagtttta	300
aatgatgtta ataacaaact agatgcgata aatacgatgc ttaacatata tctacctaaa	360
attacatcta tgtaagcga tgtaaatgaa caaaattatg cgctaagtct gcagatagaa	420
tacctaagta gacagtaca ggaaatttca gataagttag atgttattaa tttaaatgtt	480
ttaattaatt ctactcttac tgaaatcaca ccttcataac aacgtattaa atatgtaaat	540
gagaaatttg ataaatgaac atttgccaca gaaagcactc taagagcaaa acaaggaatt	600
tttaacgagg atagttttga taataatact cttgaaaatt taactgatct agctgaacta	660
gctaaaaagta taacaaaaa tgatgtggat agtttcgaat tttatcttca cacatttcat	720
gatgtattaa ttggaaataa tttatttggg cgtatcggtt taaaaacagc ttcagaatta	780
attactaaag acgagataaa gacaagtggg agtgagattg gaaaagtta tagtttctta	840
attgtgctaa cttctttaca agctaaagcc tttctcactt taacaacatg tagaaaatta	900
ttgggcttat cagatattga ttatacttct attatgaatg aacatctaaa taatgaaaaa	960
aatgaatttc gtgataacat acttcctgca ctgtcgaata agttttctaa ccctagctac	1020
gcaaaaacta taggtagtga caattatgca aaagtatttt tagaatctga accaggttat	1080
gctttagtgc gattcgaaat tattaatgac ccaatcccggt tattaaaagc gtataaagca	1140
aaactaaaac aaaattatca agttgataat cagtcgttat cagagattgt ttatttagat	1200
atcgataaac tattttgtcc ggaaaattca gaacaaaaat attatactaa aaatctgaca	1260
tttctgatg gctatgttat tactaaaatt acctttgaaa aaaaactgaa caacctaat	1320
tatgaagcaa cagcaaatat ttatgaccca tctacaggag atattgactt aaataagaag	1380
caagtggat ctacttttcc tcaaacggat tatattacta tggatatagg tgatgatgat	1440
ggtatttata tgccgttagg cgttatcagt gaaacgtttt tgactcctat taatagtttt	1500
ggattagaag ttgacgcgaa atcgaaaaca ttaaccttaa aatgtaaatc ttaoctgcga	1560

-continued

```

gaatatttat tagagtcga tctaaaaaat aaagaaacag gcttgattgc cccgccaaat 1620
gtttttatta gtaatgttgt gaaaaattgg gatatagagg aagatagtct agaaccatgg 1680
gtagcaaata acaaaaaatgc ctatgttgat aatacaggcg gtatagaaag atctaaagcc 1740
ctctttactc aagggtgacgg ggaattctca caattttattg gtgataaatt aaaacctaata 1800
acagattata ttattcaata tactgtaaaa ggaaaaccag ccattttattt aaaaaataaa 1860
agtactggat atattacgta cgaggatata aacggtaatt ctgaagaatt tcaaaactata 1920
gctgtaaaaat tcacttcaga aactgatctt tcacaaactc atttagtttt taaaagtcaa 1980
aatgggtatg aggcttgggg agacaatttt attatttttag aagctaagct atttgaaact 2040
cctgaaagtc cagaattgat aaaaatttaatt gattgggaaa gatttgggtac tacttacatt 2100
acaggaaatg agcttagaat cgatcatagt cgtggagggtt attttagaca atctcttaata 2160
atagacagtt attcaactta tgatttgagc ttttctttta gtggattatg ggctaagggt 2220
attgtgaaaa attcccaggg ggtagtattg tttgaaaaag tgaaaaacaa cggttcatca 2280
tatgaagata ttagtgaag ttttactacc gcatacaata aggatggatt ttttatagaa 2340
ctaacagccg aaagaacgag ctcaactttc catagctttc gtgatatttc tattaaggaa 2400
aagattgaa 2409

```

<210> SEQ ID NO 4

<211> LENGTH: 789

<212> TYPE: PRT

<213> ORGANISM: *Bacillus thuringiensis*

<400> SEQUENCE: 4

```

Met Asn Met Asn Asn Thr Lys Leu Asn Ala Arg Ala Leu Pro Ser Phe
  1             5             10             15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
      20             25             30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asn Leu Thr Leu
      35             40             45

Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile Ser Gly Lys
      50             55             60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
      65             70             75             80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
      85             90             95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
      100            105            110

Met Leu His Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
      115            120            125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
      130            135            140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
      145            150            155            160

Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
      165            170            175

Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
      180            185            190

Thr Leu Lys Val Lys Lys Asp Ser Ser Pro Ala Asp Ile Leu Asp Glu
      195            200            205

```

-continued

Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val	
210	215 220
Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly	
225	230 235 240
Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile	
	245 250 255
Ala Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr	
	260 265 270
Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr	
	275 280 285
Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr	
	290 295 300
Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val	
305	310 315 320
Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala	
	325 330 335
Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys	
	340 345 350
Pro Gly His Ala Leu Val Gly Phe Glu Met Ser Asn Asp Ser Ile Thr	
	355 360 365
Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp	
	370 375 380
Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu	
385	390 395 400
Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe	
	405 410 415
Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys	
	420 425 430
Thr Leu Arg Tyr Glu Val Thr Ala Asn Ser Tyr Asp Ser Ser Thr Gly	
	435 440 445
Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr	
	450 455 460
Arg Thr Leu Ser Ala Lys Asp Asp Gly Val Tyr Met Pro Leu Gly Val	
465	470 475 480
Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala	
	485 490 495
Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg	
	500 505 510
Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile	
	515 520 525
Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Asn Leu	
	530 535 540
Glu Gly Glu Asn Leu Glu Pro Trp Ile Ala Asn Asn Lys Asn Ala Tyr	
545	550 555 560
Val Asp His Thr Gly Gly Val Asn Gly Thr Arg Ala Leu Tyr Val His	
	565 570 575
Lys Asp Gly Gly Phe Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys	
	580 585 590
Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His	
	595 600 605
Leu Lys Asn Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn	

-continued

610	615	620
Asn Leu Glu Asp Tyr Gln Thr Ile Thr Lys Arg Phe Thr Thr Gly Thr 625 630 635 640		
Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu 645 650 655		
Ala Trp Gly Asp Asn Phe Thr Ile Leu Glu Ile Ser Pro Ser Glu Lys 660 665 670		
Leu Leu Ser Pro Glu Leu Ile Asn Val Asn Asn Trp Ile Arg Thr Gly 675 680 685		
Ser Thr His Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Gly 690 695 700		
Gly Asn Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg 705 710 715 720		
Val Asn Phe Ser Val Thr Gly Asp Ala Asn Val Arg Ile Arg Asn Ser 725 730 735		
Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys Asp Val 740 745 750		
Ser Glu Ile Phe Thr Thr Lys Leu Gly Lys Asp Asn Phe Tyr Ile Glu 755 760 765		
Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Leu Val Lys Phe Asn 770 775 780		
Asp Val Ser Ile Lys 785		
<210> SEQ ID NO 5		
<211> LENGTH: 795		
<212> TYPE: PRT		
<213> ORGANISM: Bacillus thuringiensis		
<400> SEQUENCE: 5		
Met Asn Met Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe 1 5 10 15		
Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp 20 25 30		
Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu 35 40 45		
Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile Ser Gly Lys 50 55 60		
Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn 65 70 75 80		
Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln 85 90 95		
Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr 100 105 110		
Met Leu His Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val 115 120 125		
Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys 130 135 140		
Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val 145 150 155 160		
Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile 165 170 175		
Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr		

-continued

180							185							190						
Asn	Leu	Lys	Val	Lys	Lys	Asp	Gly	Ser	Pro	Ala	Asp	Ile	Leu	Asp	Glu					
		195					200					205								
Leu	Thr	Glu	Leu	Thr	Glu	Leu	Ala	Lys	Ser	Val	Thr	Lys	Asn	Asp	Val					
	210					215					220									
Asp	Gly	Phe	Glu	Phe	Tyr	Leu	Asn	Thr	Phe	His	Asp	Val	Met	Val	Gly					
225					230					235					240					
Asn	Asn	Leu	Phe	Gly	Arg	Ser	Ala	Leu	Lys	Thr	Ala	Ser	Glu	Leu	Ile					
				245					250					255						
Thr	Lys	Glu	Asn	Val	Lys	Thr	Ser	Gly	Ser	Glu	Val	Gly	Asn	Val	Tyr					
			260					265					270							
Asn	Phe	Leu	Ile	Val	Leu	Thr	Ala	Leu	Gln	Ala	Lys	Ala	Phe	Leu	Thr					
		275					280					285								
Leu	Thr	Thr	Cys	Arg	Lys	Leu	Leu	Gly	Leu	Ala	Asp	Ile	Asp	Tyr	Thr					
	290				295						300									
Ser	Ile	Met	Asn	Glu	His	Leu	Asn	Lys	Glu	Lys	Glu	Glu	Phe	Arg	Val					
305					310					315					320					
Asn	Ile	Leu	Pro	Thr	Leu	Ser	Asn	Thr	Phe	Ser	Asn	Pro	Asn	Tyr	Ile					
				325					330					335						
Lys	Thr	Lys	Gly	Ser	Asp	Glu	Asp	Ala	Glu	Val	Ile	Ile	Gln	Ala	Glu					
			340					345					350							
Pro	Gly	His	Ala	Leu	Val	Gly	Phe	Glu	Met	Ile	Asn	Asp	Pro	Ser	Pro					
		355					360					365								
Ala	Leu	Lys	Val	Tyr	Gln	Ala	Lys	Leu	Thr	Thr	Asn	Tyr	Gln	Val	Asp					
	370					375						380								
Lys	Gln	Ser	Leu	Ser	Glu	Thr	Val	Tyr	Gly	Asp	Met	Asp	Lys	Ile	Leu					
385					390					395					400					
Cys	Pro	Asp	Lys	Ser	Gln	Gln	Met	Tyr	Tyr	Leu	His	Asn	Ile	Thr	Phe					
				405					410					415						
Pro	Asn	Glu	Tyr	Val	Ile	Thr	Glu	Ile	Ile	Phe	Thr	Lys	Lys	Lys	Asn					
			420					425					430							
Ser	Leu	Arg	Tyr	Glu	Val	Ile	Ala	Asn	Tyr	Tyr	Glu	Phe	Ser	Ser	Gly					
		435					440					445								
Asp	Ile	Asp	Leu	Asn	Lys	Lys	Leu	Val	Lys	Ser	Ser	Glu	Ala	Glu	Tyr					
	450					455					460									
Ser	Thr	Leu	Ser	Val	Ser	Asn	Asp	Ala	Ile	Tyr	Met	Pro	Leu	Gly	Val					
465					470					475					480					
Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	Ile	Lys	Gly	Phe	Gly	Leu	Thr	Val					
				485					490					495						
Asp	Glu	Ser	Ser	Arg	Leu	Val	Thr	Leu	Thr	Cys	Lys	Ser	Tyr	Leu	Arg					
			500					505					510							
Glu	Ile	Leu	Leu	Ala	Thr	Asp	Leu	Ser	Asn	Lys	Ala	Thr	Lys	Leu	Ile					
		515					520						525							
Val	Pro	Pro	Asn	Gly	Phe	Ile	Ser	Asn	Leu	Val	Glu	Asn	Gly	Asp	Ile					
		530				535						540								
Glu	Ala	Asp	Asn	Ile	Glu	Pro	Trp	Lys	Gly	Asn	Asn	Lys	Asn	Ala	Tyr					
545					550					555					560					
Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Lys	Ala	Leu	Tyr	Thr	Gln					
				565					570					575						
Asp	Asp	Gly	Glu	Phe	Ser	Gln	Phe	Ile	Gly	Asp	Lys	Leu	Lys	Ser	Lys					
			580					585					590							

```

<210> SEQ ID NO 6
<211> LENGTH: 803
<212> TYPE: PRT
<213> ORGANISM: Bacillus thuringiensis

<400> SEQUENCE: 6

Met Asn Met Asn Asn Thr Lys Leu Asn Ala Arg Ala Leu Pro Ser Phe
 1             5             10             15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20             25             30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
 35             40             45

Asp Glu Ile Leu Lys Asn Gln Leu Leu Asn Glu Ile Ser Gly Lys
 50             55             60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
 65             70             75             80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
 85             90             95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
100            105            110

Met Leu Asn Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
115            120            125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Arg
130            135            140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Val Ile Asn Leu Asn Val
145            150            155            160

```


-continued

Leu	Ile	Asn	Ser	Thr	Leu	Thr	Glu	Ile	Thr	Pro	Ser	Tyr	Gln	Arg	Ile	165	170	175
Lys	Tyr	Val	Asn	Glu	Lys	Phe	Asp	Lys	Leu	Thr	Phe	Ala	Thr	Glu	Ser	180	185	190
Thr	Leu	Arg	Ala	Lys	Gln	Gly	Ile	Phe	Asn	Glu	Asp	Ser	Phe	Asp	Asn	195	200	205
Asn	Thr	Leu	Glu	Asn	Leu	Thr	Asp	Leu	Ala	Glu	Leu	Ala	Lys	Ser	Ile	210	215	220
Thr	Lys	Asn	Asp	Val	Asp	Ser	Phe	Glu	Phe	Tyr	Leu	His	Thr	Phe	His	225	230	235
Asp	Val	Leu	Ile	Gly	Asn	Asn	Leu	Phe	Gly	Arg	Ser	Ala	Leu	Lys	Thr	245	250	255
Ala	Ser	Glu	Leu	Ile	Thr	Lys	Asp	Glu	Ile	Lys	Thr	Ser	Gly	Ser	Glu	260	265	270
Ile	Gly	Lys	Val	Tyr	Ser	Phe	Leu	Ile	Val	Leu	Thr	Ser	Leu	Gln	Ala	275	280	285
Lys	Ala	Phe	Leu	Thr	Leu	Thr	Thr	Cys	Arg	Lys	Leu	Leu	Gly	Leu	Ser	290	295	300
Asp	Ile	Asp	Tyr	Thr	Ser	Ile	Met	Asn	Glu	His	Leu	Asn	Asn	Glu	Lys	305	310	315
Asn	Glu	Phe	Arg	Asp	Asn	Ile	Leu	Pro	Ala	Leu	Ser	Asn	Lys	Phe	Ser	325	330	335
Asn	Pro	Ser	Tyr	Ala	Lys	Thr	Ile	Gly	Ser	Asp	Asn	Tyr	Ala	Lys	Val	340	345	350
Ile	Leu	Glu	Ser	Glu	Pro	Gly	Tyr	Ala	Leu	Val	Gly	Phe	Glu	Ile	Ile	355	360	365
Asn	Asp	Pro	Ile	Pro	Val	Leu	Lys	Ala	Tyr	Lys	Ala	Lys	Leu	Lys	Gln	370	375	380
Asn	Tyr	Gln	Val	Asp	Asn	Gln	Ser	Leu	Ser	Glu	Ile	Val	Tyr	Leu	Asp	385	390	395
Ile	Asp	Lys	Leu	Phe	Cys	Pro	Glu	Asn	Ser	Glu	Gln	Lys	Tyr	Tyr	Thr	405	410	415
Lys	Asn	Leu	Thr	Phe	Pro	Asp	Gly	Tyr	Val	Ile	Thr	Lys	Ile	Thr	Phe	420	425	430
Glu	Lys	Lys	Leu	Asn	Asn	Leu	Ile	Tyr	Glu	Ala	Thr	Ala	Asn	Phe	Tyr	435	440	445
Asp	Pro	Ser	Thr	Gly	Asp	Ile	Asp	Leu	Asn	Lys	Lys	Gln	Val	Glu	Ser	450	455	460
Thr	Phe	Pro	Gln	Thr	Asp	Tyr	Ile	Thr	Met	Asp	Ile	Gly	Asp	Asp	Asp	465	470	475
Gly	Ile	Tyr	Met	Pro	Leu	Gly	Val	Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	485	490	495
Ile	Asn	Ser	Phe	Gly	Leu	Glu	Val	Asp	Ala	Lys	Ser	Lys	Thr	Leu	Thr	500	505	510
Leu	Lys	Cys	Lys	Ser	Tyr	Leu	Arg	Glu	Tyr	Leu	Leu	Glu	Ser	Asp	Leu	515	520	525
Lys	Asn	Lys	Glu	Thr	Gly	Leu	Ile	Ala	Pro	Pro	Asn	Val	Phe	Ile	Ser	530	535	540
Asn	Val	Val	Lys	Asn	Trp	Asp	Ile	Glu	Glu	Asp	Ser	Leu	Glu	Pro	Trp	545	550	555
Val	Ala	Asn	Asn	Lys	Asn	Ala	Tyr	Val	Asp	Asn	Thr	Gly	Gly	Ile	Glu	565	570	575

-continued

Arg Ser Lys Ala Leu Phe Thr Gln Gly Asp Gly Glu Phe Ser Gln Phe
 580 585 590
 Ile Gly Asp Lys Leu Lys Pro Asn Thr Asp Tyr Ile Ile Gln Tyr Thr
 595 600 605
 Val Lys Gly Lys Pro Ala Ile Tyr Leu Lys Asn Lys Ser Thr Gly Tyr
 610 615 620
 Ile Thr Tyr Glu Asp Thr Asn Gly Asn Ser Glu Glu Phe Gln Thr Ile
 625 630 635 640
 Ala Val Lys Phe Thr Ser Glu Thr Asp Leu Ser Gln Thr His Leu Val
 645 650 655
 Phe Lys Ser Gln Asn Gly Tyr Glu Ala Trp Gly Asp Asn Phe Ile Ile
 660 665 670
 Leu Glu Ala Lys Leu Phe Glu Thr Pro Glu Ser Pro Glu Leu Ile Lys
 675 680 685
 Phe Asn Asp Trp Glu Arg Phe Gly Thr Thr Tyr Ile Thr Gly Asn Glu
 690 695 700
 Leu Arg Ile Asp His Ser Arg Gly Gly Tyr Phe Arg Gln Ser Leu Asn
 705 710 715 720
 Ile Asp Ser Tyr Ser Thr Tyr Asp Leu Ser Phe Ser Phe Ser Gly Leu
 725 730 735
 Trp Ala Lys Val Ile Val Lys Asn Ser Arg Gly Val Val Leu Phe Glu
 740 745 750
 Lys Val Lys Asn Asn Gly Ser Ser Tyr Glu Asp Ile Ser Glu Ser Phe
 755 760 765
 Thr Thr Ala Ser Asn Lys Asp Gly Phe Phe Ile Glu Leu Thr Ala Glu
 770 775 780
 Arg Thr Ser Ser Thr Phe His Ser Phe Arg Asp Ile Ser Ile Lys Glu
 785 790 795 800
 Lys Ile Glu

<210> SEQ ID NO 7

<211> LENGTH: 2388

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic nucleotide sequence encoding AXMI-005 w/c-terminal extension

<400> SEQUENCE: 7

```

atgaacatga acaacaccaa gctcaatgca agggcgctgc cgagcttcat cgactacttc      60
aatggcatct atggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagacc      120
gacaccggcg gcaacctcac cttggatgag atcctcaaga accagcagct gctgaatgag      180
atctcaggca agctggacgg cgtcaatgga agcctcaacg acctcattgc tcaaggcaac      240
ctcaacaccg agctgagcaa ggagatcctc aagattgcaa atgagcagaa ccaggtgctg      300
aatgatgtca acaacaagct ggacgccatc aacaccatgc tgcacatcta cctgccaaag      360
atcacctcaa tgctctctga tgtgatgaag cagaactacg cgctgagcct ccagattgag      420
tacctctcaa agcagctgca agagatctcc gacaagctgg acatcatcaa tgtcaatgtg      480
ctcatcaaca gcaccttgac agagatcacg ccggcctacc agaggatcaa gtatgtcaat      540
gagaagtttg aggagctcac cttcgccacc gagacaacat tgaaggtgaa gaaggacagc      600
tcgccggcgg acatcctgga tgagctcacc gagctaacag agctggccaa gagcgtcacc      660

```

-continued

```

aagaatgatg ttgatggcct cgagttctac ctcaacacct tccatgatgt gatggtgggc 720
aacaacctct tcggccgcgc gccgctcaag acggcgctcg agctgatcgc caaggagaat 780
gtcaagacaa gtggatcaga ggtgggcaat gtctacaact tcctcatcgt gctgacggcg 840
ctgcaagcca aggccttctc caccttgaca acctgccgca agttgctggg cctcgccgac 900
atcgactaca cctccatcat gaatgagcac ctcaacaagg agaaggagga gttccgcgtc 960
aacatcctgc caacattgag caacaccttc agcaacccca actacgcca ggtgaagggc 1020
tcagatgaag atgccaagat gattgtggag gccaaagctg gccatgctct ggtgggcttc 1080
gagatgagca acgacagcat caccgtgctg aaggtctacg aggccaaagt gaagcagaac 1140
taccagggtg acaagacag cttgtctgag gtgatctacg gcgacatgga caagctgcta 1200
tgtccagatc aaagcgagca gatctactac accaacaaca tcgtctttcc aaatgaatat 1260
gtcatcacca agatcgactt caccaagaag atgaaaacat tgagatatga ggtgacggcc 1320
aacagctacg acagcagcac cggcgagatc gacctcaaca agaagaaggt ggagagctca 1380
gaagctgagt acaggacgct ctccgccaa gatgatggcg tctacatgcc gctcgcgctc 1440
atctcagaaa ctttcttgac gcccatcaat ggcttcggcc tccaagctga tgagaacagc 1500
aggctcatca cttgacctg caagagctac ctacgggagc tgctgctggc caccgacctc 1560
agcaacaagg agacaagct catcgtgccg ccacagctt tcatcagcaa catcgtggag 1620
aatggcaacc tggaaggaga gaacctggag ccacggatag ccaacaacaa gaatgcttat 1680
gttgatcaca cggcgcgct caatggaaca agggcgctct atgttcacaa ggatggaggc 1740
ttcagccagt tcatcgccga caagctgaag cccaagacag aatatgtcat ccagtacacc 1800
gtcaagggca agccatcaat ccacctcaag aatgagaaca cggctacat ccactacgag 1860
gacaccaaca acaacctgga ggactaccag accatcacca agaggttcac caccggcacc 1920
gacctcaagg gcgtctacct catcttgaag agccaaaatg gagatgaagc atggggagac 1980
aacttcacca tcctggagat ctgcccata gagaagctgc tctcgccgga gctcatcaat 2040
gtcaacaact ggatcagaac tggaagcacc cacatcagcg gcaacacctt gacgctctac 2100
caaggaggag gaggcaacct caagcagaac ctccagcttg acagcttctc cactacagg 2160
gtgaacttct ccgtcaccgg cgacgccaat gtgaggatca gaaattcaag ggaggtgctc 2220
ttcgagaaga gatacatgag cggcgccaag gatgtttctg agatcttcac caccaagctg 2280
ggcaaggaca acttctacat cgagctgagc caaggcaaca acctctatgg agggccgctg 2340
gtgaagttca atgatgtgag catcaagcat caccaccatc atcactaa 2388

```

<210> SEQ ID NO 8

<211> LENGTH: 2430

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic nucleotide sequence encoding AXMI-113 w/c-terminal extension

<400> SEQUENCE: 8

```

atgaacatga acaacaccaa gctcaatgca agggcgctgc cgagcttcat cgactacttc 60
aatggcatct atggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagacc 120
gacaccggcg gcgacctcac cttggatgag atcctcaaga accagcagct gctgaatgag 180
atctcaggca agctggacgg cgtcaatgga agcctcaagc acctcattgc tcaaggcaac 240

```

-continued

ctcaacaccg agctgagcaa ggagatcctc aagattgcaa atgagcagaa ccagggtgctg	300
aatgatgtca acaacaagct ggacgccatc aacaccatgc tcaacatcta cctgccaaaag	360
atcacctcaa tgctctctga tgtgatgaag cagaactacg cgctgagcct ccagattgag	420
tacctctcaa ggcagctgca agagatctcc gacaagctgg atgtcatcaa cctcaatgtg	480
ctcatcaaca gcaccttgac agagatcacg ccaagctacc agaggatcaa gtatgtcaat	540
gagaagttcg acaagctcac ctctgccacc gagagcacc tccgcgccaa gcaaggcacc	600
ttcaatgaag attcatttga caacaacacc ttggagaact tgacagacct cgccgagctg	660
gccaaagaca tcaccaagaa tgatgtggac agcttcgagt tctacctcca caccttccat	720
gatgtgtca tcggcaacaa cctcttttga agaagcgcgc tcaagacggc atcagagctc	780
atcaccaagg atgagatcaa gacaagcggc agcgagatcg gcaaggtcta cagcttctc	840
atcgtgtgta catcattgca agccaaggcc ttctcacct tgacaacctg ccgcaagttg	900
ctgggcctct ccgacatcga ctacacctcc atcatgaatg agcacctcaa caatgagaag	960
aatgagttca gagacaacat cctgcggcg ctgagcaaca agttcagcaa cccaagctac	1020
gccaaagaca tcggctcaga caactacgcc aaggtgatcc tggagagcga gcctggctac	1080
gcgctgggtg gcttcgagat catcaatgat ccaattcctg ttctcaaggc ctacaaggcc	1140
aagctgaagc agaactacca ggtggacaac cagagcttga gcgagatcgt ctacctggac	1200
atcgacaagc tcttctgccc ggagaactca gagcagaagt actacacca gaacctcacc	1260
ttccctgatg gatatgtcat caccaagatc accttcgaga agaagctgaa caacctcacc	1320
tacgaggcca ccgccaactt ctatgatcca tcaacaggag acatcgacct caacaagaag	1380
caagtgagga gcaccttccc tcaaacagac tacatcacca tggacattgg agatgatgat	1440
ggcatctaca tgccgctcgg cgtcatctca gaaaccttct tgacgcccat caacagcttc	1500
ggcctggagg tggagcccaa gagcaagacc ttgacgctca agtgcaagag ctacctcagg	1560
gagtacctgc tggagagtga tttgaagaac aaggagacag ggctgatcgc gccgccaaat	1620
gtgttcacat gcaatgtggt gaagaactgg gacatcgagg aggattcatt ggagccatgg	1680
gtggccaaca acaagaatgc ttatgtggac aacaccggcg gcattgaaag aagcaaggcg	1740
ctcttcccc aaggagatgg agagttcagc cagttcatcg gcgacaagct aaagcccaac	1800
accgactaca tcatccagta caccgtcaag ggcaagccgg ccatctacct caagaacaag	1860
agcaccggct acatcaccta cgaggacacc aatggaaatt ctgaggagtt ccaacaatt	1920
gctgtgaagt tcacctcaga aacagatttg agccagacc acctggtgtt caagagccaa	1980
aatggatatg aagcatgggg agacaacttc atcatcctgg aggccaaagct ctctgagaca	2040
ccagaaagcc cggagctcat caagttcaat gattgggaga ggttcggcac cacctacatc	2100
accggcaatg agctgaggat tgatcattca agaggaggct acttcgccca aagcctcaac	2160
atcgacagct acagcaccta cgacctcagc ttcagcttca gggcctctg ggccaagggtg	2220
attgtgaaga acagccggcg cgtggtgctc ttcgagaagg tgaagaacaa tggaagcagc	2280
tatgaggaca tctcagagag cttcaccacc gccagcaaca aggatggctt cttcatcgag	2340
ctcaccggcg agaggacaag cagcaccttc cacagcttca gagacatcag catcaaggag	2400
aagattgaac atcaccacca tcatcactaa	2430

-continued

```

<211> LENGTH: 795
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AXMI-005 w/ c-terminal extension

<400> SEQUENCE: 9
Met Asn Met Asn Asn Thr Lys Leu Asn Ala Arg Ala Leu Pro Ser Phe
 1             5             10            15
Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20            25            30
Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asn Leu Thr Leu
 35            40            45
Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile Ser Gly Lys
 50            55            60
Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
 65            70            75            80
Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
 85            90            95
Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
100           105           110
Met Leu His Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
115           120           125
Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
130           135           140
Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
145           150           155           160
Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
165           170           175
Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
180           185           190
Thr Leu Lys Val Lys Lys Asp Ser Ser Pro Ala Asp Ile Leu Asp Glu
195           200           205
Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
210           215           220
Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
225           230           235           240
Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
245           250           255
Ala Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr
260           265           270
Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr
275           280           285
Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr
290           295           300
Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val
305           310           315           320
Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala
325           330           335
Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys
340           345           350
Pro Gly His Ala Leu Val Gly Phe Glu Met Ser Asn Asp Ser Ile Thr
355           360           365

```

-continued

Val	Leu	Lys	Val	Tyr	Glu	Ala	Lys	Leu	Lys	Gln	Asn	Tyr	Gln	Val	Asp
370						375					380				
Lys	Asp	Ser	Leu	Ser	Glu	Val	Ile	Tyr	Gly	Asp	Met	Asp	Lys	Leu	Leu
385					390					395					400
Cys	Pro	Asp	Gln	Ser	Glu	Gln	Ile	Tyr	Tyr	Thr	Asn	Asn	Ile	Val	Phe
				405					410					415	
Pro	Asn	Glu	Tyr	Val	Ile	Thr	Lys	Ile	Asp	Phe	Thr	Lys	Lys	Met	Lys
			420					425					430		
Thr	Leu	Arg	Tyr	Glu	Val	Thr	Ala	Asn	Ser	Tyr	Asp	Ser	Ser	Thr	Gly
		435					440					445			
Glu	Ile	Asp	Leu	Asn	Lys	Lys	Lys	Val	Glu	Ser	Ser	Glu	Ala	Glu	Tyr
	450					455					460				
Arg	Thr	Leu	Ser	Ala	Lys	Asp	Asp	Gly	Val	Tyr	Met	Pro	Leu	Gly	Val
465					470					475					480
Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	Ile	Asn	Gly	Phe	Gly	Leu	Gln	Ala
				485					490					495	
Asp	Glu	Asn	Ser	Arg	Leu	Ile	Thr	Leu	Thr	Cys	Lys	Ser	Tyr	Leu	Arg
			500					505					510		
Glu	Leu	Leu	Leu	Ala	Thr	Asp	Leu	Ser	Asn	Lys	Glu	Thr	Lys	Leu	Ile
		515					520					525			
Val	Pro	Pro	Ser	Gly	Phe	Ile	Ser	Asn	Ile	Val	Glu	Asn	Gly	Asn	Leu
	530					535					540				
Glu	Gly	Glu	Asn	Leu	Glu	Pro	Trp	Ile	Ala	Asn	Asn	Lys	Asn	Ala	Tyr
545					550					555					560
Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Arg	Ala	Leu	Tyr	Val	His
				565					570					575	
Lys	Asp	Gly	Gly	Phe	Ser	Gln	Phe	Ile	Gly	Asp	Lys	Leu	Lys	Pro	Lys
			580					585					590		
Thr	Glu	Tyr	Val	Ile	Gln	Tyr	Thr	Val	Lys	Gly	Lys	Pro	Ser	Ile	His
		595					600					605			
Leu	Lys	Asn	Glu	Asn	Thr	Gly	Tyr	Ile	His	Tyr	Glu	Asp	Thr	Asn	Asn
	610					615					620				
Asn	Leu	Glu	Asp	Tyr	Gln	Thr	Ile	Thr	Lys	Arg	Phe	Thr	Thr	Gly	Thr
625					630					635					640
Asp	Leu	Lys	Gly	Val	Tyr	Leu	Ile	Leu	Lys	Ser	Gln	Asn	Gly	Asp	Glu
				645					650					655	
Ala	Trp	Gly	Asp	Asn	Phe	Thr	Ile	Leu	Glu	Ile	Ser	Pro	Ser	Glu	Lys
			660					665					670		
Leu	Leu	Ser	Pro	Glu	Leu	Ile	Asn	Val	Asn	Asn	Trp	Ile	Arg	Thr	Gly
		675					680					685			
Ser	Thr	His	Ile	Ser	Gly	Asn	Thr	Leu	Thr	Leu	Tyr	Gln	Gly	Gly	Gly
	690					695					700				
Gly	Asn	Leu	Lys	Gln	Asn	Leu	Gln	Leu	Asp	Ser	Phe	Ser	Thr	Tyr	Arg
705					710					715					720
Val	Asn	Phe	Ser	Val	Thr	Gly	Asp	Ala	Asn	Val	Arg	Ile	Arg	Asn	Ser
				725					730					735	
Arg	Glu	Val	Leu	Phe	Glu	Lys	Arg	Tyr	Met	Ser	Gly	Ala	Lys	Asp	Val
			740					745					750		
Ser	Glu	Ile	Phe	Thr	Thr	Lys	Leu	Gly	Lys	Asp	Asn	Phe	Tyr	Ile	Glu
		755					760					765			
Leu	Ser	Gln	Gly	Asn	Asn	Leu	Tyr	Gly	Gly	Pro	Leu	Val	Lys	Phe	Asn
	770					775					780				

-continued

Asp Val Ser Ile Lys His His His His His His
785 790 795

<210> SEQ ID NO 10
<211> LENGTH: 809
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AXMI-113 w/ c-terminal extension

<400> SEQUENCE: 10

Met Asn Met Asn Asn Thr Lys Leu Asn Ala Arg Ala Leu Pro Ser Phe
1 5 10 15
Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
20 25 30
Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
35 40 45
Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile Ser Gly Lys
50 55 60
Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
65 70 75 80
Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
85 90 95
Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
100 105 110
Met Leu Asn Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
115 120 125
Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Arg
130 135 140
Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Val Ile Asn Leu Asn Val
145 150 155 160
Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ser Tyr Gln Arg Ile
165 170 175
Lys Tyr Val Asn Glu Lys Phe Asp Lys Leu Thr Phe Ala Thr Glu Ser
180 185 190
Thr Leu Arg Ala Lys Gln Gly Ile Phe Asn Glu Asp Ser Phe Asp Asn
195 200 205
Asn Thr Leu Glu Asn Leu Thr Asp Leu Ala Glu Leu Ala Lys Ser Ile
210 215 220
Thr Lys Asn Asp Val Asp Ser Phe Glu Phe Tyr Leu His Thr Phe His
225 230 235 240
Asp Val Leu Ile Gly Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr
245 250 255
Ala Ser Glu Leu Ile Thr Lys Asp Glu Ile Lys Thr Ser Gly Ser Glu
260 265 270
Ile Gly Lys Val Tyr Ser Phe Leu Ile Val Leu Thr Ser Leu Gln Ala
275 280 285
Lys Ala Phe Leu Thr Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ser
290 295 300
Asp Ile Asp Tyr Thr Ser Ile Met Asn Glu His Leu Asn Asn Glu Lys
305 310 315 320
Asn Glu Phe Arg Asp Asn Ile Leu Pro Ala Leu Ser Asn Lys Phe Ser
325 330 335

-continued

Asn	Pro	Ser	Tyr	Ala	Lys	Thr	Ile	Gly	Ser	Asp	Asn	Tyr	Ala	Lys	Val
			340					345					350		
Ile	Leu	Glu	Ser	Glu	Pro	Gly	Tyr	Ala	Leu	Val	Gly	Phe	Glu	Ile	Ile
		355					360					365			
Asn	Asp	Pro	Ile	Pro	Val	Leu	Lys	Ala	Tyr	Lys	Ala	Lys	Leu	Lys	Gln
	370					375					380				
Asn	Tyr	Gln	Val	Asp	Asn	Gln	Ser	Leu	Ser	Glu	Ile	Val	Tyr	Leu	Asp
385					390					395					400
Ile	Asp	Lys	Leu	Phe	Cys	Pro	Glu	Asn	Ser	Glu	Gln	Lys	Tyr	Tyr	Thr
				405					410					415	
Lys	Asn	Leu	Thr	Phe	Pro	Asp	Gly	Tyr	Val	Ile	Thr	Lys	Ile	Thr	Phe
			420					425					430		
Glu	Lys	Lys	Leu	Asn	Asn	Leu	Ile	Tyr	Glu	Ala	Thr	Ala	Asn	Phe	Tyr
		435					440					445			
Asp	Pro	Ser	Thr	Gly	Asp	Ile	Asp	Leu	Asn	Lys	Lys	Gln	Val	Glu	Ser
	450					455					460				
Thr	Phe	Pro	Gln	Thr	Asp	Tyr	Ile	Thr	Met	Asp	Ile	Gly	Asp	Asp	Asp
465					470					475					480
Gly	Ile	Tyr	Met	Pro	Leu	Gly	Val	Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro
				485					490					495	
Ile	Asn	Ser	Phe	Gly	Leu	Glu	Val	Asp	Ala	Lys	Ser	Lys	Thr	Leu	Thr
			500					505					510		
Leu	Lys	Cys	Lys	Ser	Tyr	Leu	Arg	Glu	Tyr	Leu	Leu	Glu	Ser	Asp	Leu
		515					520					525			
Lys	Asn	Lys	Glu	Thr	Gly	Leu	Ile	Ala	Pro	Pro	Asn	Val	Phe	Ile	Ser
	530					535					540				
Asn	Val	Val	Lys	Asn	Trp	Asp	Ile	Glu	Glu	Asp	Ser	Leu	Glu	Pro	Trp
545					550					555					560
Val	Ala	Asn	Asn	Lys	Asn	Ala	Tyr	Val	Asp	Asn	Thr	Gly	Gly	Ile	Glu
				565					570					575	
Arg	Ser	Lys	Ala	Leu	Phe	Thr	Gln	Gly	Asp	Gly	Glu	Phe	Ser	Gln	Phe
			580					585					590		
Ile	Gly	Asp	Lys	Leu	Lys	Pro	Asn	Thr	Asp	Tyr	Ile	Ile	Gln	Tyr	Thr
		595				600						605			
Val	Lys	Gly	Lys	Pro	Ala	Ile	Tyr	Leu	Lys	Asn	Lys	Ser	Thr	Gly	Tyr
	610					615					620				
Ile	Thr	Tyr	Glu	Asp	Thr	Asn	Gly	Asn	Ser	Glu	Glu	Phe	Gln	Thr	Ile
625					630					635					640
Ala	Val	Lys	Phe	Thr	Ser	Glu	Thr	Asp	Leu	Ser	Gln	Thr	His	Leu	Val
				645					650					655	
Phe	Lys	Ser	Gln	Asn	Gly	Tyr	Glu	Ala	Trp	Gly	Asp	Asn	Phe	Ile	Ile
			660					665					670		
Leu	Glu	Ala	Lys	Leu	Phe	Glu	Thr	Pro	Glu	Ser	Pro	Glu	Leu	Ile	Lys
		675					680					685			
Phe	Asn	Asp	Trp	Glu	Arg	Phe	Gly	Thr	Thr	Tyr	Ile	Thr	Gly	Asn	Glu
	690					695					700				
Leu	Arg	Ile	Asp	His	Ser	Arg	Gly	Gly	Tyr	Phe	Arg	Gln	Ser	Leu	Asn
705					710					715					720
Ile	Asp	Ser	Tyr	Ser	Thr	Tyr	Asp	Leu	Ser	Phe	Ser	Phe	Ser	Gly	Leu
				725					730					735	
Trp	Ala	Lys	Val	Ile	Val	Lys	Asn	Ser	Arg	Gly	Val	Val	Leu	Phe	Glu
			740					745					750		

-continued

Lys Val Lys Asn Asn Gly Ser Ser Tyr Glu Asp Ile Ser Glu Ser Phe
755 760 765

Thr Thr Ala Ser Asn Lys Asp Gly Phe Phe Ile Glu Leu Thr Ala Glu
770 775 780

Arg Thr Ser Ser Thr Phe His Ser Phe Arg Asp Ile Ser Ile Lys Glu
785 790 795 800

Lys Ile Glu His His His His His His
805

<210> SEQ ID NO 11

<211> LENGTH: 2370

<212> TYPE: DNA

<213> ORGANISM: Bacillus thuringiensis

<400> SEQUENCE: 11

```

gtgagggttaa acatgcagaa aaataataaa ttaagtgttaa gggctttacc aagtttcatt    60
gattatttta atgggattta cggattcgcc actggtatca aagatattat gaacatgatt    120
tttaaaacga atacaggagg ggatctaacc ttagacgaaa tattaaaaaa tcaacagtta    180
cttaatgaga tttctggcaa actggatgga gtgaatggca gcttaaatga tcttctcgca    240
caaggaaact tgaatactga attatctaag gaaatattaa aaattgcaaa tgaacagaat    300
agggttttaa atgatgtaaa tacaagcctt gatgcgataa atttaagtct taacacatat    360
ttacctaaaa ttacttctat gttaagtgat gtaatgaagc aaaattatgc attaggtttg    420
cagatagaat acctaagcaa acaattaaag gaaatttcag ataagctaga tgttattaat    480
gtaaatgtac tcattaactt tacacttact gaaattacac ctgcctatca aaggattaaa    540
tatgtaaatg aaaaatttga agcattaacc tctgctacag aaaccaattt aaaaacaaaa    600
caagatagct ctcatacaga tattcttgat gagttaacag agctaacgga actagcgaaa    660
agtgtacaaa aaaatgacgt ggatggcctt gaattttacc ttaatacatt ccacgatgta    720
atgattggga ataacttatt tggacgttca gctttaaaaa cagcctcgga attaattgctg    780
aaagaaaatt tgaaaacaag tggcagtgag gtaggaaatg tttataattt cttaattgta    840
ttaacagctc tgcaagcaaa agcttttctt actttaacta catgccggaa attattgggc    900
ttagcagata ttgattatac tcctattatg aatgaacacc taaataaaga aaaagaggaa    960
ttagagtgta acatccttcc tacactttct aatacttttt ctaatcctaa ttatgaaaaa   1020
gctagagggg gtgataagga tgcgaaaatc attatggaag ctaaacctgg atatgcttta   1080
gttggtattg aaataagtaa ggattcaatt gcagtattaa aagtttatca ggcaaagcta   1140
aaacacaact atcaaattga taaggattcg ttatcagaaa ttgtttatgg tgatatagat   1200
aaattattat gtccggatca atctgaacaa atgtattata caaataaaat agcattttcca   1260
aatgaatatg ttatcactaa aattgctttt actaaaaaac tgaacagttt aagatatgag   1320
gtcacagcga atttttatga ctcttctaca ggagatattg atctaaataa gaaaaaaata   1380
gaatcaagtg aggcggagtt tagtatgcta aatgctaata atgaggggtg ttatatgccg   1440
ataggtacta taagtgaaac atttttgact ccaattaatg gatttggcct cgtagtcgat   1500
gaaaattcaa gactagtaac tttgacatgt aaatcatatt taagagagac attgttagca   1560
acagacttaa gtaataaaga aactaaactg attgtccac ctaatggttt tattagcaat   1620
attgtagaaa atgggaactt agaggagaa aacttagagc cgtggaaagc aaataacaaa   1680

```

-continued

aatgcgtatg tagatcatac cggaggtgta aatggaacta aagttttata tgttcattgag	1740
gatggtgagt tctcacaatt tattggggaa aaattgaaat tgaaaacaga atatgtaatt	1800
caatatattg taaagggaaa agctgctatt tattttaaaag atgaaaaaaa tggggattac	1860
atztatgaag aaacaaataa tgaattagaa gattttcaaa ctgttactaa acgtttttatt	1920
acgggaacag attcttcaag agttcattta atttttacca gtcaaaatgg cgaggaagca	1980
tttgaggaa actttattat ttcagaaatt aggccatccg aagagttatt aagtcagaa	2040
ttgattaagt tggatgcttg ggttgatct cagggaaactt ggatctcagg aaattctctc	2100
aatattaata gtaatgtaaa tggaaacctt cgacaaaacc tttcgttaga aagttattca	2160
acctatagta tgaactttaa tgtgaatgga tttggcaagg tgacaataag aaattctcgt	2220
gaagtagtat ttgaaaggag ttatctacag ttttcctcta aatatatttc agaaaaattc	2280
acaacaacaa ccaataatac tgggttatat gtagaacttt ctctgtcttc gtctggggga	2340
gttataaatt tcggagattt ttcaatcaag	2370

<210> SEQ ID NO 12

<211> LENGTH: 2370

<212> TYPE: DNA

<213> ORGANISM: *Bacillus thuringiensis*

<400> SEQUENCE: 12

atgaacatga acaagaataa tactaaatta agcgcaagag ctttaccaag ttttattgat	60
tattttaatg gcatttatgg atttgccact ggcacaaag atattatgaa catgattttt	120
aaaacggata cagggtggtga tctaacccta gacgaaattt taaagaatca gcagttacta	180
aatgatattt ctggtaaatt ggatggggtg aatggaagct taaatgatct tatcgcacag	240
ggaaacttaa atacagaatt atctaaggaa atattaaaaa tgcacaaatga acaaaatcaa	300
gttttaaatg atgttaataa caaactcgat gcgataaata cgatgcttca tgtatatcta	360
cctaaaaatta cctctatggt aagtgatgta atgaaaccaa attatgcgct aagtatgcaa	420
atagaatacc taagtagaca attacaagaa atttcagata agctagatat tatcaacgta	480
aatgtactta ttaactctac acttactgaa attacacctg cgtatcaatg gattaaatat	540
gtgaacgaaa aatttgaaaga attaactttt gctacagaaa ctactttaaa agtaaaaaat	600
gatagcgctt ctgcagatat tcttgatgag ttaacggagt taactgaact tgcgaaaagt	660
gtaacaaaaa atgatgtgga tgggttttgaa ttttacctta atacattcca cgatgtaatg	720
gtaggaaata atttattcgg gcgttcgact ttaaaaactg cttcggaatt aattgctaaa	780
gaaaatgtga aaacaagtgg cagtgaagta ggaaatgttt ataatttctt agtgttatta	840
acagctctac aagcaaaagc ttttcttact ttaacaacat gccgaaaatt attaggccta	900
gcagatattg attatacatc tattatgaat gaacatttaa ataaggaaaa agaggaaattt	960
agagtaaaca tccttctcat actttccta actttttcta atcctaatta tgcaaaagtt	1020
aaaggaagtg atgaagatgc aaagatgatt gtggaagcta aaccaggaca tgcattgggt	1080
gggtttgaaa ttagtaatga ttcaatgaca gtattaaaag tatatgaagc taagctaaaa	1140
caaaattacc aagttgataa ggattcctta tcggaagtca tttatggtga tatggataaa	1200
ttattgtgcc cagatcaatc tgaaaaaatt tattatacaa ataatatagt atttccaaat	1260
gaatatgtaa ttactaaaat tgattttact aagaaaatga aaactttaag atatgaggta	1320
acagctaatt cttatgatcc ttctacagga gaaattgact taaataaaaa gaaagtagaa	1380

-continued

```

tcaagtaaag cggagtatag gacgttaagt gctaataatg atggagtata tatgccgtta 1440
gggtgcatca gtgaaacatt ttgactcca attaatggat ttggcctcca agctgatgaa 1500
aattcaagat taattacttt aacgtgtaaa tcatatttaa gggaactact actagcgaca 1560
gacttaagca ataaagaaac taaattgatt gtcccgccaa atagttttat tagcaatatt 1620
gtagagaatg ggtccataga agagggccac ttagagcctt ggaaagcaaa taataagaac 1680
gcttatgtag atcacacagg cggagtgaat ggtactaaag ctctatatgt tcatgaggat 1740
ggggggggtt cacaatttat gggagataaa ttaaaaccga aaactgagta tgtaattcaa 1800
tatactgtta aaggaaaacc ttctattcat ttaaaagatg aaaatactgg atatatctt 1860
tatgaagata caaataatga tttagaagat ttccaaacta taactaaaag gttcacaaca 1920
ggaaactgatt taatgagagt gtatttaatt ttaaaaagtc aaagtgggtca cgaagcttgg 1980
ggagataact ttacaatttt ggaaattaag cctgcggagg ctttagtaag cccagaattg 2040
attaatccga attcttggat tacaactcaa ggggctagca tttcaggaga taaacttttt 2100
attagcttgg ggacaaatgg gacctttaga caaaatcttt cattaacag ttattcaact 2160
tatagtataa gctttactgc atcgggacca tttaagtga cggtaagaaa ttctagggaa 2220
gtatttatatg aacgaaacaa ccttatgtct tcaactagtc atatttcttg ggaattcaaa 2280
actgaatcca ataataccgg atttatgtga gaactttccc gtcgttcttg tggtgctggt 2340
catatatcat ttgaaaacat ttctattaaa 2370

```

<210> SEQ ID NO 13

<211> LENGTH: 790

<212> TYPE: PRT

<213> ORGANISM: *Bacillus thuringiensis*

<400> SEQUENCE: 13

```

Met Arg Val Asn Met Gln Lys Asn Asn Lys Leu Ser Val Arg Ala Leu
 1             5             10             15

Pro Ser Phe Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly
 20             25             30

Ile Lys Asp Ile Met Asn Met Ile Phe Lys Thr Asn Thr Gly Gly Asp
 35             40             45

Leu Thr Leu Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile
 50             55             60

Ser Gly Lys Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Leu Ala
 65             70             75             80

Gln Gly Asn Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala
 85             90             95

Asn Glu Gln Asn Arg Val Leu Asn Asp Val Asn Thr Lys Leu Asp Ala
100            105            110

Ile Asn Leu Met Leu Asn Thr Tyr Leu Pro Lys Ile Thr Ser Met Leu
115            120            125

Ser Asp Val Met Lys Gln Asn Tyr Ala Leu Gly Leu Gln Ile Glu Tyr
130            135            140

Leu Ser Lys Gln Leu Lys Glu Ile Ser Asp Lys Leu Asp Val Ile Asn
145            150            155            160

Val Asn Val Leu Ile Asn Phe Thr Leu Thr Glu Ile Thr Pro Ala Tyr
165            170            175

Gln Arg Ile Lys Tyr Val Asn Glu Lys Phe Glu Ala Leu Thr Ser Ala

```

-continued

180						185						190					
Thr	Glu	Thr	Asn	Leu	Lys	Thr	Lys	Gln	Asp	Ser	Ser	His	Thr	Asp	Ile		
		195					200					205					
Leu	Asp	Glu	Leu	Thr	Glu	Leu	Thr	Glu	Leu	Ala	Lys	Ser	Val	Thr	Lys		
	210					215					220						
Asn	Asp	Val	Asp	Gly	Phe	Glu	Phe	Tyr	Leu	Asn	Thr	Phe	His	Asp	Val		
225					230					235					240		
Met	Ile	Gly	Asn	Asn	Leu	Phe	Gly	Arg	Ser	Ala	Leu	Lys	Thr	Ala	Ser		
			245						250					255			
Glu	Leu	Ile	Ala	Lys	Glu	Asn	Leu	Lys	Thr	Ser	Gly	Ser	Glu	Val	Gly		
			260					265					270				
Asn	Val	Tyr	Asn	Phe	Leu	Ile	Val	Leu	Thr	Ala	Leu	Gln	Ala	Lys	Ala		
		275					280					285					
Phe	Leu	Thr	Leu	Thr	Thr	Cys	Arg	Lys	Leu	Leu	Gly	Leu	Ala	Asp	Ile		
	290					295					300						
Asp	Tyr	Thr	Pro	Ile	Met	Asn	Glu	His	Leu	Asn	Lys	Glu	Lys	Glu	Glu		
305					310					315					320		
Phe	Arg	Val	Asn	Ile	Leu	Pro	Thr	Leu	Ser	Asn	Thr	Phe	Ser	Asn	Pro		
			325						330					335			
Asn	Tyr	Glu	Lys	Ala	Arg	Gly	Ser	Asp	Lys	Asp	Ala	Lys	Ile	Ile	Met		
		340						345					350				
Glu	Ala	Lys	Pro	Gly	Tyr	Ala	Leu	Val	Gly	Phe	Glu	Ile	Ser	Lys	Asp		
		355					360					365					
Ser	Ile	Ala	Val	Leu	Lys	Val	Tyr	Gln	Ala	Lys	Leu	Lys	His	Asn	Tyr		
	370					375					380						
Gln	Ile	Asp	Lys	Asp	Ser	Leu	Ser	Glu	Ile	Val	Tyr	Gly	Asp	Ile	Asp		
385					390					395					400		
Lys	Leu	Leu	Cys	Pro	Asp	Gln	Ser	Glu	Gln	Met	Tyr	Tyr	Thr	Asn	Lys		
			405						410					415			
Ile	Ala	Phe	Pro	Asn	Glu	Tyr	Val	Ile	Thr	Lys	Ile	Ala	Phe	Thr	Lys		
		420						425					430				
Lys	Leu	Asn	Ser	Leu	Arg	Tyr	Glu	Val	Thr	Ala	Asn	Phe	Tyr	Asp	Ser		
		435					440					445					
Ser	Thr	Gly	Asp	Ile	Asp	Leu	Asn	Lys	Lys	Lys	Ile	Glu	Ser	Ser	Glu		
	450					455					460						
Ala	Glu	Phe	Ser	Met	Leu	Asn	Ala	Asn	Asn	Glu	Gly	Val	Tyr	Met	Pro		
465					470					475					480		
Ile	Gly	Thr	Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	Ile	Asn	Gly	Phe	Gly		
			485						490					495			
Leu	Val	Val	Asp	Glu	Asn	Ser	Arg	Leu	Val	Thr	Leu	Thr	Cys	Lys	Ser		
		500						505					510				
Tyr	Leu	Arg	Glu	Thr	Leu	Leu	Ala	Thr	Asp	Leu	Ser	Asn	Lys	Glu	Thr		
	515						520					525					
Lys	Leu	Ile	Val	Pro	Pro	Asn	Gly	Phe	Ile	Ser	Asn	Ile	Val	Glu	Asn		
	530					535					540						
Gly	Asn	Leu	Glu	Gly	Glu	Asn	Leu	Glu	Pro	Trp	Lys	Ala	Asn	Asn	Lys		
545					550					555					560		
Asn	Ala	Tyr	Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Lys	Val	Leu		
			565					570						575			
Tyr	Val	His	Glu	Asp	Gly	Glu	Phe	Ser	Gln	Phe	Ile	Gly	Glu	Lys	Leu		
			580					585					590				

-continued

Lys Leu Lys Thr Glu Tyr Val Ile Gln Tyr Ile Val Lys Gly Lys Ala
 595 600 605
 Ala Ile Tyr Leu Lys Asp Glu Lys Asn Gly Asp Tyr Ile Tyr Glu Glu
 610 615 620
 Thr Asn Asn Glu Leu Glu Asp Phe Gln Thr Val Thr Lys Arg Phe Ile
 625 630 635 640
 Thr Gly Thr Asp Ser Ser Arg Val His Leu Ile Phe Thr Ser Gln Asn
 645 650 655
 Gly Glu Glu Ala Phe Gly Gly Asn Phe Ile Ile Ser Glu Ile Arg Pro
 660 665 670
 Ser Glu Glu Leu Leu Ser Pro Glu Leu Ile Lys Leu Asp Ala Trp Val
 675 680 685
 Gly Ser Gln Gly Thr Trp Ile Ser Gly Asn Ser Leu Asn Ile Asn Ser
 690 695 700
 Asn Val Asn Gly Thr Phe Arg Gln Asn Leu Ser Leu Glu Ser Tyr Ser
 705 710 715 720
 Thr Tyr Ser Met Asn Phe Asn Val Asn Gly Phe Gly Lys Val Thr Ile
 725 730 735
 Arg Asn Ser Arg Glu Val Val Phe Glu Arg Ser Tyr Leu Gln Phe Ser
 740 745 750
 Ser Lys Tyr Ile Ser Glu Lys Phe Thr Thr Thr Thr Asn Asn Thr Gly
 755 760 765
 Leu Tyr Val Glu Leu Ser Arg Ala Ser Ser Gly Gly Val Ile Asn Phe
 770 775 780
 Gly Asp Phe Ser Ile Lys
 785 790

<210> SEQ ID NO 14

<211> LENGTH: 790

<212> TYPE: PRT

<213> ORGANISM: Bacillus thuringiensis

<400> SEQUENCE: 14

Met Asn Met Asn Lys Asn Asn Thr Lys Leu Ser Ala Arg Ala Leu Pro
 1 5 10 15
 Ser Phe Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile
 20 25 30
 Lys Asp Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu
 35 40 45
 Thr Leu Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser
 50 55 60
 Gly Lys Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln
 65 70 75 80
 Gly Asn Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn
 85 90 95
 Glu Gln Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile
 100 105 110
 Asn Thr Met Leu His Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser
 115 120 125
 Asp Val Met Lys Pro Asn Tyr Ala Leu Ser Met Gln Ile Glu Tyr Leu
 130 135 140
 Ser Arg Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val
 145 150 155 160

-continued

Asn	Val	Leu	Ile	Asn	Ser	Thr	Leu	Thr	Glu	Ile	Thr	Pro	Ala	Tyr	Gln
				165					170					175	
Trp	Ile	Lys	Tyr	Val	Asn	Glu	Lys	Phe	Glu	Glu	Leu	Thr	Phe	Ala	Thr
			180					185					190		
Glu	Thr	Thr	Leu	Lys	Val	Lys	Asn	Asp	Ser	Ala	Ser	Ala	Asp	Ile	Leu
		195					200					205			
Asp	Glu	Leu	Thr	Glu	Leu	Thr	Glu	Leu	Ala	Lys	Ser	Val	Thr	Lys	Asn
	210					215					220				
Asp	Val	Asp	Gly	Phe	Glu	Phe	Tyr	Leu	Asn	Thr	Phe	His	Asp	Val	Met
	225				230					235					240
Val	Gly	Asn	Asn	Leu	Phe	Gly	Arg	Ser	Thr	Leu	Lys	Thr	Ala	Ser	Glu
				245					250						255
Leu	Ile	Ala	Lys	Glu	Asn	Val	Lys	Thr	Ser	Gly	Ser	Glu	Val	Gly	Asn
			260					265					270		
Val	Tyr	Asn	Phe	Leu	Val	Val	Leu	Thr	Ala	Leu	Gln	Ala	Lys	Ala	Phe
		275					280					285			
Leu	Thr	Leu	Thr	Thr	Cys	Arg	Lys	Leu	Leu	Gly	Leu	Ala	Asp	Ile	Asp
	290					295					300				
Tyr	Thr	Ser	Ile	Met	Asn	Glu	His	Leu	Asn	Lys	Glu	Lys	Glu	Glu	Phe
	305				310					315					320
Arg	Val	Asn	Ile	Leu	Pro	Ile	Leu	Ser	Asn	Thr	Phe	Ser	Asn	Pro	Asn
				325					330					335	
Tyr	Ala	Lys	Val	Lys	Gly	Ser	Asp	Glu	Asp	Ala	Lys	Met	Ile	Val	Glu
		340						345					350		
Ala	Lys	Pro	Gly	His	Ala	Leu	Val	Gly	Phe	Glu	Ile	Ser	Asn	Asp	Ser
		355					360					365			
Met	Thr	Val	Leu	Lys	Val	Tyr	Glu	Ala	Lys	Leu	Lys	Gln	Asn	Tyr	Gln
	370					375					380				
Val	Asp	Lys	Asp	Ser	Leu	Ser	Glu	Val	Ile	Tyr	Gly	Asp	Met	Asp	Lys
	385				390					395					400
Leu	Leu	Cys	Pro	Asp	Gln	Ser	Glu	Lys	Ile	Tyr	Tyr	Thr	Asn	Asn	Ile
				405					410					415	
Val	Phe	Pro	Asn	Glu	Tyr	Val	Ile	Thr	Lys	Ile	Asp	Phe	Thr	Lys	Lys
			420					425					430		
Met	Lys	Thr	Leu	Arg	Tyr	Glu	Val	Thr	Ala	Asn	Ser	Tyr	Asp	Ser	Ser
		435					440					445			
Thr	Gly	Glu	Ile	Asp	Leu	Asn	Lys	Lys	Lys	Val	Glu	Ser	Ser	Lys	Ala
	450					455					460				
Glu	Tyr	Arg	Thr	Leu	Ser	Ala	Asn	Asn	Asp	Gly	Val	Tyr	Met	Pro	Leu
	465				470					475					480
Gly	Val	Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	Ile	Asn	Gly	Phe	Gly	Leu
				485					490					495	
Gln	Ala	Asp	Glu	Asn	Ser	Arg	Leu	Ile	Thr	Leu	Thr	Cys	Lys	Ser	Tyr
			500					505					510		
Leu	Arg	Glu	Leu	Leu	Leu	Ala	Thr	Asp	Leu	Ser	Asn	Lys	Glu	Thr	Lys
		515					520					525			
Leu	Ile	Val	Pro	Pro	Asn	Ser	Phe	Ile	Ser	Asn	Ile	Val	Glu	Asn	Gly
		530				535					540				
Ser	Ile	Glu	Glu	Gly	His	Leu	Glu	Pro	Trp	Lys	Ala	Asn	Asn	Lys	Asn
	545				550					555					560
Ala	Tyr	Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Lys	Ala	Leu	Tyr
				565					570					575	

-continued

Val His Glu Asp Gly Gly Val Ser Gln Phe Met Gly Asp Lys Leu Lys
580 585 590

Pro Lys Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser
595 600 605

Ile His Leu Lys Asp Glu Asn Thr Gly Tyr Ile Leu Tyr Glu Asp Thr
610 615 620

Asn Asn Asp Leu Glu Asp Phe Gln Thr Ile Thr Lys Arg Phe Thr Thr
625 630 635 640

Gly Thr Asp Leu Met Arg Val Tyr Leu Ile Leu Lys Ser Gln Ser Gly
645 650 655

His Glu Ala Trp Gly Asp Asn Phe Thr Ile Leu Glu Ile Lys Pro Ala
660 665 670

Glu Ala Leu Val Ser Pro Glu Leu Ile Asn Pro Asn Ser Trp Ile Thr
675 680 685

Thr Gln Gly Ala Ser Ile Ser Gly Asp Lys Leu Phe Ile Ser Leu Gly
690 695 700

Thr Asn Gly Thr Phe Arg Gln Asn Leu Ser Leu Asn Ser Tyr Ser Thr
705 710 715 720

Tyr Ser Ile Ser Phe Thr Ala Ser Gly Pro Phe Asn Val Thr Val Arg
725 730 735

Asn Ser Arg Glu Val Leu Tyr Glu Arg Asn Asn Leu Met Ser Ser Thr
740 745 750

Ser His Ile Ser Gly Glu Phe Lys Thr Glu Ser Asn Asn Thr Gly Leu
755 760 765

Tyr Val Glu Leu Ser Arg Arg Ser Gly Gly Ala Gly His Ile Ser Phe
770 775 780

Glu Asn Ile Ser Ile Lys
785 790

<210> SEQ ID NO 15

<211> LENGTH: 2409

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic sequence encoding AXMI-115

<400> SEQUENCE: 15

```

atgaacatga acaacaccaa gctcaatgca agagctcttc cttccttcat cgactacttc      60
aatggcatct atggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagaca      120
gacactggag gagatctcac cttggatgag atcctcaaga accagcagct gctgaatgag      180
atctcagggg agctggatgg cgtcaatgga agcctcaatg atctcattgc tcaaggaaac      240
ctcaacacag agctctccaa ggagatcctc aagatcgcca atgagcagaa ccagggtgctg      300
aatgatgtca acaacaagct ggatgccatc aacaccatgc tgaacatcta cctccccaag      360
atcacctcaa tgctctctga tgtgatgaag cagaactatg ctctctccct ccagatcgag      420
tacctctcaa ggcagctgca agagatctcc gacaagctgg atgtcatcaa cctcaatgtg      480
ctgatcaaca gcaccttgac agagatcacc ccaagctacc agaggatcaa atatgtcaat      540
gagaagttcg acaagctcac cttcgccaca gaatcaaccc tccgcgccaa gcaaggcatc      600
ttcaatgagg acagcttcga caacaacacc ttggagaact tgacagatct tgctgagctg      660
gccaaagagc tcaccaagaa tgatgtggac agcttcgagt tctacctcca caccttccat      720

```

-continued

gatgtgctga ttggcaacaa cctcttttga agaagcgcgc tcaagacagc ttcagagctc	780
atcaccaagg atgagatcaa gacatctgga tcagagattg gcaaggtgta cagcttcttc	840
atcgtctctca ccagcctcca ggccaaggcc ttctcacc ccaccacctg ccggaagctg	900
ctgggcctct ccgacatcga ctacacctcc atcatgaatg agcacctcaa caatgagaag	960
aatgagttca gagacaacat cctgcggcg ctctccaaca agttcagcaa ccttcatat	1020
gcaaaaacca tggcagcga caactacgcc aaggtgatcc tggagagcga gcctggatat	1080
gctctggtgg gcttcgagat catcaatgat cccatccccg tgctgaaggc ctacaaggcc	1140
aagctgaagc agaactacca ggtggacaac cagagcctct cagagatcgt ctacctggac	1200
atcgacaagc tcttctgccc agagaacagc gagcagaagt actacaccaa gaacctcacc	1260
ttcctgatg gatatgtcat caccaagatc acctcgcga agaagctcaa caacctcatc	1320
tatgaagcca ccgccactt ctatgatcca tcaactggag acatcgacct gaacaagaag	1380
cagggtggaga gcaccttccc taaaacagac tacatcacca tggacattgg agatgatgat	1440
ggcatctaca tgccgctcgg cgtcatctca gaaaccttcc tccccccat caacagcttc	1500
ggcctggagg tggatgcaa gagcaagacc ctcaccttga aatgcaagag ctacctcagg	1560
gagtacctgc tggagagtga tctgaagaac aaggaaacag ggctgatcgc gccgccaaat	1620
gttttcatca gcaatgtggt gaagaactgg gacattgaag aagattcatt ggagccatgg	1680
gtggccaaca acaagaatgc ttatgtggac aacaccggcg gcattgaaag aagcaaggcg	1740
ctcttcaccc aaggagatgg agagtcttca cagttcatcg gcgacaagct gaagccaaac	1800
accgactaca tcatccagta caccgtcaag ggcaagccgg ccatctacct caagaacaag	1860
agcaccggct acatcacata tgaggacacc aatggcaaca gcgaggagtt ccaaacatt	1920
gctgtgaagt tcacctcaga aacagatctc tcccaaacc ccttggtgtt caagagccaa	1980
aatggatatg aagcatgggg agacaacttc atcatcctgg aggccaaagct ctttgaaca	2040
ccagaatctc cagagctcat caagttaat gattgggaga gatttggcac cacctacatc	2100
actggaaatg agctgaggat tgatcattca agaggaggct acttcgcga gagcttgaac	2160
atcgacagct acagcacata tgatctctcc ttctccttct ccggcctctg ggccaaggtc	2220
atcgtcaaga acagcagagg agtgggtgctc ttcgagaagg tgaagaacaa tggaagcagc	2280
tatgaggaca tctcagagag cttcaccacc gccagcaaca aggatggctt cttcatcgag	2340
ctcaccgccc agaggacaag cagcaccttc cacagcttca gagacatctc catcaaggag	2400
aagatcgag	2409

<210> SEQ ID NO 16

<211> LENGTH: 2409

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic sequence encoding AXMI-115

<400> SEQUENCE: 16

atgaacatga acaacaccaa gctcaatgca agagctcttc cttccttcat cgactacttc	60
aatggcatct atggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagaca	120
gacactggag gagatctcac cttggatgag atcctcaaga accagcagct gctgaatgag	180
atctcagggg agctggatgg cgtcaatgga agcctcaatg atctcattgc tcaaggaaac	240
ctcaacacag agctctccaa ggagatcctc aagattgcaa atgagcagaa ccaggtgctg	300

-continued

aatgatgtca acaacaagct ggaatgccatc aacaccatgc tgaacatcta cctccccaag	360
atcacctcaa tgctctctga tgtgatgaag cagaactatg ctctctccct ccagatcgag	420
tacctctcaa ggcagctgca agagatctcc gacaagctgg atgtcatcaa cctcaatgtg	480
ctgatcaaca gcaccttgac agagatcacc ccaagctacc aaaggatcaa atatgtcaat	540
gagaagtctg acaagctcac cttcgccaca gaatcaaccc tccgcgcca gcaaggcatc	600
ttcaatgagg acagcttcga caacaacacc ttggagaact tgacagatct tgctgagctg	660
gccaaagaca tcaccaagaa tgatgtggac agcttcgagt tctacctcca caccttccat	720
gatgtgctga ttggcaacaa cctcttttga agaagcgcgc tcaagacagc ttcagagctc	780
atcaccaagg atgagatcaa gacatctgga tcagaaattg gcaagggtga cagcttcttc	840
atcgtctcca ccagcttcca ggccaaggcc ttctcacc ccaccactg ccggaagctg	900
ctgggcctct ccgacatcga ctacacctcc atcatgaatg agcacctcaa caatgagaag	960
aatgagttca gagacaacat cctgcccggc ctctccaaca agttcagcaa ccttccatat	1020
gcaaaaacca tgggcagcga caactatgcc aaggatgatc tggagagcga gcctggatat	1080
gctctggtgg gcttcgagat catcaatgat cccatccctg tgctgaaggc ctacaaggcc	1140
aagctgaagc agaactacca ggtggacaac caaagcctct cagagatcgt ctacctggac	1200
atcgacaagc tcttctgccc agagaacagc gagcagaagt actacaccaa gaacctcacc	1260
ttccctgatg gatatgtcat caccaagatc accttcgaga agaagctcaa caacctcatc	1320
tatgaagcca ccgccaactt ctatgatcca tcaactggag acattgatct gaacaagaag	1380
cagggtggaga gcaccttccc taaaacagat tacatcacca tggacattgg agatgatgat	1440
ggcatctaca tgccgctcgg cgtcatctca gaaaccttcc tccccccat caacagcttc	1500
ggcctggagg tggatgcca gagcaagacc ctcaccttga aatgcaagag ctacttgagg	1560
gagtacctgc tggaaatcaga tctgaagaac aaggaaacag ggctgatcgc gccgccaaat	1620
gttttcatca gcaatgtggt gaagaactgg gacattgaag aagattcatt ggagccatgg	1680
gtggccaaca acaagaatgc ttatgtggac aacaccggcg gcattgaaag aagcaaggcg	1740
ctcttcaccc aaggagatgg agagtctcca cagttcatcg gcgacaagct gaagccaaac	1800
accgactaca tcatccagta caccgtcaag ggcaagccgg ccactctacct caagaacaag	1860
agcactggct acatcacata tgaggacacc aatggcaaca gcgaggagtt ccaacaatt	1920
gctgtgaagt tcacctcaga aacagatctc tcccaaacc ccttggtgtt caagagccaa	1980
aatggatatg aagcatgggg agacaacttc atcatcctgg aggccaaagct ctttgaacaa	2040
ccagaatctc cagagctcat caagttcaat gattgggaaa gatttggcac cacctacatc	2100
actggaaatg agctgaggat tgatcattca agaggaggct acttccgcca gagcttgaac	2160
atcgacagct acagcacata tgatctctcc ttctccttct ccggcctctg ggccaaggtc	2220
atcgtcaaga acagcagagg agtggtgctc ttcgagaagg tgaagaacaa tggaaagcagc	2280
tatgaggaca tctcagagag cttcaccacc gccagcaaca aggatggctt cttcatcgag	2340
ctcaccgccc agaggacaag cagcaccttc cacagcttca gagacatctc catcaaggag	2400
aagattgaa	2409

<210> SEQ ID NO 17

<211> LENGTH: 2364

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic sequence encoding AXMI-184

<400> SEQUENCE: 17

```

atgaacaaga acaacaccaa gctctccgcg cgcgcgctgc cctccttcat cgactacttc      60
aatggcatct atggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagaca      120
gacactggag gagatctcac cttggatgag atcctcaaga accagcagct gctgaatgac      180
atctcagggg agctggatgg cgtcaatgga agcctcaatg atctcattgc tcaaggaaac      240
ctcaacacag agctctccaa ggagatcctc aagatcgcca atgagcagaa ccaggtgctg      300
aatgatgtca acaacaagct ggatgccatc aacaccatgc tgcattgtcta cctccccaag      360
atcacctcaa tgctctctga tgtgatgaag ccaaactatg ctctctccat gcaaattgag      420
tacctctcaa ggcagctgca agagatctcc gacaagctgg acatcatcaa tgtcaatgtg      480
ctgatcaaca gcaccttgac agagatcacg cgggectacc aatggatcaa atatgtcaat      540
gagaagtttg aggagctcac cttcgccaca gaaacaacct tgaagggtgaa gaatgattca      600
gcttctgctg acatcctgga tgagctgaca gagctgacag agctggccaa gagcgtcacc      660
aagaatgatg ttgatggctt cgagttctac ctcaacacct tccatgatgt gatggtgggc      720
aacaacctct ttggaagaag caccctcaag acagcttcag agctgatcgc caaggagaat      780
gtcaagacat ctggaagcga ggtgggaaat gtctacaact tcctgggtgt gctgacggcg      840
ctgcaagcaa aggccttctc caccctcacc acctgccgga agctgctggg cctcgccgac      900
atcgactaca cctcaatcat gaatgagcac ctcaacaagg agaaggagga gttcagagtg      960
aacatcctcc ccatcctctc caacaccttc agcaacccca actacgccaa ggtgaagggc      1020
tcagatgaag atgccaagat gattgtggag gccaaagctg gccatgctct ggtgggcttc      1080
gagatcagca atgacagcat gacggtgctg aaggtgtatg aagcaaagct gaagcagAAC      1140
taccaggtgg acaagacag cctctccgag gtgatctatg gagacatgga caagctgctc      1200
tgcccagatc aatcagagaa gatctactac accaacaaca tcgtctttcc aatgaatat      1260
gtcatcacca agatcgactt caccaagaag atgaaaacct tgagatatga agtcaccgcc      1320
aacagctatg attcttcaac tggagagatc gacctcaaca agaagaaggt ggagagcagc      1380
aaggcagagt acaggacgct ctccgccaac aatgatggcg tctacatgcc gctcggcgtc      1440
atctcagaaa cttctctgac gcccatcaat ggcttcggcc tccaagctga tgagaacagc      1500
aggctcatca ccctcacctg caagagctac ctcagggagc tgctgctggc caccgacctc      1560
tccaacaagg aaacaagct gattgttctc ccaaacagct tcatcagcaa catcgtggag      1620
aatggaagca ttgaagaagg ccacctagag ccatggaagg ccaacaacaa gaatgcatat      1680
gtggaccaca cggcgcgctt caatggcacc aaggcgctct atgttcatga agatggagga      1740
gtttcacagt tcatgggaga caagctgaag ccaaaaacag aatatgtcat ccagtacacc      1800
gtcaagggca agccaagcat ccacctcaag gatgagaaca cgggctacat cctctacgag      1860
gacaccaaca atgatttaga agatttccaa accatcacca agagggtcac cactggaaca      1920
gatctgatga ggggtgtacct catctcaag agccaagtg gtcattgaagc atggggagac      1980
aacttcacca tcctggagat caagccagca gaagctctgg tgcgcccga gctcatcaac      2040
cccaacagct ggataacaac acaaggagct tcaattttctg gtgacaagct cttcatctcc      2100
cttggaaaca atggcacctt ccgccagAAC ctctccctca acagctacag cacctacagc      2160

```

-continued

atcagcttca cgcctcagg ccccttcaat gtcaccgtca ggaacagcag ggaggtgctg	2220
tatgaaagga acaacttgat gagcagcacc agccacatct ctggagagtt caagacagaa	2280
agcaacaaca cgggcctcta tgtggagctc tcaagaagaa gcggcggcgc cggccacatc	2340
agcttcgaga acatctccat caag	2364

<210> SEQ ID NO 18
 <211> LENGTH: 2364
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic sequence encoding AXMI-184

<400> SEQUENCE: 18

atgaacaaga acaacaccaa gctctccgcg cgcgcgctgc catccttcat cgactacttc	60
aatggcatct atggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagaca	120
gacactggag gatgctctac cttggatgag atcctcaaga accagcagct gctgaatgac	180
atctcagggg agctggatgg cgtcaatgga agcctcaatg atctcattgc tcaaggaaac	240
ctcaacacag agctctccaa ggagatcctc aagattgcaa atgagcagaa ccaggtgctg	300
aatgatgtca acaacaagct ggatgccatc aacaccatgc tgcattgtct cctccccaag	360
atcacctcaa tgctctctga tgtgatgaag ccaaattatg ctctctccat gcaaattgag	420
tacctctcaa ggcagctgca agagatctcc gacaagctgg acatcatcaa tgtcaatggt	480
ctcatcaaca gcaccttgac agagatcacg ccggcctacc aatggatcaa atatgtcaat	540
gagaagtttg aggagctcac cttcgccaca gaaacaacat tgaaggtgaa gaatgattca	600
gcttctgctg acatcctgga tgagctgaca gagctgacag agctggccaa gagcgtcacc	660
aagaatgatg ttgatggctt cgagttctac ctcaacacct tccatgatgt gatggtgggc	720
aacaacctct ttggaagaag caccctcaag acagcttcag agctgatcgc caaggaaaat	780
gtcaagacat ctggatcaga ggttggaat gtctacaact tcttggtggt gctgacggcg	840
ctgcaagcaa aggccttctt caccctcacc acctgccgga agctgctggg cctcgccgac	900
atcgactaca cctcaatcat gaatgagcac ctcaacaagg agaaggagga gttcagagtg	960
aacatcctcc ccatcctctc caacaccttc agcaacccca actatgccaa ggtgaagggc	1020
tcagatgaag atgccaagat gattgtggag gccaaagctg gccatgctct ggtgggcttc	1080
gagatcagca atgacagcat gacagtgtg aaggtttatg aagcaaagct gaagcagaac	1140
taccaggtgg acaaggacag cctctcagag gtgatctatg gagacatgga caagctgctc	1200
tgcccagatc aatcagagaa gatctactac accaacaaca tcgtctttcc aatgaatat	1260
gtcatcacca agatcgactt caccaagaag atgaaaacat tgagatatga agtcaccgcc	1320
aacagctatg attcttcaac tggagagatc gacctcaaca agaagaaggt ggagagcagc	1380
aaggcagagt acaggacgct ctccgccaac aatgatggcg tctacatgcc gctcggcgctc	1440
atctcagaaa ctttcttgac gcccatcaat ggcttcggcc tccaagctga tgaaaacagc	1500
aggctcatca ccctcacctg caagagctac ttgagggagc tgctgctggc caccgacctc	1560
tccaacaagg aaacaaagct gattgttctt ccaaacagct tcatcagcaa cattgtggag	1620
aatggaagca ttgaagaagg ccacctagag ccatggaagg ccaacaacaa gaatgcatat	1680
gttgatcaca ccggcgcgct caatggcacc aaggcgctct atgttcatga agatggagga	1740

-continued

gtttcacagt tcatgggaga caagctgaag ccaaaaacag aatatgtcat ccagtacacc 1800
gtcaagggca agccaagcat ccacctcaag gatgaaaaca ctggctacat cctctatgag 1860
gacaccaaca atgatttaga agattttcaa accatcacca agagggtcac cactggaaca 1920
gatctgatga ggggtgtacct catcctcaag agccaaagtg gtcatgaagc atggggagac 1980
aacttcacca tcctggagat caagccagca gaagctctgg tgcgccgga gtcctcaaac 2040
cccaacagct ggataacaac acaaggagct tcaatttctg gtgacaagct cttcatctcc 2100
cttggaaaca atggcacctt ccgccagaac ctctccctca acagctacag cacctacagc 2160
atcagcttca ccgcctcagg acccttcaat gtcaccgtca ggaacagcag ggaggtgctg 2220
tatgaaagga acaacttgat gagcagcacc agccacatct ctggagagtt caagacagaa 2280
agcaacaaca cgggcctcta tgtggagctc tcaagaagaa gcggcggcgc cggccacatc 2340
agcttcgaga acatctccat caag 2364

<210> SEQ ID NO 19
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: endoplasmic reticulum targeting peptide

<400> SEQUENCE: 19

Lys Asp Glu Leu
1

<210> SEQ ID NO 20
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 20

aacaccggcg gcgtcaatgg aacaaggcgc ctcttcaccc a 41

<210> SEQ ID NO 21
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 21

tgggtgaaga gcgcccttgt tccattgacg ccgccgtgt t 41

<210> SEQ ID NO 22
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 22

gccccgagct catcaatgtc aacaactgga tcagaactgg caccacctac atcac 55

<210> SEQ ID NO 23
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 23

gtgatgtagg tgggtgccagt tctgatccag ttgttgacat tgatgagctc cgggc 55

<210> SEQ ID NO 24

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 24

atgattggga gaggttcgga agcaccacaca tcagcggcaa tgagctgagg 50

<210> SEQ ID NO 25

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 25

cctcagctca ttgccgtga tgtgggtgct tccgaacctc tcccaatcat 50

<210> SEQ ID NO 26

<211> LENGTH: 55

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 26

ctacatcacc ggcaatacct tgacgtctta ccaaggagga ggaggctact tccgc 55

<210> SEQ ID NO 27

<211> LENGTH: 55

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 27

gcggaagtag cctcctcctc cttggtagag cgtcaaggta ttgccggtga tntag 55

<210> SEQ ID NO 28

<211> LENGTH: 57

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 28

cgacagctac agcacctaca gggatgaactt ctccgtcacc ggctggggcca aggtgat 57

<210> SEQ ID NO 29

<211> LENGTH: 57

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 29

atcaccttgg cccagccggt gacggagaag ttcaccctgt aggtgctgta gctgtcg 57

-continued

<210> SEQ ID NO 30
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 30
gcttcagcgg cctcgacgcc aatgtgagga tcagaaacag ccgcggc 47

<210> SEQ ID NO 31
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 31
gccgcggctg tttctgatcc tcacattggc gtcgaggccg ctgaagc 47

<210> SEQ ID NO 32
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 32
gtgaagaaca gccgcgaggt gctcttcgag aagagataca tgaatggaag cagctatga 59

<210> SEQ ID NO 33
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 33
tcatagctgc ttccattcat gtatctcttc tcgaagagca cctcgcggtt gttcttcac 59

<210> SEQ ID NO 34
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 34
ttcgagaagg tgaagaacag cggcgccaag gatgtttcag agagcttcac cacc 54

<210> SEQ ID NO 35
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 35
ggtggtgaag ctctctgaaa catccttggc gccgctgttc ttcaccttct cgaa 54

<210> SEQ ID NO 36
<211> LENGTH: 53
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 36

gcttcttcat cgagctcagc caaggcaaca acctctatag cagcaccttc cac 53

<210> SEQ ID NO 37
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 37

gtggaaggty ctgctataga ggttggtgcc ttggctgagc tcgatgaaga agc 53

<210> SEQ ID NO 38
<211> LENGTH: 51
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 38

gaagcaaggc gctctatggt cacaaggatg gaggcttcag ccagttcatc g 51

<210> SEQ ID NO 39
<211> LENGTH: 51
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 39

cgatgaactg gctgaagcct ccatccttgt gaacatagag cgccttgctt c 51

<210> SEQ ID NO 40
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 40

ccgccgagag gacaggaggg ccgctgggtga agttcagaga catcagcatc 50

<210> SEQ ID NO 41
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 41

gatgctgatg tctctgaact tcaccagcgg ccctcctgtc ctctcgcgcg 50

<210> SEQ ID NO 42
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 42

-continued

agcaccttcc acagcttcaa tgatgtgagc atcaagtaag gcgcgccg 48

<210> SEQ ID NO 43
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 43

cgggcgccct tacttgatgc tcacatcatt gaagctgtgg aaggtgct 48

<210> SEQ ID NO 44
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 44

cgacaagcta aagcccaaga cagaatatgt catccagtac accgtcaag 49

<210> SEQ ID NO 45
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 45

cttgacgggtg tactggatga catattctgt cttgggcttt agcttgctg 49

<210> SEQ ID NO 46
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 46

cctacgagga caccaataac aacaacctgg aggactacca aacaattgct gtgaag 56

<210> SEQ ID NO 47
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 47

cttcacagca attgtttggt agtcctccag gttgttgta ttggtgtcct cgtagg 56

<210> SEQ ID NO 48
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 48

gaggagtcc aaacaattac caagaggttc accaccggca cagatttgag ccagacc 57

<210> SEQ ID NO 49

-continued

<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 49

ggctcggctc aaatctgtgc cggcgggtgaa cctcttggtta attgtttgga actcctc 57

<210> SEQ ID NO 50
<211> LENGTH: 58
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 50

cacctcagaa acagatttga agggcgtcta cctcatcttg aagagccaaa atggatat 58

<210> SEQ ID NO 51
<211> LENGTH: 58
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 51

atatccattt tggtcttcca agatgaggta gacgcccttc aaatctgttt ctgagggtg 58

<210> SEQ ID NO 52
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 52

tcctggaggc caagccatca gagaagctgc tcagcccgga gctca 45

<210> SEQ ID NO 53
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 53

tgagctccgg gctgagcagc ttctctgatg gcttggcctc cagga 45

<210> SEQ ID NO 54
<211> LENGTH: 67
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 54

atcattcaag aggaggcaac ctcaagcaga acctccagct tgacagcttc agcacctacg 60
acctcag 67

<210> SEQ ID NO 55
<211> LENGTH: 67
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 55

ctgaggtcgt aggtgctgaa gctgtcaagc tggaggttct gcttgagggt gectcctctt    60
gaatgat                                                                    67

<210> SEQ ID NO 56
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 56

gctatgagga catctcagag atcttcacca ccaagctggg caaggacaac ttctacatcg    60
agctcaccgc                                                                70

<210> SEQ ID NO 57
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 57

gcggtgagct cgatgtagaa gttgtccttg cccagcttgg tggatgaagat ctctgagatg    60
tcctcatagc                                                                70

<210> SEQ ID NO 58
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 58

caagggaag ccgtcaatcc acctcaagaa tgagaacacc ggctacatcc actacgagga    60
caccaatgg                                                                69

<210> SEQ ID NO 59
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 59

ccattggtgt cctcgtagt gatgtagccg gtgttctcat tcttgagggt gattgacggc    60
ttgcccttg                                                                69

<210> SEQ ID NO 60
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 60

caagagccaa aatggagatg aagcatgggg agacaacttc accatcctgg agatctcgct    60

```

-continued

cttcgagaca ccagaa

76

<210> SEQ ID NO 61

<211> LENGTH: 76

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer for domain swapping

<400> SEQUENCE: 61

ttctggtgtc tcgaagagcg agatctccag gatggtgaag ttgtctcccc atgcttcac

60

tccattttgg ctcttg

76

That which is claimed:

1. An isolated or recombinant nucleic acid molecule comprising a nucleotide sequence encoding an insecticidal polypeptide having at least 95% sequence identity SEQ ID NO:6.

2. The nucleic acid molecule of claim 1, wherein said nucleotide sequence is selected from the group consisting of SEQ ID NO:15 and 16.

3. An expression cassette comprising the nucleic acid molecule of claim 1.

4. The expression cassette of claim 3, further comprising a nucleic acid molecule encoding a heterologous polypeptide.

5. A host cell that contains the recombinant nucleic acid molecule of claim 1.

6. An isolated polypeptide having at least 95% sequence identity SEQ ID NO:6, wherein said polypeptide has insecticidal activity.

7. The polypeptide of claim 6 further comprising heterologous amino acid sequences.

8. An antibody that selectively binds to the polypeptide of claim 6.

9. A composition comprising the polypeptide of claim 6.

10. The composition of claim 9, wherein said composition is selected from the group consisting of a powder, dust, pellet, granule, spray, emulsion, colloid, and solution.

11. The composition of claim 9, wherein said composition is prepared by desiccation, lyophilization, homogenization,

extraction, filtration, centrifugation, sedimentation, or concentration of a culture of *Bacillus thuringiensis* cells.

12. The composition of claim 9, comprising from about 1% to about 99% by weight of said polypeptide.

13. A plant having stably incorporated into its genome a DNA construct comprising a nucleotide sequence encoding an insecticidal polypeptide having at least 95% sequence identity SEQ ID NO:6, wherein said nucleotide sequence is operably linked to a promoter that drives expression of a coding sequence in a plant cell.

14. The plant of claim 13, wherein said plant is a plant cell.

15. A transgenic seed of the plant of claim 13.

16. The plant of claim 13, wherein said plant is selected from the group consisting of maize, sorghum, wheat, cabbage, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oil-seed rape.

17. A method for protecting a plant from an insect pest, comprising introducing into said plant or cell thereof at least one expression vector comprising the nucleic acid molecule of claim 1.

18. The nucleic acid molecule of claim 1, wherein said nucleotide sequence is operably linked to a promoter capable of directing expression of said nucleotide sequence in a plant cell.

* * * * *