



US 20220275087A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2022/0275087 A1**
(43) **Pub. Date: Sep. 1, 2022**
BRUENKER et al.(54) **PROTEASE-ACTIVATED T CELL
BISPECIFIC MOLECULES**(30) **Foreign Application Priority Data**


Mar. 22, 2016 (EP) 16161740.2

(71) Applicant: **Hoffmann-La Roche Inc.**, Little Falls,
NJ (US)**Publication Classification**(72) Inventors: **Peter BRUENKER**, Hittnau (CH);
Rebecca CROASDALE-WOOD,
Preston (GB); **Christian KLEIN**,
Bonstetten (CH); **Juergen Michael**
SCHANZER, München (DE);
Kay-Gunnar STUBENRAUCH,
Penzberg (DE); **Pablo UMANA**,
Wollerau (CH); **Martina GEIGER**,
Obfelden (CH); **Eric SULLIVAN**,
Pleasanton, CA (US); **Jigar PATEL**,
Pleasanton, CA (US)(51) **Int. Cl.**
C07K 16/28 (2006.01)
A61P 35/00 (2006.01)
C07K 16/42 (2006.01)
(52) **U.S. Cl.**
CPC **C07K 16/2809** (2013.01); **C07K 16/28**
(2013.01); **A61P 35/00** (2018.01); **C07K**
16/2863 (2013.01); **C07K 16/4208** (2013.01);
C07K 2317/31 (2013.01); **A61K 2039/505**
(2013.01)(73) Assignee: **Hoffmann-La Roche Inc.**, Little Falls,
NJ (US)(57) **ABSTRACT**(21) Appl. No.: **17/541,021**(22) Filed: **Dec. 2, 2021****Related U.S. Application Data**(60) Division of application No. 16/138,417, filed on Sep.
21, 2018, now Pat. No. 11,242,390, which is a con-
tinuation of application No. PCT/EP2017/056556,
filed on Mar. 20, 2017.(60) Provisional application No. 62/433,327, filed on Dec.
13, 2016.

The present invention generally relates to novel protease-activatable T cell activating bispecific molecules and idio-
type-specific polypeptides. The present invention also
relates to polynucleotides encoding such protease-activat-
able T cell activating bispecific molecules and idio-
type-specific polypeptides, and vectors and host cells comprising
such polynucleotides. The invention further relates to meth-
ods for producing the protease-activatable T cell activating
bispecific molecules and idio-
type-specific polypeptides of
the invention, and to methods of using these protease-
activatable T cell activating bispecific molecules and idio-
type-specific polypeptides in the treatment of disease.


Specification includes a Sequence Listing.

FIG. 1A




7859

FIG. 1B




7860

FIG. 1C




7857

FIG. 1D



7858

FIG. 1E



7861

FIG. 2

Clone ID	Isotype	Temp.	k_a (1/Ms)	k_{el} (1/s)	$t_{1/2}$ diss. (min)	K_D (nM)
4.32.63	IgG1, kappa	25°C	1.96E+05	1.05E-04	110	1
		37°C	3.46E+05	6.52E-04	18	2
4.35.77	IgG1, kappa	25°C	1.66E+05	2.09E-04	55	1
		37°C	2.68E+05	6.96E-04	17	3
4.33.88	IgG2b, kappa	25°C	2.71E+05	1.87E-03	6	7
		37°C	4.28E+05	3.24E-03	4	8
4.24.72	IgG1, kappa	25°C	8.72E+04	1.18E-03	10	14
		37°C	1.55E+05	4.45E-03	3	29
4.21.78	IgG1, kappa	25°C	1.31E+05	3.06E-03	4	23
		37°C	4.25E+05	9.28E-03	1	22
4.15.64	IgG2b, kappa	25°C	2.61E+04	2.50E-03	5	96
		37°C	1.44E+05	1.35E-02	1	94

FIG. 3B




FIG. 3A




FIG. 3D




FIG. 3C




FIG. 4A

- 7859: anti ID CH2527 scFv 4.15.64 CD3 Fc
- 7859: anti ID CH2527 scFv 4.15.64 CD3 Fc_treated
- 7857: anti ID CH2527 scFv 4.15.64 non cleavable linker CD3 Fc
- 7861: Monovalent CH2527 IgG
- 7760: bispecific CH1A1A 98/99-CD3




FIG. 4B





FIG. 4C

EC50			
7859scFv 4.15.64 MK062 protease site	4710	16 fold more than 7861	
7861	286	16 fold less than 7859	
7860scFv 4.32.63 MK062 protease site	15538	54 fold more than 7861	
7861	286	54 fold less than 7860	

FIG. 5A



7344






FIG. 5C

7496



7611

FIG. 5E



6298

FIG. 5F





FIG. 5G



Anti CD3 Fab 4.15.64 ,
crossed chains




FIG. 5H

FIG. 6




FIG. 7





FIG. 8



9
6
L




FIG. 10




FIG. 11




FIG. 12A




FIG. 12B




FIG. 12C




FIG. 12D




FIG. 13

Human cell lines	FoIR1 density
Hela	2'240'716
Skov3	91'510
HT29	10'135
MKN45	54
Human Renal Cortical Epithelial Cells (HRCEpiC)	312

FIG. 14A





FIG. 14B






FIG. 15A

FIG. 15B




FIG. 16





FIG. 17



- 7344: anti ID CH2527 scFv 4.15.64 MK062 CD3 16D6 Fc
- 7344: anti ID CH2527 scFv 4.15.64 MK062 CD3 16D6 Fc_treated
- 7676: anti ID CH2527 scFv 4.15.64 non cleavable CD3 16D6 Fc
- 6298 GA916-D-16D5-02 sf W(1)
- 6182 DP47GS TCB sf CHO W(9a)

FIG. 18A





FIG. 18B



- 7496: anti ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc
- 7496: anti ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc treated
- 7611: anti ID CH2527 scFv 4.32.63 non cleavable linker CD3 16D6 Fc
- 6298 GA916-D-16D5-02 sf W(1)
- 6182 DP47GS TCB sf CHO W(9a)

FIG. 19A





FIG. 19B



- 7344: anti ID CH2527 scFv 4.15.64 MK062 CD3 16D6 Fc
- 7344: anti ID CH2527 scFv 4.15.64 MK062 CD3 16D6 Fc_treated
- 7676: anti ID CH2527 scFv 4.15.64 non cleavable CD3 16D6 Fc
- 6298 GA916-D-16D5-02 sf V(1)
- 6182 DP47GS TCB sf CHO V(9a)

FIG. 20A





FIG. 20B



- 7496: anti ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc
- 7496: anti ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc treated
- 7611: anti ID CH2527 scFv 4.32.63 non cleavable linker CD3 16D6 Fc
- 6298 GA916-D-16D5-02 sf VW(1)
- 6182 DP47GS TCB sf CHO VW(9a)




FIG. 21A




FIG. 22A




FIG. 22B




FIG. 23





FIG. 24







FIG. 25

FIG. 26






FIG. 27

FIG. 28




FIG. 29




FIG. 30B




FIG. 30A




FIG. 31




FIG. 32

Molecule	Isotype	Temp.	k_a (L/Ms)	k_d (1/s)	K_D (nM)
GA201	IgG1, kappa	37 °C	1.68E+05	1.20E-04	1
GA201-anti-GA201-scFv MIMP2 cleaved	IgG1, kappa	37 °C	1.03E+05	1.63E-04	2
GA201-anti-GA201-scFv without MIMP2	IgG1, kappa	37 °C	-	-	-




FIG. 33A

8364






FIG. 33B


FIG. 33C



8365

FIG. 33D








FIG. 33F

FIG. 33G




FIG. 33H





FIG. 33I



8505

FIG. 33J




FIG. 34




FIG. 35A




FIG. 35B




FIG. 36A





FIG. 36B











FIG. 37A




8365




8409



8408



7235



8366




FIG. 37B









FIG. 37C













FIG. 37D




8408




8409




8363



7235



8364



6298




FIG. 38A




FIG. 38B




FIG. 39A








FIG. 39B








FIG. 40










FIG. 41A




FIG. 41B





FIG. 41C






FIG. 42A

ID 8955





FIG. 42B


ID 8957

FIG. 42C



ID 8959

ID 8997






FIG. 42F




FIG. 43




FIG. 44




FIG. 45A




FIG. 45B




FIG. 45C




FIG. 45D




FIG. 45E




FIG. 45F



FIG. 45G





FIG. 45H



G. 45I

Protease site with GS linker
Anti CD3 scFv 4.32.63
anti-CD3 VH
CH1
Common Light Chain
VH FoIR1 16D5 VH
CH1 CH1
Common Light Chain
Common Light Chain
Fc(Hole)
P329G
LALA
Fc(Knob)
P329G
LALA

FIG. 45J

GS linker

Anti CD3 scFv 4.32.63

anti-CD3 VH

CH1

Common Light Chain

VH: FolR1 16D5 VH

CH1

Common Light Chain

Fc(Hole) P329G LALA

Fc(Knob) P329G LALA

[illegible]

FIG. 45L




FIG. 45M





FIG. 45N



PROTEASE-ACTIVATED T CELL BISPECIFIC MOLECULES

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a divisional of U.S. patent application Ser. No. 16/138,417, filed on Sep. 21, 2018, which is a continuation of International Patent Application No. PCT/EP2017/056556, filed on Mar. 20, 2017, which claims priority to European Patent Application No. 16161740.2, filed on Mar. 22, 2016, and to U.S. Patent Application No. 62/433,327, filed on Dec. 13, 2016, the disclosures of which are incorporated herein by reference in their entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 14, 2022, is named 51177-025002_Sequence_Listing_2_14_22_ST25 and is 309,161 bytes in size.

FIELD OF THE INVENTION

[0003] The present invention generally relates to novel protease-activatable antigen-binding molecules that comprise an anti-idiotypic-binding moiety which reversibly masks antigen binding of the molecule. Specifically, the invention relates to T cell binding molecules having an anti-idiotypic-binding moiety that masks the CD3-binding moiety until cleaved by a protease. This allows the CD3-binding moiety to be inaccessible or “masked” until it is in proximity to a target tissue, such as a tumor, e.g., tumor-infiltrating T cells. In addition, the present invention relates to polynucleotides encoding such protease-activated T cell binding molecules and idiotype-specific polypeptides, and vectors and host cells comprising such polynucleotides. The invention further relates to methods for producing the protease-activated T cell binding molecules of the invention, and to methods of using the same, e.g., in the treatment of disease.

BACKGROUND

[0004] The selective destruction of an individual target cell or a specific target cell type is often desirable in a variety of clinical settings. For example, it is a primary goal of cancer therapy to specifically destroy tumor cells, while leaving healthy cells and tissues intact and undamaged.

[0005] An attractive way of achieving this is by inducing an immune response against the tumor, to make immune effector cells such as natural killer (NK) cells or cytotoxic T lymphocytes (CTLs) attack and destroy tumor cells. In this regard, bispecific antibodies designed to bind with one “arm” to a surface antigen on target cells, and with the second “arm” to an activating, invariant component of the T cell receptor (TCR) complex, have become of interest in recent years. The simultaneous binding of such an antibody to both of its targets will force a temporary interaction between target cell and T cell, causing activation of any cytotoxic T cell and subsequent lysis of the target cell. Hence, the immune response is re-directed to the target cells and is independent of peptide antigen presentation by the

target cell or the specificity of the T cell as would be relevant for normal MHC-restricted activation of CTLs.

[0006] In this context it is crucial that CTLs are activated only when in close proximity to a target cell, i.e., the immunological synapse is mimicked. Particularly desirable are T cell activating bispecific molecules that do not require lymphocyte preconditioning or co-stimulation in order to elicit efficient lysis of target cells. Several bispecific antibody formats have been developed and their suitability for T cell mediated immunotherapy investigated. These include BiTE (bispecific T cell engager) molecules (Nagorsen and Bäuerle, *Exp Cell Res* 317, 1255-1260 (2011)), diabodies (Holliger et al., *Prot Eng* 9, 299-305 (1996)) and derivatives thereof, such as tandem diabodies (Kipriyanov et al., *J Mol Biol* 293, 41-66 (1999)), DART (dual affinity retargeting) molecules, (Moore et al., *Blood* 117, 4542-51 (2011)), and triomabs (Seimetz et al., *Cancer Treat Rev* 36, 458-467 (2010)).

[0007] The task of generating bispecific molecules suitable for treatment provides several technical challenges related to efficacy, toxicity, applicability and producibility that have to be met. In instances where the bispecific molecule targets an antigen on a target cell, e.g., a cancer cell, that is also expressed in non-target tissue, toxicity can occur. There is thus a need for efficacious T cell activating bispecific molecules that unleash full T cell activation in the presence of target cells but not in the presence of normal cells or tissue.

SUMMARY OF THE INVENTION

[0008] The invention generally relates to T cell activating bispecific molecules that are activated selectively in the presence of a target cell.

[0009] In one aspect, the invention relates to a protease-activatable T cell activating bispecific molecule comprising

[0010] (a) a first antigen binding moiety capable of specific binding to CD3;

[0011] (b) a second antigen binding moiety capable of specific binding to a target cell antigen; and

[0012] (c) a masking moiety covalently attached to the T cell bispecific binding molecule through a protease-cleavable linker, wherein the masking moiety is capable of specific binding to the idiotype of the first or the second antigen binding moiety thereby reversibly concealing the first or second antigen binding moiety.

[0013] In one embodiment, the masking moiety of the protease-activatable T cell activating bispecific molecule is covalently attached to the first antigen binding moiety. In one embodiment the masking moiety is covalently attached to the heavy chain variable region of the first antigen binding moiety.

[0014] In one embodiment the masking moiety is covalently attached to the light chain variable region of the first antigen binding moiety. In one embodiment the masking moiety is an anti-idiotypic scFv.

[0015] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a second masking moiety reversibly concealing the second antigen binding moiety.

[0016] In one embodiment the protease capable of cleaving the protease-cleavable linker is expressed by the target cell. In one embodiment the second antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy

chain are exchanged. In one embodiment the second antigen binding moiety is a crossover Fab molecule wherein the constant regions of the Fab light chain and the Fab heavy chain are exchanged. In one embodiment the first antigen binding moiety is a conventional Fab molecule. In one embodiment the protease-activatable T cell activating bispecific molecule comprises not more than one antigen binding moiety capable of specific binding to CD3. In one embodiment the protease-activatable T cell activating bispecific molecule comprises a third antigen binding moiety which is a Fab molecule capable of specific binding to a target cell antigen. In one particular embodiment the third antigen binding moiety is identical to the second antigen binding moiety. In one particular embodiment the third antigen binding moiety is not identical to the second antigen binding moiety. In one embodiment the second antigen binding moiety is capable of specific binding to FolR1 or HER1. In one embodiment the second antigen binding moiety is capable of specific binding to FolR1, HER1 or Mesothelin. In one embodiment the second antigen binding moiety is capable of specific binding to FolR1, HER1, HER2 or Mesothelin.

[0017] In one embodiment the first and the second antigen binding moiety are fused to each other, optionally via a peptide linker. In one particular embodiment the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In one particular embodiment the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. In one particular embodiment the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety are fused to each other, optionally via a peptide linker.

[0018] In one embodiment the protease-activatable T cell activating bispecific molecule additionally comprises an Fc domain composed of a first and a second subunit capable of stable association. In one embodiment the Fc domain is an IgG, specifically an IgG₁ or IgG₄, Fc domain. In one embodiment the Fc domain is a human Fc domain. In one embodiment the Fc domain exhibits reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain. In one embodiment the Fc domain comprises one or more amino acid substitution that reduces binding to an Fc receptor and/or effector function. In one particular embodiment the one or more amino acid substitution is at one or more position selected from the group of L234, L235, and P329 (Kabat numbering). In one particular embodiment each subunit of the Fc domain comprises three amino acid substitutions that reduce binding to an activating Fc receptor and/or effector function wherein said amino acid substitutions are L234A, L235A and P329G. In one particular embodiment the Fc receptor is an Fcγ receptor. In one embodiment the effector function is antibody-dependent cell-mediated cytotoxicity (ADCC). In one embodiment, the target cell is a human cell.

[0019] In one embodiment the masking moiety comprises a heavy chain variable region comprising at least one of:

[0020] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);

[0021] (b) a CDR H2 amino acid sequence of WIN-TETGEPAYADDFKG (SEQ ID NO:21); and

[0022] (c) a CDR H3 amino acid sequence of PYDYDVLDY (SEQ ID NO:22).

[0023] In one embodiment the masking moiety comprises a light chain variable region comprising at least one of:

[0024] (a) a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO:23);

[0025] (b) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and

[0026] (c) a CDR L3 amino acid sequence of QHS-REFPWT (SEQ ID NO:25).

[0027] In one embodiment the masking moiety comprises a heavy chain variable region comprising:

[0028] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);

[0029] (b) a CDR H2 amino acid sequence of WIN-TETGEPAYADDFKG (SEQ ID NO:21);

[0030] (c) a CDR H3 amino acid sequence of PYDYDVLDY (SEQ ID NO:22); and a light chain variable region comprising:

[0031] (d) a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO:23);

[0032] (e) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and

[0033] (f) a CDR L3 amino acid sequence of QHS-REFPWT (SEQ ID NO:25).

[0034] In one embodiment the masking moiety comprises a heavy chain variable region comprising at least one of:

[0035] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);

[0036] (b) a CDR H2 amino acid sequence of IIWGDG-STNYHSALIS (SEQ ID NO:27); and

[0037] (c) a CDR H3 amino acid sequence of GIT-TVVDDYYAMDY (SEQ ID NO:28).

[0038] In one embodiment the masking moiety comprises a light chain variable region comprising at least one of:

[0039] (a) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);

[0040] (b) a CDR L2 amino acid sequence of AAT-FLAD (SEQ ID NO:30); and

[0041] (c) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).

[0042] In one embodiment the masking moiety comprises a heavy chain variable region comprising:

[0043] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);

[0044] (b) a CDR H2 amino acid sequence of IIWGDG-STNYHSALIS (SEQ ID NO:27);

[0045] (c) a CDR H3 amino acid sequence of GIT-TVVDDYYAMDY (SEQ ID NO:28); and a light chain variable region comprising:

[0046] (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);

[0047] (e) a CDR L2 amino acid sequence of AAT-FLAD (SEQ ID NO:30); and

[0048] (f) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).

[0049] In one embodiment the protease cleavable linker comprises at least one protease recognition sequence. In one embodiment the protease cleavable linker comprises a pro-

tease recognition sequence. In one embodiment the protease recognition sequence is selected from the group consisting of:

(a)	(SEQ ID NO: 36)
RQARVVNG;	
(b)	(SEQ ID NO: 37)
VHMLGFLGPGRSRGSFP;	
(c)	(SEQ ID NO: 38)
RQARVVNGXXXXXVPLSLYSYG;	
and	
(d)	(SEQ ID NO: 39)
RQARVVNGVPLSLYSYG	
(e)	(SEQ ID NO: 40)
PLGLWSQ, wherein X is any amino acid.	

[0050] In one embodiment the protease cleavable linker comprises a protease recognition sequence. In one embodiment the protease recognition sequence is selected from the group consisting of:

(a)	(SEQ ID NO: 36)
RQARVVNG;	
(b)	(SEQ ID NO: 37)
VHMLGFLGPGRSRGSFP;	
(c)	(SEQ ID NO: 38)
RQARVVNGXXXXXVPLSLYSYG;	
(d)	(SEQ ID NO: 39)
RQARVVNGVPLSLYSYG;	
(e)	(SEQ ID NO: 40)
PLGLWSQ;	
(f)	(SEQ ID NO: 97)
VHMLGFLGPGRQARVVNG;	
(g)	(SEQ ID NO: 98)
FVGGTG;	
(h)	(SEQ ID NO: 99)
KKAAPVNG;	
(i)	(SEQ ID NO: 100)
PMACKVNG;	
(j)	(SEQ ID NO: 101)
QARAKVNG;	
(k)	(SEQ ID NO: 102)
VHMLGFLGP;	

-continued

(l)	(SEQ ID NO: 103)
QARAK;	
(m)	(SEQ ID NO: 104)
VHMLGFLGPMPMAKK;	
(n)	(SEQ ID NO: 105)
KKAAP;	
and	
(o)	(SEQ ID NO: 106)
PMACK, wherein X is any amino acid.	

[0051] In one embodiment the protease capable of cleaving the protease-cleavable linker is selected from the group consisting of metalloproteinase, e.g., matrix metalloproteinase (MMP) 1-28 and A Disintegrin And Metalloproteinase (ADAM) 2, 7-12, 15, 17-23, 28-30 and 33, serine protease, e.g., urokinase-type plasminogen activator and Matriptase, cysteine protease, aspartic protease, and cathepsin protease. In one specific embodiment the protease is MMP9 or MMP2. In a further specific embodiment, the protease is Matriptase. In one embodiment the protease cleavable linker comprises the protease recognition sequence RQARVVNG (SEQ ID NO:36).

[0052] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the first antigen binding moiety comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0053] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the first antigen binding moiety comprises the heavy chain complementarity determining region (CDRs) of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and the light chain CDRs of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0054] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the first antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0055] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the first antigen binding moiety comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.

[0056] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino

acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0057] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 151, SEQ ID NO: 152 and SEQ ID NO: 153 and at least one light chain CDR comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 154, SEQ ID NO: 155 and SEQ ID NO: 156.

[0058] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0059] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 151, SEQ ID NO: 152 and SEQ ID NO: 153 and at least one light chain CDR selected from the group of SEQ ID NO: 154, SEQ ID NO: 155 and SEQ ID NO: 156.

[0060] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0061] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 158.

[0062] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.

[0063] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 158.

[0064] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to Mesothelin and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 107, SEQ ID NO: 108 and SEQ ID NO: 109 and at least one light chain CDR comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 110, SEQ ID NO: 111 and SEQ ID NO: 112.

[0065] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to Mesothelin and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 107, SEQ ID NO: 108 and SEQ ID NO: 109 and at least one light chain CDR selected from the group of SEQ ID NO: 110, SEQ ID NO: 111 and SEQ ID NO: 112.

[0066] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 114.

[0067] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 114.

[0068] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to HER1 and comprises at least one heavy chain complementarity determining region (CDR) of any one of the antibodies disclosed in WO/2006/082515 incorporated herein by reference in its entirety.

[0069] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to HER1 and comprises at least one heavy chain complementarity determining region (CDR) comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 61.

[0070] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety is capable of specific binding to HER1 and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 61.

[0071] In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 32 and a light chain comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 33. In one embodiment of the protease-activatable T cell activating bispecific molecule described herein the second antigen binding moiety comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 32 and a light chain comprising the amino acid sequence of SEQ ID NO: 33.

[0072] In one embodiment, the first antigen binding moiety is capable of specific binding to CD3, and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, and the second and third antigen binding moieties are capable of specific binding to HER2, wherein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 160 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 161, wherein the third antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 159 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 161.

[0073] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0074] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:2;

[0075] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and

[0076] (c) a light chain comprising the amino acid sequence of SEQ ID NO:1.

[0077] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0078] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:4;

[0079] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and

[0080] (c) a light chain comprising the amino acid sequence of SEQ ID NO:1.

[0081] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0082] (a) at least one heavy chain comprising the amino acid sequence of SEQ ID NO:32;

[0083] (b) at least one light chain comprising the amino acid sequence of SEQ ID NO:34.

[0084] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0085] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:72;

[0086] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and

[0087] (c) a light chain comprising an amino acid sequence of SEQ ID NO:1.

[0088] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0089] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:85;

[0090] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and

[0091] (c) a light chain comprising an amino acid sequence of SEQ ID NO:1.

[0092] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0093] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:73;

[0094] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3;

[0095] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:1; and

[0096] (d) a second light chain comprising an amino acid sequence of SEQ ID NO: 74.

[0097] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0098] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:77;

[0099] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:82;

[0100] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:78; and

[0101] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:81.

[0102] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

[0103] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:76;

[0104] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:77;

[0105] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:78; and

[0106] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:79.

[0107] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises

- [0108] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:132;
- [0109] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:136;
- [0110] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:81; and
- [0111] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:133.
- [0112] In one particular embodiment the protease-activatable T cell activating bispecific molecule described herein comprises
- [0113] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:137;
- [0114] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:139;
- [0115] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:81; and
- [0116] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:138.
- [0117] In one aspect, the invention relates to an idiotype-specific polypeptide for reversibly concealing an anti-CD3 antigen binding site of a molecule. In one embodiment the idiotype-specific polypeptide is an anti-idiotypic scFv. In one embodiment the idiotype-specific polypeptide is covalently attached to the molecule through a linker. In one embodiment the linker is a peptide linker. In one embodiment the linker is a protease-cleavable linker. In one embodiment the peptide linker comprises at least one protease recognition site. In one embodiment the protease is selected from the group consisting of metalloproteinase, e.g., matrix metalloproteinase (MMP) 1-28 and A Disintegrin And Metalloproteinase (ADAM) 2, 7-12, 15, 17-23, 28-30 and 33, serine protease, e.g., urokinase-type plasminogen activator and Matrilysin, cysteine protease, aspartic protease, and cathepsin protease. In one specific embodiment the protease is MMP9 or MMP2. In a further specific embodiment, the protease is Matrilysin. In one embodiment the idiotype-specific polypeptide is covalently attached to the molecule through more than one linker. In one embodiment the idiotype-specific polypeptide is covalently attached to the molecule through two linkers.
- [0118] In one embodiment the molecule which comprises the anti-CD3 antigen binding site is a T-cell activating bispecific molecule. In one particular embodiment the idiotype-specific polypeptide comprises a heavy chain variable region comprising at least one of:
- [0119] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);
- [0120] (b) a CDR H2 amino acid sequence of WIN-TETGEPAYADDFKG (SEQ ID NO:21); and
- [0121] (c) a CDR H3 amino acid sequence of PYDYDVLDY (SEQ ID NO:22).
- [0122] In one particular embodiment the idiotype-specific polypeptide comprises a light chain variable region comprising at least one of:
- [0123] (a) a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO:23);
- [0124] (b) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and
- [0125] (c) a CDR L3 amino acid sequence of QHS-REFPWT (SEQ ID NO:25).
- [0126] In one particular embodiment the idiotype-specific polypeptide comprises:
- [0127] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);
- [0128] (b) a CDR H2 amino acid sequence of WIN-TETGEPAYADDFKG (SEQ ID NO:21);
- [0129] (c) a CDR H3 amino acid sequence of PYDYDVLDY (SEQ ID NO:22); and a light chain variable region comprising:
- [0130] (d) a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO:23);
- [0131] (e) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and
- [0132] (f) a CDR L3 amino acid sequence of QHS-REFPWT (SEQ ID NO:25).
- [0133] In one particular embodiment the idiotype-specific polypeptide comprises a heavy chain variable region comprising at least one of:
- [0134] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- [0135] (b) a CDR H2 amino acid sequence of IIWGDG-STNYHSALIS (SEQ ID NO:27); and
- [0136] (c) a CDR H3 amino acid sequence of GIT-TVVDDYYAMDY (SEQ ID NO:28).
- [0137] In one particular embodiment the idiotype-specific polypeptide comprises a light chain variable region comprising at least one of:
- [0138] (a) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- [0139] (b) a CDR L2 amino acid sequence of AAT-FLAD (SEQ ID NO:30); and
- [0140] (c) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).
- [0141] In one particular embodiment the idiotype-specific polypeptide comprises a heavy chain variable region comprising:
- [0142] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- [0143] (b) a CDR H2 amino acid sequence of IIWGDG-STNYHSALIS (SEQ ID NO:27);
- [0144] (c) a CDR H3 amino acid sequence of GIT-TVVDDYYAMDY (SEQ ID NO:28); and a light chain variable region comprising:
- [0145] (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- [0146] (e) a CDR L2 amino acid sequence of AAT-FLAD (SEQ ID NO:30); and
- [0147] (f) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).
- [0148] According to another aspect of the invention, an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof is provided. The invention also encompasses polypeptides encoded by the polynucleotides of the invention. The invention further provides an expression vector comprising the isolated polynucleotide of the invention, and a host cell comprising the isolated polynucleotide or the expression vector of the invention. In some embodiments the host cell is a eukaryotic cell, particularly a mammalian cell. In another aspect is provided a method of producing the protease-activated T cell molecule of the invention, comprising the steps of a) culturing the host cell of the invention under conditions suitable for the expression

of the protease-activated T cell molecule and b) recovering the protease-activated T cell molecule. The invention also encompasses a protease-activated T cell molecule produced by the method of the invention.

[0149] In another aspect is provided a method of producing the idiotypic-specific polypeptide of the invention, comprising the steps of a) culturing the host cell of the invention under conditions suitable for the expression of the protease-activated T cell molecule and b) recovering the idiotypic-specific polypeptide. The invention also encompasses a idiotypic-specific polypeptide produced by the method of the invention.

[0150] The invention further provides a pharmaceutical composition comprising the protease-activatable T cell activating bispecific molecule of the invention and a pharmaceutically acceptable carrier.

[0151] Also encompassed by the invention are methods of using the protease-activated T cell molecule and pharmaceutical composition of the invention. In one aspect the invention provides a protease-activated T cell molecule or a pharmaceutical composition of the invention for use as a medicament. In one aspect is provided a protease-activated T cell molecule or a pharmaceutical composition according to the invention for use in the treatment of a disease in an individual in need thereof. In a specific embodiment the disease is cancer.

[0152] Also provided is the use of a protease-activated T cell molecule of the invention for the manufacture of a medicament for the treatment of a disease in an individual in need thereof; as well as a method of treating a disease in an individual, comprising administering to said individual a therapeutically effective amount of a composition comprising the protease-activated T cell molecule according to the invention in a pharmaceutically acceptable form. In a specific embodiment the disease is cancer. In any of the above embodiments the individual preferably is a mammal, particularly a human.

[0153] The invention also provides a method for inducing lysis of a target cell, particularly a tumor cell, comprising contacting a target cell with a protease-activated T cell molecule of the invention in the presence of a T cell, particularly a cytotoxic T cell.

[0154] In another aspect the invention also provides a composition comprising a protease-activatable T cell activating bispecific molecule described herein and a pharmaceutically acceptable carrier.

[0155] In another aspect the invention also provides a composition comprising an idiotypic-specific polypeptide as described herein and a pharmaceutically acceptable carrier.

[0156] In another aspect the invention also provides a protease-activatable T cell activating bispecific molecule or an idiotypic-specific polypeptide as described herein, or the composition described herein, for use as a medicament. In one embodiment the medicament is for treating or delaying progression of cancer, treating or delaying progression of an immune related disease, and/or enhancing or stimulating an immune response or function in an individual.

[0157] In another aspect the invention also provides a protease-activatable T cell activating bispecific molecule or idiotypic-specific polypeptide as described herein for use in the treatment of a disease in an individual in need thereof. In one embodiment, the disease is a proliferative disorder, particularly cancer.

[0158] In another aspect the invention also provides a method of treating a disease in an individual, comprising administering to said individual a therapeutically effective amount of a composition comprising the protease-activatable T cell activating bispecific molecule or composition as described herein.

[0159] In another aspect the invention also provides a method for inducing lysis of a target cell, comprising contacting a target cell with the protease-activatable T cell activating bispecific molecule or composition as described herein in the presence of a T cell. In one embodiment the method for inducing lysis of a target cell is an in vitro method. In one embodiment the target cell is a cancer cell. In one embodiment the target cell expresses a protease capable of activating the protease-activatable T cell activating bispecific molecule.

[0160] In another aspect the invention also provides an anti-idiotypic CD3 antibody or antigen-binding fragment thereof specific for an idiotypic of an anti-CD3 antigen-binding molecule, wherein the anti-idiotypic CD3 antibody or fragment thereof when bound to the anti-CD3 antigen-binding molecule specifically blocks binding of the anti-CD3 antigen-binding molecule to CD3.

[0161] In one embodiment, the anti-idiotypic CD3 antibody or antigen-binding fragment thereof is reversibly associated with the anti-CD3 antigen-binding molecule through a peptide linker comprising a protease recognition site. In one embodiment, the CD3 is a mouse, monkey or human CD3.

[0162] In another aspect the invention provides a method of reducing in vivo toxicity of a T cell activating bispecific molecule comprising attaching an idiotypic-specific polypeptide as described herein to the T cell activating bispecific molecule with a protease-cleavable linker to form a protease-activatable T cell activating bispecific molecule, wherein the protease-activatable T cell activating bispecific molecule has reduced in vivo toxicity compared to the T cell activating bispecific molecule.

SHORT DESCRIPTION OF THE FIGURES

[0163] FIGS. 1A-1E depict schematics of different CD3 binders with masking moieties. FIG. 1A: 7859 anti-ID CH2527 scFv 4.15.64 MK062 Matritase site CD3 Fc. FIG. 1B: 7860 anti-ID CH2527 scFv 4.32.63 MK062 Matritase site CD3 Fc. FIG. 1C: 7857 anti-ID CH2527 scFv 4.15.64 non-cleavable linker CD3 Fc. FIG. 1D: ID 7858 anti-ID CH2527 scFv 4.32.63 non-cleavable linker CD3 Fc. FIG. 1E: 7861 monovalent CD3 Fc.

[0164] FIG. 2 shows a table summarizing the affinities of the anti-idiotypic masking moieties to the CD3 binder (CH2527).

[0165] FIGS. 3A-3D shows Capillary Electrophoresis-SDS analysis of the molecules depicted in FIGS. 1A and 1B. FIGS. 3A and 3B: Capillary Electrophoresis-SDS analysis of the molecule depicted in FIG. 1A under non reducing (FIG. 3A) and reducing conditions (FIG. 3B). Comparison of the untreated (I) and treated molecule (III) shows complete cleavage of the anti-ID scFv after rhMatritase/ST14 treatment for 48 h at 37° C. One sample (II) was untreated but incubated at 37° C. for 48 h. FIGS. 3C and 3D: Capillary Electrophoresis-SDS analysis of the molecule depicted in FIG. 1B under non-reducing (FIG. 3C) and reducing conditions (FIG. 3D). Comparison of the untreated (I) and treated molecule (III) shows complete cleavage of the anti-

ID scFv after rhMatriptase/ST14 treatment for 48 h at 37° C. One sample (II) was untreated but incubated at 37° C. for 48 h.

[0166] FIGS. 4A-4C show the effect of anti-idiotypic masking of CD3 binding. FIGS. 4A and 4B depict results of Jurkat NFAT reporter assays to show the masking effect of anti-idiotypic CD3 scFv 4.15.64 (FIG. 4A) or anti-idiotypic CD3 scFv 4.32.63 (FIG. 4B). Monovalent CD3 IgGs were crosslinked via an anti-human Fc antibody (coated on assay plate) before Jurkat NFAT (acute lymphatic leukemia reporter cell line with a NFAT promoter, expressing human CD3 ϵ) were added. The Jurkat-NFAT reporter cell line (Promega) is a human acute lymphatic leukemia reporter cell line with a NFAT promoter, expressing human CD3 ϵ . If CD3 binder binds CD3 ϵ Luciferase is expressed and this can be measured in Luminescence after addition of One-Glo substrate (Promega). FIG. 4C shows a comparison of EC50 values of CD3 ϵ binding for masked and unmasked monovalent CD3 binder.

[0167] FIGS. 5A-5H depict schematics of different T cell bispecific molecules with masking moieties. FIG. 5A: 7344 anti-ID CH2527 scFv 4.15.64 MK062 Matriptase site CD3 16D5 Fc. FIG. 5B: 7676 anti-ID CH2527 scFv 4.15.64 non-cleavable linker CD3 16D5 Fc. FIG. 5C: 7496 anti-ID CH2527 scFv 4.32.63 MK062 Matriptase site CD3 16D5 Fc. FIG. 5D: 7611 anti-ID CH2527 scFv 4.32.63 non-cleavable linker CD3 16D5 Fc. FIG. 5E: 6298 GA916-D-16D5-02 sf W(1). FolR1 16D5 classic 2+1 TCB with common light chain. FIG. 5F: 6100 GA916-D-16D5 sf W(3a). FolR1 16D5 inverted 2+1 TCB with common light chain. FIG. 5G: ID 6182 DP47GS TCB sf CHO W(9a). DP47 inverted 2+1 TCB. FIG. 5H: 7494 anti-ID CH2527 Fab 4.15.64 MK062 Matriptase site CD3 16D5 Fc.

[0168] FIG. 6 shows a first plasmid ratios used for transfection by size exclusion chromatography (1(hole): 1 (knob): 3 (CLC)).

[0169] FIG. 7 shows a second plasmid ratios used for transfection by size exclusion chromatography. (1(hole): 2 (knob): 3 (CLC)).

[0170] FIG. 8 shows CE-SDS analysis of the TCB molecule depicted in FIG. 5A (ID 7344) (final purified preparation): Lane A=non-reduced, lane B=reduced, lane C=Protein standard.

[0171] FIG. 9 shows CE-SDS analysis of the TCB molecule depicted in FIG. 5B (ID 7676) (final purified preparation): Lane A=non-reduced, lane B=reduced, lane C=Protein standard.

[0172] FIG. 10 shows CE-SDS analysis of the TCB molecule depicted in FIG. 5C (ID 7496) (final purified preparation): Lane A=non-reduced, lane B=reduced, lane C=Protein standard.

[0173] FIG. 11 shows CE-SDS analysis of the TCB molecule depicted in FIG. 5D (ID 7611) (final purified preparation): Lane A=non-reduced, lane B=reduced, lane C=Protein standard.

[0174] FIG. 12A-12D show shows Capillary Electrophoresis-SDS analysis of the molecules depicted in FIGS. 5A and 5C. FIGS. 12A and 12B show Capillary Electrophoresis of the molecules depicted in FIG. 5A (ID 7344) anti-ID CH2527 scFv 4.15.64 MK062 CD3 16D6 Fc under non reducing (FIG. 12A) and reducing conditions (FIG. 12B). Comparison of the untreated (I) and treated molecule (III) shows complete cleavage of the anti-ID scFv after rhMatriptase/ST14 treatment for 48 h at 37° C. One sample

(II) was untreated but incubated at 37° C. for 48 h. Pre-stained protein Marker (IV) Mark 12 (Invitrogen) was used for estimation of correct molecule weight. FIGS. 12C and 12D shows Capillary Electrophoresis of the molecule depicted in FIG. 5C (ID 7496) anti-ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc under non reducing (FIG. 12C) and reducing conditions (FIG. 12D). Comparison of the untreated (I) and treated molecule (III) shows complete cleavage of the anti-ID scFv after rhMatriptase/ST14 treatment for 48 h at 37° C. One sample (II) was untreated but incubated at 37° C. for 48 h. Pre-stained protein Marker (IV) Mark 12 (Invitrogen) was used for estimation of correct molecule weight.

[0175] FIG. 13 shows FolR1 expression level quantification done by Qifikit (Dako). Antibody for FolR1: #LS-C125620-100 (LifeSpan BioSciences Inc); used at 20 μ g/ml; mouse IgG1 isotype: #554121 (BD).

[0176] FIGS. 14A and 14B show T cell activation by protease activated TCBs. FIG. 14A shows killing of Skov3 cells induced by protease-activated TCB molecules at a concentration of 10 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, molecules pre-treated with purified rhMatriptase/ST14) and human PBMCs after 48 h of incubation (E:T=7:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. FIG. 14B shows T cell activation of human PBMCs induced by protease activated TCB binding of 10 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, treated molecules) on Skov3 cells after 48 h of incubation (E:T=7:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0177] FIGS. 15A and 15B show T cell activation by protease activated TCBs. FIG. 15A shows killing of Mkn-45 cells induced by protease activated TCB molecules at a concentration of 100 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, treated molecules) and human PBMCs after 48 h of incubation (E:T=7:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. FIG. 15B shows T cell activation of human PBMCs induced by protease activated TCB binding of 100 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, treated molecules) on Mkn-45 cells after 48 h of incubation (E:T=7:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0178] FIG. 16 shows killing of HT29 cells induced by protease activated TCB molecules at a concentration of 10 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, treated molecules) and human PBMCs after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. Bars from left to right are 7344: anti-ID CH2527 scFv 4.15.64 MK062 CD3 16D6Fc; 7344: anti-ID CH2527 scFv 4.15.64 MK062 CD3 16D6Fc_treated; 7676: anti-ID CH2527 scFv 4.15.64 non-cleavable CD3 16D6Fc; 7496 anti-ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc; 7496 anti-ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc_treated; 7611: ID anti CH2527 scFv 4.32.63 non-cleavable linker CD3 16D6 Fc; 6298 GA916-D-16D5-02 sf W(1); 6182 DP47GS TCB sf CHO W(9a).

[0179] FIG. 17 shows killing of Skov3 cells induced by protease activated TCB molecules at a concentration of 10 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, treated molecules) and human PBMCs (from a different donor than PBMCs used for FIG. 14A) after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. Bars from left to right are 7344: anti-ID CH2527 scFv 4.15.64 MK062 CD3 16D6Fc; 7344: anti-ID CH2527 scFv 4.15.64 MK062 CD3 16D6Fc_treated; 7676: anti-ID CH2527 scFv 4.15.64 non-cleavable CD3 16D6Fc; 7496 anti-ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc; 7496 anti-ID CH2527 scFv 4.32.63 MK062 CD3 16D6 Fc_treated; 7611: ID anti CH2527 scFv 4.32.63 non-cleavable linker CD3 16D6 Fc; 6298 GA916-D-16D5-02 sf W(1); 6182 DP47GS TCB sf CHO W(9a).

[0180] FIGS. 18A and 18B show T cell activation by protease activated TCBs. FIG. 18A shows dose-dependent killing of HeLa cells induced by protease activated TCB molecules (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from buffy coat) after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. FIG. 18B shows dose-dependent T cell activation of human PBMCs induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) on HeLa cells after 48 h of incubation (E:T=10:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0181] FIGS. 19A and 19B show T cell activation by protease activated TCBs. FIG. 19A shows dose-dependent killing of HeLa cells induced by protease activated TCB molecules (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from buffy coat) after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. FIG. 19B shows dose-dependent T cell activation of human PBMCs induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) on HeLa cells after 48 h of incubation (E:T=10:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0182] FIGS. 20A and 20B show T cell activation by protease activated TCBs. FIG. 20A shows dose-dependent killing of Skov3 cells induced by protease activated TCB molecules (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from buffy coat) after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. FIG. 20B shows dose-dependent T cell activation of human PBMCs induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) on Skov3 cells after 48 h of incubation (E:T=10:1, effectors are

human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0183] FIGS. 21A and 21B show T cell activation by protease activated TCBs. FIG. 21A shows dose-dependent killing of Skov3 cells induced by protease activated TCB molecules (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from buffy coat) after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C. FIG. 21B shows dose-dependent T cell activation of human PBMCs induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) on Skov3 cells after 48 h of incubation (E:T=10:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0184] FIGS. 22A and 22B show T cell activation by protease activated TCBs. FIG. 22A shows dose-dependent T cell activation of human PBMCs (different donor than in experiments described above) induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) on HT29 cells after 48 h of incubation (E:T=10:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated. FIG. 22B shows dose-dependent T cell activation of human PBMCs (different donor than in FIG. 16) induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) on HT29 cells after 48 h of incubation (E:T=10:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0185] FIG. 23 shows dose-dependent T cell activation of human PBMCs (different donor than in experiments described above) induced by protease activated TCB binding (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) on HRCepiC cells after 48 h of incubation (E:T=10:1, effectors are human PBMCs). T cell activation markers CD25 (left panels) and CD69 (right panels). CD4+ and CD8+ T cells as indicated.

[0186] FIG. 24 shows killing of Ovar3 cells induced by protease activated TCB molecules at a concentration of 50 nM (TCBs with different anti-idiotypic CD3 masks, cleavable and non-cleavable linker, treated molecules) and human PBMCs after 48 h of incubation (E:T=10:1, effectors are human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 10 min at 37° C. (not fully cleaved).

[0187] FIG. 25 shows killing of Skov3 cells induced by 10 nM of protease activated TCB molecules (TCB with anti-idiotypic CD3 4.15.64 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from buffy coat) after 48 h of incubation (E:T=10:1, effectors are three different Donors for human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C.

[0188] FIG. 26 shows killing of Skov3 cells induced by 10 nM of protease activated TCB molecules (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from

buffy coat) after 48 h of incubation (E:T=10:1, effectors are three different Donors for human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C.

[0189] FIG. 27 shows killing of HeLa cells induced by 100 pM of protease activated TCB molecules (TCB with anti-idiotypic CD3 4.32.63 mask, cleavable and non-cleavable linker, treated molecule) and human PBMCs (isolated from buffy coat) after 48 h of incubation (E:T=10:1, effectors are three different Donors for human PBMCs). Pre-treatment with rhMatriptase/ST14 (R&D Systems) was done for 24 h at 37° C.

[0190] FIG. 28 depicts a schematic of anti-ID GA201 scFv Matrix Metalloprotease site GA201 Fc (GA201-anti-GA201-scFv).

[0191] FIG. 29 depicts a schematics of the anti HER1 antibody GA201.

[0192] FIGS. 30A and 30B show capillary Electrophoresis-SDS analysis of the molecule depicted in FIG. 28 under non-reducing (FIG. 30A) and reducing conditions (FIG. 30B). The molecule depicted in FIG. 28 was purified to homogeneity by Protein A and Size Exclusion chromatography and subjected to Capillary electrophoresis-SDS analysis.

[0193] FIG. 31 shows FACS analysis of GA201-anti-GA201-scFv and GA201 binding to HER1 expressed on H322M cells to confirm masking effect of anti-idiotypic GA201 scFv. GA201-anti-GA201-scFv was incubated overnight with the Matrix Metalloprotease MMP-2 and binding to H322M cells was compared to uncleaved GA201-anti-GA201-scFv, GA201 and an isotype IgG1 control antibody. Binding to HER1 on H322M cells was detected with a F(ab')₂-goat anti-human IgG Fc secondary antibody FITC conjugate and analyzed by FACS using the BD FACS Canto II. The median fluorescence intensity (MFI) was used for analysis.

[0194] FIG. 32 shows surface plasmon resonance analysis of HER1 binding of masked and unmasked GA201, before and after MMP2 cleavage.

[0195] FIGS. 33A-33J depict schematics of different T cell bispecific molecules with masking moieties. FIG. 33A: ID 8364. 16D5 TCB, classic format, anti ID CH2527 scFv 4.32.63 MMP9-MK062 Matriptase site N-terminally fused to CD3. FIG. 33B: ID 8363. 16D5 TCB, classic format, anti ID CH2527 scFv 4.32.63 Cathepsin S/B site N-terminally fused to CD3. FIG. 33C: ID 8365. 16D5 TCB, inverted format, anti ID CH2527 scFv 4.32.63 MK062 Matriptase site N-terminally fused to CD3. FIG. 33D: ID 8366. 16D5 TCB, inverted format, anti ID CH2527 scFv 4.32.63 non-cleavable linker N-terminally fused to common light chain. FIG. 33E: ID 8672. aMesothelin RG7787 charged residues TCB, classic format, anti ID CH2527 scFv 4.32.63 MMP9-MK062 Matriptase site N-terminally fused to CD3xFab. FIG. 33F: ID 8673. aMesothelin RG7787 charged residues TCB, classic format, anti ID CH2527 scFv 4.32.63 non-cleavable linker N-terminally fused to CD3xFab. FIG. 33G: ID 8674. aMesothelin RG7787 charged residues TCB, inverted format, anti ID CH2527 scFv 4.32.63 MMP9-MK062 Matriptase site N-terminally fused to CD3 XFab. FIG. 33H: 8675. aMesothelin RG7787 charged residues TCB, inverted format, anti ID CH2527 scFv 4.32.63 non-cleavable linker N-terminally fused to CD3 XFab. FIG. 33I: ID 8505. aMesothelin RG7787 charged residues

CD3 XFab TCB, inverted format. FIG. 33J: ID 8676. CD3 XFab aMesothelin RG7787 charged residues TCB, classic format.

[0196] FIG. 34 depicts CE-SDS analysis of the TCB ID 8365 and TCB ID 8366 (final purified preparation): Lane A=Protein standard, lane B=protein stored at 4° C., lane C=protein pretreated with rhMatriptase/ST14 (R&D Systems), lane D=protein incubated for 72 h at 37° C. and lane E=molecule 3.

[0197] FIGS. 35A and 35B. depicts CE-SDS analysis of the TCB depicted in FIG. 33A (ID 8364) and the TCB depicted in FIG. 33B (ID 8363). FIG. 35A: CE-SDS analysis of the TCB 8364 (final purified preparation): Lane A=Protein standard, lane B=protein stored at 4° C., lane C=protein pretreated with rhMatriptase/ST14 (R&D Systems), lane D=protein incubated for 72 h at 37° C. and lane E=non-cleavable linker construct. FIG. 35B: CE-SDS analysis of the TCB 8363 (final purified preparation): Lane A=Protein standard, lane B=protein stored at 4° C., lane C=protein pretreated with rhCathepsin B (R&D Systems), lane D=protein pretreated with rhCathepsin S (R&D Systems), lane E=protein incubated for 72 h at 37° C. and lane F=non-cleavable linker construct.

[0198] FIGS. 36A and 36B. depicts Jurkat NFAT activation assay using HeLa and Skov-3 cells as target cells. Each point represents the mean value of triplicates. Standard deviation is indicated by error bars. Jurkat-NFAT reporter cell line (Promega) is a human acute lymphatic leukemia reporter cell line with a NFAT promoter, expressing human CD3c. If the CD3 binder of the TCB binds the tumor target and the CD3 (cross-linkage is necessary) binds CD3ε the Luciferase expression can be measured in Luminescence after addition of One-Glo substrate (Promega). The FolR1 TCB (black triangles pointing down) and the pretreated protease activated TCB (8364, grey filled squares) with N-terminally fused anti ID CD3 4.32.63 scFv and MMP9-Matriptase MK062 site were compared. The molecule was treated with rhMatriptase/ST14 (R&D Systems) for about 20 h at 37° C. The masked TCB (containing a GS non-cleavable linker, grey triangles pointing up) and the non-targeted TCB control (empty triangle pointing down) are shown as well. The dotted line shows the Luminescence of target cells and effector cells without any TCB.

[0199] FIG. 36A shows a Jurkat NFAT activation assay using HeLa cells as target cells.

[0200] FIG. 36B shows a Jurkat NFAT activation assay using Skov-3 cells as target cells.

[0201] FIGS. 37A-37D depict tumor cell cytotoxicity mediated by FolR1 TCBs and human PBMCs (Effector: Target=10:1). Each point represents the mean value of triplicates. Standard deviation is indicated by error bars. FIG. 37A: HeLa target cell cytotoxicity. Comparison of two different formats of the Protease activated TCBs both containing an anti idiotypic CD3 scFv linked with a MK062 Matriptase linker. FIG. 37B: Skov-3 target cell cytotoxicity. Comparison of two different formats of the Protease activated TCBs both containing an anti idiotypic CD3 scFv linked with a MK062 Matriptase linker. FIG. 37C: HeLa target cell cytotoxicity. Comparison of classic Protease activated TCB containing an anti idiotypic CD3 scFv and GS linkers with different protease sites. Protease activated TCB containing the MMP9-Matriptase MK062 linker (8364, grey squares), FolR1 TCB (light grey triangles pointing down), protease activated TCB containing only

Matriptase MK062 (light grey rhomb)/Cathepsin site (grey circles) or non-cleavable linker (black triangles pointing down). FIG. 37D: Skov-3 target cell cytotoxicity. Comparison of classic Protease activated TCB containing an anti idiotypic CD3 scFv and GS linkers with different protease sites. Protease activated TCB containing the MMP9-Matriptase MK062 linker (8364, grey squares), FolR1 TCB (light grey triangles pointing down), protease activated TCB containing only Matriptase MK062 (light grey rhomb)/Cathepsin site (grey circles) or non-cleavable linker (black triangles pointing down).

[0202] FIGS. 38A and 38B depict quantification of CD69 of CD8 positive cells after co-incubation of primary human renal epithelial cortical cells (FIG. 38A) or human bronchial epithelial cells (FIG. 38B) with 200 nM of the different TCBs and three different donors of human PBMCs. T cells were stained after 48 h of incubation. (E:T=10:1, effectors are human PBMCs). Median fluorescence intensity of T cell activation marker CD69 for CD8⁺ T cells is shown. Each point represents the mean value of triplicates of three different human PBMC donors. Standard deviation is indicated in error bars. Unpaired t test was used for statistical analysis.

[0203] FIGS. 39A and 39B depict tumor cell cytotoxicity mediated by MSLN TCBs and human PBMCs (Effector: Target=10:1). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB. Each point represents the mean value of triplicates. Standard deviation is indicated by error bars. FIG. 39A: NCI H596 target cell cytotoxicity. Protease activated MSLN TCB containing an anti idiotypic CD3 scFv linked with a MMP9-MK062 Matriptase linker. The protease activated TCB (8672, light grey circles), the MSLN TCB (dark grey triangles pointing down) and the protease activated TCB containing a non-cleavable linker (8673, grey triangles pointing up) are compared. FIG. 39B: AsPC-1 target cell cytotoxicity. Protease activated MSLN TCB containing an anti idiotypic CD3 scFv linked with a MMP9-MK062 Matriptase linker. The protease activated TCB (8672, light grey circles), the MSLN TCB (dark grey triangles pointing down) and the protease activated TCB containing a non-cleavable linker (8673, grey triangles pointing up) are compared.

[0204] FIG. 40 depicts a Jurkat-NFAT activation assay with primary tumor samples and Protease activated FolR1 TCBs. Jurkat NFAT reporter cells are activated after co-incubation with FolR1 TCB (6298) and Protease activated FolR1 TCB containing MMP9-Matriptase cleavage site (8364). Protease activated FolR1 TCBs (8363, 8408) and control TCBs (8409, 7235) do not induce Luciferase expression. The dotted line indicates the baseline Luminescence for Jurkat NFAT cells co-incubated with tumor.

[0205] FIGS. 41A-41C: Capillary electrophoresis of protease activated TCBs after incubation in human serum. Molecules were incubated for 0 or 14 days in human IgG depleted serum at 37° C. in a humidified incubator (5% CO₂). All molecules were purified by affinity chromatography (ProteinA) and then analyzed by Capillary electrophoresis. FIG. 41A: CE-SDS analysis of serum, FolR1 TCB (6298) in serum at day 0 and day 14. FIG. 41B: CE-SDS analysis of serum, Protease activated FolR1 TCB with MMP9-Matriptase linker (8364) in serum at day 0 and day

14. FIG. 41C: CE-SDS analysis of serum, Protease activated FolR1 TCB with Matriptase linker (8408) in serum at day 0 and day 14 and the precleaved molecule in serum.

[0206] FIGS. 42A-42F depict schematics of different T cell bispecific molecules with masking moieties. FIG. 42A: ID 8955. Herceptarg TCB, classic format, anti ID CH2527 scFv 4.32.63 MK062 MMP9 linker N-terminally fused to VH. FIG. 42B: ID 8957. Herceptarg TCB, classic format, anti ID CH2527 scFv 4.32.63 non cleavable linker N-terminally fused to VH. FIG. 42C: ID 8959. Herceptarg TCB, classic format. FIG. 42D: ID 8997. FolR1 36F2 TCB, classic format, anti ID CH2527 scFv 4.32.63 MK062 MMP9 linker N-terminally fused to VH. FIG. 42E: ID 8998. FolR1 36F2 TCB, classic format, anti ID CH2527 scFv 4.32.63 non cleavable linker N-terminally fused to VH. FIG. 42F: ID 8996. FolR1 36F2 TCB, classic format.

[0207] FIG. 43 depicts Human Bronchial Epithelial Cell toxicity mediated by human PBMCs and 100 nM or 10 nM of TCBs. Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB. Each point represents the mean value of triplicates. Standard deviation is indicated by error bars.

[0208] FIG. 44 depicts FolR1 negative target cell (Mkn-45) cytotoxicity mediated by 100 nM of FolR1 TCBs and human PBMCs. Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB. Each point represents the mean value of triplicates. Standard deviation is indicated by error bars.

[0209] FIG. 45A to FIG. 45N depict schematic diagrams of exemplary antibody constructs.

DETAILED DESCRIPTION

Definitions

[0210] Terms are used herein as generally used in the art, unless otherwise defined in the following. As used herein, the term “antigen binding molecule” refers in its broadest sense to a molecule that specifically binds an antigenic determinant. Examples of antigen binding molecules are immunoglobulins and derivatives, e.g., fragments, thereof.

[0211] The term “bispecific” means that the antigen binding molecule is able to specifically bind to at least two distinct antigenic determinants. Typically, a bispecific antigen binding molecule comprises two antigen binding sites, each of which is specific for a different antigenic determinant. In certain embodiments the bispecific antigen binding molecule is capable of simultaneously binding two antigenic determinants, particularly two antigenic determinants expressed on two distinct cells.

[0212] The term “valent” as used herein denotes the presence of a specified number of antigen binding sites in an antigen binding molecule. As such, the term “monovalent binding to an antigen” denotes the presence of one (and not more than one) antigen binding site specific for the antigen in the antigen binding molecule.

[0213] An “antigen binding site” refers to the site, i.e. one or more amino acid residues, of an antigen binding molecule which provides interaction with the antigen. For example, the antigen binding site of an antibody comprises amino acid residues from the complementarity determining regions

(CDRs). A native immunoglobulin molecule typically has two antigen binding sites, a Fab molecule typically has a single antigen binding site.

[0214] As used herein, the term “antigen binding moiety” refers to a polypeptide molecule that specifically binds to an antigenic determinant. In one embodiment, an antigen binding moiety is able to direct the entity to which it is attached (e.g., a second antigen binding moiety) to a target site, for example to a specific type of tumor cell or tumor stroma bearing the antigenic determinant. In another embodiment an antigen binding moiety is able to activate signaling through its target antigen, for example a T cell receptor complex antigen. Antigen binding moieties include antibodies and fragments thereof as further defined herein. Particular antigen binding moieties include an antigen binding domain of an antibody, comprising an antibody heavy chain variable region and an antibody light chain variable region. In certain embodiments, the antigen binding moieties may comprise antibody constant regions as further defined herein and known in the art. Useful heavy chain constant regions include any of the five isotypes: α , δ , ϵ , γ , or μ . Useful light chain constant regions include any of the two isotypes: κ and λ .

[0215] As used herein, the term “antigenic determinant” is synonymous with “antigen” and “epitope,” and refers to a site (e.g., a contiguous stretch of amino acids or a conformational configuration made up of different regions of non-contiguous amino acids) on a polypeptide macromolecule to which an antigen binding moiety binds, forming an antigen binding moiety-antigen complex. Useful antigenic determinants can be found, for example, on the surfaces of tumor cells, on the surfaces of virus-infected cells, on the surfaces of other diseased cells, on the surface of immune cells, free in blood serum, and/or in the extracellular matrix (ECM). The proteins referred to as antigens herein (e.g., FolR1, HER1, HER2, CD3, Mesothelin) can be any native form of the proteins from any vertebrate source, including mammals such as primates (e.g., humans) and rodents (e.g., mice and rats), unless otherwise indicated. In a particular embodiment the antigen is a human protein. Where reference is made to a specific protein herein, the term encompasses the “full-length”, unprocessed protein as well as any form of the protein that results from processing in the cell. The term also encompasses naturally occurring variants of the protein, e.g., splice variants or allelic variants. Exemplary human proteins useful as antigens include, but are not limited to: FolR1, HER1 and CD3, particularly the epsilon subunit of CD3 (see UniProt no. P07766 (version 130), NCBI RefSeq no. NP_000724.1, SEQ ID NO: 54 for the human sequence; or UniProt no. Q95LI5 (version 49), NCBI GenBank no. BAB71849.1 for the cynomolgus [*Macaca fascicularis*] sequence). In certain embodiments the protease-activatable T cell activating bispecific molecule of the invention binds to an epitope of CD3 or a target cell antigen that is conserved among the CD3 or target antigen from different species. In certain embodiments the protease-activatable T cell activating bispecific molecule of the invention binds to CD3 and FolR1, but does not bind to FolR2 or FolR3. In certain embodiments the protease-activatable T cell activating bispecific molecule of the invention binds to CD3 and HER1. In certain embodiments the protease-activatable T cell activating bispecific molecule of the invention binds to CD3 and Mesothelin. In certain embodiments the protease-activatable T cell activating bispecific

molecule of the invention binds to CD3 and HER2. By “specific binding” is meant that the binding is selective for the antigen and can be discriminated from unwanted or non-specific interactions. The ability of an antigen binding moiety to bind to a specific antigenic determinant can be measured either through an enzyme-linked immunosorbent assay (ELISA) or other techniques familiar to one of skill in the art, e.g., surface plasmon resonance (SPR) technique (analyzed on a BIAcore instrument) (Liljeblad et al., Glyco J 17, 323-329 (2000)), and traditional binding assays (Heeley, Endocr Res 28, 217-229 (2002)). In one embodiment, the extent of binding of an antigen binding moiety to an unrelated protein is less than about 10% of the binding of the antigen binding moiety to the antigen as measured, e.g., by SPR. In certain embodiments, an antigen binding moiety that binds to the antigen, or an antigen binding molecule comprising that antigen binding moiety, has a dissociation constant (K_D) of $\leq 1 \mu\text{M}$, $\leq 100 \text{ nM}$, $\leq 10 \text{ nM}$, $\leq 1 \text{ nM}$, $\leq 0.1 \text{ nM}$, $\leq 0.01 \text{ nM}$, or $\leq 0.001 \text{ nM}$ (e.g., 10^{-8} M or less, e.g., from 10^{-8} M to 10^{-13} M , e.g., from 10^{-9} M to 10^{-13} M). “Affinity” refers to the strength of the sum total of non-covalent interactions between a single binding site of a molecule (e.g., a receptor) and its binding partner (e.g., a ligand). Unless indicated otherwise, as used herein, “binding affinity” refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., an antigen binding moiety and an antigen, or a receptor and its ligand). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_D), which is the ratio of dissociation and association rate constants (k_{off} and k_{on} , respectively). Thus, equivalent affinities may comprise different rate constants, as long as the ratio of the rate constants remains the same. Affinity can be measured by well-established methods known in the art, including those described herein. A particular method for measuring affinity is Surface Plasmon Resonance (SPR).

[0216] “Reduced binding”, for example reduced binding to an Fc receptor, refers to a decrease in affinity for the respective interaction, as measured for example by SPR. For clarity the term includes also reduction of the affinity to zero (or below the detection limit of the analytic method), i.e. complete abolishment of the interaction. Conversely, “increased binding” refers to an increase in binding affinity for the respective interaction.

[0217] “T cell activation” as used herein refers to one or more cellular response of a T lymphocyte, particularly a cytotoxic T lymphocyte, selected from: proliferation, differentiation, cytokine secretion, cytotoxic effector molecule release, cytotoxic activity, and expression of activation markers. The protease-activatable T cell activating bispecific molecules of the invention are capable of inducing T cell activation. Suitable assays to measure T cell activation are known in the art described herein.

[0218] A “target cell antigen” as used herein refers to an antigenic determinant presented on the surface of a target cell, for example a cell in a tumor such as a cancer cell or a cell of the tumor stroma. As used herein, the terms “first” and “second” with respect to antigen binding moieties etc., are used for convenience of distinguishing when there is more than one of each type of moiety. Use of these terms is not intended to confer a specific order or orientation of the protease-activatable T cell activating bispecific molecule unless explicitly so stated.

[0219] A “Fab molecule” refers to a protein consisting of the VH and CH1 domain of the heavy chain (the “Fab heavy chain”) and the VL and CL domain of the light chain (the “Fab light chain”) of an immunoglobulin.

[0220] By “fused” is meant that the components (e.g., a Fab molecule and an Fc domain subunit) are linked by peptide bonds, either directly or via one or more peptide linkers.

[0221] As used herein, the term “single-chain” refers to a molecule comprising amino acid monomers linearly linked by peptide bonds. In certain embodiments, one of the antigen binding moieties is a single-chain Fab molecule, i.e. a Fab molecule wherein the Fab light chain and the Fab heavy chain are connected by a peptide linker to form a single peptide chain. In a particular such embodiment, the C-terminus of the Fab light chain is connected to the N-terminus of the Fab heavy chain in the single-chain Fab molecule.

[0222] By a “crossover” Fab molecule (also termed “Crossfab”) is meant a Fab molecule wherein either the variable regions or the constant regions of the Fab heavy and light chain are exchanged, i.e. the crossover Fab molecule comprises a peptide chain composed of the light chain variable region and the heavy chain constant region, and a peptide chain composed of the heavy chain variable region and the light chain constant region. For clarity, in a crossover Fab molecule wherein the variable regions of the Fab light chain and the Fab heavy chain are exchanged, the peptide chain comprising the heavy chain constant region is referred to herein as the “heavy chain” of the crossover Fab molecule. Conversely, in a crossover Fab molecule wherein the constant regions of the Fab light chain and the Fab heavy chain are exchanged, the peptide chain comprising the heavy chain variable region is referred to herein as the “heavy chain” of the crossover Fab molecule.

[0223] In contrast thereto, by a “conventional” Fab molecule is meant a Fab molecule in its natural format, i.e. comprising a heavy chain composed of the heavy chain variable and constant regions (VH-CH1), and a light chain composed of the light chain variable and constant regions (VL-CL).

[0224] The term “immunoglobulin molecule” refers to a protein having the structure of a naturally occurring antibody. For example, immunoglobulins of the IgG class are heterotetrameric glycoproteins of about 150,000 daltons, composed of two light chains and two heavy chains that are disulfide-bonded. From N- to C-terminus, each heavy chain has a variable region (VH), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3), also called a heavy chain constant region. Similarly, from N- to C-terminus, each light chain has a variable region (VL), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain, also called a light chain constant region. The heavy chain of an immunoglobulin may be assigned to one of five types, called α (IgA), δ (IgD), ϵ (IgE), γ (IgG), or μ (IgM), some of which may be further divided into subtypes, e.g., γ_1 (IgG₁), γ_2 (IgG₂), γ_3 (IgG₃), γ_4 (IgG₄), α_1 (IgA₁) and α_2 (IgA₂). The light chain of an immunoglobulin may be assigned to one of two types, called kappa (κ) and lambda (λ), based on the amino acid sequence of its constant domain. An immunoglobulin essentially consists of two Fab molecules and an Fc domain, linked via the immunoglobulin hinge region.

[0225] The term “antibody” herein is used in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, and antibody fragments so long as they exhibit the desired antigen-binding activity.

[0226] An “antibody fragment” refers to a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')₂, diabodies, linear antibodies, single-chain antibody molecules (e.g., scFv), and single-domain antibodies. For a review of certain antibody fragments, see Hudson et al., Nat Med 9, 129-134 (2003). For a review of scFv fragments, see e.g., Plückthun, in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994); see also WO 93/16185; and U.S. Pat. Nos. 5,571,894 and 5,587,458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Pat. No. 5,869,046. Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. See, for example, EP 404,097; WO 1993/01161; Hudson et al., Nat Med 9, 129-134 (2003); and Hollinger et al., Proc Natl Acad Sci USA 90, 6444-6448 (1993). Triabodies and tetrabodies are also described in Hudson et al., Nat Med 9, 129-134 (2003). Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (Domantis, Inc., Waltham, Mass.; see e.g., U.S. Pat. No. 6,248,516 B 1). Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (e.g., *E. coli* or phage), as described herein.

[0227] The term “antigen binding domain” refers to the part of an antibody that comprises the area which specifically binds to and is complementary to part or all of an antigen. An antigen binding domain may be provided by, for example, one or more antibody variable domains (also called antibody variable regions). Particularly, an antigen binding domain comprises an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH).

[0228] The term “variable region” or “variable domain” refers to the domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. The variable domains of the heavy chain and light chain (VH and VL, respectively) of a native antibody generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three hypervariable regions (HVRs). See, e.g., Kindt et al., Kuby Immunology, 6th ed., W.H. Freeman and Co., page 91 (2007). A single VH or VL domain may be sufficient to confer antigen-binding specificity. The term “hypervariable region” or “HVR”, as used herein, refers to each of the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops (“hypervariable loops”). Generally, native four-chain antibodies comprise six HVRs; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). HVRs generally comprise amino acid residues from the hypervariable loops and/or from the complementarity determining regions (CDRs), the latter being of highest sequence

variability and/or involved in antigen recognition. With the exception of CDR1 in VH, CDRs generally comprise the amino acid residues that form the hypervariable loops. Hypervariable regions (HVRs) are also referred to as “complementarity determining regions” (CDRs), and these terms are used herein interchangeably in reference to portions of the variable region that form the antigen binding regions. This particular region has been described by Kabat et al., U.S. Dept. of Health and Human Services, Sequences of Proteins of Immunological Interest (1983) and by Chothia et al., J Mol Biol 196:901-917 (1987), where the definitions include overlapping or subsets of amino acid residues when compared against each other. Nevertheless, application of either definition to refer to a CDR of an antibody or variants thereof is intended to be within the scope of the term as defined and used herein. The appropriate amino acid residues which encompass the CDRs as defined by each of the above cited references are set forth below in Table 1 as a comparison. The exact residue numbers which encompass a particular CDR will vary depending on the sequence and size of the CDR. Those skilled in the art can routinely determine which residues comprise a particular CDR given the variable region amino acid sequence of the antibody.

TABLE 1

CDR Definitions ¹			
CDR	Kabat	Chothia	AbM ²
V _H CDR1	31-35	26-32	26-35
V _H CDR2	50-65	52-58	50-58
V _H CDR3	95-102	95-102	95-102
V _L CDR1	24-34	26-32	24-34
V _L CDR2	50-56	50-52	50-56
V _L CDR3	89-97	91-96	89-97

¹ Numbering of all CDR definitions in Table 1 is according to the numbering conventions set forth by Kabat et al. (see below).
² “AbM” with a lowercase “b” as used in Table 1 refers to the CDRs as defined by Oxford Molecular’s “AbM” antibody modeling software.

[0229] Kabat et al. also defined a numbering system for variable region sequences that is applicable to any antibody. One of ordinary skill in the art can unambiguously assign this system of “Kabat numbering” to any variable region sequence, without reliance on any experimental data beyond the sequence itself. As used herein, “Kabat numbering” refers to the numbering system set forth by Kabat et al., U.S. Dept. of Health and Human Services, “Sequence of Proteins of Immunological Interest” (1983). Unless otherwise specified, references to the numbering of specific amino acid residue positions in an antibody variable region are according to the Kabat numbering system.

[0230] The polypeptide sequences of the sequence listing are not numbered according to the Kabat numbering system. However, it is well within the ordinary skill of one in the art to convert the numbering of the sequences of the Sequence Listing to Kabat numbering.

[0231] “Framework” or “FR” refers to variable domain residues other than hypervariable region (HVR) residues. The FR of a variable domain generally consists of four FR domains: FR1, FR2, FR3, and FR4. Accordingly, the HVR and FR sequences generally appear in the following sequence in VH (or VL): FR1-H1(L1)-FR2-H2(L2)-FR3-H3(L3)-FR4.

[0232] The “class” of an antibody or immunoglobulin refers to the type of constant domain or constant region possessed by its heavy chain. There are five major classes of

antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α, δ, ε, γ, and μ, respectively.

[0233] The term “Fc domain” or “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain that contains at least a portion of the constant region. The term includes native sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an IgG heavy chain might vary slightly, the human IgG heavy chain Fc region is usually defined to extend from Cys226, or from Pro230, to the carboxyl-terminus of the heavy chain. However, the C-terminal lysine (Lys447) of the Fc region may or may not be present. Unless otherwise specified herein, numbering of amino acid residues in the Fc region or constant region is according to the EU numbering system, also called the EU index, as described in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., 1991. A “subunit” of an Fc domain as used herein refers to one of the two polypeptides forming the dimeric Fc domain, i.e. a polypeptide comprising C-terminal constant regions of an immunoglobulin heavy chain, capable of stable self-association. For example, a subunit of an IgG Fc domain comprises an IgG CH2 and an IgG CH3 constant domain.

[0234] A “modification promoting the association of the first and the second subunit of the Fc domain” is a manipulation of the peptide backbone or the post-translational modifications of an Fc domain subunit that reduces or prevents the association of a polypeptide comprising the Fc domain subunit with an identical polypeptide to form a homodimer. A modification promoting association as used herein particularly includes separate modifications made to each of the two Fc domain subunits desired to associate (i.e. the first and the second subunit of the Fc domain), wherein the modifications are complementary to each other so as to promote association of the two Fc domain subunits. For example, a modification promoting association may alter the structure or charge of one or both of the Fc domain subunits so as to make their association sterically or electrostatically favorable, respectively. Thus, (hetero)dimerization occurs between a polypeptide comprising the first Fc domain subunit and a polypeptide comprising the second Fc domain subunit, which might be non-identical in the sense that further components fused to each of the subunits (e.g., antigen binding moieties) are not the same. In some embodiments the modification promoting association comprises an amino acid mutation in the Fc domain, specifically an amino acid substitution. In a particular embodiment, the modification promoting association comprises a separate amino acid mutation, specifically an amino acid substitution, in each of the two subunits of the Fc domain.

[0235] The term “effector functions” refers to those biological activities attributable to the Fc region of an antibody, which vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity (CDC), Fc receptor binding, antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP), cytokine secretion, immune complex-mediated antigen uptake by antigen presenting cells, down regulation of cell surface receptors (e.g., B cell receptor), and B cell activation.

[0236] As used herein, the terms “engineer, engineered, engineering”, are considered to include any manipulation of the peptide backbone or the post-translational modifications of a naturally occurring or recombinant polypeptide or fragment thereof. Engineering includes modifications of the amino acid sequence, of the glycosylation pattern, or of the side chain group of individual amino acids, as well as combinations of these approaches.

[0237] The term “amino acid mutation” as used herein is meant to encompass amino acid substitutions, deletions, insertions, and modifications. Any combination of substitution, deletion, insertion, and modification can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics, e.g., reduced binding to an Fc receptor, or increased association with another peptide. Amino acid sequence deletions and insertions include amino- and/or carboxy-terminal deletions and insertions of amino acids. Particular amino acid mutations are amino acid substitutions. For the purpose of altering e.g., the binding characteristics of an Fc region, non-conservative amino acid substitutions, i.e. replacing one amino acid with another amino acid having different structural and/or chemical properties, are particularly preferred. Amino acid substitutions include replacement by non-naturally occurring amino acids or by naturally occurring amino acid derivatives of the twenty standard amino acids (e.g., 4-hydroxyproline, 3-methylhistidine, ornithine, homoserine, 5-hydroxylysine). Amino acid mutations can be generated using genetic or chemical methods well known in the art. Genetic methods may include site-directed mutagenesis, PCR, gene synthesis and the like. It is contemplated that methods of altering the side chain group of an amino acid by methods other than genetic engineering, such as chemical modification, may also be useful. Various designations may be used herein to indicate the same amino acid mutation. For example, a substitution from proline at position 329 of the Fc domain to glycine can be indicated as 329G, G329, G₃₂₉, P329G, or Pro329Gly.

[0238] As used herein, term “polypeptide” refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). The term “polypeptide” refers to any chain of two or more amino acids, and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides, “protein,” “amino acid chain,” or any other term used to refer to a chain of two or more amino acids, are included within the definition of “polypeptide,” and the term “polypeptide” may be used instead of, or interchangeably with any of these terms. The term “polypeptide” is also intended to refer to the products of post-expression modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, or modification by non-naturally occurring amino acids. A polypeptide may be derived from a natural biological source or produced by recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It may be generated in any manner, including by chemical synthesis. A polypeptide of the invention may be of a size of about 3 or more, 5 or more, 10 or more, 20 or more, 25 or more, 50 or more, 75 or more, 100 or more, 200 or more, 500 or more, 1,000 or more, or 2,000 or more amino acids. Polypeptides may have a defined three-dimensional structure, although they do not necessarily have such structure. Polypeptides

with a defined three-dimensional structure are referred to as folded, and polypeptides which do not possess a defined three-dimensional structure, but rather can adopt a large number of different conformations, and are referred to as unfolded.

[0239] By an “isolated” polypeptide or a variant, or derivative thereof is intended a polypeptide that is not in its natural milieu. No particular level of purification is required. For example, an isolated polypeptide can be removed from its native or natural environment. Recombinantly produced polypeptides and proteins expressed in host cells are considered isolated for the purpose of the invention, as are native or recombinant polypeptides which have been separated, fractionated, or partially or substantially purified by any suitable technique.

[0240] “Percent (%) amino acid sequence identity” with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available from Genentech, Inc., South San Francisco, Calif., or may be compiled from the source code. The ALIGN-2 program should be compiled for use on a UNIX operating system, including digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary. In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program’s alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. Unless specifically stated otherwise, all % amino acid sequence

identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

[0241] The term “polynucleotide” refers to an isolated nucleic acid molecule or construct, e.g., messenger RNA (mRNA), virally-derived RNA, or plasmid DNA (pDNA). A polynucleotide may comprise a conventional phosphodiester bond or a non-conventional bond (e.g., an amide bond, such as found in peptide nucleic acids (PNA). The term “nucleic acid molecule” refers to any one or more nucleic acid segments, e.g., DNA or RNA fragments, present in a polynucleotide.

[0242] By “isolated” nucleic acid molecule or polynucleotide is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, a recombinant polynucleotide encoding a polypeptide contained in a vector is considered isolated for the purposes of the present invention. Further examples of an isolated polynucleotide include recombinant polynucleotides maintained in heterologous host cells or purified (partially or substantially) polynucleotides in solution. An isolated polynucleotide includes a polynucleotide molecule contained in cells that ordinarily contain the polynucleotide molecule, but the polynucleotide molecule is present extrachromosomally or at a chromosomal location that is different from its natural chromosomal location. Isolated RNA molecules include in vivo or in vitro RNA transcripts of the present invention, as well as positive and negative strand forms, and double-stranded forms. Isolated polynucleotides or nucleic acids according to the present invention further include such molecules produced synthetically. In addition, a polynucleotide or a nucleic acid may be or may include a regulatory element such as a promoter, ribosome binding site, or a transcription terminator. By a nucleic acid or polynucleotide having a nucleotide sequence at least, for example, 95% “identical” to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence. As a practical matter, whether any particular polynucleotide sequence is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs, such as the ones discussed above for polypeptides (e.g., ALIGN-2).

[0243] The term “expression cassette” refers to a polynucleotide generated recombinantly or synthetically, with a series of specified nucleic acid elements that permit transcription of a particular nucleic acid in a target cell. The recombinant expression cassette can be incorporated into a plasmid, chromosome, mitochondrial DNA, plastid DNA,

virus, or nucleic acid fragment. Typically, the recombinant expression cassette portion of an expression vector includes, among other sequences, a nucleic acid sequence to be transcribed and a promoter. In certain embodiments, the expression cassette of the invention comprises polynucleotide sequences that encode bispecific antigen binding molecules of the invention or fragments thereof.

[0244] The term “vector” or “expression vector” is synonymous with “expression construct” and refers to a DNA molecule that is used to introduce and direct the expression of a specific gene to which it is operably associated in a target cell. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. The expression vector of the present invention comprises an expression cassette. Expression vectors allow transcription of large amounts of stable mRNA. Once the expression vector is inside the target cell, the ribonucleic acid molecule or protein that is encoded by the gene is produced by the cellular transcription and/or translation machinery. In one embodiment, the expression vector of the invention comprises an expression cassette that comprises polynucleotide sequences that encode bispecific antigen binding molecules of the invention or fragments thereof.

[0245] The terms “host cell”, “host cell line,” and “host cell culture” are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include “transformants” and “transformed cells,” which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell, but may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein. A host cell is any type of cellular system that can be used to generate the bispecific antigen binding molecules of the present invention. Host cells include cultured cells, e.g., mammalian cultured cells, such as CHO cells, BHK cells, NS0 cells, SP2/0 cells, YO myeloma cells, P3X63 mouse myeloma cells, PER cells, PER.C6 cells or hybridoma cells, yeast cells, insect cells, and plant cells, to name only a few, but also cells comprised within a transgenic animal, transgenic plant or cultured plant or animal tissue.

[0246] An “activating Fc receptor” is an Fc receptor that following engagement by an Fc domain of an antibody elicits signaling events that stimulate the receptor-bearing cell to perform effector functions. Human activating Fc receptors include FcγRIIIa (CD16a), FcγRI (CD64), FcγRIIa (CD32), and FcαRI (CD89).

[0247] Antibody-dependent cell-mediated cytotoxicity (ADCC) is an immune mechanism leading to the lysis of antibody-coated target cells by immune effector cells. The target cells are cells to which antibodies or derivatives thereof comprising an Fc region specifically bind, generally via the protein part that is N-terminal to the Fc region. As used herein, the term “reduced ADCC” is defined as either a reduction in the number of target cells that are lysed in a given time, at a given concentration of antibody in the medium surrounding the target cells, by the mechanism of ADCC defined above, and/or an increase in the concentration of antibody in the medium surrounding the target cells, required to achieve the lysis of a given number of target cells in a given time, by the mechanism of ADCC. The reduction

in ADCC is relative to the ADCC mediated by the same antibody produced by the same type of host cells, using the same standard production, purification, formulation and storage methods (which are known to those skilled in the art), but that has not been engineered. For example the reduction in ADCC mediated by an antibody comprising in its Fc domain an amino acid substitution that reduces ADCC, is relative to the ADCC mediated by the same antibody without this amino acid substitution in the Fc domain. Suitable assays to measure ADCC are well known in the art (see e.g., PCT publication no. WO 2006/082515 or PCT publication no. WO 2012/130831).

[0248] An “effective amount” of an agent refers to the amount that is necessary to result in a physiological change in the cell or tissue to which it is administered.

[0249] A “therapeutically effective amount” of an agent, e.g., a pharmaceutical composition, refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result. A therapeutically effective amount of an agent for example eliminates, decreases, delays, minimizes or prevents adverse effects of a disease.

[0250] An “individual” or “subject” is a mammal. Mammals include, but are not limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). Particularly, the individual or subject is a human.

[0251] The term “pharmaceutical composition” refers to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

[0252] A “pharmaceutically acceptable carrier” refers to an ingredient in a pharmaceutical composition, other than an active ingredient, which is nontoxic to a subject. A pharmaceutically acceptable carrier includes, but is not limited to, a buffer, excipient, stabilizer, or preservative.

[0253] As used herein, “treatment” (and grammatical variations thereof such as “treat” or “treating”) refers to clinical intervention in an attempt to alter the natural course of a disease in the individual being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, protease-activatable T cell activating bispecific molecules of the invention are used to delay development of a disease or to slow the progression of a disease.

[0254] The term “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, combination therapy, contraindications and/or warnings concerning the use of such therapeutic products.

[0255] An “idiotype-specific polypeptide” as used herein refers to a polypeptide that recognizes the idiotype of an antigen-binding moiety, e.g., an antigen-binding moiety specific for CD3. The idiotype-specific polypeptide is

capable of specifically binding to the variable region of the antigen-binding moiety and thereby reducing or preventing specific binding of the antigen-binding moiety to its cognate antigen. When associated with a molecule that comprises the antigen-binding moiety, the idiotype-specific polypeptide can function as a masking moiety of the molecule. Specifically disclosed herein are anti-idiotypic antibodies or anti-idiotypic-binding antibody fragments specific for the idiotype of anti-CD3 binding molecules.

[0256] “Protease” or “proteolytic enzyme” as used herein refers to any proteolytic enzyme that cleaves the linker at a recognition site and that is expressed by a target cell. Such proteases might be secreted by the target cell or remain associated with the target cell, e.g., on the target cell surface. Examples of proteases include but are not limited to metalloproteinases, e.g., matrix metalloproteinase 1-28 and A Disintegrin And Metalloproteinase (ADAM) 2, 7-12, 15, 17-23, 28-30 and 33, serine proteases, e.g., urokinase-type plasminogen activator and Matriptase, cysteine protease, aspartic proteases, and members of the cathepsin family.

[0257] “Protease activatable” as used herein, with respect to the T cell activating bispecific molecule, refers to a T cell activating bispecific molecule having reduced or abrogated ability to activate T cells due to a masking moiety that reduces or abrogates the T cell activating bispecific molecule’s ability to bind to CD3. Upon dissociation of the masking moiety by proteolytic cleavage, e.g., by proteolytic cleavage of a linker connecting the masking moiety to the T cell activating bispecific molecule, binding to CD3 is restored and the T cell activating bispecific molecule is thereby activated.

[0258] “Reversibly concealing” as used herein refers to the binding of a masking moiety or idiotype-specific polypeptide to an antigen-binding moiety or molecule such as to prevent the antigen-binding moiety or molecule from its antigen, e.g., CD3. This concealing is reversible in that the idiotype-specific polypeptide can be released from the antigen-binding moiety or molecule, e.g., by protease cleavage, and thereby freeing the antigen-binding moiety or molecule to bind to its antigen.

DETAILED DESCRIPTION

[0259] In one aspect, the invention relates to a protease-activatable T cell activating bispecific molecule comprising

[0260] (a) a first antigen binding moiety capable of specific binding to CD3;

[0261] (b) a second antigen binding moiety capable of specific binding to a target cell antigen; and

[0262] (c) a masking moiety covalently attached to the T cell bispecific binding molecule through a protease-cleavable linker, wherein the masking moiety is capable of specific binding to the idiotype of the first or the second antigen binding moiety thereby reversibly concealing the first or second antigen binding moiety.

[0263] The first antigen binding moiety capable of specific binding to CD3 comprises an idiotype. In one embodiment, the masking moiety of the protease-activatable T cell activating bispecific molecule is covalently attached to the first antigen binding moiety. In one embodiment the masking moiety is covalently attached to the heavy chain variable region of the first antigen binding moiety. In one embodiment the masking moiety is covalently attached to the light chain variable region of the first antigen binding moiety. This covalent bond is separate from the specific binding,

which is preferably non-covalent, of the masking moiety to the idiotype first antigen binding site. The idiotype of the first antigen binding moiety comprises its variable region. In one embodiment the masking moiety binds to amino acid residues that make contact with CD3 when the first antigen binding moiety is bound to CD3. In a preferred embodiment, the masking moiety is not the cognate antigen or fragments thereof of the first antigen binding moiety, i.e., the masking moiety is not a CD3 or fragments thereof. In one embodiment the masking moiety is an anti-idiotypic antibody or fragment thereof. In one embodiment, the masking moiety is an anti-idiotypic scFv. Exemplary embodiments of masking moieties which are anti-idiotypic scFv, and protease activatable T cell activating molecules comprising such masking moieties, are described in detail in the examples. In one embodiment the protease-activatable T cell activating bispecific molecule comprises a second masking moiety reversibly concealing the second antigen binding moiety.

[0264] In one embodiment the protease-activatable T cell activating bispecific molecule comprises

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19;

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to a target cell antigen.

[0265] In one embodiment the first antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence of SEQ ID NO: 55.

[0266] In one embodiment the first antigen binding moiety comprises the heavy chain variable region comprising an amino acid sequence of SEQ ID NO: 43 and the light chain variable region comprising an amino acid sequence of SEQ ID NO: 55.

[0267] In a specific embodiment the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0268] In another specific embodiment, the second antigen binding moiety is capable of specific binding to FolR1 and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0269] In a specific embodiment the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 151, SEQ ID NO: 152 and SEQ ID NO: 153

and at least one light chain CDR selected from the group of SEQ ID NO: 154, SEQ ID NO: 155 and SEQ ID NO: 156.

[0270] In another specific embodiment, the second antigen binding moiety is capable of specific binding to FolR1 and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 158.

[0271] In another specific embodiment, the second antigen binding moiety is capable of specific binding to HER1 and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 61.

[0272] In another specific embodiment, the second antigen binding moiety is capable of specific binding to HER1 and comprises a heavy chain comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence of SEQ ID NO: 32, and a light chain comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to an amino acid sequence of SEQ ID NO: 33.

[0273] In another specific embodiment, the second antigen binding moiety is capable of specific binding to HER1 and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 115 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 116.

[0274] In a specific embodiment the second antigen binding moiety is capable of specific binding to Mesothelin and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 107, SEQ ID NO: 108 and SEQ ID NO: 109 and at least one light chain CDR selected from the group of SEQ ID NO: 110, SEQ ID NO: 111 and SEQ ID NO: 112.

[0275] In another specific embodiment, the second antigen binding moiety is capable of specific binding Mesothelin and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 114.

[0276] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19;

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to FolR1 comprising at least one heavy chain complementarity determining region (CDR)

selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0277] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55,

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to FolR1 comprising heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0278] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19;

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to FolR1 comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 151, SEQ ID NO: 152 and SEQ ID NO: 153 and at least one light chain CDR selected from the group of SEQ ID NO: 154, SEQ ID NO: 155 and SEQ ID NO: 156.

[0279] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55,

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to FolR1 comprising heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 158.

[0280] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising at least one

heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19;

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to HER1 comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 61.

[0281] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to HER1 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 115 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 116.

[0282] In a particular embodiment, the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy chain are exchanged.

[0283] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19;

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Mesothelin comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 107, SEQ ID NO: 108 and SEQ ID NO: 109 and at least one light chain CDR selected from the group of SEQ ID NO: 110, SEQ ID NO: 111 and SEQ ID NO: 112.

[0284] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55,

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Mesothelin comprising heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 114.

[0285] In one embodiment, the second antigen binding moiety is a conventional Fab molecule.

[0286] In a particular embodiment, the first antigen binding moiety is a crossover Fab molecule wherein the constant regions of the Fab light chain and the Fab heavy chain are exchanged, and the second antigen binding moiety is a conventional Fab molecule. In a further particular embodiment, the first and the second antigen binding moiety are fused to each other, optionally through a peptide linker.

[0287] In particular embodiments, the protease-activatable T cell activating bispecific molecule further comprises an Fc domain composed of a first and a second subunit capable of stable association. In a further particular embodiment, not more than one antigen binding moiety capable of specific binding to CD3 is present in the protease-activatable T cell activating bispecific molecule (i.e. the protease-activatable T cell activating bispecific molecule provides monovalent binding to CD3).

Protease-Activatable T Cell Activating Bispecific Molecule Formats

[0288] The components of the protease-activatable T cell activating bispecific molecule can be fused to each other in a variety of configurations. Exemplary configurations are depicted in FIGS. 1A-1E and 5A-5H. Further exemplary configurations are depicted in FIGS. 33A-33K.

[0289] In particular embodiments, the protease-activatable T cell activating bispecific molecule comprises an Fc domain composed of a first and a second subunit capable of stable association. In some embodiments, the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain.

[0290] In one such embodiment, the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. In a specific such embodiment, the protease-activatable T cell activating bispecific molecule essentially consists of a first and a second antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety, and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

[0291] In another such embodiment, the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain. In a specific such embodiment, the protease-activatable T cell activating bispecific molecule essentially

consists of a first and a second antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first and the second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain.

[0292] In other embodiments, the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain.

[0293] In a particular such embodiment, the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In a specific such embodiment, the protease-activatable T cell activating bispecific molecule essentially consists of a first and a second antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

[0294] The antigen binding moieties may be fused to the Fc domain or to each other directly or through a peptide linker, comprising one or more amino acids, typically about 2-20 amino acids. Peptide linkers are known in the art and are described herein. Suitable, non-immunogenic peptide linkers include, for example, $(G_4S)_n$, $(SG_4)_n$, $(G_4S)_n$ or $G_4(SG_4)_n$ peptide linkers. "n" is generally a number between 1 and 10, typically between 2 and 4. A particularly suitable peptide linker for fusing the Fab light chains of the first and the second antigen binding moiety to each other is $(G_4S)_2$. An exemplary peptide linker suitable for connecting the Fab heavy chains of the first and the second antigen binding moiety is EPKSC(D)- $(G_4S)_2$ (SEQ ID NOs 105 and 106). Additionally, linkers may comprise (a portion of) an immunoglobulin hinge region. Particularly where an antigen binding moiety is fused to the N-terminus of an Fc domain subunit, it may be fused via an immunoglobulin hinge region or a portion thereof, with or without an additional peptide linker.

[0295] A protease-activatable T cell activating bispecific molecule with a single antigen binding moiety capable of specific binding to a target cell antigen is useful, particularly in cases where internalization of the target cell antigen is to be expected following binding of a high affinity antigen binding moiety. In such cases, the presence of more than one antigen binding moiety specific for the target cell antigen may enhance internalization of the target cell antigen, thereby reducing its availability.

[0296] In many other cases, however, it will be advantageous to have a protease-activatable T cell activating bispecific molecule comprising two or more antigen binding moieties specific for a target cell antigen (see examples in shown in FIGS. 5A-5H), for example to optimize targeting to the target site or to allow crosslinking of target cell antigens.

[0297] Accordingly, in certain embodiments, the protease-activatable T cell activating bispecific molecule of the

[0299] In a particular embodiment, the first antigen binding moiety is capable of specific binding to CD3, and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19; and the second and third antigen binding moieties are capable of specific binding to FolR1, wherein the second and third antigen binding moieties comprise at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0301] In a particular embodiment, the first antigen binding moiety is capable of specific binding to CD3, and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, and the second and third antigen binding moieties are capable of specific binding to FoIR1, wherein the second and third antigen binding moi-

eties comprise a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0303] In one embodiment, the first antigen binding moiety is capable of specific binding to CD3, and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19; and the second and third antigen binding moieties are capable of specific binding to HER1, wherein the second and third antigen binding moieties comprise at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 61.

[0304] In one embodiment, the first antigen binding moiety is capable of specific binding to CD3, and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19; and the second and third antigen binding moieties are capable of specific binding to HER1, wherein the second and third antigen binding moieties comprise at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 61.

[0305] In one embodiment, the first antigen binding moiety is capable of specific binding to CD3, and comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, and the second and third antigen binding moieties are capable of specific binding to HER1, wherein the second and third antigen binding moieties comprise a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 115 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 116.

[0306] In one embodiment, the first antigen binding moiety is capable of specific binding to CD3, and the second and

N-terminus of the first or second subunit of the Fc domain. In a more specific embodiment, the second and the third antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

[0315] The second and the third antigen binding moiety may be fused to the Fc domain directly or through a peptide linker. In a particular embodiment the second and the third antigen binding moiety are each fused to the Fc domain through an immunoglobulin hinge region. In a specific embodiment, the immunoglobulin hinge region is a human IgG₁ hinge region. In one embodiment the second and the third antigen binding moiety and the Fc domain are part of an immunoglobulin molecule. In a particular embodiment the immunoglobulin molecule is an IgG class immunoglobulin. In an even more particular embodiment the immunoglobulin is an IgG₁ subclass immunoglobulin. In another embodiment the immunoglobulin is an IgG₄ subclass immunoglobulin. In a further particular embodiment the immunoglobulin is a human immunoglobulin. In other embodiments the immunoglobulin is a chimeric immunoglobulin or a humanized immunoglobulin. In one embodiment, the protease-activatable T cell activating bispecific molecule essentially consists of an immunoglobulin molecule capable of specific binding to a target cell antigen, and an antigen binding moiety capable of specific binding to CD3 wherein the antigen binding moiety is a Fab molecule, particularly a crossover Fab molecule, fused to the N-terminus of one of the immunoglobulin heavy chains, optionally via a peptide linker.

[0316] In a particular embodiment, the first and the third antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain, and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In a specific such embodiment, the protease-activatable T cell activating bispecific molecule essentially consists of a first, a second and a third antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain, and wherein the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the second subunit of the Fc domain. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

[0317] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising the heavy chain complementarity determining region (CDR) 1 of SEQ

ID NO: 44, the heavy chain CDR 2 of SEQ ID NO: 45, the heavy chain CDR 3 of SEQ ID NO: 46, the light chain CDR 1 of SEQ ID NO: 17, the light chain CDR 2 of SEQ ID NO: 18 and the light chain CDR 3 of SEQ ID NO: 19, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to FoIR1 comprising the heavy chain CDR 1 of SEQ ID NO: 14, the heavy chain CDR 2 of SEQ ID NO: 15, the heavy chain CDR 3 of SEQ ID NO: 16, the light chain CDR 1 of SEQ ID NO: 17, the light chain CDR 2 of SEQ ID NO: 18 and the light chain CDR3 of SEQ ID NO: 19.

[0318] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to FoIR1 comprising heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0319] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising the heavy chain complementarity determining region (CDR) 1 of SEQ ID NO: 44, the heavy chain CDR 2 of SEQ ID NO: 45, the heavy chain CDR 3 of SEQ ID NO: 46, the light chain CDR 1 of SEQ ID NO: 17, the light chain CDR 2 of SEQ ID NO: 18 and the light chain CDR 3 of SEQ ID NO: 19, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to FoIR1 comprising the heavy chain CDR 1 of SEQ ID NO: 151, the heavy chain CDR 2 of SEQ ID NO: 152, the heavy chain CDR 3 of SEQ ID NO: 153, the light chain CDR 1 of SEQ ID NO: 154, the light chain CDR 2 of SEQ ID NO: 155 and the light chain CDR3 of SEQ ID NO: 156.

[0320] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to FolR1 comprising heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 158.

[0321] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising the heavy chain complementarity determining region (CDR) 1 of SEQ ID NO: 44, the heavy chain CDR 2 of SEQ ID NO: 45, the heavy chain CDR 3 of SEQ ID NO: 46, the light chain CDR 1 of SEQ ID NO: 17, the light chain CDR 2 of SEQ ID NO: 18 and the light chain CDR 3 of SEQ ID NO: 19, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to HER1 comprising the heavy chain CDR 1 of SEQ ID NO: 56, the heavy chain CDR 2 of SEQ ID NO: 57, the heavy chain CDR 3 of SEQ ID NO: 58, the light chain CDR 1 of SEQ ID NO: 59, the light chain CDR 2 of SEQ ID NO: 60 and the light chain CDR3 of SEQ ID NO: 61.

[0322] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to HER1 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 115 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%,

97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 116.

[0323] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising the heavy chain complementarity determining region (CDR) 1 of SEQ ID NO: 44, the heavy chain CDR 2 of SEQ ID NO: 45, the heavy chain CDR 3 of SEQ ID NO: 46, the light chain CDR 1 of SEQ ID NO: 17, the light chain CDR 2 of SEQ ID NO: 18 and the light chain CDR 3 of SEQ ID NO: 19, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to HER2, wherein the second antigen binding moiety comprises the heavy chain CDR 1 of SEQ ID NO: 142, the heavy chain CDR 2 of SEQ ID NO: 143, the heavy chain CDR 3 of SEQ ID NO: 144, the light chain CDR 1 of SEQ ID NO: 148, the light chain CDR 2 of SEQ ID NO: 149 and the light chain CDR3 of SEQ ID NO: 150, and wherein the third antigen binding moiety comprises the heavy chain CDR 1 of SEQ ID NO: 145, the heavy chain CDR 2 of SEQ ID NO: 146, the heavy chain CDR 3 of SEQ ID NO: 148, the light chain CDR 1 of SEQ ID NO: 148, the light chain CDR 2 of SEQ ID NO: 149 and the light chain CDR3 of SEQ ID NO: 150.

[0324] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to HER2, wherein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 160 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 161, and wherein the third antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 159 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 161.

[0325] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, comprising the heavy chain complementarity determining region (CDR) 1 of SEQ ID NO: 44, the heavy chain CDR 2 of SEQ ID NO: 45, the heavy chain CDR 3 of SEQ ID NO: 46, the light chain CDR 1 of SEQ ID NO: 17, the light chain CDR 2 of SEQ ID NO: 18 and the light chain CDR 3 of SEQ ID NO: 19, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to Mesothelin comprising the heavy chain CDR 1 of SEQ ID NO: 107, the heavy chain CDR 2 of SEQ ID NO: 108, the heavy chain CDR 3 of SEQ ID NO: 109, the light chain CDR 1 of SEQ ID NO: 110, the light chain CDR 2 of SEQ ID NO: 111 and the light chain CDR3 of SEQ ID NO: 112.

[0326] In one embodiment the present invention provides a protease-activatable T cell activating bispecific molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55, wherein the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions, particularly the constant regions, of the Fab light chain and the Fab heavy chain are exchanged;

(ii) a second and a third antigen binding moiety each of which is a Fab molecule capable of specific binding to Mesothelin comprising heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 114. The protease-activatable T cell activating bispecific molecule according to any of the ten above embodiments may further comprise (iii) an Fc domain composed of a first and a second subunit capable of stable association, wherein the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain, and wherein the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the second subunit of the Fc domain. In some of the protease-activatable T cell activating bispecific molecule of the invention, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety are fused to each other, optionally via a linker peptide. Depending on the configuration of the first and the second antigen binding moiety, the Fab light chain of the first antigen binding moiety may be fused at its C-terminus

to the N-terminus of the Fab light chain of the second antigen binding moiety, or the Fab light chain of the second antigen binding moiety may be fused at its C-terminus to the N-terminus of the Fab light chain of the first antigen binding moiety. Fusion of the Fab light chains of the first and the second antigen binding moiety further reduces mispairing of unmatched Fab heavy and light chains, and also reduces the number of plasmids needed for expression of some of the protease-activatable T cell activating bispecific molecule of the invention. In certain embodiments the protease-activatable T cell activating bispecific molecule comprises a polypeptide wherein the Fab light chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety (i.e. a the first antigen binding moiety comprises a crossover Fab heavy chain, wherein the heavy chain variable region is replaced by a light chain variable region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VL₍₁₎-CH1₍₁₎-CH2-CH3(-CH4)), and a polypeptide wherein a the Fab heavy chain of the second antigen binding moiety shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₂₎-CH1₍₂₎-CH2-CH3(-CH4)). In some embodiments the protease-activatable T cell activating bispecific molecule further comprises a polypeptide wherein the Fab heavy chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety (VH₍₁₎-CL₍₁₎) and the Fab light chain polypeptide of the second antigen binding moiety (VL₍₂₎-CL₍₂₎). In certain embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

[0327] In alternative embodiments the protease-activatable T cell activating bispecific molecule comprises a polypeptide wherein the Fab heavy chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety (i.e. the first antigen binding moiety comprises a crossover Fab heavy chain, wherein the heavy chain constant region is replaced by a light chain constant region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₁₎-CL₍₁₎-CH2-CH3(-CH4)), and a polypeptide wherein the Fab heavy chain of the second antigen binding moiety shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₂₎-CH1₍₂₎-CH2-CH3(-CH4)). In some embodiments the protease-activatable T cell activating bispecific molecule further comprises a polypeptide wherein the Fab light chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety (VL₍₁₎-CH1₍₁₎) and the Fab light chain polypeptide of the second antigen binding moiety (VL₍₂₎-CL₍₂₎). In certain embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

[0328] In some embodiments, the protease-activatable T cell activating bispecific molecule comprises a polypeptide wherein the Fab light chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety (i.e. the first antigen binding moiety comprises a crossover Fab heavy chain, wherein the heavy chain variable region is replaced by a light chain variable region), which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain of the second antigen binding

moiety, which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VL₍₁₎-CH1₍₁₎-VH₍₂₎-CH1₍₂₎-CH2-CH3(-CH4)). In other embodiments, the protease-activatable T cell activating bispecific molecule comprises a polypeptide wherein the Fab heavy chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety (i.e. the first antigen binding moiety comprises a crossover Fab heavy chain, wherein the heavy chain constant region is replaced by a light chain constant region), which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain of the second antigen binding moiety, which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₁₎-CL₍₁₎-VH₍₂₎-CH1₍₂₎-CH2-CH3(-CH4)). In still other embodiments, the protease-activatable T cell activating bispecific molecule comprises a polypeptide wherein the Fab heavy chain of the second antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain variable region of the first antigen binding moiety which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety (i.e. the first antigen binding moiety comprises a crossover Fab heavy chain, wherein the heavy chain variable region is replaced by a light chain variable region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₂₎-CH1₍₂₎-VL₍₁₎-CH1₍₁₎-CH2-CH3(-CH4)). In other embodiments, the protease-activatable T cell activating bispecific molecule comprises a polypeptide wherein the Fab heavy chain of the second antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain variable region of the first antigen binding moiety which in turn shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety (i.e. the first antigen binding moiety comprises a crossover Fab heavy chain, wherein the heavy chain constant region is replaced by a light chain constant region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₂₎-CH1₍₂₎-VH₍₁₎-CL₍₁₎-CH2-CH3(-CH4)).

[0329] In some of these embodiments the protease-activatable T cell activating bispecific molecule further comprises a crossover Fab light chain polypeptide of the first antigen binding moiety, wherein the Fab heavy chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety (VH₍₁₎-CL₍₁₎), and the Fab light chain polypeptide of the second antigen binding moiety (VL₍₂₎-CL₍₂₎). In others of these embodiments the protease-activatable T cell activating bispecific molecule further comprises a crossover Fab light chain polypeptide, wherein the Fab light chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety (VL₍₁₎-CH1₍₁₎), and the Fab light chain polypeptide of the second antigen binding moiety (VL₍₂₎-CL₍₂₎). In still others of these embodiments the protease-activatable T cell activating bispecific molecule further comprises a polypeptide wherein the Fab light chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety which in turn shares a carboxy-terminal peptide bond with the Fab light chain polypeptide of the

second antigen binding moiety (VL₍₁₎-CH1₍₁₎-VL₍₂₎-CL₍₂₎), a polypeptide wherein the Fab heavy chain variable region of the first antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety which in turn shares a carboxy-terminal peptide bond with the Fab light chain polypeptide of the second antigen binding moiety (VH₍₁₎-CL₍₁₎-VL₍₂₎-CL₍₂₎), a polypeptide wherein the Fab light chain polypeptide of the second antigen binding moiety shares a carboxy-terminal peptide bond with the Fab light chain variable region of the first antigen binding moiety which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the first antigen binding moiety (VL₍₂₎-CL₍₂₎-VL₍₁₎-CH1₍₁₎), or a polypeptide wherein the Fab light chain polypeptide of the second antigen binding moiety shares a carboxy-terminal peptide bond with the Fab heavy chain variable region of the first antigen binding moiety which in turn shares a carboxy-terminal peptide bond with the Fab light chain constant region of the first antigen binding moiety (VL₍₂₎-CL₍₂₎-VH₍₁₎-CL₍₁₎). The protease-activatable T cell activating bispecific molecule according to these embodiments may further comprise (i) an Fc domain subunit polypeptide (CH2-CH3(-CH4)), or (ii) a polypeptide wherein the Fab heavy chain of a third antigen binding moiety shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₃₎-CH1₍₃₎-CH2-CH3(-CH4)) and the Fab light chain polypeptide of a third antigen binding moiety (VL₍₃₎-CL₍₃₎). In certain embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

[0330] According to any of the above embodiments, components of the protease-activatable T cell activating bispecific molecule (e.g., antigen binding moiety, Fc domain) may be fused directly or through various linkers, particularly peptide linkers comprising one or more amino acids, typically about 2-20 amino acids, that are described herein or are known in the art. Suitable, non-immunogenic peptide linkers include, for example, (G₄S)_n, (SG₄)_n, (G₄S)_n or G₄(SG₄)_n peptide linkers, wherein n is generally a number between 1 and 10, typically between 2 and 4.

Fc Domain

[0331] The Fc domain of the protease-activatable T cell activating bispecific molecule consists of a pair of polypeptide chains comprising heavy chain domains of an immunoglobulin molecule. For example, the Fc domain of an immunoglobulin G (IgG) molecule is a dimer, each subunit of which comprises the CH2 and CH3 IgG heavy chain constant domains. The two subunits of the Fc domain are capable of stable association with each other. In one embodiment the protease-activatable T cell activating bispecific molecule of the invention comprises not more than one Fc domain.

[0332] In one embodiment according to the invention the Fc domain of the protease-activatable T cell activating bispecific molecule is an IgG Fc domain. In a particular embodiment the Fc domain is an IgG₁ Fc domain. In another embodiment the Fc domain is an IgG₄ Fc domain. In a more specific embodiment, the Fc domain is an IgG₄ Fc domain comprising an amino acid substitution at position S228 (Kabat numbering), particularly the amino acid substitution S228P. This amino acid substitution reduces in vivo Fab arm exchange of IgG₄ antibodies (see Stubenrauch et al., Drug Metabolism and Disposition 38, 84-91 (2010)). In a further particular embodiment the Fc domain is human.

Fc Domain Modifications Promoting Heterodimerization

[0333] Protease-activatable T cell activating bispecific molecules according to the invention comprise different antigen binding moieties, fused to one or the other of the two subunits of the Fc domain, thus the two subunits of the Fc domain are typically comprised in two non-identical polypeptide chains. Recombinant co-expression of these polypeptides and subsequent dimerization leads to several possible combinations of the two polypeptides. To improve the yield and purity of protease-activatable T cell activating bispecific molecules in recombinant production, it will thus be advantageous to introduce in the Fc domain of the protease-activatable T cell activating bispecific molecule a modification promoting the association of the desired polypeptides.

[0334] Accordingly, in particular embodiments the Fc domain of the protease-activatable T cell activating bispecific molecule according to the invention comprises a modification promoting the association of the first and the second subunit of the Fc domain. The site of most extensive protein-protein interaction between the two subunits of a human IgG Fc domain is in the CH3 domain of the Fc domain. Thus, in one embodiment said modification is in the CH3 domain of the Fc domain.

[0335] In a specific embodiment said modification is a so-called “knob-into-hole” modification, comprising a “knob” modification in one of the two subunits of the Fc domain and a “hole” modification in the other one of the two subunits of the Fc domain.

[0336] The knob-into-hole technology is described e.g., in U.S. Pat. Nos. 5,731,168; 7,695,936; Ridgway et al., *Prot Eng* 9, 617-621 (1996) and Carter, *J Immunol Meth* 248, 7-15 (2001). Generally, the method involves introducing a protuberance (“knob”) at the interface of a first polypeptide and a corresponding cavity (“hole”) in the interface of a second polypeptide, such that the protuberance can be positioned in the cavity so as to promote heterodimer formation and hinder homodimer formation. Protuberances are constructed by replacing small amino acid side chains from the interface of the first polypeptide with larger side chains (e.g., tyrosine or tryptophan). Compensatory cavities of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). Accordingly, in a particular embodiment, in the CH3 domain of the first subunit of the Fc domain of the protease-activatable T cell activating bispecific molecule an amino acid residue is replaced with an amino acid residue having a larger side chain volume, thereby generating a protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and in the CH3 domain of the second subunit of the Fc domain an amino acid residue is replaced with an amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable.

[0337] The protuberance and cavity can be made by altering the nucleic acid encoding the polypeptides, e.g., by site-specific mutagenesis, or by peptide synthesis.

[0338] In a specific embodiment, in the CH3 domain of the first subunit of the Fc domain the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in the CH3 domain of the second subunit of the Fc

domain the tyrosine residue at position 407 is replaced with a valine residue (Y407V). In one embodiment, in the second subunit of the Fc domain additionally the threonine residue at position 366 is replaced with a serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A).

[0339] In yet a further embodiment, in the first subunit of the Fc domain additionally the serine residue at position 354 is replaced with a cysteine residue (S354C), and in the second subunit of the Fc domain additionally the tyrosine residue at position 349 is replaced by a cysteine residue (Y349C). Introduction of these two cysteine residues results in formation of a disulfide bridge between the two subunits of the Fc domain, further stabilizing the dimer (Carter, *J Immunol Methods* 248, 7-15 (2001)).

[0340] In a particular embodiment the antigen binding moiety capable of binding to CD3 is fused (optionally via the antigen binding moiety capable of binding to a target cell antigen) to the first subunit of the Fc domain (comprising the “knob” modification). Without wishing to be bound by theory, fusion of the antigen binding moiety capable of binding to CD3 to the knob-containing subunit of the Fc domain will (further) minimize the generation of antigen binding molecules comprising two antigen binding moieties capable of binding to CD3 (steric clash of two knob-containing polypeptides).

[0341] In an alternative embodiment a modification promoting association of the first and the second subunit of the Fc domain comprises a modification mediating electrostatic steering effects, e.g., as described in PCT publication WO 2009/089004. Generally, this method involves replacement of one or more amino acid residues at the interface of the two Fc domain subunits by charged amino acid residues so that homodimer formation becomes electrostatically unfavorable but heterodimerization electrostatically favorable.

Fc Domain Modifications Reducing Fc Receptor Binding and/or Effector Function

[0342] The Fc domain confers to the protease-activatable T cell activating bispecific molecule favorable pharmacokinetic properties, including a long serum half-life which contributes to good accumulation in the target tissue and a favorable tissue-blood distribution ratio. At the same time it may, however, lead to undesirable targeting of the protease-activatable T cell activating bispecific molecule to cells expressing Fc receptors rather than to the preferred antigen-bearing cells. Moreover, the co-activation of Fc receptor signaling pathways may lead to cytokine release which, in combination with the T cell activating properties and the long half-life of the antigen binding molecule, results in excessive activation of cytokine receptors and severe side effects upon systemic administration. Activation of (Fc receptor-bearing) immune cells other than T cells may even reduce efficacy of the protease-activatable T cell activating bispecific molecule due to the potential destruction of T cells e.g., by NK cells.

[0343] Accordingly, in particular embodiments the Fc domain of the protease-activatable T cell activating bispecific molecules according to the invention exhibits reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain. In one such embodiment the Fc domain (or the protease-activatable T cell activating bispecific molecule comprising said Fc domain) exhibits less than 50%, preferably less than 20%, more preferably less than 10% and most preferably less than

5% of the binding affinity to an Fc receptor, as compared to a native IgG₁ Fc domain (or a protease-activatable T cell activating bispecific molecule comprising a native IgG₁ Fc domain), and/or less than 50%, preferably less than 20%, more preferably less than 10% and most preferably less than 5% of the effector function, as compared to a native IgG₁ Fc domain (or a protease-activatable T cell activating bispecific molecule comprising a native IgG₁ Fc domain). In one embodiment, the Fc domain (or the protease-activatable T cell activating bispecific molecule comprising said Fc domain) does not substantially bind to an Fc receptor and/or induce effector function. In a particular embodiment the Fc receptor is an Fcγ receptor. In one embodiment the Fc receptor is a human Fc receptor. In one embodiment the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. In one embodiment the effector function is one or more selected from the group of CDC, ADCC, ADCP, and cytokine secretion. In a particular embodiment the effector function is ADCC. In one embodiment the Fc domain domain exhibits substantially similar binding affinity to neonatal Fc receptor (FcRn), as compared to a native IgG₁ Fc domain domain. Substantially similar binding to FcRn is achieved when the Fc domain (or the protease-activatable T cell activating bispecific molecule comprising said Fc domain) exhibits greater than about 70%, particularly greater than about 80%, more particularly greater than about 90% of the binding affinity of a native IgG₁ Fc domain (or the protease-activatable T cell activating bispecific molecule comprising a native IgG₁ Fc domain) to FcRn.

[0344] In certain embodiments the Fc domain is engineered to have reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a non-engineered Fc domain. In particular embodiments, the Fc domain of the protease-activatable T cell activating bispecific molecule comprises one or more amino acid mutation that reduces the binding affinity of the Fc domain to an Fc receptor and/or effector function. Typically, the same one or more amino acid mutation is present in each of the two subunits of the Fc domain. In one embodiment the amino acid mutation reduces the binding affinity of the Fc domain to an Fc receptor. In one embodiment the amino acid mutation reduces the binding affinity of the Fc domain to an Fc receptor by at least 2-fold, at least 5-fold, or at least 10-fold. In embodiments where there is more than one amino acid mutation that reduces the binding affinity of the Fc domain to the Fc receptor, the combination of these amino acid mutations may reduce the binding affinity of the Fc domain to an Fc receptor by at least 10-fold, at least 20-fold, or even at least 50-fold. In one embodiment the protease-activatable T cell activating bispecific molecule comprising an engineered Fc domain exhibits less than 20%, particularly less than 10%, more particularly less than 5% of the binding affinity to an Fc receptor as compared to a protease-activatable T cell activating bispecific molecule comprising a non-engineered Fc domain. In a particular embodiment the Fc receptor is an Fcγ receptor. In some embodiments the Fc receptor is a human Fc receptor. In some embodiments the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human

FcγRIIIa. Preferably, binding to each of these receptors is reduced. In some embodiments binding affinity to a complement component, specifically binding affinity to C1q, is also reduced. In one embodiment binding affinity to neonatal Fc receptor (FcRn) is not reduced. Substantially similar binding to FcRn, i.e. preservation of the binding affinity of the Fc domain to said receptor, is achieved when the Fc domain (or the protease-activatable T cell activating bispecific molecule comprising said Fc domain) exhibits greater than about 70% of the binding affinity of a non-engineered form of the Fc domain (or the protease-activatable T cell activating bispecific molecule comprising said non-engineered form of the Fc domain) to FcRn. The Fc domain, or protease-activatable T cell activating bispecific molecules of the invention comprising said Fc domain, may exhibit greater than about 80% and even greater than about 90% of such affinity. In certain embodiments the Fc domain of the protease-activatable T cell activating bispecific molecule is engineered to have reduced effector function, as compared to a non-engineered Fc domain. The reduced effector function can include, but is not limited to, one or more of the following: reduced complement dependent cytotoxicity (CDC), reduced antibody-dependent cell-mediated cytotoxicity (ADCC), reduced antibody-dependent cellular phagocytosis (ADCP), reduced cytokine secretion, reduced immune complex-mediated antigen uptake by antigen-presenting cells, reduced binding to NK cells, reduced binding to macrophages, reduced binding to monocytes, reduced binding to polymorphonuclear cells, reduced direct signaling inducing apoptosis, reduced crosslinking of target-bound antibodies, reduced dendritic cell maturation, or reduced T cell priming. In one embodiment the reduced effector function is one or more selected from the group of reduced CDC, reduced ADCC, reduced ADCP, and reduced cytokine secretion. In a particular embodiment the reduced effector function is reduced ADCC. In one embodiment the reduced ADCC is less than 20% of the ADCC induced by a non-engineered Fc domain (or a protease-activatable T cell activating bispecific molecule comprising a non-engineered Fc domain). In one embodiment the amino acid mutation that reduces the binding affinity of the Fc domain to an Fc receptor and/or effector function is an amino acid substitution. In one embodiment the Fc domain comprises an amino acid substitution at a position selected from the group of E233, L234, L235, N297, P331 and P329. In a more specific embodiment the Fc domain comprises an amino acid substitution at a position selected from the group of L234, L235 and P329. In some embodiments the Fc domain comprises the amino acid substitutions L234A and L235A. In one such embodiment, the Fc domain is an IgG₁ Fc domain, particularly a human IgG₁ Fc domain. In one embodiment the Fc domain comprises an amino acid substitution at position P329. In a more specific embodiment the amino acid substitution is P329A or P329G, particularly P329G. In one embodiment the Fc domain comprises an amino acid substitution at position P329 and a further amino acid substitution at a position selected from E233, L234, L235, N297 and P331. In a more specific embodiment the further amino acid substitution is E233P, L234A, L235A, L235E, N297A, N297D or P331S. In particular embodiments the Fc domain comprises amino acid substitutions at positions P329, L234 and L235. In more particular embodiments the Fc domain comprises the amino acid mutations L234A, L235A and P329G ("P329G LALA"). In one such embodiment, the Fc domain is an IgG₁

Fc domain, particularly a human IgG₁ Fc domain. The “P329G LALA” combination of amino acid substitutions almost completely abolishes Fc γ receptor (as well as complement) binding of a human IgG₁ Fc domain, as described in PCT publication no. WO 2012/130831, incorporated herein by reference in its entirety. WO 2012/130831 also describes methods of preparing such mutant Fc domains and methods for determining its properties such as Fc receptor binding or effector functions. IgG₄ antibodies exhibit reduced binding affinity to Fc receptors and reduced effector functions as compared to IgG₁ antibodies. Hence, in some embodiments the Fc domain of the protease-activatable T cell activating bispecific molecules of the invention is an IgG₄ Fc domain, particularly a human IgG₄ Fc domain. In one embodiment the IgG₄ Fc domain comprises amino acid substitutions at position 5228, specifically the amino acid substitution S228P. To further reduce its binding affinity to an Fc receptor and/or its effector function, in one embodiment the IgG₄ Fc domain comprises an amino acid substitution at position L235, specifically the amino acid substitution L235E. In another embodiment, the IgG₄ Fc domain comprises an amino acid substitution at position P329, specifically the amino acid substitution P329G. In a particular embodiment, the IgG₄ Fc domain comprises amino acid substitutions at positions 5228, L235 and P329, specifically amino acid substitutions S228P, L235E and P329G. Such IgG₄ Fc domain mutants and their Fc γ receptor binding properties are described in PCT publication no. WO 2012/130831, incorporated herein by reference in its entirety.

[0345] In a particular embodiment the Fc domain exhibiting reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain, is a human IgG₁ Fc domain comprising the amino acid substitutions L234A, L235A and optionally P329G, or a human IgG₄ Fc domain comprising the amino acid substitutions S228P, L235E and optionally P329G.

[0346] In certain embodiments N-glycosylation of the Fc domain has been eliminated. In one such embodiment the Fc domain comprises an amino acid mutation at position N297, particularly an amino acid substitution replacing asparagine by alanine (N297A) or aspartic acid (N297D).

[0347] In addition to the Fc domains described hereinabove and in PCT publication no. WO 2012/130831, Fc domains with reduced Fc receptor binding and/or effector function also include those with substitution of one or more of Fc domain residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Pat. No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called “DANA” Fc mutant with substitution of residues 265 and 297 to alanine (U.S. Pat. No. 7,332,581).

[0348] Mutant Fc domains can be prepared by amino acid deletion, substitution, insertion or modification using genetic or chemical methods well known in the art. Genetic methods may include site-specific mutagenesis of the encoding DNA sequence, PCR, gene synthesis, and the like. The correct nucleotide changes can be verified for example by sequencing.

[0349] Binding to Fc receptors can be easily determined e.g., by ELISA, or by Surface Plasmon Resonance (SPR) using standard instrumentation such as a BIAcore instrument (GE Healthcare), and Fc receptors such as may be obtained by recombinant expression. A suitable such binding assay is described herein. Alternatively, binding affinity of

Fc domains or cell activating bispecific antigen binding molecules comprising an Fc domain for Fc receptors may be evaluated using cell lines known to express particular Fc receptors, such as human NK cells expressing Fc γ IIIa receptor.

[0350] Effector function of an Fc domain, or a protease-activatable T cell activating bispecific molecule comprising an Fc domain, can be measured by methods known in the art. A suitable assay for measuring ADCC is described herein. Other examples of in vitro assays to assess ADCC activity of a molecule of interest are described in U.S. Pat. No. 5,500,362; Hellstrom et al. Proc Natl Acad Sci USA 83, 7059-7063 (1986) and Hellstrom et al., Proc Natl Acad Sci USA 82, 1499-1502 (1985); U.S. Pat. No. 5,821,337; Bruggemann et al., J Exp Med 166, 1351-1361 (1987). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTITM non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, Calif.); and CytoTox 96® non-radioactive cytotoxicity assay (Promega, Madison, Wis.)). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed in vivo, e.g., in a animal model such as that disclosed in Clynes et al., Proc Natl Acad Sci USA 95, 652-656 (1998).

[0351] In some embodiments, binding of the Fc domain to a complement component, specifically to C1q, is reduced. Accordingly, in some embodiments wherein the Fc domain is engineered to have reduced effector function, said reduced effector function includes reduced CDC. C1q binding assays may be carried out to determine whether the protease-activatable T cell activating bispecific molecule is able to bind C1q and hence has CDC activity. See e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (see, for example, Gazzano-Santoro et al., J Immunol Methods 202, 163 (1996); Cragg et al., Blood 101, 1045-1052 (2003); and Cragg and Glennie, Blood 103, 2738-2743 (2004)).

Antigen Binding Moieties

[0352] The antigen binding molecule of the invention is bispecific, i.e. it comprises at least two antigen binding moieties capable of specific binding to two distinct antigenic determinants. According to the invention, the antigen binding moieties are Fab molecules (i.e. antigen binding domains composed of a heavy and a light chain, each comprising a variable and a constant region). In one embodiment said Fab molecules are human. In another embodiment said Fab molecules are humanized. In yet another embodiment said Fab molecules comprise human heavy and light chain constant regions.

[0353] At least one of the antigen binding moieties is a crossover Fab molecule. Such modification prevent mispairing of heavy and light chains from different Fab molecules, thereby improving the yield and purity of the protease-activatable T cell activating bispecific molecule of the invention in recombinant production. In a particular crossover Fab molecule useful for the protease-activatable T cell activating bispecific molecule of the invention, the constant regions of the Fab light chain and the Fab heavy chain are exchanged. In another crossover Fab molecule useful for the protease-activatable T cell activating bispecific molecule of

the invention, the variable regions of the Fab light chain and the Fab heavy chain are exchanged.

[0354] In a particular embodiment according to the invention, the protease-activatable T cell activating bispecific molecule is capable of simultaneous binding to a target cell antigen, particularly a tumor cell antigen, and CD3. In one embodiment, the protease-activatable T cell activating bispecific molecule is capable of crosslinking a T cell and a target cell by simultaneous binding to a target cell antigen and CD3. In an even more particular embodiment, such simultaneous binding results in lysis of the target cell, particularly a tumor cell. In one embodiment, such simultaneous binding results in activation of the T cell. In other embodiments, such simultaneous binding results in a cellular response of a T lymphocyte, particularly a cytotoxic T lymphocyte, selected from the group of: proliferation, differentiation, cytokine secretion, cytotoxic effector molecule release, cytotoxic activity, and expression of activation markers. In one embodiment, binding of the protease-activatable T cell activating bispecific molecule to CD3 without simultaneous binding to the target cell antigen does not result in T cell activation.

[0355] In one embodiment, the protease-activatable T cell activating bispecific molecule is capable of re-directing cytotoxic activity of a T cell to a target cell. In a particular embodiment, said re-direction is independent of MHC-mediated peptide antigen presentation by the target cell and and/or specificity of the T cell.

[0356] Particularly, a T cell according to any of the embodiments of the invention is a cytotoxic T cell. In some embodiments the T cell is a CD4⁺ or a CD8⁺ T cell, particularly a CD8⁺ T cell.

CD3 Binding Moiety

[0357] The protease-activatable T cell activating bispecific molecule of the invention comprises at least one antigen binding moiety capable of binding to CD3 (also referred to herein as an “CD3 antigen binding moiety” or “first antigen binding moiety”). In a particular embodiment, the protease-activatable T cell activating bispecific molecule comprises not more than one antigen binding moiety capable of specific binding to CD3. In one embodiment the protease-activatable T cell activating bispecific molecule provides monovalent binding to CD3. The CD3 antigen binding is a crossover Fab molecule, i.e. a Fab molecule wherein either the variable or the constant regions of the Fab heavy and light chains are exchanged. In embodiments where there is more than one antigen binding moiety capable of specific binding to a target cell antigen comprised in the protease-activatable T cell activating bispecific molecule, the antigen binding moiety capable of specific binding to CD3 preferably is a crossover Fab molecule and the antigen binding moieties capable of specific binding to a target cell antigen are conventional Fab molecules.

[0358] In a particular embodiment CD3 is human CD3 or cynomolgus CD3, most particularly human CD3. In a particular embodiment the CD3 antigen binding moiety is cross-reactive for (i.e. specifically binds to) human and cynomolgus CD3. In some embodiments, the first antigen binding moiety is capable of specific binding to the epsilon subunit of CD3.

[0359] The CD3 antigen binding moiety comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO:

11, SEQ ID NO: 12 and SEQ ID NO: 13 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19.

[0360] In one embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 11, the heavy chain CDR2 of SEQ ID NO: 12, the heavy chain CDR3 of SEQ ID NO: 13, the light chain CDR1 of SEQ ID NO: 17, the light chain CDR2 of SEQ ID NO: 18, and the light chain CDR3 of SEQ ID NO: 19.

[0361] In one embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 44, the heavy chain CDR2 of SEQ ID NO: 45, the heavy chain CDR3 of SEQ ID NO: 46, the light chain CDR1 of SEQ ID NO: 17, the light chain CDR2 of SEQ ID NO: 18, and the light chain CDR3 of SEQ ID NO: 19.

[0362] In one embodiment the CD3 antigen binding moiety comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43, and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0363] In one embodiment the CD3 antigen binding moiety comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.

[0364] In one embodiment the CD3 antigen binding moiety comprises the heavy chain variable region sequence of SEQ ID NO: 43 and the light chain variable region sequence of SEQ ID NO: 55.

Target Cell Antigen Binding Moiety

[0365] The protease-activatable T cell activating bispecific molecule of the invention comprises at least one antigen binding moiety capable of binding to a target cell antigen (also referred to herein as a “target cell antigen binding moiety” or “second” or “third” antigen binding moiety). In certain embodiments, the protease-activatable T cell activating bispecific molecule comprises two antigen binding moieties capable of binding to a target cell antigen. In a particular such embodiment, each of these antigen binding moieties specifically binds to the same antigenic determinant. In an even more particular embodiment, all of these antigen binding moieties are identical. In one embodiment, the protease-activatable T cell activating bispecific molecule comprises an immunoglobulin molecule capable of specific binding to a target cell antigen. In one embodiment the protease-activatable T cell activating bispecific molecule comprises not more than two antigen binding moieties capable of binding to a target cell antigen.

[0366] In a preferred embodiment, the target cell antigen binding moiety is a Fab molecule, particularly a conventional Fab molecule that binds to a specific antigenic determinant and is able to direct the Protease-activatable T cell activating bispecific molecule to a target site, for example to a specific type of tumor cell that bears the antigenic determinant.

[0367] In certain embodiments the target cell antigen binding moiety specifically binds to a cell surface antigen. In a particular embodiment the target cell antigen binding moiety specifically binds to a Folate Receptor 1 (FolR1) on the surface of a target cell. In another specific such embodiment the target cell antigen binding moiety specifically

binds to an epidermal growth factor receptor (EGFR), specifically, a human EGFR, e.g., HER1. In another specific such embodiment the target cell antigen binding moiety specifically binds to HER2. In another specific such embodiment the target cell antigen binding moiety specifically binds to Mesothelin, specifically, to human Mesothelin.

[0368] In certain embodiments the target cell antigen binding moiety is directed to an antigen associated with a pathological condition, such as an antigen presented on a tumor cell or on a virus-infected cell. Suitable antigens are cell surface antigens, for example, but not limited to, cell surface receptors. In particular embodiments the antigen is a human antigen. In a specific embodiment the target cell antigen is selected from Folate Receptor 1 (FolR1) and epidermal growth factor receptor (EGFR), specifically, a human EGFR, e.g., HER1. In a further specific embodiment the target cell antigen is HER2. In a further specific embodiment the target cell antigen is Mesothelin.

[0369] In some embodiments the protease-activatable T cell activating bispecific molecule comprises at least one antigen binding moiety that is specific for HER1. In one embodiment, the antigen binding moiety that is specific for HER1 comprises at least one heavy chain complementarity determining region (CDR) of selected from the group consisting of SEQ ID NO: 56, SEQ ID NO: 57 and SEQ ID NO: 58 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60, and SEQ ID NO: 61.

[0370] In one embodiment, the antigen binding moiety that is specific for HER1 comprises the heavy chain CDR1 of SEQ ID NO: 56, the heavy chain CDR2 of SEQ ID NO: 57, the heavy chain CDR3 of SEQ ID NO: 58, the light chain CDR1 of SEQ ID NO: 59, the light chain CDR2 of SEQ ID NO: 60, and the light chain CDR3 of SEQ ID NO: 61.

[0371] In one embodiment, the antigen binding moiety that is specific for HER1 comprises the heavy chain and light chain CDR sequences of an anti-HER1 antibody disclosed in PCT Application Publication Number WO2006/082515.

[0372] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 comprises at least one of a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 32, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 33, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 34. In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 comprises the polypeptide sequence of SEQ ID NO: 32, the polypeptide sequence of SEQ ID NO: 33, and the polypeptide sequence of SEQ ID NO: 34.

[0373] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1

of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25.

[0374] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31.

[0375] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 further comprises an anti-idiotypic CD3 scFv comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 41 or 42. In one embodiment, the anti-idiotypic scFv comprises the polypeptide sequence of SEQ ID NO: 41 or 42.

[0376] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 further comprises an anti-idiotypic HER1 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 48, the heavy chain CDR2 of SEQ ID NO: 49, the heavy chain CDR3 of SEQ ID NO: 50, the light chain CDR1 of SEQ ID NO: 51, the light chain CDR2 of SEQ ID NO: 52, and the light chain CDR3 of SEQ ID NO: 53. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 48, the heavy chain CDR2 of SEQ ID NO: 49, the heavy chain CDR3 of SEQ ID NO: 50, the light chain CDR1 of SEQ ID NO: 51, the light chain CDR2 of SEQ ID NO: 52, and the light chain CDR3 of SEQ ID NO: 53.

[0377] In one embodiment the protease-activatable T cell activating bispecific molecule that comprises at least one antigen binding moiety that is specific for HER1 further comprises a linker comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 35.

[0378] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 further comprises a linker having a protease recognition site comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 97. In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER1 further comprises a linker comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment,

the linker comprises the polypeptide sequence of SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 86.

[0379] In some embodiments the protease-activatable T cell activating bispecific molecule comprises at least one antigen binding moiety that is specific for HER2. In one embodiment, the antigen binding moiety that is specific for HER2 comprises at least one heavy chain complementarity determining region (CDR) of selected from the group consisting of SEQ ID NO: 142, SEQ ID NO: 143 and SEQ ID NO: 144 and at least one light chain CDR selected from the group of SEQ ID NO: 148, SEQ ID NO: 149, and SEQ ID NO: 150. In a further one embodiment, the antigen binding moiety that is specific for HER2 comprises at least one heavy chain complementarity determining region (CDR) of selected from the group consisting of SEQ ID NO: 145, SEQ ID NO: 146 and SEQ ID NO: 147 and at least one light chain CDR selected from the group of SEQ ID NO: 148, SEQ ID NO: 149, and SEQ ID NO: 150.

[0380] In one embodiment, the antigen binding moiety that is specific for HER2 comprises the heavy chain CDR1 of SEQ ID NO: 142, the heavy chain CDR2 of SEQ ID NO: 143, the heavy chain CDR3 of SEQ ID NO: 144, the light chain CDR1 of SEQ ID NO: 148, the light chain CDR2 of SEQ ID NO: 149, and the light chain CDR3 of SEQ ID NO: 150. In a further embodiment, the antigen binding moiety that is specific for HER2 comprises the heavy chain CDR1 of SEQ ID NO: 145, the heavy chain CDR2 of SEQ ID NO: 146, the heavy chain CDR3 of SEQ ID NO: 147, the light chain CDR1 of SEQ ID NO: 148, the light chain CDR2 of SEQ ID NO: 149, and the light chain CDR3 of SEQ ID NO: 150.

[0381] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER2 further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25.

[0382] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER2 further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31.

[0383] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one anti-

gen binding moiety that is specific for HER2 further comprises an anti-idiotypic CD3 scFv comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 41 or 42. In one embodiment, the anti-idiotypic scFv comprises the polypeptide sequence of SEQ ID NO: 41 or 42.

[0384] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER2 further comprises an anti-idiotypic HER2 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 48, the heavy chain CDR2 of SEQ ID NO: 49, the heavy chain CDR3 of SEQ ID NO: 50, the light chain CDR1 of SEQ ID NO: 51, the light chain CDR2 of SEQ ID NO: 52, and the light chain CDR3 of SEQ ID NO: 53. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 48, the heavy chain CDR2 of SEQ ID NO: 49, the heavy chain CDR3 of SEQ ID NO: 50, the light chain CDR1 of SEQ ID NO: 51, the light chain CDR2 of SEQ ID NO: 52, and the light chain CDR3 of SEQ ID NO: 53.

[0385] In one embodiment the protease-activatable T cell activating bispecific molecule that comprises at least one antigen binding moiety that is specific for HER2 further comprises a linker comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 35.

[0386] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER2 further comprises a linker having a protease recognition site comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 97. In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for HER2 further comprises a linker comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 86.

[0387] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 132, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 136, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 81 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 133.

[0388] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 132, the polypeptide sequence of

SEQ ID NO: 136, the polypeptide sequence of SEQ ID NO: 81 and the polypeptide sequence of SEQ ID NO: 133.

[0389] In particular embodiments the protease-activatable T cell activating bispecific molecule comprises at least one antigen binding moiety that is specific for FolR1. In one embodiment the FolR1 is a human FolR1. In one embodiment, the protease-activatable T cell activating bispecific molecule comprises at least one antigen binding moiety that is specific for human FolR1 and does not bind to human FolR2 or human FolR3. In one embodiment, the antigen binding moiety that is specific for FolR1 comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0390] In one embodiment, the antigen binding moiety that is specific for FolR1 comprises the heavy chain CDR1 of SEQ ID NO: 14, the heavy chain CDR2 of SEQ ID NO: 15, the heavy chain CDR3 of SEQ ID NO: 16, the light chain CDR1 of SEQ ID NO: 17, the light chain CDR2 of SEQ ID NO: 18, and the light chain CDR3 of SEQ ID NO: 19.

[0391] In a further embodiment, the antigen binding moiety that is specific for FolR1 comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 47 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 55, or variants thereof that retain functionality.

[0392] In one embodiment, the antigen binding moiety that is specific for FolR1 comprises the heavy chain variable region comprising an amino acid sequence of SEQ ID NO: 47 and the light chain variable region comprising an amino acid sequence of SEQ ID NO: 55.

[0393] In one embodiment, the antigen binding moiety that is specific for FolR1 comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 151, SEQ ID NO: 152 and SEQ ID NO: 153 and at least one light chain CDR selected from the group of SEQ ID NO: 154, SEQ ID NO: 155 and SEQ ID NO: 156.

[0394] In one embodiment, the antigen binding moiety that is specific for FolR1 comprises the heavy chain CDR1 of SEQ ID NO: 151, the heavy chain CDR2 of SEQ ID NO: 152, the heavy chain CDR3 of SEQ ID NO: 153, the light chain CDR1 of SEQ ID NO: 154, the light chain CDR2 of SEQ ID NO: 155, and the light chain CDR3 of SEQ ID NO: 156.

[0395] In a further embodiment, the antigen binding moiety that is specific for FolR1 comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 157 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 158, or variants thereof that retain functionality.

[0396] In one embodiment, the antigen binding moiety that is specific for FolR1 comprises the heavy chain variable region comprising an amino acid sequence of SEQ ID NO: 157 and the light chain variable region comprising an amino acid sequence of SEQ ID NO: 158.

[0397] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 2, and a polypeptide

sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 1.

[0398] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 2, and the polypeptide sequence of SEQ ID NO: 1.

[0399] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for FolR1 further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25.

[0400] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for FolR1 further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31.

[0401] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for FolR1 further comprises an anti-idiotypic CD3 scFv comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 41 or 42. In one embodiment, the anti-idiotypic scFv comprises the polypeptide sequence of SEQ ID NO: 41 or 42.

[0402] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for FolR1 further comprises a linker having a protease recognition site comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 97. In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for FolR1 further comprises a linker comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96.

In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 86.

[0403] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 1, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 3 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 72.

[0404] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 1, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 3 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 72.

[0405] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 1, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 3 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 85.

[0406] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 1, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 3, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 73 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 74.

[0407] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 1, the polypeptide sequence of SEQ ID NO: 3 and the polypeptide sequence of SEQ ID NO: 72.

[0408] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 1, the polypeptide sequence of SEQ ID NO: 3 and the polypeptide sequence of SEQ ID NO: 85.

[0409] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 1, the polypeptide sequence of SEQ ID NO: 3, the polypeptide sequence of SEQ ID NO: 73 and the polypeptide sequence of SEQ ID NO: 74.

[0410] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 137, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 139, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 81 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 138.

[0411] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide

sequence of SEQ ID NO: 137, the polypeptide sequence of SEQ ID NO: 139, the polypeptide sequence of SEQ ID NO: 81 and the polypeptide sequence of SEQ ID NO: 138.

[0412] In particular embodiments the protease-activatable T cell activating bispecific molecule comprises at least one antigen binding moiety that is specific for Mesothelin. In one embodiment the Mesothelin is human Mesothelin. In one embodiment, the protease-activatable T cell activating bispecific molecule comprises at least one antigen binding moiety that is specific for human Mesothelin. In one embodiment, the antigen binding moiety that is specific for Mesothelin comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 107, SEQ ID NO: 108 and SEQ ID NO: 109 and at least one light chain CDR selected from the group of SEQ ID NO: 110, SEQ ID NO: 111 and SEQ ID NO: 112.

[0413] In one embodiment, the antigen binding moiety that is specific for Mesothelin comprises the heavy chain CDR1 of SEQ ID NO: 107, the heavy chain CDR2 of SEQ ID NO: 108, the heavy chain CDR3 of SEQ ID NO: 109, the light chain CDR1 of SEQ ID NO: 110, the light chain CDR2 of SEQ ID NO: 111, and the light chain CDR3 of SEQ ID NO: 112.

[0414] In a further embodiment, the antigen binding moiety that is specific for Mesothelin comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 113 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 114, or variants thereof that retain functionality.

[0415] In one embodiment, the antigen binding moiety that is specific for Mesothelin comprises the heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 113 and the light chain variable region comprising the amino acid sequence of SEQ ID NO: 114.

[0416] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for Mesothelin further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25.

[0417] In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25.

[0418] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for Mesothelin further comprises an anti-idiotypic CD3 scFv comprising at least one of the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1

of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31.

[0419] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for Mesothelin further comprises an anti-idiotypic CD3 scFv comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 41 or 42. In one embodiment, the anti-idiotypic scFv comprises the polypeptide sequence of SEQ ID NO: 41 or 42.

[0420] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for Mesothelin further comprises a linker having a protease recognition site comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 97.

[0421] In one embodiment the protease-activatable T cell activating bispecific molecule comprising at least one antigen binding moiety that is specific for Mesothelin further comprises a linker comprising a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 86.

[0422] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 77, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 78, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 81 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 82.

[0423] In one embodiment the protease-activatable T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 76, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 77, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 78 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 79.

[0424] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 77, the polypeptide sequence of SEQ ID NO: 78, the polypeptide sequence of SEQ ID NO: 81 and the polypeptide sequence of SEQ ID NO: 82.

[0425] In one embodiment the protease-activatable T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 76, the polypeptide sequence of

SEQ ID NO: 77, the polypeptide sequence of SEQ ID NO: 78 and the polypeptide sequence of SEQ ID NO: 79.

[0426] In one embodiment, provided is a T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 76, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 77, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 78 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 81.

[0427] In one embodiment the T cell activating bispecific molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 77, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 78, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 81 and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 84.

[0428] In one embodiment the T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 76, the polypeptide sequence of SEQ ID NO: 77, the polypeptide sequence of SEQ ID NO: 78 and the polypeptide sequence of SEQ ID NO: 81.

[0429] In one embodiment the T cell activating bispecific molecule comprises the polypeptide sequence of SEQ ID NO: 77, the polypeptide sequence of SEQ ID NO: 78, the polypeptide sequence of SEQ ID NO: 81 and the polypeptide sequence of SEQ ID NO: 84.

Masking Moiety

[0430] The protease-activatable T cell activating bispecific molecule of the invention comprises at least one masking moiety. Others have tried to mask binding of an antibody by capping the binding moiety with a fragment of the antigen recognized by the binding moiety (e.g., WO2013128194). This approach has several limitations. For example, using the antigen allows for less flexibility in reducing the affinity of the binding moiety. This is so because the affinity has to be high enough to be reliably masked by the antigen mask. Also, dissociated antigen could potentially bind to and interact with its cognate receptor(s) in vivo and cause undesirable signals to the cell expressing such receptor. In contrast, the approach described herein uses an anti-idiotypic antibody or fragment thereof as a mask. Two countervailing considerations for designing an effective masking moiety are 1. effectiveness of the masking and 2. reversibility of the masking. If the affinity is too low, masking would be inefficient. However, if the affinity is too high, the masking process might not be readily reversible. It was not predictable whether a high affinity anti-idiotypic mask or a low affinity anti-idiotypic mask would work better. As described herein, higher affinity masking moieties performed overall better in masking the antigen binding side and, at the same time, could be effectively removed for activation of the molecule. In one embodiment, the anti-idiotypic mask has a KD of 1-8 nM. In one embodiment, anti-idiotypic mask has a KD of 2 nM at 37° C. In one specific embodiment, the masking moiety recognizes the idio type of the first antigen binding moiety capable of specific binding to a CD3, e.g., a human CD3. In one specific

embodiment, the masking moiety recognizes the idiotype of the second antigen binding moiety capable of binding to a target cell antigen.

[0431] In one embodiment, the masking moiety masks a CD3-binding moiety and comprises at least one of the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25. In one embodiment, the masking moiety comprises the heavy chain CDR1 of SEQ ID NO: 20, the heavy chain CDR2 of SEQ ID NO: 21, the heavy chain CDR3 of SEQ ID NO: 22, the light chain CDR1 of SEQ ID NO: 23, the light chain CDR2 of SEQ ID NO: 24, and the light chain CDR3 of SEQ ID NO: 25.

[0432] In one embodiment, the masking moiety masks a CD3-binding moiety and comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 41. In one embodiment, the masking moiety masks a CD3-binding moiety and comprises the polypeptide sequence of SEQ ID NO: 41.

[0433] In one preferred embodiment, the masking moiety masks a CD3-binding moiety and comprises at least one of the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31. In one embodiment, the masking moiety comprises the heavy chain CDR1 of SEQ ID NO: 26, the heavy chain CDR2 of SEQ ID NO: 27, the heavy chain CDR3 of SEQ ID NO: 28, the light chain CDR1 of SEQ ID NO: 29, the light chain CDR2 of SEQ ID NO: 30, and the light chain CDR3 of SEQ ID NO: 31. In one embodiment, the masking moiety masks a CD3-binding moiety and comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 42. In one preferred embodiment, the masking moiety masks a CD3-binding moiety and comprises the polypeptide sequence of SEQ ID NO: 42.

[0434] In one embodiment, the masking moiety masks a HER1-binding moiety and comprises at least one of the heavy chain CDR1 of SEQ ID NO: 48, the heavy chain CDR2 of SEQ ID NO: 49, the heavy chain CDR3 of SEQ ID NO: 50, the light chain CDR1 of SEQ ID NO: 51, the light chain CDR2 of SEQ ID NO: 52, and the light chain CDR3 of SEQ ID NO: 53. In one embodiment, the anti-idiotypic scFv comprises the heavy chain CDR1 of SEQ ID NO: 48, the heavy chain CDR2 of SEQ ID NO: 49, the heavy chain CDR3 of SEQ ID NO: 50, the light chain CDR1 of SEQ ID NO: 51, the light chain CDR2 of SEQ ID NO: 52, and the light chain CDR3 of SEQ ID NO: 53.

[0435] In one aspect, the invention relates to an idiotype-specific polypeptide for reversibly concealing antigen binding of an antigen-binding of a molecule. In one embodiment, the invention relates to an idiotype-specific polypeptide for reversibly concealing an anti-CD3 antigen binding site of a molecule. Such idiotype-specific polypeptide for reversibly concealing an anti-CD3 antigen binding site must be capable of specific binding to the anti-CD3 antigen binding site's idiotype and thereby reducing or abrogating binding of the anti-CD3 antigen binding site to CD3. In one embodiment, the invention relates to an idiotype-specific polypeptide for reversibly concealing an anti-HER1 antigen binding site of a molecule. In one embodiment the idiotype-specific poly-

peptide is an anti-idiotypic scFv. In one embodiment the idiotype-specific polypeptide is covalently attached to the molecule through a linker. In one embodiment the idiotype-specific polypeptide is covalently attached to the molecule through more than one linker. In one embodiment the idiotype-specific polypeptide is covalently attached to the molecule through two linkers. In one embodiment the linker is a peptide linker. In one embodiment the linker is a protease-cleavable linker. In one embodiment, the linker comprises the sequence of SEQ ID NO: 7, 8, 9, or 10. In one embodiment, the linker comprises the sequence of SEQ ID NO: 7, 8, 9, 10, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95 or 96. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 7. In one embodiment, the linker comprises the polypeptide sequence of SEQ ID NO: 86. In one embodiment the peptide linker comprises at least one protease recognition site. In one embodiment, the protease recognition site comprises the polypeptide sequence of SEQ ID NO: 36, 37, 38, 39, 40, 97, 98, 99, 100, 101, 102, 103, 104, 105 or 106. In one preferred embodiment, the protease recognition site comprises the protease recognition sequence RQARVVNG (SEQ ID NO: 36). In further embodiment, the linker comprises more than one protease recognition site. In one preferred embodiment, the protease recognition site comprises the protease recognition sequence VHM-PLGFLGPRQARVVNG (SEQ ID NO: 97). In one embodiment the protease is selected from the group consisting of metalloproteinase, e.g., matrix metalloproteinase (MMP) 1-28 and A Disintegrin And Metalloproteinase (ADAM) 2, 7-12, 15, 17-23, 28-30 and 33, serine protease, e.g., urokinase-type plasminogen activator and Matriptase, cysteine protease, aspartic protease, and cathepsin protease. In one specific embodiment the protease is MMP9 or MMP2. In a further specific embodiment, the protease is Matriptase.

[0436] In one embodiment the molecule which comprises the anti-CD3 antigen binding site is a T-cell activating bispecific molecule. In one particular embodiment the idiotype-specific polypeptide comprises a heavy chain variable region comprising at least one of a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO: 20); CDR H2 amino acid sequence of WINTETGEPAYADDFKG (SEQ ID NO: 21); and a CDR H3 amino acid sequence of PYDYDVLDY (SEQ ID NO: 22). In one particular embodiment the idiotype-specific polypeptide comprises a light chain variable region comprising at least one of: a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO: 23); a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO: 24); and a CDR L3 amino acid sequence of QHSREFPWT (SEQ ID NO: 25). In one particular embodiment the idiotype-specific polypeptide comprises: a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO: 20); a CDR H2 amino acid sequence of WINTETGEPAYADDFKG (SEQ ID NO: 21); a CDR H3 amino acid sequence of PYDYDVLDY (SEQ ID NO: 22); and a light chain variable region comprising: a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO: 23); a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO: 24); and a CDR L3 amino acid sequence of QHSREFPWT (SEQ ID NO: 25). In one particular embodiment the idiotype-specific polypeptide comprises a heavy chain variable region comprising at least one of: a heavy chain complementarity determining region (CDR H) 1 amino acid

sequence of SYGVS (SEQ ID NO:26); a CDR H2 amino acid sequence of IIWGDGSTNYHSALIS (SEQ ID NO:27); and a CDR H3 amino acid sequence of GITTVDDYYAMDY (SEQ ID NO:28). In one particular embodiment the idiotype-specific polypeptide comprises a light chain variable region comprising at least one of: a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29); a CDR L2 amino acid sequence of AATFLAD (SEQ ID NO:30); and a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31). In one particular embodiment the idiotype-specific polypeptide comprises a heavy chain variable region comprising: a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26); a CDR H2 amino acid sequence of IIWGDGSTNYHSALIS (SEQ ID NO:27); a CDR H3 amino acid sequence of GITTVDDYYAMDY (SEQ ID NO:28); and a light chain variable region comprising: a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29); a CDR L2 amino acid sequence of AATFLAD (SEQ ID NO:30); and a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31). In one embodiment, the idiotype-specific polypeptide comprises a heavy chain variable region comprising at least one of: a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SEQ ID NO:48; a CDR H2 amino acid sequence of SEQ ID NO:49; and a CDR H3 amino acid sequence of SEQ ID NO:50. In one embodiment, the idiotype-specific polypeptide comprises a light chain variable region comprising at least one of: a light chain complementarity determining region (CDR L) 1 amino acid sequence of SEQ ID NO:51; a CDR L2 amino acid sequence of SEQ ID NO:52; and a CDR L3 amino acid sequence of SEQ ID NO:53. In one embodiment, the idiotype-specific polypeptide comprises a heavy chain variable region comprising a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SEQ ID NO:48; a CDR H2 amino acid sequence of SEQ ID NO:49; and a CDR H3 amino acid sequence of SEQ ID NO:50, and a light chain variable region comprising a light chain complementarity determining region (CDR L) 1 amino acid sequence of SEQ ID NO:51; a CDR L2 amino acid sequence of SEQ ID NO:52; and a CDR L3 amino acid sequence of SEQ ID NO:53.

Polynucleotides

[0437] The invention further provides isolated polynucleotides encoding a protease-activatable T cell activating bispecific molecule as described herein or a fragment thereof. In some embodiments, said fragment is an antigen binding fragment.

[0438] Polynucleotides of the invention include those that are at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequences set forth in SEQ ID NOs 62-71 or SEQ ID NOs including functional fragments or variants thereof. Polynucleotides of the invention further include those that are at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequences set forth in SEQ ID NOs 117-131 including functional fragments or variants thereof. Polynucleotides of the invention further include those that are at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the sequences set forth in SEQ ID NOs 162-170 including functional fragments or variants thereof.

[0439] The polynucleotides encoding protease-activatable T cell activating bispecific molecules of the invention may be expressed as a single polynucleotide that encodes the entire protease-activatable T cell activating bispecific molecule or as multiple (e.g., two or more) polynucleotides that are co-expressed. Polypeptides encoded by polynucleotides that are co-expressed may associate through, e.g., disulfide bonds or other means to form a functional protease-activatable T cell activating bispecific molecule. For example, the light chain portion of an antigen binding moiety may be encoded by a separate polynucleotide from the portion of the protease-activatable T cell activating bispecific molecule comprising the heavy chain portion of the antigen binding moiety, an Fc domain subunit and optionally (part of) another antigen binding moiety. When co-expressed, the heavy chain polypeptides will associate with the light chain polypeptides to form the antigen binding moiety. In another example, the portion of the protease-activatable T cell activating bispecific molecule comprising one of the two Fc domain subunits and optionally (part of) one or more antigen binding moieties could be encoded by a separate polynucleotide from the portion of the protease-activatable T cell activating bispecific molecule comprising the the other of the two Fc domain subunits and optionally (part of) an antigen binding moiety. When co-expressed, the Fc domain subunits will associate to form the Fc domain.

[0440] In some embodiments, the isolated polynucleotide encodes the entire protease-activatable T cell activating bispecific molecule according to the invention as described herein. In other embodiments, the isolated polynucleotide encodes a polypeptides comprised in the protease-activatable T cell activating bispecific molecule according to the invention as described herein.

[0441] In another embodiment, the present invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes a variable region sequence. In another embodiment, the present invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule or fragment thereof, wherein the polynucleotide comprises a sequence that encodes a polypeptide sequence as shown in SEQ ID NOs 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 47, or 55. In another embodiment, the present invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule or fragment thereof, wherein the polynucleotide comprises a sequence that encodes a polypeptide sequence as shown in SEQ ID NOs 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 47, 55, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85. In another embodiment, the present invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule or fragment thereof, wherein the polynucleotide comprises a sequence that encodes a polypeptide sequence as shown in SEQ ID NOs 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 47, 55, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 132, 133, 134, 135, 136, 137, 138, 139, 140 or 141. In another embodiment, the invention is further directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence

that is at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleotide sequence shown in SEQ ID NOs 62, 63, 64, 65, 66, 67, 68, 69, 70, or 71. In another embodiment, the invention is further directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that is at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleotide sequence shown in SEQ ID NOs 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130 or 131. In another embodiment, the invention is further directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that is at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleotide sequence shown in SEQ ID NOs 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 162, 163, 164, 165, 166, 167, 168, 169 or 170. In another embodiment, the invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises the nucleic acid sequence shown in SEQ ID NOs 62, 63, 64, 65, 66, 67, 68, 69, 70, or 71. In another embodiment, the invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises the nucleic acid sequence shown in SEQ ID NOs 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130 or 131. In another embodiment, the invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises the nucleic acid sequence shown in SEQ ID NOs 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 162, 163, 164, 165, 166, 167, 168, 169 or 170. In another embodiment, the invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes a variable region sequence that is at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence in SEQ ID NOs 43, 47, or 55. The invention encompasses an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes the variable region sequence of SEQ ID NOs SEQ ID NOs 43, 47, or 55 with conservative amino acid substitutions.

[0442] In another embodiment, the invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes a variable region sequence that is at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence in SEQ ID NOs 43, 47, 55, 113, 114, 115 or 116. The invention encompasses an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a

sequence that encodes the variable region sequence of SEQ ID NOs SEQ ID NOs 43, 47, 55, 113, 114, 115 or 116 with conservative amino acid substitutions.

[0443] In another embodiment, the invention is directed to an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes a variable region sequence that is at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence in SEQ ID NOs 43, 47, 55, 113, 114, 115, 116, 157, 158, 159, 160 or 161. The invention encompasses an isolated polynucleotide encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes the variable region sequence of SEQ ID NOs SEQ ID NOs 43, 47, 55, 113, 114, 115, 116, 157, 158, 159, 160 or 161 with conservative amino acid substitutions.

[0444] In certain embodiments the polynucleotide or nucleic acid is DNA. In other embodiments, a polynucleotide of the present invention is RNA, for example, in the form of messenger RNA (mRNA). RNA of the present invention may be single stranded or double stranded.

[0445] The invention further provides isolated polynucleotides encoding an idio-type-specific polypeptide as described herein or a fragment thereof. In some embodiments, said fragment is an idio-type binding, i.e., anti-idio-type specific antibody or fragment thereof. In one embodiment the idio-type-specific polypeptide is an anti-idiotypic scFv.

[0446] The invention also encompasses an isolated polynucleotide encoding an idio-type-specific polypeptide of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes the polypeptide sequence of one or more of SEQ ID NOs 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 48, 49, 50, 51, 52, and 53. The invention also encompasses an isolated polynucleotide encoding an idio-type-specific polypeptide of the invention or a fragment thereof, wherein the polynucleotide comprises a sequence that encodes the polypeptide sequence of one or more of SEQ ID NOs 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 48, 49, 50, 51, 52, and 53 with conservative amino acid substitutions.

[0447] The polynucleotides encoding idio-type-specific polypeptides of the invention may be expressed as a single polynucleotide that encodes the entire idio-type-specific polypeptide or as multiple (e.g., two or more) polynucleotides that are co-expressed. Polypeptides encoded by polynucleotides that are co-expressed may associate through, e.g., disulfide bonds or other means to form a functional idio-type-specific polypeptide, e.g., a masking moiety. For example, in one embodiment the idio-type-specific polypeptide is an anti-idiotypic scFv (single chain variable fragment) wherein the light chain variable portion of the anti-idiotypic scFv may be encoded by a separate polynucleotide from the portion of the anti-idiotypic scFv comprising the heavy chain variable portion of the anti-idiotypic scFv. When co-expressed, the heavy chain polypeptides will associate with the light chain polypeptides to form the anti-idiotypic scFv. In some embodiments, the isolated polynucleotide encodes the idio-type-specific polypeptide according to the invention as described herein. In certain embodiments the polynucleotide or nucleic acid is DNA. In other embodiments, a polynucleotide of the present inven-

tion is RNA, for example, in the form of messenger RNA (mRNA). RNA of the present invention may be single stranded or double stranded.

Recombinant Methods

[0448] protease-activatable T cell activating bispecific molecules of the invention may be obtained, for example, by solid-state peptide synthesis (e.g., Merrifield solid phase synthesis) or recombinant production. For recombinant production one or more polynucleotide encoding the protease-activatable T cell activating bispecific molecule (fragment), e.g., as described above, is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such polynucleotide may be readily isolated and sequenced using conventional procedures. In one embodiment a vector, preferably an expression vector, comprising one or more of the polynucleotides of the invention is provided. Methods which are well known to those skilled in the art can be used to construct expression vectors containing the coding sequence of a protease-activatable T cell activating bispecific molecule (fragment) along with appropriate transcriptional/translational control signals. These methods include in vitro recombinant DNA techniques, synthetic techniques and in vivo recombination/genetic recombination. See, for example, the techniques described in Maniatis et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, Cold Spring Harbor Laboratory, N.Y. (1989); and Ausubel et al., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, Greene Publishing Associates and Wiley Interscience, N.Y. (1989). The expression vector can be part of a plasmid, virus, or may be a nucleic acid fragment. The expression vector includes an expression cassette into which the polynucleotide encoding the protease-activatable T cell activating bispecific molecule (fragment) (i.e. the coding region) is cloned in operable association with a promoter and/or other transcription or translation control elements. As used herein, a “coding region” is a portion of nucleic acid which consists of codons translated into amino acids. Although a “stop codon” (TAG, TGA, or TAA) is not translated into an amino acid, it may be considered to be part of a coding region, if present, but any flanking sequences, for example promoters, ribosome binding sites, transcriptional terminators, introns, 5' and 3' untranslated regions, and the like, are not part of a coding region. Two or more coding regions can be present in a single polynucleotide construct, e.g., on a single vector, or in separate polynucleotide constructs, e.g., on separate (different) vectors. Furthermore, any vector may contain a single coding region, or may comprise two or more coding regions, e.g., a vector of the present invention may encode one or more polypeptides, which are post- or co-translationally separated into the final proteins via proteolytic cleavage. In addition, a vector, polynucleotide, or nucleic acid of the invention may encode heterologous coding regions, either fused or unfused to a polynucleotide encoding the protease-activatable T cell activating bispecific molecule (fragment) of the invention, or variant or derivative thereof. Heterologous coding regions include without limitation specialized elements or motifs, such as a secretory signal peptide or a heterologous functional domain. An operable association is when a coding region for a gene product, e.g., a polypeptide, is associated with one or more regulatory sequences in such a way as to place expression of the gene product under the influence or control of the regulatory sequence(s). Two DNA fragments (such as a polypeptide coding region and a

promoter associated therewith) are “operably associated” if induction of promoter function results in the transcription of mRNA encoding the desired gene product and if the nature of the linkage between the two DNA fragments does not interfere with the ability of the expression regulatory sequences to direct the expression of the gene product or interfere with the ability of the DNA template to be transcribed. Thus, a promoter region would be operably associated with a nucleic acid encoding a polypeptide if the promoter was capable of effecting transcription of that nucleic acid. The promoter may be a cell-specific promoter that directs substantial transcription of the DNA only in predetermined cells. Other transcription control elements, besides a promoter, for example enhancers, operators, repressors, and transcription termination signals, can be operably associated with the polynucleotide to direct cell-specific transcription. Suitable promoters and other transcription control regions are disclosed herein. A variety of transcription control regions are known to those skilled in the art. These include, without limitation, transcription control regions, which function in vertebrate cells, such as, but not limited to, promoter and enhancer segments from cytomegaloviruses (e.g., the immediate early promoter, in conjunction with intron-A), simian virus 40 (e.g., the early promoter), and retroviruses (such as, e.g., Rous sarcoma virus). Other transcription control regions include those derived from vertebrate genes such as actin, heat shock protein, bovine growth hormone and rabbit α -globin, as well as other sequences capable of controlling gene expression in eukaryotic cells. Additional suitable transcription control regions include tissue-specific promoters and enhancers as well as inducible promoters (e.g., promoters inducible tetracyclins). Similarly, a variety of translation control elements are known to those of ordinary skill in the art. These include, but are not limited to ribosome binding sites, translation initiation and termination codons, and elements derived from viral systems (particularly an internal ribosome entry site, or IRES, also referred to as a CITE sequence). The expression cassette may also include other features such as an origin of replication, and/or chromosome integration elements such as retroviral long terminal repeats (LTRs), or adeno-associated viral (AAV) inverted terminal repeats (ITRs).

[0449] Polynucleotide and nucleic acid coding regions of the present invention may be associated with additional coding regions which encode secretory or signal peptides, which direct the secretion of a polypeptide encoded by a polynucleotide of the present invention. For example, if secretion of the protease-activatable T cell activating bispecific molecule is desired, DNA encoding a signal sequence may be placed upstream of the nucleic acid encoding a protease-activatable T cell activating bispecific molecule of the invention or a fragment thereof. According to the signal hypothesis, proteins secreted by mammalian cells have a signal peptide or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Those of ordinary skill in the art are aware that polypeptides secreted by vertebrate cells generally have a signal peptide fused to the N-terminus of the polypeptide, which is cleaved from the translated polypeptide to produce a secreted or “mature” form of the polypeptide. In certain embodiments, the native signal peptide, e.g., an immunoglobulin heavy chain or light chain signal peptide is used, or

a functional derivative of that sequence that retains the ability to direct the secretion of the polypeptide that is operably associated with it. Alternatively, a heterologous mammalian signal peptide, or a functional derivative thereof, may be used. For example, the wild-type leader sequence may be substituted with the leader sequence of human tissue plasminogen activator (TPA) or mouse β -glucuronidase.

[0450] DNA encoding a short protein sequence that could be used to facilitate later purification (e.g., a histidine tag) or assist in labeling the protease-activatable T cell activating bispecific molecule may be included within or at the ends of the protease-activatable T cell activating bispecific molecule (fragment) encoding polynucleotide.

[0451] In a further embodiment, a host cell comprising one or more polynucleotides of the invention is provided. In certain embodiments a host cell comprising one or more vectors of the invention is provided. The polynucleotides and vectors may incorporate any of the features, singly or in combination, described herein in relation to polynucleotides and vectors, respectively. In one such embodiment a host cell comprises (e.g., has been transformed or transfected with) a vector comprising a polynucleotide that encodes (part of) a protease-activatable T cell activating bispecific molecule of the invention. As used herein, the term “host cell” refers to any kind of cellular system which can be engineered to generate the protease-activatable T cell activating bispecific molecules of the invention or fragments thereof. Host cells suitable for replicating and for supporting expression of protease-activatable T cell activating bispecific molecules are well known in the art. Such cells may be transfected or transduced as appropriate with the particular expression vector and large quantities of vector containing cells can be grown for seeding large scale fermenters to obtain sufficient quantities of the protease-activatable T cell activating bispecific molecule for clinical applications. Suitable host cells include prokaryotic microorganisms, such as *E. coli*, or various eukaryotic cells, such as Chinese hamster ovary cells (CHO), insect cells, or the like. For example, polypeptides may be produced in bacteria in particular when glycosylation is not needed. After expression, the polypeptide may be isolated from the bacterial cell paste in a soluble fraction and can be further purified. In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for polypeptide-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been “humanized”, resulting in the production of a polypeptide with a partially or fully human glycosylation pattern. See Gerngross, *Nat Biotech* 22, 1409-1414 (2004), and Li et al., *Nat Biotech* 24, 210-215 (2006). Suitable host cells for the expression of (glycosylated) polypeptides are also derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures can also be utilized as hosts. See e.g., U.S. Pat. Nos. 5,959,177, 6,040,498, 6,420,548, 7,125,978, and 6,417,429 (describing PLANTIBODIES™ technology for producing antibodies in transgenic plants). Vertebrate cells may also be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line

transformed by SV40 (COS-7); human embryonic kidney line (293 or 293T cells as described, e.g., in Graham et al., *J Gen Virol* 36, 59 (1977)), baby hamster kidney cells (BHK), mouse sertoli cells (TM4 cells as described, e.g., in Mather, *Biol Reprod* 23, 243-251 (1980)), monkey kidney cells (CV1), African green monkey kidney cells (VERO-76), human cervical carcinoma cells (HELA), canine kidney cells (MDCK), buffalo rat liver cells (BRL 3A), human lung cells (W138), human liver cells (Hep G2), mouse mammary tumor cells (MMT 060562), TRI cells (as described, e.g., in Mather et al., *Annals N.Y. Acad Sci* 383, 44-68 (1982)), MRC 5 cells, and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including dhfr⁻ CHO cells (Urlaub et al., *Proc Natl Acad Sci USA* 77, 4216 (1980)); and myeloma cell lines such as YO, NS0, P3X63 and Sp2/0. For a review of certain mammalian host cell lines suitable for protein production, see, e.g., Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, N.J.), pp. 255-268 (2003). Host cells include cultured cells, e.g., mammalian cultured cells, yeast cells, insect cells, bacterial cells and plant cells, to name only a few, but also cells comprised within a transgenic animal, transgenic plant or cultured plant or animal tissue. In one embodiment, the host cell is a eukaryotic cell, preferably a mammalian cell, such as a Chinese Hamster Ovary (CHO) cell, a human embryonic kidney (HEK) cell or a lymphoid cell (e.g., YO, NS0, Sp20 cell).

[0452] Standard technologies are known in the art to express foreign genes in these systems. Cells expressing a polypeptide comprising either the heavy or the light chain of an antigen binding domain such as an antibody, may be engineered so as to also express the other of the antibody chains such that the expressed product is an antibody that has both a heavy and a light chain. In one embodiment, a method of producing a protease-activatable T cell activating bispecific molecule according to the invention is provided, wherein the method comprises culturing a host cell comprising a polynucleotide encoding the protease-activatable T cell activating bispecific molecule, as provided herein, under conditions suitable for expression of the protease-activatable T cell activating bispecific molecule, and recovering the protease-activatable T cell activating bispecific molecule from the host cell (or host cell culture medium).

[0453] The components of the protease-activatable T cell activating bispecific molecule are genetically fused to each other. Protease-activatable T cell activating bispecific molecules can be designed such that its components are fused directly to each other or indirectly through a linker sequence. The composition and length of the linker may be determined in accordance with methods well known in the art and may be tested for efficacy. Examples of linker sequences between different components of protease-activatable T cell activating bispecific molecules are found in the sequences provided herein. Additional sequences may also be included to incorporate a cleavage site to separate the individual components of the fusion if desired, for example an endopeptidase recognition sequence. In certain embodiments the one or more antigen binding moieties of the protease-activatable T cell activating bispecific molecules comprise at least an antibody variable region capable of binding an antigenic determinant. Variable regions can form part of and be derived from naturally or non-naturally occurring antibodies and fragments thereof. Methods to produce polyclonal anti-

bodies and monoclonal antibodies are well known in the art (see e.g., Harlow and Lane, "Antibodies, a laboratory manual", Cold Spring Harbor Laboratory, 1988). Non-naturally occurring antibodies can be constructed using solid phase-peptide synthesis, can be produced recombinantly (e.g., as described in U.S. Pat. No. 4,186,567) or can be obtained, for example, by screening combinatorial libraries comprising variable heavy chains and variable light chains (see e.g., U.S. Pat. No. 5,969,108 to McCafferty).

[0454] Any animal species of antibody, antibody fragment, antigen binding domain or variable region can be used in the protease-activatable T cell activating bispecific molecules of the invention. Non-limiting antibodies, antibody fragments, antigen binding domains or variable regions useful in the present invention can be of murine, primate, or human origin. If the protease-activatable T cell activating bispecific molecule is intended for human use, a chimeric form of antibody may be used wherein the constant regions of the antibody are from a human. A humanized or fully human form of the antibody can also be prepared in accordance with methods well known in the art (see e.g. U.S. Pat. No. 5,565,332 to Winter). Humanization may be achieved by various methods including, but not limited to (a) grafting the non-human (e.g., donor antibody) CDRs onto human (e.g., recipient antibody) framework and constant regions with or without retention of critical framework residues (e.g., those that are important for retaining good antigen binding affinity or antibody functions), (b) grafting only the non-human specificity-determining regions (SDRs or a-CDRs; the residues critical for the antibody-antigen interaction) onto human framework and constant regions, or (c) transplanting the entire non-human variable domains, but "cloaking" them with a human-like section by replacement of surface residues. Humanized antibodies and methods of making them are reviewed, e.g., in Almagro and Fransson, *Front Biosci* 13, 1619-1633 (2008), and are further described, e.g., in Riechmann et al., *Nature* 332, 323-329 (1988); Queen et al., *Proc Natl Acad Sci USA* 86, 10029-10033 (1989); U.S. Pat. Nos. 5,821,337, 7,527,791, 6,982, 321, and 7,087,409; Jones et al., *Nature* 321, 522-525 (1986); Morrison et al., *Proc Natl Acad Sci* 81, 6851-6855 (1984); Morrison and Oi, *Adv Immunol* 44, 65-92 (1988); Verhoeven et al., *Science* 239, 1534-1536 (1988); Padlan, *Molec Immun* 31(3), 169-217 (1994); Kashmiri et al., *Methods* 36, 25-34 (2005) (describing SDR (a-CDR) grafting); Padlan, *Mol Immunol* 28, 489-498 (1991) (describing "resurfacing"); Dall'Acqua et al., *Methods* 36, 43-60 (2005) (describing "FR shuffling"); and Osbourn et al., *Methods* 36, 61-68 (2005) and Klimka et al., *Br J Cancer* 83, 252-260 (2000) (describing the "guided selection" approach to FR shuffling). Human antibodies and human variable regions can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk and van de Winkel, *Curr Opin Pharmacol* 5, 368-74 (2001) and Lonberg, *Curr Opin Immunol* 20, 450-459 (2008). Human variable regions can form part of and be derived from human monoclonal antibodies made by the hybridoma method (see e.g., *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)). Human antibodies and human variable regions may also be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge (see e.g., Lon-

berg, *Nat Biotech* 23, 1117-1125 (2005). Human antibodies and human variable regions may also be generated by isolating Fv clone variable region sequences selected from human-derived phage display libraries (see e.g., Hoogenboom et al. in *Methods in Molecular Biology* 178, 1-37 (O'Brien et al., ed., Human Press, Totowa, N.J., 2001); and McCafferty et al., *Nature* 348, 552-554; Clackson et al., *Nature* 352, 624-628 (1991)). Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments.

[0455] In certain embodiments, the antigen binding moieties useful in the present invention are engineered to have enhanced binding affinity according to, for example, the methods disclosed in U.S. Pat. Appl. Publ. No. 2004/0132066, the entire contents of which are hereby incorporated by reference. The ability of the protease-activatable T cell activating bispecific molecule of the invention to bind to a specific antigenic determinant can be measured either through an enzyme-linked immunosorbent assay (ELISA) or other techniques familiar to one of skill in the art, e.g., surface plasmon resonance technique (analyzed on a BIA-CORE T100 system) (Liljeblad, et al., *Glyco J* 17, 323-329 (2000)), and traditional binding assays (Heeley, *Endocr Res* 28, 217-229 (2002)). Competition assays may be used to identify an antibody, antibody fragment, antigen binding domain or variable domain that competes with a reference antibody for binding to a particular antigen, e.g., an antibody that competes with the V9 antibody for binding to CD3. In certain embodiments, such a competing antibody binds to the same epitope (e.g., a linear or a conformational epitope) that is bound by the reference antibody. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in *Methods in Molecular Biology* vol. 66 (Humana Press, Totowa, N.J.). In an exemplary competition assay, immobilized antigen (e.g., CD3) is incubated in a solution comprising a first labeled antibody that binds to the antigen (e.g., V9 antibody, described in U.S. Pat. No. 6,054,297) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to the antigen. The second antibody may be present in a hybridoma supernatant. As a control, immobilized antigen is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first antibody to the antigen, excess unbound antibody is removed, and the amount of label associated with immobilized antigen is measured. If the amount of label associated with immobilized antigen is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to the antigen. See Harlow and Lane (1988) *Antibodies: A Laboratory Manual* ch.14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.).

[0456] Protease-activatable T cell activating bispecific molecules prepared as described herein may be purified by art-known techniques such as high performance liquid chromatography, ion exchange chromatography, gel electrophoresis, affinity chromatography, size exclusion chromatography, and the like. The actual conditions used to purify a particular protein will depend, in part, on factors such as net charge, hydrophobicity, hydrophilicity etc., and will be apparent to those having skill in the art. For affinity chromatography purification an antibody, ligand, receptor or

antigen can be used to which the protease-activatable T cell activating bispecific molecule binds. For example, for affinity chromatography purification of protease-activatable T cell activating bispecific molecules of the invention, a matrix with protein A or protein G may be used. Sequential Protein A or G affinity chromatography and size exclusion chromatography can be used to isolate a protease-activatable T cell activating bispecific molecule essentially as described in the Examples. The purity of the protease-activatable T cell activating bispecific molecule can be determined by any of a variety of well-known analytical methods including gel electrophoresis, high pressure liquid chromatography, and the like. For example, the heavy chain fusion proteins expressed as described in the Examples were shown to be intact and properly assembled as demonstrated by reducing SDS-PAGE (see, e.g., FIGS. 8-12). Three bands were resolved at approximately Mr 25,000, Mr 50,000 and Mr 75,000, corresponding to the predicted molecular weights of the protease-activatable T cell activating bispecific molecule light chain, heavy chain and heavy chain/light chain fusion protein.

Assays

[0457] protease-activatable T cell activating bispecific molecules provided herein may be identified, screened for, or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

Affinity Assays

[0458] The affinity of the protease-activatable T cell activating bispecific molecule for an Fc receptor or a target antigen can be determined in accordance with the methods set forth in the Examples by surface plasmon resonance (SPR), using standard instrumentation such as a BIAcore instrument (GE Healthcare), and receptors or target proteins such as may be obtained by recombinant expression. Alternatively, binding of protease-activatable T cell activating bispecific molecules for different receptors or target antigens may be evaluated using cell lines expressing the particular receptor or target antigen, for example by flow cytometry (FACS). A specific illustrative and exemplary embodiment for measuring binding affinity is described in the following and in the Examples below. According to one embodiment, K_D is measured by surface plasmon resonance using a BIAcore® T100 machine (GE Healthcare) at 25° C.

[0459] To analyze the interaction between the Fc-portion and Fc receptors, His-tagged recombinant Fc-receptor is captured by an anti-Penta His antibody (Qiagen) immobilized on CMS chips and the bispecific constructs are used as analytes. Briefly, carboxymethylated dextran biosensor chips (CMS, GE Healthcare) are activated with N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Anti Penta-His antibody is diluted with 10 mM sodium acetate, pH 5.0, to 40 mg/ml before injection at a flow rate of 5 μ l/min to achieve approximately 6500 response units (RU) of coupled protein. Following the injection of the ligand, 1 M ethanolamine is injected to block unreacted groups. Subsequently the Fc-receptor is captured for 60 s at 4 or 10 nM. For kinetic measurements, four-fold serial dilutions of the bispecific construct (range between 500 nM and 4000 nM) are injected in HBS-EP (GE Health-

care, 10 mM HEPES, 150 mM NaCl, 3 mM EDTA, 0.05% Surfactant P20, pH 7.4) at 25° C. at a flow rate of 30 μ l/min for 120 s.

[0460] To determine the affinity to the target antigen, bispecific constructs are captured by an anti-human Fab specific antibody (GE Healthcare) that is immobilized on an activated CMS-sensor chip surface as described for the anti Penta-His antibody. The final amount of coupled protein is approximately 12000 R U. The bispecific constructs are captured for 90 s at 300 nM. The target antigens are passed through the flow cells for 180 s at a concentration range from 250 to 1000 nM with a flowrate of 30 μ l/min. The dissociation is monitored for 180 s.

[0461] Bulk refractive index differences are corrected for by subtracting the response obtained on reference flow cell. The steady state response was used to derive the dissociation constant K_D by non-linear curve fitting of the Langmuir binding isotherm. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIAcore® T100 Evaluation Software version 1.1.1) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (K_D) is calculated as the ratio k_{off}/k_{on} . See, e.g., Chen et al., J Mol Biol 293, 865-881 (1999).

Activity Assays

[0462] Biological activity of the protease-activatable T cell activating bispecific molecules of the invention can be measured by various assays as described in the Examples. Biological activities may for example include the induction of proliferation of T cells, the induction of signaling in T cells, the induction of expression of activation markers in T cells, the induction of cytokine secretion by T cells, the induction of lysis of target cells such as tumor cells, and the induction of tumor regression and/or the improvement of survival.

Compositions, Formulations, and Routes of Administration

[0463] In a further aspect, the invention provides pharmaceutical compositions comprising any of the protease-activatable T cell activating bispecific molecules provided herein, e.g., for use in any of the below therapeutic methods. In one embodiment, a pharmaceutical composition comprises any of the protease-activatable T cell activating bispecific molecules provided herein and a pharmaceutically acceptable carrier. In another embodiment, a pharmaceutical composition comprises any of the protease-activatable T cell activating bispecific molecules provided herein and at least one additional therapeutic agent, e.g., as described below.

[0464] Further provided is a method of producing a protease-activatable T cell activating bispecific molecule of the invention in a form suitable for administration in vivo, the method comprising (a) obtaining a protease-activatable T cell activating bispecific molecule according to the invention, and (b) formulating the protease-activatable T cell activating bispecific molecule with at least one pharmaceutically acceptable carrier, whereby a preparation of protease-activatable T cell activating bispecific molecule is formulated for administration in vivo.

[0465] Pharmaceutical compositions of the present invention comprise a therapeutically effective amount of one or more protease-activatable T cell activating bispecific molecule dissolved or dispersed in a pharmaceutically accept-

able carrier. The phrases “pharmaceutical or pharmacologically acceptable” refers to molecular entities and compositions that are generally non-toxic to recipients at the dosages and concentrations employed, i.e. do not produce an adverse, allergic or other untoward reaction when administered to an animal, such as, for example, a human, as appropriate. The preparation of a pharmaceutical composition that contains at least one protease-activatable T cell activating bispecific molecule and optionally an additional active ingredient will be known to those of skill in the art in light of the present disclosure, as exemplified by Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, incorporated herein by reference. Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biological Standards or corresponding authorities in other countries. Preferred compositions are lyophilized formulations or aqueous solutions. As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, buffers, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, antioxidants, proteins, drugs, drug stabilizers, polymers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-1329, incorporated herein by reference). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated. The composition may comprise different types of carriers depending on whether it is to be administered in solid, liquid or aerosol form, and whether it need to be sterile for such routes of administration as injection. Protease-activatable T cell activating bispecific molecules of the present invention (and any additional therapeutic agent) can be administered intravenously, intradermally, intraarterially, intraperitoneally, intrasplenicly, intracranially, intraarticularly, intraprostatically, intrasplenicly, intrarectally, intrapleurally, intratracheally, intranasally, intravitreally, intravaginally, intrarectally, intratumorally, intramuscularly, intraperitoneally, subcutaneously, subconjunctivally, intravesicularly, mucosally, intrapericardially, intraumbilically, intraocularly, orally, topically, locally, by inhalation (e.g., aerosol inhalation), injection, infusion, continuous infusion, localized perfusion bathing target cells directly, via a catheter, via a lavage, in cremes, in lipid compositions (e.g., liposomes), or by other method or any combination of the foregoing as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, incorporated herein by reference). Parenteral administration, in particular intravenous injection, is most commonly used for administering polypeptide molecules such as the protease-activatable T cell activating bispecific molecules of the invention.

[0466] Parenteral compositions include those designed for administration by injection, e.g., subcutaneous, intradermal, intrasplenic, intravenous, intraarterial intramuscular, intrathecal or intraperitoneal injection. For injection, the protease-activatable T cell activating bispecific molecules of the invention may be formulated in aqueous solutions,

preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. The solution may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the protease-activatable T cell activating bispecific molecules may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use. Sterile injectable solutions are prepared by incorporating the protease-activatable T cell activating bispecific molecules of the invention in the required amount in the appropriate solvent with various of the other ingredients enumerated below, as required. Sterility may be readily accomplished, e.g., by filtration through sterile filtration membranes. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and/or the other ingredients. In the case of sterile powders for the preparation of sterile injectable solutions, suspensions or emulsion, the preferred methods of preparation are vacuum-drying or freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered liquid medium thereof. The liquid medium should be suitably buffered if necessary and the liquid diluent first rendered isotonic prior to injection with sufficient saline or glucose. The composition must be stable under the conditions of manufacture and storage, and preserved against the contaminating action of microorganisms, such as bacteria and fungi. It will be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein. Suitable pharmaceutically acceptable carriers include, but are not limited to: buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g., Zn-protein complexes); and/or non-ionic surfactants such as polyethylene glycol (PEG). Aqueous injection suspensions may contain compounds which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, dextran, or the like. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes.

[0467] Active ingredients may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmeth-

acylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences (18th Ed. Mack Printing Company, 1990). Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the polypeptide, which matrices are in the form of shaped articles, e.g., films, or microcapsules. In particular embodiments, prolonged absorption of an injectable composition can be brought about by the use in the compositions of agents delaying absorption, such as, for example, aluminum monostearate, gelatin or combinations thereof.

[0468] In addition to the compositions described previously, the protease-activatable T cell activating bispecific molecules may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the protease-activatable T cell activating bispecific molecules may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt. Pharmaceutical compositions comprising the protease-activatable T cell activating bispecific molecules of the invention may be manufactured by means of conventional mixing, dissolving, emulsifying, encapsulating, entrapping or lyophilizing processes. Pharmaceutical compositions may be formulated in conventional manner using one or more physiologically acceptable carriers, diluents, excipients or auxiliaries which facilitate processing of the proteins into preparations that can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

[0469] The protease-activatable T cell activating bispecific molecules may be formulated into a composition in a free acid or base, neutral or salt form. Pharmaceutically acceptable salts are salts that substantially retain the biological activity of the free acid or base. These include the acid addition salts, e.g., those formed with the free amino groups of a proteinaceous composition, or which are formed with inorganic acids such as for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric or mandelic acid. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as for example, sodium, potassium, ammonium, calcium or ferric hydroxides; or such organic bases as isopropylamine, trimethylamine, histidine or procaine. Pharmaceutical salts tend to be more soluble in aqueous and other protic solvents than are the corresponding free base forms.

Therapeutic Methods and Compositions

[0470] Any of the protease-activatable T cell activating bispecific molecules provided herein may be used in therapeutic methods. Protease-activatable T cell activating bispecific molecules of the invention can be used as immunotherapeutic agents, for example in the treatment of cancers.

[0471] For use in therapeutic methods, protease-activatable T cell activating bispecific molecules of the invention would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being

treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners.

[0472] In one aspect, protease-activatable T cell activating bispecific molecules of the invention for use as a medicament are provided. In further aspects, protease-activatable T cell activating bispecific molecules of the invention for use in treating a disease are provided. In certain embodiments, protease-activatable T cell activating bispecific molecules of the invention for use in a method of treatment are provided. In one embodiment, the invention provides a protease-activatable T cell activating bispecific molecule as described herein for use in the treatment of a disease in an individual in need thereof. In certain embodiments, the invention provides a protease-activatable T cell activating bispecific molecule for use in a method of treating an individual having a disease comprising administering to the individual a therapeutically effective amount of the protease-activatable T cell activating bispecific molecule. In certain embodiments the disease to be treated is a proliferative disorder. In a particular embodiment the disease is cancer. In certain embodiments the method further comprises administering to the individual a therapeutically effective amount of at least one additional therapeutic agent, e.g., an anti-cancer agent if the disease to be treated is cancer.

[0473] In further embodiments, the invention provides a protease-activatable T cell activating bispecific molecule as described herein for use in inducing lysis of a target cell, particularly a tumor cell. In certain embodiments, the invention provides a protease-activatable T cell activating bispecific molecule for use in a method of inducing lysis of a target cell, particularly a tumor cell, in an individual comprising administering to the individual an effective amount of the protease-activatable T cell activating bispecific molecule to induce lysis of a target cell. An "individual" according to any of the above embodiments is a mammal, preferably a human.

[0474] In a further aspect, the invention provides for the use of a protease-activatable T cell activating bispecific molecule of the invention in the manufacture or preparation of a medicament. In one embodiment the medicament is for the treatment of a disease in an individual in need thereof. In a further embodiment, the medicament is for use in a method of treating a disease comprising administering to an individual having the disease a therapeutically effective amount of the medicament. In certain embodiments the disease to be treated is a proliferative disorder. In a particular embodiment the disease is cancer. In one embodiment, the method further comprises administering to the individual a therapeutically effective amount of at least one additional therapeutic agent, e.g., an anti-cancer agent if the disease to be treated is cancer. In a further embodiment, the medicament is for inducing lysis of a target cell, particularly a tumor cell. In still a further embodiment, the medicament is for use in a method of inducing lysis of a target cell, particularly a tumor cell, in an individual comprising administering to the individual an effective amount of the medicament to induce lysis of a target cell. An "individual" according to any of the above embodiments may be a mammal, preferably a human.

[0475] In a further aspect, the invention provides a method for treating a disease. In one embodiment, the method

comprises administering to an individual having such disease a therapeutically effective amount of a protease-activatable T cell activating bispecific molecule of the invention. In one embodiment a composition is administered to said individual, comprising the protease-activatable T cell activating bispecific molecule of the invention in a pharmaceutically acceptable form. In certain embodiments the disease to be treated is a proliferative disorder. In a particular embodiment the disease is cancer. In certain embodiments the method further comprises administering to the individual a therapeutically effective amount of at least one additional therapeutic agent, e.g., an anti-cancer agent if the disease to be treated is cancer. An “individual” according to any of the above embodiments may be a mammal, preferably a human.

[0476] In a further aspect, the invention provides a method for inducing lysis of a target cell, particularly a tumor cell. In one embodiment the method comprises contacting a target cell with a protease-activatable T cell activating bispecific molecule of the invention in the presence of a T cell, particularly a cytotoxic T cell. In a further aspect, a method for inducing lysis of a target cell, particularly a tumor cell, in an individual is provided. In one such embodiment, the method comprises administering to the individual an effective amount of a protease-activatable T cell activating bispecific molecule to induce lysis of a target cell. In one embodiment, an “individual” is a human.

[0477] In certain embodiments the disease to be treated is a proliferative disorder, particularly cancer. Non-limiting examples of cancers include bladder cancer, brain cancer, head and neck cancer, pancreatic cancer, lung cancer, breast cancer, ovarian cancer, uterine cancer, cervical cancer, endometrial cancer, esophageal cancer, colon cancer, colorectal cancer, rectal cancer, gastric cancer, prostate cancer, blood cancer, skin cancer, squamous cell carcinoma, bone cancer, and kidney cancer. Other cell proliferation disorders that can be treated using a protease-activatable T cell activating bispecific molecule of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous system (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic region, and urogenital system. Also included are pre-cancerous conditions or lesions and cancer metastases. In certain embodiments the cancer is chosen from the group consisting of renal cell cancer, skin cancer, lung cancer, colorectal cancer, breast cancer, brain cancer, head and neck cancer. A skilled artisan readily recognizes that in many cases the protease-activatable T cell activating bispecific molecule may not provide a cure but may only provide partial benefit. In some embodiments, a physiological change having some benefit is also considered therapeutically beneficial. Thus, in some embodiments, an amount of protease-activatable T cell activating bispecific molecule that provides a physiological change is considered an “effective amount” or a “therapeutically effective amount”. The subject, patient, or individual in need of treatment is typically a mammal, more specifically a human. In some embodiments, an effective amount of a protease-activatable T cell activating bispecific molecule of the invention is administered to a cell. In other embodiments, a therapeutically effective amount of a protease-activatable T cell activating bispecific molecule of the invention is administered to an individual for the treatment of disease.

[0478] For the prevention or treatment of disease, the appropriate dosage of a protease-activatable T cell activating bispecific molecule of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the type of disease to be treated, the route of administration, the body weight of the patient, the type of T cell activating bispecific antigen binding molecule, the severity and course of the disease, whether the T cell activating bispecific antigen binding molecule is administered for preventive or therapeutic purposes, previous or concurrent therapeutic interventions, the patient’s clinical history and response to the protease-activatable T cell activating bispecific molecule, and the discretion of the attending physician. The practitioner responsible for administration will, in any event, determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

[0479] The protease-activatable T cell activating bispecific molecule is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, about 1 µg/kg to 15 mg/kg (e.g., 0.1 mg/kg-10 mg/kg) of protease-activatable T cell activating bispecific molecule can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of disease symptoms occurs. One exemplary dosage of the T cell activating bispecific antigen binding molecule would be in the range from about 0.005 mg/kg to about 10 mg/kg. In other non-limiting examples, a dose may also comprise from about 1 microgram/kg body weight, about 5 microgram/kg body weight, about 10 microgram/kg body weight, about 50 microgram/kg body weight, about 100 microgram/kg body weight, about 200 microgram/kg body weight, about 350 microgram/kg body weight, about 500 microgram/kg body weight, about 1 milligram/kg body weight, about 5 milligram/kg body weight, about 10 milligram/kg body weight, about 50 milligram/kg body weight, about 100 milligram/kg body weight, about 200 milligram/kg body weight, about 350 milligram/kg body weight, about 500 milligram/kg body weight, to about 1000 mg/kg body weight or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from the numbers listed herein, a range of about 5 mg/kg body weight to about 100 mg/kg body weight, about 5 microgram/kg body weight to about 500 milligram/kg body weight, etc., can be administered, based on the numbers described above. Thus, one or more doses of about 0.5 mg/kg, 2.0 mg/kg, 5.0 mg/kg or 10 mg/kg (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, e.g., every week or every three weeks (e.g., such that the patient receives from about two to about twenty, or e.g., about six doses of the protease-activatable T cell activating bispecific molecule). An initial higher loading dose, followed by one or more lower doses may be administered.

However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

[0480] The protease-activatable T cell activating bispecific molecule of the invention will generally be used in an amount effective to achieve the intended purpose. For use to treat or prevent a disease condition, the protease-activatable T cell activating bispecific molecules of the invention, or pharmaceutical compositions thereof, are administered or applied in a therapeutically effective amount. Determination of a therapeutically effective amount is well within the capabilities of those skilled in the art, especially in light of the detailed disclosure provided herein.

[0481] For systemic administration, a therapeutically effective dose can be estimated initially from in vitro assays, such as cell culture assays. A dose can then be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture. Such information can be used to more accurately determine useful doses in humans.

[0482] Initial dosages can also be estimated from in vivo data, e.g., animal models, using techniques that are well known in the art. One having ordinary skill in the art could readily optimize administration to humans based on animal data.

[0483] Dosage amount and interval may be adjusted individually to provide plasma levels of the protease-activatable T cell activating bispecific molecules which are sufficient to maintain therapeutic effect. Usual patient dosages for administration by injection range from about 0.1 to 50 mg/kg/day, typically from about 0.5 to 1 mg/kg/day. Therapeutically effective plasma levels may be achieved by administering multiple doses each day. Levels in plasma may be measured, for example, by HPLC. In cases of local administration or selective uptake, the effective local concentration of the protease-activatable T cell activating bispecific molecules may not be related to plasma concentration. One having skill in the art will be able to optimize therapeutically effective local dosages without undue experimentation.

[0484] A therapeutically effective dose of the protease-activatable T cell activating bispecific molecules described herein will generally provide therapeutic benefit without causing substantial toxicity. Toxicity and therapeutic efficacy of a protease-activatable T cell activating bispecific molecule can be determined by standard pharmaceutical procedures in cell culture or experimental animals. Cell culture assays and animal studies can be used to determine the LD_{50} (the dose lethal to 50% of a population) and the ED_{50} (the dose therapeutically effective in 50% of a population). The dose ratio between toxic and therapeutic effects is the therapeutic index, which can be expressed as the ratio LD_{50}/ED_{50} . Protease-activatable T cell activating bispecific molecule that exhibit large therapeutic indices are preferred. In one embodiment, the protease-activatable T cell activating bispecific molecule according to the present invention exhibits a high therapeutic index. The data obtained from cell culture assays and animal studies can be used in formulating a range of dosages suitable for use in humans. The dosage lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage may vary within this range depending upon a variety of factors, e.g., the dosage form employed, the route of administration utilized, the condition of the subject,

and the like. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition (see, e.g., Fingl et al., 1975, in: *The Pharmacological Basis of Therapeutics*, Ch. 1, p. 1, incorporated herein by reference in its entirety). The attending physician for patients treated with protease-activatable T cell activating bispecific molecules of the invention would know how and when to terminate, interrupt, or adjust administration due to toxicity, organ dysfunction, and the like. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate (precluding toxicity). The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated, with the route of administration, and the like. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency will also vary according to the age, body weight, and response of the individual patient.

Other Agents and Treatments

[0485] The protease-activatable T cell activating bispecific molecules of the invention may be administered in combination with one or more other agents in therapy. For instance, a protease-activatable T cell activating bispecific molecule of the invention may be co-administered with at least one additional therapeutic agent. The term "therapeutic agent" encompasses any agent administered to treat a symptom or disease in an individual in need of such treatment. Such additional therapeutic agent may comprise any active ingredients suitable for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. In certain embodiments, an additional therapeutic agent is an immunomodulatory agent, a cytostatic agent, an inhibitor of cell adhesion, a cytotoxic agent, an activator of cell apoptosis, or an agent that increases the sensitivity of cells to apoptotic inducers. In a particular embodiment, the additional therapeutic agent is an anti-cancer agent, for example a microtubule disruptor, an antimetabolite, a topoisomerase inhibitor, a DNA intercalator, an alkylating agent, a hormonal therapy, a kinase inhibitor, a receptor antagonist, an activator of tumor cell apoptosis, or an antiangiogenic agent.

[0486] Such other agents are suitably present in combination in amounts that are effective for the purpose intended. The effective amount of such other agents depends on the amount of protease-activatable T cell activating bispecific molecule used, the type of disorder or treatment, and other factors discussed above. The protease-activatable T cell activating bispecific molecule are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

[0487] Such combination therapies noted above encompass combined administration (where two or more therapeutic agents are included in the same or separate compositions), and separate administration, in which case, administration of the protease-activatable T cell activating bispecific molecule of the invention can occur prior to, simultaneously, and/or following, administration of the additional therapeutic agent and/or adjuvant. Protease-activatable T cell activating bispecific molecules of the invention can also be used in combination with radiation therapy.

Articles of Manufacture

[0488] In another aspect of the invention, an article of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above is provided. The article of manufacture comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, IV solution bags, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is a protease-activatable T cell activating bispecific molecule of the invention. The label or package insert indicates that the composition is used for treating the condition of choice. Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises a protease-activatable T cell activating bispecific molecule of the invention; and (b) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

EXEMPLARY EMBODIMENTS

- [0489]** 1. A protease-activatable T cell activating bispecific molecule comprising
- [0490]** (a) a first antigen binding moiety capable of specific binding to CD3;
- [0491]** (b) a second antigen binding moiety capable of specific binding to a target cell antigen; and
- [0492]** (c) a masking moiety covalently attached to the T cell bispecific binding molecule through a protease-cleavable linker, wherein the masking moiety is capable of specific binding to the idiotype of the first or the second antigen binding moiety thereby reversibly concealing the first or the second antigen binding moiety.
- [0493]** 2. The protease-activatable T cell activating bispecific molecule of embodiment 1, wherein the masking moiety is covalently attached to the first antigen binding moiety and reversibly conceals the first antigen binding moiety.
- [0494]** 3. The protease-activatable T cell activating bispecific molecule of embodiment 1 or 2, wherein the masking moiety is covalently attached to the heavy chain variable region of the first antigen binding moiety.
- [0495]** 4. The protease-activatable T cell activating bispecific molecule of embodiment 1 or 2, wherein the masking moiety is covalently attached to the light chain variable region of the first antigen binding moiety.
- [0496]** 5. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 4, wherein the masking moiety is an anti-idiotypic scFv.
- [0497]** 6. The protease-activatable T cell activating bispecific molecule of any one of embodiments 2 to 5, wherein the protease-activatable T cell activating bispecific molecule comprises a second masking moiety reversibly concealing the second antigen binding moiety.
- [0498]** 7. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 6, wherein the protease is expressed by the target cell.
- [0499]** 8. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 7, wherein the second antigen binding moiety is a cross-over Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy chain are exchanged.
- [0500]** 9. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 8, wherein the second antigen binding moiety is a cross-over Fab molecule wherein the constant regions of the Fab light chain and the Fab heavy chain are exchanged.
- [0501]** 10. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 9, wherein the first antigen binding moiety is a conventional Fab molecule.
- [0502]** 11. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 10, comprising not more than one antigen binding moiety capable of specific binding to CD3.
- [0503]** 12. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 11, comprising a third antigen binding moiety which is a Fab molecule capable of specific binding to a target cell antigen.
- [0504]** 13. The protease-activatable T cell activating bispecific molecule of embodiment 12, wherein the third antigen binding moiety is identical to the second antigen binding moiety.
- [0505]** 14. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 13, wherein the second antigen binding moiety is capable of specific binding to FolR1 or HER.
- [0506]** 15. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 13, wherein the second antigen binding moiety is capable of specific binding to a target cell antigen selected from the group consisting of FolR1, HER1 and Mesothelin.
- [0507]** 16. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 13, wherein the second antigen binding moiety is capable of specific binding to a target cell antigen selected from the group consisting of FolR1, HER1, HER2 and Mesothelin.
- [0508]** 17. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 16, wherein the first and the second antigen binding moiety are fused to each other, optionally via a peptide linker.
- [0509]** 18. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 17, wherein the second antigen binding moiety is fused at

- the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety.
- [0510] 19. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 17, wherein the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety.
- [0511] 20. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 19, wherein the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety are fused to each other, optionally via a peptide linker.
- [0512] 21. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 20, additionally comprising an Fc domain composed of a first and a second subunit capable of stable association.
- [0513] 22. The protease-activatable T cell activating bispecific molecule of embodiment 21, wherein the Fc domain is an IgG, specifically an IgG₁ or IgG₄, Fc domain.
- [0514] 23. The protease-activatable T cell activating bispecific molecule of embodiment 21 or 22, wherein the Fc domain is a human Fc domain.
- [0515] 24. The protease-activatable T cell activating bispecific molecule of any one of embodiments 21 to 23, wherein the Fc domain exhibits reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain.
- [0516] 25. The protease-activatable T cell activating bispecific molecule of embodiment 24, wherein the Fc domain comprises one or more amino acid substitution that reduces binding to an Fc receptor and/or effector function.
- [0517] 26. The protease-activatable T cell activating bispecific molecule of embodiment 25, wherein said one or more amino acid substitution is at one or more position selected from the group of L234, L235, and P329 (Kabat numbering).
- [0518] 27. The protease-activatable T cell activating bispecific molecule of embodiment 26, wherein each subunit of the Fc domain comprises three amino acid substitutions that reduce binding to an activating Fc receptor and/or effector function wherein said amino acid substitutions are L234A, L235A and P329G.
- [0519] 28. The protease-activatable T cell activating bispecific molecule of any one of embodiments 24 to 27, wherein the Fc receptor is an Fcγ receptor.
- [0520] 29. The protease-activatable T cell activating bispecific molecule of any one of embodiments 24 to 28, wherein the effector function is antibody-dependent cell-mediated cytotoxicity (ADCC).
- [0521] 30. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 29, wherein the masking moiety comprises a heavy chain variable region comprising at least one of:
- [0522] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);
- [0523] (b) a CDR H2 amino acid sequence of WIN-TETGEPAYADDFKG (SEQ ID NO:21); and
- [0524] (c) a CDR H3 amino acid sequence of PYDYPDVL DY (SEQ ID NO:22).
- [0525] 31. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 30, wherein the masking moiety comprises a light chain variable region comprising at least one of:
- [0526] (d) a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO:23);
- [0527] (e) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and
- [0528] (f) a CDR L3 amino acid sequence of QHS-REFPWT (SEQ ID NO:25).
- [0529] 32. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 31, wherein the masking moiety comprises a heavy chain variable region comprising:
- [0530] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);
- [0531] (b) a CDR H2 amino acid sequence of WIN-TETGEPAYADDFKG (SEQ ID NO:21);
- [0532] (c) a CDR H3 amino acid sequence of PYDYPDVL DY (SEQ ID NO:22); and a light chain variable region comprising:
- [0533] (d) a light chain (CDR L)1 amino acid sequence of RASKSVSTSNYSYIH (SEQ ID NO:23);
- [0534] (e) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and
- [0535] (f) a CDR L3 amino acid sequence of QHS-REFPWT (SEQ ID NO:25).
- [0536] 33. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 29, wherein the masking moiety comprises a heavy chain variable region comprising at least one of:
- [0537] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- [0538] (b) a CDR H2 amino acid sequence of IIWGDGSTNYHSALIS (SEQ ID NO:27); and
- [0539] (c) a CDR H3 amino acid sequence of GIT-TVVDDYYAMDY (SEQ ID NO:28).
- [0540] 34. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 29 and 33, wherein the masking moiety comprises a light chain variable region comprising at least one of:
- [0541] (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- [0542] (e) a CDR L2 amino acid sequence of AAT-FLAD (SEQ ID NO:30); and
- [0543] (f) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).
- [0544] 35. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 29 and 33 to 34, wherein the masking moiety comprises a heavy chain variable region comprising:
- [0545] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- [0546] (b) a CDR H2 amino acid sequence of IIWGDGSTNYHSALIS (SEQ ID NO:27);

- [0547] (c) a CDR H3 amino acid sequence of GIT-TVVDDYYAMDY (SEQ ID NO:28); and a light chain variable region comprising:
- [0548] (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- [0549] (e) a CDR L2 amino acid sequence of AAT-FLAD (SEQ ID NO:30); and
- [0550] (f) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).
- [0551] 36. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 35, wherein the masking moiety comprises a heavy chain variable region comprising at least one of:
- [0552] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYYIN (SEQ ID NO:48);
- [0553] (b) a CDR H2 amino acid sequence of VINPDSGGTDYNQNFKG (SEQ ID NO:49); and
- [0554] (c) a CDR H3 amino acid sequence of RDSYGFYD (SEQ ID NO:50).
- [0555] 37. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 36, wherein the masking moiety comprises a light chain variable region comprising at least one of:
- [0556] (a) a light chain (CDR L)1 amino acid sequence of KASLSVTNDVA (SEQ ID NO:51);
- [0557] (b) a CDR L2 amino acid sequence of YASNRNA (SEQ ID NO:52); and
- [0558] (c) a CDR L3 amino acid sequence of QQDYTSPT (SEQ ID NO:53).
- [0559] 38. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 37, wherein the masking moiety comprises a heavy chain variable region comprising:
- [0560] a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYYIN (SEQ ID NO:48);
- [0561] b) a CDR H2 amino acid sequence of VINPDSGGTDYNQNFKG (SEQ ID NO:49); and
- [0562] c) a CDR H3 amino acid sequence of RDSYGFYD (SEQ ID NO:50); and a light chain variable region comprising:
- [0563] d) a light chain (CDR L)1 amino acid sequence of KASLSVTNDVA (SEQ ID NO:51);
- [0564] e) a CDR L2 amino acid sequence of YASNRNA (SEQ ID NO:52); and
- [0565] f) a CDR L3 amino acid sequence of QQDYTSPT (SEQ ID NO:53).
- [0566] 39. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 38, wherein the protease cleavable linker comprises at least one protease recognition sequence.
- [0567] 40. The protease-activatable T cell activating bispecific molecule of embodiment 39, wherein the protease cleavable linker comprises a protease recognition sequence.
- [0568] 41. The protease-activatable T cell activating bispecific molecule of embodiment 40, wherein the protease recognition sequence is selected from the group consisting of:

- (a) (SEQ ID NO: 36)
RQARVVNG;
- (b) (SEQ ID NO: 37)
VHMLGFLGPGSRGSFP;
- (c) (SEQ ID NO: 38)
RQARVVNGXXXXVPLSLYSG;
and
- (d) (SEQ ID NO: 39)
RQARVVNGVPLSLYSG
- (e) (SEQ ID NO: 40)
PLGLWSQ,
wherein X is any amino acid.
- [0569] 42. The protease-activatable T cell activating bispecific molecule of embodiment 40, wherein the protease recognition sequence is selected from the group consisting of:
- (a) (SEQ ID NO: 36)
RQARVVNG;
- (b) (SEQ ID NO: 37)
VHMLGFLGPGSRGSFP;
- (c) (SEQ ID NO: 38)
RQARVVNGXXXXVPLSLYSG;
- (d) (SEQ ID NO: 39)
RQARVVNGVPLSLYSG;
- (e) (SEQ ID NO: 40)
PLGLWSQ;
- (f) (SEQ ID NO: 97)
VHMLGFLGPRQARVVNG;
- (g) (SEQ ID NO: 98)
FVGGTG;
- (h) (SEQ ID NO: 99)
KKAAPVNG;
- (i) (SEQ ID NO: 100)
PMAKKVNG;
- (j) (SEQ ID NO: 101)
QARAKVNG;
- (k) (SEQ ID NO: 102)
VHMLGFLGP;
- (l) (SEQ ID NO: 103)
QARAK;

-continued

(m) (SEQ ID NO: 104)
VHMPGLGFLGPPMAKK;

(n) (SEQ ID NO: 105)
KKAAP;
and

(o) (SEQ ID NO: 106)
PMAKK,
wherein X is any amino acid.

[0570] 43. The protease-activatable T cell activating bispecific molecule of embodiment 39 or 40, wherein the protease cleavable linker comprises the protease recognition sequence RQARVVNG (SEQ ID NO:36).

[0571] 44. The protease-activatable T cell activating bispecific molecule of embodiment 39 or 40, wherein the protease cleavable linker comprises the protease recognition sequence VHMPGLGFLGPRQARVVNG (SEQ ID NO:97).

[0572] 45. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 45, wherein the protease cleavable linker comprises the protease recognition sequence RQARVVNG (SEQ ID NO:36) or the protease recognition sequence VHMPLGFLGPRQARVVNG (SEQ ID NO:97).

[0573] 46. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 45, wherein the protease is selected from the group consisting of metalloproteinase, serine protease, cysteine protease, aspartic proteases, and cathepsin protease.

[0574] 47. The protease-activatable T cell activating bispecific molecule of embodiment 46, wherein the metalloproteinase is a matrix metalloproteinase (MMP), preferably MMP9 or MMP2.

[0575] 48. The protease-activatable T cell activating bispecific molecule of embodiment 46, wherein the serine protease is Matriptase.

[0576] 49. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, wherein the first antigen binding moiety comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 44, SEQ ID NO: 45 and SEQ ID NO: 46 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0577] 50. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 49, wherein the first antigen binding moiety is capable of specific binding to CD3 and comprises a heavy chain variable region comprising:

[0578] a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of TYAMN (SEQ ID NO:44);

[0579] b) a CDR H2 amino acid sequence of RIR-SKYNNYATYYADSVKG (SEQ ID NO:45); and

[0580] c) a CDR H3 amino acid sequence of HGNFGNSYVSWFAY (SEQ ID NO:46); and a light chain variable region comprising:

[0581] d) a light chain (CDR L)1 amino acid sequence of GSSTGAVTTSNYAN (SEQ ID NO:17);

[0582] e) a CDR L2 amino acid sequence of GTNK-RAP (SEQ ID NO:18); and

[0583] f) a CDR L3 amino acid sequence of ALWYSNLWV (SEQ ID NO:19).

[0584] 51. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 50, wherein the first antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0585] 52. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 50, wherein the first antigen binding moiety is capable of specific binding to CD3 and comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.

[0586] 53. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52, wherein the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16 and at least one light chain CDR selected from the group of SEQ ID NO: 17, SEQ ID NO: 18 and SEQ ID NO: 19.

[0587] 54. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 53, wherein the second antigen binding moiety is capable of specific binding to FolR1 and comprises a heavy chain variable region comprising:

[0588] a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of NAWMS (SEQ ID NO:14);

[0589] b) a CDR H2 amino acid sequence of RIK-SKTDGGTTDYAAPVKG (SEQ ID NO:15); and

[0590] c) a CDR H3 amino acid sequence of PWEWSWYDY (SEQ ID NO:16); and a light chain variable region comprising:

[0591] d) a light chain (CDR L)1 amino acid sequence of GSSTGAVTTSNYAN (SEQ ID NO:17);

[0592] e) a CDR L2 amino acid sequence of GTNK-RAP (SEQ ID NO:18); and

[0593] f) a CDR L3 amino acid sequence of ALWYSNLWV (SEQ ID NO:19).

[0594] 55. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 54, wherein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 55.

[0595] 56. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 54, wherein the second antigen binding moiety is capable

- of specific binding to FolR1 and comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 47 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.
- [0596] 57. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52, wherein the second antigen binding moiety is capable of specific binding to FolR1 and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 151, SEQ ID NO: 152 and SEQ ID NO: 153 and at least one light chain CDR selected from the group of SEQ ID NO: 154, SEQ ID NO: 155 and SEQ ID NO: 156.
- [0597] 58. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52 or 57, wherein the second antigen binding moiety is capable of specific binding to FolR1 and comprises a heavy chain variable region comprising:
- [0598] a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYVMH (SEQ ID NO:151);
- [0599] b) a CDR H2 amino acid sequence of IINPSGGSTSYAQKFQG (SEQ ID NO:152); and
- [0600] c) a CDR H3 amino acid sequence of SFFTGFHLDY (SEQ ID NO:153); and a light chain variable region comprising:
- [0601] d) a light chain (CDR L)1 amino acid sequence of RASQSVSSSYLA (SEQ ID NO:154);
- [0602] e) a CDR L2 amino acid sequence of GASSRAT (SEQ ID NO:155); and
- [0603] f) a CDR L3 amino acid sequence of QQYT-NEHYT (SEQ ID NO:156).
- [0604] 59. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52, 57 or 58, wherein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 158.
- [0605] 60. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52 or 57 to 59, wherein the second antigen binding moiety wherein the second antigen binding moiety is capable of specific binding to ForR1 and comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 157 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 158.
- [0606] 61. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52, wherein the second antigen binding moiety is capable of specific binding to Mesothelin and comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 107, SEQ ID NO: 108 and SEQ ID NO: 109 and at least one light chain CDR selected from the group of SEQ ID NO: 110, SEQ ID NO: 111 and SEQ ID NO: 112.
- [0607] 62. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52 or 61, wherein the second antigen binding moiety is capable of specific binding to Mesothelin and comprises a heavy chain variable region comprising:
- [0608] a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of GYTMN (SEQ ID NO:107);
- [0609] b) a CDR H2 amino acid sequence of LITPYNGASSYNQKFRG (SEQ ID NO:108); and
- [0610] c) a CDR H3 amino acid sequence of GGYDGRGFDY (SEQ ID NO:109); and a light chain variable region comprising:
- [0611] d) a light chain (CDR L)1 amino acid sequence of SASSSVSYM (SEQ ID NO:110);
- [0612] e) a CDR L2 amino acid sequence of DTSK-LAS (SEQ ID NO:111); and
- [0613] f) a CDR L3 amino acid sequence of QQWSKHPLT (SEQ ID NO:112).
- [0614] 63. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52, 61 or 62, wherein the second antigen binding moiety comprises a heavy chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 114.
- [0615] 64. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52 or 61 to 63, wherein the second antigen binding moiety is capable of specific binding to Mesothelin and comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 113 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 114.
- [0616] 65. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52, wherein the second antigen binding moiety comprises a heavy chain comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 32 and a light chain comprising an amino acid sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to the amino acid sequence of SEQ ID NO: 33.
- [0617] 66. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 52 or 65, wherein the second antigen binding moiety is capable of specific binding to HER1 and comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 115 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 116.
- [0618] 67. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0619] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:2;
- [0620] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and

- [0621] (c) a light chain comprising an amino acid sequence of SEQ ID NO:1.
- [0622] 68. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0623] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:4;
- [0624] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and
- [0625] (c) a light chain comprising an amino acid sequence of SEQ ID NO:1.
- [0626] 69. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0627] (a) at least one heavy chain comprising the amino acid sequence of SEQ ID NO:32;
- [0628] (b) at least one light chain comprising the amino acid sequence of SEQ ID NO:34.
- [0629] 70. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0630] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:72;
- [0631] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and
- [0632] (c) a light chain comprising an amino acid sequence of SEQ ID NO:1.
- [0633] 71. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0634] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:85;
- [0635] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3; and
- [0636] (c) a light chain comprising an amino acid sequence of SEQ ID NO:1.
- [0637] 72. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0638] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:73;
- [0639] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:3;
- [0640] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:1; and
- [0641] (d) a second light chain comprising an amino acid sequence of SEQ ID NO: 74.
- [0642] 73. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0643] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:77;
- [0644] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:82;
- [0645] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:78; and
- [0646] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:81.
- [0647] 74. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0648] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:76;
- [0649] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:77;
- [0650] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:78; and
- [0651] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:79.
- [0652] 75. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0653] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:132;
- [0654] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:136;
- [0655] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:81; and
- [0656] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:133.
- [0657] 76. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 48, comprising
- [0658] (a) a first heavy chain comprising the amino acid sequence of SEQ ID NO:137;
- [0659] (b) a second heavy chain comprising the amino acid sequence of SEQ ID NO:139;
- [0660] (c) a first light chain comprising an amino acid sequence of SEQ ID NO:81; and
- [0661] (d) a second light chain comprising an amino acid sequence of SEQ ID NO:138.
- [0662] 77. An idiotype-specific polypeptide for reversibly concealing an anti-CD3 antigen binding site of a molecule.
- [0663] 78. The idiotype-specific polypeptide of embodiment 77, wherein the idiotype-specific polypeptide is an anti-idiotype scFv.
- [0664] 79. The idiotype-specific polypeptide of embodiment 77 or 78, wherein the idiotype-specific polypeptide is covalently attached to the molecule through a linker.
- [0665] 80. The idiotype-specific polypeptide of embodiment 79, wherein the linker is a peptide linker.
- [0666] 81. The idiotype-specific polypeptide of embodiment 79 or 80, wherein the linker is a protease-cleavable linker.
- [0667] 82. The idiotype-specific polypeptide of any one of embodiments 79 to 81, wherein the peptide linker comprises at least one protease recognition site.
- [0668] 83. The idiotype-specific polypeptide of embodiment 82, wherein the protease is selected from the group consisting of metalloproteinase, serine protease, cysteine protease, aspartic proteases, and cathepsin protease.
- [0669] 84. The idiotype-specific polypeptide of embodiment 83, wherein the metalloproteinase is a matrix metalloproteinase (MMP), preferably MMP9 or MMP2.
- [0670] 85. The idiotype-specific polypeptide of embodiment 83, wherein the serine protease is Matriptase.
- [0671] 86. The idiotype-specific polypeptide of embodiment 82, wherein the protease cleavable linker comprises the protease recognition sequence RQARVVNG (SEQ ID NO:36) or the protease recognition sequence VHMPLGFLGPRQARVVNG (SEQ ID NO:97).

- [0672] 87. The idiotype-specific polypeptide of any one of embodiments 77 to 86, wherein the molecule is a T-cell activating bispecific molecule.
- [0673] 88. The idiotype-specific polypeptide of any one of embodiments 77 to 87, comprising a heavy chain variable region comprising at least one of:
- [0674] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);
- [0675] (b) a CDR H2 amino acid sequence of WINTETGEPAYADDFKG (SEQ ID NO:21); and
- [0676] (c) a CDR H3 amino acid sequence of PYDYDVLVDY (SEQ ID NO:22).
- [0677] 89. The idiotype-specific polypeptide of any one of embodiments 77 to 88, comprising a light chain variable region comprising at least one of:
- [0678] (d) a light chain (CDR L)1 amino acid sequence of RASKSVTSNYSYIH (SEQ ID NO:23);
- [0679] (e) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and
- [0680] (f) a CDR L3 amino acid sequence of QHSREFPWT (SEQ ID NO:25).
- [0681] 90. The idiotype-specific polypeptide of any one of embodiments 77 to 87, comprising a heavy chain variable region comprising:
- [0682] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of DYSIH (SEQ ID NO:20);
- [0683] (b) a CDR H2 amino acid sequence of WINTETGEPAYADDFKG (SEQ ID NO:21);
- [0684] (c) a CDR H3 amino acid sequence of PYDYDVLVDY (SEQ ID NO:22); and a light chain variable region comprising:
- [0685] (d) a light chain (CDR L)1 amino acid sequence of RASKSVTSNYSYIH (SEQ ID NO:23);
- [0686] (e) a CDR L2 amino acid sequence of YVSYLES (SEQ ID NO:24); and
- [0687] (f) a CDR L3 amino acid sequence of QHSREFPWT (SEQ ID NO:25).
- [0688] 91. The idiotype-specific polypeptide of any one of embodiments 77 to 87, comprising a heavy chain variable region comprising at least one of:
- [0689] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- [0690] (b) a CDR H2 amino acid sequence of IIWGDGSTNYHSALIS (SEQ ID NO:27); and
- [0691] (c) a CDR H3 amino acid sequence of GITTVVDDYYAMDY (SEQ ID NO:28).
- [0692] 92. The idiotype-specific polypeptide of any one of embodiments 77 to 87 and 91, comprising a light chain variable region comprising at least one of:
- [0693] (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- [0694] (e) a CDR L2 amino acid sequence of AATFLAD (SEQ ID NO:30); and
- [0695] (f) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).
- [0696] 93. The idiotype-specific polypeptide of any one of embodiments 77 to 87, comprising a heavy chain variable region comprising:
- [0697] (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- [0698] (b) a CDR H2 amino acid sequence of IIWGDGSTNYHSALIS (SEQ ID NO:27);
- [0699] (c) a CDR H3 amino acid sequence of GITTVVDDYYAMDY (SEQ ID NO:28); and a light chain variable region comprising:
- [0700] (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- [0701] (e) a CDR L2 amino acid sequence of AATFLAD (SEQ ID NO:30); and
- [0702] (f) a CDR L3 amino acid sequence of QHYYSTPYT (SEQ ID NO:31).
- [0703] 94. The idiotype-specific polypeptide of embodiments 77 to 93, wherein the anti-CD3 antigen binding site comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.
- [0704] 95. An isolated polynucleotide encoding the protease-activatable T cell activating bispecific antigen binding molecule of any one of embodiments 1 to 76 or the idiotype-specific polypeptide of any one of embodiments 77 to 94.
- [0705] 96. A polypeptide encoded by the polynucleotide of embodiment 95.
- [0706] 97. A vector, particularly an expression vector, comprising the polynucleotide of embodiment 95.
- [0707] 98. A host cell comprising the polynucleotide of embodiment 95 or the vector of embodiment 97.
- [0708] 99. A method of producing a protease-activatable T cell activating bispecific molecule, comprising the steps of a) culturing the host cell of embodiment 98 under conditions suitable for the expression of the protease-activatable T cell activating bispecific molecule and b) recovering the protease-activatable T cell activating bispecific molecule.
- [0709] 100. A protease-activatable T cell activating bispecific molecule produced by the method of embodiment 99.
- [0710] 101. A method of producing an idiotype-specific polypeptide, comprising the steps of a) culturing the host cell of embodiment 98 under conditions suitable for the expression of the idiotype-specific polypeptide and b) recovering the an idiotype-specific polypeptide.
- [0711] 102. An idiotype-specific polypeptide produced by the method of embodiment 101.
- [0712] 103. A pharmaceutical composition comprising the protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 76 and a pharmaceutically acceptable carrier.
- [0713] 104. A pharmaceutical composition comprising the idiotype-specific polypeptide of any one of embodiments 77 to 94 and a pharmaceutically acceptable carrier.
- [0714] 105. A protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 76, the idiotype-specific polypeptide of any one of embodiments 77 to 94 or the composition of embodiment 103 for use as a medicament.
- [0715] 106. The protease-activatable T cell activating bispecific molecule for use according to embodiment 105, wherein the medicament is for treating or delaying

- progression of cancer, treating or delaying progression of an immune related disease, or enhancing or stimulating an immune response or function in an individual.
- [0716] 107. The protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 76 or the idiotype-specific polypeptide of any one of embodiments 77 to 94 for use in the treatment of a disease in an individual in need thereof.
- [0717] 108. The protease-activatable T cell activating bispecific molecule or the idiotype-specific polypeptide for use in the treatment of a disease in an individual in need thereof of embodiment 107, wherein the disease is a cancer.
- [0718] 109. Use of the protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 76 or the idiotype-specific polypeptide of any one of embodiments 77 to 94 for the manufacture of a medicament.
- [0719] 110. The use of embodiment 109, wherein the disease is a cancer.
- [0720] 111. A method of treating a disease in an individual, comprising administering to said individual a therapeutically effective amount of a composition comprising the protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 76 or composition of embodiment 103.
- [0721] 112. A method for inducing lysis of a target cell, comprising contacting a target cell with the protease-activatable T cell activating bispecific molecule of any one of embodiments 1 to 76 or composition of embodiment 103 in the presence of a T cell.
- [0722] 113. The method of embodiment 112 wherein the target cell is a cancer cell.
- [0723] 114. The method of embodiment 112 or 113, wherein the target cell expresses a protease capable of activating the protease-activatable T cell activating bispecific molecule.
- [0724] 115. An anti-idiotype CD3 antibody or antigen-binding fragment thereof specific for an idiotype of an anti-CD3 antigen-binding molecule, wherein the anti-idiotype CD3 antibody or fragment thereof when bound to the anti-CD3 antigen-binding molecule specifically blocks binding of the anti-CD3 antigen-binding molecule to CD3.
- [0725] 116. The anti-idiotype CD3 antibody or antigen-binding fragment thereof of embodiment 115, wherein the anti-idiotype CD3 antibody or fragment thereof is reversibly associated with the anti-CD3 antigen-binding molecule through a peptide linker comprising a protease recognition site.
- [0726] 117. The anti-idiotype CD3 antibody or antigen-binding fragment thereof of embodiment 115 or 116, wherein the CD3 is a mouse, monkey or human CD3.
- [0727] 118. A method of reducing in vivo toxicity of a T cell activating bispecific molecule comprising attaching an idiotype-specific polypeptide of any one of embodiments 77 to 94 to the T cell activating bispecific molecule with a protease-cleavable linker to form a protease-activatable T cell activating bispecific molecule, wherein the in vivo toxicity of the protease-activatable T cell activating bispecific molecule is reduced compared to toxicity of the T cell activating bispecific molecule.
- [0728] 119. The invention as described hereinbefore.

EXAMPLES

[0729] The following are examples of methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

Example 1

Synthesis of Monovalent Anti-CD3 IgG Molecules with Anti-Idiotypic scFv

[0730] Described herein are CD3 binders that are masked with an N-terminally linked anti-idiotypic CD3 scFv. These constructs include a protease recognition site which is recognized by a tumor specific protease. In the presence of protease-expressing tumor cells, the linker connecting the masking moiety will be cleaved and, thereby, CD3 binding by the CD3 binder is recovered. Several monovalent anti-CD3 IgG molecules with various anti-idiotypic scFv were produced and are schematically depicted in FIGS. 1A-1E with their respective ID number. The following molecules were prepared:

- [0731] Identification No. 7859: monovalent CD3 IgG, (anti-idiotypic scFv 4.15.64—MK062 Matriptase site—CD3—N-terminal fused to CD3 Fab—inert Fc) with N-terminal fused anti CD3 scFv 4.15.64 and protease-cleavable linker.
- [0732] Identification No. 7860: monovalent CD3 IgG, (anti-idiotypic scFv 4.32.63—MK062 Matriptase site—CD3—N-terminal fused to CD3 Fab—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and protease-cleavable linker.
- [0733] Identification No. 7857: monovalent CD3 IgG, (anti-idiotypic scFv 4.15.64—non-cleavable linker—CD3—N-terminal fused to CD3 Fab—inert Fc) with N-terminal fused anti CD3 scFv 4.15.64 and protease-cleavable linker.
- [0734] Identification No. 7858: monovalent CD3 IgG, (anti-idiotypic scFv 4.32.63—non-cleavable linker—CD3—N-terminal fused to CD3 Fab—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and protease-cleavable linker.
- [0735] Identification No. 7861: monovalent CD3 IgG, (CD3 Fab—inert Fc) with N-terminal fused anti CD3 scFv 4.15.64/4.32.63 and protease linker.
- [0736] Anti-idiotypic (ID) binder sequences were obtained by RACE-PCR (rapid amplification of cDNA ends) from RNA of Hybridoma cells. Hybridoma cells were obtained by immunization of mice with CH2527 (VL₇-46 (13)/VH₃-23(12)) Fab-fragment. Single chain Fv (scFv) sequence synthesis was ordered from Invitrogen including the necessary restriction sites for cloning. Six different anti-idiotypic CH2527 binders were compared for their affinities (FIG. 2, result Biacore-Analytics (AG M. Schröml) at 25° C./37° C. (Analyt: MAK<CEA/CD3>rH)) and two of them were cloned as N-terminal fusions at the heavy chain of CD3 Fab-Fc.

[0737] The anti-ID single chain Fv DNA sequences were subcloned in frame with the CD3 VH chain pre-inserted into the respective recipient mammalian expression vector. Protein expression is driven by an MPSV promoter and a synthetic polyA signal sequence is present at the 3' end of the CDS. In addition each vector contains an EBV OriP sequence.

[0738] The molecules were produced by co-transfecting HEK293-EBNA cells growing in suspension with the mammalian expression vectors using polyethylenimine (PEI). The cells were transfected with the corresponding expression vectors in a 1:1:2 ratio (“Fc hole (CH2-CH3)”：“common light chain (CLC)”：“vector heavy chain knob (scFv-VH-CH1-CH2-CH3)”).

[0739] For transfection, HEK293 EBNA cells were cultivated in serum free ExCell culture medium containing 6 mM L-glutamine and 250 mg/l G418. For the production in 600 ml tubespin flasks (max. working volume 400 mL) 800 million HEK293 EBNA cells were seeded 24 hours before transfection without G418. For transfection 800 mio cells were centrifuged for 5 min at 210xg and supernatant was replaced by 40 ml pre-warmed CD CHO medium containing 6 mM L-Glutamine. Expression vectors were mixed with 40 ml CD CHO medium containing 6 mM L-Glutamine to a total amount of 400 µg DNA. After addition of 1080 µl PEI solution (2.7 µg/ml) the mixture was vortexed for 15 s and subsequently incubated for 10 min at room temperature. Afterwards cells were mixed with the DNA/PEI solution, transferred to a 600 ml tubespin flask and incubated for 3 hours at 37° C. in an incubator with a 5% CO2 atmosphere. After incubation, 320 ml ExCell+6 mM L-glutamine+5 g/L

(GE Healthcare) equilibrated with 20 mM histidine, 140 mM sodium chloride, pH 6.0, 0.01% Tween20.

[0741] For analytics after size exclusion chromatography the purity and molecular weight of the molecules in the single fractions were analyzed by SDS-PAGE in the absence of a reducing agent and staining with Coomassie (InstantBlue™, Expedeon). The NuPAGE® Pre-Cast gel system (4-12% Bis-Tris, Invitrogen or 3-8% Tris-Acetate, Invitrogen) was used according to the manufacturer’s instruction. The protein concentration of purified protein samples was determined by measuring the optical density (OD) at 280 nm divided by the molar extinction coefficient calculated on the basis of the amino acid sequence.

[0742] Purity and molecular weight of the molecules after the final purification step were analyzed by CE-SDS analyses in the presence and absence of a reducing agent. The Caliper LabChip GXII system (Caliper Lifescience) was used according to the manufacturer’s instruction. The aggregate content of the molecules was analyzed using a TSKgel G3000 SW XL analytical size-exclusion column (Tosoh) in 25 mM K2HPO4, 125 mM NaCl, 200 mM L-arginine monohydrochloride, 0.02% (w/v) NaN3, pH 6.7 running buffer at 25° C. The final quality of all molecules was good, with ≥92% monomer content.

TABLE 2

Summary of production and purification of protease activated monovalent CD3 IgG molecules.			
Molecule	Titer [mg/l]	Yield [mg/l]	Analytical SEC (HMW/Monomer/LMW) [%]
1	12	3.38	2.21/95.5/2.29
2	9	1.75	4.86/95.14/0
3	15	4.8	6.93/93.07/0
4	4.5	0.26	4.88/95.12/0
5	105.3	26.3	0/100/0

Pepsoy+1.0 mM VPA+3 g/l glucose medium was added and cells were cultivated for 24 hours prior to feeding with 7% Feed 7. After 6-7 days cultivation supernatant was collected for purification by centrifugation for 20-30 min at 210xg (Sigma 8K centrifuge). The solution was sterile filtered (0.22 µm filter) and sodium azide in a final concentration of 0.01% w/v was added. The solution was kept at 4° C. until purification. The secreted protein was purified from cell culture supernatants by affinity chromatography using ProteinA affinity chromatography, followed by one to two size exclusion chromatographic steps.

[0740] For affinity chromatography supernatant was loaded on a HiTrap Protein A FF column (CV=5 mL, GE Healthcare) equilibrated with 25 ml 20 mM sodium phosphate, 20 mM sodium citrate, 0.5M sodium chloride, 0.01% Tween-20 pH 7.5. Unbound protein was removed by washing with at least 10 column volumes 20 mM sodium phosphate, 20 mM sodium citrate, 0.5M sodium chloride, 0.01% Tween-20 pH 7.5 and target protein was eluted in 20 column volumes (gradient from 0%-100%) 20 mM sodium citrate, 0.5M sodium chloride, 0.01% Tween-20 pH 2.5. Protein solution was neutralized by adding 1/10 of 2 M Tris pH 10.5. Target protein was concentrated with Amicon® Ultra-15 Ultracel 30K (Merck Millipore Ltd.) to a volume of 4 ml maximum prior loading on a HiLoad Superdex 200 column

Example 2

Cleavage and Stability of Protease Activated IgGs

[0743] Capillary Electrophoresis of protease activated IgG molecules. Comparison of untreated sample and treated sample showed that the anti-ID scFv was completely cleaved off after treatment with rhMatriptase/ST14 (R&D Systems) indicated by the size shift in the SDS page analysis (FIG. 3). Analysis of samples incubated for 48 h at 37° C. confirmed stability of the molecules in formulation buffer (FIG. 3A-D).

Example 3

Masking Effect of Anti-Idiotypic scFv for CD3 IgG

[0744] The efficiency of masking the CD3 binder by N-terminal fusion of an anti-idiotypic CD3 scFv was shown by a Jurkat-NFAT reporter assay. Jurkat-NFAT reporter cells (a human acute lymphatic leukemia reporter cell line with a NFAT promoter-regulated luciferase expression, GloResponse Jurkat NFAT-RE-luc2P, Promega #CS176501) express active firefly luciferase if the NFAT promoter is activated by binding of CD3ε. The intensity of the luminescence signal upon addition of luciferase substrate is proportional to the intensity of CD3 activation and signaling.

Completely unmasked monovalent CD3 molecules served as a positive control. The treatment was done with rhMatriptase/ST14 (R&D Systems) for 48 h at 37° C. In parallel Bug/ml Anti human Fc Antibody (BioLegends) were coated in 0.025 ul/well PBS for 48 h at 4° C. in white-walled, clear bottom 96-well (flat)-plate (Greiner Bio-One). PBS was removed by pipetting before monovalent IgGs were added at the indicated concentration range of 200 nM-2.56 pM. Plates were incubated for about 30 min at 4° C. Subsequently, Jurkat-NEAT reporter cells were harvested and viability assessed using ViCell. Cells were resuspended in Jurkat medium (RPMI1640, 2 g/l Glucose, 2 g/l NaHCO₃, 10% FCS, 25 mM HEPES, 2 mM L-Glutamin, 1xNEAA, 1xSodium-pyruvate) without Hygromycine and 100 µl per well (25.000 cells/well) were added to the crosslinked monovalent CD3 IgGs. Cells were incubated for 3 h at 37° C. in a humidified incubator. Plates were taken out of the incubator for about 10 min to adapt to room temperature prior to Luminescence read out. 100 µl/well of ONE-Glo solution (1:1 ONE-Glo and assay medium volume per well) were added to wells and incubated for 10 min at room temperature in the dark. Luminescence was detected using WALLAC Victor3 ELISA reader (PerkinElmer2030), 1 sec/well as detection time. 7857 (4.15.64 mask with non-cleavable linker) and 7859 (untreated) show significantly reduced CD3ε binding compared to unmasked (7861) and pretreated molecule (7859 treated) (FIG. 4A). 7760 was included as a control to show that N-terminal linkage does not block CD3 binding itself. 7858 (4.32.63 mask with non-cleavable linker) and 7860 (untreated) show significantly reduced CD3ε binding compared to unmasked (7861) and pretreated molecule (7860 treated) (FIG. 4B). In line with the affinities of the anti-idiotypic CD3 binders the 4.32.63 mask is much more efficient than the 4.15.64. In terms of EC50 values (FIG. 4C) the 4.32.63 masked CD3 binder binds 54 fold less than the unmasked CD3 binder 7861. For the 4.15.64 mask it is only 16 fold less binding than for 7861. Depending on the tumor target and the target binder the best mask can be evaluated.

Example 4

Preparation of Anti FoIR1/Anti-CD3 T Cell Bispecific (TCB) Molecules with Anti CD3 scFv

[0745] Several T cell bispecific (TCB) molecules with various anti-idiotypic scFv were produced and are schematically depicted in FIGS. 5A-5H with their respective ID number. The following molecules were prepared:

[0746] ID7344: FoIR1 16D5 2+1 IgG, classic format (anti-idiotypic scFv 4.15.64—MK062 Matriptase site—CD3—N-terminal fused to FoIR1 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.15.64 and protease linker (FIG. 5A, SEQ ID NOs 1, 2 and 3).

[0747] ID7496: FoIR1 16D5 2+1 IgG, classic format (anti-idiotypic scFv 4.32.63—MK062 Matriptase site—CD3—N-terminal fused to FoIR1 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and protease linker (FIG. 5C, SEQ ID NOs 1, 3 and 4).

[0748] ID7676: FoIR1 16D5 2+1 IgG, classic format (anti-idiotypic scFv 4.15.64—non-cleavable GS linker—CD3—N-terminal fused to FoIR1 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.15.64 and protease linker (FIG. 5B, SEQ ID NOs 1, 3 and 6).

[0749] ID7611: FoIR1 16D5 2+1 IgG, classic format (anti-idiotypic scFv 4.32.63—non-cleavable GS linker—CD3—N-terminal fused to FoIR1 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and protease linker (FIG. 5D, SEQ ID NOs 1, 3 and 5).

[0750] Anti-idiotypic (ID) binder sequences were obtained by RACE-PCR (rapid amplification of cDNA ends) from RNA of Hybridoma cells. Hybridoma cells were obtained by immunization of mice. Single chain Fv (ScFv) sequence synthesis was ordered at Invitrogen including the necessary restriction sites for cloning. Six different anti-idiotypic CH2527 binders were compared for their affinities (FIG. 2, result Biacore-Analytics (AG M. Schröml) at 25° C./37° C. (Analyt: MAK<CEA/CD3>rH)) and four of them were cloned as N-terminal fusions at the HC of CD3-FoIR1 16D5 TCB.

[0751] The anti-ID single chain Fv DNA sequences were subcloned in frame with the CD3 VH chain pre-inserted into the respective recipient mammalian expression vector. Protein expression is driven by an MPSV promoter and a synthetic polyA signal sequence is present at the 3' end of the coding sequence (CDS). In addition each vector contains an EBV OriP sequence.

[0752] The molecules were produced by co-transfecting HEK293-EBNA cells growing in suspension with the mammalian expression vectors using polyethylenimine (PEI). The cells were transfected with the corresponding expression vectors in a 1:3:2 ratio ("vector heavy chain hole (VH-CH1-CH2-CH3)": "common light chain (CLC)": "vector heavy chain knob (scFv-VH-CH1-VH-CH1-CH2-CH3)").

[0753] For transfection HEK293 EBNA cells were cultivated in serum free ExCell culture medium containing 6 mM L-glutamine and 250 mg/l G418. For the production in 600 ml tubespin flasks (max. working volume 400 mL) 800 million HEK293 EBNA cells were seeded 24 hours before transfection without G418. For transfection 800 mio cells were centrifuged for 5 min at 210xg and supernatant was replaced by 40 ml pre-warmed CD CHO medium containing 6 mM L-Glutamine. Expression vectors were mixed with 40 ml CD CHO medium containing 6 mM L-Glutamine to a total amount of 400 µg DNA. After addition of 1080 µl PEI solution (2.7 µg/ml) the mixture was vortexed for 15 s and subsequently incubated for 10 min at room temperature. Afterwards cells were mixed with the DNA/PEI solution, transferred to a 600 ml tubespin flask and incubated for 3 hours at 37° C. in an incubator with a 5% CO₂ atmosphere. After incubation, 320 ml ExCell+6 mM L-glutamine+5 g/L Pepsoy+1.0 mM VPA+3 g/l glucose medium was added and cells were cultivated for 24 hours prior to feeding with 7% Feed 7. After 6-7 days cultivation supernatant was collected for purification by centrifugation for 20-30 min at 210xg (Sigma 8K centrifuge). The solution was sterile filtered (0.22 µm filter) and sodium azide in a final concentration of 0.01% w/v was added. The solution was kept at 4° C. until purification.

[0754] The secreted protein was purified from cell culture supernatants by affinity chromatography using ProteinA affinity chromatography, followed by one to two size exclusion chromatographic steps. For affinity chromatography supernatant was loaded on a HiTrap Protein A FF column (CV=5 mL, GE Healthcare) equilibrated with 25 ml 20 mM sodium phosphate, 20 mM sodium citrate, 0.5M sodium chloride, 0.01% Tween-20 pH 7.5. Unbound protein was

removed by washing with at least 10 column volumes 20 mM sodium phosphate, 20 mM sodium citrate, 0.5M sodium chloride, 0.01% Tween-20 pH 7.5 and target protein was eluted in 20 column volumes (gradient from 0%-100%) 20 mM sodium citrate, 0.5M sodium chloride, 0.01% Tween-20 pH 2.5. Protein solution was neutralized by adding 1/10 of 2 M Tris pH 10.5. Target protein was concentrated with Amicon® Ultra-15 Ultracel 30K (Merck Millipore Ltd.) to a volume of 4 ml maximum prior loading on a HiLoad Superdex 200 column (GE Healthcare) equilibrated with 20 mM histidine, 140 mM sodium chloride, pH 6.0, 0.01% Tween20.

[0755] For analytics after size exclusion chromatography the purity and molecular weight of the molecules in the single fractions were analyzed by SDS-PAGE in the absence of a reducing agent and staining with Coomassie (InstantBlue™, Expedeon). The NuPAGE® Pre-Cast gel system (4-12% Bis-Tris, Invitrogen or 3-8% Tris-Acetate, Invitrogen) was used according to the manufacturer's instruction. The protein concentration of purified protein samples was determined by measuring the optical density (OD) at 280 nm divided by the molar extinction coefficient calculated on the basis of the amino acid sequence.

[0756] Purity and molecular weight of the molecules after the final purification step were analyzed by CE-SDS analyses in the presence and absence of a reducing agent. The Caliper LabChip GXII system (Caliper Lifescience) was used according to the manufacturer's instruction.

[0757] The aggregate content of the molecules was analyzed using a TSKgel G3000 SW XL analytical size-exclusion column (Tosoh) in 25 mM K2HPO4, 125 mM NaCl, 200 mM L-arginine monohydrochloride, 0.02% (w/v) NaN3, pH 6.7 running buffer at 25° C. The final quality of all molecules was good, with ≥92% monomer content.

TABLE 2

Summary of production and purification of protease activated TCB molecules.			
Molecule	Titer [mg/l]	Yield [mg/l]	Analytical SEC (HMW/Monomer/LMW) [%]
1	33	3.7	0.98/92.7/6.32
2	11	0.55	3.76/96.24/0
3	12.9	0.89	2.9/93.82/2.19
4	6.7	0.35	4.59/95.41/0

Example 5

Transient Expression of Protease Activated TCBs

[0758] Different plasmid ratios used for transfection were compared by size exclusion chromatography as the knob chain was suspected to be expressed in lower levels compared to the hole chain and the light chain. As shown in FIGS. 6 and 7, using a plasmid ratio of 1(hole):2 (knob):3 (CLC) (FIG. 7) instead of 1(hole):1 (knob):3 (CLC) (FIG. 6) increased the yield of correct molecule (left peak) and decreased the amount of hole hole homodimers (right peak).

Example 6

Cleavage and Stability of Protease Activated TCB

[0759] Protease activated TCBs were analyzed by capillary electrophoresis. Comparison of untreated sample and

treated sample showed that the anti-idiotypic scFv moiety was completely cleaved off after treatment with rhMatriptase/ST14. Analysis of samples incubated for 48 h at 37° C. confirmed stability of the molecules in formulation buffer (FIGS. 12A-12D).

Example 7

Cell Killing Using Target Cell Lines that Express Different Levels of FolR1

[0760] T-cell-mediated cell killing induced by protease activated TCB molecules was assessed using target cell lines expressing different levels of FolR1 (FIG. 13). Human PBMCs were used as effector cells and cell killing was detected at 48 h of incubation with the protease activated TCB molecules. Human Peripheral blood mononuclear cells (PBMCs) were isolated from fresh taken blood or from buffy coats obtained from healthy human donors. For fresh blood 50 ml Leucosep tubes (GreinerBioOne) were used for preparation. For enriched lymphocyte preparations (buffy coats) Histopaque-1077 density preparation was used. Blood/buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature), the plasma above the PBMC-containing interphase was discarded and PBMCs transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant discarded and the PBMC pellet resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation at 37° C. for about 2-3 minutes the tubes were filled with sterile PBS to 50 ml and centrifuged at 350×g for 10 minutes. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 2% FCS and 1× GlutaMax at 37° C., 5% CO₂ in cell incubator until further use. Briefly, adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended at 0.4×10⁶ cells/ml in assay medium (RPMI1640, 2% FCS, 1× GlutaMax). Target cells were plated at a density of 20 000 cells/well using round-bottom 96-well plates. For the killing assay, the molecules were added at the indicated concentrations in triplicates. FolR1 16D5 TCB was included as positive control and an untargeted TCB molecule (binding to CD3 but not to a target cell antigen) was included as negative control. PBMCs were added to target cells at final E:T ratio of 10:1. Target cell killing was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 1 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB.

[0761] The results (FIGS. 14A, 15A, 16, 17, 18A, 19A and 20A) show that the protease activated TCB with anti-idiotypic CD3 scFv moiety N-terminally linked by a non-cleavable linker (#7676 and #7611, FIGS. 5B and D, respectively) were able to significantly reduce cell lysis on Skov3 and HT29 cells. #7611 (FIG. 5D) led to reduced killing on Hela cells while anti-idiotypic CD3 scFv 4.15.64 in #7676 (FIG. 5B) was less efficient in reduction of cell lysis. This is in line with affinities of the anti-idiotypic CD3 scFv N moiety. The higher affinity scFv moiety masks more efficiently. Comparable potency of treated and untreated TCBs

suggests Matriptase expression of Hela and Skov3 cells. Expression of Matriptase seems to be lower in HT29 cells. Treatment of Mkn-45, a FolR1 negative cell line, shows only weak killing with all molecules used herein (FIG. 15A).

Example 8

T-Cell Activation after Co-Incubation of Tumor Cell Lines with Human PBMCs

[0762] T-cell activation mediated by protease activated TCB molecules was assessed on Hela, Skov3 and HT29 cells. Human PBMCs were used as effector cells and the T cell activation was detected at 48 h of incubation with target cells and the antibodies. Target cells were plated at a density of 20 000 cells/well using round-bottom 96-well plates. Molecules were added at the indicated concentrations in triplicates. FolR1 16D5 TCB was included as positive control and an untargeted TCB molecule (binding to CD3 but not to a target cell antigen) was included as negative control. PBMCs were added to target cells at final E:T ratio of 10:1. T-cell activation was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of CD25 and CD69 on CD4 positive and CD8 positive T cells. T cell activation results are consistent with the results observed in the previous example assessing target cell killing (Example 7).

Example 9

T-Cell Activation Mediated by Protease-Activated TCBs and Target Cell Lines Expressing Low Antigen Levels

[0763] T-cell activation mediated by protease activated TCB molecules was assessed on HT29 cells expressing only low levels of FolR1 (FIG. 13). Human PBMCs isolated from buffy coat were used as effector cells. For enriched lymphocyte preparations (buffy coats) Histopaque-1077 density preparation was used. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature), the plasma above the PBMC-containing interphase was discarded and PBMCs transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant discarded and the PBMC pellet resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation at 37° C. for about 2-3 minutes the tubes were filled with sterile PBS to 50 ml and centrifuged at 350×g for 10 minutes. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 2% FCS and 1× GlutaMax at 37° C., 5% CO₂ in cell incubator until further use. Briefly, adherent target cells were harvested with Trypsin/EDTA, counted, assessed for viability and resuspended at 0.4×10⁶ cells/ml in assay medium (RPMI1640, 2% FCS, 1× GlutaMax). Target cells were plated at a density of 20 000 cells/well using round-bottom 96-well plates. Molecules were added at the indicated concentrations in triplicates. FolR1 16D5 TCB was included as positive control and an untargeted TCB molecule (binding to CD3 but not to a target cell antigen) was included as negative control. PBMCs were added to target cells at final E:T ratio of 10:1. T-cell activation was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of CD25 and CD69 on CD4 positive and CD8-positive T

cells. The potency of treated protease activated TCB is comparable to 16D5 TCB (6298). The 16D5 TCB (inverted format) show higher potency than the classic format. Masked TCBs with non-cleavable linker or without Matriptase pre-treatment do not induce T cell activation on this cell line. For cell lines with low or medium FolR1 expression levels both anti-idiotypic scFvs are sufficient in masking the CD3 Fab (FIGS. 22A and B).

Example 10

T-Cell Activation Mediated by Protease Activated TCB with Primary Cell Line HRCEpiC

[0764] T-cell activation mediated by protease activated TCB molecules was assessed on primary Human Renal Cortical Epithelial Cell (ScienceCell) cells expressing only very little amounts of FolR1 (FIG. 13). Human PBMCs isolated from buffy coat were used as effector cells. For enriched lymphocyte preparations (buffy coats) Histopaque-1077 density preparation was used. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, without break at room temperature), the plasma above the PBMC-containing interphase was discarded and PBMCs transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant discarded and the PBMC pellet resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation at 37° C. for about 2-3 minutes the tubes were filled with sterile PBS to 50 ml and centrifuged at 350×g for 10 minutes. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 2% FCS and 1×GlutaMax at 37° C., 5% CO₂ in cell incubator until further use. Briefly, adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended at 0.4×10⁶ cells/ml in assay medium (RPMI1640, 2% FCS, 1×GlutaMax). Target cells were plated at a density of 20 000 cells/well using round-bottom 96-well plates. Protease activatable TCB molecules were added at the indicated concentrations in triplicates. FolR1 16D5 TCB was included as positive control and an untargeted TCB molecule (binding to CD3 but not to a target cell antigen) was included as negative control. PBMCs were added to target cells at final E:T ratio of 10:1. T-cell activation was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of CD25 and CD69 on CD4 positive and CD8 positive T cells. Masked 16D5 TCB does not induce T cell activation upon incubation with primary human renal cortical epithelial cells despite low level FolR1 expression at the highest concentration of 10.000 pM of TCB, demonstrating the effectiveness of the anti-idiotypic masking moiety. Little T cell activation can be observed for the 16D5 TCBs (inverted and classic format) (FIG. 23).

Example 11

Anti-ID CD3 Fab Masking CD3 Binder of 16D5 TCB. Killing on Ovar3 Cells

[0765] T-cell-mediated target cell killing mediated by protease activated TCB molecules was assessed on OVCAR3 cells (FIG. 24). Human PBMCs were used as effector cells and cell killing was detected at 48 h of incubation with the molecules. Human Peripheral blood mononuclear cells (PBMCs) were isolated from fresh taken blood of a healthy

donor. 50 ml Leucosep tubes (GreinerBioOne) were used for preparation. Blood was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature), the plasma above the PBMC-containing interphase was discarded and PBMCs transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant discarded and the PBMC pellet resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation at 37° C. for about 2-3 minutes the tubes were filled with sterile PBS to 50 ml and centrifuged at 350×g for 10 minutes. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 2% FCS and 1×GlutaMax at 37° C., 5% CO₂ in cell incubator until further use. Briefly, adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended at 0.4×10⁶ cells/ml in assay medium (RPMI1640, 2% FCS, 1×GlutaMax). Target cells were plated at a density of 20 000 cells/well using round-bottom 96-well plates. For the killing assay, the molecules were added at the indicated concentrations in triplicates. FolR1 16D5 TCB was included as positive control and an untargeted TCB molecule (binding to CD3 but not to a target cell antigen) was included as negative control. PBMCs were added to target cells at final E:T ratio of 10:1. Target cell killing was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells and PBMCs with 1% Triton X-100 1 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB. The result (FIG. 24) shows that protease activated TCB with anti-idiotypic CD3 4.15.64 crossed Fab N—terminally linked by a non-cleavable linker is not significantly masking the CD3 binder. Further, Ovar3 cells appear to express Matriptase because untreated molecule also induces killing of these cells.

Example 12

Killing on Skov3 and HeLa Cells with Three Different Human PBMC Donors

[0766] T-cell killing mediated by protease activated TCB molecules was assessed on two different cell lines expressing different levels of FolR1 (FIGS. 25-27). Human PBMCs were used as effector cells and cell killing was detected at 48 h of incubation with the molecules. Human Peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats obtained from healthy human donors. For enriched lymphocyte preparations (buffy coats) Histopaque-1077 density preparation was used. Blood/buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature), the plasma above the PBMC-containing interphase was discarded and PBMCs transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant discarded and the PBMC pellet resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation at 37° C. for about 2-3 minutes the tubes were filled with sterile PBS to 50 ml and centrifuged at 350×g for 10 minutes. This washing step was repeated

once prior to resuspension of PBMCs in RPMI1640 medium containing 10% FCS and 1×GlutaMax. PBMCs were resuspended in RPMI1640 medium containing 10% FCS, 1×GlutaMax and 10% DMSO. PBMCs were frozen overnight at -80° C. in Cool Cell boxes before they were transferred to liquid nitrogen. 24 h before assay start, PBMCs were thawed and kept in RPMI1640 medium containing 10% FCS and 1×GlutaMax at 37° C., 5% CO₂ in cell incubator. The day before assay start adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended at 0.4×10⁶ cells/ml in appropriate medium. Target cells were plated at a density of 20 000 cells/well using flat-bottom 96-well plates. On the day of assay start PBMCs were counted and checked for viability. PBMCs were centrifuged at 350 g for 5 min and resuspended in assay medium (RPMI1640, 2% FCS, 1×GlutaMax). The medium of target cells was removed and PBMCs were added to the target cells before diluted antibodies were added at the indicated concentrations in triplicates. FolR1 16D5 TCB was included as positive control and an untargeted TCB molecule (binding to CD3 but not to a target cell antigen) was included as negative control. PBMCs were added to target cells at E:T ratio of 10:1. Target cell killing was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 2 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB.

[0767] The results (FIGS. 25-27) show that FolR1 TCB with scFv 4.32.63 N-terminally linked by a non-cleavable linker (FIG. 5D) induced reduced killing on HeLa cells at concentration of 100 pM and on Skov3 cells at a concentration of 10 nM. FolR1 TCB with scFv 4.15.64 N-terminally linked by a non-cleavable linker (FIG. 5B) was less efficient in reducing killing on Skov3 cells at a concentration of 10 nM. The stronger mask, meaning the anti-idiotypic scFv 4.32.63 with the higher affinity, is more efficient in masking the CD3 binder than the weak anti-idiotypic scFv 4.15.64. Comparable potency of treated and untreated TCBs suggests protease, e.g. Matriptase, expression by HeLa and Skov3 cells.

Example 13

Preparation of the HER1 Binding Antibody GA201 Masked with an Anti-Idiotypic GA201 scFv

[0768] The following molecules were prepared in this example:

[0769] 1: GA201 IgG1 antibody with N-terminal fusion of an anti-idiotypic GA201 scFv and Matrix Metalloprotease site in glycine serine linker (SEQ ID NOs 32 and 34); and

[0770] 2: HER1-binding IgG1 antibody GA201 (SEQ ID NOs 32 and 33).

[0771] Schematic illustrations thereof are shown in FIGS. 28 and 29. The GA201 anti-idiotypic (ID) binder sequence was obtained by RT-PCR (reverse transcription) from RNA of Hybridoma cells using degenerated primers binding to the ends of the variable light and heavy chain, respectively. Hybridoma cells were obtained by immunization of mice. Single chain Fv (scFv) DNA sequence synthesis with flank-

ing singular restriction endonuclease sites was ordered at Geneart and cloned as N-terminal fusion at the GA201 light chain.

[0772] A Roche expression vector was used for the construction of all heavy and light chain scFv fusion protein encoding expression plasmids. The vector is composed of the following elements:

[0773] a hygromycin resistance gene as a selection marker,

[0774] an origin of replication, oriP, of Epstein-Barr virus (EBV),

[0775] an origin of replication from the vector pUC18 which allows replication of this plasmid in *E. coli*

[0776] a beta-lactamase gene which confers ampicillin resistance in *E. coli*,

[0777] the immediate early enhancer and promoter from the human cytomegalovirus (HCMV),

[0778] the human I-immunoglobulin polyadenylation ("poly A") signal sequence, and

[0779] unique BamHI and XbaI restriction sites.

[0780] The molecules were produced by co-transfecting human embryonic kidney 293-F cells growing in suspension with the mammalian expression vectors using the FreeStyle™ 293 Expression System according to the manufacturer's instruction (Invitrogen, USA). Briefly, suspension FreeStyle™ 293-F cells were cultivated in FreeStyle™ 293 Expression medium at 37° C./8% CO₂ and the cells were seeded in fresh medium at a density of 1-2×10⁶ viable cells/ml on the day of transfection. DNA-293fectin™ complexes were prepared in Opti-MEM I medium (Invitrogen, USA) using 325 µl of 293fectin™ (Invitrogen, Germany) and 250 µg of heavy ("GA201 heavy chain") and light chain ("anti-GA201 VH-VL scFv MMP cleavable linker G4S GA201 light chain" or "GA201 light chain") plasmid DNA in a 1:1 molar ratio for a 250 ml final transfection volume. Antibody containing cell culture supernatants were harvested 7 days after transfection by centrifugation at 14000 g for 30 minutes and filtered through a sterile filter (0.22 µm). Supernatants were stored at -20° C. until purification.

[0781] The secreted protein was purified from cell culture supernatants by affinity chromatography using ProteinA affinity chromatography, followed by size exclusion chromatography. Briefly, sterile filtered cell culture supernatants were applied to a HiTrap ProteinA HP (5 ml) column equilibrated with PBS buffer (10 mM Na2HPO4, 1 mM KH2PO4, 137 mM NaCl and 2.7 mM KCl, pH 7.4). Unbound proteins were washed out with equilibration buffer. Antibody and antibody variants were eluted with 0.1 M citrate buffer, pH 2.8, and the protein containing fractions were neutralized with 0.1 ml 1 M Tris, pH 8.5. Then, the eluted protein fractions were pooled, concentrated with an Amicon Ultra centrifugal filter device (MWCO: 30 K, Millipore) to a volume of 3 ml and loaded on a Superdex200 HiLoad 120 ml 16/60 gel filtration column (GE Healthcare, Sweden) equilibrated with 20 mM Histidin, 140 mM NaCl, pH 6.0. Fractions containing purified GA201-anti-GA201-scFv or GA201 with less than 5% high molecular weight aggregates were pooled and stored as 1.0 mg/ml aliquots at -80° C.

[0782] For Protein analytics after size exclusion chromatography, the purity and molecular weight of the molecules in the single fractions were analyzed by SDS-PAGE in the absence of a reducing agent and staining with Coomassie (InstantBlue™, Expedeon). The NuPAGE® Pre-Cast gel

system (4-12% Bis-Tris, Invitrogen or 3-8% Tris-Acetate, Invitrogen) was used according to the manufacturer's instruction.

[0783] The protein concentration of purified protein samples was determined by measuring the optical density (OD) at 280 nm divided by the molar extinction coefficient calculated on the basis of the amino acid sequence. Purity and molecular weight of the molecules after the final purification step were analyzed by CE-SDS analyses in the presence and absence of a reducing agent. The Caliper LabChip GXII system (Caliper Lifescience) was used according to the manufacturer's instruction. The aggregate content of the molecules was analyzed by high-performance SEC using a Superdex 200 analytical size-exclusion column (GE Healthcare, Sweden) in 200 mM KH2PO4, 250 mM KCl, pH 7.0 running buffer at 25° C. 25 µg protein were injected on the column at a flow rate of 0.5 ml/min and eluted isocratic over 50 minutes.

[0784] The final purity of all molecules was ≥95% monomer content as detected by high performance SEC. The molecular weight of the anti-idiotypic scFv masked GA201 was determined by CE-SDS analysis as 216.3 kDa under non reducing conditions (FIG. 1A) and under reducing conditions as 58.3 kDa for the GA201 heavy chain and 60.3 kDa for the scFv linked GA201 light chain (FIG. 30B), respectively. The molecular weight based on the amino acid sequence was calculated as 49.2 kDa for the heavy chain and 51.9 kDa for the scFv fused GA201 light chain, which indicates glycosylation of both chains in HEK293 cells.

TABLE 3

Summary of production and purification of protease-activated GA201 IgG (FIG. 28) and GA201 (FIG. 29) control molecules.			
Molecule	Supernatant	Protein A-Yield	SEC-Yield
1	1.0 L	1.3 mg	0.4 mg
2	1.0 L	26.4 mg	24 mg

Example 14

Masking Effect of an Anti-Idiotypic scFv for GA201 IgG

[0785] The efficiency of masking the HER1 binding of GA201 by N-terminal fusion of an anti-idiotypic GA201 scFv was shown by FACS analysis on HER1 expressing H322M cells and Surface Plasmon Resonance (SPR) analysis on a HER1 coated chip surface. For proteolytic cleavage of GA201-anti-GA201-scFv recombinant active human MMP2 (Calbiochem) was used. 1 mg of GA201 anti-idiotypic scFv fused to GA201 by a glycine serine linker containing a MMP cleavage site was incubated with 1.2 µg MMP2 overnight at 37° C. in PBS.

[0786] For FACS analysis of HER1 binding of cleaved and uncleaved GA201-anti-GA201-scFv, the non-small cell lung cancer line H322M was used. Cells were adjusted to 1×10⁶/ml and distributed to a 96-well round-bottom plate. The molecules were added and incubated on ice for 30 minutes. Cells were washed once with FACS buffer (PBS+ 2% FCS+0.1% sodium azide) and re-suspended with a F(ab')₂-goat anti-human IgG Fc secondary antibody FITC conjugate (ThermoFisher Scientific). After another 20 minutes on ice, cells were washed twice and re-suspended in

FACS buffer and analyzed in a BD FACS Canto II. 10000 cells were measured and the median of the fluorescence signal was used for analysis. Before MMP-2 cleavage of GA201-anti-GA201-scFv no binding to HER1 on H322M cells was measurable, indicating complete masking of the GA201 binding domains by the anti-idiotypic scFv (FIG. 31). Binding of uncleaved GA201-anti-GA201-scFv was comparable to an unspecific isotype IgG control antibody (FIG. 31). In contrast, MMP cleavage of the anti-idiotypic scFv leads to activation of GA201 and binding to HER1 on H322M cells was restored to similar levels as the unmasked parental antibody GA201 (FIG. 31).

[0787] To confirm the FACS binding data of masked GA201 binding after MMP cleavage, we also performed a SPR experiment as second analytical method using a Biacore T100 instrument (GE Healthcare Biosciences AB, Uppsala, Sweden). HER1 was immobilized on the surface of a CMS biosensorchip using standard amine-coupling chemistry. The HER1 extracellular domain was injected in sodium acetate, pH 5.0 at 1 µg/ml. Reference control flow cells were treated in the same way but with vehicle buffer only. GA201-anti-GA201-scFv, before and after an overnight MMP cleavage, and GA201 were diluted in 1×PBS pH 7.4, 0.05% Tween20 Roche Diagnostics GmbH) and injected at increasing concentrations between 3.125 and 50 nM with a flow rate of 30 µl/min. The association phase was 3 minutes and the dissociation time was 10 minutes. HER1 binding was regenerated with an inject of 0.85% phosphoric acid for 30 s at a flow rate of 5 µl/min. Kinetic rate constants and equilibrium dissociation constants were calculated by using the 1:1 Langmuir binding model within the Biaevaluation software. A K_D value of 1 nM for binding of HER1 was determined for the GA201 parental unmasked antibody (FIG. 32). After an overnight MMP-2 incubation of GA201-anti-GA201-scFv, a K_D value of 2 nM was measured with similar k_a and k_d rate constants for association and dissociation as the unmasked control antibody, indicating complete restoration of HER1 binding by protease cleavage (FIG. 32). Uncleaved GA201-anti-GA201-scFv did not show any binding to HER1 in SPR analysis (FIG. 32). In summary, we have demonstrated a complete loss of binding to HER1 by fusion of an anti-idiotypic scFv to the N-terminus of the IgG₁ antibody GA201 with two independent analytical methods. Furthermore, binding to HER1 was fully restored by removal of the scFv through protease cleavage in the MMP cleavage site in the glycine serine linker.

Example 15

Preparation of Anti FolR1/Anti-CD3 and antiMesothelin/Anti-CD3 T Cell Bispecific (TCB) Molecules with Anti CD3 scFv

[0788] Several T cell bispecific (TCB) molecules with various anti-idiotypic scFv were produced and are schematically depicted in FIGS. 33A-33J with their respective ID number. The following molecules were prepared:

[0789] ID 8364: “FolR1 16D5 2+1 IgG, classic format (anti idiotypic scFv 4.32.63—MMP9—MK062 Matriptase site—CD3—N-terminal fused to FolR1 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and MMP9—MK062 protease linker” (FIG. 33A, SEQ ID NOs 1, 3 and 72).

[0790] ID 8363: “FolR1 16D5 2+1 IgG, classic format (anti idiotypic scFv 4.32.63—Cathepsin S/B site—

CD3—N-terminal fused to FolR1 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and Cathepsin S/B protease linker” (FIG. 33B, SEQ ID NOs 1, 3 and 85).

[0791] ID 8365: “FolR1 16D5 2+1 IgG, inverted format, (anti idiotypic scFv 4.32.63—MK062 Matriptase linker—CD3—N-terminal fused to CD3 VL—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and MK062 Matriptase linker” (FIG. 33C, SEQ ID NOs 1, 3, 73 and 74).

[0792] ID 8366: “FolR1 16D5 2+1 IgG, inverted format, (anti idiotypic scFv 4.32.63—non-cleavable GS linker—CD3—N-terminal fused to CD3 VL—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and non-cleavable GS linker” (FIG. 33D).

[0793] ID 8672: “aMesothelin 2+1 IgG, classic format, MSLN charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—MMP9—MK062 Matriptase—CD3—N-terminal fused to aMesothelin VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and MMP9—MK062 Matriptase” (FIG. 33E, SEQ ID NOs 77, 78, 81, 82).

[0794] ID 8673: “aMesothelin 2+1 IgG, classic format, MSLN charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—non-cleavable GS linker—CD3—N-terminal fused to aMesothelin VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 non-cleavable GS linker” (FIG. 33F).

[0795] ID 8674: “aMesothelin 2+1 IgG, inverted format, MSLN charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—MMP9—MK062 Matriptase—CD3—N-terminal fused to CD3 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and MMP9—MK062 Matriptase” (FIG. 33G, SEQ ID NOs 76, 77, 78, 79).

[0796] ID 8675: “aMesothelin 2+1 IgG, inverted format, MSLN charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—non-cleavable GS linker—CD3—N-terminal fused to CD3 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and non-cleavable GS linker” (FIG. 33H).

[0797] ID 8505: “aMesothelin 2+1 IgG, inverted format, MSLN charged variants, CD3 (aMesothelin HC N-terminally fused to CD3 VL—inert Fc)” (FIG. 33I).

[0798] ID 8676: “aMesothelin 2+1 IgG, classic format, MSLN charged variants, CD3 crossed (aMesothelin IgG with CD3—N-terminal fused to aMesothelin VH—inert Fc)” (FIG. 33J)

[0799] The variable domains were subcloned in frame with the pre-inserted domains into the respective recipient mammalian expression vector. Protein expression is driven by an MPSV promoter and a synthetic polyA signal sequence is present at the 3' end of the CDS. In addition each vector contains an EBV OriP sequence.

[0800] The molecules (except 8505, this molecule was produced by co-transfecting CHO cells growing in suspension with the mammalian expression vectors. Transient transfection was done at Evitria AG (Switzerland).) were produced by co-transfecting HEK293-EBNA cells growing in suspension with the mammalian expression vectors using polyethylenimine (PEI). For transfection HEK293 EBNA cells were cultivated in serum free ExCell culture medium containing 6 mM L-glutamine and 250 mg/l G418. For the production in 600 ml tubespin flasks (max. working volume

400 ml) 800 million HEK293 EBNA cells were seeded 24 hours before transfection without G418. For transfection 800 mio cells were centrifuged for 5 min at 210×g and supernatant was replaced by 40 ml pre-warmed CD CHO medium containing 6 mM L-Glutamine. Expression vectors were mixed with 40 ml CD CHO medium containing 6 mM L-Glutamine to a total amount of 400 µg DNA. After addition of 1080 µl PEI solution (2.7 µg/ml) the mixture was vortexed for 15 s and subsequently incubated for 10 min at room temperature. Afterwards cells were mixed with the DNA/PEI solution, transferred to a 600 ml tubespin flask and incubated for 3 hours at 37° C. in an incubator with a 5% CO₂ atmosphere. After incubation, 320 ml ExCell+6 mM L-glutamine+5 g/L Pepsoy+1.0 mM VPA+3 g/l glucose medium was added and cells were cultivated for 24 hours prior to feeding with 7% Feed 7. After 6-7 days the cultivation supernatant was collected for purification by centrifugation for 20-30 min at 210×g (Sigma 8K centrifuge). The solution was sterile filtered (0.22 µm filter) and sodium azide in a final concentration of 0.01% w/v was added. The solution was kept at 4° C. until purification.

[0801] The secreted protein was purified from cell culture supernatants by affinity chromatography using ProteinA affinity chromatography, followed by one to two size exclusion chromatographic steps. For affinity chromatography supernatant was loaded on a Protein A MabSelectSure (CV=5 mL, GE Healthcare) equilibrated with 20 mM Sodium Citrate, 20 mM Sodium Phosphate, pH 7.5. Unbound protein was removed by washing with at least 10 column volumes 20 mM Sodium Citrate, 20 mM Sodium Phosphate, pH 7.5 and target protein was eluted in 20 column volumes (gradient from 0%-100%) 20 mM Sodium Citrate, 100 mM Sodium Chloride, 100 mM Glycine, pH 3.0. Protein solution was neutralized by adding 1/10 of 0.5 M Na₂HPO₄ pH 8.0. Target protein was concentrated with Amicon® Ultra-15 Ultracel 30K (Merck Millipore Ltd.) to a volume of 4 ml maximum prior loading on a HiLoad Superdex 200 column (GE Healthcare) equilibrated with 20 mM Histidine, 140 mM NaCl, 0.01% Tween pH 6.0.

[0802] The protein concentration of purified protein samples was determined by measuring the optical density (OD) at 280 nm divided by the molar extinction coefficient calculated on the basis of the amino acid sequence.

[0803] Purity and molecular weight of the molecules after the final purification step were analyzed by CE-SDS analyses in the presence and absence of a reducing agent. The Caliper LabChip GXII system (Caliper Lifescience) was used according to the manufacturer's instruction.

[0804] The aggregate content of the molecules was analyzed using a TSKgel G3000 SW XL analytical size-exclusion column (Tosoh) in 25 mM K₂HPO₄, 125 mM NaCl, 200 mM L-arginine monohydrochloride, 0.02% (w/v) Na₃N, pH 6.7 running buffer at 25° C.

[0805] The final quality of all molecules was good, with ≥95% monomer content.

TABLE 4

Summary of production and purification of protease activated TCB molecules.			
Molecule	Titer [mg/l]	Yield [mg/l]	Analytical SEC (HMW/Monomer/LMW) [%]
1 (8364)	34.55	1.72	0.68/99.32/0
2 (8363)	33.75	1.59	4.02/95.98/0

TABLE 4-continued

Summary of production and purification of protease activated TCB molecules.			
Molecule	Titer [mg/l]	Yield [mg/l]	Analytical SEC (HMW/Monomer/LMW) [%]
3 (8365)	5.35	0.24	2.71/96.46/0.83
4 (8366)	4.2	0.43	4.908/96.02/0
5 (8672)	13.8	1.59	3.96/96.04/0
6 (8673)	14	1.99	2.15/97.85/0
7 (8674)	3.6	0.96	6.27/93.73/0
8 (8675)	5.2	0.59	5.81/90.63/3.57
9 (8505)	120	20.46	0.47/99.32/0.22
10 (8676)	22.5	3.84	1.98/96.21/1.81

Example 16

Quality Control and Stability—Capillary Electrophoresis SDS Analysis of Different TCB Molecules

[0806] Purity and molecular weight of the molecules after the final purification step were analyzed by CE-SDS analyses in the presence and absence of a reducing agent. The Caliper LabChip GXII system (Caliper Lifescience) was used according to the manufacturer's instruction. Comparison of untreated molecules (stored at 4° C.), treated molecules (treated with appropriate recombinant protease (R&D Systems) for 24 h at 37° C. and molecule incubated for 72 h at 37° C. (FIGS. 34, 35A, and 35B). Comparison of the untreated and treated molecule shows complete cleavage of the anti ID scFv after rhMatriptase/ST14 treatment for the inverted format containing MK062 Matriptase linker but incomplete cleavage of MMP9-MK062 Matriptase linker. rhCathepsin B and rhCathepsin S treatment is incomplete as well. The conditions for the purified enzymes have not been optimal. Molecules incubated at 37° C. for 72 h are running on the same height than pure molecules suggesting that the molecules are stable at 37° C. for the time of in vitro assay duration. Pre-stained protein Marker Mark 12 (Invitrogen) was used for estimation of correct molecule weight.

Example 17

Comparison of Different Linkers and Formats of Protease Activated FolR1 TCBs

[0807] Jurkat NFAT activation assay. Jurkat NFAT activation assay for comparison of different formats and linkers of protease activated TCB. Jurkat-NFAT reporter cell line (Promega) is a human acute lymphatic leukemia reporter cell line with a NFAT promoter, expressing human CD3c. If the TCB binds the tumor target and the CD3 binder (crosslinkage) binds the CD3ε Luciferase expression can be measured in Luminescence after addition of One-Glo substrate (Promega). 20.000 target cells were seeded in 96-well white walled clear bottom plate (Greiner BioOne) in 50 µl/well Jurkat medium (RPMI1640, 2 g/l Glucose, 2 g/l NaHCO₃, 10% FCS, 25 mM HEPES, 2 mM L-Glutamin, 1×NEAA, 1×Sodium-pyruvate) without Hygromycine. Plates were incubated for about 20 hours at 37° C. Jurkat-NFAT reporter cells were harvested and viability was assessed using ViCell. Cells were resuspended in Jurkat medium without Hygromycine and 50 µl per well (50.000 cells/well) were added.

The E:T ratio was 2.5:1 (based on cell number seeded). Antibodies were diluted in Jurkat medium without Hygromycin and 50 μ l/well were added. Cells were incubated at 37° C. for 6 h in a humidified incubator before they were taken out of the incubator for about 10 min to adapt to room temperature prior to Luminescence read out. 50 μ l/well of ONE-Glo solution were added to wells and incubated for 10 min at room temperature in the dark. Luminescence was detected using WALLAC Victor3 ELISA reader (PerkinElmer2030), 1 sec/well as detection time. Comparison of the pretreated protease activated TCB (8364, grey filled squares) and FolR1 TCB (black triangles pointing down) showed that potency after cleavage is recovered completely. No Luminescence was detectable for cells incubated with the masked TCB (containing a GS non-cleavable linker, grey triangles pointing up) and the non-targeted TCB control (empty triangle pointing down) for both cell lines in this concentration range. The dotted line shows the Luminescence of target cells and effector cells without any TCB (FIGS. 36A and 36B).

Example 18

Tumor Cell Cytotoxicity Mediated by Different Formats of Protease Activated TCB

[0808] T-cell killing mediated by protease activated TCB molecules was assessed on cell lines expressing different levels of FolR1. Human Peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats obtained from healthy human donors. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450 $\times$ g, 30 minutes, w/o break, room temperature) the PBMC-containing interphase was transferred in a new falcon tube that was subsequently filled with 50 ml of PBS. The mixture was centrifuged (400 $\times$ g, 10 minutes, room temperature), the supernatant was discarded and the PBMC pellet was resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation for about 2-3 minutes at 37° C. the tubes were filled with sterile PBS to 50 ml and centrifuged for 10 minutes at 350 $\times$ g. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 10% FCS, 1 $\times$ GlutaMax and 10% DMSO. PBMCs were slowly frozen in CoolCell® Cell Freezing Containers (BioCision) at -80° C. and then transferred to liquid nitrogen. One day before assay start adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended in assay medium (RPMI1640, 2% FCS, 1 $\times$ GlutaMax). Target cells were plated at a density of 20 000 cells/well using 96-well flat-bottom plates and incubated for about 20 h at 37° C. in a humidified incubator. About 20 h before assay start PBMCs were thawed in RPMI1640 medium (10% FCS, 1 $\times$ GlutaMax). PBMCs were centrifuged at 350 g for 7 min. The pellet was resuspended in fresh medium (RPMI1640, 10% FCS, 1 $\times$ GlutaMax) and incubated for max 24 h at 37° C. in a humidified incubator. On the day of the assay start PBMCs were harvested and centrifuged at 350 g for 7 min. The pellet was resuspended in assay medium and 0.2 mio PBMCs in 100 μ l/well (E:T 10:1, based on the number of seeded target cells) were added to the target cells. The molecules were diluted in assay medium (RPMI1640, 2% FCS, 1 $\times$ GlutaMax) and 50 μ l/well were added at the indicated concentrations in triplicates before the plates were incubated for about 48 h at 37° C. in a humidified incubator.

Target cell killing was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB.

[0809] The results (FIGS. 37A and 37B) show the comparison of two different formats of the Protease activated TCBs both containing the anti idiotypic CD3 scFv 4.32.63 linked with a MK062 Matriptase linker. The inverted format of the protease activated TCB (8365, grey circles) seems to be more potent in killing (HeLa and Skov-3 target cells) than the classic format of the protease activated TCB (8408, dark grey triangles pointing up). However the inverted molecule containing the non-cleavable linker (8366, light grey squares) is less efficient in masking than the classic molecule (8409, dark grey triangles pointing down).

[0810] FIG. 37C HeLa target cell cytotoxicity. Comparison of classic Protease activated TCB containing the anti idiotypic CD3 scFv 4.32.63 and GS linkers with different protease sites. Protease activated TCB containing the MMP9-Matriptase MK062 linker (8364, grey squares) reaches the potency of FolR1 TCB (light grey triangles pointing down) whereas the protease activated TCB containing only Matriptase MK062 (light grey rhomb) is less potent in killing HeLa cells. Molecules containing Cathepsin site (grey circles) or non-cleavable linker (black triangles pointing down) are comparable. FIG. 37D Skov-3 target cell cytotoxicity. Comparison of classic Protease activated TCB containing the anti idiotypic CD3 scFv 4.32.63 and GS linkers with different protease sites. Protease activated TCB containing the MMP9-Matriptase MK062 linker (8364, grey squares) nearly reaches the potency of FolR1 TCB (light grey triangles pointing down) whereas the protease activated TCB containing only Matriptase MK062 (light grey rhomb) is less potent in killing Skov-3 cells. The molecule containing Cathepsin site (grey circles) is less potent than the molecule containing only the Matriptase MK062 site and the molecule containing the non-cleavable linker (black triangles pointing down) only induces killing below 10% in the indicated concentration range for Skov-3 cells.

Example 19

T-Cell Activation after Co-Incubation of Human Renal Epithelial Cortical Cells or Human Bronchial Epithelial Cells with TCBs and Human PBMCs

[0811] T-cell activation mediated by protease activated TCB molecules was assessed for HRCEpi (Human renal cortical epithelial cells) and HBEPiC (human bronchial epithelial cells expressing only little amounts of FolR1). Human PBMCs were used as effector cells and T cell activation markers were stained after 48 h of incubation with the molecules and cells. Human Peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats obtained from healthy human donors. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450 $\times$ g, 30 minutes, w/o break, room temperature) the PBMC-containing interphase was transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400 $\times$ g, 10 minutes, room temperature), the supernatant was discarded

and the PBMC pellet was resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation for about two minutes at 37° C. the tubes were filled with sterile PBS to 50 ml and centrifuged for 10 minutes at 350×g. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 10% FCS, 1×GlutaMax and 10% DMSO. PBMCs were slowly frozen in CoolCell® Cell Freezing Containers (BioCision) at -80° C. and then transferred to liquid nitrogen. One day before the assay was started adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended in assay medium (RPMI1640, 2% FCS, 1×GlutaMax). Target cells were plated at a density of 20 000 cells/well using 96-well flat-bottom plates and incubated for about 20 h at 37° C. in a humidified incubator. About 20 h before assay start PBMCs were thawed in RPMI1640 medium (10% FCS, 1×GlutaMax). PBMCs were centrifuged for 7 min at 350 g. The pellet was resuspended in fresh medium (RPMI1640, 10% FCS, 1×GlutaMax) and incubated for max 24 h at 37° C. in a humidified incubator. On the day of the assay start PBMCs were harvested and centrifuged for 7 min at 350 g. The pellet was resuspended in assay medium and 0.2 mio PBMCs in 100 ul/well (E:T 10:1, based on the number of seeded target cells) were added to the target cells. The molecules were diluted in assay medium (RPMI1640, 2% FCS, 1×GlutaMax) and added at the indicated concentrations in triplicates before the plates were incubated for about 48 h at 37° C. in a humidified incubator.

[0812] T-cell activation was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of CD25 and CD69 on CD4 positive and CD8 positive T cells. FolR1 16D5 TCB (6298) and an untargeted TCB (binding to CD3 but not to a target cell antigen, 7235) were included as controls. Each point represents the mean value of triplicates of three different human PBMC donors. Standard deviation is indicated in error bars. Unpaired t test was used for statistical analysis. The results show an increase in CD69 for CD8 positive cells for the FolR1 TCB that is significantly higher than the median fluorescence intensity for the protease activated TCBs (FIGS. 38A and 38B).

Example 20

Tumor Cell Cytotoxicity Mediated by Different Formats of Protease Activated Mesothelin (MSLN) TCB

[0813] T-cell killing mediated by protease activated TCB molecules was assessed on cell lines expressing different levels of Mesothelin (MSLN). Human Peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats obtained from healthy human donors. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature) the PBMC-containing interphase was transferred in a new falcon tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant was discarded and the PBMC pellet was resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation for about two minutes at 37° C. the tubes were filled with sterile PBS to 50 ml and centrifuged for 10 minutes at 350×g. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 10% FCS, 1×GlutaMax and 10% DMSO. PBMCs were slowly frozen

in CoolCell® Cell Freezing Containers (BioCision) at -80° C. and then transferred to liquid nitrogen. Adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended in assay medium (RPMI1640, 2% FCS, 1×GlutaMax) one day before the assay was started. Target cells were plated at a density of 20 000 cells/well using 96-well flat-bottom plates and incubated for about 20 h at 37° C. in a humidified incubator. PBMCs were thawed in RPMI1640 medium (10% FCS, 1×GlutaMax) about 20 h before assay start. PBMCs were centrifuged for 7 min at 350 g. The pellet was resuspended in fresh medium (RPMI1640, 10% FCS, 1×GlutaMax) and incubated for max 24 h at 37° C. in a humidified incubator. On the day of the assay start PBMCs were harvested and centrifuged for 7 min at 350 g. The pellet was resuspended in assay medium and 0.2 mio PBMCs in 100 ul/well (E:T 10:1, based on the number of seeded target cells) were added to the target cells. The molecules were diluted in assay medium (RPMI1640, 2% FCS, 1×GlutaMax) and added at the indicated concentrations in triplicates before the plates were incubated for about 48 h at 37° C. in a humidified incubator. Target cell killing was assessed after 48 h of incubation at 37° C., 5% CO₂ by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB.

[0814] The results (FIGS. 39A and 39B) show target cell killing mediated by Protease activated MSLN TCB (8672) for NCI H596 and AsPC-1 cell lines. The protease activated TCBs nearly reaches the potency of MSLN TCB (8676) for NCI H596 and AsPC-1. The molecule containing the non-cleavable GS linker (8673) does not induce killing in the indicated concentration range for both cell lines.

Example 21

Jurkat-NFAT Reporter Assay to Monitor Target Expression (FOLR1 TCB) and Protease Activity (Protease Activated FOLR1 TCB) in Primary Tumor Samples

[0815] The intention of this assay was to show tumor target antigen (FolR1) expression and activity of tumor specific proteases like MMP9, Matrilysin or Cathepsin in human tumor samples.

[0816] Jurkat-NFAT reporter cell line (Promega) is a human acute lymphatic leukemia reporter cell line with a NFAT promoter, expressing human CD3c. Luciferase expression can be measured, if the T cell bispecific molecule binds the tumor target and the CD3c (crosslinkage). Luminescence is measured after addition of One-Glo substrate (Promega).

[0817] Primary tumor samples were received from Indivumed GmbH, Germany. Samples were shipped over night in transport medium. About 24 h after surgery the sample was cut in small pieces. 96-well white walled, flat (clear) bottom plate was prepared by adding 18 ul cold Matrigel (Matrigel (734-1101, Corning/VWR). Plate was incubated for 2 min at 37° C. before tumor pieces were added (triplicates). 33 ul of cold Matrigel were added per well and plate was incubated again for 2 min at 37° C. 50 ul of antibody dilution (in Jurkat medium without Hygromycin

but containing 2×Penicillin/Streptomycin) was added per well and plate was incubated for about 48 hours at 37° C., 5% CO₂.

[0818] Jurkat-NFAT reporter cells were harvested and viability was assessed using ViCell. Cells were centrifuged at 350×g, 7 min before they were resuspended in Jurkat medium without Hygromycin and 50 µl per well (50,000 cells/well) were added. Plate was incubated for 5 h at 37° C. in a humidified incubator before it was taken out for Luminescence read out. 80 µl of each well were transferred into a white walled 96-well plate. 27 µl/well of ONE-Glo solution were added to each well and incubated for 10 min at room temperature in the dark. Luminescence was detected using WALLAC Victor3 ELISA reader (PerkinElmer2030), 1 sec/well as detection time.

[0819] Jurkat NFAT reporter cells are activated after co-incubation with FolR1 TCB (6298) and Protease activated FolR1 TCB containing MMP9-Matriptase cleavage site (8364). Protease activated FolR1 TCBs (8363, 8408) and control TCBs (8409, 7235) do not induce Luciferase expression. The dotted line indicates the baseline Luminescence for Jurkat NFAT cells co-incubated with tumor (FIG. 40).

Example 22

Serum Stability of Protease Activated TCBs

[0820] Capillary electrophoresis of protease activated TCBs after incubation in human serum. Molecules were incubated for 0 or 14 days in human IgG depleted serum at 37° C. in a humidified incubator (5% CO₂). All molecules were purified by affinity chromatography (ProteinA) and then analyzed by Capillary electrophoresis.

[0821] 100 µg of each molecule was added either in buffer (Histidine buffer (Bichsel) with 0.01% Tween-20) or in human serum (IgG depleted, SP1839, TL-15216, 16FSP63814). The concentration of the molecules was higher than 2 mg/ml and the final concentration was 0.5 mg/ml. The pretreatment for one molecule (8408) was done with rhMatriptase (R&D Systems) for 24 h at 37° C., 5% CO₂ in a humidified incubator (otherwise pH of serum could change). The samples for day 0 were directly frozen in liquid nitrogen and stored at -80° C. until analysis. Samples for day 14 were incubated for 14 days at 37° C., 5% CO₂ in a humidified incubator until they were also snap frozen.

[0822] Prior to CE-SDS analysis all samples were purified via HPLC affinity chromatography (Agilent technologies 1200series, column: Upchurch scientific C-130B, packaging material: Applied Biosystems POROS 20A 60 µl, buffer: 10 mM Tris, 50 mM Glycine, 500 mM NaCl pH 8.0 and pH 2.0, injection volume: 100 µl, flow rate 1 ml/min, collection: peak based, neutralization: 0.5 M Na-phosphate pH 8.0 10% volume). Protease activated TCB is stable in human IgG depleted serum for a minimum of 14 days (FIGS. 41A-41C).

Example 23

Design of Anti Her2/Anti-CD3 and AntiFolR1/Anti-CD3 T Cell Bispecific (TCB) Molecules with Anti CD3 scFv

[0823] Several T cell bispecific (TCB) molecules designed and are schematically depicted in FIGS. 42A-42F with their respective ID number. The following molecules were designed:

[0824] ID 8955: "Herceptarg 2+1 IgG, classic format, Herceptarg charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—MMP9—MK062 Matriptase—CD3—N-terminal fused to Herceptarg VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and MMP9—MK062 Matriptase" (FIG. 42A, SEQ ID NOS 81, 132, 133 and 136).

[0825] ID 8957: "Herceptarg 2+1 IgG, classic format, Herceptarg charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—non cleavable GS linker—CD3—N-terminal fused to Herceptarg VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and non cleavable GS linker" (FIG. 42B, SEQ ID NOS 81, 132, 133 and 135).

[0826] ID 8959: "Herceptarg 2+1 IgG, classic format, Herceptarg charged variants, CD3 crossed (Herceptarg IgG with CD3—N-terminal fused to Herceptarg VH—inert Fc)" (FIG. 42C, SEQ ID NOS 81, 132, 133 and 134).

[0827] ID 8997: "FolR1 36F2 2+1 IgG, classic format, FolR1 36F2 charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—MMP9—MK062 Matriptase—CD3—N-terminal fused to FolR1 36F2 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and MMP9—MK062 Matriptase" (FIG. 42D, SEQ ID NOS 81, 137, 138 and 139).

[0828] ID 8998: "FolR1 36F2 2+1 IgG, classic format, FolR1 36F2 charged variants, CD3 crossed (anti idiotypic scFv 4.32.63—non cleavable GS linker—CD3—N-terminal fused to FolR1 36F2 VH—inert Fc) with N-terminal fused anti CD3 scFv 4.32.63 and non cleavable GS linker" (FIG. 42E, SEQ ID NOS 81, 137, 138 and 140).

[0829] ID 8996: "FolR1 36F2 2+1 IgG, classic format, FolR1 36F2 charged variants, CD3 crossed (FolR1 36F2 IgG with CD3—N-terminal fused to FolR1 36F2 VH—inert Fc)" (FIG. 42F, SEQ ID NOS 81, 137, 138 and 141).

[0830] The variable domains were subcloned in frame with the pre-inserted domains into the respective recipient mammalian expression vector. Protein expression is driven by an MPSV or CMV (for Herceptarg) promoter and a synthetic polyA signal sequence is present at the 3' end of the CDS. In addition each vector contains an EBV OriP sequence.

Example 24

Primary Cell Cytotoxicity Mediated by Protease Activated FolR1 TCB

[0831] T-cell killing mediated by protease activated FolR1 TCB molecule was assessed on primary cell lines expressing low levels of FolR1 (FIG. 43). Human Peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats obtained from healthy human donors. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature) the PBMC-containing interphase was transferred in a new falcon tube that was subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant was discarded and the PBMC pellet was resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation for about 2-3 minutes at 37° C. the tubes were

filled with sterile PBS to 50 ml and centrifuged for 10 minutes at 350×g. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 10% FCS, 1×GlutaMax and 10% DMSO. PBMCs were slowly frozen in CoolCell® Cell Freezing Containers (BioCision) at -80° C. and then transferred to liquid nitrogen. One day before assay start adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended in assay medium (RPMI1640, 2% FCS, 1×GlutaMax). Target cells were plated at a density of 20 000 cells/well using 96-well flat-bottom plates and incubated for about 20 h at 37° C. in a humidified incubator. About 20 h before assay start PBMCs were thawed in RPMI1640 medium (10% FCS, 1× GlutaMax). PBMCs were centrifuged at 350 g for 7 min. The pellet was resuspended in fresh medium (RPMI1640, 10% FCS, 1×GlutaMax) and incubated for max 24 h at 37° C. in a humidified incubator. On the day of the assay start PBMCs were harvested and centrifuged at 350 g for 7 min. The pellet was resuspended in assay medium and 0.2 mio PBMCs in 100 ul/well (E:T 10:1, based on the number of seeded target cells) were added to the target cells. The molecules were diluted in assay medium (RPMI1640, 2% FCS, 1×GlutaMax) and 50 ul/well were added at the indicated concentrations in triplicates before the plates were incubated for about 48 h, 72 h or 96 h at 37° C. in a humidified incubator. Target cell killing was assessed after 48 h, 72 h and 96 h of incubation at 37° C., 5% CO2 by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB.

[0832] Human Bronchial Epithelial Cell toxicity mediated by human PBMCs and 100 nM or 10 nM of FolR1 TCB is higher compared to Protease activated TCB.

Example 25

FolR1 Negative Target Cell Cytotoxicity Mediated by Protease Activated FolR1 TCB

[0833] T-cell killing mediated by protease activated FolR1 TCB molecule was assessed on FolR1 negative Mkn-45 cell line (FIG. 44). Human Peripheral blood mononuclear cells

(PBMCs) were isolated from buffy coats obtained from healthy human donors. Buffy coat was diluted 1:1 with sterile PBS and layered over Histopaque gradient (Sigma, #H8889). After centrifugation (450×g, 30 minutes, w/o break, room temperature) the PBMC-containing interphase was transferred in a new falcon tube that was subsequently filled with 50 ml of PBS. The mixture was centrifuged (400×g, 10 minutes, room temperature), the supernatant was discarded and the PBMC pellet was resuspended in 2 ml ACK buffer for Erythrocytes lysis. After incubation for about 2-3 minutes at 37° C. the tubes were filled with sterile PBS to 50 ml and centrifuged for 10 minutes at 350×g. This washing step was repeated once prior to resuspension of PBMCs in RPMI1640 medium containing 10% FCS, 1×GlutaMax and 10% DMSO. PBMCs were slowly frozen in CoolCell® Cell Freezing Containers (BioCision) at -80° C. and then transferred to liquid nitrogen. One day before assay start adherent target cells were harvested with Trypsin/EDTA, counted, checked for viability and resuspended in assay medium (RPMI1640, 2% FCS, 1×GlutaMax). Target cells were plated at a density of 20 000 cells/well using 96-well flat-bottom plates and incubated for about 20 h at 37° C. in a humidified incubator. About 20 h before assay start PBMCs were thawed in RPMI1640 medium (10% FCS, 1×GlutaMax). PBMCs were centrifuged at 350 g for 7 min. The pellet was resuspended in fresh medium (RPMI1640, 10% FCS, 1x GlutaMax) and incubated for max 24 h at 37° C. in a humidified incubator. On the day of the assay start PBMCs were harvested and centrifuged at 350 g for 7 min. The pellet was resuspended in assay medium and 0.2 mio PBMCs in 100 ul/well (E:T 10:1, based on the number of seeded target cells) were added to the target cells. The molecules were diluted in assay medium (RPMI1640, 2% FCS, 1×GlutaMax) and 50 ul/well were added at the indicated concentrations in triplicates before the plates were incubated for about 48 h and 72 h at 37° C. in a humidified incubator. Target cell killing was assessed after 48 h, 72 h and 96 h of incubation at 37° C., 5% CO2 by quantification of LDH release into cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science, #11 644 793 001). Maximal lysis of the target cells (=100%) was achieved by incubation of target cells with 1% Triton X-100 20 h before LDH readout. Minimal lysis (=0%) refers to target cells co-incubated with effector cells without any TCB. Protease activated TCB did not induce target cell killing at 100 nM.

EXAMPLARY SEQUENCES

Construct	Amino acid Sequence	SEQ ID No
LC Common light chain pETR13197	QAVVTQEPSTVSPGGTVTLTGSSSTGAVTTSNYANWVQEK PGQAFRLIGGTNKRAGTPARFSGSLGGKAALTLGAQP EDEAEYYCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETT TPSKQSNNKYAASSYLSLTPEQWKSHRSYSQVTHEGSTVE KTVAPTECS	1
anti CD3 (CH2527 VH_3-23 (12) VL7-46 (13)) scFv15-Matriptase MK062 CH2527 VH3_23-VH12 CH1 FolR1 16D5 VH	QIQLVQSGPELKKPGGETVRISCKASGYTFTDYSIHVKQAPG KCLKWMGWINTETGEPAYADDFKGRFAPSLSTSASTAYLQI NNLKNETATFFCAHPYDYDVLVDYWGQGSVTVTSSGGGGS GGGGSGGGSGGGSDTVLTQSPASLVSLGQRATISCRRA SKSVSTSNYSYIHWYQQKPGQPPLKLLIKYVSYLESGVPARFS GSGSGTDFTLNHPVEEEDAATYYCQHSREFPWTFGCSTKL	2

-continued

EXAMPLARY SEQUENCES		
CH1 hum Fc knob PG LALA, pETR15422 (FIG. 45A)	<u>EIKGGGGSGGGSRQARVVNNGGGSGGGSGGGSEV</u> <u>QLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWRQAPG</u> <u>KGLEWVSRIKSKYNNYATYYADSVKGRFTISRDDSNTLYLQ</u> <u>MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVS</u> <u>SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTS</u> <u>WNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYI</u> <u>CNVNHKPSNTKVDKKVEPKSCDGGGSGGGGSEVQLVES</u> <u>GGGLVKPGGSLRLSCAASGFTFSTYAMNWRQAPGKGLE</u> <u>WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSNTLYLQMN</u> <u>LKTEDTAVYYCTTPWEWSWYDYGQGLTVTVSSASTKGPS</u> <u>VFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTSWNSGALTS</u> <u>GVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPS</u> <u>NTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPK</u> <u>DTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT</u> <u>KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA</u> <u>PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGF</u> <u>YPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD</u> <u>KSRWQQGNVFPSCVMHEALHNHYTQKSLSLSPGK</u>	
Fo1R1 16D5 VH CH1 Fc hole P329G LALA HRYF, pETR15214	<u>EVQLVESGGGLVKPGGSLRLSCAASGFTFSTYAMNWRQAPG</u> <u>PKGLEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSNTLY</u> <u>YLQMNLSLKTEDTAVYYCTTPWEWSWYDYGQGLTVTVSSA</u> <u>STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTSWN</u> <u>SGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICN</u> <u>VNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFL</u> <u>FPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVE</u> <u>VHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS</u> <u>NKALGAPIEKTISKAKGQPREPQVCTLPPSRDELTKNQVSL</u> <u>CAVKGFPYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLV</u> <u>SKLTVDKSRWQQGNVFPSCVMHEALHNRYTQKSLSLSPGK</u>	3
anti CD3 (CH2527 VH 3-23(12) VL7- 46(13)) scFv 4.32.63 Matriptase MK062 CH2527 VH3 23-VH12 CH1 Fo1R1 16D5 VH CH1 hum Fc knob PG LALA, pETR15599 (FIG. 45B)	<u>QVQLKESGPGLVAPSSQLSITCTVSGFSLTSYGVSWVRQPP</u> <u>GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKQVFLKLNS</u> <u>LQDDTATYYCAKGIITVVDDYIAMDYWGQGTSTVTVSSGG</u> <u>GGSGGGSGGGSGGGSGGGSDIQTQSPASLASVGETVTIT</u> <u>CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS</u> <u>GGSGGTQYSLKINSLSQSEVARYYCOHYSTPYTFGCCTKL</u> <u>EIKGGGGSGGGSRQARVVNNGGGSGGGSGGGSEV</u> <u>QLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWRQAPG</u> <u>KGLEWVSRIKSKYNNYATYYADSVKGRFTISRDDSNTLYLQ</u> <u>MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVS</u> <u>SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTS</u> <u>WNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYI</u> <u>CNVNHKPSNTKVDKKVEPKSCDGGGSGGGGSEVQLVES</u> <u>GGGLVKPGGSLRLSCAASGFTFSTYAMNWRQAPGKGLE</u> <u>WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSNTLYLQMN</u> <u>LKTEDTAVYYCTTPWEWSWYDYGQGLTVTVSSASTKGPS</u> <u>VFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTSWNSGALTS</u> <u>GVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPS</u> <u>NTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPK</u> <u>DTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT</u> <u>KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA</u> <u>PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGF</u> <u>YPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD</u> <u>KSRWQQGNVFPSCVMHEALHNHYTQKSLSLSPGK</u>	4
anti CD3 (CH2527 VH 3-23(12) VL7- 46(13)) scFv 4.32.63 non-cleavable linker CH2527 VH3 2-VH12 CH1 Fo1R1 16D5 VH CH1 hum Fc knob PG LALA, pETR15603 (FIG. 45C)	<u>QVQLKESGPGLVAPSSQLSITCTVSGFSLTSYGVSWVRQPP</u> <u>GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKQVFLKLNS</u> <u>LQDDTATYYCAKGIITVVDDYIAMDYWGQGTSTVTVSSGG</u> <u>GGSGGGSGGGSGGGSGGGSDIQTQSPASLASVGETVTIT</u> <u>CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS</u> <u>GGSGGTQYSLKINSLSQSEVARYYCOHYSTPYTFGCCTKL</u> <u>EIKGGGGSGGGSGGGSGGGSGGGSGGGSGGGSEV</u> <u>QLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWRQAPG</u> <u>KGLEWVSRIKSKYNNYATYYADSVKGRFTISRDDSNTLYLQ</u> <u>MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVS</u> <u>SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTS</u> <u>WNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYI</u> <u>CNVNHKPSNTKVDKKVEPKSCDGGGSGGGGSEVQLVES</u> <u>GGGLVKPGGSLRLSCAASGFTFSTYAMNWRQAPGKGLE</u> <u>WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSNTLYLQMN</u> <u>LKTEDTAVYYCTTPWEWSWYDYGQGLTVTVSSASTKGPS</u> <u>VFPLAPSSKSTSGGTAALGCLVKDYFPEPTVTSWNSGALTS</u> <u>GVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPS</u>	5

EXAMPLARY SEQUENCES		
	NTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPK DTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD KSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	
anti CD3 (CH2527 VH_3-23(12) VL7- 46(13)) scFv15 non- cleavable linker CH2527 VH3_23-VH12 CH1 FcR1 16D5 VH CH1 hum Fc knob PG LALA, pETR14759 (FIG. 45D)	<u>QIQLVQSGPELKKPGETVIRISCKASGYTFTDYSIHVWKQAPG</u> <u>KCLKWMGWINTETGEPAYADDFKGRFAFSLETSASTAYLQI</u> <u>NNLKNE DTATFFCAHPYDYDVLVDYWGQGTSTVTVSSGGGGS</u> <u>GGGSGGGSGGGSGGGSDTVLTQSPASLGLVSLGQRATISCRA</u> <u>SKSVSTSNYSYIHVYQ</u> <u>QKPGQPPKLLIKYVSYLESGVPPARFSGSGSGTDTLNIHPVE</u> <u>EEDAATYYCQHSREFPWTFGCGTKLEIKGGGSGGGGSGG</u> <u>GGSGGGGGGGSGGGSGGGGSEVQLLES GGGLVQPGGS</u> <u>LRLSCAASGFTFSTYAMNWRQAPGKGLEWVSRIRSKYNN</u> <u>YATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCV</u> <u>RHGNFGNSYVSWFAYWGQGLTVTVSSASTKGPSVFPLAPS</u> <u>SKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPA</u> <u>VLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKK</u> <u>VEPKSCDGGGSGGGGSEVQLVESGGGLV KPGGSLRLSC</u> <u>AASGFTFSNAWMSWVRQAPGKGLEWVGRIKSKTDGGTDDY</u> <u>AAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTPWE</u> <u>WSWYDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA</u> <u>ALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYS</u> <u>LSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKT</u> <u>HTCPPCPAPEAAGGPSVFLFPPKPKDTLMI SRTPEVTCVVVD</u> <u>VSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSV</u> <u>LTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQ</u> <u>VYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE</u> <u>NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVM</u> <u>HEALHNHYTQKSLSLSPGK</u>	6
MK062 Protease linker	GGGSGGGGSRQARVVNGGGGSGGGSGGGGSGGGGS	7
Combined NF9/Mat5 linker	GGGSGVHMLPLGLFGPGRSRGSPFGGGGS	8
Combined MK062 MMP9	GGGSGGGGSRQARVVNGGGGSGVPLSLYSGGGGSGGGGSGGS	9
Combined MK062 MMP9	GGGSGGGGSRQARVVNGVPLSLYSGGGGSGGGGSGGGGS	10
H2527 CDR H1 Kabat	TYAMN	11
CH252 CDR H2 Kabat	RIRSKYNNYATYYADSVKG	12
CH252 CDR H3 Kabat	HGNFGNSYVSWFAY	13
FcR1 CDR H1 Kabat	NAWMS	14
FcR1 CDR H2 Kabat	RIKSKTDGGTDDYAAPVKG	15
FcR1 CDR H3 Kabat	PWEWSWYDY	16
CLC CDR1 L1 Kabat	GSSTGAVTTSNYAN	17
CLC CDR L2 Kabat	GTNKRAP	18
CLC CDR L3 Kabat	ALWYSNLWV	19
Anti-ID 4.15.64 CDR H1 Kabat	DYSIH	20
Anti-ID 4.15.64 CDR H2 Kabat	WINTETGEPAYADDFKG	21
Anti-ID 4.15.64 CDR H3 Kabat	PYDYDVLVDY	22
Anti-ID 4.15.64 CDR L1 Kabat	RASKSVSTSNYSYIH	23

-continued

EXAMPLARY SEQUENCES		
Anti-ID 4.15.64 CDR L2 Kabat	YVSYLES	24
Anti-ID 4.15.64 CDR L3 Kabat	QHSREFPWT	25
Anti-ID 4.32.63 CDR H1 Kabat	SYGVS	26
Anti-ID 4.32.63 CDR H2 Kabat	IIWGDGSTNYHSALIS	27
Anti-ID 4.32.63 CDR H3 Kabat	GITTVDDYYAMDY	28
Anti-ID 4.32.63 CDR L1 Kabat	RASENIDSYLA	29
Anti-ID 4.32.63 CDR L2 Kabat	AATFLAD	30
Anti-ID 4.32.63 CDR L3 Kabat	QHYSTPYT	31
anti HER1 (GA201 heavy chain, pUC-Exp- GA201-HC) (FIG. 45E)	QVQLVQSGAEVKKPGSSVKVSCKASGFTFTDYKIHWRQAP QGLEWMGYFNPNSGYSTYAQKPGQGRVTITADKSTSTAYM ELSSLRSEDTAVYYCARLSPGGYYVMDAWQGTTTVVSSA STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICN VNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFL FPPKPKDTLMI SRTPETCVVVDVSHEDPEVKFNWYVDGVE VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKV NKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLT CLVKGFYPSDIAVEWESNGOPENNYKTTPPVLDSDGSFFLY SKLTVDKSRWQQGNVFCSCVMHEALHNYHTQKSLSLSPGK	32
anti HER1 (GA201 light chain, pUC-Exp-GA201- LC) (FIG. 45F)	DIQMTQSPSSLSASVGDRVTITCRASQGINNYLNWYQQKPG KAPKRLIYNTNNLTGTGVPSPRFGSGSGTEFTLTISLQPEDF ATYYCLQHNFPFTFGQGTKEIKRTVAAPSVFIFPPSDEQLKS GTASVVCCLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGEC	33
anti HER1 (anti-GA201 VH-VL scFv MMP cleavable linker G4S GA201 light chain, pUC- I_GA201 MMP_LC) (FIG. 45G)	EVQLEQSGPVLVKPGTSVKMSCKASGYTFTDYYINWIIQSHG KCLEWIGVINPDSSGTDYNNFNGKATLTVDKSSSTAYMELT SLTSEDSAVYYCARRDSYGFQYWGQGTTLTVSSGGGGSGG GGSGGGGGGGGSDIVLTQTPKFLVPAVDRIITMTCKASLS VTNDVAWYQQKPGQSPKLLLYASNRNAGVPDRFTGSGYG TDFTFTITTLQAEDLAVYFCQDYTSPTPTFGCGTKLEIRGGG GGGGGGGSLGLWSQGGGGGGGGGGGGGGGGDIQMTQ SPSSLSASVGDRVTITCRASQGINNYLNWYQQKPGKAPKRLI YNTNNLTGTGVPSPRFGSGSGTEFTLTISLQPEDFATYYCLQ HNSFPTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVV CLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDY SLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC	34
MMP Protease linker	GGGGSGGGSGPLGLWSQGGGGGGGGGGGGGGGGGGGG	35
Protease recognition site 1	RQARVVNG	36
Protease recognition site 2	VHMPGLGLGPGRSRGSFP	37
Protease recognition site 3	RQARVVNGXXXXVPLSLYSG	38
Protease recognition site 4	RQARVVNGVPLSLYSG	39
Protease recognition site 5	PLGLWSQ	40

-continued

EXAMPLARY SEQUENCES		
4.15.64 Anti-idiotypic scFv	<u>QIQLVQSGPELKKPGETVIRISCKASGYTFTDYSIHVWKQAPG</u> <u>KCLKWMGWINTETGEPAYADDFKGRFAFSLETSASTAYLQI</u> <u>NNLKNEATATFFCAHPYDYDVLVDYWGQTSVTVSSGGGGS</u> <u>GGGSGGGSGGGSGGSDTVLTQSPASLGVSLGQRATISCR</u> <u>SKSVSTSNYSYIHVYQKPGQPKLLIKYVSYLESGVPARFS</u> <u>SGSGTDFTLNIHPVEEDAATYYCQHSREFPWTFGCGTKL</u> <u>EIK</u>	41
4.32.63 Anti-idiotypic scFv	<u>QVQLKESGPGLVAPSQSLITCTVSGFSLTSYGVSWVRQPP</u> <u>GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSQVFLKLS</u> <u>LQDDDTATYYCAKGIITVVDDYAMDYWGQTSVTVSSGG</u> <u>GGSGGGSGGGSGGGSDIQTQSPASLSASVGETVTIT</u> <u>CRASENIDSYLAWYQKQKSPQLLVYAATFLADDVPSRFS</u> <u>SGSGTQYSLKINLSQSEDVARYYCQHYSTPYTFGCGTKL</u> <u>EIK</u>	42
Anti-CD3 variable heavy chain (VH)	<u>EVQLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRQA</u> <u>PGKGLEWVSRIRSKYNNYATYYADSVKGRTISRDDSKNTLY</u> <u>LQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQTLVT</u> <u>VSS</u>	43
CD3 heavy chain (VH)_CDR1	TYAMN	44
CD3 heavy chain (VH)_CDR2	RIRSKYNNYATYYADSVKG	45
CD3 heavy chain (VH)_CDR3	HGNFGNSYVSWFAY	46
Anti-FolR1 16D5 variable region	<u>EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQA</u> <u>PGKGLEWVGRIKSKTDGGTDDYAAPVKGRFTISRDDSKNTLY</u> <u>YLQMNSLKTEDTAVYYCTTPWEWSYDYWGQTLVTVSS</u>	47
anti-idiotypic GA201 CDR H1 Kabat	DYYIN	48
anti-idiotypic GA201 CDR H2 Kabat	VINPDSGGTDYNQNFKG	49
anti-idiotypic GA201 CDR H3 Kabat	RDSYGFDY	50
anti-idiotypic GA201 CDR L1 Kabat	KASLSVTNDVA	51
anti-idiotypic GA201 CDR L2 Kabat	YASNRNA	52
anti-idiotypic GA201 CDR L3 Kabat	QQDYTSPTT	53
hu CD3E	<u>MQSGTHWRVLGLCLLSVGWVGQDGNEMGGITQTPYKVS I</u> <u>SGTTVILTCPPYPSGELLWQHNDKNIGGDEDDKNIGSDEHL</u> <u>SLKEFSELEQSGYYVCYPRGSKPEDANFYLYLRARVCENC</u> <u>MEMDVMSVATIVIVDICITGGLLLVYYWSKNRKAKAKPVTRG</u> <u>AGAGGRQRGQNKERPPVPNPDIPIKRGQRLYSGLNQR</u> <u>RI</u>	54
LC Common light chain pETR13197 V region	<u>QAVVTQEPSTLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK</u> <u>PGQAFRGLIGGTNKRAPGTPARFSGSLGGAALTLGSAQP</u> <u>EDEAEYYCALWYSNLWVFGGKTLTVL</u>	55
GA201 CDR H1 Kabat	DYKIH	56
GA201 CDR H2 Kabat	YFNPNSGYSTYAQKFQG	57
GA201 CDR H3 Kabat	LSPGGYYVMDA	58

-continued

EXAMPLARY SEQUENCES		
GA201 CDR L1 Kabat	RASQGINNYLN	59
GA201 CDR L2 Kabat	NTNNLQT	60
GA201 CDR L3 Kabat	LQHNSFPPT	61
Construct	DNA Sequence	SEQ ID No
LC Common light chain pETR13197	CAGGCCGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGC GGCACCGTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACC ACCAGCAACTACGCCAACTGGGTGCAGGAAAAGCCCGCCAGGCC TTCAGAGGACTGATCGCGGCACCAACAAGAGAGCCCTGGCACCC CCTGCCAGATTACGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTG ACACTGTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGC GCCCTGTGGTACAGCAACCTGTGGGTGTTTCGCGGAGGCACCAAG CTGACAGTCTTAGGTCAACCCAAAGGCTGCCCCAGCGTGACCCTG TTCCTCCAGCAGCGAGGAACTGCAGGCCAACAAAGGCCACCTG GTCTGCCTGATCAGCGACTTCTACCCAGGCGCCGTGACCGTGGCC TGGAAGGCCGACAGCAGCCCCGTGAAGGCCGCGGTGGAGACCAC CACCCCCAGCAAGCAGAGCAACAACAAGTACGCCCGCCAGCAGCTA CCTGAGCCTGACCCCGAGCAGTGAAGAGCCACAGGTCTTACAG CTGCCAGGTGACCCAGAGGCGAGCACCGTGGAGAAAACCGTGG CCCCACCGAGTGACAGCTGA	62
anti CD3 (CH2527 VH_3-23 (12) VL7-46 (13)) scFv15-Matriptase MK062 CH2527 VH3_23-VH12 CH1_Fo1R1 16D5 VH CH1 hum Fc knob PG LALA, pETR15422 (FIG. 45H)	CAGATCCAGCTGGTGCAGAGCGGCCCTGAGCTGAAGAAAACCCGGC GAGACAGTGCAGGATCAGCTGCAAGGCCAGCGGCTACACCTTCACC GACTACAGCATCCACTGGGTCAAGCAGGCCCTTGGCAAGTGCCCTG AAGTGGATGGGCTGGATCAACACCGAGACAGGCGAGCCCGCTAC GCCGACGATTTCAAGGGCAGATTTCGCTTTCAGCCTGGAAACAGC GCCAGCACCGCCTACCTGCAGATCAACAACCTGAAGAAGCAGGAG ACCGCCACCTTTTTCTGCGCCACCCCTACGACTACGACGTGCTG GATTATTGGGGCCAGGGCACCAGCGTGACCGTGTCTAGCGGAGGC GGAGGATCTGGCGCGGAGGAAGTGGCGGAGGGGATCTGGGG GAGGCGGATCTGATACCGTGTGACACAGAGCCCTGCCAGCCTGG GAGTGTCTTGGGACAGAGAGCCACCATCAGCTGTGCGGCCAGCA AGAGCGTGTCCACCAGCAACTACAGCTATATCCACTGGTATCAGCA GAAGCCCGGCCAGCCCCCAAGTGTCTGATCAAATACGTGTCTTA CCTGGAAGAGCGCGTGCCCGCCAGATTTCTGGCTCTGGCAGCGG CACCGACTTCACCTGAACATCCACCCCGTGGAAGAGGAAGATGC CGCCACCTACTACTGCCAGCACAGCAGAGTTCCCTTGGACCTTC GGTGCAGCCACCAAGCTGGAATCAAAAGCGGGGAGGCTCCCG AGGCGCGGAAGTAGACAGGCCAGAGTCTGTAAACGGGGAGGGG GGGGAAGTGGGGCGGAGGAGTGGGGGGGAGGATCCGAGGT GCAGCTGTGGAATCTGGCGCGGAGTGGTGACGCTGGCGGAT CTCTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCACCT ACGCCATGAATGGGTGCGCCAGGCCCTGGCAAAGGCCCTGGAAT GGGTGTCTCCGATCAGAAGCAAGTACAACAATACGCCACCTACTA CGCCGACAGCGTGAAGGCCGCTTACCATCAGCCGGGACGACA GCAAGAACACCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGG ACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGAACA GCTATGTGTCTTGGTTTGCTTACTGGGGCCAGGGCACCTCTGTA CCGTGTCAAGCGCTAGCACAAGGGCCCTAGCGTGTTCCTCTGG CCCCAGCAGCAAGAGCACAAAGCGCGGAACAGCCGCCCTGGGC TGCCTCGTGAAGGACTACTTCCCCAGGCCGTGACAGTGTCTTGG AACAGCGGAGCCCTGACAAGCGCGGTGCACACCTTCCCTGCCGTG CTGCAGAGCAGCGGCTGTACTCCCTGAGCAGCGTGGTCAACGTG CCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAAC CACAAGCCCAGCAACACCAAGTGGACAAGAAGGTGGAGCCCAAG AGCTGTGATGGCGGAGGAGGGTCCGAGGCGGAGGATCCGAGGT GCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGGTTT CCTGCGTCTGAGCTGCGCGGCTTCCGATTACCTTCTCCAACGC GTGGATGAGCTGGGTTTCGCGAGGCCCGGGCAAGGCCTCGAGT GGGTTGGTCTGATCAAGTCTAAACTGACGGTGGCACCAACGATT CGCGGCTCCAGTTAAGGTCGTTTACCATTTCGCGCAGCATAGC AAAACACTCTGTATCTGCAGATGAACCTCTGAAAACTGAAGACAC CGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTGGTACGAT TATTGGGGCCAGGGCACGCTGGTTACGGTGTCTAGCGCTAGTACC AAGGGCCCCAGCGTGTTCCTCCCTGGCACCCAGCAGCAAGAGCACA TCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAAGACTACTTC CCCGAGCCCGTGACCGTGTCTTGGAACTCTGGCGCCCTGACCAGC GGCGTGACACCTTTCCAGCCGTGTGACAGCAGCGGCTGTAC TCCCTGTCTCCGTGGTGACCGTGCCCTTAGTCTCCCTGGGAACA	63

-continued

EXAMPLARY SEQUENCES

	CAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACACCAAAGT CGATAAGAAAGTCGAGCCCAAGAGCTGCGACAAAACCTCACACATG CCCACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAATCTT CCTCTTCCCCCAAACCAAGGACACCTCATGATCTCCCGGACC CCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCTT GAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAAT GCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCAGTACCGT GTGGTCAGCGTCTTACCGTCTTGCAACAGGACTGGTGAATGGC AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCC ATCGAGAAAACCATCTCAAAGCCAAGGGCAGCCCCGAGAACCA CAGGTGTACACCTGCCCCCATGCGGGATGAGCTGACCAAGAAC CAGGTGAGCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCGAC ATCGCGTGGAGTGGGAGAGCAATGGGCAGCGGAGAACAACTAC AAGACCAGCCTCCCGTGTGAGCTCCGACGGCTCTTCTTCTCT ACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAAC GTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA	
FolR1 16D5 VH CH1 Fc hole P329G LALA HRYF, pETR15214	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGC GGTTCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCC AACCGCTGGATGAGCTGGGTTCGCCAGGCCCCGGGCAAGGCCT CGAGTGGGTGGTGGTATCAAGTCTAAAACCTGACGGTGGCACCAC GGATTACGCGCTCCAGTTAAAGGTGTTTTACCATTTCCCGCGAC GATAGCAAAAACACTCTGTATCTGCAGATGAACCTCTGAAAACCTGA AGACACCGCAGTCTACTACTGTACTACCCGTGGGAATGGTCTTGG TACGATTATTGGGGCCAGGGCAGCGTGGTTACGGTGTCTTCCGCT AGCACCAGGGCCCCCTCCGTGTTCCCGCTGGCCCCAGCAGCAAG AGCACCAGCGCGGCAGCAGCGCTCTGGGCTGCTGGTCAAGGA CTACTTCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCT GACCTCCGGCGTGACACCTTCCCGCGCTGCTGCAGAGTTCTGG CCTGTATAGCCTGAGCAGCGTGGTCAACGTGCTTCTAGCAGCCT GGCACCAGACCTACATCTGCAACGTGAACCAAGCCAGCAA CACCAGGTGGACAAGAAGTGGAGCCCAAGAGCTGCGCAAAAC TCACATGCCCCACCGTGCCAGCAGCTGAAGCTGCAAGGGGAGC GTGATCTTCTCTTCTCCCCCAAAACCAAGGACACCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCTGGAG GTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGC ACGTACCGTGTGGTACGCTCTTCCCGTCTGCACAGGACTGG CTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGCCCTC GGCGCCCCATCGAGAAAACCATCTCAAAGCCAAGGGCAGCCC CGAGAACCACAGGTGTGCACCCCTGCCCCCATCCCGGGATGAGCTG ACCAAGAACCAGGTGAGCCTCTCGTGCAGCTCAAAGGCTTCTATC CCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCC TTCTTCTCTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAG CAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCAC AACCGCTTACCGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA	64
anti CD3 (CH2527 VH 3-23 (12) VL7-46 (13)) scFv 4.32.63 Matriptase MK062 CH2527 VH3 23-VH12 CH1 FolR1 16D5 VH CH1 hum Fc knob PG LALA, pETR15599 (FIG. 451)	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAAGCTGACC AGCTACGGCGTGTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGGCATCACCAACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGAGGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCGAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTGCGGCACCAAGCTGGAATCAAAGCGGGGGAGGCTCCG GAGGCGGCGGAAGTAGACAGGCAGAGTCTGTAACGGGGGAGGG GGGGGAAGTGGGGCGGAGGAGTGGGGCGGAGGATCCGAGG TGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCTGGCGGA TCTCTGAGACTGAGCTGTGCCGCCAGCGCTTCACTTTCAGCACC TACGCCATGAACGGGTGCGCCAGGCCCTGGCAAAGGCCTGGAA TGGGTGTCCCGGATCAGAAGCAAGTACAACAACTACGCCACCTACT ACGCCGACAGCGTGAAGGGCCGTTTACCATCAGCCGGGACGAC AGCAAGAACACCTGTACCTGCAGATGAACAGCCTGCGGGCCGAG	65

-continued

EXAMPLARY SEQUENCES

	GACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAAC AGCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTG ACCGTGTCAAGCGCTAGCACAAAGGGCCCTAGCGTGTCCCTCTG GCCCCAGCAGCAAGAGCACAAAGCGGCGAACAGCCGCCCTGGG TGCCCTCGTGAAGGACTACTTCCCCGAGCCCGTGACAGTGTCTTG GAACAGCGGAGCCCTGACAAGCGGCGTGACACCTTCCCTGCCGT GCTGCAGAGCAGCGGCTGTACTCCCTGAGCAGCGTGGTACCCTG GCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAA CCACAAGCCAGCAACACCAAGTGGACAAGAAGGTGGAGCCCAA GAGCTGTGATGGCGGAGGAGGGTCCGGGGCGGAGGATCCGAG GTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGGT TCCCTGCGTCTGAGCTGCGCGGCTTCCGGGTTACCTTCTCCAAC GCGTGGATGAGCTGGGTTCCGCCAGGCCCCGGGCAAAGGCTCGA GTGGGTGGTCTGATCAAGTCTAAACTGACGGTGGCACCCAGGA TTACGCGGCTCCAGTTAAAGGTGTTTTACCATTTCCCGGACGAT AGCAAAACACTCTGTATCTGCAGATGAACCTCTGAAACTGAAG ACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGTA CGATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTAGCGCTAG TACCAAGGGCCCCAGCGTGTCCCCCTGGCACCCAGCAGCAAGAG CACATCTGGCGAACAGCCGCTCTGGGCTGTCTGGTGAAGACTA CTTCCCCGAGCCCGTGACCGTGTCTTGGAACTCTGGCGCCCTGAC CAGCGCGTGACACACCTTTCCAGCCGTGTCTGCAGAGCAGCGGCT GTACTCCCTGTCTCGTGGTACCGTGCCCTCTAGCTCCCTGGG AACACAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACACCA AAGTCGATAAGAAAGTCGAGCCCAAGAGCTGCAGCAAACTCACAC ATGCCACCGTGCCAGCACCTGAAGCTGCAGGGGGACCGTCAGT CTTCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCCCGG ACCCCTGAGGTACATGCGTGGTGGTGGAGTGGAGCCACGAAGAC CTGAGGTCAAGTTCAACTGGTACGTGGACGCGGTGGAGTGCAT AATGCCAAGACAAGCCGCGGGAGGAGCAGTACAACAGCAGTAC CGTGTGGTCAAGCTCTACCCGCTCTGCACCAAGGACTGGTGAAT GGCAAGGAGTACAAGTGCAGGTCTCCAACAAGCCCTCGCGGCC CCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAA CCACAGGTGTACACCCCTGCCCCATGCCGGGATGAGCTGACCAAG AACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGC GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA CTACAAGACACGCTCCCGTGTGGACTCCGACGCTCTCTTCTT CCTCTACAGCAAGCTCACCGTGACAGAGCAGGTGGCAGCAGGG GAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCAC TACACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	
anti CD3 (CH2527 VH ₃ -23(12) VL ₇ -46(13)) scFv 4.32.63 non- cleavable linker CH2527 VH ₃ -23-VH12 CH1 Fc1R1 16D5 VH CH1 hum Fc knob PG LALA, pETR15603 (FIG. 45J)	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTACGCTGACC AGCTACGGCGTGTATGGGTGCGCCAGCCTTCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAAGGCGATCACCAACCGTGGTGGACGAC TACTACGCTATGAGTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTGCCGCCACCAAGCTGGAAATCAAAGCGGGGGAGGCTCCG GAGGCGGGAAGTGGAGGCGGCGGAAGTGGCGGAGGCGGAGG GGGGGGAAGTGGGGGCGGAGGAGTGGGGGGGAGGATCCGAG GTGCAGCTGTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGG ATCTCTGAGACTGAGCTGTGCCCGCAGCGGCTTACCTTTCAGCAC CTACGCCATGAAGTGGGTGCGCCAGGCCCTGGCAAAGGCTTGA ATGGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACCTAC TACGCCGACAGCGTGAAGGGCCGGTTACCATCAGCCGGGACGA CAGCAAGAACACCCCTGTACCTGCAGATGAACAGCCTGCGGGCCGA GGACACCGCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAA CAGCTATGTGTCTTGGTTTGGCTACTGGGGCCAGGGCACCCCTCGT GACCGTGTCAAGCGCTAGCACAAAGGGCCCTAGCGTGTTCCTCT GGCCCCCAGCAGCAAGAGCACAAAGCGGCGGAACAGCCGCCCTGG GCTGCCCTCGTGAAGGACTACTTCCCCGAGCCGTGACAGTGTCTT GGAACAGCGGAGCCCTGACAAGCGGCGTGACACCTTCCCTGCC GTGCTGCAGAGCAGCGGCTGTACTCCCTGAGCAGCGTGGTCAAC GTGCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCACCGTG	66

-continued

EXAMPLARY SEQUENCES

anti CD3 (CH2527 VH 3-23 (12) VL7-46 (13)) scFv15 non- cleavable linker CH2527 VH3_23-VH12 CH1 F01R1 16D5 VH CH1 hum Fc knob PG LALA, pETR14759 (FIG. 45K)	AACCACAAGCCAGCAACACCAAGTGGACAAGAAGTGGAGCCC AAGAGCTGTGATGGCGGAGGAGGTTCCGAGGCGGAGGCTCCGA GGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAAACGGGCGG TTCCCTGCGTCTGAGCTGCGCGGCTTCCGATTACCTTCTCCAAC CGTGGATGAGCTGGGTTCCGAGGCCCGGGCAAGGCCTCGA GTGGGTTGGTCTGATCAAGTCTAAACTGACGGTGGCACCACGGA TTACGCGGCTCCAGTTAAAGGTCGTTTACCATTTCGCGGACGAT AGCAAAAACACTCTGTATCTGCAGATGAACCTCTCTGAAACTGAAG ACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTA CGATTATTGGGGCCAGGGCAGCTGGTTACGGTGTCTAGCGCTAG TACCAAGGGCCCCAGCGTGTTCGCCCTGGCACCAGCAGCAAGAG CACATCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAGACTA CTTCCCGAGCCCGTGACCGTGTCTTGAACCTCTGGCGCCCTGAC CAGCGCGTGCACACCTTTCAGCCGTGCTGCAGAGC AGCGGCCTGTACTCCCTGTCTCCGTGGTACCGTGCCCTCTAGC TCCCTGGGAACACAGACATATATCTGTAATGTCAATCACAAGCCTTC CAACACCAAGTCGATAAGAAAGTCGAGCCCAAGAGCTGCGACAA AACTCACACATGCCCCACCGTGCCCGAGCACCTGAAGCTGCAGGGGG ACCGTCAGTCTTCTCTTCCCCCAAAACCAAGGACACCCCTCATG ATCTCCCGACCCCTGAGGTCAATGCGTGGTGGTGGAGCTGAGC CAGGAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTG GAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAAC AGCAGTACCGTGTGGTACGCGTCTCACCGTCTGCACAGGAC TGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCC CTCGGCGCCCCATCGAAGAAACCATCTCCAAGCCAAAGGGCAG CCCCGAGAACCACAGGTGTACACCTGCCCCCATGCCGGGATGAG CTGACCAAGAACCAGGTGACCTGTGGTGGCTGGTCAAAGGCTTC TATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCGAGC GGAGAACAACACTACAAGACACGCTCCCGTGTGGACTCCGACGG CTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGG CAGCAGGGGAACGTCTTCTCATGCTCCGTGATGATGAGGCTCTG CACAACTACTACGCGAGAAGGCTCTCCCTGTCTCCGGTAAAT GA	67

-continued

EXAMPLARY SEQUENCES

	AAAAACAACCTGTATCTGCAGATGAACTCTCTGAAAACTGAAGACAC CGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACGAT TATTGGGGCCAGGGCACGCTGGTTACGGTGTCTAGCGCTAGTACC AAGGGCCCCAGCGTGTTCCTCCCTGGCACCAGCAGCAAGAGCACA TCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAAGACTACTTC CCGAGCCCCGTGACCGTGTCTTGAAGCTCTGGCGCCCTGACCAGC GGCGTGACACCTTTCCAGCCGCTGCTGCAGAGCAGCGCCTGTAC TCCCTGTCTCCGTGGTACCCGTGCCCTCTAGCTCCTTGGGAACA CAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACACCAAAGT CGATAAGAAAAGTCGAGCCCAGAGCTGCGACAAAACCTCACACATG CCCACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAGTCTT CCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCCCGGACC CCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCTT GAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAAT GCCAAGACAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT GTGGTCAGCGTCTTACCCTGCTGCACAGGACTGGCTGAATGGC AAGGAGTACAAGTGAAGGTCTCCAACAAGCCCTCGGCGCCCC ATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCA CAGGTGTACACCTGCCCCCATGCCGGGATGAGCTGACCAAGAAC CAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCGAC ATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTAC AAGACCAGCCTCCCGTGTGGACTCCGACGGCTCCTTCTTCTCTCT ACAGCAAGCTCACCGTGGACAAAGAGCAGGTGGCAGCAGGGGAAC GTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA CGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	
MK062 Protease linker	GCGGGGGAGGGCTCCGGAGGCGGCGGAAGTAGACAGGCCAGAG TCGTGAACGGGGGAGGGGGGGAAGTGGGGCGGAGGCAGTGG GGGCGAGGATCC	68
anti HER1 (GA201 heavy chain, pUC- Exp-GA201- HC)	CAGGTGCAGCTGGTCCAGAGCGGCGCCGAGGTGAAGAAACCCGG GTCTCTGTCAAGGTGTCATGCAAGGCTAGCGGATTACCTTTACA GACTACAAAATCCACTGGGTTAGGCAGGCACCTGGCCAAGGACTC GAATGGATGGGGTATTTCAACCCAAATTCCGGCTACTCTACCTATG CCCAGAAGTTTCAGGGAAGAGTGACTATTACAGCTGATAAGAGTAC CAGCACTGCATACATGGAGCTGTCTCTCTTCTCGCTCAGAGGACACC GCCGTCTACTATTGTGCTCGGCTGAGCCCCGGTGGCTACTATGTG ATGGATGCATGGGGCAGGGAACAACCGTAAACAGTGTCTCTGCG TCGACTAAGGGCCCTTCAGTTTTTCCACTCGCCCCAGTAGCAAGT CCACATCTGGGGGTACCGCTGCCCTGGGCTGCCTTGTGAAGACT ATTTCCCTGAACCAAGTCACTGTGTGATGGAATAGCGGAGCCCTGAC CTCGGGTGTACACACATTTCCCGCTGTGTGACGTCTAGTGGCCTG TACAGCCTCTCCTCTGTTGTGACCGTCCCTTCAAGCTCCTTGGGGA CACAGACCTATATCTGTAACGTGAATCATAAGCCATCTAACACTAAA GTAGATAAAAAAGTGGAGCCCAAGAGTTGCGACAAAACACACACCT GTCCCCCTTGCCAGCCCCCGAGCTTCTGGGAGGCCCTAGCGTCT TTCTCTTCCCAACCAAGCCTAAGGATACTCTGATGATATCCAGGAC CCCAGAAGTTACATGCGTGGTCTGGACGTCTCACACGAGGACCC CGAAGTGAAATTTAACTGGTACGTTGATGGTGTGGAAGTCCATAAT GCCAAGACCAGCCTAGAGAGGAGCAATACAACAGTACATATCGC GTGGTAAGCGTGTGACCGTTCTCCACAGGACTGGCTCAATGGG AAAGAATACAAGTGTAAGTGTCCAACAAGCTCTGCCAGCACCCA TCGAGAAGACTATTTCTAAGGCCAAAGGCCAGCCCCGGGAGCCTC AGGTCTATACACTTCCACCTCAAGGGATGAAGTACCAAGAACCA AGTGAGCTTGACTTGCTGGTAAAGGGTTCTACCCCTCCGACATC GCTGTGGAGTGGGAGTCTAATGGACAACCAAGAAAACAATTACAAAA CCACACCCCTGTCTCGACAGTGATGGCAGCTTTTTCTGTATAG CAAACTTACGTTGACAAGTCCAGATGGCAGCAGGGAAACGTGTTC TCATGTAGCGTCATGCACGAAGCTTTCATAACCACTACACACAGA AAAGCCTCAGCCTGAGTCCAGGGAAG	69
anti HER1 (GA201 light chain, pUC- Exp-GA201- LC)	GACATCCAATGACCCAGTCACCTAGTAGCCTCTCCGCCTCTGTTG GCGACAGGGTGACAATTACATGCAGAGCTTCACAGGGTATCAACAA TTACCTGAACTGGTATCAGCAGAAACAGGGAAGGCCCAAGCG CTTGATATATAACACCAATAACCTGCAAACTGGCGTCCCTAGCCGG TTCTCCGGATCTGGTAGTGGCACCGAATTTACACTCACCATCAGCT CCTTGCAGCCAGAGGATTTTCGCACATACTATTGTCTTACGCATAA TTCTTTCCCACTTTGGGCAAGGAACCTAACTGGAGATTAAGCGT ACTGTGCGCCGCTCCCTCTGTGTTCAATTTTCTTCCAAGTGATGAGC AGCTCAAAGCGGTACCGCATCCGTTGTGTGCTGCTTAACAACCTT CTATCCCGGGAAGCAAGTCCAATGGAAGGTGGACAATGTCTCT GCAGTCAGGAAACAGTCAGGAGAGCGTAACCGAGCAGGATTCCAA AGACTCTACTACTCATTGAGCTCCACCTGACACTCTCTAAGGCA	70

-continued

EXAMPLARY SEQUENCES		
	GACTATGAAAAGCATAAAGTGTACGCCTGTGAGGTTACCCACCAGG GCCTGAGTAGCCCTGTGCACAAAGTCCTTCAATAGGGGAGAGTGC	
HER1 (anti- GA201 VH- VL scFv MMP cleavable linker G4S GA201 light chain, pUC- I_GA201_MMP- LC)	GAGGTTTCAGCTGGAGCAGTCAGGACCTGTGCTGGTGAAGCCTGGG ACTTCAGTGAAGATGTCCTGTAAAGCTTCTGGATACACATTCAGT ACTACTATATAAACTGGATAATACAGAGCCATGGAAAGTGTCTTGAG TGGATTGGAGTTATTAATCCTGACAGCGGTGGTACTGACTACAACC AGAACTTCAAGGGCAAGGCCACATTGACTGTTGACAAAGTCTCCAC CACAGCCTACATGGAACCTACTAGCCTGACATCTGAGGACTCTGCA GTCTATTATTGTGCAAGAAGGGATTCTTACGGCTTTGACTACTGGG GCCAAGGCACCACTCTCACAGTCTCCTCAGGCGGAGGTGGCTCAG GGGGAGGCGGTAGCGGCGGAGGTGGCTCAGGGGGAGGCGGTAG CGACATTGTGCTGACCCAGACTCCCAAATTCCTGCTTGTGCCAGCA GGAGACAGGATTACCATGACCTGCAAGGCCAGTCTGAGTGTGACT AATGATGTAGCTTGGTATCAACAGAAACCAGGGCAGTCTCCTAAAC TGCTGTTATACTATGTCATCCAATCGCAACGCTGGAGTCCCTGATCG CTTCACTGGCAGTGGATATGGGACGGATTTCACCTTACCATCACC ACTTTCAGGCTGAAGACCTGGCAGTTTATTTCTGTCAGCAGGATT ATACCTCTCCTCCGACGTTCTGGTGTGGCACCAGCTAGAAATCCG TGGTGGCGGCGGTTCTGGCGGAGGGGTTCTGGCCCCCTGGGCG TATGGAGCCAGGTTGGCGGCGGTTCTGGCGGAGGGGTTCTGGC GGTGGTGGCTCTGGCGGTGACATCCAAATGACCCAGTCACCTAGT AGCCTCTCCGCTCTGTTGGCGACAGGGTGACAATTACATGCAGA GCTTCACAGGGTATCAACAATTACCTGAAGTGGTATCAGCAGAAAC CAGGGAAGGCCCCCAAGCGCTTGATATATAACCAATAACCTGCA AACTGGCGTCCCTAGCCGGTTCTCCGATCTGGTAGTGGCACCAG ATTTACACTCACCATCAGCTCCCTGCAGCCAGAGGATTTCCGCCACA TACTATTGTCTTACAGATAATTCTTCCACCTTTGGGCAAGGAAC TAAACTGGAGATTAAGCGTACTGTGCGCGCTCCCTCTGTGTTTATT TTTCTCCAAGTATGAGCAGCTCAAAGCGGTACCGCATCCGTTG TGTGCTGCTTAAACACTTCTATCCCGGGAAGCCAGGTCCAATG GAAGGTGGACAATGCTCTGCAGTCAGGAACAGTCAGGAGAGCGT AACCAGCAGGATTCCAAAGACTCTACTTACTCATTGAGCTCCACC CTGACACTCTCTAAGGCAGACTATGAAAAGCATAAAGGTACGCCT GTGAGGTTACCCACCAGGGCTGAGTAGCCCTGTGACAAAGTCCT TCAATAGGGGAGAGTGC	71
Construct	Amino acid Sequence	SEQ ID No
anti CD3 (CH2527 VH_3-23(12) VL7- 46(13)) scFv 4.32.63 MMP9 Matriptase MK062 CH2527 VH3_23-VH12 CH1 FolR1 16D5 VH CH1 hum Fc knob PG LALA, pETR16546 (FIG. 45L)	QVQLKESGPGLVAPSSQLSITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQDDDTATYYCAKGIITVDDYYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSGGDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLSQSEVARYYCYHYSTPYTPFCGKTL EIKGGGSGVHMPGLGFLGPRQARVVNNGGGSGGGSEVQ LLESGLLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGK GLEWVSRIISKYNNYATYYADSVKGRFTISRDDSKNTLYLQ MNSLR AEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVS SASTKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPKSCDGGGSGGGGSEVQLVES GGGLVKPGGSLRLSCAASGFTFSNANMSWVRQAPGKGLE WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMN LKTEDTAVYYCTTPWEWSWYDYWGQGLVTVSSASTKGPS VFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVSWNSGALT GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPS NTKVDKKVEPKSCDKTHTCPCPAPEAAGGSPVFLFPPKPK DTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD KSRWQQGNVFPSCSVMEALHNHYTQKSLSLSPGK	72
FolR1 16D5 HC CH2527-VH3_23-12 HC Fc knob PG LALA, pCON999 (FIG. 45M)	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNANMSWVRQA PGKGLEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTL YLQMNLSLKTEDTAVYYCTTPWEWSWYDYWGQGLVTVSSA STKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICN VNHKPSNTKVDKKVEPKSCDGGGSGGGGSEVQLLESGL GLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGKLEWVS RIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAE DTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVSSASTKGP	73

-continued

EXAMPLARY SEQUENCES		
	SVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKP SNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKP KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALG APIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKG FYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD KSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	
anti ID CD3 scFv 4.32.63 MK062 protease site CD3 VL CLambda, pETR16544	QVQLKESGPGLVAPSQSLSTICTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQTDATATYYCAKGIITVVDYYAMDYWGQGTSTVTVSSGG GSGGGGGSGGGGGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLSQSEDVARYYCQHYYSTPYTFGCCTKL EIKGGGGSGGGGSRQARVVNGGGGGSGGGSGGGGSQA VVTQEPSTLTVSPGGTIVTLTCSSTGAVTTSNYANWVQEKPG QAFRGLIGGTNKRAPGTPARFSGSLGKKAALTLGAQPED EAEYYCALWYSLWVFGGGTKLTVLGQPKAAPSVTLFPPSS EELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTP SKQSNNKYAASSYLSLTPEQWKSHRYSYSCQVTHEGSTVEKT VAPTECS	74
anti ID CD3 scFv 4.32.63 non-cleavable linker CD3 VL CLambda, pETR16545	QVQLKESGPGLVAPSQSLSTICTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQTDATATYYCAKGIITVVDYYAMDYWGQGTSTVTVSSGG GSGGGGGSGGGGGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLSQSEDVARYYCQHYYSTPYTFGCCTKL EIKGGGGSGGGGGSGGGGGGGSGGGGGSGGGGSQA AVVTQEPSTLTVSPGGTIVTLTCSSTGAVTTSNYANWVQEK QAFRGLIGGTNKRAPGTPARFSGSLGKKAALTLGAQPE DEAEYYCALWYSLWVFGGGTKLTVLGQPKAAPSVTLFPPS SEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTP PSKQSNNKYAASSYLSLTPEQWKSHRYSYSCQVTHEGSTVE KTVAPTECS	75
aMSLN RG7787 VH CH1 EE CD3 CH2527- VH3 23-12 VL CH1 Fc knob PG LALA, pETR15445	QVQLVQSGAEVKKPGASVKVSCASGYSTGYTMNWRQA PGQGLEWMGLITPYNGASSYNQKFRGKATMTVDTSTSTVY MELSSLRSEDAVYYCARGGYDGRGPDYWGQGTIVTVSSA STKGPSVFPPLAPSSKSTSGGTAALGCLVEDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICN VNHKPSNTKVDKVEPKSCDGGGGSGGGGSAVVTQEPSTL TVSPGGTIVTLTCSSTGAVTTSNYANWVQEKQAFRGLIG GTNKRAPGTPARFSGSLGKKAALTLGAQPEDEAEYYCAL WYSLWVFGGGTKLTVLSSASTKGPSVFPPLAPSSKSTSGGT AALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKVEPKSCDK THTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV VLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREP QVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCV MHEALHNHYTQKSLSLSPGK	76
aMSLN RG7787 VH CH1 EE Fc hole P329G LALA, pETR15444	QVQLVQSGAEVKKPGASVKVSCASGYSTGYTMNWRQA PGQGLEWMGLITPYNGASSYNQKFRGKATMTVDTSTSTVY MELSSLRSEDAVYYCARGGYDGRGPDYWGQGTIVTVSSA STKGPSVFPPLAPSSKSTSGGTAALGCLVEDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICN VNHKPSNTKVDKVEPKSCDKTHTCPPCPAPEAAGGPSVFL FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE VHNATKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS NKALGAPIEKTISKAKGQPREPQVCTLPSPRDELTKNQVSL CAVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLV SKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	77
aMSLN RG7787 VL Ck RK, pETR15443	DIQMTQSPSSLSASVGDRTITCSASSSVSYMHYQQKSGK APKLLIYDTSKLASGVPSRFSGSGSGTDFLTITSSLPEDFAT YYCQQWSKHPLTFGQGTKEIKRTVAAPSVFIFPPSDRKLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYSLSSLTITLSKADYEKKHVKYACEVTHQGLSSPVTKSF NRGEC	78

-continued

EXAMPLARY SEQUENCES		
anti ID CH2527 4.32.63 CD3 CH2527 VH 23-12 Ck, MMP9-MK062 site, pETR16758	QVQLKESGPGLVAPSQSLITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQTDDETATYYCAKGIITVVDYDYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGGSVHMPGLGFLGPRQARVNVGGGGSGGGGSEVQ LLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWRQAPGK GLEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQ MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTSLTVTS SASVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSSTYLSSTLTLSKADYEKH KVYACEVTHQGLSSPVTKSFNRGEC	79
anti ID CH2527 4.32.63 CD3 CH2527 VH 23-12 Ck, non-cleavable linker, pETR16759	QVQLKESGPGLVAPSQSLITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQTDDETATYYCAKGIITVVDYDYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGGSVHMPGLGFLGPRQARVNVGGGGSGGGGSEV QLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWRQAPGK KLEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQ MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTSLTVTS SASVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSSTYLSSTLTLSKADYEKH KVYACEVTHQGLSSPVTKSFNRGEC	80
CD3 CH2527 VH 23- 12-Ck, pETR13811	EVQLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWRQA PGKLEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLY LQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTSLTV VSSASVAAPSVFIFPPSDEQLKSGTASVVCCLNNFYPREAKV QWKVDNALQSGNSQESVTEQDSKDSSTYLSSTLTLSKADYE KHKVYACEVTHQGLSSPVTKSFNRGEC	81
anti CD3 (CH2527 VH 3-23(12) VL7- 46(13)) scFv 4.32.63 MMP9 Matritptase MK062 aMSLN VH CH1 EE CH2527- VL7_46-13 CH1 hum Fc knob PG LALA, pETR16751	QVQLKESGPGLVAPSQSLITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQTDDETATYYCAKGIITVVDYDYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGGSVHMPGLGFLGPRQARVNVGGGGSGGGGSQAV VTQEPSTLTVSPGGTTLTCSGSTGAVTTSNYANWVQEKPG QAFRGLIGGTNKRAPGTTPARFSGSLGGAALTLGAQPED EAEYYCALWYSNLWVFGGKTLTVLSSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSKVHTFPA VLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDK VEPKSCDGGGGSGGGGSGVQLVQSGAEVKKPGASVKVSC KASGYSFTGYTMNWRQAPGQGLEWMGLITPYNGASSYNQ KFRGKATMTVDTSSTVYMESSLRSEDVAVYYCARGGYDG RGFDYWGQGTSLTVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVEDYFPEPVTVSWNSGALTSKVHTFPAVLQSSGLYSL SVTVPSSSLGTQTYICNVNHKPSNTKVDKVEPKSCDKTHT CPPCPAPEAAGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKALGAPIEKTIISKAKGQPREPQV YTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPEN NYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMH EALHNHYTQKSLSLSPGK	82
anti CD3 (CH2527 VH_3-23(12) VL7- 46(13)) scFv 4.32.63 non-cleavable linker aMSLN VH CH1 EE CH2527-VL7_46-13 CH1 hum Fc knob PG LALA, pETR16752	QVQLKESGPGLVAPSQSLITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKSVFLKLNS LQTDDETATYYCAKGIITVVDYDYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINSLQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGGSVHMPGLGFLGPRQARVNVGGGGSGGGGSQ AVVTQEPSTLTVSPGGTTLTCSGSTGAVTTSNYANWVQEK QAFRGLIGGTNKRAPGTTPARFSGSLGGAALTLGAQPE DEAEYYCALWYSNLWVFGGKTLTVLSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSKVHTFP AVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDK KVEPKSCDGGGGSGGGGSGVQLVQSGAEVKKPGASVKVS CKASGYSFTGYTMNWRQAPGQGLEWMGLITPYNGASSYN	83

-continued

EXAMPLARY SEQUENCES		
	QKFRGKATMTVDSTSTVYMESSLRSEDVAVYVCARGGYD GRGFDYWGQGLVTVSSASTKGPSVFPLAPSSKSTSGGTA ALGCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKT HTCPPCPAPEAAGGPSVFLFPPKPKDTLMI SRTPEVTCVVD VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALGAPIEKTI SKAKGQPREPQ VYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVM HEALHNHYTQKSLSLSPGK	
CH2527 XFab aMSLN RG7787 HC EE Fc knob PG LALA, pETR16764	QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK PGQAFRGLIGGTNKRAPGTPARFSGSLGGKAALTLSGAQF EDEAEYCALWYNLWVFGGKTLTVLSSASTKGPSVFPLA PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTF PAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDGGGSGGGGSGVQLVQSGAEVKKPGASVKV SCKASGYSFTGYTMNWVRQAPGGLEWMGLITPYNGASSY NQKFRGKATMTVDSTSTVYMESSLRSEDVAVYVCARGGY DGRGFDYWGQGLVTVSSASTKGPSVFPLAPSSKSTSGGT AALGCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDK THTCPPCPAPEAAGGPSVFLFPPKPKDTLMI SRTPEVTCVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCKVSNKALGAPI EKTISKAKGQPREP QVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSV MHEALHNHYTQKSLSLSPGK	84
anti CD3 (CH2527 VH 3-23(12) VL7- 46(13)) scFv 4.32.63 Cathepsin S/B site CH2527 VH3_23-VH12 CH1 FcL1 16D5 VH CH1 hum Fc knob PG LALA, pETR16550	QVQLKESGPGLVAPSQSLSTCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSI SKDNKSQVFLKLN LQTDATATYYCAKGIITVVDYVAMDYWGQGTSTVSSGG GGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADVP SRFS GSGSGTQYSLKINLSQSEDEVARYYQCQHYSTPYTFGCSTKL EIKGGGSGGGSGGGSGFVGGTGGGSGGGSGGGSEV QLLESGLLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPG KGLEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQ MNSLR AEDTAVYYCVRHGNFGNSYVSWFAYWGQGLVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPKSCDGGGSGGGSGGGSEVQLVES GGGLVKPGGSLRLSCAASGFTFSTYAMNWVRQAPGKLE WVGRIKSKTDGTTDYAAPVKGRFTISRDDSKNTLYLQMN LKTEDTAVYYCTTPWEWSWYDYWGQGLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPS NTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPK DTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD KSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK	85
Combined MMP9 MK062, 33 AA for CD3	GGGGSVHMLGFLGPRQARVNVNGGGGSGGGGSGGS	86
Combined MMP9 MK062, 35 AA for Her1	GGGGSVHMLGFLGPRQARVNVNGGGGSGGGGSGG	87
Cathepsin S/B	GGGSGGGSGGGSGGGSGFVGGTGGGSGGGSGGGSGGS	88
KKAAPVNG	GGGSGGGSGGSKKAAPVNGGGGSGGGSGGGSGGS	89
PMACKVNG	GGGSGGGSGSPMAKKVNGGGGSGGGSGGGSGGS	90
QARAKVNG	GGGSGGGSGSQARAKVNGGGGSGGGSGGGSGGS	91
MMP9	GGGSGGGSGSVHMLGFLGPGGGGSGGGSGGGSGGS	92
QARAK	GGGSGGGSGSQARAKGGGSGGGSGGGSGGGSGGS	93
MMP9-PMACK	GGGGSVHMLGFLGPPMAKKGGGSGGGSGGGSGGS	94

-continued

EXAMPLARY SEQUENCES		
KKAAP	GGGSGGGGSKKAAPGGGSGGGSGGGSGGS	95
PMAKK	GGGSGGGGSPMAKKGGGSGGGSGGGSGGS	96
Protease recognition site 6	VHMP LGFLGPRQARVNG	97
Protease recognition site 7	FVGGTG	98
Protease recognition site 8	KKAAPVNG	99
Protease recognition site 9	PMAKKVNG	100
Protease recognition site 10	QARAKVNG	101
Protease recognition site 11	VHMP LGFLGP	102
Protease recognition site 12	QARAK	103
Protease recognition site 13	VHMP LGFLGPPMAKK	104
Protease recognition site 14	KKAAP	105
Protease recognition site 15	PMAKK	106
aMSLN CDR H1 Kabat	GYTMN	107
aMSLN CDR H2 Kabat	LITPYNGASSYNQKFRG	108
aMSLN CDR H3 Kabat	GGYDGRGFDY	109
aMSLN CDR L1 Kabat	SASSSVSYMH	110
aMSLN CDR L2 Kabat	DTSKLAS	111
aMSLN CDR L3 Kabat	QQWSKHPLT	112
aMSLN VH	QVQLVQSGAEVKKPGASVKVSKASGYSTGYTMNWVRQA PGQGLEWMGLITPYNGASSYNQKFRGKATMTVDTSSTVY MELSSLRSEDAVYYCARGGYDGRGFDYWGQGT LVTVSS	113
aMSLN VL	DIQMTQSPSSLSASVGDRTITCSASSSVSYMHVYQQKSGK APKLLIYDTSKLASGVPSRFSGSGSGTDFTLTISSLQPEDFAT YYCQQWSKHPLTFGQGTKLEIK	114
aHER1 VH	QVQLVQSGAEVKKPGSSVKVSKASGFTFTDYKIHVVRQAP GQGLEWMGYFNPNSGYSTYAQKFQGRVTITADKSTSTAYM ELSSLRSEDAVYYCARLSPGGYYVMDAWGQGT TTVTVSS	115
aHER1 VL	DIQMTQSPSSLSASVGDRTITCRASQGINNYLNWYQQKPGKA PKRLIYNTNNLQTGVPSRFSGSGSGTEFTLTISLQPEDFATYC LQHNSFPTFGQGTKLEIK	116
Construct	DNA Sequence	SEQ ID No
LC Common light chain pETR13197	CAGGCCGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGC GGCACCGTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACC ACCAGCAACTACGCCAAGTGGGTGCAGGAAAAGCCCGCCAGGCC TTCAGAGGACTGATCGGCGGCACCAACAAGAGAGCCCTGGCACC CCTGCCAGATTAGCGGATCTCTGCTGGGAGGAAAGCCGCCCTG ACACTGTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGC GCCCTGTGTTACAGCAACCTGTGGGTGTCGGCGGAGGCACCAAG CTGACAGTCTAGGTCAACCAAGGCTGCCCCAGCGTGACCCCTG TTCACCCAGCAGCGAGGAAGTGCAGGCCAACAAGGCCACCTG	117

-continued

EXAMPLARY SEQUENCES		
	GTCTGCCTGATCAGCGACTTCTACCCAGGCGCCGTGACCGTGGCC TGGAAGGCCGACAGCAGCCCCGTGAAGCCGCGCTGGAGACCAC CACCCCCAGCAAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTA CCTGAGCCTGACCCCGAGCAGTGGAAGAGCCACAGGTCTACAG CTGCCAGGTGACCCACGAGGCGAGCACCCTGGAGAAAACCGTGG CCCCCACCAGTGCAGCTGA	
anti_CD3 (CH2527 VH_3-23(12) VL7-46(13)) scFv 4.32.63 MMP9 Matriptase MK062 CH2527 VH3_23-VH12 CH1_F01R1 16D5 VH CH1 hum Fc knob PG LALA, pETR16546 (FIG. 45N)	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTCTGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGCGAGCGGAGCACCATTACAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGCATCACCACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGAGGAAGTGGCGGAGG GGGATCTGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCGAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCTACACCTT CGGCTGCGGCACCAAGCTGGAATCAAAGGAGGCGGCGGAAGTG TGACATGCCCTGGGCTTCTGGGCGCCAGACAGGCCAGAGTCG TGAACGGGGGGGCGGAGGCGAGTGGGGGGGAGGATCCGAGGT GCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGGAT CTCTGAGACTGAGCTGTGCCGCGAGCGGCTTCACTTCAGCACCT ACGCCATGAACCTGGGTGCGCCAGGCCCTGGCAAAGGCCCTGGAAT GGGTGTCCTCGGATCAGAAGCAAGTACAACAACCTACGCCACCTACTA CGCCGACAGCGTGAAGGCCCGGTTCACTATCAGCCGGGACGACA GCAAGAACAACCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGG ACACCGCGTGTACTATTGTGTGCGGCAGCGCAACTTCGGCAACA GCTATGTGTCTTGGTTTGCTACTGGGGCCAGGGCACCTCTGTGA CCGTGTCAAGCGCTAGCACAAAGGGCCCTAGCGTGTTCCTCTGG CCCCAGCAGCAAGAGCACAAAGCGGCGGAACAGCCGCCCTGGGC TGCCTCGTGAAGGACTACTTCCCGAGCCCGTGACAGTGTCTTGG AACAGCGGAGCCCTGACAAGCGCGCTGCACACCTTCCCTGCCGTG CTGCAGAGCAGCGGCCTGTACTCCCTGAGCAGCGTGGTCACCGTG CCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAAC CACAAGCCCAGCAACACCAAAGTGGACAAGAAGTGGAGCCCAAG AGCTGTGATGGCGGAGGAGGTCGCGGGCGGAGGATCCGAGGT GCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGGTTT CCTGCGTCTGAGCTGCGCGGCTTCCGGGTTCACTTCTCCAACGC GTGGATGAGCTGGGTTTCGCGAGGCCCGGGCAAGGCCCTCGAGT GGGTTGGTTCGTATCAAGTCTAAACTGACCGTGGCACCACGGATTA CGCGGCTCCAGTTAAGGTCGTTTACCAATTCCCGCAGCATAGC AAAACACTCTGTATCTGCAGATGAACCTCTGAAAACTGAAGACAC CGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACGAT TATTGGGGCCAGGGCACGCTGGTTACGGTGTCTAGCGCTAGTACC AAGGGCCCCAGCGTGTTCCTCCCTGGCACCCAGCAGCAAGAGCACA TCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAAGACTACTTC CCCGAGCCCGTGACCGTGTCTTGGAACTCTGGCGCCCTGACCAGC GGCGTGCACACCTTCCAGCCGCTGCTGCAGAGCAGCGGCTGTAC TCCCTGTCTCCGTGGTCAACCGTGCCCTCTAGCTCCTTGGGAACA CAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACACCAAAGT CGATAAGAAAGTCGAGCCCAAGAGCTGCGACAAAACCTCACACATG CCACCGTGCCAGCACCTGAAGCTGCAGGGGAGCCGTAGTCTT CCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGACC CCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAGGAAGACCT GAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAAT GCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT GTGGTCAGCGTCTCACCCTCTGCACCAAGGACTGGTGAATGGC AAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCGGCGCCCCC ATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCA CAGGTGTACACCTTCCCGCATGCCGGATGAGCTGACCAAGAAC CAGGTAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCGAC ATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTAC AAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTCTCTCTCT ACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAAC GTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA CGCAGAAGGCTCTCCCTGTCTCCGGGTAAATGA	118

-continued

EXAMPLARY SEQUENCES

FolR1 16D5 HC cH2527- VH3 23-12 HC Fc knob PG LALA, pCON999	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGC GGTTCCCTGCGTCTGAGCTGCGCGGGCTTCCGGATTACCTTCTCC AACGCGTGGATGAGCTGGGTTCGCCAGGCCCCGGGCAAAGGCCT CGAGTGGGTTGGTCGTATCAAGTCTAAACTGACGGTGGCACCAC GGATTACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCTCGGAC GATAGCAAAACACTCTGTATCTGCAGATGAACCTCTCTGAAAACGA AGACCCCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGG TACGATTATTGGGGCCAGGGCAGCTGGTTACGGTGTCTTCCGCT AGCACAAAGGGCCCTAGCGTGTTCCTCTGGCCCCCAGCAGCAAG AGCACAAAGCGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGA CTACTTCCCCGAGCCCGTGACAGTGTCTTGAACAGCGGAGCCCT GACAAGCGCGGTGCACACTTTCCTTCCCTGTGCTGCAGAGCAGCGG CCTGTACTCCCTGAGCAGCGTGGTCACCGTGCTAGCAGCAGCCT GGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCAGCAA CACCAAGTGGACAAGAAGTGGAGCCCAAGAGCTGTGATGGCGG AGGAGGGTCCGAGGCGGAGGATCCGAGGTGCAGCTGCTGGAAT CTGGCGGCGGACTGGTGCAGCCTGGCGGATCTCTGAGACTGAGCT GTGCCGCCAGCGGCTTCACCTTCAGCACCTACGCCATGAACCTGGG TGCGCCAGGCCCCCTGGCAAAGGCCTGGAATGGGTGTCCCGGATCA GAAGCAAGTACAACAACACGCCACCTACTACGCCGACAGCGTGA AGGGCCGGTTCACCATCAGCCGGGACGACAGCAAGAACACCCCTGT ACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCCCGTGTACT ATTGTGTGCGGCACGGCAACTTCGGCAACAGCTATGTGTCTTGGTT TGCCTACTGGGGCCAGGGCACCCCTCGTGACCGGTGTCAAGCGCTAG TACCAAGGGCCCCAGCGTGTTCCTCCCTGGCACCCAGCAGCAAGAG CACATCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAGAGCTA CTTCCCCGAGCCCGTGACCGTGTCTTGAACCTTGCGGCCCTTGAC CAGCGCGGTGCACACTTTCAGCCGTGTCTGACAGCAGCGGCCT GTACTCCCTGTCTCCGTGGTCACCGTGCCCTCTAGCTCCCTGGG AACACAGACATATATCTGTAATGTCAATCACAAGCCTTCCAAACACCA AAGTCGATAAGAAAGTCGAGCCCAAGAGCTGCGACAAAACCTCACAC ATGCCCCACCGTGCCCGACCTGAAGCTGCAGGGGGACCGTCAGT CTTCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCCCGG ACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGAC CCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGTGCAT AATGCCAAGACAAGGCGCGGGAGGAGCAGTACAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAGGACTGGCTGAAT GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCGGCGCC CCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAA CCACAGGTGTACACCTGCCCCCATGCCGGGATGAGCTGACCAAG AACCAGGTGACCTGTGGTGCCTGGTCAAAGGCTTCTATCCCAGC GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA CTACAAGACCACGCCCTCCGTGCTGGACTCCGACGGCTCCTTCTT CCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGG GAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCA TACACGAGAAGAGCCTCTCCCTGTCTCCGGGTAATGA	119
anti ID CD3 scFv 4.32.63 MK062 protease site CD3 VL CLambda, pETR16544	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAGGGCATCACACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGGAGGCGGAAGCAGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTACGCGGCAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTGCGGCACCAAGCTGGAAATCAAAGCGGGGGAGGCTCCG GAGGCGCGGAAGTAGACAGGCCAGAGTCGTGAACGGGGGAGGG GGGGGAAGTGGGGCGGAGGAGTGGGGCGGAGGATCCAGG CCGTCTGTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGCA CCGTGACCCCTGACATGTGGCAGTTCTACAGGCGCCGTGACACCA GCAACTACGCCAACTGGGTGAGGAAAAGCCCGGCCAGGCCTTCA GAGGACTGATCGGCGGCACCAACAAGAGAGCCCTGGCACCCCT GCCAGATTACGCGGATCTCTGTGGGAGGAAAGGCGCCCTGACA CTGTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCC CTGTGGTACAGCAACCTGTGGGTGTTCCGGCGGAGGCCAACGCTG	120

-continued

EXAMPLARY SEQUENCES		
	ACAGTCCTAGGTCAACCCAGGCTGCCCCAGCGTGACCTGTTC CCCCCAGCAGCGAGGAAGTGCAGGCCAACAGGCCACCTGGT CTGCCTGATCAGCGACTTCTACCCAGGCGCCGTGACCGTGGCCTG GAAGGCCGACAGCAGCCCCGTGAAGGCCGCGCTGGAGACCACCA CCCCCAGCAAGCAGAGCAACAACAGTACGCCGCCAGCAGCTACC TGAGCCTGACCCCGAGCAGTGAAGAGCCACAGGTCTACAGCT GCCAGGTGACCCACGAGGGCAGCACCGTGGAGAAAACCGTGGCC CCACCGAGTGCAGCTGA	
anti ID CD3 scFv 4.32.63 non-cleavable linker CD3 VL CLambda, pETR16545	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTCTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC AGCGCCTGATCAGCAGACTGAGCATCTCCAAGGCAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAGGGCATCACACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGCGGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT TCTGGCCGACGATGTGCCAGCAGATTACGCCGCGAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCCGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGCTGCGGCACCAAGCTGGAATCAAAGCGGGGGAGGCTCCG GAGGCGGCGGAAGTGGAGGCGGCGGAAGTGGCGGAGGCGGAGG GGGGGGAAGTGGGGGCGGAGGCGAGTGGGGGGGAGGATCCAG GCCGTCTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGC ACCGTGACCTGACATGTGGCAGTCTACAGGCGCGTGACCAACC AGCAACTACGCCAACTGGGTGCGAGGAAAGCCGGCCAGGCCCTC AGAGGACTGATCGCGGCGACCAACAAGAGAGCCCCCTGGCACCCCT GCCAGATTCAGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTGACA CTGTCTGGCGCCAGCCAGAAAGATGAGGCCGAGTACTACTGCGCC CTGTGTTACAGCAACCTGTGGGTGTTTCGGCGGAGGCAACAGCTG ACAGTCCTAGGTCAACCCAGGCTGCCCCAGCGTGACCTGTTC CCCCCAGCAGCGAGGAACTGCAGGCCAACAGGCCACCTGGT CTGCCGTGATCAGCGACTTCTACCCAGGCGCCGTGACCGTGGCCTG GAAGGCCGACAGCAGCCCCGTGAAGGCCGCGCTGGAGACCACCA CCCCCAGCAAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACC TGAGCCTGACCCCCGAGCAGTGAAGAGCCACAGGTCTACAGCT GCCAGGTGACCCACGAGGGCAGCACCGTGGAGAAAACCGTGGCC CCACCGAGTGCAGCTGA	121
aMSLN RG7787 VH CH1 EE CD3 cH2527- VH3 23-12 VL CH1 Fc knob PG LALA, pETR15445	CAGGTGCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACAGGC GCCAGCGTGAAGGTGTCTTCAAGGCCAGCGGCTACAGTTACCC GGCTACACCATGAACGTGGGTGCGCCAGGCTCTGGACAGGGCCTG GAATGGATGGGCTGATCACCCCTACAACGGCGCCAGCAGCTAC AACCAGAAGTTCGGGGCAGGGCCACCATGACCGTGGACACCAGC ACCTCCACCGTGTATATGGAACGTAGCAGCCTGCGGAGCGAGGAC ACCGCCGTGTACTATTGTGCCAGAGGCGGCTACGACGGCAGAGGC TTCGATTATTGGGGCCAGGGCACCCCTGTCGACCGTGTCTCTGCTA GCACCAAGGGCCCTCCGTGTTTCTCTGGCCCTTCCAGCAAGT CCACCTCTGGCGGAAGTCCGCTCTGGGCTGCCTGGTGAAGATT ACTTCCCGAGCCCGTGACCGTGTCTTGAATTCTGGCGCTCTGA CCTCCGCGGTGCACACCTTCCAGCTGTGCTGCAGTCTCTCCGGCC TGTAATCCCTGTCTCTCCGTGTCAGTGCCTCCAGCTCTCTGGG CACCCAGACCTACATCTGCAACGTGAACCAAGCCCTCCAACACC AAGGTGACGAGAAGGTGGAACCAAGTCTGCGACGCTGGCGG AGGTTCCGGAGGCGGAGGATCCAGGCTGTCTGTGACCCAGGAAC CCTCCCTGACAGTGTCTCTGGCGGCACCGTGACCTGACCTGTG GATCTTCTACCGCGCTGTGACCACTCCAACTACGCCAATTGGGT GCAGGAAAAGCCCGCCAGGCCCTCAGAGGACTGATCGGCGGCA CCAACAAGAGAGCCCCCTGGCACCCCTGCCAGATTCTCCGGTTCTC TGCTGGGCGGCAAGGCTGCCCTGACTCTGTCTGGTGTCTCAGCCTG AGGACGAGGCCGAGTACTACTGCGCCCTGTGGTACTCCAACCTGT GGGTGTTCCGGCGGAGGCAACAGCTGACCGTGTCTGTCAGCGCTT CCACCAAGGAGACCAAGTGTGTTCCCTTGGCCCCAGCTCCAAGT CTACATCCGGTGGCACAGCTGCCCTGGGATGTCTCTGAAGGACT ACTTCTCTGAGCCTGTGACAGTGTCTTGAACAGCGGAGCCCTGA CCAGCGGAGTGCACACATTCTTGCAGTGTCTGAGAGCAGCGGCC TGTATAGCCTGAGCAGCGCTGTGACCGTGCCTTCTCTAGCCTGG GAACACAGACATATATCTGTAATGTGAATCATAAGCCAGTAATACC AAAGTGGATAAGAAAGTGAACCTAAGAGCTGCGATAAGACCCACA	122

-continued

EXAMPLARY SEQUENCES		
	CCTGTCCCCCTGCCCTGCTCCTGAAGCTGCTGGTGGCCCTAGCG TGTTCTGTTCCTCCCAAGCCCAAGGACACCTGATGATCTCCCG GACCCCGAAGTGACCTGCGTGGTGGTGGATGTGTCCACGAGGA CCCTGAAGTGAAGTTCAATTGGTACGTGGACGGCGTGAAGTGCA CAACGCCAAGACCAAGCCTAGAGAGGAACAGTACAACCTCCACCTA CCGGGTGGTGTCCGTGCTGACAGTGTGCACAGGACTGGCTGAA CGGCAAAGAGTACAAGTGCAAGGTGTCCAACAAGGCCCTGGGCGC TCCCATCGAAAAGACCATCTCCAAGGCCAAGGGCCAGCCCCGGGA ACCCCAGGTGTACACCTGCCCCCATGCCGGGATGAGCTGACCAA GAACCAAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA ACTACAAGACCAAGCCTCCCGTGTGGACTCCGACGGCTCCTTCTT CCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGG GAACGTCTTCTCATGTCCGTGATGCATGAGGCTCTGCACAACCA TACACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	
aMSLN RG7787 VH CH1 EE Fc hole P329G LALA, pETR15444	CAGGTGCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACAGGC GCCAGCGTGAAGGTGTCTGCAAGGCCAGCGGCTACAGCTTCACC GGCTACACCATGAAGTGGGTGCGCCAGGCTCCTGGACAGGGCCTG GAATGGATGGGCCTGATCACCCCTACAACGGCGCCAGCAGCTAC AACCAGAAGTTCGGGGGCAAGGCCACCATGACCGTGGACACCAGC ACCTCCACCGTGTATATGGAAGTGAAGCAGCCTGCGGAGCGAGGAC ACCGCCGTGTACTATTGTGCCAGAGCGGCTACGACGGCAGAGGC TTCGATTATTGGGGCCAGGGCACCCCTCGTACCGTGTCTCTGCTA GCACCAAGGGCCCTCCGTGTTCCCTCGGCCAGCAGCAAGA GCACAGCGCGGCACAGCCGCTCTGGGCTGCTGCTCGAGGAC TACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTG ACCTCCGGCGTGACACCTTCCCGCCGCTGCTGCAGAGTCTTGGC CTGTATAGCCTGAGCAGCGTGGTCAACCGTGCCTTCTAGCAGCCTG GGCACCCAGACCTACATCTGCAACGTGAACCAAGCCAGCAAC ACCAAGGTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAAC CACACATGCCCAACCGTGCCAGCAGCTGAAGCTGACGGGGGACC GTCAGTCTTCTCTTCCCCCAAACCAAGGACACCCCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGAGCTGAGCCAC AAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAG GTGCATAATGCCAAGACAAGCCGCGGGAGGAGCAGTACAACAGC ACGTACCGTGTGGTCAAGCTCCTCACCGTCTGCACCAAGGACTGG CTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAGCCCTC GCGGCCCCCATCGAGAAAACCATCTCCAAGCCAAGGGCAGCCC CGAGAACCACAGGTGTGACCCCTGCCCCCATCCCGGGATGAGCTG ACCAAGAACCAGGTGAGCCTCTCGTGCGAGTCAAAGGCTTCTATC CCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAATACAGACCAAGCCCTCCCGTGTGGACTCCGACGGCTCC TCTTCTCTCGTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAG CAGGGGAACGTCTTCTCATGTCTCCGTGATGCATGAGGCTCTGCAC AACCCTACACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	123
aMSLN RG7787 VL Ck RK, pETR15443	GACATCCAGATGACCCAGAGCCCCAGCAGCCTGTCTGCCAGCGTG GGCAGCAGAGTGACCATCACCTGTAGCGCCAGCAGCAGCGTGTCC TACATGCACTGGTATCAGCAGAAGTCCGGCAAGGCCCCCAAGCTG CTGATCTACGACACCAGCAAGCTGGCCTCCGGCGTCCCCAGCAGA TTTTCTGGCAGCGGCTCCGGCACCGACTTCAACCTGACAAATCAGCT CCTTCCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTGGT CCAAGCACCCCTGACCTTTGGCCAGGGCACCAAGCTGGAAATCA AGCGTACCGTGGGTGCACCATCTGTCTTCTATCTTCCCGCATCTGA TCGGAAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCTGCTGAAT AACTTCTATCCAGAGAGGCCAAAGTACAGTGAAGGTGGATAACG CCTTCCAATCGGGTAACTCCAGGAGAGTGTACAGAGCAGGACA GCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGCTGAGCA AAGCAGACTACGAGAAACACAAAGTCTACGCTGCGAAGTACCCCA TCAGGGCCTGAGGTGCGCCGTCACAAGAGCTTCAACAGGGGAGA GTGTTAG	124
anti ID CH2527 4.32.63 CD3 CH2527 VH 23-12 Ck, MMP9-MK062 site, pETR16758	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGGCATCACACCGTGGTGGACGAC TACTACGCTATGAGTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGAGGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT	125

-continued

EXAMPLARY SEQUENCES

	GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTGCGGCACCAAGCTGGAAATCAAAGGAGGCGGCGGAAGTG TGCACATGCCCCCTGGGCTTCTGGGCCCCAGACAGGCCAGAGTCG TGAACGGGGGGGCGGAGGCAGTGGGGGGGAGGATCCGAGGT GCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGGAT CTCTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCACCT ACGCCATGAACTGGGTGCGCCAGGCCCTTGGCAAAGGCTGGAAT GGGTGTCCCGGATCAGAAGCAAGTACAACAACACGCCACCTACTA CGCCGACAGCGTGAAGGGCCGGTTCAACATCAGCCGGGACGACA GCAAGAACAACCTGTACTTGCAGATGAACAGCCTGCGGGCCGAGG ACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACA GCTATGTGTCTTGGTTTGCTTACTGGGGCCAGGGCACCCCTCGTGA CCGTGTCAAGCGCTAGCGTGGCGCTCCCTCCGTGTTTATCTTCC CACCTTCCGACGAGCAGCTGAAGTCCGGCACCGCTTCTGTCTGT GCCTGTGAACAACCTTCTACCCCGCGAGGCCAAGGTGCAGTGGA AGGTGGACAACGCCCTGCAGTCCGGCAACAGCCAGGAATCCGTGA CCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCCACCC TGACCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCCT CGAAGTGACCCACCGGGCTGTCTAGCCCCGTGACCAAGTCTT TCAACCGGGCGAGTGCTGA	
anti ID	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC	126
CH2527	CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC	
4.32.63 CD3	AGCTACGGCGTGTCTGGGTGCGCCAGCTTCCAGGCAAGTGTCTG	
CH2527 VH	GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC	
23-12 Ck,	AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG	
non-cleavable	AGCCAGGTGTTCTTGAAGCTGAACAGCCTGCAGACCGACGACACC	
linker,	GCCACCTACTACTGCGCCAAGGGCATCACACCGTGGTGGACGAC	
pETR16759	TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG	
	TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG	
	GGGATCTGGGGAGGCGGAAGCATATCCAGATGACCCAGAGCC	
	CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT	
	GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC	
	AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT	
	TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG	
	GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG	
	TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT	
	CGGCTGCGGCACCAAGCTGGAAATCAAAGCGGGGGAGGCTCCG	
	GAGGCGGCGGAAGTGGAGGCGGCGGAAGTGGCGGAGGCGGAGG	
	GGGGGGAAGTGGGGGCGGAGGCAGTGGGGGGGAGGATCCGAG	
	GTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGG	
	ATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAC	
	CTACGCCATGAACTGGGTGCGCCAGGCCCTGGCAAAGGCCTGGA	
	ATGGGTGTCCCGGATCAGAAGCAAGTACAACAACACGCCACCTAC	
	TACGCCGACAGCGTGAAGGGCCGGTTCAACATCAGCCGGGACGA	
	CAGCAAGAACAACCTGTACTTGCAGATGAACAGCCTGCGGGCCGA	
	GGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAA	
	CAGCTATGTGTCTTGGTTTGCTTACTGGGGCCAGGGCACCCCTCGT	
	GACCGTGTCAAGCGCTAGCGTGGCCGCTCCCTCCGTGTTTATCTT	
	CCCACCTTCCGACGAGCAGCTGAAGTCCGGCACCGCTTCTGTCTGT	
	GTGCTGTCTGAACAACCTTACCCCCGCGAGGCCAAGGTGCAGTG	
	GAAGGTGGACAACGCCCTGCAGTCCGGCAACAGCCAGGAATCCGT	
	GACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCCAC	
	CCTGACCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGC	
	CTGCGAAGTGACCCACCGGGCTGTCTAGCCCCGTGACCAAGTCT	
	TTTCAACCGGGCGAGTGCTGA	
CD3 CH2527	GAAGTGCAGCTGCTGGAATCCGGCGGAGGACTGGTGCAGCCTGG	127
VH 23-12-	CGGATCTCTGAGACTGTCTTGTGCCGCCCTCCGGCTTACCTTCTCC	
Ck,	ACCTACGCCATGAACTGGGTGCGACAGGCTCCTGGCAAGGGCCGTG	
pETR13811	GAATGGGTGTCCCGGATCAGATCCAAGTACAACAACACGCCACCT	
	ACTACGCCGACTCCGTGAAGGGCCGGTTCAACATCTCTCGGGACG	
	ACTCCAAGAACAACCTGTACTTGCAGATGAATCCCTGCGGGCCG	
	AGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCA	
	ACTCTATGTGTCTTGGTTTGCTTACTGGGGCCAGGGCACCCCTCGT	
	GACCGTGTCTATCTGTAGCGTGGCCGCTCCCTCCGTGTTTATCTTC	
	CCACCTTCCGACGAGCAGCTGAAGTCCGGCACCGCTTCTGTCTGTG	
	TGCCTGCTGAACAACCTTACCCCCGCGAGGCCAAGGTGCAGTGG	
	AAGGTGGACAACGCCCTGCAGTCCGGCAACAGCCAGGAATCCGTG	
	ACCGAGCAGGACTCCAAGGACAGCACCTACTCCCTGTCTCCAC	

-continued

EXAMPLARY SEQUENCES		
	CTGACCCCTGTCCAAGGCCGACTACGAGAAGCACAAGGTGTACGCC TGCGAAGTGACCCACAGGGCTGTCTAGCCCCGTGACCAAGTCT TTCAACCGGGCGAGTGCTGA	
anti CD3 (CH2527 VH_3-23(12) VL7-46(13)) scFv 4.32.63 MMP9 Matriptase MK062 aMSLN VH CH1 EE CH2527- VL7_46-13 CH1 hum Fc knob PG LALA, pETR16751	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTCTGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGCATCACCACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAATCATGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCGCAGCACTACTACAGCACCCCTACACCTT CGGCTGCGGCACCAAGCTGGAATCAAAGGAGGCGGCGGAAGTG TGACATGCCCTGGGCTTCTGGGCCCCAGACAGGCCAGAGTCG TGAACGGGGGGGCGGAGGCAGTGGGGGGGAGGATCCCAGGC CGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGCAC CGTGACCTGACATGTGGCAGTTCTACAGGCGCCGTGACCAACAG CAACTACGCCAACTGGGTGCAGGAAAGCCCCGCCAGGCCTTACAG AGGACTGATCGGCGGCACCAACAAGAGAGCCCCCTGGCACCCCTGC CAGATTACGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTGACACT GTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCT GTGGTACAGCAACCTGTGGTGTTCGGCGGAGGCACCAAGCTGAC AGTGCTGAGCAGCGCTTCCACCAAGGGACCCAGTGTGTTCCTT GGCCCCAGCTCCAAGTCTACATCCGTGGCAGAGCTGCCCTGGG ATGTCTCTGTAAGGACTACTTTCTGAGCCTGTGACAGTGTCTTGG AACAGCGGAGCCCTGACAGCGGAGTGACACATTCCTGCGAGTG TGCAGAGCAGCGCCGTGTATAGCCTGAGCAGCGTGTGACCGTG CCTTCTCTAGCCTGGGAACACAGACATATATCTGTAATGTGAATCA TAAGCCCAAGTAATACCAAGTGGATAAAGAAAGTGAACCTAAGAGC TGCGATGGCGGAGGAGGGTCCGGAGGCGGAGGGTCCAGGTGCA GCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACAGGCGCCAGCG TGAAGGTGTCTGCAAGGCCAGCGGTACAGCTTACCGGCTACA CCATGAACTGGGTGCGCCAGGCTCCTGGACAGGGCTGGAATGGA TGGGCTGATCACCCCTACAACGGCGCCAGCAGCTACAACCAGA AGTTCCGGGGCAAGGCCACCATGACCGTGACACAGCACCTCCA CCGTGTATATGGAACAGCAGCCTGCGGAGCGAGGACACCGCCG TGTAATTTGTGCCAGAGGCGGTACGACGGCAGAGGCTTCGATT ATTGGGGCCAGGCGACCTCTGTGACCGTGTCTCTGCTAGCACCA AGGGCCCTCCGTGTTCCTTGGCCCCAGCAGCAAGAGCACCA GCGGCGGCACAGCCGCTCTGGGCTGCCTGGTGCAGGACTACTTC CCCGAGCCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCC GGCGTGACACCTTCCCCGCGTGTGTCAGAGTTCTGGCCTGTAT AGCCTGAGCAGCGTGGTACCGTGTCTTAGCAGCCTGGGCACC CAGACCTACATCTGCAACGTGAACCAAGCCAGCAACACCAAG GTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACACA TGCCCAACCGTGCCAGCCTGAAGCTGCAGGGGGACCGTCAGTC TTCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGA CCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACC CTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATA ATGCCAAGACAAAGCCGCGGAGGAGCAGTACAACAGCACGTACC GTGTGGTCAAGCTCTTACCCTCTGCAACAGGACTGGCTGAATG GCAAGGAGTACAAGTGAAGGTCTCAACAAAGCCCTCGGCGCCC CCATCGAGAAAACATCTCCAAGCCAAAGGGCAGCCCCGAGAAC CACAGGTGTACACCTGCCCCCATGCCGGATGAGCTGACCAAGA ACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCG ACATCGCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAAC ACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTCTTCC TCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGA ACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTA CACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	128
anti CD3 (CH2527 VH_3-23(12) VL7-46(13)) scFv 4.32.63	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTCTGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGCATCACCACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAATCATGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCGCAGCACTACTACAGCACCCCTACACCTT CGGCTGCGGCACCAAGCTGGAATCAAAGGAGGCGGCGGAAGTG TGACATGCCCTGGGCTTCTGGGCCCCAGACAGGCCAGAGTCG TGAACGGGGGGGCGGAGGCAGTGGGGGGGAGGATCCCAGGC CGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGCAC CGTGACCTGACATGTGGCAGTTCTACAGGCGCCGTGACCAACAG CAACTACGCCAACTGGGTGCAGGAAAGCCCCGCCAGGCCTTACAG AGGACTGATCGGCGGCACCAACAAGAGAGCCCCCTGGCACCCCTGC CAGATTACGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTGACACT GTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCT GTGGTACAGCAACCTGTGGTGTTCGGCGGAGGCACCAAGCTGAC AGTGCTGAGCAGCGCTTCCACCAAGGGACCCAGTGTGTTCCTT GGCCCCAGCTCCAAGTCTACATCCGTGGCAGAGCTGCCCTGGG ATGTCTCTGTAAGGACTACTTTCTGAGCCTGTGACAGTGTCTTGG AACAGCGGAGCCCTGACAGCGGAGTGACACATTCCTGCGAGTG TGCAGAGCAGCGCCGTGTATAGCCTGAGCAGCGTGTGACCGTG CCTTCTCTAGCCTGGGAACACAGACATATATCTGTAATGTGAATCA TAAGCCCAAGTAATACCAAGTGGATAAAGAAAGTGAACCTAAGAGC TGCGATGGCGGAGGAGGGTCCGGAGGCGGAGGGTCCAGGTGCA GCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACAGGCGCCAGCG TGAAGGTGTCTGCAAGGCCAGCGGTACAGCTTACCGGCTACA CCATGAACTGGGTGCGCCAGGCTCCTGGACAGGGCTGGAATGGA TGGGCTGATCACCCCTACAACGGCGCCAGCAGCTACAACCAGA AGTTCCGGGGCAAGGCCACCATGACCGTGACACAGCACCTCCA CCGTGTATATGGAACAGCAGCCTGCGGAGCGAGGACACCGCCG TGTAATTTGTGCCAGAGGCGGTACGACGGCAGAGGCTTCGATT ATTGGGGCCAGGCGACCTCTGTGACCGTGTCTCTGCTAGCACCA AGGGCCCTCCGTGTTCCTTGGCCCCAGCAGCAAGAGCACCA GCGGCGGCACAGCCGCTCTGGGCTGCCTGGTGCAGGACTACTTC CCCGAGCCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCC GGCGTGACACCTTCCCCGCGTGTGTCAGAGTTCTGGCCTGTAT AGCCTGAGCAGCGTGGTACCGTGTCTTAGCAGCCTGGGCACC CAGACCTACATCTGCAACGTGAACCAAGCCAGCAACACCAAG GTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACACA TGCCCAACCGTGCCAGCCTGAAGCTGCAGGGGGACCGTCAGTC TTCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGA CCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACC CTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATA ATGCCAAGACAAAGCCGCGGAGGAGCAGTACAACAGCACGTACC GTGTGGTCAAGCTCTTACCCTCTGCAACAGGACTGGCTGAATG GCAAGGAGTACAAGTGAAGGTCTCAACAAAGCCCTCGGCGCCC CCATCGAGAAAACATCTCCAAGCCAAAGGGCAGCCCCGAGAAC CACAGGTGTACACCTGCCCCCATGCCGGATGAGCTGACCAAGA ACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCG ACATCGCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAAC ACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTCTTCC TCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGA ACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTA CACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	129

-continued

EXAMPLARY SEQUENCES	
non-cleavable	AGCCAGGTGTTCTCTGAAGCTGAACAGCCTGCAGACCGACGACACC
linker aMSLN	GCCACCTACTACTGCGCCAAGGCATCACCACCGTGGTGACGAC
VH CH1 EE	TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG
CH2527-	TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG
VL7_46-13	GGGATCTGGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC
CH1_hum Fc	CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT
knob PG	GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC
LALA,	AGAAGCAGGGCAAGAGCCCCAGCTGTGGTGTACGCCGCCACCT
pETR16752	TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG
	GCACACAGTACAGCCTGAAGATCAATCCCTGCAGAGCGAGGACG
	TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT
	CGGCTGCGGCACCAAGCTGGAATCAAAGCGGGGGAGGCTCCG
	GAGGCGGCGGAAGTGGAGGCGGCGGAAGTGGCGGAGGCGGAGG
	GGGGGGAAGTGGGGCGGAGGCAGTGGGGGGGAGGATCCAG
	GCCGTGCTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCGGC
	ACCGTGACCCCTGACATGTGGCAGTTCTACAGGCGCGGTGACCAAC
	AGCAACTACGCCAACTGGGTGAGGAAAGCCCGGCCAGGCCTTC
	AGAGGACTGATCGGCGGCACCAACAAGAGAGCCCCCTGGCACCCCT
	GCCAGATTACGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTGACA
	CTGTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCC
	CTGTGGTACAGCAACCTGTGGGTGTTCTGGCGGAGGCACCAAGCTG
	ACAGTGCTGAGCAGCGCTTCCACCAAGGGACCCAGTGTGTTCCCC
	CTGGCCCCCAGCTCCAAGTCTACATCCGGTGGCACAGCTGCCCTG
	GGATGTCTCGTGAAGGACTACTTTCTGAGCCTGTGACAGTGTCTT
	GGAACAGCGGAGCCCTGACCAGCGGAGTGACACATTCCTTCGAG
	TGCTGCAGAGCAGCGCCTGTATAGCCTGAGCAGCGTCTGTACCG
	TGCCTTCCTCTAGCCTGGGAACACAGACATATATCTGTAATGTGAAT
	CATAAGCCCAGTAATACCAAGTGGATAAGAAAGTGGAACTTAAGA
	GCTGCGATGGCGGAGGAGGGTCCGGAGGCGGAGGGTCCAGGT
	GCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACAGGCGCCA
	GCGTGAAGGTGTCTGCAAGGCCAGCGGCTACAGCTTCACCGGCT
	ACACCATGAAGTGGGTGCGCCAGGCTCCTGGACAGGCGCTGGAAT
	GGATGGGCTGATCACCCCCACACGCGCCAGCAGCTACAACC
	AGAAGTTCGGGGCAAGGCCACCATGACCGTGGACACACAGCACCT
	CCACCGTGTATATGGAAGTGAAGCAGCTGCGGAGCGAGGACACCG
	CCGTGTACTATTGTGCCAGAGGCGGCTACGACGGCAGAGGCTTCG
	ATTATTGGGGCCAGGGCACCTCGTGACCGTGTCTCTGTAGTACA
	CCAAGGGCCCCCTCCGTGTTCCCTTGGCCCCCAGCAGCAAGAGCA
	CCAGCGGCGGCACAGCCGCTCTGGGCTGCCTGGTCTGAGGACTAC
	TTCCCCGAGCCCGTGACCGTGTCTTGGAAACAGCGGAGCCCTGACC
	TCCGCGGTGCACACCTTCCCCGCGGTGCTGCAGAGTTCTGGCCTG
	TATAGCCTGAGCAGCGTGGTCAACCGTGCCTTCTAGCAGCCTGGGC
	ACCCAGACCTACATCTGCAACGTGAACCAAGCCAGCAACACCA
	AGGTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTACA
	CATGCCACCGTGCCAGCACCTGAAGCTGCAGGGGGACCGTCA
	GTCTTCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCC
	GGACCCCTGAGGTCAATGCGTGGTGGTGGACGTGAGCCACGAA
	GACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTG
	CATAATGCCAAGACAAGCCGCGGGAGGAGCAGTACAACAGCAGC
	TACCGTGTGGTACGCGTCTTCAACGCTCTGCACAGGACTGGCTG
	AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCGGC
	GCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGA
	GAACCAAGGTGTACACCTGCCCCCATGCCGGGATGAGCTGACC
	AAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCCA
	GCGACATCGCCGTGGAGTGGGAGAGCAATGGGACGCCGAGAAC
	AACTACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTC
	TTCTCTACAGCAAGCTACCGTGGACAAGAGCAGGTGGCAGCAG
	GGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACC
	ACTACACGCAAGAGCCTCTCCCTGTCTCCGGTAAATGA
CH2527 XFab	CAGGCCGTGCTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGC
aMSLN	GGCACCGTGACCTGACATGTGGCAGTTCTACAGGCGCCGTGACC
RG7787 HC	ACCAGCAACTACGCCAACTGGGTGAGGAAAGCCCGGCCAGGCC
EE Fc knob	TTAGAGGACTGATCGGCGGCACCAACAAGAGAGCCCTGGCACCC
PG LALA,	CCTGCCAGATTCTCCGGTCTCTGCTGGGCGGCAAGGCTGCCCTG
pETR16764	ACTCTGTCTGGTGTCTAGCCTGAGGACGAGGCGGAGTACTACTGC
	GCCCTGTGGTACTCCAACCTGTGGGTGTTCCGGCGAGGCACCAAG
	CTGACCGTGTCTGTCAGCGCTTCCACCAAGGGACCCAGTGTGTTT
	CCCCTGGCCCCAGCTCCAAGTCTACATCCGGTGGCACAGCTGCC
	CTGGGATGTCTCGTGAAGGACTACTTTCTGAGCCTGTGACAGTGT
	CTTGGAACAGCGGAGCCCTGACCAGCGGAGTGACACATTCCTCTG
	CAGTGTGTCAGAGCAGCGCCTGTATAGCCTGAGCAGCGTCTGTGA
	CCGTGCCCTTCTCTAGCCTGGGAACACAGACATATATCTGTAATGT
	GAATCATAAGCCCAGTAATACCAAGTGGATAAGAAAGTGGAACT

130

-continued

EXAMPLARY SEQUENCES

anti CD3 (CH2527 VH ₃ -23 (12) VL ₇ -46 (13)) scFv 4.32.63 Cathepsin S/B site CH2527 VH ₃ 23-VH12 CH1 FcL1 16D5 VH CH1 hum Fc knob PG LALA, pETR16550	AAGAGCTGCGATGGCGGAGGAGGGTCCGGAGGCGGAGGGTCCCA GGTGCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAAACAGGCG CCAGCGTGAAGGTGTCTGCAAGGCCAGCGGTACAGCTTACCCG GCTACACCATGAAGTGGGTGCGCCAGGCTCCTGGACAGGGCTGG AATGGATGGGCTGATCACCCCTACAACGGCGCCAGCAGCTACA ACCAGAAGTTCCGGGGCAAGGCCACCATGACCGTGGACACCAGCA CCTCCACCGTGTATATGGAAGTGAAGCGCTGCGGAGCGAGGACA CCGCCGTGTACTATTGTGCCAGAGGGCGGTACGACGGCAGAGGCT TCGATTATTGGGGCCAGGGCACCCCTCGTGACCGTGTCTCTGTCTA GCACCAAGGGCCCTCCGTGTTCCCGCTGGCCCCAGCAGCAAGA GCACAGCGGCGGCACAGCCGCTCTGGGCTGCCTGGTTCGAGGAC TACTTCCCCGAGCCCGTGACCGTGTCTCTGGAACAGCGAGCCCTG ACCTCCGGCGTGCACACCTTCCCGCGCTGCTGCAGAGTTCTTGGC CTGTATAGCCTGAGCAGCGTGGTCAACGTGCTTCTAGCAGCCGTG GGCACCCAGACCTACATCTGCAACGTGAACCAAGCCAGCAAC ACCAAGGTGGACGAGAAGGTGGAGGCCAAGAGCTGCGCAAAACT CACACATGCCACCGTGCACAGCACTGAAGTGCAGGGGGACC GTCAGTCTTCTCTTCCCCCAAACCAAGGACACCCCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAG GTGCATAATGCCAAGACAAGCCGCGGGAGGAGCAGTACAACAGC ACGTACCGTGTGGTCAAGCGTCTTCCCGTCTGCAACAGGACTGG CTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGCCCTC GGCGCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCC CGAAGAACACAGGTGTACACCCCTGCCCGCATGCCGGGATGAGCTG ACCAAGAACCAGGTGAGCTGTGGTGCCTGGTCAAAGGCTTCTATC CCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAATACAGACACCGCTCCCGTGTGGACTCCGACGGCTCC TTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAG CAGGGGAACGTCTTCTCATGCTCCGTGATGATGAGGCTCTGCAC AACCCTACACGAGAAGAGCCCTCTCCCTGTCTCCGGGTAAATGA CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAAGCTGACC AGCTACGGCGTGTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCAATTACCAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGGCATCACCAAGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGCAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGAGGAAGTGGCGGAGG GGGATCTGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCTCTGTGGGCGAGACAGTACCATCACAT GCCGGGCAGCGAGAACATCGACAGCTACCTGGCTTGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTACGCGGCAGCGGAAGCG GCACACAGTACAGCTGAAGATCAACTTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTGCCGCCACCAAGCTGGAATCAAAGGCGGGGGAGGCTCCG GAGGCGGCGAAGTGGAGGCGCGGAAGTTTCTGGGGGGGAC CGGGGGCGGAGGAGTGGGGGGGAGGATCCGGGGGATCCGAG GTGCAGCTGTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGG ATCTCTGAGACTGAGCTGTGCCGCGAGCGGCTTCACTTTCAGCAC CTACGCCATGAAGTGGGTGCGCCAGGCCCCCTGGCAAAGGCTTGA ATGGGTGTCCCGGATCAGAAGCAAGTACAACAATACGCCACCTAC TACGCCGACAGCGTGAAGGGCCGGTTACCATCAGCCGGGACGA CAGCAAGAACACCCCTGTACCTGCAGATGAACAGCCTGCGGGCCGA GGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAA CAGCTATGTGTCTTGGTTTGCCCTACTGGGGCAGGGCACCCCTGT GACCGTGTCAAGCGCTAGCACAAAGGGCCCTAGCGTGTCTCCCTCT GGCCCCCAGCAGCAAGAGCAAGAGCGGCGGAACAGCCGCCCTGG GCTGCTGTGAAGGACTACTTCCCCGAGCCCGTGACAGTGTCTT GGAACAGCGAGCCCTGACAAGCGCGGTGCACACCTTCCCTGCC GTGCTGCAGAGCAGCGGCTGTACTCCCTGAGCAGCGTGGTCAAC GTGCCCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTG AACCACAAGCCAGCAACACCAAGTGGACAAGAAGTGGAGGCC AAGAGCTGTGATGGCGGAGGAGGCTCCGGGGCGGAGGATCCGA GGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGG TTCCCTGCGTCTGAGCTGCGCGGCTTCCGGGTTCACTTCTCCAAC GCGTGGATGAGCTGGGTTGCCAGGCCCCGGGCAAAGGCTCGA GTGGGTGGTTCGTATCAAGTCTAAACTGACGGTGGCACCACGGA TTACGCGGCTCCAGTTAAAGGTCGTTTACCATTTCGCGCAGCAT AGCAAAAACACTCTGTATCTGCAGATGAAGTCTGAAACTGAAG ACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGTA	131

-continued

EXAMPLARY SEQUENCES		
	CGATTATTGGGGCCAGGGCAGCTGGTTACGGTGTCTAGCGCTAG TACCAAGGGCCCCAGCGTGTCCCCCTGGCACCCAGCAGCAAGAG CACATCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAGACTA CTTCCCCGAGCCCGTGACCGTGTCTTGAACCTCTGGCGCCCTGAC CAGCGCGCTGCACACCTTTCCAGCCGTGCTGCAGAGCAGCGGCCT GTACTCCCTGTCTCTCGTGGTCACCGTGCCCTCTAGCTCCCTGGG AACACAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACACCA AAGTCGATAAGAAAGTCGAGCCCAAGAGCTGCGACAAAACCTCACAC ATGCCCCACCGTGCCCCAGCACCTGAAGCTGCAGGGGGACCGTCAGT CTTCCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCCCGG ACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGAC CCTGAGGTCAAGTTCAACTGGTACGTGGACGCGGTGGAGGTGCAT AATGCCAAGACAAGGCGCGGGAGGAGCAGTACAACAGCACGTAC CGTGTGGTCAGCGTCTTACCCTCCTGCACCAGGACTGGCTGAAT GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCC CCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCGAGAA CCACAGGTGTACACCTGCCCCCATGCGGGATGAGCTGACCAAG AACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGC GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA CTACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTCTT CCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGG GAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCAACAACCA TACACGCAAGAGCCTCTCCCTGTCTCCGGGTAATGA	
Construct	Amino acid Sequence	SEQ ID No
pETR16859 Omnitarg aff.mat variant Fab cv- Fc hole PG LALA	EVQLVESGGGLVQPGGSLRLSCAASGFTFNDYTMQWVRQA PGKLEWVADVNPNSGGSIVNRRFKGRFTLSVDRSKNTLYL QMNSLRAEDTAVYYCARNLGPFFYFDYWGQTLVTVSSAS TKGPSVFPLAPSSKSTSGGTAAALGCLVEDYFPEPVTVSWNS GALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV NHKPSNTKVDEKVEPKSCDKTHTCPPCPAPEAAGGPSVFLF PPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRVVSVLTVHLQDNLNGKEYKCKVS NKGALGAPIEKTISKAKGQPREPQVCTLPPSRDELTKNQVSL SLVKGFPYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFPLV SKLTVDKSRWQQGNVFSQVMHEALHNHYTQKSLSLSPGK	132
pETR16860 Herceptarg common CLkRK	DIQMTQSPSSLSASVGDRTVITCKASQDVSTAVAWYQQKPG KAPKLLIYSASFRYTGVPSRFSRSGRTDFTLTISLQPEDFA TYYCQQHYTTPPTFGQGTKEIKRTVAAPSVFIFPPPSDRKLK SGTASVVCCLNNFYPREAKVQWKVDNALQSGNSQESVTEQ DSKDSYSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC	133
pETR17605 CD3 X Fab Herceptin HC charged variants Fc knob PG LALA	QAVVTQEPSTLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK PGQAFRGLIGGTNKRAPGTPARFSGSLGGAALTLGAQP EDEAEYYCALWYSLNLFVGGGTGKLTVLSSASTKGPSVFPLA PSSKSTSGGTAAALGCLVKDYFPEPVTVSWNSGALTSGVHTF PAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKGGGSGGGGSEVQLVESGGGLVQPGGSLRL SCAASGFNIKDTYIHWVRQAPGKLEWVARIYPTNGYTRYA DSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCSRWGGE GFYAMDYWGQTLVTVSSASTKGPSVFPLAPSSKSTSGGT AALGCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDK THTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV VLTVLHQLDNLNGKEYKCKVSNKALGAPIEKTISKAKGQPREP QVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSLKLTVDKSRWQQGNVFSQV MHEALHNHYTQKSLSLSPGK	134
pETR17606 anti CD3 (CH2527 VH 3-23(12) VL7-46(13)) scFv 4.32.63 non cleavable linker aHerceptin VH CH1 EE CH2527- VL7 46-13 CH1 hum Fc knob PG LALA	QVQLKESGPGLVAPQSLSITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSIKDNKSKVFLKLS LQTDATATYYCAKGIITVVDYAMDYWGQTSVTVSSGG GGSGGGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDPVSRFS GSGSGTQYSLKINSLSQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGQ AVVTQEPSTLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK GQAFRGLIGGTNKRAPGTPARFSGSLGGAALTLGAQPE DEAEYYCALWYSLNLFVGGGTGKLTVLSSASTKGPSVFPLAP	135

-continued

EXAMPLARY SEQUENCES		
	SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDK KVEPKSCDGGGSGGGSEVQLVESGGGLVQPGGSLRLS CAASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGYTRYAD SVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCSRWGGE FYAMDYWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAA LGCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKVEPKSCDKTH TCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQ VYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQCSVM HEALHNHYTQKSLSLSPGK	
pETR17607 anti CD3 (CH2527 VH 3-23(12) VL7-46(13)) scFv 4.32.63 MMP9 Matriptase MK062 aHerceptin VH CH1 EE CH2527-VL7 46-13 CH1 hurn Fc knob PG LALA	QVQLKESGPGLVAPSSQSLITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKQVFLKLNS LQTDATATYYCAKGIITVVDYYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVITIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINLSQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGGSVHMPGLGFLGPRQARVNVGGGGSGGGGSAV VTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPG QAFRGLIGGTNKRAPGTPARFSGSLGKKAALTLGAQPED EAEEYCALWYSLWVFGGGTKLTVLSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDK VEPKSCDGGGSGGGSEVQLVESGGGLVQPGGSLRLS AASGFNIKDTYIHWVRQAPGKGLEWVARIYPTNGYTRYAD VKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCSRWGGE YAMDYWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKVEPKSCDKTH TCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQ VYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQCSVM HEALHNHYTQKSLSLSPGK	136
FolR1 36F2 VH CH1 EE Fc hole PG LALA pETR14797	QVQLVQSGAEVKKPGASVKVSCASGYTFTSYMHWRQA PGQGLEWMGIINPSGGSTSYAQKQGRTMTHTSTSTVY MELSSLRSEDTAVYYCARSFFTGFHLDYWGQGTLLTVSSAS TKGPSVFPLAPSSKSTSGGTAALGCLVEDYFPEPVTVSWNS GALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNV NHKPSNTKVDKVEPKSCDKTHTCPPCPAPEAAGGPSVFLF PPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS NKALGAPIEKTISKAKGQPREPQVTLTPPSRDELTKNQVSL CAVKGFPYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLV SKLTVDKSRWQQGNVFSQCSVMHEALHNHYTQKSLSLSPGK	137
FolR1 36F2 VL Ck RK, pETR14798	EIVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKP GQAPRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDF AVYYCQQTNEHYTFTGQGTKEIKRTVAAPSVFIFPPSDRK LKSQTASVCLLNNFYPRKAVQWQVDNALQSGNSQESVTE QDSKDSYSLSTLTLSKADYEKHKVYACEVTHQGLSPVTK SPNRGEC	138
anti CD3 (CH2527 VH 3-23(12) VL7- 46(13)) scFv 4.32.63 MMP9 Matriptase MK062 aFolR1 36F2 VH CH1 EE CH2527- VL7 46-13 CH1 hum Fc knob PG LALA pETR17621	QVQLKESGPGLVAPSSQSLITCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKQVFLKLNS LQTDATATYYCAKGIITVVDYYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSDIQMTQSPASLSASVGETVITIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GSGSGTQYSLKINLSQSEDVARYYCQHYYSTPYTFGCGTKL EIKGGGGSVHMPGLGFLGPRQARVNVGGGGSGGGGSAV VTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPG QAFRGLIGGTNKRAPGTPARFSGSLGKKAALTLGAQPED EAEEYCALWYSLWVFGGGTKLTVLSASTKGPSVFPLAPS SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA VLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDK VEPKSCDGGGSGGGGSAVQLVQSGAEVKKPGASVKVSC KASGYTFTSYMHWRQAPGQGLEWMGIINPSGGSTSYAQ KQGRVTMTHTSTSTVYMESSLRSEDTAVYYCARSFFTG	139

-continued

EXAMPLARY SEQUENCES		
	FHLDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SVVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKHT CPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL VLHQDWLNGKEYKCKVSNKALGAPIEKTIKAKGQPREPQV YTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE NYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVMH EALHNHYTQKSLSLSPGK	
anti CD3 (CH2527 VH_3-23(12) VL7- 46(13)) scFv 4.32.63 non cleavable linker aFolR1 36F2 VH CH1 EE CH2527-VL7_46-13 CH1 hum Fc knob PG LALA pETR17622	QVQLKESGPGLVAPSQSLSTCTVSGFSLTSYGVSWVRQPP GKCLEWLGIIWGDGSTNYHSALISRLSISKDNSKQVFLKLNS LQDDDTATYYCAKGIITVVDYIYAMDYWGQGTSTVTVSSGG GGSGGGSGGGSGGGSGGGSDIQTQSPASLSASVGETVTIT CRASENIDSYLAWYQQKQKSPQLLVYAATFLADDVPSRFS GGSGGTQYSLKINSLQSEVARYYCOHYSTPYTFGCGTKL EIKGGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGQ AVVTQEPSTLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK GQAFRGLIGGTNKRAPGTPARFSGSLGGKAALTLGAQPE DEAEYYCALWYSNLWVFGGKTLTVLSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVTVPSLGTQTYICNVNHKPSNTKVDEK KVEPKSCDGGGGSGGGSGVQLVQSGAEVKKPGASVKVS CKASGYTFTSYMHWRQAPGGGLEWMGIINPSGGSTSYA QKPFQGRVTMTHTDSTSTVYMELSLRSSEDAVYICARSFFT GFHLDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA LGCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKH TCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVD SHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKALGAPIEKTIKAKGQPREPQ VYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVM HEALHNHYTQKSLSLSPGK	140
FolR1 36F2 classic format: CH2527 XFab 36F2 HC EE Fc knob PG LALA pETR17623	QAVVTQEPSTLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK PGQAFRGLIGGTNKRAPGTPARFSGSLGGKAALTLGAQPE EDEAEYYCALWYSNLWVFGGKTLTVLSSASTKGPSVFPLA PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTF PAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDGGGGSGGGSGVQLVQSGAEVKKPGASVKV SCKASGYTFTSYMHWRQAPGGGLEWMGIINPSGGSTSY AQKPFQGRVTMTHTDSTSTVYMELSLRSSEDAVYICARSFF TGFLHDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA ALGCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKT HTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVD VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALGAPIEKTIKAKGQPREPQ VYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPE NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVM HEALHNHYTQKSLSLSPGK	141
Herceptin/Omnitarg CDR H1 Kabat	DYTM	142
Herceptin/Omnitarg CDR H2 Kabat	DVNPNSGGSIIVNRRFKG	143
Herceptin/Omnitarg CDR H3 Kabat	NLGPFYFDY	144
Perjeta CDR H1 Kabat	TSNYANW	145
Perjeta CDR H2 Kabat	GTNKRAPGTPARFSGSLGG	146
Perjeta CDR H3 Kabat	TKLTV	147
CLC CDR L1 Kabat	KASQDVSTAVA	148
CLC CDR L2 Kabat	SASFRYT	149
CLC CDR L3 Kabat	QQHYTTPPT	150

-continued

EXAMPLARY SEQUENCES		
36F2 CDR H1 Kabat	SYMH	151
36F2 CDR H2 Kabat	IINPSGGSTSYAQKFQG	152
36F2 CDR H3 Kabat	SFFTGFHLDY	153
36F2 CDR L1 Kabat	RASQSVSSSYLA	154
36F2 CDR L2 Kabat	GASSRAT	155
36F2 CDR L3 Kabat	QQTNEHYT	156
Anti-FolR1 36F2 variable region VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYMHVVRQA PGQGLEWMGIINPSGGSTSYAQKFQGRVTMTHTSTSTVY MELSSLRSEDTAVYYCARSFPTGFHLDYWGQGLTVTVSS	157
Anti-FolR1 36F2 variable region VL	EIVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKP GQAPRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDF AVYYCQQTNEHYTTFGQGTKEIK	158
Herceptarg variable region VH1	EVQLVESGGGLVQPGGSLRLSCAASGFTFNDYTMQWVRQA PGKLEWVADVNPNSGGSIIVNRFRKGRFTLSVDRSKNTLYL QMNSLRSEDTAVYYCARNLGPFFYFDYWGQGLTVTVSS	159
Herceptarg variable region VH2	EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAP GKGLEWVARIYPTNGYTRYADSVKGRFTISADTSKNTAYLQ MNSLRSEDTAVYYCSRWGGEGFYAMDYWGQGLTVTVSS	160
Herceptarg common variable region VL	DIQMTQSPSSLSASVGRVTITCKASQDVSTAVAWYQQKPG KAPKLLIYSASFRYTGVPSRFSGSGSGTDFTLTISLQPEDFA TYYCQQHYTTPPTFGQGTKEIK	161
Construct	DNA Sequence	SEQ ID No
pETR16859 Omnitarg aff.mat variant Fab cv-Fc hole PG LALA	GAAGTTCAGCTGGTTGAAAGCGGTGGTGGTCTGGTTCAGCCTGGT GGTAGCCTGCGTCTGAGCTGTGCAGCAAGCGGTTTACCTTTAACG ATTATACCATGGATTGGGTTTCGTACAGCACCGGGTAAAGGTCTGGA ATGGGTTGCAGATGTTAATCCGAATAGCGGTGGTAGCATTGTTAAC CGTCGTTTTAAAGGTCGTTTTACCCGTGAGCGTTGATCGTAGCAAAA ATACCCGTGATCTGCAAATGAATAGTCTGCGTGCAGAGGATACCGC AGTGTATTATTGTGCACGTAACCTGGGTCCGTTCTTCTACTTTGATT ATTGGGGTCAGGGCACCTGGTTACCGTTAGCAGCGCTAGCACCA AGGGCCCTCCGTGTTCCCTTGGCCCCAGCAGCAAGAGCACCA GCGGCGGCACAGCCGCTCTGGGCTGCCTGGTTCGAGGACTACTTC CCCGAGCCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCC GGCGTGCACACCTTCCCGCGGTGCTGCAGAGTTCTGGCCTGTAT AGCCTGAGCAGCGTGGTACCGTGCCTTCTAGCAGCCTGGGCACC CAGACCTACATCTGCAACGTGAACCAAGCCAGCAACACCAAG GTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACACA TGCCCAACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAGTC TTCCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGA CCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACC CTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATA ATGCCAAGACAAAGCCGCGGAGGAGCAGTACAACAGCACGTACC GTGTGGTCAGCGTCTTCAACGCTCTGCACCAAGGACTGGCTGAATG GCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCC CCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAAC CACAGGTGTGCACCTGCCCCATCCCGGATGAGCTGACCAAGA ACCAGGTACGCTCTCGTGCAGCTCAAAGGCTTCTATCCCAGCG ACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACT ACAAGACCAGCCTCCCGTGTGGACTCCGACGGCTCTTCTTCC TCGTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGG AACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACT ACACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	162
pETR16860 Herceptarg common CLkRK	GACATCCAGATGACCCAGAGCCCAAGCAGCCTGTCTGCCAGCGTG GGCGACAGAGTGACCATCACATGCAAGGCCAGCCAGGACGTGTCC ACAGCCGTGGCTGGTATCAGCAGAAGCTTGGCAAGGCCCAAG CTGCTGATCTACAGCGCCAGCTTCCGGTACACCGCGGTGCCAGC AGATTACGCGCAGCAGATCCGGCACCGACTTACCTTGACCATC AGCTCCCTGCAGCCCGAGGACTTCGCCACCTACTACTGCCAGCAG CACTACACCACCCCCCAATTGGCCAGGGACCAAGGTGGAA ATCAAGCGTACGGTGGCTGCACCATGTCTTCTTCTTCCCGCAT	163

-continued

EXAMPLARY SEQUENCES		
	CTGATCGGAAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCT GAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAGGTGGAT AACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAG GACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCCTGCCAAGTCA CCCATCAGGGCCTGAGCTCGCCCGTCCAAAGAGCTTCAACAGGG GAGAGTGTAG	
pETR17606 anti CD3 (CH2527 VH_3-23(12) VL7-46(13)) scFv 4.32.63 non cleavable linker aHerceptin VH CH1 EE CH2527- VL7_46-13 CH1 hum Fc knob PG LALA	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC AGCTACGGCGTGTATGGGTGCGGCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGCGCAGCGCAGCACCATTACAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAAGGCATCACACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGCGCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGCGGAGGAAGTGGCGGAGG GGGATCTGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT GCCGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTACGCGGCGAGCGAAGCG GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTCGCGCACCAAGCTGGAATCAAGGCGGGGGAGGCTCCG GAGGCGGCGAAGTGGAGGCGGCGAAGTGGCGGAGGCGGAGG GGGGGGAAGTGGGGCGGAGGAGTGGGGGGGAGGATCCAG GCCGTGCTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGC ACCGTGACCTGACATGTGGCAGTTCTACAGGCGCGTGACACC AGCAACTACGCCAACTGGGTGACGGAAGCCCGGCCAGGCCTTC AGAGGACTGATCGCGGCACCAACAGAGAGCCCCTGGCACCCCT GCCAGATTACGCGGATCTCTGCTGGGAGGAAAGGCGCCCTGACA CTGTCTGGCGCCAGCCAGAAGATGAGGCGGAGTACTACTGCGCC CTGTGGTACAGCAACCTGTGGGTGTTGCGGCGGAGGACCAAGCTG ACAGTGCTGAGCAGCGCTTCCACCAAGGGACCCAGTGTGTTCCCC CTGGCCCCCAGCTCCAAGTCTACATCCGGTGGCAGAGCTGCCCTG GGATGTCTCTGTAAGGACTACTTTCTGAGCCTGTGACAGTGTCTT GGAACAGCGAGCCCTGACCAGCGGAGTGACACATTCCTGCAG TGCTGCAGAGCAGCGCCCTGTATAGCCTGAGCAGCGTCTGTACCG TGCCTTCTCTAGCCTGGGAACACAGACATATATCTGTAATGTGAAT CATAAGCCCAGTAATACCAAAGTGGATAAGAAAGTGGAACTTAAGA GCTGCGATGGCGGAGGAGGGTCCGAGGCGGAGGGTCCGAGGT CCAGCTGGTTCGAGTCTGGAGGAGGACTGGTGCAGCCAGGCGGAT CTCTGAGACTGAGCTGCGCGCCAGCGGATTCAACATCAAGGACA CCTACATCCACTGGGTGAGGCGAGGCCCTGGAAGGGACTGGAGT GGGTGGCCAGAATCTACCCACCAACGGCTACACAAGATACGCCG ACAGCGTGAAGGGCAGATTACCATCAGCGCGACACCAAGCAAGA ACACCGCCTACTGTCAGATGAACAGCCTGAGAGCCGAGGACACAG CCGTGTACTACTGCTCTAGATGGGAGGCGAGGGCTTCTACGCCA TGGACTACTGGGACAGGGCACACTGGTGACCGTGTCCAGCGCTA GCACCAAGGGCCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGA GCACAGCGCGCGCACAGCCGCTCTGGGCTGCCTGGTTCGAGGAC TACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTG ACCTCCGGCGTGCACACCTTCCCCGCGTGTCTGCAGAGTTCGGC CTGTATAGCCTGAGCAGCGTGGTACCGTGCCTTCTAGCAGCCTG GGCACCCAGACTACATCTGCAACGTGAACCACAAGCCAGCAAC ACCAAGTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAAC CACACATGCCCCACCGTGGCCAGCACCTGAAGCTGCAGGGGAGC GTAGTCTTCTCTTCCCCCAAAACCAAGGACACCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAG GTGCAATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAAACAGC ACGTACCGTGTGGTACGCTCTCACCGTCTGACACAGGACTGG CTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGCCCTC GGCGCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCC CGAGAACCAAGGTGTACACCTTGGCCCCATGCCGGGATGAGCTG ACCAAGAACCAGGTACGCTGTGGTGCCTGGTCAAAGGCTTCTATC CCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGACGCGGAG AACAACATACAGACACGCTCCCGTGTGGACTCCGACGGCTCC TTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAG CAGGGGAACGTCTTCTCATGCTCCGTGATGATGAGGCTCTGCAC AACCATACACGAGAAGAGCCTCTCCCTGTCTCCGGGTAATGA	164

-continued

EXAMPLARY SEQUENCES

pETR17607	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC	165
anti CD3	CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAGCCTGACC	
(CH2527	AGTACGGCGTGTCTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG	
VH_3-23 (12)	GAATGGCTGGGCATCATCTGGGGCGACGGCAGCACCATTACCAC	
VL7-46 (13))	AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG	
scFv 4.32.63	AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC	
MMP9	GCCACCTACTACTGCGCCAAGGCATCACCACCGTGGTGGACGAC	
Matriptase	TACTACGCTATGGACTACTGGGGCCAGGGCACCAGCGTGACAGTG	
MK062	TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG	
aHerceptin VH	GGGATCTGGGGGAGGCGGAAGCGATATCCAGATGACCCAGAGCC	
CH1 EE	CTGCCAGCCTGTCTGCCTCTGTGGGCGAGACAGTGACCATCACAT	
CH2527-	GCCGGGCCAGCGAGAATCATGACAGCTACCTGGCCTGGTATCAGC	
VL7_46-13	AGAAGCAGGGCAAGAGCCCCAGCTGTGGTGTACGCCGCCACCT	
CH1 hum Fc	TTCTGGCCGACGATGTGCCAGCAGATTAGCGGCAGCGGAAGCG	
knob PG	GCACACAGTACAGCCTGAAGATCAACTCCCTGCAGAGCGAGGACG	
LALA	TGGCCCGGTACTACTGCGCAGCACTACTACAGCACCCCTACACCTT	
	CGGCTGCGGCACCAAGCTGGAATCAAAGGAGGCGGCGGAAGTG	
	TGCACATGCCCCCTGGGCTTCTGGGCCCCAGACAGGCCAGAGTCG	
	TGAACGGGGGGGCGGAGGCAGTGGGGGGGAGGATCCCAGGC	
	CGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGCAC	
	CGTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCACCAG	
	CAACTACGCCCAACTGGGTGCAGGAAAGCCCCGCCAGGCCTCAG	
	AGGACTGATCGGCGGCACCAACAAGAGAGCCCCCTGGCACCCCTGC	
	CAGATTGAGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTGACACT	
	GTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCT	
	GTGGTACAGCAACCTGTGGGTGTTTCGGCGGAGGCACCAAGCTGAC	
	AGTGCTGAGCAGCGCTTCCACCAAGGGAGCCAGTGTGTTCCCCCT	
	GGCCCCAGCTCCAAGTCTACATCCGGTGGCAGAGTGCCTGGG	
	ATGTCTCTGTAAGGACTACTTTCTGAGCCTGTGACAGTGTCTTGG	
	AACAGCGGAGCCCTGACAGCGGAGTGACACATTCCTGCGAGTG	
	GTGCAGAGCAGCGCCCTGTATAGCCTGAGCAGCGTGTGACCGTG	
	CCTTCTCTAGCCTGGGAACACAGACATATATCTGTAATGTGAATCA	
	TAAGCCCAAGTAATACCAAGTGGATAAGAAAGTGGAACTTAAGAGC	
	TGCGATGGCGGAGGAGGTTCCGGAGGCGGAGGGTCCGAGGTCCA	
	GCTGGTTCAGTCTGGAGGAGGACTGGTGACCCAGGCGGATCTCT	
	GAGACTGAGCTGCGCCGCCAGCGGATTCACATCAAGGACACCTA	
	CATCCACTGGGTGAGGCAGGCCCTTGGAAAGGAGTGGAGTGGG	
	TGGCCAGAACTTACCCACCAACGGCTACACAAGATACGCCGACA	
	GCGTGAAGGGCAGATTACCATCAGCGCCGACACAGCAGAAACA	
	CCGCCCTACCTGCAGATGAACAGCCTGAGAGCCGAGGACACAGCCG	
	TGTACTACTGCTCTAGATGGGGAGGCGAGGGCTTCTACGCCATGG	
	ACTACTGGGGACAGGGCACACTGGTGACCGTGTCCAGCGCTAGCA	
	CCAAGGGCCCTCCGTGTTCCCTTGGCCCCAGCAGCAAGAGCA	
	CCAGCGGCGGCACAGCCGCTCTGGGCTGCCTGGTTCGAGGACTAC	
	TTCCCCGAGCCCGTGACCGTGTCTTGGAAAGCGGAGCCCTGACC	
	TCCGGCGTGACACCTTCCCCGCCGTGCTGCAGAGTTCTGGCCGTG	
	TATAGCCTGAGCAGCGTGGTACCCGTGCCTTCTAGCAGCCTGGGC	
	ACCCAGACCTACATCTGCAACGTGAACCAAGCCAGCAACACCA	
	AGGTGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACA	
	CATGCCACCGTGGCCAGCACCTGAAGCTGCAGGGGGACCGTCA	
	GTCTTCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCCC	
	GGACCCCTGAGGTCAATGCGTGGTGGTGGACGTGAGCCACGAA	
	GACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTG	
	CATAATGCCAAGACAAGCCGCGGGAGGAGCAGTACAACAGCAGC	
	TACCGTGTGGTACGCGTCTTACCCTCTTGCACCAAGGACTGGCTG	
	AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCGGC	
	GCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGA	
	GAACCAAGGTGTACACCTGCCCCCATGCCGGATGAGCTGACC	
	AAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCCA	
	GCGACATCGCCGTGGAGTGGGAGAGCAATGGGAGCCGGAGAAAC	
	AACACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTC	
	TTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAG	
	GGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACC	
	ACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGTAAATGA	
Fo1R1 36F2	CAGGTGCAATTGGTTCAATCTGGTGTGAAGTAAAAAACCGGGCG	166
VH CH1 EE	CTTCCGTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC	
Fc hole PG	CTATTACATGCACCTGGGTTCTGTCAAGCCCCGGGCCAGGGTCTGGA	
LALA	ATGGATGGGCATCATTAACCAAGCGGTGGCTCTACCTCTACGC	
pETR14797	GCAGAAATCCAGGGTTCGCTCACGATGACCCATGACACTAGCAC	
	CTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACT	
	CGAGTGTACTACTGTGCACGCTCTTCTTCACTGGTTTCCATCTGG	
	ACTATTGGGGTCAAGGCACCTCGTAACGGTTTCTTCTGTAGCAC	

-continued

EXAMPLARY SEQUENCES

	CAAGGGCCCCCTCCGTGTTCCCCCTGGCCCCAGCAGCAAGAGCAC CAGCGCGGCACAGCCGCTCTGGCTGCTTGGTCGAGGACTACTT CCCCGAGCCCGTGACCGTGCTTGGAAACAGCGAGCCCTGACCTC CGGCGTGACACCTTCCCCGCGTGCTGCAGAGTTCTGGCTGTA TAGCCTGAGCAGCGTGGTCACCGTGCTTCTAGCAGCCTGGGCAC CCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAA GGTGGACGAGAAGGTGGAGCCCCAAGAGCTGCGACAAAACCTCACAC ATGCCACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAGT CTTCCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCCCGG ACCCCTGAGGTACATGCGTGGTGGTGGAGCTGAGCCACGAAGAC CCTGAGGTCAAGTTCAACTGGTACGTGGACGCGTGGAGGTGCAT AATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCAGTAC CGTGTGGTCAGCGTCTCACCCTGCTGCACAGGACTGGCTGAAT GGCAAGGAGTACAAGTGCAAGGTCTCCACAAAGCCCTCGCGCC CCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAA CCACAGGTGTGCACCCCTGCCCCCATCCCGGATGAGCTGACCAAG AACCAGGTACGCTCTCGTGCGCAGTCAAAGGCTTCTATCCAGC GACATCGCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA CTACAAGACACGCTCCCGTGCTGGACTCCGACGCTCTCTTCTT CCTCGTGAGCAAGCTCACCCTGGACAAGAGCAGGTGGCAGCAGG GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCA CTACACGAGAAAGGCTCTCCCTGTCTCCGGTAAATGA	
FolR1 36F2 VL Ck RK, pETR14798	GAAATCGTGTTAACGCAGTCTCCAGGCACCCGTCTTGTCTCCAG GGGAAAGAGCCACCCCTCTTTCAGGGCCAGTCAGAGTGTAGCA GCAGCTACTTAGCCTGGTACCAGCAGAACTGGCCAGGCTCCCA GGCTCCTCATCTATGGAGCATCCAGCAGGGCCACTGGCATCCAG ACAGGTTAGTGGCAGTGGATCCGGGACAGACTTCACTCTCACCAT CAGCAGACTGGAGCTGAAGATTGTCAGTGTATTACTGTCAGCAG TATACCAACGAACATTATTATACGTTTCGGCCAGGGGACCAAGTGG AAATCAAACGTAACGGTGGCTGCACCATCTGTCTTCTCATCTTCCGCC ATCTGATCGGAAGTTGAAATCTGGAACCTGCTTGTGTGTGCTG CTGAATAACTTCTATCCAGAGAGGCCAAAGTACAGTGAAGGTGG ATAACGCCCCCCTCAATCGGGTAACTCCAGGAGAGTGTACAGAGC AGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCCGACGC TGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCTCGCAAGT CACCCATCAGGGCTGAGCTCGCCCGTCACAAAGAGCTTCAACAG GGAGAGGTGTTAG	167
anti CD3 (CH2527 VH_3-23(12) VL7-46(13)) scFv 4.32.63 MMP9 Matriptase MK062 aFolR1 36F2 VH CH1 EE CH2527- VL7_46-13 CH1 hum Fc knob PG LALA pETR17621	CAAGTGCAGCTGAAAGAGTCCGGCCCTGGACTGGTGGCCCTAGC CAGAGCCTGAGCATCACCTGTACCGTGTCCGGCTTCAAGCTGACC AGCTACGGCGTGTATGGGTGCGCCAGCCTCCAGGCAAGTGTCTG GAATGGCTGGGCATCATCTGGGCGCAGCGCAGCACAATTAACAC AGCGCCCTGATCAGCAGACTGAGCATCTCCAAGGACAACAGCAAG AGCCAGGTGTTCTGAAGCTGAACAGCCTGCAGACCGACGACACC GCCACCTACTACTGCGCCAGGGCATCACACCGTGGTGGACGAC TACTACGCTATGGACTACTGGGGCAGGGCACCAGCGTGACAGTG TCTAGCGGAGGCGGAGGATCTGGCGGCGGAGGAAGTGGCGGAGG GGGATCTGGGGGAGGCGGAAGCAGATATCCAGATGACCCAGAGCC CTGCCAGCCTGTCTGCCTCTGTGGCGGAGACAGTGACCATCACAT GCCGGGCCAGCGAGAACATCGACAGCTACCTGGCCTGGTATCAGC AGAAGCAGGGCAAGAGCCCCAGCTGCTGGTGTACGCCGCCACCT TTCTGGCCGACGATGTGCCAGCAGATTACGCGGCAGCGGAAGCG GCACACAGTACAGCCTGAAGATCACTCCCTGCAGAGCGAGGACG TGGCCCGGTACTACTGCCAGCACTACTACAGCACCCCTACACCTT CGGCTGCGGCACCAAGCTGGAATCAAAGGAGGCGCGGAAGTG TGACATGCCCCCTGGGCTTCTGGGCCCCAGACAGGCCAGAGTCG TGAACGGGGGGGGCGGAGGAGTGGGGGGGAGGATCCAGGC CGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGCGGCAC CGTGACCCCTGACATGTGGCAGTTCTACAGCGCGCTGACACCCAG CAACTACGCCAAGTGGGTGCAGGAAAGCCCGGCCAGGCCTTCAAG AGGACTGATCGGCGGCACCAACAGAGAGCCCTGGCACCCCTGC CAGATTACAGCGGATCTCTGCTGGGAGGAAGGCCGCCCTGACACT GTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCT GTGGTACAGCAACCTGTGGTGTTCGGCGGAGGCACCAAGCTGAC AGTGTGAGCAGCGCTTCCACCAAGGAGCCAGTGTGTTCCCTT GGCCCCAGCTCCAAGTCTACATCCGGTGGCACAGCTGCCCTGGG ATGTCTCGTGAAGGACTACTTCTGAGCCTGTGACAGTGTCTTGG AACAGCGAGCCCTGACAGCGGAGTGCACACATTCCCTGCAGTG CTGCAGAGCAGCGGCTGTATAGCCTGAGCAGCGTGTGACCGTG CCTTCTCTAGCCTGGGAACACAGACATATCTGTAATGTGAATCA TAAGCCAGTAATACCAAGTGGATAAAGAAAGTGAACCTAAGAGC TGCGATGGCGGAGGAGGTCCGAGGCGGAGGTCCAGGTGCA ATTGTTCAATCTGGTGTGAAGTAAAAAACCGGGCGCTTCCGTT	168

-continued

EXAMPLARY SEQUENCES

anti CD3 (CH2527 VH ₃ -23(12) VL7-46(13)) scFv 4.32.63 non cleavable linker aFolR1 36F2 VH CH1 EE CH2527- VL7_46-13 CH1 hum Fc knob PG LALA pETR17622	AAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACAT GCACTGGGTTTCGTCAGCCCCGGGCCAGGGTCTGGAATGGATGG GCATCATTAACCCAAGCGGTGGCTCTACCTCTACGCGCAGAAATT CCAGGGTCGCGTCACGATGACCCATGACACTAGCACCTCTACCGT TTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCAGTGTAC TACTGTGCACGCTCTTTCTTCACTGGTTTCCATCTGGACTATTGGG GTCAAGGCACCCCTCGTAACGGTTTCTTCTGCTAGCACCAAGGGCC CCTCCGTGTTCCCGCTGGCCCCAGCAGCAAGAGCACCAAGCGGC GGCACAGCCGCTCTGGGCTGCCTGGTCGAGGACTACTTCCCCGAG CCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCTCCGGCGT GCACACCTTCCCGCGCTGCTGCAGAGTTTGGCCTGTATAGCCT GAGCAGCGTGGTCACCGTGCCTTCTAGCAGCCTGGGCACCCAGAC CTACATCTGCAACGTGAACCACAAAGCCAGCAACACCAAGGTGGA CGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTACACATGCC ACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAGTCTTCCCT CTTCCCCCAAAACCAAGGACACCTCATGATCTCCCGGACCCCT GAGGTCAATGCGTGGTGGTGGACGTGAGCCACGAAGACCTGAG GTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCC AAGACAAAGCCGCGGAGGAGCAGTACAAAGCAGCAGTACCGTGTG GTCAGCGTCTTACCGTCTGACCCAGGACTGGCTGAATGGCAAG GAGTACAAGTGCAAGGTCTCCAACAAGCCCTCGGCGCCCCCATC GAGAAAACCATCTCAAAGCCAAAGGGCAGCCCCGAGAACCAACAG GTGTACACCTGCCCCCATGCCGGGATGAGCTGACCAAGAACCCAG GTCAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGAGAAACAACTACAA GACCACGCCCTCCCGTGTGGACTCCGACGGCTCTTCTTCTCTCTA CAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACG TCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACTACTACAC CGAGAAGAGCCTCTCCCTGTCTCCGGTAATGA	169

-continued

EXAMPLARY SEQUENCES

Fo1R1 36F2 classic format: CH2527 XFab 36F2 HC EE Fc knob PG LALA pETR17623	TGTCCTGGAACAGCGGAGCCCTGACCTCCGGCGTGACACCTTCC CCGCCGTGCTGCAGAGTTCTGGCCTGTATAGCCTGAGCAGCGTGG TCACCGTGCTTCTAGCAGCCTGGGCACCCAGACCTACATCTGCAA CGTGAACCACAAGCCCAGCAACACCAAGGTGGACGAGAAGGTGGA GCCCAAGAGCTGCGCAAAACTCACACATGCCACCGTGCCCGAGC ACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAA ACCCAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACATG CGTGGTGGTGGACGTGAGCCACGAAGACCCCTGAGGTCAAGTTCAA CTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCC GCGGGAGGAGCAGTACAACAGCAGTACCGTGTGGTCAGCGTCCT CACCGTCTGACACAGGACTGGCTGAATGGCAAGGAGTACAAGTG CAAGGTCTCCAACAAAGCCCTCGGCGCCCCATCGAGAAAACCAT CTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCT GCCCCCATGCCGGGATGAGCTGACCAAGAACCAGGTACAGCTGTG GTGCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTG GGAGAGCAATGGGCAGCCGGAGAAACAATAAGAACACGCTTCC CGTGTGGACTCCGACGGCTCCTTCTCTCTACAGCAAGCTCACC GTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCC GTGATGCATGAGGCTCTGCACAACCACTACACGAGAAGAGCCTC TCCTGTCTCCGGTAAATGA CAGGCCGTCTGTGACCCAGGAACCCAGCCTGACAGTGTCTCTGGC GGCACCGTGACCCGTGACATGTGGCAGTTCTACAGGCGCCGTGACC ACCAGCAACTACGCCAACTGGGTGCAGGAAAAGCCCGGCCAGGCC TTCAGAGGACTGATCGGCGGCACCAACAAGAGAGCCCTGGCACC CCTGCCAGATTAGCGGATCTCTGCTGGGAGGAAAGCCGCCCTG ACACTGTCTGGCGCCAGCCAGAGATGAGGCCGAGTACTACTGC GCCCTGTGGTACAGCAACCTGTGGGTGTTCGGCGGAGGCACCAAG CTGACAGTGTGAGCAGCGCTTCCACCAAGGGACCCAGTGTGTTC CCCCTGGCCCCAGCTCCAAGTCTACATCCGGTGGCACAGCTGCC CTGGGATGTCTCGTGAAGGACTACTTCTCTGAGCCTGTGACAGTGT CTTGGAACAGCGGAGCCCTGACCAGCGGAGTGCACACATTCCTG CAGTGTGTCAGAGCAGCGCTGTATAGCCTGAGCAGCGTCTGTGA CCGTGCTTCTCTAGCCTGGGAACACAGACATATATCTGTAATGT GAATCATAAGCCAGTAATACCAAGTGGATAAGAAAGTGAACCT AAGAGCTGCGATGGCGGAGGAGGTCTGGAGCGGAGGGTCCCA GGTGCAATTGGTTCAATCTGGTGTGAAGTAAAAAACCGGGCGCT TCCGTTAAAGTGAGCTGCAAGCATCCGGATACACCTTCACTTCCT ATTACATGCACCTGGGTTCTGCAAGCCCCGGGCCAGGGTCTGGAAT GGATGGGCATCATTAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCAGGGTCGCGTCACGATGACCCATGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTCTGGAAGATACTGCA GTGTACTACTGTGCACGCTCTTCTTCACTGGTTTCCATCTGGACTA TTGGGTCAAGGCACCTCGTAACGGTTTCTTCTGCTAGCACCAAG GGCCCTCCGTGTTCCCTTGGCCCCCAGCAGCAAGAGCACCAGC GGCAGCACAGCCGCTCTGGGCTGCCTGGTCGAGGACTACTTCCCC GAGCCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCCGG CGTGACACCTTCCCCGCCGTGTGACAGTCTTGCCCTGTATAG CCTGAGCAGCGTGGTCACCGTGCTTCTAGCAGCCTGGGCACCCA GACCTACATCTGCAACGTGAACACAAGCCAGCAACACCAAGGT GGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACACATG CCCACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAGTCTT CCTTCTCCCCCAAACCAAGGACACCTCATGATCTCCCGGACC CCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCTT GAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAAT GCCAAGACAAAGCCGCGGAGGAGCAGTACAACAGCACGTACCGT GTGGTCAGCGTCTCACCCTCTGCACACGAGTGGCTGAATGGC AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCC ATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCA CAGGTGTACACCTGCCCCCATGCGGGATGAGTGAACAGAAGC CAGGTACGCTGTGGTGCCTGGTCAAAGGCTTCTATCCAGCGAC ATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGAGAGAACAATAC AAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTTCTCTCTCT ACAGCAAGCTACCGTGGACAAGAGCAGGTGGCAGCAGGGGAAC GTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA CGCAGAAGAGCCTCTCCTGTCTCCGGTAAATGA	170

[0834] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and

examples should not be construed as limiting the scope of the invention. The disclosures of all patent and scientific literature cited herein are expressly incorporated in their entirety by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 170

<210> SEQ ID NO 1

<211> LENGTH: 215

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 1

```
Gln Ala Val Val Thr Gln Glu Pro Ser Leu Thr Val Ser Pro Gly Gly
1      5      10      15
Thr Val Thr Leu Thr Cys Gly Ser Ser Thr Gly Ala Val Thr Thr Ser
20     25     30
Asn Tyr Ala Asn Trp Val Gln Glu Lys Pro Gly Gln Ala Phe Arg Gly
35     40     45
Leu Ile Gly Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe
50     55     60
Ser Gly Ser Leu Leu Gly Gly Lys Ala Ala Leu Thr Leu Ser Gly Ala
65     70     75     80
Gln Pro Glu Asp Glu Ala Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn
85     90     95
Leu Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln Pro
100    105    110
Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu
115    120    125
Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro
130    135    140
Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys Ala
145    150    155    160
Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala
165    170    175
Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Arg
180    185    190
Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr
195    200    205
Val Ala Pro Thr Glu Cys Ser
210    215
```

<210> SEQ ID NO 2

<211> LENGTH: 971

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 2

```
Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
1      5      10      15
Thr Val Arg Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20     25     30
```

Ser	Ile	His	Trp	Val	Lys	Gln	Ala	Pro	Gly	Lys	Cys	Leu	Lys	Trp	Met
35															
Gly	Trp	Ile	Asn	Thr	Glu	Thr	Gly	Glu	Pro	Ala	Tyr	Ala	Asp	Asp	Phe
50															
Lys	Gly	Arg	Phe	Ala	Phe	Ser	Leu	Glu	Thr	Ser	Ala	Ser	Thr	Ala	Tyr
65															
Leu	Gln	Ile	Asn	Asn	Leu	Lys	Asn	Glu	Asp	Thr	Ala	Thr	Phe	Phe	Cys
Ala	His	Pro	Tyr	Asp	Tyr	Asp	Val	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
115															
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Thr	Val	Leu	Thr	Gln
130															
Ser	Pro	Ala	Ser	Leu	Gly	Val	Ser	Leu	Gly	Gln	Arg	Ala	Thr	Ile	Ser
145															
Cys	Arg	Ala	Ser	Lys	Ser	Val	Ser	Thr	Ser	Asn	Tyr	Ser	Tyr	Ile	His
Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	Lys	Leu	Leu	Ile	Lys	Tyr
Val	Ser	Tyr	Leu	Glu	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly
195															
Ser	Gly	Thr	Asp	Phe	Thr	Leu	Asn	Ile	His	Pro	Val	Glu	Glu	Glu	Asp
210															
Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	His	Ser	Arg	Glu	Phe	Pro	Trp	Thr	Phe
225															
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly
Gly	Gly	Ser	Arg	Gln	Ala	Arg	Val	Val	Asn	Gly	Gly	Gly	Gly	Gly	Ser
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Leu	Glu
275															
Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys
290															
Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Thr	Tyr	Ala	Met	Asn	Trp	Val	Arg
305															
Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Arg	Ile	Arg	Ser	Lys
Tyr	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe
340															
Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn
355															
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Arg	His	Gly
370															
Asn	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly
385															
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp

-continued

435					440					445					
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
450						455					460				
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
465					470					475					480
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
				485					490					495	
Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Gly
			500					505					510		
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Val	Glu	Ser
		515					520					525			
Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala
530						535					540				
Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Ala	Trp	Met	Ser	Trp	Val	Arg	Gln
545					550					555					560
Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Gly	Arg	Ile	Lys	Ser	Lys	Thr
			565						570					575	
Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	Pro	Val	Lys	Gly	Arg	Phe	Thr
			580					585					590		
Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser
		595					600					605			
Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Thr	Thr	Pro	Trp	Glu
610						615					620				
Trp	Ser	Trp	Tyr	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser
625				630						635					640
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser
			645						650					655	
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp
		660					665						670		
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr
		675					680					685			
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr
690						695					700				
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln
705				710						715					720
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp
			725						730					735	
Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro
		740						745					750		
Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
		755					760					765			
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
		770				775					780				
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn
785					790					795					800
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
			805						810					815	
Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
			820					825					830		
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser
		835					840					845			

-continued

```

Asn Lys Ala Leu Gly Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
 850                      855                      860

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Cys Arg Asp
 865                      870                      875                      880

Glu Leu Thr Lys Asn Gln Val Ser Leu Trp Cys Leu Val Lys Gly Phe
                      885                      890                      895

Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
          900                      905                      910

Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
          915                      920                      925

Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly
 930                      935                      940

Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr
 945                      950                      955                      960

Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
          965                      970

```

```

<210> SEQ ID NO 3
<211> LENGTH: 450
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 3

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1          5          10          15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20          25          30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35          40          45

Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50          55          60

Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
 65          70          75          80

Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85          90          95

Tyr Cys Thr Thr Pro Trp Glu Trp Ser Trp Tyr Asp Tyr Trp Gly Gln
 100         105         110

Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115         120         125

Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130         135         140

Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 145         150         155         160

Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 165         170         175

Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180         185         190

Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195         200         205

Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp
 210         215         220

```

-continued

Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly
 225 230 235 240
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 245 250 255
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 290 295 300
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 305 310 315 320
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Gly Ala Pro Ile Glu
 325 330 335
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Cys
 340 345 350
 Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
 355 360 365
 Ser Cys Ala Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 385 390 395 400
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Val Ser Lys Leu Thr Val Asp
 405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 420 425 430
 Glu Ala Leu His Asn Arg Phe Thr Gln Lys Ser Leu Ser Leu Ser Pro
 435 440 445
 Gly Lys
 450

<210> SEQ ID NO 4
 <211> LENGTH: 971
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 4

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
 1 5 10 15
 Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu
 35 40 45
 Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile
 50 55 60
 Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
 65 70 75 80
 Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala
 85 90 95
 Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp
 100 105 110

Gly 115	Gln	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
Gly 130	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
Gln 145	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Thr
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Ala
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Ala
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Gly
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Asp
Val 225	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Phe
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly
Gly	Gly	Ser	Arg	Gln	Ala	Arg	Val	Val	Asn	Gly	Gly	Gly	Gly	Ser
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Glu
Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Cys
Ala 305	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Thr	Tyr	Ala	Met	Asn	Trp	Arg
Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Arg	Ile	Arg	Lys
Tyr	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Phe
Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Asn
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Arg	Gly
Asn 385	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gly
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Phe
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Leu
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Trp
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Leu
Gln 465	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Ser
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Pro
Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Gly

-continued

Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Val	Glu	Ser
		515						520				525			
Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala
	530					535					540				
Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Ala	Trp	Met	Ser	Trp	Val	Arg	Gln
545					550					555					560
Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Gly	Arg	Ile	Lys	Ser	Lys	Thr
			565						570					575	
Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	Pro	Val	Lys	Gly	Arg	Phe	Thr
			580					585					590		
Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser
		595					600					605			
Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Thr	Thr	Pro	Trp	Glu
	610					615					620				
Trp	Ser	Trp	Tyr	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser
625					630						635				640
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser
				645					650					655	
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp
			660					665					670		
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr
		675					680					685			
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr
	690					695					700				
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln
705					710					715					720
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp
				725					730					735	
Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro
		740						745					750		
Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
		755					760					765			
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
	770					775					780				
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn
785					790					795					800
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
				805					810					815	
Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
				820				825					830		
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser
		835					840					845			
Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys
		850				855					860				
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg	Asp
865					870					875					880
Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly	Phe
				885				890						895	
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu
			900					905					910		
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe

-continued

915	920	925
Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly		
930	935	940
Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr		
945	950	955 960
Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys		
	965	970

<210> SEQ ID NO 5
 <211> LENGTH: 971
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 5

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln		
1	5	10 15
Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr		
	20	25 30
Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu		
	35	40 45
Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile		
	50	55 60
Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu		
	65	70 75 80
Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala		
	85	90 95
Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp		
	100	105 110
Gly Gln Gly Thr Ser Val Thr Val Ser Ser Gly Gly Gly Ser Gly		
	115	120 125
Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser Asp Ile		
	130	135 140
Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly Glu Thr		
	145	150 155 160
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala		
	165	170 175
Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val Tyr Ala		
	180	185 190
Ala Thr Phe Leu Ala Asp Asp Val Pro Ser Arg Phe Ser Gly Ser Gly		
	195	200 205
Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser Glu Asp		
	210	215 220
Val Ala Arg Tyr Tyr Cys Gln His Tyr Tyr Ser Thr Pro Tyr Thr Phe		
	225	230 235 240
Gly Cys Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Ser Gly Gly		
	245	250 255
Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Gly Gly Ser Gly Ser		
	260	265 270
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu		
	275	280 285
Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys		

-continued

290					295					300					
Ala 305	Ala	Ser	Gly	Phe	Thr 310	Phe	Ser	Thr	Tyr	Ala 315	Met	Asn	Trp	Val	Arg 320
Gln	Ala	Pro	Gly	Lys 325	Gly	Leu	Glu	Trp	Val 330	Ser	Arg	Ile	Arg	Ser 335	Lys
Tyr	Asn	Asn	Tyr	Ala 340	Thr	Tyr	Tyr	Ala 345	Asp	Ser	Val	Lys	Gly 350	Arg	Phe
Thr	Ile	Ser 355	Arg	Asp	Asp	Ser	Lys 360	Asn	Thr	Leu	Tyr	Leu 365	Gln	Met	Asn
Ser 370	Leu	Arg	Ala	Glu	Asp	Thr 375	Ala	Val	Tyr	Tyr	Cys 380	Val	Arg	His	Gly
Asn 385	Phe	Gly	Asn	Ser	Tyr 390	Val	Ser	Trp	Phe	Ala 395	Tyr	Trp	Gly	Gln	Gly 400
Thr	Leu	Val	Thr	Val 405	Ser	Ser	Ala	Ser	Thr 410	Lys	Gly	Pro	Ser	Val 415	Phe
Pro	Leu	Ala	Pro	Ser 420	Ser	Lys	Ser	Thr 425	Ser	Gly	Gly	Thr 430	Ala	Ala	Leu
Gly	Cys	Leu 435	Val	Lys	Asp	Tyr	Phe 440	Pro	Glu	Pro	Val	Thr 445	Val	Ser	Trp
Asn 450	Ser	Gly	Ala	Leu	Thr 455	Ser	Gly	Val	His	Thr	Phe 460	Pro	Ala	Val	Leu
Gln 465	Ser	Ser	Gly	Leu	Tyr 470	Ser	Leu	Ser	Ser	Val 475	Val	Thr	Val	Pro	Ser 480
Ser	Ser	Leu	Gly	Thr 485	Gln	Thr	Tyr	Ile	Cys 490	Asn	Val	Asn	His	Lys 495	Pro
Ser	Asn	Thr	Lys 500	Val	Asp	Lys	Lys	Val 505	Glu	Pro	Lys	Ser 510	Cys	Asp	Gly
Gly	Gly	Gly 515	Ser	Gly	Gly	Gly	Gly 520	Ser	Glu	Val	Gln	Leu 525	Val	Glu	Ser
Gly 530	Gly	Gly	Leu	Val	Lys 535	Pro	Gly	Gly	Ser	Leu	Arg 540	Leu	Ser	Cys	Ala
Ala 545	Ser	Gly	Phe	Thr	Phe 550	Ser	Asn	Ala	Trp	Met 555	Ser	Trp	Val	Arg	Gln 560
Ala	Pro	Gly	Lys	Gly 565	Leu	Glu	Trp	Val	Gly 570	Arg	Ile	Lys	Ser	Lys 575	Thr
Asp	Gly	Gly	Thr	Thr 580	Asp	Tyr	Ala	Ala 585	Pro	Val	Lys	Gly 590	Arg	Phe	Thr
Ile	Ser	Arg 595	Asp	Asp	Ser	Lys	Asn 600	Thr	Leu	Tyr	Leu 605	Gln	Met	Asn	Ser
Leu 610	Lys	Thr	Glu	Asp	Thr 615	Ala	Val	Tyr	Tyr	Cys 620	Thr	Thr	Pro	Trp	Glu
Trp 625	Ser	Trp	Tyr	Asp	Tyr 630	Trp	Gly	Gln	Gly	Thr 635	Leu	Val	Thr	Val	Ser 640
Ser	Ala	Ser	Thr	Lys 645	Gly	Pro	Ser	Val	Phe 650	Pro	Leu	Ala	Pro	Ser 655	Ser
Lys	Ser	Thr	Ser	Gly 660	Gly	Thr	Ala	Ala 665	Leu	Gly	Cys	Leu 670	Val	Lys	Asp
Tyr	Phe	Pro 675	Glu	Pro	Val	Thr	Val 680	Ser	Trp	Asn	Ser	Gly 685	Ala	Leu	Thr
Ser 690	Gly	Val	His	Thr	Phe 695	Pro	Ala	Val	Leu	Gln	Ser 700	Ser	Gly	Leu	Tyr

-continued

Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln
 705 710 715 720
 Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp
 725 730 735
 Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro
 740 745 750
 Cys Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe Leu Phe Pro
 755 760 765
 Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr
 770 775 780
 Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn
 785 790 795 800
 Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg
 805 810 815
 Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val
 820 825 830
 Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser
 835 840 845
 Asn Lys Ala Leu Gly Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
 850 855 860
 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Cys Arg Asp
 865 870 875 880
 Glu Leu Thr Lys Asn Gln Val Ser Leu Trp Cys Leu Val Lys Gly Phe
 885 890 895
 Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
 900 905 910
 Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
 915 920 925
 Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly
 930 935 940
 Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr
 945 950 955 960
 Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 965 970

<210> SEQ ID NO 6
 <211> LENGTH: 971
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 6

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15
 Thr Val Arg Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Ser Ile His Trp Val Lys Gln Ala Pro Gly Lys Cys Leu Lys Trp Met
 35 40 45
 Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Ala Tyr Ala Asp Asp Phe
 50 55 60
 Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80

-continued

Leu	Gln	Ile	Asn	Asn	Leu	Lys	Asn	Glu	Asp	Thr	Ala	Thr	Phe	Phe	Cys
			85						90					95	
Ala	His	Pro	Tyr	Asp	Tyr	Asp	Val	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
			100					105					110		
Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
		115				120						125			
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Thr	Val	Leu	Thr	Gln
	130					135					140				
Ser	Pro	Ala	Ser	Leu	Gly	Val	Ser	Leu	Gly	Gln	Arg	Ala	Thr	Ile	Ser
145				150						155					160
Cys	Arg	Ala	Ser	Lys	Ser	Val	Ser	Thr	Ser	Asn	Tyr	Ser	Tyr	Ile	His
			165						170					175	
Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	Lys	Leu	Leu	Ile	Lys	Tyr
		180						185					190		
Val	Ser	Tyr	Leu	Glu	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly
		195					200					205			
Ser	Gly	Thr	Asp	Phe	Thr	Leu	Asn	Ile	His	Pro	Val	Glu	Glu	Glu	Asp
	210					215					220				
Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	His	Ser	Arg	Glu	Phe	Pro	Trp	Thr	Phe
225					230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly
			245						250					255	
Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ser
		260						265					270		
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Leu	Glu
	275					280						285			
Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys
	290					295					300				
Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Thr	Tyr	Ala	Met	Asn	Trp	Val	Arg
305					310					315					320
Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Arg	Ile	Arg	Ser	Lys
			325						330					335	
Tyr	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe
		340						345					350		
Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn
	355					360						365			
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Arg	His	Gly
	370				375						380				
Asn	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly
385				390						395					400
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
			405						410					415	
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
			420					425					430		
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
		435					440					445			
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
	450					455					460				
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
465				470						475					480

-continued

Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
				485					490					495	
Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Gly
			500					505					510		
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Val	Glu	Ser
		515					520					525			
Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala
	530					535					540				
Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Ala	Trp	Met	Ser	Trp	Val	Arg	Gln
545					550					555					560
Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Gly	Arg	Ile	Lys	Ser	Lys	Thr
			565						570					575	
Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	Pro	Val	Lys	Gly	Arg	Phe	Thr
			580					585					590		
Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser
		595					600					605			
Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Thr	Thr	Pro	Trp	Glu
	610					615					620				
Trp	Ser	Trp	Tyr	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser
625				630						635					640
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser
			645						650					655	
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp
		660						665					670		
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr
		675					680					685			
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr
	690					695					700				
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln
705				710						715					720
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp
			725						730					735	
Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro
		740						745					750		
Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
		755					760					765			
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
	770				775						780				
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn
785				790						795					800
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
			805						810					815	
Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
		820						825					830		
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser
	835					840						845			
Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys
	850					855					860				
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg	Asp
865				870						875					880
Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly	Phe

-continued

885								890					895				
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu		
			900						905							910	
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe		
			915						920							925	
Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly		
			930						935							940	
Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr		
			945						950							955	
Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys							
			965						970								

```
<210> SEQ ID NO 7
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 7

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Arg Gln Ala Arg Val Val
1 5 10 15

Asn Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly
20 25 30

Ser

```
<210> SEQ ID NO 8
<211> LENGTH: 28
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 8

Gly Gly Gly Gly Ser Val His Met Pro Leu Gly Phe Leu Gly Pro Gly
1 5 10 15

Arg Ser Arg Gly Ser Phe Pro Gly Gly Gly Gly Ser
20 25

```
<210> SEQ ID NO 9
<211> LENGTH: 41
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 9

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Arg Gln Ala Arg Val Val
1 5 10 15

Asn Gly Gly Gly Gly Gly Ser Val Pro Leu Ser Leu Tyr Ser Gly Gly
20 25 30

Gly Gly Gly Ser Gly Gly Gly Gly Ser
35 40

```
<210> SEQ ID NO 10
<211> LENGTH: 36
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
```

-continued

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 10

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Arg Gln Ala Arg Val Val
1 5 10 15

Asn Gly Val Pro Leu Ser Leu Tyr Ser Gly Gly Gly Gly Gly Ser Gly
20 25 30

Gly Gly Gly Ser
35

<210> SEQ ID NO 11

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 11

Thr Tyr Ala Met Asn
1 5

<210> SEQ ID NO 12

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 12

Arg Ile Arg Ser Lys Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp Ser
1 5 10 15

Val Lys Gly

<210> SEQ ID NO 13

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 13

His Gly Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe Ala Tyr
1 5 10

<210> SEQ ID NO 14

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 14

Asn Ala Trp Met Ser
1 5

<210> SEQ ID NO 15

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 15

-continued

Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala Pro
1 5 10 15

Val Lys Gly

<210> SEQ ID NO 16
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 16

Pro Trp Glu Trp Ser Trp Tyr Asp Tyr
1 5

<210> SEQ ID NO 17
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 17

Gly Ser Ser Thr Gly Ala Val Thr Thr Ser Asn Tyr Ala Asn
1 5 10

<210> SEQ ID NO 18
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 18

Gly Thr Asn Lys Arg Ala Pro
1 5

<210> SEQ ID NO 19
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 19

Ala Leu Trp Tyr Ser Asn Leu Trp Val
1 5

<210> SEQ ID NO 20
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 20

Asp Tyr Ser Ile His
1 5

<210> SEQ ID NO 21
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 21

Trp Ile Asn Thr Glu Thr Gly Glu Pro Ala Tyr Ala Asp Asp Phe Lys
1 5 10 15

Gly

<210> SEQ ID NO 22
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 22

Pro Tyr Asp Tyr Asp Val Leu Asp Tyr
1 5

<210> SEQ ID NO 23
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 23

Arg Ala Ser Lys Ser Val Ser Thr Ser Asn Tyr Ser Tyr Ile His
1 5 10 15

<210> SEQ ID NO 24
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 24

Tyr Val Ser Tyr Leu Glu Ser
1 5

<210> SEQ ID NO 25
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 25

Gln His Ser Arg Glu Phe Pro Trp Thr
1 5

<210> SEQ ID NO 26
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 26

Ser Tyr Gly Val Ser
1 5

-continued

<210> SEQ ID NO 27
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 27

Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile Ser
1 5 10 15

<210> SEQ ID NO 28
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 28

Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr
1 5 10

<210> SEQ ID NO 29
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 29

Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala
1 5 10

<210> SEQ ID NO 30
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 30

Ala Ala Thr Phe Leu Ala Asp
1 5

<210> SEQ ID NO 31
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 31

Gln His Tyr Tyr Ser Thr Pro Tyr Thr
1 5

<210> SEQ ID NO 32
<211> LENGTH: 450
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 32

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Phe	Thr	Phe	Thr	Asp	Tyr	
			20							25				30		
Lys	Ile	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
			35				40				45					
Gly	Tyr	Phe	Asn	Pro	Asn	Ser	Gly	Tyr	Ser	Thr	Tyr	Ala	Gln	Lys	Phe	
			50				55				60					
Gln	Gly	Arg	Val	Thr	Ile	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr	
			65				70				75				80	
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85							90				95		
Ala	Arg	Leu	Ser	Pro	Gly	Gly	Tyr	Tyr	Val	Met	Asp	Ala	Trp	Gly	Gln	
			100							105				110		
Gly	Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	
			115				120				125					
Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	
			130				135				140					
Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	
			145				150				155				160	
Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	
			165							170				175		
Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	
			180							185				190		
Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	
			195				200				205					
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	
			210				215				220					
Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	
			225				230				235				240	
Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	
			245							250				255		
Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	
			260							265				270		
Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	
			275				280				285					
Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	
			290				295				300					
Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	
			305				310				315				320	
Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	
			325							330				335		
Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	
			340							345				350		
Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	
			355				360				365					
Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	
			370				375				380					
Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	
			385				390				395				400	
Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	

-continued

Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
420 425 430

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
435 440 445

Gly Lys
450

<210> SEQ ID NO 33
<211> LENGTH: 213
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 33

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Asn Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
35 40 45

Tyr Asn Thr Asn Asn Leu Gln Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Phe Pro Thr
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130 135 140

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145 150 155 160

Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
165 170 175

Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
180 185 190

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
195 200 205

Asn Arg Gly Glu Cys
210

<210> SEQ ID NO 34
<211> LENGTH: 492
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 34

Glu Val Gln Leu Glu Gln Ser Gly Pro Val Leu Val Lys Pro Gly Thr
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Tyr	Ile	Asn	Trp	Ile	Ile	Gln	Ser	His	Gly	Lys	Cys	Leu	Glu	Trp	Ile
35															
Gly	Val	Ile	Asn	Pro	Asp	Ser	Gly	Gly	Thr	Asp	Tyr	Asn	Gln	Asn	Phe
50															
Lys	Gly	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Thr	Thr	Ala	Tyr
65															
Met	Glu	Leu	Thr	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
85															
Ala	Arg	Arg	Asp	Ser	Tyr	Gly	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Thr
100															
Leu	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly
115															
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile	Val	Leu	Thr	Gln	Thr
130															
Pro	Lys	Phe	Leu	Leu	Val	Pro	Ala	Gly	Asp	Arg	Ile	Thr	Met	Thr	Cys
145															
Lys	Ala	Ser	Leu	Ser	Val	Thr	Asn	Asp	Val	Ala	Trp	Tyr	Gln	Gln	Lys
165															
Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Leu	Tyr	Tyr	Ala	Ser	Asn	Arg	Asn
180															
Ala	Gly	Val	Pro	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Tyr	Gly	Thr	Asp	Phe
195															
Thr	Phe	Thr	Ile	Thr	Thr	Leu	Gln	Ala	Glu	Asp	Leu	Ala	Val	Tyr	Phe
210															
Cys	Gln	Gln	Asp	Tyr	Thr	Ser	Pro	Pro	Thr	Phe	Gly	Cys	Gly	Thr	Lys
225															
Leu	Glu	Ile	Arg	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Pro
245															
Leu	Gly	Leu	Trp	Ser	Gln	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
260															
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser
275															
Ser	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala
290															
Ser	Gln	Gly	Ile	Asn	Asn	Tyr	Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly
305															
Lys	Ala	Pro	Lys	Arg	Leu	Ile	Tyr	Asn	Thr	Asn	Asn	Leu	Gln	Thr	Gly
325															
Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu
340															
Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Leu
355															
Gln	His	Asn	Ser	Phe	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile
370															
Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp
385															
Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn
405															
Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu
420															
430															

-continued

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
 435 440 445

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
 450 455 460

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 465 470 475 480

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 485 490

<210> SEQ ID NO 35
 <211> LENGTH: 35
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 35

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Pro Leu Gly Leu Trp
 1 5 10 15

Ser Gln Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 20 25 30

Ser Gly Gly
 35

<210> SEQ ID NO 36
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 36

Arg Gln Ala Arg Val Asn Gly
 1 5

<210> SEQ ID NO 37
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 37

Val His Met Pro Leu Gly Phe Leu Gly Pro Gly Arg Ser Arg Gly Ser
 1 5 10 15

Phe Pro

<210> SEQ ID NO 38
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct
 <220> FEATURE:
 <221> NAME/KEY: X
 <222> LOCATION: (9)..(13)
 <223> OTHER INFORMATION: X is an amino acid
 <220> FEATURE:
 <221> NAME/KEY: X
 <222> LOCATION: (9)..(13)
 <223> OTHER INFORMATION: X or Xaa is an amino acid

<400> SEQUENCE: 38

-continued

Arg Gln Ala Arg Val Val Asn Gly Xaa Xaa Xaa Xaa Xaa Val Pro Leu
1 5 10 15

Ser Leu Tyr Ser Gly
20

<210> SEQ ID NO 39
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 39

Arg Gln Ala Arg Val Val Asn Gly Val Pro Leu Ser Leu Tyr Ser Gly
1 5 10 15

<210> SEQ ID NO 40
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 40

Pro Leu Gly Leu Trp Ser Gln
1 5

<210> SEQ ID NO 41
<211> LENGTH: 249
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 41

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
1 5 10 15

Thr Val Arg Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Ser Ile His Trp Val Lys Gln Ala Pro Gly Lys Cys Leu Lys Trp Met
35 40 45

Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Ala Tyr Ala Asp Asp Phe
50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80

Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Phe Phe Cys
85 90 95

Ala His Pro Tyr Asp Tyr Asp Val Leu Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Ser Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
115 120 125

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Thr Val Leu Thr Gln
130 135 140

Ser Pro Ala Ser Leu Gly Val Ser Leu Gly Gln Arg Ala Thr Ile Ser
145 150 155 160

Cys Arg Ala Ser Lys Ser Val Ser Thr Ser Asn Tyr Ser Tyr Ile His
165 170 175

-continued

Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	Lys	Leu	Leu	Ile	Lys	Tyr
		180						185					190		
Val	Ser	Tyr	Leu	Glu	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly
		195					200					205			
Ser	Gly	Thr	Asp	Phe	Thr	Leu	Asn	Ile	His	Pro	Val	Glu	Glu	Glu	Asp
	210					215					220				
Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	His	Ser	Arg	Glu	Phe	Pro	Trp	Thr	Phe
225					230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys							
				245											

<210> SEQ ID NO 42
 <211> LENGTH: 249
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 42

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1				5					10					15	
Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Ser	Tyr
			20					25					30		
Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
		35					40					45			
Gly	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
	50					55					60				
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
65					70					75					80
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
			85						90					95	
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
		100						105					110		
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
		115					120					125			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
	130					135						140			
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
145					150					155					160
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala
			165						170					175	
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala
		180						185					190		
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
		195					200					205			
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
	210					215					220				
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
225					230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys							
				245											

<210> SEQ ID NO 43
 <211> LENGTH: 125

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 43

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr
20 25 30
Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Arg Ile Arg Ser Lys Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp
50 55 60
Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
65 70 75 80
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85 90 95
Tyr Cys Val Arg His Gly Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe
100 105 110
Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> SEQ ID NO 44
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 44

Thr Tyr Ala Met Asn
1 5

<210> SEQ ID NO 45
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 45

Arg Ile Arg Ser Lys Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp Ser
1 5 10 15
Val Lys Gly

<210> SEQ ID NO 46
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 46

His Gly Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe Ala Tyr
1 5 10

<210> SEQ ID NO 47
<211> LENGTH: 120
<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 47

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30
Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
50 55 60
Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
65 70 75 80
Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
85 90 95
Tyr Cys Thr Thr Pro Trp Glu Trp Ser Trp Tyr Asp Tyr Trp Gly Gln
100 105 110
Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 48
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 48

Asp Tyr Tyr Ile Asn
1 5

<210> SEQ ID NO 49
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 49

Val Ile Asn Pro Asp Ser Gly Gly Thr Asp Tyr Asn Gln Asn Phe Lys
1 5 10 15
Gly

<210> SEQ ID NO 50
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 50

Arg Asp Ser Tyr Gly Phe Asp Tyr
1 5

<210> SEQ ID NO 51
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 51

Lys Ala Ser Leu Ser Val Thr Asn Asp Val Ala
1 5 10

<210> SEQ ID NO 52
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 52

Tyr Ala Ser Asn Arg Asn Ala
1 5

<210> SEQ ID NO 53
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 53

Gln Gln Asp Tyr Thr Ser Pro Pro Thr
1 5

<210> SEQ ID NO 54
<211> LENGTH: 207
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 54

Met Gln Ser Gly Thr His Trp Arg Val Leu Gly Leu Cys Leu Leu Ser
1 5 10 15

Val Gly Val Trp Gly Gln Asp Gly Asn Glu Glu Met Gly Gly Ile Thr
20 25 30

Gln Thr Pro Tyr Lys Val Ser Ile Ser Gly Thr Thr Val Ile Leu Thr
35 40 45

Cys Pro Gln Tyr Pro Gly Ser Glu Ile Leu Trp Gln His Asn Asp Lys
50 55 60

Asn Ile Gly Gly Asp Glu Asp Asp Lys Asn Ile Gly Ser Asp Glu Asp
65 70 75 80

His Leu Ser Leu Lys Glu Phe Ser Glu Leu Glu Gln Ser Gly Tyr Tyr
85 90 95

Val Cys Tyr Pro Arg Gly Ser Lys Pro Glu Asp Ala Asn Phe Tyr Leu
100 105 110

Tyr Leu Arg Ala Arg Val Cys Glu Asn Cys Met Glu Met Asp Val Met
115 120 125

Ser Val Ala Thr Ile Val Ile Val Asp Ile Cys Ile Thr Gly Gly Leu
130 135 140

Leu Leu Leu Val Tyr Tyr Trp Ser Lys Asn Arg Lys Ala Lys Ala Lys
145 150 155 160

Pro Val Thr Arg Gly Ala Gly Ala Gly Gly Arg Gln Arg Gly Gln Asn
165 170 175

Lys Glu Arg Pro Pro Pro Val Pro Asn Pro Asp Tyr Glu Pro Ile Arg

-continued

180	185	190
Lys Gly Gln Arg Asp Leu Tyr	Ser Gly Leu Asn Gln Arg	Arg Ile
195	200	205

<210> SEQ ID NO 55
<211> LENGTH: 109
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 55

Gln Ala Val Val Thr Gln Glu Pro Ser Leu Thr Val Ser Pro Gly Gly
1 5 10 15
Thr Val Thr Leu Thr Cys Gly Ser Ser Thr Gly Ala Val Thr Thr Ser
20 25 30
Asn Tyr Ala Asn Trp Val Gln Glu Lys Pro Gly Gln Ala Phe Arg Gly
35 40 45
Leu Ile Gly Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe
50 55 60
Ser Gly Ser Leu Leu Gly Gly Lys Ala Ala Leu Thr Leu Ser Gly Ala
65 70 75 80
Gln Pro Glu Asp Glu Ala Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn
85 90 95
Leu Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu
100 105

<210> SEQ ID NO 56
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 56

Asp Tyr Lys Ile His
1 5

<210> SEQ ID NO 57
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 57

Tyr Phe Asn Pro Asn Ser Gly Tyr Ser Thr Tyr Ala Gln Lys Phe Gln
1 5 10 15

Gly

<210> SEQ ID NO 58
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 58

Leu Ser Pro Gly Gly Tyr Tyr Val Met Asp Ala
1 5 10

-continued

<210> SEQ ID NO 59
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 59

Arg Ala Ser Gln Gly Ile Asn Asn Tyr Leu Asn
1 5 10

<210> SEQ ID NO 60
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 60

Asn Thr Asn Asn Leu Gln Thr
1 5

<210> SEQ ID NO 61
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 61

Leu Gln His Asn Ser Phe Pro Thr
1 5

<210> SEQ ID NO 62
<211> LENGTH: 648
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 62

caggccgctcg tgaccaggga acccagcctg acagtgtctc ctggcggcac cgtgaccctg 60
acatgtggca gttctacagg cgccgtgacc accagcaact acgccaactg ggtgcaggaa 120
aagcccggcc aggccttcag aggactgac gccggcacca acaagagagc ccctggcacc 180
cctgccagat tcagcggatc tctgctggga ggaaaggccg ccctgacact gtctggcgcc 240
cagccagaag atgaggccga gtactactgc gccctgtggt acagcaacct gtgggtgttc 300
ggcggaggca ccaagctgac agtcctaggt caaccaagg ctgccccag cgtgaccctg 360
ttcccccca gcagcgagga actgcaggcc aacaaggcca ccctggtctg cctgatcagc 420
gacttctacc caggcgccgt gaccgtggcc tggaaggccg acagcagccc cgtgaaggcc 480
ggcgtggaga ccaccacccc cagcaagcag agcaacaaca agtacgccgc cagcagctac 540
ctgagcctga ccccccagca gtggaagagc cacaggctct acagctgccca ggtgaccac 600
gagggcagca ccgtggagaa aaccgtggcc cccaccgagt gcagctga 648

<210> SEQ ID NO 63
<211> LENGTH: 2916
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 63

```

cagatccagc tgggtgcagag cggccctgag ctgaagaaac ccggcgagac agtgcggatc    60
agctgcaagg ccagcggcta caccttcacc gactacagca tccactgggt caagcaggcc    120
cctggcaagt gcctgaagtg gatgggctgg atcaacacgg agacaggcga gcccgctac    180
gccgacgatt tcaagggcag attcgcttc agcctggaaa ccagcgccag caccgcctac    240
ctgcagatca acaacctgaa gaacgaggac accgccacct tttctgcgc ccaccctac    300
gactacgacg tgctggatta ttggggccag ggcaccagcg tgaccgtgc tagcggaggc    360
ggaggatctg gcggcggagg aagtggcgga ggggatctg ggggaggcgg atctgatacc    420
gtgctgacac agagccctgc cagcctggga gtgtccctgg gacagagagc caccatcagc    480
tgctggggcca gcaagagcgt gtccaccagc aactacagct atatccactg gtatcagcag    540
aagcccggcc agccccccaa gctgctgac aaatacgtgt cctacctgga aagcggcgtg    600
cccgccagat tttctggctc tggcagcggc accgacttca ccctgaacat ccaccctg    660
gaagaggaag atgccgccac ctactactgc cagcacagca gagagttccc ttggaccttc    720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtaga    780
caggccagag tcgtgaacgg gggagggggg ggaagtgggg gcggaggcag tgggggggga    840
ggatccgagg tgcagctgct ggaatctggc ggcggactgg tgcagcctgg cggatctctg    900
agactgagct gtgccggcag cgcttcacc ttcagcacct acgccaatgaa ctgggtgcgc    960
caggccccctg gcaaaaggcct ggaatgggtg tcccgatca gaagcaagta caacaactac   1020
gccacctact acgccgacag cgtgaagggc cggttcacca tcagccggga cgacagcaag   1080
aacacctgt acctgcagat gaacagcctg cgggcccagg acaccgccgt gtactattgt   1140
gtgcggcacg gcaactctcg caacagctat gtgtcttggg ttgcctactg gggccagggc   1200
acctctgtga ccgtgtcaag cgctagcaca aagggcccta gcgtgttccc tctggcccc   1260
agcagcaaga gcacaagcgg cggaacagcc gccctgggct gcctcgtgaa ggactacttc   1320
cccgagcccc tgacagtgtc ttggaacagc ggagccctga caagcggcgt gcacaccttc   1380
cctgccgtgc tgcagagcag cggcctgtac tccctgagca gcgtggtcac cgtgcctagc   1440
agcagcctgg gcacccagac ctacatctgc aacgtgaacc acaagcccag caacacccaaa   1500
gtggacaaga aggtggagcc caagagctgt gatggcggag gagggtcagg aggcggaggga   1560
tccgaggtgc aattgggtga atctggtggg ggtctggtaa aaccgggccc ttcctgcgt   1620
ctgagctgcg cggttccgg attcaccttc tccaacgcgt ggatgagctg gggtcgcag   1680
gccccgggca aaggcctcga gtgggttggg cgtatcaagt ctaaaactga cgggtggcacc   1740
acggattacg cggctccagt taaaggtcgt ttaccattt ccgcgacga tagcaaaaac   1800
actctgtatc tgcagatgaa ctctctgaaa actgaagaca ccgcagtcta ctactgtact   1860
accccgctggg aatggctctg gtacgattat tggggccagg gcacgctggg tacgggtgtc   1920
agcgctagta ccaagggccc cagcgtgttc cccctggcac ccagcagcaa gagcacatct   1980
ggcggaacag ccgctctggg ctgtctgggt aaagactact tccccgagcc cgtgaccgtg   2040
tcttggaaact ctggcgccct gaccagcggc gtgcacacct ttccagccgt gctgcagagc   2100

```

-continued

agcggcctgt actccctgtc ctccgtggtc accgtgccct ctactccct gggaacacag	2160
acatatatct gtaatgtcaa tcacaagcct tccaacacca aagtcgataa gaaagtcgag	2220
cccaagagct gcgacaaaac tcacacatgc ccaecgtgcc cagcacctga agctgcaggg	2280
ggaccgtcag tcttctcttt ccccccaaaa cccaaggaca ccctcatgat ctcccggacc	2340
cctgaggtca catgctgggt ggtggacgtg agccacgaag accctgaggt caagtccaac	2400
tggtagctgg acggcgtgga ggtgcataat gccaaagaca agccgcggga ggagcagtac	2460
aacagcacgt accgtgtgggt cagcgtcttc accgtctgc accaggactg gctgaatggc	2520
aaggagtaca agtgcaaggt ctccaacaaa gccctcggcg ccccatcgga gaaaaccatc	2580
tccaagcca aagggcagcc ccgagaacca caggtgtaca ccctgcccc atgccgggat	2640
gagctgacca agaaccaggt cagcctgtgg tgcctggta aaggttcta tcccagcgac	2700
atcgccgtgg agtgggagag caatgggcag ccggagaaca actacaagac cagcctccc	2760
gtgctggact ccgacggctc cttctctctc tacagcaagc tcaccgtgga caagagcagg	2820
tggcagcagg ggaacgtctt ctcatgtctc gtgatgcag aggtctctga caaccactac	2880
acgcagaaga gcctctccct gtctccgggt aatga	2916

<210> SEQ ID NO 64

<211> LENGTH: 1353

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 64

gaggtgcaat tgggtgaatc tgggtgggtg ctggtaaaac cgggcgggtc cctgcgtctg	60
agctcgcggg cttccggatt cacctctctc aacgcgtgga tgagctgggt tgcgcaggcc	120
ccgggcaaaag gcctcgagtg ggttgggtcgt atcaagtcta aaactgacgg tggcaccacg	180
gattacgcgg ctccagttaa aggtcgtttt accatttccc gcgacgatag caaaaacact	240
ctgtatctgc agatgaactc tctgaaaact gaagacaccg cagtctacta ctgtactacc	300
ccgtgggaat ggtcttggtg cgattattgg ggccagggca cgctgggttac ggtgtcttcc	360
gctagcacca agggcccttc cgtgttcccc ctggccccc gcagcaagag caccagcggc	420
ggcacagccg ctctgggctg cctgggtcaag gactacttcc ccgagcccg gaccgtgtcc	480
tggaaacagc gagccctgac ctccggcgtg cacaccttcc ccgcgtgct gcagagttct	540
ggcctgtata gcctgagcag cgtgggtcacc gtgccttcta gcagcctggg caccagacc	600
tacatctgca acgtgaacca caagcccagc aacaccaagg tggacaagaa ggtggagccc	660
aagagctgcg acaaaactca cacatgccca ccgtgccag cacctgaagc tgcaggggga	720
ccgtcagttc tcctcttccc cccaaaaccc aaggaccccc tcatgatctc ccggacccct	780
gaggtcacat gcgtgggtgt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg	840
tacgtggaag gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac	900
agcacgtacc gtgtggctag cgtcctcacc gtctgcacc aggactgggt gaatggcaag	960
gagtacaagt gcaaggtctc caacaaagcc ctccggcccc ccacgagaa aaccatctcc	1020
aaagccaaag ggcagccccc agaaccacag gtgtgcaccc tgccccatc ccgggatgag	1080
ctgaccaaga accaggtcag cctctcgtgc gcagtcaaag gcttctatcc cagcgacatc	1140

-continued

gccgtggagt gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccgtg	1200
ctggactccg acggctcctt ctctcctcgtg agcaagctca ccgtggacaa gagcagggtg	1260
cagcagggga acgtctcttc atgctcctg atgcatgagg ctctgcacaa ccgcttcacg	1320
cagaagagcc tctccctgtc tccgggtaaa tga	1353

<210> SEQ ID NO 65
 <211> LENGTH: 2916
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 65

caagtgcagc tgaagagtc cgccctgga ctggtggccc ctaccagag cctgagcatc	60
acctgtaccg tgtccggtt cagcctgacc agctacggcg tgtcatgggt gcgccagcct	120
ccaggcaagt gtctggaatg gctgggcac atctggggcg acggcagcac caattaccac	180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctg	240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcaa gggcacaccc	300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagt	360
tctagcggag gcggaggatc tggcgcgga ggaagtggcg gagggggatc tgggggaggc	420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca	480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag	540
aagcagggca agagccccc gctgctggtg tacgcgcca cctttctggc cgacgatgtg	600
cccagcagat tcagcggcag cggaaagcggc acacagtaca gcctgaagat caactccctg	660
cagagcgagg acgtggcccg gtactactgc cagcactact acagcaccac ctacaccttc	720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtaga	780
caggccagag tcgtgaacgg gggagggggg ggaagtgggg gcggaggcag tgggggcgga	840
ggatccgagg tgcagctgct ggaatctggc ggcggactgg tgcagcctgg cggatctctg	900
agactgagct gtgcgcccag cggcttcacc ttcagcact acgccaatgaa ctgggtgcgc	960
caggccccctg gcaaaaggcct ggaatgggtg tcccggatca gaagcaagta caacaactac	1020
gccacctact acgcccagac cgtgaagggc cggttcacca tcagccggga cgacagcaag	1080
aacacctgt acctgcagat gaacagcctg cgggcccagg acaccgccgt gtactattgt	1140
gtgcggcacg gcaacttcgg caacagctat gtgtcttgggt ttgcctactg gggccagggc	1200
accctcgtga ccgtgtcaag cgtagcaca aagggcccta gcgtgttccc tctggccccc	1260
agcagcaaga gcacaagcgg cggaaacagc gccctgggct gcctcgtgaa ggactacttc	1320
cccagccccg tgacagtgtc ttggaacagc ggagccctga caagcggcgt gcacaccttc	1380
cctgccgtgc tgcagagcag cggcctgtac tccctgagca gcgtggtcac cgtgcctagc	1440
agcagcctgg gcacccagac ctacatctgc aacgtgaacc acaagcccag caacacccaa	1500
gtggacaaga aggtggagcc caagagctgt gatggcggag gaggggtccg gggcggagga	1560
tccgagggtc aattgggtga atctggtggt ggtctggtaa aaccgggcgg ttccctgcgt	1620
ctgagctgcg cggttcctcg gtacaccttc tccaacgcgt ggatgagctg ggttcgccag	1680
gccccgggca aaggccctga gtgggttgggt cgtatcaagt ctaaaaactga cgggtggcac	1740

-continued

acggattacg	cggtccagt	taaaggtcgt	tttaccattt	cccgcgacga	tagcaaaaac	1800
actctgtatc	tgcagatgaa	ctctctgaaa	actgaagaca	cgcagtceta	ctactgtact	1860
accccggtggg	aatggtcttg	gtacgattat	tggggccagg	gcacgctggt	tacgggtgtct	1920
agcgctagta	ccaagggccc	cagcgtgttc	cccctggcac	ccagcagcaa	gagcacatct	1980
ggcggaacag	cgcctctggg	ctgtctggtg	aaagactact	tccccgagcc	cgtgaccgtg	2040
tcttggaact	ctggcgccct	gaccagcggc	gtgcacacct	ttccagcgtg	gctgcagagc	2100
agcggcctgt	actccctgtc	ctcctgtgtc	accgtgccct	ctagctccct	gggaacacag	2160
acatatatct	gtaatgtcaa	tcacaagcct	tccaacacca	aagtgcataa	gaaagtcgag	2220
cccaagagct	gcgacaaaac	tcacacatgc	ccaccgtgcc	cagcacctga	agctgcaggg	2280
ggaccgtcag	tcttcctctt	ccccccaaaa	cccaaggaca	ccctcatgat	ctcccgacc	2340
cctgaggtea	catgctgggt	ggtggacgtg	agccacgaag	accctgaggt	caagttcaac	2400
tgggtacgtg	acggcgtgga	ggtgcataat	gccaaagaaa	agccgcggga	ggagcagtac	2460
aacagcacgt	accgtgtggg	cagcgtcttc	accgtcctgc	accaggactg	gctgaatggc	2520
aaggagtaca	agtgcagggt	ctccaacaaa	gccctcgccg	cccccatcga	gaaaaccatc	2580
tccaaagcca	aagggcagcc	ccgagaacca	caggtgtaca	ccctgcccc	atgccgggat	2640
gagctgacca	agaaccaggt	cagcctgtgg	tgcttggtca	aaggtctcta	tcccagcgac	2700
atcgccgtgg	agtgaggag	caatgggcag	ccggagaaca	actacaagac	cacgcctccc	2760
gtgctggact	ccgacggctc	cttctctctc	tacagcaagc	tcaccgtgga	caagagcagg	2820
tggcagcagg	ggaacgtctt	ctcatgctcc	gtgatgcatg	aggctctgca	caaccactac	2880
acgcagaaga	gcctctccct	gtctccgggt	aatga			2916

<210> SEQ ID NO 66

<211> LENGTH: 2916

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 66

caagtgcagc	tgaaagagtc	cggccctgga	ctggtggccc	ctagccagag	cctgagcatc	60
acctgtaccg	tgtccggctt	cagcctgacc	agctacggcg	tgtcatgggt	gcgccagcct	120
ccaggcaagt	gtctggaatg	gctgggcata	atctggggcg	acggcagcac	caattaccac	180
agcgccctga	tcagcagact	gagcatctcc	aaggacaaca	gcaagagcca	ggtgttcctg	240
aagctgaaca	gcctgcagac	cgacgacacc	gccacctact	actgcgcca	gggcataccc	300
accgtggtgg	acgactacta	cgctatggac	tactggggcc	agggcaccag	cgtgacagtg	360
tctagcggag	gcggaggatc	tggcggcgga	ggaagtggcg	gagggggatc	tgggggaggc	420
ggaagcgata	tccagatgac	ccagagccct	gccagcctgt	ctgcctctgt	gggcgagaca	480
gtgaccatca	catgccgggc	cagcgagaac	atcgacagct	acctggcctg	gtatcagcag	540
aagcagggca	agagccccca	gctgctggtg	tacgcgcgca	cctttctggc	cgacgatgtg	600
cccagcagat	tcagcggcag	cggaagcggc	acacagtaca	gcctgaagat	caactccctg	660
cagagcgagg	acgtggcccg	gtactactgc	cagcactact	acagcacccc	ctacaccttc	720
ggctgcggca	ccaagctgga	aatcaaaggc	gggggaggct	ccggaggcgg	cggaagtgga	780

-continued

ggcgccgga	gtggcggagg	cggagggggg	ggaagtgggg	gcggaggcag	tgggggggga	840
ggatccgagg	tgcagctgct	ggaatctggc	ggcggactgg	tgcagcctgg	cggatctctg	900
agactgagct	gtgccgccag	cggcttcacc	ttcagcacct	acgccatgaa	ctgggtgctg	960
caggccccctg	gcaaaggcct	ggaatgggtg	tcccggatca	gaagcaagta	caacaactac	1020
gccacctact	acgccgacag	cgtgaagggc	cggttcacca	tcagccggga	cgacagcaag	1080
aacaccctgt	acctgcagat	gaacagcctg	cgggccgagg	acaccgcgt	gtactattgt	1140
gtgcggcacg	gcaacttcgg	caacagctat	gtgtcttgg	ttgcctactg	gggccagggc	1200
accctcgtga	cctgtgtcaag	cgctagcaca	aaggggcccta	gcgtgttccc	tctggccccc	1260
agcagcaaga	gcacaagcgg	cggaaacagc	gccctgggct	gcctcgtgaa	ggactacttc	1320
cccgagcccg	tgacagtgtc	ttggaacagc	ggagccctga	caagcggcgt	gcacaccttc	1380
cctgccgtgc	tgcagagcag	cggcctgtac	tccctgagca	gcgtggtcac	cgtgcctagc	1440
agcagcctgg	gcacccagac	ctacatctgc	aacgtgaacc	acaagcccag	caacacccaaa	1500
gtggacaaga	aggtggagcc	caagagctgt	gatggcggag	gagggtccgg	aggcggaggc	1560
tccgaggtgc	aattgggtga	atctggtgg	ggtctggtaa	aaccgggcgg	ttccctgcgt	1620
ctgagctgcg	cggcttcogg	attcaccttc	tccaacgcgt	ggatgagctg	ggttcgccag	1680
gccccgggca	aaggcctcga	gtgggttgg	cgtatcaagt	ctaaaactga	cggtggcacc	1740
acggattacg	cggctccagt	taaaggctgt	tttaccattt	ccgcgcagca	tagcaaaaac	1800
actctgtatc	tgcagatgaa	ctctctgaaa	actgaagaca	ccgcagtcta	ctactgtact	1860
accccgctgg	aatggtcttg	gtacgattat	tggggccagg	gcacgctggt	tacggtgtct	1920
agcgtagta	ccaagggccc	cagcgtgttc	cccctggcac	ccagcagcaa	gagcacatct	1980
ggcggaacag	ccgtctggg	ctgtctggtg	aaagactact	tccccgagcc	cgtgaccgtg	2040
tcttggaact	ctggcgccct	gaccagcggc	gtgcacacct	ttccagccgt	gctgcagagc	2100
agcggcctgt	actccctgtc	ctccgtggtc	accgtgccct	ctagctccct	gggaacacag	2160
acatatatct	gtaatgtcaa	tcacaagcct	tccaacacca	aagtcgataa	gaaagtcgag	2220
cccaagagct	gcgacaaaac	tcacacatgc	ccaccgtgcc	cagcacctga	agctgcaggg	2280
ggaccgtcag	tcttctctct	ccccccaaaa	cccaaggaca	ccctcatgat	ctcccgacc	2340
cctgaggtea	catgcgtgg	ggtggacgtg	agccacgaag	accctgagg	caagttcaac	2400
tggtagctgg	acggcgtgga	ggtgcataat	gccaaagcaa	agccgcggga	ggagcagtac	2460
aacagcacgt	accgtgtgg	cagcgtcctc	accgtcctgc	accaggactg	gctgaatggc	2520
aaggagtaca	agtgcaaggt	ctccaacaaa	gccctcggcg	cccccatcga	gaaaaccatc	2580
tccaaagcca	aagggcagcc	ccgagaacca	caggtgtaca	ccctgcccc	atgccgggat	2640
gagctgacca	agaaccaggt	cagcctgtgg	tgcctggtea	aaggcttcta	tcccagcgac	2700
atcgccgtgg	agtgaggag	caatgggcag	ccggagaaca	actacaagac	cacgcctccc	2760
gtgctggact	ccgacggctc	cttcttcttc	tacagcaagc	tcaccgtgga	caagagcagg	2820
tggcagcagg	ggaacgtctt	ctcatgtctc	gtgatgcatg	aggctctgca	caaccactac	2880
acgcagaaga	gcctctccct	gtctccgggt	aaatga			2916

<210> SEQ ID NO 67

<211> LENGTH: 2916

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 67

```

cagatccagc tgggtcagag cggccctgag ctgaagaaac ccggcgagac agtgcggatc    60
agctgcaagg ccagcggcta caccttcacc gactacagca tccactgggt caagcaggcc    120
cctggcaagt gcctgaagtg gatgggctgg atcaacacog agacaggcga gcccgctac    180
gccgacgatt tcaagggcag attcgcttc agcctggaaa ccagcgccag caccgcctac    240
ctgcagatca acaacctgaa gaacgaggac accgccacct tttctgcgc ccaccctac    300
gactacgacg tgtcggatta ttggggccag ggcaccagcg tgaccgtgc tagcggaggc    360
ggaggatctg gcggcggagg aagtggcgga ggggatctg ggggaggcgg atctgatacc    420
gtgctgacac agagccctgc cagcctggga gtgtccctgg gacagagagc caccatcagc    480
tgtcggggcca gcaagagcgt gtccaccagc aactacagct atatccactg gtatcagcag    540
aagcccggcc agccccccaa gctgctgac aaatacgtgt cctacctgga aagcggcgtg    600
cccgccagat tttctggctc tggcagcggc accgacttca ccctgaacat ccaccctgtg    660
gaagaggaag atgccgccac ctactactgc cagcacagca gagagttccc ttggaccttc    720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtgga    780
ggcggcgga gtggcggagg cggagggggg ggaagtggg gcggaggcag tgggggggga    840
ggatccgagg tgcagctgct ggaatctggc ggcggactgg tgcagcctgg cggatctctg    900
agactgagct gtgcggccag cgcttcacc ttcagcacct acgccaatgaa ctgggtgcgc    960
caggccccct gcaaaaggcct ggaatgggtg tcccgatca gaagcaagta caacaactac   1020
gccacctact acgcccagag cgtgaagggc cggttcacca tcagccggga cgacagcaag   1080
aacacctgt acctgcagat gaacagcctg cgggcccagg acaccgccgt gtactattgt   1140
gtgcggcacg gcaactctcg caacagctat gtgtcttggg ttgcctactg gggccagggc   1200
acctctgtga ccgtgtcaag cgtagcaca aaggggcccta gcgtgttccc tctggcccc   1260
agcagcaaga gcacaagcgg cggaacagcc gccctgggct gcctcgtgaa ggactacttc   1320
cccgagcccg tgacagtgtc ttggaacagc ggagccctga caagcggcgt gcacaccttc   1380
cctgcctgct tgcagagcag cggcctgtac tccctgagca gcgtgggtcac cgtgcctagc   1440
agcagcctgg gcacccagac ctacatctgc aacgtgaacc acaagcccag caacacccaaa   1500
gtggacaaga aggtggagcc caagagctgt gatggcggag gaggtccgg aggcggaggc   1560
tccgaggtgc aattggttga atctggtggt ggtctggtaa aaccgggccc ttcctcgt   1620
ctgagctgct cggttccgg attcaccttc tccaacgcgt ggatgagctg ggttcgccag   1680
gccccgggca aaggcctcga gtgggttggg cgtatcaagt ctaaaactga cgggtggcacc   1740
acggattacg cggctccagt taaaggtcgt ttaccattt ccgcgacga tagcaaaaac   1800
actctgtatc tgcagatgaa ctctctgaaa actgaagaca ccgcagtcta ctactgtact   1860
accccgctgg aatggtcttg gtacgattat tggggccagg gcacgctggt tacggtgtct   1920
agcgctagta ccaagggccc cagcgtgttc cccctggcac ccagcagcaa gagcacatct   1980
ggcggaacag ccgctctggg ctgtctggtg aaagactact tccccgagcc cgtgaccgtg   2040
tcttggaaact ctggcgccct gaccagcggc gtgcacacct ttccagccgt gctgcagagc   2100

```

-continued

agcggcctgt actccctgtc ctccgtggtc accgtgcct ctactccct gggaacacag	2160
acatatatct gtaatgtcaa tcacaagcct tccaacacca aagtcgataa gaaagtcgag	2220
cccaagagct gcgacaaaac tcacacatgc ccaccgtgcc cagcacctga agctgcagg	2280
ggaccgtcag tcttctcttt ccccccaaaa cccaaggaca ccctcatgat ctcccgacc	2340
cctgaggtca catgctgggt ggtggacgtg agccacgaag accctgaggt caagtccaac	2400
tggtagctgg acggcgtgga ggtgcataat gccaaagaca agccgcggga ggagcagta	2460
aacagcacgt accgtgtgggt cagcgtcttc accgtctgc accaggactg gctgaatggc	2520
aaggagtaca agtgcaaggt ctccaacaaa gccctcggcg ccccatcgga gaaaaccatc	2580
tccaagcca aagggcagcc ccgagaacca caggtgtaca ccctgcccc atgccgggat	2640
gagctgacca agaaccaggt cagcctgtgg tgcctggta aaggttcta tcccagcag	2700
atcgccgtgg agtgggagag caatgggcag ccggagaaca actacaagac cagcctccc	2760
gtgctggact ccgacggctc cttctctctc tacagcaagc tcaccgtgga caagagcag	2820
tggcagcagg ggaacgtctt ctcatgctcc gtgatgcatg aggtctctga caaccactac	2880
acgcagaaga gcctctccct gtctccgggt aatga	2916

<210> SEQ ID NO 68
 <211> LENGTH: 99
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 68

ggcgggggag gctccggagg cggcggaagt agacaggcca gagtcgtgaa cggggggagg	60
gggggaagtg gggcgaggag cagtgggggc ggagatcc	99

<210> SEQ ID NO 69
 <211> LENGTH: 1350
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 69

caggtgcagc tggtcagag cggcgccgag gtgaagaaac ccgggtcttc tgtcaaggtg	60
tcatgcaagg ctacggatt cacctttaca gactacaaaa tccactgggt taggcaggca	120
cctggccaag gactcgaatg gatggggat ttcaacccaa attccggcta ctctacctat	180
gccagaagt ttcaggaag agtgactatt acagctgata agagtaccag cactgcatac	240
atggagctgt cctctcttcg ctacaggac accgcctct actattgtgc tcggctgagc	300
cccggtggct actatgtgat gtagcatgg ggcaggga caaccgtaac agtgcctct	360
gcgtcgacta agggcccttc agtttttcca ctgcaccca gtagcaagtc cacatctggg	420
ggtaaccgtg ccctgggctg ccttgtgaaa gactatttcc ctgaaccagt cactgtgtca	480
tggaaatagc gagccctgac ctccggtgta cacacattcc ccgctgtgtt gcagtctagt	540
ggcctgtaca gcctctcttc tgtgtgacc gtcccttcaa gctccctggg gacacagacc	600
tatatctgta acgtgaatca taagccatct aacactaaag tagataaaaa agtggagccc	660
aagagttgcy aaaaaacaca cacctgtccc cctgcccag ccccgagct tctgggaggc	720

-continued

cctagcgtct ttctcttccc acccaagcct aaggatactc tgatgatatc caggacccca	780
gaagttacat gcgtggctcgt ggacgtctca cacgaggacc ccgaagtga atttaactgg	840
tacgttgatg gtgtggaagt ccataatgcc aagaccaagc ctagagagga gcaatacaac	900
agtacatatc gcgtggtaag cgtgttgacc gttctccacc aggactggct caatgggaaa	960
gaatacaagt gtaaagtgtc caacaagct ctgccagcac ccatcgagaa gactatttct	1020
aaggccaaag gccagccccc ggagcctcag gtctatacac ttccaccctc aagggatgaa	1080
ctgaccaaga accaagttag cttgacttgc ctggtaaagg ggttctaccc ttccgacatc	1140
gctgtggagt gggagtctaa tggacaacca gaaaacaatt acaaaaccac accccctgtc	1200
ctcgacagtg atggcagctt ttctctgtat agcaaaacta ccgttgacaa gtccagatgg	1260
cagcagggaa acgtgttctc atgtagcgtc atgcacgaag ctttgcataa ccactacaca	1320
cagaaaagcc tcagcctgag tccagggaag	1350

<210> SEQ ID NO 70
 <211> LENGTH: 639
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct
 <400> SEQUENCE: 70

gacatccaaa tgaccagtc acctagtagc ctctccgcct ctgttgga cagggtgaca	60
attacatgca gagcttcaca gggatcaac aattaccta actggatca gcagaaacca	120
gggaaggccc ccaagcgtt gatataaac accaataacc tgcaaaactgg cgtccctagc	180
cgggtctccg gatctgtag tggcacgaa ttacactca ccatcagctc cctgcagcca	240
gaggatttgc ccacatacta ttgtcttcag cataattctt tccccacctt tgggcaagga	300
actaaactgg agattaagcg tactgtgcc gctccctctg tgttcatttt tcctccaagt	360
gatgagcagc tcaaaagcgg taccgcatcc gttgtgtgcc tgcttaacaa cttctatccc	420
cgggaagcca aggtccaatg gaaggtggac aatgctctgc agtcaggaaa cagtcaggag	480
agcgtaaccg agcaggattc caaagactct acttactcat tgagctccac cctgacactc	540
tctaaggcag actatgaaa gcataaagtg tacgcctgtg aggttaccca ccagggcctg	600
agtagccctg tgacaaagtc cttcaatagg ggagagtgc	639

<210> SEQ ID NO 71
 <211> LENGTH: 1476
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct
 <400> SEQUENCE: 71

gagggttcagc tggagcagtc aggacctgtg ctggtgaagc ctgggacttc agtgaagatg	60
tcctgtaagg cttctggata cacattcact gactactata taaactggat aatacagagc	120
catggaaagt gtcttgagtg gattggagtt attaatcctg acagcggtagg tactgactac	180
aaccagaact tcaagggcaa ggccacattg actgttgaca agtcctccac cacagcctac	240
atggaactca ctagccctgac atctgaggac tctgcagtct attattgtgc aagaagggat	300
tcttacggct ttgactactg gggccaaggc accactctca cagtctctc aggcggaggt	360

-continued

```

ggctcagggg gaggcggtag cggcggaggt ggctcagggg gaggcggtag cgacattgtg 420
ctgacccaga ctcccaaatt cctgcttggt ccagcaggag acaggattac catgacctgc 480
aaggccagtc tgagtgtgac taatgatgta gcttggtatc aacagaaacc agggcagtct 540
cctaaactgc tgttatacta tgcattcaat cgcaacgctg gagtcacctga tcgcttact 600
ggcagtggtat atgggacgga ttctactttc accatcacca ctttgaggc tgaagacctg 660
gcagtttatt tctgtcagca ggattatacc tctctccga cgttcggttg tggcaccaag 720
ctagaaatcc gtggtggcgg cggttctggc ggagggggtt ctggccccct ggggctatgg 780
agccagggtg gcggcggttc tggcggaggg ggttctggcg gtggtggctc tggcggtgac 840
atccaaatga ccagtcacc tagtagctc tccgcctctg ttggcgacag ggtgacaatt 900
acatgcagag cttcacaggg tatcaacaat tacctgaact ggtatcagca gaaaccaggg 960
aaggcccca agcgcttgat atataacacc aataacctgc aaactggcgt ccctagccgg 1020
ttctccggat ctggtagtgg caccgaattt aactcacca tcagctccct gcagccagag 1080
gatttcgcca catactattg tcttcagcat aattctttcc ccacctttgg gcaaggaact 1140
aaactggaga ttaagcgtag tgcgcgcgt cctctgtgt tcatttttcc tccaagtgat 1200
gagcagctca aaagcgtag cgcacggtt gtgtgcctgc ttaacaaact ctatccccgg 1260
gaagccaagg tccaatggaa ggtggacaat gctctgcagt caggaaacag tcaggagagc 1320
gtaaccgagc aggtattcaa agactctact tactcattga gctccacct gacactctct 1380
aaggcagact atgaaaagca taaagtgtac gcctgtgagg ttaccaccca gggcctgagt 1440
agccctgtga caaagtcctt caatagggga gagtgc 1476

```

<210> SEQ ID NO 72

<211> LENGTH: 971

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 72

```

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
1           5           10          15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20          25          30

Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu
35          40          45

Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile
50          55          60

Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
65          70          75          80

Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala
85          90          95

Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp
100         105         110

Gly Gln Gly Thr Ser Val Thr Val Ser Ser Gly Gly Gly Ser Gly
115         120         125

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Asp Ile
130         135         140

Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly Glu Thr

```

-continued

145	150	155	160
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala	165	170	175
Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val Tyr Ala	180	185	190
Ala Thr Phe Leu Ala Asp Asp Val Pro Ser Arg Phe Ser Gly Ser Gly	195	200	205
Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser Glu Asp	210	215	220
Val Ala Arg Tyr Tyr Cys Gln His Tyr Tyr Ser Thr Pro Tyr Thr Phe	225	230	235
Gly Cys Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Val His	245	250	255
Met Pro Leu Gly Phe Leu Gly Pro Arg Gln Ala Arg Val Val Asn Gly	260	265	270
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu	275	280	285
Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys	290	295	300
Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr Ala Met Asn Trp Val Arg	305	310	315
Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ser Arg Ile Arg Ser Lys	325	330	335
Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe	340	345	350
Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn	355	360	365
Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Val Arg His Gly	370	375	380
Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe Ala Tyr Trp Gly Gln Gly	385	390	395
Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe	405	410	415
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu	420	425	430
Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp	435	440	445
Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu	450	455	460
Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser	465	470	475
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro	485	490	495
Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Gly	500	505	510
Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Val Glu Ser	515	520	525
Gly Gly Gly Leu Val Lys Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala	530	535	540
Ala Ser Gly Phe Thr Phe Ser Asn Ala Trp Met Ser Trp Val Arg Gln	545	550	555
			560

-continued

Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Gly	Arg	Ile	Lys	Ser	Lys	Thr	565	570	575
Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	Pro	Val	Lys	Gly	Arg	Phe	Thr	580	585	590
Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	595	600	605
Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Thr	Thr	Pro	Trp	Glu	610	615	620
Trp	Ser	Trp	Tyr	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	625	630	635
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	645	650	655
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	660	665	670
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	675	680	685
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	690	695	700
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	705	710	715
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	725	730	735
Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	740	745	750
Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	755	760	765
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	770	775	780
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	785	790	795
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	805	810	815
Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	820	825	830
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	835	840	845
Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	850	855	860
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg	Asp	865	870	875
Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly	Phe	885	890	895
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	900	905	910
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	915	920	925
Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	930	935	940
Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	945	950	955

-continued

Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 965 970

<210> SEQ ID NO 73
 <211> LENGTH: 689
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 73

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50 55 60
 Pro Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
 65 70 75 80
 Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95
 Tyr Cys Thr Thr Pro Trp Glu Trp Ser Trp Tyr Asp Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130 135 140
 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 145 150 155 160
 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 165 170 175
 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
 Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp
 210 215 220
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu
 225 230 235 240
 Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys
 245 250 255
 Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr Ala Met Asn Trp Val Arg
 260 265 270
 Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ser Arg Ile Arg Ser Lys
 275 280 285
 Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe
 290 295 300
 Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn
 305 310 315 320
 Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Val Arg His Gly
 325 330 335

-continued

Asn	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly	340	345	350	
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	355	360	365	
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	370	375	380	
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	385	390	395	400
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	405	410	415	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	420	425	430	
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	435	440	445	
Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	450	455	460	
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	465	470	475	480
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	485	490	495	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	500	505	510	
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	515	520	525	
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	530	535	540	
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	545	550	555	560
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	565	570	575	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	580	585	590	
Leu	Pro	Pro	Cys	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	595	600	605	
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	610	615	620	
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	625	630	635	640
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	645	650	655	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	660	665	670	
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	675	680	685	

Lys

<210> SEQ ID NO 74

<211> LENGTH: 497

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 74

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1				5					10					15	
Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Ser	Tyr
			20					25					30		
Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
		35					40					45			
Gly	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
	50					55					60				
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
65					70					75					80
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
				85					90					95	
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
			100					105					110		
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
		115					120					125			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
	130					135					140				
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
145					150					155					160
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala
				165					170					175	
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala
			180					185					190		
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
		195					200					205			
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
	210					215					220				
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
225					230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly
				245					250					255	
Gly	Gly	Ser	Arg	Gln	Ala	Arg	Val	Val	Asn	Gly	Gly	Gly	Gly	Gly	Ser
		260						265					270		
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln
		275					280					285			
Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys
	290					295					300				
Gly	Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val
305					310					315					320
Gln	Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn
				325					330					335	
Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly
		340						345					350		
Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala
		355					360					365			
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly
	370					375					380				
Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln	Pro	Lys	Ala	Ala	Pro	Ser	Val
385					390					395					400

-continued

Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln Ala Asn Lys Ala Thr
405 410 415

Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly Ala Val Thr Val Ala
420 425 430

Trp Lys Ala Asp Ser Ser Pro Val Lys Ala Gly Val Glu Thr Thr Thr
435 440 445

Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala Ser Ser Tyr Leu Ser
450 455 460

Leu Thr Pro Glu Gln Trp Lys Ser His Arg Ser Tyr Ser Cys Gln Val
465 470 475 480

Thr His Glu Gly Ser Thr Val Glu Lys Thr Val Ala Pro Thr Glu Cys
485 490 495

Ser

<210> SEQ ID NO 75
<211> LENGTH: 497
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 75

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20 25 30

Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu
35 40 45

Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile
50 55 60

Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
65 70 75 80

Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala
85 90 95

Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp
100 105 110

Gly Gln Gly Thr Ser Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly
115 120 125

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Asp Ile
130 135 140

Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly Glu Thr
145 150 155 160

Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala
165 170 175

Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val Tyr Ala
180 185 190

Ala Thr Phe Leu Ala Asp Asp Val Pro Ser Arg Phe Ser Gly Ser Gly
195 200 205

Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser Glu Asp
210 215 220

Val Ala Arg Tyr Tyr Cys Gln His Tyr Tyr Ser Thr Pro Tyr Thr Phe
225 230 235 240

-continued

Gly Cys Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Gly Gly
 245 250 255
 Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Gly Gly Gly Ser
 260 265 270
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Ala Val Val Thr Gln
 275 280 285
 Glu Pro Ser Leu Thr Val Ser Pro Gly Gly Thr Val Thr Leu Thr Cys
 290 295 300
 Gly Ser Ser Thr Gly Ala Val Thr Thr Ser Asn Tyr Ala Asn Trp Val
 305 310 315 320
 Gln Glu Lys Pro Gly Gln Ala Phe Arg Gly Leu Ile Gly Gly Thr Asn
 325 330 335
 Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe Ser Gly Ser Leu Leu Gly
 340 345 350
 Gly Lys Ala Ala Leu Thr Leu Ser Gly Ala Gln Pro Glu Asp Glu Ala
 355 360 365
 Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn Leu Trp Val Phe Gly Gly
 370 375 380
 Gly Thr Lys Leu Thr Val Leu Gly Gln Pro Lys Ala Ala Pro Ser Val
 385 390 395 400
 Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln Ala Asn Lys Ala Thr
 405 410 415
 Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly Ala Val Thr Val Ala
 420 425 430
 Trp Lys Ala Asp Ser Ser Pro Val Lys Ala Gly Val Glu Thr Thr Thr
 435 440 445
 Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala Ser Ser Tyr Leu Ser
 450 455 460
 Leu Thr Pro Glu Gln Trp Lys Ser His Arg Ser Tyr Ser Cys Gln Val
 465 470 475 480
 Thr His Glu Gly Ser Thr Val Glu Lys Thr Val Ala Pro Thr Glu Cys
 485 490 495
 Ser

<210> SEQ ID NO 76

<211> LENGTH: 674

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 76

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Gly Tyr
 20 25 30
 Thr Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Leu Ile Thr Pro Tyr Asn Gly Ala Ser Ser Tyr Asn Gln Lys Phe
 50 55 60
 Arg Gly Lys Ala Thr Met Thr Val Asp Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys

-continued

85								90					95				
Ala	Arg	Gly	Gly	Tyr	Asp	Gly	Arg	Gly	Phe	Asp	Tyr	Trp	Gly	Gln	Gly		
		100						105					110				
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe		
		115					120					125					
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu		
	130					135					140						
Gly	Cys	Leu	Val	Glu	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp		
145					150					155					160		
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu		
				165					170					175			
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser		
			180					185					190				
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro		
		195					200					205					
Ser	Asn	Thr	Lys	Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Gly		
	210					215					220						
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln	Glu		
225					230					235					240		
Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys	Gly		
				245					250					255			
Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val	Gln		
			260					265					270				
Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn	Lys		
		275					280					285					
Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly	Gly		
	290					295					300						
Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala	Glu		
305					310					315					320		
Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly	Gly		
			325					330						335			
Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val		
			340					345					350				
Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala		
		355					360					365					
Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser		
	370					375					380						
Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val		
385					390					395					400		
Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro		
				405					410					415			
Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys		
			420					425					430				
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp		
		435					440					445					
Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly		
	450					455					460						
Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile		
465					470					475					480		
Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu		
				485				490						495			

-continued

Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 500 505 510
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 515 520 525
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 530 535 540
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Gly Ala Pro Ile Glu
 545 550 555 560
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 565 570 575
 Thr Leu Pro Pro Cys Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
 580 585 590
 Trp Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 595 600 605
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 610 615 620
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 625 630 635 640
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 645 650 655
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 660 665 670
 Gly Lys

<210> SEQ ID NO 77
 <211> LENGTH: 449
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 77

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Gly Tyr
 20 25 30
 Thr Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Leu Ile Thr Pro Tyr Asn Gly Ala Ser Ser Tyr Asn Gln Lys Phe
 50 55 60
 Arg Gly Lys Ala Thr Met Thr Val Asp Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Gly Tyr Asp Gly Arg Gly Phe Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125
 Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
 130 135 140
 Gly Cys Leu Val Glu Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160

-continued

Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
				165					170					175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
		195					200					205			
Ser	Asn	Thr	Lys	Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro
225					230					235					240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
			245						250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
			260					265					270		
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
		275					280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
305					310					315					320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys
			325						330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Cys	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Ser
		355					360					365			
Cys	Ala	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
	370					375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
385					390					395					400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Val	Ser	Lys	Leu	Thr	Val	Asp	Lys
			405						410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
		435					440					445			

Lys

<210> SEQ ID NO 78
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 78

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5					10					15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Ser	Ala	Ser	Ser	Ser	Val	Ser	Tyr	Met
		20					25					30			
His	Trp	Tyr	Gln	Gln	Lys	Ser	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr
	35					40					45				
Asp	Thr	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser

-continued

50	55	60
Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu		
65	70	75 80
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Lys His Pro Leu Thr		
	85	90 95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro		
	100	105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Arg Lys Leu Lys Ser Gly Thr		
	115	120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys		
	130	135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu		
145	150	155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser		
	165	170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala		
	180	185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe		
	195	200 205
Asn Arg Gly Glu Cys		
210		

<210> SEQ ID NO 79

<211> LENGTH: 514

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 79

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln		
1	5	10 15
Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr		
	20	25 30
Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu		
	35	40 45
Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile		
	50	55 60
Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu		
65	70	75 80
Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala		
	85	90 95
Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp		
	100	105 110
Gly Gln Gly Thr Ser Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly		
	115	120 125
Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Asp Ile		
	130	135 140
Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly Glu Thr		
145	150	155 160
Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala		
	165	170 175
Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val Tyr Ala		

-continued

180					185					190					
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
	195						200					205			
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
	210					215					220				
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
	225				230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Val	His
			245						250					255	
Met	Pro	Leu	Gly	Phe	Leu	Gly	Pro	Arg	Gln	Ala	Arg	Val	Val	Asn	Gly
		260						265					270		
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Leu	Glu
	275					280						285			
Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys
	290					295					300				
Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Thr	Tyr	Ala	Met	Asn	Trp	Val	Arg
	305				310					315					320
Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Arg	Ile	Arg	Ser	Lys
			325						330					335	
Tyr	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe
	340							345					350		
Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn
	355						360					365			
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Arg	His	Gly
	370				375						380				
Asn	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly
	385				390					395					400
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Val	Ala	Ala	Pro	Ser	Val	Phe
			405						410					415	
Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val
		420						425					430		
Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp
	435					440						445			
Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr
	450				455						460				
Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr
	465				470					475					480
Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val
			485						490					495	
Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly
		500						505					510		

Glu Cys

<210> SEQ ID NO 80

<211> LENGTH: 514

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 80

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1			5					10						15	

Ser 20	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Ser	Tyr
Gly 35	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
Gly 50	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
Ser 65	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
Gly 130	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
Gln 145	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
Val 225	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly
Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ser
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Leu	Glu
Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys
Ala 305	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Thr	Tyr	Ala	Met	Asn	Trp	Val	Arg
Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Arg	Ile	Arg	Ser	Lys
Tyr	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe
Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Arg	His	Gly
Asn 385	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Val	Ala	Ala	Pro	Ser	Val	Phe

-continued

```

<210> SEQ ID NO 81
<211> LENGTH: 232
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 81

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr
20          25          30
Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ser Arg Ile Arg Ser Lys Tyr Asn Asn Tyr Ala Thr Tyr Tyr Ala Asp
50          55          60
Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Thr
65          70          75          80
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85          90          95
Tyr Cys Val Arg His Gly Asn Phe Gly Asn Ser Tyr Val Ser Trp Phe
100         105         110
Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Val
115         120         125
Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
130         135         140
Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
145         150         155         160
Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
165         170         175
Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser
180         185         190
Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
195         200         205
Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
210         215         220
Lys Ser Phe Asn Arg Gly Glu Cys
225         230

```


-continued

<210> SEQ ID NO 82
<211> LENGTH: 956
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 82

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20 25 30

Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu
35 40 45

Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile
50 55 60

Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
65 70 75 80

Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala
85 90 95

Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp
100 105 110

Gly Gln Gly Thr Ser Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly
115 120 125

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile
130 135 140

Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly Glu Thr
145 150 155 160

Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala
165 170 175

Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val Tyr Ala
180 185 190

Ala Thr Phe Leu Ala Asp Asp Val Pro Ser Arg Phe Ser Gly Ser Gly
195 200 205

Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser Glu Asp
210 215 220

Val Ala Arg Tyr Tyr Cys Gln His Tyr Tyr Ser Thr Pro Tyr Thr Phe
225 230 235 240

Gly Cys Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Val His
245 250 255

Met Pro Leu Gly Phe Leu Gly Pro Arg Gln Ala Arg Val Val Asn Gly
260 265 270

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Ala Val Val Thr Gln
275 280 285

Glu Pro Ser Leu Thr Val Ser Pro Gly Gly Thr Val Thr Leu Thr Cys
290 295 300

Gly Ser Ser Thr Gly Ala Val Thr Thr Ser Asn Tyr Ala Asn Trp Val
305 310 315 320

Gln Glu Lys Pro Gly Gln Ala Phe Arg Gly Leu Ile Gly Gly Thr Asn
325 330 335

Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe Ser Gly Ser Leu Leu Gly
340 345 350

Gly Lys Ala Ala Leu Thr Leu Ser Gly Ala Gln Pro Glu Asp Glu Ala

-continued

355					360					365					
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly
370						375					380				
Gly	Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser
385					390					395					400
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala
				405					410					415	
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val
			420					425					430		
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala
		435					440					445			
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val
450						455					460				
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His
465					470					475					480
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys
				485					490					495	
Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Val	Gln	Leu	Val
			500					505					510		
Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser
		515					520					525			
Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Gly	Tyr	Thr	Met	Asn	Trp	Val
530						535					540				
Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	Gly	Leu	Ile	Thr	Pro
545					550					555					560
Tyr	Asn	Gly	Ala	Ser	Ser	Tyr	Asn	Gln	Lys	Phe	Arg	Gly	Lys	Ala	Thr
				565					570					575	
Met	Thr	Val	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr	Met	Glu	Leu	Ser	Ser
			580					585					590		
Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Gly	Gly	Tyr
		595					600					605			
Asp	Gly	Arg	Gly	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val
610					615						620				
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser
625					630					635					640
Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Glu
				645					650					655	
Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu
			660					665					670		
Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu
		675					680					685			
Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr
690						695					700				
Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val
705					710					715					720
Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro
				725					730					735	
Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe
			740					745					750		
Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val
		755					760					765			

-continued

Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe
770						775					780				
Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro
785					790					795					800
Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr
				805					810					815	
Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val
			820					825					830		
Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala
		835					840					845			
Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg
	850					855					860				
Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly
865					870					875					880
Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro
				885					890					895	
Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser
			900					905					910		
Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln
		915					920					925			
Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His
	930					935					940				
Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys				
945					950					955					

<210> SEQ ID NO 83
 <211> LENGTH: 956
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 83

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1			5					10						15	
Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Ser	Tyr
	20					25						30			
Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
	35					40						45			
Gly	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
	50				55					60					
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
65				70					75					80	
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
		85						90						95	
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
	100							105					110		
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
	115					120						125			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
	130					135						140			
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
145					150					155					160

Val	Thr	Ile	Thr	Cys 165	Arg	Ala	Ser	Glu	Asn 170	Ile	Asp	Ser	Tyr	Leu 175	Ala	
Trp	Tyr	Gln	Gln	Lys 180	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala	
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp	
Val 225	Ala	Arg	Tyr	Tyr	Cys 230	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe	240
Gly	Cys	Gly	Thr	Lys 245	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly	
Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ser	
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln	
Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys	
Gly 305	Ser	Ser	Thr	Gly	Ala 310	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val	320
Gln	Glu	Lys	Pro	Gly 325	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn	
Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly	
Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala	
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly	
Gly 385	Thr	Lys	Leu	Thr	Val 390	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	400
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	
Val 450	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	
Pro 465	Ser	Ser	Ser	Leu	Gly 470	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	480
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	
Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Val	Gln	Leu	Val	
Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser	
Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Gly	Tyr	Thr	Met	Asn	Trp	Val	
Arg 545	Gln	Ala	Pro	Gly	Gln 550	Gly	Leu	Glu	Trp	Met	Gly	Leu	Ile	Thr	Pro	560

-continued

Tyr	Asn	Gly	Ala	Ser	Ser	Tyr	Asn	Gln	Lys	Phe	Arg	Gly	Lys	Ala	Thr	565	570	575
Met	Thr	Val	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr	Met	Glu	Leu	Ser	Ser	580	585	590
Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Gly	Gly	Tyr	595	600	605
Asp	Gly	Arg	Gly	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	610	615	620
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	625	630	635
Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Glu	645	650	655
Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	660	665	670
Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	675	680	685
Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	690	695	700
Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	705	710	715
Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	725	730	735
Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	740	745	750
Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	755	760	765
Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	770	775	780
Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	785	790	795
Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	805	810	815
Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	820	825	830
Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	835	840	845
Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg	850	855	860
Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly	865	870	875
Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	885	890	895
Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	900	905	910
Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	915	920	925
Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	930	935	940
Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys					945	950	955

-continued

```

<210> SEQ ID NO 84
<211> LENGTH: 674
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 84

Gln Ala Val Val Thr Gln Glu Pro Ser Leu Thr Val Ser Pro Gly Gly
1          5          10          15

Thr Val Thr Leu Thr Cys Gly Ser Ser Thr Gly Ala Val Thr Thr Ser
20          25          30

Asn Tyr Ala Asn Trp Val Gln Glu Lys Pro Gly Gln Ala Phe Arg Gly
35          40          45

Leu Ile Gly Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe
50          55          60

Ser Gly Ser Leu Leu Gly Gly Lys Ala Ala Leu Thr Leu Ser Gly Ala
65          70          75          80

Gln Pro Glu Asp Glu Ala Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn
85          90          95

Leu Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Ser Ser Ala
100         105         110

Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser
115         120         125

Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe
130         135         140

Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly
145         150         155         160

Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu
165         170         175

Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr
180         185         190

Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys
195         200         205

Val Glu Pro Lys Ser Cys Asp Gly Gly Gly Gly Ser Gly Gly Gly Gly
210         215         220

Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly
225         230         235         240

Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Gly
245         250         255

Tyr Thr Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp
260         265         270

Met Gly Leu Ile Thr Pro Tyr Asn Gly Ala Ser Ser Tyr Asn Gln Lys
275         280         285

Phe Arg Gly Lys Ala Thr Met Thr Val Asp Thr Ser Thr Ser Thr Val
290         295         300

Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr
305         310         315         320

Cys Ala Arg Gly Gly Tyr Asp Gly Arg Gly Phe Asp Tyr Trp Gly Gln
325         330         335

Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
340         345         350

Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala

```

-continued

355	360	365
Leu Gly Cys Leu Val Glu Asp Tyr Phe Pro Glu Pro Val Thr Val Ser		
370	375	380
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val		
385	390	395
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro		
	405	410
		415
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys		
	420	425
		430
Pro Ser Asn Thr Lys Val Asp Glu Lys Val Glu Pro Lys Ser Cys Asp		
	435	440
		445
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly		
	450	455
		460
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile		
	465	470
		475
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu		
	485	490
		495
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His		
	500	505
		510
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg		
	515	520
		525
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys		
	530	535
		540
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Gly Ala Pro Ile Glu		
	545	550
		555
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr		
	565	570
		575
Thr Leu Pro Pro Cys Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu		
	580	585
		590
Trp Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp		
	595	600
		605
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val		
	610	615
		620
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp		
	625	630
		635
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His		
	645	650
		655
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro		
	660	665
		670

Gly Lys

<210> SEQ ID NO 85

<211> LENGTH: 971

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 85

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
 1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30

-continued

Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
	35						40					45			
Gly	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
	50					55					60				
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
65					70					75					80
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
				85					90					95	
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
			100					105					110		
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
		115					120					125			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
	130					135					140				
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
145					150					155					160
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala
				165					170					175	
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala
			180					185					190		
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
		195					200					205			
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
	210					215					220				
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
225					230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly
				245					250					255	
Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Phe	Val	Gly	Gly	Thr	Gly	Gly	Gly
			260					265					270		
Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Ser	Glu	Val	Gln	Leu	Leu	Glu
		275					280					285			
Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys
	290					295					300				
Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Thr	Tyr	Ala	Met	Asn	Trp	Val	Arg
305					310					315					320
Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Arg	Ile	Arg	Ser	Lys
				325					330					335	
Tyr	Asn	Asn	Tyr	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe
			340					345					350		
Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn
		355					360					365			
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Arg	His	Gly
	370					375					380				
Asn	Phe	Gly	Asn	Ser	Tyr	Val	Ser	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly
385					390					395					400
Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
				405					410					415	
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
			420					425					430		

-continued

Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
	435						440					445			
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
450						455					460				
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
465					470					475					480
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
				485					490					495	
Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Gly
			500					505					510		
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Val	Glu	Ser
		515					520					525			
Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala
	530					535					540				
Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Ala	Trp	Met	Ser	Trp	Val	Arg	Gln
545					550					555					560
Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Gly	Arg	Ile	Lys	Ser	Lys	Thr
				565					570					575	
Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	Pro	Val	Lys	Gly	Arg	Phe	Thr
			580					585					590		
Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser
		595					600					605			
Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Thr	Thr	Pro	Trp	Glu
	610					615					620				
Trp	Ser	Trp	Tyr	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser
625					630						635				640
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser
				645					650					655	
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp
			660					665					670		
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr
		675					680					685			
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr
	690					695					700				
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln
705					710					715					720
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp
				725					730					735	
Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro
			740					745					750		
Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
		755					760					765			
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
	770					775					780				
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn
785					790					795					800
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
				805					810					815	
Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
			820					825					830		
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser

-continued

835					840					845					
Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys
850						855					860				
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg	Asp
865						870					875				880
Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly	Phe
				885					890					895	
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu
			900					905					910		
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe
		915					920					925			
Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly
	930					935					940				
Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr
945						950					955				960
Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys					
				965					970						

<210> SEQ ID NO 86
 <211> LENGTH: 33
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 86

Gly	Gly	Gly	Gly	Ser	Val	His	Met	Pro	Leu	Gly	Phe	Leu	Gly	Pro	Arg
1				5					10					15	

Gln	Ala	Arg	Val	Val	Asn	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly
			20				25						30		

Ser

<210> SEQ ID NO 87
 <211> LENGTH: 35
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 87

Gly	Gly	Gly	Gly	Ser	Val	His	Met	Pro	Leu	Gly	Phe	Leu	Gly	Pro	Arg
1				5					10					15	

Gln	Ala	Arg	Val	Val	Asn	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly
			20				25						30		

Ser	Gly	Gly
		35

<210> SEQ ID NO 88
 <211> LENGTH: 33
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 88

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Phe
1				5					10					15	

-continued

Val	Gly	Gly	Thr	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly
			20					25					30		

Ser

<210> SEQ ID NO 89
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 89

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Lys	Lys	Ala	Ala	Pro	Val
1				5					10					15	

Asn	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly
			20					25					30		

Ser

<210> SEQ ID NO 90
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 90

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Pro	Met	Ala	Lys	Lys	Val
1				5					10					15	

Asn	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly
			20					25					30		

Ser

<210> SEQ ID NO 91
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 91

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Arg	Ala	Lys	Val
1				5					10					15	

Asn	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly
			20					25					30		

Ser

<210> SEQ ID NO 92
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 92

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Val	His	Met	Pro	Leu	Gly
1				5					10					15	

Phe	Leu	Gly	Pro	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly
			20					25					30		

Ser

-continued

<210> SEQ ID NO 93
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 93

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Ala Arg Ala Lys Gly
1 5 10 15
Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
20 25 30

Ser

<210> SEQ ID NO 94
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 94

Gly Gly Gly Gly Ser Val His Met Pro Leu Gly Phe Leu Gly Pro Pro
1 5 10 15
Met Ala Lys Lys Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
20 25 30

Ser

<210> SEQ ID NO 95
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 95

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Lys Lys Ala Ala Pro Gly
1 5 10 15
Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
20 25 30

Ser

<210> SEQ ID NO 96
<211> LENGTH: 33
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 96

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Pro Met Ala Lys Lys Gly
1 5 10 15
Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
20 25 30

Ser

<210> SEQ ID NO 97
<211> LENGTH: 18

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 97

Val His Met Pro Leu Gly Phe Leu Gly Pro Arg Gln Ala Arg Val Val
1 5 10 15

Asn Gly

<210> SEQ ID NO 98
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 98

Phe Val Gly Gly Thr Gly
1 5

<210> SEQ ID NO 99
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 99

Lys Lys Ala Ala Pro Val Asn Gly
1 5

<210> SEQ ID NO 100
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 100

Pro Met Ala Lys Lys Val Asn Gly
1 5

<210> SEQ ID NO 101
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 101

Gln Ala Arg Ala Lys Val Asn Gly
1 5

<210> SEQ ID NO 102
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 102

Val His Met Pro Leu Gly Phe Leu Gly Pro
1 5 10

-continued

<210> SEQ ID NO 103
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 103

Gln Ala Arg Ala Lys
1 5

<210> SEQ ID NO 104
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 104

Val His Met Pro Leu Gly Phe Leu Gly Pro Pro Met Ala Lys Lys
1 5 10 15

<210> SEQ ID NO 105
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 105

Lys Lys Ala Ala Pro
1 5

<210> SEQ ID NO 106
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 106

Pro Met Ala Lys Lys
1 5

<210> SEQ ID NO 107
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 107

Gly Tyr Thr Met Asn
1 5

<210> SEQ ID NO 108
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 108

-continued

Leu	Ile	Thr	Pro	Tyr	Asn	Gly	Ala	Ser	Ser	Tyr	Asn	Gln	Lys	Phe	Arg
1				5					10					15	

Gly

<210> SEQ ID NO 109
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 109

Gly	Gly	Tyr	Asp	Gly	Arg	Gly	Phe	Asp	Tyr
1				5				10	

<210> SEQ ID NO 110
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 110

Ser	Ala	Ser	Ser	Ser	Val	Ser	Tyr	Met	His
1				5				10	

<210> SEQ ID NO 111
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 111

Asp	Thr	Ser	Lys	Leu	Ala	Ser
1				5		

<210> SEQ ID NO 112
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 112

Gln	Gln	Trp	Ser	Lys	His	Pro	Leu	Thr
1				5				

<210> SEQ ID NO 113
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 113

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
1				5					10					15	

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Gly	Tyr
				20				25					30		

Thr	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
		35					40					45			

-continued

Gly Leu Ile Thr Pro Tyr Asn Gly Ala Ser Ser Tyr Asn Gln Lys Phe
50 55 60

Arg Gly Lys Ala Thr Met Thr Val Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Gly Tyr Asp Gly Arg Gly Phe Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ser
115

<210> SEQ ID NO 114
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 114

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Ser Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Lys His Pro Leu Thr
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> SEQ ID NO 115
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 115

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Tyr
20 25 30

Lys Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Tyr Phe Asn Pro Asn Ser Gly Tyr Ser Thr Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Leu Ser Pro Gly Gly Tyr Tyr Val Met Asp Ala Trp Gly Gln
100 105 110

-continued

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 116
<211> LENGTH: 106
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 116

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Asn Asn Tyr
20 25 30
Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
35 40 45
Tyr Asn Thr Asn Asn Leu Gln Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80
Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Phe Pro Thr
85 90 95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> SEQ ID NO 117
<211> LENGTH: 648
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 117

caggccgctcg tgacccagga acccagcctg acagtgtctc ctggcggcac cgtgaccctg 60
acatgtggca gttctacagg cgccgtgacc accagcaact acgccaactg ggtgcaggaa 120
aagcccgccc aggccttcag aggactgac ggccggcacca acaagagagc ccctggcacc 180
cctgccagat tcagcggatc tctgctggga ggaaaggccg ccctgacact gtctggcgcc 240
cagccagaag atgaggccga gtactactgc gccctgtggt acagcaacct gtgggtgttc 300
ggcggaggca ccaagctgac agtcctaggt caacccaagg ctgccccag cgtgaccctg 360
ttcccccca gcagcgagga actgcaggcc aacaaggcca ccctggtctg cctgacagc 420
gacttctacc caggcgccgt gaccgtggcc tggaaggccg acagcagccc cgtgaaggcc 480
ggcgtggaga ccaccacccc cagcaagcag agcaacaaca agtacgccgc cagcagctac 540
ctgagcctga ccccccagca gtggaagagc cacaggtcct acagctgcca ggtgaccac 600
gagggcagca ccgtggagaa aaccgtggcc cccaccgagt gcagctga 648

<210> SEQ ID NO 118
<211> LENGTH: 2916
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 118

-continued

caagtgcagc	tgaagagtc	cggccctgga	ctggtggccc	ctagccagag	cctgagcacc	60
acctgtaccg	tgtccggett	cagcctgacc	agctacggcg	tgtcatgggt	gcgccagcct	120
ccaggcaagt	gtctggaatg	gctgggcacc	atctggggcg	acggcagcac	caattaccac	180
agcgcctga	tcagcagact	gagcatctcc	aaggacaaca	gcaagagcca	ggtgttctctg	240
aagctgaaca	gcctgcagac	cgacgacacc	gccacctact	actgcgcca	gggcaccacc	300
accgtggtgg	acgactacta	cgctatggac	tactggggcc	agggcaccag	cgtgacagtg	360
tctagcggag	gcggaggatc	tggcggcgga	ggaagtggcg	gagggggatc	tgggggaggc	420
ggaagcgata	tccagatgac	ccagagccct	gccagcctgt	ctgcctctgt	gggcgagaca	480
gtgaccatca	catgccgggc	cagcgagAAC	atcgacagct	acctggcctg	gtatcagcag	540
aagcagggca	agagccccc	gctgctgggt	tacgcccga	cctttctggc	cgacgatgtg	600
cccagcagat	tcagcggcag	cggaagcggc	acacagtaca	gcctgaagat	caactccctg	660
cagagcgagg	acgtggcccg	gtactactgc	cagcactact	acagcacc	ctacaccttc	720
ggctcggca	ccaagctgga	aatcaaagga	ggcggcgga	gtgtgcacat	gcccctgggc	780
ttcctgggcc	ccagacaggc	cagagtcgtg	aacggggggg	gcggaggcag	tgggggggga	840
ggatccgagg	tgcagctgct	ggaatctggc	ggcggactgg	tgcagcctgg	cggatctctg	900
agactgagct	gtgccgccag	cggcttcacc	ttcagcacct	acgccatgaa	ctgggtgcgc	960
caggcccctg	gcaaaggcct	ggaatgggtg	tcccggatca	gaagcaagta	caacaactac	1020
gccacctact	acgccgacag	cgtgaagggc	cggttcacca	tcagccggga	cgacagcaag	1080
aacaccctgt	acctgcagat	gaacagcctg	cgggcggagg	acaccgccgt	gtactattgt	1140
gtgcggcacg	gcaacttcgg	caacagctat	gtgtcttggg	ttgcctactg	gggcaggggc	1200
accctcgtga	cctgtgaag	cgctagcaca	aagggcccta	gcgtgttccc	tctggccccc	1260
agcagcaaga	gcacaagcgg	cgaacagccc	gccctgggct	gcctcgtgaa	ggactacttc	1320
cccagcccg	tgacagtgtc	ttggaacagc	ggagccctga	caagcggcgt	gcacaccttc	1380
cctgccgtgc	tgcagagcag	cggcctgtac	tccctgagca	gcgtggtcac	cgtgcctagc	1440
agcagcctgg	gcacccagac	ctacatctgc	aacgtgaacc	acaagcccag	caacacccaa	1500
gtggacaaga	aggtggagcc	caagagctgt	gatggcggag	gagggtcggg	gggcggagga	1560
tccgaggtgc	aattgggtga	atctggtggg	ggtctggtaa	aaccgggcgg	ttccctgcgt	1620
ctgagctcgg	cggcttcggg	gttcaccttc	tccaacgcgt	ggatgagctg	ggttcgccag	1680
gccccgggca	aaggcctcga	gtgggttggg	cgtatcaagt	ctaaaactga	cgggtggcacc	1740
acggattacg	cggctccagt	taaaggtcgt	tttaccattt	cccgcgacga	tagcaaaaac	1800
actctgtatc	tgcagatgaa	ctctctgaaa	actgaagaca	ccgcagtcta	ctactgtact	1860
accccggtgg	aatggtcttg	gtacgattat	tggggccagg	gcacgctggg	tacggtgtct	1920
agcgtagta	ccaagggccc	cagcgtgttc	cccctggcac	ccagcagcaa	gagcacatct	1980
ggcggaaacg	cgcctctggg	ctgtctgggt	aaagactact	tccccgagcc	cgtgaccgtg	2040
tcttggaact	ctggcgccct	gaccagcggc	gtgcacacct	ttccagccgt	gctgcagagc	2100
agcggcctgt	actccctgtc	ctccgtgggc	accgtgccct	ctagctccct	gggaacacag	2160
acatatatct	gtaatgtcaa	tcacaagcct	tccaacacca	aagtcgataa	gaaagtcgag	2220
cccaagagct	gcgacaaaac	tcacacatgc	ccaccgtgcc	cagcacctga	agctgcaggg	2280

-continued

```

ggaccgtcag tcttctcttt cccccaaaa cccaaggaca ccctcatgat ctcccggacc 2340
cctgaggtea catgcgtggt ggtggacgtg agccacgaag accctgaggt caagttcaac 2400
tggtacgtgg acggcgtgga ggtgcataat gccaaagaca agccgcggga ggagcagtac 2460
aacagcacgt accgtgtggt cagcgtcttc accgtctgc accaggactg gctgaatggc 2520
aaggagtaca agtgcaaggt ctccaacaaa gccctcggcg ccccatcga gaaaaccatc 2580
tccaaagcca aagggcagcc ccgagaacca caggtgtaca ccctgcccc atgccgggat 2640
gagctgacca agaaccaggt cagcctgtgg tgcttggtca aaggcttcta tcccagcgac 2700
atcgccgtgg agtgggagag caatgggcag ccggagaaca actacaagac cagcctccc 2760
gtgctggact ccgacggctc cttcttcttc tacagcaagc tcaccgtgga caagagcagg 2820
tggcagcagg ggaacgtctt ctcatgctcc gtgatgcatg aggctctgca caaccactac 2880
acgcagaaga gcctctccct gtctccgggt aatatga 2916

```

<210> SEQ ID NO 119

<211> LENGTH: 2070

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 119

```

gagggtgcaat tgggtgaatc tgggtggtgt ctggtaaaac cgggcgggtc cctgcgtctg 60
agctgcgcgg cttccggatt caccttctcc aacgcgtgga tgagctgggt tcgccaggcc 120
ccgggcгааг gcctcgagtг ggttggtcgt atcaagteta aaactgacgg tggcaccacg 180
gattacgcgg ctccagttaa aggtcgtttt accatttccc gcgacgatag caaaaacact 240
ctgtatctgc agatgaactc tctgaaaact gaagacaccg cagtctacta ctgtactacc 300
ccgtgggaat ggtcttggtg cgattattgg ggcagggca cgctggttac ggtgtcttcc 360
gctagcacia agggccctag cgtgttcctt ctggcccccа gcagcaagag cacaagcggc 420
ggaacagccg ccctgggctg cctcgtgaag gactacttcc ccgagcccgт gacagtgtct 480
tggaacagcg gagccctgac aagcggcgtg cacactttcc ctgccgtgct gcagagcagc 540
ggcctgtact ccctgagcag cgtggtcacc gtgcctagca gcagcctggg caccagacc 600
tacatctgca acgtgaacca caagcccagc aacaccaaag tggacaagaa ggtggagccc 660
aagagctgtg atggcggagg aggggtccgga ggcggaggat ccgaggtgca gctgctggaa 720
tctggcggcg gactggtgca gcctggcgga tctctgagac tgagctgtgc cgccagcggc 780
ttcaccttea gcacctacgc catgaactgg gtgcgccagg cccctggcaa aggcctggaa 840
tgggtgtccc ggatcagaag caagtacaac aactacgcca cctactacgc cgacagcgtg 900
aagggccggt tcaccatcag ccgggacgac agcaagaaca ccctgtacct gcagatgaac 960
agcctgcggg ccgaggacac cgccgtgtac tattgtgtgc ggcacggcaa cttcgcaac 1020
agctatgtgt cttggtttgc ctactggggc cagggcaccс tcgtgaccgt gtcaagcgct 1080
agtaccaagg gccccagcgt gttccccctg gcaccacgca gcaagagcac atctggcgga 1140
acagcgcctc tgggtgtctt ggtgaaagac tacttccccг agcccgtagc cgtgtcttgg 1200
aactctggcg ccctgaccag cggcgtgcac acctttccag ccgtgctgca gagcagcggc 1260
ctgtactccc tgtctccgtg ggtcaccgтг ccctctagct ccctgggaac acagacatat 1320

```

-continued

atctgtaatg tcaatcacia gccttccaac accaaagtgc ataagaaagt cgagcccaag	1380
agctgcgaca aaactcacac atgccaccg tgcccagcac ctgaagctgc agggggaccg	1440
tcagtcttcc tcttcccccc aaaaccaag gacaccctca tgatctcccc gaccctgag	1500
gtcacatgcg tgggtggtga cgtgagccac gaagaccctg aggtcaagtt caactggtac	1560
gtggacggcg tggaggtgca taatgccaag acaaagccgc gggaggagca gtacaacagc	1620
acgtaccgtg tggtcagcgt cctcaccgtc ctgcaccagg actggctgaa tggcaaggag	1680
tacaagtgca aggtctccaa caaagccctc ggcgccccca tcgagaaaac catctccaaa	1740
gccaaggggc agccccgaga accacaggtg tacaccctgc ccccatgccg ggatgagctg	1800
accaagaacc aggtcagcct gtggtgcctg gtcaaaggct tctatcccag cgacatcgcc	1860
gtggagtggg agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgctgtg	1920
gactccgacg gctccttctt cctctacagc aagctcaccg tggacaagag caggtggcag	1980
caggggaacg tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgcag	2040
aagagcctct cctgtctctc gggtaaatga	2070

<210> SEQ ID NO 120

<211> LENGTH: 1494

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 120

caagtgcagc tgaagagtc cggccctgga ctggtggccc ctagccagag cctgagcacc	60
acctgtaccg tgtccggett cagcctgacc agctacggcg tgtcatgggt gcgccagcct	120
ccaggcaagt gtctggaatg gctgggcacc atctggggcg acggcagcac caattaccac	180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctctg	240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcca gggcaccacc	300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg	360
tctagcggag gcggaggatc tggcggcgga ggaagtggcg gagggggatc tgggggaggc	420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca	480
gtgaccatca catgccgggc cagcgagAAC atcgacagct acctggcctg gtatcagcag	540
aagcagggca agagccccca gctgctggtg tacgcgcca cctttctggc cgacgatgtg	600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactccctg	660
cagagcgagg acgtggcccg gtactactgc cagcactact acagcaccac ctacaccttc	720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtaga	780
caggccagag tcgtgaacgg gggagggggg ggaagtgggg gcggaggcag tgggggcgga	840
ggatcccagg ccgtcgtgac ccaggaaacc agcctgacag tgtctcctgg cggcaccgtg	900
accctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg	960
cagggaaaag ccggccaggc cttcagagga ctgatcgcg gcaccaaaa gagagcccct	1020
ggcaccctcg ccagattcag cggatctctg ctgggaggaa aggcgcctct gacactgtct	1080
ggcgcccagc cagaagatga ggccgagtag tactgcgccc tgtggtacag caacctgtgg	1140
gtgttcggcg gaggcaccaa gctgacagtc ctaggtcaac ccaaggctgc cccagcgtg	1200

-continued

```

accctgttcc cccccagcag cgaggaactg caggccaaca aggccaccct ggtctgctg 1260
atcagcgact tctaccagcag cgccgtgacc gtggcctgga aggccagacag cagccccgtg 1320
aaggccggcg tggagaccac cccccagc aagcagagca acaacaagta cgccgccagc 1380
agctacctga gcctgacccc cgagcagtgg aagagccaca ggtcctacag ctgccaggtg 1440
accacgagg gcagcacctg ggagaaaacc gtggccccc cagagtgcag ctga 1494

```

```

<210> SEQ ID NO 121
<211> LENGTH: 1494
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 121

```

```

caagtgcagc tgaagagtc cgccctgga ctggtggccc ctaccagag cctgagcatc 60
acctgtaccg tgtccggtt cagcctgacc agctacggcg tgtcatgggt gcgccagcct 120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac 180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctg 240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcaa ggcatcacc 300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg 360
tctagcggag gcggaggatc tggcgcgga ggaagtggcg gagggggatc tgggggaggc 420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca 480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag 540
aagcagggca agagccccc gctgctggtg tacgcgcca cctttctggc cgacgatgtg 600
cccagcagat tcagcggcag cgaagcggc acacagtaca gcctgaagat caactccctg 660
cagagcgagg acgtggccc gtactactgc cagcactact acagcaccac ctacaccttc 720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccgagggcgg cggaagtgga 780
ggcgccggaa gtggcgagg cgagggggg ggaagtggg gcggaggcag tgggggggga 840
ggatcccagg ccgtcgtgac ccaggaaccc agcctgacag tgtctcctgg cggcaccgtg 900
accctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg 960
caggaaaagc ccggccaggc cttcagagga ctgatcgcg gcaccaaaa gagagcccct 1020
ggcacccttg ccagattcag cggatctctg ctgggaggaa aggccgccct gacactgtct 1080
ggcgcccagc cagaagatga ggccgagtag tactgcgccc tgtggtacag caacctgtgg 1140
gtgttcggcg gaggcaccaa gctgacagtc ctaggccaac ccaaggctgc ccccagcgtg 1200
accctgttcc cccccagcag cgaggaactg caggccaaca aggccaccct ggtctgctg 1260
atcagcgact tctaccagcag cgccgtgacc gtggcctgga aggccagacag cagccccgtg 1320
aaggccggcg tggagaccac cccccagc aagcagagca acaacaagta cgccgccagc 1380
agctacctga gcctgacccc cgagcagtgg aagagccaca ggtcctacag ctgccaggtg 1440
accacgagg gcagcacctg ggagaaaacc gtggccccc cagagtgcag ctga 1494

```

```

<210> SEQ ID NO 122
<211> LENGTH: 2025
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 122

caggtgcagc tgggtgcagtc tggcgccgaa gtgaagaaac caggcgccag cgtgaaggtg	60
tcttgcaagg ccagcggcta cagcttcacc ggctacacca tgaactgggt gcgccaggct	120
cctggacagg gcctggaatg gatgggectg atcacccctt acaacggcgc cagcagctac	180
aaccagaagt tccggggcaa ggccaccatg accgtggaca ccagcacctc caccgtgtat	240
atggaactga gcagcctgcg gagcgaggac accgcccgtg actattgtgc cagaggcggc	300
tacgacggca gaggtctoga ttattggggc cagggcacc cctgtaccgt gtcctctgct	360
agcaccaagg gccctccgtg gtttccctctg gcccttcca gcaagtccac ctctggcgga	420
actgcccgtc tgggtgcctt ggtggaagat tacttcccc agcccgtgac cgtgtcctgg	480
aattctggcg ctctgacctc cggcgtgcac acctttccag ctgtgctgca gtcctccggc	540
ctgtactccc tgtctccgtg cgtgacagtg cctccagct ctctgggcac ccagacctac	600
atctgcaacg tgaaccacaa gccctccaac accaaggtgg acgagaaggt ggaacccaag	660
tctgcgacg gtggcggagg ttccggaggc ggaggatccc aggtgtcgtg gaccaggaa	720
ccctccctga cagtgtctcc tggcggcacc gtgacctga cctgtggatc ttctaccggc	780
gctgtgacca cctccaacta cgccaattgg gtgcaggaaa agcccggcca ggccttcaga	840
ggactgatcg gcgcaccaa caagagagcc cctggcacc ctgccagatt ctccggttct	900
ctgctgggcg gcaaggctgc cctgactctg tctggtgctc agcctgagga cgaggccgag	960
tactactgcg cctgtgggta ctccaacctg tgggtgttcg gcggaggcac caagctgacc	1020
gtgctgtcca gcgcttccac caagggacc agtgtgttcc cctggcccc cagctccaag	1080
tctacatccg gtggcacagc tgccctggga tgtctcgtga aggactactt tctgagcct	1140
gtgacagtgt cttggaacag cggagccctg accagcggag tgcacacatt cctgacgtg	1200
ctgcagagca gcggcctgta tagcctgagc agcgtcgtga ccgtgccttc ctctagcctg	1260
ggaacacaga catatatctg taatgtgaat cataagccca gtaataccaa agtggataag	1320
aaagtggaa ctaagagctg cgataagacc cacacctgtc cccctgccc tgetcctgaa	1380
gctgctgggt gccctagcgt gttcctgttc cccccaagc ccaaggacac cctgatgatc	1440
tcccggaacc ccgaagtgc ctgcgtgggt gtggatgtgt ccacagagga cctgaagtg	1500
aagttcaatt ggtacgtgga cggcgtggaa gtgcacaacg ccaagaccaa gcctagagag	1560
gaacagtaca actccaccta ccgggtgggt tccgtgctga cagtgtgca ccaggactgg	1620
ctgaacggca aagagtacaa gtgcaagggt tccaacaagg ccctgggcgc tccatcgaa	1680
aagaccatct ccaaggccaa gggccagccc cgggaacccc aggtgtacac cctgccccca	1740
tgcgggatg agctgaccaa gaaccaggtc agcctgtggt gcctggtcaa aggtctctat	1800
cccagcgaca tcgccgtgga gtgggagagc aatgggcagc cggagaacaa ctacaagacc	1860
acgcctcccg tctgtgactc cgacggctcc ttcttccct acagcaagct caccgtggac	1920
aagagcaggt ggcagcaggg gaacgtcttc tcatgctccg tgatgcatga ggctctgcac	1980
aaccactaca cgcagaagag cctctccctg tctccgggta aatga	2025

<210> SEQ ID NO 123

<211> LENGTH: 1350

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 123
caggtgcagc tggcgcagtc tggcgccgaa gtgaagaaac caggcgccag cgtgaaggctg      60
tcctgcaagg ccagcggeta cagcttcacc ggctacacca tgaactgggt gcgccaggct      120
cctggacagg gcctggaatg gatgggcctg atcacccctt acaacggcgc cagcagctac      180
aaccagaagt tccggggcaa ggccaccatg accgtggaca ccagcacctc caccgtgtat      240
atggaactga gcagcctgcg gagcgaggac accgcccgtg actattgtgc cagaggcggc      300
tacgacggca gaggtctoga ttattggggc cagggcaccc tcgtgaccgt gtcctctgct      360
agcaccaagg gccctccgtt gttccccctg gccccacga gcaagagcac cagcggcggc      420
acagccgctc tgggtctgct ggtcagggac tacttccccg agcccgtagc cgtgtcctgg      480
aacagcggag ccctgacctc cggcgtgcac accttccccg ccgtgctgca gagttctggc      540
ctgtatagcc tgagcagcgt ggtcaccgtg ccttctagca gcctgggcac ccagacctac      600
atctgcaacg tgaaccacaa gccacgaac accaagggtg acgagaaggt ggagcccaag      660
agctgcgaca aaactcacac atgcccaccg tgcccagcac ctgaagctgc agggggaccg      720
tcagtcttcc tcttcccccc aaaacccaag gacacctca tgatctcccg gacctctgag      780
gtcacatgcg tgggtggtga cgtgagccac gaagaccctg aggtcaagtt caactggtac      840
gtggacggcg tggaggtgca taatgccaaag acaaagccgc gggaggagca gtacaacagc      900
acgtaccgtg tggtcagcgt cctcaccgtc ctgcaccagg actggctgaa tggcaaggag      960
tacaagtgca aggtctccaa caaagccctc ggcgccccca tcgagaaaac catctccaaa    1020
gccaaagggc agccccgaga accacagggtg tgcaccctgc ccccatcccc ggatgagctg    1080
accaagaacc aggtcagcct ctcgtgcgca gtcaaaggct tctatcccag cgacatcgcc    1140
gtggagtggg agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgctgctg    1200
gactccgacg gctcctctct cctcgtgagc aagctcaccg tggacaagag cagggtggcag    1260
caggggaaacg tcttctcatg ctcctgtgat catgaggctc tgcacaacca ctacacgcag    1320
aagagcctct ccctgtctcc gggtaaatga                                     1350

```

```

<210> SEQ ID NO 124
<211> LENGTH: 642
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 124
gacatccaga tgaccagag cccacagcag ctgtctgcca gcgtgggcca cagagtgacc      60
atcacctgta gcgccagcag cagcgtgtcc tacatgcaat ggtatcagca gaagtccggc      120
aaggccccc aagctgctgat ctacgacacc agcaagctgg cctccggcgt gccacgcaga      180
ttttctggca gcggctccgg caccgacttc accctgacaa tcagctccct ccagcccagag      240
gacttcgcca cctactactg ccagcagtgg tccaagcacc ccctgacctt tggccagggc      300
accaagctgg aaatcaagcg tacggtggct gcaccatctg tcttcatctt cccgccatct      360
gatcggaagt tgaatctgga aactgcctct gttgtgtgcc tgctgaataa cttctatccc      420

```

-continued

```

agagaggcca aagtacagtg gaagggtgat aacgcctccc aatcgggtaa ctcccaggag 480
agtgtcacag agcaggacag caaggacagc acctacagcc tcagcagcac cctgacgctg 540
agcaaagcag actacgagaa acacaaagtc tacgcctgcg aagtcaccca tcagggcctg 600
agctcgcccg tcacaaagag cttcaacagg ggagagtgtt ag 642

```

```

<210> SEQ ID NO 125
<211> LENGTH: 1545
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 125

```

```

caagtgcagc tgaagagatc cgccctgga ctggtggccc ctaccagag cctgagcatc 60
acctgtaccg tgtccggett cagcctgacc agctacggcg tgtcatgggt gcgccagcct 120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac 180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctctg 240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcca gggcatcacc 300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg 360
tctagcggag gcggaggatc tggcgccgga ggaagtggcg gagggggatc tgggggaggc 420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca 480
gtgaccatca catgccgggc cagcgagAAC atcgacagct acctggcctg gtatcagcag 540
aagcagggca agagccccca gctgctgggt tacgcgcgca cctttctggc cgacgatgtg 600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactccctg 660
cagagcgagg acgtggcccg gtactactgc cagcactact acagcacccc ctacaccttc 720
ggctcgggca ccaagctgga aatcaaagga ggcggcgga gtgtgcacat gcccctgggc 780
ttcctgggcc ccagacaggc cagagtctgt aacggggggg gcggaggcag tgggggggga 840
ggatccgagg tgcagctgct ggaatctggc ggcggactgg tgcagcctgg cggatctctg 900
agactgagct gtgcgcgcag cggcttcacc ttcagcacct acgccatgaa ctgggtgcgc 960
caggcccctg gcaaaggcct ggaatgggtg tcccggatca gaagcaagta caacaactac 1020
gccacctact acgcccagag cgtgaagggc cggttcacca tcagccggga cgacagcaag 1080
aacaccctgt acctgcagat gaacagcctg cgggcccagg acaccgccgt gtactattgt 1140
gtgcggcacg gcaacttcgg caacagctat gtgtcttggc ttgcctactg gggccagggc 1200
accctcgtga ccgtgtcaag cgctagcgtg gccgctccct ccgtgttcat cttcccacct 1260
tccgacgagc agctgaagtc cggcacccgt tctgtctgtg gcctgctgaa caactctac 1320
ccccgcgagg ccaagggtgca gtggaagggt gacaacgccc tgcagtccgg caacagccag 1380
gaatccgtga ccgagcagga ctccaaggac agcacctact ccctgtcctc caccctgacc 1440
ctgtccaagg ccgactacga gaagcacaag gtgtacgcct gcgaagtgac ccaccagggc 1500
ctgtctagcc ccgtgaccaa gtctttcaac cggggcgagt gctga 1545

```

```

<210> SEQ ID NO 126
<211> LENGTH: 1545
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```


-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 126

```

caagtgcagc tgaagagtc cggccctgga ctggtggccc ctagccagag cctgagcatc    60
acctgtaccg tgtecggtt cagcctgacc agctacggcg tgatcatgggt gcgccagcct    120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac    180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttcctg    240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcca ggccatcacc    300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg    360
tctagcggag gcggaggatc tggcgcgga ggaagtggcg gagggggatc tggggggaggc    420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca    480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag    540
aagcagggca agagccccca gctgctggtg tacgcgccca cctttctggc cgacgatgtg    600
cccagcagat tcagcggcag cggaaaggcg acacagtaca gcctgaagat caactccctg    660
cagagcgagg acgtggcccg gtactactgc cagcactact acagacccc ctacaccttc    720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtgga    780
ggcgcgcgaa gtggcgaggc cggagggggg ggaagtgggg gcggaggcag tgggggggga    840
ggatccgagg tgcagctgct ggaatctggc ggcgactgg tgcagcctgg cggatctctg    900
agactgagct gtgccgcag cggcttcacc ttcagcacct acgccatgaa ctgggtgcgc    960
caggccccctg gcaaaggcct ggaatgggtg tcccgatca gaagcaagta caacaactac   1020
gccacctact acgccgacag cgtgaagggc cggttcacca tcagccggga cgacagcaag   1080
aacaccctgt acctgcagat gaacagcctg cgggcccagg acaccgccgt gtactattgt   1140
gtgcggcacg gcaacttcgg caacagctat gtgtcttggg ttgcctactg gggccagggc   1200
accctcgtga ccgtgtcaag cgctagcgtg gccgctccct ccgtgttcat ctteccacct   1260
tccgacgagc agctgaagtc cggcaccgct tctgtcgtgt gcctgtcga caactctac   1320
cccccgaggc ccaaggtgca gtggaagggt gacaacgccc tgcagtccgg caacagccag   1380
gaatccgtga ccgagcagga ctccaaggac agcacctact ccctgtcctc caccctgacc   1440
ctgtccaagg ccgactacga gaagcacaag gtgtacgcct gcgaagtgac ccaccagggc   1500
ctgtctagcc ccgtgaccaa gtctttcaac cggggcgaggt gctga                    1545

```

<210> SEQ ID NO 127

<211> LENGTH: 699

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 127

```

gaagtgcagc tgctggaatc cggcggagga ctggtgcagc ctggcggatc tctgagactg    60
tcttctgccc cctccggctt cactttctcc acctaccca tgaactgggt gcgacaggct    120
cctggcaagg gcctggaatg ggtgtcccg atcagatcca agtacaacaa ctacgccacc    180
tactacgccg actccgtgaa gggccgggtc accatctctc gggacgactc caagaacacc    240
ctgtacctgc agatgaatc cctgcgggcc gaggacaccg ccgtgtacta ttgtgtgcgg    300

```

-continued

cacggcaact tgcgcaactc ctatgtgtct tggtttgcc actggggcca gggcaccctc	360
gtgaccgtgt catctgctag cgtggccgct ccctccgtgt tcatcttccc accttccgac	420
gagcagctga agtcggcgac cgcttctgtc gtgtgcctgc tgaacaactt ctacccccgc	480
gaggccaagg tgcagtggaa ggtggacaac gccctgcagt ccggcaacag ccaggaatcc	540
gtgaccgagc aggactccaa ggacagcacc tactccctgt cctccacct gacctgtcc	600
aaggccgact acgagaagca caaggtgtac gcctgcgaag tgaccacca gggcctgtct	660
agccccgtga ccaagtcttt caaccggggc gagtgctga	699

<210> SEQ ID NO 128

<211> LENGTH: 2871

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 128

caagtgcagc tgaagagtc cggccctgga ctggtggccc ctagccagag cctgagcatc	60
acctgtaccg tgtccggett cagcctgacc agctacggcg tgtcatgggt gcgccagcct	120
ccaggcaagt gtctggaatg gctgggcata atctggggcg acggcagcac caattaccac	180
agcgcctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctctg	240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcca gggcatcacc	300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagt	360
tctagcggag gcggaggatc tggcggcgga ggaagtggcg gagggggatc tgggggaggc	420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca	480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag	540
aagcagggca agagccccca gctgctggtg tacgcgcgca cctttctggc cgacgatgtg	600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactccctg	660
cagagcgagg acgtggcccg gtactactgc cagcactact acagcacccc ctacaccttc	720
ggctgcggca ccaagctgga aatcaaagga ggcggcgga gtgtgcacat gcccctgggc	780
ttcctgggcc ccagacaggc cagagtctgt aacggggggg gcggaggcag tgggggggga	840
ggatcccagg ccgtcgtgac ccaggaaccc agcctgacag tgtctcctgg cggcaccgtg	900
acctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg	960
caggaaaagc ccggccaggc cttcagagga ctgatcggcg gcaccaaaa gagagccct	1020
ggcaccctg ccagattcag cggatctctg ctgggaggaa aggcgcctt gacactgtct	1080
ggcgccagc cagaagatga ggccgagtag tactgcgccc tgtggtacag caacctgtgg	1140
gtgttcggcg gaggcaccaa gctgacagt ctgagcagcg cttccacca gggacccagt	1200
gtgttcccc tggccccag ctccaagtct acatcgggtg gcacagctgc cctgggatgt	1260
ctcgtgaagg actactttcc tgagcctgtg acagtgtctt ggaacagcgg agccctgacc	1320
agcggagtgc acacattccc tgcagtgtg cagagcagcg gcctgtatag cctgagcagc	1380
gtcgtgaccg tgccttctc tagcctggga acacagacat atatctgtaa tgtgaatcat	1440
aagcccagta atacaaaagt ggataagaaa gtggaaccta agagctgcga tggcggagga	1500
gggtccggag gcggagggtc ccaggtgcag ctggtgcagt ctggcgccga agtgaagaaa	1560

-continued

```

ccaggcgcca gcgtaagggt gtctgcaag gccagcggt acagcttcac cggtacacc 1620
atgaactggg tgcgccaggc tcttgacag gccctggaat ggatgggct gatcacccc 1680
tacaacggcg ccagcagcta caaccagaag ttccggggca aggccacat gaccgtggac 1740
accagcacct ccaccgtgta tatggaactg agcagcctgc ggagcgagga caccgctgt 1800
tactattgtg ccagaggcgg ctacgacggc agaggtctcg attattgggg ccagggcacc 1860
ctcgtgacgg tgtctctgct tagcaccaag ggccctccg tgttccccct ggccccagc 1920
agcaagagca ccagcgggcg cacagcgct ctgggctgcc tggcgagga ctactcccc 1980
gagcccgta cgtgtctctg gaacagcgga gccctgacct cggcggtgca cacttcccc 2040
gccgtgctgc agagtcttgg cctgtatagc ctgagcagcg tggtcacct gccttctagc 2100
agcctgggca ccagaccta catctgcaac gtgaaccaca agccagcaa caccaagggt 2160
gacgagaagg tggagcccaa gagctgcgac aaaactcaca catgccacc gtgccagca 2220
cctgaagctg cagggggacc gtcagtcttc ctcttcccc caaaaccaa ggacaccctc 2280
atgatctccc ggacctga ggtcacatgc gtgggtgtgg acgtgagcca cgaagacct 2340
gaggtaagt tcaactggta cgtggacggc gtggaggtgc ataagccaa gacaaagccg 2400
cgggaggagc agtacaacag cacgtacct gtggtcagcg tcctcacct cctgcaccag 2460
gactggctga atggcaagga gtacaagtgc aaggtctcca acaagacct cggcgcccc 2520
atcgagaaaa ccctctcaa agccaaaggg cagccccgag aaccacaggt gtacacctg 2580
ccccatgcc gggatgagct gaccaagaac caggtcagcc tgtgggtgct ggtcaaaggc 2640
ttctatccca gcgacatgc cgtggagtgg gagagcaatg ggcagcgga gaacaactac 2700
aagaccacgc ctccgtgct ggactccgac ggctcctct tcctctacag caagctcacc 2760
gtggacaaga gcaggtggca gcagggaac gtcttctcat gctccgtgat gcatgaggct 2820
ctgcacaacc actacacgca gaagagctc tcctgtctc cgggtaaatg a 2871

```

<210> SEQ ID NO 129

<211> LENGTH: 2871

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 129

```

caagtgcagc tgaagagtc cggcctgga ctgggtggcc ctagccagag cctgagcatc 60
acctgtaccg tgtccggtt cagcctgacc agctacggcg tgtcatgggt gcgccagcct 120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac 180
agcgcctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctg 240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcaa gggcacacc 300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagt 360
tctagcggag gcggaggatc tggcgcgga ggaagtggcg gagggggatc tgggggaggc 420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca 480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag 540
aagcagggca agagcccca gctgctggtg tacgccgcca cctttctggc cgacgatgtg 600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactcctg 660

```

-continued

cagagcgagg acgtggcccg gtactactgc cagcactact acagcacccc ctacaccttc	720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccgagggcgg cggaagtgga	780
ggcgccggaa gtggcggagg cggagggggg ggaagtgggg gcgagggcag tgggggggga	840
ggatcccagg ccgtcgtgac ccaggaaccc agcctgacag tgtctcctgg cggcacctgt	900
accctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg	960
caggaaaagc ccggccaggc cttcagagga ctgacggcg gcaccaacaa gagagcccct	1020
ggcacccctg ccagattcag cggatctctg ctgggaggaa aggcgcgcct gacactgtct	1080
ggcgcccagc cagaagatga ggccgagtag tactgcgccc tgtggtacag caacctgtgg	1140
gtgttcggcg gaggcaccaa gctgacagtg ctgagcagcg cttccaccaa gggacccagt	1200
gtgttccccc tggcccccag ctccaagtct acatccggtg gcacagctgc cctgggatgt	1260
ctcgtgaagg actactttcc tgagcctgtg acagtgtctt ggaacagcgg agccctgacc	1320
agcggagtgc acacattccc tgcagtgtg cagagcagcg gcctgtatag cctgagcagc	1380
gtcgtgacgg tgccttcttc tagcctggga acacagacat atatctgtaa tgtgaatcat	1440
aagcccagta ataccaaagt ggataagaaa gtggaaccta agagctgcga tggcggagga	1500
gggtccggag gcggaggggc ccagggtgcag ctggtgcagt ctggcgccga agtgaagaaa	1560
ccagcgccca gcgtgaaggt gtccctgcaag gccacgggct acagcttcac cggctacacc	1620
atgaactggg tgcgccaggc tcctggacag gccctggaat ggatgggcct gatcaccccc	1680
tacaacggcg ccagcagcta caaccagaag ttccggggca aggccaccat gaccgtggac	1740
accagcacct ccaccgtgta tatggaactg agcagcctgc ggagcgagga caccgccgtg	1800
tactattgtg ccagagggcg ctacgacggc agaggcttcg attattgggg ccagggcacc	1860
ctcgtgacgg tgtcctctgc tagcaccaag ggcccccctg tgttccccct ggcccccagc	1920
agcaagagca ccagcgggcg cacagccgct ctgggctgcc tggctgagga ctacttcccc	1980
gagcccgtga ccgtgtcctg gaacagcgga gccctgacct ccggcgtgca cacttcccc	2040
gccgtgtgc agagtctctg cctgtatagc ctgagcagcg tggtcacctg gccttctagc	2100
agcctgggca ccagaccta catctgcaac gtgaaccaca agcccagcaa caccaaggtg	2160
gacgagaagg tggagcccaa gagctgcgac aaaactcaca catgccacc gtgccagca	2220
cctgaagctg cagggggacc gtcagtcttc ctcttcccc caaaaccaa ggacaccctc	2280
atgatctccc ggacccctga ggtcacatgc gtggtgggtg acgtgagcca cgaagacct	2340
gaggtcaagt tcaactggta cgtggacggc gtggagggtg ataatgcaa gacaaagccg	2400
cgggaggagc agtacaacag cacgtaccgt gtggtcagcg tcctcacctg cctgcaccag	2460
gactggctga atggcaagga gtacaagtgc aaggtctcca acaaagccct cggcgccccc	2520
atcgagaaaa ccattctcaa agccaaaggg cagccccgag aaccacaggt gtacaccctg	2580
cccccatgcc gggatgagct gaccaagaac caggtcagcc tgtggtgcct ggtcaaaggc	2640
ttctatccca gcgacatgc cgtggagtgg gagagcaatg ggcagccgga gaacaactac	2700
aagaccacgc ctcccgtgct ggactccgac ggctccttct tcctctacag caagctcacc	2760
gtggacaaga gcagggtggc gcaggggaac gtcttctcat gctccgtgat gcatgaggct	2820
ctgcacaacc actacacgca gaagagcctc tccctgtctc cgggtaaatg a	2871

-continued

<210> SEQ ID NO 130
<211> LENGTH: 2025
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 130

caggccgtcg tgaccagga acccagcctg acagtgtctc ctggcggcac cgtgaccctg	60
acatgtggca gttctacagg cgccgtgacc accagcaact acgccaactg ggtgcaggaa	120
aagcccgccc aggccttcag aggactgatc ggccggcacca acaagagagc cctgggcacc	180
cctgccagat tctccggttc tctgctgggc ggcaaggctg ccctgactct gtctggtgct	240
cagcctgagg acgaggccga gtactactgc gccctgtggt actccaacct gtgggtgttc	300
ggcggaggca ccaagctgac cgtgctgtcc agcgcttcca ccaagggacc cagtgtgttc	360
cccctggccc ccagctccaa gtctacatcc ggtggcacag ctgccctggg atgtctctgtg	420
aaggactact ttcttgagcc tgtgacagtg tcttggaaca gcggagccct gaccagcgga	480
gtgcacacat tccttgacgt gctgcagagc agcgccctgt atagcctgag cagcgtcgtg	540
accgtgcctt cctctagcct gggaacacag acatatatct gtaatgtgaa tcataagccc	600
agtaatacca aagtggataa gaaagtggaa cctaagagct gcgatggcgg aggaggggtcc	660
ggagggcgag ggtcccaggt gcagctgggt cagtctggcg ccgaagtgaa gaaaccaggc	720
gccagcgtga aggtgtcctg caaggccagc ggctacagct tcaccggcta caccatgaac	780
tgggtgcccc aggtcctcgg acagggcctg gaatggatgg gcctgatcac cccctacaac	840
ggcgccagca gctacaacca gaagttccgg ggcaaggcca ccatgaccgt ggacaccagc	900
acctccacgg tgtatatgga actgagcagc ctgcggagcg aggacaccgc cgtgtactat	960
tgtgccagag gcggctacga cggcagaggc ttcgattatt ggggccaggg caccctcgtg	1020
accgtgtcct ctgctagcac caagggtccc tccgtgttcc ccctggcccc cagcagcaag	1080
agcaccagcg gcggcacagc cgctctgggc tgccctggctg aggactactt ccccgagccc	1140
gtgaccgtgt cctggaacag cggagccctg acctccggcg tgcacacctt ccccgccgtg	1200
ctgcagagtt ctggcctgta tagcctgagc agcgtgggtc ccgtgccttc tagcagcctg	1260
ggcaccagca cctacatctg caacgtgaac cacaagccca gcaacaccaa ggtggacgag	1320
aaggtggagc ccaagagctg cgacaaaact cacacatgcc caccgtgccc agcacctgaa	1380
gctgcagggg gaccgtcagt ctctctcttc cccccaaaac ccaaggacac cctcatgatc	1440
tcccggaccc ctgaggtcac atcgctgggt gtggacgtga gccacgaaga ccctgaggtc	1500
aagttcaact ggtacgtgga cggcgtggag gtgcataatg ccaagacaaa gccgcgggag	1560
gagcagtaca acagcacgta ccgtgtgggt agcgtcctca ccgtcctgca ccaggactgg	1620
ctgaatggca aggagtacaa gtgcaaggtc tccaacaaag ccctcggcgc ccccatcgag	1680
aaaaccatct ccaagccaa agggcagccc cgagaaccac aggtgtacac cctgccccca	1740
tgccgggatg agctgaccaa gaaccaggtc agcctgtggt gcctggtcaa aggtctctat	1800
cccagcgaca tcgccgtgga gtgggagagc aatgggcagc cggagaacaa ctacaagacc	1860
acgcctcccg tgctggactc cgacggctcc ttcttctct acagcaagct caccgtggac	1920
aagagcaggt ggcagcaggg gaacgtcttc tcatgctccg tgatgcatga ggctctgcac	1980
aaccactaca cgcagaagag cctctccctg tctccgggta aatga	2025

-continued

```

<210> SEQ ID NO 131
<211> LENGTH: 2916
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 131
caagtgcagc tgaaagagtc cgccctgga ctggtggccc ctagccagag cctgagcatc      60
acctgtaccg tgtccggctt cagcctgacc agctacggcg tgtcatgggt gcgccagcct      120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac      180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttcctg      240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcaa gggcatcacc      300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg      360
tctagcggag gcggaggatc tggcggcgga ggaagtggcg gagggggatc tgggggaggc      420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca      480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag      540
aagcagggca agagccccca gctgctggtg tacgcgcgca cctttctggc cgacgatgtg      600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactccctg      660
cagagcgagg acgtggcccg gtactactgc cagcactact acagacccc ctacaccttc      720
ggctcgcgca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtgga      780
ggcgcgcgaa gtttcgtggg ggggacgggg ggcggaggca gtgggggggg aggatccggg      840
ggatccgagg tgcagctgct ggaatctggc ggcggactgg tgcagcctgg cggatctctg      900
agactgagct gtgccgcag cggtctcacc ttcagcacct acgccatgaa ctgggtgctg      960
caggccccctg gcaaaggcct ggaatgggtg tcccggatca gaagcaagta caacaactac      1020
gccacctact acgccgacag cgtgaagggc cggttcacca tcagccggga cgacagcaag      1080
aacaccctgt acctgcagat gaacagcctg cgggcccagg acaccgccgt gtactattgt      1140
gtgcggcacg gcaacttcgg caacagctat gtgtcttggt ttgcctactg gggccagggc      1200
accctcgtga ccgtgtcaag cgctagcaca aagggcccta gcgtgttccc tctggcccc      1260
agcagcaaga gcacaagcgg cggaacagcc gccctgggct gcctcgtgaa ggactacttc      1320
cccgagcccc tgacagtgtc ttggaacagc ggagccctga caagcggcgt gcacaccttc      1380
cctgcccgtg tgcagagcag cggcctgtac tccctgagca gcgtggtcac cgtgcctagc      1440
agcagcctgg gcacccagac ctacatctgc aacgtgaacc acaagcccag caacaccaaa      1500
gtggacaaga aggtggagcc caagagctgt gatggcggag gaggggtccg gggcggaggga      1560
tccgaggtgc aattggttga atctggtggt ggtctggtaa aaccgggcgg ttccctgctg      1620
ctgagctgcg cggtctccgg gttcaccttc tccaacgcgt ggatgagctg ggttcgccag      1680
gccccgggca aaggcctcga gtgggttggc cgtatcaagt ctaaaaactga cgggtggcacc      1740
acggattacg cggtctcagt taaaggtcgt ttaccattt cccgcgacga tagcaaaaac      1800
actctgtatc tgcagatgaa ctctctgaaa actgaagaca ccgcagtcta ctactgtact      1860
accccgctgg aatggtcttg gtacgattat tggggccagg gcacgctggt tacggtgtct      1920
agcgctagta ccaagggccc cagcgtgttc cccctggcac ccagcagcaa gagcacatct      1980

```

-continued

```

ggcggaacag cgcctctggg ctgtctgggtg aaagactact tccccgagcc cgtgaccgtg 2040
tcttggaact ctggcgccct gaccagcggc gtgcacacct ttccagccgt gctgcagagc 2100
agcggcctgt actccctgtc ctccgtgggtc accgtgcctt ctagctccct gggaacacag 2160
acatatatct gtaatgtcaa tcacaagcct tccaacacca aagtcgataa gaaagtcgag 2220
cccaagagct gcgacaaaac tcacacatgc ccaccgtgcc cagcacctga agctgcaggg 2280
ggaccgtcag tcttcctctt ccccccaaaa cccaaggaca ccctcatgat ctcccgacc 2340
cctgaggtea catgcgtggt ggtggacgtg agccacgaag accctgaggt caagttcaac 2400
tggtacgtgg acggcgtgga ggtgcataat gccaaagaaa agccgcggga ggagcagtac 2460
aacagcacgt accgtgtggt cagcgtcctc accgtcctgc accaggactg gctgaatggc 2520
aaggagtaca agtgcaaggt ctccaacaaa gccctcggcg ccccatcgga gaaaaccatc 2580
tccaaagcca aagggcagcc ccgagaacca caggtgtaca ccctgcccc atgccgggat 2640
gagctgacca agaaccaggt cagcctgtgg tgcctggtea aaggcttcta tcccagcgac 2700
atcgccgtgg agtgggagag caatgggcag ccggagaaca actacaagac cagcctccc 2760
gtgctggact ccgacggctc cttctctctc tacagcaagc tcaccgtgga caagagcagg 2820
tggcagcagg ggaacgtctt ctcatgctcc gtgatgcatg aggtctctga caaccactac 2880
acgcagaaga gcctctccct gtctccgggt aatga 2916

```

```

<210> SEQ ID NO 132
<211> LENGTH: 449
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 132

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Asp Tyr
20          25          30
Thr Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ala Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Val Asn Arg Arg Phe
50          55          60
Lys Gly Arg Phe Thr Leu Ser Val Asp Arg Ser Lys Asn Thr Leu Tyr
65          70          75          80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95
Ala Arg Asn Leu Gly Pro Phe Phe Tyr Phe Asp Tyr Trp Gly Gln Gly
100          105          110
Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115          120          125
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130          135          140
Gly Cys Leu Val Glu Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145          150          155          160
Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165          170          175

```

-continued

Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
		195					200					205			
Ser	Asn	Thr	Lys	Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
		210				215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro
225					230					235					240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
			245						250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
			260					265					270		
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
		275					280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
		290				295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
305					310					315					320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys
			325						330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Cys	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Ser
		355					360					365			
Cys	Ala	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
		370				375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
385					390					395					400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Val	Ser	Lys	Leu	Thr	Val	Asp	Lys
			405						410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
		435					440					445			

Lys

<210> SEQ ID NO 133

<211> LENGTH: 214

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 133

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5					10					15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Ala	Ser	Gln	Asp	Val	Ser	Thr	Ala
		20						25					30		
Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
		35				40						45			
Tyr	Ser	Ala	Ser	Phe	Arg	Tyr	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55				60					
Ser	Arg	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro

-continued

65	70	75	80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro	85	90	95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala	100	105	110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Arg Lys Leu Lys Ser Gly	115	120	125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala	130	135	140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln	145	150	155
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser	165	170	175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr	180	185	190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser	195	200	205
Phe Asn Arg Gly Glu Cys	210		

<210> SEQ ID NO 134

<211> LENGTH: 675

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 134

Gln Ala Val Val Thr Gln Glu Pro Ser Leu Thr Val Ser Pro Gly Gly	1	5	10	15
Thr Val Thr Leu Thr Cys Gly Ser Ser Thr Gly Ala Val Thr Thr Ser	20	25	30	
Asn Tyr Ala Asn Trp Val Gln Glu Lys Pro Gly Gln Ala Phe Arg Gly	35	40	45	
Leu Ile Gly Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe	50	55	60	
Ser Gly Ser Leu Leu Gly Gly Lys Ala Ala Leu Thr Leu Ser Gly Ala	65	70	75	80
Gln Pro Glu Asp Glu Ala Glu Tyr Tyr Cys Ala Leu Trp Tyr Ser Asn	85	90	95	
Leu Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Ser Ser Ala	100	105	110	
Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser	115	120	125	
Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe	130	135	140	
Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly	145	150	155	160
Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu	165	170	175	
Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr	180	185	190	
Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys				

-continued

195					200					205				
Val	Glu	Pro	Lys	Ser	Cys	Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly
210						215					220			
Ser	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro
225					230					235				240
Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Asn	Ile	Lys
				245						250				255
Thr	Tyr	Ile	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu
				260						265				270
Val	Ala	Arg	Ile	Tyr	Pro	Thr	Asn	Gly	Tyr	Thr	Arg	Tyr	Ala	Asp
				275										285
Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Ala	Asp	Thr	Ser	Lys	Asn	Thr
						295					300			Ala
Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr
305					310						315			320
Cys	Ser	Arg	Trp	Gly	Gly	Glu	Gly	Phe	Tyr	Ala	Met	Asp	Tyr	Trp
				325						330				335
Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro
				340						345				350
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr
				355										365
Ala	Leu	Gly	Cys	Leu	Val	Glu	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr
				370			375				380			Val
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro
385					390					395				400
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr
				405						410				415
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn
				420						425				430
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser
				435										Cys
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Gly
				450						455				460
Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
465					470					475				480
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
				485						490				495
Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
				500						505				510
His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
				515						520				525
Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
				530						535				540
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro
				545						550				555
Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln
				565						570				575
Tyr	Thr	Leu	Pro	Pro	Cys	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val
				580						585				590
Leu	Trp	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
				595						600				605

-continued

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
610 615 620

Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
625 630 635 640

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
645 650 655

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
660 665 670

Pro Gly Lys
675

<210> SEQ ID NO 135
<211> LENGTH: 957
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 135

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20 25 30

Gly Val Ser Trp Val Arg Gln Pro Gly Lys Cys Leu Glu Trp Leu
35 40 45

Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile
50 55 60

Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
65 70 75 80

Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala
85 90 95

Lys Gly Ile Thr Thr Val Val Asp Asp Tyr Tyr Ala Met Asp Tyr Trp
100 105 110

Gly Gln Gly Thr Ser Val Thr Val Ser Ser Gly Gly Gly Ser Gly
115 120 125

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Asp Ile
130 135 140

Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly Glu Thr
145 150 155 160

Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala
165 170 175

Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val Tyr Ala
180 185 190

Ala Thr Phe Leu Ala Asp Asp Val Pro Ser Arg Phe Ser Gly Ser Gly
195 200 205

Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser Glu Asp
210 215 220

Val Ala Arg Tyr Tyr Cys Gln His Tyr Tyr Ser Thr Pro Tyr Thr Phe
225 230 235 240

Gly Cys Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Ser Gly Gly
245 250 255

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Gly Gly Ser
260 265 270

-continued

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln	
	275						280					285				
Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys	
	290					295					300					
Gly	Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val	
305				310						315					320	
Gln	Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn	
				325					330					335		
Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly	
		340						345					350			
Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala	
	355					360						365				
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly	
370						375					380					
Gly	Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	
385					390					395					400	
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	
				405				410						415		
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	
		420						425					430			
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	
	435					440					445					
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	
450					455					460						
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	
465				470					475					480		
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	
			485					490						495		
Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Val	
		500						505					510			
Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	
	515					520					525					
Cys	Ala	Ala	Ser	Gly	Phe	Asn	Ile	Lys	Asp	Thr	Tyr	Ile	His	Trp	Val	
530					535					540						
Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ala	Arg	Ile	Tyr	Pro	
545				550					555					560		
Thr	Asn	Gly	Tyr	Thr	Arg	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	
		565					570						575			
Ile	Ser	Ala	Asp	Thr	Ser	Lys	Asn	Thr	Ala	Tyr	Leu	Gln	Met	Asn	Ser	
		580					585					590				
Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ser	Arg	Trp	Gly	Gly	
	595				600					605						
Glu	Gly	Phe	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	
610					615					620						
Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	
625				630					635					640		
Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	
			645					650					655			
Glu	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	
	660						665						670			

-continued

Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly
		675					680					685			
Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly
	690					695					700				
Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys
705					710					715					720
Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys
				725					730					735	
Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu
			740					745					750		
Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu
		755					760					765			
Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys
	770					775					780				
Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys
785					790					795					800
Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu
				805					810					815	
Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys
			820					825					830		
Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys
		835					840					845			
Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys
	850					855					860				
Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys
865					870					875					880
Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln
				885					890					895	
Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly
			900					905					910		
Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln
		915					920					925			
Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn
	930					935					940				
His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys			
945					950					955					

<210> SEQ ID NO 136

<211> LENGTH: 957

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 136

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1				5				10					15		
Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Ser	Tyr
			20				25					30			
Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
		35				40					45				
Gly	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
	50				55				60						

-continued

Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
65					70					75					80
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
			85						90					95	
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
		100						105					110		
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
		115					120					125			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
	130					135					140				
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
145					150					155					160
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala
			165						170					175	
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala
		180						185					190		
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
		195					200					205			
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
	210					215					220				
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
225					230					235					240
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Val	His
			245						250					255	
Met	Pro	Leu	Gly	Phe	Leu	Gly	Pro	Arg	Gln	Ala	Arg	Val	Val	Asn	Gly
		260						265					270		
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln
		275					280					285			
Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys
	290					295					300				
Gly	Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val
305					310					315					320
Gln	Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn
			325						330					335	
Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly
		340						345					350		
Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala
		355					360					365			
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly
	370					375					380				
Gly	Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser
385					390					395					400
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala
			405					410						415	
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val
			420				425						430		
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala
	435						440					445			
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val
	450					455					460				
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His

465					470					475					480				
Lys	Pro	Ser	Asn	Thr	485					Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	495	
Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Val				
500					505					510					515				
Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser				
515					520					525					530				
Cys	Ala	Ala	Ser	Gly	Phe	Asn	Ile	Lys	Asp	Thr	Tyr	Ile	His	Trp	Val				
530					535					540					545				
Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ala	Arg	Ile	Tyr	Pro				
545					550					555					560				
Thr	Asn	Gly	Tyr	Thr	Arg	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr				
565					570					575					580				
Ile	Ser	Ala	Asp	Thr	Ser	Lys	Asn	Thr	Ala	Tyr	Leu	Gln	Met	Asn	Ser				
580					585					590					595				
Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ser	Arg	Trp	Gly	Gly				
595					600					605					610				
Glu	Gly	Phe	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr				
610					615					620					625				
Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro				
625					630					635					640				
Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val				
645					650					655					660				
Glu	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala				
660					665					670					675				
Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly				
675					680					685					690				
Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly				
690					695					700					705				
Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys				
705					710					715					720				
Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys				
725					730					735					740				
Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu				
740					745					750					755				
Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu				
755					760					765					770				
Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys				
770					775					780					785				
Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys				
785					790					795					800				
Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu				
805					810					815					820				
Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys				
820					825					830					835				
Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys				
835					840					845					850				
Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys				
850					855					860					865				
Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys				
865					870					875					880				

-continued

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
885 890 895

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
900 905 910

Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln
915 920 925

Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
930 935 940

His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
945 950 955

<210> SEQ ID NO 137
<211> LENGTH: 449
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 137

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Ile Ile Asn Pro Ser Gly Gly Ser Thr Ser Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Met Thr His Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ser Phe Phe Thr Gly Phe His Leu Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130 135 140

Gly Cys Leu Val Glu Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
180 185 190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
195 200 205

Ser Asn Thr Lys Val Asp Glu Lys Val Glu Pro Lys Ser Cys Asp Lys
210 215 220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly Pro
225 230 235 240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
245 250 255

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
260 265 270

-continued

Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
 275 280 285
 Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
 290 295 300
 Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
 305 310 315 320
 Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Gly Ala Pro Ile Glu Lys
 325 330 335
 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Cys Thr
 340 345 350
 Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Ser
 355 360 365
 Cys Ala Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
 370 375 380
 Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
 385 390 395 400
 Asp Ser Asp Gly Ser Phe Phe Leu Val Ser Lys Leu Thr Val Asp Lys
 405 410 415
 Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 420 425 430
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

Lys

<210> SEQ ID NO 138
 <211> LENGTH: 216
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 138

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Thr Asn Glu His
 85 90 95
 Tyr Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val
 100 105 110
 Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Arg Lys Leu Lys
 115 120 125
 Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
 130 135 140
 Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
 145 150 155 160

-continued

Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser
				165						170				175	
Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys
			180					185					190		
Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr
		195					200					205			
Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys								
	210					215									

<210> SEQ ID NO 139
 <211> LENGTH: 956
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 139

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1				5				10						15	
Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Ser	Tyr
			20					25					30		
Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Cys	Leu	Glu	Trp	Leu
			35				40					45			
Gly	Ile	Ile	Trp	Gly	Asp	Gly	Ser	Thr	Asn	Tyr	His	Ser	Ala	Leu	Ile
	50					55					60				
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
65					70					75				80	
Lys	Leu	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
			85					90						95	
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp
			100					105					110		
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly
			115				120					125			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile
	130					135					140				
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr
145					150					155				160	
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala
			165						170					175	
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala
			180					185					190		
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly
			195				200					205			
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp
	210					215					220				
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe
225					230					235				240	
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Val	His
				245					250					255	
Met	Pro	Leu	Gly	Phe	Leu	Gly	Pro	Arg	Gln	Ala	Arg	Val	Val	Asn	Gly
			260					265					270		
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln
			275				280						285		

-continued

Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys
290						295					300				
Gly	Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val
305				310						315					320
Gln	Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn
				325					330					335	
Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly
			340					345					350		
Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala
		355					360					365			
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly
370						375					380				
Gly	Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser
385					390					395					400
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala
				405					410					415	
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val
			420					425					430		
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala
		435					440					445			
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val
450						455					460				
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His
465					470					475					480
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys
				485					490					495	
Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Val	Gln	Leu	Val
			500					505					510		
Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser
		515					520					525			
Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Tyr	Met	His	Trp	Val
530						535					540				
Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	Gly	Ile	Ile	Asn	Pro
545					550					555					560
Ser	Gly	Gly	Ser	Thr	Ser	Tyr	Ala	Gln	Lys	Phe	Gln	Gly	Arg	Val	Thr
				565					570					575	
Met	Thr	His	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr	Met	Glu	Leu	Ser	Ser
			580					585					590		
Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Ser	Phe	Phe
		595					600					605			
Thr	Gly	Phe	His	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val
610						615					620				
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser
625					630					635				640	
Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Glu
				645					650					655	
Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu
			660					665					670		
Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu
		675					680					685			
Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr

-continued

690	695	700
Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val		
705	710	715 720
Asp Glu Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro		
	725	730 735
Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe Leu Phe		
	740	745 750
Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val		
	755	760 765
Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe		
	770	775 780
Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro		
	785	790 795 800
Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr		
	805	810 815
Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val		
	820	825 830
Ser Asn Lys Ala Leu Gly Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala		
	835	840 845
Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Cys Arg		
	850	855 860
Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Trp Cys Leu Val Lys Gly		
	865	870 875 880
Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro		
	885	890 895
Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser		
	900	905 910
Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln		
	915	920 925
Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His		
	930	935 940
Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys		
	945	950 955

<210> SEQ ID NO 140

<211> LENGTH: 956

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 140

Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
1 5 10 15
Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20 25 30
Gly Val Ser Trp Val Arg Gln Pro Pro Gly Lys Cys Leu Glu Trp Leu
35 40 45
Gly Ile Ile Trp Gly Asp Gly Ser Thr Asn Tyr His Ser Ala Leu Ile
50 55 60
Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
65 70 75 80
Lys Leu Asn Ser Leu Gln Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala

-continued

85								90					95				
Lys	Gly	Ile	Thr	Thr	Val	Val	Asp	Asp	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp		
			100					105					110				
Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gly		
		115					120					125					
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile		
	130					135					140						
Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Ala	Ser	Val	Gly	Glu	Thr		
145					150					155					160		
Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Glu	Asn	Ile	Asp	Ser	Tyr	Leu	Ala		
				165					170					175			
Trp	Tyr	Gln	Gln	Lys	Gln	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Val	Tyr	Ala		
		180						185					190				
Ala	Thr	Phe	Leu	Ala	Asp	Asp	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly		
		195					200					205					
Ser	Gly	Thr	Gln	Tyr	Ser	Leu	Lys	Ile	Asn	Ser	Leu	Gln	Ser	Glu	Asp		
	210					215					220						
Val	Ala	Arg	Tyr	Tyr	Cys	Gln	His	Tyr	Tyr	Ser	Thr	Pro	Tyr	Thr	Phe		
225					230					235					240		
Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	Gly	Gly		
				245					250					255			
Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ser		
		260						265					270				
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Ala	Val	Val	Thr	Gln		
		275					280					285					
Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly	Thr	Val	Thr	Leu	Thr	Cys		
	290					295					300						
Gly	Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser	Asn	Tyr	Ala	Asn	Trp	Val		
305					310					315					320		
Gln	Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly	Leu	Ile	Gly	Gly	Thr	Asn		
			325						330					335			
Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Leu	Leu	Gly		
		340						345					350				
Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala	Gln	Pro	Glu	Asp	Glu	Ala		
		355					360					365					
Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn	Leu	Trp	Val	Phe	Gly	Gly		
	370					375					380						
Gly	Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser		
385					390					395					400		
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala		
			405						410					415			
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val		
			420					425					430				
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala		
		435					440					445					
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val		
	450					455						460					
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His		
465					470					475					480		
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys		
			485						490					495			

-continued

Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Val	Gln	Leu	Val
			500					505					510		
Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser
		515					520					525			
Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Tyr	Met	His	Trp	Val
	530					535						540			
Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	Gly	Ile	Ile	Asn	Pro
545					550					555					560
Ser	Gly	Gly	Ser	Thr	Ser	Tyr	Ala	Gln	Lys	Phe	Gln	Gly	Arg	Val	Thr
				565					570					575	
Met	Thr	His	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr	Met	Glu	Leu	Ser	Ser
			580						585				590		
Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Ser	Phe	Phe
		595					600					605			
Thr	Gly	Phe	His	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val
	610					615					620				
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser
625					630					635					640
Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Glu
			645						650					655	
Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu
			660					665					670		
Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu
		675					680					685			
Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr
	690					695					700				
Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val
705					710					715					720
Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro
			725						730					735	
Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe
			740					745					750		
Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val
		755					760					765			
Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe
	770					775					780				
Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro
785				790					795						800
Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr
			805						810					815	
Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val
		820						825					830		
Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala
		835					840					845			
Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Cys	Arg
	850					855					860				
Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Trp	Cys	Leu	Val	Lys	Gly
865				870						875					880
Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro
				885					890						895

-continued

Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser
			900					905					910		
Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln
		915					920					925			
Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His
	930					935					940				
Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys				
945					950					955					

<210> SEQ ID NO 141
 <211> LENGTH: 674
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 141

Gln	Ala	Val	Val	Thr	Gln	Glu	Pro	Ser	Leu	Thr	Val	Ser	Pro	Gly	Gly
1				5					10					15	
Thr	Val	Thr	Leu	Thr	Cys	Gly	Ser	Ser	Thr	Gly	Ala	Val	Thr	Thr	Ser
			20					25					30		
Asn	Tyr	Ala	Asn	Trp	Val	Gln	Glu	Lys	Pro	Gly	Gln	Ala	Phe	Arg	Gly
		35					40					45			
Leu	Ile	Gly	Gly	Thr	Asn	Lys	Arg	Ala	Pro	Gly	Thr	Pro	Ala	Arg	Phe
	50					55					60				
Ser	Gly	Ser	Leu	Leu	Gly	Gly	Lys	Ala	Ala	Leu	Thr	Leu	Ser	Gly	Ala
65					70					75				80	
Gln	Pro	Glu	Asp	Glu	Ala	Glu	Tyr	Tyr	Cys	Ala	Leu	Trp	Tyr	Ser	Asn
			85					90						95	
Leu	Trp	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Ser	Ser	Ala
			100					105					110		
Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser
		115					120					125			
Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe
	130					135					140				
Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly
145					150					155				160	
Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu
			165					170						175	
Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr
		180					185						190		
Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys
	195						200					205			
Val	Glu	Pro	Lys	Ser	Cys	Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly
	210					215					220				
Ser	Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly
225					230					235				240	
Ala	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser
			245						250					255	
Tyr	Tyr	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp
		260					265						270		
Met	Gly	Ile	Ile	Asn	Pro	Ser	Gly	Gly	Ser	Thr	Ser	Tyr	Ala	Gln	Lys
	275						280					285			

-continued

Phe	Gln	Gly	Arg	Val	Thr	Met	Thr	His	Asp	Thr	Ser	Thr	Ser	Thr	Val
290						295					300				
Tyr	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr
305					310					315					320
Cys	Ala	Arg	Ser	Phe	Phe	Thr	Gly	Phe	His	Leu	Asp	Tyr	Trp	Gly	Gln
				325					330					335	
Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val
		340						345					350		
Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala
		355					360					365			
Leu	Gly	Cys	Leu	Val	Glu	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser
	370					375					380				
Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val
385					390					395					400
Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro
				405					410					415	
Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys
			420					425					430		
Pro	Ser	Asn	Thr	Lys	Val	Asp	Glu	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp
		435					440					445			
Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Ala	Gly	Gly
	450					455					460				
Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile
465					470					475					480
Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu
				485					490					495	
Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His
		500						505					510		
Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg
		515					520					525			
Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys
	530					535					540				
Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Gly	Ala	Pro	Ile	Glu
545					550					555					560
Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr
				565					570					575	
Thr	Leu	Pro	Pro	Cys	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu
		580						585					590		
Trp	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp
		595					600					605			
Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val
	610					615					620				
Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp
625					630					635					640
Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His
				645					650					655	
Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro
		660						665					670		
Gly	Lys														

-continued

<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 142

Asp Tyr Thr Met Asp
1 5

<210> SEQ ID NO 143
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 143

Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Val Asn Arg Arg Phe Lys
1 5 10 15

Gly

<210> SEQ ID NO 144
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 144

Asn Leu Gly Pro Phe Phe Tyr Phe Asp Tyr
1 5 10

<210> SEQ ID NO 145
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 145

Thr Ser Asn Tyr Ala Asn Trp
1 5

<210> SEQ ID NO 146
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 146

Gly Thr Asn Lys Arg Ala Pro Gly Thr Pro Ala Arg Phe Ser Gly Ser
1 5 10 15

Leu Leu Gly Gly
20

<210> SEQ ID NO 147
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 147

Thr Lys Leu Thr Val
1 5

<210> SEQ ID NO 148

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 148

Lys Ala Ser Gln Asp Val Ser Thr Ala Val Ala
1 5 10

<210> SEQ ID NO 149

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 149

Ser Ala Ser Phe Arg Tyr Thr
1 5

<210> SEQ ID NO 150

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 150

Gln Gln His Tyr Thr Thr Pro Pro Thr
1 5

<210> SEQ ID NO 151

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 151

Ser Tyr Tyr Met His
1 5

<210> SEQ ID NO 152

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 152

Ile Ile Asn Pro Ser Gly Gly Ser Thr Ser Tyr Ala Gln Lys Phe Gln
1 5 10 15

Gly

<210> SEQ ID NO 153

<211> LENGTH: 10

<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 153

Ser Phe Phe Thr Gly Phe His Leu Asp Tyr
1 5 10

<210> SEQ ID NO 154
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 154

Arg Ala Ser Gln Ser Val Ser Ser Ser Tyr Leu Ala
1 5 10

<210> SEQ ID NO 155
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 155

Gly Ala Ser Ser Arg Ala Thr
1 5

<210> SEQ ID NO 156
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 156

Gln Gln Tyr Thr Asn Glu His Tyr Tyr Thr
1 5 10

<210> SEQ ID NO 157
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 157

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Ile Ile Asn Pro Ser Gly Gly Ser Thr Ser Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Met Thr His Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

-continued

Ala Arg Ser Phe Phe Thr Gly Phe His Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Leu Val Thr Val Ser Ser
 115

<210> SEQ ID NO 158
 <211> LENGTH: 109
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 158

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30

Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45

Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80

Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Thr Asn Glu His
 85 90 95

Tyr Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> SEQ ID NO 159
 <211> LENGTH: 119
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 159

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Asp Tyr
 20 25 30

Thr Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Val Asn Arg Arg Phe
 50 55 60

Lys Gly Arg Phe Thr Leu Ser Val Asp Arg Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Asn Leu Gly Pro Phe Phe Tyr Phe Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Leu Val Thr Val Ser Ser
 115

<210> SEQ ID NO 160
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 160

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys Asp Thr
 20 25 30
 Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ser Arg Trp Gly Gly Glu Gly Phe Tyr Ala Met Asp Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 161

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 161

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Thr Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ser Ala Ser Phe Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Arg Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> SEQ ID NO 162

<211> LENGTH: 1350

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 162

gaagttcagc tgggtgaaag cgggtggtggt ctggttcagc ctggtggttag cctgcgtctg 60
 agctgtgcag caagcgggtt tacctttaac gattatacca tggattgggt tcgtcaggca 120
 ccgggttaaag gtctggaatg ggttcagat gttaatccga atagcgggtg tagcattgtt 180

-continued

aaccgctggt ttaaaggtcg ttttaccctg agcgttgatc gtagcaaaaa taccctgtat	240
ctgcaaatga atagtctgcg tgcagaggat accgcagtggt attattgtgc acgtaacctg	300
ggctcgttct tctactttga ttattggggt cagggcaccc tggttaccgt tagcagcgct	360
agcaccaagg gcccctccgt gttccccctg gccccagca gcaagagcac cagcggcggc	420
acagccgctc tgggctgcct ggtcgaggac tacttccccg agcccgtagc cgtgtcctgg	480
aacagcggag ccctgacctc cggcgtgcac accttccccg ccgtgctgca gagttctggc	540
ctgtatagcc tgagcagcgt ggtcacctg ctttctagca gcctgggcac ccagacctac	600
atctgcaacg tgaaccacaa gccagcaac accaaggtgg acgagaaggt ggagcccaag	660
agctgcgaca aaactcacac atgccaccg tgcccagcac ctgaagctgc agggggaccg	720
tcagtccttc tcttcccccc aaaacccaag gacacctca tgatctcccg gacctctgag	780
gtcacatgcg tgggtgggga cgtgagccac gaagacctg aggtcaagtt caactggtac	840
gtggacggcg tggaggtgca taatgccaa gcaaagccgc gggaggagca gtacaacagc	900
acgtacctg tggtcagcgt cctcacctg ctgcaccagg actggctgaa tggcaaggag	960
tacaagtgca aggtctccaa caaagccctc ggcgccccca tcgagaaaac catctccaaa	1020
gccaaggggc agccccgaga accacagggtg tgcacctgc ccccatcccg ggatgagctg	1080
accaagaacc aggtcagcct ctctgctgca gtcaaaggct tctatcccag cgacatcgcc	1140
gtggagtggg agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgctgctg	1200
gactccgacg gctccttctt cctcgtgagc aagctcaccg tggacaagag cagggtggcag	1260
caggggaacg tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgcag	1320
aagagcctct ccctgtcttc gggtaaatga	1350

<210> SEQ ID NO 163

<211> LENGTH: 645

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 163

gacatccaga tgaccagag cccagcagc ctgtctgcca gcgtgggcca cagagtgacc	60
atcacatgca aggccagcca ggacgtgtcc acagccgtgg cctgggtatca gcagaagcct	120
ggcaaggccc ccaagctgct gatctacagc gccagcttcc ggtacaccgg cgtgcccagc	180
agattcagcg gcagcagatc cggcaccgac ttcacctga ccatcagctc cctgcagccc	240
gaggacttcg ccacctacta ctgccagcag cactacacca cccccccac atttggccag	300
ggcaccaagg tggaaatcaa gcgtacgggt gctgcaccat ctgtcttcat cttcccgcca	360
tctgatcgga agttgaaatc tggaaatgcc tctgttgtgt gcctgctgaa taactttat	420
cccagagagg ccaaagtaca gtggaagggt gataacgccc tccaatcggg taactcccag	480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg	540
ctgagcaaa gagactacga gaaacacaaa gtctacgcct gcgaagtac ccatcagggc	600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gttag	645

<210> SEQ ID NO 164

<211> LENGTH: 2874

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 164

```

caagtgcagc tgaagagtc cggccctgga ctggtggccc ctagecagag cctgagcatc   60
acctgtaccg tgtccggctt cagccctgacc agctacggcg tgtcatgggt gcgccagcct   120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac   180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctctg   240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcaa gggcatcacc   300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg   360
tctagcggag gcggaggatc tggcgcgga ggaagtggcg gagggggatc tgggggagggc   420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca   480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag   540
aagcagggca agagcccca gctgctggtg tacgcgcca cctttctggc cgacgatgtg   600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactccctg   660
cagagcgagg acgtggcccg gtactactgc cagcactact acagcaccac ctacaccttc   720
ggctgcggca ccaagctgga aatcaaaggc gggggaggct ccggaggcgg cggaagtgga   780
ggcgcgcgaa gtggcgaggc cgaggggggg ggaagtgggg gcggaggcag tgggggggga   840
ggatcccagg ccgtcgtgac ccaggaaacc agcctgacag tgtctcctgg cggcacctgtg   900
accctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg   960
caggaaaagc ccggccaggc cttcagagga ctgatcgcg gcaccaacaa gagagcccct   1020
ggcacccttg ccagattcag cggatctctg ctgggaggaa agggccgccc gacactgtct   1080
ggcgcccagc cagaagatga ggccgagtag tactgcgccc tgtggtacag caacctgtgg   1140
gtgttcggcg gaggcaccaa gctgacagtg ctgagcagcg cttccaccaa gggacccagt   1200
gtgttcccc tggcccccag ctccaagtct acatccgggtg gcacagctgc cctgggatgt   1260
ctcgtgaagg actactttcc tgagcctgtg acagtgtctt ggaacagcgg agccctgacc   1320
agcggagtgc acacattccc tgcagtgtg cagagcagcg gcctgtatag cctgagcagc   1380
gtcgtgacgg tgccttcttc tagcctggga acacagacat atatctgtaa tgtgaatcat   1440
aagcccagta atacaaaagt ggataagaaa gtggaaccta agagctgcga tggcggagga   1500
gggtccggag gcggagggtc cgagggtccag ctggtcagat ctggaggagg actgggtgcag   1560
ccaggcggat ctctgagact gagctgcgcc gccagcggat tcaacatcaa ggacacctac   1620
atccactggg tgaggcaggc ccttgaaaag ggactggagt gggtgggcag aatctacccc   1680
accaacggct acacaagata cgccgacagc gtgaagggca gattcaccat cagcgccgac   1740
accagcaaga acaccgccta cctgcagatg aacagcctga gagccgagga cacagccgtg   1800
tactactgct ctatagggg aggcgagggc ttctacgcca tggactactg gggacagggc   1860
aactggtga ccgtgtccag cgctagcacc aagggccccc ccgtgttccc cctggccccc   1920
agcagcaaga gcaccagcgg cggcacagcc gctctgggct gcctggtcga ggactacttc   1980
cccgagcccg tgaccgtgtc ctggaacagc ggagccctga cctccggcgt gcacaccttc   2040
cccgcctgct tgcagagttc tggcctgtat agcctgagca gcgtggtcac cgtgccttct   2100

```

-continued

```

agcagcctgg gcacccagac ctacatctgc aacgtgaacc acaagcccag caacaccaag 2160
gtggacgaga aggtggagcc caagagctgc gacaaaactc acacatgccc accgtgccc 2220
gcacctgaag ctgcaggggg accgtcagtc ttcctcttcc ccccaaaacc caaggacacc 2280
ctcatgatct cccggacccc tgaggtcaca tgcgtggagg tggacgtgag ccacgaagac 2340
cctgaggtca agttcaactg gtacgtggac ggcgtggagg tgcataatgc caagacaaag 2400
ccgcgggagg agcagtacaa cagcacgtac cgtgtgggta gcgtcctcac cgtcctgcac 2460
caggactggc tgaatggcaa ggagtacaag tgcaaggctc ccaacaaagc cctcggcgcc 2520
cccatcgaga aaaccatctc caaagccaaa gggcagcccc gagaaccaca ggtgtacacc 2580
ctgcccccat gccgggatga gctgaccaag aaccaggta gcctgtgggtg cctgggtaaa 2640
ggcttctatc ccagcgacat cgccgtggag tgggagagca atgggcagcc ggagaacaac 2700
tacaagacca cgctcccggt gctggactcc gacggctcct tcttctctca cagcaagctc 2760
accgtggaca agagcagggtg gcagcagggg aacgtcttct catgctccgt gatgcatgag 2820
gctctgcaca accactacac gcagaagagc ctctccctgt ctccgggtaa atga 2874

```

<210> SEQ ID NO 165

<211> LENGTH: 2874

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 165

```

caagtgcagc tgaagagtc cgccctgga ctggtggccc ctaccagag cctgagcatc 60
acctgtaccg tgtccggctt cagcctgacc agctacggcg tgtcatgggt gcgccagcct 120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac 180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctctg 240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcaa gggcatcacc 300
accgtgggtg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg 360
tctagcggag gcggaggatc tggcgcgga ggaagtggcg gagggggatc tgggggagggc 420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca 480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag 540
aagcagggca agagccccc gctgctgggt tacgcgcca cctttctggc cgacgatgtg 600
cccagcagat tcagcggcag cggaagcggc acacagtaca gcctgaagat caactccctg 660
cagagcgagg acgtggcccg gtactactgc cagcactact acagcaccac ctacaccttc 720
ggctgcggca ccaagctgga aatcaaagga ggcggcgga gtgtgcacat gccctggggc 780
ttcctgggcc ccagacaggc cagagtcgtg aacggggggg gcggaggcag tgggggggga 840
ggatcccagg ccgtcgtgac ccaggaaccc agcctgacag tgtctcctgg cggcaccgtg 900
accctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg 960
cagggaaaagc ccggccaggc cttcagagga ctgatcggcg gcaccaacaa gagagcccct 1020
ggcaccctcg ccagattcag cggatctctg ctgggaggaa aggcggccct gacactgtct 1080
ggcgccagc cagaagatga ggcgagtag tactgcgccc tgtggtacag caacctgtgg 1140
gtgttcggcg gaggcaccaa gctgacagtg ctgagcagcg cttccacca gggacccagt 1200

```


-continued

gtgttccccc	tggcccccag	ctccaagtct	acatccggtg	gcacagctgc	cctgggatgt	1260
ctcgtgaagg	actactttcc	tgagcctgtg	acagtgtctt	ggaacagcgg	agccctgacc	1320
agcggagtgc	acacattccc	tgacgtgtg	cagagcagcg	gcctgtatag	cctgagcagc	1380
gtcgtgacgg	tgcccttcctc	tagcctggga	acacagacat	atatctgtaa	tgtgaatcat	1440
aagcccagta	ataccaaaagt	ggataagaaa	gtggaacctt	agagctgcga	tggcggaggga	1500
gggtccggag	gcggagggtc	cgaggccag	ctgggtcgagt	ctggaggagg	actgggtcag	1560
ccaggcggat	ctctgagact	gagctgcgcc	gccagcggat	tcaacatcaa	ggacacctac	1620
atccactggg	tgaggcaggc	ccttggaag	ggactggagt	gggtggccag	aatctacccc	1680
accaacggct	acacaagata	cgcgcagcgc	gtgaaggcca	gattccaccat	cagcgcgcag	1740
accagcaaga	acaccgccta	cctgcagatg	aacagcctga	gagccgagga	cacagccgtg	1800
tactactgct	ctagatgggg	aggcggggc	ttctacgcca	tggactactg	gggacagggc	1860
acactggtga	ccgtgtccag	cgctagcacc	aaggggccct	ccgtgttccc	cctggccccc	1920
agcagcaaga	gcaccagcgg	cgccacagcc	gctctgggct	gcctggtcga	ggactacttc	1980
cccagcccg	tgaccgtgtc	ctggaacagc	ggagccctga	cctccggcgt	gcacaccttc	2040
cccgcctg	tgacaggttc	tggcctgtat	agcctgagca	gcgtggtcac	cgtgccttct	2100
agcagcctgg	gcaccagac	ctacatctgc	aacgtgaacc	acaagcccag	caacaccaag	2160
gtggacgaga	agggtggagc	caagagctgc	gacaaaactc	acacatgccc	accgtgcccc	2220
gcacctgaag	ctgcaggggg	accgtcagtc	ttcctcttcc	ccccaaaacc	caaggacacc	2280
ctcatgatct	cccggacccc	tgaggtcaca	tgcgtgggtg	tggacgtgag	ccacgaagac	2340
cctgaggtca	agttcaactg	gtacgtggac	ggcgtggagg	tgcataatgc	caagacaaag	2400
ccgcgggagg	agcagtacaa	cagcacgtac	cgtgtgggtc	gcgtccctac	cgtcctgcac	2460
caggactggc	tgaatggcaa	ggagtacaag	tgcaaggctc	ccaacaaagc	cctcgccgcc	2520
cccacgaga	aaaccatctc	caaagccaaa	gggcagcccc	gagaaccaca	ggtgtacacc	2580
ctgcccccat	gccgggatga	gctgaccaag	aaccagggtc	gcctgtgggtg	cctgggtcaaa	2640
ggcttctatc	ccagcgacat	cgccgtggag	tgggagagca	atgggcagcc	ggagaacaac	2700
tacaagacca	cgcctcccg	gctggactcc	gacggctcct	tcttctctca	cagcaagctc	2760
accgtggaca	agagcagggtg	gcagcagggg	aacgtcttct	catgctccgt	gatgcatgag	2820
gctctgcaca	accactacac	gcagaagagc	ctctccctgt	ctccgggtaa	atga	2874

<210> SEQ ID NO 166

<211> LENGTH: 1350

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 166

cagggtgcaat	tgggttcaatc	tgggtgctgaa	gtaaaaaaac	cgggcgcttc	cggttaaagtg	60
agctgcaaag	catccggata	caccttcaact	tcctattaca	tgcactgggtg	tcgtcaagcc	120
ccgggccagg	gtctggaatg	gatgggcac	attaacccaa	gcggtggctc	tacctcctac	180
gcgcagaaat	tccagggtcg	cgctcacgatg	acccatgaca	ctagcacctc	taccgtttat	240
atggagctgt	ccagcctg	ttctgaagat	actgcagtg	actactgtgc	acgctctttc	300

-continued

ttcactgggt tccatctgga ctattgggggt caaggcaccc tcgtaacggt ttcttctgct	360
agcaccaagg gcccctccgt gtccccctg gccccagca gcaagagcac cagcggcggc	420
acagccgctc tgggtgcct ggctgaggac tacttccccg agcccgtagc cgtgtcctgg	480
aacagcggag ccctgacctc cggcgtgcac accttccccg ccgtgctgca gagttctggc	540
ctgtatagcc tgagcagcgt ggtaacccgt ccttctagca gcctgggcac ccagacctac	600
atctgcaacg tgaaccacaa gcccgcaac accaaggtgg acgagaaggt ggagcccaag	660
agctgcgaca aaactcacac atgccaccg tgcccagcac ctgaagctgc agggggaccg	720
tcagtcttcc tcttcccccc aaaaccaag gacacctca tgatctcccc gacctctgag	780
gtcacatgcg tgggtgggga cgtgagccac gaagacctg aggtcaagtt caactggtag	840
gtggacggcg tggaggtgca taatgccaag acaaagccgc gggaggagca gtacaacagc	900
acgtaccgtg tggtcagcgt cctcacctc ctgcaccagg actggctgaa tggcaaggag	960
tacaagtgca aggtctccaa caaagccctc ggcgccccca tcgagaaaac catctccaaa	1020
gccaagggc agccccgaga accacaggtg tgcacctgc ccccatcccc ggatgagctg	1080
accaagaacc aggtcagcct ctctgctgca gtcaaaggct tctatcccag cgacatcgcc	1140
gtggagtggg agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg	1200
gactccgacg gctccttctt cctcgtgagc aagctcaccg tggacaagag caggtggcag	1260
caggggaacg tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgcag	1320
aagagcctct cctgtctctc gggtaaatga	1350

<210> SEQ ID NO 167
 <211> LENGTH: 651
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 167

gaaatcgtgt taacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagccacc	60
ctctcttgca gggccagtca gagtgttagc agcagctact tagcctggta ccagcagaaa	120
cctggccagg ctcccaggct cctcatctat ggagcatcca gcagggccac tggcatccca	180
gacagggtta gtggcagtgg atccgggaca gacttcactc tcaccatcag cagactggag	240
cctgaagatt ttgcagtgtg ttactgtcag cagtatacca acgaacatta ttatacgttc	300
ggccagggga ccaaagtgga aatcaaacgt acggtggctg caccatctgt ctctcatctc	360
ccgccatctg atcggaagtt gaaatctgga actgcctctg ttgtgtgcct gctgaataac	420
ttctatccca gagaggccaa agtacagtgg aaggtggata acgcctccca atcgggtaac	480
tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc	540
ctgacgctga gcaaagcaga ctacgagaaa cacaaagtct acgcctgcga agtcacccat	600
cagggcctga gctcgcccggt cacaaagagc ttcaacaggg gagagtgtta g	651

<210> SEQ ID NO 168
 <211> LENGTH: 2871
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 168

caagtgcagc tgaagagtc cgccctgga ctggtggccc ctagccagag cctgagcatc	60
acctgtaccg tgtccggtt cagcctgacc agctacggcg tgtcatgggt gcgccagcct	120
ccaggcaagt gtctggaatg gctgggcatc atctggggcg acggcagcac caattaccac	180
agcgccctga tcagcagact gagcatctcc aaggacaaca gcaagagcca ggtgttctctg	240
aagctgaaca gcctgcagac cgacgacacc gccacctact actgcgcca gggcatcacc	300
accgtggtgg acgactacta cgctatggac tactggggcc agggcaccag cgtgacagtg	360
tctagcggag gcggaggatc tggcggcgga ggaagtggcg gagggggatc tgggggaggc	420
ggaagcgata tccagatgac ccagagccct gccagcctgt ctgcctctgt gggcgagaca	480
gtgaccatca catgccgggc cagcgagaac atcgacagct acctggcctg gtatcagcag	540
aagcagggca agagccccca gctgctggtg tacgcgcgca cctttctggc cgacgatgtg	600
cccagcagat tcagcggcag cggaaagcggc acacagtaca gcctgaagat caactccctg	660
cagagcgagg acgtggcccg gtactactgc cagcactact acagaccccc ctacaccttc	720
ggctgcggca ccaagctgga aatcaaagga ggccggcgaa gtgtgcacat gccctggggc	780
ttctctgggc ccagacaggc cagagtcgtg aacggggggg gcggaggcag tgggggggga	840
ggatcccagg ccgtcgtgac ccaggaaccc agcctgacag tgtctcctgg cggcaccgtg	900
accctgacat gtggcagttc tacaggcgcc gtgaccacca gcaactacgc caactgggtg	960
cagggaaaagc ccggccaggc cttcagagga ctgatcgcg gcaccaaca gagagcccct	1020
ggcaccctg ccagattcag cggatctctg ctgggaggaa aggcgcgccct gacactgtct	1080
ggcgcccgag cagaagatga ggccgagtag tactgcgccc tgtggtacag caacctgtgg	1140
gtgttcggcg gaggcacca gctgacagtg ctgagcagcg cttccacca gggacccagt	1200
gtgttcccc tggccccag ctccaagtct acatccggtg gcacagctgc cctgggatgt	1260
ctcgtgaagg actactttcc tgagcctgtg acagtgtctt ggaacagcgg agccctgacc	1320
agcggagtgc acacattccc tgcagtgtg cagagcagcg gcctgtatag cctgagcagc	1380
gtcgtgacgg tgccttcctc tagcctggga acacagacat atatctgtaa tgtgaatcat	1440
aagcccagta atacaaaagt ggataagaaa gtggaaccta agagctgcga tggcggaggga	1500
gggtccggag gcggagggtc ccagggtgcaa ttggttcaat ctggtgctga agtaaaaaa	1560
ccggcgctt ccgttaaagt gagctgcaaa gcacccggat acaccttcac ttctattac	1620
atgcactggg ttctgcaagc cccgggccag ggtctggaat ggatgggcat cattaaccca	1680
agcgtgggt ctacctcta cgcgcagaaa ttccagggtc gcgtcacgat gaccatgac	1740
actagcacct ctaccgttta tatggagctg tccagcctgc gttctgaaga tactgcagtg	1800
tactactgtg cacgtctttt cttcactggt ttccatctgg actattgggg tcaaggcacc	1860
ctcgtaacgg tttctctgc tagcaccaag ggccctccg tgttccccct ggccccagc	1920
agcaagagca ccagcggcgg cacagccgct ctgggctgcc tggtcgagga ctacttcccc	1980
gagcccgta ccgtgtcctg gaacagcggga gccctgacct ccggcgtgca caccttcccc	2040
gccgtgctgc agagtctctg cctgtatagc ctgagcagcg tggtaaccgt gccttctagc	2100
agcctgggca ccagaccta catctgcaac gtgaaccaca agcccagcaa caccaagggtg	2160
gacgagaagg tggagcccaa gagctgcgac aaaactcaca catgcccacc gtgcccagca	2220

-continued

cctgaagctg cagggggacc gtcagtcttc ctcttccccc caaaacccaa ggacaccctc	2280
atgatctccc ggaccctga ggtcacatgc gtggagggtg acgtgagcca cgaagaccct	2340
gaggtaagtc tcaactggta cgtggacggc gtggagggtg ataatgccc gacaaagccg	2400
cgggaggagc agtacaacag cacgtaccgt gtggtcagcg tcctcaccgt cctgcaccag	2460
gactggctga atggcaagga gtacaagtgc aaggctctcca acaaagccct cggcgccccc	2520
atcgagaaaa ccatctccaa agccaaaggg cagcccgag aaccacaggt gtacaccctg	2580
cccccatgcc gggatgagct gaccaagaac caggtcagcc tgtgggtgctt ggtcaaaggc	2640
ttctatccca gcgacatgc cgtggagtgg gagagcaatg ggcagccgga gaacaactac	2700
aagaccagc ctcccgctgt ggactccgac ggctccttct tcctctacag caagctcacc	2760
gtggacaaga gcaggtggca gcaggggaac gtcttctcat gctccgtgat gcatgaggct	2820
ctgcacaacc actacacgca gaagagcctc tcctgtctc cgggtaaatg a	2871

<210> SEQ ID NO 169

<211> LENGTH: 2857

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 169

agagtcggc cctggactgg tggccctag ccagagcctg agcatcacct gtaccgtgtc	60
cggcttcagc ctgaccagct acggcgtgtc atgggtgcgc cagcctccag gcaagtgtct	120
ggaatggctg ggcacatctt ggggcgacgg cagcaccat taccacagcg ccctgatcag	180
cagactgagc atctccaagg acaacagcaa gagccaggtg ttctgaagc tgaacagcct	240
gcagaccgac gacaccgcca cctactactg cgccaagggc atcaccaccg tggaggagca	300
ctactacgct atggactact ggggccaggg caccagcgtg acagtgtcta gcggaggcgg	360
aggatctggc ggcggaggaa gtggcggagg gggatctggg ggaggcggaa gcgatatcca	420
gatgaccagc agccctgcca gcctgtctgc ctctgtgggc gagacagtga ccatcacatg	480
ccgggccagc gagaacatcg acagctacct ggccctggtat cagcagaagc agggcaagag	540
ccccagctg ctggtgtacg ccgccacctt tctggccgac gatgtgccc gcagattcag	600
cggcagcggc agcggcacac agtacagcct gaagatcaac tcctgcaga gcgaggacgt	660
ggcccggtac tactgccagc actactacag caccacctac accttcggct gcggcaccac	720
gctggaaatc aaaggcgggg gaggtccgg aggcggcgga agtggaggcg gcggaagtgg	780
cggaggcgga ggggggggaa gtggggggcg aggcagtggt gggggaggat cccaggccgt	840
cgtgaccagc gaaccagcc tgacagtgtc tcctggcggc accgtgaccc tgacatgtgg	900
cagttctaca ggcgcctgta ccaccagcaa ctacgccaac tgggtgcagg aaaagcccg	960
ccaggccttc agaggactga tcggcggcac caacaagaga gccctggca cccctgccag	1020
attcagcgga tctctgtctg gaggaaggc cgccctgaca ctgtctggcg cccagccaga	1080
agatgaggcc gactactact gcgccctgtg gtacagcaac ctgtgggtgt tcggcggagg	1140
caccaagctg acagtgtgta gcagcgttc caccaaggga cccagtgtgt tccccctggc	1200
ccccagctcc aagtctacat ccggtggcac agctgccctg ggatgtctcg tgaaggacta	1260
ctttcctgag cctgtgacag tgtcttgaa cagcggagcc ctgaccagcg gactgcacac	1320

-continued

attccctgca	gtgctgcaga	gcagcggcct	gtatagcctg	agcagcgtcg	tgaccgtgcc	1380
ttcctctagc	ctgggaacac	agacatatat	ctgtaatgtg	aatcataagc	ccagtaatac	1440
caaagtggat	aagaaagtgg	aacctaaag	ctgcgatggc	ggaggagggt	ctggaggcgg	1500
agggtcccag	gtgcaattgg	ttcaatctgg	tgtgaagta	aaaaaacgg	gcgttccgt	1560
taaagtgagc	tgcaaagcat	ccggatacac	cttcacttcc	tattacatgc	actgggttcg	1620
tcaagccccg	ggccagggtc	tggatggat	gggcatcatt	aacccaagcg	gtggctctac	1680
ctcctacgcg	cagaaattcc	agggtcgcgt	cacgatgacc	catgacacta	gcacctctac	1740
cgtttatatg	gagctgtcca	gcctgcgttc	tgaagatact	gcagtgtact	actgtgcacg	1800
ctctttcttc	actggtttcc	atctggacta	ttggggctca	ggcaccctcg	taacggtttc	1860
ttctgctagc	accaagggcc	cctccgtgtt	ccccctggcc	cccagcagca	agagcaccag	1920
cggcggcaca	gccgctctgg	gctgcctgg	cgaggactac	ttccccgagc	ccgtgaccgt	1980
gtcctggaac	agcggagccc	tgacctccgg	cgtgcacacc	ttccccgcgg	tgtgcagag	2040
ttctggcctg	tatagcctga	gcagcgtgg	caccgtgcct	tctagcagcc	tgggcacca	2100
gacctacatc	tgcaacgtga	accacaagcc	cagcaacacc	aaggtggacg	agaaggtgga	2160
gcccagagc	tgcgacaaaa	ctcacacatg	cccaccgtgc	ccagcacctg	aagctgcagg	2220
gggaccgtca	gtcttctctt	tcccccaaaa	acccaaggac	accctcatga	tctccggac	2280
ccctgaggtc	acatgcgtgg	tgttgagcgt	gagccacgaa	gacctgagg	tcaagttcaa	2340
ctggtacgtg	gacggcgtgg	aggtgcataa	tgccaagaca	aagccgcggg	aggagcagta	2400
caacagcagc	taccgtgtgg	tcagcgtcct	caccgtcctg	caccaggact	ggctgaatgg	2460
caaggagtac	aagtgcgaag	tctccaacaa	agccctcggc	gccccatcg	agaaaacat	2520
ctccaaagcc	aaagggcagc	cccgagaacc	acaggtgtac	accctgcccc	catgcgggga	2580
tgagctgacc	aagaaccagg	tcagcctgtg	gtgcctggtc	aaaggcttct	atcccagcga	2640
catcgccgtg	gagtgggaga	gcaatgggca	gccggagaac	aactacaaga	ccacgcctcc	2700
cgtgtgggac	tccgacggct	ccttcttctt	ctacagcaag	ctcaccgtgg	acaagagcag	2760
gtggcagcag	gggaacgtct	tctcatgtct	cgtgatgcac	gaggctctgc	acaaccacta	2820
cacgcagaag	agcctctccc	tgtctccggg	taaatga			2857

<210> SEQ ID NO 170

<211> LENGTH: 2025

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 170

caggccgtcg	tgaccagga	accagcctg	acagtgtctc	ctggcggcac	cgtgaccctg	60
acatgtggca	gtttacagg	cgccgtgacc	accagcaact	acgccaactg	ggtgcaggaa	120
aagcccggcc	aggccttcag	aggactgatc	ggcggcacca	acaagagagc	ccctggcacc	180
cctgccagat	tcagcggatc	tctgctggga	ggaaaggccg	ccctgacact	gtctggcgcc	240
cagccagaag	atgaggccga	gtactactgc	gccctgtgg	acagcaacct	gtgggtgttc	300
ggcggaggca	ccaagctgac	agtgtgtgac	agcgttcca	ccaagggacc	cagtgtgttc	360
ccccctggcc	ccagctccaa	gtctacatcc	ggtggcacag	ctgcctggg	atgtctcgtg	420

-continued

aaggactact	ttcctgagcc	tgtgacagt	tcttggaaaca	gcgagagccct	gaccagcggga	480
gtgcacacat	tccttgagct	gctgcagagc	agcggcctgt	atagcctgag	cagcgtcgtg	540
accgtgcctt	cctctagcct	gggaacacag	acatatatct	gtaatgtgaa	tcataagccc	600
agtaatacca	aagtggataa	gaaagtggaa	cctaagagct	gcgatggcgg	aggaggggtct	660
ggaggcggag	ggteccaggt	gcaattggtt	caatctgggt	ctgaagtaaa	aaaaccgggc	720
gcttcctgta	aagtgcagct	caaagcatcc	ggatacacct	tcacttccta	ttacatgcac	780
tgggttcgtc	aagccccggg	ccagggtctg	gaatggatgg	gcateattaa	cccaagcggg	840
ggctctacct	cctacgcgca	gaaattccag	ggtcgcgtca	cgatgaccca	tgacactagc	900
acctctaccg	tttatatgga	gctgtccagc	ctgcgttctg	aagatactgc	agtgtactac	960
tgtgcacgct	ctttcttcac	tggtttccat	ctggactatt	ggggtcaagg	caccctcgta	1020
acggtttctt	ctgctagcac	caagggccccc	tccgtgttcc	ccctggcccc	cagcagcaag	1080
agcaccagcg	gcggcacagc	cgtctctggc	tgcttggtcg	aggactactt	ccccgagccc	1140
gtgaccgtgt	cctggaacag	cggagccctg	acctccggcg	tgacacacct	ccccgcgctg	1200
ctgcagagtt	ctggcctgta	tagcctgagc	agcgtggtea	ccgtgccttc	tagcagcctg	1260
ggcaccacga	cctacatctg	caactgtaac	cacaagccca	gcaacaccaa	ggaggacgag	1320
aaggtggagc	ccaagagctg	cgacaaaact	cacacatgcc	caccgtgccc	agcacctgaa	1380
gctgcagggg	gaccgtcagt	cttcctcttc	cccccaaac	ccaaggacac	cctcatgatc	1440
tcccggaacc	ctgaggtcac	atgcgtgggt	gtggacgtga	gccacgaaga	ccctgaggtc	1500
aagttcaact	ggtacgtgga	cggcgtggag	gtgcataatg	ccaagacaaa	gccgcgggag	1560
gagcagtaca	acagcacgta	ccgtgtggtc	agcgtcctca	ccgtcctgca	ccaggactgg	1620
ctgaatggca	aggagtacaa	gtgcaaggtc	tccaacaaag	ccctcggcgc	ccccatcgag	1680
aaaaccatct	ccaaagccaa	agggcagccc	cgagaaccac	aggtgtacac	cctgccccca	1740
tgccgggatg	agctgaccaa	gaaccaggtc	agcctgtggt	gcctggctaa	aggcttctat	1800
cccagcgaca	tcgccgtgga	gtgggagagc	aatgggcagc	cggagaacaa	ctacaagacc	1860
acgcctcccg	tgtctgactc	cgacggctcc	ttcttctct	acagcaagct	caccgtggac	1920
aagagcaggt	ggcagcaggg	gaacgtcttc	tcatgctccg	tgatgcatga	ggctctgcac	1980
aaccactaca	cgcagaagag	cctctccctg	tctccgggta	aatga		2025

1. A protease-activatable T cell activating bispecific molecule comprising

- (a) a first antigen binding moiety capable of specific binding to CD3;
- (b) a second antigen binding moiety capable of specific binding to a target cell antigen; and
- (c) a masking moiety covalently attached to the T cell bispecific binding molecule through a protease-cleavable linker, wherein the masking moiety is capable of specific binding to the idiotype of the first or the second antigen binding moiety thereby reversibly concealing the first or the second antigen binding moiety.

2. The protease-activatable T cell activating bispecific molecule of claim 1, wherein the masking moiety is:

- (a) covalently attached to the first antigen binding moiety and reversibly conceals the first antigen binding moiety; and/or

(b) an anti-idiotypic scFv.

3. The protease-activatable T cell activating bispecific molecule of claim 2, wherein the masking moiety is covalently attached to the heavy chain variable region of the first antigen binding moiety.

4. (canceled)

5. The protease-activatable T cell activating bispecific molecule of claim 1, wherein:

- (a) the second antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy chain are exchanged; and/or

- (b) the first antigen binding moiety is a conventional Fab molecule.

6. (canceled)

7. The protease-activatable T cell activating bispecific molecule of claim 1, comprising not more than one antigen binding moiety capable of specific binding to CD3.

8. The protease-activatable T cell activating bispecific molecule of claim 1, comprising a third antigen binding moiety, wherein:

- (a) the third antigen binding moiety is a Fab molecule capable of specific binding to a target cell antigen; and/or
- (b) the third antigen binding moiety is identical to the second antigen binding moiety.

9. (canceled)

10. The protease-activatable T cell activating bispecific molecule of claim 1, wherein the second antigen binding moiety is capable of specific binding to a target cell antigen selected from the group consisting of FolR1, HER1, HER2 and Mesothelin.

11. The protease-activatable T cell activating bispecific molecule of claim 1, wherein the first and the second antigen binding moiety are fused to each other via a peptide linker.

12. The protease-activatable T cell activating bispecific molecule of claim 1, wherein:

- (a) the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety;
- (b) the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety; and/or
- (c) the T cell activating bispecific molecule additionally comprises an Fc domain composed of a first and a second subunit capable of stable association.

13-14. (canceled)

15. The protease-activatable T cell activating bispecific molecule of claim 12, wherein the Fc domain is an IgG Fc domain.

16. The protease-activatable T cell activating bispecific molecule of claim 15, wherein the Fc domain exhibits reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain.

17. The protease-activatable T cell activating bispecific molecule of claim 1, wherein the masking moiety comprises a heavy chain variable region comprising:

- (a) a heavy chain complementarity determining region (CDR H) 1 amino acid sequence of SYGVS (SEQ ID NO:26);
- (b) a CDR H2 amino acid sequence of IIWGDGSTNYH-SALIS (SEQ ID NO:27);
- (c) a CDR H3 amino acid sequence of GITTVD-DYYAMDY (SEQ ID NO:28); and a light chain variable region comprising:
- (d) a light chain (CDR L)1 amino acid sequence of RASENIDSYLA (SEQ ID NO:29);
- (e) a CDR L2 amino acid sequence of AATFLAD (SEQ ID NO:30); and
- (f) a CDR L3 amino acid sequence of QHYSTPYT (SEQ ID NO:31).

18. The protease-activatable T cell activating bispecific molecule of claim 1, wherein the protease cleavable linker comprises at least one protease recognition sequence.

19. The protease-activatable T cell activating bispecific molecule of claim 18, wherein the protease cleavable linker comprises the protease recognition sequence RQARVNG (SEQ ID NO:36).

20. The protease-activatable T cell activating bispecific molecule of claim 1, wherein:

- (i) the first antigen binding moiety is capable of specific binding to CD3 and comprises a heavy chain variable region comprising:
 - a) a CDR H1 amino acid sequence of TYAMN (SEQ ID NO:44);
 - b) a CDR H2 amino acid sequence of RIRSKYN-NYATYYADSVKG (SEQ ID NO:45); and
 - c) a CDR H3 amino acid sequence of HGNFNG-SYVSWFAY (SEQ ID NO:46); and a light chain variable region comprising:
 - d) a CDR L1 amino acid sequence of GSST-GAVTTSNYAN (SEQ ID NO:17);
 - e) a CDR L2 amino acid sequence of GTNKRAP (SEQ ID NO:18); and
 - f) a CDR L3 amino acid sequence of ALWYSNLWV (SEQ ID NO:19); and/or
- (ii) the first antigen binding moiety is capable of specific binding to CD3 and comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 43 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 55.

21. (canceled)

22. The protease-activatable T cell activating bispecific molecule of claim 1, wherein:

- (i) the second antigen binding moiety is capable of specific binding to FolR1 and comprises a heavy chain variable region comprising:
 - a) a CDR H1 amino acid sequence of NAWMS (SEQ ID NO:14);
 - b) a CDR H2 amino acid sequence of RIK-SKTDGGTTDYAAPVKG (SEQ ID NO:15); and
 - c) a CDR H3 amino acid sequence of PWEWSWYDY (SEQ ID NO:16); and a light chain variable region comprising:
 - d) a CDR L1 amino acid sequence of GSST-GAVTTSNYAN (SEQ ID NO:17);
 - e) a CDR L2 amino acid sequence of GTNKRAP (SEQ ID NO:18); and
 - f) a CDR L3 amino acid sequence of ALWYSNLWV (SEQ ID NO:19); or
- (ii) the second antigen binding moiety is capable of specific binding to Mesothelin and comprises a heavy chain variable region comprising:
 - a) a CDR H1 amino acid sequence of GYTMN (SEQ ID NO:107);
 - b) a CDR H2 amino acid sequence of LITPYNGAS-SYNQKFRG (SEQ ID NO:108); and
 - c) a CDR H3 amino acid sequence of GGYDGRGFDY (SEQ ID NO:109); and a light chain variable region comprising:
 - d) a CDR L1 amino acid sequence of SASSSVSYM (SEQ ID NO:110);
 - e) a CDR L2 amino acid sequence of DTSKLAS (SEQ ID NO:111); and
 - f) a CDR L3 amino acid sequence of QQWSKHPLT (SEQ ID NO:112).

23. (canceled)

24. An idiotype-specific polypeptide for reversibly concealing an anti-CD3 antigen binding site of a molecule.

25-32. (canceled)

33. A pharmaceutical composition comprising the protease-activatable T cell activating bispecific molecule of claim **1** and a pharmaceutically acceptable carrier.

34-37. (canceled)

38. A method of treating a disease in an individual, comprising administering to said individual a therapeutically effective amount of a composition comprising the protease-activatable T cell activating bispecific molecule of claim **1**.

39. The method of claim **38** for treating or delaying progression of cancer, treating or delaying progression of an immune related disease, or enhancing or stimulating an immune response or function in an individual.

* * * * *