





(22) Date de dépôt/Filing Date: 2019/09/03
(41) Mise à la disp. pub./Open to Public Insp.: 2020/03/04
(45) Date de délivrance/Issue Date: 2023/10/10
(30) Priorités/Priorities: 2018/09/04 (US62/726,562);
2018/09/04 (US62/726,583)

(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
A61K 31/437 (2006.01), *A61K 31/55* (2006.01),
A61P 11/00 (2006.01), *A61P 37/06* (2006.01),
C07D 519/00 (2006.01)
(72) Inventeurs/Inventors:
LONG, DANIEL D., US;
SMITH, CAMERON, US;
THOMPSON, CORBIN, US
(73) Propriétaire/Owner:
THERAVANCE BIOPHARMA R&D IP, LLC, US
(74) Agent: TORYS LLP

(54) Titre : AMIDES DE DIMETHYL AMINO AZETIDINE ET AMIDES HETEROCYCLIQUES A 5 A 7 CHAINONS EN TANT QU'INHIBITEURS DE JAK
(54) Title: DIMETHYL AMINO AZETIDINE AMIDES AND 5 TO 7 MEMBERED HETEROCYCLIC AMIDES AS JAK INHIBITORS



(I)



(IB)

(57) Abrégé/Abstract:

The invention provides compounds of formula (I): (see formula I) and compounds of formula (IB): (see formula IB) where the variables are defined in the specification, or a pharmaceutically-acceptable salt thereof, that are useful as JAK kinase inhibitors. The invention also provides pharmaceutical compositions comprising such compounds and methods of using such compounds to treat respiratory diseases.

**DIMETHYL AMINO AZETIDINE AMIDES AND 5 TO 7 MEMBERED
HETEROCYCLIC AMIDES AS JAK INHIBITORS**

5

10

BACKGROUND OF THE INVENTION

Field of the Invention

The invention is directed to compounds useful as JAK kinase inhibitors. The
15 invention is also directed to pharmaceutical compositions comprising such compounds and
methods of using such compounds to treat respiratory diseases.

State of the Art

Asthma is a chronic disease of the airways for which there are no preventions or
20 cures. The disease is characterized by inflammation, fibrosis, hyper-responsiveness, and
remodeling of the airways, all of which contribute to airflow limitation. An estimated 300
million people worldwide suffer from asthma and it is estimated that the number of people
with asthma will grow by more than 100 million by 2025. In the United States, asthma
afflicts about 6 % to 8 % of the population, making it one of the most common chronic
25 diseases in the country. Although most patients can achieve control of asthma symptoms
with the use of inhaled corticosteroids that may be combined with a leukotriene modifier
and/or a long acting beta agonist, there remains a subset of patients with severe asthma
whose disease is not controlled by conventional therapies. Severe persistent asthma is
defined as disease that remains uncontrolled on high doses of inhaled corticosteroids. While
30 severe asthmatics are estimated to account for approximately 5 % of all asthma sufferers,
they have a high risk of morbidity and mortality and are responsible for a disproportionate

share of health care resource utilization among asthmatics. There remains a need for novel therapies to treat these patients.

Cytokines are intercellular signaling molecules which include chemokines, interferons, interleukins, lymphokines, and tumor necrosis factor. Cytokines are critical for normal cell growth and immunoregulation but also drive immune-mediated diseases and contribute to the growth of malignant cells. Elevated levels of many cytokines have been implicated in the pathology of asthma inflammation. For example, antibody-based therapies targeted at interleukins (IL)-5, and 13 have been shown to provide clinical benefit in subsets of severe asthma patients. Among the cytokines implicated in asthma inflammation, many act through signaling pathways dependent upon the Janus family of tyrosine kinases (JAKs), which signal through the Signal Transducer and Activator of Transcription (STAT) family of transcription factors. Cytokines implicated in asthma inflammation which signal through the JAK-STAT pathway include IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-11, IL-13, IL-23, IL-31, IL-27, thymic stromal lymphopoietin (TSLP), interferon- γ (IFN γ) and granulocyte-macrophage colony-stimulating factor (GM-CSF).

The JAK family comprises four members, JAK1, JAK2, JAK3, and tyrosine kinase 2 (TYK2). Binding of cytokine to a JAK-dependent cytokine receptor induces receptor dimerization which results in phosphorylation of tyrosine residues on the JAK kinase, effecting JAK activation. Phosphorylated JAKs, in turn, bind and phosphorylate various STAT proteins which dimerize, internalize in the cell nucleus and directly modulate gene transcription, leading, among other effects, to the downstream effects associated with inflammatory disease. The JAKs usually associate with cytokine receptors in pairs as homodimers or heterodimers. Specific cytokines are associated with specific JAK pairings. Each of the four members of the JAK family is implicated in the signaling of at least one of the cytokines associated with asthma inflammation. Consequently, a chemical inhibitor with pan-activity against all members of the JAK family could modulate a broad range of pro-inflammatory pathways that contribute to severe asthma.

However, the broad anti-inflammatory effect of such inhibitors could suppress normal immune cell function, potentially leading to increased risk of infection. Evidence of increased infection risk has been observed with the JAK inhibitor tofacitinib, which is dosed orally for the treatment of rheumatoid arthritis. In asthma, inflammation is localized to the


respiratory tract. Inflammation of the airways is characteristic of other respiratory diseases in addition to asthma. Chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, and sarcoidosis are also respiratory tract diseases in which the pathophysiology is believed to be related to JAK-
5 signaling cytokines. Local administration of a JAK inhibitor to the lungs by inhalation offers the potential to be therapeutically efficacious by delivering a potent anti-cytokine agent directly to the site of action, limiting systemic exposure and therefore limiting the potential for adverse systemic immunosuppression. The need remains for a potent JAK inhibitor
10 suitable for local administration to the lungs for treatment of respiratory disease.

JAK-signaling cytokines also play a major role in the activation of T cells, a sub-type of immune cells that is central to many immune processes. Pathological T cell activation is critical in the etiology of multiple respiratory diseases. Autoreactive T cells play a role in bronchiolitis obliterans organizing pneumonia (also termed COS). Similar to COS the
15 etiology of lung transplant rejections is linked to an aberrant T cell activation of the recipient's T cells by the transplanted donor lung. Lung transplant rejections may occur early as Primary Graft Dysfunction (PGD), organizing pneumonia (OP), acute rejection (AR) or lymphocytic bronchiolitis (LB) or they may occur years after lung transplantation as Chronic Lung Allograft Dysfunction (CLAD). CLAD was previously known as bronchiolitis
20 obliterans (BO) but now is considered a syndrome that can have different pathological manifestations including BO, restrictive CLAD (rCLAD or RAS) and neutrophilic allograft dysfunction. Chronic lung allograft dysfunction (CLAD) is a major challenge in long-term management of lung transplant recipients as it causes a transplanted lung to progressively lose functionality (Gauthier et al., *Curr. Transplant. Rep.*, **2016**, 3(3), 185–191). CLAD is
25 poorly responsive to treatment and therefore, there remains a need for effective compounds capable of preventing or treating this condition. Several JAK-dependent cytokines such as IFN γ and IL-5 are up-regulated in CLAD and lung transplant rejection (Berastegui et al, *Clin. Transplant.* **2017**, 31, e12898). Moreover, high lung levels of CXCR3 chemokines such as CXCL9 and CXCL10 which are downstream of JAK-dependent IFN signaling, are linked to
30 worse outcomes in lung transplant patients (Shino et al, *PLOS One*, **2017**, 12 (7), e0180281). Systemic JAK inhibition has been shown to be effective in kidney transplant rejection

The invention also provides a method of treating respiratory disease, in particular, asthma and CLAD, in a mammal, the method comprising administering to the mammal a therapeutically effective amount of a compound, or of a pharmaceutical composition of the invention.

5 The invention also provides a compound of the invention as described herein for use in medical therapy, as well as the use of a compound of the invention in the manufacture of a formulation or medicament for treating respiratory disease in a mammal.

The invention also provides a compound of formula (IB):



10

(IB)

wherein:

A is a 6 to 7 membered monocyclic heterocyclyl ring containing two nitrogen atoms, wherein A is connected to the carbonyl group in (I) through a nitrogen atom, A is substituted with 1 to 3 R² groups and optionally two R² groups form a -(CH₂)- bridged ring with A; or

15 A is pyrrolidinyl, wherein the pyrrolidinyl is connected to the carbonyl group in (I) through a nitrogen atom and wherein the pyrrolidinyl is substituted with NR³R⁴;

R¹ is C₁₋₃ alkyl;

each R² is independently C₁₋₃ alkyl optionally substituted with -OH;

R³ is C₁₋₃alkyl;

20 and R⁴ is C₁₋₃alkyl;

or a pharmaceutically-acceptable salt thereof.

The invention also provides a pharmaceutical composition comprising a compound of the invention and a pharmaceutically-acceptable carrier.

25 The invention also provides a method of treating respiratory disease, in particular, asthma or CLAD, in a mammal, the method comprising administering to the mammal a therapeutically effective amount of a compound, or of a pharmaceutical composition of the invention.


The invention also provides a compound of the invention as described herein for use in medical therapy, as well as the use of a compound of the invention in the manufacture of a formulation or medicament for treating respiratory disease in a mammal.

5

DETAILED DESCRIPTION OF THE INVENTION

In one aspect, the invention provides compounds having activity as a JAK kinase inhibitor. Accordingly, the invention provides a compound of formula (I):

10



wherein:

n is 0,1 or 2;

R^1 is C_{1-3} alkyl; and


15

each R^2 is independently C_{1-3} alkyl;


or a pharmaceutically-acceptable salt thereof.

In some embodiments, the compound has the formula (II):

20




In some embodiments, the compound has the formula (III):



(III)

In some embodiments, R¹ is selected from the group consisting of ethyl, propyl, and isopropyl. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 1 and R² is methyl.


In another embodiment, the invention provides a compound of formula 1:



1


or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula 1:



1


In another embodiment, the invention provides a compound of formula 2



2


or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula 2



2

In another embodiment, the invention provides a compound of formula 3




3

5

or a pharmaceutically-acceptable salt thereof.


In another embodiment, the invention provides a compound of formula 3



3

10


In another embodiment, the invention provides a compound of formula 4



4

or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula 4




8

15

4


In another embodiment, the invention provides a compound of formula 5



5


5 or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula 5



5

In another embodiment, the invention provides a compound of formula 6




10

6

or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula 6




15

6

In another embodiment, the invention provides novel compounds having activity as JAK kinase inhibitors. Accordingly, the invention provides a compound of formula (IB):

9



wherein:

A is a 6 to 7 membered monocyclic heterocyclyl ring containing two nitrogen atoms,
 5 wherein A is connected to the carbonyl group in (I) through a nitrogen atom, A is substituted
 with 1 to 3 R² groups and optionally two R² groups form a -(CH₂)- bridged ring with A; or

A is pyrrolidinyl, wherein the pyrrolidinyl is connected to the carbonyl group in (I)
 through a nitrogen atom and wherein the pyrrolidinyl is substituted with NR³R⁴;

R¹ is C₁₋₃ alkyl;


10 each R² is independently C₁₋₃ alkyl optionally substituted with -OH;

R³ is C₁₋₃alkyl;

and R⁴ is C₁₋₃alkyl;

or a pharmaceutically-acceptable salt thereof.

In some embodiments, compound (IB) has the formula (IB'):




15

In some embodiments, R¹ is selected from the group consisting of methyl, propyl, and
 isopropyl.


In some embodiments, A is selected from the group consisting of piperazinyl, 2,5-
 20 diazabicyclo[2.2.1]heptanyl, and 1,4-diazacycloheptanyl, each of which is substituted with 1
 or 2 R² groups, wherein each R² is independently C₁₋₃ alkyl optionally substituted with -OH,
 or A is pyrrolidinyl substituted with NR³R⁴, wherein R³ is C₁₋₃alkyl and R⁴ is C₁₋₃alkyl.

In some embodiments, A is selected from the group consisting of:




wherein R^2 is C_{1-3} alkyl optionally substituted with $-OH$, R^3 is C_{1-3} alkyl and R^4 is C_{1-3} alkyl.

5 In another embodiment, the invention provides a compound of formula:




or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula:




10 or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula:




or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula:




or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula:




5 or a pharmaceutically-acceptable salt thereof.

In another embodiment, the invention provides a compound of formula:



or a pharmaceutically-acceptable salt thereof.


10 In another embodiment, the disclosure provides a compound of formula (IIB):




wherein:

R¹ is C₁₋₃ alkyl;

15 A¹ is selected from the group consisting of:



and




, wherein R² is C₁₋₃ alkyl optionally

substituted with -OH;

or a pharmaceutically-acceptable salt thereof.

In some embodiments, the compound of Formula (IIB) has the formula (IIB'):


5




(IIB')

In some embodiments, R¹ is selected from the group consisting of methyl and propyl.

In some embodiments, A¹ is selected from the group consisting of:




and




10

In some embodiments, A¹ is selected from the group consisting of:




and



and R¹ is selected the group consisting of methyl and propyl.

In another embodiment, the disclosure provides a compound of formula (IIIB):

15




(IIIB)

wherein:


R¹ is C₁₋₃ alkyl;

A² is selected from the group consisting of:



wherein each R^2 is independently C_{1-3} alkyl optionally substituted with $-OH$, R^3 is C_{1-3} alkyl and R^4 is C_{1-3} alkyl;
or a pharmaceutically-acceptable salt thereof.


5 In some embodiments, the compound of Formula (IIIB) has the formula (IIIB'):




(IIIB')

In some embodiments, R^1 is selected from the group consisting of methyl, isopropyl and propyl.


10 In some embodiments, A^2 is selected from the group consisting of:



In some embodiments, R^1 is selected from the group consisting of methyl, isopropyl and propyl, and A^2 is selected from the group consisting of:



15 In another embodiment, the disclosure provides a compound of formula (IVB):




(IVB)

wherein:


R^1 is C_{1-3} alkyl;

A³ is selected from the group consisting of:



or a pharmaceutically-acceptable salt thereof.


In some embodiments, the compound of Formula (IVB) has the formula (IVB'):




(IVB')

In some embodiments, R¹ is selected from the group consisting of methyl, isopropyl and propyl.

In some embodiments, A³ is selected from the group consisting of:




In some embodiments, R¹ is selected from the group consisting of methyl, isopropyl and propyl, and A³ is selected from the group consisting of:



Chemical structures are named herein according to IUPAC conventions as implemented in ChemDraw software (PerkinElmer, Inc., Cambridge, MA). Compound 1 is designated as (*S*)-(3-(dimethylamino)azetid-1-yl)(2-(6-(2-ethyl-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)methanone.

5 Furthermore, the imidazo portion of the tetrahydroimidazopyridine moiety exists in tautomeric forms, illustrated below for a fragment of the compounds of the disclosure



According to the IUPAC convention, these representations give rise to different numbering of the atoms of the imidazole portion: (1*H*-indazol-3-yl)-4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine (structure A) vs. (1*H*-indazol-3-yl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine (structure B). It will be understood that although structures are shown, or named, in a particular form, the invention also includes the tautomer thereof.

The compounds of the disclosure may contain one or more chiral centers and therefore, such compounds (and intermediates thereof) can exist as racemic mixtures; pure stereoisomers (i.e., enantiomers or diastereomers); stereoisomer-enriched mixtures and the like. Chiral compounds shown or named herein without a defined stereochemistry at a chiral center are intended to include any or all possible stereoisomer variations at the undefined stereocenter unless otherwise indicated. The depiction or naming of a particular stereoisomer means the indicated stereocenter has the designated stereochemistry with the understanding that minor amounts of other stereoisomers may also be present unless otherwise indicated, provided that the utility of the depicted or named compound is not eliminated by the presence of another stereoisomer.

The compounds of the disclosure may also contain several basic groups (e.g., amino groups) and therefore, such compounds can exist as the free base or in various salt forms, such a mono-protonated salt form, a di-protonated salt form, a tri-protonated salt form, or mixtures thereof. All such forms are included within the scope of this invention, unless otherwise indicated.

This invention also includes isotopically-labeled compounds of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) and (IVB'), i.e., compounds of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) and (IVB'), where one or more atom has been replaced or enriched with an atom having the same atomic number but an atomic mass different from the atomic mass that predominates in nature. Examples of isotopes that may be incorporated into a compound of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) and (IVB'), include, but are not limited to, ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , and ^{18}O . Of particular interest are compounds of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) and (IVB'), enriched in tritium or carbon-14, which compounds can be used, for example, in tissue distribution studies. Also of particular interest are compounds of formula (I), (II) and (III) enriched in deuterium especially at a site of metabolism, which compounds are expected to have greater metabolic stability. Additionally of particular interest are compounds of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) and (IVB'), enriched in a positron emitting isotope, such as ^{11}C , ^{15}O and ^{13}N , which compounds can be used, for example, in Positron Emission Tomography (PET) studies.

Definitions

When describing this invention including its various aspects and embodiments, the following terms have the following meanings, unless otherwise indicated.

The term "alkyl" means a monovalent saturated hydrocarbon group which may be linear or branched or combinations thereof. Unless otherwise defined, such alkyl groups typically contain from 1 to 10 carbon atoms. Representative alkyl groups include, by way of example, methyl (Me), ethyl (Et), n-propyl (n-Pr) or (nPr), isopropyl (i-Pr) or (iPr), n-butyl (n-Bu) or (nBu), sec-butyl, isobutyl, tert-butyl (t-Bu) or (tBu), n-pentyl, n-hexyl, 2,2-dimethylpropyl, 2-methylbutyl, 3-methylbutyl, 2-ethylbutyl, 2,2-dimethylpentyl, 2-propylpentyl, and the like.

When a specific number of carbon atoms are intended for a particular term, the number of carbon atoms is shown preceding the term. For example, the term " C_{1-3} alkyl" means an alkyl group having from 1 to 3 carbon atoms wherein the carbon atoms are in any chemically-acceptable configuration, including linear or branched configurations..

The term "cycloalkyl" means a monovalent saturated carbocyclic group which may be monocyclic or multicyclic. Unless otherwise defined, such cycloalkyl groups typically contain from 3 to 10 carbon atoms. Representative cycloalkyl groups include, by way of example, cyclopropyl (cPr), cyclobutyl (cBu), cyclopentyl, cyclohexyl, cycloheptyl,
5 cyclooctyl, adamantyl, and the like.

The term "heterocyclyl", "heterocycle", "heterocyclic", or "heterocyclic ring" means a monovalent saturated or partially unsaturated cyclic non-aromatic group, having from 3 to 10 total ring atoms, wherein the ring contains from 2 to 9 carbon ring atoms and from 1 to 4 ring heteroatoms selected from nitrogen, oxygen, and sulfur. Heterocyclic groups may be
10 monocyclic or multicyclic (i.e., fused or bridged). Representative heterocyclyl groups include, by way of example, pyrrolidinyl, piperidinyl, piperazinyl, imidazolidinyl, morpholinyl, thiomorpholyl, indolin-3-yl, 2-imidazoliny, tetrahydropyranyl, 1,2,3,4-tetrahydroisoquinolin-2-yl, quinuclidinyl, 7-azanorbornanyl, nortropanyl, and the like, where the point of attachment is at any available carbon or nitrogen ring atom. Where the context
15 makes the point of attachment of the heterocyclic group evident, such groups may alternatively be referred to as a non-valent species, i.e. pyrrolidine, piperidine, piperazine, imidazole, tetrahydropyran etc.

The term "halo" means fluoro, chloro, bromo or iodo.

The term "therapeutically effective amount" means an amount sufficient to effect
20 treatment when administered to a patient in need of treatment.

The term "treating" or "treatment" means preventing, ameliorating or suppressing the medical condition, disease or disorder being treated (e.g., a respiratory disease) in a patient (particularly a human); or alleviating the symptoms of the medical condition, disease or disorder.

The term "pharmaceutically acceptable salt" means a salt that is acceptable for
25 administration to a patient or a mammal, such as a human (e.g., salts having acceptable mammalian safety for a given dosage regime). Representative pharmaceutically acceptable salts include salts of acetic, ascorbic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, edisyllic, fumaric, gentisic, gluconic, glucuronic, glutamic, hippuric,
30 hydrobromic, hydrochloric, isethionic, lactic, lactobionic, maleic, malic, mandelic, methanesulfonic, mucic, naphthalenesulfonic, naphthalene-1,5-disulfonic, naphthalene-2,6-

disulfonic, nicotinic, nitric, orotic, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic and xinafoic acid, and the like.


The term "salt thereof" means a compound formed when the hydrogen of an acid is replaced by a cation, such as a metal cation or an organic cation and the like. For example, the cation can be a protonated form of a compound of formula (I), (II) or (III), i.e. a form where one or more amino groups have been protonated by an acid. Typically, the salt is a pharmaceutically acceptable salt, although this is not required for salts of intermediate compounds that are not intended for administration to a patient.

General Synthetic Procedures

Compounds of this invention, and intermediates thereof, can be prepared according to the following general methods and procedures using commercially-available or routinely-prepared starting materials and reagents. The substituents and variables (e.g., A, R¹, R², R³, n, etc.) used in the following schemes have the same meanings as those defined elsewhere herein unless otherwise indicated. Additionally, compounds having an acidic or basic atom or functional group may be used or may be produced as a salt unless otherwise indicated (in some cases, the use of a salt in a particular reaction will require conversion of the salt to a non-salt form, e.g., a free base, using routine procedures before conducting the reaction).


Although a particular embodiment of the present invention may be shown or described in the following procedures, those skilled in the art will recognize that other embodiments or aspects of the present invention can also be prepared using such procedures or by using other methods, reagents, and starting materials known to those skilled in the art. In particular, it will be appreciated that compounds of the invention may be prepared by a variety of process routes in which reactants are combined in different orders to provide different intermediates en route to producing final products.

A general method of preparing certain final compounds of the invention is illustrated in the following Scheme.



Compound **K-18** is reacted with a ketone or aldehyde in the presence of a reducing agent to provide the compound of formula **K-19**. Compound **K-19** is then reacted with a substituted azetidine under typical amide bond formation conditions to give compound (I). Typically, the carboxylic acid is contacted with between about 1 and about 4 equivalents of the substituted azetidine in the presence of an excess of base. The amide bond formation reaction may utilize coupling agents, such as *N,N,N',N'*-tetramethyl-*O*-(7-azabenzotriazol-1-yl)uronium hexafluorophosphate (HATU) or other amide coupling agents known in the art. Hydrazine can be added to cleave undesired byproducts. The reaction is typically conducted at room temperature for between about 5 minutes and about 24 hours or until the reaction is substantially complete.

A general method of preparing certain final compounds of the invention is illustrated in the following Scheme.



Compound **K-18** is reacted with a ketone or aldehyde in the presence of a reducing agent to provide the compound of formula **K-19**. Compound **K-19** is then reacted with amine **A** under typical amide bond formation conditions to give compound (IB). Typically, the carboxylic acid is contacted with between about 1 and about 4 equivalents of amine **A** in the presence of an excess of base. The amide bond formation reaction may utilize coupling agents, such as *N,N,N',N'*-tetramethyl-*O*-(7-azabenzotriazol-1-yl)uronium hexafluorophosphate (HATU) or other amide coupling agents known in the art. Hydrazine can be added to cleave undesired byproducts. The reaction is typically conducted at room temperature for between about 5 minutes and about 24 hours or until the reaction is substantially complete.

Pharmaceutical Compositions

The compounds of the invention and pharmaceutically-acceptable salts thereof are typically used in the form of a pharmaceutical composition or formulation. Such pharmaceutical compositions may advantageously be administered to a patient by inhalation. In addition, pharmaceutical compositions may be administered by any acceptable route of

administration including, but not limited to, oral, rectal, nasal, topical (including transdermal) and parenteral modes of administration.

Accordingly, in one of its compositions aspects, the invention is directed to a pharmaceutical composition comprising a pharmaceutically-acceptable carrier or excipient and a compound of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) or (IVB'), where, as defined above, "compound of formula (I), (II) or (III)" means a compound of formula (I), (II), (III), (IB), (IB'), (IIB), (IIB'), (IIIB), (IIIB'), (IVB) or (IVB'), or a pharmaceutically-acceptable salt thereof. Optionally, such pharmaceutical compositions may contain other therapeutic and/or formulating agents if desired. When discussing compositions and uses thereof, the "compound of the invention" may also be referred to herein as the "active agent". As used herein, the term "compound of the invention" is intended to include all compounds encompassed by formula (I) and (IB) as well as the species embodied in formula (I) and (IB) and pharmaceutically-acceptable salts thereof.

The pharmaceutical compositions of the invention typically contain a therapeutically effective amount of a compound of the present invention. Those skilled in the art will recognize, however, that a pharmaceutical composition may contain more than a therapeutically effective amount, i.e., bulk compositions, or less than a therapeutically effective amount, i.e., individual unit doses designed for multiple administration to achieve a therapeutically effective amount.

Typically, such pharmaceutical compositions will contain from about 0.01 to about 95% by weight of the active agent; including, for example, from about 0.05 to about 30% by weight; and from about 0.1 % to about 10% by weight of the active agent.

Any conventional carrier or excipient may be used in the pharmaceutical compositions of the invention. The choice of a particular carrier or excipient, or combinations of carriers or excipients, will depend on the mode of administration being used to treat a particular patient or type of medical condition or disease state. In this regard, the preparation of a suitable pharmaceutical composition for a particular mode of administration is well within the scope of those skilled in the pharmaceutical arts. Additionally, the carriers or excipients used in the pharmaceutical compositions of this invention are commercially-available. By way of further illustration, conventional formulation techniques are described in Remington: The Science and Practice of Pharmacy, 20th Edition, Lippincott Williams &

White, Baltimore, Maryland (2000); and H.C. Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th Edition, Lippincott Williams & White, Baltimore, Maryland (1999).

Representative examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, the following: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, such as microcrystalline cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances employed in pharmaceutical compositions.

Pharmaceutical compositions are typically prepared by thoroughly and intimately mixing or blending the active agent with a pharmaceutically-acceptable carrier and one or more optional ingredients. The resulting uniformly blended mixture can then be shaped or loaded into tablets, capsules, pills and the like using conventional procedures and equipment.

In one aspect, the pharmaceutical composition is suitable for inhaled administration. Pharmaceutical compositions for inhaled administration are typically in the form of an aerosol or a powder. Such compositions are generally administered using inhaler delivery devices, such as a dry powder inhaler (DPI), a metered-dose inhaler (MDI), a nebulizer inhaler, or a similar delivery device.

In a particular embodiment, the pharmaceutical composition is administered by inhalation using a dry powder inhaler. Such dry powder inhalers typically administer the pharmaceutical composition as a free-flowing powder that is dispersed in a patient's air-stream during inspiration. In order to achieve a free-flowing powder composition, the therapeutic agent is typically formulated with a suitable excipient such as lactose, starch, mannitol, dextrose, polylactic acid (PLA), polylactide-co-glycolide (PLGA) or combinations

thereof. Typically, the therapeutic agent is micronized and combined with a suitable carrier to form a composition suitable for inhalation.

A representative pharmaceutical composition for use in a dry powder inhaler comprises lactose and a compound of the invention in micronized form. Such a dry powder composition can be made, for example, by combining dry milled lactose with the therapeutic agent and then dry blending the components. The composition is then typically loaded into a dry powder dispenser, or into inhalation cartridges or capsules for use with a dry powder delivery device.

Dry powder inhaler delivery devices suitable for administering therapeutic agents by inhalation are described in the art and examples of such devices are commercially available. For example, representative dry powder inhaler delivery devices or products include Aeolizer (Novartis); Airmax (IVAX); ClickHaler (Innovata Biomed); Diskhaler (GlaxoSmithKline); Diskus/Accuhaler (GlaxoSmithKline); Ellipta (GlaxoSmithKline); Easyhaler (Orion Pharma); Eclipse (Aventis); FlowCaps (Hovione); Handihaler (Boehringer Ingelheim); Pulvinal (Chiesi); Rotahaler (GlaxoSmithKline); SkyeHaler/Certihaler (SkyePharma); Twisthaler (Schering-Plough); Turbuhaler (AstraZeneca); Ultrahaler (Aventis); and the like.

In another particular embodiment, the pharmaceutical composition is administered by inhalation using a metered-dose inhaler. Such metered-dose inhalers typically discharge a measured amount of a therapeutic agent using a compressed propellant gas. Accordingly, pharmaceutical compositions administered using a metered-dose inhaler typically comprise a solution or suspension of the therapeutic agent in a liquefied propellant. Any suitable liquefied propellant may be employed including hydrofluoroalkanes (HFAs), such as 1,1,1,2-tetrafluoroethane (HFA 134a) and 1,1,1,2,3,3,3-heptafluoro-*n*-propane, (HFA 227); and chlorofluorocarbons, such as CCl₃F. In a particular embodiment, the propellant is hydrofluoroalkanes. In some embodiments, the hydrofluoroalkane formulation contains a co-solvent, such as ethanol or pentane, and/or a surfactant, such as sorbitan trioleate, oleic acid, lecithin, and glycerin.

A representative pharmaceutical composition for use in a metered-dose inhaler comprises from about 0.01% to about 5% by weight of a compound of the invention; from about 0% to about 20% by weight ethanol; and from about 0% to about 5% by weight surfactant; with the remainder being an HFA propellant. Such compositions are typically

prepared by adding chilled or pressurized hydrofluoroalkane to a suitable container containing the therapeutic agent, ethanol (if present) and the surfactant (if present). To prepare a suspension, the therapeutic agent is micronized and then combined with the propellant. The composition is then loaded into an aerosol canister, which typically forms a portion of a metered-dose inhaler device.

Metered-dose inhaler devices suitable for administering therapeutic agents by inhalation are described in the art and examples of such devices are commercially available. For example, representative metered-dose inhaler devices or products include AeroBid Inhaler System (Forest Pharmaceuticals); Atrovent Inhalation Aerosol (Boehringer Ingelheim); Flovent (GlaxoSmithKline); Maxair Inhaler (3M); Proventil Inhaler (Schering); Serevent Inhalation Aerosol (GlaxoSmithKline); and the like.

In another particular aspect, the pharmaceutical composition is administered by inhalation using a nebulizer inhaler. Such nebulizer devices typically produce a stream of high velocity air that causes the pharmaceutical composition to spray as a mist that is carried into the patient's respiratory tract. Accordingly, when formulated for use in a nebulizer inhaler, the therapeutic agent can be dissolved in a suitable carrier to form a solution. Alternatively, the therapeutic agent can be micronized or nanomilled and combined with a suitable carrier to form a suspension.

A representative pharmaceutical composition for use in a nebulizer inhaler comprises a solution or suspension comprising from about 0.05 µg/mL to about 20 mg/mL of a compound of the invention and excipients compatible with nebulized formulations. In one embodiment, the solution has a pH of about 3 to about 8.

Nebulizer devices suitable for administering therapeutic agents by inhalation are described in the art and examples of such devices are commercially available. For example, representative nebulizer devices or products include the Respimat Softmist Inhaler (Boehringer Ingelheim); the AERx Pulmonary Delivery System (Aradigm Corp.); the PARI LC Plus Reusable Nebulizer (Pari GmbH); and the like.

In yet another aspect, the pharmaceutical compositions of the invention may alternatively be prepared in a dosage form intended for oral administration. Suitable pharmaceutical compositions for oral administration may be in the form of capsules, tablets, pills, lozenges, cachets, dragees, powders, granules; or as a solution or a suspension in an

aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil liquid emulsion; or as an elixir or syrup; and the like; each containing a predetermined amount of a compound of the present invention as an active ingredient.

When intended for oral administration in a solid dosage form, the pharmaceutical compositions of the invention will typically comprise the active agent and one or more pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate. Optionally or alternatively, such solid dosage forms may also comprise: fillers or extenders, binders, humectants, solution retarding agents, absorption accelerators, wetting agents, absorbents, lubricants, coloring agents, and buffering agents. Release agents, wetting agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the pharmaceutical compositions of the invention.

Alternative formulations may also include controlled release formulations, liquid dosage forms for oral administration, transdermal patches, and parenteral formulations. Conventional excipients and methods of preparation of such alternative formulations are described, for example, in the reference by Remington, *supra*.

The following non-limiting examples illustrate representative pharmaceutical compositions of the present invention.

Dry Powder Composition

A micronized compound of formula (I) or (IB) (1 g) is blended with milled lactose (25 g). This blended mixture is then loaded into individual blisters of a peelable blister pack in an amount sufficient to provide between about 0.1 mg to about 4 mg of the compound of formula (I) or (IB) per dose. The contents of the blisters are administered using a dry powder inhaler.

Dry Powder Composition

A micronized compound of formula (I) or (IB) (1 g) is blended with milled lactose (20 g) to form a bulk composition having a weight ratio of compound to milled lactose of 1:20. The blended composition is packed into a dry powder inhalation device capable of delivering between about 0.1 mg to about 4 mg of the compound of formula (I) or (IB) per dose.

Metered-Dose Inhaler Composition

A micronized compound of formula (I) or (IB) (10 g) is dispersed in a solution prepared by dissolving lecithin (0.2 g) in demineralized water (200 mL). The resulting suspension is spray dried and then micronized to form a micronized composition comprising
5 particles having a mean diameter less than about 1.5 μm . The micronized composition is then loaded into metered-dose inhaler cartridges containing pressurized 1,1,1,2-tetrafluoroethane in an amount sufficient to provide about 0.1 mg to about 4 mg of the compound of formula (I) or (IB) per dose when administered by the metered dose inhaler.

Nebulizer Composition

10 A compound of formula (I) or (IB) (25 mg) is dissolved in a solution containing 1.5-2.5 equivalents of hydrochloric acid, followed by addition of sodium hydroxide to adjust the pH to 3.5 to 5.5 and 3% by weight of glycerol. The solution is stirred well until all the components are dissolved. The solution is administered using a nebulizer device that provides about 0.1 mg to about 4 mg of the compound of formula (I) or (IB) per dose.

Utility

15 The JAK inhibitors of the invention have been designed for the treatment of inflammatory and fibrotic disease of the respiratory tract. In particular, the compounds have been designed to enable delivery of a potent anti-cytokine agent directly to the site of action of respiratory disease in the lung while limiting systemic exposure.

20 The compounds of the invention have been shown to be potent inhibitors of the JAK family of enzymes: JAK1, JAK2, JAK3, and TYK2. In addition, the compounds have demonstrated potent inhibition of pro-inflammatory and pro-fibrotic cytokines. It has been recognized that the broad anti-inflammatory effect of JAK inhibitors could suppress normal immune cell function, potentially leading to increased risk of infection. The present
25 compounds have therefore been optimized to limit absorption from the lung into the plasma, thus minimizing the risk of immunosuppression.

30 As described in the experimental section below, the absorption and distribution of compounds 1-6 have been profiled in preclinical assays. Compounds 1-6 and 1b-6b were tested in mice and showed at 5 hours post-dosing high concentration in lung tissue and low absorption into plasma. Compounds of the invention have been shown to inhibit an effect of the pro-inflammatory cytokine IL-13 in mouse lung tissue. Specifically, the compounds have

demonstrated inhibition of IL-13-induced phosphorylation of STAT6 in lung tissue which provides evidence of local lung JAK target engagement *in vivo*. This effect has been observed when the pro-inflammatory cytokine IL-13 is administered 4 hours after administration of the test compound, providing further evidence of significant retention in the lung.

The compounds of the invention have been demonstrated to exhibit both potent inhibitory activity at the cellular level and significant retention in lung tissue. Extensive investigation by the present inventors has determined that while it is possible to identify compounds that are potent at the cellular level or compounds that show significant retention in the lung, it is far more difficult to discover compounds that exhibit both desirable characteristics at the same time.

The anti-inflammatory activity of JAK inhibitors has been robustly demonstrated in preclinical models of asthma (Malaviya et al., *Int. Immunopharmacol.*, **2010**, *10*, 829,-836; Matsunaga et al., *Biochem. and Biophys. Res. Commun.*, **2011**, *404*, 261-267; Kudlacz et al., *Eur. J. Pharmacol.*, **2008**, *582*, 154-161). Cytokines implicated in asthma inflammation which signal through the JAK-STAT pathway include IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-11, IL-13, IL-23, IL-31, IL-27, thymic stromal lymphopoietin (TSLP), interferon- γ (IFN γ) and granulocyte-macrophage colony-stimulating factor (GM-CSF). Accordingly, the compounds of the invention are expected to be useful for the treatment of inflammatory respiratory disorders, in particular, asthma. Inflammation and fibrosis of the lung is characteristic of other respiratory diseases in addition to asthma such as chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, and sarcoidosis. The present compounds, therefore, are also expected to be useful for the treatment of chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, and sarcoidosis.

When compared to their corresponding fluoro analog (compound **C-1** to **C-6** and **C-1B** to **C-6B**), compounds **1-6** and **1B-6B** have been shown to have similar JAK activity. However, they have the advantage of giving rise to significantly less sulfation metabolism, as

demonstrated in Assay 5. This is significant as sulfation metabolism occurs in the lungs, which could lead to a rapid decrease in exposure of the active parent compound.

The compounds of the disclosure have demonstrated inhibition of cytokines associated with inflammation. Therefore, the compounds of the disclosure are likely to be
5 useful for the treatment of certain specific respiratory diseases, as detailed below.

Eosinophilic airway inflammation is a characteristic feature of diseases collectively termed eosinophilic lung diseases (Cottin et al., *Clin. Chest. Med.*, **2016**, 37(3), 535-56). Eosinophilic diseases have been associated with IL-4, IL-13 and IL-5 signaling. Eosinophilic lung diseases include infections (especially helminthic infections), drug-induced
10 pneumonitis (induced for example by therapeutic drugs such as antibiotics, phenytoin, or l-tryptophan), fungal-induced pneumonitis (e.g. allergic bronchopulmonary aspergillosis), hypersensitivity pneumonitis and eosinophilic granulomatosis with polyangiitis (formerly known as Churg-Strauss syndrome). Eosinophilic lung diseases of unknown etiology include idiopathic acute eosinophilic pneumoni, idiopathic chronic eosinophilic pneumonia,
15 hypereosinophilic syndrome, and Löffler syndrome.

A polymorphism in the IL-6 gene has been associated with elevated IL-6 levels and an increased risk of developing pulmonary arterial hypertension (PAH) (Fang et al., *J. Am. Soc. Hypertens.*, **2017**, 11(3), 171-177). Corroborating the role of IL-6 in PAH, inhibition of the IL-6 receptor chain gp130 ameliorated the disease in a rat model of PAH (Huang et al.,
20 *Can. J. Cardiol.*, **2016**, 32(11), 1356.e1-1356.e10).

Cytokines such as IFN γ , IL-12 and IL-6 have been implicated in a range of non-allergic lung diseases such as sarcoidosis, and lymphangioleiomyomatosis (El-Hashemite et al., *Am. J. Respir. Cell. Mol. Biol.*, **2005**, 33, 227-230, and El-Hashemite et al., *Cancer Res.*, **2004**, 64, 3436-3443). The compounds of the invention have also been shown to inhibit IL-6
25 and IFN γ signaling.

Bronchiectasis and infiltrative pulmonary diseases are diseases associated with chronic neutrophilic inflammation.

Pathological T cell activation is critical in the etiology of multiple respiratory diseases. Autoreactive T cells play a role in bronchiolitis obliterans organizing pneumonia
30 (also termed COS). Similar to COS the etiology of lung transplant rejections is linked to an aberrant T cell activation of the recipients T cells by the transplanted donor lung. Lung

transplant rejections may occur early as Primary Graft Dysfunction (PGD), organizing pneumonia (OP), acute rejection (AR) or lymphocytic bronchiolitis (LB) or they may occur years after lung transplantation as Chronic Lung Allograft Dysfunction (CLAD). CLAD was previously known as bronchiolitis obliterans (BO) but now is considered a syndrome that can have different pathological manifestations including BO, restrictive CLAD (rCLAD or RAS) and neutrophilic allograft dysfunction. Chronic lung allograft dysfunction (CLAD) is a major challenge in long-term management of lung transplant recipients as it causes a transplanted lung to progressively lose functionality (Gauthier et al., *Curr Transplant Rep.*, **2016**, 3(3), 185–191). CLAD is poorly responsive to treatment and therefore, there remains a need for effective compounds capable of preventing or treating this condition. Several JAK-dependent cytokines such as IFN γ and IL-5 are up-regulated in CLAD and lung transplant rejection (Berastegui et al, *Clin. Transplant.* **2017**, 31, e12898). Moreover, high lung levels of CXCR3 chemokines such as CXCL9 and CXCL10 which are downstream of JAK-dependent IFN signaling, are linked to worse outcomes in lung transplant patients (Shino et al, *PLOS One*, **2017**, 12 (7), e0180281). Systemic JAK inhibition has been shown to be effective in kidney transplant rejection (Vicenti et al., *American Journal of Transplantation*, **2012**, 12, 2446-56). Therefore, JAK inhibitors have the potential to be effective in treating or preventing lung transplant rejection and CLAD. Similar T cell activation events as described as the basis for lung transplant rejection also are considered the main driver of lung graft-versus-host disease (GVHD) which can occur post hematopoietic stem cell transplants. Similar to CLAD, lung GVHD is a chronic progressive condition with extremely poor outcomes and no treatments are currently approved. A retrospective, multicenter survey study of 95 patients with steroid-refractory acute or chronic GVHD who received the systemic JAK inhibitor ruxolitinib as salvage therapy demonstrated complete or partial response to ruxolitinib in the majority of patients including those with lung GVHD (Zeiser et al, *Leukemia*, **2015**, 29, 10, 2062-68). As systemic JAK inhibition is associated with serious adverse events and a small therapeutic index, the need remains for an inhaled lung-directed, non-systemic JAK inhibitor to prevent and/or treat lung transplant rejection or lung GVHD. The compounds of the invention have the characteristics required to meet this need. More recently, immune-checkpoint inhibitor induced pneumonitis, another T cell mediated lung disease emerged with the increased use of

immune-checkpoint inhibitors. In cancer patients treated with these T cell stimulating agents, fatal pneumonitis can develop.

The mixed lymphocyte reaction assay is an in-vitro assay that mimics transplant rejection. Compound 1 was shown to effectively inhibit IFN γ secretion.

5 In one aspect, therefore, the invention provides a method of treating a respiratory disease in a mammal (e.g., a human), the method comprising administering to the mammal a therapeutically-effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, or of a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a compound of the invention, or a pharmaceutically acceptable salt
10 thereof.

In one aspect, the respiratory disease is asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema,
15 bronchiolitis obliterans, or sarcoidosis. In another aspect, the respiratory disease is asthma or chronic obstructive pulmonary disease.

In one aspect, the respiratory disease is a lung infection, an eosinophilic disease, a helminthic infection, pulmonary arterial hypertension, sarcoidosis, lymphangioliomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced
20 pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome, bronchiolitis obliterans organizing pneumonia, acute and chronic lung transplant rejections (including PGD, OP, LB, AR and CLAD, BO, restrictive
25 CLAD and neutrophilic allograft dysfunction), lung graft-versus-host disease bronchiolitis obliterans organizing pneumonia, pulmonary arterial hypertension, bronchiectasis, or immune-checkpoint-inhibitor induced pneumonitis.

The invention further provides a method of treating asthma in a mammal, the method comprising administering to the mammal a therapeutically-effective amount of a compound
30 of the invention, or a pharmaceutically acceptable salt thereof, or of a pharmaceutical

composition comprising a pharmaceutically-acceptable carrier and a compound of the invention, or a pharmaceutically acceptable salt thereof.

When used to treat asthma, the compounds of the invention will typically be administered in a single daily dose or in multiple doses per day, although other forms of administration may be used. The amount of active agent administered per dose or the total amount administered per day will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

The disclosure further provides a method of treating a respiratory disease (including but not limited to the disease described herein) in a mammal, the method comprising administering to the mammal a therapeutically-effective amount of a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or of a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a compound of the disclosure, or a pharmaceutically acceptable salt thereof.

When used to treat a respiratory disease (including but not limited to the disease described herein), the compounds of the disclosure will typically be administered in a single daily dose or in multiple doses per day, although other forms of administration may be used. The amount of active agent administered per dose or the total amount administered per day will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

As JAK inhibitors, the compounds of the disclosure may also be useful for a variety of other diseases. The compounds of the disclosure may be useful for a variety of gastrointestinal inflammatory indications that include, but are not limited to, inflammatory bowel disease, ulcerative colitis (proctosigmoiditis, pancolitis, ulcerative proctitis and left-sided colitis), Crohn's disease, collagenous colitis, lymphocytic colitis, Behcet's disease, celiac disease, immune checkpoint inhibitor induced colitis, ileitis, eosinophilic esophagitis, graft versus host disease-related colitis, and infectious colitis. Ulcerative colitis (Reimund et al., *J. Clin. Immunology*, 1996, 16, 144-150), Crohn's disease (Woywodt et al., *Eur. J.*

Gastroenterology Hepatology, **1999**, *11*, 267-276), collagenous colitis (Kumawat et al., *Mol. Immunology*, **2013**, *55*, 355-364), lymphocytic colitis (Kumawat et al., **2013**), eosinophilic esophagitis (Weinbrand-Goichberg et al., *Immunol. Res.*, **2013**, *56*, 249-260), graft versus host disease-related colitis (Coghill et al., *Blood*, **2001**, *117*, 3268-3276), infectious colitis (Stallmach et al., *Int. J. Colorectal Dis.*, **2004**, *19*, 308-315), Behcet's disease (Zhou et al., *Autoimmun. Rev.*, **2012**, *11*, 699-704), celiac disease (de Nitto et al., *World J. Gastroenterol.*, **2009**, *15*, 4609-4614), immune checkpoint inhibitor induced colitis (e.g., CTLA-4 inhibitor-induced colitis; (Yano et al., *J. Translation. Med.*, **2014**, *12*, 191), PD-1- or PD-L1-inhibitor-induced colitis), and ileitis (Yamamoto et al., *Dig. Liver Dis.*, **2008**, *40*, 253-259) are characterized by elevation of certain pro-inflammatory cytokine levels. As many pro-inflammatory cytokines signal via JAK activation, compounds described in this application may be able to alleviate the inflammation and provide symptom relief. In particular, the compounds of the disclosure may be useful for the induction and maintenance of remission of ulcerative colitis, and for the treatment of Crohn's disease, immune checkpoint inhibitor induced colitis, and the gastrointestinal adverse effects in graft versus host disease. In one aspect, therefore, the disclosure provides a method of treating a gastrointestinal inflammatory disease in a mammal (e.g., a human), the method comprising administering to the mammal a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a compound of the disclosure, or a pharmaceutically acceptable salt thereof.

Atopic dermatitis and other inflammatory skin diseases have been associated with elevation of proinflammatory cytokines that rely on the JAK-STAT pathway. Therefore, the compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may be beneficial in a number of dermal inflammatory or pruritic conditions that include, but are not limited to atopic dermatitis, alopecia areata, vitiligo, psoriasis, dermatomyositis, cutaneous T cell lymphoma (Netchiporouk et al., *Cell Cycle* **2014**; *13*, 3331-3335) and subtypes (Sezary syndrome, mycosis fungoides, pagetoid reticulosis, granulomatous slack skin, lymphomatoid papulosis, pityriasis lichenoides chronica, pityriasis lichenoides et varioliformis acuta, CD30+ cutaneous T-cell lymphoma, secondary cutaneous CD30+ large cell lymphoma, non-mycosis fungoides CD30- cutaneous large T-cell lymphoma, pleomorphic T-cell lymphoma, Lennert lymphoma, subcutaneous T-cell lymphoma, angiocentric lymphoma, blastic NK-cell

lymphoma), prurigo nodularis, lichen planus, primary localized cutaneous amyloidosis, bullous pemphigoid, skin manifestations of graft versus host disease, pemphigoid, discoid lupus, granuloma annulare, lichen simplex chronicus, vulvar/scrotal/perianal pruritus, lichen sclerosus, post herpetic neuralgia itch, lichen planopilaris, and folliculitis decalvans. In particular, atopic dermatitis (Bao et al., *JAK-STAT*, **2013**, *2*, e24137), alopecia areata (Xing et al., *Nat. Med.* **2014**, *20*, 1043-1049), vitiligo (Craiglow et al, *JAMA Dermatol.* **2015**, *151*, 1110-1112), prurigo nodularis (Sonkoly et al., *J. Allergy Clin. Immunol.* **2006**, *117*, 411-417), lichen planus (Welz-Kubiak et al., *J. Immunol. Res.* **2015**, ID:854747), primary localized cutaneous amyloidosis (Tanaka et al., *Br. J. Dermatol.* **2009**, *161*, 1217-1224), bullous pemphigoid (Feliciani et al., *Int. J. Immunopathol. Pharmacol.* **1999**, *12*, 55-61), and dermal manifestations of graft versus host disease (Okiyama et al., *J. Invest. Dermatol.* **2014**, *134*, 992-1000) are characterized by elevation of certain cytokines that signal via JAK activation. Accordingly, compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may be able to alleviate associated dermal inflammation or pruritus driven by these cytokines. In particular, compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may be expected to be useful for the treatment of atopic dermatitis and other inflammatory skin diseases. In one aspect, therefore, the disclosure provides a method of treating an inflammatory skin disease in a mammal (e.g., a human), the method comprising applying a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof and a pharmaceutical carrier to the skin of the mammal. In one aspect, the inflammatory skin disease is atopic dermatitis.

Many ocular diseases have been shown to be associated with elevations of proinflammatory cytokines that rely on the JAK-STAT pathway. The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, therefore, may be useful for the treatment of a number of ocular diseases that include, but are not limited to, uveitis, diabetic retinopathy, diabetic macular edema, dry eye disease, age-related macular degeneration, and atopic keratoconjunctivitis. In particular, uveitis (Horai and Caspi, *J. Interferon Cytokine Res.*, **2011**, *31*, 733-744), diabetic retinopathy (Abcouwer, *J. Clin. Cell. Immunol.*, **2013**, *Suppl 1*, 1-12), diabetic macular edema (Sohn et al., *American Journal of Ophthalmology*, **2011**, *152*, 686-694), dry eye disease (Stevenson et al, *Arch. Ophthalmol.*, **2012**, *130*, 90-100), and age-related macular degeneration (Knickelbein et al, *Int. Ophthalmol. Clin.*, **2015**,

55(3), 63-78) are characterized by elevation of certain pro-inflammatory cytokines that signal via the JAK-STAT pathway. Accordingly, compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may be able to alleviate the associated ocular inflammation and reverse disease progression or provide symptom relief. In one aspect, therefore, the disclosure provides a method of treating an ocular disease in a mammal, the method comprising administering a pharmaceutical composition comprising a compound of the disclosure or a pharmaceutically-acceptable salt thereof and a pharmaceutical carrier to the eye of the mammal. In one aspect, the ocular disease is uveitis, diabetic retinopathy, diabetic macular edema, dry eye disease, age-related macular degeneration, or atopic keratoconjunctivitis. In one aspect, the method comprises administering the compound of the disclosure, or a pharmaceutically acceptable salt thereof by intravitreal injection. Compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may also be used in combination with one or more compound useful to ocular diseases.

The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may also be useful to treat other diseases such as other inflammatory diseases, autoimmune diseases or cancers. The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may be useful to treat one or more of arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, transplant rejection, xerophthalmia, psoriatic arthritis, diabetes, insulin dependent diabetes, motor neurone disease, myelodysplastic syndrome, pain, sarcopenia, cachexia, septic shock, systemic lupus erythematosus, leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, ankylosing spondylitis, myelofibrosis, B-cell lymphoma, hepatocellular carcinoma, Hodgkins disease, breast cancer, Multiple myeloma, melanoma, non-Hodgkin lymphoma, non-small-cell lung cancer, ovarian clear cell carcinoma, ovary tumor, pancreas tumor, polycythemia vera, Sjogrens syndrome, soft tissue sarcoma, sarcoma, splenomegaly, T-cell lymphoma, and thalassemia major.

Combination therapy

Compounds of the disclosure or a pharmaceutically acceptable salt thereof may be used in combination with one or more agents which act by the same mechanism or by different mechanisms to treat a disease. The different agents may be administered sequentially or simultaneously, in separate compositions or in the same composition. Useful

classes of agents for combination therapy include, but are not limited to, a beta 2 adrenoceptor agonist, a muscarinic receptor antagonist, a glucocorticoid agonist, a G-protein coupled receptor-44 antagonist, a leukotriene D4 antagonist, a muscarinic M3 receptor antagonist, a histamine H1 receptor antagonist, an immunoglobulin E antagonist, a PDE 4 inhibitor, an IL-4 antagonist, a muscarinic M1 receptor antagonist, a histamine receptor antagonist, an IL-13 antagonist, an IL-5 antagonist, a 5-Lipoxygenase inhibitor, a beta adrenoceptor agonist, a CCR3 chemokine antagonist, a CFTR stimulator, an immunoglobulin modulator, an interleukin 33 ligand inhibitor, a PDE 3 inhibitor, a phosphoinositide-3 kinase delta inhibitor, a thromboxane A2 antagonist, an elastase inhibitor, a Kit tyrosine kinase inhibitor, a leukotriene E4 antagonist, a leukotriene antagonist, a PGD2 antagonist, a TNF alpha ligand inhibitor, a TNF binding agent, a complement cascade inhibitor, an eotaxin ligand inhibitor, a glutathione reductase inhibitor, an histamine H4 receptor antagonist, an IL-6 antagonist, an IL2 gene stimulator, an immunoglobulin gamma Fc receptor IIB modulator, an interferon gamma ligand, an interleukin 13 ligand inhibitor, an interleukin 17 ligand inhibitor, a L-Selectin antagonist, a leukocyte elastase inhibitor, a leukotriene C4 antagonist, a Leukotriene C4 synthase inhibitor, a membrane copper amine oxidase inhibitor, a metalloprotease-12 inhibitor, a metalloprotease-9 inhibitor, a mite allergen modulator, a muscarinic receptor modulator, a nicotinic acetylcholine receptor agonist, a nuclear factor kappa B inhibitor, a p-Selectin antagonist, a PDE 5 inhibitor, a PDGF receptor antagonist, a phosphoinositide-3 kinase gamma inhibitor, a TLR-7 agonist, a TNF antagonist, an Abl tyrosine kinase inhibitor, an acetylcholine receptor antagonist, an acidic mammalian chitinase inhibitor, an ACTH receptor agonist, an actin polymerization modulator, an adenosine A1 receptor antagonist, an adenylate cyclase stimulator, an adrenoceptor antagonist, an adrenocorticotrophic hormone ligand, an alcohol dehydrogenase 5 inhibitor, an alpha 1 antitrypsin stimulator, an alpha 1 proteinase inhibitor, an androgen receptor modulator, an angiotensin converting enzyme 2 stimulator, an ANP agonist, a Bcr protein inhibitor, a beta 1 adrenoceptor antagonist, a beta 2 adrenoceptor antagonist, a beta 2 adrenoceptor modulator, a beta amyloid modulator, a BMP10 gene inhibitor, a BMP15 gene inhibitor, a calcium channel inhibitor, a cathepsin G inhibitor, a CCL26 gene inhibitor, a CCR3 chemokine modulator, a CCR4 chemokine antagonist, a cell adhesion molecule inhibitor, a chaperonin stimulator, a chitinase inhibitor, a collagen I antagonist, a complement C3 inhibitor, a CSF-1

antagonist, a CXCR2 chemokine antagonist, a cytokine receptor common beta chain modulator, a cytotoxic T-lymphocyte protein-4 stimulator, a deoxyribonuclease I stimulator, a deoxyribonuclease stimulator, a dipeptidyl peptidase I inhibitor, a DNA gyrase inhibitor, a DP prostanoid receptor modulator, an E-Selectin antagonist, an EGFR family tyrosine kinase receptor inhibitor, an elastin modulator, an Endothelin ET-A antagonist, an Endothelin ET-B antagonist, an epoxide hydrolase inhibitor, a FGF3 receptor antagonist, a Fyn tyrosine kinase inhibitor, a GATA 3 transcription factor inhibitor, a Glucosylceramidase modulator, a Glutamate receptor modulator, a GM-CSF ligand inhibitor, a Guanylate cyclase stimulator, a H⁺ K⁺ ATPase inhibitor, an hemoglobin modulator, an Heparin agonist, an Histone deacetylase inhibitor, an Histone deacetylase-2 stimulator, an HMG CoA reductase inhibitor, an I-kappa B kinase beta inhibitor, an ICAM1 gene inhibitor, an IL-17 antagonist, an IL-17 receptor modulator, an IL-23 antagonist, an IL-4 receptor modulator, an Immunoglobulin G modulator, an Immunoglobulin G1 agonist, an Immunoglobulin G1 modulator, an Immunoglobulin epsilon Fc receptor IA antagonist, an Immunoglobulin gamma Fc receptor IIB antagonist, an Immunoglobulin kappa modulator, an Insulin sensitizer, an Interferon beta ligand, an Interleukin 1 like receptor antagonist, an Interleukin 18 ligand inhibitor, an Interleukin receptor 17A antagonist, an Interleukin-1 beta ligand inhibitor, an Interleukin-5 ligand inhibitor, an Interleukin-6 ligand inhibitor, a KCNA voltage-gated potassium channel-3 inhibitor, a Kit ligand inhibitor, a Laminin-5 agonist, a Leukotriene CysLT1 receptor antagonist, a Leukotriene CysLT2 receptor antagonist, a LOXL2 gene inhibitor, a Lyn tyrosine kinase inhibitor, a MARCKS protein inhibitor, a MDR associated protein 4 inhibitor, a Metalloprotease-2 modulator, a Metalloprotease-9 modulator, a Mineralocorticoid receptor antagonist, a Muscarinic M2 receptor antagonist, a Muscarinic M4 receptor antagonist, a Muscarinic M5 receptor antagonist, a Natriuretic peptide receptor A agonist, a Natural killer cell receptor modulator, a Nicotinic ACh receptor alpha 7 subunit stimulator, a NK cell receptor modulator, a Nuclear factor kappa B modulator, an opioid growth factor receptor agonist, a P-Glycoprotein inhibitor, a P2X3 purinoceptor antagonist, a p38 MAP kinase inhibitor, a Peptidase 1 modulator, a phospholipase A2 inhibitor, a phospholipase C inhibitor, a plasminogen activator inhibitor 1 inhibitor, a platelet activating factor receptor antagonist, a PPAR gamma agonist, a prostacyclin agonist, a protein tyrosine kinase inhibitor, a SH2 domain inositol phosphatase 1 stimulator, a signal transduction

inhibitor, a sodium channel inhibitor, a STAT-3 modulator, a Stem cell antigen-1 inhibitor, a superoxide dismutase modulator, a T cell surface glycoprotein CD28 inhibitor, a T-cell surface glycoprotein CD8 inhibitor, a TGF beta agonist, a TGF beta antagonist, a thromboxane synthetase inhibitor, a thymic stromal lymphoprotein ligand inhibitor, a
5 thymosin agonist, a thymosin beta 4 ligand, a TLR-8 agonist, a TLR-9 agonist, a TLR9 gene stimulator, a Topoisomerase IV inhibitor, a Troponin I fast skeletal muscle stimulator, a Troponin T fast skeletal muscle stimulator, a Type I IL-1 receptor antagonist, a Type II TNF receptor modulator, an ion channel modulator, a uteroglobin stimulator, and a VIP agonist.

Specific agents that may be used in combination with the present JAK inhibitor
10 compounds include, but are not limited to rosiptor acetate, umeclidinium bromide, secukinumab, metenkefalin acetate, tridecactide acetate, fluticasone propionate, alpha-cyclodextrin-stabilized sulforaphane, tezepelumab, mometasone furoate, BI-1467335, dupilumab, acclidinium, formoterol, AZD-1419, HI-1640V, rivipansel, CMP-001, mannitol, ANB-020, omalizumab, tregalizumab, Mitizax, benralizumab, golimumab, roflumilast,
15 imatinib, REGN-3500, masitinib, apremilast, RPL-554, Actimmune, adalimumab, rupatadine, parogrelil, MK-1029, beclometasone dipropionate, formoterol fumarate, mogamulizumab, seratrodist, UCB-4144, nemiralisib, CK-2127107, fevipiprant, danirixin, bosentan, abatacept, EC-18, duvelisib, dociparstat, ciprofloxacin, salbutamol HFA, erdosteine, PrEP-001, nedocromil, CDX-0158, salbutamol, enobosarm, R-TPR-022,
20 lenzilumab, fluticasone furoate, vilanterol trifenate, fluticasone propionate, salmeterol, PT-007, PRS-060, remestemcel-L, citrulline, RPC-4046, nitric oxide, DS-102, gerilimzumab, Actair, fluticasone furoate, umeclidinium, vilanterol, AG-NPP709, Gamunex, infliximab, Ampion, acumapimod, canakinumab, INS-1007, CYP-001, sirukumab, fluticasone propionate, mepolizumab, pitavastatin, solithromycin, etanercept, ivacaftor, anakinra, MPC-
25 300-IV, glycopyrronium bromide, acclidinium bromide, FP-025, risankizumab, glycopyrronium, formoterol fumarate, Adipocell, YPL-001, tiotropium bromide, glycopyrronium bromide, indacaterol maleate, andecaliximab, olodaterol, esomeprazole, dust mite vaccine, mugwort pollen allergen vaccine, vamorolone, gefapixant, revefenacin, gefitinib, ReJoin, tielukast, bedoradrine, SCM-CGH, SHP-652, RNS-60, brodalumab, BIO-
30 11006, umeclidinium bromide, vilanterol trifenate, ipratropium bromide, tralokinumab, PUR-1800, VX-561, VX-371, olopatadine, tulobuterol, formoterol fumarate, triamcinolone

acetonide, reslizumab, salmeterol xinafoate, fluticasone propionate, beclometasone dipropionate, formoterol fumarate, tiotropium bromide, ligelizumab, RUTI, bertilimumab, omalizumab, glycopyrronium bromide, SENS-111, beclomethasone dipropionate, CHF-5992, LT-4001, indacaterol, glycopyrronium bromide, mometasone furoate, fexofenadine, 5 glycopyrronium bromide, azithromycin, AZD-7594, formoterol, CHF-6001, batefenterol, OATD-01, olodaterol, CJM-112, rosiglitazone, salmeterol, setipiprant, inhaled interferon beta, AZD-8871, plecanatide, fluticasone, salmeterol, eicosapentaenoic acid monoglycerides, lebrikizumab, RG-6149, QBKPN, Mometasone, indacaterol, AZD-9898, sodium pyruvate, zileuton, CG-201, imidafenacin, CNTO-6785, CLBS-03, mometasone, RGN-137, procaterol, 10 formoterol, CCI-15106, POL-6014, indacaterol, beclomethasone, MV-130, GC-1112, Allergovac depot, MEDI-3506, QBW-251, ZPL-389, udenafil, GSK-3772847, levocetirizine, AXP-1275, ADC-3680, timapiprant, abediterol, AZD-7594, ipratropium bromide, salbutamol sulfate, tadekinig alfa, ACT-774312, dornase alfa, iloprost, batefenterol, fluticasone furoate, alicaforsen, ciclesonide, emeramide, arformoterol, SB-010, 15 Ozagrel, BTT-1023, Dectrekumab, levalbuterol, pranlukast, hyaluronic acid, GSK-2292767, Formoterol, NOV-14, Lucinactant, salbutamol, prednisolone, ebastine, dexamethasone cipeclilate, GSK-2586881, BI-443651, GSK-2256294, VR-179, VR-096, hdm-ASIT+, budesonide, GSK-2245035, VTX-1463, Emedastine, dexpramipexole, levalbuterol, N-6022, dexamethasone sodium phosphate, PIN-201104, OPK-0018, TEV-48107, suplatast, BI- 20 1060469, Gemilukast, interferon gamma, dalazatide, bilastine, fluticasone propionate, salmeterol xinafoate, RP-3128, bencycloquidium bromide, reslizumab, PBF-680, CRTH2 antagonist, Pranlukast, salmeterol xinafoate, fluticasone propionate, tiotropium bromide monohydrate, masilukast, RG-7990, Doxofylline, abediterol, glycopyrronium bromide, TEV-46017, ASM-024, fluticasone propionate, glycopyrronium bromide, salmeterol xinafoate, 25 salbutamol, TA-270, Flunisolide, sodium chromoglycate, Epsi-gam, ZPL-521, salbutamol, aviptadil, TRN-157, Zafirlukast, Stempeucel, pemirolast sodium, nadolol, fluticasone propionate + salmeterol xinafoate, RV-1729, salbutamol sulfate, carbon dioxide + perfluorooctyl bromide, APL-1, dectrekumab + VAK-694, lysine acetylsalicylate, zileuton, TR-4, human allogenic adipose-derived mesenchymal progenitor cell therapy, MEDI-9314, 30 PL-3994, HMP-301, TD-5471, NKTT-120, pemirolast, beclomethasone dipropionate, trantinterol, monosodium alpha luminol, IMD-1041, AM-211, TBS-5, ARRY-502,

seratrodast, recombinant midismase, ASM-8, deflazacort, bambuterol, RBx-10017609, ipratropium + fenoterol, fluticasone + formoterol, epinastine, WIN-901X, VALERGEN-DS, OligoG-COPD-5/20, tulobuterol, oxis Turbuhaler, DSP-3025, ASM-024, mizolastine, budesonide + salmeterol, LH-011, AXP-E, histamine human immunoglobulin, YHD-001, theophylline, ambroxol + erdosteine, ramatroban, montelukast, pranlukast, AG-1321001, tulobuterol, ipratropium + salbutamol, tranilast, methylprednisolone suleptanate, colforsin daropate, repirinast, and doxofylline.

Also provided, herein, is a pharmaceutical composition comprising a compound of the disclosure or a pharmaceutically acceptable salt thereof and one or more other therapeutic agents. The therapeutic agent may be selected from the class of agents specified above and from the list of specific agent described above. In some embodiments, the pharmaceutical composition is suitable for delivery to the lungs. In some embodiments, the pharmaceutical composition is suitable for inhaled or nebulized administration. In some embodiments, the pharmaceutical composition is a dry powder or a liquid composition.

Further, in a method aspect, the disclosure provides a method of treating a disease or disorder in a mammal comprising administering to the mammal a compound of the disclosure or a pharmaceutically acceptable salt thereof and one or more other therapeutic agents.

When used in combination therapy, the agents may be formulated in a single pharmaceutical composition, or the agents may be provided in separate compositions that are administered simultaneously or at separate times, by the same or by different routes of administration. Such compositions can be packaged separately or may be packaged together as a kit. The two or more therapeutic agents in the kit may be administered by the same route of administration or by different routes of administration.

EXAMPLES

The following synthetic and biological examples are offered to illustrate the invention, and are not to be construed in any way as limiting the scope of the invention. In the examples below, the following abbreviations have the following meanings unless otherwise indicated. Abbreviations not defined below have their generally accepted meanings.

	ACN	=	acetonitrile
	DCM	=	dichloromethane
	DIPEA	=	<i>N,N</i> -diisopropylethylamine
	DMF	=	<i>N,N</i> -dimethylformamide
5	EtOAc	=	ethyl acetate
	h	=	hour(s)
	HATU	=	<i>N,N,N',N'</i> -tetramethyl- <i>O</i> -(7-azabenzotriazol-1-yl)uronium hexafluorophosphate
	IPA	=	isopropyl alcohol
10	IPAc	=	isopropylacetate
	MeOH	=	methanol
	min	=	minute(s)
	Pd(PPh ₃) ₄	=	tetrakis(triphenylphosphine)palladium(0)
	RT	=	room temperature
15	TFA	=	trifluoroacetic acid
	THF	=	tetrahydrofuran
	bis(pinacolato)diboron	=	4,4,5,5,4',4',5',5'-octamethyl- [2,2']bi[[1,3,2]dioxaborolanyl]

Reagents and solvents were purchased from commercial suppliers (Aldrich, Fluka,
 20 Sigma, etc.), and used without further purification. Progress of reaction mixtures was
 monitored by thin layer chromatography (TLC), analytical high performance liquid
 chromatography (anal. HPLC), and mass spectrometry. Reaction mixtures were worked up
 as described specifically in each reaction; commonly they were purified by extraction and
 other purification methods such as temperature-, and solvent-dependent crystallization, and
 25 precipitation. In addition, reaction mixtures were routinely purified by column
 chromatography or by preparative HPLC, typically using C18 or BDS column packings and
 conventional eluents. Typical preparative HPLC conditions are described below.

Characterization of reaction products was routinely carried out by mass and ¹H-NMR
 spectrometry. For NMR analysis, samples were dissolved in deuterated solvent (such as
 30 CD₃OD, CDCl₃, or *d*₆-DMSO), and ¹H-NMR spectra were acquired with a Varian Gemini
 2000 instrument (400 MHz) under standard observation conditions. Mass spectrometric

(a) 1-(benzyloxy)-3-ethylbenzene (**I-2**)

To a stirred solution of 3-ethylphenol (**I-1**) (25.0 g, 204.0 mmol) in ACN (250 mL, 10 vol) was added potassium carbonate (42.0 g, 306 mmol) at room temperature. The resulting reaction mass was stirred at room temperature for 15 minutes, followed by the addition of benzyl bromide (24.0 mL, 204 mmol) in drop wise manner. The resulting reaction mixture was stirred for 6 hours at room temperature. After completion of the reaction (TLC monitoring), the resulting reaction mass was poured into water (1.0 L) followed by the extraction of compound with EtOAc (2 x 2L). The combined organics were washed with cold water, brine solution and dried over sodium sulfate, filtered and evaporated under reduced pressure. The crude product was then purified by column chromatography over silica gel (100-200M) by using eluents 2% EtOAc in hexane to get the desired product (**I-2**) as light yellow oily compound (35.0 g, 81%). ¹H NMR (400 MHz, chloroform-d) δ 7.46-7.44 (m, 2H), 7.39 (t, J = 7.6 Hz, 2H), 7.34-7.31 (m, 1H), 7.21 (t, J = 7.6 Hz), 6.86-6.80 (m, 3H), 5.07 (s, 2H), 2.64 (q, J = 7.6 Hz, 2H), 1.24 (t, J = 7.6 Hz, 3H).

(b) 4-(benzyloxy)-1-bromo-2-ethylbenzene (**I-3**)

To an ice cold stirred solution of 1-(benzyloxy)-3-ethylbenzene (**I-2**) (35.0 g, 164 mmol) in ACN (525 mL, 15 vol) was added N-bromosuccinimide (32.0 g 181 mmol) in portions over a period of 15 minutes. The resulting reaction mixture was stirred for next 1 hour at room temperature. After completion of reaction (TLC monitoring), the resulting reaction mass was poured into ice cold water (1.50 L) followed by the extraction of compound with EtOAc (2 x 1L). The combined organics were washed with water and dried over sodium sulfate, filtered and evaporated under reduced pressure to obtain the crude product. n-Hexane (250 mL) was added to the crude material, resulting in a slurry, followed by filtration through a sintered funnel. Mother liquor was evaporated under reduced pressure to obtain the desired product **I-3** as light yellow oily compound (42.0 g, 87%). ¹H NMR (400 MHz, chloroform-d) δ 7.52-7.29 (m, 7H), 6.88 (s, 1H), 6.68 (d, J = 6.0 Hz, 1H), 5.04 (s, 2H), 2.69 (q, J = 7.6 Hz, 2H), 1.20 (t, J = 7.5 Hz, 3H).

(c) 2-(4-(benzyloxy)-2-ethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**I-4**)


A stirred solution of 4-(benzyloxy)-1-bromo-2-ethylbenzene (**I-3**) (42.0 g, 144 mmol), bis(pinacolato) diboron (44.0 g, 173 mmol), and potassium acetate (28 g, 288 mmol) in dioxane (440 mL) was degassed by purging N₂ (g) for 15 min followed by addition of

PdCl₂(dppf).DCM complex (11.0 g, 15 mmol). The resulting reaction mixture was heated up to 80°C for next 16h. After completion of the reaction (TLC monitoring), the reaction mass was filtered through celite bed and mother liquor was evaporated under reduced pressure to obtain the crude product. Crude residue was purified by column chromatography over silica gel (100-200M) by using eluents 1% EtOAc in hexane to get the desired product (I-4) as light yellow oily compound (32.0 g, 66%). ¹H NMR (400 MHz, chloroform-d) δ 7.74 (d, J = 8.4 Hz, 1H), 7.45-7.36 (m, 5H), 6.84-6.78 (m, 2H), 5.08 (s, 2H), 2.91 (q, J = 7.6 Hz), 1.33 (s, 12H), 1.19 (t, J = 7.6 Hz, 3H).

(d) (4-(benzyloxy)-2-ethylphenyl)trifluoro-λ⁴-borane, potassium salt (I-5)

To a stirred solution of compound 2-(4-(benzyloxy)-2-ethylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (I-4) (20 g, 59.0 mmol), in acetone:methanol (200 mL, 1:1 ratio, 10 vol), was added a 3M solution of potassium hydrogen fluoride (23.0 g, 295 mmol, dissolved in 98.0 mL of water). The resulting reaction mixture was stirred at room temperature for 16 hours. After completion of the reaction (TLC monitoring), the resulting reaction mass was evaporated under reduced pressure. The solid thus obtained was taken up in water (100 mL) and stirred at room temperature for 30 min. The resulting reaction mass was filtered through a sintered funnel, washed with n-hexane and dried under reduced pressure to provide the desired product (I-5) as a white solid (14.0 g, 74%). ¹H NMR (400 MHz, chloroform-d) δ 7.43 (d, J = 7.2 Hz, 2H), 7.37 (t, J = 7.5 Hz, 2H), 7.30 (t, J = 7.1 Hz, 1H), 7.22 (d, J = 8.0 Hz), 6.58 (s, 1H), 6.53 (d, J = 7.9 Hz, 1H), 5.00 (s, 2H), 2.65 (q, J = 7.5 Hz, 2H), 1.07 (t, J = 7.4 Hz, 3H).

Preparation 2: 6-benzyl 5-(tert-butyl) (S)-3-benzyl-3,4,6,7-tetrahydro-5H-imidazo[4,5-c]pyridine-5,6-dicarboxylate (I-11)



25

(a) (*S*)-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid, hydrochloride salt (**I-7**)

To an ice cold stirred suspension of L-histidine (**I-6**) (5.0 kg, 32.14mol) in water (40 L, 8 vol.) was added concentrated hydrochloric acid (3.93 L, 33.75 mol), followed by the addition of formaldehyde (5.50 L, 67.5 mol, 37% aq. solution) in drop wise manner. The resulting solution was stirred for 30 minute at same temperature and then heated at 80°C for 8 hours. Reaction progress was monitored by LCMS. Water was removed under reduced pressure to obtain the crude product, and the resulting crude was stirred for 2 hours in Toluene (20 L). Organics were removed under reduced pressure to remove excess water and the compound was azeotropically dried. The resulting material was then taken in diethyl ether (20 L) and stirred for 2 hours. The solid material was then filtered and air dried to obtain the desired product (**I-7**) as an off-white solid (6.50 Kg, 85%). ¹H NMR (400 MHz, D2O) δ 8.69 (s, 1H), 4.56 (d, J = 15.4 Hz, 1H), 4.42 (d, J = 15.5 Hz, 1H), 4.20 (dd, J = 5.5, 5.2 Hz, 1H), 3.42 (dd, J = 5.0, 17.0 Hz, 1H), 3.11 (dd, J = 10.2, 16.8 Hz, 1H).

(b) (*S*)-3,5-bis(*tert*-butoxycarbonyl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid (**I-8**)

To an ice cold stirred solution of (*S*)-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid di-hydrochloride (**I-7**) (6.10 Kg, 25.40mol) in 1,4-dioxane (48 L, 8 vol) and water (48 L, 8 vol) was added triethylamine (12.36 L, 89 mol) drop wise followed by the addition of di-*tert*-butyl dicarbonate (18.07 L, 78.74 mol, dissolved in 5 L of 1,4-dioxane) over a period of 30 min. The resulting reaction mixture was stirred at room temperature for next 16 hours. After completion of reaction (TLC & LCMS monitoring), the yellowish reaction mixture was diluted with water (10 L) and washed successively with diethyl ether (2 x 10 L) and EtOAc (2 x 7.50 L). The organic phase was discarded. The aqueous layer was cooled and brought to pH ~3 with 6N HCl solution; the aqueous phase was extracted with EtOAc (3 x 10 L). The combined organics were washed with brine solution, dried over sodium sulfate, and concentrated under reduced pressure. The oily residue was crystallized from 30% EtOAc:Hexanes to afford the desired product (**I-8**) as off-white solid (5.1 Kg, 55%). (m/z): [M+H]⁺ calcd for C₁₇H₂₅N₃O₆ 368.18 found 368.21.

(c) 6-benzyl 3,5-di-*tert*-butyl (*S*)-6,7-dihydro-3H-imidazo[4,5-*c*]pyridine-3,5,6(4H)-tricarboxylate (**I-9**)

To an ice cold solution of (*S*)-3,5-bis(*tert*-butoxycarbonyl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid (**I-8**) (5.1 Kg, 13.88 mol) in DCM (51 L, 10 vol) was added sequentially saturated aqueous sodium bicarbonate (41.0 L, 8 vol), tetra-butyl ammonium iodide (5.13 Kg, 13.88 mol) and benzyl bromide (2.47 L, 20.82 mol). The resulting reaction mixture was stirred at room temperature for next 16 hours. After completion of the reaction (TLC & LCMS monitoring), the biphasic solution was separated. The aqueous layer was extracted with DCM (3 x 10 L). The combined organics were washed with brine solution, dried over sodium sulfate, filtered, and concentrated under reduced pressure to obtain the crude product, which was purified by column chromatography through silica gel (100-200M) by using eluents 40% EtOAc in hexane to get the desired product (**I-9**) as viscous oil (4.50 Kg, 72%). (m/z): [M+H]⁺ calcd for C₂₄H₃₁N₃O₆ 458.22 found 458.60.

(d) 6-benzyl 5-(*tert*-butyl) (*S*)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-10**)


To an ice cold solution of 6-benzyl 3,5-di-*tert*-butyl (*S*)-6,7-dihydro-3H-imidazo[4,5-*c*]pyridine-3,5,6(4H)-tricarboxylate (**I-9**) (4.50 Kg, 9.84 mol) in IPA (45 L, 10 vol) was added ammonium hydroxide (36 L, 8 vol) drop wise. The resulting reaction mixture was further stirred at room temperature for the next 16 hours. After completion of the reaction (TLC & LCMS monitoring), the resulting mixture was diluted with water (25 L) followed by extraction with EtOAc (3 x 20 L). The combined organics were washed with brine solution, dried over sodium sulfate, filtered and concentrated under reduced pressure to yield the crude product which was purified by column chromatography through silica gel (100-200M) by using eluents 2% MeOH in DCM to obtain the desired product (**I-10**) as a thick viscous oil (2.70 Kg, 77%). (m/z): [M+H]⁺ calcd for C₁₉H₂₃N₃O₄ 358.17 found 358.33.

(e) 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-11**)

To an ice cold solution of 6-benzyl 5-(*tert*-butyl) (*S*)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-10**) (2.70 kg, 7.55 mol) in DCM (32.4 L, 12 vol) was added aqueous 1N sodium hydroxide (24.3 L, 9 vol) followed by the sequential addition of tetra-butyl ammonium iodide (2.80 Kg, 7.55 mol) and benzyl bromide (0.99 L, 8.31 mol). The resulting reaction mixture was stirred at room temperature for next 2 hours. After completion of the reaction (TLC & LCMS monitoring), the biphasic solution was separated.

The aqueous layer was extracted with DCM (3 x 10 L). The combined organics were washed with brine solution, dried over sodium sulfate, filtered, and concentrated under reduced pressure to yield the crude product which was purified by column chromatography over silica gel (100-200M) by using eluents 40% EtOAc in hexane to obtain the desired product (I-11) as a viscous oil (1.70 Kg, 63%). (m/z): [M+H]⁺ calcd for C₂₆H₂₉N₃O₄ 448.22 found 448.20.

Preparation 3: 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(6-(4-(benzyloxy)-2-ethylphenyl)-1H-indazol-3-yl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (I-16)



10

(a) 4-bromo-2-fluorobenzoyl chloride (I-13)

To an ice cold stirred solution of 4-bromo-2-fluorobenzoic acid (I-12) (1.25 Kg, 5.71 mol) in DCM (12.5 L, 15 vol), was added oxalyl chloride (0.98 L, 11.42 mol) in a drop wise manner. The resulting reaction mixture was stirred for 10 min at the same temperature. DMF (150 mL) was then added in a drop wise manner to the reaction mixture. The resulting reaction mass was allowed to warm to room temperature and stirred for 2 hours. After completion of the reaction (by TLC monitoring), excess oxalyl chloride was removed under reduced pressure under a nitrogen atmosphere to obtain the crude product (I-13) (1.08 Kg, 80%), which was used in the next step without further purification.

20

(b) 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(4-bromo-2-fluorobenzoyl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (I-14)

To a stirred solution of 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-11**) (1.70 Kg, 3.80 mol) in ACN (13.6 L, 8 vol) was added triethylamine (2.11 L, 15.2 mol) followed by the addition of 4-bromo-2-fluorobenzoyl chloride (**I-13**) (1.08 Kg, 4.56 mol in 3.4 L ACN, 2 vol) at room temperature. After completion of addition, the resulting reaction mixture color turned brown from light yellow. The resulting reaction mixture was stirred at same temperature for 30 min, and reaction progress was monitored by TLC. The resulting reaction mixture was quenched with ice cold water (10 L), followed by extraction with EtOAc (3 x 5 L) and combined organics were washed with brine solution. The organics were dried over sodium sulfate, filtered, and concentrated under reduced pressure to yield the crude product which was purified by column chromatography over silica gel (100-200M) by using eluents 20% EtOAc in hexane to obtain the desired product (**I-14**) (1.74 Kg, 71%). (m/z): [M+H]⁺ calcd for C₃₃H₃₁BrFN₃O₅ 648.14 found 648.20.

(c) 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(6-bromo-1H-indazol-3-yl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-15**)

To a stirred solution of 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(4-bromo-2-fluorobenzoyl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-14**) (1.74 Kg, 2.68 mol) in THF (26.0 L, 15 vol) was added hydrazine hydrate (0.705 L, 13.4 mol) at room temperature. The resulting reaction mixture was heated at 60°C for 3 hours. After completion of the reaction (TLC monitoring), the resulting reaction mass was poured into ice cold water (10 L) followed by extraction of compound with EtOAc (3 x 10L). The combined organics were washed with brine and dried over sodium sulfate, filtered, and evaporated under reduced pressure to yield the crude product, which was purified by column chromatography over silica gel (100-200M) by using eluents 20% EtOAc in hexane to obtain the desired product (**I-15**) as an off-white solid (1.12 Kg, 65%). (m/z): [M+H]⁺ calcd for C₃₃H₃₂BrN₅O₄ 642.16 found 642.21.

(d) 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(6-(4-(benzyloxy)-2-ethylphenyl)-1H-indazol-3-yl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-16**)


Bis(pinacolato)diboron (250 g, 984 mmol) was charged to a 5 L 3-neck single walled flask previously etched using fluoride chemistry, along with propan-2-ol (1882 mL,

2.46E+04 mmol) and the mixture was stirred until fully dissolved. Dissolution was endothermic (-4 °C). In a 4 L Erlenmeyer flask, previously etched using fluoride chemistry, potassium fluoride hydrofluoride (538 g, 6891 mmol) was dissolved in water (2.306 L, 1.28E+05 mmol) to form a 3M solution. The dissolution was endothermic (-12 °C). The solution was then filtered to remove a small amount of insoluble material from the potassium fluoride hydrofluoride. Once both solutions were fully dissolved, the contents of the Erlenmeyer flask were charged into the single walled flask portion-wise over 15 minutes. A moderate exotherm was observed (+10 °C). The solution became a thick and translucent semi-opaque gray slurry during the addition and stirring was increased to keep the contents well mixed. The mixture was stirred for 1.5 h, and then filtered through a coarse glass fritted funnel (4 L, previously etched). The filtration required 30-45 minutes to complete. The clear biphasic filtrate was discarded. The white solids were dried for 10 minutes on the filter (cracking of the cake was observed). The solids were transferred back into a cleaned 5 L 3-neck single walled flask and re-slurried with water (1.33 L, 7.38E+04 mmol). The slurry was stirred for 2 h after which time it formed a clear homogenous hydrogel. The solution was stirred for another 1 h whereupon the solids/gel were filtered out using a 4 L coarse glass funnel (previously etched). The solids were allowed to dry on the filter for 30 minutes. The solids were transferred back to a cleaned 5 L 3-neck single walled flask and reslurried in acetone (1.084 L, 1.48E+04 mmol). The white/gray slurry was stirred for 1 h and was then filtered on a 4 L coarse glass funnel (previously etched). The filtration required 20 minutes to complete, and was then dried on the funnel for another 1 h. During this time, the solids were occasionally agitated to ensure homogenous drying. A light white powder remained after drying on the filter. The solids were dried for 20 h at 55 °C under vacuum with a slow nitrogen bleed to afford a fluffy white solid (200.3 g were collected).

To a stirred solution of 6-benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(6-bromo-1H-indazol-3-yl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (**I-15**) (10.0 g, 16.0 mmol) in 2-methyl tetrahydrofuran (100 mL, 10 vol) was added (4-(benzyloxy)-2-ethylphenyl)trifluoro- λ^4 -borane, potassium salt (**I-5**) (8.0 g, 20 mmol) and the fluffy white solid obtained above (0.20 g). The resulting reaction mixture was degassed with nitrogen gas for 30 minutes. To this solution, a prepared aqueous solution of cesium carbonate (20.0 g, 62.0 mmol in 60 mL water, 6 vol) was added. The resulting reaction mixture was further

degassed for 15 minutes followed by addition of bis(*di-tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (0.66 g, 0.93 mmol), and the reaction mixture was evacuated under vacuum and flushed by nitrogen. The resulting reaction mixture was heated at 110°C for 20 hours. After completion of the reaction (TLC & LCMS monitoring), the resulting reaction mixture was cooled to room temperature and filtered through a celite bed, then further washed with EtOAc (3 x 0.5 L). The combined organics were washed with 1N sodium hydroxide solution (3 x 0.5 L). The combined organics were then washed with brine and dried over sodium sulfate, filtered, and evaporated under reduced pressure to yield the crude product, which was purified by column chromatography over silica gel (100-200M) by using eluents 20% EtOAc in hexane to obtain the desired product (I-16) (as mixture of *N*-benzyl regioisomers) as light yellow solid (8.0 g, 66%). (m/z): [M+H]⁺ calcd for C₄₈H₄₇N₅O₅ 774.36 found 774.59.

Preparation 4: (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, hydrochloride (I-18)




(a) benzyl (*S*)-3-benzyl-2-(6-(4-(benzyloxy)-2-ethylphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylate, hydrochloride (I-17)

20 6-Benzyl 5-(*tert*-butyl) (*S*)-3-benzyl-2-(6-(4-(benzyloxy)-2-ethylphenyl)-1H-indazol-3-yl)-3,4,6,7-tetrahydro-5H-imidazo[4,5-c]pyridine-5,6-dicarboxylate (I-16) (1.0 g, 1.292 mmol) was dissolved in dioxane (8 mL) and water (1.5 mL), then hydrogen chloride solution, 4 M in dioxane (7 mL, 28.0 mmol) was added and the reaction mixture was stirred at room temperature for 3 hours (reaction progress monitored by LCMS). The reaction mixture was then frozen and lyophilized, and the crude product (I-17) was used directly in the next reaction (quantitative yield was assumed). (m/z): [M+H]⁺ calcd for C₄₃H₃₉N₅O₃ 674.31 found 674.3.

(b) (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, hydrochloride (I-18)

Benzyl (*S*)-3-benzyl-2-(6-(4-(benzyloxy)-2-ethylphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylate, hydrochloride (**I-17**) (0.918 g, 1.292 mmol) was dissolved in 2-propanol (15 mL), hydrogen chloride solution, 5 M in water (0.258 mL, 1.292 mmol), and water (0.25 mL) at 50 °C, then palladium, 10% wt. on carbon, 50%
5 water (0.138 g, 0.065 mmol) was added. The reaction flask was then purged with nitrogen, a hydrogen balloon was attached, and the reaction mixture was stirred at 50 °C for 4 days with the hydrogen balloon being replenished as needed (reaction progress monitored by LCMS). All solids were then removed by filtration and the resulting solution was concentrated. The residue was dissolved in 1:1 ACN/Water, frozen, and lyophilized. The resulting powder (**I-18**) was used without further purification (quantitative yield was assumed). (*m/z*): [M+H]⁺
10 calcd for C₂₂H₂₁N₅O₃ 404.17 found 404.2.

Preparation 5: (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid (I-19**)**




15

I-19

(*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid, HCl (**I-18**) (0.25 g, 0.568 mmol) was suspended in DMF (2.5 mL) and acetone (2.5 mL), then acetic acid (0.098 mL, 1.705 mmol) and sodium cyanoborohydride (0.179 g, 2.84 mmol) were added and the reaction mixture was stirred at
20 room temperature for 24 hours (reaction progress was monitored by LCMS). The reaction mixture was concentrated, then the crude product was purified by reverse phase chromatography (5-70% ACN/Water gradient, 50 g C18aq column) to provide the TFA salt of the title compound (149 mg, 47% yield). (*m/z*): [M+H]⁺ calcd for C₂₅H₂₇N₅O₃ 446.21 found 446.3.

Example 1: (S)-(3-(dimethylamino)azetid-1-yl)(2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone




5

1

(S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (**I-19**) (50 mg, 0.089 mmol), 3-(Dimethylamino)azetidine dihydrochloride (23.20 mg, 0.134 mmol), and DIPEA (0.078 mL, 0.447 mmol) were dissolved in DMF (1.5 mL), then HATU (51.0 mg, 0.134 mmol) was added and the reaction mixture was stirred at room temperature for 3 hours (reaction progress was monitored by LCMS). Hydrazine (0.014 mL, 0.447 mmol) was added to cleave undesired byproducts, and the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (25 mg, 37% yield). (m/z): [M+H]⁺ calcd for C₃₀H₃₇N₇O₂ 528.30 found 528.3. ¹H NMR (400 MHz, DMSO-d₆) δ 13.09 (s, 1H), 9.40 (s, 1H), 8.27 (d, J = 8.31, 1H), 7.30 (s, 1H), 7.04 (m, 2H), 6.71 (d, J = 2.54, 1H), 6.64 (dd, J = 2.53, 8.26, 1H), 4.26 (m, 1H), 4.06 (m, 2H), 3.82 (m, 2H), 3.64 (m, 2H), 3.03 (m, 2H), 2.74 (m, 2H), 2.47 (q, J = 7.56, 2H), 2.07 (d, J = 3.69, 6H), 1.07 (m, 6H), 1.00 (t, J = 7.50, 3H).

20

Preparation 6: (S)-5-ethyl-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid




(S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, HCl (0.100 g, 0.227 mmol) (**I-18**) and acetaldehyde (0.019 mL, 0.341 mmol) were dissolved in methanol (3.0 mL), then sodium

25


cyanoborohydride (0.057 g, 0.909 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours (reaction progress was monitored by LCMS). Sodium borohydride (9 mg, 0.227 mmol) was added to quench any remaining acetaldehyde, then the reaction mixture was concentrated. The crude product was then purified by reverse phase chromatography (5-70% ACN/Water gradient, 40 g C18 column) to provide the TFA salt of the title compound (62 mg, 50% yield). (m/z): [M+H]⁺ calcd for C₂₄H₂₅N₅O₃ 432.20 found 432.1.

Example 2: (S)-(3-(dimethylamino)azetid-1-yl)(5-ethyl-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone



(S)-5-ethyl-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.055 mmol), 3-(dimethylamino)azetidine dihydrochloride (14.28 mg, 0.082 mmol), and DIPEA (0.048 mL, 0.275 mmol) were dissolved in DMF (1.50 mL), then HATU (31.4 mg, 0.082 mmol) was added and the reaction mixture was stirred at room temperature for 1 hour (reaction progress was monitored by LCMS). Hydrazine (5.18 μ l, 0.165 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-60% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (25 mg, 63% yield). (m/z): [M+H]⁺ calcd for C₂₉H₃₅N₇O₂ 514.29 found 514.2.


Example 3: (S)-(3-(dimethylamino)-3-methylazetid-1-yl)(5-ethyl-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone



3

(*S*)-5-ethyl-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.055 mmol), N,N,3-trimethylazetidin-3-amine hydrochloride (12.43 mg, 0.082 mmol), and DIPEA (0.048 mL, 0.275 mmol) were dissolved in DMF (1.50 mL), then HATU (31.4 mg, 0.082 mmol) was added and the reaction mixture was stirred at room temperature for 1 hour (reaction progress was monitored by LCMS). Hydrazine (5.18 μ l, 0.165 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-60% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (25 mg, 62% yield). (m/z): [M+H]⁺ calcd for C₃₀H₃₇N₇O₂ 528.30 found 528.2.

Example 4: (*S*)-(3-(dimethylamino)-3-methylazetidin-1-yl)(2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone




4

(*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid (40 mg, 0.090 mmol) (**I-19**) N,N,3-trimethylazetidin-3-amine hydrochloride (20.29 mg, 0.135 mmol), and DIPEA (0.047 mL, 0.269 mmol) were dissolved in DMF (1.50 mL), then HATU (51.2 mg, 0.135 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours (reaction progress was monitored by LCMS). Hydrazine (8.45 μ l, 0.269 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative

HPLC (5-60% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (26 mg, 38% yield). (m/z): [M+H]⁺ calcd for C₃₁H₃₉N₇O₂ 542.32 found 542.2.

Preparation 7: (S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid (I-20)




I-20

5

(S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, HCl (**I-18**) (0.160 g, 0.364 mmol) and propionaldehyde (0.039 mL, 0.546 mmol) were dissolved in methanol (3.0 mL), then sodium cyanoborohydride (0.069 g, 1.091 mmol) was added and the reaction mixture was stirred at room temperature for 24 hours (reaction progress was monitored by LCMS). The reaction mixture was concentrated and the crude product was purified by reverse phase chromatography (5-70% ACN/Water gradient, 50g C18 column) to provide the TFA salt of the title compound (78 mg, 38% yield). (m/z): [M+H]⁺ calcd for C₂₅H₂₇N₅O₃ 446.21 found 446.3.

10

Example 5: (S)-3-(dimethylamino)azetid-1-yl(2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone




5

(S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.054 mmol), (**I-20**) 3-(dimethylamino)azetidine dihydrochloride (13.92 mg, 0.080 mmol), and DIPEA (0.047 mL, 0.268 mmol) were dissolved in DMF (1.50 mL), then HATU (30.6 mg, 0.080 mmol) was added and the reaction mixture was stirred at room temperature for 1 hour (reaction progress was monitored by LCMS). Hydrazine (5.05 μ l, 0.161 mmol) was added to cleave

54

undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-60% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (26 mg, 63% yield). (m/z): [M+H]⁺ calcd for C₃₀H₃₇N₇O₂ 528.30 found 528.2.


5 **Example 6: (S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone**



6

10 (S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.054 mmol), (I-20) N,N,3-Trimethylazetidin-3-amine hydrochloride (12.12 mg, 0.080 mmol), and DIPEA (0.047 mL, 0.268 mmol) were dissolved in DMF (1.50 mL), then HATU (30.6 mg, 0.080 mmol) was added and the reaction mixture was stirred at room temperature for 1 hour (reaction
15 progress was monitored by LCMS). Hydrazine (5.05 μl, 0.161 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-60% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (18 mg, 44% yield). (m/z): [M+H]⁺ calcd for C₃₁H₃₉N₇O₂ 542.32 found 542.2.

20 **Preparation 8: 2-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane**



(a) 1-(Benzyloxy)-4-bromo-5-ethyl-2-fluorobenzene


To a solution of 4-bromo-5-ethyl-2-fluorophenol (20 g, 910.32 mmol) in ACN (250
25 mL) was added K₂CO₃ (31.55 g, 228.3 mmol) followed by benzyl bromide (13.10 mL,

109.58 mmol) drop wise. The resulting reaction mixture was stirred at 80 °C for 2 h. The aqueous layer was extracted with EtOAc (three times), combined and washed with brine. The organic layer was dried over Na₂SO₄ and evaporated under reduced pressure to afford the title intermediate as a pale yellow oily liquid (25 g, 89 % yield). ¹H NMR (400 MHz, Chloroform-*d*) δ 7.48 – 7.30 (m, 5H), 7.27 (d, *J* = 10.5 Hz, 1H), 6.87 (d, *J* = 8.7 Hz, 1H), 5.12 (s, 2H), 2.66 (q, *J* = 7.5 Hz, 2H), 1.16 (t, *J* = 7.5 Hz, 3H).

(b) 2-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

To a solution of the product of the previous step (12.5 g, 40.45 mmol) in dioxane (100 mL) was added bis(pinacolato)diboron (15.40 g, 60.67 mmol) and KOAc (11.9 g, 121.35 mmol). The reaction mixture was purged with nitrogen for 15 min followed by addition of [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (1.65 g, 2.023 mmol). The resulting reaction mixture was stirred and heated at 110 °C for 3 h, filtered through Celite and the residue washed with EtOAc. The filtrate was diluted with excess EtOAc (200 mL) and washed with water (100 mL) followed by brine (100 mL), dried over sodium sulfate and concentrated in vacuo to get crude product which was purified by column chromatography over (100-200) silica gel, eluted with 3-5% EtOAc: Hexane to afford the desired product as an off-white solid (9.50 g, 66 % yield). ¹H NMR (400 MHz, Chloroform-*d*) δ 7.54 – 7.27 (m, 6H), 6.81 (d, *J* = 7.9 Hz, 1H), 5.16 (s, 2H), 2.84 (q, *J* = 7.5 Hz, 2H), 1.32 (s, 12H), 1.14 (t, *J* = 7.5 Hz, 3H).

Preparation 9: 6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2H-pyran-2-yl)-3-(trimethylstannyl)-1H-indazole



(a) 6-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazole

To a solution of 6-bromo-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazole (50 g, 178.57 mmol) and 2-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (76.3 g, 214.29 mmol) in DMF:H₂O (480:120 mL) was added K₃PO₄ (94.64 g, 446.86 mmol). The reaction mixture was degassed with nitrogen for 15 min, then Pd(PPh₃)₂Cl₂ catalyst (6.26 g, 8.93 mmol) was added and the mixture was again degassed with nitrogen for 5 min stirred, and heated at 100-110 °C for 5 h. The reaction mixture was filtered through CeliteTM and the residue was washed with EtOAc. The filtrate was diluted with EtOAc, washed with cold water and brine, dried over sodium sulfate and concentrated in vacuo to provide crude product which was purified by flash column chromatography to afford the title intermediate as a white solid (65 g, 86 % yield). (*m/z*): [M+H]⁺ calcd for C₂₇H₂₇FN₂O₂ 431.21 found 431.46. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.06 – 7.98 (m, 2H), 7.70 (d, *J* = 8.2 Hz, 1H), 7.51 – 7.32 (m, 5H), 7.08 (dd, *J* = 8.3 Hz, 1H), 7.03 (d, *J* = 11.9 Hz, 1H), 6.95 (d, *J* = 8.5 Hz, 1H), 5.76 – 5.64 (m, 1H), 5.20 (s, 2H), 4.04 (d, *J* = 10.1 Hz, 1H), 3.72 (t, *J* = 9.7 Hz, 1H), 2.52 (q, *J* = 7.5 Hz, 2H), 2.22 – 2.02 (m, 3H), 1.80 – 1.71 (m, 3H), 1.06 (t, *J* = 7.5 Hz, 3H).

(b) 6-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-1*H*-indazole

To a solution of the product of the previous step (65 g, 151.16 mmol) in methanol (700 mL) was added conc. HCl (120 mL) and the resulting solution was heated at 60-65 °C for 3 h, cooled to RT, and concentrated in vacuo. The residue was dissolved in EtOAc and

washed with saturated NaHCO₃ aqueous solution and water. The organic layer was dried over anhydrous Na₂SO₄ and concentrated in vacuo to afford the title intermediate as a white solid (52 g, 99 % (crude)). ¹H NMR (400 MHz, Chloroform-*d*) δ 8.13 (s, 1H), 7.77 (d, *J* = 8.3 Hz, 1H), 7.59 – 7.30 (m, 6H), 7.10 (d, *J* = 8.3 Hz, 1H), 7.01 (d, *J* = 11.8 Hz, 1H), 6.96 (d, *J* = 8.4 Hz, 1H), 5.21 (s, 2H), 2.53 (q, *J* = 7.5 Hz, 2H), 1.05 (t, *J* = 7.5 Hz, 3H).

(c) 6-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-3-iodo-1*H*-indazole

To a solution of 6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1*H*-indazole (56 g, 161.18 mmol) in DMF (400 mL) was added KOH (36.2 g, 647.39 mmol) and the mixture was stirred for 5 min. A solution of iodine (82.2 g, 323.69 mmol) in DMF (100 mL) was added slowly at 0 °C and stirred at RT for 30 min, diluted with water (3 x 150 mL) and extracted with EtOAc (3 x 200 mL). The organic layer was washed with saturated sodium metabisulfite aqueous solution (3 x 200 mL) and water (400 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to get crude product which was purified by flash column chromatography to afford the title intermediate as a brownish semi-solid (64 g, 84 % yield). ¹H NMR (400 MHz, Chloroform-*d*) δ 10.49 (s, 1H), 7.57 – 7.32 (m, 7H), 7.16 (d, *J* = 8.3 Hz, 1H), 7.04 – 6.91 (m, 2H), 5.20 (s, 2H), 2.51 (q, *J* = 7.4 Hz, 2H), 1.04 (t, *J* = 7.5 Hz, 3H).


(d) 6-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-3-iodo-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazole

To an ice-cold solution of the product of the previous step (60 g, 127.12 mmol) in DCM (700 mL) was added *p*-toluensulfonic acid (4.84 g, 25.423 mmol) followed by 3,4-dihydro-2*H*-pyran (17.43 mL, 190.68 mmol) drop wise. The reaction mixture was stirred at RT overnight, diluted with DCM and washed with saturated NaHCO₃ aqueous solution and brine. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to provide crude product which was purified by flash chromatography (silica gel) to afford the title intermediate as an off white solid (64 g, 91 % yield). (*m/z*): [M+H]⁺ calcd for C₂₇H₂₆FIN₂O₂ 557.10 found 557.30. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.56 – 7.31 (m, 7H), 7.14 (d, *J* = 8.3 Hz, 1H), 7.01 (d, *J* = 11.8 Hz, 1H), 6.95 (d, *J* = 8.5 Hz, 1H), 5.68 (d, *J* = 9.3 Hz, 1H), 5.20 (s, 2H), 4.08 – 3.99 (m, 1H), 3.77 – 3.64 (m, 1H), 2.50 (q, *J* = 7.2 Hz, 2H), 2.23 – 1.97 (m, 3H), 1.81 – 1.68 (m, 3H), 1.06 (t, *J* = 7.4 Hz, 3H).

(e) 6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-3-(trimethylstannyl)-1*H*-indazole

To a solution of 6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-3-iodo-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazole (20 g, 35.97 mmol) in toluene (150 mL) was added hexamethylditin (9.2 mL, 43.17 mmol). The reaction mixture was degassed with nitrogen for 20 min followed by addition of tetrakis (2.0 g, 1.80 mmol) and then stirred at 100 °C for 2 h, cooled to RT, filtered through CeliteTM and residue washed with EtOAc. The filtrate was concentrated and purified by column chromatography (over neutral alumina), eluted with 2-5% EtOAc:Hexane to afford the title compound (17.50 g, 82 % yield). (*m/z*): [M+H]⁺ calcd for C₃₀H₃₅FN₂O₂Sn 595.17, 593.17 found 595.49, 593.55. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.68 (d, *J* = 8.0 Hz, 1H), 7.57 – 7.29 (m, 6H), 7.13 – 7.00 (m, 2H), 6.96 (d, *J* = 8.4 Hz, 1H), 5.81 – 5.68 (m, 1H), 5.21 (s, 2H), 4.13 – 4.00 (m, 1H), 3.81 – 3.66 (m, 1H), 2.54 (q, *J* = 7.3 Hz, 2H), 2.23 – 2.00 (m, 2H), 1.87 – 1.59 (m, 4H), 1.08 (t, *J* = 7.5 Hz, 3H), 0.47 (s, 9H).

Preparation 10: 5-(*tert*-butyl) 6-methyl (*S*)-2-iodo-3-((2-trimethylsilyl)ethoxy methyl)- 3,4,6,7-tetrahydro -5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate



(a) (*S*)-4,5,6,7-Tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid

To a stirred suspension of L-histidine (50 g, 322.24 mmol) in water (420 mL) was added conc. HCl (29 mL) drop wise at 0 °C followed by formaldehyde (55 mL, 676.72 mmol) in one portion at 0 °C. The resulting reaction mixture was stirred for 30 min and then heated at 75 °C for 6 h and concentrated. The resulting crude was stirred for 2 h with diethyl ether, filtered and washed with IPA:THF (100:300 mL) to provide the HCl salt of the title intermediate as an off white solid (75 g 99 % yield (crude)). (*m/z*): [M+H]⁺ calcd for C₇H₉N₃O₂ 168.07 found 168.17.

(b) Methyl (*S*)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylate

To a stirred solution of the product of the previous step (75.0 g, 312.5 mmol) in methanol (1500 mL) was added SOCl₂ (45.6 mL, 625 mmol) dropwise at 0 °C and stirred at RT for 16 h, then heated up to reflux (70 °C) for 1 h. The solvent was removed by distillation and the crude product was triturated with methanol followed by diethyl ether to provide the
5 crude HCl salt of the title intermediate as an off white solid (80 g crude). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (s, 1H), 4.71 (dd, *J* = 9.4, 5.2 Hz, 1H), 4.36 (d, *J* = 15.5 Hz, 1H), 4.30 (d, *J* = 15.6 Hz, 1H), 3.82 (s, 3H), 3.44 – 3.21 (m, 2H).

(c) 5-(*tert*-Butyl) 6-methyl (*S*)-3,4,6,7-tetrahydro-5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate

10 To a stirred solution of the product of the previous step (80.0 g, 314.96 mmol) in methanol (1000 mL) was added DIPEA (282 mL, 1574 mmol) followed by di-*tert*-butyl dicarbonate (172 mL, 787.48 mmol) at 0 °C. The reaction mixture was stirred at RT for 16 h and then liquid NH₃ (150 mL, 25 % in water) was added and the reaction mixture was stirred again for 16 h at RT, methanol was removed by distillation and the residue was extracted in
15 DCM (3 x 200 mL). Combined organic extracts were dried over anhydrous Na₂SO₄, concentrated and purified by flash chromatography (100-200 mesh silica gel), eluted with 5% MeOH:DCM to afford the title intermediate (41 g, 46 % yield). (*m/z*): [M+H]⁺ calcd for C₁₃H₁₉N₃O₄ 282.14 found 282.21. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.85 (s, 1H), 7.50 (s, 1H), 5.18 (dd, *J* = 49.3, 5.1 Hz, 1H), 4.51 (t, *J* = 14.2 Hz, 1H), 4.09 (dd, *J* = 43.9, 16.1 Hz,
20 1H), 3.59 (s, 3H), 3.08 (d, *J* = 15.5 Hz, 1H), 2.94 (d, *J* = 15.1 Hz, 1H), 1.45 (s, 9H).


(d) 5-(*tert*-Butyl) 6-methyl (*S*)-2-iodo-3,4,6,7-tetrahydro-5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate

To a solution of the product of the previous step (41.0 g, 145.9 mmol) in THF (500 mL) was added *N*-iodosuccinimide (66.0 g, 291.8 mmol) at 0 °C and the resulting solution
25 was stirred at RT for 4 h, diluted with water and extracted with ethyl acetate. The organic portion was washed with 10% sodium thiosulphate solution (3 x 200 mL). The combined organic layer was dried over anhydrous sodium sulfate, and concentrated to provide the title compound 60 g (crude), which was used in the next step without further purification.
(*m/z*): [M+H]⁺ calcd for C₁₃H₁₈IN₃O₄ 408.03 found 408.31. ¹H NMR (400 MHz, DMSO-*d*₆)
30 δ 12.48 (s, 1H), 5.34 – 4.97 (m, 1H), 4.67 – 4.35 (m, 1H), 4.12 – 3.95 (m, 1H), 3.60 (s, 3H), 3.14 – 2.82 (m, 2H), 1.44 (s, 9H).

(e) 5-(*tert*-Butyl) 6-methyl (*S*)-2-iodo-3-((2-trimethylsilyl)ethoxy) methyl)- 3,4,6,7-tetrahydro -5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate

To a stirred solution of 5-(*tert*-butyl) 6-methyl (*S*)-2-iodo-3,4,6,7-tetrahydro-5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (40 g, 0.098 mol) in DMF (150 mL) was added
5 DIPEA (35.1 mL, 0.19 mol) at 0 °C. The reaction mixture was stirred for 10 min then 2-(trimethylsilyl)-ethoxymethyl chloride (19.1 mL, 0.10 mol) was added drop wise at 0 °C. The resulting reaction mixture was stirred for 3 h at RT. After 4 h chilled water was added and the reaction mixture was extracted with EtOAc (2 x 200 mL). The organic layer was dried over anhydrous sodium sulphate, concentrated, and purified by flash column
10 chromatography, eluted with 20-35% EtOAc:Hexane, to afford the title product as a pale yellow viscous liquid (27 g). (*m/z*): [M+H]⁺ calcd for C₁₉H₃₂IN₃O₅Si 538.12 found 538.42. ¹H NMR (400 MHz, DMSO-*d*₆) δ 5.33 – 5.04 (m, 3H), 4.79 – 4.56 (m, 1H), 4.54 – 4.14 (m, 1H), 3.60 (s, 3H), 3.47 (t, *J* = 7.8 Hz, 2H), 3.31 – 3.16 (m, 1H), 2.97 (t, *J* = 18.9 Hz, 1H), 1.44 (s, 9H), 0.92 – 0.74 (m, 2H), -0.03 (s, 9H).

15 **Preparation 11: (6*S*)-5-(*tert*-butoxycarbonyl)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazol-3-yl)-3-((2-(trimethylsilyl)ethoxy)methyl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid**



(a) 5-(*tert*-Butyl) 6-methyl (6*S*)-2-(6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazol-3-yl)-3-((2-(trimethylsilyl) ethoxy) methyl)-3,4,6,7-tetrahydro-5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate

To a stirred solution of 5-(*tert*-butyl) 6-methyl (*S*)-2-iodo-3-((2-trimethylsilyl)ethoxy) methyl)- 3,4,6,7-tetrahydro -5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate (17.0 g, 31.65 mmol) in toluene (500 mL) was added 6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-3-(trimethylstannyl)-1*H*-indazole (20 g, 34.82 mmol). The reaction mixture was purged with argon for 15 min, Pd(PPh₃)₄ (3.6 g, 3.16 mmol) and copper iodide (1.20 g, 6.33 mmol) were added and the reaction mixture was stirred at 120 °C for 16 h. The reaction mixture was filtered through CeliteTM, the filtrate was concentrated under reduced pressure and purified by silica gel column chromatography (Redisep 80 g column, eluted with DCM for 10 min and then 15-20% EtOAc in Hexane to afford the title intermediate as a yellow solid (15.10 g, 58 % yield). (*m/z*): [M+H]⁺ calcd for C₄₆H₅₈FN₅O₇Si 840.41 found 840.54. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.43 (s, 1H), 7.54 – 7.33 (m, 6H), 7.20 (s, 1H), 7.05 (d, *J* = 11.4 Hz, 1H), 6.95 (d, *J* = 8.5 Hz, 1H), 6.09 – 5.69 (m, 3H), 5.59 – 5.36 (m, 1H), 5.20 (s, 2H), 4.97 – 4.80 (m, 1H), 4.12 – 3.90 (m, 1H), 3.68 (s, 3H), 3.57 – 3.47 (m, 2H), 3.40 (d, 1H), 3.21 – 3.05 (m, 1H), 2.74 – 2.34 (m, 4H), 2.25 – 2.07 (m, 2H), 1.94 – 1.65 (m, 4H), 1.54 (s, 9H), 1.12 – 0.99 (m, 3H), 0.91 – 0.75 (m, 2H), -0.12 (s, 9H).


(b) 6-Benzyl 5-(*tert*-butyl) (6*S*)-2-(6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazol-3-yl)-3-((2-(trimethylsilyl) ethoxy) methyl)-3,4,6,7-tetrahydro-5*H*-imidazo[4,5-*c*]pyridine-5,6-dicarboxylate

To a round bottom flask was added the product of the previous step (15.0 g, 17.85 mmol) in toluene (400 mL), benzyl alcohol (46.3 mL) and Ti(OEt)₄ (7.15 mL, 35.70 mmol) and the reaction mixture was refluxed vigorously (140 °C) for 48 h, diluted with water and extracted with DCM. The suspension was filtered, filtrate was dried over Na₂SO₄, concentrated under reduced pressure and purified by silica gel column chromatography (Redisep 80 g column, 0-5% EtOAc in hexanes) for 20 min to remove excess benzyl alcohol, then eluted with 10-15% EtOAc in Hexane) to provide the title intermediate. ¹H NMR consistent with structure. (*m/z*): [M+H]⁺ calcd for C₅₂H₆₂FN₅O₇Si 916.44 found 916.86.

(c) (6*S*)-5-(*tert*-butoxycarbonyl)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazol-3-yl)-3-((2-(trimethylsilyl)ethoxy)methyl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid

To a stirred solution of the product of the previous step (21.0 g, 22.92 mmol) in 1:1
5 IPA:THF (400 mL)) was added Pd(OH)₂ (5.0 g). The reaction mixture was stirred at RT for
16 h under a hydrogen balloon, filtered through CeliteTM, concentrated under reduced pressure,
and purified by silica gel column chromatography (Rediseq 80 g column, eluted with 25-40%
EtOAc in Hexane) to provide the title compound (6.1 g, 8.29 mmol) as an off-white solid.
(*m/z*): [M+H]⁺ calcd for C₃₈H₅₀FN₅O₇Si 736.35 found 736.5. ¹H NMR consistent with
10 structure. (*m/z*): [M+H]⁺ calcd for C₃₈H₅₀FN₅O₇Si 736.35 found 736.5. ¹H NMR (400 MHz,
DMSO-*d*₆) δ 12.94 (s, 1H), 9.86 (s, 1H), 8.34 (t, *J* = 7.6 Hz, 1H), 7.66 (s, 1H), 7.20 (d, *J* =
8.7 Hz, 1H), 7.03 (d, *J* = 11.8 Hz, 1H), 6.93 (d, *J* = 9.1 Hz, 1H), 6.11 – 5.77 (m, 3H), 5.33 –
5.06 (m, 1H), 4.87 – 4.56 (m, 1H), 4.52 – 4.14 (m, 1H), 3.97 – 3.69 (m, 2H), 3.53 – 3.40 (m,
2H), 3.23 – 3.11 (m, 1H), 3.11 – 2.93 (m, 1H), 2.47 – 2.44 (m, 2H), 2.13 – 1.96 (m, 2H), 1.68
15 (d, *J* = 70.9 Hz, 4H), 1.48 (s, 9H), 1.02 (t, *J* = 7.5 Hz, 3H), 0.86 – 0.68 (m, 2H), -0.17 (s, 9H).


Preparation 12: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid



To a stirred solution of (6*S*)-5-(*tert*-butoxycarbonyl)-2-(6-(2-ethyl-5-fluoro-4-
20 hydroxyphenyl)-1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-indazol-3-yl)-3-((2-
(trimethylsilyl)ethoxy)-methyl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic
acid (5.7 g, 7.75 mmol) in 5:1 dioxane:water (60 mL) was added conc. HCl (20 mL) drop
wise at 0°C. The reaction mixture was warmed and stirred at 90 °C for 16 h and distilled
under vacuum to provide the crude residue, which was sequentially triturated with chilled
25 diethyl ether and acetonitrile to provide the HCl salt of the title compound (3.6 g, 95 % yield)
as a light brown solid. (*m/z*): [M+H]⁺ calcd for C₂₂H₂₀FN₅O₃ 422.16 found 422.24. ¹H
NMR (400 MHz, D₂O/DMSO-*d*₆) δ 8.22 (d, *J* = 8.4 Hz, 1H), 7.49 (s, 1H), 7.19 (d, *J* = 8.1
Hz, 1 H), 6.99 (d, *J* = 11.9 Hz, 1 H), 6.91 (d, *J* = 9.0 Hz, 1H), 4.56 – 4.51 (m, 1H), 4.36 (d, *J*

= 15.5 Hz, 1H), 4.30 (d, J = 15.5 Hz, 1H), 3.35 – 3.25 (m, 1H), 3.15 – 3.05 (m, 1H), 2.4 – 2.55 (m, 2H), 0.97 (t, J = 7.5 Hz, 3H).

Preparation 13: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid




5

To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, HCl (400 mg, 0.874 mmol), acetone (0.192 mL, 2.62 mmol), and acetic acid (0.150 mL, 2.62 mmol) in DMF (7 mL), was added sodium cyanoborohydride (274 mg, 4.37 mmol) and the reaction mixture was stirred at
10 RT overnight. Sodium borohydride (33 mg, 0.874 mmol) was added, the solution was concentrated, and purified by preparative HPLC to provide the TFA salt of the title compound (115 mg, 23 % yield). (m/z): $[M+H]^+$ calcd for $C_{25}H_{26}FN_5O_3$ 464.20 found 464.5.

Example 7: (*S*)-(3-(dimethylamino)azetid-1-yl)(2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)methanone C-1

15




C-1

To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (179 mg, 0.310 mmol), *N,N*-dimethylazetid-3-amine, 2 HCl (107 mg, 0.465 mmol), and DIPEA (0.162 mL 0.930 mmol) in DMF (4 mL), was added HATU (177 mg, 0.465 mmol) and the
20 reaction mixture was stirred at RT overnight. Hydrazine (5 eq) was added, the reaction

mixture was concentrated and purified by preparative HPLC to provide the TFA salt of the title compound (63 mg, 26 % yield). (*m/z*): [M+H]⁺ calcd for C₃₀H₃₆FN₇O₂ 546.29 found 546.7. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.90 (s, 1H), 8.29 (dd, 1H), 7.34 (s, 1H), 7.07 (d, 1H), 7.01 (d, 1H), 6.89 (d, 1H), 4.35 – 4.18 (m, 1H), 4.11 – 3.94 (m, 1H), 3.94 – 3.73 (m, 3H), 3.70 – 3.57 (m, 2H), 3.06 – 2.94 (m, 2H), 2.87 – 2.66 (m, 2H), 2.48 – 2.40 (m, 2H), 2.13 – 2.00 (m, 6H), 1.07 (t, 3H), 1.03 – 0.93 (m, 6H).


Example 8: (*S*)-3-(dimethylamino)-3-methylazetididin-1-yl(2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone



C-4

(*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.052 mmol), N,N,3-trimethylazetididin-3-amine, 2HCl (29.2 mg, 0.156 mmol), and DIPEA (0.045 mL, 0.260 mmol) were dissolved in DMF (1.0 mL), then HATU (29.6 mg, 0.078 mmol) was added and the reaction mixture was stirred at room temperature for 20 hours (reaction progress was monitored by LCMS). Hydrazine (4.90 μl, 0.156 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (20 70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (13 mg, 33% yield). (*m/z*): [M+H]⁺ calcd for C₃₁H₃₈FN₇O₂ 560.31 found 560.2.


Preparation 14: (*S*)-5-ethyl-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid



65

(S)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, HCl (450 mg, 0.983 mmol) and acetaldehyde (0.083 mL, 1.474 mmol) were dissolved in DMF (7 mL), then sodium cyanoborohydride (247 mg, 3.93 mmol) was added and the reaction mixture was stirred at room temperature for 24 hours (reaction progress was monitored by LCMS). Sodium borohydride (112 mg, 2.95 mmol) was added to quench any remaining acetaldehyde, then the reaction mixture was concentrated. The crude product was then purified by reverse phase chromatography (5-65% ACN/Water gradient, 100 g C18aq column) to provide the TFA salt of the title compound (165 mg, 30% yield). (m/z): [M+H]⁺ calcd for C₂₄H₂₄FN₅O₃ 450.19 found 450.2.


Example 9: (S)-(3-(dimethylamino)azetid-1-yl)(5-ethyl-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone



C-2

(S)-5-ethyl-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.053 mmol) and HATU (30.4 mg, 0.080 mmol) were combined in DMF (1.0 mL). To the solution, N,N-dimethylazetid-3-amine (16 mg, 0.160 mmol) and DIPEA (0.037 mL, 0.213 mmol) were added and the reaction mixture was stirred at room temperature for 6 hours (reaction progress was monitored by LCMS). Hydrazine (4.92 μ l, 0.160 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (10-70% ACN/Water gradient, Zorbax Bonus-RP column) to provide the TFA salt of the title compound (27 mg, 67% yield). (m/z): [M+H]⁺ calcd for C₂₉H₃₄FN₇O₂ 532.28 found 532.2.


Example 10: (S)-(3-(dimethylamino)-3-methylazetid-1-yl)(5-ethyl-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone



C-3


(*S*)-5-ethyl-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.053 mmol) and
 5 HATU (30.4 mg, 0.080 mmol) were combined in DMF (1.0 mL). To the solution, *N,N*,3-trimethylazetididin-3-amine (18 mg, 0.160 mmol) and DIPEA (0.037 mL, 0.213 mmol) were added and the reaction mixture was stirred at room temperature for 6 hours (reaction progress was monitored by LCMS). Hydrazine (4.92 μ l, 0.160 mmol) was added to cleave undesired
 10 byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (10-70% ACN/Water gradient, Zorbax Bonus-RP column) to provide the TFA salt of the title compound (28 mg, 68% yield). (m/z): [M+H]⁺ calcd for C₃₀H₃₆FN₇O₂ 546.29 found 546.2.

Preparation 15: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid



15 (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, HCl (300 mg, 0.655 mmol) and propionaldehyde (0.071 mL, 0.983 mmol) were dissolved in DMF (7 mL), then sodium cyanoborohydride (124 mg, 1.966 mmol) was added and the reaction mixture was stirred at room temperature
 20 for 72 hours (reaction progress was monitored by LCMS). Sodium borohydride (26 mg, 0.655 mmol) was added to quench any remaining aldehyde, then the reaction mixture was concentrated. The crude product was then purified by preparative HPLC (2-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (117 mg, 31% yield). (m/z): [M+H]⁺ calcd for C₂₅H₂₆FN₅O₃ 464.21 found 464.2.

Example 11: (S)-(3-(dimethylamino)azetid-1-yl)(2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone




5

C-5

(S)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (33 mg, 0.057 mmol), N,N-dimethylazetid-3-amine, 2HCl (29.7 mg, 0.171 mmol), and DIPEA (0.060 mL, 0.343 mmol) were dissolved in DMF (2.0 mL), then HATU (28.2 mg, 0.074 mmol) was added and the reaction mixture was stirred at room temperature for 18 hours (reaction progress was monitored by LCMS). Hydrazine (8.97 μ l, 0.286 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (2-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (27 mg, 61% yield). (m/z): [M+H]⁺ calcd for C₃₀H₃₆FN₇O₂ 546.29 found 546.5.

15

Example 12: (S)-(3-(dimethylamino)-3-methylazetid-1-yl)(2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone




20

C-6

(S)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (30 mg, 0.052 mmol), N,N,3-trimethylazetid-3-amine, 2HCl (29.2 mg, 0.156 mmol), and DIPEA (0.045 mL, 0.260 mmol) were dissolved in DMF (1.0 mL), then HATU (29.6 mg, 0.078 mmol) was added and

the reaction mixture was stirred at room temperature for 20 hours (reaction progress was monitored by LCMS). Hydrazine (8.89 μ l, 0.260 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (2-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (15 mg, 37% yield). (m/z): [M+H]⁺ calcd for C₃₁H₃₈FN₇O₂ 560.31 found 560.5.


Preparation 16: (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid (I-21)



10 (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, HCl (I-18) (0.160 g, 0.364 mmol) and formaldehyde solution, 37 wt% in water (0.032 mL, 0.436 mmol) were dissolved in methanol (3.0 mL), then sodium cyanoborohydride (0.069 g, 1.091 mmol) was added and the reaction mixture was stirred at room temperature for 4 hours (reaction progress was monitored by

15 LCMS). Sodium borohydride (14 mg, 0.364 mmol) was added to quench any excess formaldehyde, then the reaction mixture was concentrated. The crude product was purified by reverse phase chromatography (5-70% ACN/Water gradient, 50 g C18 column) to provide the TFA salt of the title compound (110 mg, 57% yield). (m/z): [M+H]⁺ calcd for C₂₃H₂₃N₅O₃ 418.18 found 418.2.

20 **Preparation 17: (*R*)-2-(3-methylpiperazin-1-yl)ethan-1-ol, dihydrochloride (I-24)**



(a) *tert*-butyl (*R*)-4-(2-hydroxyethyl)-2-methylpiperazine-1-carboxylate (I-23)


(*R*)-1-boc-2-methyl-piperazine (0.2 g, 0.999 mmol), DIPEA (0.698 mL, 3.99 mmol), and 2-bromoethanol (0.085 mL, 1.198 mmol) were dissolved in DMF (5 mL) and the

reaction mixture was stirred at 60 °C for 16 hours (reaction progress was monitored by LCMS). The reaction mixture was concentrated, then 10 mL of diethyl ether was added to the residue. The solution was sonicated until the residue gel had disappeared and a solid had formed. The ether solution was then decanted away from the solid. The solid was then used
5 directly in the next reaction (quantitative yield was assumed). (m/z): [M+H]⁺ calcd for C₁₂H₂₄N₂O₃ 245.18 found 245.4.

(b) (*R*)-2-(3-methylpiperazin-1-yl)ethan-1-ol, dihydrochloride (**I-24**)

Tert-butyl (*R*)-4-(2-hydroxyethyl)-2-methylpiperazine-1-carboxylate (0.244 g, 0.999 mmol) was dissolved in Dioxane (3.0 mL) and Water (0.5 mL), then hydrogen chloride
10 solution, 4 M in Dioxane (2.0 mL, 65.8 mmol) was added and the reaction mixture was stirred at room temperature for 3 hours (reaction progress was monitored by LCMS). The reaction mixture was frozen and lyophilized, and the resulting product was used without purification (quantitative yield was assumed). (m/z): [M+H]⁺ calcd for C₇H₁₆N₂O 145.13 found 145.4.

15 **Example 1B: (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-6-yl)(4-(2-hydroxyethyl)piperazin-1-yl)methanone**




1B

20 (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (**I-20**) (40 mg, 0.071 mmol), *n*-(2-hydroxyethyl)piperazine (0.018 mL, 0.143 mmol), and DIPEA (0.025 mL, 0.143 mmol) were dissolved in DMF (1.5 mL), then HATU (40.8 mg, 0.107 mmol) was added and the reaction
25 LCMS). Hydrazine (0.011 mL, 0.357 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (31 mg, 56% yield).

(m/z): [M+H]⁺ calcd for C₃₁H₃₉N₇O₃ 558.31 found 558.3. ¹H NMR (400 MHz, DMSO-d₆) δ 13.10 (s, 1H), 12.27 (d, J = 48.92 Hz, 1H), 9.40 (s, 1H), 8.28 (t, J = 8.10 Hz, 1H), 7.30 (s, 1H), 7.04 (m, 2H), 6.71 (d, J = 2.46 Hz, 1H), 6.64 (dd, J = 2.50, 8.23 Hz, 1H), 4.44 (t, J = 5.31 Hz, 1H), 4.11 (q, J = 5.26, 2H), 3.96 (m, 1H), 3.86-3.52 (m, 6H), 3.49 (q, J = 6.01 Hz, 2H), 2.95 (m, 2H), 2.48 (q, J = 7.48 Hz, 2H), 2.42-2.21 (m, 4H), 2.37 (t, J = 6.20 Hz, 2H), 1.42 (m, 2H), 1.00 (t, J = 7.52 Hz, 3H), 0.81 (m, 3H).

Example 2B: ((S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)((1S,4S)-5-methyl-2,5-diazabicyclo[2.2.1]heptan-2-yl)methanone



10

2B


(S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (**I-20**) (40 mg, 0.071 mmol), (1S,4S)-5-Methyl-2,5-diazabicyclo[2.2.1]heptane dihydrobromide (29.4 mg, 0.107 mmol), and DIPEA (0.062 mL, 0.357 mmol) were dissolved in DMF (1.5 mL), then HATU (40.8 mg, 0.107 mmol) was added and the reaction mixture was stirred at room temperature for 3 hours (reaction progress was monitored by LCMS). Hydrazine (0.011 mL, 0.357 mmol) was added to cleave undesired byproducts, then the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (32 mg, 59% yield). (m/z): [M+H]⁺ calcd for C₃₁H₃₇N₇O₂ 540.30 found 540.3.

15

20

Example 3B: ((S)-2,4-dimethylpiperazin-1-yl)((S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone

25



(a) *Tert*-butyl (*S*)-4-((*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carbonyl)-3-methylpiperazine-1-carboxylate (**I-25**)

5

(*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (**I-21**) (55 mg, 0.103 mmol), (*S*)-4-*n*-*boc*-2-methylpiperazine (41.5 mg, 0.207 mmol), and DIPEA (0.036 mL, 0.207 mmol) were dissolved in DMF (1.5 mL), then HATU (59.0 mg, 0.155 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours (reaction progress was monitored by LCMS). Hydrazine (0.016 mL, 0.517 mmol) was added to cleave undesired byproducts, then the reaction mixture was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by reverse phase chromatography (5-70% ACN/Water gradient, 50 g C18 column) to provide the TFA salt of the title compound (54 mg, 72% yield). (*m/z*): [*M*+*H*]⁺ calcd for C₃₃H₄₁N₇O₄ 600.33 found 600.3.

15

(b) ((*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)((*S*)-2-methylpiperazin-1-yl)methanone (**I-26**)


Tert-butyl (*S*)-4-((*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carbonyl)-3-methylpiperazine-1-carboxylate, TFA (**I-25**) (0.126 g, 0.177 mmol) was dissolved in dioxane (1.5 mL) and water (0.3 mL), then hydrogen chloride solution, 4 M in dioxane (1.5 mL, 6.00 mmol) was added and the reaction mixture was stirred at room temperature for 2 hours (reaction progress was monitored by LCMS). The reaction mixture was frozen and lyophilized and the resulting powder was used directly in the next reaction (quantitative yield was assumed). (*m/z*): [*M*+*H*]⁺ calcd for C₂₈H₃₃N₇O₂ 500.27 found 500.3.

25

(c) ((*S*)-2,4-dimethylpiperazin-1-yl)((*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-6-yl)methanone

((*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-6-yl)((*S*)-2-methylpiperazin-1-yl)methanone, dihydrochloride
5 (0.101 g, 0.176 mmol) and formaldehyde solution, 37 wt% in water (0.016 mL, 0.212 mmol) were dissolved in Methanol (3.0 mL), then sodium cyanoborohydride (0.055 g, 0.882 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours (reaction progress was monitored by LCMS). Sodium borohydride (7 mg, 0.176 mmol) was added to quench any remaining formaldehyde. The reaction mixture was concentrated, then the crude
10 product was purified by preparative HPLC (5-60% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (28 mg, 21% yield). (*m/z*): [M+H]⁺ calcd for C₂₉H₃₅N₇O₂ 514.29 found 514.3.

Example 4B: (*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridin-6-yl)(4-methyl-1,4-diazepan-1-yl)methanone
15




4B

((*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (**I-19**) (50 mg, 0.089 mmol), 1-
20 methylhomopiperazine (0.022 mL, 0.179 mmol), and DIPEA (0.031 mL, 0.179 mmol) were dissolved in DMF (1.5 mL), then HATU (51.0 mg, 0.134 mmol) was added and the reaction mixture was stirred at room temperature for 3 hours (reaction progress was monitored by LCMS). Hydrazine (0.014 mL, 0.447 mmol) was added to cleave undesired byproducts, and the solution was stirred at room temperature for 10 minutes. The reaction mixture was then
25 concentrated and the crude product was purified by preparative HPLC (2-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (29 mg, 42% yield). (*m/z*): [M+H]⁺ calcd for C₃₁H₃₉N₇O₂ 542.32 found 542.3. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.08 (s, 1H), 12.21 (d, *J* = 29.9 Hz, 1H), 9.40 (s, 1H), 8.27 (d, *J* = 8.33 Hz, 1H), 7.30 (s,

1H), 7.04 (t, J = 8.05, 2H), 6.71 (d, J = 2.53 Hz, 1H), 6.64 (dd, J = 2.54, 8.23 Hz, 1H), 4.11 (m, 3H), 3.91-3.52 (m, 6H), 2.97 (m, 1H), 2.91-2.53 (m, 4H), 2.49 (q, J = 7.46, 2H), 2.23 (d, J = 13.9 Hz, 3H), 1.76 (m, 2H), 1.0 (m, 9H).


Example 5B: ((S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)((R)-4-(2-hydroxyethyl)-2-methylpiperazin-1-yl)methanone



5B

(S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (**I-21**) (55 mg, 0.103 mmol), *(R)*-2-(3-methylpiperazin-1-yl)ethan-1-ol, dihydrochloride (**I-24**) (33.7 mg, 0.155 mmol), and DIPEA (0.090 mL, 0.517 mmol) were dissolved in DMF (1.5 mL), then HATU (59.0 mg, 0.155 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours (reaction progress was monitored by LCMS). Hydrazine (0.016 mL, 0.517 mmol) was added to cleave undesired byproducts, then the reaction mixture was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated, and the crude product was purified by preparative HPLC (5-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (22 mg, 28% yield). (m/z): [M+H]⁺ calcd for C₃₀H₃₇N₇O₃ 544.30 found 544.3.


Example 6B: ((S)-3-(dimethylamino)pyrrolidin-1-yl)((S)-2-(6-(2-ethyl-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone



6B


(*S*)-2-(6-(2-ethyl-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (**I-19**) (50 mg, 0.089 mmol), (*S*)-(-)-3-(dimethylamino)pyrrolidine (0.023 mL, 0.179 mmol), and DIPEA (0.031 mL, 0.179 mmol) were dissolved in DMF (1.5 mL), then HATU (51.0 mg, 0.134 mmol) was added and the reaction mixture was stirred at room temperature for 3 hours (reaction progress was monitored by LCMS). Hydrazine (0.014 mL, 0.447 mmol) was added to cleave undesired byproducts, and the solution was stirred at room temperature for 10 minutes. The reaction mixture was then concentrated and the crude product was purified by preparative HPLC (5-70% ACN/Water gradient, C18 column) to provide the TFA salt of the title compound (37 mg, 53% yield). (*m/z*): [M+H]⁺ calcd for C₃₁H₃₉N₇O₂ 542.32 found 542.3.

Preparation 18: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid




To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, HCl (400 mg, 0.874 mmol) and propionaldehyde (0.095 mL, 1.310 mmol) in DMF (7 mL), was added sodium cyanoborohydride (165 mg, 2.62 mmol) and the reaction mixture was stirred at RT overnight. Sodium borohydride (33 mg, 0.874 mmol) was added, the solution was concentrated, and purified by preparative HPLC to provide the TFA salt of the title compound (179 mg, 37 % yield). (*m/z*): [M+H]⁺ calcd for C₂₅H₂₆FN₅O₃ 464.20 found 464.5.

Preparation 19: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid




To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-
5 4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, HCl (400 mg, 0.874 mmol),
acetone (0.192 mL, 2.62 mmol), and acetic acid (0.150 mL, 2.62 mmol) in DMF (7 mL), was
added sodium cyanoborohydride (274 mg, 4.37 mmol) and the reaction mixture was stirred at
RT overnight. Sodium borohydride (33 mg, 0.874 mmol) was added, the solution was
concentrated, and purified by preparative HPLC to provide the TFA salt of the title
10 compound (115 mg, 23 % yield). (*m/z*): [M+H]⁺ calcd for C₂₅H₂₆FN₅O₃ 464.20 found
464.5.

Preparation 20: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid



15 To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-
4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, HCl (**8'**) (300 mg,
0.655 mmol) and 37 wt. % formaldehyde in water (0.059 mL, 0.786 mmol) DMF (5 mL),
was added sodium cyanoborohydride (165 mg, 2.62 mmol) and the reaction mixture was
stirred at RT overnight. Sodium borohydride 25 mg, 0.655 mmol) was added, the solution
20 was concentrated, and purified by flash chromatography (100 g column, 5-75 % ACN/water),
to provide the TFA salt of the title compound (85 mg, 24 % yield). (*m/z*): [M+H]⁺ calcd for
C₂₃H₂₂FN₅O₃ 436.17 found 436.45.

Example C-1B: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)(4-(2-hydroxyethyl)piperazin-1-yl)methanone C-1B




5

C-1B

To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (30 mg, 0.052 mmol), 2-(piperazin-1-yl)ethanol, 2HCl (0.19 mL, 0.156 mmol), and DIPEA (0.027 mL, 0.156 mmol) in DMF (1.5 mL), was added HATU (29.6 mg, 0.078 mmol) and the reaction mixture was stirred at RT overnight. Hydrazine (5 eq) was added, the reaction mixture was concentrated and purified by preparative HPLC to provide the TFA salt of the title compound (15.4 mg, 37 % yield). (*m/z*): [M+H]⁺ calcd for C₃₁H₃₈FN₇O₃ 576.30 found 576.2.

15 **Example C-2B: ((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)((1*S*,4*S*)-5-methyl-2,5-diazabicyclo-[2.2.1]heptan-2-yl)methanone C-2B**




C-2B

20 To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-propyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (30 mg, 0.052 mmol), (1*S*,4*S*)-5-methyl-2,5-diazabicyclo[2.2.1]heptane dihydrobromide (42.7 mg, 0.156

mmol), and DIPEA (0.064 mL, 0.364 mmol) in DMF (1.5 mL), was added HATU (29.6 mg, 0.078 mmol) and the reaction mixture was stirred at RT overnight. Hydrazine (5 eq) was added, the reaction mixture was concentrated and purified by preparative HPLC to provide the TFA salt of the title compound (27 mg, 66 % yield). (*m/z*): [M+H]⁺ calcd for

5 C₃₁H₃₆FN₇O₂ 558.29 found 558.3. ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.17 (dt, 1H), 7.59 – 7.50 (m, 1H), 7.32 (dd, 1H), 6.95 (d, 1H), 6.90 (d, 1H), 5.03 – 4.91 (m, 2H), 4.56 – 4.34 (m, 2H), 4.30 – 3.88 (m, 4H), 3.76 – 3.55 (m, 1H), 3.28 – 3.10 (m, 1H), 3.10 – 2.96 (m, 4H), 2.81 – 2.62 (m, 2H), 2.53 (q, 2H), 2.47 – 2.33 (m, 1H), 2.31 – 2.14 (m, 1H), 1.79 – 1.57 (m, 2H), 1.07 (t, 3H), 0.97 (td, 3H).

10 **Example C-3B: ((*S*)-2,4-dimethylpiperazin-1-yl)((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone C-3B**



15 (a) *Tert*-butyl (*S*)-4-((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carbonyl)-3-methylpiperazine-1-carboxylate
(*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid (0.120 g, 0.276 mmol), (*S*)-4-*n*-boc-
20 2-methylpiperazine (0.110 g, 0.551 mmol), and DIPEA (0.096 mL, 0.551 mmol) were dissolved in DMF (3.0 mL), then HATU (0.157 g, 0.413 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours (reaction progress was monitored by LCMS). Hydrazine (0.043 mL, 1.378 mmol) was added to cleave undesired byproducts, then the reaction mixture was stirred at room temperature for 10 minutes. The reaction mixture
25 was then concentrated and the crude product was purified by reverse phase chromatography

(5-70% ACN/Water gradient, 50 g C18 column) to provide the TFA salt of the title compound (126 mg, 62% yield). (m/z): [M+H]⁺ calcd for C₃₃H₄₀FN₇O₄ 618.32 found 618.3.


(b) ((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)((*S*)-2-methylpiperazin-1-yl)methanone

5 Tert-butyl (*S*)-4-((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carbonyl)-3-methylpiperazine-1-carboxylate, TFA (0.126 g, 0.172 mmol) was dissolved in dioxane (1.5 mL) and water (0.3 mL), then hydrochloric acid, 4 M in dioxane (1.5 mL, 49.4 mmol) was added and the reaction mixture was stirred at room temperature for 1 hour (reaction progress was monitored
10 by LCMS). The reaction mixture was frozen and lyophilized and the resulting powder was used directly in the next reaction (quantitative yield was assumed). (m/z): [M+H]⁺ calcd for C₂₈H₃₂FN₇O₂ 518.26 found 518.3.

(c) ((*S*)-2,4-dimethylpiperazin-1-yl)((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)methanone

15 ((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)((*S*)-2-methylpiperazin-1-yl)methanone, dihydrochloride (0.102 g, 0.173 mmol) and formaldehyde solution, 37 wt% in water (0.015 mL, 0.207 mmol) were dissolved in methanol (3.0 mL), then sodium cyanoborohydride
20 (0.054 g, 0.864 mmol) was added and the reaction mixture was stirred at room temperature for 20 hours (reaction progress was monitored by LCMS). Sodium borohydride (7 mg, 0.173 mmol) was added to quench any remaining formaldehyde. The reaction mixture was concentrated, then the crude product was purified by preparative HPLC (5-60% ACN/Water gradient, C18 column) to provide the title compound as the TFA salt (59 mg, 45% yield).
25 (m/z): [M+H]⁺ calcd for C₂₉H₃₄FN₇O₂ 532.28 found 532.3.


Example C-4B: (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)(4-methyl-1,4-diazepan-1-yl)methanone C-4B



C-4B

To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (30 mg, 0.052 mmol), 1-methylhomopiperazine (0.019 mL, 0.156 mmol), and DIPEA (0.036 mL, 0.208 mmol) in DMF (1 mL), was added HATU (29.6 mg, 0.078 mmol) and the reaction mixture was stirred at RT for 3 h. Hydrazine (5 eq) was added, the reaction mixture was stirred at RT for 10 min, concentrated and purified by preparative HPLC to provide the TFA salt of the title compound (26.9 mg, 66 % yield). (*m/z*): [M+H]⁺ calcd for C₃₁H₃₈FN₇O₂ 560.31 found 560.2.


Example C-5B: ((*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridin-6-yl)((*R*)-4-(2-hydroxyethyl)-2-methyl-piperazin-1-yl)methanone C-5B



C-5B

To a solution of (*S*)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1*H*-indazol-3-yl)-5-methyl-4,5,6,7-tetrahydro-3*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, TFA (30 mg, 0.052 mmol), (*R*)-2-(3-methylpiperazin-1-yl)ethanol, 2 HCl (35.6 mg, 0.164 mmol), and DIPEA (0.057 mL, 0.328 mmol) in DMF (1 mL), was added HATU (31.1 mg, 0.082 mmol) and the reaction mixture was stirred at RT overnight. Hydrazine (8.57 μL, 0.273 mmol) was added, the reaction mixture was concentrated and purified by preparative HPLC to provide the TFA salt of the title compound (15.6 mg, 36 % yield). (*m/z*): [M+H]⁺ calcd for C₃₀H₃₆FN₇O₃ 562.29 found 562.2.

Example C-6B: ((S)-3-(dimethylamino)pyrrolidin-1-yl)((S)-5-isopropyl-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridin-6-yl)methanone C-6B



C-6B

5

To a solution of (S)-2-(6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-indazol-3-yl)-5-isopropyl-4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine-6-carboxylic acid, TFA (179 mg, 0.310 mmol), (S)-N,N-dimethylpyrrolidin-3-amine (0.079 mL, 0.620 mmol), and DIPEA (0.162 mL 0.930 mmol) in DMF (4 mL), was added HATU (177 mg, 0.465 mmol) and the reaction mixture was stirred at RT overnight. Hydrazine (5 eq) was added, the reaction mixture was concentrated and purified by preparative HPLC to provide the TFA salt of the title compound (107 mg, 44 % yield). (m/z): [M+H]⁺ calcd for C₃₁H₃₈FN₇O₂ 560.31 found 560.2. ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.21 (d, 1H), 7.50 (s, 1H), 7.26 (d, 1H), 6.94 (d, 1H), 6.90 (d, 1H), 4.83 – 4.66 (m, 1H), 4.48 – 4.25 (m, 2H), 4.23 – 4.12 (m, 1H), 4.12 – 3.93 (m, 2H), 3.93 – 3.63 (m, 3H), 3.62 – 3.48 (m, 1H), 3.26 – 3.09 (m, 1H), 2.98 (d, 6H), 2.67 – 2.57 (m, 1H), 2.53 (q, 2H), 2.44 – 2.12 (m, 1H), 1.41 (t, 3H), 1.31 (d, 3H), 1.05 (t, 3H).

15

Biological Assays

The compounds of the invention have been characterized in one or more of the following biological assays.

20

Assay 1: Biochemical JAK Kinase Assays

A panel of four LanthaScreen JAK biochemical assays (JAK1, 2, 3 and Tyk2) were carried in a common kinase reaction buffer (50 mM HEPES, pH 7.5, 0.01% Brij-35, 10 mM MgCl₂, and 1 mM EGTA). Recombinant GST-tagged JAK enzymes and a GFP-tagged STAT1 peptide substrate were obtained from Life Technologies.

25

Serially diluted compounds were pre-incubated with each of the four JAK enzymes and the substrate in white 384-well microplates (Corning) at ambient temperature for 1h. ATP was subsequently added to initiate the kinase reactions in 10 μL total volume, with 1%

DMSO. The final enzyme concentrations for JAK1, 2, 3 and Tyk2 are 4.2 nM, 0.1 nM, 1 nM, and 0.25 nM respectively; the corresponding Km ATP concentrations used are 25 μM, 3 μM, 1.6 μM, and 10 μM; while the substrate concentration is 200 nM for all four assays. Kinase reactions were allowed to proceed for 1 hour at ambient temperature before a 10 μL
5 preparation of EDTA (10mM final concentration) and Tb-anti-pSTAT1 (pTyr701) antibody (Life Technologies, 2 nM final concentration) in TR-FRET dilution buffer (Life Technologies) was added. The plates were allowed to incubate at ambient temperature for 1h before being read on the EnVision reader (Perkin Elmer). Emission ratio signals (520nm/495nm) were recorded and utilized to calculate the percent inhibition values based
10 on DMSO and background controls.

For dose-response analysis, percent inhibition data were plotted vs. compound concentrations, and IC₅₀ values were determined from a 4-parameter robust fit model with the Prism software (GraphPad Software). Results were expressed as pIC₅₀ (negative logarithm of IC₅₀) and subsequently converted to pK_i (negative logarithm of dissociation
15 constant, K_i) using the Cheng-Prusoff equation.

Test compounds having a lower K_i value or higher pK_i value in the four JAK assays show greater inhibition of JAK activity.

Assay 2: Inhibition of IL-2 Stimulated pSTAT5 in Tall-1 T cells

The potency of test compounds for inhibition of interleukin-2 (IL-2) stimulated
20 STAT5 phosphorylation was measured in the Tall-1 human T cell line (DSMZ) using AlphaLisa. Because IL-2 signals through JAK1/3, this assay provides a measure of JAK1/3 cellular potency.

Phosphorylated STAT5 was measured via the AlphaLISA SureFire Ultra pSTAT5 (Tyr694/699) kit (PerkinElmer).

25 Human T cells from the Tall-1 cell line were cultured in a 37°C, 5% CO₂ humidified incubator in RPMI (Life Technologies) supplemented with 15% Heat Inactivated Fetal Bovine Serum (FBS, Life Technologies), 2mM Glutamax (Life Technologies), 25mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). Compounds were serially diluted in DMSO and dispensed aoustically to empty wells. Assay media (phenol
30 red-free DMEM (Life Technologies) supplemented with 10% FBS (ATCC)) was dispensed (4 μL/well) and plates shaken at 900rpm for 10 mins. Cells were seeded at 45,000 cells/well

in assay media (4 μ L/well), and incubated at 37°C, 5% CO₂ for 1 hour, followed by the addition of IL-2 (R&D Systems; final concentration 300 ng/mL) in pre-warmed assay media (4 μ L) for 30 minutes. After cytokine stimulation, cells were lysed with 6ul of 3x AlphaLisa Lysis Buffer (PerkinElmer) containing 1x PhosStop and Complete tablets (Roche). The lysate was shaken at 900rpm for 10 minutes at room temperature (RT). Phosphorylated STAT5 was measured via the pSTAT5 AlphaLisa kit (PerkinElmer). Freshly prepared acceptor bead mixture was dispensed onto lysate (5 μ L) under green filtered <100 lux light. Plates were shaken at 900rpm for 2mins, briefly spun down, and incubated for 2hrs at RT in the dark. Donor beads were dispensed (5 μ L) under green filtered <100 lux light. Plates were shaken at 900rpm for 2 minutes, briefly spun down, and incubated overnight at RT in the dark. Luminescence was measured with excitation at 689 nm and emission at 570 nm using an EnVision plate reader (PerkinElmer) under green filtered <100 lux light.

To determine the inhibitory potency of test compounds in response to IL-2, the average emission intensity of beads bound to pSTAT5 was measured in a human T cell line. IC₅₀ values were determined from analysis of the inhibition curves of signal intensity versus compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values (mean \pm standard deviation).

In Vitro Assay Results

Table 1

Example Number	JAK1 pK_i	JAK2 pK_i	JAK3 pK_i	Tyk2 pK_i	Tall-1 pIC₅₀
1	10.2	10.5	10.2	9.1	8.6
C-1	10.4	10.8	10.1	9.5	8.8
2	10.3	10.7	10.3	9.2	8.7
C-2	10.2	10.9	9.5	9.4	8.7
3	10.4	10.6	10.2	9.1	8.6
C-3	10.3	10.9	9.7	9.4	8.7
4	10.0	10.8	9.9	9.1	8.7
C-4	10.1	10.9	9.5	9.4	8.6
5	10.5	10.6	10.3	9.0	8.7

C-5	10.1	10.8	10.0	9.4	8.7
6	10.3	10.5	9.8	9.0	8.6
C-6	10.3	10.7	10.0	9.3	8.6
1B	10.2	10.3	9.8	9.0	8.8
C-1B	10.2	10.5	9.7	9.1	8.7
2B	10.3	10.2	9.8	8.9	8.4
C-2B	10.3	10.8	9.9	9.5	8.6
3B	10.2	10.4	10.2	9.3	8.8
C-3B	10.2	10.5	9.7	9.3	8.7
4B	10.1	10.4	10.1	9.0	8.6
C-4B	10.0	10.6	9.8	9.3	8.7
5B	10.3	10.5	10.1	9.2	8.8
C-5B	10.1	10.7	9.9	9.5	8.6
6B	10.2	10.4	10.0	9.0	8.7
C-6B	10.2	10.7	9.8	9.3	8.6

Assay 3: Murine (Mouse) model of IL-13 induced pSTAT6 induction in lung tissue

IL-13 is an important cytokine underlying the pathophysiology of asthma (Kudlacz et al. *Eur. J. Pharmacol*, 2008, 582,154-161). IL-13 binds to cell surface receptors activating members of the Janus family of kinases (JAK) which then phosphorylate STAT6 and subsequently activates further transcription pathways. In the described model, a dose of IL-13 was delivered locally into the lungs of mice to induce the phosphorylation of STAT6 (pSTAT6) which is then measured as the endpoint.

Adult Balb/c mice from Harlan were used in the assay. On the day of study, animals were lightly anesthetized with isoflurane and administered either vehicle or test compound (1 mg/mL, 50 μ L total volume over several breaths) via oral aspiration. Animals were placed in lateral recumbency post dose and monitored for full recovery from anesthesia before being returned to their home cage. Four hours later, animals were once again briefly anesthetized and challenged with either vehicle or IL-13 (0.03 μ g total dose delivered, 50 μ L total volume) via oral aspiration before being monitored for recovery from anesthesia and returned

to their home cage. One hour after vehicle or IL-13 administration, whole blood and lungs were collected for both pSTAT6 detection in lung homogenates using a Perkin Elmer AlphaLISA® SureFire® Ultra™ HV p-STAT6 (Tyr641) assay kit and for total drug concentration analysis in both lung and plasma. Blood samples were centrifuged (Eppendorf centrifuge, 5804R) for 4 minutes at approximately 12,000 rpm at 4°C to collect plasma. Lungs were rinsed in DPBS (Dulbecco's Phosphate-Buffered Saline), padded dry, flash frozen, weighed, and homogenized at a dilution of 1:3 in 0.1% formic acid in HPLC water. Plasma and lung levels of test compound were determined by LC-MS analysis against analytical standards constructed into a standard curve in the test matrix. A lung to plasma ratio was determined as the ratio of the lung concentration in ng/g to the plasma concentration in ng/mL at 5 hours.

Activity in the model is evidenced by a decrease in the level of pSTAT6 present in the lungs of treated animals at 5 hours compared to the vehicle treated, IL-13 challenged control animals. The difference between the control animals which were vehicle-treated, IL-13 challenged and the control animals which were vehicle-treated, vehicle challenged dictated the 0% and 100% inhibitory effect, respectively, in any given experiment. The compounds tested in the assay exhibited inhibition of STAT6 phosphorylation at 5 hours after IL-13 challenge as documented below.

Table 2: pSTAT6 Inhibition and Plasma/Lung Exposure Observed

Compound	Lung Concentration (ng/g) at 5 hr	Plasma Concentration (ng/mL) at 5 hr	Lung to Plasma ratio at 5 hr	pSTAT6 inhibition at 5 hours
1	10155±1979	24.0±16.2	423	75
2	7750±1652	23±6	339	72
3	5130±2205	24±6	216	73
4	15000±3349	55±21	271	72
5	6940±4248	25±9	281	55
6	7465±3084	23±2.5	330	66
1B	13000±5720	10.3±3.9	1262	67
2B	23450±12528	17.5±25	1340	72

Compound	Lung Concentration (ng/g) at 5 hr	Plasma Concentration (ng/mL) at 5 hr	Lung to Plasma ratio at 5 hr	pSTAT6 inhibition at 5 hours
3B	6330±1131	15.4±7	411	76
4B	17350±6625	22.0±23	788	67
5B	5445±1862	13.0±3.7	418	73
6B	11600±4682	11.3±5.7	1026	37

Observation of significant compound concentration in the mouse lung confirmed that the observed inhibition of IL-13 induced pSTAT6 induction was a result of the activity of the test compound. The lung to plasma ratio at 5 hours showed that compound 1 exhibited significantly more exposure in the lung than exposure in plasma in mice.

Assay 4: Inhibition of TSLP-Evoked TARC Release in Human Peripheral Blood Mononuclear Cells

Thymic stromal lymphopoietin (TSLP) and thymus and activation-regulated chemokine (TARC) are overexpressed in asthmatic airways, and correlate with disease severity. In the lungs, TSLP may be released by bronchial epithelial cells in response to allergens and viral infections. TSLP signals through an IL-7R α /TSLPR heterodimer found in a broad range of tissues and cell types, including epithelial cells, endothelial cells, neutrophils, macrophages, and mast cells. The binding of TSLP to its receptor induces a conformational change that activates JAK1 and JAK2 to phosphorylate various transcription factors, including STAT3 and STAT5. In immune cells, this triggers a cascade of intracellular events that result in cell proliferation, anti-apoptosis, dendritic cell migration, and production of Th2 cytokines and chemokines. In peripheral blood mononuclear cells (PBMC), TSLP has a proinflammatory effect by activating myeloid dendritic cells to attract and stimulate T cells, a process mediated by the chemoattractant TARC.

In this assay, it was shown that TSLP stimulation induces TARC release from PBMCs, and that this response is attenuated in a dose-dependent manner upon treatment with compound. The potencies of the test compounds were measured for inhibition of TARC release.

PBMC aliquots (previously isolated from whole blood and frozen in aliquots at -80°C) from 3 to 5 donors were thawed at 37°C and added dropwise to 40 mL pre-warmed, sterile-filtered, complete RPMI media in 50 mL Falcon tubes. Cells were pelleted and resuspended in complete media at 2.24×10^6 cells/mL. Cells were seeded at 85 μ L (190,000 cells) per well in a tissue culture treated 96-well flat bottom microplate. Cells were allowed to rest for 1 hour at 37°C with 5% CO₂.

Compounds were received as 10 mM stock solutions in DMSO. 3.7-fold serial dilutions were performed to generate 9 concentrations of test compound in DMSO at 300X the final assay test concentration. 150-fold intermediate dilutions were performed in complete media to generate compound at 2X the final assay test concentration with 0.2% DMSO. After the 1 hour rest period, 95 μ L of 2X compound was added to each well of PBMC, for a final assay concentration range of 33.33 μ M to 0.95 μ M. 95 μ L of 0.2% DMSO in complete media was added to the untreated control wells. Cells were pre-treated with compound for 1 hour at 37°C with 5% CO₂ prior to stimulation.

Recombinant human TSLP protein was reconstituted at 10 μ g/mL in sterile DPBS with 0.1% BSA and stored in aliquots at -20°C. Immediately prior to use, an aliquot was thawed and prepared at 20X the final assay concentration in complete media. 10 μ L of 20X TSLP was added to each well of PBMC, for a final assay concentration of 10 ng/mL. 10 μ L of complete media was added to the unstimulated control wells. Cells were stimulated in the presence of compound for 48 hours at 37°C with 5% CO₂.

Following stimulation, the cell culture supernatants were harvested and TARC levels were detected by enzyme-linked immunosorbent assay (ELISA), using Human CCL17/TARC Quantikine ELISA Kit (R&D Systems #DDN00) according to the manufacturer's instructions.

For dose response analysis, the log [test compound (M)] was plotted versus the percent response values for each donor, and IC₅₀ values were determined using a nonlinear regression analysis with GraphPad Prism Software using the 4-parameter sigmoidal dose-response algorithm with variable slope. Data are expressed as mean pIC₅₀ (negative decadic logarithm IC₅₀) values calculated from pIC₅₀ values of individual donors and rounded to one decimal place. The potency values for inhibition by original compounds and their des-fluoro modified analogues are summarized in Table 3.

Table 3: Potency (pIC₅₀) Values of Test Compounds for Inhibition of TSLP-evoked TARC Release in Human Peripheral Blood Mononuclear Cells

Compound	pIC₅₀ ± st. dev.
1	7.2 ± 0.1
C-1	7.0 ± 0.1
1B	7.1 ± 0.1
C-1B	7.0 ± 0.3
2B	6.8 ± 0.2
C-2B	6.6 ± 0.2
3B	7.3 ± 0.2
C-3B	7.0 ± 0.1
4B	6.8 ± 0.1
C-4B	7.0 ± 0.4
5B	7.5 ± 0.6
C-5B	7.3 ± 0.1
6B	6.8 ± 0.2
C-6B	6.8 ± 0.2

Assay 5: Lung S9 Metabolism

5 The in vitro metabolic stability of test compounds were evaluated in human lung S9 fraction (1 μM compound; 1 mg/mL S9 protein). The time 0, 15, 30, and 60 minute samples were analyzed for parent compound by high resolution LC-MS/MS. Lung S9 fractions from human (lot 1410245) were purchased from XenoTech LLC (Lenexa, KS). NADPH (Sigma Aldrich, N1630) and 3-phosphoadenosine 5-phosphosulfate (PAPS) (Sigma Aldrich, A1651) were purchased from Sigma Aldrich (St. Louis, MO). Acetonitrile and water were obtained from VWR (Radnor, PA) and were of HPLC grade or better. Raloxifene and formic acid was purchased from Sigma Aldrich (St. Louis, MO). Lung S9 incubations were performed in a water bath at 37°C in a 96-well polypropylene plate. Lung S9 solutions consisted of 100mM potassium phosphate buffered to pH 7.4 (BD Biosciences, Woburn, MA) supplemented with 15 1 mM NADPH (Sigma-Aldrich, St. Louis, MO), 3 mM magnesium chloride (Sigma Aldrich,

M1028) and in the presence of 100 μ M PAPS (Sigma-Aldrich, St. Louis, MO) cofactor, with final incubation protein concentrations of 1mg/mL. 10mM DMSO stocks of Raloxifene (n=1) and compound (n=1) were diluted in buffer and spiked into the incubations to yield 1 μ M substrate concentrations (0.001% DMSO v/v). Incubation volumes consisted of 400 μ L and time points were taken at 0, 15, 30, and 60 minutes by the removal of a 70 μ L aliquot and dilution into 140 μ L acetonitrile (0% formic acid). All samples were centrifuged at 2250 g for 10 minutes at 5°C. Supernatant (50 μ L) was taken from the centrifuged samples and diluted into 100 μ L HPLC water containing internal standard. The samples were run on a Dionex Ultimate 3000 Auto sampler and analyzed using a Thermo Q-Exactive High Resolution Mass Spectrometer (Thermo, Waltham, MA) in Full Scan mode in conjunction with an Atlantis T3 column 3 μ M – 2.1 x 50 mm (Waters Inc., 186003717). Mobile Phase A consisted of Water+0.2% formic acid and Mobile Phase B consisted of acetonitrile +0.2% formic acid. Peak integration was accomplished using Gubbs GMSU software (Gubbs Inc., Alpharetta, GA). For each sample, peak area ratios were calculated by dividing the analyte peak area by the internal standard peak area. For each incubation, the peak area ratios of the analytes in each t0 was set to 100%, and the peak areas ratios from the 60 minute samples were converted to percentages remaining relative to the corresponding t0. Determination of sulfate metabolite formation was made qualitatively by observation of early-eluting peak in the parent ion channel which, based on historical internal data, corresponded to the O-sulfate metabolite of each parent compound. The results of the assay are summarized in Table 4 (n=2 replicate).

Table 4: Metabolic Stability in Human Lung S9 Fraction

Compound	Clearance (μ L/min/mg)	Compound Remaining at 60 min (%)	Sulfate Appearance
1	3.5	81	Yes
C-1	49.0	6	Yes
2	0.6	98	Yes
C-2	46.7	6	Yes
3	0.8	97	Yes
C-3	50.6	5	Yes
4	0.0	100	Yes

C-4	53.2	4	Yes
5	1.6	96	Yes
C-5	50.5	5	Yes
6	1.8	90	Yes
C-6	53.5	4	Yes
1B	4.2	78	Yes
C-1B	20.8	29	Yes
2B	3.5	81	Yes
C-2B	61.0	3	Yes
3B	2.6	86	Yes
C-3B	33.2	14	Yes
4B	2.1	88	Yes
C-4B	53.4	4	Yes
5B	3.7	80	Yes
C-5B	42.2	8	Yes
6B	3.4	82	Yes
C-6B	51.4	5	Yes

When compared to their corresponding fluoro analog (compounds **C-1** to **C-6** and **C-1B** to **C-6B**), compounds **1-6** and **1B-6B** gave rise to significantly less sulfation metabolism.

Assay 6: Pharmacokinetics in Plasma and Lung in Mouse

5 Plasma and lung concentrations of test compounds and ratios thereof were determined in the following manner. BALB/c mice from Charles River Laboratories were used in the assay. Test compounds were individually formulated in 20 % propylene glycol in pH 4 citrate buffer at a concentration of 0.2 mg/mL and 50 µL of the dosing solution was introduced into the trachea of a mouse by oral aspiration. At various time points (typically
10 0.167, 2, 6, 24 hr) post dosing, blood samples were removed via cardiac puncture and intact lungs were excised from the mice. Blood samples were centrifuged (Eppendorf centrifuge, 5804R) for 4 minutes at approximately 12,000 rpm at 4°C to collect plasma. Lungs were padded dry, weighed, and homogenized at a dilution of 1:3 in sterile water. Plasma and lung concentrations of test compound were determined by LC-MS analysis against analytical

standards constructed into a standard curve in the test matrix. A lung-to-plasma ratio was determined as the ratio of the lung AUC in $\mu\text{g hr/g}$ to the plasma AUC in $\mu\text{g hr/mL}$, where AUC is conventionally defined as the area under the curve of test compound concentration vs. time.

5

Table 5: Plasma and Lung Tissue Exposure Following a Single Oral Aspiration Administration of Test Compounds

Compound	Plasma AUC ₍₀₋₂₄₎ ($\mu\text{g hr/mL}$)	Lung Tissue AUC ₍₀₋₂₄₎ ($\mu\text{g hr/g}$)	Lung Tissue:Plasma AUC ratio
1	0.943	54.5	57.8
1B	0.581	64.6	111
5B	0.599	29.9	49.9

Assay 7: IL-5 mediated eosinophil survival assay

10

The potency of test compounds for IL-5 mediated eosinophil survival can be measured in human eosinophils isolated from human whole blood (AllCells). Because IL-5 signals through JAK, this assay provides a measure of JAK cellular potency.

15

Human eosinophils are isolated from fresh human whole blood (AllCells) of healthy donors. Blood is mixed with 4.5% Dextran (Sigma-Aldrich) in a 0.9% sodium chloride solution (Sigma-Aldrich). Red blood cells are left to sediment for 35 minutes. The leukocyte rich upper layer is removed and layered over Ficoll-Paque (GE Healthcare) and centrifuged at 600g for 30 minutes. The plasma and mononuclear cell layers are removed before the granulocyte layer is lysed with water to remove any contaminating red blood cells. Eosinophils are further purified using a human eosinophil isolation kit (Miltenyi Biotec). A fraction of the purified eosinophils is incubated with anti-CD16 FITC (Miltenyi Biotec) for 10 minutes at 4°C in the dark. Purity is analyzed using a LSRII flow cytometer (BD Biosciences).

20

25

Cells are cultured in a 37°C, 5% CO₂ humidified incubator in RPMI 1640 (Life Technologies) supplemented with 10% Heat Inactivated Fetal Bovine Serum (FBS, Life Technologies), 2mM Glutamax (Life Technologies), 25mM HEPES (Life Technologies) and

1X Pen/Strep (Life Technologies). Cells are seeded at 10,000 cells/well in media (50 μ L). The plate is centrifuged at 300g for 5 minutes and supernatants removed. Compounds are serially diluted in DMSO and then diluted another 500-fold to a 2x final assay concentration in media. Test compounds (50 μ L/well) are added to cells, and incubated at 37 $^{\circ}$ C, 5 % CO₂ for 1 hour, followed by the addition of IL-5 (R&D Systems; final concentrations 1 ng/mL and 10 pg/mL) in pre-warmed assay media (50 μ L) for 72 hours.

After cytokine stimulation, cells are centrifuged at 300 g for 5 min and washed twice with cold DPBS (Life Technologies). To assess viability and apoptosis, cells are incubated with Propidium Iodide (Thermo Fisher Scientific) and APC Annexin V (BD Biosciences) and analyzed using a LSRII flow cytometer (BD Biosciences). IC₅₀ values are determined from analysis of the viability curves of percent cell viability vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values.

Assay 8: Inhibition of IFN γ and IL-27 induced chemokines CXCL9 and CXCL10 in human 3D airway cultures

EpiAirway tissue cultures were obtained from Mattek (AIR-100). Cultures were derived from asthmatic donors. In a cell culture insert, human derived tracheal/bronchial epithelial cells were grown and differentiated on a porous membrane support, allowing an air-liquid interface with warmed culture medium below the cells and a gaseous test atmosphere above. Tissues were cultured in maintenance media (Mattek, AIR-100-MM) in a 37 $^{\circ}$ C, 5% CO₂ humidified incubator. Four donors were tested. On Day 0, tissue cultures were treated with test compounds in liquid interface at 10 μ M, 1 μ M and/or 0.1 μ M. Compounds were diluted in dimethyl sulfoxide (DMSO, Sigma) to a final concentration of 0.1%. DMSO at 0.1% was used as a vehicle control. Test compounds were incubated with cultures for 1 hour at 37 $^{\circ}$ C, 5% CO₂, followed by the addition of pre-warmed media containing IFN γ (R&D Systems) or IL-27 (R&D Systems) at a final concentration of 100ng/ml. Tissue cultures were maintained for 8 days. Media was replaced every 2 days with fresh media containing compounds and IFN γ or IL-27. On Day 8, tissue cultures and supernatants were collected for analysis. Supernatant samples were assayed for CXCL10 (IP-10) and CXCL9 (MIG) using luminex analysis (EMD Millipore). Data is expressed as % Inhibition +/- standard deviation (\pm STDV). Percent inhibition was determined by compound inhibitory potency against IFN γ or IL-27 induced CXCL10 or CXCL9 secretion compared to

vehicle treated cells. Data is the average from 4 donors. Compound 1 was able to inhibit IFN γ induced CXCL10 secretion by 100% \pm 1.0 (at 10 μ M), 76% \pm 13 (at 1 μ M) and 18% \pm 22 (at 0.1 μ M) when compared to vehicle control. Compound 1 was able to inhibit IFN γ induced CXCL9 secretion by 100% \pm 0.1 (at 10 μ M), 93% \pm 6.9 (at 1 μ M) and 16% \pm 41 (at 0.1 μ M) when compared to vehicle. Compound 1 was able to inhibit IL-27 induced CXCL10 secretion by 100% \pm 0.0 (at 10 μ M), 98% \pm 1.0 (at 1 μ M) and 25% \pm 26 (at 0.1 μ M) when compared to vehicle control. Compound 1 was able to inhibit IL-27 induced CXCL9 secretion by 100% \pm 0.0 (at 10 μ M), 97% \pm 2.0 (at 1 μ M) and 52% \pm 18 (at 0.1 μ M) when compared to vehicle control.

10 **Assay 9: Cellular JAK Potency Assay: Inhibition of IL-2/anti-CD3 Stimulated IFN γ in human PBMCs**

The potency of the test compounds for inhibition of interleukin-2 (IL-2)/anti-CD3 stimulated interferon gamma (IFN γ) can be measured in human peripheral blood mononuclear cells (PBMCs) isolated from human whole blood (Stanford Blood Center).

15 Because IL-2 signals through JAK, this assay provides a measure of JAK cellular potency.

(1) Human peripheral blood mononuclear cells (PBMC) are isolated from human whole blood of healthy donors using a ficoll gradient. Cells are cultured in a 37 °C, 5 % CO₂ humidified incubator in RPMI (Life Technologies) supplemented with 10 % Heat Inactivated Fetal Bovine Serum (FBS, Life Technologies), 2 mM Glutamax (Life Technologies), 25 mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). Cells are seeded at 200,000 cells/well in media (50 μ L) and cultured for 1 h. Compounds are serially diluted in DMSO and then diluted another 500-fold (to a 2x final assay concentration) in media. Test compounds (100 μ L/well) are added to cells, and incubated at 37 °C, 5 % CO₂ for 1 h, followed by the addition of IL-2 (R&D Systems; final concentration 100 ng/mL) and anti-CD3 (BD Biosciences; final concentration 1 μ g/mL) in pre-warmed assay media (50 μ L) for 24 h.

(2) After cytokine stimulation, cells are centrifuged at 500 g for 5 min and supernatants removed and frozen at -80 °C. To determine the inhibitory potency of the test compound in response to IL-2/anti-CD3, supernatant IFN γ concentrations are measured via ELISA (R&D Systems). IC₅₀ values are determined from analysis of the inhibition curves of 30

concentration of IFN γ vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values.

Assay 10: Cellular JAK Potency Assay: Inhibition of IL-2 Stimulated pSTAT5 in CD4+ T cells

5 The potency of test compounds for inhibition of interleukin-2 (IL-2)/anti-CD3 stimulated STAT5 phosphorylation can be measured in CD4-positive (CD4+) T cells in human peripheral blood mononuclear cells (PBMCs) isolated from human whole blood (Stanford Blood Center) using flow cytometry. Because IL-2 signals through JAK, this assay provides a measure of JAK cellular potency.

10 CD4+ T cells are identified using a phycoerythrin (PE) conjugated anti-CD4 antibody (Clone RPA-T4, BD Biosciences), while an Alexa Fluor 647 conjugated anti-pSTAT5 antibody (pY694, Clone 47, BD Biosciences) is used to detect STAT5 phosphorylation.

(1) The protocol of Assay 9 paragraph (1) is followed with the exception that the
15 cytokine stimulation with anti-CD3 is performed for 30 min instead of 24 h.

(2) After cytokine stimulation, cells are fixed with pre warmed fix solution (200 μ L; BD Biosciences) for 10 min at 37 $^{\circ}$ C, 5 % CO₂, washed twice with DPBS buffer (1 mL, Life Technologies), and resuspended in ice cold Perm Buffer III (1000 μ L, BD Biosciences) for 30 min at 4 $^{\circ}$ C. Cells are washed twice with 2 % FBS in DPBS (FACS buffer), and then
20 resuspended in FACS buffer (100 μ L) containing anti-CD4 PE (1:50 dilution) and anti-CD3 anti-CD3Alexa Fluor 647 (1:5 dilution) for 60 min at room temperature in the dark. After incubation, cells are washed twice in FACS buffer before being analyzed using a LSRII flow cytometer (BD Biosciences). To determine the inhibitory potency of test compounds in response to IL-2/anti-CD3, the median fluorescent intensity (MFI) of pSTAT5 is measured in
25 CD4+ T cells. IC₅₀ values are determined from analysis of the inhibition curves of MFI vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values.

Assay 11: Cellular JAK Potency Assay: Inhibition of IL-4 Stimulated pSTAT6 in CD3+ T cells

30 The potency of test compounds for inhibition of interleukin-4 (IL-4) stimulated STAT6 phosphorylation can be measured in CD3-positive (CD3+) T cells in human

peripheral blood mononuclear cells (PBMCs) isolated from human whole blood (Stanford Blood Center) using flow cytometry. Because IL-4 signals through JAK, this assay provides a measure of JAK cellular potency.

CD3⁺ T cells are identified using a phycoerythrin (PE) conjugated anti-CD3 antibody (Clone UCHT1, BD Biosciences), while an Alexa Fluor 647 conjugated anti-pSTAT6 antibody (pY641, Clone 18/P, BD Biosciences) is used to detect STAT6 phosphorylation.

Human peripheral blood mononuclear cells (PBMC) are isolated from human whole blood of healthy donors as in Assays 9 and 10. Cells are seeded at 250,000 cells/well in media (200 μ L), cultured for 1 h and then resuspended in assay media (50 μ L) (RPMI supplemented with 0.1% bovine serum albumin (Sigma), 2mM Glutamax, 25mM HEPES and 1X Penstrep) containing various concentrations of test compounds. Compounds are serially diluted in DMSO and then diluted another 500-fold (to a 2x final assay concentration) in assay media. Test compounds (50 μ L) are incubated with cells at 37°C, 5% CO₂ for 1 h, followed by the addition of IL-4 (50 μ L) (R&D Systems; final concentration 20 ng/mL) in pre-warmed assay media for 30 min. After cytokine stimulation, cells are fixed with pre-warmed fix solution (100 μ L) (BD Biosciences) for 10 min at 37°C, 5% CO₂, washed twice with FACS buffer (1 mL) (2% FBS in DPBS), and resuspended in ice cold Perm Buffer III (1000 μ L) (BD Biosciences) for 30 min at 4°C. Cells are washed twice with FACS buffer, and then resuspended in FACS buffer (100 μ L) containing anti-CD3 PE (1:50 dilution) and anti-pSTAT6 Alexa Fluor 647 (1:5 dilution) for 60 min at room temperature in the dark. After incubation, cells are washed twice in FACS buffer before being analyzed using a LSRII flow cytometer (BD Biosciences).

To determine the inhibitory potency of the test compound in response to IL-4, the median fluorescent intensity (MFI) of pSTAT6 is measured in CD3⁺ T cells. IC₅₀ values are determined from analysis of the inhibition curves of MFI vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀).

Assay 12: Cellular JAK Potency Assay: Inhibition of IL-6 Stimulated pSTAT3 in CD3⁺ T cells

A protocol analogous to that of Assay 11 can be used to determine the potency of the test compound for inhibition of interleukin-6 (IL-6) stimulated STAT3 phosphorylation. An

Alexa Fluor 647 conjugated anti-pSTAT3 antibody (pY705, Clone 4/P, BD Biosciences) is used to detect STAT3 phosphorylation.

Assay 13: Murine model of *Alternaria alternata*-induced eosinophilic inflammation of the lung

5 Airway eosinophilia is a hallmark of human asthma. *Alternaria alternata* is a fungal aeroallergen that can exacerbate asthma in humans and induces eosinophilic inflammation in the lungs of mice (Havaux et al. *Clin Exp Immunol.* **2005** Feb;139(2):179-88). In mice, it has been demonstrated that alternaria indirectly activates tissue resident type 2 innate lymphoid cells in the lung, which respond to (e.g. IL-2 and IL-7) and release JAK-dependent cytokines
10 (e.g. IL-5 and IL-13) and coordinate eosinophilic inflammation (Bartemes et al. *J Immunol.* **2012** Feb 1;188(3):1503-13).

 Seven- to nine-week old male C57 mice from Taconic can be used in the study. On the day of study, animals are lightly anesthetized with isoflurane and administered either vehicle or test compound (0.03 – 1.0 mg/mL, 50 µL total volume over several breaths) via
15 oropharyngeal aspiration. Animals are placed in lateral recumbency post dose and monitored for full recovery from anesthesia before being returned to their home cage. One hour later, animals are once again briefly anesthetized and challenged with either vehicle or alternaria extract (200 µg total extract delivered, 50 µL total volume) via oropharyngeal aspiration before being monitored for recovery from anesthesia and returned to their home cage. Forty-
20 eight hours after alternaria administration, bronchoalveolar lavage fluid (BALF) is collected and eosinophils are counted in the BALF using the Advia 120 Hematology System (Siemens). Activity in the model is evidenced by a decrease in the level of eosinophils present in the BALF of treated animals at forty-eight hours compared to the vehicle treated, alternaria challenged control animals. Data are expressed as percent inhibition of the vehicle
25 treated, alternaria challenged BALF eosinophils response. To calculate percent inhibition, the number of BALF eosinophils for each condition is converted to percent of the average vehicle treated, alternaria challenged BALF eosinophils and subtracted from one-hundred percent.

Assay 14: Cellular JAK Potency Assay: Inhibition of IFN γ -Induced pSTAT1

30 The potency of test compounds for inhibition of interferon gamma (IFN γ) stimulated STAT1 phosphorylation was measured in CD14-positive (CD14+) monocytes derived from

human whole blood (Stanford Blood Center) using flow cytometry. Because IFN γ signals through JAK, this assay provides a measure of JAK cellular potency.

Monocytes were identified using a fluorescein isothiocyanate (FITC) conjugated anti-CD14 antibody (Clone RM052, Beckman Coulter), and an Alexa Fluor 647 conjugated anti-pSTAT1 antibody (pY701, Clone 4a, BD Biosciences) was used to detect STAT1 phosphorylation.

Human peripheral blood mononuclear cells (PBMC) were isolated from human whole blood of healthy donors using a ficoll gradient. Cells were cultured in a 37 °C, 5 % CO₂ humidified incubator in RPMI (Life Technologies) supplemented with 10 % Fetal Bovine Serum (FBS, Life Technologies), 2 mM Glutamax (Life Technologies), 25 mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). Cells were seeded at 250,000 cells/well in media (200 μ L), cultured for 2 h and re-suspended in assay media (50 μ L) (RPMI supplemented with 0.1 % bovine serum albumin (Sigma), 2 mM Glutamax, 25 mM HEPES and 1X Penstrep) containing various concentrations of test compound. The compound was serially diluted in DMSO and then diluted another 1000-fold in media to bring the final DMSO concentration to 0.1 %. Test compound dilutions were incubated with cells at 37 °C, 5 % CO₂ for 1 h, followed by the addition of pre-warmed IFN γ (R&D Systems) in media (50 μ L) at a final concentration of 0.6 ng/mL for 30 min. After cytokine stimulation, cells were fixed with pre-warmed fix solution (100 μ L) (BD Biosciences) for 10 min at 37 °C, 5 % CO₂, washed twice with FACS buffer (1 mL) (1% BSA in PBS), re-suspended in 1:10 anti-CD14 FITC:FACS buffer (100 μ L), and incubated at 4 °C for 15 min. Cells were washed once, and then re-suspended in ice cold Perm Buffer III (BD Biosciences) (100 μ L) for 30 min at 4 °C. Cells were washed twice with FACS buffer, and then re-suspended in 1:10 anti-pSTAT1 Alexa Fluor 647:FACS buffer (100 μ L) for 30 min at RT in the dark, washed twice in FACS buffer, and analyzed using a MACSQuant flow cytometer (Miltenyi).

To determine the inhibitory potency of the test compound, the median fluorescent intensity (MFI) of pSTAT1 was measured in CD14+ monocytes. IC₅₀ values were determined from analysis of the inhibition curves of MFI vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values. Compound 1 exhibited a

pIC₅₀ value of 7.5 in this assay. Compound **1B** exhibited a pIC₅₀ value of 7.5 in this assay. Compound **4B** exhibited a pIC₅₀ value of 7.3 in this assay.

Assay 15: Cellular JAK Potency Assay: Inhibition of GM-CSF-Induced pSTAT5

5 The potency of test compounds for inhibition of granulocyte-macropage colony-stimulating factor (GM-CSF) stimulated STAT5 phosphorylation was measured in CD14-positive (CD14+) monocytes derived from human whole blood (Stanford Blood Center) using flow cytometry. Because GM-CSF signals through JAK, this assay provides a measure of JAK cellular potency.

10 Monocytes were identified using a fluorescein isothiocyanate (FITC) conjugated anti-CD14 antibody (Clone RM052, Beckman Coulter), and an Alexa Fluor 647 conjugated anti-pSTAT5 antibody (pY694, BD Biosciences) was used to detect STAT5 phosphorylation.

Human peripheral blood mononuclear cells (PBMC) were isolated from human whole blood of healthy donors using a ficoll gradient. Cells were cultured in a 37 °C, 5 % CO₂ humidified incubator in RPMI (Life Technologies) supplemented with 10 % Fetal Bovine Serum (FBS, Life Technologies), 2 mM Glutamax (Life Technologies), 25 mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). Cells were seeded at 250,000 cells/well in media (200 µL), cultured for 2 h and re-suspended in assay media (50 µL) (RPMI supplemented with 0.1 % bovine serum albumin (Sigma), 2 mM Glutamax, 25 mM HEPES and 1X Penstrep) containing various concentrations of test compounds. The compound was serially diluted in DMSO and then diluted another 1000-fold in media to bring the final DMSO concentration to 0.1 %. Test compound dilutions were incubated with cells at 37 °C, 5 % CO₂ for 1 h, followed by the addition of pre-warmed GM-CSF (R&D Systems) in media (50 µL) at a final concentration of 0.3 ng/mL for 15 min. After cytokine stimulation, cells were fixed with pre-warmed fix solution (100 µL) (BD Biosciences) for 10 min at 37 °C, 5 % CO₂, washed twice with FACS buffer (1 mL) (1% BSA in PBS), re-suspended in 1:10 anti-CD14 FITC:FACS buffer (100 µL), and incubated at 4 °C for 15 min. Cells were washed once, and then re-suspended in ice cold Perm Buffer III (BD Biosciences) (100 µL) for 30 min at 4 °C. Cells were washed twice with FACS buffer, and then re-suspended in 1:10 anti-pSTAT1 Alexa Fluor 647:FACS buffer (100 µL) for 30 min at RT in

the dark, washed twice in FACS buffer, and analyzed using a MACSQuant flow cytometer (Miltenyi).

To determine the inhibitory potency of the test compound, the median fluorescent intensity (MFI) of pSTAT5 was measured in CD14+ monocytes. IC₅₀ values were
5 determined from analysis of the inhibition curves of MFI vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values. Compound 1 exhibited a pIC₅₀ value of 6.9 in this assay. Compound 1B exhibited a pIC₅₀ value of 6.7 in this assay. Compound 4B exhibited a pIC₅₀ value of 6.7 in this assay.

Assay 16: Cellular JAK Potency Assay: Inhibition of IL-12-Induced pSTAT4

10 The potency of test compounds for inhibition of interleukin-12 (IL-12) stimulated STAT4 phosphorylation was measured in CD3-positive (CD3+) T cells derived from human whole blood (Stanford Blood Center) using flow cytometry. Because IL-12 signals through JAK, this assay provides a measure of JAK cellular potency.

CD3+ T cells were identified using a phycoerythrin (PE) conjugated anti-CD3
15 antibody (clone UCHT1, BD Biosciences), and an Alexa Fluor 647 conjugated anti-pSTAT4 antibody (clone 38/p-Stat4, BD Biosciences) was used to detect STAT4 phosphorylation.

Human peripheral blood mononuclear cells (PBMC) were isolated from human whole blood of healthy donors using a ficoll gradient. Cells were cultured in RPMI (Life Technologies) supplemented with 10 % Fetal Bovine Serum (FBS, Life Technologies), 2
20 mM Glutamax (Life Technologies), 25 mM HEPES (Life Technologies), 1X Pen/Strep (Life Technologies), plate bound purified anti-CD3 antibody (5µg/ml, clone UCHT1, BD Biosciences) and soluble anti-CD28 antibody (1µg/ml, clone CD28.2, BD Biosciences) for 3 days in a 37 °C, 5 % CO₂ humidified incubator. Cells were harvested, washed with media and then re-suspended in media containing interleukin-2 (IL-2, 10ng/ml, R&D Systems).

25 Cells were cultured for 3 days in a 37°C, 5 % CO₂ humidified incubator. Cells were harvested, washed with RPMI and seeded at 250,000 cells/well in media (200 µL), cultured for 2 h and re-suspended in assay media (50 µL) (RPMI supplemented with 0.1 % bovine serum albumin (Sigma), 2 mM Glutamax, 25 mM HEPES and 1X Penstrep) containing various concentrations of test compounds. The compound was serially diluted in DMSO and
30 then diluted another 1000-fold in media to bring the final DMSO concentration to 0.1 %. Test compound dilutions were incubated with cells at 37 °C, 5 % CO₂ for 1 h, followed by

the addition of pre-warmed IL-12 (R&D Systems) in media (50 μ L) at a final concentration of 10 ng/mL for 30 min. After cytokine stimulation, cells were fixed with pre-warmed fix solution (100 μ L) (BD Biosciences) for 10 min at 37 $^{\circ}$ C, 5 % CO₂, washed twice with FACS buffer (1 mL) (1% BSA in PBS), and re-suspended in ice cold Perm Buffer III (1000 μ L) (BD Biosciences) for 30 min at 4 $^{\circ}$ C. Cells were washed twice with FACS buffer, and then re-suspended in FACS buffer (100 μ L) containing anti-CD3 PE (1:50 dilution) and anti-pSTAT4 Alexa Fluor 647 (1:10 dilution) for 45 min at room temperature in the dark. After incubation, cells were washed twice in FACS buffer before being analyzed using a MACSQuant flow cytometer (Miltenyi). To determine the inhibitory potency of the test compound, the median fluorescent intensity (MFI) of pSTAT4 was measured in CD3+ T cells. IC₅₀ values were determined from analysis of the inhibition curves of MFI vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values. Compound **1** exhibited a pIC₅₀ value of 6.0 in this assay. Compound **1B** exhibited a pIC₅₀ value of 6.2 in this assay. Compound **4B** exhibited a pIC₅₀ value of 6.0 in this assay.

15 **Assay 17: Inhibition of IFN γ secretion in a Mixed Lymphocyte Reaction Assay**

The mixed lymphocyte reaction assay is an in-vitro assay that mimics transplant rejection. T cells from one donor are cultured with allogeneic dendritic cells from another donor. This reaction induces a cellular immune response such as IFN γ secretion.

CD14+ monocytes were isolated from human whole blood (Stanford blood center) of donor A using a ficoll gradient and magnetic separation (CD14 microbeads, Miltenyi). Monocytes were differentiated into dendritic cells by culturing cells in RPMI (Life Technologies) supplemented with 10 % Fetal Bovine Serum (FBS, Life Technologies), 2 mM Glutamax (Life Technologies), 1X Pen/Strep (Life Technologies), interleukin-4 (IL-4, 50ng/ml, R&D Systems) and granulocyte-macrophage colony-stimulating factor (GM-CSF, 50ng/ml, R&D Systems) for 6 days in a 37 $^{\circ}$ C, 5 % CO₂ humidified incubator. Dendritic cells were harvested, washed with media and then activated by culturing cells in media containing lipopolysaccharide from *Escherichia coli* (LPS, 100ng/ml, Sigma) for 24 hours in a 37 $^{\circ}$ C, 5 % CO₂ humidified incubator. Cells were harvested, washed with media, re-suspended to 400,000 cells/ml in media and plated at 10,000 cells/well/25 μ l. CD4+ T cells were freshly isolated from human whole blood (Stanford blood center) of donor B using a ficoll gradient and magnetic separation (CD4+ T cell isolation kit, Miltenyi). T cells were re-

suspended to 4,000,000 cells/ml in RPMI (Life Technologies) supplemented with 10 % Fetal Bovine Serum (FBS, Life Technologies), 2 mM Glutamax (Life Technologies), 25 mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). CD4⁺ T cells were mixed with the dendritic cells at 100,000 cells/well/25 μ l. Cells were treated with test compounds (50 μ l at 20 μ M, 2 μ M and/or 0.2 μ M) to a final concentration of 10 μ M, 1 μ M and/or 0.1 μ M. Compounds were diluted in dimethyl sulfoxide (DMSO, Sigma) to a final concentration of 0.1%. DMSO at 0.1% was used as a vehicle control. Cells were maintained for 5 days in a 37°C, 5 % CO₂ humidified incubator. On day 5, supernatants were collected and measured for interferon gamma (IFN γ) using enzyme linked immunosorbent assay (ELISA). Percent inhibition was determined by compound inhibitory potency against IFN γ secretion compared to vehicle treated cells. Data is the average from 4 donors. Compound 1 was able to inhibit IFN γ secretion by 99% \pm 0.4 (at 10 μ M), 76% \pm 10 (at μ M) and 43% \pm 12 (at 0.1 μ M) when compared to vehicle control.

Assay 18: Inhibition of spontaneous periostin and IL-6 secretion in human 3D airway cultures derived from asthmatic donors

EpiAirway tissue cultures were obtained from Mattek (AIR-100). Cells were derived from asthmatic donors that spontaneously secrete periostin, a matricellular protein associated with Th2 mediated asthma (eosinophilic), and interleukin-6 (IL-6), an inflammatory cytokine that plays a role in both Th2 and non-Th2 related asthma. In a cell culture insert, human derived tracheal/bronchial epithelial cells were grown and differentiated on a porous membrane support, allowing an air-liquid interface with warmed culture medium below the cells and a gaseous test atmosphere above. Tissues were cultured in maintenance media (Mattek, AIR-100-MM) in a 37°C, 5% CO₂ humidified incubator. Four donors were tested. On Day 0, tissue cultures were treated with test compounds in liquid interface at 10 μ M, 1 μ M and/or 0.1 μ M. Compounds were diluted in dimethyl sulfoxide (DMSO, Sigma) to a final concentration of 0.1%. DMSO at 0.1% was used as a vehicle control. Tissue cultures were maintained for 8 days. Media was replaced every 2 days with fresh media containing compounds. On Day 8, supernatants were collected for analysis. Supernatant samples were assayed for periostin and interleukin-6 (IL-6) using luminex analysis (EMD Millipore). Data is expressed as % Inhibition \pm standard deviation (\pm STDV). Percent inhibition was determined by compound inhibitory potency against spontaneous secretion of periostin and

IL-6 compared to vehicle treated cells. Data is the average from 3 or 4 donors. Compound 1 was able to inhibit spontaneous periostin secretion by 62% ±25 (at 10µM) and 40% ±28 (at 1µM) when compared to vehicle control. Compound 1 was able to inhibit spontaneous IL-6 secretion by 91% ±9.0 (at 10µM), 70% ±33 (at 1µM) and 10% ±40 (at 0.1µM) when compared to vehicle.


Crystal Structure

A co-crystal structure was obtained of compound C-1 bound to human JAK1 at a resolution of 2.28 Å. The ligand was observed to bind in the ATP binding site. Seven specific hydrogen bonding interactions were identified based upon a distance of 3.5 Å or less between donor and acceptor atoms. Of particular note, a hydrogen bonding interaction was identified between the carbonyl of the exocyclic amide of the compound of C-1 and the sidechain of Arg879 of JAK1. A similar interaction can be expected for the compounds of the invention. In earlier modeling studies this interaction had been proposed as a way to provide selectivity for JAK1 over other tyrosine kinases, as otherwise closely related kinases (e.g. TRKA, VEGFR, ABL1) did not possess an arginine residue at the equivalent location. The observed results of the hydrogen bonding interaction in the crystal structure and improved kinome selectivity compared to series not possessing the exocyclic amide validate this design hypothesis.

While the present invention has been described with reference to specific aspects or embodiments thereof, it will be understood by those of ordinary skilled in the art that various changes can be made or equivalents can be substituted without departing from the true spirit and scope of the invention.

WHAT IS CLAIMED IS:

1. A compound of formula (I):



5

(I)

wherein:


n is 0, 1 or 2;

R¹ is C₁₋₃ alkyl; and

each R² is independently C₁₋₃ alkyl;

10 or a pharmaceutically-acceptable salt thereof.


2. The compound of Claim 1, or a pharmaceutically-acceptable salt thereof, having the formula (II):



15

(II)

3. The compound of Claim 1, or a pharmaceutically-acceptable salt thereof, having the formula (III):



20

(III)


4. The compound of any one of Claims 1 to 3, or a pharmaceutically-acceptable salt thereof, wherein R¹ is selected from the group consisting of ethyl, propyl, and isopropyl.

5. The compound of claim 1 or 2, or a pharmaceutically-acceptable salt thereof, wherein n is 0.

6. The compound of claim 1 or 2, or a pharmaceutically-acceptable salt thereof, wherein n is 1.

7. The compound of claim 1 or 2, or a pharmaceutically-acceptable salt thereof, wherein n is 1 and R² is methyl.

8. A compound of formula 1:




15

1

or a pharmaceutically-acceptable salt thereof.


9. A compound of formula 1:



20

1


10. A compound of formula 2:



2

or a pharmaceutically-acceptable salt thereof.


5 11. A compound of formula 3:



3

or a pharmaceutically-acceptable salt thereof.


10 12. A compound of formula 4:



4

or a pharmaceutically-acceptable salt thereof.


15 13. A compound of formula 5:



5

or a pharmaceutically-acceptable salt thereof.

14. A compound of formula 6:



or a pharmaceutically-acceptable salt thereof.

5

15. A pharmaceutical composition comprising a compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, and a pharmaceutically-acceptable carrier.

16. A compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal.

10

17. The compound of Claim 16 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal, wherein the respiratory disease is selected from the group consisting of asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, idiopathic pulmonary fibrosis, acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, sarcoidosis, an eosinophilic disease, a helminthic infection, pulmonary arterial hypertension, lymphangiomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome, bronchiolitis obliterans organizing pneumonia, lung graft-versus-host disease, and immune-checkpoint-inhibitor induced pneumonitis.

15

20

18. The compound of Claim 16 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal, wherein the respiratory disease is asthma or chronic obstructive pulmonary disease.

25

19. A compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, for use in the treatment of lung transplant rejection in a mammal.

20. The compound of Claim 19 or a pharmaceutically acceptable salt thereof, for use
5 in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is selected from the group consisting of primary graft dysfunction, organizing pneumonia, acute rejection, lymphocytic bronchiolitis, and chronic lung allograft dysfunction.

21. The compound of Claim 19 or a pharmaceutically acceptable salt thereof, for use
10 in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is acute lung transplant rejection.

22. The compound of Claim 19 or a pharmaceutically acceptable salt thereof, for use
15 in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is chronic lung allograft dysfunction.

23. The compound of Claim 19 or a pharmaceutically acceptable salt thereof, for use
in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is selected from the group consisting of bronchiolitis obliterans, restrictive chronic lung allograft
20 dysfunction, and neutrophilic allograft dysfunction.

24. Use of a compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a respiratory disease in a
25 mammal.

25. The use of Claim 24, wherein the respiratory disease is selected from the group consisting of asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, idiopathic pulmonary fibrosis, acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, sarcoidosis, an eosinophilic disease, a helminthic infection,
30 pulmonary arterial hypertension, lymphangiomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced pneumonitis, fungal induced pneumonitis, allergic

bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome, bronchiolitis obliterans organizing pneumonia, lung graft-versus-host disease, and immune-checkpoint-inhibitor induced
5 pneumonitis.

26. The use of Claim 24, wherein the respiratory disease is asthma or chronic obstructive pulmonary disease.

10 27. Use of a compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of lung transplant rejection in a mammal.

15 28. The use of Claim 27, wherein the lung transplant rejection is selected from the group consisting of primary graft dysfunction, organizing pneumonia, acute rejection, lymphocytic bronchiolitis, and chronic lung allograft dysfunction.

20 29. The use of Claim 27, wherein the lung transplant rejection is acute lung transplant rejection.

30 30. The use of Claim 27, wherein the lung transplant rejection is chronic lung allograft dysfunction.

25 31. The use of Claim 27, wherein the lung transplant rejection is selected from the group consisting of bronchiolitis obliterans, restrictive chronic lung allograft dysfunction, and neutrophilic allograft dysfunction.

30 32. Use of a compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, for treating a respiratory disease in a mammal.

33. The use of Claim 32, wherein the respiratory disease is selected from the group consisting of asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, idiopathic pulmonary fibrosis, acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, sarcoidosis, an eosinophilic disease, a helminthic infection, pulmonary arterial hypertension, lymphangiomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome, bronchiolitis obliterans organizing pneumonia, lung graft-versus-host disease, and immune-checkpoint-inhibitor induced pneumonitis.

34. The use of Claim 32, wherein the respiratory disease is asthma or chronic obstructive pulmonary disease.

35. Use of a compound of any one of Claims 1 to 14 or a pharmaceutically acceptable salt thereof, for treating lung transplant rejection in a mammal.


36. The use of Claim 35, wherein the lung transplant rejection is selected from the group consisting of primary graft dysfunction, organizing pneumonia, acute rejection, lymphocytic bronchiolitis, and chronic lung allograft dysfunction.

37. The use of Claim 35, wherein the lung transplant rejection is acute lung transplant rejection.

38. The use of Claim 35, wherein the lung transplant rejection is chronic lung allograft dysfunction.

39. The use of Claim 35, wherein the lung transplant rejection is selected from the group consisting of bronchiolitis obliterans, restrictive chronic lung allograft dysfunction, and neutrophilic allograft dysfunction.

40. A compound of formula (IB):



5

(IB)

wherein:

A is a 6 to 7 membered monocyclic heterocyclyl ring containing two nitrogen atoms, wherein A is connected to the carbonyl group in (I) through a nitrogen atom, A is substituted with 1 to 3 R² groups and optionally two R² groups form a -(CH₂)- bridged ring with A; or

10

A is pyrrolidinyl, wherein the pyrrolidinyl is connected to the carbonyl group in (I) through a nitrogen atom and wherein the pyrrolidinyl is substituted with NR³R⁴;

R¹ is C₁₋₃ alkyl;

each R² is independently C₁₋₃ alkyl optionally substituted with -OH;


R³ is C₁₋₃alkyl;

15

and R⁴ is C₁₋₃alkyl;

or a pharmaceutically-acceptable salt thereof.

41. The compound of Claim 40, or a pharmaceutically-acceptable salt thereof, having the formula (IB'):




20

(IB')

42. The compound of any one of Claims 40 or 41, or a pharmaceutically-acceptable salt thereof, wherein R¹ is selected from the group consisting of methyl, propyl, and isopropyl.


43. The compound of any one of Claims 40 to 42, or a pharmaceutically-acceptable salt thereof, wherein A is selected from the group consisting of piperazinyl, 2,5-diazabicyclo[2.2.1]heptanyl, and 1,4-diazacycloheptanyl, each of which is substituted with 1 or 2 R² groups, wherein each R² is independently C₁₋₃ alkyl optionally substituted with -OH, or A is pyrrolidinyl substituted with NR³R⁴, wherein R³ is C₁₋₃alkyl and R⁴ is C₁₋₃alkyl.

44. The compound of any one of Claims 40 to 43, or a pharmaceutically-acceptable salt thereof, wherein A is selected from the group consisting of:




wherein R² is C₁₋₃ alkyl optionally substituted with -OH, R³ is C₁₋₃alkyl and R⁴ is C₁₋₃alkyl.

45. A compound of formula:




or a pharmaceutically-acceptable salt thereof.

20 46. A compound of formula:



or a pharmaceutically-acceptable salt thereof.


47. A compound of formula:



5


or a pharmaceutically-acceptable salt thereof.

48. A compound of formula:



10 or a pharmaceutically-acceptable salt thereof.


49. A compound of formula:



or a pharmaceutically-acceptable salt thereof.

15

50. A compound of formula:



or a pharmaceutically-acceptable salt thereof.

51. A pharmaceutical composition comprising a compound of any one of Claims 40
5 to 50 or a pharmaceutically acceptable salt thereof, and a pharmaceutically-acceptable carrier.

52. A compound of any one of Claims 40 to 50 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal.

10 53. The compound of Claim 52 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal, wherein the respiratory disease is selected from the group consisting of asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, idiopathic pulmonary fibrosis, acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, sarcoidosis, an eosinophilic disease,
15 a helminthic infection, pulmonary arterial hypertension, lymphangiomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome,
20 bronchiolitis obliterans organizing pneumonia, lung graft-versus-host disease, and immune-checkpoint-inhibitor induced pneumonitis.

54. The compound of Claim 52 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal, wherein the respiratory disease is asthma
25 or chronic obstructive pulmonary disease.

55. A compound of any one of Claims 40 to 50 or a pharmaceutically acceptable salt thereof, for use in the treatment of lung transplant rejection in a mammal.

56. The compound of Claim 55 or a pharmaceutically acceptable salt thereof, for use in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is selected from the group consisting of primary graft dysfunction, organizing pneumonia, acute rejection, lymphocytic bronchiolitis, and chronic lung allograft dysfunction.

57. The compound of Claim 55 or a pharmaceutically acceptable salt thereof, for use in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is acute lung transplant rejection.

10

58. The compound of Claim 55 or a pharmaceutically acceptable salt thereof, for use in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is chronic lung allograft dysfunction.

59. The compound of Claim 55 or a pharmaceutically acceptable salt thereof, for use in the treatment of lung transplant rejection in a mammal, wherein the lung transplant rejection is selected from the group consisting of bronchiolitis obliterans, restrictive chronic lung allograft dysfunction, and neutrophilic allograft dysfunction.

60. Use of a compound of any one of Claims 40 to 50 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a respiratory disease in a mammal.

61. The use of Claim 60, wherein the respiratory disease is selected from the group consisting of asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, idiopathic pulmonary fibrosis, acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, sarcoidosis, an eosinophilic disease, a helminthic infection, pulmonary arterial hypertension, lymphangiomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic

pneumonia, hypereosinophilic syndrome, Löffler syndrome, bronchiolitis obliterans organizing pneumonia, lung graft-versus-host disease, and immune-checkpoint-inhibitor induced pneumonitis.

5 62. The use of Claim 60, wherein the respiratory disease is asthma or chronic obstructive pulmonary disease.

10 63. Use of a compound of any one of Claims 40 to 50 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of lung transplant rejection in a mammal.

15 64. The use of Claim 63, wherein the lung transplant rejection is selected from the group consisting of primary graft dysfunction, organizing pneumonia, acute rejection, lymphocytic bronchiolitis, and chronic lung allograft dysfunction.

 65. The use of Claim 63, wherein the lung transplant rejection is acute lung transplant rejection.

20 66. The use of Claim 63, wherein the lung transplant rejection is chronic lung allograft dysfunction.

25 67. The use of Claim 63, wherein the lung transplant rejection is selected from the group consisting of bronchiolitis obliterans, restrictive chronic lung allograft dysfunction, and neutrophilic allograft dysfunction.

 68. Use of a compound of any one of Claims 40 to 50 or a pharmaceutically acceptable salt thereof, for treating a respiratory disease in a mammal.

30 69. The use of Claim 68, wherein the respiratory disease is selected from the group consisting of asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, idiopathic pulmonary fibrosis, acute lung injury, acute respiratory distress syndrome, bronchitis,

emphysema, bronchiolitis obliterans, sarcoidosis, an eosinophilic disease, a helminthic infection, pulmonary arterial hypertension, lymphangiomyomatosis, bronchiectasis, an infiltrative pulmonary disease, drug-induced pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome, bronchiolitis obliterans organizing pneumonia, lung graft-versus-host disease, and immune-checkpoint-inhibitor induced pneumonitis.

10 70. The use of Claim 68, wherein the respiratory disease is asthma or chronic obstructive pulmonary disease.

 71. Use of a compound of any one of Claims 40 to 50 or a pharmaceutically acceptable salt thereof, for treating lung transplant rejection in a mammal.

15

 72. The use of Claim 71, wherein the lung transplant rejection is selected from the group consisting of primary graft dysfunction, organizing pneumonia, acute rejection, lymphocytic bronchiolitis, and chronic lung allograft dysfunction.

20 73. The use of Claim 71, wherein the lung transplant rejection is acute lung transplant rejection.

 74. The use of Claim 71, wherein the lung transplant rejection is chronic lung allograft dysfunction.

25


 75. The use of Claim 71, wherein the lung transplant rejection is selected from the group consisting of bronchiolitis obliterans, restrictive chronic lung allograft dysfunction, and neutrophilic allograft dysfunction.

76. The compound of Claim 16 or Claim 52 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal, wherein the respiratory disease is acute lung injury.


5 77. The compound of Claim 16 or Claim 52 or a pharmaceutically acceptable salt thereof, for use in the treatment of a respiratory disease in a mammal, wherein the respiratory disease is acute respiratory distress syndrome.

10 78. The use of any one of Claims 24, 32, 60 and 68, wherein the respiratory disease is acute lung injury.

79. The use of any one of Claims 24, 32, 60 and 68, wherein the respiratory disease is acute respiratory distress syndrome.



(I)



(IB)