

(19) **DANMARK**

(10) **DK/EP 2205593 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 D 403/04 (2006.01)** **A 61 K 31/517 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-04-18**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-01-13**
- (86) Europæisk ansøgning nr.: **08862034.9**
- (86) Europæisk indleveringsdag: **2008-10-03**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-07-14**
- (86) International ansøgning nr.: **FR2008001384**
- (87) Internationalt publikationsnr.: **WO2009077680**
- (30) Prioritet: **2007-10-03 FR 0706931**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **SANOFI, 54 rue La Boétie, 75008 Paris, Frankrig**
- (72) Opfinder: **CLAUSS, Annie, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, 75013 Paris, Frankrig**
GLAESS, Christophe, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, F-75013 Paris, Frankrig
MARCINIAK, Gilbert, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, F-75013 Paris, Frankrig
MUZET, Nicolas, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, F-75013 Paris, Frankrig
NAVE, Jean-François, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, F-75013 Paris, Frankrig
SEYER, André, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, F-75013 Paris, Frankrig
VIVET, Bertrand, c/o Sanofi-aventis, Département Brevets, 174 avenue de France, F-75013 Paris, Frankrig
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **QUINAZOLINDIONDERIVATER, FREMSTILLING OG TERAPEUTISKE ANVENDELSER HERAF**
- (56) Fremdragne publikationer:
WO-A-93/19068
PERRY M J ET AL: "CHEMOTHERAPEUTIC POTENTIAL OF PHOSPHODIESTERASE INHIBITORS" CURRENT OPINION IN CHEMICAL BIOLOGY, CURRENT BIOLOGY LTD, LONDON, GB, vol. 2, no. 4, 1 janvier 1998 (1998-01-01), pages 472-481, XP000856012 ISSN: 1367-5931
PITTS W J ET AL: "Identification of purine inhibitors of phosphodiesterase 7 (PDE7)" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 14, no. 11, 7 juin 2004 (2004-06-07), pages 2955-2958, XP004841322 ISSN: 0960-894X
LOWE J.A. ET AL: "Structure-Activity Relationship of Quinazolinone Inhibitors of Calcium-Independent Phosphodiesterase" JOURNAL OF MEDICINAL CHEMISTRY, vol. 34, 1991, pages 624-628, XP002483510
SALDOU N. ET AL: "Comparison of Recombinant Human PDE4 Isoforms: Interaction with Substrate and Inhibitors" CELLULAR SIGNALING, vol. 10, no. 6, 1998, pages 427-440, XP002483511

Fortsættes ...

DESCRIPTION

A subject-matter of the invention is quinazolinedione derivatives, the processes whereby they are obtained and their
5 therapeutic applications.

The invention relates to quinazolinedione derivatives which are inhibitors of phosphodiesterase 7 (PDE7). Some of these derivatives also inhibit phosphodiesterase 8 (PDE8).

10

Phosphodiesterases (PDEs) are intracellular enzymes responsible for the hydrolysis of cAMP (cyclic adenosine 3',5'-monophosphate) and cGMP (cyclic guanosine 3',5'-monophosphate) secondary messengers to give inactive
15 5'-monophosphate nucleotides. cAMP and cGMP play an essential role in cell signalling pathways and are involved in numerous physiological processes.

The inhibition of phosphodiesterases is reflected by an
20 increase in intracellular concentrations of cAMP and cGMP, resulting in the specific activation of phosphorylation pathways involved in varied functional responses. The increase in the intracellular concentrations of cAMP or of cGMP using selective inhibitors of phosphodiesterases appears to be a
25 promising approach in the treatment of various diseases (Bender and Beavo, Pharmacol. Rev., (2006) 58, 488-520). The inhibitors of phosphodiesterases are thus of interest as therapeutic agents and as pharmacological tools.

30 To date, eleven families of phosphodiesterases have been identified. They are distinguished by their primary structure, their substrate specificity and their sensitivity with regard to various effectors and inhibitors specific for PDEs. Each family is composed of one or more genes which are expressed in
35 various tissues in the form of splicing variants (Bender and Beavo, Pharmacol. Rev., (2006) 58, 488-520; Lugnier, Pharmacol. Therapeut., (2006) 109, 366-398).

PDE4, PDE7 and PDE8 specifically hydrolyse cAMP and PDE5, PDE6 and PDE9 specifically hydrolyse cGMP.

The family PDE7 is represented by the isoforms PDE7A and PDE7B
5 originating from two distinct genes.

Human PDE7A (Michaeli et al., J. Biol. Chem., (1993) 268, 12925-12932; Han et al., J. Biol. Chem., (1997) 272, 16152-16157; Wang et al., Biochem. Biophys. Res. Commun., (2000)
10 276, 1271-1277) and human PDE7B (Sasaki et al., Biochem. Biophys. Res. Commun., (2000), 271, 575-583; Gardner et al., Biochem. Biophys. Res. Commun., (2000) 272, 186-192) selectively hydrolyse cAMP with Michaelis constants (Km) of 0.1 to 0.2 μ M and 0.13 to 0.2 μ M respectively. The catalytic
15 part of PDE7B exhibits approximately 67% homology with that of PDE7A.

Three splicing variants are known for PDE7A. PDE7A1 and PDE7A3 are expressed mainly in the cells of the immune system and of
20 the lungs, while PDE7A2 is above all expressed in the muscles of the skeleton, the heart and the kidneys. For PDE7B, three variants have also recently been identified (Giembycz and Smith, Drugs Future, (2006) 31, 207-229).

25 The tissue distribution profiles for PDE7A and PDE7B are very different, suggesting that these two isoforms have distinct functions from the physiological viewpoint. While PDE7A is copiously expressed in haematopoietic cells, the lungs, the placenta, Leydig cells, the spleen, the collecting tubes of
30 the kidneys, and the adrenal glands, strong expression of PDE7B is detected in the pancreas, the heart, the thyroid and the muscles of the skeleton (Giembycz and Smith, Drugs Future, (2006) 31, 207-229). However, coexpression of the messenger RNAs (mRNAs) of PDE7A and PDE7B is observed in some tissues.
35 This is the case in the osteoblasts (Ahlstrom et al., Cell Mol. Biol. Lett., (2005) 10, 305-319) and in some regions of the brain: several areas of the cortex, the dentate gyrus, the majority of the components of the olfactory system, the

striatum, numerous nuclei of the thalamus and the pyramidal cells of the hippocampus (Miro et al., Synapse, (2001) 40, 201-214; Reyes-Irisarri et al., Neuroscience, (2005) 132, 1173-1185). In contrast, in some areas of the brain, only one
5 of the two isoforms is expressed. Thus, only the mRNAs of PDE7A are present in many nuclei of the brain stem. Likewise, the mRNAs of PDE7B are present at high concentrations in the nucleus accumbens and the dorsal motor nucleus of the vagus nerve, while the mRNAs of PDE7A are not detected there (Miro
10 et al., Synapse, (2001) 40, 201-214; Reyes-Irisarri et al., Neuroscience, (2005) 132, 1173-1185).

The protein PDE7A1 is well expressed in blood T lymphocytes, bronchial epithelial cell lines, lung fibroblasts and
15 eosinophils (Smith et al., Am. J. Physiol. Lung Cell Mol. Physiol., (2003) 284: L279-L289). Several reports suggest that PDE7A might play a role in the activation of T lymphocytes (Li et al., Science, (1999) 283, 848-851; Glavas et al., PNAS, (2001) 98, 6319-6324; Nakata et al., Clin. Exp. Immunol.,
20 (2002), 128, 460-466; Smith et al., Mol. Pharmacol., (2004) 66, 1679-1689). The protein PDE7A1 is also expressed in cells which play a central role in the pathogenesis of asthma and chronic obstructive pulmonary disease (COPD), such as T lymphocytes (CD4+ and CD8+), monocytes, neutrophils, alveolar
25 macrophages, or cells of the smooth muscles of the airways and pulmonary vessels (Smith et al., Am. J. Physiol. Lung Cell Mol. Physiol., (2003) 284: L279-L289). The localization of PDE7A in proinflammatory cells and cells of the immune system and its potential role in the activation of T lymphocytes
30 suggest that selective inhibitors of PDE7 might have applications in the field of diseases related to T lymphocytes and in that of diseases of the pulmonary tract.

The protein PDE7A is also present in B lymphocytes and in the
35 WSU-CLL B lymphocyte cell line originating from a patient suffering from chronic lymphocytic leukaemia. Treatment of the WSU-CLL cells with IC242, a specific inhibitor of PDE7, increases the expression of PDE7A (Lee et al., Cell Signal,


(2002) 14, 277-284). Furthermore, it has been shown that the expression of the protein PDE7B is approximately 5 to 90 times higher in peripheral blood mononuclear cells (PBMCs) of patients suffering from chronic lymphocytic leukaemia (CLL-PBMC) than in the PBMCs isolated from the blood of normal subjects (WO 2007/067946). BRL-50481, a selective inhibitor of PDE7, induces in a dose-dependent way the apoptosis of CLL-PBMCs but it is without effect on PBMCs of normal subjects. These observations suggest that selective inhibitors of PDE7 might be effective in the treatment of this type of leukaemia.

Selective inhibitors of PDE4 increase bone mineral density in the rat and the mouse and their effects appear to be related to a reduction in the activity of the osteoclasts and to an increase in that of the osteoblasts (Miyamoto et al., Biochem. Pharmacol., (1997) 54, 613-617; Waki et al., Jpn. J. Pharmacol., (1999) 79, 477-483; Kinoshita et al., Bone, (2000) 27, 811-817). A PDE7 activity has been detected in osteoblasts (Ahlström et al., Cell Mol. Biol. Lett., (2005) 10, 305-319). By increasing the intracellular concentration of cAMP, the inhibitors of PDE7, like the inhibitors of PDE4, might prove to be effective in the treatment of osteopenia and osteoporosis.

Furthermore, recent studies (WO2006/092691, WO2006/092692) report the pharmacological activities of various inhibitors of PDE7 in models of neuropathic pain in the rat, suggesting applications in the treatment of various types of pain and more particularly in that of neuropathic pain.

A subject-matter of the present invention is in particular quinazolinedione derivatives, which are powerful inhibitors of PDE7, or inhibitors of PDE7 and of PDE8 according to the derivatives, their preparation and their therapeutic applications.

The subject-matter of the invention is compounds of following general formula (I):



in which

5

- A represents an aryl group or a heteroaryl group;

- R₁ represents:

■ a hydrogen atom,

10 ■ -C(O)R in which R is a hydrogen atom, a (C₁-C₆) alkoxy group, an aryl group, a (C₃-C₆) cycloalkyl group or a (C₁-C₆) alkyl group, the said alkyl optionally being substituted by:

. one or more hydroxyl group(s),

. a benzyloxy group,

15 . a (C₁-C₆) alkoxy group, optionally substituted by an aryl, or

. a (C₃-C₆) cycloalkyl group,

■ an optionally substituted (C₁-C₆) alkyl group;

- R₂ represents:

20 ■ a hydrogen atom,

■ a halogen atom,

■ a cyano group,

■ a nitro group,

■ a (C₁-C₆) alkyl group optionally substituted by an -NH₂ or

25 else by an -NHC(O)R_b group,

■ an -OR_a group in which R_a represents:

. a hydrogen atom,

. a (C₁-C₆) alkyl group optionally substituted by one or more halogen atom(s), by one or more hydroxyl group(s), by an aryl

30 group and/or by one or more cyano group(s),

- . a (C₂-C₆) alkynyl group,
- . an aryl group;

— R₃ represents:

- 5 ■ a hydrogen atom,
- a halogen atom,
- a hydroxyl group,
- a cyano group,
- an -SCF₃ group,
- 10 ■ a nitro group,
- an oxo group,
- an -S(O)₀₋₂-alkyl group, an -S(O)₀₋₂-heterocycloalkyl group, an -O-SO₂-aryl group optionally substituted by one or more halogen atom(s);
- 15 ■ an -alkylaminoalkyl or -cycloalkylaminoalkyl group, each optionally substituted on the end alkyl,
- an optionally substituted sulphonamide group,
- an aryl group or a heteroaryl group, the said group being monocyclic or polycyclic and in addition optionally being
- 20 substituted by a (C₁-C₆) alkyl group, by one or more halogen atom(s) or by a (C₁-C₆) alkoxy group,
- a heterocycloalkyl group optionally substituted by a (C₁-C₆) alkyl group,
- a (C₁-C₆) alkyl group optionally substituted by:
 - 25 - one or more halogen atom(s),
 - an aryl group which can be substituted by one or more halogen atom(s) or by one or more hydroxyl group(s),
 - a heteroaryl group,
 - one or more hydroxyl group(s) which can be substituted by an
 - 30 aryl group itself optionally substituted by one or more halogen atom(s), or
 - a heterocycloalkyl group optionally substituted by a CO(O)Ra group or by a (C₁-C₆) alkyl group,
 - a -C(O)NR_bR_c group,
 - 35 ■ a -C(O)OR_c group or an -O-C(O)OR_c group,
 - a (C₁-C₆) alkoxy group, optionally substituted by
 - an aminoalkyl group,
 - an aminocycloalkyl group,

- a cycloalkyl group,
- a heterocycloalkyl group,
- a monocyclic or polycyclic heteroaryl group,
- one or more hydroxyl group(s),
- 5 - one or more halogen atom(s),
- a (C₁-C₆) alkoxy group,
- a -C(O)ORc group,
- a -C(O)NRbRc group,
- an oxo group, and/or
- 10 - an aryl group, itself optionally substituted by one or more halogen atom(s), a cyano group, a (C₁-C₆) alkoxy group, an -O-haloalkyl group and/or a haloalkyl group,
- an -O-cycloalkyl group, an -O-aryl group or an -O-heterocycloalkyl group, each optionally substituted by
- 15 - an aryl group, itself optionally substituted by one or more halogen atom(s) or by a (C₁-C₆) alkyl group,
- an oxo group,
- one or more halogen atom(s), and/or
- a (C₁-C₆) alkyl group, which can itself be substituted by an
- 20 aryl group and/or an oxo group,
- an -NH-CO-NH-aryl group, an -NH-CO-NH-heteroaryl group or an -NH-CO-NH-(C₁-C₆) alkyl group, each optionally being substituted by one or more halogen atom(s), by a cyano group, by a nitro group, by one or more hydroxyl group(s) or by a (C₁-
- 25 C₆) alkoxy group,
- an -N-(C₁-C₆) alkyl group, it being possible for the (C₁-C₆) alkyl group to be substituted by
- one or more oxo group(s), and/or
- one or more aryl group(s) optionally substituted by one or
- 30 more halogen atom(s) and/or by an SO₂ group,
- an -NH-CO-aryl group or an -NH-CO-heteroaryl group, each optionally being substituted by one or more halogen atom(s);

or else R₃ forms, with A, a polycyclic heteroaryl group

35 optionally substituted by a (C₁-C₆) alkoxy group or a (C₁-C₆) alkyl group optionally substituted by an aryl group which can itself be substituted by one or more halogen atom(s);

- R₄ represents a hydrogen atom, an oxo group or a (C₁-C₆) alkyl group;

- R_b represents:

- 5 . a hydrogen atom,
 . a (C₁-C₆) alkyl group optionally substituted by one or more halogen atom(s), by one or more hydroxyl, cyano, amino, heterocycloalkyl or (C₁-C₆) alkoxy group(s) or by an aryl group optionally substituted by one or more halogen atom(s),
10 . a (C₃-C₆) cycloalkyl group,
 . a (C₂-C₆) alkynyl group,
 . a (C₁-C₆) alkoxy group,
 . an aryl group optionally substituted by one or more halogen atom(s);

15

- R_c represents a hydrogen atom or a (C₁-C₆) alkyl group optionally substituted by one or more halogen atom(s);


or then R_b and R_c form, together with the nitrogen atom to
20 which they are attached, a polycyclic heteroaryl group or a heterocycloalkyl group;

- m and n represent, independently of one another, the value 0, 1 or 2, it being understood that $m+n \leq 3$;

25

- p and p' represent, independently of one another, the value 1, 2 or 3, it being understood that, when p is greater than or equal to 2, then the R₂ groups are on separate carbon atoms and can be different from one another and, when p' is greater than
30 or equal to 2, then the R₃ groups are on separate carbon atoms and can be different from one another;

- q represents the value 0 or 2, it being understood that, when q = 0, then the nitrogenous heterocyclic group attached
35 to the nitrogen situated in the 1 position of the 2,4-dioxo-1,2,3,4-tetrahydroquinazoline ring system is no longer bridged and is of the type:



- r represents the value 0 or 1.

5 The compounds of general formula (I) can comprise one or more asymmetric carbons. They can thus exist in the form of enantiomers or of diastereoisomers. These enantiomers or diastereoisomers, and their mixtures, including racemic mixtures, come within the invention.

10

Due to their structure, the compounds of general formula (I) can also exist in the form of isomers of rotamer or atropisomer type.

15 The compounds of formula (I) can also exist in the form of bases or addition salts with acids. Such addition salts come within the invention.

These salts are advantageously prepared with pharmaceutically
20 acceptable acids but the salts of other acids, for example of use in the purification or separation of the compounds of general formula (I) also come within the invention.

The compounds of general formula (I) can also occur in the
25 crystalline, amorphous or oily form, these forms coming within the invention.

The compounds of general formula (I) can in addition occur in the form of hydrates or of solvates, namely in the form of
30 combinations or of associations with one or more molecules of water or with a solvent. Such hydrates and solvates also come within the invention.

According to the present invention, the N-oxides of the

compounds comprising an amine also come within the invention.

The compounds of formula (I) according to the present invention also comprise those in which one or more hydrogen, 5 carbon or halogen, in particular chlorine or fluorine, atoms have been replaced by their radioactive isotopes, for example tritium, in order to replace hydrogen, or carbon-14, in order to replace carbon-12. Such labelled compounds are of use in research, metabolism or pharmacokinetic studies or in 10 biological and pharmacological assays as tools.

In the context of the invention, the following definitions apply:

15 - in (C_1-C_6) , the numerical indices determine the possible number of carbon atoms present in a chain or a ring. Thus, by way of example, C_1-C_6 represents a carbon chain which can have from 1 to 6 carbon atoms. Likewise, by way of example, (C_1-C_5) represents a carbon chain which can have from 1 to 5 carbon 20 atoms or also (C_3-C_6) can represent a saturated carbon ring which can have from 3 to 6 carbon atoms;

- alkoxy: an -O-alkyl group comprising a saturated, linear or branched, aliphatic chain;

25

- alkynyl: a mono- or polyunsaturated, linear or branched, aliphatic group comprising, for example, one or two acetylenic unsaturations. For example, a (C_2-C_6) alkynyl group can represent an ethynyl, propynyl, and the like;

30

- alkyl: a saturated, linear or branched, aliphatic group; for example, a (C_1-C_6) alkyl group represents a linear or branched carbon chain of 1 to 6 carbon atoms, in particular a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert- 35 butyl or pentyl;


- aminoalkyl: an $-NH(C_1-C_6)$ alkyl or also $-N((C_1-C_6) \text{ alkyl})_2$ group;

- aryl: an optionally substituted monocyclic aromatic system comprising from 5 to 14 members per ring, preferably from 5 to 10 members per ring. Mention may be made, as examples of
 5 monocyclic aryl groups, of phenyl or naphthyl; the aryl group can be substituted by a group which can be one or more halogen atom(s), a hydroxyl group, a cyano group, a trifluoromethylthio group, a nitro group, an alkyl group, an alkoxy group, an alkylthio group, a methylsulphonyl group, an
 10 alkylaminoalkyl group or alkylaminocycloalkyl group which is optionally substituted, an alkylaminoalkoxy or cycloalkylaminoalkoxy group or a sulphonamide group, for example:

15 - polycyclic aryl: an optionally substituted polycyclic aromatic system comprising from 5 to 14 members per ring, preferably from 5 to 10 members per ring, and comprising from 2 to 10 rings, at least one of the rings of which is aromatic. Mention may be made, as examples of polycyclic aryl groups, of
 20 aceanthrylene, anthracene, azulene, coronene, rubicene or naphthalene; the polycyclic aryl group can be substituted by a group which can be one or more halogen atom(s), a hydroxyl group, a cyano group, a trifluoromethylthio group, a nitro group, an alkyl group, an alkoxy group, an alkylthio group, a
 25 methylsulphonyl group, an alkylaminoalkyl or alkylaminocycloalkyl group which is optionally substituted, an alkylaminoalkoxy or cycloalkylaminoalkoxy group or a sulphonamide group, for example;

30 - a bridged ring: a bicyclic structure comprising, according to the invention, a nitrogen atom, in which at least two carbon atoms are connected via a single bond or a carbon chain which can comprise 2 carbon atoms. By way of example, the abovementioned ring is of the type:

35



- cycloalkyl: a saturated cyclic aliphatic group comprising from 3 to 8 carbon atoms. Mention may be made, by way of example, of a cyclopropyl, methylcyclopropyl, cyclobutyl, 5 cyclopentyl or cyclohexyl group;
- halogen: a fluorine, a chlorine, a bromine or an iodine;
- haloalkyl: (C₁-C₆) alkyl substituted by one to three halogen 10 atom(s);
- heteroaryl: a monocyclic aromatic system comprising from 5 to 14 ring members, preferably from 5 to 10 ring members, and comprising one to several heteroatoms, such as nitrogen, 15 oxygen or sulphur atoms. The nitrogen atoms can be in the form of N-oxides. For example, a monocyclic heterocycle can be a pyran, a pyrazine, a pyrazole, a pyridazine, a pyridine, a pyrimidine, a pyrrole, an isothiazole, an isoxazole, a furan, an imidazole, a morpholine, a thiophene, a piperazine, a 20 diazetidine, a dihydropyrrolidine, a piperidine, an azepine, and the like; a bicyclic heterocycle can be an isoquinoline, a pteridine, a chroman, and the like; a tricyclic heterocycle can be a phenanthroline, a xanthene, and the like;
- 25 - polycyclic heteroaryl: an optionally substituted polycyclic aromatic system comprising from 5 to 14 members per ring, preferably from 5 to 10 members per ring, and comprising from 2 to 10 rings, additionally comprising one to several heteroatoms, such as nitrogen, oxygen or sulphur atoms, in at 30 least one of the rings, and at least one of the rings of which is aromatic. Mention may be made, as examples of polycyclic heteroaryl groups, of indole, benzofuran, benzimidazole, benzothiophene, benzotriazole, benzothiazole, benzoxazole, quinoline, isoquinoline, indazole, quinazoline, phthalazine, 35 quinoxaline, naphthyridine, 2,3-dihydro-1H-indole, 2,3-dihydrobenzofuran, 2,3-dihydroindene, tetrahydroquinoline, tetrahydroisoquinoline or tetrahydroisoquinazoline;

- a heterocycloalkyl: an optionally substituted saturated ring comprising from 3 to 8 atoms and comprising one to several heteroatoms, such as nitrogen, oxygen or sulphur atoms, in at least one of the rings, or several heteroatoms which are
5 identical to or different from one another. For example, a heterocycloalkyl can be a pyrrolidine, a morpholine, a piperazine, a diazetidine, a dihydropyrrolidine, a piperidine, a piperadine, an azepan, an imidazolidine, a thiomorpholine, a tetrahydropyran, a tetrahydrothiopyran, a piperazine, a
10 diazepan, and the like;

- hydroxyl: an -OH group;

15 - nitro: an -NO₂ group;

- oxo: a -C(=O)- group;

- sulphonamide: group corresponding to the formula SO₂-N-alkyl or SO₂-N-cycloalkyl, alkyl and cycloalkyl being as defined
20 above;

- trifluoromethylthio is defined by the formula -S-CF₃.

Furthermore, it is understood that, in the present
25 description, when an atom or a group is substituted or optionally substituted by one or more defined group(s) or atom(s), the substituents can be identical to or different from one another and may be carried, if appropriate, by the same atom or different atoms.

30

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of formula (I) in which A represents an aryl group, in particular a phenyl or heteroaryl, in particular a pyridyl group, and all
35 the other substituents and indices are as defined in the general formula (I).

Mention may be made, among the compounds which are a subject-

matter of the invention, of a group of compounds of formula (I) in which $q = 0$, m and $n = 1$, and all the other substituents and indices are as defined in the general formula (I).

5

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of formula (I) in which R_2 represents a (C_1-C_6) alkyl group, in particular a methyl, substituted by an $-NHC(O)R_b$ group in which R_b and
10 the other groups and indices are as defined for the compound of general formula (I).

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of general
15 formula (I) in which R_2 represents an $-OR_a$ group, the R_a group and all the other groups being as defined for the compound of formula (I).

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of general
20 formula (I) in which R_2 is a halogen atom or a cyano or a hydrogen or a hydroxyl or a (C_1-C_6) alkyl group optionally substituted by an $-NH_2$ or else by an $-NHC(O)R_b$ group, the other groups being as defined for the compound of formula (I).

25

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of general formula (I) in which A is a phenyl, R_1 is a $-C(O)R$ group in which R represents a hydrogen atom, q is equal to 0, n and m
30 have the value 1, and R_2 is $-OR_a$, R_a being as defined in the general formula (I).

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of general
35 formula (I) in which A is a phenyl, R_1 is a $-C(O)R$ group in which R represents a hydrogen atom, q is equal to 0, n and m have the value 1, p is equal to 2, one of the R_2 groups is $-OR_a$, R_a being as defined in the general formula (I), and the

other of the R_2 groups is a halogen atom.

Mention may be made, among the compounds which are a subject-matter of the invention, of a group of compounds of general
 5 formula (I) in which A is a phenyl, R_1 is a $-C(O)R$ group in which R represents a hydrogen atom, q is equal to 0, n and m have the value 1 and R_2 is a methyl substituted by an $-NH-CO-R_b$ group, R_b being as defined in the general formula (I).

10 Advantageously, in the compounds of formula (I), the R_2 group is in the 6 position of the 2,4-dioxo-1,2,3,4-tetrahydroquinazoline ring system. The compounds of formula (I) can also have an R_2 group in the 7 position of the 2,4-dioxo-1,2,3,4-tetrahydroquinazoline ring system.

15 Advantageously, in the compounds of formula (I), $p = 1$ or 2.

In a specific form of the invention, when $p' = 2$, then the two R_3 groups are in the 3 and 4 positions of the ring system A and
 20 can be different from one another.

The combinations of the groups of compounds of the invention mentioned above also come within the invention as embodiments according to the invention.

25 Mention may be made, as examples of preferred compounds, of the following compounds:

No. 1: 2-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-
 30 2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propanenitrile

No. 2: 1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-hydroxyquinazoline-2,4(1H,3H)-dione

35 No. 3: {[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetonitrile

No. 4: 2-{[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-

2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propanenitrile

No. 6: { [3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-
2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetonitrile

5

No. 11: 4-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

10 No. 12: 1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione

No. 13: 4-[3-(3,4-dimethoxybenzyl)-2,4-dioxo-6-(2,2,2-trifluoroethoxy)-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-
15 carbaldehyde

No. 14: 1-(1-acetylpiperidin-4-yl)-6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione

20 No. 16: 4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 20: N-{ [3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-
25 yl]methyl}acetamide

No. 22: 1-(1-acetylpiperidin-4-yl)-6-(aminomethyl)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione hydrochloride

30 No. 23: N-{ [3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}formamide

No. 24: N-{ [1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-
35 yl]methyl}formamide

No. 25: N-{ [1-(1-acetylpiperidin-4-yl)-3-(3,4-

dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl)methyl}acetamide

No. 32: 4-[6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 33: 4-[3-(3,4-dichlorobenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

10 No. 34: 4-[3-(4-chlorobenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 35: methyl 4-{[6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}benzoate

No. 36: 4-{[6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}benzoic acid

20 No. 37: 4-{[6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}-N-(2-methoxyethyl)benzamide

No. 38: 4-[3-(3,4-dimethoxybenzyl)-6-methyl-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 39: 4-[6-(2,2-difluoroethoxy)-3-(3-hydroxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 40: 4-[6-(2,2-difluoroethoxy)-3-[3-(2-hydroxyethoxy)-4-methoxybenzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

35 No. 41: 4-[6-(2,2-difluoroethoxy)-3-(3-ethoxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 42: 4-[6-(2,2-difluoroethoxy)-3-[4-methoxy-3-(2-methoxyethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5 No. 43: 4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azepane-1-carbaldehyde

No. 47: 4-[6-(2,2-difluoroethoxy)-3-[3-(3-hydroxypropoxy)-4-methoxybenzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-
10 yl]piperidine-1-carbaldehyde

No. 48: 4-[5-chloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

15 No. 49: 4-{3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 50: 2-(5-{[6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl}-2-methoxyphenoxy)acetamide
20

No. 51: 4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]-3-methylpiperidine-1-
25 carbaldehyde

No. 52: 3-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]-8-azabicyclo[3.2.1]octane-8-carbaldehyde
30

No. 56: 4-{3-[4-(cyclopentyloxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35 No. 57: 4-[3-(3-chlorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 58: 4-[3-(4-chlorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5 No. 59: 4-{3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 72: 4-[3-(3,4-dimethoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde
10

No. 74: 4-[3-(3,4-dichlorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde
15

No. 76: 4-{3-[(6-chloropyridin-3-yl)methyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

20 No. 78: 4-[3-(3-chloro-4-methoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 79: 4-[3-(3,4-dimethoxybenzyl)-6-(2-fluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde
25

No. 89: 2-[5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]acetamide
30

No. 90: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-hydroxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35 No. 91: 4-[3-(3,4-dimethoxybenzyl)-6-ethoxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 97: 4-[5,7-dichloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-

dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 102: 4-[7-chloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5

No. 108: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-fluoro-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10 No. 111: 4-[6-(difluoromethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 112: 4-[3-(3,4-dimethoxybenzyl)-6-(1-methylethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

15

No. 114: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-methoxy-3-(1-methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

20 No. 116: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 117: 4-{3-[3,5-bis(trifluoromethyl)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

25

No. 118: 4-[3-(3-ethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

30

No. 124: 4-{3-[3-chloro-4-(2-methoxyethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35

No. 130: 4-[3-(3,4-diethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 131: 4-[3-(4-ethoxy-3-methoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5

No. 133: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-methoxy-3-methylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10 No. 134: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(trifluoromethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

15 No. 135: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(trifluoromethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

20 No. 143: 4-{3-[4-(benzyloxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 145: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-nitrobenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

25

No. 155: 4-[3-(4-ethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

30 No. 158: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35 No. 160: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-morpholin-4-ylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 165: 4-[3-(biphenyl-4-ylmethyl)-6-[2-fluoro-1-

(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 166: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(methylsulphanyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 167: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyridin-3-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 170: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-methylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 175: 2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]acetamide

No. 178: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-propoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 183: 2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-methylacetamide

No. 184: 2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N,N-dimethylacetamide

No. 185: 2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-methoxy-N-methylacetamide

No. 186: 4-{3-[4-(cyclopentyloxy)-3-ethoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

5 No. 188: 4-{3-[4-(cyclopentyloxy)-3-(1-methylethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 189: 4-{3-[4-(cyclopentyloxy)-3-propoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10

No. 190: 4-{3-[4-(cyclopentyloxy)-3-hydroxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

15

No. 193: 4-{3-[4-(difluoromethoxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

20

No. 194: 4-{3-[4-(difluoromethoxy)-3-ethoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

25 No. 200: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(thiophen-3-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 201: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyridin-4-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

30

No. 203: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-1H-indol-6-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35

No. 206: 4-{3-[4-(cyclopropylmethoxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-

dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 207: 2-[4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)-2-methoxyphenoxy]-N-methylacetamide

No. 212: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(1H-pyrazol-1-yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10

No. 213: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyridin-2-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

15 No. 215: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(thiophen-2-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 216: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(quinolin-7-ylmethyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 218: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(6-methoxynaphthalen-2-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 223: 4-{3-[4-(1H-benzimidazol-1-yl)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

30

No. 224: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-methylpropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35 No. 226: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(tetrahydrofuran-3-yloxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 228: 4-[3-{4-[(1-benzylpyrrolidin-3-yl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5 No. 230: 4-[3-(1-benzothiophen-5-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 232: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-
10 4-(1-methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 233: 4-[3-(3,4-dimethoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-
15 yl]piperidine-1-carbaldehyde

No. 234: 4-[3-{4-[(1-acetylpyrrolidin-3-yl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde
20

No. 239: 4-[3-{4-[(4-fluorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

25 No. 240: 4-[3-{4-[(4-chlorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 242: 4-[3-{4-[(3-chlorobenzyl)oxy]-3-methoxybenzyl}-6-[2-
30 fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 243: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(3-(thiophen-3-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-
35 yl]piperidine-1-carbaldehyde

No. 245: 4-[3-(4-ethoxy-3-methoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-

carbaldehyde

No. 246: 4-[3-{4-[2-(2,3-dihydro-1H-indol-1-yl)-2-oxoethoxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 250: 4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

10

No. 251: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-oxo-2-(piperidin-1-yl)ethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

15 No. 254: 4-{3-[3-ethoxy-4-(thiophen-2-ylmethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 258: 4-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(hydroxymethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 263: (2R)-2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]propanoic acid

No. 264: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-3-(thiophen-2-yl)-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

30

No. 270: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35 No. 275: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyrimidin-5-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 276: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-3-phenyl-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

5 No. 278: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[[6-(1H-pyrazol-1-yl)pyridin-3-yl]methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 279: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-10 [(2-(thiophen-2-yl)pyrimidin-5-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 280: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,4-dioxo-3,4-15 dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 282: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(3-methyl-1,2,4-oxadiazol-5-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde
20

No. 283: [2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]acetic acid

25 No. 285: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(thieno[2,3-b]pyridin-2-yl)methyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 286: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-30 [(6-phenylpyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 287: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(6-(morpholin-4-yl)pyridin-3-yl)methyl]-2,4-dioxo-3,4-35 dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 289: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(6-(thiophen-2-yl)pyridin-3-yl)methyl]-3,4-dihydroquinazolin-

1 (2H)-yl}piperidine-1-carbaldehyde

No. 292: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-5-phenyl-1H-pyrazol-3-yl)methyl]-2,4-dioxo-3,4-
5 dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 294: 4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)biphenyl-2-carbonitrile

10

No. 295: (2R)-2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]-N-methylpropanamide

15

No. 297: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(thiophen-2-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

20 No. 298: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 299: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(piperidin-1-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

25

No. 300: 4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

30

No. 301: 2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]-N-ethylacetamide

35

No. 302: (2S)-2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-

1,4-dihydroquinazolin-3(2H)-yl)methyl)phenoxy]propanoic acid

No. 305: 4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-{[(3R)-2-oxo-1-phenylpyrrolidin-3-yl]oxy}benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 306: 4-{3-[4-(cyclobutylmethoxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10

No. 307: 4-{3-[4-(benzyloxy)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

15 No. 308: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-hydroxy-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 309: 4-{3-[4-(cyclopropylmethoxy)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 310: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-methylpropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 311: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(1-methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 312: 4-[3-(4-ethoxy-3-methoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 315: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{[6-(3-methoxyphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-

dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 316: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-
{[6-(2-fluorophenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-

5 dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 317: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-
{[6-(4-fluorophenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-
dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10

No. 318: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-
{[6-(4-methoxyphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-
dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

15 No. 319: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-
dioxo-3-[(6-(thiophen-2-yl)pyridin-3-yl)methyl]-3,4-
dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 320: 4-{3-[3-ethoxy-4-(thiophen-2-ylmethoxy)benzyl]-7-
20 fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-
dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 321: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-
(1-methyl-1H-pyrazol-3-yl)benzyl]-2,4-dioxo-3,4-
25 dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 322: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-
dioxo-3-(4-(pyrimidin-5-yl)benzyl)-3,4-dihydroquinazolin-
1(2H)-yl}piperidine-1-carbaldehyde

30

No. 323: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-
[(1-methyl-3-(thiophen-2-yl)-1H-pyrazol-5-yl)methyl]-2,4-
dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

35 No. 324: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-
methoxy-4-(2-oxo-2-(piperidin-1-yl)ethoxy)benzyl]-2,4-dioxo-
3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 325: 4-[3-{4-[2-(2,3-dihydro-1H-indol-1-yl)-2-oxoethoxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5

No. 326: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

10 No. 327: 4-[3-{4-[(3-chlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 328: 4-[3-{[6-(3,5-dichlorophenyl)pyridin-3-yl]methyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 329: 4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)biphenyl-2-carbonitrile

No. 330: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(1H-pyrazol-1-yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

25

No. 331: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{[6-(3-fluorophenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

30 No. 332: 3-[5-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)pyridin-2-yl]benzonitrile

No. 333: 4-[3-(3,4-diethoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

35 No. 334: 4-[3-{4-[(4-chlorobenzyl)oxy]-3-methoxybenzyl}-7-

fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 335: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 336: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-{[6-(1H-pyrazol-1-yl)pyridin-3-yl]methyl}-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 337: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-(morpholin-4-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 338: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-propoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 339: 4-{3-[4-(1H-benzimidazol-1-yl)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 340: 5-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)-2-methoxybenzonitrile

No. 341: 3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-7-carbonitrile

No. 342: 4-[3-(4-bromobenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 343: 4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-(2-methoxyethoxy)benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-

yl]piperidine-1-carbaldehyde

No. 344: 4-{3-[4-(benzyloxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 345: 4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-ethoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

10

No. 349: 4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-(2-fluoroethoxy)benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

15

No. 350: 4-[3-{4-[(2-chloro-4-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

20

No. 351: 4-[3-{4-[(2,4-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

25 No. 352: 4-[3-{4-[(2-chloro-6-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

30 No. 353: 4-[3-{4-[(2,6-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 354: 4-[3-{4-[(2-chlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

35

No. 355: 4-[7-fluoro-3-{4-[(2-fluorobenzyl)oxy]-3-

methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 357: 2-[(3,4-dichlorobenzyl)oxy]-5-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)benzonitrile

No. 358: 4-[3-{4-[(3,4-dichlorophenoxy)methyl]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 360: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(2-phenylethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 362: 4-[3-{4-[(4,5-dichloro-2-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 369: 4-[3-{4-[(4-chlorophenoxy)methyl]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 371: 4-[3-{3-chloro-4-[(4-chlorobenzyl)oxy]-5-ethoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 373: 4-[3-{3-chloro-4-[(2,4-dichlorobenzyl)oxy]-5-ethoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 375: 4-[7-fluoro-3-{4-[(4-fluorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 376: 4-[3-{4-[(3,5-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

5

No. 377: 4-[3-(4-{[4-chloro-3-(trifluoromethyl)benzyl]oxy}-3-methoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

10

No. 379: 4-[3-{4-[(3-chlorophenoxy)methyl]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

15 No. 380: 4-[3-{4-[(3,5-difluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 381: 4-{3-[4-(benzyloxy)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 382: 4-[3-{4-[(3-chloro-5-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 383: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-{[4-(trifluoromethyl)benzyl]oxy}benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde

No. 384: 4-[3-{4-[(2,5-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde

No. 385: 4-{[4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-

3 (2H)-yl)methyl)-2-methoxyphenoxy)methyl}benzonitrile

No. 386: 3-{[4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-
1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-
5 3 (2H)-yl)methyl)-2-methoxyphenoxy)methyl}benzonitrile

No. 387: 4-[3-{4-[(4-chloro-2-fluorobenzyl)oxy]-3-
methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-
2,4-dioxo-3,4-dihydroquinazolin-1 (2H)-yl]piperidine-1-
10 carbaldehyde

No. 388: 4-[3-{4-[1-(3,4-dichlorophenyl)ethoxy]-3-
methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-
2,4-dioxo-3,4-dihydroquinazolin-1 (2H)-yl]piperidine-1-
15 carbaldehyde

No. 389: 4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{4-
[(3-hydroxybenzyl)oxy]-3-methoxybenzyl}-2,4-dioxo-3,4-
dihydroquinazolin-1 (2H)-yl}piperidine-1-carbaldehyde
20

No. 390: 4-[7-fluoro-3-{4-[(3-fluorobenzyl)oxy]-3-
methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-
3,4-dihydroquinazolin-1 (2H)-yl]piperidine-1-carbaldehyde

25 No. 391: 4-[3-{4-[(3,4-difluorobenzyl)oxy]-3-methoxybenzyl}-7-
fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-
dihydroquinazolin-1 (2H)-yl]piperidine-1-carbaldehyde

No. 392: 4-{3-[4-(5,6-dichloro-1H-benzimidazol-1-yl)-3-
30 methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-
2,4-dioxo-3,4-dihydroquinazolin-1 (2H)-yl}piperidine-1-
carbaldehyde

No. 393: 4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-
35 (1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-
3 (2H)-yl)methyl}phenyl 3,4-dichlorobenzenesulphonate

No. 394: 4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-

(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)-2-methoxyphenyl 3,4-dichlorobenzenesulphonate

No. 403: 3,4-dichloro-N-[4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)-2-methoxyphenyl]benzamide

Another subject-matter of the present invention is processes
10 for the preparation of a compound of general formula (I).

The compounds of the invention can be prepared by the methods illustrated in the following Schemes 1 to 4.

15 In that which follows, the term "leaving group" is understood to mean a group which can be easily substituted, with departure of an electron pair, by cleavage of a heterolytic bond. This group can thus be easily replaced by another group, for example during a substitution reaction. Such leaving
20 groups are, for example, halogens or an activated hydroxyl group, such as a mesylate, tosylate, triflate, and the like. Examples of leaving groups and references for their preparation are given in "Advanced Organic Chemistry", J. March, 3rd Edition, Wiley Interscience, pp. 310-316.

25 The term "protective group PG" is understood to mean a group which makes it possible to prevent the reactivity of a function group or position during a chemical reaction which may affect it and which restores the molecule after cleavage
30 according to methods known to a person skilled in the art.

The term "temporary protective group for amines or alcohols" is understood to mean the protective groups such as those described in Protective Groups in Organic Synthesis, Greene
35 T.W. and Wuts P.G.M., published by Wiley Intersciences, 1999, and in Protecting Groups, Kocienski P.J., 1994, Georg Thieme Verlag.

Mention may be made, for example, of temporary protective groups for amines: benzyls or carbamates (such as *tert*-butoxycarbonyl, cleavable in an acid medium, or benzyloxycarbonyl, cleavable by hydrogenolysis), temporary
5 protective groups for carboxylic acids: hydrogenolysable alkyl esters (such as methyl or ethyl or *tert*-butyl esters which can be hydrolysed in a basic or acid medium) and benzyl esters, temporary protective groups for alcohols or phenols, such as tetrahydropyranyl, methyloxymethyl or methylethoxymethyl,
10 *tert*-butyl and benzyl ethers, or temporary protective groups for carbonyl derivatives, such as linear or cyclic acetals, such as, for example, 1,3-dioxan-2-yl or 1,3-dioxolan-2-yl; and reference may be made to the well known general methods described in Protective Groups, cited above.

15

A person skilled in the art will be in the position to choose the appropriate protective groups according to the circumstances. The compounds of formula (I) can comprise precursor groups for other functional groups which are
20 generated subsequently in one or more other stages.

In the general synthetic schemes which follow, the starting compounds and the reactants, when their method of preparation is not described, are commercially available or are described
25 in the literature or else can be prepared according to methods which are described therein or which are known to a person skilled in the art.

The pure enantiomers of the compounds of the invention can be
30 obtained from enantiomerically pure precursors or else by chiral phase chromatography or else, when the compounds comprise acid or amine functional groups, by selective crystallizations of diastereoisomeric salts obtained by reaction of the compounds (I) with chiral amines or chiral
35 acids respectively.

According to one embodiment of the invention, the compounds of general formula (I) can be obtained according to the following

Schemes 1 to 4. Out of concern for clarity, the group R_4 has been chosen as being a hydrogen, p and p' represent 1 and 2 respectively and the R_2 and R_3 groups have been set as indicated in the schemes. However, it is to be understood that
5 R_4 can be as defined in the general formula (I), and R_2 and R_3 can have the positions indicated in the general formula (I) and that p and p' can be as defined in the general formula (I).


10 The synthetic routes described below serve solely as illustration and are under no circumstances limiting. A person skilled in the art can apply without difficulty the teaching below to the compounds of formula (I) for which R , R_1 , R_2 , R_3 , R_4 , R_a , R_b , R_c , m , n , p , p' and q are as defined in the general
15 formula (I).

According to Scheme 1, the compound of formula (IV) is obtained by a reductive amination reaction by reacting a compound of formula (II), in which R' represents a (C_1 - C_6)
20 alkyl group and R_2 is as defined for the compound of formula (I) with a compound of formula (III) in an acid medium and in the presence of a reducing agent, such as sodium triacetoxyborohydride. The PG group of the compound of formula (III) is a protective group for the amine functional group
25 which may advantageously be tert-butoxycarbonyl (boc). The compound of formula (IV) thus formed is subsequently acylated, according to methods well known to a person skilled in the art, with an alkyl or aryl chloroformate to give the compound of formula (V) in which R'' represents a (C_1 - C_6) alkyl group or
30 an aryl group which is substituted. A hydrolysis reaction in a basic medium makes it possible to obtain the compounds of formula (VI) which, by a coupling reaction with a compound of formula (VII), in which R_3 is as defined for the compound of formula (I), results in the compounds of formula (VIII). An
35 intramolecular cyclization reaction in the basic medium makes it possible to obtain the quinazolinedione derivatives of formula (IX). The protective group PG for the amine functional group is subsequently cleaved, for example in acid medium when

PG is a boc, to give the compounds of formula (Ia) which give, by an acylation reaction, the compounds of formula (Ib).

The compounds of formula (Ia) are compounds of formula (I) and can act as intermediate for other compounds of formula (I), such as the compounds of formula (Ib).

Scheme 1



10


The compounds of formula (I) for which R_2 represents $-\text{ORa}$, Ra being as defined for the compound of formula (I), correspond to the formula (Id). They can be obtained from the compounds of formula (X) according to the following Scheme 2. The compounds of formula (Ic), obtained by a hydrogenolysis reaction on the compounds of formula (X), are subjected, for example, to an alkylation reaction with an alkylating agent of

type Ra-X, in which Ra is as defined for the compound of formula (I) and X represents a leaving group (such as a halogen atom, for example), in the presence of a base, such as caesium carbonate (Cs_2CO_3), or also to a Mitsunobu reaction
 5 (Synthesis, 1981, 1) with an alcohol of type Ra-OH, Ra being as defined for the compound of formula (I), to give compounds of formula (Id).

The compounds of formula (X) and the compounds of formula (Ic)
 10 are compounds of formula (I) and can act as intermediate for other compounds of formula (I), such as the compounds of formula (Id).

Scheme 2

15




Alternatively, the compounds of formula (Id) can be obtained
 by following the procedure described in Scheme 3.

20

The compounds of formula (XII) are obtained by a nucleophilic aromatic substitution reaction involving a compound of formula (XI), in which R' is as defined above, and an alcohol of type Ra-OH, in which Ra is as defined for the compound of formula

(I), in the presence of a base. The reduction of the nitro group of the compounds of formula (XII) results in the corresponding anilino derivatives (XIII). A reductive amination reaction with a compound of formula (III), in which
5 PG is a protective group for amine functional groups, such as, for example, boc, results in the compounds of formula (XIV). The compounds of formula (XV) are obtained by reaction of a compound of formula (XIV) with potassium isocyanate (KNCO) in an acid medium. An intramolecular cyclization reaction in a
10 basic medium makes it possible to obtain the compounds of formula (XVI). The protective group PG is cleaved by methods well known to a person skilled in the art to give the compounds of formula (XVII). An acylation reaction results in the compounds of formula (XVIII). Finally, the compounds of
15 formula (Id) are obtained either by an alkylation reaction with a derivative of type (XIX), in which X represents a leaving group, such as a halogen atom, in the presence of a base, such as, for example, caesium carbonate, or also by a Mitsunobu reaction with a benzyl alcohol of type (XX). In the
20 compounds (XIX) and (XX), R_3 is as defined above.

Scheme 3




The compounds of formula (Ie) and (If), in which R_2 more particularly represents a group of type- $\text{CH}_2\text{-NHC(O)Rb}$, Rb being defined as in the compound of formula (I), can be prepared according to the following Scheme 4.

It is understood that, in Scheme 3, the R_2 group illustrated is of $-\text{O-Ra}$ type and is in the 6 position of the quinoazolidinedione structure (see, for example, compound (XVIII)) but that it is also possible to have a second R_2

group, as defined in the general formula (I), in the 7 position of the same quinazolinedione group.

The reduction of the nitro group of the compounds of type (XXI), in which R' and PG' are as defined above, the PG' group advantageously being boc, results in the corresponding anilino derivatives (XXII) which, by a reductive amination reaction in which they react in an acid medium and advantageously in the presence of a reducing agent, such as sodium triacetoxyborohydride, with a compound of formula (III), in which PG represents a benzyloxycarbonyl protective group for amines, give compounds of formula (XXIII). An acylation reaction with an alkyl or aryl chloroformate, in which R'' represents a (C₁-C₆) alkyl group or an aryl group which is substituted, results in the compounds of formula (XXIV). The quinazolinedione analogues of formula (XXV) can be obtained by a hydrolysis reaction in a basic medium and then by a coupling reaction with a compound of formula (VII), in which R₃ is as defined for the compound of formula (I), followed by an intramolecular cyclization reaction in a basic medium. The PG' group (preferably a boc) is subsequently cleaved in an acid medium to result in the compounds of formula (XXVI) which, by acylation, give compounds of formula (XXVII), in which R_b is as defined for the compound of formula (I). The PG protective group of (XXVII) is cleaved by a hydrogenolysis reaction to give the compounds of formula (Ie). Finally, the compounds of formula (If) are obtained by an acylation reaction on the compounds of formula (Ie).

30 **Scheme 4**



It is obvious that a person skilled in the art will be in a position to choose, in the light of his knowledge and the literature, other appropriate protective groups which make possible the introduction of all the groups described in the general formula (I).

When the compound of formula (I) comprises a bridged ring, it can be obtained without distinction by one of the synthetic routes described above.

The following procedures and examples describe the preparation of some compounds in accordance with the invention. These procedures and examples are not limiting and serve only to illustrate the present invention.

5

In the procedures and examples below:

- the mass spectra are produced on a quadrupole spectrometer of Platform LCZ type (Waters) or of ZQ 4000 type (Waters) in
10 positive electrospray ionization mode;

- the NMR (nuclear magnetic resonance) spectra are produced on a Fourier transform spectrometer (Bruker) at a temperature of 300°C (exchangeable protons not recorded);

15

- s = singlet,

- d = doublet,

20 - m = multiplet,

- br = broad signal

- t = triplet,

25

- q = quartet

- DMSO-d₆ = deuterated dimethyl sulphoxide,

30 - CDCl₃ = deuterated chloroform.

The mixtures of solvents are quantified in volumetric ratios.

The NMR spectra and mass spectra confirm the structures of the
35 compounds obtained according to the examples below.

In the examples which follow, the following abbreviations are used:

ACN: acetonitrile

AcOEt: ethyl acetate

5

AcOH: acetic acid

DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene

10 DCM: dichloromethane

DCE: 1,2-dichloroethane

DIAD: diisopropyl azodicarboxylate

15

DIEA: diisopropylamine

DMF: N,N-dimethylformamide

20 EtOH: ethanol

HBTU: O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
hexafluorophosphate

25 IBCF: isobutyl chloroformate

MeOH: methanol

NaBH(OAc)₃: sodium triacetoxyborohydride

30

AT: ambient temperature

min: minute

35 THF: tetrahydrofuran

NEt₃: triethylamine


TFA: trifluoroacetic acid

EXAMPLES

5 The following examples describe the preparation of some compounds in accordance with the invention. These examples are not limiting and serve only to illustrate the present invention. The numbers of the compounds in the examples refer to those given in the table below, in which the chemical
10 structures and the physical properties of a few compounds according to the invention are illustrated.

EXAMPLE 1: Compound No. °6

15 **Preparation of {[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetonitrile**



20


Stage 1.1:

1,1-Dimethylethyl

4-{[4-(benzyloxy)-2-(methoxy-

carbonyl)phenyl]amino}piperidine-1-carboxylate

25



A mixture of 2 g of methyl 2-amino-5-(benzyloxy)benzoate, 3.1 g of 1,1-dimethylethyl 4-oxopiperidine-1-carboxylate and 3.29 g of $\text{NaBH}(\text{OAc})_3$ in 10 ml of AcOH is irradiated under a microwave field (Biotage Initiator Sixty) at 110°C for 20 min. The same reaction is repeated with two other lots of 2 g of methyl 2-amino-5-(benzyloxy)benzoate. The three reaction media are combined. The combined product is taken up in AcOEt. The organic phase is washed with water, with a saturated NH_4Cl solution and with a saturated NaHCO_3 solution, dried over Na_2SO_4 and filtered, and the solvent is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/heptane mixture, (5/95, v/v) as far as (30/70, v/v), to give 10.2 g of the expected product.


Stage 1.2:

1,1-Dimethylethyl

4-{ [4-(benzyloxy)-2-

(methoxycarbonyl)phenyl] [(2-methyl-

propoxy) carbonyl] amino} piperidine-1-carboxylate




A mixture of 2 g of 1,1-dimethylethyl 4-{[4-(benzyloxy)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate obtained in stage 1.1, 0.87 ml of DIEA, 1.78 ml of IBCF and 1 g of NaOH in 10 ml of DCE is irradiated under a microwave field at 80°C for 30 min. The same reaction is repeated with 4 other lots of 2 g of 1,1-dimethylethyl 4-{[4-(benzyloxy)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate. The

reaction media are combined. The combined product is taken up in AcOEt and filtered, and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/heptane mixture, 5 (10/90, v/v) as far as (50/50, v/v), to give 9.3 g of the expected product.

Stage 1.3:

10 **Sodium salt of 5-(benzyloxy)-2-({1-[(1,1-dimethylethoxy)carbonyl]-piperidin-4-yl}[(2-methylpropoxy)carbonyl]amino)benzoic acid**




15 A mixture of 9.3 g of 1,1-dimethylethyl 4-{{4-(benzyloxy)-2-(methoxycarbonyl)phenyl}[(2-methylpropoxy)carbonyl]amino}piperidine-1-carboxylate obtained in stage 1.2 and 34.4 ml of 2N NaOH in 57 ml of MeOH is heated
20 at 100°C for 3h 00. The solution is evaporated under reduced pressure and DCM is added. Drying is carried out over Na₂SO₄, filtration is carried out and the solvent is evaporated under reduced pressure to give 8.7 g of the expected product.

25 Stage 1.4:

1,1-Dimethylethyl 4-{{4-(benzyloxy)-2-[(3,4-dimethoxybenzyl)carbamoyl]phenyl}(isobutoxycarbonyl)amino}piperidine-1-carboxylate


30



A mixture of 6 g of sodium salt of 5-(benzyloxy)-2-({1-[(1,1-dimethylethoxy)carbonyl]piperidin-4-yl}[(2-methylpropoxy)carbonyl]amino)benzoic acid obtained in stage 1.3 and 4.42 g of DIEA in 250 ml of DMF is stirred at AT for 15 min. 6.48 g of HBTU are added and the mixture is left stirring for 30 min. 2.48 g of veratrylamine are added and the reaction mixture is stirred for 48h 00. It is evaporated under reduced pressure, the residue is taken up in AcOEt, washed with a saturated NH_4Cl solution and with a saturated NaHCO_3 solution, dried over Na_2SO_4 and filtered, and the solvent is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/heptane mixture, (20/80, v/v) as far as (60/40, v/v), to give 7.5 g of expected product.

Stage 1.5:


20 **1,1-Dimethylethyl 4-[6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate**



A mixture of 2.5 g of 1,1-dimethylethyl 4-[[4-(benzyloxy)-2-[(3,4-dimethoxybenzyl)carbamoyl]phenyl](isobutoxycarbonyl)amino]piperidine-1-carboxylate obtained in stage 1.4 and 7.4 g of NaOH in 18.5 ml of DCE is irradiated under a microwave field at 110°C for 30 min. The same reaction is repeated with 2 other lots of 2.5 g of 1,1-dimethylethyl 4-[[4-(benzyloxy)-2-[(3,4-dimethoxybenzyl)carbamoyl]phenyl](isobutoxycarbonyl)amino]piperidine-1-carboxylate. The reaction media are combined. The combined product is taken up in DCM, washed with water, dried over Na₂SO₄ and filtered, and the solvent is evaporated under reduced pressure to give 6.6 g of expected product.

15 **Stage 1.6:**


6-(Benzyloxy)-3-(3,4-dimethoxybenzyl)-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)-dione



A mixture of 3.5 g of 1,1-dimethylethyl 4-[6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate obtained in stage 1.5 and 25 ml of TFA in 50 ml of DCM is stirred at AT for 2h 00. The mixture is neutralized with K₂CO₃. It is filtered and the filtrate is evaporated under reduced pressure. The residue is taken up in DCM and washed with a saturated NaHCO₃ solution and then with an 8% NaOH solution. The solution is dried over Na₂SO₄ and filtered, and the solvent is evaporated under reduced pressure to give 2.67 g of the expected product.

Stage 1.7:

4-[6-(Benzyloxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde




5

A mixture of 0.6 g of 6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)-dione obtained in stage 1.6 and 0.113 g of ammonium formate in 5 ml of ACN is irradiated under a microwave field at 140°C for 1h 00. The mixture is filtered and the filtrate is evaporated under reduced pressure to give 0.62 g of the expected product.

Stage 1.8: Compound No. 5:

15

4-[3-(3,4-Dimethoxybenzyl)-6-hydroxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde



20

A mixture of 0.618 g of 4-[6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde obtained in stage 1.7, 0.44 g of ammonium formate and 0.124 g of Pd/C (10%) in 10 ml of EtOH purged beforehand with nitrogen is irradiated under a microwave field at 80°C for 2h 00. The mixture is filtered and


25

the filtrate is evaporated under reduced pressure to give 0.513 g of the expected product.

Stage 1.9: Compound No. 6

5

{[3-(3,4-Dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetonitrile




10

0.17 g of 4-[3-(3,4-dimethoxybenzyl)-6-hydroxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde obtained in stage 1.8 and 0.25 g of Cs_2CO_3 in 3 ml of DMF are stirred for 15 min at AT. 0.056 g of bromoacetonitrile is added and the reaction mixture is subsequently irradiated under a microwave field at 100°C for 15 min. It is filtered and evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an MeOH/DCM mixture, (1/99, v/v) as far as (4/96, v/v), to give 0.112 g of the expected product.


EXAMPLE 2: Compound No. 3

Preparation of {[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetonitrile

25

**Stage 2.1:**

- 5 **1-(1-Acetylpiperidin-4-yl)-6-(benzyloxy)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione**




- 10 0.14 g of acetyl chloride is added to a mixture of 0.6 g of 6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)-dione obtained according to stage 1.6 and 0.24 g of NEt₃ in 10 ml of DCM cooled to 0°C. The mixture is stirred at AT overnight. It is washed twice with a
- 15 saturated NH₄Cl solution and filtered, and then the filtrate is evaporated under reduced pressure to give 0.64 g of the expected product.

Stage 2.2: Compound No. 2

20

1-(1-Acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-hydroxyquinazoline-2,4(1H,3H)-dione




A mixture of 0.64 g of 1-(1-acetylpiperidin-4-yl)-6-(benzyloxy)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-
 5 dione obtained in stage 2.1, 0.44 g of ammonium formate and 0.125 g of Pd/C (10%) in 10 ml of EtOH purged beforehand with nitrogen is irradiated under a microwave field at 80°C for 2h 00. The mixture is filtered and the filtrate is evaporated under reduced pressure to give 0.48 g of the expected product.

10

Stage 2.3: Compound No. 3

{[1-(1-Acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetonitrile

15




0.12 g of 1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-hydroxyquinazoline-2,4(1H,3H)-dione obtained in stage 2.2
 20 and 0.172 g of Cs₂CO₃ in 3 ml of DMF are stirred for 15 min at AT. 0.038 g of bromoacetonitrile is added and the reaction mixture is subsequently irradiated under a microwave field at 100°C for 15 min. It is filtered and the filtrate is evaporated under reduced pressure. The residue is
 25 chromatographed on silica gel, elution being carried out with an MeOH/DCM mixture, (1/99, v/v) as far as (4/96, v/v), to

give 0.094 g of the expected product.


EXAMPLE 3: Compound No. 34

5 Synthesis of 4-[3-(4-chlorobenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde



10 Stage 3.1:

Methyl 5-(2,2-difluoroethoxy)-2-nitrobenzoate




15

8.53 g of 2,2-difluoroethanol are added to a solution of 17.31 g of methyl 5-fluoro-2-nitrobenzoate, 9.64 g of NEt₃ and 32.71 g of 2,8,9-triisobutyl-2,5,8,9-tetraaza-1-phosphabicyclo[3.3.3]undecane in 250 ml of anhydrous THF. The mixture is stirred at AT for 30 min. The solvent is evaporated under reduced pressure. Water is added and extraction is carried out with AcOEt. The extract is washed with a 1N aqueous HCl solution, with water and then with a saturated NaCl solution. It is dried over MgSO₄ and filtered, and the solvent is evaporated under reduced pressure to give 21 g of the expected product.

Stage 3.2:

30 Methyl 2-amino-5-(2,2-difluoroethoxy)benzoate



21 g of methyl 5-(2,2-difluoroethoxy)-2-nitrobenzoate obtained
 5 in stage 3.1 and 1 g of Pd/C (10%) in a mixture of 300 ml of
 AcOEt, 50 ml of EtOH and 5 ml of AcOH are stirred at AT under
 a hydrogen atmosphere for 24h 00.

The mixture is filtered and evaporated under reduced pressure
 to give 18.6 g of the expected product.

10


Stage 3.3:

1,1-Dimethylethyl

4-{[4-(2,2-difluoroethoxy)-2-

(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate


15



A mixture of 4 g of methyl 2-amino-5-(2,2-
 difluoroethoxy)benzoate and 6.88 g of 1,1-dimethylethyl
 20 4-oxopiperidine-1-carboxylate obtained in stage 3.2 in 15 ml
 of AcOH is heated at 90°C for 10 min. It is allowed to cool to
 AT and 7.3 g of NaBH(OAc)₃ are added. The mixture is left
 stirring at AT for 12h 00. It is extracted with AcOEt and the
 extract is washed with a saturated K₂CO₃ solution and then with
 25 water. It is dried over MgSO₄, filtered and evaporated under
 reduced pressure to give 6.63 g of the expected product.

Stage 3.4:

1,1-Dimethylethyl 4-{carbamoyl[4-(2,2-difluoroethoxy)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate



5


1.95 g of potassium isocyanate in solution of 4 ml of water are added to a solution of 6.63 g of 1,1-dimethylethyl 4-{[4-(2,2-difluoroethoxy)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate obtained in stage 3.3 in 40 ml of AcOH. The mixture is stirred at AT for 12h 00. It is extracted with AcOEt and the extract is washed with a saturated K₂CO₃ solution and then with water. It is dried over MgSO₄, filtered and evaporated under reduced pressure to give 6.95 g of the expected product.

15

Stage 3.5:

1,1-Dimethylethyl 4-[6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate

20



2.5 g of 1,1-dimethylethyl 4-{carbamoyl[4-(2,2-difluoroethoxy)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate obtained in stage 3.4 in solution in a mixture of


25

10 ml of dioxane and 5 ml of a 1N aqueous NaOH solution are irradiated under a microwave field at 130°C for 30 min. Extraction is carried out with AcOEt and the extract is neutralized with a 1N aqueous HCl solution, washed with water, 5 dried over MgSO₄, filtered and evaporated under reduced pressure. The residue obtained is triturated in an AcOEt/pentane mixture to give the expected product. The same reaction is reproduced with 2 other lots of 2.5 g of 1,1-dimethylethyl 4-{carbamoyl[4-(2,2-difluoroethoxy)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate 10 obtained in stage 3.4 to give, in total, 5.63 g of expected product.

Stage 3.6:

15

6-(2,2-Difluoroethoxy)-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)-dione




20

A solution of 5.63 g of 1,1-dimethylethyl 4-[6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate obtained in stage 3.5 in 70 ml of formic acid is stirred at AT for 2h 00. The solvent is 25 evaporated under reduced pressure to give 6.13 g of the expected product in the form of the formic acid salt.

Stage 3.7:


30 **4-[6-(2,2-Difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde**



A mixture of 6.13 g of 6-(2,2-difluoroethoxy)-1-(piperidin-4-yl)quinazoline-2,4(1*H*,3*H*)-dione obtained in stage 3.6 and 3.12 g of ammonium formate in 28 ml of ACN and 28 ml of dioxane is irradiated under a microwave field at 140°C for 1h 00. The reaction mixture is run into water. The mixture is filtered and the precipitate is washed with water and then with ether to give 4.47 g of the expected product.

Stage 3.8: Compound No. 34

4-[3-(4-Chlorobenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2*H*)-yl]piperidine-1-carbaldehyde




A mixture of 0.15 g of 4-[6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2*H*)-yl]piperidine-1-carbaldehyde obtained in stage 3.7, 0.096 g of 1-(bromomethyl)-4-chlorobenzene and 0.3 g of Cs₂CO₃ in 3 ml of DMF is stirred at RT for 1h 00. AcOEt is added and washing is carried out with water and then with a saturated NaCl solution. The organic phase is dried over MgSO₄, filtered and evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with AcOEt, to give 0.116 g of

the expected product.

EXAMPLE 4: Compound No. 49

5 Synthesis of 4-{3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde




10

0.172 g of DIAD is added to a solution of 0.15 g of 4-[6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde obtained in stage 3.7, 0.142 g of [4-(cyclopentyloxy)-3-methoxyphenyl]methanol and 0.223 g of PPh₃ in 3 ml of anhydrous THF. The mixture is stirred at AT for 12h 00 and then at 60°C for 1h 00. It is evaporated under reduced pressure and the residue is purified on silica gel, elution being carried out with AcOEt, to give 0.083 g of the expected product.


20

EXAMPLE 5: Compound No. 20

25 Synthesis of N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide

**Stage 5.1:**


- 5 **Methyl 2-amino-5-({[(1,1-dimethylethoxy) carbonyl] amino}methyl) benzoate**



- 10 A mixture of 0.273 g of methyl 5-{{[(tert-butoxycarbonyl) amino]methyl}}-2-nitrobenzoate, 0.166 g of ammonium formate and 0.094 g of Pd/C (10%) in 10 ml of EtOH purged beforehand with nitrogen is irradiated under a microwave field at 120°C for 5 min. The mixture is filtered
15 and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/heptane mixture, (5/95, v/v) as far as (30/70, v/v), to give 0.2 g of the expected product.

- 20 **Stage 5.2:**


Benzyl 4-{[4-({[(1,1-dimethylethoxy) carbonyl] amino}methyl)-2-(methoxycarbonyl)phenyl] amino}piperidine-1-carboxylate



A solution of 1.66 g of benzyl 4-oxopiperidine-1-carboxylate and 1 g of methyl 2-amino-5-({[(1,1-dimethylethoxy)carbonyl]amino}methyl)benzoate obtained in stage 5.1 in 20 ml of DCM is added dropwise at AT to a suspension of 2.04 g of NaBH(OAc)₃ in a mixture of 20 ml of DCM and 0.41 ml of AcOH. The mixture is stirred at AT for 15h 00 and then a further 2.04 g of NaBH(OAc)₃ are added. After stirring for 6h 00, 1.66 g of benzyl 4-oxopiperidine-1-carboxylate are added and the mixture is stirred at AT for 48h 00. A saturated NaHCO₃ solution is added and extraction is carried out with DCM. The organic phase is washed with a saturated NaHCO₃ solution and twice with a saturated NH₄Cl solution. It is dried over Na₂SO₄ and filtered, and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/heptane mixture, (5/95, v/v) as far as (40/60, v/v), to give 1.6 g of the expected product.

Stage 5.3:


Benzyl 4-{[4-({[(1,1-dimethylethoxy)carbonyl]amino}methyl)-2-(methoxycarbonyl)phenyl] (ethoxycarbonyl)amino}piperidine-1-carboxylate



2.08 g of DIEA and then 1.745 g of ethyl chloroformate are added to a solution of 1.6 g of benzyl 4-{{[4-({[(1,1-dimethylethoxy)carbonyl]amino)methyl)-2-(methoxycarbonyl)phenyl]amino}piperidine-1-carboxylate obtained in stage 5.2 in 11 ml of DCM. The mixture is stirred at AT for 4 days. It is evaporated under reduced pressure. The residue is taken up in 10 ml of pyridine (10 ml) and 0.7 g of ethyl chloroformate is added. The mixture is stirred at AT for 4h 00. It is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/heptane mixture, (10/90, v/v) as far as (30/70, v/v), to give 0.875 g of the expected product.

Stage 5.4:

Benzyl 4-[3-(3,4-dimethoxybenzyl)-6-({[(1,1-dimethylethoxy)carbonyl]amino)methyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate




A mixture of 0.165 g of benzyl 4-{{[4-({[(1,1-dimethylethoxy)

carbonyl]amino}methyl)-2-(methoxycarbonyl)phenyl](ethoxycarbonyl)amino}piperidine-1-carboxylate obtained in stage 5.3 and 0.028 g of LiOH in 5 ml of THF/H₂O (70/30) is stirred at AT for 15h 00. The mixture is subsequently irradiated under a microwave field at 100°C for 1h 00. It is filtered and the filtrate is evaporated under reduced pressure. The residue is taken up in 5 ml of DMF. 0.108 g of DIEA is added and the mixture is stirred at AT for 10 min. 0.159 g of HBTU is added and the mixture is stirred at AT for 30 min. 0.061 g of veratrylamine is subsequently added and the mixture is stirred at AT for 1h 00. 0.5 ml of DBU is added and the mixture is stirred at AT for 48h 00. It is evaporated under reduced pressure and the residue is taken up in AcOEt. The solution is washed 3 times with a saturated NH₄Cl solution and twice with water. It is dried over Na₂SO₄ and filtered, and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an AcOEt/DCM mixture, (10/90, v/v) as far as (20/80, v/v), to give 0.104 g of the expected product.

Stage 5.5:

Benzyl 4-[6-(aminomethyl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate




A solution of 0.102 g of benzyl 4-[3-(3,4-dimethoxybenzyl)-6-({[(1,1-dimethylethoxy)carbonyl]amino}methyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate obtained

in stage 5.4 and 0.5 ml of TFA in 9.5 ml of DCM is stirred at AT for 2h 00. A saturated NaHCO₃ solution is added. The organic phase is dried over Na₂SO₄ and filtered, and the filtrate is evaporated under reduced pressure to give 0.09 g of the expected product.

Stage 5.6:


Benzyl 4-{6-[(acetylamino)methyl]-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carboxylate



0.049 g of acetic anhydride is added to a solution of 0.09 g of benzyl 4-[6-(aminomethyl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate obtained in stage 5.5 and 0.09 ml of NEt₃ in 3 ml of DCM and the mixture is stirred at AT for 1h 00. DCM is added and the solution is washed with a saturated NH₄Cl solution, then with a 1N HCl solution, then with a 2N NaOH solution and then with water. The organic phase is dried over Na₂SO₄ and filtered, and the filtrate is evaporated under reduced pressure to give 0.104 g of the expected product.

Stage 5.7:


N-{[3-(3,4-Dimethoxybenzyl)-2,4-dioxo-1-(piperidin-4-yl)-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide



A mixture of 0.1 g of benzyl 4-{6-[(acetylamino)methyl]-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carboxylate obtained in stage 5.6, 0.016 g of ammonium formate and 0.018 g of Pd/C (10%) in 2 ml of EtOH purged beforehand with nitrogen is irradiated under a microwave field at 80°C for 30 min. The mixture is filtered and the filtrate is evaporated under reduced pressure to give 0.077 g of the expected product.




Stage 5.8: Compound No. 20




***N*-{[3-(3,4-Dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide**









A mixture of 0.070 g of *N*-{[3-(3,4-dimethoxybenzyl)-2,4-dioxo-1-(piperidin-4-yl)-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide obtained in stage 5.7 and 0.028 g of ammonium formate in 2 ml of ACN is irradiated under a microwave field at 140°C for 1h 00. The mixture is filtered and the filtrate is evaporated under reduced pressure. The residue is chromatographed on silica gel, elution being carried out with an MeOH/DCM mixture, (0.5/99.5, v/v) as far as (7/93, v/v), to give 0.035 g of the expected product.




The chemical structures and the physical properties of compounds corresponding to the general formula (I) according to the invention, and also of some of their intermediates (in 5 particular compounds 32, 55, 120 and 257), are illustrated in the following table.




COMPOUND No.	STRUCTURE	NOMENCLATURE	NMR or MASS
1		2-([3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy)propanenitrile	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.71 (d, 3H) 1.79 (brdd, 2H) 2.41 (dq, 1H) 2.53 (dq, 1H) 2.80 (td, 1H) 3.25 (t, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.80 (d, 1H) 4.31 (d, 1H) 4.77 (brs, 1H) 5.03 (s, 2H) 5.60 (q, 1H) 6.83 (dd, 1H) 6.85 (d, 1H) 6.98 (s, 1H) 7.51 (dd, 1H) 7.74 (d, 1H) 7.84 (d, 1H) 8.03 (s, 1H)
2		1-(1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-hydroxyquinazolin-2,4(1H,3H)-dione	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.69 (d, 1H) 1.74 (d, 1H) 2.05 (s, 3H) 2.42 (dq, 1H) 2.57 (dq, 1H) 2.70 (t, 1H) 3.24 (t, 1H) 3.70 (br s, 3H) 3.71 (s, 3H) 3.92 (d, 1H) 4.51 (d, 1H) 4.69 (brs, 1H) 5.02 (s, 2H) 6.81 (dd, 1H) 6.86 (d, 1H) 6.97 (d, 1H) 7.21 (dd, 1H) 7.44 (d, 1H) 7.65 (d, 1H) 9.88 (brs, 1H)
3		{[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetoneitrile	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.71 (d, 1H) 1.76 (d, 1H) 2.04 (s, 3H) 2.41 (dq, 1H) 2.56 (dq, 1H) 2.70 (t, 1H) 3.24 (t, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.92 (d, 1H) 4.51 (d, 1H) 4.73 (brs, 1H) 5.04 (s, 2H) 5.29 (s, 2H) 6.82 (dd, 1H) 6.85 (d, 1H) 6.98 (d, 1H) 7.50 (dd, 1H) 7.70 (d, 1H) 7.83 (d, 1H)




4		2-([1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propanenitrile	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.71 (d, 3H) 1.72 (d, 2H) 1.77 (d, 1H) 2.04 (s, 3H) 2.41 (qd, 1H) 2.56 (dq, 1H) 2.70 (d, 1H) 3.24 (t, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.92 (d, 1H) 4.51 (d, 1H) 4.73 (brs, 1H) 5.03 (s, 2H) 5.60 (q, 1H) 6.83 (d, 1H) 6.85 (m, 1H) 6.98 (d, 1H) 7.51 (dd, 1H) 7.74 (d, 1H) 7.83 (d, 1H)
5		4-[3-(3,4-dimethoxybenzyl)-6-hydroxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.75 (d, 1H) 1.78 (d, 1H) 2.41 (qd, 1H) 2.53 (qd, 1H) 2.79 (td, 1H) 3.25 (td, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.80 (d, 1H) 4.31 (d, 1H) 4.72 (brs, 1H) 5.02 (s, 2H) 6.82 (dd, 1H) 6.86 (d, 1H) 6.97 (d, 1H) 7.21 (dd, 1H) 7.44 (d, 1H) 7.66 (d, 1H) 8.03 (s, 1H) 9.85 (brs, 1H)
6		{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetoneitrile	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.78 (d, 1H) 1.81 (d, 1H) 2.42 (qd, 1H) 2.54 (dt, 1H) 2.81 (td, 4H) 3.26 (dt, 1H) 3.71 (s, 3H) 3.72 (s, 3H) 3.81 (d, 1H) 4.32 (d, 1H) 4.77 (brs, 1H) 5.05 (s, 2H) 5.30 (s, 2H) 6.84 (dd, 1H) 6.87 (d, 1H) 6.99 (d, 1H) 7.51 (dd, 1H) 7.71 (d, 1H) 7.85 (d, 1H) 8.04 (s, 1H)




7		2-([3-(3,4-dimethoxybenzyl)-2,4-dioxo-1-(piperidin-4-yl)-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propanenitrile	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 1.69 (d, 3H) 1.75 (d, 2H) 2.65 (td, 2H) 2.87 (t, 2H) 3.21 (d, 2H) 3.69 (s, 3H) 3.70 (s, 3H) 4.68 (brs, 1H) 5.04 (s, 2H) 5.59 (q, 1H) 6.84 (m, 2H) 6.98 (d, 1H) 7.50 (dd, 1H) 7.73 (d, 1H) 7.80 (d, 1H)
8		{3-(3,4-dimethoxybenzyl)-2,4-dioxo-1-(piperidin-4-yl)-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}acetoneitrile	¹ H NMR (600 MHz, DMSO- <i>d</i> ₆) δ ppm 1.77 (d, 2H) 2.67 (dt, 2H) 2.89 (t, 2H) 3.23 (d, 2H) 3.69 (s, 3H) 3.70 (s, 3H) 4.69 (brs, 1H) 5.05 (s, 2H) 5.28 (s, 2H) 6.84 (m, 2H) 6.99 (s, 1H) 7.49 (dd, 1H) 7.70 (d, 1H) 7.80 (d, 1H)
9		2-([3-(3,4-dimethoxybenzyl)-1-(1-methylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propanenitrile	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 1.59 (d, 2H) 1.66 (d, 3H) 2.04 (t, 2H) 2.16 (s, 3H) 2.62 (dt, 2H) 2.82 (d, 2H) 3.66 (s, 3H) 3.67 (s, 3H) 4.43 (brs, 1H) 5.01 (s, 2H) 5.54 (q, 1H) 6.79 (m, 1H) 6.82 (d, 1H) 6.95 (d, 1H) 7.45 (dd, 1H) 7.68 (d, 1H) 7.70 (d, 1H)




10		3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-piperidin-4-ylquinazoline-2,4(1H,3H)-dione	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.85 (d, 2H) 2.78 (td, 2H) 3.04 (t, 2H) 3.33 (d, 2H) 3.71 (s, 6H) 4.61-4.82 (m, 4H) 5.04 (brt, 1H) 5.05 (s, 2H) 6.86 (m, 2H) 6.99 (s, 1H) 7.49 (dd, 1H) 7.68 (d, 1H) 7.76 (d, 1H) 8.05 (brs, 1H)
11		4-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.76 (d, 1H) 1.80 (d, 1H) 2.41 (qd, 1H) 2.53 (qd, 1H) 2.80 (td, 1H) 3.25 (dt, 1H) 3.70 (s, 6H) 3.80 (d, 1H) 4.31 (d, 1H) 4.61-4.83 (m, 5H) 5.02 (m, 1H) 5.03 (s, 2H) 6.83 (dd, 1H) 6.85 (d, 1H) 6.97 (d, 1H) 7.47 (dd, 1H) 7.67 (d, 1H) 7.77 (d, 1H) 8.03 (s, 1H)
12		1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.70 (d, 1H) 1.76 (d, 1H) 2.04 (s, 3H) 2.41 (dq, 1H) 2.56 (dq, 1H) 2.70 (t, 1H) 3.25 (t, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.92 (d, 1H) 4.51 (d, 1H) 4.61-4.83 (m, 5H) 5.02 (m, 1H) 5.03 (s, 2H) 6.82 (dd, 1H) 6.85 (m, 1H) 6.97 (d, 1H) 7.47 (dd, 1H) 7.67 (d, 1H) 7.77 (d, 1H)

13		4-[3-(3,4-dimethoxybenzyl)-2,4-dioxo-6-(2,2,2-trifluoroethoxy)-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (600 MHz, DMSO-d ₆) δ ppm 1.75 (d, 1H) 1.79 (d, 1H) 2.40 (dq, 1H) 2.52 (dq, 1H) 2.79 (td, 1H) 3.25 (td, 1H) 3.69 (s, 6H) 3.79 (d, 1H) 4.30 (d, 1H) 4.75 (brs, 1H) 4.88 (q, 2H) 5.02 (s, 2H) 6.82 (d, 1H) 6.85 (d, 1H) 6.97 (s, 1H) 7.49 (dd, 1H) 7.66 (d, 1H) 7.79 (d, 1H) 8.02 (s, 1H)
14		1-(1-(1-acetylpiperidin-4-yl)-6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.70 (d, 1H) 1.74 (d, 1H) 2.04 (s, 3H) 2.41 (qd, 1H) 2.56 (qd, 1H) 2.70 (t, 1H) 3.24 (t, 1H) 3.70 (s, 3H) 3.70 (s, 3H) 3.91 (d, 1H) 4.43 (td, 2H) 4.50 (d, 1H) 4.73 (brs, 1H) 5.03 (s, 2H) 6.41 (tt, 1H) 6.82 (dd, 1H) 6.85 (d, 1H) 6.97 (d, 1H) 7.46 (dd, 1H) 7.60 (d, 1H) 7.79 (d, 1H)
15		6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)-dione	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.83 (d, 2H) 2.75 (td, 2H) 3.00 (t, 2H) 3.30 (d, 2H) 3.71 (s, 6H) 4.44 (td, 2H) 4.75 (b. s, 1H) 5.06 (s, 2H) 6.42 (tt, 1H) 6.83-6.89 (m, 2H) 7.00 (s, 1H) 7.48 (dd, 1H) 7.62 (d, 1H) 7.78 (d, 1H)




16		4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 1.76 (d, 1H) 1.80 (d, 1H) 2.41 (qd, 1H) 2.53 (qd, 1H) 2.81 (td, 1H) 3.26 (td, 1H) 3.71 (s, 6H) 3.80 (d, 1H) 4.31 (d, 1H) 4.44 (td, 2H) 4.77 (brs, 1H) 5.04 (s, 2H) 6.42 (tt, 1H) 6.83 (m, 1H) 6.86 (d, 1H) 6.86 (d, 1H) 6.98 (d, 1H) 7.47 (dd, 1H) 7.61 (d, 1H) 7.80 (d, 1H) 8.03 (s, 1H)
17		1,1-dimethylethyl 4-[3-(3,4-dimethoxybenzyl)-6-hydroxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carboxylate	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 1.43 (s, 9H) 1.67 (d, 2H) 2.48 (m, 2H) 2.92 (brs, 2H) 3.70 (s, 3 H) 3.71 (s, 3H) 4.05 (br s, 2H) 4.61 (brs, 1H) 5.02 (s, 2H) 6.80 (dd, 1H) 6.86 (d, 1H) 6.97 (d, 1H) 7.20 (dd, 1H) 7.43 (d, 1H) 7.62 (d, 1H) 9.95 (brs, 1H)
18		3-(3,4-dimethoxybenzyl)-6-hydroxy-1-(1-methylpiperidin-4-yl)quinazoline-2,4(1H,3H)-dione	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ ppm 1.60 (d, 2H) 2.06 (td, 2H) 2.19 (s, 3H) 2.65 (qd, 2H) 2.85 (d, 2H) 3.70 (s, 3 H) 3.71 (s, 3H) 4.43 (br.s, 1H) 5.02 (s, 2H) 6.81 (m, 1H) 6.85 (d, 1H) 6.98 (d, 1H) 7.19 (dd, 1H) 7.41 (d, 1H) 7.55 (d, 1H) 9.82 (s, 1H)




19		3-(3,4-dimethoxybenzyl)-6-hydroxy-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)dione hydrochloride	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.86 (d, 2H) 2.83 (m, 2H) 3.12 (t, 2H) 3.37 (m, 2H) 3.70 (s, 6H) 4.76 (brs, 1H) 5.03 (s, 2H) 6.83 (dd, 1H) 6.85 (d, 1H) 6.98 (d, 1H) 7.21 (dd, 1H) 7.44 (d, 1H) 7.70 (d, 1H) 8.70 (brs, 2H) 9.91 (s, 1H)
20		N-([3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl)acetamide	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.75 (d, 1H) 1.78 (d, 1H) 1.86 (s, 3H) 2.40 (qd, 1H) 2.53 (qd, 1H) 2.79 (t, 1H) 3.25 (t, 1H) 3.69 (s, 3H) 3.70 (s, 3H) 3.80 (d, 1H) 4.29 (d, 2H) 4.31 (d, 1H) 4.77 (brs, 1H) 5.02 (s, 2H) 6.80 (dd, 1H) 6.85 (d, 1H) 6.96 (d, 1H) 7.65 (dd, 1H) 7.76 (d, 1H) 7.96 (d, 1H) 8.02 (s, 1H) 8.46 (t, 1H)
21		6-(aminomethyl)-3-(3,4-dimethoxybenzyl)-1-(piperidin-4-yl)quinazoline-2,4(1H,3H)-dione hydrochloride	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.89 (d, 2H) 2.83-2.92 (m, 2H) 3.16 (brs, 2H) 3.38-3.42 (m, 2H) 3.71 (s, 6H) 4.13 (brs, 2H) 4.82-4.97 (m, 1H) 5.07 (s, 2H) 6.83-6.88 (m, 2H) 7.00 (s, 1H) 7.91-7.99 (m, 2H) 8.24 (s, 1H) 8.42 (br s, 3H) 9.16 (brs, 1H)




22		1-(1-acetylpiperidin-4-yl)-6-(aminomethyl)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione hydrochloride	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.67-1.79 (m, 2H) 2.05 (s, 3H) 2.36-2.47 (m, 1H) 2.51-2.61 (m, 1H) 2.73 (t, 1H) 3.27 (t, 1H) 3.71 (s, 6H) 3.93 (d, 1H) 4.11-4.15 (m, 2H) 4.52 (d, 1H) 4.78 (brs, 1H) 5.05 (s, 2H) 6.82 (d, 1H) 6.86 (d, 1H) 6.98 (s, 1H) 7.87 (s, 2H) 8.24 (s, 1H) 8.32 (brs, 3H)
23		N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}formamide	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.73-1.83 (m, 2H) 2.36-2.45 (m, 1H) 2.48-2.57 (m, 1H) 2.77-2.84 (m, 1H) 3.23-3.29 (m, 1H) 3.70 (s, 1H) 3.71 (s, 3H) 3.81 (d, 1H) 4.31 (d, 1H) 4.37 (d, 2H) 4.77 (brs, 1H) 5.04 (s, 2H) 6.82 (dd, 1H) 6.86 (d, 1H) 6.98 (d, 1H) 7.67 (dd, 1H) 7.78 (d, 1H) 7.99 (d, 1H) 8.03 (s, 1H) 8.16 (s, 1H) 8.62 (t, 1H)
24		N-{[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}formamide	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.67-1.78 (m, 2H) 2.05 (s, 3H) 2.35-2.48 (m, 1H) 2.53-2.62 (m, 1H) 2.72 (t, 1H) 3.25 (t, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.93 (d, 1H) 4.37 (d, 2H) 4.52 (d, 1H) 4.75 (brs, 1H) 5.04 (s, 2H) 6.82 (dd, 1H) 6.86 (d, 1H) 6.98 (d, 1H) 7.67 (dd, 1H) 7.78 (d, 1H) 7.99 (d, 1H) 8.16 (s, 1H) 8.63 (t, 1H)




25		N-{[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.63-1.81 (m, 2H) 1.87 (s, 3H) 2.05 (s, 3H) 2.32-2.42 (m, 1H) 2.54-2.63 (m, 1H) 2.71 (dt, 1H) 3.25 (dt, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 3.93 (d, 1H) 4.30 (d, 2H) 4.52 (d, 1H) 4.76 (brs, 1H) 5.04 (s, 2H) 6.81 (dd 1H) 6.86 (d, 1H) 6.97 (d, 1H) 7.65 (dd, 1H) 7.76 (d, 1H) 7.97 (d, 1H) 8.47 (t, 1H)
26		6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-1-(piperidin-3-yl)quinazoline-2,4(1H,3H)-dione hydrochloride	¹ H NMR (600 MHz, DMSO-d ₆) δ ppm 1.86-1.97 (m, 3H) 2.54-2.62 (m, 1H) 2.85-2.92 (m, 1H) 3.31-3.35 (m, 1H) 3.37-3.42 (m, 1H) 3.71 (s, 3H) 3.72 (s, 3H) 3.73-3.78 (m, 1H) 3.76 (t, 1H) 4.45 (td, 2H) 4.87 (br s, 1H) 5.05 (s, 2H) 6.41 (tt, 1H) 6.83-6.88 (m, 2H) 6.99 (s, 1H) 7.52 (dd, 1H) 7.63 (d, 1H) 7.72 (d, 1H) 8.95 (brs, 1H)
27		6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-1-(pyrrolidin-3-yl)quinazoline-2,4(1H,3H)-dione hydrochloride	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 2.25-2.32 (m, 1H) 2.42-2.48 (m, 1H) 3.11-3.19 (m, 1H) 3.38-3.45 (m, 1H) 3.58-3.64 (m, 1H) 3.65-3.72 (m, 1H) 3.73 (s, 3H) 3.71 (s, 3H) 4.45 (td, 2H) 5.05-5.12 (m, 2H) 5.47-5.54 (m, 1H) 6.42 (tt, 1H) 6.82-6.88 (m, 2H) 7.01 (s, 1H) 7.55 (dd, 1H) 7.62 (d, 1H) 7.71 (d, 1H) 8.87 (brs, 1H)




	¹H NMR (600 MHz, DMSO-<i>d</i>₆) mixture of two isomers δ ppm 1.52-1.62 (m, 0.5H) 1.64-1.73 (m, 0.5H) 1.75-1.84 (m, 1H) 1.84-1.90 (m, 1H) 2.54-2.67 (m, 1.5H) 3.05 (t, 0.5H) 3.59 (t, 0.5H) 3.70 (s, 3.5H) 3.70 (s, 3H) 3.77 (dd, 0.5H) 4.06 (t, 0.5H) 4.17-4.27 (m, 1H) 4.31 (brs, 0.5H) 4.42 (td, 2H) 4.44 (brs, 0.5H) 5.04 (s, 2H) 6.38 (tt, 1H) 6.80-6.84 (m, 1H) 6.84-6.88 (m, 1H) 6.98 (brs, 1H) 7.43 (td, 1H) 7.60 (brs, 1H) 7.62 (d, 0.5H) 7.79 (d, 0.5H) 7.99 (s, 0.5H) 8.09 (s, 0.5H)
	¹H NMR (500 MHz, DMSO-<i>d</i>₆) mixture of two isomers δ ppm 1.55-1.65 (m, 0.5H) 1.67-1.89 (m, 2.5H) 1.99 (s, 1.5H) 2.06 (s, 1.5H) 2.52-2.65 (m, 1H) 3.06 (t, 0.5H) 3.30-3.37 (m, 0.5H) 3.52 (t, 0.5H) 3.70 (s, 3H) 3.71 (s, 3H) 3.52 (t, 0.5H) 3.84 (d, 0.5H) 3.91 (d, 0.5H) 4.09 (t, 0.5H) 4.28 (brs, 0.5H) 4.39-4.48 (m, 3H) 4.49 (brs, 0.5H) 5.05 (d, 2H) 6.41 (tt, 1H) 6.81-6.89 (m, 2H) 6.99 (d, 1H) 7.44-7.48 (m, 1H) 7.59-7.62 (m, 1H) 7.64 (d, 0.5H) 7.82 (d, 0.5H)




	3-([6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]pyrrolidine-1-carbaldehyde	¹H NMR (500 MHz, DMSO-d₆) mixture of two isomers δ ppm 2.18-2.27 (m, 1H) 2.36-2.46 (m, 1H) 3.41-3.48 (m, 0.5H) 3.53-3.60 (m, 0.5H) 3.63-3.73 (m, 1.5H) 3.85-3.95 (m, 1.5H) 4.43 (td, 2H) 5.06 (s, 2H) 5.38-5.45 (m, 0.5H) 5.46-5.54 (m, 0.5H) 6.41 (tt, 1H) 6.82-6.88 (m, 2H) 7.00 (s, 1H) 7.46 - 7.51 (m, 1H) 7.61 (d, 1H) 7.66 (d, 0.5H) 7.72 (d, 0.5H) 8.16 (s, 0.5H) 8.20 (s, 0.5H)
	1-(1-acetylpyrrolidin-3-yl)-6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	¹H NMR (500 MHz, DMSO-d₆) mixture of two isomers δ ppm 1.98 (s, 1.5H) 2.04 (s, 1.5H) 2.17-2.25 (m, 0.5H) 2.27-2.35 (m, 0.5H) 2.57-2.66 (m, 1H) 3.40-3.45 (m, 0.5H) 3.56-3.61 (m, 0.5H) 3.70-3.78 (m, 7.5H) 3.82-3.95 (m, 1.5H) 4.48 (td, 2H) 5.10 (d, 2H) 5.45-5.58 (m, 1H) 6.46 (tt, 1H) 6.87-6.92 (m, 2H) 7.05 (dd, 1H) 7.53 (td, 1H) 7.66 (t, 1H) 7.72 (dd, 1H)
	4-[6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹H NMR (600 MHz, DMSO-d₆) δ ppm 1.72-1.82 (m, 2H) 2.35-2.44 (m, 1H) 2.47-2.55 (m, 1H) 2.79 (td, 1H) 3.25 (td, 1H) 3.80 (d, 1H) 4.32 (d, 1H) 4.42 (td, 2H) 4.71 (brs, 1H) 6.40 (tt, 1H) 7.43 (dd, 1H) 7.55 (d, 1H) 7.74 (d, 1H) 8.03 (s, 1H)



33 	4-[3-(3,4-dichlorobenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹H NMR (500 MHz, DMSO-d₆) δ ppm 1.80 (dd, 2H) 2.32-2.43 (m, 1H) 2.44-2.52 (m, 1H) 2.80 (td, 1H) 3.26 (dt, 1H) 3.80 (brd, 1H) 4.31 (br d, 1H) 4.44 (td, 2H) 4.78 (brs, 1H) 5.09 (s, 2H) 6.41 (tt, 1H) 7.30 (dd, 1H) 7.49 (dd, 1H) 7.58 (d, 1H) 7.60 (d, 1H) 7.62 (d, 1H) 7.81 (d, 1H) 8.03 (s, 1H)
34 	4-[3-(4-chlorobenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹H NMR (500 MHz, DMSO-d₆) δ ppm 1.72-1.83 (m, 2H) 2.32-2.44 (m, 1H) 2.45-2.53 (m, 1H) 2.80 (dt, 1H) 3.25-3.30 (dt, 1H) 3.80 (d, 1 H) 4.30 (d, 1 H) 4.44 (td, 2H) 4.78 (brs, 1H) 5.09 (s, 2H) 6.42 (tt, 1H) 7.32-7.40 (m, 4H) 7.48 (dd, 1H) 7.61 (d, 1H) 7.81 (d, 1H) 8.03 (s, 1H)
35 	Methyl 4-{[6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl}benzoate	¹H NMR (500 MHz, DMSO-d₆) δ ppm 1.78 (dd, 1H) 2.33-2.42 (m, 1H) 2.44-2.52 (m, 1H) 2.80 (dt, 1H) 3.26 (dt, 1H) 3.79 (brd, 1H) 3.84 (s, 3H) 4.30 (brd, 1H) 4.45 (td, 2H) 4.79 (brs, 1H) 5.18 (s, 2 H) 6.42 (tt, 1H) 7.43 (d, 2H) 7.49 (dd, 1H) 7.62 (d, 1H) 7.83 (d, 1H) 7.91 (d, 2H) 8.02 (s, 1H)



36		4-([6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl)benzoic acid	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.79(dd, 2H) 2.38(qd, 1H) 2.49(qd, 1H) 2.80(dt, 1H) 3.26(dt, 1H) 3.79(d, 1H) 4.31(d, 1H) 4.45(td, 2H) 4.79(brs, 1H) 5.17(s, 2H) 6.42(tt, 1H) 7.40(d, 2H) 7.49(dd, 1H) 7.62(d, 1H) 7.83(d, 1H) 7.88(d, 2H) 8.02(s, 1H) 12.94(brs, 1H)
37		4-([6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl)-N-(2-methoxyethyl)benzamide	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.72-1.82(m, 2H) 2.32-2.42(m, 1H) 2.44-2.54(m, 1H) 2.75-2.83(m, 1H) 3.21-3.29(m, 4H) 3.37-3.45(m, 4H) 3.77(br.d., 1H) 4.29(br.d., 1H) 4.44(td, 2H) 4.78(br.s., 1H) 5.14(s, 2H) 6.41(tt, 1H) 7.35(m, 2H) 7.48(dd, 1H) 7.61(d, 1H) 7.77(m, 2H) 7.81(d, 1H) 8.01(s, 1H) 8.47(t, 1H)
38		4-[3-(3,4-dimethoxybenzyl)-6-methyl-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.76(dd, 2H) 2.37(s, 3H) 2.39-2.42(m, 1H) 2.50-2.54(m, 1H) 2.77-2.82(m, 1H) 3.19-3.30(m, 1H) 3.70(s, 6H) 3.79(d, 1H) 4.30(d, 1H) 4.76(brs, 1H) 5.02(s, 2H) 6.82(dd, 1H) 6.85(d, 1H) 6.97(d, 1H) 7.60(dd, 1H) 7.71(d, 1H) 7.89(s, 1H) 8.03(s, 1H)



39		4-[6-(2,2-difluoroethoxy)-3-(3-hydroxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.70-1.80 (m, 2H) 2.35-1.45 (m, 1H) 2.47-2.56 (m, 1H) 2.80 (dt, 1H) 3.25 (dt, 1H) 3.70 (s, 3H) 3.79 (brd, 1H) 4.30 (brd, 1H) 4.43 (td, 2H) 4.77 (brs, 1H) 4.96 (s, 2H) 6.41 (tt, 1H) 6.73 (d, 1H) 6.78-6.82 (m, 2H) 7.46 (dd, 1H) 7.60 (d, 1H) 7.79 (d, 1H) 8.03 (s, 1H) 8.95 (s, 1H)
40		4-[6-(2,2-difluoroethoxy)-3-(3-(2-hydroxyethoxy)-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (600 MHz, DMSO-d ₆) δ ppm 1.76 (dd, 2H) 2.35-2.45 (m, 1H) 2.50-2.54 (m, 1H) 2.79 (dt, 1H) 3.25 (dt, 1H) 3.68 (dt, 2H) 3.70 (s, 3H) 3.79 (brd, 1H) 3.90 (t, 2H) 4.30 (brd, 1H) 4.42 (td, 2H) 4.75 (brs, 1H) 4.77 (t, 1H) 5.01 (s, 2H) 6.39 (tt, 1H) 6.85 (m, 2H) 6.97 (s, 1H) 7.45 (dd, 1H) 7.60 (d, 1H) 7.77 (d, 1H) 8.02 (s, 1H)
41		4-[6-(2,2-difluoroethoxy)-3-(3-ethoxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (600 MHz, DMSO-d ₆) δ ppm 1.29 (t, 3H) 1.76 (dd, 2H) 2.33-2.45 (m, 1H) 2.50-2.56 (m, 1H) 2.76-2.82 (m, 1H) 3.21-3.31 (m, 1H) 3.70 (s, 3H) 3.79 (brd, 1H) 3.94 (q, 2H) 4.30 (brd, 1H) 4.41 (dt, 2H) 4.75 (brs, 1H) 5.01 (s, 2H) 6.39 (tt, 1H) 6.81-6.86 (m, 2H) 6.96 (s, 1H) 7.45 (dd, 1H) 7.60 (d, 1H) 7.77 (d, 1H) 8.02 (s, 1H)



42		4-[6-(2,2-difluoroethoxy)-3-[(4-methoxy-3-(2-methoxyethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (600 MHz, DMSO-d ₆) δ ppm 1.69-1.82 (m, 2H) 2.35-2.45 (m, 1H) 2.45-2.55 (m, 1H) 2.78 (t, 1H) 3.21-3.28 (m, 1H) 3.61 (brs, 2H) 3.70 (s, 3H) 3.78 (d, 1H) 4.00 (brs, 2H) 4.30 (d, 1H) 4.42 (t, 2H) 4.75 (brs, 1H) 5.01 (s, 2H) 6.39 (t, 1H) 6.82-6.88 (m, 2H) 6.97 (s, 1H) 7.45 (d, 1H) 7.59 (s, 1H) 7.77 (d, 1H) 8.02 (s, 1H)
43		4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azepane-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) mixture of two isomers δ ppm 1.72-1.81 (m, 1H) 1.85 (brd, 1H) 1.86-2.03 (m, 2H) 2.47 (brs, 2H) 3.30-3.40 (m, 1H) 3.42-3.52 (m, 1H) 3.58-3.64 (m, 2H) 3.70 (s, 3H) 3.71 (s, 3H) 4.42 (td, 2H) 4.45 (brs, 0.5H) 5.04 (brs, 2H) 5.21 (brs, 0.5H) 6.40 (tt, 1H) 6.78-6.83 (m, 1H) 6.86 (d, 1H) 6.98 (s, 1H) 7.38-7.48 (m, 1H) 7.59 (brs, 1H) 7.64-7.74 (m, 1H) 8.08 (brs, 0.5H) 8.12 (s, 0.5H)
44		3-(3,4-dimethoxybenzyl)-2,4-dioxo-1-(pyrrolidin-3-yl)-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	¹ H NMR (600 MHz, DMSO-d ₆) δ ppm 1.96-2.09 (m, 2H) 2.78-2.84 (m, 1H) 3.05-3.12 (m, 2H) 3.18-3.27 (m, 1H) 3.70 (s, 3H) 3.71 (s, 3H) 5.04 (s, 2H) 5.47-5.53 (m, 1H) 6.83 (brs, 2H) 6.99 (s, 1H) 8.15 (dd, 1H) 8.20 (d, 1H) 8.47 (d, 1H)



45		1-(1-acetylpyrrolidin-3-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	¹H NMR (600 MHz, DMSO-d₆) mixture of two isomers δ ppm 1.91(s, 1.5H) 1.98(s, 1.5H) 2.14-2.22(m, 0.5H) 2.25-2.32(m, 0.5H) 2.43-2.50(m, 0.5H) 2.52-2.58(m, 0.5H) 3.31-3.39(m, 0.5H) 3.50-3.55(m, 0.5H) 3.63-3.73(m, 7.5H) 3.79-3.87(m, 1.5H) 5.00-5.06(m, 2H) 5.43-5.55(m, 1H) 6.82-6.90(m, 2H) 6.98(d, 1H) 7.84(dd, 1H) 8.17(dt, 1H) 8.43(dd, 1H)
46		3-(3,4-dimethoxybenzyl)-1-(1-formylpyrrolidin-3-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	¹H NMR (500 MHz, DMSO-d₆) mixture of two isomers δ ppm 2.19-2.27(m, 1H) 2.36-2.43(m, 1H) 3.43-3.47(m, 0.4H) 3.53-3.59(m, 0.6) 3.62-3.73(m, 1.6H) 3.70(s, 3H) 3.71(s, 3H) 3.84-3.94(m, 1.4H) 5.04(s, 2H) 5.42-5.49(m, 0.4H) 5.50-5.56(m, 0.6H) 6.86(s, 2H) 6.99(s, 1H) 7.85(d, 0.6H) 7.90(d, 0.4H) 8.15-8.22(m, 2H) 8.45(s, 1H)
47		4-[6-(2,2-difluoroethoxy)-3-[3-(3-hydroxypropoxy)-4-methoxybenzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹H NMR (600 MHz, DMSO-d₆) δ ppm 1.72-1.79(m, 2H) 1.82(q, 2H) 2.36-2.44(m, 1H) 2.49-2.55(m, 1H) 2.78(td, 1H) 3.25(td, 1H) 3.51-3.54(m, 2H) 3.70(s, 3H) 3.76-3.80(m, 1H) 3.95(t, 2H) 4.28-4.32(m, 1H) 4.41(td, 2H) 4.47(t, 1H) 4.75(s, 1H) 5.01(s, 2H) 6.39(tt, 1H) 6.81-6.86(m, 2H) 6.96-6.97(m, 1H) 7.45(dd, 1H) 7.59(d, 1H) 7.77(d, 1H) 8.02(s, 1H)



48		4-[5-chloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.78-1.88 (m, 2H) 2.34-2.44 (m, 1H) 2.46-2.56 (m, 1H) 2.80 (t, 1H) 3.25 (t, 1H) 3.71 (s, 6H) 3.80 (d, 1H) 4.31 (d, 1H) 4.71 (brs, 1H) 5.00 (s, 2H) 6.83 (d, 1H) 6.87 (d, 1H) 6.97 (s, 1H) 7.37 (d, 1H) 7.69 (dd, 1H) 7.77 (d, 1H) 8.03 (s, 1H)
49		4-{3-[3-(cyclopentyloxy)-4-(methoxybenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.5-1.6 (m, 2H) 1.63-1.71 (m, 3H) 1.68-1.78 (m, 1H) 1.78-1.98 (m, 3H) 2.35-2.46 (m, 1H) 2.47-2.57 (m, 1H) 2.79 (dt, 1H) 3.26 (dt, 1H) 3.69 (s, 3H) 3.79 (br.d., 1H) 4.30 (br.d., 1H) 4.43 (td, 2H) 4.66-4.70 (m, 1H) 4.76 (br.s., 1H) 5.01 (s, 2H) 6.41 (tt, 1H) 6.82 - 6.86 (m, 2H) 6.96 (s, 1H) 7.46 (dd, 1H) 7.6 (d, 1H) 7.80 (d, 1H) 8.03 (s, 1H)





50		2-(5-{[6-(2,2-difluoroethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]-2-methyl}-2-methoxyphenoxy)-acetamide	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 1.72-1.82 (m, 2H) 2.35-2.45 (m, 1H) 2.46-2.56 (m, 1H) 2.79 (td, 1H) 3.25 (td, 1H) 3.74 (s, 3H) 3.79 (d, 1H) 4.31 (d, 1H) 4.35 (s, 2H) 4.43 (td, 2H) 4.76 (brs, 1H) 5.00 (s, 2H) 6.41 (tt, 1H) 6.88-6.95 (m, 3H) 7.31 (brs, 1H) 7.37 (brs, 1H) 7.46 (dd, 1H) 7.59 (d, 1H) 7.79 (d, 1H) 8.02 (s, 1H)
51		4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]-3-methylpiperidine-1-carbaldehyde	¹ H NMR (500 MHz, DMSO-d ₆) mixture of two isomers δ ppm 0.92 (t, 3H) 1.72-1.82 (m, 1H) 2.23 (brs, 1H) 2.70-2.77 (m, 0.5H) 3.00-3.10 (m, 0.5H) 3.12-3.20 (m, 1.5H) 3.44 (d, 0.5H) 3.63 (dd, 0.5H) 3.69 (s, 6H) 3.80 (d, 0.5H) 4.04 (d, 0.5H) 4.29 (d, 0.5H) 4.43 (td, 2H) 4.62-4.70 (m, 1H) 5.03 (s, 2H) 6.41 (tt, 1H) 6.80 (dd, 1H) 6.85 (d, 1H) 6.97 (d, 1H) 7.46 (dd, 1H) 7.60 (d, 1H) 7.64 (dd, 1H) 7.97 (s, 0.5H) 8.11 (s, 0.5H)




52		3-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]-8-azabicyclo[3.2.1]octane-8-carbaldehyde	¹H NMR (600 MHz, DMSO-<i>d</i>₆) δ ppm 1.85-1.95 (m, 4H) 2.18 (t, 2H) 2.27-2.33 (m, 1H) 2.34-2.41 (m, 1H) 3.70 (s, 1H) 3.71 (s, 1H) 4.25-4.29 (m, 1H) 4.40 (td, 2H) 4.50-4.54 (m, 1H) 4.55 (br.s., 1H) 5.05 (s, 2H) 6.38 (tt, H) 6.80 (d, 1H) 6.85 (d, 1H) 6.98 (s, 1H) 7.45 (dd, 1H) 7.56 (d, 1H) 7.58 (d, 1H) 8.12 (s, 1H)
53		3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-3-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	¹H NMR (500 MHz, DMSO-<i>d</i>₆) mixture of two isomers δ ppm 1.52-1.63 (m, 0.5H) 1.65-1.73 (m, 0.5H) 1.74-1.83 (m, 1H) 1.87-1.93 (m, 1H) 2.54-2.64 (m, 1.5H) 3.06 (dt, 0.5H) 3.57 (t, 0.5H) 3.70 (s, 3H) 3.71 (s, 3H) 3.79 (dd, 0.5H) 4.03 (t, 0.5H) 4.20 (d, 0.5H) 4.30 (d, 0.5H) 4.38 (brs, 0.5H) 4.53 (brs, 0.5H) 4.99-5.07 (m, 2H) 6.84-6.88 (m, 2H) 6.98 (s, 1H) 7.86 (d, 0.5H) 8.01 (d, 0.5H) 7.98 (s, 0.5H) 8.08 (s, 0.5H) 8.14-8.18 (m, 1H) 8.44 (s, 1H)




54 	1-(1-acetylpiperidin-3-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	¹H NMR (500 MHz, DMSO-d₆) mixture of two isomers δ ppm 1.56-1.66(m, 0.5H) 1.70-1.82(m, 1.5H) 1.87(brd, 1H) 1.97(s, 1.5H) 2.05(s, 1.5H) 2.54(dd, 1H) 3.05(brt, 0.5H) 3.48(t, 0.5H) 3.70-3.72(m, 6H) 3.83(d, 0.5H) 3.94(brd, 0.5H) 4.04(t, 0.5H) 4.33(brs, 0.5H) 4.45(dd, 1H) 4.53(brs, 0.5H) 4.99-5.07(m, 2H) 6.83-6.87(m, 2H) 6.98(d, 1H) 7.84(d, 0.5H) 8.02(d, 0.5H) 8.16(dt, 1H) 8.44(dd, 1H)
55 	4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	¹H NMR (600 MHz, DMSO-d₆) δ ppm 1.73-1.82(m, 2H) 2.36-2.44(m, 1H) 2.44-2.55(m, 1H) 2.79(td, 1H) 3.25(td, 1H) 3.80(brd, 1H) 4.32(brd, 1H) 4.62-4.67(m, 1H) 4.68-4.76(m, 3H) 4.76-4.82(m, 1H) 4.96-5.06(m, 1H) 7.44(dd, 1H) 7.61(d, 1H) 7.73(d, 1H) 8.03(s, 1H) 11.46(brs, 1H)




56		4-{3-[4-(cyclopentyloxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 572
57		4-[3-(3-chlorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 492




58		4-[3-(4-chlorobenzyl)-6-[2-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] + 492
59		4-{3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-[2-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	LCMS m/z [MH] + 572
60		6-bromo-3-(3,4-dimethoxybenzyl)-1-(1-(hydroxyacetyl)piperidin-3-yl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] + 532




61		1-(1-acetylpiperidin-3-yl)-6-bromo-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 516
62		3-(3,4-dimethoxybenzyl)-1-(hydroxyacetyl)piperidin-3-yl]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 454
63		6-bromo-3-(3,4-dimethoxybenzyl)-1-[piperidin-3-yl]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 474
64		N-{[1-(1-acetylpiperidin-3-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide	LCMS m/z [MH] ⁺ + 495





65		3-[6-bromo-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] + 502
66		6-bromo-1-[1-(cyclopropyl)-carbonyl]piperidin-3-yl]-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] + 542
67		1-{1-[(benzyloxy)acetyl]piperidin-3-yl}-6-bromo-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] + 622




68		1-{1-[(benzyloxy)acetyl]piperidin-3-yl}-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 638
69		3-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ + 518
70		3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-[1-(hydroxyacetyl)piperidin-3-yl]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 548




71		3-[3-(3,4-dimethoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 484
72		4-[3-(3,4-dimethoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 484
73		1-[1-(cyclopropylcarbonyl)-piperidin-3-yl]-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 558




74		4-[3-(3,4-dichlorobenzyl)-6-(2-fluoro-1-(fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 526
75		3-(3,4-dimethoxybenzyl)-6-(2-fluoro-1-(fluoromethyl)ethoxy)-1-(piperidin-3-yl)quinazolin-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 490
76		4-{3-[(6-chloropyridin-3-yl)methyl]-6-(2-fluoro-1-(fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 493



77		N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpyrrolidin-3-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide	LCMS m/z [MH] ⁺ 481
78		4-[3-(3-chloro-4-methoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)-ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 522
79		4-[3-(3,4-dimethoxybenzyl)-6-(2-fluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 486



80		6-bromo-3-(3,4-dimethoxybenzyl)-1-[pyrrolidin-3-yl]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 460
81		6-bromo-1-[1-(cyclopropyl)pyrrolidin-3-yl]-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 528
82		1-(1-acetylpyrrolidin-3-yl)-6-bromo-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 502
83		3-[6-bromo-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]pyrrolidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 488




84		1-[1-(cyclopropylcarbonyl)-piperidin-3-yl]-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	LCMS m/z [MH] ⁺ 489
85		3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(pyrrolidin-3-yl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 476
86		3-(3,4-dimethoxybenzyl)-1-[1-(hydroxyacetyl)piperidin-3-yl]-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	LCMS m/z [MH] ⁺ 479




87		1-[1-(cyclopropylcarbonyl)-pyrrolidin-3-yl]-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carbonitrile	LCMS m/z [MH] ⁺ + 475
88		1-[1-[(benzyloxy)acetyl]pyrrolidin-3-yl]-6-bromo-3-(3,4-dimethoxybenzyl)quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 608
89		2-[5-({6-[2-fluoro-1-(methyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]acetamide	LCMS m/z [MH] ⁺ + 561




90		4-{6-[2-fluoro-1-(fluoro-methyl)ethoxy]-3-(3-hydroxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 504
91		4-[3-(3,4-dimethoxybenzyl)-6-ethoxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 468
92		1-[1-(cyclopropylcarbonyl)-pyrrolidin-3-yl]-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 544




93		1-{1-[(benzyloxy) acetyl] -pyrrolidin-3-yl }-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ + 624
94		3-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]pyrrolidine-1-carbaldehyde	LCMS m/z [MH] ⁺ + 504

95		N-({1-[(1-cyclopropylcarbonyl)-pyrrolidin-3-yl]-3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl}methyl)acetamide	LCMS m/z [MH] ⁺ 521
96		3-({3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1H-[1-(hydroxyacetyl)pyrrolidin-3-yl]quinazoline-2,4(1H,3H)-dione	LCMS m/z [M+Na] ⁺ 534



97		4-[5,7-dichloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ + 492
98		1,1-dimethylethyl 3-{6-[(acetylamino)methyl]-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carboxylate	LCMS m/z [M+Na] ⁺ + 589
99		N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-3-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide	LCMS m/z [MH] ⁺ + 495



100		N-({1-[(1-acetyl-3-(4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide)	LCMS m/z [MH] ⁺ 509
101		N-({1-[(1-(cyclopropylcarbonyl)-3-(4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide)	LCMS m/z [MH] ⁺ 535
102		4-[7-chloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 458




103		N-({3-(3,4-dimethoxybenzyl)-1-(hydroxyacetyl)piperidin-3-yl}-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl)methyl)acetamide	LCMS m/z [MH] ⁺ 525
104		3-(3,4-dimethoxybenzyl)-1-(hydroxyacetyl)pyrrolidin-3-yl]-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-carbonitrile	LCMS m/z [MH] ⁺ 465
105		6-bromo-3-(3,4-dimethoxybenzyl)-1-(hydroxyacetyl)pyrrolidin-3-yl]quinazoline-2,4(1H,3H)-dione	LCMS m/z [MH] ⁺ 518







106		3-[3-(3,4-dimethoxybenzyl)-6-methyl-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]pyrrolidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 424
107		N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpyrrolidin-3-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}formamide	LCMS m/z [MH] ⁺ 467
108		4-{6-[2-fluoro-1-(fluoromethyl)-ethoxy]-3-(3-fluoro-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 506









109		4-[6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] + 530
110		1,1-dimethylethyl 3-[6-(benzyloxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azetidine-1-carboxylate	LCMS m/z [MH] + 574
111		4-[6-(difluoromethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] + 490









112		4-[3-(3,4-dimethoxybenzyl)-6-(1-methylethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 482
113		1,1-dimethylethyl 3-[3-(3,4-dimethoxybenzyl)-6-(2-fluoro-1-(fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azetidine-1-carboxylate	LCMS m/z [MH] ⁺ 562









114		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-methoxy-3-(1-methylethoxy)-benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	LCMS m/z [MH] ⁺ 546
115		1,1-dimethylethyl 3-[3-(3,4-dimethoxybenzyl)-6-hydroxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azetidine-1-carboxylate	LCMS m/z [MH] ⁺ 484









116		4-{6-[2-fluoro-1-(fluoromethyl)-3-(3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-yl}carbaldehyde	LCMS m/z [MH] ⁺ 488
117		4-{3-[3,5-bis(trifluoromethyl)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-yl}carbaldehyde	LCMS m/z [MH] ⁺ 564
118		4-[3-(3-ethoxybenzyl)-6-[2-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-yl}carbaldehyde	LCMS m/z [MH] ⁺ 502









COMPOUND No.	STRUCTURE	NOMENCLATURE	MASS LCUVM S MH+
119		4-[3-(3,4-dimethoxybenzyl)-6-hydroxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	440
120		1,1-dimethylethyl 3-({4-(benzyloxy)-2-[(3,4-dimethoxybenzyl)carbamoyl]phenyl}amino)azetidine-1-carboxylate	548
121		4-{3-[3-chloro-4-(cyclopentyloxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	576
122		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-fluoro-4-methylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	490
123		4-[3-(3,5-difluorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	494
124		4-{3-[3-chloro-4-(2-methoxyethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	566









125		4-{3-[(2-chloropyridin-4-yl)methyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	493
126		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	542
127		4-[3-(3,5-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	518
128		4-[3-(3-fluorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	476
129		4-[3-(4-fluorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	476
130		4-[3-(3,4-diethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	546
131		4-[3-(4-ethoxy-3-methoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	532
132		4-[3-(2,3-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	518








133		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-methoxy-3-methylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	502
134		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(trifluoromethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	526
135		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(3-phenoxybenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	550
136		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(3-fluoropropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	534
137		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-nitrobenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	503
138		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-nitrobenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	503
139		1-(1-acetylazetidin-3-yl)-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	504
140		4-[3-(2,5-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	518








141		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	488
142		4-{3-[3-(benzyloxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	564
143		4-{3-[4-(benzyloxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	594
144		4-[3-(3,5-dichlorobenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	526
145		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-nitrobenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	533
146		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-{4-[(trifluoromethyl)sulphanyl]benzyl}-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	558
147		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(pyrrolidin-1-ylmethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	541
148		1-[1-(cyclopropylcarbonyl)azetidin-3-yl]-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	530









149		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(piperidin-1-ylmethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	555
150		3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-[1-(hydroxyacetyl)azetidin-3-yl]quinazoline-2,4(1H,3H)-dione	520
151		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)benzonitrile	483
152		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[3-(trifluoromethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	526
153		3-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azetidine-1-carbaldehyde	490
154		1-{1-[(benzyloxy)acetyl]azetidin-3-yl}-3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazoline-2,4(1H,3H)-dione	610
155		4-[3-(4-ethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	502
156		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[3-(piperidin-1-ylmethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	555

157		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	557
158		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	557
159		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-(2-(morpholin-4-ylethoxy)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	587
160		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(morpholin-4-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	543
161		4-[3-(1-benzofur-6-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	498
162		1,1-dimethylethyl 4-[4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)benzyl]piperazine-1-carboxylate	656
163		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(6-methoxypyridin-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	489
164		4-{3-[3-chloro-4-(1-methylethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	550









		1 (2H)-yl}piperidine-1-carbaldehyde	
165		4-[3-(biphenyl-4-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	534
166		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(methylsulphanyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	504
167		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyridin-3-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	535
168		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(3-(pyrrolidin-1-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	527
169		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-fluoro-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	506
170		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-methylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	502
171		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-methyl-3-(trifluoromethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	540









172		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{4-[(4-methylpiperazin-1-yl)methyl]benzyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl)piperidine-1-carbaldehyde	570
173		4-[3-{4-(cyclopentylloxy)-3-[2-fluoro-1-(fluoromethyl)ethoxy]benzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	636
174		3-[3,4-bis(cyclopentylloxy)benzyl]-1-(1-ethenylpiperidin-4-yl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]quinazolin-2,4(1H,3H)-dione	626
175		2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]acetamide	615
176		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl)piperidine-1-carbaldehyde	540
177		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(methylsulphonyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	536
178		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-propoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	546









179		4-[3-(3-chloro-4-ethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	536
180		4-{3-[4-(1,1-dimethylethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	530
181		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[3-(trifluoromethoxy)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	542
182		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-fluoro-4-(trifluoromethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	544
183		2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-methylacetamide	629
184		2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N,N-dimethylacetamide	643
185		2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-methoxy-N-methylacetamide	659








186		4-{3-[4-(cyclopentyloxy)-3-ethoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	586
187		4-{3-[3-(cyclobutylmethoxy)-4-(cyclopentyloxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	626
188		4-{3-[4-(cyclopentyloxy)-3-(1-methylethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	600
189		4-{3-[4-(cyclopentyloxy)-3-propoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	600
190		4-{3-[4-(cyclopentyloxy)-3-hydroxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	558
191		4-{3-[4-(cyclopentyloxy)-3-(2-methylpropoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	614
192		4-[3-(1,3-benzodioxol-5-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	502
193		4-{3-[4-(difluoromethoxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	554









194		4-{3-[4-(difluoromethoxy)-3-ethoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	568
195		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-hydroxy-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	504
196		4-[3-(3,4-dihydro-2H-1,5-benzodioxepin-7-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	530
197		4-[3-(2,3-dihydro-1-benzofur-5-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	500
198		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-fluoro-5-(trifluoromethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	544
199		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-1H-benzotriazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	513
200		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(thiophen-3-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	540
201		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyridin-4-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	535









202		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(piperazin-1-ylmethyl)benzyl])-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	555
203		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-1H-indol-6-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	511
204		4-{3-[3-chloro-4-(difluoromethoxy)-5-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	588
205		4-{3-[2-(difluoromethoxy)-3-ethoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	568
206		4-{3-[4-(cyclopropylmethoxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	558
207		2-[4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]-N-methylacetamide	575
208		4-[3-(biphenyl-3-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	534
209		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{3-[(4-methylpiperazin-1-yl)methyl]benzyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	570








210		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[3-(1H-1,2,4-triazol-1-yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	525
211		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(2-(morpholin-4-yl)ethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	587
212		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(1H-pyrazol-1-yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	524
213		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyridin-2-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	535
214		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(3-(piperidin-1-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	541
215		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(thiophen-2-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	540
216		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(quinolin-7-ylmethyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	509
217		4-{3-[4-(difluoromethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	524









218		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(6-methoxynaphth-2-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	538
219		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(3-(1H-pyrazol-1-yl)benzyl)-2,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	524
220		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-(morpholin-4-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	543
221		4-[3-{4-[2-(dimethylamino)ethoxy]benzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	545
222		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-(2-methyl-1,3-thiazol-4-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	555
223		4-{3-[4-(1H-benzimidazol-1-yl)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	574
224		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-methylpropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	560
225		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(piperidin-1-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	541









226		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(tetrahydrofuran-3-yloxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	574
227		4-{3-[4-(4-ethylpiperazin-1-yl)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	570
228		4-[3-{4-[(1-benzylpyrrolidin-3-yl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	663
229		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenyl 1-methylethyl carbonate	590
230		4-[3-(1-benzothiophen-5-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	514
231		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(pyrrolidin-3-yloxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	573
232		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	546
233		4-[3-(3,4-dimethoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	536

234		4-[3-{4-[(1-acetylpyrrolidin-3-yl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	615
235		4-{3-[3-chloro-4-(cyclopropylmethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	562
236		4-{3-[3-chloro-4-(cyclobutylmethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	576
237		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-(morpholin-4-ylsulphonyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	607
238		4-[3-(3,4-dihydroquinazolin-1(2H)-ylpiperidine-1-carbaldehyde)-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	518
239		4-[3-{4-[(4-fluorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	612
240		4-[3-{4-[(4-chlorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	628
241		4-[3-(1,3-benzothiazol-6-ylmethyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	515








242		4-[3-{4-[(3-chlorobenzyl)oxy]-3-methoxybenzyl}-6-{2-fluoro-1-(fluoromethyl)ethoxy}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	628
243		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(3-(thiophen-3-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	540
244		4-[3-(3,4-diethoxybenzyl)-6-(2,2-difluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	532
245		4-[3-(4-ethoxy-3-methoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	498
246		4-[3-{4-[2-(2,3-dihydro-1H-indol-1-yl)-2-oxoethoxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	663
247		4-[6-(2,2-difluoroethoxy)-3-(4-ethoxy-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	518
248		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-(pyridin-3-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	565
249		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde)	516









		3 (2H)-yl}methyl}benzoate	
250		4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	662
251		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-oxo-2-(piperidin-1-yl)ethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	629
252		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-(pyrimidin-5-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	566
253		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(4'-fluoro-2-methoxybiphenyl-4-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	582
254		4-{3-[3-ethoxy-4-(thiophen-2-ylmethoxy)benzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	614
255		4-[3-(3,4-diethoxybenzyl)-6-(2-fluoroethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	514
256		4-[3-(3,4-dimethoxybenzyl)-6-(3-fluoro-2-hydroxypropoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	516







257		1,1-dimethylethyl 4-({4-[2-fluoro-1-(fluoromethyl)ethoxy]-2-(methoxycarbonyl)phenyl}amino)piperidine-1-carboxylate	429
258		4-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(hydroxymethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	516
259		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[3-(1H-pyrazol-1-ylmethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	538
260		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-(thiophen-2-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	570
261		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)benzoic acid	502
262		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[3-(1H-1,2,4-triazol-1-ylmethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	539
263		(2R)-2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]propanoic acid	630
264		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-3-(thiophen-2-yl)-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	544








265		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(1H-1,2,4-triazol-1-ylmethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	539
266		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-N-(2-(morpholin-4-yl)ethyl)benzamide	614
267		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(5-(thiophen-3-yl)-1,2,4-oxadiazol-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	532
268		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(4-methyl-2-phenylpyrimidin-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	550
269		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(5-(thiophen-2-yl)-1,2,4-oxadiazol-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	532
270		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	540
271		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(2-(thiophen-2-yl)-1,3-thiazol-4-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	547
272		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(2-fluoro-4-(thiophen-2-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	558








273		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-5-(thiophen-2-yl)-1H-pyrazol-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	544
274		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(3-(thiophen-2-yl)-1,2,4-oxadiazol-5-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	532
275		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyrimidin-5-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	536
276		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-3-phenyl-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	538
277		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(5-phenyl-1,2,4-oxadiazol-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	526
278		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-{[6-(1H-pyrazol-1-yl)pyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	525
279		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(2-(thiophen-2-yl)pyrimidin-5-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	542
280		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	538









281		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(imidazo[1,2-a]pyridin-7-yl)methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	598
282		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(3-methyl-1,2,4-oxadiazol-5-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	540
283		[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]acetic acid	616
284		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(5-(thiophen-2-yl)isoxazol-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	531
285		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(thieno[2,3-b]pyridin-2-ylmethyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	514
286		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(6-phenylpyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	535
287		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(6-(morpholin-4-yl)pyridin-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	544






288		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(2-phenyl-1,3-thiazol-4-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	541
289		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(6-(thiophen-2-yl)pyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	541
290		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[(4-(thiophen-2-yl)-1,3-oxazol-4-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	531
291		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(4-methoxyphenyl)-1,2,4-oxadiazol-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	556
292		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(1-methyl-5-phenyl-1H-pyrazol-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	538
293		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(5-(2-methoxyphenyl)-1,2,4-oxadiazol-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	556
294		4-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}biphenyl-2-carbonitrile	559
295		(2R)-2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]-N-methyl-2-carbonitrile	643









		methylpropanamide	
296		(2R)-2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-phenylpropanamide	705
297		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(thiophen-2-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	558
298		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	587
299		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(piperidin-1-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	585
300		4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
301		2-[2-(cyclopentylloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-ethylacetamide	643









302		(2S)-2-[2-(cyclopentyloxy)-5-({6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)phenoxy]propanoic acid	630
303		4-[3-(3-fluoro-4,5-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	536
304		4-[3-(2-fluoro-4,5-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	536
305		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-{[(3R)-2-oxo-1-phenylpyrrolidin-3-yl]oxy}benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	663
306		4-{3-[4-(cyclobutylmethoxy)-3-methoxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	572
307		4-{3-[4-(benzyloxy)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	612
308		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-hydroxy-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	522





309		4-{3-[4-(cyclopropylmethoxy)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	576
310		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-methylpropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	578
311		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(1-methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	564
312		4-[3-(4-ethoxy-3-methoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	550
313		3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(6-oxopiperidin-3-yl)quinazoline-2,4(1H,3H)-dione	504
314		4-{6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(5-methoxy-1-methyl-4-oxo-1,4-dihydropyridin-2-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	519
315		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[(6-(3-methoxyphenyl)pyridin-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	583









316		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{[6-(2-fluorophenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	571
317		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{[6-(4-fluorophenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	571
318		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{[6-(4-methoxyphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	583
319		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-{[6-(thiophen-2-yl)pyridin-3-yl]methyl}-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	559
320		4-{3-[3-ethoxy-4-(thiophen-2-ylmethoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	632
321		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	556
322		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-(4-(pyrimidin-5-yl)benzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	554
323		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[1-methyl-3-(thiophen-2-yl)-1H-pyrazol-5-yl]methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	562









		carbaldehyde	
324		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-oxo-2-(piperidin-1-yl)ethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	647
325		4-[3-{4-[2-(2,3-dihydro-1H-indol-1-yl)-2-oxoethoxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	681
326		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	558
327		4-[3-{4-[(3-chlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	646
328		4-[3-{[6-(3,5-dichlorophenyl)pyridin-3-yl]methyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	603
329		4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)biphenyl-2-carbonitrile	577








330		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(1H-pyrazol-1-yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	542
331		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-([6-(3-fluorophenyl)pyridin-3-yl]methyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	571
332		3-[5-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl)pyridin-2-yl]benzonitrile	578
333		4-[3-(3,4-diethoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	564
334		4-[3-{4-[(4-chlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	646
335		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	575
336		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-([6-(1H-pyrazol-1-yl)pyridin-3-yl]methyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	543
337		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-(morpholin-4-yl)benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	561









338		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-propoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	564
339		4-{3-[4-(1H-benzimidazol-1-yl)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	592
340		5-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxybenzonitrile	531
341		3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-7-carbonitrile	543
342		4-[3-(4-bromobenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	554
343		4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-(2-methoxyethoxy)benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	724
344		4-{3-[4-(benzyloxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	582
345		4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-ethoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	694

		dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	
346		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-hydroxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	492
347		4-[3-{4-[(3,4-dichlorobenzyl)oxy]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	650
348		4-{7-fluoro-3-[3-(2-fluoroethoxy)-4-hydroxybenzyl]-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	554
349		4-[3-{4-[(3,4-dichlorobenzyl)oxy]-3-(2-fluoroethoxy)benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	712
350		4-[3-{4-[(2-chloro-4-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	664
351		4-[3-{4-[(2,4-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
352		4-[3-{4-[(2-chloro-6-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	664








353		4-[3-{4-[(2,6-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
354		4-[3-{4-[(2-chlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	646
355		4-[7-fluoro-3-{4-[(2-fluorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	630
356		4-[3-{3-[(3,4-dichlorobenzyl)oxy]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	650
357		2-[{(3,4-dichlorobenzyl)oxy]-5-[(7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl]benzonitrile	675
358		4-[3-{4-[(3,4-dichlorophenoxy)methyl]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
359		4-{3-benzyl-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	476
360		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3-[4-(2-phenylethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	580







361		4-[3-(2-chloro-4,5-dimethoxybenzyl)-6-(2-fluoro-1-(fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	552
362		4-[3-{4-[(4,5-dichloro-2-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	698
363		4-[3-{3-[(3,4-dichlorobenzyl)oxy]-4-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
364		4-[3-(4-bromo-3-methoxybenzyl)-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	584
365		4-[3-{4-[(3,4-dichlorophenoxy)methyl]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	650
366		4-[3-{4-[(4-chlorophenoxy)methyl]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	616
367		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[3-methoxy-4-(2-phenylethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	610
368		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	506









		yl)piperidine-1-carbaldehyde	
369		4-[3-{4-[(4-chlorophenoxy)methyl]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	646
370		4-{3-[4-(benzyloxy)-3-chlorobenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	616
371		4-[3-{3-chloro-4-[(4-chlorobenzyl)oxy]-5-ethoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	694
372		4-[3-{4-[(2,4-dichlorobenzyl)oxy]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	650
373		4-[3-{3-chloro-4-[(2,4-dichlorobenzyl)oxy]-5-ethoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	728
374		4-{3-[4-(benzyloxy)-3-chloro-5-ethoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	660
375		4-[7-fluoro-3-{4-[(4-fluorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	630




376		4-[3-{4-[(3,5-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
377		4-[3-{4-[4-chloro-3-(trifluoromethyl)benzyl]oxy}-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	714
378		4-[3-{4-[(3-chlorophenoxy)methyl]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	616
379		4-[3-{4-[(3-chlorophenoxy)methyl]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	646
380		4-[3-{4-[(3,5-difluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	648
381		4-[3-{4-(benzyloxy)-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680
382		4-[3-{4-[(3-chloro-5-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	664
383		4-[7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(3-methoxy-4-{[4-(trifluoromethyl)benzyl]oxy}benzyl)-2,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	680

		dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	
384		4-[3-{4-[(2,5-dichlorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	680
385		4-{[4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]methyl}benzonitrile	637
386		3-{[4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]methyl}benzonitrile	637
387		4-[3-{4-[(4-chloro-2-fluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	664
388		4-[3-{4-[1-(3,4-dichlorophenyl)ethoxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	694
389		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-{4-[(3-hydroxybenzyl)oxy]-3-methoxybenzyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	696
390		4-[7-fluoro-3-{4-[(3-fluorobenzyl)oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	630

391		4-[3-{4-[(3,4-difluorobenzyl)oxy]-3-methoxybenzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	648
392		4-{3-[4-(5,6-dichloro-1H-benzimidazol-1-yl)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	660
393		4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenyl 3,4-dichlorobenzenesulphonate	700
394		4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenyl dichlorobenzenesulphonate	730
395		4-{3-[4-(3,4-dichlorophenoxy)-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	666
396		N-(3,4-dichlorophenyl)-4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)benzamide	645
397		N-(3,4-dichlorobenzyl)-4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)benzamide	659

398		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-(4-{[3-fluoro-4-(trifluoromethyl)benzyl]oxy}-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	698
399		4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenyl 1-methylethyl carbonate	608
400		4-{3-[4-(4-chlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	602
401		4-{3-[4-(3-chlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	602
402		4-[3-{[1-(3,4-dichlorobenzyl)-1H-pyrazol-4-yl]methyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	624
403		3,4-dichloro-N-[4-({7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenyl]benzamide	693
404		4-[3-{4-[(4-chlorobenzyl)oxy]benzyl}-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	616

405		4-{7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-3-[4-(4-fluorophenoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	586
406		4-{3-[4-(3,4-dichlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	636
407		4-{3-[4-(3,4-dichlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	651
408		4-{3-[4-(3,4-dichlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	616
409		4-{3-[4-(3,4-dichlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	624
410		4-{3-[4-(3,4-dichlorophenoxy)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	673
411		3-{4-[3,4-dichlorophenoxy]oxy]-3-methoxybenzyl}-6-[2-fluoro-1-(fluoromethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-7-carbonitrile	687
412		4-{3-[4-(5-chloro-6-fluoro-1H-benzimidazol-1-yl)benzyl]-7-fluoro-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidine-1-carbaldehyde	644

413		4-{3-[4-(5-chloro-1H-benzimidazol-1-yl)benzyl]-7-fluoro-6-[2-fluoro-1-((fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyde}	626
414		4-{3-[4-[(4-chloro-2-methoxybenzyl)oxy]-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-((fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyde}	676
415		4-{3-[4-[(3-chloro-4-hydroxybenzyl)oxy]-3-methoxybenzyl]-7-fluoro-6-[2-fluoro-1-((fluoromethyl)ethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyde}	730

The compounds of the invention form the subject of pharmacological trials which have shown their advantage as therapeutically active substances.

5 **1) Measurement of the inhibitory activity of the compounds according to the invention with regard to PDE7**

The ability of the compounds of formula (I) to inhibit PDE7 is measured using an enzymatic assay based on the separation of
10 radioactive cAMP (substrate of PDE7) from the radioactive 5'-AMP (product of the enzymatic reaction) by thin layer chromatography on polyethyleneimine (PEI) cellulose, after halting the enzymatic reaction. The 5'-AMP is extracted quantitatively from the PEI cellulose and its radioactivity is
15 measured using a liquid scintillation counter.

The inhibitory activity of the compounds of formula (I) with regard to PDE7 is represented by the inhibition constant IC_{50} , defined as the concentration of the compound (inhibitor)
20 tested in the assay which makes it possible to reduce the enzymatic activity of PDE7 by 50%. The lower the IC_{50} values, the greater the inhibitory power of the compounds.

Materials

25 The [3H]-cAMP (NET 275; 25 to 40 Ci/mmol) was purchased from Perkin-Elmer (NEN Life Sciences, Boston, United States), the rolipram was purchased from Sigma (St Louis, Mo, United States) and the sheets of polyethyleneimine cellulose F made
30 of plastic for thin layer chromatography were purchased from Merck (Darmstadt, Germany). All the other products used are of commercial origin.

Enzyme

35 Human PDE7 was partially purified from the HUT-78 cell line by following a method analogous to that described by Bloom and Beavo (Proc. Natl. Acad. Sci. USA, (1996) 93, 14188-14192).

The enzyme preparation obtained is stored at -80°C in a buffer comprising 20 mM of Tris-HCl (pH 7.0), 5 mM of MgCl_2 , 4 mM of EDTA, 1 mM of dithiothreitol and 20% of glycerol. As the partially purified PDE7 is contaminated by PDE4, it is necessary to add 10 μM of rolipram (selective inhibitor of PDE4) in the enzymatic assay in order to completely inhibit the PDE4 activity. The Michaelis constant (K_m) of PDE7 for cAMP, measured using the radiochemical assay described below, is 21 nM.

10

Solutions of compounds according to the invention

The compounds of formula (I) to be tested as PDE7 inhibitors are dissolved in DMSO at a concentration of 10 mM. These solutions are subsequently diluted in cascade in DMSO in order to obtain solutions with the desired concentrations. The latter are subsequently diluted to one-twentieth in the assay buffer to give 5% DMSO solutions. The latter are finally diluted to one-fifth in the enzymatic assay.

20

The solution of rolipram (added in the assay in order to completely inhibit the contaminating PDE4 activity) is prepared in an identical way and contributes 1% of DMSO to the enzymatic assay.

25

PDE7 Enzymatic assay

The assay is carried out in 1.5 ml Eppendorf tubes comprising 40 mM of Tris-HCl (pH 7.5), 15 mM of MgCl_2 , 1 mM of EGTA, 0.5 mg/ml of bovine serum albumin, 0.063 μCi of $[^3\text{H}]$ -cAMP (corresponding to a cAMP concentration of between 15 and 25 nM), 10 μM of rolipram and PDE7 in a final volume of 100 μl . The assay is carried out in the absence (control sample) or in the presence (treated sample) of the compounds tested as PDE7 inhibitors. The final concentration of DMSO in the assay is 2%. The reaction is initiated by addition of enzyme and the samples are maintained at ambient temperature for 30 minutes. The enzymatic dilution is adjusted so as to

35

obtain a degree of conversion of 10 to 15%. The enzymatic reaction is halted by immersion of the stoppered Eppendorf tubes in a water bath at 100°C for 3 minutes. Blanks (reaction halted immediately after addition of the enzyme) are included in each experiment. The samples are subsequently centrifuged at 10 000xg for 1 minute and an aliquot portion of 10 µl of supernatant is deposited 2 cm from the bottom edge of a sheet of PEI cellulose on which 10 µg of cAMP and 10 µg of 5'-AMP have been deposited beforehand. In order to facilitate the migration and to make it easier to cut out the strips of PEI cellulose comprising the 5'-AMP, which take place subsequently, 18 migration lanes with a width of 1 cm are delimited per plate by scraping the cellulose with a spatula over a width of 1 mm. The plates are developed over their entire length with a 0.30 M solution of LiCl in water by ascending chromatography. The 5'-AMP ($R_f = 0.20$) and the cAMP ($R_f = 0.47$) are visualized under UV light at 254 nm. The strips of PEI cellulose comprising the 5'-AMP are cut out and the nucleotide is quantitatively extracted in counting flasks with 2 ml of a solution which is 16M in formic acid and 2M in ammonium formate in water (rotary stirrer for 15 min). After addition of 10 ml of scintillation liquid (OptiPhase HiSafe 3 from Perkin-Elmer/Wallac), the radioactivity is counted using a liquid scintillation counter (model 1414, Perkin-Elmer/Wallac). Each assay is carried out in duplicate. The radioactivity specifically associated with the 5'-AMP formed in the enzymatic reaction is obtained by subtracting the mean value of blanks from the mean value of the controls (or treated samples).

30

The percentage of inhibition of PDE7 at a given concentration of the compound tested (inhibitor) is calculated using the equation : $I\% = [\text{mean value of the controls} - \text{mean value of the treated samples}] \times 100 / [\text{mean value of the controls} - \text{mean value of the blanks}]$.

35

The IC_{50} is the concentration of the compound (inhibitor) tested in the assay which makes it possible to reduce the

enzymatic activity of PDE7 by 50%.

Results

5 As illustrative and nonlimiting examples, the following quinazolinediones inhibit PDE7 with the IC₅₀ values indicated below :

Compound	IC ₅₀ (μM)
No. 16: 4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	0.015
No. 11: 4-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	0.039
No. 72: 4-[3-(3,4-dimethoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	0.089
No. 77: N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpyrrolidin-3-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamide	0.82
No. 97: 4-[5,7-dichloro-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	0.061
No. 111: 4-[6-(difluoromethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	0.0067

10 **2) Measurement of the inhibitory activity of the compounds according to the invention with regard to PDE8**

By using for PDE8 an enzymatic assay equivalent to that described for PDE7, the following IC₅₀ values are obtained :

15

Compound	IC ₅₀ PDE7 (μM)	IC ₅₀ PDE8 (μM)
No. 11	0.039	0.14
No. 251	0.046	0.39

No. 294	0.037	0.015
---------	-------	-------

3) Inhibition of T cell proliferation

Isolation of peripheral blood mononuclear cells (=PBMNCs)

5

Human peripheral blood mononuclear cells (PBMNCs) were separated from the blood of healthy donors using a centrifuging gradient on a Ficoll-Paque™ Plus (GE Healthcare Bio-Sciences AB code 17-1440-03, Uppsala). A mixture of 5.0 ml of blood and 5.0 ml of phosphate buffered saline (PBS Gibco 14190) was deposited on 10.0 ml of Ficoll-Paque in a 50 ml Blue max tube (Becton Dickinson 352070) and centrifuged with 900 g for 30 min. The band of PBMNCs was withdrawn, washed in PBS and then centrifuged with 900 g for 10 minutes. Finally, the PBMNCs were suspended (2.5×10^5 cells/ml) in an RPMI 1640 complete medium + Glutamax (Gibco 61870) additivated with 10% of FCS, the complement of which has been heat inactivated, and with penicillin-streptomycin (Gibco 15140).

20 Measurement of the proliferation by incorporation of [³H]-thymidine.

The PBMNCs were cultured in flat-bottomed 96-well plates, in a proportion of 0.5×10^5 cells/well in 200 µl of RPMI 1640 complete medium. The cells were activated by two monoclonal antibodies, one of anti-CD3 specificity, 20 ng/ml (Pharmingen 555329), and the other of anti-CD28 specificity, 200 ng/ml (Pharmingen 555725), in the presence of PDE7 inhibitors at five different concentrations, 10 µM, 3.3 µM, 1.1 µM, 0.36 µM and 0.12 µM, or of cyclosporin A, which acts as positive control of the halting of proliferation. The two monoclonal antibodies employed simultaneously bring about an increased intensity of the transcriptional response of the T cells activated via CD3 alone (Diehn M. et al., 2002. Genomic expression programs and the integration of the CD28 costimulatory signal in T cell activation. Proc. Natl. Acad. Sci. USA, 99, 11796-11801). Furthermore, the two antibodies

employed mimic the optimum condition of the in vivo activation, which employs the TCR and the CD28 coreceptor. After 72 hours at 37°C in 5% of CO₂ and 90% of humidified air, the cultures were labelled with 1 µCi [methyl-³H]-thymidine (Amersham TRK120). After incubating for an additional 8 hours, the PBMNCs were collected on a Tomtec cell collector (Wallac) using a glass fibre filter (Wallac 1450-421), and incorporation of the [³H]-thymidine in the nucleus of the cells was measured using a Microbeta 1450-Trilux (Wallac). The inhibition of the proliferation, expressed as percentage, was calculated using the following equation : $100 - (\text{cpm mean value of the cells in the presence of the compound} / \text{cpm mean value of the cells in the absence of the compound}) \times 100$.

15 Results

The inhibitory activities of the compounds of formula (I) according to the invention for the proliferation correspond to IC₅₀ values (concentration of the inhibitor which brings about 50% reduction in the incorporation of [³H]-thymidine in the nuclei of the cells). The lower the IC₅₀ values, the more the compounds are powerful inhibitors of the proliferation of T lymphocytes.

25 The following quinazolin-2(1H)-ones inhibit the proliferation of PBMNCs with IC₅₀ values indicated below (mean of the three experiments) :

Compound	IC ₅₀ (µM)
No. 1: 2-([3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy)propanenitrile	0.4 µM
No. 11: 4-[3-(3,4-dimethoxybenzyl)-6-[2-fluoro-1-(fluoromethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidine-1-carbaldehyde	0.9 µM
No. 16: 4-[6-(2,2-difluoroethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-	1.3 µM

1 (2H)-yl]piperidine-1-carbaldehyde	
-------------------------------------	--

The compounds according to the invention exhibit good pharmacological properties and are particularly capable of being used in the preparation of medicaments, in particular of
5 medicaments which are inhibitors of PDE7 or, for some of the compounds of the invention, of medicaments which are inhibitors of PDE7 and of PDE8.

According to another of its aspects, the subject matter of the
10 invention is thus medicaments which comprise at least one compound of formula (I).

The compounds according to the invention can be of use in particular in the treatment and/or prevention of inflammatory
15 or immunoinflammatory diseases including, for example, asthma, chronic obstructive pulmonary disease (COPD), allergic rhinitis, allergies, Crohn's disease, ulcerative colitis, myasthenia gravis, atopic dermatitis, psoriasis, lupus erythematosus disseminatus, rheumatoid arthritis, diabetes or
20 multiple sclerosis, in the treatment of organ transplants, in the treatment and/or prevention of certain types of cancers, such as osteosarcoma or adenocarcinoma, in the treatment and/or prevention of bone diseases, such as osteopenia or osteoporosis, in the treatment and/or prevention of acute
25 insufficiency, or in the treatment and/or prevention of pain, in particular neuropathic pain and visceral pain.

The use of the compounds of general formula (I) in the preparation of a medicament intended to treat the
30 abovementioned pathologies forms an integral part of the invention, and in particular compounds of formula (I) chosen from compounds Nos. 1 to 6, 11 to 14, 16, 20, 22 to 25, 32 to 43, 47 to 52, 55 to 59, 72, 74, 76, 78, 79, 89 to 91, 97, 102, 108, 111, 112, 114, 116 to 118, 124, 130, 131, 133 to 135,
35 143, 145, 155, 158, 160, 165 to 167, 170, 175, 178, 183 to 186, 188, 189, 190, 193, 194, 200, 201, 203, 206, 207, 212, 213, 215, 216, 218, 223, 224, 226, 228, 230, 232 to 234, 239,

240, 242, 243, 245, 246, 250, 251, 254, 258, 263, 264, 270, 275, 276, 278 to 280, 282, 283, 285 to 287, 289, 292, 294, 295, 287 to 302, 305 to 312, 315 to 345, 349 to 355, 357, 358, 360, 362, 369, 371, 373, 375 to 377, 379 to 394 and 403.

5

According to another of its aspects, the present invention relates to pharmaceutical compositions comprising, as active principle, at least one compound according to the invention. These pharmaceutical compositions comprise an effective dose
10 of at least one compound of formula (I) according to the invention and also at least one pharmaceutically acceptable excipient.

The said excipients are chosen, according to the
15 pharmaceutical form and the method of administration desired, from the usual excipients which are known to a person skilled in the art.

In the pharmaceutical compositions of the present invention
20 for oral, sublingual, subcutaneous, intramuscular, intravenous, topical, local, intratracheal, intranasal, transdermal or rectal administration, the active principle of formula (I) above or its optional salt, solvate or hydrate can be administered in unit administration form, as a mixture with
25 conventional pharmaceutical excipients, to animals and human beings for the prophylaxis or treatment of the above disorders or diseases.


The appropriate unit administration forms comprise forms by
30 the oral route, such as tablets, soft or hard gelatin capsules, powders, granules and oral solutions or suspensions, sublingual, buccal, intratracheal, intraocular or intranasal administration forms, forms for administration by inhalation, topical, transdermal, subcutaneous, intramuscular or
35 intravenous administration forms, rectal administration forms and implants. Use may be made, for topical application, of the compounds according to the invention in creams, gels, ointments or lotions.

By way of example, a unit administration form of a compound according to the invention in the form of a tablet can comprise the following components :

- 5
- | | | | | |
|----------|------------------------------|---|---|-----------|
| Compound | No. | 1 | (2-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propanenitrile) | 50.0 mg |
| Mannitol | | | | 223.75 mg |
| 10 | Croscarmellose sodium | | | 6.0 mg |
| | Maize starch | | | 15.0 mg |
| | Hydroxypropylmethylcellulose | | | 2.25 mg |
| | Magnesium stearate | | | 3.0 mg |
- 15 The said unit forms comprise doses in order to make possible daily administration of 0.5 mg to 800 mg of active principle per individual, more particularly of 0.5 mg to 200 mg, according to the pharmaceutical dosage form.
- 20 There may be cases where higher or lower dosages are appropriate ; such dosages do not depart from the scope of the invention. According to the usual practice, the dosage appropriate to each patient is determined by the physician according to the method of administration and the weight and
- 25 response of the said patient.

Patentkrav

1. Forbindelse med den almene formel (I)



5 hvor

- A betegner en arylgruppe eller en heteroarylgruppe;

- R₁ betegner:

▪ et hydrogenatom,

▪ -C(O)R, hvor R er et hydrogenatom, en (C₁-C₆)-

10 alkoxygruppe, en arylgruppe, en (C₃-C₆)-cycloalkylgruppe
 eller en (C₁-C₆)-alkylgruppe, hvilken alkylgruppe eventuelt
 er substitueret med:

. en eller flere hydroxygrupper,

. en benzyloxygruppe,

15 . en (C₁-C₆)-alkoxygruppe, der eventuelt er substitueret med
 en arylgruppe, eller

. en (C₃-C₆)-cycloalkylgruppe,▪ en (C₁-C₆)-alkylgruppe, der eventuelt er substitueret;- R₂ betegner:

20 ▪ et hydrogenatom,

▪ et halogenatom,

▪ en cyanogruppe,

▪ en nitrogruppe,

▪ en (C₁-C₆)-alkylgruppe, der eventuelt er substitueret med

25 en -NH₂-gruppe eller en -NHC(O)R_b-gruppe,

▪ en -OR_a-gruppe, hvor R_a betegner:

. et hydrogenatom,


. en (C₁-C₆)-alkylgruppe, der eventuelt er substitueret med
 et eller flere halogenatomer, en eller flere

30 hydroxygrupper, en arylgruppe og/eller en eller flere
 cyanogruupper,

- . en (C₂-C₆)-alkynylgruppe,
- . en arylgruppe;
- R₃ betegner:
 - et hydrogenatom,
 - 5 ▪ et halogenatom,
 - en hydroxygruppe,
 - en cyanogruppe,
 - en -SCF₃-gruppe,
 - en nitrogruppe,
 - 10 ▪ en oxogruppe,
 - en -S(O)₀₋₂alkylgruppe, en -S(O)₀₋₂-heterocycloalkylgruppe, en -O-SO₂-arylgruppe, der eventuelt er substitueret med et eller flere halogenatomer;
 - en -alkyl-amino-alkyl- eller -cycloalkyl-amino-
 - 15 alkylgruppe, der hver eventuelt er substitueret på den terminale alkylgruppe,
 - en sulfonamidgruppe, der eventuelt er substitueret,
 - en arylgruppe eller en heteroarylgruppe, hvilken gruppe er monocyklisk eller polycyklisk og desuden eventuelt er
 - 20 substitueret med en (C₁-C₈)-alkylgruppe, et eller flere halogenatomer eller en (C₁-C₆)-alkoxygruppe,
 - en heterocycloalkylgruppe, der eventuelt er substitueret med en (C₁-C₆)-alkylgruppe,
 - en (C₁-C₆)-alkylgruppe, der eventuelt er substitueret med:
 - 25 - et eller flere halogenatomer,
 - en arylgruppe, der kan være substitueret med et eller flere halogenatomer eller med en eller flere hydroxygrupper,
 - en heteroarylgruppe,
 - 30 - en eller flere hydroxygrupper, der kan være substitueret med en arylgruppe, som igen eventuelt er substitueret med et eller flere halogenatomer, eller
 - en heterocycloalkylgruppe, der eventuelt er substitueret med en CO(O)R_a-gruppe eller en (C₁-C₆)-alkylgruppe,
 - 35 ▪ en -C(O)NR_bR_c-gruppe,
 - en -C(O)OR_c-gruppe eller en -O-C(O)OR_c-gruppe,
 - en (C₁-C₆)-alkoxygruppe, der eventuelt er substitueret med:

- en amino-alkylgruppe,
- en amino-cycloalkylgruppe,
- en cycloalkylgruppe,
- en heterocycloalkylgruppe,
- 5 - en monocyklisk eller polycyklisk heteroarylgruppe,
- en eller flere hydroxygrupper,
- et eller flere halogenatomer,
- en (C₁-C₆)-alkoxygruppe,
- en -C(O)ORc-gruppe,
- 10 - en -C(O)NRbRc-gruppe,
- en oxogruppe og/eller
- en arylgruppe, der igen eventuelt er substitueret med et eller flere halogenatomer, en cyanogruppe, en (C₁-C₆)-alkoxygruppe, en -O-halogenalkylgruppe og/eller en
- 15 halogenalkylgruppe,
 - en -O-cycloalkylgruppe, en -O-arylgruppe eller en -O-heterocycloalkylgruppe, der hver eventuelt er substitueret med
 - en arylgruppe, der igen eventuelt er substitueret med et
 - 20 eller flere halogenatomer eller en (C₁-C₆)-alkylgruppe,
 - en oxogruppe,
 - et eller flere halogenatomer, og/eller
 - en (C₁-C₆)-alkylgruppe, der igen kan være substitueret med en arylgruppe og/eller en oxogruppe,
 - 25 ▪ en -NH-CO-NH-arylgruppe, en -NH-CO-NH-heteroarylgruppe eller en -NH-CO-NH-(C₁-C₆)-alkylgruppe, der hver eventuelt er substitueret med et eller flere halogenatomer, en cyanogruppe, en nitrogruppe, en eller flere hydroxygrupper eller en (C₁-C₆)-alkoxygruppe,
 - 30 ▪ en -N-(C₁-C₆)-alkylgruppe, hvor (C₁-C₆)-alkylgruppen kan være substitueret med
 - en eller flere oxogrupper og/eller
 - en eller flere arylgrupper, der eventuelt er substitueret med et eller flere halogenatomer og/eller en SO₂-gruppe,
 - 35 ▪ en -NH-CO-arylgruppe, en -NH-CO-heteroarylgruppe, der hver eventuelt er substitueret med et eller flere halogenatomer;

- eller R_3 danner sammen med A en polycyklisk heteroarylgruppe, der eventuelt er substitueret med en (C_1-C_6) -alkoxygruppe, en (C_1-C_6) -alkylgruppe, som eventuelt er substitueret med en arylgruppe, der igen kan være
- 5 substitueret med et eller flere halogenatomer;
- R_4 betegner et hydrogenatom, en oxogruppe eller en (C_1-C_6) alkylgruppe;
 - Rb betegner:
 - . et hydrogenatom,
- 10 . en (C_1-C_6) -alkylgruppe, der eventuelt er substitueret med et eller flere halogenatomer, en eller flere hydroxy-, cyano-, amino-, heterocycloalkyl-, (C_1-C_6) -alkoxygrupper eller en arylgruppe, der eventuelt er substitueret med et eller flere halogenatomer,
- 15 . en (C_3-C_6) -cycloalkylgruppe,
- . en (C_2-C_6) -alkynylgruppe,
 - . en (C_1-C_6) -alkoxygruppe,
 - . en arylgruppe, der eventuelt er substitueret med et eller flere halogenatomer;
- 20 - R_c betegner et hydrogenatom eller en (C_1-C_6) -alkylgruppe, der eventuelt er substitueret med et eller flere halogenatomer;
- eller Rb og R_c danner sammen med det nitrogenatom, de er bundet til, en polycyklisk heteroarylgruppe eller en
- 25 heterocycloalkylgruppe;
- m og n betegner uafhængigt af hinanden værdien 0, 1 eller 2, idet det forudsættes, at $m+n \leq 3$;
 - p og p' betegner uafhængigt af hinanden værdien 1, 2 eller 3, idet det forudsættes, at når p er større end eller
- 30 lig med 2, så sidder grupperne R_2 på adskilte carbonatomer og kan være forskellige fra hinanden, og når p' er større end eller lig med 2, så sidder grupperne R_3 på adskilte carbonatomer og kan være forskellige fra hinanden;
- q betegner værdien 0 eller 2, idet det forudsættes, at
- 35 når q = 0, så er den heterocykliske nitrogenholdige gruppe, der er bundet til nitrogenatomet i position 1 på 2,4-dioxo-1,2,3,4-tetrahydroquinazolin-kernen, ikke længere broforbundet, og den er af typen:



- r betegner værdien 0 eller 1.

2. Forbindelse med den almene formel (I) ifølge krav 1, kendetegnet ved, at A betegner en phenylgruppe eller en pyridylgruppe.

3. Forbindelse med den almene formel (I) ifølge krav 1 eller 2, kendetegnet ved, at $q = 0$, m og $n = 1$.

10

4. Forbindelse med den almene formel (I) ifølge et hvilket som helst af kravene 1 til 3, kendetegnet ved, at R_2 betegner en (C_1-C_6) -alkylgruppe, især en methylgruppe, der er substitueret med en $-NH-CO-R_b$ -gruppe, hvor R_b er som defineret i krav 1.

5. Forbindelse med den almene formel (I) ifølge et hvilket som helst af kravene 1 til 3, kendetegnet ved, at R_2 betegner en $-OR_a$ -gruppe, hvor R_a er som defineret i krav 1.

20

6. Forbindelse med den almene formel (I) ifølge et hvilket som helst af kravene 1 til 3, kendetegnet ved, at R_2 er et halogenatom eller en cyanogruppe eller et hydrogenatom eller en hydroxygruppe eller en (C_1-C_6) -alkylgruppe, der eventuelt er substitueret med en $-NH_2$ -gruppe eller en $-NHC(O)R_b$ -gruppe.

7. Forbindelse med den almene formel (I) ifølge krav 1, kendetegnet ved, at A er en phenylgruppe, R_1 er en $-C(O)R$ -gruppe, hvor R betegner et hydrogenatom, q er lig med 0, n og m har værdien 1, og R_2 er $-OR_a$.

8. Forbindelse med den almene formel (I) ifølge krav 1, kendetegnet ved, at A er en phenylgruppe, R_1 er en $-C(O)R$ -gruppe, hvor R betegner et hydrogenatom, q er lig med 0, n

35

og m har værdien 1, og R_2 er en methylgruppe, der er substitueret med en $-NH-CO-R_b$ -gruppe, hvor R_b er som defineret i krav 1.

5 9. Forbindelse med den almene formel (I) ifølge krav 1, kendetegnet ved, at A er en phenylgruppe, R_1 er en $-C(O)R$ -gruppe, hvor R betegner et hydrogenatom, q er lig med 0, n og m har hver værdien 1, p er lig med 2, en af grupperne R_2 er $-OR_a$, hvor R_a er som defineret i den almene formel (I),
10 og den anden af grupperne R_2 er et halogenatom.

10. Forbindelse med den almene formel (I) ifølge et hvilket som helst af kravene 1 til 9, kendetegnet ved, at gruppen R_2 er i position 6 på 2,4-dioxo-1,2,3,4-
15 tetrahydroquinazolinkernen, og ved, at der desuden kan være en gruppe R_2 , som kan være identisk eller forskellig, i position 7 på 2,4-dioxo-1,2,3,4-tetrahydroquinazolinkernen; i form af en base, et hydrat, et solvat, isomerer eller blandinger heraf.

20

11. Forbindelse med den almene formel (I) ifølge krav 1, der er valgt blandt følgende forbindelser:

- forbindelse nr. 1: 2-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-
25 tetrahydroquinazolin-6-yl]oxy}propannitril
- forbindelse nr. 2: 1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-hydroxyquinazolin-2,4(1H,3H)-dion
- forbindelse nr. 3: {[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-
30 yl]oxy}acetonitril
- forbindelse nr. 4: {[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]oxy}propannitril
- forbindelse nr. 6: 2-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-
35 tetrahydroquinazolin-6-yl]oxy}acetonitril

- forbindelse nr. 11: 4-[3-(3,4-dimethoxybenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 12: 1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]quinazolin-2,4(1H,3H)-dion
- forbindelse nr. 13: 4-[3-(3,4-dimethoxybenzyl)-2,4-dioxo-6-(2,2,2-trifluorethoxy)-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 10 - forbindelse nr. 14: 1-(1-acetylpiperidin-4-yl)-6-(2,2-difluorethoxy)-3-(3,4-dimethoxybenzyl)quinazolin-2,4(1H,3H)-dion
- forbindelse nr. 16: 4-[6-(2,2-difluorethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 15 - forbindelse nr. 20: N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamid
- forbindelse nr. 22: 1-(1-acetylpiperidin-4-yl)-6-(aminomethyl)-3-(3,4-dimethoxybenzyl)quinazolin-2,4(1H,3H)-dionchlorhydrat
- 20 - forbindelse nr. 23: N-{[3-(3,4-dimethoxybenzyl)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}formamid
- 25 - forbindelse nr. 24: {[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}formamid
- forbindelse nr. 25: N-{[1-(1-acetylpiperidin-4-yl)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-6-yl]methyl}acetamid
- 30 - forbindelse nr. 32: 4-[6-(2,2-difluorethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 33: 4-[3-(3,4-dichlorbenzyl)-6-(2,2-difluorethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 35 - forbindelse nr. 34: 4-[3-(4-chlorbenzyl)-6-(2,2-difluorethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 35: methyl-4-{[6-(2,2-difluorethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl}benzoat
- forbindelse nr. 36: 4-{[6-(2,2-difluorethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl}benzoesyre
- forbindelse nr. 37: 4-{[6-(2,2-difluorethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl}-N-(2-methoxyethyl)benzamid
- 10 - forbindelse nr. 38: 4-[3-(3,4-dimethoxybenzyl)-6-methyl-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 39: 4-[6-(2,2-difluorethoxy)-3-(3-hydroxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1
- 15 (2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 40: 4-[6-(2,2-difluorethoxy)-3-[3-(2-hydroxyethoxy)-4-methoxybenzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 41: 4-[6-(2,2-difluorethoxy)-3-(3-ethoxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1 (2H)-yl]piperidin-1-carbaldehyd
- 20 - forbindelse nr. 42: 4-[6-(2,2-difluorethoxy)-3-[4-methoxy-3-(2-methoxyethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 25 - forbindelse nr. 43: 4-[6-(2,2-difluorethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]azepan-1-carbaldehyd
- forbindelse nr. 47: 4-[6-(2,2-difluorethoxy)-3-[3-(3-hydroxypropoxy)-4-methoxybenzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 30 - forbindelse nr. 48: 4-[5-chlor-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 49: 4-{3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(2,2-difluorethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 35

- forbindelse nr. 50: 2-(5-{[6-(2,2-difluorethoxy)-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl]methyl}-2-methoxyphenoxy)acetamid
- forbindelse nr. 51: 4-[6-(2,2-difluorethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]-3-methylpiperidin-1-carbaldehyd
- forbindelse nr. 52: 3-[6-(2,2-difluorethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]-8-azabicyclo[3.2.1]octan-8-carbaldehyd
- 10 - forbindelse nr. 56: 4-{3-[4-(cyclopentyloxy)-3-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 57: 4-[3-(3-chlorbenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 15 - forbindelse nr. 58: 4-[3-(4-chlorbenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 59: 4-{3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 20 - forbindelse nr. 72: 4-[3-(3,4-dimethoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 74: 4-[3-(3,4-dichlorbenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 76: 4-{3-[(6-chlorpyridin-3-yl)methyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 30 - forbindelse nr. 78: 4-{3-[3-chlor-4-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 79: 4-[3-(3,4-dimethoxybenzyl)-6-(2-fluorethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 35 - forbindelse nr. 89: 2-[5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-

- 1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]acetamid
- forbindelse nr. 90: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-hydroxy-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 5 - forbindelse nr. 91: 4-[3-(3,4-dimethoxybenzyl)-6-ethoxy-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 97: 4-[5,7-dichlor-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 10 - forbindelse nr. 102: 4-[7-chlor-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 15 - forbindelse nr. 108: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-fluor-4-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 111: 4-[6-(difluormethoxy)-3-(3,4-dimethoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 20 - forbindelse nr. 112: 4-[3-(3,4-dimethoxybenzyl)-6-(1-methylethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 114: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-methoxy-3-(1-methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 25 - forbindelse nr. 116: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 30 - forbindelse nr. 117: 4-{3-[3,5-bis(trifluormethyl)benzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 35 - forbindelse nr. 118: 4-[3-(3-ethoxybenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 124: 4-{3-[3-chlor-4-(2-methoxyethoxy)benzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 5 nr. 130: 4-[3-(3,4-diethoxybenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- nr. 131: 4-{3-[4-ethoxy-3-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 10 - forbindelse nr. 133: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(4-methoxy-3-methylbenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 15 - forbindelse nr. 134: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[4-(trifluormethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 135: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[4-(trifluormethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 20 - forbindelse nr. 143: 4-{3-[4-(benzyloxy)-3-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 25 - forbindelse nr. 145: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-methoxy-4-nitrobenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 155: 4-[3-(4-ethoxybenzyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 30 - forbindelse nr. 158: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(morpholin-4-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 35 - forbindelse nr. 160: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(morpholin-4-ylbenzyl)benzyl]-

2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 165: 4-[3-(biphenyl-4-ylmethyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-

5 dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 166: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(methylsulfanyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

10 - forbindelse nr. 167: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-pyridin-3-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 170: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-methoxy-4-methylbenzyl)-2,4-

15 dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 175: 2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-

20 yl)methyl}phenoxy]acetamid

- forbindelse nr. 178: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-methoxy-4-propoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

25 - forbindelse nr. 183: 2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]-N-methylacetamid

- forbindelse nr. 184: 2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]-N,N-dimethylacetamid

30 - forbindelse nr. 185: 2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-

35 2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl}phenoxy]-N-methoxy-N-methylacetamid

- forbindelse nr. 186: 4-{3-[4-(cyclopentyloxy)-3-ethoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 188: 4-{3-[4-(cyclopentyloxy)-3-(1-methylethoxy)benzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 189: 4-{3-[4-(cyclopentyloxy)-3-propoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 190: 4-{3-[4-(cyclopentyloxy)-3-hydroxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 193: 4-{3-[4-(difluormethoxy)-3-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 194: 4-{3-[4-(difluormethoxy)-3-ethoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 200: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-thiophen-3-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 201: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-pyridin-4-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 203: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(1-methyl-1H-indol-6-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 206: 4-{3-[4-(cyclopropylmethoxy)-3-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 207: 2-[4-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]-N-methylacetamid
- forbindelse nr. 212: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[4-(1H-pyrazol-1-

yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 213: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-pyridin-2-ylbenzyl)-

5 3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 215: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-thiophen-2-ylbenzyl)-

3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 216: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(quinolein-7-ylmethyl)-

10 3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 218: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(6-methoxynaphthalen-2-yl)methyl]-

2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-

15 carbaldehyd

- forbindelse nr. 223: 4-{3-[4-(1H-benzimidazol-1-yl)benzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-

dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 224: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(2-

20 methylpropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 226: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(tetrahydrofuran-3-

25 yloxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 228: 4-[3-{4-[(1-benzylpyrrolidin-3-yl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-

2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-

30 carbaldehyd

- forbindelse nr. 230: 4-[3-(1-benzothiophen-5-ylmethyl)-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-

dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 232: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(1-

35 methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 233: 4-[3-(3,4-dimethoxybenzyl)-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 234: 4-[3-{4-[(1-acetylpyrrolidin-3-yl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 239: 4-[3-{4-[(fluorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 240: 4-[3-{4-[(chlorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 242: 4-[3-{4-[(3-chlorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 243: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(3-thiophen-3-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 245: 4-[3-(4-ethoxy-3-methoxybenzyl)-6-(2-hydroxyethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 246: 4-[3-{4-[2-(2,3-dihydro-1H-indol-1-yl)-2-oxoethoxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 250: 4-[3-{4-[(3,4-dichlorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 251: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(2-oxo-2-piperidin-1-ylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 254: 4-{3-[3-ethoxy-4-(thiophen-2-ylmethoxy)benzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 258: 4-[3-(3,4-dimethoxybenzyl)-6-[2-fluor-1-(hydroxymethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 263: (2R)-2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]propansyre
- forbindelse nr. 264: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(1-methyl-3-thiophen-2-yl-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 270: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 275: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-pyrimidin-5-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 276: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(1-methyl-3-phenyl-2-yl-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 278: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-{[6-(1H-pyrazol-1-yl)pyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 279: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[(2-thiophen-2-ylpyrimidin-5-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 280: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 282: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(3-methyl-1,2,4-oxadiazol-5-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 283: 2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]eddikesyre
- 5 - forbindelse nr. 285: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(thieno-[2,3-b]-pyridin-2-ylmethyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 286: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[(6-phenylpyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 10 - forbindelse nr. 287: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(6-morpholin-4-ylpyridin-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 15 - forbindelse nr. 289: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[(6-thiophen-2-ylpyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 20 - forbindelse nr. 292: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(1-methyl-5-phenyl-2-yl-1H-pyrazol-3-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 25 - forbindelse nr. 294: 4-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)biphenyl-2-carbonitril
- forbindelse nr. 295: (2R)-2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-methylpropanamid
- 30 - forbindelse nr. 297: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-thiophen-2-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 35 - forbindelse nr. 298: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(morpholin-4-

- ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 - forbindelse nr. 299: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(piperidin-1-ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 5 ylmethyl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 - forbindelse nr. 300: 4-[3-{4-[(3,4-dichlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 10 carbaldehyd
 - forbindelse nr. 301: 2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]-N-ethylacetamid
 15 - forbindelse nr. 302: (2S)-2-[2-(cyclopentyloxy)-5-({6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenoxy]propansyre
 - forbindelse nr. 305: 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-methoxy-4-{[(3R)-2-oxo-1-phenylpyrrolidin-3-yl]oxy}benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 20 (fluormethyl)ethoxy]-3-(3-methoxy-4-{[(3R)-2-oxo-1-phenylpyrrolidin-3-yl]oxy}benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 - forbindelse nr. 306: 4-{3-[4-(cyclobutylmethoxy)-3-methoxybenzyl]-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 25 3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 - forbindelse nr. 307: 4-{3-[4-(benzyloxy)-3-methoxybenzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 30 - forbindelse nr. 308: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-(4-hydroxy-3-methoxybenzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 - forbindelse nr. 309: 4-{3-[4-(cyclopropylmethoxy)-3-methoxybenzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
 35 methoxybenzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 310: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(2-methylpropoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 5 - forbindelse nr. 311: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(1-methylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 312: 4-[3-(4-ethoxy-3-methoxybenzyl)-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 10 - forbindelse nr. 315: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-{[6-(3-methoxyphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 15 - forbindelse nr. 316: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-{[6-(2-fluorphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 20 - forbindelse nr. 317: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-{[6-(4-fluorphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 318: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-{[6-(4-methoxyphenyl)pyridin-3-yl]methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 25 - forbindelse nr. 319: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[(6-thiophen-2-yl)pyridin-3-yl]methyl}-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 30 - forbindelse nr. 320: 4-{3-[3-ethoxy-4-(thiophen-2-ylmethoxy)benzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 35 - forbindelse nr. 321: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(1-methyl-1H-pyrazol-3-

- yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 322: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-(4-pyrimidin-5-ylbenzyl)-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 5 - forbindelse nr. 323: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[(1-methyl-3-thiophen-2-yl-1H-pyrazol-5-yl)methyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 10 - forbindelse nr. 324: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[3-methoxy-4-(2-oxo-2-piperidin-1-ylethoxy)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 325: 4-[3-{4-[2-(2,3-dihydro-1H-indol-1-yl)-2-oxoethoxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 15 - forbindelse nr. 326: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[4-(5-methyl-1,2,4-oxadiazol-3-yl)benzyl]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 20 - forbindelse nr. 327: 4-[3-{4-[(3-chlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 25 - forbindelse nr. 328: 4-[3-{[6-(3,5-dichlorphenyl)pyridin-3-yl]methyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 329: 4-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)biphenyl-2-carbonitril
- 30 - forbindelse nr. 330: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[4-(1H-pyrazol-1-yl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 35 - forbindelse nr. 331: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-[6-(3-fluorphenyl)pyridin-3-

yl)methyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-
yl}piperidin-1-carbaldehyd

- forbindelse nr. 332: 3-[5-({7-fluor-6-[2-fluor-1-
(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-
5 1,4-dihydroquinazolin-3(2H)-yl)methyl}pyrimidin-2-
yl]benzonitril

- forbindelse nr. 333: 4-[3-(3,4-diethoxybenzyl)-7-fluor-6-
[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-
dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

10 - forbindelse nr. 334: 4-[3-{4-[(4-chlorbenzyl)oxy]-3-
methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-
2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-
carbaldehyd

- forbindelse nr. 335: 4-{7-fluor-6-[2-fluor-1-
15 (fluormethyl)ethoxy]-3-[4-(morpholin-4-ylmethyl)benzyl]-
2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-
carbaldehyd

- forbindelse nr. 336: 4-{7-fluor-6-[2-fluor-1-
(fluormethyl)ethoxy]-2,4-dioxo-3-{[6-(1H-pyrazol-1-
20 yl)pyridin-3-yl)methyl]-3,4-dihydroquinazolin-1(2H)-
yl}piperidin-1-carbaldehyd

- forbindelse nr. 337: 4-{7-fluor-6-[2-fluor-1-
(fluormethyl)ethoxy]-3-[4-(morpholin-4-ylbenzyl)benzyl]-
2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-
25 carbaldehyd

- forbindelse nr. 338: 4-{7-fluor-6-[2-fluor-1-
(fluormethyl)ethoxy]-3-(3-methoxy-4-propoxybenzyl)-2,4-
dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-
carbaldehyd

30 - forbindelse nr. 339: 4-{3-[4-(1H-benzimidazol-1-yl)benzyl]-
7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-
dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 340: 5-({7-fluor-6-[2-fluor-1-
(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-

35 1,4-dihydroquinazolin-3(2H)-yl)methyl)-2-methoxybenzonitril

- forbindelse nr. 341: 3-(3,4-dimethoxybenzyl)-6-[2-fluor-
1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-
dioxo-1,2,3,4-tetrahydroquinazolin-7-carbonitril

- forbindelse nr. 342: 4-[3-(4-brombenzyl)-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 343: 4-[3-{4-[(3,4-dichlorbenzyl)oxy]-3-(2-methoxyethoxy)benzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 344: 4-{3-[4-(benzyloxy)benzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- forbindelse nr. 345: 4-[3-{4-[(3,4-dichlorbenzyl)oxy]-3-ethoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 349: 4-[3-{4-[(3,4-dichlorbenzyl)oxy]-3-(2-fluorethoxy)benzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 350: 4-[3-{4-[(2-chlor-4-fluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 351: 4-[3-{4-[(2,4-dichlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 352: 4-[3-{4-[(2-chlor-6-fluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 353: 4-[3-{4-[(2,6-dichlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 354: 4-[3-{4-[(2-chlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 355: 4-[7-fluor-3-{4-[(2-fluorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 5 - forbindelse nr. 357: 2-[(3,4-dichlorbenzyl)oxy]-5-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl)methyl)benzonitril
- forbindelse nr. 358: 4-[3-{4-[(3,4-dichlorphenoxy)methyl]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 10 - forbindelse nr. 360: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3-[4-(2-phenylethyl)benzyl]-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 15 - forbindelse nr. 362: 4-[3-{4-[(4,5-dichlor-2-fluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 20 - forbindelse nr. 369: 4-[3-{4-[(4-chlorphenoxy)methyl]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 371: 4-[3-{3-chlor-4-[(4-chlorbenzyl)oxy]-5-ethoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 25 - forbindelse nr. 373: 4-[3-{3-chlor-4-[(2,4-dichlorbenzyl)oxy]-5-ethoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 30 - forbindelse nr. 375: 4-[7-fluor-3-{4-[(4-fluorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 35 - forbindelse nr. 376: 4-[3-{4-[(3,5-dichlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-

2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 377: 4-[3-(4-{(4-chlor-3-(trifluormethyl)benzyl]oxy}-3-methoxybenzyl)-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 379: 4-[3-{4-[(3-chlorphenoxy)methyl]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-

10 carbaldehyd

- forbindelse nr. 380: 4-[3-{4-[(3,5-difluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

15 - forbindelse nr. 381: 4-{3-[4-(benzyloxy)-3-methoxybenzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd

- forbindelse nr. 382: 4-[3-{4-[(3-chlor-5-fluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 383: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-(3-methoxy-4-{[4-

25 (trifluormethyl)benzyl]oxy}benzyl)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd

- forbindelse nr. 384: 4-[3-{4-[(2,5-dichlorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-


30 carbaldehyd

- forbindelse nr. 385: 4-{[4-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]methyl}benzonitril


35 - forbindelse nr. 386: 3-{[4-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenoxy]methyl}benzonitril

- forbindelse nr. 387: 4-[3-{4-[(4-chlor-2-fluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 5 - forbindelse nr. 388: 4-[3-{4-[1-(3,4-dichlorphenyl)ethoxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- forbindelse nr. 389: 4-{7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-3-{4-[(3-hydroxybenzyl)oxy]-3-methoxybenzyl}-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 10 - forbindelse nr. 390: 4-[7-fluor-3-{4-[(3-fluorbenzyl)oxy]-3-methoxybenzyl}-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 15 - forbindelse nr. 391: 4-[3-{4-[(3,4-difluorbenzyl)oxy]-3-methoxybenzyl}-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd
- 20 - forbindelse nr. 392: 4-{3-[4-(5,6-dichlor-1H-benzimidazol-1-yl)-3-methoxybenzyl]-7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd
- 25 - forbindelse nr. 393: 4-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)phenyl-3,4-dichlorbenzensulfonat
- forbindelse nr. 394: 4-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenyl-3,4-dichlorbenzensulfonat
- 30 - forbindelse nr. 403: 3,4-dichlor-N-[4-({7-fluor-6-[2-fluor-1-(fluormethyl)ethoxy]-1-(1-formylpiperidin-4-yl)-2,4-dioxo-1,4-dihydroquinazolin-3(2H)-yl}methyl)-2-methoxyphenyl]benzamid
- 35

12. Fremgangsmåde til fremstilling af en forbindelse ifølge et hvilket som helst af kravene 1 til 11, kendetegnet ved, at en forbindelse med formlen




5 hvor R, Ra, m og n er som defineret i krav 1, og hvor der desuden kan være en gruppe R2 som defineret for forbindelserne med formlen (I) i position 7 på quinazolinindionstrukturen, omsættes med en forbindelse med formlen



ved henholdsvis en alkylerings- eller en Mitsunobureaktion.

13. Forbindelse med den almene formel (XVIII)



15 hvor R, Ra, m og n er som defineret i krav 1, og hvor der desuden kan være en gruppe R2 som defineret for forbindelserne med formlen (I) i position 7 på quinazolinindionstrukturen,

20 14. Forbindelse med den almene formel (XVIII) ifølge krav 12, der er valgt blandt forbindelse nr. 32 4-[6-(2,2-difluorethoxy)-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl]piperidin-1-carbaldehyd og forbindelse nr. 55 4-{6-[2-fluor-1-(fluormethyl)ethoxy]-2,4-dioxo-3,4-dihydroquinazolin-1(2H)-yl}piperidin-1-carbaldehyd, i form af en base, et hydrat, et solvat eller blandinger heraf.

15. Lægemiddel, kendetegnet ved, at det omfatter mindst en forbindelse med formlen (I) ifølge et hvilket som helst af kravene 1 til 11.

5 16. Farmaceutisk præparat, kendetegnet ved, at det omfatter mindst en forbindelse med formlen (I) ifølge et hvilket som helst af kravene 1 til 11 samt mindst et farmaceutisk acceptabelt excipiens.

10 17. Anvendelse af en forbindelse med formlen (I) ifølge et hvilket som helst af kravene 1 til 11 til fremstilling af et lægemiddel, der er beregnet til forebyggelse og/eller behandling af inflammatoriske eller immuno-inflammatoriske lidelser, herunder astma, kronisk obstruktiv
15 bronkopneumopati (COPD), allergisk rhinit, allergi, Crohns sygdom, colitis ulcerosa, myasthenia gravis, atopisk dermatit, psoriasis, systemisk lupus erythematosus, reumatoid polyarthrit, diabetes, dissemineret sklerose, til behandling af organtransplantationer, til behandling
20 og/eller forebyggelse af cancer, især osteosarkom eller adenokarcinom, til behandling og/eller forebyggelse af knoglesygdomme, herunder osteopeni og osteoporose, til behandling og/eller forebyggelse af akut insufficiens, til behandling og/eller forebyggelse af smerter, neuropatiske
25 smerter og viscerale smerter.

18. Anvendelse af en forbindelse med formlen (I) ifølge krav 17 til fremstilling af et lægemiddel, der er beregnet til behandling og/eller forebyggelse af neuropatiske
30 smerter.