

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2011220788 B2**

(54) Title
Isolated bacterial strain of the genus Burkholderia and pesticidal metabolites therefrom

(51) International Patent Classification(s)
C12N 1/20 (2006.01) **A01N 63/02** (2006.01)
A01N 43/14 (2006.01) **A01P 7/00** (2006.01)
A01N 43/74 (2006.01)

(21) Application No: **2011220788** (22) Date of Filing: **2011.02.24**

(87) WIPO No: **WO11/106491**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/406,541	2010.10.25	US
61/308,287	2010.02.25	US

(43) Publication Date: **2011.09.01**

(44) Accepted Journal Date: **2015.03.12**

(71) Applicant(s)
Marrone Bio Innovations, Inc.

(72) Inventor(s)
Asolkar, Ratnakar;Koivunen, Marja;Marrone, Pamela;Huang, Huazhang;Cordova-Kreylos, Ana Lucia

(74) Agent / Attorney
Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000

(56) Related Art
VANDAMME, P., et al. Journal of Clinical Microbiology, 2000, vol. 38, no. 3, pages 1042-1047

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 September 2011 (01.09.2011)

(10) International Publication Number
WO 2011/106491 A3

(51) International Patent Classification:

C12N 1/20 (2006.01) A01N 43/14 (2006.01)
A01N 63/02 (2006.01) A01P 7/00 (2006.01)
A01N 43/74 (2006.01)

(74) Agent: AGRIS, Cheryl, H.; The Law Offices Of Cheryl H. Agris, Phd, PC, P.O. Box 8495, Pelham, NY 10803 (US).

(21) International Application Number:

PCT/US2011/026016

(22) International Filing Date:

24 February 2011 (24.02.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/308,287 25 February 2010 (25.02.2010) US
61/406,541 25 October 2010 (25.10.2010) US

(71) Applicant (for all designated States except US): **MARRONE BIO INNOVATIONS, INC.** [US/US]; 2121 Second Street, Suite 107B, Davis, CA 95618 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ASOLKAR, Ratnakar** [IN/US]; 4005 Cowell Blvd, Davis, CA 95618 (US). **KOIVUNEN, Marja** [FI/US]; 1236 Evans Court, Davis, CA 95618 (US). **MARRONE, Pamela** [US/US]; 3333 Victoria Place, Davis, CA 95616 (US). **HUANG, Huazhang** [CN/US]; 2674 Briscoe Place, Woodland, CA 95776 (US). **CORDOVA-KREYLOS, Ana Lucia** [MX/US]; 18 Simmons Way, Davis, CA 95616 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GI, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

[Continued on next page]

(54) Title: ISOLATED BACTERIAL STRAIN OF THE GENUS BURKHOLDERIA AND PESTICIDAL METABOLITES THEREFROM

Figure 7



(57) Abstract: A species of *Burkholderia sp* with no known pathogenicity to vertebrates but with pesticidal activity (e.g., plants, insects, fungi, weeds and nematodes) is provided. Also provided are natural products derived from a culture of said species and methods of controlling pests using said natural products.

WO 2011/106491 A3



-
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))* — *with sequence listing part of description (Rule 5.2(a))*

(88) Date of publication of the international search report:

15 March 2012

Published:

- *with international search report (Art. 21(3j))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

ISOLATED BACTERIAL STRAIN OF THE GENUS *BURKHOLDERIA* AND PESTICIDAL METABOLITES THEREFROM

5

TECHNICAL FIELD

Provided herein is a species of *Burkholderia sp* with no known pathogenicity to vertebrates, such as mammals, fish and birds but pesticidal activity against plants, insects, fungi and nematodes. Also provided are natural products derived from a culture of said species and methods of controlling germination and growth of dicotyledenous, monocotyledonous and sedge weeds, modulating growth of fungi and controlling pests such as insects and nematodes using said natural products.

BACKGROUND

Natural products are substances produced by microbes, plants, and other organisms. Microbial natural products offer an abundant source of chemical diversity, and there is a long history of utilizing natural products for pharmaceutical purposes. One such compound is FR901228 isolated from *Chromobacterium* and has been found to be useful as an antibacterial agent and antitumor agent (see, for example, Ueda et al., US Patent No. 7,396,665).

However, secondary metabolites produced by microbes have also been successfully found to have uses for weed and pest control in agricultural applications (see, for example, Nakajima et al. 1991; Duke et al., 2000; Lydon & Duke, 1999; Gerwick et al., US Patent No. 7,393,812). Microbial natural products have been also successfully developed into agricultural insecticides (see, for example, Salama et al. 1981; Thompson et al., 2000; Krieg et al. 1983). Sometimes, such natural products have been combined with chemical pesticides (see, for example, Gottlieb, US Patent No. 4,808,207).

Burkholderia

The *Burkholderia* genus, β -subdivision of the proteobacteria, comprises more than 40 species that inhabit diverse ecological niches (Compant et al., 2008). The bacterial species in the genus *Burkholderia* are ubiquitous organisms in soil and rhizosphere (Coenye and Vandamme, 2003; Parke and Gurian-Sherman, 2001). Traditionally, they have been known as plant pathogens, *B. cepacia* being the first one discovered and identified as the pathogen causing disease in onions (Burkholder, 1950). Several *Burkholderia* species have developed beneficial interactions with their plant hosts (see, for example, Cabballero-Mellado et al., 2004, Chen et al., 2007). Some *Burkholderia* species have also been found to be opportunistic human pathogens (see, for example, Cheng and Currie, 2005 and Nierman et al., 2004). Additionally, some *Burkholderia* species have been found to have

potential as biocontrol products (see for example, Burkhead et al., 1994; Knudsen et al., 1987; Jansiewicz et al., 1988; Gouge et al., US Patent Application No. 2003/0082147; Parke et al., US Patent No. 6,077,505; Casida et al., US Patent No. 6,689,357; Jeddelloh et al., WO2001055398; Zhang et al., US Patent No. 7,141,407). Some species of in this genus have been effective in bioremediation to decontaminate polluted soil or groundwater (see, for example, Leahy et al. 1996). Further, some *Burkholderia* species have been found to secrete a variety of extracellular enzymes with proteolytic, lipolytic and hemolytic activities, as well as toxins, antibiotics, and siderophores (see, for example, Ludovic et al., 2007; Nagamatsu, 2001).

Oxazoles, Thiazoles and Indoles

Oxazoles, thiazoles and indoles are widely distributed in plants, algae, sponges, and microorganisms. A large number of natural products contain one or more of the five-membered oxazole, thiazole and indole nucleus/moieties. These natural products exhibit a broad spectrum of biological activity of demonstrable therapeutic value. For example, bleomycin A (Tomohisa et al.), a widely prescribed anticancer drug, effects the oxidative degradation of DNA and uses a bithiazole moiety to bind its target DNA sequences (Vanderwall et al., 1997). Bacitracin (Ming et al., 2002), a thiazoline-containing peptide antibiotic, interdicts bacterial cell wall new biosynthesis by complexation with C55-bactoprenolpyrophosphate. Thiangazole (Kunze et al., 1993) contains a tandem array of one oxazole and three thiazolines and exhibits antiviral activity (Jansen et al., 1992). Yet other oxazole/thiazole-containing natural products such as thiostrepton (Anderson et al., 1970) and GE2270A (Selva et al., 1997) inhibit translation steps in bacterial protein synthesis. More than 1000 alkaloids with the indole skeleton have been reported from microorganisms. One-third of these compounds are peptides with masses beyond 500 Da where the indole is tryptophan derived. The structural variety of the remaining two-thirds is higher, and their biological activity seems to cover a broader range, including antimicrobial, antiviral, cytotoxic, insecticidal, antithrombotic, or enzyme inhibitory activity.

Brief Summary

According to a first aspect of the present invention, there is provided a composition comprising an isolated strain of a *Burkholderia* sp. which has the following characteristics:

(A) a 16S rRNA gene sequence comprising the forward sequence having at least 99% identity to the sequence set forth in SEQ ID NO: 8, 11, 12 and a reverse sequence having at least 99% identity to the sequence set forth in SEQ ID NO:9, 10, 13, 14 and 15;

(B) pesticidal activity;

(C) produces a pesticidal compound selected from the group consisting of: templazole A, templazole B, templamide A, templamide B, and FR90128; and at least one of a carrier, diluent, surfactant, or adjuvant.

According to a second aspect of the present invention, there is provided a composition comprising substantially pure culture, cell fraction, whole cell broth, or supernatant derived from the strain in accordance with the first aspect of the present invention or extract thereof.

According to a third aspect of the present invention, there is provided a method for producing a compound selected from the group consisting of

(i) templazole A;

(ii) teplazole B;

(iii) templamide A;

(iv) templamide B; and

(v) FR90128;

wherein the method comprises the steps of culturing the strain of in accordance with the first aspect of the present invention and producing said compound.

According to a fourth aspect of the present invention, there is provided a method for isolating a compound from a strain of *Burkholderia* comprising

(a) producing one or more said compound according to the method in accordance with the fifth aspect of the present invention;

(b) isolating the one or more said compound produced in step (a) from the supernatant of said culture.

According to a fifth aspect of the present invention, there is provided a composition comprising

(a) a pure culture, cell fraction, whole cell broth, or supernatant derived from the strain in accordance with the first aspect of the present invention or extract thereof;

(b) a second substance, wherein said second substance is a chemical or biological pesticide and

(c) optionally at least one of a carrier, diluent, surfactant, adjuvant.

According to a sixth aspect of the present invention, there is provided a method for modulating pest infestation in a plant comprising applying to the plant and/or seeds thereof and/or substrate used for growing said plant an amount of the composition in accordance with any one of the first, second and fifth aspects of the present invention effective to modulate said pest infestation.

According to a seventh aspect of the present invention, there is provided a method for modulating emergence and/or growth of monocotyledonous, sedge or dicotyledonous weeds comprising applying to said weed or soil an amount of the composition in accordance with any one of the first, second and fifth aspects of the present invention effective to modulate said emergence and/or growth of said weeds.

According to an eighth aspect of the present invention, there is provided a seed comprising the composition in accordance with any one of the first, second and fifth aspects of the present invention.

Provided herein is an isolated strain of a non-*Burkholderia cepacia*, non-*Burkholderia plantari*, non-*Burkholderia gladioli*, *Burkholderia sp.* which has the following characteristics:

- a. Has a 16S rRNA gene sequence comprising a forward sequences having at least 99.0% identity to the sequences set forth in SEQ ID NO:8, 11 and 12 and a

reverse sequence having at least 99.0% identity to SEQ ID NO:9, 10, 13-15;

b. Has pesticidal, in particular, herbicidal, insecticidal, fungicidal and nematocidal activity;

c. Produces at least one of the compounds selected from the group consisting of:

5 (i) a compound having the following properties: (a) a molecular weight of about 525-555 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS); (b) ¹H NMR values of 6.22, 5.81, 5.69, 5.66, 5.65, 4.64, 4.31, 3.93, 3.22, 3.21, 3.15, 3.10, 2.69, 2.62, 2.26, 2.23, 1.74, 1.15, 1.12, 1.05, 1.02; (c) has ¹³C NMR values of 172.99, 172.93, 169.57, 169.23, 167.59, 130.74, 130.12, 129.93, 128.32, 73.49, 62.95, 59.42, 57.73, 38.39, 10 38.00, 35.49, 30.90, 30.36, 29.26, 18.59, 18.38, 18.09, 17.93, 12.51 and (c) an High Pressure Liquid Chromatography (HPLC) retention time of about 10-15 minutes, on a reversed phase C-18 HPLC column using a water:acetonitrile (CH₃CN) gradient;

(ii) a compound having an oxazolyl-indole structure comprising at least one indole moiety, at least one oxazole moiety, at least one substituted alkyl group and at least one carboxylic ester group; at least 17 carbons and at least 3 oxygen and 2 nitrogens;

15 (iii) a compound having an oxazolyl-benzyl structure comprising at least one benzyl moiety, at least one oxazole moiety, at least one substituted alkyl group and at least one amide group; at least 15 carbons and at least 2 oxygen and 2 nitrogens;

(iv) a compound having at least one ester, at least one amide, at least three methylene groups, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least twenty five carbons and at least eight oxygen and one nitrogen and

d. is non-pathogenic (non-infectious) to vertebrate animals, such as mammals, birds and fish;

25 e. is susceptible to kanamycin, chloramphenicol, ciprofloxacin, piperacillin, imipenem, and a combination of sulphamethoxazole and trimethoprim and

f. contains the fatty acids 16:0, cyclo 17:0, 16:0 3- OH, 14:0, cyclo 19:0 ω8c, 18:0.

In a particular embodiment, the strain has the identifying characteristics of a *Burkholderia* A396 strain (NRRL Accession No. B-50319).

30 Disclosed herein are isolated compounds which are optionally obtainable or derived from *Burkholderia* species, or alternatively, organisms capable of producing these compounds that can be used to control various pests, particularly plant phytopathogenic pests, examples of which include but are not limited to insects, nematodes, bacteria, fungi. These compounds may also be used as herbicides.

35 In particular, the isolated pesticidal compounds may include but are not limited to:

(A) a compound having the following properties: (i) a molecular weight of about

525-555 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS); (ii) ¹H NMR δ values of 6.22, 5.81, 5.69, 5.66, 5.65, 4.64, 4.31, 3.93, 3.22, 3.21, 3.15, 3.10, 2.69, 2.62, 2.26, 2.23, 1.74, 1.15, 1.12, 1.05, 1.02; (iii) has ¹³C NMR δ values of 172.99, 172.93, 169.57, 169.23, 167.59, 130.74, 130.12, 129.93, 128.32, 73.49, 62.95, 59.42, 57.73, 38.39, 38.00, 35.49, 30.90, 30.36, 29.26, 18.59, 18.38, 18.09, 17.93, 12.51 and (iv) an High Pressure Liquid Chromatography (HPLC) retention time of about 10-15 minutes, on a reversed phase C-18 HPLC column using a water:acetonitrile (CH₃CN) gradient;

(B) a compound having an oxazolyl-indole structure comprising at least one indole moiety, at least one oxazole moiety, at least one substituted alkyl group and at least one carboxylic ester group; at least 17 carbons and at least 3 oxygen and 2 nitrogens;

(C) a compound having an oxazolyl-benzyl structure comprising at least one benzyl moiety, at least one oxazole moiety, at least one substituted alkyl group and at least one amide group; at least 15 carbons and at least 2 oxygen and 2 nitrogens;

(D) a compound having at least one ester, at least one amide, at least three methylene groups, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least twenty five carbons and at least eight oxygen and one nitrogen and

(E) a compound having at least one ester, at least one amide, an epoxide methylene group, at least one tetrahydropyranose moiety, at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least 25 carbons, at least 8 oxygens and at least 1 nitrogen.

In a particular embodiment, the isolated compounds may include but are not limited to:

(A) a compound having an oxazolyl-indole structure comprising at least one indole moiety, at least one oxazole moiety, at least one substituted alkyl group, at least one carboxylic ester group, at least 17 carbons, at least 3 oxygens and at least 2 nitrogens; and which has at least one of the following: (i) a molecular weight of about 275-435; (ii) ¹H NMR δ values at 8.44, 8.74, 8.19, 7.47, 7.31, 3.98, 2.82, 2.33, 1.08; (iii) ¹³C NMR δ values of δ 163.7, 161.2, 154.8, 136.1, 129.4, 125.4, 123.5, 123.3, 121.8, 121.5, 111.8, 104.7, 52.2, 37.3, 28.1, 22.7, 22.7; (iv) an High Pressure Liquid Chromatography (HPLC) retention time of about 10-20 minutes on a reversed phase C-18 HPLC column using a water:acetonitrile (CH₃CN) with a gradient solvent system and UV detection of 210 nm; (v) UV absorption bands at about 226, 275, 327 nm.;

(B) a compound having an oxazolyl-benzyl structure comprising at least one benzyl moiety, at least one oxazole moiety, at least one substituted alkyl group and at least one amide group; at least 15 carbons and at least 2 oxygens, at least 2 nitrogens; and at least one of the following characteristics: (i) a molecular weight of about 240-290 as determined by
5 Liquid Chromatography/Mass Spectroscopy (LC/MS); (ii) ^1H NMR δ values at about 7.08, 7.06, 6.75, 3.75, 2.56, 2.15, 0.93, 0.93; (iii) ^{13}C NMR δ values of 158.2, 156.3, 155.5, 132.6, 129.5, 129.5, 127.3, 121.8, 115.2, 115.2, 41.2, 35.3, 26.7, 21.5, 21.5; (iv) a High Pressure Liquid Chromatography (HPLC) retention time of about 6-15 minutes, on a reversed phase C-18 HPLC column using a water:acetonitrile (CH_3CN) gradient and (v) UV
10 absorption bands at about 230, 285, 323 nm;


(C) a non-epoxide compound comprising at least one ester, at least one amide, at least three methylene groups, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least twenty five carbons, at least eight oxygens and one nitrogen and at least one of the
15 following characteristics: (i) a molecular weight of about 530-580 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS); (ii) ^1H NMR δ values of 6.40, 6.39, 6.00, 5.97, 5.67, 5.54, 4.33, 3.77, 3.73, 3.70, 3.59, 3.47, 3.41, 2.44, 2.35, 2.26, 1.97, 1.81, 1.76, 1.42, 1.37, 1.16, 1.12, 1.04; (iii) ^{13}C NMR δ values of 173.92, 166.06, 145.06, 138.76, 135.71, 129.99, 126.20, 123.35, 99.75, 82.20, 78.22, 76.69, 71.23, 70.79, 70.48, 69.84,
20 60.98, 48.84, 36.89, 33.09, 30.63, 28.55, 25.88, 20.37, 18.11, 14.90, 12.81, 9.41; (iv) a High Pressure Liquid Chromatography (HPLC) retention time of about 7-12 minutes, on a reversed phase C-18 HPLC column using a water:acetonitrile (CH_3CN) with a gradient solvent system and UV detection of 210 nm; (v) a molecular formula of $\text{C}_{28}\text{H}_{45}\text{NO}_{10}$ which was determined by interpretation of the ESIMS and NMR data analysis; (vi) UV absorption
25 bands between about 210-450 nm;

(D) a compound comprising (i) at least one ester, at least one amide, an epoxide methylene group, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least 25 carbons, at least 8 oxygens and at least 1 nitrogen, (ii) ^{13}C NMR δ values of 174.03, 166.12, 143.63, 137.50,
30 134.39, 128.70, 126.68, 124.41, 98.09, 80.75, 76.84, 75.23, 69.87, 69.08, 68.69, 68.60, 48.83, 41.07, 35.45, 31.67, 29.19, 27.12, 24.55, 19.20, 18.95, 13.48, 11.39, 8.04, (iii) a molecular formula of $\text{C}_{28}\text{H}_{43}\text{NO}_9$, and at least one of: (i) ^1H NMR δ values at about 6.41, 6.40, 6.01, 5.97, 5.67, 5.55, 4.33, 3.77, 3.75, 3.72, 3.64, 3.59, 3.54, 3.52, 2.44, 2.34, 2.25, 1.96, 1.81, 1.76, 1.42, 1.38, 1.17, 1.12, 1.04; (ii) an High Pressure Liquid Chromatography

(HPLC) retention time of about 6-15 minutes, on a reversed phase C-18 HPLC column using a water:acetonitrile (CH₃CN) gradient; (iii) UV absorption band between about 210-450 nm and most particularly at about 234 nm.


5 In a more particular embodiment, provided are compounds including but not limited to:

(A) a compound having the structure ##STR001##



10 or a pesticidally acceptable salt or stereoisomers thereof, wherein M is 1, 2, 3 or 4; n is 0, 1, 2, or 3; p and q are independently 1 or 2; X is O, NH or NR; R₁, R₂ and R₃ are the same or different and independently an amino acid side-chain moiety or an amino acid side-chain derivative and R is a lower chain alkyl, aryl or arylalkyl moiety;

(B) a compound having the structure ##STR002##




##STR002##

15 wherein X, Y and Z are each independently --O, --NR₁, or --S, wherein R₁ is --H or C₁-C₁₀ alkyl; R₁, R₂ and m are each independently --H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido,


carboxyl, --C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfuryl and "m" may be located anywhere on the oxazole ring;

(C) a compound having the structure ##STR002a##




5 wherein R₁ is --H or C₁-C₁₀ alkyl; R₂ is an alkyl ester;

(D) a compound having the structure ##STR003##




wherein: X and Y are each independently --OH, --NR₁, or --S, wherein R₁ is --H or C₁-C₁₀ alkyl; R₁, R₂ and m, a substituent on the oxazole ring, are each independently --H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, --C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfuryl;

15 (E) a compound having the structure ##STR003a##




wherein R₁ is --H or C₁-C₁₀ alkyl;

(F) a compound having the structure ##STR004a##



Wherein X, Y and Z are each independently -O, -NR, or -S, wherein R is H or C₁-C₁₀ alkyl; R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, and R₁₃ are each independently H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, -C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfuryl.


(G) a compound having the structure ##STR004b##



10

wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, and R₁₃ are each independently H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, -C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfuryl;


(H) a compound having the structure ##STR004c##



20 wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₁₁, are each independently H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy,

halogen, amino, amido, carboxyl, -C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfonyl;

(I) a compound having the structure ##STR005##




5

wherein X and Y are each independently --OH, --NR₁, or --S, wherein R₁, R₂ are each independently --H, alkyl (e.g., C₁-C₁₀ alkyl), substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,

10 heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, --C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfonyl;

(J) a compound having the structure ##STR006a##



15 Wherein X, Y and Z are each independently -O, -NR, or -S, wherein R is H or C₁-C₁₀ alkyl; R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₁₁, R₁₂, and R₁₃ are each independently H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, 20 halogen, amino, amido, carboxyl, -C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfonyl.

In a most particular embodiment, the compounds may include but are not limited to

(i) templazole A;


(ii) templazole B;

25 (iii) templamide A;


(iv) templamide B;

(v) FR90128;


(vi)



(vii)




(viii)



5

(ix)




(x)




10


(xi)




(xii)




(xiii)




(xiv)




(xv)



(xvi)




(xvii)




5

10

(xviii)




(xix)




5

(xx)




(xxi)




10


(xxii)



(xxiii)



5



(XL) FR901465

Also provided are methods of obtaining the compounds set forth above. In particular, the method comprises culturing the *Burkholderia* strain disclosed herein and producing the compound. Further provided is a method for isolating these compounds by isolating the compound(s) produced by a *Burkholderia* strain comprising isolating compounds produced from a supernatant of a culture of said *Burkholderia* strain.

Further provided is a combination comprising (a) a first substance selected from the group consisting of (i) a pure culture, cell fraction or supernatant derived from the *Burkholderia* strain set forth above or extract thereof for use optionally as a pesticide; (ii) one or more of the compounds set forth above (b) optionally a second substance, wherein said second substance is a chemical or biological pesticide and (c) optionally at least one of a carrier, diluent, surfactant, adjuvant, or pesticide. In a particular embodiment, the combination is a composition. In a related aspect, provided herein is a seed coated with said composition.

In a related aspect, disclosed is a method for modulating pest infestation in a plant comprising applying to the plant and/or seeds thereof and/or substrate used for growing said plant and/or a method for modulating emergence and/or growth of monocotyledonous, sedge or dicotyledonous weeds comprising applying to said weed or soil an amount of

- (I) (a) the isolated compounds set forth above and (b) optionally another substance, wherein said substance is a pesticide (e.g. nematocide, herbicide, fungicide, insecticide) or
- (II) the composition or combination set forth above

in an amount effective to modulate pest infestation and/or emergence or growth of monocotyledonous, sedge or dicotyledonous weeds.

In another related aspect, provided is the use of the strains, cultures, extracts, supernatants, combinations, compounds set forth above for modulating pest infestation in a plant comprising applying to the plant and/or seeds thereof and/or substrate used for growing said plant and/or a method for modulating emergence and/or growth of monocotyledonous, sedge or dicotyledonous weeds.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the comparison of the growth rate of *Burkholderia* A396 to *Burkholderia multivorans* ATCC 17616.

Figure 2 shows the effect of *Burkholderia* A396 extract on bindweed.

Figure 3 shows the effect of *Burkholderia* A396 extract on pigweed.

Figure 4 shows the effect of *Burkholderia* A396 extract on Cabbage looper (*Tricoplusia ni*).

Figure 5 shows the effect of *Burkholderia* A396 culture broth on Beet armyworm

(*Spodoptera exigua*).

Figure 6 shows the effect of *Burkholderia* A396 culture broth on the motility of juvenile root-knot nematodes (*Meloidogyne incognita*).

Figure 7 is a schematic representation of purification scheme for obtaining the
5 templazole and templamide compounds.

Figure 8 shows results of an *in vitro* assay to test the fungicidal effect of FR90128 on *Botrytis cinerea* (left) and *Phytophthora* sp. (right).

Figure 9 shows the effect of *Burkholderia* A396 culture broth on the average gall index (% control) of cucumber roots cv. Toschka inoculated with 3000 eggs of *Meloidogyne*
10 *sp.* 14 days after inoculation and application.

Figure 10 Effect of *Burkholderia* A396 culture broth on the average gall index of cucumber roots cv. Toschka inoculated with 3000 eggs of *Meloidogyne sp.* 14 days after inoculation and application.

15 DETAILED DESCRIPTION OF EMBODIMENTS

While the compositions and methods heretofore are susceptible to various modifications and alternative forms, exemplary embodiments will herein be described in detail. It should be understood, however, that there is no intent to limit the invention to the particular forms disclosed, but on the contrary, the intention is to cover all modifications,
20 equivalents, and alternatives falling within the spirit and scope of the invention as defined by the appended claims.

Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that
25 stated range, is included therein. Smaller ranges are also included. The upper and lower limits of these smaller ranges are also included therein, subject to any specifically excluded limit in the stated range.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this
30 invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described.

It must be noted that as used herein and in the appended claims, the singular forms "a," "and" and "the" include plural references unless the context clearly dictates otherwise.

35 As defined herein, "derived from" means directly isolated or obtained from a particular source or alternatively having identifying characteristics of a substance or organism isolated or obtained from a particular source.

As defined herein, an "isolated compound" is essentially free of other compounds or substances, e.g., at least about 20% pure, preferably at least about 40% pure, more preferably about 60% pure, even more preferably about 80% pure, most preferably about 90% pure, and even most preferably about 95% pure, as determined by analytical methods, including but not limited to chromatographic methods, electrophoretic methods.

As used herein, the term "alkyl" refers to a monovalent straight or branched chain hydrocarbon group having from one to about 12 carbon atoms, including methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-hexyl, and the like.

As used herein, "substituted alkyl" refers to alkyl groups further bearing one or more substituents selected from hydroxy, alkoxy, mercapto, cycloalkyl, substituted cycloalkyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, aryloxy, substituted aryloxy, halogen, cyano, nitro, amino, amido, --C(O)H, acyl, oxyacyl, carboxyl, sulfonyl, sulfonamide, sulfuryl, and the like.

As used herein, "alkenyl" refers to straight or branched chain hydrocarbyl groups having one or more carbon-carbon double bonds, and having in the range of about 2 up to 12 carbon atoms, and "substituted alkenyl" refers to alkenyl groups further bearing one or more substituents as set forth above.

As used herein, "alkynyl" refers to straight or branched chain hydrocarbyl groups having at least one carbon-carbon triple bond, and having in the range of about 2 up to 12 carbon atoms, and "substituted alkynyl" refers to alkynyl groups further bearing one or more substituents as set forth above.

As used herein, "aryl" refers to aromatic groups having in the range of 6 up to 14 carbon atoms and "substituted aryl" refers to aryl groups further bearing one or more substituents as set forth above.

As used herein, "heteroaryl" refers to aromatic rings containing one or more heteroatoms (e.g., N, O, S, or the like) as part of the ring structure, and having in the range of 3 up to 14 carbon atoms and "substituted heteroaryl" refers to heteroaryl groups further bearing one or more substituents as set forth above.

As used herein, "alkoxy" refers to the moiety --O-alkyl-, wherein alkyl is as defined above, and "substituted alkoxy" refers to alkoxy groups further bearing one or more substituents as set forth above.

As used herein, "thioalkyl" refers to the moiety --S-alkyl-, wherein alkyl is as defined above, and "substituted thioalkyl" refers to thioalkyl groups further bearing one or more substituents as set forth above.

As used herein, "cycloalkyl" refers to ring-containing alkyl groups containing in the range of about 3 up to 8 carbon atoms, and "substituted cycloalkyl" refers to cycloalkyl groups further bearing one or more substituents as set forth above.

As used herein, "heterocyclic", refers to cyclic (i.e., ring-containing) groups containing one or more heteroatoms (e.g., N, O, S, or the like) as part of the ring structure, and having in the range of 3 up to 14 carbon atoms and "substituted heterocyclic" refers to heterocyclic groups further bearing one or more substituent's as set forth above.

The *Burkholderia* Strain

The *Burkholderia* strain set forth herein is a non-*Burkholderia cepacia* complex, non-*Burkholderia plantari*, non-*Burkholderia gladioli*, *Burkholderia sp* and non-pathogenic to vertebrates, such as birds, mammals and fish. This strain may be isolated from a soil sample using procedures known in the art and described by Lorch et al., 1995. The *Burkholderia* strain may be isolated from many different types of soil or growth medium. The sample is then plated on potato dextrose agar (PDA). The bacteria are gram negative, and it forms round, opaque cream-colored colonies that change to pink and pinkish-brown in color and mucoid or slimy over time.

Colonies are isolated from the potato dextrose agar plates and screened for those that have biological, genetic, biochemical and/or enzymatic characteristics of the *Burkholderia* strain of the present invention set forth in the Examples below. In particular, the *Burkholderia* strain has a 16S rRNA gene comprising a forward sequence that is at least about 99.0%, preferably about 99.5%, more preferably about 99.9% and most preferably about 100% identical to the sequence set forth in SEQ ID NO: 8, 11 and 12 and a forward sequence that is at least about 99.0%, preferably about 99.5%, more preferably about 99.9% and most preferably about 100% identical to the sequence set forth in SEQ ID NO: 9, 10, 13, 14 and 15 as determined by clustal analysis. Furthermore, as set forth below, this *Burkholderia* strain may, as set forth below, have pesticidal activity, particularly, virucidal, herbicidal, germicidal, fungicidal, nematocidal, bactericidal and insecticidal and more particularly, herbicidal, insecticidal, fungicidal and nematocidal activity. It is not pathogenic to vertebrate animals, such as mammals, birds, and fish.

Additionally, the *Burkholderia* strain produces at least the pesticidal compounds set forth in the instant disclosure.

The *Burkholderia* strain is susceptible to kanamycin, chloramphenicol, ciprofloxacin, piperacillin, imipenem, and a combination of sulphamethoxazole and trimethoprim and contains the fatty acids 16:0, cyclo 17:0, 16:0 3-OH, 14:0, cyclo 19:0, 18:0.


This *Burkholderia* strain may be obtained by culturing a microorganism having the

identifying characteristics of *Burkholderia* A396 (NRRL Accession No. B-50319) on Potato Dextrose Agar (PDA) or in a fermentation medium containing defined carbon sources such as glucose, maltose, fructose, galactose, and undefined nitrogen sources such as peptone, tryptone, soytone, and NZ amine.

5

Pesticidal Compounds


The pesticidal compound disclosed herein may have the following properties: (a) is obtainable from a novel *Burkholderia* species, e.g., A396; (b) is, in particular, toxic to most common agricultural insect pests; (c) has a molecular weight of about 525-555 and more particularly, 540 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS); (d) has ^1H NMR values of 6.22, 5.81, 5.69, 5.66, 5.65, 4.64, 4.31, 3.93, 3.22, 3.21, 3.15, 3.10, 2.69, 2.62, 2.26, 2.23, 1.74, 1.15, 1.12, 1.05, 1.02; (e) has ^{13}C NMR values of 172.99, 172.93, 169.57, 169.23, 167.59, 130.74, 130.12, 129.93, 128.32, 73.49, 62.95, 59.42, 57.73, 38.39, 38.00, 35.49, 30.90, 30.36, 29.26, 18.59, 18.38, 18.09, 17.93, 12.51 (e) has an High Pressure Liquid Chromatography (HPLC) retention time of about 10-15 minutes, more specifically about 12 minutes and even more specifically about 12.14 min on a reversed phase C-18 HPLC (Phenomenex, Luna 5μ C18 (2) 100A, 100 x 4.60 mm) column using a water:acetonitrile (CH_3CN) with a gradient solvent system (0-20 min 90 - 0 % aqueous CH_3CN , 20-24 min 100% CH_3CN , 24-27 min, 0-90 % aqueous CH_3CN , 27-30 min 90% aqueous CH_3CN) at 0.5 mL/min flow rate and UV detection of 210 nm (f) has a molecular formula, $\text{C}_{24}\text{H}_{36}\text{N}_4\text{O}_6\text{S}_2$, which is determined by interpretation of ^1H , ^{13}C NMR and LC/MS data (g) a ^{13}C NMR spectrum with signals for all 24 carbons, including 5 methyl, 4 methylene, 9 methine, and 6 quaternary carbons and (g) ^1H NMR spectrum displaying characteristics of a typical depsipeptide, illustrating three -amino protons [4.63, 4.31, 3.93], and one ester carbinol proton [5.69]. In a particular embodiment, the compound has the structure ##STR001##:




Or a pesticidally acceptable salt or stereoisomers thereof, wherein M is 1, 2, 3 or 4; n is 0, 1, 2, or 3; p and q are independently 1 or 2; X is O, NH or NR; R1, R2 and R3 are the same or

different and independently an amino acid side-chain moiety or an amino acid side-chain derivative and R is a lower chain alkyl, aryl or arylalkyl moiety.

In an even more particular embodiment, the compound has the structure of FR90128:




Provided herewith are compounds set forth in ##STR002##:




##STR002##

- 10 wherein: X, Y and Z are each independently --O, --NR₁, or --S, wherein R₁ is --H or C₁-C₁₀ alkyl; R₁, R₂ and m are each independently --H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, --C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfonyl.
- 15


In an even another particular embodiment, Family ##STR002## compounds may be the compounds set forth in (vi)-(xix).



(vii)




(vii)




5

(ix)




(x)




10


(xi)




(xii)




(xiii)



(xiv)




(xv)




5

(xvi)




(xvii)




10

(xviii)




(xix)



These are from either natural materials or compounds obtained from commercial
 5 sources or by chemical synthesis. Natural sources of Family ##STR002## compounds
 include, but are not limited to, microorganisms, alga, and sponges. In a more particular
 embodiment, microorganisms which include the Family ##STR002## compounds include
 but are not limited to, or alternatively, Family ##STR002## compounds may be derived
 from species such as *Streptoverticillium waksmanii* (compound **vi**) (Umehara, et al., 1984),
 10 *Streptomyces pimprina* (compound **vii**) (Naiket al., 2001), *Streptoverticillium olivoreticuli*
 (compounds **viii**, **ix**, **x**) (Koyama Y., et al., 1981), *Streptomyces* sp (compounds **xi**, **xii**)
 (Watabe et al., 1988), *Pseudomonas syringae* (compounds **xiii**, **xiv**) (Pettit et al., 2002).
 Family ##STR002## compounds may also be derived from algae including but not limited
 to red alga (compound **xv**) (N'Diaye, et al., 1996), red alga *Martensia fragilis* (compound
 15 **xvi**) (Takahashi S. et al., 1998), *Diazona chinensis* (compounds **xvii** & **xviii**) (Lindquist N.
 et al., 1991), *Rhodophycota haraldiophyllum* sp (compound **xix**) (Guella et al., 1994).


Also provided is ##STR003##:



wherein: X and Y are each independently --OH, --NR₁, or --S, wherein R₁ is --H or C₁-C₁₀
 20 alkyl; R₁, R₂ and m, a substituent on the oxazole ring, are each independently --H, alkyl,
 substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted

aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, --C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfonyl.


5 Further provided is ##STR005##:



wherein X and Y are each independently --OH, --NR₁, or --S, wherein R₁, R₂ are each independently --H, alkyl (e.g., C₁-C₁₀ alkyl), substituted alkyl, alkenyl, substituted alkenyl, 10 alkyne, substituted alkyne, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, --C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfonyl.


In a particular embodiment, Family ##STR005## compounds such as compounds 15 from **xx-xxiii** set forth below may be derived from natural or commercial sources or by chemical synthesis:

(xx)




20


(xxi)



(xxii)



(xxiii)



5 Natural sources of Family ##STR005## compounds include, but are not limited to plants, corals, microorganisms, and sponges. The microorganisms include, but are not limited to *Streptomyces griseus* (compound **xx**) (Hirota et al., 1978), *Streptomyces albus* (compound **xxi**) (Werner et al., 1980). Family STR004 compounds may also be derived from algae including but not limited to *Haraldiophyllum sp* (compound **xxii**) (Guella et al.,
10 2006), and red algae (compound **xxiii**) (N'Diaye et al., 1994).

In one embodiment, the compound may be derived from or is obtainable from a microorganism, and in particular from *Burkholderia* species and characterized as having a structure comprising at least one ester, at least one amide, at least three methylene groups, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six
15 methyl groups, at least three hydroxyl groups, at least twenty five carbons and at least eight oxygen and one nitrogen. The compound further comprises at least one of the following characteristics:

(a) pesticidal properties and in particular, nematicidal, fungicidal, insecticidal and herbicidal properties;

20 (b) a molecular weight of about 530-580 and more particularly, 555 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS);

(c) ¹H NMR values of δ 6.40, 6.39, 6.00, 5.97, 5.67, 5.54, 4.33, 3.77, 3.73, 3.70, 3.59, 3.47, 3.41, 2.44, 2.35, 2.26, 1.97, 1.81, 1.76, 1.42, 1.37, 1.16, 1.12, 1.04;

(d) ¹³C NMR values of δ 173.92, 166.06, 145.06, 138.76, 135.71, 129.99, 126.20,
25 123.35, 99.75, 82.20, 78.22, 76.69, 71.23, 70.79, 70.48, 69.84, 60.98, 48.84, 36.89, 33.09, 30.63, 28.55, 25.88, 20.37, 18.11, 14.90, 12.81, 9.41;

(e) an High Pressure Liquid Chromatography (HPLC) retention time of about 7-12 minutes, more specifically about 10 minutes and even more specifically about 10.98 min on a reversed phase C-18 HPLC (Phenomenex, Luna 5 μ C18(2) 100 A, 100 x 4.60 mm)
30 column using a water:acetonitrile (CH₃CN) with a gradient solvent system (0-20 min; 90 - 0


% aqueous CH₃CN, 20-24 min; 100% CH₃CN, 24-27 min; 0 - 90 % aqueous CH₃CN, 27-30 min; 90% aqueous CH₃CN) at 0.5 mL/min flow rate and UV detection of 210 nm;

(f) ¹³C NMR spectrum which exhibits 28 discrete carbon signals which may be attributed to six methyls, four methylene carbons, and thirteen methines including five *sp*²,
5 four quaternary carbons;

(g) a molecular formula of C₂₈H₄₅NO₁₀ which was determined by interpretation of the ESIMS and NMR data analysis;


(h) UV absorption bands between about 210-450 nm and most particularly at about 234 nm.

10 Also provided are compounds having the structure ##STR004a##:




Wherein X, Y and Z are each independently -O, -NR, or -S, wherein R is H or C₁-C₁₀ alkyl; R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, and R₁₃ are each independently H, alkyl,
15 aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, -C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide, or sulfuryl.

In a particular embodiment, the compound has the structure set forth in
20 ##STR004b##:




wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, and R₁₃ are as previously defined for ##STR004a##.

In a more particular embodiment, the compound is Templamide A with the
25 following structure:



In another embodiment, provided is a compound having formula ##STR004c##:



Wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₁₁ are as previously defined for ##STR004a##.

- 5 In another embodiment, provided is a compound which may be derived from *Burkholderia* species and characterized as having a structure comprising at least one ester, at least one amide, an epoxide methylene group, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least 25 carbons and at least 8 oxygen and 1 nitrogen, and pesticide activity. The
- 10 compound further comprises at least one of the following characteristics:

(a) pesticidal properties and in particular, insecticidal, fungicidal, nematocidal and herbicidal properties;

(b) a molecular weight of about 520-560 and particularly 537 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS);

- 15 (c) ¹H NMR δ values at about 6.41, 6.40, 6.01, 5.97, 5.67, 5.55, 4.33, 3.77, 3.75, 3.72, 3.64, 3.59, 3.54, 3.52, 2.44, 2.34, 2.25, 1.96, 1.81, 1.76, 1.42, 1.38, 1.17, 1.12, 1.04;

(d) ¹³C NMR values of δ 174.03, 166.12, 143.63, 137.50, 134.39, 128.70, 126.68, 124.41, 98.09, 80.75, 76.84, 75.23, 69.87, 69.08, 68.69, 68.60, 48.83, 41.07, 35.45, 31.67, 29.19, 27.12, 24.55, 19.20, 18.95, 13.48, 11.39, 8.04;


- 20 (e) High Pressure Liquid Chromatography (HPLC) retention time of about 6-15 minutes, more specifically about 8 minutes on a reversed phase C-18 HPLC column using a water:acetonitrile (CH₃CN) gradient, particularly, an High Pressure Liquid Chromatography (HPLC) retention time of about 8-15 minutes, more specifically about 11 minutes and even more specifically about 11.73 min on a reversed phase C-18 HPLC (Phenomenex, Luna 5μ
- 25 C18(2) 100 A, 100 x 4.60 mm) column using a water:acetonitrile (CH₃CN) with a gradient solvent system (0-20 min; 90 - 0 % aqueous CH₃CN, 20-24 min; 100% CH₃CN, 24-27 min;

0 - 90 % aqueous CH₃CN, 27-30 min; 90% aqueous CH₃CN) at 0.5 mL/min flow rate and UV detection of 210 nm;

(f) a molecular formula of C₂₈H₄₃NO₉, which was determined by interpretation of the ESIMS and NMR data analysis;


- 5 (g) UV absorption bands at about 210-450 nm and most particularly at about 234 nm.

In a particular embodiment, the compound has the structure ##STR006a##:




- Wherein X, Y and Z are each independently -O-, -NR-, or -S-, wherein R is H or C₁-C₁₀ alkyl;
- 10 R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₁₁, R₁₂, and R₁₃ are each independently H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkyl, substituted cycloalkyl, alkoxy, substituted alkoxy, thioalkyl, substituted thioalkyl, hydroxy, halogen, amino, amido, carboxyl, -C(O)H, acyl, oxyacyl, carbamate, sulfonyl, sulfonamide,
- 15 or sulfuryl.

In a particular embodiment, the compound has the structure:




Templamide A

In another embodiment, provided is a compound having formula ##STR006b##:



- 20 Wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₁₁ are as previously defined for ##STR006a##.

In a more particular embodiment, the compound is Templamide B with the following structure:



Templamide B

In yet another particular embodiment, the compound may be derived from *Burkholderia* species and characterized as having a structure comprising at least one ester, at least one amide, an epoxide methylene group, at least one tetrahydropyranose moiety and at least three olefinic double bonds, at least six methyl groups, at least three hydroxyl groups, at least 25 carbons and at least 8 oxygens and at least 1 nitrogen. The compound further comprises at least one of the following characteristics:

(a) pesticidal properties and in particular, insecticidal, fungicidal, nematocidal and herbicidal properties;

(b) a molecular weight of about 510-550 and particularly about 523 as determined by Liquid Chromatography/Mass Spectroscopy (LC/MS);

(c) ^1H NMR δ values at about 6.41, 6.40, 6.01, 5.98, 5.68, 5.56, 4.33, 3.77, 3.75, 3.72, 3.65, 3.59, 3.55, 3.50, 2.44, 2.26, 2.04, 1.96, 1.81, 1.75, 1.37, 1.17, 1.04;


(d) ^{13}C NMR δ values of 172.22, 167.55, 144.98, 138.94, 135.84, 130.14, 125.85, 123.37, 99.54, 82.19, 78.28, 76.69, 71.31, 70.13, 69.68, 48.83, 42.52, 36.89, 33.11, 30.63, 25.99, 21.20, 20.38, 18.14, 14.93, 12.84;

(e) an High Pressure Liquid Chromatography (HPLC) retention time of about 6-15 minutes, more specifically about 8 minutes on a reversed phase C-18 HPLC column using a water:acetonitrile (CH_3CN) gradient, particularly, an High Pressure Liquid Chromatography (HPLC) retention time of about 8-15 minutes, more specifically about 10 minutes and even more specifically about 10.98 min on a reversed phase C-18 HPLC (Phenomenex, Luna 5 μ C18(2) 100 A, 100 x 4.60 mm) column using a water:acetonitrile (CH_3CN) with a gradient solvent system (0-20 min; 90 - 0 % aqueous CH_3CN , 20-24 min; 100% CH_3CN , 24-27 min; 0 - 90 % aqueous CH_3CN , 27-30 min; 90% aqueous CH_3CN) at 0.5 mL/min flow rate and UV detection of 210 nm;

(f) a molecular formula of $\text{C}_{27}\text{H}_{41}\text{NO}_9$, which was determined by interpretation of the ESIMS and NMR data analysis;

(g) UV absorption bands at about 210-450 nm and most particularly at about 234 nm.

In a more particular embodiment, the compound is a known compound FR901465 which was isolated earlier from culture broth of a bacterium of *Pseudomonas* sp. No. 2663 (Nakajima et al. 1996) and had been reported to have anticancer activity with the following structure:




FR901465

5

In an even another particular embodiment, Family ##STR006a## compounds may be the compounds set forth in **xxiv** to **xxxix**. These are from either natural materials or compounds obtained from commercial sources or by chemical synthesis. Natural sources of Family ##STR006a## compounds include, but are not limited to, microorganisms, alga, and sponges. In a more particular embodiment, microorganisms which include the Family ##STR006a## compounds which may be derived from species such as *Pseudomonas* sp. No. 2663 (compounds **xxiv-xxvi**) (Nakajima et al., 1996). The synthetic analogues of the FR901464 (**xxvii-xxxix**) which have been synthesized and patented as anticancer compounds (see Koide et al., US Patent Application No. 2008/0096879 A1).

10



Compositions

A substantially pure culture, cell fraction or supernatant and compounds produced by the *Burkholderia* strain of the present invention, may be formulated into pesticidal

compositions.

The substances set forth above can be formulated in any manner. Non-limiting formulation examples include but are not limited to emulsifiable concentrates (EC), wettable powders (WP), soluble liquids (SL), aerosols, ultra-low volume concentrate solutions (ULV), soluble powders (SP), microencapsulation, water dispersed granules, flowables (FL), microemulsions (ME), nano-emulsions (NE), etc. In particular, the concentrate, powders, granules and emulsions may be freeze-dried. In any formulation described herein, percent of the active ingredient is within a range of 0.01% to 99.99%.

The compositions may be in the form of a liquid, gel or solid. Liquid compositions comprise pesticidal compounds derived from said *Burkholderia* strain, e.g. a strain having the identifying characteristics of *Burkholderia* A396 (NRRL Accession No. B-50319).

A solid composition can be prepared by suspending a solid carrier in a solution of pesticidal compounds and drying the suspension under mild conditions, such as evaporation at room temperature or vacuum evaporation at 65°C or lower.

A composition of the invention may comprise gel-encapsulated compounds derived from the *Burkholderia* strain of the present invention. Such gel-encapsulated materials can be prepared by mixing a gel-forming agent (e.g., gelatin, cellulose, or lignin) with a solution of pesticidal compounds used in the method of the invention; and inducing gel formation of the agent.

The composition may additionally comprise a surfactant to be used for the purpose of emulsification, dispersion, wetting, spreading, integration, disintegration control, stabilization of active ingredients, and improvement of fluidity or rust inhibition. In a particular embodiment, the surfactant is a non-phytotoxic non-ionic surfactant which preferably belongs to EPA List 4B. In another particular embodiment, the nonionic surfactant is polyoxyethylene (20) monolaurate. The concentration of surfactants may range between 0.1-35% of the total formulation, preferred range is 5-25%. The choice of dispersing and emulsifying agents, such as non-ionic, anionic, amphoteric and cationic dispersing and emulsifying agents, and the amount employed is determined by the nature of the composition and the ability of the agent to facilitate the dispersion of these compositions.

The composition may further comprise another microorganism and/or pesticide (e.g., nematocide, fungicide, insecticide). The microorganism may include but is not limited to an agent derived from *Bacillus* sp., *Pseudomonas* sp., *Brevibacillus* sp., *Lecanicillium* sp., non-*Ampelomyces* sp., *Pseudozyma* sp., *Streptomyces* sp, *Burkholderia* sp, *Trichoderma* sp, *Gliocladium* sp. Alternatively, the agent may be a natural oil or oil-product having fungicidal and/or insecticidal activity (e.g., paraffinic oil, tea tree oil, lemongrass oil, clove oil, cinnamon oil, citrus oil, rosemary oil).

The composition, in particular, may further comprise an insecticide. The insecticide may include but is not limited to avermectin, *Bacillus thuringiensis*, neem oil and

azadiractin, spinosads, *Chromobacterium subsugae*, eucalyptus extract, entomopathogenic bacterium or fungi such a *Beauveria bassiana*, and *Metarrhizium anisopliae* and chemical insecticides including but not limited to organochlorine compounds, organophosphorous compounds, carbamates, pyrethroids, and neonicotinoids.

5 The composition may further comprise a nematicide. The nematicide may include, but is not limited to chemical nematicides such as fenamiphos, aldicarb, oxamyl, carbofuran, natural product nematicide, avermectin, the fungi *Paecilomyces lilacinus* and *Muscodor spp.*, the bacteria *Bacillus firmus* and other *Bacillus spp.* and *Pasteuria penetrans*.

10 The composition may further comprise a biofungicide such as extract of *R. sachalinensis* (Regalia) or a fungicide. Such fungicides include, but are not limited to, a single site anti-fungal agent which may include but is not limited to benzimidazole, a demethylation inhibitor (DMI) (e.g., imidazole, piperazine, pyrimidine, triazole), morpholine, hydroxypyrimidine, anilinopyrimidine, phosphorothiolate, quinone outside inhibitor, quinoline, dicarboximide, carboximide, phenylamide, anilinopyrimidine, phenylpyrrole, aromatic
15 hydrocarbon, cinnamic acid, hydroxyanilide, antibiotic, polyoxin, acylamine, phthalimide, benzenoid (xylylalanine). In yet a further embodiment, the antifungal agent is a demethylation inhibitor selected from the group consisting of imidazole (e.g., triflumizole), piperazine, pyrimidine and triazole (e.g., bitertanol, myclobutanil, penconazole, propiconazole, triadimefon, bromuconazole, cyproconazole, diniconazole, fenbuconazole, hexaconazole, tebuconazole,
20 tetraconazole, propiconazole).

 The antimicrobial agent may also be a multi-site non-inorganic, chemical fungicide selected from the group consisting of a nitrile (e.g., chloronitrile or fludioxonil), quinoxaline, sulphamide, phosphonate, phosphite, dithiocarbamate, chloralkythios, phenylpyridin-amine,
25 cyano-acetamide oxime.

 The compositions may be applied using methods known in the art. Specifically, these compositions may be applied to plants or plant parts. Plants are to be understood as meaning in the present context all plants and plant populations such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants
30 can be plants which can be obtained by conventional plant breeding and optimization methods or by biotechnological and genetic engineering methods or by combinations of these methods, including the transgenic plants and including the plant cultivars protectable or not protectable by plant breeders' rights. Plant parts are to be understood as meaning all parts and organs of plants above and below the ground, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stalks, stems, flowers, fruit
35 bodies, fruits, seeds, roots, tubers and rhizomes. The plant parts also include harvested material, and vegetative and generative propagation material, for example cuttings, tubers, rhizomes, offshoots and seeds.

 Treatment of the plants and plant parts with the compositions set forth above may be

carried out directly or by allowing the compositions to act on their surroundings, habitat or storage space by, for example, immersion, spraying, evaporation, fogging, scattering, painting on, injecting. In the case that the composition is applied to a seed, the composition may be applied to the seed as one or more coats prior to planting the seed using one or more coats using methods known in the art.

As noted above, the compositions may be herbicidal compositions. The composition may further comprise one or more herbicides. These may include, but are not limited to, a bioherbicide and/or a chemical herbicide. The bioherbicide may be selected from the group consisting of clove, cinnamon, lemongrass, citrus oils, orange peel oil, tentoxin, cornexistin, AAL-toxin, leptospermone, thaxtomin, sarmentine, momilactone B, sorgoleone, ascaulatoxin and ascaulatoxin aglycone. The chemical herbicide may include, but is not limited to, diflufenzopyr and salts thereof, dicamba and salts thereof, topramezone, tembotrione, S-metolachlor, atrazine, mesotrione, primisulfuron-methyl, 2,4-dichlorophenoxyacetic acid, nicosulfuron, thifensulfuron-methyl, asulam, metribuzin, diclofop-methyl, fluazifop, fenoxaprop-p-ethyl, asulam, oxyfluorfen, rimsulfuron, mecoprop, and quinclorac, thiobencarb, clomazone, cyhalofop, propanil, bensulfuron-methyl, penoxsulam, triclopyr, imazethapyr, halosulfuron-methyl, pendimethalin, bispyribac-sodium, carfentrazone ethyl, sodium bentazon/sodium acifluorfen, glyphosate, glufosinate and orthosulfamuron.

Herbicidal compositions may be applied in liquid or solid form as pre-emergence or post-emergence formulations.

For pre-emergence dry formulations, the granule size of the carrier is typically 1-2 mm (diameter) but the granules can be either smaller or larger depending on the required ground coverage. Granules may comprise porous or non-porous particles.

For post-emergence formulations, the formulation components used may contain smectite clays, attapulgite clays and similar swelling clays, thickeners such as xanthan gums, gum Arabic and other polysaccharide thickeners as well as dispersion stabilizers such as nonionic surfactants (for example polyoxyethylene (20) monolaurate).

Uses

The compositions and pesticidal compounds derived from the *Burkholderia* strain set forth herein may be used as pesticides, particularly as insecticides, nematocides, fungicides and herbicides.

Specifically, nematodes that may be controlled using the method set forth above include but are not limited to parasitic nematodes such as root-knot, ring, sting, lance, cyst, and lesion nematodes, including but not limited to *Meloidogyne*, *Heterodera* and *Globodera* spp; particularly *Meloidogyne incognita* (root knot nematodes), as well as *Globodera rostochiensis* and *globodera pailida* (potato cyst nematodes); *Heterodera glycines* (soybean cyst nematode); *Heterodera schachtii* (beet cyst nematode); and *Heterodera avenae* (cereal

cyst nematode).

Phytopathogenic insects controlled by the method of the present invention include but are not limited to insects from the order (a) Lepidoptera, for example, *Acleris* spp., *Adoxophyes* spp., *Aegeria* spp., *Agrotis* spp., *Alabama argillaceae*, *Amylois* spp., *Anticarsia gemmatalis*, *Archips* spp., *Argyrotaenia* spp., *Autographa* spp., *Busseola fusca*, *Cadra cautella*, *Carposina nipponensis*, *Chilo* spp., *Choristoneura* spp., *Clysia ambiguella*, *Cnaphalocrocis* spp., *Cnephasia* spp., *Cochylis* spp., *Coleophora* spp., *Crocidolomia binotalis*, *Cryptophlebia leucotreta*, *Cydia* spp., *Diatraea* spp., *Diparopsis castanea*, *Earias* spp., *Ephestia* spp., *Eucosma* spp., *Eupoecilia ambiguella*, *Euproctis* spp., *Euxoa* spp., *Grapholita* spp., *Hedya nubiferana*, *Heliothis* spp., *Hellula undalis*, *Hyphantria cunea*, *Keiferia lycopersicella*, *Leucoptera scitella*, *Lithocollethis* spp., *Lobesia botrana*, *Lymantria* spp., *Lyonetia* spp., *Malacosoma* spp., *Mamestra brassicae*, *Manduca sexta*, *Operophtera* spp., *Ostrinia nubilalis*, *Pammene* spp., *Pandemis* spp., *Panolis flammea*, *Pectinophora gossypiella*, *Phthorimaea operculella*, *Pieris rapae*, *Pieris* spp., *Plutella xylostella*, *Prays* spp., *Scirpophaga* spp., *Sesamia* spp., *Sparganothis* spp., *Spodoptera* spp., *Synanthedon* spp., *Thaumetopoea* spp., *Tortrix* spp., *Trichoplusia ni* and *Yponomeuta* spp.;

(b) *Coleoptera*, for example, *Agriotes* spp., *Anthonomus* spp., *Atomaria linearis*, *Chaetocnema tibialis*, *Cosmopolites* spp., *Curculio* spp., *Dermestes* spp., *Diabrotica* spp., *Epilachna* spp., *Eremnus* spp., *Leptinotarsa decemlineata*, *Lissorhoptrus* spp., *Melolontha* spp., *Oryzaephilus* spp., *Otiorhynchus* spp., *Phlyctinus* spp., *Popillia* spp., *Psylliodes* spp., *Rhizopertha* spp., *Scarabeidae*, *Sitophilus* spp., *Sitotroga* spp., *Tenebrio* spp., *Tribolium* spp. and *Trogoderma* spp.; (c) *Orthoptera*, for example, *Blatta* spp., *Blattella* spp., *Gryllotalpa* spp., *Leucophaea maderae*, *Locusta* spp., *Periplaneta* spp. and *Schistocerca* spp.; (d) *Isoptera*, for example, *Reticulitermes* spp.; (e) *Psocoptera*, for example, *Liposcelis* spp.; (f) *Anoplura*, for example, *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Pemphigus* spp. and *Phylloxera* spp.; (g) *Mallophaga*, for example, *Damalinea* spp. and *Trichodectes* spp.; (h) *Thysanoptera*, for example, *Frankliniella* spp., *Hercinotrrips* spp., *Taeniothrips* spp., *Thrips palmi*, *Thrips tabaci* and *Scirtothrips aurantii*; (i) *Heteroptera*, for example, *Cimex* spp., *Distantiella theobroma*, *Dysdercus* spp., *Euchistus* spp., *Eurygaster* spp., *Leptocorisa* spp., *Nezara* spp., *Piesma* spp., *Rhodnius* spp., *Sahlbergella singularis*, *Scotinophara* spp. and *Tniatoma* spp.; (j) *Homoptera*, for example, *Aleurothrixus floccosus*, *Aleyrodes brassicae*, *Aonidiella* spp., *Aphididae*, *Aphis* spp., *Aspidiotus* spp., *Bemisia tabaci*, *Ceroplaster* spp., *Chrysomphalus aonidium*, *Chrysomphalus dictyospermi*, *Coccus hesperidum*, *Empoasca* spp., *Eriosoma larigerum*, *Erythroneura* spp., *Gascardia* spp., *Laodelphax* spp., *Lecanium corni*, *Lepidosaphes* spp., *Macrosiphus* spp., *Myzus* spp., *Nephotettix* spp., *Nilaparvata* spp., *Paratoria* spp., *Pemphigus* spp., *Planococcus* spp., *Pseudaulacaspis* spp., *Pseudococcus* spp., *Psylla* spp., *Pulvinaria aethiopica*, *Quadraspidiotus* spp., *Rhopalosiphum* spp., *Saissetia* spp., *Scaphoideus* spp., *Schizaphis* spp., *Sitobion* spp., *Trialeurodes vaporariorum*, *Trioza*

erytrae and *Unaspis citri*; (k) Hymenoptera, for example, *Acromyrmex*, *Atta* spp., *Cephus* spp., *Diprion* spp., *Diprionidae*, *Gilpinia polytoma*, *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis*, *Neodiprion* spp., *Solenopsis* spp. and *Vespa* spp.; (l) Diptera, for example, *Aedes* spp., *Antherigona soccata*, *Bibio hortulanus*, *Calliphora erythrocephala*,
 5 *Ceratitis* spp., *Chrysomyia* spp., *Culex* spp., *Cuterebra* spp., *Dacus* spp., *Drosophila melanogaster*, *Fannia* spp., *Gastrophilus* spp., *Glossina* spp., *Hypoderma* spp., *Hyppobosca* spp., *Liriomyza* spp., *Lucilia* spp., *Melanagromyza* spp., *Musca* spp., *Oestrus* spp., *Orseolia* spp., *Oscinella frit*, *Pegomyia hyoscyami*, *Phorbia* spp., *Rhagoletis pomonella*, *Sciara* spp., *Stomoxys* spp., *Tabanus* spp., *Tannia* spp. and *Tipula* spp.; (m) Siphonaptera, for example,
 10 *Ceratophyllus* spp. and *Xenopsylla cheopis* and (n) from the order Thysanura, for example, *Lepisma saccharina*. The active ingredients according to the invention may further be used for controlling crucifer flea beetles (*Phyllotreta* spp.), root maggots (*Delia* spp.), cabbage seedpod weevil (*Ceutorhynchus* spp.) and aphids in oil seed crops such as canola (rape), mustard seed, and hybrids thereof, and also rice and maize.

15 In a particular embodiment, the insect may be a member of the *Spodoptera*, more particularly, *Spodoptera exigua*, *Myzus persicae*, *Plutella xylostella* or *Euschistus* sp.

The substances and compositions may also be used to modulate emergence in either a pre-emergent or post-emergent formulation of monocotyledonous, sedge or dicotyledonous weeds. In a particular embodiment, the weeds may be *Chenopodium album*,
 20 *Abutilon theophrasti*, *Helianthus annuus*, *Ambrosia artemesifolia*, *Amaranthus retroflexus*, *Convolvulus arvensis*, *Brassica kaber*, *Taraxacum officinale*, *Solanum nigrum*, *Malva neglect.*, *Setaria lutescens*, *Bromus tectorum*, *Poa annua*, *Poa pratensis*, *Lolium perenne* L. var. Pace, *Festuca arundinaceae* Schreb. var. Aztec II, Anthem II, LS1100, *Echinochloa crus-galli*, *Lactuca sativa*. The *Burkholderia* strain, compounds and compositions set forth above may
 25 also be used as a fungicide. The targeted fungus may be a *Fusarium* sp., *Botrytis* sp., *Monilinia* sp., *Colletotrichum* sp., *Verticillium* sp.; *Microphomina* sp., *Phytophthora* sp., *Mucor* sp., *Podosphaera* sp., *Rhizoctonia* sp., *Peronospora* sp., *Geotrichum* sp., *Phoma*, and *Penicillium*. In another most particular embodiment, the bacteria are *Xanthomonas*.

30 The invention will now be described in greater detail by reference to the following non-limiting examples.

EXAMPLES

35 The compositions and methods set forth above will be further illustrated in the following, non-limiting Examples. The examples are illustrative of various embodiments only and do not limit the claimed invention regarding the materials, conditions, weight ratios, process parameters and the like recited herein.

1. Example 1. Isolation and identification of the microbe

1.1 Isolation of the microorganism

The microbe is isolated using established techniques known to the art from a soil sample collected under an evergreen tree at the Rinnoji Temple, Nikko, Japan. The isolation is done using potato dextrose agar (PDA) using a procedure described in detail by Lorch et al., 1995. In this procedure, the soil sample is first diluted in sterile water, after which it is plated in a solid agar medium such as potato dextrose agar (PDA). The plates are grown at 25°C for five days, after which individual microbial colonies are isolated into separate PDA plates. The isolated bacterium is gram negative, and it forms round, opaque cream-colored colonies that change to pink and pinkish-brown in color and mucoid or slimy over time.

1.2. Identification of the microorganism

The microbe is identified based on gene sequencing using universal bacterial primers to amplify the 16S rRNA region. The following protocol is used: *Burkholderia* sp A396 is cultured on potato-dextrose agar plates. Growth from a 24 hour-old plate is scraped with a sterile loop and re-suspended in DNA extraction buffer. DNA is extracted using the MoBio Ultra Clean Microbial DNA extraction kit. DNA extract is checked for quality/quantity by running 5 µl on a 1% agarose gel.

PCR reactions are set up as follows: 2 µl DNA extract, 5 µl PCR buffer, 1 µl dNTPs (10 mM each), 1.25 µl forward primer (27F; 5'-AGAGTTTGATCCTGGCTCAG-3' (SEQ ID NO:1), 1.25 µl reverse primer (907R; 5'-CCGTCAATTCCTTTGAGTTT-3' (SEQ ID NO:2)) and 0.25 µl Taq enzyme. The reaction volume is made up to 50 µl using sterile nuclease-free water. The PCR reaction includes an initial denaturation step at 95°C for 10 minutes, followed by 30 cycles of 94°C/30 sec, 57°C/20 sec, 72°C/30 sec, and a final extension step at 72°C for 10 minutes.

The product's approximate concentration and size is calculated by running a 5 µl volume on a 1% agarose gel and comparing the product band to a mass ladder.

Excess primers, dNTPs and enzyme are removed from the PCR product with the MoBio PCR clean up kit. The cleaned PCR product is directly sequenced using primers 27F (same as above), 530F (5'-GTGCCAGCCGCGCGG-3' (SEQ ID NO:3)), 1114F (5'-GCAACGAGCGCAACCC (SEQ ID NO:4)) and 1525R (5'-AAGGAGGTGWTCARCC-3' (SEQ ID NO:5)), 1100R (5'-GGGTTGCGCTCGTTG-3' (SEQ ID NO:6)), 519R (5'-GWATTACCGCGGCKGCTG-3' (SEQ ID NO:7)).

The 16S rRNA gene sequence of strain A396 is compared with the available 16S rRNA gene sequences of representatives of the β-proteobacteria using BLAST. Strain A396 is closely related to members of the *Burkholderia cepacia* complex, with 99% or higher similarity to several isolates of *Burkholderia multivorans*, *Burkholderia vietnamensis*, and *Burkholderia cepacia*. A BLAST search excluding the *B. cepacia*

complex, showed 98% similarity to *B. plantarii*, *B. gladioli* and *Burkholderia sp.* isolates.

A distance tree of results using the neighbor joining method, showed that A396 is related to *Burkholderia multivorans* and other *Burkholderia cepacia* complex isolates. *Burkholderia plantarii* and *Burkholderia glumae* grouped in a separate branch of the tree.

5 The isolated *Burkholderia* strain was found to contain the following sequences:
forward sequence, DNA sequence with 27F primer, 815 nucleotides (SEQ ID NO:8);
reverse sequence, 1453 bp, using primers 1525R, 1100R, 519R (SEQ ID NO:9);
reverse sequence 824 bp using primer 907R (SEQ NO: 10); forward sequence 1152 bp
using primer 530F (SEQ ID NO:11); forward sequence 1067 bp using 1114F primer (SEQ
10 ID NO:12); reverse sequence 1223 bp using 1525R primer (SEQ NO:13); reverse sequence
1216 bp using 1100R primer (SEQ ID NO:14); reverse sequence 1194 bp using 519R
primer (SEQ ID NO:15).

1.3. Proof that *Burkholderia* A396 does not belong to *Burkholderia cepacia* complex

15 1.3.1 Molecular Biology work using specific PCR primers

In order to confirm the identification of *Burkholderia* A396 as *Burkholderia multivorans*, additional sequencing of housekeeping genes is performed. *Burkholderia multivorans* is a known member of the *Burkholderia cepacia* complex. Efforts are focused on PCR of *recA* genes, as described by Mahenthiralingam et al., 2000. The following
20 primers are used: (a) BCR1 and BCR2 set forth in Mahenthiralingam et al., 2000 to confirm *B. cepacia* complex match and (b) BCRBM1 and BCRBM2 set forth Mahenthiralingam et al, 2000 to confirm *B. multivorans* match. A product-yielding PCR reaction for the first primer set would confirm that the microbe belongs to the *B. cepacia* complex. A product-yielding PCR reaction for the second primer set would confirm that the microbe is indeed *B.*
25 *multivorans*.

No PCR product is obtained for either pair of primers. The performance of the PCR reaction and primers is tested using *Burkholderia multivorans* ATCC 17616 (positive control) and *Pseudomonas fluorescens* (negative control). Strong bands are observed both for *B. multivorans* using both sets of primers. No bands are observed for *Pseudomonas fluorescens*. The results indicate that A396 is a *Burkholderia*, but not a member of the *B. cepacia* complex, and not *Burkholderia multivorans*. This is also demonstrated in a comparative culture experiment in which both A396 and a type culture of *B. multivorans* are grown side-by-side in a shake culture, and the growth is monitored daily using optical density measurements at 600 nm. Under the set conditions, the novel species A396 grew
30 much faster than the *B. multivorans* type strain (Figure 1).

1.3.2 DNA-DNA Hybridization

In order to confirm that isolate A396 is a new species of *Burkholderia*, a DNA-DNA hybridization experiment with *Burkholderia multivorans* (the closest 16S rRNA sequence

match) is conducted. Biomass for both A396 and *B. multivorans* is produced in ISP2 broth, grown over 48 hours at 200 rpm/25°C in Fernbach flasks. The biomass is aseptically harvested by centrifugation. The broth is decanted and the cell pellet is resuspended in a 1:1 solution of water: isopropanol. DNA-DNA hybridization experiments are performed by the DSMZ, the German Collection of Microorganisms and Cell Cultures in Germany. DNA is isolated using a French pressure cell (Thermo Spectronic) and is purified by chromatography on hydroxyapatite as described by Cashion et al., 1977. DNA-DNA hybridization is carried out as described by De Ley et al., 1970 under consideration of the modifications described by Huss et al., 1983 using a model Cary 100 Bio UV/VIS-spectrophotometer equipped with a Peltier thermostatted 6x6 multicell changer and a temperature controller with *in-situ* temperature probe (Varian). DSMZ reported % DNA-DNA similarity between A396 and *Burkholderia multivorans* of 37.4%. The results indicate that *Burkholderia sp* strain A396 does not belong to the species *Burkholderia multivorans* when the recommendations of a threshold value of 70% DNA-DNA similarity for the definition of bacterial species by the ad hoc committee (Wayne et al., 1987) are considered.

1.4. Biochemical profile using Biolog GN2 plates

For the carbon source utilization profile, A396 is grown overnight on Potato Dextrose Agar (PDA). The culture is transferred to BUG agar to produce an adequate culture for Biolog experiments as recommended by the manufacturer (Biolog, Hayward, CA).

The biochemical profile of the microorganism is determined by inoculating onto a Biolog GN2 plate and reading the plate after a 24-hour incubation using the MicroLog 4-automated microstation system. Identification of the unknown bacteria is attempted by comparing its carbon utilization pattern with the Microlog 4 Gram negative database.

No clear definitive matches are found to the Biolog profile. The closest matches all had less than 35% similarity with A396: *Pseudomonas spinosa* (*Burkholderia*), *Burkholderia cepacia*, and *Burkholderia pseudomallei*. The results are shown in Table I.

Table 1. Biochemical Profile of A396

Substrate	Result	Substrate	Result
Cyclodextrin	-	L-arabinose	-
Dextrin	-	D-arabitol	-
Glycogen	-	D-cellobiose	-
Tween 40	+	Erythritol	-
Tween 80	+	D-Fructose	-
N-acetyl-D-Galactoseamine	-	L-Fucose	-
N-acetyl-D-glucosamine	-	D-Galactose	+/-
Adonitol	-	Gentibiose	-
Succinic Acid Mon-methyl ester	-	D-Glucose	+
Acetic acid	-	m-Inositol	-
Cis-aconitic acid	-	D-Lactose	-
Citric acid	-	Lactulose	-
Formic acid	+	Maltose	-
D-Galactonic Acid Lactone	-	D-Mannitol	-
D-Galacturonic Acid	-	D-Mannose	-
D-Gluconic acid	-	D-Melibiose	-
D-Glucosaminic acid	-	β -methyl-D-glucoside	-
D-Glucuronic Acid	-	D- Psicose	-
α -hydroxybutyric acid	-	D-Raffinose	-
β -hydroxybutyric acid	+	L-Rhamnose	-
γ -hydroxybutyric acid	-	D-Sorbitol	-
p-hydroxyphenylacetic acid	-	Sucrose	-
Itaconic acid	-	D-Trehalose	+
α -keto butyric acid	-	Turanose	-
α -keto glutaric acid	-	Xylitol	-
α -ket valeric acid	-	Pyruvic Acid Methyl ester	-
D,L-Lactic acid	-	Uridine	-
Malonic acid	-	Thymidine	-
Propionic acid	+	Phenyethyl-amine	-
Quinic acid	-	Putrescine	-
D-Saccharic acid	-	2-aminoethanol	-
Sebacic acid	-	2,3-Butanediol	-
Succinic Acid	+	Glycerol	+/-
Bromosuccinic acid	-	D,L-a-glycerol phosphate	+/-
Succinamic acid	-	α -D-Glucose-1-phosphate	-
Glucuronamide	-	D-glucose-6-phosphate	+
L-alaninamide	+	γ -amino butyric acid	+
D-Alanine	-	Urocanic acid	-
L-alanine	+	Inosine	-
L-alanyl-glycine	-	L-phenylalanine	+
L-asparagine	+	L-proline	-
L-aspartic acid	+/-	L-pyroglutamic acid	-
L-glutamic acid	+	D-serine	-
Glycyl-L-Aspartic acid	-	L-serine	-
Glycyl-L-glutamic acid	-	L-threonine	-
L-histidine	-	D,L-carnitine	-
Hydroxy-L-proline	+	L-ornithine	-
L-leucine	-	-	-

1.5. Fatty acid composition

After incubation for 24 hours at 28°C, a loopful of well-grown cells are harvested
 5 and fatty acid methyl esters are prepared, separated and identified using the Sherlock

Microbial Identification System (MIDI) as described (see Vandamme et al., 1992). The predominant fatty acids present in the *Burkholderia* A396 are as follows: 16:0 (24.4%), cyclo 17:0 (7.1%), 16:0 3-OH (4.4%), 14:0 (3.6%), 19:0 ω 8c (2.6%) cyclo, 18:0 (1.0%).

Summed feature 8 (comprising 18:1 ω 7c) and summed feature 3 (comprising of 16:1 ω 7c and 16:1 ω 6c) corresponded to 26.2% and 20.2 % of the total peak area, respectively.

Summed feature 2 comprising 12:0 ALDE, 16:1 iso I, and 14:0 3-OH) corresponded to 5.8% of the total peak area while summed feature 5 comprising 18:0 ANTE and 18:2 ω 6,9c corresponded to 0.4%. Other fatty acids detected in A396 in minor quantities included: 13:1

at 12-13 (0.2%), 14:1 ω 5c (0.2%), 15:0 3-OH (0.13%), 17:1 ω 7c (0.14%), 17:0 (0.15%), 16:0 iso 3-OH (0.2%), 16:0 2-OH (0.8%), 18:1 ω 7c 11-methyl (0.15%), and 18:1 2-OH (0.4%).

A comparison of the fatty acid composition of A396 with those of known microbial strains in the MIDI database suggested that the fatty acids in the novel strain A396 were most similar with those of *Burkholderia cenocepacia*.

1.6 Resistance to Antibiotics

Antibiotic susceptibility of *Burkholderia A396* is tested using antibiotic disks on Muller-Hinton medium as described in PML Microbiological's technical data sheet #535. Results obtained after 72-hour incubation at 25°C are presented in Table 2 below.

Table 2: Susceptibility of MBI-206 to various antibiotics. +++ very susceptible, ++ susceptible, - resistant

	Concentration (ug)	Susceptible
Tetracycline	30	-
Kanamycin	30	+++
Erythromycin	15	-
Streptomycin	10	-
Penicillin	10	-
Ampicillin	10	-
Oxytetracycline	30	-
Chloramphenicol	30	++
Ciprofloxacin	5	++
Gentamicin	10	-
Piperacillin	100	+++
Cefuroxime	30	-
Imipenem	10	+++
Sulphamethoxazole- Trimethoprim	23.75/ 25	++

5

The results indicate that the antibiotic susceptibility spectrum of *Burkholderia A396* is quite different from pathogenic *B. cepacia* complex strains. *Burkholderia A396* is susceptible to kanamycin, chloramphenicol, ciprofloxacin, piperacillin, imipenem, and a combination of sulphamethoxazole and trimethoprim. As a comparison, Zhou et al., 2007 tested the susceptibility of 2,621 different strains in *B. cepacia* complex isolated from cystic fibrosis patients, and found that only 7% and 5% of all strains were susceptible to imipenem or ciprofloxacin, respectively. They also found 85% of all strains to be resistant to chloramphenicol (15% susceptible), and 95% to be resistant (5% susceptible) to the combination of sulphamethoxazole and trimethoprim. Results of Zhou et al., 2007 are similar to those of Pitt et al., 1996 who determined antibiotic resistance among 366 *B. cepacia* isolates and reported that most of them are resistant to ciprofloxacin, cefuroxime, imipenem, chloramphenicol, tetracycline, and sulphamethoxazole.

10

15

2. Example 2. *Burkholderia* sp. as an Herbicide

20

2.1 Study #1

To confirm the activity found in the initial herbicide screen, an *in vivo* study is conducted using the Amberlite 7 XAD resin extract derived from a 5-day old whole cell

broth of the novel *Burkholderia* species. The dried crude extract is resuspended in 4% ethanol and 0.2 % non-ionic surfactant (glycosperse) at a concentration of 10 mg/mL, and further diluted to a concentration of 5.0 mg/mL. The two samples are sprayed on 4-week old plants of bindweed (*Convolvulus arvensis*), and the plants are kept under growth lights at 25°C for 2 weeks, at which point, the phytotoxicity evaluations are performed. In the same study, 2-week old redroot pigweed plants are sprayed with increasing concentrations of the crude extract derived from the bacterial culture. The test concentrations are 1.25, 2.5, 5.0 and 10.0 mg/mL, and the plants are incubated as described above before phytotoxicity evaluations.

Results presented in Figures 2 (bindweed) and 3 (pigweed) show the phytotoxic effect of *Burkholderia* crude extract at different concentrations, and they show good herbicidal effect on pigweed even at low treatment concentrations. Both extract treatments (5 and 10 mg/mL) result in stunting on bindweed.

2.2 Study #2

A novel strain of *Burkholderia* sp. A396 is grown in an undefined mineral medium for 5 days (25°C, 200 rpm). The whole cell broth is extracted using XAD7 resin. The dried crude extract is resuspended in 4% ethanol and 0.2 % non-ionic surfactant at a concentration of 10 mg/mL, and further diluted to concentrations of 5.0, 2.5, and 1.25 mg/mL. All four test solutions are then tested on the following broadleaf and grass weed species listed in Table 3:

Table 3. Broadleaf and Grass Weed Species Tested

Common Name	Scientific Name
Lambsquarter	<i>Chenopodium album</i>
Horseweed	<i>Conyza canadensis</i>
Curlydock	<i>Rumex crispus</i>
Crabgrass	<i>Digitaria sanguinalis</i>
Bluegrass	<i>Poa annua</i>
Dandelion	<i>Taraxacum officinale</i>
Nightshade	<i>Solanum nigrum</i>
Mustard	<i>Brassica kaber</i>
Mallow	<i>Malva neglecta</i>
Cocklebur	<i>Xanthium pensylvanicum</i>
Bermuda Grass	<i>Cynodon dactylon</i>
Foxtail	<i>Setaria lutescens</i>
Sowthistle	<i>Sonchus oleraceus</i>

A solution of 0.2 % glycosperse and Roundup at 6 fl oz per gallon rate is used as negative and positive controls, respectively.

All plant species are tested in 4"x4" plastic pots in three replicates. The untreated control plants are sprayed with the carrier solution (4% Ethanol, 0.2% glycosperse) and the positive control plants with Roundup at a rate corresponding to 6 fl. oz/acre. Treated plants are kept in a greenhouse under 12h light/12h dark conditions. Phytotoxicity data taken 22 days after treatment for species #1-8 and 12 days for species #9-12 are presented in Tables 5

and 6, respectively. The rating scale for both tables is shown in Table 4:

Table 4. Rating Scale

Rating Scale	% Control
0	0
1	<10
2	25
3	50
4	75
5	100

5 **Table 5. Phytotoxicity Data for Species #1-8**

Treatment	Horseweed	Lambsquarter	Dandelion	Curlydock	Crabgrass	Mustard	Nightshade	Bluegrass
UTC	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0
1.25 mg/mL	0.0	4.7	0.0	0.0	0.0	4.3	0.0	0.0
2.5 mg/mL	0.7	4.5	0.0	0.0	0.0	4.7	0.0	0.0
5.0 mg/mL	4.3	5.0	0.0	0.0	0.0	5.0	0.0	0.0
10.0 mg/mL	4.7	5.0	0.0	0.0*	0.0	5.0	1.5	0.0
Roundup	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0

* stunting that resulted in plants approximately half the size of untreated plants

Table 6. Phytotoxicity Data for Species #9-12

Treatment	Cocklebur	Foxtail	Bermuda Grass	Sowthistle	Mallow
UTC	0.0	0.7	0.0	0.0	2.8
1.25 mg/mL	0.5	0.3	0.3	0.0	2.0
2.5 mg/mL	0.5	0.7	0.5	0.0	2.7
5.0 mg/mL	0.8	0.3	0.2	0.0	2.2
10.0 mg/mL	0.7	0.7	0.3	0.2	1.7
Roundup	4.7	4.8	4.7	5.0	5.0

10

Based on the results obtained in these studies, the compounds extracted from fermentation broths of the isolated *Burkholderia* species had herbicidal activity against several weed species are tested. Of the twelve species tested, Lambsquarters and mustard are most susceptible, followed by mallow and horseweed. Extract concentration as low as 1.25 mg/mL is able to provide almost complete control of Lambsquarters and mustard, whereas higher concentration is required for the mallow and horseweed.

15

In a separate experiment, using the same design as described above, systemic activity is tested. A 10 mg/ml crude extract supernatant of *Burkholderia sp. A396* is painted onto first true leaves of Ragweed, Mustard, Nightshade, Crabgrass, Wheat and Barnyard Grass. Seedlings are evaluated 7 days after treatment. Observed symptoms include: burning, warping, bleaching Herbicidal activity is observed in the next leaf above the treated leaf in Ragweed, Mustard and Nightshade. No systemic activity is observed in the

20

tested grasses. In a second experiment, five fractions of the same crude extract (10 mg/ml) are evaluated using the same experimental design as described above. Seedlings of Mustard, Wheat and Crabgrass are treated. Seven and 20 days after treatment, symptoms of herbicidal activity are observed in Mustard from four out of the five fractions (091113B4F6, 091113B4F7, 091113B4F8 and 091113B4F9) using a C-18 column (Phenomenex Septra C18-E, 50 μ m, 65Å). Symptoms are observed in the next leaf above the treated leaf. No systemic activity is observed in the tested grasses.

3. Example 3. *Burkholderia* sp. as an Insecticide

3.1. Contact Activity Studies

The following assay is used in the initial screening phase to determine if the compounds derived from a culture of the novel *Burkholderia* species has contact activity against a Lepidopteran pest (larvae). It is further used as a tool for the bioassay-guided fractionation to determine the active fractions and peaks derived from the whole-cell-broth extract. The test is conducted in individual 1.25 oz plastic cups using either Cabbage looper (*Trichoplusia ni*) late third instar larvae or Beet Armyworm (*Spodoptera exigua*) early third instar larvae. A 1cm x 1cm piece of solid Beet armyworm diet is placed in the center of each cup together with one larvae. A 1 μ l aliquot of each treatment (whole cell broth or extract from a 5-day-old *Burkholderia* A396 culture) is injected on each larvae thorax (dorsal side) using a Hamilton Precision Syringe. Each treatment is replicated ten times. Water is used as a negative control treatment and malathion as the positive control treatment. After injection, each cup is covered with parafilm with an airhole, and the cups are incubated for three days at 26°C. Mortality evaluations are done daily, starting 24 hours after the treatment.

Figures 4 and 5 present the results from contact activity tests. According to the results, the filter-sterilized broth from a *Burkholderia* sp culture killed about 40% of all test insects within 3 days. Diluted broth (50%) has lower activity, resulting in about 10% control in both insects tested.

3.2. Activity Against Larvae Through Feeding

Direct toxicity via feeding is tested using the diet-overlay tests with following 96-well plate assay format using microtiter plates with 200 μ l of solid, artificial Beet Armyworm diet in each well. One hundred (100) microliters of each test sample is pipetted on the top of the diet (one sample in each well), and the sample is let dry under flowing air until the surface is dry. Each sample (filter-sterilized through a 0.2 micron filter) is tested in six replicates, and water and a commercial Bt (*B. thuringiensis*) product are used as negative and positive controls, respectively. One third instar larvae of the test insect (Cabbage looper-*Trichoplusia ni*; Beet armyworm – *Spodoptera exigua*; Diamondback

Moth – *Plutella xylostella*) is placed in each well, and the plate is covered with plastic cover with airholes. The plates with insects are incubated at 26°C for 6 days with daily mortality evaluations.

Figure 5 represents data from a diet overlay study with Beet Armyworm

- 5 (*Spodoptera exigua*) early third instar larvae tested at four different broth concentrations: 1x (100%), 1/4x (25%), 1/8x (12.5%), 1/16x (6.125%). The data shows that the undiluted, filter-sterilized broth is able to give 100% control at the end of the 7-day incubation period. Similar control is obtained with a 4-fold dilution of the broth, and in the end of the study, both undiluted and 4-fold diluted broths are comparable to Bt used as a positive control.
- 10 However, the effect of Bt is significantly faster than that of the *Burkholderia* broths. Efficacy against armyworm larvae is dependent on broth concentration, and the two lowest broth concentrations (12.5% and 6.125%) provided less control than the two highest ones. However, the performance of the 12.5% dilution is not much lower than the 25% dilution. The 16-fold dilution of broth is clearly not efficient enough, and it only provided partial
- 15 (33%) control of armyworm larvae during this 7-day study. The corresponding mortality rates for the same broth dilution used on cabbage loopers and diamondback moth larvae are a little higher with 6.125% broth killing 80% and 50% of larvae, respectively.

3.3. *In vitro* activity against sucking insects

- 20 Five stinkbug (*Euschistus* sp.) adults are placed in each 16 oz plastic container lined with a piece of paper towel. A microcentrifuge tube containing 2 mL of each test sample (filter sterilized whole broth) is capped with a cotton ball, and laid down on the bottom of the plastic container. One sunflower seed is placed next to the tube as bait. Water and a commercial product with a mixture of pyrethrin and PBO at a recommended rate are used as
- 25 negative and positive controls, respectively. Each container is closed with a lid, and they are incubated at 25°C for 7 days with daily mortality checks.

- Results are presented below in Table 7 and they show about 80% control of sucking insect (stinkbug) by day 7 in this *in vitro* system with 50% diluted broth. In this study, the diluted fermentation broth of *Burkholderia* A396 is more effective in controlling stinkbugs
- 30 than the commercial product used as a positive control. Interestingly, the non-diluted broth resulted in lower insect control, which might be an indication of antifeedant (feeding inhibition) properties of the active secondary metabolites produced by this new species of *Burkholderia*.

35 **Table 7. Effect of A396 on Stinkbugs**

Treatment	% control (Day 3)	% control (Day 5)	% control (Day 7)
A396 undil. broth (1x)	0	0	40
A396 broth dil. 50% (0.5x)	20	20	80
Pyrethrin+PBO (pos control)	0	0	40
Water (neg control)	0	0	0

4. Example 4. Sucking insect test *in vivo*

The *in vivo* efficacy of the filtered whole cell broth is tested in a plant assay with mustard plants and green peach aphid (*Myzus persicae*) as the test insect. Approximately one-month-old Florida Broadleaf mustard (*Brassica* sp.) plants are sprayed with two different concentrations (1x and 0.5x) of the filter sterilized whole cell broth of *Burkholderia* sp. using a Paasche airbrush. Water and a commercial product of avermectin (Avid) are used as negative and positive controls, respectively. The plants are allowed to dry on the benchtop, after which they are placed in a 6-cup plastic container with a lid with airholes. Ten aphids at various developmental stages are placed on each test plant, and the plants are incubated under growth lamps for 7 days at 25°C. Daily evaluations for the number of aphids on each plant (summarized in table Table 8 below) are made and recorded in a notebook.

Table 8. In vivo Efficacy of A396 on Green Peach Aphids

Treatment	# live aphids Day 0	# live aphids Day 2	# live aphids Day 4	# live aphids Day 7
A396 undiluted broth (1x)	10	36	88	145
A396 broth diluted (0.5x)	10	47	138	217
Avermectin (pos control)	10	0	0	0
Water (neg control)	10	140	364	393

According to the results, both concentrations of the filter-sterilized broth derived from a culture of a novel species of *Burkholderia* are able to control the population growth of a sucking insect, *M. persicae*.

5. Example 5. Nematocidal Activity

5.1 Study #1

To assess the effect of filter-sterilized *Burkholderia* sp A396 culture broth on the motility (and subsequent recovery) of juvenile (J2) root-knot nematodes (*Meloidogyne incognita* VW6), the following test is conducted on 24-well plastic cell-culture plates:

A 300- μ l aliquot of each test solution (either 1x or 0.5x filter-sterilized broth) is added into appropriate wells after which, fifteen nematodes dispensed in 10 μ l of DI water are added into each well, plate is closed with a lid, and incubated at 25°C for 24 hours. Water and Avid at 20,000x dilution are used as negative and positive controls, respectively. Effect of each compound on nematode mobility is checked after 24 hours by probing each nematode with a needle, and the proportion of immobile nematodes in each treatment is recorded in a notebook using a % scale. To assess the recovery of mobility in each treatment, a volume of 200 μ l is removed from each well, and the remaining solution in each well is diluted by adding 2 mL of DI water. Plates are again incubated for 24 hours as

described above, after which the second mobility evaluation is performed.

The results presented in Figure 6 show the filter-sterilized broth at both test concentrations can immobilize the free-living juvenile root-knot nematodes. This effect lasts at least for 24-hours, which suggests that *Burkholderia A396* broth can be used to prevent plants from nematode infections.

5.2 Study #2

Materials and Methods

Mini Drench Test: *Burkholderia A396* whole cell broth is tested in a greenhouse assay conducted in 45 ml pots. Cucumber seeds cv. Toshka are sown directly into pots filled with a sandy loam soil. Ten days later, pots were each treated with 5 ml of a suspension. Specific amounts used are shown in Table 9:

Table 9

Compounds	<i>Burkholderia</i> strain A396 Fosthiazate (Standard, EC 150) (positive control)
Test species	<i>Meloidogyne sp.</i> applied at 3000 eggs per mini drench pot (in 2 ml)
Test plant	<i>Cucumis sativus</i> (cucumber cv. Toschka)
Test formulation	100% liquid formulation
Test concentrations	100, 50, 25, 12.5, 6, 3, 1.5 ml/L
Test application	Drench application

As indicated in Table 9, pots are inoculated with 3000 eggs of *M. incognita*. Four replicates were prepared for each treatment and rate. The trial was harvested fourteen days after trial application and inoculation. Root galling was assessed according to Zeck's gall index (Zeck, 1971). Phytotoxicity was measured as a reduction of root galling in comparison to the control. The results are shown in Figures 9 and 10.

In *Mini Drench Test no. 1* (see Figure 9), the activity of the treatment was very high and a reduction of almost 100% was observed when applied at a concentration of 100 ml/L *Burkholderia A396*. Fosthiazate performed as usual (100% control at 20 ppm).

In *Mini Drench Test no. 2* (see Figure 10) a 100% reduction of root galling was achieved at the highest concentration of 100 ml/L dropping to approximately 50% at 1.5 ml/L. Fosthiazate performed as usual (100% control at 20 ppm).

5.3 Study #3

To demonstrate the nematicidal activity of *Burkholderia A396*, a greenhouse study on cucumber (*Cucumis sativus*) is performed using a whole cell broth of *Burkholderia A396* as the test product to control root knot nematodes (*Meloidogyne incognita*). One cucumber plant per pot is planted in soil and grown in a greenhouse under artificial lights at 28°C. Each pot with a plant is treated with an aliquot (about 80 mL) of either the undiluted test product or a test product diluted to 5% with water. Each *Burkholderia A396* treatment as

well as a positive control treatment with Temik (at a label rate) and a negative control with no additions consisted of five replicates. Plants are grown in a greenhouse for 60 days, after which each plant was harvested and evaluated for fresh shoot and root weights. Number of nematode eggs in each pot was recorded and a parameter indicating the number of eggs per a gram of root mass was calculated. Statistical analysis (ANOVA) is performed, and the statistical differences among treatment means at $p < 0.1$ was calculated. Data presented in Table 10 below shows that even though not statistically different from the untreated control, the pots treated with A396 whole cell broth contained less nematode eggs than the untreated control pots. The effect calculated as number of eggs per root mass is more clear when undiluted broth is used as a treatment.

Table 10. The effect of A396 whole cell broth on the cucumber shoot and root weight, total number of *M. incognita* eggs per pot and the number of eggs per gram of root mass.

	shoot fresh wt		root fresh wt		# of eggs		# of eggs/g of root	
untreated	15.22	b	11.76	bc	67693	a	5252.0	ab
A396 5% v/v	11.89	b	6.914	c	56084	a	8419.4	a
A 396 undiluted	15.66	b	11.09	bc	40463	a	3929.2	ab
Temik 15 G 5 lb/a	29.54	a	29.74	a	68907	a	2604.4	b
LSD at $p < 0.1$	5.34		6.9879		36509.2		3317.07	

6. Example 6. Isolation of Templazole A and B

Methods and Materials

The following procedure is used for the purification of Templazole A and B extracted from cell culture of *Burkholderia sp* (see Figure 7):

The culture broth derived from the 10-L fermentation *Burkholderia* (A396) in Hy soy growth medium is extracted with Amberlite XAD-7 resin (Asolkar et al., 2006) by shaking the cell suspension with resin at 225 rpm for two hours at room temperature. The resin and cell mass are collected by filtration through cheesecloth and washed with DI water to remove salts. The resin, cell mass, and cheesecloth are then soaked for 2 h in acetone after which the acetone is filtered and dried under vacuum using rotary evaporator to give the crude extract. The crude extract is then fractionated by using reversed-phase C18 vacuum liquid chromatography (H_2O/CH_3OH ; gradient 90:20 to 0:100%) to give 10 fractions. These fractions are then concentrated to dryness using rotary evaporator and the resulting dry residues are screened for biological activity using 96 well plate lettuce seeding assay. The active fractions are then subjected to reversed phase HPLC (Spectra System

P4000 (Thermo Scientific) to give pure compounds, which are then screened in above mentioned bioassays to locate/identify the active compounds. To confirm the identity of the compound, additional spectroscopic data such as LC/MS and NMR is recorded.

The active fraction 4 is purified further by using HPLC C-18 column (Phenomenex, Luna 10u C18(2) 100 A, 250 x 30), water:acetonitrile gradient solvent system (0-10 min; 80% aqueous CH₃CN, 10-25 min; 80 - 65% aqueous CH₃CN, 25-50 min; 65 - 50 % aqueous CH₃CN, 50-60 min; 50-70% CH₃CN, 60-80 min; 70-0% aqueous CH₃CN, 80-85 min; 0 - 20% aqueous CH₃CN) at 8 mL/min flow rate and UV detection of 210 nm, to give templazole B, retention time 46.65 min. The other active fraction 6 is also purified using HPLC C-18 column (Phenomenex, Luna 10u C18(2) 100 A, 250 x 30), water:acetonitrile gradient solvent system (0-10 min; 80 % aqueous CH₃CN, 10-25 min; 80 - 60 % aqueous CH₃CN, 25-50 min; 60 - 40% aqueous CH₃CN, 50-60 min; 40% CH₃CN, 60-80 min; 40-0% aqueous CH₃CN, 80-85 min; 0-20 % aqueous CH₃CN) at 8 mL/min flow rate and UV detection of 210 nm, to give templazole A, retention time 70.82 min.

Mass spectroscopy analysis of pure compounds is performed on a Thermo Finnigan LCQ Deca XP Plus electrospray (ESI) instrument using both positive and negative ionization modes in a full scan mode (m/z 100-1500 Da) on a LCQ DECA XP^{plus} Mass Spectrometer (Thermo Electron Corp., San Jose, CA). Thermo high performance liquid chromatography (HPLC) instrument equipped with Finnigan Surveyor PDA plus detector, autosampler plus, MS pump and a 4.6 mm x 100 mm Luna C18 5 μ m column (Phenomenex). The solvent system consists of water (solvent A) and acetonitrile (solvent B). The mobile phase begins at 10% solvent B and is linearly increased to 100% solvent B over 20 min and then kept for 4 min, and finally returned to 10% solvent B over 3 min and kept for 3 min. The flow rate is 0.5 mL/min. The injection volume was 10 μ L and the samples are kept at room temperature in an auto sampler. The compounds are analyzed by LC-MS utilizing the LC and reversed phase chromatography. Mass spectroscopy analysis of the present compounds is performed under the following conditions: The flow rate of the nitrogen gas was fixed at 30 and 15 arb for the sheath and aux/sweep gas flow rate, respectively. Electrospray ionization was performed with a spray voltage set at 5000 V and a capillary voltage at 35.0 V. The capillary temperature was set at 400°C. The data was analyzed on Xcalibur software. The active compound templazole A has a molecular mass of 298 and showed m/z ion at 297.34 in negative ionization mode. The LC-MS chromatogram for templazole B suggests a molecular mass of 258 and exhibited m/z ion at 257.74 in negative ionization mode.

^1H , ^{13}C and 2D NMR spectra were measured on a Bruker 500 MHz & 600 MHz gradient field spectrometer. The reference is set on the internal standard tetramethylsilane (TMS, 0.00 ppm).

For structure elucidation of templazole A, the purified compound with a molecular weight 298 is further analyzed using a 500 MHz NMR instrument, and has ^1H NMR δ values at 8.44, 8.74, 8.19, 7.47, 7.31, 3.98, 2.82, 2.33, 1.08 and has ^{13}C NMR δ values of 163.7, 161.2, 154.8, 136.1, 129.4, 125.4, 123.5, 123.3, 121.8, 121.5, 111.8, 104.7, 52.2, 37.3, 28.1, 22.7, 22.7. Templazole A has UV absorption bands at 226, 275, 327 nm, which suggested the presence of indole and oxazole rings. The molecular formula, $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_3$, was determined by interpretation of ^1H , ^{13}C NMR and HRESI MS data m/z 299.1396 ($\text{M}+\text{H}$) $^+$ (Calcd for $\text{C}_{17}\text{H}_{19}\text{N}_2\text{O}_3$, 299.1397), which entails a high degree of unsaturation shown by 10 double bond equivalents. The ^{13}C NMR spectrum revealed signals for all 17 carbons, including two methyls, a methoxy, a methylene carbon, an aliphatic methine, an ester carbonyl, and eleven aromatic carbons. The presence of 3'-substituted indole was revealed from ^1H - ^1H COSY and HMBC spectral data. The ^1H - ^1H COSY and HMBC also indicated the presence of a carboxylic acid methyl ester group and a $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$ side chain. From the detailed analysis of ^1H - ^1H COSY, ^{13}C , and HMBC data it was derived that the compound contained an oxazole nucleus. From the 2D analysis it was found that the isobutyl side chain was attached at C-2 position, a carboxylic acid methyl ester at C-4 position and the indole unit at C-5 position to give templazole A.

The second herbicidally active compound, templazole B, with a molecular weight 258 is further analyzed using a 500 MHz NMR instrument, and has ^1H NMR δ values at 7.08, 7.06, 6.75, 3.75, 2.56, 2.15, 0.93, 0.93 and ^{13}C NMR values of δ 158.2, 156.3, 155.5, 132.6, 129.5, 129.5, 127.3, 121.8, 115.2, 115.2, 41.2, 35.3, 26.7, 21.5, 21.5. The molecular formula, is assigned as $\text{C}_{15}\text{H}_{18}\text{N}_2\text{O}_2$, which is determined by interpretation of ^1H , ^{13}C NMR and mass data. The ^{13}C NMR spectrum revealed signals for all 15 carbons, including two methyls, two methylene carbons, one aliphatic methine, one amide carbonyl, and nine aromatic carbons. The general nature of the structure was deduced from ^1H and ^{13}C NMR spectra that showed a *para*-substituted aromatic ring [δ 7.08 (2H, d, $J = 8.8$ Hz), 6.75 (2H, d, $J = 8.8$ Hz), and 132.7, 129.5, 115.2, 127.3, 115.2, 129.5]. The ^1H NMR spectrum of this structure together with the ^1H - ^1H COSY and HSQC spectra, displayed characteristic signals for an isobutyl moiety [δ 0.93 (6H, d, $J = 6.9$ Hz), 2.15 (1H, sept., $J = 6.9$ Hz), 2.57 (2H, d, $J = 6.9$ Hz). In addition, an olefinic/aromatic proton at (δ 7.06, s), and a carbonyl carbon group (δ 158.9) were also found in the ^1H and ^{13}C NMR spectra. On inspection of the HMBC spectrum, the H-1' signal in the isobutyl moiety correlated with the olefinic carbon (C-2, δ 156.3), and the olefinic proton H-4 correlated with (C-5, δ 155.5; C-2, 156.3 & C-1'', 41.2). The methylene signal at δ 3.75 correlated with C-5, C-4 as well as the C-2'' of

the para-substituted aromatic moiety. All these observed correlations suggested the connectivity among the isobutyl, and the para-substituted benzyl moieties for the skeleton of the structure as shown. In addition, the carboxamide group is assigned at the para position of the benzyl moiety based on the HMBC correlation from the aromatic proton at H-4'' & H-6'' position. Thus, based on the above data, the structure was designated as templazole B.

7. Example 7. Isolation of FR90128

The whole cell broth from the fermentation of *Burkholderia sp.* in an undefined growth medium is extracted with Amberlite XAD-7 resin (Asolkar et al., 2006) by shaking the cell suspension with resin at 225 rpm for two hours at room temperature. The resin and cell mass are collected by filtration through cheesecloth and washed with DI water to remove salts. The resin, cell mass, and cheesecloth are then soaked for 2 h in acetone after which the acetone is filtered and dried under vacuum using rotary evaporator to give the crude extract. The crude extract is then fractionated by using reversed-phase C18 vacuum liquid chromatography (H₂O/CH₃OH; gradient 90:20 to 0:100%) to give 10 fractions. These fractions are then concentrated to dryness using rotary evaporator and the resulting dry residues are screened for biological activity using both insect bioassay as well as herbicidal bioassay. The active fractions are then subjected to reversed/normal phase HPLC (Spectra System P4000; Thermo Scientific) to give pure compounds, which are then screened in herbicidal, insecticidal and nematocidal bioassays described below to locate/identify the active compounds. To confirm the identity of the compound, additional spectroscopic data such as LC/MS and NMR is recorded.

Mass spectroscopy analysis of active peaks is performed on a Thermo Finnigan LCQ Deca XP Plus electrospray (ESI) instrument using both positive and negative ionization modes in a full scan mode (m/z 100-1500 Da) on a LCQ DECA XP^{plus} Mass Spectrometer (Thermo Electron Corp., San Jose, CA). Thermo high performance liquid chromatography (HPLC) instrument equipped with Finnigan Surveyor PDA plus detector, autosampler plus, MS pump and a 4.6 mm x 100 mm Luna C18 5 μm column (Phenomenex). The solvent system consists of water (solvent A) and acetonitrile (solvent B). The mobile phase begins at 10% solvent B and is linearly increased to 100% solvent B over 20 min and then kept for 4 min, and finally returned to 10% solvent B over 3 min and kept for 3 min. The flow rate is 0.5 mL/min. The injection volume is 10 μL and the samples are kept at room temperature in an auto sampler. The compounds are analyzed by LC-MS utilizing the LC and reversed phase chromatography. Mass spectroscopy analysis of the present compounds is performed under the following conditions: The flow rate of the nitrogen gas is fixed at 30 and 15 arb for the sheath and aux/sweep gas flow rate, respectively. Electrospray ionization is performed with a spray voltage set at 5000 V and a

capillary voltage at 35.0 V. The capillary temperature is set at 400°C. The data is analyzed on Xcalibur software. Based on the LC-MS analysis, the active insecticidal compound from fraction 5 has a molecular mass of 540 in negative ionization mode.

For structure elucidation, the purified insecticidal compound from fraction 5 with molecular weight 540 is further analyzed using a 500 MHz NMR instrument, and has ¹H NMR values at 6.22, 5.81, 5.69, 5.66, 5.65, 4.64, 4.31, 3.93, 3.22, 3.21, 3.15, 3.10, 2.69, 2.62, 2.26, 2.23, 1.74, 1.15, 1.12, 1.05, 1.02; and has ¹³C NMR values of 172.99, 172.93, 169.57, 169.23, 167.59, 130.74, 130.12, 129.93, 128.32, 73.49, 62.95, 59.42, 57.73, 38.39, 38.00, 35.49, 30.90, 30.36, 29.26, 18.59, 18.38, 18.09, 17.93, 12.51. The NMR data indicates that the compound contains amino, ester, carboxylic acid, aliphatic methyl, ethyl, methylene, oxymethylene, methine, oxymethine and sulfur groups. The detailed 1D and 2D NMR analysis confirms the structure for the compound as FR90128 as a known compound.

8. Example 8. Herbicidal activity of FR90128

The herbicidal activity of the active compound FR90128 (MW 540) is tested in a laboratory assay using one-week old barnyard grass (*Echinochloa crus-galli*) seedlings in a 96-well plate platform. One grass seedling was placed in each of the wells containing 99 microliters of DI water. One microliter aliquot of the pure compound in ethanol (10 mg/mL) is added into each well, and the plate is sealed with a lid. One microliter of ethanol in 99 microliters of water is used as a negative control. The treatments were done in eight replicates, and the sealed plate is incubated in a greenhouse under artificial lights (12 hr light/dark cycle). After five days, the results are read. The grass seedlings in all eight wells that received the active compound are dead with no green tissue left, whereas the seedlings in the negative control wells were actively growing.

9. Example 9. Insecticidal activity of FR90128

The insecticidal activity of the active compound FR90128 (MW 540) is tested in a laboratory assay using a contact bioassay system. The compound is dissolved in 100% ethanol to concentrations of 0.001, 0.005, 0.01, 0.025, 0.05, 0.1, 0.25, and 0.5 µg/µL. Individual early 3rd instar Beet Armyworm, *Spodoptera exigua*, larvae are placed in 1.25 ounce plastic cups with a 1 cm² piece of artificial diet (Bio-Serv). A Hamilton Micropipette is used to apply 1 µL of compound to the thorax of each larvae. Cups are covered with stretched parafilm and a single hole is cut into the parafilm for aeration. Ten larvae per concentration are treated. The assay is incubated at 25°C, 12h light/12h dark. Larvae are scored at 48 and 72 hours after application. Probit analysis is performed to assess LC₅₀ value which is found for compound (MW 540) as 0.213.

10. Example 10. Isolation of Templamide A, B, FR901465 and FR90128

Methods and Materials

The following procedure is used for the purification of compounds extracted from cell culture of *Burkholderia sp* (see Figure 7):

5 The culture broth derived from the 10-L fermentation *Burkholderia* (A396) in Hy soy growth medium is extracted with Amberlite XAD-7 resin (Asolkar et al., 2006) by shaking the cell suspension with resin at 225 rpm for two hours at room temperature. The resin and cell mass are collected by filtration through cheesecloth and washed with DI water to remove salts. The resin, cell mass, and cheesecloth are then soaked for 2 h in acetone
10 after which the acetone is filtered and dried under vacuum using rotary evaporator to give the crude extract. The crude extract is then fractionated by using reversed-phase C18 vacuum liquid chromatography (H₂O/CH₃OH; gradient 90:20 to 0:100%) to give 10 fractions. These fractions are then concentrated to dryness using rotary evaporator and the resulting dry residues are screened for biological activity using 96 well plate lettuce seeding
15 (herbicidal) and early 3rd instar Beet Armyworm (insecticidal) assay. The active fractions are then subjected to repeatedly to reversed phase HPLC separation (Spectra System P4000 (Thermo Scientific) to give pure compounds, which are then screened in above-mentioned bioassays to locate/identify the active compounds. To confirm the identity of the compound, additional spectroscopic data such as LC/MS, HRMS and NMR are recorded.

20 The active fraction 5 is purified further by using HPLC C-18 column (Phenomenex, Luna 10u C18(2) 100 A, 250 x 30), water:acetonitrile gradient solvent system (0-10 min; 80 % aqueous CH₃CN, 10-25 min; 80 - 65 % aqueous CH₃CN, 25-50 min; 65 - 50 % aqueous CH₃CN, 50-60 min; 50 - 70 % aqueous CH₃CN, 60-80 min; 70 - 0 % aqueous CH₃CN, 80-85 min; 0 - 20 % aqueous CH₃CN) at 8 mL/min flow rate and UV detection of
25 210 nm, to give templamide A, retention time 55.64 min and FR901465, retention time 63.59 min and FR90128, retention time 66.65 min respectively. The other active fraction 6 is also purified using HPLC C-18 column (Phenomenex, Luna 10u C18(2) 100 A, 250 x 30), water:acetonitrile gradient solvent system (0-10 min; 70-60 % aqueous CH₃CN, 10-20 min; 60-40 % aqueous CH₃CN, 20-50 min; 40 - 15 % aqueous CH₃CN, 50-75 min; 15 - 0 %
30 CH₃CN, 75-85 min; 0 - 70 % aqueous CH₃CN) at 8 mL/min flow rate and UV detection of 210 nm, to give templamide B, retention time 38.55 min.

Mass spectroscopy analysis of pure compounds is performed on a Thermo Finnigan LCQ Deca XP Plus electrospray (ESI) instrument using both positive and negative

ionization modes in a full scan mode (m/z 100-1500 Da) on a LCQ DECA XP^{plus} Mass Spectrometer (Thermo Electron Corp., San Jose, CA). Thermo high performance liquid chromatography (HPLC) instrument equipped with Finnigan Surveyor PDA plus detector, autosampler plus, MS pump and a 4.6 mm x 100 mm Luna C18 5 μ m column

5 (Phenomenex) is used. The solvent system consists of water (solvent A) and acetonitrile (solvent B). The mobile phase begins at 10% solvent B and is linearly increased to 100% solvent B over 20 min and then kept for 4 min, and finally returns to 10% solvent B over 3 min and kept for 3 min. The flow rate is 0.5 mL/min. The injection volume is 10 μ L and the samples are kept at room temperature in an auto sampler. The compounds are analyzed
10 by LC-MS utilizing the LC and reversed phase chromatography. Mass spectroscopy analysis of the present compounds is performed under the following conditions: The flow rate of the nitrogen gas is fixed at 30 and 15 arb for the sheath and aux/sweep gas flow rate, respectively. Electrospray ionization is performed with a spray voltage set at 5000 V and a capillary voltage at 45.0 V. The capillary temperature is set at 300°C. The data is analyzed
15 on Xcalibur software. The active compound templamide A has a molecular mass of 555 based on the m/z peak at 556.41 $[M + H]^+$ and 578.34 $[M + Na]^+$ in positive ionization mode. The LC-MS analysis in positive mode ionization for templamide B suggests a molecular mass of 537 based m/z ions at 538.47 $[M + H]^+$ and 560.65 $[M + Na]^+$. The molecular weight for the compounds FR901465 and FR90128 are assigned as 523 and 540
20 respectively on the basis of LCMS analysis.

¹H, ¹³C and 2D NMR spectra are measured on a Bruker 600 MHz gradient field spectrometer. The reference is set on the internal standard tetramethylsilane (TMS, 0.00 ppm).

For structure elucidation of templamide A, the purified compound with molecular
25 weight 555 is further analyzed using a 600 MHz NMR instrument, and has ¹H NMR δ values at 6.40, 6.39, 6.00, 5.97, 5.67, 5.54, 4.33, 3.77, 3.73, 3.70, 3.59, 3.47, 3.41, 2.44, 2.35, 2.26, 1.97, 1.81, 1.76, 1.42, 1.37, 1.16, 1.12, 1.04 and has ¹³C NMR values of δ 173.92, 166.06, 145.06, 138.76, 135.71, 129.99, 126.20, 123.35, 99.75, 82.20, 78.22, 76.69, 71.23, 70.79, 70.48, 69.84, 60.98, 48.84, 36.89, 33.09, 30.63, 28.55, 25.88, 20.37, 18.11,
30 14.90, 12.81, 9.41. The ¹³C NMR spectrum exhibits 28 discrete carbon signals which are attributed to six methyls, four methylene carbons, and thirteen methines including five sp^2 , four quaternary carbons. The molecular formula, C₂₈H₄₅NO₁₀, is determined by interpretation of ¹H, ¹³C NMR and HRESI MS data. The detailed analysis of ¹H-¹H COSY, HMBC and HMQC spectral data reveals the following substructures (I – IV) and two
35 isolated methylene & singlet methyl groups. These substructures are connected later using

3.21, 3.15, 3.10, 2.69, 2.62, 2.26, 2.23, 1.74, 1.15, 1.12, 1.05, 1.02; and has ^{13}C NMR values of 172.99, 172.93, 169.57, 169.23, 167.59, 130.74, 130.12, 129.93, 128.32, 73.49, 62.95, 59.42, 57.73, 38.39, 38.00, 35.49, 30.90, 30.36, 29.26, 18.59, 18.38, 18.09, 17.93, 12.51.

The NMR data indicates that the compound contains amino, ester, carboxylic acid, aliphatic methyl, ethyl, methylene, oxymethylene, methine, oxymethine and sulfur groups. The detailed 1D and 2D NMR analysis confirm the structure for the compound as FR90128 as a known compound.

11. Example 11. Herbicidal activity of Templamide A, Templamide B, FR901465 and FR90128

The herbicidal activity of templamide A, B, FR901465 and FR90128 are tested in a laboratory assay using one-week old barnyard grass (*Echinochloa crus-galli*) and lettuce (*Lactuca sativa* L.) seedlings in a 96-well plate platform. One seedling is placed in each of the wells containing 99 microliters of DI water. Into each well, a one microliter aliquot of the pure compound in ethanol (10 mg/mL) is added, and the plate is sealed with a lid. One microliter of ethanol in 99 microliters of water is used as a negative control. The treatments are done in eight replicates, and the sealed plate is incubated in a greenhouse under artificial lights (12 hr light/dark cycle). After five days, the results are read. The grass seedlings in all eight wells that received the active compound are dead with no green tissue left, whereas the seedlings in the negative control wells are actively growing. The herbicidal activity of templamide A against lettuce seedlings is slightly lower than for the grass. On the other hand, templamide B provides a better (100%) control of lettuce seedlings (used as a model system for broadleaf weeds) than templamide A (Table 11).

Table 11: Herbicidal Bioassay data for Templamide A, B, FR901465 and FR90128

Compounds ¹	Grass seedlings (% Mortality)	Lettuce seedlings (% Mortality)
Templamide A	100	88
Templamide B	0	75
FR901465	88	100
FR90128	100	88
Control	0	0

¹10 µg/mL concentration per well

12. Example 12. Insecticidal activity of active compounds

The insecticidal activity of templamide A, B, FR901465 and FR90128 are tested in a laboratory assay using a 96-well diet overlay assay with 1st instar Beet Armyworm larvae using microtiter plates with 200 µl of solid, artificial Beet Armyworm diet in each well. One hundred (100) µl of each test sample is pipetted on the top of the diet (one sample in each well), and the sample is let dry under flowing air until the surface is dry. Each sample

was tested in six replicates, and water and a commercial Bt (*B. thuringiensis*) product are used as negative and positive controls, respectively. One first instar larvae of the test insect (Beet armyworm – *Spodoptera exigua*) was placed in each well, and the plate was covered with plastic cover with airholes. The plates with insects were incubated at 26°C for 6 days with daily mortality evaluations. Based on the results presented in Table 12, templamide A and B results in 40% and 80% mortality, respectively.

Table 12: Insecticidal Bioassay data for Templamide A, B, FR901465 and FR90128 against 1st instar Beet Army Worm (*Spodoptera exigua*).

Compounds ¹	(% Mortality)
Templamide A	40
Templamide B	80
FR901465	50
FR90128	90
Bt	100
Control	0

¹ 10 µg/mL concentration per well

Example 11. Fungicidal activity of FR90128 (MW 540)

Fungicidal activity of FR90128 (MW 540) against three plant pathogenic fungi (*Botrytis cinerea*, *Phytophthora* sp., *Monilinia fructicola*) is tested in an in vitro PDA (potato dextrose agar) plate assay. Plates are inoculated with the fungus using a plug method. After the fungus had established and started to grow on the growth medium, eight sterile filter paper disks are placed on each plate about 1 cm from the edge in a circle. Ten microliters of ethanol solution containing 20, 15, 10, 7.5, 5, 2.5 1.25 mg FR90128/mL is added into filter paper disks, and the solution is left to evaporate. One disk imbedded with 10 µL of pure ethanol is used as a negative control. The assay is done with three replicates. Plates are incubated at room temperature for 5 days, after which the fungicidal activity is recorded by measuring the inhibition zone around each filter paper disk corresponding to different concentrations of FR90128. According to the results, FR90128 has no effect on the growth of *Monilinia* but it is effective in controlling the hyphal growth of both *Botrytis* and *Phytophthora*. There seems to be a clear dose-response in inhibition with threshold concentrations of 10 mg/mL and 1.25 mg/mL for *Botrytis* and *Phytophthora*, respectively (Figure 8).

DEPOSIT OF BIOLOGICAL MATERIAL

The following biological material has been deposited under the terms of the Budapest Treaty with the Agricultural Research Culture Collection (NRRL), 1815 N. University Street, Peoria, Illinois 61604 USA, and given the following number:

<u>Deposit</u>	<u>Accession Number</u>	<u>Date of Deposit</u>
<i>Burkholderia sp.</i> A396	NRRL B-50319	September 15, 2009

The strain has been deposited under conditions that assure that access to the culture will be available during the pendency of this patent application to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 C.F.R. §1.14 and 35 U.S.C. §122. The deposit represents a substantially pure culture of the deposited strain. The deposit is available as required by foreign patent laws in countries wherein counterparts of the subject application, or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by government action.

The invention described and claimed herein is not to be limited in scope by the specific aspects herein disclosed, since these aspects are intended as illustrations of several aspects of the invention. Any equivalent aspects are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. In the case of conflict, the present disclosure including definitions will control.

Various references are cited herein, the disclosures of which are incorporated by reference in their entireties.

Literature cited:

Anderson, et al. "The structure of thiostrepton," *Nature* 225: 233-235. 1970.

Andra, "Endotoxin-like properties of a rhamnolipid exotoxin from *Burkholderia (Pseudomonas) plantarii*: immune cell stimulation and biophysical characterization." *Biol. Chem.* 387: 301-310. 2006.

Arena, et al. "The mechanism of action of avermectins in *Caenorhabditis elegans* - correlation between activation of glutamate-sensitive chloride current, membrane binding and biological activity." *J Parasitol.* 81: 286-294. 1995.

Asolkar, et al., "Weakly cytotoxic polyketides from a marine-derived Actinomycete of the genus *Streptomyces* strain CNQ-085." *J. Nat. Prod.* 69:1756-1759. 2006.

Burkhead, et al., "Pyrrolnitrin production by biological control agent *Pseudomonas cepacia* B37w in culture and in colonized wounds of potatoes." *Appl. Environ. Microbiol.* 60: 2031-2039. 1994.

Burkholder, W. H. "Sour skin, a bacterial rot of onion bulbs." *Phytopathology* 40: 115-117. 1950.

Caballero-Mellado et al., "*Burkholderia unamae* sp. nov., an N₂-fixing rhizospheric and endophytic species." *Int. J. Syst. Evol. Microbiol.* 54: 1165-1172. 2004.

Cashion et al. "Rapid method for base ratio determination of bacterial DNA." *Anal. Biochem.* 81: 461-466. 1977.

Casida, et al., US Patent No. 6,689,357.

Chen et al. , "*Burkholderia nodosa* sp. nov., isolated from root nodules of the woody Brazilian legumes *Mimosa bimucronata* and *Mimosa scabrella* " *Int. J. Syst. Evol. Microbiol.* 57: 1055-1059. 2007.

Cheng, A. C. and Currie, B. J. "Melioidosis: epidemiology, pathophysiology, and management." *Clin. Microbiol.* 18: 383-416. 2005.

Coenye, T. and Vandamme, P. "Diversity and significance of *Burkholderia* species occupying diverse ecological niches." *Environ. Microbiol.* 5: 719-729. 2003.

Compant, et al. "Diversity and occurrence of *Burkholderia* spp. in the natural environment." *FEMS Microbiol. Rev.* 32: 607-626. 2008.

De Ley et al. "The quantitative measurement of DNA hybridization from renaturation rates." *Eur. J. Biochem.* 12: 133-142. 1970.

Duke et al. "Natural products as sources for herbicides: current status and future trends." *Weed Res* 40: 99-111. 2000.

Gerwick et al., US Patent No. 7,393,812.

Gottlieb et al., US Patent No. 4,808,207.

Gouge et al., US Patent Application Pub. No. 2003/0082147.

Guella et al. "Almazole C, a new indole alkaloid bearing an unusually 2,5-disubstituted oxazole moiety and its putative biogenetic precursors, from a Senegalese Delesseriacean sea weed." *Helv. Chim. Acta* 77: 1999-2006. 1994.

5 Guella et al. "Isolation, synthesis and photochemical properties of almazolone, a new indole alkaloid from a red alga of Senegal." *Tetrahedron*. 62: 1165-1170. 2006.

Henderson, P. J. and Lardy H. A. "Bongkrekiic acid. An inhibitor of the adenine nucleotide translocase of mitochondria." *J. Biol. Chem.* 245: 1319-1326. 1970.

Hirota et al. "Isolation of indolmycin and its derivatives as antagonists of L-tryptophan." *Agri. Biol Chem.* 42: 147-151. 1978.

10 Hu, F.-P. and Young, J. M. "Biocidal activity in plant pathogenic *Acidovorax*, *Burkholderia*, *Herbaspirillum*, *Ralstonia*, and *Xanthomonas* spp." *J. Appl. Microbiol.* 84: 263-271. 1998.

Huss et al. "Studies of the spectrophotometric determination of DNA hybridization from renaturation rates." *System. Appl. Microbiol.* 4: 184-192. 1983.

15 Jansiewicz, W. J. and Roitman J. "Biological control of blue mold and gray mold on apple and pear with *Pseudomonas cepacia*." *Phytopathology* 78: 1697-1700. 1988.

Jeddeloh et al., WO2001/055398.

Jansen et al. "Thiangazole: a novel inhibitor of HIV-1 from *Polyangium* Spec." *Liebigs Ann. Chem.* 4: 357-3359. 1992.

20 Jeong et al. "Toxoflavin produced by *Burkholderia glumae* causing rice grain rot is responsible for inducing bacterial wilt in many field crops." *Plant Disease* 87: 890-895. 2003.

Knudsen, G. R. and Spurr, J. "Field persistence and efficacy of five bacterial preparations for control of peanut leaf spot." *Plant Disease* 71: 442-445. 1987.

25 Koga-Ban et al. "cDNA sequences of three kinds of beta-tubulins from rice." *DNA Research* 2: 21-26. 1995.

Koide et al. US Patent Application Pub. No. 2008/0096879.

30 Koyama et al. "Isolation, characterization, and synthesis of pimprinine, pimrinrthine, and pimprinaphine, metabolites of *Streptoverticillium olivoreticuli*." *Agri. Biol. Chem.* 45: 1285-1287. 1981.

Krieg et al. "*Bacillus thuringiensis* var. tenebrionis: Ein neuer, gegenüber Larven von Coleopteren wirksamer Pathotyp." *Z. Angew. Entomol.* 96:500-508. 1983.

35 Kunze et al. "Thiangazole, a new thiazoline antibiotic from *Polyangium* sp (Myxobacteria Production, antimicrobial activity and mechanism of action." *J. Antibiot.*, 46: 1752-1755. 1993.

Leahy et al. "Comparison of factors influencing trichloroethylene degradation by toluene-oxidizing bacteria." *Appl. Environ. Microbiol.* 62: 825-833. 1996.

Lessie et al. "Genomic complexity and plasticity of *Burkholderia cepacia*." FEMS Microbiol. Lett. 144: 117-128. 1996.

Lindquist, N. et al. "Isolation and structure determination of diazonamides A and B, unusual cytotoxic metabolites from the marine ascidian *Diazona chinensis*." J. Am Chem. Soc. 113: 2303-2304. 1991.

Lorch, H et al. "Basic methods for counting microorganisms in soil and water. In Methods in applied soil microbiology and biochemistry. K. Alef and P. Nannipieri, Eds. San Diego, CA, Academic Press: pp. 146-161. 1995.

Ludovic et al. "Burkholderia diversity and versatility: An inventory of the extracellular products." J. Microbiol. Biotechnol. 17: 1407-1429. 2007.

Lydon, J. and Duke, S. "Inhibitors of glutamine biosynthesis." in Plant amino acids: Biochemistry and Biotechnology. B. Singh., Ed. New York, USA, Marcel Decker. pp. 445-464. 1999.

Mahenthalingam et al. "DNA-based diagnostic approaches for identification of *Burkholderia cepacia* complex, *Burkholderia vietnamiensis*, *Burkholderia multivorans*, *Burkholderia stabilis*, and *Burkholderia cepacia* genomovars I and III." J.Clin. Microbiol. 38: 3165-3173. 2000.

Ming, L.-J. and Epperson. "Metal binding and structure-activity relationship of the metalloantibiotic peptide bacitracin." Biochemistry 91: 46-58. 2002.

Morita et al. "Biological activity of tropolone." Biol. Pharm. Bull. 26: 1487-1490. 2003.

Nagamatsu, T. "Syntheses, transformation, and biological activities of 7-azapteridine antibiotics: toxoflavin, fervenulin, reumycin, and their analogs". Recent Res. Devel. Org. Bioorg. Chem. 4: 97-121. 2001.

Naik et al., "Pimprine, an extracellular alkaloid produced by *Streptomyces* CDRIL-312: fermentation, isolation and pharmacological activity." J. Biotech. 88: 1-10. 2001.

Nakajima et al., "Antitumor Substances, FR901463, FR901464 and FR901465. I. Taxonomy, Fermentation, Isolation, Physico-chemical Properties and Biological Activities." J. Antibiot. 49: 1196-1203. 1996.

Nakajima et al. US Patent No. 5,545,542.

Nakajima et al., "Hydantocidin: a new compound with herbicidal activity." J Antibiot. 44: 293-300. 1991.

N'Diaye, I. et al., "Almazole A and amazole B, unusual marine alkaloids of an unidentified red seaweed of the family *Delesseriaceae* from the coasts of Senegal." Tet Lett. 35: 4827-4830. 1994.

N'Diaye, I. et al., "Almazole D, a new type of antibacterial 2,5-disubstituted oxazolic dipeptide from a red alga of the coast of Senegal." Tet Lett. 37: 3049-3050. 1996.

Nierman et al., "Structural flexibility in the *Burkholderia mallei* genome." Proc. Natl. Acad. Sci. USA 101: 14246-14251. 2004.

Okazaki et al., "Rhizobial strategies to enhance symbiotic interaction: Rhizobitoxine and 1-aminocyclopropane-1-carboxylate deaminase." Microbes Environ. 19: 99-111. 2004.

5 Parke, J. L. and D. Gurian-Sherman, D. 2001. "Diversity of the *Burkholderia cepacia* complex and implications for risk assessment of biological control strains." Annual Reviews in Phytopathology 39: 225-258. 2001.

Parke, et al. US Patent No. 6,077,505.

10 Pettit, G. et al. "Isolation of Labradorins 1 and 2 from *Pseudomonas syringae*." J. Nat. Prod. 65: 1793-1797. 2002.

Pitt, et al., "Type characterization and antibiotic susceptibility of *Burkholderia (Pseudomonas) cepacia* isolates from patients with cystic fibrosis in the United Kingdom and the Republic of Ireland." J. Med. Microbiol. 44: 203-210. 1996.

15 Ramette et al., "Species abundance and diversity of *Burkholderia cepacia* complex in the environment." Appl. Environ. Microbiol. 71: 1193-1201. 2005.

Resi et al., "*Burkholderia tropica* sp. nov., a novel nitrogen-fixing, plant-associated bacterium." Int. J. Syst. Evol. Microbiol. 54: 2155-2162. 2004.

Salama et al. "Potency of spore-gamma-endotoxin complexes of *Bacillus thuringiensis* against some cotton pests." Z. Angew. Entomol. 91: 388-398. 1981.

20 Selva et al., "Targeted screening for elongation factor Tu binding antibiotics." J. Antibiot. 50: 22-26. 1997.

Takahashi, S. et al. "Martefragin A, a novel indole alkaloid isolated from a red alga, inhibits lipid peroxidation." Chem Pharm. Bull. 46: 1527-1529. 1998.

25 Thompson et al. "Spinosad - a case study: an example from a natural products discovery programme." Pest Management Science 56: 696-702. 2000.

Takita et al., "Chemistry of Bleomycin. XIX Revised structures of bleomycin and phleomycin." J. Antibiot. 31: 801-804. 1978.

30 Tran Van et al., "Repeated beneficial effects of rice inoculation with a strain of *Burkholderia vietnamiensis* on early and late yield component in low fertility sulphate acid soils of Vietnam." Plant and Soil 218: 273-284. 2000.

Tsuruo et al., "Rhizoxin, a macrocyclic lactone antibiotic, as a new antitumor agent against human and murine tumor cells and their vincristine-resistant sublines." Cancer Res. 46: 381-385. 1986.

Ueda et al., US Patent No. 7,396,665.

35 Umehara, K. et al. "Studies of new antiplatelet agents WS-30581 A and B." J. Antibiot. 37: 1153-1160. 1984.

Vandamme et al. Polyphasic taxonomic study of the emended genus *Arcobacter* with *Arcobacter butzleri* comb. nov. and *Arcobacter skirrowii* sp. nov., an aerotolerant bacterium isolated from veterinary specimens." *Int. J. Syst. Bacteriol.* 42: 344-356. 1992.

5 Vanderwall et al., "A model of the structure of HOO-Co•bleomycin bound to d(CCAGTACTGG): recognition at the d(GpT)site and implications for double-stranded DNA cleavage, *Chem. Biol.* 4: 373-387. 1997.

Vermis K., et al. "Evaluation of species-specific recA-based PCR tests for genomovar level identification within the *Burkholderia cepacia* complex." *J. Med. Microbiol* 51: 937-940, 2002.

10 Watanabe, H. et al. "A new antibiotic SF2583A, 4-chloro-5-(3'indoly)oxazole, produced by *Streptomyces*." *Meiji Seika Kenkyu Nenpo* 27: 55-62. 1988.

Wayne et al., "Report of the Ad Hoc committee on reconciliation of approaches to bacterial systematics." *Int. J. Syst. Evol. Microbiol.* 37: 463-464. 1987.

15 Werner et al., "Uptake of indolmycin in gram-positive bacteria." *Antimicrob Agents Chemotherapy* 18: 858-862. 1980.

Wilson et al. "Toxicity of rhizonin A, isolated from *Rhizopus microsporus*, in laboratory animals." *Food Chem. Toxicol.* 22: 275-281. 1984.

Zeck W.M. "Ein Bonitierungsschema zur Felddauswertung von Wurzelgallenbefall. *Pflanzenschutznachrichten.*" *Bayer* 24,1: 144-147. 1971.

20 Zhang et al., US Patent No. 7,141,407.

Zhou et al., "Antimicrobial susceptibility and synergy studies of *Burkholderia cepacia* complex isolated from patients with cystic fibrosis." *Antimicrobial Agents and Chemotherapy* 51: 1085-088. 2007.

CLAIMS

1. A composition comprising an isolated strain of a *Burkholderia* sp. which has the following characteristics:

(A) a 16S rRNA gene sequence comprising the forward sequence having at least 99% identity to the sequence set forth in SEQ ID NO: 8, 11, 12 and a reverse sequence having at least 99% identity to the sequence set forth in SEQ ID NO:9, 10, 13, 14 and 15;

(B) pesticidal activity;

(C) produces a pesticidal compound selected from the group consisting of: templazole A, templazole B, templamide A, templamide B, and FR90128; and at least one of a carrier, diluent, surfactant, or adjuvant.

2. The composition of claim 1, wherein the *Burkholderia* species is a *Burkholderia* strain having the identifying characteristics of *Burkholderia* A396 (NRRL Accession No. B-50319).

3. A composition comprising substantially pure culture, cell fraction, whole cell broth, or supernatant derived from the strain of claims 1-2 or extract thereof.

4. A method for producing a compound selected from the group consisting of

(i) templazole A;

(ii) templazole B;

(iii) templamide A;

(iv) templamide B; and

(v) FR90128;

wherein the method comprises the steps of culturing the strain of claims 1-2 and producing said compound.

5. A method for isolating a compound from a strain of *Burkholderia* comprising

(a) producing one or more said compound according to the method of claim 4;

(b) isolating the one or more said compound produced in step (a) from the supernatant of said culture.

6. A composition comprising

(a) a pure culture, cell fraction, whole cell broth, or supernatant derived from the strain of claims 1-2 or extract thereof;

(b) a second substance, wherein said second substance is a chemical or biological pesticide and

(c) optionally at least one of a carrier, diluent, surfactant, adjuvant.

7. The composition according to claim 6, wherein said second substance is a pesticide selected from the group consisting of nematicide, herbicide, fungicide and insecticide.

8. A method for modulating pest infestation in a plant comprising applying to the plant and/or seeds thereof and/or substrate used for growing said plant an amount of the composition of claims 1-3, 6 or 7 effective to modulate said pest infestation.

9. A method for modulating emergence and/or growth of monocotyledonous, sedge or dicotyledonous weeds comprising applying to said weed or soil an amount of the composition of claims 1-3, 6 or 7 effective to modulate said emergence and/or growth of said weeds.

10. A seed comprising the composition of claims 1-3, 6 or 7.

Marrone Bio Innovations, Inc.

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON

Figure 1




Figure 2

UTC (back)
A396 @ 5 mg/mL (middle)
A396 @ 10 mg/mL (front)



Figure 3




Figure 4




Figure 5

% Mortality in contact assay , Beet armyworm and A396




Figure 6




Figure 7




Figure 8



Figure 9





Figure 10



6669556Seq. TXT
SEQUENCE LISTING

<110> Marrone Bio Innovations, Inc.
<120> ISOLATED BACTERIAL STRAIN OF THE GENUS BURKHOLDERIA
<130> MCI - 42016- US
<150> 61308287
<151> 2010-02-25
<150> 61406541
<151> 2010-10-25
<160> 15
<170> Patent In version 3.5
<210> 1
<211> 20
<212> DNA
<213> primer
<400> 1
agagtttgat cctggctcag 20

<210> 2
<211> 20
<212> DNA
<213> primer
<400> 2
ccgtcaattcctttgagttt 20

<210> 3
<211> 16
<212> DNA
<213> primer
<400> 3
gtgccagccg ccgcgg 16

<210> 4
<211> 16
<212> DNA
<213> primer
<400> 4
gcaacgagcg caacc 16

<210> 5
<211> 17
<212> DNA
<213> primer
<400> 5
aaggagtgwtccarcc 17

<210> 6
<211> 15
<212> DNA
<213> primer

<400> 6
gggt t gcgct cgt t g 15

<210> 7
<211> 18
<212> DNA
<213> pr i m e r

<400> 7
gwat t accgc ggckgct g 18

<210> 8
<211> 871
<212> DNA
<213> Bur khol der i a A396

<400> 8
t gcagt cgaa cggcagcacg ggt gct t gca cct ggt ggcg agt ggcgaac ggggtagt aa 60
t acat cggaa cat gt cct gt agt ggggat agcccggcga aagccggat t aat accgcat 120
acgat ct acg gat gaaagcg ggggat ct t c ggacct cgcg ct at agggg t ggccgat ggc 180
t gat t agct a gt t ggt gggg t aaaggcct a ccaaggcgac gat cagt agc t ggt ct gaga 240
ggacgat cag ccacact ggg act gagacac ggcccagact cct acgggag gcagcagt gg 300
ggaat t t t gg acaat ggggg aaaccct gat ccagcaat gc cgcgt gt gt g aagaaggcct 360
t cgggt t gt a aagcact t t t gt ccggaaag aaat cct t t g ggct aat acc ccggggggat 420
gacggg accg gaagaat aag caccggct aa ct acgt gcca gcagccgcgg t aat acgt ag 480
ggg gcgagcg t t aat cggaa t t act gggcg t aaagcgt gc gcaggcggg t t gt t aagaca 540
gat gt gaaat ccccgggct t aacct gggaa ct gcat t t gt gact ggcaag ct agagt at g 600
gcagaggggg gt agaat t cc acgt gt agca gt gaaat gcg t agagat gt g gaggaat acc 660
gat ggcgaag gcagccccct gggccaat ac t gacgct cat gcacgaaagc gt ggggagca 720
aacaggat t a gat acct gg t agt ccacgc cct aaacgat gt caact agt t gt t ggggat 780
t cat t t cct t agt aacgt ag ct acgct ga agt t gaccgc ct ggggagt a cggt cgcaag 840
at t aaat mja gggg kgkkt g kkggggggaa a 871

<210> 9
<211> 1453
<212> DNA
<213> Bur khol der i a A396

<400> 9
gt cat gaat c ct accgt ggt gaccgt cct c ct t gcggt t a gact agccac t t ct ggt aaa 60
accact ccc at ggt gt gac gggcggg gt g t acaagacct gggaacgt at t caccgcggc 120
at gct gat cc gcgat t act a gcgat t ccag ct t cat gcac t cgagt t gca gagt gcaat c 180
cggact acga t cggg t t t ct gggat t agct cccct cgcg ggt t ggcaac cct ct gt t cc 240
gaccat t gt a t gacgt gt ga agccct accc at aagggccca t gaggact t g acgt cat ccc 300
cacct t cct c cggg t t gt ca ccggcagt ct cct t agagt g ct ct t gcgt a gcaact aagg 360

acaagggt t g cgct cgt t gc gggact t aac ccaacat ct c acgacacgag ct gacgacag 420
 ccat gcagca cct gt gt at c ggt t ct ct t t cgagcact cc cgaat ct ct t caggat t ccg 480
 accat gt caa ggggt aggt aa ggt t t t t cgc gt t gcat cga at t aat ccac at cat ccacc 540
 gct t gt gcgg gt ccccg t ca at t cct t t ga gt t t t aat ct t gcgaccgt a ct ccccagggc 600
 ggt caact t c acgcgt t agc t acgt t act a aggaaat gaa t cccaacaa ct agt t gaca 660
 t cgt t t aggg cgt ggact ac cagggt at ct aat cct gt t t gct ccccacg ct t t cgt gca 720
 t gagcgt cag t at t ggccca gggggct gcc t t cgccat cg gt at t cct cc acat ct ct ac 780
 gcat t t cact gct acacgt g gaat t ct acc ccct ct gcc at act ct agc t t gccagt ca 840
 caaat gcagt t cccaggt t a agcccgggga t t t cacat ct gt ct t aacaa accgcct gcg 900
 cacgct t t ac gccaggt aat t ccgat t aac gct cgcaccc t acgt at t ac cgcggct gct 960
 ggcacgt agt t agccggt gc t t at t ct t cc ggt accgt ca t cccccggg gt at t agccc 1020
 aaaggat t t c t t t cggaca aaagt gct t t acaaccgaa ggcct t ct t c acacacgcgg 1080
 cat t gct gga t cagggt t t c ccccat t gt c caaaat t ccc cact gct gcc t cccgt agga 1140
 gt ct gggccg t gt ct cagt c ccagt gt ggc t gat cgt cct ct cagaccag ct act gat cg 1200
 t cgct t ggt aggcct t t ac cccaccaact agct aat cag ccat cggcca accct at agc 1260
 gcgaggt ccg aagat ccccc gct t t cat cc gt agat cgt a t gcggt at t a at ccggct t t 1320
 cgccgggct a t cccccact a caggacat gt t ccgat gt at t act cacccg t t cgccact c 1380
 gccaccaggt gcaagcacc gt gct gccgt t cgact t gca t gt gt aaggc at gccgccag 1440
 cgt t caat ct gag 1453

<210> 10
 <211> 860
 <212> DNA
 <213> Burkholderia A396

<400> 10
 ccaggcggc act t cacgcg t t agct acgt t act aaggaa at gaat cccc aacaact agt 60
 t gacat cgt t t agggcgt gg act accaggg t at ct aat cc t gt t t gct cc ccacgct t t c 120
 gt gcat gagg gt cagt at t g gccaggggg ct gcct t cgc cat cgggt at t cct ccacat c 180
 t ct acgcat t t cact gct ac acgt ggaat t ct acccccct ct gccat act ct agct t gcc 240
 agt cacaaat gcagt t ccca ggt t aagccc ggggat t t ca cat ct gt ct t aacaaaccgc 300
 ct gcgcacgc t t t acgcca gt aat t ccga t t aacgct cg caccct acgt at t accgagg 360
 ct gct ggcac gt agt t agcc ggt gct t at t ct t ccgggt ac cgt cat cccc ccgggggt at t 420
 agcccaaagg at t t ct t t cc ggacaaaagt gct t t acaac ccgaaggcct t ct t cacaca 480
 cgcggcatt g ct ggat cagg gt t t ccccca t t gt ccaaaa t t ccccact g ct gcct cccg 540
 t aggagt ct g ggccgt gt ct cagt cccagt gt ggct gat c gt cct ct cag accagct act 600
 gat cgt cgcc t t ggt aggcc t t t accccac caact agct a at cagccat c ggccaaccct 660
 at agcgcgag gt ccgaagat cccccgct t t cat ccgt aga t cgt at gcgg t at t aat ccg 720

6669556Seq. TXT

gct t t cgccg ggct at cccc cact acagga cat gt t ccga t gt at t act c acccgt t cgc 780
 cact cgccac caggt gcaag caccctgt gct gccgt t cgac t t gcat gt gt aaggcat gcc 840
 gccagcgt t c aat ct gagt g 860

<210> 11
 <211> 1152
 <212> DNA
 <213> Burkhol der i a A396

<400> 11
 t cggat t act gggcgt aagc gt ggcgagc ggt t t gt t aa gacagat gt g aaat ccccgg 60
 gct t aacct g ggaact gcat t t gt gact gg caagct agag t at ggcagag gggggg agaa 120
 t t ccacgt gt agcagt gaaa t gcgt agaga t gt ggaggaa t accgat ggc gaagggagcc 180
 ccct gggcct at act gacct t cat gct cga aagcgt gagg acccaaccgg at t agat gcc 240
 ct gat aggcc at gcccaca ccat gccat g t gt t aggggc ccat t t cct t agggaggcag 300
 ct at ggggaa t t t t ggacaa t gt gggaaac cct gat ccaa caat gccgcg t gt gt gaat a 360
 aggcc t cgg gt t gt aaagc act t t t at cc ggat agat t c ct t t t gggct aaacct ccgt 420
 aggggat gac ggt accggaa gaat aaccac cgggt aact a cgt gccagca gccgcggt aa 480
 t acgt aggt gcgagcgt t a at cggaat t a ct gggcgt aa agcgt gcgca ggcggt t t gt 540
 t aagacagat gt gaaat ccc cgggct t aac ct gggaact g cat t t gt gac t ggcaagct a 600
 gagt at ggca gacgggggt a gaat t ccacg t gt agcagt g aaat gcgt ag agat gt ggag 660
 gaat accgat gggcgaagca gct cct gggg caat act gac gct cat gcac aagat cgt gc 720
 gaaacaaaca ggat aaaacc cct gt at t cc acgccccaaa cgat gt ccac caagt t gt t g 780
 gcgat cct t t cct t cgt at c gt agct acgc gggaaat t t ga cccct gggg act aggccgc 840
 at at aaaact caagggat t ccggggacc ccagagct gt gt at gat gt g at t at t ccga 900
 t gcgcgaaa acct t cct t a t ct t t gaat g gcggt act cc t gaaaat t gc ggagt gct cg 960
 aaaacaccga acccgggt ct t t ct gcgt gt cct ccct cgt gt gggat at g ct ggat at cc 1020
 cgcagacgca t ct t t gact t agt gct ccca aaact gagag ct gggaggac t cgagagggg 1080
 at ccct gcct ccccggt t g ggt gct cccc t t at ggggga aacaggt aca cggggggat c 1140
 at cccat acc t a 1152

<210> 12
 <211> 1067
 <212> DNA
 <213> Burkhol der i a A396

<400> 12
 t ct aaggaga ct gccggt ga caaaccggag gaaggt gggg at gacgt caa gt cct cat gg 60
 ccct t at ggg t agggct t ca cacgt cat ac aat ggt cgga acagagggt t gccaacccgc 120
 gagggggagc t aat cccaga aaaccgat cg t agt ccggat t gcact ct gc aact cgagt g 180
 cat gaagct g gaat cgct ag t aat cgcgga t cagcat gcc gcggt gaat a cgt t cccggg 240

6669556Seq. TXT

t ct t gt acac accgcccgt c acacat ggg agt ggg t t t accagaagt g gct agt ct aa 300
 ccgcaaggag gacggg cacc acggg aggat t cat gact gg ggt gaagt cg t aacaaggt a 360
 gccgt at cgg aaggt gcggc t ggat cacct cct t aaaccc t t t ggcct aa t aaccccggg 420
 ggaat aagt a ccgaaaaaaaa aaaaaact gg at aact t ccg t gccacaacc cgcggaaaaa 480
 t ct agggggg gggagct t aa at ggaaat t t acggggccgt aaagcgt gcg caggcggg t t 540
 gt aaacacag at gt gaaat c cccgggct t a acct gggaac t gcat t t gt g act ggcaagc 600
 t agagt at gg cacagggggg t agaat t cca cgt gt agcat t gaat gcat a gagat gagag 660
 gat accgat g gagaagggcg cccccgggga caat at gacg cct at gccac aaagct gt gg 720
 cacaat aggt t aaat acct g t gt t gt cccc gcct aaacag at t aact t g t t gt gggg at 780
 t t t ct cat aa aat act acac acgggagaat aact ggggg gct t cgt caa t t at cacaac 840
 aat gat t gcg ggcaccacg ggggt agat g ggt aat aaat cgacggcaac t at ct act t a 900
 ct t ggat gat cgcacagat t gggcgggaga gaagagaaca gcgt gt gt gt gct cct ccgc 960
 gagt gat agg t aat cggaca at act t t gac aggact t aac t gggg agcgg gat cgagt gg 1020
 at t cccgt cg gat ggcct cc gcaggt acgg cagct gggga t t acat c 1067

<210> 13
 <211> 1223
 <212> DNA
 <213> Bur khol der i a A396

<400> 13
 t t gct t acga ct t cacccca gt cat gaat c ct accgt ggt gaccgt cct c ct t gcggt t a 60
 gact agccac t t ct ggt aaa acccact ccc at ggt gt gac gggcggg gt g t acaagacc 120
 gggaacgt at t caccgcggc at gct gat cc gcgat t act a gcgat t ccag ct t cat gcac 180
 t cgagt t gca gagt gcaat c cggact acga t cggg t t t ct gggat t agct cccct cgcg 240
 ggt t ggcaac cct ct gt t cc gaccat t gt a t gacgt gt ga agccct accc at aagggcc 300
 t gaggact t g acgt cat ccc cacct t cct c cggg t t gt ca ccggcagt ct cct t agagt g 360
 ct ct t gcgt a gcaact aagg acaagggt t g cgct cgt t gc gggact t aac ccaacat ct c 420
 acgacacgag ct gacgacag ccat gcagca cct gt gt at c ggt t ct ct t t cgagcact cc 480
 cgaat ct ct t caggat t ccg accat gt caa gggg aggt aa ggt t t t t cgc gt t gcat cga 540
 at t aat ccac at cat ccacc gct t gt gggg gt ccccgct ca at t cct t t ga gt t t t aat ct 600
 t ggcaccgt a ct ccccaggc ggt caact t c acgcgt t agc t acgt t act a aggaaat gaa 660
 t cccaacaa ct agt t gaca t cgt t t aggg cgt ggact ac cagggt at ct aat cct gt t t 720
 gct ccccacg ct t t cgt gca t gagcgt cag t at t ggccca gggggct gcc t t cgccat cg 780
 gt at t cct cc acat ct ct ac gcat t t cact gct acacgt g gaat t ct acc ccct ct gcc 840
 at act ct agc t t gccagt ca caaat gcagt t cccaggt t a agcccgggga t t t cacat ct 900
 gt ct t aaaa accgcct gcg cacgct t t ac gccaggt aat t ccgat t aac gct cgcacc 960

6669556Seq. TXT

t acgt at t ac cgcggt gct ggcacgt agt t agccggt gc t t at t ct gcg gt accgt cat 1020
 cccccgggt a t agcccaaag gat t ct t t cg acaaagt gct t t acaccga t gt ct ct cac 1080
 acacgcgcat gct gat cagg t t t ccccat g t caaagt cca ct gct gct cg t aggt ct gga 1140
 cgggt t cagt t caat gt gac t gat cgt ct t t cgacaact a ct gaacgt cc ct gt agct t a 1200
 cccaccaact agct at agca t gc 1223

<210> 14
 <211> 1216
 <212> DNA
 <213> Burkhol der i a A396

<400> 14
 ccgagct gac gacagccat g cagcacct gt gt at cgggt t c t ct t t cgagc act cccgaat 60
 ct ct t cagga t t ccgacat gt caagggt a ggt aaggt t t t t cgct t gc at cgaat t aa 120
 t ccacat cat ccaccgct t g t gcgggt ccc cgt caat t cc t t t gagt t t t aat ct t gcga 180
 ccgt act ccc caggcgggt ca act t cacgcg t t agct acgt t act aaggaa at gaat cccc 240
 aacaact agt t gacat cgt t t agggcgt gg act accaggg t at ct aat cc t gt t t gct cc 300
 ccacgct t t c gt gcat gacg gt cagt at t g gcccaggggg ct gcct t cgc cat cgggt at t 360
 cct ccacat c t ct acgcat t t cact gct ac acgt ggaat t ct acccccct ct gccat act 420
 ct agct t gcc agt cacaaat gcagt t ccca ggt t aagccc ggggat t t ca cat ct gt ct t 480
 aacaaaccgc ct gcgcacgc t t t acgcca gt aat t ccga t t aacgct cg caccct acgt 540
 at t accgcgg ct gct ggcac gt agt t agcc ggt gct t at t ct t ccggt ac cgt cat cccc 600
 ccgggggt at t agcccaaagg at t t ct t t cc ggacaaaagt gct t t acaac ccgaaggcct 660
 t ct t cacaca cgcggtat t g ct ggat cagg gt t t ccccca t t gt ccaaaa t t cccact g 720
 ct gcct cccg t aggagt ct g ggccgt gt ct cagt cccagt gt ggct gat c gt cct ct cag 780
 accagct act gat cgt cgcc t t ggt aggcc t t t accccac caact agct a at cagccat c 840
 ggccaaccct at agcgcgag gt ccgaagat cccccgct t t cat ccgt aga t cgt at gcgg 900
 t at t aat ccg gct t t cgccg ggct at cccc cact acagga cat gt t ccga t gt at t act c 960
 acccgt t cgc cact cgcccc aggt gcaagc acccgt gct g ccgt t cgact t gcat gt gt a 1020
 gcat gcgcag cgt cat ct ac t aaat aaaca act ct aagaa t t t t t gcccg agggcct ct a 1080
 aacact cggg gcgt cgagag agact acgga t gaggagcat ccct ct gt ct ct aggt at gt 1140
 gt t gt cgct ct ct cacaga ggaggggacg cagcagggag ccat cgggga cgacaacat g 1200
 t acgat at ac t at ct a 1216

<210> 15
 <211> 1194
 <212> DNA
 <213> Burkhol der i a A396

<400> 15
 t t ct t cgggt a ccgt cat ccc cccgggggt at t agcccaaag gat t t ct t t c cggacaaaag 60

6669556Seq. TXT

t gct t t acaa cccgaaggcc t t ct t cacac acgcggcat t gct ggat cag ggt t t ccccc	120
at t gt ccaaa at t cccact gct gcct ccc gt aggagt ct gggccgt gt c t cagt cccag	180
t gt ggct gat cgt cct ct ca gaccagct ac t gat cgt cgc ct t ggt aggc ct t t acccca	240
ccaact agct aat cagccat cggccaacce t at agcgcga ggt ccgaaga t cccccgct t	300
t cat ccgt ag at cgt at gcg gt at t aat cc ggct t t cgcc gggct at ccc cact acagg	360
acat gt t ccg at gt at t act caccct t cg ccact cgcca ccaggt gcaa gcaccct gc	420
t gccgt t cga ct t gcat gt g t aaggcat gc cgccagcgt t caat ct gage cat gat caaa	480
ct ct gagggg gggggcct t c aacggaacga ct gggcaaaa agcgt gccca ggcgt t t t gt	540
t aagacagat gt gaaacccc ggggct t aac ct ggaaact g cat t t gt gac t ggaaagct a	600
gagt at ggca gaggggggt a gaat t ccacg t gt agcat t g aaat gcgt ag aaat ggagag	660
gaat accgat gggagagggc agccccct g ggcaaat act ggcgct t at g aacaaagt t g	720
gggcgcgccg ccgggat at g t t cccct ggg at at ccccc cct aaact gc t t acaaat at	780
t gt gt gggaa act t t t t ct c t aaaaaat ag aacacaacgg gagat at cac ccccgggggg	840
ccaccgccag at t aaacccc caaaaagt at t t ggcgggca ccccccggg gggg gagat g	900
gggt aaaat a aat ccgt gcg acgagcaaac cct cccaca cct gggat gg t cgcgaccac	960
agat gagat g cgggcggaga gaacgat acc caagcgt ggt t gt t t gcct g cat cccct cc	1020
gt cgggagt g gat at agt ag agt aat t acg gcacgact gc at t t t t t t t ct t cagt aca	1080
cct t at caca ct gt t ggat g caccgcgaga aat ccggagg t gt gagt act cccccct ct	1140
cct cgggat g t gt cggcgt ccct t ct ccc gt t caggggt gggg aagcac cgcg	1194