



(11) (21) (C) **2,258,095**
(86) 1998/01/24
(87) 1998/08/13
(45) 2000/02/22

(72) SHAH, Navnit Hargovindas, US

(72) ZELLER, Max, CH

(73) F. HOFFMANN-LA ROCHE AG, CH

(51) Int.Cl.⁶ A61K 31/365, A61K 9/48, A61K 9/20, A61K 9/16

(30) 1997/02/05 (60/037,384) US

(30) 1998/01/06 (09/003,137) US

(54) **COMPOSITIONS CONTENANT DE LA
TETRAHYDROLIPSTATINE**

(54) **TETRAHYDROLIPSTATIN CONTAINING COMPOSITIONS**

(57) L'invention concerne un produit contenant de la tetrahydrolipstatine comme ingrédient actif et des excipients acceptables sur le plan pharmaceutique, ce produit étant caractérisé par le fait qu'il se présente sous la forme de particules ayant un diamètre de 0,25 à 2 mm. L'invention concerne également des préparations ou des compositions pharmaceutiques contenant ce produit, pour une administration orale.

(57) Product containing tetrahydrolipstatin as the active ingredient and pharmaceutically acceptable excipients, characterized in that it is in the form of particles with a diameter of 0.25 to 2 mm; and pharmaceutical preparations or compositions for oral administration containing the product.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/365, 9/16	A1	(11) International Publication Number: WO 98/34607 (43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/EP98/00395 (22) International Filing Date: 24 January 1998 (24.01.98) (30) Priority Data: 60/037,384 5 February 1997 (05.02.97) US 09/003,137 6 January 1998 (06.01.98) US (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH). (72) Inventors: SHAH, Navnit, Hargovindas; 203 Beverly Hill Road, Clifton, NJ 07012 (US). ZELLER, Max; Violaweg 81, CH-4303 Kaiseraugst (CH). (74) Agent: MAHE, Jean; Grenzacherstrasse 124, CH-4070 Basle (CH).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: TETRAHYDROLIPSTATIN CONTAINING COMPOSITIONS (57) Abstract Product containing tetrahydrolipstatin as the active ingredient and pharmaceutically acceptable excipients, characterized in that it is in the form of particles with a diameter of 0.25 to 2 mm; and pharmaceutical preparations or compositions for oral administration containing the product.		

- 1 -

TETRAHYDROLIPSTATIN CONTAINING COMPOSITIONS

5 Tetrahydrolipstatin („THL“) is an inhibitor of pancreatic lipase and is known by the generic name orlistat. The use of THL as medicament, particularly as anti-obesity agent, and pharmaceutical compositions containing THL as active agent are described in US Patent No. 4,598,089.

10 Due to its low melting point of about 44°C, THL undergoes both hydrolytic and thermal degradation, particularly when stored in a humid atmosphere or above 35°C in a dry atmosphere. Furthermore, conventional dosage forms such as described in US Patent No. 4,598,089, for example, tablets or hard gelatin capsules, cannot easily be formulated from powder mix or by conventional wet granulation procedure due to picking and sticking phenomena during tablet
15 compression or encapsulation. Thus, there was a need for THL containing products and dosage forms which would be stable against moisture and heat during production and storage.

20 In one of its aspects, the present invention relates to a product containing THL as active ingredient, stabilizers, pharmaceutically acceptable excipients, characterized in that it is in the form of particles with a diameter in the range of from about 0.25 mm to about 2 mm.

25 These particles are typically in the form of particles or pellets. The preferred pellet form requires the presence of microcrystalline cellulose. Preferably, the diameter of the pellets ranges from about 0.5 to about 1.5 mm.

Surprisingly, it was found that THL containing particles do not feature the sticking and picking phenomena and exhibit superior THL stability.

5 The subject invention will now be described in terms of its preferred embodiments. These embodiments are set forth to aid in understanding the invention but are not to be construed as limiting.

10 The subject invention provides particles, such as granules and pellets, useful in producing pharmaceutical compositions, such as a unit dosage form. The use of particles in the form of pellets is preferred.

15 Surprisingly, it was found that THL containing particles (also called multiple units) minimize the sticking and picking phenomena encountered during tablet compression or encapsulation. In one of its aspects, the present invention relates to a unit dosage form comprising a plurality of pellets having a diameter in the range of from 0.25 to 2 mm wherein each particle comprises tetrahydrolipstatin, a stabilizer and at least one pharmaceutically acceptable excipient. When the particles are in the form of pellets, it is critical to employ microcrystalline cellulose.

20 The term „stabilizer“ refers to an agent having a rate of moisture uptake greater than the rate of moisture uptake for THL. Such stabilizer acts to retard hydrolytic degradation of the THL. Preferably, the stabilizer has a moisture content of greater than five percent (5%) at fifty percent (50%) relative humidity. The selection of a
25 stabilizer having the above qualities is within the skill of the artisan having read the present specification. Examples of such stabilizers include, but are not limited to, hydroxypropylmethyl cellulose, hydroxypropyl cellulose, polyvinylpyrrolidone, and lactose.

30 Preferably, the average diameter of the particles ranges from 0.5 to 1.5 mm. Diameters referred to throughout the specification are average diameters. Although it is preferred that all particles are within the

recited ranges, it is acceptable for very minor or trace amounts of undersize or oversize particles to be present.

5 A further object of the invention is a product as described above which comprises a specific combination of THL with stabilizers, and excipients, particularly a product which contains polyvinylpyrrolidone and/or lactose as stabilizers.

Preferably, the product contains either (a) lactose and at least 3% by weight of the product is polyvinylpyrrolidone or (b) at least 5% by weight of the product is polyvinylpyrrolidone.

10 Preferred compositions typically contain from about 20% to about 75% by weight THL, and from about 3% to about 60% by weight stabilizer. Preferred pellets additionally contain from about 10% to about 60% by weight microcrystalline cellulose. More preferably, such pellets contain about 25% to about 75% by weight THL; from about 20% to
15 about 60% by weight microcrystalline cellulose; from about 1% to about 10% by weight sodium starch glycolate; from about 1% to about 8% by weight sodium lauryl sulfate; from about 1% to about 10% by weight polyvinylpyrrolidone; and from about 0% to about 1% by weight talc.

20 Most preferably, such pellets contain about 50% by weight THL; about 39% by weight microcrystalline cellulose; about 3% by weight sodium starch glycolate; about 3% by weight sodium lauryl sulfate; about 5% polyvinyl pyrrolidone; and about 0.1% by weight talc.

25 Such products are chemically stable and can be filled on fast running encapsulation machines without presenting the sticking and picking phenomena.

30 In addition to the preferred stabilizers, polyvinylpyrrolidone and/or lactose, the product of the invention (the particles) contains other excipients, such as diluting agents, for example, sucrose or preferably microcrystalline cellulose (mandatory for pellets of the present invention); binders, for example, starch paste; surfactants, for example, sodium lauryl sulfate or sodium dioctylsulfosuccinate; and/or

disintegrants, for example, sodium starch glycolate. The pellets can also contain corn starch as a diluent and disintegrant, and triglycerides.

5 Pellets are preferably prepared by extrusion of a wet mass followed by spheronization. They can also be prepared by granulation on a rotating base plate in fluidized bed equipment, by agglomeration granulation on an inclined disc equipment, or in high shear mixers.

10 The process for preparing the pellets is known *per se*. See, for example, J.W. Conine and H.R. Hadley, D & CI, April 1970, p. 38-41: Small Solid Pharmaceutical Spheres; A.D. Reynold, Manufacturing Chemist & Aerosol News, June 1970, p. 40-43: A new technique for the production of spherical particles; C.W. Woodruff and N.O. Nuessle, J. Pharm. Sci. 61 (5), p. 787-790 (1972): Effect of Processing Variables on
15 Particles obtained by Extrusion-Spheronization Processing; and H.J. Malinowski and W.E. Smith, J. Pharm. Sci. 64 (10), p. 1688-1692 (1975); Use of Factorial Design to Evaluate Granulations Prepared by Spheronization.

Typically, the pelletization process by extrusion and spheronization comprises the following unit operations:

20 - wetting and kneading a powder mass containing the active substance THL and additional excipients with an appropriate, mostly aqueous solution of an appropriate binder to get a wet, pasty mass. Alternatively, the binder may be contained in the powder mix before wetting and kneading;

25 - forcing this wetted mass by means of an appropriate equipment (extruder) through tiny holes of a perforated plate, to get spaghetti-like, wet strands. The diameter of the holes may vary within wide limits, for example, between 0.4 and 1.0 mm, according to the specific product to be
30 obtained.

- breaking the strands into short pieces which are at the same time formed to more or less spherical particles by means of a fast spinning, mostly structured plate, in a vertical cylinder (spheronizer).
- 5 - drying the wet spheres by means of any suitable drying equipment, for example, a fluid bed dryer or a tray drier.
- optionally, the dried spheres may be fractionated into appropriate sized fractions, or the main material may be freed from undersized or
10 oversized material by an appropriate sieving or screening process. Preferably, the temperature is maintained below 35°C during the whole process.

15 A further aspect of the invention are pharmaceutical preparations or compositions for oral administration, comprising the particles. They can be simply filled in a PVC container from which the particles can be taken with a dosing spoon. Other oral dosage forms are sachets in which the particles are filled, alone or together with appropriate excipients, such as skim milk powder, microcrystalline cellulose,
20 sodium carboxymethylcellulose and talc, to form a powder for reconstitution. Another possibility is to embed the particle in a matrix excipient, for example, microcrystalline cellulose, followed by compression to tablets, particularly chewable tablets. The particle, can also be filled in capsules, for example, hard gelatin capsules.

25 The following Examples are illustrative but in no way limit the invention.

Example 1Preparation of Pellets

The following applies to a batch size of 4.0 kg pellets.

- 5 a) 120 g sodium lauryl sulfate and 200 g polyvinylpyrrolidone (Povidone) are dissolved under stirring in q.s. demineralized water. The solution is cooled to 10-15°C (solution A).
- 10 b) 2000 g THL, 1560 g microcrystalline cellulose and 120 g sodium starch glycolate are charged to a high speed mixer (DIOSNA, type P50) and pre-mixed for 1 minute with mixer and chopper at the low speed position "I" (pre-mix B).
- 15 c) Solution A is charged to the pre-mix B and kneaded for 2.5 minutes at mixer and chopper position "I", then the chopper is set to position "II". After 3 minutes the chopper position is again reduced to "I" for 1 minute, then the material is discharged through the outlet valve; its temperature is then between 20 and 25°C.
- 20 d) The material is fed to an extruder (NICA Lab, type E-140). The extruder is equipped with a screen with 0.8 mm nominal mesh size and thickness 1.0 mm; the screen is surrounded by a cooling device. The material is extruded to spaghettis of appropriate length. The temperatures of the extrudate and of the extruder screen are below 35°C
- 25 (extrudate D).
- e) The extrudate D is transferred in sub-batches of approx. 800 g to a spheronizer (NICA Lab type S-320) and spheronized for 3/4 - 2 minutes at 700 rpm (wet pellets E).
- 30 f) The cylinder-shaped, wet pellets E are transferred to a fluidized bed dryer (AEROMATIC, type MP-1), equipped with a dry air supply,

and dried at an incoming air temperature below 35°C to a low final moisture content of the pellets (F).

- 5 g) The dried pellets are sieved by means of a square sieve with sieve inserts 0.50 mm and 1.25 mm mesh size; the undersize-and oversize fractions are discarded; the fractions 0.5-1.25 mm are collected as THL pellets in tightly closed containers. See pellets A in Table 1 below.

Example 2

10

Preparation of Pellets

The following applies to a batch size of 6.0 kg pellets.

- 15 a) 180 g sodium lauryl sulfate and 300 g Povidone are dissolved in q.s. demineralized water by means of a stirrer. The solution is cooled down to approx. 10-15°C by putting the beaker into a mixture of ice and water (A).
- 20 b) 1800 g THL, 3120 g microcrystalline cellulose and 600 g sodium starch glycolate are given to a high speed mixer (DIOSNA, type P50) and pre-mixed for 1 minute with mixer and chopper at the low speed position "I" (B).
- 25 c) Solution A is given to the pre-mix B and kneaded for 2.5 minutes at mixer and chopper position "I", then the chopper is set to position "II". After 3 minutes, the chopper position is again reduced to "I" for 1 minute, then the material is discharged through the outlet valve; its temperature is then between 20 and 25°C (C).
- 30 d) The material is fed to an extruder and treated as in Example 1d).
- e) The resulting extrudate is transferred in sub-batches of approx. 800 g to a fluidized bed dryer (AEROMATIC, type MP-1), equipped with

a roto-granulation device and a dry air supply, and spheronized for 3 minutes at 500 rpm (E).

5 f) The cylinder-shaped, wet pellets E are then dried in the same equipment, at an incoming air temperature below 35°C, to a low final moisture content of the pellets (F).

10 g) The dried pellets F are sieved by means of a square sieve with sieve inserts of 0.50 mm and 1.25 mm mesh size; the undersize- and oversize-fractions are discarded; the fractions 0.5-1.25 mm are collected as THL pellets in tightly closed containers. See pellets B below.

The compositions of pellets A and pellets B are given in Table 1:

Table 1

15

Pellet Formulation (in weight-%)	A	B
<u>Active Drug:</u>		
THL finely milled	50.0	30.0
<u>Excipients:</u>		
Microcrystalline cellulose	39.0	52.0
Sodium starch glycolate	3.0	10.0
Sodium lauryl sulfate	3.0	3.0
Povidone	5.0	5.0
Total	100.0	100.0

Example 3

Preparation of Pellets

20

Pellets with the following compositions were prepared in a manner similar to that of pellets A and B:

Table 2

Pellet Formulation (in weight-%)	F1	F2	F3	F4	F5	F6
THL	29.1	30.2	50	50	50	50
Lactose	33.5	34.0	7	21	12	
Microcrystalline cellulose	23.6	20.8	36	19.5	21	41
Sodium starch glycolate	9.1	9.4	3	3	9.5	3
Sodium lauryl sulfate	1.1	1.1	1	1	3	1
(Povidone)	3.6	4.5	3	3		5
Polyvinylpyrrolidone					4.5	
Medium chain triglyceride				2.5		

Example 4

5

Preparation of Final Dosage Form

The pellets A or B (of Examples 1 and 2 above) are either mixed with talc and encapsulated on equipment provided with a pellet fill station (formulations C, E);

10 or a mixture of pellets A or B with skim milk powder, microcrystalline cellulose and talc is filled into sachets (formulation D).

A similar mixture is compressed to chewable tablets (formulation F).

15 The compositions of the final dosage forms C, D, E, F are given in Table 3.

Table 3: Final Dosage Form Compositions (in mg per dose)

Formulation	C	D	E	F	
Final Active Drug Content	120.0	60.0	30.0	30.0	mg
<u>Active Drug in Form of Pellets:</u>					
Pellets A	240.0	120.0	--	--	mg
Pellets B	--	--	100.0	100.0	mg
<u>Excipients:</u>					
Skim milk powder (granulated)		3875.0			mg
Microcrystalline cellulose +_sodium carboxymethyl- cellulose (AVICEL RC- types)		1000.0			mg
Microcrystalline cellulose				1888.0	mg
Talc	0.24	5.0	0.1	2.0	mg
Magnesium stearate				10.0	mg
Total Hard Gelatin Capsule Fill Weight	240.24		100.1		mg
Total Sachet Fill Weight		5000.0			mg
Chewable Tablet Weight				2000.0	mg

The detailed composition of formulation C for capsules filled with pellets A is as in Table 4:

Table 4: Production Batch Size = 160.160 kg = 666,666 capsules

Fill Weight = 240.24 mg

Ingredient	Quantity mg/Capsule	666,666 Capsules Contain (kg)
THL	120.00	80.000
Microcrystalline cellulose (AVICEL PH-101)	93.60	62.400
Sodium starch glycolate (PRIMOJEL)	7.20	4.800
Sodium lauryl sulfate	7.20	4.800
Polyvinylpyrrolidone (Povidone) (K-30)	12.00	8.000
Talc	0.24	0.160
Total	240.24 mg	160.160 kg

5

Example 5Preparation of Granules and Formulations containing THL

Ingredient	mg/capsule
1. THL	120
2. Microcrystalline cellulose	93.6
3. Sodium starch glycolate	7.2
4. Polyvinyl pyrrolidone	12.0
5. Sodium lauryl sulfate	7.2
Total	240.00

1. Polyvinyl pyrrolidone and sodium lauryl sulfate is dissolved in water.
2. THL, microcrystalline cellulose, and sodium starch glycolate are mixed for 10 minutes and granulated with the solution of step 1.
- 5 3. Granules are dried at or below 30°C and passed through # 20 mesh screen.
4. Granules are filled in a # 1 hard gelatin capsule.

Example 6

10

Preparation of Granules and Formulations containing THL

Ingredient	mg/capsule
1. THL	120
2. Lactose anhydrous	93.6
3. Sodium starch glycolate	7.2
4. Polyvinyl pyrrolidone	12.0
5. Sodium lauryl sulfate	7.2
Total	240.00

1. Polyvinyl pyrrolidone and sodium lauryl sulfate is dissolved in water.
- 15 2. THL, lactose anhydrous, and sodium starch glycolate is mixed for 10 minutes and granulated with the solution of step 1.
3. Granules are dried at or below 30°C and passed through # 20 mesh screen.
4. Granules are filled in a # 1 hard gelatin capsule.

The subject invention has been described in terms of its preferred embodiments. Upon reading the specification, a skilled artisan will become cognizant of various alternative embodiments. These variations are to be considered within the scope and spirit of the invention which is
5 only to be limited by the claims that follow and their equivalents.

Claims

1. A pharmaceutical composition comprising a plurality of particles having an average diameter of from about 0.25 mm to about 2 mm, each particle comprising tetrahydrolipstatin, a stabilizer, and at least one pharmaceutically acceptable excipient, provided that when the particles are in the form of pellets, each pellet contains microcrystalline cellulose.

2. The composition of claim 1, wherein the particles are pellets or granules, particularly wherein the pellets or granules have an average diameter of from about 0.5 mm to about 1.5 mm.

3. The composition of claim 2, wherein the stabilizer is selected from the group consisting of polyvinylpyrrolidone, lactose, a combination of polyvinylpyrrolidone and lactose, hydroxypropylmethyl cellulose, and hydroxypropyl cellulose.

4. The composition of claim 3, wherein at least 5% by weight of the composition is polyvinylpyrrolidone.

5. The composition of claim 3, wherein the stabilizer is a combination of lactose and polyvinylpyrrolidone wherein at least 3% by weight of the composition is polyvinylpyrrolidone.

6. The composition of claim 2, wherein the pharmaceutically acceptable excipient is a surfactant, diluent, or disintegrant.

7. The composition of claim 6, wherein the pharmaceutically acceptable excipient is a surfactant, particularly wherein the surfactant is sodium lauryl sulfate or sodium dioctylsulfosuccinate.

8. The composition of claim 6, wherein the pharmaceutically acceptable excipient is a diluent, particularly wherein the diluent is microcrystalline cellulose, sucrose or corn starch.

WO 98/34607

PCT/EP98/00395

- 15 -

9. The composition of claim 6, wherein the excipient is a disintegrant, particularly wherein the disintegrant is sodium starch glycolate, cross-linked polyvinyl pyrrolidone, cross-carmolose sodium, or low-substituted hydroxypropyl cellulose.

5 10. The composition of claim 2 which comprises from about 20% to about 75% by weight tetrahydrolipstatin; from about 3% to about 60% by weight stabilizer; and from about 10% to about 60% by weight microcrystalline cellulose.

10 11. The composition of claim 10 which comprises about 25% to about 75% by weight tetrahydrolipstatin; from about 20% to about 60% by weight microcrystalline cellulose; from about 1% to about 10% by weight sodium starch glycolate; from about 1% to about 8% by weight sodium lauryl sulfate; from about 1% to about 10% by weight polyvinylpyrrolidone; and from about 0% to about 1% by weight talc.

15 12. The composition of claim 11 which comprises about 50% by weight tetrahydrolipstatin; about 39% by weight microcrystalline cellulose; about 3% by weight sodium starch glycolate; about 3% by weight sodium lauryl sulfate; about 5% by weight polyvinylpyrrolidone; and about 0.1% by weight talc.

20 13. The composition of any one of claims 1, 2 and 10-12 which is in unit dosage form.

25 14. A pharmaceutical composition as in claim 1, which comprises about 120 mg of tetrahydrolipstatin; about 93.6 mg of microcrystalline cellulose; about 7.2 mg of sodium starch glycolate; about 7.2 mg of sodium lauryl sulfate; about 12 mg polyvinylpyrrolidone; and about 0.24 mg talc.

15. The composition of claim 14 which is in unit dosage form.

-16-

16. The use of a pharmaceutical composition as set out in any one of claims 1-15 for oral administration, as an inhibitor of pancreatic lipase.

17. The use of a pharmaceutical composition as set out in any one of claims 1-15 for oral administration, as an anti-obesity agent.