

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 2675794 T3**

-
- (51) Int.Cl.: **C 07 D 239/42 (2006.01)** **A 61 K 31/506 (2006.01)** **A 61 P 35/00 (2006.01)**
C 07 D 401/12 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2019-05-06**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-02-13**
- (86) Europæisk ansøgning nr.: **12705384.1**
- (86) Europæisk indleveringsdag: **2012-02-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-12-25**
- (86) International ansøgning nr.: **GB2012000176**
- (87) Internationalt publikationsnr.: **WO2012110774**
- (30) Prioritet: **2011-02-17 US 201161443773 P** **2011-08-15 US 201161523489 P**
2011-12-23 US 201161579729 P
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Cancer Therapeutics Crc Pty Limited, 4 Research Avenue, Bundoora, VIC 3083, Australien**
- (72) Opfinder: **HOLMES, Ian, Peter, Cancer Therapeutics CRC Pty Limited, 4 Research Avenue, Bundoora, VIC 3083, Australien**
BERGMAN, Ylva, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
LUNNISS, Gillian Elizabeth, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
NIKAC, Marica, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
CHOI, Neil, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
HEMLEY, Catherine Fae, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
WALKER, Scott Raymond, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
FOITZIK, Richard Charles, Monash Institute of Pharmaceutical Sciences, 381 Royal Parade, Parkville, VIC 3052, Australien
GANAME, Danny, WEHI Biotechnology Centre, 4 Research Avenue, Bundoora, VIC 3086, Australien
LESSENE, Romina, WEHI Biotechnology Centre, 4 Research Avenue, Bundoora, VIC 3086, Australien
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **SELEKTIVE FAK-INHIBITORER**
- (56) Fremdragne publikationer:
WO-A1-2004/080980
WO-A1-2010/055117

Fortsættes ...

WO-A1-2012/022408

WO-A2-2010/058032

Kinoshita et al. Am J Respir Crit Care Med 183, 2011: A6056.

Wong et al. Nat Med. Author manuscript; 18(1): 148-152; available in PCM 2015 June 05.

Mimura et al. Journal Investigative Dermatology 124, 2005, 886-892.

Clemente et al. Circ Res. 2007; 101: 1339-1348.

Vittal et al. AJP 2005, 166, 367-375.

Dun et al. World J Gastroenterol 2010; 16: 4100-4106.

DESCRIPTION

[0001] This invention relates to 2,4,5-substituted pyrimidines that inhibit Focal Adhesion Kinase (FAK), also known as protein tyrosine kinase 2 (PTK2), and to pharmaceutical compositions containing such compounds. This disclosure also relates to a method of using such compounds for the prevention and/or treatment of proliferative diseases, such as cancer.

Background

[0002] Directional cell migration is important in many physiological and pathological processes including embryonic development, wound healing, angiogenesis, tumour invasion and metastasis. Transduction of extracellular signals, that stimulate cells to move directionally, may be induced by a number of processes including transmembrane integrins binding to extra cellular matrix proteins and the action of growth factors (for example EGF, IGF and VEGF) on the extracellular domains of their cognate receptors.

[0003] FAK is a non receptor tyrosine kinase that mediates signals from both transmembrane integrins and growth factor receptors. FAK has been reported to play a central role in coordinating these diverse extra cellular signals, integrating them in a fashion that results in directional movement of cells through their external environment (Tomar and Schlaepfer. Current Opinion in Cell Biology: 2009, 21, 676-683).

[0004] Integrin clustering or the activation of a growth factor receptor (for example EGFR, IGF-1R, Her2 and VEGFR) promotes FAK autophosphorylation at Y397.

[0005] Phosphorylated Y397 FAK then binds to c-Src (referred to as Src herein) and Src mediated phosphorylation of FAK at Y576 and Y577 occurs to give rise to an active FAK-Src complex. Active FAK-Src then facilitates signaling via a number of biochemical pathways which influence processes such as cell adhesion, migration, invasion, cell survival, proliferation, acquisition of chemotherapy resistance and metastasis (Brunton and Frame. Current Opinion in Pharmacology: 2008, 8, 437-432 and Chatzizacharias et al. Expert Opinion in Therapeutic Targets: 2007, 11(10), 1315-1328).

Cell adhesion

[0006] Functional studies addressing the role of FAK in cell adhesion suggest that it contributes to both focal adhesion assembly (Richardson and Parsons. Nature: 1996, 380, 538-540) and focal adhesion turnover (Fincham et al. Oncogene: 1995, 10(11), 2247-2252). Inhibition of FAK by RNAi in both human and mouse cell lines, resulting in decreased FAK protein levels, has been shown to reduce cell adhesion to a fibronectin/laminin-coated plate *in*

vitro (Tsutsumi et al. International Journal of Oncology: 2008, 33(1), 215-224).

Cell migration

[0007] There is strong evidence that FAK is a key regulator of cell migration (Angelucci and Bologna. Current Pharmaceutical Design: 2007, 13, 2129-2145 and Mitra et al. Nature Reviews Molecular Cell Biology: 2005, 6, 56-68). Cells derived from FAK $-/-$ mouse embryos exhibit reduced migration as a result of impaired adhesion turnover (Ilić et al. Nature: 1995, 377, 539-544). Moreover, displacement of FAK from focal adhesions reduces cell migration (Gilmore and Romer. Molecular Biology of the Cell: 1996, 7(8), 1209-1224), whilst over-expression in CHO cells stimulates migration (Cary et al. Journal of Cell Science: 1996, 7, 1787-1794). In addition, inhibition of FAK by RNAi in both human and mouse cell lines, resulting in decreased FAK protein levels, has been shown to reduce cell migration in an *in vitro* haptotactic migration assay (Tsutsumi et al. International Journal of Oncology: 2008, 33(1), 215-224).

Cell invasion

[0008] FAK activation has been shown to enhance matrix degrading invasive behaviour. FAK-Src signaling through cellular apoptosis susceptibility protein (CAS) (Liao et al. Journal of Experimental and Clinical Cancer Research: 2008, 27:15) leads to the expression of matrix metalloproteases (MMPs) including MMP2 and MMP9. FAK-Src activation also promotes cell surface expression of MMP14 via phosphorylation of endophilin A2. MMP14 then activates MMP2 by cleavage of pro-MMP2 to its active form (Siesser and Hanks. Clinical Cancer Research: 2006, 12(11), 3233-3237). Highly invasive cancer cells form specialized actin-rich extra cellular matrix degrading membrane protrusions known as invadopodia which are rich in matrix-degrading proteases such as MMPs. Both FAK and Src have been shown to be instrumental in the formation of invadopodia (Chan et al. Journal of Chemical Biology: 2009, 185(2), 357-370).

Cell survival

[0009] FAK has been shown to play an important role in cell survival. Activation of FAK has been shown to result in suppression of anoikis (apoptosis in response to an inappropriate extra cellular matrix environment) (Frisch et al Journal of Cell Biology. 1996, 134(3), 793-799 and Xu et al Cell Growth and Differentiation. 1996, 7(4), 413-418). Studies have demonstrated that FAK activates multiple downstream pathways to suppress anoikis in both fibroblasts and epithelial cells (Zouq et al. Journal of Cell Science: 2008, 122, 357-367). In human intestinal crypt cells signalling via the association of FAK with $\beta 1$ integrin and subsequent binding with Src up regulates expression of the anti-apoptotic proteins Bcl-X_L and Mcl-1 via PI3-K/Akt-1


signalling. PI3-K/Akt-1 signalling also down regulates expression of the pro-apoptotic activators Bax and Bak, causes phosphorylation of the pro-apoptotic sensitizer Bad and antagonizes p38 β activation. Dissociation of FAK/Src results in a sustained/enhanced activation of p38 β which is an apoptosis/anoikis driver (Bouchard et al. Apoptosis: 2008, 13, 531-542).

Cell proliferation

[0010] Reduction in the expression of either FAK or β 1 integrin and hence disruption of the β 1-FAK signalling axis results in decreased initial proliferation of micro-metastatic cells distributed in the lung. Using 3D cultured D2 cells a strong correlation was observed between FAK Y397 and Y861 phosphorylation and proliferative ability (Shibue and Weinberg. PNAS 2009, 106(25), 10290-10295). HL-60 Cells, transfected to over express FAK, have been shown to double at a rate 1.5 times faster than control HL-60 cells. Studies revealed a marked induction of cyclin D3 expression and CDK activity in the cells over expressing FAK. Activation of PI3-K/Akt-1 signalling, a process associated with FAK activation in a number of studies, was identified as a probable cause of the cyclin expression/activation (Yamamoto et al. Cellular Signaling: 2003, 15. 575-583).

Acquisition of chemotherapy resistance

[0011] Exposure of the cisplatin sensitive ovarian cancer cell line OAW42 to repeated cycles of cisplatin treatment and subsequent recovery resulted in the formation of chemo-resistant OAW42-R cells. Studies aimed at identifying the cause of this chemo-resistance revealed that FAK was constitutively active in both the sensitive and chemo-resistant cells. However, inhibition of phosphorylation of Y397 FAK was induced by treatment with cisplatin in OAW42 cells but not in OAW42-R cells (Poulain and co-workers. Gynaecologic oncology: 2006, 101, 507-519). The effects of FAK inhibition on chemo-resistance has also been studied *in vitro* and *in vivo* using the FAK inhibitor TAE226, alone and in combination with docetaxel, in taxane-sensitive (SKOV3ip1 and HeyA8) and taxane-resistant (HeyA8-MDR) ovarian cancer cell lines. TAE226 has the structure:



and is described in WO 2004/080980 and WO 2005/016894. *In vitro*, TAE226 inhibited the phosphorylation of FAK at both Y397 and Y861 sites, inhibited cell growth in a time- and dose-dependent manner, and enhanced docetaxel-mediated growth inhibition by 10- and 20-fold in the taxane-sensitive and taxane-resistant cell lines, respectively. *In vivo*, FAK inhibition by TAE226 significantly reduced tumour burden in the HeyA8, SKOV3ip1, and HeyA8-MDR models (46-64%) compared with vehicle-treated controls. However, the greatest efficacy was

observed with concomitant administration of TAE226 and docetaxel in all three models (85-97% reduction). In addition, TAE226 in combination with docetaxel significantly prolonged survival in tumour-bearing mice (Halder et al. Cancer Res: 2007, 67(22), 10976-10983).

Metastatic potential

[0012] Several studies have examined the role of FAK protein levels and its relation to tumor progression in animal models. In a mouse skin carcinogenesis model using FAK +/- mice, reduced FAK protein expression correlated with decreased papilloma formation (46%), compared with FAK +/+ wild-type control mice (McLean et al. Cancer Research: 2001, 61, 8385-8389). Using human breast carcinoma cells, researchers showed that FAK siRNA treated cells were inhibited from metastasizing to the lung after orthotopic implantation in nude mice (Benlimame et al. Journal of Cell Biology: 2005, 171, 505-516). Similar experiments using short hairpin RNA (shRNA) against FAK in 4T1 mouse breast carcinoma cells resulted in an inhibition of metastasis to the lungs after orthotopic implantation in mammary pads (Mitra et al. Oncogene: 2006, 25, 4429-4440). Inhibition of FAK by dominant negative expression in 4T1 mouse breast carcinoma cells reduced tumour growth and angiogenesis in mice (Mitra et al. Oncogene: 2006, 25, 5969-5984). Use of a Cre/loxP recombination system to disrupt FAK function in the mammary epithelium of a transgenic model of breast cancer has demonstrated that FAK expression is required for the transition of premalignant hyperplasias to carcinomas and their subsequent metastases. The observed decrease in tumor progression was further correlated with impaired mammary epithelial proliferation suggesting that FAK plays a critical role in mammary tumor progression (Lahlou et al. PNAS USA: 2007, 104(51), 20302-20307).


[0013] In accordance with the above observations over expression of FAK mRNA and/or protein has been reported in numerous human cancers including colorectal cancer (de Heer. European Journal of Surgical Oncology: 2008, 34(11), 1253-1261), prostate cancer (Tremblay, L., W. Hauck, et al. International Journal of Cancer: 1996, 68(2), 164-171), breast cancer (Watermann et al. British Journal of Cancer 2005, 93(6), 694-698) and melanomas (Hess et al. Cancer Research: 2005, 65(21), 9851-60). Furthermore FAK over expression is frequently correlated with more aggressive phenotypes of these cancers.

[0014] Thus, there is strong evidence to suggest that a FAK inhibitor would have application for the reduction of cell adhesion, cell migration, cell invasion, cell proliferation and chemoresistance. Furthermore, a FAK inhibitor would have applicability to induce apoptosis for cells in inappropriate extra cellular matrix environments and reduce angiogenesis.


[0015] A compound which is a selective FAK inhibitor would enable the targeting of specific biological pathways, without any potential issues caused by the inhibition of any targets, such as other protein kinases.

[0016] Accordingly, compounds that selectively inhibit FAK would be useful for the treatment of proliferative diseases, such as cancer.

[0017] Two compounds reported to inhibit FAK are PF-562,271 and PF-573,228.



PF-562,271



PF-573,228

[0018] PF-562,271 is described in WO2004/056786, WO2004/056807, WO2005/023780, WO2007/063384 and Roberts et al. Cancer Res 2008, 68(6), 1935-1944.


[0019] PF-573,228 is described in Slack-Davis et al. J. Biol. Chem. 2007, 282(20), 14845-14852.

[0020] In addition to these specifically described compounds, further classes of FAK inhibitors are disclosed in WO2008/129380, WO2008/115369, WO2009/105498, US2010/113475, WO2009/143389, WO2009/071535, WO2010/055117, WO2010/058030, WO2010/058032, WO2007/140222, and WO2009/024332.

Summary of the invention


[0021] The present inventors have discovered a particular class of compounds which are effective as selective FAK inhibitors. These compounds may exhibit selectivity for FAK over kinases such as IGF-1R (Insulin-like growth factor 1 receptor), IR (insulin receptor) and CDKs (cyclin-dependent kinases), as well as over other kinases, such as VEGFR1, VEGFR2 and VEGFR3. Additionally, the compounds of the invention may have enhanced selectivity for the inhibition of cytochrome p450 enzymes, specifically the 2C9 and 3A4 isoforms. Furthermore, the compounds of the invention may be less prone to the formation of adducts with glutathione.

[0022] In a first aspect, the present invention provides compounds of the following formula (I):



wherein:

R¹ is selected from: H and



wherein:


R^{N1} is selected from H, C_{1-3} alkyl and $C(=O)Me$;

R^{N2} is selected from H, C_{1-3} alkyl and $C(=O)Me$;

R^{N3} is selected from H, C_{1-3} alkyl and $C(=O)Me$;

R^{N4} is selected from H and CH_3 ;

R^2 is selected from H and



wherein:

R^{N5} is selected from H, C_{1-3} alkyl and $C(=O)Me$;


R^{N6} is selected from H, C_{1-3} alkyl and $C(=O)Me$;

and wherein only one of R^1 and R^2 is H;

R^3 is selected from O- C_{1-2} alkyl, C_{1-2} alkyl, halo, cyano, where the C_{1-2} alkyl group may be substituted by one or more fluoro groups;

R^4 is selected from CF_3 , halo, CF_2H and CN; and

R^5 is selected from groups of the following formulae:



wherein:

R^6 is selected from H, $(CHR^{C1})_{n1}C(O)N(R^{N7})Z^1$ and $(CH_2)_{n2}C(O)OZ^2$; wherein:

$n1$ is 1;

R^{C1} is H or Me;

R^{N7} is H or CH_3 ;

Z^1 is H, CH_3 or OCH_3 ;

n_2 is 1; and

Z^2 is CH_3 ;

and where only one of R^{N7} and Z^1 can be CH_3 ,

R^7 is selected from H, and $(\text{CH}_2)_{m1}\text{C}(\text{O})\text{N}(\text{R}^{M1})\text{Y}^1$, wherein:

m_1 is 0 or 1;

R^{M1} is H; and

Y^1 is H, Me or OCH_3 ;

and only one of R^6 and R^7 is H; and

R^8 is H or, when R^7 is $\text{C}(=\text{O})\text{NH}_2$, R^8 is selected from H and C_{1-2} alkyl.

[0023] A second aspect of the present invention provides a composition comprising a compound of the first aspect and a pharmaceutically acceptable carrier or diluent.

[0024] A third aspect of the invention provides a compound of the first aspect for use in a method of therapy.

[0025] A further aspect of the invention provides a compound of the first aspect for use in the method of treatment of cancer.


[0026] Each of the groups R^1 to R^8 will be discussed in more detail below.

R^1


R^1 may have one of the following structures:




(R^{1a})




(R^{1b})



(R^{1c})



(R^{1d})



(R^{1e})


When R^1 is H, R^2 (discussed below) is not H.

Each of R^{N1} , R^{N2} and R^{N3} is independently selected from H, C_{1-3} alkyl (i.e. methyl, ethyl, prop-

1-yl and prop-2-yl) and C(=O)Me and R^{N4} is selected from either H or methyl.

R^2

R^2 may have one of the following structures:



When R^2 is H, R^1 (discussed above) is not H.

R^{N5} and R^{N6} are independently selected from H, C_{1-3} alkyl (i.e. methyl, ethyl, prop-1-yl and prop-2-yl) and C(=O)Me.

R^3

R^3 is selected from $O-C_{1-2}$ alkyl, C_{1-2} alkyl, halo, cyano, where the C_{1-2} alkyl group may be substituted by one or more fluoro groups. Thus, R^3 may be:

1. a) halo: F, Cl, Br, I;
2. b) cyano: CN
3. c) C_{1-2} alkyl, optionally substituted by one or more fluoro groups: CH_3 , CH_2F , CHF_2 , CF_3 , CH_2CH_3 , CH_2CH_2F , CH_2CHF_2 , CH_2CF_3 , $CHFCH_3$, $CHFCH_2F$, $CHFCHF_2$, $CHFCHF_3$, CF_2CH_3 , CF_2CH_2F , CF_2CHF_2 , CF_2CF_3 ;
4. d) $O-C_{1-2}$ alkyl, wherein the C_{1-2} alkyl group is optionally substituted by one or more fluoro groups: $O-CH_3$, $O-CH_2F$, $O-CHF_2$, $O-CF_3$, $O-CH_2CH_3$, $O-CH_2CH_2F$, $O-CH_2CHF_2$, $O-CH_2CF_3$, $O-CHFCH_3$, $O-CHFCH_2F$, $O-CHFCHF_2$, $O-CHFCHF_3$, $O-CF_2CH_3$, $O-CF_2CH_2F$, $O-CF_2CHF_2$, $O-CF_2CF_3$.


R^4

R^4 is selected from CF_3 , halo (i.e. F, Cl, Br, I), CF_2H and CN.

In some embodiments, the halo group is either Cl or Br.

R^5

R^5 is selected from groups of the following formulae:



R^6

R^6 is selected from H, $(CHR^{C1})_{n1}C(O)N(R^{N7})Z^1$ and $(CH_2)_{n2}C(O)OZ^2$; wherein:

n_1 is 1;

R^{C1} is H or Me;

R^{N7} is H or CH_3 ;

Z^1 is H, CH_3 or OCH_3 ;

n_2 is 1; and

Z^2 is CH_3 ;

wherein only one of R^{N7} and Z^1 may be CH_3 .

When R^6 is H, R^7 (discussed below) is not H.

If R^6 is $(CHR^{C1})_{n1}C(O)N(R^{N7})Z^1$, it may be selected from: $CH_2C(O)NH_2$, $CH_2C(O)NHCH_3$, $CH_2C(O)NHOCH_3$, $CH_2C(O)NCH_3OCH_3$, $CHCH_3C(O)NH_2$, $CHCH_3C(O)NHCH_3$, $CHCH_3C(O)NHOCH_3$, and $CHCH_3C(O)NCH_3OCH_3$.

If R^6 is $(CH_2)_{n2}C(O)OZ^2$, it is $CH_2C(O)OCH_3$.

R^7

H, and $(CH_2)_{m1}C(O)N(R^{M1})Y^1$, wherein:

m_1 is 0 or 1;

R^{M1} is H; and

Y^1 is H, Me or OCH_3 ;

When R^7 is H, R^6 (discussed above) is not H.

When R^7 is $(CH_2)_{m1}C(O)N(R^{M1})Y^1$, it may be selected from $C(O)NH_2$, $C(O)NHCH_3$, $C(O)NHOCH_3$, $CH_2C(O)NH_2$, $CH_2C(O)NHCH_3$ and $CH_2C(O)NHOCH_3$.

R^8

R^8 is H, except for when R^7 is $C(=O)NH_2$, it may alternatively be C_{1-2} alkyl. i.e. methyl or ethyl.

Includes Other Forms

[0027] Included in the above are the well known ionic, salt, and solvate forms of these substituents. For example, a reference to carboxylic acid ($-COOH$) also includes the anionic


(carboxylate) form (-COO^-), as well as a salt or solvate thereof. Similarly, a reference to an amino group includes the protonated form ($\text{-N}^+\text{HR}^1\text{R}^2$), as well as a salt or solvate of the amino group, for example, a hydrochloride salt. Similarly, a reference to a hydroxyl group also includes the anionic form (-O^-), as well as a salt or solvate thereof.

Isomers, Salts, and Solvates

[0028] Certain compounds may exist in one or more particular geometric, optical, enantiomeric, diastereomeric, epimeric, stereoisomeric, tautomeric, conformational, or anomeric forms, including but not limited to, cis- and trans-forms; E- and Z-forms; c-, t-, and r-forms; endo- and exo-forms; R-, S-, and meso-forms; D- and L-forms; d- and l-forms; (+) and (-) forms; keto-, enol-, and enolate-forms; syn- and anti-forms; synclinal- and anticlinal-forms; α - and β -forms; axial and equatorial forms; boat-, chair-, twist-, envelope-, and halfchair-forms; and combinations thereof, hereinafter collectively referred to as "isomers" (or "isomeric forms").

[0029] Note that, except as discussed below for tautomeric forms, specifically excluded from the term "isomers", as used herein, are structural (or constitutional) isomers (i.e. isomers which differ in the connections between atoms rather than merely by the position of atoms in space). For example, a reference to a methoxy group, -OCH_3 , is not to be construed as a reference to its structural isomer, a hydroxymethyl group, $\text{-CH}_2\text{OH}$. Similarly, a reference to ortho-chlorophenyl is not to be construed as a reference to its structural isomer, meta-chlorophenyl. However, a reference to a class of structures may well include structurally isomeric forms falling within that class (e.g., C_{1-7} alkyl includes n-propyl and iso-propyl; butyl includes n-, iso-, sec-, and tert-butyl; methoxyphenyl includes ortho-, meta-, and para-methoxyphenyl).

[0030] The above exclusion does not pertain to tautomeric forms, for example, keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, N-nitroso/hydroxyazo, and nitro/aci-nitro.



Note that specifically included in the term "isomer" are compounds with one or more isotopic substitutions. For example, H may be in any isotopic form, including ^1H , ^2H (D), and ^3H (T); C may be in any isotopic form, including ^{12}C , ^{13}C and ^{14}C ; O may be in any isotopic form, including ^{16}O and ^{18}O ; and the like.

[0031] Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including (wholly or partially) racemic and other mixtures thereof. Methods for

the preparation (e.g. asymmetric synthesis) and separation (e.g., fractional crystallisation and chromatographic means) of such isomeric forms are either known in the art or are readily obtained by adapting the methods taught herein, or known methods, in a known manner.

[0032] Unless otherwise specified, a reference to a particular compound also includes ionic, salt, and solvate forms thereof, for example, as discussed below.

[0033] It may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the active compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge et al. J. Pharm. Sci., 66, 1-19 (1977).

[0034] For example, if the compound is anionic, or has a functional group which may be anionic (e.g., -COOH may be -COO^-), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Na^+ and K^+ , alkaline earth cations such as Ca^{2+} and Mg^{2+} , and other cations such as Al^{3+} . Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e., NH_4^+) and substituted ammonium ions (e.g., NH_3R^+ , NH_2R_2^+ , NHR_3^+ , NR_4^+). Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $\text{N}(\text{CH}_3)_4^+$.

[0035] If the compound is cationic, or has a functional group which may be cationic (e.g., -NH_2 may be -NH_3^+), then a salt may be formed with a suitable anion. Examples of suitable inorganic anions include, but are not limited to, those derived from the following inorganic acids: hydrochloric, hydrobromic, hydroiodic, sulphuric, sulphurous, nitric, nitrous, phosphoric, and phosphorous. Examples of suitable organic anions include, but are not limited to, those derived from the following organic acids: acetic, propionic, succinic, glycolic, stearic, palmitic, lactic, malic, pantoic, tartaric, citric, gluconic, ascorbic, maleic, hydroxymaleic, phenylacetic, glutamic, aspartic, benzoic, cinnamic, pyruvic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, phenylsulfonic, toluenesulfonic, methanesulfonic, ethanesulfonic, ethane disulfonic, oxalic, pantothenic, isethionic, valeric, lactobionic, and gluconic. Examples of suitable polymeric anions include, but are not limited to, those derived from the following polymeric acids: tannic acid, carboxymethyl cellulose.

[0036] It may be convenient or desirable to prepare, purify, and/or handle a corresponding solvate of the active compound. The term "solvate" is used herein in the conventional sense to refer to a complex of solute (e.g. active compound, salt of active compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a tri-hydrate, etc.

Selectivity

[0037] The selectivity of the compounds for inhibiting FAK over other kinases such as VEGFR1, VEGFR2 and/or VEGFR3, IGF-1R, IR and CDKs can be demonstrated by biochemical assay results (see, for example, the FAK kinase assay and VEGFR3 assays described below).

[0038] The selectivity of the compounds for FAK over the inhibition of cytochrome p450 enzymes, specifically the 2C9 and 3A4 isoforms may be determined using standard inhibition assays.


[0039] How prone the compounds of the invention may be to the formation of adducts with glutathione may be determined by the protocol described in Walker, et al. *Biorg. Med. Chem. Letts.* 2008, 18, 6071-6077.

[0040] The selectivity of compounds for inhibiting FAK over other enzymes may be expressed as ratio of other enzyme's inhibition (IC_{50}) to the FAK inhibition (IC_{50}). For example, to determine the selectivity of a compound for inhibiting FAK over VEGFR3, the compound's IC_{50} for VEGFR3 is divided by the compound's IC_{50} for FAK to give a ratio. The higher the ratio, the more selective the compound is.

Further Embodiments


[0041] The following embodiments and preferences may be combined with one another as appropriate.

[0042] In some embodiments, R^2 is H and R^1 is:




wherein R^{N1} is selected from H, C_{1-3} alkyl (i.e. methyl, ethyl, prop-1-yl and prop-2-yl) and $C(=O)Me$. In some of these embodiments, it may be preferred that R^{N1} is $C(=O)Me$. In others of these embodiments, it may be preferred that R^{N1} is H, methyl or ethyl. In further of these embodiments, it may be preferred that R^{N1} is H.

[0043] In other embodiments, R^2 is H and R^1 is:




wherein R^{N2} is selected from H and C_{1-3} alkyl (i.e. methyl, ethyl, prop-1-yl and prop-2-yl). In these embodiments, it may be preferred that R^{N2} is selected from H and methyl. In these embodiments, it may also be preferred that R^{N2} is ethyl.

[0044] In other embodiments, R^2 is H and R^1 is:




wherein R^{N3} is selected from H and C_{1-3} alkyl (i.e. methyl, ethyl, prop-1-yl and prop-2-yl). In these embodiments, it may be preferred that R^{N3} is selected from H and methyl.

[0045] In other embodiments, R^2 is H and R^1 is:




wherein R^{N4} is selected from H and methyl. In these embodiments, it may be preferred that R^{N4} is H.

[0046] In some embodiments, R^1 is H and R^2 is:




where R^{N5} is selected from H and C_{1-3} alkyl (i.e. methyl, ethyl, prop-1-yl and prop-2-yl). In these embodiments, it may be preferred that R^{N5} is selected from H and methyl.

[0047] It may be preferred that R^1 is selected from :





[0048] In some embodiments, R^1 is H and R^2 is:



where R^{N6} is selected from H and C_{1-3} alkyl (i.e. methyl, ethyl, prop-1-yl and prop-2-yl). In these embodiments, it may be preferred that R^{N6} is selected from H and methyl.

[0049] It may be further preferred that R^1 is H and R^2 is:






[0050] In some embodiments, R^3 is selected from F, Me, Et, OMe and OCF_3 . In some of these embodiments, R^3 is OMe.

[0051] In some embodiments, R^4 is selected from CF_3 , Cl, Br, CF_2H , and CN.

[0052] In further embodiments, R^4 is selected from CF_3 , Cl and CF_2H . In further embodiments, R^4 is selected from CF_3 and Cl. It may be preferred that R^4 is CF_3 .

[0053] In some embodiments, it may be preferred that R^5 is a group of the following formulae:



[0054] In some embodiments, R^7 is H and R^6 is $(CHR^{C1})_{n1}C(O)N(R^{N7})Z^1$.

[0055] In further embodiments, R^7 is H and R^6 is selected from $CH_2C(O)NH_2$, $CH_2C(O)NHCH_3$, $CHCH_3C(O)NH_2$ and $CHCH_3C(O)NHCH_3$.

[0056] It may be preferred that R^7 is H and R^6 is selected from $CH_2C(O)NH_2$, $CHCH_3C(O)NH_2$ and $CH_2C(O)NHCH_3$, and more preferably from $CH_2C(O)NH_2$, and $CHCH_3C(O)NH_2$. Most preferably R^6 is $CH_2C(O)NH_2$.

[0057] In some embodiments, R^6 is H and R^7 is $(CH_2)_{m1}C(O)N(R^{M1})Y^1$.


[0058] In further embodiments, R^6 is H and R^7 is selected from $C(O)NH_2$, $C(O)NHCH_3$, $CH_2C(O)NH_2$ and $CH_2C(O)NHCH_3$.

[0059] It may be preferred that R^6 is H and R^7 is $C(O)NH_2$.


[0060] In some embodiments where R^6 is H and R^7 is $C(O)NH_2$, R^8 is methyl.

[0061] In some embodiments, it may be preferred that R^5 is a group of the following formula:






[0062] In selected embodiments of the invention, the compounds may of formula Ia:




wherein R^{1a} is selected from:




and R^{3a} is selected from Me, Et, OMe, OCF₃, and F.

[0063] In selected embodiments of the invention, the compounds may of formula Ib:



wherein R^{3b} is selected from OMe and OCF₃, and

[0064] R^{5b} is selected from:




[0065] Embodiments of the inventions are compounds of the examples, including compounds 1 to 13. Embodiments of particular interest include compounds 3, 6 and 11.

General synthesis methods

[0066] The compounds of the invention can be prepared employing the following general methods and using procedures described in detail in the experimental section. The reaction conditions referred to are illustrative and non-limiting.

[0067] Compounds of formula I, as described above, can be prepared by synthetic strategies outlined below, wherein the definitions above apply:

Scheme A




[0068] Compounds of formula F1 may be reacted with substituted synthetic anilines of formula F2 (as prepared in scheme C, D, E, F, G and H) to form intermediates of formula F3 where L¹ and L² may be the same or different and include Cl, Br, I, SMe, SO₂Me and R⁴ = CF₃, halogen, CF₂H or CN.

[0069] Compounds of the formula F1 may be prepared where L¹ and L² are different (see scheme B) to allow regioselective substitution or when L¹=L² suitable reaction conditions can be employed (choice of solvent, reaction temperature, addition of a Lewis acid, for example ZnCl₂ in diethyl ether) to allow L¹ to be selectively displaced over L². Where regiochemical mixtures and di-substitution are obtained the regioisomers may be separated by chromatography.

[0070] Compounds of the formula F1 where L¹=L² are either commercially available, for example 2,4-dichloro-5-(trifluoromethyl)pyrimidine, 2,4-dichloro-5-fluoropyrimidine, 2,4,5-trichloropyrimidine, 2,4-dichloro-5-bromopyrimidine, 2,4-dichloro-5-iodopyrimidine, 2,4-dichloro-5-cyanopyrimidine or may be prepared readily from commercial starting materials. Where R⁴ = CF₃ and differentiation of L¹ and L² is desirable, the method outlined in scheme B may be employed.


Scheme B



[0071] Commercially available 2,4-dichloro-5-(trifluoromethyl)pyrimidine (G1) can be selectively reacted with sodium thiomethoxide in the presence of zinc(II) chloride to give 2-thiomethyl-4-chloro-5-(trifluoromethyl)pyrimidine (G2). 2-Thiomethyl-4-chloro-5-(trifluoromethyl)pyrimidine (G2) can be further reacted, for example by conversion to 2-thiomethyl-4-iodo-5-(trifluoromethyl)pyrimidine (G3) under Finkelstein conditions and/or by oxidation with mCPBA to give the corresponding sulfone if further differentiation of the 2 and 4-


position is required or if additional activation is desirable.

Scheme C




[0072] Commercially available *tert*-butyl piperazine-1-carboxylate (G5) and 4-fluoro-2-substituted-1-nitrobenzenes of the formula F4 can be reacted in an S_NAr reaction to give *tert*-butyl 4-(3-substituted-4-nitrophenyl)piperazine-1-carboxylates of the formula F5. Subsequent reduction *via* hydrogenation in the presence of a catalyst, for example palladium on charcoal or platinum oxide when R^3 is Cl, Br or I, gives the corresponding anilines of the formula F6.


Scheme D



[0073] The corresponding 4-piperidine analogues of F6 can be prepared by a sequence of reactions starting with the conversion of commercially available *tert*-butyl 4-oxopiperidine-1-carboxylate (G6) to vinyl triflate G7. Coupling of G7 in a Suzuki type reaction with aryl boronic acids, or analogous boronic esters, of the formula F7 gives tetrahydropyridines of the formula F8. Subsequent reduction *via* hydrogenation in the presence of a catalyst, for example palladium on charcoal, gives anilino-piperidines of the formula F9. Alternatively, protected aniline analogues of compounds of the formula F7, for example Boc or CBz protected anilines, can be employed in the Suzuki coupling step. This negates the need for a later reduction step and may be beneficial when R^3 is halo.


Scheme E






[0074] 3-Piperidine analogues can be prepared by reaction of commercially available compounds of the formula F10, where $L^3=I$ or Br, with pyridin-3-ylboronic acid (G8) in a Suzuki type reaction to form intermediates of the formula F11. Reduction of compounds of the formula F11 with hydrogen in the presence of a catalyst, for example platinum oxide, gives intermediates of the formula F12 which may be protected using Boc anhydride to give compounds of the formula F13.

Scheme F



2-Piperidine analogues can be prepared by reaction of commercially available compounds of the formula F14, where $L^3=I$ or Br, with pyridin-2-ylboronic acid (G9) in a Suzuki type reaction to form intermediates of the formula F15. Reduction of compounds of the formula F15 with hydrogen in the presence of a catalyst, for example platinum oxide, gives intermediates of the formula F16 which may be protected using Boc anhydride to give compounds of the formula F17.


Scheme G



[0075] Compounds of the formula F20 can be prepared by coupling of commercially available *tert*-butyl piperazine-1-carboxylate (G5) and compounds of the formula F18, where $L^3=I$ or Br, in a Buchwald type reaction to give intermediates of the formula F19. Compounds of the formula F19 can be reduced with hydrogen in the presence of a catalyst, for example


palladium on charcoal or platinum oxide, to give anilines of the formula F20.

Scheme H




[0076] Compounds of the formula F21 may be reacted to form esters of the formula F22 where X=Br or I, R^8 =H or Me and Y is selected from a single bond, $-\text{CH}_2-$ and $-\text{CHCH}_3-$. When R^{01} = *t*-Bu, Boc anhydride may be employed or where R^{01} =Me methanol in the presence of an acid, for example sulfuric acid, may be used to form the desired ester. Esters of the formula F22 can be reacted with terminal acetylenes of the formula F23 in a Sonogashira type coupling to give acetylenes of the formula F24 where R^9 =TMS, TES or $(\text{CH}_3)_2\text{C}^*\text{OH}$. R^9 may then be removed to generate compounds of the formula F25. When R^9 =TMS or TES potassium carbonate or tetra-*n*-butyl ammonium fluoride may be employed to induce this transformation. When R^9 = $(\text{CH}_3)_2\text{C}^*\text{OH}$, sodium hydride in refluxed toluene may be used.

Scheme I




[0077] Pyrimidines of the formula F3 may be reacted with terminal acetylenes of the formula F25 to give acetylenes of the formula F26 in a Sonagashira type coupling. The acetylene in compounds of the formula F26 may be reduced to an alkane of the formula F27 using hydrogen gas in the presence of a transition metal catalyst. The exact choice of catalyst and conditions employed is dependant on the nature of R^4 . For example, where $R^4=CF_3$, 10% Pd/C may be used, where $R^4=Cl$, platinum oxide is employed. Esters of the formula F27 may then be deprotected to give carboxylic acids of the formula F28. Where $R^{O1}=Me$ or Et, lithium hydroxide solutions may be employed. Where $R^{O1}=t-Bu$, acidic solutions, for example trifluoroacetic acid in dichloromethane may be used. It will be appreciated that under acidic conditions Boc protecting groups in R^1 and R^2 will also be cleaved.

Scheme J




[0078] Carboxylic acids of the formula F28 can be converted to amides of the formula F29 using a suitable amine or ammonia salt in the presence of a peptide coupling agent, for example HATU.

Scheme K



[0079] Alternatively, when $R^{O1}=Me$, esters of the formula F27 may be directly converted to amides of the formula F29 by reaction with an amine at elevated temperatures.


Scheme L



Where molecules with lactams fused to the right hand side aromatic ring are required compounds of the formula F30 can be reacted with terminal acetylenes of the formula F23 in a Sonagashira type coupling to give acetylenes of the formula F31 where $R^9 = \text{TMS}$, TES or $(\text{CH}_3)_2\text{C}^*\text{OH}$. R^9 may then be removed to generate compounds of the formula F32. When $R^9 = \text{TMS}$ or TES, potassium carbonate or tetra-*n*-butyl ammonium fluoride may be employed to induce this transformation. When $R^9 = (\text{CH}_3)_2\text{C}^*\text{OH}$, sodium hydride in refluxed toluene may be used.


[0080] Compounds of the formula F32 can then be coupled to compounds of the formula F3 and further elaborated as described above and below.

Scheme M



[0081] Compounds of the formula F29, or analogues containing lactams, with Boc protecting groups present in R^1 or R^2 (in the place of R^{N1} , R^{N2} , R^{N3} , R^{N4} and R^{N5}) may then be deprotected under acidic conditions, for example using trifluoroacetic acid in dichloromethane solutions, to give the corresponding parent piperazine or piperidine compounds of the formula F33.


Scheme N



[0082] Compounds of the formula F33 may then be further modified by derivitisation of the amine functionality. For example, compounds of the formula F34 where R^{N1} , R^{N2} , R^{N3} , R^{N4} , R^{N5} or $R^{N6} = \text{Me}$ may be prepared by reductive alkylation with formaldehyde in the presence of sodium triacetoxyborohydride. Derivatives where R^{N1} , R^{N2} , R^{N3} , R^{N5} or $R^{N6} = \text{Et}$ may be prepared by reductive alkylation with acetaldehyde in the presence of sodium triacetoxyborohydride. Compounds of the formula F34 where R^{N1} , R^{N2} , R^{N3} , R^{N5} or $R^{N6} = \text{acetyl}$ may be prepared by reaction of compounds of the formula F33 with a suitable acylating agent, for example acetic anhydride.

[0083] An alternate strategy for the formation of compounds of the formula F27, where $R^3=CF_3$ and $R^7=H$, is to prepare compounds of the formula F37, as outlined in scheme N.

Scheme N



[0084] Coupling of esters of the formula F25, where $R^8=H$, with 4-iodo-2-(methylthio)-5-(trifluoromethyl)pyrimidine (G3) under Sonagashira conditions gives acetylenes of the formula F35. Oxidation, using MCPBA, gives sulfones of the formula F36. Reduction of the acetylene using hydrogen, in the presence of a catalyst, for example 10% palladium on charcoal, gives compounds of the formula F37.

[0085] Compounds of the formula F37 can be reacted with anilines of the formula F2 under acidic conditions, for example in the presence of trifluoro acetic acid to give compounds of the formula F27 which can then be further elaborated as described above.

Use of Compounds of the Invention

[0086] The present invention provides active compounds, specifically, active 2,4,5-substituted pyrimidines.

[0087] The term "active", as used herein, pertains to compounds which are capable of inhibiting FAK activity, and specifically includes both compounds with intrinsic activity (drugs) as well as prodrugs of such compounds, which prodrugs may themselves exhibit little or no intrinsic activity.

[0088] Assays which may be used in order to assess the FAK inhibition offered by a particular compound are described in the examples below.

[0089] Active compounds may also be used as part of an *in vitro* assay, for example, in order

to determine whether a candidate host is likely to benefit from treatment with the compound in question.

[0090] Also described herein are active compounds for use in a method of treatment of the human or animal body. Such a method may comprise administering to such a subject a therapeutically-effective amount of an active compound, preferably in the form of a pharmaceutical composition.

[0091] The term "treatment", as used herein in the context of treating a condition, pertains generally to treatment and therapy, whether of a human or an animal (e.g. in veterinary applications), in which some desired therapeutic effect is achieved, for example, the inhibition of the progress of the condition, and includes a reduction in the rate of progress, a halt in the rate of progress, amelioration of the condition, and cure of the condition. Treatment as a prophylactic measure (i.e. prophylaxis) is also included.

[0092] The term "therapeutically-effective amount" as used herein, pertains to that amount of an active compound, or a material, composition or dosage from comprising an active compound, which is effective for producing some desired therapeutic effect, commensurate with a reasonable benefit/risk ratio.

Cancer

[0093] The present invention provides active compounds which are anticancer agents. One of ordinary skill in the art is readily able to determine whether or not a candidate compound treats a cancerous condition for any particular cell type, either alone or in combination.

[0094] Examples of cancers include, but are not limited to, bone cancer, brain stem glioma, breast Cancer, cancer of the adrenal gland, cancer of the anal region, cancer of the bladder, cancer of the endocrine system, cancer of the oesophagus, cancer of the head or neck, cancer of the kidney or ureter, cancer of the liver, cancer of the parathyroid gland, cancer of the penis, cancer of the small intestine, cancer of the thyroid gland, cancer of the urethra, carcinoma of the cervix, carcinoma of the endometrium, carcinoma of the fallopian tubes, carcinoma of the renal pelvis, carcinoma of the vagina, carcinoma of the vulva, chronic or acute leukemia, colon cancer, cutaneous or intraocular melanoma, haemetological malignancies, Hodgkin's disease, lung cancer, lymphocytic lymphomas, neoplasms of the central nervous system (CNS), ovarian cancer, pancreatic cancer, pituitary adenoma, primary CNS lymphoma, prostate cancer, rectal cancer, renal cell carcinoma, sarcoma of soft tissue, skin cancer, spinal axis tumors, stomach cancer and uterine cancer.

[0095] Any type of cell may be treated, including but not limited to, lung, gastrointestinal (including, e.g., bowel, colon), breast (mammary), ovarian, prostate, liver (hepatic), kidney (renal), bladder, pancreas, brain, and skin.

[0096] The anti cancer treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents:-

1. (i) other antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cisplatin, oxaliplatin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan, temozolamide and nitrosoureas); antimetabolites (for example gemcitabine and antifolates such as fluoropyrimidines like 5 fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, and hydroxyurea); antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and docetaxel (Taxotere) and polokinese inhibitors); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);
2. (ii) cytostatic agents such as antioestrogens (for example tamoxifen, fulvestrant, toremifene, raloxifene, droloxifene and idoxifene), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;
3. (iii) anti-invasion agents (for example c-Src kinase family inhibitors like 4-(6-chloro-2,3-methylenedioxyanilino)-7-[2-(4-methylpiperazin-1-yl)ethoxy]-5-tetrahydropyran-4-yloxyquinazoline (AZD0530; International Patent Application WO 01/94341), N-(2-chloro-6-methylphenyl)-2-{6-[4-(2-hydroxyethyl)piperazin-1-yl]-2-methylpyrimidin-4-ylamino}thiazole-5-carboxamide (dasatinib, BMS-354825; J. Med. Chem., 2004, 47, 6658-6661 and 4-((2,4-dichloro-5-methoxyphenyl)amino)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinoline-3-carbonitrile (bosutinib, SKI-606; Cancer research (2003), 63(2), 375-81), and metalloproteinase inhibitors like marimastat, inhibitors of urokinase plasminogen activator receptor function or antibodies to Heparanase);
4. (iv) inhibitors of growth factor function: for example such inhibitors include growth factor antibodies and growth factor receptor antibodies (for example the anti erbB2 antibody trastuzumab [Herceptin^T], the anti-EGFR antibody panitumumab, the anti erbB1 antibody cetuximab [Erbix^T, C225] and any growth factor or growth factor receptor antibodies disclosed by Stern et al. Critical reviews in oncology/haematology, 2005, Vol. 54, pp11-29); such inhibitors also include tyrosine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, ZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI 774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)-quinazolin-4-amine (CI 1033), erbB2

tyrosine kinase inhibitors such as lapatinib, inhibitors of the hepatocyte growth factor family, inhibitors of the platelet-derived growth factor family such as imatinib, inhibitors of serine/threonine kinases (for example Ras/Raf signalling inhibitors such as farnesyl transferase inhibitors, for example sorafenib (BAY 43-9006)), inhibitors of cell signalling through MEK and/or AKT kinases, inhibitors of the hepatocyte growth factor family, c-kit inhibitors, abl kinase inhibitors, IGF receptor (insulin-like growth factor) kinase inhibitors; aurora kinase inhibitors (for example AZD1152, PH739358, VX-680, MLN8054, R763, MP235, MP529, VX-528 AND AX39459) and cyclin dependent kinase inhibitors such as CDK2 and/or CDK4 inhibitors;

5. (v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, [for example the anti vascular endothelial cell growth factor antibody bevacizumab (AvastinT) and VEGF receptor tyrosine kinase inhibitors such as 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (ZD6474; Example 2 within WO 01/32651), 4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (AZD2171; Example 240 within WO 00/47212), vatalanib (PTK787; WO 98/35985) and SU11248 (sunitinib; WO 01/60814), compounds such as those disclosed in International Patent Applications WO97/22596, WO 97/30035, WO 97/32856 and WO 98/13354 and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha_v\beta_3$ function and angiostatin)];
6. (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 and WO 02/08213;
7. (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
8. (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene directed enzyme pro drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi drug resistance gene therapy; and
9. (ix) immunotherapy approaches, including for example ex vivo and in vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte macrophage colony stimulating factor, approaches to decrease T cell anergy, approaches using transfected immune cells such as cytokine transfected dendritic cells, approaches using cytokine transfected tumour cell lines and approaches using anti idiotypic antibodies

[0097] A combination of particular interest is with docetaxel. Other possible combinations of interest include with gemcitabine, cisplatin and the camptothecin prodrug irinotecan.

Administration

[0098] The active compound or pharmaceutical composition comprising the active compound may be administered to a subject by any convenient route of administration, whether systemically/ peripherally or at the site of desired action, including but not limited to, oral (e.g. by ingestion); topical (including e.g. transdermal, intranasal, ocular, buccal, and sublingual); pulmonary (e.g. by inhalation or insufflation therapy using, e.g. an aerosol, e.g. through mouth or nose); rectal; vaginal; parenteral, for example, by injection, including subcutaneous, intradermal, intramuscular, intravenous, intraarterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal; by implant of a depot, for example, subcutaneously or intramuscularly. The subject may be a eukaryote, an animal, a vertebrate animal, a mammal, a rodent (e.g. a guinea pig, a hamster, a rat, a mouse), murine (e.g. a mouse), canine (e.g. a dog), feline (e.g. a cat), equine (e.g. a horse), a primate, simian (e.g. a monkey or ape), a monkey (e.g. marmoset, baboon), an ape (e.g. gorilla, chimpanzee, orang-utan, gibbon), or a human.

Formulations

[0099] While it is possible for the active compound to be administered alone, it is preferable to present it as a pharmaceutical composition (e.g. formulation) comprising at least one active compound, as defined above, together with one or more pharmaceutically acceptable carriers, adjuvants, excipients, diluents, fillers, buffers, stabilisers, preservatives, lubricants, or other materials well known to those skilled in the art and optionally other therapeutic or prophylactic agents.

[0100] Thus, the present invention further provides pharmaceutical compositions, as defined above, and methods of making a pharmaceutical composition comprising admixing at least one active compound, as defined above, together with one or more pharmaceutically acceptable carriers, excipients, buffers, adjuvants, stabilisers, or other materials, as described herein.

[0101] The term "pharmaceutically acceptable" as used herein pertains to compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of a subject (e.g. human) without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. Each carrier, excipient, etc. must also be "acceptable" in the sense of being compatible with the other ingredients of the formulation.

[0102] Suitable carriers, excipients, etc. can be found in standard pharmaceutical texts, for example, Remington's Pharmaceutical Sciences, 18th edition, Mack Publishing Company, Easton, Pa., 1990.

[0103] The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods include the step of

bringing into association the active compound with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active compound with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.

[0104] Formulations may be in the form of liquids, solutions, suspensions, emulsions, elixirs, syrups, tablets, lozenges, granules, powders, capsules, cachets, pills, ampoules, suppositories, pessaries, ointments, gels, pastes, creams, sprays, mists, foams, lotions, oils, boluses, electuaries, or aerosols.

[0105] Formulations suitable for oral administration (e.g. by ingestion) may be presented as discrete units such as capsules, cachets or tablets, each containing a predetermined amount of the active compound; as a powder or granules; as a solution or suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion; as a bolus; as an electuary; or as a paste.

[0106] A tablet may be made by conventional means, e.g., compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active compound in a free-flowing form such as a powder or granules, optionally mixed with one or more binders (e.g. povidone, gelatin, acacia, sorbitol, tragacanth, hydroxypropylmethyl cellulose); fillers or diluents (e.g. lactose, microcrystalline cellulose, calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc, silica); disintegrants (e.g. sodium starch glycolate, cross-linked povidone, cross-linked sodium carboxymethyl cellulose); surface-active or dispersing or wetting agents (e.g. sodium lauryl sulfate); and preservatives (e.g. methyl p-hydroxybenzoate, propyl p-hydroxybenzoate, sorbic acid). Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active compound therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach

[0107] Formulations suitable for topical administration (e.g. transdermal, intranasal, ocular, buccal, and sublingual) may be formulated as an ointment, cream, suspension, lotion, powder, solution, past, gel, spray, aerosol, or oil. Alternatively, a formulation may comprise a patch or a dressing such as a bandage or adhesive plaster impregnated with active compounds and optionally one or more excipients or diluents.

[0108] Formulations suitable for topical administration in the mouth include lozenges comprising the active compound in a flavoured basis, usually sucrose and acacia or tragacanth; pastilles comprising the active compound in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active compound in a suitable liquid carrier.

[0109] Formulations suitable for topical administration to the eye also include eye drops wherein the active compound is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active compound.

[0110] Formulations suitable for nasal administration, wherein the carrier is a solid, include a coarse powder having a particle size, for example, in the range of about 20 to about 500 microns which is administered in the manner in which snuff is taken, i.e. by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid for administration as, for example, nasal spray, nasal drops, or by aerosol administration by nebuliser, include aqueous or oily solutions of the active compound.

[0111] Formulations suitable for administration by inhalation include those presented as an aerosol spray from a pressurised pack, with the use of a suitable propellant, such as dichlorodifluoromethane, trichlorofluoromethane, dichloro-tetrafluoroethane, carbon dioxide, or other suitable gases.

[0112] Formulations suitable for topical administration via the skin include ointments, creams, and emulsions. When formulated in an ointment, the active compound may optionally be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active compounds may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example, at least about 30% w/w of a polyhydric alcohol, i.e., an alcohol having two or more hydroxyl groups such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or penetration of the active compound through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogues.

[0113] When formulated as a topical emulsion, the oily phase may optionally comprise merely an emulsifier (otherwise known as an emulgent), or it may comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabiliser. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabiliser(s) make up the so-called emulsifying wax, and the wax together with the oil and/or fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

[0114] Suitable emulgents and emulsion stabilisers include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate and sodium lauryl sulphate. The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations may be very low. Thus the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or

other containers. Straight or branched chain, mono- or dibasic alkyl esters such as diisoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required.

[0115] Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

[0116] Formulations suitable for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or a salicylate.

[0117] Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active compound, such carriers as are known in the art to be appropriate. Formulations suitable for parenteral administration (e.g. by injection, including cutaneous, subcutaneous, intramuscular, intravenous and intradermal), include aqueous and non-aqueous isotonic, pyrogen-free, sterile injection solutions which may contain anti-oxidants, buffers, preservatives, stabilisers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents, and liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. Examples of suitable isotonic vehicles for use in such formulations include Sodium Chloride Injection, Ringer's Solution, or Lactated Ringer's Injection. Typically, the concentration of the active compound in the solution is from about 1 ng/ml to about 10 µg/ml, for example from about 10 ng/ml to about 1 µg/ml. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules, and tablets. Formulations may be in the form of liposomes or other microparticulate systems which are designed to target the active compound to blood components or one or more organs.

Dosage

[0118] It will be appreciated that appropriate dosages of the active compounds, and compositions comprising the active compounds, can vary from patient to patient. Determining the optimal dosage will generally involve the balancing of the level of therapeutic benefit against any risk or deleterious side effects of the treatments of the present invention. The selected dosage level will depend on a variety of factors including, but not limited to, the activity of the particular compound, the route of administration, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds, and/or materials used in combination, and the age, sex, weight, condition, general health, and prior

medical history of the patient. The amount of compound and route of administration will ultimately be at the discretion of the physician, although generally the dosage will be to achieve local concentrations at the site of action which achieve the desired effect without causing substantial harmful or deleterious side-effects.

[0119] Administration *in vivo* can be effected in one dose, continuously or intermittently (e.g. in divided doses at appropriate intervals) throughout the course of treatment. Methods of determining the most effective means and dosage of administration are well known to those of skill in the art and will vary with the formulation used for therapy, the purpose of the therapy, the target cell being treated, and the subject being treated. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician.

[0120] In general, a suitable dose of the active compound is in the range of about 100 µg to about 250 mg per kilogram body weight of the subject per day. Where the active compound is a salt, an ester, prodrug, or the like, the amount administered is calculated on the basis of the parent compound and so the actual weight to be used is increased proportionately.

EXAMPLES

[0121] The following examples are provided solely to illustrate the present invention and are not intended to limit the scope of the invention, as described herein.

Acronyms

[0122] For convenience, many chemical moieties are represented using well known abbreviations, including but not limited to, methyl (Me), ethyl (Et), n-propyl (nPr), iso-propyl (iPr), n-butyl (nBu), tert-butyl (tBu), n-hexyl (nHex), cyclohexyl (cHex), phenyl (Ph), biphenyl (biPh), benzyl (Bn), naphthyl (naph), methoxy (MeO), ethoxy (EtO), benzoyl (Bz), and acetyl (Ac).

[0123] For convenience, many chemical compounds are represented using well known abbreviations, including but not limited to, methanol (MeOH), ethanol (EtOH), iso-propanol (i-PrOH), methyl ethyl ketone (MEK), ether or diethyl ether (Et₂O), acetic acid (AcOH), dichloromethane (methylene chloride, DCM), trifluoroacetic acid (TFA), dimethylformamide (DMF), tetrahydrofuran (THF), dimethylsulfoxide (DMSO), meta-chloroperoxybenzoic acid (mCPBA), tert-butyloxycarbonyl (Boc), trimethylsilyl (TMS), triethylsilyl (TES), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU), diphenylphosphoryl azide (DPPA), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDCI), 4-dimethylaminopyridine (DMAP), tetra-n-butylammonium fluoride (TBAF), *N,N*-Diisopropylethylamine (DIPEA), 1-hydroxybenzotriazole (HOBt), and 1,2-dichloroethene (DCE).

General Experimental Details

[0124] Unless otherwise stated the following generalisations apply.

[0125] ^1H NMR spectra were recorded on either a Bruker Avance DRX300 (300 MHz), a Bruker Ultrashield plus (400 MHz) or a Varian Unity Inova 600 (600 MHz) spectrometer. The multiplicity of a signal is designated by the following abbreviations: s, singlet; d, doublet; t, triplet; q, quartet; br, broad; m, multiplet. All observed coupling constants, J , are reported in Hertz. ^{13}C NMR were recorded on a Bruker Avance DRX300 (75 MHz), a Bruker Ultrashield plus (100 MHz) or a Varian Unity Inova 600 (150 MHz) spectrometer in a broad band decoupled mode.

[0126] LC/MS data was generated using either a Finnigan LCQ Advantage Max (LCMS-A), a Waters ZQ 3100 system (LCMS-B) or an Agilent 6100 Series Single Quad LC/MS (LCMS-C).

LCMS Method A (LCMS-A)

[0127] Instrument: Finnigan LCQ Advantage Max

Pump: Finnigan Surveyor LC Pump

Finnigan Surveyor Autosampler

Finnigan Surveyor PDA Detector

LC conditions:

[0128] Reverse Phase HPLC analysis

Column: Gemini 3 μ C18 20x4.0mm 110A

Injection Volume 10 μ L

Solvent A: Water 0.1% Formic Acid

Solvent B: Acetonitrile 0.1% Formic Acid

Gradient: 10-100% B over 10min

Detection: 100-600nm

MS conditions:

[0129] Ion Source: Ion trap

Ion Mode: ES positive

Temp: 300°C

Capillary V- 25

Detection: Ion counting
Scan Range: 80-1000A mu
Scan Time: 0.2 sec
Acquisition time: 10min

LCMS Method B (LCMS-B)

[0130] Instrument: Waters ZQ 3100 -Mass Detector
Waters 2545-Pump
Waters SFO System Fluidics Organizer
Waters 2996 Diode Array Detector
Waters 2767 Sample Manager

LC conditions:

[0131] Reverse Phase HPLC analysis
Column: XBridge TM C18 5µm 4.6x100mm
Injection Volume 10µL
Solvent A: Water 0.1% Formic Acid
Solvent B: Acetonitrile 0.1% Formic Acid
Gradient: 10-100% B over 10min
Flow rate: 1.5 mL/min
Detection: 100-600nm

MS conditions:

[0132] Ion Source: Single-quadrupole
Ion Mode: ES positive
Source Temp: 150°C
Desolvation Temp: 350°C
Detection: Ion counting
Cpillary (KV)-3.00
Cone (V): 30
Extractor (V):3
RF Lens (V): 0.1
Scan Range: 100-1000 Amu
Scan Time: 0.5 sec
Acquisition time: 10min
Gas Flow
Desolvation L/hr-650

LCMS Method C (LCMS-C)

[0133] Instrument: Agilent 6100 Series Single Quad LC/MS
Agilent 1200 Series HPLC
Pump: 1200 Series G1311A Quaternary pump
Autosampler: 1200 Series G1329A Thermostatted Autosampler
Detector: 1200 Series G1314B Variable Wavelength Detector

LC conditions:

[0134] Reverse Phase HPLC analysis
Column: Luna C8(2) 5 μ 50X 4.6mm 100A
Column temperature: 30°C
Injection Volume: 5 μ L
Solvent A: Water 0.1% Formic Acid
Solvent B: Acetonitrile 0.1% Formic Acid
Gradient: 5-100% B over 10min
Detection: 254 nm or 214 nm

MS conditions:

[0135] Ion Source: Quadrupole
Ion Mode: Multimode-ES
Drying gas temp: 300°C
Vaporizer temperature: 200°C
Capillary voltage (V): 2000 (positive)
Capillary voltage (V): 4000 (negative)
Scan Range: 100-1000
Step size: 0.1 sec
Acquisition time: 10min


[0136] Analytical thin-layer chromatography was performed on Merck silica gel 60F254 aluminium-backed plates which were visualised using fluorescence quenching under UV light or acidic anisaldehyde or a basic potassium permanganate dip. Flash chromatography was performed using either a Teledyne Isco CombiFlash Rf purification system using standard RediSep® cartridges or a Biotage Isolera purification system using either Grace or Biotage silica cartridges.

[0137] Where necessary, anhydrous solvents were prepared using a Braun purification system

or purchased from Sigma-Aldrich.

Example 1: 2-(2-(2-((2-Methoxy-4-(piperazin-1-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (1)

[0138]



(a) *tert*-Butyl 4-(3-methoxy-4-nitrophenyl)piperazine-1-carboxylate (I1)

[0139] A mixture of 5-fluoro-2-nitroanisole (3.00 g, 17.5 mmol), *tert*-butyl piperazine-1-carboxylate (4.57 g, 24.5 mmol) and potassium carbonate (3.63 g, 26.3 mmol) in dimethyl sulfoxide (20 mL) was heated at 90 °C for 1.5 hours under a nitrogen atmosphere. The reaction mixture was cooled and diluted with water (100 mL). The resulting precipitate was filtered and dried to give a bright yellow solid. The solid was suspended in a 5:1 solution of petroleum ether/diethyl ether (50 mL), sonicated for 2 minutes, filtered and dried to give the title compound (I1) (3.95 g, 66%) as a bright yellow solid; ¹H NMR (300 MHz, *d*₆-DMSO) δ 1.42 (s, 9 H), 3.46 (brs, 8 H), 3.90 (s, 3 H), 6.52 (s, 1 H), 6.57 (d, 1 H, *J* = 9.5 Hz) and 7.89 (d, 1 H, *J* = 9.3 Hz). LCMS Method B: rt 7.45 min; *m/z* 338.2 [M+H]⁺.

(b) *tert*-Butyl 4-(4-amino-3-methoxyphenyl)piperazine-1-carboxylate (I2)

[0140] To a solution of *tert*-butyl 4-(3-methoxy-4-nitrophenyl)piperazine-1-carboxylate (*11*) (1.90 g, 5.63 mmol) in methanol (20 mL) and ethyl acetate (80 mL) was added 10% Pd/C (0.500 g). The mixture was evacuated and backfilled three times with hydrogen gas then stirred under a hydrogen atmosphere for 18 hours. The crude reaction mixture was filtered through a plug of Celite, washing with ethyl acetate and the filtrate evaporated *in vacuo* then dried under high vacuum to give a dark red viscous oil. The oil was suspended in a 2:1 solution of diethyl ether and petroleum ether (15 mL) and sonicated for 3 minutes. The resulting mixture was evaporated *in vacuo* and dried under high vacuum to give the title compound (*12*) (1.55 g, 89%) as a dark red solid; ¹H NMR (300 MHz, *d*₆-DMSO) δ 1.47 (s, 9 H), 2.97 (t, 4 H, *J* = 4.7 Hz), 3.57 (t, 4 H, *J* = 4.7 Hz), 3.83 (s, 3 H), 6.40 (dd, 1 H, *J* = 8.3, 2.1 Hz), 6.50 (d, 1 H, *J* = 2.0 Hz) and 6.64 (d, 1 H, *J* = 8.3 Hz). LCMS Method B: rt 5.02 min; *m/z* 308.3 [M+H]⁺.

(c) *tert*-Butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperazine-1-carboxylate (*13*)

[0141] To a stirred chilled solution (0 °C) of 2,4-dichloro-5-(trifluoromethyl)pyrimidine (0.800 g, 3.68 mmol) under a stream of dry N₂ gas in 1:1 dichloroethane/*tert*-butanol (50 mL) was added a solution of zinc chloride (1 M in diethyl ether; 4.42 mL, 4.42 mmol). After 1 hour of stirring at 0 °C, a solution of *tert*-butyl 4-(4-amino-3-methoxyphenyl)piperazine-1-carboxylate (*12*) in 1:1 dichloroethane/*tert*-butanol (15 mL) (1.13 g, 3.68 mmol, 1 eq.) was added drop wise over 30 minutes followed by the drop wise addition of a solution of triethylamine (0.565 mL, 0.410 mmol, 1.1 eq.) in 1:1 dichloroethane/*tert*-butanol (5 mL) over 30 minutes. The resulting mixture was then stirred vigorously at room temperature for 18 hours. The volatiles were evaporated under reduced pressure and the residue then dried under high vacuum. Ethyl acetate and water were added and the aqueous layer was extracted with ethyl acetate (3x30 mL). The combined organic extracts were washed with water, brine, dried (magnesium sulfate), filtered and evaporated to give the crude product as a dark yellow foam. The crude product was purified by flash column chromatography on silica gel (0-20% ethyl acetate/petroleum ether) to yield the title compound (*13*) (1.25 g, 69%) as a yellow solid; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (s, 9 H), 3.11 (t, 4 H, *J* = 5.2 Hz), 3.59 (t, 4 H, *J* = 5.0 Hz), 3.88 (s, 3 H), 6.54 (s, 1 H), 6.57 (d, 1 H, *J* = 9.4 Hz), 7.83 (s, 1 H), 8.17 (d, 1 H, *J* = 8.5 Hz) and 8.53 (s, 1 H). LCMS Method B: rt 9.24 min; *m/z* 488.3 [M+H]⁺.

(d) *tert*-Butyl 4-(3-methoxy-4-((4-((2-(2-methoxy-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperazine-1-carboxylate (*15*)

[0142] To a solution of *tert*-butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperazine-1-carboxylate (*172*) (0.350 g, 0.717 mmol) in dimethylformamide

(2.5 mL) and triethylamine (0.450 mL) was added methyl 2-(2-ethynylphenyl)acetate (14: prepared according to the procedure of Peng, C. et al.; Adv. Synth. Catal. 2008, 350, 2359 - 2364 or as described below) (0.137 g, 0.789 mmol), trans-dichlorobis(triphenylphosphine) palladium(II) (0.025 g, 0.036 mmol), CuI (0.014 g, 0.072 mmol) and triphenylphosphine (0.019 g, 0.072 mmol). The reaction mixture was heated under microwave irradiation at 120 °C whilst stirring for 15 minutes. The reaction mixture was diluted with ethyl acetate and water then the resulting mixture was extracted with ethyl acetate (3x15 mL). The combined organic extracts were washed with water, brine, dried (magnesium sulfate), filtered and evaporated to give a brown solid. The solid was purified by flash column chromatography on silica gel (10-50% ethyl acetate/petroleum ether) to give the title compound (15) (0.360 g, 80%) as an orange crystalline solid; ¹H NMR (300 MHz, CDCl₃) δ 1.49 (s, 9 H), 3.11 (s, 4 H), 3.60 (s, 4 H), 3.71 (d, 3 H, *J* = 1.0 Hz), 3.89 (s, 3 H), 3.96 (s, 2 H), 6.55 (s, 1 H), 6.59 (s, 1 H), 7.26-7.45 (m, 3 H), 7.68 (d, 1 H, *J* = 7.6 Hz), 7.83 (s, 1 H), 8.27 (d, 1 H, *J* = 7.3 Hz), and 8.60 (s, 1 H). LCMS Method B: rt 9.55 min; *m/z* 626.4 [M+H]⁺.

(e) *tert*-Butyl 4-(3-methoxy-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperazine-1-carboxylate (16)

[0143] To a solution of *tert*-butyl 4-(3-methoxy-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperazine-1-carboxylate (15) (0.358 g, 0.572 mmol) in ethyl acetate (25 mL) and dimethylformamide (5 mL) was added 10% Pd/C (0.200 g). The mixture was evacuated and backfilled three times with hydrogen gas then stirred under a hydrogen atmosphere for 18 hours. The reaction mixture was filtered through a plug of Celite, washing with ethyl acetate then the filtrate was evaporated *in vacuo* and dried under high vacuum to give the title compound (16) (0.360 g, 100%) as a viscous yellow oil; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (s, 9 H), 3.00-3.20 (m, 8 H), 3.59 (t, 4 H, *J* = 4.8 Hz), 3.67 (s, 3 H), 3.75 (s, 2 H), 3.90 (s, 3 H), 6.55-6.57 (m, 2 H), 7.19-7.26 (m, 4 H), 7.75 (s, 1 H), 8.26 (d, 1 H, *J* = 9.4 Hz) and 8.50 (s, 1 H). LCMS Method B: rt 6.55 min; *m/z* 630.5 [M+H]⁺.

(f) Lithium 2-(2-(2-(2-((4-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-2-methoxyphenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (17)

[0144] To a solution of *tert*-butyl 4-(3-methoxy-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperazine-1-carboxylate (16) (0.360 g, 0.572 mmol) in tetrahydrofuran (17 mL), methanol (3 mL) and water (5 mL) was added lithium hydroxide (0.042 g, 1.77 mmol) and the resulting mixture was stirred at room temperature for 20 hours. The reaction volatiles were evaporated *in vacuo* and the resulting residue was diluted with ethyl acetate and saturated aqueous sodium hydrogen carbonate. The aqueous phase was extracted with ethyl acetate (3x20 mL) and the combined organic extracts were

washed with saturated aqueous sodium hydrogen carbonate (20 mL), dried (magnesium sulfate), filtered, evaporated *in vacuo* and dried to give the title compound (*17*) (0.356 g, 100%) as a yellow foam; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (s, 9 H), 2.96-3.06 (m, 8 H) 3.56-3.62 (m, 6 H), 3.77 (s, 3 H), 6.47-6.51 (m, 2 H), 7.00-7.11 (m, 2 H), 8.00 (brs, 2 H) and 8.43 (s, 1 H). LCMS Method B: rt 9.00 min; m/z 616.4 [M+H]⁺.

(g) *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperazine-1-carboxylate (*18*)

[0145] To a stirred solution of lithium 2-(2-(2-(2-((4-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-2-methoxyphenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*17*) (0.352 g, 0.672 mmol), 1-hydroxybenzotriazole (0.093 g, 0.686 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide)hydrochloride (0.121 g, 0.629 mmol) in dry tetrahydrofuran (16 mL) and dimethylformamide (7 mL) was added *N,N*-diisopropylethylamine (0.498 mL, 2.86 mmol) under a nitrogen atmosphere. The reaction mixture was stirred for 10 minutes at room temperature then ammonium carbonate (0.263 g, 2.859 mmol) was added and the resulting suspension was stirred at room temperature for 18 hours. The reaction volatiles were evaporated *in vacuo* and the resulting residue was diluted with ethyl acetate (15 mL) and water (10 mL). The layers were separated and the aqueous layer was extracted with ethyl acetate (2x15 mL). The combined organic extracts were washed with water, brine, dried (magnesium sulfate) and concentrated *in vacuo* to give a yellow foam which was purified by flash column chromatography on silica gel (55-85% ethyl acetate/petroleum ether) to give the title compound (*18*) (0.276 g, 87%) as a yellow foam; ¹H NMR (300 MHz, CDCl₃) δ 1.48 (d, 9 H, *J* = 1.1 Hz), 3.09 (d, 8 H, *J* = 3.9 Hz), 3.58 (brs, 4 H), 3.66 (s, 2 H), 3.88 (d, 3 H, *J* = 1.0 Hz), 5.48 (brs, 1 H), 5.78 (brs, 1 H), 6.54 (d, 1 H, *J* = 1.3 Hz), 6.57 (s, 1 H), 7.23-7.25 (brs, 4 H), 7.72 (s, 1 H), 8.13 (d, 1 H, *J* = 8.6 Hz) and 8.48 (s, 1 H). LCMS Method B: rt 8.52 min; m/z 615.4 [M+H]⁺.


(f) 2-(2-(2-(2-((2-methoxy-4-(piperazin-1-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (*1*)

[0146] To a solution of *tert*-butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperazine-1-carboxylate (*18*) (0.276 g, 0.449 mmol) in dichloromethane (8 mL), was added trifluoroacetic acid (0.669 mL, 8.98 mmol) and the resulting mixture was stirred under a nitrogen atmosphere for 18 hours. The solvents were evaporated *in vacuo* to give a yellow residue which was dissolved in 3:1 chloroform/2-propanol and 1 M aqueous sodium hydroxide solution. The aqueous layer was extracted with 3:1 chloroform/2-propanol (3x15 mL) and the combined organic extracts were washed with water, brine, dried (magnesium sulfate), filtered and evaporated to give the title compound (*1*) (0.229 g, 99%) as a yellow foam; ¹H NMR (300 MHz, *d*₆-DMSO) δ 2.87-3.10 (m, 12 H), 3.46 (s,

2 H), 3.78 (s, 3 H), 4.32 (d, 1 H, $J = 4.2$ Hz), 6.48 (dd, 1 H, $J = 2.2, 8.7$ Hz), 6.61 (d, 1 H, $J = 2.1$ Hz), 6.88 (brs, 1 H), 7.15-7.30 (m, 4 H), 7.38 (brs, 1 H), 7.46 (brs, 1 H), 8.51 (s, 1 H) and 8.81 (s, 1 H). LCMS Method B: rt 5.65 min; m/z 515.4 $[M+H]^+$.

Synthesis of Intermediate I4: Methyl 2-(2-Ethynylphenyl)acetate (I4).

[0147]



(a) Methyl 2-(2-iodophenyl)acetate (I9)

[0148] 2-(2-Iodophenyl)acetic acid (5.00 g, 19.1 mmol) was placed into a reaction flask and dissolved in MeOH (150 mL). Sulfuric acid (250 μ L) was added and reaction mixture was stirred and heated at 80 °C under nitrogen for 16 hours. The resulting mixture was cooled to room temperature and the volatiles removed by evaporation under reduced pressure. The residue was taken up in ethyl acetate (100 mL), washed with 10% NaHCO_3 (100 mL), dried (MgSO_4) and evaporated under reduced pressure to give the title compound (I9) (5.20g, 99%) as a clear liquid; ^1H NMR (400 MHz, CDCl_3) δ 7.85 (dd, $J = 7.9, 1.0$ Hz, 1H), 7.35 - 7.27 (m, 2H), 6.97 (ddd, $J = 7.9, 7.0, 2.1$ Hz, 1H), 3.81 (s, 2H), 3.72 (s, 3H).

(b) Methyl 2-(2-((trimethylsilyl)ethynyl)phenyl)acetate (I10)


[0149] Methyl 2-(2-iodophenyl)acetate (I9) (4.65 g, 16.8 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (295 mg, 421 μ mol) and Cu(I)I (80.0 mg, 421 μ mol) were placed into an oven dried reaction flask under nitrogen. (Trimethylsilyl)acetylene (2.80 mL, 20.2 mmol), dry degassed THF (20 mL) and triethylamine (20 mL) were added and the reaction mixture was stirred at room temperature for 16 hours. The volatiles were removed under reduced pressure to give a black residue which was adsorbed onto silica then chromatographed on silica gel (0-5% ethyl acetate/petroleum benzene 40-60 °C) to give the title compound (I10) (4.63 g, 99%) as a light brown liquid; ^1H NMR (400 MHz, CDCl_3) δ 7.48 (dd, $J = 7.5, 0.8$ Hz, 1H), 7.32 - 7.14 (m, 3H), 3.84 (s, 2H), 3.71 (s, 3H), 0.26 (s, 9H). LCMS Method C: rt 6.64 min.

(c) Methyl 2-(2-ethynylphenyl)acetate (I4)

[0150] Methyl 2-(2-((trimethylsilyl)ethynyl)phenyl)acetate (*I10*) (4.63 g, 19.0 mmol) was dissolved in DCM (200 mL) and TBAF (1.0 M in THF) (28.5 mL, 28.5 mmol, 1.5 eq) was added at 0 °C. The resulting solution was stirred at room temperature for 1 hour before washing with 10% NaHCO₃ (100 mL). The organic layer was dried (MgSO₄) then evaporated under reduced pressure to give a dark brown/black residue. The residue was adsorbed onto silica and then chromatographed on silica gel (0-10% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I4*) (2.76 g, 83%) as a red liquid; ¹H NMR (400 MHz, CDCl₃) δ 7.52 (dd, *J* = 7.6, 1.1 Hz, 1H), 7.43 - 7.16 (m, 3H), 3.88 (d, *J* = 9.6 Hz, 2H), 3.77 - 3.52 (m, 3H), 3.28 (s, 1H).

Synthesis of Intermediate I15: Methyl 2-(2-(2-(2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (I15)

[0151]



(a) 4-Chloro-2-(methylthio)-5-(trifluoromethyl)pyrimidine (I11)

[0152] To a solution of the 2,4-dichloro-5-(trifluoromethyl)pyrimidine (2.50 g, 11.5 mmol) in THF (50 mL) in an ice bath under nitrogen was added zinc(II) chloride (1.0 M in ether, 13.8 mL, 13.8 mmol) dropwise. The mixture was stirred in the ice bath for two hours, then sodium methanethiolate (0.888 g, 12.7 mmol) was added. The mixture was stirred overnight, allowing the reaction to slowly come to room temperature. After 18 hours the reaction was quenched with 2 M HCl (15 mL), and the organics removed by evaporation under reduced pressure. The aqueous residue was diluted with brine (15 mL), and extracted with DCM (3x30 mL). The combined organic phases were dried (phase separator) and carefully evaporated to give a pale yellow oil. Chromatography (2x40g silica cartridge, 0-20% DCM/n-hexane) followed by carefully evaporation of solvent (40 °C @ 400 mmHg then room temperature @ 200 mmHg) gave the title compound (*I11*) (2.149 g, 82% yield) as a colourless oil; ¹H NMR (600 MHz, CDCl₃) δ 8.66 (s, 1H), 2.61 (s, 3H). LCMS Method C: rt: 7.95 min; *m/z* 229.1[M+H]⁺. Note: *I11* is volatile.

(b) 4-Iodo-2-(methylthio)-5-(trifluoromethyl)pyrimidine (I12)

[0153] 4-Chloro-2-(methylthio)-5-(trifluoromethyl)pyrimidine (*I11*) (5.00 g, 21.9 mmol) was placed into a reaction flask then sodium iodide (9.80 g, 65.6 mmol) and hydroiodic acid (58%) (70 mL) were added. The reaction mixture was stirred for 48 hours in darkness then diluted with water (200 mL) where upon a colourless solid precipitated. The precipitate was collected by filtration and was washed with 10% NaHCO₃ solution until neutral. The resulting solid was washed with water (100 mL) then suction dried for 2 hours to give the title compound (*I12*) (3.956 g, 57%) as a pale yellow solid; ¹H NMR (400 MHz, CDCl₃) δ 8.42 (s, 1H), 2.58 (s, 3H). LCMS Method C: rt 6.30 min; *m/z* 321.0 [M+H]⁺.

(c) Methyl 2-(2-((2-(methylthio)-5-(trifluoromethyl)pyrimidin-4-yl)ethynyl)phenyl)acetate (I13)

[0154] 4-Iodo-2-(methylthio)-5-(trifluoromethyl)pyrimidine (*I12*) (2.00 g, 6.24 mmol), PdCl₂(PPh₃)₂ (438 mg, 625 μmol), CuI (119 mg, 625 μmol) and triphenylphosphine (164 mg, 625 μmol) were placed into an oven dried microwave reaction vial under nitrogen. Methyl 2-(2-ethynylphenyl)acetate (*I4*) (1.31 g, 7.49 mmol), THF (20 mL) and triethylamine (10 mL) were added and the resulting mixture was stirred at 100 °C under microwave irradiation for 10 min. The volatiles were evaporated under reduced pressure then the residue was adsorbed onto silica from DCM. The pre-adsorbed material was chromatographed on silica gel (0-25% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I13*) (1.571 g, 69%) as an orange solid; ¹H NMR (400 MHz, CDCl₃) δ 8.71 (d, *J* = 0.8 Hz, 1H), 7.68 (dd, *J* = 7.7, 1.1 Hz, 1H), 7.50 - 7.29 (m, 3H), 3.93 (s, 2H), 3.71 (d, *J* = 3.4 Hz, 3H), 2.62 (d, *J* = 3.4 Hz, 3H).

(d) Methyl 2-(2-((2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethynyl)phenyl)acetate (I14)

[0155] Methyl 2-(2-((2-(methylthio)-5-(trifluoromethyl)pyrimidin-4-yl)ethynyl)phenyl)acetate (*I13*) (3.14 g, 8.57 mmol) was dissolved in DCM (150 mL) and the resulting solution cooled to 0 °C. *m*CPBA (70%; 4.65 g, 18.9 mmol) was added then the reaction mixture was allowed to warm to room temperature, at which, stirring was continued overnight. The crude mixture was washed with 10% NaHCO₃ (200 mL) and the layers were separated. The organics were dried (MgSO₄) then evaporated under reduced pressure to give a light yellow solid. The solid was adsorbed onto silica then chromatographed on silica gel (0-50% ethyl acetate/ petroleum benzine 40-60 °C) to give the title compound (*I14*) (2.876 g, 84%) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ 9.13 (d, *J* = 0.7 Hz, 1H), 7.73 (dd, *J* = 7.6, 0.9 Hz, 1H), 7.54 - 7.46 (m, 1H), 7.44 - 7.32 (m, 2H), 3.94 (s, 2H), 3.77 - 3.67 (m, 3H), 3.43 (s, 3H). LCMS Method C: rt


5.90 min; m/z 421.0 (M+Na), 399.1 (M+1), 367.0 (M-OMe), 339.1 (M-COOMe).

(e) **Methyl 2-(2-(2-(2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethynyl)phenyl)acetate (I15)**

[0156] Methyl 2-(2-((2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethynyl)phenyl)acetate (I14) (1.50 g, 3.76 mmol) was taken up in DMF (30 mL) then 10% Pd/C (750 mg) was added. The resulting suspension was stirred under H₂ (1 atm) for 16 hours at room temperature. The crude reaction mixture was filtered through Celite, washing with MeOH. The filtrate was evaporated under reduced pressure to give a yellow liquid which was adsorbing onto silica. The silica adsorbed material was chromatographed on silica gel (0-100% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (I15) (1.38 g, 91%) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ 9.07 (d, J = 0.7 Hz, 1H), 7.30 - 7.12 (m, 4H), 3.72 (s, 2H), 3.68 (s, 3H), 3.41 - 3.35 (m, 2H), 3.35 (s, 3H), 3.20 (dd, J = 9.6, 6.3 Hz, 2H). LCMS Method C: rt 5.92 min; m/z 425.1 (M+Na), 403.1 (M+1), 401.1 (M-1), 371.1 (M-OMe), 343.1 (M-COOMe).

Example 2: 2-(2-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (2)

[0157]



(a) *tert*-Butyl 4-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1(2H)-carboxylate (I16)

[0158] Lithium diisopropylamide (2 M in heptane/THF/ethylbenzene; 15.1 mL, 30.1 mmol) was added drop wise to a solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (3.00 g, 15.1 mmol) in THF (50 mL) at -78 °C and the mixture left to stir for 30 minutes. A solution of *N*-phenyl-bis(trifluoromethanesulfonimide) (6.46 g, 18.1 mmol) in THF (60 mL) was then added dropwise over 30 minutes to the reaction and mixture left to stir for 30 minutes at -78 °C. The resulting mixture was then allowed to warm to room temperature and was stirred for 24 hours. The solvent was partially removed (ca 80 mL) and the reaction mixture quenched with saturated NaHCO₃ solution (50 mL). DCM (50 mL) was added to the solution and the layers separated. The aqueous layer was then extracted with DCM (2 x 50 mL). The organic layers were combined and washed with 0.2 M citric acid solution (50 mL), 1 M NaOH (50 mL), brine (50 mL) and dried over Na₂SO₄. The solvent was removed under reduced pressure to give a brown oil which was purified by column chromatography on silica gel (0-10% diethyl ether in petroleum benzine 40-60 °C) to afford the title compound (*I16*) (2.48 g, 50%) as an orange oil which crystallized on cooling to -18 °C; ¹H NMR (400 MHz, CDCl₃) δ 5.76 (s, 1H), 4.05 - 4.04 (m, 2H), 3.63 (t, *J* = 5.6 Hz, 2H), 2.46 - 2.43 (m, 2H), 1.47 (s, 9H).

(b) *tert*-Butyl (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)carbamate (I17)

[0159] 2-Methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (1.00 g, 4.01 mmol), Boc anhydride (1.75 g, 8.02 mmol) and toluene (10 mL) were stirred under nitrogen at 110 °C for 20 hours. The volatiles were evaporated under reduced pressure then the residue adsorbed onto silica gel. Chromatography (SiO₂, 0-20% ethyl acetate/petroleum benzine 40-60 °C) gave the title compound (*I17*) (1.40 g, 100%) as a colourless solid; ¹H NMR (400 MHz, CDCl₃) δ 8.11 (d, *J* = 7.9 Hz, 1H), 7.43 (dd, *J* = 8.0, 0.8 Hz, 1H), 7.26 (dd, *J* = 7.8, 6.9 Hz, 1H), 3.92 (s, 3H), 1.53 (s, 9H), 1.35 (s, 12H). LCMS Method C: rt 6.85 min.

(c) *tert*-Butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (I18)

[0160] To a solution of *tert*-butyl 4-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1(2H)-carboxylate (*I16*) (474 mg, 1.43 mmol), *tert*-butyl (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)carbamate (*I17*) (500 mg, 1.43 mmol) and Pd(PPh₃)₄ (165 mg, 0.143 mmol) in DME (15 mL) was added a solution of 3.5 M aqueous NaHCO₃ (2.00 mL, 5.0

eq) and the resulting suspension was heated at 80 °C for 16 hours. On cooling to room temperature ethyl acetate (70 mL) was added and the resulting solution was washed with water (50 mL). The organic layer was dried (MgSO₄) and evaporated to dryness. The residue was chromatographed (SiO₂, 0-20% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*118*) (520 mg, 90%) as a pale yellow liquid; ¹H NMR (400 MHz, CDCl₃) δ 8.03 (d, *J* = 8.1 Hz, 1H), 7.09 (s, 1H), 6.96 (dd, *J* = 8.4, 1.3 Hz, 1H), 6.88 (d, *J* = 1.7 Hz, 1H), 5.98 (s, 1H), 4.08 (d, *J* = 1.9 Hz, 2H), 3.89 (s, 3H), 3.64 (t, *J* = 5.6 Hz, 2H), 2.52 (s, 2H), 1.54 (s, 9H), 1.51 (s, 9H). LCMS Method C: rt 6.87 min; *m/z* 349.1 [M-*t*-Bu+2]⁺.

(d) *tert*-Butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*119*)

[0161] A suspension of *tert*-butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (*118*) (500 mg, 1.23 mmol) and 10% Pd/C (50 mg) in MeOH (30 mL) was stirred under an atmosphere of hydrogen for 16 hours. The resulting mixture was filtered through celite, washing with ethyl acetate (70 mL) then the filtrate evaporated to dryness. The residue was chromatographed (SiO₂, 0-20% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*119*) (411 mg, 82%) as a pale yellow liquid; ¹H NMR (400 MHz, CDCl₃) δ 7.94 (d, *J* = 7.4 Hz, 1H), 6.99 (s, 1H), 6.76 (d, *J* = 8.3 Hz, 1H), 6.67 (s, 1H), 4.21 (s, 2H), 3.84 (s, 3H), 2.77 (t, *J* = 12.2 Hz, 2H), 2.57 (ddd, *J* = 12.1, 9.0, 3.3 Hz, 1H), 1.79 (d, *J* = 12.6 Hz, 2H), 1.68 - 1.54 (m, 2H), 1.50 (s, 9H), 1.47 (s, 9H).

(e) Methyl 2-(2-(2-(2-((2-methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*120*)

[0162] Methyl 2-(2-(2-(2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*115*) (200 mg, 0.497 mmol) and *tert*-butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*119*) (303 mg, 0.745 mmol) were dissolved in trifluoroethanol (4 mL). Trifluoroacetic acid (200 µL) was added and the resulting mixture was heated at 100 °C under microwave irradiation for 1 hour. The crude reaction mixture was adsorbed onto silica gel and separated by silica gel chromatography (0-20% MeOH/DCM) to give the title compound (*120*) (215 mg, 82%) as a light brown liquid; ¹H NMR (400 MHz, *d*₆-Acetone) δ 8.64 (s, 1H), 8.37 (dd, *J* = 8.3, 3.7 Hz, 1H), 7.29 (ddd, *J* = 13.2, 6.9, 1.7 Hz, 3H), 7.22 (ddd, *J* = 8.9, 6.1, 1.7 Hz, 1H), 7.10 (d, *J* = 1.5 Hz, 1H), 6.97 (dd, *J* = 8.3, 1.5 Hz, 1H), 3.98 (d, *J* = 7.4 Hz, 3H), 3.82 (s, 2H), 3.74 - 3.60 (m, 5H), 3.27 (t, *J* = 11.7 Hz, 2H), 3.22 - 3.06 (m, 4H), 3.00 (s, 1H), 2.34 - 2.17 (m, 2H), 2.12 (dd, *J* = 4.0, 3.5 Hz, 2H), 2.08 (dt, *J* = 6.6, 2.2 Hz, 3H). LCMS Method C: rt 5.23 min; *m/z* 529.1 [M+1]⁺.

(f) *tert*-Butyl 4-(3-methoxy-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-

(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (I21)

[0163] A solution of methyl 2-(2-(2-(2-((2-methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (I20) (300 mg, 0.567 mmol) and Boc anhydride (247 mg, 1.13 mmol) in DCM (10 mL) was stirred at room temperature under nitrogen for 20 hours. The volatiles were evaporated under reduced pressure to give a brown liquid which was adsorbed onto silica gel. Chromatography (SiO₂, 0-15% MeOH/DCM) gave the title compound (I21) (210 mg, 59% yield) as a brown liquid; ¹H NMR (400 MHz, CDCl₃) δ 8.56 (s, 1H), 8.40 (d, *J* = 8.3 Hz, 1H), 7.95 (s, 1H), 7.31 - 7.18 (m, 4H), 6.87 (dd, *J* = 8.4, 1.7 Hz, 1H), 6.79 (d, *J* = 1.7 Hz, 1H), 4.28 (d, *J* = 6.5 Hz, 2H), 3.95 (d, *J* = 4.0 Hz, 3H), 3.78 (s, 2H), 3.70 (s, 3H), 3.26 - 3.02 (m, 4H), 2.84 (t, *J* = 12.2 Hz, 2H), 2.66 (t, *J* = 3.4 Hz, 1H), 1.87 (d, *J* = 12.7 Hz, 2H), 1.66 (dd, *J* = 12.9, 3.4 Hz, 2H), 1.52 (s, 9H). LCMS Method C: rt 7.24 min; *m/z* 629.2 [M+1]⁺, 627.0 [M-1]⁻.

(g) Lithium 2-(2-(2-(2-((4-(1-(tert-butoxycarbonyl)piperidin-4-yl)-2-methoxyphenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)ecetate (I22)

[0164] A solution of *tert*-butyl 4-(3-methoxy-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (I21) (210 mg, 0.334 mmol) and LiOH.H₂O (42 mg, 1.0 mmol) in THF (10 mL), water (2 mL) and MeOH (1 mL) was stirred at room temperature for 20 hours. The volatiles were evaporated under reduced pressure to give the title compound (I22) as a light yellow solid, 300 mg, excess mass due to inorganic salts present. LCMS Method C: rt 7.31 min; *m/z* 615.1 [M+1]⁺, 613.2 [M-1]⁻.

(h) *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (I23)

[0165] A solution of lithium 2-(2-(2-(2-((4-(1-(tert-butoxycarbonyl)piperidin-4-yl)-2-methoxyphenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (I22) (210 mg, 0.338 mmol), HATU (257 mg, 0.676 mmol), ammonium chloride (362 mg, 6.76 mmol) and DIPEA (115 µL) in dry DMF (4 mL) was stirred at room temperature overnight. The volatiles were evaporated under reduced pressure and the residue diluted with ethyl acetate. The resulting solution was washed with 10% NaHCO₃ then the organic layer dried (MgSO₄). The volatiles were removed under reduced pressure and the residue chromatographed (SiO₂, 0-100% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (I23) (185 mg, 89%) as a colourless solid; ¹H NMR (400 MHz, CDCl₃) δ 8.54 (s, 1H), 8.26 (d, *J* = 8.3 Hz, 1H), 7.88 (s, 1H), 7.25 (dd, *J* = 4.4, 2.4 Hz, 2H), 6.86 (dd, *J* = 8.3, 1.7 Hz, 1H), 6.76 (d, *J* = 1.7 Hz, 1H), 5.49 (d, *J* = 39.2 Hz, 2H), 4.24 (s, 2H), 3.92 (s, 3H), 3.70 (s, 2H), 3.20 - 3.02 (m, 4H), 2.80


(m, 2H), 2.64 (s, 2H), 1.84 (d, $J = 12.8$ Hz, 2H), 1.62 (dd, $J = 12.3, 3.5$ Hz, 2H), 1.49 (s, 9H). LCMS Method C: rt 6.64 min; m/z 636.2 $[M+Na]^+$, 614.1 $[M+1]^+$, 612.2 $[M-1]^-$, 558.2 $[M-t-Bu+2]^+$.

(i) 2-(2-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (2)

[0166] Trifluoroacetic acid (1 mL) was added to a stirred solution of *tert*-butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*123*) (185 mg, 0.301 mmol) in DCM (10 mL) and the resulting solution was stirred at room temperature for 2 hours. DCM (20 mL) and 10% $NaHCO_3$ solution (10 mL) were added then the organic layer was dried ($MgSO_4$) and evaporated to dryness under reduced pressure to give the title compound (**2**) (125 mg, 81%) as a colourless solid; ^1H NMR (400 MHz, CD_2Cl_2) δ 8.49 (s, 1H), 8.31 (d, $J = 8.3$ Hz, 1H), 7.91 (s, 1H), 7.28 - 7.12 (m, 3H), 6.94 (bs, 2H), 6.85 (d, $J = 8.2$ Hz, 1H), 6.78 (dd, $J = 11.8, 3.1$ Hz, 1H), 6.35 (s, 1H), 5.94 (s, 1H), 3.88 (s, 3H), 3.65 (s, 2H), 3.42 (d, $J = 12.3$ Hz, 2H), 3.24 - 3.02 (m, 4H), 2.91 (t, $J = 11.3$ Hz, 2H), 2.75 - 2.61 (m, 1H), 2.10 - 1.83 (m, 4H). LCMS Method C: rt 4.92 min; m/z 514.1 $[M+1]^+$, 521.1 $[M-1]^-$.

Example 3: 2-(2-(2-(2-((2-Methoxy-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (3)

[0167]




[0168] Formaldehyde solution (37% aq.; 32 μ L, 0.39 mmol) was added to a stirred solution of 2-(2-(2-(2-((2-methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (**2**) (40 mg, 78 μ mol) in dry MeOH (2 mL). Sodium triacetoxyborohydride (83 mg, 0.39 mmol) was added under nitrogen and the resulting mixture was stirred at room temperature for 2 hours. The crude mixture was diluted with ethyl acetate and adsorbed onto silica gel. Chromatography (SiO_2 , 0-20% MeOH/DCM) gave the title compound (**3**) (25 mg, 61%) as a solid; ^1H NMR (400 MHz, CD_2Cl_2) δ 8.53 (s, 1H), 8.34 (d, $J = 8.3$ Hz, 1H), 7.94 (s, 1H), 7.29 - 7.15 (m, 4H), 6.92 (dd, $J = 8.3, 1.4$ Hz, 1H), 6.82 (d, $J = 1.7$ Hz, 1H), 6.12 (bs, 1H), 5.84 (bs, 1H), 3.91 (s, 3H), 3.68 (s, 2H), 3.60 (d, $J = 10.6$ Hz, 2H), 3.24 - 3.03 (m, 4H), 2.80 (s, 3H), 2.79 (m, 2H), 2.71 (m, 1H), 2.42 - 2.20 (m, 2H), 2.01 (m, 2H).

LCMS Method C: rt 4.92 min; m/z 528.1 $[M+1]^+$, 526.1 $[M-1]^-$.

Example 3A: Alternative synthesis of 2-(2-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (2) and 2-(2-(2-(2-((2-Methoxy-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (3)

[0169]



(a) 4-Bromo-2-methoxy-1-nitrobenzene (I24)

[0170] Sodium metal (3.14 g, 136 mmol) was added portion-wise to methanol (300 mL) under a nitrogen atmosphere. Once a homogeneous solution was obtained 4-bromo-2-fluoro-1-nitrobenzene (20.0 g, 91 mmol) was added and the resulting mixture stirred at 60 °C for 1 h. The volatiles were evaporated and the solid residue suspended in water (400 mL). The resulting suspension was filtered washing with water (2x50 mL). The filter cake was air dried to give the title compound (I24) (20.1 g, 95% yield) as a pale yellow solid; ^1H NMR (400 MHz, CDCl_3) δ 7.75 (d, J = 8.6 Hz, 1H), 7.24 (d, J = 1.9 Hz, 1H), 7.18 (dd, J = 8.6, 1.9 Hz, 1H), 3.97 (s, 3H). LCMS Method C: rt 5.85 min.

(b) *tert*-Butyl 4-(3-methoxy-4-nitrophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (I25)

[0171] A mixture of 4-bromo-2-methoxy-1-nitrobenzene (*I24*) (2.364 g, 10.2 mmol), *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (3.00 g, 9.70 mmol), potassium carbonate (4.02 g, 29.1 mmol) and DMF (60 mL) was degassed with three vacuum/nitrogen cycles then PdCl₂(dppf)-DCM solvate (450 mg, 6 mol%) added under nitrogen in a Schlenk tube. A second tube was prepared in the same manner (ie. a total of 6.00 g starting boronate), and both tubes were heated to 85 °C under nitrogen. After 17 hours both tubes were cooled under nitrogen and added to 5% w/v aqueous lithium chloride (600 mL). The resulting mixture was extracted with ether (300 mL) and ethyl acetate (3x300 mL). The combined organic extracts were washed with water (600 mL) and brine (600 mL), dried (sodium sulphate) then evaporated to dryness. The residue was chromatographed (120 g silica cartridge, 0-60% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I25*) (5.863 g, 91% yield) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ 7.86 (d, J = 8.5 Hz, 1H), 7.04 - 6.97 (m, 2H), 6.17 (s, 1H), 4.12 (d, J = 2.5 Hz, 2H), 3.98 (s, 3H), 3.65 (t, J = 5.6 Hz, 2H), 2.52 (s, 2H), 1.49 (s, 9H). LCMS Method C: rt 6.27 min; *m/z* 279.0 [M-*t*Bu+2H]⁺, 235.1 [M-Boc+2H]⁺.

(c) *tert*-Butyl 4-(4-amino-3-methoxyphenyl)piperidine-1-carboxylate (I26) *tert*-Butyl 4-(3-methoxy-4-nitrophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (I25)

[0172] (5.609 g, 16.78 mmol) was dissolved in 1:1 ethanol: ethyl acetate (500 mL) and Pd/C (2.50 g) was added. The mixture was stirred vigorously under hydrogen for five hours then filtered through celite. The celite was washed with ethyl acetate (500 mL) and the combined filtrates evaporated to give the title compound (*I26*) (4.93 g, 96% yield) as an off-white solid; ¹H NMR (400 MHz, CDCl₃) δ 6.69 - 6.60 (m, 3H), 4.22 (s, 2H), 3.84 (s, 3H), 2.78 (t, J = 12.4 Hz, 2H), 2.54 (tt, J = 12.1, 3.5 Hz, 1H), 1.80 (d, J = 13.1 Hz, 2H), 1.65 - 1.53 (m, 2H), 1.48 (s, 9H). LCMS Method C: rt 4.91 min; *m/z* 251.1 [M-*t*Bu+2H]⁺, 207.2 [M-Boc+2H]⁺.

(d) *tert*-Butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (I27)

[0173] A solution of 2,4-dichloro-5-(trifluoromethyl)pyrimidine (981 mg, 4.52 mmol) in 1:1 *tert*-butanol: dichloroethane (40 mL) was cooled to 0 °C under nitrogen then 1.0 M zinc(II) chloride in ether (4.52 mL, 4.52 mmol) was added dropwise. After one hour a solution of *tert*-butyl 4-(4-amino-3-methoxyphenyl)piperidine-1-carboxylate (*I26*) (1.26 g, 4.11 mmol) and triethylamine (0.860 mL, 6.17 mmol) in 1:1 *tert*-butanol: dichloroethane (50 mL) was added dropwise at -10

°C then the mixture was stirred for 16 hours allowing the temperature to rise to room temperature. The mixture was concentrated, evaporated onto silica and chromatographed (120 g silica cartridge, 0-10% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I27*) (1.277 g, 64% yield) as a white solid; ¹H NMR (400 MHz, CDCl₃) δ 8.57 (s, 1H), 8.28 (d, J = 8.3 Hz, 1H), 8.01 (s, 1H), 6.86 (dd, J = 8.3, 1.8 Hz, 1H), 6.76 (d, J = 1.8 Hz, 1H), 4.25 (br s, 2H), 3.91 (s, 3H), 2.87 - 2.73 (m, 2H), 2.64 (tt, J = 12.1, 3.6 Hz, 1H), 1.83 (d, J = 12.5 Hz, 2H), 1.69 - 1.54 (m, overlaps with water), 1.49 (s, 9H). LCMS Method C: rt 7.12 min; *m/z* 431.0 [M-tBu+2H]⁺, 387.1 [M-Boc+2H]⁺; *m/z* 485.1 [M-H]⁻.

(e) *tert*-Butyl 4-(3-methoxy-4-((4-((2-(2-methoxy-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*I28*)

[0174] A mixture of methyl 2-(2-ethynylphenyl)acetate (*I4*) (0.472 g, 2.71 mmol), *tert*-butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*I27*) (1.10 g, 2.26 mmol), triphenylphosphine (59 mg, 10 mol%), copper(I) iodide (43 mg, 10 mol%), bis(triphenylphosphine)palladium(II) chloride (79 mg, 5 mol%), DMF (12 mL) and triethylamine (1.26 mL, 9.04 mmol) was degassed with nitrogen in a microwave tube then heated under microwave irradiation at 120°C for 15 minutes. Three additional tubes were prepared and heated as described above (i.e. a total of 4.40 g of *I27*). The cooled mixtures were combined, and poured into water (600 mL). The resulting mixture was extracted with dichloromethane (3x 250 mL) then the combined organic phases were washed with water (2x 300 mL), brine (300 mL), dried over sodium sulfate and evaporated to dryness. The residue was chromatographed (120 g silica cartridge, 0-50% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I28*) (5.52 g, 98% yield) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ 8.63 (s, 1H), 8.38 (d, J = 8.3 Hz, 1H), 8.00 (s, 1H), 7.69 (dd, J = 7.7, 0.9 Hz, 1H), 7.43 (td, J = 7.6, 1.4 Hz, 1H), 7.36 (d, J = 7.9 Hz, 1H), 7.32 (dd, J = 7.5, 1.4 Hz, 1H), 6.87 (dd, J = 8.4, 1.8 Hz, 1H), 6.76 (d, J = 1.8 Hz, 1H), 4.26 (s, 2H), 3.97 (s, 2H), 3.91 (s, 3H), 3.71 (s, 3H), 2.81 (t, J = 12.6 Hz, 2H), 2.64 (tt, J = 12.2, 3.6 Hz, 1H), 1.84 (d, J = 12.6 Hz, 2H), 1.69 - 1.55 (m, overlaps with water), 1.49 (s, 9H). LCMS Method C: rt 7.05 min; *m/z* (625.1 [M+H]⁺, 569.1 [M-tBu+2H]⁺; *m/z* 623.2 [M-H]⁻.

(f) *tert*-Butyl 4-(3-methoxy-4-((4-((2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*I21*)

[0175] *tert*-Butyl 4-(3-methoxy-4-((4-((2-(2-methoxy-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*I28*) (5.52 g, 8.84 mmol) was dissolved in DMF (250 mL) then triethylamine (1 mL) and Pd/C (2.50 g) were added and the mixture stirred at 30 °C under hydrogen. After 16 hours the mixture was filtered through celite and the celite washed with DMF (250 mL). Triethylamine (1 mL) and Pd/C (2.50 g) were added to the combined filtrates and the mixture stirred at 30 °C under hydrogen. After 18

hours the resulting mixture was filtered through celite washing with ethyl acetate (300 mL). The filtrate was divided into two lots; each was poured into water (800 mL) and the resulting mixture extracted with ethyl acetate (3x 300 mL). The ethyl acetate extracts from each lot were combined, washed with water (2x 500 mL) and brine, dried (sodium sulfate) and evaporated. The two workups were combined and chromatographed (120 g silica cartridge, 0-20% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I21*) (3.620 g, 65% yield) as a white solid; ¹H NMR (400 MHz, CDCl₃) δ 8.55 (s, 1H), 8.37 (d, J = 8.3 Hz, 1H), 7.91 (s, 1H), 7.29 - 7.18 (m, overlaps with CHCl₃), 6.85 (dd, J = 8.3, 1.8 Hz, 1H), 6.77 (d, J = 1.8 Hz, 1H), 4.26 (s, 2H), 3.93 (s, 3H), 3.68 (s, 3H), 3.19 - 3.01 (m, 4H), 2.81 (t, J = 12.7 Hz, 2H), 2.64 (tt, J = 12.0, 3.5 Hz, 1H), 1.85 (d, J = 13.3 Hz, 2H), 1.70 - 1.57 (m, overlaps with water), 1.49 (s, 9H). LCMS Method C: rt 7.14 min; *m/z* 629.2 [M+H]⁺, 573.1 [M-tBu+2H]⁺; *m/z* 627.2 [M-H].

(g) *tert*-Butyl 4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*I23*)

[0176] *tert*-Butyl 4-(3-methoxy-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*I21*) (3.62 g, 5.76 mmol) was dissolved in THF (200 mL) then a solution of lithium hydroxide monohydrate (1.21 g, 28.8 mmol) in water (100 mL) was added. The resulting mixture was stirred for 18 hours at room temperature then concentrated. The residue was poured into saturated sodium bicarbonate (200 mL) and water (300 mL). The resulting mixture was extracted with ethyl acetate (3x300 mL) and the combined organic phases were washed with brine, dried over sodium sulfate and evaporated. The residue was evaporated from toluene then dissolved in THF (70 mL) and DMF (12 mL) at 30 °C under nitrogen. The mixture was stirred vigorously while HOBT (1.012 g, 7.49 mmol), EDCI.HCl (1.330 g, 7.49 mmol) and DIPEA (5.02 mL, 28.8 mmol) were added. After five minutes ammonium carbonate (2.77 g, 28.8 mmol) was added and stirring continued for 16 hours. The resulting mixture was poured into saturated sodium bicarbonate (300 mL) and extracted with ethyl acetate (3x300 mL). The combined organic phases were washed with 1:1 saturated brine: water (2x300 mL), brine (300 mL), dried over sodium sulfate and evaporated. The residue was chromatographed (120 g silica cartridge, 0-80% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I23*) (2.698 g, 76% over two steps) as a white solid; ¹H NMR (400 MHz, CDCl₃) δ 8.54 (s, 1H), 8.28 (d, J = 8.3 Hz, 1H), 7.86 (s, 1H), 7.29 - 7.22 (m, overlaps with residual CHCl₃), 6.86 (dd, J = 8.3, 1.9 Hz, 1H), 6.76 (d, J = 1.9 Hz, 1H), 5.37 (d, J = 15.9 Hz, 2H), 4.25 (s, 2H), 3.93 (s, 3H), 3.71 (s, 2H), 3.18 - 3.04 (m, 4H), 2.81 (t, J = 12.5 Hz, 2H), 2.64 (tt, J = 12.0, 3.4 Hz, 1H), 1.84 (d, J = 12.9 Hz, 2H), 1.69 - 1.55 (m, overlaps with water), 1.49 (s, 9H). LCMS Method C: rt 6.59 min; *m/z* 614.2 [M+H]⁺, 558.1 [M-tBu+2H]⁺; *m/z* 612.2 [M-H].

(h) 2-(2-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (2)


[0177] A solution of *tert*-butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*123*) (2.694 g, 4.39 mmol) was dissolved in dichloromethane (150 mL), and TFA (15 mL) was added. The resulting mixture was stirred for 16 hours then concentrated. The residue was suspended in 10% sodium hydroxide (200 mL) then extracted with ethyl acetate (5x200 mL). The combined organic phases were washed with brine (300 mL), dried over sodium sulfate and evaporated to give the title compound (**2**) (2.252 g, 100% yield) as a white solid; ^1H NMR (400 MHz, CDCl_3) δ 8.53 (s, 1H), 8.26 (d, J = 8.3 Hz, 1H), 7.85 (s, 1H), 7.29 - 7.21 (m, overlaps with CHCl_3), 6.87 (d, J = 8.3 Hz, 1H), 6.80 (s, 1H), 5.43 (d, J = 10.1 Hz, 2H), 3.92 (d, J = 1.3 Hz, 3H), 3.70 (s, 2H), 3.20 (d, J = 12.7 Hz, 2H), 3.15 - 3.05 (m, 4H), 2.75 (t, J = 12.2 Hz, 2H), 2.61 (tt, J = 12.0, 3.5 Hz, 1H), 1.89 - 1.76 (m, overlaps with water), 1.67 (qd, J = 12.7, 3.9 Hz, 2H). LCMS Method C: rt 4.92 min; m/z 514.1 $[\text{M}+\text{H}]^+$, 536.1 $[\text{M}+\text{Na}]^+$; m/z 512.2 $[\text{M}-\text{H}]^-$.

(i) **2-(2-(2-(2-((2-Methoxy-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (3)**

[0178] 2-(2-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (**2**) (2.249 g, 4.38 mmol) was dissolved in methanol (220 mL) and 37% formaldehyde solution (0.483 mL, 17.5 mmol) was added. After five minutes sodium tris(acetoxy)borohydride (4.641 g, 21.9 mmol) was added and stirring continued at room temperature for two hours. The volatiles were evaporated and the residue suspended in 5% aqueous sodium hydroxide (200 mL). The resulting mixture was extracted with ethyl acetate (5x200 mL) then the combined organic phases were washed with brine (500 mL), dried over sodium sulfate and evaporated. The residue was evaporated from ether and the solvent traces removed under high vacuum to give the title compound (**3**) (2.196 g, 95% yield) as a white solid; ^1H NMR (400 MHz, CDCl_3) δ 8.53 (s, 1H), 8.24 (d, J = 8.2 Hz, 1H), 7.85 (s, 1H), 7.28 - 7.22 (m, overlaps with CHCl_3), 6.88 (dd, J = 8.3, 1.9 Hz, 1H), 6.81 (d, J = 1.8 Hz, 1H), 5.41 (s, 2H), 3.90 (s, 3H), 3.70 (s, 2H), 3.16 - 3.06 (m, 4H), 2.99 (d, J = 12.1 Hz, 2H), 2.48 (tt, J = 10.5, 5.9 Hz, 1H), 2.34 (s, 3H), 2.07 (td, J = 11.1, 4.1 Hz, 2H), 1.89 - 1.77 (m, overlaps with water). LCMS Method C: rt 4.94 min; m/z 528.1 $[\text{M}+\text{H}]^+$, 550.1 $[\text{M}+\text{Na}]^+$; m/z 526.2 $[\text{M}-\text{H}]^-$.

Example 4: 2-(2-(2-(2-((4-(1-Ethylpiperidin-4-yl)-2-methoxyphenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (4)


[0179]



[0180] Acetaldehyde (18 mg, 0.39 mmol) was added to a stirred solution of 2-(2-(2-((2-methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (2) (40 mg, 78 μ mol) in dry MeOH (2 mL). Sodium triacetoxyborohydride (83 mg, 0.39 mmol) was added under nitrogen and the resulting mixture was stirred at room temperature for 2 hours. The crude mixture was diluted with ethyl acetate and adsorbed onto silica gel. Chromatography (SiO₂, 0-20% MeOH/DCM) gave the title compound (4) (10 mg, 24%) as a light coloured solid; ¹H NMR (400 MHz, CD₂Cl₂) δ 8.52 (s, 1H), 8.34 (d, J = 8.3 Hz, 1H), 7.92 (s, 1H), 7.28 - 7.14 (m, 4H), 6.92 (d, J = 8.3 Hz, 1H), 6.82 (d, J = 1.7 Hz, 1H), 5.91 (bs, 1H), 5.87 (bs, 1H), 3.91 (s, 3H), 3.66 (s, 2H), 3.63 (m, 2H), 3.20 - 3.01 (m, 6H), 2.73 (d, J = 11.7 Hz, 2H), 2.42 - 2.21 (m, 3H), 2.02 (d, J = 18.0 Hz, 4H), 1.37 (t, J = 7.3 Hz, 3H). LCMS Method C: rt 4.96 min; m/z 542.2 [M+1]⁺, 540.1 [M-1]⁻.

Example 5: 3-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)benzamide (5)

[0181]



(a) 3-Iodobenzamide (I29)

[0182] To a stirred solution of 3-iodobenzoic acid (2.00 g, 8.06 mmol) and DIPEA (5.62 mL, 32.3 mmol) in MeCN (100 mL) at room temperature was added HOBT (1.63 g, 12.1 mmol) and

EDCI.HCl (2.32 g, 12.1 mmol). After stirring for 10 minutes, ammonium carbonate (4.65 g, 48.4 mmol) was added and the resulting solution was stirred overnight. The volatiles were removed *in vacuo* to yield a crude solid which was suspended in water (100 mL). The resulting suspension was sonicated for 10 minutes then the solid collected by filtration. The filter cake was washed with water (20 mL) and dried to yield the title compound (*I29*) (1.65 g 83%) as a brown solid; ^1H NMR (400 MHz, CDCl_3) δ 8.13 (t, J = 1.6 Hz, 1H), 7.83 (ddd, J = 7.9, 1.7, 1.1 Hz, 1H), 7.73 (ddd, J = 7.8, 1.7, 1.1 Hz, 1H), 7.16 (t, J = 7.8 Hz, 1H), 6.18 - 5.49 (m, J = 137.3 Hz, 2H). LCMS Method C: r_t = 5.04 min; m/z = 248 $[\text{M}+1]^+$.

(b) 3-((Trimethylsilyl)ethynyl)benzamide (*I30*)

[0183] A solution of 3-iodobenzamide (*I29*) (1.65 g, 6.66 mmol), Cu(I)I (127 mg, 0.666 mmol) and triphenylphosphine (524 mg, 2.00 mmol) in THF (100 mL) and triethylamine (4.64 mL, 33.3 mmol) was sonicated for 10 minutes under a nitrogen atmosphere. $\text{PdCl}_2(\text{PPh}_3)_2$ (642 mg, 0.67 mmol) and TMS acetylene (1.88 mL, 13.3 mmol) were then added and the reaction stirred at room temperature for 3 days. The volatiles were evaporated *in vacuo* and the residue chromatographed on silica gel (0-50% ethyl acetate/petroleum benzine 40-60 °C) to give the title compound (*I30*) (1.53 g); ^1H NMR (400 MHz, CDCl_3) δ 7.78 (t, J = 1.5 Hz, 1H), 7.67 (ddd, J = 7.8, 1.8, 1.2 Hz, 1H), 7.53 - 7.48 (m, 1H), 7.29 (td, J = 7.8, 0.5 Hz, 1H), 0.15 (s, 9H).

(c) 3-Ethynylbenzamide (*I31*)

[0184] Potassium carbonate (1.84 g, 13.3 mmol) was added to a stirred solution of 3-((trimethylsilyl)ethynyl)benzamide (*I30*) (1.45 g, 6.66 mmol) in MeOH (15 mL) at room temperature. The resulting suspension was stirred for 30 minutes then water (100 mL) and EtOAc (100 mL) were added. The resulting precipitate was collected via filtration and after air drying this was suspended in a 1:1 mix of acetone and methanol (15 mL). After sonication the resulting suspension was filtered to give the title compound (*I31*) (335 mg, 35%) which was used without additional purification.

(d) 2-Methoxy-4-(1,2,3,6-tetrahydropyridin-4-yl)aniline (*I32*)

[0185] To a solution of *tert*-butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (*I18*) (6.7 g, 0.016 mol) in DCM (60 mL) was added TFA (6.0 mL, 0.081 mol) and the resulting solution was stirred at room temperature for 14 hours. An additional 5 mL of TFA was added and the reaction was stirred for a further 6 hours. The volatiles were removed *in vacuo* and the residue taken up in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ and lyophilised to give the title compound (*I32*) as the TFA salt (6.38 g, 91%) as a brown oil; ^1H NMR (400 MHz,

*d*₄-MeOH) δ 7.31 (dd, *J* = 7.3, 2.8 Hz, 1H), 7.21 (t, *J* = 2.0 Hz, 1H), 7.13 - 7.09 (m, 1H), 6.20 (tt, *J* = 3.4, 1.6 Hz, 1H), 3.83 (dd, *J* = 5.7, 2.4 Hz, 2H), 3.43 (t, *J* = 6.1 Hz, 2H), 2.77 (ddd, *J* = 8.0, 3.9, 1.9 Hz, 2H). LCMS Method C: rt 1.38 min; *m/z* 201.1 [M+H]⁺.

(e) *tert*-Butyl 4-(4-amino-3-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (I33)

[0186] To a stirred solution of 2-methoxy-4-(1,2,3,6-tetrahydropyridin-4-yl)aniline (*I*32) (6.38 g, 14.8 mmol) in MeOH (50 mL) at room temperature was added a solution of (Boc)₂O (3.27 g, 15.0 mmol) in MeOH (20 mL) drop-wise over 5 minutes and the resultant solution was stirred for 1 hour. The mixture was concentrated *in vacuo* and the residue was taken up in 2 M NaOH (50 mL) solution. The aqueous layer was extracted with CH₂Cl₂ (4×50 mL) and the combined organic extracts were washed with brine (50 mL) then dried (MgSO₄). The solvent was removed *in vacuo* to give the title compound (*I*33) (1.97 g, 43%) as a brown oil; ¹H NMR (400 MHz, CDCl₃) δ 6.85 - 6.80 (m, 1H), 6.67 (d, *J* = 8.1 Hz, 1H), 5.90 (s, 1H), 4.05 (d, *J* = 2.7 Hz, 1H), 3.87 (s, 3H), 3.62 (t, *J* = 5.7 Hz, 1H), 2.49 (s, 2H), 1.49 (s, 9H).

(f) *tert*-Butyl 4-(4-amino-3-methoxyphenyl)piperidine-1-carboxylate (I26)

[0187] To a suspension of 10% Pd/C (0.068 g) in DMF (2 mL) under nitrogen was added a solution of *tert*-butyl 4-(4-amino-3-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (*I*33) (2.0 g, 6.5 mmol) in EtOH (50 mL) and the resulting suspension was stirred under an atmosphere of hydrogen at room temperature for 24 hours. The reaction mixture was filtered through celite, washing with ethyl acetate (100 mL). The filtrate was evaporated to dryness to give the title compound (*I*26) (1.42 g, 72%) as a light brown solid; ¹H NMR (400 MHz, CDCl₃) δ 6.66 - 6.58 (m, 3H), 4.21 (s, 2H), 3.76 - 3.66 (m, 4H), 2.53 (tt, *J* = 12.1, 3.5 Hz, 1H), 1.84 - 1.75 (m, 2H), 1.55 (tdd, *J* = 11.5, 6.4, 3.1 Hz, 1H), 1.47 (s, 9H).

(g) *tert*-Butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (I34)

[0188] Zinc chloride (1.0 M in Et₂O) (8.56 mL, 8.56 mmol) was added to a solution of 2,4-dichloro-5-(trifluoromethyl)pyrimidine (1.06 mL, 7.85 mmol) in 1:1 DCE/*t*-BuOH (10 mL) at 0 °C under nitrogen. The mixture was stirred for 1 hour at 0 °C and then *tert*-butyl 4-(4-amino-3-methoxyphenyl)piperidine-1-carboxylate (*I*26) (1.97 g, 7.13 mmol) in 1:1 DCE/*t*-BuOH (30 mL) was added. A solution of triethylamine (600 μ L, 0.432 mmol, 1.1 eq) in 1:1 DCE/*t*-BuOH (10 mL) was next added drop-wise at 0 °C. The reaction mixture was vigorously stirred for a further 30 minutes at 0 °C after the final addition and then at room temperature for 24 hours. The solvent was removed *in vacuo* to afford a brown oily residue which was purified by column

chromatography on silica gel (0-45% EtOAc in petroleum benzine 40-60 °C) to give a pale yellow oil which solidified upon standing. This was triturated sequentially with water, 1:1 water/MeOH and MeOH and the precipitate was filtered to afford a white solid. The solid was recrystallised from water to give the title compound (*134*) (0.92 g, 30%) as a white solid; ¹H NMR (400 MHz, CDCl₃) δ 8.57 (s, 1H), 8.28 (d, *J* = 8.3 Hz, 1H), 8.01 (s, 1H), 6.86 (dd, *J* = 8.4, 1.7 Hz, 1H), 6.76 (d, *J* = 1.8 Hz, 1H), 4.25 (s, 2H), 3.91 (s, 3H), 2.81 (t, *J* = 11.8 Hz, 2H), 2.64 (tt, *J* = 12.1, 3.6 Hz, 1H), 1.83 (d, *J* = 13.5 Hz, 2H), 1.69 - 1.57 (m, 2H), 1.49 (s, 9H). LCMS Method C: rt 7.00 min; *m/z* = 487.0 [M+H]⁺, 431.0 [M-^tButyl+H]⁺, 485.0 [M-H]⁻.

(h) *tert*-Butyl 4-(4-((4-((3-carbamoylphenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*135*)

[0189] To a mixture of *tert*-butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*134*) (0.098 g, 0.20 mmol), 3-ethynylbenzamide (*131*) (0.038 g, 0.26 mmol), PdCl₂(PPh₃)₂ (9 mg, 0.01 mmol), triphenylphosphine (0.010 g, 0.04 mmol) and copper(I) iodide (0.009 g, 0.05 mmol) in DMF (2 mL), that had been degassed with nitrogen for 10 minutes, was added triethylamine (0.090 mL, 0.65 mmol). The mixture was degassed with nitrogen then heated to 120 °C for 25 minutes under microwave irradiation. On cooling to room temperature the resulting mixture was left overnight, then an additional portion of 3-ethynylbenzamide (*126*) (0.047 g, 0.32 mmol) was added. After degassing with nitrogen the reaction was heated at 120 °C for 25 minutes under microwave irradiation. The volatiles were evaporated under reduced pressure and the residue purified using silica gel column chromatography (20-70% EtOAc/petroleum benzine 40-60 °C) to give the title compound (*135*) (0.073 g, 61%) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ 8.66 (s, 1H), 8.35 (d, *J* = 8.3 Hz, 1H), 8.04 (dd, *J* = 1.4, 1.4 Hz, 1H), 8.02 (s, 1H), 7.93 (ddd, *J* = 7.9, 1.8, 1.2 Hz, 1H), 7.80 (m, 1H), 7.53 (m, 1H), 6.88 (dd, *J* = 8.3, 1.6 Hz, 1H), 6.76 (d, *J* = 1.8 Hz, 1H), 4.27 (m, 2H), 3.91 (s, 3H), 2.81 (m, *J* = 12.2 Hz, 2H), 2.64 (tt, *J* = 12.2, 3.4 Hz, 1H), 1.84 (m, 2H), 1.65 (m, 2H), 1.49 (s, 9H). LCMS Method C: rt 6.59 min; *m/z* 496.1 [(M-Boc)+H]⁺, 540.0 [(M-*t*-Bu)+H]⁺.

(i) *tert*-Butyl 4-(4-((4-(3-carbamoylphenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*136*)


[0190] A mixture of *tert*-butyl 4-(4-((4-((3-carbamoylphenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (*135*) (0.073 g, 0.12 mmol) and 10% palladium on activated carbon (0.042 g) in DMF (3 mL) was stirred under a hydrogen atmosphere for 20 hours. The resulting mixture was filtered and filtrate evaporated to dryness under reduced pressure. The residue was chromatographed on silica gel (20-100% acetone/petroleum benzine 40-60 °C) to give the title compound (*136*) (0.070 g, 95%).

(j) 3-(2-(2-((2-Methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)benzamide (5)

[0191] A mixture of *tert*-butyl 4-(4-((4-(3-carbamoylphenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-methoxyphenyl)piperidine-1-carboxylate (I36) (0.070 g, 0.12 mmol) and TFA (0.200 mL, 2.61 mmol) in THF (3 mL) was stirred 16 hours at room temperature. The mixture was concentrated under reduced pressure and azeotroped with toluene. The residue was then purified using a SCX cartridge (MeOH, 0.5% NH₃/MeOH). The methanolic ammonia eluent was concentrated then dissolved in hot acetonitrile which was allowed to cool to room temperature. The cooled acetonitrile solution was filtered and the filtrate was concentrated, taken up in hot acetonitrile (2 mL) and water (1 mL) then freeze dried to give the title compound (5) (49.4 mg, 85%) as a white solid. A portion of this material was further purified by mass directed auto preparative HPLC to give the title compound (5) (4.5 mg) as a white solid; ¹H NMR (400 MHz, d₆-DMSO) δ 8.90 (m, 1H), 8.57 (s, 1H), 8.41 (s, 1H), 7.93 (s, 1H), 7.77 (s, 1H), 7.71 (m, 1H), 7.62 (dd, J = 7.5, 7.5 Hz, 1H), 7.34 (m, 2H), 6.94 (dd, J = 7.3, 1.6 Hz, 1H), 6.82 (ddd, J = 7.9, 3.4, 1.5 Hz, 1H), 3.82 (s, 3H), 3.12 (m, 2H), 3.05 (m, 4H), 2.89 (m, 1H), 2.68 (m, 2H), 2.20 (m, 1H), 1.78 (m, 2H), 1.64 (m, 2H). LCMS Method C: rt 4.89 min; m/z 500.1[M+H]⁺.

Example 6: 3-(2-(2-((2-Methoxy-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)benzamide (6)

[0192]




[0193] To a solution of 3-(2-(2-((2-methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)benzamide (5) (0.021 g, 0.042 mmol) in anhydrous methanol (2.00 mL) was added 37% aqueous formaldehyde (0.012 mL, 0.16 mmol) and sodium triacetoxyborohydride (0.044 g, 0.21 mmol) under a N₂ atmosphere. The mixture was then stirred for 1.5 hours. The resulting mixture was concentrated under reduced pressure and diluted with EtOAc (10 mL) and washed with sat. aqueous NaHCO₃ (10 mL). The Aqueous was further extracted with EtOAc (10 mL) and the combined organic layers were washed with brine (10 mL) and water (10 mL) then dried using a phase separation cartridge. The organics were concentrated under reduced pressure and purified using silica gel column chromatography (0-30% MeOH/EtOAc + 1% 2 M ethanolic NH₃). The product was then taken up in minimal hot acetonitrile and water and freeze dried to give the title compound (6) (0.016 g, 74%) as a white

solid; ^1H NMR (400 MHz, d_6 -Acetone) δ 8.62 (s, 1H) 8.27 (dd, J = 8.3, 4.1 Hz, 1H), 8.15 (s, 1H), 7.90 (s, 1H), 7.78 (d, J = 7.5 Hz, 1H), 7.41 (m, 3H), 6.99 (d, J = 1.7 Hz, 1H), 6.89 (dd, J = 8.2, 1.6 Hz, 1H), 6.55 (s, 1H), 3.98 (s, 3H), 3.20 (m, 4H), 2.90 (m, 2H), 2.57 (m, 1H), 2.22 (s, 3H), 1.98 (td, J = 11.2, 3.8 Hz, 2H), 1.78 (m, 4H). LCMS Method C: rt 4.97 min; m/z 514.2 $[\text{M}+\text{H}]^+$.

Example 7: 3-(2-(2-((4-(1-Ethylpiperidin-4-yl)-2-methoxyphenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)benzamide (7)


[0194]




[0195] To a solution of 3-(2-(2-((2-methoxy-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)benzamide (5) (0.021 g, 0.04 mmol) in methanol (2.00 mL) was added acetaldehyde (0.010 mL, 0.18 mmol) followed by sodium triacetoxyborohydride (0.044 g, 0.21 mmol) under a N_2 atmosphere. The mixture was then stirred at room temperature for 5 hours. The resulting mixture was concentrated under reduced pressure and purified using silica gel column chromatography (0-30% MeOH/EtOAc + 1% 2 M ethanolic NH_3). The product was further purified using mass directed auto-preparative HPLC to give the title compound (7) (3.5 mg, 16%); ^1H NMR (400 MHz, d_6 -Acetone) δ 8.62 (s, 1H), 8.24 (m, 2H), 7.91 (s, 1H), 7.78 (ddd, J = 7.5, 1.4, 1.4 Hz, 1H), 7.41 (m, 3H), 7.01 (d, J = 1.7 Hz, 1H), 6.89 (dd, J = 8.3, 1.7 Hz, 1H), 6.60 (m, 1H), 3.97 (s, 3H), 3.21 (m, 4H), 3.12 (m, 2H), 2.55 (m, 3H), 2.16 (m, 2H), 1.85 (m, 4H), 1.11 (t, J = 7.2 Hz, 3H). LCMS Method C: rt 4.98 min; m/z 528.2 $[\text{M}+\text{H}]^+$.

Example 8: 2-(2-(2-((2-Ethyl-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (8)

[0196]





(a) *tert*-Butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (137)

[0197] Bis(pinacolato)diboron (0.511 g, 2.01 mmol), potassium acetate (0.592 g, 6.04 mmol), dppf (56 mg, 5 mol%), and PdCl₂(dppf) dichloromethane solvate (83 mg, 5 mol%) were loaded into a Schlenk tube, and purged with nitrogen. A solution *tert*-butyl 4-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1(2H)-carboxylate (116) (1.00 g, 3.02) in dioxane (5 mL) was added, the mixture degassed with 3x vacuum/nitrogen cycles then brought to 80 °C under nitrogen. After 16 hours the mixture was cooled, and added to water (100 mL). The resulting mixture was extracted with dichloromethane (3x50 mL), and the combined DCM extracts washed with brine (50 mL), dried over sodium sulfate and evaporated. Chromatography (40 g silica cartridge, 0-100% ethyl acetate/petroleum benzine 40-60 °C) gave the title compound (137) (383 mg, 62% yield) as a crystalline white solid; ¹H NMR (400 MHz, CDCl₃) δ 6.45 (s, 1H), 3.94 (d, J = 2.7 Hz, 2H), 3.43 (t, J = 5.5 Hz, 2H), 2.22 (s, 2H), 1.45 (s, 9H), 1.26 (s). LCMS Method C: rt 6.48 min; *m/z* 254.2 [M-*t*Bu+2H]⁺, 210.2 [M-Boc+2H]⁺.

(b) *Benzyl* (4-bromo-2-ethylphenyl)carbamate (138)

[0198] 4-Bromo-2-ethylaniline (500 mg, 2.50 mmol) was dissolved in toluene (25 mL), sodium carbonate (397 mg, 3.75 mmol) and benzyl chloroformate (0.428 mL, 3.00 mmol) were added and the mixture stirred under nitrogen at room temperature. After 20 hours water (25 mL) was added, the aqueous phase separated and washed with ethyl acetate (2x25 mL). The combined organic extracts were washed with brine, dried over sodium sulfate and evaporated. Chromatography (40 g silica cartridge, 0-100% ethyl acetate/petroleum benzine 40-60 °C) gave the title compound (*138*) (708 mg, 85%) as a pink solid; ¹H NMR (400 MHz, CDCl₃) δ 7.72 (s, 1H), 7.44 - 7.28 (m, 7H), 6.43 (s, 1H), 5.20 (s, 2H), 2.54 (q, J = 7.6 Hz, 2H), 1.21 (t, J = 7.6 Hz, 3H). LCMS Method C: rt 6.46 min; *m/z* 258.0 [M-PhCH₂O+CH₃OH]⁺.

(c) *tert*-Butyl 4-(4-(((benzyloxy)carbonyl)amino)-3-ethylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (*139*)

[0199] Potassium carbonate (215 mg, 1.55 mmol) the *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (*137*) (160 mg, 0.517 mmol), benzyl (4-bromo-2-ethylphenyl)carbamate (*134*) (182 mg, 0.543 mmol), PdCl₂(dppf)-DCM solvate (22 mg, 5 mol%) and DMF (5 mL) were loaded into a Schlenk tube, and degassed with 3x vacuum/nitrogen cycles. The mixture was brought to 80 °C under nitrogen then after 18 hours cooled and poured into water (100 mL). DCM (75 mL) and brine (50 mL) were added, the aqueous phase was washed with further DCM (2x75 mL), and the combined DCM extracts washed with brine, dried and evaporated. Chromatography (40 g silica cartridge, 0-80% ethyl acetate/petroleum benzine 40-60 °C) gave the title compound (*139*) (129 mg, 57% yield) as a colourless oil; ¹H NMR (400 MHz, CDCl₃) δ 7.77 (s, 1H), 7.45 - 7.31 (m, 5H), 7.22 (dd, J = 8.5, 2.1 Hz, 1H), 7.18 (d, J = 2.1 Hz, 1H), 6.48 (s, 1H), 5.99 (s, 1H), 5.21 (s, 2H), 4.06 (d, J = 2.5 Hz, 2H), 3.63 (t, J = 5.7 Hz, 2H), 2.58 (q, J = 7.6 Hz, 2H), 2.50 (s, 2H), 1.49 (s, 9H), 1.22 (t, J = 7.6 Hz, 3H). LCMS Method C: rt 6.67 min; *m/z* 337.1 [M-Boc+2H]⁺, 381.1 [M-*t*Bu+2H]⁺.

(d) *tert*-Butyl 4-(4-amino-3-ethylphenyl)piperidine-1-carboxylate (*140*)

[0200] *tert*-Butyl 4-(4-(((benzyloxy)carbonyl)amino)-3-ethylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (*139*) (0.500 g, 1.14 mmol) was dissolved in ethanol (20 mL), and a slurry of 10% Pd/C (0.25 g) in ethanol (2 mL) was added. The mixture was stirred under hydrogen for 18 hours, then filtered through celite, washing the celite with ethanol (30 mL). The combined filtrates were evaporated, and chromatography (12 g silica cartridge, 0-80% ethyl acetate/petroleum benzine 40-60 °C) gave the title compound (*140*) (0.212 g, 61% yield) as a pink syrup; ¹H NMR (400 MHz, CDCl₃) δ 6.90 (d, J = 2.0 Hz, 1H), 6.87 (dd, J = 8.0, 2.1 Hz, 1H), 6.63 (d, J = 8.0 Hz, 1H), 4.21 (s, 2H), 3.55 (s, 2H), 2.78 (t, J = 12.2 Hz, 2H), 2.59 - 2.46 (m, 3H), 1.78 (d, J = 13.2 Hz, 2H), 1.66 - 1.53 (m, overlaps with water), 1.48 (s, 9H), 1.25 (t, J = 7.5 Hz, 3H). LCMS Method C: rt 5.33 min; *m/z* 249.2 [M-*t*Bu+2H]⁺, 205.2 [M-Boc+2H]⁺.

(e) *tert*-Butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-ethylphenyl)piperidine-1-carboxylate (I41)

[0201] A solution of 2,4-dichloro-5-(trifluoromethyl)pyrimidine (166 mg, 0.766 mmol) in 1:1 DCE: *t*-BuOH (5 mL) was stirred at 0 °C under nitrogen. 1.0 M Zinc(II) chloride in ether (0.77 mL, 0.77 mmol) was added and the mixture stirred for one hour. *tert*-Butyl 4-(4-amino-3-ethylphenyl)piperidine-1-carboxylate (*I40*) (212 mg, 0.70 mmol) in 1:1 DCE: *t*-BuOH (5 mL) was added dropwise, and after 30 minutes triethylamine (146 µL, 1.05 mmol) in 1:1 DCE: *t*-BuOH (5 mL) was added and the mixture allowed to slowly come to room temperature. After 18 hours the mixture was concentrated, and chromatography (12 g silica cartridge, 0-50% ethyl acetate/cyclohexane) gave the title compound (*I41*) (283 mg, 84% yield) as a pale pink solid; ¹H NMR (400 MHz, CDCl₃) δ 8.51 (d, J = 0.6 Hz, 1H), 7.60 (d, J = 8.3 Hz, 1H), 7.15 - 7.07 (m, 3H), 4.25 (s, 2H), 2.80 (t, J = 12.1 Hz, 2H), 2.69 - 2.59 (m, 3H), 1.83 (d, J = 12.9 Hz, 2H), 1.69 - 1.59 (m, 2H), 1.49 (s, 9H), 1.23 (t, J = 7.6 Hz, 3H). LCMS Method C: rt 6.94 min; *m/z* 429.1 [M-*t*Bu+2H]⁺, 385.1 [M-Boc+2H]⁺; *m/z* 483.1 [M-H]⁻.

(f) *tert*-Butyl 4-(3-ethyl-4-((4-((2-(2-methoxy-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (I42)

[0202] A mixture of *tert*-butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-ethylphenyl)piperidine-1-carboxylate (*I41*) (283 mg, 0.584 mmol), methyl 2-(2-ethynylphenyl)acetate (*I4*) (122 mg, 0.700 mmol), copper(I) iodide (17 mg, 15 mol%), triphenylphosphine (23 mg, 15 mol%), bis(triphenylphosphine)palladium(II) chloride (41 mg, 10 mol%), triethylamine (0.33 mL, 2.3 mmol) and DMF (3 mL) was degassed with a nitrogen then heated under microwave irradiation (120 °C/15 min). The cooled mixture was poured into water (30 mL) and extracted with ethyl acetate (3x25 mL). The combined organic extracts were washed with brine (2x50 mL), dried over sodium sulfate and evaporated. Chromatography (12 g silica cartridge, 0-80% ethyl acetate/petroleum benzine 40-60 °C) gave the title compound (*I42*) (311 mg, 86% yield) as a brown oil; ¹H NMR (400 MHz, CDCl₃) δ 8.57 (s, 1H), 7.73 - 7.65 (m, 2H), 7.46 - 7.28 (m), 7.15 - 7.08 (m), 4.25 (s, 2H), 3.95 (s, J = 9.6 Hz, 2H), 3.70 (s, 3H), 2.80 (t, J = 12.1 Hz, 2H), 2.70 - 2.59 (m, 3H), 1.84 (d, J = 12.9 Hz, 2H), 1.65 (td, J = 12.8, 4.1 Hz, 2H), 1.49 (s, 9H), 1.25 (t, J = 7.6 Hz, 3H). LCMS: rt 7.01 min; *m/z* 623.1 [M+H]⁺, 567.1 [M-*t*Bu+2H]⁺, 523.1 [M-Boc+2H]⁺; *m/z* 621.2 [M-H]⁻.

(g) *tert*-Butyl 4-(3-ethyl-4-((4-((2-(2-methoxy-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (I43)

[0203] *tert*-Butyl 4-(3-ethyl-4-((4-((2-(2-methoxy-2-oxoethyl)phenyl)ethynyl)-5-

(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*142*) (311 mg, 0.499 mmol) was dissolved in ethanol (20 mL), and a slurry of Pd/C (150 mg) in ethanol (2 mL) was added. The mixture was stirred under hydrogen for 18 hours, filtered through celite, washing the celite with ethanol (20 mL) and the filtrate concentrated. The mixture was taken up in ethanol (20 mL), and a slurry of Pd/C (150 mg) in ethanol (2 mL) was added followed by triethylamine (20 μ L). The mixture was stirred under hydrogen for 18 hours, filtered through celite, the celite washed with ethanol (10 mL) and the combined filtrates evaporated. The residue was taken up in DMF (10 mL), a slurry of Pd/C (150 mg) in DMF (2 mL) was added and the mixture stirred under hydrogen. After 16 hours the mixture was filtered through celite, and the celite washed with ethyl acetate (100 mL). The combined filtrate was evaporated to give a pale green oil. Chromatography (12 g silica cartridge, 0-60% ethyl acetate/petroleum benzene 40-60 °C) gave the title compound (*143*) (177.1 mg, 57% yield) as a pale yellow oil. ^1H NMR (400 MHz, CDCl_3) δ 8.49 (d, J = 0.5 Hz, 1H), 7.73 (d, J = 8.1 Hz, 1H), 7.25 - 7.16 (m, 4H), 7.12 - 7.07 (m, 2H), 7.06 (s, 1H), 4.25 (br s, 2H), 3.72 (s, 2H), 3.67 (s, 3H), 3.14 - 3.01 (m, 4H), 2.81 (t, J = 12.2 Hz, 2H), 2.72 - 2.59 (m, 3H), 1.84 (d, J = 13.0 Hz, 2H), 1.69 - 1.56 (m, overlaps with water in solvent), 1.49 (s, 9H), 1.26 (t, J = 7.1 Hz, 3H). LCMS: rt 7.16 min; m/z 627.2 $[\text{M}+\text{H}]^+$; 571.1 $[\text{M}-\text{tBu}+2\text{H}]^+$; m/z 625.2 $[\text{M}-\text{H}]^-$.

(h) *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-ethylphenyl)piperidine-1-carboxylate (*144*)

[0204] *tert*-Butyl 4-(3-ethyl-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*143*) (177 mg, 0.282 mmol) was dissolved in THF (10 mL), and a solution of lithium hydroxide monohydrate (59.0 mg, 1.41 mmol) in water (2 mL) was added. The mixture was stirred for 18 hours then concentrated. The residue was suspended in saturated sodium bicarbonate (20 mL), and extracted with ethyl acetate (3x20 mL). The combined ethyl acetate phases were washed with brine, dried (sodium sulphate) and evaporated. The residue was dissolved in THF (10 mL) and DMF (1 mL) at 30 °C and HOBt (50 mg, 0.37 mmol), EDCI (65 mg, 0.37 mmol) and DIPEA (0.246 mL, 1.41 mmol) were added. After ten minutes ammonium carbonate (135 mg, 1.41 mmol) was added and the mixture was stirred for 18 hours at 30 °C. The mixture was concentrated, the residue diluted with saturated sodium bicarbonate (25 mL) and extracted with ethyl acetate (3x 25 mL). The combined ethyl acetate phases were washed with brine (25 mL), dried (sodium sulfate) and evaporated. Chromatography (12 g silica cartridge, 0-100% ethyl acetate/petroleum benzene 40-60 °C) gave the title compound (*144*) (122 mg, 71% over two steps) as a white solid; ^1H NMR (400 MHz, CDCl_3) δ 8.49 (s, 1H), 7.56 (d, J = 8.1 Hz, 1H), 7.28 - 7.22 (m, overlaps with CHCl_3), 7.22 - 7.15 (m, 1H), 7.13 - 7.05 (m, 3H), 5.38 (s, 1H), 5.15 (s, 1H), 4.25 (s, 2H), 3.64 (s, 2H), 3.11 - 2.99 (m, 4H), 2.80 (t, J = 12.1 Hz, 2H), 2.69 - 2.60 (m, 3H), 1.84 (d, J = 12.3 Hz, 2H), 1.69 - 1.60 (m, 2H), 1.49 (s, 9H), 1.23 (t, J = 7.6 Hz, 3H). LCMS: rt 6.65 min; m/z 612.2 $[\text{M}+\text{H}]^+$; 556.1 $[\text{M}-\text{tBu}+2\text{H}]^+$; m/z 610.2 $[\text{M}-\text{H}]^-$.


(i) 2-(2-(2-(2-((2-Ethyl-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-

yl)ethyl)phenyl)acetamide (8)

[0205] *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-ethylphenyl)piperidine-1-carboxylate (*144*) (120 mg, 0.20 mmol) was dissolved in DCM (20 mL), TFA (2 mL) was added and the mixture stirred for 16 hours at room temperature. The mixture was concentrated and the residue suspended in 10% sodium hydroxide (10 mL) and brine (10 mL). The mixture was extracted with ethyl acetate (4x20 mL), the combined extracts washed with brine, dried (sodium sulphate) evaporated and the residue evaporated for ether to give the title compound (**8**) (93 mg, 93% yield) as a white solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 9.47 (s, 1H), 8.49 (s, 1H), 7.37 (s, 1H), 7.26 - 7.18 (m, 2H), 7.17 - 7.09 (m, 4H), 7.05 (dd, *J* = 8.2, 1.9 Hz, 1H), 6.89 (s, 1H), 3.44 (s, 2H), 3.09 - 2.88 (m, 6H), 2.62 - 2.54 (m, overlaps with DMSO), 1.70 (d, *J* = 11.4 Hz, 2H), 1.52 (qd, *J* = 12.3, 3.7 Hz, 2H), 1.10 (t, *J* = 7.5 Hz, 3H). LCMS Method C: 4.96 min; *m/z* 512.2 [*M*+H]⁺, 534.2 [*M*+Na]⁺; *m/z* 510.2 [*M*-H]⁻.

Example 9: 2-(2-(2-(2-((2-Ethyl-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (9)

[0206]




[0207] 2-(2-(2-(2-((2-Ethyl-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (**8**) (84 mg, 0.16 mmol) was dissolved in methanol (8 mL), 37% formaldehyde solution (53 µL, 0.66 mmol) was added and the mixture stirred for ten minutes at room temperature. Sodium tris(acetoxy)borohydride (174 mg, 0.821 mmol) was added, and after two hours the mixture was concentrated. The residue was diluted with 10% sodium hydroxide (15 mL) and brine (15 mL), and the mixture extracted with ethyl acetate (4x25 mL). The combined ethyl acetate phases were washed with brine, dried over sodium sulfate, evaporated and the residue evaporated from DCM to give the title compound (**9**) (72 mg, 84% yield) as a white solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 9.46 (s, 1H), 8.49 (s, 1H), 7.36 (s, 1H), 7.21 (t, *J* = 4.6 Hz, 2H), 7.18 - 7.10 (m, 4H), 7.06 (dd, *J* = 8.2, 1.6 Hz, 1H), 6.88 (s, 1H), 3.44 (s, 2H),

[0208] 3.07 - 2.90 (m, 4H), 2.86 (d, *J* = 11.3 Hz, 2H), 2.57 (q, *J* = 7.5 Hz, 2H), 2.48 - 2.37 (m, 1H), 2.19 (s, 3H), 2.02 - 1.89 (m, 2H), 1.79 - 1.59 (m, 4H), 1.09 (t, *J* = 7.5 Hz, 3H). LCMS

Method C: rt 5.04 min; m/z 526.2 $[M+H]^+$; m/z 524.2 $[M-H]^-$.

Example 10: 2-(2-(2-(2-((4-(Piperidin-4-yl)-2-(trifluoromethoxy)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (10)

[0209]



(a) 2-(2-Ethynylphenyl)acetamide (145)

[0210] To a stirred solution of methyl 2-(2-ethynylphenyl)acetate (14) (0.200 g, 1.15 mmol) in methanol (2 mL) at 0 °C in a thick-walled Schlenk tube was added magnesium nitride (0.290 g, 2.87 mmol) in a single portion. The tube was sealed immediately and allowed to warm to room

temperature in a water bath over 1 hour and then heated at 80 °C for 22 hours. After cooling to room temperature the resulting mixture was diluted with EtOAc (100 mL) and saturated aqueous NaHCO₃ (80 mL). The layers were separated and the aqueous layer was extracted with EtOAc (80 mL), neutralised with 2 M aq. HCl and extracted with EtOAc (2 x 80 mL). The organic layers were combined, washed with brine (100 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to give a yellow solid. Silica gel chromatography (40 g Si Cartridge, 0-50% EtOAc in dichloromethane) gave the title compound (*I45*) (0.093 g, 51% yield) as a white solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 7.44 (dd, *J* = 7.6, 1.0 Hz, 1H), 7.40 - 7.28 (m, 3H), 7.24 (td, *J* = 7.4, 1.7 Hz, 1H), 6.93 (s, 1H), 4.32 (s, 1H), 3.59 (s, 2H). LCMS Method C: rt 4.68 min; *m/z* 160.2 [M+H]⁺.

(b) Benzyl (4-bromo-2-(trifluoromethoxy)phenyl)carbamate (*I46*)

[0211] 4-Bromo-2-(trifluoromethoxy)aniline (1.0 g, 3.9 mmol) was dissolved in dry toluene (25 mL), sodium carbonate (0.621 g, 5.86 mmol) and benzyl chloroformate (0.669 mL, 4.69 mmol) was added and the mixture stirred under nitrogen at room temperature for 22 hours. The reaction was then heated to 80 °C and stirred at this temperature for 17 hours and then heated further to reflux and stirred for 22 hours. Water (100 mL) was added to the cooled mixture, the aqueous phase separated and washed with ethyl acetate (2x100 mL). The combined organic extracts were washed with 0.5 M aq. citric acid (70 mL), water (70 mL), brine (70 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to give a pink solid. The crude material was purified by silica gel chromatography (40 g Si cartridge, 0-20% EtOAc in 40-60 °C petroleum benzene) to give the title compound (*I46*) (1.153 g, 76% yield) as a white solid; ¹H NMR (400 MHz, CDCl₃) δ 8.15 (d, *J* = 8.7 Hz, 1H), 7.44 - 7.36 (m, 7H), 6.95 (s, 1H), 5.22 (s, 2H). LCMS Method C: rt 6.67 min; *m/z* 389.9 [M-H]⁻.

(c) *tert*-Butyl 4-(4-amino-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (*I47*)

[0212] A solution of 2 M aq. Na₂CO₃ (1.85 mL, 3.70 mmol) was added to a mixture of *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (*I37*) (~50% pure, 0.915 g, 1.480 mmol), benzyl (4-bromo-2-(trifluoromethoxy)phenyl)carbamate (*I46*) (0.635 g, 1.63 mmol), LiCl (0.125 g, 2.96 mmol), Pd(PPh₃)₂Cl₂ (0.052 g, 0.074 mmol) and TBAB (0.048 g, 0.15 mmol) in dry 1,4-dioxane (15 mL). The reaction mixture was stirred at 80 °C for 17 hours, then filtered through celite, which was washed with EtOAc and MeOH and the filtrate was concentrated *in vacuo* to give a pale yellow gum. The crude material was purified by silica gel chromatography (40 g Si Cartridge, 0-15% EtOAc in petroleum benzene 40-60 °C) to give a pale yellow gum (0.515 g). To a solution of the intermediate in EtOAc (20 mL) was added 10% Pd/C (80 mg) in EtOAc (5 mL). The reaction was then stirred at room temperature for 18 hours under an atmosphere of hydrogen and then filtered through a pad of

celite, which was washed with EtOAc (100 mL). The solvent was removed *in vacuo* to give crude product which was purified by silica gel chromatography (40 g Si Cartridge, 0-30% EtOAc in petroleum benzine 40-60 °C) to give the title compound (*147*) (80% purity, 0.220 g, 33% yield over 2 steps) as a pale yellow gum; ¹H NMR (400 MHz, CDCl₃) δ 6.98 - 6.95 (m, 1H), 6.92 (dd, *J* = 8.2, 1.8 Hz, 1H), 6.74 (d, *J* = 8.2 Hz, 1H), 4.30 - 4.13 (m, 2H), 3.77 (s, 2H), 2.77 (t, *J* = 12.5 Hz, 2H), 2.54 (tt, *J* = 12.1, 3.5 Hz, 1H), 1.78 (d, *J* = 13.1 Hz, 2H), 1.57 - 1.50 (m, 2H, obscured by water signal), 1.48 (s, 9H).

(d) *tert*-Butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (*148*)

[0213] 2,4-Dichloro-5-(trifluoromethyl)pyrimidine (0.110 g, 0.506 mmol) was stirred in a 1:1 *t*-BuOH:1,2-dichloroethane mixture (10 mL) at 0 °C. A 1.0 M ZnCl₂ solution in diethyl ether (0.578 mL, 0.578 mmol) was added cautiously, after addition the reaction was left stirring at 0 °C for 30 minutes. A solution of *tert*-butyl 4-(4-amino-3-(trifluoromethoxy) phenyl)piperidine-1-carboxylate (*147*) (impure, ~80% pure, 0.217 g, 0.482 mmol) in 1:1 *t*-BuOH:1,2-dichloroethane (5 mL) was added drop-wise at 0 °C followed by a solution of NEt₃ (0.081 mL, 0.58 mmol) in 1:1 *t*-BuOH:1,2-dichloroethane (5 mL) and the reaction was allowed to warm to room temperature and was stirred for 18 hours, then at 60 °C for 24 hours. The organic solvents were evaporated *in vacuo* and the crude gum was purified by silica gel chromatography (40 g Si cartridge, 0-40% EtOAc in petroleum benzine 40-60 °C) to give the title compound (*148*) (0.085 g, 33% yield) as a pale yellow oily solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 10.38 (s, 1H), 8.72 (s, 1H), 7.52 (d, *J* = 8.1 Hz, 1H), 7.33 - 7.29 (m, 2H), 4.13-4.04 (m, 2H), 2.88 - 2.72 (m, 3H), 1.79 (d, *J* = 12.1 Hz, 2H), 1.56 - 1.44 (m, 2H), 1.42 (s, 9H). LCMS Method C: rt 7.02 min; *m/z* 539.0, 541.0 [M-H]⁻.

(e) *tert*-Butyl 4-(4-((4-((2-(2-amino-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (*149*)

[0214] To a nitrogen de-gassed solution of 2-(2-ethynylphenyl)acetamide (*141*) (0.029 g, 0.18 mmol) and *tert*-butyl 4-(4-((4-chloro-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (*148*) (0.082 g, 0.15 mmol) in dry DMF (4 mL) were added triethylamine (0.085 mL, 0.61 mmol), triphenylphosphine (6.0 mg, 0.023 mmol), trans-dichlorobis (triphenylphosphine)palladium (II) (0.011 g, 0.015 mmol) and Cu(I)I (4.0 mg, 0.023 mmol). The reaction mixture was heated under microwave irradiation at 120 °C for 20 minutes, concentrated to dryness *in vacuo* and purified by silica gel chromatography (12 g Si cartridge, 0-100% EtOAc in petroleum benzine 40-60 °C) to give the title compound (*149*) (0.073 g, 73% yield) as a pale yellow solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 10.11 (s, 1H), 8.74 (s, 1H), 7.58 (dd, *J* = 7.7, 6.2 Hz, 2H), 7.53 - 7.47 (m, 1H), 7.42 - 7.27 (m, 5H), 6.99 (s, 1H),

4.14 - 4.03 (m, 2H), 3.67 (s, 2H), 2.87 - 2.72 (m, 3H), 1.79 (d, $J = 12.8$ Hz, 2H), 1.50 (qd, $J = 12.5, 4.1$ Hz, 2H), 1.42 (s, 9H). LCMS Method C: rt 6.63 min; m/z 564.0 $[M-Boc+2H]^+$.

(f) *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (I50)


[0215] *tert*-Butyl 4-(4-((4-((2-(2-amino-2-oxoethyl)phenyl)ethynyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (I49) (72.0 mg, 0.108 mmol) was dissolved in dry DMF (7 mL) under an atmosphere of nitrogen, and a slurry of 10% Pd(OH)₂/C (0.050 g) in EtOAc (2 mL) was added. The mixture then was stirred vigorously under hydrogen for 24 hours. Upon completion, the reaction was filtered through a pad of celite, which was washed with EtOAc (40 mL). The combined filtrates were concentrated *in vacuo* to give a pale yellow oil. The crude product was purified by silica gel chromatography (12 g Si Cartridge, 0-100% EtOAc in petroleum benzine 40-60 °C) to give the title compound (I50) (0.064 g, 88% yield) as an off-white solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 9.80 (s, 1H), 8.60 (s, 1H), 7.61 (d, $J = 8.1$ Hz, 1H), 7.39 (s, 1H), 7.32 - 7.26 (m, 2H), 7.21 (dt, $J = 6.7, 3.4$ Hz, 1H), 7.18 - 7.08 (m, 3H), 6.90 (s, 1H), 4.09 (d, $J = 12.1$ Hz, 2H), 3.45 (s, 2H), 3.09 - 2.94 (m, 4H), 2.88 - 2.72 (m, 3H), 1.79 (d, $J = 12.3$ Hz, 2H), 1.59 - 1.38 (m, 11H). LCMS Method C: rt 6.75 min, m/z 668.1 $[M+H]^+$.

(g) 2-(2-(2-(2-((4-(Piperidin-4-yl)-2-(trifluoromethoxy)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (10)

[0216] *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-(trifluoromethoxy)phenyl)piperidine-1-carboxylate (I50) (0.062 g, 0.093 mmol) was dissolved in dry DCM (6 mL) under an atmosphere of nitrogen. Trifluoroacetic acid (0.142 mL, 1.86 mmol) was added to the solution and the reaction was stirred at room temperature for 20 hours. Volatiles were removed *in vacuo*, EtOAc (50 mL) and 2 M aq. NaOH (50 mL) were added to the residue and the layers were separated. The aqueous layer was extracted with EtOAc (2x70 mL), the combined organics were washed with water (50 mL), brine (50 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to give an off-white solid. The crude material was purified by silica gel chromatography (12 g Si Cartridge, 0-100% EtOAc in petroleum benzine 40-60 °C, then 0-100% methanol in EtOAc, then 1% ammonia in methanol), to give the title compound (10) (0.042 g, 80% yield) as a white solid; ¹H NMR (400 MHz, *d*₆-DMSO) δ 9.78 (s, 1H), 8.59 (s, 1H), 7.59 (d, $J = 8.3$ Hz, 1H), 7.39 (s, 1H), 7.29 - 7.17 (m, 3H), 7.18 - 7.06 (m, 3H), 6.90 (s, 1H), 3.45 (s, 2H), 3.09 - 2.92 (m, 6H), 2.71 - 2.53 (m, 3H), 1.71 (d, $J = 11.7$ Hz, 2H), 1.49 (qd, $J = 12.1, 3.7$ Hz, 2H). LCMS Method C: rt 5.07 min; m/z 568.1 $[M+H]^+$.

Example 11: 2-(2-(2-(2-((4-(1-Methylpiperidin-4-yl)-2-(trifluoromethoxy)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (11)


[0217]




[0218] To a suspension of 2-(2-(2-(2-((4-(piperidin-4-yl)-2-(trifluoromethoxy)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (10) (0.039 g, 0.069 mmol) in anhydrous methanol (4 mL) was added a 37% aq. solution of formaldehyde (0.020 mL, 0.28 mmol) under an atmosphere of nitrogen followed by sodium triacetoxyborohydride (0.073 g, 0.34 mmol). The reaction was stirred at room temperature for 3 hours. The volatiles were then removed *in vacuo* and the residue was diluted with EtOAc (50 mL) and sat. aq. NaHCO₃ (30 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2x30 mL), the combined organic layers were washed with brine (40 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to give a gum which was taken up in DCM (~ 10 mL) and methanol (~ 1 mL) and concentrated *in vacuo*. The process was repeated with only DCM twice. The resulting gum was then suspended in diethyl ether (5 mL) and the solvent was removed *in vacuo*. The procedure was repeated to give the title compound (11) (0.037 g, 93% yield) as a white fluffy solid: ¹H NMR (400 MHz, *d*₆-DMSO) δ 9.79 (s, 1H), 8.59 (s, 1H), 7.60 (d, *J* = 8.3 Hz, 1H), 7.39 (s, 1H), 7.28 (dd, *J* = 8.4, 1.9 Hz, 1H), 7.26 - 7.23 (m, 1H), 7.23 - 7.19 (m, 1H), 7.18 - 7.08 (m, 3H), 6.89 (s, 1H), 3.45 (s, 2H), 3.08 - 2.93 (m, 4H), 2.90 - 2.83 (m, 2H), 2.57 - 2.52 (m, 1H, obscured by residual solvent signal), 2.19 (s, 3H), 1.96 (td, *J* = 11.5, 2.0 Hz, 2H), 1.80 - 1.72 (m, 2H), 1.65 (ddd, *J* = 24.6, 12.4, 3.7 Hz, 2H). LCMS Method C: rt 5.11 min; *m/z* 582.1 [M+H]⁺.

Example 12: 2-(2-(2-(2-((2-methyl-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (12)

[0219]





(a) Benzyl (4-bromo-2-methylphenyl)carbamate (151)

[0220] 2-Bromo-4-methylaniline (5.00 g, 26.9 mmol), benzyl chloroformate (5.75 mL, 40.3 mmol), Na_2CO_3 (4.27 g, 40.3 mmol) and toluene (100 mL) were stirred under nitrogen at room temperature for 20 hours. The resulting mixture was diluted with ethyl acetate and washed with water (100 mL). The organic layer was dried (MgSO_4) and the volatiles removed by evaporation under reduced pressure. Petroleum benzene 40-60 °C was added and the resulting precipitate collected by filtration to give the title compound (151) as colourless needles (8.50 g, 99%); ^1H NMR (400 MHz, COCl_3) δ 7.80 - 7.67 (m, 1H), 7.44 - 7.28 (m, 6H), 6.40 (s, 1H), 5.20 (s, 2H), 2.21 (s, 3H). LCMS Method C: rt 6.36 min; m/z 320 $[\text{M}+1]^+$.

(b) *tert*-Butyl 4-(4-(((benzyloxy)carbonyl)amino)-3-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (152)

[0221] A suspension of benzyl (4-bromo-2-methylphenyl)carbamate (151) (1.00 g, 3.12 mmol), *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (37) (1.16 g, 3.74 mmol), $\text{PdCl}_2(\text{dppf})\text{-DCM}$ complex (310 mg, 0.374 mmol) and potassium carbonate (1.29 g, 9.36 mmol) in DMF (30 mL) was stirred under nitrogen at 80 °C for 16 hours. The volatiles were evaporated under reduced pressure and the residue adsorbed onto silica gel. Chromatography on silica gel (0-30% ethyl acetate/petroleum benzene 40-60 °C) gave the title compound (152) (949 mg, 72%) as a yellow liquid; ^1H NMR (400 MHz, CDCl_3) δ 7.83 - 7.70 (m, 1H), 7.41 - 7.34 (m, 5H), 7.21 (dd, J = 8.5, 1.9 Hz, 1H), 7.16 (d, J = 1.8 Hz, 1H), 6.56 (s, 1H), 5.97 (s, 1H), 5.20 (s, 2H), 4.05 (d, J = 2.9 Hz, 2H), 3.61 (t, J = 5.7 Hz, 2H), 2.48 (d, J = 1.5 Hz, 2H), 2.23 (s, 3H), 1.50 (d, J = 2.6 Hz, 9H).

(c) *tert*-Butyl 4-(4-amino-3-methylphenyl)piperidine-1-carboxylate (153)

[0222] A suspension of *tert*-butyl 4-(4-(((benzyloxy)carbonyl)amino)-3-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (152) (798 mg, 2.49 mmol) and 10% Pd/C (250 mg) in MeOH (50 mL) was stirred under hydrogen (1 atm) for 16 hours at room temperature. The

resulting mixture was filtered through celite, washing with MeOH, then the volatiles were removed by evaporation under reduced pressure to give the title compound (*I53*) (550 mg, 76%) as a purple/brown liquid; ^1H NMR (400 MHz, CDCl_3) δ 6.95 (d, J = 6.7 Hz, 3H), 4.22 (s, 2H), 2.77 (s, 2H), 2.54 (s, 1H), 2.29 (s, 3H), 1.77 (d, J = 12.7 Hz, 2H), 1.56 (dd, J = 12.6, 3.7 Hz, 9H), 1.47 (s, 9H). LCMS Method C rt: 5.09 min; m/z 235.1 $[\text{M}-\text{tBu}+2]^+$, 191.2 $[\text{M}-\text{Boc}+2]^+$.

(d) Methyl 2-(2-(2-(2-((2-methyl-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*I54*)

[0223] A solution of methyl 2-(2-(2-(2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*I15*) (450 mg, 1.11 mmol), *tert*-butyl 4-(4-amino-3-methylphenyl)piperidine-1-carboxylate (*I53*) (390 mg, 1.34 mmol) and TFA (0.25 mL) in 2,2,2-TFE (3 mL) was heated at 120 °C under microwave irradiation for 30 minutes. The resulting mixture was adsorbed onto silica gel and chromatographed (0-50% MeOH/DCM) to give the title compound (*I54*) (478 mg, 83%) as a viscous liquid; ^1H NMR (400 MHz, CDCl_3) δ 9.14 (bs, 1H), 8.76 (bs, 1H), 8.51 (s, 1H), 7.58 (d, J = 8.2 Hz, 1H), 7.21 (ddt, J = 10.7, 7.4, 4.3 Hz, 4H), 7.15 - 7.06 (m, 1H), 3.65 (s, 2H), 3.60 (d, J = 12.6 Hz, 2H), 3.49 (s, 3H), 3.15 - 3.02 (m, 6H), 2.87 - 2.71 (m, 1H), 2.32 (s, 3H), 2.07 (d, J = 7.8 Hz, 4H). LCMS Method C: rt 5.24 min; m/z 513.2 $[\text{M}+1]^+$.

(e) Methyl 2-(2-(2-(2-((2-methyl-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*I55*)


[0224] Methyl 2-(2-(2-(2-((2-methyl-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*I54*) (1.10 g, 2.15 mmol) was dissolved in dry MeOH (20 mL) and formaldehyde solution (37% aq; 348 μL , 4.29 mmol) was added. Sodium triacetoxyborohydride (2.27 g, 10.7 mmol) was added under nitrogen and the resultant mixture was stirred at room temperature for 16 hours. Ethyl acetate (50 mL) was added and the mixture was washed with 10% NaHCO_3 solution (20 mL). The organic layer was separated, dried (MgSO_4) then the volatiles removed by evaporation under reduced pressure. Chromatography (SiO_2 , 0-50% MeOH/DCM) gave the title compound (*I55*) (480 mg, 42%) as a cream solid; ^1H NMR (400 MHz, CDCl_3) δ 8.54 (s, 1H), 7.83 - 7.73 (m, 1H), 7.35 (s, 1H), 7.32 - 7.19 (m, 3H), 7.19 - 7.09 (m, 2H), 7.07 - 6.97 (m, 1H), 3.76 (s, 2H), 3.70 (s, 3H), 3.23 (d, J = 11.5 Hz, 2H), 3.17 - 3.04 (m, 4H), 2.61 - 2.51 (m, 1H), 2.50 (s, 3H), 2.35 (s, 3H), 2.33 - 2.24 (m, 2H), 2.02 - 1.84 (m, 4H). LCMS Method C: rt 5.25 min; m/z 527.2 $[\text{M}+1]^+$.

(f) 2-(2-(2-(2-((2-Methyl-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (12)

[0225] Methyl 2-(2-(2-(2-((2-methyl-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*155*) (476 mg, 0.906 mmol) and LiOH.H₂O (113 mg, 2.71 mmol) in a mixture of THF (20 mL), water (4 mL) and MeOH (2 mL) were stirred at room temperature for 24 hours. The volatiles were evaporated under reduced pressure to give a light yellow solid which was dissolved in dry DMF (10 mL) and dry THF (10 mL). HOBT (171 mg, 1.26 mmol), EDCI (196 mg, 1.26 mmol), ammonium carbonate (458 mg, 4.87 mmol) and DIPEA (829 μ L, 4.87 mmol) were added and the resulting mixture was stirred at room temperature for 16 hours. The volatiles were evaporated under reduced pressure and the residue taken up in ethyl acetate. The resulting solution was washed with 10% NaHCO₃, the layers separated and the organic layer was dried (MgSO₄). The volatiles were evaporated under reduced pressure and the residue chromatographed (SiO₂, 0-100% MeOH/DCM) to give the title compound (*12*) (358 mg, 52%) as a cream solid; ¹H NMR (400 MHz, CDCl₃) δ 8.51 (s, 1H), 7.58 (d, *J* = 8.9 Hz, 1H), 7.34 (s, 1H), 7.28 - 7.19 (m, 4H), 7.14 - 7.09 (m, 2H), 5.66 (s, 1H), 5.49 (s, 1H), 3.65 (s, 2H), 3.13 - 3.04 (m, 4H), 3.04 - 2.97 (m, 2H), 2.54 - 2.41 (m, 1H), 2.35 (s, 3H), 2.30 (s, 3H), 2.08 (td, *J* = 11.3, 3.8 Hz, 2H), 1.88 - 1.80 (m, 4H). LCMS Method C: rt 4.91 min; *m/z* 512.2 [M+1]⁺, 510.2 [M-1]⁻.

Example 13: 2-(2-(2-(2-((2-Fluoro-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (*13*)

[0226]



(a) *tert*-Butyl (2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)carbamate (*156*)

[0227] A mixture of 2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (1.00 g, 4.22 mmol) and Boc_2O (1.14 g, 5.21 mmol) in toluene (15 mL) was heated at reflux for 16 hours. A further portion of Boc_2O (1.04 g, 4.8 mmol) was added and the mixture heated for a further 20 hours then another portion of Boc_2O (1.4 g, 6.4 mmol) was added. After heating at reflux for a further 24 hours a catalytic amount of DMAP was added and the mixture heated at reflux for 30 minutes before concentrating under reduced pressure to give the title compound (**56**) (1.42 g, 99%); ^1H NMR (400 MHz, CDCl_3) δ 7.52 (m, 2H), 7.14 (dd, J = 7.5, 7.5 Hz, 1H), 1.40 (s, 12H), 1.35 (s, 9H).

(b) *tert*-Butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (157**)**

[0228] To a mixture of *tert*-butyl 4-(((trifluoromethyl)sulfonyl)oxy)-5,6-dihydropyridine-1(2H)-carboxylate (**116**) (0.504 g, 1.52 mmol), *tert*-butyl (2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)carbamate (**162**) (0.496 g, 1.47 mmol), and $\text{PdCl}_2(\text{PPh}_3)_2$ (0.055 g, 0.078 mmol) under a N_2 atmosphere was added dioxane (15 mL) and the mixture was bubbled with N_2 for 5 minutes before addition of 3.0 M aqueous sodium carbonate (1.5 mL, 4.5 mmol). The reaction was heated at reflux for 6 hours then concentrated under reduced pressure. The residue was chromatographed on silica gel (0-25% EtOAc/petroleum benzine 40-60 °C) to give the title compound (**157**) (0.409 g, 71%); ^1H NMR (400 MHz, CDCl_3) δ 7.12 (m, 3H), 6.09 (s, 1H), 4.08 (m, J = 2.8 Hz, 2H), 3.63 (dd, J = 5.6, 5.6 Hz, 2H), 2.49 (m, 2H), 1.42 (s, 18H).

(c) *tert*-Butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-fluorophenyl)piperidine-1-carboxylate (154**)**

[0229] A mixture of *tert*-butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (**153**) (0.409 g, 1.04 mmol) and 10% Pd/C (0.043 g) in EtOAc (20 mL) was stirred for 16 hours at room temperature under a H_2 atmosphere. The mixture was filtered through celite and the filtrate concentrated under reduced pressure to give the title compound (**154**) (0.376 g, 92%) as a white foam; ^1H NMR (400 MHz, CDCl_3) δ 7.09 (dd, J = 8.1, 8.1 Hz, 1H), 6.95 (m, 2H), 4.25 (m, 2H), 2.79 (dd, J = 12.0, 12.0 Hz, 2H), 2.65 (m, 1H), 1.82 (m, 2H), 1.43 (s, 18H).

(d) *Methyl* 2-(2-(2-(2-((2-fluoro-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (158**)**

[0230] To a mixture of *tert*-butyl 4-(4-((*tert*-butoxycarbonyl)amino)-3-fluorophenyl)piperidine-1-carboxylate (*I57*) (0.376 g, 0.954 mmol) and methyl 2-(2-(2-(2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*I15*) (0.268 g, 0.667 mmol) in 2,2,2-trifluoroethanol (15 mL) was added TFA (0.8 mL). The resulting mixture was heated under microwave irradiation at 100 °C for 20 minutes then 15 minutes then concentrated under reduced pressure to give a crude sample of the title compound (*I58*) (0.344 g) which was used without purification.

(e) *tert*-Butyl 4-(3-fluoro-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*I60*)

[0231] A mixture of crude methyl 2-(2-(2-(2-((2-fluoro-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetate (*I59*) (0.344 g), Boc₂O (0.214 g, 0.98 mmol) and a catalytic amount of DMAP in DCM (15 mL) was stirred for 20 hours. Triethylamine (0.10 mL, 0.72 mmol) was then added and the mixture stirred for 20 hours at 40 °C. Further portions of triethylamine (1.50 mL, 10.8 mmol) and Boc₂O (0.260 g, 1.19 mmol) were added and stirring continued at 40 °C for 4 hours. The mixture was concentrated under reduced pressure and chromatographed on silica gel (0-75% EtOAc/petroleum benzine 40-60 °C) to give the title compound (*I60*) (0.206 g, 50%) in 79% purity by LCMS, this material was used subsequently without further purification; ¹H NMR (400 MHz, COCl₂) δ 8.57 (s, 1H), 8.28 (dd, J = 8.6, 8.6 Hz, 1H), 7.48 (d, J = 2.4 Hz, 1H), 7.22 (m, 4H), 7.00 (m, 2H), 4.25 (m, 2H), 3.75 (s, 2H), 3.68 (s, 3H), 3.11 (m, 4H), 2.81 (m, 2H), 2.64 (m, 1H), 1.84 (m, 2H), 1.62 (dd, J = 12.6, 4.5 Hz, 2H), 1.49 (s, 9H); LCMS Method C: rt 7.05 min; *m/z* 617.2 [M+H]⁺, 561.1 [(M-*t*-Bu)+H]⁺.

(f) *tert*-Butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-fluorophenyl)piperidine-1-carboxylate (*I61*)

[0232] To a solution of *tert*-butyl 4-(3-fluoro-4-((4-(2-(2-methoxy-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)phenyl)piperidine-1-carboxylate (*I60*) (0.206 g) in THF (5 mL) was added aqueous 2 M LiOH (0.500 mL, 1.00 mmol) and water (0.5 mL) and the resulting solution was stirred for 16 hours at room temperature. Aqueous LiOH (1.5 M; 0.5 mL) was added and the reaction stirred for a further 20 hours at room temperature. The mixture was then heated to 50 °C and methanol (1 mL) added before stirring for 20 hours. The volatiles were evaporated under reduced pressure and the residue azeotroped with toluene twice before addition of THF (4 mL), solid lithium hydroxide (0.021 g, 0.87 mmol), water (1 mL) and methanol (1 mL). The resulting mixture was stirred at room temperature for 5 hours and then heated to 50 °C for 16 hours. The volatiles were evaporated under reduced pressure and the residue diluted with water. The aqueous solution was extracted with EtOAc (2x20 mL) before acidifying to a pH of 3 with aqueous HCl. The resulting precipitate was collected as a pellet by centrifugation then dissolved in methanol and concentrated under reduced pressure.

The residue was azeotroped with toluene then taken up in DMF (6 mL) to which EDCI (0.080 g, 0.42 mmol), HOBT (0.061 g, 0.45 mmol) and DIPEA (0.29 mL, 1.67 mmol) were added. The mixture was stirred at room temperature for 30 minutes under nitrogen before addition of ammonium carbonate (0.129 g, 1.66 mmol). The resulting mixture was stirred at 30 °C for 20 hours before the volatiles were evaporated under reduced pressure. Water (20 mL) was added and the resulting suspension was extracted with EtOAc (3x20 mL). The combined organic extracts were concentrated under reduced pressure and the residue chromatographed on silica gel (0-100% EtOAc/petroleum benzine 40-60 °C then 70-100% EtOAc/petroleum benzine 40-60 °C) to give the title compound (*l61*) (0.049 g, 24%); ¹H NMR (400 MHz, *d*₆-Acetone) δ 8.71 (s, 1H), 8.60 (s, 1H), 7.98 (dd, *J* = 8.4, 8.4 Hz, 1H), 7.29 (m, 1H), 7.17 (m, 5H), 6.66 (s, 1H), 6.22 (s, 1H), 4.21 (m, 1H), 3.63 (s, 2H), 3.16 (m, 4H), 2.09 (m, 2H), 1.87 (m, 3H), 1.59 (qd, *J* = 12.7, 4.4 Hz, 2H), 1.46 (s, 9H); LCMS Method C: rt 6.46 min; *m/z* 602.1 [M+H]⁺, 624.1 [M+Na]⁺, 546.1 [(M-*t*-Bu)+H]⁺, 502.1 [(M-Boc)+H]⁺.

(g) 2-(2-(2-(2-((2-Fluoro-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (*l62*)

[0233] To a solution of *tert*-butyl 4-(4-((4-(2-(2-amino-2-oxoethyl)phenethyl)-5-(trifluoromethyl)pyrimidin-2-yl)amino)-3-fluorophenyl)piperidine-1-carboxylate (*l61*) (0.049 g, 0.081 mmol) in THF (3 mL) was added TFA (0.20 mL, 2.6 mmol). The resulting mixture was then stirred at room temperature for 20 hours then 40 °C for 24 hours. The volatiles were removed by evaporation under reduced and DCM (2 mL) and TFA (1.00 mL, 13.1 mmol) were added. After stirring at 30 °C for 18 hours the mixture was concentrated under reduced pressure and the residue azeotroped twice with toluene (10 mL) to give the title compound (*l62*) (0.037 g, 91%). LCMS Method C: rt 4.84 min; *m/z* 502.1[M+H]⁺,

(h) 2-(2-(2-(2-((2-Fluoro-4-(1-methylpiperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (*13*)

[0234] To a solution of 2-(2-(2-(2-((2-fluoro-4-(piperidin-4-yl)phenyl)amino)-5-(trifluoromethyl)pyrimidin-4-yl)ethyl)phenyl)acetamide (*l62*) (0.037 g, 0.074 mmol) in anhydrous methanol (3 mL) was added 37% aqueous formaldehyde (0.022 mL, 0.30 mmol) and sodium triacetoxyborohydride (0.081 g, 0.38 mmol) under a nitrogen atmosphere. The resulting mixture was stirred at room temperature for 1 hour. The volatiles were evaporated under reduced pressure and the residue taken up saturated aqueous NaHCO₃ solution (25 mL). The resulting mixture was extracted with EtOAc (2x25 mL) then the combined organic layers were washed with brine (25 mL), dried (phase separation cartridge) and evaporated to dryness under reduced pressure. The residue was purified using silica gel column chromatography (0-50% MeOH/EtOAc with 1% 2 M ethanolic ammonia) to give the title compound (*13*) (0.020 g, 52%); ¹H NMR (400 MHz, *d*₆-Acetone + *d*₄-MeOH) δ 8.54 (s, 1H),

7.88 (dd, $J = 8.5$ Hz, 1H), 7.16 (m, 6H), 3.80 (s, 2H), 3.63 (s, 2H), 3.10 (m, 4H), 2.95 (m, 2H), 2.54 (m, 1H), 2.27 (s, 3H), 1.79 (m, 4H). LCMS Method C: rt 4.93 min; m/z 516.1 $[M+H]^+$.

Biological Assays

[0235] The activity of compounds of the invention can be profiled using biochemical and cellular assays.

[0236] Primary potency at FAK can be assessed using an Alpha Screen™ technology biochemical assay.

[0237] The kinetics of this binding may be further studied using a surface plasmon resonance (SPR) technology assay using a Biacore™ S51 sensor to establish K_a , k_d and consequently K_D . When off rates from the protein greatly exceed on rates, as may occur for highly potent compounds, K_D gives an accurate measure of protein-ligand binding affinity.

[0238] The ability of compounds of the invention to inhibit FAK within cells can be assessed with an ELISA -type assay performed using a Meso Scale Discovery SECTOR Imager 6000 instrument. In this assay the ability of compounds of the invention to inhibit phosphorylation of Y397-FAK is determined.

[0239] The effect of compounds of the invention on inhibition of cellular proliferation resulting from non-FAK activity may be assessed using a 2D proliferation assay using a suitable cell line. This gives an indication of off-target activities and potential toxicity arising from them. Therefore, comparing inhibition of phosphorylation of Y397-FAK and 2D proliferation gives a measure of FAK specific mediated effects and also of potential toxicity resulting from off-target activity.

[0240] Primary potency at VEGFR3 can be assessed using an Alpha Screen™ technology biochemical assay.

FAK biochemical Alpha Screen™ assay

[0241] A biotin labeled peptide is used as substrate (amino acid sequence: Biotin-Glu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-Glu-Ala-Tyr-Gly-Trp-Met-Asp-Phe-NH₂). FAK enzyme was expressed in insect cells as catalytic domain (amino acids 411-686) N-terminally tagged with six histidine amino acids and a Tobacco Etch Virus (TeV) cleavage sequence. After lysing the cells by sonication, the kinase was purified by Ni-Immobilised Metal Affinity Chromatography chromatography, TeV cleavage leaving a N-terminal glycine, and gel filtration. The 15 μ l assay reactions are run in Greiner brand white 384-well low volume plates. All reactions contained 10 mM HEPES pH 7.4, 25 mM NaCl, 10 mM MgCl₂, 0.01 % (v/v) Tween-20, 50 μ M Na₃VO₄,

0.01% (w/v) albumin from chicken egg white, 111 nM peptide substrate, 80 μ M ATP, and 4 ng/reaction FAK enzyme, with the enzyme being omitted from negative control reactions. Compounds were added in a volume of 100 nL from dilution series made up in DMSO, positive and negative control reactions receiving the same volume DMSO without compounds. The plates were sealed with adhesive seals and incubated for 90 minutes at 30°C. The reactions were stopped with the detection reagents added at the same time. Product formation was quantified as amplified luminescence between PerkinElmer AlphaScreen™ beads, using Streptavidin-coated donor and anti-phosphotyrosine (P-Tyr-100) acceptor beads. To each reaction, 5 μ L containing 10 mM HEPES pH 7.4, 25 mM NaCl, 100 mM EDTA, 0.01 % (v/v) Tween-20, and 6.25 μ g/mL of each bead type were added. Plates were incubated for 6 hours before being read on a PerkinElmer EnVision™ plate reader in HTS Alphascreen™ mode. IC₅₀ values were obtained by calculating percent inhibition (%) for each reaction relative to controls on the same plate ($\%I = (I - CN) / (CP - CN)$ where CN/ CP are the averages of the negative/ positive reactions, respectively), then fitting the %I data vs. compound concentration [I] to $\%I = (A + ((B - A) / (1 + ((C / [I])^D))))$ where A is the lower asymptote, B is the upper asymptote, C is the IC₅₀ value, and D is the slope factor.

Results

[0242]

Compound	IC ₅₀ (nM)
1	1.9
2	7.1
3	2.2
4	3.6
5	7.2
6	2.1
7	9.4
8	5.2
9	3.6
10	2.7
11	2.2
12	3.6
13	5.6

FAK Biacore™ SPR assay

[0243] Binding parameters of compounds were determined using a Biacore™ S51 sensor. An anti-GST antibody was immobilized onto a CM5 chip by primary amine-coupling in accordance with the manufacturer's recommendations.

[0244] In running buffer (0.01 M HEPES pH 7.4, 0.15 M NaCl, 0.005% Surfactant P20, 10 mM MgCl₂, and 1% DMSO) N-terminally GST-fused purified FAK enzyme was captured on both spot 1 and 2. Spot 1 was subsequently blocked by loading with 30 nM PF-562,271 at the beginning of each cycle. Concentration series' of the test compounds were injected over the spots at 25°C. The specific binding was calculated as difference between spot 2 and 1 signals followed by solvent correction. Fitting to a one site binding model yielded the kinetic rate constants k_d and k_a and the equilibrium binding constant $K_D = k_d/k_a$.

[0245] For compounds with an expected $K_D < 5$ nM N-terminally GST-fused purified FAK enzyme was captured on spot 2 of the anti-GST antibody coated chip only. After the injection cycle of a compound the chip surface was regenerated with 10 mM glycine-HCl, pH2.2 before capturing the enzyme again. The binding sensorgrams were analysed as described before.

Results

[0246]

Compound	K_D (nM)
1	0.29
2	0.74
3	0.21 *
4	1.5
6	2.9
8	0.81
9	1.5
10	0.79
11	1.7
*This result was measured using an extended washing step of 30 minutes.	

P397Y-FAK Inhibition MSD platform cellular biomarker assay

[0247] Compounds of the invention may be tested for *in vitro* activity in the following assay: 96-well plates (cat#MA6000, Meso Scale Discovery) are coated with 30µL/well of mouse monoclonal FAK antibody [63D5] (cat#ab72140, Abcam) pre-diluted in PBS to a concentration

of 1mg/mL. The plates are sealed with adhesive film and incubated for 16 hours at 4°C. The antibody is then flicked out of the plates and 150µL of 3% [w/v] Blocker A (cat#R93AA-1, Meso Scale Discovery) is added. The plates are resealed with adhesive film and incubated at room temperature on a shaker set at medium speed for 2 hours. The plates are then washed three times with a solution containing 50mM Tris-HCl pH 7.5, 0.15M NaCl and 0.02% Tween-20, before cell lysate addition described below.

[0248] Cells are split 1:2 into T150 cell culture flasks 2 days prior to compound treatment. On the day prior to compound treatment, 200µL media containing 20,000 cells is seeded into all wells of white, clear-bottom, TC treated, µclear, 96-well microtitre plates (cat#655098, Greiner Bio-One), and the plates are incubated at 37°C and 5% CO₂ for 36 hours. 1µL/well of compound is then added from dilution series prepared in DMSO. Negative control wells receive the same volume of DMSO without compounds, and positive control wells receive 2µM of a control compound in the same volume of DMSO. Cells are treated for 1 hour at 37°C and 5% CO₂. The media/compounds are then flicked off and 55µL/well of ice-cold complete lysis buffer is added. Complete lysis buffer is prepared by adding 1 tablet PhosSTOP complete phosphatase inhibitor (cat#04906837001, Roche) and 1 tablet Complete, Mini, EDTA-free, protease inhibitor (cat#04693159001, Roche) per 10mL of incomplete lysis buffer (150mM NaCl, 20mM Tris-HCl pH 7.5, 1mM EDTA, 1mM EGTA, 1% Triton-X 100). Plates are incubated on ice for 30 minutes, with 30 seconds high speed plate shaking every 5 minutes. 40µL/well of cell lysate is transferred to the coated, blocked and washed 96-well microtitre plates described above. The 96-well plates are sealed with adhesive film and incubated for 16 hours at 4°C. The plates are then washed three times with a solution containing 50mM Tris-HCl pH 7.5, 0.15M NaCl and 0.02% Tween-20 and tapped dry. 25µL/well of detection solution (1% [w/v] Blocker A (cat#R93AA-1, Meso Scale Discovery) in 50mM Tris-HCl pH 7.5, 0.15M NaCl and 0.02% Tween-20, with 1:600 rabbit polyclonal FAK phospho Y397 antibody (cat#ab39967, Abcam), 1:1000 anti-rabbit sulfo-tag antibody (cat#R32AB-1 Meso Scale Discovery) and 1:40 reconstituted Blocker D-M (cat#D609-0100, Rockland Immunochemicals for Research)) is added, and the plates resealed with adhesive film and incubated for 1 hour at room temperature on a plate shaker set to medium speed. Plates are then washed three times with a solution containing 50mM Tris-HCl pH 7.5, 0.15M NaCl and 0.02% Tween-20 and tapped dry. 150µL/well of Read Buffer T + Surfactant (cat#R92TC-1, Meso Scale Discovery) is then added, and pFAK-397 levels quantified using a Meso Scale Discovery SECTOR Imager 6000 instrument.

[0249] IC₅₀ values are determined by first calculating percent inhibition (%I) for each lysate relative to controls on the same plate ($\%I = (S - CP) / (CN - CP)$) where S is the sample result, CN is the average result of DMSO only treated negative controls, and CP is the average result of 2µM treated positive controls. %I is plotted against compound concentration [I] and the data fitted using the following equation, $\%I = (A + ((B - A) / (1 + ((C / [I])^D))))$, where A is the lower asymptote, B is the upper asymptote, C is the IC₅₀ value, and D is the slope factor.

Results for MDA-231-LNA cells

[0250]

Compound	IC ₅₀ (nM)	% response of control at 2 μ M
1	9	122
2	12	91
3	7	102
4	13	108
6	264	80
9	59	112
10	114	116
11	16	117
12	390	77
13	14	90

2D Cellular proliferation assay

[0251] Cells are split 1:4 into T75 cell culture flasks two days prior to cell seeding. A variety of cancer cell lines can be utilized in this assay.

[0252] On the day of cell seeding 100 μ L/well of media containing 1000-5000 cells are added to 96-well microtitre plates (Cat.#655 180, greiner bio-one) except wells G12 and H12 to which 100 μ L of media is added. In a second plate, a single row of cells is seeded at the same concentration. This second plate is known as the t=0 plate and is used to calculate the relative cell number prior to addition of test agent. The plates containing cells are incubated for 24 hours at 37°C/ 5%CO₂. 0.5 μ L/well of compound is then added from dilution series prepared in DMSO. A compound with known potency is included for each set of plates in order to assess assay performance. Negative control wells receive the same volume of DMSO without compounds. Background signal is determined from wells containing media alone. The t=0 plate is read using addition of a resazurin-based reagent (see below) on the day that other plates have compound added to them. Plates containing cells to which compound has been added are then incubated for 3 days at 37°C and 5% CO₂.

[0253] After 3 days of incubation, cell proliferation is quantified by addition of 20 μ L/well of a resazurin-based reagent with a typical composition as follows: Resazurin, Sigma# R7017-1G, 0.015% w/v; methylene blue, Sigma# MB-1(25g), 0.0025% w/v; potassium hexacyanoferrate (III), Sigma# P8131-100G, 0.033 w/v; potassium hexacyanoferrate (II) trihydrate, Sigma# P9387-100G, 0.042% w/v; in PBS buffer. Plates are incubated with resazurin-based reagent for 1-4 hours (37°C, 5% CO₂) prior to the determination of fluorescence at, or near

(579_{Ex}/584_{Em}).

[0254] Percentage inhibition of proliferation (%) for each treated well relative to controls on the same plate is calculated using the equation $\%I = (S-B) - (T_0-B) / (CN-B) - (T_0-B)$ where S is the sample result B is the background fluorescence, T_0 is the t=0 value and CN is the average result of DMSO only treated negative controls. For IC₅₀ determination, %I is plotted against compound concentration [I] and the data fitted using the following equation, $\%I = (A + ((B-A) / (1 + ((C/[I])^D))))$, where A is the lower asymptote, B is the upper asymptote, C is the IC₅₀ value, and D is the slope factor.

Results for MDA-231-LNA cells

[0255]

Compound	IC ₅₀ (μM)
1	1.43
2	2.77
3	6.62
4	1.91
5	3.06
6	2.24
7	4.93
8	2.48
9	2.90
10	1.69
11	2.58
12	4.83
13	2.76

VEGFR3 Biochemical assay

[0256] Compounds of the invention may be tested for *in vitro* activity in the following assay: A biotin labeled peptide is used as substrate (amino acid sequence: Biotin-Glu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-Glu-Glu-Ala-Tyr-Gly-Trp-Met-Asp-Phe-NH₂). VEGFR3 cytoplasmic domain (amino acids 798-1298) was purchased as N-terminal GST-fusion protein ("the enzyme"). The 15 μl assay reactions are run in Greiner brand white 384-well low volume plates. All reactions contained 10 mM HEPES pH 7.4, 10 mM MgCl₂, 0.01 % (v/v) Tween-20, 50 μM Na₃VO₄,

0.01% (w/v) albumin from chicken egg white, 1 mM Dithiothreitol, 111 nM peptide substrate, 500 μ M ATP, and 3.8 ng/reaction enzyme, with the enzyme being omitted from negative control reactions. Compounds were added in a volume of 100 nl from dilution series prepared in DMSO, positive and negative control reactions receiving the same volume DMSO without compound. The plates were sealed with adhesive seals and incubated for 90 minutes at 30 degree Celsius. The reactions were stopped with the detection reagents added at the same time as follows: Product formation was quantified as amplified luminescence between PerkinElmer AlphaScreen™ beads, using Streptavidin-coated donor and anti-phosphotyrosine (P-Tyr-100) acceptor beads. To each reaction, 5 μ l containing 10 mM HEPES pH 7.4, 25 mM NaCl, 100 mM EDTA, 0.01 % (v/v) Tween-20, and 6.25 μ g/ml of each bead type were added. Plates were incubated for 6 hours before being read on a PerkinElmer EnVision™ plate reader in HTS Alphascreen™ mode. IC₅₀ values were obtained by calculating percent inhibition (%I) for each reaction relative to controls on the same plate ($\%I = (I - CN) / (CP - CN)$ where CN/ CP are the averages of the negative/ positive reactions, respectively), then fitting the %I data vs. compound concentration [I] to $\%I = (A + ((B - A) / (1 + ((C / [I])^D))))$ where A is the lower asymptote, B is the upper asymptote, C is the IC₅₀ value, and D is the slope factor.

[0257] The above assay was also run in a slightly modified form in some cases (indicated below with *). In these cases, VEGFR3 cytoplasmic domain (amino acids 818-1177, lacking 949-1002 of UniProt accession number P35916) was expressed and purified as N-terminal Hexa-His-fusion protein ("the enzyme"), rather than using the N-terminal GST-fusion protein.

Results

[0258]

Compound	IC ₅₀ (nM)
1	275*
2	3685*
3	538
5	5750*
6	10440
7	13260*
8	1176
9	1010
10	>66000
11	59763
12	240*
13	205

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2004080980A \[0011\]](#)
- [WO2005016894A \[0011\]](#)
- [WO562271A \[0018\]](#)
- [WO2004056786A \[0018\]](#)
- [WO2004056807A \[0018\]](#)
- [WO2005023780A \[0018\]](#)
- [WO2007063384A \[0018\]](#)
- [WO573228A \[0019\]](#)
- [WO2008129380A \[0020\]](#)
- [WO2008115369A \[0020\]](#)
- [WO2009105498A \[0020\]](#)
- [US2010113475A \[0020\]](#)
- [WO2009143389A \[0020\]](#)
- [WO2009071535A \[0020\]](#)
- [WO2010055117A \[0020\]](#)
- [WO2010058030A \[0020\]](#)
- [WO2010058032A \[0020\]](#)
- [WO2007140222A \[0020\]](#)
- [WO2009024332A \[0020\]](#)
- [WO0194341A \[0096\]](#)
- [WO0132651A \[0096\]](#)
- [WO0047212A \[0096\]](#)
- [WO9835985A \[0096\]](#)
- [WO0160814A \[0096\]](#)
- [WO9722596A \[0096\]](#)
- [WO9730035A \[0096\]](#)
- [WO9732856A \[0096\]](#)
- [WO9813354A \[0096\]](#)
- [WO9902166A \[0096\]](#)
- [WO0040529A \[0096\]](#)
- [WO0041669A \[0096\]](#)

- [WO0192224A](#) [\[0096\]](#)
- [WO0204434A](#) [\[0096\]](#)
- [WO0208213A](#) [\[0096\]](#)


Non-patent literature cited in the description

- **TOMARSCHLAEPPER** Current Opinion in Cell Biology, 2009, vol. 21, 676-683 [\[0003\]](#)
- **BRUNTONFRAME** Current Opinion in Pharmacology, 2008, vol. 8, 437-432 [\[0005\]](#)
- **CHATZIZACHARIAS et al.** Expert Opinion in Therapeutic Targets, 2007, vol. 11, 101315-1328 [\[0005\]](#)
- **RICHARDSONPARSONS** Nature, 1996, vol. 380, 538-540 [\[0006\]](#)
- **FINCHAM et al.** Oncogene, 1995, vol. 10, 112247-2252 [\[0006\]](#)
- **TSUTSUMI et al.** International Journal of Oncology, 2008, vol. 33, 1215-224 [\[0006\]](#) [\[0007\]](#)
- **ANGELUCCIBOLOGNA** Current Pharmaceutical Design, 2007, vol. 13, 2129-2145 [\[0007\]](#)
- **MITRA et al.** Nature Reviews Molecular Cell Biology, 2005, vol. 6, 56-68 [\[0007\]](#)
- **ILIĆ et al.** Nature, 1995, vol. 377, 539-544 [\[0007\]](#)
- **GILMOREROMER** Molecular Biology of the Cell, 1996, vol. 7, 81209-1224 [\[0007\]](#)
- **CARY et al.** Journal of Cell Science, 1996, vol. 7, 1787-1794 [\[0007\]](#)
- **LIAO et al.** Journal of Experimental and Clinical Cancer Research, 2008, vol. 27, 15- [\[0008\]](#)
- **SIESSERHANKS** Clinical Cancer Research, 2006, vol. 12, 113233-3237 [\[0008\]](#)
- **CHAN et al.** Journal of Chemical Biology, 2009, vol. 185, 2357-370 [\[0008\]](#)
- **FRISCH et al.** Journal of Cell Biology, 1996, vol. 134, 3793-799 [\[0009\]](#)
- **XU et al.** Cell Growth and Differentiation, 1996, vol. 7, 4413-418 [\[0009\]](#)
- **ZOUQ et al.** Journal of Cell Science, 2008, vol. 122, 357-367 [\[0009\]](#)
- **BOUCHARD et al.** Apoptosis, 2008, vol. 13, 531-542 [\[0009\]](#)
- **SHIBUEWEINBERG** PNAS, 2009, vol. 106, 2510290-10295 [\[0010\]](#)
- **YAMAMOTO et al.** Cellular Signaling, 2003, vol. 15, 575-583 [\[0010\]](#)
- **POULAINCO-WORKERS** Gynaecologic oncology, 2006, vol. 101, 507-519 [\[0011\]](#)
- **HALDER et al.** Cancer Res, 2007, vol. 67, 2210976-10983 [\[0011\]](#)
- **MCLEAN et al.** Cancer Research, 2001, vol. 61, 8385-8389 [\[0012\]](#)
- **BENLIMAME et al.** Journal of Cell Biology, 2005, vol. 171, 505-516 [\[0012\]](#)
- **MITRA et al.** Oncogene, 2006, vol. 25, 4429-4440 [\[0012\]](#)
- **MITRA et al.** Oncogene, 2006, vol. 25, 5969-5984 [\[0012\]](#)
- **LAHLOU et al.** PNAS USA, 2007, vol. 104, 5120302-20307 [\[0012\]](#)
- **HEER** European Journal of Surgical Oncology, 2008, vol. 34, 111253-1261 [\[0013\]](#)
- **TREMBLAY, L.W. HAUCK et al.** International Journal of Cancer, 1996, vol. 68, 2164-171 [\[0013\]](#)

- **WATERMANN et al.** British Journal of Cancer, 2005, vol. 93, 6694-698 [0013]
- **HESS et al.** Cancer Research, 2005, vol. 65, 219851-60 [0013]
- **ROBERTS et al.** Cancer Res, 2008, vol. 68, 61935-1944 [0018]
- **SLACK-DAVIS et al.** J. Biol. Chem., 2007, vol. 282, 2014845-14852 [0019]
- **BERGE et al.** J. Pharm. Sci., 1977, vol. 66, 1-19 [0033]
- **WALKER et al.** Biorg. Med. Chem. Letts., 2008, vol. 18, 6071-6077 [0039]
- J. Med. Chem., 2004, vol. 47, 6658-6661 [0096]
- Cancer research, 2003, vol. 63, 2375-81 [0096]
- **STERN et al.** Critical reviews in oncology/haematology, 2005, vol. 54, 11-29 [0096]
- Remington's Pharmaceutical Sciences Mack Publishing Company 19900000 [0102]
- **PENG, C. et al.** Adv. Synth. Catal., 2008, vol. 350, 2359-2364 [0142]


Patentkrav

1. Forbindelse med formel (I):



5 eller en ion-, salt- eller solvatform deraf, hvor:

R¹ er udvalgt blandt: H og



hvor:


R^{N1} er udvalgt blandt H, C₁₋₃-alkyl og C(=O)Me;

10 R^{N2} er udvalgt blandt H, C₁₋₃-alkyl og C(=O)Me;

R^{N3} er udvalgt blandt H, C₁₋₃-alkyl og C(=O)Me;

R^{N4} er udvalgt blandt H og CH₃;

R² er udvalgt blandt H og



15 hvor:

R^{N5} er udvalgt blandt H, C₁₋₃-alkyl og C(=O)Me;


R^{N6} er udvalgt blandt H, C₁₋₃-alkyl og C(=O)Me;

og hvor kun ét af R¹ og R² er H;

20 R³ er udvalgt blandt O-C₁₋₂-alkyl, C₁₋₂-alkyl, halogen, cyano, hvor C₁₋₂-alkyl-gruppen kan være substitueret med en eller flere fluorgrupper;

R⁴ er udvalgt blandt CF₃, halogen, CF₂H og CN; og

R⁵ er udvalgt blandt grupper med følgende formler:



hvor:

25 R⁶ er udvalgt blandt H, (CHR^{C1})_{n1}C(O)N(R^{N7})Z¹ og (CH₂)_{n2}C(O)OZ²; hvor:

n_1 er 1;

R^{C1} er H eller Me;

R^{N7} er H eller CH_3 ;

Z^1 er H, CH_3 eller OCH_3 ;

5 n_2 er 1; og

Z^2 er CH_3 ;

og hvor kun ét af R^{N7} og Z^1 kan være CH_3 ,

R^7 er udvalgt blandt H og $(CH_2)_{m1}C(O)N(R^{M1})Y^1$, hvor:

m_1 er 0 eller 1;


10 R^{M1} er H; og

Y^1 er H, Me eller OCH_3 ;


og kun ét af R^6 og R^7 er H; og

R^8 er H, eller, når R^7 er $C(=O)NH_2$, er R^8 udvalgt blandt H og C_{1-2} alkyl.


15 **2. Forbindelse ifølge krav 1, hvor R^2 er H, og R^1 er:**




hvor R^{N1} er $C(=O)Me$; eller



20 hvor R^{N1} er H, methyl eller ethyl; eller




hvor R^{N2} er udvalgt blandt H, methyl og ethyl; eller



hvor R^{N3} er udvalgt blandt H og methyl; eller

(e)




hvor R^{N4} er udvalgt blandt H og methyl.

5


3. Forbindelse ifølge krav 1, hvor R^1 er H, og R^2 er:

(a)



hvor R^{N5} er udvalgt blandt H og methyl; eller

(b)



hvor R^{N6} er udvalgt blandt H og methyl.

10


4. Forbindelse ifølge et hvilket som helst af kravene 1 til 3, hvor R^3 er udvalgt blandt F, Me, Et, OMe og OCF_3 , fortrinsvis OMe.

15

5. Forbindelse ifølge et hvilket som helst af kravene 1 til 4, hvor R^4 er udvalgt blandt CF_3 , Cl og CF_2H , fortrinsvis CF_3 .

6. Forbindelse ifølge et hvilket som helst af kravene 1 til 5, hvor R^5 er en gruppe med følgende formel:

20

(R^{5a})

7. Forbindelse ifølge krav 6, hvor R^7 er H, og R^6 er udvalgt blandt $CH_2C(O)NH_2$, $CH_2C(O)NHCH_3$, $CHCH_3C(O)NH_2$ og $CHCH_3C(O)NHCH_3$.


25

8. Forbindelse ifølge krav 6, hvor R^6 er H, og R^7 er udvalgt blandt $C(O)NH_2$, $C(O)NHCH_3$, $CH_2C(O)NH_2$ og $CH_2C(O)NHCH_3$.

9. Forbindelse ifølge krav 8, hvor R^6 er H, og R^7 er $C(O)NH_2$.

10. Forbindelse ifølge krav 9, hvor R^8 er methyl.

5 11. Forbindelse ifølge et hvilket som helst af kravene 1 til 5, hvor R^5 er en gruppe med følgende formel:



(R^{5b})

10 12. Sammensætning omfattende en forbindelse ifølge et hvilket som helst af kravene 1 til 11 og et farmaceutisk acceptabelt bærerstoff eller fortyndingsmiddel.

13. Forbindelse ifølge et hvilket som helst af kravene 1 til 11 til anvendelse i en fremgangsmåde til terapi.

15 14. Forbindelse ifølge et hvilket som helst af kravene 1 til 11 til anvendelse i fremgangsmåden til behandling af cancer.