



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2361925 A1 2000/08/24

(21) **2 361 925**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2000/02/16
(87) Date publication PCT/PCT Publication Date: 2000/08/24
(85) Entrée phase nationale/National Entry: 2001/08/01
(86) N° demande PCT/PCT Application No.: GB 00/00544
(87) N° publication PCT/PCT Publication No.: WO 00/48615
(30) Priorité/Priority: 1999/02/16 (9903539.6) GB

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 35/74, A61P 31/12

(71) Demandeur/Applicant:
STANFORD ROOK LIMITED, GB


(72) Inventeurs/Inventors:
ROOK, GRAHAM ARTHUR WILLIAM, GB;
STANFORD, JOHN LAWSON, GB

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : TRAITEMENT D'INFECTIONS VIRALES CHRONIQUES A BASE DE M. VACCAE
(54) Title: TREATMENT OF CHRONIC VIRAL INFECTIONS WITH M. VACCAE

(57) **Abrégé/Abstract:**

The present invention provides the use of an M. vaccae preparation for the manufacture of a medicament for use in the treatment of a chronic viral infection, excluding an HIV infection. Chronic viral infections include HPV infection, such as HPV infection associated with cervical dysplasia, herpes virus infection, subacute sclerosing pan-encephalitis and hepatitis virus infection.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 August 2000 (24.08.2000)

PCT

(10) International Publication Number
WO 00/48615 A3

(51) International Patent Classification⁷: **A61K 35/74,**
A61P 31/12

(21) International Application Number: PCT/GB00/00544

(22) International Filing Date: 16 February 2000 (16.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9903539.6 16 February 1999 (16.02.1999) GB

(71) Applicant (*for all designated States except US*): **STANFORD ROOK LIMITED** [GB/GB]; Glen House, 200-208 Tottenham Court Road, London W1P 9LA (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **STANFORD, John, Lawson** [GB/GB]; The Mill House, Claygate, Marden, Kent TN12 9PD (GB). **ROOK, Graham, Arthur, William** [GB/GB]; 29 Belitha Villas, London N1 1PE (GB).

(74) Agents: **BRASNETT, Adrian, H.** et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *With international search report.*

(88) Date of publication of the international search report:
14 December 2000

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TREATMENT OF CHRONIC VIRAL INFECTIONS WITH *M. VACCAE*

(57) Abstract: The present invention provides the use of an *M. vaccae* preparation for the manufacture of a medicament for use in the treatment of a chronic viral infection, excluding an HIV infection. Chronic viral infections include HPV infection, such as HPV infection associated with cervical dysplasia, herpes virus infection, subacute sclerosing pan-encephalitis and hepatitis virus infection.



WO 00/48615 A3

TREATMENT OF CHRONIC VIRAL INFECTIONS WITH M.VACCAEField of the Invention.

The present invention relates to the use of *M.vaccae* in the
5 treatment of viral infections, particularly chronic viral
infections.

Background to the invention.

10 British Specification No. 2156673 (International Patent
Specification W085/03639) describes immunotherapeutic agents
comprising killed cells of *M. vaccae*. These agents are
useful in the immunotherapy of mycobacterial disease,
especially tuberculosis and leprosy. It is stated that use
15 of this immunotherapeutic agent facilitates the removal of
the persisting bacilli responsible for tuberculosis or
leprosy which, as is well known, it is difficult to remove by
chemotherapy alone.

20 International Patent Specification PCT/GB85/00183
(W085/05034) describes compositions for the alleviation of
the symptoms of, and for the treatment or diagnosis of,
arthritic disease which comprise as active ingredient the
whole organism of *M. vaccae*. It is stated that the
25 preparations of *M. vaccae* are useful for the treatment of
various autoimmune diseases and especially arthritic
conditions including rheumatoid arthritis, ankylosing
spondylitis or Reiter's syndrome.

30 Other uses of *M.vaccae* are described in, *inter alia*,
W091/02542, which refers to compositions comprising antigenic
and immunoregulatory material derived from *M. vaccae* as being
generally useful in the treatment of pathological conditions
in which the proportion of agalactosyl IgG (i.e. IgG which
35 lacks terminal galactose from the N-linked oligosaccharides

-2-

on the heavy chains) is increased, W092/08484, which refers to the use of *M. vaccae* for the treatment of uveitis, W094/06466, which provides the use of antigenic and/or immunoregulatory materials derived from *M. vaccae* for the manufacture of a medicament useful in the therapy of AIDS and also in the therapy of HIV-positive asymptomatic patients, and South African patent application 95/2644, which teaches that immunotherapy with *M. vaccae* is expected to be effective against tumours of mesodermal, endodermal and ectodermal origin, including breast and bronchial tumours.

The therapeutic agent conveniently, and therefore preferably, comprises dead cells of *M. vaccae*, most preferably cells which have been killed by autoclaving, although W092/08488 further discloses that killed cells of *M. vaccae* can be used to stimulate and/or modify in a favourable way the immune response to antigens which are not endogenous to *M. vaccae*.

International application PCT/GB98/03346 teaches that cold-shocked *M. vaccae* preparations may be obtained which comprise proteins induced by cold-shock treatment and that such preparations have a variety of uses, including but not limited to the treatment of diseases such as Raynaud's syndrome or hypothermia.

Disclosure of the invention.

We have surprisingly found that treatment of chronic viral infections with *M. vaccae* is beneficial to the course of the infection, in either stabilizing the infection or providing partial or complete clinical improvements.

The action of *M. vaccae* against viral diseases in humans or other animals is surprising in that the action of this bacterial antigen induces a response to cells infected at a

-3-

distant site by a virus.

The present invention thus provides a method of treating a chronic viral infection, excluding an HIV infection, which method comprises administering to a human or animal subject in need of treatment an effective amount of an *M.vaccae* preparation. The action against these non-HIV chronic viruses may be distinguished over the action against HIV in that HIV directly attacks the host immune system which is believed to be modulated by the action of *M.vaccae*.

The invention further provides the use of an *M.vaccae* preparation for the manufacture of a medicament for use in the treatment of a chronic viral infection, excluding an HIV infection.

Detailed Description of the Invention.

Certain viruses can enter specific cells, change the function of the cells to produce viral proteins and express stress proteins, without killing them. Such viruses slowly replicate within the infected host cells and such infection may cause chronic or relapsing disease, and may also produce clinical lesions. Thus by "chronic viral infection", is meant a viral infection of humans or other animals which is able to infect a host and reproduce within the cells of a host over a prolonged period of time - usually weeks, months or years, without proving fatal.

Amongst viruses giving rise to chronic infections and which may be treated in accordance with the present invention are the human papilloma viruses (HPV), Herpes simplex and other herpes viruses, the viruses of hepatitis B and C as well as other hepatitis viruses, and the measles virus, all of which can produce important clinical diseases. Prolonged infection

-4-

may ultimately lead to the induction of disease which may be, e.g. in the case of hepatitis C virus liver cancer, fatal to the patient.

5 Papilloma viruses in man give rise to four different presentations of warts. Sporadic warts, usually on the hands, plantar warts of the foot, multiple warts in which scattered large areas of skin are involved in confluent warty excrescences, often involving the pudenda (venereal warts)
10 and papilloma virus associated dysplasia of the uterine cervix, quite often without other lesions. The first two of these are usually minor, self-resolving or easily treated by topical measures. The second two are much more difficult to treat and usually persist lifelong to varying degrees.
15 Cervical dysplasia is a well known premalignant condition and is the major reason for the cervical smear programme. There are at least 75 types of HPV that can be distinguished. Solitary lesions are usually associated with a single type, multiple lesions are often found to be due to several
20 different types of virus at the same time, venereal warts and cervical dysplasia are often associated with types 16, 18 or 31 virus. These types, as well as other HPV subtypes, e.g. HPV2a and HPV21 may be treated in accordance with the present invention.

25 The herpes simplex virus HSV-1 causes the common condition known as "cold sores" in which small raised or ulcerated lesions occur several times a year, at the mucocutaneous junction between the buccal or nasal mucosa and the
30 surrounding skin. Frequently the lesions occur when immunity is impaired by acute virus infections. The viruses survive longterm in small nerve endings. The HSV-2 virus can also give rise to genital herpes in which extensive lesions occur around the mucocutaneous junction between the glans penis and
35 prepuce or around the vulva. In both sexes genital herpetic

-5-

lesions tend to be much more extensive and severe than are cold sores. Fluctuating from time to time, genital herpes is much more persistent than are cold sores. Again, the virus becomes established in small nerves from which it continuously reinfects the pudendal regions. The present invention provides for treatment of both HSV-1 and HSV-2 infection, as well as for the treatment of other herpes viruses.

10 Hepatitis B and C viruses are usually, though perhaps not solely, transmitted from person to person with blood products. Both give rise to an acute hepatitis that in some people resolves without further problems and in others becomes chronic. Chronic hepatitis results in slow cirrhosis
15 of the liver that may lead to death due to liver failure or to development of malignant hepatomas. Neither disease is at all easily treated.

Very rarely the measles virus can become established
20 chronically in the brain and give rise to a progressive and fatal condition called subacute sclerosing pan-encephalitis (SSPE), for which there is no known treatment.

Other chronic viral infections which may be treated in
25 accordance with the present invention include Epstein Barr virus (EBV), as well as other viruses such as those which may be associated with tumours, or in the case of animals, various veterinary viral diseases, for example those of domestic pets or farmyard animals important in agriculture.

30 It will be understood that the action of treatment of the infection is moderate, alleviate or remove the symptoms of the infection, or delay or prevent the onset of secondary changes such as malignancies or other morphological changes,
35 by reducing the severity of infection by removal or reduction

-6-

of the viral load or otherwise.

The *Mycobacterium vaccae* preparation may be material which can be or include dead cells of *M. vaccae*. Such cells may be killed, for instance using irradiation, e.g. from ⁶⁰Cobalt at a dose of 2.5 megarads, chemically, or by any other means, although autoclaving is preferred, e.g. at 69kPa for 10 minutes at 115°C-125°C. Autoclaving may yield a more effective preparation than irradiation.

Prior to being killed, *M. vaccae* cells may be grown on a suitable solid medium. A modified Sauton's liquid medium may be preferred (Boyden et al.), solidified with agar, preferably 1.3% agar. After aerobic incubation, generally at 32°C for 10 days, the organisms may be harvested, then weighed and suspended in diluent, ready for administration. Storage, if required before use, may be at 4°C.

Instead of growing the cells on a solid medium, a liquid medium, such as the modified Sauton's medium (Boyden et al.), may be employed, for instance in a fermentor.

During growth, the *M. vaccae* may be treated by a cold-shock or a heat-shock in order to induce the production of cold- or heat-shocked proteins. Suitable heat-shock conditions include a period of growth at an elevated temperature, e.g. 37, 42 or 45°C for a period of time from 10 minutes to several, e.g. 3 or more hours. Cold-shock treatment may be performed for similar periods of time at reduced temperatures, e.g. from 10 to 25°C.

The diluent may be unbuffered saline, pyrogen-free. Preferably, the diluent is borate-buffered, preferably containing a surfactant such as Tween 80®. A suitable borate buffer is: Na₂B₄O₇·10H₂O - 3.63 g, H₃BO₃ - 5.25 g, NaCl - 6.19

-7-

g, Tween 80[®] 0.0005%, distilled water to 1 litre. These diluents are pharmaceutically acceptable.

5 The human results mentioned above have been obtained by administration of *M. vaccae* as the GMP preparation, SRL172, which is available for human use under several investigator IND's from the Federal Drug Administration, and CTX's from the Medicines Control Agency in the UK. GLP acute toxicology has been performed by Huntingdon Research. Phase 1 and Phase 10 2 safety data have been obtained in the USA, and lodged with the FDA. SRL172 is in Phase 3 trials for the immunotherapy of tuberculosis. SRL172 may be preferred for use in the present invention.

15 SRL172 is a *M. vaccae* formulation derived from the strain denoted R877R which was deposited under the Budapest Convention at the National Collection of Type Cultures (NCTC) Central Public Health Laboratory, Colindale Avenue, London NW9 5HT, United Kingdom, on 13 February 1984 under the number 20 NCTC 11659. R877R was originally isolated from mud samples from the Lango district of Central Uganda (Stanford and Paul).

Other *M. vaccae* strains may be used instead of SRL172. An 25 organism can be identified as belonging to *M. vaccae* by biochemical and antigenic criteria (Bonicke et al.).

It is preferred for the present invention that the *M. vaccae* material is administered free or substantially free from non- 30 *M. vaccae* antigenic or immunoregulatory material. In other words the medicament or composition to be administered may include, or may consist essentially of, *M. vaccae* antigenic and/or immunoregulatory material, such as dead cells, an extract or derivative thereof, and a pharmaceutically 35 acceptable diluent.

- 8 -

Administration is preferably in a "therapeutically effective amount", this being sufficient to show benefit to a patient. Such benefit may be at least amelioration of at least one symptom. The actual amount administered, and rate and time-
5 course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, eg decisions on dosage etc, is within the responsibility of general practioners and other medical doctors.

10 A single dosage (where dead cells are to be administered) will generally contain from 10^7 to 10^{10} killed *M. vaccae* microorganisms. Patients may be administered a single dose of 10^8 to 10^9 killed *M. vaccae*, though the dose may be
15 repeated if need be, for instance at intervals from 2 weeks to 6 months.

A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially
20 dependent upon the condition to be treated.

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may include, in addition to active ingredient, a
25 pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend
30 on the route of administration, which may be by any convenient route for those of skill in the art, though is preferably by injection, e.g. cutaneous, subcutaneous or intra-dermal.

35 For injection, the active ingredient will be in the form of a

- 9 -

parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Suitable diluents, which are pharmaceutically acceptable and may be preferred, have been discussed already above.

5

Oral administration may be used, in which case the pharmaceutical composition may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

15

M. vaccae may be administered by aerosol to the airways, using a suitable formulation, e.g. including particles of a size which travels to the appropriate parts of the airways. This may be a dried powder rather than aqueous suspension.

20

Instead of killed cells, material derived from *M. vaccae* may be used, in particular an extract or a synthetic molecule which has the requisite activity.

25 The invention is illustrated by the following non-limiting examples, which report case studies of patients receiving one or more standard doses (10^8 - 10^9 cells) of *M. vaccae*.

Example 1

30 Two individuals receiving immunotherapy with killed *M. vaccae* for other indications had longstanding solitary warts on their hands, presumed to be of viral origin. During treatment, complete disappearance of the warts was observed.

35 Example 2.

-10-

Three patients with extensive multiple warts caused by a variety of HPV subtypes have received series' of injections of killed *M.vaccae*.

5 The first of these was a 15 year old boy with an immune defect following bone marrow reconstitution from a parental source for severe combined immunodeficiency, aged 5 months. He is infected with HPV2a and 21, and had developed severe intractable warts 8 years ago. Over the past year he has
10 received 7 injections of *M.vaccae*, following some of which increases in IL-2 production and CD8 cytotoxic cells occurred. However, these were not maintained. Progress of the clinical lesions is ongoing and being monitored for improvements.

15 The second case was of a woman aged 29 years who first developed a finger wart aged 9, had large-scale involvement of her hands and feet by the age of 15 and developed genital warts by the time she was 19. Cervical cytology, first
20 carried out when she was 20 years old, has continued to show borderline to moderate dyskaryosis ever since, for which she has received several local diathermy and cold coagulation treatments. She has received two courses of 3 doses of *M.vaccae*, the first at monthly intervals between March and
25 May 1998, and the second more recently at 2-week intervals. She is currently being followed-up for signs of clinical benefit.

The third case is a woman aged 54, who has received three
30 monthly doses of *M.vaccae*, and is now being monitored for response.

Example 3.

A woman of 38 years had a history of 6-monthly cervical
35 smears showing dysplasia for the past 20 years, during which

-11-

time she had borne 3 children and received local laser therapy. Following a single injection of killed *M.vaccae* in 1995 her next 2 regular cervical examinations were normal for the first time in the 22 years. Subsequently she has
5 received 4 further annual injections of killed *M.vaccae* and cervical examinations, the last one being on 19 May 1999. Since commencing treatment all smear tests have been clear.

Example 4

10 A woman of 34 years, with a history of development of genital herpes 3 months previously, received two-monthly injections of killed *M.vaccae* and a third three months later. The patient was examined 15 months after the first injection and reported that she had been, and remained, free of symptoms
15 since receiving the treatment. Over this period she was expected to have suffered from 2 or 3 recurrent episodes of the disease, despite continuous treatment with antiviral chemotherapy.

20 Example 5

A 9 year old girl with a history of congenital toxoplasmosis presented a year ago with progressive SSPE following vaccination against measles. With the lack of any alternative treatment, a trial of immunotherapy with killed
25 *M.vaccae* was started in October 1997. Over the first 15 months of treatment she received repeated injections of *M.vaccae* and her disease progressed little, if at all, i.e. there was no significant deterioration in her condition. Over the following 12 months she received twice weekly
30 injections and considerable improvements were observed. The patient regained the ability to sit by herself. Large movements with eyes, head, neck arms, and legs were reported, and she was observed to play purposively. Analysis of the cerebrospinal fluid antibodies revealed a change from
35 polyclonal antibodies against measles virus (associated with

-12-

progressive disease) to oligoclonal (associated with recovery from virus infection).

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

10 Although the foregoing invention has been described in some detail by way of illustration and example for the purposes of clarity and understanding, it will be readily apparent to those of skill in the art that certain changes and modifications may be made thereto without departing from the
15 spirit or scope of the appended claims.

References:

- Bonicke *et al.*, (1964) Zentr albl. Bakteriол. Parasitenkd. Infection skr. Hyg. Abt. 1, Orig., 192: 133
- 20 Boyden *et al.*, (1955) J. Immunol. 75: 15.
- Stanford and Paul, (1973) Ann. Soc. Belge Med. Trop. 53: 141-389.

CLAIMS

1. The use of an *Mycobacterium vaccae* (*M.vaccae*) preparation, said preparation being substantially free of non-*M.vaccae* antigenic material, for the manufacture of a medicament for use in the treatment of a chronic viral infection, excluding an HIV infection.
2. The use of an *M.vaccae* preparation for the manufacture of a medicament for the treatment of a chronic viral infection of a patient, excluding an HIV infection, said preparation enhancing the immune response of the patient to said infection.
3. The use according to claim 1 or 2 wherein said chronic viral infection is HPV infection.
4. The use according to claim 3 wherein said HPV infection is associated with cervical dysplasia.
5. The use according to claim 1 or 2 wherein said chronic viral infection is a herpes virus infection.
6. The use according to claim 5 wherein said herpes virus infection is a herpes simplex virus infection.
7. The use according to claim 1 or 2 wherein said chronic viral infection is a subacute sclerosing pan-encephalitis.
8. The use according to claim 1 or 2 wherein said chronic viral infection is a hepatitis virus infection.
9. A method of treating a chronic viral infection, excluding an HIV infection, which method comprises administering to a human or animal subject in need of treatment an effective amount of an *M.vaccae* preparation.

10. A method according to claim 9 wherein said chronic viral infection is HPV infection.

11. A method according to claim 10 wherein said HPV infection is associated with cervical dysplasia.

12. A method according to claim 9 wherein said chronic viral infection is a herpes virus infection.

13. A method according to claim 12 wherein said herpes virus infection is a herpes simplex virus infection.

14. A method according to claim 9 wherein said chronic viral infection is a subacute sclerosing pan-encephalitis.

15. A method according to claim 9 wherein said chronic viral infection is a hepatitis virus infection.