

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(10) International Publication Number

WO 2018/107058 A1

(43) International Publication Date  
14 June 2018 (14.06.2018)

(51) International Patent Classification:  
*C07K 16/28* (2006.01)      *A61K 39/395* (2006.01)

(74) Agent: SCARR, Rebecca B. et al.; MCNEILL BAUR PLLC, 125 Cambridge Park Drive, Suite 301, Cambridge, Massachusetts 02140 (US).

(21) International Application Number:

PCT/US2017/065366

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:

08 December 2017 (08.12.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/432,503      09 December 2016 (09.12.2016) US


(71) Applicant: ALECTOR LLC [US/US]; 151 Oyster Point Blvd., Suite #300, South San Francisco, California 94080 (US).

(72) Inventors: PINCETIC, Andrew; C/O Alector LLC, 151 Oyster Point Blvd., Suite #300, South San Francisco, California 94080 (US). ROSENTHAL, Arnon; C/O Alector LLC, 151 Oyster Point Blvd., Suite #300, South San Francisco, California 94080 (US). LEE, Seung-Joo; C/O Alector LLC, 151 Oyster Point Blvd., Suite #300, South San Francisco, California 94080 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: ANTI-SIRP-ALPHA ANTIBODIES AND METHODS OF USE THEREOF

FIGURE 3A



(57) Abstract: The invention provides anti-SIRPA antibodies, methods of generating such antibodies, and therapeutic uses and methods employing the antibodies.

WO 2018/107058 A1

[Continued on next page]

**Published:**

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

## ANTI-SIRP-ALPHA ANTIBODIES AND METHODS OF USE THEREOF

### CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/432,503, filed December 09, 2016, which is hereby incorporated by reference in its entirety.

### SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

[0002] The content of the following submission of ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing, file name 099061-1069197\_SL.TXT, 70,619 bytes, created December 7, 2017.

### FIELD OF THE INVENTION

[0003] This invention relates to anti-SIRPA antibodies and therapeutic uses of such antibodies.

### BACKGROUND OF THE INVENTION

[0004] Phagocytic cells, such as macrophages (MΦ) and dendritic cells (DCs), distinguish healthy from abnormal cells through an intricate array of cell surface receptors that modulate cellular activation status, proliferation, and/or effector functions. Many of these receptors recognize diverse ligands that either mark unwanted cells for removal (so-called “eat-me” signals) or protect normal cells from destruction (so called “don’t-eat-me” signals). In recent years, the SIRPa—CD47 axis has emerged as a critical determinant in programmed cell removal by macrophages in various clinical settings ranging from cancer cell survival to successful engraftment of hematopoietic cell transplantation. Therapeutic agents that impact this pathway may meet a relevant medical need to ameliorate disease with particular relevance in many types of human cancers.

[0005] SIRPa (signal regulatory protein- $\alpha$ , SIRPA) belongs to the SIRP family of transmembrane receptors, which are primarily expressed within the myeloid cell lineage (including MΦ, DCs, granulocytes, etc.) and characterized by an extracellular region containing 2 membrane-proximal IgC domains and a distal IgV domain. Unique among this

family, SIRPA contains an intracellular, cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM). Upon receptor cross-linking, tyrosine-phosphorylated ITIM sites recruit and activate SHP phosphatases to negatively regulate cellular functions, such as phagocytosis or inflammatory cytokine release. CD47 serves as the principal ligand for SIRPA, and its broad expression in most cell types, including endothelial/epithelial cells, leukocytes, and erythrocytes, suggests that it mediates a “don’t-eat-me” signal to protect healthy cells from phagocyte-dependent clearance. In support of this view, several studies show that adoptive transfer of red blood cells or leukocytes from CD47-knockout mice into wild-type recipients results in rapid clearance of CD47-deficient cells. Conversely, positional genetic analysis of multiple strains of immune-compromised mice receiving human hematopoietic cells identified the *Sirpa* allele in NOD mice as the causal factor for successful engraftment in xenotransplantation models. Subsequent studies demonstrated that the allelic variant of SIRPA expressed only in NOD mice retained the ability to bind human CD47 expressed on human hematopoietic stem cells, and thus, suppress macrophage-dependent graft rejection.

**[0006]** Regulated expression of SIRPA and CD47 establishes a homeostatic control mechanism to modulate phagocytic cell activity. For example, apoptotic cells downregulate expression of CD47 to facilitate engulfment by resident macrophages while live cells remain unharmed. Likewise, inflammatory stimuli, such as LPS, decrease SIRPA expression in MΦ and DCs to potentiate their activation during inflammation. However, dysregulation of SIRPA and CD47 expression contributes to immune-associated diseases, as seen in cancer. Several tumors significantly augment expression of CD47 relative to non-cancerous cells in order to evade immune surveillance mechanisms that normally eliminate malignant cells. Preclinical studies reveal that genetic knockdown of CD47 in syngeneic tumor models, such as B16F10 melanoma, is sufficient to inhibit tumor growth in immune-competent mice. Similar results have been observed with CD47-knocked down human cancer cell lines transplanted into immune-compromised mice. Alternatively, biologic agents that disrupt SIRPA—CD47 interaction, such as anti-CD47 antibodies, also enhance tumor clearance in mouse models. When combined with commercial anti-tumor antigen antibodies, such as trastuzumab or rituximab, anti-CD47 antibodies facilitate a synergistic increase in the anti-tumor response compared to standard monotherapy. Yet, given the ubiquitous expression of CD47, anti-CD47 antibodies risk severe toxicity burdens due to off-target effects limiting their therapeutic efficacy. Nevertheless, these studies establish a crucial role for the SIRPA-

CD47 pathway in regulating myeloid cells with potential applications in cancer immunotherapy.

#### BRIEF SUMMARY OF ASPECTS OF THE DISCLOSURE

**[0007]** In certain aspects, the present disclosure provides agents that down-regulate SIRPA, *e.g.*, anti-SIRPA antibodies. Such agents can be used for treating, preventing, or reducing risk of a disease or pathology associated with SIRPA expression, activity, or signaling. In some aspects, the disclosure relates to the identification of anti-SIRPA antibodies that are capable of downregulating, *i.e.*, decreasing levels of, SIRPA on human macrophages and dendrocytes, as well as cell lines that express SIRPA. In some aspects, the disclosure relates to anti-SIRPA antibodies that antagonize the immune suppressive SIRPA-CD47 interaction and facilitate phagocytosis of CD47-expressing tumor cells. In a further aspect, the present disclosure provides unique SIRPA-specific antibodies that disrupt CD47 binding through non-competitive inhibition.

**[0008]** Thus, in one aspect, the disclosure relates to a SIRPA antibody that selectively binds SIRPA and down-regulates SIRPA expressed on the cell surface. In some embodiments, the anti-SIRPA antibody decreases cell surface levels of SIRPA, decreases intracellular levels of SIRPA, decreases total levels of SIRPA, or any combination thereof. In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody induces SIRPA degradation, SIRPA cleavage, SIRPA internalization, SIRPA shedding, downregulation of SIRPA expression, or any combination thereof. In some embodiments, which may be combined with any of the preceding embodiments, the antibody decreases cellular levels of SIRPA *in vivo*. In some embodiments that may be combined with any of the preceding embodiments, the anti-SIRPA antibody inhibits cell surface clustering of SIRPA. In further embodiments that may be combined with any of the preceding embodiments, the anti-SIRPA antibody inhibits one or more SIRPA activities; or counteracts, one or more SIRPA activities, which may be selected from the group consisting of: (a) SIRPA binding to one or more SIRPA ligands, optionally wherein the one or more SIRPA ligands are selected from the group consisting of CD47, surfactant protein A and D and any combination thereof; (b) decreasing proliferation of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes,

neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia; (c) inhibiting migration of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia; (d) inhibiting one or more functions of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia; (e) inhibition of one or more types of clearance selected from the group consisting of apoptotic neuron clearance, nerve tissue debris clearance, dysfunctional synapse clearance, non-nerve tissue debris clearance, bacteria clearance, other foreign body clearance, disease-causing protein clearance, disease-causing peptide clearance, and tumor cell clearance; optionally wherein the disease-causing protein is selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides and the tumor cell is from a cancer selected from the group consisting of bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal cell cancer, renal pelvis cancer, leukemia, lung cancer, melanoma, non-Hodgkin's lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, fibrosarcoma, and thyroid cancer; (f) inhibition of tumor cell killing by one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells; (g) inhibiting anti-

tumor cell proliferation activity of one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells; (h) modulated expression of one or more inflammatory receptors, optionally wherein the one or more inflammatory receptors comprise CD86 and the one or more inflammatory receptors are expressed on one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells; (i) promoting or rescuing functionality of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid-derived suppressor cells, tumor-associated macrophages, tumor-associated neutrophils, tumor-associated NK cells, and regulatory T cells; (j) increasing infiltration of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid-derived suppressor cells, tumor-associated macrophages, tumor-associated neutrophils, tumor-associated NK cells, non-tumorigenic CD45+CD14+ myeloid cells, and regulatory T cells into tumors; (k) increasing the number of tumor-promoting myeloid/granulocytic immune-suppressive cells and/or non-tumorigenic CD45+CD14+ myeloid cells in a tumor, in peripheral blood, or other lymphoid organ; (l) enhancing tumor-promoting activity of myeloid-derived suppressor cells and/or non-tumorigenic CD45+CD14+ myeloid cells; (m) enhancing survival of non-tumorigenic myeloid-derived suppressor cells and/or non-tumorigenic CD45+CD14+ myeloid cells; (n) decreasing activation of tumor-specific T lymphocytes with tumor killing potential; (o) decreasing infiltration of tumor-specific NK cells with tumor killing potential; (p) increasing tumor volume; (q) increasing tumor growth rate; and (r) decreasing efficacy of one or more immune-therapies that modulate anti-tumor T cell responses, optionally wherein the one or more immune-therapies are immune-therapies that target one or more target proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, TREM1, TREM2, CD39, CD73, CSF-1 receptor, and any combination thereof, or of one or more cancer vaccines. In some embodiments that may be combined with any of the preceding embodiments, the anti-SIRPA antibody induces one or more of the activities that are selected from the group consisting of: (a) increasing the number of tumor infiltrating CD3+ T cells; (b) decreasing cellular levels of SIRPA in non-tumorigenic CD14+myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally

wherein the non-tumorigenic CD14+ myeloid cells are present in blood; (c) reducing the number of non-tumorigenic CD14+ myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood; (d) reducing PD-L1 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (e) reducing PD-L2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (f) reducing B7-H2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (g) reducing B7-H3 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (h) reducing CD200R levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (i) reducing CD163 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (j) reducing CD206 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (k) decreasing tumor growth rate of solid tumors; (l) reducing tumor volume; (m) increasing efficacy of one or more PD-1 inhibitors; (n) increasing efficacy of one or more checkpoint inhibitor therapies and/or immune-modulating therapies, optionally wherein the one or more checkpoint inhibitor therapies and/or immune-modulating therapies target one or more of CTLA4, the adenosine pathway, PD-L1, PD-L2, OX40, TIM3, LAG3, or any combination thereof; (o) increasing efficacy of one or more chemotherapy agents, optionally wherein the one or more of the chemotherapy agents are gemcitabine, capecitabine, anthracyclines, doxorubicin (Adriamycin®), epirubicin (Ellence®), taxanes, paclitaxel (Taxol®), docetaxel (Taxotere®), 5-fluorouracil (5-FU), cyclophosphamide (Cytoxan®), carboplatin (Paraplatin®), and any combination thereof; (p) increasing proliferation of T cells in the presence of non-tumorigenic myeloid-derived suppressor cells (MDSC); (l) inhibiting differentiation, survival, and/or one or more functions of non-tumorigenic myeloid-derived suppressor cells (MDSC); and (r) killing CD33-expressing immunosuppressor non-tumorigenic myeloid cells and/or non-tumorigenic CD14-expressing cells in solid tumors and associated blood vessels when conjugated to a chemical or radioactive toxin.

**[0009]** In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody inhibits interaction between SIRPA and one or more

SIRPA ligands. In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody decreases cellular levels of SIRPA and inhibits interaction between SIRPA and one or more SIRPA ligands. In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody blocks binding of CD47 to human SIRPA.

**[0010]** In some embodiments, which may be combined with any of the preceding embodiments, the antibody selectively binds human SIRPA and does not substantially block binding of CD47 binding to human SIRPA expressed on cells and further, wherein binding to human SIRPA decreases the level of SIRPA on the cell surface. In some embodiments, the antibody binds to the D1 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D2 domain of SIRPA *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D3 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, such an anti-SIRPA antibody competes with an antibody comprising a  $V_H$  sequence comprising the amino acid sequence of SEQ ID NO:2 and a  $V_L$  sequence comprising the amino acid sequence of SEQ ID NO:3. In some embodiments, such an anti-SIRPA antibody comprises a  $V_H$  region comprising: a CDR3 comprising the amino acid sequence of SEQ ID NO:11, a CDR1 comprising the amino acid sequence of SEQ ID NO:9, or a CDR2 comprising the amino acid sequence of SEQ ID NO:10. In some embodiments, the anti-SIRPA antibody comprises a  $V_H$  region comprising: a) a CDR1 that comprises the amino acid sequence of SEQ ID NO:9, a CDR1 that comprises the amino acid sequence of SEQ ID NO:9 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:9; (b) a CDR2 that comprises the amino acid sequence of SEQ ID NO:10 or a CDR2 that comprises the amino acid sequence of SEQ ID NO:10 with no more than two amino acid substitutions; or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:10; and (c) a CDR3 that comprises the amino acid sequence of SEQ ID NO:11, a CDR3 that comprises the amino acid sequence of SEQ ID NO:11 with no more than two amino acid substitutions; or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:11. In some embodiments, the anti-SIRPA comprises a  $V_H$  region comprising: a CDR1 comprising the amino acid sequence of SEQ ID NO:9 or a CDR1 comprising the amino acid sequence of SEQ ID NO:9 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:10 or a CDR2 comprising the amino acid sequence of SEQ ID NO:10 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ

ID NO:11 or a CDR3 comprising the amino acid sequence of SEQ ID NO:11 with no more than one amino acid substitution. In some embodiments, the anti-SIRPA antibody comprises a VH region that comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:9, a CDR2 comprising the amino acid sequence of SEQ ID NO:10, and a CDR3 comprising the amino acid sequence of SEQ ID NO:11. In some embodiments that may be combined with any of the preceding embodiments, the antibody comprises a  $V_H$  region comprising the amino acid sequence of a  $V_H$  region shown in Figure 14A or comprises a  $V_H$  region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a  $V_H$  region of Figure 14A. In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody comprises a  $V_L$  region that comprises a CDR3 comprising the amino acid sequence of SEQ ID NO:8, a CDR1 comprising the amino acid sequence of SEQ ID NO:6, or a CDR2 comprising the amino acid sequence of SEQ ID NO:7. In some embodiments, the  $V_L$  region comprises: (a) a CDR1 comprising the amino acid sequence of SEQ ID NO:6, a CDR1 comprising the amino acid sequence of SEQ ID NO:6 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:6; (b) a CDR2 comprising the amino acid sequence of SEQ ID NO:7, a CDR2 comprising the amino acid sequence of SEQ ID NO:7 with no more than two amino acid substitutions, or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:7; and (c) a CDR3 comprising the amino acid sequence of SEQ ID NO:8, a CDR3 comprising the amino acid sequence of SEQ ID NO:8 with no more than two amino acid substitutions, or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:8. In some embodiments, the  $V_L$  region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:6 or a CDR1 comprising the amino acid sequence of SEQ ID NO:6 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:7 or a CDR2 comprising the amino acid sequence of SEQ ID NO:7 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ ID NO:8 or a CDR3 comprising the amino acid sequence of SEQ ID NO:8 with no more than one amino acid substitution. In some embodiments, the  $V_L$  region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:6, a CDR2 comprising the amino acid sequence of SEQ ID NO:7, and a CDR3 comprising the amino acid sequence of SEQ ID NO:8. In some embodiments, which may be combined with any of the preceding embodiments, the  $V_L$  region comprises the amino acid sequence of a  $V_L$  region shown in Figure 14B; or comprises

a V<sub>L</sub> region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a V<sub>L</sub> region of Figure 14B. In some embodiments, the antibody comprises an Fc region that decreases the levels of Fc $\gamma$ R expressed on the surface of cells. In some embodiments, the antibody comprises an Fc region that decreases the levels of Fc $\gamma$ RBII on the surface of cells.

**[0011]** In some embodiments, which may be combined with any of the preceding embodiments, the antibody selectively binds human SIRPA, but not murine SIRPA, and does not substantially block binding of CD47 binding to human SIRPA expressed on cells and further, wherein binding to human SIRPA decreases the level of SIRPA on the cell surface. In some embodiments, such an anti-SIRPA antibody competes with an antibody comprising a V<sub>H</sub> sequence comprising the amino acid sequence of SEQ ID NO:2 and a V<sub>L</sub> sequence comprising the amino acid sequence of SEQ ID NO:3. In some embodiments, the antibody binds to the D1 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D2 domain of SIRPA *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D3 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, the anti-SIRPA antibody comprises a V<sub>H</sub> region that comprises a CDR3 comprising the amino acid sequence of SEQ ID NO:17, a CDR1 comprising the amino acid sequence of SEQ ID NO:15, or a CDR2 comprising the amino acid sequence of SEQ ID NO:16. In some embodiments, the V<sub>H</sub> region comprises: a) a CDR1 comprising the amino acid sequence of SEQ ID NO:15, a CDR1 comprising the amino acid sequence of SEQ ID NO:15 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:15; (b) a CDR2 comprising the amino acid sequence of SEQ ID NO:16, a CDR2 comprising the amino acid sequence of SEQ ID NO:16 with no more than two amino acid substitutions, or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:16; and (c) a CDR3 comprising the amino acid sequence of SEQ ID NO:17, a CDR3 comprising the amino acid sequence of SEQ ID NO:17 with no more than two amino acid substitutions, or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:17. In some embodiments, the V<sub>H</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:15 or a CDR1 comprising the amino acid sequence of SEQ ID NO:15 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:16 or a CDR2 comprising the amino acid sequence of SEQ ID NO:16 with no more than one amino acid substitution; and a CDR3 comprising the

amino acid sequence of SEQ ID NO:16 or a CDR3 comprising the amino acid sequence of SEQ ID NO:16 with no more than one amino acid substitution. In some embodiments, the V<sub>H</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:15, a CDR2 comprising the amino acid sequence of SEQ ID NO:16, and a CDR3 comprising the amino acid sequence of SEQ ID NO:17. In some embodiments, which may be combined with any of the preceding embodiments, the antibody comprises a V<sub>H</sub> region that comprises the amino acid sequence of a V<sub>H</sub> region of Figure 14C; or comprises a V<sub>H</sub> region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a V<sub>H</sub> region of Figure 14C. In some embodiments, which may be combined with any of the preceding embodiments, the V<sub>L</sub> region comprises a CDR3 comprising the amino acid sequence of SEQ ID NO:14, a CDR1 comprising the amino acid sequence of SEQ ID NO:12, or a CDR2 comprising the amino acid sequence of SEQ ID NO:13. In some embodiments, the V<sub>L</sub> region comprises: a) a CDR1 comprising the amino acid sequence of SEQ ID NO:12, a CDR1 comprising the amino acid sequence of SEQ ID NO:12 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:12; (b) a CDR2 comprising the amino acid sequence of SEQ ID NO:13, a CDR2 comprising the amino acid sequence of SEQ ID NO:13 with no more than two amino acid substitutions, or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:13; and (c) a CDR3 comprising the amino acid sequence of SEQ ID NO:14, a CDR3 comprising the amino acid sequence of SEQ ID NO:14 with no more than two amino acid substitutions, or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:14. In some embodiments, the V<sub>L</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:12 or a CDR1 comprising the amino acid sequence of SEQ ID NO:12 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:13 or a CDR2 comprising the amino acid sequence of SEQ ID NO:13 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ ID NO:4 or a CDR3 comprising the amino acid sequence of SEQ ID NO:14 with no more than one amino acid substitution. In some embodiments, the V<sub>L</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:6, a CDR2 comprising the amino acid sequence of SEQ ID NO:7, and a CDR3 comprising the amino acid sequence of SEQ ID NO:8. In some embodiments, which may be combined with any of the preceding embodiments, the V<sub>L</sub> region comprises the amino acid sequence of a VL region of Figure 14D; or comprises a V<sub>L</sub> region having at least 90%, at least 91%, at least 92%, at

least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a  $V_L$  region of Figure 14D. In some embodiments, the antibody comprises an Fc region that decreases levels of Fc $\gamma$ R on the surface of cells. In some embodiments, the antibody comprises an Fc region that decreases levels of Fc $\gamma$ RBII on the surface of cells.

**[0012]** In a further aspect, which may be combined with any of the preceding embodiments, an isolated anti-SIRPA of the present disclosure competes with one or more antibodies selected from the group consisting of 3F9, 9C2, 8A9, 8F4, 1E2, 7H9, and 4D8 for binding to SIRPA. In some embodiments, the antibody binds to the D1 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D2 domain of SIRPA *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D3 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, the isolated anti-SIRPA antibody binds to essentially the same epitope as one or more antibodies selected from the group consisting of 3F9, 9C2, 8A9, 8F4, 1E2, 7H9, and 4D8. In some embodiments, the isolated anti-SIRPA antibody comprises a  $V_H$  region and a  $V_L$  region, wherein the  $V_H$  region, the  $V_L$  region, or both, comprise at least one, two, three, four, five, or six CDRs of a monoclonal antibody selected from the group consisting of 3F9, 9C2, 8A9, 8F4, 1E2, 7H9, and 4D8.

**[0013]** In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody is a monoclonal antibody. In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody is a humanized antibody. In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody is an Fab, Fab', Fab'-SH, F(ab')<sub>2</sub>, Fv or scFv fragment; or a multivalent antibody, an antibody is of the IgG class, the IgM class, or the IgA class.

**[0014]** In some embodiments that may be combined with any of the preceding embodiments, the anti-SIRPA antibody is of the IgG class the IgM class, or the IgA class. In some embodiments that may be combined with any of the preceding embodiments, the anti-SIRPA antibody has an IgG1, IgG2, IgG3, or IgG4 isotype. In some embodiments that may be combined with any of the preceding embodiments, the antibody binds an inhibitory Fc receptor. In some embodiments that may be combined with any of the preceding embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (Fc $\gamma$ RIIB). In some embodiments, the antibody decreases the level of Fc $\gamma$ RIIB on the cell surface. In some

embodiments that may be combined with any of the preceding embodiments: (a) the anti-SIRPA antibody has a human or mouse IgG1 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: N297A, D265A, D270A, L234A, L235A, G237A, P238D, L328E, E233D, G237D, H268D, P271G, A330R, C226S, C229S, E233P, L234V, L234F, L235E, P331S, S267E, L328F, A330L, M252Y, S254T, T256E, N297Q, P238S, P238A, A327Q, A327G, P329A, K322A, T394D, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering, or comprises an amino acid deletion in the Fc region at a position corresponding to glycine 236; (b) the anti-SIRPA antibody has an IgG1 isotype and comprises an IgG2 isotype heavy chain constant domain 1(CH1) and hinge region, optionally wherein the IgG2 isotype CH1 and hinge region comprises the amino acid sequence of ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGVHTFPALQSS GLYSLSSVVT VPSSNFGTQT YTCNVDHKPS NTKVDKTVERKCCVECPPCP (SEQ ID NO:34), and optionally wherein the antibody Fc region comprises a S267E amino acid substitution, a L328F amino acid substitution, or both, and/or a N297A or N297Q amino acid substitution, wherein the numbering of the residues is according to EU numbering; (c) the anti-SIRPA antibody has an IgG2 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: P238S, V234A, G237A, H268A, H268Q, V309L, A330S, P331S, C214S, C232S, C233S, S267E, L328F, M252Y, S254T, T256E, H268E, N297A, N297Q, A330L, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering; (d) the anti-SIRPA antibody has a human or mouse IgG4 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: L235A, G237A, S228P, L236E, S267E, E318A, L328F, M252Y, S254T, T256E, E233P, F234V, L234A/F234A, S228P, S241P, L248E, T394D, N297A, N297Q, L235E, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering; or (e) the anti-SIRPA antibody has a hybrid IgG2/4 isotype, and optionally wherein the antibody comprises an amino acid sequence comprising amino acids 118 to 260 of human IgG2 and amino acids 261 to 447 of human IgG4, wherein the numbering of the residues is according to EU or, Kabat numbering. In some embodiments that may be combined with any of the preceding embodiments: (a) the anti-SIRPA antibody has a human or mouse IgG1 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: N297A, N297Q, D270A, D265A, L234A, L235A, C226S, C229S, P238S, E233P, L234V,

P238A, A327Q, A327G, P329A, K322A, L234F, L235E, P331S, T394D, A330L, M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering; (b) the anti-SIRPA antibody has an IgG2 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: P238S, V234A, G237A, H268A, H268Q, H268E, V309L, N297A, N297Q, A330S, P331S, C232S, C233S, M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering; or (c) the anti-SIRPA antibody has an IgG4 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: E233P, F234V, L234A/F234A, L235A, G237A, E318A, S228P, L236E, S241P, L248E, T394D, M252Y, S254T, T256E, N297A, N297Q, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering. In some embodiments that may be combined with any of the preceding embodiments: (a) the Fc region further comprises one or more additional amino acid substitutions at a position selected from the group consisting of A330L, L234F; L235E, P331S, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering; (b) the Fc region further comprises one or more additional amino acid substitutions at a position selected from the group consisting of M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU or Kabat numbering; or (c) the Fc region further comprises a S228P amino acid substitution according to EU or Kabat numbering. In some embodiments that may be combined with any of the preceding embodiments, the antibody has an IgG4 isotype. In some embodiments that may be combined with any of the preceding embodiments, the anti-SIRPA antibody comprises an S228P amino acid substitution at residue position 228, an F234A amino acid substitution at residue position 234, and an L235A amino acid substitution at residue position 235, wherein the numbering of the residue position is according to EU or Kabat numbering.

**[0015]** In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody is a bispecific antibody. In some embodiments, the anti-SIRPA antibody recognizes a first and a second antigen, wherein the first antigen is SIRPA and the second antigen is: (a) an antigen facilitating transport across the blood-brain-barrier; (b) an antigen facilitating transport across the blood-brain-barrier selected from the group consisting of transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and

2), diphtheria toxin receptor, CRM197, a llama single domain antibody, TMEM 30(A), a protein transduction domain, TAT, Syn-B, penetratin, a poly-arginine peptide, an angiopep peptide, and ANG1005; (c) a disease-causing agent selected from the group consisting of disease-causing peptides or proteins or, disease-causing nucleic acids, wherein the disease-causing nucleic acids are antisense GGCCCC (G2C4) repeat-expansion RNA, the disease-causing proteins are selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides; and (d) ligands and/or proteins expressed on immune cells, wherein the ligands and/or proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, and phosphatidylserine; and a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells.

**[0016]** In some embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody is a conjugated antibody. For example, the anti-SIRPA antibody may be conjugated to a detectable marker, a toxin, or a therapeutic agent. In some embodiments, the anti-SIRPA antibody is conjugated to a toxin selected from the group consisting of ricin, ricin A chain, doxorubicin, daunorubicin, a maytansinoid, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, *Saponaria officinalis* inhibitor, glucocorticoid, auristatin, auromycin, yttrium, bismuth, combrestatin, duocarmycins, dolastatin, cc1065, and a cisplatin.

**[0017]** In further embodiments, which may be combined with any of the preceding embodiments, the anti-SIRPA antibody is used in combination with one or more antibodies that specifically bind a disease-causing protein selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides, and any combination thereof; or with one or more antibodies that bind an immunomodulatory protein selected from the group consisting of: PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, TREM1, TREM2, CD33, Siglec-5, Siglec-7, Siglec-9, Siglec-11, phosphatidylserine, disease-causing nucleic acids, antisense GGCCCC (G2C4) repeat-expansion RNA, and any combination thereof

**[0018]** In a further aspect, the disclosure provides a method of decreasing the activity, functionality, or survival of regulatory T cells, tumor-imbedded immunosuppressor dendritic cells, tumor-imbedded immunosuppressor macrophages, myeloid-derived suppressor cells, tumor-associated macrophages, acute myeloid leukemia (AML) cells, chronic lymphocytic leukemia (CLL) cell, or chronic myeloid leukemia (CML) cells in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of an agent that binds or interacts with SIRPA, *e.g.*, an antibody of any of the embodiments described above.

**[0019]** In an additional aspect, the disclosure provides a method of inducing or promoting the survival, maturation, functionality, migration, or proliferation of one or more immune cells in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of an agent, *e.g.*, an antibody of any of the embodiments described above, that decreases cellular levels of SIRPA, inhibits interaction between SIRPA

and one or more SIRPA ligands, or both. In some embodiments, the one or more immune cells are selected from the group consisting of dendritic cells, macrophages, neutrophils, NK cells, microglia, T cells, T helper cells, cytotoxic T cells, and any combination thereof.

**[0020]** In another aspect, the disclosure provides a method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any of the embodiments described above to a patient that has a tumor that expresses CD47.

**[0021]** In an additional aspect, the invention provides a method of treating cancer, the method comprising administering a therapeutically effective amount of an agent, *e.g.*, an anti-SIRPA antibody of any of the embodiments described above, that decreases the cellular levels of SIRPA. In some embodiments, the method further comprises administering a therapeutic agent that inhibits PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73. In some embodiments, the therapeutic agent is an antibody that inhibits PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73.

**[0022]** In an additional aspect, the invention provides a method of treating cancer, the method comprising administering a therapeutically effective amount of an agent, *e.g.*, an anti-SIRPA antibody of any of the embodiments described above, that decreases the cellular levels of SIRPA. In some embodiments, the method further comprises administering to the individual at least one antibody that specifically binds to an inhibitory checkpoint molecule, and/or one or more standard or investigational anti-cancer therapies. In some embodiments, the at least one antibody that specifically binds to an inhibitory checkpoint molecule is administered in combination with the anti-SIRPA antibody. In some embodiments, the at least one antibody that specifically binds to an inhibitory checkpoint molecule is selected from the group consisting of an anti-PD-L1 antibody, an anti-CTLA4 antibody, an anti-PD-L2 antibody, an anti-PD-1 antibody, an anti-B7-H3 antibody, an anti-B7-H4 antibody, and anti-HVEM antibody, an anti-B- and T-lymphocyte attenuator (BTLA) antibody, an anti-Killer inhibitory receptor (KIR) antibody, an anti-GAL9 antibody, an anti-TIM-1 antibody, an anti-TIM3 antibody, an anti-TIM-4 antibody, an anti-A2AR antibody, an anti-CD39 antibody, an anti-CD73 antibody, an anti-LAG-3 antibody, an anti-phosphatidylserine

antibody, an anti-CD27 antibody, an anti-CD30 antibody, an anti-TNF $\alpha$  antibody, an anti-CD33 antibody, an anti-Siglec-5 antibody, an anti-Siglec-7 antibody, an anti-Siglec-9 antibody, an anti-Siglec-11 antibody, an antagonistic anti-TREM1 antibody, an antagonistic anti-TREM2 antibody, an anti-TIGIT antibody, an anti-VISTA antibody, an anti-CD2 antibody, an anti-CD5 antibody, and any combination thereof. In some embodiments, which may be combined with any of the preceding embodiments, the one or more standard or investigational anti-cancer therapies are selected from the group consisting of radiotherapy, cytotoxic chemotherapy, targeted therapy, imatinib therapy, trastuzumab therapy, etanercept therapy, adoptive cell transfer (ACT) therapy, chimeric antigen receptor T cell transfer (CAR-T) therapy, vaccine therapy, and cytokine therapy.

**[0023]** In some embodiments, which may be combined with any of the preceding method embodiments, the method further comprises administering to the individual at least one antibody that specifically binds to an inhibitory cytokine. In some embodiments, the at least one antibody that specifically binds to an inhibitory cytokine is administered in combination with an anti-SIRPA antibody of any one of the preceding embodiments. In some embodiments, the at least one antibody that specifically binds to an inhibitory cytokine is selected from the group consisting of an anti-CCL2 antibody, an anti-CSF-1 antibody, an anti-IL-2 antibody, and any combination thereof. In some embodiments that may be combined with any of the preceding embodiments, the method further comprises administering to the individual at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein. In some embodiments, the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is administered in combination with an anti-SIRPA antibody of any of the preceding embodiments. In some embodiments, the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is selected from the group consisting of an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonistic anti-TREM1 antibody, an agonistic anti-TREM2 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, an agonist anti-CD30 antibody, an agonist anti-BTLA antibody, an agonist anti-HVEM antibody, an agonist anti-CD2 antibody, an agonist anti-CD5 antibody, and any combination thereof. In some embodiments, which may be combined with any of the preceding embodiments, the method further comprises administering to the individual at least one stimulatory cytokine. In some embodiments, the stimulatory cytokine

is selected from the group consisting of IFN- $\alpha$ 4, IFN- $\beta$ , IL-1 $\beta$ , TNF- $\alpha$ , IL-6, IL-8, CRP, IL-20 family members, LIF, IFN- $\gamma$ , OSM, CNTF, GM-CSF, IL-11, IL-12, IL-15, IL-17, IL-18, IL-23, CXCL10, IL-33, MCP-1, MIP-1-beta, and any combination thereof.

**[0024]** In a further aspect, the disclosure provides a method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any one of the preceding embodiments to a subject that has cancer cells of a myeloid lineage that expresses SIRPA.

**[0025]** In another aspect, the disclosure provides a method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any one of the preceding embodiments to a subject that has a cancer, wherein the cancer is selected from the group consisting of sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, melanoma, lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, and fibrosarcoma; or wherein the cancer is selected from the group consisting of glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma; diffuse large B-cell lymphoma; lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, non-Hodgkin's lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thymoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma. In some embodiments, the anti-SIRPa antibody is conjugated to a cytotoxic agent and/or induces ADCC.

**[0026]** In some embodiments, the disclosure provides a pharmaceutical composition comprising an anti-SIRPA antibody of any one of the preceding embodiments and a physiologically acceptable carrier. In some embodiments, the disclosure provides an anti-

SIRPA antibody of any one of the preceding embodiments for use in the treatment of cancer; and/or for use in a method of preparing a medicament for the treatment of cancer.

**[0027]** In a further aspect, the disclosure provides a method of preventing, reducing risk, or treating a disease, disorder, or injury selected from the group consisting of dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, taupathy disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, traumatic brain injury, stroke, frontotemporal dementia, spinal cord injury, Huntington's disease, infections, and cancer comprising administering to an individual in need thereof a therapeutically effective amount of an agent that decreases cellular levels of SIRPA, inhibits interaction between SIRPA and one or more SIRPA ligands, or both. In some embodiments, the disease, disorder, or injury is cancer and wherein the agent inhibits one or more SIRPA activities selected from the group consisting of: (a) promoting proliferation, maturation, migration, differentiation, and/or functionality of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid derived suppressor cells, tumor-associated macrophages, tumor-associated suppressor neutrophils, tumor-associated suppressor NK cells, non-tumorigenic CD14+ myeloid cells, and regulatory T cells; (b) enhancing infiltration of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid derived suppressor cells, tumor-associated macrophages, tumor-associated suppressor neutrophils, tumor-associated suppressor NK cells, and regulatory T cells into tumors; (c) increasing number of tumor-promoting myeloid/granulocytic immune-suppressive cells and/or non-tumorigenic CD14+ myeloid cells in a tumor, in peripheral blood, or other lymphoid organ; (d) enhancing tumor-promoting activity of myeloid-derived suppressor cells (MDSC) and/or non-tumorigenic CD14+ myeloid cells; (e) increasing expression of tumor-promoting cytokines in a tumor or in peripheral blood, optionally wherein the tumor-promoting cytokines are TGF-beta or IL-10; (f) increasing tumor infiltration of tumor-promoting FoxP3+ regulatory T lymphocytes; (g) decreasing activation of tumor-specific T lymphocytes with tumor killing potential; (h) decreasing infiltration of tumor-specific T lymphocytes with tumor killing potential; (i) decreasing infiltration of tumor-specific NK cells with tumor killing potential; (j) decreasing the tumor killing potential of NK cells; (k) decreasing infiltration of tumor-specific B lymphocytes with potential to enhance immune response; (l) increasing tumor volume; (m) increasing tumor growth rate; (n) increasing metastasis; (o) increasing rate of tumor

recurrence; (p) decreasing efficacy of one or more immune-therapies that modulate anti-tumor T cell responses, optionally wherein the one or more immune-therapies are immune-therapies that target one or more target proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, and any combination thereof, or one or more cancer vaccines; (q) inhibition of PLC $\gamma$ /PKC/calcium mobilization; and (r) inhibition of PI3K/Akt, Ras/MAPK signaling. In some embodiments, which may be combined with any of the preceding embodiments, the disease, disorder, or injury is cancer, and wherein the agent exhibits one or more SIRPA activities selected from the group consisting of: (a) increasing the number of tumor infiltrating CD3+ T cells; (b) decreasing cellular levels of CD33 in non-tumorigenic CD14+myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood; (c) reducing the number of non-tumorigenic CD14+ myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood; (d) reducing PD-L1 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (e) reducing PD-L2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (f) reducing B7-H2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (g) reducing B7-H3 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (h) reducing CD200R levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (i) reducing CD163 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (j) reducing CD206 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); (k) decreasing tumor growth rate of solid tumors; (l) reducing tumor volume; (m) increasing efficacy of one or more PD-1 inhibitors; (n) increasing efficacy of one or more checkpoint inhibitor therapies and/or immune-modulating therapies, optionally wherein the one or more checkpoint inhibitor therapies and/or immune-modulating therapies target one or more of CTLA4, the adenosine pathway, PD-L1, PD-L2, OX40, TIM3, LAG3, or any combination thereof; (o) increasing efficacy of one or more chemotherapy agents, optionally wherein the

one or more of the chemotherapy agents are gemcitabine, capecitabine, anthracyclines, doxorubicin (Adriamycin®), epirubicin (Ellence®), taxanes, paclitaxel (Taxol®), docetaxel (Taxotere®), 5-fluorouracil (5-FU), cyclophosphamide (Cytoxan®), carboplatin (Paraplatin®), and any combination thereof; (p) increasing proliferation of T cells in the presence of non-tumorigenic myeloid-derived suppressor cells (MDSC); (q) inhibiting differentiation, survival, and/or one or more functions of non-tumorigenic myeloid-derived suppressor cells (MDSC); and (r) killing CD33-expressing immunosuppressor myeloid cells and/or CD14-expressing cells in solid tumors and associated blood vessels when conjugated to a chemical or radioactive toxin. In some embodiments, the cancer expresses SIRPA or one or more SIRPA ligands.

**[0028]** In a further aspect, the disclosure provides a method of treating, preventing or decreasing risk of a disease, disorder, or injury, comprising an agent that down-regulates SIRPA, wherein the disease, disorder, or injury is selected from the group consisting of dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, taupathy disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, traumatic brain injury, stroke, frontotemporal dementia, spinal cord injury, and Huntington's disease. In some embodiments, the agent is an anti-SIRPA antibody, *e.g.*, of any one of the preceding embodiments, that downregulates SIRPA.

**[0029]** In a further aspect, the disclosure provides: a polynucleotide comprising a nucleic acid sequence encoding a V<sub>H</sub> region of an anti-SIRPA antibody of any one of the embodiments described herein; and/or a polynucleotide comprising a nucleic acid sequence encoding a V<sub>L</sub> region of an anti-SIRPA antibody of any one of the embodiments described herein. In a further embodiment, the disclosure provides an expression vector comprising the polynucleotide comprising the nucleic acid sequence encoding the V<sub>H</sub> region or an expression vector comprising the polynucleotide comprising the nucleic acid sequence encoding the V<sub>L</sub> region. In some embodiments, the disclosure provides an expression vector that comprises the polynucleotide encoding the V<sub>H</sub> region and the polynucleotide encoding the V<sub>L</sub> region. In an additional aspect, the disclosure provides a host cell comprising the polynucleotide comprising the nucleic acid sequence encoding the V<sub>H</sub> region or host cell comprising the polynucleotide comprising the nucleic acid sequence encoding the V<sub>L</sub> region. In some embodiments, the disclosure provides a host cell that comprises the polynucleotide encoding the V<sub>H</sub> region and the polynucleotide encoding the V<sub>L</sub> region. In some embodiments, the disclosure provides a host cell comprising an expression vector of any one of the preceding

embodiments. In a further aspect, the disclosure provides a method of producing an anti-SIRPA antibody, the method comprising culturing a host cell of any of the preceding embodiments under conditions in which the antibody is expressed. In some embodiments, the host cell is a mammalian host cell.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0030] FIG. 1A shows an amino acid sequence alignment between the two most common alleles of human SIRPA protein (v1 (SEQ ID NO:1 and v2 (SEQ ID NO:45)) depicting the divergent residues within the ligand-binding domain. Accession numbers are NP542970 and CAA71403, respectively.

[0031] FIG. 1B shows an amino acid sequence alignment between the human SIRPA v1 protein (SEQ ID NO:1) and the human SIRPB1 protein (SEQ ID NO:46), depicting the homology between the two proteins. Accession numbers are NP542970 and O00241, respectively.

[0032] FIG. 2 shows an amino acid sequence alignment between the human SIRPA protein (SEQ ID NO:1) and the mouse SIRPA protein (SEQ ID NO:47), depicting the homology between the two proteins. Accession numbers are NP542970 and Q6P6I8, respectively.

[0033] FIG. 3A shows FACS histograms on the left panel of selected SIRPA antibodies binding to the rodent Chinese hamster ovary cell line (CHO) expressing either human SIRPA (HuSIRPA) or mouse SIRPA (MuSIRPA). Shaded histograms represent the CHO-MuSIRPA cells. Black outlined histograms represent the CHO-HuSIRPA cells. The right panel presents the relative MFI values of SIRPA antibodies binding HuSIRPA compared to MuSIRPA. Results are expressed as fold over background. The background level is set to 1 on y-axis. Antibody mIgG is the isotype negative control. FIG. 3B shows FACS histograms of selected SIRPA antibodies binding to primary human macrophages. Antibody mIgG represents a negative isotype control. Shaded histograms represent the cells stained with anti-mouse IgG secondary antibody only. Black outlined histograms represent the SIRPA positive cell population. FIG. 3C shows surface plasmon resonance sensorgrams of indicated anti-SIRPA antibodies binding to recombinant soluble HuSIRPA protein. An anti-mouse IgG antibody immobilized on a CM5 chip captured anti-SIRPA antibodies and serial dilutions of His-tagged soluble HuSIRPA protein flowed over the antibody. Kd values were determined by curve fitting analysis. FIG. 3D shows binding of increasing concentrations of anti-SIRPA

antibodies to human SIRPA overexpressed on CHO cells. EC50 values were calculated by fitting data to a sigmoidal curve with Graph Pad Prism.

**[0034]** FIG. 4A shows FACS histograms of recombinant soluble human CD47 (HuCD47) binding to CHO-HuSIRPA cells in the presence of either anti-SIRPA antibodies (dashed line histogram) or mouse IgG1 isotype control (solid black outlined histogram). His-tagged HuCD47 was detected with PE-labeled anti-HIS tag secondary antibody. As a negative control (shaded histogram), CHO-HuSIRPA cells were stained with anti-HIS tag PE secondary antibody in the absence of HuCD47. FIG. 4B shows the relative MFI values of HuCD47 binding to CHO-HuSIRPA cells in the presence of indicated anti-SIRPA antibodies or mouse IgG1 isotype control. Results are depicted as fold over background by dividing MFI values of samples treated with HuCD47 and antibodies by the MFI value of cells stained with anti-HIS tag PE in the absence of HuCD47.

**[0035]** FIG. 5A shows induction of human SIRPA-dependent luciferase expression in a cell-based reporter assay. BWZ/NFAT-luciferase reporter cells (BWZ) were engineered to stably express human SIRPA-DAP12 chimera (BWZ-HuSIRPA). Cells were stimulated with increasing concentrations of plate-bound, recombinant HuCD47. Only cells expressing HuSIRPA chimera induced luciferase expression in a dose-dependent manner, as measured by luminescence signal. Results are expressed as fold over background. The background level is set to 1 on y-axis. FIG. 5B shows the ability of CD47-blocking and CD47-non-blocking anti-SIRPA antibodies to affect HuSIRPA-dependent luciferase expression in a cell-based reporter assay. BWZ-HuSIRPA cells were seeded on wells with or without plate-bound CD47 protein. All CD47-blocking antibodies (1B3, 12D6, 1H11, 5F7) potently suppress luminescence signal. Two CD47-non-blocking anti-SIRPA antibodies did not reduce luciferase expression. Results are expressed as fold over background. The background level is set to 1 on y-axis.

**[0036]** FIG. 6A shows induction of human SIRPA-dependent or human SIRPB1-dependent luciferase expression in a cell-based reporter assay. BWZ-HuSIRPA and BWZ-HuSIRPB1 reporter cells were stimulated with plate-bound, full-length anti-SIRPA antibodies or mIgG1 isotype control. CD47-blocking anti-SIRPA antibodies activated both SIRPA-expressing and SIRPB1-expressing reporter cells, whereas the CD47-non-blocking anti-SIRPA antibodies (3F9 and 9C2) specifically activated only BWZ-HuSIRPA cells. FIG. 6B shows surface plasmon resonance sensorgrams of indicated anti-SIRPA antibodies binding to recombinant

soluble HuSIRPA antigen or HuSIRPB1 antigen. An anti-mouse IgG antibody immobilized on a CM5 chip captured anti-SIRPA antibodies and equimolar concentration of antigen flowed over the captured antibody.

[0037] FIG. 7A shows SIRPA receptor down-regulation in primary human macrophages in response to antibody stimulation. Cells were treated with either soluble full-length isotype control or soluble full-length anti-SIRPA antibodies and subsequently stained with a DyLight650-conjugated anti-SIRPA reference antibody (SA56-DyL650) that binds to a distinct epitope bin. FIG. 7B shows SIRPA receptor down-regulation in primary human macrophages treated with CD47-non-blocking antibodies. For comparison, macrophages were also treated with 2 CD47-blocking antibodies (12D6 and 5F7). Results are presented as percent of reference antibody binding by dividing MFI value of samples treated with anti-SIRPA antibodies by the MFI value of samples treated with the isotype control.

[0038] FIG. 8A establishes a live cell phagocytosis assay with macrophages as effector cells and pHrodo-labeled tumor cells as targets. Biotinylated Lens culinaris agglutinin (LCA), a mannose binding lectin, was complexed with avidin-conjugated pHrodo red dye. LCA-pHrodo complexes were then mixed with Raji cells (human B-cell lymphoma line) in order to coat cell surface with pHrodo through LCA binding carbohydrate structures on the cell membrane. Labeled Raji cells (Raji-Red) either alone or opsonized with anti-CD20 antibody were mixed with macrophages at a 2:1 ratio and incubated for 2 hours to allow phagocytosis of cells. Phagocytic activity was measured by counting percent of CD14-APC+/PE+ macrophages by FACS analysis. FIG. 8B shows enhanced phagocytic activity of macrophages treated with CD47-non-blocking anti-SIRPA antibodies. Macrophages were cultured overnight in 2.5%FBS RPMI media with 5  $\mu$ g/mL of 3F9, 9C2, 1B3 (CD47-blocker), or isotype control. Raji-Red cells either alone or opsonized with anti-CD20 antibody were mixed with macrophages at a 2:1 ratio and phagocytic activity was determined as previously described. FIG. 8C shows enhanced phagocytic activity of macrophages treated with CD47-blocking anti-SIRPA antibodies. Macrophages were cultured overnight in 2.5%FBS RPMI media with 5  $\mu$ g/mL of 12D6, 9C5, 1H11, 5F7, 1B3, 3F9 (CD47-non-blocker) or isotype control. Phagocytic activity was measured as described above.

[0039] FIG. 9A shows SIRPA receptor down-regulation in primary human monocytes in response to antibody stimulation. Cells were treated with either soluble full-length isotype control or anti-SIRPA antibody, 3F9, and subsequently stained with a DyLight650-

conjugated anti-SIRPA reference antibody (SA56-DyL650) that binds to a distinct epitope bin. FIG. 9B shows respiratory burst from primary human monocytes isolated from 2 healthy donors (HD). Cells were stimulated with soluble full-length mouse IgG1 isotype control or the anti-SIRPA antibodies 3F9 and 9C2. In all experiments, production of reactive oxygen species (ROS) was monitored by labeling cells with 2  $\mu$ M of the fluorescent indicator, CM-H2DCFDA. FIG. 9C shows IL-8 secretion from primary human monocytes stimulated overnight with CD47-non-blocking antibodies. Supernatants were collected and cytokine concentration determined by standard ELISA protocols as instructed by manufacturer (eBioscience).

**[0040]** FIG. 10A shows the expression of mouse and human SIRPA in peripheral blood monocytes (solid line) and granulocytes (dashed line) by FACS staining in huSIRPA-tg mice. Human SIRPA was detected with anti-hSIRP $\alpha$ / $\beta$ -APC (clone SE5A5, Biolegend); mouse SIRPA was detected with anti-mSIRP $\alpha$ -APC (clone p84, Biolegend). Isotype staining is shown as a shaded histogram. FIG. 10B shows tumor volume measurements of huSIRPA-tg mice implanted subcutaneously with Raji B cell lymphoma cells. Three mice per group received either  $5 \times 10^5$  or  $1 \times 10^6$  Raji cells. Solid tumor formation was determined by caliper measurements twice per week. FIG. 10C shows huSIRPA expression in peripheral blood cells from mice administered 10 mg/kg of either 3F9 (solid line histograms) or isotype control (shaded histograms) antibody. The top panel of FIG. 10C shows detection of huSIRPA with a commercial anti-hSIRP $\alpha$ / $\beta$ -APC (clone SE5A5, Biolegend), an antibody which binds to a different epitope than 3F9. The bottom panel of FIG. 10C shows the detection of huSIRPA with an internally generated anti-hSIRP $\alpha$ -DyLight 650 (clone 9C2), an antibody which binds to the same epitope as 3F9. FIG. 10D shows the downregulation of huSIRPA expression in splenocytes following antibody treatment *in vivo*. The top panel of FIG. 10D shows the gating strategy of single-cell suspensions from mouse spleens stained with anti-mouse F4/80 FITC and anti-mouse CD11b Pacific Blue. The bottom panel of FIG. 10D shows huSIRPA expression from two splenic myeloid populations (F4/80LoCD11bLo and F4/80HiCD11bHi). Solid line histograms represent huSIRPA expression in mice administered isotype control antibody, whereas the dashed line histograms represent huSIRPA expression in mice administered 3F9.

**[0041]** FIG. 11A shows the downregulation of huSIRPA expression in tumor-associated myeloid cells following antibody treatment *in vivo*. The top panel of FIG. 11A shows the gating strategy of single-cell suspensions from tumors stained with anti-mouse F4/80 FITC

and anti-mouse CD11b Pacific Blue. The bottom panel of FIG. 11A shows huSIRPA expression from two splenic myeloid populations (F4/80+ and CD11b+). Solid line histograms represent huSIRPA expression in mice administered isotype control antibody, whereas the dashed line histograms represent huSIRPA expression in mice administered 3F9. FIG. 11B shows the radiance values of Raji-Luciferase lymphoma cells injected subcutaneously into huSIRPA-tg mice. On Day 10, mice were randomized into treatment or control groups based on radiance values and were dosed with i.p. injections of 3F9 or mouse IgG1 antibody at 10 mg/kg every 3-4 days until study termination. Tumor luminescence values post dosing initiation were corrected for with luminescence values at day of randomization and analyzed by linear regression for significance.

**[0042]** FIG. 12A shows the downregulation of huSIRPA expression huCD45+huCD14+ cells harvested from MDA-MB-231 tumor-bearing humanized mice following antibody treatment *in vivo*. The top panel of FIG. 12A shows huSIRPA expression level in peripheral blood huCD45+huCD14+ cells from mice administered i.p. injections of either isotype control, 3F9, or Keytruda (pembrolizumab, Merck). The bottom panel of FIG. 12A shows huSIRPA expression level in tumor infiltrating huCD45+huCD14+ cells from mice administered i.p. injections of either isotype control, 3F9, or Keytruda (pembrolizumab, Merck). FIG. 12B shows the percent of huCD45+ huCD14+ cells present in peripheral blood (FIG. 12B, top panel) or within tumors (FIG. 12B, bottom panel) from mice administered i.p. injections of either isotype control, 3F9, or Keytruda (pembrolizumab, Merck). FIG. 12C provides data showing that the % of human CD45<sup>+</sup> cells in blood of humanized mice is decreased after dosing with SIRPA antibody 3F9. Data are corrected for donor, initial blood parameters (CD45, CD33, CD3), initial animal weight, and initial tumor volume.  
\*\*\*p<0.002 by multiple linear regression (Rlm() function) vs control group (muIgG1).

**[0043]** FIG. 13A plots mean tumor volume in NSG mice engrafted with human immune stem cells from various cord blood donors (donors 5031, 5048, 129). Humanized mice were implanted subcutaneously with the human breast cancer cell line MDA-MB-231 and randomized into treatment or control groups based on tumor volume at Day -1, huCD34+ stem cell donor, body weight before randomization, and huC45+ engraftment rate before randomization. Mice were dosed with i.p. injections of either mouse IgG1 or 3F9 at 40 mg/kg every 4 days or Keytruda at 10 mg/kg every 5 days. Solid gray line represents mean tumor volume of isotype control-treated mice, solid black line represents mean tumor volume of Keytruda-treated mice, and dashed black line represents mean tumor volume of 3F9-

treated mice. FIG. 13B plots mean tumor volume in humanized NSG mice by huCD34+ stem cell donor. The top panels of FIG. 13B show mean tumor volume from treatment and control mice engrafted with stem cells from donors 5031 and 5048. The bottom panels of FIG. 13B show mean tumor volume from treatment and control mice engrafted with stem cells from donor 129. Solid gray line represents mean tumor volume of isotype control-treated mice, solid black line represents mean tumor volume of Keytruda-treated mice, and dashed black line represents mean tumor volume of 3F9-treated mice.

[0044] FIG. 14A lists potential humanized sequences of the heavy chain variable domain of 3F9. Humanized sequence is based on IGHV3-23\*01 acceptor framework andIGHJ4\*01 joining region. FIG. 14A discloses SEQ ID NOS 48-53, respectively, in order of appearance. FIG. 14B lists potential humanized sequences of the light chain variable domain of 3F9. Humanized sequence is based on IGKV3-11\*01 acceptor framework and IGKJ2\*01 joining region. FIG. 14B discloses SEQ ID NOS 54-60, respectively, in order of appearance. FIG. 14C lists potential humanized sequences of the heavy chain variable domain of 9C2. Humanized sequence is based on IGHV1-46\*01 acceptor framework and IGHJ4\*01 joining region. FIG. 14C discloses SEQ ID NOS:61, 49 and 62-67, respectively, in order of appearance. FIG. 14D lists potential humanized sequences of the light chain variable domain of 9C2. Humanized sequence is based on IGKV3-11\*01 acceptor framework and IGKJ2\*01 joining region. FIG. 14D discloses SEQ ID NOS:68, 55 and 69-74, respectively, in order of appearance. CDR sequences noted in bold. CDR definitions are AbM from website [www.bioinf.org.uk/abs/](http://www.bioinf.org.uk/abs/). “b” notes buried sidechain; “p” notes partially buried; “i” notes sidechain at interface between VH and VL domains. Sequence differences between human and murine germlines noted by asterisk (\*). Potential additional mutations in frameworks are noted below sequence. Potential changes in CDR sequences noted below each CDR sequence. These may prevent asparagine (N) deamidation.

[0045] FIG. 15A and 15B show deglycosylation of 3F9 by treatment with EndoS (16A) and that deglycosylation did not have an impact on antigen recognition (16B)

[0046] FIG. 16 provides data illustrating that both glycoforms of 3F9 significantly downregulated surface expression of SIRPA relative to isotype control-treated macrophages, but that the deglycosylated form exhibited partially reduced activity compared to the glycosylated form.

**[0047]** FIG. 17A and 17B provide data illustrating surface expression levels of Fc $\gamma$ RIIA (panel 18A, CD16) and Fc $\gamma$ RIIA/B (panel 18B, CD32A/B) on macrophages treated with control or 3F9 antibody. The antibody used to detect Fc $\gamma$ RII for this analysis does not distinguish the activating receptor (Fc $\gamma$ RIIA) from the inhibitory receptor (Fc $\gamma$ RIIB).

**[0048]** FIG. 18 provides data illustrating cell surface levels of Fc $\gamma$ RIIA (left panel) and Fc $\gamma$ RIIB (right panel) using receptor-specific antibodies on macrophages treated with glycosylated and deglycosylated forms of 3F9.

## DETAILED DESCRIPTION OF THE INVENTION

### Terminology

**[0049]** As used in herein, the singular forms ", "an" and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to "an antibody" optionally includes a combination of two or more such molecules, and the like.

**[0050]** The term "about" as used herein refers to the usual error range for the respective value readily known to the skilled person in this technical field.

**[0051]** The term "antibody" is used herein in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies, such as bispecific antibodies, and antibody fragments so long as they exhibit the desired antigen-binding activity.

**[0052]** The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variant antibodies, *e.g.*, containing naturally occurring mutations or arising during production of a monoclonal antibody preparation, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including but not limited to the hybridoma method, recombinant DNA methods,

phage-display methods, and methods utilizing transgenic animals containing all or part of the human immunoglobulin loci.

**[0053]** An "antibody fragment" refers to a molecule other than an intact antibody that comprises a portion of an intact antibody and that binds the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')<sub>2</sub>; diabodies; linear antibodies; single-chain antibody molecules, such as scFv molecules; and multispecific antibodies formed from antibody fragments.

**[0054]** An "antibody that binds to the same epitope" or that "has the same binding specificity" as a reference antibody refers to an antibody that blocks binding of the reference antibody to its antigen in a competition assay by 50% or more, and conversely, the reference antibody blocks binding of the antibody to its antigen in a competition assay by 50% or more. An antibody that binds to the same epitope may bind to the same epitope as a reference antibody, or may bind to a portion of the epitope. An exemplary competition assay is provided herein.

**[0055]** As used herein, "V-region" refers to an antibody variable region domain comprising the segments of Framework 1, CDR1, Framework 2, CDR2, and Framework 3, including CDR3 and Framework 4, which segments are added to the V-segment as a consequence of rearrangement of the heavy chain and light chain V-region genes during B-cell differentiation.

**[0056]** As used herein, "complementarity-determining region (CDR)" refers to the three hypervariable regions (HVRs) in each chain that interrupt the four "framework" regions established by the light and heavy chain variable regions. The CDRs are the primary contributors to binding to an epitope of an antigen. The CDRs of each chain are referred to as CDR1, CDR2, and CDR3, numbered sequentially starting from the N-terminus, and are also identified by the chain in which the particular CDR is located. Thus, a VH CDR3 is located in the variable domain of the heavy chain of the antibody in which it is found, whereas a VL CDR1 is the CDR1 from the variable domain of the light chain of the antibody in which it is found. The term "CDR" may be used interchangeably with "HVR".

**[0057]** The amino acid sequences of the CDRs and framework regions can be determined using various well known definitions in the art, *e.g.*, Kabat, Chothia, international ImMunoGeneTics database (IMGT), and AbM (see, *e.g.*, Johnson et al., *supra*; Chothia & Lesk, 1987, Canonical structures for the hypervariable regions of immunoglobulins. *J. Mol.*

Biol. 196, 901-917; Chothia C. et al., 1989, Conformations of immunoglobulin hypervariable regions. Nature 342, 877-883; Chothia C. et al., 1992, structural repertoire of the human VH segments J. Mol. Biol. 227, 799-817; Al-Lazikani et al., J.Mol.Biol 1997, 273(4)).

Definitions of antigen combining sites are also described in the following: Ruiz et al., IMGT, the international ImMunoGeneTics database. Nucleic Acids Res., 28, 219–221 (2000); and Lefranc,M.-P. IMGT, the international ImMunoGeneTics database. Nucleic Acids Res. Jan 1;29(1):207-9 (2001); MacCallum et al, Antibody-antigen interactions: Contact analysis and binding site topography, J. Mol. Biol., 262 (5), 732-745 (1996); and Martin et al, Proc. Natl Acad. Sci. USA, 86, 9268–9272 (1989); Martin, et al, Methods Enzymol., 203, 121–153, (1991); Pedersen et al, Immunomethods, 1, 126, (1992); and Rees et al, In Sternberg M.J.E. (ed.), Protein Structure Prediction. Oxford University Press, Oxford, 141–172 1996).

Reference to CDRs as determined by Kabat numbering are based, for example, on Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institute of Health, Bethesda, MD (1991)). Chothia CDRs are determined as defined by Chothia (see, *e.g.*, Chothia and Lesk J. Mol. Biol. 196:901-917 (1987)).

**[0058]** “Epitope” or “antigenic determinant” refers to a site on an antigen to which an antibody binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, *e.g.*, Epitope Mapping Protocols in Methods in Molecular Biology, Vol. 66, Glenn E. Morris, Ed (1996).

**[0059]** An “Fc region” refers to a C-terminal region of an immunoglobulin heavy chain, excluding the first constant region of a native immunoglobulin. The term includes refers to native Fc regions and variant Fc regions. An “Fc region” in the context of native immunoglobulins thus typically refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, and the last three constant region immunoglobulin domains of IgE and IgM, and the flexible hinge N-terminal to these domains. For IgA and IgM, the Fc may include the J chain. For IgG, a native Fc comprises immunoglobulin domains C $\gamma$ 2 and C $\gamma$ 3 and the hinge between C $\gamma$ 1 and C $\gamma$ . It is understood in the art that the boundaries of the

Fc region may vary, however, the human IgG heavy chain Fc region is usually defined to comprise residues C226 or P230 to its carboxyl-terminus, using the numbering according to the EU index as in Kabat et al. (1991, NIH Publication 91-3242, National Technical Information Service, Springfield, Va.). The C-terminal lysine (residue 447 according to the EU or, Kabat numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. Suitable native-sequence Fc regions for use in the antibodies of the present disclosure include human IgG1, IgG2, IgG3 and IgG4. The term "Fc region" includes naturally occurring allelic variants of the Fc region as well as modifications that modulate effector function. Fc regions also include variants that don't result in alterations to biological function. For example, one or more amino acids can be deleted from the N-terminus or C-terminus of the Fc region of an immunoglobulin without substantial loss of biological function.

**[0060]** The term "Fc receptor" or "FcR" describes a receptor that binds to the Fc region of an antibody. An FcR suitable for use in the present invention is typically a native human FcR or variant.

**[0061]** A "native sequence Fc region" comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof.

**[0062]** A "variant Fc region" comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification, preferably one or more amino acid substitution(s). Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, *e.g.* from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% identity with a native sequence Fc region and/or with an Fc region of a parent

polypeptide, and most preferably at least about 90% identity therewith, more preferably at least about 95% identity therewith.

**[0063]** An “antagonist” antibody, or an “inhibitory” antibody is an antibody, such as an anti-SIRPA antibody of the present disclosure, that inhibits or reduces (*e.g.*, decreases) one or more activities or functions of the antigen after the antibody binds the antigen. In some embodiments, an antagonist antibody may block binding of one or more ligands to the antigen. In some embodiments antagonist antibodies or inhibitory antibodies substantially or completely inhibit one or more activities or functions of the antigen; and/or binding of a ligand to the antigen.

**[0064]** The term “equilibrium dissociation constant” abbreviated ( $K_D$ ), refers to the dissociation rate constant ( $k_d$ , time $^{-1}$ ) divided by the association rate constant ( $k_a$ , time $^{-1}$  M $^{-1}$ ). Equilibrium dissociation constants can be measured using any method. Thus, in some embodiments antibodies of the present disclosure have a  $K_D$  of less than about 50 nM, typically less than about 25 nM, or less than 10 nM, *e.g.*, less than about 5 nM or than about 1 nM and often less than about 100 pM as determined by surface plasmon resonance analysis using a biosensor system such as a Biacore® system performed at 37°C. In some embodiments, an antibody of the present disclosure has a  $K_D$  of less than  $5 \times 10^{-5}$  M, less than  $10^{-5}$  M, less than  $5 \times 10^{-6}$  M, less than  $10^{-6}$  M, less than  $5 \times 10^{-7}$  M, less than  $10^{-7}$  M, less than  $5 \times 10^{-8}$  M, less than  $10^{-8}$  M, less than  $5 \times 10^{-9}$  M, less than  $10^{-9}$  M, less than  $5 \times 10^{-10}$  M, less than  $10^{-10}$  M, less than  $5 \times 10^{-11}$  M, less than  $10^{-11}$  M, less than  $5 \times 10^{-12}$  M, less than  $10^{-12}$  M, less than  $5 \times 10^{-13}$  M, less than  $10^{-13}$  M, less than  $5 \times 10^{-14}$  M, less than  $10^{-14}$  M, less than  $5 \times 10^{-15}$  M, or less than  $10^{-15}$  M or lower as measured as a bivalent antibody. In the context of the present invention, an “improved”  $K_D$  refers to a lower  $K_D$ .

**[0065]** The term "bivalent molecule" as used herein refers to a molecule that has two antigen-binding sites. In some embodiments, a bivalent molecule of the present invention is a bivalent antibody or a bivalent fragment thereof. In some embodiments, a bivalent molecule of the present invention is a bivalent antibody. In some embodiments, a bivalent molecule of the present invention is an IgG. In general monoclonal antibodies have a bivalent basic structure. IgG and IgE have only one bivalent unit, while IgA and IgM consist of multiple bivalent units (2 and 5, respectively) and thus have higher valencies. This bivalence increases the avidity of antibodies for antigens.

**[0066]** The terms "bivalent binding" or "bivalently binds to" as used herein refer to the binding of both antigen-binding sites of a bivalent molecule to its antigen. Preferably both antigen-binding sites of a bivalent molecule share the same antigen specificity.

**[0067]** The term "valency" as used herein refers to the number of different binding sites of an antibody for an antigen. A monovalent antibody comprises one binding site for an antigen. A bivalent antibody comprises two binding sites for the same antigen.

**[0068]** The phrase "specifically (or selectively) binds" to an antigen or target or "specifically (or selectively) immunoreactive with," when referring to a protein or peptide, refers to a binding reaction whereby the antibody binds to the antigen or target of interest. In the context of this invention, the antibody typically binds to SIRPA with a  $K_D$  that is at least 100-fold greater than its affinity for other antigens. In some embodiments, the antibody binds to human SIRPA with a  $K_D$  that is at least 100-fold greater than its affinity for other antigens. In some embodiments, the antibody binds to mouse and human SIRPA. As used herein "specific binding" or "selective binding" thus does not necessarily require (although it can include) exclusive binding. An antibody that specifically binds to a target may have an association constant of at least about  $10^3 \text{ M}^{-1}$  or  $10^4 \text{ M}^{-1}$ , sometimes about  $10^5 \text{ M}^{-1}$  or  $10^6 \text{ M}^{-1}$ , in other instances about  $10^6 \text{ M}^{-1}$  or  $10^7 \text{ M}^{-1}$ , about  $10^8 \text{ M}^{-1}$  to  $10^9 \text{ M}^{-1}$ , or about  $10^{10} \text{ M}^{-1}$  to  $10^{11} \text{ M}^{-1}$  or higher. A variety of immunoassay formats can be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See, *e.g.*, Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

**[0069]** An anti-SIRPA antibody of the present invention "downregulates" the level of SIRPA present on the cell surface of cells that express SIRPA. Thus, as used in the present disclosure "down-regulation" refers to the ability of the antibody to decrease the level of SIRPA present on the cell surface of cells that express SIRPA, *e.g.*, human macrophages. An anti-SIRPA antibody of the present invention is considered to down-regulate SIRPA when the level of SIRPA detected on the cell surface is decreased by at least 75% at least 80%, at least 85%, or at least 90% compared to an isotype-matched control antibody.

**[0070]** An "isolated" antibody, such as an anti-SIRPa antibody of the present disclosure, is one that has been identified, separated and/or recovered from a component of its production

environment (*e.g.*, naturally or recombinantly). Preferably, the isolated polypeptide is free of association with all other contaminant components from its production environment. Contaminant components from its production environment, such as those resulting from recombinant transfected cells, are materials that would typically interfere with research, diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified: (1) to greater than 95% by weight of antibody as determined by, for example, the Lowry method, and in some embodiments, to greater than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant T cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, an isolated polypeptide or antibody will be prepared by at least one purification step.

**[0071]** The terms "identical" or percent "identity," in the context of two or more polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues that are the same (*e.g.*, at least 70%, at least 75%, at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher) identity over a specified region, when compared and aligned for maximum correspondence over a comparison window or designated region. Alignment for purposes of determining percent amino acid sequence identity can be performed in various methods, including those using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity the BLAST 2.0 algorithms, which are described in Altschul et al., Nuc. Acids Res. 25:3389-3402 (1977) and Altschul et al., J. Mol. Biol. 215:403-410 (1990). Thus, for purposes of this invention, BLAST 2.0 can be used with the default parameters described to determine percent sequence for nucleic acid sequences or polypeptide sequences.

#### **Overview of certain aspects of the invention.**

**[0072]** The present disclosure relates to agents (*e.g.*, anti-SIRPA antibodies) that SIRPA and/or inhibit interaction between SIRPA and one or more SIRPA ligands; methods of making and using such agents (*e.g.*, anti-SIRPA antibodies); pharmaceutical compositions

containing such agents (e.g., anti-SIRPA antibodies); nucleic acids encoding such agents (e.g., anti-SIRPA antibodies); and host cells containing nucleic acids encoding such agents (e.g., anti-SIRPA antibodies).

**[0073]** An agent of the present disclosure that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands is a molecule having one or more of the following characteristics: (1) inhibits or reduces one or more SIRPA activities; (2) the ability to inhibit or reduce binding of SIRPA to one or more of its ligands; (3) the ability to reduce SIRPA expression (such as at the mRNA level and/or at protein level) in SIRPA-expressing cells; (4) the ability to interact, bind, or recognize a SIRPA protein; (5) the ability to specifically interact with or bind to a SIRPA protein; and (6) the ability to treat, ameliorate, or prevent any aspect of a disease or disorder described or contemplated herein.

**[0074]** Illustrative agents that inhibit the production of SIRPA include, without limitation, compounds that specifically inhibit SIRPA synthesis and/or release, antisense molecules directed to SIRPA, or a short interfering RNA (siRNA) molecule directed to a nucleic acid encoding a SIRPA. Additional exemplary agents that inhibit one or more SIRPA activities include, without limitation, anti-SIRPA antibodies that specifically bind to a SIRPA protein, compounds that specifically inhibit one or more SIRPA activities such as small molecule inhibitors and/or peptide inhibitors, compounds that specifically inhibit SIRPA binding to one or more ligands, a SIRPA structural analog, or an RNA or DNA aptamer that binds SIRPA. In some embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands is an allosteric inhibitor. In some embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands is an orthosteric inhibitor.

**[0075]** In certain embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands is a small molecule inhibitor, including, without limitation, small peptides or peptide-like molecules, soluble peptides, and synthetic non-peptidyl organic or inorganic compounds. A small molecule inhibitor may have a molecular weight of any of about 100 to about 20,000 daltons (Da), about 500 to about 15,000 Da, about 1000 to about 10,000 Da. Methods for making and testing the inhibitory effect a small molecule has on one or more SIRPA activities are well known in the art and such methods can be used to assess the effect of the small molecule inhibitor on SIRPA activity. For example, any of the methods and assays disclosed herein

may be used to screen for small molecule inhibitors that decrease cellular levels of SIRPA and/or inhibit interaction between SIRPA and one or more SIRPA ligands.

**[0076]** In certain embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands comprises at least one antisense molecule capable of blocking or decreasing the expression of a functional SIRPA by targeting nucleic acids encoding a SIRPA. Nucleic acid sequences of SIRPA are known in the art. For example, a human SIRPA can have a nucleic acid sequence as shown in NCBI Accession number NM\_080792 or Y10375.1 a mouse SIRPA can have a nucleic acid sequence as shown in NCBI Accession No. BC062197. Methods are known for the preparation of antisense oligonucleotide molecules and such methods can be used to prepare antisense oligonucleotides that will specifically bind one or more of a SIRPA mRNA without cross-reacting with other polynucleotides. Exemplary sites of targeting include, but are not limited to, the initiation codon, the 5' regulatory regions, the coding sequence, including any conserved consensus regions, and the 3' untranslated region. In certain embodiments, the antisense oligonucleotides are about 10 to about 100 nucleotides in length, about 15 to about 50 nucleotides in length, about 18 to about 25 nucleotides in length, or more. In certain embodiments, the oligonucleotides further comprise chemical modifications to increase nuclease resistance and the like, such as, for example, phosphorothioate linkages and 2'-O-sugar modifications known to those of ordinary skill in the art.

**[0077]** In certain embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands comprises at least one siRNA molecule capable of blocking or decreasing the expression of a functional SIRPA by targeting nucleic acids encoding a SIRPA. Methods for preparation of siRNA molecules are well known in the art and such methods can be used to prepare siRNA molecules that will specifically target a SIRPA mRNA without cross-reacting with other polynucleotides. siRNA molecules may be generated by methods such as by typical solid phase oligonucleotide synthesis, and often will incorporate chemical modifications to increase half-life and/or efficacy of the siRNA agent, and/or to allow for a more robust delivery formulation. Alternatively, siRNA molecules are delivered using a vector encoding an expression cassette for intracellular transcription of siRNA.

**[0078]** In certain embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands is an RNA or DNA

aptamer that binds or physically interacts with a SIRPA, and blocks interactions between a SIRPA and one or more of its ligands. In certain embodiments, the aptamer comprises at least one RNA or DNA aptamer that binds to a mature form of SIRPA.

**[0079]** In certain embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands comprises at least one Siglec-9 structural analog. The term “SIRPA structural analog” refers to compounds that have a similar three dimensional structure as part of that of a SIRPA and which bind to one or more CD3 ligands under physiological conditions *in vitro* or *in vivo*, wherein the binding at least partially inhibits a SIRPA biological activity. Suitable SIRPA structural analogs can be designed and synthesized through molecular modeling of SIRPA binding to a ligand, such as a SIRPA ligand of the present disclosure. The SIRPA structural analogs can be monomers, dimers, or higher order multimers in any desired combination of the same or different structures to obtain improved affinities and biological effects. In some embodiments, the agent binds to or interacts with an amino acid sequence of a SIRPA.

**[0080]** In certain embodiments, an agent that decreases cellular levels of SIRPA and/or inhibits interaction between SIRPA and one or more SIRPA ligands comprises a soluble SIRPA receptor protein, a soluble SIRPA-Fc fusion protein. In certain embodiments, such agents bind one or more SIRPA ligands and thereby prevent the interaction between the SIRPA ligand and SIRPA receptor.

#### *Assays*

**[0081]** Agents that decrease cellular levels of SIRPA and/or inhibit interaction between SIRPA and one or more SIRPA ligands may be identified and/or characterized using methods well known in the art, such as, for example, radiolabeled inhibitor assays, optical assays, protein binding assays, biochemical screening assays, immunoassays, mass shift measurement assays, fluorescence assays, and/or fluorogenic peptide cleavage assays.

#### *Binding assays and other assays*

**[0082]** In certain embodiments, agents that decrease cellular levels of SIRPA and/or inhibit interaction between SIRPA and one or more SIRPA ligands can be identified by techniques well known in the art for detecting the presence of a SIRPA agent candidate’s interaction and/or binding affinity to a SIRPA.

**[0083]** In certain embodiments, agents that interact with SIRPA can be identified using a radiolabeled inhibitor assay. For example, a known amount of a radiolabeled agent candidate may be incubated with a known amount of immobilized SIRPA and a buffer. Subsequently, the immobilized SIRPA may be washed with a buffer and the immobilized SIRPA may be measured for the remaining presence of the radiolabeled SIRPA agent candidate using techniques known in the art, such as, for example, a gamma counter. A measurement indicating the presence of a radiolabeled substance may indicate the radiolabeled agent candidate is capable of interacting with and/or binding to SIRPA.

**[0084]** In certain embodiments, an agent that interacts with a SIRPA may be identified using an optical technique. An exemplary optical technique to detect a SIRPA-interacting agent may include, *e.g.*, attaching SIRPA to a colorimetric resonant grafting surface, thereby shifting the wavelength of reflected light due to changes in the optical path the light must take, and subsequently measuring additional changes in the wavelength of reflected light when a candidate agent is allowed to interact with SIRPA. For example, no change in the measured wavelength of reflected light when an agent is incubated with SIRPA may indicate that the agent candidate is unable to interact with SIRPA. Changes in the measured wavelength of reflected light when an agent candidate is incubated with SIRPA may indicate that the agent candidate is capable of binding and/or interacting with SIRPA.

**[0085]** In certain embodiments, an agent that interacts with SIRPA may be identified using a protein-binding assay. An exemplary protein-binding assay to detect a SIRPA-binding agent may include, *e.g.*, co-immunoprecipitation of SIRPA in the presence of the agent candidate. For example, SIRPA may be incubated with the agent candidate in buffer, and subsequently an immobilized molecule specific to capture SIRPA, such as, for example, an anti-SIRPA antibody, may be used to capture SIRPA in the presence of the agent candidate and bind the SIRPA, potentially with an interacting agent candidate, during wash procedures known in the art. Subsequently, SIRPA, potentially with an interacting agent candidate, can be released and the presence of an agent candidate may be detected, based on the agent candidate characteristics, by techniques, such as, for example, mass spectrometry and/or Western blot.

**[0086]** In certain embodiments, an agent that interacts with a SIRPA may be identified using a biochemical and/or an immunoassay assay well known in the art. An exemplary technique may include, *e.g.*, an assay to quantitatively measure changes in SIRPA

concentration and/or protein half-life using techniques, such as, for example, Western blot, immunostaining, and co-immunoprecipitation. For example, an agent candidate may be incubated with a sample containing a SIRPA, such as a cell expressing SIRPA, and subsequently SIRPA protein quantity and/or cellular levels may be measured at points during a time course study. Changes in protein quantity, cellular levels, and/or protein half-life in comparison to a control treatment may indicate that the SIRPA agent candidate may be capable of altering SIRPA half-life and/or activity.

**[0087]** In certain embodiments, a mass shift measurement assay may be used to identify an agent that interacts with a SIRPA. An exemplary mass shift measurement assay may include, *e.g.*, detecting the presence of a strongly and/or covalently bound SIRPA agent by measuring a change in SIRPA mass when the agent candidate is interacting with SIRPA by using instruments, such as, but not limited to, a mass spectrometer. For example, a mass shift assay may be performed on a whole protein and/or a peptide-based analysis, depending on the nature of the agent candidate interaction. Detection of a mass shift correlating with the addition of said agent candidate to SIRPA may indicate that the agent candidate may be capable of interacting with or otherwise inhibiting a SIRPA. Additionally, an exemplary mass shift measurement assay may include, *e.g.*, detecting the addition of mass to SIRPA correlating with the respective agent candidate mass when the agent candidate is interacting with SIRPA using techniques, such as, for example, surface plasmon resonance. For example, the change in the refractive index of light may be measured and correlated with a change in mass of SIRPA attached to a sensor surface.

**[0088]** In certain embodiments, a chemical cross-linking assay may be used to identify a SIRPA agent that interacts with a SIRPA. For example, an agent candidate may be incubated with a SIRPA, *in vivo* or *in vitro*, with a molecule cross-linker capable of covalently linking an agent candidate interacting with SIRPA to said SIRPA molecule. Subsequently, techniques, such as, but not limited to, mass spectrometry and/or Western blot, may be used to identify an agent candidate that may be capable of interacting with or otherwise inhibiting SIRPA. For example, detection of SIRPA covalently cross-linked with the agent candidate may indicate that the agent candidate may be capable of interacting with or otherwise inhibiting SIRPA.

**[0089]** In certain embodiments, agents that interact with a SIRPA may be identified using a fluorescence assay. For example, a known amount of a fluorescent agent candidate may be

incubated with a known amount of immobilized SIRPA and a buffer. Subsequently, the immobilized SIRPA may be washed with a buffer and the immobilized SIRPA may be measured for the remaining presence of a fluorescent SIRPA agent candidate using techniques known in the art, such as, but not limited to, fluorescence detection. A measurement indicating the presence of a fluorescent substance may indicate the fluorescent agent candidate is capable of interacting with and/or binding to SIRPA.

**[0090]** Assays known in the art and described herein (e.g., Examples 2-11) can be used for identifying and testing biological activities of SIRPA agents of the present disclosure. In some embodiments, assays for testing the ability of SIRPA agents for modulating one or more Siglec-9 activities are provided.

#### **Anti-SIRP-alpha (SIRPA) Antibodies**

*Brief overview of aspects of certain anti-SIRPA antibodies of the present disclosure*

**[0091]** In some embodiments, anti-SIRPA antibodies of the present disclosure have one or more antagonistic activities that are due, at least in part, to the ability of the antibodies to down regulate cellular SIRPA. In some embodiments, an isolated SIRPA antibody of the present disclosure selectively binds SIRPA and down-regulates SIRPA. In some embodiments, the antibody does not block binding of a SIRPA ligand, e.g., CD47, to SIRPA expressed on cells. In alternative embodiments, the antibody blocks binding of a SIRPA ligand, e.g., CD47, to SIRPA. In some embodiments, the antibody is a human antibody, a humanized antibody, a bispecific antibody, a multivalent antibody, or a chimeric antibody. Exemplary descriptions of such antibodies are found throughout the present disclosure. In some embodiments, the antibody is a bispecific antibody recognizing a first antigen and a second antigen.

**[0092]** In some embodiments, anti-SIRPA antibodies of the present disclosure selectively bind to human SIRPA, including human allelic variants, also referred to herein as “polymorphic” variants, but not mouse SIRPA, and do not bind to SIRPB. FIG. 1A shows an amino acid sequence alignment between the two most common alleles of human SIRPA protein (v1 and v2, accession numbers are NP542970 and CAA71403, respectively) depicting the divergent residues within the ligand-binding domain. Thus, in some embodiments, an anti-SIRPA antibody of the present disclosure binds to a linear or conformational epitope that is present in allelic variants of human SIRPA, but not in SIRPB or mouse SIRPA. FIG. 1B shows an amino acid sequence alignment between the human SIRPA v1 protein and the

human SIRPB1 protein, accession numbers are NP542970 and O00241, respectively, depicting the homology between the two proteins. FIG. 2 shows an amino acid sequence alignment between the human SIRPA protein and the mouse SIRPA protein, accession numbers are NP542970 and Q6P6I8, respectivelydepicting the homology between the two proteins. In some embodiments, the antibodies of the present disclosure selectively bind to human and mouse SIRPA and do not bind to SIRPB.

**[0093]** SIRPA is a single-pass type I membrane protein. Within the amino acid sequence of human SIRPA (SEQ ID NO:1), an extracellular domain is located at amino acid residues 31-373; a transmembrane domain is located at amino acid residues 374-394; and an intracellular domain is located at amino acid residues 395-504.

**[0094]** Human SIRPA comprises a single V-set and two C1-sets of Ig super family (IgSF) domains, referred to as the D1 domain, the D2 domain, and the D3 domain, respectively. The D1 domain comprises amino acid residues 32-137 of human SIRPA; the D2 domain comprises amino acid residues 148-247 of human SIRPA; and the D3 domain comprises amino acid residues 254-348 of human SIRPA.

**[0095]** In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the D1 domain of SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the D1 domain of human SIRPA comprising amino acid residues 32-137 of human SIRPA amino acid sequence of SEQ ID NO:1. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to an epitope within the D1 domain of human SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to an epitope within the D1 domain of human SIRPA, wherein the epitope comprises an amino acid sequence selected from the group consisting of amino acid residues 32-137, amino acid residues 32-52, amino acid residues 55-121, amino acid residues 58-73, amino acid residues 68-83, amino acid residues 78-93, amino acid residues 88-103, amino acid residues 98-113, amino acid residues 108-123, and amino acid residues 118-133 of the human SIRPA amino acid sequence of SEQ ID NO:1.

**[0096]** In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the D2 domain of SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the D2 domain of human SIRPA comprising amino acid residues 148-247 of the human SIRPA amino acid sequence of SEQ ID NO:1. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to an epitope within the D2 domain of human

SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to an epitope within the D2 domain of human SIRPA, wherein the epitope comprises an amino acid sequence selected from the group consisting of amino acid residues 148-247, amino acid residues 148-168, amino acid residues 158-173, amino acid residues 168-183, amino acid residues 170-228, amino acid residues 178-193, amino acid residues 188-203, amino acid residues 198-213, amino acid residues 208-223, amino acid residues 218-233, and amino acid residues 228-243 of the human SIRPA amino acid sequence of SEQ ID NO:1.

**[0097]** In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the D3 domain of SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the D3 domain of human SIRPA comprising amino acid residues 254-348 of the human SIRPA amino acid sequence of SEQ ID NO:1. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to an epitope within the D3 domain of human SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to an epitope within the D3 domain of human SIRPA, wherein the epitope comprises an amino acid sequence selected from the group consisting of amino acid residues 254-348, amino acid residues 254-274, amino acid residues 264-279, amino acid residues 274-289, amino acid residues 273-331, amino acid residues 281-315, amino acid residues 281-337, amino acid residues 284-299, amino acid residues 294-309, amino acid residues 304-319, amino acid residues 314-329, amino acid residues 324-339, and amino acid residues 334-348 of the human SIRPA amino acid sequence of SEQ ID NO:1.

**[0098]** In some embodiments, the antibody binds to the D1 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D2 domain of SIRPA *e.g.*, human SIRPA. In some embodiments, the antibody binds to the D3 domain of SIRPA, *e.g.*, human SIRPA. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the same SIRPA epitope or part of the SIRPA epitope bound by an antibody having the CDRs of the antibody designated as 3F9 in Table 2. In some embodiments, an anti-SIRPA antibody of the present disclosure binds to the same SIRPA epitope or part of the SIRPA epitope bound by an antibody having the CDRs of the antibody designated as 9C2 in Table 2, which competes with 3F9 for binding to SIRPA and binds to all of part of the same epitope as 3F9. Accordingly, in some embodiments, an antibody of the present disclosure binds to the same SIRPA epitope or part of the SIRPA epitope bound by an antibody having the CDRs of the antibody designated as 3F9 in Table 2 and binds to the same epitope or part of the SIRPA epitope bound by an antibody having the CDRs of the antibody designated as 9C2 in Table 2.

**[0099]** In some embodiments, an anti-SIRPA antibody of the present disclosure competes with 3F9 and 9C2 for binding to human SIRPA.

**[0100]** In preferred embodiments, an antibody of each of the preceding three paragraphs does not block CD47 binding to SIRPA.

*SIRPA down-regulation*

**[0101]** Certain aspects of the present disclosure relate to anti-SIRPA antibodies that down-regulate, *i.e.*, decrease cellular levels of SIRPA. In some embodiments, the anti-SIRPA antibody decreases cellular levels of SIRPA without inhibiting the interaction (*e.g.*, binding) between SIRPA and one or more SIRPA ligands, *e.g.*, CD47. In some embodiments, the anti-SIRPA antibody decreases cellular levels of SIRPA and inhibits the interaction (*e.g.*, binding) between SIRPA and one or more SIRPA ligands, *e.g.*, CD47.

**[0102]** Cellular levels of SIRPA may refer to, without limitation, cell surface levels of SIRPA, intracellular levels of SIRPA, and total levels of SIRPA. In some embodiments, a decrease in cellular levels of SIRPA comprises decrease in cell surface levels of SIRPA. As used herein, an anti-SIRPA antibody decreases cell surface levels of SIRPA if it induces a decrease of 25% or more in cell surface levels of SIRPA as measured by any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art, for example utilizing flow cytometry, such as fluorescence-activated cell sorting (FACS), to measure cell surface levels of SIRPA. In some embodiments, a decrease in cellular levels of SIRPA comprises a decrease in intracellular levels of SIRPA. As used herein, an anti-SIRPA antibody decreases intracellular levels of Siglec-9 if it induces a decrease of 25% or more in intracellular levels of SIRPA as measured by any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art, for example immunostaining, Western blot analysis, co-immunoprecipitation, and cell cytometry. In some embodiments, a decrease in cellular levels of SIRPA comprises a decrease in total levels of SIRPA. As used herein, an anti-SIRPA antibody decreases total levels of SIRPA if it induces a decrease of 25% or more in total levels of SIRPA as measured by any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art, for example immunostaining, Western blot analysis, co-immunoprecipitation, and cell cytometry. In some embodiments, the anti-SIRPA antibodies induce SIRPA degradation, SIRPA cleavage, SIRPA internalization, SIRPA shedding, downregulation of SIRPA expression, or any combination thereof. In some embodiments, cellular levels of SIRPA are measured on primary cells (*e.g.*, dendritic cells,

bone marrow-derived dendritic cells, monocytes, microglia, and macrophages) or on cell lines utilizing an SIRPA cell assay.

**[0103]** In some embodiments, a downregulating anti-SIRPA antibody has an IC<sub>50</sub> of 200 nM of less, typically 100 nM or less (50% of SIRPA expressed on the cell surface is downregulated), after 4 hours of exposure of human macrophages to the antibody at 37°C. In some embodiments, SIRPA remains down-regulated for at least 24 hours of exposure to an antibody of the present invention. Cells may be analyzed for SIRPA surface expression using any technology, *e.g.*, flow cytometry.

**[0104]** In some embodiments, anti-SIRPA antibodies of the present disclosure decrease cellular levels of SIRPA by at least 25%, at least 26%, at least 27%, at least 28%, at least 29%, at least 30%, at least 31%, at least 32%, at least 33%, at least 34%, at least 35%, at least 36%, at least 37%, at least 38%, at least 39%, at least 40%, at least 41%, at least 42%, at least 43%, at least 44%, at least 45%, at least 46%, at least 47%, at least 48%, at least 49%, at least 50%, at least 51%, at least 52%, at least 53%, at least 54%, at least 55%, at least 56%, at least 57%, at least 58%, at least 59%, at least 60%, at least 61%, at least 62%, at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68%, at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or more as compared to cellular levels of SIRPA in the absence of the anti-SIRPA antibody.

**[0105]** In some embodiments, which may be combined with any of the down-regulation activities summarized in the preceding paragraphs, an anti-SIRPA antibody of the present disclosure inhibits cell surface clustering of SIRPA.

**[0106]** In some embodiments, an anti-SIRPA antibody of the present disclosure down-regulates SIRPA, but does not block binding of a SIRPA ligand, *e.g.*, CD47 to SIRPA. In the context of the present invention, an antibody that does not block binding of CD47 to SIRPA refers to an antibody that does not result in a significant decrease in CD47 binding to SIRPA when the antibody is incubated with CD47 and cells expressing SIRPA. A “significant decrease” in the context of CD47 binding to SIRPA refers to a decrease in binding of 30% or less, typically at least 25%, at least 20%, at least 15%, or at least 10% or less compared to

CD47 binding to SIRPA in the presence of an isotype-matched control antibody that does not bind SIRPA. An illustrative assay for assessing blocking activity is set forth in the examples. For example, cells that express human SIRPA, *e.g.*, human macrophages are cells such as CHO cells that are modified to express human SIRPA, are plated at  $10^5$  cells/well in a 96-well plate, washed, and incubated in 100  $\mu$ l buffer for fluorescent activate cell sorting containing 1.0  $\mu$ g/ml of monoclonal antibody or isotype control. Cells are then washed and incubated in with soluble human CD47 for 30 minutes on ice. Cells are then analyzed for surface-bound CD47.

**[0107]** Alternatively, in some embodiments, an anti-SIRPA antibody of the present disclosure down-regulates SIRPA, but blocks binding of CD47 to SIRPA. An antibody that blocks CD47 binding typically blocks CD47 binding by 50% or greater, typically 75%, or 90% or greater.

*Inhibition of SIRPA activities*

**[0108]** In some embodiments, anti-SIRPA antibodies of the present disclosure inhibit one or more activities of SIRPA, including, without limitation: SIRPA binding to one or more SIRPA ligands, optionally wherein the one or more SIRPA ligands are selected from the group consisting of CD47, surfactant protein A and D and any combination thereof; decreasing proliferation of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia; inhibiting migration of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia; inhibiting one or more functions of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2

macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia; inhibition of one or more types of clearance selected from the group consisting of apoptotic neuron clearance, nerve tissue debris clearance, dysfunctional synapse clearance, non-nerve tissue debris clearance, bacteria clearance, other foreign body clearance, disease-causing protein clearance, disease-causing peptide clearance, and tumor cell clearance; optionally wherein the disease-causing protein is selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides and the tumor cell is from a cancer selected from the group consisting of bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal cell cancer, renal pelvis cancer, leukemia, lung cancer, melanoma, non-Hodgkin's lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, fibrosarcoma, and thyroid cancer; inhibition of tumor cell killing by one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells; inhibiting anti-tumor cell proliferation activity of one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells; modulated expression of one or more inflammatory receptors, optionally wherein the one or more inflammatory receptors comprise CD86 and the one or more inflammatory receptors are expressed on one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells; promoting or rescuing functionality of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid-derived suppressor cells, tumor-associated macrophages, tumor-associated neutrophils, tumor-associated NK cells, and regulatory T

cells; increasing infiltration of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid-derived suppressor cells, tumor-associated macrophages, tumor-associated neutrophils, tumor-associated NK cells, non-tumorigenic CD45+CD14+ myeloid cells, and regulatory T cells into tumors; increasing the number of tumor-promoting myeloid/granulocytic immune-suppressive cells and/or non-tumorigenic CD45+CD14+ myeloid cells in a tumor, in peripheral blood, or other lymphoid organ; enhancing tumor-promoting activity of myeloid-derived suppressor cells and/or non-tumorigenic CD45+CD14+ myeloid cells; enhancing survival of non-tumorigenic myeloid-derived suppressor cells and/or non-tumorigenic CD45+CD14+ myeloid cells; decreasing activation of tumor-specific T lymphocytes with tumor killing potential; decreasing infiltration of tumor-specific NK cells with tumor killing potential; increasing tumor volume; increasing tumor growth rate; and decreasing efficacy of one or more immune-therapies that modulate anti-tumor T cell responses, optionally wherein the one or more immune-therapies are immune-therapies that target one or more target proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, TREM1, TREM2, CD39, CD73, CSF-1 receptor, and any combination thereof, or of one or more cancer vaccines.

**[0109]** In some embodiments, which may be combined with any of the other embodiments above, an anti-SIRPA antibody of the present disclosure induces one or more of the activities that are selected from the group consisting of increasing the number of tumor infiltrating CD3+ T cells; decreasing cellular levels of SIRPA in non-tumorigenic CD14+myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood; reducing the number of non-tumorigenic CD14+ myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood; reducing PD-L1 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); reducing PD-L2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); reducing B7-H2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); reducing B7-H3 levels in one or

more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); reducing CD200R levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); reducing CD163 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); reducing CD206 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC); decreasing tumor growth rate of solid tumors; reducing tumor volume; increasing efficacy of one or more PD-1 inhibitors; increasing efficacy of one or more checkpoint inhibitor therapies and/or immune-modulating therapies, optionally wherein the one or more checkpoint inhibitor therapies and/or immune-modulating therapies target one or more of CTLA4, the adenosine pathway, PD-L1, PD-L2, OX40, TIM3, LAG3, or any combination thereof; increasing efficacy of one or more chemotherapy agents, optionally wherein the one or more of the chemotherapy agents are gemcitabine, capecitabine, anthracyclines, doxorubicin (Adriamycin®), epirubicin (Ellence®), taxanes, paclitaxel (Taxol®), docetaxel (Taxotere®), 5-fluorouracil (5-FU), cyclophosphamide (Cytoxan®), carboplatin (Paraplatin®), and any combination thereof; increasing proliferation of T cells in the presence of non-tumorigenic myeloid-derived suppressor cells (MDSC); inhibiting differentiation, survival, and/or one or more functions of non-tumorigenic myeloid-derived suppressor cells (MDSC); and killing CD33-expressing immunosuppressor non-tumorigenic myeloid cells and/or non-tumorigenic CD14-expressing cells in solid tumors and associated blood vessels when conjugated to a chemical or radioactive toxin.

**[0110]** In some embodiments, an anti-SIRPA antibody of the present disclosure decreases the activity, functionality, or survival of regulatory T cells, tumor-imbedded immunosuppressor dendritic cells, tumor-imbedded immunosuppressor macrophages, myeloid-derived suppressor cells, tumor-associated macrophages, acute myeloid leukemia (AML) cells, chronic lymphocytic leukemia (CLL) cell, or chronic myeloid leukemia (CML).

**[0111]** In some embodiments, an anti-SIRPA antibody of the present disclosure induces or promotes the survival, maturation, functionality, migration, or proliferation of one or more immune cells, *e.g.*, one or more immune cells are selected from the group consisting of dendritic cells, macrophages, neutrophils, NK cells, microglia, T cells, T helper cells, cytotoxic T cells, and any combination thereof in an individual.

[0112] As used herein, levels of SIRPA may refer to expression levels of the gene encoding SIRPA; to expression levels of one or more transcripts encoding SIRPA; to expression levels of SIRPA protein; and/or to the amount of SIRPA protein present within cells and/or on the cell surface. Any methods known in the art for measuring levels of gene expression, transcription, translation, and/or protein abundance or localization may be used to determine the levels of SIRPA.

[0113] In some embodiments, an isolated anti-SIRPA antibody of the present disclosure is a murine antibody. In some embodiments, an isolated anti-SIRPA antibody of the present disclosure is a human antibody, a humanized antibody, a bispecific antibody, a monoclonal antibody, a multivalent antibody, or a chimeric antibody. Exemplary descriptions of such antibodies are found throughout the present disclosure.

[0114] In some embodiments, anti-SIRPA antibodies of the present disclosure bind to a human SIRPA including human allelic variants (FIG. 1A, accession numbers NP542970 and CAA71403). In some embodiments, anti-SIRPA antibodies specifically bind to primate SIRPA, including human SIRPA. In some embodiments, anti-SIRPA antibodies of the present disclosure specifically bind to both human SIRPA and primate SIRPA. In some embodiments, anti-SIRPA antibodies of the present disclosure specifically bind to human SIRPA and cross-react with murine SIRPA.

*HVR sequences of antibodies that down-regulate SIRPA that do not block CD47 binding*

[0115] In some embodiments, an anti-SIRPA antibody of the present disclosure down-regulates SIRPA and does not block CD47 binding to SIRPA. In some embodiments, such an antibody comprises a heavy chain variable region that comprises an HVR3 of antibody 3F9 as set forth in SEQ ID NO:11. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:11 in which 1, 2, 3, 4, or 5 amino acids are substituted, e.g., conservatively substituted. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:11 in which 1, 2, 3, or 4 amino acids are substituted, e.g., conservatively substituted. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:11 in which 1, 2, or 3 amino acids are substituted, e.g., conservatively substituted. In some embodiments the HVR3 has 1 or 2 amino acids substituted compared to the sequence set forth in SEQ ID NO:11. In some embodiments, 1 or 2 amino acids are deleted, relative to SEQ ID NO:11. In some embodiments, the HVR3 has at least 65% identity or at least 75% identity to the amino acid sequence of SEQ ID NO:11.

**[0116]** In some embodiments, a heavy chain variable region of an anti-SRPA antibody of the invention comprises an HVR3 as set forth in the preceding paragraph and an HVR1 and/or an HVR2 of antibody 3F9 as set forth in Table 3. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:9 in which 1, 2, 3, or 4 amino acid are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:9 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 has at least 70%, at least 80%, or at least 90% identity to the amino acid sequence of SEQ ID NO:9. In some embodiments, the HVR2 comprises the sequence of SEQ ID NO:10 in which 1, 2, 3, or 4 amino acid are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR2 comprises the sequence of SEQ ID NO:10 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR2 has at least 70%, at least 80%, or at least 90% identity to the amino acid sequence of SEQ ID NO:10.

**[0117]** In some embodiments, an anti-SIRPA antibody comprises a heavy chain variable region comprising an HVR3 of SEQ ID NO:11, an HVR1 of SEQ ID NO:9, and an HVR2 of SEQ ID NO:10.

**[0118]** In some embodiments, an anti-SIRPA antibody comprises a light chain variable region that comprises an HVR3 of antibody 3F9 as set forth in Table 2. In some embodiments, the HVR3 comprises the sequence the sequence set forth in SEQ ID NO:8 in which 1, 2, 3, or 4 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:8 in which 1, 2, or 3 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments the HVR3 has 1 or 2 amino acid substituted compared to the sequence set forth in SEQ ID NO:8. In some embodiments, 1 or 2 amino acids are deleted, relative to SEQ ID NO:8. In some embodiments, the HVR3 has at least 65% identity to the amino acid sequence of SEQ ID NO:8. In some embodiments, the HVR3 has at least 85% identity to the amino acid sequence of SEQ ID NO:8.

**[0119]** In some embodiments, a light chain variable region of an anti-SRPA antibody of the invention comprises an HVR3 as set forth in the preceding paragraph and an HVR1 and/or an HVR2 of antibody 3F9 as set forth in Table 2. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:6 in which 1, 2, 3, 4, 5, or 6; or 1, 2, 3, 4, or 5; amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 comprises the

sequence of SEQ ID NO:6 in which 1, 2, 3, or 4 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:6 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids, are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 has at least 70%, at least 80%, or at least 90% identity to the amino acid sequence of SEQ ID NO:6. In some embodiments, the HVR2 comprises the sequence of SEQ ID NO:7 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR2 has at least 70%, at least 85%, identity to the amino acid sequence of SEQ ID NO:7.

**[0120]** In some embodiments, an anti-SIRPA antibody comprises a light chain variable region having an HVR3 of SEQ ID NO:8, an HVR1 of SEQ ID NO:6, and an HVR2 of SEQ ID NO:7.

**[0121]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises a heavy chain variable region comprising an HVR3, HVR2, and HVR1 of antibody 3F9 as set forth in Table 3 and a light chain variable region comprising an HVR3, HVR2, and HVR1 of antibody 3F9 as set forth in Table 2. In some embodiments, an anti-SIRPA antibody comprises the six CDRs of 3F9 where at least one HVR differs from the HVR of 3F9 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR of 3F9. In some embodiments, such an antibody comprises two HVRs that differ from the corresponding HVR of 3F9 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 3F9. In some embodiments, the antibody comprises three HVRs that differ from the corresponding HVR of 3F9 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 3F9. In some embodiments, the antibody comprises four HVRs that differ from the corresponding HVR of 3F9 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 3F9. In some embodiments, the antibody comprises five HVRs that differ from the corresponding HVR of 3F9 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 3F9. In some emodiments, the antibody comprises one, two, or 3 amino acid changes; or one or two amino acid changes, in each HVR compared to the corresponding HVR of 3F9.

**[0122]** In some embodiments, an anti-SIRPA antibody comprises a heavy chain variable region that comprises an HVR3 of antibody 9C2 as set forth in SEQ ID NO:17. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:17 in which 1, 2, 3,

or 4 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:7 in which 1, 2, or 3 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments the HVR3 has 1 or 2 amino acid substituted compared to the sequence set forth in SEQ ID NO:17. In some embodiments, 1 or 2 amino acids are deleted, relative to SEQ ID NO:17. In some embodiments, the HVR3 has at least 65% identity to the amino acid sequence of SEQ ID NO:17. In some embodiments, the HVR3 has at least 85% identity to the amino acid sequence of SEQ ID NO:17.

**[0123]** In some embodiments, a heavy chain variable region of an anti-SRPA antibody of the invention comprises an HVR3 as set forth in the preceding paragraph and an HVR1 and/or an HVR2 of antibody 9C2 as set forth in Table 3. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:15 in which 1, 2, 3, or 4 amino acid are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:15 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 has at least 70%, at least 80%, or at least 90% identity to the amino acid sequence of SEQ ID NO:15. In some embodiments, the HVR2 comprises the sequence of SEQ ID NO:16 in which 1, 2, 3, or 4 amino acid are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR2 comprises the sequence of SEQ ID NO:16 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR has at least 70%, at least 80%, or at least 90% identity to the amino acid sequence of SEQ ID NO:16.

**[0124]** In some embodiments, an anti-SIRPA antibody comprises a heavy chain variable region comprising an HVR3 of SEQ ID NO:17, an HVR1 of SEQ ID NO:15, and an HVR2 of SEQ ID NO:16.

**[0125]** In some embodiments, an anti-SIRPA antibody comprises a light chain variable region that comprises an HVR3 of antibody 9C2 as set forth in Table 2. In some embodiments, the HVR3 comprises the sequence the sequence set forth in SEQ ID NO:14 in which 1, 2, 3, or 4 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR3 comprises the sequence set forth in SEQ ID NO:4 in which 1, 2, or 3 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments the HVR3 has 1 or 2 amino acid substituted compared to the sequence set forth in SEQ ID

NO:14. In some embodiments, 1 or 2 amino acids are deleted, relative to SEQ ID NO:14. In some embodiments, the HVR3 has at least 65% identity to the amino acid sequence of SEQ ID NO:14. In some embodiments, the HVR3 has at least 85% identity to the amino acid sequence of SEQ ID NO:14.

**[0126]** In some embodiments, a light chain variable region of an anti-SIRPA antibody of the invention comprises an HVR3 as set forth in the preceding paragraph and an HVR1 and/or an HVR2 of antibody 9C2 as set forth in Table 2. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:12 in which 1, 2, 3, or 4 amino acid are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 comprises the sequence of SEQ ID NO:12 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids, are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR1 has at least 70% identity, at least 80% identity, or at least 90% identity to the amino acid sequence of SEQ ID NO:12. In some embodiments, the HVR2 comprises the sequence of SEQ ID NO:13 in which 1, 2, or 3 amino acids; or 1 or 2 amino acids are substituted, *e.g.*, conservatively substituted. In some embodiments, the HVR2 has at least 70%, at least 85%, identity to the amino acid sequence of SEQ ID NO:13.

**[0127]** In some embodiments, an anti-SIRPA antibody comprises a light chain variable region having an HVR3 of SEQ ID NO:14, an HVR1 of SEQ ID NO:12, and an HVR2 of SEQ ID NO:13.

**[0128]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises a heavy chain variable region comprising an HVR3, HVR2, and HVR1 of antibody 9C2 as set forth in Table 3 and a light chain variable region comprising an HVR3, HVR2, and HVR1 of antibody 9C2 as set forth in Table 2. In some embodiments, an anti-SIRPA antibody comprises at least one HVR that differs from the HVR of 9C2 by one, two or three amino acids; or one or two amino acids, compared to the corresponding HVR of 9C2. In some embodiments, such an antibody comprises two HVRs that differ from the corresponding HVR of 9C2 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 9C2. In some embodiments, the antibody comprises three HVRs that differ from the corresponding HVR of 9C2 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 9C2. In some embodiments, the antibody comprises four HVRs that differ from the corresponding HVR of 9C2 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of

3F9. In some embodiments, the antibody comprises five HVRs that differ from the corresponding HVR of 9C2 by one, two, or three amino acids; or one or two amino acids, compared to the corresponding HVR in of 9C2. In some emodiments, the antibody comprises one, two, or 3 amino acid changes; or one or two amino acid changes, in each HVR compared to the corresponding HVR of 9C2.

**[0129]** In some embodiments, an N residue present in a light chain CDR2 (SEQ ID NO:7) from 3F9 may be substituted with Q, S, A, or D. In some embodiments, the N residue in a light chain CDR3 (SEQ ID NO:8) from 3F9 in Table 2 may be substituted with Q, S, A, or D. In some instances, the N residues in both the light chain CDR2 and CDR3 are substituted with Q, S, A, or D. In some embodiments, the C in a light chain CDR3 (SEQ ID NO:8) from 3F9 may be substituted with an A, S, or L.

**[0130]** In some embodiments, an N residue present in a heavy chain CDR1 (SEQ ID NO:15) from 9C2 may be substituted with Q, S, or A. In some embodiments, one or both N residues present in a heavy chain CDR2 (SEQ ID NO:16) from 9C2 may be substituted with Q, S, or A. In some embodiments, the D of a heavy chain CDR3 residues Asp-Gly (DG) of SEQ ID NO:17 may be substituted with an A, S, or E. In some embodiments, an N residue in a light chain CDR2 (SEQ ID NO:13) from 9C2 may be substituted with Q, S, D, or A. In some embodiments, an N residue in a light chain CDR3 (SEQ ID NO:14) from 9C2 may be substituted with Q, S, D, or A. A light chain CDR3 of SEQ ID NO:14 may also contain a H, Y, or F residue substituted for the Trp residue in the 9C2 light chain CDR3.

#### *Antibody frameworks*

**[0131]** Any of the antibodies described herein further include a framework, preferably a human immunoglobulin framework. For example, in some embodiments, an antibody comprises HVRs as in any of the above embodiments and further comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework. Human immunoglobulin frameworks may be part of the human antibody, or a non-human antibody may be humanized by replacing one or more endogenous frameworks with human framework region(s). Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the "best-fit" method (see, *e.g.*, Sims et al. *J. Immunol.* 151:2296 (1993)); framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (see, *e.g.*, Carter et al. *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); and

Presta et al. *J. Immunol.*, 151:2623 (1993)); human mature (somatically mutated) framework regions or human germline framework regions (see, *e.g.*, Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008)); and framework regions derived from screening FR libraries (see, *e.g.*, Baca et al., *J. Biol. Chem.* 272:10678-10684 (1997) and Rosok et al., *J. Biol. Chem.* 271:22611-22618 (1996)).

**[0132]** In some embodiments, an antibody of the present disclosure has the binding specificity of 3F9 and comprises heavy chain HVR1, HVR2, and HVR3 sequences as described above, and further, comprises at least one heavy chain framework as shown in Figure 14A, *e.g.*, hSB-3F9-H1 or hSB-3F9-H2 sequence of Figure 14A. A “framework” in this context refers to the FR1, FR2, FR3, and FR4 sequences and excludes the CDR sequence. In some embodiments, an anti-SIRPA antibody has a framework that has at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino acid sequence identity to a framework shown in Figure 14A, where the percent identity is determined based on the FR1, FR2, FR3, and FR4 sequences excluding the CDRs.

**[0133]** In some embodiments, an antibody of the present disclosure has the binding specificity of 3F9 and comprises light chain HVR1, HVR2, and HVR3 sequences as described above, and further, comprises at least one light chain framework as shown in Figure 14B, *e.g.*, an hSB-3F9-L1, hSB-3F9-L2, or hsB-3F9-L3 sequence of Figure 14B. A “framework” in this context refers to the FR1, FR2, FR3, and FR4 sequences and excludes the CDR sequence. In some embodiments, an anti-SIRPA antibody has a framework that has at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino acid sequence identity to a framework shown in Figure 14B, where the percent identity is determined based on the FR1, FR2, FR3, and FR4 sequences excluding the CDRs.

**[0134]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises a V<sub>H</sub> region and V<sub>L</sub> region as set forth in the two preceding paragraphs.

**[0135]** In some embodiments, an antibody of the present disclosure has the binding specificity of 9C2 and comprises heavy chain HVR1, HVR2, and HVR3 sequences as described above, and further, comprises at least one heavy chain framework as shown in Figure 14C, *e.g.*, the hSB-9C2-H1, hSB-9C2-H2, hSB-9C2-H3, or hSB-9C2-H4 sequence of Figure 14C. A “framework” in this context refers to the FR1, FR2, FR3, and FR4 sequences

and excludes the CDR sequence. In some embodiments, an anti-SIRPA antibody has a framework that has at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino acid sequence identity to a framework shown in Figure 14C, where the percent identity is determined based on the FR1, FR2, FR3, and FR4 sequences excluding the CDRs.

**[0136]** In some embodiments, an antibody of the present disclosure has the binding specificity of 3F9 and comprises light chain HVR1, HVR2, and HVR3 sequences as described above, and further, comprises at least one light chain framework as shown in Figure 14D, *e.g.*, an hSB-9C2-L1, hSB-9C2-L2, hSB-9C2-L3, or hSB-9C2-L sequence of Figure 14D. A “framework” in this context refers to the FR1, FR2, FR3, and FR4 sequences and excludes the CDR sequence. In some embodiments, an anti-SIRPA antibody has a framework that has at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino acid sequence identity to a framework shown in Figure 14D, where the percent identity is determined based on the FR1, FR2, FR3, and FR4 sequences excluding the CDRs.

**[0137]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises a V<sub>H</sub> region and V<sub>L</sub> region as set forth in the two preceding paragraphs.

**[0138]** In some embodiments an anti-SIRPA antibody of the present invention comprises one or more e substitutions relative to the CDR and framework region sequences shown in Figures 10A-10D. In some embodiments, substitutions are conservative substitutions. Illustrative substitutions are provided below:

#### Amino Acid Substitutions

Original Residue	Illustrative Substitutions	Frequent Substitution
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; asp, lys; arg	gln
Asp (D)	glu; asn	glu
Cys (C)	ser; ala	ser
Gln (Q)	asn; glu	asn
Glu (E)	asp; gln	asp
Gly (G)	ala	ala

Original Residue	Illustrative Substitutions	Frequent Substitution
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	tyr
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	Ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

[0139] In some embodiments, substitutions may be non-conservative substitutions.

Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

In some embodiments, non-conservative substitutions entail exchanging a member of one of these classes for another class.

[0140] Any cysteine residue not involved in maintaining the proper conformation of the antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment, such as an Fv fragment). In some embodiments, an anti-SIRPA antibody comprises a substitution at one or more C residues of a sequence shown in Figure 14A-14D.

[0141] In some embodiments, an anti-SIRPA antibody may also comprise substitutions at N residues, which are potential deamidation sites. In some embodiments, an anti-SIRPA

antibody of the invention comprises a substitution at one or more N residues of a sequence shown in Figure 14A-14D.

**[0142]** In some embodiments, an anti-SIRPA antibody comprises a substitution at a W residue, as W residues may be susceptible to oxidation. In some embodiments, the substitution is at one or more W residues of a sequence shown in Figure 14A-14D.

**[0143]** In some embodiments, an anti-SIRPA antibody that contains an Asp-Gly (DG) sequence, which may be susceptible to isoaspartate formation, may have an A or S substituted for a Gly or an E substituted for Asp.

*Anti-SIRPA1 antibody binding affinity*

**[0144]** An anti-SIRPA of the present disclosure may have nanomolar or even picomolar affinities for SIRPA. In certain embodiments, the dissociation constant ( $K_D$ ) of the antibody is about 0.05 to about 100 nM. For example,  $K_D$  of the antibody is any of about 100 nM, about 50 nM, about 10 nM, about 1 nM, about 900 pM, about 800 pM, about 790 pM, about 780 pM, about 770 pM, about 760 pM, about 750 pM, about 740 pM, about 730 pM, about 720 pM, about 710 pM, about 700 pM, about 650 pM, about 600 pM, about 590 pM, about 580 pM, about 570 pM, about 560 pM, about 550 pM, about 540 pM, about 530 pM, about 520 pM, about 510 pM, about 500 pM, about 450 pM, about 400 pM, about 350 pM about 300 pM, about 290 pM, about 280 pM, about 270 pM, about 260 pM, about 250 pM, about 240 pM, about 230 pM, about 220 pM, about 210 pM, about 200 pM, about 150 pM, about 100 pM, or about 50 pM to any of about 2 pM, about 5 pM, about 10 pM, about 15 pM, about 20 pM, or about 40 pM.

**[0145]** In some embodiments, the  $K_D$  of an anti-SIRPA for binding to human SIRPA may be about 200 nM or less, about 100 nM or less, about 50 nM or less, about 20 nM or less, about 10 nM or less, about 1 nM or less. In some embodiments, the  $K_D$  of an anti-SIRPA antibody for human SIRPA is about 100 pM or less or about 50 pM or less, less than about 10 pM, or less than about 1 pM. In some embodiments, the binding affinity is in the range of about 1 pM to about 200 nM. In some embodiments, the  $K_D$  is in the range of about 1 pM to about 100 nM.

**[0146]** In some embodiments, the  $K_D$  of an anti-SIRPA antibody for human SIRPA is less than 15 nM, less than 14.5 nM, less than 14 nM, less than 13.5 nM, less than 13 nM, less than 12.9 nM, less than 12.8 nM, less than 12.7 nM, less than 12.6 nM, less than 12.5 nM, less

than 12.4 nM, less than 12.3 nM, less than 12.2 nM, less than 12.1 nM, less than 12 nM, less than 11.5 nM, less than 11 nM, less than 10.9 nM, less than 10.8 nM, less than 10.7 nM, less than 10.6 nM, less than 10.5 nM, less than 10.4 nM, less than 10.3 nM, less than 10.2 nM, less than 10.1 nM, less than 10 nM, less than 9.5 nM, less than 9 nM, less than 8.5 nM, less than 8 nM, less than 7.5 nM, less than 7 nM, less than 6.9 nM, less than 6.8 nM, less than 6.7 nM, less than 6.6 nM, less than 6.5 nM, less than 6.4 nM, less than 6.3 nM, less than 6.2 nM, less than 6.1 nM, less than 6 nM, less than 5.5 nM, less than 5 nM, less than 4.5 nM, less than 4 nM, less than 3.5 nM, less than 3.4 nM, less than 3.3 nM, less than 3.2 nM, less than 3.1 nM, less than 3 nM, less than 2.9 nM, less than 2.8 nM, less than 2.7 nM, less than 2.6 nM, less than 2.5 nM, less than 2.4 nM, less than 2.3 nM, less than 2.2 nM, less than 2.1 nM, less than 2 nM, less than 1.9 nM, less than 1.8 nM, less than 1.7 nM, less than 1.6 nM, less than 1.5 nM, less than 1.4 nM, less than 1.3 nM, less than 1.2 nM, less than 1.1 nM, less than 1 nM, less than 0.95 nM, or less than 0.9 nM. In some embodiments, dissociation constants range from about 50 nM to about 100 pM.

**[0147]** Dissociation constants may be determined through any analytical technique, including any biochemical or biophysical technique such as ELISA, surface plasmon resonance (SPR), bio-layer interferometry (see, e.g., Octet System by ForteBio), isothermal titration calorimetry (ITC), differential scanning calorimetry (DSC), circular dichroism (CD), stopped-flow analysis, and colorimetric or fluorescent protein melting analyses.

#### *Antibody fragments*

**[0148]** Certain aspects of the present disclosure relate to a fragment of a SIRPA antibody as described herein where the fragment retains SIRPA binding activity. In some embodiments, the antibody fragment is an Fab, Fab', Fab'-SH, F(ab')<sub>2</sub>, Fv or scFv fragment. In some embodiments, an antibody fragment is provided in a multivalent format.

#### *Multivalent antibodies.*

**[0149]** In some embodiments, an anti-SIRPA antibody of the present invention may be in a multivalent format that is internalized faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The anti-SIRPA antibodies of the present disclosure or antibody fragments thereof can be multivalent antibodies (which are other than of the IgM class) with three or more antigen binding sites (e.g., tetravalent antibodies), which can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody. The multivalent antibody can comprise a dimerization domain and three or

more antigen binding sites. In typical embodiments, the dimerization domain comprises an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fc region. In some embodiments, a multivalent antibody contains three to eight, *e.g.*, four, antigen binding sites. The multivalent antibody contains at least one polypeptide chain (and preferably two polypeptide chains), wherein the polypeptide chain or chains comprise two or more variable domains.

*Bispecific and multi-specific antibodies*

[0150] Certain aspects of the present disclosure relate to bispecific antibodies, or multi-specific antibodies that comprise an anti-SIRPA antibody as described herein and an antibody that binds to a second antigen or a second SIRPA epitope. Bispecific and multi-specific antibodies may be generated using any method.

[0151] In some embodiment, the antibody is a bispecific antibody comprising a variable region of an anti-SIRPA antibody as described in the present disclosure and an antibody that binds to a second antigen. In some embodiments the second antigen is a protein selected from the group consisting of PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73. In some embodiments, the second antigen is an antigen facilitating transport across the blood-brain-barrier; an antigen facilitating transport across the blood-brain-barrier selected from the group consisting of transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), diphtheria toxin receptor, CRM197, a llama single domain antibody, TMEM 30(A), a protein transduction domain, TAT, Syn-B, penetratin, a poly-arginine peptide, an angiopep peptide, and ANG1005; a disease-causing agent selected from the group consisting of disease-causing peptides or proteins or, disease-causing nucleic acids, wherein the disease-causing nucleic acids are antisense GGCCCC (G2C4) repeat-expansion RNA, the disease-causing proteins are selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin,

lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides; or ligands and/or proteins expressed on immune cells, wherein the ligands and/or proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, and phosphatidylserine; and a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells.

*Fc regions*

**[0152]** In some embodiments, an antibody of the present disclosure comprises an Fc region. For example, the antibody may be of the IgG class, the IgM class, or the IgA class. In some embodiments, the has an IgG1, IgG2, IgG3, or IgG4 isotype. Typically, the Fc region is a native human Fc region or variant thereof.

**[0153]** In some embodiments, anti-SIRPA antibodies of the present disclosure retain the ability to bind Fc gamma receptors. In some embodiments, such antibodies may have features that result in clustering and transient stimulation of SIRPA. Such antibodies may subsequently act as longer-term inhibitors of SIRPA expression and/or one or more activities of SIRPA by inducing SIRPA degradation, SIRPAdesensitization, SIRPAcleavage, SIRPA internalization, SIRPA shedding, lysosomal degradation of SIRPA or otherwise down-regulating SIRPA. In some embodiments, anti-SIRPA antibodies decrease the level of Fc gamma receptors on the surface of cells.

**[0154]** In some embodiments, the Fc region is an Fc region that binds receptors such as Fc $\gamma$ RI, Fc $\gamma$ RII, and Fc $\gamma$ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors, Fc $\gamma$ RII receptors include Fc $\gamma$ RIIA (an “activating receptor”) and Fc $\gamma$ RIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor Fc $\gamma$ RIIA contains an immunoreceptor tyrosine-based activation motif (“ITAM”) in its cytoplasmic domain. Inhibiting receptor Fc $\gamma$ RIIB contains an immunoreceptor tyrosine-based inhibition motif (“ITIM”) in its cytoplasmic domain. (see, e.g., M. Daëron, Annu. Rev. Immunol. 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, Annu. Rev. Immunol. 9:457-92 (1991);

Capel et al., Immunomethods 4:25-34 (1994); and de Haas et al., J. Lab. Clin. Med. 126: 330-41 (1995). FcRs can also increase the serum half-life of antibodies.

**[0155]** An Fc region can include one or more mutations that influence activity of the Fc region, *e.g.*, in binding an Fc receptor.

**[0156]** In some embodiments, an antibody of the present disclosure binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (Fc $\gamma$ IIB). In some embodiments, an antibody of the present disclosure decreases the level of expression of inhibitory Fc-gamma receptor IIB on the surface of cells. In some embodiments, the Fc region contains one or more modifications. For example, in some embodiments, the Fc region contains one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments, the one or more amino acid substitutions are selected from V234A (Alegre et al., (1994) Transplantation 57:1537-1543. 31; Xu et al., (2000) Cell Immunol, 200:16-26), G237A (Cole et al. (1999) Transplantation, 68:563-571), H268Q, V309L, A330S, P331S (US 2007/0148167; Armour et al. (1999) Eur J Immunol 29: 2613-2624; Armour et al. (2000) The Haematology Journal 1(Suppl.1):27; Armour et al. (2000) The Haematology Journal 1(Suppl.1):27), C232S, and/or C233S (White et al. (2015) Cancer Cell 27, 138-148), S267E, L328F (Chu et al., (2008) Mol Immunol, 45:3926-3933), M252Y, S254T, and/or T256E, where the amino acid position is according to the EU or Kabat numbering convention.

**[0157]** In some embodiments, an antibody of the invention has an IgG2 isotype with a heavy chain constant domain that in some embodiments, contains a C127S or C2214S amino acid substitution, where the amino acid position is according to the EU or Kabat numbering convention (White et al., (2015) Cancer Cell 27, 138-148; Lightle et al., (2010) PROTEIN SCIENCE 19:753-762; and WO2008079246).

**[0158]** In certain embodiments, an antibody of the present disclosure has an IgG1 isotype. In some embodiments, the Fc gamma receptor-binding antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (Fc $\gamma$ IIB). In some embodiments, an antibody of the present disclosure decreases the level of inhibitory Fc-gamma receptor IIB expressed on the surface of cells. In some embodiments, the Fc region contains one or more modifications. For example, in some embodiments, the Fc region contains one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments, the one or more amino acid

substitutions are selected from N297A (Bolt S et al. (1993) Eur J Immunol 23:403-411), D265A (Shields et al. (2001) R. J. Biol. Chem. 276, 6591-6604), D270A, L234A, L235A (Hutchins et al. (1995) Proc Natl Acad Sci USA, 92:11980-11984; Alegre et al., (1994) Transplantation 57:1537-1543. 31; Xu et al., (2000) Cell Immunol, 200:16-26), G237A (Alegre et al. (1994) Transplantation 57:1537-1543. 31; Xu et al. (2000) Cell Immunol, 200:16-26), P238D, L328E, E233D, G237D, H268D, P271G, A330R, C226S, C229S, E233P, L234V, L234F, L235E (McEarchern et al., (2007) Blood, 109:1185-1192), P331S (Sazinsky et al., (2008) Proc Natl Acad Sci USA 2008, 105:20167-20172), S267E, L328F, A330L, M252Y, S254T, T256E, N297Q, P238S, P238A, A327Q, A327G, P329A, K322A, and/or T394D, where the amino acid position is according to the EU or Kabat numbering convention.

**[0159]** In some embodiments, an antibody of the present disclosure has an IgG1 isotype and includes an IgG2 isotype heavy chain constant domain 1 (CH1) and hinge region (White et al., (2015) Cancer Cell 27, 138-148). In certain embodiments, the IgG2 isotype CH1 and hinge region contain the amino acid sequence of

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSNFGTQTYTCNVDHKPSNTKVDKTVRKCCVECPPC (SEQ ID NO:34). In some embodiments, the antibody Fc region contains a S267E amino acid substitution, a L328F amino acid substitution, or both, and/or a N297A or N297Q amino acid substitution, where the amino acid position is according to the EU or Kabat numbering convention.

**[0160]** In some embodiments, an anti-SIRPA antibody has an IgG2 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: P238S, V234A, G237A, H268A, H268Q, V309L, A330S, P331S, C214S, C232S, C233S, S267E, L328F, M252Y, S254T, T256E, H268E, N297A, N297Q, A330L, and any combination thereof, wherein the numbering of the residues is according to EU numbering;

**[0161]** In certain embodiments, an antibody of the present disclosure has an IgG4 isotype. In some embodiments, the contains a human IgG4 constant region and comprises an Fc region that contains one or more amino acid substitutions (e.g., relative to a wild-type Fc region of the same isotype). In some embodiments, the one or more amino acid substitutions are selected from L235A, G237A, S228P, L236E (Reddy et al., (2000) J Immunol, 164:1925-

1933), S267E, E318A, L328F, M252Y, S254T, T256E, E233P, F234V, L234A/F234A, S228P, S241P, L248E, T394D, N297A, N297Q, L235E, and any combination thereof, wherein the numbering of the residues is according to EU numbering

**[0162]** In some embodiments, an anti-SIRPA antibody of the present disclosure has a hybrid IgG2/4 isotype. In certain embodiments the antibody comprises an amino acid sequence comprising amino acids 118 to 260 of human IgG2 and amino acids 261 to 447 of human IgG4, wherein the numbering of the residues is according to EU numbering.

**[0163]** In some embodiments, an anti-SIRPA antibody of the present disclosure has a human or mouse IgG1 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: N297A, N297Q, D270A, D265A, L234A, L235A, C226S, C229S, P238S, E233P, L234V, P238A, A327Q, A327G, P329A, K322A, L234F, L235E, P331S, T394D, A330L, M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU numbering.

**[0164]** In some embodiments, an anti-SIRPA antibody of the present disclosure has an IgG2 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: P238S, V234A, G237A, H268A, H268Q, H268E, V309L, N297A, N297Q, A330S, P331S, C232S, C233S, M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU numbering.

**[0165]** In some embodiments, an anti-SIRPA antibody of the present disclosure has an IgG4 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: E233P, F234V, L234A/F234A, L235A, G237A, E318A, S228P, L236E, S241P, L248E, T394D, M252Y, S254T, T256E, N297A, N297Q, and any combination thereof, wherein the numbering of the residues is according to EU numbering.

**[0166]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises an Fc region that further comprises one or more additional amino acid substitutions at a position selected from the group consisting of A330L, L234F; L235E, P331S, and any combination thereof, wherein the numbering of the residues is according to EU numbering.

**[0167]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises an Fc region that further comprises one or more additional amino acid substitutions at a

position selected from the group consisting of M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU numbering.

**[0168]** In some embodiments, an anti-SIRPA antibody of the present disclosure comprises an Fc region that further comprises a S228P amino acid substitution according to EU numbering.

**[0169]** In some embodiments, an anti-SIRPA antibody of the present disclosure has an IgG4 isotype and comprises an S228P amino acid substitution at residue position 228, an F234A amino acid substitution at residue position 234, and an L235A amino acid substitution at residue position 235, wherein the numbering of the residue position is according to EU numbering

**[0170]** In some embodiments, an anti-SIRPA antibody of the present disclosure may be modified to modulate effector function and/or to increase serum half-life of the antibody. For example, the Fc receptor binding site on the constant region may be modified or mutated to remove or reduce binding affinity to certain Fc receptors, such as Fc $\gamma$ RI, Fc $\gamma$ RII, and/or Fc $\gamma$ RIII to reduce antibody-dependent cell-mediated cytotoxicity. In some embodiments, the effector function is inhibited by removing N-glycosylation of the Fc region (*e.g.*, in the CH 2 domain of IgG) of the antibody. In some embodiments, the effector function is inhibited by modifying regions such as 233-236, 297, and/or 327-331 of human IgG as described in PCT WO 99/58572 and Armour et al., Molecular Immunology 40: 585-593 (2003); Reddy et al., J. Immunology 164:1925-1933 (2000). In other embodiments, it may also be desirable to modify an anti-SIRPA antibody of the present disclosure to modify effector function to increase binding selectivity toward the ITIM-containing Fc $\gamma$ RIIb (CD32b) to increase clustering of SIRPA antibodies on adjacent cells without activating effector functions such as ADCC.

**[0171]** In some embodiments, to increase the serum half-life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Patent 5,739,277, for example. As used herein, the term “salvage receptor binding epitope” refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG1, IgG2, IgG3, or IgG4) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

*Other amino acid sequence modifications*

[0172] Amino acid sequence modifications of anti-SIRPA antibodies of the present disclosure, or antibody fragments thereof, are also contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibodies or antibody fragments.

[0173] In some embodiments, additional amino acid sequences can be fused to the amino terminal or carboxy terminal of an anti-SIRPA antibody. Examples include, but are not limited to, an antibody with an N-terminal methionyl residue, fusion to a cytotoxic polypeptide, or fusion to an enzyme or a polypeptide that increases the serum half-life of the antibody.

[0174] In some embodiments, an antibody of the present invention may be mutated to alter the original glycosylation pattern of the antibody, *e.g.*, by deleting one mutating or more sites to prevent glycosylation by certain carbohydrate moieties and/or adding one or more glycosylation sites to introduce desired carbohydrate moieties.

[0175] Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylglucosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0176] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

*Other antibody modifications*

[0177] Anti-SIRPA antibodies of the present disclosure, or antibody fragments thereof, can be further modified to contain additional moieties, *e.g.*, moieties for derivitazation of the

antibody, drug moieties to be conjugated to the antibody and the like. Examples of moieties suitable for derivatization of an antibody are water-soluble polymers such as polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, polypropylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (*e.g.*, glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer is attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in a therapy under defined conditions, etc. Such techniques and other suitable formulations are disclosed in Remington: The Science and Practice of Pharmacy, 20th Ed., Alfonso Gennaro, Ed., Philadelphia College of Pharmacy and Science (2000).

**[0178]** In some embodiments, a cytotoxic agent or drug may be conjugated to an anti-SIRPA antibody of the present invention, *e.g.*, for the treatment of cancers, such as multiple myeloma or other cancers, that express SIRPA on the cell surface. Techniques to conjugate antibodies are disclosed are known in the art (see, *e.g.*, Jane de Lartigue, *OncLive* July 5, 2012; ADC Review on antibody-drug conjugates; and Ducry et al., (2010). *Bioconjugate Chemistry* 21 (1): 5–13). In some embodiments, the anti-SIRPA antibody is conjugated to a toxin selected from the group consisting of ricin, ricin A chain, doxorubicin, daunorubicin, a maytansinoid, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, *Saponaria officinalis* inhibitor, glucocorticoid, auristatin, auromycin, yttrium, bismuth, combrestatin, duocarmycins, dolastatin, cc1065, and a cisplatin.

**Nucleic acids, vectors, and host cells**

**[0179]** Anti-SIRPA antibodies of the present disclosure are commonly produced using recombinant methods. Accordingly, in some aspects, the invention provides, isolated nucleic acids comprising a nucleic acid sequence *e* encoding any of the anti-SIRPA antibodies as described herein; vectors comprising such nucleic acids and host cells into which the nucleic acids are introduced that are used to replicate the antibody-encoding nucleic acids and/or to express the antibodies. Such nucleic acids may encode an amino acid sequence containing the  $V_L$  and/or an amino acid sequence containing the  $V_H$  of the anti-SIRPA antibody (*e.g.*, the light and/or heavy chains of the antibody). In some embodiments, the host cell contains (1) a vector containing a polynucleotide that encodes the  $V_L$  amino acid sequence and a polynucleotide that encodes the  $V_H$  amino acid sequence, or (2) a first vector containing a polynucleotide that encodes the  $V_L$  amino acid sequence and a second vector containing a polynucleotide that encodes the  $V_H$  amino acid sequence. In some embodiments, the host cell is eukaryotic, *e.g.*, a Chinese Hamster Ovary (CHO) cell; or a human cell. In some embodiments, the host cell is a lymphoid cell (*e.g.*, Y0, NS0, Sp20 cell). Host cells of the present disclosure also include, without limitation, isolated cells, in vitro cultured cells, and ex vivo cultured cells.

**[0180]** In a further aspect, the invention provides a method of making an anti-SIRPA antibody as described herein. In some embodiments, the method includes culturing a host cell as described in the preceding paragraph under conditions suitable for expression of the antibody. In some embodiments, the antibody is subsequently recovered from the host cell (or host cell culture medium).

**[0181]** Suitable vectors containing polynucleotides encoding antibodies of the present disclosure, or fragments thereof include cloning vectors and expression vectors. While the cloning vector selected may vary according to the host cell intended to be used, useful cloning vectors generally have the ability to self-replicate, may possess a single target for a particular restriction endonuclease, and/or may carry genes for a marker that can be used in selecting clones containing the vector. Examples include plasmids and bacterial viruses, *e.g.*, pUC18, pUC19, Bluescript (*e.g.*, pBS SK+) and its derivatives, mpl8, mpl9, pBR322, pMB9, ColE1, pCR1, RP4, phage DNAs, and shuttle vectors such as pSA3 and pAT28. These and many other cloning vectors are available from commercial vendors such as BioRad, Strategene, and Invitrogen.

**[0182]** Expression vectors generally are replicable polynucleotide constructs that contain a nucleic acid of the present disclosure. The expression vector may be replicable in the host cells either as episomes or as an integral part of the chromosomal DNA. Suitable expression vectors include but are not limited to plasmids, viral vectors, including adenoviruses, adeno-associated viruses, retroviruses, and any other vector.

**[0183]** Suitable host cells for expressing an anti-SIRPA antibody as described herein include both prokaryotic or eukaryotic cells. For example, anti-SIRPA antibodies may be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified. Alternatively, the host cell may be a eukaryotic host cell, including eukaryotic microorganisms, such as filamentous fungi or yeast, including fungi and yeast strains whose glycosylation pathways have been “humanized,” resulting in the production of an antibody with a partially or fully human glycosylation pattern, vertebrate, invertebrate, and plant cells. Examples of invertebrate cells include insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells. Plant cell cultures can also be utilized as host cells.

**[0184]** In some embodiments, vertebrate host cells are used for producing anti-SIRPA antibodies of the present disclosure. For example, mammalian cell lines such as a monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, *e.g.*, in Graham et al., *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, *e.g.*, in Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK; buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep G2); mouse mammary tumor (MMT 060562); TRI cells, as described, *e.g.*, in Mather et al., *Annals N.Y. Acad. Sci.* 383:44-68 (1982); MRC 5 cells; and FS4 cells may be used to express anti-SIRPA antibodies. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR- CHO cells (Urlaub et al., *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); and myeloma cell lines such as Y0, NS0 and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, *e.g.*, Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ), pp. 255-268 (2003).

## Pharmaceutical Composition and Treatment Using an Anti-SIRPA Antibody

### *Pharmaceutical compositions*

**[0100]** Anti-SIRPA antibodies can be incorporated into a variety of formulations for therapeutic administration by combining the antibodies with appropriate pharmaceutically acceptable carriers or diluents, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms. Examples of such formulations include, without limitation, tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Pharmaceutical compositions can include, depending on the formulation desired, pharmaceutically-acceptable, non-toxic carriers of diluents, which are vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the combination. Examples of such diluents include, without limitation, distilled water, buffered water, physiological saline, PBS, Ringer's solution, dextrose solution, and Hank's solution. A pharmaceutical composition or formulation of the present disclosure can further include other carriers, adjuvants, or non-toxic, nontherapeutic, nonimmunogenic stabilizers, excipients and the like. The compositions can also include additional substances to approximate physiological conditions, such as pH adjusting and buffering agents, toxicity adjusting agents, wetting agents and detergents.

**[0101]** A pharmaceutical composition of the present disclosure can also include any of a variety of stabilizing agents, such as an antioxidant for example. When the pharmaceutical composition includes a polypeptide, the polypeptide can be complexed with various well-known compounds that enhance the *in vivo* stability of the polypeptide, or otherwise enhance its pharmacological properties (*e.g.*, increase the half-life of the polypeptide, reduce its toxicity, and enhance solubility or uptake). Examples of such modifications or complexing agents include, without limitation, sulfate, gluconate, citrate and phosphate. The polypeptides of a composition can also be complexed with molecules that enhance their *in vivo* attributes. Such molecules include, without limitation, carbohydrates, polyamines, amino acids, other peptides, ions (*e.g.*, sodium, potassium, calcium, magnesium, manganese), and lipids.

**[0102]** Further examples of formulations that are suitable for various types of administration can be found in Remington's *Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 22nd ed. (2012).

**[0103]** For oral administration, the active ingredient can be administered in solid dosage forms, such as capsules, tablets, and powders, or in liquid dosage forms, such as elixirs,

syrups, and suspensions. The active component(s) can be encapsulated in gelatin capsules together with inactive ingredients and powdered carriers, such as glucose, lactose, sucrose, mannitol, starch, cellulose or cellulose derivatives, magnesium stearate, stearic acid, sodium saccharin, talcum, magnesium carbonate. Examples of additional inactive ingredients that may be added to provide desirable color, taste, stability, buffering capacity, dispersion or other known desirable features are red iron oxide, silica gel, sodium lauryl sulfate, titanium dioxide, and edible white ink. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric-coated for selective disintegration in the gastrointestinal tract. Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

**[0104]** Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives.

**[0105]** The components used to formulate the pharmaceutical compositions are preferably of high purity and are substantially free of potentially harmful contaminants (e.g., at least National Food (NF) grade, generally at least analytical grade, and more typically at least pharmaceutical grade). Moreover, compositions intended for *in vivo* use are usually sterile. To the extent that a given compound must be synthesized prior to use, the resulting product is typically substantially free of any potentially toxic agents, particularly any endotoxins, which may be present during the synthesis or purification process. Compositions for parenteral administration are also sterile, substantially isotonic and made under GMP conditions.

**[0106]** Formulations may be optimized for retention and stabilization in the brain or central nervous system. When the agent is administered into the cranial compartment, it is desirable for the agent to be retained in the compartment, and not to diffuse or otherwise cross the blood brain barrier. Stabilization techniques include cross-linking, multimerizing, or linking to groups such as polyethylene glycol, polyacrylamide, neutral protein carriers, *etc.* in order to achieve an increase in molecular weight.

**[0107]** Other strategies for increasing retention include the entrapment of the antibody, such as an anti-SIRPA antibody of the present disclosure, in a biodegradable or bioerodible

implant. The rate of release of the therapeutically active agent is controlled by the rate of transport through the polymeric matrix, and the biodegradation of the implant. The transport of drug through the polymer barrier will also be affected by compound solubility, polymer hydrophilicity, extent of polymer cross-linking, expansion of the polymer upon water absorption so as to make the polymer barrier more permeable to the drug, geometry of the implant, and the like. The implants are of dimensions commensurate with the size and shape of the region selected as the site of implantation. Implants may be particles, sheets, patches, plaques, fibers, microcapsules and the like and may be of any size or shape compatible with the selected site of insertion.

**[0108]** The implants may have the active agent distributed through the polymeric matrix, or encapsulated, where a reservoir of active agent is encapsulated by the polymeric matrix. The selection of the polymeric composition to be employed will vary with the site of administration, the desired period of treatment, patient tolerance, the nature of the disease to be treated and the like. Characteristics of the polymers will include biodegradability at the site of implantation, compatibility with the agent of interest, ease of encapsulation, a half-life in the physiological environment.

**[0109]** Biodegradable polymeric compositions which may be employed may be organic esters or ethers, which when degraded result in physiologically acceptable degradation products, including the monomers. Anhydrides, amides, orthoesters or the like, by themselves or in combination with other monomers, may find use. The polymers will be condensation polymers. The polymers may be cross-linked or non-cross-linked. Of particular interest are polymers of hydroxyaliphatic carboxylic acids, either homo- or copolymers, and polysaccharides. Included among the polyesters of interest are polymers of D-lactic acid, L-lactic acid, racemic lactic acid, glycolic acid, polycaprolactone, and combinations thereof. By employing the L-lactate or D-lactate, a slowly biodegrading polymer is achieved, while degradation is substantially enhanced with the racemate. Copolymers of glycolic and lactic acid are of particular interest, where the rate of biodegradation is controlled by the ratio of glycolic to lactic acid. The most rapidly degraded copolymer has roughly equal amounts of glycolic and lactic acid, where either homopolymer is more resistant to degradation. The ratio of glycolic acid to lactic acid will also affect the brittleness of in the implant, where a more flexible implant is desirable for larger geometries. Among the polysaccharides of interest are calcium alginate, and functionalized celluloses, particularly carboxymethylcellulose esters characterized by being water insoluble, a molecular weight of about 5 kD to 500 kD, *etc.* Biodegradable hydrogels may also be

employed in the implants of the subject invention. Hydrogels are typically a copolymer material, characterized by the ability to imbibe a liquid. Exemplary biodegradable hydrogels which may be employed are described in Heller in: *Hydrogels in Medicine and Pharmacy*, N. A. Peppes ed., Vol. III, CRC Press, Boca Raton, Fla., 1987, pp 137-149.

[0110] Pharmaceutical compositions of the present disclosure containing an anti-SIRPA antibody of the present disclosure may be administered to an individual in need of treatment with the anti-SIRPA antibody, preferably a human, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, intracranial, intraspinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes.

[0111] Dosages and desired drug concentration of pharmaceutical compositions of the present disclosure may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary artisan. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles described in Mordini, J. and Chappell, W. "The Use of Interspecies Scaling in Toxicokinetics," In *Toxicokinetics and New Drug Development*, Yacobi et al., Eds, Pergamon Press, New York 1989, pp.42-46.

[0112] For *in vivo* administration of any of the anti-SIRPA antibodies of the present disclosure, normal dosage amounts may vary from about 10 ng/kg up to about 100 mg/kg of an individual's body weight or more per day, preferably about 1 mg/kg/day to 10 mg/kg/day, depending upon the route of administration. For repeated administrations over several days or longer, depending on the severity of the disease, disorder, or condition to be treated, the treatment is sustained until a desired suppression of symptoms is achieved.

[0113] An exemplary dosing regimen may include administering an initial dose of an anti-SIRPA antibody, of about 2 mg/kg, followed by a weekly maintenance dose of about 1 mg/kg every other week. Other dosage regimens may be useful, depending on the pattern of pharmacokinetic decay that the physician wishes to achieve. For example, dosing an individual from one to twenty-one times a week is contemplated herein. In certain embodiments, dosing ranging from about 3  $\mu$ g/kg to about 2 mg/kg (such as about 3  $\mu$ g/kg, about 10  $\mu$ g/kg, about 30  $\mu$ g/kg, about 100  $\mu$ g/kg, about 300  $\mu$ g/kg, about 1 mg/kg, and about 2 mg/kg) may be used. In certain embodiments, dosing frequency is three times per day, twice per day, once per day, once every other day, once weekly, once every two weeks, once every four weeks, once every five weeks, once every six weeks, once every seven weeks,

once every eight weeks, once every nine weeks, once every ten weeks, or once monthly, once every two months, once every three months, or longer. Progress of the therapy is easily monitored by conventional techniques and assays. The dosing regimen, including the anti-SIRPA antibody administered, can vary over time independently of the dose used.

**[0114]** Dosages for a particular anti-SIRPA antibody may be determined empirically in individuals who have been given one or more administrations of the anti-SIRPA antibody. Individuals are given incremental doses of an anti-SIRPA antibody. To assess efficacy of an anti-SIRPA antibody, a clinical symptom of the diseases, disorders, or conditions of the present disclosure (*e.g.*, cancer) can be monitored.

**[0115]** Administration of an anti-SIRPA antibody of the present disclosure can be continuous or intermittent, depending, for example, on the recipient's physiological condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to skilled practitioners. The administration of an anti-SIRPA antibody may be essentially continuous over a preselected period of time or may be in a series of spaced doses.

**[0185]** It is within the scope of the present disclosure that different formulations will be effective for different treatments and different disorders, and that administration intended to treat a specific organ or tissue may necessitate delivery in a manner different from that to another organ or tissue. Moreover, dosages may be administered by one or more separate administrations, or by continuous infusion. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

**[0186]** In one aspect of the invention, an agent that down-regulates SIRPA, *e.g.*, an anti-SIRPA antibody, is used as a therapeutic agent. Such agents are administered to treat, alleviate, and/or prevent a disease or pathology associated with SIRPA expression, activity and/or signaling in a subject. A therapeutic regimen is carried out by identifying a subject, *e.g.*, a human patient suffering from (or at risk of developing) a disease or disorder associated with SIRPA expression, activity and/or signaling, *e.g.*, a cancer or other neoplastic disorder, using standard methods. In some embodiments, cells having the pathology associated with SIRPA expression, activity, and/or signaling, express a SIRPA ligand, *e.g.*, CD47. In some embodiments, cells having the pathology associated with SIRPA expression, activity, and/or signaling, express SIRPA.

**[0187]** As further detailed below an agent that down-regulates SIRPA, *e.g.*, an anti-SIRPA antibody can be used in combination with an additional therapeutic agent that is used to treat the disease or pathology associated with SIRPA expression, activity, or signaling. The terms “in combination” and “in conjunction” are used interchangeably in the present disclosure. The additional therapeutic agent may be administered before, after, or concurrently with the agent that down-regulates SIRPA, *e.g.*, an anti-SIRPA antibody.

**[0188]** In one aspect of the present disclosure, an anti-SIRPA antibody preparation, *e.g.*, comprising an anti-SIRPA antibody that decreases expression of SIRPA on the cell surface, but does not substantially block binding of ligand, *e.g.*, CD47, to SIRPA, is administered to a human subject. Administration of the antibody may abrogate or inhibit or interfere with the expression, activity and/or signaling function of SIRPA that is mediated by ligand binding, *e.g.*, CD47 binding. In one embodiment the disease or disorder associated with SIRPA expression is cancer. In some embodiments, an anti-SIRPA antibody is administered to a patient that has a cancer, such as a hematological proliferative disorder of myeloid cells, that express SIRPA. In typical embodiments, an anti-SIRPA antibody is administered to a patient that has a cancer that expresses CD47.

**[0189]** In certain embodiments, the cancer is squamous cell carcinoma, small-cell lung cancer, non-small cell lung cancer, squamous non-small cell lung cancer (NSCLC), non-squamous NSCLC, glioma, gastrointestinal cancer, renal cancer (*e.g.* clear cell carcinoma), ovarian cancer, liver cancer, colorectal cancer, endometrial cancer, kidney cancer (*e.g.*, renal cell carcinoma (RCC)), prostate cancer (*e.g.* hormone refractory prostate adenocarcinoma), thyroid cancer, neuroblastoma, pancreatic cancer, glioblastoma (glioblastoma multiforme), cervical cancer, stomach cancer, bladder cancer, hepatoma, breast cancer, colon carcinoma, and head and neck cancer (or carcinoma), gastric cancer, germ cell tumor, pediatric sarcoma, sinonasal natural killer, melanoma (*e.g.*, metastatic malignant melanoma, such as cutaneous or intraocular malignant melanoma), bone cancer, skin cancer, uterine cancer, cancer of the anal region, testicular cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, solid tumors of childhood, cancer of the ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma,

epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally-induced cancers including those induced by asbestos, virus-related cancers (*e.g.*, human papilloma virus (HPV)-related tumor), and hematologic malignancies derived from either of the two major blood cell lineages, *i.e.*, the myeloid cell line (which produces granulocytes, erythrocytes, thrombocytes, macrophages and mast cells) or lymphoid cell line (which produces B, T, NK and plasma cells), such as all types of leukemias, lymphomas, and myelomas, *e.g.*, acute, chronic, lymphocytic and/or myelogenous leukemias, such as acute leukemia (ALL), acute myelogenous leukemia (AML), chronic lymphocytic leukemia (CLL), and chronic myelogenous leukemia (CML), undifferentiated AML (M0), myeloblastic leukemia (M1), myeloblastic leukemia (M2; with cell maturation), promyelocytic leukemia (M3 or M3 variant [M3V]), myelomonocytic leukemia (M4 or M4 variant with eosinophilia [M4E]), monocytic leukemia (M5), erythroleukemia (M6), megakaryoblastic leukemia (M7), isolated granulocytic sarcoma, and chloroma; lymphomas, such as Hodgkin's lymphoma (HL), non-Hodgkin's lymphoma (NHL), B cell hematologic malignancy, *e.g.*, B-cell lymphomas, T-cell lymphomas, lymphoplasmacytoid lymphoma, monocytoid B-cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphoma, anaplastic (*e.g.*, Ki 1+) large-cell lymphoma, adult T-cell lymphoma/leukemia, mantle cell lymphoma, angio immunoblastic T-cell lymphoma, angiocentric lymphoma, intestinal T-cell lymphoma, primary mediastinal B-cell lymphoma, precursor T-lymphoblastic lymphoma, T-lymphoblastic; and lymphoma/leukaemia (T-Lbly/T-ALL), peripheral T-cell lymphoma, lymphoblastic lymphoma, post-transplantation lymphoproliferative disorder, true histiocytic lymphoma, primary central nervous system lymphoma, primary effusion lymphoma, lymphoblastic lymphoma (LBL), hematopoietic tumors of lymphoid lineage, acute lymphoblastic leukemia, diffuse large B-cell lymphoma, Burkitt's lymphoma, follicular lymphoma, diffuse histiocytic lymphoma (DHL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, cutaneous T-cell lymphoma (CTLC) (also called mycosis fungoides or Sezary syndrome), and lymphoplasmacytoid lymphoma (LPL) with Waldenstrom's macroglobulinemia; myelomas, such as IgG myeloma, light chain myeloma, nonsecretory myeloma, smoldering myeloma (also called indolent myeloma), solitary plasmacytoma, and multiple myelomas, chronic lymphocytic leukemia (CLL), hairy cell lymphoma; hematopoietic tumors of myeloid lineage, tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; seminoma, teratocarcinoma, tumors of the central and peripheral nervous, including astrocytoma, schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and

other tumors, including melanoma, xeroderma pigmentosum, keratoacanthoma, seminoma, thyroid follicular cancer and teratocarcinoma, hematopoietic tumors of lymphoid lineage, for example T-cell and B-cell tumors, including but not limited to T-cell disorders such as T-prolymphocytic leukemia (T-PLL), including of the small cell and cerebriform cell type; large granular lymphocyte leukemia (LGL) preferably of the T-cell type; a/d T-NHL hepatosplenic lymphoma; peripheral/post-thymic T cell lymphoma (pleomorphic and immunoblastic subtypes); angiocentric (nasal) T-cell lymphoma; cancer of the head or neck, renal cancer, rectal cancer, cancer of the thyroid gland; acute myeloid lymphoma, as well as any combinations of said cancers. Anti-SIRPA antibodies of the present invention may also be used to treat metastatic cancer.

**[0190]** In some embodiments, the cancer is selected from the group consisting of sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, melanoma, lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, and fibrosarcoma.

**[0191]** In some embodiments, the cancer is selected from the group consisting of glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma; diffuse large B-cell lymphoma; lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, non-Hodgkin's lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thyumoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma

**[0192]** In some embodiments, an anti-SIRPA antibody of the present disclosure may be administered in conjunction with a therapeutic agent that acts as a checkpoint inhibitor. In some embodiments, the checkpoint inhibitor targets PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA,

CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, and CD73. In typical embodiments, the therapeutic agent is an antibody to a checkpoint inhibitor selected from D1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73. In some embodiments, a combination of antibodies to checkpoint inhibitors is administered in conjunction in an anti-SIRPA antibody of the present invention.

**[0193]** In some embodiments, an anti-SIRPA antibody of the present disclosure may be administered in conjunction with at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein, *e.g.*, an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonistic anti-TREM1 antibody, an agonistic anti-TREM2 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, an agonist anti-CD30 antibody, an agonist anti-BTLA antibody, an agonist anti-HVEM antibody, an agonist anti-CD2 antibody, an agonist anti-CD5 antibody, and any combination thereof.

**[0194]** In some embodiments, an anti-SIRPA antibody of the present invention is administered in combination with radiation therapy and/or a chemotherapeutic agents. Chemotherapeutic agents include, for example, the following groups: anti-metabolites/anti-cancer agents, such as pyrimidine analogs (5-fluorouracil, floxuridine, capecitabine, gemcitabine and cytarabine) and purine analogs, folate antagonists and related inhibitors (methotrexate, pemetrexed, mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine (cladribine)); antiproliferative/antimitotic agents including natural products such as vinca alkaloids (vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane (paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones, eribulin and navelbine; epidipodophyllotoxins (etoposide, teniposide); DNA damaging agents (actinomycin, amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin, chlorambucil, cisplatin, cyclophosphamide, Cytoxin, dactinomycin, daunorubicin, doxorubicin, epirubicin, hexamethylmelamineoxaliplatin, iphosphamide, melphalan, mechlorethamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine, taxol, taxotere, temozolamide, teniposide, triethylenethiophosphoramide and etoposide (VP 16)); DNA methyltransferase inhibitors (azacytidine); antibiotics such as dactinomycin (actinomycin D), daunorubicin, doxorubicin

(adriamycin), idarubicin, anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin) and mitomycin; enzymes (L-asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and analogs, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepa), alkylsulfonates (busulfan), nitrosoureas (carmustine (BCNU) and analogs, streptozocin), triazenes (dacarbazine (DTIC)); antiproliferative/antimitotic antimetabolites such as folic acid analogs (methotrexate); platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogs (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase), aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine, mycophenolate mofetil); anti-angiogenic compounds (TNP470, genistein, pomalidomide) and growth factor inhibitors (vascular endothelial growth factor (VEGF) inhibitors, such as ziv-aflibercept; fibroblast growth factor (FGF) inhibitors); inhibitors of apoptosis protein (IAP) antagonists (birinapant); histone deacetylase (HDAC) inhibitors (vorinostat, romidepsin, chidamide, panobinostat, mocetinostat, abexinostat, belinostat, entinostat, resminostat, givinostat, quisinostat, SB939); proteasome inhibitors (ixazomib); angiotensin receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab, panitumumab, pertuzumab, cetuximab, adalimumab, golimumab, infliximab, rituximab, ocrelizumab, ofatumumab, obinutuzumab, alemtuzumab, abciximab, atlizumab, daclizumab, denosumab, efalizumab, elotuzumab, rovelizumab, ruplizumab, ustekinumab, visilizumab, gemtuzumab ozogamicin, brentuximab vedotin); chimeric antigen receptors; cell cycle inhibitors (flavopiridol, roscovitine, bryostatin-1) and differentiation inducers (tretinoin); mTOR inhibitors, topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin, irinotecan (CPT-11) and mitoxantrone, topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prenisolone); PARP inhibitors (niraparib, olaparib); focal adhesion kinase (FAK) inhibitors (defactinib (VS-6063), VS-4718, VS-6062, GSK2256098); growth factor signal transduction kinase inhibitors (cediranib, galunisertib, rociletinib, vandetanib, afatinib, EGF816,

AZD4547); c-Met inhibitors (capmatinib, INC280); ALK inhibitors (ceritinib, crizotinib); mitochondrial dysfunction inducers, toxins such as Cholera toxin, ricin, *Pseudomonas* exotoxin, *Bordetella* pertussis adenylate cyclase toxin, or diphtheria toxin, and caspase activators; and chromatin disruptors. In some embodiments, a chemotherapeutic agent is a B-Raf inhibitor, a MEK inhibitor, a VEGF inhibitor, a VEGFR inhibitor, a tyrosine kinase inhibitor, an anti-mitotic agent, or any combination thereof.

**[0195]** In some embodiments, an anti-SIRPA antibody of the present disclosure is administered in combination with adoptive cell transfer (ACT) therapy, chimeric antigen receptor T cell transfer (CAR-T) therapy, vaccine therapy, and/or cytokine therapy.

**[0196]** In some embodiments, an anti-SIRPA antibody of the present disclosure is administered in combination with at least one antibody that specifically binds to an inhibitory cytokine, *e.g.*, an inhibitory cytokine such as an anti-CCL2 antibody, an anti-CSF-1 antibody, or an anti-IL-2 antibody.

**[0197]** In some embodiments, an anti-SIRPA antibody of the present disclosure is administered in combination with at least one stimulatory cytokine. In some embodiments that may be combined with any of the preceding embodiments, the at least one stimulatory cytokine is selected from the group consisting of IFN- $\alpha$ 4, IFN- $\beta$ , IL-1 $\beta$ , TNF- $\alpha$ , IL-6, IL-8, CRP, IL-20 family members, LIF, IFN- $\gamma$ , OSM, CNTF, GM-CSF, IL-11, IL-12, IL-15, IL-17, IL-18, IL-23, CXCL10, IL-33, MCP-1, MIP-1-beta, and any combination thereof.

**[0198]** In some embodiments, an agent that down-regulates SIRPA, *e.g.*, an anti-SIRPA antibody, is administered to a patient that has a neurological disorder, or is administered to reduce risk, slow onset, or prevent a neurological disorder. In some embodiments, the neurological disorder is dementia, including frontotemporal dementia, Alzheimer's disease, or vascular dementia. In some embodiments, the patient has mild cognitive impairment.

**[0199]** In some embodiments, an agent that down-regulates SIRPA, *e.g.*, an anti-SIRPA antibody, is administered to a patient that has Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, Taupathy diseases, or multiple sclerosis. In some embodiments, the agent is administered to a patient that has Creutzfeldt-Jakob disease, normal pressure hydrocephalus, Nasu-Hakola disease, stroke, an infection, traumatic brain injury, progressive supranuclear palsy, dementia pugilistica (chronic traumatic encephalopathy), Parkinsonism linked to chromosome 17, Lytico-Bodig disease (Parkinson-dementia complex of Guam), tangle-predominant dementia, ganglioglioma and

gangliocytoma, meningioangiomatosis, subacute sclerosing panencephalitis, lead encephalopathy, tuberous sclerosis, Hallervorden-Spatz disease, lipofuscinosis, Pick's disease, corticobasal degeneration, Argyrophilic grain disease (AGD), frontotemporal lobar degeneration, dementia with Lewy bodies, multiple system atrophy, Shy-Drager syndrome, progressive supranuclear palsy, or cortical basal ganglionic degeneration.

## EXAMPLES

**[0200]** The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of non-critical parameters that could be changed or modified to yield essentially similar results.

### Example 1: Production of anti-SIRPA antibodies

**[0201]** The amino acid sequence of the human SIRPA preprotein is set forth below in SEQ ID NO:1. Human SIRPA contains a signal peptide located at amino residues 1-30 of SEQ ID NO:1. Human SIRPA contains an extracellular immunoglobulin-like variable-type (IgV) domain located at amino residues 32-137 of SEQ ID NO:1; additional extracellular immunoglobulin-like constant-type (IgC) domain sequences located at amino residues 148-247 and 254-348 of SEQ ID NO:1; a transmembrane domain located at amino residues 374-394 of SEQ ID NO:1; and an intracellular domain located at amino residues 395-504 of SEQ ID NO:1.

### SIRPAv1 amino acid sequence (SEQ ID NO:1):

10	20	30	40	50
MEPAGPAPGR	LGPLLCLLLA	ASCAWSGVAG	EEELQVIQPD	KSVLVAAGET
60	70	80	90	100
ATLRCTATSL	IPVGPIQWFR	GAGPGRELIY	NQKEGHFPRV	TTVSDLTKRN
110	120	130	140	150
NMDFSIRIGN	ITPADAGTYY	CVKFRKGSPD	DVEFKSGAGT	ELSVRAKPSA
160	170	180	190	200
PVVSGPAARA	TPQHTVSTFC	ESHGFSPRDI	TLKWFKNNGNE	LSDFQTNVDP
210	220	230	240	250
VGESVSYSIH	STAKVVLTRE	DVHSQVICEV	AHVTLQGDPL	RGTANLSETI
260	270	280	290	300
RVPPTLEVTQ	QPVRAENQVN	VTCQVRKFYP	QRLQLTWLEN	GNVSRTETAS
310	320	330	340	350
TVTENKDGTY	NWMSWLLVNV	SAHRDDVKLT	CQVEHDGQPA	VSKSHDLKVS
360	370	380	390	400
AHPKEQGSNT	AAENTGSNER	NIYIVVGVVC	TLLVALLMAA	LYLVRIRQKK
410	420	430	440	450
AQGSTSSTRL	HEPEKNAREI	TQDTNDITYA	DLNLPKGKKP	APQAAEPNNH
460	470	480	490	500
TEYASIQTSP	QPASEDTLTY	ADLDMVHLNR	TPKQPAPKPE	PSFSEYASVQ

VPRK

**[0202]** Crystal structure analyses of SIRPA-CD47 complexes resolve the ligand binding site to the variable loops that link the  $\beta$ -sheet strands in the IgV domain of SIRPA. The CD47-binding interface consists of amino acid residues S59-P65, L96-F104, and K123-D130.

**[0203]** Multiple polymorphisms of SIRPA have been identified in humans. An alignment of the amino acid sequences of the two most common variants, referred to as SIRPA v1 and v2, was generated by 2-way blast (**FIG. 1A**). Since most variations in sequence lie beyond the ligand binding site, both SIRPA variants are reported to bind CD47 with similar affinities. Alternatively, another member of the SIRP family, SIRPB1, shares high sequence homology with SIRPA but fails to bind CD47. An alignment of the amino acid sequences of SIRPAv1 and SIRPB1 was generated by 2-way blast (**FIG. 1B**) and shows that the extracellular domain of both proteins (excluding leader sequence) shares ~90% identity. However a single A57M substitution is sufficient to rearrange the S59-P65 ligand-binding interface to prevent SIRPB1 binding to CD47. Furthermore, CD47 binding is highly species-specific with human CD47 recognizing a single allelic variant of mouse SIRPA expressed only by NOD mice. An alignment of the amino acid sequences of human SIRPAv1 and C57BL6 SIRPA was generated by 2-way blast (**FIG. 2**) and shows that the extracellular domain of both proteins (excluding leader sequence) shares ~60% identity.

#### *Anti-SIRPA antibody production*

##### *Immunization procedure*

**[0204]** Rapid prime method: Four 50-day old female BALB/c mice were immunized with using the following procedure. A series of subcutaneous aqueous injections containing human SIRPA antigen but no adjuvant were given over a period of 19 days. Mice were housed in a ventilated rack system from Lab Products. All four mice were euthanized on Day 19 and lymphocytes were harvested for hybridoma cell line generation.

**[0205]** Standard method: Four 50-day old female BALB/c or NZB/W mice were immunized using the following procedure. Mice were housed in a ventilated rack system from Lab Products. Mice were injected intraperitoneally every 3 weeks with a human SIRPA antigen mixed in CpG-ODN adjuvant at 25  $\mu$ g protein antigen per mouse (total volume 125  $\mu$ L per mouse). Test bleeds were done by saphenous vein lancing seven days after the second boost. The test bleed (immune sera) was tested by indirect ELISA assay to determine the best

two responding mice for the fusion. The mice may require a 3rd and 4th boost and another test bleed 7 days after boost to assess titre before fusion. When the antibody titre is high enough the best two responding mice are given a final intravenous boost via lateral tail vein. Four days after the IV boost the mice were euthanized for fusion. The spleens were harvested and lymphocytes isolated from the spleen were used in the fusion process to produce hybridomas.

*Hybridoma development*

[0206] Lymphocytes were isolated and fused with murine SP2/0 myeloma cells in the presence of poly-ethylene glycol (PEG 1500) as per standard Roche Protocol. Fused cells were cultured using a single-step cloning method (HAT selection). This method uses a semi-solid methylcellulose-based HAT selective medium to combine the hybridoma selection and cloning into one step. Single cell-derived hybridomas grow to form monoclonal colonies on the semi-solid media. Ten days after the fusion event, 948 of the resulting hybridoma clones were transferred to 96-well tissue culture plates and grown in HT containing medium until mid-log growth was reached (5 days).

*Hybridoma screening*

[0207] Tissue culture supernatants from the 948 hybridomas were tested by indirect ELISA on screening antigen (Primary Screening) and probed for both IgG and IgM antibodies using a Goat anti-IgG/IgM(H&L)-HRP secondary and developed with TMB substrate. Clones  $>0.2$  OD in this assay were taken to the next round of testing. Positive cultures were retested on screening antigen to confirm secretion and on an irrelevant antigen (Human Transferrin) to eliminate non-specific or “sticky” mAbs and rule out false positives. All clones of interest were isotyping by antibody trapping ELISA to determine if they are IgG or IgM isotype.

*Hybridoma cell culture*

[0208] The hybridoma cell lines of interest were maintained in culture in 24-well culture plates for 32 days post transfer to 96-well plates. This is referred to as the stability period and tests whether clones remain stable and secreting. During this stability period time temporary frozen cell line back up is made of all the clones of interest for -80°C storage (viable 6 months). Hybridomas were periodically tested during this time period for secretion and specificity.

Subcloning

**[0209]** The top hybridoma cell lines (clones) were subcloned to ensure monoclonality. Subcloning was performed by plating parental clones out again using the single-step cloning system. Between 24 and 90 subclones were transferred to 96-well culture plates. Subclones were screened by indirect ELISA and antibody trapping ELISA. The top subclones for each parent were taken for expansion in culture. Any parental clones that were <50% clonal had a second round of subcloning performed.

**[0210]** The antibodies were then screened for SIRPA binding. Antibodies that were positive for binding to human SIRPA were tested for ability to block ligand binding and ability to inhibit ligand-induced SIRPA activity in multiple cell types. The isotype and bin category of each of the antibodies are listed in **Table 1**. In Table 1, “ND” refers to antibodies for which the Bin category has not been determined.

**Table 1: Isotype and epitope bin category for anti-human SIRPA antibodies**

AB ID	Isotype	Bin
3F9	mIgG1	3
9C2	mIgG1	3
8A9	mIgG	3
12D6	mIgG	1
8F4	mIgG	2
1E2	mIgG	2
7H9	mIgG	2
4D8	mIgG	3

Antibody heavy chain and light chain variable domain sequences

**[0211]** Using standard techniques, the amino acid sequences encoding the light chain variable and the heavy chain variable domains of the generated antibodies were determined. The EU or Kabat light chain HVR sequences of the antibodies are set forth in **Table 2-5**. The EU or Kabat light chain HVR sequences of the antibodies are set forth in **Table 2**. The EU or Kabat heavy chain HVR sequences of the antibodies are set forth in **Table 3**. The EU or Kabat light chain framework (FR) sequences of the antibodies are set forth in **Table 4**. The EU or Kabat heavy chain framework (FR) sequences of the antibodies are set forth in **Table 5**.

3F9: Heavy chain variable domain sequence

EVKLVESGGGLVKPGGLSLKLSCAASGFTFSSYAMSWVRQT  
 PEKRLEWVATISDYGGSYTYPDSVKGRFTISRDNAKYTLYLQMSSLRSEDTAL  
 YYCARPPYDDYYGGFAYWGQGTLTVSA (SEQ ID NO:2)

3F9: Light chain variable domain sequence

DIVLTQSPASLAWSLGQRATISCRASKVSSSGYSYMHWY  
 QQKPGQPPKLLIYLASNLESGVPARFSGSGSGTDFTLNIHPVEEEDAATYYCQH  
 NRELPCTFGGGTKLEIK (SEQ ID NO:3)

9C2: Heavy chain variable domain sequence

EFQLQQSGAELVKPGASVKISCKASGYSLTGYNMNWVKQS  
 RGKSLEWIGNINPHYGSSTYNQNFKDATALTVDKSSAAYMQFNSLTSEDSAVY  
 YCAREGYDGVFDYWGQGTTLVSS (SEQ ID NO:4)

9C2: Light chain variable domain sequence

QIVLSQSPAILSASPGEVKVTMTCRASSVSYSYMHWYQQKPG  
 SSPKPWIYVTSNLASGVPTRFSGSGSGTSLTISRVEAEDAATYYCQQWSSNP  
 RTFGGGTKLEIK (SEQ ID NO:5)

8A9: Heavy chain variable domain sequence

QVQLQQPGAAELVKPGASVKMSCKASGYTFTSYWMHWVKQR  
 PGQGLEWIGVIDPSDSYTNYNQKFKGKATLTVDTSSSTAYMQLSSLTSEDSAVY  
 YCTRSGYGKYDFDYWGQGTTLVSS (SEQ ID NO:35)

8A9: Light chain variable domain sequence

DIVLTQSPASLAWSLGQRATISCRASQSVSTSSYSYMHWY  
 QQKPGQPPKLLIYKASNLESGVPARFSGSGSGTDFTLNIHPVEEEDTATYYCQH  
 NWEIPWTFGGGTKLEIK (SEQ ID NO:36)

8F4: Heavy chain variable domain sequence

QIQLVQSGPELKKPGETVKISCKASDYTFDYSMHWVKQA  
 PGKDLKWMGWINTETGEPTYADDFKGRFAFSLEASASTAYLQINNLKNEDTATY  
 FCARHGYPHYFYFDYWGQGTTLVSS (SEQ ID NO:37)

8F4: Light chain variable domain sequence

DIVMTQSQKFMSTSVGDRVSITCKASQNVPTAVAWYQQKP  
 GQSPKALIYLASNRHTGVPDFRTGSGSGTDFTLTITNVQSEDLADYFCLQHWNY  
 PRTFGGGTKLEIK (SEQ ID NO:38)

1E2: Heavy chain variable domain sequence

EVQLVESGGDLVKPGGLSLKLSCAASGFSFSSYAMSWVRQT  
 PAKRLEWVATISGSGGYTYPDSMKGRTISRDNAKDILYLQMSSLRSEDTAMY  
 YCARDPRYTTLYAMDYWGQGTSVTVSS (SEQ ID NO:39)

1E2: Light chain variable domain sequence

NIMMTQSPSFLAVSAGEKVTMSCKSSQSIFSGSNQKNYLA  
 WYQQKPGQSPKLLIYWASTRESGVPDFRTGSGSGTDFTLTISSVQAEDLAVYYC  
 HQHLSSCTFGGGTKLEIK (SEQ ID NO:40)

*7H9: Heavy chain variable domain sequence*

DVQLQESGPGLVKPQSQLSLTCTVTGFSISRGYDWHWIRH  
 FPGNILEWMGYITYSGISNYNPSLKSRSITHDTSKNHFFRLNSVTAEDTATY  
 YCARGGGAWFTYWGQGTLTVSA (SEQ ID NO:41)

*7H9: Light chain variable domain sequence*

DIVMTQSPATLSVTPGDRVSLSCRASQSISSLHWYHQKS  
 HESPRLLIKYASQSIISGIPSRFSAGGSGSDFTLTINSVEPEDVGVYYCQNGHSL  
 PWTFGGGTKLEIK (SEQ ID NO:42)

*4D8: Heavy chain variable domain sequence*

EVKLEESGGGLVKPGGSMKLSCAASGFTFSDAWMDWVRQS  
 PEKGLEWVAEIRGKTTNYATYYAESVKGRFTISRDDSKSSVYLQMNSFSTEDTG  
 IYYCTRRNWGFAYWGQGTLTVSA (SEQ ID NO:43)

*4D8: Light chain variable domain sequence*

DILLTQSPAILSVPGERVSFSCRASQTIGTSIHWYQQRT  
 NGSPRLLIKYASESISGIPSRFSAGGSGSGTDFTLSINSVESEDIADYYCQQTNSW  
 PLTFGAGTKLELK (SEQ ID NO:44)

**Table 2: EU or Kabat light chain HVR sequences of anti-SIRPA antibodies**

Ab ID	HVR L1	HVR L2	HVR L3
3F9	RASKSVSSSGY SY MH (SEQ ID NO:6)	LASNLES (SEQ ID NO:7)	QHNREL PCT (SEQ ID NO:8)
9C2	RASSSVS-YMH (SEQ ID NO:12)	VTSNLAS (SEQ ID NO:13)	QQWSSNPRT (SEQ ID NO:14)

**Table 3: EU or Kabat heavy chain HVR sequences of anti-SIRPA antibodies**

Ab ID	HVR H1	HVR H2	HVR H3
3F9	GFTFSSYAMS (SEQ ID NO:9)	TISDYGGSYTY (SEQ ID NO:10)	PPYDDYYGGFAY (SEQ ID NO:11)
9C2	GYSLTGYNMN (SEQ ID NO:15)	NINPHYGSST (SEQ ID NO:16)	EGYDGVFDY (SEQ ID NO:17)

**Table 4: EU or Kabat light chain Framework sequences of anti-SIRPA antibodies**

Ab ID	VL FR1	VL FR2	VL FR3	VL FR4
3F9	DIVLTQSPASLAV SLGQRATISC (SEQ ID NO:22)	WYQQKPGQPP KLLIY (SEQ ID NO:23)	GVPARFSGSGSGTD FTLNIHPVEEDAAT YYC (SEQ ID NO:24)	FGGGTKEIK (SEQ ID NO:25)

Ab ID	VL FR1	VL FR2	VL FR3	VL FR4
9C2	QIVLSQSPAILSAS PGEKVTMTC (SEQ ID NO:30)	WYQQKPGSSP KPWIY (SEQ ID NO:31)	GVPTRFSGSGSGTSY SLTISRVEAEDAATY YC (SEQ ID NO:32)	FGGGTKEIK (SEQ ID NO:33)

**Table 5: EU or Kabat heavy chain Framework sequences of anti-SIRPA antibodies**

Ab ID	VH FR1	VH FR2	VH FR3	VH FR4
3F9	EVKLVESGGGLV KPG GSLKLSCAAS (SEQ ID NO:18)	WVRQTPEKRL EWVA (SEQ ID NO:19)	YPDSVKGRFTISRDN AKYTLYLQMSSLRS EDTALYYCAR (SEQ ID NO:20)	WGQGTLVT VSA (SEQ ID NO:21)
9C2	EFQLQQSGAELV KPGASVKISCKAS (SEQ ID NO:26)	WVKQSRGKSL EWIG (SEQ ID NO:27)	YNQNFKDKATLTV DKSSAAYMQFNSL TSEDSAVYYCAR (SEQ ID NO:28)	WGQGTTLT VSS (SEQ ID NO:29)

Example 2: Characterization of anti-SIRPA antibody

**[0212]** Initial characterization of SIRPA antibodies involved screening their ability to bind the human receptor ectopically expressed on the rodent Chinese hamster ovary cell line, henceforth referred to as CHO-huSIRPA, followed by screening on primary human macrophages. Cells were harvested, plated at  $10^5$  cells/well in a 96-well plate, washed, and incubated in 100  $\mu$ l FACS buffer containing Fc blocking reagent and 1.0  $\mu$ g/ml of indicated monoclonal antibody. Cells were then washed twice and incubated in FACS buffer containing APC-conjugated secondary antibody diluted 1:200 for 30 minutes on ice. Cells were washed twice in cold FACS buffer and acquired on a BD FACS Canto. Data analysis and calculation of mean fluorescence intensity (MFI) values or % positive cells was performed with FlowJo (TreeStar) software version 10.0.7.

**[0213]** Several antibodies, 3F9 and 9C2 for example, demonstrated binding to CHO-huSIRPA as indicated by positive SIRPA antibody staining detected via FACS analysis (black outlined histograms) (**FIG. 3A**). The negative isotype control (not shown) did not bind to cells. Likewise, 3F9 and 9C2 did not bind to CHO cells highly overexpressing mouse SIRPA (CHO-mSIRPA) (**FIG. 3A**, shaded histograms) confirming the specificity of the antibodies to the human antigen. Importantly, 3F9 and 9C2 also bound to primary human macrophages (**FIG. 3B**), the principal target cell population for in vivo efficacy. MFI values

for cell lines bound by SIRPA antibodies are graphed in **Fig. 3A** and listed on **Table 6**, and typically show MFI values >100-fold over background levels.

**Table 6: MFI values of anti-huSIRPA antibodies binding to cell surface receptor listed as fold over background**

AB ID	CHO-HuSIRPa	CHO-MuSIRPa
mIgG	1	1
1B3	129.9542	0.985036
3F9	136.6835	0.998266
9C2	127.6125	0.981093
9C5	89.64852	0.98305
12D6	128.3982	0.979942
1H11	149.9567	0.972255

**[0214]** Antigen affinity measurements for 3F9 and 9C2 were acquired with standard surface plasmon resonance (SPR) techniques (**FIG. 3C**). Binding studies were performed using a Biacore T200 (GE). An anti-mouse IgG capture antibody was amine coupled to a CM5 sensor chip using standard NHS/EDC activation. SIRPA antibodies were diluted to 50 nM in 1x HBS-EP+ running buffer and captured onto sensor chip surface. Serial dilutions of recombinant soluble human SIRPA antigen were injected over captured SIRPA antibodies to record sensorgram traces. Data were processed by subtracting RU values from the reference cell as well as the buffer injections. Binding curves were globally fit to a 1:1 interaction model to yield kinetic constants listed on **Table 7**. 3F9 and 9C2 bound to monomeric human SIRPA antigen with a  $K_D$  of  $1.0 \times 10^{-8}$  and  $8.0 \times 10^{-8}$  M, respectively.

**Table 7: Association rates, dissociation rates, and equilibrium binding constants of anti-huSIRPA antibodies**

AB ID	$k_{on}$	$k_{off}$	$K_D$
3F9	$5.5 \times 10^4 \text{ (Ms)}^{-1}$	$5.7 \times 10^{-4} \text{ s}^{-1}$	10 nM
9C2	$5.4 \times 10^4 \text{ (Ms)}^{-1}$	$4.5 \times 10^{-3} \text{ s}^{-1}$	80 nM

**[0215]** Cell-based affinity measurements were also performed to ascertain the apparent affinities of 3F9 and 9C2 to cell surface antigen. Serial dilutions of monoclonal antibodies were added to  $10^5$  CHO-huSIRPA cells and allowed to achieve binding equilibrium at 4°C. After addition of fluorescently labeled secondary antibody and brief washing steps, MFI values as a function of titrated antibody concentration was recorded via FACS analysis (**FIG. 3D**). Curves were fit using nonlinear regression analysis with Graphpad Prism 6 software.

Cell-based titration experiments with 3F9 and 9C2 yielded EC50 values of 2.6 nM and 1.6 nM, respectively.

Example 3: Identifying CD47-blocking and non-blocking SIRPA antibodies

[0216] Given the role of the SIRPA—CD47 pathway in suppressing phagocytic cell effector functions, all antagonistic therapies described to date rely on competitive inhibition to block receptor-ligand interaction. Similarly, SIRPA antibodies in this application were screened for their ability to block CD47 binding to CHO-huSIRPA. Cells were harvested, plated at  $10^5$  cells/well in a 96-well plate, washed, and incubated in 100  $\mu$ l FACS buffer containing 1.0  $\mu$ g/ml of indicated monoclonal antibody or isotype control. Cells were then washed and incubated in FACS buffer containing 250 nM His-tagged, soluble human CD47 for 30 minutes on ice. Cells were washed again and stained with PE-conjugated anti-His tag monoclonal antibody to detect surface bound CD47. Data analysis and calculation of MFI values or % positive cells was performed with FlowJo (TreeStar) software version 10.0.7.

[0217] As shown in **FIG. 4A**, soluble CD47 specifically bound CHO-huSIRPA cells as indicated by positive PE-staining via FACS analysis (black outlined histograms). In the absence of CD47-His, anti-His tag antibody failed to bind cells (shaded histograms). When CHO-huSIRPA cells were pre-incubated with indicated SIRPA antibodies, several clones, for example 12D6 and 1B3, exhibited near complete blockade of soluble CD47 binding (dashed line histograms). However, 3F9 and 9C2 represent two unique clones that do not inhibit soluble CD47 binding to CHO-huSIRPA cells. MFI values for cells bound by soluble CD47 are graphed as fold-over-background in **Fig. 4B**, and confirm that 3F9 and 9C2 do not interfere with CD47 interaction.

Example 4: SIRPA antibodies modulate SIRPA-dependent gene expression

[0218] In addition to ligand blockade, SIRPA antibodies were also screened for ability to inhibit CD47-induced gene expression using a luciferase reporter gene under the control of an NFAT (nuclear factor of activated T-cells) promoter. The cell line BW5147.G.1.4 (ATCC® TIB48™), derived from mouse thymus lymphoma T lymphocytes, was infected with Cignal Lenti NFAT-luciferase virus (Qiagen) and a lentivirus expressing human SIRPA-DAP12 chimera, in which the intracellular ITIM motif of SIRPA was substituted with the intracellular ITAM motif of DAP12. Soluble human CD47 protein was serially diluted in PBS and adsorbed onto tissue culture plates. After washing,  $10^5$  NFAT-luciferase reporter cells expressing the huSIRPA/DAP12 chimera (BWZ-huSIRPA) were seeded onto plates and

incubated overnight at 37C. Luciferase activity was measured by adding OneGlo Reagent (Promega) to each well and incubating samples for 3 min at room temperature on a plate shaker. The luminescence signal was quantified using a BioTek Synergy™ Microplate Reader using GEN5™ 2.04 software.

[0219] As shown in **FIG. 5A**, plate-bound human CD47 induced luciferase activity in reporter cells expressing chimeric human SIRPA/DAP12 in a dose-dependent fashion. Importantly, the parental BWZ reporter cells, which lack SIRPA/DAP12 expression, did not emit a luminescence signal in response to CD47 verifying that the chimeric receptor mimics the signaling events initiated through ligand binding. Next, anti-SIRPA antibodies were assessed for their ability to block CD47-dependent luciferase activity in BWZ-huSIRPA reporter cells. As described above, soluble human CD47 protein was diluted in PBS and adsorbed onto 96-well tissue culture plates. After washing,  $10^5$  BWZ-huSIRPA reporter cells were seeded onto plates with either isotype control antibody or the indicated anti-SIRPA antibody and incubated overnight at 37C. **FIG. 5B** demonstrates that, in accordance with the CD47 binding assays described previously, anti-SIRPA antibodies that block CD47 binding to CHO-huSIRPA cells, such as 12D6 and 5F7, also inhibit CD47-dependent luciferase activity in reporter cells. Likewise, anti-SIRPA antibodies that do not block CD47 binding to CHO-huSIRPA cells, such as 9C2 and 3F9, also do not inhibit CD47-dependent luciferase activity in BWZ-huSIRPA cells. Furthermore, anti-SIRPA antibodies do not induce signaling in solution since reporter cells incubated with soluble SIRPA antibodies do not emit luminescence signal in the absence of plate-bound CD47.

#### Example 5: Identification of SIRPA-specific antibodies

[0220] The initial characterization of SIRPA antibodies identified a class of CD47-blocking and non-blocking antibodies capable of binding primary human myeloid cells. However, given the high sequence homology between SIRP $\alpha$  and SIRP $\beta$ 1 (~90% identity), SIRPA-specific binding remains a critical feature of an ideal anti-SIRPA lead antibody. In order to screen SIRPA antibodies for SIRP $\beta$ 1 cross-reactivity, BWZ-NFAT/luciferase reporter cells were transduced with a lentivirus expressing human SIRP $\beta$ 1. Unlike SIRP $\alpha$ , SIRP $\beta$ 1 requires co-expression of DAP12 adaptor for full cell surface localization. As a result, BWZ-huSIRP $\beta$ 1 cells were also transduced with a lentivirus separately expressing human DAP12. To test luciferase activation, selected SIRPA antibodies or isotype control were diluted in PBS at 10  $\mu$ g/mL and adsorbed onto tissue culture plates. After washing,  $10^5$  NFAT-luciferase reporter cells expressing either huSIRPA/DAP12 chimera (BWZ-huSIRPA) or

huSIRP $\beta$ 1 + DAP12 (BWZ-huSIRP $\beta$ 1) were seeded onto plates and incubated overnight at 37C. Luciferase activity was measured by adding OneGlo Reagent (Promega) to each well and incubating samples for 3 min at room temperature on a plate shaker. The luminescence signal was quantified using a BioTek Synergy<sup>TM</sup> Microplate Reader using GEN5<sup>TM</sup> 2.04 software.

[0221] As shown in **FIG. 6A**, plate-bound SIRPA antibodies induced luciferase activity in reporter cells expressing chimeric human SIRPA/DAP12 to a similar extent as previously observed with plate-bound CD47. However, most SIRPA antibodies also induced luciferase activity in BWZ-huSIRP $\beta$ 1 reporter cells indicating that these antibodies cross-react with both SIRP $\alpha$  and SIRP $\beta$ 1. Interestingly, two antibody clones, 3F9 and 9C2, specifically activated BWZ-huSIRPA cells but not BWZ- huSIRP $\beta$ 1 suggesting that these 2 clones represent unique SIRPA-specific antibodies. To confirm this observation, we performed SPR-based binding studies with Biacore T200 (GE). An anti-mouse IgG capture antibody was amine coupled to a CM5 sensor chip using standard NHS/EDC activation. SIRPA antibodies, either 3F9 or 9C2, were diluted to 50 nM in 1x HBS-EP+ running buffer and captured onto sensor chip surface. Equimolar concentrations of recombinant soluble human SIRPA antigen or human SIRPB1 antigen were injected over captured SIRPA antibodies to record sensorgram traces. Data were processed by subtracting RU values from the reference cell as well as the buffer injections. The sensorgrams in **FIG. 6B** clearly show an increase in response units following injection of SIRPA antigen over captured antibodies 3F9 and 9C2. In contrast, flowing SIRPB1 antigen over captured antibodies barely records a binding response above background. Thus, the results from **FIG. 6A and 6B** identify clones 3F9 and 9C2 as SIRPA-specific antibodies.

Example 6: SIRPA-specific antibodies decrease cell surface expression of SIRP $\alpha$  in human macrophages

[0222] It is frequently observed that antibodies targeting certain ITIM/ITAM receptors expressed on the surface of immune cells can reduce the surface levels of said receptor on monocytes, macrophages, dendritic cells, neutrophils, and/or microglia.

[0223] The ability of anti-SIRPA antibodies to reduce cell surface expression of SIRP $\alpha$  was evaluated on primary human macrophages (huMacs). Human monocytes were isolated from peripheral blood of healthy donors and differentiated into macrophages in vitro. Following differentiation, 10<sup>5</sup> huMacs were harvested and seeded onto 96-well tissue culture plates with

either 1-5  $\mu$ g/ml of isotype control or soluble anti-SIRPA antibodies. Cells were analyzed by flow cytometry for SIRP $\alpha$  surface expression following 4 hr treatment or overnight incubation. SIRP $\alpha$  expression was detected using a DyLight650-conjugated anti-human SIRPA antibody belonging to a separate epitope bin than 9C2 and 3F9.

[0224] As shown in **FIG. 7A**, the SIRPA-specific antibodies, 3F9 and 9C2, significantly reduce SIRP $\alpha$  expression by ~90% relative to isotype control-treated macrophages. FACS analysis reveals that receptor down-regulation occurs within hours after antibody addition and is sustained through overnight treatment. This is in contrast to the CD47-blocking antibodies, for example 1B3 or 3D2, which only reduced receptor expression by 50% or less. Since antibody clones 3F9 and 9C2 are also CD47-non-blocking antibodies, other CD47-non-blocking antibodies were screened for receptor down-regulation. **FIG. 7B** shows that, in most cases, CD47-non-blocking antibodies as a class significantly reduced SIRP $\alpha$  expression by ~90% or more. Again, consistent with previous observations, CD47-blocking antibodies, in this example 5F7 and 12D6, were less effective at receptor downregulation by comparison. Thus, **FIG. 7A and 7B** establishes the downregulation of SIRP $\alpha$  as a defining characteristic of the non-ligand blocking SIRPA antibodies. By reducing receptor expression, these antibodies may antagonize the SIRP $\alpha$ —CD47 signaling pathway through non-competitive inhibition, a novel mechanism not previously explored in the field.

Example 7: SIRP $\alpha$  down-regulation enhances phagocytosis of tumor cells by human macrophages

[0225] Tumor cells evade immune surveillance through the upregulation of CD47 thereby transmitting an inhibitory signal to phagocytic cells. Antagonistic antibodies therefore counteract this inhibition to enhance tumor cell phagocytosis. In order to determine if SIRPA antibodies effectively inhibit SIRP $\alpha$  signaling by receptor downregulation, a tumor cell phagocytosis assay was developed based on the acquisition of pHrodo fluorescence. Red Avidin (Invitrogen) is a streptavidin molecule conjugated with pHrodo Red dye, a fluorogenic marker that acquires fluorescence in acidic environments, such as the phagosome. For target tumor cell labeling, 500 nM Red Avidin was mixed with 15 nM biotinylated Lens Culinaris Agglutinin (LCA; Vector Labs). Red Avidin-LCA complexes were then mixed in a 1:1 volumetric ratio with 250,000 Raji cells in serum-free RPMI media on ice. The sugar-binding properties of LCA links Red Avidin to carbohydrate structures on the tumor cell surface. After brief washing steps, Red Avidin-LCA-labeled Raji cells were mixed with monocyte-derived human macrophages in serum-free RPMI media and incubated

at 37°C for 2 hours. Macrophages were then collected and stained on ice with anti-CD14 APC in FACS buffer containing Fc $\gamma$ R-blocking antibodies. Phagocytic activity was measured by counting percent of APC/pHrodo-double positive macrophages. As a control, unlabeled Raji cells were mixed with macrophages to establish background fluorescence.

[0226] **FIG. 8A(i-ii)** establishes the validity of this assay. Monocyte-derived macrophages were seeded onto 96-well tissue culture plates at 10<sup>5</sup> cells/well and treated overnight with isotype control antibody. The following day, 250,000 Red Avidin-labeled Raji cells or unlabeled Raji cells were mixed with macrophages for 2 hours and subsequently analyzed by flow cytometry. The histograms in **FIG. 8A(i)** demonstrate the shift in pHrodo-fluorescence observed when macrophages are co-cultured with Red Avidin-labeled Raji cells (solid black outlined histogram) compared to unlabeled cells (shaded histogram). However, this shifted population only represents ~5% of total CD14<sup>+</sup> macrophages (**FIG. 8Aii**). Opsonization of Red Avidin-labeled Raji cells with an anti-CD20 antibody (Rituximab) shifts pHrodo<sup>+</sup> macrophage population even further (dashed outline histogram) consistent with antibody-dependent phagocytosis enhancing tumor cell clearance. As a result of adding Rituximab, pHrodo<sup>+</sup> macrophages represent ~20% of total CD14<sup>+</sup> macrophages, a nearly 4-fold increase in phagocytic activity.

[0227] To test SIRPA antibodies, macrophages were treated overnight with the indicated candidate antibody or isotype control. The next day, labeled Raji cells were added to treated macrophages followed by quantification of phagocytic activity. As shown in **FIG. 8B**, both 3F9 and 9C2 increased the population of CD14<sup>+</sup>/pHrodo<sup>+</sup>-macrophages 2.5-fold and 1.5-fold, respectively, over isotype treated macrophages. Combination therapy, in which rituximab-opsonized Raji cells were added to 3F9- or 9C2-treated macrophages, further enhanced tumor cell engulfment relative to isotype-treated macrophages. Whereas rituximab alone increased phagocytic activity ~4-fold over untreated cells, rituximab + 3F9 or rituximab + 9C2 treatment augmented phagocytosis 7-fold and 6-fold, respectively. Since 3F9 and 9C2 are SIRPA-specific antibodies that do not competitively inhibit CD47 binding, **FIG. 8C** compares the phagocytic activity of macrophages treated with CD47-blocking versus CD47-non-blocking antibodies. Among the CD47-blocking antibodies, only 12D6 and 5F7 significantly increased tumor cell uptake by ~30-40% above isotype treated macrophages. By comparison, phagocytic activity of 3F9-treated macrophages increased 2-fold. Thus the results from **FIG. 8A-C** establish that antibody-mediated downregulation of SIRPA on macrophages enhances phagocytic uptake of tumor cells. Combining SIRPA antibodies with

anti-tumor antigen antibodies further potentiates tumor cell clearance by effector cells.

Finally, in comparison to anti-SIRPA antibodies that competitively inhibit CD47 interaction, antibodies that non-competitively inhibit CD47 binding by reducing SIRP $\alpha$  expression demonstrate a superior capacity to stimulate engulfment of tumor cells by macrophages.

Example 8: SIRP $\alpha$  down-regulation activates primary human monocytes

**[0228]** Though macrophages may be the principal effector cell population driving tumor cell clearance in response to anti-SIRPA therapy, SIRPA antibodies will engage multiple myeloid cell lineages expressing SIRP $\alpha$ . Among these cells are monocytes, which populate the peripheral blood, and thus, are easily accessible to assay target engagement upon antibody administration in vivo. In order to identify potential biomarkers, primary monocytes were isolated from peripheral blood of healthy donors and assayed for activation markers following antibody treatment.

**[0229]** The ability of anti-SIRPA antibodies to reduce surface expression of SIRP $\alpha$  was verified on monocytes. Following isolation,  $10^5$  monocytes were seeded onto 96-well tissue culture plates with either 5  $\mu$ g/ml of isotype control or soluble anti-SIRPA antibodies. Cells were analyzed by flow cytometry for SIRP $\alpha$  surface expression after overnight incubation. SIRP $\alpha$  expression was detected using a DyLight650-conjugated anti-human SIRPA antibody belonging to a distinct epitope bin. **FIG. 9A** shows that 3F9 reduces surface expression of SIRP $\alpha$  by 50% relative to isotype control treated cells. Though receptor downregulation appears less robust in monocytes than previously observed in macrophages, monocytes were assayed for production of inflammatory mediators, for example production of reactive oxygen species (ROS) and pro-inflammatory cytokines. To detect ROS production,  $10^5$  monocytes were seeded onto 96-well tissue culture plates with either 10  $\mu$ g/ml of isotype control or soluble anti-SIRPA antibodies. Subsequently, cells were labeled with 2  $\mu$ M of the fluorescent dye, CM-H2DCFDA. Following 1 hour of antibody-mediated stimulation at 37°C, the relative fluorescence units in cells were measured at excitation wavelength 495 nm and emission wavelength 530 nm. Specific fluorescence index of stimulated cells was obtained by subtraction of background fluorescence of labeled cells incubated in medium alone and/or with isotype control antibody. Plates were read with a BioTek Synergy™ Microplate Reader using GEN5™ 2.04 software. **FIG. 9B** shows that SIRPA-specific antibodies, 3F9 and 9C2, stimulated ROS production in monocytes isolated from 2 healthy donors. Additionally, **FIG. 9C** shows that  $10^5$  monocytes treated overnight with SIRP $\alpha$ -downregulating antibodies produces elevated amounts of IL-8. Thus, the results from **FIG.**

9A-C suggest that, in addition to reducing receptor surface expression, anti-SIRPA antibodies may also polarize cells towards a more active phenotype.

Example 9: SIRPA-specific antibodies decrease cell surface expression of SIRPa *in vivo*

[0230] In order to determine if anti-SIRPA antibodies reduce cell surface expression of the receptor in *in vivo* model systems, human BAC transgenic mice encoding the human SIRPA gene in a RAG2-deficient and IL2R  $\gamma$  chain-deficient background were obtained. The expression level of huSIRPA was analyzed on mouse myeloid cells by flow cytometry. As shown in **FIG. 10A**, monocytes and granulocytes isolated from mouse peripheral blood expressed human SIRPA, as well as endogenous mouse SIRPA. Macrophages and dendritic cells derived from bone marrow cells also express huSIRPA. Thus, the huSIRPA-tg mice faithfully recapitulate the expression pattern of human SIRPA in mouse cells. Furthermore, to determine if huSIRPA retained its inhibitory function, huSIRPA-tg mice were implanted with Raji cells, a human B cell lymphoma cell line that overexpresses human CD47. As shown in **FIG. 10B**, subcutaneous administration of Raji cells results in solid tumor formation suggesting that huSIRPA-tg mice support engraftment of CD47+ human cells.

[0231] To test antibody-mediated receptor downregulation *in vivo*, huSIRPA-tg mice received a single intraperitoneal (i.p.) injection of 10 mg/kg of 3F9 (anti-SIRPA antibody) or MOPC21 (mouse IgG1 isotype control). The following day, blood samples were drawn from mice into heparin-coated collection tubes and processed for FACS analysis. Additionally, spleens were also harvested and processed for FACS analysis. Briefly, blood and splenocyte samples were incubated for 5 minutes in ACK lysis buffer to lyse red blood cells and then washed extensively with cold PBS. Cells were then resuspended in FACS buffer (PBS +2% FBS + Fc receptor blocking solution). Peripheral blood myeloid cells were stained with anti-mouse CD11b-Pacific Blue and, either, anti-human SIRPa/β-APC (clone SE5A5) or DyLight 650-conjugated 9C2, the human SIRPA-specific antibody identified through hybridoma screens. Data were acquired on a BD FACS CANTO™ II cytometer (Becton Dickinson) and analyzed with FlowJo software. As shown in **FIG. 10C**, gating on CD11b+ blood monocytes and granulocytes labeled with anti-human SIRPa/β-APC reveals that 3F9 treatment fails to decrease cell surface levels of huSIRPA on both cell types when compared to isotype control-treated mice. However, 3F9 treatment blocks 9C2-DyLight 650 from binding huSIRPA on peripheral blood cells. Since 3F9 and 9C2 bind to the same epitope, this blockade verifies that 3F9 occupies the receptor on peripheral blood cells without downregulating expression.

[0232] Single-cell suspensions from mouse spleens were also obtained from isotype control- and 3F9-treated animals. Splenocytes were stained with anti-mouse CD11b-Pacific Blue, anti-mouse F4/80-FITC, and anti-human SIRP $\alpha$ / $\beta$ -APC (clone SE5A5). Data were acquired on a BD FACS CANTO<sup>TM</sup> II cytometer (Becton Dickinson) and analyzed with FlowJo software. As shown in **FIG. 10D**, two major myeloid cell populations were identified in spleens based on F4/80 and CD11b markers: an F4/80<sup>Lo</sup> CD11b<sup>+/−</sup> population (likely red pulp macrophages) and an F4/80<sup>Hi</sup> CD11b<sup>Hi</sup> population. Though both populations express huSIRPA, as demonstrated in control-treated mice, 3F9 treatment downregulated huSIRPA expression primarily in F4/80<sup>Lo</sup> CD11b<sup>+/−</sup> cells. Additionally, an F4/80<sup>Lo</sup> CD11b<sup>−</sup> population expands in the spleens of 3F9-treated mice only. Marginal decrease in huSIRPA is observed in the F4/80<sup>Hi</sup> CD11b<sup>Hi</sup> splenic population.

[0233] These results demonstrated that when utilizing huSIRPA-tg mice, anti-SIRPA antibodies engage huSIRPA *in vivo* and functionally downregulate the receptor on myeloid cells. The results further demonstrate that the huSIRPA antibody, 3F9, engages huSIRPA on peripheral blood cells and splenic myeloid cells, but internalizes the receptor in a cell type-dependent or a context-dependent manner.

Example 10: Anti-tumor effects of anti-SIRPA antibodies in BAC-transgenic mouse models

[0234] Pilot experiments with huSIRPA-tg mice were performed to assess the anti-tumor effects of anti-SIRPA antibodies. Twelve huSIRPA-tg female mice, approximately 8-12 weeks of age, were implanted unilaterally on the right flank with 500,000 Raji-Luciferase cells mixed in Matrigel solution. Tumor engraftment was monitored beginning seven to ten days post-implantation by caliper measurements of tumor volume and bioluminescence imaging. On Day 10, when tumors reached approximately 80-120 mm<sup>3</sup> in volume, mice were administered D-luciferin substrate by i.p. injection and imaged with an *in vivo* imaging system. Mice were subsequently randomized into treatment or control groups (6 mice per group) based on the average radiance (photons/second/cm<sup>2</sup>/sr) values of the luciferase signal from Raji cells. Beginning on Day 10, mice received i.p. injections at 10 mg/kg of either 3F9 (anti-SIRPA) or mouse IgG1 control antibody 2x/week for the duration of the study. Mice were observed daily and weighed twice weekly using a digital scale. The study was concluded when the mean tumor volume of the control group reached 1500 mm<sup>3</sup>. At study termination tumors were harvested and processed for FACS analysis. Briefly, tumor samples were treated with collagenase for 30 min at 37°C. Samples were dissociated through a cell

strainer and resuspended in 2% FBS in PBS. Red blood cells in samples were lysed using ACK lysis buffer and cells were then washed in 2% FBS in PBS. Cells were counted using a hemocytometer and one million cells were stained with fluorochrome-conjugated antibodies for 30 minutes on ice, then washed with 2% FBS in PBS. Cells were fixed with 4% paraformaldehyde in PBS. All the stained cells were analyzed on a FACS Canto (BD Biosciences) and the data analyzed with FlowJo software (TreeStar). Tumor-infiltrating myeloid cells were stained with anti-mouse CD11b-Pacific Blue, anti-mouse F4/80-FITC, and anti-human SIRP $\alpha$ / $\beta$ -APC (clone SE5A5). As shown in **FIG. 11A**, two major myeloid populations were identified based on F4/80 and CD11b markers: an F4/80 $^+$  CD11b $^+$  population (F4/80 $^+$  cells) and an F4/80 $^-$  CD11b $^+$  population (CD11b $^+$  cells). As shown in isotype control-treated mice, both populations express huSIRPA. However, 3F9 treatment downregulated huSIRPA expression only in F4/80 $^-$  CD11b $^+$  cells, whereas huSIRPA expression in F4/80 $^+$  CD11b $^+$  cells was not decreased.

**[0235]** As shown in **FIG. 11B**, administering the anti-SIRPA antibody, 3F9, appeared to inhibit tumor growth *in vivo* compared to vehicle control-treated animals when measuring tumor burden by bioluminescent imaging. Linear regression analysis of average radiance values indicates that a near-significant trend for efficacy emerges at Day 17 (p=0.06) when correcting for pre-treatment radiance values at Day 10. This trend continues in subsequent measurements (with p-values of 0.16, 0.77 and 0.18), but given the variability in tumor growth and limited number of huSIRPA-tg mice available, this study is statistically underpowered to reach desired significance levels.

Example 11: Anti-tumor effects of anti-SIRPA antibodies in humanized mouse models

**[0236]** Immunocompromised female NSG mice (Jax) engrafted with human cord blood-derived CD34 $^+$  hematopoietic stem cells to reconstitute human immune cell lineages, including the myeloid and lymphoid cell compartments, served as a platform to measure the immune modulating ability of anti-SIRPA antibodies. Successful engraftment of mature human immune cells is defined as >25% of huCD45 $^+$  cells in peripheral blood 12 weeks post-injection. Humanized mice were additionally screened for high cell counts of human CD14 $^+$ , human CD11b $^+$ , and human CD33 $^+$  cells in peripheral blood.

**[0237]** For immuno-oncology efficacy studies, humanized mice were implanted subcutaneously on the right flank with MDA-MB-231 cells, a triple-negative human breast cancer cell line responsive to checkpoint inhibitor therapy in this model system. Pre-

treatment tumor volumes were measured by digital calipers when tumors became palpable, and mice were randomized into treatment or control groups (12 mice per group) when tumor volumes reach 60-120 mm<sup>3</sup> on Day -1. Beginning on Day 0, mice received i.p. injections at 40 mg/kg of either 3F9 (anti-SIRPA) or mouse IgG1 control antibody every 4 days for the duration of the study. A third group instead received i.p. injections of pembrolizumab (Keytruda, Merck) at 10 mg/kg every 5 days for the duration of the study. Body weights, clinical observations, and digital caliper measurements were recorded twice weekly post dose initiation. The study was concluded when the mean tumor volume of the control group reached 2000 mm<sup>3</sup>. At termination, blood, spleen, and tumors were harvested and processed for FACS analysis. Briefly, tumor samples were treated with collagenase for 30 min at 37°C. Spleen and tumor samples were dissociated through a cell strainer and resuspended in 2% FBS in PBS. Red blood cells in samples were lysed using ACK lysis buffer and cells were then washed in 2% FBS in PBS and stained with fluorochrome-conjugated antibodies for 30 minutes on ice. Cells were fixed with 4% paraformaldehyde in PBS. All the stained cells were analyzed on a FACS Canto (BD Biosciences) and the data analyzed with FlowJo software (TreeStar).

[0238] As shown in **FIG. 12A**, treatment with the SIRPA antibody, 3F9, reduced cell surface levels of SIRPA in peripheral blood huCD45+ huCD14+ myeloid cells in tumor-bearing humanized mice when compared to either isotype control-treated or Keytruda-treated mice. However, cell surface expression levels of SIRPA was not reduced on intratumoral huCD45+ huCD14+ myeloid cells. These results resemble previous observations in huSIRPA-tg mice in which antibody-mediated receptor downregulation occurred in a cell type-dependent or context-dependent manner.

[0239] As shown in **FIG. 12B**, treatment with the SIRPA antibody, 3F9, reduced the percentage of peripheral blood huCD45+ huCD14+ myeloid cells cells in tumor-bearing humanized mice when compared to either isotype control-treated or Keytruda-treated mice. In contrast, both 3F9 and Keytruda increased the percentage of intratumoral huCD45+ huCD14+ myeloid cells. Furthermore, 3F9 treatment decreased overall percentage of human CD45+ leukocytes in peripheral blood (**FIG.12C**) of tumor-bearing humanized mice when compared to the isotype control group.

[0240] To account for various factors other than treatment modality that influence tumor growth in this model system, a multiple linear regression analysis with R's *lm()* function was

utilized to correct tumor volumes for differences in 1) huCD34+ stem cell donor, 2) tumor volume at Day -1, 3) animal body weight before randomization, and 4) engraftment rate of huCD45+ cells before randomization. **FIG. 13A** plots the mean tumor volumes per group for each time point. Both 3F9 and Keytruda treatment groups significantly reduce tumor volume in early and late time points compared to the isotype control group, though the effects are mostly observed between Days 22 and 28. Graphing the tumor volume measurements by huCD34+ stem cell donor, as shown in **FIG. 13B**, reveals that mice engrafted with human immune cells from donors 5031 and 5048 significantly inhibited tumor growth when treated with either 3F9 or Keytruda compared to isotype control. In contrast, mice engrafted with human immune cells from donor 129 did not record any significant reduction in tumor volume in either treatment group compared to isotype control group. Note, however, that mean tumor volume in the control group from donor 129 recipients was lower than the control group from donors 5031 and 5048. Such donor-to-donor variability in tumor growth underscores the necessity for appropriate controls to adequately interpret results in this platform.

**[0241]** The data presented above establishes that the SIRPA antibody, 3F9, engages the receptor in vivo and induces SIRPA downregulation in specific cell populations. Analysis of both circulating and tumor infiltrating immune cells reveals that 3F9 treatment decreased CD14+ myeloid cells in peripheral blood with a concomitant increase of CD14+ cells in tumors. Unlike Keytruda, which decreased CD4+ and CD8+ T cells in blood and tumors, 3F9 did not significantly impact T cell numbers suggesting that it primarily acts on the myeloid compartment. Importantly, receptor downregulation and changes in myeloid cell populations with 3F9 correlated with significant inhibition of tumor growth comparable to Keytruda therapy. Taken together, these studies support the pre-clinical efficacy of anti-SIRPA antibodies as a therapeutic for treating human cancer.

Example 12: *In silico* antibody humanization of 3F9 and 9C2

**[0242]** Antibody humanization is used to transform antibodies generated in a different species to best resemble a human antibody through sequence and structural relationships in order to prevent immunogenicity in human administration. Antibodies from different species share characteristic sequence and structural features that allow the grafting of the specificity-determining regions (SDRs) of the non-human antibody onto a human antibody framework. This results in retention of the specificity of the non-human antibody. The humanization process involves identification of the non-human antibody sequence and features, including

the framework regions and SDRs. The following criteria are used to humanize an antibody: 1) percent similarity in framework regions between non-human and known human antibodies, 2) length similarity in SDRs between non-human and known human antibodies, 3) genes used to generate the framework regions of the human antibody, and 4) previous use of human antibody frameworks in humanizations and as therapeutics. Similarity in framework regions and SDR lengths are important because differences can generate structural differences in the antibody that can alter the specificity of the antibody. Specific genes used to generate the framework of human antibodies are known to be beneficial or detrimental to the stability or specificity of the antibody and are selectively used or avoided, accordingly. Lastly, previously successful humanization frameworks, including those used in human therapeutics, which are well tolerated with good half-lives, are likely candidates for future successful humanizations.

**[0243]** As shown in **FIG. 14A-D**, humanized light and heavy chain variable region sequences were identified for SIRPA antibodies, 3F9 and 9C2. The first humanized sequence for 3F9 heavy chain variable domain (hSB-3F9-H1; **FIG. 14A**) is a “CDR-swap” with no changes to human framework. The subsequent humanized heavy chain sequence (hSB-3F9-H2) alters framework residues (changes shown in bold compared to sequence above it). In **FIG. 14B**, hSB-3F9-L1 is a “CDR-swap” of the light chain variable domain with no changes to human framework. Subsequent humanized light chain sequences alter framework residues (changes shown in bold compared to sequence above it; gray boxed residues are from a previous version). Light chain CDRs from 3F9 also contain potential deamidation sites (marked with #), which may be substituted with Q, S, A, or D. Additionally, the variable domain for 3F9 contains a potential free Cys at position 96, which may potentially lead to problems during manufacture. This site may be substituted with an A, S, or L residue as long as antigen binding is not altered. In **FIG. 14C**, hSB-9C2-H1 is a “CDR-swap” of the heavy chain variable domain with no changes to human framework. Subsequent humanized heavy chain sequences alter framework residues (changes shown in bold compared to sequence above it; gray boxed residues are from a previous version). Heavy chain CDRs from 9C2 also contain potential deamidation sites (marked with #), which may be substituted with Q, S, or A. 9C2 also contains an Asp-Gly (DG) sequence in CDR-H3 (marked with @), which may be susceptible to isoaspartate formation. This site may be substituted with an A, S, or E residue as long as antigen binding is not altered. In **FIG. 14D**, hSB-9C2-L1 is a “CDR-swap” of the light chain variable domain with no changes to human framework. Subsequent

humanized light chain sequences alter framework residues (changes shown in bold compared to sequence above it; gray boxed residues are from a previous version). Light chain CDRs from 9C2 contain potential deamidation sites (marked with #), which may be substituted with Q, S, D, or A. 9C2 also contains a Trp residue in CDR-L3 (marked with ^), which may be susceptible to oxidation. This site may be substituted with an H, Y, or F residue as long as antigen binding is not altered.

Example 13: Epitope mapping of anti-SIRPA antibody binding sites

[0244] Epitope mapping of anti-SIRPA antibodies is performed using an alanine-scanning library created by shotgun mutagenesis of the human SIRPA cDNA sequence. A SIRPA expression construct encoding a C-terminal V5 epitope tag is subjected to high-throughput alanine scanning mutagenesis (outlined in Davidson and Doranz, 2014 Immunology 143, 13-20) to generate a comprehensive mutation library. Each of the residues representing the SIRPA extracellular domain (amino acids 31-374) is mutated, most to alanine, while alanine codons were mutated to serine.

[0245] The SIRPA mutant library clones, arrayed in a 384-well microplate, are transfected individually into HEK-293T cells and allowed to express for 22 hours. Antibodies are digested to generate Fabs, after which cells are incubated with Fabs diluted in 10% normal goat serum (NGS) (Sigma-Aldrich, St. Louis, MO). Prior to library screening, primary Fab concentrations are determined using an independent immunofluorescence titration curve against cells expressing wild type SIRPA to ensure that signals are within the linear range of detection. Fabs are detected using 7.5 µg/ml AlexaFluor488-conjugated secondary antibody (Jackson ImmunoResearch Laboratories, Westgrove, PA) in 10% NGS. Cells are washed twice with PBS and resuspended in Cellstripper (Cellgro, Manassas, VA) with 0.1% BSA (Sigma-Aldrich, St. Louis, MO). In some cases, higher stringency conditions are used, including increased pH, increased temperature, and increased dissociation time. Mean cellular fluorescence is detected using the Intellicyt high throughput flow cytometer (HTFC, Intellicyt, Albuquerque, NM). Fab reactivities against each mutant clone are calculated relative to wild-type SIRPA protein reactivity by subtracting the signal from mock-transfected controls, and normalizing to the signal from wild-type SIRPA transfected controls.

[0246] Mutated residues within library clones are identified as “critical” to the Fab binding epitope if they do not support reactivity of the test Fab but do support reactivity of

commercially available reference antibody, MAB4546 (R&D Systems), or additional anti-SIRPA Fabs. This counter-screen strategy facilitates the exclusion of SIRPA mutants that are locally misfolded or that have an expression defect.

Example 14: Fc $\gamma$ RIIB downregulation by anti-SIRPA antibodies

[0247] In addition to the target antigen of interest, cells of the myeloid lineage also express multiple Fc receptors capable of binding to the Fc domain of therapeutic antibodies. The Fc $\gamma$  receptors (Fc $\gamma$ R) constitute the best characterized and most potent receptor class to mediate Fc-dependent effector functions. Fc $\gamma$ Rs include both ITAM-associated activating receptors (Fc $\gamma$ RI, Fc $\gamma$ RIIA, and Fc $\gamma$ RIIIA) and an ITIM-bearing inhibitory receptor (Fc $\gamma$ RIIB), and co-expression of activating/inhibitory receptors on the same cell establishes a threshold for cellular activation. In general, ligation of activating Fc $\gamma$ Rs by immune complexes initiates several signaling cascades that lead to cellular activation and subsequent induction of effector functions. These activities vary between myeloid cell types, but may include antibody-dependent cellular cytotoxicity, antibody-dependent cellular phagocytosis, and upregulation of several pro-inflammatory cytokines and chemokines, etc. In contrast, ligation of the inhibitory receptor, Fc $\gamma$ RIIB, by immune complexes counteracts the immunostimulatory signals of activating Fc $\gamma$ Rs supporting the maintenance of tissue homeostasis. For example, several studies establish that genetic knockout of Fc $\gamma$ RIIB results in enhanced pro-inflammatory macrophage activity in murine models of immune complex-mediated inflammation. Since Fc $\gamma$ RIIB is the only Fc $\gamma$ R with inhibitory activity, it plays a central role in regulating Fc $\gamma$ R-mediated inflammation by myeloid cells. In the context of the tumor microenvironment, Fc $\gamma$ RIIB expression levels may determine the polarization state of tumor-associated macrophages and the regulation of macrophage effector function *in vivo*.

[0248] To assess whether Fc $\gamma$ Rs participate in the *in vitro* activity of anti-SIRPA antibodies, antibody 3F9, was treated with EndoS (New England Biolabs) to remove the Fc-linked glycan. The enzymatic reaction completely cleaved the carbohydrate structure as shown by the LCA blot in **FIG. 15A**, which detects the mannose residues on the Fc glycan. Importantly, the deglycosylation reaction did not impact antigen recognition as both 3F9 and deglycosylated 3F9 bound SIRPA comparably in cell-based binding assays (**FIG. 15B**). Subsequently, the ability of deglycosylated 3F9 to reduce cell surface expression of SIRPA on primary human macrophages (huMacs) was compared to glycosylated 3F9. Briefly, human monocytes were isolated from peripheral blood of two healthy donors (HD 1 and HD 2) and differentiated into macrophages *in vitro*. Following differentiation, 10<sup>5</sup> huMacs were

harvested and seeded onto 96-well tissue culture plates with increasing concentrations of anti-SIRPA antibodies. Cells were analyzed by flow cytometry for SIRPA surface expression following overnight incubation. Receptor expression was detected using a DyLight650-conjugated anti-human SIRPA antibody belonging to a separate epitope bin than 9C2 and 3F9.

**[0249]** As shown in **FIG. 16**, both glycoforms of 3F9 significantly downregulated surface expression of SIRPA relative to isotype control-treated macrophages. However, in both donors, the deglycosylated 3F9 variant exhibited partially reduced activity compared to the glycosylated antibody. For example, 3F9 downregulated SIRPA expression by as much as 90% and 85% in HD 1 and HD 2, respectively; whereas deglycosylated 3F9 only achieved 70% and 75% receptor downregulation in the same donor macrophages, respectively. This finding suggests that anti-SIRPA antibodies such as 3F9 need Fc $\gamma$ R engagement for maximal activity.

**[0250]** To determine which Fc $\gamma$ R contributes to the *in vitro* activity of 3F9, monocyte-derived macrophages obtained from two healthy donors were treated overnight with either isotype control antibody or anti-SIRPA antibody, 3F9, and assessed for surface expression levels of Fc $\gamma$ RIIA (CD16) and Fc $\gamma$ RIIA/B (CD32A/B). As shown in **FIG. 17A**, 3F9 treatment moderately reduced surface expression of Fc $\gamma$ RIIA relative to isotype control-treated macrophages. In contrast, substantial downregulation of Fc $\gamma$ RIIA/B was evident on 3F9-treated macrophages relative to isotype control-treated cells (**FIG. 17B**).

**[0251]** Since the detection antibody used to measure surface levels of Fc $\gamma$ RII (clone FUN-2; Biolegend) does not distinguish the activating receptor (Fc $\gamma$ RIIA) from the inhibitory receptor (Fc $\gamma$ RIIB), this assay was repeated with receptor-specific antibodies. As previously described, monocyte-derived macrophages obtained from two healthy donors were treated overnight with either isotype control antibody or the indicated glycoform of 3F9. **FIG. 18** shows that 3F9 significantly downregulated Fc $\gamma$ RIIA in macrophages by ~70-85% relative to isotype control-treated cells. This effect was dependent on the Fc domain since deglycosylation of the antibody abrogated receptor downregulation. However, when assessing surface expression of Fc $\gamma$ RIIB, 3F9 treatment reduced expression of the inhibitory receptor to near undetectable levels relative to isotype control-treated macrophages (**FIG. 18**). Even the deglycosylated form of 3F9 exhibited robust downregulation of Fc $\gamma$ RIIB suggesting that the murine IgG1 isoform of 3F9 may preferentially associate with human

Fc $\gamma$ RIIB. Not to be bound by theory, by targeting two ITIM-bearing receptors for downregulation (SIRPA and Fc $\gamma$ RIIB), 3F9 may polarize macrophages towards an activated phenotype. In the context of tumor biology, reprogramming tumor-associated macrophages in the tumor microenvironment from a pro-tumor phenotype towards an anti-tumor phenotype with an anti-SIRPA antibody thus represents a promising mode of cancer immunotherapy.

**[0252]** All patents, patent applications, accession numbers, and other published reference materials cited in this specification are hereby incorporated herein by reference in their entirety for their disclosures of the subject matter in whose connection they are cited herein.

**WHAT IS CLAIMED IS:**

1. An isolated anti-signal regulatory protein  $\alpha$  (SIRPA) antibody that selectively binds SIRPA and down-regulates SIRPA expressed on the cell surface.
2. The isolated anti-SIRPA antibody of claim 1, wherein the anti-SIRPA antibody binds one or more polymorphic variants of human SIRPA.
3. The anti-SIRPA antibody of claim 1 or 2, wherein the anti-SIRPA antibody decreases cell surface levels of SIRPA, decreases intracellular levels of SIRPA, decreases total levels of SIRPA, or any combination thereof.
4. The anti-SIRPA antibody of claim 1 to 3, wherein the anti-SIRPA antibody induces SIRPA degradation, SIRPA cleavage, SIRPA internalization, SIRPA shedding, downregulation of SIRPA expression, or any combination thereof.
5. The anti-SIRPA antibody of any one of claims 1 to 4, wherein the antibody decreases cellular levels of SIRPA *in vivo*.
6. The anti-SIRPA antibody of any one of claims 1 to 5, wherein the anti-SIRPA antibody inhibits cell surface clustering of SIRPA.
7. The anti-SIRPA antibody of any one of claims 1 to 6, wherein the anti-SIRPA antibody inhibits one or more SIRPA activities.
8. The anti-SIRPA antibody of any one of claims 1 to 7, wherein the antibody counteracts one or more SIRPA activities that are selected from the group consisting of:
  - (a) SIRPA binding to one or more SIRPA ligands, optionally wherein the one or more SIRPA ligands are selected from the group consisting of CD47, surfactant protein A and D and any combination thereof;
  - (b) decreasing proliferation of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia;

(c) inhibiting migration of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia;

(d) inhibiting one or more functions of one or more cells selected from the group consisting of dendritic cells, bone marrow-derived dendritic cells, macrophages, neutrophils, NK cells, M1 macrophages, M1 neutrophils, M1 NK cells, activated M1 macrophages, activated M1 neutrophils, activated M1 NK cells, M2 macrophages, M2 neutrophils, M2 NK cells, monocytes, osteoclasts, T cells, T helper cells, cytotoxic T cells, granulocytes, neutrophils, microglia, M1 microglia, activated M1 microglia, and M2 microglia;

(e) inhibition of one or more types of clearance selected from the group consisting of apoptotic neuron clearance, nerve tissue debris clearance, dysfunctional synapse clearance, non-nerve tissue debris clearance, bacteria clearance, other foreign body clearance, disease-causing protein clearance, disease-causing peptide clearance, and tumor cell clearance; optionally wherein the disease-causing protein is selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides and the tumor cell is from a cancer selected from the group consisting of bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal cell cancer, renal pelvis cancer, leukemia, lung cancer, melanoma, non-Hodgkin's lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, fibrosarcoma, and thyroid cancer;

- (f) inhibition of tumor cell killing by one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells;
- (g) inhibiting anti-tumor cell proliferation activity of one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells;
- (h) modulated expression of one or more inflammatory receptors, optionally wherein the one or more inflammatory receptors comprise CD86 and the one or more inflammatory receptors are expressed on one or more of microglia, macrophages, neutrophils, NK cells, dendritic cells, bone marrow-derived dendritic cells, neutrophils, T cells, T helper cells, or cytotoxic T cells;
- (i) promoting or rescuing functionality of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid-derived suppressor cells, tumor-associated macrophages, tumor-associated neutrophils, tumor-associated NK cells, and regulatory T cells;
- (j) increasing infiltration of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid-derived suppressor cells, tumor-associated macrophages, tumor-associated neutrophils, tumor-associated NK cells, non-tumorigenic CD45+CD14+ myeloid cells, and regulatory T cells into tumors;
- (k) increasing the number of tumor-promoting myeloid/granulocytic immune-suppressive cells and/or non-tumorigenic CD45+CD14+ myeloid cells in a tumor, in peripheral blood, or other lymphoid organ;
- (l) enhancing tumor-promoting activity of myeloid-derived suppressor cells and/or non-tumorigenic CD45+CD14+ myeloid cells;
- (m) enhancing survival of non-tumorigenic myeloid-derived suppressor cells and/or non-tumorigenic CD45+CD14+ myeloid cells;
- (n) decreasing activation of tumor-specific T lymphocytes with tumor killing potential;
- (o) decreasing infiltration of tumor-specific NK cells with tumor killing potential;
- (p) increasing tumor volume;
- (q) increasing tumor growth rate; and

(r) decreasing efficacy of one or more immune-therapies that modulate anti-tumor T cell responses, optionally wherein the one or more immune-therapies are immune-therapies that target one or more target proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, TREM1, TREM2, CD39, CD73, CSF-1 receptor, and any combination thereof, or of one or more cancer vaccines.

9. The anti-SIRPA antibody of any one of claims 1 to 8, wherein the antibody induces one or more of the activities that are selected from the group consisting of:

- (a) increasing the number of tumor infiltrating CD3+ T cells;
- (b) decreasing cellular levels of SIRPA in non-tumorigenic CD14+myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood;
- (c) reducing the number of non-tumorigenic CD14+ myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood;
- (d) reducing PD-L1 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (e) reducing PD-L2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (f) reducing B7-H2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (g) reducing B7-H3 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (h) reducing CD200R levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (i) reducing CD163 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (j) reducing CD206 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (k) decreasing tumor growth rate of solid tumors;
- (l) reducing tumor volume;

- (m) increasing efficacy of one or more PD-1 inhibitors;
- (n) increasing efficacy of one or more checkpoint inhibitor therapies and/or immune-modulating therapies, optionally wherein the one or more checkpoint inhibitor therapies and/or immune-modulating therapies target one or more of CTLA4, the adenosine pathway, PD-L1, PD-L2, OX40, TIM3, LAG3, or any combination thereof;
- (o) increasing efficacy of one or more chemotherapy agents, optionally wherein the one or more of the chemotherapy agents are gemcitabine, capecitabine, anthracyclines, doxorubicin (Adriamycin®), epirubicin (Ellence®), taxanes, paclitaxel (Taxol®), docetaxel (Taxotere®), 5-fluorouracil (5-FU), cyclophosphamide (Cytoxan®), carboplatin (Paraplatin®), and any combination thereof;
- (p) increasing proliferation of T cells in the presence of non-tumorigenic myeloid-derived suppressor cells (MDSC);
  - (1) inhibiting differentiation, survival, and/or one or more functions of non-tumorigenic myeloid-derived suppressor cells (MDSC); and
- (r) killing CD33-expressing immunosuppressor non-tumorigenic myeloid cells and/or non-tumorigenic CD14-expressing cells in solid tumors and associated blood vessels when conjugated to a chemical or radioactive toxin.

10. The anti-SIRPA antibody of any one of claims 1 to 9 wherein the antibody inhibits interaction between SIRPA and one or more SIRPA ligands.

11. The anti-SIRPA antibody of claim 10, wherein the anti-SIRPA antibody decreases cellular levels of SIRPA and inhibits interaction between SIRPA and one or more SIRPA ligands.

12. The anti-SIRPA antibody of any one of claims 1 to 9, wherein the antibody blocks binding of CD47 to human SIRPA.

13. The anti-SIRPA antibody of any one of claims 1 to 9, wherein the antibody selectively binds SIRPA and does not substantially block binding of CD47 to SIRPA expressed on cells and further, wherein binding of the antibody to SIRPA decreases the level of SIRPA on the cell surface, optionally wherein the SIRPA is human SIRPA.

14. The anti-SIRPA antibody of claim 13, wherein the antibody binds to the D1 domain of SIRPA, the D2 domain of SIRPA, or the D3 domain of SIRPA.

15. The anti-SIRPA antibody of claim 13 or 14, wherein the antibody (a) competes with an antibody comprising a  $V_H$  sequence comprising the amino acid sequence of SEQ ID NO:2 and a  $V_L$  sequence comprising the amino acid sequence of SEQ ID NO:3.

16. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a  $V_H$  region comprising: a CDR3 comprising the amino acid sequence of SEQ ID NO:11, a CDR1 comprising the amino acid sequence of SEQ ID NO:9, or a CDR2 comprising the amino acid sequence of SEQ ID NO:10.

17. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a  $V_H$  region comprising:

(a) a CDR1 that comprises the amino acid sequence of SEQ ID NO:9, a CDR1 that comprises the amino acid sequence of SEQ ID NO:9 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:9;

(b) a CDR2 that comprises the amino acid sequence of SEQ ID NO:10 or a CDR2 that comprises the amino acid sequence of SEQ ID NO:10 with no more than two amino acid substitutions; or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:10; and

(c) a CDR3 that comprises the amino acid sequence of SEQ ID NO:11, a CDR3 that comprises the amino acid sequence of SEQ ID NO:11 with no more than two amino acid substitutions; or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:11.

18. The anti-SIRPA antibody of claim 17, wherein the  $V_H$  region comprises: a CDR1 comprising the amino acid sequence of SEQ ID NO:9 or a CDR1 comprising the amino acid sequence of SEQ ID NO:9 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:10 or a CDR2 comprising the amino acid sequence of SEQ ID NO:10 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ ID NO:11 or a CDR3 comprising the amino acid sequence of SEQ ID NO:11 with no more than one amino acid substitution.

19. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a  $V_H$  region that comprises a CDR1 comprising the amino acid sequence

of SEQ ID NO:9, a CDR2 comprising the amino acid sequence of SEQ ID NO:10, and a CDR3 comprising the amino acid sequence of SEQ ID NO:11.

20. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a V<sub>H</sub> region comprising the amino acid sequence of a V<sub>H</sub> region shown in Figure 14A; or comprises a V<sub>H</sub> region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a V<sub>H</sub> region of Figure 14A.

21. The anti-SIRPA antibody of any one of claims 13 to 20, wherein the antibody comprises a V<sub>L</sub> region that comprises a CDR3 comprising the amino acid sequence of SEQ ID NO:8, a CDR1 comprising the amino acid sequence of SEQ ID NO:6, or a CDR2 comprising the amino acid sequence of SEQ ID NO:7.

22. The anti-SIRPA antibody of any one of claims 13 to 20, wherein the antibody comprises a V<sub>L</sub> region that comprises:

(a) a CDR1 comprising the amino acid sequence of SEQ ID NO:6, a CDR1 comprising the amino acid sequence of SEQ ID NO:6 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:6;

(b) a CDR2 comprising the amino acid sequence of SEQ ID NO:7, a CDR2 comprising the amino acid sequence of SEQ ID NO:7 with no more than two amino acid substitutions, or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:7; and

(c) a CDR3 comprising the amino acid sequence of SEQ ID NO:8, a CDR3 comprising the amino acid sequence of SEQ ID NO:8 with no more than two amino acid substitutions, or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:8.

23. The anti-SIRPA antibody of claim 22, wherein the V<sub>L</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:6 or a CDR1 comprising the amino acid sequence of SEQ ID NO:6 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:7 or a CDR2 comprising the amino acid sequence of SEQ ID NO:7 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ ID NO:8 or a CDR3

comprising the amino acid sequence of SEQ ID NO:8 with no more than one amino acid substitution.

24. The anti-SIRPA antibody of any one of claims 13 to 20, wherein the V<sub>L</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:6, a CDR2 comprising the amino acid sequence of SEQ ID NO:7, and a CDR3 comprising the amino acid sequence of SEQ ID NO:8.

25. The anti-SIRPA antibody of any one of claims 13 to 20, wherein the V<sub>L</sub> region comprises the amino acid sequence of a V<sub>L</sub> region shown in Figure 14B; or comprises a V<sub>H</sub> region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a V<sub>L</sub> region of Figure 14B.

26. The anti-SIRPA antibody of claim any one of claims 13 to 15, wherein the antibody comprises a V<sub>H</sub> region that comprises a CDR3 comprising the amino acid sequence of SEQ ID NO:17, a CDR1 comprising the amino acid sequence of SEQ ID NO:15, or a CDR2 comprising the amino acid sequence of SEQ ID NO:16.

27. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a V<sub>H</sub> region that comprises:

a) a CDR1 comprising the amino acid sequence of SEQ ID NO:15, a CDR1 comprising the amino acid sequence of SEQ ID NO:15 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:15;

(b) a CDR2 comprising the amino acid sequence of SEQ ID NO:16, a CDR2 comprising the amino acid sequence of SEQ ID NO:16 with no more than two amino acid substitutions, or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:16; and

(c) a CDR3 comprising the amino acid sequence of SEQ ID NO:17, a CDR3 comprising the amino acid sequence of SEQ ID NO:17 with no more than two amino acid substitutions, or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:17.

28. The anti-SIRPA antibody of claim 27, wherein the V<sub>H</sub> region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:15 or a CDR1 comprising the amino acid sequence of SEQ ID NO:15 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:16 or a CDR2 comprising the amino acid sequence of SEQ ID NO:16 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ ID NO:16 or a CDR3 comprising the amino acid sequence of SEQ ID NO:16 with no more than one amino acid substitution.

29. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a V<sub>H</sub> region that comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:15, a CDR2 comprising the amino acid sequence of SEQ ID NO:16, and a CDR3 comprising the amino acid sequence of SEQ ID NO:17.

30. The anti-SIRPA antibody of any one of claims 13 to 15, wherein the antibody comprises a V<sub>H</sub> region that comprises the amino acid sequence of a V<sub>H</sub> region of Figure 14C; or comprises a V<sub>H</sub> region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a V<sub>H</sub> region of Figure 14C.

31. The anti-SIRPA antibody of any one of claims 26 to 30, wherein the V<sub>L</sub> region comprises a CDR3 comprising the amino acid sequence of SEQ ID NO:14, a CDR1 comprising the amino acid sequence of SEQ ID NO:12, or a CDR2 comprising the amino acid sequence of SEQ ID NO:13.

32. The anti-SIRPA antibody of any one of claims 26 to 30, wherein the antibody comprises a V<sub>L</sub> region that comprises:

a) a CDR1 comprising the amino acid sequence of SEQ ID NO:12, a CDR1 comprising the amino acid sequence of SEQ ID NO:12 with no more than two amino acid substitutions, or a CDR1 having at least about 90% identity to the amino acid sequence of SEQ ID NO:12;

(b) a CDR2 comprising the amino acid sequence of SEQ ID NO:13, a CDR2 comprising the amino acid sequence of SEQ ID NO:13 with no more than two amino acid substitutions, or a CDR2 having at least about 90% identity to the amino acid sequence of SEQ ID NO:13; and

(c) a CDR3 comprising the amino acid sequence of SEQ ID NO:14, a CDR3 comprising the amino acid sequence of SEQ ID NO:14 with no more than two amino acid substitutions, or a CDR3 having at least about 90% identity to the amino acid sequence of SEQ ID NO:14.

33. The anti-SIRPA antibody of claim 32, wherein the  $V_L$  region comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:12 or a CDR1 comprising the amino acid sequence of SEQ ID NO:12 with no more than one amino acid substitution; a CDR2 comprising the amino acid sequence of SEQ ID NO:13 or a CDR2 comprising the amino acid sequence of SEQ ID NO:13 with no more than one amino acid substitution; and a CDR3 comprising the amino acid sequence of SEQ ID NO:4 or a CDR3 comprising the amino acid sequence of SEQ ID NO:14 with no more than one amino acid substitution.

34. The anti-SIRPA antibody of any one of claims 13 to 20, wherein the antibody comprises a  $V_L$  region that comprises a CDR1 comprising the amino acid sequence of SEQ ID NO:6, a CDR2 comprising the amino acid sequence of SEQ ID NO:7, and a CDR3 comprising the amino acid sequence of SEQ ID NO:8.

35. The anti-SIRPA antibody of any one of claims 26 to 30, wherein the  $V_L$  region comprises the amino acid sequence of a  $V_L$  region of Figure 14D; or comprises a  $V_L$  region having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of a  $V_H$  region of Figure 14D.

36. An isolated anti-SIRPA antibody of claim 13 or 14, wherein the anti-SIRPA competes with one or more antibodies for binding to SIRPA, wherein the antibody is selected from the group consisting of 8A9, 8F4, 1E2, 7H9, and 4D8.

37. An isolated anti-SIRPA antibody of claim 13 or 14, wherein the anti-SIRPA binds to essentially the same epitope as one or more antibodies selected from the group consisting of 3F9, 9C2, 8A9, 8F4, 1E2, 7H9, and 4D8.

38. The isolated anti-SIRPA antibody of claim 36 or 37, wherein the antibody comprises a  $V_H$  region and a  $V_L$  region, wherein the  $V_H$  region, the  $V_L$  region, or

both comprise at least one, two, three, four, five, or six CDRs of a monoclonal antibody selected from the group consisting of 3F9, 9C2, 8A9, 8F4, 1E2, 7H9, and 4D8.

39. The anti-SIRPA antibody of any one of claims 1 to 38, wherein the antibody is a monoclonal antibody.

40. The anti-SIRPA antibody of any one of claims 1 to 39, wherein the antibody is a humanized antibody.

41. The anti-SIRPA antibody of any one of claims 1 to 40, wherein the antibody is an Fab, Fab', Fab'-SH, F(ab')<sub>2</sub>, Fv or scFv fragment.

42. The anti-SIRPA antibody of any one of claims 1 to 41, wherein the antibody is a multivalent antibody.

43. The anti-SIRPA antibody of any one of claims 1 to 42, wherein the anti-SIRPA antibody is of the IgG class, the IgM class, or the IgA class.

44. The anti-SIRPA antibody of claim 43, wherein the anti-SIRPA antibody has an IgG1, IgG2, IgG3, or IgG4 isotype.

45. The anti-SIRPA antibody of claim 44, wherein the antibody binds to an inhibitory Fc receptor.

46. The anti-SIRPa antibody of claim 45, wherein the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (Fc $\gamma$ RIIB).

47. The anti-SIRPA antibody of of any one of claims 1 to 42, wherein the antibody decreases cellular levels of Fc $\gamma$ R.

48. The anti-SIRPa antibody of claim 47, wherein the antibody decreases cellular levels of Fc $\gamma$ RIIB.

49. The anti-SIRPA antibody of claim 46, wherein:

(a) the anti-SIRPA antibody has a human or mouse IgG1 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: N297A, D265A, D270A, L234A, L235A, G237A, P238D, L328E, E233D, G237D, H268D, P271G, A330R, C226S, C229S, E233P, L234V, L234F, L235E, P331S, S267E, L328F, A330L, M252Y, S254T, T256E, N297Q, P238S, P238A, A327Q, A327G, P329A, K322A, T394D, and any combination thereof, wherein the numbering of the residues is according to EU numbering, or comprises an amino acid deletion in the Fc region at a position corresponding to glycine 236;

(b) the anti-SIRPA antibody has an IgG1 isotype and comprises an IgG2 isotype heavy chain constant domain 1(CH1) and hinge region, optionally wherein the IgG2 isotype CH1 and hinge region comprises the amino acid sequence of ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGVHTFPALQSS GLYSLSSVVT VPSSNFGTQT YTCNVDHKPS NTKVDKTVERKCCVECPPCP (SEQ ID NO:34), and optionally wherein the antibody Fc region comprises a S267E amino acid substitution, a L328F amino acid substitution, or both, and/or a N297A or N297Q amino acid substitution, wherein the numbering of the residues is according to EU numbering;

(c) the anti-SIRPA antibody has an IgG2 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: P238S, V234A, G237A, H268A, H268Q, V309L, A330S, P331S, C214S, C232S, C233S, S267E, L328F, M252Y, S254T, T256E, H268E, N297A, N297Q, A330L, and any combination thereof, wherein the numbering of the residues is according to EU numbering;

(d) the anti-SIRPA antibody has a human or mouse IgG4 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: L235A, G237A, S228P, L236E, S267E, E318A, L328F, M252Y, S254T, T256E, E233P, F234V, L234A/F234A, S228P, S241P, L248E, T394D, N297A, N297Q, L235E, and any combination thereof, wherein the numbering of the residues is according to EU numbering; or

(e) the anti-SIRPA antibody has a hybrid IgG2/4 isotype, and optionally wherein the antibody comprises an amino acid sequence comprising amino acids 118 to 260 of human IgG2 and amino acids 261 to 447 of human IgG4, wherein the numbering of the residues is according to EU numbering.

50. The anti-SIRPA antibody of claim 44, wherein:

(a) the anti-SIRPA antibody has a human or mouse IgG1 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: N297A, N297Q, D270A, D265A, L234A, L235A, C226S, C229S, P238S, E233P, L234V, P238A, A327Q, A327G, P329A, K322A, L234F, L235E, P331S, T394D, A330L, M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU numbering;

(b) the anti-SIRPA antibody has an IgG2 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: P238S, V234A, G237A, H268A, H268Q, H268E, V309L, N297A, N297Q, A330S, P331S, C232S, C233S, M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU numbering; or

(c) the anti-SIRPA antibody has an IgG4 isotype and comprises one or more amino acid substitutions in the Fc region at a residue position selected from the group consisting of: E233P, F234V, L234A/F234A, L235A, G237A, E318A, S228P, L236E, S241P, L248E, T394D, M252Y, S254T, T256E, N297A, N297Q, and any combination thereof, wherein the numbering of the residues is according to EU numbering.

51. The anti-SIRPA antibody of claim 50, wherein:

(a) the Fc region further comprises one or more additional amino acid substitutions at a position selected from the group consisting of A330L, L234F; L235E, P331S, and any combination thereof, wherein the numbering of the residues is according to EU numbering;

(b) the Fc region further comprises one or more additional amino acid substitutions at a position selected from the group consisting of M252Y, S254T, T256E, and any combination thereof, wherein the numbering of the residues is according to EU numbering; or

(c) the Fc region further comprises a S228P amino acid substitution according to EU numbering.

52. The anti-SIRPA antibody of any one of claims 1 to 40, wherein the antibody has an IgG4 isotype.

53. The anti-SIRPA antibody of claim 40, wherein the anti-SIRPA antibody comprises an S228P amino acid substitution at residue position 228, an F234A amino acid substitution at residue position 234, and an L235A amino acid substitution at residue position 235, wherein the numbering of the residue position is according to EU numbering.

54. The anti-SIRPA antibody of any one of claims 1 to 53, wherein the antibody is a bispecific antibody.

55. The anti-SIRPA antibody of claim 54, wherein the antibody recognizes a first and a second antigen, wherein the first antigen is SIRPA and the second antigen is:

- (a) an antigen facilitating transport across the blood-brain-barrier;
- (b) an antigen facilitating transport across the blood-brain-barrier selected from the group consisting of transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), diphtheria toxin receptor, CRM197, a llama single domain antibody, TMEM 30(A), a protein transduction domain, TAT, Syn-B, penetratin, a poly-arginine peptide, an angiopep peptide, and ANG1005;
- (c) a disease-causing agent selected from the group consisting of disease-causing peptides or proteins or, disease-causing nucleic acids, wherein the disease-causing nucleic acids are antisense GGCCCC (G2C4) repeat-expansion RNA, the disease-causing proteins are selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides; and
- (d) ligands and/or proteins expressed on immune cells, wherein the ligands and/or proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28,

CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, and phosphatidylserine; and a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells.

56. The anti-SIRPA antibody of any one of claims 1 to 55, wherein the anti-anti-SIRPA antibody is a conjugated antibody.

57. The anti-SIRPA antibody of claim 56, wherein the anti-SIRPA antibody is conjugated to a detectable marker, a toxin, or a therapeutic agent.

58. The anti-SIRPA antibody of claim 57, wherein the anti-SIRPA antibody is conjugated to a toxin selected from the group consisting of ricin, ricin A chain, doxorubicin, daunorubicin, a maytansinoid, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, *Pseudomonas* exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, *Saponaria officinalis* inhibitor, glucocorticoid, auristatin, auromycin, yttrium, bismuth, combrestatin, duocarmycins, dolastatin, cc1065, and a cisplatin.

59. The anti-SIRPA antibody of any one of claims 1 to 58, wherein the anti-SIRPA antibody is used in combination with one or more antibodies that specifically bind a disease-causing protein selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides, and any combination thereof; or with one or more antibodies that bind an immunomodulatory protein selected from the group consisting of: PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-

1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, TREM1, TREM2, CD33, Siglec-5, Siglec-7, Siglec-9, Siglec-11, phosphatidylserine, disease-causing nucleic acids, antisense GGCCCC (G2C4) repeat-expansion RNA, and any combination thereof

60. A method of decreasing the activity, functionality, or survival of regulatory T cells, tumor-imbedded immunosuppressor dendritic cells, tumor-imbedded immunosuppressor macrophages, myeloid-derived suppressor cells, tumor-associated macrophages, acute myeloid leukemia (AML) cells, chronic lymphocytic leukemia (CLL) cell, or chronic myeloid leukemia (CML) cells in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of an agent that binds or interacts with SIRPA.

61. A method of inducing or promoting the survival, maturation, functionality, migration, or proliferation of one or more immune cells in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of an agent that decreases cellular levels of SIRPA, inhibits interaction between SIRPA and one or more SIRPA ligands, or both.

62. The method of claim 61, wherein the one or more immune cells are selected from the group consisting of dendritic cells, macrophages, neutrophils, NK cells, microglia, T cells, T helper cells, cytotoxic T cells, and any combination thereof.

63. A method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any one of claims 1 to 58 to a patient that has a tumor that expresses CD47.

64. A method of treating cancer, the method comprising administering a therapeutically effective amount of an agent that decreases the cellular levels of SIRPA.

65. The method of claim 64, wherein the agent is an anti-SIRPA antibody of any one of claims 1 to 58.

66. The method of claim 63, 64, or 65, wherein the method further comprises administering a therapeutic agent that inhibits PD1, PDL1, CD40, OX40, ICOS,

CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73.

67. The method of claim 66, wherein the therapeutic agent is an antibody that inhibits PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73.

68. The method of claim 63, 64, or 65, further comprising administering to the individual at least one antibody that specifically binds to an inhibitory checkpoint molecule, and/or one or more standard or investigational anti-cancer therapies.

69. The method of claim 66, wherein the at least one antibody that specifically binds to an inhibitory checkpoint molecule is administered in combination with the anti-SIRPA antibody.

70. The method of claim 66 or 67, wherein the at least one antibody that specifically binds to an inhibitory checkpoint molecule is selected from the group consisting of an anti-PD-L1 antibody, an anti-CTLA4 antibody, an anti-PD-L2 antibody, an anti-PD-1 antibody, an anti-B7-H3 antibody, an anti-B7-H4 antibody, and anti-HVEM antibody, an anti- B- and T-lymphocyte attenuator (BTLA) antibody, an anti-Killer inhibitory receptor (KIR) antibody, an anti-GAL9 antibody, an anti-TIM-1 antibody, an anti-TIM3 antibody, an anti-TIM-4 antibody, an anti-A2AR antibody, an anti-CD39 antibody, an anti-CD73 antibody, an anti-LAG-3 antibody, an anti-phosphatidylserine antibody, an anti-CD27 antibody, an anti-CD30 antibody, an anti-TNF $\alpha$  antibody, an anti-CD33 antibody, an anti-Siglec-5 antibody, an anti-Siglec-7 antibody, an anti-Siglec-9 antibody, an anti-Siglec-11 antibody, an antagonistic anti-TREM1 antibody, an antagonistic anti-TREM2 antibody, an anti-TIGIT antibody, an anti-VISTA antibody, an anti-CD2 antibody, an anti-CD5 antibody, and any combination thereof.

71. The method of claim 66, wherein the one or more standard or investigational anti-cancer therapies are selected from the group consisting of radiotherapy, cytotoxic chemotherapy, targeted therapy, imatinib therapy, trastuzumab therapy, etanercept therapy, adoptive cell transfer (ACT) therapy, chimeric antigen receptor T cell transfer (CAR-T) therapy, vaccine therapy, and cytokine therapy.

72. The method of any one of claims 63 to 71, further comprising administering to the individual at least one antibody that specifically binds to an inhibitory cytokine.

73. The method of claim 72, wherein the at least one antibody that specifically binds to an inhibitory cytokine is administered in combination with the anti-SIRPA antibody.

74. The method of claim 66 to 67, wherein the at least one antibody that specifically binds to an inhibitory cytokine is selected from the group consisting of an anti-CCL2 antibody, an anti-CSF-1 antibody, an anti-IL-2 antibody, and any combination thereof.

75. The method of any one of claims 63 to 74, further comprising administering to the individual at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein.

76. The method of claim 66, wherein the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is administered in combination with the anti-SIRPA antibody.

77. The method of claim 75 or 76, wherein the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is selected from the group consisting of an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonistic anti-TREM1 antibody, an agonistic anti-TREM2 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, an agonist anti-CD30 antibody, an agonist anti-BTLA antibody, an agonist anti-HVEM antibody, an agonist anti-CD2 antibody, an agonist anti-CD5 antibody, and any combination thereof.

78. The method of any one of claims 63 to 77, further comprising administering to the individual at least one stimulatory cytokine, optionally IFN- $\alpha$ 4, IFN- $\beta$ , IL-1 $\beta$ , TNF- $\alpha$ , IL-6, IL-8, CRP, IL-20 family members, LIF, IFN- $\gamma$ , OSM, CNTF, GM-CSF, IL-11, IL-12, IL-15, IL-17, IL-18, IL-23, CXCL10, IL-33, MCP-1, MIP-1-beta, and any combination thereof.

79. A method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any one of claims 1 to 58 to a patient that has cancer cells of a myeloid lineage that expresses SIRPA.

80. A method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any one of claims 1 to 58 to a subject that has a cancer, wherein the cancer is selected from the group consisting of sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, melanoma, lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, and fibrosarcoma.

81. A method of treating cancer, the method comprising administering a therapeutically effective amount of an anti-SIRPA antibody of any one of claims 1 to 58 to a subject that has cancer, wherein the cancer is selected from the group consisting of glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma; diffuse large B-cell lymphoma; lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, non-Hodgkin's lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thyumoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma.

82. The method of claim 80 or 81, wherein the anti-SIRPA antibody is conjugated to a cytotoxic agent and/or induces ADCC.

83. A pharmaceutical composition comprising an anti-SIRPA antibody of any one of claims 1 to 58 and a physiologically acceptable carrier.

84. anti-SIRPA antibody of any one of claims 1 to 58 for use in the treatment of cancer.

85. An antibody of any one of claims 1 to 58 for use in a method of preparing a medicament for the treatment of cancer.

86. A method of preventing, reducing risk, or treating a disease, disorder, or injury selected from the group consisting of dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, taupathy disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, traumatic brain injury, stroke, frontotemporal dementia, spinal cord injury, Huntington's disease, infections, and cancer comprising administering to an individual in need thereof a therapeutically effective amount of an agent that decreases cellular levels of SIRPA, inhibits interaction between SIRPA and one or more SIRPA ligands, or both.

87. The method of claim 86, wherein the disease, disorder, or injury is cancer and wherein the agent inhibits one or more SIRPA activities selected from the group consisting of:

(a) promoting proliferation, maturation, migration, differentiation, and/or functionality of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid derived suppressor cells, tumor-associated macrophages, tumor-associated suppressor neutrophils, tumor-associated suppressor NK cells, non-tumorigenic CD14+ myeloid cells, and regulatory T cells;

(b) enhancing infiltration of one or more of immunosuppressor dendritic cells, immunosuppressor macrophages, immunosuppressor neutrophils, immunosuppressor NK cells, myeloid derived suppressor cells, tumor-associated macrophages, tumor-associated suppressor neutrophils, tumor-associated suppressor NK cells, and regulatory T cells into tumors;

(c) increasing number of tumor-promoting myeloid/granulocytic immune-suppressive cells and/or non-tumorigenic CD14+ myeloid cells in a tumor, in peripheral blood, or other lymphoid organ;

(d) enhancing tumor-promoting activity of myeloid-derived suppressor cells (MDSC) and/or non-tumorigenic CD14+ myeloid cells;

- (e) increasing expression of tumor-promoting cytokines in a tumor or in peripheral blood, optionally wherein the tumor-promoting cytokines are TGF-beta or IL-10;
- (f) increasing tumor infiltration of tumor-promoting FoxP3+ regulatory T lymphocytes;
- (g) decreasing activation of tumor-specific T lymphocytes with tumor killing potential;
- (h) decreasing infiltration of tumor-specific T lymphocytes with tumor killing potential;
- (i) decreasing infiltration of tumor-specific NK cells with tumor killing potential;
- (j) decreasing the tumor killing potential of NK cells;
- (k) decreasing infiltration of tumor-specific B lymphocytes with potential to enhance immune response;
- (l) increasing tumor volume;
- (m) increasing tumor growth rate;
- (n) increasing metastasis;
- (o) increasing rate of tumor recurrence;
- (p) decreasing efficacy of one or more immune-therapies that modulate anti-tumor T cell responses, optionally wherein the one or more immune-therapies are immune-therapies that target one or more target proteins selected from the group consisting of PD1/PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, CD73, and any combination thereof, or one or more cancer vaccines;
- (q) inhibition of PLC $\gamma$ /PKC/calcium mobilization; and
- (r) inhibition of PI3K/Akt, Ras/MAPK signaling.

88. The method of claim 86, wherein the disease, disorder, or injury is cancer, and wherein the agent exhibits one or more SIRPA activities selected from the group consisting of:

- (a) increasing the number of tumor infiltrating CD3+ T cells;
- (b) decreasing cellular levels of CD33 in non-tumorigenic CD14+myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood;

- (c) reducing the number of non-tumorigenic CD14+ myeloid cells, optionally wherein the non-tumorigenic CD14+ myeloid cells are tumor infiltrating cells or optionally wherein the non-tumorigenic CD14+ myeloid cells are present in blood;
- (d) reducing PD-L1 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (e) reducing PD-L2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (f) reducing B7-H2 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (g) reducing B7-H3 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (h) reducing CD200R levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (i) reducing CD163 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (j) reducing CD206 levels in one or more cells, optionally wherein the one or more cells are non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (k) decreasing tumor growth rate of solid tumors;
- (l) reducing tumor volume;
- (m) increasing efficacy of one or more PD-1 inhibitors;
- (n) increasing efficacy of one or more checkpoint inhibitor therapies and/or immune-modulating therapies, optionally wherein the one or more checkpoint inhibitor therapies and/or immune-modulating therapies target one or more of CTLA4, the adenosine pathway, PD-L1, PD-L2, OX40, TIM3, LAG3, or any combination thereof;
- (o) increasing efficacy of one or more chemotherapy agents, optionally wherein the one or more of the chemotherapy agents are gemcitabine, capecitabine, anthracyclines, doxorubicin (Adriamycin®), epirubicin (Ellence®), taxanes, paclitaxel (Taxol®), docetaxel (Taxotere®), 5-fluorouracil (5-FU), cyclophosphamide (Cytoxan®), carboplatin (Paraplatin®), and any combination thereof;
- (p) increasing proliferation of T cells in the presence of non-tumorigenic myeloid-derived suppressor cells (MDSC);
- (q) inhibiting differentiation, survival, and/or one or more functions of non-tumorigenic myeloid-derived suppressor cells (MDSC); and

(r) killing CD33-expressing immunosuppressor myeloid cells and/or CD14-expressing cells in solid tumors and associated blood vessels when conjugated to a chemical or radioactive toxin.

89. The method of claim 87 or 88, wherein the cancer expresses SIRPA or one or more SIRPA ligands.

90. The method of claim 85, wherein the disease, disorder, or injury is selected from the group consisting of dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, taupathy disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, traumatic brain injury, stroke, frontotemporal dementia, spinal cord injury, and Huntington's disease; and wherein the agent is an anti-SIRPA antibody that downregulates SIRPA.

91. A polynucleotide comprising a nucleic acid sequence encoding a V<sub>H</sub> region of an anti-SIRPA antibody of any one of claims 1 to 55.

92. A polynucleotide comprising a nucleic acid sequence encoding a V<sub>L</sub> region of an anti-SIRPA antibody of any one of claims 1 to 55.

93. An expression vector comprising the polynucleotide of claim 91 or the polynucleotide of claim 92.

94. An expression vector comprising the polynucleotide of claim 91 and the polynucleotide of claim 92.

95. A host cell comprising the polynucleotide of claim 91 or the polynucleotide of claim 92.

96. A host cell comprising the polynucleotide of claim 91 and the polynucleotide of claim 92.

97. A host cell comprising the expression vector of claim 93 or 94.

98. A method of producing an anti-SIRPA antibody, the method comprising culturing a host cell of any one of claims 95 to 97 under conditions in which the antibody is expressed.

99. The method of claim 98, wherein the host cell is a mammalian host cell.

1/37

## FIGURE 1A

2/37

FIGURE 1B

## FIGURE 2

4/37







FIGURE 3A

5/37

FIGURE 3B



6/37

**FIGURE 3C****FIGURE 3D**

7/37





FIGURE 4B



9/37






FIGURE 5A

10/37

**FIGURE 5B**

11/37



12/37





FIGURE 6B

13/37

FIGURE 7A



14/37





FIGURE 7B

FIGURE 8A


Enhanced phagocytosis of phrodo-labeled Raji cells with CD20

15/37



16/37

FIGURE 8B



17/37




FIGURE 8C

18/37





FIGURE 9A

19/37

FIGURE 9B



20/37




FIGURE 9C

21/37

FIGURE 10A





FIGURE 10B



22/37





FIGURE 10C


23/37

**FIGURE 10D** Spleen:



24/37

FIGURE 11A



25/37




FIGURE 11B

26/37

FIGURE 12A






FIGURE 12B




27/37


Figure 12C



28/37

**FIGURE 13A**

29/37

**FIGURE 13B**

30/37

## FIGURE 14A

Potential humanized sequence based on IgHV3-23\*01 acceptor framework (AbM CDR definition)

IgHV3-23\*01 EVQLESGGGLVQPGGSLRLSCAASGCPYSSYAMSWVRQAPGKGLEWVSAISGGSTYADSVKG

RPTLSRDNSKWLNLQMMNLRAEDTAVVYCAK

Joining region IgM3 J80256 | IgH34\*01 | YFDYW3QG3LIVVYSS

seq	10	20	30	40	50	60
AbM	10	20	30	40	50	ab
b b	p	p b	b	b	b	b
SB-3P9	EVKLVESGGGLVQPGGSLRLSCAAS GFTYSSYAMS WVRQAPGKGLEWVATISDYGSSYV	*	*	*	*	*
IgHV3-23*01	EVQLESGGGLVQPGGSLRLSCAAS GFTYSSYAMS WVRQAPGKGLEWVMSG-SGSSY	*	*	*	*	*
hsB-3P9-H1	EVQLESGGGLVQPGGSLRLSCAAS GFTYSSYAMS WVRQAPGKGLEWVATISDYGSSYV	*	*	*	*	*
hsB-3P9-H2	EVQLESGGGLVQPGGSLRLSCAAS GFTYSSYAMS WVRQAPGKGLEWVATISDYGSSYV	*	*	*	*	*
seq	70	80	90	100	110	120
AbM	60	70	80	90	100	110
i b	b b x	b b b	b b b	bibbb	100abed	110
SB-3P9	YDGSVKGRFTISRDNSKNTLYLQMMNLRAEDTAVVYCAK	YDGSVKGRFTISRDNSKNTLYLQMMNLRAEDTAVVYCAK	PPYDDYGGFAY	WQQGTIVVSA	*	*
IgHV3-23*01	YADSVVKGRFTISRDNSKNTLYLQMMNLRAEDTAVVYCAK	*	*	*	*	*
hsB-3P9-H1	YADSVVKGRFTISRDNSKNTLYLQMMNLRAEDTAVVYCAK	PPYDDYGGFAY	WQQGTIVVSS	*	*	*
hsB-3P9-H2	YADSVVKGRFTISRDNSKNTLYLQMMNLRAEDTAVVYCAK	p	PPYDDYGGFAY	WQQGTIVVSS	*	*

## FIGURE 14B

Potential humanized sequence based on IgKV3-11\*01 acceptor framework (AbM CDR definition)  
 IgKV3-11\*01 EIVLQSPATLSSLSPGERATLSC RAKSVSSGGSYMM WYQQKPGQAPRLIY LASNLIS  
 GIPARFSGGGGGCTDFTLTISSLEPDEAVYCCQRSWPP  
 Joining region IgKV3-11\*01 IgKV3-12 | IgKV3-13 | YIPGQXKLEIK

seq	10	20	30	40	50	60
AbM	10	20	30abcd	40	50	60
b b b p p b b b	b b b b b	bi bi i ii ibbi	i			
SB-3F9	DIVLQSPATLSSLSPGERATLSC RAKSVSSGGSYMM WYQQKPGQAPRLIY LASNLIS	*	*	*	*	*
3-11*01	EIVLQSPATLSSLSPGERATLSC RAKSVSSGGSYMM WYQQKPGQAPRLIY LASNLIS					
hSB-3F9-11	EIVLQSPATLSSLSPGERATLSC RAKSVSSGGSYMM WYQQKPGQAPRLIY LASNLIS					
hSB-3F9-12	EIVLQSPATLSSLSPGERATLSC RAKSVSSGGSYMM WYQQKPGQAPRLIY LASNLIS					
hSB-3F9-13	EIVLQSPATLSSLSPGERATLSC RAKSVSSGGSYMM WYQQKPGQAPRLIY LASNLIS					
		V	P	#		

seq	70	80	90	100	110
AbM	60	70	80	90	100
b b b b b b b	ib ibi ibi ibi ibi	i b b b			
SB-3F9	GIPARFSGGGGGCTDFTLTISSLEPDEAVYCCQRSWPP	*	*	*	*
3-11*01	GIPARFSGGGGGCTDFTLTISSLEPDEAVYCCQRSWPP				
hSB-3F9-11	GIPARFSGGGGGCTDFTLTISSLEPDEAVYCCQRSWPP				
hSB-3F9-12	GIPARFSGGGGGCTDFTLTISSLEPDEAVYCCQRSWPP				
hSB-3F9-13	GIPARFSGGGGGCTDFTLTISSLEPDEAVYCCQRSWPP				
	A	#	^		



32/37

FIGURE 14C

33/37

FIGURE 14D

34/37

**Figure 15A****Figure 15B**

35/37







Figure 16

36/37

**Figure 17A**

37/37

**Figure 18**

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2017/065366

**A. CLASSIFICATION OF SUBJECT MATTER**  
INV. C07K16/28 A61K39/395  
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, FSTA, CHEM ABS Data, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 2013/056352 A1 (UNIV HEALTH NETWORK [CA]; HOSPITAL FOR SICK CHILDREN [CA]; WANG JEAN C) 25 April 2013 (2013-04-25) the whole document</p> <p style="text-align: center;">-----</p> <p style="text-align: center;">-/-</p>	1-99

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
12 March 2018	26/03/2018
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Sirim, Pinar

## INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/065366

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DE VRIES HELGA E ET AL:          "Signal-regulatory protein alpha-CD47 interactions are required for the transmigration of monocytes across cerebral endothelium",          THE JOURNAL OF IMMUNOLOGY, THE AMERICAN ASSOCIATION OF IMMUNOLOGISTS, US,          vol. 168, no. 11, 1 June 2002 (2002-06-01),          , pages 5832-5839, XP002527047,          ISSN: 0022-1767          the whole document          see in particular abstract; page 5835,          right-column, last paragraph; page 5836,          left-column, paragraph 2</p> <p>-----</p>	1-99
X	<p>J. ALBLAS ET AL: "Signal Regulatory Protein alpha Ligation Induces Macrophage Nitric Oxide Production through JAK/STAT- and Phosphatidylinositol 3-Kinase/Rac1/NAPDH Oxidase/H2O2-Dependent Pathways",          MOLECULAR AND CELLULAR BIOLOGY.,          vol. 25, no. 16,          15 August 2005 (2005-08-15), pages 7181-7192, XP055261468,          US          ISSN: 0270-7306, DOI:          10.1128/MCB.25.16.7181-7192.2005          the whole document          see in particular abstract; full paragraph bridging pages 7184-85; Fig. 2</p> <p>-----</p>	1-99
X	<p>Per-Arne Oldenborg: "Cd47-Signal Regulatory Protein [alpha] (Sirp[alpha]) Regulates Fc[gamma] and Complement Receptor-Mediated Phagocytosis",          The Journal of Experimental Medicine, vol.193, no. 7,          2 April 2001 (2001-04-02), pages 855-861,          XP055283803,          Retrieved from the Internet:          URL:<a href="http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2193364/pdf/010027.pdf">http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2193364/pdf/010027.pdf</a>          [retrieved on 2016-06-27]          the whole document          see in particular abstract; Figures 2/3/5/7</p> <p>-----</p> <p style="text-align: center;">-/-</p>	1-99

## INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/065366

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Anonymous: "MABS164   Anti-SHPS-1 Antibody, clone P84", 1 January 2016 (2016-01-01), pages 1-4, XP055261233, Retrieved from the Internet: URL: <a href="http://www.merckmillipore.com/NL/en/product/Anti-SHPS-1-Antibody,-clone-P84,MM_NF-MABS164#anchor">http://www.merckmillipore.com/NL/en/product/Anti-SHPS-1-Antibody,-clone-P84,MM_NF-MABS164#anchor</a> Packaging%20Information [retrieved on 2016-03-29] the whole document -----	1-99
X	M. SEIFFERT ET AL: "Signal-regulatory protein alpha (SIRPalpha) but not SIRPbeta is involved in T-cell activation, binds to CD47 with high affinity, and is expressed on immature CD34+CD38- hematopoietic cells", BLOOD, vol. 97, no. 9, 1 May 2001 (2001-05-01), pages 2741-2749, XP055056736, ISSN: 0006-4971, DOI: 10.1182/blood.V97.9.2741 the whole document see in particular page 2744 and table 1 -----	1-99
A	LEE ET AL: "Novel structural determinants on SIRP alpha that mediate binding to CD47.", JOURNAL OF IMMUNOLOGY, vol. 179, no. 11, 1 January 2007 (2007-01-01), pages 7741-7750, XP055081468, the whole document see in particular abstract; pages 7747-49; Fig. 4, 5, 6 -----	1-99
X, P	TADAHIKO YANAGITA ET AL: "Anti-SIRP[alpha] antibodies as a potential new tool for cancer immunotherapy", JCI INSIGHT, vol. 2, no. 1, 12 January 2017 (2017-01-12), XP055421877, ISSN: 2379-3708, DOI: 10.1172/jci.insight.89140 the whole document see in particular abstract; page 2 last sentence; page 8, right-column, paragraph 1; -----	1-99

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No  
PCT/US2017/065366

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013056352 A1	25-04-2013	US 2014242095 A1 WO 2013056352 A1	28-08-2014 25-04-2013