



(86) Date de dépôt PCT/PCT Filing Date: 2007/01/25  
(87) Date publication PCT/PCT Publication Date: 2008/08/02  
(45) Date de délivrance/Issue Date: 2014/06/17  
(85) Entrée phase nationale/National Entry: 2008/07/22  
(86) N° demande PCT/PCT Application No.: EP 2007/050752  
(87) N° publication PCT/PCT Publication No.: 2007/085638  
(30) Priorité/Priority: 2006/01/26 (EP06100914.8)


(51) Cl.Int./Int.Cl. *C07D 239/94* (2006.01),  
*C07D 405/12* (2006.01), *C07F 9/6512* (2006.01)

(72) Inventeurs/Inventors:  
SCHROEDER, JUERGEN, DE;  
DZIEWAS, GEORG, DE;  
FACHINGER, THOMAS, DE;  
JAEGER, BURKHARD, DE;  
REICHEL, CARSTEN, DE;  
RENNER, SVENJA, DE

(73) Propriétaire/Owner:  
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,  
DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PROCEDE DE PREPARATION DE DERIVES QUINAZOLINE A SUBSTITUTION AMINOCROTONYLAMINO  
(54) Title: PROCESS FOR PREPARING AMINOCROTONYLAMINO-SUBSTITUTED QUINAZOLINE DERIVATIVES



(57) Abrégé/Abstract:


The invention provides a process for producing aminocrotonylamino-substituted quinazoline derivatives of general formula I (see formula I) and the invention also provides sulphonyl derivatives of formula XIII (see formula XIII). The compound of formula XIII is an intermediate that may be used in production of the compound of formula I, and the compound of formula I may be used as an inhibitor of signal transduction mediated by tyrosinekinases and by the Epidermal Growth Factor-Receptor (EGF-R) and are therefore suitable for treatment of tumoral diseases.

25771-1542


**PROCESS FOR PREPARING AMINOCROTONYLAMINO-SUBSTITUTED  
QUINAZOLINE DERIVATIVES**

ABSTRACT

The invention provides a process for producing aminocrotonylamino-substituted  
5 quinazoline derivatives of general formula I



and the invention also provides sulphonyl derivatives of formula XIII




The compound of formula XIII is an intermediate that may be used in production of  
10 the compound of formula I, and the compound of formula I may be used as an  
inhibitor of signal transduction mediated by tyrosinekinases and by the Epidermal  
Growth Factor-Receptor (EGF-R) and are therefore suitable for treatment of tumoral  
diseases.

25771-1542

- 1 -


**Process for preparing aminocrotonylamino-substituted quinazoline derivatives**

The invention relates to a process for preparing aminocrotonylamino-substituted quinazoline derivatives of general formula (I)



5 wherein the groups  $R_a$ ,  $R_b$ ,  $R_c$  and  $R_d$  have the meanings given below.

According to one aspect of the present invention, there is provided a process for preparing a compound of general formula



$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group,

10  $R_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

$R_c$  denotes a methyl, ethyl or 2-methoxyethyl group or

15  $R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denotes a morpholino or homomorpholino group optionally substituted by one or two  $C_{1-3}$ -alkyl groups, and


25771-1542

- 1a -

$R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

comprising the following process steps:


- 5 a) reacting 7-chloro-6-nitro-3*H*-quinazolin-4-one



(VII)

with a primary amine of formula  $R_a-NH_2$  (XV), wherein  $R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group, in the presence of  $POCl_3$ ,


- b) converting the resulting compound of general formula



(X)

10

into the sulphonyl derivative of formula



(XIII)

wherein


25771-1542

- 1b -

$R^3$  denotes a  $C_{1-4}$ -alkyl group wherein the hydrogen atoms are optionally wholly or partly replaced by fluorine atoms, or

a phenyl group optionally substituted by one to three substituents selected from the group consisting of  $C_{1-3}$ -alkyl groups, halogen atoms, cyano groups and nitro groups, wherein the substituents are identical or different, and wherein  $R_a$  in the two formulae (X) and (XIII) has the meanings given under a),


c) converting the sulphonyl derivative of formula (XIII) into a compound of formula



by reacting with an alcohol of formula  $R_d$ -OH (XVI) in the presence of a base,

wherein  $R_a$  has the meanings given under a) and  $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

d) reducing the compound of formulae (XI) thus obtained to the amino derivative of formula




25771-1542

- 1c -

wherein  $R_a$  has the meanings given under a) and  $R_d$  has the meanings given under c),


e) converting the amino derivatives of formula (XII) into the phosphonic ester of formula



5

wherein  $R_a$  has the meanings given under a) and  $R_d$  has the meanings given under c),  $R^1$  and  $R^2$  each independently of one another denote a  $C_{1-4}$ -alkyl group,

f) reacting the resulting phosphonic ester of formula (III) with a hydrogen sulphite adduct of formula



10

wherein

$M^+$  denotes a cation or a proton and

$R_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

15

$R_c$  denotes a methyl, ethyl or 2-methoxyethyl group or


25771-1542

- 1d -

$R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denote a morpholino or homomorpholino group optionally substituted by one or two  $C_{1-3}$ -alkyl groups,

in the manner of a Wittig-Horner-Emmons reaction.

- 5 According to another aspect of the invention, there is provided a sulphonyl derivative of formula



wherein

$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group and

- 10  $R^3$  denotes a  $C_{1-4}$ -alkyl group wherein the hydrogen atoms are optionally wholly or partly replaced by fluorine atoms, or
- a phenyl group optionally substituted by one to three substituents selected from the group consisting of  $C_{1-3}$ -alkyl groups, halogen atoms, cyano groups and nitro groups, wherein the substituents are identical or different.

15 Background to the invention

- Quinazoline derivatives of general formula (I) are known from WO 02/50043 and WO 04/074263, which describe compounds with valuable pharmacological properties, including in particular an inhibitory effect on signal transduction mediated by tyrosinekinases and an inhibitory effect on signal transduction mediated by Epidermal
- 20 Growth Factor-Receptor (EGF-R). Therefore, compounds of this type are suitable for the treatment of diseases, particularly for the treatment of tumoral diseases, diseases

25771-1542


- 1e -

of the lungs and airways and diseases of the gastro-intestinal tract and the bile ducts and gall bladder.

WO 2002/50043 discloses a method of production in which aminocrotonylamino-substituted quinazolines (I) are prepared in a one-pot reaction from the corresponding  
5 aniline component (II), bromocrotonic acid, oxalyl chloride and a secondary amine (see Diagram 1).

- 2 -

Diagram 1:




- 5 The process is not well suited to technical use on an industrial scale, as the yields obtained are at most 50% and as a rule laborious purification by column chromatography is needed. Moreover the educt bromocrotonic acid is not commercially available in large amounts and the corresponding methyl bromocrotonate is only available with a purity of about 80%.

10

WO 2005/037824 describes an alternative process for preparing aminocrotonylamino-substituted quinazoline derivatives of general formula (I) by Wittig-Horner-Emmons reaction of dialkyl-phosphonoacetamido-substituted quinazolines (III) with a 2-aminoacetaldehyde (IV) (Diagram 2), while instead of the aldehyde (IV) it is possible to use the corresponding hydrate or an acetal (e.g. the diethylacetal corresponding to (IV)), from which the aldehyde is liberated (beforehand or *in situ*).

15

Diagram 2:




20

The educts of formula III may be obtained according to WO 2005/037824 as follows:

- 3 -

Diagram 3:



5

Both in the prior art described above and within the scope of the invention described hereinafter, the groups  $\text{R}_a$ ,  $\text{R}_b$ ,  $\text{R}_c$ ,  $\text{R}_d$ ,  $\text{R}^1$  and  $\text{R}^2$  have the following meanings:

$\text{R}_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group,

10

$\text{R}_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

15

$\text{R}_c$  denotes a methyl, ethyl or 2-methoxyethyl group or

$\text{R}_b$  and  $\text{R}_c$  together with the nitrogen atom to which these groups are bound denote a morpholino or homomorpholino group optionally substituted by one or two  $\text{C}_{1-3}$ -alkyl groups,

- 4 -


R<sub>d</sub> denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group, and

5

R<sup>1</sup> and R<sup>2</sup> each independently of one another denote a C<sub>1-4</sub>-alkyl group, for example each denote an ethyl group.

By a homomorpholino group is meant the next highest ring-homologue of the morpholino group, namely the group of formula

10



Starting from commercially obtainable 4-chloro-anthranilic acid (V), reacting with formamidine acetate (step a) produces the quinazolinone (VI), which is then nitrated using sulphuric acid and concentrated nitric acid (step b). The desired regioisomer (VII) is then chlorinated using thionyl chloride in acetonitrile (step c) and the chlorination product (IX) is reacted *in situ* with the corresponding amine R<sub>a</sub>-NH<sub>2</sub> (step d). The compound of formula (X) thus obtained is reacted by base-catalysed nucleophilic substitution with R<sub>d</sub>-OH to form the compound (XI) (step e), which is in turn converted by hydrogenation into the corresponding aminoquinazoline (XII) (step f). The aminoquinazoline (XII) is then converted by reaction with a di-(C<sub>1-4</sub>-alkyl)-phosphonoacetic acid, e.g. with diethylphosphonoacetic acid, in suitable solvents such as tetrahydrofuran (THF), dimethylformamide (DMF) or ethyl acetate, after corresponding activation, for example with 1,1-carbonyldiimidazole, 1,1-carbonylditriazole or propanephosphonic anhydride, into the dialkyl-phosphonoacetamido-substituted quinazoline (III) needed for the Wittig-Horner-Emmons reaction.

20

25

The process described in WO 2005/037824 also has a number of serious disadvantages for technical use. For example, the use of thionyl chloride in Step (c) is problematic for safety reasons. The cyclic or heterocyclic alcohols needed to

30


- 5 -

introduce the group  $R_d$  in an excess of about 2 equivalents (eq) are starting materials which are difficult to obtain or expensive, and phase transfer catalysis, for example using 18-crown-6, is also needed to react them according to step (e) in Diagram 3 on an industrial scale. The reaction product has to be purified by recrystallisation to  
 5 eliminate the phase transfer catalyst. The hydrogenation of step (f) is carried out with the addition of acetic acid, if the educt contains a chlorine atom, so as to prevent the formation of dechlorinated by-products which are difficult to strip out. The addition of acetic acid causes traces of the nickel needed as catalyst to be dissolved, and this is then entrained into the final step and gives rise to a heavy metal problem which is  
 10 unacceptable for pharmaceutical use. Moreover, the throughput of individual partial reactions is in need of improvement; for example the throughput in step (e) according to Diagram 3 is only 1/60 (1kg of starting material require a reactor volume of 60 l).

In the light of the disadvantages of the known production method as described  
 15 above, the problem of the present invention is to provide an improved method, suitable for synthesis on an industrial scale, which permits the safe preparation of aminocrotonylamino-substituted quinazoline derivatives (I) using easily obtainable starting materials of high purity and at a lower technical cost.

#### 20 Detailed description of the invention


The problem stated above is solved according to the invention by the following process for preparing a compound of general formula



wherein  $R_a$  to  $R_d$  are as hereinbefore defined, comprising the following steps  
 25 **(embodiment A):**

- 6 -


a) reacting 7-chloro-6-nitro-3*H*-quinazolin-4-one



(VII)


with a primary amine of formula  $R_a-NH_2$  (XV), wherein  $R_a$  denotes a benzyl,  
5 1-phenylethyl or 3-chloro-4-fluorophenyl group, in the presence of  $POCl_3$ ,

b) converting the resulting compound of general formula



(X)

into the sulphonyl derivative of formula



(XIII)

10


wherein

$R^3$  denotes a  $C_{1-4}$ -alkyl group wherein the hydrogen atoms may be wholly or partly replaced by fluorine atoms, or

15 a phenyl group optionally substituted by one to three substituents selected from  $C_{1-3}$ -alkyl groups, halogen atoms, particularly fluorine, chlorine or bromine atoms, cyano or nitro groups, while the substituents may be identical or different, and wherein  $R_a$  in the two formulae (X) and (XIII) has the meanings given under a),

- 7 -

c) converting the sulphonyl derivative of formula (XIII) into a compound of formula




by reacting with an alcohol of formula R<sub>d</sub>-OH (XVI) in the presence of a base,

5

wherein R<sub>a</sub> has the meanings given under a) and R<sub>d</sub> denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,


10

d) reducing the compound of formulae (XI) thus obtained to the amino derivative of formula



15 wherein R<sub>a</sub> has the meanings given under a) and R<sub>d</sub> has the meanings given under c),

e) converting the amino derivatives of formula (XII) into the phosphonic ester of formula




20

- 8 -

wherein  $R_a$  has the meanings given under a),  $R_d$  has the meanings given under c), and  $R^1$  and  $R^2$  each independently of one another denote a  $C_{1-4}$ -alkyl group, but preferably ethyl groups,

- 5 f) reacting the resulting phosphonic ester of formula (III) with a hydrogen sulphite adduct of formula



wherein

$M^+$  denotes a cation, for example the sodium ion, or a proton and

10

$R_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,


- 15  $R_c$  denotes a methyl, ethyl or 2-methoxyethyl group or

$R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound represent a morpholino or homomorpholino group optionally substituted by one or two  $C_{1-3}$ -alkyl groups,

20

in the manner of a Wittig-Horner-Emmons reaction.

In another aspect the invention relates to the sulphonyl derivatives of formula



25 wherein

$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group and

R<sup>3</sup> denotes a C<sub>1-4</sub>-alkyl group wherein the hydrogen atoms may be wholly or partly replaced by fluorine atoms, or  
a phenyl group optionally substituted by one to three substituents selected from C<sub>1-3</sub>-  
5 alkyl groups, halogen atoms, particularly fluorine, chlorine or bromine atoms, cyano  
or nitro groups, while the substituents may be identical or different,

which are valuable synthesis components for preparing the pharmacologically active  
quinazoline derivatives of general formula (I).

10

For example R<sup>3</sup> in formula (XIII) may denote the p-toluenesulphonyl, p-bromo-  
benzenesulphonyl, phenyl, p-nitro-benzenesulphonyl, methylsulphonyl,  
trifluoromethylsulphonyl, nonafluorobutylsulphonyl or 2,2,2-trifluoroethanesulphonyl  
group.

15

The following are mentioned as particularly preferred compounds of formula (XIII):

(1) 4-(3-chloro-4-fluoro-phenylamino)-7-(4-methyl-phenylsulphonyl)-6-nitro-  
quinazoline,

20

(2) 4-(3-chloro-4-fluoro-phenylamino)-7-(4-bromo-phenylsulphonyl)-6-nitro-  
quinazoline,

(3) 4-(3-chloro-4-fluoro-phenylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline,

25

(4) 4-(3-chloro-4-fluoro-phenylamino)-7-(4-nitro-phenylsulphonyl)-6-nitro-  
quinazoline,

(5) 4-(3-chloro-4-fluoro-phenylamino)-7-(methylsulphonyl)-6-nitro-quinazoline,

30

(6) 4-(3-chloro-4-fluoro-phenylamino)-7-(trifluoromethylsulphonyl)-6-nitro-  
quinazoline,

- (7) 4-(3-chloro-4-fluoro-phenylamino)-7-(nonafluorobutylsulphonyl)-6-nitro-quinazoline,
- (8) 4-(3-chloro-4-fluoro-phenylamino)-7-(2,2,2-trifluoroethanesulphonyl)-6-nitro-quinazoline
- 5 (9) 4-(benzylamino)-7-(4-methyl-phenylsulphonyl)-6-nitro-quinazoline,
- (10) 4-(benzylamino)-7-(4-bromo-phenylsulphonyl)-6-nitro-quinazoline,
- 10 (11) 4-(benzylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline,
- (12) 4-(benzylamino)-7-(4-nitro-phenylsulphonyl)-6-nitro-quinazoline,
- 15 (13) 4-(benzylamino)-7-(methylsulphonyl)-6-nitro-quinazoline,
- (14) 4-(benzylamino)-7-(trifluoromethylsulphonyl)-6-nitro-quinazoline,
- (15) 4-(benzylamino)-7-(nonafluorobutylsulphonyl)-6-nitro-quinazoline,
- 20 (16) 4-(benzylamino)-7-(2,2,2-trifluoroethanesulphonyl)-6-nitro-quinazoline,
- (17) 4-(1-phenylethylamino)-7-(4-methyl-phenylsulphonyl)-6-nitro-quinazoline,
- 25 (18) 4-(1-phenylethylamino)-7-(4-bromo-phenylsulphonyl)-6-nitro-quinazoline,
- (19) 4-(1-phenylethylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline,
- (20) 4-(1-phenylethylamino)-7-(4-nitro-phenylsulphonyl)-6-nitro-quinazoline,
- 30 (21) 4-(1-phenylethylamino)-7-(methylsulphonyl)-6-nitro-quinazoline,
- (22) 4-(1-phenylethylamino)-7-(trifluoromethylsulphonyl)-6-nitro-quinazoline,

(23) 4-(1-phenylethylamino)-7-(nonafluorobutylsulphonyl)-6-nitro-quinazoline and

(24) 4-(1-phenylethylamino)-7-(2,2,2-trifluoroethanesulphonyl)-6-nitro-quinazoline.


5

The educts used in the process according to the invention and not described in more detail are either known from the literature or may be prepared from precursors known from the literature by simple analogous methods. For example, the sodium hydrogen sulphite adduct of formula (XIV) may be obtained from the corresponding acetal by liberating the corresponding aldehyde in hydrochloric acid solution and subsequent precipitation by the addition of NaHSO<sub>3</sub> solution. The diethylacetal may be prepared from the corresponding aldehyde by conventional methods.

Specifically, the invention described in embodiment A has the following partial aspects B, C and D:

15

### B) Preparation of the synthesis component of formula



wherein


20 R<sub>a</sub> denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group and

R<sub>d</sub> denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group, by

25

a) reacting 7-chloro-6-nitro-3*H*-quinazolin-4-one

- 12 -




(VII)

with a primary amine of formula  $R_a-NH_2$  (XV), wherein  $R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group, in the presence of  $POCl_3$ ,


5

b) converting the resulting compound of general formula



(X)

into the sulphonyl derivative of formula



(XIII)

10


wherein

$R^3$  denotes a  $C_{1-4}$ -alkyl group wherein the hydrogen atoms may be wholly or partly replaced by fluorine atoms, or

15 a phenyl group optionally substituted by one to three substituents selected from  $C_{1-3}$ -alkyl groups, halogen atoms, particularly fluorine, chlorine or bromine atoms, cyano or nitro groups, wherein the substituents may be identical or different, and wherein  $R_a$  in the two formulae (X) and (XIII) has the meanings given under a),

c) converting the sulphonyl derivative of formula (XIII) into a compound of formula


- 13 -



by reacting with an alcohol of formula  $R_d\text{-OH}$  (XVI) in the presence of a base,

- 5 wherein  $R_a$  has the meanings given under a) and  $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group.

10 **C) Preparing the synthesis component of formula**



wherein

$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group,


- 15  $R_d$  denotes a cyclopropylmethoxy, cyclobutyloxy, cyclopentyloxy, tetrahydrofuran-3-yl-oxy, tetrahydrofuran-2-yl-methoxy, tetrahydrofuran-3-yl-methoxy, tetrahydro-pyran-4-yl-oxy or tetrahydropyran-4-yl-methoxy group, and

$R^1$  and  $R^2$  each independently of one another denote a  $C_{1-4}$ -alkyl group,

20

by preparing the synthesis component of formula

- 14 -



wherein


$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group and

- 5  $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

according to the process described under B) and subsequently

10

d) reducing the compound of formulae (XI) thus obtained to the amino derivative of formula




15 wherein

$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group and

- $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or  
 20 tetrahydropyran-4-yl-methyl group, and


e) converting the amino derivatives of formula (XII) into the phosphonic ester of formula

- 15 -



wherein  $R_a$  and  $R_d$  have the meanings given under d), and  $R^1$  and  $R^2$  each independently of one another denote a  $C_{1-4}$ -alkyl group.

5 **D) Preparing a compound of general formula**



wherein

$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group,

10  $R_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

$R_c$  denotes a methyl, ethyl or 2-methoxyethyl group or


15

$R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denote a morpholino or homomorpholino group optionally substituted by one or two  $C_{1-3}$ -alkyl groups and

20  $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group, by

f) reacting a phosphonic ester of formula

- 16 -



wherein


$R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group,

- 5  $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group and

$R^1$  and  $R^2$  each independently of one another denote a  $C_{1-4}$ -alkyl group,

10

with the hydrogen sulphite adduct of formula



wherein

$M^+$  denotes a cation, for example the sodium ion or a proton,

15

$R_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

- 20  $R_c$  denotes a methyl, ethyl or 2-methoxyethyl group or

$R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denote a morpholino or homomorpholino group optionally substituted by one or two  $C_{1-3}$ -alkyl groups,

25

in the manner of a Wittig-Horner-Emmons reaction.

The reaction (a) of 7-chloro-6-nitro-3*H*-quinazolin-4-one (VII) with a primary amine of formula (XV) for preparing a compound of formula (X) is carried out in a suitable solvent, for example in acetonitrile, dioxane, THF, or mixtures thereof, e.g. in a mixture of acetonitrile, dioxane and in the presence of 1-2 equivalents POCl<sub>3</sub>, at a temperature of 50 to 80 °C.

The conversion (b) of a compound of formula (X) into the sulphonyl derivative of formula (XIII) is carried out in a suitable solvent, for example in acetonitrile, dioxane, THF, DMF, dimethylacetamide (DMA), N-methylpyrrolidone (NMP) or mixtures thereof, e.g. in a mixture of DMF and NMP by the addition of 1 to 2 equivalents of a corresponding sulphinic acid salt, for example the sodium salt of benzenesulphinic acid, at a temperature between 50°C and the boiling temperature of the solvent used, preferably at a temperature between 80 and 100°C.

The conversion (c) of the sulphonyl derivative of formula (XIII) into a compound of formula (XI) is carried out in a suitable solvent, for example in acetonitrile, dioxane, THF, DMF, DMA, diglyme, tert-butanol or mixtures thereof, by the addition of 1 to 1.5 equivalents of an alcohol of formula R<sub>d</sub>-OH (XVI) and subsequent batchwise addition of 2 to 4 equivalents of a strong base, for example powdered NaOH, KOH or LiOH, or potassium-tert.-butoxide, sodium-tert.-butoxide, lithium-tert.-butoxide, potassium-tert.-amylate, sodium-tert.-amylate or lithium-tert.-amylate as a solid or by dropwise addition of a solution of these bases in tert. butanol, THF or DMF, at a temperature between 0 and 100°C, preferably between 10 and 50°C, initially choosing a low temperature and raising the temperature after the addition of the base to complete the reaction.

The reduction (d) of a compound of formula (XI) to the amino derivative of formula (XII) is preferably carried out by catalytic hydrogenation with hydrogen in the presence of a catalyst such as Raney nickel, palladium/charcoal or platinum in a solvent such as methanol, ethanol, ethyl acetate, DMF, DMA, NMP, DMF/acetone or DMF/methanol, optionally with the addition of an acid such as acetic acid or an acid

salt at temperatures between 0 and 100°C, for example at temperatures between 0 and 80°C, but preferably at a temperature between 20 and 50°C, and under a hydrogen pressure of 1 to 10 bar, for example 1 to 7 bar, but preferably at a pressure of from 3 to 5 bar.

5

The conversion (e) of the amino derivative of formula (XII) into the phosphonic ester of formula (III) is carried out by reacting with 1.0 – 2.0 equivalents of a di-(C<sub>1-4</sub>-alkyl)-phosphonoacetic acid, preferably with diethylphosphonoacetic acid, in a suitable solvent such as THF, DMF, toluene, ethyl acetate, methyl-tert.-butylether (MTBE) or mixtures thereof, e.g. in MTBE/THF, after corresponding activation at temperatures between 0°C and 100°C. The activation may be carried out using any of the current methods of amide linking, i.e. for example with 1,1-carbonyldiimidazole, 1,1-carbonylditriazole, DCC (N,N-dicyclohexylcarbodiimide), EDC (N'-(dimethylaminopropyl)-N-ethylcarbodiimide), TBTU O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate, thiazolidine-2-thione, or by conversion into the corresponding acid chloride, possibly using thionyl chloride or phosphorus oxychloride. Optionally the activation is carried out using organic bases such as triethylamine or pyridine, while DMAP (dimethylaminopyridine) may additionally be added.

20

The reaction (f) of the phosphonic ester of formula (III) with the hydrogen sulphite adduct of formula (XIV) is carried out in a suitable solvent such as methanol, ethanol, THF, DMF, toluene, ethyl acetate and acetonitrile or mixtures thereof or in a binary or ternary mixture with water, preferably in ethanol or ethanol/water, with the addition of a suitable base, e.g. sodium carbonate, lithium hydroxide, sodium hydroxide or potassium hydroxide, optionally with the addition of a stabilising salt such as lithium chloride, at a temperature of 0°C- 50°C, using 1 - 2 equivalents, preferably 1.2 -1.6, e.g. 1.4 equivalents of the hydrogen sulphite adducts in aqueous solution.

25  
30

Preferred embodiments of the process according to the invention with all the partial steps (a) to (f) as well as the partial aspects B, C and D relate to the preparation of compounds of formula (I), wherein

- 19 -

$R_a$  denotes a 3-chloro-4-fluorophenyl group and  $R_b$ ,  $R_c$  and  $R_d$  are defined as mentioned in embodiment A (**embodiment F**), or

5  $R_b$  and  $R_c$  in each case denote a methyl group and  $R_a$  and  $R_d$  are defined as mentioned in embodiment A (**embodiment H**), or

$R_a$  denotes a 3-chloro-4-fluorophenyl group,  $R_b$  and  $R_c$  each denote a methyl group and  $R_d$  are defined as mentioned in embodiment A (**embodiment I**), or

10  $R_d$  denotes a tetrahydrofuran-3-yl or tetrahydropyran-4-yl group and  $R_a$ ,  $R_b$  and  $R_c$  are defined as mentioned in embodiment A (**embodiment J**), or

$R_a$  denotes a 3-chloro-4-fluorophenyl group,  $R_b$  and  $R_c$  in each case denote a methyl group and

15  $R_d$  denotes a tetrahydrofuran-3-yl or tetrahydropyran-4-yl group (**embodiment K**), or

$R_a$  denotes a 3-chloro-4-fluorophenyl group,  $R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denotes a morpholine group and

20  $R_d$  denotes a tetrahydrofuran-3-yl or tetrahydropyran-4-yl group (**embodiment L**), or


$R_a$  denotes a 3-chloro-4-fluorophenyl group,  $R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denotes a homomorpholine group and

$R_d$  denotes a tetrahydrofuran-3-yl or tetrahydropyran-4-yl group (**embodiment M**),

25 while  $R_a$ ,  $R_b$ ,  $R_c$  and  $R_d$  in partial steps (a) to (f), in each case corresponding to embodiments F to M, assume the meanings given therein.

The process according to the invention is used to particular advantage to prepare the compound

- 20 -



4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline,

5 or the corresponding R-enantiomer.

The aminocrotonylamino-substituted quinazoline derivatives of formula (I) obtained by the process according to the invention may subsequently be converted by known methods into the salts thereof, particularly into physiologically acceptable salts, for example into fumarates, tartrates or maleates. The conversion of the compound 4-  
 10 [(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-dimethylamino)-1-oxo-2-buten-1-yl]amino]-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline obtained according to the invention into the corresponding dimaleate is preferred, as described in WO 2005/037824.

15

The following Examples are intended to illustrate the invention in more detail:

**Example 1**4-(3-Chloro-4-fluoro-phenylamino)-7-chloro-6-nitro-quinazoline

20 g 7-chloro-6-nitro-3H-quinazolin-4-one are suspended in 80 ml acetonitrile and combined with 16.5 g phosphorus oxychloride. Then 10.8 g triethylamine are slowly  
5 added dropwise and the mixture is heated to about 80°C. After 5 hours a solution of 15.5 g 3-chloro-4-fluoroaniline in 100 ml dioxane is added dropwise and the mixture is stirred for another hour. Then 80 ml of water is added, the mixture is cooled to 20°C and made slightly alkaline with KOH solution. The suspension is suction filtered, washed with water and ethanol and dried at 50°C in vacuo.  
10 Yield: 29.07 g (89.5% of theoretical/dioxane solvate)  
m.p.: 272 – 274°C

**Example 2**4-(3-Chloro-4-fluoro-phenylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline

15 500 g 4-(3-chloro-4-fluoro-phenylamino)-7-chloro-6-nitro-quinazoline and 302 g (1.3 eq) benzenesulphinic acid sodium salt are suspended at 20°C in 1500 ml DMF, heated to 90°C and kept for 6h at this temperature. After cooling the reaction mixture the suspension is suction filtered and the residue is rinsed with 1.5 l methanol, 10 l water and 0.5 l methanol. The residue is dried at 50°C for about 12 h  
20 under reduced pressure.  
Yield: 631.2 g (86.2 % of theoretical/DMF solvate).  
m.p.: 284 -286°C

**Example 3**25 4-[(3-Chloro-4-fluorophenyl)amino]-6-nitro-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline

810 g of 4-(3-chloro-4-fluoro-phenylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline and 175.5 g (S)-3-hydroxytetrahydrofuran (1.3 eq) are placed at 20°C in 1.04 l tert-butanol and 198 ml DMF, 2556 g K-tert.-butoxide in THF (24 %) (3.6 eq) are added dropwise at 20°C and then stirred for 4 h at 25°C. After a further 2 h at 40°C the  
30 mixture is heated to 45°C for about 2 h. 2.8 l of water are added and then about 3 l solvent are distilled off under reduced pressure. 2.8 l water are added again and about 900 ml solvent are distilled off under reduced pressure. After the addition of 1.6 l methanol the mixture is cooled to 20°C. The suspension is suction filtered and

- 22 -

rinsed with a mixture of 3.2 l water and 1.6 l methanol. The residue is dried overnight at 50°C under reduced pressure.

Yield: 598.6 g (89.6 % of theoretical).

m.p.: 238 – 240°C

5

#### Example 4

##### 4-[(3-Chloro-4-fluorophenyl)amino]-6-amino-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline

100 g of 4-[(3-chloro-4-fluorophenyl)amino]-6-nitro-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline are hydrogenated in 400 ml DMF in the presence of 33.1 g Raney nickel and 18.7 g ammonium chloride at 40°C, until the calculated amount of hydrogen has been taken up. The catalyst is filtered off and the filtrate is added dropwise to 1.2 l water. The suspension is stirred for 2.5 h at 0°C, suction filtered and washed with 500 ml of water. The residue is dried overnight at 55°C under reduced pressure.


Yield: 84.36 g (97.1 % of theoretical).

m.p.: 120 – 130°C

#### Example 5

##### Diethyl {[4-(3-chloro-4-fluoro-phenylamino)-7-((S)-tetrahydrofuran-3-yloxy)-quinazolin-6-ylcarbamoyl]-methyl}-phosphonate

20



3.58 kg 1,1-carbonyldiimidazole (22.16 mol) are placed in 12.8 l of tetrahydrofuran and combined at 40°C with 4.52 kg (22.16 mol) diethylphosphonoacetic acid, dissolved in 6.5 l tetrahydrofuran. The mixture is stirred for 30 minutes at 40°C. The solution thus obtained is designated solution A.

25

- 23 -

6.39 kg (17.05 mol) of N<sup>4</sup>-(3-chloro-4-fluoro-phenyl)-7-(tetrahydrofuran-3-yloxy)quinazoline-4,6-diamine are placed in 26.5 l tetrahydrofuran and combined at 40°C with solution A and stirred for 2 hours at 30°C. 64 l of tert.-butylmethylether are added to the suspension and after cooling to 20°C the precipitate is removed by centrifuging. It is washed with a mixture of 16 l tetrahydrofuran and 16 l tert.-butylmethylether and then with 32 l water and dried at 50°C.

Yield: 6.58 kg (69.8%) white crystals. Content: HPLC 99.1 FI%

### Example 6

#### 10 Dimethylaminoacetaldehyde-hydrogen sulphite adduct


40 g of dimethylaminoacetaldehyde diethylacetal are heated to 40°C in a mixture of 48 g conc. hydrochloric acid and 20 ml of water for 3 h. Then a solution of 42.4 g sodium pyrosulphite in 72 ml of water (sodium hydrogen sulphite solution) is added dropwise and the mixture is stirred for 1 h. 200 ml of ethanol are added and then the mixture is stirred for 2 h at 0°C. The suspension is suction filtered, washed with 160 ml of ethanol and dried at 45°C in vacuo.

Yield: 42.5 g (89.6 % of theoretical)

decomp.: from 180°C

### 20 Example 7

#### (E)-4-Dimethylamino-but-2-enoic acid-[4-(3-chloro-4-fluoro-phenylamino)-7-((S)-tetrahydrofuran-3-yloxy)-quinazolin-6-yl]-amide



25 10 g of diethyl {[4-(3-chloro-4-fluoro-phenylamino)-7-((S)-tetrahydrofuran-3-yloxy)-quinazolin-6-ylcarbamoyl]-methyl}-phosphonate and 0.8 g lithium chloride are suspended in 60 ml of ethanol and cooled to -5°C. 11 g of 45% potassium hydroxide

- 24 -

solution is added dropwise first of all and then 4.8 g dimethylaminoacetaldehyde-hydrogen sulphite adduct in 48 ml of water is added. The reaction solution is stirred for 1 h and then 60 ml of water are added. The suspension is suction filtered, washed with 40 ml of water and dried in vacuo at 45°C.

5 Yield: 8 g (91 % of theoretical)


m.p.: 100 -102°C

25771-1542

- 25 -

CLAIMS:

1. A process for preparing a compound of general formula



R<sub>a</sub> denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group,

- 5 R<sub>b</sub> denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,


R<sub>c</sub> denotes a methyl, ethyl or 2-methoxyethyl group or

- 10 R<sub>b</sub> and R<sub>c</sub> together with the nitrogen atom to which these groups are bound denotes a morpholino or homomorpholino group optionally substituted by one or two C<sub>1-3</sub>-alkyl groups, and

R<sub>d</sub> denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

- 15 comprising the following process steps:

a) reacting 7-chloro-6-nitro-3*H*-quinazolin-4-one




25771-1542


- 26 -

with a primary amine of formula  $R_a\text{-NH}_2$  (XV), wherein  $R_a$  denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group, in the presence of  $\text{POCl}_3$ ,

b) converting the resulting compound of general formula



5 into the sulphonyl derivative of formula



wherein


$R^3$  denotes a  $C_{1-4}$ -alkyl group wherein the hydrogen atoms are optionally wholly or partly replaced by fluorine atoms, or

10 a phenyl group optionally substituted by one to three substituents selected from the group consisting of  $C_{1-3}$ -alkyl groups, halogen atoms, cyano groups and nitro groups, wherein the substituents are identical or different, and wherein  $R_a$  in the two formulae (X) and (XIII) has the meanings given under a),

c) converting the sulphonyl derivative of formula (XIII) into a compound of formula

25771-1542

- 27 -




by reacting with an alcohol of formula  $R_d\text{-OH}$  (XVI) in the presence of a base,

wherein  $R_a$  has the meanings given under a) and  $R_d$  denotes a cyclopropylmethyl, cyclobutyl, cyclopentyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl,

5 tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

d) reducing the compound of formulae (XI) thus obtained to the amino derivative of formula




10 wherein  $R_a$  has the meanings given under a) and  $R_d$  has the meanings given under c),

e) converting the amino derivatives of formula (XII) into the phosphonic ester of formula


25771-1542

- 28 -



wherein  $R_a$  has the meanings given under a) and  $R_d$  has the meanings given under c),  $R^1$  and  $R^2$  each independently of one another denote a  $C_{1-4}$ -alkyl group,

f) reacting the resulting phosphonic ester of formula (III) with a hydrogen sulphite adduct of formula



wherein

$M^+$  denotes a cation or a proton and

$R_b$  denotes a methyl, ethyl, isopropyl, cyclopropyl, 2-methoxyethyl, tetrahydrofuran-3-yl, tetrahydrofuran-2-yl-methyl, tetrahydrofuran-3-yl-methyl, tetrahydropyran-4-yl or tetrahydropyran-4-yl-methyl group,

$R_c$  denotes a methyl, ethyl or 2-methoxyethyl group or


$R_b$  and  $R_c$  together with the nitrogen atom to which these groups are bound denote a morpholino or homomorpholino group optionally substituted by one or two  $C_{1-3}$ -alkyl groups,

in the manner of a Wittig-Horner-Emmons reaction.

25771-1542


- 29 -

2. A process according to claim 1, wherein, in step b), the one to three optional halogen atom substituents on the phenyl group of R<sup>3</sup> are independently fluorine, chlorine or bromine.
3. A process according to claim 1 or 2, wherein, in step f), the cation of M<sup>+</sup> is a sodium ion.
4. A process according to claim 1, wherein the compound of general formula I is:



- 4-[(3-chloro-4-fluorophenyl)amino]-6-[[4-(N,N-di-methyl-amino)-1-oxo-2-buten-1-yl]amino]-7-((S)-tetrahydrofuran-3-yloxy)-quinazoline,
- or the corresponding R-enantiomer.

5. A sulphonyl derivative of formula



wherein

- R<sub>a</sub> denotes a benzyl, 1-phenylethyl or 3-chloro-4-fluorophenyl group and

25771-1542

- 30 -

R<sup>3</sup> denotes a C<sub>1-4</sub>-alkyl group wherein the hydrogen atoms are optionally wholly or partly replaced by fluorine atoms, or

a phenyl group optionally substituted by one to three substituents selected from the group consisting of C<sub>1-3</sub>-alkyl groups, halogen atoms, cyano groups and nitro groups,  
5 wherein the substituents are identical or different.

6. A sulphonyl derivative according to claim 5, wherein the one to three optional halogen substituents or the phenyl group of R<sup>3</sup> are independently fluorine, chlorine or bromine.

7. A sulphonyl derivative of formula (XIII) according to claim 5, wherein R<sup>3</sup>  
10 taken together with the sulphonyl group to which the R<sup>3</sup> group is attached denotes a p-toluenesulphonyl, p-bromo-benzenesulphonyl, phenylsulphonyl, p-nitro-benzenesulphonyl, methylsulphonyl, trifluoromethylsulphonyl, nonafluorobutylsulphonyl or 2,2,2-trifluoroethanesulphonyl group.

8. A sulphonyl derivative of formula (XIII) according to claim 5 selected  
15 from the group consisting of:

- (1) 4-(3-chloro-4-fluoro-phenylamino)-7-(4-methyl-phenylsulphonyl)-6-nitro-quinazoline,
- (2) 4-(3-chloro-4-fluoro-phenylamino)-7-(4-bromo-phenylsulphonyl)-6-nitro-quinazoline,
- 20 (3) 4-(3-chloro-4-fluoro-phenylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline,
- (4) 4-(3-chloro-4-fluoro-phenylamino)-7-(4-nitro-phenylsulphonyl)-6-nitro-quinazoline,
- (5) 4-(3-chloro-4-fluoro-phenylamino)-7-(methylsulphonyl)-6-nitro-quinazoline,
- (6) 4-(3-chloro-4-fluoro-phenylamino)-7-(trifluoromethylsulphonyl)-6-nitro-quinazoline,

25771-1542

- 31 -

- (7) 4-(3-chloro-4-fluoro-phenylamino)-7-(nonafluorobutylsulphonyl)-6-nitro-quinazoline,
- (8) 4-(3-chloro-4-fluoro-phenylamino)-7-(2,2,2-trifluoroethanesulphonyl)-6-nitro-quinazoline
- 5 (9) 4-(benzylamino)-7-(4-methyl-phenylsulphonyl)-6-nitro-quinazoline,
- (10) 4-(benzylamino)-7-(4-bromo-phenylsulphonyl)-6-nitro-quinazoline,
- (11) 4-(benzylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline,
- (12) 4-(benzylamino)-7-(4-nitro-phenylsulphonyl)-6-nitro-quinazoline,
- (13) 4-(benzylamino)-7-(methylsulphonyl)-6-nitro-quinazoline,
- 10 (14) 4-(benzylamino)-7-(trifluoromethylsulphonyl)-6-nitro-quinazoline,
- (15) 4-(benzylamino)-7-(nonafluorobutylsulphonyl)-6-nitro-quinazoline,
- (16) 4-(benzylamino)-7-(2,2,2-trifluoroethanesulphonyl)-6-nitro-quinazoline,
- (17) 4-(1-phenylethylamino)-7-(4-methyl-phenylsulphonyl)-6-nitro-quinazoline,
- (18) 4-(1-phenylethylamino)-7-(4-bromo-phenylsulphonyl)-6-nitro-quinazoline,
- 15 (19) 4-(1-phenylethylamino)-7-(phenylsulphonyl)-6-nitro-quinazoline,
- (20) 4-(1-phenylethylamino)-7-(4-nitro-phenylsulphonyl)-6-nitro-quinazoline,
- (21) 4-(1-phenylethylamino)-7-(methylsulphonyl)-6-nitro-quinazoline,
- (22) 4-(1-phenylethylamino)-7-(trifluoromethylsulphonyl)-6-nitro-quinazoline,
- (23) 4-(1-phenylethylamino)-7-(nonafluorobutylsulphonyl)-6-nitro-quinazoline and
- 20 (24) 4-(1-phenylethylamino)-7-(2,2,2-trifluoroethanesulphonyl)-6-nitro-quinazoline.

