



US00RE46211E

(19) **United States**
 (12) **Reissued Patent**
Weisbart

(10) **Patent Number:** **US RE46,211 E**
 (45) **Date of Reissued Patent:** ***Nov. 22, 2016**

(54) **DELIVERY SYSTEM USING MAB 3E10 AND
 MUTANTS AND/OR FUNCTIONAL
 FRAGMENTS THEREOF**

(75) Inventor: **Richard Weisbart**, Los Angeles, CA
 (US)

(73) Assignees: **The Regents of the University of
 California**, Oakland, CA (US); **The
 United States of America as
 represented by the Department of
 Veterans Affairs Office of the General
 Counsel**, Washington, DC (US)

(*) Notice: This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/404,154**

(22) Filed: **Mar. 13, 2009**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **7,189,396**
 Issued: **Mar. 13, 2007**
 Appl. No.: **10/192,315**
 Filed: **Jul. 10, 2002**

U.S. Applications:

(63) Continuation-in-part of application No. 09/142,142,
 filed as application No. PCT/US97/03785 on Mar. 7,
 1997, now abandoned.

(60) Provisional application No. 60/013,297, filed on Mar.
 8, 1996.

(51) **Int. Cl.**
A61K 39/395 (2006.01)
C07K 16/18 (2006.01)
C07K 16/44 (2006.01)

(52) **U.S. Cl.**
 CPC **C07K 16/44** (2013.01); **C07K 2317/565**
 (2013.01); **C07K 2317/622** (2013.01); **C07K**
2317/77 (2013.01); **C07K 2317/82** (2013.01)

(58) **Field of Classification Search**
 CPC **C07K 16/44**; **C07K 2317/565**; **C07K**
2317/82; **C07K 2317/77**; **C07K 2317/622**
 USPC **424/133.1**, **134.1**, **135.1**, **141.1**, **152.1**;
530/391.7

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,623,627 A	11/1986	Huang et al.	435/240
4,812,397 A *	3/1989	Weisbart	435/6
6,130,065 A	10/2000	Belt et al.	
6,232,444 B1 *	5/2001	Weisbart	530/387.2
7,381,799 B2 *	6/2008	Papathanassiou	530/387.1
2006/0003362 A1	1/2006	Zerangue	

FOREIGN PATENT DOCUMENTS

WO	WO 97/32602	*	3/1997
WO	WO 98/16247 A1	*	4/1998
WO	WO 98/29437 A2		7/1998
WO	WO 2005/117562 A2		12/2005

OTHER PUBLICATIONS

Arnon et al., "Targeted chemotherapy: drugs conjugated to anti-tumour antibodies", *Cancer Surveys* 1(3): 429-449 (1982).

Rudikoff, S. et al., *Proc. Nat. Acad. Sci. (USA)* [1982] 79:1979-1983.

Kong et al., "Mammalian Nucleoside Transporters", *Current Drug Metabolism*, vol. 5: 63-84 (2004).

* cited by examiner

Primary Examiner — Sharon Tucker

(74) *Attorney, Agent, or Firm* — DLA Piper LLP (US)

(57) **ABSTRACT**

A monoclonal antibody, 3E10, and active fragments thereof that selectively are transported in vivo to the nucleus of mammalian cells without cytotoxic effect are provided. The antibody and other molecules that bind to a variant of myosin IIb heavy chain found in the nucleus of skeletal muscle cells are useful as a non-viral delivery vector to target skeletal muscle in vivo. By contrast, in vitro the monoclonal antibody penetrates and is transported to the nucleus of multiple cell lines derived from different tissue types and can be used in screening tests to identify molecules that modulate growth of cells, such as cancer cells. Non-cytotoxic vectors for delivering a drug, polynucleotide or polypeptide selectively to skeletal muscle cells are also provided.

11 Claims, 5 Drawing Sheets

FIGURE 1A. Design of Constructs

SP-Vk Linker VH p53 peptide



FIGURE 1B. PCR Primers

Primer	Nucleotide Sequence (5'-3')
--------	-----------------------------

(for *pSG5*)

3E10-Fv-SP 5'-GAATTCATGGAGTCAGACACACTGCTG
CTATGGGTG-3'(SEQ ID NO:14)

3E10-Fv-HI 5'-GGATCCATGATGATGATGATGATGGTC-3'
(SEQ ID NO:15)

p53A-S 5'-GGATCCGCTCACTCCAGCCACCT-3'
(SEQ ID NO:16)

p53A-AS 5'-AGATCTTCAGTCTGAGTCAGGCCC-3'
(SEQ ID NO:17)

p53B-S 5'-GGATCCGAGCCAGGGGGGAGCA-3'
(SEQ ID NO:18)

p53B-AS 5'-AGATCTTCAGTCTGAGTCAGGCCC-3'
(SEQ ID NO:19)

p53C-S 5'-GGATCCCTGAATGAGGCCTTGGAAC-3'
(SEQ ID NO:20)

p53C-AS 5'-AGATCTTCAGTCTGAGTCAGGCCC-3'
(SEQ ID NO:21)

p53N-S 5'-GGATCCGAGGAGCCGCAGTCAGAT-3'
(SEQ ID NO:22)

p53N-AS 5'-AGATCTTCAAATATCGTCCGGGGACAG-3'
(SEQ ID NO:23)

(for *pPICZαA*)

3E10-Fv-S 5'-GAATCCGACATTGTGCTGACACAGT-3'
(SEQ ID NO:24)

3E10-Fv-AS 5'-CCGCGGTCAATGATGATGATGATGATG
ATGGGAGAC-3'(SEQ ID NO:25)

3E10-Fvp53A-S 5'-GAATCCGACATTGTGCTGACACAGT-3'
(SEQ ID NO:26)

3E10-Fvp53A-AS 5'-CCGCGGTCAGTCTGAGTCAGGCCC-3'
(SEQ ID NO:27)

3E10-Fvp53N-S 5'-GAATCCGACATTGTGCTGACACAGT-3'
(SEQ ID NO:28)

3E10-Fvp53N-AS 5'-CCGCGGTCAGTCTGAGTCAGGCCC-3'
(SEQ ID NO:29)

FIGURE 1C. Plasmid Constructs

<u>Plasmid</u>	<u>Host</u>	<u>p53 Peptide Plasmid</u>	<u>Insert</u>
<i>pScFv</i>	<i>pSG5</i>	---	
<i>pScFvp53A</i>	<i>pSG5</i>		aa 364-393
<i>pScFv-p53B</i>	<i>pSG5</i>		aa 358-393
<i>pScFv-p53C</i>	<i>pSG5</i>		aa 344-393
<i>pScFv-Mut-p53A*</i>	<i>pSG5</i>		aa 364-393
<i>pScFv-p53N</i>	<i>pSG5</i>		aa 2- 49
<i>pScFv</i>	<i>pPICZaA</i>		
<i>pScFv-p53A</i>	<i>pPICZaA</i>		aa 364-393
<i>pScFv-p53N</i>	<i>pPICZaA</i>		aa 2- 49

* *pScFv-Mut-p53A* contains the mutation R95Q in mAb 3E10 VH that prevents mAb 3E10Fv from penetrating living cells.




Figure 2






Figure 3

**Figure 4****Figure 5**

MYOSIN HEAVY CHAIN IIB (Homo Sapiens)

1 mqqgtledqii sanplleafg naktvrndns srfgkfirih fgatgklasa dietylleks
 61 rvtfqikaer syhifyqils nkkpelieml littncpydfa fvsqgeitvp siddqeelma
 121 tdsavdilgf tadekvaiyk ltgavmhygn mkfkqkgree qaepdgteva dkaayltsln
 181 sadllkslcy prvkvgnefv tkgqtvqqvy navgalakai yekmflwmvt ringqldtkq
 241 prqyfigvld iagfeifdfn sleqlcinfnt neklqqffnh hmfvleqeey kkegiwefi
 301 dfgmldlaaci eliekpmgif sileeeecmf katdtsfknk lyeqhlgsn nfgkpkpakg
 361 kpeahfslvh yagtdynia gwldknkdpl netvvglyqk samktlaflf sgaqtaeaeg
 421 gggkkggkkk gssfqtvsa frenlnklmt nlrsthphfv rciipnetkt pgamehelvl
 481 hqlrcngvle giricrkqfp srilyadfkq rykvlnasai peggfidskk aseklsgsie
 541 idhtqykfgh tkvffkagll gtleemrdek laqlitrtqa icrgflmrve frkmmerrres
 601 ifciqynira fmnvkhwpwm klyfkikp11 ksaetekema nmkeefektk eelakteakr
 661 keleekmvtl mgekndlqlq vqaeadalad aeercdqlik tkiqleakik evteraedee
 721 einaeltakk rkledccsel kkdiddlelt lakvekekha tenkvknlte emagldetia
 781 kltkekka1q eahqqtlddl qmeedkvnt1 tkaktkleqg vddlegsleg ekk1cmdler
 841 akrklegdlk laqestmdte ndkqqlnekl kkkefemsnl qgkiedeqal aiqlqkkike
 901 lqarieelee eieaerasra kaekqrsdls releeiserl eeaggatsaq iemnkcreae
 961 fqkmrrdlee stlqheataa alrkkhadsv aelgeqidsl qrvkqkleke kselkmeind
 1021 lasnmetvsk akanfekmcr tledqlseik tkeeeqqrli nelsagkarl htesgcfsrq
 1081 ldekdamvsq lsrgkqaftq qieelkrqle eetkakstla halqsarhdc dllreqyeee
 1141 qeakaelqrg mskansevaq wrtkyetdai qrteeleeak kklaqrlqda eehveavnsk
 1201 caslektkqr lqnevedlmi dversnaaci aldkkqrnfd kvlaewkqky eetqaeleas
 1261 qkesrsiste lfkvknayee sldhletlkr enkn1qqeis dlteqiaegg khihelekvk
 1321 kqldheksel qtsleeaeas leheegkilr iqlelnqvks eidrkiaekd eeldqlkrnh
 1381 lrvvesmqst ldaeirsrnd alrikkkmeg dlnemeiqln hanrqaaeal rnlrntqgil
 1441 kdtqlhllda irgqddlkeq lamverranl mqaeevelra slertergrk maeqelllas
 1501 ervql1htqn tslintkkkl etdisqigge medivqearn aekakkait daammaeelk
 1561 keqdtisahle rmkknmeqtv kdlqlrldea eqlalkggkk qiqklearvr eleseveseq
 1621 khnveavkql rkherrevkel tygteedrkn ilrlqdlvdk lqtkvkaykr qaceaceqsn
 1681 vn1akfrklq heleeakera diaesgvnkl rvksrevhtk visee

Figure 6

1

DELIVERY SYSTEM USING MAB 3E10 AND MUTANTS AND/OR FUNCTIONAL FRAGMENTS THEREOF

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in *italics* indicates the additions made by reissue; a claim printed with ~~strikethrough~~ indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

This application is a Continuation-in-Part Application of U.S. application Ser. No. 09/142,142, filed Nov. 9, 1998 now abandoned; which was the National Stage of International Application No. PCT/US97/03785 under 37 CFR § 371, filed Mar. 7, 1997 and relies for priority upon U.S. Provisional Application Ser. No. 60/013,297, filed Mar. 8, 1996.

This invention was made with Government support from the Department of Veterans Affairs. The Government has certain rights in this application.

FIELD OF THE INVENTION

The present invention relates to methods for the delivery of biologically active materials into cells, and compositions useful therefor.

BACKGROUND OF THE INVENTION

Autoantibodies to double stranded deoxyribose nucleic acid (dsDNA) are relatively specific for systemic lupus erythematosus (SLE) and are implicated in disease pathogenesis. Certain anti-DNA autoantibodies have been shown to penetrate cells and localize to the cell nucleus. Cellular penetration by anti-DNA antibodies was initially demonstrated in peripheral blood T-lymphocytes (see, for example, Okudaira, et al., in *Arthritis Rheum.* 30:669 (1987) and Alarcon-Segovia, et al., in *Clin. exp. Immunol.* 35:364 (1979)) and, subsequently, was shown to affect their function (see, for example, Okudaira, et al., *supra*, Alarcon-Segovia, et al., in *J. Immunol.* 122:1855 (1979), Alarcon-Segovia, et al., in *Clin. Immunol. Immunopath.* 23:22 (1982), Alarcon-Segovia and Llorente in *Clin. exp. Immunol.* 52:365 (1983), and Alarcon-Segovia, in *Clinics in Immunology and Allergy* 1:117 (1981)).

In some studies, antibody penetration was thought to be mediated by Fc receptors (see, for example, Llerena, et al., in *Immunology* 43:249 (1981) and Alarcon-Segovia, et al., in *Nature* 271:67 (1978)). For other anti-DNA antibodies, cellular penetration and translocation to the cell nucleus was thought to require the presence of DNA (see, for example, Okudaira, et al., *supra*). More recently, penetration of anti-DNA antibodies has been demonstrated in mesangial cells (Vlahakos, et al., in *T. Am. Soc. Nephrol.* 2(8):1345 (1992)). Anti-DNA antibodies have been shown to enter the nucleus of cultured mesangial and hepatoma cells in a time and temperature dependent manner (Yanase, et al., in *Lab. Invest.* 71:52 (1994)).

There are multiple mechanisms by which anti-DNA antibodies are thought to penetrate cells. Indeed, different antibodies may use different pathways. Since some anti-DNA antibodies have been shown to bind membrane proteins cross reactive with DNA, these proteins may be instrumental in cellular penetration (see, for example, Brentjens and Andres in *Kidney International* 35:954 (1989), Raz, et al., *J.*

2

Immunol. 142:3076 (1989), Madaio, et al., in *J. Immunol.* 138:2883 (1987), Faaber, et al., in *J. Clin. Invest.* 77:1824 (1986), Ben-Chetrit, et al., in *Clin. exp. Immunol.* 60:159 (1985), Jacob, et al., in *Proc. Natl. Acad. Sci. USA* 81:3843 (1984), Jacob, et al., in *Proc. Natl. Acad. Sci. USA* 86:4669.4669 (1989), Raz, et al., in *Eur. J. Immunol.* 23:383.383 (1993), and Jacob, et al., in *J. Clin. Invest.* 75:315 (1985)). In other cases, DNA binding proteins usually thought of as intracellular have been described in association with the membrane of some cells (see, for example, Bennett, et al., in *J. Clin. Invest.* 76:2182 (1985) and Refeneider, et al., in *Clin. Immunol. Immunopath.* 63:245 (1992)). Anti-DNA antibodies could form complexes with these proteins through their mutual binding to DNA.

For additional background information, see U.S. Pat. No. 4,812,397 and "DNA Mimics a Self-Protein That May Be a Target for Some Anti-DNA Antibodies in Systemic Lupus Erythematosus", *Journal Of Immunology*, pages 1987-1994 (Feb. 15, 1995), the contents of each of which are hereby incorporated by reference in their entirety.

Mutations in the p53 tumor suppressor protein are a frequent cause of cancer. Methods to regulate and restore the function of p53 are promising approaches to the treatment of cancer. Restoring p53 function by delivering functional p53 into cancer cells and delivering p53 peptides have been investigated in several studies. Various delivery vehicles have been used to deliver p53 and p53 peptides into cancer cells for restoration of p53 function. These include VP 22, a herpes simplex virus 1 protein (15,16), and the third alpha helix of Antennapedia homeodomain (8). The potential disadvantage of these vectors is that they are foreign proteins that may be immunogenic in humans.

In view of the availability of antibodies which are capable of penetrating cells, it would be desirable to selectively utilize such cell penetrating properties for the directed manipulation of biological materials.

BRIEF DESCRIPTION OF THE INVENTION

The invention overcomes these and other problems in the art by providing methods for transporting a biologically active molecule into the nucleus of a target cell in vitro. In the in vitro invention methods the biologically active molecule is combined with mAb 3E10 as produced by a hybridoma having ATCC accession number PTA 2439, or mutant or functional fragment thereof, and the target cell is contacted in vitro under suitable conditions with the resulting combination, thereby transporting the biologically active molecule into the nucleus of the target cell.

In another embodiment the invention provides in vivo methods for selectively transporting a biologically active molecule to the nucleus of skeletal muscle cells in a living subject. In the in vivo methods, the subject is administered a biologically active molecule combined with a transport molecule that specifically binds to the nuclear isoform of the heavy chain of myosin IIb, thereby selectively transporting the biologically active molecule into skeletal muscle cells of the subject.

In yet another embodiment, the invention provides vectors for selectively transporting a biologically active molecule into skeletal muscle cells contained in a mammalian subject. The invention vector comprises a transport molecule that, in a live subject, specifically binds to a nuclear isoform of myosin heavy chain IIb and a biologically active molecule chemically associated with the monoclonal antibody, or mutant or functional fragment thereof.

In still another embodiment, the invention provides antibodies that specifically bind to the newly discovered nuclear isoform of myosin IIb heavy chain and not to other isoforms of myosin IIb heavy chain.

DESCRIPTION OF THE FIGURES

FIG. 1A is a schematic representation of the plasmid constructs used in construction of fusion proteins encoding an scFv of mAb 3E10 and various polymers of the C-terminal region of p53 protein.

FIG. 1B shows the sense (S) and antisense (AS) nucleotide primers used for construction of the various plasmids illustrated in FIG. 1A and used in Example 8.

FIG. 1C shows the host plasmid and amino acid segments of p53 contained in the p53 Peptide insert in the various plasmids illustrated in FIG. 1A and used in Example 8.

FIG. 2 is a graph showing cytotoxicity of p53 transfection from pscFv-p53A, B, and C transfected into COS-7 cells.

FIG. 3 is a graph showing cytotoxicity of p53 transfection from pscFv-p53A, B, and C transfected into CHO cells.

FIG. 4 is a graph showing cytotoxicity of p53 from pscFv-p53A, B, and C transfected into SW480 cancer cells.

FIG. 5 is a graph showing the dose response of the cytotoxic effect of scFv-p53A.

FIG. 6 shows the amino acid sequence of a 200 kDa human skeletal muscle protein (SEQ ID NO: 13) identified as myosin IIb by Nano-LC/MS/MS. Underlining shows the 28 peptides identified by Nano-LC/MS/MS of rat skeletal muscle as identical with the human skeletal muscle protein.

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there are provided methods for the transport of biologically active molecules into a target cell. Invention methods comprise combining a biologically active molecule with mAb 3E10, or mutants or functional fragments thereof, and administering the resulting combination to said cell.

In accordance with another embodiment of the present invention, there are provided methods for the transport of biologically active molecules into a target cell, said method comprising: combining said biologically active molecule with a non-pathogenic monoclonal antibody, wherein said antibody promotes transport into said cell in an energy independent manner, and wherein said antibody is not anti-RNP, and administering the resulting combination to said cell.

Monoclonal antibodies (mAb) useful in the practice of the present invention, e.g., non-pathogenic monoclonal antibodies which promote transport into cells in an energy independent manner, and which are not anti-RNP antibodies, are capable of penetrating renal tubular epithelial cells in vivo and primary cultured neurons. Upon penetration, mAbs according to the invention (e.g., 3E10, as well as mutants and/or functional fragments thereof) localize in the cell nucleus.

In accordance with the present invention, it has been discovered that there is a class of monoclonal antibodies (e.g., mAb 3E10 and mutants and/or functional fragments thereof) which can be utilized to transport a wide variety of biologically important molecules into target cells, such as kidney cells, brain cells, ovarian cells, bone cells, and the like. Monoclonal antibody 3E10 is produced by a hybridoma 3E10 placed permanently on deposit with the American Type Culture Collection, 10801 University Blvd., Manassas,

Va. 20110-2209, USA, on Aug. 31, 2000, according to the terms of the Budapest Treaty under ATCC accession number PTA-2439. Antibodies according to the invention (e.g., mAb 3E10 or mutants and/or functional fragments thereof) can be conjugated to the biological molecule of interest to form an antibody conjugate that is capable of being transported into the cell. Upon entry into the cell, the antibody conjugate localizes in and around the cell nucleus. Antibody conjugates in accordance with the present invention may be used in the same manner as other conjugated delivery systems where an antibody or other targeting vehicle is conjugated to the biological molecule of interest to provide delivery to desired cells in the in vivo or in vitro environment.

Antibodies according to the invention (e.g., mAb 3E10 and mutants and/or functional fragments thereof) can be utilized to transport a wide variety of biologically active materials, e.g., nuclear transcription factors, enzymes, enzyme inhibitors, genes, and the like, to the cell nucleus for a variety of therapeutic effects. Pharmacologically active molecules including inorganic and organic molecules, pharmaceutical agents, drugs, peptides, proteins, genetic material, and the like, may be conjugated to antibodies according to the invention (e.g., mAb 3E10 and mutants and/or functional fragments thereof) for delivery thereof.

Naturally occurring antibodies are generally tetramers containing two light chains and two heavy chains. Experimentally, antibodies can be cleaved with the proteolytic enzyme papain, which causes each of the heavy chains to break, producing three separate subunits. The two units that consist of a light chain and a fragment of the heavy chain approximately equal in mass to the light chain are called the Fab fragments (i.e., the "antigen binding" fragments). The third unit, consisting of two equal segments of the heavy chain, is called the Fc fragment. The Fc fragment is typically not involved in antigen-antibody binding, but is important in later processes involved in ridding the body of the antigen.

As used herein, reference to mutants of mAb 3E10 includes variants of 3E10 which retain the same cell penetration characteristics as mAb 3E10, as well as variants modified by mutation to improve the utility thereof (e.g., improved ability to target specific cell types, improved ability to penetrate the cell membrane, improved ability to localize to the cellular DNA, and the like). Such mutants include variants wherein one or more conservative substitutions are introduced into the heavy chain, the light chain and/or the constant region(s) of the antibody.

As used herein, reference to functional fragments of mAb 3E10 includes portions of 3E10 that retain the same cell penetration characteristics as mAb 3E10. Such functional fragments include fragments containing at least the antigen-binding portion of mAb 3E10.

As readily recognized by those of skill in the art, altered antibodies (e.g., chimeric, humanized, CDR-grafted, bifunctional, antibody polypeptide dimers (i.e., an association of two polypeptide chain components of an antibody, e.g., one arm of an antibody comprising a heavy chain and a light chain, or an Fab fragment comprising V_L , V_H , C_L and C_H1 antibody domains, or an Fv fragment comprising a V_L domain and a V_H domain), single chain antibodies (e.g., an scFv (i.e., single chain Fv) fragment comprising a V_L domain linked to a V_H domain by a linker, and the like) can also be produced by methods well known in the art. Such antibodies can also be produced by hybridoma, chemical synthesis or recombinant methods described, for example, in (Sambrook et al., *Molecular Cloning: A Laboratory Manual* 2d Ed. (Cold Spring Harbor Laboratory, 1989); incorporated herein by reference and Harlow and Lane, *Antibodies: A*

Laboratory Manual (Cold Spring Harbor Laboratory 1988), which is incorporated herein by reference). Both anti-peptide and anti-fusion protein antibodies can be used (see, for example, Bahouth et al., Trends Pharmacol. Sci. 12:338 (1991); Ausubel et al., Current Protocols in Molecular Biology (John Wiley and Sons, NY 1989) which are incorporated herein by reference).

A presently preferred mutant contemplated for use in the practice of the present invention is a mAb 3E10 VH mutant involving a single change of the aspartic acid residue at position 31 to asparagine (3E10-31). The preparation of this mutant and a demonstration of its cell penetration ability is set forth in Example 5. This particular mAb 3E10 mutant is especially well suited for delivery of biological molecules to kidney and brain cells. Other 3E10 mutants and/or functional fragments thereof may be used to provide targeting of biological molecules. A wide variety of mutants and/or functional fragments thereof are possible provided that they exhibit substantially the same cell penetration characteristics as mAb 3E10 and 3E10-31 after conjugation to a selected biological agent for delivery.

MAb 3E10 heavy or light chains can be produced as fusion proteins with a variety of biologically active materials, e.g., nuclear transcription factors, enzymes, enzyme inhibitors, and the like, thereby enabling the transport of these proteins into the cell nucleus of target cells. In addition, mAb 3E10 can be produced in the form of a fusion protein with other proteins that bind DNA (such as, for example, poly-L-lysine). The poly-L-lysine fusion protein with mAb 3E10 would bind DNA (e.g., plasmids containing genes of interest) and transport the DNA into the nucleus of target cells.

Fusion proteins can be designed to place the protein of interest at the amino or carboxy terminus of either the antibody heavy or light chain. Since the antigen binding fragments (Fab's) of mAb 3E 10 have been shown to penetrate cells and localize in the nucleus, the entire heavy chain is not required. Therefore, potential configurations include the use of truncated portions of the heavy and light chain with or without spacer sequences as needed to maintain the functional integrity of the attached protein.

As an alternative to producing fusion proteins as described hereinabove, a universal carrier system can be devised. For example, various proteins or DNA can be conjugated to a common carrier such as protein A, poly-L-lysine, hex-histidine, and the like. The conjugated carrier will then form a complex with antibody according to the invention. A small portion of the carrier molecule that is responsible for binding immunoglobulin could be used as the carrier. Other similar configurations include design of carriers that interact with proteins engineered into the antibody heavy or light chain.

The mode of delivery chosen for administration of antibody conjugates according to the present invention to a live subject, such as a human patient or mammalian animal, will depend in large part on the particular biological molecule present in the conjugate and the target cells. In general, the same dosages and administration routes used to administer the biological molecule alone will also be used as the starting point for the antibody conjugate. However, it is preferred that smaller doses be used initially due to the expected increase in cellular penetration of the biological molecule. The actual final dosage for a given route of administration is easily determined by routine experimentation. In general the same procedures and protocols that have been previously used for other antibody-based target-

ing conjugates (e.g., parenterally, intravenous, intrathecal, and the like) are also suitable for the antibody conjugates of the present invention.

Many anti-DNA antibodies can penetrate several types of cells and localize to the cell nucleus. Recent work indicates that cellular penetration requires complexes of antibody and DNA. MAb 3E10 can penetrate many different types of cell lines in tissue culture. In contrast, mAb 3E10 may be restricted in the cells it can bind and penetrate in vivo, where there is an absence of free DNA to facilitate its penetration. It appears that the penetration of mAb 3E10 into kidney cells in vivo occurs by a different mechanism by which certain antibodies can penetrate cells and localize to the cell nucleus. The use of mAb 3E10 to penetrate kidney cells and brain cells involves a mechanism which is not common to other anti-DNA antibodies which require the presence of DNA or antibody Fc binding.

Many different autoantibodies have been shown to penetrate cells, including antibodies to RNP (see, for example, Alarcon-Segovia et al. (1978) supra, Ma et al., in Clin. exp. Immunol. 93:396 (1993) and Galoppin and J. Invest. Dermatol. 76:264 (1981)), RNA (see, for example, Varesio, et al., in Cancer Res. 35:3558 (1975)), Ro (see, for example, Lee, et al., in Arthritis Rheum. 29:782 (1986)), Proteinase 3 (see, for example, Csernok, et al., in Adv. Exp. Med. Biol. 336:45 (1993)), ribosomal protein P (see, for example, Reichlin, et al., in J. Clin. Invest. 93:443 (1994) and Koren, et al., in J. Immunol. 154:4857 (1995)), lymphocytes (see, for example, Okudaira, et al., in J. Clin. Invest. 69:1026 (1982)), synaptosomes (see, for example, Fabian in Neurology 38:1775 (1988)), and neurons (see, for example, Dalmau, et al., in Neurology 41:1757 (1991) and Hormigo and Lieberman in J. Neuroimmunol. 55:205 (1994)), and some have been shown to localize in the cell nucleus. Antibodies to ribosomal protein P have been shown to penetrate pig renal cells, localize to the nucleus, and induce cell injury (see, for example, Reichlin et al., supra and Koren et al., supra), but the presence of DNA was not required.

The requirement for free DNA and the role of Fc binding for cellular penetration appears to be different for different antibodies, but the antibodies studied appear to have distinct specificities for binding antigen, and they target different cell types. Therefore, multiple mechanisms may be operative in cellular penetration and nuclear localization. In preliminary studies to determine the mechanisms for cellular penetration and nuclear localization of mAb 3E10, DNase treatment and Fc blocking experiments were difficult to reproduce, indicating the complexity and potential artifacts involved in these procedures. Therefore, it was decided to produce mutants of mAb 3E10 variable heavy (VH) region and variable light (Vk for "variable kappa") region to definitively establish the relationship between antigen binding specificity and cellular penetration. The requirement for Fc binding and multivalent binding for cellular penetration was approached by producing molecular Fabs that are free of any contamination by undigested antibody or Fc fragments present in Fab prepared by enzyme digestion.

Monoclonal antibody 3E10 has recently been shown to cross react with a newly identified extracellular matrix protein, HP8/HEVIN (see Zack et al., in Journal of Immunology 154:1987-1994 (Feb. 15, 1995)). However, the present studies indicate that HP8/HEVIN is not involved in the penetration of mAb 3E10 into COS-7 or 3T3 cells. Indeed, it has been unequivocally shown that cellular penetration by mAb 3E10 correlates with DNA binding but is independent of Fc binding. Moreover, multivalent binding is not required. These results suggest that cellular penetration of

mAb 3E10 may occur through the formation of complexes containing antibody and DNA, but the possibility that mAb 3E10 bound to a membrane determinant that precisely resembles DNA cannot be excluded. Furthermore, the cell lines studied are penetrated by other (but not all) anti-DNA autoantibodies, suggesting a DNA dependent mechanism of penetration that may not reflect the specificity of binding and internalization of mAb 3E10 to renal tubular cells in vivo.

In addition to cellular penetration by anti-DNA antibodies, antineuronal antibodies have been shown to penetrate neurons, and the binding of intracellular targets has been proposed as a mechanism of disease pathogenesis (see, for example, Fabian in *Neurology* 40:419 (1990)). Moreover, a non-immunoglobulin protein has been shown to penetrate neurons and translocate to the nucleus. A 60 amino acid polypeptide corresponding to the homeodomain of the *Drosophila* protein Antennapedia was recently shown to penetrate neural cells, translocate to the cell nucleus, bind DNA, and regulate neural morphogenesis (see, for example, Joliet, et al., in *Proc. Natl. Acad. Sci. USA* 88:1864 (1991), La Roux, et al., in *Proc. Natl. Acad. Sci. USA* 90:9120 (1993), Bloch-Gallego, et al., in *The Journal of Cell Biology* 120:485 (1993) and Derossi, et al., in *J. Biol. Chem.* 269:10444 (1994)). Recovery of intact peptide suggests that targeting was not to the lysosomal compartment. Sequence homology between the Antennapedia homeodomain peptide and mAb 3E10 VH or Vk is not apparent.

The mechanism for the nuclear transport of anti-DNA antibodies remains unknown, but it has been suggested that the anti-DNA antibodies might be transported to the nucleus of cells as a result of arginine-rich sequences similar to the nuclear transport signals associated with nuclear transcription factors and other proteins (see, for example, Hanover in *The FASEB Journal* 6:2288 (1992)). Although the amino acid sequences of mAb 3E10 VH and Vk do not show linear sequences similar to known nuclear transport signals, these signals are quite diverse and may not be easily recognized. A novel binding domain of mAb 3E10 VH that is shared only by certain anti-DNA antibodies and is composed of conserved amino acid sequences in FR1 and FR3 has been described (see, for example, Zack, et al., in *Immunology and Cellular Biology* 72:513 (1994)). These regions have many arginine and lysine residues that could form a nuclear transport signal by their proximity in the three dimensional structure. Therefore, mAb 3E10 has several potential determinants that could serve as nuclear transport signals. These may bind other proteins (such as the recently described hSRP α) that act as functional receptors for some nuclear localization sequences and assist in transport across nuclear membranes (see, for example, Weis, et al., in *Science* 268:1049 (1995)). Alternatively, in some cases, carbohydrates are also used as nuclear transport signals (see, for example, Duverger, et al., in *Exp. Cell Res.* 207:197 (1993)). Glycosylation of the variable regions of the heavy or light chains could serve as a nuclear transport signal.

To assess the mechanism of nuclear transport of mAb 3E10, heavy and light chain cDNA devoid of signal peptide sequences were transfected into COS-7 cells. The engineered antibody was expressed in the cytoplasm and translocated to the cell nucleus (see, for example, Biocca, et al., in *EMBO J* 9 (1):101 (1990)). In contrast, mAb 3E10 was not translocated from the cytoplasm to the nucleus. Therefore, either the primary sequence alone is unable to initiate transfer of the antibody into the nucleus, or transport to the nucleus utilizes a pathway initiated by binding to the cell membrane. In either case, the mechanism for the nuclear localization of mAb 3E10 may be different than the transport

mechanism used for cytoplasmic proteins such as nuclear transcription factors. Since mAb 3E10 was not found in the nucleus of COS-7 cells that produced and secreted antibody, the antibody secretory and nuclear transport pathways must also be separate.

The present invention demonstrates the usefulness of specific antibodies for the introduction of biologically active molecules into cells, as well as the usefulness of producing molecular mutants and/or functional fragments of such autoantibodies in studying the cellular pathways of autoantibody penetration and nuclear localization.

It has been discovered that a single mutation in VH of monoclonal antibody 3E10 enhanced penetration to permit detection of nuclear localization at concentrations of only a few ng/ml. An scFv fragment was shown to be as effective as whole antibody in penetrating living cells, and cellular penetration and nuclear localization occurred in the absence of cellular toxicity. It has previously been shown that a complex of whole mAb 3E10 and catalase delivered neuroprotective amounts of catalase into primary neurons (R. H. Weisbart et al. *J. Immunol.* 164:6020, 2000). The antibody was now tested as a transport vehicle for restoration of p53 function in cancer cells using a polynucleotide encoding a fusion protein.

SW480 cells that contain two mutations in p53, His-273/Ser-309 were studied because they were shown by others to be sensitive to killing by p53 peptides. In the experiments described in Example 8, the results show that only a 30-mer C-terminal peptide of p53 produced significant cytotoxicity in SW480 cells, but not in COS-7 and CHO cells containing wild type p53. Cytotoxicity was modest even though the fusion protein was shown to penetrate all of the cells. However, the amount of fusion protein that penetrated cells was variable. Longer C-terminal peptides consisting of 36 and 50 C-terminal amino acids did not produce significant cytotoxicity.

Transcriptional activation of p53 was examined with the use of a CAT reporter attached to a consensus DNA binding domain for p53. Transfection of SW480 cells with pscFvp-53A and incubation of SW480 cells with scFv-p53A polypeptide did not increase the production of CAT, indicating that cytotoxicity of the p53A peptide was not due to restoration of p53 transcriptional activation. This finding is consistent with a recent report challenging the concept that the C-terminal of p53 is a negative regulatory domain.

The 30-mer C-terminal peptide of p53 expressed from a fusion protein containing the 3E10 monoclonal scFv fragment as nuclear delivery protein was cytotoxic for SW480 cells in vitro, but not as a result of restoring transcriptional activation of p53. Furthermore, peptide-induced cytotoxicity was modest. These results may differ from other studies of p53 peptide because a different nuclear delivery system was used and different C-terminal peptides of p53 were used. In spite of the apparent limitations of p53 peptide-induced cytotoxicity in SW480 cells in these studies, efficacy of the 3E10 monoclonal scFv fragment as a nuclear delivery vehicle in cancer cells in vitro is fully demonstrated by the data shown in Example 8 below.

In vitro, mAb 3E10 penetrates multiple types of living cells and localizes in the cell nucleus. scFv fragments of mAb 3E10 were produced and shown to penetrate living cells comparable to whole antibody. When injected intravenously into a mouse, however, mAb 3E10 scFv fragments were found localized primarily in the nucleus of skeletal muscle cells. To identify proteins responsible for cellular penetration and nuclear localization, the antibody was used to probe Western blots containing proteins derived from

nuclear and cytoplasmic lysates of different tissues. The antibody bound a 200 kDa protein found primarily in nuclear lysates of skeletal muscle cells. The specificity of antigen binding was consistent with the specificity of tissue penetration *in vivo*. The protein antigen was identified as myosin IIb by direct amino acid sequence analysis of the 200 kDa protein purified by antibody affinity chromatography and by the absence of the protein antigen in mice null for myosin IIb. The nuclear location of myosin IIb reactive with mAb 3E10 was confirmed microscopically in fixed muscle samples, indicating the presence of a nuclear isoform of myosin IIb not previously described.

Studies were conducted with lysates from multiple cell lines derived from different tissues in which mAb 3E10 penetrates. However, in these cell lines the studies failed to elucidate a protein responsible for transport of the antibody into cells grown in tissue culture. Therefore, it appears that the mechanisms for cellular penetration and nuclear localization of antibody into cell lines *in vitro* may be different than for antibody transport into skeletal muscle *in vivo*.

A variety of mechanisms responsible for autoantibody transport into cell lines have been implicated, including requirement for Fc receptors, and antibody binding to brush border myosin I (Alarcon-Segovia et al., 1978; Yanase et al., 1997). As shown in this study, mAb 3E10 did not bind myosin I, and scFv fragments devoid of the Fc region were able to penetrate living cells. It appears, therefore, that different antibodies may penetrate living cells by different mechanisms. It is possible that dsDNA may be a molecular mimic of diverse protein epitopes, some of which are molecular motors or messenger molecules.

Anti-DNA antibodies are only one small group of proteins identified that penetrate living cells. It is well known that a series of small protein transduction domains cross biological membranes independently of transporters or specific receptors, including a nine amino acid peptide of the HIV transactivation factor, TAT, VP22, a herpes simplex virus 1 protein, and the third alpha helix of Antennapedia homeodomain. It seems less likely that antibody molecules cross biological membranes through similar transduction domains because of their large size. Moreover, in contrast to small protein transduction domains that penetrate all cells, different antibodies demonstrate different tissue and cell specificities. Therefore, the motor function of myosin IIb is a more likely candidate for the transport of mAb 3E10 into skeletal muscle cells. However, further studies are required to define the precise pathway by which myosin IIb facilitates antibody transport into skeletal muscle cells.

Antibodies that penetrate living cells are frequently toxic or injurious and may explain some of the pathologic manifestations of the autoimmune diseases in which they are found. In contrast, mAb 3E10 shows no harm to cells that it penetrates in tissue culture (Weisbart et al., 1998). Moreover, studies *in vitro* have shown that mAb 3E10 and scFv fragments of mAb 3E10 can transport relatively large proteins, such as catalase, into the nucleus of cells in tissue culture (Weisbart et al., 2000). Despite this result of the *in vitro* studies, it has now surprisingly been discovered that *in vivo* (i.e., in live mammals, for example murine animals and humans), mAb 3E10 and functional fragments thereof selectively penetrate in living animals (i.e., *in vivo*) to the nucleus only of skeletal muscle cells. This surprising finding indicates that both the 3E10 antibody and the scFv fragment disclosed herein (and the elements of its transport pathway) are useful in the design of therapeutic, non-viral delivery vectors to selectively target skeletal muscle *in vivo*. Such studies also show that other molecules, such as other anti-

bodies, that will specifically and selectively bind to the newly discovered nuclear isoform of myosin IIb heavy chain (i.e., that do not bind to other isoforms of myosin IIb heavy chain) can be used as the transport molecule in an invention vector to selectively deliver biologically active molecules, such as therapeutic molecules, proteins and polynucleotides encoding biologically active molecules, to skeletal muscle *in vivo*.

Although myosin IIb is primarily a cytoplasmic protein, studies described in Example 9 below identify a heretofore unknown variant of myosin IIb (SEQ ID NO: 13) that is localized in the nucleus of skeletal muscle cells. The structure of the isoform of myosin IIb reactive with mAb 3E10 remains to be elucidated, but may be due to a post-translational modification of the molecule. It has now been discovered that this newly recognized isoform of myosin IIb selectively facilitates the transport of mAb 3E 10 into the nucleus of skeletal muscle cells.

Therefore, based on the studies described herein, in one embodiment, the invention provides methods for transporting a biologically active molecule into a target cell *in vitro* by combining the biologically active molecule with mAb 3E10 as produced by a hybridoma having ATCC accession number PTA 2439, or mutant or functional fragment thereof, and contacting the target cell *in vitro* under suitable conditions with the resulting combination, for example by culturing the target cells under growth conditions in the presence of the combination, for example as described in the Examples herein. By this method, the biologically active molecule is transported into the nucleus of the target cell. This embodiment of the invention is particularly useful as a research tool to discover molecules that will modulate the growth of cells in culture. For example, the invention method can be used as a screening tool to identify molecules that will inhibit growth of cancer cells in culture. Those molecules that pass this screening test can then be further utilized in additional screening tests *in vivo* to determine such attributes of selected molecules as *in vivo* function, side effects and safety.

For use in the invention methods, in addition to conjugating the antibody to the biologically active molecule, the latter can be attached to or associated with mAb 3E10 by any method known in the art. For example an scFv fragment of mAb 3E10, as described herein, can be expressed in a host cell as a fusion protein additionally containing a biologically active polypeptide for screening. Alternatively, the monoclonal antibody, or active fragment thereof, can be chemically linked to a polypeptide by a peptide bond or by a chemical or peptide linker molecule of the type well known in the art. Method for attaching a drug or other small molecule pharmaceutical to an antibody fragment are well known and include bifunctional chemical linkers such as N-succinimidyl (4-iodoacetyl)-aminobenzoate; sulfosuccinimidyl(4-iodoacetyl)-aminobenzoate; 4-succinimidyl-oxy-carbonyl- Ψ -(2-pyridyldithio) toluene; sulfosuccinimidyl-6-[α -methyl- Ψ -(pyridyldithio)-toluamido] hexanoate; N-succinimidyl-3-(2-pyridyldithio)-propionate; succinimidyl-6-[3 (-(-2-pyridyldithio)-propionamido] hexanoate; sulfosuccinimidyl-6-[3 (-(-2-pyridyldithio)-propionamido] hexanoate; 3-(2-pyridyldithio)-propionyl hydrazide, Ellman's reagent, dichlorotriazinic acid, S-(2-thiopyridyl)-L-cysteine, and the like. Further bifunctional linking molecules are disclosed in U.S. Pat. Nos. 5,349,066, 5,618,528, 4,569,789, 4,952,394, and 5,137,877, each of which is incorporated herein by reference in its entirety.

In yet another embodiment the invention provides methods for selectively transporting a biologically active mol-

ecule to the nucleus of skeletal muscle cells in a living subject by administering to the subject the biologically active molecule associated with a transport molecule that specifically and selectively binds to the nuclear isoform of the heavy chain of myosin IIb, thereby selectively transporting the biologically active molecule into the nucleus of skeletal muscle cells of the subject (i.e., the transport molecule does not bind to other isoforms of myosin IIb heavy chain). Therefore specific and selective binding of the transport molecule to the nuclear isoform of the heavy chain of myosin IIb in the skeletal muscle cells transports the associated biologically active molecule selectively into skeletal muscle cells. For example, the transport molecule can be either whole antibody or functional fragment or mutant thereof that specifically binds to the nuclear isoform of the heavy chain of myosin IIb in the skeletal muscle cells and not to other isoforms of myosin IIb. A presently preferred antibody for use as the transport molecule is mAb 3E10, or mutant or functional fragment thereof. The conjugate of the functional fragment and biologically active molecule can be a fusion protein expressed from a host cell, for example an scFv fragment comprising the variable region of the heavy chain (VH) and variable region of the kappa light chain (Vk) of mAb 3E10. For increased expression in the polynucleotide from which the scFv is expressed, the nucleic acids encoding the chains of mAb E310 are placed in reverse order with the Vk cDNA being placed 5' of VH.

A polynucleotide, such as one encoding a therapeutic protein, can also be selectively transfected into skeletal muscle cells by the invention vector by chemically bonding the polynucleotide to a transport molecule as disclosed herein, such as mAb 3E10 or a function fragment thereof, for example an scFv. Polynucleotides delivered into skeletal muscle cells in the subject using the invention vectors become stably integrated into the nucleus of the skeletal muscle cells. If the polynucleotide contains a gene rather than a regulatory molecule, the gene can be selectively expressed in the skeletal muscle cells of the subject.

In still another embodiment, the invention provides vectors for selectively transporting a biologically active molecule into skeletal muscle cells contained in a mammalian subject. The invention vectors comprise a transport molecule, for example a monoclonal antibody, or mutant or functional fragment thereof, that, in a live subject, specifically and selectively binds to a nuclear isoform of myosin heavy chain IIb and a biologically active molecule chemically associated with the transport molecule. A presently preferred candidate monoclonal antibody for use in the invention vectors is mAb 3E10 as produced by a hybridoma having ATCC accession number PTA 2439, or mutant or functional fragment thereof. In one embodiment, the functional antibody fragment in the invention vector is an scFv fragment. If the biologically active molecule is a polypeptide, such as a therapeutic polypeptide, the scFv fragment and the biologically active polypeptide can be contained in a fusion protein. Alternatively a functional fragment can be used to deliver a gene encoding a therapeutic protein or functional fragment thereof. A preferred functional fragment comprises the VH and Vk of mAb E310 and may optionally further comprise the signal peptide of the Vk. Such an scFv fragment is conveniently expressed from a polynucleotide having the nucleic acid encoding the Vk 5' of the nucleic acid encoding the VH.

For expression of polynucleotides encoding an invention vector, such as one comprising an scFv, the polynucleotide can be ligated into an expression construct, with each expression construct comprising a vector containing one or

more of the polynucleotides. The vector may further comprise expression regulatory sequences operably associated with the polynucleotide that can control and regulate the production in an appropriate host cell of a polypeptide(s) encoded by the polynucleotide.

In still another embodiment, the invention provides antibodies that specifically bind to the newly discovered nuclear isoform of myosin IIb heavy chain and not to other isoforms of myosin IIb heavy chain. Antibodies raised against myosin IIb heavy chain that bind selectively to the nuclear isoform of the protein can be prepared by those of skill in the art using conventional methods as disclosed herein and are known in the art, with the additional step that antibodies that bind specifically and selectively to the nuclear isoform of myosin IIb heavy chain are separated from antibodies that bind to other isoforms of myosin IIb heavy chain by attaching the nuclear isoform in its native 3-D configuration to a separation column as is known in the art and obtaining from the selection column only the antibodies that selectively bind to the nuclear isoform of the myosin IIb heavy chain. In addition to antibodies, antibody fragments and mutants of either, other transport molecules useful in the invention methods and vectors can be obtained in a similar fashion, i.e., by separating out, for example by means of a separation column, the putative transport molecules that bind to the nuclear isoform of myosin IIb heavy chain in its native 3-D configuration and do not bind to other isoforms of myosin IIb heavy chain.

Vectors suitable for use in preparation of polypeptides such as the invention antibody-containing vector include those selected from baculovirus, phage, plasmid, phagemid, cosmid, fosmid, bacterial artificial chromosome, viral DNA, P1-based artificial chromosome, yeast plasmid, and yeast artificial chromosome. For example, the viral DNA vector can be selected from vaccinia, adenovirus, fowl pox virus, pseudorabies and a derivative of SV40. Suitable bacterial vectors for use in practice of the invention methods include pQE70, pQE60, pQE-9, pBLUESCRIPT SK, pBLUESCRIPT KS, pTRC99a, pKK223-3, pDR540, PAC and pRIT2T. Suitable eukaryotic vectors for use in practice of the invention methods include pWLNEO, pXTI, pSG5, pSVK3, pBPV, pMSG, and pSVLSV40. Suitable eukaryotic vectors for use in practice of the invention methods include pWLNEO, pXTI, pSG5, pSVK3, pBPV, pMSG, and pSVLSV40.

Those of skill in the art can select a suitable regulatory region to be included in such a vector, for example from lacI, lacZ, T3, T7, apt, lambda PR, PL, trp, CMV immediate early, HSV thymidine kinase, early and late SV40, retroviral LTR, and mouse metallothionein-I regulatory regions.

Host cells in which the vectors containing the polynucleotides can be expressed include a bacterial cell, a eukaryotic cell, a yeast cell, an insect cell, or a plant cell. For example, *E. coli*, *Bacillus*, *Streptomyces*, *Salmonella typhimurium*, *Drosophila* S2, *Spodoptera* SJ9, CHO, COS (e.g. COS-7), or Bowes melanoma cells are all suitable host cells for use in practice of the invention methods.

The invention will now be described in greater detail by reference to the following non-limiting examples.

EXAMPLE 1

Monoclonal Antibodies

mAbs 3E10, 5C6, and 4H2 are IgG2a anti-dsDNA antibodies which were produced from spleen cells of MRL-mpj/lpr mice by fusion with cells from the FOX-NY cell line as

13

previously described (see Weisbart, et al., in J. Immunol. 144:2653 (1990)). mAb PP102 (Chemicon International, Temecula, Calif.), a murine IgG2a antibody that does not bind DNA, was used as a non-anti-DNA antibody isotype-matched control.

EXAMPLE 2

Monoclonal anti-DNA Antibody Binding in Vitro to Tissues of Human Organs

The monoclonal antibodies were purified from ascites by affinity chromatography using protein A-Sepharose and tested for binding kidney. mAb 3E10 was also tested for the ability to bind tissues from 19 other human organs, including blood vessels, nerve trunks, liver, connective tissues, lung, pancreas, gut, cardiac muscle, striated muscle, spleen, ovary, testis, thyroid, skin, eye, adrenal, brain, pituitary, and bone. Binding of monoclonal antibodies was detected with peroxidase conjugated affinity purified rabbit antibodies specific for mouse IgG Fc as previously described (see Taylor and Lote in Immunomicroscopy: A diagnostic tool for the surgical pathologist. Saunders, W. B., Philadelphia (1994)). mAb 3E10 Binds Human Renal Tubular Cells in Vitro

Three anti-dsDNA antibodies, i.e., mAbs 3E10, 5C6, and 4H2, were studied in vitro for binding tissue from fixed normal human kidney. All of the antibodies bound cell nuclei consistent with their anti-DNA reactivity. Thus, low magnification of renal tubular cells incubated with mAb 3E10 shows reactivity with renal tubular cell membranes and nuclei. Higher magnification emphasizes linear binding of mAb 3E10 to membranes of renal tubular cells. Only mAb 3E10 is seen to bind the cell surface of renal tubular cells.

The cell surface binding appeared consistent with binding the cell membrane. While mAb 3E10 bound tubular cells in five of five normal human kidneys studied, there was some variability in the intensity of staining. The results of incubating another anti-DNA antibody, mAb 5C6, with normal human kidney tubules shows binding to nuclei, but absence of binding to tubular cell membranes. The anti-DNA reactivity of mAb 5C6 is evident by the nuclear staining. However, in contrast to mAb 3E10, mAb 5C6 did not bind to renal tubular membranes.

Similarly, mAb 4H2 did not bind renal tubular membranes. None of the monoclonal anti-dsDNA antibodies were observed to bind antigens in renal glomeruli.

The specificity of binding of mAb 3E10 to kidney tubules was evaluated by studying its binding to tissues from other human organs. mAb 3E10 did not bind membranes or cytoplasmic antigens in tissues from multiple other organs, including blood vessels, nerve trunks, liver, connective tissues, lung, pancreas, gut, cardiac muscle, striated muscle, spleen, testis, thyroid, skin, eye, adrenal, pituitary, and bone. However, results of binding ovary and brain were inconclusive.

EXAMPLE 3

Monoclonal anti-DNA Antibody Binding in Vivo to Tissues of Normal BALB/c Mice

Normal BALB/c mice were primed with pristane and injected intraperitoneally with 2×10^7 hybridoma cells. After two weeks, the animals developed ascites containing antibodies with anti-dsDNA reactivity. Heart, liver, and kidney tissues were obtained and preserved in liquid nitrogen for studies of tissue histology. Binding of the anti-DNA anti-

14

bodies to tissues was detected with peroxidase-conjugated affinity-purified rabbit antibodies specific for mouse IgG Fc as previously described (see Taylor and Lote supra).

mAb 3E10 Binds Murine Renal Tubular Cells in Vivo

Thus, to determine if mAb 3E10 was reactive with mouse renal tubular cells in vivo, kidneys were examined from normal BALB/c mice two weeks after intraperitoneal injection with 3E10 cells to establish ascites containing high concentrations of mAb 3E10. mAb 3E10 did not localize in the membrane, cytoplasm, or nuclei of liver, cardiac muscle, or renal glomerular cells. Examination of renal tubular cells, however, showed nuclear staining, indicating that mAb 3E10 was selectively internalized and transported to the nucleus in renal tubular cells in vivo. The same results were observed in three of three BALB/c mice with ascites from 3E10 cells. In contrast to sections of dead, fixed tissue incubated with mAb 3E10 in vitro where the nuclei were exposed to the antibody, living, intact kidney cells in BALB/c mice would not be expected to contain intracellular antibody unless the antibody bound to cell membranes and entered the cells. In view of the fact that mAb 3E10 penetrated live renal tubular cells, it is likely that the cell surface staining observed in fixed dead cells could be attributed to binding the cell membrane.

The renal tubular cells from BALB/C mice with 3E10 ascites were examined in microscopic sections stained with either hematoxylin/eosin or Periodic Acid Schiff reagents. No significant abnormalities were observed.

The selective penetration of mAb 3E10 into intact renal tubular cells suggests that internalization was the result of specific antibody binding. Moreover, mAb 3E10 appears to have been transported across the nuclear membrane to localize in the nucleus in living renal tubular cells. In contrast, none of three BALB/c mice with ascites containing mAb 4H2 anti-dsDNA antibody showed binding of mAb 4H2 to liver, cardiac muscle or kidney. Comparable amounts of mAb 3E10 (0.8 mg/ml) and mAb 4H2 (1.0 mg/ml) were present in ascites used in these experiments. In contrast to mAb 4H2, mAb 3E10 enters the tubular cells and binds the cell nuclei.

These results suggest that mAb 3E10 is reactive with a membrane antigen on renal tubular cells in mouse as well as human kidney, and that the anti-kidney antibody binds renal tubular cells and is internalized and transported to the nucleus in vivo.

EXAMPLE 4

Cell Lines

MDCK dog kidney cells were kindly provided by Dr. Mostov, University of California, San Francisco, Calif. (see Apodaca, et al., in The Journal of Cell Biology 125:67 (1994)). Other cell lines, including 293 human embryonal kidney cells, COS-7 monkey kidney cells, NIH 3T3 cells, HT-29 colon cancer cells and LS 174T colon cancer cells were obtained from the American Type Culture Collection (ATCC, Rockville, Mass.). The cells were grown overnight in 96 or 48 well tissue culture plates in Dulbecco's modified Eagle media (DMEM, GIBCO BRL Life Technologies, Inc., Gaithersburg, Md.) in the presence of 10% horse serum (293 cells) or fetal calf serum (remaining cell lines) at 37° C., 5% CO₂, and humidified atmosphere.

After one day all media were aspirated and replaced with fresh media containing monoclonal antibodies. In preliminary studies, it was determined that a 10 µg/ml concentration of purified monoclonal antibody was required to demon-

15

strate binding to cells. In contrast, hybridoma supernatants (diluted 1:1 with fresh medium) containing 0.5 µg/ml were as effective as purified antibodies, so subsequent studies were done with purified antibodies (10 µg/ml) or hybridoma supernatants.

Cells were incubated with antibody for 1 to 2 hours and washed three times in Hanks Buffered Salt Solution and phosphate buffered saline (PBS). The cells were then fixed with 70% ethanol for 10 to 20 minutes and washed again multiple times in PBS. The cells were then incubated for 1 hour with alkaline phosphatase conjugated goat anti-mouse antibodies specific for binding IgG2a. The cells were washed three times in PBS and stained with nitro blue tetrazolium chloride/5-bromo-4-chloro-3-indolylphosphate p-toluidine salt (NBT/BCIP) in color development solution containing levamisole.

mAb 3E10 Penetrates Living Kidney Cells in Tissue Culture and is Translocated to the Nucleus

Several cell lines were examined for binding mAb 3E10. Kidney cell lines from human (293 cells), monkey (COS-7 cells), and dog (MDCK cells) were all observed to internalize mAb 3E10 and transport the antibody to the nucleus after only one hour of incubation. The kidney cells remained viable as demonstrated by the exclusion of trypan blue. In contrast, mAb 3E10 did not penetrate human colon cancer cells from LS 174T and HT-29 cell lines. Furthermore, an isotype matched control antibody, PP 102, without DNA binding reactivity did not penetrate any of the kidney cell lines. Antibody penetration and nuclear localization was not observed after only 15 minutes, but it was observed as early as 30 minutes and became maximal at 60 to 90 minutes.

EXAMPLE 5

Mutagenesis of mAb 3E10 Vk

The cloning of mAb 3E10 heavy and light chain cDNA was carried out as previously reported (see, for example, Zack, et al., 1994, supra, and Zack, et al., in J. Immunol. 154:1987 (1995)). Site directed mutagenesis of mAb 3E10 VH and Vk was performed by the method of Eckstein et al. (oligonucleotide—directed in vitro mutagenesis system, Amersham Corp., Arlington Heights, Ill.). The VH31 mutant used in these studies has been previously reported (see Zack, et al., 1995, supra). The mutated heavy and light chain cDNAs were ligated into the pSG5 expression vector (Stratagene, La Jolla, Calif.). Individual colonies were selected from transformed competent bacterial cells, and plasmids were prepared using the Wizard DNA purification system (Promega Corp., Madison, Wis.). Mutations were confirmed by dideoxynucleotide sequencing. The oligonucleotides used for mutagenesis of mAb 3E10 Vk are listed below.

Mutation	CDR	Residue	Oligonucleotide
SVST deletion	1	27A-D	5'-TGCAGGGCCAGCAAATCTA GCTATAGT-3' (SEQ ID NO: 1)
S to D	1	27C	5'-CAAAAAGTGTGATACATCT AGC-3' (SEQ ID NO: 2)
Y to F	1	32	5'-AGCTATAGTTTCATGCACTG G-3' (SEQ ID NO: 3)
Q to S	2	53	5'-TATGCATCCTCCTAGAA TCT-3' (SEQ ID NO: 4)
R to N	3	92	5'-TCAGCACAGTAATGAGTTTC CGTG-3' (SEQ ID NO: 5)

16

-continued

Mutation	CDR	Residue	Oligonucleotide
F to D	3	94	5'-CAGTAGGGAGGATCCGTGG ACG-3' (SEQ ID NO: 6)

Effect of a VH Mutant on Cell Penetration by mAb 3E10

Monoclonal antibody 3E10 heavy and light chain cDNAs were transfected into COS-7 cells and secretion of mAb 3E10 by the transfected COS-7 cells was demonstrated (see, for example, Zack et al., 1995, supra). Since mAb 3E10 can penetrate COS-7 cells, it was of interest to determine if secreted antibody was reinternalized in cultured cells. Only a small fraction of COS-7 cells produce antibody, thus the concentration of antibody in COS-7 supernatant is in the range of only 30 to 50 ng/ml. This is in marked contrast to concentrations of 10 to 50 µg/ml of purified antibody and 500 ng/ml of hybridoma supernatant required for the demonstration of antibody internalization in previous experiments. As expected, native mAb 3E10 was detected in the cytoplasm in about 1% of COS-7 cells after transfection. However, there was no evidence of translocation to the nucleus, indicating that the pathway by which antibody is secreted bypasses the pathway for nuclear localization. Moreover, the concentration of antibody in the COS-7 cell supernatant was insufficient to detect reentry of antibody into neighboring cells.

A mutant of mAb 3E10 VH has previously been produced involving a change in residue 31 in CDR1 from aspartic acid to asparagine. This mutation increases the ability of the antibody to bind to DNA (see Zack, et al., 1995, supra). Transfection of COS-7 cells with cDNA of the native 3E 10 light chain and cDNA of the VH31 mutant heavy chain resulted in an antibody that was readily observed to penetrate neighboring COS-7 cells and localize in the nucleus. The transfection efficiency was the same for the mutant and native heavy chain cDNAs, and the concentration of the mutant mAb 3E10 in COS-7 cell supernatants was the same as the native antibody.

Effect of mAb 3E10 Vk Mutations on Binding Specificity and Cellular Penetration

In previous studies, mAb 3E10 was used to identify a newly recognized extracellular matrix protein, HP8, in a cDNA expression library (see Zack, et al., 1995, supra). Moreover, DNA and HP8 were shown to share multiple binding determinants on mAb 3E10 VH. In the present studies, mutations in the CDR of mAb 3E10 Vk light chain were observed to eliminate binding to both dsDNA and HP8, dsDNA alone, and HP8 alone (Table I).

TABLE I

Effective of mAb 3E10 VH and Vk Mutations on Binding Specificity and Cellular Penetration

				Antibody Specificity		Cell entry	
				mAb 3E10	Mutation	dsDN	Cell
H chain	L chain	Location	Change	A	HP8	COS	3T3
Native	Native			0.26	2.02	No	No
31	Native	H CDR1	D to N	3.19	1.30	Yes	Yes
31	94	L CDR3	F to D	0.50	0.20	No	No
31	27 A-D	L CDR1	Delete	0.13	0.10	No	No
31	92	L CDR3	R to N	0.15	1.21	No	No
31	32	L CDR1	Y to F	0.16	2.76	No	No

17

TABLE I-continued

Effective of mAb 3E10 VH and Vk Mutations on Binding Specificity and Cellular Penetration							
mAb 3E10		Mutation		Antibody Specificity		Cell entry	
H chain	L chain	Location	Change	dsDN		Cell	
				A	HP8	COS	3T3
31	27C	L CDR1	S to D	3.22	0.26	Yes	Yes
31	53	L CDR2	Y to S	2.78	0.10	Yes	Yes
No H	No L			0.14	0.10	No	No

These results are consistent with the previous observations that dsDNA and HP8 share some but not all of binding determinants of mAb 3E10 VH. 3T3 cells have also been shown to express HP8 by Northern hybridization.

In order to further evaluate these differences, the relationship between antibody binding to dsDNA and HP8 and antibody penetration into COS-7 cells and 3T3 cells was studied. COS-7 cells and 3T3 cells were co-transfected with cDNA corresponding to the 3E10 heavy chain 31 mutant and cDNA corresponding to different kappa chain mutants. Cell penetration could not be demonstrated in antibodies containing each of four mutations in mAb 3E10 Vk that eliminated or reduced binding to dsDNA (see Table I). These mutations include residues 27A-D and 32 in CDR1, and residues 92 and 94 in CDR3. Two of the mutations, deletion of 27A-D in CDR1 and alteration of residue 94 in CDR3 eliminated antibody binding to both dsDNA and HP8. The other two mutations, residue 32 in CDR1 and residue 92 in CDR3 removed reactivity with dsDNA but did not affect binding to HP8. If reactivity with HP8 alone was removed, as in the mutation of residue 27C or of residue 53, the mutated antibody retained the ability to penetrate cells as long as the determinants essential for dsDNA binding remained intact. These results suggest that HP8 is not involved in antibody internalization in either of these cell lines. Cellular penetration by mAb 3E10 could be due to the formation of antibody-DNA complexes, or mAb 3E10 may bind a membrane determinant that precisely resembles DNA.

EXAMPLE 6

Molecular Constructs of mAb 3E10 Heavy and Light Chain cDNA

mAb 3E10 heavy and light chain cDNA without leader sequences were amplified by PCR using sense primers beginning at FR1 with the addition of the nucleotide sequence ATG. The primers used were:

Heavy chain sense primer:

5'-GCCATGGAGGTGCAGCTGGTGGAGTC-3' (SEQ ID NO:7)

Heavy chain antisense primer:

5'-AATTCTTATTTACCC(A)G(A)GAG
T(A)C(G)C(T)GGGGAA(T)(G)GC(G)TCT-3' (SEQ ID NO:8)

Light chain sense primer:

5'-GCCATGGACATTGTGCTGACACAGTC-3' (SEQ ID NO:9)

Light chain antisense primer:

5'-GAATTCTTAACACTCAATTCTTGTGAAGCTCTT-3' (SEQ ID NO:10)

18

To produce Fab of mAb 3E10, a heavy chain construct was amplified by PCR to contain the heavy chain leader sequence through CH1 and terminating in a stop codon. The primers used were:

Heavy chain sense primer:

5'-ATGGACTCCAGGCTCAATTTAGTTTTC-3' (SEQ ID NO:11)

Heavy chain antisense primer:

5'-TTATATAAATTTCTTGTCCACTTTGGTG-3' (SEQ ID NO:12)

The conditions used for PCR were: 1 minute denaturation at 95° C., 1 minute annealing at 55° C., and 1.5 minutes extension at 72° C. for 38 cycles with an additional 2 second extension time per cycle.

Localization of mAb 3E10 Devoid of Signal Peptides

To determine if mAb 3E10 is transported to the nucleus as a result of binding cytoplasmic proteins, mAb 3E10 was expressed in COS-7 cells without signal peptide sequences to prevent localization to the endoplasmic reticulum and subsequent secretion from the cell. Histological staining using antibodies to mouse kappa chains demonstrated production of mAb 3E10 and its localization to the cytoplasm. The failure to secrete antibody was shown by the absence of antibody in COS-7 supernatants as measured by ELISA. mAb 3E10 was localized in the cytoplasm, but it was not translocated to the nucleus. Sham transfected COS-7 cells were similarly stained using antibodies to mouse kappa chains as a control.

Penetration of Cells by mAb 3E10 Fab

To investigate the requirement for antibody Fc and multivalent antibody binding in cellular penetration, the cellular penetration of mAb 3E10 Fab was examined. mAb 3E10 heavy chain cDNA, including the leader sequence, VH, and CH1, was amplified by PCR from cDNA of mAb VH31 mutant. The amplified fragment was ligated in pSG5 and co-transfected into COS-7 cells along with mAb 3E10 light chain cDNA. Secretion of antibody Fab by COS-7 cells was confirmed by a capture ELISA with plates coated with goat antibodies to mouse gamma chains (CH1) and detected by goat antibodies to mouse kappa chains. Fab were reinternalized in neighboring cells and found localized in the nucleus as detected by antibodies to mouse gamma chains. Sham transfected COS-7 cells were similarly stained with antibodies to mouse gamma chains as a control. These results eliminate the requirement of Fc and multivalency of antigen binding for cellular penetration by mAb 3E10.

EXAMPLE 7

Expression of Antibodies

Purified pSG5 plasmids containing heavy and light chain gene inserts were expressed in COS-7 mammalian cells. Two micrograms each of plasmid DNA containing a heavy chain cDNA and a light chain cDNA were transfected using DEAE-dextran into 10⁵ COS-7 cells grown in DMEM and 10% fetal calf serum. After three days of culture, the supernatants were harvested and tested by ELISA for the presence of light and heavy chains. The cells were fixed with 70% ethanol for 1 to 2 minutes and washed again multiple times in PBS. The cells were then incubated with alkaline phosphatase conjugated goat anti-mouse antibodies specific for binding IgG2a. The cells were washed in PBS for 3 hours and stained with NBT/BCIP in color development solution containing levamisole.

Nuclear Delivery of p53 Peptides by Antibody scFv Fragments in Vitro

p53 is a nuclear transcription factor that protects cells from replicating damaged DNA by initiating apoptosis in response to alterations in its DNA. Mutations in the DNA binding domain of p53 impair its function permitting unregulated cell proliferation accounting for half of all human malignancies. Restoration of p53 function is an important goal in cancer research. Development of nuclear transport vehicles capable of delivering therapeutics into living cells is a promising approach to restoring p53 function in cancer cells. Experiments were conducted using a single chain antibody fragment of mAb 3E10 to deliver p53 to the nucleus of cancer cells.

Plasmid Constructs

mAb 3E10 was produced by fusion of spleen cells from an MRL/lpr/mpj mouse with the FOX-NY hybridoma cell line as described herein. A recombinant scFv fragment containing a myc tag was constructed (as shown in FIG. 1) using VH and Vk cloned from mRNA from hybridoma cells by RT-PCR as previously described (11). In contrast to the scFv reported, the scFv used in these studies was constructed with the Vk 5' of VH, and included the signal peptide of Vk. In addition, myc and his₆ tags were added to the C-terminal of VH for purposes of purification and histological localization. The scFv cDNA was constructed by PCR to produce a fusion protein with C-terminal peptides of p53 by ligating the scFv cDNA into the EcoRI and BamHI site of pSG5 (Stratagene, La Jolla, Calif.) and ligating cDNA corresponding to C-terminal peptides of p53 into the BamHI and BglII sites of pSG5. The design of the constructs is shown in FIG. 1A and the plasmid constructs are shown in FIG. 1C. PCR fragments were identified by electrophoresis for 1 hour at 60 milliamps in 0.8% agarose gels containing 0.5 µg/ml ethidium bromide.

The nucleotide primers used to amplify scFv and scFv-p53 cDNA constructs are shown in FIG. 1B. Three p53 C-terminal peptides were produced as fusion proteins with mAb 3E10 scFv, peptide A with 30 amino acids (amino acid residues 364-393); peptide B with 36 amino acids (residues 358-393); peptide C with 50 amino acids (residues 344-393); and a control N-terminal peptide with 48 amino acids (residues 2-49) was produced. In addition, peptide A was produced as a fusion protein with a R95Q mutant of mAb 3E10 VH (pFv-Mut-p53A) that eliminates penetration of mAb 3E10 scFv into living cells (10). The restriction sites of the constructs in pSG5 were changed by PCR to permit ligation of scFv, scFv-p53A, and scFv-p53N into the EcoRI and SacII restriction sites of pPICZαA for subsequent expression in *Pichia pastoris* (EasySelect *Pichia* Expression Kit, Invitrogen Corporation, Carlsbad, Calif.).

Protein Synthesis and Purification

pSG5 plasmids containing the cDNA constructs described above were transfected by electroporation into COS-7 cells, and the secreted proteins were purified from culture supernatant by affinity binding to KAPPA LOCK™ (Zymed Laboratories, Inc., San Francisco, Calif.). The purified proteins were identified in Western blots of SDS-PAGE gels developed with antibodies to the myc tag in 3E10 scFv. Similarly, pPICZαA containing the constructs of interest were transfected by electroporation into *Pichia pastoris*, and the yeast were grown in 500 ml buffered Glycerol/Methanol Complex Medium (BMGY/BMMY) in 2L baffled shaker flasks. Protein synthesis was induced with 0.5% methanol according to

the instructions of the manufacturer (EasySelect *Pichia* Expression Kit, Invitrogen Corporation, Carlsbad, Calif.). Recombinant protein was purified by the his₆ tag with NI-NTA-Agarose (Qiagen, Valencia, Calif.), eluted with imidazole, and dialyzed against PBS. Recombinant proteins were electrophoresed in 4-15% gradient SDS-PAGE gels (Biorad laboratories, Hercules, Calif.) and identified by Western blotting to nitrocellulose membranes developed with murine monoclonal antibodies to myc tag (mAb 9E10) and to C-terminal p53 (PAb 421). Western blots were developed with alkaline phosphatase conjugated goat antibodies to mouse IgG. Alkaline phosphatase activity was measured by the chromogenic substrate, nitro blue tetrazolium chloride/5-bromo-4-chloro-3-indolylphosphate p-toluidine salt in color development solution containing levamisole.

Cytotoxic Assays

scFv fragments of mAb 3E10 and scFv fusion proteins with peptides of p53 were assayed for cytotoxicity by transfection of plasmids (pSG5) containing cDNA corresponding to these proteins into Green Monkey Kidney cells (COS-7), Chinese hamster cells (CHO) with wild type p53, human colorectal cancer cells (SW480) containing two mutations in p53, His-273/Ser-309, and human osteosarcoma cells (SAOS2). Transfections were done with lipofectamine according to the instructions of the manufacturer (Invitrogen Corporation, Carlsbad, Calif.). Cytotoxicity was also assayed by incubating these cell lines with recombinant proteins produced transiently in COS-7 cells and *Pichia pastoris*. Cell death was measured by the nuclear uptake of propidium iodide (1 µg/ml). Dead cells were observed by fluorescence microscopy with the use of a standard rhodamine filter for red fluorescence. Multiple experiments were done and each experiment contained duplicate determinations. The results were expressed as average percent killed cells determined by counting 300 cells in each well. Differences in the number of killed cells between groups were determined by Student's t test.

Transcriptional Activation of p53

COS-7 cells were cotransfected by electroporation with the plasmids pCD53, pscFv, or pscFv-p53A and the chloramphenicol acetyl transferase (CAT) reporter, pIRG-CAT. 10⁵ cells were cultured in 60 mm diameter Petri dishes for 24 hours in DMEM (Cellgrow, Mediatech, Herndon, Va.) and harvested. The cells were lysed in cell lysis buffer and assayed for CAT by CAT ELISA (Boehringer Mannheim, GmbH, Germany). The results were expressed as the amount of CAT (ng/ml) produced based on a standard curve with known amounts of CAT.

Expression of Recombinant Proteins Containing mAb 3E10 scFv in COS-7 cells

cDNA fragments corresponding to p53 A, B and C were PCR amplified and verified on agarose gels. The corresponding peptides were expressed transiently in COS-7 cells, purified from COS-7 supernatant and detected by electrophoresis in 4-15% gradient SDS-PAGE gels identified by mAb 9E10 that binds the myc tag of scFv and mAb PAb421 that binds the C-terminal of p53.

Each cDNA and polypeptide obtained was of the expected size. The integrity of the fusion proteins was demonstrated by presence of scFv as measured by antibodies (mAb 9E10) to the myc tag of scFv, and the presence of the C-terminus of p53 was demonstrated by binding to PAb421, an anti-p53 antibody. The cDNA sequences were confirmed by automated DNA sequence analysis. The polypeptides were assayed for functional activity by cellular penetration as described below.

21

scFv-p53 Peptides Penetrate Living Cells

COS-7 cells and SW480 cells were incubated with supernatants of COS-7 cells transfected with the plasmids containing cDNA constructs corresponding to the scFv and scFv-p53 peptides. Staining of the cells showed that mAb 3E10 scFv and scFv-p53A penetrated COS-7 cells and SW480 cells and localized in the nucleus. In contrast, scFvmut-p53A with the mutation R95Q in 3E10 VH did not penetrate living cells. scFv-p53B and C penetrated COS-7 cells and SW480 cells comparable to scFv-p53A. mAb 3E10 scFv and each of the scFv-p53 peptides also penetrated CHO cells and localized in the nucleus.

scFv-p53 Cytotoxicity

Plasmids containing cDNA constructs of mAb 3E10 scFv and scFv-p53 fusion proteins were transfected into COS-7, CHO, and SW480 cells (FIG. 1C). Transfection efficiency was determined by transfection with pEGFP and observing the presence of Green Fluorescent Protein by fluorescence microscopy. The transfection efficiency was consistently between 30 and 35% in each cell type and in each of multiple experiments. Transfection of COS-7 and CHO cells with cDNA constructs corresponding to scFv and scFv-p53 peptides did not produce significant cytotoxicity, as shown by the results of these studies presented in FIGS. 2 and 3. In contrast, pscFv-p53A produced modest (16%) but significant ($p < 0.01$) killing of SW480 cells compared to pscFv alone, as shown in FIG. 4, where the results represent the mean of 4 separate experiments. Similar experiments in SW480 cells with pCD53 (wild type p53) produced 23% killing, as shown in FIG. 4. However, the maximum killing by p53 would not be expected to exceed the maximum transfection efficiency of 35%. Thus, pCD53 killed 23/30-35% of the cells or approximately 66-77% of transfected cells. On the other hand, scFv-p53A is secreted and re-internalized into all of the cells. Therefore, the total percentage of killing by scFv-p53A is 16% of total cells incorporating the fusion protein. Thus, killing by pscFv-p53A is quite modest in comparison to wild type p53. Fv-Mut-p53A contains the mutation R95Q in 3E10 VH that prevents 3E10 VH from penetrating living cells. pFv-Mut-p53A was not cytotoxic in SW480 cells suggesting that cytotoxicity in response to transfection with pFv-p53A occurred as a result of secreted Fv-p53A protein that was reinternalized.

scFv-p53A protein was then produced in *Pichia pastoris*, purified, and tested for cytotoxicity by adding it directly to cells in tissue culture. scFv alone and scFv-p53N protein (control N-terminal peptide of p53) produced in *Pichia* were used as controls. In a separate experiment, scFv-p53N was shown to penetrate living SW480 cells and localize in the nucleus. Increased killing of COS-7 cells did not occur with any of the scFv-p53 fusion proteins at a concentration of 100 $\mu\text{g/ml}$. In contrast, incubation of 100 $\mu\text{g/ml}$ of scFv-p53A with SW480 produced modest killing (8% to 10%) compared to 1% to 2% killing by scFv and scFv-p53N alone in three separate experiments ($p < 0.01$). A representative experiment is displayed in FIG. 5. A dose response study showed killing of SW480 cells with as little as 10-25 $\mu\text{g/ml}$ of scFv-p53A; however, no significant enhancement of killing occurred at higher doses. The results of a representative experiment are shown in FIG. 6. In a separate experiment, scFv-p53A did not kill SAOS2 cells devoid of the p53 gene.

Transcriptional Activation by scFv-p53 Peptides

scFv-p53A was tested for restoration of transcriptional activation in SW480 cells cotransfected with pscFv-p53A and the pIRG-CA T reporter construct. Whereas pCD53 (wild type p53) induced significant CAT production, no

22

increase in CAT occurred in response to pscFv-p53A. A representative experiment is shown in Table II below. SW480 cells transfected with the reporter pIRG-CA T were also incubated with 100 $\mu\text{g/ml}$ of scFv-p53A protein for 24 hours, and the cells were examined for the production of CAT. scFv-p53A did not increase production of CAT in two separate experiments. These results indicate that transfection with pscFv-p53A or incubation of cells with the scFv-p53A protein did not restore p53 transcriptional activation in SW480 cells.

TABLE II

Effect of Fusion Proteins on p53 Transcriptional Activation	
Plasmid Transfected	CAT (ng/ml)
Sham Transfection	0.00
IRG-CAT Reporter alone	0.02
pFv	0.07
IRG-CAT Reporter plus:	
pFv-P53A	0.11
pFv-P53B	0.10
pFv-P53C	0.10
PCD53	1.00

EXAMPLE 9

Cell Type Specific Targeted Intracellular Delivery
In Vivo

Because of the ability of mAb 3E10 to transport potential therapeutic proteins into intracellular compartments, the tissue distribution and intracellular localization in vivo after intravenous administration was determined. In this example it is shown that that, in contrast to studies in vitro wherein mAb 3E 10 is shown to penetrate into the nucleus of various cell types, mAb 3E10 is transported selectively into skeletal muscle cells in vivo through binding a nuclear-specific variant of the adult fast IIb myosin heavy chain.

Antibodies and scFv Fragments

Monoclonal antibody 3E10 (mAb 3E10) was established by fusion of spleen cells from an MRL/lpr/mpj mouse with the FOX-NY cell line as previously described herein. cDNA corresponding to functional 3E 10 heavy and light chains were cloned by RT-PCR from mRNA isolated from hybridoma cells, and an scFv fragment was constructed by PCR as previously described (Weisbart et al., 1998). However, the 3E10 scFv used for these studies was redesigned to reverse the order of VH and Vk and place Vk cDNA 5' of VH. The new scFv used the signal peptide of the kappa chain. This modified scFv showed enhanced secretion from COS-7 cells.

mAb 3E 10 heavy and light chain constructs and mutants used in these experiments were previously described (Zack et al., 1996). COS-7 cells were obtained from the American Type Culture Collection (Rockville, Md.) and grown in DMEM with 10% FCS. mAb 3E10 cDNA and various recombinant cDNA constructs were ligated into the cloning site of pSG5 (Stratagene, La Jolla, Calif.). Plasmids were transfected into COS-7 cells by electroporation for transient production of recombinant proteins. Supernatants of COS-7 cells were used as a source of mAb 3E 10 to identify antigens in tissue lysates reactive with the antibody by Western blotting.

scFv cDNA was ligated into the cloning site of pPICZ α A, transfected into *Pichia pastoris* by electroporation, and

selected with Zeocin according to the manufacturer's instructions (EasySelect Pichia Expression Kit, Invitrogen Corporation, Carlsbad, Calif.). Clones with the highest levels of secreted scFv were identified by antibody screening (Wung and Gascoigne, 2002). *Pichia pastoris* was grown in 500 ml buffered Glycerol/Methanol Complex Medium (BMGY/BMMY) in 2L baffled shake flasks, and protein expression was induced with 0.5% methanol. 3E10 scFv containing a 3'-terminal his₆ tag was purified with NI-NTA-Agarose (Qiagen, Valencia, Calif.) and dialyzed with PBS.

A murine monoclonal antibody was obtained that was specific for binding skeletal muscle myosin (Sigma-Aldrich Fine Chemicals, St. Louis, Mo.).

Cellular Penetration of 3E10 scFv In Vivo

3E10 scFv-his₆ (50 micrograms in 50 microliters saline vehicle) was injected into the tail veins of normal FVB mice. After 4 hours, the mice were anesthetized by inhalation and the circulation was perfused with saline through a cannula inserted into the heart. Samples of tissues were snap frozen in chilled 2-methylbutane at -70° C. in liquid nitrogen. Tissues were cut in 10 µm sections by cryostat and fixed in 75% alcohol for 20 min. Sectioned tissues were incubated with goat anti-his₆ antibody conjugated with horseradish peroxidase 1:500 (Invitrogen Corporation, Carlsbad, Calif.) at 37° C. for 1 hr, washed and developed with the Metal Enhanced DAB Substrate Kit (Pierce Chemical Company, Rockford, Ill.), and cellular localization of the dark brown deposit was identified by light microscopy.

Mice Null for Myosin Heavy Chains

Mice null for myosin IIb and IId were previously described (Allen and Leinwand, 2001). Lysates were prepared from skeletal muscle tissue from wild type (wt) mice and mice null for myosin IIb and IId. The lysates were analyzed by Western blotting for binding antibody.

Tissue Lysates

Tissues were homogenized in a Dounce Homogenizer, and cytoplasmic and nuclear lysates were prepared with NE-PER Nuclear and Cytoplasmic Extraction Reagents (Pierce Chemical Company) according to the manufacturer's protocol.

Electrophoresis and Western Blotting

Electrophoresis samples were diluted 1:2 in sample buffer containing 2% SDS, boiled, and applied to 4%-15% tris-HCl polyacrylamide gradient gels (4-15% Ready Gel Tris-HCl, Bio Rad Laboratories, Hercules, Calif.). The gels were electrophoresed at 140 volts for 90 min in tris/glycine/SDS buffer (Protein Electrophoresis Buffer, Bio Rad Laboratories, Hercules, Calif.). The gels were stained with either GELCODE BLUE® Stain Reagent (Pierce Chemical Company), or the proteins were transblotted to nitrocellulose in Blotting Transfer Buffer, 25 mM Tris, 192 mM Glycine, pH 8.3 (Bio Rad Laboratories, Hercules, Calif.). As an alternative, concentrated HCl was added to the blotting transfer buffer to adjust the final pH of the buffer to 8.1. Western blots were developed with 3E 10 hybridoma supernatant or purified mAb 3E10. mAb 3E10 was then measured by alkaline phosphatase-conjugated goat antibodies to mouse gamma chains. Alkaline phosphatase activity was measured by the chromogenic substrate, nitro blue tetrazolium chloride/5-bromo-4-chloro-3-indolylphosphate p-toluidine salt in color development solution containing levamisole.

Antibody Affinity Chromatography

Separate columns of Immunopure Immobilized rprotein-A (Pierce Chemical Company) were saturated with murine monoclonal antibodies mAb PP 102 and mAb 3E 10 and washed with 0.05 M sodium borate buffer, pH 8.0, then 0.02 M tris buffered saline containing 0.5 M NaCl, pH 7.4.

Nuclear lysates prepared from rat skeletal muscle were passed through the PP102 column, and the column was washed with 0.05 M sodium borate, pH 8.0 containing 0.1% SDS. Samples of the Sepharose beads were diluted 1:2 in sample buffer containing 2% SDS, and analyzed by SDS-PAGE and Western blotting. The effluent from the PP102 column was then passed through the 3E10 column. The column was then washed and samples of beads were processed and analyzed similarly.

Antibody Specificity for Skeletal Muscle

mAb 3E10 was assayed by Western blotting for binding proteins in cytoplasmic and nuclear lysates prepared from multiple tissues to identify target antigens responsible for its transport into living cells in vivo. In preliminary experiments different conditions were used to transblot proteins in tissue lysates from acrylamide gels to nitrocellulose. mAb 3E10 did not bind proteins transferred to nitrocellulose in standard transblot buffer, pH 8.3. However, if the transblot buffer was adjusted to pH 8.1, a 200 kDa protein from skeletal muscle nuclear lysates was transferred that showed prominent binding by mAb 3E10 at about 132 kDa. In contrast, in this test the antibody did not bind a similar protein in nuclear lysates from heart muscle or other tissues, including kidney, lung, liver, spleen, ovary, and brain, and peripheral blood cells. The 200 kDa muscle protein reactive with mAb 3E10 was not present in a control blot developed with second antibody alone. In addition, mAb 3E10 did not bind proteins in cytoplasmic lysates from any of these tissues. In similar tests, it was shown that three other murine monoclonal anti-dsDNA antibodies of the same isotype (5C5, 5C6 and 4H2) did not bind to the 200 kDa protein in skeletal muscle nuclear lysates, indicating binding specificity of mAb 3E10 to the skeletal muscle protein.

Binding Specificity of Recombinant Antibody and scFv Fragments

Single chain Fv fragments (scFv) of mAb 3E10 were produced, and in preliminary studies, the scFv fragments were shown to bind dsDNA and penetrate living cells in vitro comparable to mAb 3E10 (Weisbart et al., 1998, supra). Recombinant 3E10 and scFv were tested for binding the 200 kDa protein in nuclear lysates from skeletal muscle. Recombinant antibody was expressed transiently in COS-7 cells transfected with pSG5 plasmids containing antibody cDNA constructs, and the supernatants were used to test for binding of antibody to the 200 kDa protein by Western blotting. The heavy chain of mAb 3E10 was previously shown to be associated with two separate light chains, only one of which binds dsDNA and penetrates living cells in combination with the heavy chain (Zack et al., 1995). The Western blots in this study showed that mAb 3E10 heavy chain in combination with the anti-dsDNA kIII light chain bound the 200 kDa protein but not in combination with the kVser light chain. A single mutation in CDR3 of the heavy chain, R95Q, was previously reported to eliminate binding of mAb 3E10 to dsDNA and prevent its transport into living cells in vitro. The R95Q mutation also prevented mAb 3E 10 from binding the 200 kDa protein from skeletal muscle nuclear lysate. The scFv containing 3E10 VH and VkIII also bound the 200 kDa protein in a Western blot. These results indicate the use of shared antibody determinants for binding dsDNA and the 200 kDa protein, and demonstrate that scFv fragments bind the 200 kDa protein similar to whole antibody.

Localization of Antibody to Skeletal Muscle Nuclei In Vivo

Vehicle or 3E10 scFv fragments (50 micrograms in 50 microliters of saline) were infused into the tail veins of FVB mice to study distribution of the antibody in vivo. Four hours

after injection, the animals were anesthetized by inhalation and the circulation was perfused with saline through a cannula inserted into the heart. Samples of tissues were snap frozen and kept at -80°C . Multiple tissues were sectioned by cryostat and stained with peroxidase-conjugated antibodies directed to the his_6 tag of scFv fragments. Histological analysis of stained samples showed that scFv fragments were found predominantly in the nuclei of about 20% of skeletal muscle cells. scFv fragments were also found infrequently in nuclei of renal tubular cells. The relatively small amount of antigen in kidney may explain the failure to identify reactivity of mAb 3E10 in Western blots of lysates from kidney. scFv fragments were not found in any other tissues, including brain, lung, intestine, spleen, liver, pancreas, ovary, and skin. The heart muscle was inadequately perfused with saline, so localization of antibody in heart muscle could not be evaluated adequately. Sections from vehicle injected control mouse tissues showed no antibody label. These results demonstrate the relative specificity of mAb 3E10 scFv fragments for penetrating skeletal muscle cells in vivo.

Antibody Affinity Purification

Nuclear lysates of rat skeletal muscle were passed through a column containing immobilized recombinant protein-A saturated with an IgG 2a isotype control antibody, mAb PP102, with unknown binding specificity. The effluent was then passed through a protein-A-Sepharose column saturated with mAb 3E10. The affinity columns were washed with 0.05M borate buffer, pH 8.0 containing 0.1% SDS, and a sample of beads from each was boiled in 2% SDS, electrophoresed in a 4%-15% gradient acrylamide gel, and stained with GELCODE BLUE® stain (Pierce Chemical Company). A separate sample was transblotted to nitrocellulose and developed with mAb 3E10. The results of these studies showed the proteins contained in skeletal muscle nuclear lysate were stained with GELCODE BLUE®. The absence of non-specific binding of nuclear proteins to the column containing the control mAb PP102 antibody was also demonstrated. The antibody heavy chain (50 kDa) and light chain (25 kDa) were identified as expected, since the antibody was not cross-linked to the protein-A column. In contrast, several proteins in addition to the 3E10 heavy chain (50 kDa) and two light chains (25 kDa and 28 kDa) were identified by the staining. A 200 kDa protein visualized by staining was identified as corresponding to the 200 kDa protein visualized by Western blot (lane 4) that demonstrates binding by mAb 3E10. The 200 kDa protein was excised from the gel and identified by electrospray mass spectrometry (Nano-LC/MS/MS). The analysis was done in the Howard Hughes Institute, Columbia University, N.Y. Nano-LC/MS/MS

Analysis of the 200 kDa protein from rat skeletal muscle by Nano-LC/MS/MS resulted in 28 peptides that showed complete identity with human myosin IIb (SEQ ID NO:13, FIG. 6). There was no other identifiable protein. To confirm the identity of myosin as the protein reactive with mAb 3E10, affinity purification of the 200 kDa protein from skeletal muscle nuclear lysate was performed using mAb 3E10

bound to protein-A-Sepharose. The purified protein was examined for binding a monoclonal anti-myosin antibody in a Western blot. The anti-myosin antibody bound the 200 kDa protein affinity purified with mAb 3E10.

Antibody Binding to Myosin Heavy Chains from Mice Null for Myosin Heavy Chains Iib and IId

To further confirm that mAb 3E10 binds myosin IIb, the antibody was used to develop Western blots of skeletal muscle lysates from mice null for myosin IIb and IId. A control anti-myosin antibody confirmed the presence of myosin IIb in mice null for IId and myosin IId in mice null for myosin IIb. In contrast, mAb 3E10 bound a 200 kDa skeletal muscle protein from wild type mice and mice null for myosin IId, but not from mice null for myosin IIb. This result confirmed that mAb 3E10 binds mouse myosin IIb.

In preliminary studies, myosin was transblotted to nitrocellulose in standard transblot buffer, but the 200 kDa protein reactive with mAb 3E10 required a less alkaline buffer. This result suggested that mAb 3E10 was binding a specific isoform of myosin IIb. This result was evaluated further by performing Western blots and comparing results obtained with standard transblot buffer, pH 8.3, and standard buffer adjusted to pH 8.1. In these studies, myosin contained in muscle lysates from wild type mice and mice null for myosin heavy chain IIb and IId was transferred to nitrocellulose in standard buffer as shown by development with a known anti-myosin antibody. However mAb 3E10 did not bind myosin IIb under standard transblot conditions. In contrast, the myosin IIb reactive with mAb 3E10 was transferred successfully to nitrocellulose in transblot buffer adjusted to pH 8.1. These results confirm that mAb 3E10 binds myosin IIb. However, mAb 3E10 binds only myosin IIb transblotted at pH 8.1, indicating that mAb 3E10 likely binds a specific isoform variant of myosin IIb.

Myosin IIb Isoform in Muscle Nuclei

The isoform of myosin IIb reactive with mAb 3E10 was identified primarily in nuclear, but not cytoplasmic lysates, of skeletal muscle cells. To localize this myosin isoform in tissue, 3E10 scFv was examined for binding fixed skeletal muscle. In these studies, 3E10 scFv bound to the nucleus only, but was distributed in an unusual pattern. Examination of cross-sections of skeletal muscle showed irregular nuclear deposition of antibody compared to the characteristic "ring" pattern of nuclear binding to DNA observed in other cells. The pattern of nuclear staining usually observed was shown in cross sections of heart muscle for comparison. This finding is consistent with 3E10 binding an isoform of myosin IIb localized in the nucleus of skeletal muscle cells. There appears to be preferential binding of antibody to nuclear myosin IIb in skeletal muscle cells in contrast to binding to native DNA in the nuclei of cells from other tissues. This result indicates that mAb 3E10 has greater binding affinity to myosin IIb than to dsDNA and suggests that the binding of mAb 3E10 to dsDNA may be the result of cross-reactivity with an isoform of myosin IIb.

While the invention has been described in detail with reference to certain preferred embodiments thereof, it will be understood that modifications and variations are within the spirit and scope of that which is described and claimed.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 29

<210> SEQ ID NO 1

<211> LENGTH: 27

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide used for mutagenesis

<400> SEQUENCE: 1

tgcagggcca gcaaatctag ctatagt 27

<210> SEQ ID NO 2
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide used for mutagenesis

<400> SEQUENCE: 2

caaaagtgtc gatacatcta gc 22

<210> SEQ ID NO 3
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide used for mutagenesis

<400> SEQUENCE: 3

agctatagtt tcatgcactg g 21

<210> SEQ ID NO 4
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide used for mutagenesis

<400> SEQUENCE: 4

tatgcatcct ccctagaatc t 21

<210> SEQ ID NO 5
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide used for mutagenesis

<400> SEQUENCE: 5

tcagcacagt aatgagtttc cgtg 24

<210> SEQ ID NO 6
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide used for mutagenesis

<400> SEQUENCE: 6

cagtagggag gatccgtgga cg 22

<210> SEQ ID NO 7
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: heavy chain sense primer

<400> SEQUENCE: 7

-continued

gccatggagg tgcagctggt ggagtc

26

<210> SEQ ID NO 8
 <211> LENGTH: 41
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain antisense primer

<400> SEQUENCE: 8

aattcttatt taccagaga gtacgctggg gaatggcgtc t

41

<210> SEQ ID NO 9
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain sense primer

<400> SEQUENCE: 9

gccatggaca ttgtgctgac acagtc

26

<210> SEQ ID NO 10
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: light chain antisense primer

<400> SEQUENCE: 10

gaattcttaa cactcattct tgttgaagct ctt

33

<210> SEQ ID NO 11
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain sense primer

<400> SEQUENCE: 11

atggactcca ggctcaattt agttttc

27

<210> SEQ ID NO 12
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: heavy chain antisense primer

<400> SEQUENCE: 12

ttattaaatt ttcttggtcca ctttggtg

28

<210> SEQ ID NO 13
 <211> LENGTH: 1725
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

Met Gln Gly Thr Leu Glu Asp Gln Ile Ile Ser Ala Asn Pro Leu Leu
 1 5 10 15

Glu Ala Phe Gly Asn Ala Lys Thr Val Arg Asn Asp Asn Ser Ser Arg
 20 25 30

Phe Gly Lys Phe Ile Arg Ile His Phe Gly Ala Thr Gly Lys Leu Ala
 35 40 45

-continued

Ser	Ala	Asp	Ile	Glu	Thr	Tyr	Leu	Leu	Glu	Lys	Ser	Arg	Val	Thr	Phe	
50						55					60					
Gln	Leu	Lys	Ala	Glu	Arg	Ser	Tyr	His	Ile	Phe	Tyr	Gln	Ile	Leu	Ser	
65					70					75					80	
Asn	Lys	Lys	Pro	Glu	Leu	Ile	Glu	Met	Leu	Leu	Ile	Thr	Thr	Asn	Pro	
				85					90					95		
Tyr	Asp	Phe	Ala	Phe	Val	Ser	Gln	Gly	Glu	Ile	Thr	Val	Pro	Ser	Ile	
		100						105					110			
Asp	Asp	Gln	Glu	Glu	Leu	Met	Ala	Thr	Asp	Ser	Ala	Val	Asp	Ile	Leu	
		115					120					125				
Gly	Phe	Thr	Ala	Asp	Glu	Lys	Val	Ala	Ile	Tyr	Lys	Leu	Thr	Gly	Ala	
	130					135					140					
Val	Met	His	Tyr	Gly	Asn	Met	Lys	Phe	Lys	Gln	Lys	Gln	Arg	Glu	Glu	
145					150					155					160	
Gln	Ala	Glu	Pro	Asp	Gly	Thr	Glu	Val	Ala	Asp	Lys	Ala	Ala	Tyr	Leu	
			165						170					175		
Thr	Ser	Leu	Asn	Ser	Ala	Asp	Leu	Leu	Lys	Ser	Leu	Cys	Tyr	Pro	Arg	
		180					185						190			
Val	Lys	Val	Gly	Asn	Glu	Phe	Val	Thr	Lys	Gly	Gln	Thr	Val	Gln	Gln	
	195						200					205				
Val	Tyr	Asn	Ala	Val	Gly	Ala	Leu	Ala	Lys	Ala	Ile	Tyr	Glu	Lys	Met	
	210					215					220					
Phe	Leu	Trp	Met	Val	Thr	Arg	Ile	Asn	Gln	Gln	Leu	Asp	Thr	Lys	Gln	
225					230					235					240	
Pro	Arg	Gln	Tyr	Phe	Ile	Gly	Val	Leu	Asp	Ile	Ala	Gly	Phe	Glu	Ile	
			245						250					255		
Phe	Asp	Phe	Asn	Ser	Leu	Glu	Gln	Leu	Cys	Ile	Asn	Phe	Thr	Asn	Glu	
		260						265					270			
Lys	Leu	Gln	Gln	Phe	Phe	Asn	His	His	Met	Phe	Val	Leu	Glu	Gln	Glu	
	275					280						285				
Glu	Tyr	Lys	Lys	Glu	Gly	Ile	Glu	Trp	Glu	Phe	Ile	Asp	Phe	Gly	Met	
	290					295					300					
Asp	Leu	Ala	Ala	Cys	Ile	Glu	Leu	Ile	Glu	Lys	Pro	Met	Gly	Ile	Phe	
305					310					315					320	
Ser	Ile	Leu	Glu	Glu	Glu	Cys	Met	Phe	Pro	Lys	Ala	Thr	Asp	Thr	Ser	
			325						330					335		
Phe	Lys	Asn	Lys	Leu	Tyr	Glu	Gln	His	Leu	Gly	Lys	Ser	Asn	Asn	Phe	
		340						345					350			
Gln	Lys	Pro	Lys	Pro	Ala	Lys	Gly	Lys	Pro	Glu	Ala	His	Phe	Ser	Leu	
		355					360					365				
Val	His	Tyr	Ala	Gly	Thr	Val	Asp	Tyr	Asn	Ile	Ala	Gly	Trp	Leu	Asp	
	370					375					380					
Lys	Asn	Lys	Asp	Pro	Leu	Asn	Glu	Thr	Val	Val	Gly	Leu	Tyr	Gln	Lys	
385					390					395					400	
Ser	Ala	Met	Lys	Thr	Leu	Ala	Phe	Leu	Phe	Ser	Gly	Ala	Gln	Thr	Ala	
			405						410					415		
Glu	Ala	Glu	Gly	Gly	Gly	Gly	Lys	Lys	Gly	Gly	Lys	Lys	Lys	Gly	Ser	
		420					425						430			
Ser	Phe	Gln	Thr	Val	Ser	Ala	Leu	Phe	Arg	Glu	Asn	Leu	Asn	Lys	Leu	
		435					440					445				
Met	Thr	Asn	Leu	Arg	Ser	Thr	His	Pro	His	Phe	Val	Arg	Cys	Ile	Ile	
	450					455					460					
Pro	Asn	Glu	Thr	Lys	Thr	Pro	Gly	Ala	Met	Glu	His	Glu	Leu	Val	Leu	

-continued

465	470	475	480
His Gln Leu Arg Cys Asn Gly Val Leu Glu Gly Ile Arg Ile Cys Arg	485	490	495
Lys Gly Phe Pro Ser Arg Ile Leu Tyr Ala Asp Phe Lys Gln Arg Tyr	500	505	510
Lys Val Leu Asn Ala Ser Ala Ile Pro Glu Gly Gln Phe Ile Asp Ser	515	520	525
Lys Lys Ala Ser Glu Lys Leu Leu Gly Ser Ile Glu Ile Asp His Thr	530	535	540
Gln Tyr Lys Phe Gly His Thr Lys Val Phe Phe Lys Ala Gly Leu Leu	545	550	555
Gly Thr Leu Glu Glu Met Arg Asp Glu Lys Leu Ala Gln Leu Ile Thr	565	570	575
Arg Thr Gln Ala Ile Cys Arg Gly Phe Leu Met Arg Val Glu Phe Arg	580	585	590
Lys Met Met Glu Arg Arg Glu Ser Ile Phe Cys Ile Gln Tyr Asn Ile	595	600	605
Arg Ala Phe Met Asn Val Lys His Trp Pro Trp Met Lys Leu Tyr Phe	610	615	620
Lys Ile Lys Pro Leu Leu Lys Ser Ala Glu Thr Glu Lys Glu Met Ala	625	630	635
Asn Met Lys Glu Glu Phe Glu Lys Thr Lys Glu Glu Leu Ala Lys Thr	645	650	655
Glu Ala Lys Arg Lys Glu Leu Glu Glu Lys Met Val Thr Leu Met Gln	660	665	670
Glu Lys Asn Asp Leu Gln Leu Gln Val Gln Ala Glu Ala Asp Ala Leu	675	680	685
Ala Asp Ala Glu Glu Arg Cys Asp Gln Leu Ile Lys Thr Lys Ile Gln	690	695	700
Leu Glu Ala Lys Ile Lys Glu Val Thr Glu Arg Ala Glu Asp Glu Glu	705	710	715
Glu Ile Asn Ala Glu Leu Thr Ala Lys Lys Arg Lys Leu Glu Asp Glu	725	730	735
Cys Ser Glu Leu Lys Lys Asp Ile Asp Asp Leu Glu Leu Thr Leu Ala	740	745	750
Lys Val Glu Lys Glu Lys His Ala Thr Glu Asn Lys Val Lys Asn Leu	755	760	765
Thr Glu Glu Met Ala Gly Leu Asp Glu Thr Ile Ala Lys Leu Thr Lys	770	775	780
Glu Lys Lys Ala Leu Gln Glu Ala His Gln Gln Thr Leu Asp Asp Leu	785	790	795
Gln Met Glu Glu Asp Lys Val Asn Thr Leu Thr Lys Ala Lys Thr Lys	805	810	815
Leu Glu Gln Gln Val Asp Asp Leu Glu Gly Ser Leu Glu Gln Glu Lys	820	825	830
Lys Leu Cys Met Asp Leu Glu Arg Ala Lys Arg Lys Leu Glu Gly Asp	835	840	845
Leu Lys Leu Ala Gln Glu Ser Thr Met Asp Thr Glu Asn Asp Lys Gln	850	855	860
Gln Leu Asn Glu Lys Leu Lys Lys Lys Glu Phe Glu Met Ser Asn Leu	865	870	875
Gln Gly Lys Ile Glu Asp Glu Gln Ala Leu Ala Ile Gln Leu Gln Lys	885	890	895

-continued

Lys Ile Lys Glu Leu Gln Ala Arg Ile Glu Glu Leu Glu Glu Glu Ile
 900 905 910

Glu Ala Glu Arg Ala Ser Arg Ala Lys Ala Glu Lys Gln Arg Ser Asp
 915 920 925

Leu Ser Arg Glu Leu Glu Glu Ile Ser Glu Arg Leu Glu Glu Ala Gly
 930 935 940

Gly Ala Thr Ser Ala Gln Ile Glu Met Asn Lys Lys Arg Glu Ala Glu
 945 950 955 960

Phe Gln Lys Met Arg Arg Asp Leu Glu Glu Ser Thr Leu Gln His Glu
 965 970 975

Ala Thr Ala Ala Ala Leu Arg Lys Lys His Ala Asp Ser Val Ala Glu
 980 985 990

Leu Gly Glu Gln Ile Asp Ser Leu Gln Arg Val Lys Gln Lys Leu Glu
 995 1000 1005

Lys Glu Lys Ser Glu Leu Lys Met Glu Ile Asn Asp Leu Ala Ser
 1010 1015 1020

Asn Met Glu Thr Val Ser Lys Ala Lys Ala Asn Phe Glu Lys Met
 1025 1030 1035

Cys Arg Thr Leu Glu Asp Gln Leu Ser Glu Ile Lys Thr Lys Glu
 1040 1045 1050

Glu Glu Gln Gln Arg Leu Ile Asn Glu Leu Ser Ala Gln Lys Ala
 1055 1060 1065

Arg Leu His Thr Glu Ser Gly Glu Phe Ser Arg Gln Leu Asp Glu
 1070 1075 1080

Lys Asp Ala Met Val Ser Gln Leu Ser Arg Gly Lys Gln Ala Phe
 1085 1090 1095

Thr Gln Gln Ile Glu Glu Leu Lys Arg Gln Leu Glu Glu Glu Thr
 1100 1105 1110

Lys Ala Lys Ser Thr Leu Ala His Ala Leu Gln Ser Ala Arg His
 1115 1120 1125

Asp Cys Asp Leu Leu Arg Glu Gln Tyr Glu Glu Glu Gln Glu Ala
 1130 1135 1140

Lys Ala Glu Leu Gln Arg Gly Met Ser Lys Ala Asn Ser Glu Val
 1145 1150 1155

Ala Gln Trp Arg Thr Lys Tyr Glu Thr Asp Ala Ile Gln Arg Thr
 1160 1165 1170

Glu Glu Leu Glu Glu Ala Lys Lys Lys Leu Ala Gln Arg Leu Gln
 1175 1180 1185

Asp Ala Glu Glu His Val Glu Ala Val Asn Ser Lys Cys Ala Ser
 1190 1195 1200

Leu Glu Lys Thr Lys Gln Arg Leu Gln Asn Glu Val Glu Asp Leu
 1205 1210 1215

Met Ile Asp Val Glu Arg Ser Asn Ala Ala Cys Ile Ala Leu Asp
 1220 1225 1230

Lys Lys Gln Arg Asn Phe Asp Lys Val Leu Ala Glu Trp Lys Gln
 1235 1240 1245

Lys Tyr Glu Glu Thr Gln Ala Glu Leu Glu Ala Ser Gln Lys Glu
 1250 1255 1260

Ser Arg Ser Leu Ser Thr Glu Leu Phe Lys Val Lys Asn Ala Tyr
 1265 1270 1275

Glu Glu Ser Leu Asp His Leu Glu Thr Leu Lys Arg Glu Asn Lys
 1280 1285 1290

-continued

Asn Leu 1295	Gln Gln Glu Ile Ser 1300	Asp Leu Thr Glu Gln 1305	Ile Ala Glu
Gly Gly 1310	Lys His Ile His Glu 1315	Leu Glu Lys Val Lys 1320	Lys Gln Leu
Asp His 1325	Glu Lys Ser Glu Leu 1330	Gln Thr Ser Leu Glu 1335	Glu Ala Glu
Ala Ser 1340	Leu Glu His Glu Glu 1345	Gly Lys Ile Leu Arg 1350	Ile Gln Leu
Glu Leu 1355	Asn Gln Val Lys Ser 1360	Glu Ile Asp Arg Lys 1365	Ile Ala Glu
Lys Asp 1370	Glu Glu Leu Asp Gln 1375	Leu Lys Arg Asn His 1380	Leu Arg Val
Val Glu 1385	Ser Met Gln Ser Thr 1390	Leu Asp Ala Glu Ile 1395	Arg Ser Arg
Asn Asp 1400	Ala Leu Arg Ile Lys 1405	Lys Lys Met Glu Gly 1410	Asp Leu Asn
Glu Met 1415	Glu Ile Gln Leu Asn 1420	His Ala Asn Arg Gln 1425	Ala Ala Glu
Ala Leu 1430	Arg Asn Leu Arg Asn 1435	Thr Gln Gly Ile Leu 1440	Lys Asp Thr
Gln Leu 1445	His Leu Asp Asp Ala 1450	Ile Arg Gly Gln Asp 1455	Asp Leu Lys
Glu Gln 1460	Leu Ala Met Val Glu 1465	Arg Arg Ala Asn Leu 1470	Met Gln Ala
Glu Val 1475	Glu Glu Leu Arg Ala 1480	Ser Leu Glu Arg Thr 1485	Glu Arg Gly
Arg Lys 1490	Met Ala Glu Gln Glu 1495	Leu Leu Asp Ala Ser 1500	Glu Arg Val
Gln Leu 1505	Leu His Thr Gln Asn 1510	Thr Ser Leu Ile Asn 1515	Thr Lys Lys
Lys Leu 1520	Glu Thr Asp Ile Ser 1525	Gln Ile Gln Gly Glu 1530	Met Glu Asp
Ile Val 1535	Gln Glu Ala Arg Asn 1540	Ala Glu Glu Lys Ala 1545	Lys Lys Ala
Ile Thr 1550	Asp Ala Ala Met Met 1555	Ala Glu Glu Leu Lys 1560	Lys Glu Gln
Asp Thr 1565	Ser Ala His Leu Glu 1570	Arg Met Lys Lys Asn 1575	Met Glu Gln
Thr Val 1580	Lys Asp Leu Gln Leu 1585	Arg Leu Asp Glu Ala 1590	Glu Gln Leu
Ala Leu 1595	Lys Gly Gly Lys Lys 1600	Gln Ile Gln Lys Leu 1605	Glu Ala Arg
Val Arg 1610	Glu Leu Glu Ser Glu 1615	Val Glu Ser Glu Gln 1620	Lys His Asn
Val Glu 1625	Ala Val Lys Gly Leu 1630	Arg Lys His Glu Arg 1635	Arg Val Lys
Glu Leu 1640	Thr Tyr Gln Thr Glu 1645	Glu Asp Arg Lys Asn 1650	Ile Leu Arg
Leu Gln 1655	Asp Leu Val Asp Lys 1660	Leu Gln Thr Lys Val 1665	Lys Ala Tyr
Lys Arg 1670	Gln Ala Glu Glu Ala 1675	Glu Glu Gln Ser Asn 1680	Val Asn Leu
Ala Lys	Phe Arg Lys Leu Gln	His Glu Leu Glu Glu	Ala Lys Glu

-continued

1685	1690	1695
Arg Ala Asp Ile Ala Glu Ser Gln Val Asn Lys Leu Arg Val Lys		
1700	1705	1710
Ser Arg Glu Val His Thr Lys Val Ile Ser Glu Glu		
1715	1720	1725

<210> SEQ ID NO 14
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 14
 gaattcatgg agtcagacac actgctgcta tgggtg 36

<210> SEQ ID NO 15
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 15
 ggatccatga tgatgatgat gatggtc 27

<210> SEQ ID NO 16
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 16
 ggatccgctc actccagcca cct 23

<210> SEQ ID NO 17
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 17
 agatcttcag tctgagtcag gccc 24

<210> SEQ ID NO 18
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 18
 ggatccgagc caggggggag ca 22

<210> SEQ ID NO 19
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 19
 agatcttcag tctgagtcag gccc 24

-continued

<210> SEQ ID NO 20
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 20

ggatccctga atgaggcctt ggaact

26

<210> SEQ ID NO 21
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 21

agatcttcag tctgagtcag gcc

24

<210> SEQ ID NO 22
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 22

ggatccgagg agccgcagtc agat

24

<210> SEQ ID NO 23
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 23

agatcttcaa atatcgtcgc gggacag

27

<210> SEQ ID NO 24
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 24

gaatccgaca ttgtgctgac acagt

25

<210> SEQ ID NO 25
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 25

ccgcgggtcaa tgatgatgat gatgatgatg ggagac

36

<210> SEQ ID NO 26
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 26

gaatccgaca ttgtgctgac acagt

25

<210> SEQ ID NO 27

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 27

ccgcgggtcag tctgagtcag gcc

24

<210> SEQ ID NO 28

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 28

gaatccgaca ttgtgctgac acagt

25

<210> SEQ ID NO 29

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 29

ccgcgggtcag tctgagtcag gcc

24

That which is claimed is:

1. A method for transporting a biologically active molecule into the nucleus of a target cell in vitro, comprising:

- a) combining said biologically active molecule in vitro with an antibody selected from the group consisting of monoclonal antibody (mAb) 3E10 as produced by a hybridoma having ATCC accession number PTA 2439 [or], and an [scFv fragment thereof] antigen binding portion of mAb 3E10, to form an antibody-biologically active molecule conjugate or fusion protein[.]; and
- b) contacting the target cell in vitro, under suitable conditions with the antibody-biologically active molecule conjugate of a), thereby transporting the biologically active molecule into the nucleus of the target cell, wherein the target cell is selected from the group consisting of a kidney cell, an ovarian cell, a bone cell, and a primary cultured neuron.

2. The method of claim 1 wherein said biologically active molecule is a nuclear transcription factor, an enzyme, an enzyme inhibitor, genetic material, an inorganic or organic molecule, a pharmaceutical agent, a drug, or a polypeptide.

3. The method of claim 2, wherein the biologically active molecule is a polypeptide.

4. The method of claim 1, wherein the antibody is [the] selected from the group consisting of a single chain antibody (scFv), [fragment] an Fab, and an Fv.

5. The method of claim 4, wherein the biologically active molecule is a polypeptide.

6. The method of claim 1, wherein the contacting comprises incubating the cells in the presence of the conjugate or a fusion protein.

7. The method of claim 1, wherein the [scFv fragment] antigen binding portion of mAb 3E10 comprises the variable region of the heavy chain (VH) and variable region of the kappa light chain (Vk) of mAb 3E10.

8. The method of claim 7, wherein the [scFv fragment] antigen binding portion of mAb 3E10 further comprises the signal peptide of the Vk.

9. The method of claim 1, wherein the biologically active molecule is a p53 polypeptide.

10. [The method of claim 9.] A method for transporting a biologically active molecule into the nucleus of a target cell in vitro, comprising:

- a) combining the biologically active molecule in vitro with an antibody selected from the group consisting of monoclonal antibody (mAb) 3E10 as produced by a hybridoma having ATCC accession number PTA 2439, and an antigen binding portion of mAb 3E10, to form an antibody-biologically active molecule conjugate or fusion protein; and
- b) contacting the target cell in vitro, wherein the target cell is a cancer cell, under suitable conditions with the antibody-biologically active molecule conjugate of a), thereby transporting the biologically active molecule into the nucleus of the target cell.

11. The method of claim 10, wherein the antibody is selected from the group consisting of a single chain antibody (scFv), an Fab, and an Fv.

12. The method of claim 10, wherein the biologically active molecule is a p53 polypeptide.

* * * * *