

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 February 2011 (24.02.2011)

PCT

(10) International Publication Number  
**WO 2011/022657 A1**

- (51) **International Patent Classification:**  
C07K 14/755 (2006.01) A61K 38/36 (2006.01)
- (21) **International Application Number:**  
PCT/US20 10/046 180
- (22) **International Filing Date:**  
20 August 2010 (20.08.2010)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
61/235,570 20 August 2009 (20.08.2009) US
- (71) **Applicants (for all designated States except US):** BAXTER INTERNATIONAL INC. [-/US]; One Baxter Parkway, Deerfield, IL 60015 (US). BAXTER HEALTHCARE S.A. [CH/CH]; Thurgauerstrasse 130, CH-8152 Glattpark (Opfikon) (CH).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** MITTERER, Arthur [AT/AT]; Schwarzeckerweg 10, A-2304 Orth/Donau (AT). HASSLACHER, Meinhard [AT/AT]; Vorgartenstrasse 221/1/7, A-1020 Vienna (AT). MAYER, Christa [AT/AT]; Bahnhofstrasse 2, A-24 12 Wolfsthal (AT).
- (74) **Agents:** ALLIKIAN, Michael, J. et al; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, 6300 Willis Tower, Chicago, IL 60606-6357 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.1 7(U))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.1 7(Hi))

**Published:**

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2011/022657 A1

(54) **Title:** PURIFICATION OF VWF FOR INCREASED REMOVAL OF NON-LIPID ENVELOPED VIRUSES

(57) **Abstract:** The present invention provides methods for purifying Von Willebrand factor (VWF) for increased removal of non-lipid enveloped viruses.

# **PURIFICATION OF VWF FOR INCREASED REMOVAL OF NON-LIPID ENVELOPED VIRUSES**

## **CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims the priority benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 61/235,570, filed August 20, 2009, the disclosure of which is incorporated herein by reference in its entirety.

## **FIELD OF THE INVENTION**

[0002] Generally, the invention relates to methods of purifying VWF for increased removal of non-lipid enveloped viruses.

## **BACKGROUND OF THE INVENTION**

[0003] Von Willebrand factor (VWF) is a glycoprotein circulating in plasma as a series of multimers ranging in size from about 500 to 20,000 kD. Multimeric forms of VWF are composed of 250 kD polypeptide subunits linked together by disulfide bonds. VWF mediates initial platelet adhesion to the sub-endothelium of the damaged vessel wall. Only the larger multimers exhibit hemostatic activity. It is assumed that endothelial cells secrete large polymeric forms of VWF and those forms of VWF which have a low molecular weight (low molecular weight VWF) arise from proteolytic cleavage. The multimers having large molecular masses are stored in the Weibel-Pallade bodies of endothelial cells and liberated upon stimulation.

[0004] VWF is synthesized by endothelial cells and megakaryocytes as prepro-VWF that consists to a large extent of repeated domains. Upon cleavage of the signal peptide, pro-VWF dimerizes through disulfide linkages at its C-terminal region. The dimers serve as protomers for multimerization, which is governed by disulfide linkages between the free end termini. The assembly to multimers is followed by the proteolytic removal of the propeptide sequence (Leyte et al., Biochem. J. 274 (1991), 257-261).

[0005] The primary translation product predicted from the cloned cDNA of VWF is a 2813-residue precursor polypeptide (prepro-VWF). The prepro-VWF consists of a 22 amino acid signal peptide and a 741 amino acid propeptide, with the mature VWF comprising 2050 amino acids (Ruggeri Z.A., and Ware, J., FASEB J., 308-316 (1993)).

[0006] Defects in VWF are causal to Von Willebrand disease (VWD), which is characterized by a more or less pronounced bleeding phenotype. VWD type 3 is the most severe form in which VWF is completely missing, and VWD type 1 relates to a quantitative loss of VWF and its phenotype can be very mild. VWD type 2 relates to qualitative defects of VWF and can be as severe as VWD type 3. VWD type 2 has many sub forms, some being associated with the loss or the decrease of high molecular weight multimers. Von Willebrand syndrome type 2a (VWS-2A) is characterized by a loss of both intermediate and large multimers. VWS-2B is characterized by a loss of highest-molecular-weight multimers. Other diseases and disorders related to VWF are known in the art.

[0007] The removal or inactivation of non-lipid enveloped viruses from therapeutic protein solutions has traditionally been accomplished by treatment with physical methods like high temperature (*e.g.*, dry heat, vapor heat, pasteurization), irradiation with high energy rays (*e.g.*, ultraviolet (UV) rays or beta radiation), low pH, nanofiltration or by chromatographic procedures, in particular affinity chromatography. However, these procedures are often ineffective when purifying a high molecular weight protein such as VWF which does not pass through a nanofilter and/or loses its potency or molecular integrity upon treatment with heat or radiation.

[0008] Current regulatory guidelines ask manufacturers to address the issue of reduction and/or inactivation of both lipid enveloped and non-lipid enveloped viruses for recombinant pharmaceutical products. The ICH "Guideline on Viral Safety Evaluations of Biotechnology Products" (Federal Register, 1998, 63(185): 51074- 51084) gives manufacturers flexibility how to address viral issues taking into account the type of product, the production process and the risk of potentially contaminating viruses. These guidelines point out that the risk of viral contamination is a feature common to all biotechnology products derived from cell lines. Such contamination could have serious clinical consequences and can arise from the contamination of the source cell lines themselves (cell substrates) or from adventitious introduction of virus during production.

[0009] Whereas the inactivation of lipid-enveloped viruses can be performed very effectively by a solvent/detergent (*SfD*) treatment approach, the inactivation or removal of non-lipid-enveloped model viruses (NLEVs) can be challenging due to their small size and physical stability.

[0010] Thus there exists a need in the art to develop methods to efficiently inactivate or remove non-lipid enveloped viruses during the purification of VWF.

### SUMMARY OF THE INVENTION

[0011] The present invention provides an efficient method for purifying VWF for increased removal of non-lipid enveloped viruses. The present invention provides a novel method of purifying VWF for increased removal of NLEVs by performing the product loading step and the wash step of the purification process at a high pH.

[0012] One method known in the art for purifying polypeptides from NLEVs involves the use of nanofiltration. The principle behind efficient separation of protein and virus using nanofiltration exploits the size difference between the polypeptide and the virus; efficient separation requires the polypeptide to have an effective size smaller than the virus, which allows the polypeptide to pass through the pores of the nanofilter while the virus is retained. If the polypeptide and virus are of a comparable size relative to each other, however, separation is problematic because either the polypeptide and virus both pass through the nanofilter pores or neither do. The methods disclosed herein overcome this problem by using a cation exchange resin rather than nanofiltration and loading and/or washing the resin at a sufficiently high pH to separate the polypeptide from the virus.

[0013] Without being bound by theory, the methods disclosed herein are useful for improved removal of NLEV from polypeptide solutions wherein the polypeptide is of a certain size and/or conformation. A polypeptide of a sufficiently large size is likely to have localized charge characteristics at or above the isoelectric point of the polypeptide, *i.e.*, regions of the polypeptide can maintain localized positive or negative charges, thereby allowing the polypeptide to adsorb to the column resin while the virus flows through. This uneven charge distribution over the length of a polypeptide allows the polypeptide to remain attached to the resin despite loading and/or washing of the resin at a high pH.

[0014] The invention provides a method for removing a non-lipid enveloped virus from a protein-containing solution comprising loading a protein in the solution onto a cation

exchange resin, and washing the resin with a buffer at a pH higher than the isoelectric point of the protein to elute the virus. In one aspect, the protein is loaded onto the resin in a buffer at a pH higher than that of the isoelectric point of the protein to elute the virus. In another aspect, the protein is loaded onto the resin in a buffer that is not the buffer used in the wash step, and the resin is subsequently washed with the buffer that is at a pH higher than an isoelectric point of the protein.

[0015] In one embodiment, a method for removing a non-lipid enveloped virus from a protein-containing solution is provided comprising applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein, and washing the cation exchange resin with a first wash buffer to form an eluate, said first wash buffer having a pH that is equal to or lower than the solution applied to the cation exchange resin.

[0016] In one aspect, the pH of the solution is about 1 pH unit above the isoelectric point of the protein. In other aspects, the pH of the solution is about 1.1, or about 1.2, or about 1.3, or about 1.4, or about 1.5, or about 1.6, or about 1.7, or about 1.8, or about 1.9, or about 2.0, or about 2.1, or about 2.2, or about 2.3, or about 2.4, or about 2.5, or about 2.6, or about 2.7, or about 2.8, or about 2.9, or about 3.0, or about 3.1, or about 3.2, or about 3.3, or about 3.4, or about 3.5, or about 3.6, or about 3.7, or about 3.8, or about 3.9, or about 4.0, or about 4.1, or about 4.2, or about 4.3, or about 4.4, or about 4.5, or about 4.6, or about 4.7, or about 4.8, or about 4.9, or about 5.0, or about 5.1, or about 5.2, or about 5.3, or about 5.4, or about 5.5, or about 5.6, or about 5.7, or about 5.8, or about 5.9, or about 6.0 or more pH units or more above the isoelectric point of the protein. In these embodiments, the pH is greater than about 7. In a related aspect, the pH of the protein-containing solution is about 7.0. In other aspects, the pH of the protein-containing solution is about 7.1, or about 7.2, or about 7.3, or about 7.4, or about 7.5, or about 7.6, or about 7.7, or about 7.8, or about 7.9, or about 8.0, or about 8.1, or about 8.2, or about 8.3, or about 8.4, or about 8.5, or about 8.6, or about 8.7, or about 8.8, or about 8.9, or about 9.0, or about 9.1, or about 9.2, or about 9.3, or about 9.4, or about 9.5, or about 9.6, or about 9.7, or about 9.8, or about 9.9, or about 10.0, or about 10.1, or about 10.2, or about 10.3, or about 10.4, or about 10.5, or about 10.6, or about 10.7, or about 10.8, or about 10.9, or about 11.0, or about 11.1, or about 11.2, or about 11.3, or about 11.4, or about 11.5, or about 11.6, or about 11.7, or about 11.8, or about 11.9, or about 12.0, or about 12.1, or about 12.2, or about 12.3, or about 12.4, or about 12.5, or about 12.6, or about 12.7, or about 12.8, or about 12.9, or about 13.0 or higher.

[0017] In another embodiment, a method is provided for removing a non-lipid enveloped virus from a protein-containing solution comprising applying the solution to a cation exchange resin, washing the cation exchange resin with a first wash buffer at a pH higher than the pH of the solution applied to the cation exchange resin, and washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer. In one aspect, the pH of the first wash buffer is about 1 pH unit above the pH of the solution applied to the cation exchange resin. In other aspects, the pH of the first wash buffer is about 0.1, or about 0.2, or about 0.3, or about 0.4, or about 0.5, or about 0.6, or about 0.7, or about 0.8, or about 0.9, or about 1.1, or about 1.2, or about 1.3, or about 1.4, or about 1.5, or about 1.6, or about 1.7, or about 1.8, or about 1.9, or about 2.0, or about 2.1, or about 2.2, or about 2.3, or about 2.4, or about 2.5, or about 2.6, or about 2.7, or about 2.8, or about 2.9, or about 3.0, or about 3.1, or about 3.2, or about 3.3, or about 3.4, or about 3.5, or about 3.6, or about 3.7, or about 3.8, or about 3.9, or about 4.0, or about 4.1, or about 4.2, or about 4.3, or about 4.4, or about 4.5, or about 4.6, or about 4.7, or about 4.8, or about 4.9, or about 5.0, or about 5.1, or about 5.2, or about 5.3, or about 5.4, or about 5.5, or about 5.6, or about 5.7, or about 5.8, or about 5.9, or about 6.0, or about 6.1, or about 6.2, or about 6.3, or about 6.4, or about 6.5, or about 6.6, or about 6.7, or about 6.8, or about 6.9, or about 7.0, or about 7.1, or about 7.2, or about 7.3, or about 7.4, or about 7.5, or about 7.6, or about 7.7, or about 7.8, or about 7.9, or about 8 or about 8.1, or about 8.2, or about 8.3, or about 8.4, or about 8.5, or about 8.6, or about 8.7, or about 8.8, or about 8.9, or about 9, or about 9.1, or about 9.2, or about 9.3, or about 9.4, or about 9.5, or about 9.6, or about 9.7, or about 9.8, or about 9.9, or about 10 or more pH units or more above the isoelectric point of the protein. In these embodiments, the pH of the first wash buffer is greater than about 7. In other aspects, the pH of the first wash buffer is about 7.1, or about 7.2, or about 7.3, or about 7.4, or about 7.5, or about 7.6, or about 7.7, or about 7.8, or about 7.9, or about 8.0, or about 8.1, or about 8.2, or about 8.3, or about 8.4, or about 8.5, or about 8.6, or about 8.7, or about 8.8, or about 8.9, or about 9.0, or about 9.1, or about 9.2, or about 9.3, or about 9.4, or about 9.5, or about 9.6, or about 9.7, or about 9.8, or about 9.9, or about 10.0, or about 10.1, or about 10.2, or about 10.3, or about 10.4, or about 10.5, or about 10.6, or about 10.7, or about 10.8, or about 10.9, or about 11.0 or higher.

[0018] In an embodiment, the protein in the solution is a polypeptide having a molecular mass of at least about 150 kilodaltons. In various aspects, the protein in the solution is a polypeptide having a molecular mass of at least about 175 kilodaltons, or about 180

kilodaltons, or about 190 kilodaltons, or about 200 kilodaltons, or about 210 kilodaltons, or about 220 kilodaltons, or about 230 kilodaltons, or about 240 kilodaltons, or about 250 kilodaltons, or about 260 kilodaltons, or about 270 kilodaltons, or about 280 kilodaltons, or about 290 kilodaltons, or about 300 kilodaltons, or about 310 kilodaltons, or about 320 kilodaltons, or about 330 kilodaltons, or about 340 kilodaltons, or about 350 kilodaltons, or about 360 kilodaltons, or about 370 kilodaltons, or about 380 kilodaltons, or about 390 kilodaltons, or about 400 kilodaltons, or about 410 kilodaltons, or about 420 kilodaltons, or about 430 kilodaltons, or about 440 kilodaltons, or about 450 kilodaltons, or about 460 kilodaltons, or about 470 kilodaltons, or about 480 kilodaltons, or about 490 kilodaltons, or about 500 kilodaltons or more. As described herein, polypeptides also comprise multimeric structures and such multimeric structures, in various aspects, have a molecular mass of at least about 500 kilodaltons. In related aspects, the multimeric structures have a molecular mass of at least about 510, or about 520, or about 530, or about 540, or about 550, or about 560, or about 570, or about 580, or about 590, or about 600, or about 610, or about 620, or about 630, or about 640, or about 650, or about 660, or about 670, or about 680, or about 690, or about 700, or about 710, or about 720, or about 730, or about 740, or about 750, or about 760, or about 770, or about 780, or about 790, or about 800, or about 810, or about 820, or about 830, or about 840, or about 850, or about 860, or about 870, or about 880, or about 890, or about 900, or about 910, or about 920, or about 930, or about 940, or about 950, or about 960, or about 970, or about 980, or about 990 kilodaltons, or about 1 megadalton, or about 1.1 megadaltons, or about 1.2 megadaltons, or about 1.3 megadaltons, or about 1.4 megadaltons, or about 1.5 megadaltons, or about 1.6 megadaltons, or about 1.7 megadaltons, or about 1.8 megadaltons, or about 1.9 megadaltons, or about 2.0 megadaltons, or about 2.1 megadaltons, or about 2.2 megadaltons, or about 2.3 megadaltons, or about 2.4 megadaltons, or about 2.5 megadaltons, or about 2.6 megadaltons, or about 2.7 megadaltons, or about 2.8 megadaltons, or about 2.9 megadaltons, or about 3.0 megadaltons, or about 3.1 megadaltons, or about 3.2 megadaltons, or about 3.3 megadaltons, or about 3.4 megadaltons, or about 3.5 megadaltons, or about 3.6 megadaltons, or about 3.7 megadaltons, or about 3.8 megadaltons, or about 3.9 megadaltons, or about 4.0 megadaltons, or about 4.1 megadaltons, or about 4.2 megadaltons, or about 4.3 megadaltons, or about 4.4 megadaltons, or about 4.5 megadaltons, or about 4.6 megadaltons, or about 4.7 megadaltons, or about 4.8 megadaltons, or about 4.9 megadaltons, or about 5.0 megadaltons or more.

[0019] In some embodiments, the cation exchange resin has a negatively charged group selected from the group consisting of carboxymethyl (CM), sulfoalkyl (SP, SE), sulphate and methylsulfonate (S) as well as any other negatively charged ligand.

[0020] In a further embodiment, the protein is a blood coagulation protein. In various aspects, the blood coagulation protein is selected from the group consisting of Factor VIII, von Willebrand factor, FI (Fibrinogen), FV (Proaccelerin), FXI (plasma-thromboplastin antecedent), and FXIII (fibrin stabilizing factor).

[0021] In an embodiment, a method for removing a non-lipid enveloped virus from a von Willebrand (VWF)-containing solution is provided comprising applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein and washing the cation exchange resin with a first wash buffer to form an eluate, said first wash buffer having a pH that is equal to or lower than the solution applied to the cation exchange resin.

[0022] In another embodiment, a method for removing a non-lipid enveloped virus from a VWF-containing solution is provided comprising applying the solution to a cation exchange resin, washing the cation exchange resin with a first wash buffer at a pH higher than the pH of the solution applied to the cation exchange resin and washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer.

[0023] In a further embodiment, a method for removing a non-lipid enveloped virus from a VWF-containing solution comprising applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein and washing the cation exchange resin with a first wash buffer at a pH higher than the isoelectric point of the protein applied to the cation exchange resin; and washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer.

#### **BRIEF DESCRIPTION OF THE FIGURES**

[0024] **Figure 1** shows the result of an SDS-PAGE separation followed by silver staining (A) and Western Blot (B) analysis for residual rFVIII.

[0025] **Figure 2** shows the stained gel of the UNO S runs with MMV and REO virus spiked samples.



[0026] **Figure 3** shows the stained gel of the UNO S runs with MMV and REO virus spiked samples.

[0027] **Figure 4** shows the results of subjecting the purified preparations of rVWF obtained by the process variants to proteolytic digestion by V8 protease in the native state and separating the resulting peptides by RP-HPLC.

[0028] **Figure 5** shows the results of subjecting the purified preparations of rVWF obtained by the process variants to trypsin in the denatured state and separating the resulting peptides by RP-HPLC.

### **DETAILED DESCRIPTION OF THE INVENTION**

[0029] The present invention relates to a method for purifying VWF with increased removal of non-lipid enveloped viruses. The methods of the invention are applicable in column (*i.e.*, chromatography) as well as batch (*i.e.*, without column hardware) mode.

[0030] The method of the present invention utilizes a purification method on a cation exchange resin for the increased removal of non-lipid enveloped viruses. Previous methods of purification of VWF using cation exchange chromatography were performed at a neutral pH. These methods allowed for the manufacture of purified VWF of good yield and purity, but surprisingly the process had no capacity to remove non-lipid enveloped viruses.

#### **Definition of terms**

[0031] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton, et al., *DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY* (2d ed. 1994); *THE CAMBRIDGE DICTIONARY OF SCIENCE AND TECHNOLOGY* (Walker ed., 1988); *THE GLOSSARY OF GENETICS, 5TH ED.*, R. Rieger, et al. (eds.), Springer Verlag (1991); and Hale and Marham, *THE HARPER COLLINS DICTIONARY OF BIOLOGY* (1991).

[0032] Each publication, patent application, patent, and other reference cited herein is incorporated by reference in its entirety to the extent that it is not inconsistent with the present disclosure.

[0033] It is noted here that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise.

[0034] As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

[0035] As used herein the terms "express," "expressing" and "expression" mean allowing or causing the information in a gene or DNA sequence to become manifest, for example, producing a protein by activating the cellular functions involved in transcription and translation of a corresponding gene or DNA sequence. A DNA sequence is expressed in or by a cell to form an "expression product" such as a protein. The expression product itself, e.g. the resulting protein, may also be said to be "expressed." An expression product can be characterized as intracellular, extracellular or secreted. The term "intracellular" means inside a cell. The term "extracellular" means outside a cell, e.g., certain types of transmembrane proteins. A substance is "secreted" by a cell if it appears in significant measure outside the cell, from somewhere on or inside the cell.

[0036] As used herein a "polypeptide" refers to a polymer composed of amino acid residues, structural variants, related naturally-occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds. Synthetic polypeptides can be prepared, for example, using an automated polypeptide synthesizer. The term "protein" typically refers to large polypeptides. The term "peptide" typically refers to short polypeptides. The term "polypeptide" also includes polymeric structures. Therefore, a "polypeptide" may be a monomer, dimer, trimer, or larger multimeric structure. These multimeric structures can be up to 5 megadaltons or larger.

[0037] As used herein, the "isoelectric point" is the pH value at which the net electric charge of a polypeptide in an aqueous solution is zero.

[0038] As used herein a "fragment" of a polypeptide is meant to refer to any portion of a polypeptide or protein smaller than the full-length polypeptide or protein expression product.

[0039] As used herein an "analog" refers to any of two or more polypeptides substantially similar in structure and having the same biological activity, but can have varying degrees of activity, to either the entire molecule, or to a fragment thereof. Analogs differ in the composition of their amino acid sequences based on one or more mutations involving substitution of one or more amino acids for other amino acids. Substitutions can be

conservative or non-conservative based on the physico-chemical or functional relatedness of the amino acid that is being replaced and the amino acid replacing it.

[0040] As used herein a "variant" refers to a polypeptide, protein or analog thereof that is modified to comprise additional chemical moieties not normally a part of the molecule. Such moieties may modulate the molecule's solubility, absorption, biological half-life, etc. The moieties may alternatively decrease the toxicity of the molecule and eliminate or attenuate any undesirable side effect of the molecule, etc. Moieties capable of mediating such effects are disclosed in Remington's Pharmaceutical Sciences (1980). Procedure for coupling such moieties to a molecule are well known in the art. For example, the variant may be a blood clotting factor having a chemical modification which confers a longer half-life *in vivo* to the protein. In various aspects, polypeptides are modified by glycosylation, pegylation, and/or polysialylation.

### **Recombinant VWF**

[0041] The polynucleotide and amino acid sequences of prepro-VWF are set out in SEQ ID NO:1 and SEQ ID NO:2, respectively, and are available at GenBank Accession Nos. NM\_000552 and NP\_000543, respectively. The amino acid sequence corresponding to the mature VWF protein is set out in SEQ ID NO: 3 (corresponding to amino acids 764-2813 of the full length prepro-VWF amino acid sequence).

[0042] One form of useful rVWF has at least the property of *in vivo*-stabilizing, e.g. binding, of at least one Factor VIII (FVIII) molecule and having optionally a glycosylation pattern which is pharmacologically acceptable. Specific examples thereof include VWF without A2 domain thus resistant to proteolysis (Lankhof et al., *Thromb. Haemost.* 77: 1008-1013, 1997), and the VWF fragment from Val 449 to Asn 730 including the glycoprotein Ib-binding domain and binding sites for collagen and heparin (Pietu et al., *Biochem. Biophys. Res. Commun.* 164: 1339-1347, 1989). The determination of the ability of a VWF to stabilize at least one FVIII molecule can be carried out in VWF-deficient mammals according to methods known in the state in the art.

[0043] The rVWF of the present invention may be produced by any method known in the art. One specific example is disclosed in WO86/06096 published on Oct. 23, 1986 and U.S. Patent Application No. 07/559,509, filed on Jul. 23, 1990, which is incorporated herein by reference with respect to the methods of producing recombinant VWF. Thus, methods are known in the art for (i) the production of recombinant DNA by genetic engineering, e.g. via

reverse transcription of RNA and/or amplification of DNA, (ii) introducing recombinant DNA into procaryotic or eucaryotic cells by transfection, e.g. via electroporation or microinjection, (iii) cultivating said transformed cells, e.g. in a continuous or batchwise manner, (iv) expressing VWF, e.g. constitutively or upon induction, and (v) isolating said VWF, e.g. from the culture medium or by harvesting the transformed cells, in order to (vi) obtain purified rVWF, e.g. via anion exchange chromatography or affinity chromatography. A recombinant VWF may be made in transformed host cells using recombinant DNA techniques well known in the art. For instance, sequences coding for the polypeptide could be excised from DNA using suitable restriction enzymes.

[0044] Alternatively, the DNA molecule could be synthesized using chemical synthesis techniques, such as the phosphoramidate method. Also, a combination of these techniques could be used.

[0045] The invention also provides vectors encoding polypeptides of the invention in an appropriate host. The vector comprises the polynucleotide that encodes the polypeptide operatively linked to appropriate expression control sequences. Methods of effecting this operative linking, either before or after the polynucleotide is inserted into the vector, are well known. Expression control sequences include promoters, activators, enhancers, operators, ribosomal binding sites, start signals, stop signals, cap signals, polyadenylation signals, and other signals involved with the control of transcription or translation. The resulting vector having the polynucleotide therein is used to transform an appropriate host. This transformation may be performed using methods well known in the art.

[0046] Any of a large number of available and well-known host cells may be used in the practice of this invention. The selection of a particular host is dependent upon a number of factors recognized by the art, including, for example, compatibility with the chosen expression vector, toxicity of the peptides encoded by the DNA molecule, rate of transformation, ease of recovery of the peptides, expression characteristics, bio-safety and costs. A balance of these factors must be struck with the understanding that not all host cells are equally effective for the expression of a particular DNA sequence. Within these general guidelines, useful microbial host cells include bacteria, yeast and other fungi, insects, plants, mammalian (including human) cells in culture, or other hosts known in the art.

[0047] Next, the transformed host is cultured and purified. Host cells may be cultured under conventional fermentation conditions so that the desired compounds are expressed.

Such fermentation conditions are well known in the art. Finally, the polypeptides are purified from culture by methods well known in the art.

[0048] Depending on the host cell utilized to express a compound of the invention, carbohydrate (oligosaccharide) groups may conveniently be attached to sites that are known to be glycosylation sites in proteins. Generally, O-linked oligosaccharides are attached to serine (Ser) or threonine (Thr) residues while N-linked oligosaccharides are attached to asparagine (Asn) residues when they are part of the sequence Asn-X-Ser/Thr, where X can be any amino acid except proline. X is preferably one of the 19 naturally occurring amino acids not counting proline. The structures of N-linked and O-linked oligosaccharides and the sugar residues found in each type are different. One type of sugar that is commonly found on both is N-acetylneuraminic acid (referred to as sialic acid). Sialic acid is usually the terminal residue of both N-linked and O-linked oligosaccharides and, by virtue of its negative charge, may confer acidic properties to the glycosylated compound. Such site(s) may be incorporated in the linker of the compounds of this invention and are preferably glycosylated by a cell during recombinant production of the polypeptide compounds (e.g., in mammalian cells such as CHO, BHK, COS). However, such sites may further be glycosylated by synthetic or semi-synthetic procedures known in the art.

[0049] Alternatively, the compounds may be made by synthetic methods. For example, solid phase synthesis techniques may be used. Suitable techniques are well known in the art, and include those described in Merrifield (1973), *Chem. Polypeptides*, pp. 335-61 (Katsoyannis and Panayotis eds.); Merrifield (1963), *J. Am. Chem. Soc.* 85: 2149; Davis et al. (1985), *Biochem. Intl.* 10: 394-414; Stewart and Young (1969), *Solid Phase Peptide Synthesis*; U.S. Pat. No. 3,941,763; Finn et al. (1976), *The Proteins* (3rd ed.) 2: 105-253; and Erickson et al. (1976), *The Proteins* (3rd ed.) 2: 257-527'. Solid phase synthesis is the preferred technique of making individual peptides since it is the most cost-effective method of making small peptides.

#### **Fragments, variants and analogs of VWF**

[0050] Methods for preparing polypeptide fragments, variants or analogs are well-known in the art.

[0051] Fragments of a polypeptide are prepared using, without limitation, enzymatic cleavage (e.g., trypsin, chymotrypsin) and also using recombinant means to generate a polypeptide fragments having a specific amino acid sequence. Polypeptide fragments may be

generated comprising a region of the protein having a particular activity, such as a multimerization domain or any other identifiable VWF domain known in the art.

[0052] Variants of a polypeptide are contemplated to include human and non-human forms of VWF (*e.g.*, murine VWF). Also contemplated by the methods herein are chimeric polypeptides comprising, *e.g.*, a mouse/human fusion polypeptide.

[0053] Methods of making polypeptide analogs are also well-known. Amino acid sequence analogs of a polypeptide can be substitutional, insertional, addition or deletion analogs. Deletion analogs, including fragments of a polypeptide, lack one or more residues of the native protein which are not essential for function or immunogenic activity. Insertional analogs involve the addition of, *e.g.*, amino acid(s) at a non-terminal point in the polypeptide. This analog may include insertion of an immunoreactive epitope or simply a single residue. Addition analogs, including fragments of a polypeptide, include the addition of one or more amino acids at either of both termini of a protein and include, for example, fusion proteins.

[0054] Substitutional analogs typically exchange one amino acid of the wild-type for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide without the loss of other functions or properties. In one aspect, substitutions are conservative substitutions. By "conservative amino acid substitution" is meant substitution of an amino acid with an amino acid having a side chain of a similar chemical character. Similar amino acids for making conservative substitutions include those having an acidic side chain (glutamic acid, aspartic acid); a basic side chain (arginine, lysine, histidine); a polar amide side chain (glutamine, asparagine); a hydrophobic, aliphatic side chain (leucine, isoleucine, valine, alanine, glycine); an aromatic side chain (phenylalanine, tryptophan, tyrosine); a small side chain (glycine, alanine, serine, threonine, methionine); or an aliphatic hydroxyl side chain (serine, threonine).

[0055] Analogs may be substantially homologous or substantially identical to the recombinant VWF from which they are derived. Preferred analogs are those which retain at least some of the biological activity of the wild-type polypeptide, *e.g.* blood clotting activity.

[0056] Polypeptide variants contemplated include polypeptides chemically modified by such techniques as ubiquitination, glycosylation, including polysialation, conjugation to therapeutic or diagnostic agents, labeling, covalent polymer attachment such as pegylation (derivatization with polyethylene glycol), introduction of non-hydrolyzable bonds, and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not

normally occur in human proteins. Variants retain the same or essentially the same binding properties of non-modified molecules of the invention. Such chemical modification may include direct or indirect (e.g., via a linker) attachment of an agent to the VWF polypeptide. In the case of indirect attachment, it is contemplated that the linker may be hydrolyzable or non-hydrolyzable.

[0057] Preparing pegylated polypeptide analogs will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the binding construct polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of PEG: protein, the greater the percentage of poly-pegylated product. In some embodiments, the binding construct will have a single PEG moiety at the N-terminus. Polyethylene glycol (PEG) may be attached to the blood clotting factor to provide a longer half-life *in vivo*. The PEG group may be of any convenient molecular weight and may be linear or branched. The average molecular weight of the PEG ranges from about 2 kiloDalton ("kD") to about 100 kDa, from about 5 kDa to about 50 kDa, or from about 5 kDa to about 10 kDa. The PEG groups are attached to the blood clotting factor via acylation or reductive alkylation through a natural or engineered reactive group on the PEG moiety (e.g., an aldehyde, amino, thiol, or ester group) to a reactive group on the blood clotting factor (e.g., an aldehyde, amino, or ester group) or by any other technique known in the art.

[0058] Methods for preparing polysialylated polypeptide are described in United States Patent Publication 20060160948, Fernandes et Gregoriadis; *Biochim. Biophys. Acta* 1341: 26-34, 1997, and Saenko et al., *Haemophilia* 12:42-51, 2006. Briefly, a solution of colominic acid containing 0.1 M NaIO<sub>4</sub> is stirred in the dark at room temperature to oxidize the CA. The activated CA solution is dialyzed against, e.g., 0.05 M sodium phosphate buffer, pH 7.2 in the dark and this solution was added to a rVWF solution and incubated for 18 h at room temperature in the dark under gentle shaking. Free reagents can then be separated from the rVWF-polysialic acid conjugate by ultrafiltration/diafiltration. Conjugation of rVWF with polysialic acid may also be achieved using glutaraldehyde as cross-linking reagent (Migneault et al., *Biotechniques* 37: 790-796, 2004).

[0059] It is further contemplated that a polypeptide of the invention may be a fusion protein with a second agent which is a polypeptide. In one embodiment, the second agent

which is a polypeptide, without limitation, is an enzyme, a growth factor, an antibody, a cytokine, a chemokine, a cell-surface receptor, the extracellular domain of a cell surface receptor, a cell adhesion molecule, or fragment or active domain of a protein described above. In a related embodiment, the second agent is a blood clotting factor such as Factor VIII, Factor VII, Factor IX. The fusion protein contemplated is made by chemical or recombinant techniques well-known in the art.

[0060] It is also contemplated that prepro-VWF and pro-VWF polypeptides may provide a therapeutic benefit in the formulations of the present invention. For example, US Patent No. 7,005,502 describes a pharmaceutical preparation comprising substantial amounts of pro-VWF that induces thrombin generation in the presence of platelets *in vitro*. In addition to recombinant, biologically active fragments, variants, or analogs of the naturally-occurring mature VWF, the present invention contemplates the use of recombinant biologically active fragments, variants, or analogs of the prepro-VWF (set out in SEQ ID NO:2) or pro-VWF polypeptides (amino acid residues 23 to 764 of SEQ ID NO: 2) in the formulations described herein.

[0061] Polynucleotides encoding fragments, variants and analogs may be readily generated by a worker of skill to encode biologically active fragments, variants, or analogs of the naturally-occurring molecule that possess the same or similar biological activity to the naturally-occurring molecule. These polynucleotides can be prepared using PCR techniques, digestion/ligation of DNA encoding molecule, and the like. Thus, one of skill in the art will be able to generate single base changes in the DNA strand to result in an altered codon and a missense mutation, using any method known in the art, including, but not limited to site-specific mutagenesis. As used herein, the phrase "moderately stringent hybridization conditions" means, for example, hybridization at 42°C in 50% formamide and washing at 60°C in 0.1 x SSC, 0.1% SDS. It is understood by those of skill in the art that variation in these conditions occurs based on the length and GC nucleotide base content of the sequences to be hybridized. Formulas standard in the art are appropriate for determining exact hybridization conditions. See Sambrook et al., 9.47-9.51 in *Molecular Cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989).

### **Methods of Producing VWF**

[0062] Industrially, VWF, in particular human recombinant VWF (rVWF), is synthesized and expressed together with rFVIII in a genetically engineered CHO cell line. The function



of the co-expressed rVWF is to stabilize rFVIII in the cell culture process. rVWF is synthesized in the cell as the pro-form, containing a large pro-peptide attached to the N-terminus. Upon maturation in the endoplasmatic reticulum and Golgi apparatus, the pro-peptide is cleaved off by the action of the cellular protease furin and is secreted as a homopolymer of identical subunits, consisting of dimers of the expressed protein.

### **Purification of VWF**

**[0063]** Provided herein is a method for removing a non-lipid enveloped virus from a protein-containing solution is provided comprising applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein, and washing the cation exchange resin with a first wash buffer to form an eluate, said first wash buffer having a pH that is equal to or lower than the solution applied to the cation exchange resin.

**[0064]** In one aspect, the pH of the solution is about 1 pH unit above the isoelectric point of the protein. In other aspects, the pH of the solution is about 1.2, or about 1.4, or about 1.6, or about 1.8, or about 2.0, or about 2.2, or about 2.4, or about 2.6, or about 2.8, or about 3.0, or about 3.2, or about 3.4, or about 3.6, or about 3.8, or about 4.0, or about 4.2, or about 4.4, or about 4.6, or about 4.8, or about 5.0, or about 5.5, or about 6.0 pH units or more above the isoelectric point of the protein.

**[0065]** In these embodiments, the pH is greater than about 7.. In other aspects, the pH is about 7.1, or about 7.2, or about 7.3, or about 7.4, or about 7.5, or about 7.6, or about 7.7, or about 7.8, or about 7.9, or about 8.0, or about 8.1, or about 8.2, or about 8.3, or about 8.4, or about 8.5, or about 8.6, or about 8.7, or about 8.8, or about 8.9, or about 9.0, or about 9.1, or about 9.2, or about 9.3, or about 9.4, or about 9.5, or about 9.6, or about 9.7, or about 9.8, or about 9.9, or about 10.0, or about 10.1, or about 10.2, or about 10.3, or about 10.4, or about 10.5, or about 10.6, or about 10.7, or about 10.8, or about 10.9, or about 11.0, or about 11.1, or about 11.2, or about 11.3, or about 11.4, or about 11.5, or about 11.6, or about 11.7, or about 11.8, or about 11.9, or about 12.0, or about 12.1, or about 12.2, or about 12.3, or about 12.4, or about 12.5, or about 12.6, or about 12.7, or about 12.8, or about 12.9, or about 13.0 or higher.

**[0066]** In another embodiment, a method is provided for removing a non-lipid enveloped virus from a protein-containing solution comprising applying the solution to a cation exchange resin, washing the cation exchange resin with a first wash buffer at a pH higher than the pH of the solution applied to the cation exchange resin, and washing the cation

exchange resin with a second wash buffer to form an eluate, said second wash buffer having a pH that is equal to or lower than the first wash buffer. In one aspect, the pH of the first wash buffer is about 1 pH unit above the pH of the solution applied to the cation exchange resin. For this stage it is contemplated that the ion exchange media is UNOsphere™ S (BioRad Laboratories, Inc., Hercules, CA), but other cation exchange systems may be used in the practice of the methods. These cation exchange systems are known to those of ordinary skill in the art.

[0067] In other aspects, the pH of the first wash buffer is about 1.1, or about 1.2, or about 1.3, or about 1.4, or about 1.5, or about 1.6, or about 1.7, or about 1.8, or about 1.9, or about 2.0, or about 2.1, or about 2.2, or about 2.3, or about 2.4, or about 2.5, or about 2.6, or about 2.7, or about 2.8, or about 2.9, or about 3.0, or about 3.1, or about 3.2, or about 3.3, or about 3.4, or about 3.5, or about 3.6, or about 3.7, or about 3.8, or about 3.9, or about 4.0, or about 4.1, or about 4.2, or about 4.3, or about 4.4, or about 4.5, or about 4.6, or about 4.7, or about 4.8, or about 4.9, or about 5.0, or about 5.1, or about 5.2, or about 5.3, or about 5.4, or about 5.5, or about 5.6, or about 5.7, or about 5.8, or about 5.9, or about 6.0 pH units or more above the isoelectric point of the protein. In these embodiments, the pH of the first wash buffer is greater than about 7. In other aspects, the pH of the first wash buffer is about 7.1, or about 7.2, or about 7.3, or about 7.4, or about 7.5, or about 7.6, or about 7.7, or about 7.8, or about 7.9, or about 8.0, or about 8.1, or about 8.2, or about 8.3, or about 8.4, or about 8.5, or about 8.6, or about 8.7, or about 8.8, or about 8.9, or about 9.0, or about 9.1, or about 9.2, or about 9.3, or about 9.4, or about 9.5, or about 9.6, or about 9.7, or about 9.8, or about 9.9, or about 10.0, or about 10.1, or about 10.2, or about 10.3, or about 10.4, or about 10.5, or about 10.6, or about 10.7, or about 10.8, or about 10.9, or about 11.0, or about 11.1, or about 11.2, or about 11.3, or about 11.4, or about 11.5, or about 11.6, or about 11.7, or about 11.8, or about 11.9, or about 12.0, or about 12.1, or about 12.2, or about 12.3, or about 12.4, or about 12.5, or about 12.6, or about 12.7, or about 12.8, or about 12.9, or about 13.0 or higher.

[0068] The following examples are not intended to be limiting but only exemplary of specific embodiments of the invention.

## EXAMPLES

### Example 1

[0069] Viruses and cells used in the assays described below are as follows.

[0070] REO-3 (Family Reoviridae ; non-enveloped dsRNA virus), Strain Dearing (ATCC VR-824) was obtained from the ATCC. The virus was propagated and titrated on Vero cells obtained from ECACC (84113001). MMV (Family Parvoviridae ; non-enveloped ssDNA virus), prototype strain (ATCC VR-1346), was obtained from the American Type Culture Collection, Rockville, Maryland. The virus was propagated and titrated on A9 cells (ATCC CCL-I .4). PPV (Family Parvoviridae ; non-enveloped ssDNA virus), strain Tennessee (BRFF #PP951024), was obtained from Biological Research Faculty & Facility, Ijamsville, Maryland. The virus was propagated and titrated on PK-13 cells (ATCC CRL-6489). EMCV (Family Picornaviridae ; non-enveloped ssRNA) (ATCC #VR-129B) was obtained from the American Type Culture Collection. The virus was propagated and titrated on Vero cells (European Collection of Cell Cultures, ECACC, #84113001). HadV (Family Adenoviridae ; non-enveloped dsDNA), strain Adenoid 75 (ATCC VR-5), was obtained from the American Type Culture Collection. The virus was propagated and titrated on HeLa cells (ATCC CCL-2).

[0071] The steps involved in an exemplary VWF purification process comprise:

- Immune affinity chromatography of cell culture supernatant
  - i. Flow-through fraction
- Anion Exchange (*e.g.*, trimethylaminoethyl anion exchange column)
- Filtration (0.45/0.2  $\mu\text{m}$ )
- Anion Exchange (*e.g.*, Mustang Q (Pall Corporation))
- Virus Inactivation (*e.g.*, using solvent/detergent treatment)
- Filtration (0.8/0.65  $\mu\text{m}$ )
- Cation Exchange (*e.g.*, UNO S column)
- Ultrafiltration/Concentration
- Filtration (0.45/0.2  $\mu\text{m}$ )
- Gel Filtration (Superose 6 prep grade (GE Life Sciences))

[0072] **Optimization of UNO S Step.** During the UNO S step rVWF is bound to a strong cation exchange resin while some of the impurities pass through. After washing the column with increased conductivity buffers the bound rVWF is released from the column with a salt

step. During the initial virus removal studies, this step showed at least a significant removal rate for the model REO virus. The conditions of the applied parameters and the corresponding results are listed in Table 1, below.

**Table 1**

Parameter	Standard value	Changed value	MMV reduction factor (log10)
Conductivity load/wash	15 mS/cm	25 mS/cm	n a
pH load/wash/elution	6.5	8.0	0.9
Wash buffer 2	TQA buffer*	TQA buffer with 200mM betaine	0.7
Wash buffer 3	TQA buffer	TQA buffer with 20% ethylen glycol	
Wash buffer 4	TQA buffer	TQA buffer with 10 mM CaCl <sub>2</sub>	0.8
Wash buffer 5	TQA buffer	TQA buffer with 10 mM EDTA	
pH load/wash	6.5	9.0	2.0, 2.1 <sup>1</sup> , 2.12 and 2.12
pH load/wash	6.5	9.0	2.12 for REO virus

\*TQA Buffer: Tris, NaAc, niM NaCl in WFI  
pH 6.3 - 6.7 at 20 - 25°C

[0073] As can be seen in Table 1 the moderate changes in the process parameters (modification of conductivity, pH 8.0 and additives to wash buffers) did not result in a significant improvement of the MMV removal rates. Increasing the pH further to 9.0 reproducibly resulted in a significant removal rate of more than 2 logs for MMV as well as REO virus. This process change is technically easy to implement and the exposure of rVWF to the high pH environment can be kept relatively short (max. 6 hours). Elution of the bound rVWF is performed under neutral conditions.

[0074] The analysis of the virus inactivation capacity of the processes was carried out according to the recommendations of the CPMP guideline 268/95, using the following

formula: 
$$R = \log \left( \frac{V_1 \cdot T_1}{V_2 \cdot T_2} \right)$$

where

R = virus reduction factor

V1 = volume of starting material [ml]

T1 = concentration of virus in starting material [TCID50/ml]

V2 = volume of material after the step [ml]

T2 = concentration of virus after the step [TCID50/ml]

[0075] The volumes and the titers of each spiked sample before and after treatment were used to calculate R. Whenever virus was undetectable, the detection limit was taken as the virus titer for calculation. Calculations were carried out with virus titers ( $\log_{10}[\text{TCID}_{50}/\text{ml}]$ ) given to two decimal places, and only final results, i.e. reduction factors (R), were rounded to the first decimal place.

### Example 2

[0076] The UNO S eluate was concentrated to approximately 800  $\mu\text{g}$  rVWF antigen/ml by ultrafiltration using 30 kDa cut-off modified cellulose membranes to facilitate the trace analysis of impurities and product variants.

### Testing of rVWF

[0077] **Ristocetin Activity.** The Ristocetin Cofactor Activity is determined by a turbidimetric analyzer using a von Willebrand reagent containing stabilized thrombocytes and the antibiotic "ristocetin". The von Willebrand Factor contained in the sample (= Ristocetin Cofactor) causes agglutination of stabilized thrombocytes in the presence of ristocetin. The agglutination reduces the turbidity of the reagent preparation, and the change in optical density is measured by the turbidimetric analyzer. Calibration is performed by the WHO concentrate reference standard #00/514.

[0078] **VWF Antigen.** VWF- samples are tested for their content of vWF-Antigen in an ELISA assay - double sandwich system with two polyclonal antibodies. Measurement of the color reactions on the microtitre plates is performed with a photometer at 490 nm. The concentration of each sample is calculated towards the standard curve with a computer supported ELISA Analysis Program (curve algorithm : Cubic regression). All readings are corrected against the blank.

[0079] **FVIII Binding Activity.** FVIII binding of rVWF under static conditions was determined by an ELISA chromogenic assay (ECA) by incubating a constant amount of rFVIII with a diluted VWF-containing sample. The VWF- FVIII- complex formed was then transferred to a microtiter plate coated with a commercially available polyclonal rabbit anti-human VWF antibody. After incubation, unbound FVIII was removed by a subsequent washing step. Bound FVIII was quantified by a commercially available FVIII chromogenic assay (Technochrom FVIII : C reagent kit, Technoclone, Austria). The blank corrected optical densities (in mOD/min at 405 nm) were plotted against the VWF:Ag concentrations in logarithmic scale.

[0080] **SDS-PAGE Analysis.** Conventional 8% SDS-PAGE analysis under reducing conditions and staining of the gels with Coomassie Blue and Silver Stain can provide insight in the protein composition of rVWF. After transfer of the separated protein bands to a nitrocellulose membrane and immunological staining of the protein with appropriate antibodies against VWF, FVIII and Furin respectively, a comparison of VWF related proteins to total proteins can be made.

[0081] **Multimer Analysis.** The multimeric structure of VWF is analyzed by high-density horizontal SDS agarose gel electrophoresis. In brief, samples are diluted to the same concentration in the range of 0.3-1.0 IU/ml VWF:Ag, incubated with Tris-EDTA-SDS buffer and the multimers separated under non-reducing conditions on an agarose gel. VWF multimers were visualized by in-gel immunostaining with a polyclonal rabbit anti human VWF antibody, followed by alkaline phosphatase (ALP) conjugated goat anti-rabbit IgG using the ALP color development kit. Alternatively, agarose gels were blotted onto a blotting membrane and staining was performed by a polyclonal rabbit anti-human VWF antibody followed by horse radish peroxidase conjugated anti-rabbit IgG. For visualization, electro-chemi-luminescence was used which increases the sensitivity of detection for VWF by at least two magnitudes. Low (1 % agarose) and high resolution (2.5 % agarose) conditions were used to analyze the size distribution of VWF multimers and the multimeric structure, respectively.

[0082] **HPLC Analysis.** Recombinant VWF can be cleaved by GluC (V8 protease) under native conditions to give two main fragments (N-terminal and C-terminal homodimer fragment), which are separated on a reverse phase HPLC C4-column. The fragments are detected by monitoring the UV absorbance at 280 nm.

[0083] **Peptide Mapping.** The primary structure of rVWF was investigated using a peptide mapping approach. Samples of purified rVWF were reduced with dithiothreitol (DTT) and the free, sulfhydryl groups were blocked with 4-vinylpyridin. Sequencing grade trypsin was added to the rVWF and allowed to react for 18 hours. The resulting peptide mixture was separated by reverse phase chromatography. Eluting peptides were detected by on-line UV detection at 214 nm and on-line electrospray ionization mass spectrometry.

[0084] **Test for deamidated rVWF.** The analytical method for the detection of iso-aspartate (one reaction product originating from the de-amidation of asparagine) employs tryptic digestion, followed by the Protein Isoaspartyl Methyltransferase (PIMT) enzymatic

reaction using the ISOQUANT IsoAspartate Detection Kit supplied by Promega. PIMT catalyzes the transfer of a methyl group from the substrate S-adenosyl-L methionine (SAM) to IsoAsp at the carboxyl position, generating S-adenosyl homocysteine (SAH). The stoichiometrically released SAH is detected at a wavelength of 260 nm by a RP HPLC method.

[0085] The analytical data for the product obtained by the different processes are summarized in Table 2.



**Table 2: Analytical Data for pH 9 and Control Runs**

Sample	UNO S #1 (pH 9 run)	UNO S #2 (Control run)	UNO S #3 (Control run pilot scale)
VWF RcoF Activity (U/ml)	36.7	64.8	84.7
VWF Antigen ( $\mu$ g/ml)	715	1090	2010
Specific activity (U/mg)	51.3	59.4	42.1
Collagen binding (U/ml)	63.2	106.7	141.8
Specific collagen binding activity (U/mg)	88.4	97.9	70.5
FVIII binding (%)	51.7	64.6	63.1
CHO protein ( $\mu$ g/ml)	0.09	0.17	n.d.
CHO DNA (pg/ml)	Non detectable	10	n.d.
Furin activity (mU/mi)	< 6.25	< 6.25	<6.25
FVIII Ag (mU/ml)	< 125	152	1650
FVIII Activity (mU/ml)	71	1452	6222
Deamidation (mol%)	5.1	3.7	5.0
SDS-PAGE	See Figure 1		
Multimer pattern (low resolution)	See Figure 2		
Multimer pattern (high resolution )	See Figure 3		
RP-HPLC	See Figure 4		
Peptide Mapping	See Figure 5		

[0086] As can be seen from Table 2 the biochemical properties of rVWF purified by the different process variants are comparable.

[0087] The major band of the rVWF protein is very similar in all products whereas the extent of impurities is lower in sample # 1 (denominated as VWF#07 in figure 1) by both silver staining and western blot analysis for residual rFVIII. The banding pattern for rFVIII

is comparable between all batches which suggests that no degradation due to the pH 9.0 conditions occurred.

[0088] Low and high resolution agarose gel electrophoresis revealed the high similarity of the rVWF preparations. No differences in multimer composition by low resolution multimer analysis could be seen. Figure 2 shows the stained gel of the UNO S runs with MMV and REO virus spiked samples. Also the high resolution multimer analysis revealed the intact multimer pattern suggesting no damage to the rVWF multimers occurred due to the pH 9.0 conditions. Figure 3 shows the stained gel of the UNO S runs with MMV and REO virus spiked samples.

[0089] By the Isoquant assay no enhanced de-amidation could be detected due to the dwell time of rVWF at pH 9.0. Generally the molar percentage of deamidated rVWF is very low. Subjecting the purified preparations of rVWF obtained by the process variants to proteolytic digestion by either V8 protease in the native state (see Figure 4) or trypsin (see Figure 5) in the denatured state and separating the resulting peptides by RP-HPLC resulted in similar chromatograms for all samples.

[0090] Minor differences in the peak patterns are due to the presence of different amounts of impurities (mainly residual rVWF propeptide as can be seen in Figure 1) in the preparations which was confirmed by mass spectrometry or N-terminal sequence analysis.

### **Example 3**

[0091] Purification of rVWF by Cation Exchange chromatography at high pH with MMV spike.

[0092] A UNOsphere S resin packed into a column was activated with 1 CV of 2 M NaCl and equilibrated with 25 CV of an equilibration buffer (pH=9.0). Thereafter, a rVWF containing solution adjusted to a conductivity of 15 mS/cm and a pH of 9.0 and spiked with mouse minute virus (MMV) was loaded onto the column at a linear flow rate of about 10.0 cm/h. The column was then washed with 10 CV of equilibration buffer (pH=9.0) and the product was eluted with 3.5 CV of elution buffer (pH=7.5) at a linear flow rate of 65cm/h. The increased pH during the loading and wash phase significantly reduced the binding of the virus particles to the resin but retained full binding of the product VWF. As a result, most of the loaded virus particles were found in the non-binding (flow through) and wash fraction separated from the product that was recovered in the eluate pool at high yields. The results in

Table 3 show that by applying this procedure a virus removal capacity of 2 logs could be obtained with the non-enveloped model virus mouse minute virus (MMV).

[0093] The TCID50 assay was performed as follows. Briefly, serial 1/2 log dilutions of the samples were prepared in the appropriate tissue culture medium and 100 µl of each dilution were added to each of 8 wells of a microtiter plate seeded with the indicator cell line. The cells were then incubated for 7 days at 36°C ± 2°C before the cytopathic effect was evaluated by visual inspection of the cells under a microscope. Median tissue culture infectious doses (TCID50) were calculated according to the Poisson distribution and expressed as log<sub>10</sub>[TCID50/ml].

**Table 3: Purification of rVWF on UNOsphere S**

	<b>Volume</b>	<b>Virus titer (TCID50)</b>	<b>Virus content (TCID50)</b>	<b>Reduction</b>
	ml	Log <sub>10</sub> /ml	Log <sub>10</sub>	Log <sub>10</sub>
Load	400	5.26	7.86	-
Eluate pool	89.4	3.8	5.75	2.11

[0094] The purification was performed using a column with 15 mm diameter and a bed height of 14 cm. Data shown are virus titers of active mouse minute virus.

#### **Example 4**

[0095] Purification of rVWF by Cation Exchange chromatography at high pH with Reo Type 3 virus spike.

[0096] A UNOsphere S resin packed into a column was activated with 1 CV of 2 M NaCl and equilibrated with 25 CV of an equilibration buffer (pH=9.0). Thereafter, a rVWF containing solution adjusted to a conductivity of 15 mS/cm and a pH of 9.0 and spiked with various non-enveloped viruses was loaded onto the column at a linear flow rate of about 100 cm/h. The column was then washed with 10 CV of equilibration buffer (pH=9.0) and the product was eluted with 3.5 CV of elution buffer (pH=7.5) at a linear flow rate of 65 cm/h. The increased pH during the loading and wash phase significantly reduced the binding of the virus particles to the resin but retained full binding of the product VWF. As a result, most of the loaded virus particles were found in the non-binding (flow through) and wash fraction separated from the product that was recovered in the eluate pool at high yields. The results in Table 4 show that by applying this procedure a virus removal capacity of 2 logs could be obtained with the non-enveloped model virus mouse Reo Virus Type 3 (REO-III).

**Table 4: Purification of rVWF on UNOsphere S**

	<b>Volume</b>	<b>Virus titer (TCID50)</b>	<b>Virus content (TCID50)</b>	<b>Reduction</b>
	ml	Log <sub>10</sub> /ml	Log <sub>10</sub>	Log <sub>10</sub>
Load	897	3.73	6.68	-
Eluate pool	89.3	2.61	4.56	2.12

[0097] The purification was performed using a column with 15 mm diameter and a bed height of 14 cm. Data shown are virus titers of active mouse Reovirus Type III (REO-III).

#### **Example 5**

[0098] Purification of rVWF on UNOsphere S according the standard procedure (neutral pH).

[0099] A UNOsphere S resin packed into a column was activated with 1 CV of 2 M NaCl and equilibrated with 25 CV of an equilibration buffer (pH=6.5). Thereafter, a rVWF containing solution adjusted to a conductivity of 15 mS/cm and a pH of 6.5 and spiked with various non-enveloped viruses was loaded onto the column at a linear flow rate of about 100 cm/h. The column was then washed with 10 CV of equilibration buffer (pH=6.5) and the product was eluted with 3.5 CV of elution buffer (pH=7.5) at a linear flow rate of 65 cm/h. The virus titer of the various viruses tested were evaluated in the different chromatographic fractions (load, column flow through, wash, eluate, post eluate) and the reduction factors were calculated. The results in Table 5 show that by applying the standard purification procedure for VWF on UNOsphere S the removal capacity for non-enveloped viruses was insufficient for the different model viruses tested to claim a robust chromatographic step for removal of non-lipid enveloped viruses.

**Table 5: VWF purification according to the standard procedure and the corresponding reduction capacities for non-enveloped viruses.**

	<b>Reduction</b>	<b>Comment</b>
	Log <sub>10</sub>	Virus Characteristics
PPV (porcine Parvovirus)	<1	small, DNA virus
hAdV (human Adenovirus)	1.8	large, DNA virus
EMCV (Encephalo Myocarditis Virus)	<1	small, RNA
Reo Virus Type III	1.8	large, RNA
MMV (mouse Minute Virus)	<1	small, DNA

[0100] The reduction rate is calculated as the total virus load in the load fraction divided by the total virus load in the eluate fraction expressed in logarithmic values.

What is claimed is:

1. A method for removing a non-lipid enveloped virus from a protein-containing solution comprising:

applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein; and

washing the cation exchange resin with a first wash buffer to form an eluate, said first wash buffer having a pH that is equal to or lower than the solution applied to the cation exchange resin.

2. The method of claim 1 wherein the solution applied to the cation exchange resin is at least 1 pH unit above the isoelectric point of the protein.

3. A method for removing a non-lipid enveloped virus from a protein-containing solution comprising:

applying the solution to a cation exchange resin,

washing the cation exchange resin with a first wash buffer at a pH higher than the pH of the solution applied to the cation exchange resin; and

washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer.

4. The method of claim 3 wherein the pH of the first wash buffer is at least 1 pH unit above the isoelectric point of the protein applied to the cation exchange resin.

5. A method for removing a non-lipid enveloped virus from a protein-containing solution comprising:

applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein; and

washing the cation exchange resin with a first wash buffer at a pH higher than the isoelectric point of the protein applied to the cation exchange resin; and

washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer.

6. The method of claim 5 wherein the solution applied to the cation exchange resin is at least 1 pH unit above the isoelectric point of the protein.
7. The method of claim 5 wherein the pH of the first wash buffer is at least 1 pH unit above the pH of the solution applied to the cation exchange resin.
8. The method of claims 2, 4, 6 or 7 wherein the pH is greater than 7.0.
9. The method of any of claims 1 - 8 wherein the protein in the solution is a polypeptide having a molecular mass of at least about 150 kilodaltons.
10. The method of any of claims 1 - 9 wherein the cation exchange resin has a negatively charged group selected from the group consisting of carboxymethyl (CM), sulphoalkyl (SP, SE), sulfated esters of cellulose, heparin and methylsulfonate (S).
11. The method of any of claims 1-10 wherein the protein is a blood coagulation protein.
12. The method of claim 11 wherein the blood coagulation protein is selected from the group consisting of Factor VIII and von Willebrand factor.
13. A method for removing a non-lipid enveloped virus from a von Willebrand (VWF)-containing solution comprising:
  - applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein; and
  - washing the cation exchange resin with a first wash buffer to form an eluate, said first wash buffer having a pH that is equal to or lower than the solution applied to the cation exchange resin.
14. A method for removing a non-lipid enveloped virus from a VWF-containing solution comprising:
  - applying the solution to a cation exchange resin,
  - washing the cation exchange resin with a first wash buffer at a pH higher than the pH of the solution applied to the cation exchange resin; and
  - washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer.


15. A method for removing a non-lipid enveloped virus from a VWF-containing solution comprising:

applying the solution to a cation exchange resin at a pH higher than the isoelectric point of the protein; and


washing the cation exchange resin with a first wash buffer at a pH higher than the isoelectric point of the protein applied to the cation exchange resin; and

washing the cation exchange resin with a second wash buffer to form an eluate, said first eluant having a pH that is equal to or lower than the first wash buffer.






A) Silver stain





B) Western blot for residual rFVIII

**FIG. 1**




**FIG. 2**





**FIG. 4**



**FIG. 5**

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2010/046180

**A. CLASSIFICATION OF SUBJECT MATTER**  
INV. C07K14/755 A61K38/36  
ADD .

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal , WPI Data, BIOSIS, EMBASE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	<p>CAMERON R ET AL: "The removal of model viruses, poliovirus type 1 and canine parvovirus, during the purification of human albumin using ion-exchange chromatographic procedures.", BIOLOGICALS : JOURNAL OF THE INTERNATIONAL ASSOCIATION OF BIOLOGICAL STANDARDIZATION DEC 1997 LNKD- PUBMED :9467035, vol. 25, no. 4 , December 1997 (1997-12), pages 391-401, XP002610211, ISSN: 1045-1056, DOI: 10.1006/biol.1997.0115 the whole document * abstract page 392, right-hand column , paragraph 1 "Discussion"; page 399 - page 400</p> <p style="text-align: center;">----- -/--</p>	1-12

Further documents are listed in the continuation of Box C

See patent family annex

\* Special categories of cited documents

- <sup>1</sup>A<sup>1</sup> document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- <sup>1</sup>L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- <sup>1</sup>P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

18 November 2010

Date of mailing of the international search report

02/12/2010

Name and mailing address of the ISA/

European Patent Office, P B 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040,  
Fax (+31-70) 340-3016

Authorized officer

Gal I , Anne-Laure

## INTERNATIONAL SEARCH REPORT

International application No

PCT/US2010/046180

C(Conhnuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	Chapter 9 : "Ion Exchange Chromatography" In: Bollag DM, Rozycki MD, Edelstein SJ: "Protein Methods (2nd edition)", 1996, wiley-Liss, Inc., New York, NY, XP002609616, page 231 - page 239 -----	1-12
A	US 6 465 624 B1 (FISCHER BERNHARD [AT] ET AL) 15 October 2002 (2002-10-15) the whole document * abstract column 3 , line 45 - line 54 column 4 , line 59 - line 62 column 6 , line 11 - line 22 column 6 , line 45 - line 52 column 11 , line 14 - column 12 , line 32; example 3 claims 1, 8 claim 13 -----	13-15
A	CHANDRA SUDHISH ET AL: "Effectiveness of alternative treatments for reducing potential viral contaminants from plasma-derived products.", THROMBOSIS RESEARCH 1 MAR 2002 LNKD- PUBMED: 12062540, vol. 105, no. 5, 1 March 2002 (2002-03-01) , pages 391-400, XP002610212, ISSN: 0049-3848, DOI: 10.1016/S0049-3848(02)00044-0 the whole document * abstract section 3.2. "Purification steps that reduce viral contamination"; page 393 -----	13-15
A	SOLHEIM ET AL: "Pathogen reduction of blood components", TRANSFUSION AND APHERESIS SCIENCE, ELSEVIER SCIENCE, LONDON, GB, vol. 39, no. 1, 1 August 2008 (2008-08-01) , pages 75-82, XP023613846, ISSN: 1473-0502, DOI: DOI:10.1016/J.TRANSFI.2008.05.003 [retrieved on 2008-07-03] the whole document * abstract ----- -/--	13-15

INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2010/046180

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	<p>PROWSE ET AL: "Properties of Pathogen-Inactivated Plasma Components", TRANSFUSION MEDICINE REVIEWS, GRUNE AND STRATTON, ORLANDO, FL, US, vol. 23, no. 2, 1 April 2009 (2009-04-01), pages 124-133, XP026017877, ISSN: 0887-7963, DOI: DOI: 10.1016/J.TMRV.2008.12.004 [retrieved on 2009-03-20] the whole document * abstract table 1 table 4</p> <p style="text-align: center;">-----</p>	13-15



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2010/046180

Patent document cited in search report	Publication date	Publication date	Patent family member(s)	Publication date
US 6465624	B1	15-10-2002	AT 405403 B	25-08--1999
			WO 9838219 A1	03-09--1998
			AT 359301 T	15-05--2007
			AU 737986 B2	06-09--2001
			AU 5847098 A	18-09--1998
			CA 2282843 A1	03-09--1998
			DK 1005492 T3	13-08--2007
			EP 1005492 A1	07-06--2000
			ES 2285754 T3	16-11--2007
			JP 4250770 B2	08-04--2009
			JP 2001513088 T	28-08--2001
			NO 994138 A	26-08--1999

---