



US00PP21205P3

(12) **United States Plant Patent**
Blom

(10) **Patent No.:** **US PP21,205 P3**

(45) **Date of Patent:** **Aug. 17, 2010**

(54) **CHRYSANTHEMUM PLANT NAMED**
'ZANMUSPEN'

(50) Latin Name: *Chrysanthemum* × *morifolium* Ramat.
Varietal Denomination: **Zanmuspen**

(75) Inventor: **Wilhelmus Bernardus Blom,**
Leimuïden (NL)

(73) Assignee: **Chrysanthemum Breeders Association**
Research B.V.

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/314,961**

(22) Filed: **Dec. 19, 2008**

(65) **Prior Publication Data**

US 2010/0162452 P1 Jun. 24, 2010

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./288**

(58) **Field of Classification Search** Plt./288
See application file for complete search history.

Primary Examiner—Kent L Bell

(74) *Attorney, Agent, or Firm*—Steptoë & Johnson LLP

(57) **ABSTRACT**

A *chrysanthemum* plant named 'Zanmuspen' characterized
by its medium sized blooms with white ray florets with a
cream center and prolific branching; natural season flower
date August 25–30 (week 35); blooming for a period of 5
weeks.

3 Drawing Sheets

1

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *chrysanthemum* plant, botanically known as *Chrysanthemum*
×morifolium Ramat., and hereinafter referred to by the
cultivar denomination 'Zanmuspen'. 'Zanmuspen' is a product
of a breeding and selection program for outdoor pot mums
(garden mums) which had the objective of creating new *chry-*
santhemum cultivars with a double type inflorescence, a natural
season flower date starting at August 25–30; blooming for
a period of 5 weeks. 'Zanmuspen' is a seedling resulting from
the crossing of the female parent id 2862 and male parent id
3052. Plants of 'Zanmuspen' differ from plants of the female
parent in the color of ray-florets: yellow in the female parent
and white in the seedling. Plants of 'Zanmuspen' differ from
plants of the male parent in the natural season blooming
period; those of the seedling flower 1–2 weeks earlier.

The new and distinct cultivar was discovered and selected
as a flowering plant by Wilhelmus Bernardus Blom on a
cultivated field in Rijsenhout, The Netherlands in 2005. The
first act of asexual production of 'Zanmuspen' was accom-
plished when vegetative cuttings were used from the initial
selection in 2005 and propagated further in a controlled envi-
ronment in Rijsenhout, The Netherlands. The new cultivar
has been found to retain its distinctive characteristics through
successive propagations.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention of a new and distinct variety of
chrysanthemum is shown in the accompanying drawings, the
color being as nearly true as possible with color photographs
of this type.

FIG. 1 shows a plant of the cultivar in full bloom.

FIG. 2 shows the various stages of bloom of the new cul-
tivar.

2

FIG. 3 shows the various stages of foliage of the new
cultivar.

DESCRIPTION OF THE INVENTION

The observations and measurements were gathered from
plants grown out door in Rijsenhout, The Netherlands under
natural day length and temperature and planted in week 22 in
2008. The natural blooming date of this crop was August
25–30 (week 35). The average height of the plants was 25 cm.
No growth retardants were used. No tests were done on dis-
ease or insect resistance or susceptibility. No tests were done
on cold or drought tolerance. This new variety produces
medium sized blooms with white ray florets and a cream
center blooming for a period of 5 weeks.

From the cultivars known to inventor the most similar
existing cultivar in comparison to 'Zanmuspen' is 'Zanmus-
tarbu' (U.S. Plant Pat. No. 20,473). When 'Zanmustarbu' and
'Zanmuspen' are being compared the following differences
are noticed: The differences of 'Zanmustarbu' and 'Zanmus-
pen' are (1) Natural season blooming date. (2) Variance in
ray-floret length. And (3) Variance in ray-floret width. (1)
Under natural conditions, plants of 'Zanmustarbu' flower one
week earlier than those of 'Zanmuspen'. (2) There is a larger
variation in ray-floret length in 'Zanmustarbu' than in 'Zan-
muspen'. (3) There is a larger variation in ray-floret width in
'Zanmustarbu' than in 'Zanmuspen'.

The following is a description of the plant and character-
istics that distinguish 'Zanmuspen' as a new and distinct
variety.

The color designations are taken from the plant itself.
Accordingly, any discrepancies between the color designa-
tions and the colors depicted in the photographs are due to
photographic tolerances. The color chart used in this descrip-
tion is: The Royal Horticultural Society Colour Chart, edition
2001.

TABLE 1

Botanical Description of <i>chrysanthemum</i> plant <i>Chrysanthemum xmorifolium</i> Ramat 'Zanmuspen'	
Bud	
Size	Small; cross-section 0.4 cm, height 0.2 cm
Shape	Oblate
Texture	Pubescent
Outside Color	Greyed-green 191A
Phyllaries	
Number	30, arranged in 3 rows
Shape	Elliptic
Apex	Acute
Base	Truncate
Margin	Entire
Color	Upper side Greyed-green 191A Under side Greyed-green 191C
Length and width	4 mm; 1-2 mm
Texture	Pubescent
Inflorescence	
Type	Double
Height	1.8 cm
Diameter	6 cm
Peduncle length	7 cm
Peduncle color	Greyed-green 191 B to 191C
Peduncle diameter	1.5 mm
Peduncle surface	Pubescent
Number per branch	Approx. 10 inflorescences
Flowering period individual inflorescence	Ca. 4 weeks
Seeds	
	Produced in small quantities, ovate, Greyed-brown 199A, length 1.5 mm, diameter 0.8 mm
Fragrance	
Color	Faint <i>chrysanthemum</i> odor
Center of inflorescence (ray-florets)	
Color of upper surface of mature ray-florets	Immature stage: Yellow 10A Mature stage: Yellow 10A White 155A
Color of the lower surface of mature ray-florets	White 155D
Tonality from Distance	A garden mum with white inflorescences and a cream center
Color of the ray-florets after aging of the plant	White 155A with sometimes at edges Gey-Brown 199D
Ray florets	
Texture	Upper and lower side smooth
Number	180
Shape	Oblanceolate
Apex	Rounded
Base	Acute
Cross-section	Convex
Longitudinal axis of majonty	Straight
Length of corolla tube	5 mm
Ray-floret margin	Entire
Ray-floret length	2-3 cm
Ray-floret width	3-6 mm
Ratio length/width	High
Disc florets	Absent
Receptacle	
Color	Green 138D
Shape	Conical raised
Height	0.5 cm
Diameter	0.5 cm

TABLE 1-continued

Botanical Description of <i>chrysanthemum</i> plant <i>Chrysanthemum xmorifolium</i> Ramat 'Zanmuspen'	
Reproductive Organs	
Androecium	Absent
Gynoecium	Present in ray florets
Style colour	Yellow-green 154C
Style Length	3 mm
Stigma colour	Yellow 3A
Stigma Width	1 mm
Ovary	Enclosed in calyx
Plant	
Form	Grown as a potmum, outdoor raised and mounded
Growth habit	Spherical shape
Growth rate	Medium
Height	25 cm
Width	40 cm
Stem Color	Greyed-brown 199A
Stem Strength	Strong
Stem Brittleness	Not brittle
Stem Anthocyanin Coloration	Not observed
Internode length	1-2 cm
Length of lateral branch	From top to bottom 18 cm
Lateral branch color	Green 137C
Lateral branch brittleness	Medium
Lateral branch diameter	2 mm
Branching (average number of lateral branches)	Prolific with 8 breaks after pinching
Natural season blooming date	August 25-30 to September 22-27
Foliage	
Leaf color	Upper side: Green N138B Lower side: Green N138C
Color midvein	Upper side: Yellow-green 147D Lower side: Yellow-green 148D
Size	Small; length 4-6 cm, width 3.5-4 cm
Quantity (number per lateral branch)	24-26
Shape	Elliptic
Texture upper side	Sparsely pubescent
Texture under side	Pubescent
Venation arrangement	Palmate
Shape of the margin	Serrated
Shape of Base of Sinus Between Lateral Lobes	Rounded
Margin of Sinus Between Lateral Lobes	Converging
Shape of Base	Truncate
Apex	Mucronulate
Petiole length	0.3-1 cm
Petiole diameter	2-3 mm
Petiole color	Yellow-green 147D

TABLE 2

Differences with the comparison variety			
		'Zanmuspen'	'Zanmustarbu'
55	Natural season blooming date	August 25-30 (week 35)	August 18-23 (week 34)
	Variance in ray-floret length	2-3 cm	2-3.5 cm
	Variance in ray-floret width	3-6 mm	2-7 mm

60 I claim:
1. A new and distinct variety of *chrysanthemum* plant as described and illustrated.




FIG. 1




FIG. 2




FIG. 3