



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2673366 C 2017/12/19

(11)(21) **2 673 366**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) **Date de dépôt PCT/PCT Filing Date:** 2007/12/21
(87) **Date publication PCT/PCT Publication Date:** 2008/07/03
(45) **Date de délivrance/Issue Date:** 2017/12/19
(85) **Entrée phase nationale/National Entry:** 2009/06/19
(86) **N° demande PCT/PCT Application No.:** SE 2007/051080
(87) **N° publication PCT/PCT Publication No.:** 2008/079095
(30) **Priorités/Priorities:** 2006/12/22 (US60/876,958);
2006/12/22 (SE0602804-7)

(51) **Cl.Int./Int.Cl. A61K 39/35** (2006.01),
A61P 37/00 (2006.01), **G01N 33/531** (2006.01)
(72) **Inventeurs/Inventors:**
MATTSSON, LARS, SE;
EVERBERG, HENRIK, SE;
LIDHOLM, JONAS, SE
(73) **Propriétaire/Owner:**
PHADIA AB, SE
(74) **Agent:** RIDOUT & MAYBEE LLP

(54) **Titre : NOUVEAUX ALLERGENES DE LA KALLIKREINE DE LA PROSTATE**
(54) **Title: NOVEL PROSTATE KALLIKREIN ALLERGEN**

(57) **Abrégé/Abstract:**

Prostatic kallikrein for the manufacture of a diagnostic or pharmaceutical composition for diagnosis/treatment of type 1 allergy, especially allergy to dogs.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 July 2008 (03.07.2008)

PCT

(10) International Publication Number
WO 2008/079095 A1

(51) International Patent Classification:

A61K 39/35 (2006.01) *A61P 37/00* (2006.01)
G01N 33/531 (2006.01)

(21) International Application Number:

PCT/SE2007/051080

(22) International Filing Date:

21 December 2007 (21.12.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0602804-7 22 December 2006 (22.12.2006) SE
60/876,958 22 December 2006 (22.12.2006) US

(71) Applicant (for all designated States except US): **PHADIA AB** [SE/SE]; Box 1344, S-751 43 Uppsala (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MATTSSON, Lars** [SE/SE]; Handarbetsvägen 24, S-757 57 Uppsala (SE). **EVERBERG, Henrik** [SE/SE]; Tulegatan 23B, S-113 53 Stockholm (SE). **LIDHOLM, Jonas** [SE/SE]; Björknäsvägen 9, S-741 42 Knivsta (SE).

(74) Agent: **BRANN AB**; P.O. Box 17192, S-104 62 Stockholm (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau
- with information concerning refusal to authorize the rectification of an obvious mistake under Rule 91.1

(54) Title: NOVEL PROSTATE KALLIKREIN ALLERGEN

(57) Abstract: Prostatic kallikrein for the manufacture of a diagnostic or pharmaceutical composition for diagnosis/treatment of type 1 allergy, especially allergy to dogs.

WO 2008/079095 A1

Novel prostate Kallikrein allergen

Field of the invention

The present invention relates to the field of allergy. More specifically, the
5 invention relates to the identification of novel allergens from mammals and to
diagnosis and treatment of allergy towards mammals.

Background

Dog dander is a common cause of indoor allergy with symptoms including
10 rhinitis, conjunctivitis, bronchial inflammation and asthma. Dog allergens can
be detected not only in houses where dogs are kept as pets but also in other
places such as schools and day care centres where dogs are not present on a
regular basis (1).

Allergy to dog is accompanied and dependent of sensitization to proteins
15 released from dog hairs and dander. In cases of suspected allergy to dog, the
clinical investigation includes assessment of sensitization by skin prick or
specific IgE antibody measurement using extract of dog hair and/or dander. A
laboratory immunoassay for specific IgE, such as a Phadia ImmunoCAP, can
detect most cases of sensitization to dog using natural dog dander extract due
20 to favourable assay conditions and a large solid phase available for allergen
attachment.

Dog hair and dander extracts contain a complexity of allergenic and non-
allergenic proteins (2, 3). Three dog allergens have so far been identified and
studied in detail: Can f 1, Can f 2 and Can f 3. Can f 1, a member of the
25 lipocalin protein family, with reported molecular weight of 21-25 kD, was first
purified by de Groot et al. (4) and later cloned and expressed as a recombinant
protein (5). Can f 2 belongs to the same protein family but is a protein distinct
from Can f 1 (4, 5). Can f 3, dog serum albumin, is a relatively conserved protein
demonstrating extensive cross-reactivity to other mammalian albumins (6).

Of the known dog allergens, Can f 1 is the most important, binding IgE antibodies from approximately half of dog allergic subjects (7). About 20% of dog allergic subjects display IgE binding to Can f 2 but most of these are also sensitized to Can f 1. Although 30-40% of adult dog allergic individuals may show IgE binding to Can f 3 (2, 8), the specific clinical relevance of mammalian serum albumins is uncertain.

It has been known for a long time that major allergens relevant to allergy to rodents, such as mice and rats, are present in the animals' urine and these have been isolated and extensively characterized (9-13). IgE antibody binding activity has also been reported to exist in urine of other animals, including cats and dogs (14), but no allergen has been purified from urine of these animals and characterized at a molecular level.

Summary of the invention

As stated above, a laboratory immunoassay for specific IgE can detect most cases of sensitization to dog using natural dog dander extract due to favourable assay conditions and a large solid phase available for allergen attachment. However, in a miniaturized or non-laboratory immunoassay, such as an allergen microarray or a doctor's office test, the combination of less favourable assay conditions, lower capacity for antibody-binding allergen reagent and natural allergen extract of limited potency, has been found to cause insufficient diagnostic sensitivity. A similar situation may exist also for immunoassays for specific IgE to other animal epithelia. Thus, there is a need in some cases to use pure allergenic proteins to achieve sufficient sensitivity in diagnostic tests for specific IgE.

Furthermore, a significant proportion of dog allergic individuals also do not react to any of the known identified dog allergens and this was recently demonstrated in a Finnish population (7).

The above led the present inventors to look for additional, not yet identified, dog allergens. Such novel allergens may be useful not only as reagents for increased sensitivity in routine diagnostic tests, but also as a complement to known dog

allergens in different types of component-resolved diagnostic applications (15, 16). Pure allergenic proteins, or fragments and variants thereof with improved non-anaphylactic properties, may also be used in component-resolved immunotherapy (16-20).

5 A new major allergen has thus been purified from dog urine and identified as prostatic kallikrein. It is in all aspects distinct from previously known dog allergens. Further, a similar or identical and immunologically equivalent allergen has been found to exist in dog dander extract. Kallikrein represents an important addition to the panel of known dog allergens and will be useful in the
10 diagnosis of dog allergy. It is also anticipated that homologous proteins from other mammals, such as cat, horse and rodents, including rat and mouse, will have similar allergenic properties and diagnostic utility.

Prostatic kallikrein was found to exist not only in urine but also in the fur of dander of dogs. However, the fact that protein specifically expressed in prostate
15 tissue would be restricted to male individuals, suggests that female dogs would lack this allergen. Preliminary results in our laboratory indeed support this notion and, if corroborated by results from more extensive studies, the implication would be that dog allergic individuals sensitized exclusively to prostatic kallikrein may tolerate female dogs.

20 In a recently published report, it was demonstrated that vaginal hypersensitivity reaction to ejaculate was associated with IgE sensitization to human prostate-specific antigen, PSA, present in seminal plasma (21). As canine and human prostatic kallikrein and human prostate-specific antigen have partial sequence similarity, it is possible that sensitization to canine prostate-specific kallikrein
25 confers an elevated risk of developing such allergic reactions. It can also be envisaged that IgE-mediated immune reactions to prostate-specific kallikrein may play a role in certain cases of infertility in humans.

In one aspect the invention relates to the use of kallikrein in diagnosis of Type I allergy and the use of kallikrein for the manufacture of a composition for
30 diagnosis of Type I allergy.

In a further aspect the invention relates to an allergen composition "spiked" with kallikrein. Such an allergen composition may be an allergen extract or a

mixture of purified or recombinant allergen components having no or a low kallikrein content, wherein the kallikrein is added in order to bind IgE from patients whose IgE would not bind or bind poorly to the other allergen components in the composition. This aspect of the invention also relates to a method for producing such a composition, which method comprises the step of adding kallikrein to an allergen composition, such as an allergen extract (optionally spiked with other components) or a mixture of purified native or recombinant allergen components.

In yet a further aspect the invention relates to an *in vitro* diagnostic method for diagnosing a Type I allergy in a patient, wherein a body fluid sample such as a blood or serum sample from the patient is brought into contact with kallikrein or a composition according to the previous aspect, and it is detected whether or not the patient sample contain IgE antibodies that bind specifically to kallikrein. Such a diagnostic method may be carried out in any manner known in the art.

The kallikrein may e.g. be immobilized on a solid support, such as in a conventional laboratory immunoassay, in a microarray or in a lateral flow assay.

In a further aspect the invention relates to a diagnostic kit for performing the method according to the previous aspect, which kit includes kallikrein.

In the above mentioned aspects, the wildtype kallikrein molecule may be replaced with fragments or variants of kallikrein, natural or man-made, sharing epitopes for antibodies with wildtype kallikrein, as defined below.

The invention further relates to a method of treatment of Type I allergy comprising administering to a patient in need of such treatment a kallikrein or a modified kallikrein, as explained below. This aspect of the invention also relates to the use of kallikrein in such immunotherapy, including e.g. component-resolved immunotherapy (16). In one embodiment of this aspect, the kallikrein may be used in its natural form or in a recombinant form displaying biochemical and immunological properties similar to those of the natural molecule. In another embodiment, kallikrein may be used in a modified form, generated chemically or genetically, in order to abrogate or attenuate its IgE antibody binding capacity, while preferably being capable of eliciting an IgG response in a treated individual. Examples of modifications include, but are not

limited to, fragmentation, truncation or tandemization of the molecule, deletion of internal segment(s), substitution of amino acid residue(s), domain rearrangement, or disruption at least in part of the tertiary structure by disruption of disulfide bridges or its binding to another macromolecular structure, or by removal of the protein's ability to bind calcium ions or other low molecular weight compounds. In yet another embodiment of this aspect, the individual 10 kDa and/or the 18 kDa subunits of kallikrein, which display reduced IgE binding activity as compared to the intact molecule, are used as modified kallikrein.

- 10 In all of the above mentioned aspects of the invention, the kallikrein can be derived from any mammal producing kallikrein capable of inducing an allergic response in a patient. The kallikrein may be purified from its natural source, such as from urine, saliva or other body fluids, or from tissue, such as hair or dander, of the mammal in question. It may also be produced by recombinant
- 15 DNA technology or chemically synthesized by methods known to a person skilled in the art.

The invention also relates to canine prostatic kallikrein for use in diagnosis and therapy, such as diagnosis and therapy of Type I allergy to dog.

- 20 The invention also relates to a method for purification of kallikrein from mammalian urine, comprising the steps
- filtering the mammalian urine;
 - buffer exchange with a buffer suitable for hydrophobic interaction chromatography;
 - 25 -filtering of the buffer exchanged urine sample;
 - applying the buffer exchanged urine sample to a hydrophobic interaction chromatography column; and
 - collecting the flow-through fraction comprising kallikrein.

The mammalian urine may be canine urine.

Definitions

Kallikreins are proteolytic enzymes from the serine endopeptidase family found in normal blood and urine. In the IUBMB enzyme nomenclature system, plasma kallikrein has been assigned number EC 3.4.21.34 and tissue kallikrein

5 number EC 3.4.21.35. Urinary kallikrein from dog is a 28 kDa heterodimeric protein comprising two subunits of approximately 10 ± 2 and 18 ± 2 kDa, respectively, for the purposes of this invention referred to as the 10 and 18 kDa subunits, respectively. It has an amino acid sequence according to SEQ ID NO: 1, GenBank Accession no: P09582, and homologous proteins have been
10 described in a wide range of mammalian species, including, horse, cow, pig, mouse, rat and primates (e.g. Accession no AAQ23713-4 (horse), NP_001008416 (cow), P00752 (pig), P00755-6 and P15947 (mouse), P36373 and P00758 (rat), Q28773 (baboon), XP_001174026 (chimpanzee), Q07276 (macaque), P20151, Q07276 and AAM11874 (human).

15 Variants and fragments of a kallikrein should be construed as meaning proteins or peptides with a length of at least 10 amino acids, more preferably at least 50, even more preferably at least 75 or 100 amino acid residues, and a sequence identity to said kallikrein of at least 50 %, preferably over 60 %, 70 %, 80 %, 90 % or 95 %.

20 A modified kallikrein should in the context of the present invention be construed as meaning a kallikrein that has been chemically or genetically modified to change its immunological properties, e.g. as exemplified above in relation to the immunotherapy aspect of the invention.

Variants and fragments of kallikrein sharing epitopes for antibodies with
25 wildtype kallikrein should be construed as being those fragments and variants whose binding of IgE antibodies from a serum sample from a representative kallikrein sensitized patient can be significantly inhibited by kallikrein. Such an inhibition assay may e.g. be performed according to the protocol disclosed in Example 8.

30 A hypoallergenic modified kallikrein or variant or fragment of kallikrein should be construed as being a modified kallikrein or variant or fragment of kallikrein that is not capable of binding kallikrein reactive IgE antibodies from a serum

sample of a representative kallikrein sensitized patient, as determined e.g. by the protocol according to Example 3 or which displays no or significantly reduced biological allergen activity, as determined by a cellular activation assay such as the basophil histamine release assay (22, 23).

Brief description of drawings

Figure 1 shows the fractionation of dog urinary proteins by size exclusion chromatography. Fractions comprising each of the three peaks indicated in the figure (labeled 1-3) were pooled as indicated for analysis of IgE binding activity.

5 Figure 2 shows the purification of an IgE binding protein from peak 2 of Figure 1 by reversed phase chromatography. The peak containing the protein selected for further investigation is indicated by an arrow.

Figure 3 is an SDS-PAGE analysis of reduced (red) and non-reduced (ox) samples of the IgE binding protein purified from dog urine by size exclusion and
10 reversed phase chromatography. Lane M contains molecular weight marker proteins.

Figure 4 shows the effect of kallikrein as a fluid-phase inhibitor on specific IgE binding to immobilized dog dander extract.

Figure 5a-b is an assessment by immunoblot analysis of IgE antibody reactivity
15 to dog dander extract in 37 dog allergic subjects' sera. Prior to incubation with the membrane strips, serum samples were diluted as indicated. Lane M contains molecular weight marker proteins.

Figure 6 shows a comparison of the immunoblot signal intensity of a 28 kDa band, corrected for serum dilution, and the level of kallikrein-specific IgE, as
20 determined by experimental ImmunoCAP analysis. The ImmunoCAP and immunoblot detection limits applied are indicated by hatched lines. Immunoblot signal intensity is expressed in arbitrary units (AU).

Figure 7 shows specific immunoblot inhibition of the 28 kDa protein band by purified dog urinary kallikrein. Lane M contains molecular weight marker
25 proteins.

Figure 8 shows the first step of purification of kallikrein from dog dander, by size exclusion chromatography. Six fractions (labeled 1-6) indicated in the figure were analysed for IgE binding activity.

Figure 9 shows the second step of purification of kallikrein from dog dander, by reversed phase chromatography. Top fractions of three peaks indicated in the figure (labeled 1-3) were analysed for IgE antibody binding activity.

Figure 10 shows a comparative immunoblot analysis specific IgE antibody
5 binding to dog dander extract, purified urinary kallikrein and partially purified kallikrein from dog dander. Two kallikrein-reactive sera (no. 6 and 8) and one kallikrein non-reactive serum (no. 11) were used. Both reduced and non-reduced forms of the allergen preparations were analysed, as indicated in the legend. Lane M contains molecular weight marker proteins.

10 Figure 11 shows SDS Page analysis of purified recombinant dog urinary kallikrein.

Figure 12 shows analytical gelfiltration analysis of purified recombinant dog urinary kallikrein.

Figure 13 shows a comparison of specific IgE antibody binding activity of
15 natural and recombinant dog urinary kallikrein.

Detailed description of the invention

The examples below illustrate the present invention with the isolation and use of kallikrein from dog. The examples are only illustrative and should not be considered as limiting the invention, which is defined by the scope of the appended claims.

EXAMPLE 1: Detection and isolation of an IgE binding protein from dog urine

In order to investigate whether dog urine may contain allergens relevant to dog allergy in humans, the following experiments were performed. Urine was collected from a 7 year old male crossbreed between Siberian Husky and Vorsteh. After filtration through a 0.45 µm mixed cellulose ester filter, 10 mL of urine was applied to a Superdex 75 size exclusion chromatography (SEC) column (XK26/100, V_t = 505 mL, GE Healthcare Biosciences, Uppsala, Sweden) equilibrated with 20 mM MOPS pH 7.6, 0.5 M NaCl (MBS) and elution was performed with the same buffer at a flow rate of 2 mL/min. Fractions from three peaks were pooled as indicated in the chromatogram shown in Figure 1 and analysed for allergen activity. The protein content of each fraction was immobilized on ImmunoCAP (Phadia, Uppsala, Sweden) solid phase and its IgE antibody binding activity tested using eight sera from dog dander sensitized individuals. Most of these sera were selected as having high IgE binding to dog dander extract but relatively low binding to rCan f 1, rCan f 2 and nCan f 3. Of the three peaks tested, peak 2 was found to contain by far the highest level of IgE binding activity (Table 1). Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) using the NuPAGE MES buffer system (10% NuPAGE gel, Invitrogen, Carlsbad, CA, USA) of a reduced sample of peak 2 revealed two dominant protein bands, with apparent molecular weights of approximately 10 and 18 kDa, respectively (not shown).

Further protein purification from the pool corresponding to peak 2 was performed using a Source 15 reversed phase chromatography (RPC) column (ST4.6/100, V_t = 1.66 mL, GE Healthcare Biosciences). After addition of trifluoroacetic acid (TFA) to a final concentration of 0.065%, the pool was applied to the

column, followed by washing with 9 column volumes of 0.065% TFA in water. Elution was performed with a 0-45% linear gradient of acetonitrile in water containing 0.05% TFA, resulting in one distinctive but somewhat asymmetrical peak (Figure 2, peak indicated by an arrow). SDS-PAGE of reduced samples of fractions containing this peak revealed the presence of both the 10 and 18 kDa bands, seemingly unseparable (Figure 3). The fractions covering the entire peak were therefore pooled as indicated by horizontal bars in Figure 2. SDS-PAGE of a non-reduced sample of this pool revealed an additional band of 28 kD and a slight shift in mobility of the 10 kDa and 18 kDa bands (Figure 3). A faint protein band of approximately 55 kDa can also be seen in the non-reduced state, which may be a dimer of the 28 kDa protein. The occurrence of the 28 kDa band in the non-reduced state suggested that this protein may be made up of the 10 and 18 kDa polypeptides, joined together by one or more cysteine bridge(s). The fact that linear mass spectrometric analysis (data not shown) later showed dissociation of the 28 kDa component upon reduction and alkylation added further evidence to this notion.

EXAMPLE 2: Identification of the IgE binding protein from dog urine as prostate kallikrein

Mass spectrometry and N-terminal sequencing was used to determine the identity of the IgE binding protein isolated from dog urine.

Peptide mass fingerprint analysis by MALDI-TOF

For in-solution digestion of the RPC purified urinary protein, reduction and alkylation was performed by sequentially adding to the sample DTT and iodoacetamide at approximately 45- and 100-fold molar excess, respectively. Trypsin digestion was then performed overnight at 37°C, using porcine trypsin (Trypsin Gold, mass spectrometry grade, Promega, Madison, WI, USA). Samples containing digested peptides were spotted onto the MALDI target plate and α -cyano matrix in 50% acetonitrile, 10 mM $\text{NH}_4[\text{H}_2\text{PO}_4]$, 0.1 % TFA, was added. Following evaporation of the solvent, peptide mass fingerprinting (PMF) was performed in a Bruker Daltonics Autoflex 2 instrument (Bruker Daltonics, Bremen, Germany). To identify proteins matching PMF results obtained, the

MSDB database was searched using a Mascot server (Matrixscience, London, UK). Post source decay (PSD) analysis was performed on selected peptides. External calibration was performed using a peptide calibration standard (Bruker Daltonics).

5 In-gel digestion analysis of the individual protein bands from SDS-PAGE was performed essentially according to Shevchenko (24). In summary, the 10, 18 and 28 kDa bands described in Example 1 above were excised from a Coomassie brilliant blue stained SDS-PAGE gel. The gel pieces were sequentially washed with 50 mM ammonium bicarbonate containing 50 % acetonitrile
10 followed by shrinking in pure acetonitrile. After rehydration of the gel piece with 50 mM ammonium bicarbonate, acetonitrile was added to 50% and following a second acetonitrile wash step, the gel pieces were dried in a vacuum centrifuge. Reduction and alkylation was performed in sequence using 45 mM DTT and 100 mM iodoacetamide in 50 mM ammonium bicarbonate. After repeated washes
15 with 50 % acetonitrile in 50 mM ammonium bicarbonate and a final 100% acetonitrile wash, the gel particles were again dried down in a vacuum centrifuge. Trypsin digestion was performed overnight at 37°C using porcine trypsin as described above. The digested sample was then sonicated and peptides extracted from the gel particles in 50% acetonitrile containing 0.1%
20 TFA. Sample preparation and peptide mass fingerprinting was performed as described above.

The PMF analysis of the in-solution digested urinary protein resulted in a highly significant match ($p < 0.05$) to prostatic kallikrein from dog (Accession no P09582). PSD analysis of two peptides, $m/z = 1224.6$ and $m/z = 1632.8$, which
25 were also present in the in-gel digestion analysis of the 18 kDa band, gave significant database matches to the amino acid sequences FMLCAGVLEGGK and SHDLMLLHLEEPK, corresponding to residues 194-204 and 117-130, respectively, of the same protein database entry.

Corroborating results were obtained from analysis of the in-gel digested protein
30 bands as PMF of the 28 kDa band also yielded a highly significant database match ($p < 0.05$) to kallikrein from dog (P09582). Further evidence to the identity of the isolated urinary protein came from the analysis of in-gel digested samples

of the 10 kDa band, PSD analysis of peptide $m/z=1004.6$ gave a highly significant ($p<0.05$) database match with the amino acid sequence SFIHPLYK, corresponding to residues 95-102 of P09582.

N-terminal amino acid sequencing

5 For N-terminal sequencing, the reduced 10 kDa and 18 kDa protein bands were excised separately from a SDS-PAGE gel and extracted in 6 M guanidinium-HCl, 20 mM Tris pH 8.0, 0.5 M NaCl, using a plastic rod for homogenization. N-terminal sequence analysis of the extracted 10 kDa and 18 kDa bands, performed using a Hewlett-Packard G1000A instrument (Hewlett-Packard, Palo Alto, CA), yielded the amino acid sequences IIGGREXLKN and AVIRPGEDRS, respectively, which were found to match residues 25-34 and 108-117 in the dog prostatic kallikrein precursor sequence of Accession no P09582.

Taken together, the results described in this example demonstrate that the major constituent of the purified dog urinary protein, corresponding to the 10 and 18 kDa bands in reducing SDS-PAGE analysis, is identical to prostatic kallikrein from dog. Further, the observations suggest that the 10 and 18 kDa polypeptides are formed by posttranslational cleavage of a primary gene product and are held together by disulfide bridges to form the 28 kDa protein seen under non-reducing conditions, similar to what has previously been described for human kallikrein (25).

Prostatic kallikrein is also known as arginine esterase and carries that designation in database entries describing identical or nearly identical amino acid sequences, including NP_001003284, CAA68720 and AAA30831. Further, we note that another variant of kallikrein, expressed in renal, pancreatic and salivary gland tissues, has been identified in dog (Accession No CAA53210) and shares 68% amino acid identity with prostatic kallikrein.

Example 3: Assessment of IgE binding activity of kallikrein, rCan f 1, rCan f 2 and nCan f 3

30 *In vitro* IgE binding activity of the purified recombinant and natural dog allergens were examined using ImmunoCAP® (Phadia, Uppsala, Sweden), an

immunoassay system used for specific IgE antibody measurement in clinical diagnosis of atopic allergy. Recombinant Can f 1 and Can f 2 (5) were cloned and expressed in *E. coli* essentially as described (26). Dog albumin was purified from serum using anion exchange chromatography and Blue Sepharose affinity chromatography, essentially as described (27). Experimental ImmunoCAP tests were prepared and used for serum analysis as described (26).

Sera from 37 dog allergic patients from Sweden (n=9), Spain (n=23) and North America (n=4) were used in the study. All patients had a positive skin prick test for dog dander extract and a doctors' diagnosis of dog allergy with symptoms of asthma, rhinoconjunctivitis and/or urticaria. All of the sera had a positive specific IgE test (ImmunoCAP) to dog dander extract.

The levels of specific IgE to dog dander extract, rCan f 1, rCan f 2, nCan f 3 and purified kallikrein are shown in Table 2 and a summary of the results is shown in Table 3. Of the tested sera, 29 showed IgE reactivity to kallikrein and 18 to rCan f 1. Both rCan f 2 and nCan f 3 appeared as minor allergens among the subjects studied, binding IgE from only 8 and 6 of 37, respectively. Fourteen of the 37 sera (38%) reacted only to kallikrein. On average among the kallikrein-reactive sera, the level of IgE binding to kallikrein amounted to 64% of the IgE binding to dog dander. The corresponding relative levels of IgE binding to rCan f 1, rCan f 2 and nCan f 3 were 45%, 25% and 47%, respectively, among sera specifically reactive to those allergens. Only two of the 37 sera tested lacked IgE reactivity to all of the four dog allergens. The IgE binding to kallikrein showed no correlation to any of the other dog allergens, demonstrating that the immune response to kallikrein is an independent variable and not a result of cross reactivity to Can f 1, Can f 2 or Can f 3.

The results obtained clearly demonstrated that prostatic kallikrein from dog is a major and unique allergen among the dog allergic subjects studied here. By both prevalence and magnitude of IgE binding, kallikrein was found to be the most important dog allergen so far described and among the subjects studied, over one third reacted to kallikrein but none of the other allergens tested.

EXAMPLE 4: Demonstration of kallikrein-specific IgE antibody binding activity in dog dander extract

An IgE inhibition experiment was performed to examine whether dog dander contains epitopes capable of binding kallikrein-reactive IgE antibodies. Serum samples from three dog sensitized subjects (A-C) with IgE reactivity to kallikrein were first incubated for 2 h at room temperature with purified kallikrein at a final concentration of 100 µg/mL and, in parallel as negative controls, with serum diluent or the non-allergenic maltose binding protein (MBP) of *E. coli*. All samples were then analysed in duplicate for IgE binding to ImmunoCAP tests carrying immobilized dog dander extract to study whether preincubation with kallikrein specifically would prevent IgE binding to dander protein attached to the solid phase. As a control for specificity of kallikrein as inhibitor, a serum from a subject (D) sensitized to Can f 1 and Can f 2, but not to kallikrein, was included alongside the other sera in the experiment.

The results of the inhibition experiment are shown in Figure 4. Kallikrein purified from dog urine was found to completely inhibit the IgE binding to dog dander of two (A and B) of the three kallikrein-reactive sera and partly the binding of the third serum (C), which was known to be reactive also to other dog allergens. The negative control protein, MBP, showed no significant inhibitory effect as compared to serum diluent. In addition, no inhibition by kallikrein was observed on IgE binding of the Can f 1- and Can f 2-reactive serum (D).

The results demonstrated that epitope structures capable of binding kallikrein-reactive IgE antibodies are present in dog dander and, hence, are not confined to urine.

EXAMPLE 5: Assessment of IgE binding to a kallikrein-like 28 kDa protein from dog dander extract using immunoblot analysis

With the aim to identify a protein present in dog dander to which the observed kallikrein-like allergen activity may be attributed, 37 sera with known levels of kallikrein-reactive IgE were used in an immunoblot experiment. Immunoblot analysis was performed on non-reduced dog dander extract separated by SDS-

PAGE (12.5% Excel 2-D gel, GE Healthcare Biosciences) and electroblotted onto nitrocellulose membrane (Hybond ECL, GE Healthcare Biosciences). Protein blots were blocked for 1 h at room temperature using blocking buffer (50 mM phosphate pH 7.4, 0.1 % (v/v) Tween-20, 0.9% (w/v) NaCl, 0.3% w/v) Dextran
5 T10) and then incubated overnight with each patient's serum, diluted 1.5- to 20-fold in blocking buffer. The dilution factor for each serum is indicated in brackets at the top of its corresponding membrane strip in Figure 5. After washing in blocking buffer with 0.5 % (v/v) Tween-20, the membrane was incubated 4 hrs at room temperature with an ¹²⁵I-labelled anti-human IgE
10 antibody in blocking buffer and bound IgE was then radiographically detected using a storage phosphor screen and a Variable Mode Imager, Typhoon 9410 (GE Healthcare Biosciences).

The results of the experiment are shown in Figure 5a-b. Of the 37 sera used, 30 showed IgE binding to a 28 kDa protein while 21 showed IgE binding to a 23
15 kDa band, corresponding to Can f 1 and/or possibly Can f 2. Immunoblotting signal intensities were quantified using the Phoretix 1D software (Nonlinear Dynamics Ltd, Newcastle upon Tyne, UK). The level of IgE reactivity in each serum to individual bands was calculated by multiplying the signal intensity with the serum dilution factor. Figure 6 shows a comparison of the level of IgE
20 binding to the 28 kDa band in immunoblot analysis and the kallikrein ImmunoCAP measurements described in Example 3 above, revealing a close correlation.

In order to directly examine the relationship between urinary kallikrein and the 28 kDa band in dog dander, an immunoblot inhibition experiment was
25 performed. A serum mono-reactive to the 28 kDa band was preincubated 2 hrs at room temperature with either purified urinary kallikrein or rCan f 1, both at a final concentration of 100 µg/mL, or with serum diluent. Membrane strips carrying immunoblotted non-reduced dog dander extract were then subjected to the preincubated serum samples and IgE binding was analysed as described
30 above. The experiment revealed that IgE binding to the 28 kDa band in dog dander was completely abolished by serum preincubation with kallikrein

whereas it remained unaffected by preincubation with both rCan f 1 and buffer alone (Figure 7).

Taken together, the results described in this example demonstrated the presence in dog dander extract of a protein displaying close electrophoretic and immunological similarity to urinary kallikrein.

Example 6: Partial purification and identification of kallikrein in dog dander

The kallikrein-like protein from dog dander was purified by SEC and RPC for biochemical identification. Three grams of dog dander (Allergon, Valinge, Sweden) was extracted in a 100 mL of MBS by end-over-end rotation for 3 hrs at room temperature. After centrifugation at 20,000 x g and concentration using an Amicon filter (PM-10, Millipore, Billerica, MA, USA), the extract was applied to an XK50/100 Superdex 75 column (GE Healthcare Biosciences) and eluted using MBS (Figure 8). Fractions from six peaks (indicated 1-6 in Figure 8) were pooled and analysed for allergen activity. The protein content of each fraction was immobilized on ImmunoCAP (Phadia, Uppsala, Sweden) solid phase and its IgE antibody binding activity tested using eight sera from dog dander sensitized individuals, as indicated in Table 4. Most of these sera were selected as having high IgE binding to dog dander extract but relatively low binding to either of rCan f 1, rCan f 2 and nCan f 3. From Table 4 it is evident that peak 3 from the SEC separation contained the highest level of IgE binding activity of the six peaks tested. This pool was selected for further purification.

After adding TFA to a final concentration of 0.065%, the pool was applied to a ST4.6/100 Source 15 RPC column (GE Healthcare Biosciences) and elution was performed using a linear, 0-54% gradient of acetonitril in water containing 0.05% TFA (Figure 8). Analysis of allergen reactivity of the three peaks indicated in Figure 9 was performed using five sera, selected by the criteria described above. The results of the analysis (Table 5) clearly showed that peak 1 contained the highest level of IgE antibody binding. Reducing SDS-PAGE analysis of this peak revealed the presence of 10 kDa, 18 kDa and 23 kDa protein bands (not shown).

The three band present in peak 1 were excised from the gel and subjected to in-gel digestion and mass spectrometric analysis as described in Example 2 above. While the 23 kDa band was identified as Can f 1, both the 10 kDa and 18 kDa bands were identified as dog prostatic kallikrein (Accession no P09582) after
5 PSD analysis of selected peptides $m/z=1004.52$ and $m/z=1632.98$, respectively.

Further, the two 10 kDa and 18 kDa bands were eluted from excised gel bands and subjected to N-terminal amino acid sequencing. The resulting sequences, xIGGRExLKN and AVxRPGEDRx, where “x” represents unresolved residues, matched residues 25-34 and 108-117 of the canine prostatic kallikrein
10 precursor sequence, Accession no P09582.

The results described in this example demonstrated that a protein with a primary structure identical or closely related to prostatic kallikrein is present in dog dander.

15 **Example 7: Similar IgE antibody reactivity to kallikrein from dog dander and urine**

To compare the IgE antibody binding activity of kallikrein from dog urine and dander, two kallikrein-reactive sera (sera no 6 and 8 from Table 2) and one kallikrein non-reactive serum (no 11) were used in immunoblot analysis of non-
20 reduced samples of dog dander extract, purified urinary kallikrein and partially purified kallikrein from dog dander (Figure 10). The two kallikrein-reactive sera displayed IgE binding to a 28 kDa band in all three preparations, indicating that IgE binding at 28 kDa in dog dander extract is due to kallikrein. In addition, it was evident that the dominant reactivity to the 28 kDa band in purified urinary
25 kallikrein coincided with the Coomassie stained protein bands of the same preparation. IgE binding to a band of about 55 kDa in the non-reduced kallikrein preparations is consistent with the notion in Example 1 above, of a putative dimer of kallikrein. The serum that was kallikrein non-reactive according to ImmunoCAP showed no IgE binding to the 28 kDa band in any of
30 the three allergen preparations analysed.

The immunoblotting reactivity to reduced kallikrein-containing samples was considerably weaker than to non-reduced samples. Only the purified urinary kallikrein preparation, which had a higher kallikrein concentration than the other preparations analysed, gave rise to detectable IgE binding to the 18 kDa band formed upon reduction.

The observation that the immune reactivity to purified urinary kallikrein in immunoblot analysis was directed against the major protein band at 28 kDa served to support the validity of the experimental kallikrein ImmunoCAP test, in that its IgE binding was not caused a contaminant of the protein preparation used. The results further show that at least some IgE binding epitopes on kallikrein are sensitive to reduction of the molecule, as indicated by the weaker antibody binding to the 10 kDa and 18 kDa subunits, as compared to the 28 kDa unreduced molecule.

EXAMPLE 8: Assessment of IgE-binding properties of a modified kallikrein or a variant or fragment of kallikrein (analyte)

The analyte is immobilized to a solid support, such as ImmunoCAP (Phadia, Uppsala, Sweden). Serum samples from at least three representative human patients sensitized to the relevant species and showing IgE reactivity to kallikrein from that species are incubated for 3 h at room temperature with kallikrein at a final concentration of 100 µg/mL and, in parallel as negative controls, with buffer alone and the non-allergenic maltose binding protein (MBP) of *E. coli*. The samples are then analysed for IgE binding to ImmunoCAP (Phadia, Uppsala, Sweden) tests carrying immobilized analyte to study whether preincubation with kallikrein specifically inhibits or significantly lowers IgE binding.

Example 9: Purification of kallikrein from dog urine by hydrophobic interaction chromatography (HIC)

A pooled sample of dog urine was filtered through a 5 µm and a 0.45 µm filter under nitrogen pressure. All chromatographic operations were performed with

an ÄKTA Explorer 100 Air system (GE Healthcare Biosciences, Uppsala, Sweden). Four aliquots of 120 ml filtered dog urine was buffer exchanged using a Sephadex G-25 column (GE Healthcare Biosciences, Uppsala, Sweden) (column volume 461 ml), with cleaning of the column after each run. Buffer
5 used: 50 mM Na-phosphate, 1 M $(\text{NH}_4)_2\text{SO}_4$, 0.02% NaN_3 , pH = 7. The sample (about 505 ml) was then filtered through a 0.45 μm filter and applied to the HIC column (HiPrep Phenyl FF (high sub), 20ml, GE Healthcare Biosciences, Uppsala, Sweden). Buffers used for HIC separation were: A) 50 mM Na-phosphate, 1 M $(\text{NH}_4)_2\text{SO}_4$, 0.02% NaN_3 , pH = 7, and B) 50 mM Na-phosphate,
10 0.02% NaN_3 , pH = 7. The flow through fraction (containing kallikrein) was collected in 10 ml fractions (Frac 950) at a flow rate of 5 ml/min and the flow through fractions were then pooled. The adsorbed material was eluted in a step gradient using 100% buffer B.

15 The fractions were analyzed using a BCA (bicinchoninic acid) assay, as well as SDS-PAGE (non reduced samples, silver staining). SDS-PAGE under non-reducing conditions revealed that kallikrein was found in the flow-through fractions which thus were pooled for further processing.

20 Two aliquots of about 125 ml and one aliquot of about 87 ml of the pooled HIC flow through fractions were buffer exchanged with a Sephadex G-25 SF column (GE Healthcare Biosciences, Uppsala, Sweden) to a buffer with the composition 20 mM Na-phosphate, 0.02% NaN_3 , pH = 8. The kallikrein-pool (456 ml) was then concentrated on an Amicon cell (350 ml, Millipore filter, PBCC, cutoff 5000
25 kDa, diameter 76 mm) to a volume of about 43 ml. Using BCA assay, the protein concentration in the final pool was determined to be 0.9 mg/ml (in 43 ml = totally 38,7 mg) The sample applied to the HIC-column contained 101 mg protein which yielded a recovery of 38% of kallikrein after the HIC purification.

30 The purity of the kallikrein preparation was assessed by analytical gel filtration on a Superdex 75 HR 10/30 column in an ÄKTA purifier XT10 system. For this experiment the sample volume was 100 μl and the buffer was 10 mM Na-phosphate, 150 mM NaCl, 0.02% NaN_3 , pH = 7.4.

EXAMPLE 10: Identification and characterization of kallikrein from dog urine by the use of electrophoresis and mass spectrometry

Using electrophoresis the following samples were compared on the same gel,
5 applying colloidal coomassie brilliant blue (CBB) staining:

1. A standard molecular weight marker
2. Dog urine
3. HIC-eluted material (reduced)
4. HIC flow-through fraction (reduced)
- 10 5. HIC flow-through fraction (non-reduced)
6. HIC-eluted material (non-reduced)

For samples 2, 3 and 6, a large number of proteins were detected. However, in sample 4 (reduced HIC flow-through fraction), only two main bands could be seen. These two bands were later, by the use of MALDI-TOF(-TOF) analysis (see
15 below), found to correspond to two different variants of the kallikrein protein (as a result of proteolysis at R107, due to arginin-esterase activity). In sample 5 (non-reduced HIC flow-through fraction), seven distinct bands were detected and all bands were found to correspond to different variants of the kallikrein protein by the use of MALDI-TOF(-TOF) analysis (see below). Kallikrein
20 comprises 12 cystein-residues, therefore the formation of different variants is possible under non-reducing conditions due to formation of cystein-cystein bridges. Thus, the formation of for example dimers, trimers etc is likely to happen under non-reducing conditions.

25 SDS-PAGE conditions, trypsin digestion and MALDI-TOF-TOF analysis:

Diluted samples were prepared using a SDS-PAGE cleanup kit according to the procedure recommended in the manual from the supplier (GE Healthcare, Uppsala, Sweden). The gels were run in a MES buffer at 200 V for 35 minutes. Reduced and non-reduced samples were run in separate aggregates. Staining

was done overnight with colloidal CBB (de-staining was later done using water immersion for about 5 hours). Samples were manually picked from the gel using a pipette tip, and treated according to a standard protocol (using ethanol instead of acetonitrile), incubated with 12.5 ng/ μ l trypsin over night at 37
 5 degrees Celsius. 0.5 μ l of the digested samples was applied on the target plate for the MALDI system and mixed with 0.5 μ l MALDI matrix solution (saturated solution of HCCA in 50% acetonitril, 0.1% TFA). All samples were analyzed, using a MALDI-TOF-TOF (Bruker Daltonics, Bremen, Germany) mass spectrometer. To identify proteins matching PMF results obtained, the MSDB
 10 database was searched using a Mascot server (Matrixscience, London, UK). MS-MS analysis was performed on selected peptides. External calibration was performed using a peptide calibration standard (Bruker Daltonics). Database searches were run using the following search criteria:

Taxonomy: mammalia

15 Mass tolerance: 100 ppm

Allowing for oxidized methionines and 1 missed cleavage.

Kallikrein sequence from the database:

MWFLALCLAMSLGWTGAEPHFQPRIIGGRECLKNSQPWQVAVYHNGEFACGGVLV
 20 NPEWVLTAAH**CANSNCEVWLGR**HNLSESEDEGQLVQVRK**SFIHPLYKTKVPR**-----
 -
AVIRPGEDRSHDLMLLHLEEPAKITKAVRVMDLPKKEPPLGSTCYVSGWGSTDPEI
 FHPGSLQCVDLKLSSNNQCAKVYTQKVTK**FMLCAGVLEGK**KDTCKGDSGGPLICD
 GELVGITSWGATPCGKPQMPSLYTR**VMPHLMWIKDTMK**ANT (SEQ ID NO: 1)
 25 (Peptide sequences identified by MALDI-TOF (/TOF) MS (/MS) in bold italic.)

EXAMPLE 11: Cloning, purification and assessment of the IgE binding activity of recombinant dog kallikrein expressed in *Pichia pastoris*

In order to verify the identification and importance of urinary kallikrein as a dog
 30 allergen, the protein was produced as a recombinant allergen using *Pichia*

pastoris as expression host, purified and analysed for IgE antibody binding activity.

Preparation of synthetic gene construct encoding dog urinary kallikrein

A synthetic dog urinary kallikrein gene was designed by back-translating into
5 nucleotide sequence the part of the reported amino acid sequence of dog
prostatic arginine esterase (urinary kallikrein, Acc. No. P09582) corresponding
to the mature protein. The nucleotide sequence was designed for optimal codon
usage and synonymously adjusted to minimize secondary structures and
eliminate or add restriction enzyme sites as desired. Oligonucleotides
10 corresponding to the final coding sequence were obtained and assembled, and
the full-length synthetic gene amplified by PCR and cloned into the *Xho*I and
*Sal*I sites of vector pPICZ A (Invitrogen, Carlsbad, CA, USA), adding a C-
terminal hexahistidine tag to enable protein purification by immobilised metal
ion affinity chromatography (IMAC). The plasmid DNA construct was linearized
15 by *Sac* I digestion and transformed into *P. pastoris* strain X-33 for homologous
recombination into the chromosomal AOX1 locus.

Expression and purification of recombinant dog kallikrein-2

The recombinant protein was produced in *Pichia pastoris* strain X-33
20 (Invitrogen) using a 7 L bioreactor (Belach Bioteknik, Solna, Sweden). A rich
broth medium (20 g/L peptone, 10 g/L yeast extract, 3.4 g/L yeast nitrogen
base, 10 g/L ammonium sulfate, 0.4 mg/L biotin and 0.1 M potassium
phosphate) was used and the cultivation carried out at 30°C. Expression was
induced and maintained by feeding methanol to the culture to a steady-state
25 concentration of 0.1% (v/v). After 70 hrs of fermentation, the culture was
harvested by centrifugation at 10 000 g for 10 min at +4°C and the supernatant
recovered for protein purification.

The supernatant was conditioned for purification by adding imidazole to 5 mM
and NaCl to 0.15 M and adjusting the pH to 7.2 using Tris base (s) before
30 applying it to a Streamline 25 chelating column (GE Healthcare Biosciences),
charged with NiSO₄ according to the manufacturer's recommendation. After

loading, the column was washed in separate steps with 20 mM and 60 mM imidazole and the recombinant protein was then eluted with 500 mM imidazole, all in a buffer composed of 20 mM Tris-HCl pH 8.0 and 0.15 M NaCl.

Further purification of the recombinant protein was performed using cation
5 exchange chromatography. IMAC fractions containing recombinant kallikrein were identified by SDS-PAGE, pooled and diluted with 2 volumes of 20 mM MES pH 6.0. After adjusting pH to 6.0, the diluted pool was applied to an XK26/100 SP Sepharose FF column (GE Healthcare Biosciences). The column was then washed with 2 column volumes of 0.15 M NaCl in 20 mM MES pH 6.0 and the
10 recombinant protein eluted with 0.30 M NaCl in the same buffer. The protein concentration was determined from absorbance at 280 nm, using a calculated extinction coefficient of 1.46 per mg/mL.

Although the synthetic kallikrein gene construct was designed to direct the production of a single polypeptide chain, the protein purified from the culture
15 medium was found to have undergone a partial cleavage into 18 kDa and 12 kDa chains (Fig 11), similar to the processing of natural urinary kallikrein. Indeed, N-terminal sequencing revealed that the recombinant kallikrein had been cleaved at the same position as the natural molecule (data not shown).

To assess the aggregation state and integrity of the recombinant protein under
20 physiological conditions, a sample of the preparation was subjected to analytical size exclusion chromatography. As shown in Figure 12, the chromatogram was dominated by a single symmetrical peak, corresponding to a molecular weight of 34 kDa as defined by the LMW Calibration Kit (GE Healthcare Biosciences). The analysis demonstrated that the recombinant protein, despite its partial
25 processing, was held together in solution and existed in a homogeneous, most likely monomeric, aggregation state.

IgE binding activity of recombinant kallikrein

The immunological activity of the recombinant kallikrein produced was assessed
30 in comparison to the natural protein purified from dog urine. The two proteins were immobilised separately on ImmunoCAP[®] solid phase and their *in vitro* IgE

binding capacity was examined using the 37 serum samples from dog allergic subjects described in Example 3 above.

As can be seen in Figure 13, the two datasets showed a very strong correlation ($r=0.9988$), demonstrating that the recombinant kallikrein produced closely

5 resembled natural urinary kallikrein with respect to IgE antibody binding.

Drawing from the complete absence of any other dog-derived protein in the recombinant protein preparation, it can be further noted that the results eliminate any possible doubt as to the identity of the active component of the natural kallikrein preparations described in the previous examples.

References

1. Custovic A, Green R, Taggart SCO, Smith A, Pickering CAC, Chapman MD et al. Domestic allergens in public places II: dog (Can f 1) and
5 cockroach (Bla g 2) allergens in dust and mite, cat, dog and cockroach
allergens in the air in public buildings. *Clinical & Experimental Allergy*
1996;**26**:1246-1252.
2. Spitzauer S, Schweiger C, Anrather J, Ebner C, Scheiner O, Kraft D et al.
10 Characterisation of dog allergens by means of immunoblotting.
International Archives of Allergy and Immunology 1993;**100**:60-67.
3. Spitzauer S. Allergy to mammalian proteins: At the borderline between
foreign and self? [Review]. *International Archives of Allergy and*
Immunology 1999;**120**:259-269.
4. de Groot H, Goei KGH, van Swieten P, Aalberse RC. Affinity purification of
15 a major and a minor allergen from dog extract: serologic activity of
affinity-purified Can f I and of Can f I-depleted extract. *Journal of Allergy*
and *Clinical Immunology* 1991;**87**:1056-1065.
5. Konieczny A, Morgenstern JP, Bizinkauskas CB, Lilley CH, Brauer AW,
20 Bond JF et al. The major dog allergens, Can f 1 and Can f 2, are salivary
lipocalin proteins: cloning and immunological characterization of the
recombinant forms. *Immunology* 1997;**92**:577-586.
6. Boutin Y, Hebert H, Vrancken ER, Mourad W. Allergenicity and cross-
reactivity of cat and dog allergenic extracts. *Clinical Allergy* 1988;**18**:287-
293.
- 25 7. Saarelainen S, Taivainen A, Rytönen-Nissinen M, Auriola S, Immonen A,
Mantylarvi R et al. Assessment of recombinant dog allergens Can f 1 and
Can f 2 for the diagnosis of dog allergy. *Clinical & Experimental Allergy*
2004;**34**:1576-1582.
8. Cabanas R, Lopez-Serrano MC, Carreira J, Ventas P, Polo F, Caballero MT
30 et al. Importance of albumin in cross-reactivity among cat, dog and horse
allergens. *Journal of Investigational Allergology & Clinical Immunology*
2000;**10**:71-77.
9. Bayard C, Holmquist L, Vesterberg O. Purification and identification of
allergenic alpha (2u)-globulin species of rat urine. *Biochim Biophys Acta*
35 1996;**1290**:129-134.
10. Ohman JL. Allergy in man caused by exposure to mammals. *J Am Vet*
Med Assoc 1978;**172**:1403-1406.
11. Schumacher MJ. Characterization of allergens from urine and pelts of
laboratory mice. *Mol Immunol* 1980;**17**:1087-1095.
- 40 12. Siraganian RP, Sandberg AL. Characterization of mouse allergens.
Journal of Allergy and Clinical Immunology 1979;**63**:435-442.
13. Taylor AN, Longbottom JL, Pepys J. Respiratory allergy to urine proteins
of rats and mice. *Lancet* 1977;**2**:847-849.
14. Hoffman DR. Dog and cat allergens: urinary proteins or dander proteins?
45 *Annals of Allergy* 1980;**45**:205-206.
15. Hiller R, Laffer S, Harwanegg C, Huber M, Schmidt WM, Twardosz A et al.
Microarrayed allergen molecules: diagnostic gatekeepers for allergy
treatment. *FASEB Journal* 2002;**16**:414-416.

16. Valenta R, Lidholm J, Niederberger V, Hayek B, Kraft D, Gronlund H. The recombinant allergen-based concept of component-resolved diagnostics and immunotherapy (CRD and CRIT). *Clinical & Experimental Allergy* 1999;**29**:896-904.
- 5 17. Cromwell O, Fiebig H, Suck R, Kahlert H, Nandy A, Kettner J et al. Strategies for recombinant allergen vaccines and fruitful results from first clinical studies. *Immunol Allergy Clin North Am* 2006;**26**:261-281, vii.
18. Gafvelin G, Thunberg S, Kronqvist M, Gronlund H, Gronneberg R, Troye-Blomberg M et al. Cytokine and antibody responses in birch-pollen-allergic patients treated with genetically modified derivatives of the major birch pollen allergen Bet v 1. *International Archives of Allergy and Immunology* 2005;**138**:59-66.
- 10 19. Jutel M, Jaeger L, Suck R, Meyer H, Fiebig H, Cromwell O. Allergen-specific immunotherapy with recombinant grass pollen allergens. *Journal of Allergy and Clinical Immunology* 2005;**116**:608-613.
- 15 20. Mahler V, Vrtala S, Kuss O, Diepgen TL, Suck R, Cromwell O et al. Vaccines for birch pollen allergy based on genetically engineered hypoallergenic derivatives of the major birch pollen allergen, Bet v 1. *Clinical & Experimental Allergy* 2004;**34**:115-122.
- 20 21. Weidinger S, Mayerhofer A, Raemsch R, Ring J, Kohn FM. Prostate-specific antigen as allergen in human seminal plasma allergy. *Journal of Allergy and Clinical Immunology* 2006;**117**:213-215.
22. Demoly P, Lebel B, Arnoux B. Allergen-induced mediator release tests. *Allergy* 2003;**58**:553-558.
- 25 23. Ebo DG, Hagendorens MM, Bridts CH, Schuerwegh AJ, De Clerck LS, Stevens WJ. In vitro allergy diagnosis: should we follow the flow? [Review]. *Clinical & Experimental Allergy* 2004;**34**:332-339.
24. Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal Biochem* 1996;**68**:850-858.
- 30 25. Frenette G, Deperthes D, Tremblay RR, Lazure C, Dube JY. Purification of enzymatically active kallikrein hK2 from human seminal plasma. *Biochim Biophys Acta* 1997;**1334**:109-115.
- 35 26. Marknell DeWitt Å, Niederberger V, Lehtonen P, Spitzauer S, Sperr WR, Valent P et al. Molecular and immunological characterization of a novel timothy grass (*Phleum pratense*) pollen allergen, Phl p 11. *Clinical & Experimental Allergy* 2002;**32**:1329-1340.
- 40 27. van Eijk HM, Rooyakkers DR, van Acker BA, Soeters PB, Deutz NE. Automated isolation of high-purity plasma albumin for isotope ratio measurements. *J Chromatogr B Biomed Sci App* 1999;**731**:199-205.

28
CLAIMS

1. Use of canine kallikrein as an allergen for detecting the presence of antibodies for *in vitro* diagnosis of Type I allergy.
2. The use of canine kallikrein according to claim 1, wherein said kallikrein is purified from a mammalian canine source or recombinantly produced.
3. The use of canine kallikrein according to claim 1 or claim 2, wherein said kallikrein is prostatic kallikrein.
4. The use of canine kallikrein according to any one of claims 1-3, wherein said kallikrein has an amino acid sequence according to SEQ ID NO: 1.
5. The use of canine kallikrein according to any one of claims 1-3, wherein said kallikrein comprises amino acids 25-260 of SEQ ID NO: 1.
6. Method for producing an allergen composition, comprising the step of adding a canine kallikrein to a composition comprising an allergen extract and/or at least one purified allergen component having:
 - no kallikrein content, or
 - low kallikrein content such that its IgE does not bind to other allergen components in the composition.
7. The method according to claim 6, wherein said canine kallikrein is purified from a mammalian source or recombinantly produced.
8. The method according to claim 6 or claim 7, wherein said canine kallikrein is prostatic kallikrein.
9. The method according to any one of claims 6-8, wherein said canine kallikrein has an amino acid sequence according to SEQ ID NO: 1.

10. The method according to any one of claims 6-8, wherein said canine kallikrein comprises amino acids 25-260 of SEQ ID NO: 1.

11. Method for *in vitro* diagnosis of type I allergy comprising the steps:

- contacting a immunoglobulin-containing body fluid sample from a patient suspected of having type I allergy with canine kallikrein, or with an allergen composition obtained by the method of any one of claims 6 to 10; and
- detecting the presence, in the sample, of IgE antibodies specifically binding to said canine kallikrein,

wherein the presence of such IgE antibodies specifically binding to said canine kallikrein is indicative of a type I allergy.

12. The method according to claim 11, wherein said canine kallikrein is purified from a mammalian canine source or recombinantly produced.

13. The method according to claim 11 or claim 12, wherein said canine kallikrein is prostatic kallikrein.

14. The method according to any one of claims 11-13, wherein said canine kallikrein has an amino acid sequence according to SEQ ID NO: 1.

15. Diagnostic kit for performing the method according to any one of claims 11-14, comprising canine kallikrein according to any one of claims 1-5.

16. Pharmaceutical composition comprising canine kallikrein and pharmaceutically acceptable carriers, excipients, buffers or diluents, or a 10 kDa or 18 kDa subunit thereof, for use in immunotherapy treatment of Type I allergy.

17. The pharmaceutical composition according to claim 16, wherein said canine kallikrein subunit is the 10 kDa subunit.

30

18. Use of canine kallikrein having an amino acid sequence according to SEQ ID NO: 1 as an allergen for detecting the presence of antibodies for diagnosis of Type I allergy to dog.
19. Canine kallikrein having an amino acid sequence according to SEQ ID NO: 1 for use in therapy of Type I allergy to dog.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5a




Figure 5b




Figure 6




Figure 7






Figure 8



10/14

Figure 9






Figure 11




Figure 12




Figure 13

