



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(22) Date de dépôt/Filing Date: 2011/06/22

(41) Mise à la disp. pub./Open to Public Insp.: 2011/12/29

(62) Demande originale/Original Application: 2 803 504

(30) Priorité/Priority: 2010/06/24 (EP10167243.4)

(51) Cl.Int./Int.Cl. *A61K 47/54* (2017.01),
A61K 31/522 (2006.01), *A61K 38/28* (2006.01),
A61P 3/10 (2006.01)

(71) Demandeur/Applicant:
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,
DE

(72) Inventeurs/Inventors:
KLEIN, THOMAS, DE;
MARK, MICHAEL, DE;
THOMAS, LEO, DE

(74) Agent: SMART & BIGGAR IP AGENCY CO.

(54) Titre : THERAPIE POUR LE DIABETE

(54) Title: DIABETES THERAPY

(57) Abrégé/Abstract:

The present invention relates to methods for treating and/or preventing metabolic diseases comprising the combined administration of a DPP-4 inhibitor and a long- acting insulin. The invention further relates to a DPP-4 inhibitor for subcutaneous or transdermal use.



ABSTRACT

The present invention relates to methods for treating and/or preventing metabolic diseases comprising the combined administration of a DPP-4 inhibitor and a long-acting insulin. The invention further relates to a DPP-4 inhibitor for subcutaneous or transdermal use.

Diabetes therapy

The present invention is a divisional of Canadian patent application no. 2803504, filed on June 22, 2011.

- The present invention relates methods for treating and/or preventing metabolic diseases, especially type 2 diabetes mellitus and/or conditions related thereto (e.g. diabetic complications) comprising the combined administration of long-acting insulin (such as e.g. insulin glargine or insulin detemir) and a certain DPP-4 inhibitor, to pharmaceutical compositions and combinations comprising such active components, and to certain therapeutic uses thereof.
- Further, the present invention relates to a method for improving glycemic control and/or preventing, reducing the risk of, slowing the progression of, delaying the onset or treating of complications of diabetes mellitus, such as micro- and macrovascular diseases (e.g. diabetic nephropathy, retinopathy or neuropathy, or cerebro- or cardiovascular complications such as e.g. myocardial infarction, stroke or vascular death or hospitalization), in a patient in need thereof (type 1 diabetes, LADA or, particularly, type 2 diabetes patient) comprising the combined (e.g. separate, simultaneous or sequential) administration of a long-acting insulin (such as e.g. insulin glargine or insulin detemir) and a certain DPP-4 inhibitor.
- Moreover, the present invention relates to a certain DPP-4 inhibitor for subcutaneous or transdermal (systemic) therapeutic use, particularly in treating and/or preventing the metabolic diseases described herein.
- Furthermore, the present invention relates to a certain DPP-4 inhibitor for subcutaneous administration (particularly for subcutaneous injection), e.g. once daily, each other day, thrice weekly, twice weekly or once weekly, preferably less than once daily.
- Furthermore, the present invention relates to a certain DPP-4 inhibitor for transdermal administration, e.g. once daily, each other day, thrice weekly, twice weekly or once weekly, preferably less than once daily.
- Further, the present invention relates to a parenteral (preferably subcutaneous) delivery device, preferably a subcutaneous injection device, which may be with or without needle (e.g. a needle-based pen injector or a jet/needle-free injector), containing long-acting insulin and a certain DPP-4 inhibitor and, optionally, one or more pharmaceutically acceptable carriers and/or diluents.
- Further, the present invention relates to a transdermal delivery device (e.g., a transdermal patch or gel) containing long-acting insulin and a certain DPP-4 inhibitor and, optionally, one or more pharmaceutically acceptable carriers and/or diluents.
- Further, the present invention relates to the DPP-4 inhibitors and/or long-acting insulin, each as defined herein, for use in the combination therapies as described herein.

- 2 -

Type 2 diabetes mellitus is a common chronic and progressive disease arising from a complex pathophysiology involving the dual endocrine effects of insulin resistance and impaired insulin secretion with the consequence not meeting the required demands to maintain plasma glucose levels in the normal range. This leads to chronic hyperglycaemia and its associated micro- and macrovascular complications or chronic damages, such as e.g. diabetic nephropathy, retinopathy or neuropathy, or macrovascular (e.g. cardio- or cerebrovascular) complications. The vascular disease component plays a significant role, but is not the only factor in the spectrum of diabetes associated disorders. The high frequency of complications leads to a significant reduction of life expectancy. Diabetes is currently the most frequent cause of adult-onset loss of vision, renal failure, and amputation in the Industrialised World because of diabetes induced complications and is associated with a two to five fold increase in cardiovascular disease risk.

The treatment of type 2 diabetes typically begins with diet and exercise, followed by oral antidiabetic monotherapy, and although conventional monotherapy may initially control blood glucose in some patients, it is however associated with a high secondary failure rate. The limitations of single-agent therapy for maintaining glycemic control may be overcome, at least in some patients, and for a limited period of time by combining multiple drugs to achieve reductions in blood glucose that cannot be sustained during long-term therapy with single agents. Available data support the conclusion that in most patients with type 2 diabetes current monotherapy will fail and treatment with multiple drugs will be required.

But, because type 2 diabetes is a progressive disease, even patients with good initial responses to conventional combination therapy will eventually require an increase of the dosage or further treatment with insulin because the blood glucose level is very difficult to maintain stable for a long period of time. Although existing combination therapy has the potential to enhance glycemic control, it is not without limitations (especially with regard to long term efficacy). Further, traditional therapies may show an increased risk for side effects, such as hypoglycemia or weight gain, which may compromise their efficacy and acceptability.

Thus, for many patients, these existing drug therapies result in progressive deterioration in metabolic control despite treatment and do not sufficiently control metabolic status especially over long-term and thus fail to achieve and to maintain glycemic control in advanced or late stage type 2 diabetes, including diabetes with inadequate glycemic control despite conventional oral or non-oral antidiabetic medication.

35

- 3 -

Therefore, although intensive treatment of hyperglycemia can reduce the incidence of chronic damages, many patients with type 2 diabetes remain inadequately treated, partly because of limitations in long term efficacy, tolerability and dosing inconvenience of conventional antihyperglycemic therapies.

5

This high incidence of therapeutic failure is a major contributor to the high rate of long-term hyperglycemia-associated complications or chronic damages (including micro- and makrovascular complications such as e.g. diabetic nephropathy, retinopathy or neuropathy, or cerebro- or cardiovascular complications such as e.g. myocardial infarction, stroke or death) in patients with type 2 diabetes.

10

Oral antidiabetic drugs conventionally used in therapy (such as e.g. first- or second-line, and/or mono- or (initial or add-on) combination therapy) include, without being restricted thereto, metformin, sulphonylureas, thiazolidinediones, glinides and α -glucosidase inhibitors.

15

Non-oral (typically injected) antidiabetic drugs conventionally used in therapy (such as e.g. first- or second-line, and/or mono- or (initial or add-on) combination therapy) include, without being restricted thereto, GLP-1 or GLP-1 analogues, and insulin or insulin analogues.

20

However, the use of these conventional antidiabetic or antihyperglycemic agents can be associated with various adverse effects. For example, metformin can be associated with lactic acidosis or gastrointestinal side effects; sulfonylureas, glinides and insulin or insulin analogues can be associated with hypoglycemia and weight gain; thiazolidinediones can be associated with edema, bone fracture, weight gain and heart failure/cardiac effects; and alpha-glucosidase blockers and GLP-1 or GLP-1 analogues can be associated with gastrointestinal adverse effects (e.g. dyspepsia, flatulence or diarrhea, or nausea or vomiting).

25

30

Therefore, it remains a need in the art to provide efficacious, safe and tolerable antidiabetic therapies.

Further, within the therapy of type 2 diabetes, it is a need for treating the condition effectively, avoiding the complications inherent to the condition, and delaying disease progression.

- 4 -

Further, within the therapy of type 2 diabetes, it is a need for sustained improvements in diabetic phenotype, glycemic and/or metabolic control, and/or (blood) glucose profile (preferably over long-term and/or during chronic treatment).

- 5 Furthermore, it remains a need that antidiabetic treatments not only prevent the long-term complications often found in advanced stages of diabetes disease, but also are a therapeutic option in those diabetes patients who have developed or are at risk of developing complications, such as renal impairment.
- 10 Moreover, it remains a need to provide prevention or reduction of risk for adverse effects associated with conventional antidiabetic therapies.

The enzyme DPP-4 (dipeptidyl peptidase IV) also known as CD26 is a serine protease known to lead to the cleavage of a dipeptide from the N-terminal end of a number of proteins
15 having at their N-terminal end a prolin or alanin residue. Due to this property DPP-4 inhibitors interfere with the plasma level of bioactive peptides including the peptide GLP-1 and are considered to be promising drugs for the treatment of diabetes mellitus.

For example, DPP-4 inhibitors and their uses are disclosed in WO 2002/068420, WO
20 2004/018467, WO 2004/018468, WO 2004/018469, WO 2004/041820, WO 2004/046148, WO 2005/051950, WO 2005/082906, WO 2005/063750, WO 2005/085246, WO 2006/027204, WO 2006/029769, WO2007/014886; WO 2004/050658, WO 2004/111051, WO 2005/058901, WO 2005/097798; WO 2006/068163, WO 2007/071738, WO 2008/017670; WO 2007/128721, WO 2007/128724, WO 2007/128761, or WO 2009/121945.

25 Long-acting insulin within the meaning of this invention refers to a formulation of insulin or insulin analogue that usually starts working within about 1-6 hours and is usually effective for up to 24 hours or more (e.g. up to 36 hours). Long-acting insulin usually provides a continuous level of insulin activity (for up to 24-36 hours) and usually operates at a maximum
30 strength (with flat action profile) after about 8-12 hours, sometimes longer. Long-acting insulin is usually administered in the morning or before bed. Examples of long-acting insulin may include, but are not limited to, insulin glargine, insulin detemir or insulin degludec, which are insulin analogues, and ultralente insulin, which is regular human insulin formulated for slow absorption. Long-acting insulin is suited to provide for basal, as opposed to prandial,
35 insulin requirements (e.g. to control hyperglycemia). Long-acting insulin may be typically

- 5 -

administered ranging from twice or once daily, over thrice weekly up to once weekly (ultra long-acting insulin). The route of administration of insulin may include, without being limited, invasive delivery (such as e.g. via parenteral route, preferentially via subcutaneous injection) or non-invasive delivery (such as e.g. via oral, buccal/sublingual, pulmonary, intranasal or
5 transdermal (e.g. via iontophoresis, sonophoresis or vesicular carriers) route), with subcutaneously injectable long-acting insulin being preferred.

In one embodiment, the long-acting insulin of this invention refers to any basal insulin known in the art, preferably having a basal release profile. A basal release profile refers to the
10 kinetic, amount and rate of release of the insulin or insulin analogue from the formulation into a patient's systemic circulation. In a graph of the patient's mean plasma insulin levels over time, a basal release profile typically has a minimal peak (often referred to as "a peakless profile" or "flat profile") and slowly and continuously releases insulin for a prolonged period of time.

15 In a further embodiment, the long-acting insulin is an acylated derivative of human insulin. Acylated insulin derivatives may be such wherein a lipophilic group is attached to the lysine residue in position B29. A commercial product is Levemir® comprising Lys^{B29}(N^ε-tetradecanoyl) des(B30) human insulin (cf. insulin detemir). Another example is N^{εB29}-(N^α-(ω-carboxypentadecanoyl)-L-γ-glutamyl) des(B30) human insulin (cf. insulin degludec).
20

In a further embodiment, the long-acting insulin is such comprising positively charged amino acids such as Arg attached to the C-terminal end of the B-chain. A commercial product is Lantus® comprising Gly^{A21}, Arg^{B31}, Arg^{B32} human insulin (cf. insulin glargine).
25

Insulin glargine (marketed as LANTUS® by Sanofi-Aventis) is approved and marketed for subcutaneous administration once a day. Insulin glargine provides relatively constant glucose lowering activity over a 24-hour period and may be administered any time during the day provided it is administered at the same time every day.

30 Insulin detemir (marketed as LEVEMIR® by Novo Nordisk) is approved and marketed for subcutaneous administration either twice a day or once a day, preferably with the evening meal or at bedtime.

Insulin degludec (NN1250) is a neutral, soluble ultra-long acting insulin with a duration of action more than 24 hours. Degludec has a very flat, predictable and smooth action profile. It
35 is intended for subcutaneous administration once daily or less (e.g. three times a week).

- 6 -

- In addition, further examples of long-acting insulin may include, without being limited to,
- insulin lispro PEGylated with high molecular weight poly(ethylene glycol) derivatives especially as disclosed in WO 2009/152128 (the disclosure of which is incorporated herein),
5 such as e.g. the PEGylated insulin lispro compound of the formula P-[(A)-(B)], or a pharmaceutically acceptable salt thereof, wherein A is the A-chain of insulin lispro, B is the B-chain of insulin lispro, and P is a PEG having a molecular weight in the range from about 17,5 kDa to about 40 kDa, and wherein A and B are properly cross-linked and P is attached via a urethane covalent bond to the epsilon-amino group of the lysine at position 28 of B;
 - 10 - amidated insulin glargine especially in the form of Gly^{A21}, Arg^{B31}, Arg^{B32}-NH₂ human insulin (insulin glargine amide, i.e. the C-terminus of the B-chain of insulin glargine is amidated) as disclosed in WO 2008/006496 or WO 2008/006496 (the disclosures of which are incorporated herein); .
 - Lys^{B29}(N^ε-lithocholyl-γ-Glu) des(B30) human insulin or N^ε^{B29}-ω-carboxypentadecanoyl-γ-amino-butanoyl des(B30) human insulin; or
15 - amidated insulin analogs as disclosed in WO 2009/087082 (the disclosure of which is incorporated herein), especially one selected from claim 14, or in WO 2009/087081 (the disclosure of which is incorporated herein), especially one selected from claim 16.
- 20 Long-acting insulin analogues are typically given as basic anti-diabetic therapy to type 2 diabetes, type 1 diabetes or latent autoimmune diabetes with onset in adults (LADA) patients to control the blood sugar when no food intake occurs. As mentioned above, this type of insulin provides a continuous level of insulin activity for up to 36 hours. Long-acting insulin operates at maximum strength after about 8-12 hours. Because of their advantages, it is
25 thought that treatment with these insulin analogues can lead to a beneficial effect, for example less hypoglycaemia, less weight gain or a better metabolic control possibly resulting in less late diabetic complications such as problems with eyes, kidneys or feet and myocardial infarction, stroke or death. DPP-4 inhibitors are a diverse group of anti-diabetics, however also working via insulin increasing mechanisms and limited on still functioning β-
30 cells for triggering endogenous insulin secretion. DPP-4 inhibitors lower glucagon levels and reduce post-prandial glucose peaks during food intake via increase of GLP-1 and subsequent mechanisms. The combination of this two principles is thereof a favorable approach in controlling efficacious fasted and post-prandial glucose levels. Also, by the complementary modes of action improved glucose tolerance, improved metabolic (glycemic)
35 status and/or sustainable efficacy can be achieved and/or maintained over longer time.

- 7 -

Linagliptin is the only DPP-4 inhibitor which demonstrates a superior pharmacokinetic profile (e.g. half life up to 72 h, stoichiometric reversible binding to DPP-4 protein) to be potentially administered once daily, each other day or even longer. In addition, various DPP-4 substrates (e.g. SDF-1, BNP) have been shown to have increased half lives and actions following inhibition of DPP-4 and thus being potentially of additional benefit for cardiovascular outcome. The combination of linagliptin with insulin according to this invention thus is regarded to further decrease late stage macrovascular complications.

In the monitoring of the treatment of diabetes mellitus the HbA1c value, the product of a non-enzymatic glycation of the haemoglobin B chain, is of exceptional importance. As its formation depends essentially on the blood sugar level and the life time of the erythrocytes the HbA1c in the sense of a "blood sugar memory" reflects the average blood sugar level of the preceding 4-12 weeks. Diabetic patients whose HbA1c level has been well controlled over a long time by more intensive diabetes treatment (i.e. < 6.5 % of the total haemoglobin in the sample) are significantly better protected from diabetic microangiopathy. The available treatments for diabetes can give the diabetic an average improvement in their HbA1c level of the order of 1.0 – 1.5 %. This reduction in the HbA1C level is not sufficient in all diabetics to bring them into the desired target range of < 7.0 %, preferably < 6.5 % and more preferably < 6 % HbA1c.

Within the meaning of this invention, inadequate or insufficient glycemic control means in particular a condition wherein patients show HbA1c values above 6.5%, in particular above 7.0%, even more preferably above 7.5%, especially above 8%. An embodiment of patients with inadequate or insufficient glycemic control include, without being limited to, patients having a HbA1c value from 7.5 to 10% (or, in another embodiment, from 7.5 to 11%). A special sub-embodiment of inadequately controlled patients refers to patients with poor glycemic control including, without being limited, patients having a HbA1c value $\geq$ 9%.

Within glycemic control, in addition to improvement of the HbA1c level, other recommended therapeutic goals for type 2 diabetes mellitus patients are improvement of fasting plasma glucose (FPG) and of postprandial plasma glucose (PPG) levels to normal or as near normal as possible. Recommended desired target ranges of preprandial (fasting) plasma glucose are 70-130 mg/dL (or 90-130 mg/dL) or <110 mg/dL, and of two-hour postprandial plasma glucose are <180 mg/dL or <140 mg/dL.

- 8 -

In one embodiment, diabetes patients within the meaning of this invention may include patients who have not previously been treated with an antidiabetic drug (drug-naïve patients). Thus, in an embodiment, the therapies described herein may be used in naïve patients. In another embodiment, diabetes patients within the meaning of this invention may include patients with advanced or late stage type 2 diabetes mellitus (including patients with failure to conventional antidiabetic therapy), such as e.g. patients with inadequate glycemic control on one, two or more conventional oral and/or non-oral antidiabetic drugs as defined herein, such as e.g. patients with insufficient glycemic control despite (mono-)therapy with metformin, a thiazolidinedione (particularly pioglitazone), a sulphonylurea, a glinide, GLP-1 or GLP-1 analogue, insulin or insulin analogue, or an α -glucosidase inhibitor, or despite dual combination therapy with metformin/sulphonylurea, metformin/thiazolidinedione (particularly pioglitazone), sulphonylurea/ α -glucosidase inhibitor, pioglitazone/sulphonylurea, metformin/insulin, pioglitazone/insulin or sulphonylurea/insulin. Thus, in an embodiment, the therapies described herein may be used in patients experienced with therapy, e.g. with conventional oral and/or non-oral antidiabetic mono- or dual or triple combination medication as mentioned herein.

A further embodiment of diabetic patients within the meaning of this invention refers to patients ineligible for metformin therapy including

- patients for whom metformin therapy is contraindicated, e.g. patients having one or more contraindications against metformin therapy according to label, such as for example patients with at least one contraindication selected from:

- renal disease, renal impairment or renal dysfunction (e.g., as specified by product information of locally approved metformin),
- dehydration,
- unstable or acute congestive heart failure,
- acute or chronic metabolic acidosis, and
- hereditary galactose intolerance;

and

- patients who suffer from one or more intolerable side effects attributed to metformin, particularly gastrointestinal side effects associated with metformin, such as for example patients suffering from at least one gastrointestinal side effect selected from:

- nausea,
- vomiting,
- diarrhoea,

- 9 -

intestinal gas, and
severe abdominal discomfort.

5 A further embodiment of the diabetes patients which may be amenable to the therapies of this invention may include, without being limited, those diabetes patients for whom normal metformin therapy is not appropriate, such as e.g. those diabetes patients who need reduced dose metformin therapy due to reduced tolerability, intolerability or contraindication against metformin or due to (mildly) impaired/reduced renal function (including elderly patients, such as e.g. ≥ 60 -65 years).

10 A further embodiment of diabetic patients within the meaning of this invention refers to patients having renal disease, renal dysfunction, or insufficiency or impairment of renal function (including mild, moderate and severe renal impairment), e.g. as suggested by elevated serum creatinine levels (e.g. serum creatinine levels above the upper limit of normal
15 for their age, e.g. $\geq 130 - 150 \mu\text{mol/l}$, or $\geq 1.5 \text{ mg/dl}$ ($\geq 136 \mu\text{mol/l}$) in men and $\geq 1.4 \text{ mg/dl}$ ($\geq 124 \mu\text{mol/l}$) in women) or abnormal creatinine clearance (e.g. glomerular filtration rate (GFR) $\leq 30 - 60 \text{ ml/min}$).

In this context, for more detailed example, mild renal impairment may be e.g. suggested by a
20 creatinine clearance of $50-80 \text{ ml/min}$ (approximately corresponding to serum creatine levels of $\leq 1.7 \text{ mg/dL}$ in men and $\leq 1.5 \text{ mg/dL}$ in women); moderate renal impairment may be e.g. suggested by a creatinine clearance of $30-50 \text{ ml/min}$ (approximately corresponding to serum creatinine levels of >1.7 to $\leq 3.0 \text{ mg/dL}$ in men and >1.5 to $\leq 2.5 \text{ mg/dL}$ in women); and severe renal impairment may be e.g. suggested by a creatinine clearance of $< 30 \text{ ml/min}$
25 (approximately corresponding to serum creatinine levels of $>3.0 \text{ mg/dL}$ in men and $>2.5 \text{ mg/dL}$ in women). Patients with end-stage renal disease require dialysis (e.g. hemodialysis or peritoneal dialysis).

For other more detailed example, patients with renal disease, renal dysfunction or renal
30 impairment include patients with chronic renal insufficiency or impairment, which can be stratified according to glomerular filtration rate (GFR, ml/min/1.73m^2) into 5 disease stages: stage 1 characterized by normal GFR ≥ 90 plus either persistent albuminuria or known structural or hereditary renal disease; stage 2 characterized by mild reduction of GFR (GFR 60-89) describing mild renal impairment; stage 3 characterized by moderate reduction of
35 GFR (GFR 30-59) describing moderate renal impairment; stage 4 characterized by severe

- 10 -

reduction of GFR (GFR 15-29) describing severe renal impairment; and terminal stage 5 characterized by requiring dialysis or GFR < 15 describing established kidney failure (end-stage renal disease, ESRD).

- 5 A further embodiment of diabetic patients within the meaning of this invention refers to type 2 diabetes patients with or at risk of developing renal complications, such as diabetic nephropathy (including chronic and progressive renal insufficiency, albuminuria, proteinuria, fluid retention in the body (edema) and/or hypertension).
- 10 In a further embodiment, patients within the present invention may include type 1 diabetes, LADA or, particularly, type 2 diabetes patients, with or without obesity or overweight.

Within the scope of the present invention it has now been found that certain DPP-4 inhibitors as defined herein as well as pharmaceutical combinations, compositions or combined uses
15 according to this invention of these DPP-4 inhibitors and long-acting insulin (such as e.g. insulin glargine, insulin detemir or insulin degludec) as defined herein have properties, which make them suitable for the purpose of this invention and/or for fulfilling one or more of above needs.

20 The present invention thus relates to a combination comprising a certain DPP-4 inhibitor (particularly BI 1356) and a long-acting insulin (such as e.g. insulin glargine, insulin detemir or insulin degludec), each as defined herein, particularly for simultaneous, separate or sequential use in the therapies described herein.

25 The present invention further relates to a certain DPP-4 inhibitor (particularly BI 1356) in combination with a long-acting insulin (such as e.g. insulin glargine, insulin detemir or insulin degludec), each as defined herein, for use in the therapies described herein.

The present invention further relates to a method for treating and/or preventing metabolic
30 diseases, especially type 2 diabetes mellitus and/or conditions related thereto (e.g. diabetic complications) comprising the combined (e.g. simultaneous, separate or sequential) administration of an effective amount of a long-acting insulin (such as e.g. insulin glargine, insulin detemir or insulin degludec) as defined herein and of an effective amount of a DPP-4 inhibitor as defined herein to the patient (particularly human patient) in need thereof, such as
35 e.g. a patient as described herein.

- 11 -

The present invention further relates to at least one of the following methods:

- 5 - preventing, slowing the progression of, delaying or treating a metabolic disorder or disease, such as e.g. type 1 diabetes mellitus, type 2 diabetes mellitus, impaired glucose tolerance (IGT), impaired fasting blood glucose (IFG), hyperglycemia, postprandial hyperglycemia, postabsorptive hyperglycemia, overweight, obesity, dyslipidemia, hyperlipidemia, hypercholesterolemia, hypertriglyceridemia, hypertension, atherosclerosis, endothelial dysfunction, osteoporosis, chronic systemic inflammation, non alcoholic fatty liver disease (NAFLD), retinopathy, neuropathy, nephropathy, polycystic ovarian syndrome, and/or metabolic syndrome;
- 10 - improving and/or maintaining glycemic control and/or for reducing of fasting plasma glucose, of postprandial plasma glucose, of postabsorptive plasma glucose and/or of glycosylated hemoglobin HbA1c;
- 15 - preventing, slowing, delaying or reversing progression from pre-diabetes, impaired glucose tolerance (IGT), impaired fasting blood glucose (IFG), insulin resistance and/or from metabolic syndrome to type 2 diabetes mellitus;
- 20 - preventing, reducing the risk of, slowing the progression of, delaying or treating of complications of diabetes mellitus such as micro- and macrovascular diseases, such as nephropathy, micro- or macroalbuminuria, proteinuria, retinopathy, cataracts, neuropathy, learning or memory impairment, neurodegenerative or cognitive disorders, cardio- or cerebrovascular diseases, tissue ischaemia, diabetic foot or ulcer, atherosclerosis, hypertension, endothelial dysfunction, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable angina pectoris, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders, vascular restenosis, and/or stroke;
- 25 - reducing body weight and/or body fat and/or liver fat and/or intra-myocellular fat or preventing an increase in body weight and/or body fat and/or liver fat and/or intra-myocellular fat or facilitating a reduction in body weight and/or body fat and/or liver fat and/or intra-myocellular fat;
- 30 - preventing, slowing, delaying or treating the degeneration of pancreatic beta cells and/or the decline of the functionality of pancreatic beta cells and/or for improving, preserving and/or restoring the functionality of pancreatic beta cells and/or stimulating and/or restoring or protecting the functionality of pancreatic insulin secretion;
- 35 - preventing, slowing, delaying or treating non alcoholic fatty liver disease (NAFLD) including hepatic steatosis, non-alcoholic steatohepatitis (NASH) and/or liver fibrosis (such as e.g. preventing, slowing the progression, delaying, attenuating, treating or

- 12 -

reversing hepatic steatosis, (hepatic) inflammation and/or an abnormal accumulation of liver fat);

- preventing, slowing the progression of, delaying or treating type 2 diabetes with failure to conventional antidiabetic mono- or combination therapy;
 - 5 - achieving a reduction in the dose of conventional antidiabetic medication required for adequate therapeutic effect;
 - reducing the risk for adverse effects associated with conventional antidiabetic medication (e.g. hypoglycemia or weight gain); and/or
 - maintaining and/or improving the insulin sensitivity and/or for treating or preventing
 - 10 hyperinsulinemia and/or insulin resistance;
- in a patient in need thereof (such as e.g. a patient as described herein), said method comprising combined (e.g. simultaneous, separate or sequential) administration of a DPP-4 inhibitor as defined herein and a long-acting insulin as defined herein.

- 15 The present invention further relates to a certain DPP-4 inhibitor (particularly BI 1356) as defined herein for subcutaneous or transdermal use in the therapies described herein.

- The present invention further relates to a method for treating and/or preventing metabolic diseases, especially type 2 diabetes mellitus and/or conditions related thereto (e.g. diabetic
- 20 complications) comprising the administration of an effective amount of a DPP-4 inhibitor as defined herein, optionally in combination with one or more other therapeutic agents as described herein, to the patient (particularly human patient) in need thereof (such as e.g. a patient as described herein), wherein the administration of the DPP-4 inhibitor, and of optionally one or more other therapeutic agents, is by parenteral, such as e.g. subcutaneous
- 25 or transdermal route.

- The present invention further relates to the use of a DPP-4 inhibitor as defined herein for the manufacture of a pharmaceutical composition for subcutaneous use in the treatment and/or prevention of a metabolic disease, disorder or condition such as e.g. described herein,
- 30 especially type 2 diabetes mellitus and/or conditions related thereto (e.g. diabetic complications).

- The present invention further relates to at least one of the following methods:
- preventing, slowing the progression of, delaying or treating a metabolic disorder or
 - 35 disease, such as e.g. type 1 diabetes mellitus, type 2 diabetes mellitus, impaired glucose

- 13 -

- tolerance (IGT), impaired fasting blood glucose (IFG), hyperglycemia, postprandial hyperglycemia, postabsorptive hyperglycemia, overweight, obesity, dyslipidemia, hyperlipidemia, hypercholesterolemia, hypertriglyceridemia, hypertension, atherosclerosis, endothelial dysfunction, osteoporosis, chronic systemic inflammation, non alcoholic fatty liver disease (NAFLD), retinopathy, neuropathy, nephropathy, polycystic ovarian syndrome, and/or metabolic syndrome;
- 5 - improving and/or maintaining glycemic control and/or for reducing of fasting plasma glucose, of postprandial plasma glucose, of postabsorptive plasma glucose and/or of glycosylated hemoglobin HbA1c;
- 10 - preventing, slowing, delaying or reversing progression from pre-diabetes, impaired glucose tolerance (IGT), impaired fasting blood glucose (IFG), insulin resistance and/or from metabolic syndrome to type 2 diabetes mellitus;
- 15 - preventing, reducing the risk of, slowing the progression of, delaying or treating of complications of diabetes mellitus such as micro- and macrovascular diseases, such as nephropathy, micro- or macroalbuminuria, proteinuria, retinopathy, cataracts, neuropathy, learning or memory impairment, neurodegenerative or cognitive disorders, cardio- or cerebrovascular diseases, tissue ischaemia, diabetic foot or ulcer, atherosclerosis, hypertension, endothelial dysfunction, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable angina pectoris, peripheral arterial occlusive disease,
- 20 cardiomyopathy, heart failure, heart rhythm disorders, vascular restenosis, and/or stroke;
- 25 - reducing body weight and/or body fat and/or liver fat and/or intra-myocellular fat or preventing an increase in body weight and/or body fat and/or liver fat and/or intra-myocellular fat or facilitating a reduction in body weight and/or body fat and/or liver fat and/or intra-myocellular fat;
- 30 - preventing, slowing, delaying or treating the degeneration of pancreatic beta cells and/or the decline of the functionality of pancreatic beta cells and/or for improving, preserving and/or restoring the functionality of pancreatic beta cells and/or stimulating and/or restoring or protecting the functionality of pancreatic insulin secretion;
- 35 - preventing, slowing, delaying or treating non alcoholic fatty liver disease (NAFLD) including hepatic steatosis, non-alcoholic steatohepatitis (NASH) and/or liver fibrosis (such as e.g. preventing, slowing the progression, delaying, attenuating, treating or reversing hepatic steatosis, (hepatic) inflammation and/or an abnormal accumulation of liver fat);
- preventing, slowing the progression of, delaying or treating type 2 diabetes with failure to conventional antidiabetic mono- or combination therapy;

- 14 -

- achieving a reduction in the dose of conventional antidiabetic medication required for adequate therapeutic effect;
 - reducing the risk for adverse effects associated with conventional antidiabetic medication (e.g. hypoglycemia or weight gain); and/or
 - 5 - maintaining and/or improving the insulin sensitivity and/or for treating or preventing hyperinsulinemia and/or insulin resistance;
- in a patient in need thereof (such as e.g. a patient as described herein), said method comprising the parenteral, preferably subcutaneous or transdermal, administration of a DPP-4 inhibitor as defined herein, optionally in combination with one or more other
- 10 therapeutic agents as described herein.

Further, the present invention relates to the combination according to this invention comprising a DPP-4 inhibitor as defined herein and a long-acting insulin as defined herein for parenteral, particularly subcutaneous or transdermal, administration of one or both of the

15 active components to the patient in need thereof.

Further, the present invention relates to a pharmaceutical composition according to this invention comprising

a DPP-4 inhibitor as defined herein,

20 a long-acting insulin as defined herein,

and, optionally, one or more pharmaceutically acceptable carriers and/or diluents, said composition being for subcutaneous administration to the patient in need thereof, e.g. by injection.

25 In addition, the present invention relates to the combination according to this invention comprising a DPP-4 inhibitor as defined herein and a long-acting insulin as defined herein for use in treating and/or preventing (including slowing the progression or delaying the onset) of metabolic diseases as defined herein, particularly diabetes (especially type 2 diabetes or conditions related thereto, including diabetic complications), optionally in combination with

30 one or more other therapeutic agents as described herein.

In addition, the present invention relates to the use of a combination according to this invention comprising a DPP-4 inhibitor as defined herein and a long-acting insulin as defined herein for the manufacture of a medicament for use in a therapeutic method as described

35 hereinbefore or hereinafter.

- 15 -

In addition, the present invention relates to a combination according to this invention comprising a DPP-4 inhibitor as defined herein and a long-acting insulin as defined herein for use in a therapeutic method as described hereinbefore or hereinafter.

5

In addition, the present invention relates to a method of treating and/or preventing (including slowing the progression or delaying the onset) of a metabolic disease, particularly diabetes (especially type 2 diabetes) or conditions related thereto, including diabetic complications, comprising administering to the patient in need thereof (such as e.g. a patient as described
10 herein) a combination according to this invention comprising a DPP-4 inhibitor as defined herein and a long-acting insulin as defined herein.

In addition, the present invention relates a DPP-4 inhibitor as defined herein for use in a method as described hereinbefore or hereinafter, said method comprising administering the
15 DPP-4 inhibitor, optionally combined with one or more other active substances (e.g. which may selected from those mentioned herein, such as e.g. metformin or pioglitazone), in combination (e.g. separately, simultaneously or sequentially) with a long-acting insulin as defined herein to the patient

In addition, the present invention relates a long-acting insulin as defined herein for use in a method as described hereinbefore or hereinafter, said method comprising administering the long-acting insulin, optionally combined with one or more other active substances (e.g. which may selected from those mentioned herein, such as e.g. metformin or pioglitazone), in combination (e.g. separately, simultaneously or sequentially) with a DPP-4 inhibitor as
20 defined herein to the patient
25

In addition, the present invention relates to the use of a DPP-4 inhibitor as defined herein for the manufacture of a medicament for use in combination with a long-acting insulin as defined herein for treating and/or preventing (including slowing the progression or delaying the onset)
30 of metabolic diseases, particularly diabetes (especially type 2 diabetes) and conditions related thereto, including diabetic complications.

In addition, the present invention relates to the use of a long-acting insulin as defined herein for the manufacture of a medicament for use in combination with a DPP-4 inhibitor as defined
35 herein for treating and/or preventing (including slowing the progression or delaying the onset)

- 16 -

of metabolic diseases, particularly diabetes (especially type 2 diabetes) or conditions related thereto, including diabetic complications.

5 In addition, the present invention relates to the use of a DPP-4 inhibitor as defined herein, optionally in combination with one or more other active substances (such as e.g. metformin or pioglitazone), for the manufacture of a medicament for use in combination with a long-acting insulin as defined herein for treating and/or preventing (including slowing the progression or delaying the onset) of metabolic diseases, particularly diabetes (especially type 2 diabetes) or conditions related thereto, including diabetic complications.

10 In addition, the present invention relates a DPP-4 inhibitor as defined herein for use in a combination treatment according to the invention in a patient in need thereof (such as e.g. a patient as described herein).

15 In addition, the present invention relates a long-acting insulin as defined herein for use in a combination treatment according to the invention in a patient in need thereof (such as e.g. a patient as described herein).

20 In addition, the present invention relates a DPP-4 inhibitor as defined herein, optionally in combination with one or more other active substances (such as e.g. metformin or pioglitazone), for use in a combination treatment according to the invention in a patient in need thereof.

25 In addition, the invention relates to a DPP-4 inhibitor as defined herein for use in a method as described hereinbefore or hereinafter, said method comprising administering the DPP-4 inhibitor, optionally in combination with one or more other active substances (e.g. which may be selected from those mentioned herein, such as e.g. metformin or pioglitazone), to the patient.

30 Further, the invention relates to a DPP-4 inhibitor as defined herein for use in a method of preventing, reducing the risk of, slowing the progression of, delaying or treating weight gain associated with the therapy with a long-acting insulin as defined herein, preferably said method comprising administering the DPP-4 inhibitor and the long-acting insulin to the patient (e.g. type 1 diabetes, LADA or, particularly, type 2 diabetes patient, with or without obesity or overweight).

35

- 17 -

Further, the invention relates to a DPP-4 inhibitor as defined herein for use in a method of preventing, reducing the risk of, slowing the progression of, delaying or treating an increase in body weight and/or body fat and/or liver fat and/or intra-myocellular fat which may be associated with the therapy with a long-acting insulin as defined herein, preferably said
5 method comprising administering the DPP-4 inhibitor and the long-acting insulin to the patient (e.g. type 1 diabetes, LADA or, particularly, type 2 diabetes patient, with or without obesity or overweight).

Further, the invention relates to a DPP-4 inhibitor as defined herein for use in a method of
10 improving body fat composition and/or obesity parameters which may be associated with the therapy with a long-acting insulin as defined herein, preferably said method comprising administering the DPP-4 inhibitor and the long-acting insulin to the patient (e.g. type 1 diabetes, LADA or, particularly, type 2 diabetes patient, with or without obesity or
overweight).

15 Further, the invention relates to a DPP-4 inhibitor and a long-acting insulin each as defined herein for use in a method of improving diabetic phenotype, improving glycemic and/or metabolic control, improving (blood) glucose profile (e.g. improving the control of fasting and/or postprandial blood glucose levels) and/or improving glucagon suppression (e.g. over
20 long-term and/or during chronic treatment), preferably said method comprising administering the DPP-4 inhibitor and the long-acting insulin to the patient (e.g. type 1 diabetes, LADA or, particularly, type 2 diabetes patient, with or without obesity or overweight).

Further, the invention relates to a DPP-4 inhibitor as defined herein for use in a method of
25 sparing or reducing the amount of long-acting insulin as defined herein required for efficacious and/or safe therapeutic treatment, said method comprising administering the DPP-4 inhibitor and the long-acting insulin to the patient (e.g. type 1 diabetes, LADA or, particularly, type 2 diabetes patient, with or without obesity or overweight).

30 Further, the invention relates to a DPP-4 inhibitor as defined herein for use in a therapeutic or preventive method as described herein, said use comprising adding the DPP-4 inhibitor to long-acting insulin as defined herein alone or in combination with one or more other antidiabetic drugs (e.g. selected from metformin, pioglitazone and a sulphonylurea), e.g. for improving glycemic control in a patient (e.g. type 1 diabetes, LADA or, particularly, type 2
35 diabetes patient, with or without obesity or overweight) who may be inadequately controlled

- 18 -

on insulin alone or in combination with one or more other antidiabetic drugs (e.g. selected from metformin, pioglitazone and a sulphonylurea).

Further, the invention relates to a DPP-4 inhibitor and a long-acting insulin each as defined
5 herein for use in a therapeutic or preventive method as described herein, said use
comprising adding the DPP-4 inhibitor and the long-acting insulin to one or more other
antidiabetic drugs (e.g. selected from metformin, pioglitazone and a sulphonylurea), e.g. for
improving glycemic control in a patient (e.g. type 1 diabetes, LADA or, particularly, type 2
10 diabetes patient, with or without obesity or overweight) who may be inadequately controlled
on one or more other antidiabetic drugs (e.g. selected from metformin, pioglitazone and a
sulphonylurea).

Further, the invention relates to a long-acting insulin as defined herein for use in a
therapeutic or preventive method as described herein, said use comprising adding the long-
15 acting insulin to a DPP-4 inhibitor as defined herein alone or in combination with one or more
other antidiabetic drugs (e.g. selected from metformin, pioglitazone and a sulphonylurea),
e.g. for improving glycemic control in a patient (e.g. type 1 diabetes, LADA or, particularly,
type 2 diabetes patient, with or without obesity or overweight) who may be inadequately
controlled on a DPP-4 inhibitor alone or in combination with one or more other antidiabetic
20 drugs (e.g. selected from metformin, pioglitazone and a sulphonylurea).

Further, the invention relates to a combination or composition comprising a DPP-4 inhibitor
and a long-acting insulin each as defined herein for use in treatment and/or prevention as
described herein, optionally in combination with one or more other therapeutic agents.

25 Further, the invention relates to a DPP-4 inhibitor as defined herein with or without metformin
and/or pioglitazone in combination with a long-acting insulin as defined herein for therapeutic
or preventive use as described herein.

30 Further, the invention relates to a DPP-4 inhibitor as defined herein in combination with a
long-acting insulin as defined herein, with or without metformin, for use in a method of
therapy or prophylaxis as described herein.

Other aspects of the present invention become apparent to the skilled person from the
35 foregoing and following remarks (including the examples and claims).

- 19 -

The aspects of the present invention, in particular the pharmaceutical compounds, compositions, combinations, methods and uses, refer to DPP-4 inhibitors and/or long-acting insulins as defined hereinbefore and hereinafter.

5

A DPP-4 inhibitor within the meaning of the present invention includes, without being limited to, any of those DPP-4 inhibitors mentioned hereinabove and hereinbelow, preferably orally and/or subcutaneously active DPP-4 inhibitors.

- 10 An embodiment of this invention refers to a DPP-4 inhibitor for use in the treatment and/or prevention of metabolic diseases (particularly type 2 diabetes mellitus) in type 2 diabetes patients, wherein said patients further suffering from renal disease, renal dysfunction or renal impairment, particularly characterized in that said DPP-4 inhibitor is administered to said patients in the same dose levels as to patients with normal renal function, thus e.g. said
15 DPP-4 inhibitor does not require downward dosing adjustment for impaired renal function.

- For example, a DPP-4 inhibitor according to this invention (especially one which may be suited for patients with impaired renal function) may be such an oral DPP-4 inhibitor, which and whose active metabolites have preferably a relatively wide (e.g. about > 100 fold)
20 therapeutic window and/or, especially, that are primarily eliminated via hepatic metabolism or biliary excretion (preferably without adding additional burden to the kidney).

- In more detailed example, a DPP-4 inhibitor according to this invention (especially one which may be suited for patients with impaired renal function) may be such an orally administered
25 DPP-4 inhibitor, which has a relatively wide (e.g. > 100 fold) therapeutic window (preferably a safety profile comparable to placebo) and/or which fulfils one or more of the following pharmacokinetic properties (preferably at its therapeutic oral dose levels):

- The DPP-4 inhibitor is substantially or mainly excreted via the liver (e.g. > 80 % or even >
30 90 % of the administered oral dose), and/or for which renal excretion represents no substantial or only a minor elimination pathway (e.g. < 10 %, preferably < 7 %, of the administered oral dose measured, for example, by following elimination of a radiolabelled carbon (¹⁴C) substance oral dose);
- The DPP-4 inhibitor is excreted mainly unchanged as parent drug (e.g. with a mean of >
35 70%, or > 80%, or, preferably, 90% of excreted radioactivity in urine and faeces after oral

- 20 -

dosing of radiolabelled carbon (^{14}C) substance), and/or which is eliminated to a non-substantial or only to a minor extent via metabolism (e.g. < 30%, or < 20%, or, preferably, 10%);

- 5 - The (main) metabolite(s) of the DPP-4 inhibitor is/are pharmacologically inactive. Such as e.g. the main metabolite does not bind to the target enzyme DPP-4 and, optionally, it is rapidly eliminated compared to the parent compound (e.g. with a terminal half-life of the metabolite of ≤ 20 h, or, preferably, $\leq$ about 16 h, such as e.g. 15.9 h).

- 10 In one embodiment, the (main) metabolite in plasma (which may be pharmacologically inactive) of a DPP-4 inhibitor having a 3-amino-piperidin-1-yl substituent is such a derivative where the amino group of the 3-amino-piperidin-1-yl moiety is replaced by a hydroxyl group to form the 3-hydroxy-piperidin-1-yl moiety (e.g. the 3-(S)-hydroxy-piperidin-1-yl moiety, which is formed by inversion of the configuration of the chiral center).
- 15 Further properties of a DPP-4 inhibitor according to this invention may be one or more of the following: Rapid attainment of steady state (e.g. reaching steady state plasma levels (> 90% of the steady state plasma concentration) between second and fifth day of treatment with therapeutic oral dose levels), little accumulation (e.g. with a mean accumulation ratio $R_{A,AUC} \leq 1.4$ with therapeutic oral dose levels), and/or preserving a long-lasting effect on DPP-4
- 20 inhibition, preferably when used once-daily (e.g. with almost complete (> 90%) DPP-4 inhibition at therapeutic oral dose levels, > 80% inhibition over a 24h interval after once-daily intake of therapeutic oral drug dose), significant decrease in 2h postprandial blood glucose excursions by ≥ 80 % (already on first day of therapy) at therapeutic dose levels, and cumulative amount of unchanged parent compound excreted in urine on first day being
- 25 below 1% of the administered dose and increasing to not more than about 3-6% in steady state.

- 30 Thus, for example, a DPP-4 inhibitor according to this invention may be characterized in that said DPP-4 inhibitor has a primarily non-renal route of excretion, i.e. said DPP-4 inhibitor is excreted to a non-substantial or only to a minor extent (e.g. < 10 %, preferably < 7 %, e.g. about 5 %, of administered oral dose, preferably of oral therapeutic dose) via the kidney (measured, for example, by following elimination of a radiolabelled carbon (^{14}C) substance oral dose).

- 21 -

Further, a DPP-4 inhibitor according to this invention may be characterized in that said DPP-4 inhibitor is excreted substantially or mainly via the liver or faeces (measured, for example, by following elimination of a radiolabelled carbon (^{14}C) substance oral dose).

- 5 Further, a DPP-4 inhibitor according to this invention may be characterized in that said DPP-4 inhibitor is excreted mainly unchanged as parent drug (e.g. with a mean of > 70%, or > 80%, or, preferably, 90 % of excreted radioactivity in urine and faeces after oral dosing of radiolabelled carbon (^{14}C) substance),
said DPP-4 inhibitor is eliminated to a non-substantial or only to a minor extent via
10 metabolism, and/or
the main metabolite of said DPP-4 inhibitor is pharmacologically inactive or has a relatively wide therapeutic window.


- Further, a DPP-4 inhibitor according to this invention may be characterized in that
15 said DPP-4 inhibitor does not significantly impair glomerular and/or tubular function of a type 2 diabetes patient with chronic renal insufficiency (e.g. mild, moderate or severe renal impairment or end stage renal disease), and/or
said DPP-4 inhibitor trough levels in the blood plasma of type 2 diabetes patients with mild or moderate renal impairment are comparable to the levels in patients with normal renal
20 function, and/or
said DPP-4 inhibitor does not require to be dose-adjusted in a type 2 diabetes patient with impaired renal function (e.g. mild, moderate or severe renal impairment or end stage renal disease, preferably regardless of the stage of renal impairment).

- 25 Further, a DPP-4 inhibitor according to this invention may be characterized in that said DPP-4 inhibitor provides its minimally effective dose at that dose that results in >50% inhibition of DPP-4 activity at trough (24 h after last dose) in >80% of patients, and/or
said DPP-4 inhibitor provides its fully therapeutic dose at that dose that results in >80% inhibition of DPP-4 activity at trough (24 h after last dose) in >80% of patients.


- 30 Further, a DPP-4 inhibitor according to this invention may be characterized in that being suitable for use in type 2 diabetes patients who are with diagnosed renal impairment and/or who are at risk of developing renal complications, e.g. patients with or at risk of diabetic nephropathy (including chronic and progressive renal insufficiency, albuminuria, proteinuria,
35 fluid retention in the body (edema) and/or hypertension).

- 22 -


In a first embodiment (embodiment **A**), a DPP-4 inhibitor in the context of the present invention is any DPP-4 inhibitor of formula (I)




or formula (II)



10 or formula (III)



or formula (IV)



15

wherein **R1** denotes ([1,5]naphthyridin-2-yl)methyl, (quinazolin-2-yl)methyl, (quinoxalin-6-yl)methyl, (4-methyl-quinazolin-2-yl)methyl, 2-cyano-benzyl, (3-cyano-quinolin-2-yl)methyl, (3-cyano-pyridin-2-yl)methyl, (4-methyl-pyrimidin-2-yl)methyl, or (4,6-dimethyl-pyrimidin-2-

- 23 -

yl)methyl and **R2** denotes 3-(*R*)-amino-piperidin-1-yl, (2-amino-2-methyl-propyl)-methylamino or (2-(*S*)-amino-propyl)-methylamino, or its pharmaceutically acceptable salt.


- 5 In a second embodiment (embodiment **B**), a DPP-4 inhibitor in the context of the present invention is a DPP-4 inhibitor selected from the group consisting of sitagliptin, vildagliptin, saxagliptin, alogliptin, gemigliptin, (2*S*)-1-[[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethylamino]-acetyl]-pyrrolidine-2-carbonitrile, (2*S*)-1-[[1,1,-Dimethyl-3-(4-pyridin-3-yl-imidazol-1-yl)-propylamino]-acetyl]-pyrrolidine-2-
 10 carbonitrile, (S)-1-((2*S*,3*S*,11*bS*)-2-Amino-9,10-dimethoxy-1,3,4,6,7,11*b*-hexahydro-2*H*-pyrido[2,1-*a*]isoquinolin-3-yl)-4-fluoromethyl-pyrrolidin-2-one, (3,3-Difluoropyrrolidin-1-yl)-((2*S*,4*S*)-4-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrrolidin-2-yl)methanone,
 15 1-((3*S*,4*S*)-4-amino-1-(4-(3,3-difluoropyrrolidin-1-yl)-1,3,5-triazin-2-yl)pyrrolidin-3-yl)-5,5-difluoropiperidin-2-one, (2*S*,4*S*)-1-{2-[(3*S*,1*R*)-3-(1*H*-1,2,4-Triazol-1-ylmethyl)cyclopentylamino]-acetyl}-4-fluoropyrrolidine-2-carbonitrile, (R)-2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2*H*-pyrimidin-1-ylmethyl]-4-
 20 fluoro-benzonitrile, 5-{(S)-2-[2-((S)-2-Cyano-pyrrolidin-1-yl)-2-oxo-ethylamino]-propyl}-5-(1*H*-tetrazol-5-yl)-10,11-dihydro-5*H*-dibenzo[*a,d*]cycloheptene-2,8-dicarboxylic acid bis-dimethylamide, 3-{(2*S*,4*S*)-4-[4-(3-Methyl-1-phenyl-1*H*-pyrazol-5-yl)piperazin-1-yl]pyrrolidin-2-ylcarbonyl}thiazolidine,
 25 [(2*R*)-1-[(3*R*)-pyrrolidin-3-ylamino]acetyl]pyrrolidin-2-yl]boronic acid, (2*S*,4*S*)-1-[2-[(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile,
 2-{6-[(3*R*)-3-amino-3-methylpiperidin-1-yl]-1,3-dimethyl-2,4-dioxo-1,2,3,4-tetrahydro-5*H*-pyrrolo[3,2-*d*]pyrimidin-5-yl}methyl-4-fluorobenzonitrile,
 30 6-[(3*R*)-3-amino-piperidin-1-yl]-5-(2-chloro-5-fluoro-benzyl)-1,3-dimethyl-1,5-dihydro-pyrrolo[3,2-*d*]pyrimidine-2,4-dione, and (S)-2-methylpyrazolo[1,5-*a*]primidine-6-carboxylic acid {2-[(2-cyanopyrrolidin-1-yl)-2-oxoethylamino]-2-methylpropyl}amide,
 or its pharmaceutically acceptable salt.

35


- 24 -

Regarding the first embodiment (embodiment **A**), preferred DPP-4 inhibitors are any or all of the following compounds and their pharmaceutically acceptable salts:


- 1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-(3-(*R*)-amino-piperidin-1-yl)-xanthine (compare WO 2004/018468, example 2(142));



- 1-[[[1,5]naphthyridin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((*R*)-3-amino-piperidin-1-yl)-xanthine (compare WO 2004/018468, example 2(252));




- 1-[(Quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((*R*)-3-amino-piperidin-1-yl)-xanthine (compare WO 2004/018468, example 2(80));




- 2-((*R*)-3-Amino-piperidin-1-yl)-3-(but-2-ynyl)-5-(4-methyl-quinazolin-2-ylmethyl)-3,5-dihydro-imidazo[4,5-d]pyridazin-4-one (compare WO 2004/050658, example 136):

- 25 -




- 1-[(4-Methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(2-amino-2-methyl-propyl)-methylamino]-xanthine (compare WO 2006/029769, example 2(1)):




5

- 1-[(3-Cyano-quinolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-amino-piperidin-1-yl]-xanthine (compare WO 2005/085246, example 1(30)):




10

- 1-(2-Cyano-benzyl)-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-amino-piperidin-1-yl]-xanthine (compare WO 2005/085246, example 1(39)):




- 26 -


- 1-[(4-Methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*S*)-(2-amino-propyl)-methylamino]-xanthine (compare WO 2006/029769, example 2(4)):




- 5 • 1-[(3-Cyano-pyridin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-amino-piperidin-1-yl]-xanthine (compare WO 2005/085246, example 1(52)):



- 10 • 1-[(4-Methyl-pyrimidin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-amino-piperidin-1-yl]-xanthine (compare WO 2005/085246, example 1(81)):




- 1-[(4,6-Dimethyl-pyrimidin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-[(*R*)-3-amino-piperidin-1-yl]-xanthine (compare WO 2005/085246, example 1(82)):



15

- 27 -

- 1-[(Quinoxalin-6-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-((*R*)-3-amino-piperidin-1-yl)-xanthine (compare WO 2005/085246, example 1(83)):



- 5 These DPP-4 inhibitors are distinguished from structurally comparable DPP-4 inhibitors, as they combine exceptional potency and a long-lasting effect with favourable pharmacological properties, receptor selectivity and a favourable side-effect profile or bring about unexpected therapeutic advantages or improvements when combined with other pharmaceutical active substances. Their preparation is disclosed in the publications mentioned.

10


A more preferred DPP-4 inhibitor among the abovementioned DPP-4 inhibitors of embodiment **A** of this invention is 1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-(3-(*R*)-amino-piperidin-1-yl)-xanthine, particularly the free base thereof (which is also known as linagliptin or BI 1356).

15

As further DPP-4 inhibitors the following compounds can be mentioned:

- Sitagliptin (MK-0431) having the structural formula A below is (3*R*)-3-amino-1-[3-(trifluoromethyl)-5,6,7,8-tetrahydro-5*H*-[1,2,4]triazolo[4,3-*a*]pyrazin-7-yl]-4-(2,4,5-trifluorophenyl)butan-1-one, also named (2*R*)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-*a*]pyrazin-7(8*H*)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine,

20



- 25 In one embodiment, sitagliptin is in the form of its dihydrogenphosphate salt, i.e. sitagliptin phosphate. In a further embodiment, sitagliptin phosphate is in the form of a crystalline

- 28 -

anhydrate or monohydrate. A class of this embodiment refers to sitagliptin phosphate monohydrate. Sitagliptin free base and pharmaceutically acceptable salts thereof are disclosed in US Patent No. 6,699,871 and in Example 7 of WO 03/004498. Crystalline sitagliptin phosphate monohydrate is disclosed in WO 2005/003135 and in WO

5 2007/050485.


For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

A tablet formulation for sitagliptin is commercially available under the trade name Januvia®. A

10 tablet formulation for sitagliptin/metformin combination is commercially available under the trade name Janumet®.

- Vildagliptin (LAF-237) having the structural formula B below is (2S)-[[(3-hydroxyadamantan-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile, also named (S)-1-[(3-hydroxy-1-

15 adamantyl)amino]acetyl-2-cyano-pyrrolidine,



Vildagliptin is specifically disclosed in US Patent No. 6,166,063 and in Example 1 of WO
20 00/34241. Specific salts of vildagliptin are disclosed in WO 2007/019255. A crystalline form of vildagliptin as well as a vildagliptin tablet formulation are disclosed in WO 2006/078593. Vildagliptin can be formulated as described in WO 00/34241 or in WO 2005/067976. A modified release vildagliptin formulation is described in WO 2006/135723.


For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt
25 thereof, reference is thus made to these documents.

A tablet formulation for vildagliptin is expected to be commercially available under the trade name Galvus®. A tablet formulation for vildagliptin/metformin combination is commercially available under the trade name Eucreas®.

- 29 -

- Saxagliptin (BMS-477118) having the structural formula C below is (1S,3S,5S)-2-[(2S)-2-amino-2-(3-hydroxyadamantan-1-yl)acetyl]-2-azabicyclo[3.1.0]hexane-3-carbonitrile, also named (S)-3-hydroxyadamantylglycine-L-*cis*-4,5-methanoprolinenitrile,

5



(C)


Saxagliptin is specifically disclosed in US Patent No. 6,395,767 and in Example 60 of WO 01/68603.

- 10 In one embodiment, saxagliptin is in the form of its HCl salt or its mono-benzoate salt as disclosed in WO 2004/052850. In a further embodiment, saxagliptin is in the form of the free base. In a yet further embodiment, saxagliptin is in the form of the monohydrate of the free base as disclosed in WO 2004/052850. Crystalline forms of the HCl salt and of the free base of saxagliptin are disclosed in WO 2008/131149. A process for preparing saxagliptin is also
- 15 disclosed in WO 2005/106011 and WO 2005/115982. Saxagliptin can be formulated in a tablet as described in WO 2005/117841.

For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

- 20 - Alogliptin (SYR-322) having the structural formula E below is 2-({6-[(3R)-3-aminopiperidin-1-yl]-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl}methyl)benzonitrile

- 30 -




(E)

- Alogliptin is specifically disclosed in US 2005/261271, EP 1586571 and in WO 2005/095381. In one embodiment, alogliptin is in the form of its benzoate salt, its hydrochloride salt or its tosylate salt each as disclosed in WO 2007/035629. A class of this embodiment refers to alogliptin benzoate. Polymorphs of alogliptin benzoate are disclosed in WO 2007/035372. A process for preparing alogliptin is disclosed in WO 2007/112368 and, specifically, in WO 2007/035629. Alogliptin (namely its benzoate salt) can be formulated in a tablet and administered as described in WO 2007/033266. A solid preparation of alogliptin/pioglitazone and its preparation and use is described in WO 2008/093882. A solid preparation of alogliptin/metformin and its preparation and use is described in WO 2009/011451. For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.
- 15 - (2S)-1-[[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethylamino]-acetyl]-pyrrolidine-2-carbonitrile or a pharmaceutically acceptable salt thereof, preferably the mesylate, or (2S)-1-[[1,1,-Dimethyl-3-(4-pyridin-3-yl-imidazol-1-yl)-propylamino]-acetyl]-pyrrolidine-2-carbonitrile or a pharmaceutically acceptable salt thereof:
- 20 These compounds and methods for their preparation are disclosed in WO 03/037327. The mesylate salt of the former compound as well as crystalline polymorphs thereof are disclosed in WO 2006/100181. The fumarate salt of the latter compound as well as crystalline polymorphs thereof are disclosed in WO 2007/071576. These compounds can be formulated in a pharmaceutical composition as described in WO 2007/017423.
- 25 For details, e.g. on a process to manufacture, to formulate or to use these compounds or salts thereof, reference is thus made to these documents.

- 31 -

- (S)-1-((2S,3S,11bS)-2-Amino-9,10-dimethoxy-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-3-yl)-4-fluoromethyl-pyrrolidin-2-one (also named carmegliptin) or a pharmaceutically acceptable salt thereof:



5

This compound and methods for its preparation are disclosed in WO 2005/000848. A process for preparing this compound (specifically its dihydrochloride salt) is also disclosed in WO 2008/031749, WO 2008/031750 and WO 2008/055814. This compound can be formulated in a pharmaceutical composition as described in WO 2007/017423.

10 For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

- (3,3-Difluoropyrrolidin-1-yl)-((2S,4S)-4-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrrolidin-2-yl)methanone (also named gosogliptin) or a pharmaceutically acceptable salt thereof:


15

This compound and methods for its preparation are disclosed in WO 2005/116014 and US 7291618.

For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

20


- (1((3S,4S)-4-amino-1-(4-(3,3-difluoropyrrolidin-1-yl)-1,3,5-triazin-2-yl)pyrrolidin-3-yl)-5,5-difluoropiperidin-2-one or a pharmaceutically acceptable salt thereof:



25 This compound and methods for its preparation are disclosed in WO 2007/148185 and US 20070299076. For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

- 32 -

- (2S,4S)-1-{2-[(3S,1R)-3-(1H-1,2,4-Triazol-1-ylmethyl)cyclopentylamino]-acetyl}-4-fluoropyrrolidine-2-carbonitrile (also named melogliptin) or a pharmaceutically acceptable salt thereof:




5

This compound and methods for its preparation are disclosed in WO 2006/040625 and WO 2008/001195. Specifically claimed salts include the methanesulfonate and p-toluenesulfonate. For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

10


- (R)-2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile or a pharmaceutically acceptable salt thereof:



15 This compound and methods for its preparation and use are disclosed in WO 2005/095381, US 2007060530, WO 2007/033350, WO 2007/035629, WO 2007/074884, WO 2007/112368, WO 2008/033851, WO 2008/114800 and WO 2008/114807. Specifically claimed salts include the succinate (WO 2008/067465), benzoate, benzenesulfonate, p-toluenesulfonate, (R)-mandelate and hydrochloride. For details, e.g. on a process to manufacture, to formulate
20 or to use this compound or a salt thereof, reference is thus made to these documents.

- 5-{(S)-2-[2-((S)-2-Cyano-pyrrolidin-1-yl)-2-oxo-ethylamino]-propyl}-5-(1H-tetrazol-5-yl)-10,11-dihydro-5H-dibenzo[a,d]cycloheptene-2,8-dicarboxylic acid bis-dimethylamide or a pharmaceutically acceptable salt thereof:

- 33 -



This compound and methods for its preparation are disclosed in WO 2006/116157 and US 2006/270701. For details, e.g. on a process to manufacture, to formulate or to use this
 5 compound or a salt thereof, reference is thus made to these documents.

- 3-((2S,4S)-4-[4-(3-Methyl-1-phenyl-1H-pyrazol-5-yl)piperazin-1-yl]pyrrolidin-2-ylcarbonyl)thiazolidine (also named teneligliptin) or a pharmaceutically acceptable salt thereof:

10

This compound and methods for its preparation are disclosed in WO 02/14271. Specific salts are disclosed in WO 2006/088129 and WO 2006/118127 (including hydrochloride, hydrobromide, inter alia). Combination therapy using this compound is described in WO 2006/129785. For details, e.g. on a process to manufacture, to formulate or to use this
 15 compound or a salt thereof, reference is thus made to these documents.

- [(2R)-1-[(3R)-pyrrolidin-3-ylamino]acetyl]pyrrolidin-2-yl]boronic acid (also named dutogliptin) or a pharmaceutically acceptable salt thereof:

20 This compound and methods for its preparation are disclosed in WO 2005/047297, WO 2008/109681 and WO 2009/009751. Specific salts are disclosed in WO 2008/027273 (including citrate, tartrate). A formulation of this compound is described in WO 2008/144730. A formulation of dutogliptin (as its tartrate salt) with metformin is described in WO 2009/091663. For details, e.g. on a process to manufacture, to formulate or to use this
 25 compound or a salt thereof, reference is thus made to these documents.

- (2S,4S)-1-[2-[(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (also named bisegliptin) or a pharmaceutically acceptable salt thereof:

- 34 -

This compound and methods for its preparation are disclosed in WO 2005/075421, US 2008/146818 and WO 2008/114857. For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

5

- 2-({6-[(3R)-3-amino-3-methylpiperidin-1-yl]-1,3-dimethyl-2,4-dioxo-1,2,3,4-tetrahydro-5H-pyrrolo[3,2-d]pyrimidin-5-yl)methyl)-4-fluorobenzonitrile or a pharmaceutically acceptable salt thereof, or 6-[(3R)-3-amino-piperidin-1-yl]-5-(2-chloro-5-fluoro-benzyl)-1,3-dimethyl-1,5-dihydro-pyrrolo[3,2-d]pyrimidine-2,4-dione or a pharmaceutically acceptable salt thereof:

10

These compounds and methods for their preparation are disclosed in WO 2009/084497 and WO 2006/068163, respectively. Combination therapy using the latter of these two compounds is described in WO 2009/128360. For details, e.g. on a process to manufacture, to formulate or to use these compounds or salts thereof, reference is thus made to these documents.

15

- (S)-2-methylpyrazolo[1,5-a]primidine-6-carboxylic acid {2-[(2-cyanopyrrolidin-1-yl)-2-oxoethylamino]-2-methylpropyl}amide (also named anagliptin) or a pharmaceutically acceptable salt:

20

This compound and methods for its preparation are disclosed in WO 2004/067509. Combination therapy using this compound is described in WO 2009/139362. For details, e.g. on a process to manufacture, to formulate or to use this compound or a salt thereof, reference is thus made to these documents.

25

Preferably the DPP-4 inhibitor of this invention is selected from the group (group **G1**) consisting of linagliptin, sitagliptin, vildagliptin, alogliptin, saxagliptin, teneligliptin, anagliptin, gemigliptin and dutogliptin, or a pharmaceutically acceptable salt of one of the hereinmentioned DPP-4 inhibitors, or a prodrug thereof.

30

A particularly preferred DPP-4 inhibitor within the present invention is linagliptin. The term "linagliptin" as employed herein refers to linagliptin or a pharmaceutically acceptable salt thereof, including hydrates and solvates thereof, and crystalline forms thereof, preferably linagliptin refers to 1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-(3-(R)-amino-piperidin-1-yl)-xanthine. Crystalline forms are described in WO 2007/128721. Methods

35

- 35 -

for the manufacture of linagliptin are described in the patent applications WO 2004/018468 and WO 2006/048427 for example. Linagliptin is distinguished from structurally comparable DPP-4 inhibitors, as it combines exceptional potency and a long-lasting effect with favourable pharmacological properties, receptor selectivity and a favourable side-effect profile or bring
5 about unexpected therapeutic advantages or improvements in mono- or dual or triple combination therapy.

For avoidance of any doubt, the disclosure of each of the foregoing and following documents cited above in connection with the specified DPP-4 inhibitors is specifically incorporated
10 herein by reference in its entirety.

A long-acting insulin within the meaning of the present invention includes, without being limited to, any of those long-acting insulins mentioned hereinabove and hereinbelow, preferably subcutaneously active long-acting insulins.

15 Examples of long-acting insulins may include (group **G2**): insulin glargine; insulin detemir; insulin degludec; insulin lispro PEGylated with high molecular weight poly(ethylene glycol) derivatives as disclosed in WO 2009/152128; amidated insulin glargine in the form of Gly^{A21}, Arg^{B31}, Arg^{B32}-NH₂ human insulin; Lys^{B29}(N^ε-lithocholyl-γ-Glu) des(B30) human insulin; N^εB29-
20 ω-carboxypentadecanoyl-γ-amino-butanoyl des(B30) human insulin; amidated insulin analogs as disclosed in WO 2009/087082 (especially one selected from claim 14 therein), or amidated insulin analogs as disclosed in WO 2009/087081 (especially one selected from claim 16 therein).

25 Preferred examples of long-acting insulins of this invention are insulin glargine, insulin detemir and insulin degludec.

Unless otherwise noted, according to this invention it is to be understood that the definitions of the active agents (including the DPP-4 inhibitors and long-acting insulin analogues)
30 mentioned hereinabove and hereinbelow may also contemplate their pharmaceutically acceptable salts, and prodrugs, hydrates, solvates and polymorphic forms thereof. Particularly the terms of the therapeutic agents given herein refer to the respective active drugs. With respect to salts, hydrates and polymorphic forms thereof, particular reference is made to those which are referred to herein.

35

- 36 -

In an embodiment the combinations, compositions, methods and uses according to this invention relate to combinations wherein the DPP-4 inhibitor and the long-acting insulins are preferably selected according to the entries in the Table 1:

5 Table 1:

DPP-4 Inhibitor	long-acting insulin
selected from embodiment A	selected from group G2
selected from embodiment A	insulin glargine
selected from embodiment A	insulin detemir
selected from embodiment A	insulin degludec
selected from embodiment B	selected from group G2
selected from embodiment B	insulin glargine
selected from embodiment B	insulin detemir
selected from embodiment B	insulin degludec
selected from group G1	selected from group G2
selected from group G1	insulin glargine
selected from group G1	insulin detemir
selected from group G1	insulin degludec
selected from group G1 plus metformin	selected from group G2
selected from group G1 plus metformin	insulin glargine
selected from group G1 plus metformin	insulin detemir
selected from group G1 plus metformin	insulin degludec
selected from group G1 plus pioglitazone	selected from group G2
selected from group G1 plus pioglitazone	insulin glargine
selected from group G1 plus pioglitazone	insulin detemir
selected from group G1 plus pioglitazone	insulin degludec
linagliptin	selected from group G2
linagliptin	insulin glargine
linagliptin	insulin detemir
linagliptin	insulin degludec

- 37 -

linagliptin combined with metformin	selected from group G2
linagliptin combined with metformin	insulin glargine
linagliptin combined with metformin	insulin detemir
linagliptin combined with metformin	insulin degludec
linagliptin combined with pioglitazone	selected from group G2
linagliptin combined with pioglitazone	insulin glargine
linagliptin combined with pioglitazone	insulin detemir
linagliptin combined with pioglitazone	insulin degludec

In a particular embodiment (embodiment **E**) the combinations, compositions, methods and uses according to this invention relate to combinations wherein the DPP-4 inhibitor is linagliptin. According to this particular embodiment (embodiment **E**) the long-acting insulin is preferably selected according to the entries E1 to E3 in the Table 2:

Table 2:

Embodiment	long-acting insulin
E1	insulin glargine
E2	insulin detemir
E3	insulin degludec

10

Within this invention it is to be understood that the combinations, compositions or combined uses according to this invention may envisage the simultaneous, sequential or separate administration of the active components or ingredients.

15

In this context, "combination" or "combined" within the meaning of this invention may include, without being limited, fixed and non-fixed (e.g. free) forms (including kits) and uses, such as e.g. the simultaneous, sequential or separate use of the components or ingredients.

- 38 -

The present invention also provides a kit-of-parts or combination therapeutic product comprising

- a) a pharmaceutical composition comprising a DPP-4 inhibitor as defined herein, optionally
- 5 together with one or more pharmaceutically acceptable carriers and/or diluents, and
- b) a pharmaceutical composition comprising a long-acting insulin as defined herein.

The present invention also provides a kit comprising

- a) a DPP-4 inhibitor as defined herein, and
 - 10 b) a long-acting insulin as defined herein,
- and, optionally, instructions directing use of the DPP-4 inhibitor and the long-acting insulin in combination (e.g. simultaneously, separately, sequentially or chronologically staggered), e.g. for a purpose of this invention, such as e.g. for the treatment of type 2 diabetes in a (human) patient.

15

The present invention also provides a pharmaceutical composition or fixed dose combination comprising

- a) a DPP-4 inhibitor as defined herein, and
- b) a long-acting insulin as defined herein,
- 20 and, optionally, one or more pharmaceutically acceptable carriers and/or diluents.

The present invention also provides a transdermal or subcutaneous (injectable) pharmaceutical composition, delivery system or device for systemic use comprising

- a) a DPP-4 inhibitor as defined herein, and, optionally,
 - 25 b) a long-acting insulin as defined herein,
- and, optionally, one or more pharmaceutically acceptable carriers and/or diluents.

- The combined administration of this invention may take place by administering the active components or ingredients together, such as e.g. by administering them simultaneously in
- 30 one single or in two separate formulations or dosage forms. Alternatively, the administration may take place by administering the active components or ingredients sequentially, such as e.g. successively in two separate formulations or dosage forms.

- For the combination therapy of this invention the active components or ingredients may be
- 35 administered separately (which implies that they are formulated separately) or formulated

- 39 -

altogether (which implies that they are formulated in the same preparation or in the same dosage form). Hence, the administration of one element of the combination of the present invention may be prior to, concurrent to, or subsequent to the administration of the other element of the combination. In one embodiment, for the combination therapy according to
5 this invention the DPP-4 inhibitor and the long-acting insulin are administered in different formulations or different dosage forms. In another embodiment, for the combination therapy according to this invention the DPP-4 inhibitor and the long-acting insulin are administered in the same formulation or in the same dosage form.

10 Unless otherwise noted, combination therapy may refer to first line, second line or third line therapy, or initial or add-on combination therapy or replacement therapy.

With respect to embodiment **A**, the methods of synthesis for the DPP-4 inhibitors according to embodiment **A** of this invention are known to the skilled person. Advantageously, the DPP-
15 4 inhibitors according to embodiment **A** of this invention can be prepared using synthetic methods as described in the literature. Thus, for example, purine derivatives of formula (I) can be obtained as described in WO 2002/068420, WO 2004/018468, WO 2005/085246, WO 2006/029769 or WO 2006/048427, the disclosures of which are incorporated herein. Purine derivatives of formula (II) can be obtained as described, for example, in WO
20 2004/050658 or WO 2005/110999, the disclosures of which are incorporated herein. Purine derivatives of formula (III) and (IV) can be obtained as described, for example, in WO 2006/068163, WO 2007/071738 or WO 2008/017670, the disclosures of which are incorporated herein. The preparation of those DPP-4 inhibitors, which are specifically mentioned hereinabove, is disclosed in the publications mentioned in connection therewith.
25 Polymorphous crystal modifications and formulations of particular DPP-4 inhibitors are disclosed in WO 2007/128721 and WO 2007/128724, respectively, the disclosures of which are incorporated herein in their entireties. Formulations of particular DPP-4 inhibitors with metformin or other combination partners are described in WO 2009/121945, the disclosure of which is incorporated herein in its entirety.

30 Typical dosage strengths of the dual fixed combination (tablet) of linagliptin / metformin IR (immediate release) are 2.5/500 mg, 2.5/850 mg and 2.5/1000 mg, which may be administered 1-3 times a day, particularly twice a day.

- 40 -

Typical dosage strengths of the dual fixed combination (tablet) of linagliptin / metformin XR (extended release) are 5/500 mg, 5/1000 mg and 5/1500 mg (each one tablet) or 2.5/500 mg, 2.5/750 mg and 2.5/1000 mg (each two tablets), which may be administered 1-2 times a day, particularly once a day, preferably to be taken in the evening with meal.

5

The present invention further provides a DPP-4 inhibitor as defined herein for use in (add-on or initial) combination therapy with metformin (e.g. in a total daily amount from 500 to 2000 mg metformin hydrochloride, such as e.g. 500 mg, 850 mg or 1000 mg once or twice daily).

- 10 With respect to embodiment **B**, the methods of synthesis for the DPP-4 inhibitors of embodiment **B** are described in the scientific literature and/ or in published patent documents, particularly in those cited herein.

15 The elements of the combination of this invention may be administered by various ways, for example by oral, buccal, sublingual, enterical, parenteral (e.g., transdermal, intramuscular or subcutaneous), inhalative (e.g., liquid or powder inhalation, aerosol), pulmonary, intranasal (e.g. spray), intraperitoneal, vaginal, rectal, or topical routes of administration and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each
20 route of administration.

In a preferred embodiment, the component DPP-4 inhibitor of the combination according to the invention is preferably administered orally. In another preferred embodiment, the component long-acting insulin of the combination is preferably administered by injection
25 (preferably subcutaneously). In another embodiment, the component long-acting insulin of the combination is administered by a transdermal delivery system.

Suitable doses and dosage forms of the DPP-4 inhibitors may be determined by a person skilled in the art and may include those described herein or in the relevant references.

30

For pharmaceutical application in warm-blooded vertebrates, particularly humans, the compounds of this invention are usually used in dosages from 0.001 to 100 mg/kg body weight, preferably at 0.01-15 mg/kg or 0.1-15 mg/kg, in each case 1 to 4 times a day. For this purpose, the compounds, optionally combined with other active substances, may be
35 incorporated together with one or more inert conventional carriers and/or diluents, e.g. with

- 41 -

corn starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, polyvinylpyrrolidone, citric acid, tartaric acid, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, cetylstearyl alcohol, carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof into
5 conventional galenic preparations such as plain or coated tablets, capsules, powders, suspensions or suppositories.

The pharmaceutical compositions according to this invention comprising the DPP-4 inhibitors as defined herein are thus prepared by the skilled person using pharmaceutically acceptable
10 formulation excipients as described in the art and appropriate for the desired route of administration. Examples of such excipients include, without being restricted to diluents, binders, carriers, fillers, lubricants, flow promoters, crystallisation retardants, disintegrants, solubilizers, colorants, pH regulators, surfactants and emulsifiers.

15 Oral preparations or dosage forms of the DPP-4 inhibitor of this invention may be prepared according to known techniques.

Examples of suitable diluents for compounds according to embodiment **A** include cellulose powder, calcium hydrogen phosphate, erythritol, low substituted hydroxypropyl cellulose,
20 mannitol, pregelatinized starch or xylitol.

Examples of suitable lubricants for compounds according to embodiment **A** include talc, polyethyleneglycol, calcium behenate, calcium stearate, hydrogenated castor oil or magnesium stearate.

25 Examples of suitable binders for compounds according to embodiment **A** include copovidone (copolymerisates of vinylpyrrolidone with other vinyl derivatives), hydroxypropyl methylcellulose (HPMC), hydroxypropylcellulose (HPC), polyvinylpyrrolidone (povidone), pregelatinized starch, or low-substituted hydroxypropylcellulose (L-HPC).

30 Examples of suitable disintegrants for compounds according to embodiment **A** include corn starch or croscopovidone.

Suitable methods of preparing pharmaceutical formulations of the DPP-4 inhibitors according
35 to embodiment **A** of the invention are

- 42 -

- direct tableting of the active substance in powder mixtures with suitable tableting excipients;
- granulation with suitable excipients and subsequent mixing with suitable excipients and subsequent tableting as well as film coating; or
- 5 • packing of powder mixtures or granules into capsules.

Suitable granulation methods are

- wet granulation in the intensive mixer followed by fluidised bed drying;
- one-pot granulation;
- 10 • fluidised bed granulation; or
- dry granulation (e.g. by roller compaction) with suitable excipients and subsequent tableting or packing into capsules.

15 An exemplary composition (e.g. tablet core) of a DPP-4 inhibitor according to embodiment **A** of the invention comprises the first diluent mannitol, pregelatinized starch as a second diluent with additional binder properties, the binder copovidone, the disintegrant corn starch, and magnesium stearate as lubricant; wherein copovidone and/or corn starch may be optional.

20 A tablet of a DPP-4 inhibitor according to embodiment **A** of the invention may be film coated, preferably the film coat comprises hydroxypropylmethylcellulose (HPMC), polyethylene glycol (PEG), talc, titanium dioxide and iron oxide (e.g. red and/or yellow).

25 In a further embodiment, the component DPP-4 inhibitor of the combination according to the invention is administered by injection (preferably subcutaneously). In another embodiment, the component long-acting insulin of the combination is preferably administered by injection (preferably subcutaneously) as well. In another embodiment, the component long-acting insulin of the combination is administered by a transdermal delivery system.

30 Injectable formulations of the long-acting insulin and/or the DPP-4 inhibitor of this invention (particularly for subcutaneous use) may be prepared according to known formulation techniques, e.g. using suitable liquid carriers, which usually comprise sterile water, and, optionally, further additives such as e.g. preservatives, pH adjusting agents, buffering agents, isotoning agents, solubility aids and/or tensides or the like, to obtain injectable solutions or suspensions. In addition, injectable formulations may comprise further additives, for example
35 salts, solubility modifying agents or precipitating agents which retard release of the drug(s).

- 43 -

In further addition, injectable insulin formulations may comprise insulin stabilizing agents, such as zinc compounds.

5 In a further embodiment, the component DPP-4 inhibitor of the combination according to the invention is administered by a transdermal delivery system. In another embodiment, the component long-acting insulin of the combination is administered by a transdermal delivery system as well. In another embodiment, the component long-acting insulin of the combination is preferably administered by injection (preferably subcutaneously). In another embodiment, the component long-acting insulin of the combination is administered by
10 (subcutaneous implanted) insulin pellets.

Transdermal formulations (e.g. for transdermal patches or gels) of the long-acting insulin and/or the DPP-4 inhibitor of this invention may be prepared according to known formulation techniques, e.g. using suitable carriers and, optionally, further additives. To facilitate
15 transdermal passage, different methodologies and systems may be used, such as e.g. techniques involving formation of microchannels or micropores in the skin, such as e.g. iontophoresis (based on low-level electrical current), sonophoresis (based on low-frequency ultrasound) or microneedling, or the use of drug-carrier agents (e.g. elastic or lipid vesicles such as transfersomes) or permeation enhancers.

20 For further details on dosage forms, formulations and administration of DPP-4 inhibitors of this invention and/or long-acting insulin of this invention, reference is made to scientific literature and/or published patent documents, particularly to those cited herein.

25 The pharmaceutical compositions (or formulations) may be packaged in a variety of ways. Generally, an article for distribution includes one or more containers that contain the one or more pharmaceutical compositions in an appropriate form. Tablets are typically packed in an appropriate primary package for easy handling, distribution and storage and for assurance of proper stability of the composition at prolonged contact with the environment during storage.
30 Primary containers for tablets may be bottles or blister packs.

A suitable bottle, e.g. for a pharmaceutical composition or combination (tablet) comprising a DPP-4 inhibitor according to embodiment **A** of the invention, may be made from glass or polymer (preferably polypropylene (PP) or high density polyethylene (HD-PE)) and sealed
35 with a screw cap. The screw cap may be provided with a child resistant safety closure (e.g.

- 44 -

press-and-twist closure) for preventing or hampering access to the contents by children. If required (e.g. in regions with high humidity), by the additional use of a desiccant (such as e.g. bentonite clay, molecular sieves, or, preferably, silica gel) the shelf life of the packaged composition can be prolonged.

5

A suitable blister pack, e.g. for a pharmaceutical composition or combination (tablet) comprising a DPP-4 inhibitor according to embodiment **A** of the invention, comprises or is formed of a top foil (which is breachable by the tablets) and a bottom part (which contains pockets for the tablets). The top foil may contain a metallic foil, particularly aluminium or
10 aluminium alloy foil (e.g. having a thickness of 20µm to 45µm, preferably 20µm to 25µm) that is coated with a heat-sealing polymer layer on its inner side (sealing side). The bottom part may contain a multi-layer polymer foil (such as e.g. poly(vinyl chloride) (PVC) coated with poly(vinylidene chloride) (PVDC); or a PVC foil laminated with poly(chlorotrifluoroethylene) (PCTFE)) or a multi-layer polymer-metal-polymer foil (such as e.g. a cold-formable laminated
15 PVC/aluminium/polyamide composition).

To ensure a long storage period especially under hot and wet climate conditions an additional overwrap or pouch made of a multi-layer polymer-metal-polymer foil (e.g. a laminated polyethylene/aluminium/polyester composition) may be used for the blister packs.

20 Supplementary desiccant (such as e.g. bentonite clay, molecular sieves, or, preferably, silica gel) in this pouch package may prolong the shelf life even more under such harsh conditions.

Solutions for injection may be available in typical suitable presentation forms such as vials, cartridges or prefilled (disposable) pens, which may be further packaged.

25

The article may further comprise a label or package insert, which refer to instructions customarily included in commercial packages of therapeutic products, that may contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products. In one embodiment, the label or
30 package inserts indicates that the composition can be used for any of the purposes described herein.

With respect to the first embodiment (embodiment **A**), the dosage typically required of the DPP-4 inhibitors mentioned herein in embodiment **A** when administered intravenously is
35 0.1 mg to 10 mg, preferably 0.25 mg to 5 mg, and when administered orally is 0.5 mg to

- 45 -

100 mg, preferably 2.5 mg to 50 mg or 0.5 mg to 10 mg, more preferably 2.5 mg to 10 mg or 1 mg to 5 mg, in each case 1 to 4 times a day. Thus, e.g. the dosage of 1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyln-1-yl)-8-(3-(R)-amino-piperidin-1-yl)-xanthine when administered orally is 0.5 mg to 10 mg per patient per day, preferably 2.5 mg to 10 mg or 1 mg to 5 mg per patient per day.

A dosage form prepared with a pharmaceutical composition comprising a DPP-4 inhibitor mentioned herein in embodiment **A** contain the active ingredient in a dosage range of 0.1-100 mg. Thus, e.g. particular oral dosage strengths of 1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyln-1-yl)-8-(3-(R)-amino-piperidin-1-yl)-xanthine are 0.5 mg, 1 mg, 2.5 mg, 5 mg and 10 mg.

With respect to the second embodiment (embodiment **B**), the doses of DPP-4 inhibitors mentioned herein in embodiment **B** to be administered to mammals, for example human beings, of, for example, approximately 70 kg body weight, may be generally from about 0.5 mg to about 350 mg, for example from about 10 mg to about 250 mg, preferably 20-200 mg, more preferably 20-100 mg, of the active moiety per person per day, or from about 0.5 mg to about 20 mg, preferably 2.5-10 mg, per person per day, divided preferably into 1 to 4 single doses which may, for example, be of the same size. Single oral dosage strengths comprise, for example, 10, 25, 40, 50, 75, 100, 150 and 200 mg of the DPP-4 inhibitor active moiety.

An oral dosage strength of the DPP-4 inhibitor sitagliptin is usually between 25 and 200 mg of the active moiety. A recommended dose of sitagliptin is 100 mg calculated for the active moiety (free base anhydrate) once daily. Unit dosage strengths of sitagliptin free base anhydrate (active moiety) are 25, 50, 75, 100, 150 and 200 mg. Particular unit dosage strengths of sitagliptin (e.g. per tablet) are 25, 50 and 100 mg. An equivalent amount of sitagliptin phosphate monohydrate to the sitagliptin free base anhydrate is used in the pharmaceutical compositions, namely, 32.13, 64.25, 96.38, 128.5, 192.75, and 257 mg, respectively. Adjusted dosages of 25 and 50 mg sitagliptin are used for patients with renal failure. Typical dosage strengths of the dual combination of sitagliptin / metformin are 50/500 mg and 50/1000 mg.

An oral dosage range of the DPP-4 inhibitor vildagliptin is usually between 10 and 150 mg daily, in particular between 25 and 150 mg, 25 and 100 mg or 25 and 50 mg or 50 and 100 mg daily. Particular examples of daily oral dosage are 25, 30, 35, 45, 50, 55, 60, 80, 100 or

- 46 -

150 mg. In a more particular aspect, the daily administration of vildagliptin may be between 25 and 150 mg or between 50 and 100 mg. In another more particular aspect, the daily administration of vildagliptin may be 50 or 100 mg. The application of the active ingredient may occur up to three times a day, preferably one or two times a day. Particular dosage
5 strengths are 50 mg or 100 mg vildagliptin. Typical dosage strengths of the dual combination of vildagliptin / metformin are 50/850 mg and 50/1000 mg.

Alogliptin may be administered to a patient at an oral daily dose of between 5 mg/day and 250 mg/day, optionally between 10 mg and 200 mg, optionally between 10 mg and 150 mg,
10 and optionally between 10 mg and 100 mg of alogliptin (in each instance based on the molecular weight of the free base form of alogliptin). Thus, specific oral dosage amounts that may be used include, but are not limited to 10 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg and 100 mg of alogliptin per day. Alogliptin may be administered in its free base form or as a pharmaceutically acceptable salt.

15 Saxagliptin may be administered to a patient at an oral daily dose of between 2.5 mg/day and 100 mg/day, optionally between 2.5 mg and 50 mg. Specific oral dosage amounts that may be used include, but are not limited to 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, 30 mg , 40 mg, 50 mg and 100 mg of saxagliptin per day. Typical dosage strengths of the dual
20 combination of saxagliptin / metformin are 2.5/500 mg and 2.5/1000 mg.

A special embodiment of the DPP-4 inhibitors of this invention refers to those orally administered DPP-4 inhibitors which are therapeutically efficacious at low dose levels, e.g. at oral dose levels < 100 mg or < 70 mg per patient per day, preferably < 50 mg, more
25 preferably < 30 mg or < 20 mg, even more preferably from 1 mg to 10 mg, particularly from 1 mg to 5 mg (more particularly 5 mg), per patient per day (if required, divided into 1 to 4 single doses, particularly 1 or 2 single doses, which may be of the same size, preferentially, administered orally once- or twice daily (more preferentially once-daily), advantageously, administered at any time of day, with or without food. Thus, for example, the daily oral
30 amount 5 mg BI 1356 can be given in an once daily dosing regimen (i.e. 5 mg BI 1356 once daily) or in a twice daily dosing regimen (i.e. 2.5 mg BI 1356 twice daily), at any time of day, with or without food.

The long-acting insulin is typically administered by subcutaneous injection, e.g. ranging from
35 twice daily, once daily to once weekly injection. Suitable doses and dosage forms of the long-

- 47 -

acting insulin may be determined by a person skilled in the art. Blood glucose monitoring is essential in all patients receiving insulin therapy.

For example, insulin glargine (Lantus) is administered subcutaneously once a day. Lantus
5 may be administered at any time during the day, but at the same time every day. The dose of Lantus is individualized based on clinical response. The recommended starting dose of Lantus in patients with type 2 diabetes who are not currently treated with insulin is 10 units (or 0.2 Units/kg) once daily, which should subsequently be adjusted to the patient's needs.

10 Insulin detemir (Levemir) is administered subcutaneously once or twice a day. For patients treated with Levemir once daily, the dose is preferably administered with the evening meal or at bedtime. For patients who require twice-daily dosing, the evening dose can be administered either with evening meal, at bedtime, or 12 hours after the
15 morning dose. The dose of Levemir is individualized based on clinical response. For insulin-naïve patients with type 2 diabetes who are inadequately controlled on oral antidiabetic drugs, Levemir should be started at a dose of 0.1 to 0.2 Units/kg once-daily in the evening or 10 units once- or twice-daily, and the dose adjusted to achieve glycemic targets.

20 The dosage of the active ingredients in the combinations and compositions in accordance with the present invention may be varied, although the amount of the active ingredients shall be such that a suitable dosage form is obtained. Hence, the selected dosage and the selected dosage form shall depend on the desired therapeutic effect, the route of
25 administration and the duration of the treatment. Suitable dosage ranges for the combination are from the maximal tolerated dose for the single agent to lower doses, e.g. to one tenth of the maximal tolerated dose.

A particularly preferred DPP-4 inhibitor to be emphasized within the meaning of this invention
30 is 1-[(4-methyl-quinazolin-2-yl)methyl]-3-methyl-7-(2-butyn-1-yl)-8-(3-(*R*)-amino-piperidin-1-yl)-xanthine (also known as BI 1356 or linagliptin). BI 1356 exhibits high potency, 24h duration of action, and a wide therapeutic window. In patients with type 2 diabetes receiving multiple oral doses of 1, 2.5, 5 or 10 mg of BI 1356 once daily for 12 days, BI 1356 shows favourable pharmacodynamic and pharmacokinetic profile (see e.g. Table 3 below) with rapid
35 attainment of steady state (e.g. reaching steady state plasma levels (> 90% of the pre-dose

- 48 -

plasma concentration on Day 13) between second and fifth day of treatment in all dose groups), little accumulation (e.g. with a mean accumulation ratio $R_{A,AUC} \leq 1.4$ with doses above 1 mg) and preserving a long-lasting effect on DPP-4 inhibition (e.g. with almost complete (> 90%) DPP-4 inhibition at the 5 mg and 10 mg dose levels, i.e. 92.3 and 97.3% inhibition at steady state, respectively, and > 80% inhibition over a 24h interval after drug intake), as well as significant decrease in 2h postprandial blood glucose excursions by ≥ 80 % (already on Day 1) in doses ≥ 2.5 mg, and with the cumulative amount of unchanged parent compound excreted in urine on Day 1 being below 1% of the administered dose and increasing to not more than about 3-6% on Day 12 (renal clearance $CL_{R,ss}$ is from about 14 to about 70 mL/min for the administered oral doses, e.g. for the 5 mg dose renal clearance is about 70 mL/min). In people with type 2 diabetes BI 1356 shows a placebo-like safety and tolerability. With low doses of about ≥ 5 mg, BI 1356 acts as a true once-daily oral drug with a full 24 h duration of DPP-4 inhibition. At therapeutic oral dose levels, BI 1356 is mainly excreted via the liver and only to a minor extent (about < 7% of the administered oral dose) via the kidney. BI 1356 is primarily excreted unchanged via the bile. The fraction of BI 1356 eliminated via the kidneys increases only very slightly over time and with increasing dose, so that there will likely be no need to modify the dose of BI 1356 based on the patients' renal function. The non-renal elimination of BI 1356 in combination with its low accumulation potential and broad safety margin may be of significant benefit in a patient population that has a high prevalence of renal insufficiency and diabetic nephropathy.

Table 3: Geometric mean (gMean) and geometric coefficient of variation (gCV) of pharmacokinetic parameters of BI 1356 at steady state (Day 12)

Parameter	1 mg gMean (gCV)	2.5 mg gMean (gCV)	5 mg gMean (gCV)	10 mg gMean (gCV)
AUC_{0-24} [nmol·h/L]	40.2 (39.7)	85.3 (22.7)	118 (16.0)	161 (15.7)
$AUC_{T,ss}$ [nmol·h/L]	81.7 (28.3)	117 (16.3)	158 (10.1)	190 (17.4)
C_{max} [nmol/L]	3.13 (43.2)	5.25 (24.5)	8.32 (42.4)	9.69 (29.8)
$C_{max,ss}$ [nmol/L]	4.53 (29.0)	6.58 (23.0)	11.1 (21.7)	13.6 (29.6)
t_{max}^* [h]	1.50 [1.00 –	2.00 [1.00 –	1.75 [0.92 –	2.00 [1.50 –

- 49 -

	3.00]	3.00]	6.02]	6.00]
$t_{\max,ss}^*$ [h]	1.48 [1.00 – 3.00]	1.42 [1.00 – 3.00]	1.53 [1.00 – 3.00]	1.34 [0.50 – 3.00]
$T_{1/2,ss}$ [h]	121 (21.3)	113 (10.2)	131 (17.4)	130 (11.7)
Accumulation $t_{1/2}$ [h]	23.9 (44.0)	12.5 (18.2)	11.4 (37.4)	8.59 (81.2)
$R_{A,Cmax}$	1.44 (25.6)	1.25 (10.6)	1.33 (30.0)	1.40 (47.7)
$R_{A,AUC}$	2.03 (30.7)	1.37 (8.2)	1.33 (15.0)	1.18 (23.4)
fe_{0-24} [%]	NC	0.139 (51.2)	0.453 (125)	0.919 (115)
$fe_{r,ss}$ [%]	3.34 (38.3)	3.06 (45.1)	6.27 (42.2)	3.22 (34.2)
$CL_{R,ss}$ [mL/min]	14.0 (24.2)	23.1 (39.3)	70 (35.0)	59.5 (22.5)

* median and range [min-max]

NC not calculated as most values below lower limit of quantification

- As different metabolic functional disorders often occur simultaneously, it is quite often indicated to combine a number of different active principles with one another. Thus, depending on the functional disorders diagnosed, improved treatment outcomes may be
- 5 obtained if a DPP-4 inhibitor is combined with active substances customary for the respective disorders, such as e.g. one or more active substances selected from among the other antidiabetic substances, especially active substances that lower the blood sugar level or the lipid level in the blood, raise the HDL level in the blood, lower blood pressure or are indicated in the treatment of atherosclerosis or obesity.
- 10 The DPP-4 inhibitors mentioned above – besides their use in mono-therapy – may also be used in conjunction with other active substances, by means of which improved treatment results can be obtained. Such a combined treatment may be given as a free combination of the substances or in the form of a fixed combination, for example in a tablet or capsule.
- 15 Pharmaceutical formulations of the combination partner needed for this may either be obtained commercially as pharmaceutical compositions or may be formulated by the skilled man using conventional methods. The active substances which may be obtained commercially as pharmaceutical compositions are described in numerous places in the prior

- 50 -

art, for example in the list of drugs that appears annually, the "Rote Liste ®" of the federal association of the pharmaceutical industry, or in the annually updated compilation of manufacturers' information on prescription drugs known as the "Physicians' Desk Reference".

5

Examples of antidiabetic combination partners are metformin; sulphonylureas such as glibenclamide, tolbutamide, glimepiride, glipizide, gliquidone, glibomuride and gliclazide; nateglinide; repaglinide; mitiglinide; thiazolidinediones such as rosiglitazone and pioglitazone; PPAR gamma modulators such as metaglitazones; PPAR-gamma agonists such as e.g. rivoglitazone, mitoglitazone, INT-131 and balaglitazone; PPAR-gamma antagonists; PPAR-gamma/alpha modulators such as tesaglitazar, muraglitazar, aleglitazar, indeglitazar and KRP297; PPAR-gamma/alpha/delta modulators such as e.g. lobeglitazone; AMPK-activators such as AICAR; acetyl-CoA carboxylase (ACC1 and ACC2) inhibitors; diacylglycerol-acetyltransferase (DGAT) inhibitors; pancreatic beta cell GCRP agonists such as SMT3-receptor-agonists and GPR119, such as the GPR119 agonists 5-ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl]-piperidin-1-yl}-pyrimidine or 5-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-ylmethoxy]-2-(4-methanesulfonyl-phenyl)-pyridine; 11 β -HSD-inhibitors; FGF19 agonists or analogues; alpha-glucosidase blockers such as acarbose, voglibose and miglitol; alpha2-antagonists; insulin and insulin analogues such as human insulin, insulin lispro, insulin glulisin, r-DNA-insulinaspart, NPH insulin, insulin detemir, insulin degludec, insulin tregopil, insulin zinc suspension and insulin glargin; Gastric inhibitory Peptide (GIP); amylin and amylin analogues (e.g. pramlintide or daivalintide); GLP-1 and GLP-1 analogues such as Exendin-4, e.g. exenatide, exenatide LAR, liraglutide, taspoglutide, lixisenatide (AVE-0010), LY-2428757 (a PEGylated version of GLP-1), 25 dulaglutide (LY-2189265), semaglutide or albiglutide; SGLT2-inhibitors such as e.g. dapagliflozin, sergliflozin (KGT-1251), atigliflozin, canagliflozin, ipragliflozin, luseogliflozin or tofogliflozin; inhibitors of protein tyrosine-phosphatase (e.g. trodusquemine); inhibitors of glucose-6-phosphatase; fructose-1,6-bisphosphatase modulators; glycogen phosphorylase modulators; glucagon receptor antagonists; phosphoenolpyruvatecarboxykinase (PEPCK) 30 inhibitors; pyruvate dehydrogenasekinase (PDK) inhibitors; inhibitors of tyrosine-kinases (50 mg to 600 mg) such as PDGF-receptor-kinase (cf. EP-A-564409, WO 98/35958, US 5093330, WO 2004/005281, and WO 2006/041976) or of serine/threonine kinases; glucokinase/regulatory protein modulators incl. glucokinase activators; glycogen synthase kinase inhibitors; inhibitors of the SH2-domain-containing inositol 5-phosphatase type 2 35 (SHIP2) ; IKK inhibitors such as high-dose salicylate; JNK1 inhibitors; protein kinase C-theta

- 51 -

- inhibitors; beta 3 agonists such as ritobegron, YM 178, solabegron, talibegron, N-5984, GRC-1087, rafabegron, FMP825; aldosereductase inhibitors such as AS 3201, zenarestat, fidarestat, epalrestat, ranirestat, NZ-314, CP-744809, and CT-112; SGLT-1 or SGLT-2 inhibitors; KV 1.3 channel inhibitors; GPR40 modulators such as e.g. [(3S)-6-({2',6'-dimethyl-4'-[3-(methylsulfonyl)propoxy]biphenyl-3-yl}methoxy)-2,3-dihydro-1-benzofuran-3-yl]acetic acid; SCD-1 inhibitors; CCR-2 antagonists; dopamine receptor agonists (bromocriptine mesylate [Cycloset]); 4-(3-(2,6-dimethylbenzyloxy)phenyl)-4-oxobutanoic acid; sirtuin stimulants; and other DPP IV inhibitors.
- 10 Metformin is usually given in doses varying from about 500 mg to 2000 mg up to 2500 mg per day using various dosing regimens from about 100 mg to 500 mg or 200 mg to 850 mg (1-3 times a day), or about 300 mg to 1000 mg once or twice a day, or delayed-release metformin in doses of about 100 mg to 1000 mg or preferably 500 mg to 1000 mg once or twice a day or about 500 mg to 2000 mg once a day. Particular dosage strengths may be
- 15 250, 500, 625, 750, 850 and 1000 mg of metformin hydrochloride.

- For children 10 to 16 years of age, the recommended starting dose of metformin is 500 mg given once daily. If this dose fails to produce adequate results, the dose may be increased to 500 mg twice daily. Further increases may be made in increments of 500 mg weekly to a
- 20 maximum daily dose of 2000 mg, given in divided doses (e.g. 2 or 3 divided doses). Metformin may be administered with food to decrease nausea.

A dosage of pioglitazone is usually of about 1-10 mg, 15 mg, 30 mg, or 45 mg once a day.

- 25 Rosiglitazone is usually given in doses from 4 to 8 mg once (or divided twice) a day (typical dosage strengths are 2, 4 and 8 mg).

- Glibenclamide (glyburide) is usually given in doses from 2.5-5 to 20 mg once (or divided twice) a day (typical dosage strengths are 1.25, 2.5 and 5 mg), or micronized glibenclamide
- 30 in doses from 0.75-3 to 12 mg once (or divided twice) a day (typical dosage strengths are 1.5, 3, 4.5 and 6 mg).

- Glipizide is usually given in doses from 2.5 to 10-20 mg once (or up to 40 mg divided twice) a day (typical dosage strengths are 5 and 10 mg), or extended-release glibenclamide in doses
- 35 from 5 to 10 mg (up to 20 mg) once a day (typical dosage strengths are 2.5, 5 and 10 mg).

- 52 -

Glimepiride is usually given in doses from 1-2 to 4 mg (up to 8 mg) once a day (typical dosage strengths are 1, 2 and 4 mg).

- 5 A dual combination of glibenclamide/metformin is usually given in doses from 1.25/250 once daily to 10/1000 mg twice daily. (typical dosage strengths are 1.25/250, 2.5/500 and 5/500 mg).

A dual combination of glipizide/metformin is usually given in doses from 2.5/250 to 10/1000 mg twice daily (typical dosage strengths are 2.5/250, 2.5/500 and 5/500 mg).

- 10 A dual combination of glimepiride/metformin is usually given in doses from 1/250 to 4/1000 mg twice daily.

A dual combination of rosiglitazone/glimepiride is usually given in doses from 4/1 once or twice daily to 4/2 mg twice daily (typical dosage strengths are 4/1, 4/2, 4/4, 8/2 and 8/4 mg).

- 15 A dual combination of pioglitazone/glimepiride is usually given in doses from 30/2 to 30/4 mg once daily (typical dosage strengths are 30/4 and 45/4 mg).

A dual combination of rosiglitazone/metformin is usually given in doses from 1/500 to 4/1000 mg twice daily (typical dosage strengths are 1/500, 2/500, 4/500, 2/1000 and 4/1000 mg).

A dual combination of pioglitazone/metformin is usually given in doses from 15/500 once or twice daily to 15/850 mg thrice daily (typical dosage strengths are 15/500 and 15/850 mg).

20

The non-sulphonylurea insulin secretagogue nateglinide is usually given in doses from 60 to 120 mg with meals (up to 360 mg/day, typical dosage strengths are 60 and 120 mg);

repaglinide is usually given in doses from 0.5 to 4 mg with meals (up to 16 mg/day, typical dosage strengths are 0.5, 1 and 2 mg). A dual combination of repaglinide/metformin is

- 25 available in dosage strengths of 1/500 and 2/850 mg.

Acarbose is usually given in doses from 25 to 100 mg with meals. Miglitol is usually given in doses from 25 to 100 mg with meals.

- 30 Examples of combination partners that lower the lipid level in the blood are HMG-CoA-reductase inhibitors such as simvastatin, atorvastatin, lovastatin, fluvastatin, pravastatin, pitavastatin and rosuvastatin; fibrates such as bezafibrate, fenofibrate, clofibrate, gemfibrozil, etofibrate and etofyllinclofibrate; nicotinic acid and the derivatives thereof such as acipimox; PPAR-alpha agonists; PPAR-delta agonists such as e.g. {4-[(R)-2-ethoxy-3-(4-trifluoromethyl-phenoxy)-propylsulfanyl]-2-methyl-phenoxy}-acetic acid; inhibitors of acyl-
- 35

- 53 -

coenzyme A:cholesterolacyltransferase (ACAT; EC 2.3.1.26) such as avasimibe; cholesterol resorption inhibitors such as ezetimib; substances that bind to bile acid, such as cholestyramine, colestipol and colesevelam; inhibitors of bile acid transport; HDL modulating active substances such as D4F, reverse D4F, LXR modulating active substances and FXR modulating active substances; CETP inhibitors such as torcetrapib, JTT-705 (dalcetrapib) or compound 12 from WO 2007/005572 (anacetrapib); LDL receptor modulators; MTP inhibitors (e.g. lomitapide); and ApoB100 antisense RNA.

10 A dosage of atorvastatin is usually from 1 mg to 40 mg or 10 mg to 80 mg once a day.

Examples of combination partners that lower blood pressure are beta-blockers such as atenolol, bisoprolol, celiprolol, metoprolol and carvedilol; diuretics such as hydrochlorothiazide, chlortalidon, xipamide, furosemide, piretanide, torasemide, spironolactone, eplerenone, amiloride and triamterene; calcium channel blockers such as amlodipine, nifedipine, nitrendipine, nisoldipine, nicardipine, felodipine, lacidipine, lercanipidine, manidipine, isradipine, nilvadipine, verapamil, gallopamil and diltiazem; ACE inhibitors such as ramipril, lisinopril, cilazapril, quinapril, captopril, enalapril, benazepril, perindopril, fosinopril and trandolapril; as well as angiotensin II receptor blockers (ARBs) such as telmisartan, candesartan, valsartan, losartan, irbesartan, olmesartan, azilsartan and eprosartan.

A dosage of telmisartan is usually from 20 mg to 320 mg or 40 mg to 160 mg per day.

25 Examples of combination partners which increase the HDL level in the blood are Cholesteryl Ester Transfer Protein (CETP) inhibitors; inhibitors of endothelial lipase; regulators of ABC1; LXRalpha antagonists; LXRbeta agonists; PPAR-delta agonists; LXRalpha/beta regulators, and substances that increase the expression and/or plasma concentration of apolipoprotein A-I.

30 Examples of combination partners for the treatment of obesity are sibutramine; tetrahydrolipstatin (orlistat); alizyme (cetilistat); dexfenfluramine; axokine; cannabinoid receptor 1 antagonists such as the CB1 antagonist rimonabant; MCH-1 receptor antagonists; MC4 receptor agonists; NPY5 as well as NPY2 antagonists (e.g. velneperit); beta3-AR agonists such as SB-418790 and AD-9677; 5HT2c receptor agonists such as APD 356 (lorcaserin); myostatin inhibitors; Acrp30 and adiponectin; steroyl CoA desaturase (SCD1)

- 54 -

inhibitors; fatty acid synthase (FAS) inhibitors; CCK receptor agonists; Ghrelin receptor modulators; Pyy 3-36; orexin receptor antagonists; and tesofensine; as well as the dual combinations bupropion/naltrexone, bupropion/zonisamide, topiramate/phentermine and pramlintide/metreleptin.

5

Examples of combination partners for the treatment of atherosclerosis are phospholipase A2 inhibitors; inhibitors of tyrosine-kinases (50 mg to 600 mg) such as PDGF-receptor-kinase (cf. EP-A-564409, WO 98/35958, US 5093330, WO 2004/005281, and WO 2006/041976); oxLDL antibodies and oxLDL vaccines; apoA-1 Milano; ASA; and VCAM-1 inhibitors.

10

The present invention is not to be limited in scope by the specific embodiments described herein. Various modifications of the invention in addition to those described herein may become apparent to those skilled in the art from the present disclosure. Such modifications are intended to fall within the scope of the appended claims.

15

All patent applications cited herein are hereby incorporated by reference in their entireties.

Further embodiments, features and advantages of the present invention may become apparent from the following examples. The following examples serve to illustrate, by way of example, the principles of the invention without restricting it.

20

- 55 -

Examples

Linagliptin s.c. dosing and its DPP-4 inhibition in plasma

- 5 Linagliptin subcutaneous (s.c.) dosing and DPP-4 inhibition in plasma can be comparable in efficacy and duration of action to oral dosing, which may make it suitable for use in fixed combination with s.c. insulin:

Male ZDF rats (n=5) have been treated with different concentrations of BI 1356 in a
10 subcutaneous (s.c.) administration regimen (0.001 mg/kg, 0.01 mg/kg, 0.1 mg/kg and 1 mg/kg in 0.5 ml/kg NaCl solution) in comparison to 3 mg/kg p.o. (in 0.5% Natrosol, 5 ml/kg volume of application).

DPP-4 activity in EDTA plasma was detected 1, 3, 5, 7, 24, 31, 48, 72h following drug
administration (blood was taken by venous puncture under isofluran anesthesia from the
15 vena sublingualis).

Doses of BI 1356 from 0.01 mg/kg (s.c. administered) on demonstrated significant inhibition
of DPP-4 activity compared to control. The dose of 0.1 mg/kg and 1 mg/kg (s.c.
administration) of BI 1356 had a persistent DPP-4 inhibition of more than 64% over 7h. The 1
mg/kg s.c. dose was comparable in efficacy and duration of action to the 3 mg/kg oral dose.

20

Fig. 1: DPP-4 activity in plasma after linagliptin s.c. dosing.

25

30

35

- 56 -

Effect of linagliptin on body weight total body fat, liver fat and intramyocellular fat

In a further study the efficacy of chronic treatment with linagliptin on body weight, total body fat, intra-myocellular fat, and hepatic fat in a non-diabetic model of diet induced obesity (DIO) in comparison to the appetite suppressant sibutramine is investigated:

Rats are fed a high-fat diet for 3 months and received either vehicle, linagliptin (10 mg/kg), or sibutramine (5 mg/kg) for 6 additional weeks, while continuing the high-fat diet. Magnetic resonance spectroscopy (MRS) analysis of total body fat, muscle fat, and liver fat is performed before treatment and at the end of the study. Sibutramine causes a significant reduction of body weight (-12%) versus control, whereas linagliptin has no significant effect (-3%). Total body fat is also significantly reduced by sibutramine (-12%), whereas linagliptin-treated animals show no significant reduction (-5%). However, linagliptin and sibutramine result both in a potent reduction of intramyocellular fat (-24% and -34%, respectively). In addition, treatment with linagliptin results in a profound decrease of hepatic fat (-39%), whereas the effect of sibutramine (-30%) does not reach significance (see Table 4). Thus, linagliptin is weight neutral but improves intra-myocellular and hepatic lipid accumulation.

Table 4: Effect of linagliptin on body weight total body fat, liver fat and intramyocellular fat

	Body weight		Total body fat		Liver fat		Intra-myocellular fat	
	% contr.	% baseli.	% contr.	% baseli.	% contr.	% baseli.	% contr.	% baseli.
Control	-	+15% p=0.016	-	+11% p=0.001	-	+27% p=0.09	-	+23% p=0.49
Linagliptin	-3% p=0.56	+12% p=0.001	-5% p=0.27	+5% p=0.06	-39% p=0.022	-30% p=0.05	-36% p=0.14	-24% p=0.039
Sibutramine	-12% p=0.018	+1% p=0.64	-12% p=0.008	-0.4% p=0.86	-30% p=0.13	-29% p=0.12	-55% p=0.037	-34% p=0.007

In conclusion, linagliptin treatment provokes a potent reduction of intramyocellular lipids and hepatic fat, which are both independent of weight loss. The effects of sibutramine on muscular and hepatic fat are attributed mainly to the known weight reduction induced by this compound.

- 57 -

Delaying onset of diabetes and preserving beta-cell function in non-obese type-1 diabetes:

- Though reduced pancreatic T-cell migration and altered cytokine production is considered
- 5 important players for the onset of insulinitis the exact mechanism and effects on the pancreatic cell pool is still incompletely understood. In an attempt to evaluate the effect of linagliptin on pancreatic inflammation and beta-cell mass it is examined the progression of diabetes in the non-obese-diabetic (NOD) mice over a 60 day experimental period coupled with terminal stereological assessment of cellular pancreatic changes.
- 10 Sixty female NOD mice (10 weeks of age) sre included in the study and fed a normal chow diet or a diet containing linagliptin (0.083 g linagliptin/kg chow; corresponding to 3-10 mg/kg, p.o) throughout the study period. Bi-weekly plasma samples are obtained to determine onset of diabetes (BG >11 mmol/l). At termination, the pancreata are removed and a terminal blood sample is obtained for assessment of active GLP-1 levels.
- 15 At the end of the study period the incidence of diabetes is significantly decreased in linagliptin-treated mice (9 out of 30 mice) compared with the control group (18 of 30 mice, $p=0.021$). The subsequent stereological assessment of beta-cell mass (identified by insulin immunoreactivity) demonstrates a significantly larger beta cell mass (vehicle 0.18 ± 0.03 mg; linagliptin 0.48 ± 0.09 mg, $p<0.01$) and total islet mass (vehicle 0.40 ± 0.04 mg; linagliptin
- 20 0.70 ± 0.09 mg, $p<0.01$) in linagliptin treated mice. There is a tendency for linagliptin to reduce peri-islet infiltrating lymphocytes (1.06 ± 0.15 ; lina 0.79 ± 0.12 mg, $p=0.17$). As expected active plasma GLP-1 are higher at termination in linagliptin treated mice.
- In summary, the data demonstrate that linagliptin is able to delay the onset of diabetes in a type-1 diabetic model (NOD mouse). The pronounced beta-cell sparing effects which can be
- 25 observed in this animal model indicate that such DPP-4 inhibition not only protects beta-cells by increasing active GLP-1 levels, but may also exerts direct or indirect anti-inflammatory actions.

- 58 -

Claims

- 1) A combination comprising
5 a long-acting insulin and a DPP-4 inhibitor which is preferably linagliptin,
for separate, sequential or simultaneous therapeutic use of the active components.
- 2) The combination according to claim 1, wherein the long-acting insulin is basal insulin
preferably selected from insulin detemir, insulin glargine and insulin degludec.
10
- 3) The combination according to claim 1 or 2, wherein the long-acting insulin and the DPP-4
inhibitor are present each in separate dosage forms.
- 4) The combination according to claim 1 or 3, wherein the long-acting insulin and the DPP-4
15 inhibitor are present in the same dosage form.
- 5) The combination according to any one of claims 1 to 4, wherein the DPP-4 inhibitor is for
peroral administration.
- 20 6) The combination according to any one of claims 1 to 4, wherein the DPP-4 inhibitor is for
subcutaneous or transdermal administration.
- 7) The combination according to any one of claims 1 to 6, wherein the long-acting insulin is
for subcutaneous or transdermal administration.
25
- 8) The combination according to any one of claims 1 to 7, wherein both active components
are for subcutaneous administration.
- 9) The combination according to any one of claims 1 to 8, wherein the long-acting insulin and
30 the DPP-4 inhibitor are comprised in a pharmaceutical composition for subcutaneous
injection administration.
- 10) The combination according to any one of the claims 1 to 9, further comprising one or
more other therapeutic agents, such as e.g. metformin or pioglitazone.
35

- 59 -

11) The combination according to any one of the claims 1 to 10 for use in one or more of the following methods:

- 5 - preventing, slowing the progression of, delaying or treating a metabolic disorder or disease, such as e.g. type 1 diabetes mellitus, type 2 diabetes mellitus, impaired glucose tolerance (IGT), impaired fasting blood glucose (IFG), hyperglycemia, postprandial hyperglycemia, postabsorptive hyperglycemia, overweight, obesity, dyslipidemia, hyperlipidemia, hypercholesterolemia, hypertriglyceridemia, hypertension, atherosclerosis, endothelial dysfunction, osteoporosis, chronic systemic inflammation, non alcoholic fatty liver disease (NAFLD), retinopathy, neuropathy, nephropathy, polycystic ovarian
- 10 syndrome, and/or metabolic syndrome;
- improving and/or maintaining glycemic control and/or for reducing of fasting plasma glucose, of postprandial plasma glucose, of postabsorptive plasma glucose and/or of glycosylated hemoglobin HbA1c;
- 15 - preventing, slowing, delaying or reversing progression from pre-diabetes, impaired glucose tolerance (IGT), impaired fasting blood glucose (IFG), insulin resistance and/or from metabolic syndrome to type 2 diabetes mellitus;
- preventing, reducing the risk of, slowing the progression of, delaying or treating of complications of diabetes mellitus such as micro- and macrovascular diseases, such as nephropathy, micro- or macroalbuminuria, proteinuria, retinopathy, cataracts, neuropathy,
- 20 learning or memory impairment, neurodegenerative or cognitive disorders, cardio- or cerebrovascular diseases, tissue ischaemia, diabetic foot or ulcer, atherosclerosis, hypertension, endothelial dysfunction, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable angina pectoris, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders, vascular restenosis, and/or stroke;
- 25 - reducing body weight and/or body fat and/or liver fat and/or intra-myocellular fat or preventing an increase in body weight and/or body fat and/or liver fat and/or intra-myocellular fat or facilitating a reduction in body weight and/or body fat and/or liver fat and/or intra-myocellular fat;
- preventing, slowing, delaying or treating the degeneration of pancreatic beta cells and/or the decline of the functionality of pancreatic beta cells and/or for improving, preserving and/or restoring the functionality of pancreatic beta cells and/or stimulating and/or restoring or protecting the functionality of pancreatic insulin secretion;
- 30 - preventing, slowing, delaying or treating non alcoholic fatty liver disease (NAFLD) including hepatic steatosis, non-alcoholic steatohepatitis (NASH) and/or liver fibrosis
- 35 (such as e.g. preventing, slowing the progression, delaying, attenuating, treating or

- 60 -

reversing hepatic steatosis, (hepatic) inflammation and/or an abnormal accumulation of liver fat);

- preventing, slowing the progression of, delaying or treating type 2 diabetes with failure to conventional antidiabetic mono- or combination therapy;
- 5 - achieving a reduction in the dose of conventional antidiabetic medication required for adequate therapeutic effect;
- reducing the risk for adverse effects associated with conventional antidiabetic medication (e.g. hypoglycemia or weight gain); and/or
- maintaining and/or improving the insulin sensitivity and/or for treating or preventing
- 10 hyperinsulinemia and/or insulin resistance.

12) The combination according to any one of the claims 1 to 11 for use in a method of improving and/or maintaining control of fasted and post-prandial glucose levels in patients in need thereof, preferably type 2 diabetes patients.

15

13) The combination according to any one of the claims 1 to 12 for use in a method of preventing, decreasing, reducing the risk of, slowing the progression of, delaying the onset of, attenuating or treating complications of diabetes, such as e.g. micro- or macrovascular diabetic complications or angiopathies.

20

14) The combination according to any one of the claims 1 to 13 for therapeutic use in patients, preferably type 2 diabetes patients, with or at risk of renal impairment or renal complications, such as e.g. diabetic nephropathy.

25 15) The combination according to any one of the claims 1 to 14 for use in treating or preventing patients diagnosed with type 1 diabetes, LADA or type 2 diabetes.

16) The combination according to any one of the claims 1 to 15 for use in a method of achieving a reduction in the dose of insulin required for adequate therapeutic effect.

30

17) The combination according to any one of the claims 1 to 16 for use in a method of reducing hypoglycemia and/or weight gain associated with insulin therapy.

18) A DPP-4 inhibitor, which is preferably linagliptin, for use in a method of treating type 2

35 diabetes, said method comprising administration (e.g. separate, sequential or simultaneous

- 61 -

administration) of the DPP-4 inhibitor, a long-acting insulin selected from insulin detemir, insulin glargine and insulin degludec, and, optionally one or more further therapeutic agents (e.g. metformin or pioglitazone).

- 5 19) A long-acting basal insulin, preferably selected from insulin detemir, insulin glargine and insulin degludec, for use in a method of treating type 2 diabetes, said method comprising administration (e.g. separate, sequential or simultaneous administration) of the long-acting basal insulin, a DPP-4 inhibitor which is linagliptin, and, optionally one or more further therapeutic agents (e.g. metformin or pioglitazone).
- 10 20) A DPP-4 inhibitor, which is preferably linagliptin, for subcutaneous or transdermal administration, such as e.g. once daily, each other day, thrice weekly, twice weekly or once weekly, in the treatment of type 2 diabetes.
- 15 21) A pharmaceutical composition, preferably for subcutaneous or transdermal administration, comprising
- a) a DPP-4 inhibitor, which is preferably linagliptin, and, optionally,
- b) a long-acting basal insulin, which is preferably selected from insulin detemir, insulin glargine and insulin degludec,
- 20 and, optionally, one or more pharmaceutically acceptable carriers and/or diluents .

22. Separate, sequential or simultaneous use of a DPP-4 inhibitor, which is linagliptin, in combination with a long-acting insulin, which is basal insulin, in an elderly patient diagnosed with type 1 diabetes, LADA or type 2 diabetes.
- 5 23. The use according to claim 22, wherein the patient has type 2 diabetes.
24. The use according to claim 22 or 23, wherein the basal insulin is insulin detemir, insulin glargine or insulin degludec.
- 10 25. The use according to any one of claims 22 to 24, wherein the long-acting insulin and the DPP-4 inhibitor are each presented for use in a separate dosage form.
26. The use according to any one of claims 22 to 24, wherein the long-acting insulin and the DPP-4 inhibitor are presented for use in the same dosage form.
- 15 27. The use according to any one of claims 22 to 26, wherein the DPP-4 inhibitor is for subcutaneous administration.
28. The use according to any one of claims 22 to 26, wherein the long-acting insulin is for subcutaneous administration.
- 20 29. The use according to any one of claims 22 to 28, wherein both the DPP-4 inhibitor and the long-acting insulin are for subcutaneous administration.
- 25 30. The use according to any one of claims 22 to 29, wherein the long-acting insulin and the DPP-4 inhibitor are comprised in a pharmaceutical composition for subcutaneous administration by injection.
31. The use according to any one of claims 22 to 25, wherein the DPP-4 inhibitor is for peroral administration.
- 30 32. The use according to any one of the claims 22 to 31, further comprising use of one or more other therapeutic agents.

33. The use according to claim 32, wherein the one or more other therapeutic agents are selected from the group consisting of metformin, metformin hydrochloride, pioglitazone and pioglitazone hydrochloride.
- 5 34. The use according to any one of the claims 22 to 33 for improving or maintaining control of fasted and post-prandial glucose levels.
35. The use according to any one of the claims 22 to 33 for preventing, decreasing, reducing the risk of, slowing the progression of, delaying the onset of,
10 attenuating or treating complications of diabetes.
36. The use according to claim 35, wherein the complications of diabetes are selected from micro- and macrovascular diabetic complications, and angiopathies.
- 15 37. The use according to any one of the claims 22 to 36 in a patient with or at risk of renal impairment or renal complications.
38. The use according to any one of the claims 22 to 37 for achieving a reduction in the dose of insulin required for adequate therapeutic effect.
20
39. The use according to any one of the claims 22 to 38 for reducing hypoglycemia or weight gain associated with insulin therapy.
40. Use of a DPP-4 inhibitor, which is linagliptin, in the treatment of type 2
25 diabetes in an elderly patient, comprising use of the DPP-4 inhibitor, in combination with a long-acting insulin, which is basal insulin, and, optionally one or more further therapeutic agents.
41. The use according to claim 40, wherein the one or more further therapeutic
30 agents is metformin or metformin hydrochloride.
42. The use according to any one of claims 22 to 41, wherein the patient has inadequate glycemic control on insulin alone or in combination with one or more other antidiabetic drugs.

43. Use of a DPP-4 inhibitor, which is linagliptin, in the treatment of type 2 diabetes in an elderly patient, comprising use of the DPP-4 inhibitor as an add-on therapy to long-acting insulin, which is basal insulin, alone or in combination with one or more other antidiabetic drugs selected from metformin, metformin hydrochloride, pioglitazone, pioglitazone hydrochloride and a sulphonylurea, wherein the patient is inadequately controlled on insulin alone or in combination with one or more of the other antidiabetic drugs.
44. The use according to any one of claims 22 to 43, wherein the patient is an elderly patient of ≥ 60 -65 years.
45. The use according to any one of claims 22 to 44, wherein the patient has impaired or reduced renal function.
46. The use according to any one of claims 22 to 45, wherein the linagliptin is provided for use in a daily oral amount of 5 mg.
47. A pharmaceutical composition, for subcutaneous administration, comprising: (i) a DPP-4 inhibitor, which is linagliptin, and (ii) a long-acting basal insulin, and/or a pharmaceutically acceptable carrier or diluent.
48. Use of a DPP-4 inhibitor, which is linagliptin, in combination with a long-acting insulin, which is basal insulin, in the treatment of type 2 diabetes, wherein a reduction in the dose of insulin required for adequate therapeutic effect is achieved, or wherein hypoglycemia or weight gain associated with insulin therapy is reduced.

Figure 1

