



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2004/09/07
(87) Date publication PCT/PCT Publication Date: 2005/03/31
(85) Entrée phase nationale/National Entry: 2006/03/02
(86) N° demande PCT/PCT Application No.: US 2004/029094
(87) N° publication PCT/PCT Publication No.: 2005/028475
(30) Priorités/Priorities: 2003/09/04 (US60/500,199);
2003/12/08 (US60/527,907)

(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
C07D 487/04 (2006.01), *A61K 31/437* (2006.01),
A61K 31/519 (2006.01), *A61K 31/5025* (2006.01),
C07D 413/14 (2006.01), *C07D 413/04* (2006.01),
A61P 37/00 (2006.01), *A61P 35/00* (2006.01)
(71) Demandeur/Applicant:
VERTEX PHARMACEUTICALS INCORPORATED, US
(72) Inventeurs/Inventors:
ARONOV, ALEX, US;
LAUFFER, DAVID J., US;
LI, PAN, US;
TOMLINSON, RONALD C., US
(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSITIONS UTILES POUR INHIBER DES PROTEINES KINASES
(54) Title: COMPOSITIONS USEFUL AS INHIBITORS OF PROTEIN KINASES

(57) **Abrégé/Abstract:**

The present invention relates to compounds useful of inhibitors of protein kinases. The invention also provides pharmaceutically acceptable compositions comprising said compounds and methods of using the compositions in the treatment of various disease, conditions, or disorders.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
31 March 2005 (31.03.2005)

PCT

(10) International Publication Number
WO 2005/028475 A3

(51) International Patent Classification⁷: **C07D 471/04**,
487/04, 413/04, 413/14, A61K 31/437, 31/5025, 31/519,
A61P 35/00, 37/00

(21) International Application Number:
PCT/US2004/029094

(22) International Filing Date:
7 September 2004 (07.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/500,199 4 September 2003 (04.09.2003) US
60/527,907 8 December 2003 (08.12.2003) US

(71) Applicant (for all designated States except US): **VERTEX
PHARMACEUTICALS INCORPORATED** [US/US];
130 Waverly Street, Cambridge, MA 02139-4242 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ARONOV, Alex**
[RU/US]; 13 Circuit Lane, Watertown, MA 02472 (US).
LAUFFER, David, J. [US/US]; 254 Taylor Road, Stow,
MA 01775 (US). **LI, Pan** [US/US]; 15 Mystic View
Terrace, Arlington, MA 02474 (US). **TOMLINSON,
Ronald, C.** [US/US]; 15 Intervale Road, Sudbury, MA
01776 (US).

(74) Agent: **HALEY, James, F., Jr.**; c/o Fish & Neave, 1251
Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
9 June 2005

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS USEFUL AS INHIBITORS OF PROTEIN KINASES

(57) Abstract: The present invention relates to compounds useful of inhibitors of protein kinases. The invention also provides pharmaceutically acceptable compositions comprising said compounds and methods of using the compositions in the treatment of various disease, conditions, or disorders.



WO 2005/028475 A3

COMPOSITIONS USEFUL AS INHIBITORS
OF PROTEIN KINASES

Technical Field Of The Invention

[0001] The present invention relates to compounds useful as inhibitors of protein kinases. The invention also provides pharmaceutically acceptable compositions comprising the compounds of the invention and methods of using the compositions in the treatment of various disorders.

Background Of The Invention

[0002] The search for new therapeutic agents has been greatly aided in recent years by a better understanding of the structure of enzymes and other biomolecules associated with diseases. One important class of enzymes that has been the subject of extensive study is protein kinases.

[0003] Protein kinases constitute a large family of structurally related enzymes that are responsible for the control of a variety of signal transduction processes within the cell. (See, Hardie, G. and Hanks, S. *The Protein Kinase Facts Book, I and II*, Academic Press, San Diego, CA: 1995). Protein kinases are

- 2 -

thought to have evolved from a common ancestral gene due to the conservation of their structure and catalytic function. Almost all kinases contain a similar 250-300 amino acid catalytic domain. The

5 kinases may be categorized into families by the substrates they phosphorylate (e.g., protein-tyrosine, protein-serine/threonine, lipids, etc.). Sequence motifs have been identified that generally correspond to each of these kinase families (See, for example,

10 Hanks, S.K., Hunter, T., *FASEB J.* 1995, 9, 576-596; Knighton *et al.*, *Science* 1991, 253, 407-414; Hiles *et al.*, *Cell* 1992, 70, 419-429; Kunz *et al.*, *Cell* 1993, 73, 585-596; Garcia-Bustos *et al.*, *EMBO J.* 1994, 13, 2352-2361).

15 **[0004]** In general, protein kinases mediate intracellular signaling by effecting a phosphoryl transfer from a nucleoside triphosphate to a protein acceptor that is involved in a signaling pathway. These phosphorylation events act as molecular on/off

20 switches that can modulate or regulate the target protein biological function. These phosphorylation events are ultimately triggered in response to a variety of extracellular and other stimuli. Examples of such stimuli include environmental and chemical

25 stress signals (e.g., osmotic shock, heat shock, ultraviolet radiation, bacterial endotoxin, and H₂O₂), cytokines (e.g., interleukin-1 (IL-1) and tumor necrosis factor α (TNF- α)), and growth factors (e.g., granulocyte macrophage-colony-stimulating factor

30 (GM-CSF), and fibroblast growth factor (FGF)). An extracellular stimulus may affect one or more cellular responses related to cell growth, migration, differentiation, secretion of hormones, activation of

- 3 -

transcription factors, muscle contraction, glucose metabolism, control of protein synthesis, and regulation of the cell cycle.

[0005] Many diseases are associated with abnormal cellular responses triggered by protein kinase-mediated events as described above. These diseases include, but are not limited to, cancer and other proliferative disorders. Accordingly, there has been a substantial effort in medicinal chemistry to find protein kinase inhibitors that are effective as therapeutic agents.

[0006] The c-Met proto-oncogene encodes the Met receptor tyrosine kinase. The Met receptor is a 190kDa glycosylated dimeric complex composed of a 50kDa alpha chain disulfide-linked to a 145kDa beta chain. The alpha chain is found extracellularly while the beta chain contains transmembrane and cytosolic domains. Met is synthesized as a precursor and is proteolytically cleaved to yield mature alpha and beta subunits. It displays structural similarities to semaphorins and plexins, a ligand-receptor family that is involved in cell-cell interaction. The ligand for Met is hepatocyte growth factor (HGF), a member of the scatter factor family and has some homology to plasminogen [Longati, P. et al., *Curr. Drug Targets* 2001, 2, 41-55]; Trusolino, L. and Comoglio, P. *Nature Rev. Cancer* 2002, 2, 289-300].

[0007] Met functions in tumorigenesis and tumor metastasis. Chromosomal rearrangements forming Tpr-met fusions in an osteoclast cell line resulted in constitutively active Met receptors and transformation (Cooper, C.S. et al., *Nature* 1984, 311, 29-33). Met mutants exhibiting enhanced kinase activity have been identified in both hereditary and sporadic forms of

- 4 -

papillary renal carcinoma (Schmidt, L. et al., *Nat. Genet.* 1997, 16, 68-73; Jeffers, M. et al., *Proc. Nat. Acad. Sci.* 1997, 94, 11445-11500). Expression of Met along with its ligand HGF is transforming, tumorigenic, and metastatic (Jeffers, M. et al., *Oncogene* 1996, 13, 853-856; Michieli, P. et al., *Oncogene* 1999, 18, 5221-5231). HGF/Met has been shown to inhibit anoikis, suspension-induced programmed cell death (apoptosis), in head and neck squamous cell carcinoma cells. Anoikis resistance or anchorage-independent survival is a hallmark of oncogenic transformation of epithelial cells (Zeng, Q. et al., *J. Biol. Chem.* 2002, 277, 25203-25208).

[0008] MET is overexpressed in a significant percentage of human cancers and is amplified during the transition between primary tumors and metastasis. To investigate whether this oncogene is directly responsible for the acquisition of the metastatic phenotype, Giordano et al. exploited a single-hit oncogenic version of MET that was able to transform and to confer invasive and metastatic properties to nontumorigenic cells, both *in vitro* and in nude mice. They found a point mutation in the signal transducer docking site of MET that increased the transforming ability of the oncogene, but abolished its metastatic potential. They concluded that the metastatic potential of the MET oncogene relies on the properties of its multifunctional docking site, and that a single point mutation affecting signal transduction can dissociate neoplastic transformation from metastasis. Giordano, S., et al., *Proc. Nat. Acad. Sci.* 94: 13868-13872, 1997.

[0009] c-Met is implicated in various cancers, especially renal. It was found that the beta-subunit of the c-Met protooncogene product is the cell-surface receptor for hepatocyte growth factor. It was also
5 identified that the hepatocyte growth factor receptor is the c-met proto-oncogene product. Bottaro, D. P., et al., *Science* 251: 802-804, 1991.

[0010] The nexus between c-Met and colorectal cancer has also been established. Analysis of cMet expression
10 during colorectal cancer progression showed that 50% of the carcinoma specimens analyzed expressed 5-50-fold higher levels of cMet mRNA transcripts and protein versus the adjacent normal colonic mucosa. In addition, when compared to the primary tumor, 70% of
15 colorectal cancer liver metastasis showed cMet over expression. See Long et al., Met Receptor Overexpression and Oncogenic Ki-ras Mutation Cooperate to Enhance Tumorigenicity of Colon Cancer Cells in Vivo. *Mol Cancer Res.* 2003 Mar;1(5):393-401; Fujisaki,
20 et al., CD44 stimulation induces integrin-mediated adhesion of colon cancer cell lines to endothelial cells by up-regulation of integrins and c-Met and activation of integrins. *Cancer Res.* 1999 Sep 1;59(17):4427-34; Hiscox et al., Association of the
25 HGF/SF receptor, c-met, with the cell-surface adhesion molecule, E-cadherin, and catenins in human tumor cells. *Biochem Biophys Res Commun.* 1999 Aug 2;261(2):406-11; Herynk et al., Activation of c-Met in colorectal carcinoma cells leads to constitutive
30 association of tyrosine-phosphorylated beta-catenin. *Clin Exp Metastasis.* 2003;20(4):291-300; Wielenga et al., Expression of c-Met and heparan-sulfate proteoglycan forms of CD44 in colorectal cancer. *Am J*

- 6 -

Pathol. 2000 Nov;157(5):1563-73; Di Renzo et al., Overexpression and amplification of the Met/HGF receptor gene during the progression of colorectal cancer. *Clin. Cancer Res.*, 1: 147-154, 1995; and Mao, et al., Activation of c-Src by receptor tyrosine kinases in human colon cancer cells with high metastatic potential. *Oncogene*, 15:3083-3090, 1997.

5 [0011] The c-Met is also implicated in glioblastoma. High-grade malignant gliomas are the most common cancers of the central nervous system. Despite treatment with surgical resection, radiation therapy, and chemotherapy, the mean overall survival is <1.5 years, and few patients survive for > 3 years. A common reason for treatment failure is their innate resistance to radiation and chemotherapy.

10 [0012] Glioblastoma multiforme is the most common and most malignant glial neoplasm. Despite very aggressive treatment, these malignant gliomas are associated with an average life expectancy of only 9 months. The formation and malignant progression of human gliomas are complex processes and involve genetic mutations, chromosomal multiploidy, and aberrant epigenetic influences of multiple mitogens and angiogenic factors.

25 [0013] Human malignant gliomas frequently express both HGF and cMet, which can establish an autocrine loop of biological significance. Glioma cMet expression correlates with glioma grade, and an analysis of human tumor specimens showed that malignant gliomas have a 7-fold higher HGF content than low-grade gliomas.

30 [0014] Gliomas represent the most common form of primary central nervous system malignancy and are among

- 7 -

the tumors most tightly linked with HGF-cMet signaling abnormalities. Multiple studies have demonstrated that human gliomas frequently co-express HGF and cMet and that high levels of expression are associated with malignant progression. HGF gene transfer to glioma cell lines enhances tumorigenicity, tumor growth, and tumor-associated angiogenesis. It has also been shown that blocking HGF-cMet signaling reverses these phenotypes *in vivo*. It was further shown that HGF-cMet is able to activate Akt and protect glioma cell lines from apoptotic death, both *in vitro* and *in vivo*.

[0015] See Hirose et al., Clinical importance of cMet protein expression in high grade astrocytic tumors. *Neurol. Med.-Chir.* 38:851-859, 1998; Hirose et al., Immunohistochemical examination of cMet protein expression in astrocytic tumors. *Acta Neuropathol.* 95: 345-351, 1998; Koochekpour et al., Met and hepatocyte growth factor expression in human gliomas. *Cancer Res.* 57:5391-5398; Laterra et al., HGF expression enhances human glioblastoma tumorigenicity and growth. *Biochem. Biophys. Res. Commun.* 235:743-747; Moriyama et al., Concomitant expression of hepatocyte growth factor, HGF activator and cMet genes in human glioma cells *in vitro*. *FEBS Lett.* 372:78-82, 1995; Nabeshima et al., Expression of cMet correlates with grade of malignancy in human astrocytic tumors: an immunohistochemical study. *Histopathology* 31:436-443, 1997; Shiota et al., Coexpression of hepatocyte growth factor and its receptor (cMet) in HGL4 glioblastoma cells. *Lab. Invest.* 53:511-516, 1996; Welch et al., Hepatocyte growth factor and receptor (cMet) in normal and malignant astrocytic cells. *Anticancer Res.* 19:1635-1640, 1999; Bowers et al., HGF protects against

cytotoxic death in human glioblastoma via PI3-K and Akt-dependent pathways. *Cancer Res.* 60:4277-4283, 2000.

[0016] It was shown that the effect of NK4 (HGF antagonist), on HGF-promoted growth of a human breast cancer resulted in the reduction of tumor invasiveness and motility, weight and volume. Furthermore, in the *in-vitro* invasion assay and migration assay, both HGF and human fibroblasts, which secrete bioactive HGF, increased the invasiveness and migration of the breast cancer cells (MDA MB 231). See Growth and angiogenesis of human breast cancer in a nude mouse tumour model is reduced by NK4, the HGF antagonist. *Carcinogenesis*, May 9, 2003. Furthermore, transgenic mice harboring mutationally activated cMet developed metastatic mammary carcinoma. These same activating mutants were able to establish tumors in nude mouse NIH 3T3 xenografts (*PNAS*, Vol 95, pp 14417-14422, Nov. 1998).

[0017] Transgenic mice that overexpressed cMet in hepatocytes developed hepatocellular carcinoma (HCC), one of the human tumors in which cMet has been implicated previously. Inactivation of the transgene led to regression of even highly advanced tumors, apparently mediated by apoptosis and cessation of cellular proliferation. Numerous cells were proliferating in the liver tumors that were elicited by cMet. Removal of the stimulus from the transgenic hMet led to prompt cessation of cellular proliferation even in the cells of advanced malignancies (*The Journal of Cell Biology*, Vol. 153, 2001, p. 1023-1033).

[0018] HGF/Met signaling is involved in cell adhesion and motility in normal cells and plays a major role in the invasive growth that is found in most tissues, including cartilage, bone, blood vessels, and

- 9 -

neurons (reviewed in Comoglio, P.M. and Trusolino, L. *J. Clin. Invest.* 2002, 109, 857-862). Dysfunctional activation or increased numbers of Met is likely to contribute to the aberrant cell-cell interactions that lead to migration, proliferation, and survival of cells that is characteristic of tumor metastasis. Activation of Met induces and sustains a variety of tumors [Wang, R. et al., *J. Cell. Biol.* 2001, 153, 1023-1034; Liang, T. J. et al., *J. Clin. Invest.* 1996, 97, 2872-2877; Jeffers, M. et al., *Proc. Nat. Acad. Sci.* 1998, 95, 14417-14422] while loss of Met inhibits growth and invasiveness of tumor cells [Jiang, W.G. et al., *Clin. Cancer Res.* 2001, 7, 2555-2562; Abounader, R. et al., *FASEB J.* 2002 16, 108-110]. Increased expression of Met/HGF is seen in many metastatic tumors including colon (Fazekas, K. et al., *Clin. Exp. Metastasis* 2000, 18, 639-649), breast (Elliott, B.E. et al., 2002, *Can. J. Physiol. Pharmacol.* 80, 91-102), prostate (Knudsen, B.S. et al., *Urology* 2002, 60, 1113-1117), lung (Siegfried, J.M. et al., *Ann. Thorac. Surg.* 1998, 66, 1915-1918), and gastric (Amemiya, H. et al., *Oncology* 2002, 63, 286-296).

[0019] Further demonstration of the role Met plays in metastasis was shown by Giordano, et al. (2002) who presented evidence for cross-talk between the semaphorin 4D (SEMA4D; 601866) receptor, plexin B1 (PLXNB1; 601053), and MET during invasive growth in epithelial cells. Binding of SEMA4D to PLXNB1 stimulated tyrosine kinase activity of MET, resulting in tyrosine phosphorylation of both receptors. This effect was not found in cells lacking MET expression. Giordano, S., et al.: The Semaphorin 4D receptor

- 10 -

controls invasive growth by coupling with Met. *Nature Cell Biol.* 4: 720-724, 2002.

[0020] HGF-Met signaling has also been associated with increased risk of atherosclerosis (Yamamoto, Y. et al., *J.Hypertens.* 2001, 19,1975-1979; Morishita, R. et al., *Endocr. J.* 2002, 49, 273-284) and increased fibrosis of the lung (Crestani, B. et al., *Lab. Invest.* 2002, 82, 1015-1022.

[0021] Glycogen synthase kinase-3 (GSK-3) is a serine/threonine protein kinase comprised of a and b isoforms that are each encoded by distinct genes [Coghlan et al., *Chemistry & Biology*, 7, 793-803 (2000); Kim and Kimmel, *Curr. Opinion Genetics Dev.*, 10, 508-514 (2000)]. GSK-3 has been implicated in various diseases including diabetes, Alzheimer's disease, CNS disorders such as manic depressive disorder and neurodegenerative diseases, and cardiomyocyte hypertrophy [see, e.g., WO 99/65897; WO 00/38675; Kaytor and Orr, *Curr. Opin. Neurobiol.*, 12, 275-8 (2000); Haq et al., *J. Cell Biol.*, 151, 117-30 (2000); Eldar-Finkelman, *Trends Mol. Med.*, 8, 126-32 (2002)]. These diseases are associated with the abnormal operation of certain cell signaling pathways in which GSK-3 plays a role.

[0022] GSK-3 has been found to phosphorylate and modulate the activity of a number of regulatory proteins. These include glycogen synthase, which is the rate-limiting enzyme required for glycogen synthesis, the microtubule-associated protein Tau, the gene transcription factor b-catenin, the translation initiation factor eIF-2B, as well as ATP citrate lyase, axin, heat shock factor-1, c-Jun, c-myc, c-myb, CREB, and CEPBa. These diverse targets implicate GSK-3 in

- 11 -

many aspects of cellular metabolism, proliferation, differentiation and development.

[0023] In a GSK-3 mediated pathway that is relevant for the treatment of type II diabetes, insulin-induced signaling leads to cellular glucose uptake and glycogen synthesis. GSK-3 is a negative regulator of the insulin-induced signal in this pathway. Normally, the presence of insulin causes inhibition of GSK-3-mediated phosphorylation and deactivation of glycogen synthase. The inhibition of GSK-3 leads to increased glycogen synthesis and glucose uptake [Klein et al., PNAS, 93, 8455-9 (1996); Cross et al., Biochem. J., 303, 21-26 (1994); Cohen, Biochem. Soc. Trans., 21, 555-567 (1993); and Massillon et al., Biochem J. 299, 123-128 (1994); Cohen and Frame, Nat. Rev. Mol. Cell. Biol., 2, 769-76 (2001)]. However, where the insulin response is impaired in a diabetic patient, glycogen synthesis and glucose uptake fail to increase despite the presence of relatively high blood levels of insulin. This leads to abnormally high blood levels of glucose with acute and chronic effects that may ultimately result in cardiovascular disease, renal failure and blindness. In such patients, the normal insulin-induced inhibition of GSK-3 fails to occur. It has also been reported that GSK-3 is overexpressed in patients with type II diabetes [WO 00/38675]. Therapeutic inhibitors of GSK-3 are therefore useful for treating diabetic patients suffering from an impaired response to insulin.

[0024] Apoptosis has been implicated in the pathophysiology of ischemic brain damage (Li et al., 1997; Choi, et al., 1996; Charriaut-Marlangue et al., 1998; Grahm and Chen, 2001; Murphy et al., 1999;

- 12 -

Nicotera et al., 1999). Recent publications indicate that activation of GSK-3 β may be involved in apoptotic mechanisms (Kaytor and Orr, 2002; Culbert et al., 2001). Studies in rat models of ischemic stroke
5 induced by middle cerebral artery occlusion (MCAO) showed increased GSK-3b expression is following ischemia (Wang et al., Brain Res, 859, 381-5, 2000; Sasaki et al., Neurol Res, 23, 588-92, 2001).

Fibroblast growth factor (FGF) reduced ischemic brain
10 injury after permanent middle cerebral artery occlusion (MCO) in rats (Fisher et al. 1995; Song et al. 2002). Indeed, the neuroprotective effects of FGF demonstrated in ischemia models in rats may be mediated by a PI-3 kinase/AKT-dependent inactivation of GSK-3b (Hashimoto
15 et al., 2002). Thus, inhibition of GSK-3 β after a cerebral ischemic event may ameliorate ischemic brain damage.

[0025] GSK-3 is also implicated in myocardial infarction. See Jonassen et al., Circ Res, 89:1191,
20 2001 (The reduction in myocardial infarction by insulin administration at reperfusion is mediated via Akt dependent signaling pathway); Matsui et al., Circulation, 104:330, 2001 (Akt activation preserves cardiac function and prevents cardiomyocyte injury
25 after transient cardiac ischemia in vivo); Miao et al., J Mol Cell Cardiol, 32:2397, 2000 (Intracoronary, adenovirus-mediated Akt gene delivery in heart reduced gross infarct size following ischemia-reperfusion injury in vivo); and Fujio et al., Circulation et al.,
30 101:660, 2000 (Akt signaling inhibits cardiac myocyte apoptosis *in vitro* and protects against ischemia-reperfusion injury in mouse heart).

- 13 -

[0026] GSK-3 activity plays a role in head trauma.
See Noshita et al., Neurobiol Dis, 9:294, 2002
(Upregulation of Akt/PI3-kinase pathway may be crucial
for cell survival after traumatic brain injury) and
5 Dietrich et al., J Neurotrauma, 13:309, 1996
(Posttraumatic administration of bFGF significantly
reduced damaged cortical neurons & total contusion
volume in a rat model of traumatic brain injury).

[0027] GSK-3 is also known to play a role in
10 psychiatric disorders. See Eldar-Finkelman, Trends Mol
Med, 8:126, 2002; Li et al., Bipolar Disord, 4:137,
2002 (LiCl and Valproic acid, anti-psychotic, mood
stabilizing drugs, decrease GSK3 activities and
increase beta-catenin) and Lijam et al., Cell, 90:895,
15 1997 (Dishevelled KO mice showed abnormal social
behavior and defective sensorimotor gating.
Dishevelled, a cytoplasmic protein involved in WNT
pathway, inhibits GSK3beta activities).

[0028] It has been shown that GSK3 inhibition by
20 lithium and valproic acid induces axonal remodeling and
change synaptic connectivity. See Kaytor & Orr, Curr
Opin Neurobiol, 12:275, 2002 (Downregulation of GSK3
causes changes in microtubule-associated proteins: tau,
MAP1 & 2) and Hall et al., Mol Cell Neurosci, 20:257,
25 2002 (Lithium and valproic acid induces the formation
of growth cone-like structures along the axons).

[0029] GSK-3 activity is also associated with
Alzheimer's disease. This disease is characterized by
the presence of the well-known b-amyloid peptide and
30 the formation of intracellular neurofibrillary tangles.
The neurofibrillary tangles contain hyperphosphorylated
Tau protein, in which Tau is phosphorylated on abnormal
sites. GSK-3 has been shown to phosphorylate these

- 14 -

abnormal sites in cell and animal models. Furthermore, inhibition of GSK-3 has been shown to prevent hyperphosphorylation of Tau in cells [Lovestone et al., Curr. Biol., 4, 1077-86 (1994); and Brownlees et al., Neuroreport 8, 3251-55 (1997); Kaytor and Orr, Curr. Opin. Neurobiol., 12, 275-8 (2000)]. In transgenic mice overexpressing GSK3, significant increased Tau hyperphosphorylation and abnormal morphology of neurons were observed [Lucas et al., EMBO J, 20:27-39 (2001)].

Active GSK3 accumulates in cytoplasm of pretangled neurons, which can lead to neurofibrillary tangles in brains of patients with AD [Pei et al., J Neuropathol Exp Neurol, 58, 1010-19 (1999)]. Therefore, inhibition of GSK-3 slows or halts the generation of neurofibrillary tangles and thus treats or reduces the severity of Alzheimer's disease.

[0030] Evidence for the role GSK-3 plays in Alzheimer's disease has been shown *in vitro*. See Aplin et al. (1996), J Neurochem 67:699; Sun et al. (2002), Neurosci Lett 321:61 (GSK3b phosphorylates cytoplasmic domain of Amyloid Precursor Protein (APP) and GSK3b inhibition reduces Ab40 & Ab42 secretion in APP-transfected cells); Takashima et al. (1998), PNAS 95:9637; Kirschenbaum et al. (2001), J Biol Chem 276:7366 (GSK3b complexes with and phosphorylates presenilin-1, which is associated with gamma-secretase activity in the synthesis of Ab from APP); Takashima et al. (1998), Neurosci Res 31:317 (Activation of GSK3b by Ab(25-35) enhances phosphorylation of tau in hippocampal neurons. This observation provides a link between Ab and neurofibrillary tangles composed of hyperphosphorylated tau, another pathological hallmark of AD); Takashima et al. (1993), PNAS 90:7789 (Blockade

- 15 -

of GSK3b expression or activity prevents Ab-induced neuro-degeneration of cortical and hippocampal primary cultures); Suhara et al. (2003), Neurobiol Aging. 24:437 (Intracellular Ab42 is toxic to endothelial cells by interfering with activation of Akt/GSK-3b signaling-dependent mechanism); De Ferrari et al. (2003) Mol Psychiatry 8:195 (Lithium protects N2A cells & primary hippocampal neurons from Ab fibrils-induced cytotoxicity, & reduced nuclear translocation/destabilization of b-catenin); and Pigino et al., J Neurosci, 23:4499, 2003 (The mutations in Alzheimer's presenilin 1 may deregulate and increase GSK-3 activity, which in turn, impairs axonal transport in neurons. The consequent reductions in axonal transport in affected neurons can ultimately lead to neurodegeneration).

[0031] Evidence for the role GSK-3 plays in Alzheimer's disease has been shown in vivo. See Yamaguchi et al. (1996), Acta Neuropathol 92:232; Pei et al. (1999), J Neuropath Exp Neurol 58:1010 (GSK3b immunoreactivity is elevated in susceptible regions of AD brains); Hernandez et al. (2002), J Neurochem 83:1529 (Transgenic mice with conditional GSK3b overexpression exhibit cognitive deficits similar to those in transgenic APP mouse models of AD); De Ferrari et al. (2003) Mol Psychiatry 8:195 (Chronic lithium treatment rescued neurodegeneration and behavioral impairments (Morris water maze) caused by intrahippocampal injection of Ab fibrils.); McLaurin et al., Nature Med, 8:1263, 2002 (Immunization with Ab in a transgenic model of AD reduces both AD-like neuropathology and the spatial memory impairments); and Phiel et al. (2003) Nature 423:435 (GSK3 regulates

- 16 -

amyloid-beta peptide production via direct inhibition of gamma secretase in AD tg mice).

[0032] Presenilin-1 and kinesin-1 are also substrates for GSK-3 and relate to another mechanism for the role GSK-3 plays in Alzheimer's disease, as was recently described by Pigino, G., et al., Journal of Neuroscience (23:4499, 2003). It was found that GSK3beta phosphorylates kinesin-I light chain, which results in a release of kinesin-1 from membrane-bound organelles, leading to a reduction in fast anterograde axonal transport (Morfini et al., 2002). The authors suggest that the mutations in PS1 may deregulate and increase GSK-3 activity, which in turn, impairs axonal transport in neurons. The consequent reductions in axonal transport in affected neurons ultimately lead to neurodegeneration.

[0033] GSK-3 is also associated with amyotrophic lateral sclerosis (ALS). See Williamson and Cleveland, 1999 (Axonal transport is retarded in a very early phase of ALS in mSOD1 mice); Morfini et al., 2002 (GSK3 phosphorylates kinesin light chains and inhibit anterograde axonal transport); Warita et al., Apoptosis, 6:345, 2001 (The majority of spinal motor neurons lost the immunoreactivities for both PI3-K and Akt in the early and presymptomatic stage that preceded significant loss of the neurons in this SOD1 tg animal model of ALS); and Sanchez et al., 2001 (The inhibition of PI-3K induces neurite retraction mediated by GSK3 activation).

[0034] GSK-3 activity is also linked to spinal cord and peripheral nerve injuries. It has been shown that GSK3 inhibition by lithium and valproic acid can induce axonal remodeling and change synaptic connectivity.

- 17 -

See Kaytor & Orr, Curr Opin Neurobiol, 12:275, 2002
(Downregulation of GSK3 causes changes in
microtubule-associated proteins: tau, MAP1 & 2) and Hall
et al., Mol Cell Neurosci, 20:257, 2002 (Lithium and
5 valproic acid induces the formation of growth cone-like
structures along the axons). See also Grothe et al.,
Brain Res, 885:172, 2000 (FGF2 stimulate Schwann cell
proliferation and inhibit myelination during axonal
growth); Grothe and Nikkhah, 2001 (FGF-2 is up
10 regulated in the proximal and distal nerve stumps
within 5 hours after nerve crush); and Sanchez et al.,
2001 (The inhibition of PI-3K induces neurite
retraction mediated by GSK3 activation).

[0035] Another substrate of GSK-3 is b-catenin,
15 which is degraded after phosphorylation by GSK-3.
Reduced levels of b-catenin have been reported in
schizophrenic patients and have also been associated
with other diseases related to increase in neuronal
cell death [Zhong et al., Nature, 395, 698-702 (1998);
20 Takashima et al., PNAS, 90, 7789-93 (1993); Pei et al.,
J. Neuropathol. Exp, 56, 70-78 (1997); and Smith
et al., Bio-org. Med. Chem. 11, 635-639 (2001)].
Furthermore, b-catenin and Tcf-4 play a dual role in
vascular remodeling by inhibiting vascular smooth
25 muscle cell apoptosis and promoting proliferation (Wang
et al., Circ Res, 90:340, 2002). Accordingly, GSK-3 is
associated with angiogenic disorders. See also Liu
et al., FASEB J, 16:950, 2002 (Activation of GSK3
reduces hepatocyte growth factor, leading to altered
30 endothelial cell barrier function and diminished
vascular integrity) and Kim et al., J Biol Chem,
277:41888, 2002 (GSK3beta activation inhibits
angiogenesis in vivo using Matrigel plug assay: the

- 18 -

inhibition of GSK3beta signaling enhances capillary formation).

[0036] Association between GSK-3 and Huntington's disease has been shown. See Carmichael et al., J Biol Chem., 277:33791, 2002 (GSK3beta inhibition protect cells from poly-glutamine-induced neuronal and non-neuronal cell death via increases in b-catenin and its associated transcriptional pathway).

Overexpression of GSK3 reduced the activation of heat shock transcription factor-1 and heat shock protein HSP70 (Bijur et al., J Biol Chem, 275:7583, 2000) that are shown to decrease both poly-(Q) aggregates and cell death in *in vitro* HD model (Wytenbach et al., Hum Mol Genet, 11:1137, 2002).

[0037] GSK-3 effects the levels of FGF-2 and their receptors are increased during remyelination of brain aggregate cultures remyelinating rat brains. See Copelman et al., 2000, Messersmith, et al., 2000; and Hinks and Franklin, 2000. It was also found that FGF-2 induces process outgrowth by oligodendrocytes implicating involvement of FGF in remyelination (Oh and Yong, 1996; Gogate et al., 1994) and that FGF-2 gene therapy has shown to improve the recovery of experimental allergic encephalomyelitis (EAE) mice (Ruffini, et al., 2001).

[0038] GSK-3 has also been associated with hair growth because Wnt/beta-catenin signaling is shown to play a major role in hair follicle morphogenesis and differentiation (Kishimoto et al. Genes Dev, 14:1181, 2000; Millar, J Invest Dermatol, 118:216, 2002). It was found that mice with constitutive overexpression of the inhibitors of Wnt signaling in skin failed to develop hair follicles. Wnt signals are required for

- 19 -

the initial development of hair follicles and GSK3 constitutively regulates Wnt pathways by inhibiting beta-catenin. (Andl et al., Dev Cell 2:643, 2002). A transient Wnt signal provides the crucial initial stimulus for the start of a new hair growth cycle, by activating beta-catenin and TCF-regulated gene transcription in epithelial hair follicle precursors (Van Mater et al., Genes Dev, 17:1219, 2003)

5 [0039] Because GSK-3 activity is associated with sperm motility, GSK-3 inhibition is useful as a male contraceptive. It was shown that a decline in sperm GSK3 activity is associated with sperm motility development in bovine and monkey epididymis (Vijayaraghavan et al., Biol Reprod, 54: 709, 1996; 10 Smith et al., J Androl, 20:47, 1999). Furthermore, tyrosine & serine/threonine phosphorylation of GSK3 is high in motile compared to immotile sperm in bulls (Vijayaraghavan et al., Biol Reprod, 62:1647, 2000). This effect was also demonstrated with human sperm 15 (Luconi et al., Human Reprod, 16:1931, 2001).

20 [0040] The Janus kinases (JAK) are a family of tyrosine kinases consisting of JAK1, JAK2, JAK3 and TYK2. The JAKs play a critical role in cytokine signaling. The down-stream substrates of the JAK family of kinases include the signal transducer and 25 activator of transcription (STAT) proteins. JAK/STAT signaling has been implicated in the mediation of many abnormal immune responses such as allergies, asthma, autoimmune diseases such as transplant rejection, rheumatoid arthritis, amyotrophic lateral sclerosis and 30 multiple sclerosis as well as in solid and hematologic malignancies such as leukemias and lymphomas. The pharmaceutical intervention in the JAK/STAT pathway has

- 20 -

been reviewed [Frank Mol. Med. 5: 432-456 (1999) & Seidel, et al., Oncogene 19: 2645-2656 (2000)].

[0041] JAK1, JAK2, and TYK2 are ubiquitously expressed, while JAK3 is predominantly expressed in hematopoietic cells. JAK3 binds exclusively to the common cytokine receptor gamma chain (gc) and is activated by IL-2, IL-4, IL-7, IL-9, and IL-15. The proliferation and survival of murine mast cells induced by IL-4 and IL-9 have, in fact, been shown to be dependent on JAK3- and gc- signaling [Suzuki et al., Blood 96: 2172-2180 (2000)].

[0042] Cross-linking of the high-affinity immunoglobulin (Ig) E receptors of sensitized mast cells leads to a release of proinflammatory mediators, including a number of vasoactive cytokines resulting in acute allergic, or immediate (type I) hypersensitivity reactions [Gordon et al., Nature 346: 274-276 (1990) & Galli, N. Engl. J. Med., 328: 257-265 (1993)]. A crucial role for JAK3 in IgE receptor-mediated mast cell responses *in vitro* and *in vivo* has been established [Malaviya, et al., Biochem. Biophys. Res. Commun. 257: 807-813 (1999)]. In addition, the prevention of type I hypersensitivity reactions, including anaphylaxis, mediated by mast cell-activation through inhibition of JAK3 has also been reported [Malaviya et al., J. Biol. Chem. 274:27028-27038 (1999)]. Targeting mast cells with JAK3 inhibitors modulated mast cell degranulation *in vitro* and prevented IgE receptor/antigen-mediated anaphylactic reactions *in vivo*.

[0043] A recent study described the successful targeting of JAK3 for immune suppression and allograft acceptance. The study demonstrated a dose-dependent

- 21 -

survival of Buffalo heart allograft in Wistar Furth recipients upon administration of inhibitors of JAK3 indicating the possibility of regulating unwanted immune responses in graft versus host disease [Kirken, 5 transpl. proc. 33: 3268-3270 (2001)].

[0044] IL-4-mediated STAT-phosphorylation has been implicated as the mechanism involved in early and late stages of rheumatoid arthritis (RA). Up-regulation of proinflammatory cytokines in RA synovium and synovial 10 fluid is a characteristic of the disease. It has been demonstrated that IL-4-mediated activation of IL-4/STAT pathway is mediated through the Janus Kinases (JAK 1 & 3) and that IL-4-associated JAK kinases are expressed in the RA synovium [Muller-Ladner, et al., J. Immunol. 15 164: 3894-3901 (2000)].

[0045] Familial amyotrophic lateral sclerosis (FALS) is a fatal neurodegenerative disorder affecting about 10% of ALS patients. The survival rates of FALS mice were increased upon treatment with a JAK3 specific 20 inhibitor. This confirmed that JAK3 plays a role in FALS [Trieu, et al., Biochem. Biophys. Res. Commun. 267: 22-25 (2000)].

[0046] Signal transducer and activator of transcription (STAT) proteins are activated by, among 25 others, the JAK family kinases. Results from a recent study suggested the possibility of intervention in the JAK/STAT signaling pathway by targeting JAK family kinases with specific inhibitors for the treatment of leukemia [Sudbeck, et al., Clin. Cancer Res. 5: 30 1569-1582 (1999)]. JAK3 specific compounds were shown to inhibit the clonogenic growth of JAK3-expressing cell lines DAUDI, RAMOS, LC1; 19, NALM-6, MOLT-3 and HL-60.

- 22 -

[0047] In animal models, TEL/JAK2 fusion proteins have induced myeloproliferative disorders and in hematopoietic cell lines, introduction of TEL/JAK2 resulted in activation of STAT1, STAT3, STAT5, and
5 cytokine-independent growth [Schwaller, et al., EMBO J. 17: 5321-5333 (1998)].

[0048] Inhibition of JAK 3 and TYK 2 abrogated tyrosine phosphorylation of STAT3, and inhibited cell growth of mycosis fungoides, a form of cutaneous T cell
10 lymphoma. These results implicated JAK family kinases in the constitutively activated JAK/STAT pathway that is present in mycosis fungoides [Nielsen, et al., Proc. Nat. Acad. Sci. U.S.A. 94: 6764-6769 (1997)].
Similarly, STAT3, STAT5, JAK1 and JAK2 were
15 demonstrated to be constitutively activated in mouse T cell lymphoma characterized initially by LCK over-expression, thus further implicating the JAK/STAT pathway in abnormal cell growth [Yu, et al., J. Immunol. 159: 5206-5210 (1997)]. In addition, IL-6 -
20 mediated STAT3 activation was blocked by an inhibitor of JAK, leading to sensitization of myeloma cells to apoptosis [Catlett-Falcone, et al., Immunity 10:105-115 (1999)].

[0049] Tyrosine kinases are a class of enzymes that
25 mediate intracellular signal transduction pathways. Abnormal activity of these kinases has been shown to contribute to cell proliferation, carcinogenesis and cell differentiation. Thus, agents that modulate the activity of tyrosine kinases are useful for preventing
30 and treating proliferative diseases associated with these enzymes.

[0050] Syk is a tyrosine kinase that plays a critical role in FcεRI mediated mast cell degranulation

- 23 -

and eosinophil activation. Accordingly, Syk kinase is implicated in various allergic disorders, in particular asthma. It has been shown that Syk binds to the phosphorylated gamma chain of the FcεRI receptor via N-terminal SH2 domains and is essential for downstream signaling [Taylor et al., *Mol. Cell. Biol.* 1995, 15, 4149].

[0051] Inhibition of eosinophil apoptosis has been proposed as a key mechanism for the development of blood and tissue eosinophilia in asthma. IL-5 and GM-CSF are upregulated in asthma and are proposed to cause blood and tissue eosinophilia by inhibition of eosinophil apoptosis. Inhibition of eosinophil apoptosis has been proposed as a key mechanism for the development of blood and tissue eosinophilia in asthma. It has been reported that Syk kinase is required for the prevention of eosinophil apoptosis by cytokines (using antisense) [Yousefi et al., *J. Exp. Med.* 1996, 183, 1407].

[0052] The role of Syk in FcγR dependent and independent response in bone marrow derived macrophages has been determined by using irradiated mouse chimeras reconstituted with fetal liver cells from Syk -/- embryos. Syk deficient macrophages were defective in phagocytosis induced by FcγR but showed normal phagocytosis in response to complement [Kiefer et al., *Mol. Cell. Biol.* 1998, 18, 4209]. It has also been reported that aerosolized Syk antisense suppresses Syk expression and mediator release from macrophages [Stenton et al., *J. Immunology* 2000, 164, 3790].

[0053] KDR is a tyrosine kinase receptor that also binds VEGF (vascular endothelial growth factor) Neufeld et al., 1999, *FASEB J.*, 13, 9. The binding of

- 24 -

VEGF to the KDR receptor leads to angiogenesis, which is the sprouting of capillaries from preexisting blood vessels. High levels of VEGF are found in various cancers causing tumor angiogenesis and permitting the rapid growth of cancerous cells. Therefore, suppressing VEGF activity is a way to inhibit tumor growth, and it has been shown that this can be achieved by inhibiting KDR receptor tyrosine kinase. For example, SU5416 is a selective inhibitor of the tyrosine kinase and was reported to also suppress tumor vascularization and the growth of multiple tumors. Fong et al., 1999, Cancer Res. 59, 99. Other inhibitors of KDR tyrosine kinase for the treatment of cancer have also been reported (WO 98/54093, WO 99/16755, WO 00/12089).

[0054] Examples of cancers that may be treated by such inhibitors include brain cancer, genitourinary tract cancer, lymphatic system cancer, stomach cancer, cancer of the larynx, lung cancer, pancreatic cancer, breast cancer, Kaposi's sarcoma, and leukemia. Other diseases and conditions associated with abnormal tyrosine kinase activity include vascular disease, autoimmune diseases, ocular conditions, and inflammatory diseases.

[0055] A family of type III receptor tyrosine kinases including Flt3, c-Kit, PDGF-receptor and c-Fms play an important role in the maintenance, growth and development of hematopoietic and non-hematopoietic cells. [Scheijen, B, Griffin JD, *Oncogene*, 2002, 21, 3314-3333 and Reilly, JT, *British Journal of Haematology*, 2002, 116, 744-757]. FLT-3 and c-Kit regulate maintenance of stem cell/early progenitor pools as well the development of mature lymphoid and

- 25 -

myeloid cells [Lyman, S, Jacobsen, S, *Blood*, 1998, 91, 1101-1134]. Both receptors contain an intrinsic kinase domain that is activated upon ligand-mediated dimerization of the receptors. Upon activation, the kinase domain induces autophosphorylation of the receptor as well as the phosphorylation of various cytoplasmic proteins that help propagate the activation signal leading to growth, differentiation and survival. Some of the downstream regulators of FLT-3 and c-Kit receptor signaling include, PLC γ , PI3-kinase, Grb-2, SHIP and Src related kinases [Scheijen, B, Griffin JD, *Oncogene*, 2002, 21, 3314-3333]. Both receptor tyrosine kinases have been shown to play a role in a variety of hematopoietic and non-hematopoietic malignancies. Mutations that induce ligand independent activation of FLT-3 and c-Kit have been implicated acute-myelogenous leukemia (AML), acute lymphocytic leukemia (ALL), mastocytosis and gastrointestinal stromal tumor (GIST). These mutations include single amino acid changes in the kinase domain or internal tandem duplications, point mutations or in-frame deletions of the juxtamembrane region of the receptors. In addition to activating mutations, ligand dependent (autocrine or paracrine) stimulation of over-expressed wild-type FLT-3 or c-Kit can contribute to the malignant phenotype [Scheijen, B, Griffin JD, *Oncogene*, 2002, 21, 3314-3333].

[0056] c-fms encodes for macrophage colony stimulating factor receptor (M-CSF-1R) which is expressed predominately in the monocytes/macrophage lineage [Dai, XM et al., *Blood*, 2002, 99, 111-120]. MCSF-1R and its ligand regulate macrophage lineage growth and differentiation. Like the other family

- 26 -

members, MCSF-1R contains an intrinsic kinase domain that is activated upon ligand-induced dimerization of the receptor. MCSF-1R is also expressed in non-hematopoietic cells including mammary gland epithelial cells and neurons. Mutations in this receptor are potentially linked to myeloid leukemias and its expression is correlated with metastatic breast, ovarian and endometrial carcinomas [Reilly, JT, *British Journal of Haematology*, 2002, 116, 744-757 and Kacinski, BM, *Mol. Reprod and Devel.*, 1997, 46, 71-74]. Another possible indication for antagonists of MCSF-1R is osteoporosis [Teitelbaum, S, *Science* 2000, 289, 1504-1508.

[0057] Aurora-2 is a serine/threonine protein kinase that has been implicated in human cancer, such as colon, breast and other solid tumors. This kinase is involved in protein phosphorylation events that regulate the cell cycle. Specifically, Aurora-2 plays a role in controlling the accurate segregation of chromosomes during mitosis. Misregulation of the cell cycle can lead to cellular proliferation and other abnormalities. In human colon cancer tissue, the aurora-2 protein has been found to be overexpressed [Bischoff et al., *EMBO J.*, 17, 3052-3065 (1998); Schumacher et al., *J. Cell Biol.*, 143, 1635-1646 (1998); Kimura et al., *J. Biol. Chem.*, 272, 13766-13771 (1997)].

[0058] Transforming growth factor-beta (TGF-beta) activated kinase 1 (TAK-1) is a 67kDa ubiquitin-dependent serine-threonine kinase that functions as a mitogen-activated protein (MAP) kinase kinase kinase (MAPKKK or MEKK) (Wang, C., et al., *Nature* 2001, 412, 346 - 351).

- 27 -

- [0059] Originally described as stimulated by TGF-beta superfamily members (Yamaguchi K. et al., Science 1995, 270, 2008-2011) TAK-1 is known to also function in signaling from numerous cell modulators including proinflammatory cytokines. TAK-1 is critical for signaling from IL-1beta/ TLR ligands (Holtmann H, et al., J. Biol. Chem. 2001, 276, 3508-3516; Jiang Z, et al., J. Biol. Chem. 2003, 278, 16713-16719) and TNF-alpha (Takaesu G. et al., J. Mol. Biol. 2003, 326, 105-115). In addition TAK-1 plays a role in IL-18 (Wald, D., et al., Eur. J. Immunol. 2001, 31, 3747-3754), RANKL (Mizukami J., et al., Mol. Cell. Biol. 2002, 22, 992-1000) and ceramide (Shirakabe K., et al., J. Biol. Chem. 1997, 272, 8141-8144) signaling.
- [0060] Through interaction with corresponding cell surface receptors these ligands stimulate TAK-1 to relay signals to a variety of pathways such as IKK/NFkappaB, JNK, and p38, that are important regulators of cellular processes including apoptosis (Edlund S., et al., Mol Biol Cell. 2003, 14, 529-544), differentiation (Suzawa, M. et al., Nat Cell Biol 2003, 5, 224-230), and cell cycle progression (Bradham CA, et al., Am J Physiol Gastrointest Liver Physiol. 2001 281, G1279-89).
- [0061] Modification of signaling pathways can alter cellular processes and contribute to disease. Due to its central role in signaling from numerous cell surface receptors TAK-1 may be an important therapeutic target for a variety of diseases. The cytokines IL-1beta and TNFalpha are important mediators of inflammation in rheumatoid arthritis and other inflammatory diseases (Maini RN. and Taylor PC. Ann. Rev. Med. 2000, 51, 207-229). TAK-1 may be important

- 28 -

in regulating disease-relevant cellular responses in these cases (Hammaker DR, et al. J. Immunol. 2004, 172, 1612-1618). TAK-1 affects cellular fibrotic responses (Ono K., et al., Biochem. Biophys. Res. Commun. 2003, 307, 332-337). It may also plays a role in heart failure (Zhang, D., Nat. Med. 2000, 6, 556-563), osteoporosis (Mizukami J, et al., Mol. Cell. Biol. 2002, 22, 992-1000) and survival of hepatocellular carcinoma cells (Arsura M, et al. Oncogene 2003, 22, 412-425). TAK-1 signaling may affect neurite outgrowth (Yanagisawa M., et al. Genes Cells. 2001, 6, 1091-1099) and is involved in control of adipogenesis (Suzawa M., et al. Nat. Cell. Biol. 2003, 5, 224-230) and cardiomyocyte differentiation (Monzen K., et al. J. Cell. Biol. 2001, 153(4), 687-698).


[0062] As a result of the biological importance of protein kinases, there is current interest in therapeutically effective protein kinase inhibitors. Accordingly, there is still a great need to develop inhibitors of protein kinases that are useful in treating various diseases or conditions associated with protein kinase activation. In particular, it would be desirable to develop compounds that are useful as inhibitors of c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, or TAK-1 particularly given the inadequate treatments currently available for the majority of the disorders implicated in their activation.

Summary Of The Invention

[0063] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are effective as inhibitors of protein kinases. In certain embodiments, these compounds are

- 29 -

effective as inhibitors of c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, or TAK-1 protein kinases. These compounds have the general formula I:



5

or a pharmaceutically acceptable salt thereof, wherein W, Ring A, Ring B, and R¹ are as defined below. Bonds a and b are marked in Formula I to define the orientation of Ring A. Bond a is the bond between Ring A and the bicyclic ring, including Ring B. Bond b is the bond between Ring A and R¹. Bonds a and b are also marked on examples of Ring A shown in the following section. All formula and compounds in the present application follow the orientations indicated below by these bonds.


15 [0064] These compounds and pharmaceutically acceptable compositions thereof are useful for treating or preventing a variety of diseases, disorders or conditions, including, but not limited to, cancer and other proliferative disorders.

20 [0065] The compounds provided by this invention are also useful for the study of kinases in biological and pathological phenomena; the study of intracellular signal transduction pathways mediated by such kinases; and the comparative evaluation of new kinase
25 inhibitors.

Detailed Description Of The Invention

I. General Description of Compounds of the Invention:

[0066] The present invention relates to a compound of formula I:

















5 or a pharmaceutically acceptable salt thereof, wherein:

W is CH or N;


Ring B is an optionally substituted 5-6 membered heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

10 R^1 is an optionally substituted 6-membered aryl ring having 0-3 nitrogens; and

Ring A is an optionally substituted ring selected from:

**a****b****c****d****e****f****g****h****i****j****k****l****m****n**

- 31 -



[0067] If Ring B is a 6-membered ring having no heteroatoms (i.e., is not a phenyl ring), it may form a fused benzo ring. In one embodiment of this invention, if Ring A is (b), (d), or (e), then Ring B is not a 6-membered ring having no heteroatoms (i.e., is not a phenyl ring) and thus does not form a fused benzo ring. In another embodiment, if Ring A is (b), (d), or (e), then Ring B is a pyridyl ring (i.e., Ring B and the ring fused thereto form an azaindole). In a preferred form of this embodiment, the nitrogen atoms in these azaindoles are oriented as in a 7-azaindole (see, e.g., compound 1).

[0068] In another embodiment of this invention, if Ring A is (b), then Ring A is not substituted with -SR°.

2. Compounds and Definitions:

[0069] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.:

- 32 -

Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0070] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted", whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture

- 33 -

or other chemically reactive conditions, for at least a week.

[0071] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C₃-C₈ hydrocarbon or bicyclic C₈-C₁₂ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Suitable aliphatic groups include, but are not limited to, linear or branched, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl groups and hybrids thereof such as

- 34 -

(cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0072] The term "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" as used
5 herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members are an independently selected heteroatom. In some embodiments, the "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" group has
10 three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

[0073] The term "heteroatom" means one or more of
15 oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in
20 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR^+ (as in N-substituted pyrrolidinyl)).

[0074] The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation.

[0075] The term "alkoxy", or "thioalkyl", as used
25 herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("thioalkyl") atom.

[0076] The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the
30 case may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I.

- 35 -

[0077] The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term "aryl" may be used interchangeably with the term "aryl ring". The term "aryl" also refers to heteroaryl ring systems as defined hereinbelow.

[0078] The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

[0079] An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents. Suitable substituents on the unsaturated carbon atom of an aryl or heteroaryl group are selected from oxo; halogen; $-B(OH)_2$; $-R^\circ$; $-OR^\circ$; $-SR^\circ$; aryl; heteroaryl; 1,2-methylenedioxy; 1,2-ethylenedioxy; $-CO_2(C_{1-4} \text{ aliphatic})$; an optionally substituted 5-6 membered heterocyclic ring; phenyl optionally substituted with R° ; $-O(\text{phenyl})$ optionally substituted with R° ; $-(CH_2)_{1-2}(\text{phenyl})$, optionally substituted with R° ;

- 36 -

-CH=CH(phenyl), optionally substituted with R°; -NO₂;
 -CN; -NHR°; -N(R°)₂; -NR°C(O)R°; -NR°C(S)R°;
 -NR°C(O)N(R°)₂; -NR°C(S)N(R°)₂; -NR°CO₂R°;
 -NR°NR°C(O)R°; -NR°NR°C(O)N(R°)₂; -NR°NR°CO₂R°;
 5 -C(O)C(O)R°; -C(O)CH₂C(O)R°; -CO₂R°; -C(O)R°; -C(S)R°;
 -C(O)N(R°)₂; -C(S)N(R°)₂; -OC(O)N(R°)₂; -OC(O)R°;
 -C(O)N(OR°)R°; -C(NOR°)R°; -S(O)₂R°; -S(O)₃R°;
 -SO₂N(R°)₂; -S(O)R°; -NR°SO₂N(R°)₂; -NR°SO₂R°;
 -N(OR°)R°; -C(=NH)-N(R°)₂; or -(CH₂)₀₋₂NHC(O)R°;

10 wherein each independent occurrence of
 R° is selected from hydrogen, optionally substituted
 C₁₋₆ aliphatic, an optionally substituted 5-9 membered
 heteroaryl or heterocyclic ring, phenyl, -O(phenyl), or
 -CH₂(phenyl), an optionally substituted -O(5-6 membered
 15 heterocyclic ring), an optionally substituted -CO(5-6
 membered heterocyclic ring), or an optionally
 substituted -CH₂(5-6 membered heterocyclic ring); or,
 notwithstanding the definition above, two independent
 occurrences of R°, on the same substituent or different
 20 substituents, taken together with the atom(s) to which
 each R° group is bound, form a 5-8-membered
 heterocyclyl, aryl, or heteroaryl ring or a
 3-8-membered cycloalkyl ring having 0-3 heteroatoms
 independently selected from nitrogen, oxygen, or
 25 sulfur.

[0080] Optional substituents on the aliphatic group
 of R° are selected from aryl, phenyl, heteroaryl, NH₂,
 NH(C₁₋₄aliphatic), N(C₁₋₄aliphatic)₂, NH(CH₂)phenyl,
 halogen, -NHSO₂(C₁₋₄ aliphatic), C₁₋₄aliphatic, OH,

- 37 -

O(C₁₋₄aliphatic), NO₂, CN, CO₂H, an optionally substituted -CO(5-6 membered heterocyclic ring), an optionally substituted 5-6 membered heterocyclic ring, CO₂(C₁₋₄aliphatic), O(haloC₁₋₄ aliphatic), or

5 halo(C₁₋₄aliphatic), wherein each of the foregoing C₁₋₄aliphatic groups of R° is unsubstituted.

[0081] An aliphatic or heteroaliphatic group, or a non-aromatic heterocyclic ring may contain one or more substituents. Suitable substituents on the saturated
 10 carbon of an aliphatic or heteroaliphatic group, or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and additionally include the following: =O, =S, =NNHR*, =NN(R*)₂, =NNHC(O)R*,
 15 =NNHCO₂(alkyl), =NNHSO₂(alkyl), heterocyclic ring, -OH, -CH₂OH, NHR*, N(R*)₂, CO(heterocyclic ring), R*, NHSO₂R* or =NR*, wherein each R* is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic.

[0082] Optional substituents on the aliphatic group
 20 of R* are selected from 5-6 membered heterocyclic ring, heteroaryl, aryl, NH₂, NHSO₂R*, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), CO(5-6 membered heterocyclic ring), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic),
 25 or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄aliphatic groups of R* is unsubstituted.

[0083] Optional substituents on the nitrogen of a non-aromatic heterocyclic ring are selected from -(C₁₋₆

- 38 -

aliphatic)₂, -R⁺, -N(R⁺)₂, -C(O)R⁺, -CO₂R⁺, -C(O)C(O)R⁺,
 -C(O)CH₂C(O)R⁺, -SO₂R⁺, -SO₂N(R⁺)₂, -C(=S)N(R⁺)₂,
 -C(=NH)-N(R⁺)₂, or -NR⁺SO₂R⁺; wherein R⁺ is hydrogen, an
 optionally substituted C₁₋₆ aliphatic, optionally
 5 substituted phenyl, optionally substituted -O(phenyl),
 optionally substituted -CH₂(phenyl), optionally
 substituted -(CH₂)₁₋₂(phenyl); optionally substituted
 -CH=CH(phenyl); or an unsubstituted 5-6 membered
 heteroaryl or heterocyclic ring having one to four
 10 heteroatoms independently selected from oxygen,
 nitrogen, or sulfur, or, notwithstanding the definition
 above, two independent occurrences of R⁺, on the same
 substituent or different substituents, taken together
 with the atom(s) to which each R⁺ group is bound, form
 15 a 5-8-membered heterocyclyl, aryl, or heteroaryl ring
 or a 3-8-membered cycloalkyl ring having 0-3
 heteroatoms independently selected from nitrogen,
 oxygen, or sulfur.

[0084] Optional substituents on the aliphatic group
 20 or the phenyl ring of R⁺ are selected from NH₂, NH(C₁₋₄
 aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic,
 OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄
 aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄
 aliphatic), wherein each of the foregoing C₁₋₄aliphatic
 25 groups of R⁺ is unsubstituted.

[0085] The term "alkylidene chain" refers to a
 straight or branched carbon chain that may be fully
 saturated or have one or more units of unsaturation and


has two points of attachment to the rest of the molecule.


[0086] As detailed above, in some embodiments, two independent occurrences of R° (or R^{+} , or any other variable similarly defined herein), are taken together with the atom(s) to which each variable is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Exemplary rings that are formed when two independent occurrences of R° (or R^{+} , or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound include, but are not limited to the following:

15 a) two independent occurrences of R° (or R^{+} , or any other variable similarly defined herein) that are bound to the same atom and are taken together with that atom to form a ring, for example, $N(R^{\circ})_2$, where both occurrences of R° are taken together with the nitrogen atom to form a piperidin-1-yl, piperazin-1-yl, or morpholin-4-yl group; and b) two independent occurrences of R° (or R^{+} , or any other variable similarly defined herein) that are bound to different atoms and are taken together with both of those atoms to form a ring, for example where a phenyl group is

20

25


substituted with two occurrences of OR° , these two occurrences of R° are taken together with the oxygen atoms to which they are bound to form a fused

6-membered oxygen containing ring: . It will be appreciated that a variety of other rings can be formed when two independent occurrences of R^o (or R^+ , or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound and that the examples detailed above are not intended to be limiting.

[0087] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

3. Description of Exemplary Compounds:

[0088] One embodiment of the present invention relates to a compound wherein a compound of formula **I**:




5 or a pharmaceutically acceptable salt thereof, wherein:

W is CH or N;

Ring B is an optionally substituted 5-6 membered heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

R¹ is an optionally substituted 6-membered aryl ring having 0-3 nitrogens; and


Ring A is an optionally substituted ring selected from:



- 42 -

o p or q.

[0089] Another embodiment of the present invention relates to of formula I:



5 or a pharmaceutically acceptable salt thereof, wherein:

W is CH or N, wherein the H is optionally replaced with (C₁-C₆)-alkyl or NH₂;

10 Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;


R¹ is:

15 an optionally substituted 6-10-membered aryl or 5-10-membered heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

20 a -(C₁-C₄ aliphatic) substituted with a -6-10-membered aryl or a 5-10-membered heteroaryl ring or a C₃-C₈ cycloaliphatic or heterocyclic ring, having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur each aliphatic and each ring being optionally substituted, or

25 an optionally substituted C₃-C₈ cycloaliphatic (preferably, cycloalkyl);

Ring A is an optionally substituted ring selected from:



each aryl or heteroaryl is optionally substituted with one or more (preferably, 0, 1, 2, or 3) R^3 groups, wherein R^3 is halogen; $-B(OH)_2$; $-R^\circ$; $-OR^\circ$; $-SR^\circ$; 1,2-methylenedioxy; 1,2-ethylenedioxy; $-CO_2(C_{1-4}$ aliphatic); an optionally substituted 5-6 membered heterocyclic ring; phenyl optionally substituted with R° ; $-O(\text{phenyl})$ optionally substituted with R° ; $-(CH_2)_{1-2}(\text{phenyl})$, optionally substituted with R° ; $-CH=CH(\text{phenyl})$, optionally substituted with R° ; $-NO_2$; $-CN$; $-NHR^\circ$; $-N(R^\circ)_2$; $-NR^\circ C(O)R^\circ$; $-NR^\circ C(S)R^\circ$; $-NR^\circ C(O)N(R^\circ)_2$; $-NR^\circ C(S)N(R^\circ)_2$; $-NR^\circ CO_2R^\circ$; $-NR^\circ NR^\circ C(O)R^\circ$; $-NR^\circ NR^\circ C(O)N(R^\circ)_2$; $-NR^\circ NR^\circ CO_2R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-CO_2R^\circ$; $-C(O)R^\circ$; $-C(S)R^\circ$; $-C(O)N(R^\circ)_2$; $-C(S)N(R^\circ)_2$; $-OC(O)N(R^\circ)_2$; $-OC(O)R^\circ$;

- 44 -

-C(O)N(OR°)R°; -C(NOR°)R°; -S(O)₂R°; -S(O)₃R°;
 -SO₂N(R°)₂; -S(O)R°; -NR°SO₂N(R°)₂; -NR°SO₂R°;
 -N(OR°)R°; -C(=NH)-N(R°)₂; or -(CH₂)₀₋₂NHC(O)R°;

wherein each independent occurrence of
 5 R° is selected from hydrogen,
 C₁₋₆ aliphatic, a 5-10 membered heteroaryl or
 heterocyclic ring, phenyl, -O(phenyl), -CH₂(phenyl), 5
 membered heterocyclic ring, ;
 wherein each group of R° is optionally substituted with
 10 J, wherein J is aryl, phenyl, heteroaryl, NH₂, NH(C₁₋₄
 aliphatic), N(C₁₋₄ aliphatic)₂, NH(CH₂)phenyl, halogen,
 -NHSO₂(C₁₋₄ aliphatic), -NHCO₂(C₁₋₄ aliphatic), C₁₋₄
 aliphatic, OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, -CO(5-6
 membered heterocyclic ring), 5-6 membered heterocyclic
 15 ring, -CO₂(C₁₋₄ aliphatic), -O(haloC₁₋₄ aliphatic), or
 halo(C₁₋₄ aliphatic), , or

wherein each group of J is optionally
 substituted with J', wherein J' is NH₂, NH(C₁₋₄
 aliphatic), N(C₁₋₄ aliphatic)₂, NH(CH₂)phenyl, halogen,
 20 -NHSO₂(C₁₋₄ aliphatic), -NHCO₂(C₁₋₄ aliphatic), C₁₋₄
 aliphatic, OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, -CO(5-6
 membered heterocyclic ring), 5-6 membered heterocyclic
 ring, -CO₂(C₁₋₄ aliphatic), -O(haloC₁₋₄ aliphatic), or
 halo(C₁₋₄ aliphatic), wherein each the J' groups is
 25 optionally substituted with C₁₋₄ aliphatic, halogen,
 wherein each of the C₁₋₄ aliphatic groups of J' is
 unsubstituted;

- 45 -

two R^o are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms
 5 independently selected from nitrogen, oxygen, or sulfur;

wherein each aliphatic or heteroaliphatic group or non-aromatic heterocyclic ring is optionally substituted with R³, =O, =S, =NNHR*,
 10 =NN(R*)₂, =NNHC(O)R*, =NNHCO₂(alkyl), =NNHSO₂(alkyl), or =NR*,

wherein each R* is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic, wherein optional substituents on the
 15 aliphatic group of R* are selected from 5-6 membered heterocyclic ring, heteroaryl, aryl, NH₂, NHSO₂R*, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), CO(5-6 membered heterocyclic ring), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic),
 20 O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄aliphatic groups of R* is unsubstituted; and


wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with -(C₁₋₆ aliphatic)₂, -R⁺, -N(R⁺)₂, -C(O)R⁺, -CO₂R⁺, -C(O)C(O)R⁺,
 25 -C(O)CH₂C(O)R⁺, -SO₂R⁺, -SO₂N(R⁺)₂, -C(=S)N(R⁺)₂, -C(=NH)-N(R⁺)₂, or -NR⁺SO₂R⁺;

wherein R⁺ is hydrogen, an optionally substituted C₁₋₆ aliphatic, optionally substituted

phenyl, optionally substituted -O(phenyl), optionally substituted -CH₂(phenyl), optionally substituted - (CH₂)₁₋₂(phenyl); optionally substituted -CH=CH(phenyl); or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, wherein optional substituents on the aliphatic group or the phenyl ring of R⁺ are selected from NH₂, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄ aliphatic groups of R⁺ is unsubstituted, or

two R⁺ are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0090] Another embodiment of the present invention relates to a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

W is CH, CNH₂ or N;

Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,


5 wherein Ring B is optionally substituted with one or more oxo, halogen, -OH, -OR^o, -NHR^o, N(R^o)₂, a 5-6 membered heterocyclic ring, -COO(C1-4 aliphatic), -B(OH)₂, -CO(5-6 membered heterocyclic ring), aryl, heteroaryl, and R^o,

10 wherein each R^o is H or C1-6 aliphatic independently optionally substituted with phenyl, NH₂, NH(C1-4 aliphatic), NH(CH₂)phenyl, N(C1-4 aliphatic)₂, heteroaryl, -NHSO₂(C1-4 aliphatic), halogen, an optionally substituted -CO(5-6 membered heterocyclic ring), an optionally substituted 5-6 membered
15 heterocyclic ring, COO aliphatic and OH;


R₁ is an optionally substituted 6-membered aryl or heteroaryl ring having 0-3 nitrogens, (C1-C4 aliphatic)-aryl ring, C1-C6
20 aliphatic,

wherein R₁ is optionally substituted with one or more halogen or -OR^o, wherein each R^o is C1-4 aliphatic;


Ring A is an optionally substituted ring
25 selected from:




a




b








c




d



e



- 48 -



wherein Ring A is optionally substituted with oxo, -OH, NH₂, or -CH₃; and

each aryl or heteroaryl is optionally substituted with one or more halogen; -R^o; -OR^o; -SR^o; 1,2-methylenedioxy; 1,2-ethylenedioxy; phenyl optionally substituted with R^o; -O(phenyl) optionally substituted with R^o; -(CH₂)₁₋₂(phenyl), optionally substituted with R^o; -CH=CH(phenyl), optionally substituted with R^o; -NO₂; -CN; -N(R^o)₂; -NR^oC(O)R^o; -NR^oC(S)R^o; -NR^oC(O)N(R^o)₂; -NR^oC(S)N(R^o)₂; -NR^oCO₂R^o; -NR^oNR^oC(O)R^o; -NR^oNR^oC(O)N(R^o)₂; -NR^oNR^oCO₂R^o; -C(O)C(O)R^o; -C(O)CH₂C(O)R^o; -CO₂R^o; -C(O)R^o; -C(S)R^o; -C(O)N(R^o)₂; -C(S)N(R^o)₂; -OC(O)N(R^o)₂; -OC(O)R^o; -C(O)N(OR^o)R^o; -C(NOR^o)R^o; -S(O)₂R^o; -S(O)₃R^o; -SO₂N(R^o)₂; -S(O)R^o; -NR^oSO₂N(R^o)₂; -NR^oSO₂R^o; -N(OR^o)R^o; -C(=NH)-N(R^o)₂; or -(CH₂)₀₋₂NHC(O)R^o;

wherein each independent occurrence of R^o is selected from hydrogen, optionally substituted C₁₋₆ aliphatic, an optionally substituted 5-9 membered heteroaryl or heterocyclic ring, phenyl, -O(phenyl),

- 49 -

-CH₂(phenyl), an optionally substituted -O(5-6 membered heterocyclic ring), or an optionally substituted -CH₂(5-6 membered heterocyclic ring);

wherein aliphatic groups of R° are
 5 optionally substituted with NH₂, NH(C₁₋₄aliphatic),
 N(C₁₋₄aliphatic)₂, halogen, C₁₋₄aliphatic, OH,
 O(C₁₋₄aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄aliphatic),
 O(haloC₁₋₄ aliphatic), or haloC₁₋₄aliphatic, wherein
 each of the foregoing C₁₋₄aliphatic groups of R° is
 10 unsubstituted, or

two R° are taken together with the
 atom(s) to which each is bound to form a 5-8-membered
 heterocyclyl, aryl, or heteroaryl ring or a
 3-8-membered cycloalkyl ring having 0-3 heteroatoms
 15 independently selected from nitrogen, oxygen, or
 sulfur;

wherein each aliphatic or
 heteroaliphatic group or non-aromatic heterocyclic ring
 is optionally substituted with =O, =S, =NNHR*,
 20 =NN(R*)₂, =NNHC(O)R*, =NNHCO₂(alkyl), =NNHSO₂(alkyl),
 heterocyclic ring, -OH, -CH₂OH, NHR*, N(R*)₂, CO
 (heterocyclic ring), R*, NHSO₂R* or =NR*,

wherein each R* is independently
 selected from hydrogen or an optionally substituted C₁₋₆
 25 aliphatic, wherein optional substituents on the
 aliphatic group of R* are selected from 5-6 membered
 heterocyclic ring, heteroaryl, aryl, NH₂, NHSO₂R*,
 NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄
 aliphatic, OH, O(C₁₋₄ aliphatic), CO(5-6 membered

- 50 -

heterocyclic ring), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄aliphatic groups of R* is unsubstituted; and

5 wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with -(C₁₋₆ aliphatic)₂, -R⁺, -N(R⁺)₂, -C(O)R⁺, -CO₂R⁺, -C(O)C(O)R⁺, -C(O)CH₂C(O)R⁺, -SO₂R⁺, -SO₂N(R⁺)₂, -C(=S)N(R⁺)₂, -C(=NH)-N(R⁺)₂, or -NR⁺SO₂R⁺;


10 wherein R⁺ is hydrogen, an optionally substituted C₁₋₆ aliphatic, optionally substituted phenyl, optionally substituted -O(phenyl), optionally substituted -CH₂(phenyl), optionally substituted -(CH₂)₁₋₂(phenyl); optionally substituted
15 -CH=CH(phenyl); or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, wherein optional substituents on the aliphatic group or the phenyl ring of R⁺ are
20 selected from NH₂, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄aliphatic groups of R⁺ is
25 unsubstituted, or

 two R⁺ are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a

- 51 -

3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0091] Another embodiment of the present invention
5 relates to a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

W is CH, CNH₂ or N;

Ring B is an optionally substituted 5-
10 or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,


wherein Ring B is optionally substituted with one or more oxo, halogen, -OH, -OR^o, -NHR^o,
15 N(R^o)₂, a 5-6 membered heterocyclic ring, -COO(C1-4 aliphatic), -B(OH)₂, -CO(5-6 membered heterocyclic ring), aryl, heteroaryl, and R^o,

wherein each R^o is H or C1-6 aliphatic independently optionally substituted with phenyl, NH₂,
20 NH(CH₂)phenyl, NH(C1-4 aliphatic), N(C1-4 aliphatic)₂, heteroaryl, COO aliphatic, -NHSO₂(C1-4 aliphatic), halogen, an optionally substituted -CO(5-6 membered heterocyclic ring), an optionally substituted 5-6 membered heterocyclic ring, and OH;

25 R₁ is an optionally substituted 6-membered aryl or heteroaryl ring having 0-3 nitrogens,

wherein R1 is optionally substituted with one or more halogen or -OR°, wherein each R° is C1-4 aliphatic;

Ring A is an optionally substituted ring
5 selected from:



wherein Ring A is optionally substituted with oxo, -OH, NH₂, or -CH₃; and

each aryl or heteroaryl is optionally substituted with one or more halogen; -R°; -OR°; -CN; NO₂; -N(R°)₂; -NR°C(O)R°; -C(O)N(R°)₂; -S(O)₂R°; or
10 -NR°SO₂R°;

wherein each independent occurrence of R° is selected from hydrogen, optionally substituted C₁₋₆ aliphatic, an optionally substituted 6-9 membered

- 53 -

heteroaryl or heterocyclic ring, or an optionally substituted $-\text{CH}_2$ (5-6 membered heterocyclic ring);

wherein aliphatic groups of R° are optionally substituted with N (unsubstituted

5 C_{1-4} aliphatic) $_2$;


wherein each aliphatic or heteroaliphatic group or non-aromatic heterocyclic ring is optionally substituted with $=\text{O}$, heterocyclic ring, NHR^* , $\text{N(R}^*)_2$, NHSO_2R^* , CO (heterocyclic ring), CH_2OH , OH
 10 and $-\text{R}^*$, wherein each R^* is H or optionally substituted C_{1-6} aliphatic;

wherein the optional substituents of the aliphatic group of R^* are selected from 5-6 membered heterocyclic ring and aryl; and

15 wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with $(\text{C}_{1-6}$ aliphatic) $_2$, $-\text{R}^+$, $-\text{C(O)R}^+$;

wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, wherein optional
 20 substituents on the aliphatic group of R^+ are CN.

[0092] Another embodiment of the present invention relates to a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

25 W is CH , CNH_2 or N;

- 54 -


Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,


5 wherein Ring B is optionally substituted with one or more oxo; chloro; bromo; fluoro; -CH₂OH; -OH; -OCH₃; CH₃; -NHCH₃; -NHCH₂CH₃; -N(CH₃)₂; -NH-CH₂-tetrahydrofuranyl, pyrrolidinyl; piperidinyl; pyrazolyl; -COO(CH₃); -B(OH)₂; phenyl; benzyl; 10 pyrindinyl; pyrimidinyl; imidazolyl; H; cyclopropyl; cyclohexyl; cyclohexenyl; -CH₂CH₃; -CH₂N(CH₃)₂; propynyl substituted with N(CH₃)₂; ethenyl; ethenyl substituted with triazolyl; -CH₂CH₂-triazolyl; NH(CH₃); NH(CH₂)phenyl; N(CH₃)₂; imidazo-1,2-e-pyridinyl 15 optionally substituted with -SO₂(CH₃), -NHSO₂(CH₃); an optionally substituted -CO(piperazinyl) or -CO(pyrrolidinyl), an optionally substituted morpholinyl or triazolyl, or OH;

R₁ is an optionally substituted 20 6-membered aryl or heteroaryl ring having 0-3 nitrogens,

wherein R₁ is optionally substituted with one or more halogen or -OR^o, wherein each R^o is C1-4 aliphatic;

25 Ring A is an optionally substituted ring selected from:





wherein Ring A is optionally substituted with oxo, -OH, NH_2 , or $-\text{CH}_3$; and

each aryl or heteroaryl is optionally substituted with one or more fluoro; chloro ; $-\text{R}^\circ$;

5 $-\text{OR}^\circ$; $-\text{CN}$; NO_2 ; $-\text{N}(\text{CH}_3)_2$; $-\text{NHCH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$, $-\text{NH}_2$, $-\text{NHC}(\text{O})\text{CH}_3$; $-\text{C}(\text{O})\text{NH}_2$; $-\text{C}(\text{O})\text{N}(\text{CH}_3)_2$ $-\text{S}(\text{O})_2\text{CH}_3$; or $-\text{NHSO}_2\text{CH}_3$;

wherein each independent occurrence of R° is selected from hydrogen, CH_3 , $-\text{CH}_2\text{N}(\text{CH}_3)_2$,
 10 optionally substituted pyridinyl, piperindinyl, diazepanyl, morpholinyl, optionally substituted 3,9-diaza-bicyclo [4.2.1] nonane, piperazinyl or pyrrolidinyl, or an optionally substituted $-\text{CH}_2(\text{morpholinyl})$ or $-\text{CH}_2(\text{piperazinyl})$;

15 wherein each aliphatic or heteroaliphatic group or non-aromatic heterocyclic ring is optionally substituted with $=\text{O}$, pyrrolidinyl, OH, NHbenzyl, NH_2 , and $-\text{CO}(\text{piperazinyl})$; and

wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with $-\text{CH}_3$, $-(\text{CH}_3)_2$, $-\text{H}$, $-\text{CH}_2\text{CH}_3$, $-\text{C}(\text{O})\text{CH}_2\text{CN}$, $-\text{C}(\text{O})\text{CH}_3$.

[0093] One embodiment of the present invention
5 relates to a compound of formula I wherein W is N.

[0094] Another embodiment of the present invention relates to a compound of formula I wherein W is CH.

[0095] Another embodiment of the present invention relates to a compound of formula I wherein W is CNH_2 .

10 [0096] According to one embodiment, the present invention relates to a compound of formula I wherein Ring B is an optionally substituted 5-membered heteroaryl ring having one nitrogen and 0-2 additional heteroatoms independently selected from nitrogen,
15 oxygen, or sulfur.

[0097] According to another embodiment, Ring B of formula I is an optionally substituted benzo ring.

[0098] According to another embodiment, Ring B of formula I is an optionally substituted 6-membered
20 heteroaryl ring having 1-3 nitrogens.

[0099] According to another embodiment, Ring B of formula I is an optionally substituted pyrido ring.

[0100] According to yet another embodiment, Ring B of formula I is an optionally substituted pyrimido
25 ring.

[0101] Another aspect of the present invention relates to a compound of formula I wherein Ring B is an optionally substituted pyrazino ring.

[0102] Another aspect of the present invention
30 relates to a compound of formula I wherein Ring B is an optionally substituted pyridazo ring.

[0103] In one embodiment of this invention, substituents on Ring B, when present, include one or

- 57 -

more oxo, halogen, -OH, -OR°, -NHR°, N(R°)₂, a 5-6
membered heterocyclic ring, -COO(C₁₋₄ aliphatic),
-B(OH)₂, -CO(5-6 membered heterocyclic ring), aryl,
heteroaryl, and R°,

5 wherein each R° is H or C₁₋₆ aliphatic
independently optionally substituted with phenyl, NH₂,
NH(C₁₋₄ aliphatic), NH(CH₂)phenyl, N(C₁₋₄ aliphatic)₂,
heteroaryl, -NHSO₂(C₁₋₄ aliphatic), halogen, an
optionally substituted -CO(5-6 membered heterocyclic
10 ring), an optionally substituted 5-6 membered
heterocyclic ring, COO aliphatic and OH.

[0104] In another embodiment of this invention,
substituents on Ring B, when present, include one or
more oxo, halogen, -OH, -OR°, -NHR°, N(R°)₂, a 5-6
15 membered heterocyclic ring, -COO(C₁₋₄ aliphatic),
-B(OH)₂, -CO(5-6 membered heterocyclic ring), aryl,
heteroaryl, and R°,

 wherein each R° is H or C₁₋₆ aliphatic
independently optionally substituted with phenyl, NH₂,
20 NH(CH₂)phenyl, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂,
heteroaryl, COO aliphatic, -NHSO₂(C₁₋₄ aliphatic),
halogen, an optionally substituted -CO(5-6 membered
heterocyclic ring), an optionally substituted 5-6
membered heterocyclic ring, and OH.

25 **[0105]** According to another embodiment of this
invention, substituents on Ring B, when present,
include one or more oxo; chloro; bromo; fluoro; -CH₂OH;
-OH; -OCH₃; CH₃; -NHCH₃; -NHCH₂CH₃; -N(CH₃)₂;
-NH-CH₂-tetrahydrofuranyl, pyrrolidinyl; piperidinyl;
30 pyrazolyl; -COO(CH₃); -B(OH)₂; phenyl; benzyl;
pyrindinyl; pyrimidinyl; imidazolyl; H; cyclopropyl;

- 58 -

cyclohexyl; cyclohexenyl; $-\text{CH}_2\text{CH}_3$; $-\text{CH}_2\text{N}(\text{CH}_3)_2$; propynyl substituted with $\text{N}(\text{CH}_3)_2$; ethenyl; ethenyl substituted with triazolyl; $-\text{CH}_2\text{CH}_2$ -triazolyl; $\text{NH}(\text{CH}_3)$;

$\text{NH}(\text{CH}_2)\text{phenyl}$; $\text{N}(\text{CH}_3)_2$; imidazo-1,2-e-pyridinyl

5 optionally substituted with $-\text{SO}_2(\text{CH}_3)$, $-\text{NHSO}_2(\text{CH}_3)$; an optionally substituted $-\text{CO}(\text{piperazinyl})$ or $-\text{CO}(\text{pyrrolidinyl})$, an optionally substituted morpholinyl or triazolyl, or OH.

[0106] In one embodiment of the present invention,
10 R^1 is an optionally substituted 6-membered aryl ring.

In another embodiment of the present invention, R^1 is an optionally substituted 6-membered heteroaryl ring having 1, 2, or 3 nitrogens.

[0107] In one embodiment of the present invention,
15 R^1 is an - (C_{1-6} aliphatic) -6-membered aryl ring. In another embodiment of the present invention, R^1 is an - (C_{1-6} aliphatic) -6-membered heteroaryl ring having 1, 2, or 3 nitrogens. In another embodiment of the present invention, R^1 is an - (C_1 aliphatic) -6-membered
20 heteroaryl ring having 1, 2, or 3 nitrogens. Preferably, R^1 is an - (C_1 aliphatic) -6-membered aryl ring.

[0108] In one embodiment of the present invention, R^1 is an optionally substituted C_{1-8} aliphatic.
25 Preferably, the aliphatic is a 5-, 6-, 7-, or 8-membered cycloaliphatic (either substituted as defined herein or unsubstituted).

[0109] Yet another aspect of the present invention relates to a compound of formula I wherein R^1 is an
30 optionally substituted phenyl ring. Examples of


- 59 -

substituents on the R¹ phenyl ring, when present, include one or more halogen and -OR^o, wherein each R^o is C₁₋₄ aliphatic. According to one embodiment of the present invention, substituents on the R¹ phenyl ring, when present, include one or more chloro, fluoro and -OCH₃.


[0110] According to another embodiment, the present invention relates to a compound of formula I wherein R¹ is an optionally substituted pyridyl or pyrimidinyl ring. Examples of substituents on the ring, when present, include one or more halogen and -OR^o, wherein each R^o is C₁₋₄ aliphatic. According to one embodiment of the present invention, substituents on the ring, when present, include one or more chloro, fluoro and -OCH₃.

[0111] Another embodiment of the present invention relates to a compound of formula I wherein Ring A is an optionally substituted ring selected from isoxazolyl, imidazolyl, triazolyl, or tetrazolyl. Examples of substituents on the ring, when present, include one or more halogen and -OR^o, wherein each R^o is C₁₋₄ aliphatic. According to one embodiment of the present invention, substituents on the ring, when present, include one or more chloro, fluoro and -OCH₃.


[0112] In another embodiment, if Ring A is an

optionally substituted ring q: , then q is substituted with halo, NO₂, OPCN. In another embodiment, q is not substituted.

[0113] Alternatively, if Ring A is an optionally


substituted ring q: , then R' is an optionally substituted 6-membered aryl ring having 0 nitrogens. In yet another alternative embodiment, the aryl ring has 3 nitrogens.

[0114] Yet another embodiment relates to a compound of formula I wherein Ring A is selected from the following rings:



[0115] According to another embodiment, Ring A is isoxazolyl.

15 [0116] According to yet another embodiment, Ring A is selected from:




20 [0117] According to one embodiment, the present invention relates to a compound of formula I wherein Ring A is unsubstituted.


[0118] According to another embodiment, the present invention relates to a compound of formula I wherein Ring A is optionally substituted with one or more oxo, -OH, -NH₂, or -CH₃.

25


[0119] In certain embodiments of the present invention, substituents (on aryl, aliphatic, heteroaryl, etc.) are depicted in the exemplified compounds. Exemplary structures of formula I are set forth in Table 1, below.

Table 1. Examples of Compounds of Formula I:






Compound


Cmpd
#

33




34


Compound

Cmpd
#


35



36




37




38


Compound

Cmpd
#


39



40




41




42


Compound

Cmpd
#


43



44




45




46


Compound

Cmpd
#


47



48




49




50


Compound

Cmpd
#


51



52




53




54


Compound

Cmpd
#


55



56




57




58


Compound

Cmpd
#


59







60




61




62

Compound	Cmpd #
	63
	64
	65
	66


Compound

Cmpd
#


67



68




69




70


Compound

Cmpd
#


71



72




73




74


Compound

Cmpd
#


75



76




77




78


Compound

Cmpd
#


79



80




81




82


Compound

Cmpd
#


83



84




85




86


Compound

Cmpd
#


87



88




89




90


Compound

Cmpd
#


91



92




93




94


Compound

Cmpd
#


95



96




97




98


Compound

Cmpd
#


99



100




101




102


Compound

Cmpd
#


103



104




105




106


Compound

Cmpd
#


107



108




109




110


Compound

Cmpd
#


111



112




113




114


Compound

Cmpd
#


115



116







117




118

Compound


Cmpd
#

[illegible]


Compound

Cmpd
#


127



128




129




130


Compound

Cmpd
#


131



132




133




134


Compound

Cmpd
#


135



136




137




138


Compound

Cmpd
#


139



140




141




142


Compound

Cmpd
#


143



144




145




146


Compound

Cmpd
#


147



148




149




150


Compound

Cmpd
#


151



152




153



154


Compound

Cmpd
#

4. General Synthetic Methodology:

[0120] The compounds of this invention may be prepared in general by methods known to those skilled in the art for analogous compounds, as illustrated by the general scheme below, and the preparative examples that follow.


Scheme I



[0121] Reagents and conditions: (a) AlCl_3 , DCM; (b) AlCl_3 , CS_2 50°C ; (c) Trichloroacetyl chloride, AlCl_3 , DCM; (d) Methanol, Et_3N ; (e) (i) LHMDs, aryl-acetic acid, THF, -78°C , 1 hr. (ii) reflux (f) Bredereck's reagent, THF; (g) *i.* hydroxyl amine hydrochloride, NaHCO_3 , THF reflux, *ii.* TsOH , THF reflux.

[0122] Scheme I above shows a general synthetic route for preparing compounds of the present invention when Ring A is isoxazolyl.


10 Scheme II




15 [0123] Reagents and conditions: (a) POCl_3 , DMF, Jones' Reagent; (b) R^1NH_2 , CDI, DMA; (c) Lawesson's reagent; (d) hydrazine; (e) $\text{CH}(\text{OEt})_3$.

[0124] Scheme II above shows a general synthetic route for preparing compounds of the present invention when Ring A is triazolyl ring (b).

20


Scheme III

- 5 **[0125]** Scheme III above shows a general synthetic route for preparing compounds of the present invention when Ring A is tetrazole ring (**d**) from compound **9**, as prepared above at Scheme II, and NaNO₂.

Scheme IV

10

- 15 **[0126]** Scheme IV above shows a general synthetic route for preparing compounds of the present invention when Ring A is triazole ring (**b**), substituted with -NH₂, from compound **9** and hydrazine.


Scheme V

20

[0127] Scheme V above shows a general synthetic route for preparing compounds of the present invention when Ring A is triazole ring (b), substituted with -OH, from compound 9 and CDI.

5

Scheme VI



[0128] Reagents and Conditions: (a) (i) trichloroacetyl chloride, AlCl_3 , CH_2Cl_2 , (ii) Et_3N , H_2O , RT (b) (i) oxalyl chloride, DMF (cat.), CH_2Cl_2 , (ii) Ar-NH_2 , Et_3N , CH_2Cl_2 ; (c) Lawesson's reagent, toluene, reflux; (d) hydrazine, EtOH & CH_2Cl_2 ; (e) triethylorthoformate, HCO_2H ; (f) Suzuki coupling.

[0129] Step (f) involves a Suzuki coupling.

Optional step (f) may be used to prepare compounds having various R groups. The conditions may be modified as known to skilled practitioners. For example, if R is bromo or iodo, the reagents that may be used in the coupling reaction include R-B(OR)_2 , 2M Na_2CO_3 , $\text{PdCl}_2(\text{dppf})$, and DMF. If R is B(OH)_2 , the reagents include Ar-X (where $\text{X} = \text{Br}, \text{I}, \text{OTf}$), 2M Na_2CO_3 , $\text{PdCl}_2(\text{dppf})$, and DMF. As would be recognized, step (f) would not be used if the desired final product was one wherein R is bromo, iodo, or B(OR)_2 .

- 97 -

[0130] Although certain exemplary embodiments are depicted and described above and herein, it will be appreciated that compounds of the invention can be prepared according to the methods described generally above using appropriate starting materials by methods generally available to one of ordinary skill in the art.

5. Uses, Formulation and Administration

[0131] The compounds and compositions described herein are generally useful for the inhibition of protein kinase activity of one or more enzymes. Further information relating to kinase structure, function and their role in disease or disease symptoms is available at the Protein Kinase Resource website (<http://kinases.sdsc.edu/html/index.shtml>).

[0132] Examples of kinases that are inhibited by the compounds and compositions described herein and against which the methods described herein are useful include, but are not limited to, c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and TAK-1 and all subtypes of these kinases (e.g., Aurora-2). The compounds and compositions of the invention are therefore also particularly suited for the treatment of diseases and disease symptoms that involve one or more of the aforementioned kinases.

[0133] The activity of a compound utilized in this invention as an inhibitor of c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 may be assayed *in vitro*, *in vivo* or in a cell line. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity or ATPase activity of activated c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit,

- 98 -

Aurora, and/or TAK-1. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1. Inhibitor binding may be measured by
5 radiolabelling the inhibitor prior to binding, isolating the inhibitor/c-Met, inhibitor/GSK3, inhibitor/JAK, inhibitor/SYK, inhibitor/KDR, inhibitor/FLT-3, inhibitor/c-Kit, inhibitor/Aurora, and/or inhibitor/TAK-1 complex and determining the
10 amount of radiolabel bound. Alternatively, inhibitor binding may be determined by running a competition experiment where new inhibitors are incubated with c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 bound to known radioligands. Detailed
15 conditions for assaying a compound utilized in this invention as an inhibitor of c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 kinase are set forth in the Examples below.

[0134] According to another embodiment, the
20 invention provides a composition comprising a compound of this invention or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier, adjuvant, or vehicle. The amount of compound in the compositions of this invention is such that is
25 effective to detectably inhibit a protein kinase, particularly c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 kinase, in a biological sample or in a patient. Preferably the composition of this invention is formulated for administration to a patient
30 in need of such composition. Most preferably, the composition of this invention is formulated for oral administration to a patient.

- 99 -

[0135] The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

[0136] The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0137] The term "detectably inhibit", as used herein means a measurable change in c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 activity between a sample comprising said composition and a c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 kinase and an equivalent sample comprising c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 kinase in the absence of said composition.

[0138] As used herein, the term "JAK" is used interchangeably with the terms "JAK kinase" and "a JAK

- 100 -

family kinase". In certain embodiments, JAK refers to JAK3 kinase.

[0139] A "pharmaceutically acceptable derivative" means any non-toxic salt, ester, salt of an ester or
5 other derivative of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

10 [0140] As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of c-Met, GSK3, JAK, SYK, KDR kinase, FLT-3, c-Kit, Aurora, and/or TAK-1.

15 [0141] Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate,
20 benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride,
25 hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate,
30 succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in

- 101 -

obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

[0142] Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and N⁺(C₁-4 alkyl)₄ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

[0143] The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed

- 102 -

oils are conventionally employed as a solvent or suspending medium.

[0144] For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides.

5 Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions
10 or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other
15 commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of
20 formulation.

[0145] The pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or
25 solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried
30 cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain

- 103 -

sweetening, flavoring or coloring agents may also be added.

[0146] Alternatively, the pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0147] The pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[0148] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[0149] For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable

- 104 -

compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not
5 limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0150] For ophthalmic use, the pharmaceutically acceptable compositions may be formulated as micronized
10 suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable
15 compositions may be formulated in an ointment such as petrolatum.

[0151] The pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared
20 according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other
25 conventional solubilizing or dispersing agents.

[0152] Most preferably, the pharmaceutically acceptable compositions of this invention are formulated for oral administration.

[0153] The amount of the compounds of the present
30 invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the

- 105 -

compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

5 **[0154]** It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of
10 administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular
15 compound in the composition.

[0155] According to one embodiment, the invention relates to a method of inhibiting protein kinase activity in a biological sample comprising the step of contacting said biological sample with a compound of
20 this invention, or a composition comprising said compound.

[0156] According to another embodiment, the invention relates to a method of inhibiting c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1
25 kinase activity in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a composition comprising said compound.

[0157] The term "biological sample", as used herein,
30 includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

- 106 -

[0158] Inhibition of protein kinase, or a protein kinase selected from c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 kinase, activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, biological specimen storage, and biological assays.

[0159] Another embodiment of the present invention relates to a method of inhibiting protein kinase activity in a patient comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound.

[0160] According to another embodiment, the invention relates to a method of inhibiting c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, and/or TAK-1 kinase activity in a patient comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound.

[0161] The term "cMET-mediated disease" or "cMET-mediated condition", as used herein, means any disease state or other deleterious condition in which cMET is known to play a role. The terms "cMET-mediated disease" or "cMET-mediated condition" also mean those diseases or conditions that are alleviated by treatment with a cMET inhibitor. Such conditions include, without limitation, atherosclerosis, lung fibrosis, glioblastomas, gastric carcinomas, or a cancer selected from renal, colon, breast, prostate, liver, pancreatic, or lung cancer.

[0162] According to one embodiment, the present invention relates to a method of treating or lessening the severity of renal, colon, breast, prostate, or lung

- 107 -

cancer, atherosclerosis or lung fibrosis in a patient in need thereof, comprising administering to said patient a compound of the present invention or composition thereof.

5 **[0163]** According to another embodiment, the present invention relates to a method of treating or lessening the severity of renal cancer in a patient in need thereof, comprising administering to said patient a compound of the present invention or composition
10 thereof.

[0164] Another aspect of the present invention relates to a method of inhibiting tumor metastasis in a patient in need thereof, comprising administering to said patient a compound of the present invention or
15 composition thereof.

[0165] The term "GSK3-mediated disease" or "condition", as used herein means any disease or other deleterious condition in which GSK3 is known to play a role. Accordingly, another embodiment of the present
20 invention relates to treating or lessening the severity of one or more diseases in which GSK3 is known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from autoimmune disease,
25 an inflammatory disease, a metabolic disorder, a psychiatric disorder, diabetes, an angiogenic disorder, tauopathy, a neurological or neurodegenerative disorder, a spinal cord injury, glaucoma, baldness, or a cardiovascular disease wherein said method comprises
30 administering to a patient in need thereof a composition according to the present invention.

[0166] According to another embodiment, the present invention relates to a method for treating or lessening

- 108 -

the severity of a disease or condition selected from allergy, asthma, diabetes, Alzheimer's disease, Huntington's disease, Parkinson's disease, AIDS-associated dementia, amyotrophic lateral sclerosis (ALS, Lou Gehrig's disease), multiple sclerosis (MS), an injury due to head trauma, schizophrenia, anxiety, bipolar disorder, tauopathy, a spinal cord or peripheral nerve injury, myocardial infarction, cardiomyocyte hypertrophy, glaucoma, attention deficit disorder (ADD), depression, a sleep disorder, reperfusion/ischemia, stroke, an angiogenic disorder, or baldness, wherein said method comprises administering to a patient in need thereof a compound of the present invention or composition thereof.

15 [0167] According to a preferred embodiment, the method of the present invention relates to treating or lessening the severity of stroke.

[0168] According to another preferred embodiment, the method of the present invention relates to treating or lessening the severity of a neurodegenerative or neurological disorder.

[0169] Another aspect of the present invention relates to a method of decreasing sperm motility in a male patient comprising administering to said patient a compound of the present invention or composition thereof.

[0170] In other embodiments, the invention relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula I. This method is especially useful for diabetic patients.

- 109 -

[0171] According to another embodiment, the invention provides a method for treating or lessening the severity of a JAK-mediated disease or condition in a patient comprising the step of administering to said
5 patient a composition according to the present invention.

[0172] The term "JAK-mediated disease", as used herein means any disease or other deleterious condition in which a JAK family kinase is known to play a role.
10 Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which JAK is known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a
15 disease or condition selected from immune responses such as allergic or type I hypersensitivity reactions, asthma, autoimmune diseases such as transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, and multiple
20 sclerosis, neurodegenerative disorders such as Familial amyotrophic lateral sclerosis (FALS), as well as in solid and hematologic malignancies such as leukemias and lymphomas, wherein said method comprises administering to a patient in need thereof a
25 composition according to the present invention.

[0173] According to another embodiment, the invention provides a method for treating or lessening the severity of a SYK-mediated disease or condition in a patient comprising the step of administering to said
30 patient a composition according to the present invention.

[0174] The term "SYK-mediated disease", as used herein means any disease or other deleterious condition

- 110 -

in which a SYK family kinase is known to play a role. Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which SYK is known to play a
5 role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from allergic disorders, especially asthma.

[0175] According to another embodiment, the
10 invention provides a method for treating or lessening the severity of a KDR-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to the present invention.

15 [0176] The term "KDR-mediated disease", as used herein means any disease or other deleterious condition in which a KDR family kinase is known to play a role. Accordingly, another embodiment of the present invention relates to treating or lessening the severity
20 of one or more diseases in which KDR is known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from cancer such as brain cancer, genitourinary tract cancer, lymphatic system
25 cancer, stomach cancer, cancer of the larynx, lung cancer, pancreatic cancer, breast cancer, Kaposi's sarcoma, and leukemia; endometriosis, benign prostatic hyperplasia; vascular diseases such as restenosis and atherosclerosis; autoimmune diseases such as rheumatoid
30 arthritis and psoriasis; ocular conditions such as proliferative or angiogenic retinopathy and macular degeneration; and inflammatory diseases such as contact dermatitis, asthma and delayed hypersensitivity

- 111 -

reactions.

[0177] According to another embodiment, the invention provides a method for treating or lessening the severity of a FLT-3-mediated disease or condition in a patient comprising the step of administering to
5 said patient a composition according to the present invention.

[0178] The term "FLT-3-mediated disease", as used herein means any disease or other deleterious condition in which a FLT-3 family kinase is known to play a role.
10 Such conditions include, without limitation, hematopoietic disorders, in particular, acute-myelogenous leukemia (AML), acute-promyelocytic leukemia (APL), and acute lymphocytic leukemia (ALL).

[0179] According to another embodiment, the invention provides a method for treating or lessening the severity of a FMS-mediated disease or condition in a patient comprising the step of administering to said
15 patient a composition according to the present invention.

[0180] The term "FMS-mediated disease", as used herein means any disease or other deleterious condition in which a FMS family kinase is known to play a role.
Such conditions include, without limitation, cancer
25 (including, but not limited to, ovarian, endometrial, and breast cancer), inflammatory disorders, and hypertension.

[0181] According to another embodiment, the invention provides a method for treating or lessening the severity of a c-KIT-mediated disease or condition in a patient comprising the step of administering to
30 said patient a composition according to the present invention.

- 112 -

[0182] The term "c-KIT-mediated disease", as used herein means any disease or other deleterious condition in which a c-KIT family kinase is known to play a role. Such conditions include, without limitation, AML, chronic myelogenous leukemia (CML), mastocytosis, anaplastic large-cell lymphoma, ALL, gastrointestinal stromal tumor (GIST), T-cell lymphoma, adenoid cystic carcinoma, angiosarcoma, endometrial carcinoma, small cell lung carcinoma, prostate cancer, ovarian cancer, breast carcinoma, thyroid carcinoma, malignant melanoma and colon carcinoma.

[0183] According to another embodiment, the invention provides a method for treating or lessening the severity of an AUR-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to the present invention.

[0184] The term "AUR-mediated disease" or "AUR-mediated condition", as used herein, means any disease or other deleterious condition in which AUR protein kinase is known to play a role. Such conditions include, without limitation, allergic disorders, especially asthma.

[0185] According to another embodiment, the invention provides a method for treating or lessening the severity of a TAK-1-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to the present invention.

[0186] The term "TAK-1-mediated condition", as used herein means any disease or other deleterious condition in which TAK-1 is known to play a role. The terms "TAK-1-mediated disease" or "TAK-1-mediated condition"

- 113 -

also mean those diseases or conditions that are alleviated by treatment with an TAK inhibitor. Such conditions include, without limitation, autoimmune, inflammatory, proliferative, and hyperproliferative diseases, rheumatoid arthritis, heart failure, osteoporosis, hepatic cancer, neurite outgrowth, adipogenesis, and cardiomyocyte differentiation.

[0187] Depending upon the particular condition, or disease, to be treated, additional therapeutic agents, which are normally administered to treat that condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[0188] For example, chemotherapeutic agents or other anti-proliferative agents may be combined with the compounds of this invention to treat proliferative diseases and cancer. Examples of known chemotherapeutic agents include, but are not limited to, Gleevec™, adriamycin, dexamethasone, vincristine, cyclophosphamide, fluorouracil, topotecan, taxol, interferons, and platinum derivatives.

[0189] Other examples of agents the inhibitors of this invention may also be combined with include, without limitation: treatments for Alzheimer's Disease such as Aricept® and Exelon®; treatments for Parkinson's Disease such as L-DOPA/carbidopa, entacapone, ropinrole, pramipexole, bromocriptine, pergolide, trihexephendyl, and amantadine; agents for treating Multiple Sclerosis (MS) such as beta interferon (e.g., Avonex® and Rebif®), Copaxone®, and

mitoxantrone; treatments for asthma such as albuterol and Singulair®; agents for treating schizophrenia such as zyprexa, risperdal, seroquel, and haloperidol; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; and agents for treating immunodeficiency disorders such as gamma globulin.

[0190] Those additional agents may be administered separately from the compound-containing composition, as part of a multiple dosage regimen. Alternatively, those agents may be part of a single dosage form, mixed together with the compound of this invention in a single composition. If administered as part of a multiple dosage regime, the two active agents may be submitted simultaneously, sequentially or within a period of time from one another normally within five hours from one another.

[0191] The amount of both, the compound and the additional therapeutic agent (in those compositions

- 115 -

which comprise an additional therapeutic agent as described above)) that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode
5 of administration. Preferably, the compositions of this invention should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of a compound of formula I can be administered.

[0192] In those compositions which comprise an
10 additional therapeutic agent, that additional therapeutic agent and the compound of this invention may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions will be less than that required in a monotherapy utilizing
15 only that therapeutic agent. In such compositions a dosage of between 0.01 - 100 mg/kg body weight/day of the additional therapeutic agent can be administered.

[0193] The amount of additional therapeutic agent present in the compositions of this invention will be
20 no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about
25 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[0194] The compounds of this invention, or pharmaceutical compositions thereof, may also be
30 incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents and catheters. Vascular stents, for example, have been

- 116 -

used to overcome restenosis (re-narrowing of the vessel wall after injury). However, patients using stents or other implantable devices risk clot formation or platelet activation. These unwanted effects may be prevented or mitigated by pre-coating the device with a pharmaceutically acceptable composition comprising a kinase inhibitor. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition. Implantable devices coated with a compound of this invention are another embodiment of the present invention.

[0195] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

Synthetic Examples

[0196] As used herein, the term " R_t (min)" refers to the HPLC retention time, in minutes, associated with the compound. Unless otherwise indicated, the HPLC

method utilized to obtain the reported retention time is as follows:

Column: YMC Pro C18 S-5 120Acolumn, 2.0 x 50 mm


5 Gradient: 10-90% acetonitrile+water (0.1% Formic acid)

Flow rate: 1.0 mL/minute

Detection: 225 nm.

Preparation of azaindoles


10 General method A:



[0197] Reagents and conditions: (a) AlCl_3 , DCM; (b) AlCl_3 , CS_2 , 50°C ; (c) Trichloroacetyl chloride, AlCl_3 , DCM; (d) Methanol, Et_3N ; (e) (i) LHMDs, aryl-acetic acid, THF, -78°C , 1 hr. (ii) reflux (f) Bredereck's reagent, THF; (g). i. hydroxyl amine hydrochloride, NaHCO_3 , THF reflux, ii. TsOH , THF reflux.

Example 1

- 118 -



5 2-(2,3-Difluoro-phenyl)-1-(1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone:


Method A: (X = F)

[0198] To 7-azaindole (1 g, 8.5 mmol) and AlCl₃ (1.2 g, 9.0 mmol) in methylene chloride at 0°C was added (2,3-difluorophenyl)-acetyl chloride [prepared by
10 treating (2,3-difluoro-phenyl)-acetic acid (1.5 mg, 8.72 mmol) with oxalyl chloride (0.90 mL)] in methylene chloride. After stirring at room temperature for 2 hours, the solution was poured into ice water and extracted with methylene chloride, dried (Na₂SO₄), and
15 concentrated to give 300 mg (13% yield) of title compound used without purification. LCMS R_t = 3.00 minutes, MH⁺ 273.1, M⁻ 271.1.

Method B: (X = Cl)


[0199] To 7-azaindole (173 mg, 1.46 mmol), AlCl₃ (1.34 g, 11mmol) in carbon disulfide at 50°C, was added
20 (2,3-dichloro-phenyl)-acetyl chloride [prepared by treating (2,3-dichloro-phenyl)-acetic acid (300 mg, 1.46 mmol) with oxalyl chloride (0.14 mL)] drop wise in CS₂. After heating for 3 hours, the solution was
25 poured into water and extracted with ethyl acetate, dried (Na₂SO₄), and concentrated to give 303mg (68% yield) of title compound used without purification. LCMS: R_t = 3.88 mins.; m/e 305.1 (M+H), 303.1 (M-H).

- 119 -

**2-Phenyl-1-(1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone**

[0200] Modification of Method A: To a mixture of 1H-pyrrolo[2,3-b]pyridine (1.0 g, 8.46 mmol) and AlCl₃ (3.4 g, 25.50 mmol) in dry methylene chloride (20 mL) was added phenyl acetyl chloride (3.27 g, 21.15 mmol) at room temperature. The solution was then stirred at room temperature (RT) for 4 hrs. The mixture was poured into iced water and extracted with methylene chloride (3 x 20 mL). The combined organic layers were dried over MgSO₄, filtered, and evaporated. The residue was then dissolved in MeOH (20 mL) and treated with 6N NaOH (5 mL) at RT for 2 hrs. Evaporated most of the solvent, the residue was acidified with 6N HCl and extracted with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and evaporated. The residue was purified by flash column to give desired product (1.3 g, 65%). MS (ES⁻): m/e= 235.2 (M-H); LC/Method A/2.86 min.

20

Example 2

- 120 -


2- (2,3-Dichloro-phenyl) -3-dimethylamino-1- (1H-pyrrolo [2,3-b]pyridin-3-yl) -propenone

[0201] To

2- (2,3-Dichloro-phenyl) -1- (1H-pyrrolo [2,3-b]pyridin-3-yl) -ethanone (303mg, 0.991 mmol) in THF 50 mL, was added Bredereck's reagent 952 mg, 2.99 mmol) and the solution was heated to reflux overnight. Concentration afforded title compound used without purification. LCMS: R_t = 2.64 mins.; m/e 360.1 (M+H).

10

Example 3



2- (2,3-Difluoro-phenyl) -3-dimethylamino-1- (1H-pyrrolo [2,3-b]pyridin-3-yl) -propenone


15 [0202] To

2- (2,3-difluoro-phenyl) -1- (1H-pyrrolo [2,3-b]pyridin-3-yl) -ethanone (125mg, 0.726 mmol) in THF 40 mL, was added Bredereck's reagent (379 mg, 2.18 mmol) and the solution was heated at reflux overnight. Concentration afforded title compound used without purification. LCMS: R_t = 2.30 mins.; m/e 328.2 (M+H), 326.2 (M-H).

20

Example 4

- 121 -




3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (1)

[0203] To

5 2-(2,3-dichloro-phenyl)-3-dimethylamino-1-(1H-pyrrolo[2,3-b]pyridin-3-yl)-propenone (358 mg, 0.997 mmol) in THF 100 mL, was added hydroxyl amine hydrochloride (76 mg, 1.0 mmol) and NaHCO₃ (84 mg, 1.0 mmol) and the reaction mixture was refluxed for 4 hours. To the red
10 solution was added p-toluene sulfonic acid (189 mg, 0.99 mmol) and the solution was heated for an additional 2 hours. The reaction was poured into water, extracted with ethyl acetate, washed with brine, dried (Na₂SO₄) and concentrated. Purification by flash
15 chromatography (0 to 6% methanol in methylene chloride) to afford the title compound (157 mg, 48 % yield). ¹H NMR (500 MHz, DMSO-d₆) δ 12.38 (1H, bs), 8.85 (1H, s), 8.34-8.32 (1H, d), 7.89-7.87 (1H, d), 7.76-7.75 (1H, d), 7.51-7.50 (1H, d,d), 7.36-7.33 (1H, m), 7.31-7.28
20 (1H, m) 7.18-7.15 (1H, m).

LC/MS: R_t = 3.64 mins.; m/e 298.1 (M+H), 296.1 (M-H).

Example 5



- 122 -

3-[4-(2,3-Dichloro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (2)

[0204] To

2-(2,3-dichloro-phenyl)-3-dimethylamino-1-(1H-pyrrolo[2,3-b]pyridin-3-yl)-propenone (238 mg, 0.727 mmol) in THF
5 75 mL, was added hydroxyl amine hydrochloride (56 mg, 0.80 mmol) and NaHCO₃ (67 mg, 0.80 mmol) and the reaction mixture was refluxed for 4 hours. To the red solution was added p-toluene sulfonic acid (152 mg,
10 0.80 mmol) and the solution was heated for an additional 2 hours. The reaction was poured into water, extracted with ethyl acetate, washed with brine, dried (Na₂SO₄) and concentrated. Purification by flash chromatography (0 to 6% methanol in methylene chloride
15 to afford the title compound (77 mg, 36 % yield). ¹H NMR (500 MHz, DMSO-d₆) δ 12.32 (1H, bs), 8.80 (1H, s), 8.33-8.32 (1H, d,d), 7.94-7.93 (1H, d,d), 7.77-7.75 (1H, d,d), 7.52-7.50 (1H, d,d), 7.49-7.48 (1H, d), 7.47-7.44 (1H, m) 7.19-7.16 (1H, d,d).
20 LC/MS: R_t = 2.36 mins.; m/e 330.09 (M+H), 328.05 (M-H).


[0205] Various methods for the formation of the isoxazole ring were applied:

- 25 1. Hydroxylamine hydrochloride (5 equiv.), sodium acetate (6 equiv.), ethanol, reflux.
2. (a) Hydroxylamine hydrochloride, NaHCO₃, THF, reflux; (b) p-toluenesulfonic acid, THF, reflux.
- 30 3. Hydroxylamine hydrochloride, K₂CO₃, ethanol, reflux.

[0206] The following examples 6-21 were prepared by methods described above (Examples 1-5). A number of

starting 5-substituted 7-azaindole derivatives were prepared by similar methods as described in the literature (Heterocycles 1999, 50 (2), 1065; Tetrahedron Letters 1998, 39, 5355).


5

Example 6

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-indole (29)


[0207] M⁺ 297.1; M⁻ 295.2; R_t = 4.06 minutes; ¹H NMR (DMSO-d₆) δ 11.80 (s, 1H), 8.80 (s, 1H), 7.60 (d, 1H), 7.59 (d, 1H), 7.50 (m, 2H), 7.35 (m, 1H), 7.29 (m, 1H), 7.21 (t, 1H), 7.10 (t, 1H). LC/MS: R_t 4.06 mins.; m/e 297.1 (M+H), 295.2 (M-H).

15

Example 7

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-7-fluoro-1H-indole (33)


20 [0208] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.37 (1H, bs), 8.85 (1H, s), 7.655-7.650 (1H, d), 7.51-7.50 (1H, q), 7.39-7.37 (1H, d), 7.34-7.28 (2H, cm), 7.09-7.06 (2H, m). LC/MS: R_t 4.24 mins.; m/z 315.06 (M+H), 313.17 (M-H).

Example 8

5 3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-pyridin-4-yl-1H-indole (34)

[0209] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.13 (s, 1H),
8.88 (s, 1H), 8.78 (d, 2H), 8.03 (s, 1H), 8.01 (d, 2H),
7.80 (m, 2H), 7.70 (d, 1H), 7.51 (m, 1H), 7.35 (m, 2H)

10 LC/MS: Rt 2.3 mins.; m/z 374.0 (M+H), 372.1 (M-H).

Example 9

3-[4-Phenyl-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine
(35)


15 [0210] To a solution of
2-phenyl-1-(1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone
(200 mg, 0.85 mmol) in dry THF (5 mL) was added
tert-butoxy-*N,N,N',N'*-tetramethyl-methanediamine
(Bredereck's reagent) (440 mg, 2.52 mmol). The
20 solution was heated at 60 °C for 3 h and evaporated to
dry. The resulting residue was dissolved in ethanol (5
mL). To this ethanol solution was then added
hydroxylamine hydrochloride (300 mg, 4.32 mmol) and
sodium acetate (420 mg, 5.12 mmol). The mixture was
25 heated under reflux for 10 h, cooled, and poured into

- 125 -

aqueous NaHCO₃ solution. The crude product was collected by filtration and washed with water. After purification by Gilson HPLC, the pure product was obtained as powder (130 mg, 0.50 mmol, 59%).


5 [0211] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.34 (s, 1H), 8.88 (s, 1H), 8.11 (dd, 1H), 7.84 (d, 1H), 7.75 (dd, 1H), 7.50 (m, 2H), 7.43 (m, 2H), 7.37 (m, 1H), 7.11 (dd, 1H). LC/MS: Rt 3.19 mins.; m/e 262.2 (M+H), 260.2 (M-H)

10

Example 10

3-[4-(2,3-Difluoro-benzyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (36)

15 [0212] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.40 (s, 1H), 8.43 (s, 1H), 8.35 (m, 2H), 8.02 (d, 1H), 7.31 (m, 1H), 7.26 (dd, 1H), 7.14 (m, 1H), 7.06 (t, 1H), 4.14 (s, 2H). LC/MS: Rt 3.55 mins.; m/e 312.2 (M+H), 310.2 (M-H).

Example 11


20

5-Bromo-3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (37)

- 126 -

[0213] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.67 (s, 1H), 8.88 (s, 1H), 8.41 (d, 1H), 7.89 (d, 1H), 7.88 (d, 1H), 7.54 (m, 1H), 7.33 (m, 2H). LC/MS: Rt 4.00mins.; m/e 376 (M+H), 374 (M-H).


5

Example 12

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-methoxy-1H-pyrrolo[2,3-b]pyridine (38)

[0214] The Friedel-Craft reaction of aluminum chloride, 2,3-difluorophenylacetyl chloride, and 5-methoxy-7-azaindole was run at 0 C. The remaining procedures were carried out as described for Example 2-5.

[0215] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.31 (s, 1H), 8.85 (s, 1H), 8.07 (d, 1H), 7.76 (d, 1H), 7.51 (m, 1H), 7.32 (m, 2H), 7.21 (d, 1H), 3.69 (s, 3H). LC/MS: Rt 3.5 mins.; m/e 328 (M+H), 326.1 (M-H).


Example 13

20 3-[4-(2,3,5-Trifluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (39)

[0216] ^1H NMR (500 MHz, acetone- d_6) δ : 8.69 (d, J = 1.2 HZ, 1H), 8.44 (dd, J = 4.8, 1.2 HZ, 1H), 8.24 (dd, J =

- 127 -

8.0, 1.2 Hz, 1H), 7.97 (s, 1H) 7.33 (m, 2H), 7.24 (m, 1H). LC/MS: Rt 3.5 mins.; m/e 328 (M+H), 326.1 (M-H).


Example 14

5

3-[4-(2,3,4-Trifluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridine (40)

[0217] ¹H NMR (500 MHz, acetone-d₆) δ: 8.63 (d, J = 1.0 Hz, 1H) 8.36 (m, 1H), 8.10 (dd, J = 8.0, 1.4 Hz, 1H) 7.83 (s, 1H) 7.40 (m, 1H), 7.29 (m, 1H) 7.20 (m, 1H). LC/MS: Rt 3.4 mins.; m/e 316 (M+H), 314.1 (M-H).

10

Example 15


15

3-[4-(2,3,6-Trifluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridine (41)

[0218] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.45 (s, 1H), 8.87 (s, 1H), 8.35 (dd, 1H), 8.03 (dd, 1H), 7.71 (d, 1H), 7.65 (ddd, 1H), 7.32 (dddd, 1H), 7.21 (dd, 1H). LC/MS: Rt 3.38 mins.; m/e 316 (M+H), 314.1 (M-H).

20


- 128 -

Example 16

3-[4-(3-Chloro-2-fluoro-phenyl)-isoxazol-5-yl]-1H-pyrro
5 lo[2,3-b]pyridine (42)

[0219] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.38 (s, 1H),
8.85 (s, 1H), 8.33 (d, 1H), 7.88 (d, 1H), 7.73 (d, 1H),
7.66 (t, 1H), 7.51 (t, 1H), 7.30 (t, 1H), 7.16 (dd,
1H). LC/MS: Rt 3.37 mins.; m/e 314 (M+H), 312 (M-H).

10


Example 17

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,
3-b]pyridin-5-ol (43)

15 [0220] The 5-hydroxy derivative was obtained by the
Friedel-Craft reaction of aluminum chloride,
2,3-difluorophenylacetyl chloride and
5-methoxy-7-azaindole at room temperature. The
remaining procedures were carried out as described in
20 Examples 2-5 to yield the title product.

[0221] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.11 (s, 1H),
9.40 (br, 1H), 8.81 (s, 1H), 7.94 (d, 1H), 7.60 (d,
1H), 7.49 (m, 1H), 7.30 (m, 3H). LC/MS: Rt 2.98 mins.;
m/e 314 (M+H), 312.1 (M-H).

- 129 -


Example 18

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-chloro-1H-pyrrolo[2,3-b]pyridine (44)

[0222] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.67 (1H, s), 8.88 (1H, s), 8.35 (1H, d), 7.89 (1H, s), 7.79 (1H, d), 7.54 (1H, m), 7.37-7.28 (m, 2H).

LC/MS: Rt 4.00 mins.; m/e 331.9 (M+H), 330 (M-H).


10

Example 19

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-fluoro-1H-pyrrolo[2,3-b]pyridine (45)


15 [0223] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.59 (1H, s); 8.87 (1H, s); 8.34 (1H, s); 7.87, (1H, d); 7.61-7.59 (1H, m); 7.54-7.45 (1H, m); 7.39-7.1, (2H, m); 2.43 (3H, s). LC/MS: Rt 3.69 mins.; m/e 316.2 (M+H), 314.1 (M-H).

- 130 -

Example 20

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-methyl-1H-pyrrolo[2,3-b]pyridine (46).


5 [0224] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.24 (1H, bs), 8.83 (1H, s), 8.184-8.180 (1H, d), 7.68-7.67 (2H, m), 7.52-7.51 (1H, cm), 7.35-7.29 (2H, cm), 2.38 (3H, s). LC/MS: Rt 3.6mins.; m/e 312.0 (M+H), 310.1 (M-H).

Example 21

10

3-(4-Cyclohexyl-isoxazol-5-yl)-1H-pyrrolo[2,3-b]pyridine (47)

15 [0225] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.38 (s, 1H), 8.59 (s, 1H), 8.36 (dd, 1H), 8.28 (dd, 1H), 7.94 (d, 1H), 7.24 (dd, 1H), 2.81 (m, 1H), 1.90-1.72 (complex, 5H), 1.50-1.24 (complex, 5H). LC/MS: Rt 3.49 mins.; m/e 268.2 (M+H), 266.25 (M-H).


Example 22

- 131 -

3-[4-(2,3-Dichloro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-ol (48)

[0226] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.01 (s, 1H), 9.41 (s, 1H), 8.75 (s, 1H), 7.94 (d, J = 2.5 Hz, 1H), 7.75 (dd, J = 1.5, 7.5 Hz, 1H), 7.49 (dd, J = 1.5, 7.5 Hz, 1H), 7.45 (dd, J = 8.0, 7.5 Hz, 1H), 7.39 (d, J = 2.5 Hz, 1H), 7.30 (s, 1H). LC/MS: Rt 3.3 mins.; 345.9 (M+H), 344 (M-H).

Example 23




3-[4-(2,3-Dichloro-phenyl)-isoxazol-5-yl]-5-methoxy-1H-pyrrolo[2,3-b]pyridine (49)

[0227] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.24 (s, 1H), 8.80 (s, 1H), 8.05 (d, J = 2.5 Hz, 1H), 7.75 (dd, J = 1.5, 8.0 Hz, 1H), 7.56 (s, 1H), 7.48 (dd, J = 1.5, 7.5 Hz, 1H), 7.44 (dd, J = 8.0, 7.5 Hz, 1H), 7.17 (d, J = 2.5 Hz, 1H), 3.69 (s, 3H). LC/MS: Rt 3.9 mins.; m/e 359.9 (M+H), 358 (M-H).

Preparation of

3-(4-Pyridin-2-yl-isoxazol-5-yl)-1H-pyrrolo[2,3-b]pyridine, Example 24 (50)


Example 24



Step A: 1*H*-Pyrrolo[2,3-*b*]pyridine-3-carboxylic acid methyl ester
[0228]

5 2,2,2-Trichloro-1-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-ethanone was prepared using the procedures described in Method G. To a solution of this trichloro ketone (350 mg, 1.33 mmol) in MeOH (10 mL) was added triethylamine (2 mL) at RT. The resulting solution was stirred at RT
10 for 2 h. The solvent was evaporated under vacuum, the residue was washed with water, and the crude product was dried on the pump for direct use. (200 mg, 1.13 mmol, 85%). MS (ES⁺): *m/e*= 177.1 (M+H); LC: 2.2 min.

Example 25



Step B:


2-Pyridin-2-yl-1-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-ethanone
ne

[0229] To a solution of
20 1*H*-pyrrolo[2,3-*b*]pyridine-3-carboxylic acid methyl ester (200 mg, 1.13 mmol) and 2-pyridinyl-acetic acid hydrochloride (440 mg, 2.53 mmol) in anhydrous THF (10 mL) was added LiHMDS (1.0 M in THF, 10 mL, 10.0 mmol) at -78°C. The solution was stirred at -78°C for 30 min
25 and was allowed to warm up to RT. After stirring at RT for another 30 min, the reaction mixture was heated under reflux for 14 h. The solvent was then evaporated, the residue was taken up to ethyl acetate

- 133 -

(50 mL), and washed with aq. NaHCO₃. The organic layer was dried over MgSO₄, filtered, and evaporated. The residue was used directly for the next reaction.

Example 26



5


3-(4-Pyridin-2-yl-isoxazol-5-yl)-1H-pyrrolo[2,3-b]pyridine (50)

[0230] The residue obtained above was converted to 3-(4-pyridin-2-yl-isoxazol-5-yl)-1H-pyrrolo[2,3-b]pyridine (31 mg, 0.12 mmol) using the isoxazole formation procedures described

[0231] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.49 (s, 1H), 9.15 (s, 1H), 8.87 (d, 1H), 8.76 (d, 1H), 8.38 (dd, 1H), 8.25 (dd, 1H), 7.93 (dt, 1H), 7.80 (d, 1H), 7.41 (ddd, 1H), 7.26 (dd, 1H)

15

Example 27




3-[4-(2,3-Dimethoxy-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (51) was prepared by the procedures described for Example 24.

20

[0232] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.22 (s, 1H), 8.66 (s, 1H), 8.30 (dd, 1H), 7.93 (dd, 1H), 7.55 (d,

- 134 -

1H), 7.13 (m, 3H), 6.90 (dd, 1H), 3.87 (s, 3H), 3.58 (s, 3H). LC/MS: Rt 3.14 mins.; m/e 322.2 (M+H), 320.2 (M-H)


Example 28

5

3-(4-Pyridin-3-yl-isoxazol-5-yl)-1H-pyrrolo[2,3-b]pyridine (52) was prepared by the procedures described for Example 24.

[0233] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.40 (s, 1H), 8.97 (s, 1H), 8.71 (s, 1H), 8.57 (dd, 1H), 8.33 (dd, 1H), 7.90 (dt, 1H), 7.88 (s, 1H), 7.77 (dd, 1H), 7.45 (dd, 1H), 7.14 (dd, 1H). LC/MS: Rt 1.55 mins.; m/e 263.2 (M+H), 261.2 (M-H).

Preparation of
15 3-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-prop-2-ynyl-dimethyl-amine,
Example 30 (53)

Step A:Example 29

20

Dimethyl-[3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-prop-2-ynyl]-amine:

[0234] To a screw top tube, 8.1mg(0.0425mmoles) of CuI, 18mg(0.0256mmoles) of PdCl₂(PPh₃)₂, and 207mg (0.8483mmoles) of 5-Bromo-1H-pyrrolo[2,3-b]pyridine


25

- 135 -

were added. The dry solids were diluted with 1ml of dry DMF and a stream of N₂ was bubbled through the solution for 10 minutes and then 182.6ul (1.7mmoles) of Dimethyl-prop-2-ynyl-amine was added and the reaction was stirred in a sealed tube overnight at room temperature. The reaction was diluted with 10ml of DCM and washed with saturated ammonium chloride solution. The organic layer was separated and dried with MgSO₄, filtered, and concentrated to dryness yielding a crude material of 189mg. MS showed a M⁺ ion of 200.05 which confirmed the structure above. The crude material was taken to the next step.

Step B:

Example 30




2-(2,3-Difluoro-phenyl)-1-[5-(3-dimethylamino-prop-1-ynyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-ethanone:

[0235] 189mg of crude

Dimethyl-[3-(1H-pyrrolo[2,3-b]pyridin-5-yl)-prop-2-ynyl]-amine was stirred in 5ml of DCM with AlCl₃ (4 equivalents) for 30 minutes. Two equivalents of (2,3-Difluoro-phenyl)-acetyl chloride were added and the reaction was stirred in a sealed tube overnight. The LC/MS showed the reaction to be complete and was worked-up under standard conditions. Normal phase column chromatography yielded 101mg of 2-(2,3-Difluoro-phenyl)-1-[5-(3-dimethylamino-prop-1-ynyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-ethanone with a


- 136 -

yield over two steps of 33%. LC/MS retention time of 1.83 minutes. M^+ 354.17, M^- 352.1

Step C:Example 31

2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-[5-(3-dimethylamino-prop-1-ynyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-propenone:

[0236] Procedure as described previously. Crude material was taken to the next step. LC/MS retention time 1.99minutes. M^+ 409.18 .

Step D:Example 32

(3-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-prop-2-ynyl)-dimethylamine (53)

[0237] Procedure as described previously.

Purification by reverse phase column chromatography yielding 20mg (18.7% yield over two steps).

$^1\text{H-NMR}$ (DMSO- d_6 , 500 MHz) δ : 12.7 (s, 1H), 10.2 (s, 1H), 8.9 (s, 1H), 8.5 (s, 1H), 8.1 (m, 1H), 7.9 (m, 1H), 7.55 (m, 1H), 7.35 (m, 2H), 4.4 (s, 2H) ppm; MS (FIA)


379.35 ($M+H$); HPLC 2.16 min. LC/MS: Rt 2.16 mins.; m/e 379.35 ($M+H$), 377.3 ($M-H$).

- 137 -

Preparation of


{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b] pyridin-5-ylmethyl}-dimethyl-amine, Example 35
(54)

5

Example 33**Step A: 1H-Pyrrolo[2,3-b] pyridine-5-carbaldehyde**

[0238] To 5-Bromo-1H-pyrrolo [2,3-b] pyridine (1.5g, 7.61mmol) in THF (200 mL) at -78 C, under nitrogen, was added n-butyl lithium (2.5M in hexanes, 15.2 mmol, 6.1 mL), and the reaction solution was stirred with an overhead motor. After 1hr, the resulting orange gel was quenched with methyformate (4.56g, 76mmol) and the reaction solution was allowed to slowly warm to 23C. The solution was poured into water and extracted with ethyl acetate, dried (Na₂SO₄) to give 1H-Pyrrolo[2,3-b] pyridine-5-carbaldehyde (0.68g, 61% yield) as a yellow solid.

[0239] ¹H-NMR (DMSO-d₆, 500 MHz) δ: 12.17 (1H, bs), 10.09 (1H, s), 8.77-8.76 (1H, d), 8.49-8.48 (1H, d), 7.65-7.64 (1H, d), 6.68-6.67 (1H, d). LC/MS: Rt 2.18mins.; m/e 146.9 (M+H), 144.9 (M-H).

Example 34


25

- 138 -

Step B: Dimethyl-(1H-pyrrolo[2,3-b]pyridin-5-ylmethyl)-amine

[0240] To 1H-Pyrrolo[2,3-b]pyridine-5-carbaldehyde (114mg, 0.78mmol), in methanol (10mL) was added
 5 dimethylamine hydrochloride (127mg, 1.56mmol, NaOH (31.2 mg, 0.78mmol), sodium cyanoborohydride (49mg, 0.78mmol) and the solution was stirred under nitrogen for 12 hr. Poured into water (50mL), extracted with ethyl acetate, dried (Na₂SO₄) to afforded Dimethyl-
 10 (1H-pyrrolo[2,3-b]pyridin-5-ylmethyl)-amine (44mg, 33% yield) used without purification. MS: m/e 176.1 (M+H).

Example 35



Step C:


15 **2-(2,3-Difluoro-phenyl)-1-(5-dimethylaminomethyl-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone**

[0241] Dimethyl-(1H-pyrrolo[2,3-b]pyridin-5-ylmethyl)-amine (44mg, 0.25mmol) and AlCl₃ (167mg, 1.25mmol) in methylene chloride (10mL) were
 20 stirred for 0.5hr. To this mixture was added 2,3-Difluoro-phenyl)-acetyl chloride (96mg, 0.50mmol) and the reaction solution was stirred for 4 hr. Quenched with methanol (20mL) and water (20mL) and extracted with ethyl acetate, dried (Na₂SO₄). Flash
 25 chromatography (methylene chloride/methanol) afforded 2-(2,3-Difluoro-phenyl)-1-(5-dimethylaminomethyl-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone

- 139 -

(20mg, 24%yield). LCMS tr = 1.64min, m/z MH+ 330.1, M-328.2

Example 36



5


Step D: 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(5-dimethylaminomethyl-1H-pyrrolo [2,3-b] pyridin-3-yl) propenone

[0242] To

10 2-(2,3-Difluoro-phenyl)-1-(5-dimethylaminomethyl-1H-pyrrolo [2,3-b] pyridin-3-yl)-ethanone (20mg, 0.061mmol) in THF (5mL) was added Brederick's reagent (53mg, 0.31mmol) and the reaction was heated to 80 C in a sealed tube overnight. Concentration under reduced vacuum gave title compound a red oil used as obtained.

15 LC/MS: Rt 1.60mins.; m/e 385.2 (M+H), 358.2 (-27), 356.3 (M-H)

Example 37



20

- 140 -

Step E:

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-ylmethyl}-dimethyl-amine (54)


[0243] To 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(5-dimethylaminomethyl-1*H*-pyrrolo [2,3-*b*] pyridin-3-yl) propenone (23mg, 0.61mmol) in ethanol (5mL) was added sodium acetate (30mg, 0.37mmol) and hydroxylamine Hydrochloride (21mg, 0.30mmol) and the mixture was heated to 80C in a sealed tube. After 14hr, solution was cooled, diluted with ethyl acetate, washed with brine, dried (Na₂SO₄). Concentrated gave an amber oil. Preparative reverse phase chromatography gave 4.5mg (21% yield).

[0244] ¹H NMR (500 MHz, DMSO-*d*₆) δ: 12.60 (1H, bs), 9.65 (1H, vbs (TFA)), 8.80 (1H, s), 8.422-8.419 (1H, d), 8.382 -8.361 (1H, d), 7.778-7.771 (1H, d), 7.54-7.49 (1H, cm), 7.38-7.28 (2H, cm), 4.46-4.44 (2H, d), 2.76-2.71 (6H, d). LC/MS: Rt 1.98mins.; m/e 355.2 (M+H), 353.31 (M-H).

20 **Preparation of**

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-methanol, Example 41 (55)

Example 38



- 141 -


Step A:

1-(Toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid methyl ester

[0245] Under N₂ purge,

5 5-Bromo-1-(toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine (1.0 g, 2.8 mmol), Et₃N (0.75 mL, 5.4 mmol), Pd(OAc)₂ (64 mg, 0.28 mmol), Ph₃P (0.45 g, 1.7 mmol) and MeOH (5.0 mL, 120 mmol) were loaded in 20 mL DMF. The vessel was purged with CO 5 minutes and fixed with a condenser tube with CO balloon. The reaction was heated to 100°C for 6 hours and monitored by TLC (*r_f* 5-Bromo-1-(toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine = 0.72, *r_f* 1-(Toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid methyl ester = 0.43, CH₂Cl₂). After 6 hours, the reaction was removed from heat, diluted to 100 mL with water and extracted with EtOAc (100 mL). The phases were separated and organic was washed with additional water (2x100 mL) and brine (1x100 mL), dried over Na₂SO₄, filtered, and dried in vacuo. The resulting yellow powder was purified over a short plug of silica gel with CH₂Cl₂ until all material was recovered. Yield 0.94g, white powder.

[0246] ¹H NMR (500 MHz, CDCl₃) δ: 9.1 (s, 1H), 8.5 (s, 1H), 8.1 (d, 2H), 7.8 (d, 1H), 7.3 (d, 2H), 6.8 (d, 1H), 3.9 (s, 3H), 2.4 (s, 3H).


Example 39

Step B: 1H-Pyrrolo[2,3-b]pyridine-5-carboxylic acid methyl ester

[0247] A suspension of 1-(Toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid methyl ester (5.4 g, 16 mmol) in MeOH (100 mL) with NaOMe in MeOH (20 mL, 25% wt., excess) was heated at 65°C for 1 hour. The resulting material was concentrated from MeOH, diluted with H₂O (100 mL), and pH adjusted to 6 with 1N HCl. The aqueous solution was partitioned with EtOAc (100 mL) and the organic extraction was dried over Na₂SO₄ and dried in vacuo. The residue was purified with flash chromatography over silica gel in an elution from 99:1 CH₂Cl₂:MeOH to 92:8 CH₂Cl₂:MeOH. Yield 1.8 g, beige powder.

[0248] ¹H NMR (500 MHz, CDCl₃) δ: 9.7 (bs, 1H), 9.0 (s, 1H), 8.7 (s, 1H), 7.4 (t, 1H), 6.6 (t, 1H), 4.0 (s, 3H).

Example 40




Step C: (1H-Pyrrolo[2,3-b]pyridin-5-yl)-methanol

[0249] To 1H-Pyrrolo[2,3-b]pyridine-5-carboxylic acid methyl ester (75mg, 0.394mmol) in THF at 0 C was added lithium aluminum hydride (45mg, 1.18mmol) and the reaction mixture was slowly allowed to warm to ambient temperature. The reaction was refluxed for 12hr, allowed to cool and quenched with water. Extraction with ethyl acetate, washed with brine, dried (Na₂SO₄) to afford (1H-Pyrrolo[2,3-b]pyridin-5-yl)-methanol (57mg, 98%yield).

- 143 -

[0250] ^1H NMR (500 MHz, DMSO- d_6) δ : 11.50 (1H, bs), 8.166-8.163 (1H, d), 7.862-7.860 (1H, d), 7.43-7.42 (1H, d), 6.41-6.40 (1H, d). 5.10-5.08 (1H, t), 4.58-4.57 (2H, d). FIA MS, m/z MH^+ 149.1.

5


Example 41**Step D:**

2-(2,3-Difluoro-phenyl)-1-(5-hydroxymethyl-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone

[0251] (1H-Pyrrolo[2,3-b]pyridin-5-yl)-methanol (57mg, 0.39mmol) and AlCl_3 (160mg, 1.15mmol) in methylene chloride (5mL) were stirred for 0.5hr. To this mixture was added 2,3-Difluoro-phenyl)-acetyl chloride (219mg, 1.15mmol) and the reaction solution was stirred for 14 hr. Quenched with methanol (10mL) and water (10mL) and extracted with ethyl acetate, dried (Na_2SO_4). This afforded (2,3-Difluoro-phenyl)-acetic acid 3-[(2,3-difluoro-phenyl)-acetyl]-1H-pyrrolo[2,3-b]pyridin-5-yl (LCMS t_r = 4.04min, m/z MH^+ 457.0 M^- 455.1) that was taken up in methanol (5mL) and treated with 1N NaOH (1mL) and stirred for 3 hrs. Diluted ethyl acetate and pH adjusted to 7 with 10% NaHSO_4 . Concentration gave title compound as an off-white solid (93mg, 80% yield).

- 144 -

[0252] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.51 (1H, bs), 8.63 (1H, s), 8.41-8.40 (1H, d), 8.29-8.28 (1H, d), 7.34530 (1H, m), 7.20-7.15 (2H, cm), 5.23-5.20 (1H, q) 4.60-4.53 (2H, d), 4.39-4.38 (2H, s). LC/MS: Rt 2.55mins.; m/e 303.1 (M+H), 301.2 (M-H).

Example 42**Step E:**


10 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(5-hydroxymethyl-1H-pyrrolo[2,3-b]pyridin-3-yl)-propenone

[0253] To

2-(2,3-Difluoro-phenyl)-1-(5-hydroxymethyl-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone (93mg, 0.31mmol) in THF
15 (15mL) was added Bredereck's reagent (500 μ L, 2.4mmol) and the reaction was heated to 80 C overnight.

Concentration under reduced vacuum gave title compound a red oil, used as obtained. LC/MS: Rt 2.72mins.; m/e 359.1 (M+H), 356.2 (M-H).

20

Example 43

Step F:

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-methanol (55)


[0254] To 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(5-hydroxymethyl-1H-pyrrolo [2,3-b]pyridin-3-yl)-propenone (110mg, 0.31mmol) in tetrahydrofuran (20mL) was added sodium hydrogen carbonate (39mg, 0.46mmol) and hydroxylamine Hydrochloride (32mg, 0.46mmol) and the mixture was heated to 80C. After 5hr p-toluenesulfonic acid (catalytic amount) was added and the reaction mixture was heated for an additional 14hr. The solution was cooled, diluted with ethyl acetate, washed with brine, dried (Na₂SO₄). Concentration gave an amber oil.

Preparative reverse phase chromatography afforded {3-[4-(2-Fluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-methanol (3.0mg, 3% yield).

[0255] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.30 (1H, bs), 8.40 (1H, s), 8.295-8.292 (1H, d), 7.95 (1H, s), 7.685-7.680 (1H, d), 7.54-7.49 (1H, cm), 7.38- 7.16 (2H, cm), 7.06 (1H, t), 4.57 (2H, s). LC/MS: Rt 2.81mins.; m/e 328.1 (M+H), 326.2 (M-H).

Preparation of

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-4-yl}-dimethyl-amine, Example 44 (56):


Example 44

2-(2,3-Difluoro-phenyl)-1-(4-fluoro-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone

[0256] 4-Fluoro-1H-pyrrolo[2,3-b]pyridine (230mg, 1.69mmol) (Org. Lett. **2003**, 5(26), 5023) and AlCl₃ (678, 5.1mmol) in methylene chloride (30mL) were stirred for 0.5hr. To this mixture was added 2,3-Difluoro-phenyl)-acetyl chloride (644mg, 3.38mmol) and the reaction solution was stirred for 14hr. Quenched with methanol (50mL) and water (50mL) and extracted with ethyl acetate, dried (Na₂SO₄). Flash chromatography (methylene chloride/methanol) afforded 2-(2,3-Difluoro-phenyl)-1-(4-fluoro-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone (448mg, 91%yield). LC/MS: Rt 3.16mins.; m/e 287.1 (M+H), 285.2 (M-H).

15

Example 45



2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(4-dimethylamino-1H-pyrrolo[2,3-b]pyridin-3-yl)-propenone

[0257] To


2-(2,3-Difluoro-phenyl)-1-(4-fluoro-1H-pyrrolo[2,3-b]pyridin-3-yl)-ethanone (428mg, 1.15mmol) in THF (50mL) was added Brederick's reagent (1.6mL, 7.7mmol) and the reaction was heated to 80 C overnight. Concentration under reduced vacuum a red oil, used as obtained as a mixture (1:1) of

2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(4-dimethylam

- 147 -

ino-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-propenone and
 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(4-fluoro-1*H*-
 pyrrolo[2,3-*b*]pyridin-3-yl)-propenone LC/MS: Rt
 1.75mins.; m/e 371.0 (M+H), 342.2 (M-27) and Rt 2.58
 5 mins.; m/e 346.0 (M+H), 344.0 (M-H).

Example 46



10 {3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1*H*-pyrrolo[2,
 ,3-*b*]pyridin-4-yl}-dimethyl-amine (56)

[0258] To a mixture (1:1) of
 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(4-dimethyl-
 amino-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-propenone and
 2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-(4-fluoro-1*H*-
 15 pyrrolo[2,3-*b*]pyridin-3-yl)-propenone (110mg, 0.31mmol)
 in tetrahydrofuran (20mL) was added sodium hydrogen
 carbonate (39mg, 0.46mmol) and hydroxylamine
 Hydrochloride (32mg, 0.46mmol) and the mixture was
 heated to 80C. After 5hr p-toluenesulfonic acid
 20 (catalytic amount) was added and the reaction mixture
 was heated for an additional 4hr. The solution was
 cooled, diluted with ethyl acetate, washed with brine,
 dried (Na₂SO₄). Concentration gave an amber oil.


Preparative reverse phase chromatography afforded
 25 {3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1*H*-pyrrolo[2

,3-b]pyridin-4-yl}-dimethyl-amine (3.0mg, 3% yield).

¹H NMR (dmsO): 13.3-12.9 (1H, vbs), 9.02 (1H, s), 8.13-8.12 (1H, d), 7.75 (1H, s), 7.44-7.39 (1H, m), 7.23-7.13 (2H, cm), 6.71-6.70 (1H, d), 2.81 (6H, s).

5 LC/MS Rt 2.1mins.; m/e 341.0 (M+H), 339.1 (M-H)

General method B:




10 [0259] Reagents and Conditions: (a) (i) NaH, THF, RT, (ii) methanesulfonyl chloride; (b) R-B(OR)₂, 2M Na₂CO₃, PdCl₂(dppf), DMF; (c) (i) substituted-phenylacetyl chloride, AlCl₃, CH₂Cl₂, (ii) Brederick's Reagent, THF, reflux, (iii) H₂NOH HCl, NaOAc, THF, reflux.

15 Preparation of

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-pyridin-4-yl-1H-pyrrolo[2,3-b]pyridine, Example 47 (57)

Example 47



20 Step A:

5-Bromo-1-methanesulfonyl-1H-pyrrolo[2,3-b]pyridine


[0260] To a solution of

25 5-bromo-1H-pyrrolo[2,3-b]pyridine (600 mg, 3.06 mmol) in dry THF (30 mL) was added NaH (246 mg, 6.12 mmol) at room temperature. The suspension was stirred for 30

- 149 -

min before addition of methanesulfonyl chloride (540 mg, 4.80 mmol). The solution was stirred at room temperature for another 30 min and poured into water (50 mL). The aqueous solution was extracted with ethyl acetate (2 x 30 mL), the combined organic layers were dried over MgSO₄ and filtered, the filtrate was evaporated under vacuum to afford white solid (780 mg, 93%). The crude product was used directly for the next reaction. LC/MS: Rt 3.26 min.; (m/e= 275.0, 276.9 (M+H, M+2+H)).

Example 48




Step B: 5-Pyridin-4-yl-1H-pyrrolo [2,3-b]pyridine [0261] To a solution of 5-bromo-1-methanesulfonyl-1H-pyrrolo [2,3-b]pyridine (140 mg, 0.51 mmol) and 4-pyridinyl boronic acid (125 mg, 1.02 mmol) in DMF (4 mL) was added aqueous Na₂CO₃ (2M, 1.3 mL, 2.6 mmol). To this suspension was then added PdCl₂(dppf) (20 mg, 0.025 mmol) under N₂ atmosphere. The flask was then covered with septa, heated at 80°C for 9h, and poured into water. The precipitate was collected by filtration, washed with water, and redissolved in MeOH (10 mL). To this Methanol solution was added 6N NaOH solution (2 mL) and the resulting basic reaction mixture was heated at 50°C for 3 h. After evaporating the MeOH, the aqueous residue was acidified with 6N HCl to pH=2. The precipitate was filtered off and washed with 2N HCl.

- 150 -

The acidic filtrate was then neutralized with saturated NaHCO₃ solution. The crude product was collected by filtration, washed with water, and dried on the pump for direct use (60 mg, 0.31 mmol). LC/MS: Rt 1.96 mins.; m/e 196.1 (M+H).

Example 49




Step C:

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-pyridin-4-yl-1H-pyrrolo[2,3-b]pyridine (57)

[0262] 5-Pyridin-4-yl-1H-pyrrolo[2,3-b]pyridine (60 mg, 0.31 mmol) was converted to 3-(4-(2,3-difluoro-phenyl)-isoxazol-5-yl)-5-pyridin-4-yl-1H-pyrrolo[2,3-b]pyridine (60 mg, 0.16 mmol) by using Method A.

¹H NMR (500 MHz, DMSO-d₆) δ: 12.65 (s, 1H), 8.90 (s, 1H), 8.78 (d, 1H), 8.64 (m, 2H), 8.03 (d, 1H), 7.94 (d, 1H), 7.61 (d, 2H), 7.53 (m, 1H), 7.39 (m, 1H), 7.33 (m, 1H), (free base). LC/MS: Rt 2.5 mins.; m/e 375 (M+H), 373 (M-H).


Example 50



3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-phenyl-1H-pyrrolo[2,3-b]pyridine (58)

[0263] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.53 (s, 1H), 8.88 (s, 1H), 8.62 (d, 1H), 7.90 (d, 1H), 7.84 (d, 1H), 7.51 (m, 5H), 7.34 (m, 3H). LC/MS: Rt 4.2 mins.; m/e 374 (M+H), 372.1 (M-H).

Example 51




10 3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-pyridin-3-yl-1H-pyrrolo[2,3-b]pyridine (59)

[0264] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.58 (s, 1H), 8.89 (s, 1H), 8.75 (d, 1H), 8.67 (d, 1H), 8.58 (dd, 1H), 7.98 (m, 1H), 7.93 (m, 2H), 7.51 (m, 2H), 7.37 (dd, 1H), 7.31 (m, 1H), 1H.

LC/MS: Rt 2.5 mins.; m/e 375 (M+H), 373 (M-H).

Example 52




20 3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(1-methyl-piperidin-4-yl)-1H-pyrrolo[2,3-b]pyridine (60)

[0265] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.40 (s, 1H), 9.55 (br, 1H), 8.87 (s, 1H), 8.26 (d, 1H), 7.76 (d,

- 152 -

1H), 7.72 (d, 1H), 7.51 (m, 1H), 7.32 (m, 2H), 3.53 (d, 2H), 3.09 (m, 2H), 2.92 (m, 1H), 2.84 (d, 3H), 2.03 (d, 2H), 1.84 (m, 2H). LC/MS: Rt 2.1 mins.; m/e 395 (M+H), 393 (M-H).


5

Example 53

(3-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-phenyl)-dimethyl-amine (61)

10 [0266] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.48 (s, 1H), 8.88 (s, 1H), 8.60 (d, 1H), 7.89 (d, 1H), 7.82 (d, 1H), 7.51 (m, 1H), 7.33 (m, 2H), 7.25 (t, 1H), 6.81 (s, 1H), 6.74 (m, 2H), 2.96 (s, 6H). LC/MS: Rt 3.4 mins.; m/e 417 (M+H), 415 (M-H).

15


Example 54

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-piperidin-4-yl-1H-pyrrolo[2,3-b]pyridine (62)

20 [0267] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.40 (s, 1H), 8.86 (s, 1H), 8.66 (br, d, 1H), 8.34 (br, 1H), 8.25 (d, 1H), 7.75 (d, 1H), 7.74 (d, 1H), 7.52 (m, 1H), 7.33 (m, 2H), 3.40 (br, d, 2H), 3.00 (m, 3H), 1.93 (br, d, 2H),

- 153 -

1.78 (m, 2H). LC/MS: Rt 2.1 mins.; m/e 381 (M+H), 379 (M-H).


Example 55

5

1-(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-piperidin-1-yl)-ethanone (63)

[0268] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.35 (s, 1H), 8.85 (s, 1H), 8.24 (d, 1H), 7.84 (d, 1H), 7.48 (m, 1H), 7.40 (d, 1H), 7.32 (m, 2H), 4.51 (d, br, 1H), 3.90 (d, br, 1H), 3.11 (td, 1H), 2.85 (tt, 1H), 2.51 (td, 1H), 2.06 (s, 3H), 1.71 (t, br, 2H), 1.46-1.25 (m, 2H).
LC/MS: Rt 3.0 mins.; m/e 423 (M+H), 421 (M-H).

10

Example 56

15

3-(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-piperidin-1-yl)-3-oxo-propionitrile (64)


[0269] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.34 (s, 1H), 8.85 (s, 1H), 8.23 (d, 1H), 7.80 (d, 1H), 7.52 (d, 1H), 7.48 (m, 1H), 7.32 (m, 2H), 4.48 (d, br, 1H), 4.05 (s, 2H), 3.78 (d, br, 1H), 3.15 (td, 1H), 2.88 (tt, 1H), 2.69 (td, 1H), 1.75 (d, br, 2H), 1.56 (qd, 1H), 1.35

20

- 154 -

(qd, 1H). LC/MS: Rt 3.2 mins.; m/e 448 (M+H), 446 (M-H).


Example 57



5 N-(3-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-phenyl)-acetamide (65)

[0270] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.51 (s, 1H), 8.87 (s, 1H), 8.58 (d, 1H), 8.01 (d, 1H), 7.82 (d, overlap, 2H), 7.62 (d, 1H), 7.53 (m, 1H), 7.39 (m, 2H), 7.31 (m, 1H), 7.21 (d, 1H), 2.09 (s, 3H). LC/MS: Rt 3.5 mins; m/e 431 (M+H), 429 (M-H).


Example 58



15 3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-vinyl-1H-pyrrolo[2,3-b]pyridine (66)


[0271] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.44 (s, 1H), 8.87 (s, 1H), 8.46 (s, 1H), 7.81 (s, 1H), 7.78 (s, 1H), 7.51 (m, 1H), 7.32 (m, 2H), 6.80 (dd, J = 11.0, 17.5Hz, 1H), 5.66 (d, J = 17.5Hz, 1H), 5.23 (d, J = 11.0Hz, 1H). LC/MS: Rt 3.8 mins.; m/e 324 (M+H), 322 (M-H).

- 155 -

Example 59

5-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-1H-pyridin-2-one (67)


5 [0272] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.47 (s, 1H),
8.87 (s, 1H), 8.51 (d, $J=2.1\text{Hz}$, 1H), 7.85 (d, $J=2.7\text{Hz}$,
1H), 7.73 (d, $J=2.0\text{Hz}$, 1H), 7.67 (dd, $J=2.7$, 9.5Hz,
1H), 7.57 (d, $J=2.4\text{Hz}$, 1H), 7.54 (m, 1H), 7.32 (m, 2H),
6.45 (d, $J=9.4\text{Hz}$, 1H). LC/MS: Rt 2.8 mins.; m/e 391
10 (M+H), 389 (M-H).

Example 60

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(4-piperazi
n-1-yl-phenyl)-1H-pyrrolo[2,3-b]pyridine (68)


15 [0273] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.46 (s, 1H),
8.87 (s, 1H), 8.69 (br, 2H), 8.58 (d, 1H), 7.86 (d,
1H), 7.81 (d, 1H), 7.54 (m, 1H), 7.44 (d, $J = 8.6\text{Hz}$,
2H), 7.32 (m, 2H), 7.09 (d, $J = 8.6\text{Hz}$, 2H), 3.43 (br,
4H), 3.25 (br, 4H), 2.32 (s, 4.1H). LC/MS: Rt 2.4
20 mins; m/e 458.2 (M+H), 456.2 (M-H).

- 156 -

Example 61

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(6-Fluoro-pyridin-3-yl)-1H-pyrrolo[2,3-b]pyridine (69)


[0274] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.60 (s, 1H), 8.89 (s, 1H), 8.66 (d, 1H), 8.42 (d, 1H), 8.20 (dt, 1H), 7.96 (d, 1H), 7.91 (d, 1H), 7.52 (m, 1H), 7.35 (m, 2H), 7.29 (dd, 1H), 2.37 (s, 2.8H). LC/MS: Rt 3.8 mins.; m/e 393.1 (M+H), 391.2 (M-H).

Example 62

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-[4-(4-methyl-piperazin-1-yl)-phenyl]-1H-pyrrolo[2,3-b]pyridine methanesulfonate (70)


[0275] ^1H NMR (500 MHz, DMSO- d_6) δ : DMSO- d_6 : 12.45 (s, 1H), 8.87 (s, 1H), 8.57 (d, 1H), 7.87 (d, 1H), 7.79 (d, 1H), 7.55 (m, 1H), 7.42 (d, J = 8.5Hz, 2H), 7.34 (m, 2H), 7.07 (d, J = 8.6Hz, 2H), 3.20-3.02 (br, 4H), 2.72 (br, 4H), 2.30 (s, 6H). LC/MS: Rt 2.4 mins.; m/e 472.2 (M+H), 470.4 (M-H).

- 157 -

Example 63


3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-[6-(4-methy
 5 1-piperazin-1-yl)-pyridin-3-yl]-1H-pyrrolo[2,3-b]pyridi
 ne (71)

[0276] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.50 (s, 1H),
 9.70 (br, 1H), 8.88 (s, 1H), 8.59 (d, 1H), 8.37 (d,
 1H), 7.86 (m, 2H), 7.79 (dd, 1H), 7.52 (m, 1H), 7.33
 10 (m, 2H), 7.03 (d, 1H), 3.80 (br, 4H), 3.15 (br, 4H),
 2.75 (s, 3H), 2.31 (s, 2.8H). LC/MS: Rt 2.2 mins.;
 m/e 473.3 (M+H), 471.4 (M-H).

Example 64

15 3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(1-ethyl-pi
 peridin-4-yl)-1H-pyrrolo[2,3-b]pyridine (72)

[0277] ^1H NMR (500 MHz, CD $_3$ OD) δ : 8.60 (s, 1H),
 8.27 (d, 1H), 7.98 (d, 1H), 7.62 (s, 1H), 7.35 (m, 1H),
 7.25 (m, 2H), 3.63 (d, br, 2H), 3.18 (q, 2H), 3.06 (m,
 20 2H), 2.68 (s, 3H), 2.13 (d, br, 2H), 1.98 (m, 2H), 1.38
 (t, 3H). LC/MS: Rt 2.1 mins.; m/e 409.4 (M+H), 407.4
 (M-H).

General method C:


5 **[0278]** Reagents and Conditions: (a)

bis(pinacolato)diboron, KOAc, PdCl₂(dppf), dioxane, 80 °C;

(b) Ar-Br or Ar-I, 2M Na₂CO₃, PdCl₂(dppf), DMF, 80 °C;

(c) (i) substituted-phenylacetyl chloride, AlCl₃, CH₂Cl₂, (ii) Brederick's Reagent, THF, reflux, (iii)

10 H₂NOH HCl, NaOAc, THF, reflux.

Example 65

15 **1-Methanesulfonyl-5-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridine**

[0279] To a solution of

5-bromo-1-methanesulfonyl-1H-pyrrolo[2,3-b]pyridine

(1.40 g, 5.1 mmol), bis(pinacolato)diboron (1.42 g, 5.6 mmol), and KOAc (15.3 mmol) in dioxane (30 mL) was

20 added PdCl₂(dppf) (250 mg, 0.31 mmol) under N₂

atmosphere. The solution was heated at 80 °C for 3 h.


The solvent was removed by evaporation, the residue was taken up to hexane (50 mL), and the precipitate was

collected by filtration. The crude brownish solid was

25 used without purification. MS (ES⁺): m/e = 241.0

- 159 -

(M+H-C₄H₇); LC: 2.33 min. ¹H NMR (500 MHz, DMSO-d₆) δ: 8.62 (d, 1H), 8.38 (d, 1H), 7.74 (d, 1H), 6.83 (d, 1H), 3.72 (s, 3H), 1.34 (s, 12H) ppm.


Example 66

5

5-Pyridin-2-yl-1H-pyrrolo[2,3-b]pyridine

[0280] To a solution of 1-methanesulfonyl-5-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridine (300 mg, 0.93 mmol) and 2-bromopyridine (150 mg, 0.95 mmol) in DMF (3 mL) was added aqueous Na₂CO₃ (2M, 1.9 mL, 3.8 mmol). To this suspension was then added PdCl₂(dppf) (60 mg, 0.075 mmol) under N₂ atmosphere. The flask was then covered with septa, heated at 80°C for 9h, and poured into water (30 mL). The aqueous solution was extracted with ethylacetate (3 x 30 mL). The combined organic layers were dried over Na₂SO₄, filtered, and evaporated. The resulting residue was dissolved in MeOH (10 mL) and treated with 6N NaOH solution (2 mL) at 50°C for 3 h. After evaporating the MeOH, the aqueous residue was acidified with 6N HCl to pH=8. The precipitate was collected by filtration, washed with water, and dried on the pump for direct use (100 mg off-white solid, 0.51 mmol). MS (ES⁺): m/e= 196.1 (M+H); LC: 2.04 min.

- 160 -


Example 67

3-(4-(2,3-Difluoro-phenyl)-isoxazol-5-yl)-5-pyridin-2-yl-1H-pyrrolo[2,3-b]pyridine (73)

5 [0281] 5-Pyridin-2-yl-1H-pyrrolo[2,3-b]pyridine (100 mg, 0.51 mmol) was converted to 3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-5-pyridin-2-yl-1H-pyrrolo[2,3-b]pyridine (50 mg, 0.13 mmol) by using Method A.

10 [0282] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.60 (s, 1H), 8.89 (s, 1H), 8.69 (d, 1H), 8.34 (d, 1H), 8.31 (d, 1H), 7.95 (d, 1H), 7.93 (d, 1H), 7.57 (s, 1H), 7.50 (m, 1H), 7.35 (m, 1H), 7.30 (m, 1H), 3.93 (s, 3H). LC/MS: Rt 3.0 mins; m/e 375 (M+H), 373 (M-H).


15

Example 68

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(3-fluoro-pyridin-2-yl)-1H-pyrrolo[2,3-b]pyridine (74)

20 [0283] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.60 (s, 1H), 8.91 (s, 1H), 8.88 (s, 1H), 8.57 (m, 1H), 8.13 (s, 1H), 7.87 (d, 1H), 7.85 (ddd, 1H), 7.50 (m, 2H), 7.37 (m, 1H), 7.29 (m, 1H). LC/MS: Rt 3.7 mins.; m/e 392.9 (M+H), 391 (M-H).


- 161 -

Example 69

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-pyrimidin-2-yl-1H-pyrrolo[2,3-b]pyridine (75)

5 [0284] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.63 (s, 1H), 9.36 (d, 1H), 8.96 (d, 1H), 8.91 (d, 2H), 8.89 (s, 1H), 7.84 (d, 1H), 7.53 (q, 1H), 7.46 (t, 1H), 7.39 (t, 1H), 7.30 (q, 1H). LC/MS: Rt 3.5 mins.; m/e 375.9 (M+H), 374 (M-H).


10

Example 70

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(3H-imidazo[1,4-b]pyridin-2-yl)-1H-pyrrolo[2,3-b]pyridine (76)


15 [0285] ^1H NMR (500 MHz, DMSO- d_6) δ : 14.65 (br, 1H), 12.64 (s, 1H), 9.07 (s, 1H), 8.90 (s, 1H), 8.82 (d, 1H), 8.57 (d, 1H), 8.13 (s, 1H), 7.76 (d, 1H), 7.50 (m, 1H), 7.39 (m, 1H), 7.30 (m, 1H). LC/MS: Rt 2.1 mins; m/e 364 (M+H), 362.1 (M-H).

- 162 -

Example 71

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H,1'H-[5,5']
bi[pyrrolo[2,3-b]pyridinyl] (77)

5 [0286] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.52 (s, 1H),
11.72 (s, 1H), 8.88 (s, 1H), 8.65 (d, 1H), 8.34 (d,
1H), 8.03 (d, 1H), 7.92 (s, 1H), 7.83 (d, 1H), 7.53 (m,
2H), 7.34 (m, 2H), 6.53 (dd, 1H)


Example 72

10

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(4-methoxy-
pyridin-2-yl)-1H-pyrrolo[2,3-b]pyridine (78)


15 [0287] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.52 (s, 1H),
9.05 (d, 1H), 8.88 (s, 1H), 8.53 (d, 1H), 8.48 (d, 1H),
7.83 (s, 1H), 7.50 (q, 1H), 7.43 (d, 1H), 7.38 (t, 1H),
7.29 (m, 1H), 6.95 (dd, 1H), 3.95 (s, 3H). LC/MS: Rt
2.5 mins.; m/e 405 (M+H), 403.1 (M-H).

- 163 -

Example 73

6-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-N,N-dimethyl-nicotinamide (79)

5 [0288] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.58 (s, 1H),
9.08 (d, J = 2.0 Hz, 1H), 8.89 (s, 1H), 8.74 (d, J =
4.5 Hz, 1H), 8.48 (d, J = 2.0 Hz, 1H), 7.88 (d, J = 2.5
Hz, 1H), 7.81 (s, 1H), 7.48 (m, 1H), 7.38 (m, 1H), 7.33
10 (dd, J = 1.0, 5.0 Hz, 1H), 7.28 (m, 1H), 3.05 (s, 3H),
2.93 (s, 3H). LC/MS: Rt 3.1 mins; m/e 446 (M+H),
444.1 (M-H).

Example 74

15 (2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-pyridin-4-yl)-dimethyl-amine
(80)


[0289] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.71 (s, 1H),
8.91 (s, 2H), 8.53 (s, 1H), 8.24 (d, J = 7.0 Hz, 1H),
7.87 (s, 1H), 7.50 (m, 1H), 7.31 (m, 1H), 7.30 (m, 1H),

- 164 -

7.18 (s, 1H), 6.91 (d, J = 5.5 Hz, 1H), 3.24 (s, 6H).

LC/MS: Rt 2.4 mins; m/e 418 (M+H), 416 (M-H).

Example 75




5

(2-Chloro-6-{3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-pyridin-4-ylmethyl)-dimethylamine (81)

[0290] ¹H NMR (500 MHz, CD3OD) δ: 9.05 (d, 1H),
 10 8.79 (d, 1H), 8.62 (s, 1H), 8.04 (s, 1H), 7.70 (s, 1H),
 7.54 (s, 1H), 7.36-7.24 (m, 3H), 4.42 (s, 2H), 2.97 (s,
 6H). LC/MS: Rt 2.6 mins.; m/e 466 (M+H), 464.1
 (M-H).

Example 76




15

(2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-pyridin-4-ylmethyl)-dimethylamine (82)

20 [0291] ¹H NMR (500 MHz, DMSO-d6) δ: 12.57 (s, 1H),
 9.81 (br, s, 1H), 9.08 (s, 1H), 8.91 (s, 1H), 8.76 (br,
 1H), 8.63 (br, 1H), 8.02 (br, 1H), 7.85 (s, 1H),
 7.55-7.28 (complex, 4H), 4.15 (br, 2H), 3.28 (s, 6H),

- 165 -


2.31 (s, 2.7H). LC/MS: Rt 2.4 mins.; m/e 432.3 (M+H), 430.3 (M-H).

Example 77

5

2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-pyridin-4-ylamine (83)

[0292] ¹H NMR (500 MHz, DMSO-d₆) δ: 13.45 (s, 1H),
12.78 (s, 1H), 8.92 (s, 1H), 8.74 (d, 1H), 8.52 (d,
10 1H), 8.17 (d, br, 1H), 8.13 (br, 1H), 8.03 (br, 1H),
7.86 (d, 1H), 7.53 (q, 1H), 7.41 (t, 1H), 7.33 (q,
1H), 7.11 (d, 1H), 6.85 (dd, 1H), 2.32 (s, 3.5H).
LC/MS: Rt 2.4 mins.; m/e 390.2 (M+H), 388.3 (M-H).

Example 78

15


(6-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
o[2,3-b]pyridin-5-yl}-pyridin-3-ylmethyl)-dimethylamin
e (84)

20 [0293] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.60 (s, 1H),
9.72 (br, 1H), 9.10 (s, 1H), 8.90 (s, 1H), 8.77 (s,
1H), 8.64 (s, 1H), 8.09 (d, J = 8.2 Hz, 1H), 8.03 (dd,
J = 8.2, 1.8 Hz, 1H), 7.84 (d, J = 2.5 Hz, 1H), 7.51
(m, 1H), 7.38-7.30 (m, 2H), 4.39 (d, 2H), 2.81 (d, 6H),

- 166 -


2.34 (s, 3H). LC/MS: Rt 2.2 mins.; m/e 432.2 (M+H), 430.3 (M-H).

General method D:



5 **[0294]** Reagents and Conditions: (a) R-B(OR)₂, 2M Na₂CO₃, PdCl₂(dppf), DMF.

Example 79




10 **3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-pyrimidin-5-yl-1H-pyrrolo[2,3-b]pyridine (85)**

[0295] To a solution of 5-bromo-3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (prepared by Method A, 50 mg, 0.13 mmol) and 5-pyrimidine boronic acid (33 mg, 0.27 mmol) in DME (1 mL) was added sat. NaHCO₃ (1.2 M, 0.54 mL, 0.65 mmol) and tri-*tert*-butylphosphine (27 mg, 0.13 mmol). The suspension was stirred under N₂ atmosphere while catalyst PdCl₂(dppf) (5 mg, 0.006 mmol) was added. The reaction mixture was then heated at 80°C for 5h and diluted with ethyl acetate. The inorganic salt was removed by filtration, the filtrate was evaporated and purified by HPLC to afford the desired product as off-white solid (5.0 mg, 0.013 mmol, 10%).

- 167 -


[0296] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.62 (br, 1H), 9.19 (s, 1H), 9.04 (s, 2H), 8.88 (s, 1H), 8.72 (d, 1H), 8.06 (d, 1H), 7.92 (s, 1H), 7.50 (m, 1H), 7.36 (m, 1H), 7.30 (m, 1H) ppm. LC/MS: Rt 3.1 mins.; m/e 375.9 (M+H), 374.1 (M-H).

Example 80

3-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-phenylamine (86)

10 [0297] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.51 (s, 1H), 8.88 (s, 1H), 8.59 (d, 1H), 8.02 (d, 1H), 7.83 (s, 1H), 7.55 (m, 1H), 7.36 (m, 3H), 7.21 (s, 1H), 7.15 (d, br, 1H), 6.98 (d, br, 1H). LC/MS: Rt 2.8 mins.; m/e 389 (M+H), 387.1 (M-H).

15


Example 81

3-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-phenol (87)

20 [0298] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.49 (s, 1H), 9.52 (s, 1H), 8.87 (s, 1H), 8.56 (d, 1H), 7.88 (d, 1H), 7.86 (d, 1H), 7.54 (m, 1H), 7.38 (m, 2H), 7.26 (t, 1H),

- 168 -


6.95 (m, 2H), 6.78 (dd, 1H). LC/MS: Rt 3.6 mins.;
m/e 389.9 (M+H), 388 (M-H).

Example 82

5

(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
o[2,3-b]pyridin-5-yl}-phenyl)-dimethyl-amine (88)


[0299] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.43 (s, 1H),
8.87 (s, 1H), 8.56 (d, 1H), 7.85 (d, 1H), 7.76 (d, 1H),
10 7.55 (m, 1H), 7.40-7.30 (m, 4H) 6.92 (d, 2H), 2.98 (s,
6H). LC/MS: Rt 3.4 MINS.; M/E 417 (M+H), 415 (M-H).

Example 83

15 4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-benzonitrile (89)

[0300] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.60 (s, 1H),
8.89 (s, 1H), 8.71 (d, 1H), 7.97 (d, 1H), 7.93 (m, 3H),
7.77 (d, 2H), 7.54 (m, 1H), 7.35 (m, 2H).
20 LC/MS: Rt 4.1 mins.; m/e 398.9 (M+H), 397 (M-H).

- 169 -


Example 84

4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
5 [2,3-b]pyridin-5-yl}-phenylamine (90)

[0301] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.48 (s, 1H),
8.88 (s, 1H), 8.58 (d, 1H), 7.86 (d overlap, 2H), 7.54
(m, 1H), 7.48 (d, 2H), 7.35 (m, 2H), 7.10 (d, 2H).

LC/MS: Rt 2.7 mins.; m/e 389 (M+H), 387 (M-H).


10

Example 85

(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
o[2,3-b]pyridin-5-yl}-benzyl)-dimethyl-amine (91)

15 [0302] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.57 (s, 1H),
9.82 (br, 1H), 8.89 (s, 1H), 8.68 (d, 1H), 7.98 (d,
1H), 7.90 (d, 1H), 7.68 (d, 2H), 7.60 (d, 2H), 7.54 (m,
1H), 7.35 (m, 2H), 4.35 (s, 2H), 2.79 (s, 6H). LC/MS:
Rt 2.4 mins.; m/e 431 (M+H), 429 (M-H).


- 170 -

Example 86

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(4-morpholi
 5 n-4-ylmethyl-phenyl)-1H-pyrrolo[2,3-b]pyridine (92)

[0303] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.58 (s, 1H),
 9.95 (br, 1H), 8.89 (s, 1H), 8.68 (d, 1H), 7.99 (d,
 1H), 7.89 (d, 1H), 7.68 (d, 2H), 7.61 (d, 2H), 7.53 (m,
 1H), 7.35 (m, 2H), 4.42 (s, 2H), 3.99 (d, br, 2H), 3.65
 10 (t, br, 2H), 3.32 (d, br, 2H), 3.17 (t, br, 2H).

LC/MS: Rt 2.4 mins.; m/e 473 (M+H), 471 (M-H).


Example 87

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(4-pyrrolid
 15 in-1-ylmethyl-phenyl)-1H-pyrrolo[2,3-b]pyridine (93)

[0304] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.51 (s, 1H),
 8.87 (s, 1H), 8.61 (d, 1H), 7.90 (s, 1H), 7.82 (d, 1H),
 7.57 (m, 1H), 7.45 (d, 2H), 7.39 (d, 2H), 7.35 (m, 2H),
 3.61 (s, 2H), 2.45 (m, br, 4H), 1.71 (m, br, 4H).

20 LC/MS: Rt 2.5 mins.; 457 (M+H), 455 (M-H).


- 171 -

Example 88

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-[4-(4-methyl-
1-piperazin-1-ylmethyl)-phenyl]-1H-pyrrolo[2,3-b]pyridi
5 ne (94)

[0305] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.55 (s, 1H),
8.89 (s, 1H), 8.65 (d, 1H), 7.94 (d, 1H), 7.89 (d, 1H),
7.59 (d, 2H), 7.55 (m, 1H), 7.50 (d, 2H), 7.36 (m, 2H),
3.98 (s, 2H), 3.58-2.99 (complex, 8H), 2.84 (s, 3H).


10 LC/MS: Rt 2.2 mins.; m/e 486 (M+H), 484 (M-H).

Example 89

N-(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrr
olo[2,3-b]pyridin-5-yl}-phenyl)-acetamide (95)


15 [0306] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.49 (s, 1H),
10.03 (s, 1H), 8.88 (s, 1H), 8.60 (s, 1H), 7.88 (d,
1H), 7.83 (d, 1H), 7.67 (d, 2H), 7.54 (m, 1H), 7.46 (d,
2H), 7.35 (m, 2H), 2.07 (s, 3H). LC/MS: Rt 3.4
mins.; m/e 431 (M+H), 429 (M-H).

- 172 -

Example 90

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(5-methoxy-
 5 pyridin-3-yl)-1H-pyrrolo[2,3-b]pyridine (96)


[0307] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.60 (s, 1H),
 8.89 (s, 1H), 8.69 (d, 1H), 8.34 (d, 1H), 8.31 (d, 1H),
 7.95 (d, 1H), 7.93 (d, 1H), 7.57 (s, 1H), 7.50 (m, 1H),
 7.35 (m, 1H), 7.30 (m, 1H), 3.93 (s, 3H). LC/MS: Rt
 10 2.9 mins.; m/e 405 (M+H), 403 (M-H).

Example 91

3'-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1-methanesul-
 fonyl-1H,1'H-[5,5']bi[pyrrolo[2,3-b]pyridinyl] (97)


15 [0308] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.58 (s, 1H),
 8.89 (s, 1H), 8.71 (d, 1H), 8.66 (d, 1H), 8.24 (d, 1H),
 8.02 (d, 1H), 7.90 (d, 1H), 7.80 (d, 1H), 7.57 (m, 1H),
 7.37 (m, 1H), 7.33 (m, 1H), 6.88 (d, 1H), 3.77 (s, 3H).
 LC/MS: Rt 3.9 mins.; m/e 491.9 (M+H), 490 (M-H).

- 173 -

Example 92


N-(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-phenyl)-methane sulfonamide
(98)

[0309] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.52 (s, 1H), 9.82 (s, 1H), 8.88 (s, 1H), 8.60 (d, 1H), 7.86 (m, 2H), 7.53 (m, 3H), 7.32 (m, 4H), 3.05 (s, 3H). LC/MS: Rt 3.7 mins.; m/e 466.90 (M+H), 465.0 (M-H).


Example 93

3-[4-(2,3-Dichloro-phenyl)-isoxazol-5-yl]-5-pyridin-3-yl-1H-pyrrolo[2,3-b]pyridine (99)

[0310] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.57 (s, 1H), 8.87 (s, 1H), 8.84 (s, 1H), 8.69 (d, 1H), 8.66 (d, 1H), 8.11 (d, 1H), 7.93 (s, 1H), 7.76 (dd, 1H), 7.72 (d, 1H), 7.63 (dd, 1H), 7.51 (dd, 1H), 7.46 (dd, 1H). LC/MS: Rt 2.9 mins.; m/e 406.8 (M+H), 405 (M-H).

General method E:**Preparation of**

3 - [4 - (2,3-Difluoro-phenyl) -isoxazol-5-yl] -6-pyrrolidin-
 5 1-yl-1*H*-pyrrolo[2,3-*b*]pyridine, Example 97 (100)

Example 94**Step A: 1*H*-Pyrrolo[2,3-*b*]pyridine 7-oxide**

[0311] To a solution of 7-azaindole (2.0 g, 16.93
 10 mmol) in dry DME (60 mL) was added m-CPBA (70%) (6.3 g,
 25.55 mmol). The resulting yellow solution was stirred
 at RT for 2 h during which time the product was
 precipitated out. The mixture was cooled and the light
 yellow product was isolated by filtration and washed
 15 with ether. A suspension of this yellow solid in water
 (60 mL) was basified to pH 9 with sat. K₂CO₃ solution.
 The solution was then cooled in refrigerator for a
 weekend. A white precipitate was collected by
 filtration and the filtrate was half evaporated and

- 175 -

cooled again to repeat the crystallization procedure. The precipitates were combined and dried on the pump for the next reaction (1.5 g, 11.2 mmol, 66%). MS (ES+): m/e= 135.1 (M+H); LC: 1.37 min.

5

Example 95

Step B: 6-Chloro-pyrrolo[2,3-*b*]pyridine-1-carboxylic acid ethyl ester


[0312] To a solution of 1H-pyrrolo[2,3-*b*]pyridine 7-oxide (500 mg, 3.73 mmol) and HMDS (600 mg, 3.72 mmol) in dry THF was added ethylchloroformate (1.0 g, 9.21 mmol) dropwise at RT. The solution was stirred at RT for 1 h and evaporated. The residue was taken up to ethyl acetate and washed with sat. NaHCO₃ solution. After evaporation, the crude product was purified by flash column to afford a colorless oil (600 mg, 72%). MS (ES+): m/e= 225.1 (M+H); LC: 3.29 min.

Example 96

Step C: 6-Chloro-1H-pyrrolo[2,3-*b*]pyridine


[0313] To a solution of 6-Chloro-pyrrolo[2,3-*b*]pyridine-1-carboxylic acid ethyl ester (400 mg, 1.78 mmol) in MeOH (35 mL) was added 1N NaOH (13 mL). The solution was stirred at RT for 6 h and evaporated the solvent. The residue was neutralized with sat. NaHCO₃ and the resulting precipitate was collected by filtration. After washing with water, the solid was dried on the pump for direct use (260 mg, 1.71 mmol, 96%). MS (ES+): m/e= 153.1 (M+H); LC: 2.85 min.

- 176 -

Example 97**Step D:**

1- (6-Chloro-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl) -2- (2,3-difluoro-phenyl) -ethanone

5 [0314] 6-Chloro-1*H*-pyrrolo[2,3-*b*]pyridine (250 mg, 1.64 mmol) was converted to 1- (6-chloro-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl) -2- (2,3-difluoro-phenyl) -ethanone using Friedel-Crafts reaction as
10 described in Method A.


Example 98**Step E:**

2- (2,3-Difluoro-phenyl) -1- (6-pyrrolidin-1-yl-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl) -ethanone

15 [0315] A solution of 1- (6-chloro-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl) -2- (2,3-difluoro-phenyl) -ethanone (100 mg, 0.29 mmol) and pyrrolidine (1 mL) in NMP (2 mL) was heated in a sealed
20 tube with microwave at 220°C for 15 min. The solution was poured into water and 0.5N HCl was added to precipitate the product. The crude product was collected by filtration, washed with water, and dried

- 177 -

on the pump for direct use (80 mg, 0.23 mmol, 79%). MS (ES+): m/e = 342.2 (M+H); LC: 2.92 min.

Example 995 **Step F:**


3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-6-pyrrolidin-1-yl-1H-pyrrolo[2,3-b]pyridine (100)

[0316]

2-(2,3-Difluoro-phenyl)-1-(6-pyrrolidin-1-yl-1H-pyrrolo
10 [2,3-b]pyridin-3-yl)-ethanone (80 mg, 0.23 mmol) was converted to

3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-6-pyrrolidin-1-yl-1H-pyrrolo[2,3-b]pyridine (22 mg, 0.06 mmol, 26%) by using the isoxazole formation procedures described
15 in Method A. MS ¹H NMR (500 MHz, DMSO-d₆) δ: 11.75 (s, 1H), 8.78 (s, 1H), 7.65 (d, 1H), 7.50 (dd, 1H), 7.30 (m, 2H), 7.20 (s, 1H), 6.39 (d, 1H), 3.45 (brs, 4H), 1.95 (brs, 4H). LC/MS: Rt 3.14 mins.; m/e 367.2 (M+H), 365.4 (M-H).

20


Example 100

- 178 -

6-Chloro-3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine (101)

[0317] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.60 (s, 1H), 8.89 (s, 1H), 7.91 (d, 1H), 7.82 (s, 1H), 7.51 (q, 1H), 7.30 (m, 2H), 7.25 (d, 1H). LC/MS: Rt 4.09 mins.; m/e 332.1 (M+H), 330.1 (M-H).


Example 101



{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-6-yl}-methyl-amine (102)


[0318] ^1H NMR (500 MHz, DMSO- d_6) δ : 11.69 (s, 1H), 8.78 (s, 1H), 7.56 (d, 1H), 7.50 (m, 1H), 7.31 (m, 1H), 7.15 (d, 1H), 6.36 (d, 1H), 2.81 (s, 3H). LC/MS: Rt 2.7 mins; m/e 327.2 (M+H), 325.2 (M-H).

Example 102




6-Chloro-3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-5-pyridin-4-yl-1H-pyrrolo[2,3-b]pyridine (103)

[0319] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.84 (s, 1H), 8.88 (s, 1H), 8.77 (d, 2H), 7.99 (d, 1H), 7.79 (s, 1H), 7.61 (d, 2H), 7.51 (m, 1H), 7.32 (m, 2H). LC/MS: Rt 2.68 mins.; m/e 408.9 (M+H), 407 (M-H).

General method F:

[0320] Reagents and Conditions: (a) pyrrolidine,
 5 CO, PdCl₂(dppf), DMF, 80 °C; (b) (i)
 substituted-phenylacetyl chloride, AlCl₃, CH₂Cl₂, (ii)
 Brederick's Reagent, THF, reflux, (iii) H₂NOH HCl,
 NaOAc, THF, reflux.

Example 103

10


Pyrrolidin-1-yl-(1H-pyrrolo[2,3-b]pyridin-5-yl)-methanone

[0321] A mixture of
 5-bromo-1-methanesulfonyl-1H-pyrrolo[2,3-b]pyridine
 15 (300 mg, 1.1 mmol), PdCl₂(dppf) (55 mg, 0.07 mmol), and
 pyrrolidine (2 mL) in DMF (5 mL) was charged with CO
 balloon. The system was degassed with vacuum twice
 before it was heated to 80°C for 6 h. The solution was
 cooled and poured into water. The aqueous solution was
 20 extracted with ethyl acetate (3 x 50 mL), the combined
 organic layers were dried over Na₂SO₄, and the solvent
 was removed by vacuum evaporation. The crude product
 was treated with 6N NaOH in MeOH for 2 h. MeOH was
 removed by evaporation. The aqueous solution was
 25 neutralized with 6N HCl to pH 8 and extracted with
 ethyl acetate. The combined organic layers were dried

- 180 -

over Na₂SO₄ and the solvent was removed by vacuum evaporation to give a yellow solid (80 mg, 0.37 mmol, 34%), which was used directly for the next step. MS (ES⁺): m/e = 216.1 (M+H); LC: 2.18 min.

5


Example 104

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-pyrrolidin-1-yl-methanone (104)
[0322]

10 Pyrrolidin-1-yl-(1H-pyrrolo[2,3-b]pyridin-5-yl)-methanone (80 mg, 0.37 mmol) was converted to {3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-pyrrolidin-1-yl-methanone (20.0 mg, 0.05 mmol, 14%) by using Method A.

15 [0323] ¹H NMR (500 MHz, DMSO-d₆/CD₃OD) δ: 8.84 (s, 1H), 8.50 (d, 1H), 7.99 (d, 1H), 7.86 (s, 1H), 7.47 (m, 1H), 7.34 (m, 1H), 7.28 (m, 1H), 3.50 (br, 2H), 3.32 (br, 2H), 1.85 (br, 4H). LC/MS: Rt 3.2 mins.; m/e 395 (M+H), 393 (M-H).

20

Example 105


3 - [4 - (2,3-Difluoro-phenyl) -isoxazol-5-yl] -1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid methyl ester (105)

[0324] was prepared by a similar carbonylation step but was carried out in methanol to give the title compound.

[0325] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.84 (s, 1H), 8.91 (s, 1H), 8.89 (d, 1H), 8.34 (d, 1H), 7.96 (d, 1H), 7.54 (m, 1H), 7.34 (m, 2H), 3.87 (s, 3H). LC/MS: R_t 3.6 mins.; m/e 356 (M+H), 354 (M-H).


10 Preparation of triazolyl-azaindoles

General method G:




[0326] Reagents and Conditions: (a) (i) trichloroacetyl chloride, AlCl_3 , CH_2Cl_2 , (ii) Et_3N , H_2O , RT (b) (i) oxalyl chloride, DMF (cat.), CH_2Cl_2 , (ii) Ar-NH_2 , Et_3N , CH_2Cl_2 ; (c) Lawesson's reagent, toluene, reflux; (d) hydrazine, EtOH & CH_2Cl_2 ; (e) triethylorthoformate, HCO_2H ; optional step (f) Suzuki coupling; when $\text{R} = \text{Br}$ or I : R-B(OR)_2 , 2M Na_2CO_3 , $\text{PdCl}_2(\text{dppf})$, DMF; when $\text{R} = \text{B(OH)}_2$: Ar-X (where $\text{X} = \text{Br}$, I , OTf), 2M Na_2CO_3 , $\text{PdCl}_2(\text{dppf})$, DMF

- 182 -


Example 106

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridine-5-boronic acid (158)

5 [0327] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.32 (s, 1H), 8.85 (s, 1H), 8.69 (d, 1H), 8.52 (s, 1H), 8.13 (s, 2H), 7.64 (d, 1H), 7.50 (m, 1H), 7.34 (m, 1H), 7.28 (m, 1H). LC/MS: Rt 2.8 mins.; m/z 341.9 (M+H), 340.1 (M-H).



10


Example 107

5-Bromo-1H-pyrrolo[2,3-b]pyridine-3-carboxylic acid

15 [0328] To a solution of 5-bromo-7-azaindole (2.0 g, 10.1 mmol) in DCM (50 mL) was added AlCl_3 (6.8 g, 51.0 mmol). The suspension was stirred at RT for 10 min and trichloroacetyl chloride (2.8 g, 15.40 mmol) was added slowly. The mixture was stirred at RT for overnight


- 183 -

and then poured into iced-water. The aqueous solution was extracted with DCM three times, and organic layers were combined and evaporated. The crude solid was dissolved in THF (50 mL) and treated with water (25 mL) and triethylamine (5 mL) at RT for 6h. The solvents were then removed by evaporation and the resulting solid was poured into 1N HCl solution. The crude product was collected by filtration and washed with water. After drying on the pump for over night, a white solid was obtained (2.4 g, 9.96 mmol). MS (ES+): m/e= 241.0 (M+H); LC: 2.7 min.

Example 108


5-Bromo-1H-pyrrolo[2,3-b]pyridine-3-carboxylic acid (2,3-difluorophenyl)-amide
[0329] To a suspension of 5-bromo-1H-pyrrolo[2,3-b]pyridine-3-carboxylic acid (950 mg, 3.94 mmol) in DCM (20 mL) and DMF (0.1 mL) was added oxalyl chloride (600 mg, 4.72 mmol) slowly. The mixture was stirred at RT for 1 h. To this suspension was then added a solution of 2,3-difluorophenyl amine (610 mg, 4.72 mmol) and triethylamine (800 mg, 7.91 mmol) in DCM (5 mL). The reaction was kept at RT for another 2 h. The solvent was then evaporated, the residue was washed water, and dried for direct use. MS (ES+): m/e= 352 (M+H); LC: 3.5 min.

- 184 -

Example 109

**5-Bromo-1H-pyrrolo[2,3-b]pyridine-3-carbothioic acid
(2,3-difluoro-phenyl)-amide**

5 [0330] To a solution of
5-bromo-1H-pyrrolo[2,3-b]pyridine-3-carboxylic acid
(2,3-difluoro-phenyl)-amide (300 mg, 0.85 mmol) in
toluene (6 mL) was added Lawesson's reagent (210 mg,
0.52 mmol). The suspension was heated under reflux for
10 14 h. The solvent was removed by evaporation and the
residue was dried on the pump for the next reaction. MS
(ES⁺): m/e= 368 (M+H); LC: 3.6 min.

Example 110

15 **5-Bromo-3-[4-(2,3-difluoro-phenyl)-4H-[1,2,4]triazol-3-
yl]-1H-pyrrolo[2,3-b]pyridine**

[0331] A crude material from above was dissolved in
a co-solvents of ethanol (5 mL) and DCM (5 mL).
Hydrazine (2 mL) was added at RT. The solution was
20 stirred at RT for 4 h and evaporated. The residue was
poured into aqueous NaHCO₃ solution, filtered, washed
with water, and dried. The crude product was dissolved
in triethyl orthoformate (5 mL). To this solution was
added HCOOH (1 mL) at 0°C. The reaction was allowed to


- 185 -

warm up to RT and stayed for overnight. The solvents were removed by evaporation, the residue was taken up to ethyl acetate (50 mL) and washed with aq. NaHCO₃. After drying over Na₂SO₄, the solvent was evaporated to afford the desired triazole as yellow solid (120 mg, 0.32 mmol).

[0332] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.32 (s, 1H), 8.86 (s, 1H), 8.62 (d, 1H), 8.42 (d, 1H), 7.75 (q, 1H), 7.57 (t, 1H), 7.46 (m, 1H), 7.07 (d, 1H)

10 LC/MS: Rt 2.8 mins; m/e 377.1 (M+H).

Example 111




15 (4-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-phenyl)-dimethyl-amine (106)

[0333] The crude product (50 mg, 0.13 mmol) obtained above was converted to


20 (4-{3-[4-(2,3-difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-phenyl)-dimethyl-amine (19 mg, 0.05 mmol) by using Suzuki procedures as described in Method D.

[0334] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.15 (s, 1H), 8.91 (s, 1H), 8.60 (s, 1H), 8.54 (s, 1H), 7.76 (m, 1H), 7.62 (m, 3H), 7.48 (m, 1H), 7.08 (m, br, 3H), 3.04 (s, 6H), 2.33 (s, 3H). LC/MS: Rt 2.2 mins.; m/e 417.3 (M+H), 415.3 (M-H).

Example 112

5 3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridine (107)


[0335] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.12 (s, 1H),
 8.90 (s, 1H), 8.45 (dd, 1H), 8.35 (dd, 1H), 7.75 (m,
 1H), 7.58 (t, 1H), 7.47 (m, 1H), 7.25 (dd, 1H), 7.04
 10 (s, 1H). LC/MS: Rt 2.06 mins.; m/e 298.2 (M+H),
 296.2 (M-H).

Example 113

15 3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-methoxy-1H-pyrrolo[2,3-b]pyridine (108)


[0336] ¹H NMR (500 MHz, DMSO-d₆) δ: 11.97 (s, 1H),
 8.85 (s, 1H), 8.07 (d, 1H), 7.92 (d, 1H), 7.74 (m, 1H),
 7.58 (m, 1H), 7.47 (m, 1H), 6.96 (d, 1H), 3.87 (s, 3H).
 20 LC/MS: Rt 2.35 mins.; m/e 328 (M+H), 326 (M-H).

- 187 -

Example 114

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-(4-morpholin-4-ylmethyl-phenyl)-1H-pyrrolo[2,3-b]pyridine
 5 (109)

[0337] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.20 (s, 1H), 9.82 (brs, 1H), 8.88 (s, 1H), 8.71 (s, 1H), 8.69 (s, 1H), 7.86 (d, 2H), 7.77 (m, 1H), 7.66 (d, 2H), 7.59 (m, 1H), 7.49 (m, 1H), 7.06 (d, 1H), 4.45 (d, 2H), 3.97 (d, 10 2H), 3.65 (t, 2H), 3.35 (2H, covered by water), 3.17 (br, 2H), 2.28 (s, 3H). LC/MS: Rt 1.80 mins.; m/e 473.3 (M+H), 471.4 (M-H).

Example 115

15


3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-[4-(4-methyl-piperazin-1-yl)-phenyl]-1H-pyrrolo[2,3-b]pyr
 idine (110)

[0338] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.11 (s, 1H), 9.64 (br, 1H), 8.88 (s, 1H), 8.60 (d, 1H), 8.56 (d, 1H), 7.76 (q, 1H), 7.62 (d, 2H), 7.59 (m, 1H), 7.48 (m, 1H), 7.16 (d, 2H), 7.04 (d, 1H), 3.94 (d, 2H), 3.55 (d, 20 2H), 3.20 (q, 2H), 3.04 (t, 2H), 2.89 (d, 3H), 2.33 (s,

- 188 -

3.8H). LC/MS: Rt 1.8 mins.; m/e 472.3 (M+H), 470.4 (M-H).


Example 116



5 3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-(6-piperazin-1-yl-pyridin-3-yl)-1H-pyrrolo[2,3-b]pyridine (111)

[0339] ¹H NMR (500 MHz, DMSO-d₆) δ: 8.81 (s, 1H), 8.52 (s, 1H), 8.48 (d, 1H), 8.43 (s, 1H), 7.84 (dd, 1H), 7.55 (q, 1H), 7.58 (t, 1H), 7.48 (m, 1H), 7.03 (s, 1H), 6.93 (dd, 1H), 3.45 (m, 8H), 2.35 (s, 8H).
 10 LC/MS: Rt 1.5 mins.; m/e 459.3 (M+H), 457.4 (M-H).

Example 117




15

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-1H-pyrrolo[2,3-b]pyridine (112)

[0340] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.15 (s, 1H), 9.62 (br, 1H), 8.85 (s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 8.50 (s, 1H), 7.95 (d, 1H), 7.75 (m, 1H), 7.60 (m, 1H), 3.25 (s, 3H).
 20

- 189 -


1H), 7.48 (m, 1H), 7.06 (s, 1H), 7.04 (s, 1H),
 3.35-3.00 (mbr, 8H), 2.68 (br, 3H), 2.30 (s, 2.1H).
 LC/MS: Rt 1.6 mins.; m/e 473.3 (M+H), 471.4 (M-H).

Example 118

5

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-py
 ridin-3-yl-1H-pyrrolo[2,3-b]pyridine (113)


[0341] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.35 (s, 1H),
 9.21 (d, 1H), 8.91 (s, 1H), 8.80 (m, 3H), 8.68 (d, 1H),
 10 7.95 (dd, 1H), 7.76 (q, 1H), 7.60 (t, 1H), 7.49 (m,
 1H), 7.14 (d, 1H), 2.32 (s, 4H). LC/MS: Rt 1.6
 mins.; m/e 375.2 (M+H), 373.2 (M-H).

Example 119

15 3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-py
 ridin-4-yl-1H-pyrrolo[2,3-b]pyridine (114)

[0342] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.49 (s, 1H),
 8.98 (s, 2H), 8.92 (m, 3H), 8.41 (d, 2H), 7.75 (q,
 1H), 7.60 (t, 1H), 7.49 (m, 1H), 7.19 (d, 1H),
 20 (s, 3.5H). LC/MS: Rt 1.5 mins.; m/e 375.2 (M+H),
 373.2 (M-H).

- 190 -


Example 120

[0343]

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-(6-
 5 -fluoro-pyridin-3-yl)-1H-pyrrolo[2,3-b]-pyridine

[0344] obtained as a yellow solid (yield 75%).

MS: m/e 393.3 (M+1); LC: Rt 2.7 min


Example 121

10 [0345]

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-[6-
 -(2-pyrrolidin-1-ylmethyl-pyrrolidin-1-yl)-pyridin-3-yl]
]-1H-pyrrolo[2,3-b]pyridine (115)

15 [0346] ¹H NMR (500 MHz, DMSO-d₆) δ: 8.81 (s, 1H),
 8.52 (s, 1H), 8.48 (s, 1H), 8.38 (s, 1H), 7.78 (m, 2H),
 7.57 (m, 1H), 7.47 (m, 1H), 7.01 (s, 1H), 6.57 (d, 1H),
 4.18 (m, 1H), 3.54 (m, 2H), 3.17 (br, 2H), 2.65-2.55
 (covered by DMSO, 4H), 2.10-1.90 (complex, 4H), 1.70
 (m, 4H). LC/MS: Rt 2.00 mins.; m/e 527.30 (M+H).


- 191 -

Example 122

[0347]

5' - { 3 - [4 - (2,3-Difluoro-phenyl) - 4H - [1,2,4] triazol-3-yl] -
 5 1H-pyrrolo[2,3-b]pyridin-5-yl } - 3,4,5,6-tetrahydro-2H - [1
 , 2'] bi-pyridinyl-4-ol (116)

[0348] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.10 (br, 1H),
 8.81 (s, 1H), 8.52 (s, 1H), 8.47 (d, 1H), 8.42 (d, 1H),
 7.82 (dd, 1H), 7.74 (m, 1H), 7.58 (t, 1H), 7.47 (m,
 10 1H), 7.03 (s, 1H), 6.96 (d, 1H), 4.68 (br, 1H), 4.06
 (d, br, 2H), 3.73 (m, br, 2H), 3.14 (t, 2H), 1.81 (d,
 br, 2H), 1.39 (dt, br, 2H). LC/MS: Rt 1.80 mins.;
 m/e 474.30 (M+H).

Example 123

15


[0349]

3 - [4 - (2,3-Difluoro-phenyl) - 4H - [1,2,4] triazol-3-yl] - 5 - [6
 - (4-methyl - [1,4] diazepan-1-yl) - pyridin-3-yl] - 1H-pyrrolo
 20 [2,3-b] -pyridine (117)

[0350] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.09 (s, 1H),
 8.86 (s, 1H), 8.56 (d, 1H), 8.49 (d, 1H), 8.40 (d, 1H),
 7.81 (dd, 1H), 7.75 (m, 1H), 7.59 (t, 1H), 7.46 (m,
 1H), 7.04 (s, 1H), 6.75 (d, 1H), 3.79 (dd, 2H), 3.65

- 192 -

(t, 2H), 2.63 (dd, 2H), 2.48 (covered by DMSO, 2H), 2.27 (s, 3H), 1.92 (m, 2H). LC/MS: Rt 1.40 mins.; m/e 487.40 (M+H).

Example 124


5

[0351]

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-(6-pyrrolidin-1-yl-pyridin-3-yl)-1H-pyrrolo[2,3-b]pyridine (118)

10 [0352] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.09 (s, 1H), 8.86 (s, 1H), 8.55 (d, 1H), 8.48 (s, 1H), 8.39 (d, 1H), 7.81 (dd, 1H), 7.74 (m, 1H), 7.59 (t, 1H), 7.47 (m, 1H), 7.04 (d, 1H), 6.57 (d, 1H), 3.44 (tbr, 4H), 1.97 (tbr, 4H). LC/MS: Rt 2.0 mins.; m/e 444.3 (M+H).

15

Example 125


[0353]

20 N'-(5-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-pyridin-2-yl)-N,N-dimethyl-ethane-1,2-diamine (119)

[0354] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.17 (s, 1H), 9.72 (br, 1H), 8.87 (s, 1H), 8.59 (d, 2H), 8.40 (d, 1H), 8.00 (d, 1H), 7.76 (q, 1H), 7.59 (dd, 1H), 7.48 (m, 1H), 7.03 (d, 1H), 6.87 (d, 1H), 3.73 (t, 2H),

- 193 -


3.33 (t, 2H), 2.88 (s, 6H). LC/MS: Rt 1.3 mins.;
m/e 461.3 (M+H).

Example 126

5 [0355]

(5' - {3 - [4 - (2,3-Difluoro-phenyl) -4H- [1,2,4] triazol-3-yl] -1H-pyrrolo[2,3-b]pyridin-5-yl} -3,4,5,6-tetrahydro-2H- [1,2']bi-pyridinyl-4-yl) -methanol (120)

[0356] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.20 (s, 1H),
10 8.90 (s, 1H), 8.62 (d, 1H), 8.58 (d, 1H), 8.37 (d, 1H),
8.12 (d, 1H), 7.77 (dd, 1H), 7.61 (dd, 1H), 7.49 (m,
1H), 7.32 (br, 1H), 7.09 (s, 1H), 4.34 (m, 3H), 3.31
(d, 1H), 3.08 (m, 2H), 1.80 (d, br, 2H), 1.74 (br, 1H),
1.30 (m, 2H). LC/MS: Rt 1.90 mins.; m/e 488.3
15 (M+H).

Example 127


[0357]

3 - [4 - (2,3-Difluoro-phenyl) -4H- [1,2,4] triazol-3-yl] -5- (6
20 -morpholin-4-yl-pyridin-3-yl) -1H-pyrrolo[2,3-b]pyridine
(121)

[0358] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.20 (s, 1H),
8.90 (s, 1H), 8.62 (d, 1H), 8.57 (d, 1H), 8.43 (d, 1H),
8.13 (dd, 1H), 7.75 (dd, 1H), 7.59 (t, 1H), 7.48 (m,

- 194 -

1H), 7.21 (d, 1H), 7.09 (d, 1H), 3.77 (t, 4H), 3.60 (t, 4H). LC/MS: Rt 2.00 mins.; m/e 460.3 (M+H).


Example 128

5 [0359]

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-[6-(3,5-dimethyl-piperazin-1-yl)-pyridin-3-yl]-1H-pyrrolo[2,3-b]pyridine (122)

[0360] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.20 (s, 1H),
 10 9.12 (br, 1H), 8.90 (s, 1H), 8.61 (d, 1H), 8.56 (d, 1H), 8.50 (d, 1H), 8.01 (dd, 1H), 7.75 (dd, 1H), 7.59 (dd, 1H), 7.48 (m, 1H), 7.17 (d, 1H), 7.08 (d, 1H), 4.54 (d, 2H), 3.38 (br, 2H), 2.85 (dd, 2H), 1.31 (d, 6H). LC/MS: Rt 1.70 mins.; m/e 487.3 (M+H).

15

Example 129


[0361]

5'-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-4-pyrrolidin-1-yl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl (123)

[0362] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.16 (s, 1H),
 20 9.53 (br, 1H), 8.87 (s, 1H), 8.59 (d, 1H), 8.55 (d,

- 195 -

1H), 8.47 (d, 1H), 7.95 (dd, 1H), 7.75 (m, 1H), 7.59
 (m, 1H), 7.48 (m, 1H), 7.10 (d, 1H), 7.05 (d, 1H), 4.48
 (d, 2H), 3.55 (m, br, 2H), 3.43 (m, 1H), 3.11 (m, br,
 2H), 2.92 (t, 2H), 2.14 (d, 2H), 2.05 (m, br, 2H), 1.85
 5 (m, 2H), 1.59 (m, 2H). LC/MS: Rt 1.40 mins.; m/e
 527.3 (M+H).


Example 130

[0363]

10 (5-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-
 1H-pyrrolo[2,3-b]pyridin-5-yl}-pyridin-2-yl)-(tetrahydr
 o-furan-2-ylmethyl)-amine (124)

[0364] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.22 (s, 1H),
 8.88 (s, 1H), 8.61 (m, 2H), 8.26 (s, 1H), 8.18 (d, br,
 15 1H), 7.76 (m, 1H), 7.59 (m, 1H), 7.49 (m, 1H), 7.14 (d,
 br, 1H), 7.06 (d, 1H), 4.06 (m, 1H), 3.80 (m, 1H), 3.70
 (dd, 1H), 3.56 (dd, 1H), 3.42 (dd, 1H), 2.02 (m, 1H),
 1.87 (m, 2H), 1.61 (m, 1H). LC/MS: Rt 1.90 mins.;
 m/e 474.3 (M+H).

20

Example 131

25 [0365]

3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-[6


- 196 -

- (4-ethyl-piperazin-1-yl)-pyridin-3-yl]-1H-pyrrolo[2,3-b]pyridine (125)

[0366] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.17 (s, 1H), 9.62 (br, 1H), 8.88 (s, 1H), 8.61 (d, 1H), 8.56 (d, 1H), 8.52 (d, 1H), 8.00 (dd, 1H), 7.75 (m, 1H), 7.59 (m, 1H), 7.48 (m, 1H), 7.13 (d, 1H), 7.06 (d, 1H), 4.50 (d, br, 2H), 3.61 (d, br, 2H), 3.21 (m, br, 4H), 3.09 (q, 2H), 1.27 (t, 3H). LC/MS: Rt 1.7 mins.; m/z 487.3 (M+H), 485.4 (M-H)

10

Example 132



[0367]


15 3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-5-[6-(4-isopropyl-piperazin-1-yl)-pyridin-3-yl]-1H-pyrrolo[2,3-b]pyridine (126)

[0368] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.16 (s, 1H), 9.45 (br, 1H), 8.87 (s, 1H), 8.61 (d, 1H), 9.57 (d, 1H), 8.52 (d, 1H), 7.99 (dd, 1H), 7.75 (m, 1H), 7.58 (m, 1H), 7.48 (m, 1H), 7.12 (d, 1H), 7.05 (d, 1H), 4.53 (d, br, 2H), 3.57 (d, br, 2H), 3.16 (m, 5H), 1.30 (d, 6H). LC/MS: Rt 1.70 mins.; m/z 501.3 (M+H), 499.4 (M-H).

25

Example 133

- 197 -




[0369]

5'-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-4-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl (127)

[0370] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.21 (s, 1H), 8.88 (s, 1H), 8.62 (d, 1H), 8.58 (d, 1H), 8.33 (d, 1H), 8.15 (d, br, 1H), 7.74 (m, 1H), 7.59 (m, 1H), 7.48 (m, 1H), 7.32 (d, br, 1H), 7.08 (d, 1H), 4.27 (d, br, 2H), 3.09 (t, br, 2H), 1.76 (d, br, 2H), 1.72 (m, 1H), 1.21 (m, 2H), 0.93 (d, 3H). LC/MS: Rt 2.30 mins.; m/z 472.3 (M+H), 470.4 (M-H)

Example 134




[0371]

3-(5-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-pyridin-2-yl)-9-methyl-3,9-diaza-bicyclo[4.2.1]nonane (128)

[0372] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.13 (s, 1H), 9.75 (br, 1H), 8.87 (s, 1H), 8.59 (d, 1H), 8.55 (d, 1H), 8.43 (d, 1H), 7.93 (dd, 1H), 7.75 (m, 1H), 7.59 (m, 1H), 7.48 (m, 1H), 7.03 (d, 1H), 7.01 (d, 1H), 4.47 (d, br, 2H), 4.10 (br, 1H), 3.98 (m, br, 2H), 3.65 (dd, br, 1H), 2.87 (d, 3H), 2.42-1.90 (complex, 8H). LC/MS: Rt 1.60 mins.; 513.3 (M+H), 511.4 (M-H).


- 198 -

Example 135

[0373]

5 5-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-2-(4-methyl-piperazin-1-yl)-phenylamine (129)

[0374] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.12 (s, 1H), 9.88 (br, 1H), 8.87 (s, 1H), 8.64 (d, 1H), 8.56 (d, 1H), 7.95 (s, 1H), 7.77 (dd, 1H), 7.60 (dd, 1H), 7.49 (m, 1H), 7.18 (d, 1H), 7.11 (d, 1H), 7.04 (d, 1H), 7.00 (d, 1H), 3.56 (d, br, 2H), 3.30 (m, 4H), 2.97 (t, 2H), 2.50 (s, 3H). LC/MS: Rt 1.70 mins.; m/e 487.3 (M+H).

Example 136


15

[0375]

4-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4]triazol-3-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-N¹,N¹-dimethyl-benzene-1,2-diamine (130)

20 [0376] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.18 (s, 1H), 8.90 (s, 1H), 8.68 (d, 1H), 8.64 (d, 1H), 7.78 (m, 1H), 7.62 (m, 1H), 7.55 (d, 1H), 7.49 (m, 1H), 7.40 (d, 1H), 7.31 (d, 1H), 7.04 (d, 1H), 3.01 (s, 6H). LC/MS: Rt 1.90 mins.; m/e 432.3 (M+H).



- 199 -

Example 137

[0377]

(4-{3-[4-(2,3-Difluoro-phenyl)-4H-[1,2,4] triazol-3-yl]-
 5 1H-pyrrolo[2,3-b]pyridin-5-yl}-2-nitro-phenyl)-dimethyl
 -amine (131)

[0378] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.17 (s, 1H),
 8.88 (s, 1H), 8.64 (s, 1H), 8.57 (s, 1H), 8.07 (s, 1H),
 7.88 (d, 1H), 7.75 (m, 1H), 7.59 (m, 1H), 7.48 (m, 1H),
 10 7.32 (d, 1H), 7.07 (s, 1H), 2.89 (s, 6H). LC/MS: Rt
 3.30 mins.; m/e 462.3 (M+H).

General method H:Example 138


15

5-(2-[1,2,4]Triazol-1-yl-vinyl)-1H-pyrrolo[2,3-b]pyridi
 ne

[0379] A mixture of 5-bromo-azaindole (1 g, 5.1
 mmol), 1-vinyltriazole (600 mg, 6.3 mmol), and
 20 triethylamine (5 mL) was dissolved in DMF (25 mL). The
 solution was treated with N₂ gas and PdCl₂(dppf) (250
 mg, 0.3 mmol) was added. The reaction was heated with


- 200 -

stirring at 120°C for 16h and evaporated under vacuum. The residue DMF solution was poured into water, filtered, the solid was washed with ether. After drying on the pump for overnight, the crude solid was
5 used for the next reaction directly (800 mg, 3.8 mmol). MS (ES+): m/e= 212.1 (M+H); LC: 2.2 min.

Example 139

10 **5-(2-[1,2,4]Triazol-1-yl-ethyl)-1H-pyrrolo[2,3-b]pyridine**

[0380] A crude material from above (300 mg, 1.42 mmol) was dissolved in MeOH (15 mL). The solution was treated with hydrogen balloon in the presence of Pd/C (10%, 50 mg) for 4h. The catalyst was removed by
15 filtration through celite, the solvent was evaporated, and the residue (200 mg, 0.94 mmol) was dried on the pump for the next use. MS (ES+): m/e= 214.1 (M+H); LC: 0.5 min.

Example 140

20

- 201 -


3 - [4 - (2,3-Difluoro-phenyl) -isoxazol-5-yl] -5 - (2 - [1,2,4] triazol-1-yl-ethyl) -1H-pyrrolo[2,3-b]pyridine (132)

[0381] The crude product obtained above (200 mg, 0.94 mmol) was converted to

5 3 - [4 - (2,3-difluoro-phenyl) -isoxazol-5-yl] -5 - (2 - [1,2,4] triazol-1-yl-ethyl) -1H-pyrrolo[2,3-b]pyridine (108 mg, 0.27 mmol) using Method A.

[0382] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.28 (s, 1H), 8.84 (s, 1H), 8.29 (s, 1H), 8.05 (d, J = 2 Hz, 1H), 7.95 (s, 1H), 7.69 (d, overlap, 2H), 7.52 (m, 1H), 7.31 (m, 2H), 4.40 (t, J = 7.25 Hz, 2H), 3.17 (t, J = 7.25 Hz, 2H). LC/MS: Rt 3.1 mins.; 393 (M+H), 391.1 (M-H).

Example 141




15

3 - [4 - (2,3-Difluoro-phenyl) -isoxazol-5-yl] -5 - (2 - [1,2,4] triazol-1-yl-vinyl) -1H-pyrrolo[2,3-b]pyridine (133)

¹H NMR (500 MHz, DMSO-d₆) δ: 12.48 (s, 1H), 8.88 (s, 1H), 8.84 (s, 1H), 8.57 (d, 1H), 8.18 (s, 1H), 8.15 (d, 1H), 8.02 (d, 1H), 7.75 (d, 1H), 7.54 (m, 1H), 7.36 (d, 1H), 7.35 (m, overlap, 2H). LC/MS: Rt 3.4 mins.; m/e 390.9 (M+H), 389.1 (M-H).


20

General method I:



[0383] Reagents and Conditions: (a) (i) LHMDs,
 5 THF, -78°C, (ii) acetyl chloride (b) hydroxylamine
 hydrochloride, EtOH, reflux.

Example 142



3-[4-(2,3-Difluoro-phenyl)-3-methyl-isoxazol-5-yl]-1H-p
 10 yrrolo[2,3-b]pyridine (134)


[0384] To a solution of
 2-(2,3-difluoro-phenyl)-1-(1H-pyrrolo[2,3-b]pyridin-3-y
 l)-ethanone (200 mg, 0.73 mmol) in dry THF (5 mL) was
 added LiHDMS (1.0 M in THF, 2.2 mL, 2.2 mmol) at -78°C.
 15 The mixture was stirred at this temperature for 1.5 h
 and acetyl chloride (170 mg, 2.2 mmol) was added. The
 reaction was allowed to warm up to RT and continued to
 stir for another 2 h. The solution was then diluted
 with ethyl acetate and washed with 1N HCl. After
 20 evaporation, the crude product was purified by flash
 column to afford the desired product, which was treated
 with NH₂OH HCl (100 mg, 1.44 mmol) in ethanol (10 mL)
 under reflux for 4h to give
 3-[4-(2,3-difluoro-phenyl)-3-methyl-isoxazol-5-yl]-1H-p

- 203 -

pyrrolo[2,3-b]pyridine (45 mg, 0.14 mmol) as white solid.

[0385] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.29 (s, 1H), 8.33 (d, 1H), 7.92 (d, 1H), 7.55 (m, 2H), 7.35 (m, 2H), 7.16 (dd, 1H), 2.20 (s, 3H). LC/MS: Rt 3.3 mins.; m/e 312.1 (M+H), 310.1 (M-H).


Example 143



10 3-[4-(2,3-Difluoro-phenyl)-3-methyl-isoxazol-5-yl]-5-[4-(4-methyl-piperazin-1-yl)-phenyl]-1H-pyrrolo[2,3-b]pyr
idine (135)

[0386] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.40 (s, 1H), 9.58 (br, 1H), 8.57 (s, 1H), 7.80 (s, 1H), 7.69 (s, 1H), 7.59 (br, 1H), 7.45 (d, 2H), 7.38 (s, 1H), 7.12 (d, 2H), 3.94 (d, 2H), 3.55 (d, 2H), 3.17 (m, 2H), 3.02 (m, 2H), 2.89 (s, 3H), 2.31 (s, 4H), 2.21 (s, 3H). LC/MS: Rt 2.4 mins.; 486.3 (M+H), 484.4 (M-H).

General method J:




Example 144

3-[1-(2,3-Difluoro-phenyl)-1*H*-tetrazol-5-yl]-1*H*-pyrrolo
[2,3-*b*] pyridine (11)

[0387]


5 *N*-(2,3-Difluoro-phenyl)-*N'*-amino-1*H*-pyrrolo[2,3-*b*]pyridine-3-carboxamide (prepared by using procedures described in Method G) (20mg, 0.07 mmol) was dissolved in 2N HCl (2 mL). A solution of NaNO₂ (6 mg) in water (1 mL) was added at 0°C. The mixture was stirred at
10 0°C for 30 min and neutralized with 6N NaOH. The precipitate was collected by filtration and the crude product was purified by HPLC to afford a white solid (12 mg, 0.04 mmol).


[0388] ¹H NMR (500 MHz, DMSO-*d*₆) δ: 12.47 (s, 1H),
15 8.49 (dd, 1H), 8.40 (dd, 1H), 7.89 (m, 1H), 7.76 (m, 1H), 7.58 (m, 1H), 7.35 (d, 1H), 7.31 (dd, 1H).
LC/MS: Rt 3.00 mins.; m/e 299 (M+H), 297.1 (M-H).



Preparation of

20 Benzyl-(4-{3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-cyclohex-3-enyl)-amine
(11)






Preparation of

- 5 **Benzyl-(4-{3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclohex-3-enyl)-amine, Example 127 (136)**

Example 145



10

Step A:


Benzyl-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-cyclohex-3-enyl]-carbamic acid *tert*-butyl ester [0389] A mixture of

- 15 4-[benzyl(*tert*-butoxycarbonyl)amino]cyclohexenyl trifluoromethanesulfonate (1.66 g, 3.81 mmol, prepared according to *Tetrahedron* 53 (1997) 1391-1402), bis(pinacolato)diboron (1.06 g, 4.17 mmol), KOAc (1.11 g, 11.3 mmol) and PdCl₂(dppf) (155 mg, 0.19 mmol) in
- 20 dioxane (20 mL) was degassed at RT, stirred under N₂ at 80°C for 15 hours and then concentrated. The residue was dissolved in EtOAc, and washed with water. Purification by Flash Chromatography (FC) (hexane/EtOAc

- 206 -

50:2 to 50:5) gave the title product (1.04 g) in 66.1% yield. FIA-MS: $m/e = 414.3$ ($M+1$).

Example 146




5 **Step B:**

1-(5-Bromo-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-(2,3-difluoro-phenyl)-ethanone

[0390] 2,3-Difluorophenylacetyl chloride (58.1 mmol) in CH_2Cl_2 (150 ml) was added to a suspension of
10 5-bromo-1H-pyrrolo[2,3-b]pyridine (8.0 g, 40.6 mmol) and AlCl_3 (46 g, 345 mmol) in CH_2Cl_2 (100 ml) at 0°C . After the addition, the cooling bath was removed and the reaction mixture was stirred at room temperature for 1.5 hrs. The reaction was cooled with ice-bath and
15 methanol (100ml) was added to the reaction mixture while maintaining the temperature below 30°C . The reaction mixture was evaporated and the residue was suspended in water. The solid was collected by vacuum filtration, washed with water and hexane to give the
20 title compound (14.03 g) in 98% yield.

[0391] ^1H -NMR (500 MHz, CDCl_3): 10.93 (br.s, 1H), 8.95 (s, 1H), 8.39 (s, 1H), 8.10 (s, 1H), 7.01 (m, 3H), 4.16 (s, 2H).

- 207 -

Example 147**Step C:**

1- [5- (4-Benzylamino-cyclohexyl) -1H-pyrrolo[2,3-b]pyridi
 5 n-3-yl] -2- (2,3-difluoro-phenyl) -ethanone

[0392] A mixture of

benzyl- [4- (4,4,5,5-tetramethyl- [1,3,2] -dioxo-borolan-2-
 yl) -cyclohex-3-enyl] -carbamic acid *tert*-butyl ester
 (820 mg, 1.98 mmol from a)) and

10 1- (5-Bromo-1H-pyrrolo[2,3-b]pyridin-3-yl) -2- (2,3-difluo
 ro-phenyl) -ethanone (702 mg, 2.0 mmol, from b)),
 Pd(dppf)Cl₂ (163 mg, 0.2 mmol), 2.0M aqueous Na₂CO₃ (2
 ml, 4 mmol) in DMF (15 ml) was degassed and heated at
 80°C under N₂ for 3 days. The mixture was concentrated.


15 [0393] The residue was suspended in CH₂Cl₂, washed
 with saturated NH₄Cl. Purification by flash
 chromatography gave
 benzyl- (4- {3- [(2,3-difluoro-phenyl) -acetyl] -1H-pyrrolo[
 2,3-b]pyridin-5-yl} -cyclohex-3-enyl) -carbamic acid
 20 *tert*-butyl ester (545 mg). FIA-MS *m/e*= 558.3 (M+H),
 556.3 (M-H).

[0394] A solution of the above carbamic acid
tert-butyl ester (306 mg, 0.54 mmol) was dissolved in
 CH₂Cl₂ (1 ml), treated with TFA (3 ml) for 1 hour. The
 25 reaction was evaporated and the solid residue
 triturated with ether and hexane, and filtered to give
 the final desired product as a white solid (402 mg) in

- 208 -

80% overall yield. FIA-MS m/e = 458.2 (M+H), 456.3 (M-H).

Example 148



5 **Step D:**

Benzyl-(4-{3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclohex-3-enyl)-amine (136)


[0395] A mixture of

10 1-[5-(4-benzylamino-cyclohexyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-2-(2,3-difluoro-phenyl)-ethanone (150 mg, 0.27 mmol. Preparation: see below) and
t-butoxybis(dimethylamino)methane (0.27 ml, 1.0 mmol, Brederck's reagent) in THF (5 ml) was heated at 60°C
15 for 2 h. The reaction mixture was concentrated and dried under high vacuum for 0.5 h. The residue was dissolved in Ethanol (10 ml), refluxed with
hydroxylamine hydrogen chloride (94 mg, 1.35 mmol) and sodium acetate (1.62 mmol) for 2.5 h. The mixture was
20 concentrated and suspended in satd. NaHCO₃ and filtered. The solid was further purified by chromatography to give 50 mg of the desired product in 38.4% yield.

[0396] ¹H NMR (500 MHz, DMSO-d₆) δ : 12.41 (s, NH), 8.86 (s, 1H), 8.85 (m, 2H, NH₂), 8.45 (d, 1H), 7.81 (d, 1H), 7.70 (d, 1H), 7.5 (m, 6H), 7.32 (m, 2H), 5.98 (br.s, 1H), 4.28 (t, 2H), 3.39 (m, 1H), 2.72 (m, 1H), 2.4 (m, 4H), 1.79 (m, 1H). LC/MS: Rt 2.62 mins.; m/e 483.3 (M+H), 481.3 (M-H).

Example 149

4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-cyclohex-3-enylamine (137)



5

[0397] A suspension

1-(5-bromo-1H-pyrrolo[2,3-b]pyridin-3-yl)-2-(2,3-difluoro-phenyl)-ethanone (1.053g, 3 mmol) and bis(pinacolato)diboron (840 mg, 3.3 mmol), PdCl₂(PPh₃)₂ (62 mg, 0.15 mmol) and KOAc (882 mg, 9.0 mmol) in 1,4-dioxane (18 ml) was heated at 80°C under Argon for 3 hours, and then concentrated. The residue was suspended in water and filtration gave 1.15 g of 2-(2,3-Difluoro-phenyl)-1-[5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-ethanone in 96% yield. LC-MS: m/e = 399.3 (M+H), 397.3 (M-H).

[0398] To a solution of the above boronic ester (1.0 g, 2.5 mmol), trifluoro-methanesulfonic acid 4-tert-butoxycarbonylamino-cyclohex-1-enyl ester (690 mg, 2.0 mmol, prepared from (4-Oxo-cyclohexyl)-carbamic acid tert-butyl ester (Heterocycles 58 (2002) 471-504)) in a mixed solvent of DMF (15 ml) and DMSO (6 ml) was added aqueous 2.0M Na₂CO₃ (2ml, 4 mmol) and PdCl₂(PPh₃)₂.

20

- 210 -

The mixture was degassed and heated at 80°C under Argon for 72 h, and then concentrated under high vacuum to remove DMF. The residue was suspended in aqueous NaHCO₃, then filtered. The solid was purified for FC to give


(4-{3-[(2,3-difluoro-phenyl)-acetyl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclohex-3-enyl)-carbamic acid *tert*-butyl ester as a white solid (530 mg) in 56% yield, LC-MS: *m/e* = 468.4 (M+H), 466.4 (M-H). ¹H-NMR (500MHz, CDCl₃): 13.41 (br.s, 1H), 9.23 (s, 1H), 8.44 (s, 1H), 8.38 (s, 1H), 7.09 (m, 3H), 6.24 (s, 1H), 4.56 (br. s, 1H), 4.32 (s, 2H), 3.90 (br.s, 1H), 2.63 (m, 3H), 2.13 (m, 2H), 1.28 (m, 1H), 1.49 (s, 9H).

[0399] Treatment of the above carbamic ester (330 mg, 0.7 mmol) with Bredeek's reagent followed with hydroxylamine hydrogen chloride (according to General Method K) then with TFA gave 300 mg of the final product.

[0400] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.42 (s, 1H, NH), 8.86 (s, 1H), 8.43 (s, 1H), 7.90 (br.s, 3H, NH₃), 7.80 (s, 1H), 7.70 (s, 1H), 7.51 (m, 1H), 7.32 (m, 2H), 5.94 (s, 1H), 3.36 (m, 1H), 2.56 (m, 1H), 2.45 (m, 2H), 2.36 (s, 3H), 2.23 (m, 1H), 2.08 (m, 1H), 1.75 (m, 1H) LC/MS: Rt 2.10 mins.; *m/e* 393.3 (M+H), 391.4 (M-H).

Example 150

(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
o[2,3-b]pyridin-5-yl}-cyclohex-3-enyl)-dimethyl-amine
(138)



5 [0401] A mixture of


4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-cyclohex-3-enylamine (47.8 mg,
0.12 mmol), 37% formaldehyde (0.5 ml) and formic acid
(1 ml) was heated at 80°C for 10 h. The mixture was
10 concentrated and purified by HPLC to give 17.3 mg of
the diamine product in 34.5% yield.

[0402] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.43 (s, 1H,
NH), 9.48 (s, 1H, MeOH), 8.86 (s, 1H), 8.45 (d, 1H),
7.82 (d, 1H), 7.68 (d, 1H), 7.52 (m, 1H), 7.34 (m, 1H),
15 7.30 (m, 1H), 5.98 (s, 1H), 3.45 (m, 1H), 3.83 (s, 3H,
NCH₃), 3.82 (s, 3H, NCH₃), 2.7-2.4 (m, 4H), 2.31 (s,
3H, MeOH), 2.22 (m, 1H), 1.75 (m, 1H). LC/MS: Rt
2.22 mins.; m/e 421.4 (M+H), 419.4 (M-H).

20

Example 151

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-5-(4-morpholi
n-4-yl-cyclohex-1-enyl)-1H-pyrrolo[2,3-b]pyridine (139)



[0403] A solution of 1,4-anhydroerythritol (208 mg,
25 2 mmol) in water was treated with sodium periodate (400
mg, 4 mmol) for 12 h, transferred to a solution of


- 212 -

4-{3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
[2,3-b]pyridin-5-yl}-cyclohex-3-enylamine (50 mg, 0.12
mmol) in methanol (5 ml). The resulting solution was
treated with sodium cyanoborohydride. After the
5 reaction was completed, TFA was added, and the mixture
was concentrated and purified by HPLC to give the
morpholine product (20 mg) in 36% yield.

[0404] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.43 (s, 1H,
NH), 9.63 (br. s, MsOH), 8.86 (s, 1H), 8.46 (d, 1H),
10 7.82 (d, 1H), 7.69 (d, 1H), 7.51 (q, 1H), 7.34 (q, 1H),
7.29 (q, 1H), 5.99 (d, 1H), 4.05 (d, 2H), 3.72 (t, 2H),
3.56-3.48 (m, 3H), 3.18 (m, 2H), 2.61 (m, 2H), 2.44
(m, 2H), 2.33 (m, 1H), 2.32 (s, 3H, MsOH), 1.75 (ddd,
1H). LC/MS: Rt 2.22 mins.; m/e 463.4 (M+H), 461.4
15 (M-H).

Example 152

N-(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrr
olo[2,3-b]pyridin-5-yl}-cyclohex-3-enyl)-methanesulfona
20 mide (140)



[0405] To a solution of
4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo
25 [2,3-b]pyridin-5-yl}-cyclohex-3-enylamine (47 mg, 0.12
mmol) in DMF (2 ml) was added methanesulfonic acid
benzotriazol-1-yl ester (27 mg, 0.13 mmol) and Hunig
base (10 drops) and stirred for 1 h. TFA was added,

- 213 -


and the reaction mixture was purified by HPLC to give the methanesulfonamide (25.9 mg) in 46% yield.

[0406] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.42 (s, 1H, NH), 8.86 (s, 1H), 8.42 (s, 1H), 7.82 (s, 1H), 7.63 (s, 1H), 7.53 (m, 1H), 7.31 (m, 2H), 7.09 (br.s, 1H), 5.90 (s, 1H), 3.46 (m, 1H), 2.96 (s, 3H), 2.7-2.4 (m, 3H), 2.39 (s, 3H, MSOH), 2.14 (m, 1H), 2.01 (m, 1H), 1.66 (m, 1H). LC/MS: Rt 3.38 mins.; m/e 471.3 (M+H), 469.3 (M-H).

10

Example 153

(4-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclohexyl)-dimethylamine (141)



15

[0407] A mixture of trifluoro-methanesulfonic acid 4-*tert*-butoxycarbonylamino-cyclohex-1-enyl ester (730 mg, 20 mmol),

5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridine (490 mg, 20 mmol), Pd(dppf)Cl₂ (80 mg) and 2.0M Na₂CO₃ (3 ml, 6 mmol) in DMF (20 ml) was heated at 80°C under N₂ for 3 hours, and then concentrated. The residue was suspended in water, filtered and the solid was purified by FC to give

25

- 214 -

[4-(1H-Pyrrolo[2,3-b]pyridin-5-yl)-cyclohex-3-enyl]-carbamic acid tert-butyl ester (600mg) in 96% yield.

FIA-MS: $m/e = 314.05$ (M+H). A suspension of the above cyclohexene (300mg) and 10% Pd/C (200mg) in a 1:1 mixed solvent of EtOAc and MeOH (30ml) was shaken under 50psi H₂ for 20 h. Filtration gave

[4-(1H-pyrrolo[2,3-b]pyridin-5-yl)-cyclohexyl]-carbamic acid tert-butyl ester (280 mg). FIA-MS: $m/e = 316.3$ (M+H).

10 [0408] A solution of 2,3-difluorophenylacetyl chloride (1.0 mmol) in CH₂Cl₂ (3 ml) was added dropwise to a mixture of

[4-(1H-pyrrolo[2,3-b]pyridin-5-yl)-cyclohexyl]-carbamic acid tert-butyl ester (280 mg, 0.88 mmol) and AlCl₃ (612mg, 4.6 mmol) in CH₂Cl₂ (15 ml) at 0°C. After the addition, the reaction was stirred for 1.5 hrs at 0°C. Methanol (5ml) was added to the reaction. After 1 hr. the reaction was evaporated and the resulting residue purified by flash chromatography to afford

20 1-[5-(4-Amino-cyclohexyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-2-(2,3-difluoro-phenyl)-ethanone (229 mg) in 70.5% yield. LC-MS: $m/e = 370.3$ (M+H), 368.4 (M-H). After the treatment of the above cyclohexylamine (220 mg, 0.60 mmol) with 37% formaldehyde (2 ml) and formic acid (4 ml) at 80°C for 15 h, pure


2-(2,3-Difluoro-phenyl)-1-[5-(4-dimethylamino-cyclohexyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-ethanone was obtained by FC (167 mg) in 70.1% yield. LC-MS: $m/e: 398.4$ (M+H), 396.4 (M-H).

30 [0409] The final desired product, (4-{3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclohexyl)-dimethyl-amine, was prepared according to Method K by treatment of

- 215 -

2-(2,3-Difluoro-phenyl)-1-[5-(4-dimethylamino-cyclohexyl)-1H-pyrrolo[2,3-b]pyridin-3-yl]-ethanone (200 mg, 0.5 mmol) with Brederick's reagent followed by hydroxylamine in 12.5% yield.

- 5 [0410] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.33 (s, 1H, NH), 9.39 (br. s, OH), 8.85 (s, 1H), 8.23 (s, 1H), 7.78 (s, 1H), 7.57 (s, 1H), 7.52 (m, 1H), 7.33 (m, 2H), 3.22 (m, 1H), 2.77 (s, 6H), 2.64 (m, 1H), 2.33 (s, 3H), 2.09 (d, 2H), 1.90 (d, 2H), 1.59 (m, 2H), 1.45 (m, 2H).
- 10 LC/MS: Rt 2.10 mins.; m/e 423.4 (M+H), 421.5 (M-H).

Scheme for I-142 and I-143

- 217 -

Example 154

[0411] Experimental for 5-cyclopropylazindole compounds:

2:

5 1-(Toluene-4-sulfonyl)-5-vinyl-1H-pyrrolo[2,3-b]pyridine:

[0412]

5-Bromo-1-(toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine, (300mg, 0.85mmol), vinylboronic acid
10 dibutylester, (184mg, 1.0mmol), potassium carbonate, (420mg, 3.0mmol) and Pd(Ph₃P)₄ were combined in 3mL DME and 1mL water in a tube under nitrogen and heated in a microwave reactor to 130° C for 10 min. The organic layer was then separated and evaporated. The residue
15 was purified by silica chromatography (eluent: methylene chloride), affording 215 mg(85%)
1-(Toluene-4-sulfonyl)-5-vinyl-1H-pyrrolo[2,3-b]pyridine. MS ES+ 299.0.

3:

20 2-[1-(Toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridin-5-yl]-cyclopropanecarboxylic acid ethyl ester:

[0413]

1-(Toluene-4-sulfonyl)-5-vinyl-1H-pyrrolo[2,3-b]pyridine, (200mg, 0.67mmol), and ethyl diazoacetate, (730μL, 7.0mmol), were combined in 3mL xylene and heated to
25 95°C for 30min, then to 115°C for 4hours. The solvent was evaporated onder vacuum and the residue purified by silica chromatography, (eluent: methylene chloride), affording 140mg, (50%),
30 2-[1-(Toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridin-5-yl]

- 218 -

] -cyclopropanecarboxylic acid ethyl ester as a mixture of cis and trans isomers. MS ES+ 385.3

4:

2 - (1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl) -cyclopropanecarboxylic acid ethyl ester:

[0414]

2 - [1 - (Toluene-4-sulfonyl) -1*H*-pyrrolo[2,3-*b*]pyridin-5-yl] -cyclopropanecarboxylic acid ethyl ester, (153mg, 0.4mmol), was added to 2mL ethanol and sodium ethoxide solution, (~2.68M, 300μL, 0.8mmol), was added. The mixture was heated to 70°C for 3 hours. The mixture was cooled to rt and several mL sat. ammonium chloride solution were added. The mixture was extracted with dichloromethane and the organic layers evaporated affording 85mg essentially pure 2 - (1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl) -cyclopropanecarboxylic acid ethyl ester, which was used without further purification. MS ES+ 231.1

20 5:

2 - {3 - [(2,3-Difluoro-phenyl) -acetyl] -1*H*-pyrrolo[2,3-*b*]pyridin-5-yl} -cyclopropanecarboxylic acid ethyl ester

[0415] Was prepared from

2 - (1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl) -cyclopropanecarboxylic acid ethyl ester using a procedure similar to those described previously in this document. MS ES+ 385.2

- 219 -

6:

2-{3-[(2,3-Difluoro-phenyl)-acetyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-cyclopropanecarboxylic acid
[0416]

5 2-{3-[(2,3-Difluoro-phenyl)-acetyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-cyclopropanecarboxylic acid ethyl ester, (130mg, 0.33mmol, was dissolved in 1mL ethanol and 1mL 10% potassium carbonate solution. The mixture was heated to reflux for ~16 hours. The reaction was then
10 acidified with 6N HCl to a pH ~4-5 and extracted with dichloromethane. The essentially pure 2-{3-[(2,3-Difluoro-phenyl)-acetyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-cyclopropanecarboxylic acid was used without further purification in the next step.

15 MS ES+ 357

7(trans) and 8(cis):

2-(2,3-Difluoro-phenyl)-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-ethanone

20 [0417]

2-{3-[(2,3-Difluoro-phenyl)-acetyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-cyclopropanecarboxylic acid, (93mg, 0.26mmol), was combined with DIEA, (67mg, 0.52mmol), and HBTU, (113mg, 0.3mmol) in 1mL DMF and stirred at rt
25 for ~15 min. Then N-methyl piperazine, (26mg, 0.26mmol), was added and the reaction stirred at rt for 4 hours. The DMF was evaporated under vacuum, water added and the suspension extracted with dichloromethane. The solvent was evaporated under
30 vacuum and the residue purified by silica chromatography, (eluent: 5% methanol/DCM), affording

- 220 -

trans-2-(2,3-Difluoro-phenyl)-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-ethanone, (35mg, Rf 0.5) and the *cis* isomer (16mg, Rf 0.3).

5 **9:**

***trans*-2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-propenone**
[0418]

10 *trans*-2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-propenone was prepared from
15 *trans*-2-(2,3-Difluoro-phenyl)-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-ethanone, using a procedure similar to those described previously in this document. It was isolated, but not purified and converted directly to the corresponding isoxazole.

10:

20 ***cis*-2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-propenone**
[0419]

25 *cis*-2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-propenone was prepared from
30 *cis*-2-(2,3-Difluoro-phenyl)-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl}-ethanone, using a procedure similar to those described previously in this document. It was

- 221 -

isolated, but not purified and converted directly to the corresponding isoxazole.

11:

trans-(2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclopropyl)-(4-methyl-piperazin-1-yl)-methanone (142)

[0420]

trans-(2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclopropyl)-(4-methyl-piperazin-1-yl)-methanone (142) was prepared from trans-2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1H-pyrrolo[2,3-b]pyridin-3-yl}-propenone using a procedure similar to those described previously in this document. It was purified by prep hplc and isolated as the TFA salt. LC/MS: Rt 2.29 mins.; m/e 464.2 (M+H).

[0421] ¹H NMR (500 MHz, MeOH-d₄) δ: 8.60 (s, 1H), 8.22 (s, 1H), 7.80 (bs, 1H), 7.65 (s, 1H), 7.35 (m, 1H), 7.25 (m, 2H), 4.8-4.4 (m, 2H), 3.6-3.0 (m, 6H), 2.95 (s, 3H), 2.6 (m, 1H), 2.25 (m, 1H), 1.60 (m, 1H), 1.35 (m, 1H).

12:

cis-(2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclopropyl)-(4-methyl-piperazin-1-yl)-methanone (143)

[0422]


cis-(2-{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-5-yl}-cyclopropyl)-(4-methyl-piperazin-1-yl)-methanone (143) was prepared from cis-2-(2,3-Difluoro-phenyl)-3-dimethylamino-1-{5-[2-(4-methyl-piperazine-1-carbonyl)-cyclopropyl]-1H-pyrrolo[2

- 222 -

,3-*b*]pyridin-3-yl}-propenone using a procedure similar to those described previously in this document. It was purified by prep hplc and isolated as the TFA salt.

LC/MS: Rt 2.24 mins.; m/e 464.2 (M+H)

5 [0423] ^1H NMR (500 MHz, MeOH- d_4) δ : 8.65 (s, 1H), 8.25 (bs, 1H), 8.05 (bs, 1H), 7.55 (s, 1H), 7.35 (m, 1H), 7.25 (m, 2H), 5.0-4.0 (m, 2H), 3.6-3.0 (m, 6H), 2.85 (bs, 3H), 2.75 (m, 1H), 2.45 (m, 1H), 1.75 (m, 1H), 1.45 (m, 1H).




10

[0424] a) CO, Pd(OAc) $_2$, Ph $_3$ P, MeOH, TEA, DMF; b) NaOMe / MeOH c) LAH, THF 70 C; d) AlCl $_3$ Methylene chloride; e) 1N NaOH, Methanol; f) Bredereck's reagent, THF, 80C; g) NH $_2$ OH HCl, NaCO $_3$, THF 80C

15

- 223 -




[0425] a) 1) (2,2-Dimethoxy-ethyl)-methyl-amine,
 5 80C, acetonitrile ; b) Et₂O, POCl₃; c) Brederick's
 reagent, THF, 80C; d) NH₂OH HCl, NaCO₃, THF 80C

Preparation of

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-6-methyl-1,6-
 dihydro-pyrrolo[2,3-c]pyridin-7-one

10

Example 155




- 224 -

Step A:

4-[(2,3-Difluoro-phenyl)-acetyl]-1H-pyrrole-2-carboxylic acid (2,2-dimethoxy-ethyl)-methyl-amide
[0426]

5 2,2,2-Trichloro-1-{4-[(2,3-difluoro-phenyl)-acetyl]-1H-pyrrol-2-yl}-ethanone (300mg, 0.82mmol) and
(2,2-Dimethoxy-ethyl)-methyl-amine (116 μ L, 0.900mmol)
in acetonitrile were heated for 5 hr. Evaporation
afforded 299mg (99%) of
10 4-[(2,3-Difluoro-phenyl)-acetyl]-1H-pyrrole-2-carboxylic acid (2,2-dimethoxy-ethyl)-methyl-amide. ^1H NMR (500
MHz, CDCl_3) δ : 10.1-9.9 (1H, vbs), 7.02 - 6.96 (5H,
cm), 4.51 (1H, s), 4.05 (2H, s) 3.59 (3H, bs),
3.39-3.36 (6H, s). LC/MS: Rt 3.13 mins.; m/e 367.26
15 (M+H), 365.34 (M-H).

Example 156**Step B:**

3-[(2,3-Difluoro-phenyl)-acetyl]-6-methyl-1,6-dihydro-p
20 yrrolo[2,3-c]pyridin-7-one
[0427] To


4-[(2,3-Difluoro-phenyl)-acetyl]-1H-pyrrole-2-carboxylic
c acid (2,2-dimethoxy-ethyl)-methyl-amide (200mg,
0.55mmol) in dioxane (50 mL) was added POCl_3 (50 μ L,
25 0.55mmol) at 0C. The reaction solution was heated to
60C and stirred for 14hr. The reaction was quenched
with water and extracted with ethyl acetate, dried

- 225 -

(Na₂SO₄). Flash chromatography (methylene chloride / methanol) gave 67mg (41% yield) of title compound.

[0428] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.8 (1H, bs), 8.34-8.33 (1H, d), 7.66-7.65 (1H, m) 7.36-7.32 (2H, cm), 7.16-7.15 (2H, m), 6.94-6.92 (1H, d), 4.35 (2H, s) 3.52 (3H, bs). LC/MS: 2.61 mins.; m/e 303.2 (M+H), 301.2 (M-H).

Example 157



10

Step C:


3-[3-Dimethylamino-2-(2,3-difluoro-phenyl)-acryloyl]-6-methyl-1,6-dihydro-pyrrolo[2,3-c]pyridin-7-one

[0429] To

15 3-[(2-Fluoro-phenyl)-acetyl]-6-methyl-1,6-dihydro-pyrrolo[2,3-c]pyridin-7-one (67mg, 0.22mmol) in THF (20mL) was added Brederick's reagent (183μL, 0.89mmol) and the reaction was heated to 80 C overnight. Concentration under reduced vacuum a red oil, used as obtained.

20 LC/MS: Rt 2.32 mins.; m/e 330.3 (M⁺-27), M⁻ 256.3 (M-27) and Rt 2.60 m/e M- 329.3 (m-27).

- 226 -


Example 158

3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-6-methyl-1,6-
 5 dihydro-pyrrolo[2,3-c]pyridin-7-one (144)

[0430] To 3-[3-Dimethylamino-2-(2-fluoro-phenyl)-acryloyl]-6-methyl-1,6-dihydro-pyrrolo [2,3-c] pyridin-7-one (79mg, 0.22mmol) in tetrahydrofuran (20mL) was added sodium
 10 hydrogen carbonate (19mg, 0.0.27mmol) and hydroxylamine Hydrochloride (23mg, 0.27mmol) and the mixture was heated to 80C. After 5hr p-toluenesulfonic acid (catalytic amount) was added and the reaction mixture was heated for an additional 1hr. The solution was
 15 cooled, diluted with ethyl acetate, washed with brine, dried (Na₂SO₄). Concentration gave an amber oil. Preparative reverse phase chromatography afforded 3-[4-(2-Fluoro-phenyl)-isoxazol-5-yl]-6-methyl-1,6-dihydro-pyrrolo[2,3-c]pyridin-7-one (18.6 mg, 25% yield).

20 [0431] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.37 (1H, bs), 8.85 (1H, s), 7.655-7.650 (1H, d), 7.51-7.50 (1H, q), 7.39-7.37 (1H, d), 7.34-7.28 (2H, cm), 7.09-7.06 (2H, m). LC/MS: Rt 3.21mins.; m/e 328.1 (M+H), 326.2 (M-H).

- 227 -


Example 159

[0432]

5 3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1,6-dihydro-pyrrolo[2,3-c]pyridin-7-one (145)

[0433] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.65 (s, 0.75H), 12.31 (s, 0.25H), 11.17 (br, 0.75H), 11.09 (br, 0.25H), 9.23 (s, 0.25H), 8.82 (0.75H), 7.50 (m, 1H),
 10 7.41 (s, 1H), 7.30 (m, 2H), 6.97 (m, 1H), 6.68 (d, 0.25H), 6.42 (d, 0.75H)

LC/MS: Rt 3.06 mins.; m/z 314.1 (M+H), 312.2 (M-H)


Example 160

15

[0434]

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-7-oxo-1,7-dihydro-pyrrolo[2,3-c]pyridin-6-yl}-acetic acid methyl ester (146)

20 [0435] ¹H NMR (500 MHz, DMSO-d₆) δ: 12.73 (1H, bs), 8.84 (1H, s), 7.51-7.49 (H, cm), 7.464-7.468 (1H, d), 7.35-7.28 (2H, cm), 7.27-7.26 (1H, d), 6.47-6.46 (1H, d), 4.80 (2H, s), 3.68 (3H, s). LC/MS: Rt 3.21 mins.; m/z 385.91 (M+H), 384.05 (M-H)


Example 161

[0436]

5 6-Benzyl-3-[4-(2,3-difluoro-phenyl)-isoxazol-5-yl]-1,6-dihydro-pyrrolo[2,3-c]pyridin-7-one (147)

[0437] ^1H NMR (500 MHz, DMSO- d_6) δ : 12.71 (1H, bs), 8.83 (1H, s), 7.53-7.47 (H, m), 7.44-7.43 (1H, d), 7.37-7.35 (1H, d), 7.34-7.24 (7H, cm), 6.49-6.47 (1H, d), 5.20 (1H, s). LC/MS: Rt 4.0 mins.; m/z 403.9 (M+H), 402.1 (M-H)

10

Preparation of 31

15


[0438] Reagents and Conditions: (a) (i) LHMDs, THF -78 C, (ii) diethyl oxalate; (b) H_2NNH_2 , (i-PrO) $_4\text{Ti}$, CH_2Cl_2 ; (c) NMP, Microwave (250 C, 5 mins.); (d) LHMDs, 2,3-difluoroacetic acid, THF, 0 C; (e) (i) Bredereck's reagent, THF, reflux, (ii) hydroxylamine hydrochloride, NaOAc, THF, reflux.

20

- 229 -

3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-1H-pyrazolo[3,4-b]pyridine (31)

Example 162



5


Step A: Ethyl 2-(2-fluoropyridin-3-yl)-2-oxoacetate

[0439] To a solution of 2-fluoropyridine (1.0 mL, 11.6 mmol) in THF (30 mL) at -78 under nitrogen was added a solution of lithiumdiisopropylamide in THF/heptane/ethylbenzene. The resulting solution was allowed to stir for 1.5 hours then diethyl oxalate (1.89 mL, 13.9 mmol) was added dropwise via syringe. After 30 minutes, the resulting solution was diluted with EtOAc and washed with saturated NH₄Cl then water. The organic layer was dried over MgSO₄ and concentrated under vacuum to give an oil. Chromatography (20% to 30% EtOAc: hexane) gave 0.68 g (30% yield) of ethyl 2-(2-fluoropyridin-3-yl)-2-oxoacetate as an oil.

10

15

Example 163



20

Step B:


(Z)-Ethyl-2-(2-fluoropyridin-3-yl)-2-hydrazonoacetate & (Z)-Isopropyl-2-(2-fluoropyridin-3-yl)-2-hydrazonoacetate.

25

[0440] To a solution of ethyl 2-(2-fluoropyridin-3-yl)-2-oxoacetate (1.8 g, 9.14

- 230 -

mmol) in dichloromethane (25 mL) was added titanium isopropoxide (5.45 mL, 18.3 mmol) and hydrazine (0.89 mL, 18.3 mmol). The resulting yellow solution was allowed to stir at room temperature for 2.5 hours followed by the addition of water (2 mL). After 2.5 hours, the suspension was filtered through celite and washed with dichloromethane. Concentration of the solvent gave 1.29 grams of a 1:2 mixture of (Z)-ethyl-2-(2-fluoropyridin-3-yl)-2-hydrazonoacetate LC-MS Rt 1.5 min ES+ (212) and (Z)-isopropyl-2-(2-fluoropyridin-3-yl)-2-hydrazonoacetate LC-MS Rt 2.0 min ES+ (226) as a waxy solid.

Example 164

15

Step C: Ethyl 1H-pyrazolo[3,4-b]pyridine-3-carboxylate [0441] A solution of (1.01 g, 4.96 mmol) of a 1:2 mixture


(Z)-ethyl-2-(2-fluoropyridin-3-yl)-2-hydrazonoacetate and

(Z)-isopropyl-2-(2-fluoropyridin-3-yl)-2-hydrazonoacetate in NMP (12 mL) was divided into three 5 mL microwave vessels then heated at 250 C for 5 min. The reactions were combined, diluted with EtOAc and washed with water, saturated sodium chloride then dried (MgSO4) and concentrated to dryness. Chromatography (SiO2, 4:1 to 1:1 EtOAc:hexane) gave 0.59 g (65% yield) of a 2:1 mixture of ethyl

1H-pyrazolo[3,4-b]pyridine-3-carboxylate LC-MS Rt 2.1

- 231 -

min ES+ (192.1), ES- (190.1) and isopropyl
1H-pyrazolo[3,4-b]pyridine-3-carboxylate LC-MS Rt 2.4
min ES+ (206.1), ES- (204.2) as a pink solid.


Example 165

5

Step D:

2-(2,3-Difluorophenyl)-1-(1H-pyrazolo[3,4-b]pyridin-3-yl)ethanone

[0442] To a solution of 2,3-difluorophenylacetic
10 acid (236mg, 1.4 mmol) in THF at 0C was added 3.85 mL
(3.85 mmol) of a solution of lithium
bis-trimethylsilylamide in THF. The resulting solution
was then added to solution of a 2:1 mixture of ethyl
1H-pyrazolo[3,4-b]pyridine-3-carboxylate and isopropyl
15 1H-pyrazolo[3,4-b]pyridine-3-carboxylate (104mg, 0.55
mmol) in THF. The resulting mixture was heated at 75C
for 3 hours. The reaction progress was monitored by
TLC and HPLC and quenched with saturated NH₄Cl, diluted
with EtOAc and washed with saturated NaHCO₃ and brine.
20 The organic layer was dried over MgSO₄ and concentrated
to give 137 mg (91% yield) of
2-(2,3-difluorophenyl)-1-(1H-pyrazolo[3,4-b]pyridin-3-yl)
ethanone. LC-MS Rt= 3.3 min ES+ (274.0) ES- (272.1)


Example 166**Preparation of**

5 **3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-1H-pyrazolo[3,4-b]pyridine (31)**

[0443]


2-(2,3-Difluorophenyl)-1-(1H-pyrazolo[3,4-b]pyridin-3-yl)ethanone (137 mg, 0.5 mmol) was allowed to react
 10 according to the METHOD OF BREDERECKS to afford 112 mg (40% yield) of
 3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-1H-pyrazolo[3,4-b]pyridine.

[0444] ^1H NMR (500 MHz, CD_3OD) δ : 8.75 (s, 1H), 8.60 (d, 2H), 8.46 (d, 2H), 7.48 (m, 1H), 7.35-7.29 (m, 2H) and 7.22 (bm, 1H); ^1H NMR (Acetone- d_6) δ : 8.81 (d, J = 1.3 Hz, 1H) 8.66 (dd, J = 1.5 and 4.5 Hz, 1H) 8.48 (dd, J = 1.5 and 8.2 Hz, 1H) 7.57 (m, 1H) 7.39 (m, 2H) 7.28 (m, 1H). LC/MS Rt 3.30 mins.; m/e 299 (M^+H), 297 (M^-H)



- 233 -

[0445] Reagents and Conditions: (a) (i) LHMDs, THF
 -78 C, (ii) diethyl oxalate; (b) H₂NNH₂, (i-PrO)₄Ti,
 CH₂Cl₂; (c) NMP, Microwave (250 C, 5 mins.); (d) LHMDs,
 2,3-difluoroacetic acid, THF, 0 C; (e)
 5 (i) Brederick's reagent, THF, reflux, (ii)
 hydroxylamine hydrochloride, NaOAc, THF, reflux.


Example 167

10 5-Bromo-3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-1H-pyrazolo[3,4-b]pyridine (149)

[0446]

5-Bromo-3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-1H-pyrazolo[3,4-b]pyridine was prepared according to the method described for the preparation of
 15 3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-1H-pyrazolo[3,4-b]pyridine starting with 5-bromo-2-fluoropyridine.

[0447] ¹H NMR (500 MHz, DMSO-d₆) δ: 14.59 (s, 1H), 9.05 (s, 1H), 8.74 (d, 1H, J = 2.0 Hz), 8.52 (d, 1H, J = 2.0 Hz), 7.55-7.50 (m, 2H) and 7.34-7.30 ppm (m, 1H).
 20 LC/MS Rt 4.0 mins.; m/e 377 (M+H) 375 (M-H)


Example 168

3-(4-(2,3-Difluorophenyl)isoxazol-5-yl)-5-(pyridin-3-yl)-1H-pyrazolo[3,4-b]pyridine (150)

[0448]

3-(4-(2,3-difluorophenyl)isoxazol-5-yl)-5-(pyridin-3-yl)-1H-pyrazolo[3,4-b]pyridine was prepared according to METHOD FOR SUZUKI COUPLING to afford 3.8 mg (13 % yield)

[0449] ^1H NMR (500 MHz, DMSO- d_6) δ : 14.52 (s, 1H), 9.07 (s, 1H), 9.03 (d, $J=2.05$ Hz, 2H), 8.68 (d, $J=4.63$ Hz, 1H), 8.59 (d, $J=1.89$ Hz, 1H), 8.29 (d, $J=7.76$ Hz, 1H), 7.64-7.61 (m, 1H), 7.55-7.50 (m, 2H), and 7.35-7.31 ppm (m, 1H). LC/MS Rt 2.5 mins.; m/e 376 (M+H), 374 (M-H)




15

[0450] Reagents and Conditions: (a) (i) LHMDS, THF -78 C, (ii) diethyl oxalate; (b) H_2NNH_2 , (i-PrO) $_4\text{Ti}$, CH_2Cl_2 ; (c) NMP, Microwave (250 C, 5 mins.); (d) LHMDS, 2,3-difluoroacetic acid, THF, 0 C; (e)

20

- 235 -

(i) Bredereck's reagent, THF, reflux, (ii)
hydroxylamine hydrochloride, NaOAc, THF, reflux; (f)
HNR₁R₂, NMP.


Example 169

5

{3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrazolo[3,4-b]pyridin-6-yl}-ethyl-amine (30):

[0451] ¹H NMR (500 MHz, Acetone-d₆) δ: 1.2 (t, 3H), 3.5 (q, 2H overlap d-solvent), 6.6 (d, 1H), 7.2 (m, 1H), 7.35 (m, 1H), 7.55 (t, 1H), 7.95 (d, 1H), 8.7 (s, 1H). LC/MS: Rt 3.3 mins.; m/e 342 (M+H), 340.2 (M-H).

10

Example 170

15 {3-[4-(2,3-Difluoro-phenyl)-isoxazol-5-yl]-1H-pyrrolo[2,3-b]pyridin-6-yl}-dimethyl-amine (32):

[0452] ¹H NMR (500 MHz, Acetone-d₆) δ: 3.2 (s, 6H), 6.8 (d, 1H), 7.25 (m, 1H), 7.3 (m, 1H), 7.6 (t, 1H), 8.1 (d, 1H), 8.7 (s, 1H). LC/MS: Rt 3.6 mins.; m/e 342 (M+H), 340.1 (M-H)

20

- 236 -

Example 171K_i Determination for the Inhibition of c-Met

[0453] Compounds were screened for their ability to inhibit c-Met kinase activity using a standard coupled enzyme system (Fox *et al.*, *Protein Sci.* 1998, 7, 2249). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 300 μM NADH, 1 mM DTT, and 1.5% DMSO. Final substrate concentrations in the assay were 200 μM ATP (Sigma Chemicals, St Louis, MO) and 10 μM polyGluTyr (Sigma Chemical Company, St. Louis). Reactions were carried out at 30 °C and 80 nM c-Met. Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 300 μM NADH, 30 μg/ml pyruvate kinase and 10 μg/ml lactate dehydrogenase.

[0454] An assay stock buffer solution was prepared containing all of the reagents listed above with the exception of ATP and a test compound of the present invention. The assay stock buffer solution (175 μl) was incubated in a 96 well plate with 5 μl of the test compound of the present invention at final concentrations spanning 0.006 μM to 12.5 μM at 30°C for 10 minutes. Typically, a 12-point titration was conducted by preparing serial dilutions (from 10 mM compound stocks) with DMSO of the test compounds of the present invention in daughter plates. The reaction was initiated by the addition of 20 μl of ATP (final concentration 200 μM). Rates of reaction were obtained using a Molecular Devices Spectramax plate reader (Sunnyvale, CA) over 10 minutes at 30°C. The K_i values

- 237 -

were determined from the rate data as a function of inhibitor concentration.

[0455] Compounds of the present invention were found to be inhibitors of c-Met. Compounds 1, 2, 37, 38, 40, 42, 43, 44, 46, 48, 49, 53, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 102, 105, 106, 109, 110, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 129, 130, 131, 133, 135, 136, 137, 138, 139, 140, 141, 142, 149, 153, 154, and 155 had Ki values of <0.2 μ M. Compounds 31, 35, 39, 41, 45, 55, 103, 104, 108, 111, 132, 134, 143, 144, and 148 had Ki values of 0.2 μ M-<1.0 μ M. Compounds 11, 29, 30, 32, 36, 47, 51, 52, 54, 56, 80, 100, 101, 107, 143, 145, 146, 147, 151, and 152 had Ki values of 1.0 μ M-12.5 μ M.

Example 172

GSK-3 Inhibition Assay:

[0456] Compounds of the present invention were screened for their ability to inhibit GSK-3 β (AA 1-420) activity using a standard coupled enzyme system (Fox et al., Protein Sci. 1998, 7, 2249). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 300 μ M NADH, 1 mM DTT and 1.5% DMSO. Final substrate concentrations in the assay were 20 μ M ATP (Sigma Chemicals, St Louis, MO) and 300 μ M peptide (American Peptide, Sunnyvale, CA). Reactions were carried out at 30°C and 20 nM GSK-3 β . Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 300 μ M NADH, 30 μ g/ml pyruvate kinase and 10 μ g/ml lactate dehydrogenase.

- 238 -

[0457] An assay stock buffer solution was prepared containing all of the reagents listed above with the exception of ATP and the test compound of the present invention. The assay stock buffer solution (175 μ l) was incubated in a 96 well plate with 5 μ l of the test compound of the present invention at final concentrations spanning 0.002 μ M to 30 μ M at 30°C for 10 min. Typically, a 12 point titration was conducted by preparing serial dilutions (from 10 mM compound stocks) with DMSO of the test compounds of the present invention in daughter plates. The reaction was initiated by the addition of 20 μ l of ATP (final concentration 20 μ M). Rates of reaction were obtained using a Molecular Devices Spectramax plate reader (Sunnyvale, CA) over 10 min at 30°C. The K_i values were determined from the rate data as a function of inhibitor concentration.

[0458] Compounds of the present invention were found to inhibit GSK3. Compounds 63, 64, 67, 68, 69, 70, 72, 82, 86, 90, 97, 98, 99, 106, 109, 113, 114, 133, 135, 137, 138, 139, 140, 141, 149, and 155 had K_i values of <0.2 μ M. Compounds 36, 37, 38, 43, 44, 49, 53, 57, 58, 59, 60, 61, 62, 65, 66, 71, 76, 77, 78, 79, 81, 83, 84, 88, 89, 92, 93, 94, 96, 105, 110, 112, 115, 116, 117, 118, 119, 132, 134, 136, 142, and 145 had K_i values of 0.2-<1.0 μ M. Compounds 1, 2, 29, 33, 34, 35, 39, 40, 41, 42, 45, 46, 47, 48, 52, 55, 75, 80, 85, 87, 91, 95, 102, 103, 104, 108, 111, 143, 148, 150, 151, 153, and 154 had K_i values of 1.0-12.5 μ M.

- 239 -

Example 173JAK3 Inhibition Assay

[0459] Compound inhibition of JAK3 was assayed by the method described by G. R. Brown, et al., Bioorg. Med. Chem. Lett. 2000, vol. 10, pp 575-579 in the following manner. Into Maxisorb plates, previously coated at 4°C with Poly (Glu, Ala, Tyr) 6:3:1 then washed with phosphate buffered saline 0.05% and Tween (PBST), was added 2 mM ATP, 5 mM MgCl₂, and a solution of compound in DMSO. The reaction was started with JAK enzyme and the plates incubated for 60 minutes at 30°C. The plates were then washed with PBST, 100 mL HRP-Conjugated 4G10 antibody was added, and the plate incubated for 90 minutes at 30°C. The plate was again washed with PBST, 100 mL TMB solution is added, and the plates were incubated for another 30 minutes at 30°C. Sulfuric acid (100 mL of 1M) was added to stop the reaction and the plate is read at 450 nm to obtain the optical densities for analysis to determine IC₅₀ values.

[0460] Compounds of the present invention were found to inhibit JAK3. Compounds 37, 38, 41, 43, 48, 49, 57, 58, 59, 65, 66, 67, 68, 70, 71, 73, 75, 76, 77, 79, 82, 83, 84, 86, 87, 88, 90, 91, 92, 93, 94, 95, 96, 98, 99, 105, 133, 135, 136, 137, 138, 139, 140, 153, 154, and 155 had Ki values of <0.2 μM. Compounds 1, 2, 35, 39, 42, 44, 46, 47, 51, 53, 56, 61, 63, 64, 69, 74, 78, 81, 85, 97, 106, 109, 110, 116, 118, 124, 129, 130, 134, 141, 142, 145, and 149 had Ki values of 0.2-<1.0 μM. Compounds 31, 34, 36, 40, 45, 50, 52, 55, 60, 62, 72, 80, 102, 103, 104, 108, 111, 112, 113, 114, 115, 117,

- 240 -

119, 120, 121, 122, 123, 131, 132, 143, 144, 150, and 152 had K_i values of 1.0-12.5 μ M.

Example 174

SYK Enzyme Assay

5 **[0461]** Compounds were screened for their ability to inhibit SYK using a standard coupled enzyme assay (Fox *et al.*, *Protein Sci.* 1998, 7, 2249). Reactions were carried out in 100 mM HEPES (pH 7.5), 10 mM $MgCl_2$, 25 mM NaCl, 1 mM DTT and 1.5% DMSO. Final substrate
10 concentrations in the assay were 200 μ M ATP (Sigma chemical Co.) and 4 μ M poly Gly-Tyr peptide (Sigma Chemical Co.). Assays were carried out at 30 °C and 200 nM SYK. Final concentrations of the components of the coupled enzyme system were 2.5 mM
15 phosphoenolpyruvate, 300 μ M NADH, 30 μ g/ml pyruvate kinase and 10 μ g/ml lactate dehydrogenase.

[0462] An assay stock buffer solution was prepared containing all of the reagents listed above, with the exception of SYK, DTT, and the test compound of
20 interest of the present invention. 56 μ l of the test reaction was placed in a 96 well plate followed by the addition of 1 μ l of 2 mM DMSO stock containing the test compound of the present invention (final compound concentration 30 μ M). The plate was pre-incubated for
25 ~10 minutes at 30°C and the reaction initiated by the addition of 10 μ l of enzyme (final concentration 25 nM). Rates of reaction were obtained using a BioRad Ultramark plate reader (Hercules, CA) over a 5 minute
30 read time at 30°C, and K_i values for the compounds of the present invention were determined according to standard methods.

- 241 -

[0463] Compounds of the present invention were found to inhibit SYK. Compounds 1, 43, 67, 68, 70, 77, 86, 90, 99, 135, and 155 had K_i values of 0.2- $<1.0 \mu\text{M}$. Compounds 29, 35, 37, 38, 41, 42, 44, 61, 65, 71, 73, 76, 82, 84, 88, 91, 93, 98, 106, 133, and 134 had K_i values of 1.0-12.5 μM .

Example 175

KDR Enzyme Assay

[0464] Compounds were screened for their ability to inhibit KDR using a standard coupled enzyme assay (Fox et al., Protein Sci., (1998) 7, 2249). Assays were carried out in a mixture of 200 mM HEPES 7.5, 10 mM MgCl_2 , 25 mM NaCl, 1 mM DTT and 1.5% DMSO. Final substrate concentrations in the assay were 300 μM ATP (Sigma Chemicals) and 10 μM poly E4Y (Sigma). Assays were carried out at 37°C and 30 nM KDR. Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 200 μM NADH, 30 $\mu\text{g}/\text{ML}$ pyruvate kinase and 10 $\mu\text{g}/\text{ml}$ lactate dehydrogenase.

[0465] An assay stock buffer solution was prepared containing all of the reagents listed above, with the exception of ATP and the test compound of interest. 177 μl of the stock solution was placed in a 96 well plate followed by addition of 3 μl of 2 mM DMSO stock containing the test compound (final compound concentration 30 μM). The plate was preincubated for about 10 minutes at 37°C and the reaction initiated by addition of 20 μl of ATP (final concentration 300 μM). Rates of reaction were obtained using a Molecular Devices plate reader (Sunnyvale, CA) over a 5 minute

- 242 -

read time at 37°C. Compounds showing greater than 50% inhibition versus standard wells containing the assay mixture and DMSO without test compound were titrated to determine IC50 values determined.

5 [0466] Compounds of the present invention were found to inhibit KDR. Compounds 48, 49, 58, 65, 66, 67, 68, 69, 70, 71, 77, 78, 79, 80, 81, 82, 83, 84, 86, 88, 89, 90, 93, 94, 96, 97, 98, 99, 133, 137, 138, 140, 149, 154, and 155 had Ki values of <0.2 μ M. Compounds 2,
10 35, 37, 38, 40, 42, 43, 44, 45, 46, 53, 55, 56, 57, 59, 61, 75, 76, 87, 91, 92, 95, 104, 105, 106, 109, 110, 112, 113, 114, 116, 117, 118, 119, 135, 136, 139, 142, 148, 150, and 153 had Ki values of 0.2-<1.0 μ M. Compounds 1, 31, 36, 39, 41, 47, 51, 62, 63, 64, 73,
15 74, 85, 86, 102, 111, 115, 132, 134, and 141 had Ki values of 1.0-12.5 μ M.

Example 176

Inhibition of FLT-3

[0467] Compounds were screened for their ability to
20 inhibit FLT-3 activity using a radiometric filter-binding assay. This assay monitors the ^{33} P incorporation into a substrate poly(Glu, Tyr) 4:1 (pE4Y). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM
25 NaCl, 1 mM DTT, 0.01% BSA and 2.5% DMSO. Final substrate concentrations in the assay were 90 μ M ATP and 0.5mg/mL pE4Y (both from Sigma Chemicals, St Louis, MO). The final concentration of compounds is generally between 0.01 and 5 μ M. Typically, a 12-point titration
30 was conducted by preparing serial dilutions from 10 mM DMSO stock of test compound. Reactions were carried out at room temperature.

- 243 -

[0468] Two assay solutions were prepared. Solution 1 contains 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 1 mg/ml pE4Y and 180 μ M ATP (containing 0.3 μ Ci of [γ -³³P]ATP for each reaction). Solution 2 contains 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 2 mM DTT, 0.02% BSA and 3 nM FLT-3. The assay was run on a 96 well plate by mixing 50 μ L each of Solution 1 and 2.5 mL of the test compounds. The reaction was initiated with Solution 2. After incubation for 20 minutes at room temperature, the reaction was stopped with 50 μ L of 20% TCA containing 0.4 mM of ATP. All of the reaction volume was then transferred to a filter plate and washed with 5% TCA by a Harvester 9600 from TOMTEC (Hamden, CT). The amount of ³³P incorporation into pE4y was analyzed by a Packard TopCount Microplate Scintillation Counter (Meriden, CT). The data was fitted using Prism software to get an IC₅₀ or K_i.

[0469] Compounds of the present invention were found to inhibit FLT. Compounds 38, 57, 59, 65, 68, 70, 71, 76, 77, 79, 82, 84, 86, 87, 90, 91, 92, 93, 94, 95, 98, 99, 105, 133, 134, 137, 138, 139, 140, 142, 149, 153, and 155 had K_i values of <0.2 μ M. Compounds 1, 43, 46, 47, 48, 49, 53, 58, 61, 62, 63, 64, 66, 69, 73, 75, 78, 81, 85, 96, 103, 106, 109, 110, 112, 115, 116, 117, 118, 119, 132, 136, 141, 143, and 154 had K_i values of 0.2-1.0 μ M. Compounds 2, 31, 34, 36, 39, 41, 42, 44, 45, 54, 55, 56, 60, 72, 74, 80, 83, 102, 104, 108, 111, 144, and 152 had K_i values of 1.0-12.5 μ M.

Example 177

Inhibition of FMS

[0470] Compounds are screened for their ability to inhibit FMS activity using a radiometric filter-binding

- 244 -

assay. This assay monitors the ^{33}P incorporation into a substrate poly(Glu, Tyr) 4:1 (pE4Y). Reactions are carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl_2 , 25 mM NaCl, 1 mM DTT, 0.01% BSA and 2.5% DMSO. Final substrate concentrations in the assay are 90 μM ATP and 0.5mg/mL pE4Y (both from Sigma Chemicals, St Louis, MO). The final concentration of compounds is generally between 0.01 and 5 μM . Typically, a 12-point titration is conducted by preparing serial dilutions from 10 mM DMSO stock of test compound. Reactions were carried out at room temperature.

[0471] Two assay solutions are prepared. Solution 1 contains 100 mM HEPES (pH 7.5), 10 mM MgCl_2 , 25 mM NaCl, 1 mg/ml pE4Y and 180 μM ATP (containing 0.3 μCi of $[\gamma\text{-}^{33}\text{P}]\text{ATP}$ for each reaction). Solution 2 contains 100 mM HEPES (pH 7.5), 10 mM MgCl_2 , 25 mM NaCl, 2 mM DTT, 0.02% BSA and 3 nM FMS. The assay is run on a 96 well plate by mixing 50 μL each of Solution1 and 2.5 mL of the test compounds. The reaction is initiated with Solution2. After incubation for 20 minutes at room temperature, the reaction is stopped with 50 μL of 20% TCA containing 0.4mM of ATP. All of the reaction volume is then transferred to a filter plate and washed with 5% TCA by a Harvester9600 from TOMTEC (Hamden, CT). The amount of ^{33}P incorporation into pE4y was analyzed by a Packard TopCount Microplate Scintillation Counter (Meriden, CT). The data was fitted using Prism software to get an IC_{50} or K_i .

- 245 -

Example 178Inhibition of c-KIT

[0472] Compounds are screened for their ability to inhibit c-KIT activity using a radiometric filter-binding assay. This assay monitors the ³³P incorporation into a substrate poly(Glu, Tyr) 4:1 (pE4Y). Reactions are carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 1 mM DTT, 0.01% BSA and 2.5% DMSO. Final substrate concentrations in the assay were 700 μM ATP and 0.5 mg/mL pE4Y (both from Sigma Chemicals, St Louis, MO). The final concentration of compounds is generally between 0.01 and 5 μM. Typically, a 12-point titration is conducted by preparing serial dilutions from 10 mM DMSO stock of test compound. Reactions were carried out at room temperature.

[0473] Two assay solutions are prepared. Solution 1 contains 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 1 mg/mL pE4Y and 1.4 mM ATP (containing 0.5 μCi of [γ-³³P]ATP for each reaction). Solution 2 contains 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 2 mM DTT, 0.02% BSA and 25 nM c-KIT. The assay is run on a 96 well plate by mixing 33 μL of Solution1 and 1.65 μL of the test compounds. The reaction is initiated with 33 μL of Solution2. After incubation for 20 minutes at room temperature, the reaction was stopped with 50 μL of 10% TCA containing 0.2 mM of ATP. All of the reaction volume is then transferred to a filter plate and washed with 5% TCA by a Harvester9600 from TOMTEC (Hamden, CT). The amount of ³³P incorporation into pE4y is analyzed by a Packard TopCount Microplate

- 246 -

Scintillation Counter (Meriden, CT). The data is fitted using Prism software to get an IC_{50} or K_i .

[0474] Compounds of the present invention were found to inhibit c-KIT. Compounds 1, 38, 43, 49, 53, 57, 59, 61, 65, 67, 68, 69, 70, 71, 72, 73, 76, 77, 78, 82, 83, 84, 85, 86, 87, 88, 90, 91, 92, 93, 94, 95, 96, 99, 105, 106, 109, 110, 114, 129, 130, 131, 135, 136, 137, 138, 139, 140, 141, 142, 149, 153, and 155 had K_i values of $<0.2 \mu M$. Compounds 40, 46, 48, 55, 60, 62, 108, 111, 112, 113, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, and 144 had K_i values of $0.2-1.0 \mu M$.

Example 179

Inhibition of AUR-2

[0475] Compounds are screened in the following manner for their ability to inhibit Aurora-2 using a standard coupled enzyme assay (Fox et al. (1998) *Protein Sci* 7, 2249). To an assay stock buffer solution containing 0.1M HEPES 7.5, 10 mM $MgCl_2$, 1 mM DTT, 25 mM NaCl, 2.5 mM phosphoenolpyruvate, 300 mM NADH, 30 mg/ml pyruvate kinase, 10 mg/ml lactate dehydrogenase, 40 mM ATP, and 800 μM peptide (LRRASLG, American Peptide, Sunnyvale, CA) is added a DMSO solution of a compound of the present invention to a final concentration of 30 μM . The resulting mixture is incubated at 30 °C for 10 min. The reaction was initiated by the addition of 10 μL of Aurora-2 stock solution to give a final concentration of 70 nM in the assay. The rates of reaction are obtained by monitoring absorbance at 340 nm over a 5 minute read time at 30 °C using a BioRad Ultramark plate reader (Hercules, CA). The K_i values are determined from the rate data as a function of inhibitor concentration.

- 247 -

[0476] Compounds of the present invention were found to inhibit AUR-2. Compounds 37, 44, 49, 57, 59, 65, 68, 70, 71, 76, 86, 87, 90, 92, 93, 94, 95, 96, 98, 99, and 153 had K_i values of $< 0.2 \mu\text{M}$. Compounds 1, 2, 38, 5 42, 43, 46, 47, 48, 58, 75, 82, 83, 91, 134, and 145 had K_i values of $0.2-1.0 \mu\text{M}$.

Example 180

TAK-1 Inhibition Assay

[0477] Compounds were screened for their ability to 10 inhibit TAK1A kinase activity using a radiometric filter binding assay. Reactions were carried out in a solution containing Buffer A (100 mM HEPES (pH 7.5), 10 mM MgCl_2), 25 mM NaCl, 2 mM DTT, and 1.5% DMSO. Final substrate concentrations in the assay were $50 \mu\text{M}$ ATP (a 15 mixture of unlabeled ATP (Sigma Chemicals, St Louis, MO) and ^{33}P -labeled ATP (PerkinElmer Life Sciences, Boston, MA) for a final specific activity of 50 Ci/mol), and $12 \mu\text{M}$ bovine myelin basic protein (MBP, Vertex Pharmaceuticals, Cambridge, MA). Reactions were 20 carried out at ambient temperature ($\sim 20^\circ\text{C}$) using 20 nM TAK1A-TAB fusion protein. Under these conditions the extent of reaction is linear with time for a period of 2 hours.

[0478] A test compound of the present invention (1 25 μL in DMSO) was combined with ATP and Buffer A in a final volume of $47 \mu\text{L}$ in a 96 well plate. Typically, a 6 point titration was conducted by preparing serial dilutions (from 10 mM compound stocks) with DMSO of the test compounds of the present invention in daughter 30 plates, for final concentrations spanning $0.046 \mu\text{M}$ to $3.73 \mu\text{M}$. The reaction was initiated by the addition of $20 \mu\text{l}$ of an enzyme stock solution consisting of

- 248 -

TAK1A-TAB fusion (described by Sugita, T. et al. in *Biochem. Biophys. Res. Comm.* 2002, 297, 1277-1281), MBP, Buffer A, NaCl, and DTT. The reaction was allowed to proceed for two hours at ambient temperature, then
5 quenched with an equal volume of 10 mM unlabeled ATP in 10% trichloroacetic acid. A 110 μ L aliquot of the quenched reaction was transferred to a Multiscreen PH filter plate (Millipore, Billerica, MA) and allowed to incubate at ambient temperature overnight (typically
10 16-20 hours). Following incubation the filter plates were washed with 3 $\times$ 150 μ L aliquots of 5% trichloroacetic acid using a modified Biotek Elx405 plate washer. A 70 μ L aliquot of Microscint 20 scintillation fluid (PerkinElmer) was added to each
15 well, and the plate was then sealed and read on a TopCount NXT microplate scintillation counter (PerkinElmer). The K_i values were determined from the rate data as a function of inhibitor concentration.

[0479] Compounds of the present invention were
20 found to inhibit TAK-1. Compounds 68, 70, 71, 110, 135, 136, 137, 138, 140, and 155 had K_i values of $< 0.2 \mu$ M. Compounds 48, 49, 53, 61, 69, 77, 78, 79, 84, 97, 98, 99, 106, 109, 115, 116, 117, 118, 133, 139, 141, and 142 had K_i values of $0.2-1.0 \mu$ M. Compounds 46, 62,
25 72, 111, 112, 113, 114, and 119 had K_i values of $1.0-12.5 \mu$ M.


[0480] While we have described a number of
embodiments of this invention, it is apparent that our
basic examples may be altered to provide other
30 embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the

- 249 -

appended claims rather than by the specific embodiments that have been represented by way of example.

What is Claimed is:

1. A compound of formula I:




or a pharmaceutically acceptable salt thereof, wherein:

W is CH or N;


Ring B is an optionally substituted 5-6 membered heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

R¹ is an optionally substituted 6-membered aryl ring having 0-3 nitrogens; and


Ring A is an optionally substituted ring selected from:



- 251 -



2. A compound of formula **I**:



or a pharmaceutically acceptable salt thereof, wherein:

W is CH or N, wherein the H is optionally replaced with (C₁-C₆)-alkyl or NH₂;

Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;


R¹ is:

an optionally substituted 6-10-membered aryl or 5-10-membered heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

a -(C₁-C₄ aliphatic) substituted with a -6-10-membered aryl or a 5-10-membered heteroaryl ring or a C₃-C₈ cycloaliphatic or heterocyclic ring, having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur each aliphatic and each ring being optionally substituted, or

an optionally substituted C₃-C₈ cycloaliphatic;

Ring A is an optionally substituted ring selected from:



each aryl or heteroaryl is optionally substituted with one or more R^3 groups, wherein R^3 is halogen; $-B(OH)_2$; $-R^\circ$; $-OR^\circ$; $-SR^\circ$; 1,2-methylenedioxy; 1,2-ethylenedioxy; $-CO_2(C_{1-4} \text{ aliphatic})$; an optionally substituted 5-6 membered heterocyclic ring; phenyl optionally substituted with R° ; $-O(\text{phenyl})$ optionally substituted with R° ; $-(CH_2)_{1-2}(\text{phenyl})$, optionally substituted with R° ; $-CH=CH(\text{phenyl})$, optionally substituted with R° ; $-NO_2$; $-CN$; $-NHR^\circ$; $-N(R^\circ)_2$; $-NR^\circ C(O)R^\circ$; $-NR^\circ C(S)R^\circ$; $-NR^\circ C(O)N(R^\circ)_2$; $-NR^\circ C(S)N(R^\circ)_2$; $-NR^\circ CO_2R^\circ$; $-NR^\circ NR^\circ C(O)R^\circ$; $-NR^\circ NR^\circ C(O)N(R^\circ)_2$; $-NR^\circ NR^\circ CO_2R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-CO_2R^\circ$;

- 253 -

$-C(O)R^{\circ}$; $-C(S)R^{\circ}$; $-C(O)N(R^{\circ})_2$; $-C(S)N(R^{\circ})_2$;
 $-OC(O)N(R^{\circ})_2$; $-OC(O)R^{\circ}$; $-C(O)N(OR^{\circ})R^{\circ}$; $-C(NOR^{\circ})R^{\circ}$;
 $-S(O)_2R^{\circ}$; $-S(O)_3R^{\circ}$; $-SO_2N(R^{\circ})_2$; $-S(O)R^{\circ}$; $-NR^{\circ}SO_2N(R^{\circ})_2$;
 $-NR^{\circ}SO_2R^{\circ}$; $-N(OR^{\circ})R^{\circ}$; $-C(=NH)-N(R^{\circ})_2$; or
 $-(CH_2)_{0-2}NHC(O)R^{\circ}$;

wherein each independent occurrence of R° is selected from hydrogen, C_{1-6} aliphatic, a 5-10 membered heteroaryl or heterocyclic ring, phenyl, $-O(phenyl)$, $-CH_2(phenyl)$, 5 membered heterocyclic ring;

wherein each group of R° is optionally substituted with J , wherein J is aryl, phenyl, heteroaryl, NH_2 , $NH(C_{1-4} \text{ aliphatic})$, $N(C_{1-4} \text{ aliphatic})_2$, $NH(CH_2)phenyl$, halogen, $-NH SO_2(C_{1-4} \text{ aliphatic})$, $-NH CO_2(C_{1-4} \text{ aliphatic})$, $C_{1-4} \text{ aliphatic}$, OH , $O(C_{1-4} \text{ aliphatic})$, NO_2 , CN , CO_2H , $-CO(5-6 \text{ membered heterocyclic ring})$, 5-6 membered heterocyclic ring, $-CO_2(C_{1-4} \text{ aliphatic})$, $-O(haloC_{1-4} \text{ aliphatic})$, or $halo(C_{1-4} \text{ aliphatic})$, or

wherein each group of J is optionally substituted with J' , wherein J' is NH_2 , $NH(C_{1-4} \text{ aliphatic})$, $N(C_{1-4} \text{ aliphatic})_2$, $NH(CH_2)phenyl$, halogen, $-NH SO_2(C_{1-4} \text{ aliphatic})$, $-NH CO_2(C_{1-4} \text{ aliphatic})$, $C_{1-4} \text{ aliphatic}$, OH , $O(C_{1-4} \text{ aliphatic})$, NO_2 , CN , CO_2H , $-CO(5-6 \text{ membered heterocyclic ring})$, 5-6 membered heterocyclic ring, $-CO_2(C_{1-4} \text{ aliphatic})$, $-O(haloC_{1-4} \text{ aliphatic})$, or $halo(C_{1-4} \text{ aliphatic})$, wherein each the J' groups is optionally substituted with $C_{1-4} \text{ aliphatic}$, halogen,

- 254 -

wherein each of the C₁₋₄ aliphatic groups of J' is unsubstituted;

two R° are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

wherein each aliphatic or heteroaliphatic group or non-aromatic heterocyclic ring is optionally substituted with R³, =O, =S, =NNHR*, =NN(R*)₂, =NNHC(O)R*, =NNHCO₂(alkyl), =NNHSO₂(alkyl), or =NR*,

wherein each R* is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic, wherein optional substituents on the aliphatic group of R* are selected from 5-6 membered heterocyclic ring, heteroaryl, aryl, NH₂, NHSO₂R*, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), CO(5-6 membered heterocyclic ring), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄aliphatic groups of R* is unsubstituted; and


wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with -(C₁₋₆ aliphatic)₂, -R⁺, -N(R⁺)₂, -C(O)R⁺, -CO₂R⁺, -C(O)C(O)R⁺, -C(O)CH₂C(O)R⁺, -SO₂R⁺, -SO₂N(R⁺)₂, -C(=S)N(R⁺)₂, -C(=NH)-N(R⁺)₂, or -NR⁺SO₂R⁺;

- 255 -

wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, optionally substituted phenyl, optionally substituted $-O(\text{phenyl})$, optionally substituted $-\text{CH}_2(\text{phenyl})$, optionally substituted $-(\text{CH}_2)_{1-2}(\text{phenyl})$; optionally substituted $-\text{CH}=\text{CH}(\text{phenyl})$; or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, wherein optional substituents on the aliphatic group or the phenyl ring of R^+ are selected from NH_2 , $\text{NH}(C_{1-4} \text{ aliphatic})$, $\text{N}(C_{1-4} \text{ aliphatic})_2$, halogen, $C_{1-4} \text{ aliphatic}$, OH , $\text{O}(C_{1-4} \text{ aliphatic})$, NO_2 , CN , CO_2H , $\text{CO}_2(C_{1-4} \text{ aliphatic})$, $\text{O}(\text{halo } C_{1-4} \text{ aliphatic})$, or $\text{halo}(C_{1-4} \text{ aliphatic})$, wherein each of the foregoing C_{1-4} aliphatic groups of R^+ is unsubstituted, or

two R^+ are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

3. A compound of formula **I**:



- 256 -

I

or a pharmaceutically acceptable salt thereof, wherein:

W is CH, CNH₂ or N;

Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,


wherein Ring B is optionally substituted with one or more oxo, halogen, -OH, -OR^o, -NHR^o, N(R^o)₂, a 5-6 membered heterocyclic ring, -COO(C₁₋₄ aliphatic), -B(OH)₂, -CO(5-6 membered heterocyclic ring), aryl, heteroaryl, and R^o,

wherein each R^o is H or C₁₋₆ aliphatic independently optionally substituted with phenyl, NH₂, NH(C₁₋₄ aliphatic), NH(CH₂)phenyl, N(C₁₋₄ aliphatic)₂, heteroaryl, -NHSO₂(C₁₋₄ aliphatic), halogen, an optionally substituted -CO(5-6 membered heterocyclic ring), an optionally substituted 5-6 membered heterocyclic ring, COO aliphatic and OH;


R¹ is an optionally substituted 6-membered aryl or heteroaryl ring having 0-3 nitrogens, (C₁₋₄ aliphatic)-aryl ring, C₁₋₆ aliphatic,

wherein R¹ is optionally substituted with one or more halogen or -OR^o, wherein each R^o is C₁₋₄ aliphatic;

Ring A is an optionally substituted ring selected from:



- 257 -



wherein Ring A is optionally substituted with oxo, -OH, NH_2 , or $-\text{CH}_3$; and

each aryl or heteroaryl is optionally substituted with one or more halogen; $-\text{R}^\circ$; $-\text{OR}^\circ$; $-\text{SR}^\circ$; 1,2-methylenedioxy; 1,2-ethylenedioxy; phenyl optionally substituted with R° ; $-\text{O}(\text{phenyl})$ optionally substituted with R° ; $-(\text{CH}_2)_{1-2}(\text{phenyl})$, optionally substituted with R° ; $-\text{CH}=\text{CH}(\text{phenyl})$, optionally substituted with R° ; $-\text{NO}_2$; $-\text{CN}$; $-\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{C}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{C}(\text{S})\text{R}^\circ$; $-\text{NR}^\circ\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{C}(\text{S})\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{CO}_2\text{R}^\circ$; $-\text{NR}^\circ\text{NR}^\circ\text{C}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{NR}^\circ\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{NR}^\circ\text{CO}_2\text{R}^\circ$; $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^\circ$; $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^\circ$; $-\text{CO}_2\text{R}^\circ$; $-\text{C}(\text{O})\text{R}^\circ$; $-\text{C}(\text{S})\text{R}^\circ$; $-\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{C}(\text{S})\text{N}(\text{R}^\circ)_2$; $-\text{OC}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{OC}(\text{O})\text{R}^\circ$; $-\text{C}(\text{O})\text{N}(\text{OR}^\circ)\text{R}^\circ$; $-\text{C}(\text{NOR}^\circ)\text{R}^\circ$; $-\text{S}(\text{O})_2\text{R}^\circ$; $-\text{S}(\text{O})_3\text{R}^\circ$; $-\text{SO}_2\text{N}(\text{R}^\circ)_2$; $-\text{S}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{SO}_2\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{SO}_2\text{R}^\circ$; $-\text{N}(\text{OR}^\circ)\text{R}^\circ$; $-\text{C}(=\text{NH})-\text{N}(\text{R}^\circ)_2$; or $-(\text{CH}_2)_0-2\text{NHC}(\text{O})\text{R}^\circ$;

- 258 -

wherein each independent occurrence of R° is selected from hydrogen, optionally substituted C_{1-6} aliphatic, an optionally substituted 5-9 membered heteroaryl or heterocyclic ring, phenyl, -O(phenyl), -CH₂(phenyl), an optionally substituted -O(5-6 membered heterocyclic ring), or an optionally substituted -CH₂(5-6 membered heterocyclic ring);

wherein aliphatic groups of R° are optionally substituted with NH₂, NH(C_{1-4} aliphatic), N(C_{1-4} aliphatic)₂, halogen, C_{1-4} aliphatic, OH, O(C_{1-4} aliphatic), NO₂, CN, CO₂H, CO₂(C_{1-4} aliphatic), O(halo C_{1-4} aliphatic), or halo C_{1-4} aliphatic, wherein each of the foregoing C_{1-4} aliphatic groups of R° is unsubstituted, or

two R° are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

wherein each aliphatic or heteroaliphatic group or non-aromatic heterocyclic ring is optionally substituted with =O, =S, =NNHR*, =NN(R*)₂, =NNHC(O)R*, =NNHCO₂(alkyl), =NNHSO₂(alkyl), heterocyclic ring, -OH, -CH₂OH, NHR*, N(R*)₂, CO (heterocyclic ring), R*, NHSO₂R* or =NR*,

wherein each R* is independently selected from hydrogen or an optionally substituted C_{1-6} aliphatic, wherein optional substituents on the aliphatic group of R* are selected from 5-6 membered

- 259 -

heterocyclic ring, heteroaryl, aryl, NH_2 , NHSO_2R^* , $\text{NH}(\text{C}_{1-4} \text{ aliphatic})$, $\text{N}(\text{C}_{1-4} \text{ aliphatic})_2$, halogen, C_{1-4} aliphatic, OH , $\text{O}(\text{C}_{1-4} \text{ aliphatic})$, $\text{CO}(\text{5-6 membered heterocyclic ring})$, NO_2 , CN , CO_2H , $\text{CO}_2(\text{C}_{1-4} \text{ aliphatic})$, $\text{O}(\text{halo C}_{1-4} \text{ aliphatic})$, or $\text{halo}(\text{C}_{1-4} \text{ aliphatic})$, wherein each of the foregoing C_{1-4} aliphatic groups of R^* is unsubstituted; and

wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with $-(\text{C}_{1-6} \text{ aliphatic})_2$, $-\text{R}^+$, $-\text{N}(\text{R}^+)_2$, $-\text{C}(\text{O})\text{R}^+$, $-\text{CO}_2\text{R}^+$, $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^+$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^+$, $-\text{SO}_2\text{R}^+$, $-\text{SO}_2\text{N}(\text{R}^+)_2$, $-\text{C}(=\text{S})\text{N}(\text{R}^+)_2$, $-\text{C}(=\text{NH})-\text{N}(\text{R}^+)_2$, or $-\text{NR}^+\text{SO}_2\text{R}^+$;


wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, optionally substituted phenyl, optionally substituted $-\text{O}(\text{phenyl})$, optionally substituted $-\text{CH}_2(\text{phenyl})$, optionally substituted $-(\text{CH}_2)_{1-2}(\text{phenyl})$; optionally substituted $-\text{CH}=\text{CH}(\text{phenyl})$; or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, wherein optional substituents on the aliphatic group or the phenyl ring of R^+ are selected from NH_2 , $\text{NH}(\text{C}_{1-4} \text{ aliphatic})$, $\text{N}(\text{C}_{1-4} \text{ aliphatic})_2$, halogen, C_{1-4} aliphatic, OH , $\text{O}(\text{C}_{1-4} \text{ aliphatic})$, NO_2 , CN , CO_2H , $\text{CO}_2(\text{C}_{1-4} \text{ aliphatic})$, $\text{O}(\text{halo C}_{1-4} \text{ aliphatic})$, or $\text{halo}(\text{C}_{1-4} \text{ aliphatic})$, wherein each

- 260 -

of the foregoing C₁₋₄aliphatic groups of R⁺ is unsubstituted, or

two R⁺ are taken together with the atom(s) to which each is bound to form a 5-8-membered heterocyclyl, aryl, or heteroaryl ring or a 3-8-membered cycloalkyl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

4. A compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

W is CH, CNH₂ or N;

Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

wherein Ring B is optionally substituted with one or more oxo, halogen, -OH, -OR^o, -NHR^o, N(R^o)₂, a 5-6 membered heterocyclic ring, -COO(C₁₋₄ aliphatic), -B(OH)₂, -CO(5-6 membered heterocyclic ring), aryl, heteroaryl, and R^o,

wherein each R^o is H or C₁₋₆ aliphatic independently optionally substituted with phenyl, NH₂, NH(CH₂)phenyl, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, heteroaryl, COO aliphatic, -NHSO₂(C₁₋₄ aliphatic),


- 261 -

halogen, an optionally substituted -CO(5-6 membered heterocyclic ring), an optionally substituted 5-6 membered heterocyclic ring, and OH;

R^1 is an optionally substituted 6-membered aryl or heteroaryl ring having 0-3 nitrogens,

wherein R^1 is optionally substituted with one or more halogen or $-OR^o$, wherein each R^o is C_{1-4} aliphatic;

Ring A is an optionally substituted ring selected from:



wherein Ring A is optionally substituted with oxo, -OH, NH_2 , or $-CH_3$; and

- 262 -

each aryl or heteroaryl is optionally substituted with one or more halogen; $-R^{\circ}$; $-OR^{\circ}$; $-CN$; NO_2 ; $-N(R^{\circ})_2$; $-NR^{\circ}C(O)R^{\circ}$; $-C(O)N(R^{\circ})_2$; $-S(O)_2R^{\circ}$; or $-NR^{\circ}SO_2R^{\circ}$;

wherein each independent occurrence of R° is selected from hydrogen, optionally substituted C_{1-6} aliphatic, an optionally substituted 6-9 membered heteroaryl or heterocyclic ring, or an optionally substituted $-CH_2$ (5-6 membered heterocyclic ring);

wherein aliphatic groups of R° are optionally substituted with N (unsubstituted C_{1-4} aliphatic) $_2$;

wherein each aliphatic or heteroaliphatic group or non-aromatic heterocyclic ring is optionally substituted with $=O$, heterocyclic ring, NHR^* , $N(R^*)_2$, $NHSO_2R^*$, CO (heterocyclic ring), CH_2OH , OH and $-R^*$, wherein each R^* is H or optionally substituted C_{1-6} aliphatic;


wherein the optional substituents of the aliphatic group of R^* are selected from 5-6 membered heterocyclic ring and aryl; and

wherein each nitrogen of a non-aromatic heterocyclic ring is optionally substituted with $(C_{1-6}$ aliphatic) $_2$, $-R^+$, $-C(O)R^+$;

wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, wherein optional substituents on the aliphatic group of R^+ are CN.

5. A compound of formula I:

- 263 -



or a pharmaceutically acceptable salt thereof, wherein:

W is CH, CNH₂ or N;

Ring B is an optionally substituted 5- or 6-membered aryl, heteroaryl or heterocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,


wherein Ring B is optionally substituted with one or more oxo; chloro; bromo; fluoro; -CH₂OH; -OH; -OCH₃; CH₃; -NHCH₃; -NHCH₂CH₃; -N(CH₃)₂; -NH-CH₂-tetrahydrofuranyl; pyrrolidinyl; piperidinyl; pyrazolyl; -COO(CH₃); -B(OH)₂; phenyl; benzyl; pyrindinyl; pyrimidinyl; imidazolyl; H; cyclopropyl; cyclohexyl; cyclohexenyl; -CH₂CH₃; -CH₂N(CH₃)₂; propynyl substituted with N(CH₃)₂; ethenyl; ethenyl substituted with triazolyl; -CH₂CH₂-triazolyl; NH(CH₃); NH(CH₂)phenyl; N(CH₃)₂; imidazo-1,2-e-pyridinyl optionally substituted with -SO₂(CH₃), -NHSO₂(CH₃); an optionally substituted -CO(piperazinyl) or -CO(pyrrolidinyl), an optionally substituted morpholinyl or triazolyl, or OH;

R¹ is an optionally substituted 6-membered aryl or heteroaryl ring having 0-3 nitrogens,

wherein R¹ is optionally substituted with one or more halogen or -OR^o, wherein each R^o is C₁₋₄ aliphatic;

- 264 -

Ring A is an optionally substituted ring selected from:



wherein Ring A is optionally substituted with oxo, -OH, NH₂, or -CH₃; and

each aryl or heteroaryl is optionally substituted with one or more fluoro; chloro ; -R^o; -OR^o; -CN; NO₂; -N(CH₃)₂; -NHCH₂CH₂N(CH₃)₂, -NH₂, -NHC(O)CH₃; -C(O)NH₂; -C(O)N(CH₃)₂ -S(O)₂CH₃; or -NHSO₂CH₃;

wherein each independent occurrence of R^o is selected from hydrogen, CH₃, -CH₂(N(CH₃)₂), optionally substituted pyridinyl, piperindinyl, diazepanyl, morpholinyl, optionally substituted 3,9-diaza-bicyclo [4.2.1] nonane, piperazinyl or

- 265 -

pyrrolidinyl, or an optionally substituted
-CH₂(morpholinyl) or -CH₂(piperazinyl);

wherein each aliphatic or
heteroaliphatic group or non-aromatic heterocyclic ring
is optionally substituted with =O, pyrrolidinyl, OH,
NHbenzyl, NH₂, and -CO(piperazinyl); and

wherein each nitrogen of a non-aromatic
heterocyclic ring is optionally substituted with -CH₃,
-(CH₃)₂, -H, -CH₂CH₃, -C(O)CH₂CN, -C(O)CH₃.

6. The compound according to claim 2,
wherein Ring B is an optionally substituted 5-membered
heteroaryl ring having one nitrogen and 0-2 additional
heteroatoms independently selected from nitrogen,
oxygen, or sulfur.

7. The compound according to claim 2,
wherein Ring B is an optionally substituted 6-membered
heteroaryl ring having 1-3 nitrogens.

8. The compound according to claim 7,
wherein Ring B is an optionally substituted pyrido
ring.

9. The compound according to claim 7,
wherein Ring B is an optionally substituted pyrimido
ring.

- 266 -

10. The compound according to claim 7, wherein Ring B is an optionally substituted pyrazino ring.

11. The compound according to claim 7, wherein Ring B is an optionally substituted pyridazo ring.

12. The compound according to claim 2, wherein Ring B is an optionally substituted phenyl ring.





13. The compound according to claim 2, wherein R¹ is an optionally substituted phenyl ring.

14. The compound according to claim 2, wherein R¹ is an optionally substituted pyridyl or pyrimidinyl ring.


15. The compound according to claim 2, wherein Ring A is an optionally substituted ring selected from isoxazolyl, imidazolyl, triazolyl, or tetrazolyl.

16. The compound according to claim 15, wherein Ring A is selected from the following rings:


- 267 -

**a****b****d****o**


or

**p.**


17. The compound according to claim 16,
wherein Ring A is selected from:

**o**

or





**p.**


18. The compound according to claim 15,
wherein Ring A is:

**b.**


19. The compound according to claim 2,
wherein W is CH.

20. A compound selected from the group
consisting of:





- 269 -



21. A compound selected from the group consisting of nos. 33-155.

22. A composition comprising an effective amount of a compound according to claim 2, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

23. The composition according to claim 22, wherein said compound is in an amount sufficient to detectably inhibit c-Met, GSK3, JAK, SYK, or KDR protein kinase activity.

24. The composition according to claim 22, additionally comprising a therapeutic agent selected from a chemotherapeutic or anti-proliferative agent, an anti-inflammatory agent, an immunomodulatory or immunosuppressive agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating destructive bone disorders, an agent for treating liver disease, an anti-viral agent, an agent for treating blood disorders, an agent for treating

- 270 -

diabetes, or an agent for treating immunodeficiency disorders.

25. A method of inhibiting c-Met, GSK3, JAK, SYK, KDR, FLT-3, c-Kit, Aurora, or TAK-1 kinase activity in:

- (a) a patient; or
- (b) a biological sample;

which method comprises administering to said patient, or contacting said biological sample with:

- a) a composition according to claim 22;

or

- b) a compound according to claim 2.

26. A method of treating or lessening the severity of a cancer or a proliferative disorder in a patient in need thereof, comprising the step of administering to said patient:

- a) a composition according to claim 22;

or

- b) a compound according to claim 2.

27. The method according to claim 26, comprising the additional step of administering to said patient an additional therapeutic agent selected from a chemotherapeutic or anti-proliferative agent, wherein said additional therapeutic agent is administered together with said composition as a single dosage form

- 271 -

or separately from said composition as part of a multiple dosage form.

28. A method of treating or lessening the severity of renal cancer in a patient in need thereof, comprising administering to said patient a composition according to claim 22.

29. A method of treating or lessening the severity of a disease or condition selected from glioblastoma, a gastric carcinoma, colon cancer, breast cancer, prostate cancer, liver cancer, pancreatic cancer, or lung cancer in a patient in need thereof, comprising administering to said patient a composition according to claim 22.

30. The method according to claim 29, wherein said disease or condition is glioblastoma, breast cancer, colon cancer, or liver cancer.

31. A method of inhibiting tumor metastasis in a patient, comprising administering to said patient a composition according to claim 22.

32. A method of treating asthma in a patient in need thereof, comprising administering to said patient a composition according to claim 22.

- 272 -

33. A method of treating or lessening the severity of a disease or condition selected from allergic disorders, proliferative disorders, autoimmune disorders, conditions associated with organ transplant, inflammatory disorders, immunologically mediated disorders, viral diseases, or destructive bone disorders, comprising the step of administering to said patient:

a composition according to claim 22; or
a compound according to claim 2.

34. The method of claim 33, further comprising the step of administering to said patient an additional therapeutic agent selected from a chemotherapeutic or anti-proliferative agent, a treatment for Alzheimer's Disease, a treatment for Parkinson's Disease, an agent for treating Multiple Sclerosis (MS), a treatment for asthma, an agent for treating schizophrenia, an anti-inflammatory agent, an immunomodulatory or immunosuppressive agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating destructive bone disorders, an agent for treating liver disease, an agent for treating a blood disorder, and an agent for treating an immunodeficiency disorder, wherein:

said additional therapeutic agent is appropriate for the disease being treated; and

said additional therapeutic agent is administered together with said composition as a single

- 273 -

dosage form or separately from said composition as part of a multiple dosage form.

35. The method of claim 33, wherein the disease or condition is a cancer.

36. The method of claim 35, wherein the cancer is selected from the group consisting of breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity and pharynx (oral), lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia.

37. The method of claim 33, wherein the disease or condition is selected from autoimmune diseases, inflammatory diseases, metabolic, neurological and neurodegenerative diseases,

- 274 -

cardiovascular diseases, allergy, asthma, diabetes, Alzheimer's disease, Huntington's disease, Parkinson's disease, AIDS-associated dementia, amyotrophic lateral sclerosis (ALS, Lou Gehrig's disease), multiple sclerosis (MS), schizophrenia, cardiomyocyte hypertrophy, reperfusion/ischemia, and baldness.

38. The method of claim 33, wherein the disease or condition is selected from cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune disease.

39. The method of claim 33, wherein the disease or condition is selected from hypercalcemia, osteoporosis, osteoarthritis, cancer, bone metastasis, and Paget's disease.

40. The method of claim 33, wherein the disease or condition is selected from immune responses, allergic or type I hypersensitivity reactions, and asthma, autoimmune diseases, transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, multiple sclerosis, neurodegenerative disorders, Familial amyotrophic

- 275 -

lateral sclerosis (FALS), solid and hematologic malignancies, leukemias and lymphomas.

41. The method of claim 33, wherein the disease or condition is selected from hematopoietic disorders, acute-myelogenous leukemia (AML), acute-promyelocytic leukemia (APL), and acute lymphocytic leukemia (ALL).

42. The method of claim 33, wherein the disease or condition is an allergic disorder.

43. The method of claim 42, wherein the disease or condition is asthma.