




(86) **Date de dépôt PCT/PCT Filing Date:** 2014/01/07
(87) **Date publication PCT/PCT Publication Date:** 2014/07/17
(85) **Entrée phase nationale/National Entry:** 2015/06/25
(86) **N° demande PCT/PCT Application No.:** US 2014/010501
(87) **N° publication PCT/PCT Publication No.:** 2014/110036
(30) **Priorité/Priority:** 2013/01/09 (US61/750,695)

(51) **Cl.Int./Int.Cl.** *C07D 213/82* (2006.01),
A61K 31/05 (2006.01), *A61K 31/09* (2006.01),
C07C 233/76 (2006.01), *C07C 39/08* (2006.01),
C07C 43/205 (2006.01), *C07C 49/83* (2006.01),
C07C 61/39 (2006.01), *C07D 213/38* (2006.01),
C07D 213/89 (2006.01), *C07D 233/64* (2006.01),
C07D 401/12 (2006.01)

(71) **Demandeur/Applicant:**
AQUINOX PHARMACEUTICALS (CANADA) INC., CA

(72) **Inventeurs/Inventors:**
MACKENZIE, LLOYD F., CA;
MACRURY, THOMAS B., US;
...

(54) **Titre : MODULATEURS DE SHIP1 ET PROCEDES ASSOCIES**
(54) **Title: SHIP1 MODULATORS AND METHODS RELATED THERETO**



(57) **Abrégé/Abstract:**

Compounds of formula (I): [Formula should be inserted here]; where [Formula should be inserted here], n, R¹, R^{4a}, R^{4b}, R⁵, R⁷ and R⁸ are defined herein, or pharmaceutically acceptable salts thereof, are described herein. The disclosed compounds have activity

(72) **Inventeurs(suite)/Inventors(continued):** HARWIG, CURTIS, CA; BOGUCKI, DAVID, CA; RAYMOND, JEFFERY R., CA; PETTIGREW, JEREMY, D., CA; KHLEBNIKOV, VLADIMIR, CA; SHAN, RUDONG, CA

(74) **Agent:** TEES, SUSAN M.

(57) **Abrégé(suite)/Abstract(continued):**

as SHIP1 modulators, and thus may be used to treat any of a variety of diseases, disorders or conditions that would benefit from SHIP1 modulation. Compositions comprising a compound of formula (I) in combination with a pharmaceutically acceptable carrier or diluent are also disclosed, as are methods of SHIP1 modulation by administration of such compounds to an animal in need thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

WIPO | PCT



(10) International Publication Number

WO 2014/110036 A4

(43) International Publication Date
17 July 2014 (17.07.2014)

(51) International Patent Classification:

C07D 213/82 (2006.01) **C07D 233/64** (2006.01)
A61K 31/05 (2006.01) **C07D 401/12** (2006.01)
A61K 31/09 (2006.01) **C07C 233/76** (2006.01)
C07D 213/89 (2006.01) **C07C 49/83** (2006.01)
C07C 39/08 (2006.01) **C07D 213/38** (2006.01)
C07C 43/205 (2006.01) **C07C 61/39** (2006.01)

(21) International Application Number:

PCT/US2014/010501

(22) International Filing Date:

7 January 2014 (07.01.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/750,695 9 January 2013 (09.01.2013) US

(71) Applicant: AQUINOX PHARMACEUTICALS INC.

[CA/CA]; 450 - 887 Great Northern Way, Vancouver, British Columbia V5T 4T5 (CA).

(72) Inventors: MACKENZIE, Lyoyd, F.; 242 24th Street

West, North Vancouver, British Columbia V7M 2C4 (CA).

MACRURY, Thomas, B.; 1636 Edwards Drive, Point

Roberts, Washington 98281 (US). HARWIG, Curtis; 515

East 15th Avenue #403, Vancouver, British Columbia V5T

4S4 (CA). **BOGUCKI, David**; 14957 59th Avenue, Surrey, British Columbia V3S 4S6 (CA). **RAYMOND, Jeffery, R.**; #19 - 2658 Morningstar Crescent, Vancouver, British Columbia V5S 4P4 (CA). **PETTIGREW, Jeremy, D.**; 7328 Arcola Street #1407, Burnaby, British Columbia V5E 0A7 (CA). **KHLEBNIKOV, Vladimir**; 3803 - 111A Street, Edmonton, Alberta Y6J 1G2 (CA). **SHAN, Rudong**; 331 Magrath Blvd, N.W., Edmonton, Alberta T6R 0M7 (CA).

(74) Agents: ROTH, Carol, J. et al.; Seed Intellectual Property


Law Group PLLC, Suite 5400, 701 Fifth Avenue, Seattle, Washington 98104-7064 (US).

(81) Designated States (unless otherwise indicated, for every


kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]

(54) Title: SHIP1 MODULATORS AND METHODS RELATED THERETO



(57) **Abstract:** Compounds of formula (I): [Formula should be inserted here]; where [Formula should be inserted here], n, R¹, R^{4a}, R^{4b}, R⁵, R⁷ and R⁸ are defined herein, or pharmaceutically acceptable salts thereof, are described herein. The disclosed compounds have activity as SHIP1 modulators, and thus may be used to treat any of a variety of diseases, disorders or conditions that would benefit from SHIP1 modulation. Compositions comprising a compound of formula (I) in combination with a pharmaceutically acceptable carrier or diluent are also disclosed, as are methods of SHIP1 modulation by administration of such compounds to an animal in need thereof.



WO 2014/110036 A4

WO 2014/110036 A4

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

Published:

— *with international search report (Art. 21(3))*

— *with amended claims (Art. 19(1))*

Date of publication of the amended claims: 2 October 2014

SHIP1 MODULATORS AND METHODS RELATED THERETO

PRIORITY APPLICATION

This application claims the priority benefit of U.S. Provisional Patent Application No. 61/750,695, filed on January 9, 2013, the relevant disclosure of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention is generally directed to SHIP1 modulators, as well as to compositions and methods related to the same.

BACKGROUND OF THE INVENTION

In response to extracellular signals, phosphoinositide 3-kinase (PI3K) becomes activated and phosphorylates phosphatidylinositol-4,5-bisphosphate (PI-4,5-P₂) within the plasma membrane to generate phosphatidylinositol-3,4,5-trisphosphate (PIP₃). PIP₃ then initiates a cascade of downstream signaling pathways by interacting with pleckstrin homology (PH) domain-containing proteins, such as protein kinase B (PKB, also known as Akt), that regulate cellular activation, function, proliferation and/or survival, depending on the cell type and stimulus (Deane *et al.*, *Annu Rev Immunol* **22**, 563-598, 2004). Cellular levels of PIP₃ are normally tightly regulated by PI3K, the 5' inositol phosphatases SHIP1 (SH2 domain-containing inositol phosphatase), SHIP2, and by the 3' inositol phosphatase PTEN. SHIP1 and SHIP2 dephosphorylate PIP₃ to phosphatidylinositol-3,4-bisphosphate (PI-3,4-P₂), whereas PTEN dephosphorylates PIP₃ to PI-4,5-P₂ (Sly *et al.*, *Exp Hematol* **31**, 1170-1181, 2003; Vivanco *et al.*, *Nat Rev Cancer* **2**, 489-501, 2002). Of these three, SHIP1 is unique in that its expression is restricted primarily to immune and hematopoietic cells (Sly *et al.*, *Exp Hematol* **31**, 1170-1181, 2003; Damen *et al.*, *Proc Natl Acad Sci U S A* **93**, 1689-1693, 1996).

SHIP1's role in immune cell homeostasis is shown both by the myeloproliferative syndrome observed in SHIP1^{-/-} mice, as well as the hypersensitivity of SHIP1^{-/-} mice and cells to immune stimulation (Helgason *et al.*, *Genes Dev* **12**, 1610-1620, 1998; Sly *et al.*, *Immunity* **21**, 227-239, 2004). SHIP1 has been shown to mediate signaling from the inhibitory FcγRIIB receptor (Coggeshall *et al.*, *Mol Immunol* **39**, 521-529, 2002), and is important in terminating signal transduction from activating immune/hematopoietic cell receptor systems (Kalesnikoff *et al.*, *Rev Physiol Biochem Pharmacol* **149**, 87-103, 2003).

Diminished SHIP1 activity or expression has been observed in human inflammatory diseases (Vonakis *et al.*, *J Allergy Clin Immunol* **108**, 822-831, 2001) and hematopoietic malignancies (Liang *et al.*, *Proteomics* **6**, 4554-4564, 2006; Fukuda *et al.*, *Proc Natl Acad Sci USA* **102**, 15213-15218, 2005; Luo *et al.*, *Zhongguo Shi Yan Xue Ye Xue Za Zhi* **12**, 420-426, 2004; Vanderwinden *et al.*, *Cell Signal* **18**, 661-669, 2006; Ong, C.J. *et al.*, *Blood* (2007), Vol. 110, No. 6, pp. 1942-1949).

Because dysregulated activation of the PI3K pathway contributes to inflammatory/immune disorders and cancer, intense efforts have been invested into the development of inhibitors of PI3K itself, as well as downstream protein kinases (Workman *et al.*, *Nat Biotechnol* **24**, 794-796, 2006; Simon, *Cell* **125**, 647-649, 2006; Hennessy *et al.*, *Nat Rev Drug Discov* **4**, 988-1004, 2005; Knight *et al.*, *Cell* **125**, 733-747, 2006; Ong, C.J. *et al.*, *Blood* (2007), Vol. 110, No. 6, pp. 1942-1949). The precedent for discovery and biologic efficacy of kinase inhibitors is well established, and a number of promising new PI3K isoform-specific inhibitors have recently been developed and used in mouse models of inflammatory disease (Camps *et al.*, *Nat Med* **11**, 936-943, 2005; Barber *et al.*, *Nat Med* **11**, 933-935, 2005) and glioma (Fan *et al.*, *Cancer Cell* **9**, 341-349, 2006) with minimal toxicities. However, because of the dynamic interplay between phosphatases and kinases in regulating biologic processes, inositol phosphatase activators represent a complementary, alternative approach to reduce cellular PIP₃ levels. Of the phosphoinositol phosphatases that degrade PIP₃, SHIP1 is a particularly ideal target for development of therapeutics for treating immune and hemopoietic disorders because its hematopoietic-restricted expression (Hazen AL, *et al.* **113**, 2924-33, 2009; Rohrschneider LR, Fuller JF, Wolf I, Liu Y, Lucas DM. Structure, function, and biology of SHIP proteins. *Genes Dev.* **14**:505-20, 2000) would limit the effects of a specific SHIP1 agonist to target cells.

To date, a number of small molecule SHIP1 modulators have been disclosed, including sesquiterpene compounds such as pelorol. Pelorol is a natural product isolated from the topical marine sponge *Dactylospongia elegans* (Kwak *et al.*, *J Nat Prod* **63**, 1153-1156, 2000; Goclik *et al.*, *J Nat Prod* **63**, 1150-1152, 2000). Other reported SHIP1 modulators include the compounds set forth in PCT Published Patent Application Nos. WO 2003/033517, WO 2004/035601, WO 2004/092100 (or US 7,601,874), WO 2007/147251, WO 2007/147252 and WO 2011/069118.


While significant strides have been made in this field, there remains a need for effective small molecule SHIP1 modulators. There is also a need for pharmaceutical compositions containing such compounds, as well as for methods relating to the use

thereof to treat disorders or conditions that would benefit from SHIP1 modulation. The present invention fulfills these needs, and provides other related advantages.


SUMMARY OF THE INVENTION

The present invention is generally directed to compounds which are SHIP1 modulators and pharmaceutical compositions comprising the compounds and methods of using the compounds and the pharmaceutical compositions of the invention for the treatment of diseases, disorders or conditions that would benefit from SHIP1 modulation. As used herein, a SHIP1 modulator can serve as either an agonist or antagonist to SHIP1.


Accordingly, in one aspect, this invention is directed to compounds of formula (I):



wherein:



is selected from:



n is 1, 2, 3, 4, 5, or 6;

R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, $-R^{9a}-C(O)-R^{9b}-$, $-R^{9a}-S(O)_t-R^{9b}-$ (where t is 0, 1 or 2),
 $-R^{9a}-O-R^{9b}-$, $-R^{9a}-C(O)N(R^{11a})-R^{9b}-$, $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$ or $-R^{9a}-N(R^{11a})-R^{9b}-$;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$,

or R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

or R^{4a} and R^{4b} together form $=C(R^{13})_2$;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

each R^6 and R^8 is independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

each R^{13} is hydrogen, alkyl or haloalkyl; and

R^{14} is a straight or branched alkylene chain;

or a stereoisomer thereof or a pharmaceutically acceptable salt thereof;

provided that compounds of formula (I) do not include the following compounds:

(1S,2S,4aS,8aR)-decahydro-1-[2-(3-methoxyphenyl)ethyl]-5,5,8a-trimethyl-2-naphthalenecarboxylic acid;

(1S,2S,4aS,8aR)-decahydro-1-[2-(3-methoxyphenyl)ethyl]-5,5,8a-trimethyl-2-naphthalenemethanol;

α -(3,5-dimethoxyphenyl)decahydro-2-hydroxy-2,5,5,8a-tetramethyl-1-naphthalenemethanol;

(4aS,8aS)-3,4,4a,5,6,7,8,8a-octahydro- α -(3-methoxy-5-methylphenyl)-2,5,5,8a-tetramethyl-1-naphthalenemethanol;

(1S,2R,4aS,8aS)-decahydro-2-hydroxy- α -(3-methoxy-5-methylphenyl)-2,5,5,8a-tetramethyl-1-naphthalenemethanol;

(1R,2R,4aS,8aS)-decahydro-1-[(3-methoxy-5-methylphenyl)methyl]-2,5,5,8a-tetramethyl-2-naphthalenol;

(4aS,5S,8aS)-decahydro-5-[2-(3-methoxyphenyl)ethyl]-1,1,4a-trimethyl-6-methylene-naphthalene;

2-[(1S,4aS,8aS)-decahydro-5,5,8a-trimethyl-2-methylene-1-naphthalenyl]-1-(3-methoxyphenyl)-ethanone;

(1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;

(1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;

(1S,2R,4aS,8aS)-1-[3,5-bis(propan-2-yloxy)phenoxyethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;

(1R,2R,8aS)-1-[(3,5-dimethoxyphenylsulfonyl)methyl]-2,5,5,8a-tetramethyldecahydronaphthalen-2-ol;

(1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)sulfonyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;

(1R,2R,4aS,8aS)-1-[(3-methoxyphenyl)sulfonyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol; and

(1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)sulfonyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol.

In another aspect, this invention is directed to compositions comprising a pharmaceutically acceptable excipient, carrier and/or diluent and a compound of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

In another aspect, this invention is directed to a method for modulating SHIP1 activity in a mammal comprising administering an effective amount of a compound of

formula (I), or a pharmaceutically acceptable salt thereof or a stereoisomer thereof, or a composition comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof or a stereoisomer thereof, to the mammal in need thereof.

In another aspect, this invention is directed to methods for treating a disease, disorder or condition in a mammal comprising administering an effective amount of a compound of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, as set forth above, to the mammal in need thereof, where the disease, disorder or condition is an autoimmune disease, disorder or condition, an inflammatory disease, disorder or condition, or a neoplastic or cell proliferative disease, disorder or condition.

In another aspect, this invention is directed to methods of treating a disease, disorder or condition in a mammal comprising administering an effective amount of a compound of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, typically in the form of a composition, to the mammal in need thereof. Methods of this invention include administering an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof or a stereoisomer thereof, to the mammal in need thereof (such as a human).

In another aspect, this invention is directed to methods of preparing compounds of formula (I), or stereoisomers thereof or pharmaceutically acceptable salts thereof.

These aspects and embodiments thereof are described in more detail below. To this end, various references are set forth herein which describe in more detail certain background information, procedures, compounds and/or compositions, and are each hereby incorporated by reference in their entirety.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated:

"Oxo" refers to =O.

"Cyano" refers to -CN.

"Nitro" refers to -NO₂.

"Hydroxy" refers to -OH.

"Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to

twelve carbon atoms, preferably one to eight carbon atoms, more preferably one to six carbon atoms, and which is attached to the rest of the molecule by a single bond, *e.g.*, methyl, ethyl, *n*-propyl, 1-methylethyl (*iso*-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), 3-methylhexyl, 2-methylhexyl, and the like. When specifically stated in the specification, an alkyl group may be optionally substituted by one of the following groups: alkyl, halo, haloalkyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilyl, $-OR^{20}$, $-OC(O)-R^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{22}$, $-N(R^{20})C(O)R^{22}$, $-N(R^{20})S(O)_pR^{22}$ (where *p* is 1 to 2), $-S(O)_pOR^{22}$ (where *p* is 1 to 2), $-S(O)_tR^{22}$ (where *t* is 0 to 2), and $-S(O)_pN(R^{20})_2$ (where *p* is 1 to 2) where each R^{20} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{22} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl.

"Alkenyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one double bond, having from two to twelve carbon atoms, preferably one to eight carbon atoms and which is attached to the rest of the molecule by a single bond, *e.g.*, ethenyl, prop-1-enyl, but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like. When specifically stated in the specification, an alkenyl group may be optionally substituted by one of the following groups: alkyl, halo, haloalkyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilyl, $-OR^{20}$, $-OC(O)-R^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{22}$, $-N(R^{20})C(O)R^{22}$, $-N(R^{20})S(O)_pR^{22}$ (where *p* is 1 to 2), $-S(O)_pOR^{22}$ (where *p* is 1 to 2), $-S(O)_tR^{22}$ (where *t* is 0 to 2), and $-S(O)_pN(R^{20})_2$ (where *p* is 1 to 2) where each R^{20} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{22} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl.

"Alkoxy" refers to a radical of the formula $-OR_a$ where R_a is an alkyl radical as defined above containing one to twelve carbon atoms. The alkyl part of the alkoxy radical may be optionally substituted as defined above for an alkyl radical

"Alkylene" or "alkylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group or linking two parts of the molecule, consisting solely of carbon and hydrogen, containing no unsaturation and having from one to twelve carbon atoms, *e.g.*, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain is attached to the rest of the

molecule through a single bond and to the radical group through a single bond or is attached to two parts of the molecule through a single bond at each point of attachment. When specifically stated in the specification, an alkylene chain may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilyl, $-OR^{20}$, $-OC(O)-R^{20}$, $-N(R^{20})_2$, $-C(O)R^{20}$, $-C(O)OR^{20}$, $-C(O)N(R^{20})_2$, $-N(R^{20})C(O)OR^{22}$, $-N(R^{20})C(O)R^{22}$, $-N(R^{20})S(O)_pR^{22}$ (where p is 1 to 2), $-S(O)_pOR^{22}$ (where p is 1 to 2), $-S(O)_tR^{22}$ (where t is 0 to 2), and $-S(O)_pN(R^{20})_2$ (where p is 1 to 2) where each R^{20} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; and each R^{22} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

"Aryl" refers to a hydrocarbon ring system radical comprising hydrogen, 6 to 18 carbon atoms and at least one aromatic ring. For purposes of this invention, the aryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems. Aryl radicals include, but are not limited to, aryl radicals derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, fluoranthene, fluorene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, phenalene, phenanthrene, pleiadene, pyrene, and triphenylene. When specifically stated in the specification, an aryl group may be optionally substituted by one or more substituents independently selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, cyano, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, heteroarylalkyl, $-R^{21}-OR^{20}$, $-R^{21}-OC(O)-R^{20}$, $-R^{21}-N(R^{20})_2$, $-R^{21}-C(O)R^{20}$, $-R^{21}-C(O)OR^{20}$, $-R^{21}-C(O)N(R^{20})_2$, $-R^{21}-N(R^{20})C(O)OR^{22}$, $-R^{21}-N(R^{20})C(O)R^{22}$, $-R^{21}-N(R^{20})S(O)_pR^{22}$ (where p is 1 to 2), $-R^{21}-N=C(OR^{20})R^{20}$, $-R^{21}-S(O)_pOR^{22}$ (where p is 1 to 2), $-R^{21}-S(O)_tR^{22}$ (where t is 0 to 2), and $-R^{21}-S(O)_pN(R^{20})_2$ (where p is 1 to 2) where each R^{20} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; each R^{21} is independently a direct bond or a straight or branched alkylene chain; and each R^{22} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

"Aralkyl" refers to a radical of the formula $-R_b-R_c$ where R_b is an alkylene chain as defined above and R_c is one or more aryl radicals as defined above, for example, benzyl, diphenylmethyl and the like. When specifically stated in the specification, the

alkylene chain part of the aralkyl radical may be optionally substituted as described above for an optionally substituted alkylene chain. When specifically stated in the specification, the aryl part of the aralkyl radical may be optionally substituted as described above for an optionally substituted aryl group.

"Cycloalkyl" refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which may include fused or bridged ring systems, having from three to fifteen carbon atoms, preferably having from three to ten carbon atoms, and which is saturated or unsaturated and attached to the rest of the molecule by a single bond. Monocyclic radicals include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic radicals include, for example, adamantyl, norbornyl, decalyl, and the like. When specifically stated in the specification, a cycloalkyl group may be optionally substituted by one or more substituents independently selected from the group consisting of alkyl, halo, haloalkyl, cyano, nitro, oxo, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, heteroarylalkyl, $-R^{21}-OR^{20}$, $-R^{21}-OC(O)-R^{20}$, $-R^{21}-N(R^{20})_2$, $-R^{21}-C(O)R^{20}$, $-R^{21}-C(O)OR^{20}$, $-R^{21}-C(O)N(R^{20})_2$, $-R^{21}-N(R^{20})C(O)OR^{22}$, $-R^{21}-N(R^{20})C(O)R^{22}$, $-R^{21}-N(R^{20})S(O)_pR^{22}$ (where p is 1 to 2), $-R^{21}-N=C(OR^{20})R^{20}$, $-R^{21}-S(O)_pOR^{22}$ (where p is 1 to 2), $-R^{21}-S(O)_tR^{22}$ (where t is 0 to 2), and $-R^{21}-S(O)_pN(R^{20})_2$ (where p is 1 to 2) where each R^{20} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; each R^{21} is independently a direct bond or a straight or branched alkylene chain; and each R^{22} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl.

"Cycloalkylalkyl" refers to a radical of the formula $-R_bR_g$ where R_b is an alkylene chain as defined above and R_g is a cycloalkyl radical as defined above. When specifically stated in the specification, the alkylene chain and/or the cycloalkyl radical may be optionally substituted as defined above for optionally substituted alkylene chain and optionally substituted cycloalkyl.

"Halo" refers to bromo, chloro, fluoro or iodo.

"Haloalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, 3-bromo-2-fluoropropyl, 1-bromomethyl-2-bromoethyl, and the like. The alkyl part of the haloalkyl radical may be optionally substituted as defined above for an alkyl group.

"Heterocyclyl" refers to a stable 3- to 18-membered non-aromatic ring radical which consists of two to twelve carbon atoms and from one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. Unless stated otherwise specifically in the specification, the heterocyclyl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclyl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized; and the heterocyclyl radical may be partially or fully saturated. Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, dioxinyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazolyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, 1,2,4-thiadiazol-5(4*H*)-ylidene, tetrahydrofuryl, trioxanyl, trithianyl, triazinanyl, tetrahydropyranyl, thiomorpholyl, thiamorpholyl, 1-oxo-thiomorpholyl, and 1,1-dioxo-thiomorpholyl. When specifically stated in the specification, a heterocyclyl group may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, cyano, oxo, thioxo, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{21}-OR^{20}$, $-R^{21}-OC(O)-R^{20}$, $-R^{21}-N(R^{20})_2$, $-R^{21}-C(O)R^{20}$, $-R^{21}-C(O)OR^{20}$, $-R^{21}-C(O)N(R^{20})_2$, $-R^{21}-N(R^{20})C(O)OR^{22}$, $-R^{21}-N(R^{20})C(O)R^{22}$, $-R^{21}-N(R^{20})S(O)_pR^{22}$ (where *p* is 1 to 2), $-R^{21}-N=C(OR^{20})R^{20}$, $-R^{21}-S(O)_pOR^{22}$ (where *p* is 1 to 2), $-R^{21}-S(O)_tR^{22}$ (where *t* is 0 to 2), and $-R^{21}-S(O)_pN(R^{20})_2$ (where *p* is 1 to 2) where each R^{20} is independently hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; each R^{21} is independently a direct bond or a straight or branched alkylene chain; and each R^{22} is alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl.

"*N*-heterocyclyl" refers to a heterocyclyl radical as defined above containing at least one nitrogen. The point of attachment of the *N*-heterocyclyl to the rest of the molecule can be through a nitrogen atom or a carbon atom in the *N*-heterocyclyl. When specifically stated in the specification, an *N*-heterocyclyl radical may be optionally substituted as described above for an optionally substituted heterocyclyl radical.

"Heterocyclalkyl" refers to a radical of the formula $-R_bR_h$ where R_b is an alkylene chain as defined above and R_h is a heterocycl radical as defined above, and if the heterocycl is a nitrogen-containing heterocycl, the heterocycl may be attached to the alkyl radical at the nitrogen atom. When specifically stated in the specification, the alkylene chain of the heterocyclalkyl radical may be optionally substituted as defined above for an optionally substituted alkylene chain. When specifically stated in the specification, the heterocycl part of the heterocyclalkyl radical may be optionally substituted as defined above for an optionally substituted heterocycl group.

"Heteroaryl" refers to a 5- to 14-membered ring system radical comprising hydrogen atoms, one to thirteen carbon atoms, one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, and at least one aromatic ring. For purposes of this invention, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzo[d]imidazolyl, benzthiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzo[d]isoxazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothieryl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, benzoxazolinonyl, benzimidazolthionyl, carbazolyl, cinnolinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazo[1,2-a]pyridinyl, imidazo[1,2-a]pyrimidinyl, imidazo[1,2-a]pyrazinyl, imidazo[1,5-a]pyrazinyl, imidazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indolizinyl, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, pteridinonyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyridinonyl, pyrazinyl, pyrimidinyl, pyrimidinonyl, pyridazinyl, pyrido[2,3-d]pyrimidinonyl, pyrazolo[1,5-a]pyrimidinyl, quinazolinyl, quinazolinonyl, quinoxalinyl, quinoxalinonyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, thiazolyl, thiadiazolyl, thieno[3,2-d]pyrimidin-4-onyl, thieno[2,3-d]pyrimidin-4-onyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl (i.e. thienyl). When specifically stated in the specification, a heteroaryl group may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo,

haloalkyl, cyano, oxo, thioxo, nitro, thioxo, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{21}-OR^{20}$, $-R^{21}-OC(O)-R^{20}$, $-R^{21}-N(R^{20})_2$, $-R^{21}-C(O)R^{20}$, $-R^{21}-C(O)OR^{20}$, $-R^{21}-C(O)N(R^{20})_2$, $-R^{21}-N(R^{20})C(O)OR^{22}$, $-R^{21}-N(R^{20})C(O)R^{22}$, $-R^{21}-N(R^{20})S(O)_pR^{22}$ (where p is 1 to 2), $-R^{21}-N=C(OR^{20})R^{20}$, $-R^{21}-S(O)_pOR^{22}$ (where p is 1 to 2), $-R^{21}-S(O)_tR^{22}$ (where t is 0 to 2), and $-R^{21}-S(O)_pN(R^{20})_2$ (where p is 1 to 2) where each R^{20} is independently hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; each R^{21} is independently a direct bond or a straight or branched alkylene chain; and each R^{22} is alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl.

"*N*-heteroaryl" refers to a heteroaryl radical as defined above containing at least one nitrogen. The point of attachment of the *N*-heteroaryl to the rest of the molecule can be through a nitrogen atom or a carbon atom in the *N*-heteroaryl. When specifically stated in the specification, an *N*-heteroaryl radical may be optionally substituted as described above for an optionally substituted heteroaryl radical.

"Heteroarylalkyl" refers to a radical of the formula $-R_bR_i$ where R_b is an alkylene chain as defined above and R_i is a heteroaryl radical as defined above. When specifically stated in the specification, the heteroaryl part of the heteroarylalkyl radical may be optionally substituted as defined above for an optionally substituted heteroaryl group. When specifically stated in the specification, the alkylene chain part of the heteroarylalkyl radical may be optionally substituted as defined above for an optionally substituted alkylene chain.

"Prodrugs" is meant to indicate a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound of the invention. Thus, the term "prodrug" refers to a metabolic precursor of a compound of the invention that is pharmaceutically acceptable. A prodrug may be inactive when administered to a subject in need thereof, but is converted *in vivo* to an active compound of the invention. Prodrugs are typically rapidly transformed *in vivo* to yield the parent compound of the invention, for example, by hydrolysis in blood. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in a mammalian organism (see, Bundgard, H., *Design of Prodrugs* (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam)). A discussion of prodrugs is provided in Higuchi, T., *et al.*, "Pro-drugs as Novel Delivery Systems," A.C.S. Symposium Series, Vol. 14, and in *Bioreversible Carriers in Drug Design*, Ed. Edward B. Roche, American

Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated in full by reference herein.

The term "prodrug" is also meant to include any covalently bonded carriers, which release the active compound of the invention *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of the invention may be prepared by modifying functional groups present in the compound of the invention in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound of the invention. Prodrugs include compounds of the invention wherein a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the compound of the invention is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol or amide derivatives of amine functional groups in the compounds of the invention and the like. Further, in the case of a carboxylic acid ($-C(O)OH$), esters may be employed, such as methyl esters, ethyl esters, and the like.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

"Mammal" includes humans and both domestic animals such as laboratory animals and household pets, (e.g., cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildlife and the like.

"Optional" or "optionally" means that the subsequently described event or circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution ("unsubstituted"). When a functional group is described as "optionally substituted," and in turn, substituents on the functional group are also "optionally substituted" and so on, for the purposes of this invention, such iterations are limited to five, preferably such iterations are limited to two.

"Pharmaceutically acceptable carrier, diluent or excipient" includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been

approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

"Pharmaceutically acceptable salt" includes both acid and base addition salts.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, *p*-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

"Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine,

procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

Often crystallizations produce a solvate of the compound of the invention. As used herein, the term "solvate" refers to an aggregate that comprises one or more molecules of a compound of the invention with one or more molecules of solvent. The solvent may be water, in which case the solvate may be a hydrate. Alternatively, the solvent may be an organic solvent. Thus, the compounds of the present invention may exist as a hydrate, including a monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate and the like, as well as the corresponding solvated forms. The compound of the invention may be true solvates, while in other cases, the compound of the invention may merely retain adventitious water or be a mixture of water plus some adventitious solvent. Furthermore, some of the crystalline forms of the compounds of formula (I) may exist as polymorphs, which are included in the present invention.

A "pharmaceutical composition" refers to a formulation of a compound of the invention and a medium generally accepted in the art for the delivery of the biologically active compound to mammals, *e.g.*, humans. Such a medium includes all pharmaceutically acceptable carriers, diluents or excipients therefor.

"Therapeutically effective amount" refers to that amount of a compound of the invention which, when administered to a mammal, preferably a human, is sufficient to effect treatment, as defined below, of a disease or condition alleviated by the modulation of SHIP1 in the mammal, preferably a human. The amount of a compound of the invention which constitutes a "therapeutically effective amount" will vary depending on the compound, the condition and its severity, the manner of administration, and the age of the mammal to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

"Treating" or "treatment" as used herein covers the treatment of the disease or condition of interest in a mammal, preferably a human, having the disease or condition of interest, and includes:

(a) preventing the disease or condition from occurring in a mammal, in particular, when such mammal is predisposed to the condition but has not yet been diagnosed as having it;

(b) inhibiting the disease or condition, *i.e.*, arresting its development;

(c) relieving (or ameliorating) the disease or condition, *i.e.*, causing regression of the disease or condition; or

(d) relieving (or ameliorating) the symptoms resulting from the disease or condition, *i.e.*, relieving inflammation without addressing the underlying disease or condition.

As used herein, the terms "disease" and "condition" may be used interchangeably or may be different in that the particular malady or condition may not have a known causative agent (so that etiology has not yet been worked out) and it is therefore not yet recognized as a disease but only as an undesirable condition or syndrome, wherein a more or less specific set of symptoms have been identified by clinicians.


The compounds of the invention, or their pharmaceutically acceptable salts may contain one or more asymmetric centres and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. Compounds of formula (I) may also possess axial chirality which may result in atropisomers. The present invention is meant to include all such possible isomers, as well as their racemic and optically pure forms. Optically active (+) and (-), (*R*)- and (*S*)-, or (*D*)- and (*L*)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallisation. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centres of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers. Likewise, all tautomeric forms are also intended to be included.

A "stereoisomer" refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. The present invention contemplates various stereoisomers and mixtures thereof and includes enantiomers, which refers to two stereoisomers whose







molecules are nonsuperimposable mirror images of one another. See, for example, Smith, M.B. and J. March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 6th edition (Wiley, 2007), for a detailed description of the structure and properties of enantiomers and stereoisomers.

A "tautomer" refers to a proton shift from one atom of a molecule to another atom of the same molecule. The present invention includes tautomers of any said compounds.

Representative isomers of the compounds of formula (I) include, but are not limited to, the following structures where R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$:







or the following structures where R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3:




;

or the following structures where R^{4a} and R^{4b} together form $=C(R^{13})_2$:



The use of parentheses and brackets in substituent groups is used herein to conserve space. Accordingly, the use of parenthesis in a substituent group indicates that the group enclosed within the parentheses is attached directly to the atom preceding the parenthesis. For example, the following substituent group:




is represented herein as $\text{---R}^9\text{---N}(\text{R}^{11})\text{C}(\text{O})\text{---R}^9\text{---N}(\text{R}^{11})\text{R}^{12}$.

The chemical naming protocol and structure diagrams used herein are a modified form of the I.U.P.A.C. nomenclature system, using ChemBioDraw Ultra Version 12.0 software program, wherein the compounds of the invention are named herein as derivatives of a central core structure. For complex chemical names employed herein, a substituent group is named before the group to which it attaches. For example, cyclopropylethyl comprises an ethyl backbone with cyclopropyl substituent. In chemical structure diagrams, all bonds are identified, except for some carbon atoms, which are assumed to be bonded to sufficient hydrogen atoms to complete the valency.

Certain carbons are identified by numerals in formula (I) of the compounds of the invention. Thus, the carbon in the decahydronaphthalene central moiety to which R^1 is attached is indicated by the numeral 1. Likewise, the carbon to which R^5 is attached is indicated by numeral 3. These numerals may or may not be the same as the locants in the compound names given herein.

Thus, for example, a compound of formula (I) wherein m and n are each 1, R^1 is $-R^{9a}-C(O)-R^{9b}-$ where R^{9a} and R^{9b} are both direct bonds, R^2 and R^3 are each $-R^9-O-R^{11}$ where R^9 is a direct bond and R^{11} is hydrogen, R^{4a} is hydrogen and R^{4b} is methyl, R^5 is $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$ where each R^9 is a direct bond, each R^{11} is hydrogen and R^{12} is ethyl, and R^6 , R^7 and R^8 are each hydrogen, *i.e.*, a compound of the following structure:



is named herein as 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-ethylurea.

The following abbreviations are used herein:

Ac₂O for acetic anhydride;

AcOH for acetic acid;

AlMe₃ for trimethylaluminum;

Boc for *tert*-butoxycarbonyl;

BH₃·THF for borane tetrahydrofuran complex;


BnBr for benzyl bromide;
Bu₃SnH for tributyltin hydride;
n-BuLi for *n*-butyl lithium;
CDI for 1,1'-carbonyldiimidazole;
DABCO for 1,4-diazabicyclo[2.2.2]octane;
DBU for 1,8-diazabicyclo[5.4.0]undec-7-ene;
DCC for N,N'-dicyclohexylcarbodiimide;
DCE for dichloroethane;
Diglyme for diethylene glycol dimethyl ether;
DIPEA/DIEA for N,N-diisopropylethylamine;
DMAP for 4-dimethylaminopyridine;
DMF for N,N-dimethylformamide;
DMSO for dimethyl sulfoxide;
Et₂O for diethyl ether;
Et₃N for triethylamine;
EtNCO for ethyl isocyanate;
EtOAc for ethyl acetate;
EtOH for ethanol;
H₂/Pd/C for hydrogen on palladium on charcoal;
H₂NMe·HCl for methylamine hydrochloride;
HBTU for O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate;
IBX for 2-iodoxybenzoic acid;
i-PrOH for *iso*-propanol;
Imid. for imidazole;
KOtBu for potassium tert-butoxide;
LiAlH₄/LAH for lithium aluminum hydride;
LiEt₃BH for lithium triethylborohydride (Super hydride);
m-CPBA for m-chloroperoxybenzoic acid;
MeCN for acetonitrile;
MeI for methyl iodide;
MeNCS for methyl isocyanate;
2-MePhCO₂H for *o*-toluic acid;
3-MePhCO₂H for *m*-toluic acid;
4-MePhCO₂H for *p*-toluic acid;
Me₄Phen for 3,4,7,8-tetramethyl-[1,10]-phenanthroline;

MeOCH₂PPh₃Cl for methoxymethyl triphenylphosphonium chloride;
MeOH for methanol;
MePPh₃Br for methyl triphenylphosphonium bromide;
MeSO₃SiMe₃ for trimethylsilylmethanesulfonate;
MsCl for mesyl chloride;
MW for microwave;
NaSEt for sodium ethanethiolate;
NaBH(OAc)₃ for sodium triacetoxyborohydride;
n-BuLi for *n*-butyllithium;
NMP for N-methyl-2-pyrrolidone;
NMR for nuclear magnetic resonance;
PCC for pyridinium chlorochromate;
Pd/C for palladium metal on charcoal;
PhCO₂H for benzoic acid;
PPh₃ for triphenylphosphine;
PhMe for toluene;
p-TsOH.H₂O for *p*-toluenesulfonic acid monohydrate;
PyBOP for benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate;
Pyr for pyridine;
TBAF for tetrabutylammonium fluoride;
TBDPS for *tert*-butyldiphenylsilyl;
TBDPSCI for *tert*-butyldiphenylsilyl chloride;
TBTU for O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate;
TFA for trifluoroacetic acid;
THF for tetrahydrofuran;
TLC for thin layer chromatography;
TPSH for 2,4,6-triisopropylbenzenesulfonyl hydrazide; and
VAZO® for 1,1'-azobis(cyclohexanecarbonitrile).

Embodiments of the invention


Of the various aspects of the invention set forth above in the Summary of the Invention, certain embodiments are preferred.

One embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (Ia):



wherein n , R^1 , R^2 , R^3 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above for compounds of formula (I) as set forth above in the Summary of the Invention, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^{9a}-C(O)-R^{9b}-$, *i.e.*, a compound of formula (Ia1):



wherein R^{9a} and R^{9b} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl,

optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$,

-R⁹-N(R¹¹)C(O)N(R¹¹)-OR¹², -R⁹-N(R¹¹)C(=NR¹¹)N(R¹¹)R¹², -R⁹-N(R¹¹)S(O)_pR¹¹ (where p is 1 or 2), -R⁹-N(R¹¹)C(S)N(R¹¹)R¹² or -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2); where each R⁹ is independently a direct bond or a straight or branched alkylene chain; where each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R¹⁴ is a straight or branched alkylene chain;

R⁶, R⁷ and R⁸ are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen or methyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen;

R⁵ is independently selected from hydrogen, oxo, cyano, optionally substituted heterocyclylalkyl, optionally substituted heteroarylalkyl, -R⁹-OR¹¹, -R⁹-C(O)R¹¹, -R⁹-C(O)OR¹¹, -R⁹-N(R¹¹)R¹², -R⁹-C(O)N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)R¹², -R⁹-N(R¹¹)-R¹⁴-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)N(R¹¹)-OR¹², -R⁹-N(R¹¹)C(=NR¹¹)N(R¹¹)R¹², -R⁹-N(R¹¹)S(O)_pR¹¹ (where p is 1 or 2), -R⁹-N(R¹¹)C(S)N(R¹¹)R¹² or -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2); where each R⁹ is independently a direct bond or a straight or branched alkylene chain; where each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R¹⁴ is a straight

or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is hydrogen, oxo, cyano or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} is a direct bond or a methylene chain and R^{9b} is a direct bond.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(2R,3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol;

(2R,3R,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol;

(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol;

(2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol;

(4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbonitrile;

(3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one;

(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one;

(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one;

2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-

dimethoxyphenyl)ethan-1-one;
 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3-hydroxy-5-methoxyphenyl)ethan-1-one; and
 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dihydroxyphenyl)ethan-1-one.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen or methyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} is hydrogen and R^{4b} is methyl;

R⁵ is independently selected from -R⁹-C(O)R¹¹, -R⁹-C(O)OR¹¹ or -R⁹-C(O)N(R¹¹)R¹²; where each R⁹ is independently a direct bond or a straight or branched alkylene chain; and where each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R⁶, R⁷ and R⁸ are each hydrogen; and

R^{9a} and R^{9b} are each a direct bond.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide;

(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide;

(3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid;

(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid;

2-[[[(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbonyl]pyridine; and

5-{[(1S,2R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-yl)carbonyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen or methyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} is hydrogen and R^{4b} is methyl;

R⁵ is independently selected -R⁹-N(R¹¹)R¹² or -R⁹-N(R¹¹)-R¹⁴-N(R¹¹)R¹²; where each R⁹ is independently a direct bond or a straight or branched alkylene chain; where each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R¹⁴ is a straight or branched alkylene chain;

R⁶, R⁷ and R⁸ are each hydrogen; and

R^{9a} and R^{9b} are each a direct bond.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(3S,4S,4aS,8aS)-N-(3-aminopropyl)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine;

3-{[(1S,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}-5-methylphenol;

(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine;

(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine;

5-{[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;

5-{[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-5-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;

5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(1H-pyrrol-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;
 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;
 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-4-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;
 5-[[[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-2-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;
 (2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine;
 3-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}-5-methylphenol;
 3-[[[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-5-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}-5-methylphenol; and
 5-[[[(1S,2S,3R,4aS,8aS)-3-[(3-aminopropyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen or methyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} is hydrogen and R^{4b} is methyl;

R⁵ is independently selected from -R⁹-N(R¹¹)C(O)R¹², -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)N(R¹¹)-OR¹², -R⁹-N(R¹¹)C(=NR¹¹)N(R¹¹)R¹², -R⁹-N(R¹¹)S(O)_pR¹¹ (where p is 1 or 2), -R⁹-N(R¹¹)C(S)N(R¹¹)R¹² or -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2); where each R⁹ is independently a direct bond or a straight or branched alkylene chain; and where each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R⁶, R⁷ and R⁸ are each hydrogen;

R^{9a} and R^{9b} are each a direct bond.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide;

N-[(3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyrazine-2-carboxamide;

N-[(2R,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide;

N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide;

N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}pyridine-3-carboxamide;

N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}benzamide;

3-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea;

1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-

decahydronaphthalen-2-yl]-3-ethylurea;
 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-3-methoxyurea;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]benzamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]pyridine-3-carboxamide;
 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-3-methylurea;
 3-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]acetamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]pyridine-2-carboxamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]pyridine-4-carboxamide;
 3-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]carbamoyl}pyridin-1-ium-1-olate;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-4-methylpiperazine-1-carboxamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-3-aminopropanamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]piperidine-4-carboxamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]pyridine-3-carboxamide;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-6-aminopyridine-3-carboxamide;
 1-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]guanidine;
 N-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]methanesulfonamide;
 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]guanidine;

N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide;
 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylthiourea;
 2-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl)methyl)sulfamoyl]benzoic acid.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen or methyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} is hydrogen and R^{4b} is methyl;

R⁵ is independently selected from optionally substituted heterocyclalkyl or optionally substituted heteroarylalkyl

R⁶, R⁷ and R⁸ are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

5-[(1S,2R,3S,4aS,8aS)-3-(1H-imidazol-1-ylmethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol;

5-[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-(1H-1,2,4-triazol-1-ylmethyl)-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol; and

5-[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(4-methylpiperazin-1-yl)methyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen or methyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} is alkyl and R^{4b} is $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;
 R^5 is independently selected from hydrogen, oxo, cyano, optionally substituted heterocyclalkyl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocycl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is alkyl and R^{4b} is $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is hydrogen

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a methylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;

2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-

dimethoxyphenyl)ethan-1-one; and
 (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)carbonyl]-2,5,5,8a-tetramethyl-
 decahydronaphthalen-2-ol.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxocyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is a independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 and R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl; and

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is hydrogen or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^6 , R^7 and R^8 are each hydrogen; and
 R^{9a} and R^{9b} are each a direct bond.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from :

[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone;
 3-[[[(4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl]-5-methylphenol;
 3-[[[(4aS,8aS)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl]-5-methylphenol;
 [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone; and
 (2R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-2-ol.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} together form $=C(R^{13})_2$;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene

chain;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

each R^{13} is hydrogen, alkyl or haloalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} together form $=C(R^{13})_2$;

R^5 is independently selected from hydrogen or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;


R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

each R^{13} is hydrogen, alkyl or haloalkyl.

Another embodiment of a compound of formula (Ia1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is (2R,4R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-4a,8,8-trimethyl-3-methyldene-decahydronaphthalen-2-ol.

Another embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}$ -, *i.e.*, the compound of formula (Ia2):



wherein R^{9a} , R^{9b} and R^{10} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$,

$-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$ where R^9 is a direct bond or a methylene chain and each R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is independently hydrogen or $-OR^{11}$ where R^{11} is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$ where R^9 is a direct bond or a methylene chain and each R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, optionally substituted heterocyclalkyl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclalkyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is independently hydrogen or $-OR^{11}$ where R^{11} is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is hydrogen and R^{4b} is alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$ where R^9 is a direct bond or a methylene chain and each R^{11} is hydrogen;

R^5 is hydrogen;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} is a direct bond or a methylene chain and R^{9b} is a direct bond; and each R^{10} is independently hydrogen or $-OR^{11}$ where R^{11} is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

[(1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol;

3-[(1S,2S,4aS,8aR)-2-(hydroxymethyl)-5,5,8a-trimethyl-decahydronaphthalen-1-yl]methyl-5-methylphenol;

(1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid;

(1S,2S,4aS,8aR)-1-[(3-hydroxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid;

5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-hydroxyethyl}benzene-1,3-diol;

2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-ol;

(4aR,5S,6S,8aS)-5-[2-(3,5-dimethoxyphenyl)ethyl]-1,1,4a,6-tetramethyl-decahydronaphthalene;

5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethyl}benzene-1,3-diol; and

(1S,2S,4aS,8aR)-1-[(3,5-dihydroxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9

is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is alkyl and R^{4b} is $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, optionally substituted heterocyclalkyl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclalkyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is independently hydrogen or $-OR^{11}$ where R^{11} is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is alkyl and R^{4b} is $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is hydrogen;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} is a direct bond or a methylene chain and R^{9b} is a direct bond; and each R^{10} is independently hydrogen or $-OR^{11}$ where R^{11} is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9

is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is a independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 and R^8 are each independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is independently hydrogen or $-OR^{11}$ where R^{11} is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R⁵ is hydrogen;
 R⁶, R⁷ and R⁸ are each hydrogen;
 R^{9a} and R^{9b} are each a direct bond; and
 each R¹⁰ is hydrogen.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(4aS,8aS)-8-[(3-methoxy-5-methylphenyl)methyl]-4,4,7,8a-tetramethyl-1,2,3,4,4a,5,6,8a-octahydronaphthalene; and
 (4aS,5S,8aS)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹, provided that at least one of R² and R³ is -R⁹-OR¹¹ when R⁶ is hydrogen;

R^{4a} and R^{4b} together form =C(R¹³)₂;

R⁵ is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, -R⁹-OR¹¹, -R⁹-C(O)R¹¹, -R⁹-C(O)OR¹¹, -R⁹-N(R¹¹)R¹², -R⁹-C(O)N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)R¹², -R⁹-N(R¹¹)-R¹⁴-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)N(R¹¹)-OR¹², -R⁹-N(R¹¹)C(=NR¹¹)N(R¹¹)R¹², -R⁹-N(R¹¹)S(O)_pR¹¹ (where p is 1 or 2), -R⁹-N(R¹¹)C(S)N(R¹¹)R¹² or -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2);

R⁶ and each R⁸ are independently selected from hydrogen, alkyl, halo or haloalkyl;

R⁷ is hydrogen, alkyl, halo or haloalkyl;

each R⁹, R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R¹⁰ is independently hydrogen, alkyl, -OR¹¹, -C(O)OR¹¹, -C(O)N(R¹¹)R¹², -N(R¹¹)R¹² or -N(R¹¹)C(O)R¹¹;

each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally

substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and each R^{13} is hydrogen, alkyl or haloalkyl; and R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} together form $=C(R^{13})_2$;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 and R^8 are each independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$,
 $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{13} is hydrogen, alkyl or haloalkyl.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} together form $=C(R^{13})_2$;

R^5 is hydrogen;

R^6 , R^7 and R^8 are each hydrogen;


R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is hydrogen;

each R^{13} is hydrogen, alkyl or haloalkyl.

Another embodiment of a compound of formula (Ia2), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is (4a*S*,5*S*,8a*S*)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene.

Another embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^9-S(O)_t-R^{9b}$, *i.e.*, a compound of formula (Ia3):



wherein t , R^{9a} and R^{9b} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia3), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

t is 0, 1 or 2;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia3), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

t is 0, 1 or 2;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9

is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia3), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

t is 0, 1 or 2;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, optionally substituted

heterocyclalkyl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocycl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia3), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

t is 0, 1 or 2;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is selected from hydrogen, $-R^9-OR^{11}$ or $-R^9-N(R^{11})R^{12}$; where each R^9 is independently a direct bond or a straight or branched alkylene chain; and where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocycl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;


R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} is a methylene chain and R^{9b} is a direct bond.

Another embodiment of a compound of formula (Ia3), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(1R,2R,3R,4aS,8aS)-3-amino-1-[[[(3,5-dimethoxyphenyl)sulfonyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol; and
(4S,4aS,8aS)-4-[[[(3,5-dimethoxybenzene)sulfonyl]methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol.

Another embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^{9a}-O-R^{9b}-$, *i.e.*, a compound of formula (Ia4):



;

wherein R^{9a} and R^{9b} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$,

-R⁹-N(R¹¹)-R¹⁴-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)R¹²,
 -R⁹-N(R¹¹)C(O)N(R¹¹)-OR¹², -R⁹-N(R¹¹)C(=NR¹¹)N(R¹¹)R¹², -R⁹-N(R¹¹)S(O)_pR¹¹
 (where p is 1 or 2), -R⁹-N(R¹¹)C(S)N(R¹¹)R¹² or
 -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2);

R⁶ and each R⁸ are independently selected from hydrogen, alkyl, halo or haloalkyl;

R⁷ is hydrogen, alkyl, halo or haloalkyl;

each R⁹, R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R¹⁴ is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R² and R³ are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen, alkyl or optionally substituted aralkyl, provided that at least one of R² and R³ is -R⁹-OR¹¹;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or -R⁹-OR¹¹ where R⁹ is a direct bond and R¹¹ is hydrogen;

R⁵ is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, -R⁹-OR¹¹, -R⁹-C(O)R¹¹, -R⁹-C(O)OR¹¹, -R⁹-N(R¹¹)R¹², -R⁹-C(O)N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)R¹², -R⁹-N(R¹¹)-R¹⁴-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)R¹², -R⁹-N(R¹¹)C(O)N(R¹¹)-OR¹², -R⁹-N(R¹¹)C(=NR¹¹)N(R¹¹)R¹², -R⁹-N(R¹¹)S(O)_pR¹¹ (where p is 1 or 2), -R⁹-N(R¹¹)C(S)N(R¹¹)R¹² or -R⁹-N(R¹¹)C(O)-R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2); where each R⁹ is independently a direct bond or a straight or branched alkylene chain; where each R¹¹ and R¹² is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally

substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen, alkyl or optionally substituted aralkyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, optionally substituted heterocyclalkyl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen, alkyl or benzyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is selected from hydrogen or $-R^9-OR^{11}$ where R^9 is independently a direct bond or a straight or branched alkylene chain and R^{11} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} is a methylene chain and R^{9b} is a direct bond.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

(4S,4aS,8aS)-4-(3-methoxy-5-methylphenoxyethyl)-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol; and

(1S,2R,4aS,8aS)-1-[3,5-bis(benzyloxy)-phenoxyethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted

heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$,

$-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is a independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 and R^8 are each independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl; and

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is hydrogen;

R^6 , R^7 and R^8 are each hydrogen; and


R^{9a} is a methylene chain and R^{9b} is a direct bond.

Another embodiment of a compound of formula (Ia4), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

3-([(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy)-5-methylphenol; and

3-([(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy)-5-methoxyphenol.

Another embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^{9a}-C(O)N(R^{11a})-R^{9b}-$, *i.e.*, a compound of formula (Ia5):



wherein R^{9a} , R^{9b} and R^{11a} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally

substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or alkyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

R^{11a} is hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally

substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or alkyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is independently selected from hydrogen, halo, oxo, cyano, optionally substituted heterocyclalkyl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each independently a direct bond or a straight or branched alkylene chain; and

R^{11a} is hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

t is 0, 1 or 2;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;

R^5 is selected from hydrogen, halo or $-R^9-OR^{11}$ where R^9 is independently a direct bond or a straight or branched alkylene chain and R^{11} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 , R^7 and R^8 are each hydrogen;

R^{9a} is a methylene chain and R^{9b} is a direct bond; and;

R^{11a} is hydrogen, alkyl, optionally substituted aryl or optionally substituted aralkyl.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)acetamide; and

2-[(1S,2R,3R,4aS,8aR)-3-bromo-2,5,5,8a-tetramethyl-decahydro-naphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted

heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is a independently a direct bond or a straight or branched alkylene chain; where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and where R^{14} is a straight or branched alkylene chain;

R^6 and R^8 are each independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain and

R^{11a} is hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen or methyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} is selected from alkyl and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is hydrogen;

R^6 , R^7 and R^8 are each hydrogen;


R^{9a} is a direct bond or methylene and R^{9b} is a direct bond; and

R^{11a} is hydrogen, alkyl, optionally substituted aryl or optionally substituted aralkyl.

Another embodiment of a compound of formula (Ia5), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

2-[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide;
 (4aS,8aS)-N-(3,5-dimethoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide;
 (4aS,8aS)-N-(3-hydroxy-5-methoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide; and
 (4aS,8aS)-N-(3,5-dihydroxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide.

Another embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$, i.e., a compound of formula (Ia6):



wherein R^{9a} , R^{9b} and R^{11a} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia6), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally

substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia6), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or

$-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is

independently a direct bond or a straight or branched alkylene chain; and

where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl,

optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted

heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted

heteroaryl or optionally substituted heteroarylalkyl;

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} is a straight or branched alkylene chain;

R^{9b} is a direct bond or a straight or branched alkylene chain; and

R^{11a} is hydrogen, alkyl, optionally substituted aryl or optionally substituted aralkyl.

Another embodiment of a compound of formula (Ia6), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is $-R^9-N(R^{11})C(O)R^{12}$ where each R^9 is a direct bond, R^{11} is hydrogen or alkyl, and R^{12} is optionally substituted aryl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl or optionally substituted heteroaryl;

R^6 is hydrogen, halo or haloalkyl;

R^7 and R^8 are each hydrogen;

R^{9a} is methylene or ethylene;

R^{9b} is a direct bond; and

R^{11a} is hydrogen.

Another embodiment of a compound of formula (Ia6), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-((4-(trifluoromethyl)benzamido)methyl)-decahydronaphthalen-2-yl)nicotinamide;

4-methyl-N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-((4-(trifluoromethyl)benzamido)methyl)-decahydronaphthalen-2-yl)piperazine-1-carboxamide;


N-((2R,3S,4S,4aS)-4-((4-fluorobenzamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide;

4-methyl-N-((2R,3S,4S,4aR)-3,4a,8,8-tetramethyl-4-(2-(4-(trifluoromethyl)benzamido)ethyl)-decahydronaphthalen-2-yl)piperazine-1-carboxamide; and

N-((2R,3S,4S,4aR)-4-(2-(4-fluorobenzamido)ethyl)-3,4a,8,8-

tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide.

Another embodiment of a compound of formula (Ia), as described above, is a compound where R^1 is $-R^{9a}-N(R^{11a})-R^{9b}$, *i.e.*, a compound of formula (Ia7):



wherein R^{9a} , R^{9b} and R^{11a} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ia7), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1, 2, 3, 4, 5, or 6;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain.

Another embodiment of a compound of formula (Ia7), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen, alkyl or benzyl, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$ where R^9 is a direct bond and R^{11} is hydrogen;


R^5 is selected from hydrogen or $-R^9-OR^{11}$ where R^9 is independently a direct bond or a straight or branched alkylene chain and R^{11} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 , R^7 and R^8 are each hydrogen; and

R^{9a} is a methylene chain and R^{9b} is a direct bond.


Another embodiment of a compound of formula (Ia7), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is the compound which is (2R,4aS,8aS)-1-[[[(3-hydroxy-5-methoxyphenyl)amino]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol.

Another embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (Ib):



wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ib), as described above, is a compound wherein R^1 is $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$, *i.e.*, a compound of formula (Ib1):



wherein R^{9a} , R^{9b} and R^{11a} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ib1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; and

where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} is a straight or branched alkylene chain;

R^{9b} is a direct bond or a straight or branched alkylene chain; and

R^{11a} is hydrogen, alkyl, optionally substituted aryl or optionally substituted aralkyl.

Another embodiment of a compound of formula (Ib1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is $-R^9-N(R^{11})C(O)R^{12}$ where each R^9 is a direct bond, R^{11} is hydrogen or alkyl, and R^{12} is optionally substituted aryl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl or optionally substituted heteroaryl;

R^6 is hydrogen, halo or haloalkyl;

R^7 and R^8 are each hydrogen;

R^{9a} is methylene or ethylene;

R^{9b} is a direct bond; and

R^{11a} is hydrogen.


Another embodiment of a compound of formula (Ib1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide;

N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide; and


N-((2R,3S,4S,4aR)-4-(2-(furan-2-carboxamido)ethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide.

Another embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (Ic):



wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ic), as described above, is a compound wherein R^1 is $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$, *i.e.*, a compound of formula (Ic1):



wherein R^{9a} , R^{9b} and R^{11a} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ic1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; and where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl,

optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} is a straight or branched alkylene chain;

R^{9b} is a direct bond or a straight or branched alkylene chain; and

R^{11a} is hydrogen, alkyl, optionally substituted aryl or optionally substituted aralkyl.

Another embodiment of a compound of formula (Ic1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is $-R^9-N(R^{11})C(O)R^{12}$ where each R^9 is a direct bond, R^{11} is hydrogen or alkyl, and

R^{12} is optionally substituted aryl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl or optionally substituted heteroaryl;

R^6 is hydrogen;

R^7 and R^8 are each hydrogen;

R^{9a} is methylene or ethylene;

R^{9b} is a direct bond; and

R^{11a} is hydrogen.

Another embodiment of a compound of formula (Ic1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound selected from:

N-((2R,3S,4S,4aS)-4-(isonicotinamidomethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide;


N-(((1S,2S,3R,8aS)-2,5,5,8a-tetramethyl-3-(nicotinamido)decahydronaphthalen-1-yl)methyl)nicotinamide;

N-((2R,3S,4S,4aS)-4-(isonicotinamidomethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide;

4-methyl-N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-(nicotinamidomethyl)decahydronaphthalen-2-yl)piperazine-1-carboxamide; and


4-methyl-N-((2R,3S,4S,4aR)-3,4a,8,8-tetramethyl-4-(2-(nicotinamido)ethyl)decahydronaphthalen-2-yl)piperazine-1-carboxamide.

Another embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (Id):



wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Id), as described above, is a compound wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, a compound of formula (Id1):



wherein R^{9a} , R^{9b} and R^{10} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Id1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or

$-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; and where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain; and

each R^{10} is hydrogen or alkyl.

Another embodiment of a compound of formula (Id1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is $-R^9-N(R^{11})C(O)R^{12}$ where each R^9 is a direct bond, R^{11} is hydrogen or alkyl, and R^{12} is optionally substituted aryl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl or optionally substituted heteroaryl;

R^6 is hydrogen, halo or haloalkyl;

R^7 and R^8 are each hydrogen;


R^{9a} and R^{9b} are each a direct bond; and

each R^{10} is hydrogen.

Another embodiment of a compound of formula (Id1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is


N-((2R,3S,4S,4aS)-4-((1H-imidazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide.

Another embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (Ie):



wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ie), as described above, is a compound wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, a compound of formula (Ie1):



wherein R^{9a} , R^{9b} and R^{10} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (Ie1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; and where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl,

optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain;

and

each R^{10} is hydrogen or alkyl.

Another embodiment of a compound of formula (Ie1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is $-R^9-N(R^{11})C(O)R^{12}$ where each R^9 is a direct bond, R^{11} is hydrogen or alkyl, and

R^{12} is optionally substituted aryl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl or optionally substituted heteroaryl;

R^6 is hydrogen, halo or haloalkyl;


R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each a direct bond; and

each R^{10} is hydrogen.


Another embodiment of a compound of formula (Ie1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is N-((2R,3S,4S,4aS)-4-((1H-1,2,4-triazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide.

Another embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (If):



wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (If), as described above, is a compound wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, a compound of formula (If1):



wherein R^{9a} , R^{9b} and R^{10} are as defined above in the Summary of the Invention for compounds of formula (I), or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of a compound of formula (If1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$,

R^5 is independently selected from $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or

$-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2); where each R^9 is independently a direct bond or a straight or branched alkylene chain; and where each R^{11} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

R^6 and each R^8 are independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

R^{9a} and R^{9b} are independently a direct bond or a straight or branched alkylene chain;

and

each R^{10} is hydrogen or alkyl.

Another embodiment of a compound of formula (If1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is a compound wherein:

n is 1;

R^{4a} is hydrogen and R^{4b} is methyl;

R^5 is $-R^9-N(R^{11})C(O)R^{12}$ where each R^9 is a direct bond, R^{11} is hydrogen or alkyl, and

R^{12} is optionally substituted aryl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl or optionally substituted heteroaryl;

R^6 is hydrogen, halo or haloalkyl;


R^7 and R^8 are each hydrogen;

R^{9a} and R^{9b} are each a direct bond; and


each R^{10} is hydrogen.

Another embodiment of a compound of formula (If1), as described above, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, is N-((2R,3S,4S,4aS)-4-((1H-benzo[d]imidazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide.

Another embodiment of a compound of formula (I), as set forth above in the Summary of the Invention, is a compound of formula (I) wherein:



is selected from:



n is 1, 2, 3, 4, 5, or 6;

R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, $-R^{9a}-C(O)-R^{9b}-$, $-R^{9a}-S(O)_t-R^{9b}-$ (where t is 0, 1 or 2), $-R^{9a}-O-R^{9b}-$, or $-R^{9a}-C(O)N(R^{11a})-R^{9b}-$ or $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$,

or R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

or R^{4a} and R^{4b} together form $=C(R^{13})_2$;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl,

optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

each R^6 and R^8 is independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl;

each R^{13} is hydrogen, alkyl or haloalkyl; and

R^{14} is a straight or branched alkylene chain;

or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

One embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is an autoimmune disease, disorder or condition, an inflammatory disease, disorder or condition, or a neoplastic or cell proliferative disease, disorder or condition.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is an autoimmune disease, disorder or condition selected from idiopathic pulmonary fibrosis, an inflammatory bowel disease, rheumatoid arthritis, Still's Disease, Sjögren's Syndrome, systemic lupus erythematosus, multiple sclerosis, psoriasis and systemic sclerosis.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is an inflammatory bowel disease selected from Crohn's Disease and ulcerative colitis.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is an inflammatory disease, disorder or condition selected from acute respiratory distress syndrome, allergic rhinitis, Alzheimer's Disease, asthma, an ocular inflammatory disease, atopic dermatitis, bladder pain syndrome/ interstitial cystitis, chronic obstructive pulmonary disease (COPD) including emphysematous, bronchitic, and alpha 1 anti-trypsin deficiency related COPD; dermal contact hypersensitivity, eczema, eosinophilic gastrointestinal disorder, fibromyalgia, gout, hepatic fibrosis, irritable bowel syndrome, ischemic reperfusion disease, kidney fibrosis, pancreatitis, Parkinson's Disease, post operative inflammation, a seronegative spondyloarthropathy, and vasculitis.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is an ocular inflammatory disease selected from allergic conjunctivitis, dry eye, and uveitis.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is a seronegative spondyloarthropathy selected from ankylosing spondylitis, psoriatic arthritis, and Reiter's Syndrome.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is vasculitis selected from Wegener's Granulomatosis, polyarteritis nodosa, leukocytoclastic vasculitis, Churg-Strauss Syndrome, cryoglobulinemic vasculitis, and giant cell arteritis.

Another embodiment of the methods for treating a disease, disorder or condition in a mammal in need thereof is where the disease, disorder or condition is a neoplastic or cell proliferative disease, disorder or condition selected from acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia, basophilic leukemia, cutaneous T-cell lymphoma, Sezary Syndrome, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, hypereosinophilic syndromes, mastocytosis and thrombocythemia.

Another embodiment of the invention is a method of using the compounds of formula (I) as standards or controls in *in vitro* or *in vivo* assays in determining the efficacy of test compounds in modulating SHIP1 activity.


In another embodiment of the invention, the compounds of formula (I) are isotopically-labeled by having one or more atoms therein replaced by an atom having a different atomic mass or mass number. Such isotopically-labeled (*i.e.*, radiolabelled) compounds of formula (I) are considered to be within the scope of this invention. Examples of isotopes that can be incorporated into the compounds of formula (I) include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine, chlorine, and iodine, such as, but not limited to, ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , and ^{125}I , respectively. These isotopically-labeled compounds would be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action for SHIP1 modulation, or binding affinity to pharmacologically important site of action for SHIP1 modulation. Certain isotopically-labeled compounds of formula (I), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, *i.e.* ^3H , and carbon-14, *i.e.*, ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

Substitution with heavier isotopes such as deuterium, *i.e.* ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances.


Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds of formula (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Examples as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

In other embodiments, compounds of formula (I) exclude the following compounds:


1. (1S,2S,4aS,8aR)-decahydro-1-[2-(3-methoxyphenyl)ethyl]-5,5,8a-trimethyl-2-naphthalenecarboxylic acid, having the following formula (a):




2. (1S,2S,4aS,8aR)-decahydro-1-[2-(3-methoxyphenyl)ethyl]-5,5,8a-trimethyl-2-naphthalenemethanol, having the following formula (b):




3. α -(3,5-dimethoxyphenyl)decahydro-2-hydroxy-2,5,5,8a-tetramethyl-1-naphthalenemethanol, having the following formula (c):




4. (4aS,8aS)-3,4,4a,5,6,7,8,8a-octahydro- α -(3-methoxy-5-methylphenyl)-2,5,5,8a-tetramethyl-1-naphthalenemethanol, having the following formula (d):




5. (1S,2R,4aS,8aS)-decahydro-2-hydroxy- α -(3-methoxy-5-methylphenyl)-2,5,5,8a-tetramethyl-1-naphthalenemethanol, having the following formula (e):




6. (1R,2R,4aS,8aS)-decahydro-1-[(3-methoxy-5-methylphenyl)methyl]-2,5,5,8a-tetramethyl-2-naphthalenol, having the following formula (f):




7. (4aS,5S,8aS)-decahydro-5-[2-(3-methoxyphenyl)ethyl]-1,1,4a-trimethyl-6-methylene-naphthalene, having the following formula (g):




8. 2-[(1S,4aS,8aS)-decahydro-5,5,8a-trimethyl-2-methylene-1-naphthalenyl]-1-(3-methoxyphenyl)-ethanone, having the following formula (h):




9. (1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol, having the following formula (i):




10. (1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol, having the following formula (j):




11. (1S,2R,4aS,8aS)-1-[3,5-bis(propan-2-yloxy)phenoxy]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol, having the following formula (k):




12. (1R,2R,8aS)-1-((3,5-dimethoxyphenylsulfonyl)methyl)-2,5,5,8a-tetramethyldecahydronaphthalen-2-ol, having the following formula (m):




13. (1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)sulfonyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol, having the following formula (n):



14. (1R,2R,4aS,8aS)-1-{[(3-methoxyphenyl)sulfanylmethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol, having the following formula (o):




15. (1R,2R, 4aS,8aS)-1-{[(3,5-dimethoxyphenyl)sulfanylmethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol, having the following formula (p):



In other embodiments, compounds of formula (I) exclude all isomeric forms of compounds (a) through (p) above.

In further embodiments, compounds of formula (Ia) exclude embodiments wherein R^1 is $-CH_2-$, $-(CH_2)_2-$, $-CH(OH)-$ and/or $-CH_2C(O)-$. In still further embodiments, compounds of formula (Ia) exclude compounds wherein one or both of R^2 and R^3 is alkoxy (e.g., wherein R^2 and R^3 are independently hydrogen, hydroxy or alkyl, provided that at least one of R^2 and R^3 is hydroxy).

In other embodiments, preferred stereochemistry of the compounds of formula (I) is shown below:



Specific embodiments of the compounds of the invention are described in more detail below in the Preparation of the Compounds of the Invention.

Utility and Testing of the Compounds of the Invention

Compounds of formula (I) above have activity as SHIP1 modulators and utility over a wide range of therapeutic applications, and may be used to treat any of a variety of diseases, disorders or conditions that would benefit from SHIP1 modulation. For example, such diseases, disorders or conditions include (but are not limited to) autoimmune diseases such as idiopathic pulmonary fibrosis, inflammatory bowel disease (including Crohn's Disease and ulcerative colitis), rheumatoid arthritis, Still's Disease, Sjögren's Syndrome, systemic lupus erythematosus, multiple sclerosis, psoriasis, and systemic sclerosis; inflammatory diseases such as acute respiratory distress syndrome (ARDS), allergic rhinitis, Alzheimer's Disease, asthma, ocular inflammatory diseases (including allergic conjunctivitis, dry eye, and uveitis), chronic obstructive pulmonary disease (COPD) including emphysematous, brochitic, and COPD due to alpha 1 anti-trypsin deficiency, atopic dermatitis, dermal contact hypersensitivity, eczema, eosiniphilic gastrointestinal disorder, fibromyalgia, gout, hepatic fibrosis, irritable bowl syndrome, bladder pain syndrome/interstitial cystitis, post operative inflammation, ischemic reperfusion disease, kidney fibrosis, pancreatitis, Parkinsons Disease, , seronegative spondyloarthropathies (including anklyosing spondylitis, psoriatic arthritis, and Reiter's Syndrome), and vasculitis (including Wegener's Granulomatosis, polyarteritis nodosa, leucocytoclastic vasculitis, Churg-Strauss Syndrome, cryoglobulinemic vasculitis and giant cell arteritis); and neoplastic diseases or other cell proliferative disorders such as acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, chronic

lymphocytic leukemia, basophilic leukemia, cutaneous T-cell lymphoma, Sezary Syndrome, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, hypereosinophilic syndromes, mastocytosis and thrombocythemia.

The effectiveness of a compound as a SHIP1 modulator may be determined by any number of known techniques, including the assays set forth in Examples 120-123.

Pharmaceutical Compositions of the Invention and Administration

For the purposes of administration, the compounds of the present invention may be formulated as pharmaceutical compositions. Pharmaceutical compositions comprise one or more compounds of this invention in combination with a pharmaceutically acceptable carrier and/or diluent. The compound is present in the composition in an amount which is effective to treat a particular disorder--that is, in an amount sufficient to achieve SHIP1 modulation activity, and preferably with acceptable toxicity to the patient. Typically, the pharmaceutical compositions of the present invention may include a compound in an amount from 0.1 mg to 250 mg per dosage depending upon the route of administration, and more typically from 1 mg to 60 mg. Appropriate concentrations and dosages can be readily determined by one skilled in the art.

Pharmaceutically acceptable carrier and/or diluents are familiar to those skilled in the art. For compositions formulated as liquid solutions, acceptable carriers and/or diluents include saline and sterile water, and may optionally include antioxidants, buffers, bacteriostats and other common additives. The compositions can also be formulated as pills, capsules, granules, or tablets which contain, in addition to a compound of this invention, diluents, dispersing and surface active agents, binders, and lubricants. One skilled in this art may further formulate the compounds in an appropriate manner, and in accordance with accepted practices, such as those disclosed in *Remington's Pharmaceutical Sciences* (Mack Pub. Co., N.J. current edition).

In another embodiment, the present invention provides a method for modulation SHIP1 generally and, more specifically, to treating the diseases, disorders and conditions as discussed above. Such methods include administering of a compound of the present invention to a mammal, preferably a human, in an amount sufficient to treat the condition. In this context, "treat" includes prophylactic administration. Such methods include systemic administration of a compound of this invention, preferably in the form of a pharmaceutical composition as discussed above. As used herein,

systemic administration includes oral and parenteral methods of administration. For oral administration, suitable pharmaceutical compositions include powders, granules, pills, tablets, and capsules as well as liquids, syrups, suspensions, and emulsions. These compositions may also include flavorants, preservatives, suspending, thickening and emulsifying agents, and other pharmaceutically acceptable additives. For parenteral administration, the compounds of the present invention can be prepared in aqueous injection solutions which may contain buffers, antioxidants, bacteriostats, and other additives commonly employed in such solutions.

Preparation of the Compounds of the Invention

The following Reaction Schemes illustrate methods to make compounds of the present invention, *i.e.*, compounds of formula (I), as set forth above in the Summary of the Invention.

The compounds of the present invention may be prepared by known organic synthesis techniques, including the methods described in more detail in the Examples. In general, the compounds of formula (I) may be made by the following Reaction Schemes, wherein all substituents are as defined above unless indicated otherwise. Although not generally depicted in the following schemes, one skilled in the art will understand that appropriate protecting group strategies may be useful in preparing compounds of formula (I). Protecting group methodology is well known to those skilled in the art (*see, for example*, Greene, T.W. and Wuts, P.G.M. *Greene's Protective Groups in Organic Synthesis* (2006), 4th Ed. Wiley).

It is also understood that one skilled in the art would be able to make the compounds of the invention by similar methods or by methods known to one skilled in the art. It is also understood that one skilled in the art would be able to make in a similar manner as described below other compounds of the invention not specifically illustrated below by using the appropriate starting components and modifying the parameters of the synthesis as needed. In general, starting components may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, etc. or synthesized according to sources known to those skilled in the art (*see, e.g.*, Smith, M.B. and J. March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 6th edition (Wiley, 2007)) or prepared as described herein.


It is also understood that in the following description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of this invention may not possess pharmacological activity as such, they may be administered to a mammal and thereafter metabolized in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds of this invention are included within the scope of the invention.

A. Preparation of Compounds of Formulae (Ia1a), (Ia1b), (Ia1c), (Ia1d), (Ia1e), (Ia1f) and (Ia1g)

Compounds of formulae (Ia1a), (Ia1b), (Ia1c), (Ia1d), (Ia1e), (Ia1f) and (Ia1g) are compounds of formula (Ia1), as set forth above in the Embodiments of the Invention, where R^{9b} is a direct bond, R^{4b} is $-R^9-OR^{11}$ (where R^9 is a direct bond and R^{11} is hydrogen), R^{4a} is hydrogen, alkyl or $-R^9-OR^{11}$ (where R^9 is an alkylene chain and R^{11} is as described above for R^{11} in the Summary of the Invention for compounds of formula (I)), and n , R^2 , R^3 , R^5 , R^6 , R^7 , R^8 , R^{9a} and R^{13} are as described above in the Summary of the Invention for compounds of formula (I), and are prepared below in Reaction Scheme 1:

REACTION SCHEME 1



Compounds of formula (Ia2a) are compounds of formula (Ia2), as described above in the Embodiments, where R^{9b} is a direct bond and one R¹⁰ is hydrogen and the other R¹⁰ is hydroxy, R^{4a} is -R⁹-OR¹¹ (where R⁹ is a direct bond and R¹¹ is hydrogen), R^{4b} is hydrogen, alkyl or -R⁹-OR¹¹ (where R⁹ is an alkylene chain and R¹¹ is as described above for R¹¹ in the Summary of the Invention for compounds of formula (I)), and m, n, R², R³, R⁵, R⁶, R⁷, R⁸, R^{9a} and R¹³ are as described above in the Summary of the Invention for compounds of formula (I), and may be purchased or prepared according to methods known in the art or by methods disclosed herein.


In general, oxidation of the benzylic alcohol of compound of formula (Ia2a) with an appropriate reagent, for example pyridinium chlorochromate (PCC), results in ketone compound of formula (Ia1a). Elimination of the tertiary alcohol of compound of

formula (1a1a) under appropriate conditions, for example Tin (IV) chloride, produces compounds of formula (1a1b), compounds of formula (1a1c) or compounds of formula (1a1d). Further treatment of compound of formula (1a1b) with an appropriate reagent, for example $\text{BH}_3\text{-THF}$ followed by NaOH , yields alcohol compound of formula (1a1e). Alcohol compound of formula (1a1e) may then be oxidized, for example with PCC, to yield ketone compound of formula (1a1f). The non-benzylic keto moiety of compound of formula (1a1f) may be partially reduced to the corresponding alcohol or completely reduced to the corresponding methylene using methods known in the art to produce a compound of formula (1a1g) wherein R^5 is hydrogen or hydroxy. Alternatively, a compound of formula (1a1f) may be treated under reductive alkylation or reductive amination conditions to produce a compound of formula (1a1g) wherein, for example, R^5 is optionally substituted amino, or converted to another functional group, such as a cyano, amide or carboxylic acid, using methods known in the art. One skilled in the art will recognize that, instead of starting with a compound comprising the R^6 , R^7 and R^8 moiety, the R^6 , R^7 and R^8 moiety may be installed at a later point in the synthesis using methods known to one skilled in the art.

B. Preparation of Compounds of Formulae (1a2b), (1a2c), (1a2d), (1a2e) and (1a2f)

Compounds of formulae (1a2b), (1a2c), (1a2d), (1a2e) and (1a2f) are compounds of formula (1a2), as set forth above in the Embodiments of the Invention, where R^{9b} is a direct bond and both R^{10} 's are hydrogen, R^{4b} is $-\text{R}^9\text{-OR}^{11}$ where R^9 is a direct bond and R^{11} is hydrogen, R^{4a} is hydrogen, alkyl or $-\text{R}^9\text{-OR}^{11}$ where R^9 is an alkylene chain and R^{11} is as described above for R^{11} in the Summary of the Invention for compounds of formula (I), R^5 is hydrogen, and m, n, R^2 , R^3 , R^6 , R^7 , R^8 , R^{9a} and R^{13} are as described above in the Summary of the Invention for compounds of formula (I), and are prepared below in Reaction Scheme 2:

REACTION SCHEME 2



Compounds of formula (1a2a) are described above in Reaction Scheme 1.


In general, complete reduction of the benzylic alcohol of compound of formula (1a2a) under conditions known in the art results in alcohol compound of formula (1a2b). Elimination of the tertiary alcohol of compound of formula (1a2b) under appropriate conditions, for example, Tin (IV) chloride, produces compounds of formulae (1a2c), (1a2d) or (1a2e). Further reduction of a compound of formulae (1a2c), (1a2d) or (1a2e) under appropriate conditions, for example catalytic hydrogenation or other suitable method, yields compounds of formula (1a2f). One skilled in the art will recognize that, instead of starting with a compound comprising the

R^5 , R^6 , R^7 and/or R^8 moiety, the R^5 , R^6 , R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

C. Preparation of Compounds of Formulae (Ia3a), (Ia3b), (Ia3c), (Ia3d) and (Ia3f)

Compounds of formulae (Ia3a), (Ia3b), (Ia3c), (Ia3d) and (Ia3e) are compounds of formula (Ia3), as set forth above in the Embodiments of the Invention, where t is 0, R^{9a} is straight or branched alkylene chain, R^{4b} is $-R^9-OR^{11}$ (where R^9 is a direct bond and R^{11} is hydrogen), R^{4a} is hydrogen, alkyl or $-R^9-OR^{11}$ (where R^9 is an straight or branched alkylene chain and R^{11} is as described above in the Summary of the Invention for compounds of formula (I)), R^5 is hydrogen and m , n , R^2 , R^3 , R^6 , R^7 , R^8 , R^{9b} and R^{13} are as described above in the Summary of the Invention for compounds of formula (I), and are prepared below in Reaction Scheme 3:

REACTION SCHEME 3



Compounds of formula (A) and formula (B) may be purchased or prepared by methods known in the art.


In general, activation of the primary alcohol in compound of formula (A), for example by conversion to the mesylate using methods known in the art, and reaction of the activated compound of formula (A) with the phenyl compound of formula (B) results in compound of formula (la3a). Compounds of formulae (la3b), (la3c) and (la3d) may be produced from compounds of formula (la3a) using methods analogous to those described above in Reaction Scheme 1, and compounds of formula (la3e)

may be produced from compounds of formulae (Ia3b), (Ia3c) and (Ia3d) using methods analogous to those described above in Reaction Scheme 1. One skilled in the art will recognize that compounds of formulae (Ia3a), (Ia3b), (Ia3c), (Ia3d) and (Ia3e) may be further oxidized under the appropriate oxidizing conditions to produce the corresponding compounds of formula (Ia3) where t is 1 or 2. One skilled in the art will also recognize that, instead of starting with a compound comprising the R⁵, R⁶, R⁷ and/or R⁸ moiety, the R⁵, R⁶, R⁷ and/or R⁸ moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

D. Preparation of Compounds of Formulae (Ia4a), (Ia4b), (Ia4c), (Ia4d) and (Ia4e)

Compounds of formulae (Ia4a), (Ia4b), (Ia4c), (Ia4d) and (Ia4e) are compounds of formula (Ia4), as set forth above in the Embodiments of the invention, where R^{9b} is a direct bond, R^{4b} is -R⁹-OR¹¹ (where R⁹ is a direct bond and R¹¹ is hydrogen), R^{4a} is hydrogen, alkyl or -R⁹-OR¹¹ (where R⁹ is an alkylene chain and R¹¹ is as described above in the Summary of the Invention for compounds of formula (I)), R⁵ is hydrogen and m, n, R², R³, R⁶, R⁷, R⁸, R^{9a} and R¹³ are as described above in the Summary of the Invention for compounds of formula (I), and are prepared below in Reaction Scheme 4 where W is a suitable leaving group, such as halogen:

REACTION SCHEME 4



Compounds of formula (A) and formula (C) may be purchased or prepared by methods known in the art or by methods disclosed herein.


In general, reaction of the primary alcohol in compound of formula (A) with the phenyl compound of formula (C) results in compound of formula (la4a). Compounds of formulae (la4b), (la4c) and (la4d) may be produced from compounds of formula (la4a) using methods analogous to those described above in Reaction Scheme 1. Compounds of formula (la4e) may be produced from compounds of formulae (la4b), (la4b) and (la4c) using methods analogous to those described above in Reaction Scheme 2. One skilled in the art will recognize that, instead of starting with a

compound comprising the R^5 , R^6 , R^7 and/or R^8 moiety, the R^5 , R^6 , R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

E. Preparation of Compounds of Formulae (Ia5a), (Ia5b), (Ia5c) and (Ia5d)

Compounds of formulae (Ia5a), (Ia5b), (Ia5c) and (Ia5d) are compounds of formula (Ia5), as set forth above in the Embodiments of the invention, where R^{11a} is hydrogen, R^{4b} is $-R^9-OR^{11}$ (where R^9 is a direct bond and R^{11} is hydrogen), R^{4a} is hydrogen, alkyl or $-R^9-OR^{11}$ (where R^9 is an alkylene chain and R^{11} is as described above for R^{11} in the Summary of the Invention for compounds of formula (I)), R^5 is hydrogen and m , n , R^2 , R^3 , R^6 , R^7 , R^8 , R^{9a} , R^{9b} and R^{13} are as described above in the Summary of the Invention for compounds of formula (I), and are prepared below in Reaction Scheme 5:

REACTION SCHEME 5




Compounds of formula (D) and formula (H) may be purchased or prepared by methods known in the art.

In general, elimination of the tertiary alcohol in the compound of formula (D) under appropriate conditions, for example with p-toluene sulphonic acid, followed by oxidation of the aldehyde to the corresponding carboxylic acid under conditions known in the art results in the formation of a compound of formulae (E), (F) or (G). Activation of the carboxylic acid in these compounds under appropriate conditions, for example with oxalic acid, followed by reaction with aniline compound of formula (H) results in a compound of formulae (la5a), (la5b) or (la5c). Compounds of formula (la5d) may then be produced using methods analogous to those described above in Reaction Scheme 2. One skilled in the art will recognize that, instead of starting with a compound comprising the R^5 , R^6 , R^7 and/or R^8 moiety, the R^5 , R^6 , R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

F. Preparation of Compounds of Formulae (la1h), (la1i), (la1j), (la1k) and (la1l)

Compounds of formulae (la1h), (la1i), (la1j), (la1k) and (la1l) are compounds of formula (la1), as set forth above in the Embodiments of the Invention, where R^{9a} is methylene, ethylene or propylene and R^{9b} is a direct bond, R^{4b} is $-R^9-OR^{11}$ (where R^9 is a direct bond and R^{11} is hydrogen), R^{4a} is hydrogen, alkyl or $-R^9-OR^{11}$ (where R^9 is an alkylene chain and R^{11} is as described above for R^{11} in the Summary of the Invention for compounds of formula (I)), R^5 is hydrogen and m, n, R^2 , R^3 , R^6 , R^7 , R^8 and R^{13} are as described above in the Summary of the Invention for compounds of formula (I), and are prepared below in Reaction Scheme 6:

REACTION SCHEME 6



Compounds of formula (J) and formula (K) may be purchased or prepared by methods known in the art.


Utilizing organometallic methodologies, the reaction of a compound of formula (J) with phenyllithium compound of formula (K) results in ketone compound of formula (la1h). Compounds of formulae (la1i), (la1j), (la1k) and (la1l) may then be produced using methods analogous to those described above in Reaction Scheme 2. The keto moiety of compounds of formulae (la1h), (la1i), (la1j), (la1k) and (la1l) may also be

partially reduced to the corresponding hydroxyl compound or completely reduced to a methylene to yield compounds of formula (Ia2) wherein one R¹⁰ is hydrogen and the other is hydroxy or wherein both R¹⁰'s are hydrogen, which may be further functionalized. Methods for such reduction are well known in the art. One skilled in the art will recognize that, instead of starting with a compound comprising the R⁵, R⁶, R⁷ and/or R⁸ moiety, the R⁵, R⁶, R⁷ and/or R⁸ moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

G. Preparation of Compounds of Formulae (Ia1f2), (Ia1g1), (Ia1g2), (Ia1g3) and (Ia1g4)

Compound of formula (Ia1f2) are compounds of formula (Ia1f), as described above in Reaction Scheme 1, and compounds of formulae (Ia1g1), (Ia1g2), (Ia1g3) and (Ia1g4) are compounds of formula (Ia1g), as described above in Reaction Scheme 1, and are prepared below in Reaction Scheme 7 where R^{4a1} is alkyl, R^{11b} is optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl, and X is halo:

REACTION SCHEME 7



Compounds of formula (la1f1) are compounds of formula (la1f) as described above in Reaction Scheme 1, and can be prepared by methods disclosed herein.


In general, compounds of formula (la1g4) are prepared by first treating a compound of formula (la1f1) with acid or base isomerization conditions to give a

compound of formula (Ia1f2). Carbonyl homologation of the compound of formula (Ia1f2) under standard Wittig olefination conditions followed by acid hydrolysis affords a compound of formula (Ia1g2). Nucleophilic addition to the aldehyde carbonyl of the compound of formula (Ia1g2) yields a compound of formula (Ia1g3), which is then treated under standard oxidation conditions to give a compound of formula (Ia1g4).

H. Preparation of Compounds of Formulae (Ia1g5), (Ia1g6), (Ia1g7) and (Ia1g8)

Compounds of formulae (Ia1g5), (Ia1g6), (Ia1g7) and (Ia1g8) are compounds of formula (Ia1g), as described above in Reaction Scheme 1, and are prepared as described below in Reaction Scheme 8 where R^{4a1} is alkyl:

REACTION SCHEME 8



Compounds of formula (la1f2) are described above in Reaction Scheme 7 and are prepared by methods disclosed herein.


In general, compounds of formula (la1g8) are prepared by first treating a compound of formula (la1f2) under standard Wittig reaction conditions to form the

methyldene compound of formula (Ia1g5). Hydroboration of the methyldene of the compound of formula (Ia1g5) with oxidative work up affords the primary alcohol compound of formula (Ia1g6). In addition to direct methods, the alcohol of the compound of formula (Ia1g5) is converted to an azide compound of formula (Ia1g7) by first activation of the hydroxyl by standard conditions, such as treatment with methanesulfonyl chloride, followed by treatment with sodium azide under standard conditions. Reduction of the azide of the compound of formula (Ia1g7), for example using Staudinger conditions, affords a compound of formula (Ia1g8).

I. Preparation of Compounds of Formulae (Ia-2) and (Ia-3)

Compounds of formulae (Ia-2) and (Ia-3) are compounds of formula (Ia), as described above in the Embodiments of the Invention where R^5 is $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$ (where the first R^9 , R^{11} and R^{12} are as described above in the Summary of the Invention for compounds of formula (I) and the second R^9 is a direct bond), and are prepared as described below in Reaction Scheme 9 where n , R^1 , R^2 , R^3 , R^{4a} , R^{4b} , R^6 , R^7 , R^8 , R^{11} and R^{12} are as described above in the Summary of the Invention for compounds of formula (I):

REACTION SCHEME 9



Compounds of formula (la-1) are compounds of formula (la) as described herein and are prepared by the methods described herein or by methods known to one skilled in the art. Compounds of formula (M) can be prepared by methods known to one skilled in the art. Compounds of formula (N) are commercially available or can be prepared by the methods known to one skilled in the art.


In general, compounds of formula (la-2) are prepared by reaction of the amine compound of formula (la-1) with the anhydride activated compound of formula (M) under standard conditions.

In general, compounds of formula (la-3) are prepared by reaction of the amine compound of formula (N) with carbonyl diimidazole under standard conditions, followed by reaction of the resulting intermediate with the amine compound of formula (la-1).

J. Preparation of Compounds of Formula (Ia3f)

Compounds of formula (Ia3f) are compounds of formula (Ia3), as described above in the Embodiments of the Invention where R^5 is $-R^9-N(R^{11})R^{12}$ (where R^9 is a direct bond and R^{11} and R^{12} are as described above in the Summary of the Invention for compounds of formula (I)), and are prepared as described below in Reaction Scheme 10 where R^{23} is an acyl group such as acetyl, R^5 is hydrogen, each R^{13} is hydrogen, R^{9a} is straight or branched alkylene chain, and n , R^2 , R^3 , R^6 , R^7 , R^8 and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I):

REACTION SCHEME 10




Compounds of formulae (O) and (B) may be prepared by methods disclosed herein or by methods known to one skilled in the art.

In general, compounds of formula (Ia3f) are prepared by the method shown above in Reaction Scheme 10 by first treating a compound of formula (O) (where ----- indicates an optional double bond), under standard epoxidation conditions, such as treatment with a suitable oxidizing agent in an aprotic solvent, to form a (5:1) mixture of a compound of formula (P) and a compound of formula (Q). Activation of the primary alcohol in the compounds of formulae (P) and (Q), for example by conversion to the mesylate using methods known in the art, and reaction of the activated compounds of formulae (P) and (Q) with the phenyl compound of formula (B) results in compound of formula (R). Treatment of the compound of formula (R) with sodium azide under standard conditions produces the compound of formula (S), which is then reduced under standard conditions to form the compound of formula (Ia3f). One skilled in the art will recognize that, instead of starting with a compound comprising the R^6 , R^7 and/or R^8 moiety, the R^6 , R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein. Furthermore, one skilled in the art will recognize that the compound of formula (Ia3f) may be further functionalized to produce compounds of formula (Ia3) wherein R^5 is $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2), where each R^9 , R^{11} , R^{12} and R^{14} are as described above in the Summary of the Invention for compounds of formula (I). It is also understood that other compounds of formula (I) where R^5 is $-R^9-N(R^{11})R^{12}$ (where R^9 is a direct bond and R^{11} and R^{12} are as described above in the Summary of the Invention for compounds of formula (I)) may be prepared by this method utilizing the appropriate starting materials.

K. Preparation of Compounds of Formulae (Ia-4)

Compounds of formula (Ia-4) are compounds of formula (Ia), as described above in the Embodiments of the Invention where R^{4a} and R^{4b} together form $=C(R^{13})_2$ (where each R^{13} is hydrogen), and are prepared as described below in Reaction Scheme 11 where n , R^2 , R^3 , R^6 , R^7 , R^8 , R^{9a} and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I):

REACTION SCHEME 11




Compounds of formula (la1b1) are compounds of formula (lab1) and are prepared as described herein or by methods described by one skilled in the art.

In general, compounds of formula (la-4) are prepared by the method disclosed in the Reaction Scheme 11 by first treating a compound of formula (la1b1) with a suitable oxidizing agent in an aprotic solvent to form a compound of formula (T). The compound of formula (T) is then treated with acid under suitable conditions to promote rearrangement of the epoxide to form a compound of formula (la-4). One skilled in the art will recognize that, instead of starting with a compound comprising the R^2 , R^3 , R^6 , R^7 and/or R^8 moiety, the R^2 , R^3 , R^6 , R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

L. Preparation of Compounds of Formula (Ia5d1)

Compounds of formula (Ia5d1) are compounds of formula (Ia5), as described above in Reaction Scheme 5 where R^{4a} is methyl and R^{4b} is hydrogen, and are prepared as described below in Reaction Scheme 12 where n , R^2 , R^3 , R^6 , R^7 , R^8 , R^{9a} and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I) where Ac is an acyl group, such as acetyl:

REACTION SCHEME 12



Compounds of formulae (U) and (H) may be prepared by one skilled in the art by methods known to one skilled in the art or by methods disclosed herein.


In general, compound of formula (Ia5d1) are prepared by first treating a compound of formula (U) under standard reduction conditions to form a compound of formula (V). The primary hydroxyl in the compound of formula (V) is then protected, for

example, with an acyl group, such as acetyl, under standard conditions to form the compound of formula (W). Dehydration of the tertiary alcohol in the compound of formula (W) affords the methyldene compound of formula (X). Reduction of the methyldene of the compound of formula (X) under standard hydrogenation conditions followed by removal of the hydroxyl protecting group affords a compound of formula (Y). Oxidation of the primary alcohol of the compound of formula (Y) under standard conditions yields the carboxylic acid compound of formula (Z). The compound of formula (Z) undergoes standard amide coupling with a compound of formula (H) to afford a compound of formula (la5d1).

M. Preparation of Compounds of Formulae (la2g1), (la2g2) and (la2g3)

Compounds of formulae (la2g1), (la2g2) and (la2g3) are compounds of formula (la2g), as described above in Reaction Scheme 2 where R^5 is hydrogen, and are prepared as described below in Reaction Scheme 13 where n , R^2 , R^3 , R^6 , R^7 , R^8 , R^{9a} and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I):

REACTION SCHEME 13



Compounds of formula (la2d1) are compounds of formula (la2d), as described above in Reaction Scheme 2 where each R¹³ is hydrogen and R⁵ is hydrogen, and may be prepared by methods disclosed herein or by methods known to one skilled in the art.

In general, compounds of formulae (la2g1), (la2g2) and (la2g3) are prepared by the method shown above in Reaction Scheme 13 by first treating a compound of formula (la2d1) under hydroboration conditions with oxidative work up. The resulting primary alcohol compound of formula (la2g1) can be selectively oxidized to the aldehyde compound of formula (la2g2) and can be further oxidized to the carboxylic acid compound of formula (la2g3).


N. Preparation of Compounds of Formula (I-1)

Compounds of formula (I-1) are compounds of formula (I), as described above in the Summary of the Invention where R¹ is -R^{9a}-N(R^{11a})C(O)-R^{9b}- (where R^{11a} is hydrogen and R^{9a} and R^{9b} are as described above in the Summary of the Invention for

compounds of formula (I)), and are prepared as described below in Reaction Scheme

14 where , n, R⁷, R⁸, R^{9a} and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I), R^{4a} is methyl, R²⁴ is R¹², -R⁹-N(R¹¹)R¹², -N(R¹¹)-OR¹², -R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2 and each R⁹, R¹¹ and R¹² is as described above in the Summary of the Invention), Pg¹ is an oxygen protecting group, such as *tert*-butyldimethylphenylsilyl, and Pg² is a nitrogen protecting group, such as *para*-methoxybenzyl:

REACTION SCHEME 14




Compounds of formula (AA) may be prepared according to the methods disclosed in PCT Published Patent Application No. WO 2007/147251 or by methods known to one skilled in the art. Compounds of formula (II) and (NN) are commercially





available or can be prepared by methods known to one skilled in the art or by methods disclosed herein.

In general, compounds of formula (I-1) are prepared by the method disclosed above in Reaction Scheme 14 by first treating a compound of formula (AA) under standard dehydration conditions, for example, Tin (IV) chloride, to afford a compound of formula (BB). Standard reduction of the aldehyde of the compound of formula (BB) affords a primary alcohol compound of formula (CC). The hydroxyl is protected under standard conditions to form a compound of formula (DD), then the double bond in the compound of formula (DD) is hydroborated with oxidative work up to give the secondary alcohol compound of formula (EE). The hydroxyl in the compound of formula (EE) is oxidized to the ketone compound of formula (FF), which undergoes reductive amination with a suitably protected amine to give a compound of formula (GG). The protecting group is removed from the amine to give a compound of formula (HH) which undergoes amide coupling chemistry with a carboxylic acid compound of formula (II) to afford an amide compound of formula (JJ). The hydroxyl protecting group is removed to give a compound of formula (KK). In addition to direct methods, the hydroxyl of the compound of formula (KK) is converted to an azide compound of formula (LL) by first activation of the hydroxyl by standard conditions, such as treatment with methanesulfonyl chloride, followed by treatment with sodium azide under standard conditions. Reduction of the azide of the compound of formula (LL), for example, using Staudinger conditions, affords a compound of formula (MM). Amide compounds of formula (I-1) are prepared by amide coupling chemistry between the amine compounds of the formula (MM) with carboxylic acid compounds of the formula (NN).


O. Preparation of Compounds of Formula (I-2)

Compounds of formula (I-2) are compounds of formula (I), as described above in the Summary of the Invention where R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$ (where R^{9a} is as defined above in the Summary of the Invention for compounds of formula (I), each R^{10} is hydrogen and R^{9b} is a directed bond), and are prepared as described below in



, , ,  or , and n, R^{4a}, R⁷, R⁸ and R^{9a} are as described above in the Summary of the Invention for compounds of formula (I), and R²⁴ is R¹², -R⁹-N(R¹¹)R¹², -N(R¹¹)-OR¹², or -R⁹-N(R¹¹)S(O)_pR¹² (where p is 1 or 2 and each R⁹, R¹¹ and R¹² is as described above in the Summary of the Invention):

REACTION SCHEME 15



Compounds of formula (KK) may be prepared by methods known one skilled in the art or by methods described herein. Compounds of formula (PP) are commercially available.

In general, compounds of formula (I-2) are prepared by the method described above in Reaction Scheme 15 by first activating the hydroxyl of the compound of formula (KK) by treating the compound of formula (KK) with methanesulfonyl chloride (MsCl) under standard conditions to form the compound of formula (OO). The compound of formula (OO) is then treated with a compound of formula (PP) under


nucleophilic displacement conditions in polar aprotic solvent to form a compound of formula (I-2). One skilled in the art will recognize that, instead of starting with a compound comprising the R^7 and/or R^8 moiety, the R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

P. Preparation of Compounds of Formula (I-3)

Compounds of formula (I-3) are compounds of formula (I), as described above in the Summary of the Invention where R^1 is $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$ (where R^{11a} is hydrogen and R^{9a} and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I)), and are prepared as described below in Reaction Scheme

16 where $\textcircled{A}$, n , R^7 , R^8 , R^{9a} and R^{9b} are as described above in the Summary of the Invention for compounds of formula (I), R^{4a} is methyl, and R^{24} is R^{12} , $-R^9-N(R^{11})R^{12}$, $-N(R^{11})-OR^{12}$, or $-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2 and each R^9 , R^{11} and R^{12} is as described above in the Summary of the Invention):

REACTION SCHEME 16




Compounds of formula (OO) may be prepared as described above in Reaction Scheme 16 or by methods known to one skilled in the art. Compounds of formula (NN) are commercially available or can be prepared by methods known to ones skilled in the art.

In general, compounds of formula (I-3) are prepared by the method disclosed above in Reaction Scheme 16 by first treating a compound of formula (OO) under nucleophilic displacement conditions with potassium cyanide in polar aprotic solvent to form a compound of formula (QQ), which is then reduced under standard reduction conditions to form the compound of formula (RR). The compound of formula (RR) is then treated with the compound of formula (NN) under standard amide coupling conditions to form the compound of formula (I-3). One skilled in the art will recognize that, instead of starting with a compound comprising the R^7 and/or R^8 moiety, the R^7 and/or R^8 moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

Q. Preparation of Compounds of Formula (Ia7a)

Compounds of formula (Ia7a) are compounds of formula (Ia7), as described above in the Embodiments of the Invention, where R^{9a} is methylene and n , R^2 , R^3 , R^{4a} , R^{4b} , R^6 , R^7 , R^8 , R^{9b} and R^{11a} are as described above in the Embodiments of the Invention for compounds of formula (Ia7), and are prepared below in Reaction Scheme 17:

REACTION SCHEME 17



Compounds of formula (SS) and formula (UU) may be purchased or prepared by methods known in the art or by methods disclosed herein.


In general, compounds of formula (Ia7a) are prepared by first oxidizing a compound of formula (SS) with an appropriate reagent, for example, pyridinium chlorochromate, to form a compound of formula (TT). Treatment of the compound of formula (TT) with a compound of formula (UU) under standard reductive amination conditions results in a compound of formula (Ia7a). One skilled in the art will recognize that, instead of starting with a compound comprising the R², R³, R⁵, R⁶, R⁷ and R⁸, the R², R³, R⁵, R⁶, R⁷ and R⁸ moiety may be installed at a later point in the synthesis using methods analogous to those described herein.

R. Preparation of Compounds of Formulae (Ia5e), (Ia5f) and (Ia5g)

Compounds of formulae (Ia5e), (Ia5f) and (Ia5g) are compounds of formula (Ia5), as described above in the Embodiments of the Invention, where R^{9a} is an

alkylene chain and n , R^2 , R^3 , R^{4a} , R^6 , R^7 , R^8 , R^{9b} and R^{11a} are as described above in the Embodiments of the Invention for compounds of formula (Ia5), and are prepared below in Reaction Scheme 18:

REACTION SCHEME 18



Compounds of formula (U) and formula (H) may be purchased or prepared by methods known in the art or by methods disclosed herein.

In general, compounds of formulae (Ia5e), (Ia5f) and (Ia5g) are prepared by utilizing organometallic methodologies by first treating a compound of formula (H) with an organometallic compound, such as trimethylaluminum and then treating the resulting product with a compound of formula (U) under suitable conditions to yield a compound of formula (Ia5e). The compound of formula (Ia5e) is then treated with a brominating agent to yield a compound of formula (Ia5f), which is then treated under standard elimination conditions to yield a compound of formula (Ia5g).

All of the compounds described herein as being prepared which may exist in free base or acid form may be converted to their pharmaceutically acceptable salts by treatment with the appropriate inorganic or organic base or acid. Salts of the compounds prepared below may be converted to their free base or acid form by standard techniques. Furthermore, all compounds of the invention which contain an acid or an ester group can be converted to the corresponding ester or acid, respectively, by methods known to one skilled in the art or by methods described herein.

Representative compounds of the invention which were prepared by the methods disclosed herein include (but are not limited to) the compounds listed below in Table 1.

TABLE 1

Cpd. No.	Compound Name	Ex. #
2	(1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	1
3	[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone	2
4	3-{[(4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl}-5-methylphenol	3
5	3-{[(4aS,8aS)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl}-5-methylphenol	3
6	(2R,3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol	4
7	(3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one	5
8	(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one	6
9	(2R,3R,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol	7
11	(3S,4S,4aS,8aS)-N-(3-aminopropyl)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine dihydrochloride	8
13	(4aS,8aS)-8-[(3-methoxy-5-methylphenyl)methyl]-4,4,7,8a-tetramethyl-1,2,3,4,4a,5,6,8a-octahydronaphthalene	9

Cpd. No.	Compound Name	Ex. #
14	(4aS,5S,8aS)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene	9
15	(4aS,5S,8aS)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene	9
16	[(1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol	10
17	3-([(1S,2S,4aS,8aR)-2-(hydroxymethyl)-5,5,8a-trimethyl-decahydronaphthalen-1-yl]methyl)-5-methylphenol	11
19	(1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid	12
20	(1S,2S,4aS,8aR)-1-[(3-hydroxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid	13
22	3-([(1S,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl)-5-methylphenol hydrochloride	14
23	(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol	15
27	(1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxy)methyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	16
29	3-([(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy)-5-methylphenol	17
30	(4S,4aS,8aS)-4-(3-methoxy-5-methylphenoxy)methyl)-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol	18
33	(1S,2R,4aS,8aS)-1-[3,5-bis(benzyloxy)-phenoxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	19
34	(1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxy)methyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	20
37	(1S,2R,4aS,8aS)-1-[3,5-bis(propan-2-yloxy)phenoxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	21
38	5-([(1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]methoxy}benzene-1,3-diol	22

Cpd. No.	Compound Name	Ex. #
41	(4aS,8aS)-N-(3,5-dimethoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide	23
42	(4aS,8aS)-N-(3-hydroxy-5-methoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide	24
43	(4aS,8aS)-N-(3,5-dihydroxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide	24
45	2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one	25
46	2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one	26
47	2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3-hydroxy-5-methoxyphenyl)ethan-1-one	27
48	2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dihydroxyphenyl)ethan-1-one	27
49	(1S,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)(hydroxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	28
50	(1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	29
51	[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone	30
52	(2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol	31
53	(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one	32
54	(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine	33
55	(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine	34
56	5-{[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	35

Cpd. No.	Compound Name	Ex. #
57	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide	36
58	5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-hydroxyethyl}benzene-1,3-diol	37
59	2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-ol	38
61	(4aR,5S,6S,8aS)-5-[2-(3,5-dimethoxyphenyl)ethyl]-1,1,4a,6-tetramethyl-decahydronaphthalene	39
62	5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethyl}benzene-1,3-diol	40
63	(4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbonitrile	41
64	(3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide	42
65	(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide	43
67	(3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid	44
68	(3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid	45
75	(1R,2R,4aS,8aS)-1-{[(3-methoxy-5-methylphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	46
76	(1R,2R,4aS,8aS)-1-{[(3-methoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	47
80	(1R,2R,4aS,8aS)-1-{[(3,5-dimethoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	48
81	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide	49

Cpd. No.	Compound Name	Ex. #
82	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide	50
83	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide	51
84	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide	52
85	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide	53
86	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide	54
87	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide	55
88	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide	56
89	N-[(3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyrazine-2-carboxamide	57
94	2-[(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbonyl}pyridine	58
95	5-[(1S,2R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-yl)carbonyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	59
102	N-[(2R,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide	60
103	N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide	61
109	N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}pyridine-3-carboxamide	62
111	N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}benzamide	63

Cpd. No.	Compound Name	Ex. #
113	5-{[(1S,2R,3S,4aS,8aS)-3-(1H-imidazol-1-ylmethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	64
115	5-{[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-(1H-1,2,4-triazol-1-ylmethyl)-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	65
117	5-{[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(4-methylpiperazin-1-yl)methyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol dihydrochloride salt	66
118	3-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea	67
119	5-{[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-5-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	68
120	1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-ethylurea	69
121	1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methoxyurea	70
122	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]benzamide	71
123	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-3-carboxamide	72
124	1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylurea	73
125	5-{[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(1H-pyrrol-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	74
126	5-{[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	75
127	5-{[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-4-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	76
132	(1R,2R,3R,4aS,8aS)-3-amino-1-{[(3,5-dimethoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	77

Cpd. No.	Compound Name	Ex. #
133	5-{[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-2-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol	78
135	(2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine	79
136	3-{[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}-5-methylphenol	80
137	3-{[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-5-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}-5-methylphenol	81
138	3-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea	82
139	N-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide	83
141	(2R,4R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-4a,8,8-trimethyl-3-methylidene-decahydronaphthalen-2-ol	84
142	(2R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-2-ol	85
149	2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)acetamide	86
157	(1S,2S,4aS,8aR)-1-[(3,5-dihydroxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid	87
158	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-2-carboxamide	88
159	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-4-carboxamide	89
160	3-{[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl}pyridin-1-ium-1-olate	90
163	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylpiperazine-1-carboxamide	91

Cpd. No.	Compound Name	Ex. #
165	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-aminopropanamide	92
167	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]piperidine-4-carboxamide	93
168	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-3-carboxamide	94
170	5-[[[(1S,2S,3R,4aS,8aS)-3-[(3-aminopropyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol dihydrochloride	95
171	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-6-aminopyridine-3-carboxamide	96
173	(4S,4aS,8aS)-4-[[[(3,5-dimethoxybenzene)sulfonyl]methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol	97
175	1-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]guanidine	98
176	N-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide	99
178	1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]guanidine	100
179	N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide	101
180	1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylthiourea	102
181	2-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl}methyl)sulfamoyl]benzoic acid	103
196	N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-((4-(trifluoromethyl)benzamido)methyl)-decahydronaphthalen-2-yl)nicotinamide	104
197	N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide	105


Cpd. No.	Compound Name	Ex. #
198	N-((2R,3S,4S,4aS)-4-(isonicotinamidomethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide	106
199	N-(((1S,2S,3R,8aS)-2,5,5,8a-tetramethyl-3-(nicotinamido)decahydronaphthalen-1-yl)methyl)nicotinamide	107
200	N-((2R,3S,4S,4aS)-4-((1H-imidazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide	108
201	N-((2R,3S,4S,4aS)-4-((1H-1,2,4-triazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide	109
202	N-((2R,3S,4S,4aS)-4-((1H-benzo[d]imidazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide	110
208	4-methyl-N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-((4-(trifluoromethyl)benzamido)methyl)-decahydronaphthalen-2-yl)piperazine-1-carboxamide	111
209	N-((2R,3S,4S,4aS)-4-((4-fluorobenzamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide	112
210	N-((2R,3S,4S,4aS)-4-(isonicotinamidomethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide	113
211	4-methyl-N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-(nicotinamidomethyl)decahydronaphthalen-2-yl)piperazine-1-carboxamide	114
212	N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide	115
215	4-methyl-N-((2R,3S,4S,4aR)-3,4a,8,8-tetramethyl-4-(2-(4-(trifluoromethyl)benzamido)ethyl)-decahydronaphthalen-2-yl)piperazine-1-carboxamide	116
216	N-((2R,3S,4S,4aR)-4-(2-(4-fluorobenzamido)ethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide	117
217	N-((2R,3S,4S,4aR)-4-(2-(furan-2-carboxamido)ethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide	118
218	4-methyl-N-((2R,3S,4S,4aR)-3,4a,8,8-tetramethyl-4-(2-(nicotinamido)ethyl)decahydronaphthalen-2-yl)piperazine-1-carboxamide	119

Cpd. No.	Compound Name	Ex. #
220	3-{[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy}-5-methoxyphenol.	120
223	(2R,4aS,8aS)-1-{[(3-hydroxy-5-methoxyphenyl)amino]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol	121
225	2-[(1S,2R,3R,4aS,8aR)-3-bromo-2,5,5,8a-tetramethyl-decahydro-naphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide	122
226	2-[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide	122

The following Examples are provided for purposes of illustration, not limitation. In summary, the following Examples disclose the synthesis of representative compounds of this invention and compounds used in the preparation of compounds of the invention, as well as representative assays for the same.

EXAMPLE 1

Synthesis of (1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 2)




To a solution of (1S,2R,4aS,8aS)-1-[hydroxy(3-methoxy-5-methylphenyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 1, prepared according to Andersen, R. et al. PCT Int. Patent Appl. WO 2007/147251 A1, 5.00 g, 13.9 mmol) in CH₂Cl₂ (50 mL) under argon was added pyridinium chlorochromate (PCC) (3.32 g, 15.4 mmol) as a solid in one portion and the mixture stirred at room temperature for 1.5 h. The reaction was concentrated to ~10-15 mL and filtered through a pad of Celite and silica gel, washing with Et₂O (500 mL). The filtrate was concentrated and purified by column chromatography through a plug of silica gel (Et₂O) to afford (1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)carbonyl]-

2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 2, 4.44 g, 88%) as a yellow foam. ^1H NMR (CDCl_3): δ 7.40 (s, 1H), 7.31 (s, 1H), 6.88 (s, 1H), 3.83 (s, 3H), 3.56 (s, 1H), 2.39 (s, 3H), 1.88 (dt, $J = 12, 3.6$ Hz, 1H), 1.79-0.96 (m, 17H), 0.90 (s, 3H), 0.83 (s, 3H).

EXAMPLE 2


Synthesis of [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone (Compound No. 3)



To a solution of (1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)-carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 2, 0.49 g, 1.37 mmol) in CH_2Cl_2 (20 mL) at 0°C under argon was added SnCl_4 (0.48 mL, 4.1 mmol) and the reaction stirred at 0°C for 10 min. The mixture was warmed to room temperature and diluted with CH_2Cl_2 (40 mL) and water (30 mL). The aqueous layer was extracted with CH_2Cl_2 (2 x 20 mL) and the combined organic extracts washed with saturated aqueous NaHCO_3 (1 x 40 mL), dried (Na_2SO_4) and concentrated. The resulting crude brown oil was purified by column chromatography on silica gel (hexanes/EtOAc, 9:1) to give [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone (Compound No. 3, 0.35 g, 75%) as a clear oil. ^1H NMR (CDCl_3): δ 7.33 (s, 1H), 7.24 (m, 1H), 6.91 (m, 1H), 5.61 (m, 1H), 4.12 (m, 1H), 3.85 (s, 3H), 2.40 (s, 3H), 2.12-1.92 (m, 2H), 1.55-1.37 (m, 6H), 1.30-1.14 (m, 4H), 0.96 (s, 3H), 0.91 (s, 3H), 0.90 (s, 3H).

EXAMPLE 3


Synthesis of 3-[[[(4a*S*,8a*S*)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl]-5-methylphenol (Compound No. 4) and 3-[[[(4a*S*,8a*S*)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl]-5-methylphenol (Compound No. 5)



To a microwave (MW) vessel (2-5 mL) was added Na₂S (109 mg, 1.40 mmol) and a solution of [(1*S*,4a*S*,8a*S*)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone (Compound No. 3, 98 mg, 0.29 mmol) in NMP (1.5 mL). The reaction vessel was sealed and placed in a Biotage Microwave Initiator reactor and irradiated at 180°C for 30 min. The reaction mixture was allowed to cool to room temperature and then quenched with 10% aqueous HCl (5 mL) and diluted with EtOAc (20 mL). The organic layer was washed with brine (3 x 10 mL), dried (Na₂SO₄) and concentrated. The crude product was purified by flash chromatography on silica gel (hexanes/EtOAc, 85:15) to afford a mixture of 3-[[[(4a*S*,8a*S*)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl]-5-methylphenol and 3-[[[(4a*S*,8a*S*)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]carbonyl]-5-methylphenol (Compound No. 4 and Compound No. 5 respectively, 71 mg, 78%) as a white foam. ¹H NMR (CDCl₃): δ 7.40-7.20 (m, total 3H), 6.93-6.86 (m, total 1H), 5.72-5.61 (m, total 1H), 2.38-2.35 (m, total 3H), 2.17-1.52 (m, total 3H), 1.40-1.08 (m, 12H), 0.95-0.86 (m, total 7H). ES-MS *m/z* 327 ([M+1]⁺).

EXAMPLE 4


Synthesis of (2R,3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 6)



To a solution of [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone (Compound No. 3, 2.26 g, 6.65 mmol) in anhydrous THF (70 mL) at 0°C under argon was added a solution of BH₃-THF (1.0 M in THF, 10.0 mL, 10.0 mmol) and the reaction stirred at 0°C for 1.5 h before warming to room temperature and stirring for another 30 min. The reaction was then quenched carefully at 0°C with MeOH (2 mL), 15% aqueous NaOH (2 mL) and 50% aqueous H₂O₂ (2 mL) and stirred at room temperature for 30 min. The mixture was diluted with EtOAc (50 mL), water (40 mL) and saturated aqueous sodium thiosulfate (30 mL) and stirred for another 15 min. The layers were separated and the organic layer washed with brine (1 x 40 mL). The combined aqueous layers were extracted with EtOAc (1 x 30 mL) and the organic extracts dried (Na₂SO₄) and concentrated to afford a white foam (2.50 g). The crude product was purified by column chromatography on silica gel (hexanes/EtOAc, 3:1) to give (2R,3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 6, 1.60 g, 67%) as a white solid. ¹H NMR (CDCl₃): δ 7.32 (s, 1H), 7.26-7.24 (m, 1H), 6.90-6.87 (m, 1H), 3.96 (q, *J* = 2.1 Hz, 1H), 3.82 (s, 3H), 3.72 (d, *J* = 4.5 Hz, 1H), 2.37 (s, 3H), 2.16-2.06 (m, 1H), 1.85-1.52 (m, 7H), 1.46-1.18 (m, 3H), 1.36 (s, 3H), 0.99 (d, *J* = 8.1 Hz, 3H), 0.89 (s, 3H), 0.86 (s, 3H). ES-MS *m/z* 359 ([M+1]⁺).

EXAMPLE 5


Synthesis of (3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 7)



To a solution of (2R,3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 6, 1.05 g, 2.93 mmol) in CH_2Cl_2 (25 mL) under argon was added PCC (782 mg, 3.63 mmol) and the mixture stirred for 1 h. TLC analysis showed some remaining starting alcohol so another portion of PCC (178 mg, 0.830 mmol) was added and the mixture stirred another 1 h. The reaction was concentrated, diluted with Et_2O (150 mL), filtered and re-concentrated. The resultant brown oil was purified by column chromatography on silica gel (hexanes/ EtOAc , 3:1) to afford (3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 7, 0.93 g, 89%) as a pale yellow oil. ^1H NMR (CDCl_3): δ 7.23 (s, 1H), 7.20 (m, 1H), 6.91 (m, 1H), 3.83 (s, 3H), 3.63 (d, $J = 6.9$ Hz, 1H), 2.82 (m, $J = 7.5$ Hz, 1H), 2.60-2.41 (m, 2H), 2.38 (m, 3H), 2.17 (s, 1H), 1.87 (m, 1H), 1.72-1.40 (m, 4H), 1.35 (s, 2H), 1.27-1.08 (m, 5H), 0.90 (s, 6H).

EXAMPLE 6


Synthesis of (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 8)



To a microwave (MW) vessel (2-5 mL) was added Na₂S (65 mg, 0.83 mmol) and a solution of (3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 7, 59 mg, 0.17 mmol) in NMP (1.5 mL). The reaction vessel was sealed and placed in a Biotage Microwave Initiator reactor and irradiated at 180°C for 30 min. The reaction mixture was allowed to cool to room temperature and then quenched with 10% aqueous HCl (5 mL) and diluted with EtOAc (20 mL). The organic layer was washed with brine (3 x 10 mL), dried (Na₂SO₄) and concentrated. The crude product was purified by flash chromatography on silica gel (hexanes/EtOAc, 4:1 then 7:3) to afford (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 8, 25 mg, 44%) a white solid. ¹H NMR (CDCl₃): δ 7.16 (s, 1H), 7.09 (s, 1H), 6.85 (s, 1H), 3.30 (d, *J* = 12 Hz, 1H), 3.04-2.93 (m, 1H), 2.50-2.43 (m, 1H), 2.42 (d, *J* = 12 Hz, 1H), 2.34 (s, 3H), 1.52-1.06 (m, 8H), 1.21 (s, 3H), 0.88 (d, *J* = 6.3 Hz, 3H), 0.87 (s, 3H), 0.86 (s, 3H). ES-MS *m/z* 343 ([M+1]⁺).

EXAMPLE 7


Synthesis of (2R,3R,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 9)



methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 9, 23 mg, 37%) as a mixture of diastereomers (white solid). ^1H NMR (CDCl_3): δ 7.40-7.20 (m, total 2H), 6.92-6.83 (m, total 1H), 4.10-3.25 (m, total 2H), 2.37 and 2.35 (2s, total 3H), 2.18-2.02 (m, total 1H), 1.86-1.19 (m, total 6H), 1.42-1.20 (m, 8H), 0.99-0.94 (m, total 3H), 0.90-0.82 (m, total 6H). ES-MS m/z 345 ($[\text{M}+1]^+$).

EXAMPLE 8

Synthesis of (3S,4S,4aS,8aS)-N-(3-aminopropyl)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine dihydrochloride (Compound No. 11)




A. To a solution of (3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 7, 158 mg, 0.44 mmol) in THF (2 mL) under argon was added a solution of *tert*-butyl N-(3-aminopropyl)carbamate (90 mg, 0.52 mmol) in THF (2 mL) followed by glacial AcOH (40 μL , 0.70 mmol) and the reaction stirred at room temperature for 40 min. Solid $\text{NaBH}(\text{OAc})_3$ (135 mg, 0.636 mmol) was then added and the mixture stirred for 2 d before diluting with CH_2Cl_2 (30 mL) and saturated aqueous NaHCO_3 (20 mL). The aqueous layer was extracted with CH_2Cl_2 (2 x 15 mL) and the combined organic extracts dried (Na_2SO_4) and concentrated. The resultant yellow foam was purified by column chromatography on silica gel (EtOAc then $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 9:1) to afford *tert*-butyl N-(3-[(3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}propyl)carbamate (Compound No. 10, 103 mg, 62%) as a yellow oil.

B. To a solution of *tert*-butyl N-(3-[(3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}propyl)carbamate (Compound No. 10, 39 mg, 0.076 mmol) in MeOH (0.5 mL) was added a solution of HCl in dioxane (4.0 M, 0.30 mL, 1.2 mmol) and the mixture stirred at room temperature for 1.5 h before concentrating *in vacuo*. The product was diluted with

MeOH (2-3 mL) and concentrated and this process was repeated (2x). Et₂O (10 mL) was added to precipitate a white solid and the supernatant solution was decanted. The solid was washed by decantation with Et₂O (3 x 5 mL) and the remaining traces of solvent removed under vacuum to give (3S,4S,4aS,8aS)-N-(3-aminopropyl)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine dihydrochloride (Compound No. 11, 23 mg, 62%) as a white solid. ¹H NMR (CD₃OD): δ 7.69 (s, 1H), 7.40 (s, 1H), 7.00 (s, 1H), 3.96 (d, *J* = 9.9 Hz, 1H), 3.85 (s, 3H), 3.65-3.60 (m, 1H), 3.24-3.02 (m, 4H), 2.70-2.58 (m, 1H), 2.41 (s, 3H), 2.37-2.22 (m, 2H), 2.2 (d, *J* = 9.9 Hz, 1H), 1.92-1.78 (m, 1H), 1.64 (d, *J* = 9.9 Hz, 1H), 1.56-1.26 (m, 6H), 1.05 (s, 3H), 1.00 (s, 3H), 0.95 (d, *J* = 6.9 Hz, 3H), 0.90 (s, 3H). ES-MS *m/z* 415 ([M+1]⁺).

EXAMPLE 9

Synthesis of (4aS,8aS)-8-[(3-methoxy-5-methylphenyl)methyl]-4,4,7,8a-tetramethyl-1,2,3,4,4a,5,6,8a-octahydronaphthalene (Compound No. 13), (4aS,5S,8aS)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene (Compound No. 14) and (4aS,5S,8aS)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene (Compound No. 15)




To a solution of (1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 12, prepared according to Andersen, R. *et al.* PCT Int. Patent Appl. WO 2007/147251 A1, 470 mg, 1.37 mmol) and pyridine (1.10 mL, 13.6 mmol) in CH₂Cl₂ (20 mL) at -78°C under argon was added thionyl chloride (0.50 mL, 6.9 mmol) dropwise neat. The reaction was stirred at -78°C for 1 h then quenched with saturated aqueous sodium bicarbonate (30 mL) and diluted with CH₂Cl₂ (20 mL). The mixture was warmed to room temperature, the layers separated and the aqueous layer extracted with CH₂Cl₂ (3 x 15 mL). The combined organic extracts were washed with brine (1 x 40 mL), dried (Na₂SO₄) and

concentrated to afford a mixture of (4a*S*,8a*S*)-8-[(3-methoxy-5-methylphenyl)methyl]-4,4,7,8a-tetramethyl-1,2,3,4,4a,5,6,8a-octahydronaphthalene, (4a*S*,5*S*,8a*S*)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene and (4a*S*,5*S*,8a*S*)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene (Compound No. 13, Compound No. 14, and Compound No. 15, respectively, 0.44 g, 1.3 mmol) as a clear oil.

EXAMPLE 10

Synthesis of [(1*S*,2*S*,4a*S*,8a*R*)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol (Compound No. 16)




To a mixture of (4a*S*,8a*S*)-8-[(3-methoxy-5-methylphenyl)methyl]-4,4,7,8a-tetramethyl-1,2,3,4,4a,5,6,8a-octahydronaphthalene, (4a*S*,5*S*,8a*S*)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene and (4a*S*,5*S*,8a*S*)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene (Compound No. 13, Compound No. 14, and Compound No. 15 respectively, 0.44 g, 1.3 mmol) in anhydrous THF (11 mL) at 0°C under argon was added a solution of BH₃-THF (1.0 M in THF, 2.6 mL, 2.6 mmol) and the reaction stirred for 2 d before quenching carefully with MeOH (4 mL), 15% aqueous NaOH (2 mL) and 50% aqueous H₂O₂ (3 mL). The mixture was stirred at room temperature for 5 h then diluted with EtOAc (25 mL) and water (20 mL). The layers were separated and the aqueous layer extracted with EtOAc (2 x 20 mL). The combined organic extracts were washed with brine (1 x 40 mL), dried (Na₂SO₄) and concentrated. The resultant white foam (0.46 g) was purified by column chromatography on silica gel (hexanes/EtOAc, 3:1 then 7:3) to give [(1*S*,2*S*,4a*S*,8a*R*)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol (Compound No. 16, 191 mg, 41% from Compound No. 12) as clear oil. ¹H NMR (CDCl₃): δ 6.59 (s, 1H), 6.54 (s, 2H), 3.90 (d, *J* = 9.9 Hz, 1H), 3.77 (s, 3H), 3.57

(t, $J = 9.9$ Hz, 1H), 2.80 (dd, $J = 13.5, 3.2$ Hz, 1H), 2.37 (dd, $J = 13.5, 11.4$ Hz, 1H), 2.30 (s, 3H), 2.04 (s, 1H), 1.96-1.90 (m, 1H), 1.84-1.74 (m, 1H), 1.64-1.14 (m, 9H), 1.08-0.86 (m, 2H), 0.86 (s, 3H), 0.83 (s, 3H), 0.79 (s, 3H). ES-MS m/z 345 ($[M+1]^+$).

EXAMPLE 11

Synthesis of 3-[[[(1S,2S,4aS,8aR)-2-(hydroxymethyl)-5,5,8a-trimethyl-decahydronaphthalen-1-yl]methyl]-5-methylphenol (Compound No. 17)




To a MW vessel (2-5 mL) was added Na_2S (226 mg, 2.90 mmol) and a solution of [(1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol (Compound No. 16, 190 mg, 0.55 mmol) in NMP (3 mL). The reaction vessel was sealed and placed in a Biotage Microwave Initiator reactor and irradiated at 180°C for 3 h. The reaction mixture was allowed to cool to room temperature and then quenched with 10% aqueous HCl (10 mL) and diluted with EtOAc (30 mL). The organic layer was washed with brine (3 x 10 mL), dried (Na_2SO_4) and concentrated. The crude product was purified by flash chromatography on silica gel (hexanes/EtOAc, 3:2) to afford 3-[[[(1S,2S,4aS,8aR)-2-(hydroxymethyl)-5,5,8a-trimethyl-decahydronaphthalen-1-yl]methyl]-5-methylphenol (Compound No. 17, 90 mg, 50%) as a white foam. ^1H NMR (CDCl_3): δ 6.54 (s, 1H), 6.48 (s, 2H), 5.78 (br s, 1H), 3.91 (d, $J = 10.5$ Hz, 1H), 3.60 (t, $J = 10.2$ Hz, 1H), 2.76 (dd, $J = 13.2, 2.9$ Hz, 1H), 2.33 (dd, $J = 13.5, 11.7$ Hz, 1H), 2.26 (s, 3H), 2.05 (s, 1H), 1.94-1.86 (m, 1H), 1.84-1.74 (m, 1H), 1.68-1.14 (m, 9H), 1.09-0.86 (m, 2H), 0.86 (s, 3H), 0.82 (s, 3H), 0.77 (s, 3H). ES-MS m/z 331 ($[M+1]^+$).

Synthesis of (1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 19)

EXAMPLE 13


Synthesis of (1S,2S,4aS,8aR)-1-[(3-hydroxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 20)



To a solution of (1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 19, 105 mg, 0.293 mmol) in CH_2Cl_2 (5 mL) at -78°C under argon was added a solution of BBr_3 (1 M in CH_2Cl_2 , 1.5 mL, 1.5 mmol) and the mixture allowed to stir from -78°C to room temperature overnight. The reaction was quenched with MeOH (10 mL) and concentrated. This was repeated 2x and the residue diluted with *i*-PrOH (10 mL) and 1 N HCl (3 mL) and heated to reflux for 1 h. The mixture was cooled, concentrated and diluted with EtOAc (25 mL) and water (10 mL). The layers were separated and the aqueous layer extracted with EtOAc (1 x 10 mL). The combined organic extracts were dried (Na_2SO_4), concentrated and purified by column chromatography on silica gel (hexanes/EtOAc, 3:1) to afford (1S,2S,4aS,8aR)-1-[(3-methoxy-5-methylphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 20, 53 mg, 53%) as a brown foam. ^1H NMR (CDCl_3): δ 6.68 (s, 1H), 6.65 (s, 1H), 6.48 (s, 1H), 2.97 (t, $J = 12.6$ Hz, 1H), 2.80 (dd, $J = 13.2, 3.3$ Hz, 1H), 2.47-2.42 (m, 1H), 2.28 (s, 3H), 2.24-2.16 (m, 1H), 1.90-1.81 (m, 1H), 1.78-0.92 (m, 10H), 0.94 (s, 3H), 0.87 (s, 3H), 0.87-0.83 (m, 2H), 0.84 (s, 3H). ES-MS m/z 345 ($[\text{M}+1]^+$).

EXAMPLE 14

Synthesis of 3-[[[(1S,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenol hydrochloride (Compound No. 22)



A. To a solution of (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 8, 110 mg, 0.32 mmol) in THF (4 mL) under argon was added NH_4OAc (230 mg, 2.99 mmol), Et_3N (0.10 mL, 0.72 mmol) and $\text{NaBH}(\text{OAc})_3$ (118 mg, 0.557 mmol) and the mixture stirred at room temperature overnight before adding another portion of $\text{NaBH}(\text{OAc})_3$ (118 mg, 0.557 mmol) and stirring for 2 d. The reaction was then diluted with CH_2Cl_2 (30 mL) and saturated aqueous NaHCO_3 (25 mL). The aqueous layer was extracted with CH_2Cl_2 (2 x 10 mL) and the combined organic extracts dried (Na_2SO_4) and concentrated. The resultant yellow foam was purified by column chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 96:4:0 then 88:10:2) to afford the desired amine (40 mg) as an impure mixture (yellow oil).


B. To a solution of the crude amine from above (40 mg) in THF (1 mL) under argon was added a solution of di-*tert*-butyl dicarbonate (40 mg, 0.18 mmol) in THF (1 mL) and the reaction stirred overnight. The mixture was concentrated and purified by column chromatography on silica gel (hexanes/ EtOAc , 4:1 then 7:3) to give *tert*-butyl N-[(4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamate (Compound No. 21, 30 mg, 58%) as a white foam.

C. To a solution of *tert*-butyl N-[(4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamate (Compound No. 21, 30 mg, 0.068 mmol) in MeOH (0.5 mL) was added a solution of HCl in dioxane (4.0 M, 0.30 mL, 1.2 mmol) and the mixture stirred at room temperature for 2 h before concentrating *in vacuo*. The product was diluted with MeOH (2-3 mL) and concentrated and this process was repeated (2x). Et_2O (10 mL) was added to precipitate a white solid and the supernatant solution was decanted. The solid was

washed by decantation with Et₂O (3 x 5 mL) and the remaining traces of solvent removed under vacuum to give 3-[[[(1S,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenol hydrochloride (Compound No. 22, 10 mg, 39%) as a white solid. ¹H NMR (CD₃OD): δ 7.27 (s, 1H), 7.13 (s, 1H), 6.87 (s, 1H), 3.12-3.01 (m, 1H), 2.35 (s, 3H), 2.30-2.16 (m, 1H), 2.06-1.96 (m, 1H), 1.62-1.08 (m, 9H), 1.05 (s, 3H), 0.96 (s, 3H), 0.90 (s, 3H), 0.89 (d, *J* = 6 Hz, 3H). ES-MS *m/z* 344 ([M+1]⁺).

EXAMPLE 15


Synthesis of (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 23)



To a solution of (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 8, 202 mg, 0.590 mmol) in anhydrous THF (12 mL) at 0°C under argon was added a solution of Super-Hydride in THF (1.0 M in THF, 3.6 mL, 3.6 mmol) and the reaction stirred to room temperature overnight. The mixture was quenched with water (10 mL), diluted with EtOAc (20 mL) and brine (20 mL) and the layers separated. The organic phase was washed with brine (1 x 25 mL), dried (Na₂SO₄), concentrated and purified by column chromatography on silica gel (hexanes/EtOAc, 3:1 then 65:35) to afford (3R,4S,4aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 23, 75 mg, 37%) as a white solid. ¹H NMR (CDCl₃): δ 7.28-7.26 (m, 1H), 7.20-7.18 (m, 1H), 6.87-6.86 (m, 1H), 5.45 (s, 1H), 3.40-3.28 (m, 1H), 2.97 (d, *J* = 11.4 Hz, 1H), 2.38 (s, 3H), 2.12-1.94 (m, 2H), 1.61 (s, 3H), 1.50-1.35 (m, 2H), 1.30-1.22 (m, 1H), 1.18-1.00 (m, 3H), 1.07 (s, 3H), 0.90 (s, 3H), 0.89 (d, *J* = 6 Hz, 3H), 0.86 (s, 3H). ES-MS *m/z* 345 ([M+1]⁺).

EXAMPLE 16

Synthesis of (1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 27)




A. Cu(I)I (0.05 g, 0.25 mmol) and NaI (1.49 g, 10 mmol) were added to an oven dried sealed tube and the tube sealed with a rubber septum and flushed with a stream of nitrogen for 15 min. Anhydrous dioxane (5 mL), N,N'-dimethylethylenediamine (0.055 mL, 0.50 mmol) and 1-bromo-3-methoxy-5-methylbenzene (Compound No. 24, 1.0 g, 5 mmol) were added via a syringe with vigorous stirring under a stream of nitrogen at room temperature. The rubber septum was replaced with the Teflon cap and the sealed tube was heated at 110°C in an oil bath for 18 h. The reaction was allowed to cool to room temperature and then quenched with a saturated aqueous solution of ammonium chloride. The reaction mixture was extracted with CH₂Cl₂. The organic layer was washed with water, and then concentrated to dryness to give 1-iodo-3-methoxy-5-methylbenzene (Compound No. 25, 1.05 g, 87% yield) as a pale yellow oil.

B. (1S,2R,4aS,8aS)-1-(Hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 26, prepared according to Kuchkova *et al. Synthesis*, 1997, 1045, 938 mg, 3.90 mmol), CuI (84 mg, 0.44 mmol), 3,4,7,8-tetramethyl-[1,10]phenanthroline (188 mg, 0.796 mmol), and Cs₂CO₃ (1.91 g, 5.86 mmol) were weighed into a sealed flask and degassed with Ar for 2 min. A solution of 1-iodo-3-methoxy-5-methylbenzene (Compound No 25, 1.04 g, 4.19 mmol) in toluene (20 mL) and diglyme (4 mL) was added to the flask and the reaction heated to 120°C for 3 d. The mixture was filtered through Celite, washed with EtOAc (300 mL), dried (MgSO₄) and concentrated. The crude product was purified by chromatography on silica gel

(EtOAc/hexanes, 1:9 to 1:4) to afford (1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxy-methyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 27, 1.18g, 84%) as a yellow oil. ^1H NMR (CDCl_3): δ 6.36-6.35 (d, $J = 2.1$ Hz, 2H), 6.30 (t, $J = 2.4$ Hz, 1H), 4.24-4.16 (m, 2H), 3.77 (s, 3H), 2.30 (s, 3H), 3.23 (br s, 1H), 1.96-1.88 (m, 1H), 1.82-1.10 (m, 10H), 1.27 (s, 3H), 1.06-1.00 (m, 1H), 0.91 (s, 3H), 0.90 (s, 3H), 0.82 (s, 3H). ES-MS m/z 361 ($[\text{M}+1]^+$).

EXAMPLE 17

Synthesis of 3-[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy-5-methylphenol (Compound No. 29)




A. To a solution of (1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxy-methyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 27, 1.18 g, 32.8 mmol) in CH_2Cl_2 (70 mL) at -78°C under argon was added neat SnCl_4 (0.88 mL, 7.5 mmol) dropwise and the reaction stirred at -78°C for 3.5 h before adding another portion of SnCl_4 (0.88 mL, 7.5 mmol). The mixture was stirred another 2 h at -78°C then quenched at -78°C with water (50 mL). The reaction was warmed to room temperature, the layers separated and the aqueous layer extracted with CH_2Cl_2 (1 x 40 mL). The combined organic extracts were washed with saturated aqueous NaHCO_3 (1 x 60 mL), dried (Na_2SO_4) and concentrated to afford a yellow oil (1.1 g). Purification of the crude product by column chromatography on silica gel (hexanes/EtOAc, 98:2) gave an impure mixture containing (4aS,5S,8aS)-5-(3-methoxy-5-methylphenoxy-methyl)-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene (Compound No. 28, 1.16 g) as a clear oil.

B. To a MW vessel (2-5 mL) was added Na_2S (460 mg, 5.89 mmol) and a solution of the impure mixture containing (4aS,5S,8aS)-5-(3-methoxy-5-methylphenoxy-methyl)-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene (Compound No. 28, ~1 mmol) in NMP (6 mL). The reaction vessel was sealed and

placed in a Biotage Microwave Initiator reactor and irradiated at 180°C for 1.5 h. The reaction mixture was allowed to cool to room temperature and then quenched with 1 N HCl (20 mL) and diluted with EtOAc (40 mL). The organic layer was washed with water (1 x 40 mL), dried (Na₂SO₄) and concentrated. The crude product was purified by flash chromatography on silica gel (hexanes/EtOAc, 92:8 then 85:15) to afford 3-[[[(1S,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy]-5-methylphenol (Compound No. 29, 40 mg, 10%) as a colorless oil. ¹H NMR (CDCl₃): δ 6.31 (s, 1H), 6.24-6.20 (m, 2H), 5.50 (br s, 1H), 4.71 (br s, 1H), 4.07 (dd, *J* = 9.6, 3 Hz, 1H), 3.88 (dd, *J* = 9.6, 6.3 Hz, 1H), 2.26 (s, 3H), 2.20-1.83 (m, 4H), 1.70 (s, 3H), 1.66-1.38 (m, 6H), 0.90 (s, 3H), 0.88 (s, 6H). ES-MS *m/z* 329 ([M+1]⁺).

EXAMPLE 18

Synthesis of (4S,4aS,8aS)-4-(3-methoxy-5-methylphenoxyethyl)-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 30)




To a solution of the impure mixture containing (4aS,5S,8aS)-5-(3-methoxy-5-methylphenoxyethyl)-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene (Compound No. 28, ~0.7 mmol) in dry THF (9 mL) at 0°C under argon was added a solution of BH₃-THF (1.0 M in THF, 1.0 mL, 1.0 mmol) dropwise. The reaction was stirred under argon for 1.5 h, then allowed to warm to room temperature and stirred for an additional 1.5 h. The reaction was quenched with MeOH (7 mL), 15% aqueous NaOH (0.23 mL) and 50% aqueous H₂O₂ (0.23 mL) and allowed to stir at room temperature overnight. The mixture was diluted with EtOAc (25 mL) and water (25 mL). The layers were separated and the organic layer washed with brine (1 x 25 mL), dried (MgSO₄) and concentrated. The crude material was purified by chromatography on silica gel (EtOAc/hexanes, 91:9 then 4:1) to yield (4S,4aS,8aS)-4-(3-methoxy-5-methylphenoxyethyl)-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 30, 50 mg, 21%) as a white solid. ¹H NMR (CDCl₃): δ 6.33-6.30 (m, 2H), 6.29-6.26 (m,

1H), 4.12 (dd, $J = 9.3, 4.8$ Hz, 1H), 3.94 (br s, 1H), 3.86 (t, $J = 9.3$ Hz, 1H), 3.77 (s, 3H), 2.29 (s, 3H), 2.20-2.04 (m, 2H), 1.76-1.04 (m, 10H), 0.96 (d, $J = 8.1$ Hz, 3H), 0.94 (s, 3H), 0.88 (s, 3H), 0.84 (s, 3H). ES-MS m/z 361 ($[M+1]^+$).

EXAMPLE 19

Synthesis of (1S,2R,4aS,8aS)-1-[3,5-bis(benzyloxy)-phenoxyethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 33)



A. To a solution of 1-bromo-3,5-dimethoxybenzene (3.01 g, 13.9 mmol) in dry THF (25 mL) at -78°C under argon was added a solution of *n*-BuLi (2.2 M in hexanes, 7.00 mL, 15.4 mmol) dropwise and the mixture stirred for 40 min. A solution of iodine in dry THF (20 mL) was then added dropwise and the reaction allowed to stir for an additional 45 min at -78°C before quenching with 20% aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (45 mL) and warming to room temperature. The mixture was diluted with EtOAc (50 mL), the layers separated and the organic layer washed with brine (50 mL), dried (MgSO_4) and concentrated to a yellow solid (3.12 g). Purification of the crude product by chromatography on silica gel (20% EtOAc/hexanes) gave 1-iodo-3,5-dimethoxybenzene (Compound No. 31, 2.01 g, 55%) as a white solid.


B. To a solution of 1-iodo-3,5-dimethoxybenzene (Compound No. 31, 2.01 g, 7.61 mmol) in CH_2Cl_2 (20 mL) at 0°C under argon was added a solution of BBr_3 (1 M in CH_2Cl_2 , 11.4 mL, 11.4 mmol) and the mixture allowed to stir for 17 h. The reaction was quenched with MeOH (13 mL) and concentrated. The residue was diluted with EtOAc (25 mL) and the organic layer washed with water (3 x 20 mL) and brine (20 mL), dried (MgSO_4) and concentrated to a brown oil.

C. To a solution of the crude residue from above in DMF (20 mL) under argon was added K_2CO_3 (2.32 g, 16.8 mmol) and benzyl bromide (2.00 mL, 16.8 mmol) and the mixture was allowed to stir for 2 d. The mixture was diluted with EtOAc (40 mL) and the organic layer washed with water (3 x 30 mL), dried ($MgSO_4$) and concentrated to a dark yellow oil. The crude material was purified by chromatography on silica gel (EtOAc/hexanes, 5:95 to 15:85) to yield 1,3-bis(benzyloxy)-5-iodobenzene (Compound No. 32, 2.66 g, 84% over 2 steps) as an off-white solid.

D. (1S,2R,4aS,8aS)-1-(Hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 26, 1.54 g, 6.41 mmol), CuI (0.12 g, 0.63 mmol), 3,4,7,8-tetramethyl-[1,10]phenanthroline (0.30 g, 1.3 mmol), and Cs_2CO_3 (3.19 g, 9.79 mmol) were weighed into a sealed flask and degassed with Ar for 2 min. A solution of 1,3-bis(benzyloxy)-5-iodobenzene (Compound No. 32, 2.66 g, 6.39 mmol) in toluene (30 mL) and diglyme (6 mL) was added to the flask and the reaction heated to 120°C for 2 d. The mixture was filtered through Celite, washed with EtOAc (300 mL), dried ($MgSO_4$) and concentrated. The crude product was purified by chromatography on silica gel (EtOAc/hexanes, 8:92) to afford (1S,2R,4aS,8aS)-1-[3,5-bis(benzyloxy)-phenoxy-methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 33, 1.48 g, 44%) as a white foam. ^1H NMR ($CDCl_3$): δ 7.45-7.32 (m, 10H), 6.27 (t, J = 2.1 Hz, 1H), 6.21 (d, J = 2.1 Hz, 2H), 5.00 (s, 4H), 4.10-4.23 (m, 2H), 1.95-1.88 (m, 1H), 1.81-1.28 (m, 8H), 1.25 (s, 3H), 1.24-0.98 (m, 4H), 0.90 (s, 3H), 0.89 (s, 3H), 0.82 (s, 3H).

EXAMPLE 20

Synthesis of (1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxy-methyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 34)




In a sealed tube purged with nitrogen, 1-iodo-3,5-dimethoxybenzene (Compound No. 31, 0.78 g, 2.95 mmol), (1S,2R,4aS,8aS)-1-(hydroxymethyl)-2,5,5,8a-

tetramethyl-decahydronaphthalen-2-ol (Compound No. 26, 0.65 g, 2.68 mmol), Cs_2CO_3 (1.3 g, 3.99 mmol), 3,4,7,8-tetramethyl-[1,10]-phenylthrine (0.12 g, 0.50 mmol), CuI (0.05 g, 0.26 mmol), diglyme (2 mL) and toluene (10 mL) were added. The reaction mixture was stirred at 120°C for 18 h. The reaction was cooled to room temperature, diluted with EtOAc (300 mL) and washed with water (100 mL). The organic layer was dried (Na_2SO_4) and concentrated. Purification by column chromatography on silica gel (Hexanes/EtOAc, 2:1) gave (1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 34, 0.71 g, 70%) as a colorless oil. ^1H NMR (CDCl_3): δ 6.08 (s, 3H), 4.20 (m, 2H), 3.75 (s, 6H), 3.08 (s, 1H), 1.95-1.30 (m, 10H), 1.30 (s, 3H), 1.25-1.00 (m, 2H), 0.90 (s, 6H), 0.88 (m, 2H), 0.80 (s, 3H).

EXAMPLE 21

Synthesis of (1S,2R,4aS,8aS)-1-[3,5-bis(propan-2-yloxy)phenoxymethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 37)



A. To a solution of 1-bromo-3,5-dimethoxybenzene (15.0 g, 69.0 mmol) in CH_2Cl_2 (40 mL) at 0°C under argon was added dropwise a 1 M BBr_3 solution in CH_2Cl_2 (140 mL, 140 mmol). The ice bath was removed and the reaction was stirred at room temperature for 18 h. The reaction was quenched with MeOH and concentrated. The residue was dissolved in EtOAc (500 mL) and washed with water (300 mL). The organic layer was separated, dried (Na_2SO_4) and concentrated to give 5-bromobenzene-1,3-diol (13.03 g, 69.0 mmol) as an orange oil.


B. To a solution of 5-bromobenzene-1,3-diol (13.03 g, 69.0 mmol) in DMF (150 mL) under argon was added K_2CO_3 (38.08 g, 276 mmol) followed by 2-bromopropane (26.0 mL, 276 mmol). The mixture was heated to 60°C and stirred for 18 h. The reaction was cooled to room temperature, quenched with water (700 mL) and extracted with EtOAc (3 x 300 mL). The organic layer was separated, dried (Na_2SO_4) and concentrated. Purification by column chromatography on silica gel (Hexanes/EtOAc, 10:1) gave 1-bromo-3,5-bis(propan-2-yloxy)benzene (Compound No. 35, 17.6 g, 93%) as an orange oil.

C. To a solution of 1-bromo-3,5-bis(propan-2-yloxy)benzene (Compound No. 35, 2.0 g, 7.32 mmol) in THF (100 mL) at -78°C under argon was added dropwise a 1.6 M *n*-BuLi solution in hexanes (9.16 mL, 14.65 mmol). The temperature was maintained at -78°C for 30 min, after which a solution of iodine (3.17 g, 29.3 mmol) in THF (15 mL) was added dropwise. The mixture was warmed to -35°C and maintained at this temperature for 1.5 h. ^1H NMR indicated that the reaction was complete. The reaction mixture was diluted with EtOAc (100 mL) and washed with water (2 x 50 mL) and brine (50 mL). The organic layer was separated, dried (Na_2SO_4) and concentrated. Purification by column chromatography on silica gel (Hexanes/EtOAc, 4:1) gave 1-iodo-3,5-bis(propan-2-yloxy)benzene (Compound No. 36, 1.85 g, 79%) as a viscous orange oil.

D. In a sealed tube purged with nitrogen, 1-iodo-3,5-bis(propan-2-yloxy)benzene (Compound No. 36, 1.85 g, 5.78 mmol), (1*S*,2*R*,4*aS*,8*aS*)-1-(hydroxymethyl)-2,5,5,8*a*-tetramethyl-decahydronaphthalen-2-ol (Compound No. 26, 1.26 g, 5.25 mmol), Cs_2CO_3 (2.56 g, 7.87 mmol), 3,4,7,8-tetramethyl-[1,10]-phenylthrolin (0.25 g, 1.04 mmol), CuI (0.10 g, 0.70 mmol), diglyme (5 mL) and toluene (20 mL) were placed. The reaction mixture was stirred at 120°C for 18 h. The reaction was cooled to room temperature, diluted with EtOAc (600 mL) and washed with water (200 mL). The solid material was removed by filtration. The organic layer was separated, dried (Na_2SO_4) and concentrated. Purification by column chromatography on silica gel (Hexanes/EtOAc, 4:1) gave (1*S*,2*R*,4*aS*,8*aS*)-1-[3,5-bis(propan-2-yloxy)phenoxyethyl]-2,5,5,8*a*-tetramethyl-decahydronaphthalen-2-ol (Compound No. 37, 1.50 g, 66%) as a colorless oil. ^1H NMR ($CDCl_3$): δ 6.05 (s, 3H), 4.47 (m, 2H), 4.20 (m, 2H), 3.28 (s, 1H), 1.95-1.35 (m, 8H), 1.32 (d, 12H), 1.30 (s, 3H), 1.29-1.00 (m, 4H), 0.90 (s, 3H), 0.88 (s, 3H), 0.80 (s, 3H).

EXAMPLE 23

Synthesis of (4a*S*,8a*S*)-*N*-(3,5-dimethoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 41)




A. A solution of (1*R*,2*R*,4a*S*,8a*S*)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalene-1-carbaldehyde (Compound No. 39, prepared according to Andersen, R. *et al.* PCT Int. Patent Appl. WO 2007/147251 A1, 2.94 g, 12.3 mmol) and *p*-TsOH·H₂O (117 mg, 0.615 mmol) in toluene (100 mL) under argon was heated to reflux under a Dean-Stark trap for 1 h 45 min. The solution was washed with saturated aqueous NaHCO₃ (25 mL), and the aqueous layer was extracted with EtOAc (15 mL). The combined organic layers were dried (Na₂SO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:99 then 98:2) to give (4a*S*,8a*S*)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carbaldehyde (1.33 g, 49%) as colorless crystals.

B. A solution of NaClO₂ (440 mg, 3.89 mmol) in water (7.8 mL) was added to a mixture of (4a*S*,8a*S*)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carbaldehyde (612 mg, 2.78 mmol), aqueous NaH₂PO₄ (2.6 mL, pH 4.2), 50% aqueous H₂O₂ (0.18 mL, 3.1 mmol), CH₃CN (5.6 mL), and CH₂Cl₂ (0.5 mL) then stirred at room temperature for 19.5 h. The mixture was diluted with water (10 mL), and excess peroxide was destroyed with solid Na₂SO₃. The aqueous phase was extracted with CH₂Cl₂ (4 x 10 mL) then dried (Na₂SO₄) and concentrated to give (8a*S*)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxylic acid (Compound No. 40, 582 mg, 89%) as colorless crystals.

C. To a solution of (4a*S*,8a*S*)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxylic acid (Compound No. 40, 148 mg, 0.626 mmol) in CH₂Cl₂ (6 mL) under argon was added DMF (cat.) and oxalyl chloride (0.16 mL, 1.8 mmol) then stirred at room temperature for 1 h. The solution was concentrated then azeotroped with CH₂Cl₂ (3 x 8 mL). The acid chloride residue was dissolved in pyridine (3 mL), and 3,5-dimethoxyaniline (101 mg, 0.659 mmol) was added. The solution was stirred at room temperature for 1.5 h then concentrated. The residue was dissolved in EtOAc (30 mL), washed with saturated aqueous NaHCO₃ (2 x 10 mL) and brine (10 mL) then dried (Na₂SO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:9) to give (4a*S*,8a*S*)-N-(3,5-dimethoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 41, 56 mg, 24%) as a yellow foam.

EXAMPLE 24

Synthesis of (4a*S*,8a*S*)-N-(3-hydroxy-5-methoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 42) and (4a*S*,8a*S*)-N-(3,5-dihydroxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 43)



A. To a solution of (4a*S*,8a*S*)-N-(3,5-dimethoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 41, 56 mg, 0.15 mmol) in CH₂Cl₂ (3 mL) at -78°C under argon was added BBr₃ (1.0 M in CH₂Cl₂, 0.60 mL, 0.60 mmol). The solution was stirred at -78°C for 10 min then at room temperature for 2 h. Methanol (10 mL) was carefully added to the solution then concentrated (3x). The residue was dissolved in *i*-PrOH/10% aqueous HCl (3:1, 4 mL) and heated to 80°C for 45 min. The solution was diluted with EtOAc (30 mL) and washed with brine (2 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (hexanes/acetone, 4:1 then 3:2) to give (4a*S*,8a*S*)-N-(3-hydroxy-5-methoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-


octahydronaphthalene-1-carboxamide (Compound No. 42, 16 mg, 30%) as a yellow solid and (4aS,8aS)-N-(3,5-dihydroxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 43, 17 mg, 33%) as a yellow solid.

B. (8aS)-N-(3-hydroxy-5-methoxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydro-naphthalene-1-carboxamide (Compound No. 42): ^1H NMR (CDCl_3): δ 8.38 (br s, 1H), 7.61 (m, 1H), 7.06 (s, 1H), 6.26 (m, 1H), 6.16 (m, 1H), 3.75 (s, 3H), 2.09 (m, 2H), 0.87-1.70 (m, 21 H). ES-MS m/z 358 ($[\text{M}+1]^+$).

C. (8aS)-N-(3,5-dihydroxyphenyl)-2,5,5,8a-tetramethyl-3,4,4a,5,6,7,8,8a-octahydronaphthalene-1-carboxamide (Compound No. 43): ^1H NMR (CDCl_3): δ 7.75 (br s, 2H), 7.18 (s, 1H), 6.79 (s, 1H), 6.75 (s, 1H), 6.26 (s, 1H), 2.05 (m, 2H), 1.65-0.82 (m, 21 H). ES-MS m/z 344 ($[\text{M}+1]^+$).

EXAMPLE 25

Synthesis of 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 45)




To a solution of 1-bromo-3,5-dimethoxybenzene (3.34 g, 15.4 mmol) in THF (28 mL) under argon at -78°C was added *n*-BuLi (2.2 M in hexanes, 6.35 mL, 14.0 mmol). The solution was stirred at -78°C for 15 min then added via cannula to a solution of sclareolide (Compound No. 44, 3.50 g, 14.0 mmol) in THF (28 mL) at -78°C . After stirring at -78°C for 2.5 h, saturated aqueous NH_4Cl (25 mL) was added, and the mixture was allowed to warm to room temperature. The mixture was acidified with 1N HCl (25 mL), and the aqueous phase was extracted with EtOAc (30 mL). The combined organic layers were washed with brine (20 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:4) to give 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 45, 3.06 g, 56%) as colorless crystals. ^1H NMR (CDCl_3): δ 7.13 (d, $J = 2.4$ Hz, 2H), 6.64 (t,

$J = 2.4$ Hz, 1H), 3.84 (s, 6H), 3.06 (dd, $J = 18, 5.0$ Hz, 1H), 2.94 (dd, $J = 18, 5.1$ Hz, 1H), 2.19 (t, $J = 5.1$ Hz, 1H), 1.97 (m, 1H), 1.71 (m, 1H), 1.59-1.24 (m, 7H), 1.12 (s, 3H), 1.11 (m, 2H), 0.92 (m, 1H), 0.88 (s, 3H), 0.87 (s, 3H), 0.80 (s, 3H). ES-MS m/z 371 ($[M-17]^+$).

EXAMPLE 26

Synthesis of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 46)




A. A solution of SOCl₂ (2.64 mL, 36.3 mmol) and pyridine (4.70 mL, 58.1 mmol) in CH₂Cl₂ (18 mL) was added to a solution of 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 45, 2.82 g, 7.26 mmol) and pyridine (1.17 mL, 14.5 mmol) in CH₂Cl₂ (73 mL) at -78°C under argon. After stirring at -78°C for 55 min, saturated aqueous NaHCO₃ (30 mL) was added, and the mixture was allowed to warm to room temperature. The aqueous phase was extracted with CH₂Cl₂ (2 x 10 mL), and the combined organic layers were dried (MgSO₄), concentrated, and azeotroped with hexanes (3 x 30 mL). The residue was filtered through silica gel using 5% EtOAc/hexanes to give a colorless oil (2.43 g) that was used in the next step without further purification.

B. To a solution of the colorless oil from above (2.43 g) in EtOAc (20 mL) was added 10% Pd/C (224 mg) then shaken under an atmosphere of H₂(g) at 45 psi for 2 h. The mixture was filtered through Celite then concentrated. The residue was dissolved in THF (66 mL), cooled to 0°C under argon, and BH₃·THF (1.0 M in THF, 4.6 mL, 4.6 mmol) was added. After stirring at room temperature for 1.5 h, the solution was cooled to 0°C and 10 N NaOH (0.62 mL, 6.2 mmol) was added followed by 50% aqueous H₂O₂ (0.53 mL, 9.2 mmol). The mixture was stirred at room temperature for 0.5 h then at 60°C for 1 h. The mixture was diluted with water (20 mL), excess peroxide was destroyed with solid Na₂SO₃ and the organic layer was washed with brine (15 mL). The combined aqueous layers were extracted with EtOAc (20 mL), and

the combined organic layers were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 2:98) to give 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 46, 947 mg, 35% over 2 steps) as colorless crystals. ^1H NMR (CDCl_3): δ 7.11 (d, $J = 2.4$ Hz, 2H), 6.64 (t, $J = 2.4$ Hz, 1H), 3.84 (s, 6H), 2.85 (dd, $J = 18, 3.2$ Hz, 1H), 2.75 (dd, $J = 19, 6.2$ Hz, 1H), 1.84-1.72 (m, 2H), 1.65-0.82 (m, 20H), 0.68 (d, $J = 6.6$ Hz, 3H). ES-MS m/z 373 ($[\text{M}+1]^+$).

EXAMPLE 27

Synthesis of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3-hydroxy-5-methoxyphenyl)ethan-1-one (Compound No. 47) and 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dihydroxyphenyl)ethan-1-one (Compound No. 48)




A. A solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 46, 200 mg, 0.537 mmol) in CH_2Cl_2 (5.4 mL) was cooled to -78°C under argon. BBr_3 (1.0 M in CH_2Cl_2 , 2.15 mL, 2.15 mmol) was added, and the solution was stirred at room temperature for 2 h. Methanol (5 mL) was carefully added to the solution then concentrated (3x). The residue was dissolved in *i*-PrOH/10% aqueous HCl (3:1, 5.5 mL) and heated to 85°C for 35 min. The solution was concentrated, diluted with EtOAc (30 mL), washed with brine (2 x 10 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 3:7) to give 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3-hydroxy-5-methoxyphenyl)ethan-1-one (Compound No. 47, 37 mg, 19%) as a colorless foam and 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dihydroxyphenyl)ethan-1-one (Compound No. 48, 149 mg, 81%) as a colorless foam. 2-[(1S,2S,4aS,8aR)-2,5,5,8a-Tetramethyl-decahydronaphthalen-1-yl]-1-(3-hydroxy-5-methoxyphenyl)ethan-1-one (Compound No. 47): ^1H NMR (CDCl_3): δ 7.11 (m, 2H), 6.63 (m, 1H), 5.81 (s, 1H), 3.83 (s, 3H), 2.84 (dd, $J = 19, 3.3$ Hz, 1H), 2.45 (dd, $J = 19,$

6.0 Hz, 1H), 1.82-1.71 (m, 2H), 1.73-0.82 (m, 20H), 0.67 (d, $J = 6.3$ Hz, 3H). ES-MS m/z 359 ($[M+1]^+$). 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dihydroxyphenyl)ethan-1-one (Compound No. 48): ^1H NMR (CDCl_3): δ 7.08 (d, $J = 2.4$ Hz, 2H), 6.58 (t, $J = 2.1$ Hz, 1H), 5.43 (s, 2H), 2.83 (dd, $J = 19.3, 3.2$ Hz, 1H), 2.73 (dd, $J = 19, 6.2$ Hz, 1H), 1.76 (m, 2H), 1.66-0.82 (m, 20H), 0.66 (d, $J = 6.3$ Hz, 3H). ES-MS m/z 345 ($[M+1]^+$).

EXAMPLE 28


Synthesis of (1S,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)(hydroxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 49)



To dried magnesium turnings (530 mg, 21.8 mmol) under argon was added THF (7 mL) followed by 1,2-dibromoethane (0.18 mL, 2.1 mmol) and a few drops of a solution of 1-bromo-3,5-dimethoxybenzene (4.73 g, 21.8 mmol) in THF (15 mL). Once the reaction initiated, the remainder of the 1-bromo-3,5-dimethoxybenzene solution was added over 10 min. After stirring at room temperature for 1 h, the mixture was cooled to -15°C and a solution of (1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalene-1-carbaldehyde (Compound No. 39, 2.60 g, 10.9 mmol) in THF (20 mL) was added over 10 min. The mixture was stirred at -15°C for 0.5 h then 0°C for 2.5 h. The mixture was acidified to pH 2 using 10% aqueous HCl then diluted with water (20 mL) and EtOAc (60 mL). The organic phase was washed with brine (10 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:3 to 3:7) to give (1S,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)(hydroxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 49, 1.01 g, 25%) as a colorless foam. ^1H NMR (CDCl_3): δ 6.61 (d, $J = 2.4$ Hz, 2H), 6.37 (t, $J = 2.3$ Hz, 1H), 4.80 (dd, $J = 8.1, 3.9$ Hz, 1H), 4.02 (s, 1H), 3.81 (s, 6H), 2.76 (d, $J = 3.9$ Hz, 1H), 2.12 (d, $J = 7.8$ Hz, 1H), 1.85 (m, 1H), 1.68-1.17 (m, 9H), 1.04 (s, 3H), 0.96 (m, 2H), 0.84 (s, 3H), 0.79 (s, 3H), 0.42 (m, 1H).

EXAMPLE 29


Synthesis of (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 50)



To a solution of (1S,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)(hydroxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 49, 3.01 g, 7.99 mmol) in CH_2Cl_2 (32 mL) was added PCC (1.90 g, 8.81 mmol) and the mixture was stirred at room temperature for 2 h then filtered and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 3:17 to 1:4) to afford (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 50, 1.94 g, 65%) as a colorless foam. ^1H NMR (CDCl_3): δ 7.14 (d, $J = 2.1$ Hz, 2H), 6.61 (t, $J = 2.3$ Hz, 1H), 3.84 (s, 6H), 3.53 (s, 1H), 1.87 (m, 1H), 1.77-1.00 (m, 17H), 0.90 (s, 3H), 0.83 (s, 3H). ES-MS m/z 357 ($[\text{M}-17]^+$).

EXAMPLE 30

Synthesis of [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 51)




A solution of (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)carbonyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 50, 1.92 g, 5.13 mmol) in CH_2Cl_2 (70 mL) was cooled to 0°C under argon and SnCl_4 (1.80 mL, 15.4 mmol) was added. The solution was stirred at 0°C for 10 min then washed with water (30 mL).

The aqueous phase was extracted with CH_2Cl_2 (2 x 10 mL), and the combined organic phases were washed with saturated aqueous NaHCO_3 (30 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:19) to afford [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 51, 1.54 g, 84%) as a colorless oil. ^1H NMR (CDCl_3): δ 7.06 (d, J = 2.4 Hz, 2H), 6.63 (t, J = 2.3 Hz, 1H), 5.62 (br s, 1H), 4.09 (br s, 1H), 3.85 (s, 6H), 2.02 (m, 2H), 1.55-1.13 (m, 10H), 0.96 (s, 3H), 0.91 (s, 3H), 0.90 (s, 3H). ES-MS m/z 357 ($[\text{M}+1]^+$).

EXAMPLE 31

Synthesis of (2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 52)




To a solution of [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 51, 1.52 g, 4.26 mmol) in THF (100 mL) was added $\text{BH}_3\cdot\text{THF}$ (1.0 M in THF, 6.40 mL, 6.40 mmol) at 0°C under argon. The solution was stirred at 0°C for 10 min then room temperature for 1.5 h. The solution was cooled to 0°C , and a solution containing 15% aqueous NaOH (2.24 mL) and 50% aqueous H_2O_2 (0.74 mL, 13 mmol) was added dropwise. The mixture was stirred at room temperature for 20 min and 65°C for 1 h then allowed to cool to room temperature. To the mixture was added saturated aqueous NH_4Cl (30 mL) and acidified to pH 2 using concentrated HCl . The aqueous phase was extracted with EtOAc (30 mL), and the combined organic layers were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:4) to afford (2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 52, 0.60 g, 38%) as colorless crystals. ^1H NMR (CDCl_3): δ 7.08 (d, J = 2.4 Hz, 2H), 6.61 (t, J = 2.4 Hz, 1H), 3.95 (br s, 1H), 3.83 (s, 6H), 3.70 (d, J = 4.5 Hz,

1H), 2.14 (m, 1H), 1.84-1.18 (m, 12H), 0.98 (m, 4H), 0.89 (s, 3H), 0.86 (s, 3H). ES-MS m/z 375 ($[M+1]^+$).

EXAMPLE 32


Synthesis of (3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 53)



To a solution of (2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 52, 0.58 g, 1.5 mmol) in CH_2Cl_2 (15 mL) was added PCC (0.50 g, 2.3 mmol), and the mixture was stirred at room temperature for 4 h then filtered and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 3:17) to afford (3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 53, 543 mg, 93%) as a colorless oil. ^1H NMR (CDCl_3): δ 6.99 (d, J = 2.4 Hz, 2H), 6.63 (t, J = 2.1 Hz, 1H), 3.83 (s, 6H), 3.62 (d, J = 6.6 Hz, 1H), 2.83 (m, 1H), 2.53 (dd, J = 17, 13 Hz, 1H), 2.44 (dd, J = 17, 4.2 Hz, 1H), 1.86 (m, 1H), 1.72-1.40 (m, 4H), 1.32 (s, 3H), 1.19 (m, 2H), 1.08 (d, J = 7.5 Hz, 3H), 0.90 (s, 6H). ES-MS m/z 373 ($[M+1]^+$).

EXAMPLE 33


Synthesis of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine
(Compound No. 54)



A mixture of (3R,4S,4aS,8S)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 53, 531 mg, 1.43 mmol), 4-methoxybenzylamine (0.19 mL, 1.5 mmol), sodium triacetoxyborohydride (906 mg, 4.27 mmol), and AcOH (0.082 mL, 1.4 mmol) in CH₂Cl₂ (14 mL) was stirred at room temperature under argon for 22.5 h then washed with 1 N NaOH (10 mL). The aqueous phase was extracted with CH₂Cl₂ (3 x 10 mL), and the combined organic layers were dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:9) to afford (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 54, 443 mg, 63%) as a colorless oil. ¹H NMR (CDCl₃): δ 7.27 (m, 2H), 7.07 (d, *J* = 2.4 Hz, 2H), 6.88 (m, 2H), 6.61 (t, *J* = 2.4 Hz, 1H), 3.84 (d, 1H), 3.84 (s, 6H), 3.82 (s, 3H), 3.54 (m, 2H), 2.87 (m, 1H), 2.34 (m, 1H), 1.86 (m, 1H), 1.66-1.06 (m, 12H), 0.92 (s, 3H), 0.86 (s, 3H), 0.75 (d, *J* = 6.9 Hz, 3H). ES-MS *m/z* 494 ([M+1]⁺).

EXAMPLE 34


Synthesis of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55)



To a suspension of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 54, 433 mg, 0.877 mmol) in MeOH (10 mL) was carefully added 10% aqueous HCl (0.5 mL) and 10% Pd/C (90 mg), and the mixture was stirred under $\text{H}_2(\text{g})$ at 1 atm at room temperature for 23 h. The mixture was filtered through Celite then concentrated. To the residue was added water (10 mL) then adjusted to pH 10 with 1 N NaOH. The mixture was extracted with CH_2Cl_2 (3 x 10 mL), and the combined organic extracts were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 200:2:1-200:10:1) to afford (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 208 mg, 63%) as colorless crystals. ^1H NMR (CDCl_3): δ 7.09 (d, $J = 1.8$ Hz, 2H), 6.62 (t, $J = 2.3$ Hz, 1H), 3.84 (s, 6H), 3.47 (d, $J = 11$ Hz, 1H), 3.22 (m, 1H), 2.31 (m, 1H), 1.79-1.07 (m, 14H), 0.88 (s, 3H), 0.84 (s, 3H), 0.77 (d, $J = 7.2$ Hz, 3H). ES-MS m/z 374 ($[\text{M}+1]^+$).

EXAMPLE 35


Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56)



A solution of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 197 mg, 0.527 mmol) in CH_2Cl_2 (5.3 mL) was cooled to -78°C under argon, BBr_3 (1.0 M in CH_2Cl_2 , 2.64 mL, 2.64 mmol) was added, and the solution was stirred at room temperature for 3.5 h. Methanol (10 mL) was carefully added to the solution then concentrated (3x). The residue was dissolved in *i*-PrOH/10% aqueous HCl (3:1, 5.5 mL) and heated to 85°C for 40 min. The solution was concentrated, diluted with EtOAc (30 mL) and water (10 mL), then neutralized to pH 7 using 1N NaOH. The aqueous phase was extracted with EtOAc (10 mL), and the combined organic layers were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 100:10:1) to afford 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 162 mg, 89%) as a yellow foam. ^1H NMR (CD_3OD): δ 6.88 (d, $J = 2.1$ Hz, 2H), 6.47 (t, $J = 2.3$ Hz, 1H), 3.45 (d, $J = 12$ Hz, 1H), 3.27 (m, 1H), 2.31 (m, 1H), 1.78 (m, 2H), 1.55-1.18 (m, 6H), 1.06 (m, 4H), 0.92 (s, 3H), 0.87 (s, 3H), 0.77 (d, $J = 6.6$ Hz, 3H). ES-MS m/z 346 ($[\text{M}+1]^+$).

EXAMPLE 36

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide (Compound No. 57)




A. To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 111 mg, 0.321 mmol), DMAP (4 mg, 0.03 mmol), and Et_3N (0.16 mL, 1.1 mmol) in THF (3.2 mL) at 0°C was added Ac_2O (0.10 mL, 1.1 mmol), and the solution was stirred under argon at 0°C for 5 min then room temperature for 3 h. The solution was concentrated, dissolved in EtOAc (40 mL), washed with saturated aqueous NaHCO_3 (5 X 10 mL) and brine (10 mL) then dried (MgSO_4) and concentrated. The yellow foam (150 mg) that was obtained was used in the next step without further purification.

B. To a solution of the yellow foam from above (150 mg) in MeOH (3.5 mL) was added 10 N NaOH (0.3 mL), and the solution was stirred at room temperature for 40 min then concentrated. The residue was diluted with water (10 mL) and neutralized to pH 7 with 10% aqueous HCl then the mixture was extracted with EtOAc (35 mL). The organic extract was washed with brine (10 mL) then dried (MgSO₄) and concentrated to afford N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide (Compound No. 57, 124 mg, 99% over 2 steps) as a light yellow solid. ¹H NMR (CD₃OD): δ 8.21 (d, *J* = 9.3 Hz, 1H), 6.93 (d, *J* = 2.4 Hz, 2H), 6.49 (t, *J* = 2.1 Hz, 1H), 4.25 (m, 1H), 3.42 (d, *J* = 12 Hz, 1H), 2.36 (m, 1H), 2.04 (s, 3H), 1.69-1.21 (m, 8H), 1.07 (m, 1H), 1.02 (s, 3H), 0.85 (s, 6H), 0.68 (d, *J* = 6.6 Hz, 3H). ES-MS *m/z* 388 ([M+1]⁺).

EXAMPLE 37


Synthesis of 5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-hydroxyethyl}benzene-1,3-diol (Compound No. 58)



A solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dihydroxyphenyl)ethan-1-one (Compound No. 48, 473 mg, 1.37 mmol) in THF (14 mL) under argon was stirred at 0°C while LiAlH₄ (182 mg, 4.80 mmol) was added. The solution was stirred at 0°C for 0.5 h then room temperature for 1.5 h. The solution was treated with water (1 mL), acidified to pH 3 using 10% aqueous HCl then diluted with EtOAc (20 mL). The aqueous layer was extracted with EtOAc (2 x 20 mL), and the combined organic layers were washed with brine (10 mL) then dried (MgSO₄) and concentrated. The residue was heated to reflux in CH₂Cl₂ (15 mL) for 5 min then allowed to cool to room temperature. The insoluble material was collected by filtration to afford 5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-hydroxyethyl}benzene-1,3-diol (Compound No. 58, 375 mg, 79%) as a colorless solid (1:1 mixture of epimers). ¹H NMR (CD₃OD): δ 6.30 (m, 4H), 6.16 (t, *J* = 2.1 Hz, 1H), 6.14 (t, *J* = 2.1 Hz, 1H), 4.41 (m, 2H), 1.88-0.68 (m, 19H), 0.59 (m, 2H), 0.15 (m, 2H). ES-MS *m/z* 329 ([M-17]⁺).

EXAMPLE 38


Synthesis of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-ol (Compound No. 59)



A solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-one (Compound No. 46, 128 mg, 0.344 mmol) in THF (3.5 mL) under argon was stirred at 0°C while LiAlH₄ (13 mg, 0.34 mmol) was added. The solution was stirred at 0°C for 15 min then water (0.013 mL) was added followed by 15% aqueous NaOH (0.013 mL) and water (0.039 mL). The mixture was stirred at room temperature for 10 min then diluted with EtOAc (10 mL), dried (MgSO₄), filtered and concentrated to give 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-ol (Compound No. 59, 124 mg, 96%) as a colorless oil.

EXAMPLE 39

Synthesis of (4aR,5S,6S,8aS)-5-[2-(3,5-dimethoxyphenyl)ethyl]-1,1,4a,6-tetramethyl-decahydronaphthalene (Compound No. 61)




A. A solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-1-(3,5-dimethoxyphenyl)ethan-1-ol (Compound No. 59, 98 mg, 0.26 mmol) in THF (1.7 mL) under argon was stirred at room temperature while NaH (60% dispersion in mineral oil, 34 mg, 0.85 mmol) was added. The mixture was heated to 40°C for 1 h then cooled to 0°C and CS₂ (0.094 mL, 1.6 mmol) was added. The mixture was stirred at 0°C for 20 min then at room temperature for 25 min, and

Mel (0.098 mL, 1.6 mmol) was added. After stirring at room temperature for 1.5 h, the mixture was concentrated then partitioned between EtOAc (35 mL) and water (10 mL). The organic phase was washed with water (2 x 10 mL) and brine (10 mL) then dried (MgSO_4) and concentrated to give the desired xanthate (Compound No. 60, 147 mg) as a yellow oil that was used in the next step without further purification.

B. A solution of the yellow oil from above (Compound No. 60, 147 mg) and tributyltin hydride (0.10 mL, 0.38 mmol) in toluene (1.7 mL) was heated to reflux under argon then 1,1'-azobis(cyclohexanecarbonitrile) (VAZO®) (6 mg, 0.02 mmol) was added. The mixture was heated at reflux for 1.5 h, additional 1,1'-azobis(cyclohexanecarbonitrile) (14 mg, 0.057 mmol) was added and heating was continued for 3 h. The mixture was concentrated, and the residue was purified by chromatography on silica gel (EtOAc/hexanes, 2:98) to give (4aR,5S,6S,8aS)-5-[2-(3,5-dimethoxyphenyl)ethyl]-1,1,4a,6-tetramethyl-decahydronaphthalene as a colorless oil (Compound No. 61, 79 mg, 84% over 2 steps). ^1H NMR (CDCl_3): δ 6.34 (d, $J = 2.1$ Hz, 2H), 6.29 (t, $J = 2.4$ Hz, 1H), 3.78 (s, 6H), 2.59 (td, $J = 13, 5.3$ Hz, 1H), 2.44 (td, $J = 13, 6.0$ Hz, 1H), 1.82-0.77 (m, 26H), 0.54 (m, 1H).

EXAMPLE 40

Synthesis of 5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethyl}benzene-1,3-diol (Compound No. 62)




A solution of (4aR,5S,6S,8aS)-5-[2-(3,5-dimethoxyphenyl)ethyl]-1,1,4a,6-tetramethyl-decahydronaphthalene (Compound No. 61, 79 mg, 0.22 mmol) in CH_2Cl_2 (2.2 mL) was cooled to -78°C under argon, BBr_3 (1.0 M in CH_2Cl_2 , 1.10 mL, 1.10 mmol) was added, and the solution was stirred at room temperature for 15 h. Methanol (5 mL) was carefully added to the solution then concentrated (2x). The residue was dissolved in *i*-PrOH/10% aqueous HCl (3:1, 3.0 mL) and heated to 85°C for 45 min. The solution was concentrated and the residue was partitioned between EtOAc (30 mL) and water (5 mL). The organic layer was washed with water (5 mL) and brine (5

mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{MeOH}/\text{CH}_2\text{Cl}_2$, 1:24) to afford 5-{2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethyl}benzene-1,3-diol (Compound No. 62, 53 mg, 73%) as a colorless foam. ^1H NMR (CDCl_3): δ 6.25 (d, $J = 2.1$ Hz, 2H), 6.17 (t, $J = 2.3$ Hz, 1H), 4.64 (s, 2H), 2.53 (td, $J = 13, 5.1$ Hz, 1H), 2.38 (td, $J = 13, 6.0$ Hz, 1H), 1.80-0.77 (m, 26H), 0.53 (m, 1H). ES-MS m/z 331 ($[\text{M}+1]^+$).

EXAMPLE 41


Synthesis of (4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbonitrile (Compound No. 63)



A suspension of (3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 7, 702 mg, 1.97 mmol) and 2,4,6-triisopropylbenzenesulfonyl hydrazide (705 mg, 2.36 mmol) in MeOH (6.6 mL) was stirred at room temperature under argon for 1 h 10 min then KCN (449 mg, 6.89 mmol) was added. After heating to reflux for 2 h 45 min, the mixture was allowed to cool to room temperature then concentrated. The residue was partitioned between CH_2Cl_2 (20 mL) and water (20 mL), and the aqueous phase was extracted with CH_2Cl_2 (2 x 10 mL). The combined organic layers were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{EtOAc}/\text{hexanes}$, 1:9) to give (4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbonitrile (Compound No. 63) (contaminated with (4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one) as a colorless foam (242 mg).

EXAMPLE 42


Synthesis of (3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide (Compound No. 64)



A mixture containing (4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbonitrile (Compound No. 63) from above (196 mg) and solid KOH (450 mg, 8.0 mmol) in ethylene glycol/water (4:1, 2.5 mL) was heated to 120°C under argon for 22 h then allowed to cool to room temperature. The mixture was adjusted to pH 1 using 1 N HCl then partitioned between water (40 mL) and EtOAc (30 mL). The organic phase was washed with water (10 mL), and the combined aqueous layers were extracted with EtOAc (20 mL). The combined organic layers were washed with brine (2 x 15 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:24) to give (4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide (Compound No. 64, 116 mg, 19% over 2 steps) as a colorless solid. ¹H NMR (CDCl₃): δ 7.30 (s, 1H), 7.24 (m, 1H), 6.91 (s, 1H), 5.47 (br s, 1H), 5.36 (br s, 1H), 3.85 (s, 3H), 3.01 (d, *J* = 11 Hz, 1H), 2.43 (m, 1H), 2.41 (s, 3H), 2.01 (m, 1H), 1.86 (m, 1H), 1.66 (m, 1H), 1.51-1.37 (m, 2H), 1.30-1.00 (m, 8H), 0.90 (s, 3H), 0.86 (s, 3H), 0.80 (d, *J* = 6.6 Hz, 3H).

EXAMPLE 43


Synthesis of (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide (Compound No. 65)



A suspension of (4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide (Compound No. 64, 116 mg, 0.301 mmol) in CH_2Cl_2 (3.0 mL) was cooled to -78°C under argon, BBr_3 (1.0 M in CH_2Cl_2 , 2.30 mL, 2.30 mmol) was added, and the solution was stirred at room temperature for 23 h. Methanol (10 mL) was carefully added to the solution then concentrated. The residue was dissolved in *i*-PrOH/10% aqueous HCl (3:1, 4 mL) and heated to 80°C for 30 min. The solution was concentrated, diluted with water (10 mL) then adjusted to pH 6 using 1 N NaOH. The mixture was extracted with EtOAc (30 mL), and the organic layer was washed with brine (10 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (MeOH/ CH_2Cl_2 , 3:97) to afford (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxamide (Compound No. 65, 46 mg, 41%) as a yellow solid. ^1H NMR (CD_3OD): δ 7.24 (s, 1H), 7.12 (s, 1H), 6.85 (m, 1H), 3.11 (d, $J = 11$ Hz, 1H), 2.38 (m, 1H), 2.35 (s, 3H), 2.12 (td, $J = 12, 4.4$ Hz, 1H), 1.79 (m, 1H), 1.71-1.03 (m, 11H), 0.92 (s, 3H), 0.88 (s, 3H), 0.73 (d, $J = 6.3$ Hz, 3H). ES-MS m/z 372 ($[\text{M}+1]^+$).

EXAMPLE 44

Synthesis of (3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 67)



A. To a suspension of (methoxymethyl)triphenylphosphonium chloride (687 mg, 2.00 mmol) in THF (5.0 mL) at 0°C under argon was added potassium *t*-butoxide (1.0 M in THF, 2.00 mL, 2.00 mmol) dropwise. The mixture was stirred at 0°C for 25 min then a solution of (3R,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 7, 357 mg, 1.00 mmol) in THF (2.9 mL) was added dropwise at 0°C. The mixture was stirred at 0°C for 5 min then room temperature for 1 h 40 min. The mixture was diluted with EtOAc (30 mL), washed with brine (10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:19) to give [(1S,4aS,8aS)-3-(methoxymethylidene)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone (Compound No. 66, 373 mg, 97%) as colorless crystals.


B. To a solution of [(1S,4aS,8aS)-3-(methoxymethylidene)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3-methoxy-5-methylphenyl)methanone (Compound No. 66, 373 mg, 0.970 mmol) in THF (8 mL) was added 2 N HCl (2 mL), and the mixture was heated to reflux for 0.5 h. The mixture was diluted with EtOAc (30 mL), washed with brine (10 mL) then dried (MgSO₄) and concentrated to give a colorless oil (389 mg) that was used in the next step without further purification.

C. To a solution of the colorless oil from above (389 mg) in CH₃CN (2 mL) was added aqueous NaH₂PO₄ (pH 4.2, 1.1 mL) followed by 50% aqueous H₂O₂ (0.073 mL, 1.3 mmol) and a solution of 80% NaClO₂ (186 mg, 1.65 mmol) in water (3.3 mL). The mixture was stirred at room temperature for 27 h then diluted with EtOAc (20 mL) and water (5 mL). Excess peroxide was destroyed with solid Na₂SO₃, and the aqueous phase was acidified to pH 2 with concentrated HCl then extracted with EtOAc (2 x 10 mL). The combined organic layers were dried (MgSO₄) and concentrated to

give impure (3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 67, 395 mg) as a colorless foam.

EXAMPLE 45


Synthesis of (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 68)



A solution of impure (3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid from above (Compound No. 67, 395 mg) in CH_2Cl_2 (10 mL) was cooled to -78°C under argon, BBr_3 (1.0 M in CH_2Cl_2 , 4.9 mL, 4.9 mmol) was added, and the solution was stirred at room temperature for 4 h. Methanol (10 mL) was carefully added to the solution then concentrated (4X). The residue was dissolved in *i*-PrOH/10% aqueous HCl (2.5:1, 14 mL) and heated to 85°C for 45 min. The solution was concentrated and the residue was partitioned between EtOAc (30 mL) and water (10 mL). The aqueous phase was extracted with EtOAc (10 mL), and the combined organic layers were dried (MgSO_4) and concentrated. The residue was dissolved in 1 N NaOH (2 mL) and diluted with water (20 mL) then extracted with EtOAc (2 x 10 mL). The aqueous phase was acidified to pH 1 using 1 N HCl and extracted with CH_2Cl_2 (4 x 8 mL). The combined CH_2Cl_2 extracts were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (MeOH/ CH_2Cl_2 , 3:97) to afford (3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 68, 174 mg, 48%) as colorless crystals. ^1H NMR (CDCl_3): δ 7.29 (s, 1H), 7.20 (s, 1H), 6.87 (s, 1H), 2.98 (d, J = 11 Hz, 1H), 2.50 (m, 1H), 2.38 (s, 3H), 2.23 (td, J = 12, 4.2 Hz, 1H), 1.93 (m, 1H), 1.73-1.01 (m, 12H), 0.91 (s, 3H), 0.86 (s, 3H), 0.81 (d, J = 6.3 Hz, 3H). ES-MS m/z 373 ($[\text{M}+1]^+$).

EXAMPLE 46

Synthesis of (1R,2R,4aS,8aS)-1-[[[3-methoxy-5-methylphenyl)sulfanyl]methyl]-2,5,5,8a-tetramethyl-decahydro-naphthalen-2-ol (Compound No. 75)



A. To a mixture of 3-methoxy-5-methylphenol (Compound No. 69, 3.39 g, 24.5 mmol) and 1,4-diazabicyclo[2.2.2]octane (5.50 g, 49.0 mmol) in DMF (20 mL) under argon was added dimethylthiocarbamoyl chloride (Compound No. 70, 6.06 g, 49.0 mmol) portionwise. The reaction mixture was stirred at room temperature overnight and 500 mL of ether was added. The organics were washed with water (150 mL), brine (2 × 50 mL), dried (Na₂SO₄), and concentrated. Purification by chromatography on silica gel (Hexanes/EtOAc, 4:1) gave O-3-methoxy-5-methylphenyl dimethylcarbamothioate (Compound No. 71, 5.05 g, 91%) as a colorless solid.

B. O-3-Methoxy-5-methylphenyl dimethylcarbamothioate (Compound No. 71, 5.0 g, 22.2 mmol) was vacuum-degassed (4 cycles, flask was backfilled with nitrogen) and heated to 280°C for 4 h, then cooled to room temperature. Purification by chromatography on silica gel (Hexanes/EtOAc, 85:15) gave S-3-methoxy-5-methylphenyl dimethylcarbamothioate (Compound No. 72, 4.10 g, 82%) as a light yellow oil.


C. A mixture of S-3-methoxy-5-methylphenyl dimethylcarbamothioate (Compound No. 72, 4.10 g, 18.2 mmol) and KOH (6.0 g) in MeOH (200 mL) was refluxed for 2 h. After cooling, the mixture was concentrated and diluted with EtOAc (500 mL). The organics were washed with 1N HCl (50 mL), brine (2 × 50 mL), dried (Na₂SO₄), and concentrated to give 3-methoxy-5-methylbenzenethiol (Compound No. 73, 2.80 g, quant.) as a colorless oil.

D. To a mixture of (1S,2R,4aS,8aS)-1-(hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 26, 4.66 g, 19.39 mmol) and Et₃N (7.01 mL, 50.41 mmol) in DMF (40 mL) under argon was added methanesulfonyl chloride (2.66 g, 23.27 mmol) dropwise at 0°C. After stirring at room temperature for 3 h, ether (400 mL) was added. The organics were washed with 1N HCl (20 mL), saturated aqueous NaHCO₃ (20 mL), brine (2 × 20 mL), dried (Na₂SO₄), and concentrated to give ((1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyldecahydronaphthalen-1-yl)methyl methanesulfonate (Compound No. 74, 5.90 g, 96%) as a light yellow sticky mass.

E. A mixture of 3-methoxy-5-methylbenzenethiol (Compound No. 73, 3.14 g, 20.36 mmol), ((1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyldecahydronaphthalen-1-yl)methyl methanesulfonate (Compound No. 74, 6.48 g, 20.36 mmol), and Cs₂CO₃ (19.90 g, 61.08 mmol) in CH₃CN (300 mL) under argon was stirred at room temperature for 1 h, then refluxed for 24 h. After cooling, the mixture was concentrated. Purification by chromatography on silica gel (Hexanes/EtOAc, 4:1) gave (1R,2R,4aS,8aS)-1-[(3-methoxy-5-methylphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydro-naphthalen-2-ol (Compound No. 75, 5.70 g, purity ~75%) as a yellow sticky mass.

EXAMPLE 47

Synthesis of (1R,2R,4aS,8aS)-1-[(3-methoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 76)




A mixture of 3-methoxythiophenol (2.20 g, 15.72 mmol), [(1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]methyl methanesulfonate (Compound No. 74, 5.0 g, 15.72 mmol), and Cs₂CO₃ (15.30 g, 47.16 mmol) in CH₃CN (600 mL) under argon was stirred at room temperature for 1 h, then refluxed for 24 h. After cooling, the mixture was concentrated and purified by column chromatography on

silica gel (Hexanes/EtOAc, 10:1 to 6:1) to give (1R,2R,4aS,8aS)-1-[[[(3-methoxyphenyl)sulfanyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 76, 4.03 g, 71%) as a yellow oil. ¹H NMR (CDCl₃): δ 7.20 (t, 1H), 6.95 (m, 2H), 6.71 (d, 1H), 3.78 (s, 3H), 3.02 (ABq, 2H), 2.00 (s, 1H), 1.90 (m, 1H) 1.78-1.22 (m, 8H), 1.21 (s, 3H), 1.20- 0.98 (m, 3H), 0.90 (s, 3H), 0.85 (s, 3H), 0.80 (s, 3H).

EXAMPLE 48

Synthesis of (1R,2R,4aS,8aS)-1-[[[(3,5-dimethoxyphenyl)-sulfanyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 80)



A. To a mixture of 3,5-dimethoxyphenol (7.70 g, 50 mmol) and 1,4-diazabicyclo[2.2.2]octane (11.22 g, 100 mmol) in DMF (40 mL) under argon was added dimethylthiocarbamoyl chloride (Compound No. 70, 12.36 g, 100 mmol) portionwise. The reaction mixture was stirred at room temperature overnight and 500 mL of ether was added. The organics were washed with water (150 mL), brine (2 × 50 mL), dried (Na₂SO₄), and concentrated. Purification by chromatography on silica gel (Hexanes/EtOAc, 4:1) gave O-3,5-dimethoxyphenyl dimethylcarbamothioate (Compound No. 77, 10.88 g, 90%) as a white solid.


B. O-3,5-Dimethoxyphenyl dimethylcarbamothioate (Compound No. 77, 1.0 g, 4.14 mmol) was vacuum-degassed (4 cycles, flask was backfilled with nitrogen) and heated to 260°C for 3 h, then cooled to room temperature to give S-3,5-dimethoxyphenyl dimethylcarbamothioate (Compound No. 78, 1.0 g, quant.) as a light brown oil.

C. A mixture of S-3,5-dimethoxyphenyl dimethylcarbamothioate (Compound No. 78, 1.0 g, 4.14 mmol) and KOH (1.5 g) in MeOH (20 mL) was refluxed for 2 h. After cooling, it was concentrated. EtOAc (100 mL) was added. The organics were washed with 1N HCl (20 mL), brine (3 × 20 mL), dried (Na₂SO₄), and concentrated to give 3,5-dimethoxybenzenethiol (Compound No. 79, 0.70 g, quant.) as a light brown oil.

D. A mixture of 3,5-dimethoxybenzenethiol (Compound No. 79, 6.74 g, 39.60 mmol), ((1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyldecahydronaphthalen-1-yl)methyl methanesulfonate (Compound No. 74, 12.62 g, 39.60 mmol), and Cs₂CO₃ (38.70 g, 118.77 mmol) in CH₃CN (560 mL) under argon was stirred at room temperature for 1 h, then refluxed for 24 h. After cooling, the mixture was concentrated. Purification by chromatography on silica gel (Hexanes/EtOAc, 4:1) gave (1R,2R,4aS,8aS)-1-{[(3,5-dimethoxyphenyl)-sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 80, 9.40 g, 60%) as a light yellow sticky mass. ¹H NMR (CDCl₃): δ 6.52 (d, 2H), 6.30 (d, 1H), 3.78 (2s, 6H), 3.01 (AB, 2H), 1.90-1.30 (m, 10H), 1.22 (s, 3H), 1.20-0.98 (m, 3H), 0.90 (2s, 6H), 0.82 (s, 3H).

EXAMPLE 49

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide (Compound No. 81)




A mixture of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 125 mg, 0.33 mmol),

benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (230 mg, 0.440 mmol), *p*-toluic acid (52 mg, 0.38 mmol), *N,N*-diisopropylethylamine (0.15 mL, 0.86 mmol) and DMF (3.5 mL) under argon was allowed to stir at room temperature for 18 h. EtOAc (20 mL) was added to the solution, and the mixture washed with saturated aqueous NaHCO₃ (3 x 10 mL) and saturated aqueous NaCl (2 x 10 mL). The organic layer was concentrated and the residue was purified by flash chromatography using hexanes/EtOAc (3:1) to afford the product, N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide (Compound No. 81, 100 mg, 62%) as a clear colourless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.66 (d, *J* = 8.0 Hz, 2H), 7.26 (d, *J* = 7.9 Hz, 2H), 7.03 (s, 2H), 6.63 (s, 1H) 6.47 (d, *J* = 9.2 Hz, 1H), 4.56 (s, 1H), 3.83 (s, 6H), 3.10 (d, *J* = 11.7 Hz, 1H), 2.54 (m, 1H), 2.41 (s, 3H), 1.01 (s, 3H), 1.3 (m, 7H), 0.85 (s, 3H), 0.83 (s, 3H), 0.79 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 203.0, 167.7, 161.1, 142.9, 142.0, 132.9, 129.6, 127.0, 106.4, 104.4, 58.9, 55.8, 49.9, 49.6, 42.0, 41.5, 39.6, 33.9, 33.6, 33.2, 27.9, 22.9, 21.2, 18.6, 16.6, 13.7; MS *m/z*: 492.4 ([M+1]⁺).

EXAMPLE 50

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide (Compound No. 82)




A solution of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide (Compound No. 81, 71 mg, 0.14 mmol) in CH₂Cl₂ (1.5 mL) under argon was cooled to -78°C. Boron tribromide (1 M in CH₂Cl₂, 1.0 mL, 1.0 mmol) was added and the mixture was stirred at -78°C for 10 min. The reaction was warmed up to room temperature and stirred for 19 h. Methanol (3 mL) was carefully added and the resultant mixture was concentrated *in vacuo*. This was repeated three times. Isopropanol/1 M HCl (3:1, 2 mL) was added, and the solution was heated to 85°C for 45 min. The solution was

concentrated, EtOAc (10 mL) and water (5 mL) were added, and the pH was adjusted to approximately 7 (1 M HCl and 1 M NaOH). The mixture was extracted with EtOAc (2 x 10 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by flash chromatography using CH₂Cl₂/MeOH (20:1 to 10:1, EtOAc to load) afforded the impure product (53 mg, 82%) as a white solid. Additional purification via radial chromatography using hexanes/EtOAc (5:2) afforded pure N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylbenzamide (Compound No. 82, 27 mg, 42%). ¹H NMR (300 MHz, CD₃OD) δ 8.30 (d, *J* = 10.0 Hz, 1H), 7.70 (d, *J* = 8.1 Hz, 2H), 7.29 (d, *J* = 8.1 Hz, 2H), 6.94 (s, 2H), 6.48 (s, 1H), 4.87 (s, 2H), 3.59 (d, *J* = 12.3 Hz, 1H), 3.30 (s, 1H), 2.47 (m, 1H), 2.40 (s, 3H), 1.77 (m, 2H), 1.35 (m, 3H), 1.04 (s, 4H), 0.87 (s, 9H), 0.76 (d, *J* = 6.5 Hz, 3H); ¹³C NMR (75 MHz, CD₃OD) δ 206.4, 171.8, 159.8, 144.6, 139.4, 132.9, 129.3, 129.1, 125.7, 108.0, 58.9, 51.6, 42.9, 42.4, 40.2, 34.1, 33.9, 33.8, 28.9, 22.1, 21.4, 19.5, 16.7, 14.2; MS *m/z*: 464.3 ([M+1]⁺).

EXAMPLE 51

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide (Compound No. 83)




A mixture of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 125 mg, 0.33 mmol), benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (230 mg, 0.440 mmol), *m*-toluic acid (52 mg, 0.38 mmol), *N,N*-diisopropylethylamine (0.15 mL, 0.86 mmol) and DMF (3.5 mL) under argon was allowed to stir at room temperature for 20 h. EtOAc (20 mL) was added to the solution, and the mixture washed with saturated aqueous NaHCO₃ (3 x 10 mL) and saturated aqueous NaCl (2 x 10 mL). The organic layer was concentrated and the residue was purified by flash chromatography using hexanes/EtOAc (3:1) to afford the product N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-

dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide (Compound No. 83, 95 mg, 59%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 7.49 (s, 1H), 7.44 (d, J = 7.4 Hz, 1H), 7.21 (d, J = 4.7 Hz, 2H), 7.02 (d, J = 2.2 Hz, 2H), 6.60 (d, J = 9.6 Hz, 1H), 6.57 (d, J = 1.9 Hz, 1H), 4.54 (m, 1H), 3.75 (s, 6H), 3.18 (d, J = 12.2 Hz, 1H), 2.48 (m, 1H), 2.30 (s, 3H), 1.70 (m, 2H), 1.16 (m, 3H), 1.05 (s, 4H), 0.85 (d, J = 2.5 Hz, 6H), 0.81 (s, 3H), 0.78 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 202.9, 167.9, 160.9, 142.8, 138.8, 135.7, 132.2, 128.7, 127.9, 123.7, 106.3, 104.3, 58.7, 55.7, 49.8, 49.5, 41.9, 41.4, 39.5, 33.8, 33.5, 33.1, 27.8, 21.8, 21.5, 18.6, 16.5, 13.6; MS m/z : 492.4 ($[\text{M}+1]^+$).

EXAMPLE 52

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide (Compound No. 84)




A solution of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide (Compound No. 83, 110 mg, 0.22 mmol) in CH_2Cl_2 (2.2 mL) under argon was cooled to -78°C . Boron tribromide (1 M in CH_2Cl_2 , 1.1 mL, 1.1 mmol) was added and the mixture was stirred at -78°C for 10 min. The reaction was warmed up to room temperature and stirred overnight. Methanol (3 mL) was carefully added and the resultant mixture was concentrated *in vacuo*. This was repeated three times. Isopropanol/1 M HCl (3:1, 4 mL) was added, and the solution was heated to 85°C for 45 min. The solution was concentrated, EtOAc (20 mL) and water (10 mL) were added, and the pH was adjusted to approximately 7 (1 M HCl and 1 M NaOH). The mixture was extracted with EtOAc (2 x 20 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by flash chromatography using $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (20:1, EtOAc to load) afforded the impure product (92 mg, 89%) as a white solid. Additional purification via radial chromatography using hexanes/EtOAc (5:2) afforded pure N-

[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylbenzamide (Compound No. 84, 56 mg, 54%). ^1H NMR (300 MHz, CD_3OD) δ 8.35 (d, J = 9.3 Hz, 1H), 7.56 (s, 2H), 7.35 (s, 2H), 6.95 (s, 2H), 6.49 (d, J = 1.5 Hz, 1H), 4.88 (s, 2H), 4.52 (s, 1H), 3.59 (d, J = 12.0 Hz, 1H), 3.30 (s, 1H), 2.41 (s, 3H), 1.99 (s, 1H), 1.78 (s, 2H), 1.45 (q, J = 15.6, 6.3 Hz, 2H), 1.28 (m, 4H) 1.04 (s, 6H), 0.78 (s, 3H), 0.75 (s, 3H); ^{13}C NMR (75 MHz, CD_3OD) δ 206.4, 171.8, 159.8, 144.6, 139.4, 137.0, 132.9, 129.3, 129.1, 125.7, 107.9, 58.5, 51.6, 42.9, 42.4, 40.2, 34.1, 33.9, 28.9, 22.1, 21.4, 19.5, 16.7, 14.2; MS m/z : 464.3 ($[\text{M}+1]^+$).

EXAMPLE 53

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide (Compound No. 85)




A mixture of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 125 mg, 0.33 mmol), benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (230 mg, 0.440 mmol), *o*-toluic acid (52 mg, 0.38 mmol), *N,N*-diisopropylethylamine (0.15 mL, 0.86 mmol) and DMF (3.5 mL) under argon was allowed to stir at room temperature for 20 h. EtOAc (20 mL) was added to the solution, and the mixture washed with saturated aqueous NaHCO_3 (3 x 10 mL) and saturated aqueous NaCl (2 x 10 mL). The organic layer was concentrated and the residue was purified by flash chromatography using hexanes/EtOAc (3:1) to afford the product N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide (Compound No. 85, 100 mg, 62%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 7.66 (d, J = 8.1 Hz, 2H), 7.27 (d, J = 7.8 Hz, 2H), 7.04 (s, 2H), 6.62 (s, 1H), 6.31 (d, J = 8.1 Hz, 1H), 4.55 (m, 1H), 3.83 (s, 6H), 3.10 (d, J = 11.7 Hz, 1H), 2.53 (m, 1H), 2.41 (s, 3H), 1.80 (m, 2H), 1.41 (m, 1H), 1.27 (m, 4H), 1.15 (m, 2H), 1.10 (s, 3H), 0.86 (d, J = 4.80 Hz, 6H), 0.84 (s, 3H), 0.81 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ

203.0, 170.2, 161.1, 142.8, 137.8, 136.0, 131.2, 130.0, 126.8, 126.2, 106.5, 104.4, 55.9, 49.9, 49.3, 42.1, 41.6, 39.6, 33.9, 33.3, 33.2, 28.0, 22.0, 18.7, 16.7, 14.5, 13.8; MS m/z : 492.4 ($[M+1]^+$).

EXAMPLE 54

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide (Compound No. 86)




A solution of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide (Compound No. 85, 46 mg, 0.09 mmol) in CH_2Cl_2 (1.0 mL) under argon was cooled to -78°C . Boron tribromide (1 M in CH_2Cl_2 , 0.7 mL, 0.7 mmol) was added and the mixture was stirred at -78°C for 10 min. The reaction was warmed up to room temperature and stirred overnight. Methanol (3 mL) was carefully added and the resultant mixture was concentrated *in vacuo*. This was repeated three times. Isopropanol/1 M HCl (3:1, 2 mL) was added, and the solution was heated to 85°C for 45 min. The solution was concentrated, EtOAc (10 mL) and water (5 mL) were added, and the pH was adjusted to approximately 7 (1 M HCl and 1 M NaOH). The mixture was extracted with EtOAc (2 x 10 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by flash chromatography using $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (20:1, EtOAc to load) afforded the impure product (36 mg, 86%) as a white solid. Additional purification of this material with 20 mg from another reaction via radial chromatography using hexanes/EtOAc (5:2) afforded pure N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-2-methylbenzamide (Compound No. 86). ^1H NMR (300 MHz, CD_3OD) δ 8.67 (d, J = 9.3 Hz, 1H), 7.26 (m, 4H), 6.92 (s, 2H), 6.46 (s, 1H), 4.88 (s, 2H), 4.49 (s, 1H), 3.51 (d, J = 12.3 Hz, 1H), 3.31 (s, 1H), 2.38 (s, 3H), 2.00 (s, 2H), 1.50 (m, 1H), 1.25 (m, 6H), 0.89 (d, J = 10.5 Hz, 6H), 0.81 (s, 3H), 0.78 (s, 3H); ^{13}C NMR (75 MHz, CD_3OD) δ 206.3,

173.4, 159.7, 144.5, 139.0, 136.1, 131.4, 130.5, 128.0, 126.7, 108.0, 58.7, 51.5, 42.9, 42.3, 40.1, 34.0, 33.9, 33.8, 28.9, 22.2, 19.7, 19.5, 16.8, 14.1; MS m/z : 464.4 ($[M+1]^+$).

EXAMPLE 55


Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide
(Compound No. 87)



A mixture of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 80 mg, 0.02 mmol), benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (135 mg, 0.260 mmol), 2-naphthoic acid (45 mg, 0.26 mmol), *N,N*-diisopropylethylamine (0.10 mL, 0.52 mmol) and DMF (2.3 mL) under argon was allowed to stir at room temperature for 15 h. EtOAc (15 mL) was added to the solution, and the mixture washed with saturated aqueous NaHCO_3 (3 x 10 mL) and saturated aqueous NaCl (2 x 10 mL). The organic layer was concentrated and the residue was purified via radial chromatography using hexanes/EtOAc (5:1) to afford pure N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide (Compound No. 87, 69 mg, 66%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 8.23 (s, 1H), 7.86 (m, 4H), 7.56 (m, 2H), 7.07 (d, $J = 2.0$ Hz, 2H), 6.62 (t, $J = 4.2, 2.1$ Hz, 1H), 6.57 (d, $J = 8.5$ Hz, 1H), 4.65 (m, 1H), 3.82 (s, 6H), 3.19 (d, $J = 11.7$ Hz, 1H), 2.59 (m, 1H), 2.01 (s, 1H), 1.83 (m, 2H), 1.13 (s, 4H), 0.96 (d, $J = 6.6$ Hz, 1H), 0.85 (m, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 203.1, 168.0, 161.1, 142.9, 134.9, 133.0, 132.9, 129.1, 128.8, 128.0, 127.8, 127.4, 127.0, 123.8, 106.5, 104.3, 58.8, 55.8, 50.1, 49.6, 42.0, 41.4, 39.7, 33.9, 33.7, 33.3, 27.9, 21.9, 18.7, 16.7, 13.7; MS m/z : 528.4 ($[M+1]^+$).

EXAMPLE 56


Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide
(Compound No. 88)



A solution of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide (Compound No. 87) (19 mg, 0.04 mmol) in CH_2Cl_2 (2.0 mL) under argon was cooled to -78°C . Boron tribromide (1 M in CH_2Cl_2 , 0.3 mL, 0.3 mmol) was added and the mixture was stirred at -78°C and allowed to warm to room temperature over 2 days. Methanol (6 mL) was carefully added and the resultant mixture was concentrated *in vacuo*. This was repeated three times. Isopropanol/1 M HCl (3:1, 4 mL) was added, and the solution was heated to 85°C for 40 min. The solution was concentrated, EtOAc (25 mL) and water (10 mL) were added, and the pH was adjusted to approximately 9 (2 M NaOH). The mixture was extracted with EtOAc (3 x 10 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by radial chromatography using hexanes/EtOAc (3:1 to 1:1 to 1:3) afforded pure N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]naphthalene-2-carboxamide (Compound No. 88, 13 mg, 72%) as a white solid. ^1H NMR (300 MHz, CD_3OD) δ 8.56 (d, $J = 9.1$ Hz, 1H), 8.30 (s, 1 H), 7.97 (m, 3 H), 7.81 (dd, $J = 8.5, 1.5$ Hz, 1 H), 7.59 (m, 2 H), 6.96 (d, $J = 2.1$ Hz, 2 H), 6.48 (t, $J = 2.0$ Hz, 1 H), 4.59 (m, 1 H), 3.65 (d, $J = 12.1$ Hz, 1 H), 2.52 (m, 1 H), 1.95-1.18 (m, 9 H), 1.07 (s, 3 H), 0.92 (s, 3 H), 0.90 (s, 3 H), 0.82 (d, $J = 7.0$ Hz, 3 H); ^{13}C NMR (75 MHz, CD_3OD) δ 205.2, 170.4, 158.6, 143.4, 135.0, 133.0, 132.8, 128.8, 128.0, 127.63, 127.56, 127.4, 126.6, 124.2, 106.8, 57.7, 50.7, 41.7, 41.2, 39.0, 33.0, 32.7, 27.7, 21.0, 18.4, 15.6, 13.1; MS m/z : 500.3 ($[\text{M}+1]^+$).

EXAMPLE 57

Synthesis of N-[(3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyrazine-2-carboxamide (Compound No. 89)




A mixture of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 145 mg, 0.42 mmol), O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (519 mg, 1.62 mmol), pyrazine-2-carboxylic acid (172 mg, 1.39 mmol), *N,N*-diisopropylethylamine (0.60 mL, 3.4 mmol) and DMF (8 mL) under argon was allowed to stir at room temperature for 22 h. EtOAc (90 mL) was added to the solution, and the mixture washed with saturated aqueous NaHCO₃ (3 x 20 mL), water (20 mL) and saturated aqueous NaCl (2 x 20 mL). The organic layer was dried over anhydrous Na₂SO₄, concentrated and the residue was purified via radial chromatography using EtOAc to afford impure N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyrazine-2-carboxamide (Compound No. 89). The impure material was dissolved in MeOH (8 mL). Aqueous NaOH (2 M, 2 mL, 4 mmol) was added and the resultant mixture was stirred at room temperature for two days before being concentrated *in vacuo*. The residue was partitioned between EtOAc (90 mL) and water (30 mL). The organic layer was washed with saturated aqueous NaHCO₃ (3 x 30 mL), saturated aqueous NaCl (20 mL) and dried over anhydrous Na₂SO₄. The organic layer was concentrated and the residue was purified via radial chromatography using a gradient of 10% EtOAc/hexanes to 100% EtOAc to afford N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyrazine-2-carboxamide (Compound No. 89, 40 mg, 21%) as a white solid. ¹H NMR (300 MHz, CD₃OD) δ 9.26 (d, *J* = 1.4 Hz, 1 H), 8.85 (d, *J* = 2.4 Hz, 1 H), 8.78 (m, 1 H), 6.88 (d, *J* = 2.1 Hz, 2 H), 6.48 (m, 1 H), 4.51 (m, 1 H), 3.38 (d, *J* = 11.5 Hz, 1 H), 2.53 (m, 1 H), 1.94-1.18 (m, 9 H), 1.09 (s, 3 H), 0.88 (s, 3 H), 0.83 (s, 3 H), 0.75 (d, *J* = 6.3 Hz, 3 H); ¹³C NMR (75 MHz, CD₃OD) δ 204.2, 163.7,

163.3, 147.6, 143.7, 143.6, 143.2, 106.5, 58.1, 50.3, 41.7, 41.1, 39.2, 33.2, 32.7, 27.4, 20.9, 18.3, 15.5, 12.9; MS m/z : 452.4 ($[M+1]^+$).

EXAMPLE 58

Synthesis of 2-[[[(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbonyl]pyridine (Compound No. 94)



A. A mixture of (3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 53, 1.74 g, 4.68 mmol) and NaOH (2 M, 2.35 mL, 4.70 mmol) in MeOH/THF (2:1, 75 mL) was heated to 80°C for 210 min. After cooling, the mixture was concentrated *in vacuo* and the resulting residue was partitioned between CH₂Cl₂ and saturated aqueous NH₄Cl. The aqueous layer was extracted with CH₂Cl₂ (3x) and the combined organic layers were washed with brine (1x), dried over anhydrous Na₂SO₄ and concentrated to afford (3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 90, 1.59 g, 91%) as a white solid.

B. To a suspension of methoxymethyl triphenylphosphonium chloride (1.37 g, 4.00 mmol) in THF (15 mL) at room temperature under argon was added potassium *t*-butoxide (448 mg, 4.00 mmol) and resultant mixture was stirred for 20 min. To the resulting red mixture was added (3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 90, 500 mg, 1.34 mmol). After 1 h at room temperature, the reaction was quenched with saturated aqueous NaHCO₃ (20 mL) and EtOAc (60 mL) was added. The organic layer was

washed with brine (3 x 20 mL), dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. Purification by flash chromatography using EtOAc/hexanes (gradient from 5% to 10% to 20%) afforded a mixture of the desired enol ethers, [(1S,2S,4aS,8aS)-3-(methoxymethylidene)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone, (Compound No. 91, 424 mg, 81%) as a white foam.

C. A mixture of the enol ethers (Compound No. 91, 183 mg, 0.457 mmol) and HCl (1M, 0.5 mL, 0.5 mmol) in THF (2 mL) was heated at reflux for 30 min. After cooling to room temperature, the mixture was diluted with EtOAc (25 mL), washed with brine (3 x 5 mL), dried over anhydrous Na₂SO₄ and concentrated *in vacuo* to afford the aldehyde, (3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbaldehyde, (Compound No. 92, dr ~ 10:1, 167 mg, 95%) as a white foam.


D. To a solution of 2-bromopyridine (91 μ L, 0.95 mmol) in THF (5 mL) at -78°C under argon was added *n*-butyllithium (1.9 M in hexanes, 0.45 mL, 0.86 mmol). After 10 min, a solution of (3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbaldehyde (Compound No. 92, 167 mg, 0.432 mmol) in THF (2 mL) at -78°C was added via cannula. After 3 h at this temperature, a saturated aqueous solution of NH₄Cl (10 mL) was added and the mixture was allowed to warm to room temperature. The mixture was extracted with EtOAc (3 x 15 mL), dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. Purification by radial chromatography with EtOAc/hexanes (gradient from 20% to 35% to 50%) afforded a ca. 1:1 mixture of the diastereomeric alcohols, [(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl](pyridin-2-yl)methanol, (Compound No. 93, 145 mg, 72%) as a thick oil.

E. A mixture of the alcohols from above (Compound No. 93, 145 mg, 0.311 mmol) and manganese dioxide (1.45 g) in CH₂Cl₂ (15 mL) under argon was stirred at room temperature overnight. Additional manganese dioxide (1.73 g) was added and the resultant mixture was stirred for the weekend. The mixture was filtered through a pad of celite and concentrated *in vacuo*. Purification by radial chromatography with EtOAc/hexanes (1:3) afforded 2-([(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl)carbonyl]pyridine (Compound No. 94, 108 mg, 75%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 8.72 (d, *J* = 4.2 Hz, 1 H), 8.10 (d, *J* = 7.8 Hz, 1 H), 7.86 (td, *J* = 7.8, 1.6 Hz, 1 H), 7.48 (m, 1 H), 7.09 (d, *J* = 2.7 Hz, 2 H), 6.63 (m, 1 H), 3.99 (td, *J* = 11.3, 3.8 Hz, 1 H), 3.85 (s, 6 H), 3.19 (d, *J* = 11.9 Hz, 1 H), 2.76 (m, 1 H), 1.87 (m, 1 H), 1.59-1.00 (m, 7 H), 0.96 (d, *J* = 6.4 Hz, 1H),

1.16 (s, 3 H), 0.86 (s, 3 H), 0.80 (s, 3 H), 0.67 (d, $J = 6.3$ Hz, 3 H); ^{13}C NMR (75 MHz, CDCl_3) δ 204.7, 204.1, 161.0, 153.6, 149.2, 143.2, 137.2, 127.2, 122.7, 106.3, 104.8, 62.8, 55.8, 54.5, 50.7, 42.2, 41.2, 39.4, 33.7, 33.6, 32.1, 31.8, 26.3, 22.0, 19.2, 18.7, 14.3; MS m/z : 464.1 ($[\text{M}+1]^+$).

EXAMPLE 59

Synthesis of 5-[[[(1S,2R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-yl)carbonyl]-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 95)




A solution of 2-[[[(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbonyl]pyridine (Compound No. 94, 73 mg, 0.16 mmol) in CH_2Cl_2 (10 mL) under argon was cooled to -78°C . Boron tribromide (1 M in CH_2Cl_2 , 0.8 mL, 0.8 mmol) was added and the mixture was stirred at -78°C and allowed to warm to room temperature overnight. Methanol (10 mL) was carefully added and the resultant mixture was concentrated *in vacuo*. This was repeated three times. Isopropanol/1 M HCl (3:1, 8 mL) was added, and the solution was heated to 85°C for 45 min. The solution was concentrated, EtOAc (40 mL) and water (20 mL) were added, and the pH was adjusted to approximately 9 (2 M NaOH). The mixture was extracted with EtOAc (3 x 20 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. ^1H NMR of the resultant residue indicated that the reaction had not gone to completion. The residue was dissolved in CH_2Cl_2 (10 mL) at room temperature under argon. Boron tribromide (1 M in CH_2Cl_2 , 1.6 mL, 1.6 mmol) was added and the resultant mixture was stirred for 24 h. Methanol (10 mL) was carefully added and the resultant mixture was concentrated *in vacuo*. This was repeated three times. Isopropanol/1 M HCl (3:1, 8 mL) was added, and the solution was heated to 85°C for 45 min. The solution was concentrated, EtOAc (40 mL) and water (20 mL) were added, and the pH was adjusted to approximately 9 (2 M NaOH). The mixture was extracted with EtOAc (3 x 20 mL), dried over anhydrous Na_2SO_4 , filtered, and

concentrated *in vacuo*. Purification by radial chromatography using hexanes/EtOAc (gradient from 3:1 to 2:1) afforded pure 5-{[(1S,2R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-yl)carbonyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 95, 57 mg, 84%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 8.75 (dm, $J = 4.2$ Hz, 1 H), 8.16 (d, $J = 7.3$ Hz, 1 H), 7.93 (td, $J = 7.8, 1.3$ Hz, 1 H), 7.57 (m, 1 H), 7.21 (br s, 2 H), 6.99 (d, $J = 1.7$ Hz, 2 H), 6.70 (m, 1 H), 3.88 (td, $J = 11.4, 3.5$ Hz, 1 H), 2.95 (d, $J = 11.4$ Hz, 1 H), 2.71 (m, 1 H), 1.84 (m, 1 H), 1.69 (m, 2 H), 1.51-0.96 (m, 6 H), 1.08 (s, 3 H), 0.84 (s, 3 H), 0.76 (s, 3 H), 0.62 (d, $J = 8.0$ Hz, 3 H); ^{13}C NMR (75 MHz, CDCl_3) δ 205.9, 204.4, 157.8, 153.1, 148.9, 143.3, 138.2, 127.9, 123.4, 108.4, 108.2, 63.0, 54.1, 50.6, 42.0, 41.2, 39.4, 33.6, 32.0, 26.2, 22.0, 19.0, 18.6, 14.3; MS m/z : 436.3 ($[\text{M}+1]^+$).

EXAMPLE 60

Synthesis of N-[(2R,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide (Compound No. 102)



A. A solution of Ph₃PMeBr (863 mg, 2.42 mmol), KO^tBu (277 mg, 2.42 mmol) and THF (10 mL) was stirred at room temperature under argon. After 1 h, (3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 90, 300 mg, 0.805 mmol) was added to the bright yellow reaction mixture. After 2 h, the reaction mixture was cooled in ice then saturated NaHCO₃ solution (5 mL) was added. The mixture was diluted with EtOAc (75 mL) and was washed with half-saturated NaCl solution (2 x 15 mL). The EtOAc solution was dried over MgSO₄, filtered and concentrated. Purification using radial chromatography, eluting with 5% EtOAc/Hexanes afforded [(1S,2S,4aS,8aS)-2,5,5,8a-tetramethyl-3-methylidene-decahydronaphthalen-1-yl](3,5-

dimethoxyphenyl)methanone as a colourless film (Compound No. 96, 258 mg, 87% yield).

B. Borane (1.4 mL of a 1 M solution in THF, 1.4 mmol) was added to a -78°C solution of [(1S,2S,4aS,8aS)-2,5,5,8a-tetramethyl-3-methylenedecahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 96, 258 mg, 0.69 mmol) in THF (5 mL) under argon. The reaction was allowed to warm to room temperature. After 2 h, the reaction was cooled in ice then water (5 mL) was slowly added followed by NaBO₃ (215 mg). After 15 min, the ice bath was removed and the reaction was allowed to continue at room temperature for 2 h. The reaction mixture was diluted with EtOAc (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The EtOAc solution was dried over MgSO₄, filtered and concentrated. Purification using radial chromatography, eluting with 25% EtOAc/Hexanes afforded a mixture of diastereomers of [(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanol as a white foam (Compound No. 97, 216 mg, 80% yield).

C. Methanesulfonyl chloride (0.13 mL, 1.67 mmol) was added to an ice cold solution of [(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanol (Compound No. 97, 216 mg, 0.55 mmol) in pyridine (5 mL) under argon. After 1 h, saturated NaHCO₃ solution (5 mL) was added and the resulting solution was stirred at room temperature for 20 min. The mixture was diluted with EtOAc (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The EtOAc solution was dried over MgSO₄, filtered and concentrated. Residual pyridine was co-distilled with toluene (3 x 15 mL) to afford the desired mesylates as a white foam (253 mg, 97% yield).

D. A solution of the above mesylates (253 mg, 0.54 mmol), NaN₃ (106 mg, 1.6 mmol) and DMF (5 mL) under argon was heated at 60°C for 2 days. The reaction mixture was cooled in ice, then saturated NaHCO₃ solution (2 mL) was added. The mixture was diluted with Et₂O (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The Et₂O solution was dried over MgSO₄, filtered and concentrated to afford [(1S,2R,4aS,8aS)-3-(azidomethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone as a brown oil (Compound No. 98, 236 mg, quantitative yield).

E. A solution of [(1S,2R,4aS,8aS)-3-(azidomethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 98, 263 mg, 0.54 mmol), triphenylphosphine (715 mg, 2.7 mmol), water (0.05 mL 2.7 mmol)

and THF (5 mL) was stirred at room temperature for 19 h then was heated at reflux for 5 h. The reaction was cooled to below room temperature then was concentrated. Purification using radial chromatography, eluting with 5% MeOH/CH₂Cl₂ then 5% MeOH/CH₂Cl₂ with 2% Et₃N afforded [(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanamine as an off-white solid (Compound No. 99, 263 mg, quantitative yield).


F. Acetic anhydride (0.15 mL, 1.6 mmol) was added to room temperature solution of [(3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanamine (Compound No. 99, 263 mg, 0.54 mmol) in pyridine (5 mL) under argon. After 17 h, the reaction mixture was cooled in ice then saturated NaHCO₃ solution (1.5 mL) was added. The mixture was diluted with EtOAc (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The EtOAc solution was dried over MgSO₄, filtered and concentrated. Residual pyridine was co-distilled with toluene (3 x 15 mL). Purification using radial chromatography, eluting with 2% MeOH/CH₂Cl₂ afforded N-[(2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide as a white solid (Compound No. 100, 104 mg, 44% yield from Compound No. 97) and N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide as a white solid (Compound No. 101, 34 mg, 16% yield from Compound No. 97).

G. BBr₃ (1.2 mL of a 1 M solution in CH₂Cl₂) was added to a -78°C solution of N-[(2R,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide (Compound No. 100, 102 mg, 0.237 mmol) in CH₂Cl₂ (5 mL) under argon. The reaction mixture was allowed to warm to room temperature while allowed to react for 24 h. MeOH (10 mL) was added and the resulting mixture concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL), then was heated at 85°C for 45 min. The mixture was cooled to below room temperature and then was concentrated. CH₂Cl₂ (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH₂Cl₂ and EtOAc extracts were combined and dried over MgSO₄, filtered and concentrated. Purification using radial chromatography, eluting with 80% EtOAc/Hexanes afforded N-[(2R,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}acetamide as a white solid

(Compound No. 102, 87 mg, 91% yield). ^1H NMR (CDCl_3): δ 6.80 (s, 2H), 6.55 (br s, 1H), 6.45 (s, 1H), 3.25 (m, 2H), 3.15 (d, 1H), 2.35 (m, 1H), 1.90 (s, 3H), 1.90 (1H), 1.6-1.0 (7H), 0.95 (s, 3H), 0.78 (s, 3H), 0.76 (s, 3H), 0.70 (d, 3H). MS m/z 400.1 ($[\text{M}-1]^-$).

EXAMPLE 61


Synthesis of N- $\{[(2\text{S},3\text{R},4\text{S},4\text{aS},8\text{aS})-4-[(3,5\text{-dihydroxyphenyl})\text{carbonyl}]-3,4\text{a},8,8\text{-tetramethyl-decahydronaphthalen-2-yl}]methyl\}$ acetamide (Compound No. 103)



BBr_3 (0.8 mL of a 1 M solution in CH_2Cl_2) was added to an ice cooled solution of N- $\{[(2\text{S},3\text{R},4\text{S},4\text{aS},8\text{aS})-4-[(3,5\text{-dimethoxyphenyl})\text{carbonyl}]-3,4\text{a},8,8\text{-tetramethyl-decahydronaphthalen-2-yl}]methyl\}$ acetamide (Compound No. 101, 34 mg, 0.079 mmol) in CH_2Cl_2 (10 mL) under argon. After 5 min, the ice bath was removed then after 10 min the reaction vessel was fitted with a reflux condenser and the solution was heated in a 50°C oil bath for 6.5 h. The reaction was then allowed to continue at room temperature for 17 h. MeOH (10 mL) was added and the resulting mixture was concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL) then was heated at 85°C for 45 min. The mixture was cooled to below room temperature then was concentrated. CH_2Cl_2 (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH_2Cl_2 and EtOAc extracts were combined and dried over MgSO_4 , filtered and concentrated. Purification using radial chromatography, eluting with 80% EtOAc/Hexanes then 1% MeOH/EtOAc afforded N- $\{[(2\text{S},3\text{R},4\text{S},4\text{aS},8\text{aS})-4-[(3,5\text{-dihydroxyphenyl})\text{carbonyl}]-3,4\text{a},8,8\text{-tetramethyl-decahydronaphthalen-2-yl}]methyl\}$ acetamide as a white solid (Compound No. 103, 32 mg, 100% yield). ^1H NMR (CDCl_3): δ 7.10 (1H), 6.70 (s, 2H), 6.40 (s, 1H), 3.35 (m, 1H), 2.95 (m, 1H), 2.85 (d, 1H), 1.85 (s, 3H), 1.75 (m, 1H), 1.50 (m, 1H), 1.4-0.9 (7H), 0.90 (s, 3H), 0.70 (s, 3H), 0.65 (6H). MS m/z 402.1 ($[\text{M}+1]^+$).

EXAMPLE 62

Synthesis of N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}pyridine-3-carboxamide
(Compound No. 109)



A. NaBH_4 (117 mg, 3.02 mmol) was added to an ice cold solution of (3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalene-2-carbaldehyde (Compound No. 92, 1.17 g, 3.02 mmol) in MeOH (15 mL) under argon. After 5 min, the ice bath was removed and the reaction was allowed to continue at room temperature for 30 min. The reaction mixture was cooled in ice then saturated NaHCO_3 solution (1.5 mL) was added. The solution was allowed to stir at room temperature for 20 min then was concentrated to remove most of the MeOH. The residue was diluted with EtOAc (150 mL) and was washed with half-saturated NaCl solution (3 x 30 mL). The EtOAc solution was dried over MgSO_4 , filtered and concentrated. Purification using flash chromatography, eluting with 25% EtOAc/Hexanes, afforded [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanol as a white foam (Compound No. 104, 1.13 g, 96% yield).

D. Methanesulfonyl chloride (0.31 mL, 3.9 mmol) was added to an ice cold solution of [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanol (Compound No. 104, 505 mg, 1.3 mmol) in pyridine (6.5 mL) under argon. After 1.5 h, saturated NaHCO₃ solution (5 mL) was added and the resulting solution was stirred at room temperature for 40 min. The mixture was diluted with EtOAc (150 mL) and was washed with half-saturated NaCl solution (3 x 30 mL). The EtOAc solution was dried over MgSO₄, filtered and concentrated. Residual pyridine was co-distilled with toluene (3 x 40 mL) to afford [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl methanesulfonate as a white foam (Compound No. 105, 601 mg, 99% yield).

E. A solution of [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl methanesulfonate (Compound No. 105, 290 mg, 0.62 mmol), NaN₃ (121 mg, 1.86 mmol) and DMF (5 mL) under argon was heated at 60°C for 18 h. The reaction mixture was cooled to room temperature then was diluted with Et₂O (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The Et₂O solution was dried over MgSO₄, filtered and concentrated to afford [(1S,2R,3S,4aS,8aS)-3-(azidomethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone as a yellow oil (Compound No. 106, 260 mg, quantitative yield).

F. A solution of [(1S,2R,3S,4aS,8aS)-3-(azidomethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 106, 257mg, 0.62 mmol), triphenylphosphine (815 mg, 3.1 mmol), water (0.05 mL 2.7 mmol) and THF (5 mL) was stirred at room temperature for 19 h then was heated at reflux for 6 h. The reaction was cooled to room temperature then was concentrated. Purification using radial chromatography, eluting with 5% MeOH/CH₂Cl₂ then 5% MeOH/CH₂Cl₂ with 2% Et₃N afforded [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanamine as a foam (Compound No. 107, 221 mg, 92%).


G. A solution of [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanamine (Compound No. 107, 50 mg, 0.13 mmol), nicotinic acid (21 mg, 0.17 mmol) and dicyclohexylcarbodiimide (DCC) (31 mg, 0.15 mmol) in CH₂Cl₂ (5 mL) was stirred at room temperature under argon. After 22 h, the reaction mixture was filtered through a plug of silica gel, eluting with CH₂Cl₂ then EtOAc. The filtrate was concentrated and purified using radial

chromatography, eluting with 40%, 60% then 80% EtOAc/Hexanes to afford N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}pyridine-3-carboxamide as a white solid (Compound No. 108, 50 mg, 78% yield).

H. BBr_3 (1.0 mL of a 1 M solution in CH_2Cl_2) was added to a room temperature solution of N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}pyridine-3-carboxamide (Compound No. 108, 50 mg, 0.10 mmol) in CH_2Cl_2 (10 mL) under argon. The reaction mixture was heated at reflux for 2 h then allowed to react at room temperature for 3 days. Added MeOH (10 mL) then concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL) then was heated at 85°C for 45 min. The mixture was cooled to below room temperature then was concentrated. CH_2Cl_2 (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH_2Cl_2 and EtOAc extracts were combined and dried over MgSO_4 , filtered and concentrated. Purification using radial chromatography, eluting with 2% MeOH/EtOAc afforded N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}pyridine-3-carboxamide as a white solid (Compound No. 109, 33 mg, 70% yield). ^1H NMR (CDCl_3): δ 8.90 (s, 1H), 8.60 (m, 1H), 8.15 (m, 1H), 7.70 (m, 1H), 7.40 (m, 1H), 6.90 (s, 2H), 6.45 (s, 1H), 3.70 (m, 1H), 3.25 (m, 1H), 2.97 (d, 1H), 1.90 (m, 1H), 1.70 (m, 1H), 1.4-1.0 (9H), 0.95 (s, 3H), 0.75 (9H). MS m/z 465.3 ($[\text{M}+1]^+$).

EXAMPLE 63

Synthesis of N-[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}benzamide (Compound No. 111)




A. A solution of [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanamine (Compound No. 107, 56

mg, 0.14 mmol), benzoic acid (23 mg, 0.19 mmol) and DCC (40 mg, 0.19 mmol) in CH₂Cl₂ (5 mL) was stirred at room temperature under argon. After 19 h, the reaction mixture was filtered through a plug of silica gel, eluting with CH₂Cl₂ then EtOAc. The filtrate was concentrated and purified using radial chromatography, eluting with 20% EtOAc/Hexanes to afford N-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}benzamide as a white solid (Compound No. 110, 49 mg, 69% yield).

B. BBr₃ (0.5 mL of a 1 M solution in CH₂Cl₂) was added to an ice cooled solution of N-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}benzamide (Compound No. 110, 49 mg, 0.10 mmol) in CH₂Cl₂ (10 mL) under argon. The reaction mixture was allowed to react at room temperature for 6 days. Added MeOH (10 mL) then concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL) then was heated at 85°C for 45 min. The mixture was cooled to below room temperature then was concentrated. CH₂Cl₂ (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH₂Cl₂ and EtOAc extracts were combined and dried over MgSO₄, filtered and concentrated. Purification using radial chromatography, eluting with 40% EtOAc/Hexanes afforded N-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}benzamide as a white solid (Compound No. 111, 26 mg, 56% yield). ¹H NMR (CDCl₃): δ 8.90 (s, 1H), 7.70 (m, 2H), 7.40 (m, 3H), 7.10 (1H), 6.78 (s, 2H), 6.42 (s, 1H), 3.65 (m, 1H), 3.20 (m, 1H), 2.95 (d, 1H), 1.85 (m, 1H), 1.65 (m, 1H), 1.4-1.0 (7H), 0.90 (s, 3H), 0.70 (9H). MS *m/z* 478.2 ([M+1]⁺).

EXAMPLE 64

Synthesis of 5-[[[(1S,2R,3S,4aS,8aS)-3-(1H-imidazol-1-ylmethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 113)




A. NaH (86 mg of a 60% dispersion in mineral oil, 2.1 mmol) was added to an ice cold solution of imidazole (146 mg, 2.1 mmol) in DMF (5 mL) under argon. After 1 h, [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl methanesulfonate (Compound No. 105, 100 mg, 0.21 mmol) was added and the reaction was allowed to warm to room temperature. After 17 h, the reaction mixture was cooled in ice and saturated NaHCO₃ solution (5 mL) was added. The solution was diluted with Et₂O (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The Et₂O solution was dried over MgSO₄, filtered and concentrated. Purification using radial chromatography, eluting with 5% MeOH/CH₂Cl₂, afforded 1-[[[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl]-1H-imidazole as a white solid (Compound No. 112, 79 mg, 84% yield).

B. BBr₃ (0.74 mL of a 1 M solution in CH₂Cl₂) was added to a -78°C solution of 1-[[[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl]-1H-imidazole (Compound No. 112, 65 mg, 0.15 mmol) in CH₂Cl₂ (5 mL) under argon. The reaction mixture was allowed to warm to room temperature while allowed to react for 21 h. Added MeOH (10 mL) then concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL) then was heated at 85°C for 45 min. The mixture was cooled to room temperature then was concentrated. CH₂Cl₂ (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH₂Cl₂ and EtOAc extracts were combined and dried over MgSO₄, filtered and concentrated to afford 5-[[[(1S,2R,3S,4aS,8aS)-3-(1H-imidazol-1-ylmethyl)-2,5,5,8a-tetramethyl-

decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol as a tan solid (Compound No. 113, 40 mg, 65% yield). ^1H NMR (CDCl_3): δ 7.42 (s, 1H), 6.95 (s, 1H), 6.85 (s, 1H), 6.78 (s, 2H), 6.42 (s, 1H), 4.15 (m, 1H), 3.72 (m, 1H), 2.97 (d, 1H), 1.95 (m, 1H), 1.50 (m, 1H), 1.4-1.1 (4H), 1.0 (6H), 0.8 (7H), 0.65 (6H). MS m/z 409.3 ($[\text{M}-1]^+$).

EXAMPLE 65

Synthesis of 5-{[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-(1H-1,2,4-triazol-1-ylmethyl)-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 115)




A. NaH (86 mg of a 60% dispersion in mineral oil, 2.1 mmol) was added to an ice cold solution of 1,2,4-triazole (148 mg, 2.1 mmol) in DMF (5 mL) under argon. After 1 h, [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl methanesulfonate (Compound No. 105, 100 mg, 0.21 mmol) was added and the reaction was allowed to warm to room temperature. After 3 days, the reaction mixture was cooled in ice and saturated NaHCO_3 solution (3 mL) was added. The solution was diluted with Et_2O (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The Et_2O solution was dried over MgSO_4 , filtered and concentrated. Purification using radial chromatography, eluting with 40% then 60% EtOAc /Hexanes, afforded 1-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}-1H-1,2,4-triazole as a white solid (Compound No. 114, 84 mg, 89% yield).

B. BBr_3 (0.93 mL of a 1 M solution in CH_2Cl_2) was added to a -78°C solution of 1-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}-1H-1,2,4-triazole (Compound No. 114, 82 mg, 0.19 mmol) in CH_2Cl_2 (5 mL) under argon. The reaction mixture was allowed to warm to room temperature while allowed to react for 18 h. Added MeOH (10 mL) then concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL) then was heated at 85°C for 45 min. The mixture was cooled to room temperature then was concentrated.

CH₂Cl₂ (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH₂Cl₂ and EtOAc extracts were combined and dried over MgSO₄, filtered and concentrated. ¹H NMR indicated the reaction was not complete therefore the reaction was set up again with the modification that 10 mL CH₂Cl₂ was used as reaction solvent. After 2 days, the work up procedure was repeated. Purification using radial chromatography, eluting with 50% then 80% EtOAc/Hexanes afforded 5-[[[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-(1H-1,2,4-triazol-1-ylmethyl)-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol as a white solid ((Compound No. 115, 51 mg, 66% yield). ¹H NMR (CDCl₃): δ 8.10 (s, 1H), 7.90 (2, 1H), 6.78 (s, 2H), 6.45 (s, 1H), 4.35 (m, 1H), 3.98 (m, 1H), 2.97 (d, 1H), 1.90 (m, 1H), 1.70 (m, 1H), 1.4-1.1 (4H), 1.0 (6H), 0.80 (7H), 0.65 (6H). MS *m/z* 412.3 ([M+1]⁺).

EXAMPLE 66

Synthesis of 5-[[[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(4-methylpiperazin-1-yl)methyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol dihydrochloride salt (Compound No. 117)




A. NaH (86 mg of a 60% dispersion in mineral oil, 2.1 mmol) was added to an ice cold solution of 1-methylpiperazine ("NMP") (0.24 mL, 2.1 mmol) in DMF (5 mL) under argon. After 1 h, [(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl methanesulfonate (Compound No. 105, 100 mg, 0.21 mmol) was added and the reaction was allowed to warm to room temperature. After 1 day, the reaction was heated at 50°C for 23 h then the mixture was cooled in ice and saturated NaHCO₃ solution (3 mL) was added. The solution was diluted with Et₂O (75 mL) and was washed with half-saturated NaCl solution (3 x 15 mL). The Et₂O solution was dried over MgSO₄, filtered and concentrated. Purification using radial chromatography, eluting with 2% then 5%, then

10% MeOH/CH₂Cl₂, afforded 1-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}-4-methylpiperazine as a colourless film (Compound No. 116, 54 mg, 53% yield).

B. BBr₃ (0.6 mL of a 1 M solution in CH₂Cl₂) was added to a -78°C solution of 1-{[(2S,3R,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methyl}-4-methylpiperazine (Compound No. 116, 54 mg, 0.12 mmol) in CH₂Cl₂ (5 mL) under argon. The reaction mixture was allowed to warm to room temperature while allowed to react for 23 h. Added MeOH (10 mL) then concentrated. The MeOH addition/concentration cycle was repeated twice more. The residue was taken up into *i*-PrOH (4.5 mL) and 1 N HCl (1.5 mL) then was heated at 85°C for 45 min. The mixture was cooled to room temperature then was concentrated. CH₂Cl₂ (30 mL) and water (10 mL) were added then the pH was adjusted to approximately 9 by addition of 2 N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc (25 mL). The CH₂Cl₂ and EtOAc extracts were combined and dried over MgSO₄, filtered and concentrated. ¹H NMR indicated the reaction was not complete therefore the reaction was set up again with the modifications that the reaction was begun at 0°C and that 10 mL CH₂Cl₂ was used as reaction solvent. After 3 days, the work up procedure was repeated. Purification using radial chromatography, eluting with 5% MeOH/CH₂Cl₂ with 2% Et₃N afforded 26 mg of a white solid. The material was taken up into MeCN (6 mL) then HCl (0.15 mL of a 1.25 M solution in *i*-PrOH) was added resulting in formation of a white precipitate. Most of the liquid was decanted then MeCN (5 mL) was added to the solid. The suspension was concentrated and dried to afford 5-{[(1S,2R,3S,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(4-methylpiperazin-1-yl)methyl]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol dihydrochloride salt as a white solid (Compound No. 117, 26 mg, 44% yield). ¹H NMR (CD₃OD): δ 6.85 (2H), 6.50 (1H), 3.8 (br s, 10H), 3.45 (m, 1H), 3.2 (m, 1H), 3.15 (d, 1H), 3.0 (s, 3H), 2.15 (m, 1H), 1.90 (m, 1H), 1.78 (m, 1H), 1.6-1.1 (6H), 1.00 (s, 3H), 0.97 (s, 3H), 0.90 (s, 3H), 0.85 (d, 3H). MS *m/z* 443.4 ([M+1]⁺).

EXAMPLE 67


Synthesis of 3-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea (Compound No. 118)



To a suspension of 5-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 61 mg, 0.18 mmol) and pyridin-3-yl-carbamic acid isopropenyl ester (39 mg, 0.22 mmol, prepared according to Gallou, I. *et al. J. Org. Chem.*, 2005, 70 (17), pp 6960–6963) in THF (1.00 mL) was added 1-methylpyrrolidine (0.018 mL, 0.17 mmol) under argon then heated to 55°C for 24 h. The suspension was allowed to cool to room temperature and stirred for another 17 h. The mixture was concentrated, and the residue was purified by chromatography on silica gel (MeOH/EtOAc, 1:9) to give 3-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea as a colourless powder (Compound No. 118, 59 mg, 72%). ¹H NMR (CD₃OD): δ 8.56 (s, 1H), 8.14 (m, 1H), 7.97 (m, 1H), 7.34 (dd, *J* = 8.3, 4.7 Hz, 1H), 6.88 (d, *J* = 2.4 Hz, 1H), 6.51 (t, *J* = 2.3 Hz, 1H), 4.13 (m, 1H), 3.23 (d, *J* = 12 Hz, 1H), 2.42 (m, 1H), 1.11–1.85 (m, 10H), 1.08 (s, 3H), 0.88 (s, 6H), 0.77 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 204.6, 158.7 (2C), 156.3, 143.4, 142.1, 139.3, 126.3, 124.1, 106.8, 106.5 (2C), 58.2, 50.3, 41.9, 41.3, 39.2, 33.3, 32.8, 32.7, 27.8, 20.9, 18.4, 15.5, 12.7. ES-MS *m/z* 466 ([*M*+1]⁺).

EXAMPLE 68


Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-5-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 119)



A. To a suspension of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 81 mg, 0.23 mmol) and 4-imidazolecarboxaldehyde (34 mg, 0.35 mmol) in CH_2Cl_2 (2.3 mL) was added acetic acid (0.013 mL, 0.23 mmol) and sodium triacetoxyborohydride (158 mg, 0.745 mmol) under argon. The solution was stirred for 72 h, diluted with 10% MeOH/ CH_2Cl_2 (20 mL), and washed with saturated aqueous NaHCO_3 (20 mL). The aqueous layer was extracted with 10% MeOH/ CH_2Cl_2 (3 x 10 mL), and the combined organic layers were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 100:10:1) to give 5-[[[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-5-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol as a colourless powder (Compound No. 119, 24 mg, 24%). ^1H NMR (CD_3OD): δ 7.69 (s, 1H), 7.07 (s, 1H), 6.87 (d, J = 2.1 Hz, 2H), 6.46 (t, J = 2.3 Hz, 1H), 3.93 (d, J = 14 Hz, 1H), 3.74 (d, J = 14 Hz, 1H), 3.53 (d, J = 12 Hz, 1H), 3.00 (br s, 1H), 2.33 (m, 1H), 1.95 (m, 1H), 1.19-1.60 (m, 7H), 1.06 (m, 1H), 1.04 (s, 3H), 0.96 (s, 3H), 0.89 (s, 3H), 0.77 (d, J = 6.6 Hz, 3H). ^{13}C NMR (CD_3OD): δ 206.5, 159.9 (2C), 144.5, 136.7 (2C), 118.2, 108.1, 107.8 (2C), 58.4, 58.0, 47.3, 44.2, 42.9, 42.1, 40.5, 35.0, 34.0 (2C), 25.4, 22.2, 19.6, 16.5, 14.1. ES-MS m/z 426 ($[\text{M}+1]^+$).

EXAMPLE 69


Synthesis of 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-ethylurea (Compound No. 120)



To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 50 mg, 0.14 mmol) in THF (1.46 mL) was added ethyl isocyanate (0.023 mL, 0.29 mmol) under argon. The solution was stirred at room temperature for 6 h then another equivalent of ethyl isocyanate (0.010 mL, 0.13 mmol) was added. The solution was stirred for another 60 h then concentrated. The residue was purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:2) to give 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-ethylurea as a colourless solid (Compound No. 120, 45 mg, 75%). ¹H NMR (CD₃OD): δ 6.87 (d, *J* = 1.8 Hz, 2H), 6.49 (s, 1H), 6.29 (d, *J* = 9.0 Hz, 1H), 4.02 (m, 1H), 3.18 (m, 3H), 2.34 (m, 1H), 1.03-1.75 (m, 15H), 0.85 (s, 6H), 0.70 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.1, 161.0, 159.8 (2C), 144.6, 108.0, 107.7 (2C), 59.2, 51.2, 49.2, 43.0, 42.4, 40.3, 35.7, 34.6, 33.9 (2C), 29.2, 22.1, 19.5, 16.7, 15.9, 14.0. ES-MS *m/z* 417 ([M+1]⁺).

EXAMPLE 70


Synthesis of 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methoxyurea (Compound No. 121)



To a suspension of methoxylamine hydrochloride (291 mg, 3.48 mmol) in THF (5.0 mL) was added 1,1'-carbonyldiimidazole (564 mg, 3.48 mmol) and *N,N*-diisopropylethylamine (1.10 mL, 6.31 mmol) under argon. The solution was stirred at room temperature for 1 h then 5-[[[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 109 mg, 0.316 mmol) in THF (10.8 mL) was added, and the solution was heated to 60°C overnight. After 20 h, the solution was allowed to cool to room temperature and saturated aqueous NaHCO₃ (20 mL) was added. The aqueous layer was extracted with EtOAc (3 x 20 mL), and the combined organic layers were dried (MgSO₄) and concentrated. The residue was partially purified by chromatography on silica gel (MeOH/EtOAc, 1:49) to give a colourless solid (112 mg). To a solution of the colourless solid (112 mg) in MeOH (5.0 mL) was added 10N NaOH (0.50 mL, 5.0 mmol) at room temperature. The solution was stirred for 1 h then concentrated, and the residue was partitioned between EtOAc (40 mL) and H₂O (20 mL). The aqueous layer was adjusted to pH 9 and extracted with EtOAc (2 x 30 mL). The combined organic layers were dried (MgSO₄) and concentrated. The residue was suspended in CH₂Cl₂, heated to a boil, and allowed to cool to room temperature. 1-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methoxyurea, a colourless solid (Compound No. 121, 19 mg, 14%), was collected by filtration. ¹H NMR (CD₃OD): δ 6.86 (d, *J* = 1.8 Hz, 2H), 6.53 (d, *J* = 9.9 Hz, 1H), 6.49 (s, 1H), 4.15 (m, 1H), 3.74 (s, 3H), 3.25 (d, *J* = 12 Hz, 1H), 2.41 (m, 1H), 1.75 (m, 2H), 1.06-1.57 (m, 10H), 0.87 (s, 6H), 0.72 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 205.7, 162.1, 159.9 (2C), 144.4, 108.1, 107.7 (2C), 64.7, 59.3, 51.2, 49.5, 42.9, 42.3, 40.3, 34.4, 34.0, 33.9, 29.0, 22.1, 19.5, 16.7, 14.0. ES-MS *m/z* 419 ([*M*+1]⁺).

EXAMPLE 71


Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]benzamide (Compound No. 122)



To a solution of 5-[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 149 mg, 0.431 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (170 mg, 0.448 mmol), and benzoic acid (63 mg, 0.52 mmol) in DMF (4.32 mL) was added *N,N*-diisopropylethylamine (0.11 mL, 0.63 mmol) under argon. The solution was stirred at room temperature for 66 h then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (5 x 10 mL) and brine (2 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:24) to give *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]benzamide as a colourless powder (Compound No. 122, 130 mg, 67%). ¹H NMR (CD₃OD): δ 8.41 (d, *J* = 9.3 Hz, 1H), 7.75 (d, *J* = 8.1 Hz, 2H), 7.52 (m, 3H), 6.95 (d, *J* = 1.8 Hz, 2H), 6.48 (s, 1H), 4.51 (m, 1H), 3.60 (d, *J* = 12 Hz, 1H), 2.49 (m, 1H), 1.79 (m, 2H), 1.28-1.56 (m, 6H), 1.08 (m, 4H), 0.88 (s, 6H), 0.78 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.3, 171.5, 159.6 (2C), 144.5, 136.9, 132.2, 129.3 (2C), 128.5 (2C), 108.0, 107.9 (2C), 58.8, 51.6, 48.6, 42.8, 42.3, 40.1, 34.0, 33.9, 33.8, 28.8, 22.1, 19.5, 16.7, 14.2. ES-MS *m/z* 450 ([*M*+1]⁺).

EXAMPLE 72


Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-3-carboxamide (Compound No. 123)



To a solution of 5-[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 147 mg, 0.425 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (164 mg, 0.432 mmol), and nicotinic acid (64 mg, 0.52 mmol) in DMF (4.27 mL) was added *N,N*-diisopropylethylamine (0.10 mL, 0.57 mmol) under argon. The solution was stirred at room temperature for 66 h then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (5 x 10 mL) and brine (2 x 10 mL) then dried (MgSO₄) and concentrated. The residue was suspended in CH₂Cl₂, heated to a boil, and allowed to cool to room temperature. A pink solid was collected by filtration then purified by chromatography on silica gel (EtOAc) to give *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-3-carboxamide as a colourless solid (Compound No. 123, 72 mg, 38%). ¹H NMR (CD₃OD): δ 8.88 (s, 1H), 8.70 (d, *J* = 3.6 Hz, 1H), 8.16 (d, *J* = 8.7 Hz, 1H), 7.57 (m, 1H), 6.94 (d, *J* = 2.1 Hz, 2H), 6.48 (s, 1H), 4.54 (m, 1H), 3.57 (d, *J* = 13 Hz, 1H), 2.50 (m, 1H), 1.81 (m, 2H), 1.24-1.56 (m, 6H), 1.09 (m, 4H), 0.89 (s, 6H), 0.78 (d, *J* = 6.3 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.2, 169.0, 164.5, 159.8 (2C), 152.3, 149.1, 144.6, 137.3, 133.4, 125.0, 107.9 (2C), 58.9, 52.0, 48.8, 42.9, 42.5, 40.2, 34.0, 33.9 (2C), 28.8, 22.1, 19.5, 16.7, 14.2. ES-MS *m/z* 451 ([*M*+1]⁺).

EXAMPLE 73


Synthesis of 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylurea (Compound No. 124)



To a suspension of methylamine hydrochloride (130 mg, 1.93 mmol) and 1,1'-carbonyldiimidazole (306 mg, 1.89 mmol) in THF (5.0 mL) was added *N,N*-diisopropylethylamine (0.61 mL, 3.5 mmol) under argon. The solution was stirred at room temperature for 1 h then 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 62 mg, 0.18 mmol) in THF (3.68 mL) was added, and the solution was heated to 60°C overnight. After 17 h, the solution was allowed to cool to room temperature and saturated aqueous NaHCO₃ (20 mL) was added. The aqueous layer was extracted with EtOAc (3 x 20 mL), and the combined organic layers were dried (MgSO₄) and concentrated to give a yellow oil (229 mg). To a solution of the yellow oil (229 mg) in MeOH (5.69 mL) was added 10N NaOH (0.57 mL, 5.7 mmol) at room temperature. The solution was stirred for 2 h then concentrated, and the residue was partitioned between EtOAc (20 mL) and H₂O (30 mL). The aqueous layer was adjusted to pH 8 and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc) to give 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylurea as a colourless solid (Compound No. 124, 38 mg, 53%). ¹H NMR (CD₃OD): δ 6.87 (d, *J* = 2.1 Hz, 2H), 6.50 (t, *J* = 2.1 Hz, 1H), 6.34 (d, *J* = 8.4 Hz, 1H), 4.03 (m, 1H), 3.21 (d, *J* = 12 Hz, 1H), 2.73 (s, 3H), 2.34 (m, 1H), 1.17-1.74 (m, 8H), 1.08 (m, 4H), 0.86 (s, 6H), 0.70 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.1, 161.8, 159.8 (2C), 144.6, 108.0, 107.7 (2C), 59.2, 51.3, 49.2, 43.1, 42.4, 40.3, 34.6, 33.9 (2C), 29.3, 26.9, 22.1, 19.5, 16.7, 14.0. ES-MS *m/z* 403 ([M+1]⁺).

EXAMPLE 74


Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(1H-pyrrol-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol
(Compound No. 125)



To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 70 mg, 0.20 mmol) in THF (2.05 mL) was added pyrrole-2-carboxaldehyde (25 mg, 0.26 mmol), acetic acid (0.016 mL, 0.28 mmol) and sodium triacetoxyborohydride (141 mg, 0.665 mmol) under argon. The mixture was stirred for 17 h, diluted with EtOAc (30 mL), and washed with saturated aqueous NaHCO₃ (20 mL). The aqueous layer was extracted with EtOAc (3 x 20 mL), and the combined organic layers were dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/EtOAc, 1:49) to give 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(1H-pyrrol-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol as a colourless powder (Compound No. 125, 29 mg, 34%). ¹H NMR (CD₃OD): δ 6.87 (d, *J* = 2.1 Hz, 2H), 6.71 (m, 1H), 6.46 (m, 1H), 6.04 (m, 2H), 3.93 (d, *J* = 14 Hz, 1H), 3.73 (d, *J* = 14 Hz, 1H), 3.51 (d, *J* = 12 Hz, 1H), 2.95 (br s, 1H), 2.31 (m, 1H), 1.93 (m, 1H), 1.19-1.54 (m, 7H), 1.06 (m, 4H), 0.97 (s, 3H), 0.88 (s, 3H), 0.74 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.8, 159.9 (2C), 144.5, 130.2, 118.5, 108.6, 108.1, 108.0, 107.8 (2C), 58.1, 57.9, 47.3, 44.8, 43.0, 42.2, 40.5, 35.1, 34.1, 33.9, 25.6, 22.2, 19.6, 16.5, 14.1. ES-MS *m/z* 423 ([M-1]).

EXAMPLE 75


Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol
(Compound No. 126)



To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 72 mg, 0.21 mmol) in THF (2.09 mL) was added 2-pyridinecarboxaldehyde (0.024 mL, 0.25 mmol), acetic acid (0.012 mL, 0.21 mmol) and sodium triacetoxyborohydride (144 mg, 0.679 mmol) under argon. The mixture was stirred for 16 h, diluted with saturated aqueous NaHCO₃ (20 mL), and the aqueous layer was extracted with EtOAc (3 x 30 mL). The combined organic layers were dried (MgSO₄) and concentrated. The residue was partially purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:2) to give a yellow solid (63 mg). The yellow solid (63 mg) was suspended in CH₂Cl₂, heated to a boil, and allowed to cool to room temperature. 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-Tetramethyl-3-[(pyridin-2-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol, a colourless solid (Compound No. 126, 49 mg, 54%), was collected by filtration. ¹H NMR (CD₃OD): δ 8.52 (d, *J* = 4.2 Hz, 1H), 7.83 (m, 1H), 7.51 (d, *J* = 7.8 Hz, 1H), 7.32 (m, 1H), 6.87 (s, 2H), 6.46 (s, 1H), 4.00 (d, *J* = 14 Hz, 1H), 3.81 (d, *J* = 14 Hz, 1H), 3.58 (d, *J* = 11 Hz, 1H), 2.87 (br s, 1H), 2.31 (m, 1H), 1.93 (m, 1H), 1.18-1.67 (m, 7H), 1.06 (m, 4H), 0.93 (s, 3H), 0.88 (s, 3H), 0.81 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.7, 160.6, 159.4 (2C), 149.6, 144.2, 138.2, 124.0, 123.3, 107.9, 107.6 (2C), 58.6, 57.8, 53.4, 47.0, 42.7, 41.9, 40.3, 35.2, 34.0, 33.7, 25.9, 22.2, 19.4, 16.6, 14.0. ES-MS *m/z* 437 ([M+1]⁺).

EXAMPLE 76


Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-4-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol
(Compound No. 127)



To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 70 mg, 0.20 mmol) in THF (2.03 mL) was added 4-pyridinecarboxaldehyde (0.023 mL, 0.24 mmol), acetic acid (0.012 mL, 0.21 mmol) and sodium triacetoxyborohydride (139 mg, 0.656 mmol) under argon. The mixture was stirred for 14 h, diluted with saturated aqueous NaHCO₃ (30 mL), and the aqueous layer was extracted with EtOAc (3 x 30 mL). The combined organic layers were dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/EtOAc, 1:49) to give 5-[[[(1S,2S,3R,4aS,8aS)-2,5,5,8a-tetramethyl-3-[(pyridin-4-ylmethyl)amino]-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol as a colourless solid (Compound No. 127, 40 mg, 45%). ¹H NMR (CD₃OD): δ 8.48 (d, *J* = 5.4 Hz, 2H), 7.50 (d, *J* = 5.4 Hz, 2H), 6.87 (d, *J* = 1.8, 2H), 6.47 (m, 1H), 3.96 (d, *J* = 14 Hz, 1H), 3.77 (d, *J* = 15 Hz, 1H), 3.56 (d, *J* = 12 Hz, 1H), 2.86 (m, 1H), 2.30 (m, 1H), 1.88 (m, 1H), 1.18-1.66 (m, 7H), 1.06 (m, 4H), 0.89 (s, 3H), 0.87 (s, 3H), 0.80 (d, *J* = 6.9 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.8, 159.6 (2C), 152.9, 149.8 (2C), 144.4, 124.8 (2C), 107.9, 107.6 (2C), 58.3, 58.0, 51.2, 47.4, 42.9, 42.1, 40.4, 35.4, 34.0, 33.8, 26.0, 22.2, 19.5, 16.7, 14.0. ES-MS *m/z* 437 ([M+1]⁺).

EXAMPLE 77

Synthesis of (1R,2R,3R,4aS,8aS)-3-amino-1-[(3,5-dimethoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 132)



A. A solution of the mixture of [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methyl acetate (8 parts) and [(1S,4aS,8aS)-5,5,8a-trimethyl-2-methyldene-decahydronaphthalen-1-yl]methyl acetate (1 part) (Compound No. 128, prepared according to Kuchkova *et al. Chem. Nat. Comp.* 2007, 43(4), pp 412-416, 2.02 g, 7.64 mmol) and *m*-chloroperoxybenzoic acid (3.33 g, 14.9 mmol) in CH₂Cl₂ (52 mL) under argon was stirred at room temperature for 1 h. Water (30 mL) and as solution of 10% aqueous Na₂SO₃ (30 mL) were added and the mixture stirred at room temperature for 15 min. The layers were partitioned, and the organic layer washed with saturated aqueous NaHCO₃ (3 x 30 mL) and brine (30 mL) then dried (MgSO₄) and concentrated to afford a colourless oil (2.01 g) that was used in the next step without further purification.

B. A solution of the colourless oil from above (2.01 g) in 6:1 MeOH/10 N NaOH was stirred at room temperature for 2 h. Water (50 mL) was added, and the mixture was extracted with EtOAc (4 x 50 mL). The organic layers were combined, washed with brine (50 mL) then dried (MgSO₄) and concentrated. The residue was purified by silica gel chromatography (25% - 35 % EtOAc/hexanes) to afford a 5:1 mixture of [(1aR,2S,2aS,6aS,7aS)-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxiren-2-yl]methanol and [(1aS,2S,2aS,6aS,7aR)-1a,2a,6,6-tetramethyl-

decahydronaphtho[2,3-b]oxiren-2-yl]methanol as a colourless solid (Compounds No. 129a and 129b, 1.71 g, 81% over 2 steps).

C. To a solution of the 5:1 mixture of [(1aR,2S,2aS,6aS,7aS)-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxiren-2-yl]methanol and [(1aS,2S,2aS,6aS,7aR)-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxiren-2-yl]methanol from above (Compounds No. 129a and 129b, 1.53 g, 6.42 mmol) and triethylamine (1.20 mL, 8.61 mmol) in CH₂Cl₂ at 0°C under argon was added methanesulfonyl chloride (0.55 mL, 7.1 mmol). The solution was stirred at 0°C for 5 min then at room temperature for 1.5 h. The solution was diluted with ethyl acetate (180 mL) and washed with saturated aqueous NaHCO₃ (5 x 35 mL) and brine (35 mL) then dried (MgSO₄) and concentrated, affording a yellow solid (1.97 g), that was used in the next step without further purification.

D. A mixture of the yellow solid from above (1.97 g), Cs₂CO₃ (6.34 g, 19.4 mmol), and 3,5-dimethoxythiophenol (Compound No. 79, 1.13 g, 6.65 mmol) in DMF (32 mL) under argon was stirred at 80°C for 17 h. The mixture was diluted with EtOAc (300 mL) and saturated aqueous NH₄Cl (100 mL). The aqueous layer was extracted with EtOAc (100 mL), and the organic layers combined, washed with brine (5 x 100 mL), dried (MgSO₄) and concentrated. The residue was purified on silica gel (15% EtOAc/hexanes) to afford (1aR,2R,2aS,6aS,7aS)-2-{[(3,5-dimethoxyphenyl)sulfanyl]methyl}-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxirene as a green oil (Compound No. 130, 2.73 g, 82% over 2 steps).


E. To a solution of (1aR,2R,2aS,6aS,7aS)-2-{[(3,5-dimethoxyphenyl)sulfanyl]methyl}-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxirene (Compound No. 130, 507 mg, 1.30 mmol) in DMF (13.0 mL) under argon was added sodium azide (549 mg, 8.45 mmol) and ammonium chloride (209 mg, 3.90 mmol). The mixture was heated to 80°C for 5 days then allowed to cool to room temperature and diluted with EtOAc (150 mL). The organic layer was washed with H₂O (20 mL) and brine (5 x 40 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 15:185) to give (1R,2R,3R,4aS,8aS)-3-azido-1-{[(3,5-dimethoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol as a yellow oil (Compound No. 131, 299 mg, 53%).

F. To a solution of (1R,2R,3R,4aS,8aS)-3-azido-1-{[(3,5-dimethoxyphenyl)sulfanyl]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 131, 299 mg, 0.690 mmol) in THF (6.88 mL) was added lithium

aluminum hydride (2.0 M in THF, 0.69 mL, 1.4 mmol) dropwise at 0°C under argon. The mixture was stirred for 20 min at 0°C then at room temperature for 3 h. The mixture was cooled to 0°C, and H₂O (0.06 mL) was added followed by 15% aqueous NaOH (0.06 mL) and H₂O (0.16 mL). The mixture was stirred for 30 min at 0 °C then dried (MgSO₄) and stirred at room temperature for 10 min. The mixture was filtered and concentrated, and the residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:19) to give (1R,2R,3R,4aS,8aS)-3-amino-1-[[[(3,5-dimethoxyphenyl)sulfanyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol as a colourless foam (Compound No. 132, 163 mg, 58%). ¹H NMR (CDCl₃): δ 6.46 (s, 2H), 6.24 (s, 1H), 3.78 (s, 6H), 3.55 (br s, 1H), 3.06 (dd, *J* = 12, 4.5 Hz, 1H), 2.83 (m, 1H), 0.97-1.87 (m, 16H), 0.88 (s, 3H), 0.82 (s, 3H). ¹³C NMR (CDCl₃): δ 161.0 (2C), 141.0, 106.1 (2C), 97.9, 77.5, 56.1, 55.5 (2C), 53.6, 46.8, 41.9, 39.4, 39.2, 33.1, 32.9, 29.1, 27.9, 25.5, 21.7, 18.2, 15.5. ES-MS *m/z* 408 ([M+1]⁺).

EXAMPLE 78

Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-2-ylmethyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 133)




To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 75 mg, 0.22 mmol) in EtOH (2.2 mL) under argon was added 2-imidazolecarboxaldehyde (26 mg, 0.27 mmol), and the mixture was heated to reflux for 2 h. After allowing the solution to cool to room temperature, sodium borohydride (26 mg, 0.69 mmol) was added, and the mixture was stirred for 30 min then partitioned between H₂O (10 mL) and EtOAc (20 mL). The aqueous layer was adjusted to pH 10 and extracted with EtOAc (2 x 10 mL). The combined organic layers were dried (MgSO₄) and concentrated. The residue was suspended in CH₂Cl₂, heated to a boil, and allowed to cool to room temperature. 5-[[[(1S,2S,3R,4aS,8aS)-3-[(1H-imidazol-2-ylmethyl)amino]-2,5,5,8a-tetramethyl-

decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol, a colourless solid (Compound No. 133, 49 mg, 53%), was collected by filtration. ^1H NMR (CD_3OD): δ 6.99 (s, 2H), 6.86 (d, $J = 2.1$ Hz, 2H), 6.46 (m, 1H), 3.92 (d, $J = 14$ Hz, 1H), 3.77 (d, $J = 14$ Hz, 1H), 3.52 (d, $J = 12$ Hz, 1H), 2.80 (br s, 1H), 2.28 (m, 1H), 1.90 (m, 1H), 1.17-1.62 (m, 7H), 1.06 (m, 4H), 0.92 (s, 3H), 0.87 (s, 3H), 0.76 (d, $J = 6.9$ Hz, 3H). ^{13}C NMR (CD_3OD): δ 206.9, 159.8 (2C), 148.7, 144.5, 122.5 (2C), 108.1, 107.8 (2C), 58.5, 58.0, 47.3, 45.1, 43.0, 42.2, 40.5, 35.4, 34.1, 33.9, 25.9, 22.3, 19.6, 16.7, 14.1. ES-MS m/z 426 ($[\text{M}+1]^+$).

EXAMPLE 79

Synthesis of (2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 135)




A. Sodium triacetoxyborohydride (6.96 g, 32.8 mmol) was added to a solution of (3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-one (Compound No. 134, prepared from Compound No. 7 in similar manner as described for Compound No. 90, 3.87 g, 10.9 mmol), 4-methoxybenzylamine (1.56 mL, 12.0 mmol), and acetic acid (0.63 mL, 11 mmol) in CH_2Cl_2 (110 mL), under argon and the mixture was stirred at room temperature for 25 h. The mixture was diluted with CH_2Cl_2 (70 mL) then washed with 1 N NaOH (100 mL). The aqueous layer was extracted with CH_2Cl_2 (3 x 75 mL), and the organic layers were combined, dried (MgSO_4) and concentrated. The residue was partially purified by chromatography on silica gel ($\text{EtOAc}/\text{CH}_2\text{Cl}_2$, 1:19-1:9) to give (2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine as a yellow foam (3.72 g).

B. Methanol (80 mL) was carefully added under argon to (2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-N-[(4-

methoxyphenyl)methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (3.72 g) and 10% Pd/C (788 mg) followed by the addition of 10% aqueous HCl (4.0 mL). The mixture was stirred at room temperature under H₂(g) at 1 atm for 16 h then filtered through Celite and concentrated. The residue was purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 200:10:1) to afford (2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine as a colourless solid (Compound No. 135, 1.62 g, 42% over 2 steps). ¹H NMR: (CDCl₃): δ 7.36 (s, 1H), 7.27 (s, 1H), 6.90 (s, 1H), 3.84 (s, 3H), 3.51 (d, *J* = 11.6 Hz, 1H), 3.25 (m, 1H), 2.40 (s, 3H), 2.30 (m, 1H), 1.75 (m, 1H), 1.62-1.09 (m, 8H), 1.07 (s, 3H), 0.89 (s, 3H), 0.84 (s, 3H), 0.78 (d, *J* = 6.8 Hz, 3H).

EXAMPLE 80


Synthesis of 3-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenol (Compound No. 136)



Boron tribromide (1.0 M in CH₂Cl₂, 18.1 mL, 18.1 mmol) was added to a solution of (2R,3S,4S,4aS,8aS)-4-[(3-methoxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 135, 1.62 g, 4.53 mmol) in CH₂Cl₂ (45 mL) at -78°C under argon. The solution was stirred for 5 min at -78°C then at room temperature for 3.25 h. The solution was carefully quenched with MeOH (60 mL) and concentrated. The residue was dissolved in *i*-PrOH (45 mL) and 10% aqueous HCl (15 mL), and the solution was stirred at 85°C for 100 min. The solution was concentrated, diluted with EtOAc (180 mL) and H₂O (60 mL), then neutralized to pH 7 with 1 N NaOH. The aqueous layer was extracted with EtOAc (2 x 60 mL), and the organic layers were combined, dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:0.9) to give 3-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenol as a colourless solid (Compound No. 136, 1.51 g, 97%).

EXAMPLE 82


Synthesis of 3-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea (Compound No. 138)



To a suspension of 3-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenol (Compound No. 136, 61 mg, 0.18 mmol) and pyridin-3-yl-carbamic acid isopropenyl ester (39 mg, 0.22 mmol) in THF (0.35 mL) was added 1-methylpyrrolidine (0.020 mL, 0.19 mmol) under argon then heated to 55°C for 16 h. The mixture was concentrated, and the residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:9-15:85) to give 3-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-1-(pyridin-3-yl)urea as a light yellow solid (Compound No. 138, 70 mg, 85%). ¹H NMR (CD₃OD): δ 8.55 (br s, 1H), 8.14 (d, *J* = 4.5 Hz, 1H), 7.98 (m, 1H), 7.35 (dd, *J* = 8.4, 4.9 Hz, 1H), 7.28 (s, 1H), 7.17 (m, 1H), 6.87 (m, 1H), 6.67 (d, *J* = 9.3 Hz, 1H), 4.14 (m, 1H), 3.28 (m, 1H), 2.44 (m, 1H), 2.38 (s, 3H), 1.87-1.12 (m, 9H), 1.08 (s, 3H), 0.89 (s, 3H), 0.88 (s, 3H), 0.77 (d, *J* = 6.7 Hz, 3H). ¹³C NMR (CD₃OD): δ 205.8, 158.9, 157.4, 143.8, 143.4, 141.0, 140.6, 127.5, 125.3, 121.6, 121.1, 113.1, 59.3, 51.5, 43.1, 42.7, 40.3, 34.4, 34.0, 33.9, 29.0, 22.1, 21.5, 19.5, 16.7, 14.0. ES-MS *m/z* 464 ([*M*+1]⁺).

EXAMPLE 83

Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide (Compound No. 139)




A. Acetic anhydride (0.040 mL, 0.42 mmol) was added to a solution of 3-[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl-5-methylphenol (Compound No. 136, 62 mg, 0.18 mmol), DMAP (2.2 mg, 0.018 mmol), and Et₃N (0.059 mL, 0.42 mmol) in THF (1.8 mL) at 0°C under argon. The solution was stirred for 5 min at 0°C then at room temperature for 3 h. The solution was concentrated, dissolved in EtOAc (40 mL), washed with saturated aqueous NaHCO₃ (5 x 10 mL) and brine (10 mL) then dried (MgSO₄) and concentrated. The colourless crystals (75 mg) that were obtained were used in the next step without further purification.

B. The colourless crystals from above (75 mg) were dissolved in MeOH (2 mL) and 10 N NaOH (0.2 mL) was added. The solution was stirred for 35 min at room temperature then concentrated. The residue was diluted with H₂O (10 mL) and adjusted to pH 6 with 10% aqueous HCl. The mixture was extracted with EtOAc (40 mL), and the organic extract was dried (MgSO₄) and concentrated to give *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]acetamide as a colourless solid (Compound No. 139, 70 mg, quantitative over 2 steps). ¹H NMR (CDCl₃): δ 7.26 (br s, 1H), 6.88 (s, 1H), 6.55 (br s, 1H), 5.99 (br s, 1H), 4.37 (m, 1H), 3.03 (d, *J* = 11.9 Hz, 1H), 2.43 (m, 1H), 2.37 (s, 3H), 2.14 (s, 3H), 1.77-1.02 (m, 12H), 0.84 (s, 3H), 0.82 (s, 3H), 0.71 (d, *J* = 6.7 Hz, 3H). ¹³C NMR (CDCl₃): δ 203.5s, 171.2, 156.2, 142.3, 140.5, 121.3, 121.0, 113.0, 58.2, 50.1, 48.6, 41.6, 41.3, 39.3, 33.5, 33.0, 32.8, 27.7, 23.6, 21.8, 21.6, 18.5, 16.5, 13.5. ES-MS *m/z* 386 ([*M*+1]⁺).

EXAMPLE 84

Synthesis of (2R,4R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-4a,8,8-trimethyl-3-methylidene-decahydronaphthalen-2-ol (Compound No. 141)




A. A solution of [(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methanone (Compound No. 51, 208 mg, 0.583 mmol) and *m*-chloroperoxybenzoic acid (154 mg, 0.687 mmol) in CH_2Cl_2 (5.8 mL) was stirred under argon at 0°C for 2 h. The reaction was quenched with 10% aqueous Na_2SO_3 (3.0 mL) and allowed to stir at room temperature for 1 h. The mixture was extracted with EtOAc (20 mL), and the organic layer was washed with saturated aqueous NaHCO_3 (5 x 10 mL) and brine (10 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:9) to afford (1aS,2R,2aS,6aS,7aR)-2-[(3,5-dimethoxyphenyl)carbonyl]-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxirene as a colourless oil (Compound No. 140, 158 mg, 73%).

B. A solution of (1aS,2R,2aS,6aS,7aR)-2-[(3,5-dimethoxyphenyl)carbonyl]-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxirene (Compound No. 140, 103 mg, 0.277 mmol) and *p*-toluenesulfonic acid (2.6 mg, 0.014 mmol) in 1,4-dioxane/ H_2O (9:1, 2.0 mL) was heated to reflux for 2 h. The solution was diluted with saturated aqueous NaHCO_3 (10 mL) and EtOAc (30 mL), and the aqueous layer was extracted with EtOAc (2 x 10 mL). The organic layers were combined, washed with brine (10 mL), then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:9-2:3) to afford (2R,4R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-4a,8,8-trimethyl-3-methylidene-decahydronaphthalen-2-ol as colourless crystals (Compound No. 141, 40 mg, 39%). ^1H NMR (CDCl_3): δ 7.11 (d, J = 2.2 Hz, 2H), 6.61 (t, J = 2.2 Hz, 1H), 5.03 (s, 1H), 4.58 (s, 1H), 4.47 (br s, 1H), 4.37 (s, 1H), 3.82 (s, 6H), 1.96-1.06 (m, 12H), 0.91 (s, 3H), 0.86 (s, 3H). ^{13}C NMR (CDCl_3): δ 200.5, 160.9 (2C), 146.9, 141.2, 112.6, 106.2 (2C), 105.0, 73.5, 57.6, 55.7 (2C), 48.0, 42.2, 41.1, 38.8, 33.5, 33.2, 31.0, 21.8, 19.1, 14.1. ES-MS m/z 373 ($[\text{M}+1]^+$).

EXAMPLE 85


Synthesis of (2R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-2-ol (Compound No. 142)



A solution of (1aS,2R,2aS,6aS,7aR)-2-[(3,5-dimethoxyphenyl)carbonyl]-1a,2a,6,6-tetramethyl-decahydronaphtho[2,3-b]oxirene (Compound No. 140, 55 mg, 0.15 mmol) and KOH (29 mg, 0.52 mmol) in MeOH (1.5 mL) was heated to reflux for 18 h then saturated aqueous NH_4Cl (10 mL) was added. The mixture was extracted with 10% MeOH/ CH_2Cl_2 (2 x 10 mL), and the organic layers were combined, washed with brine (10 mL), dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 3:7-3:2) to afford (2R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-2-ol as a colourless solid (Compound No. 142, 28 mg, 51%). ^1H NMR (CDCl_3): δ 7.09 (d, $J = 2.3$ Hz, 2H), 6.65 (t, $J = 2.3$ Hz, 1H), 4.07 (m, 1H), 3.84 (s, 6H), 1.90 (m, 2H), 1.59-1.22 (m, 13H), 0.96 (s, 3H), 0.88 (s, 3H). ^{13}C NMR (CDCl_3): δ 200.3, 161.1 (2C), 146.2, 139.9, 130.5, 107.4 (2C), 105.6, 69.4, 55.7 (2C), 45.4, 41.7, 39.1, 37.1, 33.2, 33.0, 29.0, 21.7, 19.5, 18.9, 18.7. ES-MS m/z 355 ($[\text{M}-17]^+$).

EXAMPLE 86

Synthesis of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)acetamide (Compound No. 149)



A. A solution of sclareolide (Compound No. 44, 5.76 g, 23.0 mmol) in THF (115 mL) was cooled to 0°C under argon and lithium aluminum hydride (1.92 g, 50.6 mmol) was added. After stirring at room temperature for 2 h, the mixture was cooled to 0°C and H_2O (1.9 mL) was carefully added followed by 15% aqueous NaOH (1.9 mL) and H_2O (5.7 mL). The mixture was stirred for 1 h at room temperature then dried (MgSO_4), filtered and concentrated to give (1R,2R,4aS,8aS)-1-(2-hydroxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol as a colourless solid (Compound No. 143, 5.51 g, 94%).

B. Acetic anhydride (2.04 mL, 21.6 mmol) was added to a solution of (1R,2R,4aS,8aS)-1-(2-hydroxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 143, 5.51 g, 21.7 mmol), DMAP (132 mg, 1.08 mmol), and Et_3N (3.32 mL, 23.8 mmol) in CH_2Cl_2 (108 mL) at 0°C under argon. The solution was stirred at 0°C for 20 min then at room temperature for 40 min. The solution was concentrated, dissolved in EtOAc (100 mL), washed with saturated aqueous NaHCO_3 (5 x 30 mL) and brine (20 mL) then dried (MgSO_4) and concentrated. The colourless oil, 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethyl

acetate, (Compound No. 144, 7.03 g) that was obtained was used in the next step without further purification.

C. To a solution of 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethyl acetate (Compound No. 144, 7.03 g) and pyridine (3.51 mL, 434 mmol) in CH₂Cl₂ (217 mL) was added a solution of thionyl chloride (7.89 mL, 108 mmol) and pyridine (14.0 mL, 173 mmol) in CH₂Cl₂ (54 mL) at -78°C under argon over 30 min. The solution was stirred at -78°C for 40 min then saturated aqueous NaHCO₃ (50 mL) was added and allowed to warm to room temperature. The organic phase was washed with H₂O (30 mL) and concentrated followed by azeotropic removal of pyridine using hexanes (50 mL). The residue was partitioned between EtOAc (100 mL) and saturated aqueous NaHCO₃ (40 mL). The organic phase was washed with saturated aqueous NaHCO₃ (4 x 40 mL) and brine (15 mL) then dried (MgSO₄) and concentrated. The residue was filtered through silica gel (EtOAc/hexanes, 1:19) to give 2-[(1S,4aS,8aS)-5,5,8a-trimethyl-2-methylidene-decahydronaphthalen-1-yl]ethyl acetate as a light yellow oil (Compound No. 145, 6.01 g) that was used in the next step without further purification.

D. To a solution of 2-[(1S,4aS,8aS)-5,5,8a-trimethyl-2-methylidene-decahydronaphthalen-1-yl]ethyl acetate (Compound No. 145, 6.01 g) in EtOAc (40 mL) was added 10% Pd/C (600 mg) then shaken under an atmosphere of H₂(g) at 45 psi for 1.5 h. The mixture was filtered through Celite then concentrated to give a colourless oil (5.53 g). The colourless oil (5.53 g) was dissolved in MeOH (100 mL) and 10 N NaOH (9.9 mL) was slowly added. The solution was stirred for 20 min at room temperature then cooled to 0°C and adjusted to pH 2 with concentrated HCl. The mixture was concentrated then partitioned between CH₂Cl₂ (40 mL) and H₂O (30 mL). The aqueous layer was extracted with CH₂Cl₂ (2 x 20 mL), and the combined organic layers were dried (MgSO₄) and concentrated to give a pale oil, 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethan-1-ol (Compound No. 146, 5.21 g) that was used in the next step without further purification.

E. Jones reagent was freshly prepared by adding CrO₃ (6.56 g, 65.6 mmol) to H₂SO₄ (5.89 mL, 105 mmol) at 0°C then slowly diluting the mixture with H₂O (24 mL) at 0°C. The Jones reagent was added dropwise over 10 min to a solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]ethan-1-ol (Compound No. 146, 5.21 g) in acetone (73 mL) at 0°C, and the mixture was stirred at 0°C for 1 h then at room temperature for 2 days. The mixture was diluted with *i*-PrOH (40 mL) then concentrated. The residue was diluted with H₂O (100 mL) and extracted

with CH₂Cl₂ (50 mL then 3 x 15 mL). The combined organic extracts were dried (MgSO₄) and concentrated, and the residue was partially purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:24) to give 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]acetic acid as yellow crystals (Compound No. 147, 4.87 g).


F. To a solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]acetic acid (Compound No. 147, 302 mg) in CH₂Cl₂ (6 mL) under argon was added DMF (cat.) and oxalyl chloride (0.31 mL, 3.6 mmol) then stirred at room temperature for 45 min. The solution was concentrated followed by azeotropic removal of excess oxalyl chloride with CH₂Cl₂ (3 X 10 mL). The acid chloride residue was dissolved in pyridine (3 mL), and 3,5-dimethoxyaniline (202 mg, 1.32 mmol) was added. The solution was stirred at room temperature for 17 h then concentrated. The residue was partitioned between EtOAc (35 mL) and saturated aqueous NaHCO₃ (15 mL), and the organic layer was washed with saturated aqueous NaHCO₃ (2 x 15 mL) and brine (10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 15:85) to give 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dimethoxyphenyl)acetamide as a colourless solid (Compound No. 148, 151 mg, 29% over 6 steps).

G. A solution of 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dimethoxyphenyl)acetamide (Compound No. 148, 151 mg, 0.390 mmol) in CH₂Cl₂ (4 mL) under argon was stirred at room temperature while BBr₃ (1.0 M in CH₂Cl₂, 2.5 mL, 2.5 mmol) was added then the solution was stirred at room temperature for 3 h. Methanol (10 mL) was carefully added to the solution then concentrated (3X). The residue was dissolved in *i*-PrOH/10% aqueous HCl (3:1, 4 mL) and heated to 85°C for 40 min. The solution was concentrated, and the residue was partitioned between EtOAc (20 mL) and H₂O (10 mL). The aqueous phase was adjusted to pH 9 using 1 N NaOH and extracted with EtOAc (20 mL). The combined organic layers were dried (MgSO₄) and concentrated, and the residue was purified by chromatography on silica gel (acetone/hexanes, 33:67) to give 2-[(1S,2S,4aS,8aR)-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)acetamide as a tan solid (Compound No. 149, 116 mg, 83%). ¹H NMR (CD₃OD): δ 6.55 (d, *J* = 2.4 Hz, 2H), 6.01 (t, *J* = 2.1 Hz, 1H), 2.46 (dd, *J* = 16.2, 2.1 Hz, 1H), 2.07 (dd, *J* = 16.4, 6.2 Hz, 1H), 1.80-0.94 (m, 13H), 0.87 (m, 12H). ¹³C NMR (CD₃OD): δ 175.1, 159.5 (2C), 141.6, 100.2 (2C), 99.5, 56.3, 54.1, 43.1, 40.0,

38.8, 38.0, 37.3, 35.3, 34.2, 33.9, 23.0, 22.2, 21.2, 19.8, 15.0. ES-MS m/z 360 ($[M+1]^+$).

EXAMPLE 87

Synthesis of (1S,2S,4aS,8aR)-1-[(3,5-dihydroxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 157)



A. Acetic anhydride (0.52 mL, 5.5 mmol) was added to a solution of (1S,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)(hydroxy)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 49, 2.09 g, 5.55 mmol), DMAP (34 mg, 0.28 mmol), and Et_3N (0.85 mL, 6.1 mmol) in CH_2Cl_2 (55 mL) at 0°C under argon. The solution was stirred at 0°C for 15 min then at room temperature for 45 min. The solution was concentrated, dissolved in EtOAc (100 mL), washed with saturated aqueous NaHCO_3 (5 x 20 mL) and brine (15 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:3) to give [(1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methyl acetate as a light yellow oil (Compound No. 150, 2.26 g, 97%).

B. To a solution of [(1S,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl](3,5-dimethoxyphenyl)methyl acetate (Compound No. 150,

2.26 g, 5.40 mmol) in MeOH (20 mL) was carefully added 10% Pd/C (228 mg) portionwise then shaken under an atmosphere of H₂(g) at 45 psi for 17 h. The mixture was filtered through Celite then concentrated to give a colourless oil, (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-yl acetate (Compound No. 151, 2.10 g).

C. A solution of (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-yl acetate (Compound No. 151, 2.10 g) in THF (35 mL) was cooled to 0°C under argon and lithium aluminum hydride (297 mg, 7.83 mmol) was added. After stirring at room temperature for 2 h, the mixture was cooled to 0°C and H₂O (0.3 mL) was carefully added followed by 15% aqueous NaOH (0.3 mL) and H₂O (0.9 mL). The mixture was stirred for 30 min at room temperature then dried (MgSO₄), filtered and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:3) to give (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol as a colourless oil (Compound No. 152, 1.53 g, 78% over 2 steps).

D. To a solution of (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 152, 1.53 g, 4.24 mmol) and pyridine (0.69 mL, 8.5 mmol) in CH₂Cl₂ (42 mL) was added a solution of thionyl chloride (1.54 mL, 21.2 mmol) and pyridine (2.75 mL, 34.0 mmol) in CH₂Cl₂ (11 mL) at -78°C under argon over 30 min. The solution was stirred at -78°C for 40 min then saturated aqueous NaHCO₃ (20 mL) was added and allowed to warm to room temperature. The aqueous phase was extracted with CH₂Cl₂ (20 mL), and the combined organic layers were concentrated followed by azeotropic removal of pyridine using hexanes (3 x 10 mL). The residue was partitioned between EtOAc (80 mL) and saturated aqueous NaHCO₃ (15 mL). The organic phase was washed with saturated aqueous NaHCO₃ (3 x 15 mL) and brine (10 mL) then dried (MgSO₄) and concentrated to give a yellow oil, (4aS,5S,8aS)-5-[(3,5-dimethoxyphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene (Compound No. 153, 1.50 g) that was used in the next step without further purification.

E. To a solution of (4aS,5S,8aS)-5-[(3,5-dimethoxyphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene (Compound No. 153, 1.50 g) in THF (42 mL) under argon was added BH₃·THF (1.0 M in THF, 6.3 mL, 6.3 mmol) at 0°C, and the solution was stirred at 0°C for 5 min then at room temperature for 1.8 h. The solution was cooled to 0°C, and 10 N NaOH (0.84 mL) was added followed by dropwise addition of 50% aqueous H₂O₂ (0.73 mL, 13 mmol). The mixture was stirred

at room temperature for 20 min and 65°C for 50 min then allowed to cool to room temperature. The mixture was diluted with H₂O (25 mL), and saturated aqueous Na₂SO₃ was added until excess peroxide was destroyed. The organic layer was washed with brine (15 mL), and the combined aqueous layers were extracted with EtOAc (20 mL). The combined organic layers were dried (MgSO₄) and concentrated, and the residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:3) to give [(1S,2S,4aS,8aR)-1-[(3,5-dimethoxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol as a colourless oil (Compound No. 154, 0.59 g, 39% over 2 steps).

F. To a solution of [(1S,2S,4aS,8aR)-1-[(3,5-dimethoxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalen-2-yl]methanol (Compound No. 154, 0.59 g, 1.6 mmol) in acetone (11 mL) under argon was added 2-iodoxybenzoic acid (1.37 g, 4.89 mmol), and the mixture was heated to 60°C for 3 h. The mixture was allowed to cool to room temperature then filtered and concentrated to give a yellow oil, (1S,2S,4aS,8aR)-1-[(3,5-dimethoxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carbaldehyde (Compound No. 155, 605 mg) that was used in the next step without further purification.


G. To a solution of (1S,2S,4aS,8aR)-1-[(3,5-dimethoxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carbaldehyde (Compound No. 155, 605 mg) and sulfamic acid (544 mg, 5.60 mmol) in THF/DMSO (10:1, 18 mL) at 0°C was added a solution of 80% sodium chlorite (543 mg, 4.80 mmol) in H₂O (16 mL), and the mixture was stirred at 0°C for 3 h. The mixture was diluted with EtOAc (50 mL), H₂O (10 mL) and saturated aqueous NH₄Cl (10 mL). The organic layer was washed with brine (3 x 15 mL) then dried (MgSO₄) and concentrated. The residue was partially purified by chromatography on silica gel (EtOAc/hexanes, 1:4) to give (1S,2S,4aS,8aR)-1-[(3,5-dimethoxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid as a yellow oil (Compound No. 156, 374 mg).

H. A solution of (1S,2S,4aS,8aR)-1-[(3,5-dimethoxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid (Compound No. 156, 99 mg) in CH₂Cl₂ (2.6 mL) under argon was stirred at 0°C while BBr₃ (1.0 M in CH₂Cl₂, 1.6 mL, 1.6 mmol) was added then the solution was stirred at room temperature for 23 h. Methanol (5 mL) was carefully added to the solution then concentrated (2X). The residue was diluted with *i*-PrOH/10% aqueous HCl (2:1, 3 mL) and heated to 90°C for 45 min. The solution was concentrated, and the residue was partitioned between EtOAc (30 mL) and brine (15 mL). The organic layer was dried (MgSO₄) and

concentrated, and the residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 3:97) to give (1S,2S,4aS,8aR)-1-[(3,5-dihydroxyphenyl)methyl]-5,5,8a-trimethyl-decahydronaphthalene-2-carboxylic acid as a pale solid (Compound No. 157, 11 mg, 7% over 3 steps). ¹H NMR (CD₃OD): δ 6.28 (d, *J* = 2.4 Hz, 2H), 6.08 (t, *J* = 2.3 Hz, 1H), 2.96 (m, 1H), 2.68 (dd, *J* = 13.2, 3.0 Hz, 1H), 2.40 (m, 1H), 2.19 (m, 1H), 1.85 (m, 1H), 1.79-0.95 (m, 13H), 0.87 (s, 3H), 0.85 (s, 3H). ¹³C NMR (CD₃OD): δ 179.2, 159.0 (2C), 146.3, 109.1 (2C), 101.0, 58.1, 58.0, 43.3, 40.7, 40.0 (2C), 34.3, 34.0, 32.7, 30.9, 21.9, 20.5, 19.8, 15.3. ES-MS *m/z* 345 ([M-1]⁻).

EXAMPLE 88

Synthesis of *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-2-carboxamide (Compound No. 158)




To a solution of 5-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 79 mg, 0.23 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (88 mg, 0.27 mmol), and 2-picolinic acid (34 mg, 0.28 mmol) in DMF (1.1 mL) under argon was added *N,N*-diisopropylethylamine (0.056 mL, 0.32 mmol), and the solution was stirred at room temperature for 19.5 h then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (10 mL) and brine (5 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/CH₂Cl₂, 2:3) to give *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-2-carboxamide as a colourless foam (Compound No. 158, 73 mg, 71%). ¹H NMR (CD₃OD): δ 8.74 (d, *J* = 4.2 Hz, 1H), 8.14 (d, *J* = 7.5 Hz, 1H), 8.00 (m, 1H), 7.61 (m, 1H), 6.90 (d, *J* = 2.1 Hz, 2H), 6.50 (t, *J* = 2.3 Hz, 1H), 4.47 (m, 1H), 3.37 (d, *J* = 12.0 Hz, 1H), 2.52 (m, 1H), 1.84 (m, 2H), 1.61-1.09 (m, 10H), 0.87 (s, 3H), 0.81 (s, 3H), 0.75 (d, *J* = 6.9 Hz, 3H). ¹³C NMR (CD₃OD): δ 205.4, 165.9, 159.9 (2C), 150.8,

149.8, 144.3, 139.2, 128.0, 123.3, 108.2, 107.6 (2C), 59.3, 51.3, 49.8, 42.8, 42.3, 40.3, 34.4, 33.9, 33.8, 28.6, 22.0, 19.5, 16.8, 14.0. ES-MS m/z 451 ($[M+1]^+$).

EXAMPLE 89


Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-4-carboxamide (Compound No. 159)



To a solution of 5-[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 79 mg, 0.23 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (88 mg, 0.27 mmol), and isonicotinic acid (34 mg, 0.28 mmol) in DMF (1.1 mL) under argon was added *N,N*-diisopropylethylamine (0.056 mL, 0.32 mmol), and the solution was stirred at room temperature for 19.5 h then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (10 mL) and brine (5 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/EtOAc, 1:99) to give *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-4-carboxamide as a colourless solid (Compound No. 159, 53 mg, 51%). ¹H NMR (CD₃OD): δ 8.70 (d, J = 5.7 Hz, 2H), 7.68 (m, 2H), 6.94 (d, J = 2.4 Hz, 2H), 6.48 (t, J = 2.0 Hz, 1H), 4.53 (m, 1H), 3.57 (d, J = 12.0 Hz, 1H), 2.50 (m, 1H), 1.80 (m, 2H), 1.60-1.29 (m, 6H), 1.08 (m, 4H), 0.88 (s, 6H), 0.77 (d, J = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.2, 169.3, 159.8 (2C), 150.7 (2C), 145.4, 144.5, 123.3 (2C), 107.9 (3C), 58.8, 52.1, 48.8, 42.9, 42.5, 40.1, 33.9 (3C), 28.7, 22.1, 19.5, 16.6, 14.2. ES-MS m/z 451 ($[M+1]^+$).

EXAMPLE 90

Synthesis of 3-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl]pyridin-1-ium-1-olate (Compound No. 160)




To a solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 74 mg, 0.21 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (185 mg, 0.58 mmol), and nicotinic acid *N*-oxide (80 mg, 0.58 mmol) in DMF (1.1 mL) under argon was added *N,N*-diisopropylethylamine (0.12 mL, 0.69 mmol), and the solution was stirred at room temperature for 21 h then concentrated. The residue was partitioned between EtOAc (40 mL) and saturated aqueous NaHCO₃ (10 mL). The organic phase was washed with brine (5 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:2) to give 3-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl]pyridin-1-ium-1-olate as a pale solid (Compound No. 160, 39 mg, 39%).

¹H NMR (DMSO-d₆): δ 9.59 (s, 2H), 8.60 (s, 1H), 8.50 (d, *J* = 9.3 Hz, 1H), 8.39 (d, *J* = 5.7 Hz, 1H), 7.67 (d, *J* = 7.8 Hz, 1H), 7.55 (m, 1H), 6.89 (d, *J* = 1.8 Hz, 2H), 6.47 (s, 1H), 4.40 (br s, 1H), 3.60 (d, *J* = 11.7 Hz, 1H), 2.31 (m, 1H), 1.66-1.15 (m, 8H), 0.94 (s, 4H), 0.79 (s, 6H), 0.61 (d, *J* = 6.3 Hz, 3H). ¹³C NMR (DMSO-d₆): δ 203.8, 163.2, 158.3 (2C), 142.9, 140.4, 137.5, 134.5, 126.5, 124.6, 106.7, 106.3 (2C), 55.8, 49.5, 48.6, 46.0, 41.5, 38.4, 33.1, 32.4, 32.1, 27.3, 21.4, 18.0, 15.9, 13.2. ES-MS *m/z* 467 ([*M*+1]⁺).

EXAMPLE 91

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylpiperazine-1-carboxamide (Compound No. 163)



A. A solution of isopropenyl chloroformate (51 mg, 0.42 mmol) in THF (1.0 mL) was added to a solution of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 122 mg, 0.327 mmol) and Et₃N (0.059 mL, 0.42 mmol) in THF (0.6 mL) at 0°C under argon then stirred at room temperature for 23 h. The mixture was diluted with EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (5 x 10 mL) and brine (10 mL) then dried (MgSO₄) and concentrated. After azeotropic distillation using toluene (3 x 10 mL), prop-1-en-2-yl N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamate was obtained as a yellow solid (Compound No. 161, 144 mg, 97%) and used in the next step without further purification.


B. A solution of prop-1-en-2-yl N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamate (Compound No. 161, 144 mg, 0.315 mmol), 1-methylpiperazine (38 mg, 0.38 mmol) and 1-methylpyrrolidine (29 mg, 0.34 mmol) in toluene (1.0 mL) was heated to 100°C for 25 h under argon. The mixture was allowed to cool to room temperature and partitioned between EtOAc (35 mL) and saturated aqueous NaHCO₃ (10 mL). The

organic layer was washed with saturated aqueous NaHCO_3 (2 x 10 mL) and brine (10 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{MeOH}/\text{CH}_2\text{Cl}_2$, 3:47) to give *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylpiperazine-1-carboxamide as a light yellow film (Compound No. 162, 128 mg, 82%).

C. A solution of *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylpiperazine-1-carboxamide (Compound No. 162, 128 mg, 0.256 mmol) in CH_2Cl_2 (2.6 mL) under argon was stirred at -78°C while BBr_3 (1.0 M in CH_2Cl_2 , 1.5 mL, 1.5 mmol) was added then the solution was stirred at room temperature for 6 h. Methanol (10 mL) was carefully added to the solution at 0°C then concentrated. After azeotropic distillation using MeOH (3 x 10 mL), the residue was diluted with *i*-PrOH (3.5 mL) and 10% aqueous HCl (1 mL) then heated to 80°C for 40 min. The solution was concentrated, diluted with H_2O (1 mL) then adjusted to pH 8 using saturated aqueous NaHCO_3 . The mixture was extracted with EtOAc (20 mL then 2 x 10 mL), and the organic extracts were combined, dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 100:10:1) to afford *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-4-methylpiperazine-1-carboxamide as a colourless solid (Compound No. 163, 51 mg, 42%). ^1H NMR (CD_3OD): δ 6.92 (d, J = 2.1 Hz, 2H), 6.49 (s, 1H), 6.15 (d, J = 7.5 Hz, 1H), 4.19 (m, 1H), 3.52 (m, 4H), 3.42 (d, J = 12.0 Hz, 1H), 2.49 (m, 4H), 2.38 (m, 1H), 2.34 (s, 3H), 1.68 (m, 2H), 1.59-1.22 (m, 6H), 1.07 (m, 1H), 1.02 (s, 3H), 0.85 (s, 6H), 0.70 (d, J = 6.9 Hz, 3H). ^{13}C NMR (CD_3OD): δ 206.3, 160.4, 159.8 (2C), 144.6, 107.9, 107.8 (2C), 59.1, 55.7 (2C), 52.3, 48.6, 46.1, 45.2 (2C), 42.9, 42.4, 40.1, 34.3, 33.9 (2C), 29.4, 22.2, 19.6, 16.8, 14.2. ES-MS m/z 472 ($[\text{M}+1]^+$).

EXAMPLE 92

Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-aminopropanamide (Compound No. 165)




A. To a solution of 5-[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 76 mg, 0.22 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (85 mg, 0.26 mmol), and *N*-(*tert*-butoxycarbonyl)-*beta*-alanine (50 mg, 0.26 mmol) in DMF (1.1 mL) under argon was added *N,N*-diisopropylethylamine (0.054 mL, 0.31 mmol), and the solution was stirred at room temperature for 16 h then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (10 mL) and brine (5 x 10 mL) then dried (MgSO₄) and concentrated. The residue was partially purified by chromatography on silica gel (EtOAc/CH₂Cl₂, 3:1) to give *tert*-butyl *N*-(2-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl)ethyl)carbamate as a colourless solid (Compound No. 164, 103 mg).

B. Trifluoroacetic acid (1 mL) was added to a solution of *tert*-butyl *N*-(2-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl)ethyl)carbamate (Compound No. 164, 103 mg) in CH₂Cl₂ (3 mL) under argon, and the solution was stirred at room temperature for 50 min. The solution was cooled to 0°C and adjusted to pH 8 using 10 N NaOH then diluted with H₂O (10 mL) and EtOAc (15 mL). The aqueous layer was extracted with EtOAc (15 mL), and the combined organic layers were dried (MgSO₄) and concentrated. The residue was stirred in H₂O (2 mL), and *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-aminopropanamide, a pale solid (Compound No. 165, 52 mg, 57% over 2 steps), was collected by filtration. ¹H NMR (CD₃OD): δ 6.93 (d, *J* = 1.8 Hz, 2H), 6.49 (s, 1H), 4.28 (br s, 1H), 3.42 (d, *J* = 12.3 Hz, 1H), 3.02 (br s, 2H), 2.57 (m, 2H), 2.37 (m, 1H), 1.68 (m, 2H), 1.55-1.22 (m, 6H), 1.08 (m, 1H), 1.03 (s, 3H), 0.85 (s, 6H), 0.69 (d, *J* = 6.3 Hz,

3H). ^{13}C NMR (CD_3OD): δ 206.3, 173.7, 160.0 (2C), 144.5, 108.2, 107.9 (2C), 58.8, 50.9, 48.8, 42.9, 42.3, 40.2, 38.7, 36.9, 33.9, 33.8 (2C), 28.7, 22.1, 19.5, 16.6, 14.0. ES-MS m/z 415 ($[\text{M}-1]^-$).

EXAMPLE 93

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]piperidine-4-carboxamide (Compound No. 167)




A. To a solution of 5-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 73 mg, 0.21 mmol), O-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (81 mg, 0.25 mmol), and 1-(*tert*-butoxycarbonyl)piperidine-4-carboxylic acid (58 mg, 0.25 mmol) in DMF (1.1 mL) under argon was added *N,N*-diisopropylethylamine (0.052 mL, 0.30 mmol), and the solution was stirred at room temperature for 17 h then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO_3 (10 mL) and brine (5 x 10 mL) then dried (MgSO_4) and concentrated. The residue was partially purified by chromatography on silica gel (EtOAc/ CH_2Cl_2 , 3:2) to give *tert*-butyl 4-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl}piperidine-1-carboxylate as a colourless solid (Compound No. 166, 112 mg).

B. *tert*-Butyl 4-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl}piperidine-1-carboxylate (Compound No. 166, 112 mg) was dissolved in a solution of 1.25 M HCl in *i*-PrOH (4 mL) and stirred at room temperature for 24 h. The solution was concentrated, and the residue was partitioned between EtOAc (20 mL) and saturated aqueous NaHCO_3 (20 mL). The aqueous layer was extracted with EtOAc (2 x 20 mL), and the combined organic layers were washed with brine (10 mL) then dried (MgSO_4) and concentrated. The residue was briefly stirred as a suspension in CH_2Cl_2 (2 mL), and *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-

decahydronaphthalen-2-yl]piperidine-4-carboxamide, a light yellow solid (Compound No. 167, 61 mg, 64% over 2 steps), was collected by filtration. ^1H NMR (CD_3OD): δ 8.14 (d, $J = 7.8$ Hz, 1H), 6.96 (d, $J = 2.1$ Hz, 2H), 6.50 (s, 1H), 4.24 (m, 1H), 3.45 (d, $J = 12.0$ Hz, 1H), 3.31 (m, 2H), 2.89 (m, 2H), 2.74 (m, 1H), 2.39 (m, 1H), 1.88-1.22 (m, 12H), 1.08 (m, 1H), 1.02 (s, 3H), 0.85 (s, 6H), 0.69 (d, $J = 6.3$ Hz, 3H). ^{13}C NMR (CD_3OD): δ 206.2, 176.6, 160.0 (2C), 144.5, 108.2, 107.9 (2C), 58.9, 50.8, 48.6, 45.2 (2C), 42.8, 42.3, 41.9, 40.1, 33.8 (3C), 28.6, 28.5, 28.2, 22.1, 19.5, 16.6, 14.1. ES-MS m/z 457 ($[\text{M}+1]^+$).

EXAMPLE 94

Synthesis of *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-3-carboxamide (Compound No. 168)




To a solution of (2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-amine (Compound No. 55, 72 mg, 0.19 mmol), *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (74 mg, 0.23 mmol), and nicotinic acid (28 mg, 0.23 mmol) in DMF (1.0 mL) under argon was added *N,N*-diisopropylethylamine (0.047 mL, 0.27 mmol), and the solution was stirred at room temperature for 18 h then concentrated. The residue was partitioned between EtOAc (40 mL) and saturated aqueous NaHCO_3 (10 mL). The organic phase was washed with brine (5 x 10 mL) then dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/ CH_2Cl_2 , 3:7) to give *N*-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]pyridine-3-carboxamide as a colourless solid (Compound No. 168, 69 mg, 75%). ^1H NMR (CDCl_3): δ 8.98 (br s, 1H), 8.76 (br s, 1H), 8.11 (d, $J = 8.1$ Hz, 1H), 7.43 (dd, $J = 7.7, 5.0$ Hz, 1H), 7.02 (d, $J = 1.8$ Hz, 2H), 6.62 (s, 1H), 6.36 (d, $J = 9.0$ Hz, 1H), 4.60 (m, 1H), 3.83 (s, 6H), 3.09 (d, $J = 12.3$ Hz, 1H), 2.57 (m, 1H), 1.91-1.02 (m, 12H), 0.86 (s, 3H), 0.85 (s, 3H), 0.80 (d, $J = 6.6$ Hz, 3H). ^{13}C NMR

(CDCl₃): δ 202.7, 165.8, 161.0 (2C), 152.2, 147.6, 142.6, 135.4, 131.4, 123.8, 106.4 (2C), 104.2, 58.7, 55.7 (2C), 50.2, 49.4, 41.9, 41.4, 39.5, 33.8, 33.3, 33.1, 27.8, 21.8, 18.5, 16.5, 13.6. ES-MS m/z 479 ([M+1]⁺).

EXAMPLE 95

Synthesis of 5-[[[(1S,2S,3R,4aS,8aS)-3-[(3-aminopropyl)amino]-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol dihydrochloride
(Compound No. 170)




A. A solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 73 mg, 0.21 mmol) and *tert*-butyl N-(3-oxopropyl)carbamate (45 mg, 0.26 mmol) in MeOH (4.2 mL) was heated to reflux under argon for 35 min. After allowing the solution to cool to room temperature, sodium borohydride (16 mg, 0.42 mmol) was added, and the mixture was stirred for 2 h then concentrated. The residue was dissolved in EtOAc (30 mL) and washed with brine (10 mL). The aqueous phase was extracted with EtOAc (10 mL), and the combined organic layers were dried (MgSO₄) and concentrated. The residue was partially purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:5:1) to give a colourless solid (87 mg). The colourless solid (87 mg) was dissolved in EtOAc (30 mL) and washed with H₂O (18 x 15 mL) and brine (10 mL) then dried (MgSO₄) and concentrated to give *tert*-butyl N-(3-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}propyl)carbamate as a colourless solid (Compound No. 169, 78 mg, 74%).

B. *tert*-Butyl N-(3-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}propyl)carbamate (Compound No. 169, 78 mg, 0.16 mmol) was dissolved in a solution of 1.25 M HCl in *i*-PrOH (4 mL) and stirred at room temperature for 18.5 h. The solution was concentrated followed by azeotropic distillation using MeOH (3 mL). The residue was dissolved in MeOH (0.5 mL), and Et₂O (40 mL) was added to precipitate a colourless solid. The supernatant was decanted, and the remaining solid was washed by decantation with Et₂O (4 x 10

mL). Remaining traces of solvent were removed under vacuum to afford 5-
 {[(1S,2S,3R,4aS,8aS)-3-[(3-aminopropyl)amino]-2,5,5,8a-tetramethyl-
 decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol dihydrochloride as a colourless
 solid (Compound No. 170, 57 mg, 77%). ¹H NMR (CD₃OD): δ 6.99 (d, *J* = 2.1 Hz, 2H),
 6.49 (t, *J* = 2.0 Hz, 1H), 3.73 (d, *J* = 12.3 Hz, 1H), 3.63 (br s, 1H), 3.26-3.07 (m, 4H),
 2.63 (m, 1H), 2.28 (m, 2H), 2.04 (m, 1H), 1.84 (m, 1H), 1.63-1.31 (m, 6H), 1.06 (m,
 4H), 1.01 (s, 3H), 0.94 (d, *J* = 7.2 Hz, 3H), 0.90 (s, 3H). ¹³C NMR (CD₃OD): δ 204.4,
 160.0 (2C), 143.8, 108.6, 108.0 (2C), 61.5, 57.5, 46.9, 44.2, 42.4, 41.7, 40.1, 38.2,
 34.2, 33.8, 33.5, 24.9, 23.1, 21.9, 19.3, 16.2, 14.2. ES-MS *m/z* 403 ([M+1]⁺).

EXAMPLE 96

Synthesis of N-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-
 tetramethyl-decahydronaphthalen-2-yl]-6-aminopyridine-3-carboxamide (Compound
 No. 171)




To a solution of 5-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-
 decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 69 mg, 0.20
 mmol), O-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (77 mg,
 0.24 mmol), and 6-aminonicotinic acid (33 mg, 0.24 mmol) in DMF (1.0 mL) under
 argon was added *N,N*-diisopropylethylamine (0.049 mL, 0.28 mmol), and the solution
 was stirred at room temperature for 21 h then concentrated. The residue was
 partitioned between EtOAc (30 mL) and H₂O (10 mL). The organic phase was washed
 with brine (3 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified
 by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:1) to give *N*-
 [(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-
 decahydronaphthalen-2-yl]-6-aminopyridine-3-carboxamide as a colourless solid
 (Compound No. 171, 40 mg, 43%). ¹H NMR (CD₃OD): δ 8.39 (d, *J* = 2.1 Hz, 1H), 8.18
 (d, *J* = 9.3 Hz, 1H), 7.86 (dd, *J* = 8.9, 2.3 Hz, 1H), 6.95 (d, *J* = 1.8 Hz, 2H), 6.61 (d, *J* =

8.7 Hz, 1H), 6.49 (t, $J = 2.0$ Hz, 1H), 4.50 (m, 1H), 3.58 (d, $J = 12.0$ Hz, 1H), 2.47 (m, 1H), 1.77 (m, 2H), 1.60-1.24 (m, 6H), 1.08 (m, 1H), 1.05 (s, 3H), 0.87 (s, 6H), 0.75 (d, $J = 6.9$ Hz, 3H). ^{13}C NMR (CD_3OD): δ 206.4, 169.5, 169.4, 162.6, 159.8 (2C), 148.6, 144.6, 138.7, 121.2, 109.1, 107.9 (2C), 58.9, 51.5, 51.4, 42.8, 42.4, 40.1, 34.1, 33.9 (2C), 29.0, 22.1, 19.5, 16.7, 14.2. ES-MS m/z 466 ($[\text{M}+1]^+$).

EXAMPLE 97

Synthesis of (4S,4aS,8aS)-4-[[[(3,5-dimethoxybenzene)sulfonyl]methyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol (Compound No. 173)



A. A solution of (1R,2R,4aS,8aS)-1-[(3,5-dimethoxyphenyl)sulfanylmethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 80, 40 mg, 0.1 mmol) and *m*-chloroperoxybenzoic acid (77%, 53 mg, 0.24 mmol) in CH_2Cl_2 (3 mL) under argon was stirred at room temperature for 4 h. The mixture was concentrated, diluted with saturated aqueous NaHCO_3 (15 mL) and EtOAc (15 mL). The organic layer was washed with brine (1 x 15 mL), dried (Na_2SO_4), concentrated and purified by silica gel chromatography (hexanes:EtOAc, 3:1 to 1:1) to afford (1R,2R,4aS,8aS)-1-[(3,5-dimethoxybenzene)sulfonylmethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol as a red solid (29 mg, 67%).


B. To a solution of (1R,2R,4aS,8aS)-1-[(3,5-dimethoxybenzene)sulfonylmethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol from above (50 mg, 0.118 mmol) in MeCN (0.3 mL) under argon was added trimethylsilylmethanesulfonate (0.03 mL, 0.177 mmol) and the reaction mixture stirred for 24 h. The reaction mixture was diluted with water (5 mL), washed with saturated aqueous NaHCO_3 (5 mL), and water (5 mL), dried over Na_2SO_4 , filtered and concentrated to afford Compound No. 172 as a mixture of double bond isomers (46 mg, 96%, ~1:1) as a red oil.

C. To the reaction mixture containing Compound No. 172 (46 mg, 0.113 mmol) in THF (1 mL) under argon at 0°C was slowly added $\text{BH}_3\cdot\text{THF}$ (1.0 M, 0.2 mL,

0.170 mmol) and stirred for 15 min. The reaction mixture was then warmed to room temperature and stirred for another 90 min. The mixture was cooled to 0°C and aqueous NaOH (10 M, 0.1 mL, 0.226 mmol) was carefully added, followed by the addition of H₂O₂ (50% wt, 0.02 mL, 0.339 mmol), and stirred for 45 min. The reaction was allowed to warm to room temperature and was stirred for 24 h after which water (5 mL) and solid Na₂SO₃ (excess) were added. The resultant mixture was extracted using EtOAc (5 mL), then washed with brine (5 mL), dried over Na₂SO₄, filtered and concentrated under vacuum. The crude product was then purified using column chromatography (hexanes:EtOAc, 4:1 to 3:1 to 2:1 to 1:1 to 1:2) to afford (4S,4aS,8aS)-4-[(3,5-dimethoxybenzene)sulfonylmethyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-ol as a colourless oil (Compound No. 173, 11 mg, 23%). ¹H NMR (CDCl₃): δ 7.05 (s, 1H), 6.70 (s, 1H), 3.90 (s, 6H), 3.10 (m, 2H), 2.30 (d, 1H), 1.75-1.45 (m, 6H), 1.44 (s, 3H), 1.42-1.02 (m, 4H), 0.98 (s, 3H), 0.89 (s, 3H), 0.85 (s, 3H). ES-MS m/z 442.3 ([M+18]⁺).

EXAMPLE 98

Synthesis of 1-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]guanidine (Compound No. 175)




A. A solution of 3-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl-5-methylphenol (Compound No. 136, 60 mg, 0.17 mmol), *N,N'*-bis(*tert*-butoxycarbonyl)-1*H*-pyrazole-1-carboxamide (65 mg, 0.21 mmol), and *N,N*-diisopropylethylamine (0.035 mL, 0.20 mmol) in DMF (0.25 mL) under argon was stirred at room temperature for 18 h then concentrated. The residue was dissolved in EtOAc (30 mL) and washed with H₂O (10 mL), saturated aqueous NaHCO₃ (4 x 10 mL) and brine (10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:99) to give *tert*-butyl *N*-[(1*Z*)-{[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}{[(*tert*-

butoxy)carbonyl]imino}}methyl]carbamate as a colourless solid (Compound No. 174, 86 mg, 84%).

B. To a suspension of *tert*-butyl *N*-[(1*Z*)-{[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}{[(*tert*-butoxy)carbonyl]imino}}methyl]carbamate (Compound No. 174, 85 mg, 0.15 mmol) in CH₂Cl₂ (1.5 mL) was added trifluoroacetic acid (0.75 mL, 10 mmol) under argon at room temperature. The resulting homogeneous solution was stirred at room temperature for 2 h then concentrated, and the residue was partitioned between EtOAc (40 mL) and saturated aqueous NaHCO₃ (10 mL). The aqueous phase was extracted with EtOAc (2 x 10 mL), and the organic layers were combined, dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:1), giving a green solid (53 mg). The solid was suspended in 10% aqueous HCl and ether, then filtered and washed with ether. The solid was dissolved in MeOH and concentrated. The residue was diluted with EtOAc (12 mL) and washed with saturated aqueous NaHCO₃ (10 mL). The aqueous layer was extracted with EtOAc (5 mL), and the organic layers were combined, dried (MgSO₄) and concentrated to afford 1-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]guanidine as a colourless solid (Compound No. 175, 34 mg, 61%). ¹H NMR (CD₃OD): δ 7.66 (s, 1H), 7.22 (s, 1H), 6.82 (s, 1H), 3.81 (m, 1H), 3.58 (d, *J* = 12.1 Hz, 1H), 2.45 (m, 1H), 2.33 (s, 3H), 1.82-1.21 (m, 8H), 1.01 (m, 4H), 0.86 (s, 6H), 0.80 (d, *J* = 6.7 Hz, 3H). ES-MS *m/z* 386 ([*M*+1]⁺).

EXAMPLE 99

Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide (Compound No. 176)




A. To a solution of 3-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenol (Compound No. 136, 73 mg, 0.21 mmol) and Et₃N (0.070 mL, 0.50 mmol) in CH₂Cl₂ (2.1 mL) under argon was added methanesulfonyl chloride (0.035 mL, 0.45 mmol) at 0°C. The solution was stirred for 30 min at 0°C then at room temperature for 21 h. The solution was concentrated, and the residue was dissolved in EtOAc (40 mL) then washed with saturated aqueous NaHCO₃ (3 x 10 mL) and brine (10 mL). The organic phase was dried (MgSO₄) and concentrated, and the residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:99-CH₂Cl₂/MeOH/NH₄OH, 100:10:0.6) to give 3-[[[(1S,2S,3R,4aS,8aS)-3-methanesulfonamido-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenyl methanesulfonate as a colourless film (42 mg, 40%).

B. A solution of 3-[[[(1S,2S,3R,4aS,8aS)-3-methanesulfonamido-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]-5-methylphenyl methanesulfonate (42 mg, 0.084 mmol) and sodium hydroxide (25 mg, 0.63 mmol) in 10:1 MeOH/H₂O (4.4 mL) was heated to reflux for 20 h. The solution was concentrated, and the residue was partitioned between 10% MeOH/CH₂Cl₂ (40 mL) and H₂O (10 mL). The aqueous layer was adjusted to pH 8 with 10% aqueous HCl and extracted with 10% MeOH/CH₂Cl₂ (2 x 10 mL). The organic layers were combined, dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:4-3:2) to give *N*-[(2R,3S,4S,4aS,8aS)-4-[(3-hydroxy-5-methylphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide as a light yellow solid (Compound No. 176, 33 mg, 94%). ¹H NMR (CD₃OD): δ 7.38 (s, 1H), 7.16 (s, 1H), 6.85 (s, 1H), 3.74 (m, 1H), 3.39 (d, *J* = 12.0 Hz, 1H), 2.97 (s, 3H), 2.37 (m, 4H), 1.90-1.15 (m, 8H), 1.02 (m, 4H), 0.92 (s, 3H), 0.87 (s, 3H), 0.79 (d, *J* = 6.6 Hz, 3H). ES-MS *m/z* 422 ([*M*+1]⁺).

EXAMPLE 100

Synthesis of 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]guanidine (Compound No. 178)




A. A solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 148 mg, 0.428 mmol), *N,N'*-bis(*tert*-butoxycarbonyl)-1*H*-pyrazole-1-carboxamidine (162 mg, 0.522 mmol), and *N,N*-diisopropylethylamine (0.09 mL, 0.5 mmol) in DMF (1.1 mL) under argon was stirred at room temperature for 19 h then concentrated. The residue was dissolved in EtOAc (30 mL) and washed with H₂O (10 mL), saturated aqueous NaHCO₃ (2 x 10 mL) and brine (5 x 10 mL) then dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:99) to give *tert*-butyl N-[(1*Z*)-{[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}{[(*tert*-butoxy)carbonyl]imino})methyl]carbamate as a colourless solid (Compound No. 177, 179 mg, 71%).

B. To a suspension of *tert*-butyl N-[(1*Z*)-{[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]amino}{[(*tert*-butoxy)carbonyl]imino})methyl]carbamate (Compound No. 177, 179 mg, 0.305 mmol) in CH₂Cl₂ (3.2 mL) was added trifluoroacetic acid (1.59 mL) under argon at room temperature. The resulting homogeneous solution was stirred at room temperature for 1.5 h then concentrated. After azeotropic distillation using CH₂Cl₂ (2 x 5 mL), the residue was partitioned between 10% MeOH/CH₂Cl₂ (40 mL) and saturated aqueous NaHCO₃ (20 mL). The aqueous phase was extracted with 10% MeOH/EtOAc (4 x 15 mL), and the organic layers were combined, dried (MgSO₄) and concentrated to afford 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]guanidine as a colourless solid (Compound No. 178, 115 mg, 97%). ¹H NMR (CD₃OD): δ 6.91 (d, *J* = 2.1 Hz, 2H), 6.51 (t, *J* = 2.3 Hz, 1H), 3.84 (m, 1H), 3.31 (m, 1H), 2.47 (m, 1H), 1.74 (m, 2H), 1.56-1.09 (m, 7H), 1.04 (s, 3H), 0.88 (s, 6H), 0.77 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 205.3, 160.2 (2C), 158.6, 144.2, 108.4, 107.8 (2C), 58.3, 54.1, 48.9, 42.7, 42.3, 40.2, 34.0, 33.9, 33.7, 28.4, 22.1, 19.4, 16.5, 13.9. ES-MS *m/z* 388 ([*M*+1]⁺).

EXAMPLE 101

Synthesis of *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide (Compound No. 179)




A. To a solution of 5-[(1*S*,2*S*,3*R*,4*aS*,8*aS*)-3-amino-2,5,5,8*a*-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 118 mg, 0.342 mmol) and Et₃N (0.16 mL, 1.1 mmol) in CH₂Cl₂ (3.4 mL) under argon was added methanesulfonyl chloride (0.08 mL, 1 mmol) at 0°C. The solution was stirred for 17 h at room temperature then concentrated. The residue was dissolved in EtOAc (40 mL) and washed with saturated aqueous NaHCO₃ (5 x 10 mL) and brine (10 mL) then dried (MgSO₄) and concentrated to give a brown oil (169 mg).

B. To a solution of the brown oil from above (169 mg) in MeOH (5.8 mL) was added 10 N NaOH (0.29 mL, 2.9 mmol), and the solution was heated to reflux for 72 h then concentrated. The residue was partitioned between 10% MeOH/CH₂Cl₂ (40 mL) and H₂O (10 mL), and the aqueous layer was adjusted to pH 9 with 10% aqueous HCl. The aqueous layer was extracted with 10% MeOH/CH₂Cl₂ (3 x 10 mL), and the organic layers were combined, dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (MeOH/CH₂Cl₂, 1:19) to give *N*-[(2*R*,3*S*,4*S*,4*aS*,8*aS*)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4*a*,8,8-tetramethyl-decahydronaphthalen-2-yl]methanesulfonamide as a colourless solid (Compound No. 179, 23 mg, 16% over 2 steps). ¹H NMR (CD₃OD): δ 6.90 (d, *J* = 1.9 Hz, 2H), 6.48 (m, 1H), 3.73 (m, 1H), 3.30 (m, 1H), 2.97 (s, 3H), 2.35 (m, 1H), 1.85 (m, 1H), 1.71 (m, 1H), 1.59-1.19 (m, 6H), 1.05 (m, 1H), 1.03 (s, 3H), 0.91 (s, 3H), 0.86 (s, 3H), 0.79 (d, *J* = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 206.1, 159.8 (2C), 144.5, 108.0, 107.9 (2C), 58.6, 56.3, 48.1, 42.9, 42.1 (2C), 40.2, 34.7, 33.8 (2C), 30.0, 22.2, 19.5, 17.1, 14.0. ES-MS *m/z* 424 ([*M*+1]⁺).

EXAMPLE 102


Synthesis of 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylthiourea (Compound No. 180)



To a solution of 5-[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl}benzene-1,3-diol (Compound No. 56, 54 mg, 0.16 mmol) in THF (1.0 mL) was added methyl isothiocyanate (24 mg, 0.33 mmol) under argon. The solution was stirred at room temperature for 66 h then concentrated. The residue was purified by chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$, 100:10:2) to give 1-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]-3-methylthiourea as a colourless solid (Compound No.180, 46 mg, 71%). ^1H NMR (CD_3OD): δ 6.89 (m, 2H), 6.50 (s, 1H), 4.60 (br s, 1H), 3.29 (m, 1H), 3.04 (s, 3H), 2.41 (m, 1H), 1.87 (m, 1H), 1.68-1.04 (m, 11H), 0.86 (s, 3H), 0.84 (s, 3H), 0.73 (d, $J = 6.6$ Hz, 3H). ^{13}C NMR (CD_3OD): δ 205.9, 184.5, 159.8 (2C), 144.5, 108.1, 107.7 (2C), 59.4, 55.3, 49.3, 42.9, 42.4, 40.3, 34.4, 33.9 (2C), 31.6, 28.3, 22.1, 19.5, 16.6, 14.0. ES-MS m/z 419 ($[\text{M}+1]^+$).

EXAMPLE 103

Synthesis of 2-[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl)methyl)sulfamoyl]benzoic acid (Compound No. 181)




A. A solution of 5-[[[(1S,2S,3R,4aS,8aS)-3-amino-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]carbonyl]benzene-1,3-diol (Compound No. 56, 143 mg, 0.414 mmol), 2-(1,1,3-trioxo-2,3-dihydro-1 λ^6 ,2-benzothiazol-2-yl)acetic acid (120 mg, 0.497), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (237 mg, 0.455 mmol) and *N,N*-diisopropylethylamine (0.22 mL, 1.3 mmol) in CH₂Cl₂ (2.8 mL) under argon was stirred at room temperature for 42 h then concentrated. The residue was partitioned between EtOAc (40 mL) and saturated aqueous NaHCO₃ (10 mL). The organic layer was washed with saturated aqueous NaHCO₃ (3 x 10 mL) and brine (10 mL) then dried (MgSO₄) and concentrated. The residue was partially purified by chromatography on silica gel (CH₂Cl₂/MeOH/NH₄OH, 100:10:1) then filtered through silica gel (acetone/CH₂Cl₂, 1:4) to give a colourless film (137 mg).

B. The colourless film from above (137 mg) was dissolved in 1 N NaOH (2 mL) and stirred at room temperature for 1 h. The solution was diluted with MeOH (1 mL) and stirred at room temperature for 1.2 h then adjusted to pH 2 with concentrated HCl. The mixture was diluted with EtOAc (40 mL) and H₂O (10 mL), and the organic layer was washed with brine (2 x 10 mL) then dried (MgSO₄) and concentrated to give 2-[[[(2R,3S,4S,4aS,8aS)-4-[(3,5-dihydroxyphenyl)carbonyl]-3,4a,8,8-tetramethyl-decahydronaphthalen-2-yl]carbamoyl]methyl)sulfamoyl]benzoic acid as a colourless solid (Compound No. 181, 134 mg, 55% over 2 steps). ¹H NMR (DMSO-d₆): δ 9.58 (s, 2H), 8.07 (d, J = 9.6 Hz, 1H), 7.92 (m, 1H), 7.78 (m, 1H), 7.67 (m, 2H), 7.20 (m, 1H), 6.83 (d, J = 1.5 Hz, 2H), 6.47 (t, J = 2.1 Hz, 1H), 4.00 (m, 1H), 3.70 (m, 2H), 3.31 (m, 1H), 2.14 (m, 1H), 1.56-1.06 (m, 8H), 0.89 (m, 4H), 0.74 (s, 3H), 0.72 (s, 3H), 0.39 (d, J = 6.6 Hz, 3H). ¹³C NMR (CD₃OD): δ 205.9, 170.2 (2C), 159.8 (2C), 144.4, 139.3, 134.0, 133.3, 132.5, 132.0, 130.5, 108.0, 107.9 (2C), 58.9, 51.2, 46.7, 42.8, 42.3, 40.2, 34.0, 33.9, 33.8, 28.4, 22.1, 19.5, 16.5, 13.9. ES-MS m/z 585 ([M-1]⁻).

EXAMPLE 104

Synthesis of N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-((4-(trifluoromethyl)benzamido)methyl)-decahydronaphthalen-2-yl)nicotinamide
(Compound No. 196)



A. To a solution of Compound No. 39 (3.85 g, 16.2 mmol) in CH_2Cl_2 (75 mL) at 0°C under argon was added SnCl_4 (3.8 mL, 32.3 mmol) and the reaction stirred at 0°C for 1 hour then at room temperature for 2 hours. The mixture was cooled to 0°C and diluted with water (25 mL) then was stirred vigorously for 30 minutes while warming to room temperature. To the mixture was added brine (25 mL) then the layers were separated. The aqueous layer was further extracted with CH_2Cl_2 (2 x 75 mL) and the combined organic extracts dried (MgSO_4) and concentrated to a green oil.

B. To solution of the above crude green oil (Compound No. 183, 3.56 g, 16.2 mmol) in MeOH (75 mL) at 0°C under argon was added NaBH₄ (625 mg, 16.2 mmol) and the reaction was allowed to warm to room temperature while stirring for 3 days. The reaction was cooled to 0°C then was added saturated NaHCO₃ solution (10 mL). The mixture was concentrated to remove most of the MeOH. The residue was taken up in EtOAc (300 mL) and washed with half-saturated NaCl solution (3 x 30 mL), dried (MgSO₄) and concentrated to a yellow oil.

C. To a solution of the above crude yellow oil (Compound No. 184, 3.59 g, 16.2 mmol) and imidazole (4.4 g, 64.8 mmol) in DMF (40 mL) at 0°C under argon was added TBDPSCI (8.3 mL, 32.4 mmol) and the reaction was allowed to warm to room temperature while stirring for 18 hours. The mixture was diluted with Et₂O (300 mL) and washed with half-saturated NaCl solution (3 x 30 mL), dried (MgSO₄) and concentrated. The resulting yellow oil was filtered through a silica gel plug, eluting with 10% EtOAc/Hexanes (300 mL) and concentrated to a yellow oil.

D. To a solution of the above yellow oil (Compound No. 185, 7.44 g, 16.2 mmol) in THF (40 mL) at 0°C under argon was added BH₃-THF (1.0 M in THF, 48.6 mL, 48.6 mmol) and the reaction was allowed to warm to room temperature while stirring for 20 hours. The mixture was cooled to 0°C then water (25 mL) was slowly added followed by NaBO₃·4H₂O (7.5 g). The mixture was stirred at room temperature for 30 minutes then heated at reflux for 2.5 hours. The mixture was cooled to room temperature and was concentrated to remove most of the THF. The residue was taken up in EtOAc (300 mL) and washed with half-saturated NaCl solution (6 x 30 mL), dried (MgSO₄) and concentrated to a colourless oil.

E. A solution of the above colourless oil (Compound No. 186, 7.90 g, 16.2 mmol) and IBX (9.07 g, 32.4 mmol) in MeCN (100 mL) under argon was heated at 75°C for 3.5 hours. The mixture was cooled to room temperature then was filtered through a silica gel plug eluting with 30% EtOAc/Hexanes (300 mL) and concentrated. The crude product was purified by flash chromatography on silica gel (5% EtOAc/hexanes) to afford Compound No. 187 (3.95 g, 51%) as a colourless oil.

F. A solution of Compound No. 187 (3.95 g, 8.28 mmol) and NaOH (2 M in water, 4.1 mL, 8.3 mmol) in MeOH (28 mL) and THF (14 mL) was heated at reflux for 1.5 hours. The mixture was cooled to 0°C then diluted with saturated NaHCO₃ solution (40 mL) and concentrated to remove most of the MeOH and THF. The residue was taken up in EtOAc (350 mL), washed with half-saturated NaCl solution (3 x 30 mL),

dried (MgSO_4) and concentrated to afford Compound No. 188 (3.86 g, 98%) as slightly yellow oil.

G. A mixture of Compound No. 188 (3.86 g, 8.10 mmol), 4-methoxybenzylamine (3.20 mL, 24.3 mmol), sodium triacetoxyborohydride (15.5 mg, 72.9 mmol), and AcOH (1.4 mL, 24 mmol) in CH_2Cl_2 (100 mL) under argon was stirred at room temperature for 17 hours. The mixture was cooled to 0°C then was added saturated NaHCO_3 solution (120 mL). After stirring 15 minutes at 0°C and 45 minutes at room temperature the mixture was extracted with EtOAc (300 mL). The organic extract was washed with half-saturated NaCl solution (3 x 30 mL), dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hexanes, 1:9) to afford Compound No. 189 (3.80 g, 78%) as a slightly yellow oil.

H. To a solution of Compound No. 189 (3.80 g, 6.35 mmol) and $\text{NH}_4\text{CO}_2\text{H}$ (4.13 g, 63.5 mmol) in MeOH (100 mL) and EtOAc (12 mL) flushed thoroughly with Argon was added a slurry of Pd (10% on carbon, 675 mg, 0.63 mmol) in MeOH (15 mL). The reaction was heated at reflux for 18 hours then was cooled to room temperature. The mixture was filtered through a Celite pad eluting with EtOAc (200 mL) then was concentrated. The residue was taken up in 0.5 M aqueous NaOH solution (300 mL) and extracted with CH_2Cl_2 (3 x 100 mL). The combined organic extracts were dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (30% EtOAc/hexanes then 10% MeOH/EtOAc with 1% NH_4OH) to afford Compound No. 190 (2.44 g, 80%) as a viscous oil.

I. A mixture of Compound No. 190 (466 mg, 0.975 mmol), nicotinic acid (245 mg, 1.95 mmol) and DCC (366 mg, 1.76 mmol) in CH_2Cl_2 (20 mL) under argon was stirred at room temperature for 18 hours. The mixture was filtered through a silica gel plug eluting with EtOAc (150 mL) and concentrated to a white solid.

J. A mixture of crude Compound No. 191 (568 mg, 0.975 mmol) and TBAF (1.0 M in THF, 2.9 mL, 2.9 mmol) in THF (20 mL) under argon was heated at reflux for 2 hours then was cooled and concentrated. The residue was purified by chromatography on silica gel (50% then 80% EtOAc/hexanes then 10% MeOH/EtOAc) to afford Compound No. 192 (308 mg, 92%) as a white solid.

K. To a solution of Compound No. 192 (241 mg, 0.70 mmol) in pyridine (3.5 mL) and CH_2Cl_2 (3 mL) at 0°C under argon was added MsCl (0.17 mL, 2.1 mmol). After 45 minutes saturated NaHCO_3 solution (3 mL) was added and the mixture stirred at room temperature for 30 minutes. The mixture was diluted with EtOAc (75 mL) and

washed with half-saturated NaCl solution (3 x 15 mL), dried (MgSO_4) and concentrated to afford Compound No. 193 (289 mg, 97%) as a white solid.


L. A mixture of Compound No. 193 (284 mg, 0.67 mmol) and NaN_3 (131 mg, 2.0 mmol) in DMF (6 mL) at 60°C under argon was stirred for 16 hours. The mixture was cooled, diluted with Et_2O (75 mL), washed with half-saturated NaCl solution (3 x 15 mL), dried (MgSO_4) and concentrated. The residue was purified by chromatography on silica gel (50% then 80% EtOAc/hexanes) to afford Compound No. 194 (211 mg, 85%) as a white foam.

M. A solution of Compound No. 194 (207 mg, 0.56 mmol), triphenylphosphine (742 mg, 2.8 mmol) and water (0.05 mL, 2.8 mmol) in THF (10 mL) was stirred at room temperature for 24 hours. The mixture was concentrated and the residue was purified by chromatography on silica gel (10% MeOH/EtOAc with 1% NH_4OH then 15% MeOH/EtOAc with 3% NH_4OH) to afford Compound No. 195 (175 mg, 91%) as a white foam.

N. A mixture of Compound No. 195 (25 mg, 0.073 mmol), 4-trifluoromethyl benzoic acid (43 mg, 0.22 mmol) and DCC (38 mg, 0.18 mmol) in CH_2Cl_2 (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (50% then 66% EtOAc/hexanes) to afford Compound No. 196 (41 mg, 100%) as a white solid. ^1H NMR (CDCl_3): δ 8.95 (m, 1H), 8.73 (m, 1H), 8.08 (m, 1H), 7.80 (m, 2H), 7.67 (m, 2H), 7.40 (m, 1H), 6.30 (m, 1H), 6.18 (m, 1H), 4.50 (m, 1H), 4.10 (m, 1H), 3.75 (m, 1H), 3.43 (m, 1H), 3.25 (m, 1H), 2.1-1.0 (12H), 0.98 (s, 3H), 0.85 (s, 3H), 0.80 (s, 3H). ES-MS m/z 516 ($[\text{M}+1]^+$).

EXAMPLE 105


Synthesis of N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide (Compound No. 197)



A mixture of Compound No. 195 (25 mg, 0.073 mmol), 2-furoic acid (25 mg, 0.22 mmol) and DCC (38 mg, 0.18 mmol) in CH₂Cl₂ (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (EtOAc) to afford Compound No. 197 (33 mg, 100%) as a white solid. ¹H NMR (CDCl₃): δ 8.95 (m, 1H), 8.73 (m, 1H), 8.08 (m, 1H), 7.40 (2H), 7.06 (m, 1H), 6.45 (m, 1H), 6.28 (2H), 4.50 (m, 1H), 4.05 (m, 1H), 3.75 (m, 1H), 3.45 (m, 1H), 3.19 (m, 1H), 2.1-1.0 (12H), 0.98 (s, 3H), 0.85 (s, 3H), 0.80 (s, 3H). ES-MS *m/z* 438 ([M+1]⁺).

EXAMPLE 106


Synthesis of N-((2R,3S,4S,4aS)-4-(isonicotinamidomethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide (Compound No. 198)



A mixture of Compound No. 195 (25 mg, 0.073 mmol), isonicotinic acid (27 mg, 0.22 mmol) and DCC (38 mg, 0.18 mmol) in CH₂Cl₂ (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (EtOAc then 10% MeOH/EtOAc) to afford Compound No. 198 (31 mg, 94%) as a white solid. ¹H NMR (CDCl₃): δ 8.95 (m, 1H), 8.73 (3H), 8.08 (m, 1H), 7.55 (2H), 7.40 (m, 1H), 6.23 (m, 1H), 6.10 (m, 1H), 4.50 (m, 1H), 3.78 (m, 1H), 3.5 (m, 1H), 3.25 (m, 1H), 2.1-1.4 (5H), 1.3-1.0 (6H), 0.98 (s, 3H), 0.85 (s, 3H), 0.80 (s, 3H). ES-MS *m/z* 449 ([M+1]⁺).

EXAMPLE 107


Synthesis of N-(((1S,2S,3R,8aS)-2,5,5,8a-tetramethyl-3-(nicotinamido)decahydronaphthalen-1-yl)methyl)nicotinamide (Compound No. 199)



A mixture of Compound No. 195 (25 mg, 0.073 mmol), nicotinic acid (27 mg, 0.22 mmol) and DCC (38 mg, 0.18 mmol) in CH_2Cl_2 (5 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (EtOAc then 10% MeOH/EtOAc) to afford Compound No. 198 (36 mg, 100%) as a white solid. ^1H NMR (CDCl_3): δ 8.92 (2H), 8.70 (2H), 8.08 (2H), 7.39 (2H), 6.38 (m, 1H), 6.08 (m, 1H), 4.50 (m, 1H), 3.78 (m, 1H), 3.5 (m, 1H), 3.25 (m, 1H), 2.05 (2H), 1.90 (m, 1H), 1.8-1.4 (3H), 1.3-1.0 (6H), 0.98 (s, 3H), 0.87 (s, 3H), 0.82 (s, 3H). ES-MS m/z 449 ($[\text{M}+1]^+$).

EXAMPLE 108

Synthesis of N-((2R,3S,4S,4aS)-4-((1H-imidazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide (Compound No. 200)




To a solution of imidazole (81 mg, 1.2 mmol) in DMF (3 mL) at 0°C under argon was added NaH (60% in mineral oil, 48 mg, 1.2 mmol). After 1.5 hours Compound No. 193 (50 mg, 0.12 mmol) was added and the reaction was stirred at room temperature. After 17 hours the reaction was stirred at 60°C for 4 hours then cooled to 0°C , quenched with saturated NaHCO_3 solution (3 mL), diluted with Et_2O (40 mL), washed with half-saturated NaCl solution (3 x 8 mL), dried (MgSO_4) and concentrated. The

residue was purified by chromatography (50% then 100% EtOAc/hexanes then 10% then 20% MeOH/EtOAc) to afford Compound No. 200 (24 mg, 51%) as a white solid. ^1H NMR (CDCl_3): δ 8.85 (m, 1H), 8.62 (m, 1H), 8.00 (m, 1H), 7.33 (2H), 7.10 (m, 1H), 6.85 (2H), 4.53 (m, 1H), 4.02 (m, 1H), 3.70 (m, 1H), 2.02 (1H), 1.85 (2H), 1.8-1.4 (5H), 1.30-1.05 (3H), 1.00 (d, 3H), 0.95 (s, 3H), 0.85 (6H). ES-MS m/z 395 ($[\text{M}+1]^+$).

EXAMPLE 109


Synthesis of N-((2R,3S,4S,4aS)-4-((1H-1,2,4-triazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide (Compound No. 201)



To a solution of 1,2,4-triazole (85 mg, 1.2 mmol) in DMF (3 mL) at 0°C under argon was added NaH (60% in mineral oil, 48 mg, 1.2 mmol). After 1.5 hours Compound No. 193 (50 mg, 0.12 mmol) was added and the reaction was stirred at room temperature for 18 hours. The reaction cooled to 0°C, quenched with saturated NaHCO_3 solution (3 mL), diluted with Et_2O (40 mL), washed with half-saturated NaCl solution (3 x 8 mL), dried (MgSO_4) and concentrated. The residue was purified by chromatography (50% then 100% EtOAc/hexanes then 10% then 20% MeOH/EtOAc) to afford Compound No. 201 (37 mg, 79%) as a white solid. ^1H NMR (CDCl_3): δ 8.92 (m, 1H), 8.72 (m, 1H), 8.02 (2H), 7.92 (s, 1H), 7.40 (m, 1H), 6.35 (m, 1H), 4.48 (m, 1H), 4.25 (m, 1H), 3.98 (m, 1H), 2.10 (m, 1H), 1.95-1.45 (6H), 1.35-1.05 (4H), 0.98 (s, 3H), 0.90 (d, 3H), 0.85 (s, 3H), 0.80 (s, 3H). ES-MS m/z 396 ($[\text{M}+1]^+$).

EXAMPLE 110


Synthesis of N-((2R,3S,4S,4aS)-4-((1H-benzo[d]imidazol-1-yl)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide (Compound No. 202)



To a solution of benzimidazole (145 mg, 1.2 mmol) in DMF (3 mL) at 0°C under argon was added NaH (60% in mineral oil, 48 mg, 1.2 mmol). After 1.5 hours Compound No. 193 (50 mg, 0.12 mmol) was added and the reaction was stirred at room temperature for 18 hours. The reaction cooled to 0°C, quenched with saturated NaHCO₃ solution (3 mL), diluted with Et₂O (40 mL), washed with half-saturated NaCl solution (3 x 8 mL), dried (MgSO₄) and concentrated. The residue was purified by chromatography (80% then 100% EtOAc/hexanes then 20% MeOH/EtOAc) to afford Compound No. 201 (34 mg, 64%) as a white solid. ¹H NMR (CDCl₃): δ 8.82 (m, 1H), 8.59 (m, 1H), 7.95 (m, 1H), 7.87 (s, 1H), 7.75 (m, 1H), 7.42 (m, 1H), 7.25 (3H), 6.65 (m, 1H), 4.50 (m, 1H), 4.25 (m, 1H), 3.95 (m, 1H), 2.15 (m, 1H), 1.90 (2H), 1.75 (2H), 1.60 (m, 1H), 1.50 (2H), 1.1 (3H), 1.05 (s, 3H), 0.95 (9H). ES-MS *m/z* 445 ([M+1]⁺).

EXAMPLE 111

Synthesis of 4-methyl-N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-((4-(trifluoromethyl)benzamido)methyl)-decahydronaphthalen-2-yl)piperazine-1-carboxamide (Compound No. 208)



A. To a solution of Compound No. 190 (1.97 g, 4.12 mmol) and Et₃N (2.9 mL, 20.6 mmol) in CH₂Cl₂ (20 mL) at 0°C under argon was added 4-methyl-1-piperazinecarboxyl chloride (0.93 g, 4.54 mmol). After 10 minutes the cold bath was

removed and the reaction was stirred at room temperature for 18 hours. The reaction was cooled to 0°C and quenched with saturated NaHCO₃ solution (3 mL) and stirred at room temperature for 30 minutes, diluted with CH₂Cl₂ (120 mL), washed with half-saturate NaCl solution (2 x 20 mL), dried (MgSO₄) and concentrated to afford Compound No. 203 (2.59 g, 100%) as a white foam.

B. A mixture of crude Compound No. 203 (353 mg, 0.58 mmol) and tetrabutylammonium fluoride (TBAF) (1.0 M in THF, 1.8 mL, 1.8 mmol) in THF (10 mL) under argon was heated at reflux for 2 hours then was cooled and concentrated. The residue was purified by chromatography on silica gel (10% MeOH/EtOAc with 1% NH₄OH then 20% MeOH/EtOAc with 2% NH₄OH) to afford Compound No. 204 (215 mg, 100%) as a white solid.

C. To a solution of Compound No. 204 (211 mg, 0.58 mmol) in pyridine (3 mL) and CH₂Cl₂ (3 mL) at 0°C under argon was added MsCl (137 µL, 1.73 mmol). After 1 hour saturated NaHCO₃ solution (3 mL) was added and the mixture stirred at room temperature for 30 minutes. The mixture was diluted with EtOAc (75 mL), washed with half-saturated NaCl solution (3 x 15 mL), dried (MgSO₄) and concentrated to afford Compound No. 205 (245 mg, 96%) as an orange foam.

D. A mixture of Compound No. 205 (190 mg, 0.43 mmol) and NaN₃ (83 mg, 1.3 mmol) in DMF (3 mL) at 50°C under argon was stirred for 7 hours then was continued at room temperature. The mixture was diluted with Et₂O (75 mL), washed with half-saturated NaCl solution (3 x 8 mL), dried (MgSO₄) and concentrated. The residue was purified by chromatography on silica gel (10% MeOH/EtOAc with 1% NH₄OH) to afford Compound No. 206 (141 mg, 84%) as a colourless film.


E. A solution of Compound No. 206 (141 mg, 0.36 mmol), triphenylphosphine (956 mg, 3.6 mmol) and water (65 µL, 3.6 mmol) in THF (10 mL) was stirred at room temperature for 4 days. The mixture was concentrated and the residue was purified by chromatography on silica gel (40% MeOH/EtOAc with 4% NH₄OH) to afford Compound No. 207 (126 mg, 96%) as a white solid.

F. A mixture of Compound No. 207 (20 mg, 0.055 mmol), 4-trifluoromethyl benzoic acid (32 mg, 0.16 mmol) and DCC (29 mg, 0.14 mmol) in CH₂Cl₂ (4 mL) under argon was stirred at room temperature for 22 hours. The mixture was purified by chromatography on silica gel (5% MeOH/EtOAc with 0.5% NH₄OH then 10% MeOH/EtOAc with 1% NH₄OH) to afford Compound No. 208 (26 mg, 90%) as a white solid. ¹H NMR (CDCl₃): δ 7.85 (m, 2H), 7.70 (m, 2H), 6.15 (m, 1H), 4.58 (m, 1H), 4.20

(m, 1H), 3.79 (m, 1H), 3.39 (4H), 3.20 (m, 1H), 2.39 (4H), 2.30 (s, 3H), 2.05 (m, 1H), 1.95 (m, 1H), 1.8-1.4 (6H), 1.3-0.8 (15H). ES-MS m/z 537 ($[M+1]^+$).

EXAMPLE 112


Synthesis of N-((2R,3S,4S,4aS)-4-((4-fluorobenzamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide (Compound No. 209)



A mixture of Compound No. 207 (25 mg, 0.069 mmol), 4-fluorobenzoic acid (30 mg, 0.21 mmol) and DCC (36 mg, 0.17 mmol) in CH_2Cl_2 (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (5% MeOH/EtOAc with 0.5% NH_4OH then 10% MeOH/EtOAc with 1% NH_4OH) to afford Compound No. 209 (18 mg, 55%) as a white solid. ^1H NMR (CDCl_3): δ 7.75 (m, 2H), 7.10 (m, 2H), 6.05 (m, 1H), 4.59 (m, 1H), 4.20 (m, 1H), 3.78 (m, 1H), 3.39 (4H), 3.19 (m, 1H), 2.39 (4H), 2.30 (s, 3H), 2.05 (m, 1H), 1.95 (m, 1H), 1.8-1.4 (6H), 1.3-0.8 (15H). ES-MS m/z 487 ($[M+1]^+$).

EXAMPLE 113


Synthesis of N-((2R,3S,4S,4aS)-4-(isonicotinamidomethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide (Compound No. 210)



A mixture of Compound No. 207 (25 mg, 0.069 mmol), isonicotinic acid (30 mg, 0.21 mmol) and DCC (36 mg, 0.17 mmol) in CH_2Cl_2 (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (20% MeOH/EtOAc with 2% NH_4OH) to afford Compound No. 210 (22 mg, 69%) as a white solid. ^1H NMR (CDCl_3): δ 8.77 (m, 2H), 7.59 (m, 2H), 6.35 (m, 1H), 4.62 (m, 1H), 4.18 (m, 1H), 3.78 (m, 1H), 3.40 (4H), 3.20 (m, 1H), 2.40 (4H), 2.30 (s, 3H), 2.0 (3H), 1.8-1.4 (5H), 1.3-0.8 (15H). ES-MS m/z 470 ($[\text{M}+1]^+$).

EXAMPLE 114


Synthesis of 4-methyl-N-((2R,3S,4S,4aS)-3,4a,8,8-tetramethyl-4-(nicotinamidomethyl)decahydronaphthalen-2-yl)piperazine-1-carboxamide (Compound No. 211)



A mixture of Compound No. 207 (25 mg, 0.069 mmol), nicotinic acid (26 mg, 0.21 mmol) and DCC (36 mg, 0.17 mmol) in CH_2Cl_2 (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (10% MeOH/EtOAc with 1% NH_4OH then 20% MeOH/EtOAc with 2% NH_4OH) to afford Compound No. 211 (25 mg, 78%) as a white solid. ^1H NMR (CDCl_3): δ 8.99 (m, 1H), 8.72 (m, 1H), 8.15 (m, 1H), 7.40 (m, 1H), 6.40 (m, 1H), 4.72 (m, 1H), 4.18 (m, 1H), 3.78 (m, 1H), 3.42 (4H), 3.20 (m, 1H), 2.42 (4H), 2.32 (s, 3H), 2.0 (2H), 1.8-1.4 (4H), 1.3-0.8 (17H). ES-MS m/z 470 ($[\text{M}+1]^+$).

EXAMPLE 115


Synthesis of N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide (Compound No. 212)



A mixture of Compound No. 207 (25 mg, 0.069 mmol), 2-furoic acid (24 mg, 0.21 mmol) and DCC (36 mg, 0.17 mmol) in CH₂Cl₂ (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (5% MeOH/EtOAc with 0.5% NH₄OH then 10% MeOH/EtOAc with 1% NH₄OH) to afford Compound No. 212 (19 mg, 61%) as a white solid. ¹H NMR (CDCl₃): δ 7.42 (m, 1H), 7.18 (m, 1H), 6.52 (m, 1H), 6.30 (m, 1H), 4.59 (m, 1H), 4.19 (m, 1H), 3.78 (m, 1H), 3.39 (4H), 3.15 (m, 1H), 2.40 (4H), 2.28 (s, 3H), 2.05 (m, 1H), 1.95 (m, 1H), 1.8-1.4 (4H), 1.3-0.8 (17H). ES-MS *m/z* 459 ([M+1]⁺).

EXAMPLE 116

Synthesis of 4-methyl-N-((2R,3S,4S,4aR)-3,4a,8,8-tetramethyl-4-(2-(4-(trifluoromethyl)benzamido)ethyl)-decahydronaphthalen-2-yl)piperazine-1-carboxamide (Compound No. 215)



A. A mixture of Compound No. 205 (744 mg, 1.68 mmol) and KCN (342 mg, 5.04 mmol) in DMSO (5 mL) at 60°C under argon was stirred for 16 hours. The mixture was cooled to room temperature, diluted with Et₂O (50 mL) and washed with water (20 mL). The aqueous wash was extracted with Et₂O (50 mL) and the combined Et₂O layers was washed with brine (10 mL), dried (MgSO₄) and concentrated. The residue was purified by chromatography (10% MeOH/EtOAc with 1% NH₄OH) to afford Compound No. 213 (327 mg, 52%) as a white foam.


B. To a solution of Compound No. 213 (98 mg, 0.26 mmol) in THF at 0°C under argon was added LiAlH₄ (2.0 M in THF, 0.65 mL, 1.3 mmol) over 1 minute. The reaction was heated at reflux for 2 hours then stirred at room temperature for 16 hours. The mixture was cooled to 0°C then NaBO₃·4H₂O was added and the mixture stirred at room temperature for 1 hour. The mixture was filtered through Celite eluting with EtOAc (100 mL) and concentrated. The residue was purified by chromatography on silica gel (5% NH₄OH/MeOH) to afford Compound No. 214 (75 mg, 76%) as a white film.

C. A mixture of Compound No. 214 (42 mg, 0.11 mmol), 4-trifluoromethyl benzoic acid (65 mg, 0.33 mmol) and DCC (57 mg, 0.28 mmol) in CH₂Cl₂ (10 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by

chromatography on silica gel (5% MeOH/EtOAc with 0.5% NH₄OH then 10% MeOH/EtOAc with 1% NH₄OH) to afford Compound No. 215 (49 mg, 80%) as a white solid. ¹H NMR (CDCl₃): δ 7.88 (m, 2H), 7.70 (m, 2H), 6.55 (m, 1H), 4.72 (m, 1H), 4.05 (m, 1H), 3.6-3.3 (6H), 2.39 (4H), 2.28 (s, 3H), 2.0-0.7 (25H). ES-MS *m/z* 551 ([M+1]⁺).

EXAMPLE 117


Synthesis of N-((2R,3S,4S,4aR)-4-(2-(4-fluorobenzamido)ethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide (Compound No. 216)



A mixture of Compound No. 214 (25 mg, 0.066 mmol), 4-fluorobenzoic acid (28 mg, 0.20 mmol) and DCC (34 mg, 0.17 mmol) in CH₂Cl₂ (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (5% MeOH/EtOAc with 0.5% NH₄OH then 10% MeOH/EtOAc with 1% NH₄OH) to afford Compound No. 216 (28 mg, 85%) as a white solid. ¹H NMR (CDCl₃): δ 7.78 (m, 2H), 7.10 (m, 2H), 6.25 (m, 1H), 4.72 (m, 1H), 4.05 (m, 1H), 3.50 (m, 1H), 3.40 (4H), 3.35 (m, 1H), 2.40 (4H), 2.0-0.7 (25H). ES-MS *m/z* 501 ([M+1]⁺).

EXAMPLE 118


Synthesis of N-((2R,3S,4S,4aR)-4-(2-(furan-2-carboxamido)ethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide (Compound No. 217)



A mixture of Compound No. 214 (25 mg, 0.066 mmol), 2-furoic acid (23 mg, 0.20 mmol) and DCC (34 mg, 0.17 mmol) in CH_2Cl_2 (4 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (5% MeOH/EtOAc with 0.5% NH_4OH then 10% MeOH/EtOAc with 1% NH_4OH) to afford Compound No. 217 (26 mg, 84%) as a white solid. ^1H NMR (CDCl_3): δ 7.45 (m, 1H), 7.05 (m, 1H), 6.50 (m, 1H), 6.40 (m, 1H), 4.70 (m, 1H), 4.05 (m, 1H), 3.45 (5H), 3.33 (m, 1H), 2.40 (4H), 2.30 (s, 3H), 1.9-1.2 (10H), 1.12 (m, 1H), 1.00 (d, 3H), 0.9-0.6 (11H). ES-MS m/z 473 ($[\text{M}+1]^+$).

EXAMPLE 119


Synthesis of 4-methyl-N-((2R,3S,4S,4aR)-3,4a,8,8-tetramethyl-4-(2-(nicotinamido)ethyl)decahydronaphthalen-2-yl)piperazine-1-carboxamide (Compound No. 218)



A mixture of Compound No. 214 (25 mg, 0.066 mmol), nicotinic acid (25 mg, 0.20 mmol) and DCC (34 mg, 0.17 mmol) in CH_2Cl_2 (3 mL) under argon was stirred at room temperature for 18 hours. The mixture was purified by chromatography on silica gel (10% MeOH/EtOAc with 1% NH_4OH then 20% MeOH/EtOAc with 2% NH_4OH) to afford Compound No. 218 (27 mg, 84%) as a white solid. ^1H NMR (CDCl_3): δ 8.98 (m, 1H), 8.75 (m, 1H), 8.10 (m, 1H), 7.40 (m, 1H), 6.48 (m, 1H), 4.70 (m, 1H), 4.05 (m, 1H), 3.55 (m, 1H), 3.40 (4H), 3.20 (m, 1H), 2.40 (4H), 2.30 (s, 3H), 2.00 (m, 1H), 1.77 (4H), 1.6-1.2 (5H), 1.2-0.9 (4H), 0.9 (11H). ES-MS m/z 484 ($[\text{M}+1]^+$).

EXAMPLE 120

Synthesis of 3-[[[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy]-5-methoxyphenol (Compound No. 220)



A. A solution of 1-bromo-3,5-dimethoxybenzene (2.17 g, 10.0 mmol) in anhydrous THF (50 mL) was cooled to -78 °C under N₂. To the mixture, a 1.6 M solution of n-BuLi in hexanes (12.5 mL, 20.0 mmol) was added, and the mixture was stirred at -78 °C for 30 min. A solution of iodine (5.1 g, 40 mmol) in anhydrous THF (50 mL) was added dropwise to the reaction mixture and the mixture was stirred for 1 h while slowly warming to -30 °C. The reaction was quenched with water and the organic layer was separated and concentrated to dryness. The residue was purified by flash column chromatography on silica gel (Hexanes/EtOAc, 4:1) to yield 1-iodo-3,5-dimethoxybenzene (Compound No. 31) (1.45 g, 55% yield) as a pale brown solid.

B. (1S,2R,4aS,8aS)-1-(hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 36) (prepared according to Kuchkova *et al.*, *Synthesis*, 1045-1048, 1997) (0.59 g, 2.4 mmol), Cu(I)I (0.04 g, 0.21 mmol) and 3,4,7,8-tetramethyl-[1,10]phenanthroline (0.10 g, 0.50 mmol) were added to an oven


dried tube, which was sealed with a rubber septum, and flushed with a stream of nitrogen for 15 min. Anhydrous toluene (10 mL), anhydrous diglyme (2.0 mL) and 1-iodo-3,5-dimethoxybenzene (Compound No. 31) (0.71 g, 2.7 mmol) were added via a syringe with vigorous stirring under a stream of nitrogen at room temperature. The rubber septum was replaced with the teflon cap and the sealed tube was heated at 130 °C in an oil bath for 18 h. The reaction mixture was allowed to cool to room temperature, then quenched with water and extracted with EtOAc. The organic layer was separated, dried with Na₂SO₄ and concentrated to dryness under reduced pressure. The residue was purified by flash column chromatography on silica gel (Hexanes/EtOAc, 4:1) to yield (1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 34) (0.65 g, 70% yield) as a pale yellow oil.

C. To a stirred solution of (1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 34) (0.65 g, 1.7 mmol) in CH₂Cl₂ (20 mL), cooled in a dry ice-acetone bath to -78 °C, a solution of SnCl₄ (0.8 mL, 6.9 mmol) in CH₂Cl₂ (10 mL) was added dropwise under N₂. As soon as the addition ended, the dry ice-acetone bath was replaced with an ice bath. The mixture was warmed to room temperature and stirred for 3 h. The reaction mixture was quenched with water and the organic layer was separated and concentrated to dryness. The residue was purified by flash column chromatography on silica gel (Hexanes/Toluene, 1:2) to yield (4aS,5S,8aS)-5-(3,5-dimethoxyphenoxyethyl)-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene (Compound 219) (0.15 g, 24% yield) as a white solid.

D. A mixture of (4aS,5S,8aS)-5-(3,5-dimethoxyphenoxyethyl)-1,1,4a,6-tetramethyl-1,2,3,4,4a,5,8,8a-octahydronaphthalene (5) (0.15 g, 0.45 mmol), Na₂S (0.22 g, 2.7 mmol) and NMP (5.0 mL) were sealed in a microwave vessel and heated at 180 °C with stirring for 1 h. The reaction mixture was cooled to room temperature, quenched with water and extracted with EtOAc. The organic layer was separated and concentrated to dryness. The residue was purified by flash column chromatography on silica gel (Hexanes/EtOAc, 6:1) to yield 3-([(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]methoxy)-5-methoxyphenol (Compound No. 220) (0.058 g, 37% yield) as a pale brown oil. ¹H NMR(CDCl₃): δ 6.05 (s, 1H), 6.00 (s, 2H), 5.50 (s, 1H), 4.60 (m, 1H), 4.05 (m, 1H), 3.85 (m, 1H), 3.75 (s, 3H), 2.15 (m, 1H), 2.00–1.80 (m, 3H), 1.68 (s, 3H), 1.61–0.95 (m, 6H), 0.88 (s, 3H), 0.84 (s, 6H); MS m/z 345 (C₂₂H₃₂O₃ + 1).

EXAMPLE 121

Synthesis of (2R,4aS,8aS)-1-[[[(3-hydroxy-5-methoxyphenyl)amino]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 223)



A. (1S,2R,4aS,8aS)-1-(hydroxymethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 26) (prepared according to Kuchkova *et al.*, *Synthesis*, 1045 - 1048, 1997) (2.0 g, 8.3 mmol) was added to a mixture of pyridinium chlorochromate (3.0 g, 9.5 mmol) in anhydrous CH₂Cl₂ (40 mL). The mixture was stirred at room temperature for 4 h under N₂. The reaction was filtered through a plug of silica and the filtrate concentrated to dryness. The crude product was purified by flash chromatography on silica gel (Hexanes/EtOAc, 6:1) to yield (2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalene-1-carbaldehyde (Compound No. 221) (0.73 g, 37% yield) as a pale green solid.


B. A mixture of (2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalene-1-carbaldehyde (Compound No. 221) (0.25 g, 1.1 mmol), 3,5-dimethoxyaniline (0.25 g, 1.6 mmol) and sodium triacetoxyborohydride (0.34 g, 1.6 mmol) in dichloroethane (10 mL) was stirred at room temperature for 18 h under N₂. The reaction mixture was concentrated to dryness and the residue was purified by flash chromatography on silica gel (Hexanes/EtOAc, 2:1) to yield (2R,4aS,8aS)-1-

{[(3,5-dimethoxyphenyl)amino]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 222) (0.10 g, 26 %) as a colorless oil.

C. A mixture of (2R,4aS,8aS)-1-{[(3,5-dimethoxyphenyl)amino]-methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 223) (0.10 g, 0.27 mmol), EtSNa (0.44 g, 5.3 mmol) and DMF (3.0 mL) were sealed in a microwave vessel and heated at 160 °C with stirring for 2 h. The reaction mixture was cooled to room temperature, quenched with water and extracted with EtOAc. The organic layer was separated and then concentrated to dryness. The crude product was purified by flash chromatography on silica gel (Hexanes/EtOAc, 2:1) to yield (2R,4aS,8aS)-1-{[(3-hydroxy-5-methoxyphenyl)amino]methyl}-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol (Compound No. 223) (0.080 g, 83 % yield) as a pale brown solid. ¹H NMR (CDCl₃): δ 5.84 (s, 2H), 5.78 (s, 1H), 3.75 (s, 3H), 3.30-3.12 (m, 3H), 2.10-1.80 (m, 3H), 1.70-0.90 (m, 12H), 0.88 (s, 3H), 0.83 (s, 3H), 0.78 (s, 3H); MS m/z 362 (C₂₂H₃₅NO₃ + 1).

EXAMPLE 122

Synthesis of 2-[(1S,2R,3R,4aS,8aR)-3-bromo-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide (Compound No. 225) and 2-[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide (Compound No. 226)



A. To a stirred solution of N-(3,5-dimethoxyphenyl)-N-methylamine (0.43 g, 2.6 mmol) in anhydrous toluene (10 mL) at room temperature, trimethylaluminum (2 M

in hexanes, 1.9 mL, 3.8 mmol) was added dropwise. The mixture was stirred at room temperature for 55 minutes. A solution of (3aR,5aS,9aS,9bR)-3a,6,6,9a-tetramethyl-dodecahydronaphtho[2,1-b]furan-2-one (Compound No. 44) (0.71 g, 2.8 mmol) in anhydrous toluene (3.0 mL) was added dropwise and the mixture was heated at 80 °C for 17 h. The reaction was cooled to 0 °C and quenched with 1 M HCl, diluted with water and extracted with EtOAc. The organic phase was separated and concentrated to dryness. The crude product was purified by flash chromatography on silica gel (Hexanes/EtOAc, 1:1) to yield 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dimethoxyphenyl)-N-methylacetamide (Compound No. 224) (0.98 g, 91% yield) as a white semi-solid.

B. A solution of 2-[(1R,2R,4aS,8aS)-2-hydroxy-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dimethoxyphenyl)-N-methylacetamide (Compound No. 224) (0.16 g, 0.38 mmol) in anhydrous CH₂Cl₂ (3.0 mL) was prepared at -28 °C under N₂. To the mixture, BBr₃ (1 M in CH₂Cl₂, 2.0 mL, 2.0 mmol) was added dropwise. The mixture was stirred at -28 °C to -20 °C for 8 h and then was allowed to slowly warm to room temperature overnight. The solution was cooled to 0 °C and quenched with anhydrous MeOH. The solution was evaporated and the residue co-evaporated with anhydrous MeOH. The residue was dissolved in CH₂Cl₂ and was shaken with 1M HCl. The pH was adjusted to 8 using NaHCO₃ and the organic phase was separated. The aqueous phase was extracted with CH₂Cl₂ and the organic extracts were combined and concentrated to dryness. The residue was purified by flash chromatography on silica gel (Hexanes/EtOAc, 3:2) to yield 2-[(1S,2R,3R,4aS,8aR)-3-bromo-2,5,5,8a-tetramethyl-decahydro-naphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide (Compound No. 225) (0.12 g, 69% yield) as an off-white solid. ¹H NMR (CD₃OD): δ 6.26 (s, 1H), 6.14 (d, 2H), 4.60 (d, 1H), 3.16 (s, 3H), 2.21 – 2.00 (m, 3H), 1.93 – 1.83 (m, 2H), 1.73 – 1.67 (d, 1H), 1.63 – 1.52 (m, 2H), 1.51 – 1.37 (m, 3H), 1.30 – 1.18 (m, 1H), 1.11 – 0.99 (m, 1H), 0.87 (d, 3H), 0.84 (s, 3H), 0.81 (s, 3H), 0.70 (s, 3H). MS *m/z* 450, 452 (C₂₃H₃₄BrNO₃ - 1).

C. A solution of 2-[(1S,2R,3R,4aS,8aR)-3-bromo-2,5,5,8a-tetramethyl-decahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide (Compound No. 225) (0.054 g, 0.119 mmol) in anhydrous toluene (2.0 mL) and DBU (0.10 mL, 0.66 mmol) were heated at 118 °C for 23 h. The mixture was cooled in ice, diluted with CH₂Cl₂ and washed with 1 M HCl. The acidic aqueous layer was extracted with CH₂Cl₂. The organic phase was washed with diluted brine and this aqueous washing was back extracted with CH₂Cl₂. The combined organic extracts were concentrated to

dryness. The crude solid was triturated with CH₂Cl₂ and collected by filtration to yield 2-[(1S,4aS,8aS)-2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8,8a-octahydronaphthalen-1-yl]-N-(3,5-dihydroxyphenyl)-N-methylacetamide (Compound No. 226) (0.024 g, 56% yield) as an off-white solid. ¹H NMR (CD₃OD): δ 6.27 (dd, 1H), 6.15 (d, 2H), 5.35 (br.s, 1H), 3.18 (s, 3H), 2.63 – 2.56 (m, 1H), 2.20 – 2.14 (m, 1H), 2.08 – 1.93 (m, 2H), 1.87 – 1.75 (m, 1H), 1.71 – 1.63 (m, 1H), 1.58 – 1.38 (m, 3H), 1.29 – 1.15 (m, 2H), 1.04 (ddd, 1H), 0.87 (s, 6H), 0.86 (s, 3H), 0.61 (s, 3H). MS *m/z* 372 (C₂₃H₃₃NO₄ + 1).

EXAMPLE 123

His-hSHIP1 Activity of Representative Compounds

Test compounds are dissolved in 95% ethanol to form stock solutions. Before screening, the stock solutions are diluted with Phosphatase Assay Buffer (20 mM Tris-HCL, 10 mM MgCl₂ pH 7.5, 0.02% Tween 20) to form working assay solutions that contain 10% ethanol. The assay is carried out on 96-well microtiter plates using a modified procedure of that reported by Ong *et al.*, *Blood* **110**, 1942-1949, 2007 and Yang *et al.*, *Org Lett* **7**, 1073-1076, 2005, both of which references are incorporated herein by reference in their entirety.

Data were generated by one of two variant methods. Test compounds are screened by one of two approaches described as follows. In the first approach, each reaction contains 5 μL of His-hSHIP1 enzyme (5-20 ng), 10 μL of the substrate, 1,3,4,5-inositol tetrakisphosphate (IP₄; 50 μM final), 5 μL of Phosphatase Assay Buffer, and 5L of test compound at various concentrations in 10% ethanol (0 – 300 μM final). Control blanks are also prepared by replacing His-hSHIP1 enzyme, IP₄, or test compounds with Phosphatase Assay Buffer. After adding the reaction components in a 96-well microtiter plate on ice, the reaction is mixed by briefly shaking the plate vigorously. The reaction is then incubated at 37 °C for 15 min with gentle shaking followed by addition of 100 μL of Biomol Green Reagent (BIOMOL, PA, USA) to terminate the reaction. The free phosphate released from IP₄ by His-hSHIP1 binds to the Biomol Green Reagent, which turns the dye to green color. After incubating the mixture for 20 min at room temperature for color development, the absorbance is read with SpectraMax Plus 96-well plate reader (Molecular Devices, Sunnyvale, California, USA) at a wavelength of 650 nm.

In the second approach, a master mix is prepared that contains 50 μL of His-hSHIP1 enzyme (0.4-1.6 ng/μL final), 25 μL of the substrate, IP₄ (25 or 250 μM final),

and 50 μ L of test compound in 3.3% ethanol (100 μ L final). Control blanks are also prepared by replacing His-hSHIP1 enzyme, IP4, or test compounds with Phosphate Assay Buffer. A reference compound that has been shown to activate SHIP1 is also included. Each reaction component is preincubated at 37°C for 30 min before adding to a 96-well microtiter plate. The master mix is then incubated at 37°C. At time 0, 20, 25 and 30 min, 25 μ L of the master mix is removed and transferred to a new 96-well microtiter plate, to which 100 μ L of Biomol Green Reagent is added to stop the reaction. After incubating the mixture for 20 min at room temperature for color development, the absorbance is read with SpectraMax Plus 96-well plate reader at a wavelength of 650 nm. Phosphate released is plotted against time to calculate the initial velocities (*i.e.*, slope of the graph) at each IP4 concentration. The initial velocities are baseline corrected and the ratio of initial velocities (IP4₂₅/IP4₂₅₀) is calculated and used to rate the test compounds.

According to the above assay(s), the representative compounds listed in Table 2 below were found to activate His-hSHIP1 enzyme at concentrations $\leq 300 \mu$ M. Scoring of the compound is expressed as follows:

Scoring	Approach 1 (% increase relative to background)	Approach 2 (Initial velocity Ratio)
+++	$x \geq 65$	higher than reference or reference - 15%
++	$50 \leq x < 65$	reference - 16% to reference - 30%
+	$0 < x < 50$	reference - 31% or lower

Data generated by the second method is distinguished by an asterisk.

Table 2

Cpd. No.	Scoring
4	++
6	+
8	+
9	+
11	+++
17	++
22	+
23	+

Cpd. No.	Scoring
27	++
29	+++
30	+
38	+
42	+
43	+
45	++
46	++
47	++
48	+++
50	++
51	++
52	+++
53	++
54	++
55	+
56	+
57	+++
58	+++
62	+++
65	+++
49*	+++
68*	+++
20*	+++
136*	++
175*	++
176*	++
149*	++
137*	++
137*	++
138*	++
139*	++

Cpd. No.	Scoring
157*	++
181*	+++
141*	+
142*	++
173*	+++
118*	+++
178*	+++
119*	++
179*	+

EXAMPLE 124

Activity of Representative Compounds on Akt Phosphorylation in Lymphocytes

Phosphorylation of AKT has been shown to be modulated by SHIP1 (Helgason *et al.*, *J Exp Med* **191**, 781-794, 2000). Jurkat (PTEN-/SHIP1-) or Molt-4 (PTEN-/SHIP1+) cells are starved in serum free RPMI for overnight. In a 15 mL conical tube, 2-3 million serum starved cells (1 million cells per mL) are treated with various concentrations of test compound (0.1, 1, or 10 μ M final in 0.1% DMSO) for 30 min at 37°C followed by stimulation with 100 ng/mL of IGF-1 for 1 hour at 37°C. After stimulation, cells are washed once with ice-cold DPBS and lysed with Lysis Buffer (20 mM Tris-HCl, pH 7.5, 140 mM NaCl, 1% NP-40, Complete Mini Protease Inhibitor Cocktail, 10 mM NaF, 1 mM Na₃VO₄, 1 mM β -glycerolphosphate) on ice for 30 min with vortexing every 10 min. Samples are then centrifuged at 13,000 rpm for 20 min, and supernatants are collected as total cell lysate samples. Protein concentration is determined using bicinchonic acid assay, and about 15 μ g of total protein from each sample is loaded and separated on a 4-12% Tris-Glycine gel. After SDS-PAGE, proteins are transferred from the gel to a nitrocellulose membrane. The membrane is blocked in 5% BSA in PBS containing 0.1% Tween-20 (PBS-T) for 1 hour at room temperature before probing with primary antibodies for overnight at 4°C. The following antibodies are used: mouse anti-SHIP1 (1:500 dilution; Santa Cruz, CA, USA), rabbit anti-phospho-Akt(Ser473) (1:1000 dilution; Cell Signaling Technologies, MA, USA), rabbit anti-Akt (1:1000; Cell Signaling Technologies, MA, USA), and rabbit anti-actin (1:2000; Cell Signaling Technologies, MA, USA). The membrane is then incubated with goat anti-rabbit or anti-mouse secondary antibodies (1:3000) for 1 hour at room

temperature. Target proteins on the membrane are detected with ECL solution and exposed on a film.

According to the above assay, the representative compounds listed in Table 3 below were found to inhibit Akt phosphorylation at ≤ 10 μ M in Molt-4 (SHIP1 +), but not Jurkat (SHIP1 -) lymphocytes. Scoring in Table 3 is expressed as follows: + (inhibits Akt phosphorylation at 10 μ M); - (no effect on Akt phosphorylation at 10 μ M).

Table 3

Cpd. No.	Molt-4 (SHIP +)	Jurkat (SHIP1 -)
22	+	-
57	+	-
22	+	-
57	-	-
68	-	-
20	-	-

EXAMPLE 125

Activity of Representative Compounds on Passive Cutaneous Anaphylaxis in Mice

The activity of representative compounds on passive cutaneous anaphylaxis in mice was evaluated according to the procedures disclosed by Ovary, *J Immunol* **81**, 355-357, 1958 and Halpern *et al.*, *Br J Pharmacol Chemother* **20**, 389-398, 1963, both of which are incorporated herein by reference in their entirety.

To induce a passive cutaneous anaphylaxis, mice undergo intradermal ear inoculation on their right ear with 25 ng in 20 μ L of anti-DNP-IgE. The left ears are untreated and serve as negative controls. Twenty-four hours after inoculation, all mice are administered test compound by oral gavage (PO). Sixty minutes after oral administration, mice are given a tail vein injection of 2% Evan's Blue (0.2 μ m filtered, in 200 μ L saline) followed by a second tail IV injection of 100 μ g DNP-HSA (in 200 μ L). Sixty minutes following the DNP-HAS injection, mice are euthanized using CO₂ inhalation. Subsequently, ear biopsies are performed by taking four millimetre punches from both ears, which then undergo Evan's Blue extraction using formamide incubation in 96 well plates. Eighty μ L of eluents are transferred to flat-bottom 96-well plates and absorbance read using SpectraMax M5 spectrophotometer (Molecular Devices, Sunnyvale, California, USA) at 620 nm. Background readings from all

samples are taken at 740 nm and subtracted from the 620 nm readings. Data are reported as OD.

According to the above assay, the representative compounds listed in Table 4 below were found to inhibit allergen-induced passive cutaneous anaphylaxis at doses less than 20 mg/kg. Scoring in Table 4 is expressed as follows: + (1-30% inhibition); ++ (31-50% inhibition); +++ (>50% inhibition).

Table 4

Cpd. No.	% Inhibition
38	+++
57	+++
58	+++

EXAMPLE 126

Activity of Representative Compounds on Carrageenan Paw Edema in Mice

The activity of representative compounds on carrageenan paw edema in mice was evaluated according to the procedures disclosed by Winter *et al.*, *Proc Soc Exp Biol Med* **111**, 544-547, 1962 which is incorporated herein by reference in its entirety. To induce edema in the paw, test compounds were administered orally one hour before intraplantar injection of the right hind paw with carrageenan (50 μ L of 1% suspension). Hind paw edema, as a measure of inflammation, was recorded using a plethysmometer (Ugo Basile, Italy) 4 hours after λ -carrageenan administration. When tested in this assay, representative Compound No. 48 was found to inhibit allergen-induced carrageenan-induced paw edema by 21% at 30 mg/kg.

* * * * *


All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification are incorporated herein by reference in their entireties.

Although the foregoing invention has been described in some detail to facilitate understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims. Accordingly, the described embodiments are to be considered as illustrative and not restrictive, and the invention is not to be limited to the details given herein, but may be modified within the scope and equivalents of the appended claims.


AMENDED CLAIMS

received by the International Bureau on 08 August 2014 (08.08.2014)


1. A compound of formula (I):



wherein:



is selected from:



n is 1, 2, 3, 4, 5, or 6;

R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, $-R^{9a}-C(O)-R^{9b}-$, $-R^{9a}-S(O)_t-R^{9b}-$ (where t is 0, 1 or 2),

$-R^{9a}-O-R^{9b}-$, $-R^{9a}-C(O)N(R^{11a})-R^{9b}-$, $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$ or $-R^{9a}-N(R^{11a})-R^{9b}-$;

R^2 and R^3 are each independently selected from hydrogen, alkyl or $-R^9-OR^{11}$, provided that at least one of R^2 and R^3 is $-R^9-OR^{11}$ when R^6 is hydrogen;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$,

or R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or $-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);

each R^6 and R^8 is independently selected from hydrogen, alkyl, halo or haloalkyl;

R^7 is hydrogen, alkyl, halo or haloalkyl;

each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;

each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;

each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and

R^{14} is a straight or branched alkylene chain;

or a stereoisomer thereof or a pharmaceutically acceptable salt thereof;

provided that compounds of formula (I) do not include the following compounds:

(1S,2S,4aS,8aR)-decahydro-1-[2-(3-methoxyphenyl)ethyl]-5,5,8a-trimethyl-2-naphthalenecarboxylic acid;

(1S,2S,4aS,8aR)-decahydro-1-[2-(3-methoxyphenyl)ethyl]-5,5,8a-trimethyl-2-naphthalenemethanol;

α -(3,5-dimethoxyphenyl)decahydro-2-hydroxy-2,5,5,8a-tetramethyl-1-naphthalenemethanol;


(4aS,8aS)-3,4,4a,5,6,7,8,8a-octahydro- α -(3-methoxy-5-methylphenyl)-2,5,5,8a-tetramethyl-1-naphthalenemethanol;

(1S,2R,4aS,8aS)-decahydro-2-hydroxy- α -(3-methoxy-5-methylphenyl)-2,5,5,8a-tetramethyl-1-naphthalenemethanol;

(1R,2R,4aS,8aS)-decahydro-1-[(3-methoxy-5-methylphenyl)methyl]-2,5,5,8a-


- tetramethyl-2-naphthalenol;
- (1S,2R,4aS,8aS)-1-(3-methoxy-5-methylphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;
- (1S,2R,4aS,8aS)-1-(3,5-dimethoxyphenoxyethyl)-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;
- (1S,2R,4aS,8aS)-1-[3,5-bis(propan-2-yloxy)phenoxyethyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;
- (1R,2R,8aS)-1-((3,5-dimethoxyphenylsulfonyl)methyl)-2,5,5,8a-tetramethyldecahydronaphthalen-2-ol;
- (1R,2R,4aS,8aS)-1-[[3-methoxy-5-methylphenyl)sulfonyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol;
- (1R,2R,4aS,8aS)-1-[[3-methoxyphenyl)sulfonyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol; and
- (1R,2R, 4aS,8aS)-1-[[3,5-dimethoxyphenyl)sulfonyl]methyl]-2,5,5,8a-tetramethyl-decahydronaphthalen-2-ol.

2. The compound of Claim 1 having the formula (Ia):




wherein n, R¹, R², R³, R^{4a}, R^{4b}, R⁵, R⁶, R⁷ and R⁸ are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

3. The compound of Claim 2 where R¹ is -R^{9a}-C(O)-R^{9b}-, *i.e.*, a compound of formula (Ia1):




wherein R^{9a} and R^{9b} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

4. The compound of Claim 2 wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, the compound of formula (Ia2):




wherein R^{9a} , R^{9b} and R^{10} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

5. The compound of Claim 2, wherein R^1 is $-R^9-S(O)_t-R^9-$, *i.e.*, a compound of formula (Ia3):



wherein t , R^{9a} and R^{9b} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.


6. The compound of Claim 2, wherein R^1 is $-R^{9a}-O-R^{9b}-$, *i.e.*, a compound of formula (Ia4):



;

wherein R^{9a} and R^{9b} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.


7. The compound of Claim 2, wherein R^1 is $-R^{9a}-C(O)N(R^{11a})-R^{9b}-$, *i.e.*, a compound of formula (Ia5):



;


wherein R^{9a} , R^{9b} and R^{11a} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

8. The compound of Claim 2 where R^1 is $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$, *i.e.*, a compound of formula (Ia6):




wherein R^{9a} , R^{9b} and R^{11a} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

9. The compound of Claim 2 wherein R^1 is $-R^{9a}-N(R^{11a})-R^{9b}-$, *i.e.*, a compound of formula (Ia7):




wherein R^{9a} , R^{9b} and R^{11a} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

10. The compound of Claim 1 having the formula (Ic):




wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

11. The compound of Claim 10 wherein R^1 is $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$, *i.e.*, a compound of formula (lc1):




wherein R^{9a} , R^{9b} and R^{11a} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

12. The compound of Claim 1 having the formula (ld):




wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

13. The compound of Claim 12 wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, a compound of formula (ld1):




wherein R^{9a} , R^{9b} and R^{10} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

14. The compound of Claim 1 having the formula (Ie):




wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

15. The compound of Claim 14 wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, a compound of formula (Ie1):




wherein R^{9a} , R^{9b} and R^{10} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

16. The compound of Claim 1 having the formula (If):




wherein n , R^1 , R^{4a} , R^{4b} , R^5 , R^6 , R^7 and R^8 are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

17. The compound of Claim 16 wherein R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, *i.e.*, a compound of formula (If1):




wherein R^{9a} , R^{9b} and R^{10} are as defined above in Claim 1, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

18. The compound of Claim 1 wherein:



is selected from:



n is 1, 2, 3, 4, 5, or 6;

R^1 is $-R^{9a}-C(R^{10})_2-R^{9b}-$, $-R^{9a}-C(O)-R^{9b}-$, $-R^{9a}-S(O)_t-R^{9b}-$ (where t is 0, 1 or 2), $-R^{9a}-O-R^{9b}-$, or $-R^{9a}-C(O)N(R^{11a})-R^{9b}-$ or $-R^{9a}-N(R^{11a})C(O)-R^{9b}-$;

R^{4a} and R^{4b} are each independently selected from hydrogen, alkyl, $-R^9-OR^{11}$ or $-C(O)OR^{11}$,

or R^{4a} is selected from hydrogen, alkyl, $-R^9-OR^{11}$, or $-C(O)OR^{11}$ and R^{4b} is a direct bond to the carbon at numeral 1 or a direct bond to the carbon at numeral 3;

R^5 is independently selected from hydrogen, oxo, cyano, nitro, halo, alkyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^9-OR^{11}$, $-R^9-C(O)R^{11}$, $-R^9-C(O)OR^{11}$, $-R^9-N(R^{11})R^{12}$, $-R^9-C(O)N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)R^{12}$, $-R^9-N(R^{11})-R^{14}-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)-R^9-N(R^{11})R^{12}$, $-R^9-N(R^{11})C(O)N(R^{11})-OR^{12}$, $-R^9-N(R^{11})C(=NR^{11})N(R^{11})R^{12}$, $-R^9-N(R^{11})S(O)_pR^{11}$ (where p is 1 or 2), $-R^9-N(R^{11})C(S)N(R^{11})R^{12}$ or

$-R^9-N(R^{11})C(O)-R^9-N(R^{11})S(O)_pR^{12}$ (where p is 1 or 2);
 each R^6 and R^8 is independently selected from hydrogen, alkyl, halo or haloalkyl;
 R^7 is hydrogen, alkyl, halo or haloalkyl;
 each R^9 , R^{9a} and R^{9b} is independently a direct bond or a straight or branched alkylene chain;
 each R^{10} is independently hydrogen, alkyl, $-OR^{11}$, $-C(O)OR^{11}$, $-C(O)N(R^{11})R^{12}$, $-N(R^{11})R^{12}$ or $-N(R^{11})C(O)R^{11}$;
 each R^{11} , R^{11a} and R^{12} is independently hydrogen, alkyl, alkenyl, haloalkyl, optionally substituted cycloalkyl, optionally substituted cycloalkylalkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl or optionally substituted heteroarylalkyl; and
 R^{14} is a straight or branched alkylene chain;
 or a stereoisomer thereof or a pharmaceutically acceptable salt thereof.


19. A compound selected from the group consisting of:
 (2R,4R,4aS,8aS)-4-[(3,5-dimethoxyphenyl)carbonyl]-4a,8,8-trimethyl-3-methylidene-decahydronaphthalen-2-ol;
 (4aS,5S,8aS)-5-[(3-methoxy-5-methylphenyl)methyl]-1,1,4a-trimethyl-6-methylidene-decahydronaphthalene;
 N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)nicotinamide;
 N-((2R,3S,4S,4aS)-4-((furan-2-carboxamido)methyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide; and
 N-((2R,3S,4S,4aR)-4-(2-(furan-2-carboxamido)ethyl)-3,4a,8,8-tetramethyldecahydronaphthalen-2-yl)-4-methylpiperazine-1-carboxamide;
 or a pharmaceutically acceptable salt thereof.

20. A composition comprising a compound of Claim 1 or Claim 19, or a stereoisomer thereof or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

21. A method for modulating SHIP1 comprising administering an effective amount of a compound of Claim 1 or Claim 19, or a pharmaceutically acceptable salt thereof or a stereoisomer thereof, or a composition of Claim 20 to a mammal in need

thereof.

22. A method for treating a disease, disorder or condition comprising administering an effective amount of a compound of Claim 1 or Claim 19, or a pharmaceutically acceptable salt or stereoisomer thereof, or a composition of Claim 20 to a mammal in need thereof, where the disease, disorder or condition is an autoimmune disease, disorder or condition, an inflammatory disease, disorder or condition, or a neoplastic or cell proliferative disease, disorder or condition.



;

