

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2013/139694 A1

(43) International Publication Date
26 September 2013 (26.09.2013)

WIPO | PCT

(51) International Patent Classification:

A61K 38/26 (2006.01) *A61K 47/00* (2006.01)
A61P 3/10 (2006.01)

(21) International Application Number:

PCT/EP2013/055362

(22) International Filing Date:

15 March 2013 (15.03.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12160743.6	22 March 2012 (22.03.2012)	EP
61/748,840	4 January 2013 (04.01.2013)	US
13153459.6	31 January 2013 (31.01.2013)	EP

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).

(72) Inventors: VILHELMSEN, Thomas; Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd (DK). ELIASSEN, Helle; Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd (DK). HANSEN, Tue; Novo Nordisk, Novo Allé, DK-2880 Bagsværd (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2013/139694 A1

(54) Title: COMPOSITIONS OF GLP-1 PEPTIDES AND PREPARATION THEREOF

(57) Abstract: The invention relates to pharmaceutical compositions comprising a first type of granules and a second type of granules, wherein said first type of granules comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and no GLP-1 peptide, and wherein said second type of granules comprises a GLP-1 peptide and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, as well as the intermediate granules, processes for the preparation of the granules and compositions, and use thereof in medicine.

Compositions of GLP-1 peptides and preparation thereof

The present invention is directed to compositions comprising pharmacologically active agents, such as GLP-1, and a delivery agent as well as processes for their preparation and use thereof in medicine.

5 BACKGROUND

One of the main challenges in oral delivery of proteins and peptides is the inability of these compounds to be readily transported across the membranes of the gastrointestinal tract. The delivery agent SNAC has previously been shown to improve the bioavailability of orally administered peptides. The present invention relates to further improvements of the 10 bioavailability by oral administration of compositions of such peptides, in particular of GLP-1 peptides.

SUMMARY

In one embodiment the invention relates to a pharmaceutical composition comprising a first type and a second type of granules, wherein said first type of granules 15 comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and no GLP-1 peptide, and wherein said second type of granules comprises a GLP-1 peptide and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the term "granule" refers to small particles gathered into a large mass.

In one embodiment the invention relates to a process of producing a pharmaceutical 20 composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide with a filler and/or a binder; b) dry granulation of the mixture of step a; c) mixing the granules obtained in step b with a composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino) caprylic acid; and d) optionally adding further lubricant, wherein the mixture of step a does not comprise a 25 salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

In one embodiment the invention relates to a process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler; b) dry granulation of the 30 mixture of step a; c) mixing the granules obtained in step b with a composition comprising a GLP-1 peptide; and d) optionally adding further lubricant, wherein the mixture of step a does not comprise a GLP-1 peptide.

In one embodiment the invention relates to a process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide with a filler and/or a binder; b) dry granulation of the mixture of step a; c) mixing a salt of N-(8-(2-

5 hydroxybenzoyl)amino)caprylic acid, optionally a lubricant and/or a filler; d) dry granulation of the mixture of step c; e) mixing the granules obtained in step b with the granules obtained in step d; and f) optionally adding further lubricant, wherein the mixture of step a does not comprise a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and wherein the mixture of step c does not comprise a GLP-1 peptide.

10 In one embodiment the invention relates to a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, a lubricant, optionally a filler, and no GLP-1 peptide. In one embodiment the invention relates to a process of producing a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, wherein the process comprises the steps: a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid (such as SNAC) with a lubricant 15 and/or a filler; and b) dry granulation of the mixture of step a, wherein the mixture of step a does not comprise a GLP-1 peptide.

20 In one embodiment the invention relates to a granule comprising a GLP-1 peptide, a filler, a binder, and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In one embodiment the invention relates to a process of producing a granule comprising a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide, optionally a 25 filler and/or a binder; and b) roller compacting the mixture of step a, wherein the mixture of step a does not comprise a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

In one embodiment the invention relates to a pharmaceutical composition or a granule obtained by the process as defined herein.

30 In one embodiment the invention relates to a composition or a granule as defined herein for use in medicine, such as for treatment of diabetes or obesity, wherein said composition is administered orally. In one embodiment the invention relates to a method for treatment of diabetes or obesity comprising administering the composition as defined herein to a patient in need thereof, wherein said composition is administered orally.

DESCRIPTION

The present inventors surprisingly found that the dissolution properties of tablets manufactured from the same composition (i.e. same type and the same amounts of

excipients and delivery agent) was determined by the design of the granules from which the tablets were formed.

Furthermore, the present inventors found that the dissolution behaviour surprisingly had a marked effect on the bioavailability of GLP-1 peptide from the composition, such as the 5 tablet. Thus, the present inventors have shown that the bioavailability of solid tablets manufactured from various granule designs can be predicted from *in vitro* data, i.e. dissolution data. In one embodiment this invention provides compositions, granules and methods for their preparation with improved bioavailability of the GLP-1 peptide.

Generally, the term "bioavailability" as used herein refers to the fraction of an 10 administered dose of an active pharmaceutical ingredient (API) and/or active moieties, such as a GLP-1 peptide as defined herein, which reaches the systemic circulation un-changed or in another active form. By definition, when an API and/or active moieties are administered intravenously, its bioavailability is 100%. However, when it is administered via other routes (such as orally), its bioavailability decreases (due to incomplete absorption and/or first-pass 15 metabolism). Knowledge about bioavailability is important when calculating dosages for non-intravenous routes of administration.

Absolute oral bioavailability is calculated as the relative exposure of the API and/or active moieties in systemic circulation following oral administration (estimated as the area under the plasma concentration versus time curve) compared to the exposure of the API 20 following intravenous administration.

Pharmaceutical compositions

In some embodiments the invention relates to a pharmaceutical composition comprising a first type and a second type of granules, wherein said first type of granules comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and said second type of 25 granules comprises a GLP-1 peptide. In some embodiments the first type of granules further comprises a lubricant, such as magnesium stearate. In some embodiments the first type of granules further comprises a filler, such as microcrystalline cellulose. Accordingly, the first type of granules may further comprise a lubricant and optionally a filler. In some embodiments the second type of granules further comprises a filler, such as microcrystalline 30 cellulose. In some embodiments the second type of granules further comprises a binder, such as povidone. Accordingly, the second type of granules may further comprise a filler and optionally a binder. In some embodiments the composition further comprises an extragranular lubricant, such as magnesium stearate.

In some embodiments N-(8-(2-hydroxybenzoyl)amino)caprylic acid is referred to as "NAC".

In some embodiments the first type of granules comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid does not contain a GLP-1 peptide. In some embodiments 5 the second type of granules comprising a GLP-1 peptide does not contain a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the pharmaceutical composition comprises a first type and a second type of granules, wherein said first type of granules comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and no GLP-1 peptide, and wherein said second type of granules comprises a GLP-1 peptide and no salt of N-(8-(2-10 hydroxybenzoyl)amino)caprylic acid.

In some embodiments the term "granule" refers to particles gathered into larger particles.

In some embodiments the invention relates to a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, 15 wherein the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is simultaneous with or faster than the release of said GLP-1 peptide as determined by dissolution testing using Assay (I). In some embodiments the invention relates to a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the release of said salt of N-(8-(2-20 hydroxybenzoyl)amino)caprylic acid is simultaneous with or faster than the release of said GLP-1 peptide.

In some embodiments the term "release" when used with reference to N-(8-(2-hydroxybenzoyl)amino)caprylic acid and optionally compared to release of GLP-1 peptide is determined within 30 minutes, such as within 25, 20, 15 minutes, or such as within 10 or 5 25 minutes, as determined by dissolution testing using Assay (I). Accordingly, the release may be determined within 30 minutes of dissolution testing using Assay (I). In some embodiments the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid when referred to as "faster than" the release of said GLP-1 peptide is determined within 30 minutes, such as within 25, 20, 15 minutes, or such as within 10 or 5 minutes, as determined by dissolution 30 testing using Assay (I). In some embodiments the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid when referred to as "faster than" the release of said GLP-1 peptide is determined within 30 minutes, such as within 25, 20, 15 minutes, or such as within 10 or 5 minutes, as determined by dissolution testing using Assay (I) at pH 2.5.

In some embodiments the release of said salt of N-(8-(2-35 hydroxybenzoyl)amino)caprylic acid when referred to as "simultaneous with" the release of

5 said GLP-1 peptide is determined within 30 minutes, such as within 25, 20, 15 minutes, or such as within 10 or 5 minutes, as determined by dissolution testing using Assay (I). In some embodiments the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid when referred to as "simultaneous with" the release of said GLP-1 peptide is determined within 30 minutes, such as within 25, 20, 15 minutes, or such as within 10 or 5 minutes, as determined by dissolution testing using Assay (I) at pH 2.5.

10 In some embodiments the invention relates to a pharmaceutical composition according to any one of the preceding claims, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within the first 60 minutes as determined by dissolution testing using Assay (I) at pH 2.5. In some embodiments the invention relates to a pharmaceutical composition, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within the first 30 minutes as determined by dissolution testing using Assay (I) at pH 2.5. In some embodiments the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 55 minutes, such as within 15 50, 45 or 40 minutes, as determined by dissolution testing using Assay (I) at pH 2.5. In some embodiments the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 39 or 38 minutes, such as within 37, 36, or 35 minutes, as determined by dissolution testing using Assay (I) at pH 2.5. In some embodiments the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 34 minutes, such as within 20 33, 32, or 31 minutes, as determined by dissolution testing using Assay (I) at pH 2.5. In some embodiments the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 31 minutes, such as within 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, 50, 55, or 60 minutes as determined by dissolution testing using Assay (I) at pH 2.5.

25 In some embodiments the composition comprises granules which have been manufactured by dry granulation. In some embodiments the composition comprises granules which have been manufactured by roller compaction. In some embodiments the moldings from the roller compactions process are comminuted into granules. As used herein the term "composition" refers to a pharmaceutical composition.

30 In some embodiments the composition is in the form of a solid dosage form. In some embodiments the composition is in the form of a tablet. In some embodiments the composition is in the form of a capsule. In some embodiments the composition is in the form of a sachet.

35 In some embodiments the composition or granule comprises at least one pharmaceutically acceptable excipient. The term "**excipient**" as used herein broadly refers

to any component other than the active therapeutic ingredient(s). The excipient may be an inert substance, which is inert in the sense that it substantially does not have any therapeutic and/or prophylactic effect per se. The excipient may serve various purposes, e.g. as a delivery agent, absorption enhancer, vehicle, filler (also known as diluents), binder, lubricant, 5 glidant, disintegrant, crystallization retarders, acidifying agent, alkalizing agent, preservative, antioxidant, buffering agent, chelating agent, complexing agents, surfactant agent, emulsifying and/or solubilizing agents, sweetening agents, wetting agents stabilizing agent, colouring agent, flavouring agent, and/or to improve administration, and/or absorption of the active substance. A person skilled in the art may select one or more of the aforementioned 10 excipients with respect to the particular desired properties of the solid oral dosage form by routine experimentation and without any undue burden. The amount of each excipient used may vary within ranges conventional in the art. Techniques and excipients which may be used to formulate oral dosage forms are described in *Handbook of Pharmaceutical Excipients*, 6th edition, Rowe et al., Eds., American Pharmaceuticals Association and the 15 Pharmaceutical Press, publications department of the Royal Pharmaceutical Society of Great Britain (2009); and *Remington: the Science and Practice of Pharmacy*, 21th edition, Gennaro, Ed., Lippincott Williams & Wilkins (2005).

In some embodiments the composition or granule comprises a **filler**, such as lactose (e.g. spray-dried lactose, α -lactose, β -lactose, Tabletose $^{\circledR}$, various grades of 20 Pharmatose $^{\circledR}$, Microtose $^{\circledR}$ or Fast-FloC $^{\circledR}$), microcrystalline cellulose (various grades of Avicel $^{\circledR}$, Elcema $^{\circledR}$, Vivacel $^{\circledR}$, Ming Tai $^{\circledR}$ or Solka-Floc $^{\circledR}$), other cellulose derivatives, sucrose, sorbitol, mannitol, dextrins, dextrose, maltodextrins, dextrose, fructose, kaolin, mannitol, sorbitol, sucrose, sugar, starches or modified starches (including potato starch, maize starch and rice starch), calcium phosphate (e.g. basic calcium phosphate, calcium 25 hydrogen phosphate, dicalcium phosphate hydrate), calcium sulphate, calcium carbonate, or sodium alginate. In some embodiments the filler is microcrystalline cellulose, such as Avicel PH 101.

In some embodiments the composition or granule comprises a **binder**, such as lactose (e.g. spray-dried lactose, α -lactose, β -lactose, Tabletose $^{\circledR}$, various grades of 30 Pharmatose $^{\circledR}$, Microtose $^{\circledR}$ or Fast-FloC $^{\circledR}$), microcrystalline cellulose (various grades of Avicel $^{\circledR}$, Elcema $^{\circledR}$, Vivacel $^{\circledR}$, Ming Tai $^{\circledR}$ or Solka-Floc $^{\circledR}$), hydroxypropylcellulose, L- hydroxypropylcellulose (low-substituted), hypromellose (HPMC) (e.g. Methocel E, F and K, Metolose SH of Shin-Etsu, Ltd, such as, e.g., the 4,000 cps grades of Methocel E and Metolose 60 SH, the 4,000 cps grades of Methocel F and Metolose 65 SH, the 4,000, 15,000 35 and 100,000 cps grades of Methocel K; and the 4,000, 15,000, 39,000 and 100,000 grades

of Metolose 90 SH), methylcellulose polymers (such as, e.g., Methocel A, Methocel A4C, Methocel A15C, Methocel A4M), hydroxyethylcellulose, ethylcellulose, sodium carboxymethylcellulose, other cellulose derivatives, sucrose, dextrins, maltodextrins, starches or modified starches (including potato starch, maize starch and rice starch), calcium 5 lactate, calcium carbonate, acacia, sodium alginate, agar, carrageenan, gelatin, guar gum, pectin, PEG, or povidone. In some embodiments the binder is povidone, such as povidone K 90.

In some embodiments the composition or granule comprises a **disintegrant**, such as alginic acid, alginates, microcrystalline cellulose, hydroxypropyl cellulose, other cellulose 10 derivatives, croscarmellose sodium, crospovidone, polacrilin potassium, sodium starch glycolate, starch, pregelatinized starch, or carboxymethyl starch (e.g. Primogel® and Explotab®).

In some embodiments the composition or granule comprises a **lubricant**, such as stearic acid, magnesium stearate, calcium stearate or other metallic stearate, talc, waxes, 15 glycerides, light mineral oil, glycerylbehenate, hydrogenated vegetable oils, sodium stearylfumarate, polyethylene glycols, alkyl sulfates, or sodium benzoate. In some embodiments the composition or granule comprises a lubricant, such as magnesium silicate, talc, or colloidal silica. In some embodiments the lubricant is magnesium stearate.

In some embodiments the composition or granule comprises one or more excipients 20 selected from **crystallization retarders**, such as Povidone, etc.; **solubilizing agents** (also known as surfactants), such as anionic surfactants (e.g. Pluronic or Povidone), cationic surfactants, nonionic surfactants, and/or zwitterionic surfactants; **colouring agents**, including dyes and pigments, such as Iron Oxide Red or Yellow, titanium dioxide, and/or talc; and/or **pH control agents**, such as citric acid, tartaric acid, fumaric acid, sodium citrate, 25 dibasic calcium phosphate, and/or dibasic sodium phosphate.

In some embodiments the composition comprises at least 60% (w/w) delivery agent, less than 10% (w/w) binder, 5-40% (w/w) filler, and less than 10% (w/w) lubricant.

In some embodiments the composition comprises at least 60% (w/w), such as 65-75 % (w/w), 60-80 % (w/w), or 50-90 % (w/w), delivery agent. In some embodiments the 30 composition comprises at least 70% (w/w), such as 70-80 % (w/w), delivery agent.

In some embodiments the composition comprises 0.1-10% (w/w), such as 0.2-4% (w/w) or 0.5-3% (w/w), binder. In some embodiments the composition comprises 1.5-2.5 % (w/w), such as 1.7-2.3 % (w/w), 1.8-2.2 % (w/w), or 1.9-2.1 % (w/w), binder. In some 35 embodiments the composition comprises 1% (w/w) or 2% (w/w) binder.

In some embodiments the composition comprises 5-40% (w/w), such as 10-30% (w/w) or 5-25% (w/w), filler. In some embodiments the composition comprises 10-25 % (w/w), such as 17-23 % (w/w), 18-22 % (w/w), or 19-21 % (w/w), filler. In some embodiments the composition comprises 10.9% (w/w) or 18% (w/w) filler, or comprises 19.5% (w/w) or 20.5% (w/w) filler.

In some embodiments the composition comprises 0.1-10% (w/w) or 0.5-5% (w/w), such as 1-3.5% (w/w) or 1% (w/w), lubricant. In some embodiments the composition comprises 1.5-3 % (w/w), such as 2.1-2.7 % (w/w), 2.2-2.6 % (w/w) or 2.3-2.5 % (w/w), lubricant.

10 Still further, the composition or granule of the invention may be formulated as is known in the art of oral formulations of insulinotropic compounds.

15 The composition or granule may be administered in several dosage forms, for example as a tablet; a capsule such as hard capsules, sachet or a powder. The composition or granule may further be compounded in a drug carrier or drug delivery system, e.g. in order to improve stability and/or solubility or further improve bioavailability.

In some embodiments the weight of the tablet is in the range of 150 mg to 1000 mg, such as in the range of 300-600 mg or 350-450 mg.

20 In some embodiments the invention relates to a first granule comprising at least 75 % (w/w) delivery agent, less than 10% (w/w) lubricant, and optionally less than 20 % filler and no GLP-1 peptide. In some embodiments the invention relates to a first granule comprising at least 80% (w/w) delivery agent, less than 10% (w/w) lubricant, and optionally less than 20 % filler and no GLP-1 peptide. In some embodiments the first granule comprises 75-90 % (w/w), such as 78-88 % (w/w), 80-86 % (w/w) or 82-84 % (w/w), delivery agent. In some embodiments the first granule comprises less than 10% (w/w), such as 1-3 % (w/w), 1.5-2.5 % (w/w) or 1.9-2.3 % (w/w), lubricant. In some embodiments the first granule comprises less than 20 %, such as 10-20 % (w/w), 12-18 % (w/w) or 14-17 % (w/w), filler. In some embodiments the first granule comprises no GLP-1 peptide. In some embodiments the granule comprises at least 80% (w/w) delivery agent, less than 10% (w/w) lubricant, and optionally less than 20% filler.

30 In some embodiments the invention relates to a second granule comprising a GLP-1 peptide, at least 15 % (w/w) filler and less than 40% (w/w) binder and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the second granule comprises at least 1 %, such as 1-70 % (w/w), 2-40% (w/w) or 4-30% (w/w), GLP-1 peptide. In some embodiments the second granule comprises at least 20 %, such as 40-80% (w/w) or 50-75% (w/w), filler. In some embodiments the second granule comprises less than 30 %, such as 5-

30% (w/w), 10-28% (w/w) or 15-25% (w/w),binder. In some embodiments the second granule comprises no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the granule comprises a GLP-1 peptide, at least 15 % (w/w) filler and less than 40% (w/w) binder. In some embodiments the granule comprises a GLP-1 peptide, at least 50% (w/w) 5 filler and less than 40% (w/w) binder.

In some embodiments the invention relates to a composition comprising a first and a second type of granules, wherein the first type of granule comprises at least 75 % (w/w) delivery agent, less than 10% (w/w) lubricant, optionally less than 20 % filler and no GLP-1 peptide, and wherein the second type of granule comprises a GLP-1 peptide, at least 15 % 10 (w/w) filler, less than 40% (w/w) binder and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the invention relates to a composition comprising a first and a second type of granules, wherein the first type of granule comprises at least 75 % (w/w) delivery agent, less than 10% (w/w) lubricant, less than 20 % filler and no GLP-1 peptide, and wherein the second type of granule comprises a GLP-1 peptide, at least 15 % (w/w) filler, 15 less than 40% (w/w) binder and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the invention relates to a composition comprising a first and a second type of granules, wherein the first type of granule comprises at least 75 % (w/w) delivery agent, less than 10% (w/w) lubricant, and no GLP-1 peptide, and wherein the second type of granule comprises a GLP-1 peptide, at least 15 % (w/w) filler, less than 40% (w/w) binder and 20 no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

Methods of preparation of pharmaceutical compositions

The composition of the invention may be prepared as is known in the art. In some embodiments the composition or the granule may be prepared as described in the examples 25 herein. In some embodiments the composition may be granulated prior to being compressed into tablets. In some embodiments the granules of the invention are manufactured by dry granulation, such as by roller compactioncompaction. In some embodiments the moldings from the roller compactions process are comminuted into granules. The composition may comprise one or more intragranular parts and an extragranular part, wherein the 30 intragranular parts have been granulated, and wherein the extragranular part has been added after granulation. A first intragranular part may comprise the GLP-1 peptide and one or more excipients, and a second intragranular part may comprise the delivery agent and optionally one or more excipients. A first intragranular part may comprise the GLP-1 peptide, filler and/or a binder and a second intragranular part may comprise the delivery agent,

lubricant and filler. The first intragranular part may comprise the GLP-1 peptide, filler and/or binder and a second intragranular part may comprise the delivery agent, lubricant and/or filler. In some embodiments the firstintragranular part comprises the GLP-1 agonist (i.e. the GLP-1 peptide), microcrystalline cellulose and/or povidone and the secondintragranular part 5 comprises the delivery agent, magnesium stearate and microcrystalline cellulose. In some embodiments the firstintragranular part comprises the GLP-1 agonist (i.e. the GLP-1 peptide), microcrystalline cellulose and/or povidone and the secondintragranular part comprises the delivery agent, magnesium stearate and/or microcrystalline cellulose. The extragranular part may comprise a lubricant. In some embodiments the extragranular part 10 comprises magnesium stearate. In some embodiments the term "filler and/or binder" or "filler and/or a binder" refers to a filler and optionally a binder. In some embodiments the term "lubricant and filler" or "lubricant and/or filler" refers to a lubricant and optionally a filler.

15 To prepare a dry **blend** of tabletting material, the various components are weighed, optionally delumped and then combined. The mixing of the components may be carried out until a homogeneous blend is obtained.

If granules are to be used in the tabletting material, granules may be produced in a manner known to a person skilled in the art, for example by **dry granulation** techniques in which the pharmaceutically active agent and/or delivery agents are compacted with the excipients to form relatively large moldings, for example slugs or ribbons, which are 20 comminuted by grinding, and the ground material serves as the tabletting material to be later compressed into tablets. Suitable equipment for dry granulation includes but is not limited to roller compaction equipment from Gerteis, such as Gerteis MINI-PACTOR.

25 To **compress** the tabletting material into a solid oral dosage form, for example a tablet, a tablet press may be used. In a tabletting press, the tabletting material is filled (e.g. force fed or gravity fed) into a die cavity. The tabletting material is then compressed by a punch with pressure. Subsequently, the resulting compact, or tablet is ejected from the tabletting press. The above mentioned compression process is subsequently referred to herein as the "compression process". Suitable **tablet presses** include, but are not limited to, rotary tablet presses and eccentric tablet presses. Examples of tablet presses include, but 30 are not limited to, the Fette 102i (Fette GmbH), the Korsch XL100, the Korsch PH 106 rotary tablet press (Korsch AG, Germany), the Korsch EK-O eccentric tabletting press (Korsch AG, Germany) and the Manesty F-Press (Manesty Machines Ltd., United Kingdom).

35 In some embodiments the method of preparation of the tablet comprises i) dry granulation of a mixture comprising the GLP-1 agonist (i.e. the GLP-1 peptide), filler and a binder; ii) dry granulation of a mixture comprising the delivery agent, lubricant and filler; iii)

mixing of the granules with a lubricant; and then iv) compression of the blend into tablets. In some embodiments the method of preparation of the tablet comprises i) dry granulation of a mixture comprising the GLP-1 agonist (i.e. the GLP-1 peptide), filler and a binder; ii) dry granulation of a mixture comprising the delivery agent, lubricant and/or filler; iii) mixing of the 5 granules with a lubricant; and then iv) compression of the blend into tablets.

In some embodiments the invention relates to a process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide with a filler and/or a binder; b) dry granulation of the mixture of step a; c) mixing the granules obtained in step b with a composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid; and d) optionally adding further lubricant. In some embodiments the invention relates to a process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant 10 and/or a filler; b) dry granulation of the mixture of step a; c) mixing the granules obtained in step b with a composition comprising a GLP-1 peptide; and d) optionally adding further lubricant. In some embodiments the invention relates to a process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide 15 with a filler and/or a binder; b) dry granulation of the mixture of step a; c) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, optionally a lubricant and/or a filler; d) dry granulation of the mixture of step c; e) mixing the granules obtained in step b with the granules obtained in step d; and f) optionally adding further lubricant.

In one embodiment the invention relates to a process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide with a filler and/or a binder; b) dry granulation of the mixture of step a; c) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, optionally a lubricant and/or a filler; d) dry granulation of the mixture of step c; e) mixing the granules obtained in step b with the granules obtained 20 in step d; and f) optionally adding further lubricant, wherein the mixture of step a does not comprise a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and wherein the mixture of step c does not comprise a GLP-1 peptide.

In one embodiment the invention relates to a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, a lubricant, optionally a filler, and no GLP-1 peptide. In 35 one embodiment the invention relates to a process of producing a granule comprising a salt

of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, wherein the process comprises the steps: a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid (such as SNAC) with a lubricant and/or a filler; and b) dry granulation of the mixture of step a, wherein the mixture of step a does not comprise a GLP-1 peptide.

5 In one embodiment the invention relates to a granule comprising a GLP-1 peptide, a filler, a binder, and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In one embodiment the invention relates to a process of producing a granule comprising a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide, optionally a filler and/or a binder; and b) roller compacting the mixture of step a, wherein the mixture of
10 step a does not comprise a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

In some embodiments the mixing step comprising mixing of a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler prior to dry granulation has a duration of at least 20 minutes. In some embodiments the mixing step comprising mixing of a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler prior to dry
15 granulation has a duration of at least 30 minutes or at least 40 minutes, such as 50 minutes. In some embodiments the mixing step comprising mixing of the first type of granules comprising a GLP-1 peptide with the second type of granules comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid has a duration of at least 15 minutes or at least 20 minutes, such as at least 25 minutes or at least 30 minutes.

20 In some embodiments the invention relates to a pharmaceutical composition obtained by the process as defined herein.

In some embodiments the invention relates to a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a lubricant. In some embodiments the granule comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, a lubricant, and a filler. In
25 some embodiments the invention relates to a granule comprising a GLP-1 peptide, a filler and a binder. In some embodiments the granule is prepared according to the process as defined herein.

In some embodiments the invention relates to a process of producing a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, wherein the process
30 comprises the steps: a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid (such as SNAC) with a lubricant and/or a filler; and b) dry granulation of the mixture of step a. In some embodiments the mixing step comprising mixing of a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler prior to dry granulation has a duration of at least 20 minutes. In some embodiments the mixing step comprising mixing of a salt of N-(8-(2-

hydroxybenzoyl)amino) caprylic acid with a lubricant and/or a filler prior to dry granulation has a duration of at least 30 minutes or at least 40 minutes, such as 50 minutes.

In some embodiments the invention relates to a process of producing a granule comprising a GLP-1 peptide, wherein the process comprises the steps: a) mixing a GLP-1 peptide, optionally a filler and/or a binder; and b) roller compacting the mixture of step a.

In some embodiments the invention relates to a granule obtained by the process as defined herein.

In some embodiments the lubricant is magnesium stearate. In some embodiments the filler is microcrystalline cellulose. In some embodiments the binder is povidone.

10

In some embodiments the term "resistance to crushing of tablets" has the meaning defined in section 2.9.8 in the European Pharmacopoeia 7.5, 7th edition 2012; resistance to crushing may be measured inter alia in Newton (N) or kilopond (kP) using a jaw speed of 20 N/s (1 kP equals 9.807 N).

15

In some embodiments the term "roller compaction force" means the force between the rolls of the roller compactor when compacting materials into a continuous strip of compressed material as determined by a pressure transducer that converts the hydraulic pressure into electrical signal; the roller compaction force may be measured in kiloNewton (kN) or in kiloNewton per roll width (kN/cm).

20

In vitro methods

Dissolution of compositions of the invention may be determined as described in Assay (I) herein. In some embodiments the dissolution testing using Assay (I) is carried out at any one of pH 1.0-8.0. In some embodiments the dissolution testing using Assay (I) is carried out at pH 1.0. In some embodiments the dissolution testing using Assay (I) is carried out at pH 2.5. In some embodiments the dissolution testing using Assay (I) is carried out at pH 6.8.

GLP-1 peptides

In one embodiment the composition of the invention comprises a GLP-1 peptide. The term "GLP-1 peptide" as used herein refers to a compound, which fully or partially activates the human GLP-1 receptor. In some embodiments the "GLP-1 peptide" binds to a GLP-1 receptor, e.g., with an affinity constant (K_D) or activate the receptor with a potency (EC_{50}) of below 1 μ M, e.g. below 100 nM as measured by methods known in the art (see e.g. WO 98/08871) and exhibits insulinotropic activity, where insulinotropic activity may be

30

measured *in vivo* or *in vitro* assays known to those of ordinary skill in the art. For example, the GLP-1 peptide may be administered to an animal with increased blood glucose (e.g. obtained using an Intravenous Glucose Tolerance Test (IVGTT), a person skilled in the art will be able to determine a suitable glucose dosage and a suitable blood sampling regime, 5 e.g. depending on the species of the animal, for the IVGTT) and the plasma insulin concentration measured over time. In some embodiments the GLP-1 peptide is referred to as a GLP-1 agonist.

In some embodiments the GLP-1 peptide is a GLP-1 analogue, optionally comprising one substituent. The term "analogue" as used herein referring to a GLP-1 peptide 10 (hereafter "peptide") means a peptide wherein at least one amino acid residue of the peptide has been substituted with another amino acid residue and/or wherein at least one amino acid residue has been deleted from the peptide and/or wherein at least one amino acid residue has been added to the peptide and/or wherein at least one amino acid residue of the peptide has been modified. Such addition or deletion of amino acid residues may take place at the N- 15 terminal of the peptide and/or at the C-terminal of the peptide. In some embodiments a simple nomenclature is used to describe the GLP-1 peptide, e.g., [Aib8] GLP-1(7-37) designates an analogue of GLP-1(7-37) wherein the naturally occurring Ala in position 8 has been substituted with Aib. In some embodiments the GLP-1 peptide comprises a maximum 20 of twelve, such as a maximum of 10, 8 or 6, amino acids which have been altered, e.g., by substitution, deletion, insertion and/or modification, compared to e.g. GLP-1(7-37). In some embodiments the analogue comprises up to 10 substitutions, deletions, additions and/or insertions, such as up to 9 substitutions, deletions, additions and/or insertions, up to 8 substitutions, deletions, additions and/or insertions, up to 7 substitutions, deletions, additions 25 and/or insertions, up to 6 substitutions, deletions, additions and/or insertions, up to 5 substitutions, deletions, additions and/or insertions, up to 4 substitutions, deletions, additions and/or insertions or up to 3 substitutions, deletions, additions and/or insertions, compared to e.g. GLP-1(7-37). Unless otherwise stated the GLP-1 comprises only L-amino acids.

In some embodiments the term "GLP-1 analogue" or "analogue of GLP-1" as used 30 herein refers to a peptide, or a compound, which is a variant of the human Glucagon-Like Peptide-1 (GLP-1(7-37)). GLP-1(7-37) has the sequence HAEGTFTSDV SSYLEGQAAKEFIAWLVKGRG (SEQ ID No: 1). In some embodiments the term "variant" refers to a compound which comprises one or more amino acid substitutions, deletions, additions and/or insertions.

In one embodiment the GLP-1 peptide exhibits at least 60%, 65%, 70%, 80% or 35 90% sequence identity to GLP-1(7-37) over the entire length of GLP-1(7-37). As an example


of a method for determination of sequence identity between two analogues the two peptides [Aib8]GLP-1(7-37) and GLP-1(7-37) are aligned. The sequence identity of [Aib8]GLP-1(7-37) relative to GLP-1(7-37) is given by the number of aligned identical residues minus the number of different residues divided by the total number of residues in GLP-1(7-37).

5 Accordingly, in said example the sequence identity is (31-1)/31.

In one embodiment the C-terminal of the GLP-1 peptide is an amide.

In some embodiments the GLP-1 peptide is GLP-1(7-37) or GLP-1(7-36)amide. In some embodiments the GLP-1 peptide is exendin-4, the sequence of which is HGEGTFITSDL SKQMEEEAVRLFIEWLKNGGPSSGAPPSS (SEQ ID No: 2).

10 In some embodiments the GLP-1 peptide comprises one substituent which is covalently attached to the peptide. In some embodiments the substituent comprises a fatty acid or a fatty diacid. In some embodiments the substituent comprises a C16, C18 or C20 fatty acid. In some embodiments the substituent comprises a C16, C18 or C20 fatty diacid. In some embodiments the substituent comprises formula (X)



15 (X), wherein n is at least 13, such as n is 13, 14, 15, 16, 17, 18 or 19.

In some embodiments the substituent comprises formula (X), wherein n is in the range of 13 to 19, such as in the range of 13 to 17. In some embodiments the substituent comprises formula (X), wherein n is 13, 15 or 17. In some embodiments the substituent comprises formula (X), wherein n is 13. In some embodiments the substituent comprises formula (X), wherein n is 15. In some embodiments the substituent comprises formula (X), wherein n is 17. In some embodiments the substituent comprises one or more 8-amino-3,6-dioxaoctanoic acid (OEG), such as two OEG.

20 In some embodiments the substituent is [2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxyheptadecanoylamino) butyrylamino]ethoxy}ethoxy)acetylamino]ethoxy)acetyl].

25 In some embodiments the substituent is [2-(2-{2-[2-(2-{(S)-4-carboxy-4-(trans-4-[(19-carboxynonadecanoylamino)methyl]cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetylamino]ethoxy)acetyl].

30 In some embodiments the GLP-1 peptide is semaglutide, also known as N-epsilon26-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butyrylamino]ethoxy}ethoxy)acetylamino]ethoxy}ethoxy)acetyl][Aib8,Arg34]GLP-1(7-37), which may be prepared as described in WO2006/097537, Example 4.

In some embodiments the composition comprises the GLP-1 peptide or a pharmaceutically acceptable salt, amide, or ester thereof. In some embodiments the composition comprises the GLP-1 peptide one or more pharmaceutically acceptable counter ions.

5 In some embodiments the dosage of GLP-1 peptide is in the range of 0.01 mg to 100 mg. In some embodiments the composition or granule comprises an amount of a GLP-1 peptide in the range of at least 1 mg, such as at least 5 mg or at least 10 mg. In some embodiments the composition or granule comprises 10 mg GLP-1 peptide.

In some embodiments the composition comprises an amount of a GLP-1 peptide in
10 the range of 0.05 to 25 μ mol, such as in the range of 0.5 to 20 μ mol.

In some embodiments the GLP-1 peptide is selected from one or more of the GLP-1 peptides mentioned in WO93/19175, WO96/29342, WO98/08871, WO99/43707, WO99/43706, WO99/43341, WO99/43708, WO2005/027978, WO2005/058954, WO2005/058958, WO2006/005667, WO2006/037810, WO2006/037811, WO2006/097537, 15 WO2006/097538, WO2008/023050, WO2009/030738, WO2009/030771 and WO2009/030774.

cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]ethoxy}ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-37)amide; N-epsilon37-[2-(2-{2-[2-(2-[2-[S]-4-carboxy-4-({trans-4-[(19-carboxy-nonadecanoylamino)methyl]cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]ethoxy)acetyl][Aib8,Glu22,Arg26,5 Arg34,Lys37]GLP-1-(7-37)amide; N-epsilon37-[2-(2-{2-[2-(2-[2-[S]-4-carboxy-4-({trans-4-[(19-carboxy-nonadecanoylamino)methyl]cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]butyrylamino]ethoxy)acetyl][DesaminoHis7,Glu22,Arg26,Arg34,Lys37]GLP-1-(7-37)amide; N-epsilon37-[2-(2-{2-[2-(2-{2-[S]-4-carboxy-4-({trans-4-[(19-carboxy-nonadecanoylamino)methyl]cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]butyrylamino]ethoxy)acetyl][DesaminoHis7,Glu22,Arg26,Arg34,Lys37]GLP-1-(7-37); N-epsilon26[2-(2-{2-[2-(2-[S]-4-carboxy-4-({4-[(19-carboxy-15 nonadecanoylamino)methyl]cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]butyrylamino]ethoxy)acetyl][Aib8, Lys 26]GLP-1 (7-37)amide; N-epsilon26 [2-(2-[2-(2-[2-((S)-2-[trans-4-((9-carboxy nonadecanoylamino)methyl]cyclohexylcarbonyl)amino]-4-carboxybutanoylamino]ethoxy)ethoxy]acetyl]amino]ethoxy)acetyl][Aib8, Lys26] GLP-1 (7-37)amide; N-epsilon37-[2-(2-{2-[2-(2-[S]-4-carboxy-4-({trans-4-[(19-carboxy-20 nonadecanoylamino)methyl]cyclohexane-carbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]butyrylamino]ethoxy)acetyl][DesaminoHis7,Arg26,Arg34,Lys37]GLP-1-(7-37); N-epsilon37-[2-(2-{2-[2-(2-[S]-4-carboxy-4-({trans-4-[(19-carboxy-nonadecanoylamino)methyl]cyclohexanecarbonyl}amino)butyrylamino]ethoxy}ethoxy)acetyl]amino]butyrylamino]ethoxy)acetyl][DesaminoHis7,Glu22,25 Arg26,Glu30,Arg34,Lys37]GLP-1-(7-37); N-epsilon26[2-(2-{2-[S]-4-carboxy-4-{4-[4-(16-(1H-tetrazol-5-yl)-hexadecanoylsulfamoyl)butyrylamino]-butyrylamino}butyrylamino]ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-37); N-epsilon26[2-(2-{2-[S]-4-carboxy-4-((S)-4-carboxy-4-{12-[4-(16-(1H-tetrazol-5-yl)hexadecanoyl-sulfamoyl)butyrylamino]dodecanoyl}butyrylamino)butyrylamino]ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-37); N-epsilon26[2-(2-{2-[S]-4-carboxy-4-{6-[4-(16-(1H-tetrazol-5-yl)hexadecanoyl-sulfamoyl)butyrylamino]hexanoyl}butyrylamino]butyrylamino]ethoxy)ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-37); N-epsilon26[2-(2-{2-[S]-4-carboxy-4-((S)-4-carboxy-4-{4-(16-(1H-tetrazol-5-yl)hexadecanoylsulfamoyl)butyrylamino]butyrylamino}butyrylamino]ethoxy)ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-34); N-35

carboxyheptadecanoylamino)-4(S)-carboxybutyrylamino]ethoxy]ethoxy]ethoxy}
 ethoxy)ethoxy]ethoxy]ethoxy]propionyl}[Aib8,Arg34]GLP-1-(7-37); N-epsilon26-[2-(2-
 (2-(2-[2-(2-(4-(17-carboxyheptadecanoylamino)-4-carboxybutyrylamino)
 ethoxy)ethoxy]acetyl)ethoxy)ethoxy)acetyl}]-[Aib8,22,27,30,35,Arg34,Pro37, Lys26] GLP-1
 (7-37)amide; N-epsilon26-[2-(2-[2-[4-(21-carboxyuneicosanoylamino)-4(S)-
 carboxybutyrylamino]ethoxy]ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-37); and N-epsilon26-[2-(2-
 [2-(2-[2-(2-[4-(21-carboxyuneicosanoylamino)-4(S)-carboxybutyrylamino]
 ethoxy)ethoxy]acetyl)ethoxy]ethoxy)acetyl][Aib8,Arg34]GLP-1-(7-37). In one
 embodiment the GLP-1 peptide is N-epsilon26-[2-[2-(2-[2-(2-{(S)-4-carboxy-4-[10-(4-
 carboxyphenoxy) decanoylamino]butyrylamino}ethoxy)ethoxy] acetyl)ethoxy]ethoxy)
 ethoxy]acetyl}, N-epsilon37-[2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-[10-(4-carboxy-phenoxy)
 decanoylamino] butyrylamino}ethoxy)ethoxy]acetyl)ethoxy]ethoxy]acetyl}-
 [Aib8,Arg34,Lys37]GLP-1(7-37)-OH (Compound A).

In one embodiment the GLP-1 peptide is N-epsilon26-[2-(2-{2-[2-(2-{2-[(S)-4-carboxy-4-(17-carboxyheptadecanoylamino) butyrylamino]ethoxy}ethoxy)acetyl]amino]ethoxy}ethoxy)acetyl]Aib8,Arg34]GLP-1(7-37), also known as semaglutide.

In one embodiment GLP-1 peptides can be produced by appropriate derivatisation of an appropriate peptide backbone which has been produced by recombinant DNA technology or by peptide synthesis (e.g., Merrifield-type solid phase synthesis) as known in the art of peptide synthesis and peptide chemistry.


In one embodiment the production of peptides like GLP-1(7-37) and GLP-1 analogues is well known in the art. The GLP-1 moiety of the GLP-1 peptide of the invention (or fragments thereof) may for instance be produced by classical peptide synthesis, e.g., solid phase peptide synthesis using t-Boc or Fmoc chemistry or other well established techniques, see, e.g., Greene and Wuts, "Protective Groups in Organic Synthesis", John Wiley & Sons, 1999, Florencio Zaragoza Dörwald, "Organic Synthesis on solid Phase", Wiley-VCH Verlag GmbH, 2000, and "Fmoc Solid Phase Peptide Synthesis", Edited by W.C. Chan and P.D. White, Oxford University Press, 2000.

In one embodiment GLP-1 peptides may be produced by recombinant methods, viz. by culturing a host cell containing a DNA sequence encoding the GLP-1 peptide and capable of expressing the peptide in a suitable nutrient medium under conditions permitting the expression of the peptide. Non-limiting examples of host cells suitable for expression of these peptides are: *Escherichia coli*, *Saccharomyces cerevisiae*, as well as mammalian BHK or CHO cell lines.

In one embodiment GLP-1 peptides of the invention which include non-natural amino acids and/or a covalently attached N-terminal mono- or dipeptide mimetic may e.g. be produced as described in the experimental part. Or see e.g., Hodgson et al: "The synthesis of peptides and proteins containing non-natural amino acids", Chemical Society Reviews, 5 vol. 33, no. 7 (2004), p. 422-430; and WO 2009/083549 A1 entitled "Semi-recombinant preparation of GLP-1 analogues".

Salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid

The delivery agent used in the present invention is a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid. In some embodiments the delivery agent is an 10 absorption enhancer. The structural formula of N-(8-(2-hydroxybenzoyl)amino)caprylate is shown in formula (I).



In some embodiments the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is in 15 the caprylic acid form and/or the caprylate form. In some embodiments the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid comprises one monovalent cation, two monovalent cations or one divalent cation. In some embodiments the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is selected from the group consisting of the sodium salt, potassium salt and calcium salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

Salts of N-(8-(2-hydroxybenzoyl)amino)caprylate may be prepared using the method 20 described in e.g. WO96/030036, WO00/046182, WO01/092206 or WO2008/028859.

The salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid may be crystalline and/or 25 amorphous. In some embodiments the delivery agent comprises the anhydrate, monohydrate, dihydrate, trihydrate, a solvate or one third of a hydrate of the salt of N-(8-(2-hydroxybenzoyl)amino) caprylic acid as well as combinations thereof. In some embodiments the delivery agent is a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid as described in WO2007/121318. The salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid may be any polymorph thereof.

In some embodiments the delivery agent is sodium N-(8-(2-hydroxybenzoyl)amino)caprylate (referred to as "SNAC" herein), also known as sodium 8-(30 salicyloylamino) octanoate.

In some embodiments the amount of the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid in the composition is in the range of 0.6-3.5mmol. In some embodiments

the amount of the salt of N-(8-(2-hydroxybenzoyl) amino)caprylic acid in the composition is at least 0.6 mmol, such as selected from the group at least 0.8 mmol or at least 0.9 mmol. In some embodiments the amount of the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid in the composition is up to 2.5mmol. In some embodiments the amount of the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid in the composition is 0.6-2.0mmol. In some 5 embodiments the amount of the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is 1 mmol, such as 1.08 mmol.

In some embodiments the amount of SNAC in the composition is in the range of 100-1000 mg. In some embodiments the amount of SNAC in the composition is at least 150 10 mg or at least 250 mg. In some embodiments the amount of SNAC in the composition is up to 800 mg, such as up to 700 mg or up to 600 mg. In some embodiments the amount of SNAC in the composition is 300 mg.

In some embodiments the molar ratio between GLP-1 agonist (i.e. GLP-1 peptide) and delivery agent in the composition is less than 10, such as less than 5 or less than 1. In 15 some embodiments the molar ratio between GLP-1 agonist(i.e. GLP-1 peptide) and delivery agent in the composition is less than 1/10, such as less than 1/100 or less than 5/1000.

Pharmaceutical Indications

The present invention also relates to a composition or a granule of the invention for use as a medicament. In one embodiment the composition or the granule is administered 20 orally.

In particular embodiments, the composition or a granule of the invention may be used for the following medical treatments, all preferably relating one way or the other to diabetes:

(i) prevention and/or treatment of all forms of diabetes, such as hyperglycemia, type 25 2 diabetes, impaired glucose tolerance, type 1 diabetes, non-insulin dependent diabetes, MODY (maturity onset diabetes of the young), gestational diabetes, and/or for reduction of HbA1C;

(ii) delaying or preventing diabetic disease progression, such as progression in type 2 diabetes, delaying the progression of impaired glucose tolerance (IGT) to insulin requiring 30 type 2 diabetes, and/or delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes;

(iii) improving β -cell function, such as decreasing β -cell apoptosis, increasing β -cell function and/or β -cell mass, and/or for restoring glucose sensitivity to β -cells;

(iv) prevention and/or treatment of cognitive disorders;

(v) prevention and/or treatment of eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; and/or delaying

5 gastric emptying;

(vi) prevention and/or treatment of diabetic complications, such as neuropathy, including peripheral neuropathy; nephropathy; or retinopathy;

(vii) improving lipid parameters, such as prevention and/or treatment of dyslipidemia, lowering total serum lipids; lowering HDL; lowering small, dense LDL; lowering VLDL:

10 lowering triglycerides; lowering cholesterol; increasing HDL; lowering plasma levels of lipoprotein a (Lp(a)) in a human; inhibiting generation of apolipoprotein a (apo(a)) in vitro and/or in vivo;

(iix) prevention and/or treatment of cardiovascular diseases, such as syndrome X; atherosclerosis; myocardial infarction; coronary heart disease; stroke, cerebral ischemia; an early cardiac or early cardiovascular disease, such as left ventricular hypertrophy; coronary artery disease; essential hypertension; acute hypertensive emergency; cardiomyopathy; heart insufficiency; exercise tolerance; chronic heart failure; arrhythmia; cardiac dysrhythmia; syncopy; atherosclerosis; mild chronic heart failure; angina pectoris; cardiac bypass reocclusion; intermittent claudication (atherosclerosisoblitterens); diastolic dysfunction; and/or systolic dysfunction;

(ix) prevention and/or treatment of gastrointestinal diseases, such as inflammatory bowel syndrome; small bowel syndrome, or Crohn's disease; dyspepsia; and/or gastric ulcers;

(x) prevention and/or treatment of critical illness, such as treatment of a critically ill patient, a critical illness poly-nephropathy (CIPNP) patient, and/or a potential CIPNP patient; prevention of critical illness or development of CIPNP; prevention, treatment and/or cure of systemic inflammatory response syndrome (SIRS) in a patient; and/or for the prevention or reduction of the likelihood of a patient suffering from bacteraemia, septicaemia, and/or septic shock during hospitalisation; and/or

(xi) prevention and/or treatment of polycystic ovary syndrome (PCOS).

In a particular embodiment, the indication is selected from the group consisting of (i)-(iii) and (v)-(iix), such as indications (i), (ii), and/or (iii); or indication (v), indication (vi), indication (vii), and/or indication (iix).

5 In another particular embodiment, the indication is (i). In a further particular embodiment the indication is (v). In a still further particular embodiment the indication is (iix).

In one embodiment the invention relates to a composition or a granule of the invention for treatment of diabetes or obesity, wherein said granule is administered orally. In one embodiment the invention relates to a method for treatment of diabetes or obesity 10 comprising oral administration of a composition comprising a composition or a granule of the invention to a patient in need thereof.

The following indications are particularly preferred: Type 2 diabetes, and/or obesity.

PARTICULAR EMBODIMENTS

15 The following are particular embodiments of the invention

1. A pharmaceutical composition comprising a first type and a second type of granules, wherein said first type of granules comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, and wherein said second type of granules comprises a GLP-1 peptide.

20 2. A pharmaceutical composition according to any one of the preceding embodiments, wherein said first type of granules does not comprise GLP-1 peptide.

3. A pharmaceutical composition according to any one of the preceding embodiments, wherein said second type of granules does not comprise salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

25 4. A pharmaceutical composition according to any one of the preceding embodiments, wherein said first type of granules further comprises a lubricant, such as magnesium stearate.

5. A pharmaceutical composition according to any one of the preceding embodiments, wherein said first type of granules further comprises a filler, such as microcrystalline cellulose.

30 6. A pharmaceutical composition according to any one of the preceding embodiments, wherein said second type of granules further comprises a filler, such as microcrystalline cellulose.

7. A pharmaceutical composition according to any one of the preceding embodiments, wherein said second type of granules further comprises a binder, such as povidone.

8. A pharmaceutical composition according to any one of the preceding embodiments, wherein the granules are manufactured by dry granulation, such as by roller compaction.

9. A pharmaceutical composition according to any one of the preceding embodiments, wherein the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is

5 simultaneous with or faster than the release of said GLP-1 peptide.

10. A pharmaceutical composition according to any one of the preceding embodiments, wherein said the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and/or said GLP-1 peptide is determined by dissolution testing using Assay (I) and wherein said release is optionally determined within 30 minutes of said dissolution testing.

10 11. A pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is simultaneous with or faster than the release of said GLP-1 peptide as determined by dissolution testing using Assay (I).

15 12. A pharmaceutical composition according to any one of the preceding embodiments, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within the first 60 minutes as determined by dissolution testing using Assay (I) at pH 2.5.

13. A pharmaceutical composition according to any one of the preceding embodiments, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 55 minutes, such as within 50, 45, 40, 39, 38, 37, 36, 35, 34, 33, 32, or 31 minutes as 20 determined by dissolution testing using Assay (I) at pH 2.5.

14. A pharmaceutical composition according to any one of the preceding embodiments, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within the first 30 minutes as determined by dissolution testing using Assay (I) at pH 2.5.

15. A pharmaceutical composition according to any one of the preceding embodiments, 25 wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 31 minutes, such as within 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, 50, 55, or 60 minutes as determined by dissolution testing using Assay (I) at pH 2.5.

16. A pharmaceutical composition according to any one of the preceding embodiments, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks 30 within 30 minutes, such as within 25, 20, 15, 10, or 5 minutes as determined by dissolution testing using Assay (I) at pH 2.5.

17. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at any one of pH 1.0-8.0.

18. A pharmaceutical composition according to any one of the preceding embodiments, 35 wherein said dissolution testing using Assay (I) is carried out at pH 1.0.

19. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at pH 2.5.
20. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at pH 6.8.
- 5 21. A pharmaceutical composition according to any one of the preceding embodiments, wherein the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is SNAC.
22. A pharmaceutical composition according to any one of the preceding embodiments, wherein the GLP-1 peptide comprises an albumin binding moiety.
- 10 23. A pharmaceutical composition according to any one of the preceding embodiments, wherein the GLP-1 peptide is N-epsilon26-[2-(2-{2-[2-(2-{[(S)-4-carboxy-4-(17- carboxyheptadecanoylamino) butyrylamino]ethoxy}ethoxy)acetyl]amino]ethoxy}ethoxy)acetyl][Aib8,Arg34]GLP-1(7-37).
24. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition further comprises one or more pharmaceutically acceptable excipients.
- 15 25. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a solid dosage form.
26. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a tablet.
- 20 27. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a capsule.
28. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a sachet.
- 25 29. A pharmaceutical composition according to any one of the preceding embodiments further comprising any combination of the features according to embodiments 1-28.
- 30 30. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:
 - a) mixing a GLP-1 peptide with a filler and/or a binder;
 - b) dry granulation of the mixture of step a;
 - c) mixing the granules obtained in step b with a composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid; and
 - d) optionally adding further lubricant.

31. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

- 5 a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler;
- b) dry granulation of the mixture of step a;
- c) mixing the granules obtained in step b with a composition comprising a GLP-1 peptide; and
- d) optionally adding further lubricant.

10 32. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a GLP-1 peptide with a filler and/or a binder;
- b) dry granulation of the mixture of step a;
- 15 c) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, optionally a lubricant and/or a filler;
- d) dry granulation of the mixture of step c;
- e) mixing the granules obtained in step b with the granules obtained in step d; and
- 20 f) optionally adding further lubricant.

33. A process according to any one of embodiments 30-32, wherein said dry granulation is roller compaction.

34. A process according to any one of embodiments 30-33, wherein said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is SNAC.

25 35. A process according to any one of embodiments 30-34, wherein said GLP-1 peptide is semaglutide.

36. A process according to any one of embodiments 30-35, wherein said lubricant and/or said further lubricant is magnesium stearate.

37. A process according to any one of embodiments 30-36, wherein said filler is 30 microcrystalline cellulose.

38. A process according to any one of embodiments 30-37, wherein said binder is povidone.

39. A process according to any one of embodiments 30-38, wherein said mixing step comprising mixing of a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant

and/or a filler prior to dry granulation has a duration of at least 20 minutes, such as at least 30 minutes or at least 40 minutes, such as 50 minutes.

40. A process according to any one of embodiments 30-39, wherein said mixing comprising mixing of the first type of granules comprising a GLP-1 peptide with the second

5 type of granules comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid has a duration of at least 15 minutes or at least 20 minutes, such as 25 minutes or at least 30 minutes.

41. A pharmaceutical composition obtained by the process as defined in any one of embodiments 30-40.

10 42. A pharmaceutical composition as defined in any one of embodiments 1-29 or 41 for use in medicine.

43. A pharmaceutical composition as defined in any one of embodiments 1-29 or 41 for treatment of diabetes or obesity.

15 44. A pharmaceutical composition according to embodiment 42 or 43, wherein said pharmaceutical composition is administered orally.

45. A method for treatment of diabetes or obesity comprising administration of a composition comprising the pharmaceutical composition as defined in any one of embodiments 1-29 or 41 to a patient in need thereof.

46. A method according to embodiment 45, wherein said composition is administered 20 orally.

47. A granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, a lubricant, and a filler.

48. A granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a lubricant.

25 49. A granule comprising a GLP-1 peptide, a filler and a binder.

50. A granule according to any one of embodiments 47-49, wherein said lubricant is magnesium stearate.

51. A granule according to any one of embodiments 47-50, wherein said filler is microcrystalline cellulose.

30 52. A granule according to any one of embodiments 47-51, wherein said binder is povidone.

53. A granule according to any one of embodiments 47-52, wherein said granule is prepared according to the process as defined in any one of embodiments 30-40.

35 54. A process of producing a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, wherein the process comprises the steps:

- a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid (such as SNAC) with a lubricant and/or a filler; and
- b) dry granulation of the mixture of step a.

55. A process according to embodiment 54, wherein said salt of N-(2-

5 hydroxybenzoyl)amino)caprylic acid is SNAC.

56. A process according to embodiment 54 or 55, wherein said lubricant is magnesium stearate.

57. A process according to any one of embodiments 54-56, wherein said mixing step comprising mixing of a salt of N-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant 10 and/or a filler prior to dry granulation has a duration of at least 20 minutes, such as at least 30 minutes or at least 40 minutes, such as 50 minutes.

58. A process of producing a granule comprising a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a GLP-1 peptide, optionally a filler and/or a binder; and
- 15 b) roller compacting the mixture of step a.

59. A process according to embodiment 58, wherein said GLP-1 peptide is semaglutide.

60. A process according to any one of embodiments 58-59, wherein said binder is povidone.

61. A process according to any one of embodiments 58-60, wherein said filler is 20 microcrystalline cellulose.

62. A process according to any one of embodiments 58-61, wherein said dry granulation is roller compaction.

63. A granule obtained by the process as defined in any one of embodiments 58-62.

25 64. A method for increasing oral bioavailability of a GLP-1 peptide comprising administration to a patient in need thereof of a composition as defined in any of embodiments 1-29 or 41.

65. A pharmaceutical composition according to any one of the preceding embodiments 30 wherein said simultaneous or faster release is determined within 30 minutes, such as within 25, 20, 15 or 10 minutes, as determined by dissolution testing using Assay (I).

66. A pharmaceutical composition according to any one of the preceding embodiments wherein said simultaneous or faster release is determined within 30 minutes, such as within 25, 20, 15 or 10 minutes, as determined by dissolution testing using Assay (I) at pH 2.5.

FURTHER PARTICULAR EMBODIMENTS

The following are further particular embodiments of the invention:

1. A pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is simultaneous with or faster than the release of said GLP-1 peptide as determined by dissolution testing using Assay (I).
5
2. A pharmaceutical composition according to any one of the preceding embodiments, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within the first 30 minutes as determined by dissolution testing using Assay (I) at pH 2.5.
10
3. A pharmaceutical composition according to any one of the preceding embodiments, wherein the amount of dissolved salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within 31 minutes, such as within 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, 50, 55, or 60 minutes as determined by dissolution testing using Assay (I) at pH 2.5.
15
4. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at any one of pH 1.0-8.0.
5.
5. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at pH 1.0.
6.
6. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at pH 2.5.
20
7. A pharmaceutical composition according to any one of the preceding embodiments, wherein said dissolution testing using Assay (I) is carried out at pH 6.8.
8.
8. A pharmaceutical composition according to any one of the preceding embodiments, wherein the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is SNAC.
9.
9. A pharmaceutical composition according to any one of the preceding embodiments, wherein the GLP-1 peptide comprises an albumin binding moiety.
25
10. A pharmaceutical composition according to any one of the preceding embodiments, wherein the GLP-1 peptide is N-epsilon26-[2-(2-{2-[2-{2-[S]-4-carboxy-4-(17-carboxyheptadecanoylamino)butyrylamino]ethoxy}ethoxy)acetyl]amino]ethoxy)acetyl][Aib8,Arg34]GLP-1(7-37).
30
11. A pharmaceutical composition comprising a first type and a second type of granules, wherein said first type of granules comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and said second type of granules comprises a GLP-1 peptide.
12. A pharmaceutical composition according to embodiment 11, wherein said first type of granules further comprises a lubricant, such as magnesium stearate.

13. A pharmaceutical composition according to embodiment 11 or 12, wherein said first type of granules further comprises a filler, such as microcrystalline cellulose.

14. A pharmaceutical composition according to any one of embodiments 11-13, wherein said second type of granules further comprises a filler, such as microcrystalline cellulose.

5 15. A pharmaceutical composition according to any one of embodiments 11-14, wherein said second type of granules further comprises a binder, such as povidone.

16. A pharmaceutical composition according to any one of embodiments 11-15, wherein the granules are manufactured by dry granulation, such as by roller compaction.

10 17. A pharmaceutical composition according to any one of embodiments 11-16 further comprising any combination of the features according to embodiments 1-10.

18. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition further comprises one or more pharmaceutically acceptable excipients.

15 19. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a solid dosage form.

20. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a tablet.

21. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a capsule.

20 22. A pharmaceutical composition according to any one of the preceding embodiments, wherein said composition is in the form of a sachet.

23. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

25 a) mixing a GLP-1 peptide with a filler and/or a binder;
b) dry granulation of the mixture of step a;
c) mixing the granules obtained in step b with a composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid; and
d) optionally adding further lubricant.

30 24. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler;
35 b) dry granulation of the mixture of step a;

- c) mixing the granules obtained in step b with a composition comprising a GLP-1 peptide; and
- d) optionally adding further lubricant.

25. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-

5 hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a GLP-1 peptide with a filler and/or a binder;
- b) dry granulation of the mixture of step a;
- c) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, optionally a 10 lubricant and/or a filler;
- d) dry granulation of the mixture of step c;
- e) mixing the granules obtained in step b with the granules obtained in step d; and
- f) optionally adding further lubricant.

15 26. A process according to any one of embodiments 23-25, wherein said dry granulation is roller compaction.

27. A process according to any one of embodiments 23-26, wherein said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is SNAC.

28. A process according to any one of embodiments 23-27, wherein said GLP-1 peptide 20 is semaglutide.

29. A process according to any one of embodiments 23-28, wherein said lubricant and/or said further lubricant is magnesium stearate.

30. A process according to any one of embodiments 23-29, wherein said filler is microcrystalline cellulose.

25 31. A process according to any one of embodiments 23-30, wherein said binder is povidone.

32. A process according to any one of embodiments 23-31, wherein said mixing step comprising mixing of a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler prior to dry granulation has a duration of at least 30 minutes or at least 40 30 minutes, such as 50 minutes.

33. A process according to any one of embodiments 23-32, wherein said mixing comprising mixing of the first type of granules comprising a GLP-1 peptide with the second type of granules comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid has a duration of at least 15 minutes or at least 20 minutes, such as 25 minutes.

34. A pharmaceutical composition obtained by the process as defined in any one of embodiments 23-33.

35. A pharmaceutical composition as defined in any one of embodiments 1-22 or 34 for use in medicine.

5 36. A pharmaceutical composition as defined in any one of embodiments 1-22 or 34 for treatment of diabetes or obesity.

37. A pharmaceutical composition according to embodiment 35 or 36, wherein said pharmaceutical composition is administered orally.

38. A method for treatment of diabetes or obesity comprising administration of a 10 composition comprising the pharmaceutical composition as defined in any one of embodiments 1-22 or 34 to a patient in need thereof.

39. A method according to embodiment 38, wherein said composition is administered orally.

15 40. A granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, a lubricant, and a filler.

41. A granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a lubricant.

42. A granule comprising a GLP-1 peptide, a filler and a binder.

43. A granule according to any one of embodiments 40-42, wherein said lubricant is 20 magnesium stearate.

44. A granule according to any one of embodiments 40-43, wherein said filler is microcrystalline cellulose.

45. A granule according to any one of embodiments 40-44, wherein said binder is povidone.

25 46. A granule according to any one of embodiments 40-45, wherein said granule is prepared according to the process as defined in any one of embodiments 23-34.

47. A process of producing a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, wherein the process comprises the steps:

30 a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid (such as SNAC) with a lubricant and/or a filler; and

b) dry granulation of the mixture of step a.

48. A process according to embodiment 47, wherein said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is SNAC.

49. A process according to embodiment 47 or 48, wherein said lubricant is magnesium 35 stearate.

50. A process according to any one of embodiments 47-49, wherein said mixing step comprising mixing of a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler prior to dry granulation has a duration of at least 30 minutes or at least 40 minutes, such as 50 minutes.

5 51. A process of producing a granule comprising a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a GLP-1 peptide, optionally a filler and/or a binder; and
- b) roller compacting the mixture of step a.

52. A process according to embodiment 51, wherein said GLP-1 peptide is semaglutide.

10 53. A process according to any one of embodiments 51-52, wherein said binder is povidone.

54. A process according to any one of embodiments 47-53, wherein said filler is microcrystalline cellulose.

15 55. A process according to any one of embodiments 47-54, wherein said dry granulation is roller compaction.

56. A granule obtained by the process as defined in any one of embodiments 47-55.

57. A granule as defined in any one of embodiments 40-46 or 56 for use in medicine.

58. A granule as defined in any one of embodiments 40-46 or 56 for treatment of diabetes or obesity.

20 59. A granule according to embodiment 57 or 58, wherein said granule is administered orally in a pharmaceutical composition.

60. A method for treatment of diabetes or obesity comprising oral administration of a pharmaceutical composition comprising the granule as defined in any one of embodiments 40-46 or 56 to a patient in need thereof.

25 61. A method according to embodiment 60, wherein said pharmaceutical composition is administered orally.

EXAMPLES

Materials and Methods

General Methods of Preparation

Dry granulation

5 Dry granulation was carried out by roller compaction on a Gerteis MINI-PACTOR using smooth rolls and the settings listed in Table 1.

Table 1. Settings for dry granulation on the roller compactor

Parameter	Setting
Agitator speed	5.0 rpm
Roll speed	1.5 or 3.0 rpm
Gap	1.0 mm
Force	5.0 or 6.0 kN/cm
Granulator screen	0.63 mm wire screen
Granulator speed	60 rpm

10 Roller compaction force, i.e. the force between the rolls of the roller compactor when compacting materials into a continuous strip of compressed material, was determined by a pressure transducer that converts the hydraulic pressure into electrical signal; the roller compaction force may be measured in kiloNewton (kN) or in kiloNewton per roll width (kN/cm).

15 Subsequent to dry granulation comminution of the moldings into granules was carried out.

Tablet preparation

Tablets were produced on a Korsch PH106 or a Fette 102i mounted with a gravity feeder and a single set of punches resulting in 13 mm x 7.5 mm convex oval tablets having 20 no score. The press speed of the Korsch PH106 was set around 25 rpm and the counter pressure was adjusted to 40 kN. The press speed of the Fette 102i was set around 20 rpm. The fill volume was adjusted to obtain tablets having a target weight 407.7 mg. The compression force was set to obtain tablets with a crushing strength of around 180 ± 20 N for the Korsch PH106 and of around 128 N for the Fette 102i.

Resistance to crushing of tablets

Resistance to crushing of tablets was determined according to section 2.9.8 in the European Pharmacopoeia 7.5, 7th edition 2012 and at a jaw speed of 20 N/s.

General Methods of Detection and Characterisation5 *Assay (I): Dissolution testing*

The dissolution test was conducted with apparatus 2 in accordance with United States Pharmacopoeia 35 using a paddle rotation speed of 50 rpm. For testing at pH 1.0, 2.5 or 6.8, the 500 mL dissolution medium of 0.1 N hydrochloric acid (pH 1.0), 0.05 M phthalate buffer (pH 2.5), or 0.05 M phosphate buffer (pH 6.8), respectively, was used at a temperature of 37 °C. All dissolution media had a content of 0.1 % Tween80. Sample aliquots were removed at appropriate intervals and samples with acidic medium were neutralized with tribasic sodium phosphate to prevent precipitation. Sample contents were determined using a RP-HPLC method for dual detection of SNAC and GLP-1 (e.g. semaglutide) The HPLC method was based on gradient elution on a C8 column. The solvent system was trifluoroacetic acid and acetonitrile with UV detection at 210 and 335 nm. The sample contents were calculated based on the peak area of the SNAC and GLP-1 (e.g. semaglutide) peaks in the chromatogram relative to the peak areas of the SNAC and GLP-1 (e.g. semaglutide) references, respectively. The released amounts of SNAC and GLP-1 (e.g. semaglutide) were calculated as percentages of the nominal contents in the tablet i.e. 300 mg/tablet SNAC and 10 mg/tablet GLP-1 (e.g. semaglutide) and then optionally corrected for the actual contents in the tablets. The actual contents in the tablets were determined using Assay (III). Assay (II):

15 *Bioavailability in dogs*

20 Animals, Dosing and Blood Sampling: Male and female beagle dogs, weighing approx. 6-22 kg during the study period, were included in the study. The dogs were dosed in fasting state and the dogs were fed approx. 4 hours post dosing. The formulations were administered by oral administration to the dogs in groups of typically 8 (such as 4 male and 4 females).

25 The bioavailability studies were conducted either as single dose (SD) studies or multiple dose (MD) studies. In the MD studies the formulation was administered with five consecutive dosing occasions in each study (once a day dosing).

30 Blood samples were taken to cover the pharmacokinetic profile. An example of a SD blood sampling regimen could be the following time points: pre-dose, 0.25, 0.5, 0.75, 1,

1.5, 2, 2.5, 3, 4, 6, 8, 24, 48, 72, 96, 120, 144, 192 and 240 hours post dosing. An example of a MD blood sampling regimen could be the following time points: pre-dose, 0.5, 1.5 and 3 hr post dosing after each dosing occasion and a blood sampling regimen equivalent to a SD blood sampling regimen following the last dosing occasion.

5 Preparation of Plasma: All blood samples were collected into test tubes containing EDTA for stabilisation and kept on ice until centrifugation. Plasma was separated from whole blood by centrifugation and the plasma was stored at -20°C or lower until analysis.

10 Analysis of Plasma Samples: The plasma was analysed for semaglutide using a Luminescence Oxygen Channeling Immunoassay (LOCI). The LOCI assay employs donor beads coated with streptavidin and acceptor beads conjugated with a monoclonal antibody binding to a mid-molecular region of GLP-1 (e.g. semaglutide). The other monoclonal antibody, specific for an N-terminal epitope, was biotinylated. In the assay the three reactants were combined with the GLP-1 (e.g. semaglutide) which form a two-sited immuno-complex. Illumination of the complex releases singlet oxygen atoms from the donor beads which 15 channels into the acceptor beads and trigger chemiluminescence which was measured in the EnVision plate reader. The amount of light was proportional to the concentration of semaglutide and the lower limit of quantification (LLOQ) in plasma was 100 pM.

20 Pharmacokinetic Calculations: GLP-1 (e.g. semaglutide) plasma concentration data were subjected to non-compartmental pharmacokinetic analysis using the PC based software WinNonlin, v. 5.2 or later (Pharsight, Mountain View, CA. 94041, USA). For each individual dog the following pharmacokinetic parameters were estimated: Area Under the Curve (AUC), and dose normalized AUC (AUC/D). Bioavailability (F) was calculated as the fraction absorbed (in %) based on the dose normalised AUC (AUCinf./D) following oral and intravenous administration. Summary statistics of pharmacokinetic results were presented as 25 arithmetic mean.

Assay (III): Analysis of amount of GLP-1 and SNAC

For assay analysis tablets were dissolved using 0.05 M Na₂HPO₄ with 0.01% Tween20 as extraction buffer. Sample content was determined using a RP-HPLC method 30 for dual detection of SNAC and GLP-1 (e.g. semaglutide). The HPLC method was based on gradient elution on a C8 column. The solvent system was trifluoroacetic acid and acetonitrile with UV detection at 210 and 335 nm. The sample contents were calculated based on the peak area of the SNAC and GLP-1 (e.g. semaglutide) peaks in the chromatogram relative to

the peak areas of the SNAC and GLP-1 (e.g. semaglutide) references, respectively. The content was reported as average of 10 tablets.

Example 1: Preparation of tablet compositions comprising GLP-1 and SNAC

Tablet compositions comprising GLP-1 and SNAC were prepared with the components

5 shown in Table 2. Compound A is N-epsilon26-{2-[2-{2-[2-(2-({S})-4-carboxy-4-[10-(4-
carboxyphenoxy) decanoylamino]butyrylamino}ethoxy)ethoxy] acetylamino}ethoxy)
ethoxy]acetyl}, N-epsilon37-{2-[2-{2-[2-(2-({S})-4-carboxy-4-[10-(4-carboxy-phenoxy)
decanoylamino] butyrylamino}ethoxy)ethoxy]acetylamino}ethoxy) ethoxy]acetyl}-
[Aib8,Arg34,Lys37]GLP-1(7-37)-OH.

10

Table 2. Composition of the tablet compositions

Component	Amount (mg/tablet)	Function
Semaglutide or Compound A	10	Active ingredient
SNAC	300	Delivery agent
Microcrystalline cellulose (Avicel PH 101)	80	Filler
Povidone K 90 (Kollidon 90F)	8	Binder
Magnesium stearate	9.7	Lubricant
Total amount	407.7	

Tablet compositions were prepared by mixing the components listed in Table 2 in
15 different ways. The tablet compositions consisted of a first type of granules, in some cases a
second type of granules, as well as extragranular ingredients mixed with the first type of
granules and, if present, the second type of granules. Preparation of granules by roller
compaction and preparation of tablets from the tablet compositions were as described in the
section General Methods of Preparation. The design of the tablet compositions is shown in
20 Table 3.

Table 3. Design of tablet compositions (the amount of each component in mg/tablet is shown
in brackets)

<i>Tablet composition</i>	<i>Composition of first type of granules (mg/tablet)</i>	<i>Composition of second type of granules (mg/tablet)</i>	<i>Extrgranular ingredients (mg/tablet)</i>
B	SNAC (300), magnesium stearate (7.7)	semaglutide (10), microcrystalline cellulose (80), povidone (8)	magnesium stearate (2)
C	SNAC (300), semaglutide (10), magnesium stearate (7.7)	-	microcrystalline cellulose (80), povidone (8), magnesium stearate (2)
D	SNAC (300), semaglutide (10), povidone (8), magnesium stearate (7.7)	-	microcrystalline cellulose (80), magnesium stearate (2)
E	SNAC (300), semaglutide (10), microcrystalline cellulose (80), povidone (8), magnesium stearate (7.7)	-	magnesium stearate (2)
F	SNAC (300), microcrystalline cellulose (57), magnesium stearate (7.7)	semaglutide (10), microcrystalline cellulose (23), povidone (8)	magnesium stearate (2)
G	SNAC (300), magnesium stearate (7.7)	Compound A (10), microcrystalline cellulose (80), povidone (8)	magnesium stearate (2)
H	SNAC (300), microcrystalline cellulose (57), magnesium stearate (7.7)	Compound A (10), microcrystalline cellulose (23), povidone (8)	magnesium stearate (2)

Further details of the preparation of the tablet compositions are provided below.

Tablet composition B

Magnesium stearate for the first granule fraction was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a stainless steel bowl in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 5 60 s until the blend was visually homogenous. The remaining quantity of SNAC was transferred to a blender and was pre-mixed for 2 min at 25 rpm. The SNAC and magnesium stearate pre-mix was added to the blender and mixing was performed for 20 min at 25 rpm. The blend was roller compacted. The granules were sieved through a 180 µm mesh.

10 Semaglutide, microcrystalline cellulose and povidone for the second granule fraction were weighed directly into a stainless steel bowl in the order of decreasing amounts and mixed manually for at least 3 min until visually homogenous before transferring the pre-mix to a 1000 mL Duma bottle. The Duma bottle was closed with a lid and tumbled manually in a Turbula-like movement for 1 min. The blend was roller compacted.

15 The two types of granules were added to a blending container in order of decreasing content and mixed for 5 minutes at 32 rpm. Extragranular magnesium stearate was mixed with the granule blend by manual mixing using volume-doubling followed by 30 s mixing in the Turbula mixer at 32 rpm. Tablets were prepared from this composition.

Tablet composition C

20 Magnesium stearate for the granule fraction was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a stainless steel bowl in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 60 s until the blend was visually homogenous. The remaining quantity of SNAC was transferred to a blender and was pre-mixed for 2 min at 25 rpm. The SNAC and magnesium 25 stearate pre-mix was added to the blender and mixing was performed for 20 min at 25 rpm. Semaglutide was geometrically diluted using the mixed SNAC and magnesium stearate from the blender by mixing it manually for at least 60 s until visually homogenous. The pre-mix was then added to the mixed SNAC and magnesium stearate. The mixing step was finalized by mixing in the blender for 10 min at 25 rpm. The blend was roller compacted.

30 The granules and all other constituents except extragranular magnesium stearate were added to a blending container in order of decreasing content and mixed for 5 minutes at 32 rpm. Extragranular magnesium stearate was mixed with the granule blend by manual mixing using volume-doubling followed by 30 s mixing in the Turbula mixer at 32 rpm. Tablets were prepared from this composition.

Tablet composition D

Magnesium stearate for the granule fraction was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a stainless steel bowl in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 5 60 s until the blend was visually homogenous. The remaining quantity of SNAC was transferred to a blender and was pre-mixed for 2 min at 25 rpm. The SNAC and magnesium stearate pre-mix was added to the blender and mixing was performed for 20 min at 25 rpm. Semaglutide and povidone were weighed into a stainless steel bowl in the order of decreasing amounts and the components were mixed manually until visually homogenous. A 10 geometric dilution was then performed using the mixed SNAC and magnesium stearate from the blender by mixing it manually for at least 60 s until visually homogenous. The pre-mix was then added to the mixed SNAC and magnesium stearate. The mixing step was finalized in the blender by mixing for 10 min at 25 rpm. The blend was roller compacted.

The granules and microcrystalline cellulose were added to a blending container in 15 order of decreasing content and mixed for 5 minutes at 32 rpm. Extragranular magnesium stearate was mixed with the granule blend by manual mixing using volume-doubling followed by 30 s mixing in the Turbula mixer at 32 rpm. Tablets were prepared from this composition.

Tablet composition E

20 Magnesium stearate for the granule fraction was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a stainless steel bowl in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 60 s until the blend was visually homogenous. The remaining quantity of SNAC was transferred to a blender and was pre-mixed for 2 min at 25 rpm. The SNAC and magnesium 25 stearate pre-mix was added to the blender and mixing was performed for 20 min at 25 rpm. Semaglutide, microcrystalline cellulose and povidone were weighed into a stainless steel bowl in the order of decreasing amounts and the components were mixed manually until visually homogenous. A geometric dilution was then performed using the mixed SNAC and magnesium stearate from the blender by mixing it manually for at least 60 s until visually 30 homogenous. The pre-mix was then added to the mixed SNAC and magnesium stearate. The mixing step was finalized in the blender by mixing for 10 min at 25 rpm. The blend was roller compacted.

Extragranular magnesium stearate was mixed with the granules by manual mixing using volume-doubling followed by 30 s mixing in the Turbula mixer at 32 rpm. Tablets were 35 prepared from this composition.

Tablet composition F

Magnesium stearate for granule fraction one was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a stainless steel bowl in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 60 s until the blend was visually homogenous. The remaining quantity of SNAC was transferred to a blender and was pre-mixed for 2 min at 25 rpm. The SNAC and magnesium stearate pre-mix was added to the blender and mixing was performed for 20 min at 25 rpm. Microcrystalline cellulose was geometrically diluted using the mixed SNAC and magnesium stearate from the blender by mixing it manually for at least 60 s until visually homogenous. The pre-mix was then added to the mixed SNAC and magnesium stearate. The mixing step was finalized by mixing in the blender for 10 min at 25 rpm. The blend was roller compacted.

Semaglutide, microcrystalline cellulose (Avicel PH 101, FMC Biopolymer) and povidone (Kollidon 90F, BASF) for granule fraction two were weighed directly into a stainless steel bowl in the order of decreasing amounts and mixed manually for at least 3 min until visually homogenous before transferring the pre-mix to a 500 mL Duma bottle. The Duma bottle was closed with a lid and tumbled manually in a Turbula-like movement for 1 min. The blend was roller compacted.

The two types of granules were added to a blending container in order of decreasing content and mixed for 5 minutes at 32 rpm. Extragranular magnesium stearate was mixed with the granule blend by manual mixing using volume-doubling followed by 30 s mixing in the Turbula mixer at 32 rpm. Tablets were prepared from this composition.

Tablet composition G

Magnesium stearate for the first granule fraction was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a plastic bag in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 60 s. The SNAC and magnesium stearate pre-mix was added to the blender followed by the addition of the remaining SNAC and mixing was performed for 50 min at 25 rpm. The blend was roller compacted. The granules were sieved through a 1000 and 90 µm mesh.

Compound A, microcrystalline cellulose and povidone for the second granule fraction were weighed directly into a stainless steel bowl in the order of decreasing amounts and mixed manually for at least 3 min until visually homogenous before transferring the pre-mix to a 500 mL Duma bottle. The Duma bottle was closed with a lid and tumbled manually in a Turbula-like movement for 3 min. The blend was roller compacted.

The two types of granules were added to a blending container and mixed for 20 minutes at 25 rpm. Extragranular magnesium stearate was passed through a 355 µm sieve and mixed with the granule blend by manual mixing using volume-doubling followed by 2 min mixing in the Turbula mixer at 25 rpm. Tablets were prepared from this composition.

5

Tablet composition H

Magnesium stearate for granule fraction one was passed through a 355 µm sieve. Magnesium stearate was manually mixed with SNAC in a plastic bag in corresponding volumes. Two cycles of geometric dilution was applied by mixing for around 60 s. The SNAC and magnesium stearate pre-mix was added to the blender followed by the addition of the remaining SNAC and mixing was performed for 50 min at 25 rpm. Microcrystalline cellulose was then added to the mixed SNAC and magnesium stearate and the mixing step was finalized by mixing in the blender for 20 min at 25 rpm. The blend was roller compacted.

Compound A, microcrystalline cellulose (Avicel PH 101, FMC Biopolymer) and povidone (Kollidon 90F, BASF) for granule fraction two were weighed directly into a stainless steel bowl in the order of decreasing amounts and mixed manually for at least 3 min until visually homogenous before transferring the pre-mix to a 500 mL Duma bottle. The Duma bottle was closed with a lid and tumbled manually in a Turbula-like movement for at least 3 min. The blend was roller compacted.

The two types of granules were added to a blending container and mixed for 20 minutes at 25 rpm. Extragranular magnesium stearate was passed through a 355 µm sieve and mixed with the granule blend by manual mixing using volume-doubling followed by 2 minutes mixing in the Turbula mixer at 25 rpm. Tablets were prepared from this composition.

25 **Example 2: Dissolution of GLP-1 and SNAC from tablet composition B**

The dissolution of semaglutide and SNAC from tablet composition B was determined using Assay (I) described herein. The results are shown in Table 4. Corrected results are adjusted for the content of semaglutide or SNAC determined by analysis using Assay (III) described herein. The corrected results show that SNAC is released faster than semaglutide during the initial 20 min of dissolution. Furthermore, it shows that the difference between the faster release of SNAC compared to semaglutide is the largest in the dissolution medium with the pH-value of 2.5. Lastly, the data shows that in dissolution media with pH-values of 1.0 and 2.5 the amount of dissolved SNAC peaks within the initial 45 and 30 min, respectively.

Table 4. Dissolution of semaglutide and SNAC from tablet composition B

Dissolution time (min)	pH of dissolution media											
	1.0				2.5				6.8			
	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)
0	0	0	0	0	0	0	0	0	0	0	0	0
5	-	-	-	-	8	9	19	19	-	-	-	-
10	9	10	13	13	17	18	28	28	-	-	-	-
15	-	-	-	-	22	24	32	32	45	49	53	52
20	17	18	19	19	27	29	34	34	-	-	-	-
30	20	22	20	20	35	38	33	33	74	81	81	80
45	23	25	20	20	42	46	30	30	86	94	92	91
60	24	26	19	19	46	50	28	28	92	100	96	95
Corrected for actual content	Yes	No										

Example 3: Dissolution of GLP-1 and SNAC from tablet composition C

5 The dissolution of semaglutide and SNAC from tablet composition C was determined using Assay (I) described herein. The results are shown in Table 5. The results show that SNAC is released slower than semaglutide after the initial 5 min of dissolution in the dissolution media with low pH-values of 1.0 and 2.5. Corrected results are adjusted for the content of semaglutide or SNAC determined by analysis using Assay (III) described herein. During the initial 5 min of corrected dissolution in the dissolution medium with the pH-value of 2.5 SNAC is released equally fast assemaglutide. Furthermore, the corrected dissolution values show that SNAC is released faster than semaglutide in the dissolution medium with the pH-value of 6.8.

10

Table 5. Dissolution of semaglutide and SNAC from tablet composition C

Dissolution time (min)	pH of dissolution media									
	1.0		2.5		6.8					
	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)				
0	0	0	0	0	0	0	0	0	0	0
5	-	-	-	-	11	12	11	11	-	-
10	14	16	11	11	20	22	19	19	-	-
15	-	-	-	-	24	27	23	23	45	50
20	23	25	16	16	28	31	25	25	-	-
30	27	30	18	18	34	38	28	28	77	85
45	30	33	19	19	41	46	29	29	88	98
60	31	34	18	18	46	51	27	27	94	104
Corrected for actual content	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No

Example 4: Dissolution of GLP-1 and SNAC from tablet composition D

The dissolution of semaglutide and SNAC from tablet composition D was determined using Assay (I) described herein. The results are shown in Table 6. The results show that SNAC is released slower than semaglutide after the initial 5 min of dissolution in the dissolution media with low pH-values of 1.0 and 2.5. Corrected results are adjusted for the content of semaglutide or SNAC determined by analysis using Assay (III) described herein. During the initial 5 min of corrected dissolution in the dissolution medium with the pH-value of 2.5 SNAC is released equally fast with semaglutide. Furthermore, the corrected dissolution values show that SNAC is released faster than semaglutide in the dissolution medium with the pH-value of 6.8.

Table 6. Dissolution of semaglutide and SNAC from tablet composition D

Dissolution time (min)	pH of dissolution media					
	1.0		2.5		6.8	
	Semaglutide	SNAC	Semaglutide	SNAC	Semaglutide	SNAC

	released (% of total)		released (% of total)		released (% of total)		released (% of total)		released (% of total)		released (% of total)	
0	0	0	0	0	0	0	0	0	0	0	0	0
5	-	-	-	-	11	12	11	11	-	-	-	-
10	10	11	8	8	17	19	16	16	-	-	-	-
15	-	-	-	-	21	23	19	19	33	37	35	35
20	17	19	12	12	23	26	20	20	-	-	-	-
30	23	26	15	15	28	31	22	22	59	66	62	62
45	26	29	16	16	33	37	24	24	75	83	77	77
60	27	30	15	15	38	42	25	25	82	91	84	84
Corrected for actual content	Yes	No										

Example 5: Dissolution of GLP-1 and SNAC from tablet composition E

The dissolution of semaglutide and SNAC from tablet composition E was determined using Assay (I) described herein. The results are shown in Table 7. Corrected results are adjusted for the content of semaglutide or SNAC determined by analysis using Assay (III) described herein. The results show that SNAC is released slower than semaglutide during the entire dissolution time in a dissolution medium with a pH-value of 1.0. In a dissolution media with a pH-value of 2.5 the results show that SNAC initially is released faster than semaglutide during the initial 10 to 15 min of dissolution before becoming slower. Furthermore, the data shows that in dissolution medium with a pH-value of 2.5 the amount of dissolved SNAC peaks within the initial 20 min. Lastly, the corrected dissolution values show that SNAC is released faster than semaglutide in the dissolution medium with the pH-value of 6.8.

Table 7. Dissolution of semaglutide and SNAC from tablet composition E

Dissolution time (min)	pH of dissolution media					
	1.0		2.5		6.8	
	Semaglutide released (% of total)	SNAC released (% of	Semaglutide released (% of total)	SNAC released (% of	Semaglutide released (% of total)	SNAC released (% of

			total)				total)				total)	
0	0	0	0	0	0	0	0	0	0	0		0
5	-	-	-	-	21	23	22	22	-	-	-	-
10	15	17	12	12	38	42	40	40	-	-	-	-
15	-	-	-	-	50	56	46	46	47	53	51	51
20	23	26	16	16	57	64	43	43	-	-	-	-
30	28	31	16	16	63	71	37	37	76	85	79	79
45	32	36	17	17	68	76	31	31	86	96	89	89
60	36	40	17	17	70	78	28	28	90	101	93	93
Corrected for actual content	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No

Example 6: Dissolution of GLP-1 and SNAC from tablet composition F

The dissolution of semaglutide and SNAC from tablet composition F was determined using Assay (I) described herein. The results are shown in Table 8. Corrected results are 5 adjusted for the content of semaglutide or SNAC determined by analysis using Assay (III) described herein. The corrected results show that SNAC is released faster than semaglutide during the initial 20 to 30 min of dissolution. Furthermore, it shows that the difference between the faster release of SNAC compared to semaglutide is the largest in the dissolution medium with the pH-value of 2.5. Lastly, the data shows that in a dissolution medium with a 10 pH-value of 2.5 the amount of dissolved SNAC peaks within the initial 30 min.

Table 8. Dissolution of semaglutide and SNAC from tablet composition F

Dissolution time (min)	pH of dissolution media									
	1.0		2.5		6.8					
	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)	Semaglutide released (% of total)	SNAC released (% of total)				
0	0	0	0	0	0	0	0	0	0	0
5	-	-	-	-	9	10	17	17	-	-
10	8	9	10	10	20	22	31	31	-	-
15	-	-	-	-	28	31	40	40	51	56

20	12	13	13	13	33	36	43	43	-	-	-	-
30	15	17	15	15	38	42	42	42	75	82	79	79
45	19	21	16	16	43	47	39	39	85	93	88	88
60	21	23	16	16	48	53	38	38	89	98	93	93
Corrected for actual content	Yes	No										

Example 7: Bioavailability of GLP-1 in dogs from tablet compositions B-F

Bioavailability of GLP-1 from tablet compositions B-F was determined in dogs according to Assay (II) described herein. The results are shown in Table 9.

5

Table 9. Bioavailability of GLP-1 in dogs from tablet compositions B-F

Tablet composition	Bioavailability of GLP-1 in dogs
	(%F)
B	0.7
C	0.5
D	0.3
E	0.4
F	1.0

The results show that tablet composition F provided a bioavailability of 1.0%. The results show that tablet composition B provided a bioavailability of 0.7%. The results show 10 that tablet composition C provided a bioavailability of 0.5%. The results show that tablet composition E provided a bioavailability of 0.4%. The results show that tablet composition D provided a bioavailability of 0.3%.

Example 8: Dissolution of GLP-1 and SNAC from tablet composition G

15 The dissolution of Compound A and SNAC from tablet composition G was determined using Assay (I) described herein. The results are shown in Table 10. The results show that SNAC is released faster than Compound A during the initial 20 min of dissolution. Furthermore, it shows that the difference between the faster release of SNAC compared to Compound A is the largest in the dissolution medium with the pH-value of 2.5. Lastly, the

data shows that in dissolution media with pH-value of 2.5 the amount of dissolved SNAC peaks within the initial 45 min, respectively.

Table 10. Dissolution of Compound A and SNAC from tablet composition G

Dissolution time (min)	pH of dissolution media			
	2.5		6.8	
	Compound A released (% of total)	SNAC released (% of total)	Compound A released (% of total)	SNAC released (% of total)
0	0	0	0	0
5	9	17	-	-
10	16	24	-	-
15	21	26	38	49
20	25	28	-	-
30	32	28	70	79
45	39	28	85	92
60	43	26	91	96
120	-	-	95	101

5

Example 9: Dissolution of GLP-1 and SNAC from tablet composition H

The dissolution of Compound A and SNAC from tablet composition H was determined using Assay (I) described herein. The results are shown in Table 11. The results show that SNAC is released faster than Compound A during the initial 20 min of dissolution. Furthermore, it shows that the difference between the faster release of SNAC compared to Compound A is the largest in the dissolution medium with the pH-value of 2.5. Lastly, the data shows that in a dissolution medium with a pH-value of 2.5 the amount of dissolved SNAC peaks within the initial 30 min.

15 Table 11. Dissolution of Compound A and SNAC from tablet composition H

Dissolution time	pH of dissolution media	
	2.5	6.8

	Compound A released (% of total)	SNAC released (% of total)	Compound A released (% of total)	SNAC released (% of total)
0	0	0	0	0
5	10	19	-	-
10	19	31	-	-
15	26	37	37	47
20	31	38	-	-
30	36	34	63	75
45	42	29	74	83
60	46	27	79	87
120	-	-	89	97

Example 10: Bioavailability of GLP-1 in dogs from tablet compositions G-H

Bioavailability of GLP-1 from tablet compositions G-H was determined in dogs according to Assay (II) described herein. The results are shown in Table 12.

5

Table 12. Bioavailability of GLP-1 in dogs from tablet compositions G-H

Tablet composition	Bioavailability of GLP-1 in dogs
	(%F)
G	2.1
H	2.5

The results show that tablet composition G provided a bioavailability of 2.1%. The results show that tablet composition H provided a bioavailability of 2.5%.

10

While certain features of the invention have been illustrated and described herein, many modifications, substitutions, changes, and equivalents will now occur to those of ordinary skill in the art. It is, therefore, to be understood that the appended claims are intended to cover all such modifications and changes as fall within the true spirit of the invention.

15

CLAIMS

1. A pharmaceutical composition comprising a first type of granules and a second type of granules, wherein said first type of granules comprises a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and no GLP-1 peptide, and wherein said second type of granules comprises a GLP-1 peptide and no salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.
2. A pharmaceutical composition according to any one of the preceding claims, wherein the release of said salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is simultaneous with or faster than the release of said GLP-1 peptide as determined by dissolution testing using Assay (I) and wherein said release is optionally determined within 30 minutes of said dissolution testing.
3. A pharmaceutical composition according to any one of the preceding claims, wherein the dissolved amount of the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid peaks within the first 60 minutes, such as within 55, 50, 45, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, or 30 minutes, as determined by dissolution testing using Assay (I) at pH 2.5.
4. A pharmaceutical composition according to any one of the preceding claims, wherein said dissolution testing using Assay (I) is carried out at any one of pH 1.0-8.0, such as pH 1.0, 2.5, or 6.8.
5. A pharmaceutical composition according to any one of the preceding claims, wherein said first type of granules and said second type of granules comprises at least 50 % (w/w) of said composition.
6. A pharmaceutical composition according to any one of the preceding claims, wherein the salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid is SNAC.
7. A pharmaceutical composition according to any one of the preceding claims, wherein the GLP-1 peptide comprises an albumin binding moiety.
8. A pharmaceutical composition according to any one of the preceding claims, wherein the GLP-1 peptide is N-epsilon26-[2-(2-[2-[2-(2-[(S)-4-carboxy-4-(17-carboxy-

heptadecanoylamino) butyrylamino]ethoxy}ethoxy)acetylamino]ethoxy}ethoxy)acetyl] [Aib8,Arg34]GLP-1(7-37) or N-epsilon26-{2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-[10-(4-carboxyphenoxy) decanoylamino]butyrylamino]ethoxy}ethoxy] acetylamino}ethoxy) ethoxy]acetyl}, N-epsilon37-{2-[2-(2-{2-[2-(2-{(S)-4-carboxy-4-[10-(4-carboxy-phenoxy) decanoylamino] butyrylamino]ethoxy}ethoxy]acetylamino}ethoxy) ethoxy]acetyl}- [Aib8,Arg34,Lys37]GLP-1(7-37)-OH.

9. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a GLP-1 peptide with a filler and/or a binder;
- b) dry granulation of the mixture of step a;
- c) mixing the granules obtained in step b with a composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid; and
- 15 d) optionally adding further lubricant,

wherein the mixture of step a does not comprise a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid.

10. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid with a lubricant and/or a filler;
- b) dry granulation of the mixture of step a;
- 25 c) mixing the granules obtained in step b with a composition comprising a GLP-1 peptide; and
- d) optionally adding further lubricant

wherein the mixture of step a does not comprise a GLP-1 peptide.

30 11. A process of producing a pharmaceutical composition comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and a GLP-1 peptide, wherein the process comprises the steps:

- a) mixing a GLP-1 peptide with a filler and/or a binder;
- b) dry granulation of the mixture of step a;

- c) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, optionally a lubricant and/or a filler;
- d) dry granulation of the mixture of step c;
- e) mixing the granules obtained in step b with the granules obtained in step d;

5 and

- f) optionally adding further lubricant,

wherein the mixture of step a does not comprise a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid and wherein the mixture of step c does not comprise a GLP-1 peptide.

10

12. A pharmaceutical composition obtained by the process as defined in any one of claims 9-11.

15

13. A process of producing a granule comprising a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid, wherein the process comprises the steps:

- a) mixing a salt of N-(8-(2-hydroxybenzoyl)amino)caprylic acid (such as SNAC) with a lubricant and/or a filler; and
- b) dry granulation of the mixture of step a,

wherein the mixture of step a does not comprise a GLP-1 peptide.

20

14. A granule obtained by the process as defined in claim 13.

25

15. A composition as defined in any one of claims 1-8 or 12 or a granule as defined in claim 14 for use in medicine, such as for treatment of diabetes or obesity, wherein said composition is administered orally.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/055362

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K38/26 A61P3/10 A61K47/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, WPI Data, CHEM ABS Data, Sequence Search, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2009/124639 A1 (OYEWUMI MOSES [US] ET AL) 14 May 2009 (2009-05-14) example 8 -----	13,14
A,P	WO 2012/080471 A1 (NOVO NORDISK AS [DK]; SAUERBERG PER [DK]; BJERREGAARD SIMON [DK]; NIEL) 21 June 2012 (2012-06-21) the whole document -----	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
7 May 2013	15/05/2013
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Vandenbogaerde, Ann

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/EP2013/055362

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2009124639	A1	14-05-2009	NONE
WO 2012080471	A1	21-06-2012	NONE