



(86) Date de dépôt PCT/PCT Filing Date: 2002/05/06

(87) Date publication PCT/PCT Publication Date: 2002/12/05

(45) Date de délivrance/Issue Date: 2011/06/28

(85) Entrée phase nationale/National Entry: 2003/11/03

(86) N° demande PCT/PCT Application No.: EP 2002/005093

(87) N° publication PCT/PCT Publication No.: 2002/096292

(30) Priorité/Priority: 2001/05/30 (US60/294,461)

(51) Cl.Int./Int.Cl. *A61B 5/103* (2006.01),
A61B 10/00 (2006.01), *A61K 8/02* (2006.01),
A61Q 19/00 (2006.01), *A45D 34/00* (2006.01),
A45D 33/38 (2006.01), *A45D 44/00* (2006.01),
A45D 44/22 (2006.01)


(72) Inventeurs/Inventors:
TELESCA, JOSEPHINE, US;
MURRAY, LIAM ANTHONY, US;
GOTT, ROBERT EDWARD, US;
SLAVTCHEFF, CRAIG STEPHEN, US

(73) Propriétaire/Owner:
UNILEVER PLC, GB

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : BANDE ADHESIVE INDICATRICE DE RIDES ET PROCEDE D'UTILISATION DE CELLE-CI

(54) Title: WRINKLE INDICATOR TAPE STRIP AND METHOD OF USING THE SAME



(57) Abrégé/Abstract:

A test kit and method for visualizing fine lines and wrinkles is provided, the kit including a transparent strip (2) with an adhesive on one surface thereof and an imaging substrate with at least one darkened glossy area for receiving the transparent strip. Written instructions are provided in the kit. These advise a consumer to place the adhesive surface of the strip against a skin area requiring measurement. Thereafter the strip is removed and placed against the darkened area of the substrate. Topographical features of the skin can be viewed through the transparent strip with the transferred skin image onto the blackened background. A cosmetic anti-aging product can be applied to the skin over a period of time. Beneficial changes caused by the product are visualized through the test strip as periodic measurements are taken.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number
WO 02/096292 A1

(51) International Patent Classification⁷: **A61B 5/103**

(21) International Application Number: PCT/EP02/05093

(22) International Filing Date: 6 May 2002 (06.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/294,461 30 May 2001 (30.05.2001) US

(71) Applicant (for AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, OM, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZM, ZW only): **UNILEVER PLC** [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).

(71) Applicant (for AL, AM, AT, AZ, BA, BE, BF, BG, BJ, BR, BY, CF, CG, CH, CI, CM, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GA, GE, GN, GQ, GR, GW, HR, HU, ID, IS, IT, JP, KG, KP, KR, KZ, LR, LT, LU, LV, MA, MC, MD, MG, MK, ML, MR, MX, MZ, NE, NL, NO, PH, PL, PT, RO, RU, SE, SI, SK, SN, TD, TG, TJ, TM, TN, TR, UA, UZ, VN, YU only): **UNILEVER NV** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

(71) Applicant (for IN only): **HINDUSTAN LEVER LIMITED** [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, 400 020 Mumbai (IN).

(72) Inventors: **TELESCA, Josephine**; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT

06611 (US). **MURRAY, Liam, Anthony**; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). **GOTT, Robert, Edward**; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). **SLAVTCHEFF, Craig, Stephen**; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US).

(74) Agents: **ELLIOTT, Peter, William** et al.; Unilever PLC, Patent Department, Colworth House, Sharnbrook, Bedford, Bedfordshire MK44 1LQ (GB).


(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

[Continued on next page]

(54) Title: WRINKLE INDICATOR TAPE STRIP AND METHOD OF USING THE SAME



(57) Abstract: A test kit and method for visualizing fine lines and wrinkles is provided, the kit including a transparent strip (2) with an adhesive on one surface thereof and an imaging substrate with at least one darkened glossy area for receiving the transparent strip. Written instructions are provided in the kit. These advise a consumer to place the adhesive surface of the strip against a skin area requiring measurement. Thereafter the strip is removed and placed against the darkened area of the substrate. Topographical features of the skin can be viewed through the transparent strip with the transferred skin image onto the blackened background. A cosmetic anti-aging product can be applied to the skin over a period of time. Beneficial changes caused by the product are visualized through the test strip as periodic measurements are taken.

WO 02/096292 A1

WO 02/096292 A1



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WRINKLE INDICATOR TAPE STRIP AND METHOD OF USING THE SAME

The invention concerns a test strip for evaluating changes
5 in skin wrinkles, especially in the context of measuring the
efficacy of anti-aging cosmetic products.

A number of publications have disclosed test devices for the
lay person to self-diagnose their skin conditions. U.S.
10 Patent 3,571,947 (Maddison et al.) discloses a system for
identifying blemishes. A flexible, compliant film of plastic
is imprinted with pictorials of various types of common
blemishes. These reflect different dermal diseases. They
are cross-referenced with a handbook identifying the diseases
15 from the type of blemish. Cross-indexing treatments further
provides a suggested treatment to remedy the medical
condition.

U.S. Patent 5,727,949 (Bar-Or et al.) provides a dual ring
panel reference card. The panels are mounted for relative
20 movement whereby a selected diagnostic characteristic of a
skin problem can be aligned with a second diagnostic
characteristic and a determinable prognosis revealed from the
specific paired characteristics.

CuDerm Corporation has developed a simple diagnostic test to
25 determine the degree of skin dryness. CuDerm utilizes
adhesive discs (D-Squame) capable of removing a small section
of squameous cells (skin cells) and compares the results
against a chart. The disc is a transparent plastic with
adhesive on one side. The test involves placing the adhesive
30 surface of the disc against a user's cheek, peeling off the

- 2 -

disc and placing same on a dark background card. Flakes from the skin stick to the adhesive surface and are visualized against the dark background. Other than loose flakes, no topographical imprint is ever taken from the evaluated user's skin.

There are many cosmetic products sold which advertise certain skin benefits. Consumers usually cannot easily discern whether the claimed benefit is actually delivered. Even if perceivable, these actives impart an effect which may emerge only slowly over a period of time. Anti-aging actives are particularly illustrative. Facial fine lines and wrinkles can be minimized with actives such as alpha hydroxycarboxylic acids and/or retinol, to provide some visible improvement over an extended application period. They don't function instantaneously.

Accordingly, it is an advantage of the present invention to provide a low-cost simple test for a consumer to self-evaluate a cosmetic product's anti-aging benefits over a prolonged application period.

Another advantage of the present invention is to provide a low-cost simple self-evaluation tool for measuring changes in fine lines and wrinkles on the face or other aging susceptible parts of the human dermis.

- 3 -

In one aspect, the present invention provides a test kit for visualizing fine lines and wrinkles on a person's skin comprising:

- 5 (i) a transparent first strip provided with an adhesive on one surface thereof, the adhesive having sufficient tack to maintain an imprint of fine lines and wrinkles after removal of the strip from the skin;
- 10 (ii) an imaging substrate with at least one darkened area for receiving the transparent strip;
- (iii) a dusting powder for application against the skin prior to placement thereon of the adhesive surface of the
15 strip; and
- (iv) written instructions within the kit directing a consumer to follow a procedure in which the dusting powder is placed against the skin and the adhesive
20 surface of the strip is placed against a skin area treated with dusting powder and requiring measurement, the strip is removed and placed against the darkened area of the substrate, and the aforesaid procedure is repeated at a future time with a second strip followed by
25 comparison of patterns resultant from the first and second strip applications to the skin.

In another aspect of the invention there is provided a method for evaluating efficacy of an anti-aging cosmetic
30 product, the method comprising:

(A) providing a kit which comprises:

- 35 (i) a transparent first strip provided with an adhesive on one surface thereof, the adhesive having

- 3a -

sufficient tack to maintain an imprint of fine lines and wrinkles after removal of the strip from the skin;

(ii) a dusting powder; and

5

(iii) an imaging substrate with at least one darkened area for receiving the transparent strip;

(B) applying the cosmetic product to the skin;

10

(C) placing the dusting powder against the skin treated with the cosmetic product in step (B);

(D) placing the adhesive surface of the strip against

15

the skin treated with the dusting powder and the cosmetic product;

(E) removing the strip and placing same against one of the at least darkened areas of the substrate; and

20

(F) repeating steps (D) and (E) at a future time with a second strip followed by comparison of patterns resultant from the first and second strip applications to the skin.

- 4 -

Additional advantages, features and benefits of the present invention will become more readily apparent from consideration of the drawing in which:

5 Fig. 1 is a first embodiment of an application strip according to the present invention;

Fig. 2 is a second embodiment of an application strip according to the present invention; and

10 Fig. 3 is the application strip of the embodiment shown in Fig. 1 subsequent to being placed on the skin, removed therefrom and mounted on a darkened field reading card.

Now there has been found a simple diagnostic test for allowing a consumer to evaluate the claimed effectiveness of anti-aging cosmetic products. Effectiveness of the anti-aging result can be monitored over a period of time through an inexpensive kit. The kit employs a transparent plastic strip coated with a transparent adhesive layer. When applied to a wrinkle prone area of the face or body, the adhesive layer accepts a topological wrinkle imprint.

15

- 5 -

Removal of the strip from this wrinkle area can then be imaged by placement onto a darkened, preferably black field.

Fig. 1 illustrates a transparent strip 2 adhesively attached to a release backing 4. Strip 2 is kidney-shaped for
5 placement adjacent either the right or left eye so as to cover the periorbital canthus (crow's foot area). This curvilinear shape allows for maximum coverage around an outer corner of the eye.

A tab 6 is attached to the strip 2. The tab serves as a
10 gripping structure. Separation of the strip from the release backing is facilitated by initiating removal at the tab. The opaque, preferably black coloration of the tab in contrast to the transparency of the strip signals to a user the difference of this area and cues the user to start
15 lifting at that point.

Fig. 2 illustrates a second embodiment of a more elongate double lobed shape. Strip 2' is removably adhered onto a release backing 4'. Tab 6' is oriented between both lobes of the strip and lies along an axis of symmetry bisecting
20 the strip. The elongate nature of this embodiment even more than the first embodiment ensures that eyebrow hairs are not trapped under the adhesive when applied. It is undesirable to capture hairs. Any hairs caught in the adhesive may cause pain upon the strip being removed. This is considered
25 an undesirable ouch factor.

In the procedure for testing efficacy of various anti-aging products, the strip is removed from its release backing. Thereupon it is placed along an area of skin to be imaged for its topography. Facial areas are primarily intended for

- 6 -

evaluation, and more particularly areas surrounding the eye. Subsequently, the strip is removed and placed upon an imaging card 8. The dark, preferably black background of the card fixes the imprint while the transparent strip
5 allows a view of that imprint. Fig. 3 illustrates the strip showing fine lines and wrinkles 10 being visualized against the black background of the imaging card.

Subsequent to a baseline analysis of fine lines and wrinkles, treatment is begun with a selected cosmetic anti-
10 aging product. Treatment is continued for a period of time sufficient to allow the product to treat the signs of aging.

A second imaging field is placed adjacent to the first. After the treatment period of time, such as four weeks, another imprint is taken by a second transparent strip 21.
15 If the cosmetic product is properly functioning, fewer fine lines and wrinkles 11 will appear on the imaged second field. This procedure can then be repeated at six or eight weeks or at any further time interval. Each test will employ a fresh strip and new blackened area on the same or
20 another image card.

In a preferred embodiment, the kit includes a dusting device. Most preferred is a dusting paper which is formed of a cellulosic substrate supporting a water-dispersible titanium dioxide embedded therein. This device is available
25 from Leading Plus International, Taiwan. Prior to applying the adhesive transparent strip, the target area of the face is rubbed with the dusting paper. Powdered titanium dioxide is deposited thereon as an even film. Contact subsequently with the adhesive strip allows the latter to preferentially

- 7 -

adhere to powder deposit along ridges of the fine lines and wrinkles. An image in powder form of those fine lines and wrinkles is thereby obtained. Although a paper delivery system as described above is preferred, dusting powder can
5 also be delivered from a shaker container similar to those for the dispensing of talcum powder.

Strips for use in the present invention will be transparent articles allowing observation of any patterns on a lower surface thereof. Suitable materials for the strip are
10 plastics or cellulose of any variety which can be formed as transparent films. Typically the plastic may be selected from polyethylene, polypropylene, polystyrene, polyester, polycarbonate, polyacrylate, polyvinyl chloride, polyvinyl alcohol and polybutene. Not only homopolymers but
15 copolymers may be utilized for the strip material.

Copolymers may be formed from such monomers as C₂-C₁₀ olefins, vinyl chloride, acrylates and styrene constructed through free-radical polymerization. Condensation plastics may also be utilized in the formation of copolymers wherein
20 the monomers may be selected from C₂-C₁₀ dicarboxylic acids, C₂-C₁₀ polyols, C₂-C₆ alkoxylates and combinations thereof. Polyethylene, polypropylene and polyester terephthalate are the preferred plastic substrates for forming the strip.

The thickness of the strip may range anywhere from 0.001 to
25 2 mm, preferably from 0.01 to 1 mm, more preferably from 0.1 to 0.5 mm and optimally from 0.5 to 0.8 mm.

The backing is typically made from a material and in a manner that is generally impervious to the adhesive. The

- 8 -

backing may be elastic or non-elastic but preferably the former. Flexibility allows easier removal of the adhesive strip. The backing can be formed from a variety of materials including organic polymers and cellulose. A
5 release coating such as a silicone may be placed on an upper surface of the backing to ease removal of the adjacent adhesive strip.

The adhesive will be a pressure sensitive type preferably as a layer with an average thickness from 0.01 mm to 3 mm,
10 preferably from 0.05 mm to 2 mm, more preferably from 0.1 mm to 1 mm, optimally from 0.4 mm to 0.8 mm.

Pressure sensitive adhesives suitable for use in this invention are coatable adhesives. A wide variety of coatable pressure sensitive adhesives can be used, such as
15 solvent coatable, hot melt coatable, as well as latex PSA's that are coatable out of water. Also, solventless curable adhesives (often referred to as 100% solids) can be used. Where thicker adhesive coatings are desired, it may be desirable either to apply multiple layers of the adhesive,
20 hot melt coat, or to photopolymerize the adhesive in situ. Specific examples of pressure sensitive adhesives include acrylates, such as isooctyl acrylate/acrylic acid copolymers, tackified acrylates, and plasticizer-containing acrylates such as those disclosed in U.S. Pat. No. 4,946,742
25 (Landin); natural or synthetic rubber resins, including thermoset rubbers as well as thermoplastic rubbers and elastomers, such as nitrile rubbers (e.g., acrylonitrile-butadiene), styrene-butadiene, styrene-isoprene, styrene-butadiene-styrene, styrene-isoprene-styrene, and natural

- 9 -

rubber; silicone-based adhesives, such as polysiloxanes; polyolefins; polyesters; polyamides; and polyurethanes.

Particularly preferred are the acrylic type pressure sensitive adhesives. Most especially a pressure sensitive adhesive with a low tack value. These materials are commercially available under the Flexcon[®] brand.

Relative thickness of the strip to the adhesive may range from 1:200 to 200:1, preferably from 1:10 to 10:1, optimally from 2:1 to 1:2. Relative weight ratio of the strip to the adhesive may range from 1:200 to 200:1, preferably from 1:10 to 10:1, optimally from 2:1 to 1:2.

Anti-aging cosmetic products of this invention may contain one or more anti-aging actives and a cosmetic carrier. Illustrative actives are alpha- and beta-hydroxyacids, retinoids (e.g. retinol and retinyl palmitate), ascorbic acid and derivatives (e.g. ascorbyl tetraisoalmitate, magnesium ascorbyl phosphate), lipoic acid, green tea, tocopherol and derivatives, dihydroepiandrosterone (DHEA) and combinations thereof. Amounts may range from 0.00001 to 10% by weight of the product. Carriers may include water, silicones, natural and synthetic esters (e.g. triglycerides, lanolin and fatty acid esters), hydrocarbons, propellants, thickeners, surfactants and combinations thereof. Amounts may range from 5 to 99.9% by weight of the product.

Anti-aging cosmetic products may take various forms including creams, lotions, wipes, aerosols, powders and transdermal patches.

- 10 -

Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in this description indicating amounts of material ought to be understood as modified by the word "about".

- 5 The term "comprising" is meant not to be limiting to any subsequently stated elements but rather to encompass non-specified elements of major or minor functional importance. In other words the listed steps, elements or options need not be exhaustive. Whenever the words "including" or
- 10 "having" are used, these terms are meant to be equivalent to "comprising" as defined above.

All parts, percentages and proportions referred to herein and in the appended claims are by weight unless otherwise illustrated.

Claims:

1. A test kit for visualizing fine lines and wrinkles on a person's skin comprising:

(i) a transparent first strip provided with an adhesive on one surface thereof, the adhesive having sufficient tack to maintain an imprint of fine lines and wrinkles after removal of the strip from the skin;

(ii) an imaging substrate with at least one darkened area for receiving the transparent strip;

(iii) a dusting powder for application against the skin prior to placement thereon of the adhesive surface of the strip; and

(iv) written instructions within the kit directing a consumer to follow a procedure in which the dusting powder is placed against the skin and the adhesive surface of the strip is placed against a skin area treated with dusting powder and requiring measurement, the strip is removed and placed against the darkened area of the substrate, and the aforesaid procedure is repeated at a future time with a second strip followed by comparison of patterns resultant from the first and second strip applications to the skin.

2. The kit according to claim 1 wherein the adhesive is a pressure sensitive adhesive.

3. The kit according to claim 2 wherein the adhesive is an acrylate polymer.

4. A method for evaluating efficacy of an anti-aging cosmetic product, the method comprising:

(A) providing a kit which comprises:

(i) a transparent first strip provided with an adhesive on one surface thereof, the adhesive having sufficient tack to maintain an imprint of fine lines and wrinkles after removal of the strip from the skin;

(ii) a dusting powder; and

(iii) an imaging substrate with at least one darkened area for receiving the transparent strip;

(B) applying the cosmetic product to the skin;

(C) placing the dusting powder against the skin treated with the cosmetic product in step (B);

(D) placing the adhesive surface of the strip against the skin treated with the dusting powder and the cosmetic product;

(E) removing the strip and placing same against one of the at least darkened areas of the substrate; and

(F) repeating steps (D) and (E) at a future time with a second strip followed by comparison of patterns resultant from the first and second strip applications to the skin.

5. The method according to claim 4 wherein the dusting powder is carried to the skin on a paper.

6. The method according to claim 4 wherein the dusting powder is a water-dispersible titanium dioxide.

7. The kit according to claim 1 wherein the dusting powder is carried to the skin on a paper.

8. The kit according to claim 1 wherein the dusting powder is a water-dispersible titanium dioxide.

1/1

Fig.1.




Fig.2.




Fig.3.

