



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/12/12
(87) Date publication PCT/PCT Publication Date: 2018/06/21
(85) Entrée phase nationale/National Entry: 2019/06/05
(86) N° demande PCT/PCT Application No.: US 2017/065864
(87) N° publication PCT/PCT Publication No.: 2018/111904
(30) Priorité/Priority: 2016/12/12 (US62/433,113)


(51) Cl.Int./Int.Cl. *C07D 403/06* (2006.01),
A61K 31/415 (2006.01), *C07D 405/06* (2006.01),
C07D 405/14 (2006.01)

(71) Demandeur/Applicant:
VETTORE, LLC, US

(72) Inventeurs/Inventors:
PARNELL, KENNETH MARK, US;
MCCALL, JOHN M., US;
ROMERO, DONNA, US

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : INHIBITEURS HETEROCYCLIQUES DE MCT4
(54) Title: HETEROCYCLIC INHIBITORS OF MCT4



(57) **Abrégé/Abstract:**

Disclosed herein are compounds and compositions useful in the treatment of MCT4 mediated diseases, such as proliferative and inflammatory diseases, having the structure of Formula (I). Methods of inhibition MCT4 activity in a human or animal subject are also provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2018/111904 A1

(43) International Publication Date
21 June 2018 (21.06.2018)

(51) International Patent Classification:

C07D 403/06 (2006.01) *C07D 405/14* (2006.01)
A61K 31/415 (2006.01) *C07D 405/06* (2006.01)

(21) International Application Number:

PCT/US2017/065864

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(22) International Filing Date:

12 December 2017 (12.12.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/433,113 12 December 2016 (12.12.2016) US

(71) Applicant: VETTORE, LLC [US/US]; 1700 Owens St., Suite 515, San Francisco, CA 94110 (US).


(72) Inventors: PARNELL, Kenneth, Mark; 1700 Owens Street, Suite 515, San Francisco, CA 94110 (US). MC-CALL, John, M.; 1700 Owens Street, Suite 515, San Francisco, CA 94110 (US). ROMERO, Donna; 1700 Owens Street, Suite 515, San Francisco, CA 94110 (US).

(74) Agent: HATHAWAY, Cynthia; Global Patent Group, LLC, 17014 New College Avenue, Suite 201, St. Louis, MO 63040 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: HETEROCYCLIC INHIBITORS OF MCT4



(57) Abstract: Disclosed herein are compounds and compositions useful in the treatment of MCT4 mediated diseases, such as proliferative and inflammatory diseases, having the structure of Formula (I). Methods of inhibition MCT4 activity in a human or animal subject are also provided.

WO 2018/111904 A1

HETEROCYCLIC INHIBITORS OF MCT4

[001] This application claims the benefit of priority of United States Provisional Application No. 62/433,113, filed December 12, 2016, the contents of which are incorporated by reference as if written herein in its entirety.

[002] Disclosed herein are new heterocyclic compounds and compositions and their application as pharmaceuticals for the treatment of disease. Methods of inhibition of MCT4 activity in a human or animal subject are also provided for the treatment diseases such as cancer.

[003] Lactic acid export from glycolytic cells is typically mediated by the monocarboxylate transporter MCT4. MCT4 exhibits weak affinity for lactate ($K_m = 28$ mM) coupled with a high turnover rate, allowing rapid export of large amounts of lactic acid. MCT4 expression is normally limited to highly glycolytic tissues such as white muscle fibers, lymphocytes, astrocytes, and Sertoli cells. Though MCT4 is absent from most normal tissues, MCT4 expression is highly upregulated, and correlates with poor survival, in many cancer indications, including colorectal cancer, glioma, head and neck cancer, triple-negative breast cancer, prostate cancer, KRAS mutant lung cancer, liver cancer, and kidney cancer.

[004] The correlation of MCT4 expression and poor cancer outcome appears to be of significant functional consequence in multiple cancer models. Stable expression of MCT4 is highly tumorigenic in a respiration-impaired, Ras-transformed fibroblast xenograft model. Conversely, MCT4 silencing slows or ablates tumor growth in xenograft models of breast cancer, colorectal cancer, and glioma. MCT4 expression is required for inflammatory cytokine IL-8-mediated angiogenesis in breast and colon cancer xenograft models. MCT4 has also been shown to play important roles in cancer cell migration, invasion, and various aspects of the Warburg effect (e.g., proliferation on glucose, extracellular acidification, and lactate secretion).


[005] Inhibition of MCT4-mediated lactic acid export may be an effective strategy to impair the Warburg effect in cancer. Unfortunately, no potent and selective MCT4 inhibitors have been described. Moderate to weak MCT4 inhibitors are known (e.g., phloretin and α -CN-4-OH-cinnamate); however, these compounds promiscuously inhibit a number of other transporters, including MCT1.

[006] Thus, there is a need for potent and selective MCT4 inhibitors for use in the treatment or prevention of cancer.

DETAILED DESCRIPTION

[007] Novel compounds and pharmaceutical compositions, certain of which have been found to inhibit MCT4 have been discovered, together with methods of synthesizing and using the compounds including methods for the treatment of MCT4-mediated diseases in a patient by administering the compounds.


[008] Provided is a compound of structural Formula I



or a salt thereof, wherein:

A^1 , A^2 , and A^3 are independently chosen from N and C, wherein at least one of A^1 , A^2 , and A^3 is N;

L is chosen from a bond, methylene, and ethylene;



X is H or null, or is chosen from alkenylamino, alkyl, aminoalkenyl, and aminoalkyl, any of which is optionally substituted with one to three R^1 groups;

Y is chosen from alkenyl, alkenylamino, alkyl, aminoalkenyl, aminoalkyl, aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R^2 groups, or

X and Y together with the atoms to which they are attached may form an aryl, cycloalkyl, heteroaryl, or heterocycloalkyl ring, any of which is optionally substituted with one to three R^7 groups; and

Z is chosen from alkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl, any of which is optionally substituted with one to three R^3 groups;

each R^1 is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, heteroaryl;

each R² is independently chosen from alkyl, alkenyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, or

two R², together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

each R³ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R⁸ groups, or

two R³, together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R⁴ and R⁵ are independently chosen from H and C₁-C₆alkyl, wherein R⁴ and R⁵ together comprise no more than 6 carbons, or

R⁴ and R⁵, together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring;

R⁶ is chosen from H and C₁-C₄alkyl;

each R⁷ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl; and

each R⁸ is independently chosen from cyano, halo, hydroxy, and oxo.

[009] Certain compounds disclosed herein may possess useful MCT4 inhibiting activity, and may be used in the treatment or prophylaxis of a disease or condition in which MCT4 plays an active role. Thus, in broad aspect, certain embodiments also provide pharmaceutical compositions comprising one or more compounds disclosed herein together with a pharmaceutically acceptable carrier, as well as methods of making and using the compounds and compositions. Certain embodiments provide methods for inhibiting MCT4. Other embodiments provide methods for treating a MCT4-mediated disorder in a patient in need of such treatment, comprising administering to said patient a therapeutically effective amount of a compound or composition according to the present invention. Also provided is the use of certain compounds disclosed herein for use in the manufacture of a medicament for the treatment of a disease or condition ameliorated by the inhibition of MCT4.

[010] In certain embodiments, X is chosen from alkenylamino, alkyl, aminoalkenyl, aminoalkyl, and H; and Y is chosen from alkenyl, alkenylamino, alkyl, aminoalkenyl, aminoalkyl, aryl, cycloalkyl, and heteroaryl.



[011] In certain embodiments, X and Y together with the atoms to which they are attached may form an aryl, cycloalkyl, heteroaryl, or heterocycloalkyl ring.




[012] In certain embodiments, exactly two of A^1 , A^2 , and A^3 is N.



[013] In certain embodiments, exactly one of A^1 , A^2 , and A^3 is N.


[014] In certain embodiments, A^1 and A^2 are C; and A^3 is N.

[015] In certain embodiments, X is hydrogen.

[016] In certain embodiments, W is chosen from , ,

, , and .

[017] In certain embodiments, W is chosen from  and .

[018] In certain embodiments, W is .

[019] In certain embodiments, Y is chosen from aryl and heteroaryl, any of which is optionally substituted with one to three R^2 groups.

[020] In certain embodiments, Y is chosen from phenyl, thienyl, thiazolyl, quinoliny, isoquinoliny, cinnoliny, indazolyl, and indolyl.

[021] In certain embodiments, Y is chosen from quinoliny, isoquinoliny, cinnoliny, indazolyl, and indolyl.

[022] In certain embodiments, Y is chosen from phenyl, thienyl, and thiazolyl.


[023] In certain embodiments, Z is chosen from phenyl, pyridinyl, quinoliny, isoquinoliny, cinnoliny, indazolyl, and indolyl.

[024] In certain embodiments, Z is chosen from phenyl and pyridinyl.

[025] In certain embodiments, each R^2 is independently chosen from alkenyl, alkoxy, alkoxyalkyl, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, aryl, halo, heteroaryl, and haloalkyl.

[026] In certain embodiments, each R^2 is independently chosen from alkoxy, alkoxyalkyl, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, halo, and haloalkyl.

- [027] In certain embodiments, two R^2 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring.
- [028] In certain embodiments, each R^3 is independently chosen from alkenyl, alkoxy, alkyl, aryl, halo, heteroaryl, and haloalkyl.
- [029] In certain embodiments, each R^3 is independently chosen from alkoxy, alkyl, aryl, and heteroaryl.
- [030] In certain embodiments, two R^3 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring.
- [031] In certain embodiments, each R^3 is alkoxy.
- [032] In certain embodiments, each R^3 is methoxy.
- [033] In certain embodiments, each R^3 is halo.
- [034] In certain embodiments, each R^3 is independently selected from F, Br, and I.
- [035] In certain embodiments, each R^3 is F.
- [036] In certain embodiments, R^4 and R^5 are independently chosen from H and C₁-C₂alkyl.
- [037] In certain embodiments, R^4 and R^5 are independently chosen from H and methyl.
- [038] In certain embodiments, R^4 and R^5 , together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring.
- [039] In certain embodiments, R^6 is chosen from H and C₁-C₂alkyl.
- [040] In certain embodiments, R^6 is chosen from H and methyl.
- [041] In certain embodiments, each R^7 is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl.
- [042] In certain embodiments, each R^7 is independently chosen from alkyl and alkoxy.
- [043] In certain embodiments, the compounds have structural formula Ia:



or a salt or tautomer thereof, wherein:

A^5 is chosen from C-X and N;

A^6 is chosen from C-Y and N;


and A⁷ are and A⁸ are chosen from the following combinations:

A⁷ is C and A⁸ is NH, or

A⁷ is N and A⁸ is N;

wherein at least one of A⁵, A⁶, and A⁷ is N;

L is chosen from a bond and methylene;



X is H, or is chosen from alkenylamino, alkyl, aminoalkenyl, and aminoalkyl, any of which is optionally substituted with one to three R¹ groups;

Y is chosen from alkenyl, alkenylamino, alkyl, aminoalkenyl, aminoalkyl, aryl, cycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R² groups, or

X and Y together with the atoms to which they are attached may form an aryl, cycloalkyl, heteroaryl, or heterocycloalkyl ring, any of which is optionally substituted with one to three R⁷ groups; and

Z is chosen from alkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl, any of which is optionally substituted with one to three R³ groups;

each R¹ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, heteroaryl;

each R² is independently chosen from alkyl, alkenyl, alkoxy, alkoxyalkyl, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl;


each R³ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl;

R⁴ and R⁵ are independently chosen from H and C₁-C₆alkyl, wherein R⁴ and R⁵ together comprise no more than 6 carbons;

R⁶ is chosen from H and C₁-C₄alkyl; and


each R⁷ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl.

[044] In certain embodiments, the compounds have structural Formula II:



or a salt thereof, wherein:

L is chosen from a bond and methylene;



Y is chosen from aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R^2 groups;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^3 groups;

each R^2 is independently chosen from alkyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, or

two R^2 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

each R^3 is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R^8 groups, or

two R^3 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R^4 and R^5 are independently chosen from H and C_1 - C_6 alkyl, wherein R^4 and R^5 together comprise no more than 6 carbons and wherein at least one of R^4 and R^5 is C_1 - C_6 alkyl, or

R^4 and R^5 , together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring;

R^6 is chosen from H and C_1 - C_4 alkyl; and

each R^8 is independently chosen from cyano, halo, hydroxy, and oxo.

[045] In certain embodiments of structural formula II,

L is a bond; and

each R^2 is chosen from alkenyl, C_2 - C_6 alkoxy, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, aryl, halo, heteroaryl, and haloalkyl.

[046] In certain embodiments of structural formula II,

Z is chosen from aryl and heteroaryl, either of which is substituted with one to three R^3 groups;

each R^2 is chosen from alkenyl, C_2 - C_6 alkoxy, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, aryl, heteroaryl, and haloalkyl; and

each R^3 is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, cycloalkyl, aryl, and heteroaryl.

[047] In certain embodiments of structural formula II,

Y is heteroaryl, and is substituted with one to three R^2 groups; and

each R^2 is chosen from alkenyl, C_2 - C_6 alkoxy, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, aryl, halo, heteroaryl, and haloalkyl.

[048] In certain embodiments of structural formula II,


Y is selected from the group consisting of furyl, thienyl, pyrrolyl, imidazolyl, and oxazolyl, any of which is substituted with one to three R^2 groups; and

each R^2 is chosen from alkenyl, C_2 - C_6 alkoxy, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, aryl, halo, heteroaryl, and haloalkyl.

[049] In certain embodiments of structural formula II, at least one of R^4 and R^5 is H.

[050] In certain embodiments of structural formula II, exactly one of R^4 and R^5 is H.

[051] In certain embodiments, the compounds have structural Formula III:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one or two R³ groups;

R^{2a} and R^{2b} are independently chosen from H, alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, halo, cycloalkyl, and heterocycloalkyl;

each R³ is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, amino, aminoalkyl, carboxamido, halo, cyano, hydroxy, and cycloalkyl, and is optionally substituted with one or two R⁸ groups;

R⁴ and R⁵ are independently chosen from H and C₁-C₆alkyl, wherein R⁴ and R⁵ together comprise no more than 6 carbons and wherein at least one of R⁴ and R⁵ is C₁-C₆alkyl, or

R⁴ and R⁵, together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring;

R⁶ is chosen from H and methyl; and

each R⁸ is independently chosen from cyano, halo, hydroxy, and oxo.

[052] In certain embodiments of structural formula III,

Z is chosen from quinolinyl, isoquinolinyl, cinnolinyl, indazolyl, and indolyl, any of which is optionally substituted with one to three R³ groups; and

each R³ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl.

[053] In certain embodiments of structural formula III, Z is chosen from alkyl, cycloalkyl, and heterocycloalkyl, any of which is optionally substituted with one to three R³ groups;

[054] In certain embodiments of structural formula III, R² is chosen from alkenyl, C₂-C₆alkoxy, cycloalkoxy, cycloalkylmethoxy, haloalkoxy, alkyl, aryl, halo, heteroaryl, and haloalkyl.

[055] In certain embodiments of structural formula III, R² is selected from alkoxy, alkyl, halo, haloalkyl, and haloalkoxy.

[056] In certain embodiments of structural formula III, each R³ is independently chosen from alkenyl, alkoxy, alkyl, amino, aryl, halo, heteroaryl, and haloalkyl; and

[057] In certain embodiments of structural formula III, Z is chosen from alkyl, cycloalkyl, and heterocycloalkyl, any of which is optionally substituted with one to three R³ groups;

[058] In certain embodiments of structural formula III,

L is a bond;

Z is chosen from aryl and heteroaryl, either of which is substituted with one to three R^3 groups; and

each R^3 is independently chosen from alkenyl, alkoxy, alkyl, amino, fluoro, bromo, and iodo.

[059] In certain embodiments of structural formula III,

L is methylene;

Z is chosen from aryl and heteroaryl, either of which is substituted with one to three R^3 groups; and

each R^3 is independently chosen from alkenyl, C_2 - C_6 alkoxy, C_4 - C_8 alkyl, aryl, and iodo.

[060] In certain embodiments of structural formula III,

L is a bond; and

Z is chosen from pyridin-3-yl and pyridin-4-yl, and is optionally substituted with one to three R^3 groups.

[061] In certain embodiments of structural formula III,

L is methylene; and


Z is pyridyl, and is optionally substituted with one to three R^3 groups.

[062] In certain embodiments of structural formula III,

L is methylene; and


Z is pyridyl, and is optionally substituted with two or three R^3 groups.

[063] In certain embodiments, the compounds have structural Formula IV:



or a salt thereof, wherein:

L is chosen from a bond and methylene;



Z is chosen from alkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl, any of which is optionally substituted with one to three R³ groups;


each R³ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl;

R⁴ and R⁵ are independently chosen from H and C₁-C₆alkyl, wherein R⁴ and R⁵ together comprise no more than 6 carbons;

R⁶ is chosen from H and C₁-C₄alkyl; and

n is selected from 1, 2, and 3.

[064] In certain embodiments, the compounds have structural Formula V:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

Y is chosen from phenyl, thienyl, furyl, thiazolyl, and oxazolyl, any of which is optionally substituted with one or two R² groups, or

each R² is independently chosen from alkoxy, alkylthio, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, and halo;


R^{3a} is chosen from H, alkoxy, haloalkoxy, cycloalkoxy, amino, and halo;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl, or

R⁴ and R⁵, together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring; and

R⁶ is chosen from H and C₁-C₄alkyl.

[065] In certain embodiments, the compounds have structural Formula VI:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

R^{2a} is chosen from alkoxy, alkylthio, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, and halo;


R^{3a} is chosen from H, alkoxy, haloalkoxy, cycloalkoxy, amino, and halo;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl, or

R⁴ and R⁵, together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring; and

R⁶ is chosen from H and C₁-C₄alkyl.

[066] In certain embodiments, the compounds have structural Formula VII:



or a salt thereof, wherein:

Y is chosen from aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R² groups, or

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R³ groups;

each R² is independently chosen from alkyl, alkenyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, or

two R², together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

each R^3 is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R^8 groups, or


two R^3 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R^4 and R^5 are independently chosen from C_1 - C_6 alkyl, wherein exactly one of R^4 and R^5 is methyl;

R^6 is chosen from H and C_1 - C_4 alkyl; and


each R^8 is independently chosen from cyano, halo, hydroxy, and oxo.

[067] In certain embodiments, the compounds have structural Formula VII:



or a salt thereof, wherein:

L is chosen from a bond and methylene;



Y is chosen from alkenyl, alkenylamino, alkyl, aminoalkenyl, aminoalkyl, aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R^2 groups;

each R^2 is independently chosen from alkyl, alkenyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl;


n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R^4 and R^5 are independently chosen from C_1 - C_3 alkyl; and

R⁶ is chosen from H and C₁-C₄alkyl.

[068] In certain embodiments of structural formula VII, Y is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R² groups.

[069] In certain embodiments of structural formula VII, each R² is independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.




[070] In certain embodiments of structural formula VII W is

[071] In certain embodiments of structural formula VII, n is 1, forming azetidin-1-yl.

[072] In certain embodiments of structural formula VII, L is a bond.

[073] In certain embodiments, the compounds have structural Formula VIII:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

Y is chosen from phenyl, indazolyl, thienyl, indazolyl, and indolyl, any of which is optionally substituted with one or two R² groups;

n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

each R² is independently chosen from alkyl, alkoxy, alkylthio, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, and halo;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl; and

R⁶ is chosen from H and C₁-C₄alkyl.

[074] In certain embodiments of structural formula VIII, Y is chosen from phenyl and indazolyl, either of which is optionally substituted with one or two R² groups.

[075] In certain embodiments of structural formula VIII, each R² is independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.

[076] In certain embodiments of structural formula VIII, n is 1, forming azetidin-1-yl.


[077] In certain embodiments of structural formula VIII, L is a bond.

[078] In certain embodiments of structural formula VIII, Y is phenyl substituted with one or two R² groups independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.

[079] In certain embodiments of structural formula VIII, each R² is independently chosen from C₁-C₄alkoxy, cyclopropoxy, cyclobutoxy, and C₁-C₃alkyl.

[080] In certain embodiments of structural formula VIII, Y is indazolyl substituted with an R² groups chosen from methyl, ethyl, and propyl.

[081] In certain embodiments of structural formula VIII, Y is chosen from:




R^{2c} is chosen from C₁-C₄alkoxy and C₁-C₄cycloalkoxy;

R^{2d} is chosen from null and C₁-C₄alkoxy; and



R^{2e} is C₁-C₄alkyl.

[082] In certain embodiments, the compounds have structural Formula IX:



or a salt thereof, wherein:

L is chosen from a bond and methylene;



Y is chosen from aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R² groups;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R³ groups;

each R² is independently chosen from alkyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl,


carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl;

R^4 and R^5 are independently chosen from C₁-C₃alkyl; and

R^6 is chosen from H and C₁-C₄alkyl.

[083] In certain embodiments of structural formula IX, Y is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^2 groups.

[084] In certain embodiments of structural formula IX, each R^2 is independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.




[085] In certain embodiments of structural formula IX, W is

[086] In certain embodiments of structural formula IX, L is a bond.


[087] In certain embodiments of structural formula IX, Y is phenyl substituted with one or two R^2 groups independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.

[088] In certain embodiments of structural formula IX, each R^2 is independently chosen from C₁-C₄alkoxy, cyclopropoxy, cyclobutoxy, and C₁-C₃alkyl.

[089] In certain embodiments, the compounds have structural Formula X:



or a salt thereof, wherein:



R^{2e} is C₁-C₄alkyl;

L is chosen from a bond and methylene;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^3 groups;


each R^3 is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R^8 groups, or

two R^3 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R^4 and R^5 are independently chosen from C_1 - C_3 alkyl;

R^6 is chosen from H and C_1 - C_4 alkyl; and

each R^8 is independently chosen from cyano, halo, hydroxy, and oxo.




[090] In certain embodiments of structural formula X, W is

[091] In certain embodiments of structural formula X, L is a bond.

[092] In certain embodiments of structural formula X, Z is chosen from phenyl and indazolyl, either of which is optionally substituted with one to three R^3 groups.

[093] In certain embodiments of structural formula X, Z is chosen from:



n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R^4 and R^5 are independently chosen from C_1 - C_3 alkyl;

R^6 is chosen from H and C_1 - C_4 alkyl; and


R^9 is chosen from halo, amino, and C_1 - C_4 alkoxy.

[094] In certain embodiments of structural formula X, R^9 is chosen from chloro, dimethylamino, and ethoxy.


[095] In certain embodiments of structural formula X, R^9 is chosen from chloro and ethoxy.

[096] In certain embodiments of structural formula X, n is 1, forming azetidin-1-yl.

[097] In certain embodiments, the compounds have structural Formula XI:



or a salt thereof, wherein:



n is 1, 2, or 3 (forming a cyclopropyl, cyclobutyl, or cyclopentyl);

L is chosen from a bond and methylene;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R³ groups;

each R³ is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R⁸ groups, or


two R³, together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and

each R⁸ is independently chosen from cyano, halo, hydroxy, and oxo.


[0098] In certain embodiments of structural formula XI, W is



[0099] In certain embodiments of structural formula XI, L is a bond.

[0100] In certain embodiments of structural formula XI, Z is chosen from phenyl and indazolyl, either of which is optionally substituted with one to three R³ groups.

[0101] In certain embodiments of structural formula XI, Z is chosen from:



n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and


R⁹ is chosen from halo, amino, and C₁-C₄alkoxy.

[0102] In certain embodiments of structural formula XI, R⁹ is chosen from chloro, dimethylamino, and ethoxy.

[0103] In certain embodiments of structural formula XI, R⁹ is chosen from chloro and ethoxy.


[0104] In certain embodiments of structural formula XI, n is 1, forming azetidin-1-yl.

[0105] In certain embodiments, the compounds have structural Formula II:




or a salt thereof, wherein:

L is chosen from a bond and methylene;




Y is chosen from:



R^{2e} is C₁-C₄alkyl;

Z is chosen from:




n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and

R⁹ is chosen from halo and C₁-C₄alkoxy.




[0106] In certain embodiments of structural formula II, W is

[0107] In certain embodiments of structural formula II, R⁹ is chosen from chloro and ethoxy.

[0108] In certain embodiments of structural formula II, L is a bond.

[0109] In certain embodiments of structural formula II:



Y is

R^{2c} is chosen from C₁-C₄alkoxy and C₁-C₄cycloalkoxy; and


R^{2d} is chosen from null and C₁-C₄alkoxy.

[0110] In certain embodiments of structural formula II:

R^{2c} is chosen from methoxy, methylpropoxy, cyclopropoxy, and cyclobutoxy; and

R^{2d} is chosen from null and methoxy.


[0111] In certain embodiments of structural formula II:



Y is


R^{2e} is C₁-C₄alkyl.

[0112] In certain embodiments of structural formula II, R^{2e} is chosen from ethyl and propyl.




[0113] In certain embodiments of structural formula II, Z is

[0114] In certain embodiments of structural formula II, n is 1, forming azetidin-1-yl.



[0115] In certain embodiments of structural formula II, Z is



[0116] In certain embodiments of structural formula II, Z is

[0117] In certain embodiments of structural formula II, R⁹ is chosen from chloro and ethoxy.

[0118] Also provided are embodiments wherein any embodiment above may be combined with any one or more of these embodiments, provided the combination is not mutually exclusive.

[0119] As used herein, two embodiments are “mutually exclusive” when one is defined to be something which is different than the other. For example, an embodiment wherein two groups combine to form a cycloalkyl is mutually exclusive with an embodiment in which one group is ethyl the other group is hydrogen. Similarly, an embodiment wherein one group is CH₂ is mutually exclusive with an embodiment wherein the same group is NH.

[0120] Also provided is a compound chosen from the Examples disclosed herein.

[0121] The present invention also relates to a method of inhibiting at least one MCT4 function comprising the step of contacting MCT4 with a compound as described herein. The cell phenotype, cell proliferation, activity of MCT4, change in biochemical output produced by active MCT4, expression of MCT4, or binding of MCT4 with a natural binding partner may be monitored. Such methods may be modes of treatment of disease, biological assays, cellular assays, biochemical assays, or the like.

[0122] Also provided herein is a method of treatment of a MCT4-mediated disease comprising the administration of a therapeutically effective amount of a compound as disclosed herein, or a salt thereof, to a patient in need thereof.

[0123] In certain embodiments, the disease is chosen from proliferative inflammatory diseases.

[0124] In certain embodiments, the disease is a metabolic disease.

[0125] In certain embodiments, said metabolic disease is chosen from metabolic syndrome, diabetes, dyslipidemia, fatty liver disease, non-alcoholic steatohepatitis, obesity, and insulin resistance.

[0126] In certain embodiments, said diabetes is Type II diabetes.

[0127] In certain embodiments, said dyslipidemia is hyperlipidemia.

[0128] Further provided is a method for achieving an effect in a patient comprising the administration of a therapeutically effective amount of a compound as disclosed above to a patient, wherein the effect is selected from the group consisting of reduction of triglycerides, reduction of cholesterol, and reduction of hemoglobin A1c.

[0129] Further provided is the method as disclosed above wherein said cholesterol is chosen from LDL and VLDL cholesterol.

[0130] Further provided is the method as disclosed above wherein said triglycerides are chosen from plasma triglycerides and liver triglycerides.

[0131] Also provided herein is a compound as disclosed herein for use as a medicament.

[0132] Also provided herein is a compound as disclosed herein for use as a medicament for the treatment of a MCT4-mediated disease.

[0133] Also provided is the use of a compound as disclosed herein as a medicament.

[0134] Also provided is the use of a compound as disclosed herein as a medicament for the treatment of a MCT4-mediated disease.

[0135] Also provided is a compound as disclosed herein for use in the manufacture of a medicament for the treatment of a MCT4-mediated disease.

[0136] Also provided is the use of a compound as disclosed herein for the treatment of a MCT4-mediated disease.

[0137] Also provided herein is a method of inhibition of MCT4 comprising contacting MCT4 with a compound as disclosed herein, or a salt thereof.

[0138] Also provided herein is a method for achieving an effect in a patient comprising the administration of a therapeutically effective amount of a compound as disclosed herein, or a salt thereof, to a patient, wherein the effect is chosen from cognition enhancement.

[0139] In certain embodiments, the MCT4-mediated disease is chosen from proliferative inflammatory diseases.

[0140] Also provided is a method of modulation of a MCT4-mediated function in a subject comprising the administration of a therapeutically effective amount of a compound as disclosed herein.

[0141] Also provided is a pharmaceutical composition comprising a compound as disclosed herein, together with a pharmaceutically acceptable carrier.

[0142] In certain embodiments, the pharmaceutical composition is formulated for oral administration.

[0143] In certain embodiments, the oral pharmaceutical composition is chosen from a tablet and a capsule.

Abbreviations and Definitions

[0144] To facilitate understanding of the disclosure, a number of terms and abbreviations as used herein are defined below as follows:

[0145] When introducing elements of the present disclosure or the preferred embodiment(s) thereof, the articles "a", "an", "the" and "said" are intended to mean that there are one or more of the elements. The terms "comprising", "including" and "having" are intended to be inclusive and mean that there may be additional elements other than the listed elements.

[0146] The term "and/or" when used in a list of two or more items, means that any one of the listed items can be employed by itself or in combination with any one or more of the listed items. For example, the expression "A and/or B" is intended to mean either or both of A and B, i.e. A alone, B alone or A and B in combination. The expression "A, B and/or C" is intended to mean A alone, B alone, C alone, A and B in combination, A and C in combination, B and C in combination or A, B, and C in combination.

[0147] When ranges of values are disclosed, and the notation "from n_1 ... to n_2 " or "between n_1 ... and n_2 " is used, where n_1 and n_2 are the numbers, then unless otherwise specified, this notation is intended to include the numbers themselves and the range between them. This range may be integral or continuous between and including the end values. By way of example, the range "from 2 to 6 carbons" is intended to include two, three, four, five, and six carbons, since carbons come in integer units. Compare, by way of example, the range "from 1 to 3 μM (micromolar)," which is intended to include 1 μM , 3 μM , and everything in between to any number of significant figures (e.g., 1.255 μM , 2.1 μM , 2.9999 μM , etc.).

[0148] The term "about," as used herein, is intended to qualify the numerical values which it modifies, denoting such a value as variable within a margin of error. When no particular margin of error, such as a standard deviation to a mean value given in a chart or table of data, is recited, the term "about" should be understood to mean that range which would encompass the recited value and the range which would be included by rounding up or down to that figure as well, taking into account significant figures.

[0149] The term "acyl," as used herein, alone or in combination, refers to a carbonyl attached to an alkenyl, alkyl, aryl, cycloalkyl, heteroaryl, heterocycle, or any other moiety where the atom attached to the carbonyl is carbon. An "acetyl" group refers to a $-\text{C}(\text{O})\text{CH}_3$ group. An "alkylcarbonyl" or "alkanoyl" group refers to an alkyl group attached to the parent molecular moiety through a carbonyl group. Examples of such groups include methylcarbonyl and ethylcarbonyl. Examples of acyl groups include formyl, alkanoyl and aroyl.

[0150] The term “alkenyl,” as used herein, alone or in combination, refers to a straight-chain or branched-chain hydrocarbon radical having one or more double bonds and containing from 2 to 20 carbon atoms. In certain embodiments, said alkenyl will comprise from 2 to 6 carbon atoms. The term “alkenylene” refers to a carbon-carbon double bond system attached at two or more positions such as ethenylene $[(-CH=CH-), (-C::C-)]$. Examples of suitable alkenyl radicals include ethenyl, propenyl, 2-methylpropenyl, 1,4-butadienyl and the like. Unless otherwise specified, the term “alkenyl” may include “alkenylene” groups.

[0151] The term “alkoxy,” as used herein, alone or in combination, refers to an alkyl ether radical, wherein the term alkyl is as defined below. Examples of suitable alkyl ether radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, and the like.

[0152] The term “alkyl,” as used herein, alone or in combination, refers to a straight-chain or branched-chain alkyl radical containing from 1 to 20 carbon atoms. In certain embodiments, said alkyl will comprise from 1 to 10 carbon atoms. In further embodiments, said alkyl will comprise from 1 to 8 carbon atoms. Alkyl groups may be optionally substituted as defined herein. Examples of alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl, noyl and the like. The term “alkylene,” as used herein, alone or in combination, refers to a saturated aliphatic group derived from a straight or branched chain saturated hydrocarbon attached at two or more positions, such as methylene $(-CH_2-)$ and ethylene $(-CH_2CH_2-)$. Unless otherwise specified, the term “alkyl” may include “alkylene” groups.

[0153] The term “alkylamino,” as used herein, alone or in combination, refers to an alkyl group attached to the parent molecular moiety through an amino group. Suitable alkylamino groups may be mono- or dialkylated, forming groups such as, for example, N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-ethylmethylamino and the like.

[0154] The term “alkylidene,” as used herein, alone or in combination, refers to an alkenyl group in which one carbon atom of the carbon-carbon double bond belongs to the moiety to which the alkenyl group is attached.

[0155] The term “alkylthio,” as used herein, alone or in combination, refers to an alkyl thioether $(R-S-)$ radical wherein the term alkyl is as defined above and wherein the sulfur may be singly or doubly oxidized. Examples of suitable alkyl thioether radicals include

methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, iso-butylthio, sec-butylthio, tert-butylthio, methanesulfonyl, ethanesulfinyl, and the like.

[0156] The term “alkynyl,” as used herein, alone or in combination, refers to a straight-chain or branched chain hydrocarbon radical having one or more triple bonds and containing from 2 to 20 carbon atoms. In certain embodiments, said alkynyl comprises from 2 to 6 carbon atoms. In further embodiments, said alkynyl comprises from 2 to 4 carbon atoms. The term “alkynylene” refers to a carbon-carbon triple bond attached at two positions such as ethynylene ($-C\equiv C-$, $-C\equiv C-$). Examples of alkynyl radicals include ethynyl, propynyl, hydroxypropynyl, butyn-1-yl, butyn-2-yl, pentyn-1-yl, 3-methylbutyn-1-yl, hexyn-2-yl, and the like. Unless otherwise specified, the term “alkynyl” may include “alkynylene” groups.

[0157] The terms “amido”, “carboxamido”, and “carbamoyl,” as used herein, alone or in combination, refer to an amino group as described below attached to the parent molecular moiety through a carbonyl group, or vice versa. The term “C-amido” as used herein, alone or in combination, refers to a $-C(O)N(RR')$ group with R and R' as defined herein or as defined by the specifically enumerated “R” groups designated. The term “N-amido” as used herein, alone or in combination, refers to a $RC(O)N(R')$ - group, with R and R' as defined herein or as defined by the specifically enumerated “R” groups designated. The term “acylamino” as used herein, alone or in combination, embraces an acyl group attached to the parent moiety through an amino group. An example of an “acylamino” group is acetylamino ($CH_3C(O)NH-$).

[0158] The term “amino,” as used herein, alone or in combination, refers to $-NRR'$, wherein R and R' are independently chosen from hydrogen, alkyl, acyl, heteroalkyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl, any of which may themselves be optionally substituted. Additionally, R and R' may combine to form heterocycloalkyl, either of which may be optionally substituted.

[0159] The term “aryl,” as used herein, alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such polycyclic ring systems are fused together. The term “aryl” embraces aromatic groups such as phenyl, naphthyl, anthracenyl, and phenanthryl.

[0160] The term “arylalkenyl” or “aralkenyl,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkenyl group.

[0161] The term “arylalkoxy” or “aralkoxy,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkoxy group.

[0162] The term “arylalkyl” or “aralkyl,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkyl group.

[0163] The term “arylalkynyl” or “aralkynyl,” as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an alkynyl group.

[0164] The term “arylalkanoyl” or “aralkanoyl” or “aroyl,” as used herein, alone or in combination, refers to an acyl radical derived from an aryl-substituted alkanecarboxylic acid such as benzoyl, naphthoyl, phenylacetyl, 3-phenylpropionyl (hydrocinnamoyl), 4-phenylbutyryl, (2-naphthyl)acetyl, 4-chlorohydrocinnamoyl, and the like.

[0165] The term aryloxy as used herein, alone or in combination, refers to an aryl group attached to the parent molecular moiety through an oxy.

[0166] The terms “benzo” and “benz,” as used herein, alone or in combination, refer to the divalent radical $C_6H_4=$ derived from benzene. Examples include benzothiophene and benzimidazole.

[0167] The term “carbamate,” as used herein, alone or in combination, refers to an ester of carbamic acid ($-NHCOO-$) which may be attached to the parent molecular moiety from either the nitrogen or acid end, and which may be optionally substituted as defined herein.

[0168] The term “O-carbamyl” as used herein, alone or in combination, refers to a $-OC(O)NRR'$, group-with R and R' as defined herein.

[0169] The term “N-carbamyl” as used herein, alone or in combination, refers to a $ROC(O)NR'$ - group, with R and R' as defined herein.

[0170] The term “carbonyl,” as used herein, when alone includes formyl [$-C(O)H$] and in combination is a $-C(O)-$ group.

[0171] The term “carboxyl” or “carboxy,” as used herein, refers to $-C(O)OH$ or the corresponding “carboxylate” anion, such as is in a carboxylic acid salt. An “O-carboxy” group refers to a $RC(O)O-$ group, where R is as defined herein. A “C-carboxy” group refers to a $-C(O)OR$ groups where R is as defined herein.

[0172] The term “cyano,” as used herein, alone or in combination, refers to $-CN$.

[0173] The term “cycloalkyl,” or, alternatively, “carbocycle,” as used herein, alone or in combination, refers to a saturated or partially saturated monocyclic, bicyclic or tricyclic alkyl group wherein each cyclic moiety contains from 3 to 12 carbon atom ring members and which may optionally be a benzo fused ring system which is optionally substituted as defined

herein. In certain embodiments, said cycloalkyl will comprise from 5 to 7 carbon atoms. Examples of such cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, indanyl, octahydronaphthyl, 2,3-dihydro-1H-indenyl, adamantyl and the like. "Bicyclic" and "tricyclic" as used herein are intended to include both fused ring systems, such as decahydronaphthalene, octahydronaphthalene as well as the multicyclic (multicentered) saturated or partially unsaturated type. The latter type of isomer is exemplified in general by, bicyclo[1,1,1]pentane, camphor, adamantane, and bicyclo[3,2,1]octane.

[0174] The term "ester," as used herein, alone or in combination, refers to a carboxy group bridging two moieties linked at carbon atoms.

[0175] The term "ether," as used herein, alone or in combination, refers to an oxy group bridging two moieties linked at carbon atoms.

[0176] The term "halo," or "halogen," as used herein, alone or in combination, refers to fluorine, chlorine, bromine, or iodine.

[0177] The term "haloalkoxy," as used herein, alone or in combination, refers to a haloalkyl group attached to the parent molecular moiety through an oxygen atom.

[0178] The term "haloalkyl," as used herein, alone or in combination, refers to an alkyl radical having the meaning as defined above wherein one or more hydrogens are replaced with a halogen. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. "Haloalkylene" refers to a haloalkyl group attached at two or more positions. Examples include fluoromethylene

(-CFH-), difluoromethylene (-CF₂-), chloromethylene (-CHCl-) and the like.

[0179] The term "heteroalkyl," as used herein, alone or in combination, refers to a stable straight or branched chain, or combinations thereof, fully saturated or containing from 1 to 3 degrees of unsaturation, consisting of the stated number of carbon atoms and from one to three heteroatoms chosen from N, O, and S, and wherein the N and S atoms may optionally be oxidized and the N heteroatom may optionally be quaternized. The heteroatom(s) may be

placed at any interior position of the heteroalkyl group. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃.

[0180] The term "heteroaryl," as used herein, alone or in combination, refers to a 3 to 15 membered unsaturated heteromonocyclic ring, or a fused monocyclic, bicyclic, or tricyclic ring system in which at least one of the fused rings is aromatic, which contains at least one atom chosen from N, O, and S. In certain embodiments, said heteroaryl will comprise from 1 to 4 heteroatoms as ring members. In further embodiments, said heteroaryl will comprise from 1 to 2 heteroatoms as ring members. In certain embodiments, said heteroaryl will comprise from 5 to 7 atoms. The term also embraces fused polycyclic groups wherein heterocyclic rings are fused with aryl rings, wherein heteroaryl rings are fused with other heteroaryl rings, wherein heteroaryl rings are fused with heterocycloalkyl rings, or wherein heteroaryl rings are fused with cycloalkyl rings. Examples of heteroaryl groups include pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl, pyranyl, furyl, thienyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, thiadiazolyl, isothiazolyl, indolyl, isoindolyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl, quinoxaliny, quinazolinyl, indazolyl, benzotriazolyl, benzodioxolyl, benzopyranyl, benzoxazolyl, benzoxadiazolyl, benzothiazolyl, benzothiadiazolyl, benzofuryl, benzothienyl, chromonyl, coumarinyl, benzopyranyl, tetrahydroquinolinyl, tetrazolopyridazinyl, tetrahydroisoquinolinyl, thienopyridinyl, furopyridinyl, pyrrolopyridinyl and the like. Exemplary tricyclic heterocyclic groups include carbazolyl, benzidolyl, phenanthrolinyl, dibenzofuranyl, acridinyl, phenanthridinyl, xanthenyl and the like.

[0181] The terms "heterocycloalkyl" and, interchangeably, "heterocycle," as used herein, alone or in combination, each refer to a saturated, partially unsaturated, or fully unsaturated (but nonaromatic) monocyclic, bicyclic, or tricyclic heterocyclic group containing at least one heteroatom as a ring member, wherein each said heteroatom may be independently chosen from nitrogen, oxygen, and sulfur. In certain embodiments, said heterocycloalkyl will comprise from 1 to 4 heteroatoms as ring members. In further embodiments, said heterocycloalkyl will comprise from 1 to 2 heteroatoms as ring members. In certain embodiments, said heterocycloalkyl will comprise from 3 to 8 ring members in each ring. In further embodiments, said heterocycloalkyl will comprise from 3 to 7 ring members in each ring. In yet further embodiments, said heterocycloalkyl will comprise from 5 to 6 ring members in each ring. "Heterocycloalkyl" and "heterocycle" are intended to include sulfones, sulfoxides, N-oxides of tertiary nitrogen ring members, and carbocyclic fused and benzo

fused ring systems; additionally, both terms also include systems where a heterocycle ring is fused to an aryl group, as defined herein, or an additional heterocycle group. Examples of heterocycle groups include aziridinyl, azetidiny, 1,3-benzodioxolyl, dihydroisoindolyl, dihydroisoquinolyl, dihydrocinnolyl, dihydrobenzodioxinyl, dihydro[1,3]oxazolo[4,5-b]pyridinyl, benzothiazolyl, dihydroindolyl, dihydrodropyridinyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,3-dioxolanyl, isoindolyl, morpholyl, piperazinyl, pyrrolidinyl, tetrahydropyridinyl, piperidinyl, thiomorpholyl, and the like. The heterocycle groups may be optionally substituted unless specifically prohibited.

[0182] The term “hydrazinyl” as used herein, alone or in combination, refers to two amino groups joined by a single bond, i.e., -N-N-.

[0183] The term “hydroxy,” as used herein, alone or in combination, refers to -OH.

[0184] The term “hydroxyalkyl,” as used herein, alone or in combination, refers to a hydroxy group attached to the parent molecular moiety through an alkyl group.

[0185] The term “imino,” as used herein, alone or in combination, refers to =N-.

[0186] The term “iminohydroxy,” as used herein, alone or in combination, refers to =N(OH) and =N-O-.

[0187] The phrase “in the main chain” refers to the longest contiguous or adjacent chain of carbon atoms starting at the point of attachment of a group to the compounds of any one of the formulas disclosed herein.

[0188] The term “isocyanato” refers to a -NCO group.

[0189] The term “isothiocyanato” refers to a -NCS group.

[0190] The phrase “linear chain of atoms” refers to the longest straight chain of atoms independently selected from carbon, nitrogen, oxygen and sulfur.

[0191] The term “lower,” as used herein, alone or in a combination, where not otherwise specifically defined, means containing from 1 to and including 6 carbon atoms (i.e., C1-C6 alkyl).

[0192] The term “lower aryl,” as used herein, alone or in combination, means phenyl or naphthyl, either of which may be optionally substituted as provided.

[0193] The term “lower heteroaryl,” as used herein, alone or in combination, means either 1) monocyclic heteroaryl comprising five or six ring members, of which between one and four said members may be heteroatoms chosen from N, O, and S, or 2) bicyclic heteroaryl, wherein each of the fused rings comprises five or six ring members, comprising between them one to four heteroatoms chosen from N, O, and S.

[0194] The term “lower cycloalkyl,” as used herein, alone or in combination, means a monocyclic cycloalkyl having between three and six ring members (i.e., C3-C6 cycloalkyl). Lower cycloalkyls may be unsaturated. Examples of lower cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

[0195] The term “lower heterocycloalkyl,” as used herein, alone or in combination, means a monocyclic heterocycloalkyl having between three and six ring members, of which between one and four may be heteroatoms chosen from N, O, and S (i.e., C3-C6 heterocycloalkyl). Examples of lower heterocycloalkyls include pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidinyl, piperazinyl, and morpholinyl. Lower heterocycloalkyls may be unsaturated.

[0196] The term “lower amino,” as used herein, alone or in combination, refers to -NRR', wherein R and R' are independently chosen from hydrogen and lower alkyl, either of which may be optionally substituted.

[0197] The term “mercaptyl” as used herein, alone or in combination, refers to an RS-group, where R is as defined herein.

[0198] The term “nitro,” as used herein, alone or in combination, refers to -NO₂.

[0199] The terms “oxy” or “oxa,” as used herein, alone or in combination, refer to -O-.

[0200] The term “oxo,” as used herein, alone or in combination, refers to =O.

[0201] The term “perhaloalkoxy” refers to an alkoxy group where all of the hydrogen atoms are replaced by halogen atoms.

[0202] The term “perhaloalkyl” as used herein, alone or in combination, refers to an alkyl group where all of the hydrogen atoms are replaced by halogen atoms.

[0203] The terms “sulfonate,” “sulfonic acid,” and “sulfonic,” as used herein, alone or in combination, refer the -SO₃H group and its anion as the sulfonic acid is used in salt formation.

[0204] The term “sulfanyl,” as used herein, alone or in combination, refers to -S-.

[0205] The term “sulfinyl,” as used herein, alone or in combination, refers to -S(O)-.

[0206] The term “sulfonyl,” as used herein, alone or in combination, refers to -S(O)₂-.

[0207] The term “N-sulfonamido” refers to a RS(O)₂NR'- group with R and R' as defined herein.

[0208] The term “S-sulfonamido” refers to a -S(O)₂NRR', group, with R and R' as defined herein.

[0209] The term “sulfonamido” encompasses both N-sulfonamido and S-sulfonamido groups.

[0210] The terms “thia” and “thio,” as used herein, alone or in combination, refer to a –S– group or an ether wherein the oxygen is replaced with sulfur. The oxidized derivatives of the thio group, namely sulfinyl and sulfonyl, are included in the definition of thia and thio.

[0211] The term “thiol,” as used herein, alone or in combination, refers to an –SH group.

[0212] The term “thiocarbonyl,” as used herein, when alone includes thioformyl –C(S)H and in combination is a –C(S)– group.

[0213] The term “N-thiocarbamyl” refers to an ROC(S)NR’– group, with R and R’ as defined herein.

[0214] The term “O-thiocarbamyl” refers to a –OC(S)NRR’, group with R and R’ as defined herein.

[0215] The term “thiocyanato” refers to a –CNS group.

[0216] The term “trihalomethanesulfonamido” refers to a X₃CS(O)₂NR– group with X is a halogen and R as defined herein.

[0217] The term “trihalomethanesulfonyl” refers to a X₃CS(O)₂– group where X is a halogen.

[0218] The term “trihalomethoxy” refers to a X₃CO– group where X is a halogen.

[0219] The term “trisubstituted silyl,” as used herein, alone or in combination, refers to a silicone group substituted at its three free valences with groups as listed herein under the definition of substituted amino. Examples include trimethylsilyl, tert-butyldimethylsilyl, triphenylsilyl and the like.

[0220] Any definition herein may be used in combination with any other definition to describe a composite structural group. By convention, the trailing element of any such definition is that which attaches to the parent moiety. For example, the composite group alkylamido would represent an alkyl group attached to the parent molecule through an amido group, and the term alkoxyalkyl would represent an alkoxy group attached to the parent molecule through an alkyl group.

[0221] When a group is defined to be “null,” what is meant is that said group is absent.

[0222] The term “optionally substituted” means the antecedent group may be substituted or unsubstituted. When substituted, the substituents of an “optionally substituted” group may include, without limitation, one or more substituents independently selected from the following groups or a particular designated set of groups, alone or in combination: lower

alkyl, lower alkenyl, lower alkynyl, lower alkanoyl, lower heteroalkyl, lower heterocycloalkyl, lower haloalkyl, lower haloalkenyl, lower haloalkynyl, lower perhaloalkyl, lower perhaloalkoxy, lower cycloalkyl, phenyl, aryl, aryloxy, lower alkoxy, lower haloalkoxy, oxo, lower acyloxy, carbonyl, carboxyl, lower alkylcarbonyl, lower carboxyester, lower carboxamido, cyano, hydrogen, halogen, hydroxy, amino, lower alkylamino, arylamino, amido, nitro, thiol, lower alkylthio, lower haloalkylthio, lower perhaloalkylthio, arylthio, sulfonate, sulfonic acid, trisubstituted silyl, N₃, SH, SCH₃, C(O)CH₃, CO₂CH₃, CO₂H, pyridinyl, thiophene, furanyl, lower carbamate, and lower urea. Where structurally feasible, two substituents may be joined together to form a fused five-, six-, or seven-membered carbocyclic or heterocyclic ring consisting of zero to three heteroatoms, for example forming methylenedioxy or ethylenedioxy. An optionally substituted group may be unsubstituted (e.g., -CH₂CH₃), fully substituted (e.g., -CF₂CF₃), monosubstituted (e.g., -CH₂CH₂F) or substituted at a level anywhere in-between fully substituted and monosubstituted (e.g., -CH₂CF₃). Where substituents are recited without qualification as to substitution, both substituted and unsubstituted forms are encompassed. Where a substituent is qualified as “substituted,” the substituted form is specifically intended. Additionally, different sets of optional substituents to a particular moiety may be defined as needed; in these cases, the optional substitution will be as defined, often immediately following the phrase, “optionally substituted with.”

[0223] The term R or the term R', appearing by itself and without a number designation, unless otherwise defined, refers to a moiety chosen from hydrogen, alkyl, cycloalkyl, heteroalkyl, aryl, heteroaryl and heterocycloalkyl, any of which may be optionally substituted. Such R and R' groups should be understood to be optionally substituted as defined herein. Whether an R group has a number designation or not, every R group, including R, R' and R_n where n=(1, 2, 3, ... n), every substituent, and every term should be understood to be independent of every other in terms of selection from a group. Should any variable, substituent, or term (e.g. aryl, heterocycle, R, etc.) occur more than one time in a formula or generic structure, its definition at each occurrence is independent of the definition at every other occurrence. Those of skill in the art will further recognize that certain groups may be attached to a parent molecule or may occupy a position in a chain of elements from either end as written. For example, an unsymmetrical group such as -C(O)N(R)- may be attached to the parent moiety at either the carbon or the nitrogen.

[0224] Asymmetric centers exist in the compounds disclosed herein. These centers are designated by the symbols “R” or “S,” depending on the configuration of substituents around the chiral carbon atom. It should be understood that the invention encompasses all stereochemical isomeric forms, including diastereomeric, enantiomeric, and epimeric forms, as well as d-isomers and l-isomers, and mixtures thereof. Individual stereoisomers of compounds can be prepared synthetically from commercially available starting materials which contain chiral centers or by preparation of mixtures of enantiomeric products followed by separation such as conversion to a mixture of diastereomers followed by separation or recrystallization, chromatographic techniques, direct separation of enantiomers on chiral chromatographic columns, or any other appropriate method known in the art. Starting compounds of particular stereochemistry are either commercially available or can be made and resolved by techniques known in the art. Additionally, the compounds disclosed herein may exist as geometric isomers. The present invention includes all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof. Additionally, compounds may exist as tautomers; all tautomeric isomers are provided by this invention. Additionally, the compounds disclosed herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms.

[0225] The term “bond” refers to a covalent linkage between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure. A bond may be single, double, or triple unless otherwise specified. A dashed line between two atoms in a drawing of a molecule indicates that an additional bond may be present or absent at that position.

[0226] The term “disease” as used herein is intended to be generally synonymous, and is used interchangeably with, the terms “disorder,” “syndrome,” and “condition” (as in medical condition), in that all reflect an abnormal condition of the human or animal body or of one of its parts that impairs normal functioning, is typically manifested by distinguishing signs and symptoms, and causes the human or animal to have a reduced duration or quality of life.

[0227] The term “combination therapy” means the administration of two or more therapeutic agents to treat a therapeutic condition or disorder described in the present disclosure. Such administration encompasses co-administration of these therapeutic agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of active ingredients or in multiple, separate capsules for each active ingredient. In addition, such

administration also encompasses use of each type of therapeutic agent in a sequential manner. In either case, the treatment regimen will provide beneficial effects of the drug combination in treating the conditions or disorders described herein.

[0228] The term “MCT4 inhibitor” is used herein to refer to a compound that exhibits an IC₅₀ with respect to MCT4 activity of no more than about 100 μ M and more typically not more than about 50 μ M, as measured in the MCT4 enzyme assay described generally herein below. IC₅₀ is that concentration of inhibitor that reduces the activity of an enzyme (e.g., MCT4) to half-maximal level. Certain compounds disclosed herein have been discovered to exhibit inhibition against MCT4. In certain embodiments, compounds will exhibit an IC₅₀ with respect to MCT4 of no more than about 10 μ M; in further embodiments, compounds will exhibit an IC₅₀ with respect to MCT4 of no more than about 5 μ M; in yet further embodiments, compounds will exhibit an IC₅₀ with respect to MCT4 of not more than about 1 μ M; in yet further embodiments, compounds will exhibit an IC₅₀ with respect to MCT4 of not more than about 200 nM, as measured in the MCT4 binding assay described herein.

[0229] The phrase “therapeutically effective” is intended to qualify the amount of active ingredients used in the treatment of a disease or disorder or on the effecting of a clinical endpoint.

[0230] The term “therapeutically acceptable” refers to those compounds (or salts, prodrugs, tautomers, zwitterionic forms, etc.) which are suitable for use in contact with the tissues of patients without undue toxicity, irritation, and allergic response, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.

[0231] As used herein, reference to “treatment” of a patient is intended to include prophylaxis. Treatment may also be preemptive in nature, i.e., it may include prevention of disease. Prevention of a disease may involve complete protection from disease, for example as in the case of prevention of infection with a pathogen, or may involve prevention of disease progression. For example, prevention of a disease may not mean complete foreclosure of any effect related to the diseases at any level, but instead may mean prevention of the symptoms of a disease to a clinically significant or detectable level. Prevention of diseases may also mean prevention of progression of a disease to a later stage of the disease.

[0232] The term “patient” is generally synonymous with the term “subject” and includes all mammals including humans. Examples of patients include humans, livestock such as cows, goats, sheep, pigs, and rabbits, and companion animals such as dogs, cats, rabbits, and horses. Preferably, the patient is a human.

[0233] The term "prodrug" refers to a compound that is made more active in vivo. Certain compounds disclosed herein may also exist as prodrugs, as described in Hydrolysis in Drug and Prodrug Metabolism : Chemistry, Biochemistry, and Enzymology (Testa, Bernard and Mayer, Joachim M. Wiley-VHCA, Zurich, Switzerland 2003). Prodrugs of the compounds described herein are structurally modified forms of the compound that readily undergo chemical changes under physiological conditions to provide the compound. Additionally, prodrugs can be converted to the compound by chemical or biochemical methods in an ex vivo environment. For example, prodrugs can be slowly converted to a compound when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent. Prodrugs are often useful because, in some situations, they may be easier to administer than the compound, or parent drug. They may, for instance, be bioavailable by oral administration whereas the parent drug is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. A wide variety of prodrug derivatives are known in the art, such as those that rely on hydrolytic cleavage or oxidative activation of the prodrug. An example, without limitation, of a prodrug would be a compound which is administered as an ester (the "prodrug"), but then is metabolically hydrolyzed to the carboxylic acid, the active entity. Additional examples include peptidyl derivatives of a compound.

[0234] The compounds disclosed herein can exist as therapeutically acceptable salts. The present invention includes compounds listed above in the form of salts, including acid addition salts. Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable. However, salts of non-pharmaceutically acceptable salts may be of utility in the preparation and purification of the compound in question. Basic addition salts may also be formed and be pharmaceutically acceptable. For a more complete discussion of the preparation and selection of salts, refer to Pharmaceutical Salts: Properties, Selection, and Use (Stahl, P. Heinrich. Wiley-VCHA, Zurich, Switzerland, 2002).

[0235] The term "therapeutically acceptable salt," as used herein, represents salts or zwitterionic forms of the compounds disclosed herein which are water or oil-soluble or dispersible and therapeutically acceptable as defined herein. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting the appropriate compound in the form of the free base with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, L-ascorbate, aspartate, benzoate, benzenesulfonate

(besylate), bisulfate, butyrate, camphorate, camphorsulfonate, citrate, digluconate, formate, fumarate, gentisate, glutarate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, malonate, DL-mandelate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphonate, picrate, pivalate, propionate, pyroglutamate, succinate, sulfonate, tartrate, L-tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate (p-tosylate), and undecanoate. Also, basic groups in the compounds disclosed herein can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and benzyl and phenethyl bromides. Examples of acids which can be employed to form therapeutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric. Salts can also be formed by coordination of the compounds with an alkali metal or alkaline earth ion. Hence, the present invention contemplates sodium, potassium, magnesium, and calcium salts of the compounds disclosed herein, and the like.

[0236] Basic addition salts can be prepared during the final isolation and purification of the compounds by reacting a carboxy group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary, secondary, or tertiary amine. The cations of therapeutically acceptable salts include lithium, sodium, potassium, calcium, magnesium, and aluminum, as well as nontoxic quaternary amine cations such as ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, 1-phenamine, and N,N'-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, and piperazine.

[0237] While it may be possible for the compounds of the subject invention to be administered as the raw chemical, it is also possible to present them as a pharmaceutical formulation. Accordingly, provided herein are pharmaceutical formulations which comprise one or more of certain compounds disclosed herein, or one or more pharmaceutically

acceptable salts, esters, prodrugs, amides, or solvates thereof, together with one or more pharmaceutically acceptable carriers thereof and optionally one or more other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Proper formulation is dependent upon the route of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art. The pharmaceutical compositions disclosed herein may be manufactured in any manner known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[0238] The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous, intraarticular, and intramedullary), intraperitoneal, transmucosal, transdermal, rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Typically, these methods include the step of bringing into association a compound of the subject invention or a pharmaceutically acceptable salt, ester, amide, prodrug or solvate thereof ("active ingredient") with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

[0239] Formulations of the compounds disclosed herein suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

[0240] Pharmaceutical preparations which can be used orally include tablets, push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. Tablets may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with binders, inert diluents, or lubricating, surface active or

dispersing agents. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein. All formulations for oral administration should be in dosages suitable for such administration. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

[0241] The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in powder form or in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, saline or sterile pyrogen-free water, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

[0242] Formulations for parenteral administration include aqueous and non-aqueous (oily) sterile injection solutions of the active compounds which may contain antioxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran.

Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0243] In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0244] For buccal or sublingual administration, the compositions may take the form of tablets, lozenges, pastilles, or gels formulated in conventional manner. Such compositions may comprise the active ingredient in a flavored basis such as sucrose and acacia or tragacanth.

[0245] The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter, polyethylene glycol, or other glycerides.

[0246] Certain compounds disclosed herein may be administered topically, that is by non-systemic administration. This includes the application of a compound disclosed herein externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

[0247] Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as gels, liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient for topical administration may comprise, for example, from 0.001% to 10% w/w (by weight) of the formulation. In certain embodiments, the active ingredient may comprise as much as 10% w/w. In other embodiments, it may comprise less than 5% w/w. In certain embodiments, the active ingredient may comprise from 2% w/w to 5% w/w. In other embodiments, it may comprise from 0.1% to 1% w/w of the formulation.

[0248] For administration by inhalation, compounds may be conveniently delivered from an insufflator, nebulizer pressurized packs or other convenient means of delivering an aerosol spray. Pressurized packs may comprise a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide

or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Alternatively, for administration by inhalation or insufflation, the compounds according to the invention may take the form of a dry powder composition, for example a powder mix of the compound and a suitable powder base such as lactose or starch. The powder composition may be presented in unit dosage form, in for example, capsules, cartridges, gelatin or blister packs from which the powder may be administered with the aid of an inhalator or insufflator.

[0249] Preferred unit dosage formulations are those containing an effective dose, as herein below recited, or an appropriate fraction thereof, of the active ingredient.

[0250] It should be understood that in addition to the ingredients particularly mentioned above, the formulations described above may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

[0251] Compounds may be administered orally or via injection at a dose of from 0.1 to 500 mg/kg per day. The dose range for adult humans is generally from 5 mg to 2 g/day. Tablets or other forms of presentation provided in discrete units may conveniently contain an amount of one or more compounds which is effective at such dosage or as a multiple of the same, for instance, units containing 5 mg to 500 mg, usually around 10 mg to 200 mg.

[0252] The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

[0253] The compounds can be administered in various modes, e.g. orally, topically, or by injection. The precise amount of compound administered to a patient will be the responsibility of the attendant physician. The specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diets, time of administration, route of administration, rate of excretion, drug combination, the precise disorder being treated, and the severity of the indication or condition being treated. Also, the route of administration may vary depending on the condition and its severity.

[0254] In certain instances, it may be appropriate to administer at least one of the compounds described herein (or a pharmaceutically acceptable salt, ester, or prodrug thereof) in combination with another therapeutic agent. By way of example only, if one of the side effects experienced by a patient upon receiving one of the compounds herein is hypertension,

then it may be appropriate to administer an anti-hypertensive agent in combination with the initial therapeutic agent. Or, by way of example only, the therapeutic effectiveness of one of the compounds described herein may be enhanced by administration of an adjuvant (i.e., by itself the adjuvant may only have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, by way of example only, the benefit of experienced by a patient may be increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit. By way of example only, in a treatment for diabetes involving administration of one of the compounds described herein, increased therapeutic benefit may result by also providing the patient with another therapeutic agent for diabetes. In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

[0255] Thus, in another aspect, certain embodiments provide methods for treating MCT4-mediated disorders in a human or animal subject in need of such treatment comprising administering to said subject an amount of a compound disclosed herein effective to reduce or prevent said disorder in the subject, in combination with at least one additional agent for the treatment of said disorder that is known in the art. In a related aspect, certain embodiments provide therapeutic compositions comprising at least one compound disclosed herein in combination with one or more additional agents for the treatment of MCT4-mediated disorders.

[0256] Also provided herein is a method of treating a monocarboxylate transporter MCT4-mediated disorder in a subject in need thereof, comprising the sequential or co-administration of a compound as disclosed herein or a pharmaceutically acceptable salt thereof, and another therapeutic agent.

[0257] In certain embodiments, the therapeutic agent is a protein kinase inhibitor.

[0258] In certain embodiments, the protein kinase inhibitor is chosen from Aurora B, EGFR, PLK-1, CDKs inhibitors.

[0259] In certain embodiments, the therapeutic agent is chosen from an antimetabolite, bcr-abl inhibitor, DNA damaging agent, EGFR inhibitor, microtubule stabilizing inhibitor, mitotic arrest inhibitor, S-phase inhibitor, and a taxane.

[0260] In certain embodiments, the therapeutic agent is a DNA damaging agent chosen from an alkylating agent, anthracycline, antimetabolite agent, crosslinking agent, DNA

replication inhibitor, intercalator, microtubule disruptor, PARP inhibitor, radiomimetic agent, radiosensitizer, strand break agent, and topoisomerase II inhibitor.

[0261] In certain embodiments, the therapeutic agent is chosen from aminoglutethimide, amsacrine, anastrozole, asparaginase, barasertib, bcg, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, chloroquine, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, demethoxyviridin, dichloroacetate, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, everolimus, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine, genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, irinotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, lonidamine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, metformin, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, olaparib, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, perifosine, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, sorafenib, streptozocin, sunitinib, suramin, tamoxifen, temozolomide, temsirolimus, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, and vinorelbine.

[0262] For use in cancer and neoplastic diseases a MCT4 inhibitor may be optimally used together with one or more of the following non-limiting examples of anti-cancer agents: (1) alkylating agents, including but not limited to cisplatin (PLATIN), carboplatin (PARAPLATIN), oxaliplatin (ELOXATIN), streptozocin (ZANOSAR), busulfan (MYLERAN) and cyclophosphamide (ENDOXAN); (2) anti-metabolites, including but not limited to mercaptopurine (PURINETHOL), thioguanine, pentostatin (NIPENT), cytosine arabinoside (ARA-C), gemcitabine (GEMZAR), fluorouracil (CARAC), leucovorin (FUSILEV) and methotrexate (RHEUMATREX); (3) plant alkaloids and terpenoids, including but not limited to vincristine (ONCOVIN), vinblastine and paclitaxel (TAXOL); (4) topoisomerase inhibitors, including but not limited to irinotecan (CAMPTOSAR), topotecan (HYCAMTIN) and etoposide (EPOSIN); (5) cytotoxic antibiotics, including but not limited to actinomycin D (COSMEGEN), doxorubicin (ADRIAMYCIN), bleomycin (BLENOXANE) and mitomycin (MITOSOL); (6) angiogenesis inhibitors, including but not limited to sunitinib (SUTENT) and bevacizumab (AVASTIN); and (7) tyrosine kinase

inhibitors, including but not limited to imatinib (GLEEVEC), erlotinib (TARCEVA), lapatinib (TYKERB) and axitinib (INLYTA).

[0263] Where a subject is suffering from or at risk of suffering from an inflammatory condition, a MCT4 inhibitor compound described herein is optionally used together with one or more agents or methods for treating an inflammatory condition in any combination. Therapeutic agents/treatments for treating an autoimmune and/or inflammatory condition include, but are not limited to any of the following examples: (1) corticosteroids, including but not limited to cortisone, dexamethasone, and methylprednisolone; (2) nonsteroidal anti-inflammatory drugs (NSAIDs), including but not limited to ibuprofen, naproxen, acetaminophen, aspirin, fenoprofen (NALFON), flurbiprofen (ANSAID), ketoprofen, oxaprozin (DAYPRO), diclofenac sodium (VOLTAREN), diclofenac potassium (CATAFLAM), etodolac (LODINE), indomethacin (INDOCIN), ketorolac (TORADOL), sulindac (CLINORIL), tolmetin (TOLECTIN), meclofenamate (MECLOMEN), mefenamic acid (PONSTEL), nabumetone (RELAFEN) and piroxicam (FELDENE); (3) immunosuppressants, including but not limited to methotrexate (RHEUMATREX), leflunomide (ARAVA), azathioprine (IMURAN), cyclosporine (NEORAL, SANDIMMUNE), tacrolimus and cyclophosphamide (CYTOXAN); (4) CD20 blockers, including but not limited to rituximab (RITUXAN); (5) Tumor Necrosis Factor (TNF) blockers, including but not limited to etanercept (ENBREL), infliximab (REMICADE) and adalimumab (HUMIRA); (6) interleukin-1 receptor antagonists, including but not limited to anakinra (KINERET); (7) interleukin-6 inhibitors, including but not limited to tocilizumab (ACTEMRA); (8) interleukin-17 inhibitors, including but not limited to AIN457; (9) Janus kinase inhibitors, including but not limited to tasocitinib; and (10) syk inhibitors, including but not limited to fostamatinib.

[0264] In certain embodiments, the method further comprises administering non-chemical methods of cancer treatment.

[0265] In certain embodiments, the method further comprises administering radiation therapy.

[0266] In certain embodiments, the method further comprises administering surgery, thermoablation, focused ultrasound therapy, cryotherapy, or any combination thereof.

[0267] In any case, the multiple therapeutic agents (at least one of which is a compound disclosed herein) may be administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents may be provided in a single, unified form, or

in multiple forms (by way of example only, either as a single pill or as two separate pills). One of the therapeutic agents may be given in multiple doses, or both may be given as multiple doses. If not simultaneous, the timing between the multiple doses may be any duration of time ranging from a few minutes to four weeks.

[0268] Thus, in another aspect, certain embodiments provide methods for treating MCT4-mediated disorders in a human or animal subject in need of such treatment comprising administering to said subject an amount of a compound disclosed herein effective to reduce or prevent said disorder in the subject, in combination with at least one additional agent for the treatment of said disorder that is known in the art. In a related aspect, certain embodiments provide therapeutic compositions comprising at least one compound disclosed herein in combination with one or more additional agents for the treatment of MCT4-mediated disorders.

[0269] The present disclosure provides compounds and pharmaceutical compositions that inhibit glutaminase activity, particularly MCT4 activity and are thus useful in the treatment or prevention of disorders associated with MCT4. Compounds and pharmaceutical compositions of the present disclosure selectively modulate MCT4 and are thus useful in the treatment or prevention of a range of disorders associated with MCT4 and include, but are not limited to, proliferative and inflammatory diseases.

[0270] Accordingly, provided herein is a method for inhibiting activity of the monocarboxylate transporter MCT4, or a mutant thereof, in a biological sample comprising the step of contacting said biological sample with a compound as disclosed herein, or a salt thereof.

[0271] Also provided herein is a method for inhibiting activity of the monocarboxylate transporter MCT4, or a mutant thereof, in a patient comprising the step of administering to the patient a compound as disclosed herein, or a salt thereof.

[0272] Also provided herein is a method for selectively inhibiting activity of the monocarboxylate transporter MCT4, or a mutant thereof, over the monocarboxylate transporter MCT1, or a mutant thereof, in a patient comprising the step of administering to the patient a compound as disclosed herein, or a salt thereof.

[0273] In certain embodiments, the inhibition is at least 100-fold selective for MCT4 over MCT1.

[0274] In certain embodiments, the compounds and pharmaceutical compositions of the present disclosure may be useful in the treatment or prevention of cancer.

[0275] In certain embodiments, the compounds of the present disclosure may be used to prevent or treat cancer, wherein the cancer is one or a variant of Acute Lymphoblastic Leukemia (ALL), Acute Myeloid Leukemia (AML), Adrenocortical Carcinoma, AIDS-Related Cancers (Kaposi Sarcoma and Lymphoma), Anal Cancer, Appendix Cancer, Atypical Teratoid/Rhabdoid Tumor, Basal Cell Carcinoma, Bile Duct Cancer (including Extrahepatic), Bladder Cancer, Bone Cancer (including Osteosarcoma and Malignant Fibrous Histiocytoma), Brain Tumor (such as Astrocytomas, Brain and Spinal Cord Tumors, Brain Stem Glioma, Central Nervous System Atypical Teratoid/Rhabdoid Tumor, Central Nervous System Embryonal Tumors, Craniopharyngioma, Ependymoblastoma, Ependymoma, Medulloblastoma, Medulloepithelioma, Pineal Parenchymal Tumors of Intermediate Differentiation, Supratentorial Primitive Neuroectodermal Tumors and Pineoblastoma), Breast Cancer, Bronchial Tumors, Burkitt Lymphoma, Carcinoid Tumor, Carcinoma of Unknown Primary, Central Nervous System (such as Atypical Teratoid/Rhabdoid Tumor, Embryonal Tumors and Lymphoma), Cervical Cancer, Childhood Cancers, Chordoma, Chronic Lymphocytic Leukemia (CLL), Chronic Myelogenous Leukemia (CML), Chronic Myeloproliferative Disorders, Colon Cancer, Colorectal Cancer, Craniopharyngioma, Cutaneous T-Cell Lymphoma (Mycosis Fungoides and Sézary Syndrome), Duct, Bile (Extrahepatic), Ductal Carcinoma In Situ (DCIS), Embryonal Tumors (Central Nervous System), Endometrial Cancer, Ependymoblastoma, Ependymoma, Esophageal Cancer, Esthesioneuroblastoma, Ewing Sarcoma Family of Tumors, Extracranial Germ Cell Tumor, Extragonadal Germ Cell Tumor, Extrahepatic Bile Duct Cancer, Eye Cancer (like Intraocular Melanoma, Retinoblastoma), Fibrous Histiocytoma of Bone (including Malignant and Osteosarcoma) Gallbladder Cancer, Gastric (Stomach) Cancer, Gastrointestinal Carcinoid Tumor, Gastrointestinal Stromal Tumors (GIST), Germ Cell Tumor (Extracranial, Extragonadal, Ovarian), Gestational Trophoblastic Tumor, Glioma, Hairy Cell Leukemia, Head and Neck Cancer, Heart Cancer, Hepatocellular (Liver) Cancer, Histiocytosis, Langerhans Cell, Hodgkin Lymphoma, Hypopharyngeal Cancer, Intraocular Melanoma, Islet Cell Tumors (Endocrine, Pancreas), Kaposi Sarcoma, Kidney (including Renal Cell), Langerhans Cell Histiocytosis, Laryngeal Cancer, Leukemia (including Acute Lymphoblastic (ALL), Acute Myeloid (AML), Chronic Lymphocytic (CLL), Chronic Myelogenous (CML), Hairy Cell), Lip and Oral Cavity Cancer, Liver Cancer (Primary), Lobular Carcinoma In Situ (LCIS), Lung Cancer (Non-Small Cell and Small Cell), Lymphoma (AIDS-Related, Burkitt, Cutaneous T-Cell (Mycosis Fungoides and Sézary Syndrome), Hodgkin, Non-Hodgkin,

Primary Central Nervous System (CNS), Macroglobulinemia, Waldenström, Male Breast Cancer, Malignant Fibrous Histiocytoma of Bone and Osteosarcoma, Medulloblastoma, Medulloepithelioma, Melanoma (including Intraocular (Eye)), Merkel Cell Carcinoma, Mesothelioma (Malignant), Metastatic Squamous Neck Cancer with Occult Primary, Midline Tract Carcinoma Involving NUT Gene, Mouth Cancer, Multiple Endocrine Neoplasia Syndromes, Multiple Myeloma/Plasma Cell Neoplasm, Mycosis Fungoides, Myelodysplastic Syndromes, Myelodysplastic/Myeloproliferative Neoplasms, Myelogenous Leukemia, Chronic (CML), Myeloid Leukemia, Acute (AML), Myeloma and Multiple Myeloma, Myeloproliferative Disorders (Chronic), Nasal Cavity and Paranasal Sinus Cancer, Nasopharyngeal Cancer, Neuroblastoma, Non-Hodgkin Lymphoma, Non-Small Cell Lung Cancer, Oral Cancer, Oral Cavity Cancer, Lip and, Oropharyngeal Cancer, Osteosarcoma and Malignant Fibrous Histiocytoma of Bone, Ovarian Cancer (such as Epithelial, Germ Cell Tumor, and Low Malignant Potential Tumor), Pancreatic Cancer (including Islet Cell Tumors), Papillomatosis, Paraganglioma, Paranasal Sinus and Nasal Cavity Cancer, Parathyroid Cancer, Penile Cancer, Pharyngeal Cancer, Pheochromocytoma, Pineal Parenchymal Tumors of Intermediate Differentiation, Pineoblastoma and Supratentorial Primitive Neuroectodermal Tumors, Pituitary Tumor, Plasma Cell Neoplasm/Multiple Myeloma, Pleuropulmonary Blastoma, Pregnancy and Breast Cancer, Primary Central Nervous System (CNS) Lymphoma, Prostate Cancer, Rectal Cancer, Renal Cell (Kidney) Cancer, Renal Pelvis and Ureter, Transitional Cell Cancer, Retinoblastoma, Rhabdomyosarcoma, Salivary Gland Cancer, Sarcoma (like Ewing Sarcoma Family of Tumors, Kaposi, Soft Tissue, Uterine), Sézary Syndrome, Skin Cancer (such as Melanoma, Merkel Cell Carcinoma, Nonmelanoma), Small Cell Lung Cancer, Small Intestine Cancer, Soft Tissue Sarcoma, Squamous Cell Carcinoma, Squamous Neck Cancer with Occult Primary, Metastatic, Stomach (Gastric) Cancer, Supratentorial Primitive Neuroectodermal Tumors, T-Cell Lymphoma (Cutaneous, Mycosis Fungoides and Sézary Syndrome), Testicular Cancer, Throat Cancer, Thymoma and Thymic Carcinoma, Thyroid Cancer, Transitional Cell Cancer of the Renal Pelvis and Ureter, Trophoblastic Tumor (Gestational), Unknown Primary, Unusual Cancers of Childhood, Ureter and Renal Pelvis, Transitional Cell Cancer, Urethral Cancer, Uterine Cancer, Endometrial, Uterine Sarcoma, Waldenström Macroglobulinemia or Wilms Tumor.

[0276] In certain embodiments, the cancer to be treated is one specific to T-cells such as T-cell lymphoma and lymphoblastic T-cell leukemia.

[0277] In certain embodiments, methods described herein are used to treat a disease condition comprising administering to a subject in need thereof a therapeutically effective amount of a compound of Formula I or pharmaceutically acceptable salt thereof, wherein the condition is cancer which has developed resistance to chemotherapeutic drugs and/or ionizing radiation.

[0278] In certain embodiments, the compounds and pharmaceutical compositions of the present disclosure may be useful in the treatment or prevention of an inflammatory disease.

[0279] In certain embodiments, the compounds of the present disclosure may be used to prevent or treat inflammatory disease, wherein the inflammatory disease is one or a variant of acid-induced lung injury, acne (PAPA), acute respiratory distress syndrome, Addison's disease, adrenal hyperplasia, adrenocortical insufficiency, ageing, AIDS, alcoholic hepatitis, alcoholic liver disease, allergen induced asthma, allergic bronchopulmonary aspergillosis, allergic conjunctivitis, alopecia, Alzheimer's disease, amyloidosis, amyotrophic lateral sclerosis, angina pectoris, angioedema, anhidrotic ectodermal dysplasia (e.g. with immune deficiency), ankylosing spondylitis, anterior segment inflammation, antiphospholipid syndrome, aphthous stomatitis, appendicitis, asthma, atherosclerosis, atopic dermatitis, autoimmune diseases, autoimmune hepatitis, bee sting-induced inflammation, Behcet's disease, Bell's Palsy, berylliosis, Blau syndrome, bone pain, bronchiolitis, burns, bursitis, cardiac hypertrophy, carpal tunnel syndrome, catabolic disorders, cataracts, cerebral aneurysm, chemical irritant-induced inflammation, chorioretinitis, chronic heart failure, chronic lung disease of prematurity, chronic obstructive pulmonary disease, colitis, complex regional pain syndrome, connective tissue disease, corneal ulcer, Crohn's disease, cryopyrin-associated periodic syndromes, cryptococcosis, cystic fibrosis, deficiency of the interleukin-1-receptor antagonist, dermatitis, dermatitis endotoxemia, dermatomyositis, endometriosis, endotoxemia, epicondylitis, erythroblastopenia, familial amyloidotic polyneuropathy, familial cold urticaria, familial Mediterranean fever, fetal growth retardation, glaucoma, glomerular disease, glomerular nephritis, gout, gouty arthritis, graft-versus-host disease, gut diseases, head injury, headache, hearing loss, heart disease, hemolytic anemia, Henoch-Scholein purpura, hepatitis, hereditary periodic fever syndrome, herpes zoster and simplex, HIV-1, Huntington's disease, hyaline membrane disease, hyperammonemia, hypercalcemia, hypercholesterolemia, hyperimmunoglobulinemia D with recurrent fever, hypoplastic and other anemias, idiopathic pulmonary fibrosis, idiopathic thrombocytopenic purpura, incontinentia pigmenti, infectious mononucleosis, inflammatory bowel disease, inflammatory

lung disease, inflammatory neuropathy, inflammatory pain, insect bite-induced inflammation, iritis, ischemia/reperfusion, juvenile rheumatoid arthritis, keratitis, kidney disease, kidney injury caused by parasitic infections, kidney transplant rejection prophylaxis, leptospirosis, Loeffler's syndrome, lung injury, lupus, lupus nephritis, meningitis, mesothelioma, mixed connective tissue disease, Muckle-Wells syndrome (urticaria deafness amyloidosis), multiple sclerosis, muscle wasting, muscular dystrophy, myasthenia gravis, myocarditis, mycosis fungoides, myelodysplastic syndrome, myositis, nasal sinusitis, necrotizing enterocolitis, neonatal onset multisystem inflammatory disease (NOMID), nephrotic syndrome, neuritis, neuropathological diseases, non-allergen induced asthma, obesity, ocular allergy, optic neuritis, organ transplant, osteoarthritis, otitis media, Paget's disease, pain, pancreatitis, Parkinson's disease, pemphigus, pericarditis, periodic fever, periodontitis, pertussis, perineal or peritoneal endometriosis, pharyngitis and adenitis (PFAPA syndrome), plant irritant-induced inflammation, pneumocystis infection, pneumonia, pneumonitis, poison ivy/urushiol oil-induced inflammation, polyarteritis nodosa, polychondritis, polycystic kidney disease, polymyositis, psoriasis, psychosocial stress disease, pulmonary disease, pulmonary fibrosis, pulmonary hypertension, pyoderma gangrenosum, pyogenic sterile arthritis, renal disease, retinal disease, rheumatic disease, rheumatoid arthritis, rheumatic carditis, sarcoidosis, seborrhea, sepsis, severe pain, sickle cell, sickle cell anemia, silica-induced diseases, Sjogren's syndrome, skin diseases, sleep apnea, spinal cord injury, Stevens-Johnson syndrome, stroke, subarachnoid hemorrhage, sunburn, systemic sclerosis (scleroderma), temporal arteritis, tenosynovitis, thrombocytopenia, thyroiditis, tissue transplant, TNF receptor associated periodic syndrome (TRAPS), Toxoplasmosis, transplant, traumatic brain injury, tuberculosis, type 1 diabetes, type 2 diabetes, ulcerative colitis, urticaria, uveitis, Wegener's granulomatosis, and weight loss.

[0280] Thus, in another aspect, certain embodiments provide methods for treating a monocarboxylate transporter MCT4-mediated disorder in a subject in need thereof, comprising the step of administering to said patient a compound as disclosed herein, or a pharmaceutically acceptable salt thereof.

[0281] In certain embodiments, the subject is a human.

[0282] In certain embodiments, the monocarboxylate transporter MCT4-mediated disorder is chosen from an inflammatory disorder and a proliferative disorder.

[0283] In certain embodiments, the monocarboxylate transporter MCT4-mediated disorder is a proliferative disorder.

[0284] In certain embodiments, the proliferative disorder is cancer.

[0285] In certain embodiments, the cancer is chosen from adenocarcinoma, adult T-cell leukemia/lymphoma, bladder cancer, blastoma, bone cancer, breast cancer, brain cancer, carcinoma, myeloid sarcoma, cervical cancer, colorectal cancer, esophageal cancer, gastrointestinal cancer, glioblastoma multiforme, glioma, gallbladder cancer, gastric cancer, head and neck cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, intestinal cancer, kidney cancer, laryngeal cancer, leukemia, lung cancer, lymphoma, liver cancer, small cell lung cancer, non-small cell lung cancer, mesothelioma, multiple myeloma, ocular cancer, optic nerve tumor, oral cancer, ovarian cancer, pituitary tumor, primary central nervous system lymphoma, prostate cancer, pancreatic cancer, pharyngeal cancer, renal cell carcinoma, rectal cancer, sarcoma, skin cancer, spinal tumor, small intestine cancer, stomach cancer, T-cell lymphoma, testicular cancer, thyroid cancer, throat cancer, urogenital cancer, urothelial carcinoma, uterine cancer, vaginal cancer, and Wilms' tumor.

[0286] In certain embodiments, the monocarboxylate transporter MCT4-mediated disorder is an inflammatory disorder.

[0287] In certain embodiments, the inflammatory disorder is chosen from Crohn's disease, ulcerative colitis, idiopathic pulmonary fibrosis, muscular dystrophy, rheumatoid arthritis, and systemic sclerosis (scleroderma).

[0288] Also provided herein is a compound as disclosed herein, or a pharmaceutically acceptable salt thereof, for use in human therapy.

[0289] Also provided herein is a compound as disclosed herein, or a pharmaceutically acceptable salt thereof, for use in treating a monocarboxylate transporter MCT4-mediated disorder, for example as disclosed in any of the embodiments and paragraphs above pertaining to methods of treatment.

[0290] Also provided herein is the use of a compound as disclosed herein, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament to treat a monocarboxylate transporter MCT4-mediated disorder, for example as disclosed in any of the embodiments and paragraphs above pertaining to methods of treatment.

[0291] Metabolic syndrome (also known as metabolic syndrome X) is characterized by having at least three of the following symptoms: insulin resistance; abdominal fat - in men this is defined as a 40 inch waist or larger, in women 35 inches or larger; high blood sugar levels - at least 110 milligrams per deciliter (mg/dL) after fasting; high triglycerides - at least 150 mg/dL in the blood stream; low HDL- less than 40 mg/dL; pro-thrombotic state (e.g.

high fibrinogen or plasminogen activator inhibitor in the blood); or blood pressure of 130/85 mmHg or higher. A connection has been found between metabolic syndrome and other conditions such as obesity, high blood pressure and high levels of LDL cholesterol, all of which are risk factors for cardiovascular diseases. For example, an increased link between metabolic syndrome and atherosclerosis has been shown. People with metabolic syndrome are also more prone to developing type 2 diabetes, as well as PCOS (polycystic ovarian syndrome) in women and prostate cancer in men.

[0292] As described above, insulin resistance can be manifested in several ways, including type 2 diabetes. Type 2 diabetes is the condition most obviously linked to insulin resistance. Compensatory hyperinsulinemia helps maintain normal glucose levels- often for decades, before overt diabetes develops. Eventually the beta cells of the pancreas are unable to overcome insulin resistance through hypersecretion. Glucose levels rise, and a diagnosis of diabetes can be made. Patients with type 2 diabetes remain hyperinsulinemic until they are in an advanced stage of disease. As described above, insulin resistance can also correlate with hypertension. One half of patients with essential hypertension are insulin resistant and hyperinsulinemic, and there is evidence that blood pressure is linked to the degree of insulin resistance. Hyperlipidemia, too, is associated with insulin resistance. The lipid profile of patients with type 2 diabetes includes increased serum very-low-density lipoprotein cholesterol and triglyceride levels and, sometimes, a decreased low-density lipoprotein cholesterol level. Insulin resistance has been found in persons with low levels of high-density lipoprotein. Insulin levels have also been linked to very-low-density lipoprotein synthesis and plasma triglyceride levels.

[0293] Accordingly, also disclosed are methods of treating insulin resistance in a subject comprising selecting a subject in need of treatment for insulin resistance; and administering to the subject an effective amount of a compound that inhibits MCT4.

[0294] Specific diseases to be treated by the compounds, compositions, and methods disclosed herein are those mediated at least in part by MCT4. Accordingly, disclosed herein are methods: for reducing glycogen accumulation in a subject; for raising HDL or HDLc, lowering LDL or LDLc, shifting LDL particle size from small dense to normal LDL, lowering VLDL, lowering triglycerides, or inhibiting cholesterol absorption in a subject; for reducing insulin resistance, enhancing glucose utilization or lowering blood pressure in a subject; for reducing visceral fat in a subject; for reducing serum transaminases in a subject; or for treating disease; all comprising the administration of a therapeutic amount of a

compound as described herein, to a patient in need thereof. In further embodiments, the disease to be treated may be a metabolic disease. In further embodiment, the metabolic disease may be selected from the group consisting of: obesity, diabetes melitus, especially Type 2 diabetes, hyperinsulinemia, glucose intolerance, metabolic syndrome X, dyslipidemia, hypertriglyceridemia, hypercholesterolemia, and hepatic steatosis. In other embodiments, the disease to be treated may be selected from the group consisting of: cardiovascular diseases including vascular disease, atherosclerosis, coronary heart disease, cerebrovascular disease, heart failure and peripheral vessel disease. In preferred embodiments, the methods above do not result in the induction or maintenance of a hypoglycemic state.

[0295] Besides being useful for human treatment, certain compounds and formulations disclosed herein may also be useful for veterinary treatment of companion animals, exotic animals and farm animals, including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.


List of Abbreviations

[0296] Ac₂O = acetic anhydride; AcCl = acetyl chloride; AcOH = acetic acid; AIBN = azobisisobutyronitrile; aq. = aqueous; BAST = bis(2-methoxyethyl)aminosulfur trifluoride; BCECF = 2',7'-bis-(carboxyethyl)-5(6)-carboxyfluorescein; Bu = butyl; Bu₃SnH = tributyltin hydride; CD₃OD = deuterated MeOH; CDCl₃ = deuterated chloroform; CDI = 1,1'-carbonyldiimidazole; DAST = (diethylamino)sulfur trifluoride; dba = dibenzylideneacetone; DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene; DCM = dichloromethane; DEAD = diethyl azodicarboxylate; DIAD = diisopropyl azodicarboxylate; DIBAL-H = di-iso-butyl aluminium hydride; DIEA = DIPEA = N,N-diisopropylethylamine; DMAP = 4-dimethylaminopyridine; DMF = N,N-dimethylformamide; DMSO-d₆ = deuterated dimethyl sulfoxide; DMSO = dimethyl sulfoxide; DPPA = diphenylphosphoryl azide; dppf = 1,1'-bis(diphenylphosphino)ferrocene; EDC•HCl = EDCI•HCl = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride; Et = ethyl; Et₂O = diethyl ether; EtOAc = ethyl acetate; EtOH = ethanol; h = hour; HATU=2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium; HMDS = hexamethyldisilazane; HOBT = 1-hydroxybenzotriazole; i-Pr = isopropyl = 2-propyl; i-PrOH = isopropanol; LAH = lithium aluminium hydride; LDA =lithium diisopropyl amide; LiHMDS = Lithium bis(trimethylsilyl)amide; MeCN = acetonitrile; MeI = methyl iodide; MeOH = MeOH; MP-carbonate resin = macroporous triethylammonium methylpolystyrene carbonate resin; MsCl

= mesyl chloride; MTBE = methyl tertiary butyl ether; n-BuLi = n-butyllithium; NaHMDS = sodium bis(trimethylsilyl)amide; NaOEt = sodium ethoxide; NaOMe = sodium methoxide; NaOtBu = sodium t-butoxide; NBS = N-bromosuccinimide; NCS = N-chlorosuccinimide; NIS = N-iodosuccinimide; NMP = N-Methyl-2-pyrrolidone; Pd(PPh₃)₄ = tetrakis-(triphenylphosphine)palladium(0); Pd₂(dba)₃ = tris(dibenzylideneacetone)dipalladium(0); PdCl₂(PPh₃)₂ = bis(triphenylphosphine)palladium(II) dichloride; PG = protecting group; Ph = phenyl; prep-HPLC = preparative high-performance liquid chromatography; PMB = para-methoxybenzyl; PMBCl = para-methoxybenzyl chloride; PMBOH = para-methoxybenzyl alcohol; PyBop = (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate; Pyr = pyridine; RT = room temperature; RuPhos = 2-dicyclohexylphosphino-2',6'-diisopropoxybiphenyl; sat. = saturated; ss = saturated solution; tBu = t-Bu = tert-butyl = 1,1-dimethylethyl; TBAF = tetrabutylammonium fluoride; TBDPS = t-butyldiphenylsilyl; t-BuOH = tert-butanol; T3P = Propylphosphonic Anhydride; TEA = Et₃N = triethylamine; TFA = trifluoroacetic acid; TFAA = trifluoroacetic anhydride; THF = tetrahydrofuran; TIPS = triisopropylsilyl; Tol = toluene; TsCl = tosyl chloride; Trt = trityl = (triphenyl)methyl; Xantphos = 4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene; XPhos = 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl.


General Synthetic Methods for Preparing Compounds

[0297] The following schemes can be used to practice the present invention.




[0298] A general, but not intended to be limiting, synthetic scheme for the examples in this disclosure is depicted in Scheme I. An appropriately substituted 2,4-diketobutanoic ester is reacted with a substituted hydrazine to form the pyrazole core. Reduction of the ester gives a hydroxymethyl compound. The side chain can be completed by direct alkylation with a 2-haloester, as shown in pathway (a). Alternatively, the hydroxymethyl functionality is converted to a halomethyl group, which can undergo displacement with a 2-hydroxyester, as shown in pathway (b), or be transformed via Wittig chemistry, as shown in pathway (c) to give an alkene.

Scheme II




[0299] Scheme II, depicted above, may be used to prepare certain examples disclosed herein. In step (a), organohalide **201** is converted to an organolithium, followed by condensation with the substituted acetamide to give acetyl functionalized compound **202**. This compound is reacted in step (b) under Claisen conditions with an oxalate ester to give 2,4-diketoester **203**, which is then reacted with a substituted hydrazine in step (c) to construct the pyrazole core of **204**. The ester functionality is reduced in step (d), and in step (e) the resulting alcohol **205** is alkylated with an appropriate haloacetate ester, affording ether **206**. Finally, the ester is hydrolyzed in step (f) under basic conditions to give carboxylic acid **207**. In the scheme as depicted, R_{100} - R_{103} will be understood by one of skill in the art to be any appropriate group. For example, in certain embodiments, R_{100} and R_{101} may be independently chosen from aryl, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted. Likewise, in certain embodiments, R_{102} and R_{103} may be independently chosen from hydrogen, alkyl, and halogen. R_{100} - R_{103} may also correspond to the groups defined in Formula I, II, or any other formula disclosed herein. Finally, this scheme may be joined at any point by employing a suitable intermediate shown herein that is available either from commercial sources or alternate synthetic methods.

Scheme III




[0300] Scheme III, depicted above, can be used to prepare certain example compounds disclosed herein. Steps (a) – (d) from Scheme I are used to obtain pyrazolemethanol **301**, which is then converted to bromomethyl compound **302** in step (e), using one of various techniques available for this transformation. Alkyl halide **302** is then reacted under Williamson ether conditions with a hydroxyester in step (f), to give **303**. Synthesis is then completed in step (g) by ester hydrolysis to **304**.

Scheme IV




[0301] Scheme IV, depicted above, can be utilized for certain example compounds disclosed herein. Steps (a) – (d) from Scheme I are used to obtain pyrazolemethanol **401**, which is alkylated to give tBu ester **402**. Cleavage of the ester is accomplished under acidic conditions, to give acid **403**.

Scheme V




[0302] Scheme V, depicted above, can be utilized for certain example compounds disclosed herein. Steps (a) – (e) from Scheme III are used to obtain (bromomethyl)pyrazole **501**. Reaction with a hydroxyester under basic conditions gives tBu ester **502**. Cleavage of the ester is accomplished under acidic conditions, to give acid **503**.

Scheme VI




[0303] Scheme VI, depicted above, may be use to prepare certain example compounds disclosed herein. Phenylacetone is condensed with oxalate ester to give the 5-phenylpentanoate ester **601**. Condensation with substituted hydrazine forms the pyrazole core of **602**. Ester reduction, alcohol alkylation, and ester hydrolysis proceeds as before. Synthesis of acid **603** is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.

Scheme VII



[0304] Scheme VII, depicted above, can be used to prepare certain example compounds disclosed herein. Acetylated hydroxyaryl compound **701** reacts under Williamson ether conditions with an alkyl halide or similar compound, giving alkoxyaryl compound **702**. Condensation with oxalate ester gives a 2,4-diketoester **703**, as in Scheme II. Synthesis is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.


Scheme VIII



[0305] Scheme VIII, depicted above, can be used to synthesize certain example compounds disclosed herein. Bromoaryl compound **801** is converted to methoxyaryl compound **802** in the presence of a suitable metal catalyst in the presence of base. The methyl ether is cleaved with a Lewis acid. The resulting hydroxyl compound is reacted under Williamson ether conditions to give alkoxyaryl compound **803**. Alternatively, the bromoaryl compound is reacted directly with an alcohol of interest to afford the desired ether **803** directly. In the scheme as depicted, R_{100} - R_{106} will be understood by one of skill in the art to


be any appropriate group. Synthesis of **804** is completed by either using steps (c) – (f) from Scheme II, or steps (c) – (g) from Scheme III.

Scheme IX



[0306] Scheme IX, depicted above, can be used to synthesize certain example compounds disclosed herein. Bromoaryl compound **901** is converted to the (bromomethyl) pyrazole **904**. The compound is reacted under Williamson ether conditions with a suitable glycolic ester to give ether **905**. Transesterification to the methyl ester **905** is followed by displacement of the aryl bromide using an appropriate Pd^0 catalyst. Synthesis is completed by basic hydrolysis of the methyl ester, affording acid **906**.


Scheme X



[0307] Scheme X, depicted above, can be used to synthesize certain example compounds disclosed herein. Chloroaryl carboxylic ester **1001** is coupled with an organozinc reagent in


the presence of copper(I) iodide to give **1002**. The ester functionality is converted to methyl ketone **1003**. Condensation with diethyl oxalate gives the β -diketone compound. Synthesis of **1004** is completed by either using steps (c) – (f) from Scheme II, or steps (c) – (g) from Scheme III.

Scheme XI




[0308] Scheme XI, depicted above, can be used to synthesize certain example compounds disclosed herein. Pyrazole ester **1101** is obtained via steps (a) – (c) of Scheme II. The compound is coupled under Mitsunobu conditions with an appropriate alcohol to give alkylated product **1102**. Synthesis of acid **1103** is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.

Scheme XII




[0309] Scheme XII, depicted above, can be used to synthesize certain example compounds disclosed herein. Pyrazole ester **1201** is obtained via steps (a) – (c) of Scheme II. The compound is coupled with an alkyl halide in the presence of base to give **1202**. Synthesis of acid **1203** is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.

Scheme XIII




[0310] Scheme XIII, depicted above, may be used to prepared certain example compounds disclosed herein. A 2,4-diketoester is condensed with a nitroaromatic compound, such as 4-nitroaniline, to give pyrazole moiety **1301**. Reduction of the nitro group is followed by reductive amination to give a (dialkyl)amino compound **1302**. Synthesis of acid **1303** is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.

Scheme XIV




[0311] Scheme XIV, depicted above, can be used to synthesize certain example compounds disclosed herein. A 2,4-diketoester is condensed with hydrazine to form pyrazole **1401**, which is then alkylated with a substituted benzyl bromide to afford **1402**. Reduction to the amine is followed by reductive alkylation to give substituted amine **1403**. Synthesis of acid **1404** is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.




[0312] Scheme XV, depicted above, can be used to synthesize certain example compounds disclosed herein. (Bromomethyl) pyrazole **1501**, obtained via steps (a) – (e) of Scheme II, is converted to phosphonium salt **1502**. Treatment with base forms phosphorane ylid **1503**; Wittig reaction with an appropriate aldehyde gives alkene **1504**. The alkene can be optionally hydrogenated to give alkane **1506**. Synthesis is completed by ester hydrolysis give alkene **1505** or alkane **1507**, respectively.

Scheme XVI




[0313] Scheme XVI, depicted above, may be used to prepare certain example compounds disclosed herein. Acetonitrile is condensed with an oxalate ester to give a ketoester product, which can be isolated as its salt **1601**. Neutralization of the salt is followed by condensation with a substituted hydrazine to give a 5-aminopyrazole compound **1602**. The amino functionality is then substituted by reaction with an organohalide in a palladium (0) mediated reaction, affording **1603**, and then protected as its Boc carbamate **1604**. Ester reduction, alcohol alkylation, and ester hydrolysis proceeds as before to give acid **1605**. The Boc protecting group is removed under acidic conditions, affording **1606**.

Scheme XVII




[0314] Scheme XVII, depicted above, may be use to prepare certain example compounds disclosed herein. 2-Ketosuccinate ester is condensed with a substituted hydrazine to give 5-hydroxypyrazole compound **1701**, shown here as its keto tautomer. The hydroxy group is alkylated with an electrophilic arene, such as 2-fluoronitrobenzene, giving ether **1702**. The nitro functionality is removed by stepwise reduction with zinc(0) to amine **1703**, followed by deamination with alkyl nitrite, yielding **1704**. Synthesis of acid **1705** is completed by either using steps (d) – (f) from Scheme II, or steps (d) – (g) from Scheme III.

Scheme XVIII




[0315] Scheme XVIII, depicted above, may be used to prepare certain example compounds disclosed herein. Nitro acetophenone **1801** is reacted under Claisen conditions with an oxalate ester to give 2,4-diketoester **1802**, which is then reacted with a substituted hydrazine in step (c) to construct the pyrazole core of **1803**, which is alkylated to provide **1804**. The ester functionality is reduced to the alcohol (not shown) and alkylated to give ester **1805**. At this point, the nitro group is reduced with zinc to give amine **1806**, and acylated using standard coupling reagents to give amide **1807**. Synthesis is completed with basic hydrolysis of the ester, affording carboxylic acid **1808**.

Scheme XIX




[0316] Scheme XIX, depicted above, may be used to prepare certain example compounds disclosed herein. In pathway (a), amine **1806**, prepared using the method of Scheme XVIII, is treated with an organonitrite in the presence of a disulfide to afford a thioether, which is oxidized to sulfone **1901**. In pathway (b), amine **1806** is reacted with a sulfonyl chloride to give sulfonamide **1903**. In pathway (c), amine **1806** is treated with an aldehyde under reductive amination conditions to give amine **1905**. In pathways (a), (b), and (c), synthesis is completed with basic hydrolysis of the ester, affording carboxylic acid **1902**, **1904**, and **1906**, respectively.

Scheme XX




[0317] Scheme XX, depicted above, may be use to prepare certain example compounds disclosed herein. Benzyl ether **206** is prepared by using steps (a) – (e) of Scheme II. Treatment with acid cleaves both the ether group and the methyl ester to give acid **2001**. If desired, synthesis can be continued by Fischer esterification to **2002**, alkylation of the phenol under basic conditions, and hydrolysis of the methyl ester to give acid **2003**.

Scheme XXI




[0318] Scheme XXI, depicted above, may be use to prepare certain example compounds disclosed herein. Carboxylic acid **207** is prepared by using steps (a) – (f) of Scheme II. Treatment of the acid with an alkylsulfonamide give ester **2101**.

Scheme XXII




[0319] Scheme XXII, depicted above, may be used to prepare certain example compounds disclosed herein. Aryllithium **2201**, prepared by lithiation, halogen-metal exchange, or other methods that are available in the art, is reacted with N-methyl-N-methoxyacetamide, which affords acetyl compound **2202**, which can be transformed to the desired product through any of the previous schemes.

Scheme XXIII




[0320] Scheme XXIII, depicted above, may be used to prepare certain example compounds disclosed herein. Carboxylic acid **2301** is converted to the corresponding alkoxy amide **2302** using standard amide formation procedures. The alkoxy amide is then reacted with an appropriate organolithium reagent, which affords ketone compound **2303**, which can be transformed to the desired product through any of the previous schemes.

Scheme XXIV




[0321] Scheme XXIV, depicted above, may be used to prepare certain example compounds disclosed herein. Aryl halide **2401**, prepared using any of the methods disclosed herein, is converted to the corresponding arylamine **2402**. This transformation can be accomplished with a copper-based catalyst, or other techniques known in the art. Synthesis can then be completed using methods disclosed herein.




Scheme XXV


[0322] Scheme XXV, depicted above, may be used to prepare certain example compounds disclosed herein. Aryl halide **2501**, prepared using any of the methods disclosed herein, is converted to the corresponding arylamide **2502**. This transformation can be accomplished with a copper-based catalyst, or other techniques known in the art. Synthesis can then be completed using methods disclosed herein.

Scheme XXVI

[0323] Scheme XXVI, depicted above, may be used to prepare certain example compounds disclosed herein. Aryl halide **2601**, prepared using any of the methods disclosed herein, is converted to the corresponding aryl nitrile **2602**. This transformation can be accomplished with a Pd-based catalyst, or other techniques known in the art. Synthesis can then be completed using methods disclosed herein.

Scheme XXVII

[0324] Scheme XXVII, depicted above, may be used to prepare certain example compounds disclosed herein. Carboxylic acid **2701**, prepared using any of the methods disclosed herein, is converted to the corresponding carboxylate salt **2702**. This transformation can be accomplished with a metal hydroxide or metal oxide. The metal can be chosen from the alkali metals, alkaline earth metals, or other metals.




Scheme XXVIII

[0325] Scheme XXVIII, depicted above, may be used to prepare certain example compounds disclosed herein. An aryl or heteroaryl halide **2801** is converted to the corresponding hydrazine **2802**. Synthesis of the pyrazole core is accomplished via condensation with an appropriate carbonyl compound, as disclosed herein. Amine functionality in the aryl or heteroaryl moiety of **2801** can be protected by attachment of a suitable protecting group, such as the SEM protecting group. The protecting group can be removed at a convenient point in synthesis by using methods known in the art.

Intermediates

[0326] The following intermediates were synthesized for use in preparing the compounds of this disclosure.


Intermediate A-1: 1-(1-Methyl-1H-indol-6-yl)ethan-1-one



[0327] **6-Bromo-1-methyl-1H-indole** A mixture of 6-bromo-1H-indole (5 g, 25.50 mmol, 1.00 equiv), K₂CO₃ (7 g, 50.65 mmol, 2.00 equiv), and CH₃I (7.3 g, 51.43 mmol, 2.00 equiv) in acetone (100 mL) was stirred for 16 h at 55 °C, then cooled to rt, diluted with 100 mL H₂O, and extracted with 2x200 mL of EtOAc. The combined organic layers were washed with 100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:10) to afford 3.4 g (63%) of the title compound as a yellow oil. LC-MS: (ES, *m/z*): 208. ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.69 (dt, *J* = 1.6, 0.7 Hz, 1H), 7.49 (dd, *J* = 8.4, 0.6 Hz, 1H), 7.34 (d, *J* = 3.1 Hz, 1H), 7.13 (dd, *J* = 8.4, 1.8 Hz, 1H), 6.43 (dd, *J* = 3.1, 0.9 Hz, 1H), 3.77 (s, 3H).

[0328] **1-(1-Methyl-1H-indol-6-yl)ethan-1-one** To a solution of 6-bromo-1-methyl-1H-indole (3.4 g, 16.19 mmol, 1.00 equiv) in THF (100 mL) was added BuLi (12 mL, 2.00 equiv, 2.5 M) dropwise with stirring at -78 °C. The solution was stirred for 1 h, then *N*-methoxy-*N*-methylacetamide (4.1 g, 39.76 mmol, 2.50 equiv) was added dropwise with stirring at -78 °C. The solution was stirred for 30 min, then warmed to rt and stirred an additional 2 h. The reaction mixture was cooled to 0 °C, then quenched by the addition of 1 M HCl and extracted with 2x200 mL of EtOAc. The combined organic layers were washed with 100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:5) to afford 1.9 g (68%) of the title compound as a yellow solid. LC-MS: (ES, *m/z*): 173. ¹H NMR: (300 MHz, DMSO-*d*₆) δ 8.12 (dt, *J* = 1.5, 0.8 Hz, 1H), 7.70 – 7.59 (m, 2H), 7.57 (d, *J* = 3.0 Hz, 1H), 6.51 (dd, *J* = 3.0, 0.9 Hz, 1H), 3.88 (s, 3H), 2.63 (s, 3H).

Intermediate A-2: 1-(1-Methyl-1H-indazol-6-yl)ethan-1-one



[0329] **1-(1-methyl-1H-indazol-6-yl)ethan-1-one** To a solution of 6-bromo-1-methyl-1H-indazole (2.5 g, 11.85 mmol, 1.00 equiv) in THF (50 mL) under N₂ at -78 °C was

added n-BuLi (10 mL, 2.00 equiv) dropwise with stirring. The solution was stirred for 60 min at this temperature, then *N*-methoxy-*N*-methylacetamide (3 g, 29.09 mmol, 2.50 equiv) was added, and the solution was stirred for an additional 60 min at this temperature. The resulting solution was stirred for 1 h at rt, cooled to 0 °C, quenched by the addition of HCl (1M), and extracted with 2x100 mL of EtOAc. The combined organic layers were washed with 50 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (1:10) to afford 1.5 g (73%) of the title compound as a yellow oil. LC-MS: (ES, *m/z*): 174. ¹H NMR: (400 MHz, DMSO-*d*₆) δ 8.36 (q, *J* = 1.1 Hz, 1H), 8.16 (d, *J* = 1.0 Hz, 1H), 7.86 (dd, *J* = 8.5, 0.8 Hz, 1H), 7.69 (dd, *J* = 8.5, 1.4 Hz, 1H), 4.16 (s, 3H), 2.70 (s, 3H).


[0330] The following acetyl arenes were obtained from similar routes.

Table 1: Synthesis of acetyl arenes from aryl bromides.

Aryl bromide	Product		Spectral
1-bromo-3-(propan-2-yl)benzene	1-[3-(Propan-2-yl)phenyl]-ethan-1-one	A-3	
6-bromo-1-ethyl-1 <i>H</i> -indazole	1-(1-Ethyl-1 <i>H</i> -indazol-6-yl)-ethan-1-one	A-4	LC-MS: (ES, <i>m/z</i>): 189.
1-bromo-3-chloro-5-methoxybenzene	1-(3-chloro-5-methoxy-phenyl)ethan-1-one	A-5	LC-MS: (ES, <i>m/z</i>): 182.
5-bromo-1-methyl-1 <i>H</i> -indazole	1-(1-methyl-1 <i>H</i> -indazol-5-yl)ethan-1-one	A-6	LC-MS: (ES, <i>m/z</i>): 174.
4-bromo-1-methyl-1 <i>H</i> -indazole	1-(1-methyl-1 <i>H</i> -indazol-4-yl)ethan-1-one	A-7	LC-MS: (ES, <i>m/z</i>): 174.
6-bromo-1-ethyl-1 <i>H</i> -1,2,3-benzotriazole	1-(1-ethyl-1 <i>H</i> -1,2,3-benzotriazol-6-yl)ethan-1-one	A-8	LC-MS: (ES, <i>m/z</i>) 189.

[0331] The following acetyl arene intermediates were prepared via Scheme XXII.


Intermediate A-9: 1-(3-Cyclobutoxyphenyl)ethan-1-one



[0332] To a mixture of 1-(3-hydroxyphenyl)ethan-1-one (4.5 g, 33.05 mmol, 1.00 equiv) and Cs₂CO₃ (10.78 g, 33.09 mmol, 1.00 equiv) in DMA (50 mL) was added dropwise bromocyclobutane (9.0 g, 66.67 mmol, 2.00 equiv) with stirring at rt. The resulting mixture

was stirred for 16 h at 130 °C, then cooled and diluted with 200 mL of EtOAc. The combined organic layers were washed with 2x200 mL of H₂O and 2x200 mL of brine, then dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:10) to afford 5.0 g (80%) of the title compound as colorless oil. ¹H NMR: (300 MHz, DMSO-*d*₆) δ 7.53 (ddd, *J* = 7.6, 1.7, 1.1 Hz, 1H), 7.41 (t, *J* = 7.9 Hz, 1H), 7.32 (dd, *J* = 2.6, 1.6 Hz, 1H), 7.10 (ddd, *J* = 8.1, 2.7, 1.0 Hz, 1H), 4.83 – 4.67 (m, 1H), 2.55 (s, 3H), 2.47 – 2.31 (m, 2H), 2.14 – 1.94 (m, 2H), 1.88 – 1.72 (m, 1H), 1.72 – 1.56 (m, 1H).

Intermediate A-10: 1-[3-(Oxetan-3-ylmethoxy)phenyl]ethan-1-one



[0333] Into a flask under N₂ were combined a solution of oxetan-3-ylmethanol (3.24 g, 36.77 mmol, 1.00 equiv) in THF (100 mL), 1-(3-hydroxyphenyl)ethan-1-one (5 g, 36.72 mmol, 1.00 equiv), and PPh₃ (14.5 g, 1.50 equiv), followed by the addition of DEAD (8.3 g, 47.66 mmol, 1.30 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 12 h at rt, then extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified by silica gel chromatography with EtOAc /petroleum ether (0-20%) to afford 4.4 g (58%) of the title compound as light yellow oil. LC-MS: (ES, *m/z*): 206.90. H-NMR: δ_H (300 MHz, DMSO-*d*₆) 7.56 (1 H, ddd, *J* = 7.6, 1.6, 1.0 Hz), 7.45 (2 H, m), 7.23 (1 H, ddd, *J* = 8.2, 2.7, 1.1 Hz), 4.71 (2 H, dd, *J* = 7.9, 6.0 Hz), 4.44 (2 H, t, *J* = 6.0, 6.0 Hz), 4.26 (2 H, d, *J* = 6.7 Hz), 3.39 (1 H, m), 2.57 (3 H, s).

[0334] The following alkylated compounds were obtained from Williamson synthesis of 3-hydroxyacetophenone.





Table 2: Williamson synthesis of alkoxy acetophenones.

Alkyl halide	Base / solvent	Temp	Product		Spectral
3-bromo-oxetane	Cs ₂ CO ₃ / DMA	130 °C	1-[3-(Oxetan-3-yl-oxy)phenyl]ethan-1-one	A-11	LC-MS: (ES, <i>m/z</i>): 193.
Bromocyclopropane	Cs ₂ CO ₃ / DMA	150 °C	1-(3-cyclopropoxy-phenyl)ethan-1-one	A-12	


1-Bromo-2,2-dimethylpropane	K ₂ CO ₃ / DMF	110 °C	1-[3-(2,2-Dimethylpropoxy)phenyl]ethan-1-one	A-13	LC-MS: (ES, <i>m/z</i>): 206.9.
(Bromo-methyl)-cyclobutane	K ₂ CO ₃ / DMF	110 °C	1-[3-(Cyclobutylmethoxy)phenyl]ethan-1-one	A-14	LC-MS: (ES, <i>m/z</i>): 291.25.
(Bromo-methyl)-cyclopropane	K ₂ CO ₃ / DMF	80 °C	1-[3-(Cyclopropylmethoxy)phenyl]ethan-1-one	A-15	LC-MS: (ES, <i>m/z</i>): 191.
Iodoethane	K ₂ CO ₃ / DMF	90 °C	1-(3-Ethoxyphenyl)ethan-1-one	A-16	LC-MS: (ES, <i>m/z</i>): 165.
Benzyl bromide	K ₂ CO ₃ / DMF	80 °C	1-[3-(benzyloxy)phenyl]ethan-1-one	A-17	LC-MS: (ES, <i>m/z</i>): 226.95.
1-Bromo-2-methylpropane	K ₃ PO ₄ / DMSO	60 °C	1-[3-(2-Methylpropoxy)phenyl]ethan-1-one	A-18	

Intermediate A-19: 1-(3,5-Dimethoxyphenyl)ethan-1-one



[0335] In a flask were combined 1-(3,5-dihydroxyphenyl)ethan-1-one (2 g, 13.15 mmol, 1.00 equiv) iodomethane (6.5 g, 45.79 mmol, 4.00 equiv), K₂CO₃ (3.6 g, 25.86 mmol, 2.00 equiv) and acetone (40 mL). The resulting mixture was stirred for 16 h at 56 °C, then cooled to rt, diluted with 100 mL of water, and extracted with 2x50 mL of EtOAc. The combined organic layers were washed with 20 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (1:10), to afford 2 g (84%) of the title product as colorless oil. LC-MS: (ES, *m/z*): 181. ¹H NMR: (300 MHz, DMSO-*d*₆) δ 7.07 (d, *J* = 2.3 Hz, 2H), 6.77 (t, *J* = 2.3 Hz, 1H), 3.89 – 3.71 (m, 6H), 2.56 (s, 3H).


Intermediate A-20: 1-(3,5-Diethoxyphenyl)ethan-1-one



[0336] In a flask were combined 1-(3,5-dihydroxyphenyl)ethan-1-one (5 g, 32.86 mmol, 1.00 equiv), iodoethane (20 g, 128.23 mmol, 4.00 equiv), K₂CO₃ (18 g, 130.43 mmol, 4.00 equiv), and acetone (100 mL). The resulting mixture was heated to reflux overnight, then

cooled to rt and extracted with 3x100 mL of EtOAc. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum to afford 6 g (88%) of the title compound as a brown solid. LC-MS: (ES, *m/z*): 208.95.


Intermediate A-21: 1-(3-Ethoxy-5-methoxyphenyl)ethan-1-one



[0337] **1-(3-Hydroxy-5-methoxyphenyl)ethan-1-one** To a mixture of 1-(3,5-dihydroxyphenyl)ethan-1-one (5.0 g, 32.86 mmol, 1.00 equiv), K₂CO₃ (6.85 g, 49.56 mmol, 1.50 equiv), and acetone (100 mL) was added MeI (4.67 g, 32.89 mmol, 1.00 equiv) dropwise with stirring at rt. The resulting solution was heated to reflux for 2 h, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (1:3) to afford 2 g (37%) of the title compound as a white solid. LC-MS: (ES, *m/z*): 167.

[0338] **1-(3-Ethoxy-5-methoxyphenyl)ethan-1-one** To a mixture of 1-(3-hydroxy-5-methoxyphenyl)ethan-1-one (2 g, 12.04 mmol, 1.00 equiv), K₂CO₃ (2.5 g, 18.09 mmol, 1.50 equiv), and DMF (50 mL) was added EtI (2.8 g, 17.95 mmol, 1.00 equiv) dropwise with stirring at rt. The resulting solution was stirred for 16 h at 90 °C, diluted with 100 mL of EtOAc, washed with 1x100 mL of water and 1x100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (1:5) to afford 950 mg (41%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 195. ¹H-NMR: (DMSO, *ppm*): δ: 7.06 (dq, *J* = 2.9, 1.4 Hz, 2H), 6.75 (t, *J* = 2.3 Hz, 1H), 4.07 (q, *J* = 7.0 Hz, 2H), 2.56 (s, 3H), 1.34 (t, *J* = 6.9 Hz, 3H).

Intermediate A-22: 1-[2-(Propan-2-yloxy)-1,3-oxazol-5-yl]ethan-1-one




[0339] **2-Hydroxy-1,3-oxazole-5-carboxylic acid** A solution of ethyl 2-chloro-1,3-oxazole-5-carboxylate (3 g, 17.09 mmol, 1.00 equiv) and LiOH (1.7 g, 70.98 mmol, 4.00 equiv) in THF/H₂O (60/15 mL) was stirred for 16 h at 30 °C. The pH was then adjusted to 4 with 1 M HCl, and the resulting solution was extracted with 4x50 mL of EtOAc. The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under vacuum to afford 2 g (91%) of the title compound as a yellow solid. LC-MS: (ES, *m/z*): 129.

[0340] **2-Hydroxy-*N*-methoxy-*N*-methyl-1,3-oxazole-5-carboxamide** A solution of *N*-methoxy-*N*-methylamine hydrochloride (1.8 g, 18.45 mmol, 1.20 equiv), HATU (7.6 g, 20.54 mmol, 1.30 equiv), DIEA (6 g, 46.43 mmol, 3.00 equiv), and 2-hydroxy-1,3-oxazole-5-carboxylic acid (2 g, 15.50 mmol, 1.00 equiv) in DMF (60 mL) was stirred for 5 h at room temperature. The resulting mixture was concentrated under vacuum and purified with silica gel chromatography using CH₂Cl₂ / MeOH (50:1) to afford 1.3 g (49%) of the title compound as a yellow solid. LC-MS: (ES, *m/z*): 173.

[0341] ***N*-methoxy-*N*-methyl-2-(propan-2-yloxy)-1,3-oxazole-5-carboxamide** A mixture of 2-hydroxy-*N*-methoxy-*N*-methyl-1,3-oxazole-5-carboxamide (1.8 g, 10.46 mmol, 1.00 equiv), 2-iodopropane (3.6 g, 2.00 equiv), K₂CO₃ (2.9 g, 2.00 equiv) and DMF (15 mL) in a 20-mL sealed tube was stirred for 16 h at 90 °C. The reaction mixture was cooled to rt, diluted with 20 mL of H₂O, and extracted with 4x20 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using CH₂Cl₂ / MeOH (100:1) to afford 1.5 g (67%) of the title compound as a yellow solid. LC-MS: (ES, *m/z*): 214.

[0342] **1-[2-(Propan-2-yloxy)-1,3-oxazol-5-yl]ethan-1-one** To a solution of *N*-methoxy-*N*-methyl-2-(propan-2-yloxy)-1,3-oxazole-5-carboxamide (1.5 g, 7.00 mmol, 1.00 equiv) in THF (40 mL) under N₂ at -78 °C was added in portions MeMgBr (21 mL, 3.00 equiv). The resulting solution was stirred for 2 h at rt, cooled to 0 °C, quenched by the addition of HCl (1M), and extracted with 2x50 mL of EtOAc. The combined organic layers were washed with 30 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (1:10) to afford 670 mg (57%) of the title compound as a yellow solid. LC-MS: (ES, *m/z*): 169.

Intermediate A-23:1-[1,3-dimethyl-1H-thieno[2,3-c]pyrazol-5-yl]ethan-1-one




[0343] ***N*-methoxy-*N*,1,3-trimethyl-1H-thieno[2,3-c]pyrazole-5-carboxamide** A solution of 1,3-dimethyl-1H-thieno[2,3-c]pyrazole-5-carboxylic acid (2 g, 10.19 mmol, 1.00 equiv) methoxy(methyl)amine hydrochloride (1.06 g, 10.87 mmol, 1.10 equiv), DIEA (4 g, 30.95 mmol, 3.00 equiv), and HATU (5.8 g, 15.26 mmol, 1.50 equiv) in DMF (30 mL) was stirred for 2 h at rt, diluted with 100 mL of water, and extracted with 2x100 mL of EtOAc. The combined organic layers were washed with 100 mL brine, dried over Na₂SO₄,

concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:3) to afford 2.2 g (90%) of the title compound as a yellow solid. LC-MS: (ES, m/z): 240.

[0344] **1-[1,3-dimethyl-1H-thieno[2,3-c]pyrazol-5-yl]ethan-1-one** To a solution of the product from the previous step (2.2 g, 9.19 mmol, 1.00 equiv) in THF (50 mL) at -78 °C under N₂ was added methylmagnesium bromide (1.0 M in THF, 29 mL, 3.00 equiv) dropwise with stirring. The resulting solution was stirred for 2 h at room temperature, then cooled to 0 °C and then quenched by the addition of aq HCl. The resulting solution was diluted with 100 mL H₂O and extracted with 2x200 mL of EtOAc. The combined organic layers were washed with 50 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel column chromatography using EtOAc / petroleum ether (1:3) to afford 1.4 g (78%) of the title compound as a white solid.


[0345] LC-MS: (ES, m/z): 195. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.04 (s, 1H), 3.86 (s, 3H), 2.52 (s, 4H), 2.39 (s, 3H).

Intermediate A-24: 1-(5-Methoxythien-2-yl)ethan-1-one



[0346] A mixture of 1-(5-chlorothiophen-2-yl)-1-ethanone (11 g, 96.35 mmol, 1.00 equiv), Cs₂CO₃ (22 g, 1.20 equiv), and Brettphos Pd G3 (550 mg) in MeOH (60 mL) was heated in a sealed tube for 1 hr at 100 °C with microwave radiation. The resulting mixture was concentrated under vacuum, then extracted with 3x100 mL of EtOAc. The combined organic layers were dried over anhydrous MgSO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1/50) to afford 5 g (33%) of the title product as a yellow solid.


Intermediate A-25: 1-[2-(2-Methylpropoxy)-1,3-thiazol-5-yl]ethan-1-one



[0347] To a solution of 1-(2-bromo-1,3-thiazol-5-yl)ethan-1-one (2 g, 9.71 mmol, 1.00 equiv) in 2-methylpropan-1-ol (15 mL) in a 40 mL sealed tube under N₂ was added sequentially Cs₂CO₃ (6.4 g, 19.64 mmol, 2.00 equiv) and Brettphos Pd G3 (440 mg, 0.49 mmol, 0.05 equiv). The resulting solution was stirred for 1 h at 90 °C. The solution was cooled, diluted with 50 mL H₂O, and extracted with 2x100 mL of EtOAc. The combined

organic layers were washed with 50 mL of aq NaCl, dried over anhydrous Na₂SO₄, concentrated under vacuum, and purified with silica gel column using EtOAc / petroleum ether (1:10) to afford 550 mg (28%) of the title compound as a yellow oil.


Intermediate A-26: 1-[5-(2-Methylpropoxy)thien-2-yl]ethan-1-one



[0348] **1-(5-Hydroxythien-2-yl)ethan-1-one** To a solution of 1-(5-Methoxy-thien-2-yl)ethan-1-one was added BBr₃ (180 mL) dropwise with stirring at rt. The resulting solution was stirred for 2 days at 35 °C. The reaction mixture was cooled with a water/ice bath. The reaction was then quenched by the addition of 500 mL of water/ice. The resulting solution was diluted with 500 mL of CH₂Cl₂. The pH value of the solution was adjusted to 7-8 with (sat) NaHCO₃ (aq) (500 mol/L). The mixture was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:10). This resulted in the title compound as a red oil (5.0 g, 61%).

[0349] **1-[5-(2-Methylpropoxy)thien-2-yl]ethan-1-one** To a solution of the product from the previous step (800 mg, 5.63 mmol, 1.00 equiv) in DMF (20 mL) was added Cs₂CO₃ (3.67 g, 11.26 mmol, 2.00 equiv), followed by the addition of 1-iodo-2-methylpropane (2.07 g, 11.25 mmol, 2.00 equiv) dropwise with stirring at rt. The resulting solution was stirred for 16 h at 90 °C. The resulting solution was diluted with 100 mL of EtOAc then washed with 2x100 mL of water and 2x100 mL of brine. The mixture was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:10). This resulted in the title compound as a yellow oil (600 mg, 54%).


Intermediate A-27: 1-(4-Methoxythien-2-yl)ethan-1-one



[0350] To a solution of 1-(4-bromothien-2-yl)ethan-1-one (3.0 g, 14.63 mmol, 1.00 equiv) in MeOH (25 mL) was added NaOMe (5.4 mL, 2.00 equiv, 5.4M) at room temperature, followed by CuBr (627 mg, 4.38 mmol, 0.30 equiv) at room temperature. The resulting solution was stirred for 16 h at 100 °C, then diluted with 100 mL of EtOAc, washed with 2x100 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The

residue was purified with silica gel chromatography using EtOAc / petroleum ether (1:20) to afford 700 mg (31%) of the title product as a yellow oil.


Intermediate A-28: 1-[2-(2-Methylpropyl)-1,3-oxazol-5-yl]ethan-1-one



[0351] **Ethyl 2-(2-methylpropyl)-1,3-oxazole-5-carboxylate** To a solution of ethyl 2-chloro-1,3-oxazole-5-carboxylate (3 g, 17.09 mmol, 1.00 equiv) in THF (45 mL), under N₂, was added copper(I) iodide (1.5 g, 7.88 mmol, 0.50 equiv). This was followed by the addition of bromo(2-methylpropyl)zinc (15 mL, 2.00 equiv) at -5 °C. The resulting solution was stirred for 3 h at room temperature. The reaction was then quenched by the addition of aq 1 M HCl. The resulting solution was extracted with 2x100 mL of EtOAc, and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:5). This resulted in the title compound as a red oil (1.3g, 39%).


[0352] **2-(2-Methylpropyl)-1,3-oxazole-5-carboxylic acid** To a solution of the product from the previous step (1.3 g, 6.59 mmol, 1.00 equiv) in THF / H₂O (20/5 mL) was added LiOH (600 mg, 25.05 mmol, 4.00 equiv). The resulting solution was stirred for 3 h at room temperature. The reaction mixture was cooled with a water/ice bath. The pH value of the solution was adjusted to 1 with aq 1 M HCl. The resulting solution was extracted with 2x50 mL of EtOAc, and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. This resulted in the title compound as a red solid (1 g, 90%).

[0353] **1-[2-(2-Methylpropyl)-1,3-oxazol-5-yl]ethan-1-one** To a solution of the product from the previous step (1 g, 5.91 mmol, 1.00 equiv) in Et₂O (30 mL), under N₂, was added MeLi (7.4 mL, 2.00 equiv). The resulting solution was stirred for 3 h at 0 °C in a water/ice bath. The reaction was then quenched by the addition of aq HCl (1M). The resulting solution was extracted with 2x50 mL of EtOAc, and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:5). This resulted in the title compound as a red oil (800 mg, 81%).

Intermediate A-29: 1-(1-methyl-1H-1,3-benzodiazol-6-yl)ethan-1-one


[0354] **N-methoxy-N,1-dimethyl-1H-1,3-benzodiazole-6-carboxamide** To a solution of 1-methyl-1H-1,3-benzodiazole-6-carboxylic acid (5.0 g, 28.38 mmol, 1.00 equiv) in CH₂Cl₂ (100 mL) was added methoxy(methyl)amine hydrochloride (2.78 g, 28.50 mmol, 1.00 equiv), EDC (6.54 g, 42.13 mmol, 1.20 equiv), followed by the addition of DMAP (3.46 g, 28.32 mmol, 1.00 equiv), in portions at rt. The resulting solution was stirred for 16 h, concentrated under vacuum, and purified with silica gel chromatography using CH₂Cl₂ / MeOH (100:1) to afford 4.0 g (64%) of the title compound as a yellow solid. LC-MS: (ES, m/z): [M+1] = 220.

[0355] **1-(1-methyl-1H-1,3-benzodiazol-6-yl)ethan-1-one** To a solution of the product from the previous step (2.0 g, 9.12 mmol, 1.00 equiv) in THF (60 mL) under N₂ was added MeMgBr (9.1 mL, 3.00 equiv) dropwise with stirring at -60 °C. The resulting mixture was stirred at this temperature for 30 min, allowed to warm to rt, then stirred for 2 h at rt. The reaction was then quenched by the addition of water/ice. The resulting solution was extracted with 100 mL of EtOAc, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using CH₂Cl₂ / MeOH (50:1) to afford 1.3 g (82%) of the title compound as a yellow solid.


Intermediate A-30: 3-Acetyl-N-(propan-2-yl)benzamide

[0356] To a suspension of 3-acetylbenzoic acid (10 g, 60.92 mmol), HATU (20 g, 52.60 mmol,) and propan-2-amine (4 g, 67.67 mmol) in DMF (180 mL) at -5 °C was added dropwise DIEA. The resulting mixture was stirred at room temperature for 2 h under N₂, then poured into 100 mL H₂O, and extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with 300 mL brine, dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. This resulted in 11 g (88%) of the title product as a yellow solid.


[0357] The following 2,4-dioxobutanoate esters were synthesized for use in preparing the compounds of this disclosure, including, but not limited to, use in Scheme I.

Intermediate B-1: Methyl 4-(2-methoxyphenyl)-2,4-dioxobutanoate

[0358] To a solution of 1-(2-methoxyphenyl)ethan-1-one (1 g, 6.66 mmol, 1.00 equiv) in MeOH (20 mL) was added sequentially dimethyl oxalate (2.36 g, 19.99 mmol, 3.00 equiv), then MeONa (3.7 mL, 3.00 equiv, 5.4 M). The resulting solution was stirred overnight at 30 °C then quenched by the addition of water/ice. The pH value of the solution was adjusted to 5 with aq 1 M HCl. The resulting solution was extracted with 3x50 mL of EtOAc, and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / hexane (1/4). This resulted in the title compound as a yellow solid (1.2g, 76%).

Intermediate B-2: Methyl 4-(3-methoxyphenyl)-2,4-dioxobutanoate

[0359] To a solution of 1-(3-methoxyphenyl)ethan-1-one (10 g, 66.59 mmol, 1.00 equiv) in MeOH (200 mL) was added sequentially NaOMe (30 mL), and dimethyl oxalate (10.2 g, 86.38 mmol, 1.3 equiv). The resulting solution was stirred for 8 h at room temperature. The reaction mixture was cooled to 0 °C with a water/ice bath. The resulting solution was diluted with 200 mL of EtOAc. The pH value of the solution was adjusted to 3 with aq HCl (3 mol/L). The resulting solution was extracted with 2x300 mL of EtOAc, and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. This resulted in the title compound as a yellow solid (11 g, 70%).

Intermediate B-3: Methyl 4-[3-(oxetan-3-ylmethoxy)phenyl]-2,4-dioxobutanoate

[0360] In a 100-mL flask were combined a solution of 1-[3-(oxetan-3-ylmethoxy)phenyl]ethan-1-one (Int. A-10, 3 g, 14.55 mmol, 1.00 equiv) in MeOH (40 mL), dimethyl oxalate (3.42 g, 28.96 mmol, 2.00 equiv), and a solution of MeONa (2.36 g, 43.70 mmol, 3.00 equiv) in MeOH (8 mL). The resulting solution was stirred for 12 h at rt, then poured onto ice/water. The pH was adjusted to 3 with HCl. The solids that formed were collected by filtration, affording 3.2 g (75%) of the title compound as a light yellow solid. LC-MS:(ES,

m/z): 293.0. ¹H-NMR: δ_H (300 MHz, DMSO-*d*₆) 7.68 (1 H, d, *J*= 7.7 Hz), 7.53 (2 H, m), 7.30 (1 H, m), 7.14 (1 H, s), 4.72 (2 H, dd, *J* =7.9, 6.0 Hz), 4.45 (2 H, t, *J* 6.0, 6.0 Hz), 4.30 (2 H, d, *J* =6.7 Hz), 3.86 (3 H, s), 3.40 (1H, tt, *J*= 8.0, 8.0, 6.3, 6.3 Hz).

[0361] The following substituted 4-aryl-2,4-dioxobutanoate esters were obtained from Claisen condensation of an 1-arylethanone with an oxalic ester.





Table 3: Claisen synthesis of 4-aryl-2,4-dioxobutanoate esters.

R _B OH / time / temp	Aryl ethanone	Product 4-aryl-2,4-dioxobutanoate ester		Spectral
MeOH / 12 hr / rt	1-(3-nitrophenyl)- ethan-1-one	Methyl 4-(3-nitrophenyl)-2,4- dioxobutanoate	B-4	
MeOH / 12 hr / rt	1-(Pyridin-2- yl)ethan-1-one	Methyl 2,4-dioxo-4-(pyridin-2- yl)butanoate	B-5	LC-MS: (ES, <i>m/z</i>): 291.25.
MeOH / 12 hr / 30 °C	1-(2,3-Dihydro-1,4- benzodioxin-6-yl)- ethan-1-one	Methyl 4-(2,3-dihydro-1,4- benzodioxin-6-yl)-2,4- dioxobutanoate	B-6	LC-MS: (ES, <i>m/z</i>): 265.
MeOH / 16 hr / rt	1-(2 <i>H</i> -1,3-Benzo- dioxol-5-yl)ethan- 1-one	Methyl 4-(2 <i>H</i> -1,3-benzodioxol- 5-yl)-2,4-dioxobutanoate	B-7	LC-MS: (ES, <i>m/z</i>): 251.
MeOH / 16 hr / 40 °C	1-(1-Methyl-1 <i>H</i> - pyrazol-4-yl)ethan- 1-one	Methyl 4-(1-methyl-1 <i>H</i> -pyrazol- 4-yl)-2,4-dioxobutanoate	B-8	LC-MS: (ES, <i>m/z</i>): 211
EtOH / 5 hr / rt	1-(1-Benzothio- phen-2-yl)ethan-1- one	Ethyl 4-(1-benzothiophen-2-yl)- 2,4-dioxobutanoate	B-9	LC-MS: (ES, <i>m/z</i>): 277
MeOH / 2 hr / 60 °C	1-(4-Bromothien-2- yl)ethanone	Methyl 4-(4-bromothien-2-yl)- 2,4-dioxobutanoate	B-10	
EtOH / 16 hr / rt	A-1	Ethyl 4-(1-methyl-1 <i>H</i> -indol-6- yl)-2,4-dioxobutanoate	B-11	LC-MS: (ES, <i>m/z</i>): 273.
EtOH / 16 hr / rt	A-2	Ethyl 4-(1-methyl-1 <i>H</i> -indazol-6- yl)-2,4-dioxobutanoate	B-12	LC-MS: (ES, <i>m/z</i>): 274.
MeOH / 16 hr / rt	A-3	Methyl 4-[3-(propan-2-yl)- phenyl]2,4-dioxobutanoate	B-13	
MeOH / 16 hr / rt	A-4	Methyl 4-(1-ethyl-1 <i>H</i> -indazol-6- yl)-2,4-dioxobutanoate	B-14	LC-MS: (ES, <i>m/z</i>): 275.
EtOH / 16 hr / rt	A-4	Ethyl 4-(1-ethyl-1 <i>H</i> -indazol-6- yl)-2,4-dioxobutanoate	B-15	LC-MS: (ES, <i>m/z</i>): 288.

MeOH / 16 hr / rt	A-5	Methyl 4-(3-chloro-5-methoxyphenyl)-2,4-dioxobutanoate	B-16	LC-MS: (ES, <i>m/z</i>): 271
EtOH / 16 hr / rt	A-6	Ethyl 4-(1-methyl-1 <i>H</i> -indazol-5-yl)-2,4-dioxobutanoate	B-17	LC-MS: (ES, <i>m/z</i>): 274.
EtOH / 16 hr / rt	A-7	Ethyl 4-(1-methyl-1 <i>H</i> -indazol-4-yl)-2,4-dioxobutanoate	B-18	LC-MS: (ES, <i>m/z</i>): 274.
MeOH / 16 h / rt	A-8	Methyl 4-(1-ethyl-1 <i>H</i> -1,2,3-benzotriazol-6-yl)-2,4-dioxobutanoate	B-19	LC-MS: (ES, <i>m/z</i>): 275
EtOH / 16 hr / rt	A-9	Ethyl 4-(3-cyclobutoxyphenyl)-2,4-dioxobutanoate	B-20	LC-MS: (ES, <i>m/z</i>): 291.
EtOH / 12 hr / rt	A-11	Ethyl 4-[3-(oxetan-3-yloxy)-phenyl]-2,4-dioxobutanoate	B-21	LC-MS: (ES, <i>m/z</i>): 293.
MeOH / 12 hr / rt	A-12	Methyl 4-(3-cyclopropoxy-phenyl)-2,4-dioxobutanoate	B-22	LC-MS: (ES, <i>m/z</i>): 262.85.
MeOH / 12 hr / rt	A-13	Methyl 4-[3-(2,2-dimethylpropoxy)phenyl]-2,4-dioxobutanoate	B-23	LC-MS: (ES, <i>m/z</i>): 292.95.
MeOH / 12 hr / rt	A-14	Methyl 4-[3-(cyclobutyl-methoxy)phenyl]-2,4-dioxobutanoate	B-24	LC-MS: (ES, <i>m/z</i>): 290.95.
MeOH / 16 hr / rt	A-15	Methyl 4-[3-(cyclopropyl-methoxy)phenyl]-2,4-dioxobutanoate	B-25	LC-MS: (ES, <i>m/z</i>): 276.
EtOH / 16 hr / rt	A-16	Ethyl 4-(3-ethoxyphenyl)-2,4-dioxobutanoate	B-26	LC-MS: (ES, <i>m/z</i>): 264.
MeOH / 12 hr / rt	A-17	Methyl 4-[3-(benzyloxy)-phenyl]-2,4-dioxobutanoate	B-27	LC-MS: (ES, <i>m/z</i>): 313.1.
MeOH / 16 hr / rt	A-18	Methyl 4-[3-(2-methylpropoxy)-phenyl]-2,4-dioxobutanoate	B-28	LC-MS: (ES, <i>m/z</i>): 279.
MeOH/ 16 hr / rt	A-19	Methyl 4-(3,5-dimethoxyphenyl)-2,4-dioxobutanoate	B-29	LC-MS: (ES, <i>m/z</i>): 267.
MeOH / 12 hr / rt	A-20	Methyl 4-(3,5-diethoxyphenyl)-2,4-dioxobutanoate	B-30	LC-MS: (ES, <i>m/z</i>): 295.2.
MeOH / 16 hr / rt	A-21	Methyl 4-(3-ethoxy-5-methoxyphenyl)-2,4-dioxobutanoate	B-31	LC-MS: (ES, <i>m/z</i>): 281
EtOH / 5 hr / rt	A-22	Ethyl 2,4-dioxo-4-[2-(propan-2-yloxy)-1,3-oxazol-5-yl]-butanoate	B-32	LC-MS: (ES, <i>m/z</i>): 270


MeOH / 16 h / rt	A-23	Methyl 4-[1,3-dimethyl-1H-thieno[2,3-c]pyrazol-5-yl]-2,4-dioxobutanoate	B-33	LC-MS: (ES, m/z): 281
MeOH / 16 h / rt	A-24	Methyl 4-(5-2-methoxythien-2-yl)-2,4-dioxobutanoate	B-34	
MeOH / 16 h / rt	A-26	Methyl 4-(5-(2-methylpropoxy)-thien-2-yl)-2,4-dioxobutanoate	B-35	
EtOH / 5 h / rt	A-28	4-[2-(2-Methylpropyl)-1,3-oxazol-5-yl]-2,4-dioxobutanoate	B-36	
MeOH / 16 h / rt	A-29	Methyl 4-(1-methyl-1H-1,3-benzodiazol-6-yl)-2,4-dioxobutanoate	B-37	LC-MS: (ES, m/z): [M+1] = 261
MeOH / 16 h / rt	A-30	Methyl 4-[3-[(propan-2-yl)-carbamoyl]phenyl]-2,4-butanedioate	B-38	

Intermediate B-39: Ethyl 4-[2-(2-methylpropoxy)-1,3-thiazol-5-yl]-2,4-dioxobutanoate



[0362] To a solution of Int. **A-25** (500 mg, 2.51 mmol, 1.00 equiv) in THF (10 mL) was added sequentially diethyl oxalate (1100 mg, 7.53 mmol, 2.00 equiv), and t-BuOK (850 mg, 7.58 mmol, 3.00 equiv). The resulting solution was stirred for 4 h at room temperature followed by dilution with 20 mL of H₂O, extraction with 2x30 mL of EtOAc and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:4). This resulted in the title compound as a yellow solid (400 mg, 53%)

Intermediate B-40: Ethyl 2,4-dioxo-5-phenylpentanoate



[0363] To a solution of 1-phenyl-2-propanone (1 g, 8.32 mmol, 1.00 equiv) in THF (25 mL) was added NaH (269 mg, 11.21 mmol, 1.50 equiv), in portions at 0 °C in 30 min, followed by the addition of diethyl oxalate (1.64 g, 11.22 mmol, 1.50 equiv). The resulting solution was stirred for 1 h at room temperature, then cooled to 0 °C. The reaction was then quenched by the addition of 20 mL of water. The resulting solution was extracted with 3x50 mL of EtOAc, and the combined organic layers were dried over anhydrous Na₂SO₄,

concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:40) to afford 700 mg (36%) of the title product as a yellow liquid.


[0364] The following substituted ketoesters were also used in the synthesis of pyrazoles.

Intermediate **B-41**: Methyl 2,4-dioxo-4-(1-propyl-1H-indazol-6-yl)butanoate

Intermediate **B-42**: Methyl 4-(3-cyclobutoxyphenyl)-2,4-dioxobutanoate

[0365] The following arylhydrazine intermediates were synthesized for use in preparing the compounds of this disclosure, including, but not limited to, use in Scheme I.

Intermediate C-1: 2-(Difluoromethoxy)phenyl]hydrazine trifluoroacetate



[0366] **Boc-protected 2-(difluoromethoxy)phenyl]hydrazine** To a solution of 1-bromo-2-(difluoromethoxy)benzene (8.0 g, 35.87 mmol, 1.00 equiv), (tert-butoxy)-carbohydrazide (5.68 g, 42.98 mmol, 1.20 equiv), and BrettPhos Pd G3 (1.74 g, 1.92 mmol, 0.05 equiv) in dioxane (60 mL) under N₂ was added Cs₂CO₃ (11.70 g, 35.91 mmol, 1.00 equiv), in portions at room temperature. The resulting solution was stirred for 16 h at 100 °C, then cooled, diluted with 200 mL of EtOAc, washed with 200 mL of brine, dried over Na₂SO₄ and concentrated under vacuum. The crude product was purified with silica gel chromatography using EtOAc /petroleum ether (1:5) to afford 8.0 g (81%) of the title compound as an orange solid. LC-MS: (ES, *m/z*): 275. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.83 (s, 1H), 7.29 – 7.22 (m, 1H), 7.07 (q, *J* = 7.7 Hz, 2H), 6.80 – 6.68 (m, 2H), 1.42 (s, 9H).

[0367] **2-(Difluoromethoxy)phenyl]hydrazine trifluoroacetate** A solution of the product from the previous step (2.0 g, 7.29 mmol, 1.00 equiv) and TFA (10 mL) in CH₂Cl₂ (20 mL) was stirred for 2 h at room temperature, then concentrated under vacuum, to afford 1.0 g (48%) of the title compound as a brown oil. LC-MS: (ES, *m/z*): 175.

[0368] The following arylhydrazines were synthesized from the corresponding aryl halides using similar procedures.

Table 4: Synthesis of aryl hydrazines from aryl bromides.

[0369] The compounds were isolated as trifluoroacetate salts.

Aryl bromide	Product	Spectral
--------------	---------	----------


1-Bromo-2-ethoxy-benzene	(2-Ethoxyphenyl)hydrazine	C-2	LC-MS: (ES, m/z): 152.
7-Bromo-1-methyl-1 <i>H</i> -indazole	7-Hydrazinyl-1-methyl-1 <i>H</i> -indazole	C-3	LC-MS: (ES, m/z): 163.
4-Bromo-1-methyl-1 <i>H</i> -indazole	4-Hydrazinyl-1-methyl-1 <i>H</i> -indazole	C-4	LC-MS: (ES, m/z): 162.
7-Bromo-1-ethyl-1 <i>H</i> -indazole	1-Ethyl-7-hydrazinyl-1 <i>H</i> -indazole	C-5	LC-MS: (ES, m/z): 277.
1-Bromo-2-(propan-2-yloxy)benzene	[2-(Propan-2-yloxy)-phenyl]hydrazine	C-6	LC-MS: (ES, m/z): 274.

[0370] The free amino groups of the following reactants were first protected with SEM groups (NaH; SEM-Cl). Treatment of the coupling product with TFA for removal of the Boc groups also accomplished removal of the SEM protecting groups.

Table 5: Synthesis of aryl hydrazines from aryl bromides.

Aryl bromide	Product		Spectral
4-Bromo-1 <i>H</i> -indazole	4-Hydrazinyl-1 <i>H</i> -indazole	C-7	LC-MS: (ES, m/z): 148.9.
7-Bromo-1 <i>H</i> -indazole	7-Hydrazinyl-1 <i>H</i> -indazole	C-8	LC-MS: (ES, m/z): 379.
4-Bromo-3-methyl-1 <i>H</i> -indazole	4-Hydrazinyl-3-methyl-1 <i>H</i> -indazole	C-9	LC-MS: (ES, m/z): 163.2.
7-Bromo-1-methyl-1 <i>H</i> -1,3-benzodiazole	7-Hydrazinyl-1-methyl-1 <i>H</i> -1,3-benzodiazole trifluoroacetate	C-10	
7-Bromo-1,3-dimethyl-1 <i>H</i> -indazole	7-Hydrazinyl-1,3-dimethyl-1 <i>H</i> -indazole	C-11	LC-MS: (ES, m/z): 177.1.

Intermediate C-12: 7-Hydrazinyl-1-methyl-1*H*-1,2,3-benzotriazole trifluoroacetate



[0371] **2-Bromo-N-methyl-6-nitroaniline** A mixture of 1-bromo-2-fluoro-3-nitrobenzene (7 g, 31.82 mmol, 1.00 equiv), K₂CO₃ (8.8 g, 63.21 mmol, 2.00 equiv), and CH₃NH₂ (10 mL, 33%) in CH₂Cl₂ (50 mL) was stirred 16 h at rt, then diluted with 100 mL of water. The resulting solution was extracted with 2x200 mL of CH₂Cl₂. The combined organic layers were washed with 50 mL of brine, dried over Na₂SO₄, and concentrated under vacuum to afford 7 g (95%) of the title compound as a red oil. LC-MS: (ES, m/z): 230. ¹H NMR (400


MHz, DMSO-*d*₆) δ 7.77 (ddd, $J = 9.8, 8.0, 1.6$ Hz, 2H), 6.70 (t, $J = 8.0$ Hz, 1H), 6.43 – 6.19 (m, 1H), 2.71 (d, $J = 5.3$ Hz, 3H).

[0372] **6-Bromo-1-N-methylbenzene-1,2-diamine** A mixture of the product from the previous step (8 g, 34.62 mmol, 1.00 equiv) and Zn (4 g, 62.50 mmol, 2.00 equiv) in AcOH (120 mL) was stirred for 16 h at rt. The solids were removed by filtration, and the filtrate was concentrated under vacuum. The residue was dissolved in 300 mL of CH₂Cl₂, washed with 100 mL of NaHCO₃, dried over Na₂SO₄, and concentrated under vacuum, to afford 4 g (57%) of the title compound as a black oil. LC-MS: (ES, m/z): 201.

[0373] **7-Bromo-1-methyl-1H-1,2,3-benzotriazole** To a solution of the product from the previous step (4 g, 19.89 mmol, 1.00 equiv) in HBr / H₂O (60 mL) at 5 °C was added a solution of NaNO₂ (2.07 g, 1.50 equiv) in H₂O (10 mL) dropwise with stirring. The solution was stirred for 4 h at 15-20 °C. The pH was adjusted to 8 with NaHCO₃ (3 mol/L). The resulting solution was extracted with 2x200 mL of EtOAc. The combined organic layers were washed with 100 mL of brine, dried over anhydrous Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:5), to afford 4 g (95%) of the title compound as a black solid. LC-MS: (ES, m/z): 212. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.06 (dd, $J = 8.3, 0.8$ Hz, 1H), 7.84 – 7.73 (m, 1H), 7.31 (dd, $J = 8.4, 7.5$ Hz, 1H), 4.51 (s, 3H).


[0374] **7-Hydrazinyl-1-methyl-1H-1,2,3-benzotriazole trifluoroacetate** was prepared from the previous compound by using the procedure of **C-1**. LC-MS: (ES, m/z): 163.

Intermediate C-13: [(2-Chlorophenyl)methyl]hydrazine dihydrochloride



[0375] To a solution of hydrazine hydrate (85%) (31 g, 8.50 equiv) in EtOH (100 mL) was added dropwise a solution of 1-chloro-2-(chloromethyl)benzene (10 g, 62.10 mmol, 1.00 equiv) in EtOH (50 mL) over 1 hr at 70 °C. The resulting solution was stirred for 1 h at 70 °C, concentrated under vacuum diluted with water, and extracted with EtOAc three times. The combined organic layers were chilled to 0 °C, then a solution of 4N HCl in dioxane was added. The solids that formed were collected by filtration, affording 6.1 g (63%) of the title product as a white solid.

Intermediate D-1: Ethyl 5-[[[(tert-butoxy)carbonyl](phenyl)amino]-1-[(2-chlorophenyl)methyl]-1H-pyrazole-3-carboxylate



[0376] **Ethyl 3-cyano-2-(potassiooxy)prop-2-enoate** To a solution of diethyl oxalate (5.06 g, 34.62 mmol, 1.00 equiv) in CH₃CN (20 mL) was added t-BuOK (3.90 g, 34.76 mmol, 1.12 equiv), in portions at rt. The resulting solution was stirred for 1.5 h at rt. The solids that formed were collected by filtration. This resulted in 5.09 g (82%) of the title product as a yellow solid.


[0377] **Ethyl 5-amino-1-[(2-chlorophenyl)methyl]-1H-pyrazole-3-carboxylate** To a solution of the product from the previous step (2.0 g, 11.16 mmol, 1.00 equiv) in 1,4-dioxane (20 mL) was added TFA (2 mL) dropwise with stirring at rt. The resulting mixture was stirred at rt for 30 min. To this was added Int. **C-13** (2.0 g, 12.77 mmol, 1.15 equiv), in portions at rt. The resulting solution was stirred overnight at rt, then diluted with 200 mL of EtOAc, washed with 2x200 mL of brine, dried over anhydrous Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:3) to afford 1.8 g (58%) of the title compound as a yellow oil.

[0378] **Ethyl 1-[(2-chlorophenyl)methyl]-5-(phenylamino)-1H-pyrazole-3-carboxylate** A solution of the product from the previous step (1.0 g, 3.57 mmol, 1.00 equiv), iodobenzene (768 mg, 3.76 mmol, 1.05 equiv), Brettphos Pd G3 precatalyst (326 mg, 0.36 mmol, 0.10 equiv), and Cs₂CO₃ (1.4 g, 4.30 mmol, 1.20 equiv) in 1,4-dioxane (20 mL). The resulting solution was stirred in a sealed tube under N₂ at 90 °C for 3 h. The resulting solution was diluted with 200 mL of EtOAc, washed with 2x200 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using petroleum ether : EtOAc:DCM (3:1:0.1) to afford 1.3 g (102%) of the title product as a yellow solid.

[0379] **Ethyl 5-[[[(tert-butoxy)carbonyl](phenyl)amino]-1-[(2-chlorophenyl)methyl]-1H-pyrazole-3-carboxylate** To a solution of the product from the previous step (800 mg, 2.25 mmol, 1.00 equiv) and DMAP (548 mg, 4.49 mmol, 2.00 equiv) in toluene (20 mL) was added di-tert-butyl dicarbonate (980 mg, 4.49 mmol, 2.00 equiv), in portions at rt. The

resulting solution was heated at reflux overnight. The resulting solution was diluted with 200 mL of EtOAc, washed with 2x100 mL of brine, dried over anhydrous Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:3) to afford 1.0 g (101%) of the title product as a colorless oil.

Intermediate D-2: Methyl 5-(3-cyclopropoxyphenyl)-1H-pyrazole-3-carboxylate



[0380] A solution of Int. **B-22** (2 g, 7.63 mmol, 1.00 equiv) and hydrazine hydrate (540 mg, 10.00 mmol, 1.30 equiv) in AcOH (20 mL) was stirred for 2 h at 100 °C, then cooled. The pH was adjusted to 7 with satd NaHCO₃, and the resulting solution was extracted with 3x100 mL EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / hexane (1/1) to afford 1.5 g (76%) of the title compound as a light yellow solid. LC-MS: (ES, *m/z*): 259. ¹H NMR (300 MHz, MeOD) δ 7.35 (m, 3H), 7.13 (d, *J* = 21.3 Hz, 2H), 3.92 (s, 3H), 3.84 (dp, *J* = 6.4, 3.0, 3.0, 2.9, 2.9 Hz, 1H), 0.82 (m, 2H), 0.72 (m, 2H).

[0381] The following substituted 3-pyrazolecarboxylic esters were obtained from condensation of methyl 4-(aryl)-2,4-dioxobutanoate with hydrazine:





Table 6: Pyrazole synthesis with hydrazine.

Diketone	Product		Spectral
B-2	Methyl 5-(3-methoxyphenyl)-1H-pyrazole-3-carboxylate	D-3	
B-4	Methyl 5-(3-nitrophenyl)-1H-pyrazole-3-carboxylate	D-4	
B-6	Methyl 5-(2,3-dihydro-1,4-benzodioxin-6-yl)-1H-pyrazole-3-carboxylate	D-5	LC-MS: (ES, <i>m/z</i>): 261.
B-7	Methyl 5-(2H-1,3-benzodioxol-5-yl)-1H-pyrazole-3-carboxylate	D-6	LC-MS: (ES, <i>m/z</i>): 247.


B-8	Methyl 5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-7	LC-MS: (ES, <i>m/z</i>): 207.
B-9	Ethyl 5-(1-benzothiophen-2-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-8	LC-MS: (ES, <i>m/z</i>): 272.
B-11	Ethyl 5-(1-methyl-1 <i>H</i> -indol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-9	LC-MS: (ES, <i>m/z</i>): 269.
B-12	Ethyl 5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-10	LC-MS: (ES, <i>m/z</i>): 271.
B-14	Methyl 5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-11	LC-MS: (ES, <i>m/z</i>): 271.
B-16	Methyl 5-(3-chloro-5-methoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-12	LC-MS: (ES, <i>m/z</i>): 252.
B-17	Ethyl 5-(1-methyl-1 <i>H</i> -indazol-5-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-13	LC-MS: (ES, <i>m/z</i>): 270.
B-18	Ethyl 5-(1-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-14	LC-MS: (ES, <i>m/z</i>): 270.
B-19	Methyl 5-(1-ethyl-1 <i>H</i> -1,2,3-benzotriazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-15	LC-MS: (ES, <i>m/z</i>): 271.
B-20	Ethyl 5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-16	LC-MS: (ES, <i>m/z</i>): 286.
B-23	Methyl 5-[3-(2,2-dimethylpropoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-17	LC-MS: (ES, <i>m/z</i>): 289.1.
B-26	Ethyl 5-(3-ethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-18	LC-MS: (ES, <i>m/z</i>): 260.
B-27	Methyl 5-[3-(benzyloxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-19	LC-MS: (ES, <i>m/z</i>): 309.1.
B-28	Methyl 5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-20	
B-29	Methyl 5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-21	LC-MS: (ES, <i>m/z</i>): 263.
B-30	Methyl 5-(3,5-diethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-22	LC-MS: (ES, <i>m/z</i>): 290.95.
B-31	Methyl 5-(3-ethoxy-5-methoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-23	LC-MS: (ES, <i>m/z</i>): 277.
B-32	Ethyl 5-[2-(propan-2-yloxy)-1,3-oxazol-5-yl]-1 <i>H</i> -pyrazole-3-carboxylate	D-24	LC-MS: (ES, <i>m/z</i>): 266.
B-33	Methyl 5-[1,3-dimethyl-1 <i>H</i> -thieno[2,3- <i>c</i>]pyrazol-5-yl]-1 <i>H</i> -pyrazole-3-carboxylate	D-25	LC-MS: (ES, <i>m/z</i>): 277.
B-38	Methyl 5-[3-[(propan-2-yl)carbamoyl]phenyl]-1 <i>H</i> -pyrazol-3-carboxylate	D-26	

Intermediate D-27: Methyl 1-[(2-chlorophenyl)methyl]-5-(3-(propan-2-yl)phenyl)-1H-pyrazole-3-carboxylate



[0382] A solution of Int. **B-13** (1.0 g, 4.03 mmol, 1.00 equiv) and Int. **C-13** (1.38 g, 6.01 mmol, 1.50 equiv) in AcOH (20 mL) was stirred for 2 h at 100 °C, then concentrated under vacuum, diluted with 200 mL of EtOAc, washed with 2x100 mL of sat NaHCO₃, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The crude product was purified with silica gel chromatography using EtOAc / petroleum ether (1:5) to afford 1.05 g (71%) of the title product as a yellow solid.

Intermediate D-28: Methyl 1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazole-3-carboxylate



[0383] To a solution of Int. **B-2** (3.8 g, 16.09 mmol, 1.00 equiv) in MeOH (80 mL) was added Int. **C-13** (4.8 g, 20.91 mmol, 1.30 equiv). The resulting solution was stirred for 3 h at 65 °C, then concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:2), to afford the title compound as a white solid (5.5 g, 96%).

[0384] The following substituted 3-pyrazolecarboxylic esters were obtained from condensation of methyl 4-(aryl)-2,4-dioxobutanoate with various substituted hydrazines:




Table 7: Pyrazole synthesis with substituted hydrazines.

Ar ₁ NHNH ₂	Diketone	Product	Spectral
-----------------------------------	----------	---------	----------

Phenyl-hydrazine	B-28	Methyl 5-[3-(2-methylpropoxy)-phenyl]-1-phenyl-1 <i>H</i> -pyrazole-3-carboxylate	D-29	LC-MS: (ES, <i>m/z</i>): 357.
(2-Bromophenyl)-hydrazine	B-14	Methyl 1-(2-bromophenyl)-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-30	LC-MS: (ES, <i>m/z</i>): 425.1.
(2-Bromophenyl)-hydrazine	B-18	Ethyl 1-(2-bromophenyl)-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-31	LC-MS: (ES, <i>m/z</i>): 425.
(2-Bromophenyl)-hydrazine	B-23	Methyl 1-(2-bromophenyl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-32	LC-MS: (ES, <i>m/z</i>): 443.1.
(2-Bromophenyl)-hydrazine	B-29	Methyl 1-(2-bromophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-33	LC-MS: (ES, <i>m/z</i>): 417.
(2-Bromophenyl)-hydrazine	B-41	Methyl 1-(2-bromophenyl)-5-(1-propyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-34	LC-MS: (ES, <i>m/z</i>): 439.1.
(2-Bromophenyl)-hydrazine	B-42	Methyl 1-(2-bromophenyl)-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-35	LC-MS: (ES, <i>m/z</i>): 441.2.
(2-Fluorophenyl)-hydrazine	B-3	Methyl 1-(2-fluorophenyl)-5-[3-(oxetan-3-ylmethoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-36	LC-MS: (ES, <i>m/z</i>): 383.1.
(2-Fluorophenyl)-hydrazine	B-20	Ethyl 5-(3-cyclobutoxyphenyl)-1-(2-fluorophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-37	LC-MS: (ES, <i>m/z</i>): 381.
(2-Fluorophenyl)-hydrazine	B-21	Ethyl 1-(2-fluorophenyl)-5-[3-(oxetan-3-ylloxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-38	LC-MS: (ES, <i>m/z</i>): 383.
(2-Fluorophenyl)-hydrazine	B-28	Methyl 1-(2-fluorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-39	LC-MS: (ES, <i>m/z</i>): 369.
(2-Methylphenyl)-hydrazine	B-20	Ethyl 5-(3-cyclobutoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-40	LC-MS: (ES, <i>m/z</i>): 376.
(2-Methylphenyl)-hydrazine	B-30	Methyl 5-(3,5-diethoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-41	LC-MS: (ES, <i>m/z</i>): 381.05.
(2-Nitrophenyl)-hydrazine	B-14	Methyl 5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1-(2-nitrophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-42	LC-MS: (ES, <i>m/z</i>): 392.30.
Quinolin-8-yl-hydrazine	B-23	Methyl 5-[3-(2,2-dimethylpropoxy)-phenyl]-1-(quinolin-8-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-43	LC-MS: (ES, <i>m/z</i>): [M+1] = 416.

C-2	B-29	Methyl 5-(3,5-dimethoxyphenyl)-1-(2-ethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-44	LC-MS: (ES, <i>m/z</i>): 383.
C-3	B-15	Ethyl 5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-45	LC-MS: (ES, <i>m/z</i>): 414.
C-3	B-20	Ethyl 5-(3-cyclobutoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-46	LC-MS: (ES, <i>m/z</i>): 416.
C-3	B-23	Methyl 5-[3-(2,2-dimethylpropoxy)-phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-47	LC-MS: (ES, <i>m/z</i>): 419.10.
C-3	B-24	Methyl 5-[3-(cyclobutylmethoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-48	LC-MS: (ES, <i>m/z</i>): 417.
C-3	B-25	Methyl 5-[3-(cyclopropylmethoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-49	LC-MS: (ES, <i>m/z</i>): 402.
C-3	B-28	Methyl 1-(1-methyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-50	LC-MS: (ES, <i>m/z</i>): 405.
C-3	B-29	Methyl 5-(3,5-dimethoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-51	LC-MS: (ES, <i>m/z</i>): 393.
C-3	B-30	Methyl 5-(3,5-diethoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-52	LC-MS: (ES, <i>m/z</i>): 421.05.
C-5	B-23	Methyl 5-[3-(2,2-dimethylpropoxy)-phenyl]-1-(1-ethyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-53	LC-MS: (ES, <i>m/z</i>): 432.
C-6	B-2	Methyl 5-(3-methoxyphenyl)-1-[2-(propan-2-yloxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-54	LC-MS: (ES, <i>m/z</i>): 367.
C-11	B-23	Methyl 1-(1,3-dimethyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-55	LC-MS: (ES, <i>m/z</i>): 433.3.
C-12	B-23	Methyl 5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -1,2,3-benzotriazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-56	LC-MS: (ES, <i>m/z</i>): 420.
C-13	B-37	Methyl 1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -1,3-benzodiazol-6-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-57	LC-MS: (ES, <i>m/z</i>): [M+1] = 381.

Ar ₁ NHNH ₂	Diketone	Product	
(4-Nitrophenyl)-hydrazine	B-28	Methyl-5-[3-(2-methylpropoxy)phenyl]-1-[(4-nitrophenyl)methyl]-1H-pyrazole-3-carboxylate	D-58
(Pyridin-3-yl)-hydrazine	B-2	Methyl 5-(3-methoxyphenyl)-1-(pyridin-3-yl)-1H-pyrazole-3-carboxylate	D-59
C-13	B-1	Methyl 1-[(2-chlorophenyl)methyl]-5-(2-methoxy-phenyl)-1H-pyrazole-3-carboxylate	D-60
C-13	B-10	Methyl 5-(4-bromothien-2-yl)-1-[(2-chlorophenyl)-methyl]-1H-pyrazole-3-carboxylate	D-61
C-13	B-35	Methyl 1-[(2-chlorophenyl)methyl]-5-[5-(2-methylpropoxy)thien-2-yl]1H-pyrazole-3-carboxylate	D-62
C-13	B-36	Methyl 1-[(2-chlorophenyl)-5-[2-(2-methylpropyl)-1,3-oxazol-5-yl]-1H-pyrazole-3-carboxylate	D-63
C-13	B-39	Methyl 1-[(2-chlorophenyl)methyl]-5-[2-(2-methyl-propoxy)-1,3-thiazol-5-yl]-1H-pyrazole-3-carboxylate	D-64

[0385] The following substituted 3-pyrazolecarboxylic esters were obtained from condensation of various methyl 4-aryl-2,4-dioxobutanoates with (2-chlorophenyl) hydrazine:




Table 8: Pyrazole synthesis with 2-chlorophenyl hydrazine.

Diketone	Product		Spectral
B-2	Methyl 5-(3-methoxyphenyl)-1-(2-methylphenyl)-1H-pyrazole-3-carboxylate (This product from reaction with 2-methyl hydrazine)	D-65	LC-MS: (ES, <i>m/z</i>): 322.
B-3	Methyl 1-(2-chlorophenyl)-5-[3-(oxetan-3-yl-methoxy)phenyl]-1H-pyrazole-3-carboxylate	D-66	LC-MS: (ES, <i>m/z</i>): 399.0
B-5	Methyl 1-(2-chlorophenyl)-5-(pyridin-2-yl)-1H-pyrazole-3-carboxylate	D-67	LC-MS: (ES, <i>m/z</i>): 314.20.
B-12	Ethyl 1-(2-chlorophenyl)-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazole-3-carboxylate	D-68	LC-MS: (ES, <i>m/z</i>): 381.
B-14	Methyl 1-(2-chlorophenyl)-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazole-3-carboxylate	D-69	LC-MS: (ES, <i>m/z</i>): 381.

B-20	Ethyl 1-(2-chlorophenyl)-5-(3-cyclobutoxy-phenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-70	LC-MS: (ES, <i>m/z</i>): 396.
B-21	Ethyl 1-(2-chlorophenyl)-5-[3-(oxetan-3-yloxy)-phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-71	LC-MS: (ES, <i>m/z</i>): 398.95.
B-22	Methyl 1-(2-chlorophenyl)-5-(3-cyclopropoxy-phenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-72	LC-MS: (ES, <i>m/z</i>): 369.25.
B-23	Methyl 1-(2-chlorophenyl)-5-[3-(2,2-dimethyl-propoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-73	LC-MS: (ES, <i>m/z</i>): 398.95.
B-24	Methyl 1-(2-chlorophenyl)-5-[3-(cyclobutyl-methoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-74	LC-MS: (ES, <i>m/z</i>): 396.95.
B-28	Methyl 1-(2-chlorophenyl)-5-[3-(2-methyl-propoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-75	
B-29	Methyl 1-(2-chlorophenyl)-5-(3,5-dimethoxy-phenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-76	LC-MS: (ES, <i>m/z</i>): 373.
B-30	Methyl 1-(2-chlorophenyl)-5-(3,5-diethoxy-phenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-77	LC-MS: (ES, <i>m/z</i>): 401.3.
B-40	Ethyl [5-benzyl-1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-78	

[0386] The following substituted 3-pyrazolecarboxylic esters were obtained from condensation of Int. **B-22** with various substituted hydrazines:






Table 9: Pyrazole synthesis with methyl 4-(3-cyclopropoxyphenyl)-2,4-dioxobutanoate (Int. B-22).

ArNHNH₂, solvent	Product		Spectral
(2-Fluoro-phenyl)hydrazine hydrochloride, EtOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(2-fluorophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-79	LC-MS: (ES, <i>m/z</i>): 353.0.
(2-Bromophenyl)-hydrazine, MeOH	Methyl 1-(2-bromophenyl)-5-(3-cyclopropoxy-phenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-80	LC-MS: (ES, <i>m/z</i>): 413.
(2-Bromo-4-fluoro-phenyl)hydrazine, MeOH	Methyl 1-(2-bromo-4-fluorophenyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-81	LC-MS: (ES, <i>m/z</i>): 430.90.

(2,4-Dichloro-phenyl)hydrazine, MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(2,4-dichlorophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-82	LC-MS: (ES, <i>m/z</i>): 402.95.
(2,5-Dichloro-phenyl)hydrazine, MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(2,5-dichlorophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-83	LC-MS: (ES, <i>m/z</i>): 402.90.
[2-(Difluoromethoxy)phenyl]-hydrazine, CF ₃ COOH / MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-[2-(difluoromethoxy)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-84	LC-MS (ES, <i>m/z</i>): 401
(2-Ethoxyphenyl)-hydrazine, MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(2-ethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-85	LC-MS: (ES, <i>m/z</i>): 378.
(2-Methoxy-phenyl)hydrazine	Methyl 5-(3-cyclopropoxyphenyl)-1-(2-methoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-86	LC-MS: (ES, <i>m/z</i>): 364.9.
(2-Methylphenyl)-hydrazine dihydrochloride, MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-87	LC-MS (ES, <i>m/z</i>): 349.
(2-Nitrophenyl)-hydrazine, EtOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(2-nitrophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-88	LC-MS: (ES, <i>m/z</i>): 380.
(2-(Propan-2-yl)-phenyl)hydrazine HCl, MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-[2-(propan-2-yl)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-89	
[2-(Trifluoromethyl)phenyl]-hydrazine (HCl) ₂ , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-[2-(trifluoromethyl)phenyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-90	LC-MS: (ES, <i>m/z</i>): 403.
C-3 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-91	LC-MS (ES, <i>m/z</i>): 389
C-4 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-92	LC-MS: (ES, <i>m/z</i>): 388.
C-5 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(1-ethyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-93	LC-MS: (ES, <i>m/z</i>): 402.
C-7 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-94	LC-MS: (ES, <i>m/z</i>): 375.1.
C-8 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-95	LC-MS: (ES, <i>m/z</i>): 375.
C-9 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(3-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-96	LC-MS: (ES, <i>m/z</i>): 388.95.
C-10 , MeOH	Methyl 5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -1,3-benzodiazol-7-yl)-1 <i>H</i> -pyrazole-3-carboxylate	D-97	LC-MS: (ES, <i>m/z</i>): 389.0.

Intermediate E-1: 1-(Bromomethyl)-2-ethoxybenzene

[0387] To a solution of (2-ethoxyphenyl)methanol (15.2 g, 99.87 mmol, 1.00 equiv) and CBr_4 (50.1 g, 150.00 mmol, 1.50 equiv) in CH_2Cl_2 (200 mL), was added PPh_3 (39.45 g, 150.41 mmol, 1.50 equiv), in portions at 0 °C. The resulting solution was stirred for 16 h at rt, concentrated under vacuum, and purified with silica gel chromatography using with EtOAc /petroleum ether (1:10) to afford 14 g (65%) of the title compound as a colorless oil. ^1H -NMR: (CDCl_3 , ppm): δ : 7.51 – 7.25 (m, 1H), 7.06 – 6.86 (m, 1H), 4.65 (s, 1H), 4.16 (q, $J = 7.0$ Hz, 1H), 1.53 (t, $J = 7.0$ Hz, 1H).


Intermediate E-2: 1-(Bromomethyl)-2-cyclopropoxybenzene

[0388] **Methyl 2-cyclopropoxybenzoate** To a mixture of methyl 2-hydroxybenzoate (5.0 g, 32.86 mmol, 1.00 equiv) Cs_2CO_3 (32 g, 98.21 mmol, 3.00 equiv) in DMA (100 mL) was added dropwise bromocyclopropane (39.8 g, 328.99 mmol, 10.00 equiv) with stirring at rt. The resulting solution was stirred for 16 h at 130 °C, cooled, diluted with 200 mL of EtOAc, washed with 2x200 mL of water and 2x200 mL of brine, dried over Na_2SO_4 , and concentrated under vacuum. The residue was purified by silica gel chromatography using EtOAc /petroleum ether (1:6) to afford 1.6 g (25%) of the title compound as a yellow oil. LC-MS: (ES, m/z): 193. ^1H NMR (400 MHz, CDCl_3) δ 7.80 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.49 (ddd, $J = 8.9, 7.3, 1.8$ Hz, 1H), 7.38 (dd, $J = 8.4, 1.1$ Hz, 1H), 7.01 (td, $J = 7.5, 7.5, 1.2$ Hz, 1H), 3.89 (s, 3H), 3.83 (tt, $J = 5.9, 5.9, 3.2, 3.2$ Hz, 1H), 0.86 (m, 4H).

[0389] **(2-Cyclopropoxyphenyl)methanol** To a solution of methyl 2-cyclopropoxybenzoate (1.5 g, 7.80 mmol, 1.00 equiv) in THF (100 mL) was added LiAlH_4 (594 mg, 15.65 mmol, 2.00 equiv), in portions at 0 °C. The resulting mixture was stirred for 2 h at 0 °C in an ice/salt bath, then quenched by the addition of $\text{Na}_2\text{SO}_4 / 10 \text{ H}_2\text{O}$. The solids were removed by filtration. The resulting solution was concentrated under vacuum and purified with silica gel chromatography using EtOAc /petroleum ether (1:5) to afford 750 mg (59%) of the title compound as a yellow oil. LC-MS: (ES, m/z): 187. ^1H NMR: (400 MHz, CDCl_3) δ 7.30 (m, 3H), 6.98 (tdd, $J = 7.5, 7.5, 3.7, 1.5$ Hz, 1H), 4.65 (s, 2H), 3.80 (m, 1H), 0.82 (m, 4H).

[0390] **1-(Bromomethyl)-2-cyclopropoxybenzene** To a solution of (2-cyclopropoxy-phenyl)methanol (750 mg, 4.57 mmol, 1.00 equiv) and CBr₄ (3.06 g, 9.16 mmol, 2.00 equiv) in CH₂Cl₂ (30 mL) was added PPh₃ (2.40 g, 9.15 mmol, 2.00 equiv) in portions at 0 °C. The resulting solution was stirred for 2 h at room temperature, concentrated under vacuum, and purified with silica gel chromatography using with EtOAc /petroleum ether (1:5) to afford 1.0 g (96%) of the title compound as colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.30 (m, 3H), 6.96 (dt, J = 7.6, 1.8, 1.8 Hz, 1H), 4.52 (s, 2H), 3.84 (m, 1H), 0.83 (m, 4H).

Intermediate E-3 2-(bromomethyl)-1-ethoxy-3-fluorobenzene



[0391] **Methyl 2-ethoxy-6-fluorobenzoate** A mixture of methyl 2-fluoro-6-hydroxybenzoate (500 mg, 2.94 mmol, 1.00 equiv), K₂CO₃ (811 mg, 5.87 mmol, 2.00 equiv), EtI (911 mg, 5.84 mmol, 2.00 equiv) in DMF (15 mL) was stirred overnight at 80 °C, cooled to rt, and extracted with 3x20 mL of EtOAc. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum to afford 500 mg (86%) of the title compound as light yellow oil.

[0392] LC-MS: (ES, m/z): 199.3. ¹H NMR (300 MHz, CDCl₃) δ 7.29 (m, 1H), 6.70 (m, 2H), 4.08 (q, J = 7.0, 7.0, 7.0 Hz, 2H), 3.92 (s, 3H), 1.40 (t, J = 7.0, 7.0 Hz, 3H).

[0393] **2-Ethoxy-6-fluorobenzoic acid** A solution of the product from the previous step (2.6 g, 13.12 mmol, 1.00 equiv) and NaOH (2 g, 50.00 mmol, 4.00 equiv) in MeOH / H₂O (20/10 mL) was stirred overnight at 40 °C. The pH value of the solution was adjusted to 5 with 1 M HCl. The resulting solution was extracted with 3x50 mL of EtOAc. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum to afford 1.9 g (79%) of the title compound as a colorless oil.

[0394] LC-MS: (ES, m/z): 185.3.

[0395] **(2-Ethoxy-6-fluorophenyl)methanol** To a solution of the product from the previous step (1.9 g, 10.32 mmol, 1.00 equiv) and isobutyl chloroformate (2.1 g, 15.38 mmol, 1.50 equiv) in THF (50 mL) at 0 °C was added TEA (4.5 mL, 3.00 equiv). The reaction was stirred for 2 h at 0 °C, then filtered. To the filtrate was added NaBH₄ (780 mg, 20.62 mmol, 2.00 equiv) at 0 °C. The resulting solution was stirred for 3 h at room temperature. The reaction was then quenched by the addition of 10 mL of water/ice. The resulting solution was extracted with 3x50 mL of EtOAc. The combined organic layers were dried over Na₂SO₄,


concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (2/1) to afford 0.7 g (40%) of the title compound as a light yellow oil.

[0396] ^1H NMR (300 MHz, DMSO) δ 7.27 (td, J = 8.3, 8.3, 7.0 Hz, 1H), 6.77 (m, 2H), 4.75 (t, J = 5.5, 5.5 Hz, 1H), 4.48 (dd, J = 5.5, 1.9 Hz, 2H), 4.06 (q, J = 7.0, 7.0, 7.0 Hz, 2H), 1.35 (t, J = 6.9, 6.9 Hz, 3H).

[0397] **2-(Bromomethyl)-1-ethoxy-3-fluorobenzene** A solution of the product from the previous step (700 mg, 4.11 mmol, 1.00 equiv), PPh_3 (2.16 g, 8.24 mmol, 2.00 equiv), and CBr_4 (2.7 g, 2.00 equiv) in CH_2Cl_2 (30 mL) was stirred for 2 h at rt. The resulting solution was extracted with 3x30 mL of EtOAc. The combined organic layers were dried over Na_2SO_4 , concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (5/1) to afford 0.4 g (42%) of the title compound as a light yellow oil.

[0398] ^1H NMR (300 MHz, CDCl_3) δ 7.24 (m, 1H), 6.69 (m, 2H), 4.67 (dd, J = 32.8, 1.5 Hz, 2H), 4.13 (qd, J = 7.0, 7.0, 7.0, 2.5 Hz, 2H), 1.48 (td, J = 7.0, 6.9, 3.2 Hz, 3H).

Intermediate E-4 1-(Bromomethyl)-2-(propan-2-yloxy)benzene



[0399] **[2-(Propan-2-yloxy)phenyl]methanol** To a solution of 2-(hydroxymethyl)phenol (2 g, 16.11 mmol, 1.00 equiv) in DMF (10 mL) was added sequentially 2-iodopropane (5.48 g, 32.24 mmol, 2.00 equiv) then Cs_2CO_3 (10.5 g, 32.23 mmol, 2.00 equiv). The resulting solution was stirred overnight at 85 °C. The reaction mixture was cooled to room temperature. The resulting solution was extracted with 3x50 mL of EtOAc, and the organic layers were combined, dried over anhydrous Na_2SO_4 , and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / hexane (1/4). This resulted in the title compound as a light yellow oil (2.1 g, 78%).


[0400] **1-(Bromomethyl)-2-(propan-2-yloxy)benzene** To a solution of the product from the previous step (3 g, 18.05 mmol, 1.00 equiv) in CH_2Cl_2 (50 mL) were added sequentially CBr_4 (12.84 g, 36.17 mmol, 2.00 equiv) and PPh_3 (9.48 g, 36.14 mmol, 2.00 equiv) in an ice bath. The resulting solution was stirred for 5 h at room temperature. The resulting solution was extracted with 3x100 mL of CH_2Cl_2 , and the organic layers were combined, dried over anhydrous Na_2SO_4 , and concentrated under vacuum. The residue was applied onto a silica gel

column with EtOAc / hexane (1/4). This resulted in the title compound as a light yellow oil (1.6 g, 39%).

[0401] The following alkylating agents were reacted with pyrazoles as disclosed below:


Intermediate E-5	1-(Chloromethyl)-2-nitrobenzene
Intermediate E-6	1-Bromo-2-(bromomethyl)-benzene
Intermediate E-7	1-(Bromomethyl)-2-chlorobenzene
Intermediate E-8	2-(Bromomethyl)-1-fluoro-3-nitrobenzene
Intermediate E-9	(2-Bromoethyl)benzene
Intermediate E-10	1-(Chloromethyl)-2-chlorobenzene
Intermediate E-11	2-(Bromomethyl)pyridine
Intermediate E-12	7-(Bromomethyl)-1-methyl-1H-indazole
Intermediate E-13	1-bromo-2-methylpropane
Intermediate E-14	4-(bromomethyl)tetrahydropyran

Intermediate D-98: Methyl 1-[(3-chlorophenyl)methyl]-5-phenyl-1H-pyrazole-3-carboxylate




[0402] To a suspension of methyl 5-phenyl-1H-pyrazole-3-carboxylate (1) (1.0 g, 4.94 mmol) in toluene was added NaH (60%) (0.394 g, 9.88 mmol) portionwise under N₂ at room temperature, and stirring was continued for 30 min. To the above mixture a solution of 3-chlorobenzyl bromide (0.96 ml, 7.42 mmol) in toluene (3 mL) was added dropwise at 60 °C. The reaction mixture was stirred at 110 °C for 16 h. The mixture was cooled to room temperature and quenched with aq. NH₄Cl solution. The mixture was partitioned with EtOAc (100 mL) and the organic layer was separated. The EtOAc layer was washed with brine (2x25 mL) and dried over Na₂SO₄, and the solvent was evaporated. The residue was chromatographed over SiO₂ (ISCO CombiFlash® Rf 200) using 0-50% gradient of EtOAc in hexane to afford title compound (1.2 g, 75%). ¹H NMR (CDCl₃, 400 MHz) δ 3.95 (s, 3H), 5.36 (s, 2H), 6.85-6.92 (m, 2H), 6.95-6.99 (m, 1H), 7.17-7.29 (m, 4H), 7.35-7.48 (m, 3H).

Intermediate D-99: Methyl 1-[(2-cyclopropoxyphenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazole-3-carboxylate



[0403] To a solution of Int. **D-3** (793 mg, 3.41 mmol, 1.00 equiv) in toluene (10 mL) was added NaH (164 mg, 6.83 mmol, 2.00 equiv). The resulting solution was stirred at rt for 30 min, then warmed to 60 °C. To this was added a solution of Int. **E-2** (1 g, 4.40 mmol, 1.30 equiv) in toluene (10 mL). The resulting solution was stirred at 60 °C for 1 h, then the solution was heated to 110 °C and stirred for 6 h at this temperature. The reaction mixture was then cooled to rt, quenched by the addition of 10 mL of water/ice, and extracted with 3x100 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, and concentrated under vacuum, and purified with silica gel column using EtOAc / hexane (1/3) to afford 1 g (77%) of the title compound as a yellow oil. LC-MS: (ES, *m/z*): 379.

Intermediate D-100: Methyl 1-[(2-chlorophenyl)methyl]-5-(3,5-dimethoxyphenyl)-1H-pyrazole-3-carboxylate



[0404] To a solution of methyl Int. D-21 (1.0 g, 3.81 mmol, 1.00 equiv) in toluene (50 mL) was added NaH (304 mg, 7.60 mmol, 2.00 equiv) in portions at rt. The mixture was stirred at rt for 30 min, then Int. **E-7** (1.56 g, 7.59 mmol, 2.00 equiv) was added dropwise with stirring at 60 °C. The resulting solution was stirred for 1 h at 60 °C, then warmed to 110 °C and stirred for an additional 3 h. The reaction was then quenched by the addition of water/ice. The resulting solution was extracted with 200 mL of EtOAc, and the combined organic layers were washed with 2x100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:3) to afford 800 mg (54%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 387. ¹H-NMR: (CDCl₃, *ppm*): δ: 7.42 – 7.29 (m, 1H), 7.29 – 7.13 (m, 2H), 6.97 (s, 1H), 6.77 – 6.67

(m, 1H), 6.42 (dd, J = 30.5, 2.3 Hz, 3H), 5.55 (s, 2H), 3.97 (d, J = 1.0 Hz, 3H), 3.69 – 3.60 (m, 6H).

[0405] The following substituted pyrazoles were obtained from alkylation with an alkyl halide:


Table 10: Pyrazole alkylation.

Pyrazole	Alkyl halide	Product	Spectral
D-2	E-1	Methyl 5-(3-cyclopropoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazole-3-carboxylate	D-101 LC-MS: (ES, m/z): 393.
D-2	E-2	Methyl 5-(3-cyclopropoxyphenyl)-1-[(2-nitrophenyl)methyl]-1H-pyrazole-3-carboxylate	D-102 LC-MS: (ES, m/z): 394.
D-2	E-6	Methyl 1-[(2-bromophenyl)methyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazole-3-carboxylate	D-103 LC-MS: (ES, m/z): 429.2.
D-5	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-(2,3-dihydro-1,4-benzodioxin-6-yl)-1H-pyrazole-3-carboxylate	D-104 LC-MS: (ES, m/z): 385.
D-6	E-7	Methyl 5-(2H-1,3-benzodioxol-5-yl)-1-[(2-chlorophenyl)methyl]-1H-pyrazole-3-carboxylate	D-105 LC-MS: (ES, m/z): 371.
D-7	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-pyrazol-4-yl)-1H-pyrazole-3-carboxylate	D-106 LC-MS: (ES, m/z): 331.
D-8	E-7	Ethyl 5-(1-benzothiophen-2-yl)-1-[(2-chlorophenyl)methyl]-1H-pyrazole-3-carboxylate	D-107 LC-MS: (ES, m/z): 397.
D-9	E-7	Ethyl 1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-indol-6-yl)-1H-pyrazole-3-carboxylate	D-108 LC-MS: (ES, m/z): 393.
D-10	E-1	Ethyl 1-[(2-ethoxyphenyl)methyl]-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazole-3-carboxylate	D-109 LC-MS: (ES, m/z): 404.
D-10	E-7	Ethyl 1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazole-3-carboxylate	D-110 LC-MS: (ES, m/z): 394.
D-11	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazole-3-carboxylate	D-111 LC-MS: (ES, m/z): 395.
D-11	E-11	Methyl 5-(1-ethyl-1H-indazol-6-yl)-1-([pyridin-2-yl]methyl)-1H-pyrazole-3-carboxylate	D-112 LC-MS: (ES, m/z): 362.3.
D-12	E-1	Methyl 5-(3-chloro-5-methoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazole-3-carboxylate	D-113 LC-MS: (ES, m/z): 401.
D-13	E-7	Ethyl 1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-indazol-5-yl)-1H-pyrazole-3-carboxylate	D-114 LC-MS: (ES, m/z): 394.
D-11	E-1	Methyl 1-[(2-ethoxyphenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazole-3-carboxylate	D-115 LC-MS: (ES, m/z): 404.
D-14	E-7	Ethyl 1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-indazol-4-yl)-1H-pyrazole-3-carboxylate	D-116 LC-MS: (ES, m/z): 381.
D-15	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1H-1,2,3-benzotriazol-6-yl)-1H-pyrazole-3-carboxylate	D-117 LC-MS: (ES, m/z): 396.

D-16	E-1	Ethyl 5-(3-cyclobutoxyphenyl)-1-[(2-ethoxy-phenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-118	LC-MS: (ES, <i>m/z</i>): 421.
D-16	E-6	Ethyl 1-[(2-bromophenyl)methyl]-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-119	LC-MS: (ES, <i>m/z</i>): 454.
D-17	E-12	Methyl 5-[3-(2,2-dimethylpropoxy)phenyl]-1-[(1-methyl-1 <i>H</i> -indazol-7-yl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-120	LC-MS: (ES, <i>m/z</i>): 433.3.
D-18	E-1	Methyl 5-(3-ethoxyphenyl)-1-[(2-ethoxy-phenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-121	LC-MS: (ES, <i>m/z</i>): 394.
D-19	E-7	Methyl 5-[3-(benzyloxy)phenyl]-1-[(2-chloro-phenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-122	LC-MS: (ES, <i>m/z</i>): 433.05.
D-20	E-11	Methyl 5-[3-(2-methylpropoxy)phenyl]-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-123	LC-MS: (ES, <i>m/z</i>): 366.
D-21	E-1	Methyl 5-(3,5-dimethoxyphenyl)-1-[(2-ethoxy-phenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-124	LC-MS: (ES, <i>m/z</i>): 397.
D-21	E-9	5-(3,5-Dimethoxyphenyl)-1-(2-phenylethyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-125	LC-MS: (ES, <i>m/z</i>): 366.
D-21	E-3	Methyl 5-(3,5-dimethoxyphenyl)-1-[(2-ethoxy-6-fluorophenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-126	LC-MS: (ES, <i>m/z</i>): 415.10.
D-21	E-11	Methyl 5-(3,5-dimethoxyphenyl)-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-127	LC-MS: (ES, <i>m/z</i>): 354.
D-22	E-1	Methyl 5-(3,5-diethoxyphenyl)-1-[(2-ethoxy-phenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-128	LC-MS: (ES, <i>m/z</i>): 425.1.
D-22	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-(3,5-diethoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-129	LC-MS: (ES, <i>m/z</i>): 415.15.
D-23	E-1	Methyl 5-(3-ethoxy-5-methoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-130	LC-MS: (ES, <i>m/z</i>): 411.
D-24	E-10	Ethyl 1-[(2-chlorophenyl)methyl]-5-[2-(propan-2-yloxy)-1,3-oxazol-5-yl]-1 <i>H</i> -pyrazole-3-carboxylate	D-131	LC-MS: (ES, <i>m/z</i>): 389.
D-25	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-[1,3-dimethyl-1 <i>H</i> -thieno[2,3- <i>c</i>]pyrazol-5-yl]-1 <i>H</i> -pyrazole-3-carboxylate	D-132	LC-MS: (ES, <i>m/z</i>): 401.


Pyrazole	Alkyl Halide	Product	
D-3	E-8	Methyl 1-[(2-fluoro-6-nitrophenyl)methyl]-5-(3-methoxyphenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-133
D-3	E-4	Methyl 5-(3-methoxyphenyl)-1-[[2-(propan-2-yloxy)-phenyl]methyl]-1 <i>H</i> -pyrazole-3-carboxylate	D-134
D-3	E-13	Methyl 5-(3-methoxyphenyl)-1-(2-methylpropyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-135
D-3	E-14	Methyl 5-(3-methoxyphenyl)-1-([oxan-4-yl]methyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-136
D-4	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-(3-nitrophenyl)-1 <i>H</i> -pyrazole-3-carboxylate	D-137
D-26	E-7	Methyl 1-[(2-chlorophenyl)methyl]-5-[3-[(propan-2-yl)-carbamoyl]phenyl]-1 <i>H</i> -pyrazol-3-carboxylate	D-138

Intermediate D-139: Methyl 1-[(2,6-dimethoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazole-3-carboxylate



[0406] To a solution of Int. **D-20** (1.37 g, 4.99 mmol, 1.00 equiv) in THF (50 mL) under N₂ at 0 °C was added (2,6-dimethoxyphenyl)methanol (1.26 g, 7.49 mmol, 1.50 equiv) followed by PPh₃ (1.81 g, 6.90 mmol, 1.30 equiv). Subsequently DIAD (1.75 g, 8.65 mmol, 1.50 equiv) was added dropwise and the resulting solution was stirred for 16 h at 25 °C. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:19). This resulted in the title compound as a yellow solid (890 mg, 42%).

Intermediate D-140: Methyl 1-[(3-chloropyridin-2-yl)methyl]-5-(3-methoxyphenyl)-1H-pyrazole-3-carboxylate




[0407] **(3-Chloropyridin-2-yl)methanol** To a solution of 3-chloropyridine-2-carboxylic acid (2.1 g, 13.33 mmol, 1.00 equiv) in THF (40 mL) was added Et₃N (2.7 g, 26.68 mmol, 2.00 equiv), followed by the addition of chloro(propan-2-yloxy)methanone (2.45 g, 19.99 mmol, 1.50 equiv) dropwise with stirring at 0 °C. The solution was stirred for 1 h at room temperature. The solid was removed by filtration. To the filtrate was added NaBH₄ (1.53 g, 40.44 mmol, 3.00 equiv). The resulting solution was stirred for 2 h at rt, then diluted with 50 mL of H₂O and extracted with 2x100 mL of EtOAc. The combined organic layers were washed with 50 mL of saturated NaCl, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using with EtOAc / petroleum ether (1:6) to afford 0.9 g (47%) of the title compound as a colorless oil.

[0408] **Methyl 1-[(3-chloropyridin-2-yl)methyl]-5-(3-methoxyphenyl)-1H-pyrazole-3-carboxylate** To a solution of the product from the previous step (850 mg, 5.92 mmol, 1.00 equiv) in THF (8.5 mL), under N₂, was added Int. **D-3** (1.4 g, 6.03 mmol, 1.00 equiv) and

Ph₃P (3.1 g, 11.82 mmol, 2.00 equiv), followed by the addition of DIAD (1.8 g, 8.91 mmol, 1.50 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 5 h at rt, diluted with 30 mL of H₂O, and extracted with 2x50 mL of EtOAc. The combined organic layers were washed with 20 mL of saturated NaCl, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1:10) to afford 1.7 g (80%) of the title compound as a white solid. ¹H-NMR (400 MHz, DMSO-*d*₆) δ 8.90 (s, 1H), 8.35 (dd, *J* = 4.7, 1.5 Hz, 1H), 7.99 (dd, *J* = 8.1, 1.5 Hz, 1H), 7.51 (s, 1H), 7.45 (dt, *J* = 7.7, 1.2 Hz, 1H), 7.41 (dd, *J* = 2.6, 1.5 Hz, 1H), 7.36 – 7.30 (m, 1H), 6.91 (ddd, *J* = 8.2, 2.7, 1.0 Hz, 1H), 5.99 (s, 2H), 3.80 (s, 3H), 3.77 (s, 3H).


Intermediate D-141: Methyl 1-[[4-(dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazole-3-carboxylate



[0409] **Methyl 1-[(4-aminophenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazole-3-carboxylate** To a solution of Int. **D-58** (2 g, 4.88 mmol, 1.00 equiv) in AcOH / H₂O (100/40 mL) was added Zn (1 g, 15.29 mmol, 3.00 equiv), and the resulting solution was stirred for 5 h at 25 °C. The solids were collected by filtration. The pH value of the solution was adjusted to 7 with (sat) NaHCO₃ (aq) and the resulting solution was extracted with 200 mL of EtOAc. The combined organic layers were concentrated under vacuum to afford 1.5 g (81%) of the title compound as a yellow oil.


[0410] **Methyl 1-[[4-(dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazole-3-carboxylate** To a solution of the product from the previous step (1.35 g, 3.56 mmol, 1.00 equiv) in MeOH (50 mL) was added formaldehyde (5 mL, 1.50 equiv) in 30 min. To this was added NaCNBH₃ (800 mg, 12.73 mmol, 3.00 equiv), and the resulting solution was stirred for 5 min at 25 °C. The resulting solution was extracted with 200 mL of EtOAc, and the combined organic layers were purified with silica gel chromatography using EtOAc / petroleum ether (1:7) to afford 810 mg (56%) of the title compound as a yellow liquid.

Intermediate D-142: Methyl 1-[[2-(dimethylamino)-6-fluorophenyl]methyl]-5-(3-methoxyphenyl)-1H-pyrazole-3-carboxylate



[0411] Intermediate **D-142** was obtained from Int. **D-133** by a zinc reduction / reduction amination sequence similar to that used to prepare Int. **D-141**.

Intermediate D-143: Methyl 1-[(2-chlorophenyl)methyl]-5-[3-(oxetan-3-ylmethoxy)phenyl]-1H-pyrazole-3-carboxylate




[0412] **1-[(2-Chlorophenyl)methyl]-5-(3-hydroxyphenyl)-1H-pyrazole-3-carboxylic acid** A solution of Int. **D-122** (3.2 g, 7.39 mmol, 1.00 equiv) in AcOH/HCl (15/5 mL) was stirred for 2 h at 90 °C, then cooled, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (2/1) to afford 2 g (82%) of the title compound as a light yellow oil. LC-MS: (ES, m/z): 329.0. ^1H NMR: (300 MHz, DMSO) δ 12.83 (s, 1H), 9.72 (s, 1H), 7.46 (m, 1H), 7.29 (m, 3H), 6.81 (m, 5H), 5.48 (s, 2H).

[0413] **Methyl 1-[(2-chlorophenyl)methyl]-5-(3-hydroxyphenyl)-1H-pyrazole-3-carboxylate** A solution of the product from the previous step (2 g, 6.08 mmol, 1.00 equiv) and H_2SO_4 (1 mL) in MeOH (20 mL) was stirred for 2 h at 65 °C. The pH was adjusted to 7 with saturated NaHCO_3 . The resulting solution was extracted with 3x30 mL of EtOAc, and the combined organic layers were dried over Na_2SO_4 , concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (2/1) to afford 1.7 g (82%) of the title compound as a white solid. LC-MS: (ES, m/z): 343.1. ^1H NMR: (400 MHz, DMSO) δ 9.74 (s, 1H), 7.46 (m, 1H), 7.31 (m, 3H), 6.93 (s, 1H), 6.83 (m, 4H), 5.50 (s, 2H), 3.82(s, 3H).


[0414] **Methyl 1-[(2-chlorophenyl)methyl]-5-[3-(oxetan-3-ylmethoxy)phenyl]-1H-pyrazole-3-carboxylate** To a solution of the product from the previous step (1.7 g, 4.96 mmol, 1.00 equiv), PPh₃ (2.62 g, 9.99 mmol, 2.00 equiv), and (oxetan-3-yl)methanol (870 mg, 9.87 mmol, 2.00 equiv) in THF (30 mL) at -10 °C under an N₂ atmosphere was added DTAD (1.82 g, 9.01 mmol, 1.80 equiv). The resulting solution was stirred overnight at rt and extracted with 3x30 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc /petroleum ether (1/3) to afford 1.1 g (54%) of the title compound as a white solid. LC-MS: (ES, *m/z*): 413.1.

Intermediate D-144: Methyl 5-(3-cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-indazol-7-yl)-1H-pyrazole-3-carboxylate



[0415] To a solution of Int. **D-95** (600 mg, 1.60 mmol, 1.00 equiv) in CH₂Cl₂ (15 mL) was added NaH (128 mg, 5.33 mmol, 2.00 equiv). The solution was stirred 10 min, then SEMCl (664 mg, 2.50 equiv) was added. The resulting solution was stirred for 16 h at rt, then quenched by the addition of water, and extracted with 2x100 mL of EtOAc. The combined organic layers were washed with 100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel column using EtOAc /petroleum ether (1:2) to afford 300 mg (37%) of the title compound as a yellow oil. LC-MS: (ES, *m/z*): 504.


Intermediate D-145: Methyl 5-(3-cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-indazol-4-yl)-1H-pyrazole-3-carboxylate



[0416] To a solution of Int. **D-94** (460 mg, 1.23 mmol, 1.00 equiv) in CH₂Cl₂ (10 mL) was added NaH (57.5 mg, 2.40 mmol, 2.00 equiv) at 0 °C. The resulting solution was stirred for 30 min, then SEMCl (253 mg, 0.69 mmol, 1.30 equiv) was added. The resulting solution


was stirred for 5 h at rt, then quenched with H₂O and extracted with 3x20 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with silica gel chromatography using EtOAc / petroleum ether (1/3) to afford 300 mg (48%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 505.1.

Intermediate D-146: methyl 5-(3-cyclopropoxyphenyl)-1-(3-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-indazol-4-yl)-1H-pyrazole-3-carboxylate



[0417] To a solution of Int. **D-96** (500 mg, 1.29 mmol, 1.00 equiv) in CH₂Cl₂ (10 mL) was added NaH (62 mg, 2.58 mmol, 2.00 equiv). The resulting solution was stirred for 30 min at 0 °C, then [2-(chloromethoxy)ethyl]trimethylsilane (278 mg, 1.67 mmol, 1.30 equiv) was added. The resulting solution was stirred for 5 h at rt, then quenched with H₂O and extracted with 3x20 mL of CH₂Cl₂. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with prep-TLC using EtOAc /petroleum ether (1/3) to afford 250 mg (36%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 519.40.

Intermediate D-147: Ethyl 1-[(2-chlorophenyl)methyl]-5-phenoxy-1H-pyrazole-3-carboxylate




[0418] **Ethyl 1-[(2-chlorophenyl)methyl]-5-oxo-4,5-dihydro-1H-pyrazole-3-carboxylate** A solution of Int. **C-13** (4 g, 17.54 mmol, 1.00 equiv) and 1,4-diethyl 2-oxobutanedioate (0 mg, 1.30 equiv) in EtOH (120 mL) was heated to reflux overnight, then concentrated under vacuum. The crude product was purified using silica gel chromatography using EtOAc / petroleum ether (1:20~1:5), to afford 2.8 g (57%) of the title product as a yellow syrup.

[0419] **Ethyl 1-[(2-chlorophenyl)methyl]-5-(2-nitrophenoxy)-1H-pyrazole-3-carboxylate** A solution of the product from the previous step (2.8 g, 9.97 mmol, 1.00 equiv), 1-fluoro-2-nitrobenzene (2.8 g, 19.84 mmol, 2.00 equiv), and K₂CO₃ (2.7 g, 19.57 mmol, 2.00 equiv) in DMF (20 mL) was stirred overnight at 90 °C. The resulting solution was diluted with EtOAc, washed with water and brine, dried over anhydrous Na₂SO₄, and concentrated under vacuum. The crude product was purified with silica gel chromatography using EtOAc / petroleum ether (1:50-1:10) to afford 0.8 g (20%) of the title product as a yellow oil.

[0420] **Ethyl 5-(2-aminophenoxy)-1-[(2-chlorophenyl)methyl]-1H-pyrazole-3-carboxylate** A solution of the product from the previous step (800 mg, 1.99 mmol, 1.00 equiv) in AcOH / H₂O(3:1) (8 mL) was stirred over Zn (600 mg, 9.38 mmol, 5.00 equiv) for 3 h at 60 °C. The solids were removed by filtration, and the filtrate was concentrated under vacuum. The crude product was purified by silica gel chromatography using EtOAc / petroleum ether (1:30~1:5) to afford 540 mg (73%) of the title product as a yellow solid.

[0421] **Ethyl 1-[(2-chlorophenyl)methyl]-5-phenoxy-1H-pyrazole-3-carboxylate** To a solution of the product from the previous step (540 mg, 1.45 mmol, 1.00 equiv) in THF (15 mL) was added *tert*-butyl nitrite (450 mg, 4.36 mmol, 3.00 equiv). The resulting solution was stirred for 30 min at 60 °C, then concentrated under vacuum. The residue was purified with Prep-TLC (EtOAc:PE=1:3) to afford 220 mg (42%) of the title product as a solid.

Intermediate F-1: [1-[(3-Chlorophenyl)methyl]-5-phenyl-1H-pyrazol-3-yl]methanol



[0422] A solution of Int. **D-98** (0.80 g, 2.45 mmol) in anhydrous THF was cooled to 0 °C. To the above mixture LiAlH₄ (0.14g, 3.67 mmol) was added portionwise, and stirring was continued at 0 °C for 1.30 h. The mixture was quenched with water (0.15 mL) and 30% aq. NaOH solution (0.3 mL) at 0°C, and stirring was continued for 30 min. The reaction mixture was filtered, the filter cake was washed with THF (2x 10 mL), and the filtrates were combined and evaporated to dryness. The residue was chromatographed over SiO₂ (ISCO CombiFlash® Rf 200) using 0-40% gradient of EtOAc in DCM to afford the title product (0.58 g 79%). ¹H NMR (CDCl₃, 400 MHz) δ 4.64-4.81 (d, 2H), 5.31 (s, 2H), 6.35 (s, 1H),


6.86-6.94 (m, 1H), 6.98-7.05 (m, 1H), 7.19-7.23 (m, 2H), 7.26-7.35 (m, 2H), 7.39-7.46 (m, 3H).

[0423] The following substituted 3-(hydroxymethyl)pyrazoles were obtained from LiAlH_4 reduction of the corresponding methyl esters:

Table 11: LiAlH_4 reduction of pyrazole carboxylic esters.

Ester	Product		Spectral
D-57	[1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-1,3-benzodiazol-6-yl)-1H-pyrazol-3-yl]methanol	F-2	LC-MS: (ES, m/z): [M+1] = 353.
D-59	[5-(3-Methoxyphenyl)-1-(pyridin-3-yl)-1H-pyrazol-3-yl]methanol	F-3	
D-65	[5-(3-Methoxyphenyl)-1-(2-methylphenyl)-1H-pyrazol-3-yl]methanol	F-4	
D-117	[1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1H-1,2,3-benzotriazol-6-yl)-1H-pyrazol-3-yl]methanol	F-5	LC-MS: (ES, m/z): 368.
D-132	[1-[(2-chlorophenyl)methyl]-5-[1,3-dimethyl-1H-thieno[2,3-c]pyrazol-5-yl]-1H-pyrazol-3-yl]methanol	F-6	LC-MS: (ES, m/z): 373.
D-137	[1-[(2-chlorophenyl)methyl]-5-(3-nitrophenyl)-1H-pyrazole-3-yl]methanol	F-7	

Intermediate F-8: [1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methanol



[0424] (iii) To a solution of Int. **D-28** (5.5 g, 15.41 mmol, 1.00 equiv) in THF (100 mL) was added LiBH_4 (20 mL) dropwise with stirring at 0 °C in 5 min. The resulting solution was stirred for 16 h at room temperature. The reaction was then quenched by the addition of 5 mL of water, diluted with 200 mL of H_2O , extracted with 2x200 mL of EtOAc, and the organic layers were combined. The resulting mixture was washed with 1x100 mL of brine then dried over anhydrous Na_2SO_4 and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:8). This resulted in the title compound as a white solid (4.3 g, 85%).

[0425] The following substituted 3-(hydroxymethyl)pyrazoles were obtained from LiBH₄ reduction of the corresponding methyl esters:

Table 12: LiBH₄ reduction of pyrazole carboxylic esters.

Ester	Product		Spectral
D-29	[5-[3-(2-Methylpropoxy)phenyl]-1-phenyl-1 <i>H</i> -pyrazol-3-yl]methanol	F-9	LC-MS: (ES, <i>m/z</i>): 323.
D-30	[1-(2-Bromophenyl)-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-10	LC-MS: (ES, <i>m/z</i>): 398.9.
D-31	[1-(2-Bromophenyl)-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-11	LC-MS: (ES, <i>m/z</i>): 383.
D-32	[1-(2-Bromophenyl)-5-[3-(2,2-dimethylpropoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-12	LC-MS: (ES, <i>m/z</i>): 417.00.
D-33	[1-(2-Bromophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-13	LC-MS: (ES, <i>m/z</i>): 389.
D-34	[1-(2-bromophenyl)-5-(1-propyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-14	LC-MS: (ES, <i>m/z</i>): 413.0.
D-35	[1-(2-bromophenyl)-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-15	LC-MS: (ES, <i>m/z</i>): 399.2.
D-36	[1-(2-Fluorophenyl)-5-[3-(oxetan-3-ylmethoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-16	LC-MS: (ES, <i>m/z</i>): 355.2.
D-37	[5-(3-Cyclobutoxyphenyl)-1-(2-fluorophenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-17	LC-MS: (ES, <i>m/z</i>): 339.
D-38	[1-(2-Fluorophenyl)-5-[3-(oxetan-3-yloxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-18	LC-MS: (ES, <i>m/z</i>): 341.
D-39	[1-(2-Fluorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-19	LC-MS: (ES, <i>m/z</i>): 341.
D-40	[5-(3-cyclobutoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-20	LC-MS: (ES, <i>m/z</i>): 334.
D-41	[5-(3,5-Diethoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-21	LC-MS: (ES, <i>m/z</i>): 353.00.
D-42	[5-(1-Ethyl-1 <i>H</i> -indazol-6-yl)-1-(2-nitrophenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-22	LC-MS: (ES, <i>m/z</i>): 364.3.
D-43	[5-[3-(2,2-dimethylpropoxy)phenyl]-1-(quinolin-8-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-23	LC-MS: (ES, <i>m/z</i>): [M+1] = 388.
D-44	[5-(3,5-Dimethoxyphenyl)-1-(2-ethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-24	LC-MS: (ES, <i>m/z</i>): 355.
D-45	[5-(1-Ethyl-1 <i>H</i> -indazol-6-yl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-25	LC-MS: (ES, <i>m/z</i>): 372.

D-46	[5-(3-Cyclobutoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-26	LC-MS: (ES, <i>m/z</i>): 375.
D-47	[5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-27	LC-MS: (ES, <i>m/z</i>): 391.10.
D-48	[5-[3-(Cyclobutylmethoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-28	LC-MS: (ES, <i>m/z</i>): 389.
D-49	[5-[3-(Cyclopropylmethoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-29	LC-MS: (ES, <i>m/z</i>): 375.
D-50	[1-(1-Methyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-30	LC-MS: (ES, <i>m/z</i>): 377.
D-51	[5-(3,5-dimethoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-31	LC-MS: (ES, <i>m/z</i>): 365.
D-52	[5-(3,5-Diethoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-32	LC-MS: (ES, <i>m/z</i>): 393.05.
D-53	[5-[3-(2,2-Dimethylpropoxy)phenyl]-1-(1-ethyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-33	LC-MS: (ES, <i>m/z</i>): 404.
D-54	[5-(3-Methoxyphenyl)-1-[2-(propan-2-yloxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-34	LC-MS: (ES, <i>m/z</i>): 339.
D-55	[1-(1,3-dimethyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-35	LC-MS: (ES, <i>m/z</i>): 405.35.
D-56	[5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -1,2,3-benzotriazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-36	
D-66	[1-(2-chlorophenyl)-5-[3-(oxetan-3-yl-methoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-37	LC-MS: (ES, <i>m/z</i>): 371.0.
D-67	[1-(2-chlorophenyl)-5-(pyridin-2-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-38	LC-MS: (ES, <i>m/z</i>): 285.85.
D-68	[1-(2-Chlorophenyl)-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-39	LC-MS: (ES, <i>m/z</i>): 339.
D-69	[1-(2-chlorophenyl)-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-40	LC-MS: (ES, <i>m/z</i>): 353.
D-70	[1-(2-Chlorophenyl)-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-41	LC-MS: (ES, <i>m/z</i>): 355.
D-71	[1-(2-Chlorophenyl)-5-[3-(oxetan-3-yloxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-42	LC-MS: (ES, <i>m/z</i>): 357.
D-72	[1-(2-chlorophenyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-43	LC-MS: (ES, <i>m/z</i>): 369.25.
D-73	[1-(2-Chlorophenyl)-5-[3-(2,2-dimethylpropoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-44	LC-MS: (ES, <i>m/z</i>): 371.00.
D-74	[1-(2-Chlorophenyl)-5-[3-(cyclobutylmethoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-45	LC-MS: (ES, <i>m/z</i>): 368.95.

D-76	[1-(2-Chlorophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-46	LC-MS: (ES, <i>m/z</i>): 345.
D-77	[1-(2-Chlorophenyl)-5-(3,5-diethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-47	LC-MS: (ES, <i>m/z</i>): 373.1.
D-79	[5-(3-cyclopropoxyphenyl)-1-(2-fluorophenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-48	LC-MS: (ES, <i>m/z</i>): 325.25.
D-80	[1-(2-Bromophenyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-49	LC-MS: (ES, <i>m/z</i>): 387.
D-81	[1-(2-Bromo-4-fluorophenyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-50	LC-MS: (ES, <i>m/z</i>): 405.0.
D-82	[5-(3-Cyclopropoxyphenyl)-1-(2,4-dichlorophenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-51	LC-MS: (ES, <i>m/z</i>): 375.00.
D-83	[5-(3-Cyclopropoxyphenyl)-1-(2,5-dichlorophenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-52	LC-MS: (ES, <i>m/z</i>): 375.05.
D-84	[5-(3-Cyclopropoxyphenyl)-1-[2-(difluoromethoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-53	LC-MS: (ES, <i>m/z</i>): 373.
D-85	[5-(3-Cyclopropoxyphenyl)-1-(2-ethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-54	LC-MS: (ES, <i>m/z</i>): 350.
D-86	[5-(3-Cyclopropoxyphenyl)-1-(2-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-55	LC-MS: (ES, <i>m/z</i>): 337.
D-87	[5-(3-Cyclopropoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-56	LC-MS: (ES, <i>m/z</i>): 321.
D-88	[5-(3-Cyclopropoxyphenyl)-1-(2-nitrophenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-57	LC-MS: (ES, <i>m/z</i>): 352.
D-89	[5-(3-Cyclopropoxyphenyl)-1-[2-(propan-2-yl)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-58	LC-MS: (ES, <i>m/z</i>): 349.
D-90	[5-(3-Cyclopropoxyphenyl)-1-[2-(trifluoromethyl)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-59	LC-MS: (ES, <i>m/z</i>): 375.
D-91	[5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-60	LC-MS: (ES, <i>m/z</i>): 361.
D-92	[5-(3-Cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-61	LC-MS: (ES, <i>m/z</i>): 361.
D-93	[5-(3-Cyclopropoxyphenyl)-1-(1-ethyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-62	LC-MS: (ES, <i>m/z</i>): 375.
D-97	[5-(3-Cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -1,3-benzodiazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-63	LC-MS: (ES, <i>m/z</i>): 361.2.
D-99	[1-[(2-Cyclopropoxyphenyl)methyl]-5-(3-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-64	LC-MS: (ES, <i>m/z</i>): 351.
D-100	[1-[(2-chlorophenyl)methyl]-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-65	LC-MS: (ES, <i>m/z</i>): 359.


D-107	[5-(1-Benzothiophen-2-yl)-1-[(2-chlorophenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-66	LC-MS: (ES, <i>m/z</i>): 354.
D-110	[1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-67	LC-MS: (ES, <i>m/z</i>): 353.
D-113	[5-(3-Chloro-5-methoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-68	LC-MS: (ES, <i>m/z</i>): 373.
D-116	[1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-69	LC-MS: (ES, <i>m/z</i>): 353.
D-115	[1-[(2-Ethoxyphenyl)methyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-70	LC-MS: (ES, <i>m/z</i>): 377.
D-121	[5-(3-Ethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-71	LC-MS: (ES, <i>m/z</i>): 352.
D-130	[5-(3-Ethoxy-5-methoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-72	LC-MS: (ES, <i>m/z</i>): 383.
D-101	[5-(3-Cyclopropoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-73	LC-MS: (ES, <i>m/z</i>): 364.
D-102	[5-(3-cyclopropoxyphenyl)-1-[(2-nitrophenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-74	LC-MS: (ES, <i>m/z</i>): 366.
D-103	[1-[(2-Bromophenyl)methyl]-5-(3-cyclopropoxy-phenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-75	LC-MS: (ES, <i>m/z</i>): 399.
D-104	[1-[(2-chlorophenyl)methyl]-5-(2,3-dihydro-1,4-benzodioxin-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-76	LC-MS: (ES, <i>m/z</i>): 357.
D-105	[5-(2 <i>H</i> -1,3-benzodioxol-5-yl)-1-[(2-chlorophenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-77	LC-MS: (ES, <i>m/z</i>): 343.
D-106	[1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-78	LC-MS: (ES, <i>m/z</i>): 303.
D-108	[1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -indol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-79	LC-MS: (ES, <i>m/z</i>): 351.
D-109	[1-[(2-Ethoxyphenyl)methyl]-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-80	LC-MS: (ES, <i>m/z</i>): 363.
D-128	[5-(3,5-Diethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-81	LC-MS: (ES, <i>m/z</i>): 397.05.
D-111	[1-[(2-Chlorophenyl)methyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-82	LC-MS: (ES, <i>m/z</i>): 367.
D-112	[5-(1-Ethyl-1 <i>H</i> -indazol-6-yl)-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-83	LC-MS: (ES, <i>m/z</i>): 334.
D-114	[1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -indazol-5-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-84	LC-MS: (ES, <i>m/z</i>): 352.
D-118	[5-(3-Cyclobutoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-85	LC-MS: (ES, <i>m/z</i>): 379.

D-119	[1-[(2-Bromophenyl)methyl]-5-(3-cyclobutoxy-phenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-86	LC-MS: (ES, <i>m/z</i>): 412.
D-120	[5-[3-(2,2-dimethylpropoxy)phenyl]-1-[(1-methyl-1 <i>H</i> -indazol-7-yl)methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-87	LC-MS: (ES, <i>m/z</i>): 405.3.
D-123	[5-[3-(2-methylpropoxy)phenyl]-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-88	LC-MS: (ES, <i>m/z</i>): 338.
D-124	[5-(3,5-Dimethoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-89	LC-MS: (ES, <i>m/z</i>): 369.
D-125	[5-(3,5-Dimethoxyphenyl)-1-(2-phenylethyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-90	LC-MS: (ES, <i>m/z</i>): 339.
D-126	[5-(3,5-dimethoxyphenyl)-1-[(2-ethoxy-6-fluorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-91	LC-MS: (ES, <i>m/z</i>): 387.1.
D-127	[5-(3,5-dimethoxyphenyl)-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-92	LC-MS: (ES, <i>m/z</i>): 326.
D-129	[1-[(2-Chlorophenyl)methyl]-5-(3,5-diethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-93	LC-MS: (ES, <i>m/z</i>): 387.00.
D-131	[1-[(2-Chlorophenyl)methyl]-5-[2-(propan-2-yloxy)-1,3-oxazol-5-yl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-94	LC-MS: (ES, <i>m/z</i>): 348.
D-143	[1-[(2-Chlorophenyl)methyl]-5-[3-(oxetan-3-yl-methoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-95	LC-MS: (ES, <i>m/z</i>): 385.1.
D-144	[5-(3-Cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)-ethoxy]methyl]-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]-methanol	F-96	LC-MS: (ES, <i>m/z</i>): 477.
D-145	[5-(3-Cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)-ethoxy]methyl]-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]-methanol	F-97	LC-MS: (ES, <i>m/z</i>): 477.2.
D-146	[5-(3-Cyclopropoxyphenyl)-1-(3-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-98	LC-MS: (ES, <i>m/z</i>): 491.4.

Ester	Product	
D-60	[1-[(2-chlorophenyl)methyl]-5-(2-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methanol	F-99
D-61	[5-(4-bromothien-2-yl)-1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-100
D-62	[1-[(2-chlorophenyl)methyl]-5-[5-(2-methylpropoxy)-thien-2-yl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-101
D-63	1-[(2-chlorophenyl)-5-[2-(2-methylpropyl)-1,3-oxazol-5-yl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-102
D-64	[1-[(2-chlorophenyl)methyl]-5-[2-(2-methylpropoxy)-1,3-thiazol-5-yl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-103
D-75	[1-(2-Chlorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methanol	F-104
D-78	(5-Benzyl-1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl)-methanol	F-105

D-134	(5-(3-methoxyphenyl)-1-[[2-(propan-2-yloxy)phenyl]methyl]-1H-pyrazol-3-yl)methanol	F-106
D-135	(5-(3-methoxyphenyl)-1-(2-methylpropyl)-1H-pyrazol-3-yl)methanol	F-107
D-136	(5-(3-methoxyphenyl)-1-([oxan-4-yl]methyl)-1H-pyrazol-3-yl)methanol	F-108
D-138	(1-[(2-chlorophenyl)methyl]-5-[3-[(propan-2-yl)-carbamoyl]phenyl]-1H-pyrazol-3-yl)methanol	F-109
D-139	(1-[(2,6-dimethoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl)methanol	F-110
D-141	(1-[[4-(Dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl)methanol	F-111
D-142	(1-[[2-(dimethylamino)-6-fluorophenyl]methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl)methanol	F-112
D-147	(1-[(2-chlorophenyl)methyl]-5-phenoxy-1H-pyrazol-3-yl)methanol	F-113
D-1	(5-[[tert-Butoxy]carbonyl](phenyl)amino]-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl)methanol	F-114
D-140	(1-[(3-chloropyridin-2-yl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl)methanol	F-115

Intermediate F-116: [5-(3-Cyclopropoxyphenyl)-1-[[2-(dimethylamino)phenyl]methyl]-1H-pyrazol-3-yl]methanol



[0426] **[1-[(2-Aminophenyl)methyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]-methanol** In a 25-mL flask were combined Int. **F-74** [5-(3-cyclopropoxyphenyl)-1-[(2-nitrophenyl)methyl]-1H-pyrazol-3-yl]methanol (350 mg, 0.96 mmol, 1.00 equiv), AcOH (7.5 mL), H₂O (2.5 mL), and Zn (350 mg, 5.38 mmol, 5.60 equiv). The resulting mixture was stirred for 3 h at rt. The solids were removed. The resulting solution was extracted with 3x30 mL of EtOAc. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum to afford 300 mg (93%) of the title compound as a yellow oil. LC-MS (ES, *m/z*): 336.

[0427] **[5-(3-Cyclopropoxyphenyl)-1-[[2-(dimethylamino)phenyl]methyl]-1H-pyrazol-3-yl]methanol** Into a 15-mL flask were combined the product from the previous step (300 mg, 0.89 mmol, 1.00 equiv) MeOH (5 mL), AcOH (0.25 mL), HCHO (358 mg, 11.92 mmol, 4.00 equiv), and NaCNBH₃ (225 mg, 3.57 mmol, 4.00 equiv). The resulting solution was stirred for 2 h at 40 °C, quenched by the addition of 5 mL of water /


ice, and extracted with 3x30 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified by prep-TLC with EtOAc / hexane (1/1) to afford 300 mg (92%) of the title compound as a yellow oil. LC-MS: (ES, *m/z*): 365. ¹H NMR: (300 MHz, DMSO) δ 7.30 (dd, *J* = 8.8, 7.6 Hz, 1H), 7.22 (td, *J* = 7.6, 7.2, 1.6 Hz, 1H), 7.14 (dd, *J* = 8.1, 1.4 Hz, 1H), 7.00 (m, 2H), 6.88 (dd, *J* = 6.9, 1.5 Hz, 2H), 6.73 (dd, *J* = 7.7, 1.5 Hz, 1H), 6.42 (s, 1H), 5.35 (s, 2H), 4.46 (s, 2H), 3.67 (dt, *J* = 5.9, 3.1, 3.1 Hz, 1H), 2.55 (s, 6H), 0.62 (ddd, *J* = 5.9, 3.5, 1.3 Hz, 2H), 0.57 (m, 2H).

[0428] The following compounds were obtained via a similar zinc reduction / alkylation sequence.

Table 13: Zn reduction / alkylation of (nitroaryl) pyrazoles


Nitro Cpd	Product		Spectral
F-22	[1-[2-(Dimethylamino)phenyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]-methanol	F-117	LC-MS: (ES, <i>m/z</i>): 362.40.
F-57	[5-(3-cyclopropoxyphenyl)-1-[2-(dimethylamino)phenyl]-1H-pyrazol-3-yl]-methanol	F-118	LC-MS: (ES, <i>m/z</i>): 350.

Intermediate G-1: 3-(Bromomethyl)-1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazole



[0429] To a solution of Int. **F-8** (4.3 g, 13.08 mmol, 1.00 equiv) in CH₂Cl₂ (100 mL) was added CBr₄ (6.5 g, 1.50 equiv). This was followed by the addition of a solution of PPh₃ (7 g, 26.69 mmol, 2.00 equiv) in CH₂Cl₂ (10 mL) dropwise with stirring at 0 °C in 5 min. The resulting solution was stirred for 4 h at rt. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:5). This resulted in the title compound as a yellow oil (3.5 g, 68%).

[0430] The following substituted 3-(bromomethyl)pyrazoles were obtained from reaction of the corresponding (hydroxymethyl)pyrazole with PPh₃ / CBr₄:

Table 14: CBr₄ bromination of (hydroxymethyl) pyrazoles.


Alcohol	Bromo cpd		Spectral
F-6	3-(Bromomethyl)-1-[(2-chlorophenyl)-methyl]-5-[1,3-dimethyl-1H-thieno-[2,3-c]pyrazol-5-yl]-1H-pyrazole	G-2	LC-MS: (ES, <i>m/z</i>) 435.
F-9	3-(bromomethyl)-5-[3-(2-methylpropoxy)phenyl]-1-phenyl-1H-pyrazole	G-3	LC-MS: (ES, <i>m/z</i>): 385.
F-10	6-[3-(bromomethyl)-1-(2-bromophenyl)-1H-pyrazol-5-yl]-1-ethyl-1H-indazole	G-4	LC-MS: (ES, <i>m/z</i>): 459.
F-14	6-[3-(bromomethyl)-1-(2-bromophenyl)-1H-pyrazol-5-yl]-1-propyl-1H-indazole	G-5	LC-MS: (ES, <i>m/z</i>): 475.2.
F-19	3-(Bromomethyl)-1-(2-fluorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazole	G-6	LC-MS: (ES, <i>m/z</i>): 403.
F-27	7-[3-(bromomethyl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1H-pyrazol-1-yl]-1-methyl-1H-indazole	G-7	LC-MS: (ES, <i>m/z</i>): 453.
F-30	7-[3-(bromomethyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-1-yl]-1-methyl-1H-indazole	G-8	LC-MS: (ES, <i>m/z</i>): 441.3.
F-34	3-(Bromomethyl)-5-(3-methoxyphenyl)-1-[2-(propan-2-yloxy)phenyl]-1H-pyrazole	G-9	LC-MS: (ES, <i>m/z</i>): 401.
F-43	3-(bromomethyl)-1-(2-chlorophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazole	G-10	LC-MS: (ES, <i>m/z</i>): 405.15.
F-46	3-(bromomethyl)-1-(2-chlorophenyl)-5-(3,5-dimethoxyphenyl)-1H-pyrazole	G-11	LC-MS: (ES, <i>m/z</i>): 406. ¹ H-NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 7.63 (dt, <i>J</i> = 7.5, 2.0 Hz, 2H), 7.59 – 7.41 (m, 2H), 6.87 (s, 1H), 6.44 (t, <i>J</i> = 2.3 Hz, 1H), 6.34 (d, <i>J</i> = 2.3 Hz, 2H), 4.70 (s, 2H), 3.61(s, 6H).
F-49	3-(Bromomethyl)-1-(2-bromophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazole	G-12	LC-MS: (ES, <i>m/z</i>): 447. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.76 (dd, <i>J</i> = 7.8, 1.5 Hz, 1H), 7.50 (dddd, <i>J</i> = 29.7, 15.0, 7.5, 1.9 Hz, 3H), 7.23 (t, <i>J</i> = 8.0 Hz, 1H),

			6.97 – 6.85 (m, 2H), 6.85 – 6.75 (m, 2H), 4.78 (s, 1H), 4.68 (s, 1H), 3.55 (tt, $J = 6.0, 3.0$ Hz, 1H), 0.69 – 0.58 (m, 2H), 0.49 (q, $J = 2.7, 2.2$ Hz, 2H).
F-55	3-(Bromomethyl)-5-(3-cyclopropoxyphenyl)-1-(2-methoxyphenyl)-1H-pyrazole	G-13	LC-MS: (ES, m/z): 399
F-60	7-[3-(bromomethyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-1-yl]-1-methyl-1H-indazole	G-14	LC-MS: (ES, m/z): 425.
F-64	3-(Bromomethyl)-1-[(2-cyclopropoxyphenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazole	G-15	LC-MS: (ES, m/z): 414.
F-65	3-(Bromomethyl)-1-[(2-chlorophenyl)methyl]-5-(3,5-dimethoxyphenyl)-1H-pyrazole	G-16	LC-MS: (ES, m/z): 423. $^1\text{H-NMR}$: (CDCl_3 , ppm): δ : 7.42 – 7.30 (m, 2H), 7.29 – 7.16 (m, 4H), 6.83 – 6.71 (m, 2H), 6.52 – 6.36 (m, 8H), 5.45 (d, $J = 1.0$ Hz, 4H), 5.31 (s, 1H), 4.57 (s, 4H), 3.66 (s, 12H).
F-67	6-[3-(Bromomethyl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-5-yl]-1-methyl-1H-indazole	G-17	LC-MS: (ES, m/z): 417. $^1\text{H NMR}$: (300 MHz, CDCl_3) δ 8.01 (d, $J = 1.0$ Hz, 1H), 7.74 (dd, $J = 8.3, 0.8$ Hz, 1H), 7.42 – 7.33 (m, 1H), 7.28 (s, 1H), 7.26 (dd, $J = 3.8, 1.5$ Hz, 2H), 7.10 (dd, $J = 8.4, 1.4$ Hz, 1H), 6.95 – 6.84 (m, 1H), 6.58 (s, 1H), 5.48 (d, $J = 0.8$ Hz, 2H), 4.61 (s, 2H), 3.99 (s, 3H).
F-82	6-[3-(bromomethyl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-5-yl]-1-ethyl-1H-indazole	G-18	LC-MS: (ES, m/z): 431.
F-89	3-(Bromomethyl)-5-(3,5-dimethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazole	G-19	LC-MS: (ES, m/z): 431.0.
F-94	5-[3-(Bromomethyl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-5-yl]-2-(propan-2-yloxy)-1,3-oxazole	G-20	LC-MS: (ES, m/z): 411.
F-118	2-[3-(Bromomethyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-1-yl]- N,N -dimethylaniline	G-21	LC-MS: (ES, m/z): 413.

Alcohol	Bromo cpd	
---------	-----------	--

F-99	3-(Bromomethyl)-1-[(2-chlorophenyl)methyl]-5-(2-methoxyphenyl)-1 <i>H</i> -pyrazole	G-22
F-100	3-(bromomethyl)-5-(4-bromothien-2-yl)-[1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazole	G-23
F-106	3-(Bromomethyl)-(5-(3-methoxyphenyl)-1-[[2-(propan-2-yloxy)phenyl]methyl]-1 <i>H</i> -pyrazole	G-24
F-107	3-(Bromomethyl)-(5-(3-methoxyphenyl)-1-(2-methylpropyl)-1 <i>H</i> -pyrazole	G-25
F-108	3-(Bromomethyl)-(5-(3-methoxyphenyl)-1-([oxan-4-yl]methyl)-1 <i>H</i> -pyrazole	G-26
F-116	2-([3-(Bromomethyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-1-yl]methyl)- <i>N,N</i> -dimethylaniline	G-27

Intermediate H-1: Methyl 2-([1-[(3-chlorophenyl)methyl]-5-phenyl-1*H*-pyrazol-3-yl]methoxy)-2-methylpropanoate



[0431] To a solution of Int. **F-1** (0.288 g, 0.964 mmol) DMF was added NaH (60%) (0.077 g, 1.93 mmol) portionwise under N₂ at room temperature, and stirring was continued for 30 min. The mixture was cooled to 0 °C and methyl 2-bromo-2-methylpropanoate (0.16 mL, 1.25 mmol) was added dropwise followed by NaI (0.143 g, 0.964 mmol). The reaction mixture was warmed to room temperature gradually and stirred overnight. At the end of this period aq. NH₄Cl solution was added, and the mixture was partitioned with EtOAc (40 mL). The aqueous layer was extracted with EtOAc (2x 20 mL). The combined EtOAc layers were washed with brine and dried over Na₂SO₄, and the solvent was evaporated. The residue was chromatographed over SiO₂ (ISCO CombiFlash® Rf 200) using 0-50% gradient of EtOAc in hexanes to afford the title product as an oil (0.068 g, 38%). ¹H NMR (CDCl₃, 400 MHz): δ 1.56 (s, 6H), 3.68 (s, 3H), 4.58 (s, 2H), 5.25 (s, 2H), 6.45 (s, 1H), 6.87-6.93 (m, 1H), 7.00 (s, 1H), 7.16-7.22 (m, 2H), 7.26-7.34 (m, 2H), 7.34-7.42 (m, 3H).

[0432] The following substituted pyrazole carboxylic esters were obtained from Williamson ether synthesis of the corresponding (hydroxymethyl)pyrazole with $\text{BrC}(\text{CH}_3)_2\text{COOMe}$:




Table 15: Ether synthesis from (hydroxymethyl) pyrazoles.

Alcohol	Product		Spectral
F-10	Methyl 2-([1-(2-bromophenyl)-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-2	LC-MS: (ES, <i>m/z</i>): 499.1. ¹ H NMR: (300 MHz, DMSO) δ 8.01 (d, <i>J</i> = 1.0 Hz, 1H), 7.74 (dd, <i>J</i> = 8.0, 1.5 Hz, 1H), 7.66 (dd, <i>J</i> = 8.4, 0.8 Hz, 1H), 7.62 (dd, <i>J</i> = 7.8, 1.8 Hz, 1H), 7.53 (td, <i>J</i> = 7.7, 7.6, 1.5 Hz, 1H), 7.48 (dd, <i>J</i> = 1.5, 0.8 Hz, 1H), 7.44 (m, 1H), 6.96 (dd, <i>J</i> = 8.4, 1.4 Hz, 1H), 6.79 (s, 1H), 4.50 (s, 2H), 4.32 (q, <i>J</i> = 7.2, 7.2, 7.2 Hz, 2H), 3.72 (s, 3H), 1.47 (s, 6H), 1.24 (t, <i>J</i> = 7.2, 7.2 Hz, 3H).
F-11	Methyl 2-([1-(2-bromophenyl)-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-3	LC-MS: (ES, <i>m/z</i>): 483. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 7.99 (d, <i>J</i> = 1.0 Hz, 1H), 7.73 (dd, <i>J</i> = 7.9, 1.5 Hz, 1H), 7.67 – 7.47 (m, 4H), 7.42 (td, <i>J</i> = 7.6, 1.9 Hz, 1H), 6.86 (dd, <i>J</i> = 8.4, 1.4 Hz, 1H), 6.78 (s, 1H), 4.49 (s, 2H), 3.94 (s, 3H), 3.71 (s, 3H), 1.46 (s, 6H).
F-12	Methyl 2-([1-(2-bromophenyl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-4	LC-MS: (ES, <i>m/z</i>): 517.05. ¹ H NMR (300 MHz, DMSO) δ 7.76 (dd, <i>J</i> = 7.7, 1.3 Hz, 1H), 7.53 (m, 2H), 7.44 (ddd, <i>J</i> = 7.8, 6.6, 2.6 Hz, 1H), 7.18 (t, <i>J</i> = 8.0, 8.0 Hz, 1H), 6.81 (m, 2H), 6.70 (d, <i>J</i> = 2.4 Hz, 2H), 4.45 (s, 2H), 3.70 (s, 3H), 3.39 (s, 2H), 1.45 (s, 6H), 0.93 (s, 9H).
F-20	Methyl 2-([5-(3-cyclobutoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-5	LC-MS: (ES, <i>m/z</i>): 434. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.68 (s, 1H), 7.38 – 7.20 (m, 2H), 7.05 – 6.94 (m, 2H), 6.92 – 6.82 (m, 2H), 6.75 – 6.62 (m, 2H), 6.42 (s, 1H), 5.26 (s, 2H), 4.50 (q, <i>J</i> = 7.1 Hz, 1H), 4.42 (s, 2H), 4.02 (q, <i>J</i> = 6.9 Hz, 2H), 2.30 – 2.15 (m, 2H), 2.03 – 1.86 (m, 2H), 1.79 – 1.65 (m, 1H), 1.63 – 1.47 (m, 1H), 1.40 (s, 6H), 1.24 (t, <i>J</i> = 6.9 Hz, 3H).

F-25	Methyl 2-([5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-6	LC-MS: (ES, <i>m/z</i>): 472. ¹ H NMR: (400 MHz, DMSO- <i>d</i> ₆) δ 8.18 (s, 1H), 7.97 (d, <i>J</i> = 0.9 Hz, 1H), 7.90 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.59 (dd, <i>J</i> = 8.4, 0.8 Hz, 1H), 7.54 (q, <i>J</i> = 1.1 Hz, 1H), 7.46 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.18 (dd, <i>J</i> = 8.1, 7.3 Hz, 1H), 6.95 (dd, <i>J</i> = 8.4, 1.4 Hz, 1H), 6.91 (s, 1H), 4.55 (s, 2H), 4.26 (t, <i>J</i> = 7.2 Hz, 2H), 3.72 (s, 3H), 3.48 (s, 3H), 1.48 (s, 6H), 1.17 (t, <i>J</i> = 7.1 Hz, 3H).
F-26	Methyl 2-([5-(3-cyclobutoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-7	LC: (ES, <i>m/z</i>): 475. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 8.21 (s, 1H), 7.93 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.38 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.28 – 7.09 (m, 2H), 6.94 – 6.86 (m, 1H), 6.79 (s, 1H), 6.75 – 6.66 (m, 1H), 6.45 (dd, <i>J</i> = 2.5, 1.6 Hz, 1H), 4.50 (s, 2H), 4.19 – 4.05 (m, 1H), 3.70 (s, 3H), 3.43 (s, 3H), 2.10 – 1.95 (m, 2H), 1.89 – 1.74 (m, 2H), 1.67 (dd, <i>J</i> = 11.6, 8.3 Hz, 1H), 1.52 (d, <i>J</i> = 8.3 Hz, 2H), 1.45 (s, 6H).
F-27	Methyl 2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-8	LC-MS: (ES, <i>m/z</i>): 491.3. ¹ H NMR: (300 MHz, DMSO) δ 8.19 (s, 1H), 7.92 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.39 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.16 (m, 2H), 6.81 (m, 3H), 6.62 (m, 1H), 4.51 (s, 2H), 3.70 (s, 3H), 3.42 (s, 3H), 3.17 (d, <i>J</i> = 6.6 Hz, 2H), 1.46 (s, 6H), 0.87 (s, 9H).
F-28	Methyl 2-([5-[3-(cyclobutylmethoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-9	LC-MS: (ES, <i>m/z</i>): 489. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 8.19 (s, 1H), 7.92 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.39 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.28 – 7.02 (m, 2H), 6.94 – 6.71 (m, 3H), 6.65 (dd, <i>J</i> = 2.6, 1.6 Hz, 1H), 4.50 (s, 2H), 3.70 (s, 3H), 3.56 (d, <i>J</i> = 6.8 Hz, 2H), 3.43 (s, 3H), 3.16 (s, 1H), 2.04 – 1.91 (m, 2H), 1.89 – 1.77 (m, 2H), 1.66 (dt, <i>J</i> = 11.0, 7.6 Hz, 2H), 1.45 (s, 6H).
F-30	Methyl 2-methyl-2-([1-(1-methyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy]propanoate	H-10	LC-MS: (ES, <i>m/z</i>): 477. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): δ 8.20 (s, 1H), 7.92 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.40 (dd, <i>J</i> = 7.4, 1.0 Hz, 1H), 7.26 – 7.08 (m, 2H), 6.91 – 6.74 (m, 3H), 6.64 (dd, <i>J</i> = 2.5, 1.5 Hz, 1H), 4.52 (s, 2H), 3.71 (s, 3H), 3.44 (s, 3H), 3.35 (d, <i>J</i> = 6.6 Hz, 2H), 1.81 (dh, <i>J</i> = 13.4, 6.8 Hz, 1H), 1.47 (s, 6H), 0.85 (d, <i>J</i> = 6.6 Hz, 6H).
F-33	Methyl 2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-ethyl-1 <i>H</i> -indazol-7-yl)-	H-11	LC-MS: (ES, <i>m/z</i>): 504. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.25 (s, 1H), 7.93 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.33

	1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate		(dd, $J = 7.3, 1.0$ Hz, 1H), 7.27 – 7.07 (m, 2H), 6.87 (s, 1H), 6.84 – 6.72 (m, 2H), 6.61 (dd, $J = 2.5, 1.6$ Hz, 1H), 4.52 (s, 2H), 3.71 (s, 2H), 3.18 (s, 3H), 3.16 (s, 2H), 1.47 (s, 6H), 1.10 (t, $J = 7.1$ Hz, 3H), 0.87 (s, 9H).
F-47	Methyl 2-([1-(2-chlorophenyl)-5-(3,5-diethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]-methoxy)-2-methylpropanoate	H-12	LC-MS: (ES, m/z): 473.05. ¹ H NMR: (300 MHz, DMSO) δ 7.57 (m, 4H), 6.70 (s, 1H), 6.38 (t, $J = 2.3, 2.3$ Hz, 1H), 6.30 (d, $J = 2.2$ Hz, 2H), 4.45 (s, 2H), 3.85 (q, $J = 7.0, 7.0, 7.0$ Hz, 4H), 3.71 (s, 3H), 1.45 (s, 6H), 1.22 (t, $J = 7.0, 7.0$ Hz, 6H).
F-48	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(2-fluorophenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-13	LC-MS: (ES, m/z): 425.2. ¹ H-NMR: δ_{H} (300 MHz, DMSO- d_6) 7.54 (2 H, m), 7.31 (3 H, m), 6.90 (3 H, m), 6.67 (1 H, s), 4.45 (2 H, s), 3.69 (3 H, s), 3.61 (1 H, dp, $J = 6.1, 3.1, 3.1, 3.0, 3.0$ Hz), 1.44 (6 H, s), 0.54 (4 H, m).
F-61	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-14	LC-MS: (ES, m/z): 461. ¹ H NMR (300 MHz, DMSO- d_6) δ 7.77 – 7.62 (m, 2H), 7.42 (dd, $J = 8.6, 7.3$ Hz, 1H), 7.23 (t, $J = 7.9$ Hz, 1H), 7.01 – 6.84 (m, 4H), 6.72 (s, 1H), 4.52 (s, 2H), 4.08 (s, 3H), 3.71 (s, 3H), 3.57 (tt, $J = 6.2, 3.0$ Hz, 1H), 1.48 (s, 6H), 0.60 – 0.46 (m, 2H), 0.40 (dq, $J = 6.8, 3.4$ Hz, 2H).
F-62	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(1-ethyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-15	LC-MS: (ES, m/z): 475. ¹ H NMR (300 MHz, DMSO- d_6) δ 8.24 (s, 1H), 7.91 (dd, $J = 8.0, 1.0$ Hz, 1H), 7.30 (dd, $J = 7.3, 1.0$ Hz, 1H), 7.27 – 7.05 (m, 2H), 6.93 (dt, $J = 7.8, 1.2$ Hz, 1H), 6.89 – 6.80 (m, 2H), 6.76 (dd, $J = 2.5, 1.6$ Hz, 1H), 4.51 (s, 2H), 3.70 (s, 3H), 3.35 (d, $J = 4.6$ Hz, 2H), 3.33 – 3.29 (m, 1H), 1.45 (s, 6H), 1.10 (t, $J = 7.1$ Hz, 3H), 0.52 – 0.28 (m, 4H).
F-66	Methyl 2-([5-(1-benzothiophen-2-yl)-1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-16	LC-MS: (ES, m/z): 454. ¹ H NMR: (300 MHz, DMSO- d_6) δ 8.04 – 7.96 (m, 1H), 7.90 – 7.82 (m, 1H), 7.59 (d, $J = 0.8$ Hz, 1H), 7.53 – 7.47 (m, 1H), 7.46 – 7.39 (m, 2H), 7.32 (tt, $J = 7.4, 5.5$ Hz, 2H), 6.78 – 6.70 (m, 1H), 6.69 (s, 1H), 5.63 (s, 2H), 4.43 (s, 2H), 3.71 (s, 3H), 1.44 (s, 6H).
F-69	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-17	LC-MS: (ES, m/z): 453. ¹ H NMR (300 MHz, DMSO- d_6) δ 7.94 (d, $J = 1.0$ Hz, 1H), 7.79 – 7.65 (m, 1H), 7.45 (dd, $J = 8.5, 7.1$ Hz, 1H), 7.41 – 7.32 (m, 1H), 7.29 – 7.21 (m, 2H), 7.10 (dd, $J = 7.1, 0.8$ Hz, 1H), 6.85 – 6.73 (m, 1H), 6.58 (s,

			1H), 5.40 (s, 2H), 4.45 (s, 2H), 4.09 (s, 3H), 3.69 (s, 3H), 1.44 (s, 6H).
F-70	Methyl 2-([1-[(2-ethoxy-phenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-18	LC-MS: (ES, <i>m/z</i>): 477. ¹ H NMR: (400 MHz, DMSO- <i>d</i> ₆) δ 8.09 (d, <i>J</i> = 1.0 Hz, 1H), 7.80 (dd, <i>J</i> = 8.3, 0.8 Hz, 1H), 7.68 (q, <i>J</i> = 1.0 Hz, 1H), 7.27 – 7.19 (m, 1H), 7.16 (dd, <i>J</i> = 8.4, 1.4 Hz, 1H), 6.95 (dd, <i>J</i> = 8.2, 1.1 Hz, 1H), 6.86 (td, <i>J</i> = 7.5, 1.1 Hz, 1H), 6.72 (dd, <i>J</i> = 7.5, 1.7 Hz, 1H), 6.51 (s, 1H), 5.33 (s, 2H), 4.41 (s, 2H), 4.40 – 4.35 (m, 2H), 3.93 (d, <i>J</i> = 7.0 Hz, 2H), 3.70 (s, 3H), 1.44 (s, 6H), 1.33 (t, <i>J</i> = 7.2 Hz, 3H), 1.12 (t, <i>J</i> = 6.9 Hz, 3H).
F-71	Methyl 2-([5-(3-ethoxy-phenyl)-1-[(2-ethoxy-phenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-19	LC-MS: (ES, <i>m/z</i>): 452. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.33 (t, <i>J</i> = 7.9 Hz, 1H), 7.24 (ddd, <i>J</i> = 8.2, 7.3, 1.7 Hz, 1H), 7.04 – 6.91 (m, 3H), 6.90 – 6.79 (m, 2H), 6.69 (dd, <i>J</i> = 7.6, 1.7 Hz, 1H), 6.43 (s, 1H), 5.77 (s, 1H), 5.26 (s, 2H), 4.38 (s, 2H), 3.96 (dq, <i>J</i> = 26.5, 6.9 Hz, 4H), 3.69 (s, 3H), 1.42 (s, 6H), 1.25 (dt, <i>J</i> = 18.7, 7.0 Hz, 6H).
F-75	Methyl 2-([1-[(2-bromo-phenyl)methyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-20	LC-MS: (ES, <i>m/z</i>): 499.3. ¹ H NMR: (300 MHz, DMSO) δ 7.62 (dd, <i>J</i> = 7.9, 1.3 Hz, 1H), 7.35 (tdd, <i>J</i> = 7.5, 7.5, 2.6, 1.0 Hz, 2H), 7.23 (td, <i>J</i> = 7.6, 7.6, 1.8 Hz, 1H), 7.01 (m, 3H), 6.69 (dd, <i>J</i> = 7.7, 1.7 Hz, 1H), 6.47 (s, 1H), 5.34 (s, 2H), 4.40 (s, 2H), 3.68 (s, 4H), 1.42 (s, 6H), 0.59 (tt, <i>J</i> = 5.8, 5.8, 2.4, 2.4 Hz, 4H).
F-79	Methyl 2-([1-[(2-chloro-phenyl)methyl]-5-(1-methyl-1H-indol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-21	LC-MS: (ES, <i>m/z</i>): 451. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.59 (d, <i>J</i> = 8.1 Hz, 1H), 7.48 – 7.42 (m, 2H), 7.41 (d, <i>J</i> = 3.1 Hz, 1H), 7.34 – 7.29 (m, 2H), 7.03 (dd, <i>J</i> = 8.2, 1.5 Hz, 1H), 6.84 – 6.76 (m, 1H), 6.49 – 6.42 (m, 2H), 5.44 (s, 2H), 4.42 (s, 2H), 3.74 (s, 3H), 3.70 (s, 3H), 1.44 (s, 6H).
F-80	Methyl 2-([1-[(2-ethoxy-phenyl)methyl]-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-22	LC-MS: (ES, <i>m/z</i>): 463. ¹ H NMR: (400 MHz, DMSO- <i>d</i> ₆) δ 8.09 (d, <i>J</i> = 0.9 Hz, 1H), 7.80 (dd, <i>J</i> = 8.3, 0.8 Hz, 1H), 7.69 (q, <i>J</i> = 1.0 Hz, 1H), 7.28 – 7.20 (m, 1H), 7.16 (dd, <i>J</i> = 8.4, 1.4 Hz, 1H), 6.95 (dd, <i>J</i> = 8.4, 1.1 Hz, 1H), 6.86 (td, <i>J</i> = 7.5, 1.0 Hz, 1H), 6.75 (dd, <i>J</i> = 7.5, 1.7 Hz, 1H), 6.49 (s, 1H), 5.32 (s, 2H), 4.41 (s, 2H), 4.01 (s, 3H), 3.93 (q, <i>J</i> = 7.0 Hz, 2H), 3.70 (s, 3H), 1.43 (s, 6H), 1.12 (t, <i>J</i> = 6.9 Hz, 3H).
F-90	Methyl 2-([5-(3,5-di-methoxyphenyl)-1-(2-	H-23	LC-MS: (ES, <i>m/z</i>): 439.

	phenylethyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.29 – 7.11 (m, 3H), 7.04 – 6.94 (m, 2H), 6.54 (t, <i>J</i> = 2.3 Hz, 1H), 6.37 (d, <i>J</i> = 2.3 Hz, 2H), 6.26 (s, 1H), 4.39 (s, 2H), 4.25 (dd, <i>J</i> = 8.1, 6.6 Hz, 2H), 3.74 (s, 6H), 3.70 (s, 3H), 3.05 (t, <i>J</i> = 7.3 Hz, 2H), 1.43 (s, 6H).
--	---	--	--

Alcohol	Product		Spectral
F-13	Methyl 2-([1-(2-bromophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-24	LC-MS: (ES, <i>m/z</i>): 489.
F-14	Methyl 2-([1-(2-bromophenyl)-5-(1-propyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-25	LC-MS: (ES, <i>m/z</i>): 535.1.
F-15	Methyl 2-([1-(2-bromophenyl)-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-26	LC-MS: (ES, <i>m/z</i>): 499.3.
F-16	Methyl 2-([1-(2-fluorophenyl)-5-[3-(oxetan-3-ylmethoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-27	LC-MS: (ES, <i>m/z</i>): 455.3.
F-17	Methyl 2-([5-(3-cyclobutoxyphenyl)-1-(2-fluorophenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-28	LC-MS: (ES, <i>m/z</i>): 439.
F-18	Methyl 2-([1-(2-fluorophenyl)-5-[3-(oxetan-3-yloxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-29	LC-MS: (ES, <i>m/z</i>): 440.
F-21	Methyl 2-([5-(3,5-diethoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-30	LC-MS: (ES, <i>m/z</i>): 453.25.
F-24	Methyl 2-([5-(3,5-dimethoxyphenyl)-1-(2-ethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-31	LC-MS: (ES, <i>m/z</i>): 455.
F-29	Methyl 2-([5-[3-(cyclopropylmethoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-32	LC-MS: (ES, <i>m/z</i>): 475.
F-31	Methyl 2-([5-(3,5-dimethoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-33	LC-MS: (ES, <i>m/z</i>): 465.
F-32	Methyl 2-([5-(3,5-diethoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-34	LC-MS: (ES, <i>m/z</i>): 493.15.
F-35	Methyl 2-([1-(1,3-dimethyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-35	LC-MS: (ES, <i>m/z</i>): 505.3.
F-36	Methyl 2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -1,2,3-benzotriazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-36	LC-MS: (ES, <i>m/z</i>): 492
F-37	Methyl 2-([1-(2-chlorophenyl)-5-[3-(oxetan-3-ylmethoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-37	LC-MS: (ES, <i>m/z</i>): 471.3.
F-38	Methyl 2-([1-(2-chlorophenyl)-5-(pyridin-2-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-38	LC-MS: (ES, <i>m/z</i>): 386.00.

F-39	Methyl 2-([1-(2-chlorophenyl)-5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-39	LC-MS: (ES, <i>m/z</i>): 455.
F-40	Methyl 2-([1-(2-chlorophenyl)-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-40	LC-MS: (ES, <i>m/z</i>): 453.
F-41	Methyl 2-([1-(2-chlorophenyl)-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-41	LC-MS: (ES, <i>m/z</i>): 455.
F-42	Methyl 2-([1-(2-chlorophenyl)-5-[3-(oxetan-3-yloxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-42	LC-MS: (ES, <i>m/z</i>): 457.
F-44	Methyl 2-([1-(2-chlorophenyl)-5-[3-(2,2-dimethylpropoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-43	LC-MS: (ES, <i>m/z</i>): 471.10.
F-45	methyl 2-([1-(2-chlorophenyl)-5-[3-(cyclobutylmethoxy)-phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-44	LC-MS: (ES, <i>m/z</i>): 469.15.
F-46	Methyl 2-([1-(2-chlorophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-45	LC-MS: (ES, <i>m/z</i>): 445.
F-49	Methyl 2-([1-(2-bromophenyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-46	LC-MS: (ES, <i>m/z</i>): 486.95
F-50	Methyl 2-([1-(2-bromo-4-fluorophenyl)-5-(3-cyclopropoxy-phenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-47	LC-MS: (ES, <i>m/z</i>): 503.1.
F-51	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(2,4-dichloro-phenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-48	LC-MS: (ES, <i>m/z</i>): 475.05.
F-52	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(2,5-dichloro-phenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-49	LC-MS: (ES, <i>m/z</i>): 475.1.
F-53	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(difluoro-methoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methyl-propanoate	H-50	LC-MS: (ES, <i>m/z</i>): 473.
F-54	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(2-ethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-51	LC-MS: (ES, <i>m/z</i>): 450.
F-56	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-52	LC-MS: (ES, <i>m/z</i>): 421.
F-58	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(propan-2-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-53	LC-MS: (ES, <i>m/z</i>): 449.
F-59	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(trifluoro-methyl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methyl-propanoate	H-54	LC-MS: (ES, <i>m/z</i>): 475.
F-60	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-55	LC-MS: (ES, <i>m/z</i>): 461.
F-63	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(1-methyl-1 <i>H</i> -1,3-benzodiazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methyl-propanoate	H-56	LC-MS: (ES, <i>m/z</i>): 461.1.
F-68	Methyl 2-([5-(3-chloro-5-methoxyphenyl)-1-[(2-ethoxy-phenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-57	LC-MS: (ES, <i>m/z</i>): 473

F-72	Methyl 2-([5-(3-ethoxy-5-methoxyphenyl)-1-[(2-ethoxy-phenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-58	LC-MS: (ES, <i>m/z</i>): 483.
F-73	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-59	LC-MS: (ES, <i>m/z</i>): 465.
F-76	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(2,3-dihydro-1,4-benzodioxin-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-60	LC-MS: (ES, <i>m/z</i>): 457.
F-77	Methyl 2-([5-(2 <i>H</i> -1,3-benzodioxol-5-yl)-1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-61	LC-MS: (ES, <i>m/z</i>): 443.
F-78	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-62	LC-MS: (ES, <i>m/z</i>): 403.
F-81	Methyl 2-([5-(3,5-diethoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-63	LC-MS: (ES, <i>m/z</i>): 497.25.
F-82	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-64	LC-MS: (ES, <i>m/z</i>): 467.
F-83	Methyl 2-([5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-65	LC-MS: (ES, <i>m/z</i>): 434.
F-84	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -indazol-5-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-66	LC-MS: (ES, <i>m/z</i>): 452.
F-85	Methyl 2-([5-(3-cyclobutoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-67	LC-MS: (ES, <i>m/z</i>): 479.
F-86	Methyl 2-([1-[(2-bromophenyl)methyl]-5-(3-cyclobutoxy-phenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-68	LC-MS: (ES, <i>m/z</i>): 512.
F-87	Methyl 2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-[(1-methyl-1 <i>H</i> -indazol-7-yl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-69	LC-MS: (ES, <i>m/z</i>): 505.
F-88	Methyl 2-methyl-2-([5-[3-(2-methylpropoxy)phenyl]-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)propanoate	H-70	LC-MS: (ES, <i>m/z</i>): 438.
F-89	Methyl 2-([5-(3,5-dimethoxyphenyl)-1-[(2-ethoxyphenyl)-methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-71	LC-MS: (ES, <i>m/z</i>): 469.
F-91	Methyl 2-([5-(3,5-dimethoxyphenyl)-1-[(2-ethoxy-6-fluorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-72	LC-MS: (ES, <i>m/z</i>): 487.2.
F-92	Methyl 2-([5-(3,5-dimethoxyphenyl)-1-([pyridin-2-yl]methyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-73	LC-MS: (ES, <i>m/z</i>): 426.
F-93	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3,5-diethoxy-phenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-74	LC-MS: (ES, <i>m/z</i>): 487.2.
F-95	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[3-(oxetan-3-yl-methoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-75	LC-MS: (ES, <i>m/z</i>): 485.2.

F-96	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)ethoxy]methyl]-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]-methoxy)-2-methylpropanoate	H-76	LC-MS: (ES, <i>m/z</i>): 577.
F-97	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)ethoxy]methyl]-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]-methoxy)-2-methylpropanoate	H-77	LC-MS: (ES, <i>m/z</i>): 577.3.
F-98	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(3-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1 <i>H</i> -indazol-4-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-78	LC-MS: (ES, <i>m/z</i>): 591.3.
F-4	Methyl 2-([5-(3-methoxyphenyl)-1-(2-methylphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-79	LC-MS: (ES, <i>m/z</i>): 394.
F-2	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-methyl-1 <i>H</i> -1,3-benzodiazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-80	LC-MS: (ES, <i>m/z</i>): 453.
F-5	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1 <i>H</i> -1,2,3-benzotriazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-81	LC-MS: (ES, <i>m/z</i>): 468.
F-117	Methyl 2-([1-[2-(dimethylamino)phenyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-82	LC-MS: (ES, <i>m/z</i>): 478.3.

Alcohol	Product	
F-23	Methyl 2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-(quinolin-8-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-83
F-101	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[5-(2-methylpropoxy)thien-2-yl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-84
F-102	Methyl 2-(1-[(2-chlorophenyl)-5-[2-(2-methylpropyl)-1,3-oxazol-5-yl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-85
F-103	Methyl 2-(1-[(2-chlorophenyl)methyl]-5-[2-(2-methylpropoxy)-1,3-thiazol-5-yl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-86
F-104	Methyl 2-([1-(2-chlorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-87
F-105	2-([5-Benzyl-1-[(2-chlorophenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid	H-88
F-110	Methyl 2-[1-[(2,6-dimethoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-89
F-111	Methyl 2-([1-[[4-(dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-90
F-112	Methyl 2-(1-[[2-(dimethylamino)-6-fluorophenyl]methyl]-5-(3-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-91
F-113	Methyl (1-[(2-chlorophenyl)methyl]-5-phenoxy-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-92

F-114	Methyl 2-[(5-[[[(tert-Butoxy)carbonyl](phenyl)amino]-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl)methoxy]-2-methylpropanoate	H-93
F-115	Methyl 2-[(1-[(3-chloropyridin-2-yl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoate	H-94
F-7	Methyl 2-[(1-[(2-chlorophenyl)methyl]-5-(3-nitrophenyl)-1H-pyrazole-3-yl)methoxy]-2-methylpropanoate	H-95

[0433] The following esters of 2-hydroxy-2-methylbutanoic acid were prepared.

[0434] **Methyl 2-hydroxy-2-methylbutanoate** To a solution of 2-hydroxy-2-methylbutanoic acid (3.0 g, 25.40 mmol, 1.00 equiv) in MeOH (100 mL) was added H₂SO₄ (3 mL) dropwise with stirring at 0 °C. The resulting solution was heated to reflux for 16 h, cooled, concentrated under vacuum, diluted with 200 mL of EtOAc, washed with 1x100 mL of (sat)NaHCO₃(aq), dried over Na₂SO₄, and concentrated under vacuum to afford 780 mg (23%) of the title compound as a colorless oil.

[0435] **Methyl (2R)-2-hydroxy-2-methylbutanoate** To a solution of (2R)-2-hydroxy-2-methylbutanoic acid (1.0 g, 8.47 mmol, 1.00 equiv) in MeOH (50 mL) was added SOCl₂ (2.0 g, 2.00 equiv) dropwise with stirring at 0 °C. The resulting solution was heated to reflux for 16 h, then concentrated under vacuum, diluted with 100 mL of CH₂Cl₂. The mixture was dried over Na₂SO₄ and concentrated under vacuum, affording 700 mg (63%) of the title compound as a colorless oil. ¹H-NMR: (300 MHz, CDCl₃) δ : 3.79 (s, 3H), 3.04 (s, 1H), 1.89 – 1.61 (m, 2H), 1.41 (s, 3H), 0.88 (t, J = 7.4 Hz, 3H).

[0436] **Methyl (2S)-2-hydroxy-2-methylbutanoate** To a solution of (2S)-2-hydroxy-2-methylbutanoic acid (500 mg, 4.23 mmol, 1.00 equiv) in MeOH (30 mL) was added SOCl₂ (1.03 mg, 0.01 mmol, 2.00 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 16 h at 60 °C, then concentrated under vacuum, diluted with 200 mL of CH₂Cl₂, washed with 1x100 mL of (sat)NaHCO₃(aq), and evaporated under reduced pressure. The solid that formed was dried in an oven under reduced pressure, affording 400 mg (72%) of the title compound as colorless oil. LC-MS: (ES, *m/z*): 132. ¹H NMR (DMSO) δ: 5.10 (s, 1H), 3.62 (s, 3H), 1.73 – 1.44 (m, 2H), 1.24 (s, 3H), 0.78 (t, J = 7.4 Hz, 3H).

[0437] The following substituted pyrazole carboxylic esters were obtained from Williamson ether synthesis of the corresponding (bromomethyl)pyrazole with HO-C(CH₃)₂COOMe:





Table 16: Ether synthesis from (bromomethyl) pyrazoles.

Bromo cpd	Product		Spectral
G-3	Methyl 2-methyl-2-([5-[3-(2-methylpropoxy)phenyl]-1-phenyl-1H-pyrazol-3-yl]methoxy)propanoate	H-96	LC-MS: (ES, <i>m/z</i>): 423.
G-5	Ethyl (2 <i>R</i>)-2-([1-(2-bromophenyl)-5-(1-propyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate (This reaction was performed with Methyl (2 <i>R</i>)-2-hydroxy-2-methylbutanoate)	H-97	LC-MS: (ES, <i>m/z</i>): 541.1.
G-6	Methyl 2-([1-(2-fluorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-98	LC-MS: (ES, <i>m/z</i>): 441.
G-9	Methyl 2-([5-(3-methoxyphenyl)-1-[2-(propan-2-yl-oxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-99	LC-MS: (ES, <i>m/z</i>): 439.
G-22	Methyl 2-[3-(bromomethyl)-1-[(2-chlorophenyl)-methyl]-5-(2-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]-methoxy)-2-methylpropanoate	H-100	
G-10	Methyl 2-([1-(2-chlorophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-101	
G-12	Methyl (2 <i>R</i>)-2-([1-(2-bromophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate	H-102	LC-MS: (ES, <i>m/z</i>): 499.3.
G-13	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-(2-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-103	LC-MS: (ES, <i>m/z</i>): 437.
G-25	Methyl 2-([5-(3-methoxyphenyl)-1-(2-methylpropyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-104	
G-26	Methyl 2-([5-(3-methoxyphenyl)-1-([oxan-4-yl]-methyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-105	
G-15	Methyl 2-([1-[(2-cyclopropoxyphenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-106	LC-MS: (ES, <i>m/z</i>): 451.
G-16	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-107	LC-MS: (ES, <i>m/z</i>): 459.

G-17	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-108	LC-MS: (ES, <i>m/z</i>): 453.
G-20	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[2-(propan-2-yloxy)-1,3-oxazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-109	LC-MS: (ES, <i>m/z</i>): 448.
G-2	Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[1,3-dimethyl-1H-thieno[2,3-c]pyrazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-110	LC-MS: (ES, <i>m/z</i>): 473.
G-27	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[[2-(dimethylamino)phenyl]methyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoate	H-111	LC-MS: (ES, <i>m/z</i>): 464.
G-21	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(dimethylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-112	LC-MS: (ES, <i>m/z</i>): 450
G-24	Methyl 2-([5-(3-methoxyphenyl)-1-[[2-(propan-2-yloxy)phenyl]methyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoate	H-113	

Intermediate H-114: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(4-cyclopropoxythien-2-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0438] **tert-Butyl 2-([1-[(2-chlorophenyl)methyl]-5-(4-bromothiophen-2-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate** To a solution of tert-butyl 2-hydroxy-2-methylpropanoate (2.14 g, 13.36 mmol, 3.00 equiv) in THF (50 mL) was added NaH (535 mg, 13.38 mmol, 3.00 equiv), in portions at 0 °C. The mixture was stirred at 0 °C for 20 min. To this was added Int **G-23** (2 g, 4.48 mmol, 1.00 equiv) at 0 °C and Bu₄NI (1.0 g, 2.72 mmol, 0.50 equiv). The resulting solution was stirred for 16 h at rt then quenched by the addition of water/ice. The resulting solution was extracted with 500 mL of EtOAc, and the organic layers were combined and dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:2). This resulted in the title compound as a yellow oil (400 mg, 17%).

[0439] **Methyl 2-([5-(4-bromothiien-2-yl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate.** To a solution of the product from the previous step (400 mg, 0.76 mmol, 1.00 equiv) in MeOH (20 mL) was added aq HCl (1 mL) dropwise with stirring at rt. The resulting solution was stirred for 3 h at 65 °C, concentrated under vacuum, and diluted with 100 mL of EtOAc. The pH value of the solution was adjusted to 7-8 with (sat) NaHCO₃ (aq). The mixture was dried over anhydrous Na₂SO₄ and concentrated under vacuum. This resulted in the title compound as a yellow oil (350 mg, 95%).


[0440] **Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(4-cyclopropoxythien-2-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate** To a solution of the product from the previous step (350 mg, 0.72 mmol, 1.00 equiv) in dioxane (10 mL) was added stepwise Cs₂CO₃ (473 mg, 1.45 mmol, 2.00 equiv), 2nd Generation XantPhos precatalyst (322 mg, 0.36 mmol, 0.50 equiv), and cyclopropanol (168 mg, 2.89 mmol, 4.00 equiv). The final reaction mixture was irradiated with microwave radiation for 1 h at 85 °C. The resulting solution was diluted with 100 mL of EtOAc. The resulting mixture was washed with 2x100 mL of brine. The mixture was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:2). This resulted in the title compound as a yellow oil (150 mg, 45%).

Intermediate H-115: Methyl 2-([5-(3-aminophenyl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0441] To a solution of Int. **H-95** (400 mg, 0.90 mmol, 1.00 equiv) in AcOH / H₂O (10/1 mL) was added Zn (400 mg, 6.15 mmol), in portions at room temperature. The resulting mixture was stirred for 2 h at room temperature. The solids were removed by filtration. The pH value of the solution was adjusted to 7-8 with sat NaHCO₃. The resulting solution was extracted with 100 mL of EtOAc, and the organic layers were combined, washed with 2 x 200 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified by Prep-TLC with DCM/MeOH (30:1), affording 230 mg (62%) of the title product as a yellow oil.

Intermediate H-116: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[3-(2-methylpropanamido)-phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0442] A solution of Int. **H-115** (200 mg, 0.48 mmol, 1.00 equiv), HATU (276 mg, 0.73 mmol, 1.50 equiv), 2-methylpropanoic acid (130 mg, 1.48 mmol, 3.00 equiv) and DIEA (187 mg, 1.45 mmol, 3.00 equiv) in DMF (20 mL) was stirred overnight at rt, then diluted with 200 mL of EtOAc, washed with 2x100 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified by Prep-TLC with DCM/MeOH (30:1), to afford 155 mg (66%) of the title product as a yellow solid.

Intermediate H-117: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[3-(methylsulfanyl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0443] To a solution of Int. **H-115** (700 mg, 1.69 mmol, 1.00 equiv) in MeCN (10 mL), was added dimethyl disulfide (319 mg, 3.39 mmol, 2.00 equiv). The resulting mixture was heated 60 °C for 1 h, then cooled to rt. tert-Butyl nitrite (350 mg, 3.40 mmol, 2.00 equiv) was then added dropwise with stirring, and the resulting solution was stirred for 3 h at 60 °C, then diluted with 200 mL of EtOAc, washed with 2x100 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto Prep-TLC with EtOAc / petroleum ether (1:2). This resulted in 300 mg (40%) of the title product as a yellow oil.

Intermediate H-118: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methanesulfonylphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0444] To a solution of Int. **H-117** (300 mg, 0.67 mmol, 1.00 equiv) in MeOH / H₂O (10/5 mL) was added K₂S₂O₈ (233 mg, 1.53 mmol, 1.00 equiv), in portions at rt. The resulting solution was stirred for 2 h, then diluted with 200 mL of EtOAc, washed with 2x100 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto Prep-TLC with EtOAc / petroleum ether (1:3). This resulted in 180 mg (58%) of the title product as a yellow solid.

Intermediate H-119: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methanesulfonamidophenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0445] To a solution of Int. **H-115** (200 mg, 0.48 mmol, 1.00 equiv), pyridine (75 mg, 0.95 mmol, 2.00 equiv), and DMAP (6 mg, 0.05 mmol, 0.10 equiv) in DCM (10 mL). was added methanesulfonyl chloride (83 mg, 0.72 mmol, 1.50 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred overnight at rt, then diluted with 100 mL of DCM, washed with 2x100 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto Prep-TLC with EtOAc / petroleum ether (1:4). This resulted in 100 mg (42%) of the title product as a yellow oil.

Intermediate H-120: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[3-(methylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate



[0446] To a solution of Int. **H-115** (200 mg, 0.48 mmol, 1.00 equiv) in MeOH (10 mL) was added paraformaldehyde (107 mg, 2.43 mmol, 5.00 equiv), followed by the addition of MeONa (0.45 mL, 5.00 equiv, 5.4M) dropwise with stirring at room temperature. The resulting solution was stirred for 2 h at 60 °C, then cooled to 0 °C. NaBH₄ (92 mg, 2.43 mmol, 5.00 equiv) was added, and the solution was stirred for 1 h at 60 °C, then cooled, diluted with 100 mL of EtOAc, washed with 2x50 mL of brine, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto Prep-TLC with EtOAc /petroleum ether (1:4). This resulted in 100 mg (48%) of the title product as a yellow oil.

Intermediate H-121: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-ethylbutanoate




[0447] **Methyl 2-ethyl-2-hydroxybutanoate** To a solution of 2-ethyl-2-hydroxybutanoic acid (2.0 g, 15.13 mmol, 1.00 equiv) in MeOH (100 mL) was added H₂SO₄ (3 mL) dropwise with stirring at rt. The resulting solution was heated to reflux for 16 h, then concentrated under vacuum, diluted with 200 mL of EtOAc, washed with 2x100 mL of sat NaHCO₃, dried over anhydrous Na₂SO₄ and concentrated under vacuum, affording 800 mg (36%) of the title product as a colorless oil.

[0448] **Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-ethylbutanoate** To a solution of methyl 2-ethyl-2-hydroxybutanoate (371 mg, 2.54 mmol, 2.00 equiv) in DMF/THF (7/7 mL) was added NaH (100 mg, 4.17 mmol, 2.00 equiv) at 0 °C in 30 min. To this was added NaI (140 mg) and Int. **G-1** (500 mg,

1.28 mmol, 1.00 equiv). The resulting solution was stirred for 16 h at rt. The residue was applied onto a Prep-TLC with EtOAc / petroleum ether (1:10). The collected fractions were combined and concentrated under vacuum. This resulted in 210 mg (38%) of the title product as a white liquid.

Intermediate H-122: Methyl (2R)-2-([5-(3-cyclopropoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate



[0449] **Methyl (2R)-2-([5-(3-cyclopropoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate** To a solution of methyl (2R)-2-hydroxy-2-methylbutanoate (250 mg, 1.89 mmol, 2.00 equiv) in THF (20 mL) was added NaH (76 mg, 1.90 mmol, 2.00 equiv), in portions at 0 °C. The mixture was stirred at 0 °C for 20 min, then Int. F-10 (400 mg, 0.94 mmol, 1.00 equiv) at 0°C, Bu₄NI (200 mg, 0.54 mmol, 0.50 equiv) was added. The resulting solution was stirred for 16 h at rt, then quenched by the addition of water/ice. The resulting solution was extracted with 100 mL of ethyl acetate, and the combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with Prep-TLC using EtOAc / petroleum ether (1:1) to afford 160 mg (36%) of the title compound as a yellow solid. LC-MS: (ES, *m/z*): 475.

[0450] The following substituted pyrazole carboxylic esters were obtained from Williamson ether synthesis of the corresponding (bromomethyl)pyrazole with HO-C(CH₃)(Et)COOMe:





Table 17: Ether synthesis from (bromomethyl) pyrazoles.

Bromo cpd	Product		Spectral
--------------	---------	--	----------

G-1	Methyl (2 <i>S</i>)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-123	LC-MS: (ES, <i>m/z</i>): 443.
G-1	Methyl (2 <i>R</i>)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-124	LC-MS: (ES, <i>m/z</i>): 443.
G-4	Ethyl (2 <i>R</i>)-2-([1-(2-bromophenyl)-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-125	LC-MS: (ES, <i>m/z</i>): 525.
G-7	Methyl (2 <i>R</i>)-2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-126	LC-MS: (ES, <i>m/z</i>): 505.
G-8	Ethyl (2 <i>R</i>)-2-methyl-2-([1-(1-methyl-1 <i>H</i> -indazol-7-yl)-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy]butanoate	H-127	LC-MS: (ES, <i>m/z</i>): 505.15.
G-10	Ethyl (2 <i>R</i>)-2-([1-(2-chlorophenyl)-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-128	LC-MS: (ES, <i>m/z</i>): 469.1.
G-11	Methyl (2 <i>R</i>)-2-([1-(2-chlorophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-129	LC-MS: (ES, <i>m/z</i>): 459.
G-18	Methyl (2 <i>R</i>)-2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-130	LC-MS: (ES, <i>m/z</i>): 481.
G-18	Methyl (2 <i>S</i>)-2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-131	
G-19	Ethyl (2 <i>R</i>)-2-([5-(3,5-dimethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoate	H-132	LC-MS: (ES, <i>m/z</i>): 497.5.


Intermediate H-133: Methyl 2-([1-[2-(azetidin-1-yl)phenyl]-5-(1-ethyl-1*H*-indazol-6-yl)-1*H*-pyrazol-3-yl]methoxy)-2-methylpropanoate



[0451] Into a 20-mL sealed tube were combined methyl 2-([1-(2-bromophenyl)-5-(1-ethyl-1*H*-indazol-6-yl)-1*H*-pyrazol-3-yl]methoxy)-2-methylpropanoate (200 mg, 0.40 mmol, 1.00 equiv) L-proline (46 mg, 0.40 mmol, 1.00 equiv), CuI (38 mg, 0.20 mmol, 0.50 equiv), K₂CO₃ (166 mg, 1.20 mmol, 3.00 equiv), azetidine (69 mg, 1.21 mmol, 3.00 equiv), and DMSO (10 mL). The resulting solution was stirred overnight at 70 °C. The reaction mixture was cooled to room temperature. The resulting solution was extracted with 3x20 mL of

EtOAc and the organic layers combined and dried over Na₂SO₄ and concentrated under vacuum. The residue was purified by prep-TLC with EtOAc /petroleum ether (1/3). This resulted in 50 mg (26%) of the title compound as light yellow oil. LC-MS: (ES, *m/z*): 474. ¹H NMR: (300 MHz, CDCl₃) δ 7.90 (s, 1H), 7.62 (d, *J* = 8.4 Hz, 1H), 7.34 (s, 1H), 7.24 (m, 2H), 7.07 (dd, *J* = 7.8, 1.5 Hz, 1H), 6.76 (s, 1H), 6.69 (td, *J* = 7.6, 7.5, 1.3 Hz, 1H), 6.45 (dd, *J* = 8.2, 1.3 Hz, 1H), 4.63 (s, 2H), 4.25 (q, *J* = 7.3, 7.3, 7.3 Hz, 2H), 3.79 (s, 3H), 3.58 (dq, *J* = 24.9, 7.4, 7.3, 7.3 Hz, 4H), 2.10 (p, *J* = 7.3, 7.3, 7.3, 7.3 Hz, 2H), 1.56 (s, 6H), 1.34 (q, *J* = 8.2, 8.2, 7.2 Hz, 3H).

Intermediate H-134: Methyl 2-([1-[2-(azetidin-1-yl)phenyl]-5-[3-(2,2-dimethylpropoxy)-phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate



[0452] Into a 30-mL sealed tube were combined Int. **H-4** (200 mg, 0.39 mmol, 1.00 equiv), L-proline (44 mg, 0.38 mmol, 1.00 equiv), CuI (40 mg, 0.21 mmol, 0.50 equiv), K₂CO₃ (160 mg, 1.16 mmol, 3.00 equiv), azetidine (66 mg, 1.16 mmol, 3.00 equiv), and DMSO (10 mL). The resulting solution was stirred overnight at 70 °C, cooled, and extracted with 3x15 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified with by prep-TLC using EtOAc /petroleum ether (1/3) to afford 70 mg (37%) of the title compound as a light yellow oil. LC-MS: (ES, *m/z*): 492.30. ¹H NMR: (300 MHz, DMSO) δ 7.28 (ddd, *J* = 8.5, 7.4, 1.6 Hz, 1H), 7.20 (t, *J* = 7.9, 7.9 Hz, 1H), 6.98 (ddd, *J* = 9.9, 7.2, 1.5 Hz, 2H), 6.84 (m, 2H), 6.69 (m, 2H), 6.51 (dd, *J* = 8.3, 1.3 Hz, 1H), 4.44 (s, 2H), 3.71 (s, 3H), 3.50 (m, 2H), 3.42 (m, 2H), 2.07 (m, 2H), 1.45 (s, 6H), 0.94 (s, 9H).

[0453] The following amino substituted aryl compounds were obtained through use of similar copper(I) mediated coupling procedures.


Table 18: Arylamine synthesis from (bromoaryl) pyrazoles.

Aryl halide	Amine	Product		Spectral
-------------	-------	---------	--	----------

H-2	Pyrrolidine	Methyl 2-([5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-135	LC-MS: (ES, <i>m/z</i>): 488.15.
H-3	Pyrrolidine	Methyl 2-methyl-2-([5-(1-methyl-1 <i>H</i> -indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy]propanoate	H-136	LC-MS: (ES, <i>m/z</i>): 474.
H-24	Pyrrolidine	Methyl 2-([5-(3,5-dimethoxyphenyl)-1-[2-(pyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-137	LC-MS: (ES, <i>m/z</i>): 480.
H-24	Azetidine	Methyl 2-([1-[2-(azetidin-1-yl)phenyl]-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-138	LC-MS: (ES, <i>m/z</i>): 466.
H-25	Azetidine	Methyl 2-([1-[2-(azetidin-1-yl)phenyl]-5-(1-propyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-139	LC-MS: (ES, <i>m/z</i>): 488.1.
H-25	Pyrrolidine	Methyl 2-methyl-2-([5-(1-propyl-1 <i>H</i> -indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy]propanoate	H-140	LC-MS: (ES, <i>m/z</i>): 502.2.
H-26	Azetidine	Methyl 2-([1-[2-(azetidin-1-yl)phenyl]-5-(3-cyclobutoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-141	LC-MS: (ES, <i>m/z</i>): 476.5.
H-46	Pyrrolidine	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(pyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-142	LC-MS: (ES, <i>m/z</i>): 476.4.
H-46	Azetidine	methyl 2-([1-[2-(azetidin-1-yl)phenyl]-5-(3-cyclopropoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-143	LC-MS: (ES, <i>m/z</i>): 462.15.
H-46	(3 <i>S</i>)-3-hydroxy-pyrrolidine	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-[(3 <i>S</i>)-3-hydroxypyrrolidin-1-yl]phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-144	LC-MS: (ES, <i>m/z</i>): 492.
H-46	Morpholine	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(morpholin-4-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-145	LC-MS: (ES, <i>m/z</i>): 492.5.
H-46	2-Oxo-pyrrolidine	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(2-oxopyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-146	LC-MS: (ES, <i>m/z</i>): 490.1.
H-46	Pyrazole	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-(1 <i>H</i> -pyrazol-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-147	LC-MS: (ES, <i>m/z</i>): 473.25.
H-46	(3 <i>R</i>)-3-hydroxy-pyrrolidine	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[2-[(3 <i>R</i>)-3-hydroxypyrrolidin-1-yl]phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoate	H-148	LC-MS: (ES, <i>m/z</i>): 492.

H-47	Azetidine	Methyl 2-([1-[2-(azetidin-1-yl)-4-fluorophenyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-149	LC-MS: (ES, <i>m/z</i>): 480.40.
H-20	Pyrrolidine	Methyl 2-([5-(3-cyclopropoxyphenyl)-1-[[2-(pyrrolidin-1-yl)phenyl]methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-150	LC-MS: (ES, <i>m/z</i>): 490.15.
H-68	Pyrrolidine	Methyl 2-([5-(3-cyclobutoxyphenyl)-1-[[2-(pyrrolidin-1-yl)phenyl]methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate	H-151	LC-MS: (ES, <i>m/z</i>): 504.
H-97	Azetidine	Ethyl (2 <i>R</i>)-2-([1-[2-(azetidin-1-yl)phenyl]-5-(1-propyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate	H-152	LC-MS: (ES, <i>m/z</i>): 516.5.
H-102	Azetidine	Methyl (2 <i>R</i>)-2-([1-[2-(azetidin-1-yl)phenyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate	H-153	LC-MS: (ES, <i>m/z</i>): 476.2.
H-125	Pyrrolidine	Ethyl (2 <i>R</i>)-2-([5-(1-ethyl-1H-indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate	H-154	LC-MS: (ES, <i>m/z</i>): 516.
H-125	Azetidine	Ethyl (2 <i>R</i>)-2-([1-[2-(azetidin-1-yl)phenyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate	H-155	LC-MS: (ES, <i>m/z</i>): 502.

Intermediate H-156: Methyl 2-([1-(2-cyanophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate



[0454] To a solution of Int. **H-46** (150 g, 309.04 mmol, 1.00 equiv) and Zn(CN)₂ (72 mg, 0.62 mmol, 2.00 equiv) in DMF (15 mL) was added Pd(PPh₃)₄ (36 mg, 0.03 mmol, 0.10 equiv), in portions at rt. The resulting solution was stirred for 16 h at 100 °C, diluted with 100 mL of EtOAc, washed with 2x50 mL of (sat)FeSO₄(aq), dried over Na₂SO₄ and concentrated under vacuum. The residue was applied onto a Prep-TLC with EtOAc /petroleum ether (1:2). This resulted in 100 mg of the title compound as colorless oil. LC-MS: (ES, *m/z*): 432.

Chromatographic Procedures

[0455] All chromatography was performed on an 2#-AnalyseHPLC-SHIMADZU(HPLC-10) instrument, with detection at 254/220 nm. Columns used for purification include:


[0456] "Xselect": XSelect CSH Prep C18 OBD Column,, 19*250mm,5um.

[0457] "XBridge": XBridge Prep C18 OBD Column, 19*150mm 5um.

Examples

[0458] The invention is further illustrated by the following examples.

Example 1: 2-([1-[(3-Chlorophenyl)methyl]-5-phenyl-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid.













[0459] To a solution of Int. **H-1** (0.068 g, 0.170 mmol) in THF : MeOH : H₂O (2:1:1) (8 mL) was added LiOH * H₂O (0.035 g, 0.85 mmol) at RT and stirring was continued for further 3 h. The solvent was then evaporated and to the residue water (2mL) was added and acidified with 1M citric acid. The mixture was extracted with EtOAc (25 mL) and washed with water (10 mL) followed by brine (10 mL). The EtOAc layer was dried over Na₂SO₄, and solvent was removed by evaporation. The crude product was chromatographed over SiO₂ (ISCO CombiFlash® Rf 200) using 0-30% gradient of MeOH in DCM to afford the title product as white solid (0.032 g). ¹HNMR(CDCl₃): δ 1.56(s, 6H), 4.61 (s, 2H), 5.34 (s, 2H), 6.38(s, 1H), 6.87-6.97(m, 1H), 7.04(s, 1H), 7.18-7.24(m, 2H), 7.26-7.35(m, 2H), 7.35-7.45(m, 3H).






[0460] Examples 2-50 were prepared analogously as described in Example 1.






Table 19: Examples 2-50.






Ex.	Structure	IUPAC Name Analytical data
2		2-([5-(4-Chlorophenyl)-1-[(4-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.52(s, 6H), 4.54(s, 2H), 5.329s, 2H), 6.48(s, 1H), 6.91-7.00(m, 1H), 7.21-7.30-(m, 2H), 7.31-7.38(m, 2H), 7.40-7.48 (m 3H).






3		2-([5-(4-Chlorophenyl)-1-[(2-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.50(s, 6H), 4.58(s, 2H), 5.41(s, 2H), 6.56(s, 1H), 6.72-6.80 (m, 1H), 7.20-7.34(m, 2H), 7.34-50 (m, 5H).
4		2-([1-[(2-Chlorophenyl)methyl]-5-(4-fluorophenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.51(s, 6H), 4.58(s, 2H), 5.40(s, 2H), 6.55(s, 1H), 6.70-6.78(m, 1H), 7.14-7.18(m, 2H), 7.20-7.29(m, 2H), 7.32-7.40(m, 3H).
5		2-([5-(4-Chlorophenyl)-1-[(2,4-dichlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.48(s, 6H), 4.58(s, 2H), 5.40 (s, 2H), 6.54(s, 1H), 6.73(d, 1H), 7.20-7.38 (m, 3H), 7.40-48 (m, 3H).
6		2-([5-(4-Chlorophenyl)-1-[(2,4-dichlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)acetic acid ¹ H NMR (DMSO-d ₆ , 400 MHz) δ 3.80 (s, 2H), 4.50 (s, 2H), 5.30 (s, 2H), 6.50 (s, 1H), 6.60-80 (m, 1H), 7.20-70 (m, 6H).
7		2-([1-Benzyl-5-(4-Chlorophenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.53(s, 6H), 4.58(s, 2H), 5.38 (s, 2H), 6.51(s, 1H), 6.90(d, 2H), 7.18-7.50 (m, 7H).

8		2-([1-[(2-Chlorophenyl)methyl]-5-phenyl-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (DMSO-d ₆ , 400 MHz) δ 1.56 (s, 6H), 4.58(s, 2H), 5.41(s, 2H), 6.56(s, 1H), 6.75-6.78(m, 1H), 7.20-7.30 (m, 4H), 7.32-7.43(m, 4H). ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.54(s, 6H), 4.589s, 2H), 5.40(s, 2H), 6.55(s, 1H), 6.68-6.74(m, 1H), 7.20-7.42(m, 8H).
9		2-([5-(2-Chlorophenyl)-1-[(2-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.53(s, 6H), 4.58(s, 2H), 5.28(s, 2H), 6.43(s, 1H), 6.78-7.81(m, 1H), 7.16-7.34(m, 5H), 7.38-7.42 (m, 1H), 7.50-7.56(m, 1H).
10		2-([5-(3-Chlorophenyl)-1-[(2-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.32 (s, 6H), 4.58(s, 2H), 5.42(s, 2H), 6.57(s, 1H), 6.67-6.80 (m, 1H), 7.22-7.30 (m, 3H), 7.32-7.42(m, 4H).
11		2-([1-[(2-Chlorophenyl)methyl]-5-(m-tolyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.58 (s, 6H), 2.35 (s, 3H), 4.61 (s, 2H), 5.42 (s, 2H), 6.39 (s, 1H), 6.75-6.84 (m, 1H), 7.00-7.12 (m, 2H), 7.18-7.29 (4H), 7.29-7.38 (m, 1H).
12		2-([1-[(2-Chlorophenyl)methyl]-5-(3-fluorophenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.58(s, 6H), 4.64 (s, 2H), 5.48 (s, 2H), 6.42 (s, 1H), 6.72-6.85 (m, 1H), 6.88-7.14 (m, 3H), 7.14-7.40 (m, 4H).

13		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(trifluoromethyl)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.58 (s, 6H), 4.62 (s, 2H), 5.41 (s, 2H), 6.46 (s, 1H), 6.80-6.88 (m, 1H), 7.17-7.24 (m, 2H), 7.29-7.38 (m, 1H), 7.39-7.47 (m, 1H), 7.49-7.54 (m, 2H), 7.59-7.68 (m, 1H).
14		2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.58 (s, 6H), 3.68 (s, 3H), 4.62 (s, 2H), 5.46 (s, 2H), 6.42 (s, 1H), 6.74-6.81 (m, 2H), 6.82-6.88 (m, 1H), 6.89-6.94 (m, 1H), 7.15-7.23 (m, 2H), 7.26-7.37 (m, 2H).
15		2-([1-[(2-Chlorophenyl)methyl]-5-cyclohexyl-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.16-1.42 (m, 4H), 1.54 (s, 6H), 1.57-1.81 (m, 5H), 2.36-2.51 (m, 1H), 2.76-2.91 (m, 1H), 4.56 (s, 2H), 5.39 (s, 2H), 6.15 (s, 1H), 6.54-6.68 (m, 1H), 7.08-7.25 (m, 2H), 7.31-7.44 (m, 1H).
16		2-([1-[(2-Chlorophenyl)methyl]-5-cyclopentyl-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.22-1.64 (m, 10H), 1.64-1.80 (m, 2H), 1.82-2.00 (m, 2H), 2.80-2.92 (m, 1H), 4.74 (s, 2H), 5.40 (s, 2H), 6.10 (s, 1H), 6.58-6.62 (m, 1H), 7.10-7.22 (m, 2H), 7.38 (d, 1H).
17		1-[[5-(4-Chlorophenyl)-1-[(2-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy]cyclopropanecarboxylic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.18-1.32 (m, 3H), 1.64-1.80 (m, 1H), 4.61 (s, 2H), 5.42 (s, 2H), 6.52 (s, 1H), 6.60-6.80 (m, 1H), 7.20-7.52 (m, 7H).






18		2-([1-[(2-Chlorophenyl)methyl]-5-thiazol-2-yl-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CDCl ₃ , 400 MHz) δ 1.38 (m, 6H), 4.48 (s, 2H), 5.61 (s, 2H), 6.55-6.65 (m, 1H), 7.03 (s, 1H), 7.09-7.16 (m, 1H), 7.17-7.23 (m, 1H), 7.27-7.33 (d, 1H), 7.34-7.42 (m, 1H), 7.88-7.96 (d, 1H).
19		2-([5-(4-Chlorophenyl)-1-[(2-fluorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD, 400 MHz) δ 1.48 (s, 6H), 4.53 (s, 2H), 5.38 (s, 2H), 6.50 (s, 1H), 6.81-6.92 (m, 1H), 7.00-7.13 (m, 2H), 7.21-7.31 (m, 1H), 7.34-7.48 (m, 4H)
20		2-([5-(4-Chlorophenyl)-1-[(2-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanamide ¹ H NMR (CD ₃ OD 400 MHz) δ 1.46 (s, 6H), 4.58 (s, 2H), 5.40 (s, 2H), 6.52 (s, 1H), 6.71-6.76 (m, 1H), 7.20-7.58 (m, 6H)
21		2-([5-(4-chlorophenyl)-1-([pyridin-4-yl]methyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (CD ₃ OD 400 MHz) δ 1.52 (s, 6H), 4.58 (s, 2H), 5.56 (s, 2H), 6.59 (s, 1H), 7.21-7.52 (m, 6H), 8.50 (bs, 2H)
22		2-([5-(4-Chlorophenyl)-1-([pyridin-2-yl]methyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H NMR (DMSO-d ₆ , 400 MHz) δ 1.38 (s, 6H), 4.40 (s, 2H), 5.38 (s, 2H), 6.41 (s, 1H), 6.90-7.10 (m, 1H), 7.20-7.85 (m, 6H), 8.51 (bs, 1H)

23		2-([5-(4-Chlorophenyl)-1-([pyridin-3-yl]methyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ^1H NMR (CD_3OD 400 MHz) δ 1.52(s, 6H), 4.58(s, 2H), 5.10(s, 2H), 6.55(s, 1H), 7.30-7.41(m, 3H), 7.42-7.51(m, 3H), 8.19(s, 1H), 8.41(d, 1H).
24		2-([1-[(2-Chlorophenyl)methyl]-5-(thien-3-yl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ^1H NMR(CDCl_3): δ 1.37(s, 6H), 4.38(s, 2H), 5.43(s, 2H), 6.48(s, 1H), 6.62-6.64(m, 1H), 7.17-7.31(m, 3H), 7.44-7.46(m, 1H), 7.57-7.64(m, 2H).
25		2-([1-[(2-Chlorophenyl)methyl]-5-(thien-2-yl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ^1H NMR (CDCl_3): δ 1.37(s, 6H), 4.39(s, 2H), 5.47(s, 2H), 6.53(s, 1H), 6.59-6.61(m, 1H), 7.09-7.14(m, 2H), 7.24-7.31(m, 2H), 7.45-7.47(m, 1H), 7.62-7.64(m, 1H).
26		2-([1-[(2-Fluorophenyl)methyl]-5-(3-(2-methylpropoxy)phenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ^1H NMR(CDCl_3): δ 0.97 (s, 3H), 0.99 (s, 3H), 1.55 (s, 6H), 2.00-2.06 (m, 1H), 3.70(d, 2H), 4.62 (s, 2H), 5.39 (s, 2H), 6.37 (s, 1H), 6.79-6.80 (m, 1H), 6.87-7.09 (M, 5H), 7.22-7.30(m, 2H)
27		2-([1-[(2-Chlorophenyl)methyl]-5-(3-(2-methylpropoxy)phenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ^1H NMR (CDCl_3): δ 0.95(s, 3H), 0.96(s, 3H), 1.56(s, 6H), 1.97-2.03(m, 1H), 3.49(d, 2H), 4.63 (s, 2H), 5.43(s, 2H), 6.41(s, 1H), 6.73-6.80 (m, 2H), 6.84-6.92(m, 2H), 7.19-7.28(m, 3H), 7.33-7.35(m, 1H).

28		2-([1-[(2-Chlorophenyl)methyl]-5-(3-ethoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.34 (t, 3H), 1.56(s, 6H), 3.86(q, 2H), 4.62(s, 2H), 5.43 (s, 2H), 6.40(s, 1H), 6.76-6.78 (m, 2H), 6.83-6.85(m, 1H), 6.89-6.91(m, 1H), 7.19-7.28 (m, 3H), 7.33-7.35(m, 1H).
29		2-([1-[(2-Bromophenyl)methyl]-5-(3-isopropoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.21(s, 3H), 1.23(s, 3H), 4.28-4.32(m, 1H), 4.62(s, 2H), 5.39(s, 2H), 6.42(s, 1H), 6.71-6.74(m, 2H), 6.82-6.89(m, 2H), 7.11-7.15(m, 1H), 7.22-7.27(m, 2H), 7.51-7.54(m, 1H).
30		2-([1-[(2-Bromophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ HNMR (CDCl ₃): δ 1.56(s, 6H), 3.65(s, 3H), 4.62(s, 2H), 5.39(s, 2H), 6.41(s, 1H), 6.73-6.76(m, 2H), 6.84-6.92(m, 2H), 7.15(t, 1H), 7.13-7.29(m, 2H), 7.52-7.54(m, 1H).
31		2-([1-[(2-Isopropylphenyl)methyl]-5-(3-methoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.13 (s, 3H), 1.15(s, 3H), 1.55 (s, 6H), 2.94-2.97(m, 1H), 3.62(s, 3H), 4.63(s, 2H), 5.42(s, 2H), 6.39(s, 1H), 6.65-6.67(m, 1H), 6.75-6.76(m, 1H), 6.85-6.91(m, 2H), 7.07-7.11(m, 1H), 7.21-7.28(m, 3H)
32		2-([5-(3-Methoxyphenyl)-1-(o-tolylmethyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.55 (s, 6H), 2.18 (s, 3H), 3.64 (s, 3H), 4.62 (s, 2H), 5.31(s, 2H), 6.38(s, 1H), 6.69-6.81(m, 1H), 6.77-6.78(m, 1H), 6.86-6.92 (m, 2H), 7.10-7.15 (m, 3H), 7.25-7.29 (m, 1H).
33		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(cyclopropylmethoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 0.25-0.29(m, 2H), 0.58-0.63(m, 2H), 1.16-1.20(m, 1H), 1.55(s, 6H), 5.58(d, 2H), 4.63(s, 2H), 5.42(s, 2H), 6.40(s, 1H),

		6.374-6.78(m, 2H), 6.84-6.93(m, 2H), 7.18-7.28(m, 3H), 7.33-7.35(m, 1H).
34		2-([1-(2-Chlorophenyl)methyl]-5-[3-(trifluoromethoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.56(6H), 4.63(s, 2H), 5.41(s, 2H), 6.46(s, 1H), 6.77-6.79(m, 1H), 7.12-7.25(m, 5H), 7.32-7.35(m, 1H), 7.38-7.42 (m, 1H).
35		2-([1-(2-Chlorophenyl)methyl]-5-(3-isopropoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.22(s, 3H), 1.23(s, 3H), 1.55(s, 6H), 4.30-4.33(m, 1H), 4.62(s, 2H), 5.43(s, 2H), 6.41(s, 1H), 6.75-6.90(4H), 7.19-7.35(m, 4H).
36		2-([5-(3-Methoxyphenyl)-1-(2-methoxyphenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.55(s, 6H), 3.70(s, 3H), 3.73(s, 3H), 4.62(s, 2H), 5.34(s, 2H), 6.35(s, 1H), 6.77-6.392(m, 6H), 7.22-7.29(m, 2H).
37		2-([1-(2-Chlorophenyl)methyl]-5-[3-(cyclobutoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.49-1.61 (m, 7H), 1.75-1.82(m 1H), 2.01-2.11(m 2H), 2.17-2.24(m 2H), 4.37-4.45(m 1H), 4.62(s, 2H), 5.42(s, 2H), 6.39(s, 1H), 6.64(s, 1H), 6.75-6.86 (m, 3H), 7.19-7.35(m 4H).

38		2-([1-[[2-(Dimethylamino)phenyl]methyl]-5-(3-methoxyphenyl)]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.56(s, 6H), 2.60(s, 6H), 6.64(s, 3H), 4.63(s, 2H), 5.44(s, 2H), 6.30(s, 1H), 6.71-7.11(m, 6H), 7.20-7.25(m, 2H).
39		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(cyclopropoxy)phenyl]]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 0.55-0.68(m, 4H), 1.55(s, 6H), 3.50-3.54(m, 1H), 4.63(s, 2H), 5.44(s, 2H), 6.42(s, 1H), 6.76-7.02(m, 4H), 7.17-7.35(m, 4H).
40		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(morpholinomethyl)phenyl]]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CD ₃ OD): δ 1.50(s, 6H), 3.15-3.23(bs, 4H), 3.60-3.81(bs, 2H), 3.85-4.16(bs, 2H), 4.35(s, 2H), 4.57(s, 2H), 5.45(s, 2H), 6.60(s, 1H), 6.75(d, 1H), 7.20-7.55(m, 7H).
41		1-[[1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)]-1H-pyrazol-3-yl]methoxy]cyclobutanecarboxylic acid. ¹ HNMR (CDCl ₃): δ 1.78-1.85(m, 1H), 1.95-2.04(m, 2H), 2.18-2.25(m, 1H), 3.09-3.14(m, 1H), 3.66(s, 3H), 4.27-4.32(m, 1H), 4.55-4.65(dd, 2H), 5.43(s, 2H), 6.44(s, 1H), 6.73-6.91(m, 4H), 7.17-7.34(m, 4).
42		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(cyclobutylmethoxy)phenyl]]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.55(s, 6H), 1.75-1.82(m, 2H), 1.85-1.98(m, 2H), 2.06-2.14(m, 2H), 2.64-2.70(m, 1H), 3.71(d, 2H), 4.63(s, 2H), 5.43(s, 2H), 6.42(s, 1H), 6.75-6.92(m, 4H), 7.20-7.36(m, 4).






43		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(2,2,2-trifluoroethoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.55(s, 6H), 4.16(q, 2H), 4.63(s, 2H), 5.42(s, 2H), 6.43(s, 1H), 6.77-6.81(m, 2H), 6.96-6.98(m, 2H), 7.19-7.36(m, 4H).
44		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(cyclopentylmethoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.25-1.29(m, 2H), 1.55-1.61(m, 10H), 1.76-1.79(m, 2H), 2.24-2.31(m, 1H), 3.59(d, 2H), 4.62(s, 2H), 5.43(s, 2H), 6.41(s, 1H), 6.73-6.91(m, 4H), 7.19-7.35(m, 4H).
45		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(tetrahydropyran-4-ylmethoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.34-1.41(m, 2H), 1.55(s, 6H), 1.66-1.69(m, 2H), 1.97-1.99(m, 1H), 3.38-3.46m, 2H), 3.56(d, 2H), 3.98-4.01(m, 2H), 4.62(s, 2H), 5.42(s, 2H), 6.42(s, 1H), 6.71(s, 1H), 6.77-6.95(m, 3H), 7.19-7.35(m, 4H).
46		2-([1-[(2-Chlorophenyl)methyl]-5-(3-morpholinophenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ HNMR (CDCl ₃): δ 1.55(s, 6H), 2.96(t, 4H), 3.76(t, 4H), 4.62(s, 2H), 5.41(s, 2H), 6.41(s, 1H), 6.68(s, 1H), 6.78-6.80(m, 2H), 6.88-6.91(m, 1H), 7.19-7.35(m, 4H).
47		2-([1-[(2-Chlorophenyl)methyl]-5-(1,3-thiazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropionic acid ¹ HNMR(CDCl ₃): δ 1.56(s, 6H), 4.63(s, 2H), 5.84(s, 2H), 6.62-6.67(m, 2H), 7.10-7.19(m 2H), 7.33-7.35(m, 1H), 7.39(d, 1H), 8.83(d, 1H).

48		2-([1-[(2-Chlorophenyl)methyl]-5-(3-cyano-phenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropionic acid ¹ HNMR(CDCl ₃): δ 1.56(s, 6H), 4.62(s, 2H), 5.39(s, 2H), 6.46(s, 1H), 6.82-6.84(m, 1H), 7.18-7.24(m, 2H), 7.33-7.35(m 1H), 7.46-7.55(m, 3H), 7.65-7.68(m, 1H).
49		2-([1-[(2-Chlorophenyl)methyl]-5-(1,3-thiazol-5-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropionic acid ¹ HNMR(CDCl ₃): δ 1.56(s, 6H), 4.62(s, 2H), 5.50(s, 2H), 6.60-6.6(m, 2H), 7.14-7.24(m, 2H), 7.36-7.38(m, 1H), 7.76(s, 1H), 8.84(s, 1H).
50		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(dimethyl-amino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropionic acid MS: Calculated: 427.92; Found: 428.9 [M+H].

[0461] Examples 51-74 were prepared using the methods disclosed in the indicated Schemes.

Table 20: Examples 51-74.

Ex.	Scheme	Structure	IUPAC Name Analytical data
51	II		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(propan-2-yl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ H-NMR (300 MHz, MeOD): δ 1.12 (6 H, d), 1.53 (6 H, s), 2.83 (1 H, p), 4.60 (2 H, s), 5.40 (2 H, s), 6.54 (1 H, s), 6.72 – 6.81 (1 H, m), 7.05 – 7.45 (7 H, m).

52	II		2-([1-[(2-chlorophenyl)methyl]-5-[3-(methoxymethyl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD): δ 1.52 (6 H, s), 4.42 (2 H, s), 4.59 (2 H, s), 5.42 (2 H, s), 6.56 (1 H, s), 6.70 – 6.80 (1 H, m), 7.17 – 7.33 (4 H, m), 7.32 – 7.45 (3 H, m).
53	II		2-([1-benzyl-5-(3-methoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD) δ 1.51 (6 H, s), 3.68 (3 H, s), 4.57 (2 H, s), 5.35 (2 H, s), 6.48 (1 H, s), 6.84 (1 H, dd), 6.89 – 7.06 (4 H, m), 7.17 – 7.38 (4 H, m).
54	II		2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-4-methyl-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR:(300 MHz, MeOD) δ 1.53 (6 H, s), 2.09 (3 H, s), 3.67 (3 H, s), 4.60 (2 H, s), 5.31 (2 H, s), 6.67 – 6.86 (3 H, m), 6.97 (1 H, ddd), 7.17 – 7.40 (4 H, m).
55	II		2-([5-(3-methoxyphenyl)-1-phenyl-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD) δ 1.53 (6 H, s), 3.64 (3 H, s), 4.61 (2 H, s), 6.63 – 6.77 (2 H, m), 6.77 – 6.93 (2 H, m), 7.16 – 7.36 (3 H, m), 7.40 (3 H, dddd).
56	II		2-([5-(3-methoxyphenyl)-1-(pyridin-2-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 367 ¹ H-NMR (400 MHz, MeOD) δ 1.54 (6 H, s), 3.69 (3 H, s), 4.65 (2 H, s), 6.70 (1 H, s), 6.76 – 6.78 (2 H, m), 6.89(1 H, ddd), 7.20 – 7.25 (1H, d), 7.41-7.45 (2H, m), 7.91 (1 H, dt), 8.41 (1 H, t).


57	II		2-([1-(2-chlorophenyl)-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 401 ¹ H-NMR (400 MHz, MeOD) δ 1.55 (6 H, s), 3.64 (3 H, s), 4.63 (2 H, s), 6.73 (2 H, d), 6.86 (2 H, dd), 7.20 (1 H, dd), 7.42 – 7.60 (4 H, m).
58	II		2-([1-(3-chlorophenyl)-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 1.54 (6 H, s), 3.71 (3 H, s), 4.62 (2 H, s), 6.68 (1 H, s), 6.77 – 6.87 (2 H, m), 6.94 (1 H, ddd), 7.19 (1 H, dt), 7.27 (1 H, t), 7.32 – 7.44 (3 H, m).
59	II		2-([1-(4-chlorophenyl)-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 401 ¹ H-NMR (400 MHz, MeOD) δ 1.54 (6 H, s), 3.71 (3 H, s), 4.62 (2 H, s), 6.67 (1 H, s), 6.75 – 6.85 (2 H, m), 6.92 (1 H, ddd), 7.21 – 7.32 (3 H, m), 7.37 – 7.46 (2 H, m).
60	VII		2-([1-Benzyl-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 0.97 (6 H, d), 1.53 (6 H, s), 1.97 (1 H, dp), 3.48 (2 H, d), 3.78 (3 H, s), 4.58 (2 H, s), 5.31 (2 H, s), 6.52 (1 H, s), 6.67 – 6.74 (1 H, m), 6.76 – 6.82 (1 H, m), 6.85 – 7.00 (4 H, m), 7.22 – 7.33 (2 H, m)
61	VII		2-([1-[(2-methoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 453 ¹ H-NMR (400 MHz, MeOD) δ 0.97 (6 H, d), 1.53 (6 H, s), 1.97 (1 H, dp), 3.48 (2 H, d), 3.78 (3 H, s), 4.58 (2 H, s), 5.31 (2 H, s), 6.52 (1 H, s), 6.67 – 6.74 (1 H, m), 6.76 – 6.82 (1 H, m), 6.85 – 7.00 (4 H, m), 7.22 – 7.33 (2 H, m).

62	VII		2-([1-[(3-methoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 453 ¹ H-NMR (400 MHz, MeOD) δ 0.99 (6 H, d), 1.53 (6 H, s), 1.99 (1 H, dq), 3.55 (2 H, d), 3.73 (3 H, s), 4.59 (2 H, s), 5.33 (2 H, s), 6.50 (1 H, s), 6.56 – 6.65 (2 H, m), 6.78 – 6.86 (2 H, m), 6.92 – 7.00 (2 H, m), 7.22 (1 H, t), 7.33 (1 H, t).
63	VII		2-([1-[(4-methoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 453 ¹ H-NMR (400 MHz, MeOD) δ 1.00 (6 H, d), 1.53 (6 H, s), 2.01 (1 H, dp), 3.57 (2 H, d), 3.77 (3 H, s), 4.58 (2 H, s), 5.29 (2 H, s), 6.47 (1 H, s), 6.78 – 6.90 (3 H, m), 6.92 – 7.01 (4 H, m), 7.29 – 7.39 (1 H, m).
64	VIII		2-([1-[(2-chlorophenyl)methyl]-5-(5-methoxythien-2-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD): δ 1.51 (6 H, s), 3.86 (3 H, s), 4.54 (2 H, s), 5.50 (2 H, s), 6.19 (1 H, d), 6.51 (1 H, s), 6.55 – 6.66 (2 H, m), 7.25 (2 H, dtd), 7.43 (1 H, dd).
65	VIII		2-([1-[(2-chlorophenyl)methyl]-5-(4-methoxythien-2-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD): δ 1.51 (6 H, s), 3.76 (3 H, s), 4.56 (2 H, s), 5.54 (2 H, s), 6.51 (1 H, d), 6.56 – 6.69 (3 H, m), 7.26 (2 H, dtd), 7.44 (1 H, dd).
66	VIII		2-([1-[(2-chlorophenyl)methyl]-5-(5-methoxythien-3-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 1.53 (6 H, s), 3.85 (3 H, s), 4.57 (2 H, s), 5.52 (2 H, s), 6.26 (1 H, d), 6.52 – 6.59 (2 H, m), 6.64 – 6.71 (1 H, m), 7.28 (2 H, dtd), 7.45 (1 H, dd).

67	VII		2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-ethylbutyric acid ¹ H-NMR (400 MHz, MeOD): δ (0.91 (6 H, t), 1.87 (4 H, q), 3.68 (3 H, s), 4.52 (2 H, s), 5.43 (2 H, s), 6.67 (1 H, s), 6.72 – 6.86 (2 H, m), 6.88 – 7.00 (2 H, m), 7.21 – 7.36 (3 H, m), 7.37 – 7.48 (1 H, m).
68	VII		2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid ¹ H-NMR (400 MHz, MeOD) δ 0.97 (3 H, t), 1.51 (3 H, s), 1.89 (2 H, qd), 3.68 (3 H, s), 4.59 (2 H, s), 5.44 (2 H, s), 6.56 (1 H, s), 6.71 – 6.79 (1 H, m), 6.83 (1 H, dd), 6.89 – 7.01 (2 H, m), 7.22 – 7.36 (3 H, m), 7.37 – 7.45 (1 H, m).
69	XVIII		2-([1-[(2-Chlorophenyl)methyl]-5-[3-(2-methylpropanamido)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, DMSO) : δ 1.09 (6 H, d), 1.39 (6 H, s), 2.57 (1 H, p), 5.39 (2 H, s), 6.40 (1 H, s), 6.69 – 6.79 (1 H, m), 7.00 (1 H, dt), 7.21 – 7.48 (4 H, m), 7.53 – 7.63 (1 H, m), 7.78 (1 H, t), 9.93 (1 H, s).
70	XIX		2-([1-[(2-Chlorophenyl)methyl]-5-(3-methanesulfonylphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) : δ 1.54 (6 H, s), 3.10 (3 H, s), 4.61 (2 H, s), 5.48 (2 H, s), 6.67 (1 H, s), 6.82 (1 H, dd), 7.21 – 7.33 (2 H, m), 7.35 – 7.42 (1 H, m), 7.64 – 7.74 (2 H, m), 7.90 (1 H, q), 7.96 – 8.03 (1 H, m).
71	XIX		2-([1-[(2-Chlorophenyl)methyl]-5-(3-methanesulfonamidophenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD): δ 1.52 (6 H, s), 2.04 (1 H, s), 2.87 (3 H, s), 4.58 (2 H, s), 5.44 (2 H, s), 6.56 (1 H, s), 6.66 – 6.76 (1 H, m), 7.12 (1 H, dt), 7.25 (4 H, dtd), 7.32 – 7.44 (2 H, m).

72	XIX		2-([1-[(2-chlorophenyl)methyl]-5-[3-(methylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD) : δ 1.52 (6 H, s), 2.04 (1 H, s), 2.87 (3 H, s), 4.58 (2 H, s), 5.44 (2 H, s), 6.56 (1 H, s), 6.66 – 6.76 (1 H, m), 7.12 (1 H, dt), 7.25 (4 H, dtd), 7.32 – 7.44 (2 H, m).
73	XIX		2-([1-[(2-chlorophenyl)methyl]-5-[3-[(2-methylpropyl)amino]phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD) : δ 0.88 (6 H, d), 1.52 (6 H, s), 1.74 (1 H, dp), 2.66 (2 H, d), 4.57 (2 H, s), 5.42 (2 H, s), 6.40 – 6.77 (5 H, m), 7.11 (1 H, dd), 7.19 – 7.35 (2 H, m), 7.35 – 7.47 (1 H, m).
74	XII		2-([1-[(2-Chlorophenyl)methyl]-5-[3-[(propan-2-yl)carbamoyl]phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR: (400 MHz, DMSO- <i>d</i> ₆) δ 1.16 (6 H, d), 1.41 (6 H, s), 3.33 (5 H, s), 4.02 – 4.17 (1 H, m), 4.44 (2 H, s), 5.40 (2 H, s), 6.53 (1 H, s), 6.75 – 6.88 (1 H, m), 7.23 – 7.36 (2 H, m), 7.41 – 7.47 (1 H, m), 7.49 – 7.57 (2 H, m), 7.81 – 7.98 (2 H, m), 8.29 (1 H, d), 12.63 (1 H, s).

Example 75: 2-([1-[(2-chlorophenyl)methyl]-5-(phenylamino)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid.



[0462] **2-[(5-[[*tert*-Butoxy]carbonyl](phenyl)amino)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid** was prepared from Int. **H-93** using the procedure of Example 1. This resulted in 200 mg as yellow oil.

[0463] **2-([1-[(2-Chlorophenyl)methyl]-5-(phenylamino)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid** To a solution of the product from the previous step (200 g,

400.01 mmol, 1.00 equiv) in DCM (10 mL) was added TFA (5 mL) dropwise with stirring at rt. The resulting solution was stirred for 1 h, then concentrated under vacuum. The crude product (150 mg) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, SunFire C18 OBD Prep Column, 0.1 mM, 5 μ M, 19 mm x 150 mm, mobile phase, Waters (0.1% TFA) and ACN (61.0% ACN up to 74.0% in 6 min); Detector, UV 254 nm), to afford 19.2 mg of the title product as a white solid.

[0464] $^1\text{H-NMR}$ (300 MHz, MeOD) δ 1.50 (6 H, s), 4.48 (2 H, s), 5.34 (2 H, d), 6.18 (1 H, s), 6.68 – 6.95 (4 H, m), 7.11 – 7.30 (4 H, m), 7.33 – 7.43 (1 H, m).


[0465] The following Examples were prepared using the indicated Schemes.

Table 21: Examples 76-77.

Ex.	Scheme	Structure	IUPAC Name Analytical data
76	XVII		2-([1-[(2-Chlorophenyl)methyl]-5-phenoxy-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid $^1\text{H-NMR}$ (CD ₃ OD, ppm): δ (300 MHz, MeOD) 1.46 (6 H, s), 4.44 (2 H, s), 5.35 (2 H, s), 5.78 (1 H, s), 6.90 – 7.00 (1 H, m), 7.06 – 7.46 (8 H, m).
77	VI		2-([5-Benzyl-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid $^1\text{H-NMR}$ (300 MHz, MeOD): δ 1.48 (6 H, s), 3.92 (2 H, s), 4.50 (2 H, s), 5.32 (2 H, s), 6.23 (1 H, s), 6.42 – 6.52 (1 H, m), 7.07 – 7.30 (7 H, m), 7.39 (1 H, dd).


[0466] Example 78 is intentionally left blank.

Example 79: 2-([1-[(2-Chlorophenyl)methyl]-5-(3-hydroxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid.



[0467] (i) 2-([1-[(2-Chlorophenyl)methyl]-5-(3-hydroxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid: A solution of methyl 2-([5-(3-benzyloxyphenyl)-1-[(2-chlorophenyl)methyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoate in mixture of acetic acid and conc HCl (8 mL) (3:1, v/v) was heated at 90 °C for 6 h. At the end of this period the reaction mixture was evaporated to dryness and the residue was chromatographed over SiO₂ using 0-20% gradient of MeOH in DCM to afford the title product. ¹HNMR (CDCl₃): δ 1.55 (s, 6H), 4.61 (s, 2H), 5.40(s, 2H), 6.38 (s, 1H), 6.74-6.83(m, 4H), 7.18-7.35(m, 4H).

Example 80: 2-([1-[(2-Chlorophenyl)methyl]-5-[3-(oxetan-3-ylmethoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid.




[0468] (i) Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-hydroxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoate: To a solution of 2-([1-[(2-chlorophenyl)methyl]-5-(3-hydroxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid (Example 79) in MeOH was added 4 drops of conc H₂SO₄, and the solution was refluxed for 16 h. The product was evaporated to dryness and the residue was chromatographed over SiO₂ using 0-25% gradient of MeOH in DCM to afford the title product.

[0469] (ii) Methyl 2-([1-[(2-chlorophenyl)methyl]-5-[3-(oxetan-3-ylmethoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate. To a solution of methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-hydroxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate (0.170 g, 0.410 mmol) in DMF was added [oxetan-3-yl]methyl 4-methylbenzenesulfonate (0.149 g, 0.615 mmol) and K₂CO₃ (0.113 g, 0.820 mmol) at room temperature. The mixture

was heated at 60 °C for 16 h. The reaction mixture was cooled to room temperature, water was added, and the mixture was extracted with EtOAc (20 mLx2). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over Na₂SO₄, and concentrated under reduced pressure. The residue was chromatographed over SiO₂ using 0-100% gradient of EtOAc in hexane to afford the title product. ¹HNMR (CDCl₃): δ 1.54 (s, 6H), 3.33-3.36(m, 1H), 3.77(s, 3H), 3.97(d, 2H), 4.30 (t, 2H), 4.56(s, 2H), 4.84(t, 2H), 5.40(s, 2H), 6.53(s, 1H), 6.77-6.90(m, 4H), 7.19-7.37(m, 4H).


[0470] (iii) 2-([1-[(2-Chlorophenyl)methyl]-5-[3-(oxetan-3-ylmethoxy)phenyl]1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid was prepared analogously as described in Step iv of Example 1, yielding the title product (20 mg). ¹HNMR (CDCl₃): δ 1.55(s, 6H), 3.34-3.74 (m, 1H), 3.98 (d, 2H), 4.49 (t, 2H), 4.63(s, 2H), 4.83(t, 2H), 5.43(s, 2H), 6.44(s, 1H), 6.76-6.93(m, 4H), 7.20-7.37(m, 4H).

Example 81: 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methyl-N-methylsulfonyl-propanamide.



[0471] To a solution of Example 14 (2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid) (0.08 g, 0.192 mmol) in CH₃CN was added CDI (0.047 g, 0.288 mmol) and methanesulfonamide (0.022g, 0.23 mmol) at rt. The mixture was stirred at rt for 16 h. The solvent was evaporated and the residue was chromatographed over SiO₂ using 0-20% gradient of MeOH in DCM to afford the title product. ¹HNMR (CDCl₃): δ 1.55(s, 6H), 3.21(s, 3H), 3.67(s, 3H), 4.58(s, 2H), 5.48(s, 2H), 6.36(s, 1H), 6.75-6.92(m, 4H), 7.18-7.35(m, 4H).

Example 82: Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)1H-pyrazol-3-yl]methoxy)-2-methyl-propanoate



[0472] Example 82 was prepared analogously as described in Example 1, omitting Step iv. ¹HNMR(CDCl₃): δ 1.54(s, 6H), 3.65(s, 3H), 3.77(s, 3H), 4.56(s, 2H), 5.37(s, 2H), 6.53(s, 1H), 6.71-6.78(m, 2H), 6.85-6.90(m, 2H), 7.10-7.13(m, 1H), 7.21-7.27(m, 2H), 7.51-7.53(m, 1H).

[0473] Examples 83-100 are intentionally left blank.

[0474] The following Examples were prepared using the indicated Schemes.

Table 22. Examples 101-119.

Ex.	Scheme	Structure	IUPAC Name Analytical Data
101	II		2-([1-(2-Chlorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 443. ¹ H-NMR (400 MHz, DMSO) δ 0.90 (6 H, d), 1.25 (1 H, d), 1.43 (6 H, s), 1.88 (1 H, hept), 3.54 (2 H, d), 4.48 (2 H, s), 6.65 – 6.73 (2 H, m), 6.75 – 6.88 (2 H, m), 7.20 (1 H, t), 7.45 – 7.65 (4 H, m).
102	II		2-([1-(2-Methoxyphenyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 439 ¹ H-NMR (400 MHz, DMSO) δ 0.90 (6 H, d), 1.43 (6 H, s), 1.87 (1 H, hept), 3.45 – 3.55 (5 H, m), 4.45 (2 H, s), 6.60 (1 H, s), 6.67 (1 H, dd), 6.76 – 6.86 (2 H, m), 7.01 – 7.23 (3 H, m), 7.35 (1 H, dd), 7.44 (1 H, ddd), 12.73 (1 H, s). LC-MS (ES, <i>m/z</i>): 439.


103	II		2-([1-[2-(Dimethylamino)phenyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 452 ¹ H-NMR (300 MHz, MeOD) δ 1.02 (6 H, d), 1.53 (6 H, s), 2.04 (1 H, dt), 3.02 (6 H, s), 3.65 (2 H, d), 4.60 (2 H, s), 5.48 (2 H, s), 6.49 (1 H, s), 6.81 (1 H, d), 6.92 – 7.05 (3 H, m), 7.26 (1 H, t), 7.42 (2 H, dt), 7.54 (1H, d).
104	II		2-methyl-2-([5-[3-(2-methylpropoxy)phenyl]-1-phenyl-1H-pyrazol-3-yl]methoxy)propanoic acid LC-MS (ES, <i>m/z</i>): 409 ¹ H-NMR (300 MHz, MeOD) 7.48 – 7.32 (m, 3H), 7.35 – 7.14 (m, 3H), 6.91 – 6.78 (m, 2H), 6.73 – 6.63 (m, 2H), 4.61 (s, 2H), 3.54 (d, <i>J</i> = 6.5 Hz, 2H), 1.92 (hept, <i>J</i> = 6.7 Hz, 1H), 1.53 (s, 6H), 0.95 (d, <i>J</i> = 6.7 Hz, 6H).
105	II		2-Methyl-2-([1-(2-methylphenyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)propanoic acid. LC-MS (ES, <i>m/z</i>): 423 ¹ H-NMR (300 MHz, MeOD) δ 0.95 (d, 6H), 1.54 (s, 6H), 1.81 – 2.01 (m, 4H), 3.34 (s, 1H), 3.46 (d, 2H), 4.61 (s, 2H), 6.63 (dd, 1H), 6.71 – 6.92 (m, 3H), 7.18 (t, 1H), 7.26 – 7.47 (m, 4H).
106	II		2-([1-(2-fluorophenyl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 427 ¹ H-NMR (300 MHz, MeOD) δ 7.57 – 7.46 (m, 2H), 7.38 – 7.29 (m, 1H), 7.28 – 7.16 (m, 2H), 6.91 – 6.81 (m, 2H), 6.72 (q, 2H), 4.63 (s, 2H), 3.56 (d, 2H), 1.98 – 1.93 (m, 1H), 1.55 (s, 6H), 0.98 (d, 6H).
107	XIII		2-[(1-[[4-(Dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 0.95 (7 H, d), 1.21 – 1.27 (1 H, m), 1.40 (6 H, s), 1.96 (1 H, dq), 2.84 (6 H, s), 3.64 (2 H, d), 4.40 (2 H, s), 5.18 (2 H, s), 6.37 (1 H, s), 6.60 – 6.67 (2 H, m), 6.83 – 6.92 (3 H, m), 6.98 (2 H, ddd), 7.36 (1 H, t), 12.68 (1 H, s). LC-MS (ES, <i>m/z</i>): 466.

108	XIII		2-[(1-[[3-(Dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 466 ¹ H-NMR (300 MHz, MeOD) δ 1.01 (6 H, d), 1.54 (6 H, s), 1.95 – 2.06 (1 H, m), 3.07 (6 H, s), 3.56 (2 H, d), 4.59 (2 H, s), 5.32 (2 H, s), 6.51 (1 H, s), 6.80 – 6.87 (3 H, m), 7.10 (2 H, d), 7.24 (1 H, ddd), 7.32 – 7.37 (2 H, dd).
109	XIII		2-[(1-[[2-(Dimethylamino)phenyl]methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 466 ¹ H-NMR (300 MHz, MeOD) δ 1.02 (6 H, d), 1.53 (6 H, s), 2.04 (1 H, dt), 3.02 (6 H, s), 3.65 (2 H, d), 4.60 (2 H, s), 5.48 (2 H, s), 6.49 (1 H, s), 6.81 (1 H, d), 6.92 – 7.05 (3 H, m), 7.26 (1 H, t), 7.42 (2 H, dt), 7.54 (1H, d).
110	VII		2-[(1-(2-Chlorophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid ¹ H NMR (MeOD, ppm): δ: 0.50 – 0.59 (m, 2H), 0.63 – 0.72 (m, 2H), 1.55 (s, 6H), 3.53 (tt, 1H), 4.63 (s, 2H), 6.73 (s, 1H), 6.86 – 6.98 (m, 3H), 7.22 (t, 1H), 7.41 – 7.61 (m, 4H). LC-MS (ES, <i>m/z</i>): 427.
111	VII		2-[(5-(3-Cyclopropoxyphenyl)-1-[(2-methoxyphenyl)methyl]-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 437 ¹ H-NMR (300 MHz, CDCl ₃) δ 0.54 – 0.61 (2 H, m), 0.65 (2 H, dt), 1.53(6 H, s), 3.45 – 3.58 (1 H, m), 3.74 (3 H, s), 4.63 (2 H, s), 5.38 (2 H, s), 6.41 (1 H, s), 6.76 – 6.84 (2 H, m), 6.85 – 6.92 (1 H, m), 7.01 (2 H, dd), 7.17 – 7.35 (3 H, m).
112	VII		2-[(5-(3-cyclopropoxyphenyl)-1-(2-methoxyphenyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 423 ¹ H NMR (300 MHz, MeOD) δ 7.45 (ddd, <i>J</i> = 8.3, 7.5, 1.7 Hz, 1H), 7.38 (dd, <i>J</i> = 8.0, 1.7 Hz, 1H), 7.19 (ddd, <i>J</i> = 7.6, 6.9, 2.0 Hz, 1H), 7.07 (m, 2H), 6.90 (m, 3H), 6.65 (s, 1H), 4.60 (s, 2H), 3.51 (m, 4H), 1.53 (s, 6H), 0.63 (m, 2H), 0.54 (m, 2H).

113	VII		2-([1-(2-Methoxyphenyl)-5-[3-(propan-2-yloxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 425 ¹ H-NMR (300 MHz, MeOD) δ 1.16 (6 H, d), 1.53 (6 H, s), 3.53 (3 H, s), 4.30 (1 H, p), 4.59 (2 H, s), 6.64 (1 H, s), 6.67 – 6.72 (1 H, m), 6.79 (1 H, ddd), 6.86 (1 H, dt), 7.01 – 7.10 (2 H, m), 7.17 (1 H, t), 7.34 – 7.39 (1 H, m), 7.43 (1 H, td).
114	VII		2-([5-(3-cyclopropoxyphenyl)-1-[2-(dimethylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 436 ¹ H-NMR (300 MHz, CDCl ₃) δ 7.47 – 7.29 (m, 2H), 7.26 – 7.06 (m, 3H), 6.96 – 6.81 (m, 3H), 6.71 (s, 1H), 4.63 (s, 2H), 3.50 (tt, <i>J</i> = 6.2, 3.0 Hz, 1H), 2.44 (s, 6H), 1.54 (s, 6H), 0.70 – 0.47 (m, 4H).
115	II		2-([5-(3-Methoxyphenyl)-1-(pyridin-3-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 1.55 (s, 6H), 3.72 (s, 3H), 4.65 (s, 2H), 6.73 (s, 1H), 6.79 – 6.88 (m, 2H), 6.96 (ddd, <i>J</i> = 8.4, 2.5, 0.9 Hz, 1H), 7.25 – 7.34 (m, 1H), 7.51 (dd, <i>J</i> = 8.3, 4.8 Hz, 1H), 7.82 (ddd, <i>J</i> = 8.3, 2.5, 1.4 Hz, 1H), 8.49 (d, <i>J</i> = 2.5 Hz, 1H), 8.53 (dd, <i>J</i> = 4.9, 1.4 Hz, 1H). LC-MS (ES, <i>m/z</i>): 368.
116	II		2-([5-(3-Methoxyphenyl)-1-(pyridin-3-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 368. ¹ H-NMR (400 MHz, DMSO) δ 2.23 (s, 5H), 4.53 (s, 3H), 5.31 (s, 2H), 7.49 (s, 1H), , 7.63 (d, 1H), 7.22 (t, 1H), 7.72 (s, 1H), 7.81 – 7.83 (m, 1H), 8.07 – 8.09 (m, 2H), 8.12 – 8.16 (m, 1H), 9.38 (d, 2H).
117	II		2-([1-(2-Fluorophenyl)-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 384 ¹ H-NMR (300 MHz, MeOD) δ 1.55 (6 H, d), 3.66 (3H, s), 4.63 (2 H, s), 6.72 (1 H, s), 6.75 (1 H, dd), 6.81 – 6.92 (2 H, m), 7.17-7.28 (2 H, m), 7.28-7.37 (1 H, m), 7.46-7.56 (2H, m).

118	II		2-([5-(3-methoxyphenyl)-1-(2-methylphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 381 ¹ H-NMR (300 MHz, MeOD) δ 7.49 – 7.28 (m, 4H), 7.18 (t, 1H), 6.94 – 6.79 (m, 2H), 6.74 (s, 1H), 6.67 (dd, 1H), 4.61 (s, 2H), 3.59
119	II		22-([1-(2-Methoxyphenyl)-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, <i>m/z</i>): 397 ¹ H-NMR (300 MHz, CDCl ₃) δ 1.57 (6 H, s), 3.48 (3 H, s), 3.62 (3 H, s), 4.68 (2 H, s), 6.57 (1 H, s), 6.72 (1 H, dd), 6.76 – 6.85 (2 H, m), 6.88 (1 H, dd), 7.01 (1 H, td), 7.15 (1 H, t), 7.31 – 7.36 (1 H, m), 7.37 – 7.44 (1 H, m).

Examples 120/121: (2*S*)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1*H*-pyrazol-3-yl]methoxy)-2-methylbutanoic acid and (2*R*)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1*H*-pyrazol-3-yl]methoxy)-2-methylbutanoic acid.




[0475] *Ethyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1*H*-pyrazol-3-yl]methoxy)-2-methylbutanoate*. To a solution of Int. **G-1** (700 mg, 1.79 mmol, 1.00 equiv) in THF (10 mL), under N₂, was added NaH (200 mg, 8.33 mmol, 3.00 equiv). The solution was stirred for 10 min at 0 °C. To this was added methyl 2-hydroxy-2-methylbutanoate (700 mg, 5.30 mmol, 3.00 equiv), Bu₄NI (330 mg, 0.50 equiv). The resulting solution was stirred for 4 h at room temperature. The reaction was quenched by the addition of 1 mL of water. The resulting solution was diluted with 20 mL of H₂O, extracted with 2x30 mL of EtOAc, and the organic layers were combined. The resulting mixture was washed with 20 mL of saturated NaCl, then dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was applied onto a silica gel column with EtOAc / petroleum ether (1:5). This resulted in the title compound as a yellow oil (300 mg, 38%).

[0476] (vi) Chiral separation of *Ethyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate*. The crude product of step v was purified by Chiral-Prep-HPLC with the following conditions (Prep-HPLC-004): Column, Chiralpak IC, 2*25cm, 5um; mobile phase, Hex- and ethanol- (hold 2.0% ethanol- in 23 min); Detector, UV 254/220 nm. 100 mg product was obtained which Alpha(25 °C, 0.05 mol/L, MeOH). This resulted in ethyl (2*S*)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate as colorless oil (100 mg, 36%) and ethyl (2*R*)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate as colorless oil (100 mg, 36%).


[0477] (vii) (2*S*)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid was obtained from ethyl (2*S*)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate analogously as described in Step iv of Example 1. ¹H-NMR (400 MHz, MeOD) δ 7.45 – 7.38 (m, 1H), 7.34 (d, J = 7.9 Hz, 1H), 7.31 – 7.29 (m, 1H), 7.29 – 7.24 (m, 1H), 6.83 (dd, J = 2.5, 1.6 Hz, 1H), 6.80 – 6.72 (m, 1H), 6.56 (s, 1H), 5.44 (s, 2H), 4.59 (s, 2H), 3.69 (s, 3H), 1.89 (qd, J = 7.2, 5.0 Hz, 2H), 1.51 (s, 3H), 0.98 (t, J = 7.5 Hz, 3H). LC-MS (ES, *m/z*): 429.

[0478] (viii) (2*R*)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid was obtained from ethyl (2*R*)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate analogously as described in Step iv of Example 1. ¹H-NMR (400 MHz, MeOD) δ 7.45 – 7.38 (m, 1H), 7.37 – 7.31 (m, 1H), 7.31 – 7.29 (m, 1H), 7.29 – 7.24 (m, 1H), 6.97 (ddd, J = 8.4, 2.6, 0.9 Hz, 1H), 6.93 (dt, J = 7.6, 1.2 Hz, 1H), 6.83 (dd, J = 2.6, 1.6 Hz, 1H), 6.81 – 6.72 (m, 1H), 6.56 (s, 1H), 5.44 (s, 2H), 4.59 (s, 2H), 3.69 (s, 3H), 1.89 (qd, J = 7.2, 4.9 Hz, 2H), 1.51 (s, 3H), 0.98 (t, J = 7.5 Hz, 3H). LC-MS (ES, *m/z*): 429.

Alternative synthesis of Examples 120/121: (2S)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid and (2R)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid.




[0479] **Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate** To a solution of methyl 2-hydroxy-2-methylbutanoate (780 mg, 5.90 mmol, 1.00 equiv) in THF (50 mL) under N₂ was added NaH (236 mg, 5.90 mmol, 1.00 equiv) in portions at 0 °C. The mixture was stirred at 0 °C for 30 min, then Int. **G-1** (1.16 g, 2.96 mmol, 0.50 equiv) and Bu₄NI (600 mg, 1.63 mmol, 0.50 equiv) were added at 0 °C. The resulting solution was stirred for 16 h at rt, then then quenched by the addition of water/ice. The resulting solution was diluted with 200 mL of EtOAc, washed with 2x100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Prep OBD C18 Column, 30x150mm 5um; mobile phase, Water (0.1%FA) and ACN (55.0% ACN up to 79.0% in 7 min); Detector, UV 254/220nm, to afford 200 mg (8%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 443.



[0480] **Chiral separation of (2R)- and (2S)- Methyl 2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate** The racemic methyl ester (200 mg, 0.45 mmol, 1.00 equiv) was separated into enantiomers by Chiral-Prep-HPLC with the following conditions Column: Chiralpak IC, 2*25cm, 5um; Mobile Phase A: Hex--HPLC, Mobile Phase B: EtOH--HPLC; Flow rate: 20 mL/min; Gradient: 2 B to 2 B in 24 min; 220/254 nm ; RT1:18.171 ; RT2:20.742, affording 80 mg (40%) of methyl (2S)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate, and 70 mg (35%) of methyl (2R)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate, both as colorless oils.


Alternate synthesis of methyl (2S)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate



[0481] **Methyl (2S)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate** To a solution of methyl (2S)-2-hydroxy-2-methylbutanoate (400 mg, 3.03 mmol, 1.00 equiv) in THF (20 mL) was added NaH (122


mg, 3.05 mmol, 1.00 equiv) in portions at 0 °C. The mixture was stirred at 0 °C for 10 min, then Int. **G-1** (400 mg, 1.02 mmol, 0.34 equiv) and Bu₄NI (200 mg, 0.54 mmol, 0.50 equiv) were added at 0 °C. The resulting solution was stirred for 16 h at rt, quenched by the addition of water/ice, and extracted with 100 mL of EtOAc. The combined organic layers were washed with 1x100 mL of brine, dried over Na₂SO₄, concentrated under vacuum, and purified with Prep-TLC using EtOAc / petroleum ether (1:3) to afford 150 mg (11%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 443.

Example 120. (2S)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid



[0482] To a solution of methyl (2S)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate (70 mg, 0.16 mmol, 1.00 equiv) in THF / H₂O / MeOH (4/2/2 mL) was added LiOH (23 mg, 0.96 mmol, 6.00 equiv), in portions at rt. The resulting solution was stirred for 2 h at room temperature. The pH was adjusted to 3-4 with conc HCl. The resulting solution was extracted with 100 mL of EtOAc. The combined organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Prep OBD C18 Column, 30x150mm 5um; mobile phase, Water (0.1%FA) and ACN (40.0% ACN up to 75.0% in 7 min); Detector, UV 254/220nm to afford 40 mg (59%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 429. ¹H NMR: (400 MHz, DMSO-d₆) δ 12.60 (s, 1H), 7.50 – 7.22 (m, 4H), 7.02 – 6.84 (m, 3H), 6.78 – 6.69 (m, 1H), 6.45 (s, 1H), 5.38 (s, 2H), 4.41 (d, J = 2.2 Hz, 2H), 3.69 (s, 3H), 1.87 – 1.61 (m, J = 7.1 Hz, 2H), 1.36 (s, 3H), 0.85 (t, J = 7.4 Hz, 3H).

Example 121. (2R)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid



[0483] To a solution of methyl (2R)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoate (80 mg, 0.18 mmol, 1.00 equiv) in THF / H₂O / MeOH (4/2/2 mL) was added LiOH (26 mg, 1.09 mmol, 6.00 equiv) in portions at rt. The resulting solution was stirred for 2 h at rt. The pH value was adjusted to 3-4 with conHCl. The resulting solution was extracted with 100 mL of EtOAc, and the organic layers were combined, dried over Na₂SO₄, concentrated under vacuum, and purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Prep OBD C18 Column, 30??150mm 5um; mobile phase, Water (0.1%FA) and ACN (40.0% ACN up to 75.0% in 7 min); Detector, UV 254/220nm to afford 46.3 mg (60%) of the title compound as a colorless oil. LC-MS: (ES, *m/z*): 429. ¹H NMR: (400 MHz, DMSO-*d*₆) δ 12.61 (s, 1H), 7.49 – 7.22 (m, 4H), 7.02 – 6.84 (m, 3H), 6.78 – 6.69 (m, 1H), 6.46 (s, 1H), 5.38 (s, 2H), 4.41 (d, *J* = 2.2 Hz, 2H), 3.69 (s, 3H), 1.84 – 1.61 (m, *J* = 7.6 Hz, 2H), 1.36 (s, 3H), 0.85 (t, *J* = 7.4 Hz, 3H).

[0484] The following Examples were prepared using Scheme III.

Table 23. Examples 122-127.





Ex.	Structure	IUPAC Name Analytical data
122		(2S)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-butanoic acid. LC-MS (ES, <i>m/z</i>): 415. ¹H-NMR (400 MHz, MeOD) δ 7.46 – 7.37 (m, 1H), 7.36 – 7.21 (m, 3H), 7.00 – 6.88 (m, 2H), 6.82 (dd, <i>J</i> = 2.6, 1.5 Hz, 1H), 6.79 – 6.71 (m, 1H), 6.58 (s, 1H), 5.44 (s, 2H), 4.76 (d, <i>J</i> = 12.1 Hz, 1H), 4.47 (d, <i>J</i> = 12.1 Hz, 1H), 3.82 (dd, <i>J</i> = 7.3, 4.7 Hz, 1H), 3.68 (s, 3H), 1.89 – 1.66 (m, 2H), 1.00 (t, <i>J</i> = 7.4 Hz, 3H).

123		(2R)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-butanoic acid. LC-MS (ES, <i>m/z</i>): 415 ¹ H-NMR (300 MHz, MeOD) δ 7.46 – 7.36 (m, 1H), 7.36 – 7.22 (m, 3H), 7.00 – 6.88 (m, 2H), 6.82 (dd, <i>J</i> = 2.6, 1.5 Hz, 1H), 6.78 – 6.71 (m, 1H), 6.58 (s, 1H), 5.44 (s, 2H), 4.75 (d, <i>J</i> = 12.1 Hz, 1H), 4.46 (d, <i>J</i> = 12.1 Hz, 1H), 3.82 (dd, <i>J</i> = 7.4, 4.8 Hz, 1H), 3.68 (s, 3H), 1.90 – 1.66 (m, 2H), 1.00 (t, <i>J</i> = 7.4 Hz, 3H).
124		(2S)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-propanoic acid LC-MS (ES, <i>m/z</i>): 401 ¹ H-NMR (300 MHz, MeOD) δ 7.49 – 7.36 (m, 1H), 7.34 – 7.22 (m, 3H), 7.01 – 6.89 (m, 2H), 6.83 (dd, <i>J</i> = 2.6, 1.6 Hz, 1H), 6.79 – 6.71 (m, 1H), 6.55 (s, 1H), 5.45 (s, 2H), 4.73 (d, <i>J</i> = 11.9 Hz, 1H), 4.56 (d, <i>J</i> = 11.9 Hz, 1H), 4.17 (q, <i>J</i> = 6.9 Hz, 1H), 3.68 (s, 3H), 1.44 (d, <i>J</i> = 6.9 Hz, 3H).
125		(2R)-2-([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-propanoic acid. LC-MS (ES, <i>m/z</i>): 401 ¹ H-NMR (300 MHz, MeOD) δ 7.45 – 7.37 (m, 1H), 7.37 – 7.22 (m, 3H), 7.01 – 6.89 (m, 2H), 6.83 (dd, <i>J</i> = 2.6, 1.6 Hz, 1H), 6.79 – 6.71 (m, 1H), 6.56 (s, 1H), 5.45 (s, 2H), 4.73 (d, <i>J</i> = 11.9 Hz, 1H), 4.56 (d, <i>J</i> = 11.9 Hz, 1H), 4.17 (q, <i>J</i> = 6.9 Hz, 1H), 3.69 (s, 3H), 1.44 (d, <i>J</i> = 6.8 Hz, 3H).
126		2-([1-[(2-Chlorophenyl)methyl]-5-(2-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD) δ 1.52 (6 H, s), 3.64 (3 H, s), 4.57 (2 H, s), 5.24 (2 H, s), 6.40 (1 H, s), 6.73 – 6.83 (1 H, m), 6.89 – 7.10 (2 H, m), 7.12 – 7.24 (3 H, m), 7.25 – 7.46 (2 H, m). LC-MS (ES, <i>m/z</i>): 415.
127		2-([1-[(2-Chlorophenyl)methyl]-5-(4-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. ¹ H-NMR (400 MHz, MeOD) δ 7.44 – 7.37 (m, 1H), 7.35 – 7.07 (m, 4H), 7.03 – 6.89 (m, 2H), 6.73 (dd, <i>J</i> = 7.1, 2.1 Hz, 1H), 6.50 (s, 1H), 5.41 (s, 2H), 4.58 (s, 2H), 3.82 (s, 3H), 1.53 (s, 6H). LC-MS (ES, <i>m/z</i>): 415.



[0485] The following Examples were prepared using the indicated Schemes.

Table 24. Examples 128-142.


Ex.	Scheme	Structure	IUPAC Name Analytical Data
128	VIII		2-([1-[(2-Chlorophenyl)methyl]-5-[5-(2-methylpropoxy)thien-2-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 1.01 (dd, 6H), 1.52 (s, 6H), 2.07 (ddd, 1H), 3.83 (dd, 2H), 4.55 (s, 2H), 5.52 (s, 2H), 6.17 – 6.23 (m, 1H), 6.53 (s, 1H), 6.58 – 6.66 (m, 2H), 7.20 – 7.35 (m, 2H), 7.45 (dt, 1H). LC-MS (ES, <i>m/z</i>): 463.
129	VIII		2-([1-[(2-Chlorophenyl)methyl]-5-[5-(propan-2-yloxy)thien-2-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 1.34 (d, 6H), 1.52 (s, 6H), 4.42 (p, 1H), 4.56 (s, 2H), 5.52 (s, 2H), 6.22 (dd, 1H), 6.53 (s, 1H), 6.59 – 6.67 (m, 2H), 7.19 – 7.35 (m, 2H), 7.45 (dd, 1H). LC-MS (ES, <i>m/z</i>): 449.
130	IX		2-([1-[(2-Chlorophenyl)methyl]-5-(4-cyclopropoxythien-2-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 1.52 (s, 6H), 1.74 – 1.90 (m, 2H), 2.65 (t, 2H), 4.45 (t, 1H), 4.58 (s, 2H), 5.55 (s, 2H), 6.61 – 6.69 (m, 2H), 6.90 (d, 1H), 7.13 (d, 1H), 7.20 – 7.38 (m, 3H), 7.45 (dd, 1H). LC-MS (ES, <i>m/z</i>): 447.
131	IX		2-([1-[(2-Chlorophenyl)methyl]-5-(5-cyclopropoxythien-2-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 1.53 (s, 6H), 1.89 (qd, 2H), 2.89 (t, 2H), 4.51 (t, 1H), 4.58 (s, 2H), 5.55 (s, 2H), 6.62 (d, 2H), 6.76 – 6.88 (m, 2H), 7.27 (dt, 2H), 7.45 (dd, 1H). LC-MS (ES, <i>m/z</i>): 447.

132	VIII		2-([1-[(2-Chlorophenyl)methyl]-5-[2-(2-methylpropoxy)-1,3-thiazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 7.46 (dd, <i>J</i> = 7.9, 1.3 Hz, 1H), 7.32 (td, <i>J</i> = 7.7, 1.7 Hz, 1H), 7.25 (td, <i>J</i> = 7.6, 1.4 Hz, 1H), 7.04 (s, 1H), 6.68 – 6.60 (m, 2H), 5.52 (s, 2H), 4.57 (s, 2H), 4.18 (d, <i>J</i> = 6.6 Hz, 2H), 2.12 (hept, <i>J</i> = 6.7 Hz, 1H), 1.52 (s, 6H), 1.33 (dd, <i>J</i> = 17.9, 3.5 Hz, 1H), 1.02 (d, <i>J</i> = 6.7 Hz, 6H). LC-MS (ES, <i>m/z</i>): 464.
133	VIII		2-([1-[(2-Chlorophenyl)methyl]-5-[2-(propan-2-yloxy)-1,3-thiazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 7.46 (dd, <i>J</i> = 7.9, 1.4 Hz, 1H), 7.31 (td, <i>J</i> = 7.7, 1.7 Hz, 1H), 7.25 (td, <i>J</i> = 7.5, 1.4 Hz, 1H), 7.03 (s, 1H), 6.64 (dd, <i>J</i> = 7.6, 1.7 Hz, 1H), 6.61 (s, 1H), 5.52 (s, 2H), 5.13 (dq, <i>J</i> = 12.3, 6.2 Hz, 1H), 4.57 (s, 2H), 1.52 (s, 6H), 1.40 (d, <i>J</i> = 6.2 Hz, 6H). LC-MS (ES, <i>m/z</i>): 450.
134	X		2-([1-[(2-Chlorophenyl)methyl]-5-[2-(2-methylpropyl)-1,3-oxazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) 7.46 (dd, <i>J</i> = 7.9, 1.3 Hz, 1H), 7.29 (td, <i>J</i> = 7.5, 1.6 Hz, 1H), 7.25 – 7.18 (m, 1H), 7.17 (s, 1H), 6.81 (s, 1H), 6.59 (dd, <i>J</i> = 7.7, 1.6 Hz, 1H), 5.65 (s, 2H), 4.59 (s, 2H), 2.64 (d, <i>J</i> = 7.2 Hz, 2H), 2.09 – 1.93 (m, 1H), 1.53 (s, 6H), 0.88 (d, <i>J</i> = 6.7 Hz, 6H). LC-MS (ES, <i>m/z</i>): 442
135	XI		2-([1-[(2,6-Dimethoxyphenyl)methyl]-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 0.98 (6 H, d), 1.44 (6 H, s), 2.01 (1 H, dt), 3.16 (1 H, s), 3.32 (1 H, s), 3.76 (8 H, d), 4.66 (2 H, s), 5.28 (2 H, s), 6.59 (1 H, s), 6.69 (2 H, d), 6.79 (1 H, dt), 7.16 – 7.35 (4 H, m). LC-MS (ES, <i>m/z</i>): 483.

136	XIV		2-[(1-[[2-(Dimethylamino)-6-fluorophenyl]methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid ¹ H-NMR (400 MHz, DMSO) δ 1.44 (s, 6H), 2.48 (s, 6H), 3.83 (s, 3H), 4.49 (s, 2H), 5.44 (s, 2H), 6.43 (s, 1H), 6.75 (t, 1H), 6.87 – 7.06 (m, 4H), 7.23 (td, 1H), 7.35 (t, 1H). LC-MS (ES, <i>m/z</i>): 441.
137	XII		2-([5-(3-Methoxyphenyl)-1-(2-methylpropyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 0.76 (d, 6H), 1.28 – 1.38 (m, 3H), 1.52 (s, 6H), 2.07 (dt, 1H), 3.85 (s, 3H), 3.96 (d, 2H), 4.55 (s, 2H), 6.38 (s, 1H), 6.94 – 7.07 (m, 3H), 7.41 (t, 1H). LC-MS (ES, <i>m/z</i>): 347.
138	XII		2-([5-(3-Methoxyphenyl)-1-([oxan-4-yl]methyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 1.13 (qd, 2H), 1.28 – 1.41 (m, 5H), 1.52 (s, 6H), 2.06 (dq, 1H), 3.23 – 3.34 (m, 2H), 3.86 (s, 5H), 4.05 (d, 2H), 4.54 (s, 2H), 6.37 (d, 1H), 6.95 – 7.08 (m, 3H), 7.43 (t, 1H). LC-MS (ES, <i>m/z</i>): 389.
139	XI		2-([1-[(3-Chloropyridin-2-yl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid ¹ H-NMR (400 MHz, MeOD) δ 8.37 (dd, <i>J</i> = 4.7, 1.4 Hz, 1H), 7.89 (dd, <i>J</i> = 8.1, 1.5 Hz, 1H), 7.38 – 7.24 (m, 4H), 6.91 – 6.83 (m, 1H), 6.71 (s, 1H), 5.87 (s, 2H), 4.64 (s, 2H), 3.83 (s, 3H), 1.42 (s, 6H). LC-MS (ES, <i>m/z</i>): 416.
140	XI		2-([5-(3-Methoxyphenyl)-1-[[2-(propan-2-yloxy)phenyl]methyl]-1H-pyrazol-3-yl)methoxy]-2-methylpropanoic acid ¹ H-NMR (300 MHz, MeOD) δ 1.25 (6 H, d), 1.52 (6 H, s), 3.64 (3 H, s), 4.57 (2 H, s), 4.59 – 4.68 (1 H, m), 5.30 (2 H, s), 6.51 (1 H, s), 6.67 (1 H, dd), 6.78 – 6.88 (2 H, m), 6.90 – 7.00 (3 H, m), 7.19 – 7.26 (1 H, m), 7.29 (1 H, dd). LC-MS (ES, <i>m/z</i>): 439.

141	XI		2-([1-[(2-Ethoxyphenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid. LC-MS (ES, m/z): 425 $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.33 (3 H, t), 1.56 (6 H, s), 3.68 (3 H, s), 4.00 (2 H, q), 4.65 (2 H, s), 5.40 (2 H, s), 6.43 (1 H, s), 6.77 – 6.82 (1 H, m), 6.83 – 6.89 (3 H, m), 6.89 – 6.98 (2 H, m), 7.16 – 7.24 (1 H, m), 7.26 – 7.32 (1 H, m).
142	XI		2-([1-[(2-Cyclopropoxyphenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, m/z): 437 $^1\text{H-NMR}$ (300 MHz, MeOD) δ 7.27 (m, 3H), 6.94 (m, 2H), 6.88 (m, 1H), 6.82 (dd, J = 2.6, 1.5 Hz, 1H), 6.72 (d, J = 7.4 Hz, 1H), 6.48 (s, 1H), 5.25 (s, 2H), 4.56 (s, 2H), 3.77 (tt, J = 6.0, 6.0, 3.0, 3.0 Hz, 1H), 3.67 (s, 3H), 1.51 (s, 6H), 0.75 (m, 2H), 0.58 (m, 2H).

Example 143: (3E)-4-[1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbut-3-enoic acid.




[0486] (ii) ([1-[(2-Chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methyl)triphenylphosphonium bromide. To a solution of Int. **G-1** (2.4 g, 6.13 mmol, 1.00 equiv) in toluene (20 mL) was added PPh_3 (2.4 g, 9.15 mmol, 1.50 equiv) at room temperature. The resulting solution was stirred for 3 h at 110 °C. The solids were collected by filtration. The solid was dried in an oven under reduced pressure. This resulted in the title compound as a white solid (2.2 g, 55%).

[0487] (iii) Methyl (3E)-4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbut-3-enoate. To a solution of ([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methyl)triphenylphosphonium bromide (500 mg, 0.76 mmol, 1.00 equiv) in THF (20 mL) was added methyl 2,2-dimethyl-3-oxopropanoate (200 mg, 1.54 mmol, 2.00 equiv). This was followed by the addition of NaH (50 mg, 1.25 mmol,

2.00 equiv), in portions at 0 °C. The resulting solution was stirred for 16 h at room temperature then quenched by the addition of water/ice. The resulting solution was diluted with 100 mL of EtOAc, then washed with 2x50 mL of brine. The mixture was dried over Na₂SO₄ and concentrated under vacuum. The residue was applied onto a Prep-TLC with EtOAc / petroleum ether (1:2). This resulted in the title compound as a colorless oil (125 mg, 38%).

[0488] (iv) (3E)-4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbut-3-enoic acid was obtained analogously as described in Step iv of Example 101 using methyl (3E)-4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbut-3-enoate as starting material. ¹H-NMR (400 MHz, MeOD) δ 1.42 (s, 6H), 3.69 (s, 3H), 5.43 (s, 2H), 6.49 (d, 1H), 6.56 – 6.68 (m, 2H), 6.71 – 6.79 (m, 1H), 6.84 (dd, 1H), 6.90 – 7.06 (m, 2H), 7.23 – 7.47 (m, 5H). LC-MS (ES, *m/z*): 411.

Example 144: 4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbutanoic acid.



[0489] (i) Methyl (3E)-4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbut-3-enoate was obtained as in Steps i-iii of Example 143.


[0490] (ii) Methyl 4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbutanoate. To a solution of methyl (3E)-4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbut-3-enoate (60 mg, 0.14 mmol, 1.00 equiv) in EtOAc / MeOH (10/10 mL) was added Pd/C (20 mg). The resulting solution was stirred for 3 h under H₂ at room temperature. The solids were removed by filtration. The resulting mixture was concentrated under vacuum. The residue was purified by Prep-TLC with 20:1 CH₂Cl₂ : MeOH. This resulted in the title compound as a colorless oil (40 mg, 66%).

[0491] (iii) 4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbutanoic acid was obtained analogously as described in Step iv of Example 101 using methyl 4-[1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]-2,2-dimethylbutanoate as starting material. ¹H-NMR (400 MHz, MeOD) δ 1.26 – 1.38 (m, 6H),

1.98 (s, 2H), 2.69 (s, 2H), 3.68 (s, 3H), 5.41 (s, 2H), 6.35 (s, 1H), 6.69 (dd, 1H), 6.81 (dd, 1H), 6.87 – 7.00 (m, 2H), 7.23 – 7.36 (m, 3H), 7.38 – 7.45 (m, 1H). LC-MS (ES, m/z): 413.


[0492] Examples 145-149 are intentionally left blank.

Example 150: 2-([1-[(2-Chlorophenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid



[0493] To a solution of Int. **H-64** (120 mg, 0.26 mmol, 1.00 equiv) in THF / H₂O / MeOH (8 / 4 / 4 mL) was added LiOH (37 mg, 1.54 mmol, 6.00 equiv), in portions at room temperature. The resulting solution was stirred for 2 h at room temperature. The pH was then adjusted to 3-4 with aq HCl. The resulting solution was extracted with 100 mL EtOAc, and the combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The crude product (100 mg) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Prep OBD C18 Column, 300x150mm 5um; mobile phase, Water (0.1%FA) and ACN (38.0% ACN up to 65.0% in 7 min); Detector, uv 254/220nm, to afford 42 mg (36%) of the title compound as a white solid. LC-MS: (ES, m/z): 453. ¹H-NMR: (DMSO)δ: 12.70 (s, 1H), 8.09 (d, J = 0.9 Hz, 1H), 7.80 (dd, J = 8.4, 0.8 Hz, 1H), 7.70 (t, J = 1.1 Hz, 1H), 7.48 – 7.37 (m, 1H), 7.36 – 7.23 (m, 2H), 7.14 (dd, J = 8.4, 1.3 Hz, 1H), 6.86 – 6.75 (m, 1H), 6.55 (s, 1H), 5.47 (s, 2H), 4.46 (s, 2H), 4.41 (q, J = 0.9 Hz, 2H), 1.42 (s, 6H), 1.36 (t, J = 0.6 Hz, 3H).


Example 151: Sodium 2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate




[0494] A solution of 2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid (Example **150**, 50 mg, 0.11 mmol, 1.00

equiv) and NaOH (18 mg, 0.45 mmol, 4.00 equiv) in MeOH / H₂O (5/2 mL) was stirred for 3 h at rt, then purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Shield RP18 OBD Column,, 5um, 19*150mm; mobile phase, Water(10MMOL/L NH₄HCO₃) and MeOH- (15.0% MeOH- up to 95.0% in 7 min); Detector, UV 254/220nm, to afford 20.9 mg (40%) of the title compound as a white solid. LC-MS: (ES, *m/z*): 453.00. ¹H NMR (300 MHz, DMSO) δ 8.07 (s, 1H), 7.77 (d, *J* = 8.3 Hz, 1H), 7.65 (s, 1H), 7.41 (m, 1H), 7.28 (m, 2H), 7.10 (d, *J* = 8.4 Hz, 1H), 6.79 (m, 1H), 6.56 (s, 1H), 5.46 (s, 2H), 4.49 (s, 2H), 4.37 (q, *J* = 7.2, 7.2, 7.2 Hz, 2H), 1.34 (d, *J* = 17.1 Hz, 9H).

Example 152: 2-([5-(3-Cyclopropoxyphenyl)-1-(1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid




[0495] (i) 2-([5-(3-Cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid A solution of Int. H-76 (100 mg, 0.17 mmol, 1.00 equiv) and LiOH (20 mg, 0.84 mmol, 4.00 equiv) in THF/H₂O (4/1 mL) was stirred for 2 h at room temperature, then cooled to 0 °C. The pH was adjusted to 1 with 1 M HCl. The resulting solution was extracted with 2 x 100 mL EtOAc, and the combined organic layers were washed with 100 mL of brine, dried over Na₂SO₄ and concentrated under vacuum, to afford 50 mg (51%) of the title compound as a white solid. LC-MS: (ES, *m/z*): 563.



[0496] (ii) 2-([5-(3-Cyclopropoxyphenyl)-1-(1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid A solution of the product from the previous step (50 mg, 0.09 mmol, 1.00 equiv) and TFA (3 mL) in CH₂Cl₂ (10 mL) was stirred for 5 h at room

temperature, then concentrated under vacuum. The crude product (50 mg) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, Xselect CSH OBD Column 30*150mm 5um, n; mobile phase, Water (0.1%FA) and ACN (43.0% ACN up to 55.0% in 7 min); Detector, UV 254/220nm, to afford 15 mg (39%) of the title compound as a white solid. LC-MS: (ES, m/z): 433. ^1H NMR (300 MHz, DMSO- d_6) δ 13.09 (s, 1H), 8.18 (s, 1H), 7.81 (dd, $J = 7.0, 2.0$ Hz, 1H), 7.20 (t, $J = 7.9$ Hz, 1H), 7.14 – 7.02 (m, 2H), 6.95 – 6.85 (m, 2H), 6.85 – 6.80 (m, 1H), 6.71 (s, 1H), 4.55 (s, 2H), 3.45 (tt, $J = 6.3, 3.0$ Hz, 1H), 1.44 (s, 6H), 0.58 – 0.43 (m, 2H), 0.43 – 0.26 (m, 2H).

Example 153: 2-([5-(3-Cyclopropoxyphenyl)-1-(1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid




[0497] (i) 2-([5-(3-cyclopropoxyphenyl)-1-(1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid A solution of Int. **H-77** (160 mg, 0.28 mmol, 1.00 equiv) and LiOH (70 mg, 2.92 mmol, 10.54 equiv) in THF / MeOH / H₂O (5/1/1 mL) was stirred for 3 h at room temperature. The pH value was adjusted to 5 with 1 M HCl. The resulting solution was extracted with 3 x 20 mL EtOAc. The combined organic layers were concentrated under reduced pressure and dried, to afford 140 mg (90%) of the title compound as light yellow oil. LC-MS: (ES, m/z):563.2.

[0498] (ii) 2-([5-(3-Cyclopropoxyphenyl)-1-(1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid A solution of the product from the previous step (140 mg, 0.25 mmol, 1.00 equiv) and HCl / dioxane (5 mL) was stirred for 3 h at room temperature, then concentrated under vacuum. The crude product (100 mg) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, Xselect CSH F-Phenyl OBD Column 19*150mm 5um, n; mobile phase, Water (0.1%FA) and ACN (35.0% ACN up to 45.0% in 7 min); Detector, uv 254/220nm, to afford 18.9 mg (18%) of the title compound as a white solid. LC-MS: (ES, m/z):433.05. ^1H NMR (300 MHz, DMSO) δ 13.29 (s, 1H), 7.69 (d, $J = 1.0$ Hz, 1H), 7.57 (d, $J = 8.4$ Hz, 1H), 7.35 (dd, $J = 8.4, 7.3$ Hz, 1H), 7.21 (t, $J = 7.9, 7.9$ Hz, 1H), 6.91 (dtd, $J = 8.3, 5.8, 4.8, 3.0$ Hz, 3H), 6.81 (m, 1H), 6.70

(s, 1H), 4.53 (s, 2H), 3.52 (tt, $J = 6.2, 6.2, 3.0, 3.0$ Hz, 1H), 1.44 (s, 6H), 0.46 (m, 2H), 0.36 (m, 2H).

Example 154: 2-([5-(3-cyclopropoxyphenyl)-1-(3-methyl-1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid







[0499] (i) 2-([5-(3-cyclopropoxyphenyl)-1-(3-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid A solution of Int. H-78 (100 mg, 0.17 mmol, 1.00 equiv) and LiOH (40 mg, 1.67 mmol, 10.00 equiv) in THF / MeOH / H₂O (5/1/1 mL) was stirred for 3 h at room temperature. The pH value of the solution was adjusted to 5 with 1 M HCl. The resulting solution was extracted with 3 x 15 mL of EtOAc. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum, to afford 80 mg (82%) of the title compound as light yellow oil. LC-MS: (ES, m/z): 577.3.





[0500] (ii) 2-([5-(3-cyclopropoxyphenyl)-1-(3-methyl-1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid A solution of the product from the previous step (80 mg, 0.14 mmol, 1.00 equiv) in HCl / dioxane (5 mL) was stirred 3 h at room temperature, then concentrated under vacuum. The crude product was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, Xselect CSH OBD Column 30*150mm 5um, n; mobile phase, Water (0.1%FA) and ACN (40.0% ACN up to 50.0% in 7 min); Detector, UV 254/220nm, to afford 8.2 mg (13%) of the title compound as a white solid. LC-MS: (ES, m/z): 447.15. ¹H NMR (300 MHz, MeOD) δ 7.62 (dd, $J = 8.5, 0.8$ Hz, 1H), 7.42 (dd, $J = 8.5, 7.2$ Hz, 1H), 7.17 (t, $J = 7.9, 7.9$ Hz, 1H), 6.99 (m, 2H), 6.80 (m, 3H), 4.65 (s, 2H), 3.25 (dt, $J = 5.7, 2.6, 2.6$ Hz, 1H), 2.07 (s, 3H), 1.54 (s, 6H), 0.41 (m, 4H).

[0501] The following substituted pyrazole carboxylic acids were obtained from LiOH hydrolysis of the corresponding methyl esters:





Table 25: Examples 155-276.

Ex.	Ester	Structure	IUPAC Name Purification Analytical data
155	H-134		2-([1-[2-(Azetidin-1-yl)phenyl]-5-[3-(2,2-dimethylpropoxy)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 63% ACN up to 83% in 7 min LC-MS: (ES, <i>m/z</i>): 478.25. ¹ H NMR (300 MHz, MeOD) δ 7.30 (ddd, <i>J</i> = 8.6, 7.4, 1.6 Hz, 1H), 7.20 (t, <i>J</i> = 7.9, 7.9 Hz, 1H), 7.03 (m, 2H), 6.87 (t, <i>J</i> = 2.0, 2.0 Hz, 1H), 6.82 (ddd, <i>J</i> = 8.2, 2.5, 1.0 Hz, 1H), 6.75 (td, <i>J</i> = 7.6, 7.5, 1.3 Hz, 1H), 6.70 (s, 1H), 6.55 (dd, <i>J</i> = 8.2, 1.3 Hz, 1H), 4.59 (s, 2H), 3.57 (d, <i>J</i> = 21.2 Hz, 4H), 2.13 (p, <i>J</i> = 7.3, 7.3, 7.3, 7.3 Hz, 2H), 1.53 (s, 6H), 0.98 (s, 9H).
156	H-38		2-([1-(2-Chlorophenyl)-5-(pyridin-2-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid Xselect: 10% ACN up to 67% in 7 min LC-MS: (ES, <i>m/z</i>): 371.95. H-NMR: ¹ H NMR (300 MHz, DMSO) δ 12.68 (s, 1H), 8.31 (ddd, <i>J</i> = 4.9, 1.8, 0.9 Hz, 1H), 7.77 (td, <i>J</i> = 7.8, 7.7, 1.8 Hz, 1H), 7.51 (m, 5H), 7.24 (ddd, <i>J</i> = 7.6, 4.8, 1.1 Hz, 1H), 6.96 (s, 1H), 4.49 (s, 2H), 1.43 (s, 6H).
157	H-40		2-([1-(2-chlorophenyl)-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 30% ACN up to 57% in 7 min LC-MS: (ES, <i>m/z</i>): 381. ¹ H-NMR: (DMSO, <i>ppm</i>) δ : 8.02 (d, <i>J</i> = 1.1 Hz, 1H), 7.89 – 7.75 (m, 1H), 7.73 – 7.47 (m, 5H), 7.34 (d, <i>J</i> = 1.4 Hz, 1H), 6.95 (dt, <i>J</i> = 8.4, 1.4 Hz, 1H), 4.44 – 4.22 (m, 2H), 3.88 (d, <i>J</i> = 1.4 Hz, 3H), 1.24 (td, <i>J</i> = 7.1, 1.4 Hz, 3H).
158	H-27		2-([1-(2-Fluorophenyl)-5-[3-(oxetan-3-ylmethoxy)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XBridge: 25% ACN up to 60% in 7 min LC-MS0: (ES, <i>m/z</i>): 441. ¹ H NMR: (DMSO) δ : 12.71 (s, 1H), 7.63 – 7.47 (m, 2H), 7.40 – 7.29 (m, 2H), 7.22 (t, <i>J</i> = 8.0 Hz, 1H), 6.91 (ddd, <i>J</i> = 8.4, 2.6, 1.0 Hz, 1H), 6.85 – 6.68 (m, 3H), 4.67 (dd, <i>J</i> = 7.9, 6.0 Hz, 2H), 4.49 (s, 2H), 4.35 (t, <i>J</i> = 6.0 Hz, 2H), 4.09 (d, <i>J</i> = 6.8 Hz, 2H), 3.34 – 3.23 (m, 1H), 1.43 (s, 6H).





159	H-28		2-([5-(3-Cyclobutoxyphenyl)-1-(2-fluorophenyl)-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 47% ACN up to 67% in 7 min LC-MS: (ES, <i>m/z</i>): 425. ¹H NMR: (400 MHz, DMSO-<i>d</i>₆) δ 7.65 – 7.48 (m, 2H), 7.41 – 7.29 (m, 2H), 7.23 (t, <i>J</i> = 8.0 Hz, 1H), 6.93 – 6.74 (m, 2H), 6.67 (s, 1H), 6.57 (dd, <i>J</i> = 2.5, 1.5 Hz, 1H), 4.48 (s, 2H), 4.43 (q, <i>J</i> = 7.1 Hz, 1H), 2.23 (dddd, <i>J</i> = 9.2, 7.8, 6.5, 2.6 Hz, 2H), 1.91 (qdd, <i>J</i> = 9.8, 7.7, 2.7 Hz, 2H), 1.78 – 1.66 (m, 1H), 1.64 – 1.50 (m, 1H), 1.43 (s, 6H).
160	H-29		2-([1-(2-Fluorophenyl)-5-[3-(oxetan-3-yl-oxy)phenyl]-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 28% ACN up to 54% in 7 min LC-MS: (ES, <i>m/z</i>): 427. ¹H NMR: (300 MHz, DMSO) δ 7.58 (td, <i>J</i> = 8.0, 7.8, 1.9 Hz, 1H), 7.51 (m, 1H), 7.36 (m, 2H), 7.27 (m, 1H), 6.88 (dt, <i>J</i> = 7.7, 1.2, 1.2 Hz, 1H), 6.77 (ddd, <i>J</i> = 8.3, 2.6, 0.9 Hz, 1H), 6.68 (s, 1H), 6.49 (dd, <i>J</i> = 2.6, 1.5 Hz, 1H), 5.09 (m, 1H), 4.73 (m, 2H), 4.48 (s, 2H), 4.37 (m, 2H), 1.42 (s, 6H).
161	H-5		2-([5-(3-Cyclobutoxyphenyl)-1-(2-methylphenyl)-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 48% ACN up to 69% in 7 min LC-MS: (ES, <i>m/z</i>): 421. ¹H NMR: (400 MHz, DMSO-<i>d</i>₆) δ 12.70 (s, 1H), 7.45 – 7.25 (m, 4H), 7.20 (t, <i>J</i> = 8.0 Hz, 1H), 6.90 – 6.81 (m, 1H), 6.75 (dd, <i>J</i> = 8.2, 2.4 Hz, 1H), 6.66 (s, 1H), 6.55 – 6.45 (m, 1H), 4.48 (s, 2H), 4.34 (p, <i>J</i> = 7.2 Hz, 1H), 2.22 (dddt, <i>J</i> = 11.6, 8.7, 5.9, 2.4 Hz, 2H), 1.92 (ddd, <i>J</i> = 9.8, 7.6, 2.7 Hz, 2H), 1.88 (s, 3H), 1.71 (dd, <i>J</i> = 11.7, 8.5 Hz, 1H), 1.57 (tt, <i>J</i> = 10.1, 8.1 Hz, 1H), 1.43 (s, 6H).
162	H-30		2-([5-(3,5-Diethoxyphenyl)-1-(2-methylphenyl)-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 48% ACN up to 75% in 7 min LC-MS: (ES, <i>m/z</i>): 439.15. ¹H NMR (300 MHz, DMSO) δ 12.70 (s, 1H), 7.32 (m, 4H), 6.68 (s, 1H), 6.35 (t, <i>J</i> = 2.2, 2.2 Hz, 1H), 6.28 (d, <i>J</i> = 2.2 Hz, 2H), 4.48 (s, 2H), 3.81 (q, <i>J</i> = 6.9, 6.9, 6.9 Hz, 4H), 1.87 (s, 3H), 1.42 (s, 6H), 1.19 (t, <i>J</i> = 6.9, 6.9 Hz, 6H).

163	H-82		2-([1-[2-(dimethylamino)phenyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 35% ACN up to 65% in 7 min LC-MS (ES, m/z): 448.20. ^1H NMR (300 MHz, DMSO) δ 8.16 (s, 0.12H), 7.97 (d, $J = 0.9$ Hz, 1H), 7.60 (dd, $J = 8.4, 0.8$ Hz, 1H), 7.32 (m, 3H), 6.98 (m, 2H), 6.85 (dd, $J = 8.2, 1.3$ Hz, 1H), 6.73 (s, 1H), 4.52 (s, 2H), 4.25 (q, $J = 7.1, 7.1, 7.1$ Hz, 2H), 2.13 (s, 6H), 1.43 (s, 6H), 1.20 (t, $J = 7.2, 7.2$ Hz, 3H).
164	H-31		2-([5-(3,5-Dimethoxyphenyl)-1-(2-ethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 35% ACN up to 65% in 7 min LC-MS: (ES, m/z): 441. ^1H NMR: (300 MHz, MeOH- d_4) δ 7.44 (td, $J = 7.6, 1.5$ Hz, 2H), 7.15 – 6.99 (m, 2H), 6.68 (s, 1H), 6.44 – 6.35 (m, 3H), 4.61 (s, 2H), 3.82 (d, $J = 18.7$ Hz, 2H), 3.62 (s, 6H), 1.55 (s, 6H), 1.06 (t, $J = 7.0$ Hz, 3H).
165	H-6		2-([5-(1-Ethyl-1H-indazol-6-yl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 30% ACN up to 53% in 7 min LC-MS: (ES, m/z): 459. ^1H NMR (300 MHz, DMSO- d_6) δ 12.75 (s, 1H), 8.16 (s, 1H), 7.96 (d, $J = 0.9$ Hz, 1H), 7.89 (dd, $J = 8.2, 1.0$ Hz, 1H), 7.63 – 7.51 (m, 2H), 7.45 (dd, $J = 7.3, 1.0$ Hz, 1H), 7.17 (dd, $J = 8.1, 7.3$ Hz, 1H), 6.94 (dd, $J = 8.4, 1.4$ Hz, 1H), 6.89 (s, 1H), 4.56 (s, 2H), 4.25 (q, $J = 7.1$ Hz, 2H), 3.47 (s, 3H), 1.45 (s, 6H), 1.16 (t, $J = 7.1$ Hz, 3H).
166	H-7		2-([5-(3-Cyclobutoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 53% ACN up to 78% in 7 min LC-MS: (ES, m/z): 461. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.71 (s, 1H), 8.22 (s, 1H), 7.94 (dd, $J = 8.1, 1.0$ Hz, 1H), 7.39 (dd, $J = 7.3, 1.0$ Hz, 1H), 7.27 – 7.13 (m, 2H), 6.90 (dt, $J = 7.8, 1.2$ Hz, 1H), 6.80 (s, 1H), 6.72 (ddd, $J = 8.3, 2.6, 1.0$ Hz, 1H), 6.50 – 6.42 (m, 1H), 4.54 (s, 2H), 4.14 (p, $J = 7.0$ Hz, 1H), 3.45 (s, 3H), 2.01 (d, $J = 9.2$ Hz, 2H), 1.89 – 1.74 (m, 2H), 1.74 – 1.59 (m, 1H), 1.57 – 1.47 (m, 1H), 1.44 (s, 6H).





167	H-8		2-([5-[3-(2,2-dimethylpropoxy)phenyl]-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 80% in 7 min LC-MS: (ES, m/z): 477.20. ^1H NMR: (300 MHz, DMSO) δ 8.18 (s, 1H), 7.91 (dd, J = 8.1, 1.0 Hz, 1H), 7.39 (dd, J = 7.3, 1.0 Hz, 1H), 7.16 (m, 2H), 6.80 (m, 3H), 6.63 (dd, J = 2.5, 1.5 Hz, 1H), 4.54 (s, 2H), 3.43 (s, 3H), 3.17 (s, 2H), 1.44 (s, 6H), 0.87 (s, 9H).
168	H-9		2-([5-[3-(cyclobutylmethoxy)phenyl]-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 80% in 8 min LC-MS: (ES, m/z): 475. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.73 (s, 1H), 8.20 (s, 1H), 7.93 (dd, J = 8.1, 1.0 Hz, 1H), 7.40 (dd, J = 7.3, 1.0 Hz, 1H), 7.30 – 7.10 (m, 2H), 6.94 – 6.76 (m, 3H), 6.67 (dd, J = 2.5, 1.6 Hz, 1H), 4.54 (s, 2H), 3.57 (d, J = 6.8 Hz, 2H), 3.44 (s, 3H), 2.46 (d, J = 7.4 Hz, 1H), 2.05 – 1.76 (m, 4H), 1.74 – 1.55 (m, 2H), 1.44 (s, 6H).
169	H-32		2-([5-[3-(Cyclopropylmethoxy)phenyl]-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 40% ACN up to 70% in 7 min LC-MS: (ES, m/z): 461. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.71 (s, 1H), 8.19 (s, 1H), 7.91 (dd, J = 8.1, 1.0 Hz, 1H), 7.40 (dd, J = 7.3, 1.0 Hz, 1H), 7.28 – 7.04 (m, 2H), 6.80 (d, J = 8.0 Hz, 3H), 6.72 – 6.58 (m, 1H), 4.54 (s, 2H), 3.48 (d, J = 7.0 Hz, 2H), 3.44 (s, 3H), 1.44 (s, 6H), 1.00 (dddd, J = 15.0, 10.1, 5.1, 2.3 Hz, 1H), 0.55 – 0.43 (m, 2H), 0.25 – 0.11 (m, 2H).
170	H-10		2-methyl-2-([1-(1-methyl-1H-indazol-7-yl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy]propanoic acid XSelect: 50% ACN up to 80% in 7 min LC-MS: (ES, m/z): 463. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.73 (s, 1H), 8.18 (s, 1H), 7.91 (dd, J = 8.1, 1.0 Hz, 1H), 7.39 (dd, J = 7.3, 1.0 Hz, 1H), 7.26 – 7.05 (m, 2H), 6.94 – 6.73 (m, 3H), 6.63 (dd, J = 2.5, 1.6 Hz, 1H), 4.53 (s, 2H), 3.43 (s, 3H), 3.34 (d, J = 6.7 Hz, 2H), 1.79 (hept, J = 6.7 Hz, 1H), 1.43 (s, 6H), 0.84 (d, J = 6.7 Hz, 6H).

171	H-34		2-([5-(3,5-Diethoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 81% in 7 min LC-MS: (ES, m/z): 479.15. ^1H NMR: (300 MHz, DMSO) δ 12.70 (m, 1H), 8.18 (s, 1H), 7.91 (d, $J = 8.1$ Hz, 1H), 7.44 (d, $J = 7.2$ Hz, 1H), 7.20 (t, $J = 7.7$, 7.7 Hz, 1H), 6.81 (s, 1H), 6.31 (s, 3H), 4.53 (s, 2H), 3.73 (q, $J = 7.0$, 6.9, 6.9 Hz, 4H), 3.41 (s, 3H), 1.44 (s, 6H), 1.13 (t, $J = 6.9$, 6.9 Hz, 6H).
172	H-11		2-([5-[3-(2,2-Dimethylpropoxy)phenyl]-1-(1-ethyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 85% in 7 min LC-MS: (ES, m/z): 491. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.67 (s, 1H), 8.24 (s, 1H), 7.92 (dd, $J = 8.1$, 1.1 Hz, 1H), 7.32 (dd, $J = 7.3$, 1.1 Hz, 1H), 7.24 – 6.98 (m, 2H), 6.85 (s, 1H), 6.79 (dddd, $J = 7.2$, 6.4, 2.5, 1.0 Hz, 2H), 6.65 – 6.55 (m, 1H), 4.53 (s, 2H), 3.71 (d, $J = 36.9$ Hz, 2H), 3.15 (s, 2H), 1.43 (s, 6H), 1.09 (t, $J = 7.1$ Hz, 3H), 0.86 (s, 9H).
173	H-37		2-([1-(2-Chlorophenyl)-5-[3-(oxetan-3-ylmethoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 25% ACN up to 60% in 7 min LC-MS: (ES, m/z): 457.0. ^1H -NMR: δ_{H} (300 MHz, DMSO- d_6) 12.70 (s, 1H), 7.64 – 7.45 (m, 4H), 7.20 (t, $J = 8.0$ Hz, 1H), 6.88 (ddd, $J = 8.4$, 2.5, 1.0 Hz, 1H), 6.82 – 6.69 (m, 3H), 4.67 (dd, $J = 7.9$, 6.0 Hz, 2H), 4.49 (s, 2H), 4.35 (t, $J = 6.0$ Hz, 2H), 4.06 (d, $J = 6.8$ Hz, 2H), 3.34 – 3.22 (m, 1H), 1.43 (s, 6H).
174	H-39		2-([1-(2-Chlorophenyl)-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 25% ACN up to 58% in 7 min LC-MS: (ES, m/z): 425. ^1H NMR: (400 MHz, DMSO- d_6) δ 12.75 (s, 1H), 8.01 (d, $J = 1.0$ Hz, 1H), 7.66 – 7.60 (m, 2H), 7.58 (ddd, $J = 4.0$, 3.0, 1.9 Hz, 2H), 7.54 – 7.46 (m, 2H), 6.85 (dd, $J = 8.4$, 1.4 Hz, 1H), 6.79 (s, 1H), 4.52 (s, 2H), 3.95 (s, 3H), 1.45 (s, 6H).

175	H-41		2-([1-(2-Chlorophenyl)-5-(3-cyclobutoxy-phenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 47% ACN up to 67% in 7 min LC-MS: (ES, m/z): 441. ^1H-NMR: ^1H NMR (400 MHz, DMSO-d_6) δ 12.71 (s, 1H), 7.60 (ddd, $J = 12.7, 7.5, 2.0$ Hz, 2H), 7.56 – 7.46 (m, 2H), 7.21 (t, $J = 8.0$ Hz, 1H), 6.84 (dt, $J = 7.7, 1.2$ Hz, 1H), 6.77 (ddd, $J = 8.3, 2.5, 1.0$ Hz, 1H), 6.68 (s, 1H), 6.53 (t, $J = 2.0$ Hz, 1H), 4.48 (s, 2H), 4.38 (p, $J = 7.2$ Hz, 1H), 2.31 – 2.18 (m, 2H), 1.91 (qdd, $J = 9.6, 7.5, 2.8$ Hz, 2H), 1.73 (qt, $J = 10.1, 2.7$ Hz, 1H), 1.65 – 1.50 (m, 1H), 1.43 (s, 6H).
176	H-42		2-([1-(2-Chlorophenyl)-5-[3-(oxetan-3-yl-oxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 25% ACN up to 60% in 7 min LC-MS: (ES, m/z): 443. ^1H NMR: (300 MHz, DMSO) δ 12.72 (s, 1H), 7.60 (ddd, $J = 6.2, 4.2, 2.7$ Hz, 2H), 7.51 (ddd, $J = 6.3, 3.5, 2.2$ Hz, 2H), 7.24 (t, $J = 8.0, 8.0$ Hz, 1H), 6.87 (dt, $J = 7.8, 1.1, 1.1$ Hz, 1H), 6.74 (ddd, $J = 8.2, 2.6, 0.9$ Hz, 1H), 6.68 (s, 1H), 6.45 (dd, $J = 2.6, 1.5$ Hz, 1H), 5.06 (ddd, $J = 6.0, 4.9, 1.1$ Hz, 1H), 4.77 (m, 2H), 4.48 (s, 2H), 4.39 (m, 2H), 1.42 (s, 6H).
177	H-43		2-([1-(2-chlorophenyl)-5-[3-(2,2-dimethyl-propoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 0.1% DEA / 57.0% ACN up to 85% ACN in 7 min LC-MS: (ES, m/z): 457.05. ^1H NMR (300 MHz, DMSO) δ 12.70 (s, 1H), 7.58 (m, 2H), 7.50 (m, 2H), 7.17 (t, $J = 8.0, 8.0$ Hz, 1H), 6.83 (ddd, $J = 8.4, 2.5, 1.0$ Hz, 1H), 6.75 (dt, $J = 7.7, 1.2, 1.2$ Hz, 1H), 6.70 (d, $J = 3.1$ Hz, 2H), 4.48 (s, 2H), 3.40 (s, 2H), 1.42 (s, 6H), 0.92 (s, 9H).
178	H-44		2-([1-(2-Chlorophenyl)-5-[3-(Cyclobutyl-methoxy)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 55% up to 85% in 7 min LC-MS: (ES, m/z): 455.09. ^1H NMR: (300 MHz, DMSO) δ 12.72(s, 1H), 7.58 (ddd, $J = 6.5, 4.9, 2.5$ Hz, 2H), 7.49 (m, 2H), 7.17 (t, $J = 8.0, 8.0$ Hz, 1H), 6.82 (ddd, $J = 8.3, 2.5, 0.9$ Hz, 1H), 6.77 (m, 1H), 6.70 (d, $J = 2.9$ Hz, 2H), 4.49 (s, 2H), 3.74 (d, $J = 6.8$ Hz, 2H), 2.55 (m, 1H), 1.99





			(m, 2H), 1.85 (m, 2H), 1.73 (m, 2H), 1.43 (s, 6H).
179	H-45		2-([1-(2-Chlorophenyl)-5-(3,5-dimethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 38% ACN up to 57% in 7 min LC-MS: (ES, <i>m/z</i>): 431. ¹ H NMR: (300 MHz, MeOH- <i>d</i> ₄) δ 7.61 – 7.42 (m, 4H), 6.73 (s, 1H), 6.38 (dd, <i>J</i> = 9.8, 2.2 Hz, 3H), 4.61 (s, 2H), 3.63 (s, 6H), 1.53 (s, 6H).
180	H-12		2-([1-(2-Chlorophenyl)-5-(3,5-diethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 42% ACN up to 68% in 7 min LC-MS: (ES, <i>m/z</i>): 459.05. ¹ H NMR: (300 MHz, DMSO) δ 12.74 (s, 1H), δ 7.51 (m, 4H), 6.70 (s, 1H), 6.38 (t, <i>J</i> = 2.3, 2.3 Hz, 1H), 6.30 (d, <i>J</i> = 2.2 Hz, 2H), 4.48 (s, 2H), 3.84 (q, <i>J</i> = 7.0, 7.0, 7.0 Hz, 4H), 1.42 (s, 6H), 1.22 (t, <i>J</i> = 7.0, 7.0 Hz, 6H).
181	H-13		2-([5-(3-cyclopropoxyphenyl)-1-(2-fluorophenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoate SunFire: 69% ACN up to 71% in 8 min LC-MS: (ES, <i>m/z</i>): 411.3. ¹ H-NMR: δ _H (400 MHz, DMSO- <i>d</i> ₆) 12.73 (1 H, s), 7.54 (2 H, m), 7.35 (2 H, m), 7.26 (1 H, t, <i>J</i> = 7.9, 7.9 Hz), 6.95 (1 H, ddd, <i>J</i> = 8.3, 2.5, 1.0 Hz), 6.88 (2 H, m), 6.68 (1 H, s), 4.48 (2 H, s), 3.62 (1 H, tt, <i>J</i> = 6.0, 6.0, 3.0, 3.0 Hz), 1.43 (6 H, s), 0.62 (2 H, m), 0.50 (2 H, m).
182	H-46		2-([1-(2-Bromophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 38% ACN up to 70% in 7 min LC-MS: (ES, <i>m/z</i>): 471. ¹ H NMR: (DMSO) δ: 12.71 (s, 1H), 7.81 – 7.71 (m, 1H), 7.59 – 7.37 (m, 3H), 7.23 (t, <i>J</i> = 8.0 Hz, 1H), 6.96 – 6.78 (m, 3H), 6.67 (s, 1H), 4.47 (s, 2H), 3.55 (tt, <i>J</i> = 6.0, 2.9 Hz, 1H), 1.42 (s, 6H), 0.71 – 0.43 (m, 4H).

183	H-149		2-([1-[2-(Azetidin-1-yl)-4-fluorophenyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XBridge: 10% ACN up to 52% in 7 min LC-MS: (ES, m/z): 466.20. ^1H NMR (300 MHz, DMSO) δ 7.26 (t, J = 8.2, 8.2 Hz, 1H), 7.00 (m, 3H), 6.92 (ddd, J = 8.2, 2.4, 1.1 Hz, 1H), 6.63 (s, 1H), 6.45 (td, J = 8.4, 8.4, 2.8 Hz, 1H), 6.29 (dd, J = 11.3, 2.8 Hz, 1H), 4.46 (s, 2H), 3.62 (tt, J = 6.1, 6.1, 3.0, 3.0 Hz, 1H), 3.49 (m, 4H), 2.07 (p, J = 7.2, 7.2, 7.2, 7.2 Hz, 2H), 1.39 (s, 6H), 0.69 (m, 2H), 0.56 (m, 2H).
184	H-48		2-([5-(3-Cyclopropoxyphenyl)-1-(2,4-dichlorophenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 77% in 7 min LC-MS: (ES, m/z): 461.00. ^1H NMR (300 MHz, DMSO) δ 12.72 (s, 1H), 7.81 (d, J = 2.2 Hz, 1H), 7.60 (m, 2H), 7.25 (t, J = 7.9, 7.9 Hz, 1H), 6.89 (m, 3H), 6.68 (s, 1H), 4.48 (s, 2H), 3.63 (tt, J = 6.1, 6.1, 3.0, 3.0 Hz, 1H), 1.42 (s, 6H), 0.66 (m, 2H), 0.51 (m, 2H).
185	H-49		2-([5-(3-Cyclopropoxyphenyl)-1-(2,5-dichlorophenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 48% ACN up to 74% in 7 min LC-MS: (ES, m/z): 461.05. ^1H NMR (300 MHz, DMSO) δ 12.72 (s, 1H), 7.83 (m, 1H), 7.60 (d, J = 1.5 Hz, 2H), 7.25 (t, J = 7.9, 7.9 Hz, 1H), 6.91 (m, 3H), 6.69 (s, 1H), 4.50 (s, 2H), 3.61 (dt, J = 6.0, 3.1, 3.1 Hz, 1H), 1.43 (s, 6H), 0.65 (dd, J = 6.2, 2.0 Hz, 2H), 0.50 (tt, J = 6.2, 6.2, 3.4, 3.4 Hz, 2H).
186	H-50		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(difluoromethoxy)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XBridge: 40% ACN up to 65% in 7 min LC-MS: (ES, m/z): 459. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.72 (s, 1H), 7.59 – 7.47 (m, 2H), 7.42 – 7.13 (m, 3H), 6.95 – 6.79 (m, 3H), 6.66 (d, J = 10.0 Hz, 1H), 4.47 (s, 2H), 3.57 (tt, J = 6.1, 2.9 Hz, 1H), 1.42 (s, 6H), 0.67 – 0.42 (m, 4H).

187	H-51		2-([5-(3-cyclopropoxyphenyl)-1-(2-ethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 40% ACN up to 67% in 7 min LC-MS: (ES, <i>m/z</i>): 437. ¹ H NMR: (300 MHz, MeOH- <i>d</i> ₄) δ 7.50 – 7.37 (m, 2H), 7.26 – 7.15 (m, 1H), 7.13 – 6.99 (m, 2H), 6.97 – 6.87 (m, 3H), 6.67 (s, 1H), 4.62 (s, 2H), 3.75 (s, 2H), 3.51 (tt, <i>J</i> = 6.1, 3.0 Hz, 1H), 1.55 (s, 6H), 1.02 (t, <i>J</i> = 7.0 Hz, 3H), 0.64 (d, <i>J</i> = 6.0 Hz, 2H), 0.57 – 0.47 (m, 2H).
188	H-52		2-([5-(3-Cyclopropoxyphenyl)-1-(2-methylphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 40% ACN up to 67% in 7 min LC-MS: (ES, <i>m/z</i>): 407. ¹ H NMR: (300 MHz, MeOH- <i>d</i> ₄) δ 7.44 – 7.27 (m, 4H), 7.20 (t, <i>J</i> = 8.0 Hz, 1H), 6.92 (tt, <i>J</i> = 8.5, 1.1 Hz, 2H), 6.84 (t, <i>J</i> = 2.1 Hz, 1H), 6.73 (s, 1H), 4.61 (s, 2H), 3.46 (tt, <i>J</i> = 6.1, 3.0 Hz, 1H), 1.96 (s, 3H), 1.54 (s, 6H), 0.63 (pd, <i>J</i> = 6.0, 3.8 Hz, 2H), 0.57 – 0.48 (m, 2H).
189	H-53		2-([5-(3-cyclopropoxyphenyl)-1-[2-(propan-2-yl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 76% in 7 min LC-MS: (ES, <i>m/z</i>): 435. ¹ H-NMR: (DMSO, <i>ppm</i>): δ: 12.65 (s, 1H), 7.46 (dd, <i>J</i> = 3.7, 1.3 Hz, 2H), 7.36 – 7.16 (m, 3H), 6.94 – 6.76 (m, 3H), 6.68 (s, 1H), 4.49 (s, 2H), 3.53 (tt, <i>J</i> = 6.1, 2.9 Hz, 1H), 1.43 (s, 6H), 1.00 (s, 3H), 0.84 (s, 3H), 0.67 – 0.51 (m, 2H), 0.47 (tq, <i>J</i> = 5.5, 3.2 Hz, 2H).
190	H-54		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(trifluoromethyl)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XBridge: 40% ACN up to 63% in 7 min LC-MS: (ES, <i>m/z</i>): 461. ¹ H NMR (300 MHz, MeOH- <i>d</i> ₄) δ 7.94 – 7.83 (m, 1H), 7.79 – 7.64 (m, 2H), 7.48 – 7.37 (m, 1H), 7.20 (t, <i>J</i> = 8.0 Hz, 1H), 6.91 (dddd, <i>J</i> = 6.4, 4.0, 2.7, 1.2 Hz, 2H), 6.83 (t, <i>J</i> = 2.0 Hz, 1H), 6.73 (s, 1H), 4.61 (s, 2H), 3.49 (tt, <i>J</i> = 6.1, 3.0 Hz, 1H), 1.53 (s, 6H), 0.71 – 0.58 (m, 2H), 0.53 (ddd, <i>J</i> = 6.2, 4.3, 2.8 Hz, 2H).





191	H-55		2-([5-(3-Cyclopropoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XBridge: 38% ACN up to 62% in 7 min LC-MS: (ES, <i>m/z</i>): 447. ¹ H-NMR: (DMSO, <i>ppm</i>): δ: 12.74 (s, 1H), 8.20 (s, 1H), 7.92 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.38 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.27 – 7.13 (m, 2H), 6.99 – 6.76 (m, 4H), 4.54 (s, 2H), 3.49 (s, 3H), 3.43 – 3.34 (m, 1H), 1.44 (s, 6H), 0.52 – 0.36 (m, 4H).
192	H-14		2-([5-(3-Cyclopropoxyphenyl)-1-(1-methyl-1H-indazol-4-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 35% ACN up to 68% in 7 min LC-MS: (ES, <i>m/z</i>): 446. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 7.80 – 7.59 (m, 2H), 7.41 (dd, <i>J</i> = 8.6, 7.3 Hz, 1H), 7.21 (t, <i>J</i> = 7.9 Hz, 1H), 7.07 – 6.80 (m, 4H), 6.71 (s, 1H), 4.54 (s, 2H), 4.07 (s, 3H), 3.56 (tt, <i>J</i> = 6.1, 2.9 Hz, 1H), 1.45 (s, 6H), 0.54 – 0.45 (m, 2H), 0.43 – 0.33 (m, 2H).
193	H-15		2-([5-(3-Cyclopropoxyphenyl)-1-(1-ethyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 61% ACN up to 66% in 7 min LC-MS: (ES, <i>m/z</i>): 461. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.71 (s, 1H), 8.25 (s, 1H), 7.92 (dd, <i>J</i> = 8.1, 1.1 Hz, 1H), 7.31 (dd, <i>J</i> = 7.3, 1.1 Hz, 1H), 7.25 – 7.09 (m, 2H), 6.94 (dt, <i>J</i> = 7.7, 1.2 Hz, 1H), 6.91 – 6.81 (m, 2H), 6.78 (dd, <i>J</i> = 2.5, 1.6 Hz, 1H), 4.55 (s, 2H), 3.77 (d, <i>J</i> = 28.7 Hz, 2H), 3.41 – 3.25 (m, <i>J</i> = 3.3 Hz, 1H), 1.44 (s, 6H), 1.12 (t, <i>J</i> = 7.1 Hz, 3H), 0.57 – 0.29 (m, 4H).
194	H-56		2-([5-(3-Cyclopropoxyphenyl)-1-(1-methyl-1H-1,3-benzodiazol-7-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 36% ACN up to 55% in 7 min LC-MS: (ES, <i>m/z</i>): 447.10. ¹ H NMR (300 MHz, DMSO) δ 12.72 (s, 1H), 8.18 (s, 1H), 7.78 (dd, <i>J</i> = 7.6, 1.5 Hz, 1H), 7.22 (m, 3H), 6.89 (m, 2H), 6.79 (t, <i>J</i> = 2.1, 2.1 Hz, 1H), 6.77 (s, 1H), 4.52 (s, 2H), 3.41 (dq, <i>J</i> = 6.1, 3.1, 3.0, 3.0 Hz, 1H), 3.29 (s, 3H), 1.43 (s, 6H), 0.47 (d, <i>J</i> = 6.1 Hz, 2H), 0.39 (m, 2H).

195	H-16		2-([5-(1-Benzothiophen-2-yl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 80% in 7 min LC-MS: (ES, <i>m/z</i>): 441. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 12.68 (s, 1H), 8.02 – 7.95 (m, 1H), 7.90 – 7.81 (m, 1H), 7.57 (d, <i>J</i> = 0.8 Hz, 1H), 7.52 – 7.45 (m, 1H), 7.44 – 7.36 (m, 2H), 7.31 (pd, <i>J</i> = 7.4, 1.8 Hz, 2H), 6.77 – 6.70 (m, 1H), 6.68 (s, 1H), 5.62 (s, 2H), 4.45 (s, 2H), 1.42 (s, 6H).
196	H-57		2-([5-(3-Chloro-5-methoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 76% in 7 min LC-MS: (ES, <i>m/z</i>): 459. ¹ H-NMR-PH-VET-168-0: (DMSO): δ: 12.62 (s, 1H), 7.24 (ddd, <i>J</i> = 8.5, 7.4, 1.8 Hz, 1H), 7.11 – 6.81 (m, 5H), 6.76 (dd, <i>J</i> = 7.6, 1.7 Hz, 1H), 6.48 (s, 1H), 5.27 (s, 2H), 4.41 (s, 2H), 3.99 (q, <i>J</i> = 6.9 Hz, 2H), 3.32 (s, 1H), 1.40 (s, 6H), 1.22 (t, <i>J</i> = 6.9 Hz, 3H).
197	H-17		2-([1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1H-indazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 71% in 7 min LC-MS: (ES, <i>m/z</i>): 438. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.66 (s, 1H), 7.93 (d, <i>J</i> = 1.0 Hz, 1H), 7.70 (dt, <i>J</i> = 8.5, 0.9 Hz, 1H), 7.44 (dd, <i>J</i> = 8.5, 7.1 Hz, 1H), 7.39 – 7.30 (m, 1H), 7.30 – 7.20 (m, 2H), 7.09 (dd, <i>J</i> = 7.1, 0.8 Hz, 1H), 6.88 – 6.69 (m, 1H), 6.57 (s, 1H), 5.38 (s, 2H), 4.47 (s, 2H), 4.07 (s, 3H), 1.41 (s, 6H).
198	H-18		2-([1-[(2-Ethoxyphenyl)methyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 65% in 7 min LC-MS: (ES, <i>m/z</i>): 463. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.70 (s, 1H), 8.09 (d, <i>J</i> = 0.9 Hz, 1H), 7.79 (dd, <i>J</i> = 8.3, 0.8 Hz, 1H), 7.68 (q, <i>J</i> = 1.1 Hz, 1H), 7.30 – 7.11 (m, 2H), 6.95 (dd, <i>J</i> = 8.3, 1.1 Hz, 1H), 6.86 (td, <i>J</i> = 7.5, 1.1 Hz, 1H), 6.72 (dd, <i>J</i> = 7.6, 1.7 Hz, 1H), 6.51 (s, 1H), 5.33 (s, 2H), 4.53 – 4.32 (m, 4H), 3.93 (q, <i>J</i> = 6.9 Hz, 2H), 1.42 (s, 6H), 1.33 (t, <i>J</i> = 7.2 Hz, 3H), 1.13 (t, <i>J</i> = 6.9 Hz, 3H).





199	H-19		2-([5-(3-Ethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 80% in 7 min LC-MS: (ES, <i>m/z</i>): 439. ¹H NMR: (300 MHz, DMSO-<i>d</i>₆) δ 12.68 (s, 1H), 7.33 (t, <i>J</i> = 7.9 Hz, 1H), 7.23 (ddd, <i>J</i> = 8.5, 7.5, 1.7 Hz, 1H), 7.03 – 6.91 (m, 3H), 6.91 – 6.82 (m, 2H), 6.69 (dd, <i>J</i> = 7.6, 1.7 Hz, 1H), 6.42 (s, 1H), 5.26 (s, 2H), 4.42 (s, 2H), 3.96 (dq, <i>J</i> = 20.8, 6.9 Hz, 4H), 1.40 (s, 6H), 1.25 (dt, <i>J</i> = 12.7, 6.9 Hz, 6H).
200	H-58		2-([5-(3-Ethoxy-5-methoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 52% ACN up to 71% in 7 min LC-MS: (ES, <i>m/z</i>): 469. ¹H-NMR: (DMSO, <i>ppm</i>): δ: 12.63 (s, 1H), 7.24 (ddd, <i>J</i> = 8.6, 7.4, 1.7 Hz, 1H), 6.98 (dd, <i>J</i> = 8.3, 1.0 Hz, 1H), 6.87 (td, <i>J</i> = 7.5, 1.0 Hz, 1H), 6.67 (dd, <i>J</i> = 7.5, 1.6 Hz, 1H), 6.54 – 6.40 (m, 4H), 5.27 (s, 2H), 4.41 (s, 2H), 3.96 (dq, <i>J</i> = 28.8, 7.0 Hz, 4H), 3.67 (s, 3H), 1.40 (s, 6H), 1.25 (td, <i>J</i> = 6.9, 4.8 Hz, 6H).
201	H-59		2-([5-(3-Cyclopropoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 42% ACN up to 75% in 7 min LC-MS: (ES, <i>m/z</i>): 451. ¹H NMR (300 MHz, DMSO-<i>d</i>₆) δ 12.68 (s, 1H), 7.35 (ddd, <i>J</i> = 8.2, 7.0, 1.1 Hz, 1H), 7.29 – 7.17 (m, 1H), 7.01 (tdd, <i>J</i> = 13.0, 8.6, 1.4 Hz, 4H), 6.87 (td, <i>J</i> = 7.5, 1.1 Hz, 1H), 6.68 (dd, <i>J</i> = 7.6, 1.7 Hz, 1H), 6.43 (s, 1H), 5.27 (s, 2H), 4.42 (s, 2H), 4.00 (q, <i>J</i> = 6.9 Hz, 2H), 3.69 (tt, <i>J</i> = 6.3, 3.1 Hz, 1H), 1.40 (s, 6H), 1.23 (t, <i>J</i> = 6.9 Hz, 3H), 0.71 – 0.51 (m, 4H).
202	H-150		2-([5-(3-Cyclopropoxyphenyl)-1-[[2-(pyrrolidin-1-yl)phenyl]methyl]-1<i>H</i>-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 28% ACN up to 58% in 7 min LC-MS: (ES, <i>m/z</i>): 476.15. ¹H NMR: (400 MHz, DMSO) δ 7.31 (t, <i>J</i> = 7.9, 7.9 Hz, 1H), 7.15 (m, 1H), 7.00 (m, 2H), 6.94 (m, 2H), 6.85 (t, <i>J</i> = 7.6, 7.6 Hz, 1H), 6.60 (dd, <i>J</i> = 7.6, 1.8 Hz, 1H), 6.46 (s, 1H), 5.31 (s, 2H), 4.45 (s, 2H), 3.64 (m, 1H), 3.03 (m, 4H), 1.84 (m, 4H), 1.39 (s, 6H), 0.60 (dq, <i>J</i> = 5.3, 3.2, 2.4, 2.4 Hz, 2H), 0.55 (dq, <i>J</i> = 7.8, 3.6, 3.6, 3.4 Hz, 2H).

203	H-60		2-([1-[(2-Chlorophenyl)methyl]-5-(2,3-dihydro-1,4-benzodioxin-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 45% ACN up to 58% in 7 min LC-MS: (ES, m/z): 443. ^1H NMR: (300 MHz, MeOD) δ 7.39 (m, 1H), 7.25 (tt, $J = 7.4, 7.4, 5.6, 5.6$ Hz, 2H), 6.85 (m, 1H), 6.77 (d, $J = 7.8$ Hz, 2H), 6.68 (m, 1H), 6.46 (s, 1H), 5.40 (s, 2H), 4.56 (s, 2H), 4.24 (q, $J = 1.1, 1.0, 1.0$ Hz, 4H), 1.51 (s, 6H).
204	H-61		[[5-(2H-1,3-Benzodioxol-5-yl)-1-[(2-chlorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 45% ACN up to 60% in 7 min LC-MS: (ES, m/z): 429. ^1H NMR: (DMSO) δ : 12.67 (s, 1H), 7.49 – 7.38 (m, 1H), 7.36 – 7.21 (m, 2H), 7.00 – 6.91 (m, 2H), 6.83 (dd, $J = 7.9, 1.8$ Hz, 1H), 6.76 – 6.66 (m, 1H), 6.37 (s, 1H), 6.05 (s, 2H), 5.35 (s, 2H), 4.39 (s, 2H), 1.38 (s, 6H)
205	H-62		2-([1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1H-pyrazol-4-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 25% ACN up to 52% in 7 min ^1H NMR (DMSO) δ : 7.93 (s, 1H), 7.55 – 7.46 (m, 2H), 7.29 (dtd, $J = 21.6, 7.5, 1.6$ Hz, 2H), 6.59 (dd, $J = 7.6, 1.7$ Hz, 1H), 6.43 (s, 1H), 5.43 (s, 2H), 4.38 (s, 2H), 3.83 (s, 3H), 1.39 (s, 6H)
206	H-21		2-([1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1H-indol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 45% ACN up to 80% in 7 min LC-MS: (ES, m/z): 437. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.68 (s, 1H), 7.57 (dd, $J = 8.2, 0.7$ Hz, 1H), 7.48 – 7.36 (m, 3H), 7.34 – 7.26 (m, 2H), 7.01 (dd, $J = 8.2, 1.5$ Hz, 1H), 6.83 – 6.75 (m, 1H), 6.51 – 6.42 (m, 2H), 5.42 (s, 2H), 4.43 (s, 2H), 3.72 (s, 3H), 1.41 (s, 6H).
207	H-22		2-([1-[(2-Ethoxyphenyl)methyl]-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 37% ACN up to 65% in 7 min LC-MS: (ES, m/z): 449. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.68 (s, 1H), 8.07 (d, $J = 1.0$ Hz, 1H), 7.78 (d, $J = 8.3$ Hz, 1H), 7.67 (d, $J = 1.3$ Hz, 1H), 7.29 – 7.10 (m, 2H), 7.01 – 6.81 (m, 2H), 6.74 (dd, $J = 7.6, 1.7$ Hz, 1H), 6.48 (s, 1H), 5.31 (s,

			2H), 4.43 (s, 2H), 4.00 (s, 3H), 3.91 (q, $J = 6.9$ Hz, 2H), 1.40 (s, 6H), 1.11 (t, $J = 6.9$ Hz, 3H).
208	H-63		2-([5-(3,5-Diethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 45% ACN up to 80% in 7 min LC-MS: (ES, m/z): 483.15. ^1H NMR (300 MHz, DMSO) δ 7.22 (m, 1H), 6.97 (m, 1H), 6.86 (td, $J = 7.4, 7.4, 1.1$ Hz, 1H), 6.66 (dd, $J = 7.6, 1.7$ Hz, 1H), 6.45 (m, 3H), 6.42 (s, 1H), 5.26 (s, 2H), 4.40 (s, 2H), 4.00 (q, $J = 7.0, 6.9, 6.9$ Hz, 2H), 3.90 (q, $J = 7.0, 7.0, 7.0$ Hz, 4H), 1.39 (s, 6H), 1.24 (td, $J = 7.0, 7.0, 2.6$ Hz, 9H).
209	H-66		2-([1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1H-indazol-5-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 35% ACN up to 65% in 7 min LC-MS: (ES, m/z): 439. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.71 (s, 1H), 8.07 (d, $J = 0.9$ Hz, 1H), 7.79 (d, $J = 1.4$ Hz, 1H), 7.70 (dt, $J = 8.8, 1.0$ Hz, 1H), 7.49 – 7.36 (m, 2H), 7.32 – 7.22 (m, 2H), 6.76 (dd, $J = 5.7, 3.7$ Hz, 1H), 6.44 (s, 1H), 5.39 (s, 2H), 4.42 (s, 2H), 4.05 (s, 3H), 1.39 (s, 6H).
210	H-67		2-([5-(3-cyclobutoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 50% ACN up to 80% in 7 min LC-MS: (ES, m/z): 465. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.68 (s, 1H), 7.38 – 7.20 (m, 2H), 7.05 – 6.94 (m, 2H), 6.92 – 6.82 (m, 2H), 6.75 – 6.62 (m, 2H), 6.42 (s, 1H), 5.26 (s, 2H), 4.50 (q, $J = 7.1$ Hz, 1H), 4.42 (s, 2H), 4.02 (q, $J = 6.9$ Hz, 2H), 2.30 – 2.15 (m, 2H), 2.03 – 1.86 (m, 2H), 1.79 – 1.65 (m, 1H), 1.63 – 1.47 (m, 1H), 1.40 (s, 6H), 1.24 (t, $J = 6.9$ Hz, 3H).
211	H-71		2-([5-(3,5-Dimethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 40% ACN up to 68% in 7 min LC-MS: (ES, m/z): 455. ^1H NMR: (300 MHz, MeOH- d_4) δ 7.24 (td, $J = 7.8, 7.4, 1.7$ Hz, 1H), 6.99 – 6.81 (m, 2H), 6.69 (dd, $J = 7.6, 1.6$ Hz, 1H), 6.54 – 6.41 (m, 4H), 5.33 (s, 2H), 4.57 (s, 2H), 4.04 (q, J

			= 7.0 Hz, 2H), 3.64 (s, 6H), 1.52 (s, 6H), 1.34 (t, $J = 7.0$ Hz, 3H).
212	H-23		2-([5-(3,5-Dimethoxyphenyl)-1-(2-phenylethyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 40% ACN up to 66% in 7 min LC-MS: (ES, m/z): 425. ^1H NMR: (300 MHz, DMSO- d_6) δ 12.67 (s, 1H), 7.32 – 7.11 (m, 3H), 7.06 – 6.92 (m, 2H), 6.53 (t, $J = 2.3$ Hz, 1H), 6.37 (d, $J = 2.3$ Hz, 2H), 6.26 (s, 1H), 4.43 (s, 2H), 4.34 – 4.20 (m, 2H), 3.74 (s, 6H), 3.13 – 2.99 (m, 2H), 1.41 (s, 6H).
213	H-74		2-([1-[(2-Chlorophenyl)methyl]-5-(3,5-diethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 53% ACN up to 76% in 7 min LC-MS: (ES, m/z): 473.10. ^1H NMR: (300 MHz, DMSO) δ 12.63 (s, 1H), 7.45 (m, 1H), 7.30 (m, 2H), 6.72 (m, 1H), 6.45 (m, 4H), 5.38 (s, 2H), 4.41 (s, 2H), 3.91 (q, $J = 7.0, 7.0, 7.0$ Hz, 4H), 1.39 (s, 6H), 1.24 (t, $J = 6.9, 6.9$ Hz, 6H).
214	H-111		2-([5-(3-Cyclopropoxyphenyl)-1-[[2-(dimethylamino)phenyl]methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 67% ACN up to 68% in 7 min LC-MS: (ES, m/z): 450. ^1H -NMR: (CDCl ₃ , ppm): ^1H NMR (300 MHz, MeOD) δ 7.25 (m, 2H), 7.17 (dd, $J = 8.1, 1.4$ Hz, 1H), 7.00 (m, 2H), 6.90 (m, 2H), 6.74 (dd, $J = 7.6, 1.5$ Hz, 1H), 6.51 (s, 1H), 5.43 (s, 2H), 4.59 (s, 2H), 3.55 (m, 1H), 2.60 (s, 6H), 1.53 (s, 6H), 0.59 (m, 4H).
215	H-99		2-([5-(3-Methoxyphenyl)-1-[2-(propan-2-yloxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 38% ACN up to 61% in 7 min LC-MS: (ES, m/z): 425. ^1H NMR (300 MHz, DMSO) δ 7.46 (m, 1H), 7.38 (t, $J = 7.9, 7.9$ Hz, 1H), 7.18 (t, $J = 7.9, 7.9$ Hz, 1H), 7.03 (m, 2H), 6.81 (m, 2H), 6.68 (m, 1H), 6.58 (s, 1H), 4.45 (s, 2H), 4.34 (m, 1H), 3.57 (s, 3H), 1.41 (s, 6H), 0.96 (s, 3H), 0.59 (s, 3H).




216	H-96		2-Methyl-2-([5-[3-(2-methylpropoxy)-phenyl]-1-phenyl-1H-pyrazol-3-yl]-methoxy)propanoic acid SunFire: 69% ACN up to to 71% ACN in 8 min LC-MS: (ES, m/z): 409. H-NMR: (MeOD, ppm): δ : 7.48 – 7.32 (m, 3H), 7.35 – 7.14 (m, 3H), 6.91 – 6.78 (m, 2H), 6.73 – 6.63 (m, 2H), 4.61 (s, 2H), 3.54 (d, J = 6.5 Hz, 2H), 1.92 (hept, J = 6.7 Hz, 1H), 1.53 (s, 6H), 0.95 (d, J = 6.7 Hz, 6H).
217	H-122		(2R)-2-([5-(3-cyclopropoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect 43% ACN up to 65% in 7 min LC-MS: (ES, m/z): 461. ^1H NMR: (300 MHz, DMSO- d_6) δ : 12.67 (s, 2H), 8.18 (s, 1H), 7.91 (d, J = 8.0 Hz, 2H), 7.37 (d, J = 7.3 Hz, 2H), 7.28 – 7.12 (m, 4H), 7.03 – 6.75 (m, 8H), 4.53 (s, 4H), 3.45 (s, 7H), 2.49 (s, 1H), 1.76 (dh, J = 14.3, 7.2 Hz, 4H), 1.40 (s, 5H), 0.88 (t, J = 7.4 Hz, 6H), 0.50 – 0.35 (m, 8H).
218	H-123		(2S)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 67% ACN up to 68% in 7 min) LC-MS: (ES, m/z): 429. ^1H NMR (DMSO) δ : 12.64 (s, 1H), 7.50 – 7.22 (m, 4H), 7.02 – 6.84 (m, 3H), 6.78 – 6.69 (m, 1H), 6.45 (s, 1H), 5.38 (s, 2H), 4.41 (d, J = 1.9 Hz, 2H), 3.69 (s, 3H), 1.84 – 1.60 (m, J = 7.1 Hz, 2H), 1.36 (s, 3H), 0.85 (t, J = 7.4 Hz, 3H).
219	H-124		(2R)-2-([1-[(2-chlorophenyl)methyl]-5-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XBridge: 40% ACN up to 75% in 7 min LC-MS: (ES, m/z): 429. ^1H NMR: (400 MHz, DMSO- d_6) δ : 12.61 (s, 1H), 7.49 – 7.22 (m, 4H), 7.02 – 6.84 (m, 3H), 6.78 – 6.69 (m, 1H), 6.46 (s, 1H), 5.38 (s, 2H), 4.41 (d, J = 2.2 Hz, 2H), 3.69 (s, 3H), 1.84 – 1.61 (m, J = 7.6 Hz, 2H), 1.36 (s, 3H), 0.85 (t, J = 7.4 Hz, 3H)

220	H-130		(2 <i>R</i>)-2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1 <i>H</i> -indazol-6-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 48% ACN up to 67% in 7 min LC-MS: (ES, <i>m/z</i>): 467. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ : 12.63 (s, 1H), 8.08 (s, 1H), 7.79 (d, <i>J</i> = 8.4 Hz, 1H), 7.68 (s, 1H), 7.46 – 7.37 (m, 1H), 7.29 (q, <i>J</i> = 4.8 Hz, 2H), 7.13 (d, <i>J</i> = 8.4 Hz, 1H), 6.84 – 6.74 (m, 1H), 6.53 (s, 1H), 5.46 (s, 2H), 4.52 – 4.32 (m, 4H), 1.75 (hept, <i>J</i> = 7.0 Hz, 2H), 1.41 – 1.26 (m, 6H), 0.86 (t, <i>J</i> = 7.4 Hz, 3H).
221	H-142		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(pyrrolidin-1-yl)phenyl]-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 45% ACN up to 77% in 7 min LC-MS: (ES, <i>m/z</i>): 462.1. ¹ H NMR: (300 MHz, DMSO) δ 7.22 (m, 2H), 7.03 (dd, <i>J</i> = 7.7, 1.7 Hz, 1H), 6.93 (m, 2H), 6.86 (ddd, <i>J</i> = 8.3, 2.5, 1.0 Hz, 1H), 6.74 (dd, <i>J</i> = 8.5, 1.3 Hz, 1H), 6.67 (td, <i>J</i> = 7.5, 7.5, 1.2 Hz, 1H), 6.63 (s, 1H), 4.47 (s, 2H), 3.51 (tt, <i>J</i> = 6.1, 6.1, 3.0, 3.0 Hz, 1H), 2.75 (qq, <i>J</i> = 8.5, 8.5, 8.5, 5.0, 5.0, 4.6 Hz, 4H), 1.64 (m, 4H), 1.41 (s, 6H), 0.62 (m, 2H), 0.50 (m, 2H).
222	H-129		(2 <i>R</i>)-2-([1-(2-Chlorophenyl)-5-(3,5-dimethoxyphenyl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 45% ACN up to 64% in 7 min LC-MS: (ES, <i>m/z</i>): 445. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.67 (s, 1H), 7.74 – 7.38 (m, 4H), 6.71 (s, 1H), 6.41 (t, <i>J</i> = 2.2 Hz, 1H), 6.32 (d, <i>J</i> = 2.2 Hz, 2H), 4.47 (d, <i>J</i> = 2.0 Hz, 2H), 3.60 (s, 6H), 1.77 (hept, <i>J</i> = 7.0 Hz, 2H), 1.39 (s, 3H), 0.88 (t, <i>J</i> = 7.4 Hz, 3H).
223	H-126		(2 <i>R</i>)-2-([5-[3-(2,2-Dimethylpropoxy)phenyl]-1-(1-methyl-1 <i>H</i> -indazol-7-yl)-1 <i>H</i> -pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 55% ACN up to 81% in 7 min LC-MS: (ES, <i>m/z</i>): 491. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.70 (s, 1H), 8.20 (s, 1H), 7.92 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.41 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.29 – 7.04 (m, 2H), 6.92 – 6.75 (m, 3H), 6.63 (dd, <i>J</i> = 2.5, 1.6 Hz, 1H), 4.54 (s, 2H), 3.44 (s, 3H), 3.19 (s, 2H), 1.91 – 1.69 (m, 2H), 1.41 (s, 3H), 0.89 (d, <i>J</i> = 4.1 Hz, 12H).

224	H-107		2-([1-[(2-Chlorophenyl)methyl]-5-(3,5-dimethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 45% ACN up to 65% in 7 min LC-MS: (ES, <i>m/z</i>): 445. ¹ H-NMR: (MeOD): δ : 7.46 – 7.36 (m, 1H), 7.35 – 7.21 (m, 2H), 6.80 – 6.71 (m, 1H), 6.58 – 6.39 (m, 4H), 5.43 (s, 2H), 4.58 (s, 2H), 3.66 (s, 6H), 1.52 (s, 6H).
225	H-112		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(dimethylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 64% ACN up to 78% in 7 min LC-MS: (ES, <i>m/z</i>): 436. ¹ H-NMR: (MeOD, <i>ppm</i>): δ : 7.47 – 7.29 (m, 2H), 7.26 – 7.06 (m, 3H), 6.96 – 6.81 (m, 3H), 6.71 (s, 1H), 4.63 (s, 2H), 3.50 (tt, <i>J</i> = 6.2, 3.0 Hz, 1H), 2.44 (s, 6H), 1.54 (s, 6H), 0.70 – 0.47 (m, 4H).
226	H-108		2-([1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 38% ACN up to 61% in 7 min LC-MS: (ES, <i>m/z</i>): 439. ¹ H NMR: (300 MHz, MeOH- <i>d</i> ₄) δ 8.02 (d, <i>J</i> = 1.0 Hz, 1H), 7.77 (dd, <i>J</i> = 8.4, 0.9 Hz, 1H), 7.46 (q, <i>J</i> = 1.1 Hz, 1H), 7.43 – 7.32 (m, 1H), 7.32 – 7.20 (m, 2H), 7.13 (dd, <i>J</i> = 8.4, 1.4 Hz, 1H), 6.95 – 6.77 (m, 1H), 6.64 (s, 1H), 5.49 (s, 2H), 4.61 (s, 2H), 3.97 (s, 3H), 1.53 (s, 6H).
227	H-109		2-([1-[(2-Chlorophenyl)methyl]-5-[2-(propan-2-yloxy)-1,3-oxazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 35% ACN up to 50% in 7 min LC-MS: (ES, <i>m/z</i>): 434. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.59 (s, 1H), 7.49 (dd, <i>J</i> = 7.7, 1.6 Hz, 1H), 7.31 (dtd, <i>J</i> = 16.7, 7.5, 1.7 Hz, 2H), 6.78 (dd, <i>J</i> = 7.5, 1.9 Hz, 1H), 6.54 (s, 1H), 5.50 (s, 2H), 4.37 (s, 2H), 4.11 (p, <i>J</i> = 6.7 Hz, 1H), 1.38 (s, 6H), 1.26 (d, <i>J</i> = 6.7 Hz, 6H).





228	H-133		2-([1-[2-(Azetidin-1-yl)phenyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 38% ACN up to 64% in 7 min LC-MS: (ES, <i>m/z</i>): 460.15. ¹ H NMR (300 MHz, DMSO) δ 7.99 (d, <i>J</i> = 0.9 Hz, 1H), 7.65 (d, <i>J</i> = 8.5 Hz, 1H), 7.54 (d, <i>J</i> = 1.3 Hz, 1H), 7.27 (ddd, <i>J</i> = 8.5, 7.4, 1.6 Hz, 1H), 7.09 (ddd, <i>J</i> = 20.0, 8.1, 1.5 Hz, 2H), 6.71 (m, 2H), 6.49 (dd, <i>J</i> = 8.2, 1.3 Hz, 1H), 4.49 (s, 2H), 4.28 (q, <i>J</i> = 7.2, 7.2, 7.2 Hz, 2H), 3.47 (dt, <i>J</i> = 11.8, 7.3, 7.3 Hz, 4H), 2.05 (m, 2H), 1.43 (s, 6H), 1.23 (t, <i>J</i> = 7.1, 7.1 Hz, 3H).
229	H-135		2-([5-(1-ethyl-1H-indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 52% ACN up to 78% in 7 min LC-MS: (ES, <i>m/z</i>): 474. ¹ H NMR (300 MHz, DMSO) δ 12.66 (s, 1H), 7.98 (d, <i>J</i> = 1.0 Hz, 1H), 7.63 (d, <i>J</i> = 8.4 Hz, 1H), 7.43 (d, <i>J</i> = 1.4 Hz, 1H), 7.25 (ddd, <i>J</i> = 8.6, 7.2, 1.7 Hz, 1H), 7.07 (ddd, <i>J</i> = 8.4, 5.7, 1.5 Hz, 2H), 6.72 (m, 3H), 4.50 (s, 2H), 4.25 (q, <i>J</i> = 7.2, 7.2, 7.2 Hz, 2H), 2.77 (m, 4H), 1.70 (m, 2H), 1.60 (m, 2H), 1.43 (s, 6H), 1.22 (t, <i>J</i> = 7.2, 7.2 Hz, 3H).
230	H-136		2-Methyl-2-([5-(1-methyl-1H-indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]methoxy]propanoic acid XBridge: 18% ACN up to 39% in 7 min LC-MS: (ES, <i>m/z</i>): 460. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 7.97 (d, <i>J</i> = 0.9 Hz, 1H), 7.60 (d, <i>J</i> = 8.5 Hz, 1H), 7.47 (s, 1H), 7.25 (ddd, <i>J</i> = 8.6, 7.2, 1.7 Hz, 1H), 7.14 – 6.94 (m, 2H), 6.82 – 6.57 (m, 3H), 4.50 (s, 2H), 3.88 (s, 3H), 2.77 (q, <i>J</i> = 6.0, 4.4 Hz, 4H), 1.84 – 1.48 (m, 4H), 1.40 (s, 6H).
231	H-137		2-([5-(3,5-Dimethoxyphenyl)-1-[2-(pyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 42% ACN up to 75% in 7 min LC-MS: (ES, <i>m/z</i>): 466. H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.59 (s, 1H), 7.25 (ddd, <i>J</i> = 8.7, 7.2, 1.7 Hz, 1H), 7.00 (dd, <i>J</i> = 7.7, 1.7 Hz, 1H), 6.80 – 6.72 (m, 1H), 6.71 – 6.62 (m, 2H), 6.42 (d, <i>J</i> = 2.3 Hz, 2H), 6.37 (t, <i>J</i> = 2.2 Hz, 1H), 4.46 (s, 2H), 2.79 (t, <i>J</i> = 5.5 Hz, 4H), 1.79 – 1.55 (m, 4H), 1.41 (s, 6H).

232	H-143		2-([1-[2-(Azetidin-1-yl)phenyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 57% ACN up to 67% in 7 min LC-MS: (ES, m/z): 448.3. ^1H NMR: (300 MHz, MeOD) δ 7.29 (ddd, J = 8.2, 7.3, 1.6 Hz, 1H), 7.20 (m, 1H), 7.05 (dd, J = 7.3, 1.6 Hz, 3H), 6.89 (m, 1H), 6.74 (td, J = 7.6, 7.6, 1.3 Hz, 1H), 6.69 (s, 1H), 6.54 (dd, J = 8.2, 1.3 Hz, 1H), 4.59 (s, 2H), 3.49 (m, 5H), 2.13 (m, 2H), 1.53 (s, 6H), 0.66 (q, J = 5.7, 5.0, 5.0 Hz, 2H), 0.56 (tt, J = 6.0, 6.0, 3.1, 3.1 Hz, 2H).
233	H-144		2-([5-(3-Cyclopropoxyphenyl)-1-[2-[(3S)-3-hydroxypyrrolidin-1-yl]phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 36% ACN up to 64% in 7 min LC-MS: (ES, m/z): 478. ^1H NMR (300 MHz, DMSO) δ 7.22 (m, 2H), 7.02 (m, 2H), 6.89 (m, 2H), 6.73 (m, 1H), 6.63 (dd, J = 16.0, 5.1 Hz, 2H), 4.47 (m, 2H), 4.10 (m, 1H), 3.52 (dp, J = 8.9, 3.1, 3.1, 3.0, 3.0 Hz, 1H), 3.08 (q, J = 8.8, 8.8, 8.6 Hz, 1H), 2.95 (dt, J = 9.1, 4.4, 4.4 Hz, 1H), 2.86 (dt, J = 10.0, 4.7, 4.7 Hz, 1H), 2.74 (dt, J = 9.2, 5.4, 5.4 Hz, 1H), 2.63 (ddd, J = 11.3, 5.4, 2.6 Hz, 1H), 1.78 (ddd, J = 12.9, 8.4, 4.8 Hz, 1H), 1.64 (m, 1H), 1.41 (d, J = 2.2 Hz, 6H), 0.62 (q, J = 7.7, 6.1, 6.1 Hz, 2H), 0.48 (dq, J = 8.5, 3.2, 3.2, 3.2 Hz, 2H).
234	H-145		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(morpholin-4-yl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 40% ACN up to 65% in 7 min LC-MS: (ES, m/z): 478.1. ^1H NMR: (300 MHz, DMSO) δ 7.50 (dd, J = 7.8, 1.6 Hz, 1H), 7.39 (td, J = 7.7, 7.7, 1.7 Hz, 1H), 7.19 (dd, J = 8.4, 7.1 Hz, 2H), 6.93 (m, 2H), 6.83 (m, 2H), 6.69 (s, 1H), 4.48 (s, 2H), 3.49 (tt, J = 5.9, 5.9, 3.3, 3.3 Hz, 1H), 3.32 (dd, J = 9.8, 5.7 Hz, 2H), 3.11 (m, 2H), 2.39 (d, J = 12.6 Hz, 2H), 2.15 (m, 2H), 1.41 (s, 6H), 0.52 (m, 4H).
235	H-146		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(2-oxopyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 33% ACN up to 58% in 7 min LC-MS: (ES, m/z): 476.10. ^1H NMR (300 MHz, DMSO) δ 12.65 (s, 1H), 7.45 (m, 2H), 7.31 (m, 1H), 7.22 (t, J = 7.9, 7.9 Hz, 1H), 7.12 (m, 1H), 7.06 (m, 1H), 7.01 (dt, J = 7.6, 1.2, 1.2 Hz, 1H), 6.88 (ddd,

			J = 8.3, 2.5, 1.0 Hz, 1H), 6.57 (s, 1H), 4.45 (s, 2H), 3.56 (tt, J = 6.1, 6.1, 3.0, 3.0 Hz, 1H), 3.38 (t, J = 6 Hz, 2H), 2.11 (t, J = 7.9, 7.9 Hz, 2H), 1.92 (p, J = 7.4, 7.4, 7.2, 7.2 Hz, 2H), 1.41 (s, 6H), 0.54 (m, 2H), 0.44 (m, 2H).
236	H-147		2-([5-(3-Cyclopropoxyphenyl)-1-[2-(1H-pyrazol-1-yl)phenyl]-1H-pyrazol-3-yl]-methoxy)-2-methylpropanoic acid XSelect: 38% ACN up to 65% in 7 min LC-MS: (ES, <i>m/z</i>): 459.15. ¹ H NMR (300 MHz, DMSO) δ 7.63 (m, 3H), 7.52 (m, 2H), 7.09 (t, J = 7.9, 7.9 Hz, 1H), 6.83 (m, 2H), 6.63 (dd, J = 2.5, 1.5 Hz, 1H), 6.56 (m, 2H), 6.28 (dd, J = 2.5, 1.8 Hz, 1H), 4.48 (s, 2H), 3.52 (dt, J = 6.0, 3.1, 3.1 Hz, 1H), 1.41 (s, 6H), 0.61 (m, 2H), 0.47 (m, 2H).
237	H-148		2-([5-(3-Cyclopropoxyphenyl)-1-[2-[(3R)-3-hydroxypyrrolidin-1-yl]phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 37% ACN up to 64% in 7 min LC-MS: (ES, <i>m/z</i>): 478.15. ¹ H NMR (300 MHz, DMSO) δ 7.22 (m, 2H), 7.02 (m, 2H), 6.90 (m, 2H), 6.74 (m, 1H), 6.63 (m, 2H), 4.46 (m, 2H), 4.10 (p, J = 5.2, 5.2, 4.5, 4.5 Hz, 1H), 3.52 (m, 1H), 3.06 (p, J = 8.4, 8.4, 8.2, 8.2 Hz, 1H), 2.95 (td, J = 9.0, 8.5, 4.0 Hz, 1H), 2.86 (dt, J = 9.9, 4.7, 4.7 Hz, 1H), 2.74 (dt, J = 9.3, 5.4, 5.4 Hz, 1H), 2.63 (ddd, J = 10.6, 5.9, 2.5 Hz, 1H), 1.78 (ddd, J = 12.8, 8.3, 4.8 Hz, 1H), 1.64 (dd, J = 7.7, 4.1 Hz, 1H), 1.41 (d, J = 2.2 Hz, 6H), 0.61 (dd, J = 9.2, 4.6 Hz, 2H), 0.48 (dq, J = 10.3, 3.1, 2.7, 2.7 Hz, 2H).
238	H-151		2-([5-(3-Cyclobutoxyphenyl)-1-[[2-(pyrrolidin-1-yl)phenyl]methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 60% ACN up to 95% in 7 min LC-MS: (ES, <i>m/z</i>): 490. ¹ H NMR: (300 MHz, DMSO- <i>d</i> ₆) δ 12.67 (s, 1H), 7.29 (t, J = 8.0 Hz, 1H), 7.19 (ddd, J = 8.7, 7.2, 1.6 Hz, 1H), 7.06 (dd, J = 8.2, 1.2 Hz, 1H), 6.97 – 6.80 (m, 3H), 6.64 – 6.54 (m, 2H), 6.45 (s, 1H), 5.30 (s, 2H), 4.44 (s, 2H), 4.41 (d, J = 7.1 Hz, 1H), 3.19 – 2.87 (m, 4H), 2.26 – 2.07 (m, 2H), 1.99 – 1.88 (m, 2H), 1.88 – 1.80 (m, 4H), 1.71 (q, J = 10.1 Hz, 1H), 1.62 – 1.46 (m, 1H), 1.41 (s, 6H), 1.20 (s, 1H).





239	H-156		2-([1-(2-Cyanophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 35% ACN up to 64% in 7 min LC-MS: (ES, <i>m/z</i>):418. ¹ H-NMR: (DMSO, <i>ppm</i>): ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.73 (s, 1H), 7.99 (dd, <i>J</i> = 7.7, 1.5 Hz, 1H), 7.83 (td, <i>J</i> = 7.8, 1.6 Hz, 1H), 7.73 – 7.52 (m, 2H), 7.27 (t, <i>J</i> = 7.9 Hz, 1H), 6.97 (ddd, <i>J</i> = 8.3, 2.5, 1.0 Hz, 1H), 6.91 – 6.79 (m, 2H), 6.75 (s, 1H), 4.52 (s, 2H), 3.64 (tt, <i>J</i> = 6.1, 3.0 Hz, 1H), 1.44 (s, 6H), 0.70 – 0.44 (m, 4H).
240			2-([5-(3,5-Dimethoxyphenyl)-1-phenyl-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 395. ¹ H NMR (CDCl ₃): δ 1.61 (s, 6H), 3.66(s, 6H), 4.69 (s, 2H), 6.37 (d, <i>J</i> = 2.30, 2H), 6.41-6.44 (m, 1H), 6.55(s, 1H), 7.30-7.40 (m, 5H).
241			2-([5-(3,5-Dimethoxyphenyl)-1-(3-chlorophenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 431. ¹ H NMR (CDCl ₃): δ 1.59 (s, 6H), 3.68 (s, 6H), 4.65 (s, 2H), 6.35 (d, <i>J</i> =2.15 hZ, 2H), 6.40-6.46 (m, 1H), 6.53 (s,1H), 7.08-7.14 (m, 1H), 7.19-7.31 (m, 2H), 7.39-7.44 (m, 1H).
242			2-([5-(3,5-Dimethoxyphenyl)-1-(3-methoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 427. ¹ H NMR (CDCl ₃): δ 1.59 (s, 6H), 3.66 (s, 6H), 3.73 (s, 3H), 4.66 (s, 2H), 6.36 (d, <i>J</i> = 2.25 Hz, 2H), 6.41 (t, <i>J</i> = 2.21 Hz, 1H), 6.79-6.95 (m, 3H), 7.21 (t, <i>J</i> = 8.10 Hz, 1H).
243			2-([5-(3,5-Dimethoxyphenyl)-1-(3-ethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 441. ¹ H NMR (CDCl ₃): δ 1.33 (t, <i>J</i> = 7.00 Hz, 3H), 1.54 (s, 6H), 3.63 (s, 6H), 3.92 (q, <i>J</i> = 7.00 Hz, 2H), 4.63 (s, 2H), 6.34-6.38 (m, 3H), 6.48 (s, 1H), 6.81-6.88 (m, 3H), 7.16 (t, <i>J</i> = 8.00 Hz, 1H).





244			2-([5-(3,5-Dimethoxyphenyl)-1-[3-(dimethylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 441. ¹ H NMR (CDCl ₃): δ 1.51 (s, 6H), 2.85 (s, 6H), 3.62 (s, 6H), 4.58 (s, 2H), 6.38-6.41 (m, 3H), 6.54-6.657(m, 1H), 6.61(s, 1H), 7.74-6.76(m, 1H), 7.20 (t, J=8.11 Hz, 1H).
245			2-([5-(3,5-Dimethoxyphenyl)-1-[2-(dimethylamino)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid LC-MS (ES, <i>m/z</i>): 440. ¹ H NMR (CDCl ₃): δ 1.52 (s, 6H), 2.23(s, 6H), 3.57(s, 6H), 4.60(s, 2H), 6.31-6.36(m, 3H), 6.67(s, 1H), 6.91(d, J=8.00 Hz, 1H), 7.03-7.04(m, 1H), 7.33-7.38(m, 2H).
246	H-139		2-([1-[2-(Azetidin-1-yl)phenyl]-5-(1-propyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 40% ACN up to 70% in 7 min LC-MS: (ES, <i>m/z</i>): 474.10 ¹ H NMR (300 MHz, MeOD) δ 7.96 (d, J = 0.9 Hz, 1H), 7.71 (dd, J = 8.5, 0.8 Hz, 1H), 7.42 (s, 1H), 7.31 (m, 2H), 7.13 (d, J = 7.8 Hz, 1H), 6.80 (d, J = 13.8 Hz, 2H), 6.56 (d, J = 8.2 Hz, 1H), 4.63 (s, 2H), 4.17 (t, J = 7.0, 7.0 Hz, 2H), 3.57 (m, 4H), 2.14 (q, J = 7.2, 7.2, 7.2 Hz, 2H), 1.69 (q, J = 7.2, 7.2, 7.2 Hz, 2H), 1.54 (s, 6H), 0.79 (t, J = 7.4, 7.4 Hz, 3H).
247	H-70		2-Methyl-2-([5-[3-(2-methylpropoxy)-phenyl]-1-([pyridin-2-yl]methyl)-1H-pyrazol-3-yl]methoxy)propanoic acid XSelect: 40% ACN up to 70% in 7 min LC-MS: (ES, <i>m/z</i>): 424. ¹ H NMR (300 MHz, Methanol-d ₄) δ 8.70 (ddd, J = 5.5, 1.7, 0.8 Hz, 1H), 8.27 (td, J = 7.9, 1.7 Hz, 1H), 7.75 (ddd, J = 7.7, 5.5, 1.2 Hz, 1H), 7.51 – 7.31 (m, 2H), 7.15 – 6.88 (m, 3H), 6.57 (s, 1H), 5.65 (s, 2H), 4.58 (s, 2H), 3.71 (d, J = 6.5 Hz, 2H), 2.04 (dq, J = 13.2, 6.6 Hz, 1H), 1.52 (s, 6H), 1.03 (d, J = 6.7 Hz, 6H).



248	H-80		2-([1-[(2-Chlorophenyl)methyl]-5-(1-methyl-1H-1,3-benzodiazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 20% ACN up to 34% in 7 min LC-MS: (ES, <i>m/z</i>): 438. ¹ H NMR (DMSO- <i>d</i> ₆) δ: 8.23 (s, 1H), 7.65 (d, <i>J</i> = 8.4 Hz, 1H), 7.57 (s, 1H), 7.47 – 7.36 (m, 1H), 7.34 – 7.20 (m, 2H), 7.17 (d, <i>J</i> = 8.3 Hz, 1H), 6.83 – 6.73 (m, 1H), 6.47 (s, 1H), 5.42 (s, 2H), 4.46 (s, 2H), 3.78 (s, 3H), 1.36 (s, 6H).
249	H-33		2-([5-(3,5-Dimethoxyphenyl)-1-(1-methyl-1H-indazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 30% ACN up to 65% in 7 min LC-MS: (ES, <i>m/z</i>): 451. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.19 (s, 1H), 7.92 (s, 1H), 7.44 (dd, <i>J</i> = 7.3, 1.0 Hz, 1H), 7.21 (s, 1H), 6.84 (s, 1H), 6.35 (dt, <i>J</i> = 10.9, 2.2 Hz, 3H), 4.53 (s, 2H), 3.52 (s, 6H), 3.41 (s, 3H), 1.44 (s, 6H).
250	H-140		2-Methyl-2-([5-(1-propyl-1H-indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]methoxy)propanoic acid XBridge: 10% ACN up to 50% in 7 min LC-MS: (ES, <i>m/z</i>): 488.2. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.98 (s, 1H), 7.64 (d, <i>J</i> = 8.4 Hz, 1H), 7.37 (s, 1H), 7.25 (m, 1H), 7.08 (t, <i>J</i> = 7.5, 7.5 Hz, 2H), 6.72 (m, 3H), 4.50 (s, 2H), 4.14 (t, <i>J</i> = 7.1, 7.1 Hz, 2H), 2.77 (q, <i>J</i> = 6.4, 6.4, 5.2 Hz, 4H), 1.63 (tt, <i>J</i> = 15.4, 15.4, 7.1, 7.1 Hz, 6H), 1.40 (s, 6H), 0.72 (t, <i>J</i> = 7.4, 7.4 Hz, 3H).
251	H-138		2-((1-(2-(Azetidin-1-yl)phenyl)-5-(3,5-dimethoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XBridge: 25% ACN up to 37% in 7 min LC-MS: (ES, <i>m/z</i>): 452. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.26 (t, <i>J</i> = 7.4 Hz, 1H), 6.95 (d, <i>J</i> = 7.5 Hz, 1H), 6.67 (q, <i>J</i> = 7.0, 6.0 Hz, 2H), 6.58 – 6.43 (m, 3H), 6.38 (t, <i>J</i> = 2.2 Hz, 1H), 4.47 (s, 2H), 3.59 (s, 6H), 3.45 (dq, <i>J</i> = 31.0, 7.3 Hz, 4H), 2.17 – 1.89 (m, 2H), 1.36 (s, 6H).

252	H-72		2-([5-(3,5-Dimethoxyphenyl)-1-[(2-ethoxy-6-fluorophenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 42% ACN up to 72% in 7 min LC-MS: (ES, <i>m/z</i>): 473. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.27 (td, <i>J</i> = 8.4, 8.3, 6.9 Hz, 1H), 6.76 (dt, <i>J</i> = 8.8, 4.6, 4.6 Hz, 2H), 6.66 (d, <i>J</i> = 2.3 Hz, 2H), 6.57 (t, <i>J</i> = 2.2, 2.2 Hz, 1H), 6.25 (s, 1H), 5.26 (s, 2H), 4.27 (s, 2H), 3.88 (q, <i>J</i> = 7.0, 7.0, 7.0 Hz, 2H), 3.78 (s, 6H), 1.34 (s, 6H), 1.06 (t, <i>J</i> = 6.9, 6.9 Hz, 3H).
253	H-83		2-([5-[3-(2,2-Dimethylpropoxy)phenyl]-1-(quinolin-8-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 80% in 7 min LC-MS: (ES, <i>m/z</i>): [M+1] = 474. ¹ H-NMR: (DMSO, ppm): δ: 8.77 (dd, <i>J</i> = 4.2, 1.7 Hz, 1H), 8.46 (dd, <i>J</i> = 8.4, 1.7 Hz, 1H), 8.14 (dd, <i>J</i> = 8.2, 1.5 Hz, 1H), 7.83 (dd, <i>J</i> = 7.4, 1.5 Hz, 1H), 7.70 (dd, <i>J</i> = 8.2, 7.3 Hz, 1H), 7.55 (dd, <i>J</i> = 8.3, 4.2 Hz, 1H), 7.02 (t, <i>J</i> = 8.0 Hz, 1H), 6.79 – 6.60 (m, 3H), 6.53 (dd, <i>J</i> = 2.6, 1.5 Hz, 1H), 4.49 (s, 2H), 3.02 (s, 2H), 1.43 (s, 6H), 0.81 (s, 9H).
254	H-73		2-([5-(3,5-Dimethoxyphenyl)-1-([pyridin-2-yl]methyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 25% ACN up to 56% in 7 min LC-MS: (ES, <i>m/z</i>): 412.05. ¹ H NMR (300 MHz, DMSO) δ 12.62 (s, 1H), 8.52 (ddd, <i>J</i> = 4.8, 1.8, 0.9 Hz, 1H), 7.78 (td, <i>J</i> = 7.7, 7.7, 1.8 Hz, 1H), 7.30 (ddd, <i>J</i> = 7.6, 4.8, 1.2 Hz, 1H), 7.08 (dt, <i>J</i> = 7.8, 1.1, 1.1 Hz, 1H), 6.70 (d, <i>J</i> = 2.3 Hz, 2H), 6.52 (t, <i>J</i> = 2.3, 2.3 Hz, 1H), 6.42 (s, 1H), 5.37 (s, 2H), 4.38 (s, 2H), 3.70 (s, 6H), 1.38 (s, 6H).
255	H-35		2-([1-(1,3-Dimethyl-1H-indazol-7-yl)-5-[3-(2,2-dimethylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 80% in 7 min LC-MS: (ES, <i>m/z</i>): 405.35. ¹ H NMR (300 MHz, DMSO) δ 12.60 (s, 1H), 7.83 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 7.34 (dd, <i>J</i> = 7.3, 0.9 Hz, 1H), 7.12 (m, 2H), 6.78 (m, 3H), 6.60 (m, 1H), 4.52 (s, 2H), 3.33 (s, 3H), 3.14 (s, 2H), 2.46 (s, 3H), 1.42 (s, 6H), 0.85 (s, 9H).

256	H-69		2-([5-[3-(2,2-Dimethylpropoxy)phenyl]-1-[(1-methyl-1H-indazol-7-yl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 60% ACN up to 70% in 7 min LC-MS: (ES, m/z): 491.25. ¹ H NMR (400 MHz, DMSO) δ 8.03 (s, 1H), 7.66 (d, J = 8.0 Hz, 1H), 7.33 (t, J = 7.9, 7.9 Hz, 1H), 7.02 (m, 2H), 6.94 (dd, J = 8.3, 2.4 Hz, 1H), 6.88 (s, 1H), 6.56 (d, J = 7.2 Hz, 1H), 6.51 (s, 1H), 5.95 (s, 2H), 4.43 (s, 2H), 4.20 (s, 3H), 3.32 (s, 2H), 1.39 (s, 6H), 0.88 (s, 9H).
257	H-131		(S)-2-((1-(2-chlorobenzyl)-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl)methoxy)-2-methylbutanoic acid LC-MS: (ES, m/z): 467.10. ¹ H NMR (300 MHz, DMSO-d ₆) δ 12.64 (s, 1H), 8.07 (d, J = 0.9 Hz, 1H), 7.78 (dd, J = 8.3, 0.8 Hz, 1H), 7.67 (d, J = 1.3 Hz, 1H), 7.49 – 7.34 (m, 1H), 7.36 – 7.18 (m, 2H), 7.11 (dd, J = 8.3, 1.3 Hz, 1H), 6.84 – 6.72 (m, 1H), 6.52 (s, 1H), 5.45 (s, 2H), 4.43 (d, J = 2.2 Hz, 2H), 4.37 (q, J = 7.2 Hz, 2H), 1.74 (hept, J = 7.0 Hz, 2H), 1.37 (s, 3H), 1.31 (t, J = 7.2 Hz, 3H), 0.85 (t, J = 7.4 Hz, 3H).
258	H-36		2-([5-[3-(2,2-Dimethylpropoxy)phenyl]-1-(1-methyl-1H-1,2,3-benzotriazol-7-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 50% ACN up to 70% in 7 min LC-MS: (ES, m/z): 478. ¹ H NMR (300 MHz, DMSO-d ₆) δ 12.80 (s, 1H), δ 8.17 (dd, J = 8.2, 1.1 Hz, 1H), 7.59 – 7.31 (m, 2H), 7.12 (t, J = 8.0 Hz, 1H), 6.89 – 6.77 (m, 2H), 6.76 – 6.62 (m, 2H), 4.54 (s, 2H), 3.81 (s, 3H), 3.29 (s, 2H), 1.42 (s, 6H), 0.87 (s, 9H).
259	H-65		2-([5-(1-Ethyl-1H-indazol-6-yl)-1-([pyridin-2-yl]methyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid SunFire: 36% ACN up to 50% in 7 min LC-MS: (ES, m/z): 420.2. ¹ H NMR (300 MHz, DMSO) δ 8.50 (dd, J = 4.9, 1.7 Hz, 1H), 8.07 (d, J = 0.9 Hz, 1H), 7.87 (t, J = 1.2 Hz, 1H), 7.78 (d, J = 8.4 Hz, 1H), 7.73 (dd, J = 7.7, 1.8 Hz, 1H), 7.27 (m, 2H), 7.07 (d, J = 7.8 Hz, 1H), 6.49 (s, 1H), 5.42 (s, 2H), 4.39 (m, 4H), 1.38 (s, 6H), 1.33 (t, J = 7.2 Hz, 3H).

260	H-141		2-([1-[2-(Azetidin-1-yl)phenyl]-5-(3-cyclobutoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 55% ACN up to 70% in 7 min LC-MS: (ES, m/z): 462.1. ¹ H NMR (300 MHz, DMSO-d ₆) δ 7.27 (td, J = 7.8, 1.6 Hz, 1H), 7.19 (t, J = 7.9 Hz, 1H), 6.97 (td, J = 6.2, 3.2 Hz, 2H), 6.75 – 6.64 (m, 3H), 6.61 (s, 1H), 6.50 (dd, J = 8.3, 1.3 Hz, 1H), 4.43 (s, 2H), 4.29 (p, J = 7.1 Hz, 1H), 3.44 (dt, J = 14.4, 7.3 Hz, 4H), 2.31 – 2.13 (m, 2H), 2.03 (p, J = 7.3 Hz, 2H), 1.95 – 1.82 (m, 2H), 1.78 – 1.64 (m, 1H), 1.62 – 1.46 (m, 1H), 1.39 (s, 6H).
261	H-81		2-([1-[(2-chlorophenyl)methyl]-5-(1-ethyl-1H-1,2,3-benzotriazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 40% ACN up to 62% in 7 min LC-MS: (ES, m/z): 454. ¹ H NMR (400 MHz, DMSO-d ₆) δ 12.70 (s, 1H), 8.08 (d, J = 8.6 Hz, 1H), 8.00 (t, J = 1.1 Hz, 1H), 7.50 – 7.37 (m, 2H), 7.31 – 7.23 (m, 2H), 6.86 – 6.79 (m, 1H), 6.58 (s, 1H), 5.48 (s, 2H), 4.73 (q, J = 7.2 Hz, 2H), 4.45 (s, 2H), 1.47 (t, J = 7.3 Hz, 3H), 1.41 (s, 6H).
262	H-128		(2R)-2-([1-(2-Chlorophenyl)-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 57% ACN up to 61% in 7 min LC-MS: (ES, m/z): 440.9. ¹ H NMR (300 MHz, DMSO-d ₆) δ 12.65 (s, 1H), 7.69 – 7.38 (m, 4H), 7.22 (t, J = 8.0 Hz, 1H), 6.89 (ddt, J = 11.0, 7.6, 1.1 Hz, 2H), 6.80 (dd, J = 2.4, 1.6 Hz, 1H), 6.66 (s, 1H), 4.45 (d, J = 2.2 Hz, 2H), 3.55 (tt, J = 6.1, 2.9 Hz, 1H), 1.75 (hept, J = 7.0 Hz, 2H), 1.38 (s, 3H), 0.86 (t, J = 7.4 Hz, 3H), 0.71 – 0.57 (m, 2H), 0.52 – 0.41 (m, 2H).
263	H-127		(2R)-2-Methyl-2-([1-(1-methyl-1H-indazol-7-yl)-5-[3-(2-methylpropoxy)phenyl]-1H-pyrazol-3-yl]methoxy)butanoic acid XSelect: 55% ACN up to 71% in 7 min LC-MS: (ES, m/z): 477.1. ¹ H NMR (300 MHz, DMSO-d ₆) δ 8.17 (s, 1H), 7.89 (d, J = 8.0 Hz, 1H), 7.38 (d, J = 7.2 Hz, 1H), 7.24-7.10 (m, 2H), 6.82-6.75 (m, 3H), 6.61 (t, J = 2.0 Hz, 1H), 4.51 (s, 2H), 3.41 (s, 3H), 3.32 (d, J = 6.7 Hz, 2H), 1.87 – 1.65 (m, 3H), 1.38 (s, 3H), 0.89-0.81 (m, 9H).

264	H-132		(2R)-2-([5-(3,5-Dimethoxyphenyl)-1-[(2-ethoxyphenyl)methyl]-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 60% ACN up to 64% in 7 min LC-MS: (ES, <i>m/z</i>): 469.0. ¹H NMR (300 MHz, DMSO-d₆) δ 7.21 (ddd, <i>J</i> = 8.7, 7.5, 1.7 Hz, 1H), 6.95 (dd, <i>J</i> = 8.4, 1.0 Hz, 1H), 6.84 (td, <i>J</i> = 7.5, 1.0 Hz, 1H), 6.74 (s, 1H), 6.66 (dd, <i>J</i> = 7.5, 1.7 Hz, 1H), 6.54 – 6.38 (m, 4H), 5.24 (s, 2H), 4.38 (d, <i>J</i> = 2.4 Hz, 2H), 3.97 (q, <i>J</i> = 6.9 Hz, 2H), 3.65 (s, 7H), 1.85 – 1.59 (m, <i>J</i> = 7.4 Hz, 2H), 1.34 (s, 3H), 1.21 (t, <i>J</i> = 6.9 Hz, 4H), 0.83 (t, <i>J</i> = 7.4 Hz, 3H)
265	H-110		2-([1-[(2-Chlorophenyl)methyl]-5-[1,3-dimethyl-1H-thieno[2,3-c]pyrazol-5-yl]-1H-pyrazol-3-yl]methoxy)-2-methylpropanoic acid XSelect: 48% ACN up to 52% in 7 min LC-MS: (ES, <i>m/z</i>): 459. ¹H NMR (300 MHz, DMSO-d₆) δ 7.52 – 7.44 (m, 1H), 7.35-7.20 (m, 2H), 7.11 (s, 1H), 6.70 (dd, <i>J</i> = 6.9, 2.5 Hz, 1H), 6.48 (s, 1H), 5.50 (s, 2H), 4.38 (s, 2H), 3.81 (s, 3H), 2.29 (s, 3H), 1.37 (s, 6H).
266	H-153		(2R)-2-([1-[2-Azetidin-1-yl]phenyl]-5-(3-cyclopropoxyphenyl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid SunFire: 72% ACN up to 82% in 7 min LC-MS: (ES, <i>m/z</i>): 462.3. ¹H NMR (300 MHz, DMSO-d₆) δ 7.29-7.21 (m, 2H), 7.10 – 6.95 (m, 3H), 6.94 – 6.84 (m, 1H), 6.75 – 6.63 (m, 1H), 6.61 (s, 1H), 6.48 (d, <i>J</i> = 8.2 Hz, 1H), 4.43 (s, 2H), 3.56-3.50 (m, 5H), 2.02 (q, <i>J</i> = 7.3 Hz, 2H), 1.74 (p, <i>J</i> = 7.1 Hz, 2H), 1.37 (s, 3H), 0.85 (t, <i>J</i> = 7.4 Hz, 3H), 0.64 (d, <i>J</i> = 6.0 Hz, 2H), 0.51 (q, <i>J</i> = 4.4, 3.3 Hz, 2H).
267	H-152		(2R)-2-([1-[2-(azetidin-1-yl)phenyl]-5-(1-propyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 57% ACN up to 61% in 7 min LC-MS: (ES, <i>m/z</i>): 488.3. ¹H NMR (300 MHz, DMSO-d₆) δ 7.98 (s, 1H), 7.66 (d, <i>J</i> = 8.4 Hz, 1H), 7.46 (s, 1H), 7.26 (t, <i>J</i> = 7.8 Hz, 1H), 7.15 (d, <i>J</i> = 8.5 Hz, 1H), 7.09 – 7.00 (m, 1H), 6.70 (d, <i>J</i> = 11.2 Hz, 2H), 6.48 (d, <i>J</i> = 8.2 Hz, 1H), 4.46 (s, 2H), 4.15 (t, <i>J</i> = 7.1 Hz, 2H), 3.53 – 3.44 (m, 4H), 2.04 (q, <i>J</i> = 7.4, 6.7 Hz, 2H), 1.74 (dq, <i>J</i> = 14.6, 7.3 Hz, 2H), 1.61 (h, <i>J</i> = 7.3 Hz, 2H),

			1.38 (s, 3H), 0.86 (t, J = 7.4 Hz, 3H), 0.72 (t, J = 7.4 Hz, 3H).
268	H-154		(2R)-2-([5-(1-ethyl-1H-indazol-6-yl)-1-[2-(pyrrolidin-1-yl)phenyl]-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XSelect: 50% ACN up to 72% in 7 min LC-MS: (ES, m/z): 488. ¹H NMR (300 MHz, DMSO-d₆) δ 12.71 (s, 1H), δ 7.96 (d, J = 0.9 Hz, 1H), 7.62 (d, J = 8.4 Hz, 1H), 7.46 – 7.40 (m, 1H), 7.24 (ddd, J = 8.5, 7.1, 1.6 Hz, 1H), 7.07 (td, J = 6.7, 6.3, 3.3 Hz, 2H), 6.87 – 6.59 (m, 3H), 4.48 (s, 2H), 4.23 (q, J = 7.2 Hz, 2H), 2.76 (d, J = 7.5 Hz, 4H), 1.74 (dh, J = 14.0, 7.2 Hz, 4H), 1.62 – 1.50 (m, 2H), 1.38 (s, 3H), 1.20 (t, J = 7.2 Hz, 3H), 0.86 (t, J = 7.4 Hz, 3H).
269	H-155		(2R)-2-([1-[2-(Azetidin-1-yl)phenyl]-5-(1-ethyl-1H-indazol-6-yl)-1H-pyrazol-3-yl]methoxy)-2-methylbutanoic acid XBridge: 44% ACN up to 54% in 7 min LC-MS: (ES, m/z): 474. ¹H NMR (300 MHz, DMSO-d₆) δ 7.96 (s, 1H), 7.63 (d, J = 8.4 Hz, 1H), 7.50 (s, 1H), 7.24 (t, J = 7.6 Hz, 1H), 7.08 (dd, J = 20.7, 8.1 Hz, 2H), 6.88 – 6.58 (m, 2H), 6.46 (d, J = 8.4 Hz, 1H), 4.48 (s, 2H), 4.25 (q, J = 7.2 Hz, 2H), 3.45 (dd, J = 22.1, 7.2 Hz, 4H), 2.18 – 1.88 (m, 2H), 1.71 (dd, J = 15.0, 7.4 Hz, 2H), 1.33 (s, 3H), 1.20 (t, J = 7.1 Hz, 3H), 0.84 (t, J = 7.3 Hz, 3H).

[0502] The activity of the compounds in Examples **1-269** as MCT4 inhibitors is illustrated in the following assays. The other compounds listed above, which have not yet been made and/or tested, are predicted to have activity in these assays as well.

Biological Activity Assays

[0503] The following are assays that may be used to evaluate the biological efficacy of compounds of Formula (I) in a manner similar to that previously reported for MCT1 and MCT4 and are known to those with skill in the art. See, e.g., Murray, C. M. *et al.*, “Monocarboxylate transporter MCT1 is a target for immunosuppression,” *Nature chemical biology* **1**, 371-376 (2005); and Ovens, M. J., *et al.*, “AR-C155858 is a potent inhibitor of monocarboxylate transporters MCT1 and MCT2 that binds to an intracellular site involving transmembrane helices 7-10,” *The Biochemical Journal* **425**, 523-530, (2010).

Preparing BCECF-loaded cells:

[0504] Cells ($\sim 7 \times 10^6$) are trypsinized (0.05% Trypsin-EDTA), pelleted (300g, 5 min), and resuspended in 1 mL Tyrode's Solution, pH 7.4 (119 mM NaCl, 5 mM KCl, 25 mM HEPES, pH 7.4, 2 mM CaCl_2 , 2 mM MgCl_2 , 6 g/L glucose). 10 μL of a 30 mM DMSO stock of BCECF-AM ester (Life Technologies) is added and the cells are incubated at 37°C for 5 min. The cells are pelleted (300g, 5 min), washed once with 1 mL Tyrode's Solution, pH 7.4, re-pelleted (300g, 5 min), and resuspended in 1 mL Tyrode's Solution, pH 7.4.

Assay 1: Lactate transport in MCT4-expressing MDA-MB-453 breast cancer cells.

[001] MCT4 may be stably expressed in MDA-MB-453 breast cancer cells that do not express native MCT1 or MCT4. MCT4 activity may be assessed by monitoring the intracellular pH change that accompanies lactate/proton symport, using the pH-sensitive fluorescent dye 2',7'-bis-(carboxyethyl)-5(6)-carboxyfluorescein (BCECF), in a manner similar to that previously reported for MCT1 and MCT4. The following is an exemplary procedure for assaying MCT4 activity of the compounds of Formula (I).

Assay 2: MCT4-mediated lactate transport in NCI-H358 lung adenocarcinoma cells.

[002] NCI-H358 lung adenocarcinoma cells may be used to measure MCT4 activity in cells with high native levels of MCT4 and low levels of MCT1 and are known to those with skill in the art. Preparation of BCECF-loaded cells and lactate transport activity may be determined as described for Assay 1.

Assay 3: MCT4-mediated lactate transport in MDA-MB-231 breast cancer cells.

[0505] MDA-MB-231 breast cancer cells may be used to measure MCT4 activity in cells with high native levels of MCT4 and low levels of MCT1 and are known to those with skill in the art. MCT4 activity may be assessed by monitoring the intracellular pH change that accompanies lactate/proton symport, using the pH-sensitive fluorescent dye 2',7'-bis-(carboxyethyl)-5(6)-carboxyfluorescein (BCECF), in a manner similar to that previously reported for MCT1 and MCT4. The following is an exemplary procedure for assaying MCT4 activity of the compounds of Formula (I).

Assay 4: MCT1-mediated lactate transport in BT20 breast cancer cells.

[0506] MCT1 activity may be measured using BT-20 breast cancer cells that express high native levels of MCT1, but do not express MCT4 and are known to those with skill in the art. Preparation of BCECF loaded cells are as described for Assay 1. Lactate transport assay is as described for Assay 1, except 10 mM L-lactate (rather than 50 mM) is added.

[003] Results of the assays above are given below in Tables 26 and 27, in which “ND” means no data. As can be seen, most compounds disclosed herein are selective for MCT4 over MCT1.

Table 26: MCT Activity.

Example	MCT4 IC ₅₀ (nM) MDA-MB-453 + MCT4	MCT4 IC ₅₀ (nM) NCI-H358	MCT1 IC ₅₀ (nM) BT20
1	340	140	50,000
2	450	240	29,000
3	72	60	25,000
4	68	54	33,000
5	160	63	1,800
6	2,400	4,600	6,600
7	720	460	70,000
8	69	81	46,000
9	320	340	39,000
10	21	33	8,800
11	45	25	29,000
12	38	49	58,000
13	20	22	17,000
14	26	44	>100,000
15	900	440	>33,000
16	1,000	640	73,000
17	440	860	1,200
18	5,300	4,200	45,000
19	110	160	57,000
20	17,000	22,000	7,900
21	35,000	48,000	300,000
22	14,000	39,000	150,000
23	1,500	2,100	>300,000

Table 27: MCT Activity.

N.D. = not determined

Ex.	MCT4 IC ₅₀ (nM) MDA-MB- 231	MCT1 IC ₅₀ (nM) BT20
14	30	>100,000
24	30	>100,000
25	33	77,000
26	9.4	27,000
27	1	5,600
28	12	15,000
29	3.7	16,000
30	8.8	27,000

31	53	42,000
32	79	83,000
33	3.2	9,100
34	9.3	>11,000
35	5.2	59,000
36	82	50,000
37	7.6	8,400
38	48	60,000
39	5.4	10,000
40	520	>100,000
41	55	1,200

42	2.6	9,800
43	13	>33,000
44	1.1	8,400
45	19	12,000
46	142	47,000
47	410	>100,000
48	89	55,000
49	240	130,000
50	70	34,000
51	15	6,000
52	190	84,000
53	160	>100,000
54	37	27,000
55	260	>100,000
56	8,000	>133,000
57	110	>133,000
58	150	78,000
59	340	81,000
60	13	44,000
61	3.1	11,000
62	33	27,000
63	37	17,000
64	8.3	45,000
65	8	13,000
66	7.8	45,000
67	11	17,000
68	9.1	26,000
69	5,800	>100,000
70	9,300	>100,000
71	34,000	64,000
72	220	45,000
73	21	>11,000
74	6,400	>100,000
75	1,700	>100,000
76	400	38,000
77	340	9,000
79	59	15,000
80	81	61,000
81	6,300	6,600
82	460	16,000
101	3.4	12,000

102	15	42,000
103	2.6	28,000
104	20	12,000
105	7.6	15,000
106	12	21,000
107	170	9,900
108	36	5,200
109	2.1	>11,000
110	20	30,000
111	7.9	32,000
112	29	94,000
113	6.7	33,000
114	4.3	73,000
115	34,000	>100,000
116	11,000	61,000
117	430	>100,000
118	170	50,000
119	67	65,000
120	4	22,000
121	8	21,000
122	50	2,500
123	1,500	11,000
124	420	31,000
125	5,900	61,000
126	130	35,000
127	41	15,000
128	1.2	>11,000
129	2.8	16,000
130	98	55
131	225	650
132	8.6	8,200
133	13	>33,000
134	38	72,000
135	53	220
136	920	>33,000
137	1,400	>100,000
138	535	>100,000
139	9,700	>100,000
140	8.3	16,000
141	6.7	33,000
142	6.3	26,000

143	195	3,300
144	127	2,400
150	2.8	25,000
151	3.6	70,000
152	80	>100,000
153	830	44,000
154	240	>100,000
155	0.32	>33,000
156	1400	>100,000
157	26	86,000
158	580	>100,000
159	11	>100,000
160	270	>100,000
161	3.3	22,000
162	6.4	34,000
163	4.4	73,000
164	43	>100,000
165	85	99,000
166	5.9	53,000
167	0.59	36,000
168	1.7	61,000
169	12.2	68,000
170	2.0	32,000
171	12	32,000
172	0.59	33,000
173	230	66,000
174	101	>100,000
175	5.1	28,000
176	180	>100,000
177	1.4	35,000
178	2.5	27,000
179	13	>100,000
180	2.9	28,000
181	20	25,000
182	5.9	12,000
183	1.3	86,000
184	22	>33,000
185	9.3	17,000
186	21.5	90,000
187	43	63,000
188	7.5	15,000

189	18	22,000
190	7.8	31,000
191	17	16,000
192	190	47,000
193	18	38,000
194	3500	>100,000
195	5.3	13,000
196	1.9	38,000
197	2000	76,000
198	3.5	26,000
199	9.6	34,000
200	1.9	17,000
201	7.2	15,000
202	6.2	23,000
203	185	93,000
204	40	>100,000
205	890	>100,000
206	16	45,000
207	15	3,200
208	7.5	17,000
209	31	95,000
210	2.4	7,300
211	2.4	20,000
212	195	>100,000
213	12	12,000
214	6.0	94,000
215	360	>100,000
216	N.D.	N.D
217	8.7	18,000
218	7.5	22,000
219	4.0	14,000
220	0.54	33,000
221	3.2	38,000
222	3.9	94,000
223	0.3	21,000
224	4.1	30,000
225	N.D.	N.D.
226	15	54,000
227	690	37,000
228	5.9	31,000
229	1.9	42,000

230	8.1	92,000
231	4.5	63,000
232	2.1	29,000
233	700	>100,000
234	960	70,000
235	>30,000	>100,000
236	170	>100,000
237	160	>100,000
238	3.7	21,000
239	150	>100,000
240	28	62,000
241	25	8,500
242	59	57,000
243	23	78,000
244	68	16,000
245	7.6	77,000
246	3.10	32,000
247	23.00	>100,000
248	560.00	>100,000
249	39.00	>100,000

250	1.30	41,000
251	5.60	98,000
252	25.00	43,000
253	19.00	55,000
254	285.00	>100,000
255	2.40	7,300
256	2.60	17,000
257	0.35	19,000
258	12.00	27,000
259	850.00	>100,000
260	0.83	26,000
261	17.00	43,000
262	9.10	7,900
263	2.20	16,000
264	0.35	10,000
265	5.60	84,000
266	0.36	64,000
267	0.57	11,000
268	0.30	36,000
269	1.00	58,000

Metabolic Stability Assays

[0507] The following are assays that may be used to evaluate the metabolic stability of compounds of Formula (I) in human, rat, or mouse microsomes. The assays follow a common procedure recited below.

[0508] 1. Master solution: 200 μ L of 200 mM Phosphate buffer, 106 μ L of ultra-pure water, 40 μ L of 50 mM $MgCl_2$, and 10 μ L of 20 mg/mL liver microsomes (human, rat, or mouse).

[0509] 2. Two separated experiments were performed as follows. a) With NADPH: 10 μ L of 20 mg/mL liver microsomes and 40 μ L of 10 mM NADPH were added to the incubations. The final concentrations of microsomes and NADPH were 0.5 mg/mL and 1 mM, respectively. b) Without NADPH: 10 μ L of 20 mg/mL liver microsomes and 40 μ L of ultra-pure H_2O were added to the incubations. The final concentration of microsomes was 0.5 mg/mL.

[0510] 3. The reaction was started with the addition of 4 μ L of 200 μ M control compound or test compound solutions. Verapamil was used as positive control in this study. The final concentration of test compound or control compound was 2 μ M.

[0511] 4. Aliquots of 50 μ L were taken from the reaction solution at 0, 15, 30, 45 and 60 min. The reaction was stopped by the addition of 4 volumes of cold acetonitrile with IS (100 nM alprazolam, 200 nM labetalol and 2 μ M ketoprofen). Samples were centrifuged at 3,220 g for 40 minutes. A 90 μ L aliquot of the supernatant was mixed with 90 μ L of ultra-pure H_2O and then used for LC-MS/MS analysis.

[0512] 5. Data analysis. All calculations were carried out using Microsoft Excel. Peak areas were determined from extracted ion chromatograms. The slope value, k, was determined by linear regression of the natural logarithm of the remaining percentage of the parent drug vs. incubation time curve. The *in vitro* half-life (*in vitro* $t_{1/2}$) was determined from the slope value: *in vitro* $t_{1/2} = -(0.693)/k$.

[0513] Results are given below in Table 28, wherein “ND” means no data.

Table 28: Metabolic Stability.

Ex.	Human microsome $t_{1/2}$, min	Human microsome % 60 min	Mouse microsome $t_{1/2}$, min	Mouse microsome % 60 min	Rat microsome $t_{1/2}$, min	Rat microsome % 60 min
14	245	82	1300	95	ND	ND
27	50	43	61	50	ND	ND

35	115	67	35	31	ND	ND
36	290	85	1200	99	ND	ND
38	76	56	79	61	ND	ND
39	380	89	370	88	ND	ND
54	78	59	760	95	ND	ND
57	190	80	>2000	100	ND	ND
64	200	82	270	84	ND	ND
65	140	78	560	92	ND	ND
66	120	73	260	83	ND	ND
68	790	95	380	91	ND	ND
110	2377	99	223	86	549	94
120	198	81	410	101	51	44
121	198	82	186	81	86	61
132	116	70	98	64	112	70
133	1006	97	344	90	404	91
134	186	80	109	72	264	85
140	109	67	64	62	59	48
150	341	90	∞	103	1715	98
155	52	62	38	37	73	70
157	149	78	9639	102	748	97
161	194	82	∞	99	259	80
166	1265	94	194	82	∞	102
167	250	84	430	97	74	72
168	187	78	229	81	173	81
170	471	87	∞	99	∞	95
175	212	83	711	94	315	87
177	184	75	334	87	634	90
178	115	66	125	70	145	72
179	644	93	∞	98	318	85
180	276	88	∞	103	2293	98
183	99	66	122	73	101	71
188	∞	101	1400	92	490	87
191	1405	98	1005	97	556	94
196	88	62	37	33	38	32
198	207	80	243	84	205	82
201	83	58	83	58	45	39
202	37	32	38	32	11	2
211	180	78	49	43	73	57
220	∞	105	1038	102	292	86
221	68	56	38	34	37	33
222	∞	110	1004	94	335	90

224	736	91	829	98	132	73
226	468	95	682	95	455	88
228	130	71	767	96	395	91
229	103	64	196	81	109	69
232	97	65	73	59	70	50
243	433	88	1301	97	302	88
246	95	65	168	81	109	69
251	181	77	93	63	181	83
257	1297	96	∞	102	∞	103

Pharmacokinetic Assays in Mouse and Rat

[0514] The following are assays that may be used to evaluate the pharmacokinetic properties of Formula (I) in mouse or rat. The assays follow a common procedure recited below.

[0515] 1. Formulations: For IV, DMSO/PEG400/30%HP- β -CD (10:20:70). For PO (oral gavage), DMSO/PEG400 (10:90). Formulations freshly prepared on the day of dosing or prior to dosing. Storage at room temperature.

[0516] 2. Test Species: For Mouse: Male CD1 strain, approximately 6-8 weeks in age, approximately 20-30 g. For Rat: Male Sprague-Dawley strain, approximately 6-8 weeks in age, approximately 200-300 g.

[0517] 3. Dose Level: For IV, 3 mg/kg. For PO, 10 mg/kg.

[0518] 4. Number of animals: 3 per assay.

[0519] 5. PK time points: For IV (plasma): 5 min, 15 min, 1, 2, 4, 6, 8, 24 h post dose. For PO (plasma): 15 min, 30 min, 1, 2, 4, 6, 8, 24 h post dose.

[0520] 6. Blood Sample Collection and Processing. Blood was collected (~0.2 mL per time point) from jugular vein. Blood of each sample was transferred into plastic micro centrifuge tubes containing anticoagulant of EDTA-K₂. Blood samples were centrifuged at 2,000 g for 5 minutes at 4°C to obtain plasma.

[0521] 7. Samples Analysis: Plasma samples were analyzed using LC/MS/MS method, and pharmacokinetic parameters (half-life: $t_{1/2}$; oral bioavailability: F; volume of distribution: V_{ss}).

[0522] Results are given below in Table 29, where “ND” means no data.

Table 29: Pharmacokinetic Properties in Rodents.

Ex.	Mouse t_{1/2} (h)	Mouse F (%)	Mouse Vss (L/kg)	Rat t_{1/2} (h)	Rat F (%)	Rat Vss (L/kg)
14	5.4	100	0.79	3.8	64	1.3
110	26.7	100	0.81	4.7	84	0.34
150	4.9	89	9.9	3.6	43	1.2
161	ND	ND	ND	5.7	61	0.32
167	1.7	24	2.6	2.2	72	1.2
170	2.6	100	2.0	2.4	18	1.1
175	ND	ND	ND	4.2	48	0.40
179	ND	ND	ND	2.2	76	0.30
180	ND	ND	ND	4.4	49	0.39
191	ND	ND	ND	6.3	97	1.0
211	1.9	57	1.5	3.6	55	3.6
220	6.9	100	1.7	2.8	29	0.68
224	ND	ND	ND	2.6	53	1.3
228	4.4	100	2.0	3.1	94	1.3
229	4.2	100	1.9	2.7	100	1.0
232	3.1	56	0.21	2.6	52	0.15
246	ND	ND	ND	4.7	52	1.7
251	ND	ND	ND	3.0	98	0.38
257	3.6	76	2.8	2.7	76	1.6
260	5.2	86	0.22	3.7	85	0.20
264	2.7	94	1.1	4.6	100	2.5
266	3.8	100	0.36	2.7	100	0.23
267	3.3	64	3.9	2.6	54	7.7
268	2.7	77	0.77	2.2	57	0.76
269	3.3	100	1.3	ND	ND	ND

Adjuvant-Induced Arthritis Model in Rat

[0523] The following assays are used to evaluate anti-inflammatory properties of Formula (I) in rat. The assays follow a common procedure recited below.

[0524] 1. Formulation: 0.5% Methylcellulose in saline. Formulation freshly prepared on day of dosing.

[0525] 2. Test Species: SD rat, approximately 6-8 weeks in age, approximately 200 g.

[0526] 3: Arthritis Induction Procedure: Complete Freund's Adjuvant (CFA; 50 μ L/rat, *M. Tuberculosis* at 4 mg/mL) was administered via left paw sub-plantar injection.

[0527] 4: Dosing: PO (oral gavage), 3 days post CFA injection.

[0528] 5: Body Weight. Body weights of all animals were recorded during the course of study, at Day 1 and Day 3.

[0529] 6: Difference of Weight Bearing Test (DWB). Distribution of body weight of rats between CFA-injected and contra-lateral paws were measured by weight balance changing instrument. Animals were tested to register the weight load exerted by the hind paws by means of a force plate inserted in the floor. The mean weight bearing (in g) between CFA injected paw and contra-lateral paw were determined over 10 seconds. Measurements were taken three times, including baseline prior to dose, 1 hour, and 4 hours post dosing on Day 3.

[0530] 7: Mechanical Allodynia Test (MA). Mechanical allodynia of the left hind paw will be measured during course of study by determining withdrawal thresholds to Von Frey filament. The filament was applied perpendicularly to the plantar surface of the paw with increasing force. The threshold for paw withdrawal was calculated by taking the average of 2-3 repeated stimuli (in g) which induced a reflex paw withdrawal. Measurements were taken three times, including baseline prior to dose, 1 hour, and 4 hours post dosing on Day 3.

[0531] 8: Minimal Efficacious Dose was determined by comparison of Formula (I) compounds to positive control Dexamethasone.

Table 31: Efficacy in Rat Adjuvant-Induced Arthritis Model

Ex.	Difference in Weight Bearing, Minimal Efficacious Dose (mg/kg)	Mechanical Allodynia (Von Frey), Minimal Efficacious Dose (mg/kg)

14	30	30
110	10	10
150	10	10
161	≤ 10	≤ 10
170	3	3
175	> 10	> 10
179	≤ 10	≤ 10
180	> 10	> 10
191	> 10	> 10
220	3	3
224	≤ 10	≤ 10
229	3	3
232	1	1
260	1	1
264	1	1
266	3	3
267	3	3
268	1	1


[0532] All references, patents or applications, U.S. or foreign, cited in the application are hereby incorporated by reference as if written herein in their entireties. Where any inconsistencies arise, material literally disclosed herein controls.

[0533] From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

CLAIMS


What is claimed is:

1. A compound of structural Formula II:




or a salt thereof, wherein:

L is chosen from a bond and methylene;




Y is chosen from:



R^{2e} is C₁-C₄alkyl;

Z is chosen from:




n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);


R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and

R⁹ is chosen from halo and C₁-C₄alkoxy.



2. The compound as recited in claim 1, wherein W is R_5R_4 .
3. The compound as recited in claim 2, wherein R^9 is chosen from chloro and ethoxy.
4. The compound as recited in claim 3, wherein L is a bond.
5. The compound as recited in claim 4, wherein:



R^{2c} is chosen from C₁-C₄alkoxy and C₁-C₄cycloalkoxy; and


R^{2d} is chosen from null and C₁-C₄alkoxy.

6. The compound as recited in claim 5, wherein:

R^{2c} is chosen from methoxy, methylpropoxy, cyclopropoxy, and cyclobutoxy; and

R^{2d} is chosen from null and methoxy.


7. The compound as recited in claim 6, wherein:



R^{2e} is C₁-C₄alkyl.

8. The compound as recited in claim 7, wherein R^{2e} is chosen from ethyl and propyl.

9. A compound of structural Formula I





or a salt thereof, wherein:

A^1 , A^2 , and A^3 are independently chosen from N and C, wherein at least one of A^1 ,

A^2 , and A^3 is N;

L is chosen from a bond, methylene, and ethylene;

W is chosen from , , , ,



X is H or null, or is chosen from alkenylamino, alkyl, aminoalkenyl, and aminoalkyl, any of which is optionally substituted with one to three R^1 groups;

Y is chosen from alkenyl, alkenylamino, alkyl, aminoalkenyl, aminoalkyl, aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R² groups, or

X and Y together with the atoms to which they are attached may form an aryl, cycloalkyl, heteroaryl, or heterocycloalkyl ring, any of which is optionally substituted with one to three R⁷ groups; and

Z is chosen from alkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl, any of which is optionally substituted with one to three R³ groups;

each R¹ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, heteroaryl;

each R² is independently chosen from alkyl, alkenyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, or

two R², together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

each R³ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R⁸ groups, or

two R³, together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R⁴ and R⁵ are independently chosen from H and C₁-C₆alkyl, wherein R⁴ and R⁵ together comprise no more than 6 carbons, or


R⁴ and R⁵, together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring;

R⁶ is chosen from H and C₁-C₄alkyl;

each R⁷ is independently chosen from alkyl, alkenyl, alkoxy, haloalkyl, haloalkoxy, amino, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, aryl, and heteroaryl; and

each R⁸ is independently chosen from cyano, halo, hydroxy, and oxo.


10. The compound as recited in claim 9, or a salt thereof, wherein exactly one of A¹, A², and A³ is N.
11. The compound as recited in claim 10, or a salt thereof, wherein A¹ and A² are C; and A³ is N.
12. The compound as recited in claim 11, or a salt thereof, wherein X is hydrogen.
13. The compound as recited in claim 12, or a salt thereof, wherein



R⁴ and R⁵ are independently chosen from H and C₁-C₆alkyl, with R⁴ and R⁵ together having no more than 6 carbons, or


R⁴ and R⁵, together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring.

14. The compound as recited in claim 12, or a salt thereof, wherein:
- Y is chosen from aryl and heteroaryl, any of which is optionally substituted with one to three R² groups; and
- each R² is independently chosen from alkyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, carboxamido, sulfonamido, halo, aryl, and heteroaryl.
15. The compound as recited in claim 9, having structural Formula II:



or a salt thereof, wherein:

L is chosen from a bond, methylene, and ethylene;



Y is chosen from aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R^2 groups;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^3 groups;

each R^2 is independently chosen from alkyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, or

two R^2 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

each R^3 is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R^8 groups, or

two R^3 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R^4 and R^5 are independently chosen from H and C_1 - C_6 alkyl, wherein R^4 and R^5 together comprise no more than 6 carbons and wherein at least one of R^4 and R^5 is C_1 - C_6 alkyl, or

R^4 and R^5 , together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring;

R^6 is chosen from H and C_1 - C_4 alkyl; and


each R^8 is independently chosen from cyano, halo, hydroxy, and oxo.

16. The compound as recited in claim 15, or a salt thereof, wherein:

each R^2 is independently chosen from alkyl, alkoxy, alkoxyalkyl, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, and halo, or


two R^2 , together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring.

17. The compound as recited in claim 16, or a salt thereof, wherein Z is chosen from phenyl, pyridinyl, pyridazinyl, pyrimidinyl, and pyrazinyl, any of which is optionally substituted with one or two R³ groups.
18. The compound as recited in claim 17, or a salt thereof, wherein each R³ is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, amino, aminoalkyl, and halo, and is optionally substituted with one or two R⁸ groups.
19. The compound as recited in claim 18, or a salt thereof, wherein Y is heteroaryl, and is substituted with one or two R² groups.
20. The compound as recited in claim 19, or a salt thereof, wherein Y is chosen from pyridinyl, thienyl, furyl, pyrrolyl, oxazolyl, thiazolyl, pyrazolyl, imidazolyl, triazolyl, benzothiophenyl, benzofuranyl, quinolinyl, isoquinolinyl, cinnolinyl, indazolyl, and indolyl, any of which is optionally substituted with one or two R² groups.
21. The compound as recited in claim 20, wherein Y is chosen from thienyl, furyl, benzothiophenyl, benzofuranyl, indazolyl, and indolyl and is optionally substituted with one or two R² groups.
22. The compound as recited in claim 21, wherein each R³ is independently chosen from alkyl, haloalkyl, alkoxy, and haloalkoxy.
23. The compound as recited in claim 22, wherein each R² is independently chosen from alkyl, alkoxy, cycloalkoxy, cycloalkylmethoxy, and halo.
24. The compound as recited in claim 23 wherein



R^{3a} is chosen from H, alkyl, alkoxy, haloalkyl, haloalkoxy, amino, aminoalkyl, halo, cyano, and hydroxy.

25. The compound as recited in claim 24, wherein



R⁴ and R⁵ are independently chosen from H, methyl, and ethyl, and wherein at most one of R⁴ and R⁵ is H.

26. The compound as recited in claim 15, wherein R⁴ and R⁵ are independently chosen from C₁-C₆alkyl, wherein R⁴ and R⁵ together comprise no more than 6 carbons.
27. The compound as recited in claim 26, wherein exactly one of R⁴ and R⁵ is methyl.

28. The compound as recited in claim 27, wherein

Y is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^2 groups; and

each R^2 is independently chosen from alkyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, cycloalkyl, and heterocycloalkyl.


29. The compound as recited in claim 28, wherein each R^3 is independently chosen from alkyl, alkoxy, cycloalkoxy, amino, cycloalkyl, and heterocycloalkyl.

30. The compound as recited in claim 29, wherein Y is phenyl, and is optionally substituted with one or two R^2 groups.

31. The compound as recited in claim 30, wherein each R^2 is independently chosen from alkoxy, cycloalkoxy, and cycloalkylmethoxy.

32. The compound as recited in claim 29, wherein Z is phenyl, and is optionally substituted with one or two R^3 groups.

33. The compound as recited in claim 15, having structural Formula III:



or a salt thereof, wherein:

L is chosen from a bond, methylene, and ethylene;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one or two R^3 groups;

R^{2a} and R^{2b} are independently chosen from H, alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, halo, cycloalkyl, and heterocycloalkyl;

each R^3 is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, amino, aminoalkyl, carboxamido, halo, cyano, hydroxy, and cycloalkyl, and is optionally substituted with one or two R^8 groups;


R^4 and R^5 are independently chosen from H and C_1 - C_6 alkyl, wherein R^4 and R^5 together comprise no more than 6 carbons and wherein at least one of R^4 and R^5 is C_1 - C_6 alkyl, or

R^4 and R^5 , together with the atoms to which they are attached, form a 3-7 membered cycloalkyl or heterocycloalkyl ring;

R^6 is chosen from H and methyl; and


each R^8 is independently chosen from cyano, halo, hydroxy, and oxo.

34. The compound as recited in claim 33, wherein Z is chosen from phenyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinolinyl, isoquinolinyl, cinnolinyl, indazolyl, and indolyl, any of which is optionally substituted with one or two R^3 groups.
35. The compound as recited in claim 34 wherein R^{2a} is selected from alkyl, alkoxy, haloalkyl, haloalkoxy, and halo.
36. The compound as recited in claim 35, wherein each R^3 is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, amino, aminoalkyl, halo, cyano, and hydroxy.
37. The compound as recited in claim 36, wherein



R^4 and R^5 are independently chosen from H, methyl, and ethyl, and wherein at most one of R^4 and R^5 is H.


38. The compound as recited in claim 37, wherein Z is phenyl, and is optionally substituted with one or two R^3 groups.
39. The compound as recited in claim 38, wherein



R^{3a} is chosen from H, alkyl, alkoxy, haloalkyl, haloalkoxy, amino, aminoalkyl, halo, cyano, and hydroxy.


40. The compound as recited in claim 37, wherein Z is chosen from pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinolinyl, isoquinolinyl, cinnolinyl, indazolyl, and indolyl, any of which is optionally substituted with one or two R^3 groups.
41. The compound as recited in claim 40, wherein Z is chosen from quinolinyl, isoquinolinyl, cinnolinyl, indazolyl, and indolyl, any of which is optionally substituted with one or two R^3 groups.
42. The compound as recited in claim 41, wherein Z is chosen from indazolyl and indolyl, any of which is optionally substituted with one or two R^3 groups.
43. The compound as recited in claim 35, wherein R^{2b} is chosen from H and alkoxy.
44. The compound as recited in claim 43, wherein R^{2a} is alkoxy.

45. The compound as recited in claim 44, wherein each R^3 is independently chosen from alkyl, alkoxy, amino, aminoalkyl, haloalkyl, haloalkoxy, halo, cyano, and hydroxy.
46. The compound as recited in claim 45, wherein R^4 and R^5 are independently chosen from H, methyl, and ethyl, and wherein at most one of R^4 and R^5 is H.
47. The compound as recited in claim 46, wherein Z is phenyl, and is optionally substituted with one or two R^3 groups.
48. The compound as recited in claim 47, wherein







R^{3a} is chosen from H, alkyl, alkoxy, amino, aminoalkyl, haloalkyl, haloalkoxy, halo, cyano, and hydroxy.





49. The compound as recited in claim 48, wherein Z is chosen from indazolyl and indolyl, any of which is optionally substituted with one or two R^3 groups.
50. The compound as recited in claim 9, wherein the compound has structural Formula VII:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

W is chosen from , , , ,

, , , and ;

Y is chosen from alkenyl, alkenylamino, alkyl, aminoalkenyl, aminoalkyl, aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R^2 groups;

each R^2 is independently chosen from alkyl, alkenyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy,

amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl;


n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl; and

R⁶ is chosen from H and C₁-C₄alkyl.

51. The compound as recited in claim 50, wherein Y is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R² groups.

52. The compound as recited in claim 51, wherein each R² is independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.




53. The compound as recited in claim 52, wherein W is

54. The compound as recited in claim 53, wherein n is 1, forming azetidin-1-yl.

55. The compound as recited in claim 54, wherein L is a bond.

56. The compound as recited in claim 9, wherein compound has structural Formula VIII:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

Y is chosen from phenyl, indazolyl, thienyl, indazolyl, and indolyl, any of which is optionally substituted with one or two R² groups;

n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);


each R² is independently chosen from alkyl, alkoxy, alkylthio, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, and halo;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl; and

R⁶ is chosen from H and C₁-C₄alkyl.

57. The compound as recited in claim 56, wherein Y is chosen from phenyl and indazolyl, either of which is optionally substituted with one or two R² groups.

58. The compound as recited in claim 57, wherein each R^2 is independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.
59. The compound as recited in claim 58, wherein n is 1, forming azetidin-1-yl.
60. The compound as recited in claim 59, wherein L is a bond.
61. The compound as recited in claim 60, wherein Y is phenyl substituted with one or two R^2 groups independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.
62. The compound as recited in claim 61, wherein each R^2 is independently chosen from C_1 - C_4 alkoxy, cyclopropoxy, cyclobutoxy, and C_1 - C_3 alkyl.
63. The compound as recited in claim 60, wherein Y is indazolyl substituted with an R^2 groups chosen from methyl, ethyl, and propyl.
64. The compound as recited in claim 60, wherein Y is chosen from:




R^{2c} is chosen from C_1 - C_4 alkoxy and C_1 - C_4 cycloalkoxy;

R^{2d} is chosen from null and C_1 - C_4 alkoxy; and




R^{2e} is C_1 - C_4 alkyl.


65. The compound as recited in claim 9, wherein the compound has structural Formula IX:



or a salt thereof, wherein:

L is chosen from a bond and methylene;

W is chosen from , , , and



Y is chosen from aryl, arylmethyl, arylamino, aryloxy, cycloalkyl, heterocycloalkyl, and heteroaryl, any of which is optionally substituted with one to three R^2 groups;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^3 groups;


each R^2 is independently chosen from alkyl, alkoxy, alkoxyalkyl, alkylthio, haloalkyl, haloalkoxy, cycloalkoxy, cycloalkylmethoxy, heterocycloalkylmethoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl;

R^4 and R^5 are independently chosen from C_1 - C_3 alkyl; and

R^6 is chosen from H and C_1 - C_4 alkyl.

66. The compound as recited in claim 65, wherein Y is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R^2 groups.

67. The compound as recited in claim 66, wherein each R^2 is independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.




68. The compound as recited in claim 67, wherein W is

69. The compound as recited in claim 68, wherein L is a bond.


70. The compound as recited in claim 69, wherein Y is phenyl substituted with one or two R^2 groups independently chosen from alkoxy, cycloalkoxy, haloalkoxy, alkyl, halo, and haloalkyl.

71. The compound as recited in claim 70, wherein each R^2 is independently chosen from C_1 - C_4 alkoxy, cyclopropoxy, cyclobutoxy, and C_1 - C_3 alkyl.

72. The compound as recited in claim 9, wherein the compound has structural Formula X:



or a salt thereof, wherein:



R^{2e} is C_1 - C_4 alkyl;

L is chosen from a bond and methylene;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R³ groups;


each R³ is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R⁸ groups, or

two R³, together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and

each R⁸ is independently chosen from cyano, halo, hydroxy, and oxo.




73. The compound as recited in claim 72, wherein W is

74. The compound as recited in claim 73, wherein L is a bond.

75. The compound as recited in claim 74, wherein Z is chosen from phenyl and indazolyl, either of which is optionally substituted with one to three R³ groups.

76. The compound as recited in claim 75, wherein Z is chosen from:



n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and


R⁹ is chosen from halo, amino, and C₁-C₄alkoxy.

77. The compound as recited in claim 76, wherein R⁹ is chosen from chloro, dimethylamino, and ethoxy.


78. The compound as recited in claim 77, wherein R⁹ is chosen from chloro and ethoxy.

79. The compound as recited in claim 76, wherein n is 1, forming azetidin-1-yl.

80. The compound as recited in claim 9, wherein the compound has structural Formula XI:



or a salt thereof, wherein:



n is 1, 2, or 3 (forming a cyclopropyl, cyclobutyl, or cyclopentyl);

L is chosen from a bond and methylene;

Z is chosen from aryl and heteroaryl, either of which is optionally substituted with one to three R³ groups;


each R³ is independently chosen from alkyl, alkoxy, haloalkyl, haloalkoxy, cycloalkoxy, amino, aminoalkyl, carboxamido, sulfonamido, halo, cyano, hydroxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, and is optionally substituted with one to three R⁸ groups, or

two R³, together with the intervening atoms, form a 5-7 membered cycloalkyl or heterocycloalkyl ring;

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and

each R⁸ is independently chosen from cyano, halo, hydroxy, and oxo.




81. The compound as recited in claim 80, wherein W is

82. The compound as recited in claim 81, wherein L is a bond.

83. The compound as recited in claim 82, wherein Z is chosen from phenyl and indazolyl, either of which is optionally substituted with one to three R³ groups.

84. The compound as recited in claim 83, wherein Z is chosen from:




n is 1 or 2 (forming an azetidin-1-yl or pyrrolidin-1-yl);

R⁴ and R⁵ are independently chosen from C₁-C₃alkyl;

R⁶ is chosen from H and C₁-C₄alkyl; and

R⁹ is chosen from halo, amino, and C₁-C₄alkoxy.



R^{2c} is chosen from C₁-C₄alkoxy and C₁-C₄cycloalkoxy; and


R^{2d} is chosen from null and C₁-C₄alkoxy.

91. The compound as recited in claim 90, wherein:

R^{2c} is chosen from methoxy, methylpropoxy, cyclopropoxy, and cyclobutoxy; and

R^{2d} is chosen from null and methoxy.


92. The compound as recited in claim 89 wherein:



R^{2e} is C₁-C₄alkyl.


93. The compound as recited in claim 92, wherein R^{2e} is chosen from ethyl and propyl.

94. The compound as recited in claim 89, wherein L is a bond.




95. The compound as recited in claim 94, wherein Z is

96. The compound as recited in claim 95, wherein n is 1, forming azetidin-1-yl.




97. The compound as recited in claim 94, wherein Z is





98. The compound as recited in claim 89, wherein Z is


99. The compound as recited in claim 98, wherein R^9 is chosen from chloro and ethoxy.


100. A compound chosen from:





























101. An isolated enantiomer of a compound as recited in claim 100.

102. A pharmaceutical composition comprising a compound as recited in any of claims 1-101, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.
103. A method for inhibiting activity of the monocarboxylate transporter MCT4, or a mutant thereof, in a biological sample comprising the step of contacting said biological sample with a compound as recited in any of claims 1-101, or a salt thereof.
104. A method for inhibiting activity of the monocarboxylate transporter MCT4, or a mutant thereof, in a patient comprising the step of administering to the patient a compound as recited in any of claims 1-101, or a salt thereof.
105. A method for selectively inhibiting activity of the monocarboxylate transporter MCT4, or a mutant thereof, over the monocarboxylate transporter MCT1, or a mutant thereof, in a patient comprising the step of administering to the patient a compound as recited in any of claims 1-101, or a salt thereof.
106. The method as recited in claim 105 wherein the inhibition is at least 100-fold selective for MCT4 over MCT1.
107. A method for treating a monocarboxylate transporter MCT4-mediated disorder in a subject in need thereof, comprising the step of administering to said patient a compound as recited in any of claims 1-101, or a pharmaceutically acceptable salt thereof.
108. The method as recited in claim 107, wherein the subject is a human.
109. The method as recited in claim 107, wherein the monocarboxylate transporter MCT4-mediated disorder is chosen from an inflammatory disorder and a proliferative disorder.
110. The method as recited in claim 107, wherein the monocarboxylate transporter MCT4-mediated disorder is a proliferative disorder.
111. The method as recited in claim 110, wherein the proliferative disorder is cancer.
112. The method as recited in claim 111, wherein the cancer is chosen from adenocarcinoma, adult T-cell leukemia/lymphoma, bladder cancer, blastoma, bone cancer, breast cancer, brain cancer, carcinoma, myeloid sarcoma, cervical cancer, colorectal cancer, esophageal cancer, gastrointestinal cancer, glioblastoma multiforme, glioma, gallbladder cancer, gastric cancer, head and neck cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, intestinal cancer, kidney cancer, laryngeal cancer, leukemia, lung cancer, lymphoma, liver cancer, small cell lung cancer, non-small cell lung cancer, mesothelioma, multiple myeloma, ocular cancer, optic nerve tumor, oral cancer, ovarian cancer, pituitary tumor, primary central nervous system lymphoma, prostate cancer,

pancreatic cancer, pharyngeal cancer, renal cell carcinoma, rectal cancer, sarcoma, skin cancer, spinal tumor, small intestine cancer, stomach cancer, T-cell lymphoma, testicular cancer, thyroid cancer, throat cancer, urogenital cancer, urothelial carcinoma, uterine cancer, vaginal cancer, and Wilms' tumor.

113. The method as recited in claim 107, wherein the monocarboxylate transporter MCT4-mediated disorder is an inflammatory disorder.

114. The method as recited in claim 113, wherein the inflammatory disorder is chosen from Crohn's disease, ulcerative colitis, idiopathic pulmonary fibrosis, muscular dystrophy, rheumatoid arthritis, and systemic sclerosis (scleroderma).

115. A method of treating a monocarboxylate transporter MCT4-mediated disorder in a subject in need thereof, comprising the sequential or co-administration of a compound as recited in any of claims 1-101 or a pharmaceutically acceptable salt thereof, and another therapeutic agent.

116. The method as recited in claim 115, wherein the therapeutic agent is a protein kinase inhibitor.

117. The method as recited in claim 116, wherein the protein kinase inhibitor is chosen from Aurora B, EGFR, PLK-1, CDKs inhibitors.

118. The method as recited in claim 115, wherein the therapeutic agent is chosen from an antimetabolite, bcr-abl inhibitor, DNA damaging agent, EGFR inhibitor, microtubule stabilizing inhibitor, mitotic arrest inhibitor, S-phase inhibitor, and a taxane.

119. The method as recited in claim 118, wherein the therapeutic agent is a DNA damaging agent chosen from an alkylating agent, anthracycline, antimetabolite agent, crosslinking agent, DNA replication inhibitor, intercalator, microtubule disruptor, PARP inhibitor, radiomimetic agent, radiosensitizer, strand break agent, and topoisomerase II inhibitor.

120. The method as recited in claim 115, wherein the therapeutic agent is chosen from aminoglutethimide, amsacrine, anastrozole, asparaginase, barasertib, bcr, bicalutamide, bleomycin, buserelin, busulfan, camptothecin, capecitabine, carboplatin, carmustine, chlorambucil, chloroquine, cisplatin, cladribine, clodronate, colchicine, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, demethoxyviridin, dichloroacetate, dienestrol, diethylstilbestrol, docetaxel, doxorubicin, epirubicin, estradiol, estramustine, etoposide, everolimus, exemestane, filgrastim, fludarabine, fludrocortisone, fluorouracil, fluoxymesterone, flutamide, gemcitabine,


genistein, goserelin, hydroxyurea, idarubicin, ifosfamide, imatinib, interferon, irinotecan, ironotecan, letrozole, leucovorin, leuprolide, levamisole, lomustine, lonidamine, mechlorethamine, medroxyprogesterone, megestrol, melphalan, mercaptopurine, mesna, metformin, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, nocodazole, olaparib, octreotide, oxaliplatin, paclitaxel, pamidronate, pentostatin, perifosine, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, sorafenib, streptozocin, sunitinib, suramin, tamoxifen, temozolomide, temsirolimus, teniposide, testosterone, thioguanine, thiotepa, titanocene dichloride, topotecan, trastuzumab, tretinoin, vinblastine, vincristine, vindesine, and vinorelbine.

121. The method as recited in claim 107 wherein the method further comprises administering non-chemical methods of cancer treatment.
122. The method as recited in claim 121, wherein the method further comprises administering radiation therapy.
123. The method as recited in claim 121, wherein the method further comprises administering surgery, thermoablation, focused ultrasound therapy, cryotherapy, or any combination thereof.
124. A compound as recited in any of claims 1-101, or a pharmaceutically acceptable salt thereof, for use in human therapy.
125. A compound as recited in any of claims 1-101, or a pharmaceutically acceptable salt thereof, for use in treating a monocarboxylate transporter MCT4-mediated disorder.
126. Use of a compound as recited in any of claims 1-101, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament to treat a monocarboxylate transporter MCT4-mediated disorder.
127. The method as recited in claim 107, wherein the monocarboxylate transporter MCT4-mediated disorder is a metabolic disease.
128. The method as recited in Claim 127 wherein said disease is a metabolic disease.
129. The method as recited in claim 128 wherein said metabolic disease is chosen from metabolic syndrome, diabetes, dyslipidemia, fatty liver disease, non-alcoholic steatohepatitis, obesity, and insulin resistance.
130. The method of claim 129 wherein said diabetes is Type II diabetes.
131. The method of claim 130 wherein said dyslipidemia is hyperlipidemia.
132. A method for achieving an effect in a patient comprising the administration of a therapeutically effective amount of a compound as recited in any of claims 1-101 to a

patient, wherein the effect is selected from the group consisting of reduction of triglycerides, reduction of cholesterol, and reduction of hemoglobin A1c.

133. The method of claim 132 wherein said cholesterol is chosen from LDL and VLDL cholesterol.

134. The method of claim 133 wherein said triglycerides are chosen from plasma triglycerides and liver triglycerides.



(I)