

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2008216600 B2

(54) Title
Antibodies against ErbB3 and uses thereof

(51) International Patent Classification(s)
A61K 39/395 (2006.01) **A61P 35/00** (2006.01)

(21) Application No: **2008216600** (22) Date of Filing: **2008.02.15**

(87) WIPO No: **WO08/100624**

(30) Priority Data

(31) Number (32) Date (33) Country
60/901,904 **2007.02.16** **US**
61/009,796 **2008.01.02** **US**

(43) Publication Date: **2008.08.21**
(44) Accepted Journal Date: **2013.09.05**

(71) Applicant(s)
Merrimack Pharmaceuticals, Inc.

(72) Inventor(s)
Nielsen, Ulrik; Schoeberl, Birgit; Feldhaus, Michael; Muruganandam, Arumugam; Buckler, David

(74) Agent / Attorney
Shelston IP, L 21 60 Margaret St, Sydney, NSW, 2000

(56) Related Art
EP 1 283 053 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 August 2008 (21.08.2008)

PCT

(10) International Publication Number
WO 2008/100624 A3

(51) International Patent Classification:
A61K 39/395 (2006.01) A61P 35/00 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2008/002119

(22) International Filing Date:
15 February 2008 (15.02.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/901,904 16 February 2007 (16.02.2007) US
61/009,796 2 January 2008 (02.01.2008) US

(71) Applicant (for all designated States except US): MER-RIMACK PHARMACEUTICALS, INC. [—/US]; One Kendall Square, Building 700, 2nd Floor, Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHOEBERL, Birgit [DE/US]; 324 Windsor Street, Cambridge, MA 02141 (US). NIELSEN, Ulrik [DK/US]; 18 Thomas Street, Quincy, MA 02169 (US). FELDHAUS, Michael [US/US]; P.O. Box 1888, Grantham, NH 03753 (US).

(74) Agents: REMILLARD, Jane, E. et al.; Lahive & Cockfield, LLP, One Post Office Square, Boston, MA 02109-2127 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
15 January 2009



A3

WO 2008/100624 A3

(54) Title: ANTIBODIES AGAINST ERBB3 AND USES THEREOF

(57) Abstract: The present invention provides a novel class of monoclonal antibodies which bind ErbB3 receptor and inhibits various ErbB3 functions. For example, the antibodies described herein are capable of binding to ErbB3 and inhibiting EGF-like ligand mediated phosphorylation of the receptor.

ANTIBODIES AGAINST ERBB3 AND USES THEREOF

Background of the Invention

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

The ErbB/HER subfamily of polypeptide growth factor receptors include the epidermal growth factor (EGF) receptor (EGFR, ErbB1/HER1), the *neu* oncogene product (ErbB2/HER2), and the more recently identified ErbB3/HER3 and ErbB4/HER4 receptor proteins (see, e.g., Hynes *et. al.* (1994) *Biochim. Biophys. Acta Rev. Cancer* 1198, 165–184). Each of these receptors is predicted to consist of an extracellular ligand-binding domain, a membrane-spanning domain, a cytosolic protein tyrosine kinase (PTK) domain and a C-terminal phosphorylation domain (see, e.g., Kim *et al.*, (1998) *Biochem. J.* 334, 189-195).

Experiments *in vitro* have indicated that the protein tyrosine kinase activity of the ErbB3 protein is attenuated significantly relative to that of other ErbB/HER family members and this attenuation has been attributed, in part, to the occurrence of non-conservative amino acid substitutions in the predicted catalytic domain of ErbB3 (see, e.g., Guy *et al.* (1994) *Proc. Natl. Acad. Sci. USA.* 91, 8132–8136; Sierke *et al.* (1997) *Biochem. J.* 322, 757–763). However, the ErbB3 protein has been shown to be phosphorylated in a variety of cellular contexts. For example, ErbB3 is constitutively phosphorylated on tyrosine residues in a subset of human breast cancer cell lines overexpressing this protein (see, e.g., Kraus *et al.* (1993) *Proc. Natl. Acad. Sci. USA.* 90, 2900–2904; and Kim *et al.* *Supra*; see, also, Schaefer *et al.* (2006) *Neoplasia* 8(7):613-22 and Schaefer *et al.* *Cancer Res* (2004) 64(10):3395-405).

Although, the role of ErbB3 in cancer has been explored (see, e.g., Horst *et al.* (2005) 115, 519-527; Xue *et al.* (2006) *Cancer Res.* 66, 1418-1426), ErbB3 remains largely unappreciated as a target for clinical intervention. Current immunotherapies primarily focus on inhibiting the action of ErbB2 and, in particular, heterodimerization of ErbB2/ErbB3 complexes (see, e.g., Sliwkowski *et al.* (1994) *J. Biol. Chem.* 269(20):14661-14665 (1994)). Accordingly, in one or more preferred embodiments, the present invention provides improved immunotherapies that effectively inhibit ErbB3 signaling, and can be used to treat and diagnose a variety of cancers.

It is an object of the present invention to overcome or ameliorate at least one of the disadvantages of the prior art, or to provide a useful alternative.

Summary of the Invention

5 According to a first aspect, the present invention provides an isolated monoclonal antibody, or antigen binding portion thereof, comprising: a heavy chain variable region CDR1 comprising SEQ ID NO:7; a heavy chain variable region CDR2 comprising SEQ ID NO:8; a heavy chain variable region CDR3 comprising SEQ ID NO:9; a light chain variable region CDR1 comprising SEQ ID NO:10; a light chain variable region CDR2 comprising SEQ ID NO:11; and a light chain variable region CDR3 comprising SEQ ID NO:12.

10

According to a second aspect, the present invention provides an isolated monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 comprising:

15 (a) a heavy chain variable region CDR1 comprising SEQ ID NO:7 or an amino acid sequence which is at least 90% identical thereto;

(b) a heavy chain variable region CDR2 comprising SEQ ID NO:8 or an amino acid sequence which is at least 90% identical thereto;

(c) a heavy chain variable region CDR3 comprising SEQ ID NO:9 or an amino acid sequence which is at least 90% identical thereto;

20 (d) a light chain variable region CDR1 comprising SEQ ID NO:10 or an amino acid sequence which is at least 90% identical thereto;

(e) a light chain variable region CDR2 comprising SEQ ID NO:11 or an amino acid sequence which is at least 90% identical thereto; and

25 (f) a light chain variable region CDR3 comprising SEQ ID NO:12 or an amino acid sequence which is at least 90% identical thereto.

According to a third aspect, the present invention provides an isolated monoclonal antibody, or antigen binding portion thereof, comprising:

30 (a) a heavy chain variable region CDR1 comprising SEQ ID NO:7 or conservative sequence modifications thereof;

(b) a heavy chain variable region CDR2 comprising SEQ ID NO:8 or conservative sequence modifications thereof;

- 1b -

- (c) a heavy chain variable region CDR3 comprising SEQ ID NO:9 or conservative sequence modifications thereof;
- (d) a light chain variable region CDR1 comprising SEQ ID NO:10 or conservative sequence modifications thereof;
- 5 (e) a light chain variable region CDR2 comprising SEQ ID NO:11 or conservative sequence modifications thereof; and
- (f) a light chain variable region CDR3 comprising SEQ ID NO:12 or conservative sequence modifications thereof.

According to a fourth aspect, the present invention provides an isolated
10 monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises heavy and light chain variable regions, wherein the heavy chain variable region comprises an amino acid sequence at least 95% identical to the heavy chain variable region amino acid sequence set forth in SEQ ID NO:1

According to a fifth aspect, the present invention provides an isolated
15 monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises heavy and light chain variable regions, wherein the heavy chain variable region comprises SEQ ID NO:1.

According to a sixth aspect, the present invention provides an isolated
monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and
20 comprises heavy and light chain variable regions, wherein the light chain variable region comprises an amino acid sequence at least 95% identical to the light chain variable region amino acid sequence set forth in SEQ ID NO:2.

According to a seventh aspect, the present invention provides an isolated
monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and
25 comprises heavy and light chain variable regions, wherein the light chain variable region comprises SEQ ID NO:2.

According to an eighth aspect, the present invention provides an isolated
monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises a heavy chain variable region comprising SEQ ID NO:1 or a sequence at least
30 95% identical thereto and a light chain variable region comprising SEQ ID NO:2 or a sequence at least 95% identical thereto

According to a ninth aspect, the present invention provides an isolated
monoclonal antibody, or antigen binding portion thereof, comprising a heavy chain

variable region comprising SEQ ID NO:1 and a light chain variable region comprising SEQ ID NO:2.

According to a tenth aspect, the present invention provides an isolated antibody, or antigen binding portion thereof, that binds to an epitope which is same or overlapping 5 with the epitope bound by the antibody of the ninth aspect.

According to an eleventh aspect, the present invention provides an isolated antibody, or antigen binding portion thereof, that competes for binding to ErB3 with the antibody of the ninth aspect.

According to a twelfth aspect, the present invention provides a composition 10 comprising the antibody, or antigen binding portion thereof, of any one of the preceding aspects in a pharmaceutically acceptable carrier.

According to a thirteenth aspect, the present invention provides an isolated nucleic acid encoding a heavy chain variable region of an antibody, or antigen binding portion thereof, of the fifth aspect, wherein the nucleic acid comprises SEQ ID NO:25.

15 According to a fourteenth aspect, the present invention provides an isolated nucleic acid encoding a light chain variable region of an antibody, or antigen binding portion thereof, of the seventh aspect, wherein the nucleic acid comprises SEQ ID NO:26.

According to a fifteenth aspect, the present invention provides an isolated nucleic 20 acid encoding a heavy chain variable region of the antibody, or antigen binding portion thereof, of the fifth aspect comprising a sequence that hybridizes under high stringency to a nucleic acid comprising the nucleotide sequence SEQ ID NO:25.

According to a sixteenth aspect, the present invention provides an isolated nucleic acid encoding a light chain variable region of the antibody, or antigen binding 25 portion thereof, of the seventh aspect comprising a sequence that hybridizes under high stringency to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:26.

According to a seventeenth aspect, the present invention provides an expression vector comprising the nucleic acid of any one of the thirteenth to sixteenth aspects.

According to an eighteenth aspect, the present invention provides a host cell 30 comprising the vector of the seventeenth aspect or the nucleic acid of any one of thirteenth to sixteenth aspects.

According to a nineteenth aspect, the present invention provides a transgenic non-human mammal which expresses a monoclonal antibody, or antigen binding portion

thereof, that binds the same epitope as the antibody, or antigen binding portion thereof, of any one of the first to eleventh aspects.

According to a twentieth aspect, the present invention provides a transgenic plant which expresses a monoclonal antibody, or antigen binding portion thereof, that binds
5 the same epitope as the antibody, or antigen binding portion thereof, of any one of the first to eleventh aspects.

According to a twenty-first aspect, the present invention provides a host cell which produces an antibody, or antigen binding portion, thereof of any one of the first to eleventh aspects.

10 According to a twenty-second aspect, the present invention provides a kit comprising one or more isolated monoclonal antibodies, or antigen binding portions thereof, of any one of the first to eleventh aspects, and optionally comprising instructions for use in treating or diagnosing a disease associated with ErbB3 dependent signaling

15 According to a twenty-third aspect, the present invention provides a method of inhibiting ErbB3 signaling in a subject, comprising administering to the subject an isolated monoclonal antibody, or antigen binding portion thereof, of any one of the first to eleventh aspects, in an amount sufficient to inhibit ErbB3 signaling

According to a twenty-fourth aspect, the present invention provides a method of
20 treating cancer in a subject, comprising administering to the subject a therapeutically effective amount of an isolated monoclonal antibody or antigen binding portion thereof of any one of the first to eleventh aspects, in an amount sufficient to treat cancer

According to a twenty-fifth aspect, the present invention provides a method of diagnosing a cancer associated with ErbB3 in a subject, comprising (a) contacting *ex*
25 *vivo* or *in vivo* cells from the subject with an isolated monoclonal antibody, or antigen binding portion thereof, of any one of the first to eleventh aspects, and (b) measuring the level of binding to ErbB3 on the cells, wherein abnormally high levels of binding to ErbB3 indicate that the subject has a cancer associated with ErbB3


According to a twenty-sixth aspect, the present invention provides use of an
30 isolated monoclonal antibody, or antigen binding portion thereof, of any one of the first to eleventh aspects for the manufacture of a medicament for inhibiting ErbB3 signaling

- 1e -

According to a twenty-seventh aspect, the present invention provides use of an isolated monoclonal antibody, or antigen binding portion thereof, of any one of the first to eleventh aspects for the manufacture of a medicament for treating cancer

Unless the context clearly requires otherwise, throughout the description and the 5 claims, the words "comprise", "comprising", and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of "including, but not limited to".

The present invention provides a novel class of monoclonal antibodies which binds to the ErbB3 receptor and inhibits various ErbB3 functions. For example, the 10 antibodies described herein are capable of binding to ErbB3 and inhibiting EGF-like ligand mediated phosphorylation of the receptor. As described herein, EGF-like ligands include EGF, TGF- α , betacellulin, heparin-binding epidermal growth factor, biregulin —



and amphiregulin, which bind to EGFR and induce dimerization of EGFR with ErbB3. This dimerization, in turn, causes phosphorylation of ErbB3, and activates signaling through the receptor. Monoclonal antibodies of the present invention, thus, are useful for treating and diagnosing a variety of cancers associated with ErbB3-mediated cellular 5 signaling. Accordingly, in one embodiment, the present invention provides monoclonal antibodies (and antigen binding portions thereof) which bind to ErbB3 and inhibit EGF-like ligand mediated phosphorylation of ErbB3.

In another embodiment, the antibodies are further characterized by one or more of the following properties: (i) inhibition of ErbB3 ligand-mediated signaling, including 10 signaling mediated by binding of ErbB3 ligands, such as heregulin, epiregulin, epigen and BIR, to ErbB3; (ii) inhibition of proliferation of cells expressing ErbB3; (iii) the ability to decrease levels of ErbB3 on cell surfaces (e.g., by inducing internalization of ErbB3); (iv) inhibition of VEGF secretion of cells expressing ErbB3; (v) inhibition of the migration of cells expressing ErbB3; (vi) inhibition of spheroid growth of cells 15 expressing ErbB3; and/or (vii) binding to an epitope located on domain I (residues 20-209) of ErbB3, for example, an epitope involving or spanning residues 20-202 of the amino acid sequence of ErbB3.

Particular monoclonal antibodies and antigen binding portions thereof of the present invention exhibit a K_D of 50 nM or less, as measured by a surface plasmon 20 resonance assay or a cell binding assay

In further embodiments, particular monoclonal antibodies and antigen binding portions thereof of the present invention include a heavy chain variable region comprising an amino acid sequence which is at least 80% (e.g., 85%, 90%, 95%, 96%, 97%, 98% or 99%) identical to the heavy chain variable region amino acid sequence set 25 forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:35, or SEQ ID NO: 37. Other particular monoclonal antibodies and antigen binding portions thereof of the present invention include a light chain variable region comprising an amino acid sequence which is at least 80% (e.g., 85%, 90%, 95%, 96%, 97%, 98% or 99%) identical to the light chain variable region amino acid sequence set forth in SEQ ID NO:2, SEQ 30 ID NO:4, SEQ ID NO:6, SEQ ID NO:36, or SEQ ID NO:38. The antibodies may also include both of the aforementioned heavy chain and light chain variable regions.

The variable heavy and light chain regions of the antibodies or antigen binding portions thereof typically include one or more complementarity determining regions (CDRs). These include one or more CDR1, CDR2, and CDR3 regions. Accordingly, 35 other particular antibodies and antigen binding portions thereof of the present invention include one or more CDR sequences selected from a heavy chain variable region CDR1 comprising SEQ ID NO:7; a heavy chain variable region CDR2 comprising SEQ ID NO:8; a heavy chain variable region CDR3 comprising SEQ ID NO:9; a light chain

variable region CDR1 comprising SEQ ID NO:10; a light chain variable region CDR2 comprising SEQ ID NO:11; a light chain variable region CDR3 comprising SEQ ID NO:12; and combinations thereof.

Still other particular antibodies and antigen binding portions thereof of the 5 present invention include one or more CDR sequences selected from a heavy chain variable region CDR1 comprising SEQ ID NO:13; a heavy chain variable region CDR2 comprising SEQ ID NO:14; a heavy chain variable region CDR3 comprising SEQ ID NO:15; a light chain variable region CDR1 comprising SEQ ID NO:16; a light chain variable region CDR2 comprising SEQ ID NO:17; a light chain variable region CDR3 comprising SEQ ID NO:18; and combinations thereof.

Still other particular antibodies and antigen binding portions thereof of the present invention include; or one or more CDR sequences selected from a heavy chain variable region CDR1 comprising SEQ ID NO:19; a heavy chain variable region CDR2 comprising SEQ ID NO:20; a heavy chain variable region CDR3 comprising SEQ ID NO:21; a light chain variable region CDR1 comprising SEQ ID NO:22; a light chain variable region CDR2 comprising SEQ ID NO:23; a light chain variable region CDR3 comprising SEQ ID NO:24; and combinations thereof.

Still other particular antibodies and antigen binding portions thereof of the 20 present invention include; or one or more CDR sequences selected from a heavy chain variable region CDR1 comprising SEQ ID NO:39; a heavy chain variable region CDR2 comprising SEQ ID NO:40; a heavy chain variable region CDR3 comprising SEQ ID NO:41; a light chain variable region CDR1 comprising SEQ ID NO:42; a light chain variable region CDR2 comprising SEQ ID NO:43; a light chain variable region CDR3 comprising SEQ ID NO:44; and combinations thereof.

Still other particular antibodies and antigen binding portions thereof of the 25 present invention include; or one or more CDR sequences selected from a heavy chain variable region CDR1 comprising SEQ ID NO:45; a heavy chain variable region CDR2 comprising SEQ ID NO:46; a heavy chain variable region CDR3 comprising SEQ ID NO:47; a light chain variable region CDR1 comprising SEQ ID NO:48; a light chain variable region CDR2 comprising SEQ ID NO:49; a light chain variable region CDR3 comprising SEQ ID NO:50; and combinations thereof.

The antibodies and antigen binding portions thereof may also comprise one or more CDRs which are at least 80% (*e.g.*, 85%, 90%, 95%, 96%, 97%, 98% or 99%) identical to any of the aforementioned CDRs, or combinations of CDRs.

35 In one embodiment, the antibodies and antibody portions thereof are fully human (*i.e.*, contains human CDR and framework sequences). Particular human antibodies of the present invention include those having a heavy chain variable region that is from a

human VH3 germ line gene, and/or a light chain variable region from human VL2 germ line gene.

Also encompassed by the present invention are monoclonal antibodies and portions thereof that bind to the same or overlapping epitopes bound by any of the

5 antibodies or portions thereof described herein (e.g., an epitope located on domain I of ErbB3, such as an epitope involving or spanning, residues 20-202 of the amino acid sequence of ErbB3). Antibodies which have the same activity as the antibodies described herein, e.g., antibodies having the same sequence as Ab #6, are also encompassed by the present invention.

10 Antibodies of the present invention include all known forms of antibodies and other protein scaffolds with antibody-like properties. For example, the antibody can be a human antibody, a humanized antibody, a bispecific antibody, a chimeric antibody or a protein scaffold with antibody-like properties, such as fibronectin or Ankyrin repeats. The antibody also can be a Fab, Fab'2, ScFv, SMIP, affibody, nanobody, or a domain

15 antibody. The antibody also can have any of the following isotypes: IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgAsec, IgD, and IgE.

In yet another embodiment, the present invention further provides compositions comprising combinations of antibodies or antigen binding portions described herein, formulated with an acceptable carrier and/or adjuvant. In a particular embodiment, the

20 composition comprises two or more antibodies that bind different epitopes on ErbB3 or antibodies described herein combined with anti-cancer antibodies which do not bind ErbB3.

In still another embodiment, the present invention provides isolated nucleic acids encoding the antibodies and antigen binding portions thereof described herein. In

25 particular embodiments, the nucleic acid encodes a heavy chain variable region comprising a nucleotide sequence which is at least 80% (e.g., 85%, 90%, 95%, 96%, 97%, 98% or 99%) identical to, or which hybridizes under high stringency conditions to, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:35, or SEQ ID NO:37; or a light chain variable region comprising a nucleotide sequence which is at least 80%

30 (e.g., 85%, 90%, 95%, 96%, 97%, 98% or 99%) identical to, or which hybridizes under high stringency conditions to, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:36, or SEQ ID NO:38; or combinations of such heavy and light variable regions.

The present invention further provides transgenic non-human mammals, hybridomas, and transgenic plants that express and/or produce the antibodies and

35 antigen binding portions described herein.

Also provided by the invention are kits comprising one or more isolated monoclonal antibodies or antigen binding portions thereof described herein and,

optionally, instructions for use in treating or diagnosing a disease associated with ErbB3 dependent signaling, such as cancers.

Antibodies and antigen binding portions thereof of the present invention can be used in a broad variety of therapeutic and diagnostic applications, particularly

5 oncological applications. Accordingly, in another aspect, the invention provides method for inhibiting EGF-like ligand mediated phosphorylation of ErbB3 in a subject by administering one or more antibodies or antigen binding portions thereof described herein in an amount sufficient to inhibit EGF-like mediated phosphorylation of ErbB3. The invention further provides methods for treating a variety of cancers in a subject,

10 including, but not limited to, melanoma, breast cancer, ovarian cancer, renal carcinoma, gastrointestinal/colon cancer, lung cancer, clear cell sarcoma, and prostate cancer, by administering one or more antibodies or antigen binding portions thereof described herein in an amount sufficient to treat the cancer. The antibodies or antigen binding portions thereof can be administered alone or in combination with other therapeutic

15 agents, such as anti-cancer agents, *e.g.*, other antibodies, chemotherapeutic agents and/or radiation.

In yet other embodiments, the invention provides methods for diagnosing and prognosing diseases (*e.g.*, cancers) associated with ErbB3. In one embodiment, this is achieved by contacting antibodies or antigen binding portions of the invention (*e.g.*, *ex vivo* or *in vivo*) with cells from the subject, and measuring the level of binding to ErbB3 on the cells, wherein abnormally high levels of binding to ErbB3 indicate that the subject has a cancer associated with ErbB3.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

25

Brief Description of the Drawings

Figures 1A and 1B are bar graphs depicting the binding of various anti-ErbB3 antibody candidates (fabs, referred to as Abs herein) to ErbB3 expressed on MALME-3M melanoma cells using a goat anti-human Alexa 647 secondary antibody.

30

Figures 2A-2D are graphs depicting the K_D values of various anti-ErbB3 antibody candidates. Figures 2A and 2B are graphs depicting the K_D value of Antibody # 6 (referred to as Ab #6) and Antibody # 3 (referred to as Ab #3), respectively, as measured using surface plasmon resonance (SPR) technology. Figures 2C and 2D are graphs depicting the K_D values of Ab #6 and Ab #3, respectively, as measured using a cell binding assay using MALME-3M melanoma cells.

35 Figure 3 is a graph depicting the binding specificity of an anti-ErbB3 antibody (Ab #6) to ErbB3 using ELISA. EGFR, BSA and TGF- α were used as controls.

Figure 4 is a graph depicting the ability of an anti-ErbB3 antibody (Ab #6) to decrease total ErbB3 levels in MALME-3M melanoma cells *in vitro*, as measured using ELISA.

Figures 5A and 5B are graphs depicting the ability of an anti-ErbB3 antibody (Ab #6) to downregulate ErbB3 receptors on MALME-3M cells, measured using FACS analysis. Figure 5A shows the results using an IgG1 isotype of the antibody. Figure 5B shows the results using an IgG2 isotype of the antibody.

Figures 6A-6D are graphs depicting the timecourse of antibody-mediated ErbB3 downregulation (Ab #6), as measured using FACS analysis.

Figure 7 is a bar graph depicting the ability of various anti-ErbB3 antibodies to downregulate ErbB3 in melanoma cells *in vivo*.

Figure 8 is a bar graph depicting the ability of an anti-ErbB3 antibody (Ab #6) to downregulate ErbB3 in ADRr xenografts *in vivo*.

Figure 9 is a graph depicting the ability of an anti-ErbB3 antibody (Ab #6) to inhibit proliferation of MALME-3M cells in a Cell Titer Glow Assay.

Figure 10 is a graph depicting the ability of an anti-ErbB3 antibody (Ab #6) to inhibit cell proliferation in an ovarian cell line, ADRr.

Figure 11 is a graph depicting the ability of an anti-ErbB3 antibody (Ab #6) to inhibit proliferation of ACHN cells.

Figure 12 is a bar graph depicting the ability of an anti-ErbB3 antibody (Ab #6) to inhibit ErbB3 phosphorylation in ADRr xenografts *in vivo*.

Figures 13A-13C are graphs depicting the ability of an anti-ErbB3 antibody (Ab #6) to inhibit betacellulin and heregulin-mediated phosphorylation of ErbB3 in ADRr cells.

Figures 14A-14B are graphs depicting the ability of an anti-ErbB3 antibody (Ab #6 IgG2 isotype) to inhibit ErbB3 phosphorylation in ovarian tumor cell lines OVCAR 5 and OVCAR 8.

Figures 15A-15C are graphs depicting the ability of betacellulin (BTC) to bind ErbB1 as shown by a lack of binding to ErbB1 negative MALME-3M cells (Figure 17A); binding to ErbB1 positive ADRr cells at concentrations of 10 nM (Figure 17B) and 200 nM (Figure 17B), respectively, and the inhibition of such binding by Erbitux.

Figures 16A-16B are graphs depicting the ability of an anti-ErbB3 antibody (Ab #6 IgG2 isotype) to inhibit heregulin-mediated signaling in MALME-3M cells. Figure 16A depicts the ability of the Ab #6 to inhibit heregulin-mediated phosphorylation of ErbB3 in MALME-3M cells and 16B depicts the ability of Ab #6 to inhibit phosphorylation of AKT in MALME-3M cells.

Figures 17A-D are graphs depicting the ability of an anti-ErbB3 antibody (Ab #6) to inhibit (A) ovarian (ADRr cells), (B) prostate Du145 cells), (C) ovarian (OvCAR8 cells), and (D) pancreatic (Colo357 cells) tumor growth via xenograft studies.

Figures 18A and 18B are graphs depicting the ability of Ab #6 (Figure 18A) and 5 Fab for Ab #3 (Figure 18B) to inhibit heregulin binding to ErbB3 on MALME-3M cells, as measured using FACS analysis.

Figures 19A and 19B are graphs depicting the ability of Ab #6 to inhibit the 10 binding of epiregulin to ErbB3 on ADRr cells. Figure 19A depicts the binding of epiregulin to ADRr cells, and Figure 19B depicts the ability of both Erbitux and Ab #6 to inhibit epiregulin binding to ADRr cells.

Figures 20A and 20B are graphs depicting the ability of heparin binding 15 epidermal growth factor (HB-EGF) to bind ErbB on ADRr cells (Figure 20A) and the inability of an anti-ErbB3 antibody (Ab #6) to inhibit such binding (Figure 20B).

Figures 21A-21C show the amino acid sequences of the variable heavy and light 20 chain regions of antibodies: Ab #6, Ab #3, Ab #14, Ab #17, and Ab #19.

Figures 22A-22B show the nucleotide sequences of the variable heavy and light chain regions of antibodies: Ab #6, Ab #3, and Ab #14.

Figure 23 shows the amino acid sequences of the variable light chain regions of 25 antibodies: Ab #6, Ab #17, and Ab #19, which have been reverted to the corresponding germline amino acid sequence. Amino acid residue changes are underlined.

Figures 24A-24C are graphs showing the ability of Ab #6 to inhibit VEGF secretion of tumor cells.

Figure 25 is a graph showing the effect of Ab #6 on cell migration.

Figures 26A-C are graphs showing (A) inhibition of spheroid growth in AdrR 25 cells, (B) inhibition of HRG induced spheroid growth in AdrR, and (C) inhibition of HRG induced spheroid growth in Du145 cells.

Figures 27 A and B are graphs showing the effect of Ab #6 on (A) HRG and (B) 30 BTC binding to AdrR cells.

Figure 28 is a graph showing the effect of Ab #6 on HGF induced ErbB3 phosphorylation.

Figures 29 A and B show the effect of Ab #6 on phosphorylation of (A) pErbB1 and pErbB3 and (B) HRG induced ErbB2/3 complex formation.

Figure 30 is a graph showing Ab #6 binds amino acid residues 20-202 of ErbB3.

35 **Detailed Description of the Invention**

In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

I. Definitions

The terms "ErbB3," "HER3," "ErbB3 receptor," and "HER3 receptor," as used interchangeably herein, refer human ErbB3 protein, as described in U.S. Pat. No. 5,480,968 and Plowman *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:4905-4909 (1990); see, 5 also, Kani *et al.*, *Biochemistry* 44:15842-857 (2005), Cho and Leahy, *Science* 297:1330-1333 (2002)).

The term "EGF-like ligand," as used herein, refers to ligands of epidermal growth factor receptor (EGFR), including epidermal growth factor (EGF) and closely related proteins, such as transforming growth factor- α (TGF- α), betacellulin (BTC), 10 heparin-binding epidermal growth factor (HB-EGF), biregulin (BIR) and amphiregulin (AR), which bind to EGFR on the surface of cells and stimulate the receptor's intrinsic protein-tyrosine kinase activity. Specifically, EGF-like ligands induce formation of EGFR (also referred to as ErbB1) and ErbB3 protein complex (see *e.g.*, Kim *et al.*, (1998) *Biochem J.*, 334:189-195), which results in phosphorylation of tyrosine residues 15 in the complex.

The antibodies and antigen binding portions thereof of the present invention inhibit EGF-like ligand mediated phosphorylation of ErbB3 and, in certain embodiments, exhibit one or more of the following additional properties: (i) inhibition of one or more of heregulin, epiregulin, epigen and biregulin (BIR)-mediated signaling 20 through ErbB3; (ii) inhibition of proliferation of cells expressing ErbB3; (iii) the ability to decrease levels of ErbB3 on cell surfaces; (iv) inhibition of VEGF secretion of cells expressing ErbB3; (v) inhibition of the migration of cells expressing ErbB3; (vi) inhibition of spheroid growth of cells expressing ErbB3; and/or (vii) binding to an epitope located on domain I of ErbB3, *e.g.*, an epitope which involves or spans residues 25 20-202 of the amino acid sequence of ErbB3.

The term "inhibition" as used herein, refers to any statistically significant decrease in biological activity, including full blocking of the activity. For example, "inhibition" can refer to a decrease of about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% in biological activity.

Accordingly, the phrase "inhibition of EGF-like ligand mediated phosphorylation of ErbB3," as used herein, refers to the ability of an antibody or antigen binding portion to statistically significantly decrease the phosphorylation of ErbB3 induced by an EGF-like ligand, relative to the phosphorylation in an untreated (control) cell. The cell which expresses ErbB3 can be a naturally occurring cell or cell line or can be recombinantly 30 produced by introducing nucleic acid encoding ErbB3 into a host cell. In one embodiment, the antibody or antigen binding portion thereof inhibits EGF-like ligand mediated phosphorylation of ErbB3 by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 35 30%.

90%, or 100%, as determined, for example, by Western blotting followed by probing with an anti-phosphotyrosine antibody as described in Kim *et al.*, (1998) *Biochem J.*, 334:189-195 and the Examples *infra*.

The phrase “inhibition of heregulin, epiregulin, epigen or biregulin-mediated signaling through ErbB3,” as used herein, refers to the ability of an antibody or an antigen-binding portion thereof to statistically significantly decrease signaling mediated by an ErbB3 ligand (e.g., heregulin, epiregulin, epigen and biregulin) through ErbB3, relative to the signaling in the absence of the antibody (control). ErbB3-ligands are also referred to herein as “heregulin-like ligands.” This means that, in the presence of the antibody or antigen binding portion thereof, a signal mediated in a cell expressing ErbB3 by one or more of heregulin, epiregulin, epigen and biregulin, relative to a control (no antibody), is statistically significantly decreased. An ErbB3-ligand mediated signal can be measured by assaying for the level or activity of an ErbB3 substrate, and/or a protein which is present in a cellular cascade involving ErbB3. In one embodiment, the antibody or antigen binding portion thereof decreases the level or activity of an ErbB3 substrate and/or that of a protein in a cellular cascade involving ErbB3, by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% relative to the level or activity in the absence of such antibody or antigen binding portion thereof (control). Such ErbB3-ligand mediated signaling can be measured using art recognized techniques which measure the level or activity of a substrate of ErbB3 (e.g., SHC or PI3K) or a protein in a cellular cascade involving ErbB3 (e.g., AKT) using kinase assays for such proteins (see, e.g., Horst *et al. supra*, Sudo *et al.* (2000) *Methods Enzymol.*, 322:388-92; and Morgan *et al.* (1990) *Eur. J. Biochem.*, 191:761-767).

In a particular embodiment, the antibody or antigen binding portion thereof inhibits ErbB3-ligand (e.g., heregulin, epiregulin, epigen or biregulin) mediated signaling through ErbB3 by inhibiting the binding of the ErbB3-ligand (e.g., one or more of heregulin, epiregulin, epigen or biregulin) to ErbB3. Some ligands (e.g., biregulin or BIR) function both as EGF-like ligands (*i.e.*, bind to EGFR/ErbB1) as well as ErbB3-like ligands (*i.e.*, bind to ErbB3).

The phrase “inhibition of heregulin, epiregulin, epigen or biregulin binding to ErbB3,” as used herein, refers to the ability of an antibody or an antigen-binding portion thereof to statistically significantly decrease the binding of an ErbB3 ligand (e.g., one or more of heregulin, epiregulin, epigen or biregulin) to ErbB3, relative to the binding in the absence of the antibody (control). This means that, in the presence of the antibody or antigen binding portion thereof, the amount of the ErbB3-ligand (e.g., heregulin, epiregulin, epigen or biregulin) which binds to ErbB3 relative to a control (no antibody), is statistically significantly decreased. The amount of an ErbB3 ligand which binds

ErbB3 may be decreased in the presence of an antibody or antigen binding portion thereof of the invention by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% relative to the amount in the absence of the antibody or antigen binding portion thereof (control). A decrease in ErbB3-ligand binding can be measured using art recognized techniques which measure the level of binding of labeled ErbB3-ligand (e.g., radiolabelled heregulin, epiregulin, epigen or biregulin) to cells expressing ErbB3 in the presence or absence (control) of the antibody or antigen binding portion thereof.

The phrase "inhibition of proliferation of a cell expressing ErbB3," as used herein, refers to the ability of an antibody or an antigen-binding portion thereof to statistically significantly decrease proliferation of a cell expressing ErbB3 relative to the proliferation in the absence of the antibody. In one embodiment, the proliferation of a cell expressing ErbB3 (e.g., a cancer cell) may be decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or 15 at least 80%, or at least 90%, or 100% when the cells are contacted with the antibody or antigen binding portion thereof of the present invention, relative to the proliferation measured in the absence of the antibody or antigen binding portion thereof (control). Cellular proliferation can be assayed using art recognized techniques which measure rate of cell division, the fraction of cells within a cell population undergoing cell division, 20 and/or rate of cell loss from a cell population due to terminal differentiation or cell death (e.g., using a cell titer glow assay or thymidine incorporation).

The phrase "the ability to decrease levels of ErbB3 on cell surfaces," as used herein, refers to the ability of an antibody or antigen binding portion thereof to statistically significantly reduce the amount of ErbB3 found on the surface of a cell 25 which has been exposed to the antibody relative to an untreated (control) cell. For example, a decrease in levels of ErbB3 on cell surfaces may result from increased internalization of ErbB3 (or increased ErbB3 endocytosis). In one embodiment, the antibody or antigen binding portion thereof decreases cell surface expression of ErbB3 by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% and/or increases 30 internalization of the ErbB3 receptor by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% relative to the cell surface expression or internalization in the absence of the antibody or antigen binding portion thereof (control). The levels of ErbB3 on 35 surfaces of cells and/or internalization of the ErbB3 receptor in the absence and the presence of an antibody or antigen-binding portion thereof can be readily measured using art recognized techniques, such as those described in Horst *et al.*, *supra* and in the examples herein.

The phrase “inhibition of VEGF secretion of cells expressing ErbB3,” as used herein, refers to the ability of an antibody or an antigen-binding portion thereof to statistically significantly decrease VEGF secretion of a cell expressing ErbB3 relative to the VEGF secretion in the absence of the antibody. In one embodiment, the VEGF secretion of a cell expressing ErbB3 (e.g., a cancer cell) may be decreased by at least 5 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% when the cells are contacted with the antibody or antigen binding portion thereof of the present invention, relative to the VEGF secretion measured in the absence of the antibody or antigen binding portion thereof (control). VEGF secretion can be assayed using art recognized techniques, such as those described herein.

The phrase “inhibition of the migration of cells expressing ErbB3,” as used herein, refers to the ability of an antibody or an antigen-binding portion thereof to statistically significantly decrease the migration of a cell expressing ErbB3 relative to 15 the migration of the cell in the absence of the antibody. In one embodiment, the migration of a cell expressing ErbB3 (e.g., a cancer cell) may be decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% when the cells are contacted with the antibody or antigen binding portion thereof of the present invention, relative to cell 20 migration measured in the absence of the antibody or antigen binding portion thereof (control). Cell migration can be assayed using art recognized techniques, such as those described herein.

The phrase “inhibition of spheroid growth of cells expressing ErbB3,” as used herein, refers to the ability of an antibody or an antigen-binding portion thereof to 25 statistically significantly decrease the migration of a cell expressing ErbB3 relative to the migration of the cell in the absence of the antibody. In one embodiment, the migration of a cell expressing ErbB3 (e.g., a cancer cell) may be decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or 100% when the cells are contacted with 30 the antibody or antigen binding portion thereof of the present invention, relative to cell migration measured in the absence of the antibody or antigen binding portion thereof (control). Cell migration can be assayed using art recognized techniques, such as those described herein. The term “antibody” or “immunoglobulin,” as used interchangeably herein, includes whole antibodies and any antigen binding fragment (i.e., “antigen-binding portion”) or single chains thereof. An “antibody” comprises at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as V_H) and a heavy chain constant region. The heavy chain constant region is comprised of three 35

domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as V_L) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The V_H and V_L regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may 5 mediate the binding of the immunoglobulin to host tissues or factors, including various 10 cells of the immune system (e.g., effector cells) and the first component (Clq) of the classical complement system. Exemplary antibodies of the invention include antibodies#1, 3 and 14, and antigen-binding portions thereof.

The term “antigen-binding portion” of an antibody (or simply “antibody 15 portion”), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., ErbB3). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term “antigen-binding portion” of an antibody include (i) a Fab fragment, a monovalent fragment 20 consisting of the V_L , V_H , CL and CH1 domains; (ii) a $F(ab')_2$ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the V_H and CH1 domains; (iv) a Fv fragment consisting of the V_L and V_H domains of a single arm of an antibody, (v) a dAb including VH and VL domains; (vi) a dAb fragment (Ward *et al.* (1989) *Nature* 341, 544-546), which 25 consists of a V_H domain; (vii) a dAb which consists of a VH or a VL domain; and (viii) an isolated complementarity determining region (CDR) or (ix) a combination of two or more isolated CDRs which may optionally be joined by a synthetic linker. Furthermore, although the two domains of the Fv fragment, V_L and V_H , are coded for by separate 30 genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the V_L and V_H regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird *et al.* (1988) *Science* 242, 423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85, 5879-5883). Such single chain antibodies are also intended to be encompassed within the term “antigen-binding portion” of an antibody. These antibody fragments are obtained 35 using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies. Antigen-binding portions can be produced by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact immunoglobulins.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are

5 highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. Monoclonal antibodies can be prepared using any art recognized technique and those described herein such as, for

10 example, a hybridoma method, as described by Kohler *et al.* (1975) *Nature*, 256:495, a transgenic animal, as described by, for example, (see *e.g.*, Lonberg, *et al.* (1994) *Nature* 368(6474): 856-859), recombinant DNA methods (see, *e.g.*, U.S. Pat. No. 4,816,567), or using phage antibody libraries using the techniques described in, for example, Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991).

15 Monoclonal antibodies include chimeric antibodies, human antibodies and humanized antibodies and may occur naturally or be recombinantly produced.

The term "recombinant antibody," refers to antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (*e.g.*, a mouse) that is transgenic or transchromosomal for

20 immunoglobulin genes (*e.g.*, human immunoglobulin genes) or a hybridoma prepared therefrom, (b) antibodies isolated from a host cell transformed to express the antibody, *e.g.*, from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial antibody library (*e.g.*, containing human antibody sequences) using phage display, and

25 (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of immunoglobulin gene sequences (*e.g.*, human immunoglobulin genes) to other DNA sequences. Such recombinant antibodies may have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies can be subjected to *in vitro* mutagenesis and thus the amino acid sequences of the V_H and V_L regions of the

30 recombinant antibodies are sequences that, while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

The term "chimeric immunoglobulin" or antibody refers to an immunoglobulin or antibody whose variable regions derive from a first species and whose constant

35 regions derive from a second species. Chimeric immunoglobulins or antibodies can be constructed, for example by genetic engineering, from immunoglobulin gene segments belonging to different species.

The term "human antibody," as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from human germline immunoglobulin sequences as described, for example, by Kabat et al. (See Kabat, *et al.* (1991) *Sequences of proteins of Immunological Interest, Fifth Edition*, 5 U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Furthermore, if the antibody contains a constant region, the constant region also is derived from human germline immunoglobulin sequences. The human antibodies may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic 10 mutation *in vivo*). However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The human antibody can have at least one or more amino acids replaced with an 15 amino acid residue, e.g., an activity enhancing amino acid residue which is not encoded by the human germline immunoglobulin sequence. Typically, the human antibody can have up to twenty positions replaced with amino acid residues which are not part of the human germline immunoglobulin sequence. In a particular embodiment, these replacements are within the CDR regions as described in detail below.

20 The term "humanized immunoglobulin" or "humanized antibody" refers to an immunoglobulin or antibody that includes at least one humanized immunoglobulin or antibody chain (i.e., at least one humanized light or heavy chain). The term "humanized immunoglobulin chain" or "humanized antibody chain" (i.e., a "humanized immunoglobulin light chain" or "humanized immunoglobulin heavy chain") refers to an 25 immunoglobulin or antibody chain (i.e., a light or heavy chain, respectively) having a variable region that includes a variable framework region substantially from a human immunoglobulin or antibody and complementarity determining regions (CDRs) (e.g., at least one CDR, preferably two CDRs, more preferably three CDRs) substantially from a non-human immunoglobulin or antibody, and further includes constant regions (e.g., at 30 least one constant region or portion thereof, in the case of a light chain, and preferably three constant regions in the case of a heavy chain). The term "humanized variable region" (e.g., "humanized light chain variable region" or "humanized heavy chain variable region") refers to a variable region that includes a variable framework region substantially from a human immunoglobulin or antibody and complementarity 35 determining regions (CDRs) substantially from a non-human immunoglobulin or antibody.

A "bispecific" or "bifunctional antibody" is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific

antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, (1990) *Clin. Exp. Immunol.* 79, 315-321; Kostelny *et al.* (1992) *J. Immunol.* 148, 1547-1553. In a particular embodiment, a bispecific antibody according to the present invention includes 5 binding sites for both ErbB3 and IGF1-R (*i.e.*, insulin-like growth factor 1-receptor). In another embodiment, a bispecific antibody according to the present invention includes binding sites for both ErbB3 and C-MET. In other embodiments, a bispecific antibody includes a binding site for ErbB3 and a binding site for ErbB2, ERbB3, ErbB4, EGFR, Lewis Y, MUC-1, EpCAM, CA125, prostate specific membrane antigen, PDGFR- α , 10 PDGFR- β , C-KIT, or any of the FGF receptors.

As used herein, a “heterologous antibody” is defined in relation to the transgenic non-human organism or plant producing such an antibody.

An “isolated antibody,” as used herein, is intended to refer to an antibody which is substantially free of other antibodies having different antigenic specificities (*e.g.*, an 15 isolated antibody that specifically binds to ErbB3 is substantially free of antibodies that specifically bind antigens other than ErbB3). In addition, an isolated antibody is typically substantially free of other cellular material and/or chemicals. In one embodiment of the invention, a combination of “isolated” monoclonal antibodies having different ErbB3 binding specificities are combined in a well defined composition.

20 As used herein, “isotype” refers to the antibody class (*e.g.*, IgM or IgG1) that is encoded by heavy chain constant region genes. In one embodiment, an antibody or antigen binding portion thereof is of an isotype selected from an IgG1, an IgG2, an IgG3, an IgG4, an IgM, an IgA1, an IgA2, an IgAsec, an IgD, or an IgE antibody isotype. In some embodiments, a monoclonal antibody of the invention is of the IgG1 25 isotype. In other embodiments, a monoclonal antibody of the invention is of the IgG2 isotype.

As used herein, “isotype switching” refers to the phenomenon by which the class, or isotype, of an antibody changes from one Ig class to one of the other Ig classes.

30 As used herein, “nonswitched isotype” refers to the isotypic class of heavy chain that is produced when no isotype switching has taken place; the CH gene encoding the nonswitched isotype is typically the first CH gene immediately downstream from the functionally rearranged VDJ gene. Isotype switching has been classified as classical or non-classical isotype switching. Classical isotype switching occurs by recombination events which involve at least one switch sequence regions in a gene encoding an 35 antibody. Non-classical isotype switching may occur by, for example, homologous recombination between human σ_{μ} and human Σ_{μ} (δ -associated deletion). Alternative non-classical switching mechanisms, such as intertransgene and/or interchromosomal recombination, among others, may occur and effectuate isotype switching.

As used herein, the term “switch sequence” refers to those DNA sequences responsible for switch recombination. A “switch donor” sequence, typically a μ switch region, will be 5' (*i.e.*, upstream) of the construct region to be deleted during the switch recombination. The “switch acceptor” region will be between the construct region to be deleted and the replacement constant region (*e.g.*, γ , ϵ , etc.). As there is no specific site where recombination always occurs, the final gene sequence will typically not be predictable from the construct.

An “antigen” is an entity (*e.g.*, a proteinaceous entity or peptide) to which an antibody or antigen-binding portion thereof binds. In various embodiments of the present invention, an antigen is ErbB3 or a ErbB3-like molecule. In a particular embodiment according to the invention, an antigen is human ErbB3.

The term “epitope” or “antigenic determinant” refers to a site on an antigen to which an immunoglobulin or antibody specifically binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include techniques in the art and those described herein, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, *e.g.*, *Epitope Mapping Protocols in Methods in Molecular Biology*, Vol. 66, G. E. Morris, Ed. (1996).

Also encompassed by the present invention are antibodies that bind the same or an overlapping epitope as the antibodies of the present invention, *i.e.*, antibodies that compete for binding to ErbB3, or bind epitopes which overlap with epitopes bound by the antibodies described herein, *i.e.*, an epitope located on domain I of ErbB3.

Antibodies that recognize the same epitope can be identified using routine techniques such as an immunoassay, for example, by showing the ability of one antibody to block the binding of another antibody to a target antigen, *i.e.*, a competitive binding assay. Competitive binding is determined in an assay in which the immunoglobulin under test inhibits specific binding of a reference antibody to a common antigen, such as ErbB3. Numerous types of competitive binding assays are known, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahli *et al.*, (1983) *Methods in Enzymology* 9:242); solid phase direct biotin-avidin EIA (see Kirkland *et al.*, (1986) *J. Immunol.* 137:3614); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow and Lane, (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press); solid phase direct label RIA using I-125 label (see Morel *et al.*,

(1988) *Mol. Immunol.* 25(1):7); solid phase direct biotin-avidin EIA (Cheung *et al.*, (1990) *Virology* 176:546); and direct labeled RIA. (Moldenhauer *et al.*, (1990) *Scand. J. Immunol.* 32:77). Typically, such an assay involves the use of purified antigen (e.g., ErbB3) bound to a solid surface or cells bearing either of these, an unlabeled test immunoglobulin and a labeled reference immunoglobulin. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess. Usually, when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 50-55%, 55-60%, 60-65%, 65-70% 70-75% or more.

10 As used herein, the terms “specific binding,” “specifically binds,” “selective binding,” and “selectively binds,” mean that an antibody or antigen-binding portion thereof, exhibits appreciable affinity for a particular antigen or epitope and, generally, does not exhibit significant cross-reactivity with other antigens and epitopes.

15 “Appreciable” or preferred binding includes binding with an affinity of at least 10^6 , 10^7 , 10^8 , 10^9 M⁻¹, or 10^{10} M⁻¹. Affinities greater than 10^7 M⁻¹, preferably greater than 10^8 M⁻¹ are more preferred. Values intermediate of those set forth herein are also intended to be within the scope of the present invention and a preferred binding affinity can be indicated as a range of affinities, for example, 10^6 to 10^{10} M⁻¹, preferably 10^7 to 10^{10} M⁻¹,

20 more preferably 10^8 to 10^{10} M⁻¹. An antibody that “does not exhibit significant cross-reactivity” is one that will not appreciably bind to an undesirable entity (e.g., an undesirable proteinaceous entity). For example, in one embodiment, an antibody or antigen-binding portion thereof that specifically binds to ErbB3 will appreciably bind that ErbB3 molecule but will not significantly react with other ErbB molecules and non-25 ErbB proteins or peptides. Specific or selective binding can be determined according to any art-recognized means for determining such binding, including, for example, according to Scatchard analysis and/or competitive binding assays.

30 The term “K_D,” as used herein, is intended to refer to the dissociation equilibrium constant of a particular antibody-antigen interaction or the affinity of an antibody for an antigen. In one embodiment, the antibody or antigen binding portion thereof according to the present invention binds an antigen (e.g., ErbB3) with an affinity (K_D) of 50 nM or better (i.e., or less) (e.g., 40 nM or 30 nM or 20 nM or 10 nM or less), as measured using a surface plasmon resonance assay or a cell binding assay. In a particular embodiment, an antibody or antigen binding portion thereof according to the present invention binds 35 ErbB3 with an affinity (K_D) of 8 nM or better (e.g., 7 nM, 6 nM, 5 nM, 4 nM, 2 nM, 1.5 nM, 1.4 nM, 1.3 nM, 1nM or less), as measured by a surface plasmon resonance assay or a cell binding assay. In other embodiments, an antibody or antigen binding portion thereof binds an antigen (e.g., ErbB3) with an affinity (K_D) of approximately less than

10⁻⁷ M, such as approximately less than 10⁻⁸ M, 10⁻⁹ M or 10⁻¹⁰ M or even lower when determined by surface plasmon resonance (SPR) technology in a BIACORE 3000 instrument using recombinant ErbB3 as the analyte and the antibody as the ligand, and binds to the predetermined antigen with an affinity that is at least two-fold greater than 5 its affinity for binding to a non-specific antigen (e.g., BSA, casein) other than the predetermined antigen or a closely-related antigen.

The term “K_{off},” as used herein, is intended to refer to the off rate constant for the dissociation of an antibody from the antibody/antigen complex.

The term “EC50,” as used herein, refers to the concentration of an antibody or an 10 antigen-binding portion thereof, which induces a response, either in an *in vitro* or an *in vivo* assay, which is 50% of the maximal response, *i.e.*, halfway between the maximal response and the baseline.

As used herein, “glycosylation pattern” is defined as the pattern of carbohydrate 15 units that are covalently attached to a protein, more specifically to an immunoglobulin protein.

The term “naturally-occurring” as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory 20 is naturally-occurring.

The term “rearranged” as used herein refers to a configuration of a heavy chain or light chain immunoglobulin locus wherein a V segment is positioned immediately adjacent to a D-J or J segment in a conformation encoding essentially a complete V_H or V_L domain, respectively. A rearranged immunoglobulin gene locus can be identified by 25 comparison to germline DNA; a rearranged locus will have at least one recombined heptamer/nonamer homology element.

The term “unrearranged” or “germline configuration” as used herein in reference to a V segment refers to the configuration wherein the V segment is not recombined so as to be immediately adjacent to a D or J segment.

30 The term “nucleic acid molecule,” as used herein, is intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

The term “isolated nucleic acid molecule,” as used herein in reference to nucleic acids encoding antibodies or antibody portions (e.g., V_H, V_L, CDR3) that bind to ErbB3, 35 is intended to refer to a nucleic acid molecule in which the nucleotide sequences encoding the antibody or antibody portion are free of other nucleotide sequences encoding antibodies that bind antigens other than ErbB3, which other sequences may naturally flank the nucleic acid in human genomic DNA.

The term “modifying,” or “modification,” as used herein, is intended to refer to changing one or more amino acids in the antibodies or antigen-binding portions thereof. The change can be produced by adding, substituting or deleting an amino acid at one or more positions. The change can be produced using known techniques, such as PCR

5 mutagenesis. For example, in some embodiments, an antibody or an antigen-binding portion thereof identified using the methods of the invention can be modified, to thereby modify the binding affinity of the antibody or antigen-binding portion thereof to ErbB3.

The present invention also encompasses “conservative amino acid substitutions” in the sequences of the antibodies of the invention, *i.e.*, nucleotide and amino acid 10 sequence modifications which do not abrogate the binding of the antibody encoded by the nucleotide sequence or containing the amino acid sequence, to the antigen, *i.e.*, ErbB3. Conservative amino acid substitutions include the substitution of an amino acid in one class by an amino acid of the same class, where a class is defined by common physicochemical amino acid side chain properties and high substitution frequencies in 15 homologous proteins found in nature, as determined, for example, by a standard Dayhoff frequency exchange matrix or BLOSUM matrix. Six general classes of amino acid side chains have been categorized and include: Class I (Cys); Class II (Ser, Thr, Pro, Ala, Gly); Class III (Asn, Asp, Gln, Glu); Class IV (His, Arg, Lys); Class V (Ile, Leu, Val, Met); and Class VI (Phe, Tyr, Trp). For example, substitution of an Asp for another class 20 III residue such as Asn, Gln, or Glu, is a conservative substitution. Thus, a predicted nonessential amino acid residue in an anti-ErbB3 antibody is preferably replaced with another amino acid residue from the same class. Methods of identifying nucleotide and amino acid conservative substitutions which do not eliminate antigen binding are well-known in the art (see, *e.g.*, Brummell *et al.*, *Biochem.* 32:1180-1187 (1993); Kobayashi 25 *et al.* *Protein Eng.* 12(10):879-884 (1999); and Burks *et al.* *Proc. Natl. Acad. Sci. USA* 94:412-417 (1997)).

The term “non-conservative amino acid substitution” refers to the substitution of an amino acid in one class with an amino acid from another class; for example, substitution of an Ala, a class II residue, with a class III residue such as Asp, Asn, Glu, 30 or Gln.

Alternatively, in another embodiment, mutations (conservative or non-conservative) can be introduced randomly along all or part of an anti-ErbB3 antibody coding sequence, such as by saturation mutagenesis, and the resulting modified anti-ErbB3 antibodies can be screened for binding activity.

35 A “consensus sequence” is a sequence formed from the most frequently occurring amino acids (or nucleotides) in a family of related sequences (See *e.g.*, Winnaker, *From Genes to Clones* (Verlagsgesellschaft, Weinheim, Germany 1987). In a family of proteins, each position in the consensus sequence is occupied by the amino

acid occurring most frequently at that position in the family. If two amino acids occur equally frequently, either can be included in the consensus sequence. A “consensus framework” of an immunoglobulin refers to a framework region in the consensus immunoglobulin sequence.

5 Similarly, the consensus sequence for the CDRs of can be derived by optimal alignment of the CDR amino acid sequences of ErbB3 antibodies of the present invention.

For nucleic acids, the term “substantial homology” indicates that two nucleic acids, or designated sequences thereof, when optimally aligned and compared, are 10 identical, with appropriate nucleotide insertions or deletions, in at least about 80% of the nucleotides, usually at least about 90% to 95%, and more preferably at least about 98% to 99.5% of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to the complement of the strand.

15 The percent identity between two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between 20 two sequences can be accomplished using a mathematical algorithm, as described in the non-limiting examples below.

The percent identity between two nucleotide sequences can be determined using the GAP program in the GCG software, using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. The percent 25 identity between two nucleotide or amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch 30 (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

35 The nucleic acid and protein sequences of the present invention can further be used as a “query sequence” to perform a search against public databases to, for example, identify related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100,

wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is “isolated” or “rendered substantially pure” when purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art. See, F. Ausubel, *et al.*, ed. Current Protocols in Molecular Biology, Greene Publishing and Wiley Interscience, New York (1987).

The nucleic acid compositions of the present invention, while often in a native sequence (except for modified restriction sites and the like), from either cDNA, genomic or mixtures thereof may be mutated, in accordance with standard techniques to provide gene sequences. For coding sequences, these mutations, may affect amino acid sequence as desired. In particular, DNA sequences substantially homologous to or derived from native V, D, J, constant, switches and other such sequences described herein are contemplated (where “derived” indicates that a sequence is identical or modified from another sequence).

The term “operably linked” refers to a nucleic acid sequence placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice. A nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the

sequence. With respect to transcription regulatory sequences, operably linked means that the DNA sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame. For switch sequences, operably linked indicates that the sequences are capable of effecting switch

5 recombination.

The term "vector," as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector,

10 wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated

15 along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. The terms, "plasmid" and "vector" may be used interchangeably.

20 However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The term "recombinant host cell" (or simply "host cell"), as used herein, is intended to refer to a cell into which a recombinant expression vector has been

25 introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein.

30 The terms "treat," "treating," and "treatment," as used herein, refer to therapeutic or preventative measures described herein. The methods of "treatment" employ administration to a subject, an antibody or antigen binding portion of the present invention, for example, a subject having a disease or disorder associated with ErbB3 dependent signaling or predisposed to having such a disease or disorder, in order to

35 prevent, cure, delay, reduce the severity of, or ameliorate one or more symptoms of the disease or disorder or recurring disease or disorder, or in order to prolong the survival of a subject beyond that expected in the absence of such treatment.

The term “disease associated with ErbB3 dependent signaling,” or “disorder associated with ErbB3 dependent signaling,” as used herein, includes disease states and/or symptoms associated with a disease state, where increased levels of ErbB3 and/or activation of cellular cascades involving ErbB3 are found. It is understood that ErbB3

5 heterodimerizes with other ErbB proteins such as, EGFR and ErbB2, when increased levels of ErbB3 are found. Accordingly, the term “disease associated with ErbB3 dependent signaling,” also includes disease states and/or symptoms associated with disease states where increased levels of EGFR/ErbB3 and/or ErbB2/ErbB3 heterodimers are found. In general, the term “disease associated with ErbB3 dependent signaling,”

10 refers to any disorder, the onset, progression or the persistence of the symptoms of which requires the participation of ErbB3. Exemplary ErbB3-mediated disorders include, but are not limited to, for example, cancer.

The terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth.

15 Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastric cancer, pancreatic cancer, glial cell tumors such as glioblastoma and neurofibromatosis, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer,

20 melanoma, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer, renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer. In a particular embodiment, a cancer treated or diagnosed using the methods of the present invention is selected from melanoma, breast cancer, ovarian cancer, renal carcinoma, gastrointestinal/colon cancer, lung

25 cancer, and prostate cancer.

The term “effective amount,” as used herein, refers to that amount of an antibody or an antigen binding portion thereof that binds ErbB3, which is sufficient to effect treatment, prognosis or diagnosis of a disease associated with ErbB3 dependent signaling, as described herein, when administered to a subject. A therapeutically effective amount will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The dosages for administration can range from, for example, about 1 ng to about 10,000 mg, about 5 ng to about 9,500 mg, about 10 ng to about 30,000 mg, about 20 ng to about 8,500 mg, about 30 ng to about 7,500 mg, about 40 ng to about 7,000 mg, about 50 ng to about 6,500 mg, about 100 ng to about 6,000 mg, about 200 ng to about 5,500 mg, about 300 ng to about 5,000 mg, about 400 ng to about 4,500 mg, about 500 ng to about 4,000 mg, about 1 µg to about 3,500 mg, about 5 µg to about

3,000 mg, about 10 µg to about 2,600 mg, about 20 µg to about 2,575 mg, about 30 µg to about 2,550 mg, about 40 µg to about 2,500 mg, about 50 µg to about 2,475 mg, about 100 µg to about 2,450 mg, about 200 µg to about 2,425 mg, about 300 µg to about 2,000, about 400 µg to about 1,175 mg, about 500 µg to about 1,150 mg, about 0.5 mg to about 1,125 mg, about 1 mg to about 1,100 mg, about 1.25 mg to about 1,075 mg, about 1.5 mg to about 1,050 mg, about 2.0 mg to about 1,025 mg, about 2.5 mg to about 1,000 mg, about 3.0 mg to about 975 mg, about 3.5 mg to about 950 mg, about 4.0 mg to about 925 mg, about 4.5 mg to about 900 mg, about 5 mg to about 875 mg, about 10 mg to about 850 mg, about 20 mg to about 825 mg, about 30 mg to about 800 mg, about 40 mg to about 775 mg, about 50 mg to about 750 mg, about 100 mg to about 725 mg, about 200 mg to about 700 mg, about 300 mg to about 675 mg, about 400 mg to about 650 mg, about 500 mg, or about 525 mg to about 625 mg, of an antibody or antigen binding portion thereof, according to the invention. Dosage regiments may be adjusted to provide the optimum therapeutic response. An effective amount is also one in which any toxic or detrimental effects (*i.e.*, side effects) of an antibody or antigen binding portion thereof are minimized and/or outweighed by the beneficial effects.

The term "patient" includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

As used herein, the term "subject" includes any human or non-human animal.

20 For example, the methods and compositions of the present invention can be used to treat a subject having cancer. In a particular embodiment, the subject is a human. The term "non-human animal" includes all vertebrates, *e.g.*, mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, reptiles, *etc.*

25 The term "sample" refers to tissue, body fluid, or a cell from a patient or a subject. Normally, the tissue or cell will be removed from the patient, but *in vivo* diagnosis is also contemplated. In the case of a solid tumor, a tissue sample can be taken from a surgically removed tumor and prepared for testing by conventional techniques. In the case of lymphomas and leukemias, lymphocytes, leukemic cells, or lymph tissues can be obtained and appropriately prepared. Other patient samples, including urine, tear drops, serum, cerebrospinal fluid, feces, sputum, cell extracts *etc.* can also be useful for particular tumors.

30 The terms "anti-cancer agent" and "antineoplastic agent" refer to drugs used to treat malignancies, such as cancerous growths. Drug therapy may be used alone, or in combination with other treatments such as surgery or radiation therapy. Several classes of drugs may be used in cancer treatment, depending on the nature of the organ involved. For example, breast cancers are commonly stimulated by estrogens, and may be treated with drugs which inactive the sex hormones. Similarly, prostate cancer may

be treated with drugs that inactivate androgens, the male sex hormone. Anti-cancer agents of the present invention include, among others, the following agents:

Anti-Cancer Agent	Comments	Examples
Antibodies	Antibodies which bind IGF-1R (insulin-like growth factor type 1 receptor), which is expressed on the cell surface of most human cancers	A12 (fully humanized mAb) 19D12 (fully humanized mAb) CP751-871 (fully humanized mAb) H7C10 (humanized mAb) alphaIR3 (mouse) scFV/FC (mouse/human chimera) EM/164 (mouse)
(a) antibodies other than anti-ErbB3 antibodies; and		
(b) anti-ErbB3 antibodies which bind different epitopes		
Small Molecules Targeting IGF1R	Antibodies which bind EGFR (epidermal growth factor receptor); Mutations affecting EGFR expression or activity could result in cancer	Matuzumab (EMD72000) Erbitux® / Cetuximab (Imclone) Vectibix® / Panitumumab (Amgen) mAb 806 Nimotuzumab (TheraCIM)
	Antibodies which bind cMET (Mesenchymal epithelial transition factor); a member of the MET family of receptor tyrosine kinases	AVEO (AV299) (AVEO) AMG102 (Amgen) 5D5 (OA-5D5) (Genentech)
	Anti-ErbB3 antibodies which bind different epitopes	Ab #14 (MM 121-14) described herein Herceptin® (Trastuzumab; Genentech) 1B4C3; 2D1D12 (U3 Pharma AG) NVP-AEW541-A BMS-536,924 (1H-benzoimidazol-2-yl)-1H-pyridin-2-one) BMS-554,417 Cycloligan TAE226 PQ401
Small Molecules Targeting EGFR	IGF-1R (insulin-like growth factor type 1 receptor), which is expressed on the cell surface of most human cancers	Iressa® / Gefitinib (AstraZeneca) CI-1033 (PD 183805) (Pfizer) Lapatinib (GW-572016) (GlaxoSmithKline) Tykerb® / Lapatinib Ditosylate (SmithKline Beecham) Tarceva®/ Erlotinib HCL (OSI-774) (OSI Pharma) PKI-166 (Novartis)

		PD-158780 EKB-569 Tyrphostin AG 1478(4-(3-Chloroanillino)-6,7-dimethoxyquinazoline)
Small Molecules Targeting cMET	cMET (Mesenchymal epithelial transition factor); a member of the MET family of receptor tyrosine kinases)	PHA665752 ARQ 197
Antimetabolites	An antimetabolite is a chemical with a similar structure to a substance (a metabolite) required for normal biochemical reactions, yet different enough to interfere with the normal functions of cells, including cell division.	Flourouracil (5-FU) Capecitabine / XELODA® (HLR Roche) 5-Trifluoromethyl-2'-deoxyuridine Methotrexate sodium (Trexall) (Barr) Raltitrexed / Tomudex® (AstraZaneca) Pemetrexed / Alimta® (Lilly) Tegafur Cytosine Arabinoside (Cytarabine, Ara-C) / Thioguanine® (GlaxoSmithKline) 5-azacytidine 6-mercaptopurine (Mercaptopurine, 6-MP) Azathioprine / Azasan® (AAIPHARMA LLC) 6-thioguanine (6-TG) / Purinethol® (TEVA) Pentostatin / Nipent® (Hospira Inc.) Fludarabine phosphate / Fludara® (Bayer Health Care) Cladribine (2-CdA, 2-chlorodeoxyadenosine) / Leustatin® (Ortho Biotech)
Alkylating agents	An alkylating antineoplastic agent is an alkylating agent that attaches an alkyl group to DNA. Since cancer cells generally proliferate unrestricatively more than do healthy cells they are more sensitive to DNA damage, and alkylating agents are used clinically to treat a variety of tumours.	Ribonucleotide Reductase Inhibitor (RNR) Cyclophosphamide / Cytoxan (BMS) Neosar (TEVA) Ifosfamide / Mitoxana® (ASTA Medica) Thiotepa (Bedford, Abraxis, Teva) BCNU → 1,3-bis(2-chloroethyl)-1-nitosourea CCNU → 1,-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea (methyl CCNU) Hexamethylmelamine (Altretamine, HMM) / Hexalen® (MGI Pharma Inc.) Busulfan / Myleran (GlaxoSmithKline) Procarbazine HCL / Matulane (Sigma Tau Pharmaceuticals, Inc.) Dacarbazine (DTIC) Chlorambucil / Leukaran® (SmithKline Beecham) Melphalan / Alkeran® (GlaxoSmithKline)

		Cisplatin (Cisplatinum, CDDP) / Platinol (Bristol Myers)
		Carboplatin / Paraplatin (BMS)
		Oxaliplatin / Eloxitan® (Sanofi-Aventis US)
Topoisomerase inhibitors	Topoisomerase inhibitors are chemotherapy agents designed to interfere with the action of topoisomerase enzymes (topoisomerase I and II), which are enzymes that control the changes in DNA structure by catalyzing the breaking and rejoining of the phosphodiester backbone of DNA strands during the normal cell cycle.	Doxorubicin HCL / Doxil® (Alza) Daunorubicin citrate / Daunoxome® (Gilead) Mitoxantrone HCL/Novantrone (EMD Serono) Actinomycin D Etoposide / Vepesid® (BMS)/ Etopophos® (Hospira, Bedford, Teva Parenteral, Etc.) Topotecan HCL / Hycamtin® (GlaxoSmithKline) Teniposide (VM-26) / Vumon® (BMS) Irinotecan HCL(CPT-11) / Camptosar® (Pharmacia & Upjohn)
Microtubule targeting agents	Microtubules are one of the components of the cytoskeleton. They have diameter of ~ 24 nm and length varying from several micrometers to possibly millimeters in axons of nerve cells. Microtubules serve as structural components within cells and are involved in many cellular processes including mitosis, cytokinesis, and vesicular transport.	Vincristine / Oncovin® (Lilly) Vinblastine sulfate/Velban®(discontinued) (Lilly) Vinorelbine tartrate / Navelbine® (PierreFabre) Vindesine sulphate / Eldisine® (Lilly) Paclitaxel / Taxol® (BMS) Docetaxel / Taxotere® (Sanofi Aventis US) Nanoparticle paclitaxel (ABI-007) / Abraxane® (Abraxis BioScience, Inc.) Ixabepilone / IXEMPRA™ (BMS)
Kinase inhibitors	Tyrosine kinases are enzymes within the cell that function to attach phosphate groups to the amino acid tyrosine. By blocking the ability of protein tyrosine kinases to function, these compounds provide a tool for controlling cancerous	Imatinib mesylate / Gleevec (Novartis) Sunitinib malate / Sutent® (Pfizer) Sorafenib tosylate / Nexavar® (Bayer) Nilotinib hydrochloride monohydrate / Tasigna® (Novartis)

cell growth.

Protein synthesis inhibitors	Induces cell apoptosis	L-asparaginase / Elspar® (Merck & Co.)
Immunotherapeutic agents	Induces cancer patients to exhibit immune responsiveness	Alpha interferon Angiogenesis Inhibitor / Avastin® (Genentech) IL-2→ Interleukin 2 (Aldesleukin) / Proleukin® (Chiron) IL-12→ Interleukin 12
Hormones	Hormone therapies associated with menopause and aging seek to increase the amount of certain hormones in your body to compensate for age- or disease-related hormonal declines. Hormone therapy as a cancer treatment either reduces the level of specific hormones or alters the cancer's ability to use these hormones to grow and spread.	Toremifene citrate / Fareston® (GTx, Inc.) Fulvestrant / Faslodex® (AstraZeneca) Raloxifene HCL / Evista® (Lilly) Anastrazole / Arimidex® (AstraZeneca) Letrozole / Femara® (Novartis) Fadrozole (CGS 16949A) Exemestane / Aromasin® (Pharmacia & Upjohn) Leuprolide acetate / Eligard® (QTL USA) Lupron® (TAP Pharm.) Goserelin acetate / Zoladex® (AstraZeneca) Triptorelin pamoate / Trelstar® (Watson Labs) Buserelin / Suprefact® (Sanofi Aventis) Nafarelin Cetrorelix / Cetrotide® (EMD Serono) Bicalutamide / Casodex® (AstraZeneca) Nilutamide / Nilandron® (Aventis Pharm.) Megestrol acetate / Megace® (BMS) Somatostatin Analogs (Octreotide acetate / Sandostatin® (Novartis))
Glucocorticoids	Anti-inflammatory drugs used to reduce swelling that causes cancer pain.	Prednisolone Dexamethasone / Decadron® (Wyeth)
Aromatase inhibitors	Includes imidazoles	Ketoconazole
mTOR inhibitors	The mTOR signaling pathway was originally discovered during studies of the immunosuppressive agent rapamycin. This highly conserved pathway regulates cell proliferation and	Sirolimus (Rapamycin) / Rapamune® (Wyeth) Temsirolimus (CCI-779) / Torisel® (Wyeth) Deforolimus (AP23573) (Ariad Pharm.) Everolimus (RAD001) / Certican® (Novartis)

metabolism in response to environmental factors, linking cell growth factor receptor signaling via phosphoinositide-3-kinase (PI-3K) to cell growth, proliferation, and angiogenesis.

Chemotherapeutic agents

Adriamycin, 5-Fluorouracil, Cytoxin, Bleomycin, Mitomycin C, Daunomycin, Carminomycin, Aminopterin, Dactinomycin, Mitomycins, Esperamicins

One or more anti-cancer agents may be administered either simultaneously or before or after administration of an antibody or antigen binding portion thereof of the present invention.

Various aspects of the invention are described in further detail in the following 5 subsections.

II. Methods for Producing Antibodies of the Invention

(i) Monoclonal Antibodies

Monoclonal antibodies of the invention can be produced using a variety of 10 known techniques, such as the standard somatic cell hybridization technique described by Kohler and Milstein (1975) *Nature* 256: 495, viral or oncogenic transformation of B lymphocytes or phage display technique using libraries of human antibody genes. In particular embodiments, the antibodies are fully human monoclonal antibodies.

Accordingly, in one embodiment, a hybridoma method is used for producing an 15 antibody that binds ErbB3. In this method, a mouse or other appropriate host animal can be immunized with a suitable antigen in order to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the antigen used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes can then be fused with myeloma cells using a suitable fusing agent, such as polyethylene 20 glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution 25 procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal. The monoclonal antibodies secreted by the subclones can be separated from the culture medium, ascites

fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

In another embodiment, antibodies and antibody portions that bind ErbB3 can be

5 isolated from antibody phage libraries generated using the techniques described in, for example, McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991), Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) and Hoet *et al* (2005) *Nature Biotechnology* 23, 344-348 ; U.S. Patent Nos. 5,223,409; 5,403,484; and 5,571,698 to Ladner *et al.*; U.S. Patent Nos. 5,427,908 and 5,580,717 to Dower *et al.*;

10 U.S. Patent Nos. 5,969,108 and 6,172,197 to McCafferty *et al.*; and U.S. Patent Nos. 5,885,793; 6,521,404; 6,544,731; 6,555,313; 6,582,915 and 6,593,081 to Griffiths *et al.*. Additionally, production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage

15 libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)) may also be used.

In a particular embodiment, the monoclonal antibody or antigen binding portion thereof that binds ErbB3 is produced using the phage display technique described by Hoet *et al.*, *supra*. This technique involves the generation of a human Fab library having a unique combination of immunoglobulin sequences isolated from human donors and

20 having synthetic diversity in the heavy-chain CDRs is generated. The library is then screened for Fabs that bind to ErbB3.

In yet another embodiment, human monoclonal antibodies directed against ErbB3 can be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse system (see e.g., Lonberg, *et al.* (1994) *Nature* 368(6474): 856-859; Lonberg, N. *et al.* (1994), *supra*; reviewed in Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Lonberg, N. and Huszar, D. (1995) *Intern. Rev. Immunol.* 13: 65-93, and Harding, F. and Lonberg, N. (1995) *Ann. N.Y. Acad. Sci.* 764:536-546. See further, U.S. Patent Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; all to Lonberg and Kay; U.S. Patent No. 5,545,807 to Surani *et al.*; PCT Publication Nos. WO 92/03918, WO 93/12227, WO 94/25585, WO 97/13852, WO 98/24884 and WO 99/45962, all to Lonberg and Kay; and PCT Publication No. WO 01/14424 to Korman *et al.*).

In another embodiment, human antibodies of the invention can be raised using a mouse that carries human immunoglobulin sequences on transgenes and transchromosomes, such as a mouse that carries a human heavy chain transgene and a human light chain transchromosome (see e.g., PCT Publication WO 02/43478 to Ishida *et al.*).

Still further, alternative transgenic animal systems expressing human immunoglobulin genes are available in the art and can be used to raise anti-ErbB3 antibodies of the invention. For example, an alternative transgenic system referred to as the Xenomouse (Abgenix, Inc.) can be used; such mice are described in, for example, 5 U.S. Patent Nos. 5,939,598; 6,075,181; 6,114,598; 6, 150,584 and 6,162,963 to Kucherlapati *et al.*

Moreover, alternative transchromosomal animal systems expressing human immunoglobulin genes are available in the art and can be used to raise anti-ErbB3 antibodies of the invention. For example, mice carrying both a human heavy chain 10 transchromosome and a human light chain transchromosome can be used; as described in Tomizuka *et al.* (2000) *Proc. Natl. Acad. Sci. USA* 97:722-727. Furthermore, cows carrying human heavy and light chain transchromosomes have been described in the art (Kuroiwa *et al.* (2002) *Nature Biotechnology* 20:889-894) and can be used to raise anti-ErbB3 antibodies of the invention.

15 In yet another embodiment, antibodies of the present invention can be prepared using a transgenic plant and/or cultured plant cells (such as, for example, tobacco, maize and duckweed) that produce such antibodies. For example, transgenic tobacco leaves expressing antibodies or antigen binding portions thereof can be used to produce such antibodies by, for example, using an inducible promoter (see, e.g., Cramer *et al.*, *Curr. 20 Top. Microbiol. Immunol.* 240:95 118 (1999)). Also, transgenic maize can be used to express such antibodies and antigen binding portions thereof (see, e.g., Hood *et al.*, *Adv. Exp. Med. Biol.* 464:127 147 (1999)). Antibodies can also be produced in large amounts from transgenic plant seeds including antibody portions, such as single chain antibodies (scFv's), for example, using tobacco seeds and potato tubers (see, e.g., Conrad *et al.*, 25 *Plant Mol. Biol.* 38:101 109 (1998)). Methods of producing antibodies or antigen binding portions in plants can also be found in, e.g., Fischer *et al.*, *Biotechnol. Appl. Biochem.* 30:99 108 (1999), Ma *et al.*, *Trends Biotechnol.* 13:522 7 (1995); Ma *et al.*, *Plant Physiol.* 109:341 6 (1995); Whitelam *et al.*, *Biochem. Soc. Trans.* 22:940 944 (1994) and U.S. Patent Nos. 6,040,498 and 6,815,184.

30 The binding specificity of monoclonal antibodies or portions thereof that bind ErbB3 prepared using any technique including those disclosed here, can be determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). The binding affinity of a monoclonal antibody or portion thereof also can be determined by the Scatchard analysis 35 of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

In certain embodiments, an ErbB3 antibody or portion thereof produced using any of the methods discussed above may be further altered or optimized to achieve a

desired binding specificity and/or affinity using art recognized techniques, such as those described herein.

In one embodiment, partial antibody sequences derived from an ErbB3 antibody may be used to produce structurally and functionally related antibodies. For example, 5 antibodies interact with target antigens predominantly through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to 10 express recombinant antibodies that mimic the properties of specific naturally occurring antibodies by constructing expression vectors that include CDR sequences from the specific naturally occurring antibody grafted onto framework sequences from a different antibody with different properties (see, e.g., Riechmann, L. *et al.*, 1998, *Nature* 332:323-327; Jones, P. *et al.*, 1986, *Nature* 321:522-525; and Queen, C. *et al.*, 1989, *Proc. Natl. Acad. See. U.S.A.* 86:10029-10033). Such framework sequences can be obtained from 15 public DNA databases that include germline antibody gene sequences.

Thus, one or more structural features of an anti-ErbB3 antibody of the invention, such as the CDRs, can be used to create structurally related anti-ErbB3 antibodies that retain at least one functional property of the antibodies of the invention, e.g., inhibiting 20 EGF-like ligand mediated phosphorylation of ErbB3; inhibiting one or more of heregulin, epiregulin, epigen or biregulin-mediated signaling through ErbB3; inhibiting proliferation of cells expressing ErbB3; and/or decreasing levels of ErbB3 on cell surfaces.

In a particular embodiment, one or more CDR regions selected from SEQ ID 25 NOS:7-12, SEQ ID NOS:13-18, SEQ ID NOS:19-24, SEQ ID NOS:39-44, and SEQ ID NOS:45-50 is combined recombinantly with known human framework regions and CDRs to create additional, recombinantly-engineered, anti-ErbB3 antibodies of the invention. The heavy and light chain variable framework regions can be derived from the same or different antibody sequences.

It is well known in the art that antibody heavy and light chain CDR3 domains 30 play a particularly important role in the binding specificity/affinity of an antibody for an antigen (see, Hall *et al.*, *J. Immunol.*, 149:1605-1612 (1992); Polymenis *et al.*, *J. Immunol.*, 152:5318-5329 (1994); Jahn *et al.*, *Immunobiol.*, 193:400-419 (1995); Klimka *et al.*, *Brit. J. Cancer*, 83:252-260 (2000); Beiboer *et al.*, *J. Mol. Biol.*, 296:833-849 35 (2000); Rader *et al.*, *Proc. Natl. Acad. Sci. USA*, 95:8910-8915 (1998); Barbas *et al.*, *J. Am. Chem. Soc.*, 116:2161-2162 (1994); Ditzel *et al.*, *J. Immunol.*, 157:739-749 (1996)). Accordingly, in certain embodiments, antibodies are generated that include the heavy and/or light chain CDR3s of the particular antibodies described herein (e.g., SEQ ID

NOs:9, 15, 21, 41, 47 and/or SEQ ID NOs:12, 18, 24, 44, 50). The antibodies can further include the heavy and/or light chain CDR1 and/or CDR2s of the antibodies of the present invention (e.g., SEQ ID NOs:7-8 and/or SEQ ID NOs:10-11; SEQ ID NOs:13-14 and/or SEQ ID NOs:16-17; SEQ ID NOs:20-21 and/or SEQ ID NOs:22-23; SEQ ID NOs:39-40 and/or SEQ ID NOs:42-43; or SEQ ID NOs:45-46 and/or SEQ ID NOs:48-49).

The CDR1, 2, and/or 3 regions of the engineered antibodies described above can comprise the exact amino acid sequence(s) as those disclosed herein (e.g., CDRs of Ab #6, Ab #3, Ab #14, Ab #17, or Ab #19, set forth in SEQ ID NOs:7-12, 13-18, 19-24, 39-44, and 45-50, respectively). However, the ordinarily skilled artisan will appreciate that some deviation from the exact CDR sequences may be possible while still retaining the ability of the antibody to bind ErbB3 effectively (e.g., conservative amino acid substitutions). Accordingly, in another embodiment, the engineered antibody may be composed of one or more CDRs that are, for example, 90%, 95%, 98%, 99% or 99.5% identical to one or more CDRs of Ab #6, Ab #3 or Ab #14.

In another embodiment, one or more residues of a CDR may be altered to modify binding to achieve a more favored on-rate of binding. Using this strategy, an antibody having ultra high binding affinity of, for example, 10^{10} M^{-1} or more, can be achieved. Affinity maturation techniques, well known in the art and those described herein, can be used to alter the CDR region(s) followed by screening of the resultant binding molecules for the desired change in binding. Accordingly, as CDR(s) are altered, changes in binding affinity as well as immunogenicity can be monitored and scored such that an antibody optimized for the best combined binding and low immunogenicity are achieved.

In addition to, or instead of, modifications within the CDRs, modifications can also be made within one or more of the framework regions, FR1, FR2, FR3 and FR4, of the heavy and/or the light chain variable regions of an antibody, so long as these modifications do not eliminate the binding affinity of the antibody.

In another embodiment, the antibody is further modified with respect to effector function, so as to enhance the effectiveness of the antibody in treating cancer, for example. For example cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC).

See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shope, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.* *Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered

which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.* *Anti-Cancer Drug Design* 3:219-230 (1989).

Also encompassed by the present invention are bispecific antibodies and immunoconjugates, as discussed below.

5

(ii) Bispecific Antibodies

Bispecific antibodies of the present invention include at least one binding specificity for ErbB3 and at least one binding specificity for another antigen, such as the product of an oncogene. Bispecific antibodies can be prepared as full length antibodies 10 or antibody fragments (e.g. F(ab')₂ bispecific antibodies).

Methods for making bispecific antibodies are well known in the art (see, e.g., WO 05117973 and WO 06091209). For example, production of full length bispecific antibodies can be based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (see, e.g., Millstein *et al.*, 15 *Nature*, 305:537-539 (1983)). Further details of generating bispecific antibodies can be found, for example, in Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986) and in Brennan *et al.*, *Science*, 229: 81 (1985), which describes a chemical linkage process for making bispecific antibodies. Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For 20 example, bispecific antibodies have been produced using leucine zippers (see, e.g., Kostelny *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992)). Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported (see, e.g., Gruber *et al.*, *J. Immunol.*, 152:5368 (1994)).

In a particular embodiment, the bispecific antibody comprises a first antibody or 25 binding portion thereof which binds to ErbB3 and a second antibody or binding portion thereof which binds to ErbB2, ERBB3, ErbB4, EGFR, IGF1-R, C-MET, Lewis Y, MUC-1, EpCAM, CA125, prostate specific membrane antigen, PDGFR- α , PDGFR- β , C-KIT, or any of the FGF receptors.

30

(iii) Immunoconjugates

Immunoconjugates of the present invention can be formed by conjugating the 35 antibodies or antigen binding portions thereof described herein to another therapeutic agent. Suitable agents include, for example, a cytotoxic agent (e.g., a chemotherapeutic agent), a toxin (e.g. an enzymatically active toxin of bacterial, fungal, plant or animal origin, or fragments thereof), and/or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A

chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), *momordica charantia* inhibitor, curcin, crotin, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated anti-ErbB3 antibodies. Examples include ²¹²Bi, ¹³¹I, ¹³¹In, ⁹⁰Y and ¹⁸⁶Re.

Immunoconjugates of the invention can be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, *Science* 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody (see, *e.g.*, WO94/11026).

20

III. Methods for Screening Antibodies of the Invention

Subsequent to producing antibodies or antigen binding portions that bind ErbB3, such antibodies, or portions thereof, can be screened for various properties, such as those described herein, using a variety of assays that are well known in the art.

25

In one embodiment, the antibodies or antigen binding portions thereof are screened for the ability to inhibit EGF-like ligand mediated phosphorylation of ErbB3. This can be done by treating cells expressing ErbB3 with an EGF-like ligand in the presence and absence of the antibody or antigen binding portion thereof. The cells can then be lysed and the crude lysates can be centrifuged to remove insoluble material.

30

ErbB3 phosphorylation can be measured, for example, by Western blotting followed by probing with an anti-phosphotyrosine antibody as described in Kim *et al.*, *supra* and the Examples below.

35

In other embodiments, the antibodies and antigen binding portions are further screened for one or more of the following properties: (1) inhibition of ErbB3-ligand (e.g., heregulin, epiregulin, epigen or biregulin) mediated signaling through ErbB3; (2) inhibition of proliferation of cells expressing ErbB3; (3) the ability to decrease levels of ErbB3 on cell surface (e.g., by inducing internalization of ErbB3), (4) inhibition of VEGF secretion of cells expressing ErbB3; (5) inhibition of the migration of cells

expressing ErbB3; (6) inhibition of spheroid growth of cells expressing ErbB3; and/or (7) binding to an epitope located on domain I of ErbB3, each of which can be readily measured using art recognized techniques and those discussed herein.

Inhibition of one or more of heregulin, epiregulin, epigen or biregulin-mediated signaling through ErbB3 can be readily measured using routine assays, such as, described in Horst *et al. supra*. For example, the ability of an antibody or antigen binding portion thereof to inhibit heregulin, epiregulin, epigen or biregulin-mediated signaling through ErbB3 can be measured by kinase assays for known substrates of ErbB3 such as, for example, SHC and PI3K, as described in, for example, Horst *et al. supra*, Sudo *et al.*, (2000) *Methods Enzymol.*, 322:388-92; and Morgan *et al.* (1990) *Eur. J. Biochem.*, 191:761-767, following stimulation by one or more of heregulin, epiregulin, epigen or biregulin. Accordingly, cells expressing ErbB3 can be stimulated with one or more of heregulin, epiregulin, epigen or biregulin, and incubated with a candidate antibody or antigen-binding portion thereof. Cell lysates subsequently prepared from such cells can be immunoprecipitated with an antibody for a substrate of ErbB3 (or a protein in a cellular pathway involving ErbB3) such as, for example, an anti-JNK-1 antibody, and assayed for kinase activity (e.g., JNK kinase activity or PI3-kinase activity) using art recognized techniques. A decrease in or complete disappearance in level or activity (e.g., kinase activity) of a ErbB3 substrate or protein in a pathway involving ErbB3 in the presence of the antibody or antigen binding portion thereof, relative to the level or activity in the absence of the antibody or antigen binding portion thereof, is indicative of an antibody or antigen binding portion which inhibits one or more of heregulin, epiregulin, epigen or biregulin-mediated signaling.

In certain embodiments, the antibody or antigen binding portion thereof inhibits ErbB3-ligand (e.g., heregulin, epiregulin, epigen or biregulin) mediated signaling by decreasing the binding of one or more of heregulin, epiregulin, epigen or biregulin to ERbB3.

In order to select for those antibodies or antigen binding portions thereof which inhibit the binding of one or more of heregulin, epiregulin, epigen or biregulin to ErbB3, cells which express ErbB3 (e.g. MALME-3M cells, as described in the Examples *infra*), can be contacted with a labeled ErbB3-ligand (e.g., radiolabeled heregulin, epiregulin, epigen or biregulin) in the absence (control) or presence of the anti-ErbB3 antibody or antigen binding portion thereof. If the antibody or antigen binding portion thereof inhibits heregulin, epiregulin, epigen or biregulin binding to ErbB3, then a statistically significant decrease in the amount of label recovered (e.g., radiolabeled heregulin, epiregulin, epigen or biregulin), relative to the amount in the absence of the antibody or antigen binding portion thereof, will be observed.

The antibody or antigen binding portion thereof may inhibit the binding of the ErbB3-ligand (e.g., heregulin, epiregulin, epigen or biregulin) by any mechanism. For example, the antibody or antigen binding portion thereof may inhibit binding of the ErbB3 ligand (e.g., one or more of heregulin, epiregulin, epigen or biregulin) to ErbB3

5 by binding to the same site or an overlapping site on ErbB3 as the ErbB3 ligand. Alternatively, the antibody or antigen binding portion thereof may inhibit binding of an ErbB3 ligand by altering or distorting the conformation of ErbB3, such that it is unable to bind to the ErbB3 ligand.

Antibodies and antigen binding portions thereof that decrease levels of ErbB3 on

10 cell surfaces can be identified by their ability to downregulate ErbB3 on tumor cells. In certain embodiments, the antibodies or antigen binding portions thereof decrease ErbB3 cell surface expression by inducing internalization (or increasing endocytosis) of ErbB3. To test this, ErbB3 can be biotinylated and the number of ErbB3 molecules on the cell surface can be readily determined, for example, by measuring the amount of biotin on a

15 monolayer of cells in culture in the presence or absence of an antibody or antigen binding portion thereof, for example, as described in, e.g., Waterman *et al.*, *J. Biol. Chem.* (1998), 273:13819-27, followed by immunoprecipitation of ErbB3 and probing with streptavidin. A decrease in detection of biotinylated ErbB3 over time in the presence of an antibody or antigen binding portion is indicative of an antibody which

20 decreases ErbB3 levels on cell surfaces.

Antibodies or antigen binding portions thereof of the present invention can also be tested for their ability to inhibit proliferation of cells expressing ErbB3, for example, tumor cells, using art recognized techniques, such as the Cell Titer Glow Assay described in the Examples below (also see, e.g., Macallan *et al.*, *Proc. Natl. Acad. Sci.* (1998) 20;95(2):708-13; Perez *et al.* (1995) *Cancer Research* 55, 392-398).

In another embodiment, the antibodies or antigen binding portions thereof are screened for the ability to inhibit VEGF secretion of cells expressing ErbB3. This can be done by using well-known assays, such as the VEGF ELISA kit available from R&D Systems (Minneapolis, MN, Cat.#DY293B). Similarly, the antibodies or portions can be

30 screened for the ability to inhibit the migration of cells expressing ErbB3 (e.g., MCF-7 cells) using a trans-well assay (Millipore Corp., Billerica, MA, Cat # ECM552) as described herein.

In another embodiment, the antibodies or antigen binding portions thereof are screened for the ability to inhibit spheroid growth of cells expressing ErbB3. This can

35 be done by using an assay which approximates conditions of a developing tumor growth (see, e.g., Herman *et al.* (2007) *Journal of Biomolecular Screening Electronic publication*) as described herein.

Antibodies or antigen binding portions thereof that bind to the same or overlapping epitopes as one or more antibodies of the present invention can also be identified using standard techniques known in the art and described herein. For example, in order to screen for antibodies which bind to the same or an overlapping 5 epitope on ErbB3 bound by an antibody of interest, a cross-blocking assay, such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed.

IV. Pharmaceutical Compositions

10 In another aspect, the present invention provides a composition, *e.g.*, a pharmaceutical composition, containing one or a combination of monoclonal antibodies, or antigen-binding portion(s) thereof, of the present invention, formulated together with a pharmaceutically acceptable carrier. In one embodiment, the compositions include a combination of multiple (*e.g.*, two or more) isolated antibodies of the invention, which 15 bind different epitopes on ErbB3.

As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or 20 epidermal administration (*e.g.*, by injection or infusion). Depending on the route of administration, the active compound, *i.e.*, antibody, bispecific and multispecific molecule, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

A “pharmaceutically acceptable salt” refers to a salt that retains the desired 25 biological activity of the parent compound and does not impart any undesired toxicological effects (see *e.g.*, Berge, S.M., *et al.* (1977) *J. Pharm. Sci.* 66:1-19). Examples of such salts include acid addition salts and base addition salts. Acid addition salts include those derived from nontoxic inorganic acids, such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, phosphorous and the like, as well as from 30 nontoxic organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanoic acids, hydroxy alkanoic acids, aromatic acids, aliphatic and aromatic sulfonic acids and the like. Base addition salts include those derived from alkaline earth metals, such as sodium, potassium, magnesium, calcium and the like, as well as from nontoxic organic amines, such as N,N'-dibenzylethylenediamine, N- 35 methylglucamine, chloroprocaine, choline, diethanolamine, ethylenediamine, procaine and the like.

Pharmaceutical compositions of the invention can be administered alone or in combination therapy, *i.e.*, combined with other agents. For example, the combination

therapy can include a composition of the present invention with at least one or more additional therapeutic agents, such as the anti-cancer agents described infra. The pharmaceutical compositions of the invention can also be administered in conjunction with radiation therapy and/or surgery.

5 A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. The active compounds can be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal 10 patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. *See, e.g., Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel 15 Dekker, Inc., New York, 1978.

To administer a compound of the invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. 20 Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan *et al.* (1984) *J. Neuroimmunol.* 7:27).

25 Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the invention is contemplated. Supplementary active compounds can also be incorporated into the compositions.

30 Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid 35 polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars,

polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

5 Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from 10 those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

15 Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. For example, the human antibodies of the invention may be administered once or twice weekly by subcutaneous injection or once or twice monthly by subcutaneous injection.

20 It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

25 Examples of pharmaceutically-acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

30 For the therapeutic compositions, formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in

unit dosage form and may be prepared by any methods known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the subject being treated, and the particular mode of administration. The amount of active ingredient which can be 5 combined with a carrier material to produce a single dosage form will generally be that amount of the composition which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 0.001 per cent to about ninety percent of active ingredient, preferably from about 0.005 per cent to about 70 per cent, most preferably from about 0.01 per cent to about 30 per cent.

10 Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate. Dosage forms for the topical or transdermal administration of compositions of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and 15 inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

20 The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, 25 subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

30 Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

35 These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Particular examples of adjuvants which are well-known in the art include, for example, inorganic adjuvants (such as aluminum salts, e.g., aluminum phosphate and aluminumhydroxide), organic adjuvants (e.g., squalene), oil-based adjuvants, virosomes (e.g., virosomes which contain a membrane-bound heagglutinin and neuraminidase derived from the influenza virus).

Prevention of presence of microorganisms may be ensured both by sterilization procedures, *supra*, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they can be given alone or as a pharmaceutical composition containing, for example, 0.001 to 90% (more preferably, 0.005 to 70%, such as 0.01 to 30%) of active ingredient in combination with a pharmaceutically acceptable carrier.

Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts. A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a composition of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. It is preferred that administration be intravenous, intramuscular, intraperitoneal, or subcutaneous, preferably administered proximal to the site of the target. If desired, the effective daily dose of a therapeutic

composition may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical formulation (composition).

Therapeutic compositions can be administered with medical devices known in the art. For example, in a preferred embodiment, a therapeutic composition of the invention can be administered with a needleless hypodermic injection device, such as the devices disclosed in U.S. Patent Nos. 5,399,163, 5,383,851, 5,312,335, 5,064,413, 10 4,941,880, 4,790,824, or 4,596,556. Examples of well-known implants and modules useful in the present invention include: U.S. Patent No. 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; U.S. Patent No. 4,486,194, which discloses a therapeutic device for administering medications through the skin; U.S. Patent No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; U.S. Patent 15 No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; U.S. Patent No. 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments; and U.S. Patent No. 4,475,196, which discloses an osmotic drug delivery system. Many other such implants, delivery systems, and modules are known to those skilled in the art.

In certain embodiments, the monoclonal antibodies of the invention can be formulated to ensure proper distribution *in vivo*. For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds of the invention cross the BBB (if desired), they can be formulated, for example, in liposomes. For methods of manufacturing liposomes, see, e.g., U.S. Patents 4,522,811; 5,374,548; and 5,399,331. The liposomes may comprise one or more moieties which are selectively transported into specific cells or organs, thus enhance targeted drug delivery (see, e.g., V.V. Ranade (1989) *J. Clin. Pharmacol.* 29:685). Exemplary targeting moieties include folate or biotin (see, e.g., U.S. Patent 5,416,016 to 20 Low *et al.*); mannosides (Umezawa *et al.*, (1988) *Biochem. Biophys. Res. Commun.* 153:1038); antibodies (P.G. Bloeman *et al.* (1995) *FEBS Lett.* 357:140; M. Owais *et al.* (1995) *Antimicrob. Agents Chemother.* 39:180); surfactant protein A receptor (Briscoe *et al.* (1995) *Am. J. Physiol.* 1233:134), different species of which may comprise the formulations of the inventions, as well as components of the invented molecules; p120 (Schreier *et al.* (1994) *J. Biol. Chem.* 269:9090); see also K. Keinanen; M.L. Laukkanen 35 (1994) *FEBS Lett.* 346:123; J.J. Killion; I.J. Fidler (1994) *Immunomethods* 4:273.

V. Methods of Using Antibodies of the Invention

The present invention also provides methods of using antibodies and antigen-binding portions thereof that bind ErbB3 in a variety of *ex vivo* and *in vivo* diagnostic and therapeutic applications. For example, antibodies of the invention can be used for 5 treating a disease associated with ErbB3 dependent signaling, including a variety of cancers.

In one embodiment, the present invention provides a method for treating a disease associated with ErbB3 dependent signaling by administering to a subject an antibody or antigen binding portion thereof of the invention in an amount effective to 10 treat the disease. Suitable diseases include, for example, a variety of cancers including, but not limited to, melanoma, breast cancer, ovarian cancer, renal carcinoma, gastrointestinal cancer, colon cancer, lung cancer, and prostate cancer.

The antibody can be administered alone or with another therapeutic agent which acts in conjunction with or synergistically with the antibody to treat the disease 15 associated with ErbB3 mediated signaling. Such therapeutic agents include, for example, the anticancer agents described infra (e.g., cytotoxins, chemotherapeutic agents, small molecules and radiation).

In another embodiment, the present invention provides a method for diagnosing a disease (e.g., a cancer) associated with ErbB3 upregulation in a subject, by contacting 20 antibodies or antigen binding portions of the invention (e.g., *ex vivo* or *in vivo*) with cells from the subject, and measuring the level of binding to ErbB3 on the cells. Abnormally high levels of binding to ErbB3 indicate that the subject has a disease associated with ErbB3 upregulation.

Also within the scope of the present invention are kits comprising antibodies and 25 antigen binding portions thereof of the invention which optionally include instructions for use in treating or diagnosing a disease associated with ErbB3 upregulation and/or ErbB3 dependent signaling. The kits may include a label indicating the intended use of the contents of the kit. The term label includes any writing, marketing materials or recorded material supplied on or with the kit, or which otherwise accompanies the kit.

30 Other embodiments of the present invention are described in the following Examples.

The present invention is further illustrated by the following examples which should not be construed as further limiting. The contents of Sequence Listing, figures and all references, patents and published patent applications cited throughout this 35 application are expressly incorporated herein by reference.

Examples**Materials and Methods**

Throughout the examples, the following materials and methods were used unless 5 otherwise stated.

In general, the practice of the present invention employs, unless otherwise indicated, conventional techniques of chemistry, molecular biology, recombinant DNA technology, immunology (especially, e.g., antibody technology), and standard techniques in polypeptide preparation. See, e.g., Sambrook, Fritsch and Maniatis, 10 *Molecular Cloning: Cold Spring Harbor Laboratory Press* (1989); *Antibody Engineering Protocols (Methods in Molecular Biology)*, 510, Paul, S., Humana Pr (1996); *Antibody Engineering: A Practical Approach* (Practical Approach Series, 169), McCafferty, Ed., Irl Pr (1996); *Antibodies: A Laboratory Manual*, Harlow et al., C.S.H.L. Press, Pub. (1999); and *Current Protocols in Molecular Biology*, eds. Ausubel 15 et al., John Wiley & Sons (1992). *In vitro* and *in vivo* model systems for assaying HCV biology are described, for example, in *Cell culture models and animal models of viral hepatitis. Part II: hepatitis C, Lab. Anim. (NY)*, 34(2):39-47 (2005) and in *The chimpanzee model of hepatitis C virus infections, ILAR J.*, 42(2):117-26 (2001).

20 **Cell Lines**

All the cell lines used in the experiments described below were obtained from the National Cancer Institute or provided by investigators, as indicated.

Cell Lines:

25 MCF7- ATCC cat. No. HTB-22
T47D- ATCC cat. No. HTB-133
Colo357- These cells were obtained from an academic investigator and are described by Kolb et al. (2006) *Int. J. Cancer*, 120:514-523.
Du145- ATCC cat. No. HTB-81
OVCAR8- source already described in provisional application.
30 H1975 ATCC cat. No. CRL-5908

Pulverization of Tumor Cells

A cryopulverizer (Covaris Inc) was used for the pulverization of tumors. Tumors were stored in special bags (pre-weighed before the addition of the tumor) and placed in liquid nitrogen while handling them. For small tumors, 200 uL of Lysis buffer was first 35 added to the bag containing the tumor, frozen in liquid nitrogen and then pulverized to improve the recovery of the tumor from the bag. Pulverized tumors were transferred to 2 mL Eppendorf tubes and placed in liquid nitrogen until ready for further processing

Lysis of Tumor Cells

Tumors were lysed in Lysis buffer supplemented with protease and phosphatase inhibitors. Lysis Buffer was added to the tumor aliquots in a final concentration of about 62.5 mg/mL. Tumor samples were homogenized by vortexing for 30 sec and incubating on ice for about 30 min. The lysates were spun for about 10 min in Qiagen Qiashredder 5 columns for further homogenization of the samples. Cleared lysates were aliquoted into fresh tubes for further processing.

BCA Assay

BCA assay (Pierce) was performed following the manufacturer's protocol on all tumor samples. The total protein concentration (in mg/mL) of each tumor sample was 10 later used in the normalization of the ELISA results

ELISA Assay

All ELISA reagents for the total and phospho-ErbB3 ELISAs were purchased from R&D Systems as DuoSet kits. 96-well Nunc Maxisorb plates were coated with 50 uL of an antibody and incubated overnight at room temperature. Next morning, plates 15 were washed 3 times with 1000 µl/well in the BioTek plate washer with PBST (0.05% Tween-20). Plates were subsequently blocked for about an 1 hr at room temperature with 2 % BSA in PBS. The plates were washed 3 times with 1000 µl/well in the BioTek plate washer with PBST (0.05%Tween-20). 50 µL of cell lysates and standards diluted in 50% Lysis buffer and 1%BSA were used in duplicates for further processing. Samples 20 were incubated for 2hrs at 4°C on a plate shaker and washed 3 times with 1000 µl/well in the BioTek plate washer with PBST (0.05%Tween-20). About 50 µl of a detection antibody diluted in 2% BSA, PBST was added and incubated for about 1 hr at room temperature. For phospho-ErbB3, the detection antibody was directly conjugated to horse radish peroxidase (HRP) and incubated for 2 hrs at room temperature. The plate 25 was washed 3 times with 1000 µl/well in the BioTek plate washer with PBST (0.05%Tween-20). About 50 µl of Streptavidin-HRP was added and incubate for 30 min at room temperature (except for pErbB3). The plates were washed 3 times with 1000 µl/well in the BioTek plate washer with PBST (0.05%Tween-20). About 50 µL of Supersignal Pico ELISA substrate was added and the plate was read using a Fusion plate 30 reader. The data was analysed using EXCEL. Duplicate samples were averaged and the error bars were used to represent the standard deviation between the two replicates.

Example 1: Production of Antibodies Using Phage Display

In order to obtain human anti-ErbB3 antibodies referred to herein as Ab #6, Ab 35 #3, Ab #14, Ab #17, and Ab #19, a human Fab-phage library including a unique combination of immunoglobulin sequences obtained from human donors (Hoet *et al. supra*, incorporated by reference in its entirety herein) was initially screened for ErbB3 binders.

Using purified ErbB3 and a Chinese hamster ovary cell line expressing cell surface ErbB3, 73 unique Fab sequences from the library were identified. These 73 clones were then reformatted as Fab only without the phage. Using high throughput methods, these Fabs were expressed on a small scale and tested for binding using ELISA and the Flexchip method which is a high-throughput surface plasmon resonance (SPR) technology. The 73 Fabs without the phage were spotted on a chip surface and the binding kinetics and epitope blocking to a ErbB3-his fusion target protein or a ErbB3-Fcprotein (R & D Systems) were measured. The equilibrium binding constant and on/off rates for the Fabs were calculated from the data obtained.

10 Binding of the various Fabs to MALME-3M cells was next examined using about 500 nM of the Fabs and a 1:750 dilution of a goat anti-human Alexa 647 secondary antibody. As shown in Figures 1A and 1B, several candidate Fabs exhibited appreciable staining of MALME-3M cells.

15 **Example 2: Optimization of anti-ErbB3 Fabs**

Subsequent to the identification of Fabs which blocked the binding of ErbB3 ligand, heregulin, to ErbB3, the VH and VL sequences of the Fabs were codon-optimized as follows.

20 Specifically, the VH and VL regions were reformatted using expression constructs for expression as an IgG1 or IgG2 isotype. The constructs included a Selexis backbone which has a cassette designed for substitution of the appropriate heavy and light chain sequences. The Selexis vectors included a CMV promoter and a matching poly-A signal.

25 The nucleic acid sequences for the codon-optimized VH and VL of Ab #6 are set forth in SEQ ID NOS:25 and 26, respectively, and those for Ab #3 are set forth in SEQ ID NOS:27 and 28, respectively, as shown in Figure 22.

Example 3: Binding Affinity for ErbB3

30 The dissociation constants of the anti-ErbB3 antibodies were measured using two independent techniques, *i.e.*, a Surface Plasmon Resonance Assay and a cell binding assay using MALME-3M cells.

Surface Plasmon Resonance Assay

35 The Surface Plasmon Resonance Assay (also referred to as the Flexchip assay) was performed as described in Wassaf *et al.* (2006) *Analytical Biochem.*, 351:241-253. The K_D value was calculated based on the formula $K_D = K_d/K_a$.

The K_D values of Ab #6 and Ab #3, respectively, as measured using the Surface Plasmon Resonance Assay, are depicted in Figures 2A and 2B. Ab #6 had a K_D value of

about 4 nM and Ab #3 had a K_D value of about 8 nM, as depicted in Figures 2A and 2B, respectively.

Cell Binding Assay

5 The cell binding assay for determining the K_D values of Ab #6 and Ab #3 was performed as follows.

10 MALME-3M cells were detached with 2 mls trypsin-EDTA + 2 mls RPMI + 5mM EDTA at room temperature for 5 minutes. Complete RPMI (10 mls) was added immediately to the trypsinized cells, resuspended gently and spun down in a Beckman tabletop centrifuge at 1100 rpm for 5 minutes. Cells were resuspended in BD stain buffer (PBS + 2% fetal bovine serum + 0.1% sodium azide, Becton Dickinson) at a concentration of 2×10^6 cells per ml and 50 μ l (1×10^5 cells) aliquots were plated in a 96-well titer plate.

15 A 150 μ l solution of 200 nM anti-ErbB3 antibody (Ab #6 or Ab #3) in BD stain buffer was prepared in an eppendorf tube and serially diluted 2-fold into 75 μ l BD stain buffer. The concentrations of the diluted antibody ranged from 200 nM to 0.4 nM. 50 μ l aliquots of the different protein dilutions were then added directly to the 50 μ l cell suspension giving the final concentrations of 100 nM, 50 nM, 25 nM, 12 nM, 6 nM, 3 nM, 1.5 nM, 0.8 nM, 0.4 nM and 0.2 nM of the antibody.

20 Aliquotted cells in the 96-well plate were incubated with the protein dilutions for 30 minutes at room temperature on a platform shaker and washed 3 times with 300 μ l BD stain buffer. Cells were then incubated with 100 μ l of a 1:750 dilution of Alexa 647-labeled goat anti-human IgG in BD stain buffer for 45 minutes on a platform shaker in the cold room. Finally, cells were washed twice, pelleted and resuspended in 250 μ l 25 BD stain buffer + 0.5 μ g/ml propidium iodide. Analysis of 10,000 cells was done in a FACScalibur flow cytometer using the FL4 channel. MFI values and the corresponding concentrations of the anti-ErbB3-antibodies were plotted on the y-axis and x-axis, respectively. The K_D of the molecule was determined by GraphPad Prism using the one-site binding model for a non-linear regression curve.

25 The K_D value was calculated based on the formula $Y = B_{max} * X / K_D + X$ (B_{max} = fluorescence at saturation. X = antibody concentration. Y = degree of binding). As shown in Figures 2C and 2D, Ab # 6 and Ab #3 had K_D values of about 4 nM and 1.3 nM, respectively, in a cell binding assay using MALME-3M cells.

30 **Example 4: Binding Specificity / Epitope Binding for ErbB3**

The binding specificity of an IgG2 isotype of Ab #6 to ErbB3 was assayed using ELISA as follows. Identification of the epitope bound by Ab #6 was also analyzed.

Specifically, 96-well Nunc Maxisorb plates were coated with 50 μ l of 5 μ g/ml protein (recombinant human ErbB3, recombinant human EGFR or unrelated protein (BSA)) and incubated overnight at room temperature. The next morning, plates were washed 3 times with 1000 μ l/well of PBST (0.05% Tween-20) in the BioTek plate washer. The wells were blocked for 1 hr at room temperature with 2 % BSA in PBS. The plates were washed 3 times with 1000 μ l/well of PBST (0.05% Tween-20) in the BioTek plate washer. About 50 μ L of the Ab #6 was added at several dilutions (1 μ M and serial 2 fold dilutions) in 2% BSA, PBST. All samples were run in duplicate and incubated for 2hrs at 4°C on a plate shaker. The plates were washed plates 3 times with 1000 μ l/well of PBST (0.05% Tween-20) in the BioTek plate washer. 50 μ l of human IgG detection antibody (HRP conjugated (Bethyl Inc; 1:75000 dilution in 2% BSA, PBST)) was added and the plates were incubated for 1 hr at room temperature. The plates were washed plate 3 times with 1000 μ l/well of PBST (0.05% Tween-20) in the BioTek plate washer. 50 μ L of Supersignal Pico ELISA substrate was added and the plate was read on the Fusion plate reader. The data was analyzed using the EXCEL program. Duplicate samples were averaged and the error bars represent the standard deviation between the two replicates.

As shown in Figure 3, Ab #6 bound recombinant ErbB3 in an ELISA, but did not show any appreciable binding to EGFR, BSA or TGF- α .

20 A fragment (truncation mutant) corresponding to amino acid residues 20-202 of ErbB3 was cloned into the yeast display vector pYD2 (a modified version of pYD1 (Invitrogen) with a stop codon engineered in front of the His tag) between the Nhe and BsiWI restriction sites. The plasmid was transformed into the yeast strain EBY100 (Invitrogen) and clones containing the plasmid selected on Trp- selective medium. The 25 clone was grown in glucose containing medium overnight at 30°C and expression of the ErbB3 truncation mutant was induced by transfer to a galactose-containing medium for 2 days at 18°C. Yeast displaying the ErbB3 truncation mutant were stained with 50 nM of Ab #6, followed by a goat anti-human antibody labeled with Alexa dye-647. A separate sample was stained with the goat anti-human antibody only to show that there is 30 no non-specific binding to yeast of the secondary antibody. Analysis was performed by flow cytometry on the FACS Calibur cell sorter (BD Biosciences).

As shown in Figure 30, Ab #6 bound to the truncation mutant, i.e., amino acid residues 20-202 of ErbB3.

35 **Example 5: Downregulation of Total ErbB3 on Tumor Cells**

The ability of Ab #6 to downregulate ErbB3 expression both *in vitro* and *in vivo* in tumor cells was tested as follows.

MALME-3M cells were seeded in 96 well tissue culture plates and grown in RPMI-1640 media supplemented with antibiotics, 2mM L-glutamine and 10% fetal bovine serum (FBS) for 24 hours at 37°C and 5% carbon dioxide. Media was then switched to RPMI-1640 MEDIA with antibiotics, 2mM L-glutamine with and without 5 the antibody at concentrations of 1uM, 250nM, 63nM, 16nM, 4.0nM, 1.0nM, 240pM, 61pM and 15pM. Cells were grown for 24 hours at 37°C and 5% carbon dioxide, washed with cold PBS, then harvested with mammalian protein extract (MPER) lysis (Pierce, 78505) buffer containing 150mM NaCl, 5mM sodium pyrophosphate, 10uM bpV (phen), 50uM phenalarsine, 1mM sodium orthovanadate, and protease inhibitor 10 cocktail (Sigma, P714). Cell lysates were diluted two-fold with 4% bovine serum albumin in phosphate buffered saline with 0.1% tween-20, then analyzed by ELISA with mouse anti-human ErbB3 capture antibody and biotinylated mouse anti-human ErbB3 secondary detection antibody. Signal was generated with streptavidin conjugated to horseradish-peroxidase reacted with chemiluminescent substrate (Pierce, 37070). 15 ELISAS were visualized using a luminometer.

As shown in Figure 4, Ab #6 decreased total ErbB3 levels by about 46.9% in MALME-3M cells *in vitro*, as measured by ELISA. Media containing no serum and antibody was used as control.

In a further experiment, the downregulation of ErbB3 receptors on MALME-3M 20 cells using IgG1 and IgG2 isotypes of Ab #6 was examined using FACS analysis. MALME-3M cells were trypsinized from a 15 cm dish and washed once with RPMI + 10% fetal bovine serum. Cell pellets were resuspended at a density of 1×10^6 cells per ml. Two aliquots of 2×10^5 cells were added to a 12-well tissue culture plate and resuspended in a final volume of 800 ul RPMI + 10% fetal bovine serum. To one well, 25 Ab #6 IgG1 or Ab #6IgG2 isotype was added to a final concentration of 100 nM (treated sample) and to the other well, an equivalent volume of PBS was (untreated sample) was added.

The following day, treated and untreated cells were trypsinized, washed and 30 incubated with 100 nM of Ab #6 in BD stain buffer for 30 minutes on ice. Cells were washed twice with 1 ml BD stain buffer and incubated with 100 ul of a 1:500 dilution of Alexa 647-labeled goat anti-human Alexa 647 for 45 minutes on ice. Cells were then washed and resuspended in 300 ul BD stain buffer + 0.5 ug/ml propidium iodide. Analysis of 10,000 cells was done in a FACScalibur flow cytometer using the FL4 channel.

35 As shown in Figures 5A and 5B, both IgG1 and IgG2 isotypes of Ab #6 downregulated ErbB3 on MALME-3M cells by about 62% and about 66%, respectively.

In order to determine whether this decrease was due to internalization of the ErbB3 receptor on the surface of MALME-3M cells, the expression of ErbB3 in the

presence of the antibody over time was measured. Specifically, MALME-3M cells were trypsinized from a 15 cm dish and washed once with RPMI + 10% fetal bovine serum. Cell pellets were resuspended at a density of 1×10^6 cells per ml. Two aliquots of 2×10^5 cells were added to a 12-well tissue culture plate and resuspended in a final volume 5 of 800 μ l RPMI + 10% fetal bovine serum. To one well, the anti-ErbB3 antibody was added to a final concentration of 100 nM (treated sample) and to the other well, an equivalent volume of PBS was (untreated sample) was added. The following day, treated and untreated cells were trypsinized, washed and incubated with 100 nM anti-ErbB3 antibody in BD stain buffer for 30 minutes on ice. Cells were washed twice with 10 1 ml BD stain buffer and incubated with 100 μ l of a 1:500 dilution of Alexa 647-labeled goat anti-human Alexa 647 for 45 minutes on ice. Cells were then washed and resuspended in 300 μ l BD stain buffer + 0.5 μ g/ml propidium iodide. Analysis of 10,000 cells was done in a FACScalibur flow cytometer using the FL4 channel.

As shown in Figure 6, downregulation of ErbB3 in the presence of Ab #6 was 15 measured at 0 hour (Figure 6A), 0.5 hour (Figure 6B), 2 hour (Figure 6C) and 24 hours (Figure 6D). As shown in Figure 6A-6D, about 50% of the cell surface ErbB3 receptors were downregulated after about 30 minutes and at about 24 hours, about 93% of the cell surface receptors were downregulation.

The ability of Ab #6 to cause ErbB3 downregulation *in vivo* in melanoma cells 20 was also examined as follows.

Briefly, T-cell deficient nu/nu mice (3-4 week old female mice originated at NIH; outbred; albino background) were purchased from Charles River Labs (Wilmington, MA). MALME-3M cells for implantation were grown in culture (RPMI media, 10% FBS, L-glutamine and antibiotics, 37°C, 5% CO₂) to about 80% confluence 25 before harvesting. Cells were kept on ice until implantation. Mice were implanted via subcutaneous injection with 100ul MALME-3M cells on the right flank and allowed to recover while being monitored for initial tumor growth.

The tumors were measured (length by width) by digital caliper and the mice were dosed with IgG2a (Sigma, M7769-5MG) by intravenous injection. Mice were dosed 30 intra-peritoneally every other day with either 15 μ g or 100 μ g of antibody number 6 and tumors were measured three times per week and recorded in a Microsoft EXCEL spreadsheet.

Final tumor measurements (L x W) were taken, the mice were euthanized by CO₂ asphyxiation and tumors were excised, snap frozen in liquid nitrogen, and were 35 stored at -80°C (for biochemical analysis). Final tumor measurements were analyzed and graphed by tumor area and tumor volume as described, for example, in Burtrum *et al.*, (2003) *Cancer Res.*, 63:8912-8921. The data was also analyzed by "normalized" and "non-normalized" means for both tumor volume and tumor area. For the

“normalization” of the data, at each time point of measurement, each tumor in each group was divided by the initial tumor size determined by caliper measurement.

As shown in Figure 7, among the various antibodies tested in this assay, Ab #6 caused downregulation of total ErbB3 as soon as 24 hours post-injection in tumors 5 treated with either IgG1 or IgG2 isotype of Ab #6). PBS was used as a control

In a further experiment, the ability of Ab #6 to downregulate ErbB3 in ADR xenografts *in vivo* was examined.

Briefly, the samples were pulverized in a cryopulverizer (Covaris Inc). Tumors were stored in special bags (pre-weighed before the addition of the tumor) and placed in 10 liquid nitrogen while handling them. For small tumors, 200 μ L of Lysis buffer was first added to the bag with the tumor, frozen in liquid nitrogen and then pulverized to improve the recovery of the tumor from the bag. Pulverized tumors were transferred to 2 ml Eppendorf tubes and placed in liquid nitrogen until lysed. Tumors were lysed in Lysis buffer supplemented with protease and phosphatase inhibitors. Lysis Buffer was 15 added to the tumor aliquots in a final concentration of 62.5 mg/ml. Tumor samples were homogenized by vortexing for 30 seconds and letting them sit on ice for 30 min. The lysates were spun for 10 minutes in Qiagen Qiashredder columns for further homogenization of the samples. Cleared lysates were aliquoted into fresh tubes.

The BCA assay was performed as set forth in the materials and methods section 20 supra.

The total levels of ErbB3 were determined by ELISA. The ELISA reagents were purchased from R&D Systems as DuoSet kits. 96-well Nunc Maxisorb plates were 25 coated with 50 μ l of respective capture antibody and incubated overnight at room temperature. The next morning, the plates were washed 3 times with 1000 μ l/well in a BioTek plate washer with PBST (0.05% Tween-20) and then blocked for 1 hour at room temperature with 2 % BSA in PBS. The plates were then washed three times with 1000 μ l/well in the BioTek plate washer with PBST (0.05%Tween-20). Lysates (50 μ l) and standards were diluted in 50% Lysis buffer and 1%BSA; all samples were run in 30 duplicate. Plates were incubated for 2 hours at 4° C on a plate shaker and then washed three times with 1000 μ l/well in a BioTek plate washer with PBST (0.05%Tween-20). Fifty microliters of detection antibody diluted in 2% BSA, PBST was added and the plates were incubated for 1 hour at room temperature. Plates were washed three times with 1000 μ l/well in the BioTek plate washer with PBST (0.05%Tween-20). Fifty 35 microliters of Streptavidin-HRP was added and the plates were incubated for 30 miutes at room temperature. Plates were washed again three times with 1000 μ l/well in a BioTek plate washer with PBST (0.05%Tween-20). Fifty microliters of Supersignal Pico ELISA substrate was added and readout was performed on a Fusion plate reader.

Data was analyzed using EXCEL. Duplicate samples were averaged and the error bars represent the standard deviation between the two replicates.

The results of this experiment are shown in Figure 8. As shown in Figure 8, Ab #6 downregulated ErbB3 in ADRr xenografts *in vivo*.

5

Example 6: Inhibition of Tumor Cell Proliferation

The ability of Ab #6 to inhibit cellular proliferation of cells expressing ErbB3 (e.g., cancer cells) was examined as follows.

MALME3M, ACHN and NCI/ADRr cells were seeded in 96 well tissue culture 10 plates and grown in RPMI-1640 media supplemented with antibiotics, 2mM L-glutamine and 10% fetal bovine serum (FBS) for 24 hours at 37 degrees Celsius and 5% carbon dioxide. Media was then switched to RPMI-1640 media with antibiotics, 2mM L-glutamine and with and without the antibody at 1uM, 250nM, 63nM, 16nM, 4.0nM, 1.0nM, 240pM, 61pM and 15pM concentrations. Cells were grown for 96 hours at 37°C 15 and 5% carbon dioxide, then harvested with CellTiter-Glo® Luminescent Cell Viability Assay (Promega, G7573) and analyzed on a luminometer. Media containing no serum and antibody was used as control.

As shown in Figures 9, 10 and 11, Ab #6 inhibited proliferation of MALME-3M 20 cells (Figure 9), ADRr ovarian cancer cells (Figure 10) and ACHN cells (Figure 11) which express ErbB3. Specifically, Ab # 6 inhibited proliferation of MALME-3M cells by about 19.6%, as measured using the Cell Titer Glow assay, and inhibited proliferation of ADRr ovarian cancer cells by about 30.5%. Also, as shown in Figure 11, Ab # 6 inhibited proliferation of ACHN cells by about 25.4%.

25 **Example 7: Inhibition of ErbB3 Phosphorylation in Tumor Cells**

The ability of Ab #6 to inhibit ErbB3 phosphorylation *in vivo* was examined as follows.

The samples were pulverized using the technique described in Example 5 *supra*, 30 with respect to Figure 8. The BCA assay was performed as set forth in the Materials and Methods section *supra*, and the ELISA assay was performed as described in Example 5 *supra* with respect to Figure 8.

The results of this experiment are shown in Figure 12. As shown in Figure 12, 35 Ab #6 significantly inhibited ErbB3 phosphorylation in ADRr ovarian xenografts *in vivo*, as measured by the amount of phosphorylated ErbB3 (pErbB3) in ng/mg of total protein.

The ability of Ab #6 to inhibit betacellulin (BTC) or heregulin (HRG) induced ErbB3 phosphorylation was also examined, as follows.

Ovarian ADRr cells were preincubated with Ab #6 for 30 minutes prior to stimulation with 50 mM BTC, 10 mM HRG or 333 nM TGF- α . Following pre incubation, the media was removed and the cells were stimulated for 5 minutes at 37 $^{\circ}$ C, 5% CO₂ with 50nM BTC or 333nM TGF- α (for PE498). HRG controls (5 minutes, 5 nM), 10% serum and 0% serum controls were also used. Cells were washed with 1X cold PBS and lysed in 30 μ l cold lysis buffer (M-PER buffer plus sodium vanadate (NaVO₄, Sigma), 2-glycerophosphate, phenylarsine oxide, BpV and protease inhibitors) by incubating on ice for 30 minutes. Lysates were stored overnight at -80 $^{\circ}$ C.

As shown in Figures 13A-13C, Ab #6 significantly inhibited both betacellulin and heregulin-mediated phosphorylation of ErbB3.

In a further experiment, the ability of Ab #6 to inhibit ErbB3 phosphorylation in ovarian tumor cell lines OVCAR 5 and OVCAR 8 was examined as follows.

The OVCAR 5 and OVCAR 8 cell lines were obtained from the National Cancer Institute, Division of Cancer Treatment and Diagnostics ("DCTD"). The ELISA was performed as described in the Materials and Methods section *supra*.

The results of this experiment are depicted in Figures 14A and 14B. As depicted in Figures 14A and 14B, Ab #6 inhibited ErbB3 phosphorylation in both OVCAR 5 and OVCAR 8 ovarian cancer cell lines.

As discussed above, Ab #6 inhibits betacellulin-mediated phosphorylation of ErbB3. In order to investigate whether betacellulin-mediated phosphorylation of ErbB3 occurs through ErbB1 or ErbB3, the following experiment was performed.

ADRr cells or MALME-3M cells (1×10^5) were pre-incubated with 25 μ M of anti-ErbB3 Ab # 6 or 25 μ M of Erbitux (as control) in 50 μ l BD stain buffer for 30 minutes on ice. After 30 minutes, 50 μ l of 400 nM biotinylated BTC was added to the cells and incubated for another 30 minutes on ice. This gave a final concentration of 12.5 μ M antibodies and 200 nM BTC. Cells were then washed twice with 500 μ l BD stain buffer and incubated with 100 μ l of a 1:200 dilution of streptavidin-PE (PE=phycoerythrin) (Invitrogen) in BD stain buffer for 45 minutes. Finally, cells were washed twice, resuspended in 300 μ l of BD stain buffer and analyzed in a FACScalibur flow cytometer. As a positive control, 1×10^5 ADRr or MALME-3M cells were incubated with 200 nM BTC for 30 minutes on ice, washed twice and incubated with a 1:200 dilution of streptavidin-PE for 45 minutes. To assess background staining from the streptavidin-PE conjugate, cells were incubated with 100 μ l of a 1:200 dilution of streptavidin-PE only for 45 minutes.

The results of this experiment are depicted in Figures 15A-15C. As shown in Figure 15A, betacellulin (BTC) does not show any appreciable binding to ErbB1 negative MALME-3M cells. However, as depicted in Figures 15B and 15C, BTC does show binding to ErbB1 positive ADRr cells.

Also, as shown in Figures 15B and 15C, this binding was blocked by Erbitux, which is an anti-EGFR antibody which specifically binds EGFR and was included as a control to demonstrate that EGF-like ligands bind to EGFR, and which is described in *e.g.*, Adams *et al.* (2005), *Nature Biotechnology* 23, 1147 – 1157.

5

Example 8: Inhibition of Heregulin-Mediated Signaling in Tumor Cells

The ability of Ab # 6 to inhibit heregulin-mediated tumor cell signaling was investigated as follows.

MALME-3M cells were seeded in 96 well tissue culture plates and grown in

10 RPMI-1640 media supplemented with antibiotics, 2mM L-glutamine and 10% fetal bovine serum (FBS) for 24 hours at 37°C and 5% carbon dioxide. Cells were serum starved in RPMI-1640 media with antibiotics and 2mM L-glutamine for 24 hours at 37°C and 5% carbon dioxide. Cells were pre-treated with and without the anti-ErbB3 antibody (IgG2 isotype of Ab #6) at 1μM, 250nM, 63nM, 16nM, 4.0nM, 1.0nM, 240pM

15 and 61pM concentrations for 30 minutes then stimulated with HRG1-beta1-ECD for 10 minutes at 37°C and 5% carbon dioxide. Cells were washed with cold PBS then harvested with mammalian protein extract (MPER) lysis (Pierce, 78505) buffer containing 150mM NaCl 5mM sodium pyrophosphate, 10uM bpV (phen), 50μM phenalarsine, 1mM sodium orthovanadate, and protease inhibitor cocktail (Sigma, P714). Cell lysates were diluted two-fold with 4% bovine serum albumin in phosphate buffered saline with 0.1% tween-20, then analyzed by ELISA for AKT (a downstream effector of ErbB3) and ErbB3 phosphorylation.

20

In order to test for AKT phosphorylation, lysates were run on an ELISA plate with a capture antibody specific for AKT and biotinylated detection antibody specific to

25 the phosphorylation site on serine 473 of AKT. Signal was generated with streptavidin conjugated to horseradish-peroxidase reacted with chemiluminescent substrate (Pierce, 37070). In order to assay for ErbB3 phosphorylation, lysates were run on an ELISA plate with a capture antibody specific for ErbB3 and an anti-phosphotyrosine detection antibody conjugated to horseradish-peroxidase. This was then reacted with

30 chemiluminescent substrate (Pierce, 37070). ELISAs were visualized using a luminometer.

As shown in Figures 16A and 16B, Ab #6 was a potent inhibitor of heregulin-mediated signaling in MALME-3M cells, as measured by decreased phosphorylation of ErbB3 (Figure 16A) and AKT (Figure 16B). Notably, Ab #6 inhibited the

35 phosphorylation of AKT by almost 100%.

Example 9: Inhibition of Ovarian, Prostate, and Pancreatic Tumor Growth

To assess the efficacy of Ab #6 *in vivo*, several xenograft models of human cancer were established in nude mice and the inhibition of tumor growth was assessed at multiple doses. For example, T-cell deficient nu/nu mice (3-4 week old female mice 5 originated at NIH; outbred; albino background) were purchased from Charles River Labs (Wilmington, MA) for xenograft studies. ADRr cells for implantation were grown in culture (RPMI media, 10% FBS, L-glutamine and antibiotics, 37°C, 5% CO2) to about 85% confluence before harvesting. Cells were kept on ice until implantation. Mice were implanted via subcutaneous injection with 100 μ l ADRr cells on the right flank and 10 allowed to recover while being monitored for initial tumor growth.

Tumors were measured (length by width) by digital caliper and the mice were dosed with IgG2a (Sigma, M7769-5MG) by intravenous injection. Mice were dosed intra-peritoneally every third day with either 30 μ g or 300 μ g of Ab #6 and tumors were measured three times per week and recorded in a Microsoft Excel spreadsheet.

15 Final tumor measurements (L x W) were taken, the mice were euthanized by CO2 asphyxiation and tumors were excised, snap frozen in liquid nitrogen, and were stored at -80°C (for biochemical analysis). Final tumor measurements were analyzed and graphed by tumor area and tumor volume, as described in Burtrum *et al.*, *supra*. The data was also analyzed by "normalized" and "non-normalized" means for both tumor volume and 20 tumor area. For the "normalization" of the data, at each time point of measurement, each tumor in each group was divided by the initial tumor size determined by caliper measurement.

The data from three different models derived from human tumor cell lines, ADRr (ovarian), Du145 (prostate) and OvCAR8 (ovarian) are shown in Figures 17A-C and 25 Colo357 xenograft study is shown in Figure 17D. The data from these studies demonstrated that a 300ug dose of Ab #6 every three days (Q3d) results in significant inhibition of tumor growth ($p<0.05$ for multiple time points during the studies). Moreover, this inhibitory effect of Ab #6 was further elevated when the dose was increased to 600ug, Q3d, in the Du145 prostate cancer model as well as a renal and a 30 pancreatic carcinoma xenograft model (ACHN and COLO357). However, further elevating the dose to 1500ug Q3d did not result in increased efficacy (OvCAR8-Figure 17; COLO357) suggesting that the 600ug is saturating in regards to tumor growth inhibition. Pharmacokinetic (PK) analyses of the serum from the animals from these studies demonstrate a dose-dependent increase in the serum retention of Ab #6. 35 Similarly, biochemical analysis of the intra-tumoral levels of Ab #6 from these different studies showed a dose-dependent range of 0 to ~6 pg MM121/ug of total tumor lysate (data not shown).

Example 10: Inhibition of binding of ErbB3 Ligands to ErbB3 on Tumor Cells

5 In a further experiment, the specificity of the antibodies of the invention to inhibit the binding of ErbB3 ligands to ErbB3, and not EGF-like ligands to EGFR, was investigated as follows.

In one experiment, the specificity of Ab #6 and a Fab version of Ab #3 (Ab/Fab #3) to inhibit the binding of ErbB3 ligands (e.g., heregulin and epiregulin) to ErbB3 was investigated.

10 In order to investigate the ability of Ab #6 and Ab/Fab #3 to inhibit the binding of heregulin to ErbB3, the following experiment was performed.

15 ADRr cells (1×10^5) were incubated with 10 μ M of an anti-ErbB3 antibody (e.g., Ab #6 or Ab/Fab #3) in 50 μ l BD stain buffer for 30 minutes on ice. After 30 minutes, 50 μ l of 40 nM biotinylated heregulin EGF was added to the cells and incubated for another 10 minutes on ice. This gave a final concentration of 5 μ M antibody and 20 nM heregulin EGF. Cells were then washed twice with 500 μ l BD stain buffer and incubated with 100 μ l of a 1:200 dilution of streptavidin-PE (PE=phycoerythrin) (Invitrogen) in BD stain buffer for 45 minutes. Finally, cells were washed twice, resuspended in 300 μ l of BD stain buffer and analyzed in a FACScalibur flow cytometer. As a positive control, 1×10^5 ADRr cells were incubated with 20 nM 20 heregulin EGF for 10 minutes on ice, washed twice and incubated with a 1:200 dilution of streptavidin-PE for 45 minutes. In order to assess background staining from the streptavidin-PE conjugate, 1×10^5 ADRr cells were incubated with 100 μ l of a 1:200 dilution of streptavidin-PE only for 45 minutes.

25 The results of this experiment are shown in Figures 18A and 18B. As depicted in Figures 18A and 18B, both Ab #6 and Ab/Fab #3 were able to inhibit heregulin binding to ErbB3.

Similarly, the ability of Ab #6 to inhibit the binding of another ErbB3-ligand, epiregulin, to ErbB3, was examined as follows.

30 ADRr cells (1×10^5) were pre-incubated with 25 μ M of Ab #6 or 25 μ M of Erbitux (as control) in 50 μ l BD stain buffer for 30 minutes on ice. After 30 minutes, 50 μ l of 2 μ M biotinylated Epi was added to the cells and incubated for another 30 minutes on ice. This gave a final concentration of 12.5 μ M antibodies and 1 μ M Epi. Cells were then washed twice with 500 μ l BD stain buffer and incubated with 100 μ l of a 1:200 dilution of streptavidin-PE (PE=phycoerythrin) (Invitrogen) in BD stain buffer for 45 minutes. Finally, cells were washed twice, resuspended in 300 μ l of BD stain buffer and analyzed in a FACScalibur flow cytometer. As a positive control, 1×10^5 ADRr cells were incubated with 1 μ M Epi for 30 minutes on ice, washed twice and incubated with a 1:200 dilution of streptavidin-PE for 45 minutes. To assess background staining from

the streptavidin-PE conjugate, cells were incubated with 100 μ l of a 1:200 dilution of streptavidin-PE only for 45 minutes.

The results of this experiment are depicted in Figures 19A and 19B. As shown in Figure 19A, epiregulin binds to ErbB3 positive ADRr cells. Further, as shown in 5 Figure 19B, this binding is inhibited by both Erbitux and Ab #6, suggesting that epiregulin may bind to both EGFR and ErbB3.

A further experiment was performed to investigate whether Ab #6 is able to inhibit the binding of an EGF-like ligand (*e.g.*, HB-EGF) to tumor cells.

ADRr cells (1×10^5) were pre-incubated with 25 μ M of Ab #6 or 25 μ M of 10 Erbitux (as control) in 50 μ l BD stain buffer for 30 minutes on ice. After 30 minutes, 50 μ l of 400 nM biotinylated HB-EGF was added to the cells and incubated for another 30 minutes on ice. This gave a final concentration of 12.5 μ M antibodies and 200 nM HB-EGF. Cells were then washed twice with 500 μ l BD stain buffer and incubated with 100 μ l of a 1:200 dilution of streptavidin-PE (PE=phycoerythrin) (Invitrogen) in BD stain 15 buffer for 45 minutes. Finally, cells were washed twice, resuspended in 300 μ l of BD stain buffer and analyzed in a FACScalibur flow cytometer. As a positive control, 1×10^5 ADRr cells were incubated with 200 nM HB-EGF for 30 minutes on ice, washed twice and incubated with a 1:200 dilution of streptavidin-PE for 45 minutes. To assess background staining from the streptavidin-PE conjugate, cells were incubated with 100 20 μ l of a 1:200 dilution of streptavidin-PE only for 45 minutes.

As shown in Figures 20A and 20B, HB-EGF binds to ErbB on ADRr cells and Ab #6 does not inhibit this binding, evidencing that Ab #6 is specific for inhibiting the binding of ErbB3 ligands (*e.g.*, heregulin and epiregulin) to ErbB3.

25 **Example 11: Inhibition of VEGF Secretion in Tumor Cells**

The ability of Ab #6 to inhibit VEGF secretion of cells expressing ErbB3 (*e.g.*, cancer cells) was examined using VEGF secretion assay (VEGF ELISA kit available from R&D Systems, Minneapolis, MN, Cat.#DY293B). First, the ability of Ab #6 to inhibit VEGF secretion in the untreated and HRG -beta1 treated MCF-7, T47D, and 30 COLO-357 cells was analyzed. These studies revealed that COLO-357 secreted the highest amount of VEGF into the media. As these cells also had very high HRG levels (data not shown), addition of HRG to the media was not able to further induce VEGF secretion (Fig. 24A). In contrast, HRG was able to induce VEGF secretion in MCF-7 and T47D cells.

35 Ab #6 shows a potent inhibitory effect at high levels in all three cell lines with the highest being in COLO-357 (Fig. 24A). Ab #6 also shows a similar effect in vivo by inhibiting VEGF secretion in three different xenografts, the highest being in COLO-357 xenograft (Fig. 24B). Inhibition of VEGF correlates with inhibition of ErbB3

phosphorylation (Fig. 24 C). Inhibition of VEGF secretion also correlates with inhibition of angiogenesis of the tumor cells. In particular, it has been identified that myeloma cell-secreted factors, such as VEGF and bFGF, trigger angiogenesis (see, e.g., Leung et al. (1989) *Science* 246(4935):1306-9; Yen et al. (2000) *Oncogene* 19(31):3460-9).

Example 12: Inhibition of Cell Migration

The ability of Ab #6 to inhibit the migration of cells expressing ErbB3 (e.g., MCF-7 cells) was examined using a trans-well assay (Millipore Corp., Billerica, MA, Cat # ECM552). First, MCF-7 cells were serum-starved overnight and then incubated in the presence or absence of Ab #6 (8uM final concentration) for 15 minutes at room temperature. The cells were then transferred to an upper chamber that is separated from a lower chamber by a collagen type I-coated membrane through which the cells can migrate. 10%FBS was added to media in the lower chamber to act as a chemoattractant in the presence of absence of Ab #6. The chambers were incubated at 37°C for 16 hours and then the cells that migrated through the membrane were removed using a detachment buffer and incubated with a cell-binding fluorescent dye. Fluorescence was quantitated using a fluorescent plate reader. The average fluorescence ± SEM (n=2) is shown in Figure 25.

As shown in Figure 25, 10% FBS stimulates cell migration (lane 3) as compared to untreated control (lane 1) and 8uM Ab #6 inhibits the FBS induced cell migration (lane 4).

Example 13: Inhibition of Spheroid Growth

The ability of Ab #6 to inhibit the spheroid growth of cells expressing ErbB3 was examined using an assay which approximates conditions of a developing tumor growth (Herman et al. (2007) *Journal of Biomolecular Screening Electronic publication*). AdrR and DU145 spheroids were initiated at a frequency of 1 spheroid per well of a 96 well plate using the hanging drop method (Herrman et al., 2008). Individual spheroids were then treated with either Ab #6 (8 uM final concentration), Heregulin- β 1 EGF domain (R&D Systems, Minneapolis, MN, Cat # 396-HB, 3.4 nM final concentration), or a combination of both, as indicated. The diameters of the spheroids were measured using light microscopy (10X objective) at day 1 and day 13.

Ab #6 inhibits spheroid growth in AdrR cells (Fig. 26A). In addition, 3.4nM HRG stimulates spheroid growth and Ab #6 inhibits the HRG effect (Fig. 26B). Spheroids derived from DU145 did not increase in size during 13 days of the

experiment; however, growth was significantly stimulated by HRG1-beta 1. In these cells, 8uM Ab #6 inhibits HRG induced spheroid growth (Fig. 26C).

Example 14: Inhibition of Signaling

5 The ability of Ab #6 to inhibit the signaling induced by different ligands was examined. For example, the effect of Ab #6 on HRG and BTC binding to AdrR cells expressing ErbB3 receptor was tested. As shown in Figures 27A and B, using FACS analysis, Ab #6 competes with HRG and not BTC for binding to AdrR cells. Accordingly, blocking by Ab #6 of HRG binding to ErbB3 would prevent signaling 10 induced by HRG.

15 Additionally, various ligands were tested for inducement of ErbB3 phosphorylation. Three ligands, HRG, BTC, and HGF, were able to stimulate ErbB3 induced phosphorylation in AdrR cells, while EGF could not. As shown in Figure 28, Ab #6 inhibits HGF induced pErbB3 phosphorylation in AdrR cells (Fig. 28). Further, 15 as known in the art (see, e.g., Wallenius et al. (2000) Am J Pathol. 156 (3):821-9 10702398), enhanced HGF signaling has been found in various epithelial and non- epithelial tumors.

ErbB3/cMET interaction and the role of Ab #6 in modulating this interaction

20 It has been shown that non-small-cell lung cancers carrying activating mutations in the epidermal growth factor receptor (EGFR) develop resistance to tyrosine kinase inhibitors by recruiting MET and HER3 and thus activating the PI3K-AKT cell survival pathway (Engelmann et al. (2007) Science 316: 1039-1043; Gou (2007) PNAS: 105(2): 692-697). The association between EGFR and c-MET in cell lines that carry activating 25 EGFR mutations has been well established by co-immunoprecipitation (Engelmann et al. 2007; Gou 2007). Gou et al. recently demonstrated that c-MET and ErbB3 also exist in a complex in a gastric cell line MKN45 known to be dependent on amplified c-MET, using co-immunoprecipitation.

30 This c-MET-erbB3 interaction occurs also in AdrR cells carrying the wild type EGFR and is not dependent on amplified c-MET. HGF (Hepatocyte Growth Factor) induces ErbB3 phosphorylation in AdrR cells in a dose dependent manner as shown in Figure 28. In addition, Ab #6 inhibits HGF induced erbB3 phosphorylation.

35 The effect of HRG and BTC on both ErbB1 and ErbB3 phosphorylation has also been investigated, and HRG and BTC were found to induce phosphorylation of both ErbB1 and ErbB3. HRG was found to be a more potent inducer of ErbB3 phosphorylation while BTC was a potent inducer of ErbB1 phosphorylation (Fig. 29).

This phosphorylation is likely to be driven by the complex between ErbB1 and ErbB3. Briefly, HRG binding to ErbB3 induces complex formation between ErbB1 and ErbB3, leading to the activation of both receptors. The same phenomenon appears likely for BTC, where BTC binding to ErbB1 stimulates complex formation between ErbB1 and 5 ErbB3, leading to the phosphorylation of both ErbB1 and ErbB3.

Antibody Inhibition of ligand (HRG, BTC, EGF, and HGF) stimulated ErbB3 phosphorylation.

The ability of Ab #6 to inhibit ligand (HRG, BTC, EGF, and HGF) induced 10 ErbB3 phosphorylation was examined based on the following method:

1. AdrR cells were plated into 96 well plate at a density of 30,000 cells/well/100 uL in RPMI medium containing 10% FBS and allowed to grow overnight;
2. The next day, cells were serum-starved by changing medium to FBS-free medium and allowed to grow overnight;
- 15 3. Cells were pre-treated with different concentrations of Ab #6 (from 0.01 nM to 100nM), or buffer (control), for 2hours;
4. The cells were then stimulated with 10 nM HRG and HGF for 10 minutes, or 10 nM BTC and EGF for 5 minutes;
5. The reaction was stopped by removing the culture medium and washing the cells 20 once with ice cold PBS;
6. The cells were then lysed in 25 mM Tris, pH+7.5, 150 mM NaCl, 1mM EDTA, 1.0% Triton X-100, 1.0% CHAPS, 10% v/v glycerol, containing 1X protease inhibitor and 1X phosphatase inhibitor; and
7. ErbB3 phosphorylation was measured in cell lysates using Human Phospho- 25 ErbB3 ELISA kit (R&D Systems, Minneapolis, MN, Cat. No. DYC1769) according to manufacturer's instructions.

Antibody Inhibition of ErbB2-ErbB3 protein complex formation.

AdrR cells were pre-incubated with buffer (control), or 250 nM Ab 30 #6 for 60 minutes at room temperature, then treated with 10 nM HRG or 10 nM BTC or control buffer for 10 minutes. The cells were lysed in 25 mM Tris, pH+7.5, 150 mM NaCl, 1mM EDTA, 1.0% Triton X-100, 1.0% CHAPS, 10% v/v glycerol, containing 0.2mM PMSF, 50mTU/mL aprotinin, and 100 uM leupeptin, and the crude lysate were centrifuged briefly to 35 remove insoluble material. Supernatant was transferred to a new eppendorf tube, and anti-ErbB3 antibody (Santa Cruz sc-285) was added at 1:500 dilution. Supernatants were incubated overnight with gentle shaking at 4C.

60 ul of Immobilized Protein A/G agarose beads (Pierce, Rockford, IL, Cat# 20421) was first washed with 1X PBS. The cell lysate-antibody mixture was added to the PBS washed beads, and incubated for 2 hours with gentle shaking at 4°C. The immunoprecipitates were then washed with ice-cold

5 lysis buffer 3 times, resuspended in 30 ul of 2X SDS sample buffer, heat denatured at 95°C for 7 minutes and run on 4-12% Bis-Tris Gels. SDS-PAGE and electro-transferred to PVDF membrane in Tri-Glycine buffer with 10% MeOH. The membrane was blocked for 1hour in 10ml of blocking buffer (Li-Cor Biosciences, Lincoln, NE, Cat# 927-40000) and

10 then incubated with the anti-ErbB2 antibody at 1:1000 (Cell Signaling Technology, Danvers, MA, Cat # 29D8) in 10 ml of blocking buffer (Li-Cor Biosciences, Cat# 927-40000). The signal was detected using goat anti-rabbit IRDye800 at 1:5000 (2ul) in 10ml of blocking buffer (Li-Cor Biosciences, Cat# 927-40000).

15 Ab #6 was also shown to completely inhibit HRG stimulated ErbB2/3 complex formation (Fig. 29B).

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than

20 routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims. Any combination of the embodiments disclosed in the dependent claims are contemplated to be within the scope of the invention.

Incorporation by Reference

All publications, patents, and pending patent applications referred to herein are hereby incorporated by reference in their entirety.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. An isolated monoclonal antibody, or antigen binding portion thereof, comprising: a heavy chain variable region CDR1 comprising SEQ ID NO:7; a heavy chain variable region CDR2 comprising SEQ ID NO:8; a heavy chain variable region CDR3 comprising SEQ ID NO:9; a light chain variable region CDR1 comprising SEQ ID NO:10; a light chain variable region CDR2 comprising SEQ ID NO:11; and a light chain variable region CDR3 comprising SEQ ID NO:12.
- 10 2. An isolated monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 comprising:
 - (a) a heavy chain variable region CDR1 comprising SEQ ID NO:7 or an amino acid sequence which is at least 90% identical thereto;
 - (b) a heavy chain variable region CDR2 comprising SEQ ID NO:8 or an amino acid sequence which is at least 90% identical thereto;
 - 15 (c) a heavy chain variable region CDR3 comprising SEQ ID NO:9 or an amino acid sequence which is at least 90% identical thereto;
 - (d) a light chain variable region CDR1 comprising SEQ ID NO:10 or an amino acid sequence which is at least 90% identical thereto;
 - (e) a light chain variable region CDR2 comprising SEQ ID NO:11 or an amino acid sequence which is at least 90% identical thereto; and
 - (f) a light chain variable region CDR3 comprising SEQ ID NO:12 or an amino acid sequence which is at least 90% identical thereto.
- 25 3. An isolated monoclonal antibody, or antigen binding portion thereof, comprising:
 - (a) a heavy chain variable region CDR1 comprising SEQ ID NO:7 or conservative sequence modifications thereof;
 - (b) a heavy chain variable region CDR2 comprising SEQ ID NO:8 or conservative sequence modifications thereof;
 - 30 (c) a heavy chain variable region CDR3 comprising SEQ ID NO:9 or conservative sequence modifications thereof;
 - (d) a light chain variable region CDR1 comprising SEQ ID NO:10 or conservative sequence modifications thereof;

- (e) a light chain variable region CDR2 comprising SEQ ID NO:11 or conservative sequence modifications thereof; and
- (f) a light chain variable region CDR3 comprising SEQ ID NO:12 or conservative sequence modifications thereof.

5

4. An isolated monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises heavy and light chain variable regions, wherein the heavy chain variable region comprises an amino acid sequence at least 95% identical to the heavy chain variable region amino acid sequence set forth in SEQ ID NO:1.

10

5. An isolated monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises heavy and light chain variable regions, wherein the heavy chain variable region comprises SEQ ID NO:1.

15 6. An isolated monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises heavy and light chain variable regions, wherein the light chain variable region comprises an amino acid sequence at least 95% identical to the light chain variable region amino acid sequence set forth in SEQ ID NO:2.

20 7. An isolated monoclonal antibody, or antigen binding portion thereof, that binds to ErbB3 and comprises heavy and light chain variable regions, wherein the light chain variable region comprises SEQ ID NO:2.

8. An isolated monoclonal antibody, or antigen binding portion thereof, that binds to
25 ErbB3 and comprises a heavy chain variable region comprising SEQ ID NO:1 or a sequence at least 95% identical thereto and a light chain variable region comprising SEQ ID NO:2 or a sequence at least 95% identical thereto.

9. An isolated monoclonal antibody, or antigen binding portion thereof, comprising a
30 heavy chain variable region comprising SEQ ID NO:1 and a light chain variable region comprising SEQ ID NO:2.

10. An isolated antibody, or antigen binding portion thereof, that binds to an epitope which is same or overlapping with the epitope bound by the antibody of claim 9.
11. An isolated antibody, or antigen binding portion thereof, that competes for binding to 5 ErB3 with the antibody of claim 9.
12. The antibody, or antigen binding portion thereof, of any one of the preceding claims, wherein the antibody is selected from the group consisting of a human antibody, a humanized antibody, a bispecific antibody and a chimeric antibody.
- 10 13. The antibody, or antigen binding portion thereof, of any one of the preceding claims, wherein the antibody or antigen binding portion thereof is selected from the group consisting of a Fab, Fab'2, ScFv and a domain antibody.
- 15 14. The antibody, or antigen binding portion thereof, of any one of the preceding claims, wherein the antibody isotype is selected from the group consisting of an IgG1, an IgG2, an IgG3, an IgG4, an IgM, an IgA1, an IgA2, an IgAsec, an IgD, and an IgE antibody.
- 20 15. The antibody, or antigen binding portion thereof, of claim 14, wherein the antibody is an IgG2 isotype antibody.
16. A composition comprising the antibody, or antigen binding portion thereof, of any one of the preceding claims in a pharmaceutically acceptable carrier.
- 25 17. The composition of claim 16, wherein the composition is a sterile aqueous composition.
18. The composition of claim 16, further comprising a molecule that targets EGFR.
- 30 19. An isolated nucleic acid encoding a heavy chain variable region of an antibody, or antigen binding portion thereof, of claim 5, wherein the nucleic acid comprises SEQ ID NO:25.

20. An isolated nucleic acid encoding a light chain variable region of an antibody, or antigen binding portion thereof, of claim 7, wherein the nucleic acid comprises SEQ ID NO:26.

5

21. An isolated nucleic acid encoding a heavy chain variable region of the antibody, or antigen binding portion thereof, of claim 5 comprising a sequence that hybridizes under high stringency to a nucleic acid comprising the nucleotide sequence SEQ ID NO:25.

10 22. An isolated nucleic acid encoding a light chain variable region of the antibody, or antigen binding portion thereof, of claim 7 comprising a sequence that hybridizes under high stringency to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:26.

23. An expression vector comprising the nucleic acid of any one of claims 19 to 22.

15

24. A host cell comprising the vector of claim 23 or the nucleic acid of any one of claims 19 to 22.

20 25. A transgenic non-human mammal which expresses a monoclonal antibody, or antigen binding portion thereof, that binds the same epitope as the antibody, or antigen binding portion thereof, of any one of claims 1 to 15.

25 26. A transgenic plant which expresses a monoclonal antibody, or antigen binding portion thereof, that binds the same epitope as the antibody, or antigen binding portion thereof, of any one of claims 1 to 15.

27. A host cell which produces an antibody, or antigen binding portion, thereof of any one of claims 1 to 15.

30 28. A kit comprising one or more isolated monoclonal antibodies, or antigen binding portions thereof, of any one of claims 1 to 15, and optionally comprising instructions for use in treating or diagnosing a disease associated with ErbB3 dependent signaling.

29. The kit of claim 28, wherein the disease is a cancer.
30. A method of inhibiting ErbB3 signaling in a subject, comprising administering to the subject an isolated monoclonal antibody, or antigen binding portion thereof, of any 5 one of claims 1 to 15, in an amount sufficient to inhibit ErbB3 signaling.
31. A method of treating cancer in a subject, comprising administering to the subject a therapeutically effective amount of an isolated monoclonal antibody, or antigen binding portion thereof, of any one of claims 1 to 15, in an amount sufficient to treat cancer.
- 10 32. The method of claim 30 or 31, wherein the antibody, or antigen binding portion thereof, comprises a heavy chain variable region CDR1 comprising SEQ ID NO:7, a heavy chain variable region CDR2 comprising SEQ ID NO:8, a heavy chain variable region CDR3 comprising SEQ ID NO:9, a light chain variable region CDR1 comprising 15 SEQ ID NO:10, a light chain variable region CDR2 comprising SEQ ID NO:11, and a light chain variable region CDR3 comprising SEQ ID NO:12.
33. The method of claim 32, wherein the antibody or antigen binding portion thereof comprises a heavy chain variable region comprising SEQ ID NO:1.
- 20 34. The method of claim 32, wherein the antibody or antigen binding portion thereof comprises a light chain variable region comprising SEQ ID NO:2.
35. The method of claim 32, wherein the antibody or antigen binding portion thereof 25 comprises a heavy chain variable region comprising SEQ ID NO:1 and a light chain variable region comprising SEQ ID NO:2.
36. The method of any one of claims 31 to 35, wherein the cancer is selected from the group consisting of melanoma, breast cancer, ovarian cancer, renal carcinoma, 30 gastrointestinal/colon cancer, lung cancer, clear cell sarcoma, and prostate cancer.
37. The method of any one of claims 30 to 36, wherein the subject is human.

38. The method of any one of claims 30 to 37, wherein the antibody or antigen binding portion thereof is administered intravenously, intramuscularly, or subcutaneously to the subject.
- 5 39. The method of any one of claims 31 to 38, wherein the antibody or antigen binding portion thereof is administered in combination with a second therapeutic agent.
- 10 40. The method of claim 39, wherein the second agent is a second antibody, or antigen binding portion thereof.
41. The method of claim 39, wherein the second agent is an anti-cancer agent.
- 15 42. The method of claim 41, wherein the anti-cancer agent is selected from the group consisting of an antibody, a small molecule, an antimetabolite, an alkylating agent, a topoisomerase inhibitor, a microtubule-targeting agent, a kinase inhibitor, a protein synthesis inhibitor, an immunotherapeutic, a hormone or analog thereof, a somatostatin analog, a glucocorticoid, an aromatase inhibitor, and an mTOR inhibitor.
- 20 43. The method of claim 42, wherein the anti-cancer agent is an antibody that is an anti-IGF1R antibody, an anti-EGFR antibody, or an anti-cMET antibody.
44. The method of claim 42, wherein the small molecule targets IGF1R, EGFR, or cMET.
- 25 45. A method of diagnosing a cancer associated with ErbB3 in a subject, comprising (a) contacting *ex vivo* or *in vivo* cells from the subject with an isolated monoclonal antibody, or antigen binding portion thereof, of any of claims 1 to 15, and (b) measuring the level of binding to ErbB3 on the cells, wherein abnormally high levels of binding to ErbB3 indicate that the subject has a cancer associated with ErbB3.
- 30 46. Use of an isolated monoclonal antibody, or antigen binding portion thereof, of any one of claims 1 to 15 for the manufacture of a medicament for inhibiting ErbB3 signaling.

47. Use of an isolated monoclonal antibody, or antigen binding portion thereof, of any one of claims 1 to 15 for the manufacture of a medicament for treating cancer.
- 5 48. An isolated monoclonal antibody, or antigen binding portion thereof, of any one of claims 1 to 11; a composition of claim 16; an isolated nucleic acid of any one of claims 19 to 22; an expression vector of claim 23; a host cell of claim 24 or claim 27; a transgenic non-human mammal of claim 25; a transgenic plant of claim 26; a kit of claim 28; a method of any one of claims 30, 31 or 45; or use of claim 46 or claim 47,
- 10 substantially as herein described with reference to any one or more of the examples but excluding comparative examples.

1/43




Fig. 1A

2/43




Fig. 1B

3/43

Binding of Antibodies to ErbB3 as Measured by SPR

$$K_D = k_d/k_a$$




Fig. 2A






Fig. 2B

4/43

Cell Binding Assay for Antibodies to MALME-3M cells

$$Y = B_{max} * X / K_D + X$$

*Fig. 2C**Fig. 2D*

5/43

AB #6 IgG2 binding is specific to ErbB3.




Fig. 3

6/43

Antibody#6 causes a decrease in total ErbB3 levels in MALME 3M melanoma cells *in vitro* measured by ELISA




Fig. 4

7/43

Downregulation of ErbB3 receptors on MALME-3M (melanoma) cells with Ab #6 IgG1 and IgG2 measured by FACS





Fig. 5A

Fig. 5B

8/43

**Timecourse of Ab #6
mediated ErbB3 downregulation**

*Fig. 6A**Fig. 6B*

9/43






Fig. 6C
Fig. 6D

10/43

Ab #6 causes ErbB3 downregulation in melanoma cells in vivo*Fig. 7*

11/43

Total ErbB3 levels in ADRr xenografts**Antibody Treatment***Fig. 8*

12/43

**Ab #6 inhibits proliferation in MALME 3M
melanoma cells measured by Cell Titer Glow assay**





Fig. 9

13/43

Ab #6 inhibits cell proliferation in ADRr (ovarian) cell line*Fig. 10*

14/43





Fig. 11

15/43

Ab #6 inhibits ErbB3 phosphorylation in ADRr xenografts in vivo**pErbB3 levels in ADRr xenografts****Antibody Treatment***Fig. 12*

16/43

Ab #6 IgG2/BTC

IC50	8.37e-10 M
95% CI	5.32e-10 to 1.32e-9

Ab #6 IC50 in NCI-ADRr cells-0% serum




Fig. 13A

17/43

Ab #6 IgG2/BTC	
IC50	2.488e-009
95% CI	1.051e-009 to 5.889e-009

Ab #6 IgG2 in NCI-ADRr cells-0% serum




Fig. 13B

18/43




Fig. 13C

19/43




Fig. 14A




Fig. 14B

20/43




Fig. 15A






Fig. 15B

21/43

Blocking BTC Binding by Erbitux but not Ab #6*Fig. 15C*

22/43

Ab #6 IgG2 inhibits Heregulin-mediated signaling with similar IC50s in MALME 3M



Ab #6 Inhibition of Tumor Growth




Fig. 17A

Ab #6 Inhibition of Tumor Growth




Fig. 17B

25/43


Ab #6 Inhibition of Tumor Growth

Fig. 17C

26/43

Ab #6 Inhibition of Tumor Growth




Fig. 17D

27/43

Blocking of HRG Binding to MALME3 by Ab #6




Fig. 18A

Blocking of HRG Binding to MALME3 by Ab #3





Fig. 18B

Epiregulin Binding to ADR_r **Blocking of Epiregulin Binding to ADR_r by Ab #6 and Erbitux**

28/43

*Fig. 19A**Fig. 19B*

29/43




Fig. 20B




Fig. 20A

Ab # 6 VH amino acid sequence (SEQ ID NO:1)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSHYVMAWVRQAPGKGLEWVSSISSS
 GGWTLYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCTRGLKMATIFD
 YWGQGTLVTVSS

Ab # 6 VL amino acid sequence (SEQ ID NO:2)

QSALTQPASVSGSPGQSITISCTGTSSDVGSYNVWSWYQQHPGKAPKLIIYEVSQR
 PSGVSNRFSGSKSGNTASLTISGLQTEDEADYYCCSYAGSSIFVIFGGGTKVTL

Ab # 3 VH amino acid sequence (SEQ ID NO:3)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSAYNMRWVRQAPGKGLEWVSVIYPS
 GGATRYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARGYYYYGMDV
 WGQGTLVTVSS

Ab # 3 VL amino acid sequence (SEQ ID NO:4)

QSVLTQPPSASGTPGQRVTISCSGSDSNIGRNYIYWYQQFPGTAPKLLIYRNNQRP
 SGVPDRISGSKSGTSASLAISGLRSEDEAEYHCGTWDDSLSGPVFGGGTKLTVL

Ab # 14 VH amino acid sequence (SEQ ID NO:5)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSAYGMGWVRQAPGKGLEWVSYISPS
 GGHTKYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCAKVLETGLVD
 AFDIWGQGTMVTVSS

Ab # 14 VL amino acid sequence (SEQ ID NO:6)

QYELTQPPSVVYPGQTASITCSGDQLGSKFVSWYQQRPGQSPVLVMYKDKRRP
 SEIPERFSGNSGNTATLTISGTQAIDEADYYCQAWDSSSTYVFGTGTKVTL

Ab # 17 VH amino acid sequence (SEQ ID NO:35)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYGMGWVRQAPGKGLEWVSYISPS
 GGITVYADSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCARLNYYYGLDV
 WGQGTTVTVSS

Ab # 17 VL amino acid sequence (SEQ ID NO:36)

QDIQMTQSPSSLSASVGDRITITCQASQDIGDSLNWYQQKPGKAPRLLIYDASNLE
 TGVPPRFSGSGSGTDFTRSLQPEDIATYFCQQSANAPFTFGPGTKVDIK

Ab # 19 VH amino acid sequence (SEQ ID NO:37)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSRYGMWWVRQAPGKGLEWVSYIGSS
 GGPTYYVDSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCAGGRGTPYYFDS
 WGQGTLVTVSS

Ab # 19 VL amino acid sequence (SEQ ID NO:38)

QYELTQPASVSGSPGQSITISCTGTSSDGRWNIVSWYQQHPGKAPKLMIYDVSNRPSGVSNRF
 SGSKSGNTASLTISGLQAEDEADYYCSSYTSSTWVFGGGTKLTVL

Fig. 21A

Ab # 6 VH CDR1 (SEQ ID NO:7)
HYVMA

Ab # 6 VH CDR2 (SEQ ID NO:8)
SISSSGGWTLYADSVKG

Ab # 6 VH CDR3 (SEQ ID NO:9)
GLKMATIFDY

Ab # 6 VL CDR1 (SEQ ID NO:10)
TGTSSDVGSYNVVS

Ab # 6 VL CDR2 (SEQ ID NO:11)
EVSQRPS

Ab # 6 VL CDR3 (SEQ ID NO:12)
CSYAGSSIFVI

Ab # 3 VH CDR1 (SEQ ID NO:13)
AYNMR

Ab # 3 VH CDR2 (SEQ ID NO:14)
VIYPSGGATRYADSVKG

Ab # 3 VH CDR3 (SEQ ID NO:15)
GYYYYGMDV

Ab # 3 VL CDR1 (SEQ ID NO:16)
SGSDSNIGRNYIY

Ab # 3 VL CDR2 (SEQ ID NO:17)
RNNQRPS

Ab # 3 VL CDR3 (SEQ ID NO:18)
GTWDDSLSGPV

Fig. 21B

32/43

<u>Ab # 14 VH CDR1 (SEQ ID NO:19)</u> AYGMG	<u>Ab # 19 VH CDR1 (SEQ ID NO:45)</u> RYGMW
<u>Ab # 14 VH CDR2 (SEQ ID NO:20)</u> YISPSGGHTKYADSVKG	<u>Ab # 19 VH CDR2 (SEQ ID NO:46)</u> YIGSSGGPTYYVDSVKG
<u>Ab # 14 VH CDR3 (SEQ ID NO:21)</u> VLETGLLVDAFDI	<u>Ab # 19 VH CDR3 (SEQ ID NO:47)</u> GRGTPYYFDS
<u>Ab # 14 VL CDR1 (SEQ ID NO:22)</u> SGDQLGSKFVS	<u>Ab # 19 VL CDR1 (SEQ ID NO:48)</u> TGTSSDIGRWNIVS
<u>Ab # 14 VL CDR2 (SEQ ID NO:23)</u> YKDKRRPS	<u>Ab # 19 VL CDR2 (SEQ ID NO:49)</u> DVSNRPS
<u>Ab # 14 VL CDR3 (SEQ ID NO:24)</u> QAWDSSTYV	<u>Ab # 19 VL CDR3 (SEQ ID NO:50)</u> SSYTSSSTWV
<u>Ab # 17 VH CDR1 (SEQ ID NO:39)</u> WYGMG	
<u>Ab # 17 VH CDR2 (SEQ ID NO:40)</u> YISPSGGITVYADSVKG	
<u>Ab # 17 VH CDR3 (SEQ ID NO:41)</u> LNYYYGLDV	
<u>Ab # 17 VL CDR1 (SEQ ID NO:42)</u> QASQDIGDSLN	
<u>Ab # 17 VL CDR2 (SEQ ID NO:43)</u> DASNLET	
<u>Ab # 17 VL CDR3 (SEQ ID NO:44)</u> QQSANAPFT	

Fig. 21C

Ab # 6 VH Codon Optimized Nucleic Acid Sequence (SEQ ID NO:25)

gaggtgcagctgctggagagcggcggagggctggtccagccaggcggcagcctgaggctgtcctgcgcgcagcggcttcac
cttcagccactacgtgtatggcctgggtcgccaggcccaggcaaggccctggaatgggtgtccagcatcagcagcggcgg
ctggaccctgtacgcccacagcgtgaagggcagggtcaccatcagcaggacaacagaacaacaccctgtacactgcagatgaac
agcctgagggccgaggacaccggcgttactactgcaccaggcctgaagatggccaccatctcgactactgggcccaggc
accctggtgaccgtgagcagc

Ab # 6 VL Codon Optimized Nucleic Acid Sequence (SEQ ID NO:26)

cagtccgcctgaccaggcccagccgcagcgtgagcggcagccaggccagagcatcaccatcagctgcaccggcaccaggcagcga
cgtggcagctacaacgtgggtcctggtatcagcagcacccggcaaggccccaagctgtatctacgagggtgtcccagagg
cccagcggcgtgagcaacaggttcagcggcagcaagagcggcaacaccggcagcctgaccatcagcggcgtcagaccgagg
acgaggcccactactgctcagctacgcccagcagcatctcgtatctcggcggagggaccaaggtgaccgtccta

Ab # 3 VH Codon Optimized Nucleic Acid Sequence (SEQ ID NO:27)

gaggtgcagctgctggaaagcggcggagggctggtgcagccaggcggcagcctgaggctgtcctgcgcgcagcggcttcac
cttcagccctacaacatgagatgggtcgccaggcccaggcaaggccctggaatgggtgtccgtatctacccagcggcgg
agccaccagatacgcgcacagcgtgaagggcagggttaccatcagcaggacaacagaacaacaccctgtacactgcagatgaa
cagcctgagggccgaggacaccggcgttactactgcgccaggcctactactactacggcatggacgtgtgggcccaggc
cctggtaccgtgaggcagc

Ab # 3 VL Codon Optimized Nucleic Acid Sequence (SEQ ID NO:28)

cagagcgtgctgaccaggcccccaagcgcgcagcggcaccaggccaggccaggggtgaccatcagctgcagcggcagcagca
acatcggcaggaactacatctactggtatcagcagttccccggcaccggcccaagctgtatctacaggaacaaccaggaggccc
agcggcgtgcccacaggatcagcggcagcaagagcggcaccagcggcaccgtggccatcagcggcctgagaagcggag
gaggccgagtaccactgcggcaccctggacacagcctgagcggcccagtgttcggcggagggaccaagctgaccgtccta

Fig. 22A

Ab # 14 VH Nucleic Acid Sequence (SEQ ID NO:29)

gaagttcaatttttagagtctggcggttgcagcctgggttcttacgtcttgcgcgtctccggattcacttctct
gcttacggatgggttgggtcgcctggtaaagggttgagtggttcttatatctcttcgttgcgcataactaag
tatgtgactccgttaaagggtcgcttactatctctagagacaactctaagaactctacttgcaagatgaacagcttaaggc
tgaggacacggccgtatattactgtcgaaactggcttatgggtatgtcgatgcatttgcatactggggccaaggac
atggtcaccgtctcaagc

Ab # 14 VL Nucleic Acid Sequence (SEQ ID NO:30)

cagtacgaatttgactcagccaccctcagtgccgtgtacccaggacagacagccagcatcacctgcctggagatcaattggg
gagtaaattgtttcctggatcagcagagggcaggccagtcctgtgtggcatgtataaagataaaaggccggccgtcaga
gatccctgagcgattctctggctccaactctggaaacacagccactctgaccatcagcgggaccaggctatagatgaggct
gactattattgtcaggcgtggacagcagcacttatgtctcggactgggaccaaggtcaccgtccta

Ab # 6 VH Pre-Optimization Nucleic Acid Sequence (SEQ ID NO:31)

gaagttcaatttttagagtctggcggttgcagcctgggttcttacgtcttgcgcgtctccggattcacttctct
cattacgttatgggttgggtcgcctggtaaagggttgagtggttcttatctcttcgttgcgttgcactctt
gctgactccgttaaagggtcgcttactatctctagagacaactctaagaactctacttgcaagatgaacagcttaaggc
aggacacagccgttattactgtactagaggtctcaagatggctacaattttgactactggggcagggcaccctggtcacc
tctcaagc

Ab # 6 VL Pre-Optimization Nucleic Acid Sequence (SEQ ID NO:32)

cagagcgcttgactcagccgcctccgtctgggtctccgttgcacagtcgtcaccatctctgcactggaaaccaggcagtgt
gttgggaggatataatgttgcctctggtaacacaacacccaggcaagcccccaactcatcattatgaggtcagtgcgg
ccctcagggggttctaatcgctctggctccaagtcgttgcacacgcgcctccgtacaatctctggcactgaggac
gaggctgattattactgtctcatatgcaggtagtagtatttcgtatattcggcggagggaccaaggtgaccgtccta

Ab # 3 VH Pre-Optimization Nucleic Acid Sequence (SEQ ID NO:33)

gaagttcaatttttagagtctggcggttgcagcctgggttcttacgtcttgcgcgtctccggattcacttctct
gcttacaatatcggttgggttgcgcctggtaaagggttgagtggttcttgcgttatctatcttcgttgcgtactcg
atgtgactccgttaaagggtcgcttactatctctagagacaactctaagaactctacttgcaagatgaacagcttaaggc
aggacacggccgttattactgtcgagagggtactactacggatggacgtctggggccaaggcaccctggtcacc
tctcaagc

Ab # 3 VL Pre-Optimization Nucleic Acid Sequence (SEQ ID NO:34)

cagagcgcttgactcagccaccctcagcgttggaccccccggcagagggtcaccatctctgttgcgttgc
acatcggaagaaattatataattggtaccagcaattccaggaaacggcccccaagtcctcatataggaataatcagc
cctcaggggccctgaccgaatctctggctccaagtcgttgcacccgcctccgttgcgttgc
gaggctgaggatcatgtggacatggatgcagcctgagtggtccggattcggcggagggactaagctgaccgtccta

Fig. 22B

- **Ab #6 VLb:**
- QSALTQPASVSGSPGQSITISCTGTSSDVGSYNVVSWYQQHPGKAPK
LMIYEVSKRPSGVSNRFSGSKSGNTASLTISGLQAEDeadYYCCSYA
GSSIFVIFGGGTVT (SEQ ID NO:51)
- **Ab #17 VK1b:**
- QDIQMTQSPSSLSASVGDRITITCQASQDIDGDSLNWYQQKPGKAPRLL
IYDASNLETGVPPRFSGSGSGTDFTRSLQPEDIATYFCQQSANAP
FTFGPGTKVDIR (SEQ ID NO:52)
- **Ab #19 VL2b:**
- QYELTQPASVSGSPGQSITISCTGTSSDGRWNIVSWYQQHPGKAPK
LMIYDVSNRPSGVSNRFSGSKSGNTASLTISGLQAEDeadYYCSSYTS
SSTVFGGGTKLTV (SEQ ID NO:53)

Fig. 23

36/43




Fig. 24A

37/43




Fig. 24B

38/43

Correlation of phosphoErbB3 inhibition and inhibition of VEGF Secretion




Fig. 24C

39/43

Effect of Ab #6 on Cell Migration









Fig. 25

40/43

*Fig. 26A**Fig. 26B**Fig. 26C*

41/43

Effect of Ab #6 on HRG binding to AdrR cells*Fig. 27A***Effect of Ab #6 on BTC binding to AdrR cells***Fig. 27B*

42/43




Fig. 28




Fig. 29A




Fig. 29B

43/43

Fig. 30

