



US010517957B2

(12) **United States Patent**
Low et al.

(10) **Patent No.:** **US 10,517,957 B2**
(45) **Date of Patent:** **Dec. 31, 2019**

(54) **PSMA BINDING LIGAND-LINKER
CONJUGATES AND METHODS FOR USING**

(71) Applicant: **Purdue Research Foundation**, West
Lafayette, IN (US)

(72) Inventors: **Philip S. Low**, West Lafayette, IN
(US); **Sumith A. Kularatne**, West
Lafayette, IN (US)

(73) Assignee: **Purdue Research Foundation**, West
Lafayette, IN (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/997,451**

(22) Filed: **Jun. 4, 2018**

(65) **Prior Publication Data**

US 2018/0303950 A1 Oct. 25, 2018

Related U.S. Application Data

(63) Continuation of application No. 14/794,482, filed on
Jul. 8, 2015, now Pat. No. 10,046,054, which is a
continuation of application No. 12/673,931, filed as
application No. PCT/US2008/073375 on Aug. 15,
2008, now Pat. No. 9,193,763.

(60) Provisional application No. 60/956,489, filed on Aug.
17, 2007, provisional application No. 61/074,358,
filed on Jun. 20, 2008.

(51) **Int. Cl.**

A61K 47/64 (2017.01)
A61K 47/54 (2017.01)
A61K 51/04 (2006.01)
C07K 5/08 (2006.01)
C07C 323/52 (2006.01)
C07D 207/416 (2006.01)
A61K 49/00 (2006.01)
A61K 31/426 (2006.01)
A61K 31/4745 (2006.01)
A61K 31/475 (2006.01)
A61K 51/08 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 47/64** (2017.08); **A61K 31/426**
(2013.01); **A61K 31/475** (2013.01); **A61K**
31/4745 (2013.01); **A61K 47/54** (2017.08);
A61K 47/542 (2017.08); **A61K 47/547**
(2017.08); **A61K 47/548** (2017.08); **A61K**
49/0041 (2013.01); **A61K 49/0043** (2013.01);
A61K 49/0052 (2013.01); **A61K 49/0056**
(2013.01); **A61K 51/04** (2013.01); **A61K**
51/0402 (2013.01); **A61K 51/0489** (2013.01);
A61K 51/088 (2013.01); **C07C 323/52**
(2013.01); **C07D 207/416** (2013.01); **C07K**
5/08 (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,691,024 A 9/1987 Shirahata et al.
4,713,249 A 12/1987 Schroder
5,103,018 A 4/1992 Motomichi et al.
5,266,333 A 11/1993 Cady et al.
5,418,982 A 5/1995 Kishi
5,627,165 A 5/1997 Glazier
5,795,877 A 8/1998 Jackson et al.
5,863,536 A 1/1999 Jackson et al.
5,866,679 A 2/1999 Defeo-jones et al.
5,902,817 A 5/1999 Jackson et al.
5,948,750 A 9/1999 Garsky et al.
5,962,237 A 10/1999 Ts'o et al.
5,962,521 A 10/1999 Jackson et al.
5,968,915 A 10/1999 Jackson et al.
5,998,362 A 12/1999 Feng et al.
6,054,444 A 4/2000 Jackson et al.
6,127,333 A 10/2000 Brady et al.
6,174,858 B1 1/2001 Brady et al.
6,177,404 B1 1/2001 Defeo-jones et al.
6,232,287 B1 5/2001 Ruoslahti et al.
6,368,598 B1 4/2002 D Amico et al.
6,391,305 B1 5/2002 Feng et al.
6,428,785 B1 8/2002 Gokcen
6,479,470 B1 11/2002 Kozikowski et al.
6,528,499 B1 3/2003 Kozikowski et al.
6,692,724 B1 2/2004 Yang et al.
6,875,886 B2 4/2005 Frangioni

(Continued)

FOREIGN PATENT DOCUMENTS

AU 2008289108 A1 2/2009
AU 2008289108 B2 9/2014

(Continued)

OTHER PUBLICATIONS

Jayaprakash et al., "Design and Synthesis of a PSMA Inhibitor-Doxorubicin Conjugate for Targeted Prostate Cancer Therapy", *ChemMedChem*, 2006, vol. 1, pp. 299-302 (Year: 2006).*
"U.S. Appl. No. 14/794,482, Examiner Interview Summary dated Oct. 31, 2017", 3 pgs.
"U.S. Appl. No. 14/794,482, Non Final Office Action dated Aug. 10, 2017", 17 pgs.
"U.S. Appl. No. 14/794,482, Notice of Allowance dated Apr. 6, 2018", 8 pgs.
"U.S. Appl. No. 14/794,482, Notice of Non-Compliant Amendment dated May 22, 2017", 2 pgs.
"U.S. Appl. No. 14/794,482, Preliminary Amendment filed Jul. 8, 2015", 6 pgs.

(Continued)

Primary Examiner — Savitha M Rao
Assistant Examiner — Gregg Polansky
(74) *Attorney, Agent, or Firm* — Schwegman Lundberg & Woessner, P.A.

(57) **ABSTRACT**

Described herein are prostate specific membrane antigen (PSMA) binding conjugates that are useful for delivering therapeutic, diagnostic and imaging agents. Also described herein are pharmaceutical composition containing them and methods of using the conjugates and compositions. Also described are processes for manufacture of the conjugates and the compositions containing them.

(56)

References Cited

U.S. PATENT DOCUMENTS

6,946,133	B1	9/2005	Schlom et al.	8,859,509	B2	10/2014	Spiegel et al.
7,008,765	B1	3/2006	Bussemakers et al.	8,865,126	B2	10/2014	Leamon et al.
7,128,893	B2	10/2006	Leamon et al.	8,877,970	B2	11/2014	Zimmerman et al.
7,129,254	B2	10/2006	Berger et al.	8,907,058	B2	12/2014	Low et al.
7,147,837	B2	12/2006	Lauffer et al.	8,916,161	B2	12/2014	Buckley
7,192,586	B2	3/2007	Bander	8,916,167	B2	12/2014	Low et al.
7,232,805	B2	6/2007	Weinshenker et al.	8,926,944	B2	1/2015	Babich et al.
7,361,338	B2	4/2008	Jakovovits et al.	8,926,945	B2	1/2015	Port et al.
7,381,745	B2	6/2008	Kozikowski et al.	8,940,871	B2	1/2015	Wu et al.
7,399,460	B2	7/2008	Wedeking et al.	8,946,388	B2	2/2015	Sahin et al.
7,408,079	B2	8/2008	Pomper et al.	8,962,799	B2	2/2015	Babich et al.
7,485,299	B2	2/2009	Afar et al.	8,987,319	B2	3/2015	Miller
7,514,078	B2	4/2009	Bander et al.	9,044,468	B2	6/2015	Pomper et al.
7,534,580	B2	5/2009	Reeves et al.	9,056,841	B2	6/2015	Pomper et al.
7,601,332	B2	10/2009	Vlahov et al.	9,193,763	B2	11/2015	Low et al.
7,635,682	B2	12/2009	Denmeade et al.	9,226,981	B2	1/2016	Pomper et al.
7,638,122	B2	12/2009	Yu et al.	9,242,012	B2	1/2016	Ma et al.
7,659,395	B2	2/2010	Pajouhesh et al.	9,278,067	B2	3/2016	Boulikas
7,662,795	B2	2/2010	Rodriguez et al.	9,295,727	B2	3/2016	Zale et al.
7,696,185	B2	4/2010	Berkman	9,309,193	B2	4/2016	Babich et al.
7,713,944	B2	5/2010	Kinberger et al.	9,636,413	B2	5/2017	Vlahov et al.
7,740,847	B2	6/2010	Allan et al.	9,687,572	B2	6/2017	Babich et al.
7,767,202	B2	8/2010	Pardoll et al.	9,951,324	B2	4/2018	Low et al.
7,767,803	B2	8/2010	Diener et al.	10,046,054	B2	8/2018	Low et al.
7,794,929	B2	9/2010	Baylin et al.	2001/0031252	A1	10/2001	Low et al.
7,862,798	B2	1/2011	Leamon et al.	2002/0001782	A1	1/2002	Watanabe et al.
7,872,235	B2	1/2011	Rousoo et al.	2002/0055121	A1	5/2002	Vielkind
7,875,586	B2	1/2011	Kovbasnjuk et al.	2002/0103136	A1	8/2002	Feng
7,879,981	B2	2/2011	Obata	2002/0115596	A1	8/2002	Garsky et al.
7,910,594	B2	3/2011	Vlahov et al.	2002/0132983	A1	9/2002	Junghans
RE42,275	E	4/2011	Berkman	2003/0035804	A1	2/2003	D Amico et al.
7,990,533	B2	8/2011	Maier et al.	2003/0086900	A1	5/2003	Low et al.
8,000,773	B2	8/2011	Rousoo et al.	2003/0133927	A1	7/2003	Defeo-jones et al.
8,101,369	B2	1/2012	Nam et al.	2003/0138432	A1	7/2003	Glazier
8,101,713	B2	1/2012	Cuello et al.	2003/0207808	A1	11/2003	Savitzky et al.
8,105,568	B2	1/2012	Vlahov et al.	2003/0215456	A1	11/2003	Yao et al.
8,153,595	B2	4/2012	Chen	2003/0220241	A1	11/2003	Defeo-jones et al.
8,211,401	B2	7/2012	Babich et al.	2003/0232760	A1	12/2003	Garsky et al.
8,211,402	B2	7/2012	Babich et al.	2004/0001846	A1	1/2004	Israeli et al.
8,211,473	B2	7/2012	Troiano et al.	2004/0002478	A1	1/2004	Kozikowski et al.
8,211,635	B2	7/2012	Barton	2004/0018203	A1	1/2004	Pastan et al.
8,227,634	B2	7/2012	Pomper et al.	2004/0029778	A1	2/2004	Isaacs
8,236,330	B2	8/2012	Zale et al.	2004/0033195	A1	2/2004	Leamon et al.
8,246,968	B2	8/2012	Zale et al.	2004/0052727	A1	3/2004	Dalton et al.
8,258,111	B2	9/2012	Shen et al.	2004/0054190	A1	3/2004	Pomper et al.
8,273,363	B2	9/2012	Zale et al.	2004/0058857	A1	3/2004	Yao
8,313,128	B2	11/2012	Belyea et al.	2004/0110723	A1	6/2004	Frangioni
8,313,728	B2	11/2012	Leamon et al.	2004/0146516	A1	7/2004	Roben et al.
8,388,977	B2	3/2013	Low et al.	2004/0213791	A1	10/2004	Bander et al.
8,404,817	B2	3/2013	Sherman et al.	2004/0229845	A1	11/2004	Frangioni
8,414,898	B2	4/2013	Afar et al.	2004/0242582	A1	12/2004	Green et al.
8,445,851	B2	5/2013	Rousoo et al.	2005/0002942	A1	1/2005	Vlahov et al.
8,450,290	B2	5/2013	Worm et al.	2005/0107325	A1	5/2005	Manoharan et al.
8,465,725	B2	6/2013	Babich et al.	2005/0119166	A1	6/2005	Brady et al.
8,487,128	B2	7/2013	Weissbach et al.	2005/0158780	A1	7/2005	Lupold et al.
8,487,129	B2	7/2013	Babich et al.	2005/0234247	A1	10/2005	Klar et al.
8,507,434	B2	8/2013	Popel et al.	2005/0239138	A1	10/2005	Hess et al.
8,557,772	B2	10/2013	Karagiannis et al.	2005/0239739	A1	10/2005	Matulic-adamic et al.
8,562,945	B2	10/2013	Babich et al.	2005/0245486	A1	11/2005	Frangioni
8,603,499	B2	12/2013	Zale et al.	2005/0255042	A1*	11/2005	Lam
8,603,500	B2	12/2013	Zale et al.	2006/0024317	A1	2/2006	Boyd et al.
8,603,501	B2	12/2013	Zale et al.	2006/0052312	A1	3/2006	Erhardt et al.
8,606,349	B2	12/2013	Rousoo et al.	2006/0062793	A1	3/2006	Webb et al.
8,644,910	B2	2/2014	Rousoo et al.	2006/0105975	A1	5/2006	Pendergrast et al.
8,685,891	B2	4/2014	Muraca	2006/0106047	A1	5/2006	Jiang et al.
8,703,918	B2	4/2014	Colombatti et al.	2006/0140871	A1	6/2006	Sillerud
8,709,483	B2	4/2014	Farokhzad et al.	2006/0148718	A1	7/2006	Brady et al.
8,772,226	B2	7/2014	Denmeade et al.	2006/0155021	A1	7/2006	Lenges et al.
8,772,459	B2	7/2014	Ho et al.	2006/0155146	A1	7/2006	Lenges et al.
8,778,305	B2	7/2014	Pomper et al.	2007/0010014	A1	1/2007	Wood et al.
8,802,153	B2	8/2014	Cheng et al.	2007/0020327	A1	1/2007	Fikes et al.
8,816,095	B2	8/2014	Brown et al.	2007/0031326	A1	2/2007	Shirvan et al.
8,834,842	B2	9/2014	Leamon et al.	2007/0031438	A1	2/2007	Junghans
8,840,865	B2	9/2014	Babich et al.	2007/0041901	A1	2/2007	Diener et al.
8,852,630	B2	10/2014	Spiegel et al.	2007/0117153	A1	5/2007	Bieniarz et al.
				2007/0128670	A1	6/2007	Klatzmann et al.
				2007/0134332	A1	6/2007	Turnell et al.
				2007/0142296	A1	6/2007	Mcbride et al.
				2007/0148662	A1	6/2007	Israeli et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2007/0160617 A1 7/2007 Ma et al.
 2007/0172422 A1 7/2007 Glazier
 2007/0179100 A1 8/2007 Manoharan
 2007/0219165 A1 9/2007 Berkman
 2007/0225213 A1 9/2007 Kosak
 2007/0244055 A1 10/2007 Brady et al.
 2007/0254316 A1 11/2007 Rodriguez et al.
 2007/0254317 A1 11/2007 Busseret-michel et al.
 2008/0008649 A1 1/2008 Cappelletti et al.
 2008/0008719 A1 1/2008 Bowdish et al.
 2008/0089869 A1 4/2008 Denmeade et al.
 2008/0114153 A1 5/2008 Steeves et al.
 2008/0175789 A1 7/2008 Frangioni
 2008/0176821 A1 7/2008 Kozikowski et al.
 2008/0193381 A1 8/2008 Babich et al.
 2008/0214436 A1 9/2008 Yu et al.
 2008/0248052 A1 10/2008 Vlahov et al.
 2008/0269105 A1 10/2008 Taft et al.
 2008/0311037 A1 12/2008 Heston et al.
 2009/0117042 A1 5/2009 Pomper et al.
 2009/0123467 A1 5/2009 Bedi et al.
 2009/0180951 A1 7/2009 Zimmerman et al.
 2009/0214636 A1 8/2009 Low et al.
 2009/0247614 A1 10/2009 Manoharan et al.
 2009/0258002 A1 10/2009 Barrett et al.
 2009/0274625 A1 11/2009 Denmeade et al.
 2010/0048490 A1 2/2010 Vlahov et al.
 2010/0055735 A1 3/2010 Low et al.
 2010/0092496 A1 4/2010 Boyd et al.
 2010/0178246 A1 7/2010 Babich et al.
 2010/0183509 A1 7/2010 Babich et al.
 2010/0183517 A1 7/2010 Berkman
 2010/0209343 A1 8/2010 Bander et al.
 2010/0240701 A1 9/2010 Vlahov et al.
 2010/0324008 A1 12/2010 Low et al.
 2011/0008253 A1 1/2011 Babich et al.
 2011/0027180 A1 2/2011 Magnani
 2011/0027274 A1 2/2011 Cheng et al.
 2011/0064657 A1 3/2011 Pomper et al.
 2011/0142760 A1 6/2011 Pomper et al.
 2011/0172254 A1 7/2011 Leamon et al.
 2011/0176998 A1 7/2011 Pomper et al.
 2011/0200677 A1 8/2011 Chandran et al.
 2011/0288152 A1 11/2011 Low et al.
 2012/0009121 A1 1/2012 Pomper et al.
 2012/0276162 A1 11/2012 Zale et al.
 2012/0322741 A1 12/2012 Low et al.
 2013/0034494 A1 2/2013 Babich et al.
 2013/0172406 A1 7/2013 Zale et al.
 2013/0336888 A1 12/2013 Babich et al.
 2014/0073763 A1 3/2014 Low et al.
 2014/0107316 A1 4/2014 Vlahov et al.
 2014/0140925 A1 5/2014 Leamon et al.
 2014/0154702 A1 6/2014 Parker et al.
 2014/0187501 A1 7/2014 Bilodeau et al.
 2014/0314864 A1 10/2014 Cheng et al.
 2015/0023875 A1 1/2015 Farokhzad et al.
 2015/0079001 A1 3/2015 Pomper et al.
 2015/0104387 A1 4/2015 Pomper et al.
 2015/0110715 A1 4/2015 Eder et al.
 2015/0246144 A1 9/2015 Pomper et al.
 2015/0297735 A1 10/2015 Vlahov et al.
 2015/0315196 A1 11/2015 Howard
 2015/0366968 A1 12/2015 Basilion
 2016/0067341 A1 3/2016 Low et al.
 2016/0074526 A1 3/2016 Bilodeau et al.
 2016/0114060 A1 4/2016 Pomper et al.
 2016/0151508 A1 6/2016 Low et al.
 2016/0220694 A1 8/2016 Vlahov et al.
 2016/0287731 A1 10/2016 Vlahov et al.
 2016/0361376 A1 12/2016 Vlahov et al.
 2016/0361432 A1 12/2016 Vlahov et al.
 2016/0361433 A1 12/2016 Vlahov et al.
 2017/0258923 A1 9/2017 Low et al.
 2018/0243431 A1 8/2018 Low et al.

2018/0271988 A1 9/2018 Low et al.
 2018/0271989 A1 9/2018 Low et al.
 2018/0271990 A1 9/2018 Low et al.
 2018/0289827 A1 10/2018 Low et al.
 2018/0289828 A1 10/2018 Low et al.
 2018/0289829 A1 10/2018 Low et al.

FOREIGN PATENT DOCUMENTS

CA 2606138 A1 10/2005
 CA 2696627 A1 2/2009
 CA 2696627 C 9/2016
 CN 1662263 A 8/2005
 CN 101863924 A 10/2010
 CN 102014956 A 4/2011
 CN 102014956 B 6/2015
 CN 104873982 A 9/2015
 DE 202014008232 U1 3/2015
 EP 0116208 A1 8/1984
 EP 1177200 B1 6/2005
 EP 1472541 B1 9/2009
 EP 2187965 A1 5/2010
 EP 2318366 A2 5/2011
 EP 2136788 B1 10/2011
 EP 2373621 A2 10/2011
 EP 2389361 A2 11/2011
 EP 2408755 A2 1/2012
 EP 2644192 A1 10/2013
 EP 2644594 A1 10/2013
 EP 2648766 A1 10/2013
 EP 2436376 B1 7/2014
 EP 2759535 A1 7/2014
 EP 2240171 B1 8/2014
 EP 2170075 B1 12/2014
 EP 2823826 A2 1/2015
 EP 2097111 B1 7/2015
 EP 2921482 A2 9/2015
 EP 2938364 A1 11/2015
 EP 2942065 A1 11/2015
 EP 2958596 A1 12/2015
 EP 2706057 B1 4/2016
 EP 3038996 A1 7/2016
 EP 3388086 A1 10/2018
 JP 2002506204 A 2/2002
 JP 2004536034 A 12/2004
 JP 2005274569 A 10/2005
 JP 2006501149 A 1/2006
 JP 2006514961 A 5/2006
 JP 2006518712 A 8/2006
 JP 2007521803 A 8/2007
 JP 2009519209 A 5/2009
 JP 2010515732 A 5/2010
 JP 2010518112 A 5/2010
 JP 2010536790 A 12/2010
 JP 2014221779 A 11/2014
 JP 5902237 B2 4/2016
 JP 2018150350 A 9/2018
 WO WO-8801622 A1 3/1988
 WO WO-9107418 A1 5/1991
 WO WO-9945374 A2 9/1999
 WO WO-0064911 A1 11/2000
 WO WO-0066091 A1 11/2000
 WO WO-0243773 A2 6/2002
 WO WO-02062398 A2 8/2002
 WO WO-02098885 A1 12/2002
 WO WO-03000201 A2 1/2003
 WO WO-03060523 A1 7/2003
 WO WO-03092742 A1 11/2003
 WO WO-03097647 A1 11/2003
 WO WO-2004010957 A2 2/2004
 WO WO-2004069159 A2 8/2004
 WO WO-2005082023 A2 9/2005
 WO WO-2006012527 A1 2/2006
 WO WO-2006093991 A1 9/2006
 WO WO-2006096754 A2 9/2006
 WO 2006136564 A1 12/2006
 WO WO-2007006041 A2 1/2007
 WO WO-2007022494 A2 2/2007
 WO WO-2007042504 A2 4/2007

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO-2007106869	A1	9/2007
WO	WO-2008057437	A2	5/2008
WO	WO-2008058192	A2	5/2008
WO	WO-2008088648	A2	7/2008
WO	WO-2008098112	A2	8/2008
WO	WO-2008101231	A2	8/2008
WO	WO-2008121949	A1	10/2008
WO	WO-2009002529	A2	12/2008
WO	WO-2009002993	A1	12/2008
WO	WO-2009026177	A1	2/2009
WO	WO-2009070302	A1	6/2009
WO	WO-2009082606	A2	7/2009
WO	WO-2010014933	A2	2/2010
WO	WO-2010065899	A2	6/2010
WO	WO-2010065902	A2	6/2010
WO	WO-2010065906	A2	6/2010
WO	WO-2011014821	A1	2/2011
WO	WO-2010108125	A3	3/2011
WO	WO-2011106639	A1	9/2011
WO	WO-2012078534	A1	6/2012
WO	WO-2012166923	A2	12/2012
WO	WO-2013022797	A1	2/2013
WO	WO-2013028664	A1	2/2013
WO	WO-2013130776	A1	9/2013
WO	WO-2014062697	A2	4/2014
WO	WO-2014078484	A1	5/2014
WO	WO-2014106208	A1	7/2014
WO	WO-2014127365	A1	8/2014
WO	WO-2014134543	A1	9/2014
WO	WO-2015055318	A1	4/2015
WO	WO-2015057250	A1	4/2015
WO	WO-2015171792	A1	11/2015
WO	WO-2016030329	A1	3/2016
WO	WO-2016040179	A1	3/2016

OTHER PUBLICATIONS

"U.S. Appl. No. 14/794,482, Preliminary Amendment filed Nov. 23, 2015", 7 pgs.

"U.S. Appl. No. 14/794,482, Response filed Feb. 28, 2017 to Restriction Requirement dated Nov. 28, 2016", 2 pgs.

"U.S. Appl. No. 14/794,482, Response filed Jul. 19, 2017 to Notice of Non-Compliant Amendment dated May 22, 2017", 3 pgs.

"U.S. Appl. No. 14/794,482, Response filed Nov. 27, 2017 to Non Final Office Action dated Aug. 10, 2017", 5 pgs.

"U.S. Appl. No. 14/794,482, Restriction Requirement dated Nov. 28, 2016", 6 pgs.

"Compound summary for: CID 58099954", Pubchem, (Aug. 19, 2012), 10 pgs.

"Definition of ligand, Random House Kemerman Webster's College Dictionary", downloaded on Jan. 25, 2014 from <<http://www.thefreedictionary.com/liqand>>, 1 pg.

"Radioisotopes in Medicine", [online]. [retrieved on May 12, 2017]. Retrieved from the Internet: <URL: <http://www.word-nuclear.org/information-library/non-power-nuclearapplications/radioisotopes-research/radioisotopes-in-medicine.aspx>>. From <<http://www.word-nuclear.org/information-library/non-power-nuclearapplications/radioisotopes-research/radioisotopes-in-medicine.aspx>>, (Dec. 28, 2016), 20 pages.

Banerjee, S, et al., "Sequential SPECT and Optical Imaging of Experimental Models of Prostate Cancer with a Dual Modality Inhibitor of the Prostate-Specific Membrane Antigen", *Angewandte Chemie International Edition*, 50(39), (2011), 9167-9170.

Banerjee, S R, et al., "Synthesis and Evaluation of Technetium-99m and Rhenium-143 Labeled Inhibitors of the Prostate-Specific Membrane Antigen (PSMA)", *J Med Chem.*, 51(15), (Aug. 14, 2008), 4504-4517.

Benesova, M, et al., "Linker Modifications of DOT A-conjugated Inhibitors of the Prostate-Specific Membrane Antigen (PSMA)", Poster PW082, presented at the European Association of Nuclear Medicine Conference, (Oct. 21, 2013), 1 pg.

Benesova, M., et al., "Linker Modifications of DOT A-conjugated Inhibitors of the Prostate-Specific Membrane Antigen (PSMA)", *Abstract, Eur. J. Nucl. Med. Mol. Imaging*, 40, Suppl. 2, (Oct. 16, 2013), p. S93(2 pages).

Bennett, V. J., "Analysis of fluorescently labeled substance P analogs: binding, imaging and receptor activation", *BMC Chemical Biology*, 1:1. doi:10.1186/1472-6769-1-1, (2001).

Chen, Ying, et al., "Radiohalogenated Prostate-Specific Membrane Antigen (PSMA)-Based Ureas as Imaging Agents for Prostate Cancer", *J. Med. Chem.* 51 (24), (2008), 7933-7943.

Cole, et al., "Cancertheranostics: the rise of targeted magnetic nanoparticles", *Trends in Biotechnology*, 29(7), (2011), 323-332.

Davis, Mindy I, et al., "Crystal Structure of Prostate-Specific Membrane Antigen, A Tumor Marker and Peptidase", *PNAS*, vol. 102, No. 17, (Apr. 26, 2005), 5981-5986.

Eder, et al., "68Ga-complex lipophilicity and the targeting property of a urea-based PSMA inhibitor for PET imaging", *Bioconjugate Chemistry*, 23(4), (2012), 688-697.

Farokhzad, et al., "Nanoparticle-Aptamer Bioconjugates: A New Approach for Targeting Prostate Cancer Cells", *Cancer Research* 64, (2004), 7668-7672.

Foss, Catherine A, et al., "Radiolabeled Small-Molecule Ligands for Prostate-Specific Membrane Antigen: in Vivo Imaging in Experimental Models of Prostate Cancer", *Clinical cancer research* 11.11, (2005), 4022-4028.

Gomez-Hens, et al., "Long Wavelength Fluorophores: New Trends in Their Analytical Use", *Trends in Analytical Chemistry*, 23, (127-136), 2004.

Haseman, M K, et al., "Capromab Pendetide Imaging of Prostate Cancer", *Cancer Biother Radiopharm*, 15, (2000), 131-140.

Henne, et al., "Synthesis and activity of a folate peptide camptothecin prodrug", *ScienceDirect, Bioorganic & Medical Chemistry Letters* 16, (2006), 5350-5355.

Hillier, Shawn M, et al., "Preclinical Evaluation of Novel Glutamate-Urea-Lysine Analogues That Target Prostate-Specific Membrane Antigen as Molecular Imaging Pharmaceuticals for Prostate Cancer", *Cancer Res.* 69(17), (Sep. 1, 2009), 6932-40.

Humblet, V, et al., "An HPLC/Mass Spectrometry Platform for the Development of Multimodality Contrast Agents and Targeted Therapeutics: Prostate-Specific Membrane Antigen Small Molecule Derivatives", *Contrast Med. Mol. Imaging*, 1, (2006), 196-211.

Jackson, Paul F, et al., "Design of NAALADase Inhibitors: A Novel Neuroprotective Strategy", *Current Medicinal Chemistry*, vol. 8, No. 8, (2001), 949-957.

Kaur, G, et al., "Biological evaluation of tubulysin A: a potential anticancer and antiangiogenic natural product", *Biochem. J.*, 396, (2006), 235-242.

Kozikowski, Alan P, et al., "Design of Remarkably Simple, Yet Potent Urea-Based Inhibitors of Glutamate Carboxypeptidase II (NAALADase)", *Journal of Medicinal Chemistry*, vol. 44, No. 3, (2001), 298-301.

Kozikowski, Alan P, et al., "Synthesis of Urea-Based Inhibitors as Active Site Probes of Glutamate Carboxypeptidase II: Efficacy as Analgesic Agents", *Journal of Medicinal Chemistry*, vol. 47, No. 7, (Mar. 2, 2004), 1729-1738.

Kularatne, et al., "Prostate-Specific Membrane Antigen Targeted Imaging and Therapy of Prostate Cancer Using a PSMA Inhibitor as a Homing Ligand", *Molecular Pharmaceutics*, 6(3), (2009), 780-789.

Kularatne, S., "Synthesis and Biological Analysis of Prostate-Specific Membrane Antigen-Targeted Anticancer Prodrugs", *J. Med. Chem.* 53(21), (2010), 7767-7777.

Larson, S. M, et al., "Tumor Localization of 167-18 F-Fluoro-5 α -Dihydrotestosterone Versus 18F-FDG in Patients with Progressive, Metastatic Prostate Cancer", *J Nucl Med*, 45, (2004), 366-373.

Lu, G, et al., "Synthesis and SAR of 99mTc/Re-labeled small molecule prostate specific membrane antigen inhibitors with novel polar chelates", *Bioorganic and Medicinal Chemistry Letters*, 23, (2013), 1557-1563.

Lupold, et al., "Identification and Characterization of Nuclease-Stabilized RNA Molecules That Bind Human Prostate Cancer Cells Via the Prostate-Specific Membrane Antigen", *Cancer Res.*; 62, (2002), 4029-4033.

(56)

References Cited

OTHER PUBLICATIONS

- Majer, Pavel, et al., "Synthesis and Biological Evaluation of Thiol-Based Inhibitors of Glutamate Carboxypeptidase II: Discovery of an Orally Active GCP II Inhibitor", *Journal of Medicinal Chemistry*, vol. 46, No. 10, (2003), 1989-1996.
- Maresca, K. et al., "Molecular Targeting of Prostate Cancer With Small Molecule Inhibitors of Prostate Specific Membrane Antigen (PSMA)", *J. Nucl. Med.* 48 (Supplement 2):25P, (2007).
- Maresca, K. P., "A Series of Halogenated Heterodimeric Inhibitors of Prostate Specific Membrane Antigen (PSMA) as Radiolabeled Probes for Targeting Prostate Cancer", *J. Med. Chem.* 52(2), (2009), 347-357.
- Martin, P., "Ein Neuer Zugang Zu 2'-O-Alkylribonucleosiden und Eigenschaften deren Oligonucleotide", *Helv. Chim. Acta*, 78, no English translation, (1995), 486-504.
- Mcnamara, et al., "Cell type specific delivery of siRNAs with aptamer-siRNA chimeras", *Nature Biotechnology*; 24, (2006), 1005-1015.
- Mease, RC, et al., "N-[N-[(S)-1, 3-Dicarboxypropyl]Carbamoyl]-4-[18F]Fluorobenzyl-L-Cysteine, [18F]DCFB: A New Imaging Probe for Prostate Cancer", *Clin Cancer Res.*, 14, (2008), 3036-3043.
- Melby, et al., *Cancer Research* 53(8), (1993), 1755-1760.
- Mesters, et al., "Structure of Glutamate Carboxypeptidase II, a Drug Target in Neuronal Damage and Prostate Cancer", *The EMBO Journal*, vol. 25, No. 6, (2006), 1375-1384.
- Mier, W., et al., "Conjugation of DOTA Using Isolated Phenolic Active Elers: The Labeling and Biodistribution of Albumin as Blood Pool Marker", *Bioconjugate Chem.*, 16, (2005), 237-240.
- Nan, F, et al., "Dual Function Glutamate-Related Ligands: Discovery of a Novel, Potent Inhibitor of Glutamate Carboxypeptidase II Possessing mGluR3 Agonist Activity", *J Med Chem*, 43, (2000), 772-774.
- Olsnes, S, et al., "Immunology Today", 10, (1989), 291-295.
- Paranjpe, et al., "Tumor-targeted bioconjugate based delivery of camptothecin: design, synthesis and in vitro evaluation", *ScienceDirect Journal of Controlled Release* 100, (2004), 275-292.
- Pathalk, et al., "Enzymic protecting group techniques in organic synthesis", *Stereosel, Biocatal*, (2000), 775-797.
- Peltier, et al., "The Total Synthesis of Tubulysin D", *J. Am. Chem. Soc.* 128, (2006), 16018-19.
- Pomper, M. G, et al., "11C-MCG: synthesis, uptake selectivity, and primate PET of a probe for glutamate carboxypeptidase II (NAALADase)", *Mol Imaging*, 1, (2002), 96-101.
- Ranasinghe, M. G, et al "Facile Synthesis of Unsymmetrical Thiol-sulfonates via Sulfonylation of Mercaptans", *Synthetic Communications*, vol. 18, No. 3, (1988), 227-232.
- Reddy, et al., "PSMA-specific anti-tumor activity of the targeted-tubulysin conjugate, EC1169", *Poster, American Association for Cancer Research Annual Meeting (Apr. 8, 2013)*, (Apr. 8, 2013), 1 pg.
- Reddy, et al., "PSMA-specific anti-tumor activity of the targeted-tubulysin conjugate, EC1169", *Presentation Abstract, American Association for Cancer Research Annual Meeting (Apr. 8, 2013)*, (Apr. 8, 2013), 1 pg.
- Reske, S. N, et al., "Imaging Prostate Cancer with 11C-Choline PET/CT", *J Nucl Med*, 47, (2006), 1249-1254.
- Rinnab, L, et al., "Evaluation of [11C]-choline positron-emission/computed tomography in patients with increasing prostate-specific antigen levels after primary treatment for prostate cancer", *BJU Int*, 100, (2007), 786, 793.
- Rosenthal, S. A, et al., "Utility of Capromab Pendetide (ProstaScint) Imaging in the Management of Prostate Cancer", *Tech Urol*, 7, (2001), 27-37.
- Roy, et al., "DUPA Conjugation of a Cytotoxic Indenoisoquinoline Topoisomerase I Inhibitor for Selective Prostate Cancer Cell Targeting" *J. Med. Chem.* 58, (2015), 3094-3103.
- Schafer, et al., "A dimerized urea-based inhibitor of the prostate-specific membrane antigen for 68Ga-PET imaging of prostate cancer", *EJNMMI Research*, 2:23, (2012), 11 pgs.
- Scher, B., et al., "Value of 11 C-choline PET and PET/CT in patients with suspected prostate cancer", *Eur. J. Nucl. Med. Mol. Imaging.*, 34, (2007), 45-53.
- Schulke, N, et al., "The homodimer of prostate-specific membrane antigen is a functional target for cancer therapy", *Proc Natl Acad Sci U S A*, 100, (2003), 12590-12595.
- Schuster, D. M, et al., "Initial Experience with the Radiotracer Anti-1-Amino-3-18F-Fluorocyclobutane-1-Carboxylic Acid with PET/CT in Prostate Carcinoma", *J Nucl Med*, 48, (2007), 56-63.
- Silverman, "The Organic Chemistry of Drug Design and Drug Action", Elsevier Academic Press (2nd Ed. 2003), (2003), 1-592 (304 pages).
- Silvola, et al., "A118F-NOTA-Folate Accumulates in Atherosclerotic Plaques and Can be Detected by PET/CT", *Poster presented Nov. 7, 2015 in Orlando, FL at the 2015 American Heart Association, ReSuscitation Science Symposium*, [Online] Retrieved from the Internet: <<https://aha.scientificposters.com/epsView.cfm?Vud4d4t239pJgyhGx5DzRxHdvD8LemQoMXh31P5rozni1u5bJrEupTaG09TFyJlzABwMcSXfzcc%3D>>, (Nov. 7, 2015), 1 pg.
- Silvola, et al., "A118F-NOTA-Folate Accumulates in Atherosclerotic Plaques and Can be Detected by PET/CT", *Abstract 18873, Circulation [Online] Retrieved from the internet: <http://circ.ahajournals.org/content/132/Suppl_3/A_18873?cited-by=legid=circulationaha;132/Suppl_3/A_18873; Circulation, 2015, 132:A_18873>*, (Nov. 10, 2015), 2 pgs.
- Tasch, J, et al., "A Unique Folate Hydrolase, Prostate-Specific Membrane Antigen (PSMA): A Target for Immunotherapy?", *Crit. Rev. Immunol.*, 21, (2001), 249-261.
- Tehrani, O. S., "Tumor Imaging Using 1-(2'-deoxy-2'-18F-Fluoro-2'-D-Arabinofuranosyl)Thymine and PET", *J Nuc/ Med*, 48, (2007), 1436-1441.
- Truffert, et al., "Tetrahedron", 52, (1996).
- Vees, H, et al., "18F-choline and/or 11C-acetate positron emission tomography: detection of residual or progressive subclinical disease at very low prostate-specific antigen values (<ng/mL) after radical prostatectomy", *BJU Int*, 99, (2007), 1415-1420.
- Vlahov, et al., "Design and regioselective synthesis of a new generation of targeted chemotherapeutics. Part 1: EC145, a folic acid conjugate of desacetylvinblastine monohydrate", *ScienceDirect, Bioorganic & Medical Chemistry Letters* 16, (2006), 5093-5096.
- Wang, K., et al., "Prostate-Specific Membrane Antigen Targeted Tubulysin Conjugates for Cancer Therapy", *Poster, 246th American Chemical Society Meeting and Exposition (Sep. 8, 2013)*, (Sep. 8, 2013).
- Wiberg, et al., "A comparison of some properties of C=O and C=S bonds", *ARKIVOC*, (2011), 45-56.
- Yang, et al., "Characterization of the pH of Folate Receptor-Containing Endosomes and the Rate of Hydrolysis of Internalized Acid-Labile Folate-Drug Conjugates", *JPET* 321, (2007), 462-468.
- Zhou, J., et al., "Naag Peptidase Inhibitors And Their Potential for Diagnosis And Therapy", *Nat Rev Drug Discovery*, 4, (2005), 1015-1026.
- Zophel, K, et al., "Is 11C-choline the most appropriate tracer for prostate cancer?", *Eur J Nucl Med Mol Imaging*, 31, (2004), 756-759.
- "99mTc-MIP-1404 for Imaging Prostate Cancer: Phase I Clinical Study to Assess the Image Quality of a Simplified Kit Formulation Compared to a Multi-step Preparation of 99mTc-MIP-1404", *ClinicalTrials.gov Identifier: NCT01654874*, [online]. Retrieved from the Internet: <[URL: https://clinicaltrials.gov/ct2/show/NCT01654874](https://clinicaltrials.gov/ct2/show/NCT01654874)>, (Aug. 1, 2012), 7 pgs.
- "A Phase I Pilot Study of 99mTc-MIP-1404 SPECT/CT Imaging to Histology in Men With Prostate Cancer", *ClinicalTrials.gov Identifier: NCT01615406*, [online]. Retrieved from the internet: <[URL: https://clinicaltrials.gov/ct2/show/NCT01615406](https://clinicaltrials.gov/ct2/show/NCT01615406)>, (Jun. 8, 2012), 7 pgs.
- "A Phase 2 Study With MIP-1404 in Men With High-Risk PC Scheduled for RP and EPLND Compared to Histopathology", *ClinicalTrials.gov Identifier: NCT01667536*, [online]. Retrieved from the Internet: <[URL: https://clinicaltrials.gov/ct2/show/NCT01667536?id=NCT01667536](https://clinicaltrials.gov/ct2/show/NCT01667536?id=NCT01667536)>, (Aug. 17, 2012), 10 pgs.
- "U.S. Appl. No. 12/673,931, 312 Amendment filed Jul. 2, 2015", 165 pgs.

(56)

References Cited

OTHER PUBLICATIONS

“U.S. Appl. No. 12/673,931, Final Office Action dated Jan. 30, 2014”, 14 pgs.
 “U.S. Appl. No. 12/673,931; Non Final Office Action dated Apr. 11, 2013”; 15 pgs.
 “U.S. Appl. No. 12/673,931, Non Final Office Action dated Dec. 5, 2014”, 11 pgs.
 “U.S. Appl. No. 12/673,931; Notice of Allowance dated Apr. 9, 2015”, 8 pgs.
 “U.S. Appl. No. 12/673,931, Notice of Allowance dated Oct. 7, 2015”, 8 pgs.
 “U.S. Appl. No. 12/673,931, Preliminary Amendment filed Feb. 17, 2010”, 6 pgs.
 “U.S. Appl. No. 12/673,931, Preliminary Amendment filed Mar. 3, 2011”, 7 pgs.
 “U.S. Appl. No. 12/673,931, PTO Response to Rule 312 Communication dated Jul. 15, 2015”, 2 pgs.
 “U.S. Appl. No. 12/673,931, Response filed Mar. 5, 2015 to Non Final Office Action dated Dec. 5, 2014”, 8 pgs.
 “U.S. Appl. No. 12/673,931, Response filed May 30, 2014 to Final Office Action dated Jan. 30, 2014”, 15 pgs.
 “U.S. Appl. No. 12/673,931, Response filed Oct. 2, 2012 to Restriction Requirement dated Aug. 2, 2012”, 12 pgs.
 “U.S. Appl. No. 12/673,931, Response filed Oct. 11, 2013 to Non Final Office Action dated Apr. 11, 2013”, 21 pgs.
 “U.S. Appl. No. 12/673,931, Restriction Requirement dated Aug. 2, 2012”, 9 pgs.
 “U.S. Appl. No. 12/673,931, Supplemental Amendment filed Jan. 18, 2013”, 9 pgs.
 “U.S. Appl. No. 13/826,079, Notice of Allowance dated Jun. 2, 2014”, 8 pgs.
 “U.S. Appl. No. 13/826,079, Notice of Allowance dated Sep. 12, 2014”, 8 pgs.
 “U.S. Appl. No. 13/826,079, Preliminary Amendment filed Mar. 14, 2013”, 4 pgs.
 “U.S. Appl. No. 13/826,079, Preliminary Amendment filed May 16, 2014”, 3 pgs.
 “U.S. Appl. No. 15/606,913,312 Amendment filed Jul. 1, 2019”, 4 pgs.
 “U.S. Appl. No. 15/606,913, Advisory Action dated Apr. 16, 2019”, 3 pgs.
 “U.S. Appl. No. 15/606,913, Examiner Interview Summary dated May 8, 2019”, 3 pgs.
 “U.S. Appl. No. 15/606,913, Final Office Action dated Nov. 5, 2018”, 8 pgs.
 “U.S. Appl. No. 15/606,913, Non Final Office Action dated Apr. 3, 2018”, 10 pgs.
 “U.S. Appl. No. 15/606,913, Notice of Allowance dated Jun. 26, 2019”, 10 pgs.
 “U.S. Appl. No. 15/606,913, Preliminary Amendment filed May 26, 2017”, 4 pgs.
 “U.S. Appl. No. 15/606,913, Preliminary Amendment filed May 30, 2017”, 5 pgs.
 “U.S. Appl. No. 15/606,913, Response filed Apr. 4, 2019 to Non Final Office Action dated Nov. 5, 2018”, 6 pgs.
 “U.S. Appl. No. 15/606,913, Response filed Jul. 3, 2018 to Non Final Office Action dated Apr. 3, 2018”, 7 pgs.
 “U.S. Appl. No. 15/606,913, Response filed Dec. 29, 2017 to Restriction Requirement dated Nov. 3, 2017”, 7 pgs.
 “U.S. Appl. No. 15/606,913, Restriction Requirement dated Nov. 3, 2017”, 6 pgs.
 “U.S. Appl. No. 15/959,110, Advisory Action dated Apr. 17, 2019”, 3 pgs.
 “U.S. Appl. No. 15/959,110, Final Office Action dated Jan. 11, 2019”, 15 pgs.
 “U.S. Appl. No. 15/959,110, Non Final Office Action dated Jun. 27, 2018”, 9 pgs.
 “U.S. Appl. No. 15/959,110, Non Final Office Action dated Jun. 27, 2019”, 14 pgs.

“U.S. Appl. No. 15/959,110, Preliminary Amendment filed Apr. 20, 2018”, 4 pgs.
 “U.S. Appl. No. 15/959,110, Preliminary Amendment filed Apr. 24, 2018”, 6 pgs.
 “U.S. Appl. No. 15/959,110, Response filed Sep. 25, 2018 to Non Final Office Action dated Jun. 27, 2018”, 10 pgs.
 “U.S. Appl. No. 15/959,110, Response filed Apr. 3, 2019 to Final Office Action dated Jan. 11, 2019”, 8 pgs.
 “U.S. Appl. No. 15/990,083, Examiner Interview Summary dated Jun. 12, 2019”, 4 pgs.
 “U.S. Appl. No. 15/990,083, Non Final Office Action dated Jan. 25, 2019”, 13 pgs.
 “U.S. Appl. No. 15/990,083, Preliminary Amendment filed May 25, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,083, Preliminary Amendment filed May 29, 2018”, 5 pgs.
 “U.S. Appl. No. 15/990,083, Response filed Sep. 27, 2018 to Restriction Requirement dated Jul. 27, 2018”, 2 pgs.
 “U.S. Appl. No. 15/990,083, Restriction Requirement dated Jul. 27, 2018”, 6 pgs.
 “U.S. Appl. No. 15/990,095, Non Final Office Action dated Jan. 9, 2019”, 8 pgs.
 “U.S. Appl. No. 15/990,095, Non Final Office Action dated Jun. 13, 2019”, 7 pgs.
 “U.S. Appl. No. 15/990,095, Preliminary Amendment filed May 25, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,095, Preliminary Amendment filed May 29, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,095, Response filed Feb. 20, 2019 to Non Final Office Action dated Jan. 9, 2019”, 5 pgs.
 “U.S. Appl. No. 15/990,095, Response filed Sep. 21, 2018 to Restriction Requirement dated Jul. 23, 2018”, 2 pgs.
 “U.S. Appl. No. 15/990,095, Restriction Requirement dated Jul. 23, 2018”, 6 pgs.
 “U.S. Appl. No. 15/990,111, Examiner Interview Summary dated May 8, 2018”, 3 pgs.
 “U.S. Appl. No. 15/990,111, Non Final Office Action dated Jan. 7, 2019”, 16 pgs.
 “U.S. Appl. No. 15/990,111, Preliminary Amendment filed May 25, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,111, Preliminary Amendment filed May 29, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,111, Response filed Sep. 21, 2018 to Restriction Requirement dated Jul. 23, 2018”, 2 pgs.
 “U.S. Appl. No. 15/990,111, Restriction Requirement dated Jul. 23, 2018”, 6 pgs.
 “U.S. Appl. No. 15/990,136, Non Final Office Action dated Jan. 7, 2019”, 11 pgs.
 “U.S. Appl. No. 15/990,136, Preliminary Amendment filed May 25, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,136, Preliminary Amendment filed May 29, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,136, Response filed Sep. 27, 2018 to Restriction Requirement dated Jul. 27, 2018”, 2 pgs.
 “U.S. Appl. No. 15/990,136, Restriction Requirement dated Jul. 27, 2018”, 6 pgs.
 “U.S. Appl. No. 15/990,144, Examiner Interview Summary dated Jun. 13, 2019”, 4 pgs.
 “U.S. Appl. No. 15/990,144, Non Final Office Action dated Jan. 15, 2019”, 10 pgs.
 “U.S. Appl. No. 15/990,144, Preliminary Amendment filed May 25, 2018”, 4 pgs.
 “U.S. Appl. No. 15/990,144, Preliminary Amendment filed May 29, 2018”, 3 pgs.
 “U.S. Appl. No. 15/990,144, Response filed Sep. 21, 2018 to Restriction Requirement dated Jul. 23, 2018”, 2 pgs.
 “U.S. Appl. No. 15/990,144, Restriction Requirement dated Jul. 23, 2018”, 6 pgs.
 “U.S. Appl. No. 15/990,152, Examiner interview Summary dated May 8, 2019”, 4 pgs.
 “U.S. Appl. No. 15/990,152, Non Final Office Action dated Jan. 17, 2019”, 10 pgs.

(56)

References Cited

OTHER PUBLICATIONS

"U.S. Appl. No. 15/990,152, Preliminary Amendment filed May 25, 2018", 4 pgs.

"U.S. Appl. No. 15/990,152, Preliminary Amendment filed May 29, 2018", 4 pgs.

"U.S. Appl. No. 15/990,152, Response filed Sep. 27, 2018 to Restriction Requirement dated Jul. 27, 2018", 2 pgs.

"U.S. Appl. No. 15/990,152, Restriction Requirement dated Jul. 27, 2018", 6 pgs.

- "Australian Application Serial No. 2008289108, First Examination Report dated Aug. 15, 2013", 2 pgs.

"Australian Application Serial No. 2008289108, Response filed Aug. 15, 2014 to First Examination Report dated Aug. 15, 2013", 14 pgs.

"Australian Application Serial No. 2018200419, First Examination Report dated Oct. 24, 2018", 4 pgs.

"Canadian Application Serial No. 2,696,627, Examiners Rule 30(2) Requisition dated Jun. 22, 2015", 3 pgs.

"Canadian Application Serial No. 2,696,627, Examiners Rule 30(2) Requisition dated Oct. 9, 2014", 3 pgs.

"Canadian Application Serial No. 2,696,627, Response filed Apr. 9, 2015 to Examiner's Rule 30(2) Requisition dated Oct. 9, 2014", 22 pgs.

"Canadian Application Serial No. 2,696,627, Response filed Dec. 18, 2015 to Examiner's Rule 30(2) Requisition dated Jun. 22, 2015", 12 pgs.

"Canadian Application Serial No. 2,696,627, Voluntary Amendment filed Apr. 24, 2015", 10 pgs.

"Canadian Application Serial No. 2,696,627, Voluntary Amendment filed May 18, 2010", 8 pgs.

"Canadian Application Serial No. 2,696,627, Voluntary Amendment filed Aug. 15, 2013", 15 pgs.

"Canadian Application Serial No. 2,924,360, Office Action dated Sep. 11, 2018", 3 pgs.

"Chinese Application Serial No. 201480071256.X, Office Action dated Apr. 20, 2017", 19 pgs.

"Chinese Application Serial No. 201510221167.5, Office Action dated Mar. 9, 2018", (w/ English Translation), 17 pgs.

"Chinese Application Serial No. 201510221167.5, Office Action dated Jun. 3, 2019", W/ English Translation, 19 pgs.

"Chinese Application Serial No. 201510221167.5, Office Action dated Jun. 26, 2017", w/ English translation, 21 pgs.

"Chinese Application Serial No. 201510221167.5, Office Action dated Nov. 14, 2018", (w/ English Translation), 7 pgs.

"Chinese Application Serial No. 201510221167.5, Response filed Mar. 28, 2019 to Office Action dated Nov. 14, 2018", (w/ English Translation of Claims), 21 pgs.

"Chinese Application Serial No. 201510221167.5, Response filed Nov. 8, 2017 to Office Action dated Jun. 26, 2017", w/ English claims, 22 pgs.

"Chinese Application Serial No. 201610184873.1, Office Action dated Jul. 24, 2018", W/ English Translation, 15 pgs.

"Eurasian Application Serial No. 16904952.8, Office Action dated Dec. 20, 2018", 4 pgs.

"Eurasian Application Serial No. 201690862/28, Office Action dated May 22, 2017", 2 pgs.

"European Application Serial No. 08798020.7, Communication Pursuant to Article 94(3) EPC dated May 13, 2016", 4 pgs.

"European Application Serial No. 08798020.7, Communication Pursuant to Article 94(3) EPC dated May 16, 2018", 3 pgs.

"European Application Serial No. 08798020.7, Communication Pursuant to Article 94(3) EPC dated Jun. 21, 2017", 3 pgs.

"European Application Serial No. 08798020.7, Extended European Search Report dated Oct. 28, 2014", 4 pgs.

"European Application Serial No. 08798020.7, Intention to Grant dated Mar. 5, 2019", 102 pgs.

"European Application Serial No. 08798020.7, Response filed Apr. 27, 2010 to Communication pursuant to Rules 161(1) and 162 EPC dated Mar. 26, 2010", 14 pgs.

"European Application Serial No. 08798020.7, Response filed May 22, 2015 to Extended European Search Report dated Oct. 28, 2014", 10 pgs.

"European Application Serial No. 08798020.7, Response filed Sep. 20, 2018 to Communication Pursuant to Article 94(3) EPC dated May 16, 2018", 77 pgs.

"European Application Serial No. 08798020.7, Response filed Sep. 23, 2016 to Communication Pursuant to Article 94(3) EPC dated May 13, 2016", 12 pgs.

, Et al., "European Application Serial No. 08798020.7, Response filed Dec. 21, 2017 to Communication Pursuant to Article 94(3) EPC dated Jun. 21, 2017", 9 pgs.

"European Application Serial No. 14861854.9, extended European Search Report dated Sep. 7, 2017", 12 pgs.

"European Application Serial No. 14861854.9, Office Action dated Sep. 25, 2018", 4 pgs.

"European Application Serial No. 14861854.9, Partial Supplementary European Search Report dated May 19, 2017", 14 pgs.

"European Application Serial No. 14861854.9, Response filed Apr. 6, 2018 to Office Action dated Sep. 26, 2017", 15 pgs.

"European Application Serial No. 14861854.9, Result of Consultation dated Jan. 25, 2019", 3 pgs.

"European Application Serial No. 18175078, Extended European Search Report dated Sep. 14, 2018", 5 pgs.

"European Application Serial No. 18175078.7, Extended European Search Report dated Sep. 14, 2018", 5 pgs.

"European Application Serial No. 18175078.7, Response filed Apr. 16, 2019 to Extended European Search Report dated Sep. 14, 2018", 81 pgs.

"Indian Application Serial No. 949/KOLNP/2010, Examination Report dated Mar. 31, 2015", 2 pgs.

"Indian Application Serial No. 949/KOLNP/2010, Response filed Mar. 22, 2016 to Examination Report dated Mar. 31, 2015", 8 pgs.

"International Application Serial No. PCT/US2008/073375, International Preliminary Report Patentability dated Feb. 24, 2010", 6 pgs.

"International Application Serial No. PCT/US2008/073375, International Search Report dated Nov. 17, 2008", 1 pg.

"International Application Serial No. PCT/US2008/073375, Written Opinion dated Nov. 17, 2008", 5 pgs.

"International Application Serial No. PCT/US2009/061049, International Search Report dated Mar. 24, 2010", 5 pgs.

"International Application Serial No. PCT/US2009/061049, Written Opinion dated Mar. 24, 2010", 8 pgs.

"International Application Serial No. PCT/US2009/061067, International Search Report dated Jun. 17, 2010", 4 pgs.

"International Application Serial No. PCT/US2009/061067, Written Opinion dated Jun. 17, 2010", 6 pgs.

"International Application Serial No. PCT/US2011/026238, International Search Report dated Apr. 27, 2011", 3 pgs.

"International Application Serial No. PCT/US2011/026238, Written Opinion dated Apr. 27, 2011", 6 pgs.

"International Application Serial No. PCT/US2013/070007, International Search Report dated Mar. 5, 2014", 3 pgs.

"International Application Serial No. PCT/US2013/070007, Written Opinion dated Mar. 5, 2014", 11 pgs.

"International Application Serial No. PCT/US2014/065467, International Search Report dated Apr. 15, 2015", 5 pgs.

"International Application Serial No. PCT/US2014/065467, Written Opinion dated Apr. 15, 2015", 5 pgs.

"International Application Serial No. PCT/US2016/012653, International Search Report dated Mar. 11, 2016", 3 pgs.

"Istard Posters", Eur J Nucl Med Mol Imaging, 39 (Suppl2):, (2012), S304-S353.

"Japanese Application Serial No. 2017-210775, Office Action dated Nov. 6, 2018", w/English Translation, 4 pgs.

"Japanese Application Serial No. 2017-210775, Written Opinion and Amendment filed Apr. 22, 2019 to Office Action dated Nov. 6, 2018", (w/ English Translation), 12 pgs.

"Japanese Application Serial No. 2017-223872, Office Action dated Feb. 26, 2019", (w/ English Translation), 6 pgs.

"Japanese Application Serial No. 2017-223872, Office Action dated Oct. 9, 2018", (w/ English Translation), 6 pgs.

(56)

References Cited

OTHER PUBLICATIONS

- "Japanese Application Serial No. 2017-223872, Response filed May 22, 2019 to Office Action dated Feb. 26, 2019", w/ English Claims, 10 pgs.
- "Japanese Application Serial No. 2017-223872, Written Amendment filed Feb. 7, 2018", (w/ English Translation), 13 pgs.
- "Japanese Application Serial No. 2017-223872, Written Opinion and Amendment filed Jan. 7, 2019 in response to Office Action dated Oct. 9, 2018", (w/ English Translation), 14 pgs.
- Japanese Application Serial No. 2017-223872, Notification of Reasons for Rejection dated Jul. 2, 2019 w/ English Translation 4 pgs.
- "Japanese Application Serial No. 2018-100343, Notification of Reasons for Rejection dated Mar. 5, 2019", (w/ English Translation), 5 pgs.
- "Pilot Study of 99mTc-MIP-1404 SPECT/CT Imaging in Men With Prostate Cancer Undergoing Prostatectomy and/or Pelvic Lymph Node Dissection", ClinicalTrials.gov Identifier: NCT01572701, (Apr. 6, 2012), 7 pgs.
- "South African Application Serial No. 2010/01875, Voluntary Amendment Filed Apr. 10, 2019", 8 pgs.
- Afshar-Oromieh, A., et al., "[68Ga]Gallium-labelled PSMA ligand as superior PET tracer for the diagnosis of prostate cancer: comparison with 18F-FECH", *Eur J Nucl Med Mol Imaging*, 39, Haberkorn, (2012), 1085-1086.
- Afshar-Oromieh, A., et al., "Comparison of PET imaging with a 68Ga-labelled PSMA ligand and 18F-choline-based PET/CT for the diagnosis of recurrent prostate cancer", *Eur J Nucl Med Mol Imaging*, 41, Zechmann, (2014), 11-20.
- Afshar-Oromieh, A., et al., "Comparison of PET/CT and PET/MRI hybrid systems using a 68 Ga-labelled PSMA ligand for the diagnosis of recurrent prostate cancer: initial experience", *Eur J Nucl Med Mol Imaging*, 41, Haber, (2014), 887-897.
- Afshar-Oromieh, A., et al., "PET imaging with a [68Ga]gallium-labelled PSMA ligand for the diagnosis of prostate cancer: biodistribution in humans and first evaluation of tumour lesions", *Eur J Nucl Med Mol Imaging*, 40(4), (2013), 486-495.
- Afshar-Oromieh, A., et al., "PET/MRI with a 68Ga-PSMA ligand for the detection of prostate cancer", *Eur J Nucl Med Mol Imaging*, 40, Haberkorn, (2013), 1629-1630.
- Afshar-Oromieh, A., et al., "The diagnostic value of PET/CT imaging with the 68Ga-labelled PSMA ligand HBEDCC in the diagnosis of recurrent prostate cancer", *Eur J Nucl Med Mol Imaging*, 42(2), (2015), 197-209.
- Aggarwal, S., et al., "A Dimeric, Peptide That Binds Selectively to Prostate-Specific Membrane Antigen and Inhibits its Enzymatic Activity", *Cancer Res*, 66(18), (2006), 9171-9177.
- Alt, K., et al., "High-Resolution Animal PET Imaging of Prostate Cancer Xenografts with Three Different 64Cu-Labeled Antibodies against Native Cell-Adherent PSMA", *The Prostate* 70(13), (2010); 1413-1421.
- Ananias, H. J.K., et al., "Expression of the Gastrin-Releasing Peptide Receptor, the Prostate Stem Cell Antigen and the Prostate-Specific Membrane Antigen in Lymph Node and Bone Metastases of Prostate Cancer", *The Prostate*, 69, (2009), 1101-1108.
- Anderson, M. O., et al., "Substrate specificity of prostate-specific membrane antigen", *Bioorganic & Medicinal Chemistry* 15, (2007), 6678-6686.
- Antunes, A. A., et al., "PGC and PSMA in prostate cancer diagnosis: tissue analysis from biopsy samples", *Int Braz J Urol.*, 39(5), (2013), 649-56.
- Armor, T., et al., "A comparison of 2D and 3D regions within the same patient to derive organ and tissue kinetics", (Abstract), *J Nucl Med.*, vol. 53 No. Supplement 1 13, (May 2012), 1 pg.
- Bacich, D. J., et al., "Cloning, expression, genomic localization, and enzymatic activities of the mouse homolog of prostate-specific membrane antigen/NAALADase/folate hydrolase", *Mammalian Genome* 12(2), (2001), 117-123.
- Baiz, D., et al., "Synthesis and Characterization of a Novel Prostate Cancer-Targeted Phosphatidylinositol-3-kinase Inhibitor Prodrug", *J. Med. Chem.* 55, (2012), 8038-8046.
- Banerjee, S. R., et al., "64Cu-Labeled Inhibitors of Prostate-Specific Membrane Antigen for PET Imaging of Prostate Cancer", *J. Med. Chem.* 57, (2014), 2657-2669.
- Banerjee, S. R., et al., "68Ga-Labeled Inhibitors of Prostate-Specific Membrane Antigen (PSMA) for Imaging Prostate Cancer", *J. Med. Chem.* 53;(2010), 5333-5341.
- Banerjee, S. R., et al., "A Modular Strategy to Prepare Multivalent Inhibitors of Prostate-Specific Membrane Antigen (PSMA)", *Oncotarget*, vol. 2 No. 12, (2011), 1244-1253.
- Banerjee, S. R., et al., "Effect of Chelator on the Pharmacokinetics of 99mTc-Labeled Imaging Agents for the Prostate-Specific Membrane Antigen (PSMA)", *J. Med. Chem.* 56, (2013), 6108-6121.
- Barinka, C., et al., "A high-resolution structure of ligand-free human glutamate carboxypeptidase II", *Acta Cryst*, F63, (2007), 150-153.
- Barinka, C., et al., "Interactions between Human Glutamate Carboxypeptidase II and Urea-Based Inhibitors: Structural Characterization", *J. Med. Chem.* 51, (2008), 7737-7743.
- Barinka, C., et al., "Structural Insight into the Pharmacophore Pocket of Human Glutamate Carboxypeptidase II", *J. Med. Chem.*, 50, (2007), 3267-3273.
- Barrett, J. A., et al., "First-in-Man Evaluation of 2 High-Affinity PSMA-Avid Small Molecules for Imaging Prostate Cancer", *J Nucl Med.*, 54(3), (2013), 380-387.
- Beheshti, M., et al., "Prostate Cancer: Role of SPECT and PET in Imaging Bone Metastases", *Semin Nucl Med* 39(6), (2009), 396-407.
- Belloli, S., et al., "Characterization of preclinical models of prostate cancer using PET-based molecular imaging", *Eur J Nucl Med Mol Imaging* 36:, (2009), 1245-1255.
- Ben Jemaa, A., et al., "A Comparison of the Biological Features of Prostate Cancer with (Psa+, PSMA+) Profile according to RKIP", *BioMed Research International* vol. 2013, Article ID 409179, (2013), 7 pgs.
- Ben Jemaa, A., et al., "A novel regulation of PSMA and PSA expression by Q640X AR in 22R v1 and LNCaP prostate cancer cells", *Cell Biol Int*, 37(5), (2013), 464-470.
- Ben Jemaa, A., et al., "Cellular distribution and heterogeneity of PSA and PSMA expression in normal, hyperplasia and human prostate cancer", *La Tunisie Medicale*, 91(7), (2013), 458-463.
- Bostwick, D. G., et al., "Prostate Specific Membrane Antigen Expression in Prostatic Intraepithelial Neoplasia and Adenocarcinoma", *Cancer*, 82, (1998), 2256-2261.
- Bouchelouche, K., et al., "Image and treat: an individualized approach to urological tumors", *Curr Opin Oncol*, 22(3), (2010), 274-280.
- Bouchelouche, K., et al., "Imaging Prostate Cancer: An Update on Positron Emission Tomography and Magnetic Resonance Imaging", *Curr Urol Rep*, 11, (2010), 180-190.
- Bouchelouche, K., et al., "PET/CT Imaging and Radioimmunotherapy of Prostate Cancer", *Semin Nucl Med*, 41, (2011), 29-44.
- Bouchelouche, K., et al., "Positron emission tomography/computed tomography and radioimmunotherapy of prostate cancer", *Curr Opin Oncol*, 21(5), 469-474.
- Bouchelouche, K., et al., "Prostate Specific Membrane Antigen-A Target for Imaging and Therapy with Radionuclides", *Discov Med*. 9(44):, (Jan. 2010), 55-61.
- Bzdega, T., et al., "The cloning and characterization of a second brain enzyme with NAAG peptidase activity", *Journal of Neurochemistry*, 89(3), (2004), 627-635.
- Ceci, F., et al., "11C-Choline PET/CT in patients with hormone-resistant prostate cancer showing biochemical relapse after radical prostatectomy", *Eur J Nucl Med Mol Imaging*, 40, (2013), 149-155.
- Chandran, S. S., et al., "Characterization of a targeted nano particle functionalized with urea-based inhibitor of prostate-specific membrane antigen (PSMA)", *Cancer Biology & Therapy*, 7(6), (2008), 974-982.
- Chang, S. S., et al., "Five Different Anti-Prostate-Specific Membrane Antigen (PSMA) Antibodies Confirm PSMA Expression in Tumor-Associated Neovasculature", *Cancer Research*, 59, (Jul. 1, 1999), 3192-3198.
- Chang, S. S., et al., "The clinical role of prostate-specific membrane antigen (PSMA)", *Urologic Oncology*, 7(1), (2002), 7-12.

(56)

References Cited

OTHER PUBLICATIONS

- Chen, Y., et al., "2-(3-{1-Carboxy-5-[(6-[18F]Fluoro-Pyridine-3-Carbonyl)-Amino]-Pentyl}-Ureido)-Pentanedioic Acid, [18F]DCFPyL, a PSMA-Based PET Imaging Agent for Prostate Cancer", *Clin Cancer Res*, 17(24), (2011), 7645-7453.
- Chen, Y., et al., "A low molecular weight PSMA-based fluorescent imaging agent for cancer", *Biochemical and Biophysical Research Communications* 390, (2009), 624-629.
- Chen, Y., et al., "Synthesis and Biological Evaluation of Low Molecular Weight Fluorescent Imaging Agents for the Prostate-Specific Membrane Antigen", *Bioconjugate Chem.*, 23(12), (2012), 2377-2385.
- Chen, Z., et al., "PSMA-Targeted Theranostic Nanoplex for Prostate Cancer Therapy", *ACS Nano*, 6(9), (2012), 7752-7762.
- Chopra, A., et al., "68Ga-Labeled 2-[3-(1-carboxy-5-{7-[5-carboxy-5-(3-phenyl-2-{3-phenyl-2[2-(4,7,10-tris-carboxymethyl-1,4,7,10-tetraazacyclododec-1-yl)acetylamino]propionylamino}propionylamino)pentylcarbamoyl]heptanoylamino}pentyl)ureido]pentanedioic acid [68Ga]6", [Updated Dec. 28, 2010]. In: *Molecular Imaging and Contrast Agent Database (MICAD)* [Internet]. Bethesda (MD): National Center for Biotechnology Information (US); (Sep. 27, 2010), 1-6.
- Chopra, A., et al., "68Ga-labeled 2-[3-[5-(7-{1-benzyloxycarbonyl-5-[2-(4,7,10-tris-carboxymethyl-1,4,7,10-tetraazacyclododec-1-yl)acetylamino]pentylcarbamoyl]-heptanoylamino)-1-carboxypentyl]ureido]pentanedioic acid [68Ga]3", [Updated Dec. 28, 2010]. In: *Molecular Imaging and Contrast Agent Database (MICAD)* [Internet]. Bethesda (MD): National Center for Biotechnology Information (US); (Sep. 28, 2010), 1-6.
- Chuu, C.-P., et al., "Androgen suppresses proliferation of castration-resistant LNCaP 104-R2 prostate cancer cells through androgen receptor, Skp2, and c-Myc", *Cancer Sci*, 102(11), (2011), 2022-2028.
- Cimitan, M., et al., "[18F]fluorocholine PET/CT imaging for the detection of recurrent prostate cancer at PSA relapse: experience in 100 consecutive patients", *Eur J Nucl Med Mol Imaging*, 33, (2006), 1387-1398.
- Colabufo, N. A., et al., "PB183, a sigma receptor ligand, as a potential PET probe for the imaging of prostate adenocarcinoma", *Bioorganic & Medicinal Chemistry Letters*, 18, (2008), 1990-1993.
- Cunha, A. C., et al., "Tissue-specificity of prostate specific antigens: Comparative analysis of transcript levels in prostate and non-prostatic tissues", *Cancer Letters* 236, (2006), 229-238.
- Dahl, M., et al., "Sarcosine induces increase in HER2/neu expression in androgen-dependent prostate cancer cells", *Mol Biol Rep*, 38(7), (2011), 4237-4243.
- De Santis, M., et al., "Rolle der Chemotherapie beim kastrationsresistenten Prostatakarzinom. Gibt es neue Ansätze? [Role of chemotherapy in castration-resistant prostate cancer. Are there new approaches?]", (w/ English Abstract), *Urologe*, 51(1), (2012), 39-43 (5 pgs.).
- Degrado, T. R., et al., "Synthesis and Evaluation of 18F-Labeled Choline Analogs as Oncologic PET Tracers", *J Nucl Med*, 42(12), (2001), 1805-1814.
- Degrado, T. R., et al., "Synthesis and Evaluation of 18F-labeled Choline as an Oncologic Tracer for Positron Emission Tomography: Initial Findings in Prostate Cancer", *Cancer Research* 61(1), (2001), 110-117.
- Dimitrakopoulou-Strauss, A., et al., "PET Imaging of Prostate Cancer with 11C-Acetate", *J. Nucl Med*, 44(4), (2003), 556-558.
- Dumas, F., et al., "Molecular Expression of PSMA mRNA and Protein in Primary Renal Tumors", *Int. J. Cancer*, 80(6), (1999), 799-803.
- Eder, M., et al., "Novel Preclinical and Radiopharmaceutical Aspects of [68Ga]Ga-PSMA-HBED-CC: A New PET Tracer for Imaging of Prostate Cancer", *Pharmaceuticals*, 7(7), (2014), 779-796.
- Eder, M., et al., "Pharmacokinetic Properties of Peptidic Radiopharmaceuticals: Reduced Uptake of (EH)3-Conjugates in Important Organs", *J. Nucl Med*, 54(6), (2013), 1327-1330.
- Eder, M., et al., "Preclinical Evaluation of a Bispecific Low-Molecular Heterodimer Targeting Both PSMA and GRPR for Improved PET Imaging and Therapy of Prostate Cancer", *The Prostate*, 74, (2014), 659-668.
- Eder, M., et al., "PSMA as a target for radiolabelled small molecules", *Eur J Nucl Med Mol Imaging* 40, (2013), 819-823.
- Eiber, M., et al., "68Ga-PSMA PET/MR with multimodality image analysis for primary prostate cancer", *Abdom Imaging*, 40(6), (2015), 1769-1771.
- Elsasser-Beile, U., et al., "A New Generation of Monoclonal and Recombinant Antibodies Against Cell-Adherent Prostate Specific Membrane Antigen for Diagnostic and Therapeutic Targeting of Prostate Cancer", *The Prostate*, 66, (2006), 1359-1370.
- Elsasser-Beile, U., et al., "PET Imaging of Prostate Cancer Xenografts with a Highly Specific Antibody against the Prostate-Specific Membrane Antigen", *J Nucl Med* 50(4), (2009), 606-611.
- Elsasser-Beile, U., et al., "Targeted Therapies for Prostate Cancer Against the Prostate Specific Membrane Antigen", *Current Drug Targets*, 10(2), (2009), 118-125.
- El-Zaria, M. E., et al., "Preparation and evaluation of carborane-derived inhibitors of prostate specific membrane antigen (PSMA)", *Dalton Trans.*, 43(13), (2014), 4950-4961.
- Emonds, K. M., et al., "Do androgens control the uptake of 18F-FDG, 11C-choline and 11C-acetate in human prostate cancer cell lines?", *Eur J Nucl Med Mol Imaging*, 38, (2011), 1842-1853.
- Evans, M. J., et al., "Noninvasive measurement of androgen receptor signaling with a positron-emitting radiopharmaceutical that targets prostate-specific membrane antigen", *Proc. Natl. Acad. Sci. USA*, 108(23), (Jun. 7, 2011), 9578-9582.
- Fair, W. R., et al., "Prostate-Specific Membrane Antigen", *The Prostate*, 32, (1997), 140-148.
- Fall, K., et al., "Prostate-Specific Antigen Levels as a Predictor of Lethal Prostate Cancer", *J Natl Cancer Inst*, 99(7), (2007), 526-532.
- Fortmuller, K., et al., "Effective Targeting of Prostate Cancer by Lymphocytes Redirected by a PSMA x CD3 Bispecific Single-Chain Diabody", *The Prostate*, 71, (2011), 588-596.
- Fortuin, A. S., et al., "Value of PET/CT and MR Lymphography in Treatment of Prostate Cancer Patients With Lymph Node Metastases", *Int J Radiation Oncol Biol Phys*, 84(3), (2012), 712-718.
- Foss, C. A., et al., "GCPII Imaging and Cancer", *Current Medicinal Chemistry*, 19(9), (2012), 1346-1359.
- Franc, B. L., et al., "Detection and localization of carcinoma within the prostate using high resolution transrectal gamma imaging (TRGI) of monoclonal antibody directed at prostate specific membrane antigen (PSMA)—Proof of concept and initial imaging results", *European Journal of Radiology*, 82(11), (2013), 1877-1884.
- Frigerio, B., et al., "A single-chain fragment against prostate specific membrane antigen as a tool to build theranostic reagents for prostate cancer", *European Journal of Cancer*, 49(9), (2013), 2223-2232.
- Ghosh, A., et al., "Tumor Target Prostate Specific Membrane Antigen (PSMA) and its Regulation in Prostate Cancer", *Journal of Cellular Biochemistry*, 91(3), (2004), 528-539.
- Giovacchini, et al., "Predictive factors of [11 C]choline PET/CT in patients with biochemical failure after radical prostatectomy", *Eur J Nucl Med Mol Imaging*, 37, (2010), 301-309.
- Goodman Jr., et al., "Interaction of prostate specific membrane antigen with clathrin and the adaptor protein complex-2", *International Journal of Oncology*, 31(5), (2007), 1199-1203.
- Graham, et al., "Radiofluorinated derivatives of 2-(phosphonomethyl)pentanedioic acid as inhibitors of prostate specific membrane antigen (PSMA) for the imaging of prostate cancer", *J. Med. Chem.*, 55, (2012), 9510-9520.
- Grant, et al., "Prostate Specific Membrane Antigen (PSMA) Regulates Angiogenesis Independently of VEGF during Ocular Neovascularization", *PLoS ONE* 7(7): e41285., (2012), 1-10.
- Gregor, et al., "Induction of autoantibodies to syngeneic prostate-specific membrane antigen by xenogeneic vaccination", *Int. J. Cancer*, 116(3), (2005), 415-421.
- Haberkorn, et al., "Mechanistic and high-throughput approaches for the design of molecular imaging probes and targeted therapeutics", *Clin Transl Imaging*, 2(1), (2014), 33-41.

(56)

References Cited

OTHER PUBLICATIONS

- Haffner, M. C., et al., "Prostate-specific membrane antigen expression in the neovasculature of gastric and colorectal cancers", *Human Pathology*, 40(12), (2009), 1754-1761.
- Hain, S. F., et al., "Positron emission tomography for urological tumours", *BJU International*, 92(2), (2003), 159-164.
- Hara, T., "11C-Choline and 2-Deoxy-2-[18F]Fluoro-D-Glucose in Tumor Imaging with Positron Emission Tomography", *Molecular Imaging and Biology*, 4(4), (2002), 267-273.
- Hara, T., et al., "Development of 18F-Fluoroethylcholine for Cancer Imaging with PET: Synthesis, Biochemistry, and Prostate Cancer Imaging", *J Nucl Med*, 43(2), (2002), 187-199.
- Hara, T., et al., "PET Imaging of Prostate Cancer Using Carbon-11-Choline", *J Nucl Med*, 39(6), (1998), 990-995.
- Harada, N., et al., "Preparation of Asymmetric Urea Derivatives that Target Prostate-Specific Membrane Antigen for SPECT Imaging", *J. Med. Chem.* 56, (2013), 7890-7901.
- Heidenreich, A., "Immuntherapie beim metastasierten Prostatairzinom brauchen wir diese wirklich? [Immunotherapy for metastatic prostate cancer: do we really need this?]", (w/ English Abstract), *Urologe*, 51(1), (2012), 32-38 (8 pgs.).
- Henry, M. D., et al., "A Prostate-Specific Membrane Antigen-Targeted Monoclonal Antibody-Chemotherapeutic Conjugate Designed for the Treatment of Prostate Cancer", *Cancer Research*, 64(21), (2004), 7995-8001.
- Hillier, S., et al., "[131 I] MIP-1466, a small molecule prostate-specific membrane antigen (PSMA) inhibitor for targeted radiotherapy of prostate cancer (PCa)", *J Nucl Med*, vol. 53 No. supplement 1 170, (May 2012), 1 pg.
- Hillier, S. M., et al., "123I-MIP-1072, a Small-Molecule Inhibitor of Prostate-Specific Membrane Antigen, Is Effective at Monitoring Tumor Response to Taxane Therapy", *J Nucl Med*, 52(7), (2011), 1087-1093.
- Hillier, S. M., et al., "99mTc-Labeled Small-Molecule Inhibitors of Prostate-Specific Membrane Antigen for Molecular Imaging of Prostate Cancer", *J Nucl Med*, 54, (2013), 1369-1376.
- Hlouchova, K., et al., "Biochemical characterization of human glutamate carboxypeptidase III", *Journal of Neurochemistry*, 101(3), (2007), 682-696.
- Hlouchova, K., et al., "GCP II Variants, Paralogs and Orthologs", *Current Medicinal Chemistry*, 19, (2012), 1316-1322.
- Hlouchova, K., et al., "Structural insight into the evolutionary and pharmacologic homology of glutamate carboxypeptidases II and III", *FEBS Journal*, 276(16), (2009), 4448-4462.
- Ho, C.-L., et al., "Molecular Imaging, Pharmacokinetics, and Dosimetry of 111In-AMBA in Human Prostate Tumor-Bearing Mice", *Journal of Biomedicine and Biotechnology* vol. 2011, Article ID 101497, (2011), 8 pgs.
- Holland, J. P., et al., "89Zr-DFO-J591 for ImmunoPET of Prostate-Specific Membrane Antigen Expression In Vivo", *J Nucl Med*, 51(8), (2010), 1293-1300.
- Hong, et al., "Positron emission tomography imaging of prostate cancer", *Amino Acids*, 39(1), (2010), 11-27.
- Hospers, G. A. P., et al., "PET Imaging of Steroid Receptor Expression in Breast and Prostate Cancer", *Current Pharmaceutical Design*, 14(28), (2008), 3020-3032.
- Huang, et al., "Improving the Biodistribution of PSMA-Targeting Tracers With Highly Negatively Charged Linker", *The Prostate*, 74, (2014), 702-713.
- Huang, et al., "PSMA-Targeted Stably Linked 'Dendrimer-Glutamate Urea-Methotrexate' as a Prostate Cancer Therapeutic", *Biomacromolecules*, 15(3), (2014), 915-923.
- Humblet, et al., "High-affinity Near-infrared Fluorescent Small-molecule Contrast Agents for In Vivo Imaging of Prostate-specific Membrane Antigen", *Molecular Imaging*, 4(4), (Oct. 2005), 448-462.
- Humblet, et al., "Multivalent Scaffolds for Affinity Maturation of Small Molecule Cell Surface Binders and Their Application to Prostate Tumor Targeting", *J. Med. Chem.* 52, (2009), 544-550.
- Husarik, et al., "Evaluation of [18F]-choline PET/CT for staging and restaging of prostate cancer", *Eur J Nucl Med Mol Imaging*, 35, (2008), 253-263.
- Hwang, et al., "N-3-[18F]Fluoropropylputrescine as Potential PET Imaging Agent for prostate and Prostate Derived Tumors", *J Nucl Med*, 30(7), (1989), 1205-1210.
- Hwang, D.-R., et al., "Imaging Prostate Derived Tumors with PET and N-(3-[18F]Fluoropropyl)putrescine", *Nucl. Med. Biol.*, 17(6), (1990), 525-532.
- Igerc, I., et al., "The value of 18F-Choline PET/CT in patients with elevated PSA-level and negative prostate needle biopsy for localisation of prostate cancer", *Eur J Nucl Med Mol Imaging*, 35, (2008), 976-983.
- Jackson, P. F., et al., "Design, Synthesis, and Biological Activity of a Potent Inhibitor of the Neuropeptidase N-Acetylated a-Linked Acidic Dipeptidase", *Journal of Medicinal Chemistry*, 39, (1996), 619-622.
- Jadvar, et al., "Glucose Metabolism of Human Prostate Cancer Mouse Xenografts", *Molecular Imaging*, 4(2), (Apr. 2005), 91-97.
- Jadvar, et al., "Imaging evaluation of prostate cancer with 18F-fluorodeoxyglucose PET/CT: utility and limitations", *Eur J Nucl Med Mol Imaging* < 40 (Suppl 1), (2013), S5-S10.
- Jadvar, et al., "Molecular imaging of prostate cancer with 18F-fluorodeoxyglucose PET", *Nat. Rev. Urol.*, 6, (2009), 317-323.
- Jadvar, et al., "Molecular Imaging of Prostate Cancer: PET Radiotracers", *AJR Am J Roentgenol*, 199(2), (2012), 278-291.
- Jambor, et al., "Functional Imaging of Localized Prostate Cancer Aggressiveness Using 11C-Acetate PET/CT and 1H-MR Spectroscopy", *J Nucl Med*; 51., (2010), 1676-1683.
- James, S., "Urea based rhenium tricarbonyl dipeptide compounds as potential radiopharmaceuticals for PSMA imaging", INOR258, The 229th ACS National Meeting, San Diego, CA, (2005), 1 pg.
- Kahn, D., et al., "111Indium-CapromabPendetide in the Evaluation of Patients with Residual or Recurrent Prostate Cancer After Radical Prostatectomy", *The Journal of Urology*, 159(6), (Jun. 1998), 2041-2047.
- Kasperzyk, J. L., et al., "Prostate-Specific Membrane Antigen Protein Expression in Tumor Tissue and Risk of Lethal Prostate Cancer", *Cancer Epidemiol Biomarkers Prev*; 22(12), (2013), 2354-2363.
- Kasten, et al., "Targeting prostate cancer cells with PSMA inhibitor-guided gold nanoparticles", *Bioorganic & Medicinal Chemistry Letters* 23, (2013), 565-568.
- Kim, D., et al., "Tribody: Robust Self-Assembled Trimeric Targeting Ligands with High Stability and Significantly Improved Target-Binding Strength", *Biochemistry*, 52, (2013), 7283-7294.
- Kinoshita, Y., et al., "Expression of Prostate-Specific Membrane Antigen in Normal and Malignant Human Tissues", *World J Surg*, 30, (2006), 628-636.
- Klotz, Laurence, "Cancer overdiagnosis and overtreatment", *Curr Opin Urol.*, 22, (2012), 203-209.
- Klusak, et al., "Reaction Mechanism of Glutamate Carboxypeptidase II Revealed by Mutagenesis, X-ray Crystallography, and Computational Methods", *Biochemistry*, 48, (2009), 4126-4138.
- Kosuri, et al., "Review of Salvage Therapy for Biochemically Recurrent Prostate Cancer: The Role of Imaging and Rationale for Systemic Salvage Targeted Anti-Prostate-Specific Membrane Antigen Radioimmunotherapy", *Advances in Urology* vol. 2012, Article ID 921674, (2012), 8 pgs.
- Kothari, P., et al., "18F-labeled small molecule inhibitors of prostate specific membrane antigen (PSMA) for PET imaging of prostate cancer", *J Nucl Med*, vol. 53 No. supplement 11721, (May 2012), 2 pgs.
- Kotzerke, et al., "PET or Prostate Cancer Imaging: Still a Quandary or the Ultimate Solution?", *The Journal of Nuclear Medicine*, 43(2), (Feb. 2002), 200-202.
- Kovar, et al., "Pharmacokinetic and Biodistribution Assessment of a Near Infrared-Labeled PSMA-Specific Small Molecule in Tumor-Bearing Mice", *Prostate Cancer* vol. 2014, Article ID 104248, (2014), 10 pgs.
- Krohn, et al., "[68Ga]PSMA-HBED uptake mimicking lymph node metastasis in coeliac ganglia: an important pitfall in clinical practice", *Eur J Nucl Med Mol Imaging*, 42, (2015), 210-214.

(56)

References Cited

OTHER PUBLICATIONS

- Kularatne, et al., "Design, Synthesis, and Preclinical Evaluation of Prostate-Specific Membrane Antigen Targeted ^{99m}Tc -Radioimaging Agents", *Molecular Pharmaceuticals*, 6(3), (2009), 790-800.
- Kularatne, S. A., et al., "Comparative Analysis of Folate Derived PET Imaging Agents with [18F]-2-Fluoro-2-deoxy-D-glucose Using Rodent Inflammatory Paw Model", *Molecular Pharmaceuticals*, vol. 10, (2013), 3103-3111.
- Kuru, T. H., et al., "MRT-navigierte stereotaktische Prostatabiopsie [MRI-navigated stereotactic prostate biopsy]", (w/English Abstract), *Urologe* 51(1), (2012), 50-56 (8 pgs.).
- Kwee, et al., "18F-choline PET/CT imaging of RECIST measurable lesions in hormone refractory prostate cancer", *Ann Nucl Med* 23(6), (2009), 541-548.
- Lambert, et al., "Molecular Evolution of the Transferrin Receptor/ Glutamate Carboxypeptidase II Family", *J Mol Evol.* 64(1), (2007), 113-128.
- Lapi, S. E., et al., "Assessment of an 18F-Labeled Phosphoramidate Peptidomimetic as a New Prostate-Specific Membrane Antigen-Targeted Imaging Agent for Prostate Cancer", *J Nucl Med*; 50, (2009), 2042-2048.
- Leek, et al., "Prostate-specific membrane antigen: evidence for the existence of a second related human gene", *British Journal of Cancer*, 72(3), (1995), 583-588.
- Lees, et al., "Active surveillance in prostate cancer: patient selection and triggers for intervention", *Curr Opin Urol*, 22(3), (2012), 210-215.
- Lesche, et al., "Preclinical evaluation of BAY 1075553, a novel 18F-labelled inhibitor of prostate-specific membrane antigen for PET imaging of prostate cancer", *Enr J Nucl Med Mol Imaging*, 41, (2014), 89-101.
- Li, X., et al., "C-11 Choline PET/CT Imaging for Differentiating Malignant From Benign Prostate Lesions", *Clin Nucl Med*, 33(10), (2008), 671-676.
- Liu, HE, et al., "Constitutive and Antibody-Induced Internalization of Prostate-Specific Membrane Antigen", *Cancer Research*, 58, (Sep. 15, 1998), 4055-4060.
- Liu, M., et al., "Synthesis and Biological Evaluation of Diethylenetriamine Pentaacetic acid-Polyethylene Glycol Folate: A new Folate-Derived, ^{99m}Tc -Based Radiopharmaceutical", *Bioconjugate Chem*, 16(5), (2005), 1126-1132.
- Liu, T., et al., "A targeted low molecular weight near-infrared fluorescent probe for prostate cancer", *Bioorganic & Medicinal Chemistry Letters*, 20(23), (2010), 7124-7126.
- Liu, T., et al., "Functional prostate-specific membrane antigen is enriched in exosomes from prostate cancer cells", *International Journal of Oncology*, 44, (2014), 918-922.
- Liu, T., et al., "Prolonged androgen deprivation leads to down regulation of androgen receptor and prostate-specific membrane antigen in prostate cancer cells", *International Journal of Oncology*, 41, (2012), 2087-2092.
- Liu, T., et al., "Pseudoirreversible Inhibition of Prostate-Specific Membrane Antigen by Phosphoramidate Peptidomimetics", *Biochemistry*, 47(48), (2008), 12658-12660.
- Liu, T., et al., "Targeting prostate cancer cells with a multivalent PSMA inhibitor-guided streptavidin conjugate", *Bioorganic & Medicinal Chemistry Letters* 22, (2012), 3931-3934.
- Lord, M., et al., "18F-Fluorocholine integrated PET/MRI for the initial staging of prostate cancer", *Eur J Nucl Med Mol Imaging*, 38(12), (2011), p. 2288.
- Luboldt, W., et al., "Prostate Carcinoma: Diffusion-weighted Imaging as Potential Alternative to Conventional MR and 11C-Choline PET/CT for Detection of Bone Metastases", *Radiology*, 249(3), (2008), 1017_1025.
- Lutje, S., et al., "Dual-Modality Image-Guided Surgery of Prostate Cancer with a Radiolabeled Fluorescent Anti-PSMA Monoclonal Antibody", *J Nucl Med*, 55(6), (2014), 995-1001.
- Lutje, S., et al., "Prospects in Radionuclide Imaging of Prostate Cancer", *The Prostate*, 72, (2012), 1262-1272.
- Malik, N., et al., "One pot radiofluorination of a new potential PSMA ligand [A118F]NOTA-DUPA-Pep", *J. Label Compd. Radiopharm.* 55(2), (2012), 320-325.
- Malik, N., et al., "Radiosynthesis of a new PSMA targeting ligand ([18F]FPy-DU PA-Pep)", *Applied Radiation and Isotopes*, 69(7), (2011), 1014-1018.
- Mannweiler, S., et al., "Heterogeneity of Prostate-Specific Membrane Antigen (PSMA) Expression in Prostate Carcinoma with Distant Metastasis", *Pathol. Oncol. Res.*, 15(2), (2009), 167-172.
- Maresca, K., et al., "Influence of functionalized chelators on affinity and pharmacokinetics of $^{99m}\text{Tc}(\text{CO})_3$ -labeled small molecules targeting prostate specific membrane antigen (PSMA)", (Abstract), *J Nucl Med*, vol. 51, No. Supplement 2, (May 2010), p. 250 (1 page).
- Matthies, A., et al., "Imaging of prostate cancer metastases with 18F-fluoroacetate using PET/CT", *Eur J Nucl Med Mol Imaging*, 31(5), (May 2004), p. 797.
- Mease, R. C., et al., "PET Imaging in Prostate Cancer: Focus on Prostate-Specific Membrane Antigen", *Current Topics in Medicinal Chemistry*, 13, (2013), 951-962.
- Meighan, M. A., et al., "Recombinant Glutamate Carboxypeptidase II (Prostate Specific Membrane Antigen-PSMA)—Cellular Localization and Bioactivity Analyses", *Journal of Protein Chemistry*, 22(4), (May 2003), 317-326.
- Meinhardt, W., et al., "Laparoscopic Sentinel Lymph Node Biopsy for Prostate Cancer: The Relevance of Locations Outside the Extended Dissection Area", *Prostate Cancer*, vol. 2012, Article ID 751753, (2012), 4 pgs.
- Mertens, K., et al., "PET with 18F-labelled choline-based tracers for tumour imaging: a review of the literature", *Eur J Nucl Med Mol Imaging*, 37, (2010), 2188-2193.
- Mhaweche-Fauceglia, P., et al., "Prostate-specific membrane antigen (PSMA) protein expression in normal and neoplastic tissues and its sensitivity and specificity in prostate adenocarcinoma: an immunohistochemical study using multiple tumour tissue microarray technique", *Histopathology*, 50(4), (2007), 472-483.
- Milowsky, M.I., et al., "Phase I Trial of Yttrium-90-Labeled Anti-Prostate-Specific Membrane Antigen Monoclonal Antibody J591 for Androgen-Independent Prostate Cancer", *J Clin Oncol*, 22(13), (2004), 2522-2531.
- Minner, S., et al., "High Level PSMA Expression Is Associated With Early PSA Recurrence in Surgically Treated Prostate Cancer", *The Prostate*, 71, (2011), 281-288.
- Mlcochova, P., et al., "Mapping of the active site of glutamate carboxypeptidase II by site-directed mutagenesis", *FEBS Journal*, 274(18), (2007), 4731-4741.
- Moltzahn, F., et al., "Die ossare Metastasierung des Prostatakarzinoms [Bony metastasis of prostate cancer]", (w/ English Abstract), *Urologe*, 51, (2012), 20-26 (8 pgs.).
- Morris, M. J., et al., "11C-acetate PET imaging in prostate cancer", *Eur J Nucl Med Mol Imaging*, 34, (2007), 181-184.
- Muller, C., et al., "Synthesis and in Vitro/in Vivo Evaluation of Novel $^{99m}\text{Tc}(\text{CO})_3$ -Folates", *Bioconjugate Chem*, 17(3), (2006), 797-806.
- Murphy, G. P., et al., "Current Evaluation of the Tissue Localization and Diagnostic Utility of Prostate Specific Membrane Antigen", *Cancer*, 83(11), (1998), 2259-2269.
- Nedrow-Byers, J. R., et al., "A Phosphoramidate-Based Prostate-Specific Membrane Antigen-Targeted SPECT Agent", *The Prostate*, 72, (2012), 904-912.
- Oehr, P., et al., "Imaging of prostate cancer", *Curr Opin Oncol*, 19, (2007), 259-264.
- O'Keefe, D. S., et al., "Comparative Analysis of Prostate-Specific Membrane Antigen (PSMA) Versus a Prostate-Specific Membrane Antigen-Like Gene", *The Prostate*, 58:, (2004), 200-210.
- Omlin, A., et al., "Androgen-und Oestrogen-biosynthesehemmer beim kastrationsresistenten Prostatakarzinom [Inhibitors of androgen and estrogen biosynthesis in castration-resistant prostate cancer]", (w/ English Abstract), *Urologe*, 51, (2012), 8-14 (8 pgs.).
- Osborne, J. R., et al., "A Prospective Pilot Study of 89Zr-J591/Prostate Specific Membrane Antigen Positron Emission Tomogra-

(56)

References Cited

OTHER PUBLICATIONS

- phy in Men with Localized Prostate Cancer Undergoing Radical Prostatectomy”, *The Journal of Urology*, 191(5), (May 2014), 1439-1445.
- Oyama, et al., “PET Imaging in Prostate Cancer”, *Hinyokika Kiyo*, 52, (2006), 503-505.
- Oyama, N., et al., “¹¹C-Acetate PET Imaging of Prostate Cancer”, *J Nucl Med*, 4 2 , (2002), 181-186.
- Oyama, N., et al., “¹¹C-Acetate PET Imaging of Prostate Cancer: Detection of Recurrent Disease at PSA Relapse”, *J Nucl Med*, 44(4), (2003), 549-555.
- Panchuk-Voloshina, Nataliya, et al., “Alexa dyes, a series of new fluorescent dyes that yield exceptionally bright, photostable conjugates.”, *J Histochem Cytochem.*, 47(9), (Sep. 1999), 1179-88.
- Parker, et al., “Design, production, and characterization of a single-chain variable fragment (ScFv) derived from the prostate specific membrane antigen (PSMA) monoclonal antibody J591”, *Protein Expression and Purification*, 89(2), (2013), 136-145.
- Pavlicek, et al., “Glutamate Carboxypeptidase II: An Overview of Structural Studies and Their Importance for Structure-Based Drug Design and Deciphering the Reaction Mechanism of the Enzyme”, *Current Medicinal Chemistry*, 19(9), (2012), 1300-1309.
- Pavlicek, et al., “Structural characterization of P1'-diversified urea-based inhibitors of glutamate carboxypeptidase II”, *Bioorganic & Medicinal Chemistry Letters* 24, (2014), 2340-2345.
- Perner, et al., “Prostate-specific membrane antigen expression as a predictor of prostate cancer progression”, *Human Pathology* 38, (2007), 696-701.
- Pillarsetty, et al., “²-18F-Fluoropropionic Acid as a PET Imaging Agent for Prostate Cancer”, *J Nucl Med*, 50(10), (2009), 1709-1714.
- Pinto, F., et al., “Imaging in Prostate Cancer Staging: Present Role and Future Perspectives”, *Urol Int*; 88(2), (2012), 125-136.
- Ponde, D. E., et al., “¹⁸F-Fluoroacetate: A Potential Acetate Analog for Prostate Tumor Imaging—In Vivo Evaluation of ¹⁸F-Fluoroacetate Versus ¹¹C-Acetate”, *J Nucl Med*, 48(3), (2007), 420-428.
- Poulsen, et al., “[¹⁸F] fluoromethylcholine (FCH) positron emission tomography/computed tomography (PET/CT) for lymph node staging of prostate cancer: a prospective study of 210 patients”, *BJU International*, 110, (2012), 1666-1671.
- Poulsen, M. H., et al., “[¹⁸F]-fluorocholine positron-emission/computed tomography for lymph node staging of patients with prostate cancer: preliminary results of a prospective study”, *BJU International*, 106, (2010), 639-644.
- Preusser, S. et al., “Kastrationsresistentes Prostatakarzinom [Castration-resistant prostate cancer]”, (w/ English Abstract), *Urologe* 51(1), (2012), 27-31 (6 pgs.).
- Rais, R., et al., “Bioanalytical method for evaluating the pharmacokinetics of the GCP-II inhibitor 2-phosphonomethylpentanedioic acid (2-PMPA)”, *Journal of Pharmaceutical and Biomedical Analysis* 88, (2014), 162-169.
- Rajasekaran, et al., “A Novel Cytoplasmic Tail MXXXL Motif Mediates the Internalization of Prostate specific Membrane Antigen”, *Molecular Biology of the Cell*, vol. 14, (Dec. 2003), 4835-4845.
- Reske, et al., “[¹¹C]choline PET/CT imaging in occult local relapse of prostate cancer after radical prostatectomy”, *Eur J Nucl Med Mol Imaging* 35:, (2008), 9-17.
- Reske, et al., “[¹¹C]Choline uptake with PET/CT for the initial diagnosis of prostate cancer: relation to PSA levels, tumour stage and anti-androgenic therapy”, *Eur J Nucl Med Mol Imaging* 35(6), (2008), 1740-1741.
- Reske, S. N., et al., “Nukleomedizinische Diagnostik beim Prostatakarzinom [Nuclear imaging of prostate cancer]” (w/ English Abstract), *Urology* 46(11), (2007), 1485-1499 (16 pgs.).
- Reske, S. N., et al., “PET and PET/CT in der Rezidivdiagnostik des Prostatakarzinoms [PET and PET / CT in the recurrence diagnosis of prostate cancer]”, (w/ English Summary), *Urologe* 45(10), (2006), 1240-1250 (12 pgs.).
- Reske, S. N., et al., “Weiterentwicklung der PET und des PET/CT beim Prostatakarzinom [Advancement of PET and PET/CT in prostate carcinoma]”, (w/ English Abstract), *Urologe* 45, (2006), 707-714 (9 pgs.).
- Rinnab, et al., “[¹¹C]Choline PET/CT for Targeted Salvage Lymph Node Dissection in Patients with Biochemical Recurrence after Primary Curative Therapy for Prostate Cancer”, *Urol Int*; 81, (2008), 191-197.
- Rinnab, et al., “[¹¹C]choline PET/CT in prostate cancer patients with biochemical recurrence after radical prostatectomy”, *World J Urol* 27, (2009), 619-625.
- Rioja, et al., “Role of positron emission tomography in urological oncology”, *BJU International*, 106, 1578-1594.
- Ristau, et al., “The prostate-specific membrane antigen: Lessons and current clinical implications from 20 years of research”, *Urologic Oncology: Seminars and Original Investigations* 32, (2014), 272-279.
- Roethke, et al., “Hybrid Positron Emission Tomography-Magnetic Resonance Imaging with Gallium 68 Prostate-specific Membrane Antigen Tracer: A Next Step for Imaging of Recurrent Prostate Cancer-Preliminary Results”, (2013), 862-864.
- Rybalov, et al., “Impact of total PSA, PSA doubling time and PSA velocity on detection rates of ¹¹C-Choline positron emission tomography in recurrent prostate cancer”, *World J Urol* 31, (2013), 319-323.
- Sacha, et al., “Expression of Glutamate Carboxypeptidase II in Human Brain”, *Neuroscience* 144, (2007), 1361-1372.
- Scattoni, V., et al., “Detection of Lymph-Node Metastases with Integrated [¹¹C]Choline PET/CT in Patients with PSA Failure after Radical Retropubic Prostatectomy: Results Confirmed by Open Pelvic-Retroperitoneal Lymphadenectomy”, *European Urology*, 52, (2007), 423-429.
- Scheffel, et al., “PET Imaging of GRP Receptor Expression in Prostate Cancer”, *J. Nucl Med*, 45(8), (Aug. 2004), 1277-1278.
- Scher, et al., “PET/CT imaging of recurrent prostate cancer”, *Enr J Nucl Med Mol Imaging* 35:, (2008), 5-8.
- Shvarts, O., et al., “Positron Emission Tomography in Urologic Oncology”, *Cancer Control*, vol. 9, No. 4, (2002), 335-342.
- Silver, D. A., et al., “Prostate-specific Membrane Antigen Expression in Normal and Malignant Human Tissues”, *Clinical Cancer Research*, 3, (Jan. 1997), 81-85.
- Simone, et al., “What’s in a Label? Radioimmunotherapy for Metastatic Prostate Cancer”, *Clin Cancer Res*; 19(18), (2013), 4908-4910.
- Slusher, et al., “Immunocytochemical Localization of the N-Acetyl-Aspartyl-Glutamate (NAAG) Hydrolyzing Enzyme N-Acetylated a-Linked Acidic Dipeptidase (NAALADase)”, *The Journal of Comparative Neurology* 315, (1992), 217-229.
- Slusher, B. S., et al., “Selective inhibition of NAALADase, which converts NAAG to glutamate, reduces ischemic brain injury”, *Nature Medicine*, 5(12), (Dec. 1999), 1396-1400.
- Soloviev, D., et al., “PET imaging with ¹¹C-acetate in prostate cancer: a biochemical, radiochemical and clinical perspective”, *Eur J Nucl Med Mol Imaging* 35(5), (2008), 942-949.
- Spahn, M., et al., “We soil die Hormontherapie beim kastrationsresistenten Prostatakarzinom fortgeführt werden? [How should hormone therapy for castration-resistant prostate cancer be continued?]”, (w/ English Abstract), *Urologe* 51, (2012), 15-19 (6 pgs.).
- Sweat, S. D., et al., “Prostate-Specific Membrane Antigen Expression Is Greatest In Prostate Adenocarcinoma and Lymph Node Metastases”, *Urology* 52, (1998), 637-640.
- Tang, H., et al., “Prostate targeting ligands based on N-acetylated a-linked acidic dipeptidase”, *Biochemical and Biophysical Research Communications* 307, (2003), 8-14.
- Tang, Q.-I., et al., “Updated Application of Prostate-Specific Membrane Antigen to the Diagnosis and Treatment of Prostate Cancer”, (w/ English Abstract), *National Journal of Andrology*, vol. 14, No. 1, (Jan. 2008), 79-82.
- Taylor, R. M., et al., “Prostate Cancer Targeting Motifs: Expression of an³, Neurotensin Receptor1, Prostate Specific Membrane Antigen, and Prostate Stem Cell Antigen in Human Prostate Cancer Cell Lines and Xenografts”, *The Prostate* 72, (2012), 523-532.

(56)

References Cited

OTHER PUBLICATIONS

- Testa, C., et al., "Prostate Cancer: Sextant Localization with MR Imaging, MR Spectroscopy, and 11C-Choline PET/CT", *Radiology*, 244(3), (Sep. 2007), 797-806.
- Thalmann G., "Fortgeschrittenes Prostatakarzinom [Advanced prostate cancer]", *Urologe* 51(1), (2012), p. 7 (2 pgs.).
- Tykvtart, J., et al., "Rational design of urea-based glutamate carboxypeptidase II (GCPII) inhibitors as versatile tools for specific drug targeting and delivery", *Bioorganic & Medicinal Chemistry*, 22, (2014), 4099-4108.
- Uprimny, C., et al., "68Ga-PSMA ligand PET versus 18F-NaF PET: evaluation of response to 223Ra therapy in a prostate cancer patient", *Eur J Nucl Med Mol Imaging* 42(2), (2015), 362-363.
- Vallabhajosula, S., et al., "Radioimmunotherapy of Prostate Cancer in Human Xenografts Using Monoclonal Antibodies Specific to Prostate Specific Membrane Antigen (PSMA): Studies in Nude Mice", *The Prostate*, 58, (2004), 145-155.
- Vavere, A. L., et al., "1-11C-Acetate as a PET Radiopharmaceutical for Imaging Fatty Acid Synthase Expression in Prostate Cancer", *J Nucl Med*; 49(2), (2008), 327-334.
- Viola-Villegas, N., et al., "Targeting Gallium to Cancer Cells through the Folate Receptor", *Drug Target Insights*, vol. 3, (2008), 13-25.
- Viola-Villegas, N., et al., "Targeting the Folate Receptor (FR): Imaging and Cytotoxicity of Re1 Conjugates in FR-Overexpressing Cancer Cells", *ChemMedChem*, vol. 3, (2008), 1387-1394.
- Wang, H., et al., "Bioisosterism of urea-based GCPII inhibitors: Synthesis and structure-activity relationship studies", *Bioorganic & Medicinal Chemistry Letters* 20, (2010), 392-397.
- Wang, X., et al., "Development of Targeted Near-Infrared Imaging Agents for Prostate Cancer", *Mol Cancer Ther*, 13(11), (2014), 2595-2606.
- Weinisen, M., et al., "Synthesis and preclinical evaluation of DOTAGA-conjugated PSMA ligands for functional imaging and endoradiotherapy of prostate cancer", *EJNMMI Research*, 4:63, (2014), 15 pgs.
- Weissbach, L., "Welche Inhalte sollte eine „living guideline besetzen? [Which components should living guidelines contain?]", (w/ English Abstract), *Urologe*, 51, (2012), 57-59 (5 pgs.).
- Whitaker, H. C., et al., "N-acetyl-L-aspartyl-L-glutamate peptidase-like 2 is overexpressed in cancer and promotes a pro-migratory and pro-metastatic phenotype", *Oncogene* 33, (2014), 5274-5287.
- Wiehr, S., et al., "Pharmacokinetics and PET Imaging Properties of Two Recombinant Anti-PSMA Antibody Fragments in Comparison to their Parental Antibody", *The Prostate*, 74, (2014), 743-755.
- Wright, G. L., et al., "Expression of Prostate-Specific Membrane Antigen in Normal, Benign, and Malignant Prostate Tissues", *Urol Oncol*, 1(1), (1995), 18-28.
- Wu, L. Y., et al., "The molecular pruning of a phosphoramidate peptidomimetic inhibitor of prostate-specific membrane antigen", *Bioorganic & Medicinal Chemistry* 15(23), (2007), 7434-7443.
- Yamaguchi, T., et al., "Prostate cancer: a comparative study of 11C-choline PET and MR imaging combined with proton MR spectroscopy", *Eur J Nucl Med Mol Imaging*, 32, (2005), 742-748.
- Zaheer, A., et al., "New Agents and Techniques for Imaging Prostate Cancer", *J Nucl Med*, 50(9), (2009), 1387-1390.
- Zechmann, C. M., et al., "Radiation dosimetry and first therapy results with a 124I/131 I-labeled small molecule (MIP-1095) targeting PSMA for prostate cancer therapy", *Eur J Nucl Med Mol Imaging* 41(7), (2014), 1280-1292.
- Zhang, A. X., et al., "A Remote Arene-Binding Site on Prostate Specific Membrane Antigen Revealed by Antibody-Recruiting Small Molecules", *J. Am. Chem. Soc.*, 132(36), (2010), 12711-12716.
- Zhang, Y., et al., "Prostate Specific Membrane Antigen (PSMA): A Novel Modulator of p38 for Proliferation, Migration, and Survival in Prostate Cancer Cells", *The Prostate*, 73(8), (2013), 835-841.
- Zhou, J., "In vivo evaluation of medical device-associated inflammation using macrophage-specific position emission tomography (PET) imaging", *Bioorganic and Medicinal Chemistry Letters*, vol. 23, (2013), 2044-2047.

* cited by examiner




FIGURE 1A




FIGURE 1B




FIGURE 2




FIGURE 3A




FIGURE 3B




FIGURE 4




FIGURE 5A




FIGURE 5B




FIGURE 6A




FIGURE 6B




FIGURE 6C




FIGURE 6D




FIGURE 7A




FIGURE 7B




FIGURE 7C




FIGURE 7D




FIGURE 8A




FIGURE 8B




FIGURE 8C




FIGURE 9A




FIGURE 9B




FIGURE 10A




FIGURE 10B




FIGURE 10C




FIGURE 11




FIGURE 12A




FIGURE 12 B




FIGURE 12C




FIGURE 12D




FIGURE 13A




FIGURE 13B




FIGURE 13C




FIGURE 13D




FIGURE 14A




FIGURE 14B

PSMA BINDING LIGAND-LINKER CONJUGATES AND METHODS FOR USING

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 14/794,482, filed Jul. 8, 2015, which is a continuation of U.S. patent application Ser. No. 12/673,931, filed Feb. 17, 2010, now U.S. Pat. No. 9,193,763, issued Nov. 24, 2015, which is a U.S. national application under 37 C.F.R. § 371(b) of International Application Serial No. PCT/US2008/073375, filed Aug. 15, 2008, which claims priority under 35 U.S.C. § 119(e) to U.S. Provisional Application Ser. No. 60/956,489 filed Aug. 17, 2007, and U.S. Provisional Application Ser. No. 61/074,358, filed Jun. 20, 2008, in which all of which are expressly incorporated herein by reference.

TECHNICAL FIELD

The invention described herein pertains to compounds and methods for treating diseases of the prostate, such as prostate cancer and related diseases. More specifically, embodiments of the invention described herein pertain to conjugates of biologically active agents conjugated to PSMA binding ligands.

BACKGROUND

The prostate is one of the male reproductive organs found in the pelvis below the urinary bladder. It functions to produce and store seminal fluid which provides nutrients and fluids that are vital for the survival of sperm introduced into the vagina during reproduction. Like many other tissues, the prostate glands are also prone to develop either malignant (cancerous) or benign (non-cancerous) tumors. The American Cancer Society predicted that over 230,000 men would be diagnosed with prostate cancer and over 30,000 men would die from the disease in year 2005. In fact, prostate cancer is one of the most common male cancers in western societies, and is the second leading form of malignancy among American men. Current treatment methods for prostate cancer include hormonal therapy, radiation therapy, surgery, chemotherapy, photodynamic therapy, and combination therapy. The selection of a treatment generally varies depending on the stage of the cancer. However, many of these treatments affect the quality of life of the patient, especially those men who are diagnosed with prostate cancer over age 50. For example, the use of hormonal drugs is often accompanied by side effects such as osteoporosis and liver damage. Such side effects might be mitigated by the use of treatments that are more selective or specific to the tissue being responsible for the disease state, and avoid non-target tissues like the bones or the liver. As described herein, prostate specific membrane antigen (PSMA) represents a target for such selective or specific treatments.

PSMA is named largely due to its higher level of expression on prostate cancer cells; however, its particular function on prostate cancer cells remains unresolved. PSMA is over-expressed in the malignant prostate tissues when compared to other organs in the human body such as kidney, proximal small intestine, and salivary glands. Though PSMA is expressed in brain, that expression is minimal, and most ligands of PSMA are polar and are not capable of penetrating the blood brain barrier. PSMA is a type II cell surface membrane-bound glycoprotein with ~110 kD molecular

weight, including an intracellular segment (amino acids 1-18), a transmembrane domain (amino acids 19-43), and an extensive extracellular domain (amino acids 44-750). While the functions of the intracellular segment and the transmembrane domains are currently believed to be insignificant, the extracellular domain is involved in several distinct activities. PSMA plays a role in central nervous system, where it metabolizes N-acetyl-aspartyl glutamate (NAAG) into glutamic and N-acetyl aspartic acid. Accordingly, it is also sometimes referred to as an N-acetyl alpha linked acidic dipeptidase (NAALADase). PSMA is also sometimes referred to as a folate hydrolase I (FOLH I) or glutamate carboxypeptidase (GCP II) due to its role in the proximal small intestine where it removes γ -linked glutamate from poly- γ -glutamated folate and α -linked glutamate from peptides and small molecules.

PSMA also shares similarities with human transferrin receptor (TfR), because both PSMA and TfR are type II glycoproteins. More specifically, PSMA shows 54% and 60% homology to TfR1 and TfR2, respectively. However, though TfR exists only in dimeric form due to the formation of inter-strand sulfhydryl linkages, PSMA can exist in either dimeric or monomeric form.

Unlike many other membrane-bound proteins, PSMA undergoes rapid internalization into the cell in a similar fashion to cell surface bound receptors like vitamin receptors. PSMA is internalized through clathrin-coated pits and subsequently can either recycle to the cell surface or go to lysosomes. It has been suggested that the dimer and monomer form of PSMA are inter-convertible, though direct evidence of the interconversion is being debated. Even so, only the dimer of PSMA possesses enzymatic activity, and the monomer does not.

Though the activity of the PSMA on the cell surface of the prostate cells remains under investigation, it has been recognized by the inventors herein that PSMA represents a viable target for the selective and/or specific delivery of biologically active agents, including diagnostic agents, imaging agents, and therapeutic agents to such prostate cells.

SUMMARY OF THE INVENTION

It has been discovered that biologically active compounds that are conjugated to ligands capable of binding to prostate specific membrane antigen (PSMA) via a linker may be useful in the imaging, diagnosis, and/or treatment of prostate cancer, and related diseases that involve pathogenic cell populations expressing or over-expressing PSMA. PSMA is a cell surface protein that is internalized in a process analogous to endocytosis observed with cell surface receptors, such as vitamin receptors. Accordingly, it has been discovered that certain conjugates that include a linker having a predetermined length, and/or a predetermined diameter, and/or preselected functional groups along its length may be used to treat, image, and/or diagnose such diseases.

In one illustrative embodiment of the invention, conjugates having the formula

B-L-D

are described wherein B is a prostate specific membrane antigen (PSMA) binding or targeting ligand, L is a linker, and D is a drug. As used herein, the term drug D collectively includes therapeutic agents, cytotoxic agents, imaging agents, diagnostic agents, and the like, unless otherwise indicated or by the context. For example, in one illustrative configuration, the conjugate described herein is used to

eliminate a pathogenic population of cells and therefore the drug D is a therapeutic agent, a cytotoxic agent, and the like. In another illustrative configuration, the conjugate described herein is used to image and/or diagnose a disease or disease state, and therefore the drug D is an imaging agent, a diagnostic agent, and the like. Other configurations are also contemplated and described herein. It is to be understood that analogs and derivatives of each of the foregoing B, L, and D are also contemplated and described herein, and that when used herein, the terms B, L, and D collectively refer to such analogs and derivatives.

In one illustrative embodiment, the linker L may be a releasable or non-releasable linker. In one aspect, the linker L is at least about 7 atoms in length. In one variation, the linker L is at least about 10 atoms in length. In one variation, the linker L is at least about 14 atoms in length. In another variation, the linker L is between about 7 and about 31, between about 7 and about 24, or between about 7 and about 20 atoms in length. In another variation, the linker L is between about 14 and about 31, between about 14 and about 24, or between about 14 and about 20 atoms in length.

In an alternative aspect, the linker L is at least about 10 angstroms (Å) in length. In one variation, the linker L is at least about 15 Å in length. In another variation, the linker L is at least about 20 Å in length. In another variation, the linker L is in the range from about 10 Å to about 30 Å in length.

In an alternative aspect, at least a portion of the length of the linker L is about 5 Å in diameter or less at the end connected to the binding ligand B. In one variation, at least a portion of the length of the linker L is about 4 Å or less, or about 3 Å or less in diameter at the end connected to the binding ligand B. It is appreciated that the illustrative embodiments that include a diameter requirement of about 5 Å or less, about 4 Å or less, or about 3 Å or less may include that requirement for a predetermined length of the linker, thereby defining a cylindrical-like portion of the linker. Illustratively, in another variation, the linker includes a cylindrical portion at the end connected to the binding ligand that is at least about 7 Å in length and about 5 Å or less, about 4 Å or less, or about 3 Å or less in diameter.

In another embodiment, the linker L includes one or more hydrophilic linkers capable of interacting with one or more residues of PSMA, including amino acids that have hydrophobic side chains, such as Ser, Thr, Cys, Arg, Orn, Lys, Asp, Glu, Gln, and like residues. In another embodiment, the linker L includes one or more hydrophobic linkers capable of interacting with one or more residues of PSMA, including amino acids that have hydrophilic side chains, such as Val, Leu, Ile, Phe, Tyr, Met, and like residues. It is to be understood that the foregoing embodiments and aspects may be included in the linker L either alone or in combination with each other. For example, linkers L that are at least about 7 atoms in length and about 5 Å, about 4 Å or less, or about 3 Å or less in diameter or less are contemplated and described herein, and also include one or more hydrophilic linkers capable of interacting with one or more residues of PSMA, including Val, Leu, Ile, Phe, Tyr, Met, and like residues are contemplated and described herein.

In another embodiment, one end of the linker is not branched and comprises a chain of carbon, oxygen, nitrogen, and sulfur atoms. In one embodiment, the linear chain of carbon, oxygen, nitrogen, and sulfur atoms is at least 5 atoms in length. In one variation, the linear chain is at least 7 atoms, or at least 10 atoms in length. In another embodiment, the chain of carbon, oxygen, nitrogen, and sulfur atoms are not substituted. In one variation, a portion of the

chain of carbon, oxygen, nitrogen, and sulfur atoms is cyclized with a divalent fragment. For example, a linker (L) comprising the dipeptide Phe-Phe may include a piperazine-1,4-diyl structure by cyclizing two nitrogens with an ethylene fragment, or substituted variation thereof.

In another embodiment, pharmaceutical compositions are described herein, where the pharmaceutical composition includes the conjugates described herein in amounts effective to treat diseases and disease states, diagnose diseases or disease states, and/or image tissues and/or cells that are associated with pathogenic populations of cells expressing or over expressing PSMA. Illustratively, the pharmaceutical compositions also include one or more carriers, diluents, and/or excipients.

In another embodiment, methods for treating diseases and disease states, diagnosing diseases or disease states, and/or imaging tissues and/or cells that are associated with pathogenic populations of cells expressing or over expressing PSMA are described herein. Such methods include the step of administering the conjugates described herein, and/or pharmaceutical compositions containing the conjugates described herein, in amounts effective to treat diseases and disease states, diagnose diseases or disease states, and/or image tissues and/or cells that are associated with pathogenic populations of cells expressing or over expressing PSMA.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A. Cell bound radioactivity versus concentration of SK28-^{99m}Tc ($K_d=18.12$ nM) in the presence (▲) or absence (■) of excess PMPA.

FIG. 1B. In Vitro Binding Studies Using LNCaP Cells and SK33 (14 atom linker). LNCaP cells containing increasing concentrations of DUPA-^{99m}Tc in the presence (▲) or absence (■) of excess PMPA.

FIG. 2. Cell bound radioactivity versus concentration of SK28-^{99m}Tc; at 4° C. (■) and at 37° C. (▲).

FIG. 3A. Plot of cell bound radioactivity versus concentration of DUPA-Linker-^{99m}Tc imaging agents: (■) 0-atom linker ($K_d=171$ nM); (▲) 7-atom linker ($K_d=68$ nM); (▼) 14-atom linker ($K_d=15$ nM); (◆) 16-atom linker ($K_d=40$ nM).

FIG. 3B. K_D values for DUPA-Linker-^{99m}Tc compounds binding to LNCaP cells.

FIG. 4. Plot of days post injection versus tumor volume for LNCaP tumors: (a) 2.5 million+Matrigel; (b) 2.5 million+HC Matrigel; (c) 5 million+Matrigel; (d) 5 million+HC Matrigel.

FIG. 5A. Plot of days post injection versus tumor volume for LNCaP tumors (27 mice) and (b) KB cells (5 mice) and A549 cells (5 mice).

FIG. 5B. Plot of days post injection versus tumor volume for KB cells (5 mice) and A549 cells (5 mice).

FIG. 6A. Mice (Set 1) previously injected with LNCaP tumors, treated with 1 ng/kg SK28-^{99m}Tc (14-atom linker), the left hand image shows white light images and image shows an overlay of the radioimage with the white light image. In each panel, the right mouse was treated with 50 mg/kg PMPA (to block PSMA binding) and the left mouse was treated without added PMPA.

FIG. 6B. Mice (Set 2) previously injected with LNCaP tumors, treated with 1 ng/kg SK28-^{99m}Tc (14-atom linker), the left hand image shows white light images and image shows an overlay of the radioimage with the white light image. In each panel, the right mouse was treated with 50

mg/kg PMPA (to block PSMA binding) and the left mouse was treated without added PMPA.

FIG. 6C. Mice (Set 3) previously injected with LNCaP tumors, treated with 1 ng/kg SK28-^{99m}Tc (14-atom linker), the left hand image shows white light images and image shows an overlay of the radioimage with the white light image. In each panel, the right mouse was treated with 50 mg/kg PMPA (to block PSMA binding) and the left mouse was treated without added PMPA.

FIG. 6D. Shows a single mouse study for LNCaP tumors imaged using Kodak imager 4 hours after subcutaneous (administered through intraperitoneal) injection of 1 ng/kg SK28-^{99m}Tc showing in the left hand image an overlay of radioimage with kidney shield and white light image with no shield and in the right hand image an overlay of radioimage with kidney shield and X-ray image with no shield.

FIG. 7A. Mice previously injected with LNCaP tumors treated using SK60-^{99m}Tc (zero atom linker). The left image shows white light images, the center image shows overlay of radioimage with white light image, and the right image shows overlay of radioimage with white light image by shielding the kidney of mice.

FIG. 7B. Mice previously injected with KB cells treated using SK28-^{99m}Tc (14 atom linker). The left image shows white light images, the center image shows overlay of radioimage with white light image, and the right image shows overlay of radioimage with white light image by shielding the kidney of mice.

FIG. 7C. Mice previously injected with A549 cells treated using SK28-^{99m}Tc (14 atom linker). The left image shows white light images, the center image shows overlay of radioimage with white light image, and the right image shows overlay of radioimage with white light image by shielding the kidney of mice.

FIG. 7D. Whole body images of solid tumor xenografts in nu/nu mice taken 4 h after injection of 150 μ Ci jDUPA-^{99m}Tc. Overlay of whole-body radioimages on white light images of mice bearing LNCaP tumors that were treated with DUPA-^{99m}Tc in the absence (a, c) or presence (b, d) of 100-fold molar excess PMPA. Overlay of radioimages on white light images of mice bearing an A549 tumor (e) or a KB tumor (f) that were similarly treated with DUPA-^{99m}Tc.

FIG. 8A. Bio-distribution data as measured for direct cpm count verses tissue for SK28-^{99m}Tc with or without PMPA (as a competitor) on LNCaP, (a) without, (b) with; (c) A549; and (d) KB tumors implanted on axial of male nude mice.

FIG. 8B. Bio-distribution data just for tumor and kidney for SK28-^{99m}Tc with (b) or without (a) PMPA (as a competitor) on LNCaP tumors implant on axial of male nude mice.

FIG. 8C. Biodistribution studies of DUPA-^{99m}Tc in nu/nu mice bearing LNCaP, A549, or KB tumors.

FIG. 9A. Acute MTD (single dose) showing percentage weight change after a single dose of SK71; saline alone, 1.1 μ mol/kg, 2.3 μ mol/kg, 4.5 μ mol/kg, and 9 μ mol/kg.

FIG. 9B. Chronic MTD showing percentage weight change after 5 doses given on alternate days (M, W, F, M, W); saline alone, 2 μ mol/kg and 4 μ mol/kg.

FIG. 10A. Efficacy study showing tumor volume in animals treated with the conjugate SK71 administered in 5 doses on alternate days (M, W, F, M, W) at 1 μ mol/kg.

FIG. 10B. Efficacy study (control group) showing tumor volume in animals treated with saline alone administered in 5 doses on alternate days (M, W, F, M, W).

FIG. 10C. Efficacy study (competition) showing tumor volume in animals treated with excess PSMA and the

conjugate SK71 administered in 5 doses on alternate days (M, W, F, M, W) at 1 μ mol/kg.

FIG. 11. Efficacy Study (1 micromole/kg every other day for 10 days; i.e. 5 doses).

FIG. 12A. [³H]-Thymidine incorporation of LNCaP cells after treatment with SK71 (IC₅₀~2 nM).

FIG. 12B. [³H]-Thymidine incorporation of LNCaP cells after treatment with SK77 (IC₅₀~3 nM).

FIG. 12C. [³H]-Thymidine incorporation of LNCaP cells after treatment with SK37 (IC₅₀~33 nM).

FIG. 12D. [³H]-Thymidine incorporation of LNCaP cells after treatment with SK45 (IC₅₀~29 nM).

FIG. 13A. The effect of treatment with SK71 (1.5 μ mol/kg) on tumor volume in nu/nu mice previously treated with LNCaP cells in HC Matrigel. Treated mice (■), untreated mice (●), treated mice pre-injected with 100-fold molar excess of PMPA (▲).

FIG. 13B. The effect of treatment with SK71 (1.5 μ mol/kg) on percentage weight change in nu/nu mice previously treated with LNCaP cells in HC Matrigel. Treated mice (■), untreated mice (●), treated mice pre-injected with 100-fold molar excess of PMPA (▲).

FIG. 13C. The effect of treatment with SK71 (2.0 μ mol/kg) on tumor volume in nu/nu mice previously treated with LNCaP cells in HC Matrigel. Treated mice (■), untreated mice (●), treated mice pre-injected with 30-fold molar excess of PMPA (▼).

FIG. 13D. The effect of treatment with SK71 (2.0 μ mol/kg) on percentage weight change in nu/nu mice previously treated with LNCaP cells in HC Matrigel. Treated mice (■), untreated mice (●), treated mice pre-injected with 30-fold molar excess of PMPA ().

FIG. 14A. The effect of treatment with SK77 (2.0 μ mol/kg) on tumor volume in nu/nu mice previously treated with LNCaP cells in HC Matrigel. Untreated mice (■), treated mice (▼).

FIG. 14B. The effect of treatment with SK77 (2.0 μ mol/kg) on percentage weight change in nu/nu mice previously treated with LNCaP cells in HC Matrigel. Untreated mice (■), treated mice (▼).

FIG. 15A. MUPA 99mTc imaging agent conjugate (9-atom linker) with energy minimized computer model.

FIG. 15B. MUPA 99mTc imaging agent conjugate (syn-SK33, 14-atom linker) with energy minimized computer model.

DETAILED DESCRIPTION

Drug delivery conjugates are described herein where a PSMA binding ligand is attached to a releasable or non-releasable linker which is attached to a drug, therapeutic agent, diagnostic agent, or imaging agent.

Illustratively, the bivalent linkers described herein may be included in linkers used to prepare PSMA-binding drug conjugates, PSMA-binding imaging agent conjugates, and PSMA-binding diagnostic agent conjugates of the following formulae:

B-L-TA

B-L-IA

B-L-DA

where B is a PSMA-binding moiety, including analogs or derivatives thereof, L is a linker, TA is a therapeutic agent, including analogs or derivatives thereof, IA is an imaging agent, including analogs or derivatives thereof, and DA is a

diagnostic agent, including analogs or derivatives thereof. The linker L can comprise multiple bivalent linkers, including the bivalent linkers described herein. It is also to be understood that as used herein, TA collectively refers to therapeutic agents, and analogs and derivatives thereof, IA collectively refers to imaging agents, and analogs and derivatives thereof, and DA collectively refers to diagnostic agents, and analogs and derivatives thereof.

The linker may also include one or more spacer linkers and optionally additional releasable linkers. The spacer and releasable linkers may be attached to each other in any order or combination. Similarly, the PSMA binding ligand may be attached to a spacer linker or to a releasable linker. Similarly, the drug, therapeutic agent, diagnostic agent, or imaging agent may be attached to a spacer linker or to a releasable linker. Each of these components of the conjugates may be connected through existing or additional heteroatoms on the targeting ligand, drug, therapeutic agent, diagnostic agent, imaging agent, releasable or spacer linker. Illustrative heteroatoms include nitrogen, oxygen, sulfur, and the formulae $-(NHR^1NHR^2)-$, $-SO-$, $-(SO_2)-$, and $-N(R^3)O-$, wherein R^1 , R^2 , and R^3 are each independently selected from hydrogen, alkyl, heteroalkyl, heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and the like, each of which may be optionally substituted.

In one illustrative embodiment, compounds are described herein that include linkers having predetermined length and diameter dimensions. In one aspect, linkers are described herein that satisfy one or more minimum length requirements, or a length requirement falling within a predetermined range. In another aspect, satisfaction of a minimum length requirement may be understood to be determined by computer modeling of the extended conformations of linkers. In another aspect, satisfaction of a minimum length requirement may be understood to be determined by having a certain number of atoms, whether or not substituted, forming a backbone chain of atoms connecting the binding ligand (B) with the drug (D). In another embodiment, the backbone chain of atoms is cyclized with another divalent fragment. In another aspect, linkers are described herein that satisfy one or more maximum or minimum diameter requirements. In another aspect, satisfaction of a maximum or minimum diameter requirement may be understood to be determined by computer modeling of various conformations of linkers modeled as the space-filling, CPK, or like configurations. In another aspect, satisfaction of a maximum or minimum diameter requirement may be understood to be applied to one or more selected portions of the linker, for example the portion of the linker proximal to the binding ligand (B), or the portion of the linker proximal to the drug (D), and the like. In another aspect, linkers are described herein that satisfy one or more chemical composition requirements, such as linkers that include one or more polar groups that may positively interact with the one or more Arg or Lys side-chain nitrogens and/or Asp or Glu side chain oxygens found in the funnel portion of PSMA. In one variation, linkers are described herein that satisfy one or more chemical composition requirements, such as linkers that include one or more non-polar groups that may positively interact with the one or more Tyr or Phe side-chain carbons found in the funnel portion of PSMA.

In one embodiment, the atom-length of the linker is defined by the number of atoms separating the binding or targeting ligand B, or analog or derivative thereof, and the drug D, or analog or derivative thereof. Accordingly, in configurations where the binding ligand B, or analog or derivative thereof, is attached directly to the drug D, or

analog or derivative thereof, the attachment is also termed herein as a "0-atom" linker. It is understood that such 0-atom linkers include the configuration wherein B and D are directly attached by removing a hydrogen atom from each attachment point on B and D, respectively. It is also understood that such 0-atom linkers include the configuration wherein B and D are attached through an overlapping heteroatom by removing a hydrogen atom from one of B or D, and a heteroatom functional group, such as OH, SH, NH_2 , and the like from the other of B or D. It is also understood that such 0-atom linkers include the configuration wherein B and D are attached through a double bond, which may be formed by removing two hydrogen atoms from each attachment point on B and D, respectively, or whereby B and D are attached through one or more overlapping heteroatoms by removing two hydrogen atoms, one hydrogen and one heteroatom functional group, or two heteroatom functional groups, such as OH, SH, NH_2 , and the like, from each of B or D. In addition, B and D may be attached through a double bond formed by removing a double bonded heteroatom functional group, such as O, S, NH, and the like, from one or both of B or D. It is also to be understood that such heteroatom functional groups include those attached to saturated carbon atoms, unsaturated carbon atoms (including carbonyl groups), and other heteroatoms. Similarly, the length of linkers that are greater than 0 atoms are defined in an analogous manner.

Accordingly, in another illustrative embodiment, linkers (L) are described having a chain length of at least 7 atoms. In one variation, linkers (L) are described having a chain length of at least 14 atoms. In another variation, linkers (L) are described having a chain length in the range from about 7 atoms to about 20 atoms. In another variation, linkers (L) are described having a chain length in the range from about 14 atoms to about 24 atoms.

In another embodiment, the length of the linker (L) is defined by measuring the length of an extended conformation of the linker. Such extended conformations may be measured in art-recognized computer modeling programs, such as PC Model 7 (MMX). Accordingly, in another illustrative embodiment, linkers are described having a chain length of at least 15 Å, at least 20 Å, or at least 25 Å.

In another embodiment, linkers are described having at least one hydrophobic side chain group, such as an alkyl, cycloalkyl, aryl, arylalkyl, or like group, each of which is optionally substituted. In one aspect, the hydrophobic group is included in the linker by incorporating one or more Phe or Tyr groups, including substituted variants thereof, and analogs and derivatives thereof, in the linker chain. It is appreciated that such Phe and/or Tyr side chain groups may form positive pi-pi ($\pi-\pi$) interactions with Tyr and Phe residues found in the funnel of PSMA. In addition, it is appreciated that the presence of large side chain branches, such as the arylalkyl groups found on Phe and Tyr may provide a level of conformational rigidity to the linker, thus limiting the degrees of freedom, and reducing coiling and promoting extended conformations of the linker. Without being bound by theory, it is appreciated that such entropy restrictions may increase the overall binding energy of the bound conjugates described herein. In addition, it is appreciated that the rigidity increases that may be provided by sterically hindered side chains, such as Phe and Tyr described herein, may reduce or prevent coiling and interactions between the ligand and the imaging agent. For example, computational energy minimization of a representative 9-atom and 14-atom linker (see, for example, FIGS. 15A and 15B) shows that there are no intra-molecular interactions between the ligand

and the imaging agent. Moreover, the presence of side chain the two Phe side chains appears to promote a more extended conformation in syn-SK33 (FIG. 15B) than in the amino-hexanoic acid-containing conjugate (FIG. 15A)

It has been discovered herein that the funnel shaped tunnel leading to the catalytic site or active site of PSMA imposes length, shape, and/or chemical composition requirements on the linker portion of conjugates of PSMA binding ligands and therapeutic, diagnostic, and imaging agents that positively and negatively affect the interactions between PSMA and those conjugates. Described herein are illustrative embodiments of those conjugates that include such length, shape, and/or chemical composition requirements on the linker. Such length, shape, and/or chemical composition requirements were assessed using molecular modeling. For example, the space filling and surface model of the PSMA complex with (s)-2-(4-iodobenzensylphosphonomethyl)-pentanedioic [2-PMPA derivative] PDB ID code 2C6P were generated using PROTEIN EXPLORER. The PROTEIN EXPLORER model verified the 20 Å deep funnel, and also showed diameter features at various locations along the funnel that may be used to define linkers having favorable structural features. In addition, the model showed that close to the active site of PSMA, there are a higher number of hydrophobic residues that may provide additional binding interactions when the corresponding functional groups are included in the linker. Finally, the model showed the presence of three hydrophobic pockets that may provide additional binding interactions when the corresponding functional groups are included in the linker.

In another illustrative embodiment, the following molecular models were created for a conjugate of MUPA and a tripeptide ^{99m}Tc imaging agent connected by a 9-atom linker, as shown in FIG. 15A, and syn-SK33 including a branched 14-atom linker, as shown in FIG. 15B. The models were created using PC Model 7 (MMX) with energy minimization, and using the following bond length parameters: C—C (sp³-sp³)=1.53 Å, C—C (sp³-sp²)=1.51 Å, C—N (sp³-N)=1.47 Å, C—N (sp²-N)=1.38 Å. Such models may be used to calculate the length of the linker connecting the binding ligand (B) and the drug (D). In addition, such models may be modified to create extended conformations, and subsequently used to calculate the length of the linker connecting the binding ligand (B) and the drug (D).

The first human PSMA gene was cloned from LNCaP cells and is reported to be located in chromosome 11p11-12. In addition, there is a PSMA-like gene located at the loci 11q14.3. The crystal structure of PSMA has been reported by two different groups at different resolutions, and each shows that the active site contains two zinc atoms, confirming that PSMA is also considered a zinc metalloprotease. Davis et al, PNAS, 102:5981-86, (2005) reported the crystal structure at low resolution (3.5 Å), while Mesters et al, The EMBO Journal, 1-10 (2006) reported the crystal structure at higher resolution (2-2.2 Å), the disclosures of which are incorporated herein by reference. The crystal structures show that PSMA is a homodimer that contains a protease domain, an apical domain, a helical domain and a CPG2 dimerization domain. The protease domain of PSMA contains a binuclear zinc site, catalytic residues and a substrate binding region including three arginine residues (also referred to as a substrate binding arginine patch). In the crystal structure, the two zinc ions in the active site are each ligated to an oxygen of phosphate, or to the phosphinate moiety of the inhibitor GPI 18431 for the co-crystal structure. In the high resolution crystal structures of the extracellular domain, PSMA was co-crystallized with both potent

inhibitors, weak inhibitors, and glutamate at 2.0, 2.4, and 2.2 Å, respectively. The high resolution crystal structure shows a 20 Å deep funnel shaped tunnel leads to the catalytic site or active site of PSMA. The funnel is lined with the side chains of a number of Arg and Lys residues, Asp and Glu residues, and Tyr and Phe residues.

In another embodiment, the linker (L) is a chain of atoms selected from C, N, O, S, Si, and P. The linker may have a wide variety of lengths, such as in the range from about 7 to about 100. The atoms used in forming the linker may be combined in all chemically relevant ways, such as chains of carbon atoms forming alkylene groups, chains of carbon and oxygen atoms forming polyoxyalkylene groups, chains of carbon and nitrogen atoms forming polyamines, and others. In addition, it is to be understood that the bonds connecting atoms in the chain may be either saturated or unsaturated, such that for example, alkanes, alkenes, alkynes, cycloalkanes, arylenes, imides, and the like may be divalent radicals that are included in the linker. In addition, it is to be understood that the atoms forming the linker may also be cyclized upon each other to form divalent cyclic radicals in the linker. In each of the foregoing and other linkers described herein the chain forming the linker may be substituted with a wide variety of groups.

In another embodiment, linkers (L) are described that include at least one releasable linker. In one variation, linkers (L) are described that include at least two releasable linkers. In another variation, linkers (L) are described that include at least one self-immolative linker. In another variation, linkers (L) are described that include at least one releasable linker that is not a disulfide. In another embodiment, linkers (L) are described that do not include a releasable linker.

It is appreciated that releasable linkers may be used when the drug to be delivered is advantageously liberated from the binding ligand-linker conjugate so that the free drug will have the same or nearly the same effect at the target as it would when administered without the targeting provided by the conjugates described herein. In another embodiment, the linker L is a non-releasable linker. It is appreciated that non-releasable linkers may be used when the drug is advantageously retained by the binding ligand-linker conjugate, such as in imaging, diagnosing, uses of the conjugates described herein. It is to be understood that the choice of a releasable linker or a non-releasable linker may be made independently for each application or configuration of the conjugates, without limiting the invention described herein. It is to be further understood that the linkers L described herein comprise various atoms, chains of atoms, functional groups, and combinations of functional groups. Where appropriate in the present disclosure, the linker L may be referred to by the presence of spacer linkers, releasable linkers, and heteroatoms. However, such references are not to be construed as limiting the definition of the linkers L described herein.

The linker (L) comprising spacer and/or releasable linkers (i.e., cleavable linkers) can be any biocompatible linker. The releasable or cleavable linker can be, for example, a linker susceptible to cleavage under the reducing or oxidizing conditions present in or on cells, a pH-sensitive linker that may be an acid-labile or base-labile linker, or a linker that is cleavable by biochemical or metabolic processes, such as an enzyme-labile linker. In one embodiment, the spacer and/or releasable linker comprises about 1 to about 30 atoms, or about 2 to about 20 atoms. Lower molecular weight linkers (i.e., those having an approximate molecular weight of about 30 to about 300) are also described. Precursors to such

15

each of the releasable linkers is optionally substituted with a substituent X^2 , as defined below, and the releasable linker is bonded to the oxygen to form an acetal or ketal. Alternatively, the releasable linker may include oxygen, and the releasable linker can be methylene, wherein the methylene is substituted with an optionally-substituted aryl, and the releasable linker is bonded to the oxygen to form an acetal or ketal. Further, the releasable linker may include oxygen, and the releasable linker can be sulfonylalkyl, and the releasable linker is bonded to the oxygen to form an alkyl-sulfonate.

In another embodiment of the above releasable linker embodiment, the releasable linker may include nitrogen, and the releasable linkers can be iminoalkylidenyl, carbonylalkylideniminyl, iminocycloalkylidenyl, and carbonylcycloalkylideniminyl, wherein each of the releasable linkers is optionally substituted with a substituent X^2 , as defined below, and the releasable linker is bonded to the nitrogen to form an hydrazone. In an alternate configuration, the hydrazone may be acylated with a carboxylic acid derivative, an orthoformate derivative, or a carbamoyl derivative to form various acylhydrazone releasable linkers.

Alternatively, the releasable linker may include oxygen, and the releasable linkers can be alkylene(dialkylsilyl), alkylene(alkylarylsilyl), alkylene(diarylsilyl), (dialkylsilyl) aryl, (alkylarylsilyl)aryl, and (diarylsilyl)aryl, wherein each of the releasable linkers is optionally substituted with a substituent X^2 , as defined below, and the releasable linker is bonded to the oxygen to form a silanol.


In the above releasable linker embodiment, the drug can include a nitrogen atom, the releasable linker may include nitrogen, and the releasable linkers can be carbonylarylcarbonyl, carbonyl(carboxyaryl)carbonyl, carbonyl(biscarboxyaryl)carbonyl, and the releasable linker can be bonded to the heteroatom nitrogen to form an amide, and also bonded to the drug nitrogen to form an amide.

In the above releasable linker embodiment, the drug can include an oxygen atom, the releasable linker may include nitrogen, and the releasable linkers can be carbonylarylcarbonyl, carbonyl(carboxyaryl)carbonyl, carbonyl(biscarboxyaryl)carbonyl, and the releasable linker can form an amide, and also bonded to the drug oxygen to form an ester.

The substituents X^2 can be alkyl, alkoxy, alkoxyalkyl, hydroxy, hydroxyalkyl, amino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, halo, haloalkyl, sulfhydrylalkyl, alkylthioalkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, heteroaryl, substituted heteroaryl, carboxy, carboxyalkyl, alkyl carboxylate, alkyl alkanoate, guanidinoalkyl, R^4 -carbonyl, R^3 -carbonylalkyl, R^6 -acylamino, and R^7 -acylaminoalkyl, wherein R^4 and R^5 are each independently selected from amino acids, amino acid derivatives, and peptides, and wherein R^6 and R^7 are each independently selected from amino acids, amino acid derivatives, and peptides. In this embodiment the releasable linker can include nitrogen, and the substituent X^2 and the releasable linker can form a heterocycle.


The heterocycles can be pyrrolidines, piperidines, oxazolidines, isoxazolidines, thiazolidines, isothiazolidines, pyrrolidinones, piperidinones, oxazolidinones, isoxazolidinones, thiazolidinones, isothiazolidinones, and succinimides.

In one embodiment, the polyvalent linkers described herein are or include compounds of the following formulae:




16

-continued




where n is an integer selected from 1 to about 4; R^a and R^b are each independently selected from the group consisting of hydrogen and alkyl, including lower alkyl such as C_1 - C_4 alkyl that are optionally branched; or R^a and R^b are taken together with the attached carbon atom to form a carbocyclic ring; R is an optionally substituted alkyl group, an optionally substituted acyl group, or a suitably selected nitrogen protecting group; and (*) indicates points of attachment for the drug, vitamin, imaging agent, diagnostic agent, other polyvalent linkers, or other parts of the conjugate.

In another embodiment, the polyvalent linkers described herein are or include compounds of the following formulae




where m is an integer selected from 1 to about 4; R is an optionally substituted alkyl group, an optionally substituted acyl group, or a suitably selected nitrogen protecting group; and (*) indicates points of attachment for the drug, vitamin, imaging agent, diagnostic agent, other polyvalent linkers, or other parts of the conjugate.

In another embodiment, the polyvalent linkers described herein are or include compounds of the following formulae:



17



where m is an integer selected from 1 to about 4; R is an optionally substituted alkyl group, an optionally substituted acyl group, or a suitably selected nitrogen protecting group; and (*) indicates points of attachment for the drug, vitamin, imaging agent, diagnostic agent, other polyvalent linkers, or other parts of the conjugate.

In another embodiment, the linker L includes one or more spacer linkers. Such spacer linkers can be 1-alkylenesuccinimid-3-yl, optionally substituted with a substituent X^1 , as defined below, and the releasable linkers can be methylene, 1-alkoxyalkylene, 1-alkoxycycloalkylene, 1-alkoxyalkylenecarbonyl, 1-alkoxycycloalkylenecarbonyl, wherein each of the releasable linkers is optionally substituted with a substituent X^2 , as defined below, and wherein the spacer linker and the releasable linker are each bonded to the spacer linker to form a succinimid-1-ylalkyl acetal or ketal.

The spacer linkers can be carbonyl, thionocarbonyl, alkylene, cycloalkylene, alkylencycloalkyl, alkylenecarbonyl, cycloalkylenecarbonyl, carbonylalkylcarbonyl, 1-alkylene-succinimid-3-yl, 1-(carbonylalkyl)succinimid-3-yl, alkylene-sulfoxyl, sulfonylalkyl, alkylene-sulfoxylalkyl, alkylene-sulfonylalkyl, carbonyltetrahydro-2H-pyranyl, carbonyltetrahydrofuranlyl, 1-(carbonyltetrahydro-2H-pyranyl)succinimid-3-yl, and 1-(carbonyltetrahydrofuranlyl)succinimid-3-yl, wherein each of the spacer linkers is optionally substituted with a substituent X^1 , as defined below. In this embodiment, the spacer linker may include an additional nitrogen, and the spacer linkers can be alkylenecarbonyl, cycloalkylenecarbonyl, carbonylalkylcarbonyl, 1-(carbonylalkyl)succinimid-3-yl, wherein each of the spacer linkers is optionally substituted with a substituent X^1 , as defined below, and the spacer linker is bonded to the nitrogen to form an amide. Alternatively, the spacer linker may include an additional sulfur, and the spacer linkers can be alkylene and cycloalkylene, wherein each of the spacer linkers is optionally substituted with carboxy, and the spacer linker is bonded to the sulfur to form a thiol. In another embodiment, the spacer linker can include sulfur, and the spacer linkers can be 1-alkylenesuccinimid-3-yl and 1-(carbonylalkyl)succinimid-3-yl, and the spacer linker is bonded to the sulfur to form a succinimid-3-ylthiol.


In an alternative to the above-described embodiments, the spacer linker can include nitrogen, and the releasable linker can be a divalent radical comprising alkyleneaziridin-1-yl, carbonylalkylaziridin-1-yl, sulfoxylalkylaziridin-1-yl, or sulfonylalkylaziridin-1-yl, wherein each of the releasable linkers is optionally substituted with a substituent X^2 , as defined below. In this alternative embodiment, the spacer

18

linkers can be carbonyl, thionocarbonyl, alkylenecarbonyl, cycloalkylenecarbonyl, carbonylalkylcarbonyl, 1-(carbonylalkyl)succinimid-3-yl, wherein each of the spacer linkers is optionally substituted with a substituent X^1 , as defined below, and wherein the spacer linker is bonded to the releasable linker to form an aziridine amide.

The substituents X^1 can be alkyl, alkoxy, alkoxyalkyl, hydroxy, hydroxyalkyl, amino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, halo, haloalkyl, sulfhydrylalkyl, alkylthioalkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, heteroaryl, substituted heteroaryl, carboxy, carboxyalkyl, alkyl carboxylate, alkyl alkanoate, guanidinoalkyl, R^4 -carbonyl, R^5 -carbonylalkyl, R^6 -acylamino, and R^7 -acylaminoalkyl, wherein R^4 and R^5 are each independently selected from amino acids, amino acid derivatives, and peptides, and wherein R^6 and R^7 are each independently selected from amino acids, amino acid derivatives, and peptides. In this embodiment the spacer linker can include nitrogen, and the substituent X^1 and the spacer linker to which they are bound to form a heterocycle.

Additional illustrative spacer linkers include alkylene-amino-alkylenecarbonyl, alkylene-thio-(carbonylalkylsuccinimid-3-yl), and the like, as further illustrated by the following formulae:



where the integers x and y are 1, 2, 3, 4, or 5:

In another embodiment, linkers that include hydrophilic regions are also described. In one aspect, the hydrophilic region of the linker forms part or all of a spacer linker included in the conjugates described herein. Illustrative hydrophilic spacer linkers are described in PCT international application serial No. PCT/US2008/068093, filed Jun. 25, 2008, the disclosure of which is incorporated herein by reference.

The term “cycloalkyl” as used herein includes molecular fragments or radicals comprising a bivalent chain of carbon atoms, a portion of which forms a ring. It is to be understood that the term cycloalkyl as used herein includes fragments and radicals attached at either ring atoms or non-ring atoms, such as cyclopropyl, cyclohexyl, 3-ethylcyclopent-1-yl, cyclopropylethyl, cyclohexylmethyl, and the like.

The term “cycloalkylene” as used herein includes molecular fragments or radicals comprising a bivalent chain of carbon atoms, a portion of which forms a ring. It is to be understood that the term cycloalkyl as used herein includes fragments and radicals attached at either ring atoms or non-ring atoms, such as cycloprop-1,1-diyl, cycloprop-1,2-diyl, cyclohex-1,4-diyl, 3-ethylcyclopent-1,2-diyl, 1-methylenecyclohex-4-yl, and the like.

The terms “heteroalkyl” and “heteroalkylene” as used herein includes molecular fragments or radicals comprising monovalent and divalent, respectively, groups that are formed from a linear or branched chain of carbon atoms and heteroatoms, wherein the heteroatoms are selected from nitrogen, oxygen, and sulfur, such as alkoxyalkyl, alkyleneoxyalkyl, aminoalkyl, alkylaminoalkyl, alkyleneaminoalkyl,

alkylthioalkyl, alkylenethioalkyl, alkoxyalkylaminoalkyl, alkylaminoalkoxyalkyl, alkyleneoxyalkylaminoalkyl, and the like.

The term "heterocyclyl" as used herein includes molecular fragments or radicals comprising a monovalent chain of carbon atoms and heteroatoms, wherein the heteroatoms are selected from nitrogen, oxygen, and sulfur, a portion of which, including at least one heteroatom, form a ring, such as aziridine, pyrrolidine, oxazolidine, 3-methoxypyrrolidine, 3-methylpiperazine, and the like. Accordingly, as used herein, heterocyclyl includes alkylheterocyclyl, heteroalkylheterocyclyl, heterocyclylalkyl, heterocyclylheteroalkyl, and the like. It is to be understood that the term heterocyclyl as used herein includes fragments and radicals attached at either ring atoms or non-ring atoms, such as tetrahydrofuran-2-yl, piperidin-1-yl, piperidin-4-yl, piperazin-1-yl, morpholin-1-yl, tetrahydrofuran-2-ylmethyl, piperidin-1-ylethyl, piperidin-4-ylmethyl, piperazin-1-ylpropyl, morpholin-1-ylethyl, and the like.

The term "aryl" as used herein includes molecular fragments or radicals comprising an aromatic mono or polycyclic ring of carbon atoms, such as phenyl, naphthyl, and the like.

The term "heteroaryl" as used herein includes molecular fragments or radicals comprising an aromatic mono or polycyclic ring of carbon atoms and at least one heteroatom selected from nitrogen, oxygen, and sulfur, such as pyridinyl, pyrimidinyl, indolyl, benzoxazolyl, and the like.

The term "substituted aryl" or "substituted heteroaryl" as used herein includes molecular fragments or radicals comprising aryl or heteroaryl substituted with one or more substituents, such as alkyl, heteroalkyl, halo, hydroxy, amino, alkyl or dialkylamino, alkoxy, alkylsulfonyl, amino-sulfonyl, carboxylate, alkoxy-carbonyl, aminocarbonyl, cyano, nitro, and the like. It is to be understood that the alkyl groups in such substituents may be optionally substituted with halo.

The term "iminoalkylidenyl" as used herein includes molecular fragments or radicals comprising a divalent radical containing alkylene as defined herein and a nitrogen atom, where the terminal carbon of the alkylene is double-bonded to the nitrogen atom, such as the formulae $-(CH)=N-$, $-(CH_2)_2(CH)=N-$, $-CH_2C(Me)=N-$, and the like.

The term "amino acid" as used herein includes molecular fragments or radicals comprising an aminoalkylcarboxylate, where the alkyl radical is optionally substituted with alkyl, hydroxy alkyl, sulfhydrylalkyl, aminoalkyl, carboxyalkyl, and the like, including groups corresponding to the naturally occurring amino acids, such as serine, cysteine, methionine, aspartic acid, glutamic acid, and the like.

For example, in one embodiment, amino acid is a divalent radical having the general formula:



where R is hydrogen, alkyl, acyl, or a suitable nitrogen protecting group, R' and R'' are hydrogen or a substituent, each of which is independently selected in each occurrence, and q is an integer such as 1, 2, 3, 4, or 5. Illustratively, R' and/or R'' independently correspond to, but are not limited to, hydrogen or the side chains present on naturally occurring amino acids, such as methyl, benzyl, hydroxymethyl, thiomethyl, carboxyl, carboxylmethyl, guanidinopropyl, and the like, and derivatives and protected derivatives thereof. The above described formula includes all stereoisomeric variations. For example, the amino acid may be selected from asparagine, aspartic acid, cysteine, glutamic acid,

lysine, glutamine, arginine, serine, ornithine, threonine, and the like. In one variation, the amino acid may be selected from phenylalanine, tyrosine, and the like, derivatives thereof, and substituted variants thereof.

The terms "arylalkyl" and "heteroarylalkyl" as used herein includes molecular fragments or radicals comprising aryl and heteroaryl, respectively, as defined herein substituted with a linear or branched alkylene group, such as benzyl, phenethyl, α -methylbenzyl, picolinyl, pyrimidinylethyl, and the like.

It is to be understood that the above-described terms can be combined to generate chemically-relevant groups, such as "haloalkoxyalkyl" referring to for example trifluoro-methoxyethyl, 1,2-difluoro-2-chloroeth-1-yloxypropyl, and the like.

The term "amino acid derivative" as used herein refers generally to aminoalkylcarboxylate, where the amino radical or the carboxylate radical are each optionally substituted with alkyl, carboxylalkyl, alkylamino, and the like, or optionally protected; and the intervening divalent aryl fragment is optionally substituted with alkyl, hydroxy alkyl, sulfhydrylalkyl, aminoalkyl, carboxyalkyl, and the like, including groups corresponding to the side chains found in naturally occurring amino acids, such as are found in serine, cysteine, methionine, aspartic acid, glutamic acid, and the like.

The term "peptide" as used herein includes molecular fragments or radicals comprising a series of amino acids and amino acid analogs and derivatives covalently linked one to the other by amide bonds.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form 3-thiosuccinimid-1-ylalkyloxymethoxy, where the methyl is optionally substituted with alkyl or substituted aryl.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form 3-thiosuccinimid-1-ylalkylcarbonyl, where the carbonyl forms an acylaziridine with the drug, or analog or derivative thereof.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form 1-alkoxycycloalkylenoxy.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form alkyleneaminocarbonyl(dicarboxylarylene)carboxylate.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form dithioalkylcarbonylhydrazide, where the hydrazide forms an hydrazone with the drug, or analog or derivative thereof.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form 3-thiosuccinimid-1-ylalkylcarbonylhydrazide, where the hydrazide forms an hydrazone with the drug, or analog or derivative thereof.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form 3-thioalkylsulfonylalkyl(disubstituted silyl)oxy, where the disubstituted silyl is substituted with alkyl or optionally substituted aryl.

In another embodiment, the bivalent linker comprises a plurality of spacer linkers selected from the group consisting of the naturally occurring amino acids and stereoisomers thereof.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker

21

taken together to form 3-dithioalkyloxycarbonyl, where the carbonyl forms a carbonate with the drug, or analog or derivative thereof.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioarylklyloxycarbonyl, where the carbonyl forms a carbonate with the drug, or analog or derivative thereof, and the aryl is optionally substituted.

In another embodiment, the bivalent linker comprises a spacer linker and a releasable linker taken together to form 3-thiosuccinimid-1-ylalkyloxyalkyloxyalkylidene, where the alkylidene forms a hydrazone with the drug, or analog or derivative thereof, each alkyl is independently selected, and the oxyalkyloxy is optionally substituted with alkyl or optionally substituted aryl.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioalkyloxycarbonylhydrazide.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioalkylamino, where the amino forms a vinylogous amide with the drug, or analog or derivative thereof.


In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioalkylamino, where the amino forms a vinylogous amide with the drug, or analog or derivative thereof, and the alkyl is ethyl.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioalkylaminocarbonyl, where the carbonyl forms a carbamate with the drug, or analog or derivative thereof.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioalkylaminocarbonyl, where the carbonyl forms a carbamate with the drug, or analog or derivative thereof, and the alkyl is ethyl.

In another embodiment, the bivalent linker comprises a releasable linker, a spacer linker, and a releasable linker taken together to form 3-dithioarylklyloxycarbonyl, where the carbonyl forms a carbamate or a carbamoylaziridine with the drug, or analog or derivative thereof.

In another embodiment, the polyvalent linker includes spacer linkers and releasable linkers connected to form a polyvalent 3-thiosuccinimid-1-ylalkyloxymethyloxy group, illustrated by the following formula




where n is an integer from 1 to 6, the alkyl group is optionally substituted, and the methyl is optionally substituted with an additional alkyl or optionally substituted aryl group, each of which is represented by an independently selected group R. The (*) symbols indicate points of attachment of the polyvalent linker fragment to other parts of the conjugates described herein.

In another embodiment, the polyvalent linker includes spacer linkers and releasable linkers connected to form a


22

polyvalent 3-thiosuccinimid-1-ylalkylcarbonyl group, illustrated by the following formula




where n is an integer from 1 to 6, and the alkyl group is optionally substituted. The (*) symbols indicate points of attachment of the polyvalent linker fragment to other parts of the conjugates described herein. In another embodiment, the polyvalent linker includes spacer linkers and releasable linkers connected to form a polyvalent 3-thioalkylsulfonyl-alkyl(disubstituted silyl)oxy group, where the disubstituted silyl is substituted with alkyl and/or optionally substituted aryl groups.

In another embodiment, the polyvalent linker includes spacer linkers and releasable linkers connected to form a polyvalent dithioalkylcarbonylhydrazide group, or a polyvalent 3-thiosuccinimid-1-ylalkylcarbonylhydrazide, illustrated by the following formulae



where n is an integer from 1 to 6, the alkyl group is optionally substituted, and the hydrazide forms a hydrazone with (B), (D), or another part of the polyvalent linker (L). The (*) symbols indicate points of attachment of the polyvalent linker fragment to other parts of the conjugates described herein.

In another embodiment, the polyvalent linker includes spacer linkers and releasable linkers connected to form a polyvalent 3-thiosuccinimid-1-ylalkyloxymethyloxyalkylidene group, illustrated by the following formula












where each n is an independently selected integer from 1 to 6, each alkyl group independently selected and is optionally substituted, such as with alkyl or optionally substituted aryl, and where the alkylidene forms a hydrazone with (B), (D), or another part of the polyvalent linker (L). The (*) symbols indicate points of attachment of the polyvalent linker fragment to other parts of the conjugates described herein.

Additional illustrative linkers are described in WO 2006/012527, the disclosure of which is incorporated herein by reference. Additional linkers are described in the following

23










Table, where the (*) atom is the point of attachment of additional spacer or releasable linkers, the drug, and/or the binding ligand.

Illustrative Releasable Linkers



24

-continued



5

10

15

20

25

30

35

40

45

50













55

60

65





25

-continued



26

-continued



Each of the spacer and releasable linkers described herein is bivalent. In addition, the connections between spacer linkers, releasable linkers, drugs D and ligands B may occur at any atom found in the various spacer linkers, releasable linkers, drugs D, and ligands B.

The drug can include a nitrogen atom, and the releasable linker can be haloalkylenecarbonyl, optionally substituted with a substituent X², and the releasable linker is bonded to the drug nitrogen to form an amide.


The drug can include an oxygen atom, and the releasable linker can be haloalkylenecarbonyl, optionally substituted with a substituent X², and the releasable linker is bonded to the drug oxygen to form an ester.

The drug can include a double-bonded nitrogen atom, and in this embodiment, the releasable linkers can be alkylencarbonylamino and 1-(alkylenecarbonylamino)succinimid-3-yl, and the releasable linker can be bonded to the drug nitrogen to form a hydrazone.

The drug can include a sulfur atom, and in this embodiment, the releasable linkers can be alkylenethio and carbonylalkylthio, and the releasable linker can be bonded to the drug sulfur to form a disulfide.

In another embodiment, the binding or targeting ligand capable of binding or targeting PSMA is a phosphoric, phosphonic, or phosphinic acid or derivative thereof. In one aspect, the phosphoric, phosphonic, or phosphinic acid or derivative thereof includes one or more carboxylic acid groups. In another aspect, the phosphoric, phosphonic, or phosphinic acid or derivative thereof includes one or more thiol groups or derivatives thereof. In another aspect, the phosphoric, phosphonic, or phosphinic acid or derivative thereof includes one or more carboxylic acid bioisosteres, such as an optionally substituted tetrazole, and the like.

In another embodiment, the PSMA ligand is a derivative of pentanedioic acid. Illustratively, the pentanedioic acid derivative is a compound of the formula:



wherein X is RP(O)(OH)CH_2 — (see, e.g., U.S. Pat. No. 5,968,915 incorporated herein by reference); RP(O)(OH)N(R^1)— (see, e.g., U.S. Pat. No. 5,863,536 incorporated herein by reference); RP(O)(OH)O — (see, e.g., U.S. Pat. No. 5,795,877 incorporated herein by reference); RN(OH)C(O)Y — or RC(O)NH(OH)Y , wherein Y is $-\text{CR}_1\text{R}_2-$, $-\text{NR}_3-$ or $-\text{O}-$ (see, e.g., U.S. Pat. No. 5,962,521 incorporated herein by reference); RS(O)Y , RSO_2Y , or RS(O)(NH)Y , wherein Y is $-\text{CR}_1\text{R}_2-$, $-\text{NR}_3-$ or $-\text{O}-$ (see, e.g., U.S. Pat. No. 5,902,817 incorporated herein by reference); and RS-alkyl, wherein R is for example hydrogen, alkyl, aryl, or arylalkyl, each of which may be optionally substituted (see, e.g., J. Med. Chem. 46:1989-1996 (2003) incorporated herein by reference).

In each of the foregoing formulae, R, R_1 , R_2 , and R_3 are each independently selected from hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, and aryl. In addition, in each case, each of R, R_1 , R_2 , and R_3 may be optionally substituted, such as with one or more groups selected from C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 - C_4 alkenyloxy, phenoxy, benzyloxy, amino, aryl. In one aspect, aryl is selected from 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, benzyl, and phenyl, and in each case aryl may be optionally substituted with one or more, illustratively with one to three, groups selected from halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 - C_4 alkenyloxy, phenoxy, benzyloxy, and amino. In one variation of each of the above formulae, R is not hydrogen.

Illustrative PSMA ligands described in U.S. Pat. No. 5,968,915 include 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid; 2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid; 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid; 2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid; 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid; 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid; 2-[[2-(tetrahydrofuran-2-yl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[2-(2-tetrahydropyran-2-yl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((4-pyridyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((2-pyridyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((phenylmethyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((2-phenylethyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((3-phenylpropyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((3-phenylbutyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((2-phenylbutyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; 2-[[((4-phenylbutyl)methyl)hydroxyphosphinyl]methyl]pentanedioic acid; and 2-[[aminomethyl]hydroxyphosphinyl]methyl]pentanedioic acid.

Illustrative PSMA ligands described in U.S. Pat. No. 5,863,536 include N-[methylhydroxyphosphinyl]glutamic acid; N-[ethylhydroxyphosphinyl]glutamic acid; N-[propylhydroxyphosphinyl]glutamic acid; N-[butylhydroxyphosphinyl]glutamic acid; N-[phenylhydroxyphosphinyl]glutamic acid; N-[phenylmethyl]hydroxyphosphinyl]glutamic acid; N-[((2-phenylethyl)methyl)hydroxyphosphinyl]glutamic acid; and N-methyl-N-[phenylhydroxyphosphinyl]glutamic acid.

Illustrative PSMA ligands described in U.S. Pat. No. 5,795,877 include 2-[[methylhydroxyphosphinyl]oxy]pen-


tanedioic acid; 2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid; 2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid; 2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid; 2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid; 2-[[((4-pyridyl)methyl)hydroxyphosphinyl]oxy]pentanedioic acid; 2-[[((2-pyridyl)methyl)hydroxyphosphinyl]oxy]pentanedioic acid; 2-[[((phenylmethyl)hydroxyphosphinyl]oxy]pentanedioic acid; and 2-[[((2-phenylethyl)methyl)hydroxyphosphinyl]oxy]pentanedioic acid.

Illustrative PSMA ligands described in U.S. Pat. No. 5,962,521 include 2-[[N-hydroxycarbonyl]methyl]pentanedioic acid; 2-[[N-hydroxy-N-methyl]carbamoyl]methyl]pentanedioic acid; 2-[[N-butyl-N-hydroxycarbonyl]methyl]pentanedioic acid; 2-[[N-benzyl-N-hydroxycarbonyl]methyl]pentanedioic acid; 2-[[N-hydroxy-N-phenyl]carbamoyl]methyl]pentanedioic acid; 2-[[N-hydroxy-N-2-phenylethyl]carbamoyl]methyl]pentanedioic acid; 2-[[N-ethyl-N-hydroxycarbonyl]methyl]pentanedioic acid; 2-[[N-hydroxy-N-propyl]carbamoyl]methyl]pentanedioic acid; 2-[[N-hydroxy-N-3-phenylpropyl]carbamoyl]methyl]pentanedioic acid; 2-[[N-hydroxy-N-4-pyridyl]carbamoyl]methyl]pentanedioic acid; 2-[[N-hydroxycarbonyl]methyl]pentanedioic acid; 2-[[N-hydroxy(methyl)carboxamido]methyl]pentanedioic acid; 2-[[N-hydroxy(benzyl)carboxamido]methyl]pentanedioic acid; 2-[[N-hydroxy(phenyl)carboxamido]methyl]pentanedioic acid; 2-[[N-hydroxy(2-phenylethyl)carboxamido]methyl]pentanedioic acid; 2-[[N-hydroxy(ethyl)carboxamido]methyl]pentanedioic acid; 2-[[N-hydroxy(propyl)carboxamido]methyl]pentanedioic acid; 2-[[N-hydroxy(3-phenylpropyl)carboxamido]methyl]pentanedioic acid; and 2-[[N-hydroxy(4-pyridyl)carboxamido]methyl]pentanedioic acid.


Illustrative PSMA ligands described in U.S. Pat. No. 5,902,817 include 2-[(sulfinyl)methyl]pentanedioic acid; 2-[(methylsulfinyl)methyl]pentanedioic acid; 2-[(ethylsulfinyl)methyl]pentanedioic acid; 2-[(propylsulfinyl)methyl]pentanedioic acid; 2-[(butylsulfinyl)methyl]pentanedioic acid; 2-[(phenylsulfinyl)methyl]pentanedioic acid; 2-[[2-(2-phenylethyl)sulfinyl]methyl]pentanedioic acid; 2-[[3-phenylpropyl)sulfinyl]methyl]pentanedioic acid; 2-[[4-pyridyl)sulfinyl]methyl]pentanedioic acid; 2-[(benzylsulfinyl)methyl]pentanedioic acid; 2-[(sulfonyl)methyl]pentanedioic acid; 2-[(methylsulfonyl)methyl]pentanedioic acid; 2-[(ethylsulfonyl)methyl]pentanedioic acid; 2-[(propylsulfonyl)methyl]pentanedioic acid; 2-[(butylsulfonyl)methyl]pentanedioic acid; 2-[(phenylsulfonyl)methyl]pentanedioic acid; 2-[[2-(2-phenylethyl)sulfonyl]methyl]pentanedioic acid; 2-[[3-(3-phenylpropyl)sulfonyl]methyl]pentanedioic acid; 2-[[4-pyridyl)sulfonyl]methyl]pentanedioic acid; 2-[(benzylsulfonyl)methyl]pentanedioic acid; 2-[(sulfoximiny]methyl]pentanedioic acid; 2-[(methylsulfoximiny]methyl]pentanedioic acid; 2-[(ethylsulfoximiny]methyl]pentanedioic acid; 2-[(propylsulfoximiny]methyl]pentanedioic acid; 2-[(butylsulfoximiny]methyl]pentanedioic acid; 2-[(phenylsulfoximiny]methyl]pentanedioic acid; 2-[[2-(2-phenylethyl)sulfoximiny]methyl]pentanedioic acid; 2-[[3-(3-phenylpropyl)sulfoximiny]methyl]pentanedioic acid; 2-[[4-pyridyl)sulfoximiny]methyl]pentanedioic acid; and 2-[(benzylsulfoximiny]methyl]pentanedioic acid.

29


Pentanedioic acid derivatives described herein have been reported to have high binding affinity at PSMA, including but not limited to the following phosphonic and phosphinic acid derivatives




2 μM




700 nM




0.3 nM




185 nM




560 nM



2 nM



0.5 nM



2 nM

30


with the dissociation constants (K_d values) shown for the E-I complex (see, Current Medicinal Chem. 8:949-.957 (2001); Silverman, "The Organic Chemistry of Drug Design and Drug Action," Elsevier Academic Press (2nd Ed. 2003), the disclosures of which are incorporated herein by reference);

In another illustrative embodiment, the pentanedioic acid derivative includes a thiol group, such as compounds of the following formulae:

10

15

20



(R, S) 90 ± 26 nM
(R) 85 ± 33 nM
(S) 67 ± 29 nM

25

30

with the inhibition constants (IC_{50} values) shown for the E-I complex.

In another embodiment, the PSMA ligand is a urea of two amino acids. In one aspect, the amino acids include one or more additional carboxylic acids. In another aspect, the amino acids include one or more additional phosphoric, phosphonic, phosphinic, sulfinic, sulfonic, or boronic acids. In another aspect, the amino acids include one or more thiol groups or derivatives thereof. In another aspect, the amino acid includes one or more carboxylic acid bioisosteres, such as tetrazoles and the like.

35


40

45

In another embodiment, the PSMA ligand is an aminocarbonyl derivative of pentanedioic acid. Illustratively, the aminocarbonylpentanedioic acid derivative is a compound of the formula:

50

55




60

wherein R^1 and R^2 are each selected from hydrogen, optionally substituted carboxylic acids, such as thiolacetic acids, thiolpropionic acids, and the like; malonic acids, succinic acids, glutamic acids, adipic acids, and the like; and others. Illustrative aminocarbonylpentanedioic acid derivatives are described in J. Med. Chem. 44:298-301 (2001) and J. Med. Chem. 47:1729-38 (2004), the disclosures of which are incorporated herein by reference.

65

31

In another embodiment, the PSMA ligand is a compound of the formula:




It is appreciated that the urea compounds described herein may also be advantageous in the preparation of the ligands also described herein due to the sub-nanomolar potency, water solubility, and/or long term stability of these compounds. The urea compounds described herein may generally be prepared from commercially available starting materials as described herein.

32

It is appreciated that in each of the above illustrative pentanedioic acid compounds and urea compounds, there is at least one asymmetric carbon atom. Accordingly, the above illustrative formulae are intended to refer both individually and collectively to all stereoisomers as pure enantiomers, or mixtures of enantiomers and/or diastereomers, including but not limited to racemic mixtures, mixtures that include one epimer at a first asymmetric carbon but allow mixtures at other asymmetric carbons, including racemic mixtures, and the like.


In another illustrative embodiment, the binding agent is a urea of an amino dicarboxylic acid, such as aspartic acid, glutamic acid, and the like, and another amino dicarboxylic acid, or an analog thereof, such as a binding agent of the formulae










wherein Q is a amino dicarboxylic acid, such as aspartic acid, glutamic acid, or an analog thereof, n and m are each selected from an integer between 1 and about 6, and (*) represents the point of attachment for the linker L.

In another embodiment, the PSMA ligand includes at least four carboxylic acid groups, or at least three free carboxylic acid groups after the PSMA ligand is conjugated to the agent or linker. It is understood that as described herein, carboxylic acid groups on the PSMA ligand include bioisosteres of carboxylic acids.


Illustratively, the PSMA ligand is a compound of the formulae:



33



34



In another embodiment, the PSMA ligand is 2-[3-(1-Carboxy-2-mercapto-ethyl)-ureido]-pentanedioic acid (MUPA) or 2-[3-(1,3-Dicarboxy-propyl)-ureido]-pentanedioic acid (DUPA)

Other illustrative examples of PSMA ligands include peptide analogs such as quisqualic acid, aspartate glutamate (Asp-Glu), Glu-Glu, Gly-Glu, γ -Glu-Glu, beta-N-acetyl-L-aspartate-L-glutamate (β -NAAG), and the like.

In another illustrative embodiment, the binding agent is a urea of an amino dicarboxylic acid, such as aspartic acid, glutamic acid, and the like, and another amino dicarboxylic acid, or an analog thereof, and the linker is peptide of amino acids, including naturally occurring and non-naturally occurring amino acids. In one embodiment, the linker is a peptide comprising amino acids selected from Glu, Asp, Phe, Cys, beta-amino Ala, and aminoalkylcarboxylic acids, such as Gly, beta Ala, amino valeric acid, amino caproic acid, and the like. In another embodiment, the linker is a peptide consisting of amino acids selected from Glu, Asp, Phe, Cys, beta-amino Ala, and aminoalkylcarboxylic acids, such as Gly, beta Ala, amino valeric acid, amino caproic acid, and the like. In another embodiment, the linker is a peptide comprising at least one Phe. In variations, the linker is a peptide comprising at least two Phe residues, or at least three Phe residues. In another embodiment, the linker is a peptide comprising Glu-Phe or a dipeptide of an aminoalkylcarboxylic acid and Phe. In another embodiment, the linker is a peptide comprising Glu-Phe-Phe or a tripeptide of an aminoalkylcarboxylic acid and two Phe residues. In another embodiment, the linker is a peptide comprising one or more Phe residues, where at least one Phe is about 7 to about 11, or about 7 to about 14 atoms from the binding ligand B. In another embodiment, the linker is a peptide comprising Phe-Phe about 7 to about 11, or about 7 to about 14 atoms from the binding ligand B. It is to be understood that in each of the foregoing embodiments and variations, one or more Phe residues may be replaced with Tyr, or another substituted variation thereof.

In another illustrative embodiment, the binding agent is a urea of an amino dicarboxylic acid, such as aspartic acid, glutamic acid, and the like, and another amino dicarboxylic acid, or an analog thereof, and the linker includes one or more aryl or arylalkyl groups, each of which is optionally substituted, attached to the backbone of the linker. In another embodiment, the linker is a peptide comprising one or more


35

aryl or arylalkyl groups, each of which is optionally substituted, attached to the backbone of the linker about 7 to about 11 atoms from the binding ligand B. In another embodiment, the linker is a peptide comprising two aryl or arylalkyl groups, each of which is optionally substituted, attached to the backbone of the linker, where one aryl or arylalkyl group is about 7 to about 11, or about 7 to about 14 atoms from the binding ligand B, and the other aryl or arylalkyl group is about 10 to about 14, or about 10 to about 17 atoms from the binding ligand B.

As described herein, the conjugates are targeted to cells that express or over-express PSMA, using a PSMA binding ligand. Once delivered, the conjugates bind to PSMA. It is understood that in certain embodiments the conjugates remain on the surface of the cell for a period of time sufficient for imaging and/or diagnosis. In other embodiments, the conjugates are internalized in the cell expressing or over-expressing PSMA by endogenous cellular mechanisms, such as endocytosis, for subsequent imaging and/or diagnosis, or treatment. Once internalized, the conjugates may remain intact or be decomposed, degraded, or otherwise altered to allow the release of the agent forming the conjugate. It is appreciated that in imaging and/or diagnostic configurations, the agent may remain intact as the conjugate or be released once it has been internalized into the targeted cell. It is further appreciated that in therapeutic configurations, the agent is advantageously released from the conjugate once it has been internalized into the targeted cell.






In one illustrative embodiment, the drug is an imaging agent. In another illustrative variation, the drug is a diagnostic agent. In another illustrative variation, the drug is an chemotherapeutic agent.

In one aspect, the imaging agent is a radioisotope covalently attached to the linker. In another aspect, the imaging agent is a radioactive isotope, such as a radioactive isotope of a metal, coordinated to a chelating group. Illustrative radioactive metal isotopes include technetium, rhenium, gallium, gadolinium, indium, copper, and the like, including isotopes ^{111}In , $^{99\text{m}}\text{Tc}$, ^{64}Cu , ^{67}Cu , ^{67}Ga , ^{68}Ga , and the like. Additional illustrative examples of radionuclide imaging agents are described in U.S. Pat. No. 7,128,893, the disclosure of which is incorporated herein by reference. Additional illustrative chelating groups are tripeptide or tetrapeptides, including but not limited to tripeptides having the formula:



wherein R is independently selected in each instance H, alkyl, heteroalkyl, cycloalkyl, heterocyclyl, alkenyl, alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and the like, each of which is optionally substituted. It is to be understood that one R includes a heteroatom, such as nitro, oxygen, or sulfur, and is the point of attachment of linker L. Illustratively, the following chelating groups are described:


36




where X is oxygen, nitrogen, or sulfur, and where X is attached to linker L, and n is an integer from 1 to about 5.

In another aspect, the imaging agent is a fluorescent agent. Fluorescent agents include Oregon Green fluorescent agents, including but not limited to Oregon Green 488, Oregon Green 514, and the like, AlexaFluor fluorescent agents, including but not limited to AlexaFluor 488, AlexaFluor 647, and the like, fluorescein, and related analogs, BODIPY fluorescent agents, including but not limited to BODIPY F1, BODIPY 505, and the like, rhodamine fluorescent agents, including but not limited to tetramethylrhodamine, and the like, DyLight fluorescent agents, including but not limited to DyLight 680, DyLight 800, and the like, CW 800, Texas Red, phycoerythrin, and others. Illustrative fluorescent agent are shown in the following illustrative general structures:

37



where X is oxygen, nitrogen, or sulfur, and where X is attached to linker L; Y is OR^a, NR^a₂, or NR^a₃⁺; and Y' is O, NR^a, or NR^a₂⁺; where each R is independently selected in each instance from H, fluoro, sulfonic acid, sulfonate, and salts thereof, and the like; and R^a is hydrogen or alkyl.



where X is oxygen, nitrogen, or sulfur, and where X is attached to linker L; and each R is independently selected in each instance from H, alkyl, heteroalkyl, and the like; and n is an integer from 0 to about 4.

In another aspect, the imaging agent is a PET imaging agent, or a FRET imaging agent. PET imaging agents ¹⁸F, ¹¹C, ⁶⁴Cu, ⁶⁵Cu, and the like. FRET imaging agents include ⁶⁴Cu, ⁶⁵Cu, and the like. It appreciated that in the case of ¹⁸F, ¹¹C, the imaging isotope may be present on any part of the linker, or alternatively may be present on a structure attached to the linker. For example in the case of ¹⁸F, fluoroaryl groups, such as fluorophenyl, difluorophenyl, fluoronitrophenyl, and the like are described. For example in the case of ¹¹C, alkyl and alkyl aryl are described.

In another aspect, the chemotherapeutic agent is a cytotoxic compound. The cytotoxic compounds described herein operate by any of a large number of mechanisms of action. Generally, cytotoxic compounds disrupt cellular mechanisms that are important for cell survival and/or cell proliferation and/or cause apoptosis.

The drug can be any molecule capable of modulating or otherwise modifying cell function, including pharmaceutically active compounds. Suitable molecules can include, but are not limited to: peptides, oligopeptides, retro-inverso oligopeptides, proteins, protein analogs in which at least one non-peptide linkage replaces a peptide linkage, apoproteins, glycoproteins, enzymes, coenzymes, enzyme inhibitors, amino acids and their derivatives, receptors and other membrane proteins; antigens and antibodies thereto; haptens and antibodies thereto; hormones, lipids, phospholipids, liposomes; toxins; antibiotics; analgesics; bronchodilators; beta-blockers; antimicrobial agents; antihypertensive agents; cardiovascular agents including antiarrhythmics, cardiac glycosides, antianginals and vasodilators; central nervous system agents including stimulants, psychotropics, antiman-

38

ics, and depressants; antiviral agents; antihistamines; cancer drugs including chemotherapeutic agents; tranquilizers; anti-depressants; H-2 antagonists; anticonvulsants; antinauseants; prostaglandins and prostaglandin analogs; muscle relaxants; anti-inflammatory substances; stimulants; decongestants; antiemetics; diuretics; antispasmodics; antiasthmatics; anti-Parkinson agents; expectorants; cough suppressants; mucolytics; and mineral and nutritional additives.

Further, the drug can be any drug known in the art which is cytotoxic, enhances tumor permeability, inhibits tumor cell proliferation, promotes apoptosis, decreases anti-apoptotic activity in target cells, is used to treat diseases caused by infectious agents, enhances an endogenous immune response directed to the pathogenic cells, or is useful for treating a disease state caused by any type of pathogenic cell. Drugs suitable for use in accordance with this invention include adrenocorticoids and corticosteroids, alkylating agents, antiandrogens, antiestrogens, androgens, aclamycin and aclamycin derivatives, estrogens, antimetabolites such as cytosine arabinoside, purine analogs, pyrimidine analogs, and methotrexate, busulfan, carboplatin, chlorambucil, cisplatin and other platinum compounds, taxanes, such as tamoxifen, taxol, paclitaxel, paclitaxel derivatives, Taxotere®, and the like, maytansines and analogs and derivatives thereof, cyclophosphamide, daunomycin, doxorubicin, rhizoxin, T2 toxin, plant alkaloids, prednisone, hydroxyurea, teniposide, mitomycins, discodermolides, microtubule inhibitors, epothilones, tubulysin, cyclopropyl benz[e]indolone, seco-cyclopropyl benz[e]indolone, O—Ac-seco-cyclopropyl benz[e]indolone, bleomycin and any other antibiotic, nitrogen mustards, nitrosureas, vincristine, vinblastine, and analogs and derivative thereof such as deacetylvinblastine monohydrate, colchicine, colchicine derivatives, allocolchicine, thiocolchicine, trityl cysteine, Halicondrin B, dolastatins such as dolastatin 10, amanitins such as α-amanitin, camptothecin, irinotecan, and other camptothecin derivatives thereof, geldanamycin and geldanamycin derivatives, estramustine, nocodazole, MAP4, colcemid, inflammatory and proinflammatory agents, peptide and peptidomimetic signal transduction inhibitors, and any other art-recognized drug or toxin. Other drugs that can be used in accordance with the invention include penicillins, cephalosporins, vancomycin, erythromycin, clindamycin, rifampin, chloramphenicol, aminoglycoside antibiotics, gentamicin, amphotericin B, acyclovir, trifluridine, ganciclovir, zidovudine, amantadine, ribavirin, and any other art-recognized antimicrobial compound.

Illustrative drugs and other therapeutic agents are described in U.S. Patent Application Publication Nos. US-2005-0002942-A1, US-2001-0031252-A1, and US-2003-0086900-A1. Illustrative imaging agents and diagnostic agents are described in U.S. Patent Application Publication No. US-2004-0033195-A1 and International Patent Application Publication No. WO 03/097647. The disclosures of each of the foregoing patent application publications are incorporated herein by reference.

The invention described herein also includes pharmaceutical compositions comprising an amount of a binding ligand (B) drug delivery conjugate effective to eliminate a population of pathogenic cells in a host animal when administered in one or more doses. The binding ligand drug delivery conjugate is preferably administered to the host animal parenterally, e.g., intradermally, subcutaneously, intramuscularly, intraperitoneally, intravenously, or intrathecally. Alternatively, the binding ligand drug delivery conjugate can be administered to the host animal by other medically useful processes, such as orally, and any effective dose and

suitable therapeutic dosage form, including prolonged release dosage forms, can be used.

Examples of parenteral dosage forms include aqueous solutions of the active agent, in an isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carriers such as liquid alcohols, glycols, esters, and amides. The parenteral dosage form in accordance with this invention can be in the form of a reconstitutable lyophilizate comprising the dose of the drug delivery conjugate. In one aspect of the present embodiment, any of a number of prolonged release dosage forms known in the art can be administered such as, for example, the biodegradable carbohydrate matrices described in U.S. Pat. Nos. 4,713,249; 5,266,333; and 5,417,982, the disclosures of which are incorporated herein by reference, or, alternatively, a slow pump (e.g., an osmotic pump) can be used.

In one illustrative aspect, at least one additional composition comprising a therapeutic factor can be administered to the host in combination or as an adjuvant to the above-detailed methodology, to enhance the binding ligand drug delivery conjugate-mediated elimination of the population of pathogenic cells, or more than one additional therapeutic factor can be administered. The therapeutic factor can be selected from a chemotherapeutic agent, or another therapeutic factor capable of complementing the efficacy of the administered binding ligand drug delivery conjugate.

In one illustrative aspect, therapeutically effective combinations of these factors can be used. In one embodiment, for example, therapeutically effective amounts of the therapeutic factor, for example, in amounts ranging from about 0.1 MIU/m²/dose/day to about 15 MIU/m²/dose/day in a multiple dose daily regimen, or for example, in amounts ranging from about 0.1 MIU/m²/dose/day to about 7.5 MIU/m²/dose/day in a multiple dose daily regimen, can be used along with the binding ligand drug delivery conjugates to eliminate, reduce, or neutralize pathogenic cells in a host animal harboring the pathogenic cells (MIU=million international units; m²=approximate body surface area of an average human).

In another embodiment, chemotherapeutic agents, which are, for example, cytotoxic themselves or can work to enhance tumor permeability, are also suitable for use in the method of the invention in combination with the binding ligand drug delivery conjugates. Such chemotherapeutic agents include adrenocorticoids and corticosteroids, alkylating agents, antiandrogens, antiestrogens, androgens, aclamycin and aclamycin derivatives, estrogens, antimetabolites such as cytosine arabinoside, purine analogs, pyrimidine analogs, and methotrexate, busulfan, carboplatin, chlorambucil, cisplatin and other platinum compounds, tamoxifen, taxol, paclitaxel, paclitaxel derivatives, Taxotere®, cyclophosphamide, daunomycin, doxorubicin, rhizoxin, T2 toxin, plant alkaloids, prednisone, hydroxyurea, teniposide, mitomycins, discodermolides, microtubule inhibitors, epothilones, tubulysin, cyclopropyl benz[e]indolone, seco-cyclopropyl benz[e]indolone, O—Ac-seco-cyclopropyl benz[e]indolone, bleomycin and any other antibiotic, nitrogen mustards, nitrosureas, vincristine, vinblastine, and analogs and derivative thereof such as deacetylvinblastine monohydrate, colchicine, colchicine derivatives, allocolchicine, thiocholicine, trityl cysteine, Halicondrin B, dolastatins such as dolastatin 10, amanitins such as α -amanitin, camptothecin, irinotecan, and other camptothecin derivatives thereof, geldanamycin and geldanamycin derivatives, estramustine, nocodazole, MAP4, colcemid, inflammatory and proinflammatory agents, peptide and peptidomimetic signal transduction inhibitors, and any other art-

recognized drug or toxin. Other drugs that can be used in accordance with the invention include penicillins, cephalosporins, vancomycin, erythromycin, clindamycin, rifampin, chloramphenicol, aminoglycoside antibiotics, gentamicin, amphotericin B, acyclovir, trifluridine, ganciclovir, zidovudine, amantadine, ribavirin, maytansines and analogs and derivatives thereof, gemcitabine, and any other art-recognized antimicrobial compound.

The therapeutic factor can be administered to the host animal prior to, after, or at the same time as the binding ligand drug delivery conjugates and the therapeutic factor can be administered as part of the same composition containing the binding ligand drug delivery conjugate or as part of a different composition than the binding ligand drug delivery conjugate. Any such therapeutic composition containing the therapeutic factor at a therapeutically effective dose can be used in the present invention.

Additionally, more than one type of binding ligand drug delivery conjugate can be used. Illustratively, for example, the host animal can be treated with conjugates with different vitamins, but the same drug in a co-dosing protocol. In other embodiments, the host animal can be treated with conjugates comprising the same binding ligand linked to different drugs, or various binding ligands linked to various drugs. In another illustrative embodiment, binding ligand drug delivery conjugates with the same or different vitamins, and the same or different drugs comprising multiple vitamins and multiple drugs as part of the same drug delivery conjugate could be used.

In another illustrative aspect, any effective regimen for administering the binding ligand drug delivery conjugates can be used. For example, the binding ligand drug delivery conjugates can be administered as single doses, or can be divided and administered as a multiple-dose daily regimen. In other embodiments, a staggered regimen, for example, one to three days per week can be used as an alternative to daily treatment, and for the purpose of defining this invention such intermittent or staggered daily regimen is considered to be equivalent to every day treatment and within the scope of this invention. In one embodiment, the host is treated with multiple injections of the binding ligand drug delivery conjugate to eliminate the population of pathogenic cells. In another embodiment, the host is injected multiple times (preferably about 2 up to about 50 times) with the binding ligand drug delivery conjugate, for example, at 12-72 hour intervals or at 48-72 hour intervals. In other embodiments, additional injections of the binding ligand drug delivery conjugate can be administered to the patient at an interval of days or months after the initial injections(s) and the additional injections prevent recurrence of the disease state caused by the pathogenic cells.

Illustratively, the binding ligand drug delivery conjugates can be administered parenterally to the animal or patient suffering from the disease state, for example, intradermally, subcutaneously, intramuscularly, intraperitoneally, or intravenously in combination with a pharmaceutically acceptable carrier. In another embodiment, the binding ligand drug delivery conjugates can be administered to the animal or patient by other medically useful procedures and effective doses can be administered in standard or prolonged release dosage forms. In another aspect, the therapeutic method can be used alone or in combination with other therapeutic methods recognized for treatment of disease states mediated by activated macrophages.

Described herein is a method for imaging pathogenic cell populations that express or over-express PSMA.

Described herein is a method for diagnosing diseases and disease states that are related to pathogenic cell populations that express or over-express PSMA. The compounds described herein bind selectively and/or specifically to cells that express or over-express PSMA. In addition, they not only show selectivity between pathogenic cells and normal tissues, they show selectivity among pathogenic cell populations (see FIG. 8 where PSMA expressing LnCAP cells are preferentially visualized compared to A549 tumors or KB tumors, which are not). In addition, the response is specific to PSMA binding as indicated by competition studies conducted with the conjugates described herein where binding is determined with the conjugate alone or in the presence of excess PMPA, a known binding ligand of PSMA. Binding at both the kidney and tumor is blocked in the presence of excess PMPA (see, for example, Method Examples described herein).

In another embodiment, the conjugate has a binding constant K_d of about 100 nM or less. In another aspect, the conjugate has a binding constant K_d of about 75 nM or less. In another aspect, the conjugate has a binding constant K_d of about 50 nM or less. In another aspect, the conjugate has a binding constant K_d of about 25 nM or less.

In another embodiment, the conjugates described herein exhibit selectivity for PSMA expressing or PSMA over-expressing cells or tissues relative to normal tissues such as blood, heart, lung, liver, spleen, duodenum, skin, muscle, bladder, and prostate, with at least 3-fold selectivity, or at least 5-fold selectivity. In one variation, the conjugates described herein exhibit selectivity for PSMA expressing or PSMA over-expressing cells or tissues relative to normal tissues with at least 10-fold selectivity. It is appreciated that the selectivity observed for imaging is indicative of the selectivity that may be observed in treating disease states responsive to the selective or specific elimination of cells or cell populations that express or over-express PSMA.

The unitary daily dosage of the drug delivery conjugate can vary significantly depending on the host condition, the disease state being treated, the molecular weight of the conjugate, its route of administration and tissue distribution, and the possibility of co-usage of other therapeutic treatments such as radiation therapy. The effective amount to be administered to a patient is based on body surface area, patient weight, and physician assessment of patient condition. Effective doses can range, for example, from about 1 ng/kg to about 1 mg/kg, from about 1 μ g/kg to about 500 μ g/kg, from about 1 μ g/kg to about 100 μ g/kg, and from about 1 μ g/kg to about 10 μ g/kg.

Generally, any manner of forming a conjugate between the bivalent linker (L) and the binding ligand (B), or analog or derivative thereof, between the bivalent linker (L) and the drug, or analog or derivative thereof, including any intervening heteroatoms, can be utilized in accordance with the present invention. Also, any art-recognized method of forming a conjugate between the spacer linker, the releasable linker, and one or more heteroatoms to form the bivalent linker (L) can be used. The conjugate can be formed by direct conjugation of any of these molecules, for example, through hydrogen, ionic, or covalent bonds. Covalent bonding can occur, for example, through the formation of amide, ester, disulfide, or imino bonds between acid, aldehyde, hydroxy, amino, sulfhydryl, or hydrazo groups.

The synthetic methods are chosen depending upon the selection of the optionally included heteroatoms or the heteroatoms that are already present on the spacer linkers, releasable linkers, the drug, and/or the binding ligand. In general, the relevant bond forming reactions are described in

Richard C. Larock, "Comprehensive Organic Transformations, a guide to functional group preparations," VCH Publishers, Inc. New York (1989), and in Theodora E. Greene & Peter G. M. Wuts, "Protective Groups in Organic Synthesis," 2d edition, John Wiley & Sons, Inc. New York (1991), the disclosures of which are incorporated herein by reference.


More specifically, disulfide groups can be generally formed by reacting an alkyl or aryl sulfonylthioalkyl derivative, or the corresponding heteroarylthioalkyl derivative such as a pyridin-2-ylthioalkyl derivative, and the like, with an alkylenethiol derivative. For example, the required alkyl or aryl sulfonylthioalkyl derivative may be prepared according to the method of Ranasinghe and Fuchs, Synth. Commun. 18(3), 227-32 (1988), the disclosure of which is incorporated herein by reference. Other methods of preparing unsymmetrical dialkyl disulfides are based on a trans-thiolation of unsymmetrical heteroaryl-alkyl disulfides, such as 2-thiopyridinyl, 3-nitro-2-thiopyridinyl, and like disulfides, with alkyl thiol, as described in WO 88/01622, European Patent Application No. 0116208A1, and U.S. Pat. No. 4,691,024, the disclosures of which are incorporated herein by reference. Further, carbonates, thiocarbonates, and carbamates can generally be formed by reacting an hydroxy-substituted compound, a thio-substituted compound, or an amine-substituted compound, respectively, with an activated alkoxy-carbonyl derivative having a suitable leaving group.

EXAMPLES

The compounds described herein may be prepared by conventional organic synthetic methods. In addition, the compounds described herein may be prepared as indicated below. Unless otherwise indicated, all starting materials and reagents are available from commercial supplies. All amino acid starting materials were purchased from Chem-Impex Int (Chicago, Ill.). ^1H NMR spectra were obtained using a Bruker 500 MHz cryoprobe, unless otherwise indicated.

Example 1A


General synthesis of PSMA inhibitor intermediates for conjugation. Illustrated for specific synthesis of DUPA derivative 2-[3-(1,3-Bis-tert-butoxycarbonyl-propyl)-ureido]-pentanedioic acid 1-tert-butyl ester (I).



45


46

-continued



47

48




SK28 was synthesized using standard Fluorenylmethyl-oxycarbonyl (Fmoc) solid phase peptide synthesis (SPPS) starting from Fmoc-Cys(Trt)-Wang resin (Novabiochem; Catalog #04-12-2050). SK28 was purified using reverse phase preparative HPLC (Waters, xTerra C₁₈ 10 μm; 19×250 mm) A=0.1 TFA, B=Acetonitrile (ACN); λ=257 nm; Solvent gradient: 5% B to 80% B in 25 min, 80% B wash 30 min run, (61%). Purified compounds were analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×15 mm); A=0.1 TFA, B=ACN; λ=257 nm, 5% B to 80% B in 10 min, 80% B wash 15 min run. C₄₇H₆₅N₉O₁₇S; MW=1060.13 g/mol; white solid; R_f=7.7 min; ¹H NMR


(DMSO-d₆/D₂O) δ 0.93 (m, 2H); 1.08 (m, 5H); 1.27 (m, 5H); 1.69 (m, 2H); 1.90 (m, 2H); 1.94 (m, 2H); 2.10 (m, 2H); 2.24 (q, 2H); 2.62 (m, 2H); 2.78 (m, 4H); 2.88 (dd, 1H); 2.96 (t, 2H); 3.01 (dd, 1H); 3.31 (dd, 1H); 3.62 (dd, 1H); 3.80 (q, 1H, αH); 4.07 (m, 1H, αH); 4.37 (m, 1H, αH); 4.42 (m, 2H, αH); 4.66 (m, 1H, αH); 7.18 (m, 10H, Ar—H); LC-MS=1061 (M+H)₄⁺; ESI-MS=1061 (M+H)⁺.

Example 2AA


The following example compound was synthesized by an analogous process.



The following compounds were synthesized according to the processes described herein using Fmoc SPPS starting from Fmoc-Cys(Trt)-Wang resin (Novabiochem; Catalog #04-12-2050), and purified using reverse phase preparative HPLC (Waters, xTerra C₁₈ 10 μm; 19×250 mm) and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×15 mm):




SK60 (0-atom linker): solvent gradient A=0.1 TFA, B=ACN; λ=220 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 30 min run, (75.3%). C₂₁H₃₂N₆O₁₄S; MW=624.58 g/mol; white solid; R_f=6.3 min; ¹H NMR (DMSO-d₆/D₂O) δ 1.70 (m, 2H); 1.92 (m, 2H); 2.17 (m, 2H); 2.23 (m, 2H); 2.57 (m, 1H); 2.77 (m, 4H); 3.45 (dd, 1H); 3.54 (dd, 1H); 3.83 (t, 1H, αH); 4.06 (m, 1H, αH); 4.38 (m, 1H, α-H); 4.63 (m, 1H, α-H); ESI-MS=625 (M+H)⁺



SK62 (7 atom linker): solvent gradient A=0.1 TFA, TFA, B=ACN; λ=220, 257 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 30 min run, (72%). C₃₅H₄₈N₈O₁₈S; MW=900.86 g/mol; white solid; R_f=8.2 min; ¹H NMR (DMOS-d₆/D₂O) δ 1.62 (m, 1H); 1.70 (m, 2H); 1.79 (m,


1H); 1.90 (m, 2H); 2.09 (t, 2H); 2.16 (m, 2H); 2.24 (m, 2H); 2.60 (m, 1H); 2.75 (m, 4H); 2.81 (m, 1H); 2.97 (m, 1H); 3.33 (dd, 1H); 3.60 (dd, 1H); 3.81 (t, 1H, αH); 4.07 (m, 2H, αH); 4.33 [m, 1H, α-H]; 4.39 (t, α-H); 4.65 (m, 1H, α-H); 7.20 (m, 5H, Ar—H); ESI-MS=901 (M+H)⁺.



51

52


SK38 (16 atom linker): solvent gradient A=10 mM NH₄OAc, B=ACN; λ=257 nm; Solvent gradient: 1% B to 80% B in 25 min, 80% B wash 30 min run, (63%). C₄₃H₆₃N₉O₁₉S, MW=1042.07 g/mol; white solid; R_f=min; ¹H NMR (DMSO-d₆/D₂O) δ 0.94 (m, 2H); 1.08 (m, 5H); 1.27 (m, 5H); 1.66 (m, 2H); 1.70 (m, 2H); 1.79 (m, 1H); 1.90 (m, 2H); 2.09 (t, 2H); 2.74 (m, 2H); 2.84 (m, 1H); 2.95 (t, 3H); 3.07 (d, 2H); 3.23 (m, 1H); 3.43 (dd, 1H); 3.52 (dt, 1H); 3.78 (m, 1H, αH); 3.81 (m, 1H, αH); 3.88 (m, 1H, αH); 4.11 (m, 1H, αH); 4.39 [m, 2H, α-H]; 4.65 (m, 1H, α-H); 7.14 (m, 1H, Ar-H); 7.21 (m, 4H, Ar-H); ESI-MS=1043 (M+H)⁺.



SK57 (24 atom linker): solvent gradient A=0.1 TFA, B=ACN; λ=257 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 30 min run, (56%). C₄₅H₇₀N₈O₂₂S, MW=1107.14 g/mol; colorless solid; ¹H NMR (DMSO-d₆/D₂O) δ 1.66 (m, 2H); 2.07 (m, 4H); 2.31 (t, 1H); 2.43 (m, 1H); 2.77 (m, 2H); 2.98 (dd, 1H); 3.14 (t, 2H); 3.24 (d, 1H); 3.40 (m, 4H, PEG-H); 3.46 (s, 24H, PEG-H); 3.78 (t, 1H); 3.81 (t, 1H); 4.03 (m, 1H, αH); 4.40 (m, 2H, α-H); 7.16 (m, 1H, Ar-H); 7.22 (m, 4H, Ar-H); ESI-MS=1108 (M+H)⁺.

Example 2F

The following compound may be synthesized according to the processes described herein.




(9-atom linker)

53


Example 3A

General process for adding radionuclide to chelating group. Illustrated for radio labeling of SK28 with ^{99m}Tc to prepare SK33.




SnCl_2 /Sodium glucoheptanoate
 $\text{NaHCO}_3/\text{pH} = 6.8$

Sodium pertechnetate



+



Preparation of SK28 formulation kits. HPLC grade Milipore filtered water (50 mL) was added to a 100 mL bottle and argon was purged for at least 10 min. Sodium α -D-glucoheptanoate dihydrate (800 mg) was dissolved in argon purged water (5 mL). Stannous chloride dihydrate (10 mg) was dissolved in 0.02 M HCl (10 mL) while bubbling argon. Stannous chloride (0.8 mL) was added to the sodium glucoheptanoate solution under argon. SK28 (1.4 mg) was added

to the sodium glucoheptanoate/stannous chloride solution under argon. The pH of the reaction mixture was adjusted to 6.8 ± 0.2 using 0.1 N NaOH. Argon purged water (5.2 mL) was added to the reaction mixture to make total volume as 10 mL. 1.0 mL of reaction mixture was dispensed to each

60 vial (10 vials) under argon atmosphere and lyophilized for 36-48 h. The vials were sealed with rubber stoppers and aluminum seals under argon atmosphere to make SK28 formulation kits. The formulation kit vials were stored at -20°C . until they used.

65 Labeling SK28 with ^{99m}Tc . Radio labeling of SK28 with ^{99m}Tc may be performed according to published procedures. A formulation vial was warmed to room temperature for 10

55




min and heated in a boiling water bath for 3 min. Then 15 mCi of sodium pertechnetate ^{99m}Tc (1.0 mL) was injected and an equal volume of gas was withdrawn from the vial to normalize the pressure. The vial was heated in the boiling water bath for 15-20 min and then cooled to room temperature before using in the experiment. Radiochemical purity was analyzed by radioactive TLC (>98%), that showed syn

56

and anti isomers of the radio labeled compound (SK33/SK28- ^{99m}Tc).

Examples 3B-3E


⁵ The following Examples were prepared according to the processes described herein (both syn and anti isomers were obtained; only the syn isomer is shown):

SK60- ^{99m}Tc (0-atom linker, 1.0 mg)SK62- ^{99m}Tc (7-atom linker, 1.0 mg)SK38- ^{99m}Tc (16-atom linker, 1.4 mg)

57


58

-continued

SK57-^{99m}Tc (24-atom linker, 1.8 mg)

Example 3F

The following compound may be synthesized according²⁰ to the processes described herein.




(9-atom linker)


45

Example 4

General synthesis of PSMA imaging agent conjugates illustrated for SK59 using Universal PSMA (DUPA) resin, a 2-atom linker, and FITC. This conjugate may also be used⁵⁰ for detecting circulating tumor cells in prostate cancer patients.



1) 20% piperidine in DMF
2) PSMA-I, HATU, DIPEA/DMF



Synthesis of PSMA universal resin and SK59. Universal PSMA ligand (DUPA) resin was synthesized using Universal NovaTag™ resin (Novabiochem; Catalog #04-12-3910). Fmoc group was deprotected using 20% piperidine/DMF (N,N-dimethylformamide), after swelling the resin with DCM (CH₂Cl₂) and DMF. tert-Butyl protected DUPA was coupled using HATU [2-(1H-7-azabenzotriazol-1-yl)-1,3,3-tetramethyl uronium hexafluorophosphate] and DIPEA (N,N-diisopropylethylamine) in DMF. The pendant Mmt (4-Methoxytrityl) was removed with 1M HOBt (1-Hydroxybenzotriazole) in DCM/TFE (trifluoroethanol). The resin intermediate can be washed with DMF and used immediately in subsequent synthetic steps or washed with DCM/DMF and then with MeOH, and dried for later use.

Universal PSMA resin was reacted with commercially available FITC (1.25 equiv) in the presence of DIPEA (4 equiv) in DMF to yield SK59 (2 atom linker) construct. The final compound was cleaved from the resin using a mixture


of TFA (trifluoro acetic acid), TIPS (triisopropylsilane), and water. Purification was by reverse phase preparative HPLC (Waters, xTerra C₁₈ 5 μm; 19×150 mm) A=10 mM NH₄OAc, B=ACN; λ=488 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 40 min run, (63%). SK59 was analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×15 mm); A=10 mM NH₄OAc, B=ACN; λ=488 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run; C₃₄H₃₃N₆O₁₃ S; MW=751.72 g/mol; orange color solid, R_f=7.2 min; ESI-MS=752 (M+H)⁺; 774 (M+Na)⁺; 750 (M-H)⁻.

Example 5A

General synthesis of PSMA imaging agent conjugates illustrated for SK64 using Universal PSMA (DUPA) resin, a 16-atom linker, and FITC.

61

62




Universal PSMA resin was coupled with Fmoc-Glu-(OtBu)-OH and Fmoc-EAOA (8-aminooctanoic acid) using standard Fmoc SPPS. After conjugating with fluorescein-isothiocyanate (1.25 equiv) in the presence of DIPEA (4 equiv) in DMF, SK64 (16 atom linker) compound was cleaved from the resin using TFA/TIPS/H₂O. Purification was performed using reverse phase preparative HPLC (Waters, xTerra C₁₈ 5 μm; 19×150 mm) A=10 mM NH₄OAc, B=ACN; λ=488 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 40 min run, (57%). SK64 was analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈

5 μm; 3.0×150 mm); A=10 mM NH₄OAc, B=ACN; λ=488 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run; C₄₇H₅₅N₇O₁₇S; MW=1022.04 g/mol; orange color solid, R_f=7.8 min; ESI-MS=1022 (M+H)⁺; 1020 (M-H)⁻.


Examples 5B-5C

The following compounds were prepared using the synthetic processes described herein:

63




64



SK63 (7-atom linker, C₃₉H₄₀N₆O₁₇, Mol. Wt.: 864.76) was prepared using universal PSMA resin and standard Fmoc SPPS conjugated with Fmoc-Glu-(O^tBu)-OH. After coupling with FITC, compounds were cleaved from the resin using TFA/TIPS/H₂O cocktail and purified with reverse phase preparative HPLC (Waters, xTerra C₁₈ 5 μm; 19×150 mm) A=10 mM NH₄OAc, B=ACN; λ=488 nm;

20 Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 40 min run, (65%); analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×150 mm); A=10 mM NH₄OAc, B=ACN; λ=488 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run; SK63: C₃₉H₄₀N₆O₁₆S; MW=880.83 g/mol; orange color solid, R_f=6.8 min; ESI-MS=881 (M+H)⁺; 903 (M+Na)⁺; 863 (M-H)⁻.



50 SK58 (24-atom linker, C₄₉H₆₂N₆O₂₀S, Mol. Wt.: 1087.11) was prepared using universal PSMA resin and standard Fmoc SPPS conjugated with Fmoc-(PEG)₅-OH and purified by HPLC 1% B to 60% B in 25 min, 80% B wash 40 min run, (65%); analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×150 mm); A=10 mM NH₄OAc, B=ACN; λ=488 nm, 1% B to 60% B in 10 min, 80% B wash 15 min run; SK58: C₄₉H₆₀N₆O₂₀S; MW=1087.11 g/mol; orange color solid, R_f=7.3 min; ESI-MS=1087 (M+H)⁺; 1109 (M+Na)⁺; 1085 (M-H)⁻.


60

Example 6A

65 General synthesis of Cys-maleimide PSMA imaging agent conjugates illustrated for SK56 using Wang PSMA (DUPA) resin, a 28-atom linker, and Oregon Green 488, where n=3.


65

66



PC01SK54
 $C_{38}H_{59}N_5O_{18}S$
 Mol. Wt.: 905.96

Oregon Green 488 maleimide/THF
 Water/pH~7/Ar



PC01SK56
 $C_{62}H_{70}F_2N_6O_{25}S$
 Mol. Wt.: 1369.31

Related analogs where n is an integer from 4 to about 30 may also be prepared according to the processes described herein.

SK54 was prepared using standard Fmoc SPPS starting from Fmoc-Cys(Trt)-Wang resin (Novabiochem; Catalog #04-12-2050), purified using reverse phase preparative HPLC (Waters, xTerra C_{18} 10 μ m; 19x250 mm) A=0.1 TFA; B=ACN; λ =257 nm; Solvent gradient: 1% B to 60% B in 25 min, 80% B wash 40 min run, (63%), and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C_{18} 5 μ m; 3.0x50 mm); A=10 mM NH_4OAc , B=ACN; λ =257 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run; $C_{38}H_{59}N_5O_{18}S$, MW=905.96 g/mol; colorless solid; R_f =9.2 min, LC-MS=906.3 g/mol; ESI-MS=906 (M+H)⁺; 904 (M-H)⁻.

SK56 (24 atom linker). HPLC grade Milli-Q water and satd $NaHCO_3$ were purged with argon for 10 min. SK54 was dissolved in 1.0 mL of argon purged water while bubbling argon. The pH of the solution was increased up to 6.8 and Oregon Green 488 maleimide dissolved in 1.0 mL of THF was added to the reaction mixture. The reaction was monitored by analytical HPLC (10 mM NH_4OAc , pH=7.0; 1% B to 50% B in 10 min 80% B wash 15 min run) and reaction

was completed within 10 min. THF was evaporated and reaction mixture was diluted with 5.0 mL of 7 mM phosphate buffer. Purification was performed using reverse phase preparative HPLC (Waters, xTerra C_{18} 10 μ m; 19x250 mm) A=7 mM Phosphate buffer pH=7.2, B=ACN; λ =488 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash 40 min run, (89%); and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C_{18} 5 μ m; 3.0x150 mm); A=10 mM NH_4OAc , B=ACN; λ =488 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run; $C_{62}H_{70}F_2N_6O_{25}S$; MW=1369.31 g/mol; orange color solid, R_f =7.0 min; LC-MS=1370.2; ESI-MS=1391 (M+Na)⁺.


The following 24-atom linker compounds were prepared in an analogous manner to those described herein using the General syntheses described herein.

Example 6B

The following AlexaFluor 488 conjugate compound was prepared according to the processes described herein starting with SK55, where n=3.

67


68



Related analogs where n is an integer from 4 to about 30 may also be prepared according to the processes described herein. 40

Examples 7A-7C


The following DUPA imaging agent conjugate compounds, SK51, SK45, and SK49 were prepared according to the processes described herein, where n is 5:



SK51 (25-Atom Linker, and AlexaFluor 647, MW ~2300 (Commercially Available from Invitrogen))

65

Related analogs where n is an integer from 0 to about 12 may also be prepared according to the processes described herein.




SK45 (25 Atom Linker BODIPY 505,
C₆₇H₈₇BF₂N₁₃O₂₀S, Mol. Wt.: 1475.35)

Related analogs where n is an integer from 0 to about 12 may also be prepared according to the processes described herein.

α H); 4.66 (m, 1H, α H); 7.18 (m, 10H, Ar—H); LC-MS=1061 (M+H)₊; ESI-MS=1061 (M+H)⁺.

Synthesis of SK51 (AlexaFluor 647 conjugate), SK45 (BODIPY conjugate) and SK49 (Oregon Green 488 conjugate). HPLC grade Milli-Q water and satd NaHCO₃ were



SK49 (25 Atom Linker-Oregon Green 488,
C₇₁H₇₆F₂N₁₀O₂₄, Mol. Wt.: 1523.48)

Related analogs where n is an integer from 0 to about 12 may also be prepared according to the processes described herein.

Synthesis of the Linker. In each of the foregoing Examples, the linker was synthesized using standard Fmoc SPPS starting from Fmoc-Cys(Trt)-Wang resin (Novabiochem; Catalog #04-12-2050); C₄₇H₆₅N₂O₁₇S; MW=1060.13 g/mol; white solid; R_f=7.7 min; ¹H NMR (DMSO-d₆/D₂O) δ 0.93 (m, 2H); 1.08 (m, 5H); 1.27 (m, 5H); 1.69 (m, 2H); 1.90 (m, 2H); 1.94 (m, 2H); 2.10 (m, 2H); 2.24 (q, 2H); 2.62 (m, 2H); 2.78 (m, 4H); 2.88 (dd, 1H); 2.96 (t, 2H); 3.01 (dd, 1H); 3.31 (dd, 1H); 3.62 (dd, 1H); 3.80 (q, 1H, α H); 4.07 (m, 1H, α H); 4.37 (m, 1H, α H); 4.42 (m, 2H,

purged with argon for 10 min. Linker was dissolved in 1.0 mL of argon purged while bubbling argon. The pH of the solution was increased to 6.8 and AlexaFluor maleimide, BODIPY maleimide, or Oregon green 488 maleimide, respectively, was dissolved in 1.0 mL of tetrahydrofuran (THF) was added to the reaction mixture. Progress of the reaction was monitored by analytical HPLC (10 mM NH₄OAc, pH=7.0; 1% B to 50% B in 10 min 80% B wash 15 min run) and reaction was completed within 10 min. THF was evaporated and reaction mixture was diluted with 5.0 mL of 1 mM phosphate buffer (pH=7.2).

Compounds were purified using reverse phase preparative HPLC (Waters, xTerra C₁₈ 5 μ m; 18x150 mm) A=1 mM Phosphate buffer pH=7.2, B=ACN; λ =647 or 488 nm; Solvent gradient: 1% B to 50% B in 25 min, 80% B wash

71

40 min run; and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×50 mm); A=10 mM NH₄OAc, B=ACN; λ=588 or 488 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run.

SK51: MW~2360.13 g/mol; blue color solid, R_f=6.7 min; 5 (structure of the AlexaFluor 647 is not known);


SK45: C₆₇H₈₇BF₂N₁₃O₂₀S; MW=1475.35 g/mol; orange color solid, R_f=7.6 min; LC-MS=1475.3 (M+H)⁺;

72

SK49: C₇₁H₇₆F₂N₁₀O₂₄S; MW=1523.48 g/mol; orange color solid, R_f=6.7 min; LC-MS=1524 (M+H)⁺.

Example 8A

General synthesis of PSMA disulfide linker intermediate for releasable agent conjugate, illustrated for SK68.



PC01SK68
C₃₂H₄₂N₆O₁₇S
Mol. Wt.: 814.77

73


SK68 was synthesized using standard Fmoc SPPS starting from Fmoc-Cys(Trt)-Wang resin (Novabiochem; Catalog #04-12-2050), purified using reverse phase preparative HPLC (Waters, xTerra C₁₈ 10 μm; 19×250 mm) A=0.1 TFA, B=ACN; λ=257 nm; Solvent gradient: 1% B to 50% B in 30 min, 80% B wash 40 min run, (68%); and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C₁₈ 5 μm; 3.0×15 mm); A=0.1 TFA, B=ACN; λ=257 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run. C₃₂H₄₂N₆O₁₇S; MW=814.77 g/mol; white solid; R_f=8.2 min; ¹H NMR (DMOS-d₆/D₂O) δ 1.70 (m, 3H); 1.90 (m, 3H); 2.10 (m,

74

2H); 2.17 (m, 2H); 2.23 (m, 2H); 2.36 (m, 1H); 2.59 (dd, 1H); 2.79 (m, 3H); 3.04 (dd, 1H); 4.07 (m, 2H, αH); 4.13 (m, 1H, αH); 4.37 [m, 1H, α-H]; 4.47 (m, 2H, α-H); 7.19 (m, 5H, Ar—H); 7.87 (d, Ures-NH); 8.20 (d, 1H, Urea-NH); LC-MS=815.3 (M+H)⁺.


Example 8B

General synthesis of PSMA disulfide linker intermediate for releasable agent conjugate, illustrated for SK28L.





77


78



Tubulysin B
 $C_{42}H_{63}N_5O_{10}S$
 Mol. Wt.: 830.44




EC0312
 $C_{50}H_{72}N_8O_{11}S_3$
 Mol. Wt.: 1057.35



79

80

-continued



PC01SK71
 $C_{77}H_{109}N_{13}O_{28}S_3$
 Mol. Wt.: 1760.95


Synthesis of EC0312. Tubulysin B (30 mg, 0.036 mmol) was dissolved in ethylacetate (600 μ L) under argon at -15° C. Isobutyl chloroformate (4.7 μ L, 0.054 mmol) and diisopropylethylamine (13.2 μ L, 0.076 mmol) were added to the reaction mixture; reaction was stirred at -15° C. for 45 min under argon. EC0311 (13.4 mg, 0.054 mmol) dissolved in ethylacetate (500 μ L) was added. Reaction mixture was stirred at -15° C. for another 15 min and then at room temperature for 45 min. Solvent was evaporated and residue was purified using short column (2%-8% methanol in CH_2Cl_2) to get EC0312 (34.4 mg, 90.5%). EC0312 was characterized using NMR (Varian 300 MHz, in $CDCl_3$), and LC-MS=1058.3 (M+H)⁺.

Synthesis of SK71. HPLC grade Milli-Q water and satd $NaHCO_3$ were purged with argon for 10 min. SK68 was dissolved in 1.0 mL of argon purged water while bubbling argon through the solution. The pH of the solution was increased to 6.8 using argon purged $NaHCO_3$ and EC0312 dissolved in THF (2.0 mL) was added to the reaction

HPLC (Waters, xTerra C_{18} 10 μ m; 19 \times 250 mm) A=2 mM Phosphate buffer, B=ACN; λ =254 nm; 5% B to 80% B in 25 min 80% B wash 40 min run; and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C_{18} 5 μ m; 3.0 \times 15 mm); A=10 mM NH_4OAc , B=ACN; λ =254 nm, 1% B to 50% B in 10 min, 80% B wash 15 min run. $C_{77}H_{109}N_{13}O_{28}S_3$; MW=1760.95 g/mol; white color solid, R_f =7.6 min; 1H NMR (DMSO- d_6 /D $_2$ O) was consistent with the SK71 structure; HRMS (MALDI) (m/z): (M-H)⁻ calcd. for, $C_{77}H_{110}N_{13}O_{28}S_3$, 1758.6594; found, 1758.7033; LRMS (LCMS) (m/z): (M+H)⁺ calcd. for 1761.9; found, 1761.8; UV/Vis: λ_{max} =254 nm.

Example 9B

Similarly, the D-Cys analog of SK71 was prepared as described herein.



mixture. Progress of the reaction was monitored by analytical HPLC (10 mM NH_4OAc , pH=7.0; λ =254; 1% B to 50% B in 10 min 80% B wash 15 min run) and reaction was completed within 10 min. THF was evaporated and reaction mixture was diluted with 5.0 mL of 2 mM phosphate buffer. SK71 (61.3%) was purified using reverse phase preparative


Example 9C

General synthesis for preparing disulfide-linked conjugates, illustrated for tubulysin B conjugate SK77 (31-atom linker).

81


82

Scheme 6.4




Tubulysin B
 $C_{42}H_{63}N_5O_{10}S$
 Mol. Wt.: 830.44


1) DIPEA/Isobutylchloroformate
 EtOAc/-15° C.



EC0311
 $C_8H_{11}N_3O_2S_2$
 Mol. Wt.: 245.32



EC0312
 $C_{50}H_{72}N_8O_{11}S_3$
 Mol. Wt.: 1057.35




PC01SK28
 $C_{47}H_{65}N_9O_{17}S$
 Mol. Wt.: 1060.13

83

84

-continued



PC01SK77
 $C_{93}H_{133}N_{16}O_{28}S_3$
 Mol. Wt.: 2006.32

15


HPLC grade Milli-Q water and satd $NaHCO_3$ were purged with argon for 10 min. SK68 was dissolved in 1.0 mL of argon purged water while bubbling argon. The pH of the solution was increased to 6.8 using argon purged $NaHCO_3$ and EC0312 dissolved in THF (2.0 mL) was added to the reaction mixture. Progress of the reaction was monitored by analytical HPLC (10 mM NH_4OAc , pH=7.0; $\lambda=254$; 1% B to 50% B in 10 min 80% B wash 15 min run) and reaction was completed within 10 min. THF was evaporated and reaction mixture was diluted with 5.0 mL of 2 mM phosphate buffer. SK77 (61%) was purified using reverse phase preparative HPLC (Waters, xTerra C_{18} 10 μm ; 19x250 mm) A=2 mM phosphate buffer, B=ACN; $\lambda=254$ nm; 5% B to

80% B in 25 min 80% B wash 40 min run; and analyzed using reverse phase analytical HPLC (Waters, X-Bridge C_{18} 5 μm ; 3.0x15 mm); A=10 mM NH_4OAc , B=ACN; $\lambda=254$ nm, 1% B to 50% B in 10 min, 80% B wash 15 min run. $C_{93}H_{133}N_{16}O_{28}S_3$; MW=2006.32 g/mol; white color solid, $R_f=7.7$ min; 1H NMR (DMSO- d_6 /D $_2$ O); LC-MS=2007.0 (M+H) $^+$.

25

Example 9D


Similarly, the D-Cys analog of SK77 was prepared as described herein.



45

Example 9E

Similarly, the D-Cys, propanoic acid analog of SK77 was prepared as described herein.




Examples 9F-9G

The following DUPA vinblastine and DUPA camptothecin compounds, SK37 and SK45, respectively, were prepared according to the processes described herein.


65

85

86



SK37 (vinblastine conjugate, $C_{93}H_{123}N_{15}O_{26}S_2$, Mol. Wt.: 1931.19) prepared in 63.1% yield. $C_{93}H_{123}N_{15}O_{26}S_2$: MW=1931.19 g/mol; white color solid, $R_f=7.7$ min; 1H NMR (DMSO- d_6/D_2O); LC-MS=1932.6 (M+H) $^+$.




SK45 (camptothecin conjugate, $C_{70}H_{83}N_{11}O_{23}S_2$, Mol. Wt.: 1510.60) prepared in 66% yield. $C_{70}H_{83}N_{11}O_{23}S_2$: MW=1510.60 g/mol; white color solid, $R_f=7.5$ min; 1H NMR (DMSO- d_6/D_2O); LC-MS=1511.1 (M+H) $^+$.


Example 9H

50


Similarly, the Glu-Asp-Phe analog of SK37 was prepared as described herein.




The following compounds were prepared using the synthetic processes described herein:




SK125 (FITC conjugate)



SK131 (rhodamine conjugate)



SK179 (FITC conjugate)




FITC conjugate


89

90


-continued



DyLight 680 conjugate



DyLight 800 conjugate




PET agent conjugate


91

92



-continued




PET agent conjugate



PET agent conjugate


DOTA conjugate capable of chelating for example ^{64}Cu , ^{65}Cu , and the likeDOTA conjugate capable of chelating for example ^{64}Cu , ^{65}Cu , and the like

93



DTPA conjugate capable of chelating for example In, Ga, Ir, Yr, and the like

94



Tripeptide conjugate capable of chelating for example Tc, Tc oxides, and the like

The foregoing exemplary embodiments are intended to be illustrative of the invention, and should not be interpreted or construed as limiting in any way the invention as described herein.

METHOD EXAMPLES

Example 1A

In Vitro Binding Studies Using Lncap Cells and SK28 (14 Atom Spacer). LNCaP cells (a human prostate cancer cell line over-expressing PSMA, purchased from American Type Culture Collection (ATCC)) were seeded in two 24-well (120,000 cells/well) falcon plates and allowed to grow to adherent monolayers for 48 hours in RPMI with glutamine (2 mM)(Gibco RPMI medium 1640, catalog #22400) plus 10% FBS (Fetal Bovine Serum), 1% sodium pyruvate (100 mM) and 1% PS (penicillin streptomycin) in a 5%-CO₂ atmosphere at 37° C. Cells of one 24-well plate were incubated with increasing concentrations of SK28-99mTc from 0 nM-450 nM (triplicates for each concentration) in a 5%-CO₂ atmosphere at 37° C. for 1 hour. Cells of the second 24-well plate were incubated with 50 uM PMPA in a 5%-CO₂ atmosphere at 37° C. for 30 minutes, then incubated with increasing concentrations of SK28-99mTc from 0 nM-450 nM (triplicates for each concentration) in a 5%-CO₂ atmosphere at 37° C. for 1 hour (competition study). Cells were rinsed three times with 1.0 mL of RPMI. Cells were lysed with tris-buffer, transferred to individual gamma scintigraphy vials, and radioactivity was counted. The plot of cell bound radioactivity verses concentration of radiolabeled compound was used to calculate the K_d value. The competition study was used to determine the binding specificity of the ligand (DUPA) to the PSMA (FIG. 1A).

Example 1B

In Vitro Binding Studies Using LNCaP Cells and SK33 (14 atom spacer). LNCaP cells (150,000 cells/well) were seeded onto 24-well Falcon plates and allowed to form confluent monolayers over 48 h. Spent medium in each well was replaced with fresh medium (0.5 mL) containing increasing concentrations of DUPA-99mTc in the presence (▲) or absence (■) of excess PMPA. After incubating for 1 h at 37° C., cells were rinsed with culture medium (2×1.0 mL) and tris buffer (1×1.0 mL) to remove any unbound radioactivity. After suspending cells in tris buffer (0.5 mL), cell bound radioactivity was counted using a γ -counter (Packard, Packard Instrument Company). The dissociation constant (K_D) was calculated using a plot of cell bound radioactivity versus the concentration of the radiotracer using nonlinear regression in GraphPad Prism 4. Error bars represent 1 standard deviation (n=3). Experiment was performed three times with similar results. (FIG. 1B).

Example 2

Quantification of PSMA Molecules on LNCaP Cells. LNCaP cells were seeded in a 24-well falcon plate and allowed to grow to adherent monolayers for 48 hours in RPMI (Gibco RPMI medium 1640, catalog #22400) plus 10% FBS (Fetal Bovine Serum), 1% glutaric and 1% PS (penicillin streptomycin) in a 5%-CO₂ atmosphere at 37° C. Cells were then incubated with increasing concentrations of SK28-99mTc from 0 nM-450 nM (triplicates for each concentration) in a 5%-CO₂ atmosphere at 4° C. or at 37° C. for 1 hour. Cells were rinsed three times with 1.0 mL of RPMI. Cells were lysed with tris-buffer, transferred to individual gamma scintigraphy vials, and radioactivity was counted. The plot of cell bound radioactivity verses concentration of

radiolabeled compound was used to calculate number of PSMA/LNCaP cell. The radioactivity of a 30 nM sample of SK28-99mTc (20 uL) was counted. At 4° C. (to prevent endocytosis of PSMA), the number of moles in the 30 nM sample = $30 \text{ nM} \times 20 \text{ uL} = (30 \times 10^{-9} \text{ mol/L}) \times (20 \times 10^{-6} \text{ L}) = 6 \times 10^{-13} \text{ mol}$. The number of atoms in the 30 nM sample = $(6 \times 10^{-13} \text{ mol}) \times (6.023 \times 10^{23} \text{ atom/mol}) = 3.6 \times 10^{11} \text{ atom}$. The radio count of 20 uL of the 30 nM sample = 20477 cpm (cpm/atom = $3.6 \times 10^{11} / 20477 = 1.76 \times 10^7$). The cell bound radioactivity at the saturation point at 4° C. = 12 000 cpm. The number of atoms at the saturation point = $(1.76 \times 10^7 \text{ atom}) \times (12 \text{ 000 cpm})$. The number of cells/well = 245,000. The number of PSMA/cell at 4° C. = $(2.12 \times 10^{11}) / 2.45 \times 10^5 = 864 \text{ 396.4} \sim 0.9 \times 10^6$ PSMA/LNCaP cell.

The cell bound radioactivity at the saturation point at 37° C. = 33,000 cpm (approximately three fold higher than at 4° C.). This shows that PSMA undergoes endocytosis, unloading the drug and recycling, similar to cell surface receptors. See FIG. 2.

Example 3

Spacer-Dependent Binding Studies. LNCaP cells were seeded in 24-well (120,000 cells/plate) falcon plates (10 plates) and allowed to grow to adherent monolayers for 48 hours in RPMI (Gibco RPMI medium 1640, catalog #22400) plus 10% FBS (Fetal Bovine Serum), 1% sodium pyruvate and 1% PS (penicillin streptomycin) in a 5%-CO₂ atmosphere at 37° C. Cells were then incubated with increasing concentrations of SK60-99mTc (zero atom spacer), SK62-99mTc (7 atom spacer), SK28-99mTc (14 atom spacer), SK38-99mTc (16 atom spacer) and SK57-99mTc (24 atom spacer) from 0 nM-1280 nM (triplicates for each concentration) in a 5%-CO₂ atmosphere at 37° C. for 1 hour. Also, in separate plates, cells was incubated with 50 uM PMPA in a 5%-CO₂ atmosphere at 37° C. for 30 minutes and then incubated with increasing concentration of SK60-99mTc (zero atom spacer), SK62-99mTc (7 atom spacer), SK28-99mTc (14 atom spacer), SK38-99mTc (16 atom spacer) and SK57-99mTc (24 atom spacer) from 0 nM-1280 nM (triplicates for each concentration) in a 5%-CO₂ atmosphere at 37° C. for 1 hour (competition studies; data not shown). Cells were rinsed three times with 1.0 mL of RPMI. Cells were lysed with tris-buffer, transferred to individual gamma scintigraphy vials, and radioactivity was counted. The plot of cell bound radioactivity verses concentration of the radiolabeled compound was used to calculate the K_d value. The plot of % saturation verses concentration of the radiolabeled compound as well as the plot for K_d verses spacer length are shown (FIGS. 3A and B).

Example 4

In Vivo Growth of Human LNCaP Tumor Cells in Nude Mice. LNCaP cells were maintained in RPMI 1640 (Gibco RPMI medium 1640, catalog #22400) with glutamine (2 mM), 10% FBS (Fetal Bovine Serum), 1% sodium pyruvate (100 mM) and 1% PS (penicillin streptomycin) in a 5%-CO₂ atmosphere at 37° C. Four to five week-old athymic male nude mice (nu/nu) were obtained from the NCI Charles River and maintained in a sterile environment. Mice were housed in polycarbonate shoebox cages with wire top lids and maintained on a normal diet. Mice were allowed to acclimate for one week prior to inoculation of LNCaP cells. Matrigel and high concentrated (HC) matrigel were purchased from BD Biosciences. Nude mice were inoculated with either 2.5×10^6 or 5.0×10^6 in vitro propagated LNCaP

cells in 50% matrigel (100 uL RPMI medium+100 uL of matrigel) or 50% high concentrated matrigel (100 uL RPMI medium+100 uL of HC matrigel) to determine optimal conditions, including number of cells, vehicle, etc. Cells were subcutaneously injected into each axial and each flank of the nude mice to determine the optimal site. The volume of each tumor was measured in perpendicular directions twice a week using a caliper and body weight was measured once a week (data not shown).

The volume of each tumor was calculated as $0.5 \times L \times W^2$, where L = measurement of longest axis in millimeters and W = measurement of axis perpendicular to L in millimeters. Approximately 5.0×10^6 LNCaP cells in 50% HC matrigel on the axial gave 600 mm³ tumors within 30 days. See FIG. 4.

Example 5

Comparison of Incap, kb and a549 Cell Tumor Growth in Mice. LNCaP, KB, and A549 cells were maintained in RPMI 1640 (Gibco RPMI medium 1640, catalog #22400) with glutamine (2 mM), 10% FBS (Fetal Bovine Serum), 1% sodium pyruvate (100 mM) and 1% PS (penicillin streptomycin) in a 5%-CO₂ atmosphere at 37° C. Four-five weeks old male nude mice (nu/nu) were obtained from the NCI Charles River and maintained in a sterile environment. Mice were housed in polycarbonate shoebox cages with wire top lids and maintained on a normal diet. Mice were allowed to acclimate for one week prior to inoculation of cells.

For tumor cell inoculation, 5.0×10^6 LNCaP cells in 50% high concentrated matrigel, 1.0×10^6 KB cells in RPMI medium, or 1.0×10^6 A549 cells in RPMI medium were subcutaneously injected into the right axial (some animals were injected in both) of the nude mice. The volume of each tumor was measured in two perpendicular directions twice a week using a caliper (See FIGS. 5A and 5B) and body weight was measured once a week (data not shown). The volume of the tumors were calculated as $0.5 \times L \times W^2$, where L = measurement of longest axis in millimeters and W = measurement of axis perpendicular to L in millimeters.

Example 6A

In Vivo Imaging of Tumors in Mouse Using PSMA-99mTc. When tumors reached a volume of between 500-600 mm³, 99mTc-labeled compounds (e.g. SK28-99mTc, SK60-99mTc, etc) prepared as described, were administered through intraperitoneal injection (subcutaneously). Four hours later, animals were euthanized and blood was taken by cardio punch and transferred to individual gamma scintigraphy vials per each animal. The imaging experiments were carried out using either a Kodak or gamma scintigraphic camera imager (FIGS. 6A, 6B, 6C, 7A, 7B and 7C). [Note: PMPA was injected 30 minutes before injecting SK28-99mTc. Other than uptake in the cancerous masses, SK28-99mTc distribution was limited to kidneys (FIGS. 6A, 6B and 6C). Both mice (FIGS. 7A, 7B and 7C) were injected with SK60-99mTc and distribution was limited mostly to the kidneys (no tumor uptake even after shielding both kidneys.)]

FIGS. 6A, 6B, and 6C show images of mice with human LNCaP tumors using SK28-99mTc (radiolabeled 14 Atom spacer). FIGS. 6A-6C represent 3 separate sets of mice: the left hand image shows white light images and the right hand image shows an overlay of the radioimage with the white light image of mice with LNCaP tumors imaged using a Kodak camera imager 4 hours after subcutaneous (admin-

97

istered through intraperitoneal) injection of 1 ng/kg SK28-99mTc without [left mouse in each set of images] and with 50 mg/kg PMPA [right mouse in both sets of images] to block PSMA (as a competitor). FIG. 6D shows a single mouse study for LNCaP tumors imaged using Kodak imager 4 hours after subcutaneous (administered through intraperitoneal) injection of 1 ng/kg SK28-99mTc showing in the left hand image an overlay of radioimage with kidney shield and white light image with no shield and in the right hand image an overlay of radioimage with kidney shield and X-ray image with no shield.

Example 6B

In Vivo Imaging of Tumors in Mouse Using dupa-99mTc. To further establish the specificity of our DUPA conjugates for prostate cancer cells, DUPA-99mTc was injected intraperitoneally (i.p.) into athymic nude mice bearing LNCaP tumors on their shoulders. After 4 h to allow for clearance of unbound conjugate, the distribution of the retained DUPA-99mTc was imaged by gamma scintigraphy. As seen in FIGS. 7D(a) and 7D(c), the targeted 99mTc radiotracer accumulated mainly in the PSMA positive LNCaP tumor, with little or no radioactivity in other tissues except the kidneys. Importantly, kidney uptake may be peculiar to the mouse, since immunohistochemical and RT-PCR analyses suggest that PSMA expression is high in murine kidneys but minimal in human kidneys. In vivo specificity of the PSMA-targeted imaging agent was further tested by prior administration of excess PMPA to block all PSMA sites before DUPA-99mTc administration. As shown in FIGS. 7D(b) and 7D(d), blocked LNCaP tumors display no DUPA-99mTc uptake, confirming the specificity of the DUPA conjugate for PSMA in vivo. To further document this specificity, the radiotracer was also administered to two PSMA negative mouse xenograft models [A549 (a human lung cancer cell line) and KB (a human nasopharyngeal cancer cell line)], and again whole body images were taken. As anticipated, no radioactivity was observed in either KB or A549 tumors (FIGS. 7D(e) and 7D(f)), even after shielding of the kidneys was performed to allow detection of low levels of DUPA-99mTc in other tissues. These studies thus confirm that very little DUPA-99mTc binding occurs to sites unrelated to PSMA in vivo.

FIG. 7D shows the whole body images of solid tumor xenografts in nu/nu mice taken 4 h after injection of 150 μ Ci DUPA-99mTc. Overlay of whole-body radioimages on white light images of mice bearing LNCaP tumors that were treated with DUPA-99mTc in the absence 7D(a, c) or presence 7D(b, d) of 100-fold molar excess PMPA. Overlay of radioimages on white light images were also obtained of mice bearing an A549 tumor 7D(e) or a KB tumor 7D(f) that were similarly treated with DUPA-99mTc. Except in images 7D(a) and 7D(b), kidneys were shielded with lead pads. All images were taken using a Kodak Imaging Station 4 h after an intraperitoneal injection of DUPA-99mTc. Arrows indicate solid tumor xenografts. Similar images were obtained on all 5 mice in each treatment group.

Example 7A

Biodistribution Studies. After imaging, all animals were dissected approximately 6-7 h after administering SK28-99mTc or SK60-99mTc [or other radiolabeled compounds (data not shown)] and organs (blood, tumor, heart, liver, kidney, spleen, skin, muscle, etc) were transferred to individual gamma scintigraphy vials for each animal and radio-

98

activity was counted. Note: blood samples were collected (using cardio punch) immediately after sacrificing the animal and before imaging the animal. The plot of tumor to tissue cpm/g ratio verses tissue was used to determine bio-distribution of the imaging agent (FIGS. 8A and 8B).

Example 7B

Biodistribution studies of DUPA-99mTc in nu/nu mice bearing LNCaP, A549, or KB tumors. Tumor-bearing mice were euthanized 4 h after intraperitoneal injection of DUPA-99mTc (50 μ mol/kg, 150 μ Ci) and tissue-accumulated radioactivity was counted using a γ -counter. The percent injected dose per gram wet tissue was calculated as described in the Methods. The data were obtained in a single experiment and error bars represent s.d. (n=5). LNCaP tumors (solid bars), LNCaP tumors in mice pre-injected with 100-fold molar excess of PMPA (open bars), A549 tumors (cross-hatched bars), KB tumors (horizontally-hatched bars) (FIG. 8C).

Example 8

Single dose toxicity in live mice. Administration of SK71 was in a single dose as indicated. The data show that the MTD for the conjugate is about 4.5 μ mol/kg for single dosing. (See FIG. 9A)

Example 9

Multiple dose toxicity in live mice. SK71 was administered in 5 doses on alternate days (M, W, F, M, W). The data show that the MTD for SK71 is 2 μ mol/kg for multiple dosing, and that the conjugate is effective on LNCaP tumors (mice used for MTD 2 weeks after implantation of LNCaP cells before treatment was initiated). All 4 mice in the saline control group had large tumors, whereas no mice in the two treated groups had visible tumors after 18 days of treatment. (See FIG. 9B)

Example 10

Efficacy study compared to control group and competition group. Animals were treated with (a) the conjugate SK71 administered in 5 doses on alternate days (M, W, F, M, W) at 1 μ mol/kg, and compared to (b) vehicle treated animals (FIG. 10B), and to (c) animals treated with the conjugate in conjunction with PMPA. Treatment at 1 μ mol/kg shows a successive decrease in tumor size (starting tumor size approximately 250 mm³) during the course of treatment. At the lower dose of 1 μ mol/kg shown in FIG. 10A, tumor volumes rebounded at the cessation of dosing. At the higher dose of at least 2 μ mol/kg, complete disappearance of the tumor was observed during the testing period. The competition experiments (See FIG. 10C) indicate that the successful treatment of the implanted tumor is related to selective or specific targeting of PSMA mediated delivery.

Example 11

Efficacy Study (1 micromole/kg every other day for 10 days; i.e. 5 doses). The data (See FIG. 11) indicate that tumors in treated animals decreased in size during the duration of treatment.

Example 12

Evaluation of a PSMA-Targeted Therapeutic Agent in Vitro. Analysis of SK71 (FIG. 12A), SK77 (FIG. 12B),

99

SK37 (FIG. 12C), and SK45 (FIG. 12D), toxicity to LNCaP cells in culture. LNCaP cells were pulsed for 2 h with increasing concentrations of SK71 or SK77 in the presence (▲) or absence (■) of 100-fold molar excess PMPA. After 2× washes, cells were incubated an additional 66 h in fresh medium at 37° C. Cell viability was then analyzed using the [3H]-thymidine incorporation assay, as described herein. Data were obtained in a single experiment and error bars represent s.d. (n=3 wells per concentration).

Example 13

Potency In Vivo. Effect of SK71 on the growth of subcutaneous tumors (FIGS. 13A and 13C, and on the weights of the treated mice (FIGS. 13B and 13D). LNCaP cells in HC Matrigel were implanted subcutaneously into shoulders of nu/nu male mice. Once tumors reached either 100 mm³ (13A, 13B) or 330 mm³ (13C, 13D) in volume, animals were treated with SK71 [1.5 μmol/kg (a, b) or 2.0 μmol/kg (c, d)]. Treated mice (■), untreated mice (●), treated mice pre-injected with 100-fold (13A, 13B) or 30-fold (13C, 13D) molar excess of PMPA (▲ and ▼, respectively). Data were obtained in a single experiment and error bars represent s.d. [n=4 (13A, 13B) or 3 (13C, 13D)]. FIG. 10. SK71 potency in vivo.

Potency In Vivo. Effect of SK77 on the growth of subcutaneous tumors (FIG. 14A), and on the weights of the treated mice (FIG. 14B). LNCaP cells in HC Matrigel were implanted subcutaneously into shoulders of nu/nu male mice. Once tumors reached 100 mm³ in volume, animals were treated with SK77 (2 μmol/kg). Untreated mice (■), treated mice (▼). Data were obtained in a single experiment and error bars represent s.d. (n=4 mice/group).


What is claimed is:

1. A compound of the formula



or a salt thereof, wherein

B is a prostate specific membrane antigen binding or targeting ligand of the formula




where R1 is hydrogen and R2 is a substituted carboxylic acid;

L comprises an unsubstituted divalent carbonylalkylcarboxyl diradical and a tripeptide comprising one or more

100

optionally substituted Phe, wherein L does not comprise a cycloalkylene group; and D is a chelating group radical of the formula



where * is the point of attachment to L.

2. The compound of claim 1 or a salt thereof, wherein D further comprises a radioactive isotope of a metal coordinated to the chelating group.

3. The compound of claim 1 or a salt thereof, wherein L further comprises an amino acid or an amino acid derivative.

4. The compound of claim 1 or a salt thereof, wherein L further comprises an amino acid selected from the group consisting of Glu, Asp, Phe, Cys, beta-amino Ala, and aminoalkylcarboxylic acids.

5. The compound of claim 1 or a salt thereof, wherein L further comprises an aminoalkylcarboxylate, where the amino or the carboxylate are each optionally protected, and the alkyl is substituted with alkyl, hydroxy alkyl, sulfhydrylalkyl, aminoalkyl, or carboxy alkyl.

6. The compound of claim 1 or a salt thereof, wherein L further comprises an amino acid selected from the group consisting of Ser, Thr, Cys, Arg, Om, Lys, Asp, Glu, and Gln.

7. The compound of claim 1 or a salt thereof, wherein L further comprises an amino acid selected from the group consisting of Val, Leu, Ile, Phe, Tyr, and Met.

8. The compound of claim 1 or a salt thereof, wherein L further comprises phenylalanyl-phenylalanyl.

9. The compound of claim 1 or a salt thereof, wherein L further comprises one or more aryl or arylalkyl groups attached to the backbone of the linker about 7 to about 11 atoms from the binding ligand B.

10. The compound of claim 1 or a salt thereof, wherein L comprises a chain of at least about 14 atoms.

11. The compound of claim 1 or a salt thereof, wherein L comprises a chain of between about 14 and about 24 atoms.

12. The compound of claim 1 or a salt thereof, wherein L does not include a releasable linker.

13. A pharmaceutical composition comprising the conjugate of claim 1, and optionally one or more carriers, diluents, or excipients, or a combination thereof.

* * * * *