





(86) Date de dépôt PCT/PCT Filing Date: 2001/02/16
(87) Date publication PCT/PCT Publication Date: 2001/08/23
(45) Date de délivrance/Issue Date: 2010/05/11
(85) Entrée phase nationale/National Entry: 2002/08/16
(86) N° demande PCT/PCT Application No.: GB 2001/000655
(87) N° publication PCT/PCT Publication No.: 2001/060810
(30) Priorités/Priorities: 2000/02/16 (GB0003629.3);
2000/09/05 (GB0021774.5)

(51) Cl.Int./Int.Cl. *C07D 307/22* (2006.01),
A61K 31/341 (2006.01), *A61P 35/00* (2006.01)
(72) Inventeurs/Inventors:
HIGA, TATSUO, JP;
TANAKA, JUNICHI, JP;
GARCIA GRAVALOS, DOLORES, ES
(73) Propriétaire/Owner:
PHARMA MAR, S.A., ES
(74) Agent: MARKS & CLERK

(54) Titre : DERIVES DE TETRAHYDROFURYL OXY- ET AMINO-SUBSTITUES AYANT UNE ACTIVITE
ANTITUMORALE
(54) Title: OXY-AND AMINO-SUBSTITUTED TETRAHYDROFURYL DERIVATIVES WITH ANTITUMOUR ACTIVITY



(57) **Abrégé/Abstract:**

Compounds of formula (I) wherein the substituent groups defined by R_1 , R_2 , R_3 and R_4 are each independently selected of H, $C(=O)R$, $P(=O)R'R''$, $S(=O)R'R''$, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl; wherein each of the R' , R'' groups is independently selected from the group consisting of H, OH, NO_2 , NH_2 , SH, CN, halogen, $=O$, $C(=O)H$, $C(=O)CH_3$, CO_2H , CO_2CH_3 , substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl; wherein R_1 , R_2 , R_3 could form part of a heterocyclic ring; R_2 can be independently an internal salt; wherein X_1 , X_2 and X_3 on formula (I) are independently placed in any particular position of the chain and independently selected of H, OH, OR' , SH, SR' , SOR' , SO_2R' , NO_2 , NH_2 , NHR' , $N(R')_2$, $NHC(O)R'$, CN, halogen, $=O$, $C(=O)H$, $C(=O)CH_3$, CO_2H , CO_2CH_3 , substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic; wherein substituent groups defined by R' are each independently

(57) Abrégé(suite)/Abstract(continued):

selected from the group consisting of H, OH, OR', SH, SR', SOR', SO₂R', NO₂, NH₂, NHR', N(R')₂, NHC(O)R', CN, halogen, =O, C(=O)H, C(=O)CH₃, CO₂H, CO₂CH₃, substituted or unsubstituted C₁-C₁₈ alkyl, substituted or unsubstituted C₂-C₁₈ alkenyl, substituted or unsubstituted C₂-C₁₈ alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic; wherein x can be between 6 and 20; and wherein the dotted line is one or several double bonds placed in any particular position of the side chain; are useful as antitumor compounds and include compound IK-8-73-4 of structure (a) which was isolated from a sponge *Pachastrissa* sp.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number
WO 01/60810 A1

(51) International Patent Classification⁷: **C07D 307/22**,
A61K 31/341, A61P 35/00

[JP/JP]; 1-62-3, Shuri-Ishimine, Naha, Okinawa 903-0804 (JP). **TANAKA, Junichi** [JP/JP]; 5-15-20, Hantagawa, Naha, Okinawa 902-0071 (JP). **GARCIA GRAVALOS, Dolores** [ES/ES]; Maldonado, 63, E-28006 Madrid (ES).

(21) International Application Number: PCT/GB01/00655

(74) Agent: **RUFFLES, Graham, Keith**; 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).

(22) International Filing Date: 16 February 2001 (16.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0003629.3 16 February 2000 (16.02.2000) GB
0021774.5 5 September 2000 (05.09.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*):
PHARMA MAR, S.A. [ES/ES]; Calle de la Calera, 3, Poligono Industrial de Tres Cantos, Tres Cantos, E-28760 Madrid (ES).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for SD only*): **RUFFLES, Graham, Keith** [GB/GB]; 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).

(72) Inventors; and


Published:

(75) Inventors/Applicants (*for US only*): **HIGA, Tatsuo**

— with international search report

[Continued on next page]

(54) Title: OXY-AND AMINO-SUBSTITUTED TETRAHYDROFURYL DERIVATIVES WITH ANTITUMOUR ACTIVITY



(57) Abstract: Compounds of formula (I) wherein the substituent groups defined by R_1 , R_2 , R_3 and R_4 are each independently selected of H, C(=O)R, P(=O)R'R'', S(=O)R'R'', substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl; wherein each of the R', R'' groups is independently selected from the group consisting of H, OH, NO₂, NH₂, SH, CN, halogen, =O, C(=O)H, C(=O)CH₃, CO₂H, CO₂CH₃, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl; wherein R_1 , R_2 , R_3 could form part of a heterocyclic ring; R_2 can be independently an internal salt; wherein X_1 , X_2 and X_3 on formula (I) are independently placed in any particular position of the chain and independently selected of H, OH, OR', SH, SR', SOR', SO₂R', NO₂, NH₂, NHR', N(R')₂, NHC(O)R', CN, halogen, =O, C(=O)H, C(=O)CH₃, CO₂H, CO₂CH₃, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic; wherein substituent groups defined by R' are each independently selected from the group consisting of H, OH, OR', SH, SR', SOR', SO₂R', NO₂, NH₂, NHR', N(R')₂, NHC(O)R', CN, halogen, =O, C(=O)H, C(=O)CH₃, CO₂H, CO₂CH₃, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic; wherein x can be between 6 and 20; and wherein the dotted line is one or several double bonds placed in any particular position of the side chain; are useful as antitumor compounds and include compound IK-8-73-4 of structure (a) which was isolated from a sponge *Pachastrissa* sp.

WO 01/60810 A1

WO 01/60810 A1



— *with amended claims*

Date of publication of the amended claims: 29 November 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/60810

PCT/GB01/00655

OXY-AND AMINO-SUBSTITUTED TETRAHYDROFURYL DERIVATIVES
WITH ANTITUMOUR ACTIVITY


The present invention relates to antitumour compounds.

BACKGROUND OF THE INVENTION

Marine organisms continue to be an interesting source of biologically active molecules. Compounds such as ecteinascidin 743 are in clinical trials. Some important active compounds have been isolated from sponges, notably palau'amine. Other compounds of marine origin include spisulosines such as spisulosine 285 which is of interest for its antitumor activity, and reference is made to WO 9952521.

SUMMARY OF THE INVENTION

The present invention is directed to compounds which are analogs of the spisulosine family, having the following formula I:



I

wherein the substituent groups defined by R_1 , R_2 , R_3 and R_4 are each independently selected of H, $C(=O)R'$, $P(=O)R'R''$, $S(=O)R'R''$, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl. Wherein each of the R' , R'' groups is independently selected from the group consisting of H, OH, NO_2 , NH_2 , SH, CN, halogen, =O, $C(=O)H$, $C(=O)CH_3$, CO_2H , CO_2CH_3 , substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl. Wherein R_1 , R_2 , R_3 could form part of a heterocyclic ring (saturated: e.g. morpholine, oxazolidine or containing unsaturated double bonds: e.g. morpholinone, oxazolidinone, typically having 5 or 6 ring atoms). R_2 can be independently an internal salt. Wherein X_1 , X_2 and X_3 on formula I are independently placed in any particular position of the chain including closer to the ring on either carbon with the dotted line and independently selected of H, OH, OR' , SH, SR' , SOR' , SO_2R' , NO_2 , NH_2 , NHR' , $N(R')_2$, $NHC(O)R'$, CN, halogen, =O, $C(=O)H$, $C(=O)CH_3$, CO_2H , CO_2CH_3 , substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic. Wherein substituent groups defined by R' are each independently selected from the group consisting of H, OH, OR' , SH, SR' , SOR' , SO_2R' , NO_2 , NH_2 , NHR' , $N(R')_2$, $NHC(O)R'$, CN, halogen, =O, $C(=O)H$, $C(=O)CH_3$, CO_2H , CO_2CH_3 , substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic.

Wherein x can be between 6 and 20.


Wherein the dotted line is one or several double bonds placed in any particular position of the side chain.

Wherein the stereochemistry of the groups OR_1 and NR_2R_3 can be independently *syn* or *anti*.

Antitumoral activities of these compounds include leukaemias, lung cancer, colon cancer, kidney cancer, prostate cancer, ovarian cancer, breast cancer, sarcomas and melanomas.

The compounds are of thus of use in medicine, in particular in the treatment of tumours. The invention also relates to pharmaceutical preparations comprising such compounds for treatment of tumours, for example, solid tumours, and use of the compounds in the preparation of a medicament for the treatment of tumours.

In particular, we provide the compound we designate IK-8-73-4. The compound IK-8-73-4 is of formula $C_{18}H_{37}NO_2$ and has the structure:



The compounds of this invention can form salts, for example R^2 can be independently an internal salt. Preferred internal salts are formed using any kind of mineral or organic acid such as hydrochloric acid, hydrobromic acid, tartaric acid, succinic acid, etc. The method, use and compositions of this invention extend to such pharmaceutically acceptable salts.

PREFERRED EMBODIMENTS

In some preferred compounds, the group R_4 is hydrogen. One or both of R_2 and R_3 is suitably hydrogen. Typically R_1 is hydrogen. Preferably one, two or three of X_1 , X_2 and X_3 are hydrogen. Suitably the dotted line is absent to leave a single bond. Preferably x is 10 to 18, more preferably 10 to 14, such as about 12.

Suitable halogen substituents in the compounds of the present invention include F, Cl, Br and I.

Except where indicated, alkyl groups preferably have from 1 to about 12 carbon atoms, more preferably 1 to about 8 carbon atoms, still more preferably 1 to about 6 carbon atoms, and most preferably 1, 2, 3 or 4 carbon atoms. Methyl, ethyl and propyl including isopropyl are particularly preferred alkyl groups in the compounds of the present invention. As used herein, the term alkyl, unless otherwise modified, refers to both cyclic and noncyclic groups, although cyclic groups will comprise at least three carbon ring members.

Preferred alkenyl and alkynyl groups in the compounds of the present invention have one or more unsaturated linkages and from 2 to about 12 carbon atoms, more preferably 2 to about 8 carbon atoms, still more preferably 2 to about 6 carbon atoms, even more preferably 1, 2, 3 or 4 carbon atoms. The terms alkenyl and alkynyl as used herein refer to both cyclic and noncyclic groups, although straight or branched noncyclic groups are generally more preferred.

Preferred alkoxy groups in the compounds of the present invention include groups having one or more oxygen linkages and from 1 to about 12 carbon atoms, more preferably from 1 to about 8 carbon atoms, and still more preferably 1 to about 6 carbon atoms, and most preferably 1, 2, 3 or 4 carbon atoms.

Preferred alkylthio groups in the compounds of the present invention have one or more thioether linkages and from 1 to about 12 carbon atoms, more preferably from 1 to about 8 carbon atoms, and still more preferably 1 to about 6 carbon atoms. Alkylthio groups having 1, 2, 3 or 4 carbon atoms are particularly preferred.

Preferred alkylsulfinyl groups in the compounds of the present invention include those groups having one or more sulfoxide (SO) groups and from 1 to about 12 carbon atoms, more preferably from 1 to about 8 carbon atoms, and still more preferably 1 to about 6 carbon atoms. Alkylsulfinyl groups having 1, 2 3 or 4 carbon atoms are particularly preferred.

Preferred alkylsulfonyl groups in the compounds of the present invention include those groups having one or more sulfonyl (SO₂) groups and from 1 to about 12 carbon atoms, more preferably from 1 to about 8 carbon atoms, and still more preferably 1 to about 6 carbon atoms. Alkylsulfonyl groups having 1, 2, 3 or 4 carbon atoms are particularly preferred.

Preferred aminoalkyl groups include those groups having one or more primary, secondary and/or tertiary amine groups, and from 1 to about 12 carbon atoms, more preferably 1 to about 8 carbon atoms, still more preferably 1 to about 6 carbon atoms, even more preferably 1, 2, 3 or 4 carbon atoms. Secondary and tertiary amine groups are generally more preferred than primary amine moieties.

Suitable heteroaromatic groups in the compounds of the present invention contain one, two or three heteroatoms selected from N, O or S atoms and include, e.g., coumarinyl including 8-coumarinyl, quinolinyl including 8-quinolinyl, pyridyl, pyrazinyl, pyrimidyl, furyl, pyrrolyl, thienyl, thiazolyl, oxazolyl, imidazolyl, indolyl, benzofuranyl and benzothiazolyl. Suitable heteroalicyclic groups in the compounds of the present invention contain one, two or three heteroatoms selected from N, O or S atoms and include, e.g., tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, morpholino and pyrrolindinyl groups.

Suitable carbocyclic aryl groups in the compounds of the present invention include single and multiple ring compounds, including multiple ring compounds that contain separate and / or fused aryl groups. Typical carbocyclic aryl groups contain 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms. Specifically preferred carbocyclic aryl groups include phenyl including substituted phenyl such as 2-substituted phenyl, 3-substituted phenyl, 2, 3-substituted phenyl, 2,5-substituted phenyl, 2,3,5-substituted and 2,4,5-substituted phenyl, including where one or more of the phenyl substituents is an electron-withdrawing group such as halogen, cyano, nitro, alkanoyl, sulfinyl, sulfonyl and the like; naphthyl including 1-naphthyl and 2-naphthyl; biphenyl; phenanthryl; and anthracyl.

References herein to substituted R' groups in the compounds of the present invention refer to the specified moiety that may be substituted at one or more available positions by one or more suitable groups, e.g., halogen such as fluoro, chloro, bromo and iodo; cyano; hydroxyl; nitro; azido; alkanoyl such as a fluoro, chloro, bromo and iodo; cyano; hydroxyl; nitro; azido; alkanoyl such as a C1-6 alkanoyl group such as acyl and the like; carboxamido; alkyl groups including those groups having 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms and more preferably 1-3 carbon atoms; alkenyl and alkynyl groups including groups having one or more unsaturated linkages and from 2 to about 12 carbon or from 2 to about 6 carbon atoms; alkoxy groups having those having one or more oxygen linkages and from 1 to about 12 carbon atoms or 1 to about 6 carbon atoms; aryloxy such as phenoxy; alkylthio groups including those moieties having one or more thioether linkages and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; alkylsulfinyl groups including those moieties having one or more sulfinyl linkages and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; alkylsulfonyl groups including those moieties having one or more sulfonyl linkages and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; aminoalkyl groups such as groups having one or more N atoms and from 1 to about 12 carbon atoms or from 1 to about 6 carbon atoms; carbocyclic aryl having 6 or more carbons, particularly phenyl (e.g., R being a substituted or unsubstituted biphenyl moiety); and aralkyl such as benzyl.

Similar considerations apply to the substituted groups permitted for X₁, X₂ and X₃, including substituted C₁-C₁₈ alkyl, substituted C₂-C₁₈ alkenyl, substituted C₂-C₁₈ alkynyl, substituted aryl, substituted heteroaromatic, as well as for these groups where mentioned for R'. Preferably these definitions are interpreted to avoid unmeaningful multiple substitutions, such as alkyl substituted by alkyl, which remains an alkyl group, as well as to avoid structures which are not credible for the purpose of this invention. Preferably there are at most 10 to 50, more preferably 12 to 40, 14 to 30 or most preferably 16 to 20 carbon atoms in the molecule.

The substituents X₁, X₂ and X₃ on formula I are independently placed in any particular position of the chain including closer to the ring on either carbon with the


dotted line and independently selected. Thus, for example, there may be a substituent adjacent the ring, another further from the ring, and another remote from the ring.

Preferred internal salts are formed using any kind of mineral or organic acid such as hydrochloric acid, hydrobromic acid, tartaric acid, succinic acid, etc.

In a preferred class of compounds of this invention, R_4 is an alkyl chain with a substituent $-NR^2R^3$ and a substituent $-YR^1$ or $=YR^1$ on the next carbon towards the furan ring, where Y is selected from the group consisting of N, O, P, S and is preferably O. Suitably the amino group is on the second carbon and the amino group on the third carbon, counting the terminal carbon as the first.

There can be unsaturation in the $-(CH_2)_n-$, preferably a single unsaturation.

Preferred compounds of this invention include as side chain the partial structure:




Where the dotted line is a single or double bond. Wherein X_1 , X_2 and X_3 on formula I and formula II are independently selected of H, OH, OR' , SH, SR' , SOR' , SO_2R' , NO_2 , NH_2 , NHR' , $N(R')_2$, $NHC(O)R'$, CN, halogen, =O, $C(=O)H$, $C(=O)CH_3$, CO_2H , CO_2CH_3 , substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, substituted or unsubstituted C_2 - C_{18} alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaromatic. Wherein x and y can be between 0

and 20. Preferably x is in the range 8 to 16, more preferably 10, 11, 12, 13 or 14. Preferably y is 0, 1 or 2. Currently the most preferred is that x is 11, 12 or 13, and y is 0 or 1.

The compound IK-8-73-4 can be isolated from the sponge *Pachastrissa sp.*, Desmospongiae, Astrophorida, Calthropellidae. Accordingly, the invention also relates to compound IK-8-73-4 obtainable from *Pachastrissa sp.*, and to a method of isolating IK-8-73-4, characterised in that the starting material for the isolation is *Pachastrissa sp.*

The compound isolated from the sponge exists in the following stereochemistry:



The biological activity of the compound is illustrated by the following IC₅₀ data, µg/ml:

P388	A459	HT29	MEL28
leukemia	human lung	human colon	melanoma
0.01	0.01	0.01	0.01

Another especially preferred embodiment of the present invention is pharmaceutical compositions useful as antitumor agents which contain as active ingredient a compound or compounds of the invention, as well as the processes for their preparation.

Examples of pharmaceutical compositions include any solid (tablets, pills, capsules, granules etc.) or liquid (solutions, suspensions or emulsions) with suitable composition or oral, topical or parenteral administration.

Administration of the compounds or compositions of the present invention may be any suitable method, such as intravenous infusion, oral preparation, intraperitoneal and intravenous preparation.

The compounds of this invention can also be made by synthesis. Illustratively, the compounds can be made having regard to the procedure in Scheme 6 of Liebigs Ann. Chem., 1990, 1069-1078. In Scheme 6, (2*R*, 3*S*, 4*S*)-4-benzoylamido-3-hydroxy-2-dodecyltetrahydrofuran, compound 43, is prepared from (4*S*, 1'*S*, 2'*R*)-4,5-dihydro-4-(1',2'-dihydroxytetradecy)-2-phenyl-1,3-oxazole, compound 35. The (2*R*, 3*S*, 4*S*)-4-benzoylamido-3-hydroxy-2-dodecyltetrahydrofuran is then converted through (2*R*, 3*S*, 4*S*)-4-benzylamido-3-hydroxy-2-dodecyltetrahydrofuran and (2*R*, 3*S*, 4*S*)-4-amino-3-hydroxy-2-dodecyltetrahydrofuran to give (2*R*, 3*S*, 4*S*)-4-acetamido-3-acetoxy-2-dodecyltetrahydrofuran, compound 45. The 4-benzylamido-3-hydroxy-2-dodecyltetrahydrofuran and (2*R*, 3*S*, 4*S*)-4-amino-3-hydroxy-2-dodecyltetrahydrofuran are intermediates and are not purified.

This procedure can be used to prepare further compounds of this invention, by modifying as necessary the starting material 35 or 43 in Scheme 6, and isolating the debenzylated compound obtained at the end of the second step in the transition from compound 43 to compound 45, followed by derivatisation as necessary. Moreover, other procedures are available for the synthesis of tetrahydrofurans, and they can be adjusted as appropriate to give compounds of this invention.

The synthetic compounds of this invention are preferably provided in substantially pure form, that is free from impurity including starting material and reagents. In particular, we provide the benzyl and debenzylated compound obtained at the end of the first and second steps in the transition from compound 43 to compound 45 of the procedure in Liebigs Ann. Chem., 1990, 1069-1078. The invention extends to compositions and use of the benzoyl and acetyl compounds, compounds 43 and 45.

Examples of pharmaceutical compositions of the invention include any solid (for example tablets, pills, capsules, granules) or liquid (solutions, suspensions or emulsions) with suitable composition or oral, topical or parenteral administration, and they may contain the pure compound or in combination with any carrier or other pharmacologically active compounds. These compositions may need to be sterile when administered parenterally.

Suitably, the compound may be conjugated to a carrier protein or another suitable agent for delivery into the animal or human body. Conjugation may occur directly between a carrier and the compound, or indirectly via a suitable linker.

Administration of the compound or compositions of the present invention may be by any suitable method, such as intravenous infusion, oral preparations, intraperitoneal and intravenous administration. We prefer that infusion times of up to 24 hours are used, more preferably 2-12 hours, with 2-6 hours most preferred. Short infusion times which allow treatment to be carried out without an overnight stay in hospital are especially desirable. However, infusion may be 12 to 24 hours or even longer if required. Infusion may be carried out at suitable intervals of say 2 to 4 weeks. Pharmaceutical compositions containing compound of the invention may be delivered by liposome or nanosphere encapsulation, in sustained release formulations or by other standard delivery means.

The correct dosage of the compound will vary according to the particular formulation, the mode of application, and the particular *situs*, host and cancer or tumour being treated. Other factors like age, body weight, sex, diet, time of administration, rate of excretion, condition of the host, drug combinations, reaction sensitivities and severity of the disease shall be taken into account. Administration can be carried out continuously or periodically within the maximum tolerated dose.

The compound of this invention may be used with other drugs to provide a combination therapy. The other drugs may form part of the same composition, or be provided as a separate composition for administration at the same time or a different

time. The identity of the other drug is not particularly limited, and suitable candidates include:

- a) drugs with antimitotic effects, especially those which target cytoskeletal elements, including microtubule modulators such as taxane drugs (such as taxol, paclitaxel, taxotere, docetaxel), podophylotoxins or vinca alkaloids (vincristine, vinblastine);
- b) antimetabolite drugs such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate);
- c) alkylating agents such as nitrogen mustards (such as cyclophosphamide or ifosfamide);
- d) drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharmorubicin or epirubicin;
- e) drugs which target such as etoposide;
- f) hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuprorelin, goserelin, cyprotrone or octreotide;
- g) drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin;
- h) alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplatin) or nitrosoureas;
- i) drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors;
- j) gene therapy and antisense agents;
- k) antibody therapeutics; and
- l) other bioactive compounds of marine origin, notably the ecteinascidins such as ecteinascidin 743 and the didemnins such as aplidine.

EXAMPLE

An unidentified sponge (OP-98-18) was collected at Iriomote island, Okinawa on May 1998. The sponge (0.10 kg, wet) was extracted with acetone (0.50 L, 3

times), and the acetone solution was concentrated. The resulting aqueous suspension was partitioned with EtOAc to give 470 mg of lipophilic material (IC₅₀ 0.25 µg/mL). Most of it (310 mg) was subjected to chromatographic separation first on silica gel (active fraction, 63.0 mg, eluted with EtOAc-MeOH, 1-1) then preparative TLC (silica, EtOAc-MeOH, 3-1), and finally cation exchange column (eluted with MeOH, and MeOH+0.05% TFA). We obtained totally 10.1 mg (3.2% from extract) of compound 1.

We also found the same compound from another sponge specimen. We initially judged that the sponges were different, however, extracts showed identical ¹H NMR spectra of the same species. The original sponge was identified as *Pachastrissa* sp. (Demospongiae, Astrophorida, Calthropellidae). The sponge is deposited both at our lab (voucher no. OP-98--18 and OP-98-21) and at Queensland Museum (QM G317007). The collection was made on May 9, 1998 at Sakiyama Bay, Iriomote Island, Okinawa.

A specimen of the sponge *Pachastrissa* sp. (OP-98-21) was collected. The sponge was brought back to lab and kept frozen until extraction. The sample (0.20 kg, wet) was extracted with acetone (0.5 L, 3 times), and the filtered acetone extract was concentrated. The resulting aqueous suspension was partitioned with EtOAc. The organic layer (1.40 g) showed IC₅₀ 0.1 µg/mL. The water layer was concentrated, and the resulting solid material was washed with MeOH. After filtration, MeOH soluble material was taken. This MeOH extract (1.62 g) showed IC₅₀ 0.02 µg/mL.

Most (1.4 g) of the MeOH extract was subject to separation, first on Sephadex™ LH20 (CH₂ Cl₂-MeOH, 1-1), then on HPLC (RP18, not eluted with MeOH-H₂O (19-1), but with EtOAc-MeOH (1-1)), and finally cation exchange column (MeOH, MeOH-0.1%TFA) to give 11.5 mg (0.8 % from extract) of compound 1. We tried to separate components with MeOH-H₂O first. However, after separation we noticed that we did not recover all the material, so we washed the column with less polar solvent,

EtOAc-MeOH. And the compound was recovered in this fraction (checked by NMR and also by cytotoxicity results).

Compound 1 (free form, IK-8-73-4):

white amorphous solid;

$[\alpha]_{\text{D}21+180}$ (c 0.1, EtOH);

IR (film) ν max 3342, 2920, 2850, 2355, 1556, 1469, 1215, 1039, 758, 667 cm^{-1} ;

^1H NMR (500 MHz, CD_3OD) δ 0.90 (3H, m, H-18), 1.29 (24H, brs, H-6 to 17), 1.62 (2H, m, H-5), 3.43 (1H, dd, $J = 7.9, 8.9$ Hz, H-1), 3.51 (1H, dt, $J = 4.3, 8.9$ Hz, H-2), 3.81 (1H, dt, $J = 2.8, 6.7$ Hz, H-4), 3.86 (1H, dd, $J = 2.8, 4.3$ Hz, H-3), 3.90 (1H, dd, $J = 7.6, 7.9$ Hz, H-1);

^{13}C NMR (125 MHz, CD_3OD) δ 84.3 d (C-4), 73.3 d (C-3), 72.1 t (C-1), 56.0 d (C-2), 33.1 t, 30.9 t (C-5), 30.8 t, 30.7 t, 30.5 t, 30.5 t, 27.1 t, and 23.7 t, (C-6 to C-17), 14.4q (C-18);

ESIMS m/z 300 ($[\text{M}+\text{H}]^+$).

Compound 1 (salt form, IK-8-61-4):

white amorphous solid:


^1H NMR (500 MHz, CD_3OD) δ 0.87 (3H, m, H-18), 1.29 (24H, brs, H-6 to 17), 1.65 (2H, m, H-5), 3.71 (1H, dt, $J = 3.5, 6.6$ Hz, H-4), 3.79 (1H, dd, $J = 5.0, 8.5$ Hz, H-1), 3.88 (1H, m, H-2), 3.91 (1H, dd, $J = 7.3, 8.5$ Hz, H-1), 4.25 (1H, dd, $J = 3.5, 5.2$ Hz, H-3);

^{13}C NMR (125 MHz, CD_3OD) δ 84.3 d (C-4), 70.9 d (C-3), 68.9 t (C-1), 54.3 d (C-2), 33.0 t, 30.8 t, 30.7 t, 30.7 t, 30.4 t, 29.7 t (C-5), 27.2 t, and 23.7 t (C-6 to C17), 14.4 q (C-18);


EIMS m/z 299 (M^+).

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A compound having the formula:



2. Use, in the manufacture of a medicament for treating a tumour, of a compound of formula:



or a pharmaceutically acceptable salt thereof;

wherein R_1 is H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, or substituted or unsubstituted C_2 - C_{18} alkynyl; or R_1 together with R_2 or R_3 forms a saturated or unsaturated heterocyclic ring;

R_2 and R_3 are each independently H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, or substituted or unsubstituted C_2 - C_{18} alkynyl; or one of R_2 or R_3 together with R_1 forms a saturated or unsaturated heterocyclic ring or R_2 and R_3 together form a saturated or unsaturated heterocyclic ring; and


R_4 is H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, or substituted or unsubstituted C_2 - C_{18} alkynyl;

wherein R' is H, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, or substituted or unsubstituted C_2 - C_{18} alkynyl; and R'' is H, =O, substituted or unsubstituted C_1 - C_{18} alkyl, substituted or unsubstituted C_2 - C_{18} alkenyl, or substituted or unsubstituted C_2 - C_{18} alkynyl;


wherein X_1 , X_2 and X_3 are H; wherein x is 6 to 20 and wherein the dotted line is one or several optional double bonds placed in any particular position of the side chain.

3. The use of claim 2, wherein one of R_2 or R_3 together with R_1 forms a saturated or unsaturated heterocyclic ring and the other is H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl.


4. The use of claim 2, wherein the compound is of formula:



5. The use of claim 2, wherein the compound is:




6. A method of preparing compound IK-8-73-4 of structure:



which comprises obtaining a supply of a sponge *Pachastrissa* sp. containing said compound IK-8-73-4, and extracting the sponge with acetone, thereby isolating IK-8-73-4 from said supply of sponge.

7. A pharmaceutical composition for treatment of a tumour, said composition comprising a compound of formula:



or a pharmaceutically acceptable salt thereof;

wherein R_1 is H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl; or R_1 together with R_2 or R_3 forms a saturated or unsaturated heterocyclic ring;

R_2 and R_3 are each independently H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl; or one of R_2 or R_3 together with R_1 forms a saturated or unsaturated heterocyclic ring or R_2 and R_3 together form a saturated or unsaturated heterocyclic ring; and


R_4 is H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl;

wherein R' is H, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl; and R'' is H, $=O$, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl;

wherein X_1 , X_2 and X_3 are H ; wherein x is 6 to 20 and wherein the dotted line is one or several optional double bonds placed in any particular position of the side chain.

8. The composition of claim 7, wherein one of R_2 or R_3 together with R_1 forms a saturated or unsaturated heterocyclic ring and the other is H, $C(=O)R'$, $S(=O)R'R''$, substituted or unsubstituted C_1-C_{18} alkyl, substituted or unsubstituted C_2-C_{18} alkenyl, or substituted or unsubstituted C_2-C_{18} alkynyl.

9. The composition of 7, wherein the compound is of formula:



10. The composition of claim 7, wherein the compound is:

