

(12) STANDARD PATENT APPLICATION (11) Application No. AU 2012209050 A1
(19) AUSTRALIAN PATENT OFFICE

(54) Title
Covalent diabodies and uses thereof

(51) International Patent Classification(s)
C07K 16/00 (2006.01) **C12P 21/04** (2006.01)
A61K 39/395 (2006.01) **C12P 21/08** (2006.01)
C07H 21/04 (2006.01)

(21) Application No: **2012209050** (22) Date of Filing: **2012.08.02**

(43) Publication Date: **2012.08.23**

(43) Publication Journal Date: **2012.08.23**

(62) Divisional of:
2006236439

(71) Applicant(s)
Macrogenics, Inc.

(72) Inventor(s)
Huang, Ling;Johnson, Leslie S.

(74) Agent / Attorney
Fisher Adams Kelly, Level 29 12 Creek Street, Brisbane, QLD, 4000

ABSTRACT

The present invention is directed to diabody molecules and uses thereof in the treatment of a variety of diseases and disorders, including immunological disorders, infectious disease, intoxication and cancers. The diabody molecules of the invention comprise two polypeptide chains that associate to form at least two epitope binding sites, which may recognize the same or different epitopes on the same or differing antigens. Additionally, the antigens may be from that same or different molecules. The individual polypeptide chains of the diabody molecule may be covalently bound through non-peptide bond covalent bonds, such as, but not limited to, disulfide bonding of cysteine residues located within each polypeptide chain. In particular embodiments, the diabody molecules of the present invention further comprise an Fc region, which allows antibody-like functionality to engineer into the molecule.

AUSTRALIA

Patents Act 1990

**ORIGINAL
COMPLETE SPECIFICATION
STANDARD PATENT**

Invention Title:

"COVALENT DIABODIES AND USES THEREOF "

The following statement is a full description of this invention, including the best method of performing it known to me/us:

COVALENT DIABODIES AND USES THEREOF

[0001] This application claims the benefit of U.S. Provisional Application No. 60/671,657, filed April 15, 2005, which is hereby incorporated by reference in its entirety.

1. FIELD OF THE INVENTION

[0002] The present invention is directed to diabody molecules and uses thereof in the treatment of a variety of diseases and disorders, including immunological disorders and cancers. The diabody molecules of the invention comprise at least two polypeptide chains that associate to form at least two epitope binding sites, which may recognize the same or different epitopes. Additionally, the epitopes may be from the same or different molecules or located on the same or different cells. The individual polypeptide chains of the diabody molecule may be covalently bound through non-peptide bond covalent bonds, such as, but not limited to, disulfide bonding of cysteine residues located within each polypeptide chain. In particular embodiments, the diabody molecules of the present invention further comprise an Fc region, which allows antibody-like functionality to be engineered into the molecule.

2. BACKGROUND OF THE INVENTION

[0003] The design of covalent diabodies is based on the single chain Fv construct (scFv) (Holliger et al., 1993, Proc. Natl. Acad. Sci. USA 90:6444-6448; herein incorporated by reference in its entirety). In an intact, unmodified IgG, the VL and VH domains are located on separate polypeptide chains, *i.e.*, the light chain and the heavy chain, respectively. Interaction of an antibody light chain and an antibody heavy chain and, in particular, interaction of VL and VH domains forms one of the epitope binding sites of the antibody. In contrast, the scFv construct comprises a VL and VH domain of an antibody contained in a single polypeptide chain wherein the domains are separated by a flexible linker of sufficient length to allow self-assembly of the two domains into a functional epitope binding site. Where self assembly of the is impossible due to a linker of insufficient length (less than about 12 amino acid residues), two of the scFv constructs interact with each other to form a bivalent molecule, the VL of one chain associating with the VH of the other (reviewed in Marvin et al., 2005, Acta Pharmacol. Sin. 26:649-658). Moreover, addition of a cysteine residue to the c-terminus of the construct has been show to allow disulfide bonding of the polypeptide chains, stabilizing the resulting dimer without interfering with the binding characteristics of the bivalent molecule (see *e.g.*, Olafsen et al., 2004, Prot. Engr. Des. Sel. 17:21-27). Further, where VL and VH domains of differing

specificity are selected, not only a bivalent, but also a bispecific molecule may be constructed.

[0004] Bivalent diabodies have wide ranging applications including therapy and immunodiagnosis. Bivalency allows for great flexibility in the design and engineering of diabody in various applications, providing enhanced avidity to multimeric antigens, the cross-linking of differing antigens, and directed targeting to specific cell types relying on the presence of both target antigens. Due to their increased valency, low dissociation rates and rapid clearance from the circulation (for diabodies of small size, at or below ~50 kDa), diabody molecules known in the art have also shown particular use in the field of tumor imaging (Fitzgerald et al., 1997, Protein Eng. 10:1221). Of particular importance is the cross linking of differing cells, for example the cross linking of cytotoxic T cells to tumor cells (Staerz et al., 1985, Nature 314:628-631, and Holliger et al., 1996, Protein Eng. 9:299-305). Diabody epitope binding domains may also be directed to a surface determinant of any immune effector cell such as CD3, CD16, CD32, or CD64, which are expressed on T lymphocytes, natural killer (NK) cells or other mononuclear cells. In many studies, diabody binding to effector cell determinants, *e.g.*, Fcγ receptors (FcγR), was also found to activate the effector cell (Holliger et al., 1996, Protein Eng. 9:299-305; Holliger et al., 1999, Cancer Res. 59:2909-2916). Normally, effector cell activation is triggered by the binding of an antigen bound antibody to an effector cell via Fc-FcγR interaction; thus, in this regard, diabody molecules of the invention may exhibit Ig-like functionality independent of whether they comprise an Fc domain (*e.g.*, as assayed in any effector function assay known in the art or exemplified herein (*e.g.*, ADCC assay)). By cross-linking tumor and effector cells, the diabody not only brings the effector cell within the proximity of the tumor cells but leads to effective tumor killing (*see e.g.*, Cao and Lam, 2003, Adv. Drug. Deliv. Rev. 55:171-197, hereby incorporated by reference herein in its entirety).

2.1 EFFECTOR CELL RECEPTORS AND THEIR ROLES IN THE IMMUNE SYSTEM

[0005] In traditional immune function the interaction of antibody-antigen complexes with cells of the immune system results in a wide array of responses, ranging from effector functions such as antibody-dependent cytotoxicity, mast cell degranulation, and phagocytosis to immunomodulatory signals such as regulating lymphocyte proliferation and antibody secretion. All these interactions are initiated through the binding of the Fc domain of antibodies or immune complexes to specialized cell surface receptors on hematopoietic cells. The diversity of cellular responses triggered by antibodies and immune complexes

results from the structural heterogeneity of Fc receptors. Fc receptors share structurally related antigen binding domains which presumably mediate intracellular signaling.

[0006] The Fc γ receptors, members of the immunoglobulin gene superfamily of proteins, are surface glycoproteins that can bind the Fc γ portion of immunoglobulin molecules. Each member of the family recognizes immunoglobulins of one or more isotypes through a recognition domain on the alpha chain of the Fc γ receptor. Fc γ receptors are defined by their specificity for immunoglobulin subtypes. Fc γ receptors for IgG are referred to as Fc γ R, for IgE as Fc ϵ R, and for IgA as Fc α R. Different accessory cells bear Fc γ receptors for antibodies of different isotype, and the isotype of the antibody determines which accessory cells will be engaged in a given response (reviewed by Ravetch J.V. *et al.* 1991, *Annu. Rev. Immunol.* 9: 457-92; Gerber J.S. *et al.* 2001 *Microbes and Infection*, 3: 131-139; Billadeau D.D. *et al.* 2002, *The Journal of Clinical Investigation*, 2(109): 161-168; Ravetch J.V. *et al.* 2000, *Science*, 290: 84-89; Ravetch J.V. *et al.*, 2001 *Annu. Rev. Immunol.* 19:275-90; Ravetch J.V. 1994, *Cell*, 78(4): 553-60). The different Fc γ receptors, the cells that express them, and their isotype specificity is summarized in Table 1 (adapted from *Immunobiology: The Immune System in Health and Disease*, 4th ed. 1999, Elsevier Science Ltd/Garland Publishing, New York).

[0007] *Fc γ Receptors*

[0008] Each member of this family is an integral membrane glycoprotein, possessing extracellular domains related to a C2-set of immunoglobulin-related domains, a single membrane spanning domain and an intracytoplasmic domain of variable length. There are three known Fc γ Rs, designated Fc γ RI(CD64), Fc γ RII(CD32), and Fc γ RIII(CD16). The three receptors are encoded by distinct genes; however, the extensive homology between the three family members suggest they arose from a common progenitor perhaps by gene duplication.

[0009] *Fc γ RII(CD32)*

[0010] Fc γ RII proteins are 40 kDa integral membrane glycoproteins which bind only the complexed IgG due to a low affinity for monomeric Ig (10^6 M⁻¹). This receptor is the most widely expressed Fc γ R, present on all hematopoietic cells, including monocytes, macrophages, B cells, NK cells, neutrophils, mast cells, and platelets. Fc γ RII has only two immunoglobulin-like regions in its immunoglobulin binding chain and hence a much lower affinity for IgG than Fc γ RI. There are three human Fc γ RII genes (Fc γ RII-A, Fc γ RII-B, Fc γ RII-C), all of which bind IgG in aggregates or immune complexes.

[0011] Distinct differences within the cytoplasmic domains of Fc γ RII-A and Fc γ RII-B create two functionally heterogeneous responses to receptor ligation. The

fundamental difference is that the A isoform initiates intracellular signaling leading to cell activation such as phagocytosis and respiratory burst, whereas the B isoform initiates inhibitory signals, *e.g.*, inhibiting B-cell activation.

[0012] *FcγRIII (CD16)*

- 5 [0013] Due to heterogeneity within this class, the size of FcγRIII ranges between 40 and 80 kDa in mouse and man. Two human genes encode two transcripts, FcγRIIIA, an integral membrane glycoprotein, and FcγRIIIB, a glycosylphosphatidyl-inositol (GPI)-linked version. One murine gene encodes an FcγRIII homologous to the membrane spanning human FcγRIIIA. The FcγRIII shares structural characteristics with each of the other two FcγRs. Like FcγRII, FcγRIII binds IgG with low affinity and contains the corresponding two extracellular Ig-like domains. FcγRIIIA is expressed in macrophages, mast cells and is the lone FcγR in NK cells. The GPI-linked FcγRIIIB is currently known to be expressed only in human neutrophils.

[0014] *Signaling through FcγRs*

[0015] Both activating and inhibitory signals are transduced through the FcγRs following ligation. These diametrically opposing functions result from structural differences among the different receptor isoforms. Two distinct domains within the cytoplasmic signaling domains of the receptor called immunoreceptor tyrosine based activation motifs (ITAMs) or immunoreceptor tyrosine based inhibitory motifs (ITIMs) account for the different responses. The recruitment of different cytoplasmic enzymes to these structures dictates the outcome of the FcγR-mediated cellular responses. ITAM-containing FcγR complexes include FcγRI, FcγRIIA, FcγRIIIA, whereas ITIM-containing complexes only include FcγRIIB.

[0016] Human neutrophils express the FcγRIIA gene. FcγRIIA clustering via immune complexes or specific antibody cross-linking serves to aggregate ITAMs along with receptor-associated kinases which facilitate ITAM phosphorylation. ITAM phosphorylation serves as a docking site for Syk kinase, activation of which results in activation of downstream substrates (*e.g.*, PI₃K). Cellular activation leads to release of proinflammatory mediators.

[0017] The FcγRIIB gene is expressed on B lymphocytes; its extracellular domain is 96% identical to FcγRIIA and binds IgG complexes in an indistinguishable manner. The presence of an ITIM in the cytoplasmic domain of FcγRIIB defines this inhibitory subclass of FcγR. Recently the molecular basis of this inhibition was established. When co-ligated along with an activating FcγR, the ITIM in FcγRIIB becomes phosphorylated and attracts the SH2 domain of the inositol polyphosphate 5'-phosphatase (SHIP), which hydrolyzes

phosphoinositol messengers released as a consequence of ITAM-containing FcγR-mediated tyrosine kinase activation, consequently preventing the influx of intracellular Ca^{++} . Thus crosslinking of FcγRIIB dampens the activating response to FcγR ligation and inhibits cellular responsiveness. B cell activation, B cell proliferation and antibody secretion is thus aborted.

TABLE 1. Receptors for the Fc Regions of Immunoglobulin Isotypes

Receptor	FcγRI (CD64)	FcγRII-A (CD32)	FcγRII-B2 (CD32)	FcγRII-B1 (CD32)	FcγRIII (CD16)	FcεRI	FcαRI (CD89)
Binding	IgG1 10^8 M^{-1}	IgG1 $2 \times 10^6 \text{ M}^{-1}$	IgG1 $2 \times 10^6 \text{ M}^{-1}$	IgG1 $2 \times 10^6 \text{ M}^{-1}$	IgG1 $5 \times 10^5 \text{ M}^{-1}$	IgE 10^{10} M^{-1}	IgA1, IgA2 10^7 M^{-1}
Cell Type	Macrophages Neutrophils Eosinophils Dendritic cells	Macrophages Neutrophils Eosinophils Dendritic cells Platelets Langerhan cells	Macrophages Neutrophils Eosinophils	B cells Mast cells	NK cells Eosinophil Macrophages Neutrophils Mast Cells	Mast cells Eosinophil Basophils	Macrophages Neutrophils Eosinophils
Effect of Ligation	Uptake Stimulation Activation of respiratory burst Induction of killing	Uptake Granule release	Uptake Inhibition of Stimulation	No uptake Inhibition of Stimulation	Induction of Killing	Secretion of granules	Uptake Induction of killing

3. SUMMARY OF THE INVENTION

[0018] The present invention relates to covalent diabodies and/or covalent diabody molecules and to their use in the treatment of a variety of diseases and disorders including cancer, autoimmune disorders, allergy disorders and infectious diseases caused by bacteria, fungi or viruses. Preferably, the diabody of the present invention can bind to two different epitopes on two different cells wherein the first epitope is expressed on a different cell type than the second epitope, such that the diabody can bring the two cells together.

[0019] In one embodiment, the present invention is directed to a covalent bispecific diabody, which diabody comprises a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope, and, optionally, (iii) a third domain comprising at least one cysteine residue, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site; which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and, optionally, (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site (VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL2)(VH2) that binds the second epitope.

[0020] In another embodiment, the present invention is directed to a covalent bispecific diabody, which diabody comprises a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site; which

second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), which fourth and fifth domains are covalently linked such that the third and fourth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site (VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL2)(VH2) that binds the second epitope.

[0021] In certain aspects, the present invention is directed to diabody molecule, which molecule comprises a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site; which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site (VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL2)(VH2) that binds the second epitope.

[0022] In certain embodiments, the present invention is directed to a covalent bispecific diabody, which diabody is a dimer of diabody molecules, each diabody molecule comprising a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope and (iii) a third domain comprising an Fc domain or portion

thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site; and which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain of each diabody molecule are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain of each diabody molecule associate to form a first binding site (VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain of each diabody molecule associate to form a second binding site (VL2)(VH2) that binds the second epitope.

[0023] In yet other embodiments, the present invention is directed to a covalent tetrapeptide diabody, which diabody is a dimer of diabody molecules, the first diabody molecule comprising a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site; and which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site (VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL2)(VH2) that binds the second epitope; and the second diabody molecule comprising a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a third immunoglobulin (VL3) specific for a third epitope, (ii) a second domain comprising

a binding region of a heavy chain variable domain of a fourth immunoglobulin (VH4) specific for a fourth epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site; and which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the fourth immunoglobulin (VL4), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the third immunoglobulin (VH3), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site (VL3)(VH3) that binds the third epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL4)(VH4) that binds the fourth epitope.

[0024] In certain aspects of the invention the first epitope, second epitope, and where applicable, third epitope and fourth epitope can be the same. In other aspects, the first epitope, second epitope, and where applicable, third epitope and fourth epitope can each be different from the other. In certain aspects of the invention comprising a third epitope binding domain, the first epitope and third epitope can be the same. In certain aspects of the invention comprising a fourth epitope binding domain, the first epitope and fourth epitope can be the same. In certain aspects of the invention comprising a third epitope binding domain, the second epitope and third epitope can be the same. In certain aspects of the invention comprising a fourth epitope binding domain, the second epitope and fourth epitope can be the same. In preferred aspects of the invention, the first epitope and second epitope are different. In yet other aspects of the invention comprising a third epitope binding domain and a fourth epitope binding domain, the third epitope and fourth epitope can be different. It is to be understood that any combination of the foregoing is encompassed in the present invention.

[0025] In particular aspects of the invention, the first domain and the fifth domain of the diabody or diabody molecule can be derived from the same immunoglobulin. In another aspect, the second domain and the fourth domain of the diabody or diabody molecule can be derived from the same immunoglobulin. In yet another aspect, the first domain and the fifth domain of the diabody or diabody molecule can be derived from a different immunoglobulin. In yet another aspect, the second domain and the fourth domain of the

diabody or diabody molecule can be derived from a different immunoglobulin. It is to be understood that any combination of the foregoing is encompassed in the present invention.

[0026] In certain aspects of the invention, the covalent linkage between the first polypeptide chain and second polypeptide chain of the diabody or diabody molecule can be via a disulfide bond between at least one cysteine residue on the first polypeptide chain and at least one cysteine residue on the second polypeptide chain. The cysteine residues on the first or second polypeptide chains that are responsible for disulfide bonding can be found anywhere on the polypeptide chain including within the first, second, third, fourth, fifth and sixth domains. In a specific embodiment the cysteine residue on the first polypeptide chain is found in the first domain and the cysteine residue on the second polypeptide chain is found in the fifth domain. The first, second, fourth and fifth domains correspond to the variable regions responsible for binding. In preferred embodiments, the cysteine residues responsible for the disulfide bonding between the first and second polypeptide chains are located within the third and sixth domains, respectively. In a particular aspect of this embodiment, the third domain of the first polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23), which can be encoded by the amino acid sequence (SEQ ID NO:17). In another aspect of this embodiment, the sixth domain of the second polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23), which can be encoded by the amino acid sequence (SEQ ID NO:17). In still another aspect of this embodiment, the third domain of the first polypeptide chain comprises the amino acid sequence VEPKSC (SEQ ID NO:79), derived from the hinge domain of a human IgG, and which can be encoded by the nucleotide sequence (SEQ ID NO:80). In another aspect of this embodiment, the sixth domain of the second polypeptide chain comprises the amino acid sequence VEPKSC (SEQ ID NO:79), derived from the hinge domain of a human IgG, and which can be encoded by the nucleotide sequence (SEQ ID NO:80). In certain aspects of this embodiment, the third domain of the first polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23); and the sixth domain of the second polypeptide chain comprises the amino acid sequence VEPKSC (SEQ ID NO:79). In other aspects of this embodiment, the sixth domain of the second polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23); and the third domain of the first polypeptide chain comprises the amino acid sequence VEPKSC (SEQ ID NO:79). In yet other aspects of this embodiment, the third domain of the first polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23); and the sixth

domain of the second polypeptide chain comprises a hinge domain. In other aspects of this embodiment, the sixth domain of the second polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23); and the third domain of the first polypeptide chain comprises the hinge domain. In yet other aspects of this embodiment, the third domain of the first polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23); and the sixth domain of the first polypeptide chain comprises an Fc domain, or portion thereof. In still other aspects of this embodiment, the sixth domain of the second polypeptide chain comprises the C-terminal 6 amino acids of the human kappa light chain, FNRGEC (SEQ ID NO:23); and the third domain of the first polypeptide chain comprises an Fc domain, or portion thereof.

[0027] In other embodiments, the cysteine residues on the first or second polypeptide that are responsible for the disulfide bonding can be located outside of the first, second or third domains on the first polypeptide chain and outside of the fourth, fifth and sixth domain on the second polypeptide chain. In particular, the cysteine residue on the first polypeptide chain can be N-terminal to the first domain or can be C-terminal to the first domain. The cysteine residue on the first polypeptide chain can be N-terminal to the second domain or can be C-terminal to the second domain. The cysteine residue on the first polypeptide chain can be N-terminal to the third domain or can be C-terminal to the third domain. Further, the cysteine residue on the second polypeptide chain can be N-terminal to the fourth domain or can be C-terminal to the fourth domain. The cysteine residue on the second polypeptide chain can be N-terminal to the fifth domain or can be C-terminal to the fifth domain. Accordingly, the cysteine residue on the second polypeptide chain can be C-terminal to the sixth domain or can be N-terminal to the sixth domain. In a particular aspect, disulfide bond can be between at least two cysteine residues on the first polypeptide chain and at least two cysteine residues on the second polypeptide chain. In a particular aspect, wherein the third domain and sixth domain do not comprise an Fc domain, or portion thereof, the cysteine residue can be at the C-terminus of the first polypeptide chain and at the C-terminus of the second polypeptide chain. It is to be understood that any combination of the foregoing is encompassed in the present invention.

[0028] In specific embodiments of the invention described *supra*, the covalent diabody of the invention encompasses dimers of diabody molecules, wherein each diabody molecule comprises a first and second polypeptide chain. In certain aspects of this embodiment the diabody molecules can be covalently linked to form the dimer, with the proviso that the covalent linkage is not a peptide bond. In preferred aspects of this

embodiment, the covalent linkage is a disulfide bond between at least one cysteine residue on the first polypeptide chain of each of the diabody molecules of the dimer. In yet more preferred aspects of this invention, the covalent linkage is a disulfide bond between at least one cysteine residue on the first polypeptide chain of each of the diabody molecules forming the dimer, wherein said at least one cysteine residue is located in the third domain of each first polypeptide chain.

[0029] In certain aspects of the invention, the first domain on the first polypeptide chain can be N-terminal to the second domain or can be C-terminal to the second domain. The first domain on the first polypeptide chain can be N-terminal to the third domain or can be C-terminal to the third domain. The second domain on the first polypeptide chain can be N-terminal to the first domain or can be C-terminal to the first domain. Further, the second domain on the first polypeptide chain can be N-terminal to the third domain or can be C-terminal to the third domain. Accordingly, the third domain on the first polypeptide chain can be N-terminal to the first domain or can be C-terminal to the first domain. The third domain on the first polypeptide chain can be N-terminal to the second domain or can be C-terminal to the second domain. With respect to the second polypeptide chain, the fourth domain can be N-terminal to the fifth domain or can be C-terminal to the fifth domain. The fourth domain can be N-terminal to the sixth domain or can be C-terminal to the sixth domain. The fifth domain on the second polypeptide chain can be N-terminal to the fourth domain or can be C-terminal to the fourth domain. The fifth domain on the second polypeptide chain can be N-terminal to the sixth domain or can be C-terminal to the sixth domain. Accordingly the sixth domain on the second polypeptide chain can be N-terminal to the fourth domain or can be C-terminal to the fourth domain. The sixth domain on the second polypeptide chain can be N-terminal to the fifth domain or can be C-terminal to the fifth domain. It is to be understood that any combination of the foregoing is encompassed in the present invention.

[0030] In certain embodiments, first domain and second domain can be located C-terminal to the third domain on the first polypeptide chain; or the first domain and second domain can be located N-terminal to the third domain on the first polypeptide chain. With respect to the second polypeptide chain, the fourth domain and fifth domain can be located C-terminal to the sixth domain, or the fourth domain and fifth domain can be located N-terminal to the sixth domain. In certain aspects of this embodiment, the present invention is directed to a covalent bispecific diabody, which diabody is a dimer of diabody molecules, each diabody molecule comprising a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain

variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site and wherein the third domain is located N-terminal to both the first domain and second domain; and which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain of each diabody molecule are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain of each diabody molecule associate to form a first binding site (VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain of each diabody molecule associate to form a second binding site (VL2)(VH2) that binds the second epitope.

[0031] In yet another embodiment, the present invention is directed to a covalent tetrapeptide diabody, which diabody is a dimer of diabody molecules, the first diabody molecule comprising a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site and wherein the third domain is located N-terminal to both the first domain and second domain; and which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site

(VL1)(VH1) that binds the first epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL2)(VH2) that binds the second epitope; and the second diabody molecule comprises a first and a second polypeptide chain, which first polypeptide chain comprises (i) a first domain comprising a binding region of a light chain variable domain of a third immunoglobulin (VL3) specific for a third epitope, (ii) a second domain comprising a binding region of a heavy chain variable domain of a fourth immunoglobulin (VH4) specific for a fourth epitope and (iii) a third domain comprising an Fc domain or portion thereof, which first and second domains are covalently linked such that the first and second domains do not associate to form an epitope binding site and wherein the third domain is located N-terminal to both the first domain and second domain; and which second polypeptide chain comprises (i) a fourth domain comprising a binding region of a light chain variable domain of the fourth immunoglobulin (VL4), (ii) a fifth domain comprising a binding region of a heavy chain variable domain of the third immunoglobulin (VH3), and (iii) a sixth domain comprising at least one cysteine residue, which fourth and fifth domains are covalently linked such that the fourth and fifth domains do not associate to form an epitope binding site; and wherein the first polypeptide chain and the second polypeptide chain are covalently linked, with the proviso that the covalent link is not a peptide bond; wherein the first domain and the fifth domain associate to form a first binding site (VL3)(VH3) that binds the third epitope; wherein the second domain and the fourth domain associate to form a second binding site (VL4)(VH4) that binds the fourth epitope.

[0032] As discussed above, the domains on the individual polypeptide chains are covalently linked. In specific aspects, the covalent link between the first and second domain, first and third domain, second and third domain, fourth and fifth domain, fourth and sixth domain, and/or fifth and sixth domain can be a peptide bond. In particular, the first and second domains, and the fourth and fifth domains can be separated by the third domain and sixth domain, respectively, or by additional amino acid residues, so long as the first and second, and fourth and fifth domains do not associate to form a binding site. The number of amino acid residues can be 0, 1, 2, 3, 4, 5, 6, 7, 8 or 9 amino acid residues. In one preferred aspect, the number of amino acid residues between the domains is 8.

[0033] In certain aspects of the invention, the domains of the first and second polypeptid chain comprising an Fc domain, *i.e.*, optionally, the third and sixth domains, respectively, can further comprise a hinge domain such that the domain comprises a hinge-Fc region. In alternative embodiments, the first polypeptide chain or the second polypeptide chain can comprise a hinge domain without also comprising an Fc domain. The heavy

chains, light chains, hinge regions, Fc domains, and/or hinge-Fc domains for use in the invention can be derived from any immunoglobulin type including IgA, IgD, IgE, IgG or IgM. In a preferred aspect, the immunoglobulin type is IgG, or any subtype thereof, *i.e.*, IgG₁, IgG₂, IgG₃ or IgG₄. In other aspects, the immunoglobulin from which the light and heavy chains are derived is humanized or chimerized.

[0034] Further, the first epitope and second epitopes, and, where applicable, third epitope and fourth epitope, to which the diabody or diabody molecule binds can be different epitopes from the same antigen or can be different epitopes from different antigens. The antigens can be any molecule to which an antibody can be generated. For example, proteins, nucleic acids, bacterial toxins, cell surface markers, autoimmune markers, viral proteins, drugs, etc. In particular aspects, at least one epitope binding site of the diabody is specific for an antigen on a particular cell, such as a B-cell, a T-cell, a phagocytic cell, a natural killer (NK) cell or a dendritic cell.

[0035] In certain aspects of the present embodiment, at least one epitope binding site of the diabody or diabody molecule is specific for a Fc receptor, which Fc receptor can be an activating Fc receptor or an inhibitory Fc receptor. In particular aspects, the Fc receptor is a Fc γ receptor, and the Fc γ receptor is a Fc γ RI, Fc γ RII or Fc γ RIII receptor. In more preferred aspects, the Fc γ RIII receptor is the Fc γ RIIIA (CD16A) receptor or the Fc γ RIIIB (CD16B) receptor, and, more preferably, the Fc γ RIII receptor is the Fc γ RIIIA (CD16A) receptor. In another preferred aspect, the Fc γ RII receptor is the Fc γ RIIA (CD32A) receptor or the Fc γ RIIB (CD32B) receptor, and more preferably the Fc γ RIIB (CD32B) receptor. In a particularly preferred aspect, one binding site of the diabody is specific for CD32B and the other binding site is specific for CD16A. In a specific embodiment of the invention, at least one epitope binding site of the diabody or diabody molecule is specific for an activating Fc receptor and at least one other site is specific for an inhibitory Fc receptor. In certain aspects of this embodiment the activating Fc receptor is CD32A and the inhibitory Fc receptor is CD32B. In other aspects of this embodiment the activating Fc receptor is BCR and the inhibitory Fc receptor is CD32B. In still other aspects of this embodiment, the activating Fc receptor is IgERI and the inhibitory Fc receptor is CD32B.

[0036] In cases where one epitope binding site is specific for CD16A, the VL and VH domains can be the same as or similar to the VL and VH domains of the mouse antibody 3G8, the sequence of which has been cloned and is set forth herein. In other cases where one epitope binding site is specific for CD32A, the VL and VH domains can be the same as or similar to the VL and VH domains of the mouse antibody IV.3. In yet other cases where one epitope binding site is specific for CD32B, the VL and VH domains can be

the same as or similar to the VL and VH domains of the mouse antibody 2B6, the sequence of which has been cloned and is set forth herein. It is to be understood that any of the VL or VH domains of the 3G8, 2B6 and IV.3 antibodies can be used in any combination. The present invention is also directed to a bispecific diabody or diabody molecule wherein the first epitope is specific for CD32B, and the second epitope is specific for CD16A.

[0037] In other aspects, an epitope binding site can be specific for a pathogenic antigen. As used herein, a pathogenic antigen is an antigen involved in a specific pathogenic disease, including cancer, infection and autoimmune disease. Thus, the pathogenic antigen can be a tumor antigen, a bacterial antigen, a viral antigen, or an autoimmune antigen. Exemplary pathogenic antigens include, but are not limited to lipopolysaccharide, viral antigens selected from the group consisting of viral antigens from human immunodeficiency virus, Adenovirus, Respiratory Syncytial Virus, West Nile Virus (e.g., E16 and/or E53 antigens) and hepatitis virus, nucleic acids (DNA and RNA) and collagen. Preferably, the pathogenic antigen is a neutralizing antigen. In a preferred aspect, where one epitope binding site is specific for CD16A or CD32A, the other epitope binding site is specific for a pathogenic antigen excluding autoimmune antigens. In yet another preferred aspect, where one epitope binding site is specific for CD32B, the other epitope binding site is specific for any pathogenic antigen. In specific embodiments, the diabody molecule of the invention binds two different antigen on the same cell, for example, one antigen binding site is specific for an activating Fc receptor while the other is specific for an inhibitory Fc receptor. In other embodiments, the diabody molecule binds two distinct viral neutralizing epitopes, for example, but not limited to, E16 and E53 of West Nile Virus.

[0038] In yet another embodiment of the present invention, the diabodies of the invention can be used to treat a variety of diseases and disorders. Accordingly, the present invention is directed to a method for treating a disease or disorder comprising administering to a patient in need thereof an effective amount of a covalent diabody or diabody molecule of the invention in which at least one binding site is specific for a pathogenic antigen, such as an antigen expressed on the surface of a cancer cell or on the surface of a bacterium or virion and at least one other binding site is specific for a Fc receptor, *e.g.*, CD16A.

[0039] In yet another embodiment, the invention is directed to a method for treating a disease or disorder comprising administering to a patient in need thereof an effective amount of a diabody or diabody molecule of the invention, in which at least one binding site is specific for CD32B and at least one other binding site is specific for CD16A.

[0040] In yet another embodiment, the invention is directed to a method for inducing immune tolerance to a pathogenic antigen comprising administering to a patient in

need there an effective amount of a covalent diabody or dovalent diabody molecule of the invention, in which at least one binding site is specific for CD32B and at least one other binding site is specific for said pathogenic antigen. In aspects of this embodiment, the pathogenic antigen can be an allergen or another molecule to which immune tolerance is desired, such as a protein expressed on transplanted tissue.

[0041] In yet another embodiment, the present invention is directed to a method for detoxification comprising administering to a patient in need thereof an effective amount of a covalent diabody or diabody molecule of the invention, in which at least one binding site is specific for a cell surface marker and at least one other binding site is specific for a toxin. In particular aspects, the diabody of the invention administered is one where one binding site is specific for a cell surface marker such as an Fc and the other binding site is specific for a bacterial toxin or for a drug. In one aspect, the cell surface marker is not found on red blood cells.

3.1 DEFINITIONS

[0042] Unless otherwise defined, all terms of art, notations and other scientific terms or terminology used herein are intended to have the meanings commonly understood by those of skill in the art to which this invention pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a substantial difference over what is generally understood in the art. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, nucleic acid chemistry, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, Current Protocols in Immunology (J. E. Coligan et al., eds., 1999, including supplements through 2001); Current Protocols in Molecular Biology (F. M. Ausubel et al., eds., 1987, including supplements through 2001); Molecular Cloning: A Laboratory Manual, third edition (Sambrook and Russel, 2001); PCR: The Polymerase Chain Reaction, (Mullis et al., eds., 1994); The Immunoassay Handbook (D. Wild, ed., Stockton Press NY, 1994); Bioconjugate Techniques (Greg T. Hermanson, ed., Academic Press, 1996); Methods of Immunological Analysis (R. Masseyeff, W. H. Albert, and N. A. Staines, eds., Weinheim: VCH Verlagsgesellschaft mbH, 1993), Harlow and Lane Using Antibodies: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1999; and Beaucage et al. eds., Current Protocols in Nucleic Acid Chemistry John Wiley & Sons, Inc., New York, 2000).

[0043] As used herein, the terms “antibody” and “antibodies” refer to monoclonal antibodies, multispecific antibodies, human antibodies, humanized antibodies, synthetic antibodies, chimeric antibodies, polyclonal antibodies, camelized antibodies, single-chain Fvs (scFv), single chain antibodies, Fab fragments, F(ab') fragments, disulfide-linked bispecific Fvs (sdFv), intrabodies, and anti-idiotypic (anti-Id) antibodies (including, *e.g.*, anti-Id and anti-anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. In particular, antibodies include immunoglobulin molecules and immunologically active fragments of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site. Immunoglobulin molecules can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY), class (*e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂) or subclass.

[0044] As used herein, the terms “immunospecifically binds,” “immunospecifically recognizes,” “specifically binds,” “specifically recognizes” and analogous terms refer to molecules that specifically bind to an antigen (*e.g.*, epitope or immune complex) and do not specifically bind to another molecule. A molecule that specifically binds to an antigen may bind to other peptides or polypeptides with lower affinity as determined by, *e.g.*, immunoassays, BIAcore, or other assays known in the art. Preferably, molecules that specifically bind an antigen do not cross-react with other proteins. Molecules that specifically bind an antigen can be identified, for example, by immunoassays, BIAcore, or other techniques known to those of skill in the art.

[0045] As used herein, immune complex, refers to a structure which forms when at least one target molecule and at least one heterologous Fcγ region-containing polypeptide bind to one another forming a larger molecular weight complex. Examples of immune complexes are antigen-antibody complexes which can be either soluble or particulate (*e.g.*, an antigen/antibody complex on a cell surface.).

[0046] As used herein, the terms “heavy chain,” “light chain,” “variable region,” “framework region,” “constant domain,” and the like, have their ordinary meaning in the immunology art and refer to domains in naturally occurring immunoglobulins and the corresponding domains of synthetic (*e.g.*, recombinant) binding proteins (*e.g.*, humanized antibodies, single chain antibodies, chimeric antibodies, etc.). The basic structural unit of naturally occurring immunoglobulins (*e.g.*, IgG) is a tetramer having two light chains and two heavy chains, usually expressed as a glycoprotein of about 150,000 Da. The amino-terminal (“N”) portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal (“C”) portion of each chain defines a constant region, with light chains having a single constant

domain and heavy chains usually having three constant domains and a hinge region. Thus, the structure of the light chains of an IgG molecule is $n-V_L-C_L-c$ and the structure of IgG heavy chains is $n-V_H-C_{H1}-H-C_{H2}-C_{H3}-c$ (where H is the hinge region). The variable regions of an IgG molecule consist of the complementarity determining regions (CDRs), which contain the residues in contact with antigen and non-CDR segments, referred to as framework segments, which in general maintain the structure and determine the positioning of the CDR loops (although certain framework residues may also contact antigen). Thus, the V_L and V_H domains have the structure $n-FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4-c$.

[0047] When referring to binding proteins or antibodies (as broadly defined herein), the assignment of amino acids to each domain is in accordance with the definitions of Kabat, Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md., 1987 and 1991). Amino acids from the variable regions of the mature heavy and light chains of immunoglobulins are designated by the position of an amino acid in the chain. Kabat described numerous amino acid sequences for antibodies, identified an amino acid consensus sequence for each subgroup, and assigned a residue number to each amino acid. Kabat's numbering scheme is extendible to antibodies not included in his compendium by aligning the antibody in question with one of the consensus sequences in Kabat by reference to conserved amino acids. This method for assigning residue numbers has become standard in the field and readily identifies amino acids at equivalent positions in different antibodies, including chimeric or humanized variants. For example, an amino acid at position 50 of a human antibody light chain occupies the equivalent position to an amino acid at position 50 of a mouse antibody light chain.

[0048] As used herein, the term "heavy chain" is used to define the heavy chain of an IgG antibody. In an intact, native IgG, the heavy chain comprises the immunoglobulin domains V_H , $CH1$, hinge, $CH2$ and $CH3$. Throughout the present specification, the numbering of the residues in an IgG heavy chain is that of the EU index as in Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, NH1, MD (1991), expressly incorporated herein by references. The "EU index as in Kabat" refers to the numbering of the human IgG1 EU antibody. Examples of the amino acid sequences containing human IgG1 hinge, $CH2$ and $CH3$ domains are shown in FIGS. 1A and 1B as described, *infra*. FIGS. 1A and 1B also set forth amino acid sequences of the hinge, $CH2$ and $CH3$ domains of the heavy chains of IgG2, IgG3 and IgG4. The amino acid sequences of IgG2, IgG3 and IgG4 isotypes are aligned with the IgG1 sequence by placing the first and last cysteine residues of the respective hinge regions, which form the inter-heavy chain

S-S bonds, in the same positions. For the IgG2 and IgG3 hinge region, not all residues are numbered by the Eu index.

[0049] The “hinge region” or “hinge domain” is generally defined as stretching from Glu216 to Pro230 of human IgG1. An example of the amino acid sequence of the human IgG1 hinge region is shown in FIG. 1A (amino acid residues in FIG. 1A are numbered according to the Kabat system). Hinge regions of other IgG isotypes may be aligned with the IgG1 sequence by placing the first and last cysteine residues forming inter-heavy chain S-S binds in the same positions as shown in FIG. 1A.

[0050] As used herein, the term “Fc region,” “Fc domain” or analogous terms are used to define a C-terminal region of an IgG heavy chain. An example of the amino acid sequence containing the human IgG1 is shown in FIG. 1B. Although boundaries may vary slightly, as numbered according to the Kabat system, the Fc domain extends from amino acid 231 to amino acid 447 (amino acid residues in FIG. 1B are numbered according to the Kabat system). FIG. 1B also provides examples of the amino acid sequences of the Fc regions of IgG isotypes IgG2, IgG3, and IgG4.

[0051] The Fc region of an IgG comprises two constant domains, CH2 and CH3. The CH2 domain of a human IgG Fc region usually extends from amino acids 231 to amino acid 341 according to the numbering system of Kabat (FIG. 1B). The CH3 domain of a human IgG Fc region usually extends from amino acids 342 to 447 according to the numbering system of Kabat (FIG. 1B). The CH2 domain of a human IgG Fc region (also referred to as “C γ 2” domain) is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are interposed between the two CH2 domains of an intact native IgG.

[0052] As used herein the terms “Fc γ R binding protein,” “Fc γ R antibody,” and “anti-Fc γ R antibody”, are used interchangeably and refer to a variety of immunoglobulin-like or immunoglobulin-derived proteins. “Fc γ R binding proteins” bind Fc γ R via an interaction with V_L and/or V_H domains (as distinct from Fc γ -mediated binding). Examples of Fc γ R binding proteins include fully human, polyclonal, chimeric and humanized antibodies (*e.g.*, comprising 2 heavy and 2 light chains), fragments thereof (*e.g.*, Fab, Fab', F(ab')₂, and Fv fragments), bifunctional or multifunctional antibodies (*see, e.g.*, Lanzavecchia *et al.*, 1987, *Eur. J. Immunol.* 17:105), single chain antibodies (*see, e.g.*, Bird *et al.*, 1988, *Science* 242:423-26), fusion proteins (*e.g.*, phage display fusion proteins), “minibodies” (*see, e.g.*, U.S. Patent No. 5,837,821) and other antigen binding proteins comprising a V_L and/or V_H domain or fragment thereof. In one aspect, the Fc γ RIIIA binding protein is a “tetrameric antibody” *i.e.*, having generally the structure of a naturally

occurring IgG and comprising variable and constant domains, *i.e.*, two light chains comprising a V_L domain and a light chain constant domain and two heavy chains comprising a V_H domain and a heavy chain hinge and constant domains.

[0053] As used herein the term “FcγR antagonists” and analogous terms refer to protein and non-proteinaceous substances, including small molecules which antagonize at least one biological activity of an FcγR, *e.g.*, block signaling. For example, the molecules of the invention block signaling by blocking the binding of IgGs to an FcγR.

[0054] As used herein, the term “derivative” in the context of polypeptides or proteins refers to a polypeptide or protein that comprises an amino acid sequence which has been altered by the introduction of amino acid residue substitutions, deletions or additions. The term “derivative” as used herein also refers to a polypeptide or protein which has been modified, *i.e.*, by the covalent attachment of any type of molecule to the polypeptide or protein. For example, but not by way of limitation, an antibody may be modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular antigen or other protein, *etc.* A derivative polypeptide or protein may be produced by chemical modifications using techniques known to those of skill in the art, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, *etc.* Further, a derivative polypeptide or protein derivative possesses a similar or identical function as the polypeptide or protein from which it was derived.

[0055] As used herein, the term “derivative” in the context of a non-proteinaceous derivative refers to a second organic or inorganic molecule that is formed based upon the structure of a first organic or inorganic molecule. A derivative of an organic molecule includes, but is not limited to, a molecule modified, *e.g.*, by the addition or deletion of a hydroxyl, methyl, ethyl, carboxyl or amine group. An organic molecule may also be esterified, alkylated and/or phosphorylated.

[0056] As used herein, the term “diabody molecule” refers to a complex of two or more polypeptide chains or proteins, each comprising at least one VL and one VH domain or fragment thereof, wherein both domains are comprised within a single polypeptide chain. In certain embodiments “diabody molecule” includes molecules comprising an Fc or a hinge-Fc domain. Said polypeptide chains in the complex may be the same or different, *i.e.*, the diabody molecule may be a homo-multimer or a hetero-multimer. In specific aspects, “diabody molecule” includes dimers or tetramers or said polypeptide chains containing both a VL and VH domain. The individual polypeptide chains comprising the multimeric

proteins may be covalently joined to at least one other peptide of the multimer by interchain disulfide bonds.

[0057] As used herein, the terms “disorder” and “disease” are used interchangeably to refer to a condition in a subject. In particular, the term “autoimmune disease” is used interchangeably with the term “autoimmune disorder” to refer to a condition in a subject characterized by cellular, tissue and/or organ injury caused by an immunologic reaction of the subject to its own cells, tissues and/or organs. The term “inflammatory disease” is used interchangeably with the term “inflammatory disorder” to refer to a condition in a subject characterized by inflammation, preferably chronic inflammation. Autoimmune disorders may or may not be associated with inflammation. Moreover, inflammation may or may not be caused by an autoimmune disorder. Thus, certain disorders may be characterized as both autoimmune and inflammatory disorders.

[0058] “Identical polypeptide chains” as used herein also refers to polypeptide chains having almost identical amino acid sequence, for example, including chains having one or more amino acid differences, preferably conservative amino acid substitutions, such that the activity of the two polypeptide chains is not significantly different

[0059] As used herein, the term “cancer” refers to a neoplasm or tumor resulting from abnormal uncontrolled growth of cells. As used herein, cancer explicitly includes, leukemias and lymphomas. In some embodiments, cancer refers to a benign tumor, which has remained localized. In other embodiments, cancer refers to a malignant tumor, which has invaded and destroyed neighboring body structures and spread to distant sites. In some embodiments, the cancer is associated with a specific cancer antigen.

[0060] As used herein, the term “immunomodulatory agent” and variations thereof refer to an agent that modulates a host’s immune system. In certain embodiments, an immunomodulatory agent is an immunosuppressant agent. In certain other embodiments, an immunomodulatory agent is an immunostimulatory agent. Immunomodulatory agents include, but are not limited to, small molecules, peptides, polypeptides, fusion proteins, antibodies, inorganic molecules, mimetic agents, and organic molecules.

[0061] As used herein, the term “epitope” refers to a fragment of a polypeptide or protein or a non-protein molecule having antigenic or immunogenic activity in an animal, preferably in a mammal, and most preferably in a human. An epitope having immunogenic activity is a fragment of a polypeptide or protein that elicits an antibody response in an animal. An epitope having antigenic activity is a fragment of a polypeptide or protein to which an antibody immunospecifically binds as determined by any method well-known to

one of skill in the art, for example by immunoassays. Antigenic epitopes need not necessarily be immunogenic.

[0062] As used herein, the term “fragment” refers to a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, at least 10 contiguous amino acid residues, at least 15 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino residues, at least 70 contiguous amino acid residues, at least contiguous 80 amino acid residues, at least contiguous 90 amino acid residues, at least contiguous 100 amino acid residues, at least contiguous 125 amino acid residues, at least 150 contiguous amino acid residues, at least contiguous 175 amino acid residues, at least contiguous 200 amino acid residues, or at least contiguous 250 amino acid residues of the amino acid sequence of another polypeptide. In a specific embodiment, a fragment of a polypeptide retains at least one function of the polypeptide.

[0063] As used herein, the terms “nucleic acids” and “nucleotide sequences” include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), combinations of DNA and RNA molecules or hybrid DNA/RNA molecules, and analogs of DNA or RNA molecules. Such analogs can be generated using, for example, nucleotide analogs, which include, but are not limited to, inosine or tritylated bases. Such analogs can also comprise DNA or RNA molecules comprising modified backbones that lend beneficial attributes to the molecules such as, for example, nuclease resistance or an increased ability to cross cellular membranes. The nucleic acids or nucleotide sequences can be single-stranded, double-stranded, may contain both single-stranded and double-stranded portions, and may contain triple-stranded portions, but preferably is double-stranded DNA.

[0064] As used herein, a “therapeutically effective amount” refers to that amount of the therapeutic agent sufficient to treat or manage a disease or disorder. A therapeutically effective amount may refer to the amount of therapeutic agent sufficient to delay or minimize the onset of disease, *e.g.*, delay or minimize the spread of cancer. A therapeutically effective amount may also refer to the amount of the therapeutic agent that provides a therapeutic benefit in the treatment or management of a disease. Further, a therapeutically effective amount with respect to a therapeutic agent of the invention means the amount of therapeutic agent alone, or in combination with other therapies, that provides a therapeutic benefit in the treatment or management of a disease.

[0065] As used herein, the terms “prophylactic agent” and “prophylactic agents” refer to any agent(s) which can be used in the prevention of a disorder, or prevention of

recurrence or spread of a disorder. A prophylactically effective amount may refer to the amount of prophylactic agent sufficient to prevent the recurrence or spread of hyperproliferative disease, particularly cancer, or the occurrence of such in a patient, including but not limited to those predisposed to hyperproliferative disease, for example those genetically predisposed to cancer or previously exposed to carcinogens. A prophylactically effective amount may also refer to the amount of the prophylactic agent that provides a prophylactic benefit in the prevention of disease. Further, a prophylactically effective amount with respect to a prophylactic agent of the invention means that amount of prophylactic agent alone, or in combination with other agents, that provides a prophylactic benefit in the prevention of disease.

[0066] As used herein, the terms “prevent”, “preventing” and “prevention” refer to the prevention of the recurrence or onset of one or more symptoms of a disorder in a subject as result of the administration of a prophylactic or therapeutic agent.

[0067] As used herein, the term “in combination” refers to the use of more than one prophylactic and/or therapeutic agents. The use of the term “in combination” does not restrict the order in which prophylactic and/or therapeutic agents are administered to a subject with a disorder. A first prophylactic or therapeutic agent can be administered prior to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second prophylactic or therapeutic agent to a subject with a disorder.

[0068] “Effector function” as used herein is meant a biochemical event that results from the interaction of an antibody Fc region with an Fc receptor or an antigen. Effector functions include but are not limited to antibody dependent cell mediated cytotoxicity (ADCC), antibody dependent cell mediated phagocytosis (ADCP), and complement dependent cytotoxicity (CDC). Effector functions include both those that operate after the binding of an antigen and those that operate independent of antigen binding.

[0069] “Effector cell” as used herein is meant a cell of the immune system that expresses one or more Fc receptors and mediates one or more effector functions. Effector cells include but are not limited to monocytes, macrophages, neutrophils, dendritic cells, eosinophils, mast cells, platelets, B cells, large granular lymphocytes, Langerhans' cells,

natural killer (NK) cells, and may be from any organism including but not limited to humans, mice, rats, rabbits, and monkeys.

[0070] As used herein, the term “specifically binds an immune complex” and analogous terms refer to molecules that specifically bind to an immune complex and do not specifically bind to another molecule. A molecule that specifically binds to an immune complex may bind to other peptides or polypeptides with lower affinity as determined by, *e.g.*, immunoassays, BIAcore, or other assays known in the art. Preferably, molecules that specifically bind an immune complex do not cross-react with other proteins. Molecules that specifically bind an immune complex can be identified, for example, by immunoassays, BIAcore, or other techniques known to those of skill in the art.

[0071] A “stable fusion protein” as used herein refers to a fusion protein that undergoes minimal to no detectable level of degradation during production and/or storage as assessed using common biochemical and functional assays known to one skilled in the art, and can be stored for an extended period of time with no loss in biological activity, *e.g.*, binding to FcγR.

4. BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1 A-B AMINO ACID SEQUENCE OF HUMAN IgG CH1, HINGE and Fc REGIONS

[0072] Figure 1 provides the amino acid sequences of human IgG1, IgG2, IgG3 and IgG4 hinge (A) and Fc (B) domains. (IgG1 hinge domain (SEQ ID NO:1); IgG2 hinge domain (SEQ ID NO:2); IgG3 hinge domain (SEQ ID NO:3); IgG4 hinge domain (SEQ ID NO:4); IgG1 Fc domain (SEQ ID NO:5); IgG2 Fc domain (SEQ ID NO:6); IgG3 Fc domain (SEQ ID NO:7); IgG1 Fc domain (SEQ ID NO:8)). The amino acid residues shown in FIGS. 1A and 1B are numbered according to the numbering system of Kabat EU. Isotype sequences are aligned with the IgG1 sequence by placing the first and last cysteine residues of the respective hinge regions, which form the inter-heavy chain S-S bonds, in the same positions. For figure 1B, residues in the CH2 domain are indicated by +, while residues in the CH3 domain are indicated by ~.

FIG. 2 SCHEMATIC REPRESENTATION OF POLYPEPTIDE CHAINS OF COVALENT BIFUNCTIONAL DIABODIES

[0073] Polypeptides of a covalent, bifunctional diabody consist of an antibody VL and an antibody VH domain separated by a short peptide linker. The 8 amino acid residue linker prevents self assembly of a single polypeptide chain into scFv constructs, and, instead, interactions between the VL and VH domains of differing polypeptide chains

predominate. 4 constructs were created (each construct is described from the amino terminus ("n"), left side of the construct, to the carboxy terminus ("c"), right side of figure): construct (1) (SEQ ID NO:9) comprised, n-the VL domain Hu2B6 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu3G8 - and a C-terminal sequence (LGGC)-c; construct (2) (SEQ ID NO:11) comprised n-the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 - and a C-terminal sequence (LGGC)-c; construct (3) (SEQ ID NO:12) comprised n-the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu3G8 - and a C-terminal sequence (LGGC)-c; construct (4) (SEQ ID NO:13) comprised n-the VL domain Hu2B6 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 - and a C-terminal sequence (LGGC)-c.

FIG. 3 SDS-PAGE ANALYSIS OF AFFINITY PURIFIED DIABODIES

[0074] Affinity purified diabodies were subjected to SDS-PAGE analysis under reducing (lanes 1-3) or non-reducing (lanes 4-6) conditions. Approximate molecular weights of the standard (in between lanes 3 and 4) are indicated. Lanes 1 and 4, h3G8 CMD; Lanes 2 and 5, h2B6 CMD; and Lanes 3 and 6, h2B6-h3G8 CBD.

FIGS. 4 A-B SEC ANALYSIS OF AFFINITY PURIFIED DIABODIES

[0075] Affinity purified diabodies were subjected to SEC analysis. (A) Elution profile of known standards: full-length IgG (~150 kDa), Fab fragment of IgG (~50 kDa), and scFv (~30 kDa); (B) Elution profile of h2b6 CMD, h3G8 CMD, and h2B6-h3G8 CBD.

FIG. 5 BINDING OF h2B6-h3G8 CBD TO sCD32B AND sCD16A

[0076] The binding of h2B6-h3G8 CBD to sCD32B and sCD16A was assayed in a sandwich ELISA. sCD32B was used as the target protein. The secondary probe was HRP conjugated sCD16A. h3G8 CMD, which binds CD16A, was used as control.

FIGS. 6 A-C BIACORE ANALYSIS OF DIABODY BINDING TO sCD16A, sCD32B AND sCD32A

[0077] The binding of h2B6-h3G8 CBD, h2B6 CMD and h3G8 CMD to sCD16A, sCD32B, and sCD32A (negative control) was assayed by SPR analysis. h3G8 scFv was also tested as a control. (A) Binding to sCD16; (B) Binding to sCD32B and (C) Binding to sCD32A. Diabodies were injected at a concentration of 100 nM, and scFv at a concentration of 200 nM, over receptor surfaces at a flow rate of 50 ml/min for 60 sec.

FIGS. 7 A-C BIACORE ANALYSIS OF DIABODY BINDING TO sCD16A and sCD32B

[0078] The binding of h2B6-h3G8 CBD, h2B6 CMD and h3G8 CMD to sCD16A, and sCD32B was assayed by SPR analysis. h3G8 scFv was also tested as a control. (A) Binding of h3G8 CMD to sCD16A; (B) Binding of h2B6-h3G8 CBD to sCD16A; (C) Binding of h3G8 scFv to sCD16A; (D) Binding of h2B6 CMD to sCD32B; and (E) Binding of h2B6-h3G8 CBD to sCD32B. Diabodies were injected at concentrations of 6.25-200 nM over receptor surfaces at a flow rate of 70 ml/min for 180 sec.

FIG. 8 SCHEMATIC DEPICTING THE INTERACTION OF POLYPEPTIDE CHAINS COMPRISING VL AND VH DOMAINS TO FORM A COVALENT BISPECIFIC DIABODY MOLECULE

[0079] NH₂ and COOH represent the amino-terminus and carboxy terminus, respectively of each polypeptide chain. S represents the C-terminal cysteine residue on each polypeptide chain. VL and VH indicate the variable light domain and variable heavy domain, respectively. Dotted and dashed lines are to distinguish between the two polypeptide chains and, in particular, represent the linker portions of said chains. h2B6 Fv and h3G8 Fv indicate an epitope binding site specific for CD32B and CD16, respectively.

FIG. 9 SCHEMATIC REPRESENTATION OF POLYPEPTIDE CHAINS CONTAINING Fc DOMAINS OF COVALENT BISPECIFIC DIABODIES

[0080] Representation of polypeptide constructs of the diabody molecules of the invention (each construct is described from the amino terminus ("n"), left side of the construct, to the carboxy terminus ("c"), right side of figure). Construct (5) (SEQ ID NO:14) comprised, n-VL domain Hu2B6 - a first linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu3G8 - a second linker (LGGC)- and a C-terminal Fc domain of human IgG1-c; construct (6) (SEQ ID NO:15) comprised n-the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 - and second linker (LGGC)- and a C-terminal Fc domain of human IgG1-c; construct (7) (SEQ ID NO:16) comprised n-the VL domain Hu2B6 - a first linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu3G8 - and a C-terminal sequence (LGGCFNRGEC) (SEQ ID NO:17)-c; construct (8) (SEQ ID NO:18) comprised n-the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 - and second linker (LGGC)-and a C-terminal hinge/Fc domain of human IgG1 (with amino acid substitution A215V)-c.

FIG.10 BINDING OF DIABODY MOLECULES COMPRISING Fc DOMAINS TO sCD32B AND sCD16A

[0081] The binding of diabody molecules comprising Fc domains to sCD32B and sCD16A was assayed in a sandwich ELISA. Diabodies assayed were produced by 3 recombinant expression systems: cotransfection of pMGX669 and pMGX674, expressing constructs 1 and 6, respectively; cotransfection of pMGX667 and pMGX676, expressing constructs 2 and 5, respectively; and cotransfection of pMGX674 and pMGX676, expressing constructs 5 and 6, respectively. sCD32B was used as the target protein. The secondary probe was HRP conjugated sCD16A.

FIG. 11 SCHEMATIC DEPICTING THE INTERACTION OF TWO POLYPEPTIDE CHAINS EACH COMPRISING AN Fc DOMAIN TO FORM A BIVALENT, COVALENT DIABODY

[0082] NH₂ and COOH represent the amino-terminus and carboxy terminus, respectively of each polypeptide chain. S represents the at least one disulfide bond between a cysteine residue in the second linker sequence of each polypeptide chain. VL and VH indicate the variable light domain and variable heavy domain, respectively. Dotted and dashed lines are to distinguish between the two polypeptide chains and, in particular, represent the first linker portions of said chains. CH2 and CH3 represent the CH2 and CH3 constant domains of an Fc domain. h2B6 Fv and h3G8 Fv indicate an epitope binding site specific for CD32B and CD16, respectively.

FIG.12 BINDING OF DIABODY MOLECULES COMPRISING HINGE/Fc DOMAINS TO sCD32B AND sCD16A

[0083] The binding of diabody molecules comprising Fc domains to sCD32B and sCD16A was assayed in a sandwich ELISA. Diabodies assayed were produced by 4 recombinant expression systems: cotransfection of pMGX669 + pMGX674, expressing constructs 1 and 6, respectively; cotransfection of pMGX669 + pMGX678, expressing constructs 2 and 8, respectively; cotransfection of pMGX677 + pMGX674, expressing constructs 7 and 6, respectively; and cotransfection of pMGX677 + pMGX678, expressing constructs 7 and 8, respectively. sCD32B was used as the target protein. The secondary probe was HRP conjugated sCD16A.

FIG. 13 SCHEMATIC DEPICTING THE INTERACTION OF POLYPEPTIDE CHAINS TO FORM A TETRAMERIC DIABODY MOLECULE

[0084] NH₂ and COOH represent the amino-terminus and carboxy terminus, respectively of each polypeptide chain. S represents the at least one disulfide bond between

a cysteine residue in the second linker sequence the Fc bearing, 'heavier,' polypeptide chain and a cysteine residue in the C-terminal sequence of the non-Fc bearing, 'lighter,' polypeptide chain. VL and VH indicate the variable light domain and variable heavy domain, respectively. Dotted and dashed lines are to distinguish between polypeptide chains and, in particular, represent the first linker portions of said heavier chains or the linker of said lighter chains. CH2 and CH3 represent the CH2 and CH3 constant domains of an Fc domain. h2B6 Fv and h3G8 Fv indicate an epitope binding site specific for CD32B and CD16, respectively.

FIG. 14 SCHEMATIC REPRESENTATION OF POLYPEPTIDES CHAINS CONTAINING Fc DOMAINS WHICH Form COVALENT BISPECIFIC DIABODIES

[0085] Representation of polypeptide constructs which form the diabody molecules of the invention (each construct is described from the amino terminus ("n"), left side of the construct, to the carboxy terminus ("c"), right side of figure). Construct (9) (SEQ ID NO:19) comprised n-a Hinge/Fc domain of human IgG1 - the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 - linker (GGGSGGGG (SEQ ID NO:10))- and a C-terminal LGGC sequence-c; construct (10) (SEQ ID NO:20) comprised n-an Fc domain of human IgG1 - the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 - linker (GGGSGGGG (SEQ ID NO:10))- and a C-terminal LGGC sequence-c; construct (11) (SEQ ID NO:21) comprised n-the VL domain Hu2B6 (G105C) - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu3G8 - and a C-terminal hinge/Fc domain of human IgG1 with amino acid substitution A215V-c; construct (12) (SEQ ID NO:22) comprised n-the VL domain Hu3G8 - linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6 (G44C) - and a C-terminal FNRGEC (SEQ ID NO:23) sequence-c.

FIG. 15 A-B SDS-PAGE AND WESTERN BLOT ANALYSIS OF AFFINITY TETRAMERIC DIABODIES

[0086] Diabodies produced by recombinant expression systems cotransfected with vectors expressing constructs 10 and 1, constructs 9 and 1, and constructs 11 and 12 were subjected to SDS-PAGE analysis non-reducing conditions (A) and Western Blot analysis using goat anti-human IgG1 H+L as the probe (B). Proteins in the SDS-PAGE gel were visualized with Simply Blue Safestain (Invitrogen). For both panels A and B, diabody molecules comprising constructs 10 and 1, constructs 9 and 1, and constructs 11 and 12A are in lanes 1, 2 and 3, respectively.

FIG.16 BINDING OF DIABODY MOLECULES COMPRISING Fc DOMAINS AND ENGINEERED INTERCHAIN DISULFIDE BONDS TO sCD32B AND sCD16A

[0087] The binding of diabody molecules comprising Fc domains and engineered disulfide bonds between the 'lighter' and 'heavier' polypeptide chains to sCD32B and sCD16A was assayed in a sandwich ELISA. Diabodies assayed were produced by 3 recombinant expression systems: expressing constructs 1 and 10, expressing constructs 1 and 9, and expressing constructs 11 and 12, respectively. sCD32B was used as the target protein. The secondary probe was HRP conjugated sCD16A. Binding of h3G8 was used as control.

FIG. 17 SCHEMATIC REPRESENTATION OF POLYPROTEIN PRECURSOR OF DIABODY MOLECULE AND SHCEMATIC REPRESENTATION OF POLYPEPTIDE CHAINS CONTAINING LAMBDA LIGHT CHAIN AND/OR HINGE DOMAINS

[0088] Representation of polypeptide constructs which comprise the diabody molecules of the invention (each construct is described from the amino terminus ("n"), left side of the construct, to the carboxy terminus ("c"), right side of figure). Construct (13) (SEQ ID NO:97) comprised, n-VL domain 3G8 - a first linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of 2.4G2VH - a second linker (LGGC)- furin recognition site (RAKR (SEQ ID NO:95))-VL domain of 2.4G2- a third linker (GGGSGGGG (SEQ ID NO:10))-VH domain of 3G8- and a C-terminal LGGC domain; (nucleotide sequence encoding SEQ ID NO:97 is provided in SEQ ID NO:98). Construct (14) (SEQ ID NO:99) comprised, n-VL domain 3G8 - a first linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of 2.4G2VH - a second linker (LGGC)- furin recognition site (RAKR (SEQ ID NO:95))-FMD (Foot and Mouth Disease Virus Protease C3) site-VL domain of 2.4G2- a third linker (GGGSGGGG (SEQ ID NO:10))-VH domain of 3G8- and a C-terminal LGGC domain; (nucleotide sequence encoding SEQ ID NO:99 is provided in SEQ ID NO:100). Construct (15) (SEQ ID NO:101) comprised, n-VL domain Hu2B6 - a linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu3G8-and a C-terminal FNRGEC (SEQ ID NO:23) domain; (nucleotide sequence encoding SEQ ID NO:101 is provided in SEQ ID NO:102). Construct (16) (SEQ ID NO:103) comprised, n-VL domain Hu3G8 - a linker (GGGSGGGG (SEQ ID NO:10)) - the VH domain of Hu2B6-and a C-terminal VEPKSC (SEQ ID NO:79) domain; (nucleotide sequence encoding SEQ ID NO:103 is provided in SEQ ID NO:104).

FIG. 18 BINDING OF DIABODY MOLECULES DERIVED FROM A POLYPROTEIN PRECURSOR MOLECULE TO mCD32B AND sCD16A

[0089] The binding of diabody molecules derived from the polyprotein precursor molecule construct 13 (SEQ ID NO:97) to murine CD32B (mCD32B) and soluble CD16A (sCD16A) was assayed in a sandwich ELISA. mCD32B was used as the target protein. The secondary probe was biotin conjugated sCD16A.

FIG.19 BINDING OF DIABODY MOLECULES COMPRISING LAMBDA CHAIN AND/OR HINGE DOMAINS TO sCD32B AND sCD16A

[0090] The binding of diabody molecules comprising domains derived from the C-terminus of the human lambda light chain and/or the hinge domain of IgG to sCD32B and sCD16A was assayed and compared to the diabody comprising constructs 1 and 2 (FIG. 5) in a sandwich ELISA. Diabodies assayed were produced by the recombinant expression system expressing constructs 15 and 16 (SEQ ID NO:101 and SEQ ID NO:103, respectively). sCD32B was used as the target protein. The secondary probe was HRP conjugated sCD16A. Bars with small boxes represent the construct 15/16 combination while bars with large boxes represent construct 1/2 combination.

5. DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0091] Each polypeptide chain of the diabody molecule comprises a VL domain and a VH domain, which are covalently linked such that the domains are constrained from self assembly. Interaction of two of the polypeptide chains will produce two VL-VH pairings, forming two epitope binding sites, *i.e.*, a bivalent molecule. Neither the VH or VL domain is constrained to any position within the polypeptide chain, *i.e.*, restricted to the amino (N) or carboxy (C) terminus, nor are the domains restricted in their relative positions to one another, *i.e.*, the VL domain may be N-terminal to the VH domain and vice-versa. The only restriction is that a complimentary polypeptide chain be available in order to form functional diabody. Where the VL and VH domains are derived from the same antibody, the two complimentary polypeptide chains may be identical. For example, where the binding domains are derived from an antibody specific for epitope A (*i.e.*, the binding domain is formed from a VL_A-VH_A interaction), each polypeptide will comprise a VH_A and a VL_A. Homodimerization of two polypeptide chains of the antibody will result in the formation two VL_A-VH_A binding sites, resulting in a bivalent monospecific antibody. Where the VL and VH domains are derived from antibodies specific for different antigens, formation of a functional bispecific diabody requires the interaction of two different polypeptide chains, *i.e.*, formation of a heterodimer. For example, for a bispecific diabody,

one polypeptide chain will comprise a VL_A and a VL_B; homodimerization of said chain will result in the formation of two VL_A-VH_B binding sites, either of no binding or of unpredictable binding. In contrast, where two differing polypeptide chains are free to interact, *e.g.*, in a recombinant expression system, one comprising a VL_A and a VH_B and the other comprising a VL_B and a VH_A, two differing binding sites will form: VL_A-VH_A and VL_B-VH_B. For all diabody polypeptide chain pairs, the possibility of misalignment or mis-binding of the two chains is a possibility, *i.e.*, interaction of VL-VL or VH-VH domains; however, purification of functional diabodies is easily managed based on the immunospecificity of the properly dimerized binding site using any affinity based method known in the art or exemplified herein, *e.g.*, affinity chromatography.

[0092] In other embodiments, one or more of the polypeptide chains of the diabody comprises an Fc domain. Fc domains in the polypeptide chains of the diabody molecules preferentially dimerize, resulting in the formation of a diabody molecule that exhibits immunoglobulin-like properties, *e.g.*, Fc-FcγR, interactions. Fc comprising diabodies may be dimers, *e.g.*, comprised of two polypeptide chains, each comprising a VH domain, a VL domain and an Fc domain. Dimerization of said polypeptide chains results in a bivalent diabody comprising an Fc domain, albeit with a structure distinct from that of an unmodified bivalent antibody (FIG.11). Such diabody molecules will exhibit altered phenotypes relative to a wild-type immunoglobulin, *e.g.*, altered serum half-life, binding properties, etc. In other embodiments, diabody molecules comprising Fc domains may be tetramers. Such tetramers comprise two 'heavier' polypeptide chains, *i.e.* a polypeptide chain comprising a VL, a VH and an Fc domain, and two 'lighter' polypeptide chains, *i.e.*, polypeptide chain comprising a VL and a VH. Said lighter and heavier chains interact to form a monomer, and said monomers interact via their unpaired Fc domains to form an Ig-like molecule. Such an Ig-like diabody is tetravalent and may be monospecific, bispecific or tetraspecific.

[0093] The at least two binding sites of the diabody molecule can recognize the same or different epitopes. Different epitopes can be from the same antigen or epitopes from different antigens. In one embodiment, the epitopes are from different cells. In another embodiment, the epitopes are cell surface antigens on the same cell or virus. The epitope binding sites can recognize any antigen to which an antibody can be generated. For example, proteins, nucleic acids, bacterial toxins, cell surface markers, autoimmune markers, viral proteins, drugs, etc. In particular aspects, at least one epitope binding site of the diabody is specific for an antigen on a particular cell, such as a B-cell or T-cell, a phagocytotic cell, a natural killer (NK) cell or a dendritic cell.

[0094] Each domain of the polypeptide chain of the diabody, *i.e.*, the VL, VH and FC domain may be separated by a peptide linker. The peptide linker may be 0, 1, 2, 3, 4, 5, 6, 7, 8, or 9. amino acids. In certain embodiments the amino acid linker sequence is GGGSGGGG (SEQ ID NO:10) encoded by the nucleic acid sequence (SEQ ID NO:76).

[0095] In certain embodiments, each polypeptide chain of the diabody molecule is engineered to comprise at least one cysteine residue that will interact with a counterpart at least one cysteine residue on a second polypeptide chain of the invention to form an inter-chain disulfide bond. Said interchain disulfide bonds serve to stabilize the diabody molecule, improving expression and recovery in recombinant systems, resulting in a stable and consistent formulation as well as improving the stability of the isolated and/or purified product *in vivo*. Said at least one cysteine residue may be introduced as a single amino acid or as part of larger amino-acid sequence, *e.g.* hinge domain, in any portion of the polypeptide chain. In a specific embodiment, said at least one cysteine residue is engineered to occur at the C-terminus of the polypeptide chain. In some embodiments, said at least one cysteine residue is introduced into the polypeptide chain within the amino acid sequence LGGC. In a specific embodiment, the C-terminus of the polypeptide chain comprising the diabody molecule of the invention comprises the amino acid sequence LGGC. In another embodiment, said at least one cysteine residue is introduced into the polypeptide within an amino acid sequence comprising a hinge domain, *e.g.* SEQ ID NO:1 or SEQ ID NO:4. In a specific embodiment, the C-terminus of a polypeptide chain of the diabody molecule of the invention comprises the amino acid sequence of an IgG hinge domain, *e.g.* SEQ ID NO:1. In another embodiment, the C-terminus of a polypeptide chain of a diabody molecule of the invention comprises the amino acid sequence VEPKSC (SEQ ID NO:79), which can be encoded by nucleotide sequence (SEQ ID NO:80). In other embodiments, said at least one cysteine residue is introduced into the polypeptide chain within the amino acid sequence LGGCFNRGEC (SEQ ID NO:17), which can be encoded by the nucleotide sequence (SEQ ID NO:78). In a specific embodiment, the C-terminus of a polypeptide chain comprising the diabody of the invention comprises the amino acid sequence LGGCFNRGEC (SEQ ID NO:17), which can be encoded by the nucleotide sequence (SEQ ID NO:78). In yet other embodiments, said at least one cysteine residue is introduced into the polypeptide chain within the amino acid sequence FNRGEC (SEQ ID NO:23), which can be encoded by the nucleotide sequence (SEQ ID NO:77). In a specific embodiment, the C-terminus of a polypeptide chain comprising the diabody of the invention comprises the amino acid sequence FNRGEC (SEQ ID NO:23), which can be encoded by the nucleotide sequence (SEQ ID NO:77).

[0096] In certain embodiments, the diabody molecule comprises at least two polypeptide chains, each of which comprise the amino acid sequence LGGC and are covalently linked by a disulfide bond between the cysteine residues in said LGGC sequences. In another specific embodiment, the diabody molecule comprises at least two polypeptide chains, one of which comprises the sequence FNRGEC (SEQ ID NO:23) while the other comprises a hinge domain (containing at least one cysteine residue), wherein said at least two polypeptide chains are covalently linked by a disulfide bond between the cysteine residue in FNRGEC (SEQ ID NO:23) and a cysteine residue in the hinge domain. In particular aspects, the cysteine residue responsible for the disulfide bond located in the hinge domain is Cys-128 (as numbered according to Kabat EU; located in the hinge domain of an unmodified, intact IgG heavy chain) and the counterpart cysteine residue in SEQ ID NO:23 is Cys-214 (as numbered according to Kabat EU; located at the C-terminus of an unmodified, intact IgG light chain) (Elkabetz et al., 2005, J. Biol. Chem. 280:14402-14412; hereby incorporated by reference herein in its entirety). In yet other embodiments, the at least one cysteine residue is engineered to occur at the N-terminus of the amino acid chain. In still other embodiments, the at least one cysteine residue is engineered to occur in the linker portion of the polypeptide chain of the diabody molecule. In further embodiments, the VH or VL domain is engineered to comprise at least one amino acid modification relative to the parental VH or VL domain such that said amino acid modification comprises a substitution of a parental amino acid with cysteine.

[0097] The invention encompasses diabody molecules comprising an Fc domain or portion thereof (*e.g.* a CH2 domain, or CH3 domain). The Fc domain or portion thereof may be derived from any immunoglobulin isotype or allotype including, but not limited to, IgA, IgD, IgG, IgE and IgM. In preferred embodiments, the Fc domain (or portion thereof) is derived from IgG. In specific embodiments, the IgG isotype is IgG1, IgG2, IgG3 or IgG4 or an allotype thereof. In one embodiment, the diabody molecule comprises an Fc domain, which Fc domain comprises a CH2 domain and CH3 domain independently selected from any immunoglobulin isotype (*i.e.* an Fc domain comprising the CH2 domain derived from IgG and the CH3 domain derived from IgE, or the CH2 domain derived from IgG1 and the CH3 domain derived from IgG2, etc.). Said Fc domain may be engineered into a polypeptide chain comprising the diabody molecule of the invention in any position relative to other domains or portions of said polypeptide chain (*e.g.*, the Fc domain, or portion thereof, may be c-terminal to both the VL and VH domains of the polypeptide of the chain; may be n-terminal to both the VL and VH domains; or may be N-terminal to one domain and c-terminal to another (*i.e.*, between two domains of the polypeptide chain)).

[0098] The present invention also encompasses molecules comprising a hinge domain. The hinge domain be derived from any immunoglobulin isotype or allotype including IgA, IgD, IgG, IgE and IgM. In preferred embodiments, the hinge domain is derived from IgG, wherein the IgG isotype is IgG1, IgG2, IgG3 or IgG4, or an allotype thereof. Said hinge domain may be engineered into a polypeptide chain comprising the diabody molecule together with an Fc domain such that the diabody molecule comprises a hinge-Fc domain. In certain embodiments, the hinge and Fc domain are independently selected from any immunoglobulin isotype known in the art or exemplified herein. In other embodiments the hinge and Fc domain are separated by at least one other domain of the polypeptide chain, *e.g.*, the VL domain. The hinge domain, or optionally the hinge-Fc domain, may be engineered in to a polypeptide of the invention in any position relative to other domains or portions of said polypeptide chain. In certain embodiments, a polypeptide chain of the invention comprises a hinge domain, which hinge domain is at the C-terminus of the polypeptide chain, wherein said polypeptide chain does not comprise an Fc domain. In yet other embodiments, a polypeptide chain of the invention comprises a hinge-Fc domain, which hinge-Fc domain is at the C-terminus of the polypeptide chain. In further embodiments, a polypeptide chain of the invention comprises a hinge-Fc domain, which hinge-Fc domain is at the N-terminus of the polypeptide chain.

[0099] As discussed above, the invention encompasses multimers of polypeptide chains, each of which polypeptide chains comprise a VH and VL domain. In certain aspects, the polypeptide chains in said multimers further comprise an Fc domain. Dimerization of the Fc domains leads to formation of a diabody molecule that exhibits immunoglobulin-like functionality, *i.e.*, Fc mediated function (*e.g.*, Fc-FcγR interaction, complement binding, etc.). In certain embodiments, the VL and VH domains comprising each polypeptide chain have the same specificity, and said diabody molecule is bivalent and monospecific. In other embodiments, the VL and VH domains comprising each polypeptide chain have differing specificity and the diabody is bivalent and bispecific.

[00100] In yet other embodiments, diabody molecules of the invention encompass tetramers of polypeptide chains, each of which polypeptide chain comprises a VH and VL domain. In certain embodiments, two polypeptide chains of the tetramer further comprise an Fc domain. The tetramer is therefore comprised of two 'heavier' polypeptide chains, each comprising a VL, VH and Fc domain, and two 'lighter' polypeptide chains, comprising a VL and VH domain. Interaction of a heavier and lighter chain into a bivalent monomer coupled with dimerization of said monomers via the Fc domains of the heavier chains will lead to formation of a tetravalent immunoglobulin-like molecule (exemplified in Example

6.2 and Example 6.3). In certain aspects the monomers are the same, and the tetravalent diabody molecule is monospecific or bispecific. In other aspects the monomers are different, and the tetra valent molecule is bispecific or tetraspecific.

[00101] Formation of a tetraspecific diabody molecule as described *supra* requires the interaction of four differing polypeptide chains. Such interactions are difficult to achieve with efficiency within a single cell recombinant production system, due to the many variants of potential chain mispairings. One solution to increase the probability of mispairings, is to engineer “knobs-into-holes” type mutations into the desired polypeptide chain pairs. Such mutations favor heterodimerization over homodimerization. For example, with respect to Fc-Fc-interactions, an amino acid substitution (preferably a substitution with an amino acid comprising a bulky side group forming a ‘knob’, *e.g.*, tryptophan) can be introduced into the CH2 or CH3 domain such that steric interference will prevent interaction with a similarly mutated domain and will obligate the mutated domain to pair with a domain into which a complementary, or accommodating mutation has been engineered, *i.e.*, ‘the hole’ (*e.g.*, a substitution with glycine). Such sets of mutations can be engineered into any pair of polypeptides comprising the diabody molecule, and further, engineered into any portion of the polypeptides chains of said pair. Methods of protein engineering to favor heterodimerization over homodimerization are well known in the art, in particular with respect to the engineering of immunoglobulin-like molecules, and are encompassed herein (*see e.g.*, Ridgway et al., 1996, Protein Engr. 9:617-621, Atwell et al., 1997, J. Mol. Biol. 270: 26-35, and Xie et al., 2005, J. Immunol. Methods 296:95-101; each of which is hereby incorporated herein by reference in its entirety).

[00102] The invention also encompasses diabody molecules comprising variant Fc or variant hinge-Fc domains (or portion thereof), which variant Fc domain comprises at least one amino acid modification (*e.g.* substitution, insertion deletion) relative to a comparable wild-type Fc domain or hinge-Fc domain (or portion thereof). Molecules comprising variant Fc domains or hinge-Fc domains (or portion thereof) (*e.g.*, antibodies) normally have altered phenotypes relative to molecules comprising wild-type Fc domains or hinge-Fc domains or portions thereof. The variant phenotype may be expressed as altered serum half-life, altered stability, altered susceptibility to cellular enzymes or altered effector function as assayed in an NK dependent or macrophage dependent assay. Fc domain variants identified as altering effector function are disclosed in International Application WO04/063351, U.S. Patent Application Publications 2005/0037000 and 2005/0064514, U.S. Provisional Applications 60/626,510, filed November 10, 2004, 60/636,663, filed December 15, 2004, and 60/781,564, filed March 10, 2006, and U.S. Patent Applications

11/271, 140, filed November 10, 2005, and 11/305,787, filed December 15, 2005, concurrent applications of the Inventors, each of which is incorporated by reference in its entirety.

[00103] The bispecific diabodies of the invention can simultaneously bind two separate and distinct epitopes. In certain embodiments the epitopes are from the same antigen. In other embodiments, the epitopes are from different antigens. In preferred embodiments, at least one epitope binding site is specific for a determinant expressed on an immune effector cell (*e.g.* CD3, CD16, CD32, CD64, etc.) which are expressed on T lymphocytes, natural killer (NK) cells or other mononuclear cells. In one embodiment, the diabody molecule binds to the effector cell determinant and also activates said effector cell. In this regard, diabody molecules of the invention may exhibit Ig-like functionality independent of whether they further comprise an Fc domain (*e.g.*, as assayed in any effector function assay known in the art or exemplified herein (*e.g.*, ADCC assay). In certain embodiments the bispecific diabody of the invention binds both a cancer antigen on a tumor cell and an effector cell determinant while activating said cell. In alternative embodiments, the bispecific diabody or diabody molecule of the invention may inhibit activation of a target, *e.g.*, effector, cell by simultaneously binding, and thus linking, an activating and inhibitory receptor on the same cell (*e.g.*, bind both CD32A and CD32B, BCR and CD32B, or IgE1 and CD32B) as described *supra* (*see*, Background Section). In a further aspect of this embodiment, the bispecific diabody may exhibit anti-viral properties by simultaneously binding two neutralizing epitopes on a virus (*e.g.*, RSV epitopes; WNV epitopes such as E16 and E53).

[00104] In certain embodiments, bispecific diabody molecules of the invention offer unique opportunities to target specific cell types. For example, the bispecific diabody or diabody molecule can be engineered to comprise a combination of epitope binding sites that recognize a set of antigens unique to a target cell or tissue type. Additionally, where either or both of the individual antigens is/are fairly common separately in other tissue and/or cell types, low affinity binding domains can be used to construct the diabody or diabody molecule. Such low affinity binding domains will be unable to bind to the individual epitope or antigen with sufficient avidity for therapeutic purposes. However, where both epitopes or antigens are present on a single target cell or tissue, the avidity of the diabody or diabody molecule for the cell or tissue, relative to a cell or tissue expressing only one of the antigens, will be increased such that said cell or tissue can be effectively targeted by the invention. Such a bispecific molecule can exhibit enhanced binding to one or both of its

target antigens on cells expressing both of said antigens relative to a monospecific diabody or an antibody with a specificity to only one of the antigens.

[00105] Preferably, the binding properties of the diabodies of the invention are characterized by *in vitro* functional assays for determining binding activity and/or one or more FcγR mediator effector cell functions (mediated via Fc-FcγR interactions or by the immunospecific binding of a diabody molecule to an FcγR) (See Section 5.4.2 and 5.4.3). The affinities and binding properties of the molecules, *e.g.*, diabodies, of the invention for an FcγR can be determined using *in vitro* assays (biochemical or immunological based assays) known in the art for determining binding domain-antigen or Fc-FcγR interactions, *i.e.*, specific binding of an antigen to a binding domain or specific binding of an Fc region to an FcγR, respectively, including but not limited to ELISA assay, surface plasmon resonance assay, immunoprecipitation assays (See Section 5.4.2). In most preferred embodiments, the molecules of the invention have similar binding properties in *in vivo* models (such as those described and disclosed herein) as those in *in vitro* based assays. However, the present invention does not exclude molecules of the invention that do not exhibit the desired phenotype in *in vitro* based assays but do exhibit the desired phenotype *in vivo*.

[00106] In some embodiments, molecules of the invention are engineered to comprise an altered glycosylation pattern or an altered glycoform relative to the comparable portion of the template molecule. Engineered glycoforms may be useful for a variety of purposes, including, but not limited to, enhancing effector function. Engineered glycoforms may be generated by any method known to one skilled in the art, for example by using engineered or variant expression strains, by co-expression with one or more enzymes, for example, DI N-acetylglucosaminyltransferase III (GnTII1), by expressing a diabody of the invention in various organisms or cell lines from various organisms, or by modifying carbohydrate(s) after the diabody has been expressed and purified. Methods for generating engineered glycoforms are known in the art, and include but are not limited to those described in Umana et al, 1999, Nat. Biotechnol 17:176-180; Davies *et al.*, 2001 Biotechnol Bioeng 74:288-294; Shields et al, 2002, J Biol Chem 277:26733-26740; Shinkawa et al., 2003, J Biol Chem 278:3466-3473) US 6,602,684; USSN 10/277,370; USSN 10/113,929; PCT WO 00/61739A1; PCT WO 01/292246A1; PCT WO 02/311140A1; PCT WO 02/30954A1; Potillegent™ technology (Biowa, Inc. Princeton, NJ); GlycoMAb™ glycosylation engineering technology (GLYCART biotechnology AG, Zurich, Switzerland); each of which is incorporated herein by reference in its entirety. See, *e.g.*, WO 00061739; EA01229125; US 20030115614; Okazaki *et al.*, 2004, JMB, 336: 1239-49 each of which is incorporated herein by reference in its entirety.

[00107] The invention further encompasses incorporation of unnatural amino acids to generate the diabodies of the invention. Such methods are known to those skilled in the art such as those using the natural biosynthetic machinery to allow incorporation of unnatural amino acids into proteins, *see, e.g.*, Wang *et al.*, 2002 Chem. Comm. 1: 1-11; Wang *et al.*, 2001, Science, 292: 498-500; van Hest *et al.*, 2001. Chem. Comm. 19: 1897-1904, each of which is incorporated herein by reference in its entirety. Alternative strategies focus on the enzymes responsible for the biosynthesis of amino acyl-tRNA, *see, e.g.*, Tang *et al.*, 2001, J. Am. Chem. 123(44): 11089-11090; Kiick *et al.*, 2001, FEBS Lett. 505(3): 465; each of which is incorporated herein by reference in its entirety.

[00108] In some embodiments, the invention encompasses methods of modifying a VL, VH or Fc domain of a molecule of the invention by adding or deleting a glycosylation site. Methods for modifying the carbohydrate of proteins are well known in the art and encompassed within the invention, *see, e.g.*, U.S. Patent No. 6,218,149; EP 0 359 096 B1; U.S. Publication No. US 2002/0028486; WO 03/035835; U.S. Publication No. 2003/0115614; U.S. Patent No. 6,218,149; U.S. Patent No. 6,472,511; all of which are incorporated herein by reference in their entirety.

5.1 DIABODY BINDING DOMAINS

[00109] The diabodies of the present invention comprise antigen binding domains generally derived from immunoglobulins or antibodies. The antibodies from which the binding domains used in the methods of the invention are derived may be from any animal origin including birds and mammals (*e.g.*, human, non-human primate, murine, donkey, sheep, rabbit, goat, guinea pig, camel, horse, or chicken). Preferably, the antibodies are human or humanized monoclonal antibodies. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or libraries of synthetic human immunoglobulin coding sequences or from mice that express antibodies from human genes.

[00110] The invention contemplates the use of any antibodies known in the art for the treatment and/or prevention of cancer, autoimmune disease, inflammatory disease or infectious disease as source of binding domains for the diabodies of the invention. Non-limiting examples of known cancer antibodies are provided in section 5.7.1 as well as other antibodies specific for the listed target antigens and antibodies against the cancer antigens listed in section 5.6.1; nonlimiting examples of known antibodies for the treatment and/or prevention of autoimmune disease and inflammatory disease are provided in section 5.7.2. as well as antibodies against the listed target antigens and antibodies against the antigens

listed in section 5.6.2; in other embodiments antibodies against epitopes associated with infectious diseases as listed in Section 5.6.3 can be used. In certain embodiments, the antibodies comprise a variant Fc region comprising one or more amino acid modifications, which have been identified by the methods of the invention to have a conferred effector function and/or enhanced affinity for Fc γ RIIB and a decreased affinity for Fc γ RIIA relative to a comparable molecule comprising a wild type Fc region. A non-limiting example of the antibodies that are used for the treatment or prevention of inflammatory disorders which can be engineered according to the invention is presented in Table 9, and a non-limiting example of the antibodies that are used for the treatment or prevention of autoimmune disorder is presented in Table 10.

[00111] For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use diabodies with variable domains derived from human, chimeric or humanized antibodies. Variable domains from completely human antibodies are particularly desirable for therapeutic treatment of human subjects. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also U.S. Patent Nos. 4,444,887 and 4,716,111; and International Publication Nos. WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

[00112] A humanized antibody is an antibody, a variant or a fragment thereof which is capable of binding to a predetermined antigen and which comprises a framework region having substantially the amino acid sequence of a human immunoglobulin and a CDR having substantially the amino acid sequence of a non-human immunoglobulin. A humanized antibody may comprise substantially all of at least one, and typically two, variable domains in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin (*i.e.*, donor antibody) and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence.

[00113] The framework and CDR regions of a humanized antibody need not correspond precisely to the parental sequences, *e.g.*, the donor CDR or the consensus framework may be mutagenized by substitution, insertion or deletion of at least one residue so that the CDR or framework residue at that site does not correspond to either the consensus or the donor antibody. Such mutations, however, are preferably not extensive. Usually, at least 75% of the humanized antibody residues will correspond to those of the parental framework region (FR) and CDR sequences, more often 90%, and most preferably

greater than 95%. Humanized antibodies can be produced using variety of techniques known in the art, including but not limited to, CDR-grafting (European Patent No. EP 239,400; International Publication No. WO 91/09967; and U.S. Patent Nos. 5,225,539, 5,530,101, and 5,585,089), veneering or resurfacing (European Patent Nos. EP 592,106 and EP 519,596; Padlan, 1991, *Molecular Immunology* 28(4/5):489-498; Studnicka *et al.*, 1994, *Protein Engineering* 7(6):805-814; and Roguska *et al.*, 1994, *Proc Natl Acad Sci USA* 91:969-973), chain shuffling (U.S. Patent No. 5,565,332), and techniques disclosed in, *e.g.*, U.S. Patent Nos. 6,407,213, 5,766,886, 5,585,089, International Publication No. WO 9317105, Tan *et al.*, 2002, *J. Immunol.* 169:1119-25, Caldas *et al.*, 2000, *Protein Eng.* 13:353-60, Morea *et al.*, 2000, *Methods* 20:267-79, Baca *et al.*, 1997, *J. Biol. Chem.* 272:10678-84, Roguska *et al.*, 1996, *Protein Eng.* 9:895-904, Couto *et al.*, 1995, *Cancer Res.* 55 (23 Supp):5973s-5977s, Couto *et al.*, 1995, *Cancer Res.* 55:1717-22, Sandhu, 1994, *Gene* 150:409-10, Pedersen *et al.*, 1994, *J. Mol. Biol.* 235:959-73, Jones *et al.*, 1986, *Nature* 321:522-525, Riechmann *et al.*, 1988, *Nature* 332:323, and Presta, 1992, *Curr. Op. Struct. Biol.* 2:593-596. Often, framework residues in the framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, *e.g.*, by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, *e.g.*, Queen *et al.*, U.S. Patent No. 5,585,089; U.S. Publication Nos. 2004/0049014 and 2003/0229208; U.S. Patent Nos. 6,350,861; 6,180,370; 5,693,762; 5,693,761; 5,585,089; and 5,530,101 and Riechmann *et al.*, 1988, *Nature* 332:323, all of which are incorporated herein by reference in their entireties.).

[00114] In a most preferred embodiment, the humanized binding domain specifically binds to the same epitope as the donor murine antibody. It will be appreciated by one skilled in the art that the invention encompasses CDR grafting of antibodies in general. Thus, the donor and acceptor antibodies may be derived from animals of the same species and even same antibody class or sub-class. More usually, however, the donor and acceptor antibodies are derived from animals of different species. Typically the donor antibody is a non-human antibody, such as a rodent MAb, and the acceptor antibody is a human antibody.

[00115] In some embodiments, at least one CDR from the donor antibody is grafted onto the human antibody. In other embodiments, at least two and preferably all three CDRs of each of the heavy and/or light chain variable regions are grafted onto the human antibody. The CDRs may comprise the Kabat CDRs, the structural loop CDRs or a

combination thereof. In some embodiments, the invention encompasses a humanized FcγRIIB antibody comprising at least one CDR grafted heavy chain and at least one CDR-grafted light chain.

[00116] The diabodies used in the methods of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the diabody. For example, but not by way of limitation, the diabody derivatives include diabodies that have been modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to, specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[00117] A chimeric antibody is a molecule in which different portions of the antibody are derived from different immunoglobulin molecules such as antibodies having a variable region derived from a non-human antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See *e.g.*, Morrison, 1985, *Science* 229:1202; Oi *et al.*, 1986, *BioTechniques* 4:214; Gillies *et al.*, 1989, *J. Immunol. Methods* 125:191-202; and U.S. Patent Nos. 6,311,415, 5,807,715, 4,816,567, and 4,816,397, which are incorporated herein by reference in their entirety.

[00118] Often, framework residues in the framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, *e.g.*, by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, *e.g.*, U.S. Patent No. 5,585,089; and Riechmann *et al.*, 1988, *Nature* 332:323, which are incorporated herein by reference in their entirety.)

[00119] Monoclonal antibodies from which binding domains of the diabodies of the invention can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, *et al.*, in: *Monoclonal Antibodies and T-Cell Hybridomas*, pp. 563-681 (Elsevier, N.Y., 1981) (both of which are incorporated by reference in their entirety). The term "monoclonal antibody"

as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

[00120] Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art. In a non-limiting example, mice can be immunized with an antigen of interest or a cell expressing such an antigen. Once an immune response is detected, *e.g.*, antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells. Hybridomas are selected and cloned by limiting dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding the antigen. Ascites fluid, which generally contains high levels of antibodies, can be generated by inoculating mice intraperitoneally with positive hybridoma clones. Antigens of interest include, but are not limited to, antigens associated with the cancers provided in section 5.8.1, antigens associated with the autoimmune diseases and inflammatory diseases provided in section 5.8.2, antigens associated with the infectious diseases provided in section 5.8.3, and the toxins provided in section 5.8.4.

[00121] Antibodies can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains, such as Fab and Fv or disulfide-bond stabilized Fv, expressed from a repertoire or combinatorial antibody library (*e.g.*, human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, *e.g.*, using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage, including fd and M13. The antigen binding domains are expressed as a recombinantly fused protein to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the immunoglobulins, or fragments thereof, of the present invention include those disclosed in Brinkman *et al.*, J. Immunol. Methods, 182:41-50, 1995; Ames *et al.*, J. Immunol. Methods, 184:177-186, 1995; Kettleborough *et al.*, Eur. J. Immunol., 24:952-958, 1994; Persic *et al.*, Gene, 187:9-18, 1997; Burton *et al.*, Advances in Immunology, 57:191-280, 1994; PCT Application No. PCT/GB91/01134; PCT Publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S.

Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

[00122] Phage display technology can be used to increase the affinity of an antibody for its antigen. This technique would be useful in obtaining high affinity antibodies. The technology, referred to as affinity maturation, employs mutagenesis or CDR walking and re-selection using the cognate antigen to identify antibodies that bind with higher affinity to the antigen when compared with the initial or parental antibody (See, *e.g.*, Glaser *et al.*, 1992, J. Immunology 149:3903). Mutagenizing entire codons rather than single nucleotides results in a semi-randomized repertoire of amino acid mutations. Libraries can be constructed consisting of a pool of variant clones each of which differs by a single amino acid alteration in a single CDR and which contain variants representing each possible amino acid substitution for each CDR residue. Mutants with increased binding affinity for the antigen can be screened by contacting the immobilized mutants with labeled antigen. Any screening method known in the art can be used to identify mutant antibodies with increased avidity to the antigen (*e.g.*, ELISA) (See Wu *et al.*, 1998, Proc Natl. Acad Sci. USA 95:6037; Yelton *et al.*, 1995, J. Immunology 155:1994). CDR walking which randomizes the light chain is also possible (See Schier *et al.*, 1996, J. Mol. Bio. 263:551).

[00123] The present invention also encompasses the use of binding domains comprising the amino acid sequence of any of the binding domains described herein or known in the art with mutations (*e.g.*, one or more amino acid substitutions) in the framework or CDR regions. Preferably, mutations in these binding domains maintain or enhance the avidity and/or affinity of the binding domains for FcγRIIB to which they immunospecifically bind. Standard techniques known to those skilled in the art (*e.g.*, immunoassays) can be used to assay the affinity of an antibody for a particular antigen.

[00124] Standard techniques known to those skilled in the art can be used to introduce mutations in the nucleotide sequence encoding an antibody, or fragment thereof, including, *e.g.*, site-directed mutagenesis and PCR-mediated mutagenesis, which results in amino acid substitutions. Preferably, the derivatives include less than 15 amino acid substitutions, less than 10 amino acid substitutions, less than 5 amino acid substitutions, less than 4 amino acid substitutions, less than 3 amino acid substitutions, or less than 2 amino acid substitutions relative to the original antibody or fragment thereof. In a preferred embodiment, the derivatives have conservative amino acid substitutions made at one or more predicted non-essential amino acid residues.

5.1.1 DIABODIES COMPRISING EPTIOPE BINDING SITES WHICH IMMUNOSPECIFICALLY BIND FcγRIIB

[00125] In a particular embodiment, at least one of the binding domains of the diabodies of the invention agonizes at least one activity of FcγRIIB. In one embodiment of the invention, said activity is inhibition of B cell receptor-mediated signaling. In another embodiment, the binding domain inhibits activation of B cells, B cell proliferation, antibody production, intracellular calcium influx of B cells, cell cycle progression, or activity of one or more downstream signaling molecules in the FcγRIIB signal transduction pathway. In yet another embodiment, the binding domain enhances phosphorylation of FcγRIIB or SHIP recruitment. In a further embodiment of the invention, the binding domain inhibits MAP kinase activity or Akt recruitment in the B cell receptor-mediated signaling pathway. In another embodiment, the binding domain agonizes FcγRIIB-mediated inhibition of FcεRI signaling. In a particular embodiment, said binding domain inhibits FcεRI-induced mast cell activation, calcium mobilization, degranulation, cytokine production, or serotonin release. In another embodiment, the binding domains of the invention stimulate phosphorylation of FcγRIIB, stimulate recruitment of SHIP, stimulate SHIP phosphorylation and its association with Shc, or inhibit activation of MAP kinase family members (*e.g.*, Erk1, Erk2, JNK, p38, etc.). In yet another embodiment, the binding domains of the invention enhance tyrosine phosphorylation of p62dok and its association with SHIP and rasGAP. In another embodiment, the binding domains of the invention inhibit FcγR-mediated phagocytosis in monocytes or macrophages.

[00126] In another embodiment, the binding domains antagonize at least one activity of FcγRIIB. In one embodiment, said activity is activation of B cell receptor-mediated signaling. In a particular embodiment, the binding domains enhance B cell activity, B cell proliferation, antibody production, intracellular calcium influx, or activity of one or more downstream signaling molecules in the FcγRIIB signal transduction pathway. In yet another particular embodiment, the binding domains decrease phosphorylation of FcγRIIB or SHIP recruitment. In a further embodiment of the invention, the binding domains enhance MAP kinase activity or Akt recruitment in the B cell receptor mediated signaling pathway. In another embodiment, the binding domains antagonize FcγRIIB-mediated inhibition of FcεRI signaling. In a particular embodiment, the binding domains enhance FcεRI-induced mast cell activation, calcium mobilization, degranulation, cytokine production, or serotonin release. In another embodiment, the binding domains inhibit phosphorylation of FcγRIIB, inhibit recruitment of SHIP, inhibit SHIP phosphorylation and its association with Shc, enhance activation of MAP kinase family members (*e.g.*, Erk1,

Erk2, JNK, p38, etc.). In yet another embodiment, the binding domains inhibit tyrosine phosphorylation of p62dok and its association with SHIP and rasGAP. In another embodiment, the binding domains enhance FcγR-mediated phagocytosis in monocytes or macrophages. In another embodiment, the binding domains prevent phagocytosis, clearance of opsonized particles by splenic macrophages.

[00127] In other embodiments, at least one of the binding domains can be used to target the diabodies of the invention to cells that express FcγRIIB.

[00128] In one particular embodiment, one of the binding domains is derived from a mouse monoclonal antibody produced by clone 2B6 or 3H7, having ATCC accession numbers PTA-4591 and PTA-4592, respectively. Hybridomas producing antibodies 2B6 and 3H7 have been deposited with the American Type Culture Collection (10801 University Blvd., Manassas, VA. 20110-2209) on August 13, 2002 under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedures, and assigned accession numbers PTA-4591 (for hybridoma producing 2B6) and PTA-4592 (for hybridoma producing 3H7), respectively, and are incorporated herein by reference. In a preferred embodiment, the binding domains are human or have been humanized, preferably are derived from a humanized version of the antibody produced by clone 3H7 or 2B6.

[00129] The invention also encompasses diabodies with binding domains from other antibodies, that specifically bind FcγRIIB, preferably human FcγRIIB, more preferably native human FcγRIIB, that are derived from clones including but not limited to 1D5, 2E1, 2H9, 2D11, and 1F2 having ATCC Accession numbers, PTA-5958, PTA-5961, PTA-5962, PTA-5960, and PTA-5959, respectively. Hybridomas producing the above-identified clones were deposited under the provisions of the Budapest Treaty with the American Type Culture Collection (10801 University Blvd., Manassas, VA. 20110-2209) on May 7, 2004, and are incorporated herein by reference. In preferred embodiments, the binding domains from the antibodies described above are humanized.

[00130] In a specific embodiment, the binding domains used in the diabodies of the present invention are from an antibody or an antigen-binding fragment thereof (*e.g.*, comprising one or more complementarily determining regions (CDRs), preferably all 6 CDRs) of the antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2. In another embodiment, the binding domain binds to the same epitope as the mouse monoclonal antibody produced from clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2, respectively and/or competes with the mouse monoclonal antibody produced from clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2 as determined, *e.g.*, in an ELISA assay or other

appropriate competitive immunoassay, and also binds FcγRIIB with a greater affinity than the binding domain binds FcγRIIA.

[00131] The present invention also encompasses diabodies with binding domains comprising an amino acid sequence of a variable heavy chain and/or variable light chain that is at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of the variable heavy chain and/or light chain of the mouse monoclonal antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2. The present invention further encompasses diabodies with binding domains that specifically bind FcγRIIB with greater affinity than said antibody or fragment thereof binds FcγRIIA, and that comprise an amino acid sequence of one or more CDRs that is at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of one or more CDRs of the mouse monoclonal antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2. The determination of percent identity of two amino acid sequences can be determined by any method known to one skilled in the art, including BLAST protein searches.

[00132] The present invention also encompasses the use of diabodies containing binding domains that specifically bind FcγRIIB with greater affinity than binding domain binds FcγRIIA, which are encoded by a nucleotide sequence that hybridizes to the nucleotide sequence of the mouse monoclonal antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2 under stringent conditions. In a preferred embodiment, the binding domain specifically binds FcγRIIB with greater affinity than FcγRIIA, and comprises a variable light chain and/or variable heavy chain encoded by a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequence of the variable light chain and/or variable heavy chain of the mouse monoclonal antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2 under stringent conditions. In another preferred embodiment, the binding domains specifically bind FcγRIIB with greater affinity than FcγRIIA, and comprise one or more CDRs encoded by a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequence of one or more CDRs of the mouse monoclonal antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2. Stringent hybridization conditions include, but are not limited to, hybridization to filter-bound DNA in 6X sodium chloride/sodium citrate (SSC) at about 45°C followed by one or more washes in 0.2X SSC/0.1% SDS at about 50-65°C, highly stringent conditions such as hybridization to filter-bound DNA in 6X SSC at about 45°C followed by one or

more washes in 0.1X SSC/0.2% SDS at about 60°C, or any other stringent hybridization conditions known to those skilled in the art (see, for example, Ausubel, F.M. *et al.*, eds. 1989 Current Protocols in Molecular Biology, vol. 1, Green Publishing Associates, Inc. and John Wiley and Sons, Inc., NY at pages 6.3.1 to 6.3.6 and 2.10.3, incorporated herein by reference).

[00133] The present invention also encompasses the use of binding domains comprising the amino acid sequence of any of the binding domains described above with mutations (*e.g.*, one or more amino acid substitutions) in the framework or CDR regions. Preferably, mutations in these binding domains maintain or enhance the avidity and/or affinity of the binding domains for FcγRIIB to which they immunospecifically bind. Standard techniques known to those skilled in the art (*e.g.*, immunoassays) can be used to assay the affinity of an antibody for a particular antigen.

[00134] Standard techniques known to those skilled in the art can be used to introduce mutations in the nucleotide sequence encoding an antibody, or fragment thereof, including, *e.g.*, site-directed mutagenesis and PCR-mediated mutagenesis, which results in amino acid substitutions. Preferably, the derivatives include less than 15 amino acid substitutions, less than 10 amino acid substitutions, less than 5 amino acid substitutions, less than 4 amino acid substitutions, less than 3 amino acid substitutions, or less than 2 amino acid substitutions relative to the original antibody or fragment thereof. In a preferred embodiment, the derivatives have conservative amino acid substitutions made at one or more predicted non-essential amino acid residues.

[00135] In preferred embodiments, the binding domains are derived from humanized antibodies. A humanized FcγRIIB specific antibody may comprise substantially all of at least one, and typically two, variable domains in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin (*i.e.*, donor antibody) and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence.

[00136] The diabodies of present invention comprise humanized variable domains specific for FcγRIIB in which one or more regions of one or more CDRs of the heavy and/or light chain variable regions of a human antibody (the recipient antibody) have been substituted by analogous parts of one or more CDRs of a donor monoclonal antibody which specifically binds FcγRIIB, with a greater affinity than FcγRIIA, *e.g.*, a monoclonal antibody produced by clone 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2. In other

embodiments, the humanized antibodies bind to the same epitope as 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2, respectively.

[00137] In a preferred embodiment, the CDR regions of the humanized FcγRIIB binding domain are derived from a murine antibody specific for FcγRIIB. In some embodiments, the humanized antibodies described herein comprise alterations, including but not limited to amino acid deletions, insertions, modifications, of the acceptor antibody, *i.e.*, human, heavy and/or light chain variable domain framework regions that are necessary for retaining binding specificity of the donor monoclonal antibody. In some embodiments, the framework regions of the humanized antibodies described herein does not necessarily consist of the precise amino acid sequence of the framework region of a natural occurring human antibody variable region, but contains various alterations, including but not limited to amino acid deletions, insertions, modifications that alter the property of the humanized antibody, for example, improve the binding properties of a humanized antibody region that is specific for the same target as the murine FcγRIIB specific antibody. In most preferred embodiments, a minimal number of alterations are made to the framework region in order to avoid large-scale introductions of non-human framework residues and to ensure minimal immunogenicity of the humanized antibody in humans. The donor monoclonal antibody is preferably a monoclonal antibody produced by clones 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2.

[00138] In a specific embodiment, the binding domain encompasses variable domains of a CDR-grafted antibody which specifically binds FcγRIIB with a greater affinity than said antibody binds FcγRIIA, wherein the CDR-grafted antibody comprises a heavy chain variable region domain comprising framework residues of the recipient antibody and residues from the donor monoclonal antibody, which specifically binds FcγRIIB with a greater affinity than said antibody binds FcγRIIA, *e.g.*, monoclonal antibody produced from clones 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2. In another specific embodiment, the diabodies of the invention comprise variable domains from a CDR-grafted antibody which specifically binds FcγRIIB with a greater affinity than said antibody binds FcγRIIA, wherein the CDR-grafted antibody comprises a light chain variable region domain comprising framework residues of the recipient antibody and residues from the donor monoclonal antibody, which specifically binds FcγRIIB with a greater affinity than said antibody binds FcγRIIA, *e.g.*, monoclonal antibody produced from clones 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2.

[00139] The humanized anti- FcγRIIB variable domains used in the invention may have a heavy chain variable region comprising the amino acid sequence of CDR1 (SEQ ID NO:24 or SEQ ID NO:25) and/or CDR2 (SEQ ID NO:26 or SEQ ID NO:27) and/or CDR3 (SEQ ID NO:28 or SEQ ID NO:29) and/or a light chain variable region comprising the amino acid sequence of CDR1 (SEQ ID NO:32 or SEQ ID NO:33) and/or a CDR2 (SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, or SEQ ID NO:37) and/or CDR3 (SEQ ID NO:38 or SEQ ID NO:39).

[00140] In one specific embodiment, the diabody comprises variable domains from a humanized 2B6 antibody, wherein the VH region consists of the FR segments from the human germline VH segment VH1-18 (Matsuda *et al.*, 1998, J. Exp. Med. 188:2151062) and JH6 (Ravetch *et al.*, 1981, Cell 27(3 Pt. 2): 583-91), and one or more CDR regions of the 2B6 VH, having the amino acid sequence of SEQ ID NO:24, SEQ ID NO:26, or SEQ ID NO:28. In one embodiment, the 2B6 VH has the amino acid sequence of SEQ ID NO:40. In another embodiment the 2B6 VH domain has the amino acid sequence of Hu2B6VH, SEQ ID NO:87, and can be encoded by the nucleotide sequence of SEQ ID NO:88. In another specific embodiment, the diabody further comprises a VL region, which consists of the FR segments of the human germline VL segment VK-A26 (Lautner-Rieske *et al.*, 1992, Eur. J. Immunol. 22:1023-1029) and JK4 (Hieter *et al.*, 1982, J. Biol. Chem. 257:1516-22), and one or more CDR regions of 2B6VL, having the amino acid sequence of SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, and SEQ ID NO:38. In one embodiment, the 2B6 VL has the amino acid sequence of SEQ ID NO:41; SEQ ID NO:42, or SEQ ID NO:43. In a specific embodiment, the 2B6 VL has the amino acid sequence of Hu2B6VL, SEQ ID NO:89, and can be encoded by the nucleotide sequence provided in SEQ ID NO:90.

[00141] In another specific embodiment, the diabody has variable domains from a humanized 3H7 antibody, wherein the VH region consists of the FR segments from a human germline VH segment and the CDR regions of the 3H7 VH, having the amino acid sequence of SEQ ID NO: 37. In another specific embodiment, the humanized 3H7 antibody further comprises a VL regions, which consists of the FR segments of a human germline VL segment and the CDR regions of 3H7VL, having the amino acid sequence of SEQ ID NO:44.

[00142] In particular, binding domains immunospecifically bind to extracellular domains of native human FcγRIIB, and comprise (or alternatively, consist of) CDR sequences of 2B6, 3H7, 1D5, 2E1, 2H9, 2D11, or 1F2, in any of the following combinations: a VH CDR1 and a VL CDR1; a VH CDR1 and a VL CDR2; a VH CDR1

and a VL CDR3; a VH CDR2 and a VL CDR1; VH CDR2 and VL CDR2; a VH CDR2 and a VL CDR3; a VH CDR3 and a VH CDR1; a VH CDR3 and a VL CDR2; a VH CDR3 and a VL CDR3; a VH1 CDR1, a VH CDR2 and a VL CDR1; a VH CDR1, a VH CDR2 and a VL CDR2; a VH CDR1, a VH CDR2 and a VL CDR3; a VH CDR2, a VH CDR3 and a VL CDR1, a VH CDR2, a VH CDR3 and a VL CDR2; a VH CDR2, a VH CDR2 and a VL CDR3; a VH CDR1, a VL CDR1 and a VL CDR2; a VH CDR1, a VL CDR1 and a VL CDR3; a VH CDR2, a VL CDR1 and a VL CDR2; a VH CDR2, a VL CDR1 and a VL CDR3; a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR1; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR2; a VH CDR1, a VH CDR2, a VH CDR3 and a VL CDR3; a VH CDR1, a VH CDR2, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR2, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR2 and a VL CDR3; a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR2; a VH CDR1, a VH CDR2, a VH CDR3, a VL CDR1 and a VL CDR3; a VH CDR1, a VH CDR2, a VL CDR1, a VL CDR2, and a VL CDR3; a VH CDR1, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3; a VH CDR2, a VH CDR3, a VL CDR1, a VL CDR2, and a VL CDR3; or any combination thereof of the VH CDRs and VL CDRs disclosed herein.

[00143] Antibodies for deriving binding domains to be included in the diabodies of the invention may be further characterized by epitope mapping, so that antibodies may be selected that have the greatest specificity for FcγRIIB compared to FcγRIIA. Epitope mapping methods of antibodies are well known in the art and encompassed within the methods of the invention. In certain embodiments fusion proteins comprising one or more regions of FcγRIIB may be used in mapping the epitope of an antibody of the invention. In a specific embodiment, the fusion protein contains the amino acid sequence of a region of an FcγRIIB fused to the Fc portion of human IgG2. Each fusion protein may further comprise amino acid substitutions and/or replacements of certain regions of the receptor with the corresponding region from a homolog receptor, *e.g.*, FcγRIIA, as shown in Table 2 below. pMGX125 and pMGX132 contain the IgG binding site of the FcγRIIB receptor, the former with the C-terminus of FcγRIIB and the latter with the C-terminus of FcγRIIA and can be used to differentiate C-terminus binding. The others have FcγRIIA substitutions in

the IgG binding site and either the FcγIIA or FcγIIB N-terminus. These molecules can help determine the part of the receptor molecule where the antibodies bind.

Table 2. List of the fusion proteins that may be used to investigate the epitope of the monoclonal anti-FcγRIIB antibodies. Residues 172 to 180 belong to the IgG binding site of FcγRIIA and B. The specific amino acids from FcγRIIA sequence are in bold.

Plasmid	Receptor	N-terminus	172-180	SEQ ID NO:	C-terminus
pMGX125	RIIb	IIb	KKFSRSDPN	45	APS-----SS (IIb)
pMGX126	RIIa/b	IIa	QKFSRLDPN	46	APS-----SS (IIb)
pMGX127		IIa	QKFSRLDPT	47	APS-----SS (IIb)
pMGX128		IIb	KKFSRLDPT	48	APS-----SS (IIb)
pMGX129		IIa	QKFSHLDPT	49	APS-----SS (IIb)
pMGX130		IIb	KKFSHLDPT	50	APS-----SS (IIb)
pMGX131		IIa	QKFSRLDPN	51	VPSMGSSS(IIa)
pMGX132		IIb	KKFSRSDPN	52	VPSMGSSS(IIa)
pMGX133	RIIa-131R	IIa	QKFSRLDPT	53	VPSMGSSS(IIa)
pMGX134	RIIa-131H	IIa	QKFSHLDPT	54	VPSMGSSS(IIa)
pMGX135		IIb	KKFSRLDPT	55	VPSMGSSS(IIa)
pMGX136		IIb	KKFSHLDPT	56	VPSMGSSS(IIa)

[00144] The fusion proteins may be used in any biochemical assay for determination of binding to an anti-FcγRIIB antibody of the invention, *e.g.*, an ELISA. In other embodiments, further confirmation of the epitope specificity may be done by using peptides with specific residues replaced with those from the FcγRIIA sequence.

[00145] The antibodies can be characterized using assays for identifying the function of the antibodies of the invention, particularly the activity to modulate FcγRIIB signaling. For example, characterization assays of the invention can measure phosphorylation of tyrosine residues in the ITIM motif of FcγRIIB, or measure the inhibition of B cell receptor-generated calcium mobilization. The characterization assays of the invention can be cell-based or cell-free assays.

[00146] It has been well established in the art that in mast cells coaggregation of FcγRIIB with the high affinity IgE receptor, FcεRI, leads to inhibition of antigen-induced degranulation, calcium mobilization, and cytokine production (Metcalf D.D. *et al.* 1997, *Physiol. Rev.* 77:1033; Long E.O. 1999 *Annu Rev. Immunol* 17: 875). The molecular details of this signaling pathway have been recently elucidated (Ott V. L., 2002, *J. Immunol.* 162(9):4430-9). Once coaggregated with FcεRI, FcγRIIB is rapidly phosphorylated on tyrosine in its ITIM motif, and then recruits Src Homology-2 containing

inositol-5-phosphatase (SHIP), an SH2 domain-containing inositol polyphosphate 5-phosphatase, which is in turn phosphorylated and associates with Shc and p62^{dok} (p62^{dok} is the prototype of a family of adaptor molecules, which includes signaling domains such as an aminoterminal pleckstrin homology domain (PH domain), a PTB domain, and a carboxy terminal region containing PXXP motifs and numerous phosphorylation sites (Carpino *et al.*, 1997 Cell, 88: 197; Yamanshi *et al.*, 1997, Cell, 88:205).

[00147] The anti-FcγRIIB antibodies for use in the invention may likewise be characterized for ability to modulate one or more IgE mediated responses. Preferably, cells lines co-expressing the high affinity receptor for IgE and the low affinity receptor for FcγRIIB will be used in characterizing the anti-FcγRIIB antibodies in modulating IgE mediated responses. In a specific embodiment, cells from a rat basophilic leukemia cell line (RBL-H23; Barsumian E.L. *et al.* 1981 Eur. J. Immunol.11:317, which is incorporated herein by reference in its entirety) transfected with full length human FcγRIIB will be used. RBL-2H3 is a well characterized rat cell line that has been used extensively to study the signaling mechanisms following IgE-mediated cell activation. When expressed in RBL-2H3 cells and coaggregated with FcεRI, FcγRIIB inhibits FcεRI-induced calcium mobilization, degranulation, and cytokine production (Malbec *et al.*, 1998, J. Immunol. 160:1647; Daeron *et al.*, 1995 J. Clin. Invest. 95:577; Ott *et al.*, 2002 J. of Immunol. 168:4430-4439).

[00148] Antibodies for use in the invention may also be characterized for inhibition of FcεRI induced mast cell activation. For example, cells from a rat basophilic leukemia cell line (RBL-H23; Barsumian E.L. *et al.* 1981 Eur. J. Immunol.11:317) that have been transfected with FcγRIIB are sensitized with IgE and stimulated either with F(ab')₂ fragments of rabbit anti-mouse IgG, to aggregate FcεRI alone, or with whole rabbit anti-mouse IgG to coaggregate FcγRIIB and FcεRI. In this system, indirect modulation of downstream signaling molecules can be assayed upon addition of antibodies of the invention to the sensitized and stimulated cells. For example, tyrosine phosphorylation of FcγRIIB and recruitment and phosphorylation of SHIP, activation of MAP kinase family members, including but not limited to Erk1, Erk2, JNK, or p38; and tyrosine phosphorylation of p62^{dok} and its association with SHIP and RasGAP can be assayed.

[00149] One exemplary assay for determining the inhibition of FcεRI induced mast cell activation by the antibodies of the invention can comprise of the following: transfecting RBL-H23 cells with human FcγRIIB; sensitizing the RBL-H23 cells with IgE; stimulating RBL-H23 cells with either F(ab')₂ of rabbit anti-mouse IgG (to aggregate FcεRI alone and elicit FcεRI-mediated signaling, as a control), or stimulating RBL-H23 cells with whole rabbit anti-mouse IgG to (to coaggregate FcγRIIB and FcεRI, resulting in inhibition

of FcεRI-mediated signaling). Cells that have been stimulated with whole rabbit anti-mouse IgG antibodies can be further pre-incubated with the antibodies of the invention. Measuring FcεRI-dependent activity of cells that have been pre-incubated with the antibodies of the invention and cells that have not been pre-incubated with the antibodies of the invention, and comparing levels of FcεRI-dependent activity in these cells, would indicate a modulation of FcεRI-dependent activity by the antibodies of the invention.

[00150] The exemplary assay described above can be for example, used to identify antibodies that block ligand (IgG) binding to FcγRIIB receptor and antagonize FcγRIIB-mediated inhibition of FcεRI signaling by preventing coaggregating of FcγRIIB and FcεRI. This assay likewise identifies antibodies that enhance coaggregation of FcγRIIB and FcεRI and agonize FcγRIIB-mediated inhibition of FcεRI signaling by promoting coaggregating of FcγRIIB and FcεRI.

[00151] In some embodiments, the anti-FcγRIIB diabodies, comprising the epitope binding domains of anti-FcγRIIB antibodies identified described herein or known in the art, of the invention are characterized for their ability to modulate an IgE mediated response by monitoring and/or measuring degranulation of mast cells or basophils, preferably in a cell-based assay. Preferably, mast cells or basophils for use in such assays have been engineered to contain human FcγRIIB using standard recombinant methods known to one skilled in the art. In a specific embodiment the anti-FcγRIIB antibodies of the invention are characterized for their ability to modulate an IgE mediated response in a cell-based β-hexosaminidase (enzyme contained in the granules) release assay. β-hexosaminidase release from mast cells and basophils is a primary event in acute allergic and inflammatory condition (Aketani *et al.*, 2001 Immunol. Lett. 75: 185-9; Aketani *et al.*, 2000 Anal. Chem. 72: 2653-8). Release of other inflammatory mediators including but not limited to serotonin and histamine may be assayed to measure an IgE mediated response in accordance with the methods of the invention. Although not intending to be bound by a particular mechanism of action, release of granules such as those containing β-hexosaminidase from mast cells and basophils is an intracellular calcium concentration dependent process that is initiated by the cross-linking of FcγRIs with multivalent antigen.

[00152] The ability to study human mast cells has been limited by the absence of suitable long term human mast cell cultures. Recently two novel stem cell factor dependent human mast cell lines, designated LAD 1 and LAD2, were established from bone marrow aspirates from a patient with mast cell sarcoma/leukemia (Kirshenbaum *et al.*, 2003, Leukemia research, 27:677-82, which is incorporated herein by reference in its entirety.).

Both cell lines have been described to express FcεRI and several human mast cell markers. LAD 1 and 2 cells can be used for assessing the effect of the antibodies of the invention on IgE mediated responses. In a specific embodiment, cell-based β-hexosaminidase release assays such as those described supra may be used in LAD cells to determine any modulation of the IgE-mediated response by the anti-FcγRIIB antibodies of the invention. In an exemplary assay, human mast cells, *e.g.*, LAD 1, are primed with chimeric human IgE anti-nitrophenol (NP) and challenged with BSA-NP, the polyvalent antigen, and cell degranulation is monitored by measuring the β-hexosaminidase released in the supernatant (Kirshenbaum *et al.*, 2003, *Leukemia research*, 27:677-682, which is incorporated herein by reference in its entirety).

[00153] In some embodiments, if human mast cells have a low expression of endogenous FcγRIIB, as determined using standard methods known in the art, *e.g.*, FACS staining, it may be difficult to monitor and/or detect differences in the activation of the inhibitory pathway mediated by the anti-FcγRIIB diabodies of the invention. The invention thus encompasses alternative methods, whereby the FcγRIIB expression may be upregulated using cytokines and particular growth conditions. FcγRIIB has been described to be highly up-regulated in human monocyte cell lines, *e.g.*, THP1 and U937, (Tridandapani *et al.*, 2002, *J. Biol. Chem.*, 277(7): 5082-5089) and in primary human monocytes (Pricop *et al.*, 2001, *J. of Immunol.*, 166: 531-537) by IL4. Differentiation of U937 cells with dibutyryl cyclic AMP has been described to increase expression of FcγRII (Cameron *et al.*, 2002 *Immunology Letters* 83, 171-179). Thus the endogenous FcγRIIB expression in human mast cells for use in the methods of the invention may be up-regulated using cytokines, *e.g.*, IL-4, IL-13, in order to enhance sensitivity of detection.

[00154] The anti-FcγRIIB diabodies can also be assayed for inhibition of B-cell receptor (BCR)-mediated signaling. BCR-mediated signaling can include at least one or more down stream biological responses, such as activation and proliferation of B cells, antibody production, etc. Coaggregation of FcγRIIB and BCR leads to inhibition of cell cycle progression and cellular survival. Further, coaggregation of FcγRIIB and BCR leads to inhibition of BCR-mediated signaling.

[00155] Specifically, BCR-mediated signaling comprises at least one or more of the following: modulation of down stream signaling molecules (*e.g.*, phosphorylation state of FcγRIIB, SHIP recruitment, localization of Btk and/or PLCγ, MAP kinase activity, recruitment of Akt (anti-apoptotic signal), calcium mobilization, cell cycle progression, and cell proliferation.

[00156] Although numerous effector functions of FcγRIIB-mediated inhibition of BCR signaling are mediated through SHIP, recently it has been demonstrated that lipopolysaccharide (LPS)-activated B cells from SHIP deficient mice exhibit significant FcγRIIB-mediated inhibition of calcium mobilization, Ins(1,4,5)P₃ production, and Erk and Akt phosphorylation (Brauweiler A. *et al.*, 2001, Journal of Immunology, 167(1): 204-211). Accordingly, ex vivo B cells from SHIP deficient mice can be used to characterize the antibodies of the invention. One exemplary assay for determining FcγRIIB-mediated inhibition of BCR signaling by the antibodies of the invention can comprise the following: isolating splenic B cells from SHIP deficient mice, activating said cells with lipopolysaccharide, and stimulating said cells with either F(ab')₂ anti-IgM to aggregate BCR or with anti-IgM to coaggregate BCR with FcγRIIB. Cells that have been stimulated with intact anti-IgM to coaggregate BCR with FcγRIIB can be further pre-incubated with the antibodies of the invention. FcγRIIB-dependent activity of cells can be measured by standard techniques known in the art. Comparing the level of FcγRIIB-dependent activity in cells that have been pre-incubated with the antibodies and cells that have not been pre-incubated, and comparing the levels would indicate a modulation of FcγRIIB-dependent activity by the antibodies.

[00157] Measuring FcγRIIB-dependent activity can include, for example, measuring intracellular calcium mobilization by flow cytometry, measuring phosphorylation of Akt and/or Erk, measuring BCR-mediated accumulation of PI(3,4,5)P₃, or measuring FcγRIIB-mediated proliferation B cells.

[00158] The assays can be used, for example, to identify diabodies or anti-FcγRIIB antibodies for use in the invention that modulate FcγRIIB-mediated inhibition of BCR signaling by blocking the ligand (IgG) binding site to FcγRIIB receptor and antagonizing FcγRIIB-mediated inhibition of BCR signaling by preventing coaggregation of FcγRIIB and BCR. The assays can also be used to identify antibodies that enhance coaggregation of FcγRIIB and BCR and agonize FcγRIIB-mediated inhibition of BCR signaling.

[00159] The anti-FcγRIIB antibodies can also be assayed for FcγRII-mediated signaling in human monocytes/macrophages. Coaggregation of FcγRIIB with a receptor bearing the immunoreceptor tyrosine-based activation motif (ITAM) acts to down-regulate FcγR-mediated phagocytosis using SHIP as its effector (Tridandapani *et al.* 2002, J. Biol. Chem. 277(7):5082-9). Coaggregation of FcγRIIA with FcγRIIB results in rapid phosphorylation of the tyrosine residue on FcγRIIB's ITIM motif, leading to an enhancement in phosphorylation of SHIP, association of SHIP with Shc, and phosphorylation of proteins having the molecular weight of 120 and 60-65 kDa. In

addition, coaggregation of FcγRIIA with FcγRIIB results in down-regulation of phosphorylation of Akt, which is a serine-threonine kinase that is involved in cellular regulation and serves to suppress apoptosis.

[00160] The anti-FcγRIIB diabodies can also be assayed for inhibition of FcγR-mediated phagocytosis in human monocytes/macrophages. For example, cells from a human monocytic cell line, THP-1 can be stimulated either with Fab fragments of mouse monoclonal antibody IV.3 against FcγRII and goat anti-mouse antibody (to aggregate FcγRIIA alone), or with whole IV.3 mouse monoclonal antibody and goat anti-mouse antibody (to coaggregate FcγRIIA and FcγRIIB). In this system, modulation of downstream signaling molecules, such as tyrosine phosphorylation of FcγRIIB, phosphorylation of SHIP, association of SHIP with Shc, phosphorylation of Akt, and phosphorylation of proteins having the molecular weight of 120 and 60-65 kDa can be assayed upon addition of molecules of the invention to the stimulated cells. In addition, FcγRIIB-dependent phagocytic efficiency of the monocyte cell line can be directly measured in the presence and absence of the antibodies of the invention.

[00161] Another exemplary assay for determining inhibition of FcγR-mediated phagocytosis in human monocytes/macrophages by the antibodies of the invention can comprise the following: stimulating THP-1 cells with either Fab of IV.3 mouse anti-FcγRII antibody and goat anti-mouse antibody (to aggregate FcγRIIA alone and elicit FcγRIIA-mediated signaling); or with mouse anti-FcγRII antibody and goat anti-mouse antibody (to coaggregate FcγRIIA and FcγRIIB and inhibiting FcγRIIA-mediated signaling. Cells that have been stimulated with mouse anti-FcγRII antibody and goat anti-mouse antibody can be further pre-incubated with the molecules of the invention. Measuring FcγRIIA-dependent activity of stimulated cells that have been pre-incubated with molecules of the invention and cells that have not been pre-incubated with the antibodies of the invention and comparing levels of FcγRIIA-dependent activity in these cells would indicate a modulation of FcγRIIA-dependent activity by the antibodies of the invention.

[00162] The exemplary assay described can be used for example, to identify binding domains that block ligand binding of FcγRIIB receptor and antagonize FcγRIIB-mediated inhibition of FcγRIIA signaling by preventing coaggregation of FcγRIIB and FcγRIIA. This assay likewise identifies binding domains that enhance coaggregation of FcγRIIB and FcγRIIA and agonize FcγRIIB-mediated inhibition of FcγRIIA signaling.

[00163] The FcγRIIB binding domains of interest can be assayed while comprised I antibodies by measuring the ability of THP-1 cells to phagocytose fluoresceinated IgG-opsonized sheep red blood cells (SRBC) by methods previously described (Tridandapani *et*

al., 2000, J. Biol. Chem. 275: 20480-7). For example, an exemplary assay for measuring phagocytosis comprises of: treating THP-1 cells with the antibodies of the invention or with a control antibody that does not bind to FcγRII, comparing the activity levels of said cells, wherein a difference in the activities of the cells (*e.g.*, rosetting activity (the number of THP-1 cells binding IgG-coated SRBC), adherence activity (the total number of SRBC bound to THP-1 cells), and phagocytic rate) would indicate a modulation of FcγRIIA-dependent activity by the antibodies of the invention. This assay can be used to identify, for example, antibodies that block ligand binding of FcγRIIB receptor and antagonize FcγRIIB-mediated inhibition of phagocytosis. This assay can also identify antibodies that enhance FcγRIIB-mediated inhibition of FcγRIIA signaling.

[00164] In a preferred embodiment, the binding domains modulate FcγRIIB-dependent activity in human monocytes/macrophages in at least one or more of the following ways: modulation of downstream signaling molecules (*e.g.*, modulation of phosphorylation state of FcγRIIB, modulation of SHIP phosphorylation, modulation of SHIP and Shc association, modulation of phosphorylation of Akt, modulation of phosphorylation of additional proteins around 120 and 60-65 kDa) and modulation of phagocytosis.

5.1.2 CD16A BINDING DOMAINS

[00165] The following section discusses CD16A binding proteins which can be used as sources for light and heavy chain variable regions for covalent diabody production. In the present invention CD16A binding proteins includes molecules comprising VL and VH domains of anti-CD16A antibodies, which VH and VL domains are used in the production of the diabodies of the present invention.

[00166] A variety of CD16A binding proteins may be used in connection with the present invention. Suitable CD16A binding proteins include human or humanized monoclonal antibodies as well as CD16A binding antibody fragments (*e.g.*, scFv or single chain antibodies, Fab fragments, minibodies) and another antibody-like proteins that bind to CD16A via an interaction with a light chain variable region domain, a heavy chain variable region domain, or both.

[00167] In some embodiments, the CD16A binding protein for use according to the invention comprises a VL and/or VH domain that has one or more CDRs with sequences derived from a non-human anti-CD16A antibody, such as a mouse anti-CD16A antibody, and one or more framework regions with derived from framework sequences of one or more human immunoglobulins. A number of non-human anti-CD16A monoclonal antibodies,

from which CDR and other sequences may be obtained, are known (see, e.g., Tamm and Schmidt, 1996, *J. Imm.* 157:1576-81; Fleit et al., 1989, p.159; LEUKOCYTE TYPING II: HUMAN MYELOID AND HEMATOPOIETIC CELLS, Reinherz et al., eds. New York: Springer-Verlag; 1986; LEUCOCYTE TYPING III: WHITE CELL DIFFERENTIATION ANTIGENS McMichael A J, ed., Oxford: Oxford University Press, 1986); LEUKOCYTE TYPING IV: WHITE CELL DIFFERENTIATION ANTIGENS, Kapp et al., eds. Oxford Univ. Press, Oxford; LEUKOCYTE TYPING V: WHITE CELL DIFFERENTIATION ANTIGENS, Schlossman et al., eds. Oxford Univ. Press, Oxford; LEUKOCYTE TYPING VI: WHITE CELL DIFFERENTIATION ANTIGENS, Kishimoto, ed. Taylor & Francis. In addition, as shown in the Examples, new CD16A binding proteins that recognize human CD16A expressed on cells can be obtained using well known methods for production and selection of monoclonal antibodies or related binding proteins (e.g., hybridoma technology, phage display, and the like). See, for example, O'Connel et al., 2002, *J. Mol. Biol.* 321:49-56; Hoogenboom and Chames, 2000, *Imm. Today* 21:371078; Krebs et al., 2001, *J. Imm. Methods* 254:67-84; and other references cited herein. Monoclonal antibodies from a non-human species can be chimerized or humanized using techniques using techniques of antibody humanization known in the art.

[00168] Alternatively, fully human antibodies against CD16A can be produced using transgenic animals having elements of a human immune system (see, e.g., U.S. Pat. Nos. 5,569,825 and 5,545,806), using human peripheral blood cells (Casali et al., 1986, *Science* 234:476), by screening a DNA library from human B cells according to the general protocol outlined by Huse et al., 1989, *Science* 246:1275, and by other methods.

[00169] In a preferred embodiment, the binding donor is from the 3G8 antibody or a humanized version thereof, e.g., such as those disclosed in U.S. patent application publication 2004/0010124, which is incorporated by reference herein in its entirety. It is contemplated that, for some purposes, it may be advantageous to use CD16A binding proteins that bind the CD16A receptor at the same epitope bound by 3G8, or at least sufficiently close to this epitope to block binding by 3G8. Methods for epitope mapping and competitive binding experiments to identify binding proteins with the desired binding properties are well known to those skilled in the art of experimental immunology. See, for example, Harlow and Lane, cited supra; Stahl et al., 1983, *Methods in Enzymology* 9:242-53; Kirkland et al., 1986, *J. Immunol.* 137:3614-19; Morel et al., 1988, *Molec. Immunol.* 25:7-15; Cheung et al., 1990, *Virology* 176:546-52; and Moldenhauer et al., 1990, *Scand. J. Immunol.* 32:77-82. For instance, it is possible to determine if two antibodies bind to the same site by using one of the antibodies to capture the antigen on an ELISA plate and then

measuring the ability of the second antibody to bind to the captured antigen. Epitope comparison can also be achieved by labeling a first antibody, directly or indirectly, with an enzyme, radionuclide or fluorophore, and measuring the ability of an unlabeled second antibody to inhibit the binding of the first antibody to the antigen on cells, in solution, or on a solid phase.

[00170] It is also possible to measure the ability of antibodies to block the binding of the CD16A receptor to immune complexes formed on ELISA plates. Such immune complexes are formed by first coating the plate with an antigen such as fluorescein, then applying a specific anti-fluorescein antibody to the plate. This immune complex then serves as the ligand for soluble Fc receptors such as sFcRIIIa. Alternatively a soluble immune complex may be formed and labeled, directly or indirectly, with an enzyme radionuclide or fluorophore. The ability of antibodies to inhibit the binding of these labeled immune complexes to Fc receptors on cells, in solution or on a solid phase can then be measured.

[00171] CD16A binding proteins of the invention may or may not comprise a human immunoglobulin Fc region. Fc regions are not present, for example, in scFv binding proteins. Fc regions are present, for example, in human or humanized tetrameric monoclonal IgG antibodies. As described *supra*, in some embodiments of the present invention, the CD16A binding protein includes an Fc region that has an altered effector function, e.g., reduced affinity for an effector ligand such as an Fc receptor or C1 component of complement compared to the unaltered Fc region (e.g., Fc of naturally occurring IgG1, proteins). In one embodiment the Fc region is not glycosylated at the Fc region amino acid corresponding to position 297. Such antibodies lack Fc effector function.

[00172] Thus, the CD16A binding protein may not exhibit Fc-mediated binding to an effector ligand such as an Fc receptor or the C1 component of complement due to the absence of the Fc domain in the binding protein while, in other cases, the lack of binding or effector function is due to an alteration in the constant region of the antibody.

5.1.2.1 CD16A Binding Proteins Comprising CDR Sequences Similar to a mAb 3G8 CDR Sequences.

[00173] CD16A binding proteins that can be used in the practice of the invention include proteins comprising a CDR sequence derived from (i.e., having a sequence the same as or similar to) the CDRs of the mouse monoclonal antibody 3G8. Complementary cDNAs encoding the heavy chain and light chain variable regions of the mouse 3G8 monoclonal antibody, including the CDR encoding sequences, were cloned and sequenced as described. The nucleic acid and protein sequences of 3G8 are provided below. Using the mouse variable region and CDR sequences, a large number of chimeric and humanized

monoclonal antibodies, comprising complementary determining regions derived from 3G8 CDRs were produced and their properties analyzed. To identify humanized antibodies that bind CD16A with high affinity and have other desirable properties, antibody heavy chains comprising a VH region with CDRs derived from 3G8 were produced and combined (by coexpression) with antibody light chains comprising a VL region with CDRs derived from 3G8 to produce a tetrameric antibody for analysis. Properties of the resulting tetrameric antibodies were determined as described below. As described below, CD16A binding proteins comprising 3G8 CDRs, such as the humanized antibody proteins described herein, may be used according to the invention.

5.1.2.1.1 VH Region

[00174] In one aspect, the CD16A binding protein of the invention may comprise a heavy chain variable domain in which at least one CDR (and usually three CDRS) have the sequence of a CDR (and more typically all three CDRS) of the mouse monoclonal antibody 3G8 heavy chain and for which the remaining portions of the binding protein are substantially human (derived from and substantially similar to, the heavy chain variable region of a human antibody or antibodies).

[00175] In an aspect, the invention provides a humanized 3G8 antibody or antibody fragment containing CDRs derived from the 3G8 antibody in a substantially human framework, but in which at least one of the CDRs of the heavy chain variable domain differs in sequence from the corresponding mouse antibody 3G8 heavy chain CDR. For example, in one embodiment, the CDR(S) differs from the 3G8 CDR sequence at least by having one or more CDR substitutions shown known in the art to affect binding of 3G8 to CD16A, as known in the art or as disclosed in Tables 3 and 4A-H. Suitable CD16 binding proteins may comprise 0, 1, 2, 3, or 4, or more of these substitutions (and often have from 1 to 4 of these substitutions) and optionally can have additional substitutions as well.

Table 3. V_H Domain Substitutions

No.	Kabat Position	Region	Substitutions
1	2	FR1	Ile
2	5	FR1	Lys
3	10	FR1	Thr
4	30	FR1	Arg
5	34	CDR1	Val
6	50	CDR2	Leu
7	52	CDR2	Phe or Tyr or Asp

8	54	CDR2	Asn
9	60	CDR2	Ser
10	62	CDR2	Ser
11	70	FR3	Thr
12	94	FR3	Gln or Lys or Ala or His
13	99	CDR3	Tyr
14	101	CDR3	Asp

Table 4A. V_H Sequences Derived from 3G8 V_H *

	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
3G8VH	A	A	A	A	A	A	A
Ch3G8VH	A	A	A	A	A	A	B
HxC	B	A	B	A	A	A	B
CxH	A	A	A	A	B	A	B
Hu3G8VH-1	B	A	B	A	B	A	B
Hu3G8VH-2	C	A	B	A	B	A	B
Hu3G8VH-3	D	A	B	A	B	A	B
Hu3G8VH-4	B	A	B	A	C	B	B
Hu3G8VH-5	B	A	B	A	C	A	B
Hu3G8VH-6	B	B	B	A	B	B	B
Hu3G8VH-7	B	B	B	A	B	A	B
Hu3G8VH-8	B	A	B	A	B	C	B
Hu3G8VH-9	B	A	B	B	B	B	B
Hu3G8VH-10	B	A	B	A	B	B	B
Hu3G8VH-11	B	A	B	B	B	A	B
Hu3G8VH-12	B	A	B	C	B	A	B
Hu3G8VH-13	B	A	B	D	B	A	B
Hu3G8VH-14	B	A	B	E	B	A	B
Hu3G8VH-15	B	A	B	A	D	A	B
Hu3G8VH-16	B	A	B	A	E	A	B
Hu3G8VH-17	B	A	B	A	F	A	B
Hu3G8VH-18	B	A	B	A	G	A	B
Hu3G8VH-19	B	A	B	A	C	C	B
Hu3G8VH-20	B	B	B	C	B	A	B
Hu3G8VH-21	B	A	B	A	D	B	B
Hu3G8VH-22	B	B	B	C	B	C	B
Hu3G8VH-23	B	B	B	C	E	C	B
Hu3G8VH-24	B	B	B	C	F	C	B
Hu3G8VH-25	B	B	B	C	G	C	B
Hu3G8VH-26	B	B	B	C	C	C	B
Hu3G8VH-27	B	B	B	C	E	D	B
Hu3G8VH-28	B	B	B	C	F	D	B
Hu3G8VH-29	B	B	B	C	G	D	B
Hu3G8VH-30	B	B	B	C	C	D	B
Hu3G8VH-31	E	B	B	C	B	A	B

Hu3G8VH-32	E	B	B	H	B	A	B
Hu3G8VH-33	E	B	B	H	B	A	B
Hu3G8VH-34	E	B	B	C	B	C	B
Hu3G8VH-35	E	B	B	C	C	C	B
Hu3G8VH-36	E	B	B	H	C	D	B
Hu3G8VH-37	E	B	B	H	E	C	B
Hu3G8VH-38	E	B	B	F	B	A	B
Hu3G8VH-39	E	B	B	I	B	A	B
Hu3G8VH-40	E	B	B	G	B	A	B
Hu3G8VH-41	E	B	B	J	B	A	B
Hu3G8VH-42	E	B	B	C	H	A	B
Hu3G8VH-43	E	B	B	C	H	C	B
Hu3G8VH-44	E	B	B	C	I	D	B
Hu3G8VH-45	E	B	B	C	J	D	B

*Letters in Table 4A refer to sequences in Tables 4 B-H.

TABLE 4B
FR1

A	B	C	D	E		RESIDUE
Q	Q	Q	Q	Q		1
V	V	V	V	I		2
T	T	T	T	T		3
L	L	L	L	L		4
K	R	K	R	K		5
E	E	E	E	E		6
S	S	S	S	S		7
G	G	G	G	G		8
P	P	P	P	P		9
G	A	A	A	T		10
I	L	L	L	L		11
L	V	V	V	V		12
Q	K	K	K	K		13
P	P	P	P	P		14
S	T	T	T	T		15
Q	Q	Q	Q	Q		16
T	T	T	T	T		17
L	L	L	L	L		18
S	T	T	T	T		19
L	L	L	L	L		20
T	T	T	T	T		21
C	C	C	C	C		22
S	T	T	T	T		23
F	F	F	F	F		24
S	S	S	S	S		25
G	G	G	G	G		26
F	F	F	F	F		27
S	S	S	S	S		28

L	L	L	L	L		29
R	S	S	R	S		30
30	31	32	33	34		Se ID No

TABLE 4C
CDR1

A	B		RESIDUE
T	T		31
S	S		32
G	G		33
M	V		34
G	G		35
V	V		35A
G	G		35B
35	36		Seq ID No

TABLE 4D
FR2

A	B		RESIDUE
W	W		36
I	I		37
R	R		38
Q	Q		39
P	P		40
S	P		41
G	G		42
K	K		43
G	A		44
L	L		45
E	E		46
W	W		47
L	L		48
A	A		49
37	38		Seq ID No.

TABLE 4E
CDR2

A	B	C	D	E	F	G	H	I	J		RESIDUE
H	H	H	H	H	L	H	L	H	L		50
I	I	I	I	I	I	I	I	I	I		51
W	Y	W	Y	W	D	F	W	D	W		52
W	W	W	W	W	W	W	W	W	W		53
D	N	D	D	N	D	D	D	D	N		54
D	D	D	D	D	D	D	D	D	D		55
D	D	D	D	D	D	D	D	D	D		56

K	K	K	K	K	K	K	K	K	K		57
R	R	R	R	R	R	R	R	R	R		58
Y	Y	Y	Y	Y	Y	Y	Y	Y	Y		59
N	N	S	N	N	S	S	S	S	S		60
P	P	P	P	P	P	P	P	P	P		61
A	A	S	A	A	S	S	S	S	S		62
L	L	L	L	L	L	L	L	L	L		63
K	K	K	K	K	K	K	K	K	K		64
S	S	S	S	S	S	S	S	S	S		65
39	40	41	42	43	44	45	46	47	48		Seq ID No

TABLE 4F
FR3

A	B	C	D	E	F	G	H	I	J		RESIDUE
R	R	R	R	R	R	R	R	R	R		66
L	L	L	L	L	L	L	L	L	L		67
T	T	T	T	T	T	T	T	T	T		68
I	I	I	I	I	I	I	I	I	I		69
S	S	S	S	S	S	S	T	T	T		70
K	K	K	K	K	K	K	K	K	K		71
D	D	D	D	D	D	D	D	D	D		72
T	T	T	T	T	T	T	T	T	T		73
S	S	S	S	S	S	S	S	S	S		74
S	K	K	K	K	K	K	K	K	K		75
N	N	N	N	N	N	N	N	N	N		76.
Q	Q	Q	Q	Q	Q	Q	Q	Q	Q		77
V	V	V	V	V	V	V	V	V	V		78
F	V	V	V	V	V	V	V	V	V		79
L	L	L	L	L	L	L	L	L	L		80
K	T	T	T	T	T	T	T	T	T		81
I	M	M	M	M	M	M	M	M	M		82
A	T	T	T	T	T	T	T	T	T		82A
S	N	N	N	N	N	N	N	N	N		82B
V	M	M	M	M	M	M	M	M	M		82C
D	D	D	D	D	D	D	D	D	D		83
T	P	P	P	P	P	P	P	P	P		84
A	V	V	V	V	V	V	V	V	V		85
D	D	D	D	D	D	D	D	D	D		86
T	T	T	T	T	T	T	T	T	T		87
A	A	A	A	A	A	A	A	A	A		88
T	T	T	T	T	T	T	T	T	T		89
Y	Y	Y	Y	Y	Y	Y	Y	Y	Y		90
Y	Y	Y	Y	Y	Y	Y	Y	Y	Y		91
C	C	C	C	C	C	C	C	C	C		92
A	A	A	A	A	A	A	A	A	A		93
Q	R	Q	T	K	A	H	R	H	Q		94
49	50	51	52	53	54	55	56	57	58		Seq ID No

TABLE 4G
CDR3

A	B	C	D		RESIDUE
I	I	I	I		95
N	N	N	N		96
P	P	P	P		97
A	A	A	A		98
W	W	Y	Y		99
F	F	F	F		100
A	D	A	D		101
Y	Y	Y	Y		102
59	60	61	62		Seq ID No

TABLE 4H
FR4

A	B		RESIDUE
W	W		103
G	G		104
Q	Q		105
G	G		106
T	T		107
L	L		108
V	V		109
T	T		110
V	V		111
S	S		112
A	S		113
63	64		Seq ID No

[00176] In one embodiment, a CD16A binding protein may comprise a heavy chain variable domain sequence that is the same as, or similar to, the VH domain of the Hu3G8VH-1 construct, the sequence of which is provided in SEQ ID NO:70. For example, the invention provides a CD16A binding protein comprising a VH domain with a sequence that (1) differs from the VH domain of Hu3G8VH-1 (SEQ ID NO:70) by zero, one, or more than one of the CDR substitutions set forth in Table 1; (2) differs from the VH domain of Hu3G8VH-1 by zero, one or more than one of the framework substitutions set forth in Table 1; and (3) is at least about 80% identical, often at least about 90%, and sometimes at least about 95% identical, or even at least about 98% identical to the Hu3G8VH-1 VH sequence at the remaining positions.

[00177] Exemplary VH domains of CD16 binding proteins of the invention have the sequence of 3G8VH, Hu3G8VH-5 and Hu3G8VH-22 (SEQ ID NO:81, SEQ ID NO:71 and SEQ ID NO:72, respectively). Exemplary nucleotide sequences encoding the sequences of 3G8VH and Hu3G8VH-5 (SEQ ID NO:81 and SEQ ID NO:71, respectively) are provided by SEQ ID NO:82 and SEQ ID NO:83, respectively.

[00178] The VH domain may have a sequence that differs from that of Hu3G8VH-1 (SEQ ID NO:70) by at least one, at least two, at least three, at least four, at least five, or at least six of the substitutions shown in Table 3. These substitutions are believed to result in increased affinity for CD16A and/or reduce the immunogenicity of a CD16A binding protein when administered to humans. In certain embodiments, the degree of sequence identity with the Hu3G8VH-1 VH domain at the remaining positions is at least about 80%, at least about 90%, at least about 95% or at least about 98%.

[00179] For illustration and not limitation, the sequences of a number of CD16A binding protein VH domains is shown in Table 4. Heavy chains comprising these sequences fused to a human C γ 1 constant region were coexpressed with the hu3G8VL-1 light chain (described below) to form tetrameric antibodies, and binding of the antibodies to CD16A was measured to assess the effect of amino acid substitutions compared to the hu3G8VH-1 VH domain. Constructs in which the VH domain has a sequence of hu3G8VH-1, 2, 3, 4, 5, 8, 12, 14, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44 and 45 showed high affinity binding, with hu3G8VH-6 and -40 VH domains showing intermediate binding. CD16A binding proteins comprising the VH domains of hu3G8VH-5 and hu3G8VH-22 (SEQ ID NO:71 and SEQ ID NO:72, respectively) are considered to have particularly favorable binding properties.

5.1.2.2 VL Region

[00180] Similar studies were conducted to identify light chain variable domain sequences with favorable binding properties. In one aspect, the invention provides a CD16A binding protein containing a light chain variable domain in which at least one CDR (and usually three CDRs) has the sequence of a CDR (and more typically all three CDRs) of the mouse monoclonal antibody 3G8 light chain and for which the remaining portions of the binding protein are substantially human (derived from and substantially similar to, the heavy chain variable region of a human antibody or antibodies).

[00181] In one aspect, the invention provides a fragment of a humanized 3G8 antibody containing CDRs derived from the 3G8 antibody in a substantially human framework, but in which at least one of the CDRs of the light chain variable domain differs

in sequence from the mouse monoclonal antibody 3G8 light chain CDR. In one embodiment, the CDR(s) differs from the 3G8 sequence at least by having one or more amino acid substitutions in a CDR, such as, one or more substitutions shown in Table 2 (e.g., arginine at position 24 in CDR1; serine at position 25 in CDR1; tyrosine at position 32 in CDR1; leucine at position 33 in CDR1; aspartic acid, tryptophan or serine at position 50 in CDR2; serine at position 53 in CDR2; alanine or glutamine at position 55 in CDR2; threonine at position 56 in CDR2; serine at position 93 in CDR3; and/or threonine at position 94 in CDR3). In various embodiments, the variable domain can have 0, 1, 2, 3, 4, 5, or more of these substitutions (and often have from 1 to 4 of these substitutions) and optionally, can have additional substitutions as well.

[00182] In one embodiment, a suitable CD16A binding protein may comprise a light chain variable domain sequence that is the same as, or similar to, the VL domain of the Hu3G8VL-1 (SEQ ID NO:73) construct, the sequence of which is provided in Table 6. For example, the invention provides a CD16A binding protein comprising a VL domain with a sequence that (1) differs from the VL domain of Hu3G8VL-1 (SEQ ID NO:73) by zero, one, or more of the CDR substitutions set forth in Table 5; (2) differs from the VL domain of Hu3G8VL-1 by zero, one or more of the framework substitutions set forth in Table 5; and (3) is at least about 80% identical, often at least about 90%, and sometimes at least about 95% identical, or even at least about 98% identical to the Hu3G8VL-1 VL sequence (SEQ ID NO:73) at the remaining positions.

Table 5. 3G8 V_L Domain Substitutions

No.	Kabat Position	Region	Substitutions
1	24	CDR1	Arg
2	25	CDR1	Ser
3	32	CDR1	Tyr
4	33	CDR1	Leu
5	50	CDR2	Asp or Trp or Ser
6	51	CDR2	Ala
7	53	CDR2	Ser
8	55	CDR2	Ala or Gln
9	56	CDR2	Thr
10	93	CDR3	Ser

11	94	CDR3	Thr
----	----	------	-----

Table 6. V_L Sequences Derived from 3G8 V_L*

	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
3G8VL	A	A	A	A	A	A	A
Ch3G8VL	A	A	A	A	A	A	A
Hu3G8VL-1	B	A	A	A	B	A	B
Hu3G8VL-2	B	B	A	A	B	A	B
Hu3G8VL-3	B	C	A	A	B	A	B
Hu3G8VL-4	B	D	A	A	B	A	B
Hu3G8VL-5	B	E	A	A	B	A	B
Hu3G8VL-6	B	F	A	A	B	A	B
Hu3G8VL-7	B	G	A	A	B	A	B
Hu3G8VL-8	B	A	A	B	B	A	B
Hu3G8VL-9	B	A	A	C	B	A	B
Hu3G8VL-10	B	A	A	D	B	A	B
Hu3G8VL-11	B	A	A	E	B	A	B
Hu3G8VL-12	B	A	A	F	B	A	B
Hu3G8VL-13	B	A	A	G	B	A	B
Hu3G8VL-14	B	A	A	A	B	B	B
Hu3G8VL-15	B	A	A	A	B	C	B
Hu3G8VL-16	B	A	A	A	B	D	B
Hu3G8VL-17	B	A	A	A	B	E	B
Hu3G8VL-18	B	B	A	D	B	A	B
Hu3G8VL-19	B	B	A	D	B	D	B
Hu3G8VL-20	B	B	A	D	B	E	B
Hu3G8VL-21	B	C	A	D	B	A	B
Hu3G8VL-22	B	C	A	D	B	D	B

Hu3G8VL-23	B	C	A	D	B	E	B
Hu3G8VL-24	B	D	A	D	B	A	B
Hu3G8VL-25	B	D	A	D	B	D	B
Hu3G8VL-26	B	D	A	D	B	E	B
Hu3G8VL-27	B	E	A	D	B	A	B
Hu3G8VL-28	B	E	A	D	B	D	B
Hu3G8VL-29	B	E	A	D	B	E	B
Hu3G8VL-30	B	A	A	D	B	D	B
Hu3G8VL-31	B	A	A	D	B	E	B
Hu3G8VL-32	B	A	A	H	B	A	B
Hu3G8VL-33	B	A	A	I	B	A	B
Hu3G8VL-34	B	A	A	J	B	A	B
Hu3G8VL-35	B	B	A	H	B	D	B
Hu3G8VL-36	B	C	A	H	B	D	B
Hu3G8VL-37	B	E	A	H	B	D	B
Hu3G8VL-38	B	B	A	I	B	D	B
Hu3G8VL-39	B	C	A	I	B	D	B
Hu3G8VL-40	B	E	A	I	B	D	B
Hu3G8VL-41	B	B	A	J	B	D	B
Hu3G8VL-42	B	C	A	J	B	D	B
Hu3G8VL-43	B	E	A	J	B	D	B
Hu3G8VL-44	B	A	A	K	B	A	B

*Letters in Table 6A refer to sequences in Tables 6B-H.

TABLE 6B
FR1

A	B		RESIDUE

D	D		1
T	I		2
V	V		3
L	M		4
T	T		5
Q	Q		6
S	S		7
P	P		8
A	D		9
S	S		10
L	L		11
A	A		12
V	V		13
S	S		14
L	L		15
G	G		16
Q	E		17
R	R		18
A	A		19
T	T		20
I	I		21
S	N		22
C	C		23
65	66		Seq ID No

TABLE 6C
CDR1

A	B	C	D	E	F	G	RESIDUE
K	R	K	K	K	K	K	24
A	A	S	A	A	A	A	25
S	S	S	S	S	S	S	26
Q	Q	Q	Q	Q	Q	Q	27
S	S	S	S	S	S	S	27A
V	V	V	V	V	V	V	27B
D	D	D	D	D	D	D	27C
F	F	F	F	F	F	F	27D
D	D	D	D	D	D	D	28
G	G	G	G	G	G	G	29
D	D	D	D	D	D	D	30
S	S	S	S	S	S	S	31
F	F	F	Y	F	F	Y	32
M	M	M	M	L	M	L	33
N	N	N	N	N	A	A	34
67	68	69	70	71	72	73	Seq ID No

TABLE 6D
FR2

A		RESIDUE
W		35
Y		36
Q		37
Q		38
K		39
P		40
G		41
Q		42
P		43
P		44
K		45
L		46
L		47
I		48
Y		49
74		Seq ID No

TABLE 6E
CDR2

A	B	C	D	E	F	G	H	I	J	K		RESIDUE
T	D	W	T	D	D	S	S	S	T	T		50
T	A	A	T	A	A	A	T	T	T	T		51
S	S	S	S	S	S	S	S	S	S	S		52
N	N	N	N	N	N	N	N	N	N	S		53
L	L	L	L	L	L	L	L	L	L	L		54
E	E	E	E	E	A	Q	E	Q	Q	Q		55
S	S	S	T	T	T	S	S	S	S	S		56
75	76	77	78	79	80	81	82	83	84	85		Seq ID No

TABLE 6F
FR3

A	B		RESIDUE
G	G		57
I	V		58
P	P		59
A	D		60
R	R		61
F	F		62
S	S		63
A	G		64
S	S		65
G	G		66
S	S		67

G	G		68
T	T		69
D	D		70
F	F		71
T	T		72
L	L		73
N	T		74
I	I		75
H	S		76
P	S		77
V	L		78
E	Q		79
E	A		80
E	E		81
D	D		82
T	V		83
A	A		84
T	V		85
Y	Y		86
Y	Y		87
C	C		88
86	87		Seq ID No

TABLE 6G
CDR3

A	B	C	D	E		RESIDUE
Q	Q	Q	Q	Q		89
Q	Q	Q	Q	Q		90
S	S	S	S	S		91
N	Y	Y	N	N		92
E	S	E	S	E		93
D	T	D	D	T		94
P	P	P	P	P		95
Y	Y	Y	Y	Y		96
T	T	T	T	T		97
88	89	90	91	92		Seq ID No

TABLE 6H
FR4

A	B		RESIDUE
F	F		98
G	G		99
G	Q		100
G	G		101
T	T		102
K	K		103
L	L		104
E	E		105
I	I		106
K	K		107
93	94		Seq ID No

[00183] Exemplary VL domains of CD16 binding proteins of the invention have the sequence of 3G8VL, Hu3G8VL-1 or Hu3G8VL-43, (SEQ ID NO:84, SEQ ID NO:73 and SEQ ID NO:74, respectively) as shown in Tables 5 and 6. Exemplary nucleotide sequences encoding 3G8VL (SEQ ID NO:84) and Hu3G8VL-1 (SEQ ID NO:73) are provided in SEQ ID NO:85 and SEQ ID NO:86, respectively.

[00184] The VL domain may have a sequence that differs from that of Hu3G8VL-1 (SEQ ID NO:73) by zero, one, at least two, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, or at least 9 of the substitutions shown in Table 2. These substitutions are believed to result in increased affinity for CD16A and/or reduce the immunogenicity of a CD16A binding protein when administered to humans. In certain embodiments, the degree of sequence identity at the remaining positions is at least about 80%, at least about 90% at least about 95% or at least about 98%.

[00185] For illustration and not limitation, the sequences of a number of CD16A binding proteins VL domains is shown in Table 6. Light chains comprising these sequences fused to a human Cκ constant domain were coexpressed with a Hu3G8VH heavy chain (described above) to form tetrameric antibodies, and the binding of the antibodies to CD16A was measured to assess the effect of amino acid substitutions compared to the Hu3G8VL-1 VL domain (SEQ ID NO:73). Constructs in which the VL domain has a sequence of hu3G8VL-1, 2, 3, 4, 5, 10, 16, 18, 19, 21, 22, 24, 27, 28, 32, 33, 34, 35, 36, 37, and 42 showed high affinity binding and hu3G8VL-15, 17, 20, 23, 25, 26, 29, 30, 31, 38, 39, 40 and 41 showed intermediate binding. CD16A binding proteins comprising the VL domains of hu3G8VL-1, hu3G8VL-22, and hu3G8VL-43 are considered to have particularly favorable binding properties (SEQ ID NO:73, SEQ ID NO:75 and SEQ ID NO:74, respectively).

5.1.2.2.1 Combinations of VL and/or VH Domains

[00186] As is known in the art and described elsewhere herein, immunoglobulin light and heavy chains can be recombinantly expressed under conditions in which they associate to produce a diabody, or can be so combined in vitro. It will thus be appreciated that a 3G8-derived VL-domain described herein can be combined a 3G8-derived VH-domain described herein to produce a CD16A binding diabody, and all such combinations are contemplated.

[00187] For illustration and not for limitation, examples of useful CD16A diabodies are those comprising at least one VH domain and at least one VL domain, where the VH domain is from hu3G8VH-1, hu3G8VH-22 or hu3G8VH-5 (SEQ ID NO:70, SEQ ID NO:72 and SEQ ID NO:71, respectively) and the VL domain is from hu3G8VL-1, hu3G8VL-22 or hu3G8VL-43 (SEQ ID NO:73, SEQ ID NO:75 and SEQ ID NO:43, respectively). In particular, humanized antibodies that comprise hu3G8VH-22 (SEQ ID NO:22) and either, hu3G8VL-1, hu3G8VL-22 or hu3G8VL-43 (SEQ ID NO:73, SEQ ID NO:72 and SEQ ID NO:74, respectively), or hu3G8VH-5 (SEQ ID NO:71) and hu3G8VL-1 (SEQ ID NO:73) have favorable properties.

[00188] It will be appreciated by those of skill that the sequences of VL and VH domains described here can be further modified by art-known methods such as affinity maturation (see Schier et al., 1996, J. Mol. Biol. 263:551-67; Daugherty et al., 1998, Protein Eng. 11:825-32; Boder et al., 1997, Nat. Biotechnol. 15:553-57; Boder et al., 2000, Proc. Natl. Acad. Sci. U.S.A 97:10701-705; Hudson and Souriau, 2003, Nature Medicine 9:129-39). For example, the CD16A binding proteins can be modified using affinity maturation

techniques to identify proteins with increased affinity for CD16A and/or decreased affinity for CD16B.

[00189] One exemplary CD16 binding protein is the mouse 3G8 antibody. Amino acid sequence comprising the VH and VL domains of humanized 3G8 are described in FIGS. 2, 9, 14 and set forth in SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73 and SEQ ID NO:74.

5.2 DIABODIES COMPRISING Fc REGIONS OR PORTIONS THEREOF

[00190] The invention encompasses diabody molecules comprising Fc domains or portions thereof (*e.g.*, a CH2 or CH3 domain). In certain embodiments the Fc domain, or portion(s) thereof, comprises one or more constant domain(s) of the Fc region of IgG2, IgG3 or IgG4 (*e.g.*, CH2 or CH3). In other embodiments, the invention encompasses molecules comprising and Fc domain or portion thereof, wherein said Fc domain or portion thereof comprises at least one amino acid modification (*e.g.* substitution) relative to a comparable wild-type Fc domain or portion thereof. Variant Fc domains are well known in the art, and are primarily used to alter the phenotype of the antibody comprising said variant Fc domain as assayed in any of the binding activity or effector function assays well known in the art, *e.g.* ELISA, SPR analysis, or ADCC. Such variant Fc domains, or portions thereof, have use in the present invention by conferring or modifying the effector function exhibited by a diabody molecule of the invention comprising an Fc domain (or portion thereof) as functionally assayed, *e.g.*, in an NK dependent or macrophage dependent assay. Fc domain variants identified as altering effector function are disclosed in International Application WO04/063351, U.S. Patent Application Publications 2005/0037000 and 2005/0064514, U.S. Provisional Applications 60/626,510, filed November 10, 2004, 60/636,663, filed December 15, 2004, and 60/781,564, filed March 10, 2006, and U.S. Patent Applications 11/271, 140, filed November 10, 2005, and 11/305,787, filed December 15, 2005, concurrent applications of the Inventors, each of which is incorporated by reference in its entirety.

[00191] In other embodiments, the invention encompasses the use of any Fc variant known in the art, such as those disclosed in Duncan et al, 1988, Nature 332:563-564; Lund et al., 1991, J. Immunol 147:2657-2662; Lund et al, 1992, Mol Immunol 29:53-59; Alegre et al, 1994, Transplantation 57:1537-1543; Hutchins et al. , 1995, Proc Natl. Acad Sci U S A 92:11980-11984; Jefferis et al, 1995, Immunol Lett. 44:111-117; Lund et al., 1995, Faseb

J 9:115-119; Jefferis et al, 1996, Immunol Lett 54:101-104; Lund et al, 1996, J Immunol 157:4963-4969; Armour et al, 1999, Eur J Immunol 29:2613-2624; Idusogie et al, 2000, J Immunol 164:4178-4184; Reddy et al, 2000, J Immunol 164:1925-1933; Xu *et al.*, 2000, Cell Immunol 200:16-26; Idusogie et al, 2001, J Immunol 166:2571-2575; Shields *et al.*, 2001, J Biol Chem 276:6591-6604; Jefferis et al, 2002, Immunol Lett 82:57-65; Presta *et al.*, 2002, Biochem Soc Trans 30:487-490); US 5,624,821; US 5,885,573; US 6,194,551; PCT WO 00/42072; PCT WO 99/58572; each of which is incorporated herein by reference in its entirety.

[00192] In certain embodiments, said one or more modifications to the amino acids of the Fc region reduce the affinity and avidity of the Fc region and, thus, the diabody molecule of the invention, for one or more FcγR receptors. In a specific embodiment, the invention encompasses diabodies comprising a variant Fc region, or portion thereof, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one FcγR, wherein said FcγR is FcγRIIA. In another specific embodiment, the invention encompasses diabodies comprising a variant Fc region, or portion thereof, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one FcγR, wherein said FcγR is FcγRIIA. In another specific embodiment, the invention encompasses diabodies comprising a variant Fc region, or portion thereof, wherein said variant Fc region comprises at least one amino acid modification relative to a wild type Fc region, which variant Fc region only binds one FcγR, wherein said FcγR is FcγRIIB. In certain embodiments, the invention encompasses molecules comprising a variant Fc domain wherein said variant confers or mediates increased ADCC activity and/or an increased binding to FcγRIIA (CD32A), relative to a molecule comprising no Fc domain or comprising a wild-type Fc domain, as measured using methods known to one skilled in the art and described herein. In alternate embodiments, the invention encompasses molecules comprising a variant Fc domain wherein said variant confers or mediates decreased ADCC activity (or other effector function) and/or an increased binding to FcγRIIB (CD32B), relative to a molecule comprising no Fc domain or comprising a wild-type Fc domain, as measured using methods known to one skilled in the art and described herein.

[00193] The invention also encompasses the use of an Fc domain comprising domains or regions from two or more IgG isotypes. As known in the art, amino acid modification of the Fc region can profoundly affect Fc-mediated effector function and/or binding activity. However, these alterations in functional characteristics can be further

refined and/or manipulated when implemented in the context of selected IgG isotypes. Similarly, the native characteristics of the isotype Fc may be manipulated by the one or more amino acid modifications. The multiple IgG isotypes (*i.e.*, IgG1, IgG2, IgG3 and IgG4) exhibit differing physical and functional properties including serum half-life, complement fixation, FcγR binding affinities and effector function activities (*e.g.* ADCC, CDC) due to differences in the amino acid sequences of their hinge and/or Fc domains. In certain embodiments, the amino acid modification and IgG Fc region are independently selected based on their respective, separate binding and/or effector function activities in order to engineer a diabody with desired characteristics. In most embodiments, said amino acid modifications and IgG hinge/Fc regions have been separately assayed for binding and/or effector function activity as described herein or known in the art in the context of an IgG1. In certain embodiments, said amino acid modification and IgG hinge/Fc region display similar functionality, *e.g.*, increased affinity for FcγRIIA, when separately assayed for FcγR binding or effector function in the context of the diabody molecule or other Fc-containing molecule (*e.g.* and immunoglobulin). The combination of said amino acid modification and selected IgG Fc region then act additively or, more preferably, synergistically to modify said functionality in the diabody molecule of the invention, relative to a diabody molecule of the invention comprising a wild-type Fc region. In other embodiments, said amino acid modification and IgG Fc region display opposite functionalities, *e.g.*, increased and decreased, respectively, affinity for FcγRIIA, when separately assayed for FcγR binding and/or effector function in the context of the diabody molecule or other Fc containing molecule (*e.g.*, an immunoglobulin) comprising a wild-type Fc region as described herein or known in the art; the combination of said "opposite" amino acid modification and selected IgG region then act to selectively temper or reduce a specific functionality in the diabody of the invention relative to a diabody of the invention not comprising an Fc region or comprising a wild-type Fc region of the same isotype. Alternatively, the invention encompasses variant Fc regions comprising combinations of amino acid modifications known in the art and selected IgG regions that exhibit novel properties, which properties were not detectable when said modifications and/or regions were independently assayed as described herein.

[00194] The functional characteristics of the multiple IgG isotypes, and domains thereof, are well known in the art. The amino acid sequences of IgG1, IgG2, IgG3 and IgG4 are presented in FIGS. 1A-1B. Selection and/or combinations of two or more domains from specific IgG isotypes for use in the methods of the invention may be based on any known parameter of the parent isotypes including affinity to FcγR (Table 7; Flesch and

Neppert, 1999, J. Clin. Lab. Anal. 14:141-156; Chappel et al., 1993, J. Biol. Chem. 33:25124-25131; Chappel et al., 1991, Proc. Natl. Acad. Sci. USA 88:9036-9040, each of which is hereby incorporated by reference in its entirety). For example, use of regions or domains from IgG isotypes that exhibit limited or no binding to FcγRIIB, *e.g.*, IgG2 or IgG4, may find particular use where a diabody is desired to be engineered to maximize binding to an activating receptor and minimize binding to an inhibitory receptor. Similarly, use of Fc regions or domains from IgG isotypes known to preferentially bind C1q or FcγRIIA, *e.g.*, IgG3 (Brüggemann et al., 1987, J. Exp. Med 166:1351-1361), may be combined with Fc amino acid modifications of known in the art to enhance ADCC, to engineer a diabody molecule such that effector function activity, *e.g.*, complement activation or ADCC, is maximized.

Table 7. General characteristics of IgG binding to FcγR, adapted from Flesch and Neppert, 1999, J. Clin. Lab. Anal. 14:141-156

Receptor	Estimated Affinity for IgG (M ⁻¹)	Relative Affinity
FcγRI	10 ⁸ - 10 ⁹	IgG3>IgG1>>IgG4 no-binding: IgG2
FcγRIIA R ^{I31 A}	<10 ⁷	IgG3>IgG1 no-binding: IgG2, IgG4
FcγRIIA H ^{I31 A}	<10 ⁷	IgG3>IgG1>IgG2 no-binding: IgG4
FcγRIIB ^A	<10 ⁷	IgG3>IgG1>IgG4 no-binding: IgG2
FcγRIII	<10 ⁷	IgG3=IgG1 no-binding: IgG2, IgG4

^A binds only complexed IgG
[00195]

5.3 MOLECULAR CONJUGATES

[00196] The diabody molecules of the invention may be recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to heterologous polypeptides (*i.e.*, an unrelated polypeptide; or portion thereof, preferably at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids of the polypeptide to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences.

[00197] Further, the diabody molecules of the invention (*i.e.*, polypeptides) may be conjugated to a therapeutic agent or a drug moiety that modifies a given biological response. As an alternative to direct conjugation, owing to the multiple epitope binding sites on the multivalent, *e.g.*, tetravalent, diabody molecules of the invention, at least one binding region

of the diabody may be designed to bind the therapeutic agent or desired drug moiety without affecting diabody binding.

[00198] Therapeutic agents or drug moieties are not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin (*i.e.*, PE-40), or diphtheria toxin, ricin, gelonin, and pokeweed antiviral protein, a protein such as tumor necrosis factor, interferons including, but not limited to, α -interferon (IFN- α), β -interferon (IFN- β), nerve growth factor (NGF), platelet derived growth factor (PDGF), tissue plasminogen activator (TPA), an apoptotic agent (*e.g.*, TNF- α , TNF- β , AIM I as disclosed in PCT Publication No. WO 97/33899), AIM II (*see*, PCT Publication No. WO 97/34911), Fas ligand (Takahashi *et al.*, *J. Immunol.*, 6:1567-1574, 1994), and VEGI (PCT Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent (*e.g.*, angiostatin or endostatin), or a biological response modifier such as, for example, a lymphokine (*e.g.*, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), and granulocyte colony stimulating factor ("G-CSF"), macrophage colony stimulating factor, ("M-CSF"), or a growth factor (*e.g.*, growth hormone ("GH"); proteases, or ribonucleases.

[00199] The diabody molecules of the invention (*i.e.*, polypeptides) can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz *et al.*, 1989, *Proc. Natl. Acad. Sci. USA*, 86:821-824, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, *Cell*, 37:767 1984) and the "flag" tag (Knappik *et al.*, *Biotechniques*, 17(4):754-761, 1994).

[00200] Additional fusion proteins may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to alter the activities of molecules of the invention (*e.g.*, epitope binding sites with higher affinities and lower dissociation rates). *See*, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten *et al.*, 1997, *Curr. Opinion Biotechnol.* 8:724-33; Harayama, 1998, *Trends Biotechnol.* 16:76; Hansson, *et al.*, 1999, *J. Mol. Biol.* 287:265; and Lorenzo and

Blasco, 1998, *BioTechniques* 24:308 (each of these patents and publications are hereby incorporated by reference in its entirety). The diabody molecules of the invention, or the nucleic acids encoding the molecules of the invention, may be further altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. One or more portions of a polynucleotide encoding a molecule of the invention, may be recombined with one or more components, motifs, sections, parts, domains, fragments, *etc.* of one or more heterologous molecules.

[00201] The present invention also encompasses diabody molecules of the invention conjugated to or immunospecifically recognizing a diagnostic or therapeutic agent or any other molecule for which serum half-life is desired to be increased/decreased and/or targeted to a particular subset of cells. The molecules of the invention can be used diagnostically to, for example, monitor the development or progression of a disease, disorder or infection as part of a clinical testing procedure to, *e.g.*, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the molecules of the invention to a detectable substance or by the molecules immunospecifically recognizing the detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the molecules of the invention or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art, or the molecule may immunospecifically recognize the detectable substance: immunospecifically binding said substance. *See*, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Such diagnosis and detection can be accomplished designing the molecules to immunospecifically recognize the detectable substance or by coupling the molecules of the invention to detectable substances including, but not limited to, various enzymes, enzymes including, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic group complexes such as, but not limited to, streptavidin/biotin and avidin/biotin; fluorescent materials such as, but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent material such as, but not limited to, luminol; bioluminescent materials such as, but not limited to, luciferase, luciferin, and aequorin; radioactive material such as, but not limited to, bismuth (^{213}Bi), carbon (^{14}C), chromium (^{51}Cr), cobalt (^{57}Co), fluorine (^{18}F), gadolinium (^{153}Gd , ^{159}Gd), gallium (^{68}Ga , ^{67}Ga), germanium (^{68}Ge), holmium

(^{166}Ho), indium (^{115}In , ^{113}In , ^{112}In , ^{111}In), iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), lanthanum (^{140}La), lutetium (^{177}Lu), manganese (^{54}Mn), molybdenum (^{99}Mo), palladium (^{103}Pd), phosphorous (^{32}P), praseodymium (^{142}Pr), promethium (^{149}Pm), rhenium (^{186}Re , ^{188}Re), rhodium (^{105}Rh), ruthenium (^{97}Ru), samarium (^{153}Sm), scandium (^{47}Sc), selenium (^{75}Se), strontium (^{85}Sr), sulfur (^{35}S), technetium (^{99}Tc), thallium (^{201}Tl), tin (^{113}Sn , ^{117}Sn), tritium (^3H), xenon (^{133}Xe), ytterbium (^{169}Yb , ^{175}Yb), yttrium (^{90}Y), zinc (^{65}Zn); positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions.

[00202] The diabody molecules of the invention may immunospecifically recognize or be conjugated to a therapeutic moiety such as a cytotoxin (*e.g.*, a cytostatic or cytotoxic agent), a therapeutic agent or a radioactive element (*e.g.*, alpha-emitters, gamma-emitters, *etc.*). Cytotoxins or cytotoxic agents include any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cisdichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

[00203] Moreover, a diabody molecule of the invention can be conjugated to or be designed to immunospecifically recognize therapeutic moieties such as a radioactive materials or macrocyclic chelators useful for conjugating radiometal ions (*see above* for examples of radioactive materials). In certain embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane- $\text{N},\text{N}',\text{N}'',\text{N}'''$ -tetraacetic acid (DOTA) which can be attached to the polypeptide via a linker molecule. Such linker molecules are commonly known in the art and described in Denardo *et al.*, 1998, *Clin Cancer Res.* 4:2483-90; Peterson *et al.*, 1999, *Bioconjug. Chem.* 10:553; and Zimmerman *et al.*, 1999, *Nucl. Med. Biol.* 26:943-50 each of which is incorporated herein by reference in their entireties.

[00204] Techniques for conjugating such therapeutic moieties to polypeptides, including *e.g.*, Fc domains, are well known; *see, e.g.*, Arnon *et al.*, "Monoclonal Antibodies

For Immunotargeting Of Drugs In Cancer Therapy”, in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), 1985, pp. 243-56, Alan R. Liss, Inc.); Hellstrom *et al.*, “Antibodies For Drug Delivery”, in *Controlled Drug Delivery (2nd Ed.)*, Robinson *et al.* (eds.), 1987, pp. 623-53, Marcel Dekker, Inc.); Thorpe, “Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review”, in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), 1985, pp. 475-506); “Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy”, in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), 1985, pp. 303-16, Academic Press; and Thorpe *et al.*, *Immunol. Rev.*, 62:119-58, 1982.

[00205] The diabody molecule of the invention may be administered with or without a therapeutic moiety conjugated to it, administered alone, or in combination with cytotoxic factor(s) and/or cytokine(s) for use as a therapeutic treatment. Where administered alone, at least one epitope of a multivalent, *e.g.*, tetravalent, diabody molecule may be designed to immunospecifically recognize a therapeutic agent, *e.g.*, cytotoxic factor(s) and/or cytokine(s), which may be administered concurrently or subsequent to the molecule of the invention. In this manner, the diabody molecule may specifically target the therapeutic agent in a manner similar to direct conjugation. Alternatively, a molecule of the invention, can be conjugated to an antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety. Diabody molecules of the invention may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

5.4 CHARACTERIZATION OF BINDING OF DIABODY MOLECULES

[00206] The diabody molecules of the present invention may be characterized in a variety of ways. In particular, molecules of the invention may be assayed for the ability to immunospecifically bind to an antigen, *e.g.*, FcRIIIA or FcRIIB, or, where the molecule comprises an Fc domain (or portion thereof) for the ability to exhibit Fc-FcγR interactions, *i.e.* specific binding of an Fc domain (or portion thereof) to an FcγR. Such an assay may be performed in solution (*e.g.*, Houghten, *Bio/Techniques*, 13:412-421, 1992), on beads (Lam, *Nature*, 354:82-84, 1991, on chips (Fodor, *Nature*, 364:555-556, 1993), on bacteria (U.S. Patent No. 5,223,409), on spores (U.S. Patent Nos. 5,571,698; 5,403,484; and 5,223,409), on plasmids (Cull *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:1865-1869, 1992) or on phage (Scott and Smith, *Science*, 249:386-390, 1990; Devlin, *Science*, 249:404-406, 1990; Cwirla

et al., *Proc. Natl. Acad. Sci. USA*, 87:6378-6382, 1990; and Felici, *J. Mol. Biol.*, 222:301-310, 1991) (each of these references is incorporated by reference herein in its entirety). Molecules that have been identified to immunospecifically bind to an antigen, *e.g.*, FcγRIIIA, can then be assayed for their specificity and affinity for the antigen.

[00207] Molecules of the invention that have been engineered to comprise multiple epitope binding domains may be assayed for immunospecific binding to one or more antigens (*e.g.*, cancer antigen and cross-reactivity with other antigens (*e.g.*, FcγR)) or, where the molecules comprise an Fc domain (or portion thereof) for Fc-FcγR interactions by any method known in the art. Immunoassays which can be used to analyze immunospecific binding, cross-reactivity, and Fc-FcγR interactions include, but are not limited to, competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (*see, e.g.*, Ausubel *et al.*, eds, 1994, *Current Protocols in Molecular Biology*, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety).

[00208] The binding affinity and the off-rate of antigen-binding domain interaction or Fc-FcγR interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen, such as tetrameric FcγR (*e.g.*, ^3H or ^{125}I , see Section 5.4.1) with a molecule of interest (*e.g.*, molecules of the present invention comprising multiple epitope binding domains in the presence of increasing amounts of unlabeled epitope, such as tetrameric FcγR (see Section 5.4.1), and the detection of the molecule bound to the labeled antigen. The affinity of the molecule of the present invention for an antigen and the binding off-rates can be determined from the saturation data by Scatchard analysis.

[00209] The affinities and binding properties of the molecules of the invention for an antigen or FcγR may be initially determined using *in vitro* assays (biochemical or immunological based assays) known in the art for antigen-binding domain or Fc-FcγR, interactions, including but not limited to ELISA assay, surface plasmon resonance assay, immunoprecipitation assays. Preferably, the binding properties of the molecules of the invention are also characterized by *in vitro* functional assays for determining one or more FcγR mediator effector cell functions, as described in section 5.4.2. In most preferred

embodiments, the molecules of the invention have similar binding properties in *in vivo* models (such as those described and disclosed herein) as those in *in vitro* based assays. However, the present invention does not exclude molecules of the invention that do not exhibit the desired phenotype in *in vitro* based assays but do exhibit the desired phenotype *in vivo*.

[00210] In some embodiments, screening and identifying molecules comprising multiple epitope binding domains and , optionally, Fc domains (or portions thereof) are done functional based assays, preferably in a high throughput manner. The functional based assays can be any assay known in the art for characterizing one or more FcγR mediated effector cell functions such as those described herein in Sections 5.4.2 and 5.4.3. Non-limiting examples of effector cell functions that can be used in accordance with the methods of the invention, include but are not limited to, antibody-dependent cell mediated cytotoxicity (ADCC), antibody-dependent phagocytosis, phagocytosis, opsonization, opsonophagocytosis, cell binding, rosetting, C1q binding, and complement dependent cell mediated cytotoxicity.

[00211] In a preferred embodiment, BIAcore kinetic analysis is used to determine the binding on and off rates of molecules of the present invention to an antigen or and FcγR. BIAcore kinetic analysis comprises analyzing the binding and dissociation of an antigen or FcγR from chips with immobilized molecules (*e.g.*, molecules comprising epitope binding domains or Fc domains (or portions thereof), respectively) on their surface. BIAcore analysis is described in Section 5.4.3.

[00212] Preferably, fluorescence activated cell sorting (FACS), using any of the techniques known to those skilled in the art, is used for immunological or functional based assay to characterize molecules of the invention. Flow sorters are capable of rapidly examining a large number of individual cells that have been bound, *e.g.*, opsonized, by molecules of the invention (*e.g.*, 10-100 million cells per hour) (Shapiro *et al.*, *Practical Flow Cytometry*, 1995). Additionally, specific parameters used for optimization of diabody behavior, include but are not limited to, antigen concentration (*i.e.*, FcγR tetrameric complex, see Section 5.4.1), kinetic competition time, or FACS stringency, each of which may be varied in order to select for the diabody molecules comprising molecules of the invention which exhibit specific binding properties, *e.g.*, concurrent binding to multiple epitopes. Flow cytometers for sorting and examining biological cells are well known in the art. Known flow cytometers are described, for example, in U.S. Patent Nos. 4,347,935; 5,464,581; 5,483,469; 5,602,039; 5,643,796; and 6,211,477; the entire contents of which are incorporated by reference herein. Other known flow cytometers are the FACS Vantage™

system manufactured by Becton Dickinson and Company, and the COPAS™ system manufactured by Union Biometrica.

[00213] Characterization of target antigen binding affinity or Fc-FcγR binding affinity, and assessment of target antigen or FcγR density on a cell surface may be made by methods well known in the art such as Scatchard analysis or by the use of kits as per manufacturer's instructions, such as Quantum™ Simply Cellular ® (Bangs Laboratories, Inc., Fishers, IN). The one or more functional assays can be any assay known in the art for characterizing one or more FcγR mediated effector cell function as known to one skilled in the art or described herein. In specific embodiments, the molecules of the invention comprising multiple epitope binding domains and, optionally, and Fc domain (or portion thereof) are assayed in an ELISA assay for binding to one or more target antigens or one or more FcγRs, *e.g.*, FcγRIIIA, FcγRIIA, FcγRIIA; followed by one or more ADCC assays. In some embodiments, the molecules of the invention are assayed further using a surface plasmon resonance-based assay, *e.g.*, BIAcore. Surface plasmon resonance-based assays are well known in the art, and are further discussed in Section 5.4.3, and exemplified herein, *e.g.*, in Example 6.1.

[00214] In most preferred embodiments, the molecules of the invention comprising multiple epitope binding domains and, optionally, and Fc domain (or portion thereof) is further characterized in an animal model for interaction with a target antigen (*e.g.*, an FcγR) or for Fc-FcγR interaction. Where Fc-FcγR interactions are to be assessed, preferred animal models for use in the methods of the invention are, for example, transgenic mice expressing human FcγRs, *e.g.*, any mouse model described in U.S. Patent No. 5,877,397, and 6,676,927 which are incorporated herein by reference in their entirety. Further transgenic mice for use in such methods include, but are not limited to, nude knockout FcγRIIIA mice carrying human FcγRIIIA; nude knockout FcγRIIIA mice carrying human FcγRIIA; nude knockout FcγRIIIA mice carrying human FcγRIIB and human FcγRIIA; nude knockout FcγRIIIA mice carrying human FcγRIIB and human FcγRIIA; nude knockout FcγRIIIA and FcγRIIA mice carrying human FcγRIIIA and FcγRIIA and nude knockout FcγRIIIA, FcγRIIA and FcγRIIB mice carrying human FcγRIIIA, FcγRIIA and FcγRIIB.

5.4.1 BINDING ASSAYS COMPRISING FcγR

[00215] Characterization of binding to FcγR by molecules comprising an Fc domain (or portion thereof) and/or comprising epitope binding domain specific for an FcγR may be done using any FcγR, including but not limited to polymorphic variants of FcγR. In some embodiments, a polymorphic variant of FcγRIIIA is used, which contains a phenylalanine

at position 158. In other embodiments, characterization is done using a polymorphic variant of FcγRIIIA which contains a valine at position 158. FcγRIIIA 158V displays a higher affinity for IgG1 than 158F and an increased ADCC activity (*see, e.g., Koene et al., 1997, Blood, 90:1109-14; Wu et al., 1997, J. Clin. Invest. 100: 1059-70*, both of which are incorporated herein by reference in their entirety); this residue in fact directly interacts with the lower hinge region of IgG1 as recently shown by IgG1-FcγRIIIA co-crystallization studies, *see, e.g., Sonderman et al., 2000, Nature, 100: 1059-70*, which is incorporated herein by reference in its entirety. Studies have shown that in some cases, therapeutic antibodies have improved efficacy in FcγRIIIA-158V homozygous patients. For example, humanized anti-CD20 monoclonal antibody Rituximab was therapeutically more effective in FcγRIIIA 158V homozygous patients compared to FcγRIIIA 158F homozygous patients (*See, e.g., Cartron et al., 2002 Blood, 99(3): 754-8*). In other embodiments, therapeutic molecules comprising this region may also be more effective on patients heterozygous for FcγRIIIA-158V and FcγRIIIA-158F, and in patients with FcγRIIA-131H. Although not intending to be bound by a particular mechanism of action, selection of molecules of the invention with alternate allotypes may provide for variants that once engineered into therapeutic diabodies will be clinically more efficacious for patients homozygous for said allotype.

[00216] An FcγR binding assay was developed for determining the binding of the molecules of the invention to FcγR, and, in particular, for determining binding of Fc domains to FcγR. The assay allowed detection and quantitation of Fc-FcγR interactions, despite the inherently weak affinity of the receptor for its ligand, *e.g., in the micromolar range for FcγRIIB and FcγRIIIA*. The method is described in detail in International Application WO04/063351 and U.S. Patent Application Publications 2005/0037000 and 2005/0064514, each of which is hereby incorporated by reference in its entirety. Briefly, the method involves the formation of an FcγR complex that may be sued in any standard immunoassay known in the art, *e.g., FACS, ELISA, surface plasmon resonance, etc.* Additionally, the FcγR complex has an improved avidity for an Fc region, relative to an uncomplexed FcγR. According to the invention, the preferred molecular complex is a tetrameric immune complex, comprising: (a) the soluble region of FcγR (*e.g., the soluble region of FcγRIIIA, FcγRIIA or FcγRIIB*); (b) a biotinylated 15 amino acid AVITAG sequence (AVITAG) operably linked to the C-terminus of the soluble region of FcγR (*e.g., the soluble region of FcγRIIIA, FcγRIIA or FcγRIIB*); and (c) streptavidin-phycoerythrin (SA-PE); in a molar ratio to form a tetrameric FcγR complex (preferably in a 5:1 molar

ratio). The fusion protein is biotinylated enzymatically, using for example, the *E.coli* Bir A enzyme, a biotin ligase which specifically biotinylates a lysine residue in the 15 amino acid AVITAG sequence. The biotinylated soluble FcγR proteins are then mixed with SA-PE in a 1X SA-PE:5X biotinylated soluble FcγR molar ratio to form a tetrameric FcγR complex.

[00217] Polypeptides comprising Fc regions have been shown to bind the tetrameric FcγR complexes with at least an 8-fold higher affinity than the monomeric uncomplexed FcγR. The binding of polypeptides comprising Fc regions to the tetrameric FcγR complexes may be determined using standard techniques known to those skilled in the art, such as for example, fluorescence activated cell sorting (FACS), radioimmunoassays, ELISA assays, *etc.*

[00218] The invention encompasses the use of the immune complexes comprising molecules of the invention, and formed according to the methods described above, for determining the functionality of molecules comprising an Fc region in cell-based or cell-free assays.

[00219] As a matter of convenience, the reagents may be provided in an assay kit, *i.e.*, a packaged combination of reagents for assaying the ability of molecules comprising an Fc regions to bind FcγR tetrameric complexes. Other forms of molecular complexes for use in determining Fc-FcγR interactions are also contemplated for use in the methods of the invention, *e.g.*, fusion proteins formed as described in U.S. Provisional Application 60/439,709, filed on January 13, 2003; which is incorporated herein by reference in its entirety.

5.4.2 FUNCTIONAL ASSAYS OF MOLECULES WITH VARIANT HEAVY CHAINS

[00220] The invention encompasses characterization of the molecules of the invention comprising multiple epitope binding domains and, optionally, Fc domains (or portions thereof) using assays known to those skilled in the art for identifying the effector cell function of the molecules. In particular, the invention encompasses characterizing the molecules of the invention for FcγR-mediated effector cell function. Additionally, where at least one of the target antigens of the diabody molecule of the invention is an FcγR, binding of the FcγR by the diabody molecule may serve to activate FcγR-mediated pathways similar to those activated by FcγR-Fc binding. Thus, where at least one epitope binding domain of the diabody molecule recognizes an FcγR, the diabody molecule may elicit FcγR-mediated effector cell function without containing an Fc domain (or portion thereof), or without concomitant Fc-FcγR binding. Examples of effector cell functions that can be assayed in accordance with the invention, include but are not limited to, antibody-dependent cell

mediated cytotoxicity, phagocytosis, opsonization, opsonophagocytosis, C1q binding, and complement dependent cell mediated cytotoxicity. Any cell-based or cell free assay known to those skilled in the art for determining effector cell function activity can be used (For effector cell assays, *see* Perussia *et al.*, 2000, *Methods Mol. Biol.* 121: 179-92; Baggiolini *et al.*, 1998 *Experientia*, 44(10): 841-8; Lehmann *et al.*, 2000 *J. Immunol. Methods*, 243(1-2): 229-42; Brown EJ. 1994, *Methods Cell Biol.*, 45: 147-64; Munn *et al.*, 1990 *J. Exp. Med.*, 172: 231-237, Abdul-Majid *et al.*, 2002 *Scand. J. Immunol.* 55: 70-81; Ding *et al.*, 1998, *Immunity* 8:403-411, each of which is incorporated by reference herein in its entirety).

[00221] In one embodiment, the molecules of the invention can be assayed for FcγR-mediated phagocytosis in human monocytes. Alternatively, the FcγR-mediated phagocytosis of the molecules of the invention may be assayed in other phagocytes, *e.g.*, neutrophils (polymorphonuclear leukocytes; PMN); human peripheral blood monocytes, monocyte-derived macrophages, which can be obtained using standard procedures known to those skilled in the art (*e.g.*, *see* Brown EJ. 1994, *Methods Cell Biol.*, 45: 147-164). In one embodiment, the function of the molecules of the invention is characterized by measuring the ability of THP-1 cells to phagocytose fluoresceinated IgG-opsonized sheep red blood cells (SRBC) by methods previously described (Tridandapani *et al.*, 2000, *J. Biol. Chem.* 275: 20480-7).

[00222] Another exemplary assay for determining the phagocytosis of the molecules of the invention is an antibody-dependent opsonophagocytosis assay (ADCP) which can comprise the following: coating a target bioparticle such as *Escherichia coli*-labeled FITC (Molecular Probes) or *Staphylococcus aureus*-FITC with (i) wild-type 4-4-20 antibody, an antibody to fluorescein (*See* Bedzyk *et al.*, 1989, *J. Biol. Chem.*, 264(3): 1565-1569, which is incorporated herein by reference in its entirety), as the control antibody for FcγR-dependent ADCP; or (ii) 4-4-20 antibody harboring the D265A mutation that knocks out binding to FcγRIII, as a background control for FcγR-dependent ADCP (iii) a diabody comprising the epitope binding domain of 4-4-20 and an Fc domain and/or an epitope binding domain specific for FcγRIII; and forming the opsonized particle; adding any of the opsonized particles described (i-iii) to THP-1 effector cells (a monocytic cell line available from ATCC) at a 1:1, 10:1, 30:1, 60:1, 75:1 or a 100: 1 ratio to allow FcγR-mediated phagocytosis to occur; preferably incubating the cells and *E. coli*-FITC/antibody at 37°C for 1.5 hour; adding trypan blue after incubation (preferably at room temperature for 2-3 min.) to the cells to quench the fluorescence of the bacteria that are adhered to the outside of the cell surface without being internalized; transferring cells into a FACS buffer (*e.g.*, 0.1%, BSA in PBS, 0.1%, sodium azide), analyzing the fluorescence of the THP1 cells using

FACS (e.g., BD FACS Calibur). Preferably, the THP-1 cells used in the assay are analyzed by FACS for expression of FcγR on the cell surface. THP-1 cells express both CD32A and CD64. CD64 is a high affinity FcγR that is blocked in conducting the ADCP assay in accordance with the methods of the invention. The THP-1 cells are preferably blocked with 100 μg/mL soluble IgG1 or 10% human serum. To analyze the extent of ADCP, the gate is preferably set on THP-1 cells and median fluorescence intensity is measured. The ADCP activity for individual mutants is calculated and reported as a normalized value to the wild type chMab 4-4-20 obtained. The opsonized particles are added to THP-1 cells such that the ratio of the opsonized particles to THP-1 cells is 30:1 or 60:1. In most preferred embodiments, the ADCP assay is conducted with controls, such as *E. coli*-FITC in medium, *E. coli*-FITC and THP-1 cells (to serve as FcγR-independent ADCP activity), *E. coli*-FITC, THP-1 cells and wild-type 4-4-20 antibody (to serve as FcγR-dependent ADCP activity), *E. coli*-FITC, THP-1 cells, 4-4-20 D265A (to serve as the background control for FcγR-dependent ADCP activity).

[00223] In another embodiment, the molecules of the invention can be assayed for FcγR-mediated ADCC activity in effector cells, e.g., natural killer cells, using any of the standard methods known to those skilled in the art (See e.g., Perussia *et al.*, 2000, *Methods Mol. Biol.* 121: 179-92; Weng *et al.*, 2003, *J. Clin. Oncol.* 21:3940-3947; Ding *et al.*, *Immunity*, 1998, 8:403-11). An exemplary assay for determining ADCC activity of the molecules of the invention is based on a ⁵¹Cr release assay comprising of: labeling target cells with [⁵¹Cr]Na₂CrO₄ (this cell-membrane permeable molecule is commonly used for labeling since it binds cytoplasmic proteins and although spontaneously released from the cells with slow kinetics, it is released massively following target cell necrosis); opsonizing the target cells with the molecules of the invention comprising variant heavy chains; combining the opsonized radiolabeled target cells with effector cells in a microtitre plate at an appropriate ratio of target cells to effector cells; incubating the mixture of cells for 16-18 hours at 37°C; collecting supernatants; and analyzing radioactivity. The cytotoxicity of the molecules of the invention can then be determined, for example using the following formula: % lysis = (experimental cpm - target leak cpm)/(detergent lysis cpm - target leak cpm) x 100%. Alternatively, % lysis = (ADCC-AICC)/(maximum release-spontaneous release). Specific lysis can be calculated using the formula: specific lysis = % lysis with the molecules of the invention - % lysis in the absence of the molecules of the invention. A graph can be generated by varying either the target: effector cell ratio or antibody concentration.

[00224] Preferably, the effector cells used in the ADCC assays of the invention are peripheral blood mononuclear cells (PBMC) that are preferably purified from normal human blood, using standard methods known to one skilled in the art, *e.g.*, using Ficoll-Paque density gradient centrifugation. Preferred effector cells for use in the methods of the invention express different FcγR activating receptors. The invention encompasses, effector cells, THP-1, expressing FcγRI, FcγRIIA and FcγRIIB, and monocyte derived primary macrophages derived from whole human blood expressing both FcγRIIA and FcγRIIB, to determine if heavy chain antibody mutants show increased ADCC activity and phagocytosis relative to wild type IgG1 antibodies.

[00225] The human monocyte cell line, THP-1, activates phagocytosis through expression of the high affinity receptor FcγRI and the low affinity receptor FcγRIIA (Fleit *et al.*, 1991, *J. Leuk. Biol.* 49: 556). THP-1 cells do not constitutively express FcγRIIA or FcγRIIB. Stimulation of these cells with cytokines effects the FcR expression pattern (Pricop *et al.*, 2000 *J. Immunol.* 166: 531-7). Growth of THP-1 cells in the presence of the cytokine IL4 induces FcγRIIB expression and causes a reduction in FcγRIIA and FcγRI expression. FcγRIIB expression can also be enhanced by increased cell density (Tridandapani *et al.*, 2002, *J. Biol Chem.* 277: 5082-9). In contrast, it has been reported that IFNγ can lead to expression of FcγRIIA (Pearse *et al.*, 1993 *PNAS USA* 90: 4314-8). The presence or absence of receptors on the cell surface can be determined by FACS using common methods known to one skilled in the art. Cytokine induced expression of FcγR on the cell surface provides a system to test both activation and inhibition in the presence of FcγRIIB. If THP-1 cells are unable to express the FcγRIIB the invention also encompasses another human monocyte cell line, U937. These cells have been shown to terminally differentiate into macrophages in the presence of IFNγ and TNF (Koren *et al.*, 1979, *Nature* 279: 328-331).

[00226] FcγR dependent tumor cell killing is mediated by macrophage and NK cells in mouse tumor models (Clynes *et al.*, 1998, *PNAS USA* 95: 652-656). The invention encompasses the use of elutriated monocytes from donors as effector cells to analyze the efficiency Fc mutants to trigger cell cytotoxicity of target cells in both phagocytosis and ADCC assays. Expression patterns of FcγRI, FcγRIIA, and FcγRIIB are affected by different growth conditions. FcγR expression from frozen elutriated monocytes, fresh elutriated monocytes, monocytes maintained in 10% FBS, and monocytes cultured in FBS + GM-CSF and or in human serum may be determined using common methods known to those skilled in the art. For example, cells can be stained with FcγR specific antibodies and

analyzed by FACS to determine FcR profiles. Conditions that best mimic macrophage *in vivo* FcγR expression is then used for the methods of the invention.

[00227] In some embodiments, the invention encompasses the use of mouse cells especially when human cells with the right FcγR profiles are unable to be obtained. In some embodiments, the invention encompasses the mouse macrophage cell line RAW264.7(ATCC) which can be transfected with human FcγRIIIA and stable transfectants isolated using methods known in the art, *see, e.g.,* Ralph *et al.*, *J. Immunol.* 119: 950-4). Transfectants can be quantitated for FcγRIIIA expression by FACS analysis using routine experimentation and high expressors can be used in the ADCC assays of the invention. In other embodiments, the invention encompasses isolation of spleen peritoneal macrophage expressing human FcγR from knockout transgenic mice such as those disclosed herein.

[00228] Lymphocytes may be harvested from peripheral blood of donors (PBM) using a Ficoll-Paque gradient (Pharmacia). Within the isolated mononuclear population of cells the majority of the ADCC activity occurs via the natural killer cells (NK) containing FcγRIIIA but not FcγRIIB on their surface. Results with these cells indicate the efficacy of the mutants on triggering NK cell ADCC and establish the reagents to test with elutriated monocytes.

[00229] Target cells used in the ADCC assays of the invention include, but are not limited to, breast cancer cell lines, *e.g.,* SK-BR-3 with ATCC accession number HTB-30 (*see, e.g.,* Tremp *et al.*, 1976, *Cancer Res.* 33-41); B-lymphocytes; cells derived from Burkitts lymphoma, *e.g.,* Raji cells with ATCC accession number CCL-86 (*see, e.g.,* Epstein *et al.*, 1965, *J. Natl. Cancer Inst.* 34: 231-240), and Daudi cells with ATCC accession number CCL-213 (*see, e.g.,* Klein *et al.*, 1968, *Cancer Res.* 28: 1300-10). The target cells must be recognized by the antigen binding site of the diabody molecule to be assayed.

[00230] The ADCC assay is based on the ability of NK cells to mediate cell death via an apoptotic pathway. NK cells mediate cell death in part by FcγRIIIA's recognition of an IgG Fc domain bound to an antigen on a cell surface. The ADCC assays used in accordance with the methods of the invention may be radioactive based assays or fluorescence based assays. The ADCC assay used to characterize the molecules of the invention comprising variant Fc regions comprises labeling target cells, *e.g.,* SK-BR-3, MCF-7, OVCAR3, Raji, Daudi cells, opsonizing target cells with an antibody that recognizes a cell surface receptor on the target cell via its antigen binding site; combining the labeled opsonized target cells and the effector cells at an appropriate ratio, which can be determined by routine

experimentation; harvesting the cells; detecting the label in the supernatant of the lysed target cells, using an appropriate detection scheme based on the label used. The target cells may be labeled either with a radioactive label or a fluorescent label, using standard methods known in the art. For example the labels include, but are not limited to, [^{51}Cr] Na_2CrO_4 ; and the acetoxymethyl ester of the fluorescence enhancing ligand, 2,2':6',2''-terpyridine-6-6''-dicarboxylate (TDA).

[00231] In a specific preferred embodiment, a time resolved fluorimetric assay is used for measuring ADCC activity against target cells that have been labeled with the acetoxymethyl ester of the fluorescence enhancing ligand, 2,2':6',2''-terpyridine-6-6''-dicarboxylate (TDA). Such fluorimetric assays are known in the art, *e.g.*, *see*, Blomberg *et al.*, 1996, *Journal of Immunological Methods*, 193: 199-206; which is incorporated herein by reference in its entirety. Briefly, target cells are labeled with the membrane permeable acetoxymethyl diester of TDA (bis(acetoxymethyl) 2,2':6',2''-terpyridine-6-6''-dicarboxylate, (BATDA), which rapidly diffuses across the cell membrane of viable cells. Intracellular esterases split off the ester groups and the regenerated membrane impermeable TDA molecule is trapped inside the cell. After incubation of effector and target cells, *e.g.*, for at least two hours, up to 3.5 hours, at 37°C, under 5% CO_2 , the TDA released from the lysed target cells is chelated with Eu^{3+} and the fluorescence of the Europium-TDA chelates formed is quantitated in a time-resolved fluorometer (*e.g.*, Victor 1420, Perkin Elmer/Wallace).

[00232] In another specific embodiment, the ADCC assay used to characterize the molecules of the invention comprising multiple epitope binding sites and, optionally, an Fc domain (or portion thereof) comprises the following steps: Preferably $4\text{--}5 \times 10^6$ target cells (*e.g.*, SK-BR-3, MCF-7, OVCAR3, Raji cells) are labeled with bis(acetoxymethyl) 2,2':6',2''-terpyridine-t-6''-dicarboxylate (DELFA BATDA Reagent, Perkin Elmer/Wallac). For optimal labeling efficiency, the number of target cells used in the ADCC assay should preferably not exceed 5×10^6 . BATDA reagent is added to the cells and the mixture is incubated at 37°C preferably under 5% CO_2 , for at least 30 minutes. The cells are then washed with a physiological buffer, *e.g.*, PBS with 0.125 mM sulfinpyrazole, and media containing 0.125 mM sulfinpyrazole. The labeled target cells are then opsonized (coated) with a molecule of the invention comprising an epitope binding domain specific for $\text{Fc}\gamma\text{RIIA}$ and, optionally, an Fc domain (or portion thereof). In preferred embodiments, the molecule used in the ADCC assay is also specific for a cell surface receptor, a tumor antigen, or a cancer antigen. The diabody molecule of the invention may specifically bind any cancer or tumor antigen, such as those listed in section 5.6.1. The target cells in the

ADCC assay are chosen according to the epitope binding sites engineered into the diabody of the invention, such that the diabody binds a cell surface receptor of the target cell specifically.

[00233] Target cells are added to effector cells, *e.g.*, PBMC, to produce effector:target ratios of approximately 1:1, 10:1, 30:1, 50:1, 75:1, or 100:1. The effector and target cells are incubated for at least two hours, up to 3.5 hours, at 37°C, under 5% CO₂. Cell supernatants are harvested and added to an acidic europium solution (*e.g.*, DELFIA Europium Solution, Perkin Elmer/Wallac). The fluorescence of the Europium-TDA chelates formed is quantitated in a time-resolved fluorometer (*e.g.*, Victor 1420, Perkin Elmer/Wallac). Maximal release (MR) and spontaneous release (SR) are determined by incubation of target cells with 1% TX-100 and media alone, respectively. Antibody independent cellular cytotoxicity (AICC) is measured by incubation of target and effector cells in the absence of a test molecule, *e.g.*, diabody of the invention. Each assay is preferably performed in triplicate. The mean percentage specific lysis is calculated as: $\text{Experimental release (ADCC) - AICC} / (\text{MR} - \text{SR}) \times 100$.

[00234] The invention encompasses assays known in the art, and exemplified herein, to characterize the binding of C1q and mediation of complement dependent cytotoxicity (CDC) by molecules of the invention comprising Fc domains (or portions thereof). To determine C1q binding, a C1q binding ELISA may be performed. An exemplary assay may comprise the following: assay plates may be coated overnight at 4°C with polypeptide comprising a molecule of the invention or starting polypeptide (control) in coating buffer. The plates may then be washed and blocked. Following washing, an aliquot of human C1q may be added to each well and incubated for 2 hrs at room temperature. Following a further wash, 100 uL of a sheep anti-complement C1q peroxidase conjugated antibody may be added to each well and incubated for 1 hour at room temperature. The plate may again be washed with wash buffer and 100 ul of substrate buffer containing OPD (O-phenylenediamine dihydrochloride (Sigma)) may be added to each well. The oxidation reaction, observed by the appearance of a yellow color, may be allowed to proceed for 30 minutes and stopped by the addition of 100 ul of 4.5 N H₂SO₄. The absorbance may then read at (492-405) nm.

[00235] To assess complement activation, a complement dependent cytotoxicity (CDC) assay may be performed, *e.g.* as described in Gazzano-Santoro et al., J. Immunol. Methods 202:163 (1996), which is incorporated herein by reference in its entirety. Briefly, various concentrations of the molecule comprising a (variant) Fc domain (or portion thereof) and human complement may be diluted with buffer. Cells which express the

antigen to which the diabody molecule binds may be diluted to a density of about 1×10^6 cells/ml. Mixtures of the diabody molecules comprising a (variant) Fc domain (or portion thereof), diluted human complement and cells expressing the antigen may be added to a flat bottom tissue culture 96 well plate and allowed to incubate for 2 hrs at 37C. and 5% CO₂ to facilitate complement mediated cell lysis. 50 uL of alamar blue (Accumed International) may then be added to each well and incubated overnight at 37 C. The absorbance is measured using a 96-well fluorometer with excitation at 530 nm and emission at 590 nm. The results may be expressed in relative fluorescence units (RFU). The sample concentrations may be computed from a standard curve and the percent activity as compared to nonvariant molecule, i.e., a molecule not comprising an Fc domain or comprising a non-variant Fc domain, is reported for the variant of interest.

5.4.3 OTHER ASSAYS

[00236] The molecules of the invention comprising multiple epitope binding domain and, optionally, an Fc domain may be assayed using any surface plasmon resonance based assays known in the art for characterizing the kinetic parameters of an antigen-binding domain or Fc-FcγR binding. Any SPR instrument commercially available including, but not limited to, BIAcore Instruments, available from Biacore AB (Uppsala, Sweden); IAsys instruments available from Affinity Sensors (Franklin, MA.); IBIS system available from Windsor Scientific Limited (Berks, UK), SPR-CELLIA systems available from Nippon Laser and Electronics Lab (Hokkaido, Japan), and SPR Detector Spreeta available from Texas Instruments (Dallas, TX) can be used in the instant invention. For a review of SPR-based technology see Mullet *et al.*, 2000, *Methods* 22: 77-91; Dong *et al.*, 2002, *Review in Mol. Biotech.*, 82: 303-23; Fivash *et al.*, 1998, *Current Opinion in Biotechnology* 9: 97-101; Rich *et al.*, 2000, *Current Opinion in Biotechnology* 11: 54-61; all of which are incorporated herein by reference in their entirety. Additionally, any of the SPR instruments and SPR based methods for measuring protein-protein interactions described in U.S. Patent No.'s 6,373,577; 6,289,286; 5,322,798; 5,341,215; 6,268,125 are contemplated in the methods of the invention, all of which are incorporated herein by reference in their entirety.

[00237] Briefly, SPR based assays involve immobilizing a member of a binding pair on a surface, and monitoring its interaction with the other member of the binding pair in solution in real time. SPR is based on measuring the change in refractive index of the solvent near the surface that occurs upon complex formation or dissociation. The surface onto which the immobilization occur is the sensor chip, which is at the heart of the SPR technology; it consists of a glass surface coated with a thin layer of gold and forms the basis

for a range of specialized surfaces designed to optimize the binding of a molecule to the surface. A variety of sensor chips are commercially available especially from the companies listed *supra*, all of which may be used in the methods of the invention. Examples of sensor chips include those available from BIAcore AB, Inc., *e.g.*, Sensor Chip CM5, SA, NTA, and HPA. A molecule of the invention may be immobilized onto the surface of a sensor chip using any of the immobilization methods and chemistries known in the art, including but not limited to, direct covalent coupling via amine groups, direct covalent coupling via sulfhydryl groups, biotin attachment to avidin coated surface, aldehyde coupling to carbohydrate groups, and attachment through the histidine tag with NTA chips.

[00238] In some embodiments, the kinetic parameters of the binding of molecules of the invention comprising multiple epitope binding sites and, optionally, and Fc domain, to an antigen or an FcγR may be determined using a BIAcore instrument (*e.g.*, BIAcore instrument 1000, BIAcore Inc., Piscataway, NJ). As discussed *supra*, see section 5.4.1, any FcγR can be used to assess the binding of the molecules of the invention either where at least one epitope binding site of the diabody molecule immunospecifically recognizes an FcγR, and/or where the diabody molecule comprises an Fc domain (or portion thereof). In a specific embodiment the FcγR is FcγRIIIA, preferably a soluble monomeric FcγRIIIA. For example, in one embodiment, the soluble monomeric FcγRIIIA is the extracellular region of FcγRIIIA joined to the linker-AVITAG sequence (*see*, U.S. Provisional Application No. 60/439,498, filed on January 9, 2003 (Attorney Docket No. 11183-004-888) and U.S. Provisional Application No. 60/456,041 filed on March 19, 2003, which are incorporated herein by reference in their entireties). In another specific embodiment, the FcγR is FcγRIIB, preferably a soluble dimeric FcγRIIB. For example in one embodiment, the soluble dimeric FcγRIIB protein is prepared in accordance with the methodology described in U.S. Provisional application No. 60/439,709 filed on January 13, 2003, which is incorporated herein by reference in its entirety.

[00239] For all immunological assays, FcγR recognition/binding by a molecule of the invention may be effected by multiple domains: in certain embodiments, molecules of the invention immunospecifically recognize an FcγR via one of the multiple epitope binding domains; in yet other embodiments, where the molecule of the invention comprises an Fc domain (or portion thereof), the diabody molecule may immunospecifically recognize an FcγR via Fc-FcγR interactions; in yet further embodiments, where a molecule of the invention comprises both an Fc domain (or portion thereof) and an epitope binding site that

immunospecifically recognizes an FcγR, the diabody molecule may recognize an FcγR via one or both of an epitope binding domain and the Fc domain (or portion thereof). An exemplary assay for determining the kinetic parameters of a molecule comprising multiple epitope binding domains and, optionally, and Fc domain (or portion thereof) to an antigen and/or an FcγR using a BIAcore instrument comprises the following: a first antigen is immobilized on one of the four flow cells of a sensor chip surface, preferably through amine coupling chemistry such that about 5000 response units (RU) of said first antigen is immobilized on the surface. Once a suitable surface is prepared, molecules of the invention that immunospecifically recognize said first antigen are passed over the surface, preferably by one minute injections of a 20 μg/mL solution at a 5 μL/mL flow rate. Levels of molecules of the invention bound to the surface at this stage typically ranges between 400 and 700 RU. Next, dilution series of a second antigen (*e.g.*, FcγR) or FcγR receptor in HBS-P buffer (20mM HEPES, 150 mM NaCl, 3mM EDTA, pH 7.5) are injected onto the surface at 100 μL/min. Regeneration of molecules between different second antigen or receptor dilutions is carried out preferably by single 5 second injections of 100mM NaHCO₃ pH 9.4; 3M NaCl. Any regeneration technique known in the art is contemplated in the method of the invention.

[00240] Once an entire data set is collected, the resulting binding curves are globally fitted using computer algorithms supplied by the SPR instrument manufacturer, *e.g.*, BIAcore, Inc. (Piscataway, NJ). These algorithms calculate both the K_{on} and K_{off} , from which the apparent equilibrium binding constant, K_d is deduced as the ratio of the two rate constants (*i.e.*, K_{off}/K_{on}). More detailed treatments of how the individual rate constants are derived can be found in the BIAevaluation Software Handbook (BIAcore, Inc., Piscataway, NJ). The analysis of the generated data may be done using any method known in the art. For a review of the various methods of interpretation of the kinetic data generated *see* Myszka, 1997, *Current Opinion in Biotechnology* 8: 50-7; Fisher *et al.*, 1994, *Current Opinion in Biotechnology* 5: 389-95; O'Shannessy, 1994, *Current Opinion in Biotechnology*, 5:65-71; Chaiken *et al.*, 1992, *Analytical Biochemistry*, 201: 197-210; Morton *et al.*, 1995, *Analytical Biochemistry* 227: 176-85; O'Shannessy *et al.*, 1996, *Analytical Biochemistry* 236: 275-83; all of which are incorporated herein by reference in their entirety.

[00241] In preferred embodiments, the kinetic parameters determined using an SPR analysis, *e.g.*, BIAcore, may be used as a predictive measure of how a molecule of the invention will function in a functional assay, *e.g.*, ADCC. An exemplary method for predicting the efficacy of a molecule of the invention based on kinetic parameters obtained

from an SPR analysis may comprise the following: determining the K_{off} values for binding of a molecule of the invention to FcγRIIIA and FcγRIIB (via an epitope binding domain and/or an Fc domain (or portion thereof)); plotting (1) $K_{off}(wt)/K_{off}(mut)$ for FcγRIIIA; (2) $K_{off}(mut)/K_{off}(wt)$ for FcγRIIB against the ADCC data. Numbers higher than one show a decreased dissociation rate for FcγRIIIA and an increased dissociation rate for FcγRIIB relative to wild type; and possess and enhanced ADCC function.

5.5 METHODS OF PRODUCING DIABODY MOLECULES OF THE INVENTION

[00242] The diabody molecules of the present invention can be produced using a variety of methods well known in the art, including de novo protein synthesis and recombinant expression of nucleic acids encoding the binding proteins. The desired nucleic acid sequences can be produced by recombinant methods (e.g., PCR mutagenesis of an earlier prepared variant of the desired polynucleotide) or by solid-phase DNA synthesis. Usually recombinant expression methods are used. In one aspect, the invention provides a polynucleotide that comprises a sequence encoding a CD16A VH and/or VL; in another aspect, the invention provides a polynucleotide that comprises a sequence encoding a CD32B VH and/or VL. Because of the degeneracy of the genetic code, a variety of nucleic acid sequences encode each immunoglobulin amino acid sequence, and the present invention includes all nucleic acids encoding the binding proteins described herein.

5.5.1 POLYNUCLEOTIDES ENCODING MOLECULES OF THE INVENTION.

[00243] The present invention also includes polynucleotides that encode the molecules of the invention, including the polypeptides and antibodies. The polynucleotides encoding the molecules of the invention may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art.

[00244] Once the nucleotide sequence of the molecules that are identified by the methods of the invention is determined, the nucleotide sequence may be manipulated using methods well known in the art, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, *etc.* (*see, for example, the techniques described in Sambrook et al., 2001, Molecular Cloning, A Laboratory Manual, 3rd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY; and Ausubel et al., eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties*), to generate, for example, antibodies having a different amino acid sequence, for example by generating amino acid substitutions, deletions, and/or insertions.

[00245] In one embodiment, human libraries or any other libraries available in the art, can be screened by standard techniques known in the art, to clone the nucleic acids encoding the molecules of the invention.

5.5.2 RECOMBINANT EXPRESSION OF MOLECULES OF THE INVENTION

[00246] Once a nucleic acid sequence encoding molecules of the invention (*i.e.*, antibodies) has been obtained, the vector for the production of the molecules may be produced by recombinant DNA technology using techniques well known in the art. Methods which are well known to those skilled in the art can be used to construct expression vectors containing the coding sequences for the molecules of the invention and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. (See, for example, the techniques described in Sambrook *et al.*, 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.* eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY).

[00247] An expression vector comprising the nucleotide sequence of a molecule identified by the methods of the invention can be transferred to a host cell by conventional techniques (*e.g.*, electroporation, liposomal transfection, and calcium phosphate precipitation) and the transfected cells are then cultured by conventional techniques to produce the molecules of the invention. In specific embodiments, the expression of the molecules of the invention is regulated by a constitutive, an inducible or a tissue, specific promoter.

[00248] The host cells used to express the molecules identified by the methods of the invention may be either bacterial cells such as *Escherichia coli*, or, preferably, eukaryotic cells, especially for the expression of whole recombinant immunoglobulin molecule. In particular, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for immunoglobulins (Foecking *et al.*, 1998, *Gene* 45:101; Cockett *et al.*, 1990, *Bio/Technology* 8:2).

[00249] A variety of host-expression vector systems may be utilized to express the molecules identified by the methods of the invention. Such host-expression systems represent vehicles by which the coding sequences of the molecules of the invention may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express the molecules of

the invention *in situ*. These include, but are not limited to, microorganisms such as bacteria (*e.g.*, *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing coding sequences for the molecules identified by the methods of the invention; yeast (*e.g.*, *Saccharomyces Pichia*) transformed with recombinant yeast expression vectors containing sequences encoding the molecules identified by the methods of the invention; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing the sequences encoding the molecules identified by the methods of the invention; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus (CaMV) and tobacco mosaic virus (TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing sequences encoding the molecules identified by the methods of the invention; or mammalian cell systems (*e.g.*, COS, CHO, BHK, 293, 293T, 3T3 cells, lymphotic cells (*see* U.S. 5,807,715), Per C.6 cells (human retinal cells developed by Crucell) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).

[00250] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther *et al.*, 1983, *EMBO J.* 2:1791), in which the antibody coding sequence may be ligated individually into the vector in frame with the *lac Z* coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, *Nucleic Acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.* 24:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to a matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[00251] In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-

essential regions (*e.g.*, the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (*e.g.*, the polyhedrin promoter).

[00252] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the immunoglobulin molecule in infected hosts (*e.g.*, see Logan & Shenk, 1984, *Proc. Natl. Acad. Sci. USA* 81:355-359). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, *etc.* (see Bittner *et al.*, 1987, *Methods in Enzymol.* 153:51-544).

[00253] In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products may be important for the function of the protein. For example, in certain embodiments, the polypeptides comprising a diabody molecule of the invention may be expressed as a single gene product (*e.g.*, as a single polypeptide chain, *i.e.*, as a polyprotein precursor), requiring proteolytic cleavage by native or recombinant cellular mechanisms to form the separate polypeptides of the diabody molecules of the invention. The invention thus encompasses engineering a nucleic acid sequence to encode a polyprotein precursor molecule comprising the polypeptides of the invention, which includes coding sequences capable of directing post translational cleavage of said polyprotein precursor. Post-translational cleavage of the polyprotein precursor results in the polypeptides of the invention. The post translational cleavage of the precursor molecule comprising the polypeptides of the invention may occur *in vivo* (*i.e.*, within the host cell by native or recombinant cell systems/mechanisms, *e.g.* furin cleavage at an appropriate site) or may occur *in vitro* (*e.g.* incubation of said polypeptide chain in a composition comprising proteases or peptidases of known activity and/or in a composition comprising conditions or

reagents known to foster the desired proteolytic action). Purification and modification of recombinant proteins is well known in the art such that the design of the polyprotein precursor could include a number of embodiments readily appreciated by a skilled worker. Any known proteases or peptidases known in the art can be used for the described modification of the precursor molecule, *e.g.*, thrombin (which recognizes the amino acid sequence LVPR[^]GS (SEQ ID NO:91)), or factor Xa (which recognizes the amino acid sequence I(E/D)GR[^] (SEQ ID NO:92) (Nagani et al., 1985, PNAS USA 82:7252-7255, and reviewed in Jenny *et al.*, 2003, Protein Expr. Purif. 31:1-11, each of which is incorporated by reference herein in its entirety)), enterokinase (which recognizes the amino acid sequence DDDDK[^] (SEQ ID NO:93) (Collins-Racie et al., 1995, Biotechnol. 13:982-987 hereby incorporated by reference herein in its entirety)), furin (which recognizes the amino acid sequence RXXR[^], with a preference for RX(K/R)R[^] (SEQ ID NO:94 and SEQ ID NO:95, respectively) (additional R at P6 position appears to enhance cleavage)), and AcTEV (which recognizes the amino acid sequence ENLYFQ[^]G (SEQ ID NO:96) (Parks et al., 1994, Anal. Biochem. 216:413 hereby incorporated by reference herein in its entirety)) and the Foot and Mouth Disease Virus Protease C3. See for example, section 6.4, *supra*.

[00254] Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 293T, 3T3, WI38, BT483, Hs578T, HTB2, BT20 and T47D, CRL7030 and Hs578Bst.

[00255] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express an antibody of the invention may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, *etc.*), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded

into cell lines. This method may advantageously be used to engineer cell lines which express the antibodies of the invention. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the molecules of the invention.

[00256] A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, *Cell* 11: 223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1992, *Proc. Natl. Acad. Sci. USA* 48: 202), and adenine phosphoribosyltransferase (Lowy *et al.*, 1980, *Cell* 22: 817) genes can be employed in tk-, hgprt- or apt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci. USA* 77:357; O'Hare *et al.*, 1981, *Proc. Natl. Acad. Sci. USA* 78: 1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, *Proc. Natl. Acad. Sci. USA* 78: 2072); neo, which confers resistance to the aminoglycoside G-418 *Clinical Pharmacy* 12: 488-505; Wu and Wu, 1991, 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, 1993, *Science* 260:926-932; and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIB TECH* 11(5):155-215). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel *et al.* (eds.), 1993, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY; Kriegler, 1990, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY; and in Chapters 12 and 13, Dracopoli *et al.* (eds), 1994, Current Protocols in Human Genetics, John Wiley & Sons, NY.; Colberre-Garapin *et al.*, 1981, *J. Mol. Biol.* 150:1; and hygro, which confers resistance to hygromycin (Santerre *et al.*, 1984, *Gene* 30:147).

[00257] The expression levels of a molecule of the invention can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol. 3 (Academic Press, New York, 1987). When a marker in the vector system expressing an antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the nucleotide sequence of a polypeptide of the diabody molecule, production of the polypeptide will also increase (Crouse *et al.*, 1983, *Mol. Cell. Biol.* 3:257).

[00258] The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding the first polypeptide of the diabody molecule and the second vector encoding the second polypeptide of the diabody molecule. The two vectors may contain identical selectable markers which enable equal expression of both

polypeptides. Alternatively, a single vector may be used which encodes both polypeptides. The coding sequences for the polypeptides of the molecules of the invention may comprise cDNA or genomic DNA.

[00259] Once a molecule of the invention (*i.e.*, diabodies) has been recombinantly expressed, it may be purified by any method known in the art for purification of polypeptides, polyproteins or diabodies (*e.g.*, analogous to antibody purification schemes based on antigen selectivity) for example, by chromatography (*e.g.*, ion exchange, affinity, particularly by affinity for the specific antigen (optionally after Protein A selection where the diabody molecule comprises an Fc domain (or portion thereof)), and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of polypeptides, polyproteins or diabodies.

5.6 PROPHYLACTIC AND THERAPEUTIC METHODS

[00260] The molecules of the invention are particularly useful for the treatment and/or prevention of a disease, disorder or infection where an effector cell function (*e.g.*, ADCC) mediated by Fc γ R is desired (*e.g.*, cancer, infectious disease). As discussed *supra*, the diabodies of the invention may exhibit antibody-like functionality in eliciting effector function although the diabody molecule does not comprise an Fc domain. By comprising at least one epitope binding domain that immunospecifically recognizes an Fc γ R, the diabody molecule may exhibit Fc γ R binding and activity analogous to Fc-Fc γ R interactions. For example, molecules of the invention may bind a cell surface antigen and an Fc γ R (*e.g.*, Fc γ RIIIA) on an immune effector cell (*e.g.*, NK cell), stimulating an effector function (*e.g.*, ADCC, CDC, phagocytosis, opsonization, etc.) against said cell.

[00261] In other embodiments, the diabody molecule of the invention comprises an Fc domain (or portion thereof). In such embodiments, the Fc domain may further comprise at least one amino acid modification relative to a wild-type Fc domain (or portion thereof) and/or may comprise domains from one or more IgG isotypes (*e.g.*, IgG1, IgG2, IgG3 or IgG4). Molecules of the invention comprising variant Fc domains may exhibit conferred or altered phenotypes relative to molecules comprising the wild type Fc domain such as an altered or conferred effector function activity (*e.g.*, as assayed in an NK dependent or macrophage dependent assay). In said embodiments, molecules of the invention with conferred or altered effector function activity are useful for the treatment and/or prevention of a disease, disorder or infection where an enhanced efficacy of effector function activity is desired. In certain embodiments, the diabody molecules of the invention comprising an Fc domain (or portion thereof) mediate complement dependent cascade. Fc domain variants

identified as altering effector function are disclosed in International Application WO04/063351, U.S. Patent Application Publications 2005/0037000 and 2005/0064514, U.S. Provisional Applications 60/626,510, filed November 10, 2004, 60/636,663, filed December 15, 2004, and 60/781,564, filed March 10, 2006, and U.S. Patent Applications 11/271, 140, filed November 10, 2005, and 11/305,787, filed December 15, 2005, concurrent applications of the Inventors, each of which is incorporated by reference in its entirety.

[00262] The invention encompasses methods and compositions for treatment, prevention or management of a cancer in a subject, comprising administering to the subject a therapeutically effective amount of one or more molecules comprising one or more epitope binding sites, and optionally, an Fc domain (or portion thereof) engineered in accordance with the invention, which molecule further binds a cancer antigen. Molecules of the invention are particularly useful for the prevention, inhibition, reduction of growth or regression of primary tumors, metastasis of cancer cells, and infectious diseases. Although not intending to be bound by a particular mechanism of action, molecules of the invention mediate effector function resulting in tumor clearance, tumor reduction or a combination thereof. In alternate embodiments, the diabodies of the invention mediate therapeutic activity by cross-linking of cell surface antigens and/or receptors and enhanced apoptosis or negative growth regulatory signaling.

[00263] Although not intending to be bound by a particular mechanism of action, the diabody molecules of the invention exhibit enhanced therapeutic efficacy relative to therapeutic antibodies known in the art, in part, due to the ability of diabody to immunospecifically bind a target cell which expresses a particular antigen (*e.g.*, FcγR) at reduced levels, for example, by virtue of the ability of the diabody to remain on the target cell longer due to an improved avidity of the diabody-epitope interaction.

[00264] The diabodies of the invention with enhanced affinity and avidity for antigens (*e.g.*, FcγRs) are particularly useful for the treatment, prevention or management of a cancer, or another disease or disorder, in a subject, wherein the FcγRs are expressed at low levels in the target cell populations. As used herein, FcγR expression in cells is defined in terms of the density of such molecules per cell as measured using common methods known to those skilled in the art. The molecules of the invention comprising multiple epitope binding sites and, optionally, and FcγR (or portion thereof) preferably also have a conferred or an enhanced avidity and affinity and/or effector function in cells which express a target antigen, *e.g.*, a cancer antigen, at a density of 30,000 to 20,000 molecules/cell, at a density of 20,000 to 10,000 molecules/cell, at a density of 10,000 molecules/cell or less, at a density

of 5000 molecules/cell or less, or at a density of 1000 molecules /cell or less. The molecules of the invention have particular utility in treatment, prevention or management of a disease or disorder, such as cancer, in a sub-population, wherein the target antigen is expressed at low levels in the target cell population.

[00265] The molecules of the invention may also be advantageously utilized in combination with other therapeutic agents known in the art for the treatment or prevention of diseases, such as cancer, autoimmune disease, inflammatory disorders, and infectious diseases. In a specific embodiment, molecules of the invention may be used in combination with monoclonal or chimeric antibodies, lymphokines, or hematopoietic growth factors (such as, *e.g.*, IL-2, IL-3 and IL-7), which, for example, serve to increase the number or activity of effector cells which interact with the molecules and, increase immune response. The molecules of the invention may also be advantageously utilized in combination with one or more drugs used to treat a disease, disorder, or infection such as, for example anti-cancer agents, anti-inflammatory agents or anti-viral agents, *e.g.*, as detailed in Section 5.7.

5.6.1 CANCERS

[00266] The invention encompasses methods and compositions for treatment or prevention of cancer in a subject comprising administering to the subject a therapeutically effective amount of one or more molecules comprising multiple epitope binding domains. In some embodiments, the invention encompasses methods and compositions for the treatment or prevention of cancer in a subject with FcγR polymorphisms such as those homozygous for the FcγRIIA-158V or FcγRIIA-158F alleles. In some embodiments, the invention encompasses engineering at least one epitope binding domain of the diabody molecule to immunospecifically bind FcγRIIA (158F). In other embodiments, the invention encompasses engineering at least one epitope binding domain of the diabody molecule to immunospecifically bind FcγRIIA (158V).

[00267] The efficacy of standard monoclonal antibody therapy depends on the FcγR polymorphism of the subject (Carton *et al.*, 2002 *Blood*, 99: 754-8; Weng *et al.*, 2003 *J Clin Oncol*.21(21):3940-7 both of which are incorporated herein by reference in their entireties). These receptors are expressed on the surface of the effector cells and mediate ADCC. High affinity alleles, of the low affinity activating receptors, improve the effector cells' ability to mediate ADCC. In contrast to relying on Fc-FcγR interactions to effect effector function, the methods of the invention encompass engineering molecules to immunospecifically recognize the low affinity activating receptors, allowing the molecules to be designed for a specific polymorphism. Alternately or additionally, the molecule of the invention may be

engineered to comprise a variant Fc domain that exhibits enhanced affinity to FcγR (relative to a wild type Fc domain) on effector cells. The engineered molecules of the invention provide better immunotherapy reagents for patients regardless of their FcγR polymorphism.

[00268] Diabody molecules engineered in accordance with the invention are tested by ADCC using either a cultured cell line or patient derived PMBC cells to determine the ability of the Fc mutations to enhance ADCC. Standard ADCC is performed using methods disclosed herein. Lymphocytes are harvested from peripheral blood using a Ficoll-Paque gradient (Pharmacia). Target cells, *i.e.*, cultured cell lines or patient derived cells, are loaded with Europium (PerkinElmer) and incubated with effectors for 4 hrs at 37°C. Released Europium is detected using a fluorescent plate reader (Wallac). The resulting ADCC data indicates the efficacy of the molecules of the invention to trigger NK cell mediated cytotoxicity and establish which molecules can be tested with both patient samples and elutriated monocytes. Diabody molecules showing the greatest potential for eliciting ADCC activity are then tested in an ADCC assay using PBMCs from patients. PBMC from healthy donors are used as effector cells.

[00269] Accordingly, the invention provides methods of preventing or treating cancer characterized by a cancer antigen by engineering the diabody molecule to immunospecifically recognize said cancer antigen such that the diabody molecule is itself cytotoxic (*e.g.*, via crosslinking of surface receptors leading to increased apoptosis or downregulation of proliferative signals) and/or comprises an Fc domain, according to the invention, and/or mediates one or more effector function (*e.g.*, ADCC, phagocytosis). The diabodies that have been engineered according to the invention are useful for prevention or treatment of cancer, since they have a cytotoxic activity (*e.g.*, enhanced tumor cell killing and/or enhanced for example, ADCC activity or CDC activity).

[00270] Cancers associated with a cancer antigen may be treated or prevented by administration of a diabody that binds a cancer antigen and is cytotoxic, and/or has been engineered according to the methods of the invention to exhibit effector function. For example, but not by way of limitation, cancers associated with the following cancer antigens may be treated or prevented by the methods and compositions of the invention: KS 1/4 pan-carcinoma antigen (Perez and Walker, 1990, *J. Immunol.* 142:32-37; Bumal, 1988, *Hybridoma* 7(4):407-415), ovarian carcinoma antigen (CA125) (Yu *et al.*, 1991, *Cancer Res.* 51(2):48-475), prostatic acid phosphatase (Tailor *et al.*, 1990, *Nucl. Acids Res.* 18(1):4928), prostate specific antigen (Henttu and Vihko, 1989, *Biochem. Biophys. Res. Comm.* 10(2):903-910; Israeli *et al.*, 1993, *Cancer Res.* 53:227-230), melanoma-associated antigen p97 (Estin *et al.*, 1989, *J. Natl. Cancer Instit.* 81(6):445-44), melanoma antigen

gp75 (Vijayasardahl *et al.*, 1990, *J. Exp. Med.* 171(4):1375-1380), high molecular weight melanoma antigen (HMW-MAA) (Natali *et al.*, 1987, *Cancer* 59:55-3; Mittelman *et al.*, 1990, *J. Clin. Invest.* 86:2136-2144)), prostate specific membrane antigen, carcinoembryonic antigen (CEA) (Foon *et al.*, 1994, *Proc. Am. Soc. Clin. Oncol.* 13:294), polymorphic epithelial mucin antigen, human milk fat globule antigen, Colorectal tumor-associated antigens such as: CEA, TAG-72 (Yokata *et al.*, 1992, *Cancer Res.* 52:3402-3408), CO17-1A (Ragnhammar *et al.*, 1993, *Int. J. Cancer* 53:751-758); GICA 19-9 (Herlyn *et al.*, 1982, *J. Clin. Immunol.* 2:135), CTA-1 and LEA, Burkitt's lymphoma antigen-38.13, CD19 (Ghetie *et al.*, 1994, *Blood* 83:1329-1336), human B-lymphoma antigen-CD20 (Reff *et al.*, 1994, *Blood* 83:435-445), CD33 (Sgouros *et al.*, 1993, *J. Nucl. Med.* 34:422-430), melanoma specific antigens such as ganglioside GD2 (Saleh *et al.*, 1993, *J. Immunol.*, 151, 3390-3398), ganglioside GD3 (Shitara *et al.*, 1993, *Cancer Immunol. Immunother.* 36:373-380), ganglioside GM2 (Livingston *et al.*, 1994, *J. Clin. Oncol.* 12:1036-1044), ganglioside GM3 (Hoon *et al.*, 1993, *Cancer Res.* 53:5244-5250), tumor-specific transplantation type of cell-surface antigen (TSTA) such as virally-induced tumor antigens including T-antigen DNA tumor viruses and envelope antigens of RNA tumor viruses, oncofetal antigen-alpha-fetoprotein such as CEA of colon, bladder tumor oncofetal antigen (Hellstrom *et al.*, 1985, *Cancer Res.* 45:2210-2188), differentiation antigen such as human lung carcinoma antigen L6, L20 (Hellstrom *et al.*, 1986, *Cancer Res.* 46:3917-3923), antigens of fibrosarcoma, human leukemia T cell antigen-Gp37 (Bhattacharya-Chatterjee *et al.*, 1988, *J. of Immun.* 141:1398-1403), neoglycoprotein, sphingolipids, breast cancer antigen such as EGFR (Epidermal growth factor receptor), HER2 antigen (p185^{HER2}), polymorphic epithelial mucin (PEM) (Hilkens *et al.*, 1992, *Trends in Bio. Chem. Sci.* 17:359), malignant human lymphocyte antigen-APO-1 (Bernhard *et al.*, 1989, *Science* 245:301-304), differentiation antigen (Feizi, 1985, *Nature* 314:53-57) such as I antigen found in fetal erythrocytes and primary endoderm, I(Ma) found in gastric adenocarcinomas, M18 and M39 found in breast epithelium, SSEA-1 found in myeloid cells, VEP8, VEP9, Myl, VIM-D5, and D156-22 found in colorectal cancer, TRA-1-85 (blood group H), C14 found in colonic adenocarcinoma, F3 found in lung adenocarcinoma, AH6 found in gastric cancer, Y hapten, Le^y found in embryonal carcinoma cells, TL5 (blood group A), EGF receptor found in A431 cells, E₁ series (blood group B) found in pancreatic cancer, FC10.2 found in embryonal carcinoma cells, gastric adenocarcinoma, CO-514 (blood group Le^a) found in adenocarcinoma, NS-10 found in adenocarcinomas, CO-43 (blood group Le^b), G49, EGF receptor, (blood group ALe^b/Le^y) found in colonic adenocarcinoma, 19.9 found in colon cancer, gastric cancer mucins, T₅A₇ found in myeloid cells, R₂₄ found in

melanoma, 4.2, G_{D3}, D1.1, OFA-1, G_{M2}, OFA-2, G_{D2}, M1:22:25:8 found in embryonal carcinoma cells and SSEA-3, SSEA-4 found in 4-8-cell stage embryos. In another embodiment, the antigen is a T cell receptor derived peptide from a cutaneous T cell lymphoma (see Edelson, 1998, *The Cancer Journal* 4:62).

[00271] Cancers and related disorders that can be treated or prevented by methods and compositions of the present invention include, but are not limited to, the following: Leukemias including, but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome, chronic leukemias such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease, non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenström's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors including but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including, but not limited to, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer, including but not limited to, pheochromocytom and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer, including but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers including but not limited to, Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipius; eye cancers including but not limited to, ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers, including but not limited to, squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer, including but not

limited to, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers including but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers including but not limited to, endometrial carcinoma and uterine sarcoma; ovarian cancers including but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers including but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers including but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers including but not limited to hepatocellular carcinoma and hepatoblastoma, gallbladder cancers including but not limited to, adenocarcinoma; cholangiocarcinomas including but not limited to, papillary, nodular, and diffuse; lung cancers including but not limited to, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers including but not limited to, germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers including but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penile cancers; oral cancers including but not limited to, squamous cell carcinoma; basal cancers; salivary gland cancers including but not limited to, adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; pharynx cancers including but not limited to, squamous cell cancer, and verrucous; skin cancers including but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers including but not limited to, renal cell cancer, adenocarcinoma, hypernephroma, fibrosarcoma, transitional cell cancer (renal pelvis and/ or ureter); Wilms' tumor; bladder cancers including but not limited to, transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangi endotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas (for a review of such disorders, see Fishman *et al.*, 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy *et al.*, 1997, *Informed Decisions: The*

Complete Book of Cancer Diagnosis, Treatment, and Recovery, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

[00272] Accordingly, the methods and compositions of the invention are also useful in the treatment or prevention of a variety of cancers or other abnormal proliferative diseases, including (but not limited to) the following: carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, ovary, pancreas, stomach, prostate, cervix, thyroid and skin; including squamous cell carcinoma; hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Burkett's lymphoma; hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; other tumors, including melanoma, seminoma, teratocarcinoma, neuroblastoma and glioma; tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma, and schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including melanoma, xenoderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer and teratocarcinoma. It is also contemplated that cancers caused by aberrations in apoptosis would also be treated by the methods and compositions of the invention. Such cancers may include but not be limited to follicular lymphomas, carcinomas with p53 mutations, hormone dependent tumors of the breast, prostate and ovary, and precancerous lesions such as familial adenomatous polyposis, and myelodysplastic syndromes. In specific embodiments, malignancy or dysproliferative changes (such as metaplasias and dysplasias), or hyperproliferative disorders, are treated or prevented by the methods and compositions of the invention in the ovary, bladder, breast, colon, lung, skin, pancreas, or uterus. In other specific embodiments, sarcoma, melanoma, or leukemia is treated or prevented by the methods and compositions of the invention.

[00273] In a specific embodiment, a molecule of the invention (*e.g.*, a diabody comprising multiple epitope binding domains and, optionally, and Fc domain (or portion thereof)) inhibits or reduces the growth of cancer cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the growth of cancer cells in the absence of said molecule of the invention.

[00274] In a specific embodiment, a molecule of the invention (*e.g.*, a diabody comprising multiple epitope binding domains and, optionally, and Fc domain (or portion

thereof)) kills cells or inhibits or reduces the growth of cancer cells at least 5%, at least 10%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 100% better than the parent molecule.

5.6.2 AUTOIMMUNE DISEASE AND INFLAMMATORY DISEASES

[00275] In some embodiments, molecules of the invention comprise an epitope binding domain specific for FcγRIIB and or/ a variant Fc domain (or portion thereof), engineered according to methods of the invention, which Fc domain exhibits greater affinity for FcγRIIB and decreased affinity for FcγRIIA and/or FcγRII relative to a wild-type Fc domain. Molecules of the invention with such binding characteristics are useful in regulating the immune response, *e.g.*, in inhibiting the immune response in connection with autoimmune diseases or inflammatory diseases. Although not intending to be bound by any mechanism of action, molecules of the invention with an affinity for FcγRIIB and/or comprising an Fc domain with increased affinity for FcγRIIB and a decreased affinity for FcγRIIA and/or FcγRII may lead to dampening of the activating response to FcγR and inhibition of cellular responsiveness, and thus have therapeutic efficacy for treating and/or preventing an autoimmune disorder.

[00276] The invention also provides methods for preventing, treating, or managing one or more symptoms associated with an inflammatory disorder in a subject further comprising, administering to said subject a therapeutically or prophylactically effective amount of one or more anti-inflammatory agents. The invention also provides methods for preventing, treating, or managing one or more symptoms associated with an autoimmune disease further comprising, administering to said subject a therapeutically or prophylactically effective amount of one or more immunomodulatory agents. Section 5.7 provides non-limiting examples of anti-inflammatory agents and immunomodulatory agents.

[00277] Examples of autoimmune disorders that may be treated by administering the molecules of the present invention include, but are not limited to, alopecia areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, Behcet's disease, bullous pemphigoid, cardiomyopathy, celiac sprue-dermatitis, chronic fatigue immune dysfunction syndrome (CFIDS), chronic inflammatory demyelinating polyneuropathy, Churg-Strauss syndrome, cicatricial pemphigoid, CREST syndrome, cold

agglutinin disease, Crohn's disease, discoid lupus, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, glomerulonephritis, Graves' disease, Guillain-Barre, Hashimoto's thyroiditis, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, juvenile arthritis, lichen planus, lupus erythematosus, Ménière's disease, mixed connective tissue disease, multiple sclerosis, type 1 or immune-mediated diabetes mellitus, myasthenia gravis, pemphigus vulgaris, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, Raynaud's phenomenon, Reiter's syndrome, Rheumatoid arthritis, sarcoidosis, scleroderma, Sjögren's syndrome, stiff-man syndrome, systemic lupus erythematosus, lupus erythematosus, takayasu arteritis, temporal arteritis/ giant cell arteritis, ulcerative colitis, uveitis, vasculitides such as dermatitis herpetiformis vasculitis, vitiligo, and Wegener's granulomatosis. Examples of inflammatory disorders include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections. As described herein in Section 2.2.2, some autoimmune disorders are associated with an inflammatory condition. Thus, there is overlap between what is considered an autoimmune disorder and an inflammatory disorder. Therefore, some autoimmune disorders may also be characterized as inflammatory disorders. Examples of inflammatory disorders which can be prevented, treated or managed in accordance with the methods of the invention include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections.

[00278] Molecules of the invention comprising at least one epitope binding domain specific for FcγRIIB and/or a variant Fc domain with an enhanced affinity for FcγRIIB and a decreased affinity for FcγRIIA can also be used to reduce the inflammation experienced by animals, particularly mammals, with inflammatory disorders. In a specific embodiment, a molecule of the invention reduces the inflammation in an animal by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the inflammation in an animal, which is not administered the said molecule.

[00279] Molecules of the invention comprising at least one epitope binding domain specific for FcγRIIB and/or a variant Fc domain with an enhanced affinity for FcγRIIB and a decreased affinity for FcγRIIA can also be used to prevent the rejection of transplants.

5.6.3 INFECTIOUS DISEASE

[00280] The invention also encompasses methods for treating or preventing an infectious disease in a subject comprising administering a therapeutically or prophylactically effective amount of one or more molecules of the invention comprising at least one epitope binding domain specific for an infectious agent associated with said infectious disease. In certain embodiments, the molecules of the invention are toxic to the infectious agent, enhance immune response against said agent or enhance effector function against said agent, relative to the immune response in the absence of said molecule. Infectious diseases that can be treated or prevented by the molecules of the invention are caused by infectious agents including but not limited to viruses, bacteria, fungi, protozoa, and viruses.

[00281] Viral diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention include, but are not limited to, those caused by hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, small pox, Epstein Barr virus, human immunodeficiency virus type I (HIV-I), human immunodeficiency virus type II (HIV-II), and agents of viral diseases such as viral meningitis, encephalitis, dengue or small pox.

[00282] Bacterial diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by bacteria include, but are not limited to, mycobacteria rickettsia, mycoplasma, neisseria, S. pneumonia, Borrelia burgdorferi (Lyme disease), Bacillus anthracis (anthrax), tetanus, streptococcus, staphylococcus, mycobacterium, tetanus, pertissus, cholera, plague, diptheria, chlamydia, S. aureus and legionella.

[00283] Protozoal diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by protozoa include, but are not limited to, leishmania, kokzidioa, trypanosoma or malaria.

[00284] Parasitic diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by parasites include, but are not limited to, chlamydia and rickettsia.

[00285] According to one aspect of the invention, molecules of the invention comprising at least one epitope binding domain specific for an infectious agent exhibit an antibody effector function towards said agent, *e.g.*, a pathogenic protein. Examples of infectious agents include but are not limited to bacteria (*e.g.*, *Escherichia coli*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Enterococcus faecialis*, *Candida albicans*, *Proteus vulgaris*, *Staphylococcus viridans*, and *Pseudomonas aeruginosa*), a pathogen (*e.g.*, B-lymphotropic papovavirus (LPV); Bordatella pertussis; Borna Disease virus (BDV); Bovine coronavirus; Choriomeningitis virus; Dengue virus; a virus, *E. coli*; Ebola; Echovirus 1; Echovirus-11 (EV); Endotoxin (LPS); Enteric bacteria; Enteric Orphan virus; Enteroviruses; Feline leukemia virus; Foot and mouth disease virus; Gibbon ape leukemia virus (GALV); Gram-negative bacteria; *Helicobacter pylori*; Hepatitis B virus (HBV); Herpes Simplex Virus; HIV-1; Human cytomegalovirus; Human coronavirus; Influenza A, B & C; Legionella; Leishmania mexicana; Listeria monocytogenes; Measles virus; Meningococcus; Morbilliviruses; Mouse hepatitis virus; Murine leukemia virus; Murine gamma herpes virus; Murine retrovirus; Murine coronavirus mouse hepatitis virus; Mycobacterium avium-M; Neisseria gonorrhoeae; Newcastle disease virus; Parvovirus B19; Plasmodium falciparum; Pox Virus; Pseudomonas; Rotavirus; Salmonella typhimurium; Shigella; Streptococci; T-cell lymphotropic virus 1; Vaccinia virus).

5.6.4 DETOXIFICATION

[00286] The invention also encompasses methods of detoxification in a subject exposed to a toxin (*e.g.*, a toxic drug molecule) comprising administering a therapeutically or prophylactically effective amount of one or more molecules of the invention comprising at least one epitope binding domain specific for the toxic drug molecule. In certain embodiments, binding of a molecule of the invention to the toxin reduces or eliminates the adverse physiological effect of said toxin. In yet other embodiments, binding of a diabody of the invention to the toxin increases or enhances elimination, degradation or neutralization of the toxin relative to elimination, degradation or neutralization in the absence of said diabody. Immunotoxicotherapy in accordance with the methods of the invention can be used to treat overdoses or exposure to drugs including, but not limited to, digoxin, PCP, cocaine, colchicine, and tricyclic antidepressants.

5.7 COMBINATION THERAPY

[00287] The invention further encompasses administering the molecules of the invention in combination with other therapies known to those skilled in the art for the treatment or prevention of cancer, autoimmune disease, infectious disease or intoxication,

including but not limited to, current standard and experimental chemotherapies, hormonal therapies, biological therapies, immunotherapies, radiation therapies, or surgery. In some embodiments, the molecules of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or more agents, therapeutic antibodies or other agents known to those skilled in the art for the treatment and/or prevention of cancer, autoimmune disease, infectious disease or intoxication.

[00288] In certain embodiments, one or more molecule of the invention is administered to a mammal, preferably a human, concurrently with one or more other therapeutic agents useful for the treatment of cancer. The term "concurrently" is not limited to the administration of prophylactic or therapeutic agents at exactly the same time, but rather it is meant that a molecule of the invention and the other agent are administered to a mammal in a sequence and within a time interval such that the molecule of the invention can act together with the other agent to provide an increased benefit than if they were administered otherwise. For example, each prophylactic or therapeutic agent (*e.g.*, chemotherapy, radiation therapy, hormonal therapy or biological therapy) may be administered at the same time or sequentially in any order at different points in time; however, if not administered at the same time, they should be administered sufficiently close in time so as to provide the desired therapeutic or prophylactic effect. Each therapeutic agent can be administered separately, in any appropriate form and by any suitable route. In various embodiments, the prophylactic or therapeutic agents are administered less than 1 hour apart, at about 1 hour apart, at about 1 hour to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, no more than 24 hours apart or no more than 48 hours apart. In preferred embodiments, two or more components are administered within the same patient visit.

[00289] In other embodiments, the prophylactic or therapeutic agents are administered at about 2 to 4 days apart, at about 4 to 6 days apart, at about 1 week part, at about 1 to 2 weeks apart, or more than 2 weeks apart. In preferred embodiments, the prophylactic or therapeutic agents are administered in a time frame where both agents are still active. One skilled in the art would be able to determine such a time frame by determining the half life of the administered agents.

[00290] In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a subject. Cycling therapy involves the administration of a first agent for a period of time, followed by the administration of a second agent and/or third agent for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

[00291] In certain embodiments, prophylactic or therapeutic agents are administered in a cycle of less than about 3 weeks, about once every two weeks, about once every 10 days or about once every week. One cycle can comprise the administration of a therapeutic or prophylactic agent by infusion over about 90 minutes every cycle, about 1 hour every cycle, about 45 minutes every cycle. Each cycle can comprise at least 1 week of rest, at least 2 weeks of rest, at least 3 weeks of rest. The number of cycles administered is from about 1 to about 12 cycles, more typically from about 2 to about 10 cycles, and more typically from about 2 to about 8 cycles.

[00292] In yet other embodiments, the therapeutic and prophylactic agents of the invention are administered in metronomic dosing regimens, either by continuous infusion or frequent administration without extended rest periods. Such metronomic administration can involve dosing at constant intervals without rest periods. Typically the therapeutic agents, in particular cytotoxic agents, are used at lower doses. Such dosing regimens encompass the chronic daily administration of relatively low doses for extended periods of time. In preferred embodiments, the use of lower doses can minimize toxic side effects and eliminate rest periods. In certain embodiments, the therapeutic and prophylactic agents are delivered by chronic low-dose or continuous infusion ranging from about 24 hours to about 2 days, to about 1 week, to about 2 weeks, to about 3 weeks to about 1 month to about 2 months, to about 3 months, to about 4 months, to about 5 months, to about 6 months. The scheduling of such dose regimens can be optimized by the skilled oncologist.

[00293] In other embodiments, courses of treatment are administered concurrently to a mammal, *i.e.*, individual doses of the therapeutics are administered separately yet within a time interval such that molecules of the invention can work together with the other agent or agents. For example, one component may be administered one time per week in combination with the other components that may be administered one time every two weeks or one time every three weeks. In other words, the dosing regimens for the therapeutics are carried out concurrently even if the therapeutics are not administered simultaneously or within the same patient visit.

[00294] When used in combination with other prophylactic and/or therapeutic agents, the molecules of the invention and the prophylactic and/or therapeutic agent can act additively or, more preferably, synergistically. In one embodiment, a molecule of the invention is administered concurrently with one or more therapeutic agents in the same pharmaceutical composition. In another embodiment, a molecule of the invention is administered concurrently with one or more other therapeutic agents in separate pharmaceutical compositions. In still another embodiment, a molecule of the invention is administered prior to or subsequent to administration of another prophylactic or therapeutic agent. The invention contemplates administration of a molecule of the invention in combination with other prophylactic or therapeutic agents by the same or different routes of administration, *e.g.*, oral and parenteral. In certain embodiments, when a molecule of the invention is administered concurrently with another prophylactic or therapeutic agent that potentially produces adverse side effects including, but not limited to, toxicity, the prophylactic or therapeutic agent can advantageously be administered at a dose that falls below the threshold that the adverse side effect is elicited.

[00295] The dosage amounts and frequencies of administration provided herein are encompassed by the terms therapeutically effective and prophylactically effective. The dosage and frequency further will typically vary according to factors specific for each patient depending on the specific therapeutic or prophylactic agents administered, the severity and type of cancer, the route of administration, as well as age, body weight, response, and the past medical history of the patient. Suitable regimens can be selected by one skilled in the art by considering such factors and by following, for example, dosages reported in the literature and recommended in the *Physician's Desk Reference* (56th ed., 2002).

5.7.1 ANTI-CANCER AGENTS

[00296] In a specific embodiment, the methods of the invention encompass the administration of one or more molecules of the invention with one or more therapeutic agents used for the treatment and/or prevention of cancer. In one embodiment, angiogenesis inhibitors may be administered in combination with the molecules of the invention. Angiogenesis inhibitors that can be used in the methods and compositions of the invention include but are not limited to: Angiostatin (plasminogen fragment); antiangiogenic antithrombin III; Angiozyme; ABT-627; Bay 12-9566; Benefin; Bevacizumab; BMS-275291; cartilage-derived inhibitor (CDI); CAI; CD59 complement fragment; CEP-7055; Col 3; Combretastatin A-4; Endostatin (collagen XVIII fragment); Fibronectin fragment;

Gro-beta; Halofuginone; Heparinases; Heparin hexasaccharide fragment; HMV833; Human chorionic gonadotropin (hCG); IM-862; Interferon alpha/beta/gamma; Interferon inducible protein (IP-10); Interleukin-12; Kringle 5 (plasminogen fragment); Marimastat; Metalloproteinase inhibitors (TIMPs); 2-Methoxyestradiol; MMI 270 (CGS 27023A); MoAb IMC-1C11; Neovastat; NM-3; Panzem; PI-88; Placental ribonuclease inhibitor; Plasminogen activator inhibitor; Platelet factor-4 (PF4); Prinomastat; Prolactin 16kDa fragment; Proliferin-related protein (PRP); PTK 787/ZK 222594; Retinoids; Solimastat; Squalamine; SS 3304; SU 5416; SU6668; SU11248; Tetrahydrocortisol-S; tetrathiomolybdate; thalidomide; Thrombospondin-1 (TSP-1); TNP-470; Transforming growth factor-beta (TGF-b); Vasculostatin; Vasostatin (calreticulin fragment); ZD6126; ZD 6474; farnesyl transferase inhibitors (FTI); and bisphosphonates.

[00297] Anti-cancer agents that can be used in combination with the molecules of the invention in the various embodiments of the invention, including pharmaceutical compositions and dosage forms and kits of the invention, include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziqune; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; interleukin II (including recombinant interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1 ; interferon alfa-n3; interferon beta-I a; interferon gamma-I b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocil; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol

acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedapa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricirbine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride. Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D₃; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; ansacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorlins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8;

cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam; cypemycin;
 cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrodidemnin
 B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone;
 didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-;
 dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine;
 droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab;
 eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen
 agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole;
 fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone;
 fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin;
 fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase
 inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene
 bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofofosine;
 ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth
 factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane;
 iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B;
 itasetron; jaspilakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin;
 lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte
 alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole;
 linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds;
 lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin;
 loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine;
 mannostatin A; marimastat; masoprocil; maspin; matrilysin inhibitors; matrix
 metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase;
 metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double
 stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin
 fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal
 antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell
 wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor
 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall
 extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip;
 naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin;
 neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators;
 nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides;
 onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin;

osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer. Preferred additional anti-cancer drugs are 5-fluorouracil and leucovorin.

[00298] Examples of therapeutic antibodies that can be used in methods of the invention include but are not limited to ZENAPAX® (daclizumab) (Roche Pharmaceuticals,

Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody for the prevention of acute renal allograft rejection; PANOREX™ which is a murine anti-17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); BEC2 which is a murine anti-idiotypic (GD3 epitope) IgG antibody (ImClone System); IMC-C225 which is a chimeric anti-EGFR IgG antibody (ImClone System); VITAXIN™ which is a humanized anti- α V β 3 integrin antibody (Applied Molecular Evolution/MedImmune); Smart M195 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo); LYMPHOCIDE™ which is a humanized anti-CD22 IgG antibody (Immunomedics); ICM3 is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primatized anti-CD80 antibody (IDEC Pharm/Mitsubishi); IDEC-131 is a humanized anti-CD40L antibody (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5) antibody (Alexion Pharm); D2E7 is a humanized anti-TNF- α antibody (CAT/BASF); CDP870 is a humanized anti-TNF- α Fab fragment (Celltech); IDEC-151 is a primatized anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-CD4 IgG antibody (Medarex/Eisai/Genmab); CDP571 is a humanized anti-TNF- α IgG4 antibody (Celltech); LDP-02 is a humanized anti- α 4 β 7 antibody (LeukoSite/Genentech); OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA™ is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN™ is a humanized anti-VLA-4 IgG antibody (Elan); and CAT-152 is a human anti-TGF- β 2 antibody (Cambridge Ab Tech). Other examples of therapeutic antibodies that can be used in accordance with the invention are presented in Table 8.

Table 8: Anti-cancer therapeutic antibodies

Company	Product	Disease	Target
Abgenix	ABX-EGF	Cancer	EGF receptor
AltaRex	OvaRex	ovarian cancer	tumor antigen CA125
	BravaRex	metastatic cancers	tumor antigen MUC1
Antisoma	Theragyn (pemtumomabytrrium-90)	ovarian cancer	PEM antigen
	Therex	breast cancer	PEM antigen
Boehringer Ingelheim	Blvatuzumab	head & neck cancer	CD44

Company	Product	Disease	Target
Centocor/J&J	Panorex	Colorectal cancer	17-1A
	ReoPro	PTCA	gp IIIb/IIIa
	ReoPro	Acute MI	gp IIIb/IIIa
	ReoPro	Ischemic stroke	gp IIIb/IIIa
Corixa	Bexocar	NHL	CD20
CRC Technology	MAB, idiotypic 105AD7	colorectal cancer vaccine	gp72
Crucell	Anti-EpCAM	cancer	Ep-CAM
Cytoclonal	MAB, lung cancer	non-small cell lung cancer	NA
Genentech	Herceptin	metastatic breast cancer	HER-2
	Herceptin	early stage breast cancer	HER-2
	Rituxan	Relapsed/refractory low-grade or follicular NHL	CD20
	Rituxan	intermediate & high-grade NHL	CD20
	MAB-VEGF	NSCLC, metastatic	VEGF
	MAB-VEGF	Colorectal cancer, metastatic	VEGF
	AMD Fab	age-related macular degeneration	CD18
IDEC	E-26 (2 nd gen. IgE)	allergic asthma & rhinitis	IgE
	Zevalin (Rituxan + yttrium-90)	low grade of follicular, relapsed or refractory, CD20-positive, B-cell NHL and Rituximab-refractory NHL	CD20
ImClone	Cetuximab + innotecan	refractory colorectal carcinoma	EGF receptor
	Cetuximab + cisplatin &	newly diagnosed or recurrent head	EGF receptor

Company	Product	Disease	Target
	radiation	& neck cancer	
	Cetuximab + gemcitabine	newly diagnosed metastatic pancreatic carcinoma	EGF receptor
	Cetuximab + cisplatin + 5FU or Taxol	recurrent or metastatic head & neck cancer	EGF receptor
	Cetuximab + carboplatin + paclitaxel	newly diagnosed non-small cell lung carcinoma	EGF receptor
	Cetuximab + cisplatin	head & neck cancer (extensive incurable local- regional disease & distant metastases)	EGF receptor
	Cetuximab + radiation	locally advanced head & neck carcinoma	EGF receptor
	BEC2 + Bacillus Calmette Guerin	small cell lung carcinoma	mimics ganglioside GD3
	BEC2 + Bacillus Calmette Guerin	melanoma	mimics ganglioside GD3
	IMC-1C11	colorectal cancer with liver metastases	VEGF-receptor
ImmonoGen	nuC242-DM1	Colorectal, gastric, and pancreatic cancer	nuC242
ImmunoMedics	LymphoCide	Non-Hodgkins lymphoma	CD22
	LymphoCide Y-90	Non-Hodgkins lymphoma	CD22
	CEA-Cide	metastatic solid tumors	CEA
	CEA-Cide Y-90	metastatic solid tumors	CEA
	CEA-Scan (Tc-99m- labeled arcitumomab)	colorectal cancer (radioimaging)	CEA
	CEA-Scan (Tc-99m- labeled arcitumomab)	Breast cancer (radioimaging)	CEA

Company	Product	Disease	Target
	CEA-Scan (Tc-99m-labeled arcitumomab)	lung cancer (radioimaging)	CEA
	CEA-Scan (Tc-99m-labeled arcitumomab)	intraoperative tumors (radio imaging)	CEA
	LeukoScan (Tc-99m-labeled sulesomab)	soft tissue infection (radioimaging)	CEA
	LymphoScan (Tc-99m-labeled)	lymphomas (radioimaging)	CD22
	AFP-Scan (Tc-99m-labeled)	liver 7 gem-cell cancers (radioimaging)	AFP
Intracel	HumaRAD-HN (+ yttrium-90)	head & neck cancer	NA
	HumaSPECT	colorectal imaging	NA
Medarex	MDX-101 (CTLA-4)	Prostate and other cancers	CTLA-4
	MDX-210 (her-2 overexpression)	Prostate cancer	HER-2
	MDX-210/MAK	Cancer	HER-2
MedImmune	Vitaxin	Cancer	$\alpha v \beta_3$
Merck KGaA	MAb 425	Various cancers	EGF receptor
	IS-IL-2	Various cancers	Ep-CAM
Millennium	Campath (alemtuzumab)	chronic lymphocytic leukemia	CD52
NeoRx	CD20-streptavidin (+ biotin-yttrium 90)	Non-Hodgkins lymphoma	CD20
	Avidicin (albumin + NRLU13)	metastatic cancer	NA
Peregrine	Oncolym (+ iodine-131)	Non-Hodgkins lymphoma	HLA-DR 10 beta
	Cotara (+ iodine-131)	unresectable malignant glioma	DNA-associated proteins
Pharmacia Corporation	C215 (+ staphylococcal enterotoxin)	pancreatic cancer	NA
	MAb, lung/kidney cancer	lung & kidney cancer	NA
	nacolomab tafenatox	colon &	NA

Company	Product	Disease	Target
	(C242 + staphylococcal enterotoxin)	pancreatic cancer	
Protein Design Labs	Nuvion	T cell malignancies	CD3
	SMART M195	AML	CD33
	SMART 1D10	NHL	HLA-DR antigen
Titan	CEAVac	colorectal cancer, advanced	CEA
	TriGem	metastatic melanoma & small cell lung cancer	GD2-ganglioside
	TriAb	metastatic breast cancer	MUC-1
Trilex	CEAVac	colorectal cancer, advanced	CEA
	TriGem	metastatic melanoma & small cell lung cancer	GD2-ganglioside
	TriAb	metastatic breast cancer	MUC-1
Viventia Biotech	NovoMAb-G2 radiolabeled	Non-Hodgkins lymphoma	NA
	Monopharm C	colorectal & pancreatic carcinoma	SK-1 antigen
	GlioMAb-H (+ gelonin toxin)	glioma, melanoma & neuroblastoma	NA
Xoma	Rituxan	Relapsed/refractory low-grade or follicular NHL	CD20
	Rituxan	intermediate & high-grade NHL	CD20
	ING-1	adenomcarcinoma	Ep-CAM

5.7.2 IMMUNOMODULATORY AGENTS AND ANTI-INFLAMMATORY AGENTS

[00299] The present invention provides methods of treatment for autoimmune diseases and inflammatory diseases comprising administration of the molecules of the invention in conjunction with other treatment agents. Examples of immunomodulatory agents include, but are not limited to, methothrexate, ENBREL, REMICADE™, leflunomide, cyclophosphamide, cyclosporine A, and macrolide antibiotics (e.g., FK506 (tacrolimus)), methylprednisolone (MP), corticosteroids, steroids, mycophenolate mofetil, rapamycin (sirolimus), mizoribine, deoxyspergualin, brequinar, malononitriloamindes (e.g., leflunamide), T cell receptor modulators, and cytokine receptor modulators.

[00300] Anti-inflammatory agents have exhibited success in treatment of inflammatory and autoimmune disorders and are now a common and a standard treatment for such disorders. Any anti-inflammatory agent well-known to one of skill in the art can be used in the methods of the invention. Non-limiting examples of anti-inflammatory agents include non-steroidal anti-inflammatory drugs (NSAIDs), steroidal anti-inflammatory drugs, beta-agonists, anticholinergic agents, and methyl xanthines. Examples of NSAIDs include, but are not limited to, aspirin, ibuprofen, celecoxib (CELEBREX™), diclofenac (VOLTAREN™), etodolac (LODINE™), fenoprofen (NALFON™), indomethacin (INDOCIN™), ketoralac (TORADOL™), oxaprozin (DAYPRO™), nabumentone (RELAFEN™), sulindac (CLINORIL™), tolmentin (TOLECTIN™), rofecoxib (VIOXX™), naproxen (ALEVE™, NAPROSYN™), ketoprofen (ACTRON™) and nabumetone (RELAFEN™). Such NSAIDs function by inhibiting a cyclooxygenase enzyme (e.g., COX-1 and/or COX-2). Examples of steroidal anti-inflammatory drugs include, but are not limited to, glucocorticoids, dexamethasone (DECADRON™), cortisone, hydrocortisone, prednisone (DELTASONE™), prednisolone, triamcinolone, azulfidine, and eicosanoids such as prostaglandins, thromboxanes, and leukotrienes.

[00301] A non-limiting example of the antibodies that can be used for the treatment or prevention of inflammatory disorders in conjunction with the molecules of the invention is presented in Table 9, and a non-limiting example of the antibodies that can be used for the treatment or prevention of autoimmune disorder is presented in Table 10.

Table 9: Therapeutic antibodies for the treatment of inflammatory diseases

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	Rheumatoid Arthritis

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	SLE
5G1.1	Complement (C5)	Humanized	IgG	Alexion Pharm Inc	Nephritis
5G1.1-SC	Complement (C5)	Humanized	ScFv	Alexion Pharm Inc	Cardiopulmonary Bypass
5G1.1-SC	Complement (C5)	Humanized	ScFv	Alexion Pharm Inc	Myocardial Infarction
5G1.1-SC	Complement (C5)	Humanized	ScFv	Alexion Pharm Inc	Angioplasty
ABX-CBL	CBL	Human		Abgenix Inc	GvHD
ABX-CBL	CD147	Murine	IgG	Abgenix Inc	Allograft rejection
ABX-IL8	IL-8	Human	IgG2	Abgenix Inc	Psoriasis
Antegren	VLA-4	Humanized	IgG	Athena/Elan	Multiple Sclerosis
Anti-CD11a	CD11a	Humanized	IgG1	Genentech Inc/Xoma	Psoriasis
Anti-CD18	CD18	Humanized	Fab'2	Genentech Inc	Myocardial infarction
Anti-LFA1	CD18	Murine	Fab'2	Pasteur-Merieux/ Immunotech	Allograft rejection
Antova	CD40L	Humanized	IgG	Biogen	Allograft rejection
Antova	CD40L	Humanized	IgG	Biogen	SLE
BTI-322	CD2	Rat	IgG	Medimmune Inc	GvHD, Psoriasis
CDP571	TNF-alpha	Humanized	IgG4	Celltech	Crohn's
CDP571	TNF-alpha	Humanized	IgG4	Celltech	Rheumatoid Arthritis
CDP850	E-selectin	Humanized		Celltech	Psoriasis
Corsevin M	Fact VII	Chimeric		Centocor	Anticoagulant
D2E7	TNF-alpha	Human		CAT/BASF	Rheumatoid Arthritis
Hu23F2G	CD11/18	Humanized		ICOS Pharm Inc	Multiple Sclerosis
Hu23F2G	CD11/18	Humanized	IgG	ICOS Pharm Inc	Stroke
IC14	CD14			ICOS Pharm Inc	Toxic shock

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
ICM3	ICAM-3	Humanized		ICOS Pharm Inc	Psoriasis
IDEC-114	CD80	Primatised		IDEC Pharm/Mitsubishi	Psoriasis
IDEC-131	CD40L	Humanized		IDEC Pharm/Eisai	SLE
IDEC-131	CD40L	Humanized		IDEC Pharm/Eisai	Multiple Sclerosis
IDEC-151	CD4	Primatised	IgG1	IDEC Pharm/Glaxo SmithKline	Rheumatoid Arthritis
IDEC-152	CD23	Primatised		IDEC Pharm	Asthma/Allergy
Infliximab	TNF-alpha	Chimeric	IgG1	Centocor	Rheumatoid Arthritis
Infliximab	TNF-alpha	Chimeric	IgG1	Centocor	Crohn's
LDP-01	beta2-integrin	Humanized	IgG	Millennium Inc (LeukoSite Inc.)	Stroke
LDP-01	beta2-integrin	Humanized	IgG	Millennium Inc (LeukoSite Inc.)	Allograft rejection
LDP-02	alpha4beta7	Humanized		Millennium Inc (LeukoSite Inc.)	Ulcerative Colitis
MAK-195F	TNF alpha	Murine	Fab'2	Knoll Pharm, BASF	Toxic shock
MDX-33	CD64 (FcR)	Human		Medarex/Centcon	Autoimmune haematological disorders
MDX-CD4	CD4	Human	IgG	Medarex/Eisai / Genmab	Rheumatoid Arthritis
MEDI-507	CD2	Humanized		Medimmune Inc	Psoriasis
MEDI-507	CD2	Humanized		Medimmune Inc	GvHD
OKT4A	CD4	Humanized	IgG	Ortho Biotech	Allograft rejection

Antibody Name	Target Antigen	Product Type	Isotype	Sponsors	Indication
Orthoclone OKT4A	CD4	Humanized	IgG	Ortho Biotech	Autoimmune disease
Orthoclone/anti-CD3 OKT3	CD3	Murine	mIgG2a	Ortho Biotech	Allograft rejection
RepPro/Abciximab	gpIIbIIIa	Chimeric	Fab	Centocor/Lilly	Complications of coronary angioplasty
rhuMab-E25	IgE	Humanized	IgG1	Genentech/Novartis/Tanox Biosystems	Asthma/Allergy
SB-240563	IL5	Humanized		GlaxoSmithKline	Asthma/Allergy
SB-240683	IL-4	Humanized		GlaxoSmithKline	Asthma/Allergy
SCH55700	IL-5	Humanized		Celltech/Schering	Asthma/Allergy
Simulect	CD25	Chimeric	IgG1	Novartis Pharm	Allograft rejection
SMART a-CD3	CD3	Humanized		Protein Design Lab	Autoimmune disease
SMART a-CD3	CD3	Humanized		Protein Design Lab	Allograft rejection
SMART a-CD3	CD3	Humanized	IgG	Protein Design Lab	Psoriasis
Zenapax	CD25	Humanized	IgG1	Protein Design Lab/Hoffman-La Roche	Allograft rejection

Table 10: Therapeutic antibodies for the treatment of autoimmune disorders

Antibody	Indication	Target Antigen
ABX-RB2		antibody to CBL antigen on T cells, B cells and NK cells fully human antibody from the Xenomouse
5c8 (Anti CD-40 antigen antibody)	Phase II trials were halted in Oct. 99 examine "adverse events"	CD-40
IDEC 131	systemic lupus erythematosus (SLE)	anti CD40 humanized

Antibody	Indication	Target Antigen
IDEC 151	rheumatoid arthritis	primatized ; anti-CD4
IDEC 152	Asthma	primatized; anti-CD23
IDEC 114	Psoriasis	primatized anti-CD80
MEDI-507	rheumatoid arthritis; multiple sclerosis Crohn's disease Psoriasis	anti-CD2
LDP-02 (anti-b7 mAb)	inflammatory bowel disease Chron's disease ulcerative colitis	a4b7 integrin receptor on white blood cells (leukocytes)
SMART Anti-Gamma Interferon antibody	autoimmune disorders	Anti-Gamma Interferon
Verteportin	rheumatoid arthritis	
MDX-33	blood disorders caused by autoimmune reactions Idiopathic Thrombocytopenia Purpura (ITP) autoimmune hemolytic anemia	monoclonal antibody against FcRI receptors
MDX-CD4	treat rheumatoid arthritis and other autoimmunity	monoclonal antibody against CD4 receptor molecule
VX-497	autoimmune disorders multiple sclerosis rheumatoid arthritis inflammatory bowel disease lupus psoriasis	inhibitor of inosine monophosphate dehydrogenase (enzyme needed to make new RNA and DNA used in production of nucleotides needed for lymphocyte proliferation)
VX-740	rheumatoid arthritis	inhibitor of ICE interleukin-1 beta (converting enzyme controls pathways leading to aggressive immune response)
VX-745	specific to inflammation involved in chemical signalling of immune response onset and progression of inflammation	inhibitor of P38MAP kinase mitogen activated protein kinase
Enbrel (etanercept)		targets TNF (tumor necrosis factor)

Antibody	Indication	Target Antigen
IL-8		fully human monoclonal antibody against IL-8 (interleukin 8)
Apogen MP4		recombinant antigen selectively destroys disease associated T-cells induces apoptosis T-cells eliminated by programmed cell death no longer attack body's own cells specific apogens target specific T-cells

5.7.3 AGENTS FOR USE IN THE TREATMENT OF INFECTIOUS DISEASE

[00302] In some embodiments, the molecules of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or additional therapeutic agents known to those skilled in the art for the treatment and/or prevention of an infectious disease. The invention contemplates the use of the molecules of the invention in combination with antibiotics known to those skilled in the art for the treatment and or prevention of an infectious disease. Antibiotics that can be used in combination with the molecules of the invention include, but are not limited to, macrolide (*e.g.*, tobramycin (Tobi®)), a cephalosporin (*e.g.*, cephalexin (Keflex®), cephadrine (Velosef®), cefuroxime (Ceftin®), cefprozil (Cefzil®), cefaclor (Ceclor®), cefixime (Suprax®) or cefadroxil (Duricef®)), a clarithromycin (*e.g.*, clarithromycin (Biaxin®)), an erythromycin (*e.g.*, erythromycin (EMycin®)), a penicillin (*e.g.*, penicillin V (V-Cillin K® or Pen Vee K®)) or a quinolone (*e.g.*, ofloxacin (Floxin®), ciprofloxacin (Cipro®) or norfloxacin (Noroxin®)), aminoglycoside antibiotics (*e.g.*, apramycin, arbekacin, bambarmycins, butirosin, dibekacin, neomycin, neomycin, undecylenate, netilmicin, paromomycin, ribostamycin, sisomicin, and spectinomycin), amphenicol antibiotics (*e.g.*, azidamfenicol, chloramphenicol, florfenicol, and thiamphenicol), ansamycin antibiotics (*e.g.*, rifamide and rifampin), carbacephems (*e.g.*, loracarbef), carbapenems (*e.g.*, biapenem and imipenem), cephalosporins (*e.g.*, cefaclor, cefadroxil, cefamandole, cefatrizine, cefazedone, cefozopran, cefpimizole, cefpiramide, and cefpirome), cephamycins (*e.g.*, cefbuperazone, cefmetazole, and cefminox), monobactams (*e.g.*, aztreonam, carumonam, and tigemonam), oxacephems (*e.g.*, flomoxef, and moxalactam), penicillins (*e.g.*, amdinocillin, amdinocillin pivoxil, amoxicillin, bacampicillin, benzylpenicillinic acid,

benzylpenicillin sodium, epicillin, fenbenicillin, floxacillin, penamccillin, penethamate hydriodide, penicillin o-benethamine, penicillin O, penicillin V, penicillin V benzathine, penicillin V hydrabamine, penimepicycline, and phencihicillin potassium), lincosamides (e.g., clindamycin, and lincomycin), amphomycin, bacitracin, capreomycin, colistin, enduracidin, enviomycin, tetracyclines (e.g., apicycline, chlortetracycline, clomocycline, and demeclocycline), 2,4-diaminopyrimidines (e.g., brodimoprim), nitrofurans (e.g., furaltadone, and furazolium chloride), quinolones and analogs thereof (e.g., cinoxacin,, clinafloxacin, flumequine, and grepagloxacin), sulfonamides (e.g., acetyl sulfamethoxypyrazine, benzylsulfamide, noprylsulfamide, phthalylsulfacetamide, sulfachrysoidine, and sulfacytine), sulfones (e.g., diathymosulfone, glucosulfone sodium, and solasulfone), cycloserine, mupirocin and tuberin.

[00303] In certain embodiments, the molecules of the invention can be administered in combination with a therapeutically or prophylactically effective amount of one or more antifungal agents. Antifungal agents that can be used in combination with the molecules of the invention include but are not limited to amphotericin B, itraconazole, ketoconazole, fluconazole, intrathecal, flucytosine, miconazole, butoconazole, clotrimazole, nystatin, terconazole, tioconazole, ciclopirox, econazole, haloprogrin, naftifine, terbinafine, undecylenate, and griseofuldin.

[00304] In some embodiments, the molecules of the invention can be administered in combination with a therapeutically or prophylactically effective amount of one or more anti-viral agent. Useful anti-viral agents that can be used in combination with the molecules of the invention include, but are not limited to, protease inhibitors, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors and nucleoside analogs. Examples of antiviral agents include but are not limited to zidovudine, acyclovir, gangcyclovir, vidarabine, idoxuridine, trifluridine, and ribavirin, as well as foscarnet, amantadine, rimantadine, saquinavir, indinavir, amprenavir, lopinavir, ritonavir, the alpha-interferons; adefovir, clevadine, entecavir, pleconaril.

5.8 VACCINE THERAPY

[00305] The invention further encompasses using a composition of the invention to induce an immune response against an antigenic or immunogenic agent, including but not limited to cancer antigens and infectious disease antigens (examples of which are disclosed *infra*). The vaccine compositions of the invention comprise one or more antigenic or immunogenic agents to which an immune response is desired, wherein the one or more antigenic or immunogenic agents is coated with a variant antibody of the invention that has

an enhanced affinity to FcγRIIIA. The vaccine compositions of the invention are particularly effective in eliciting an immune response, preferably a protective immune response against the antigenic or immunogenic agent.

[00306] In some embodiments, the antigenic or immunogenic agent in the vaccine compositions of the invention comprise a virus against which an immune response is desired. The viruses may be recombinant or chimeric, and are preferably attenuated. Production of recombinant, chimeric, and attenuated viruses may be performed using standard methods known to one skilled in the art. The invention encompasses a live recombinant viral vaccine or an inactivated recombinant viral vaccine to be formulated in accordance with the invention. A live vaccine may be preferred because multiplication in the host leads to a prolonged stimulus of similar kind and magnitude to that occurring in natural infections, and therefore, confers substantial, long-lasting immunity. Production of such live recombinant virus vaccine formulations may be accomplished using conventional methods involving propagation of the virus in cell culture or in the allantois of the chick embryo followed by purification.

[00307] In a specific embodiment, the recombinant virus is non-pathogenic to the subject to which it is administered. In this regard, the use of genetically engineered viruses for vaccine purposes may require the presence of attenuation characteristics in these strains. The introduction of appropriate mutations (*e.g.*, deletions) into the templates used for transfection may provide the novel viruses with attenuation characteristics. For example, specific missense mutations which are associated with temperature sensitivity or cold adaptation can be made into deletion mutations. These mutations should be more stable than the point mutations associated with cold or temperature sensitive mutants and reversion frequencies should be extremely low. Recombinant DNA technologies for engineering recombinant viruses are known in the art and encompassed in the invention. For example, techniques for modifying negative strand RNA viruses are known in the art, see, *e.g.*, U.S. Patent No. 5,166,057, which is incorporated herein by reference in its entirety.

[00308] Alternatively, chimeric viruses with "suicide" characteristics may be constructed for use in the intradermal vaccine formulations of the invention. Such viruses would go through only one or a few rounds of replication within the host. When used as a vaccine, the recombinant virus would go through limited replication cycle(s) and induce a sufficient level of immune response but it would not go further in the human host and cause disease. Alternatively, inactivated (killed) virus may be formulated in accordance with the invention. Inactivated vaccine formulations may be prepared using conventional techniques to "kill" the chimeric viruses. Inactivated vaccines are "dead" in the sense that their

infectivity has been destroyed. Ideally, the infectivity of the virus is destroyed without affecting its immunogenicity. In order to prepare inactivated vaccines, the chimeric virus may be grown in cell culture or in the allantois of the chick embryo, purified by zonal ultracentrifugation, inactivated by formaldehyde or β -propiolactone, and pooled.

[00309] In certain embodiments, completely foreign epitopes, including antigens derived from other viral or non-viral pathogens can be engineered into the virus for use in the intradermal vaccine formulations of the invention. For example, antigens of non-related viruses such as HIV (gp160, gp120, gp41) parasite antigens (*e.g.*, malaria), bacterial or fungal antigens or tumor antigens can be engineered into the attenuated strain.

[00310] Virtually any heterologous gene sequence may be constructed into the chimeric viruses of the invention for use in the intradermal vaccine formulations. Preferably, heterologous gene sequences are moieties and peptides that act as biological response modifiers. Preferably, epitopes that induce a protective immune response to any of a variety of pathogens, or antigens that bind neutralizing antibodies may be expressed by or as part of the chimeric viruses. For example, heterologous gene sequences that can be constructed into the chimeric viruses of the invention include, but are not limited to, influenza and parainfluenza hemagglutinin neuraminidase and fusion glycoproteins such as the HN and F genes of human PIV3. In yet another embodiment, heterologous gene sequences that can be engineered into the chimeric viruses include those that encode proteins with immuno-modulating activities. Examples of immuno-modulating proteins include, but are not limited to, cytokines, interferon type 1, gamma interferon, colony stimulating factors, interleukin -1, -2, -4, -5, -6, -12, and antagonists of these agents.

[00311] In yet other embodiments, the invention encompasses pathogenic cells or viruses, preferably attenuated viruses, which express the variant antibody on their surface.

[00312] In alternative embodiments, the vaccine compositions of the invention comprise a fusion polypeptide wherein an antigenic or immunogenic agent is operatively linked to a variant antibody of the invention that has an enhanced affinity for Fc γ RIIIA. Engineering fusion polypeptides for use in the vaccine compositions of the invention is performed using routine recombinant DNA technology methods and is within the level of ordinary skill.

[00313] The invention further encompasses methods to induce tolerance in a subject by administering a composition of the invention. Preferably a composition suitable for inducing tolerance in a subject, comprises an antigenic or immunogenic agent coated with a variant antibody of the invention, wherein the variant antibody has a higher affinity to Fc γ RIIB. Although not intending to be bound by a particular mechanism of action, such

compositions are effective in inducing tolerance by activating the Fc γ RIIB mediated inhibitory pathway.

5.9 COMPOSITIONS AND METHODS OF ADMINISTERING

[00314] The invention provides methods and pharmaceutical compositions comprising molecules of the invention (*i.e.*, diabodies) comprising multiple epitope binding domains and, optionally, an Fc domain (or portion thereof). The invention also provides methods of treatment, prophylaxis, and amelioration of one or more symptoms associated with a disease, disorder or infection by administering to a subject an effective amount of a fusion protein or a conjugated molecule of the invention, or a pharmaceutical composition comprising a fusion protein or a conjugated molecule of the invention. In a preferred aspect, an antibody, a fusion protein, or a conjugated molecule, is substantially purified (*i.e.*, substantially free from substances that limit its effect or produce undesired side-effects). In a specific embodiment, the subject is an animal, preferably a mammal such as non-primate (*e.g.*, cows, pigs, horses, cats, dogs, rats *etc.*) and a primate (*e.g.*, monkey such as, a cynomolgous monkey and a human). In a preferred embodiment, the subject is a human. In yet another preferred embodiment, the antibody of the invention is from the same species as the subject.

[00315] Various delivery systems are known and can be used to administer a composition comprising molecules of the invention, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the antibody or fusion protein, receptor-mediated endocytosis (*See, e.g.*, Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432), construction of a nucleic acid as part of a retroviral or other vector, *etc.* Methods of administering a molecule of the invention include, but are not limited to, parenteral administration (*e.g.*, intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidural, and mucosal (*e.g.*, intranasal and oral routes). In a specific embodiment, the molecules of the invention are administered intramuscularly, intravenously, or subcutaneously. The compositions may be administered by any convenient route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, *etc.*) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, pulmonary administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. *See, e.g.*, U.S. Patent Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013;

WO 98/31346; and WO 99/66903, each of which is incorporated herein by reference in its entirety.

[00316] The invention also provides that the molecules of the invention, are packaged in a hermetically sealed container such as an ampoule or sachette indicating the quantity of antibody. In one embodiment, the molecules of the invention are supplied as a dry sterilized lyophilized powder or water free concentrate in a hermetically sealed container and can be reconstituted, *e.g.*, with water or saline to the appropriate concentration for administration to a subject. Preferably, the molecules of the invention are supplied as a dry sterile lyophilized powder in a hermetically sealed container at a unit dosage of at least 5 mg, more preferably at least 10 mg, at least 15 mg, at least 25 mg, at least 35 mg, at least 45 mg, at least 50 mg, or at least 75 mg. The lyophilized molecules of the invention should be stored at between 2 and 8°C in their original container and the molecules should be administered within 12 hours, preferably within 6 hours, within 5 hours, within 3 hours, or within 1 hour after being reconstituted. In an alternative embodiment, molecules of the invention are supplied in liquid form in a hermetically sealed container indicating the quantity and concentration of the molecule, fusion protein, or conjugated molecule. Preferably, the liquid form of the molecules of the invention are supplied in a hermetically sealed container at least 1 mg/ml, more preferably at least 2.5 mg/ml, at least 5 mg/ml, at least 8 mg/ml, at least 10 mg/ml, at least 15 mg/kg, at least 25 mg/ml, at least 50 mg/ml, at least 100 mg/ml, at least 150 mg/ml, at least 200 mg/ml of the molecules.

[00317] The amount of the composition of the invention which will be effective in the treatment, prevention or amelioration of one or more symptoms associated with a disorder can be determined by standard clinical techniques. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the condition, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[00318] For diabodies encompassed by the invention, the dosage administered to a patient is typically 0.0001 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.0001 mg/kg and 20 mg/kg, 0.0001 mg/kg and 10 mg/kg, 0.0001 mg/kg and 5 mg/kg, 0.0001 and 2 mg/kg, 0.0001 and 1 mg/kg, 0.0001 mg/kg and 0.75 mg/kg, 0.0001 mg/kg and 0.5 mg/kg, 0.0001 mg/kg to 0.25 mg/kg, 0.0001 to 0.15 mg/kg, 0.0001 to 0.10 mg/kg, 0.001 to 0.5 mg/kg, 0.01 to 0.25 mg/kg or 0.01 to 0.10 mg/kg of the patient's body weight. The dosage and frequency of administration of

diabodies of the invention may be reduced or altered by enhancing uptake and tissue penetration of the diabodies by modifications such as, for example, lipidation.

[00319] In one embodiment, the dosage of the molecules of the invention administered to a patient are 0.01mg to 1000mg/day, when used as single agent therapy. In another embodiment the molecules of the invention are used in combination with other therapeutic compositions and the dosage administered to a patient are lower than when said molecules are used as a single agent therapy.

[00320] In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by injection, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a molecule of the invention, care must be taken to use materials to which the molecule does not absorb.

[00321] In another embodiment, the compositions can be delivered in a vesicle, in particular a liposome (*See Langer, Science 249:1527-1533 (1990); Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, ibid., pp. 3 17-327; see generally ibid.*).

[00322] In yet another embodiment, the compositions can be delivered in a controlled release or sustained release system. Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more molecules of the invention. *See, e.g., U.S. Patent No. 4,526,938; PCT publication WO 91/05548; PCT publication WO 96/20698; Ning et al., 1996, "Intratumoral Radioimmunotherapy of a Human Colon Cancer Xenograft Using a Sustained-Release Gel," Radiotherapy & Oncology 39:179-189, Song et al., 1995, "Antibody Mediated Lung Targeting of Long-Circulating Emulsions," PDA Journal of Pharmaceutical Science & Technology 50:372-397; Cleek et al., 1997, "Biodegradable Polymeric Carriers for a bFGF Antibody for Cardiovascular Application," Pro. Int'l. Symp. Control. Rel. Bioact. Mater. 24:853-854; and Lam et al., 1997, "Microencapsulation of Recombinant Humanized Monoclonal Antibody for Local Delivery," Proc. Int'l. Symp. Control Rel. Bioact. Mater. 24:759-760, each of which is incorporated herein by reference in its entirety. In one embodiment, a pump may be used in a controlled release system (*See Langer, supra; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:20; Buchwald et al., 1980, Surgery 88:507; and Saudek et al., 1989, N. Engl. J. Med. 321:574*). In another embodiment, polymeric materials can be used to achieve*

controlled release of antibodies (*see e.g.*, Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, J., *Macromol. Sci. Rev. Macromol. Chem.* 23:61; *See also* Levy *et al.*, 1985, *Science* 228:190; During *et al.*, 1989, *Ann. Neurol.* 25:351; Howard *et al.*, 1989, *J. Neurosurg.* 71:105; U.S. Patent No. 5,679,377; U.S. Patent No. 5,916,597; U.S. Patent No. 5,912,015; U.S. Patent No. 5,989,463; U.S. Patent No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO 99/20253). Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target (*e.g.*, the lungs), thus requiring only a fraction of the systemic dose (*see, e.g.*, Goodson, in *Medical Applications of Controlled Release, supra*, vol. 2, pp. 115-138 (1984)). In another embodiment, polymeric compositions useful as controlled release implants are used according to Dunn *et al.* (*See* U.S. 5,945,155). This particular method is based upon the therapeutic effect of the in situ controlled release of the bioactive material from the polymer system. The implantation can generally occur anywhere within the body of the patient in need of therapeutic treatment. In another embodiment, a non-polymeric sustained delivery system is used, whereby a non-polymeric implant in the body of the subject is used as a drug delivery system. Upon implantation in the body, the organic solvent of the implant will dissipate, disperse, or leach from the composition into surrounding tissue fluid, and the non-polymeric material will gradually coagulate or precipitate to form a solid, microporous matrix (*See* U.S. 5,888,533).

[00323] Controlled release systems are discussed in the review by Langer (1990, *Science* 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more therapeutic agents of the invention. *See, e.g.*, U.S. Patent No. 4,526,938; International Publication Nos. WO 91/05548 and WO 96/20698; Ning *et al.*, 1996, *Radiotherapy & Oncology* 39:179-189; Song *et al.*, 1995, *PDA Journal of Pharmaceutical Science & Technology* 50:372-397; Cleek *et al.*, 1997, *Pro. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-854; and Lam *et al.*, 1997, *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-760, each of which is incorporated herein by reference in its entirety.

[00324] In a specific embodiment where the composition of the invention is a nucleic acid encoding a diabody of the invention, the nucleic acid can be administered *in vivo* to promote expression of its encoded diabody, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by use of a retroviral vector (See U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (See *e.g.*, Joliot *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88:1864-1868), *etc.* Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression by homologous recombination.

[00325] Treatment of a subject with a therapeutically or prophylactically effective amount of molecules of the invention can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with molecules of the invention in the range of between about 0.1 to 30 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. In other embodiments, the pharmaceutical compositions of the invention are administered once a day, twice a day, or three times a day. In other embodiments, the pharmaceutical compositions are administered once a week, twice a week, once every two weeks, once a month, once every six weeks, once every two months, twice a year or once per year. It will also be appreciated that the effective dosage of the molecules used for treatment may increase or decrease over the course of a particular treatment.

5.9.1 PHARMACEUTICAL COMPOSITIONS

[00326] The compositions of the invention include bulk drug compositions useful in the manufacture of pharmaceutical compositions (*e.g.*, impure or non-sterile compositions) and pharmaceutical compositions (*i.e.*, compositions that are suitable for administration to a subject or patient) which can be used in the preparation of unit dosage forms. Such compositions comprise a prophylactically or therapeutically effective amount of a prophylactic and/or therapeutic agent disclosed herein or a combination of those agents and a pharmaceutically acceptable carrier. Preferably, compositions of the invention comprise a prophylactically or therapeutically effective amount of one or more molecules of the invention and a pharmaceutically acceptable carrier.

[00327] The invention also encompasses pharmaceutical compositions comprising a diabody molecule of the invention and a therapeutic antibody (*e.g.*, tumor specific monoclonal antibody) that is specific for a particular cancer antigen, and a pharmaceutically acceptable carrier.

[00328] In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant (*e.g.*, Freund's adjuvant (complete and incomplete), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

[00329] Generally, the ingredients of compositions of the invention are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00330] The compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include, but are not limited to those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, *etc.*, and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, *etc.*

5.9.2 GENE THERAPY

[00331] In a specific embodiment, nucleic acids comprising sequences encoding molecules of the invention, are administered to treat, prevent or ameliorate one or more symptoms associated with a disease, disorder, or infection, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded antibody or fusion protein that mediates a therapeutic or prophylactic effect.

[00332] Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

[00333] For general reviews of the methods of gene therapy, *see* Goldspiel *et al.*, 1993, *Clinical Pharmacy* 12:488-505; Wu and Wu, 1991, *Biotherapy* 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIBTECH* 11(5):155-215. Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel *et al.* (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990).

[00334] In a preferred aspect, a composition of the invention comprises nucleic acids encoding a diabody of the invention, said nucleic acids being part of an expression vector that expresses the antibody in a suitable host. In particular, such nucleic acids have promoters, preferably heterologous promoters, operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, 1989, *Proc. Natl. Acad. Sci. USA* 86:8932-8935; and Zijlstra *et al.*, 1989, *Nature* 342:435-438).

[00335] In another preferred aspect, a composition of the invention comprises nucleic acids encoding a fusion protein, said nucleic acids being a part of an expression vector that expresses the fusion protein in a suitable host. In particular, such nucleic acids have promoters, preferably heterologous promoters, operably linked to the coding region of a fusion protein, said promoter being inducible or constitutive, and optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the coding sequence of the fusion protein and any other desired sequences are flanked by regions that

promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the fusion protein.

[00336] Delivery of the nucleic acids into a subject may be either direct, in which case the subject is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids *in vitro*, then transplanted into the subject. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

[00337] In a specific embodiment, the nucleic acid sequences are directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, *e.g.*, by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, *e.g.*, by infection using defective or attenuated retroviral or other viral vectors (*see* U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to an antigen subject to receptor-mediated endocytosis (*See, e.g.*, Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432) (which can be used to target cell types specifically expressing the receptors), *etc.* In another embodiment, nucleic acid-antigen complexes can be formed in which the antigen comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (*See, e.g.*, PCT Publications WO 92/06180; WO 92/22635; WO 92/20316; WO 93/14188; WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, 1989, *Proc. Natl. Acad. Sci. USA* 86:8932-8935; and Zijlstra *et al.*, 1989, *Nature* 342:435-438).

[00338] In a specific embodiment, viral vectors that contain nucleic acid sequences encoding a molecule of the invention (*e.g.*, a diabody or a fusion protein) are used. For example, a retroviral vector can be used (*See* Miller *et al.*, 1993, *Meth. Enzymol.* 217:581-599). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody or a fusion protein to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the nucleotide sequence into a subject. More detail about retroviral vectors can be found in Boesen *et al.*, (1994, *Biotherapy* 6:291-302),

which describes the use of a retroviral vector to deliver the *mdr 1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes *et al.*, 1994, *J. Clin. Invest.* 93:644-651; Klein *et al.*, 1994, *Blood* 83:1467-1473; Salmons and Gunzberg, 1993, *Human Gene Therapy* 4:129-141; and Grossman and Wilson, 1993, *Curr. Opin. in Genetics and Devel.* 3:110-114.

[00339] Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson (*Current Opinion in Genetics and Development* 3:499-503, 1993, present a review of adenovirus-based gene therapy. Bout *et al.*, (*Human Gene Therapy*, 5:3-10, 1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld *et al.*, 1991, *Science* 252:431-434; Rosenfeld *et al.*, 1992, *Cell* 68:143-155; Mastrangeli *et al.*, 1993, *J. Clin. Invest.* 91:225-234; PCT Publication W094/12649; and Wang *et al.*, 1995, *Gene Therapy* 2:775-783. In a preferred embodiment, adenovirus vectors are used.

[00340] Adeno-associated virus (AAV) has also been proposed for use in gene therapy (*see, e.g.*, Walsh *et al.*, 1993, *Proc. Soc. Exp. Biol. Med.* 204:289-300 and U.S. Patent No. 5,436,146).

[00341] Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a subject.

[00342] In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to, transfection, electroporation, microinjection, infection with a viral or bacteriophage vector, containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, *etc.* Numerous techniques are known in the art for the introduction of foreign genes into cells (*See, e.g.*, Loeffler and Behr, 1993,

Meth. Enzymol. 217:599-618, Cohen *et al.*, 1993, *Meth. Enzymol.* 217:618-644; and *Clin. Pharma. Ther.* 29:69-92, 1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

[00343] The resulting recombinant cells can be delivered to a subject by various methods known in the art. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, *etc.*, and can be determined by one skilled in the art.

[00344] Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, *etc.*

[00345] In a preferred embodiment, the cell used for gene therapy is autologous to the subject.

[00346] In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody or a fusion protein are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention (*See e.g.*, PCT Publication WO 94/08598; Stemple and Anderson, 1992, *Cell* 71:973-985; Rheinwald, 1980, *Meth. Cell Bio.* 21A:229; and Pittelkow and Scott, 1986, *Mayo Clinic Proc.* 61:771).

[00347] In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

5.9.3 KITS

[00348] The invention provides a pharmaceutical pack or kit comprising one or more containers filled with the molecules of the invention. Additionally, one or more other prophylactic or therapeutic agents useful for the treatment of a disease can also be included in the pharmaceutical pack or kit. The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[00349] The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises one or more molecules of the invention. In another embodiment, a kit further comprises one or more other prophylactic or therapeutic agents useful for the treatment of cancer, in one or more containers. In another embodiment, a kit further comprises one or more cytotoxic antibodies that bind one or more cancer antigens associated with cancer. In certain embodiments, the other prophylactic or therapeutic agent is a chemotherapeutic. In other embodiments, the prophylactic or therapeutic agent is a biological or hormonal therapeutic.

5.10 CHARACTERIZATION AND DEMONSTRATION OF THERAPEUTIC UTILITY

[00350] Several aspects of the pharmaceutical compositions, prophylactic, or therapeutic agents of the invention are preferably tested *in vitro*, in a cell culture system, and in an animal model organism, such as a rodent animal model system, for the desired therapeutic activity prior to use in humans. For example, assays which can be used to determine whether administration of a specific pharmaceutical composition is desired, include cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise contacted with a pharmaceutical composition of the invention, and the effect of such composition upon the tissue sample is observed. The tissue sample can be obtained by biopsy from the patient. This test allows the identification of the therapeutically most effective prophylactic or therapeutic molecule(s) for each individual patient. In various specific embodiments, *in vitro* assays can be carried out with representative cells of cell types involved in an autoimmune or inflammatory disorder (*e.g.*, T cells), to determine if a pharmaceutical composition of the invention has a desired effect upon such cell types.

[00351] Combinations of prophylactic and/or therapeutic agents can be tested in suitable animal model systems prior to use in humans. Such animal model systems include,

but are not limited to, rats, mice, chicken, cows, monkeys, pigs, dogs, rabbits, *etc.* Any animal system well-known in the art may be used. In a specific embodiment of the invention, combinations of prophylactic and/or therapeutic agents are tested in a mouse model system. Such model systems are widely used and well-known to the skilled artisan. Prophylactic and/or therapeutic agents can be administered repeatedly. Several aspects of the procedure may vary. Said aspects include the temporal regime of administering the prophylactic and/or therapeutic agents, and whether such agents are administered separately or as an admixture.

[00352] Preferred animal models for use in the methods of the invention are, for example, transgenic mice expressing human FcγRs on mouse effector cells, *e.g.*, any mouse model described in U.S. 5,877,396 (which is incorporated herein by reference in its entirety) can be used in the present invention. Transgenic mice for use in the methods of the invention include, but are not limited to, mice carrying human FcγRIIIA; mice carrying human FcγRIIA; mice carrying human FcγRIIB and human FcγRIIIA; mice carrying human FcγRIIB and human FcγRIIA. Preferably, mutations showing the highest levels of activity in the functional assays described above will be tested for use in animal model studies prior to use in humans. Sufficient quantities of antibodies may be prepared for use in animal models using methods described supra, for example using mammalian expression systems and purification methods disclosed and exemplified herein.

[00353] Mouse xenograft models may be used for examining efficacy of mouse antibodies generated against a tumor specific target based on the affinity and specificity of the epitope binding domains of the diabody molecule of the invention and the ability of the diabody to elicit an immune response (Wu et al., 2001, Trends Cell Biol. 11: S2-9). Transgenic mice expressing human FcγRs on mouse effector cells are unique and are tailor-made animal models to test the efficacy of human Fc-FcγR interactions. Pairs of FcγRIIIA, FcγRIIB and FcγRIIA transgenic mouse lines generated in the lab of Dr. Jeffrey Ravetch (Through a licensing agreement with Rockefeller U. and Sloan Kettering Cancer center) can be used such as those listed in the Table 11 below.

Table 11: Mice Strains

Strain Background	Human FcR
Nude / CD16A KO	None
Nude / CD16A KO	FcγRIIIA
Nude / CD16A KO	FcγR IIA
Nude / CD16A KO	FcγR IIA and IIIA
Nude / CD32B KO	None
Nude / CD32B KO	FcγR IIB

[00354] The anti-inflammatory activity of the combination therapies of invention can be determined by using various experimental animal models of inflammatory arthritis known in the art and described in Crofford L.J. and Wilder R.L., "Arthritis and Autoimmunity in Animals", in *Arthritis and Allied Conditions: A Textbook of Rheumatology*, McCarty *et al.*(eds.), Chapter 30 (Lee and Febiger, 1993). Experimental and spontaneous animal models of inflammatory arthritis and autoimmune rheumatic diseases can also be used to assess the anti-inflammatory activity of the combination therapies of invention. The following are some assays provided as examples, and not by limitation.

[00355] The principle animal models for arthritis or inflammatory disease known in the art and widely used include: adjuvant-induced arthritis rat models, collagen-induced arthritis rat and mouse models and antigen-induced arthritis rat, rabbit and hamster models, all described in Crofford L.J. and Wilder R.L., "Arthritis and Autoimmunity in Animals", in *Arthritis and Allied Conditions: A Textbook of Rheumatology*, McCarty *et al.*(eds.), Chapter 30 (Lee and Febiger, 1993), incorporated herein by reference in its entirety.

[00356] The anti-inflammatory activity of the combination therapies of invention can be assessed using a carrageenan-induced arthritis rat model. Carrageenan-induced arthritis has also been used in rabbit, dog and pig in studies of chronic arthritis or inflammation. Quantitative histomorphometric assessment is used to determine therapeutic efficacy. The methods for using such a carrageenan-induced arthritis model is described in Hansra P. *et al.*, "Carrageenan-Induced Arthritis in the Rat," *Inflammation*, 24(2): 141-155, (2000). Also commonly used are zymosan-induced inflammation animal models as known and described in the art.

[00357] The anti-inflammatory activity of the combination therapies of invention can also be assessed by measuring the inhibition of carrageenan-induced paw edema in the rat, using a modification of the method described in Winter C. A. *et al.*, "Carrageenan-Induced Edema in Hind Paw of the Rat as an Assay for Anti-inflammatory Drugs" *Proc. Soc. Exp. Biol Med.* 111, 544-547, (1962). This assay has been used as a primary *in vivo* screen for the anti-inflammatory activity of most NSAIDs, and is considered predictive of human efficacy. The anti-inflammatory activity of the test prophylactic or therapeutic agents is expressed as the percent inhibition of the increase in hind paw weight of the test group relative to the vehicle dosed control group.

[00358] Additionally, animal models for inflammatory bowel disease can also be used to assess the efficacy of the combination therapies of invention (Kim *et al.*, 1992,

Scand. *J. Gastroentrol.* 27:529-537; Strober, 1985, *Dig. Dis. Sci.* 30(12 Suppl):3S-10S).

Ulcerative colitis and Crohn's disease are human inflammatory bowel diseases that can be induced in animals. Sulfated polysaccharides including, but not limited to amylopectin, carrageen, amylopectin sulfate, and dextran sulfate or chemical irritants including but not limited to trinitrobenzenesulphonic acid (TNBS) and acetic acid can be administered to animals orally to induce inflammatory bowel diseases.

[00359] Animal models for autoimmune disorders can also be used to assess the efficacy of the combination therapies of invention. Animal models for autoimmune disorders such as type 1 diabetes, thyroid autoimmunity, systemic lupus eruthematosus, and glomerulonephritis have been developed (Flanders *et al.*, 1999, *Autoimmunity* 29:235-246; Krogh *et al.*, 1999, *Biochimie* 81:511-515; Foster, 1999, *Semin. Nephrol.* 19:12-24).

[00360] Further, any assays known to those skilled in the art can be used to evaluate the prophylactic and/or therapeutic utility of the combinatorial therapies disclosed herein for autoimmune and/or inflammatory diseases.

[00361] Toxicity and efficacy of the prophylactic and/or therapeutic protocols of the instant invention can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Prophylactic and/or therapeutic agents that exhibit large therapeutic indices are preferred. While prophylactic and/or therapeutic agents that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such agents to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[00362] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage of the prophylactic and/or therapeutic agents for use in humans. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any agent used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (*i.e.*, the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately

determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[00363] The anti-cancer activity of the therapies used in accordance with the present invention also can be determined by using various experimental animal models for the study of cancer such as the SCID mouse model or transgenic mice or nude mice with human xenografts, animal models, such as hamsters, rabbits, *etc.* known in the art and described in *Relevance of Tumor Models for Anticancer Drug Development* (1999, eds. Fiebig and Burger); *Contributions to Oncology* (1999, Karger); *The Nude Mouse in Oncology Research* (1991, eds. Boven and Winograd); and *Anticancer Drug Development Guide* (1997 ed. Teicher), herein incorporated by reference in their entireties.

[00364] Preferred animal models for determining the therapeutic efficacy of the molecules of the invention are mouse xenograft models. Tumor cell lines that can be used as a source for xenograft tumors include but are not limited to, SKBR3 and MCF7 cells, which can be derived from patients with breast adenocarcinoma. These cells have both erbB2 and prolactin receptors. SKBR3 cells have been used routinely in the art as ADCC and xenograft tumor models. Alternatively, OVCAR3 cells derived from a human ovarian adenocarcinoma can be used as a source for xenograft tumors.

[00365] The protocols and compositions of the invention are preferably tested *in vitro*, and then *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. Therapeutic agents and methods may be screened using cells of a tumor or malignant cell line. Many assays standard in the art can be used to assess such survival and/or growth; for example, cell proliferation can be assayed by measuring ³H-thymidine incorporation, by direct cell count, by detecting changes in transcriptional activity of known genes such as proto-oncogenes (*e.g.*, fos, myc) or cell cycle markers; cell viability can be assessed by trypan blue staining, differentiation can be assessed visually based on changes in morphology, decreased growth and/or colony formation in soft agar or tubular network formation in three-dimensional basement membrane or extracellular matrix preparation, *etc.*

[00366] Compounds for use in therapy can be tested in suitable animal model systems prior to testing in humans, including but not limited to in rats, mice, chicken, cows, monkeys, rabbits, hamsters, *etc.*, for example, the animal models described above. The compounds can then be used in the appropriate clinical trials.

[00367] Further, any assays known to those skilled in the art can be used to evaluate the prophylactic and/or therapeutic utility of the combinatorial therapies disclosed herein for treatment or prevention of cancer, inflammatory disorder, or autoimmune disease.

6. EXAMPLES

6.1 DESIGN AND CHARACTERIZATION OF COVALENT BISPECIFIC DIABODIES

[00368] A monospecific covalent diabody and a bispecific covalent diabody were constructed to assess the recombinant production, purification and binding characteristics of each. The affinity purified diabody molecules that were produced by the recombinant expression systems described herein were found by SDS-PAGE and SEC analysis to consist of a single dimeric species. ELISA and SPR analysis further revealed that the covalent bispecific diabody exhibited affinity for both target antigens and could bind both antigens simultaneously.

[00369] **Materials and Methods:**

[00370] Construction and Design of Polypeptide Molecules: Nucleic acid expression vectors were designed to produce four polypeptide constructs, schematically represented in FIG. 2. Construct 1 (SEQ ID NO:9) comprised the VL domain of humanized 2B6 antibody, which recognizes FcγRIIB, and the VH domain of humanized 3G8 antibody, which recognizes FcγRIIIA. Construct 2 (SEQ ID NO:11) comprised the VL domain of Hu3G8 and the VH domain of Hu2B6. Construct 3 (SEQ ID NO:12) comprised the VL domain of Hu3G8 and the VH domain of Hu3G8. Construct 4 (SEQ ID NO:13) comprised the VL domain of Hu2B6 and the VH domain of Hu2B6.

[00371] PCR and Expression Vector Construction: The coding sequences of the VL or VH domains were amplified from template DNA using forward and reverse primers designed such that the initial PCR products would contain overlapping sequences, allowing overlapping PCR to generate the coding sequences of the desired polypeptide constructs.

[00372] Initial PCR amplification of template DNA: Approximately 35 ng of template DNA, e.g. light chain and heavy chain of antibody of interest; 1 ul of 10uM forward and reverse primers; 2.5 ul of 10x pfuUltra buffer (Stratagene, Inc.); 1 ul of 10 mM dNTP; 1 ul of 2.5 units/ul of pfuUltra DNA polymerase (Stratagene, Inc.); and distilled water to 25 ul total volume were gently mixed in a microfuge tube and briefly spun in a microcentrifuge to collect the reaction mixture at the bottom of the tube. PCR reactions were performed using GeneAmp PCR System 9700 (PE Applied Biosystem) and the following settings: 94°C, 2 minutes; 25 cycles of 94°C, each 15 seconds; 58°C, 30 seconds; and 72°C, 1 minute.

[00373] The VL of Hu2B6 was amplified from the light chain of Hu2B6 using forward and reverse primers SEQ ID NO: 57 and SEQ ID NO:58, respectively. The VH of Hu2B6 was amplified from the heavy chain of Hu2B6 using forward and reverse primers

SEQ ID NO: 59 and SEQ ID NO:60, respectively. The VL of Hu3G8 was amplified from the light chain of Hu3G8 using forward and reverse primers SEQ ID NO: 57 and SEQ ID NO:61, respectively. The VH of Hu3G8 was amplified from the heavy chain of Hu3G8 using forward and reverse primers SEQ ID NO: 62 and SEQ ID NO:63, respectively.

[00374] PCR products were electrophoresed on a 1% agarose gel for 30 minutes at 120 volts. PCR products were cut from the gel and purified using MinElute Gel Extraction Kit (Qiagen, Inc.).

[00375] *Overlapping PCR*: Initial PCR products were combined as described below and amplified using the same PCR conditions described for initial amplification of template DNA. Products of overlapping PCR were also purified as described *supra*.

[00376] The nucleic acid sequence encoding construct 1, SEQ ID NO:9 (shown schematically in FIG. 2), was amplified by combining the PCR products of the amplifications of VL Hu2B6 and VH Hu3G8, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:63, respectively. The nucleic acid sequence encoding construct 2, SEQ ID NO:11 (shown schematically in FIG. 2), was amplified by combining the PCR products of the amplifications of VL Hu3G8 and VH Hu2B6, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:60, respectively. The nucleic acid sequence encoding construct 3, SEQ ID NO:12 (shown schematically in FIG. 2), was amplified by combining the PCR products of the amplifications of VL Hu3G8 and VH Hu3G8, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:63, respectively. The nucleic acid sequence encoding construct 4, SEQ ID NO:13 (shown schematically in FIG. 2), was amplified by combining the PCR products of the amplifications of VL Hu2B6 and VH Hu2B6, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:60, respectively.

[00377] The forward primers of the VL domains (*i.e.*, SEQ ID NO:57) and reverse primers of the VH domains (*i.e.*, SEQ ID NO:60 and SEQ ID NO:63) contained unique restriction sites to allow cloning of the final product into an expression vector. Purified overlapping PCR products were digested with restriction endonucleases Nhe I and EcoR I, and cloned into the pCIneo mammalian expression vector (Promega, Inc.). The plasmids encoding constructs were designated as identified in Table 12:

Table 12. PLASMID CONSTRUCTS

Encoding Construct	Plasmid Designation	Insert
1	pMGX0669	hu2B6VL-hu3G8VH
2	pMGX0667	hu3G8VL-hu2B6VH
3	pMGX0666	hu3G8VL-hu3G8VH
4	pMGX0668	hu2B6VL-hu2B6VH

[00378] *Polypeptide/diabody Expression:* pMGX0669, encoding construct 1, was cotransfected with pMGX0667, encoding construct 2, in HEK-293 cells using Lipofectamine 2000 according to the manufacturer's directions (Invitrogen). Co-transfection of these two plasmids was designed to lead to the expression of a covalent bispecific diabody (CBD) immunospecific for both FcγRIIB and FcγRIIA (the h2B6-h3G8 diabody). pMGX0666 and pMGX0668, encoding constructs 3 and 4, respectively, were separately transfected into HEK-293 cells for expression of a covalent monospecific diabody (CMD), immunospecific for FcγRIIA (h3G8 diabody) and FcγRIIB (h2B6 diabody), respectively. Following three days in culture, secreted products were purified from the conditioned media.

[00379] *Purification:* Diabodies were captured from the conditioned medium using the relevant antigens coupled to CNBr activated Sepharose 4B. The affinity Sepharose resin was equilibrated in 20 mM Tris/HCl, pH 8.0 prior to loading. After loading, the resin was washed with equilibration buffer prior to elution. Diabodies were eluted from the washed resin using 50 mM Glycine pH 3.0. Eluted diabodies were immediately neutralized with 1M Tris/HCl pH 8.0 and concentrated using a centrifugation type concentrator. The concentrated diabodies were further purified by size exclusion chromatography using a Superdex 200 column equilibrated in PBS.

[00380] *SEC:* Size exclusion chromatography was used to analyze the approximate size and heterogeneity of the diabodies eluted from the column. SEC analysis was performed on a GE healthcare Superdex 200HR 10/30 column equilibrated with PBS. Comparison with the elution profiles of a full length IgG (~150 kDa), an Fab fragment (~50 kDa) and a single chain Fv (~30 kDa) were used as controls).

[00381] *ELISA:* The binding of eluted and purified diabodies was characterized by ELISA assay, as described in 5.4.2. 50 ul/well of a 2 ug/ml solution of sCD32B-Ig was coated on 96-well Maxisorp plate in Carbonate buffer at 4°C over night. The plate was washed three times with PBS-T (PBS, 0.1% Tween 20) and blocked by 0.5% BSA in PBS-T for 30 minutes at room temperature. Subsequently, h2B6-h3G8 CBD, h2B6 CMD, or h3G8 CMD were diluted into the blocking buffer in a serial of two-fold dilutions to generate a range of diabody concentrations, from 0.5 µg/ml to 0.001 µg/ml. The plate was then incubated at room temperature for 1 hour. After washing with PBS-T three times, 50 ul/well of 0.2 ug/ml sCD16A-Biotin was added to each well. The plate was again incubated at room temperature for 1 hour. After washing with PBS-T three times, 50 ul/well of a 1:5000 dilution of HRP conjugated streptavidin (Amersham Pharmacia Biotech) was used for detection. The HRP-streptavidin was allowed to incubate for 45 minutes at room

temperature. The plate was washed with PBS-T three times and developed using 80 ul/well of TMB substrate. After a 10 minute incubation, the HRP-TMB reaction was stopped by adding 40 ul/well of 1% H₂SO₄. The OD450 nm was read by using a 96-well plate reader and SOFTmax software, and results plotted using GraphPadPrism 3.03 software.

[00382] BIAcore Assay: The kinetic parameters of the binding of eluted and purified diabodies were analyzed using a BIAcore assay (BIAcore instrument 1000, BIAcore Inc., Piscataway, N.J.) and associated software as described in section 5.4.3.

[00383] sCD16A, sCD32B or sCD32A (negative control) were immobilized on one of the four flow cells (flow cell 2) of a sensor chip surface through amine coupling chemistry (by modification of carboxymethyl groups with mixture of NHS/EDC) such that about 1000 response units (RU) of either receptor was immobilized on the surface. Following this, the unreacted active esters were "capped off" with an injection of 1M Et-NH₂. Once a suitable surface was prepared, covalent bispecific diabodies (h2B6-h3G8 CBD) or covalent monospecific diabodies (h2B6 CMD or h3G8 CMB) were passed over the surface by 180 second injections of a 6.25-200nM solution at a 70 mL/min flow rate. h3G8 scFv was also tested for comparison.

[00384] Once an entire data set was collected, the resulting binding curves were globally fitted using computer algorithms supplied by the manufacturer, BIAcore, Inc. (Piscataway, NJ). These algorithms calculate both the K_{on} and K_{off} , from which the apparent equilibrium binding constant, K_D is deduced as the ratio of the two rate constants (*i.e.*, K_{off}/K_{on}). More detailed treatments of how the individual rate constants are derived can be found in the BIAevaluation Software Handbook (BIAcore, Inc., Piscataway, NJ).

[00385] Association and dissociation phases were fitted separately. Dissociation rate constant was obtained for interval 32-34 sec of the 180 sec dissociation phase; association phase fit was obtained by a 1:1 Langmuir model and base fit was selected on the basis R_{max} and χ^2 criteria for the bispecific diabodies and scFv; Bivalent analyte fit was used for CMD binding.

[00386] Results

[00387] SDS-PAGE analysis under non-reducing conditions revealed that the purified product of the h3G8 CMD, h2B6 CMD and h2B6-h3G8 CBD expression systems were each a single species with an estimated molecular weight of approximately 50 kDa (FIG. 3, lanes 4, 5 and 6, respectively). Under reducing conditions, the product purified from either of the CMD expression systems ran as a single band (lanes 1 and 2), while the product purified from the h2B6-h3G8 CBD system was revealed to be 2 separate proteins

(FIG. 3, lane 3). All polypeptides purified from the expression system and visualized by SDS-PAGE under reducing conditions migrated at approximately 28 kDa.

[00388] SEC analysis of each of the expression system products also revealed a single molecular species (FIG. 4B), each of which eluted at the same approximate time as an Fab fragment of IgG (~50kDa) (FIG. 4A). The results indicate that affinity purified product was a homogenous covalent homodimer for the case of CMD expression system and a homogenous covalent heterodimer for the case of the h2B6-h3G8 CBD.

[00389] An ELISA sandwich assay was used to test binding of the h2B6-h3G8 CBD for specificity to either or both of CD32B and/or CD16A (FIG. 5). CD32B served as the target antigen and CD16A was used as the secondary probe. The positive signal in the ELISA revealed that the heterodimeric h2B6-h3G8 CBD had specificity for both antigens. Similar testing of the h3G8 CMD (which should not bind CD32B) showed no signal.

[00390] SPR analysis indicated that h3G8 CMD immunospecifically recognized sCD16 but not sCD32B, that h2B6 CMD immunospecifically recognized sCD32B but not sCD16, and that h2B6-h3G8 CBD immunospecifically recognized both sCD16 and sCD32B (FIGS. 6A-B). None of the diabodies tested bound the control receptor, sCD32A (FIG. 6C).

[00391] SPR analysis was also used to estimate the kinetic and equilibrium constants of the CMDs and h2B6-h3G8 CBD to sCD16 and/or sCD32B. Results were compared to the same constants calculated for an h3G8 scFv. FIGS. 7A-E show the graphical results of the SPR analysis. The kinetic on and off rates, as well as the equilibrium constant, calculated from the results depicted in FIG. 7 are provided in Table 13.

Table 13. Kinetic and Equilibrium Constants Calculated from BIAcore Data.

Receptor / Analyte	k-on	k-off	Kd
sCD16 / h3G8 diabody	2.3 x 10 ⁵	0.004	18.0
sCD16 / h2B6-h3G8 CBD	4.6 x 10 ⁵	0.010	22.7
sCD16 / h3G8 scFv	3.2 x 10 ⁵	0.013	38.7
sCD32B / h2B6-h3G8 CBD	3.6 x 10 ⁵	0.005	15.0
sCD32B / h2B6 diabody	6.2 x 10 ⁵	0.013	21.0

[00392] Coupled with the results of the ELISA analysis, the studies confirm that the h2B6-h3G8 covalent heterodimer retained specificity for both CD32B and CD16, and was capable of binding both antigens simultaneously. The molecule is schematically represented in FIG. 8.

6.2 DESIGN AND CHARACTERIZATION OF COVALENT BISPECIFIC DIABODIES COMPRISING Fc DOMAINS

[00393] In an effort to create an IgG like molecule, *i.e.*, comprising an Fc domain, one of the polypeptides comprising the heterodimeric CBD molecule presented in Example 6.1 was modified to further comprise an Fc domain (creating a 'heavier' and 'lighter' chain, analogous to an antibody heavy and light chain). The heterodimeric bispecific molecule would then contain an Fc domain that will dimerize with a homologous molecule, forming a tetrameric IgG-like molecule with tetravalency (*i.e.*, formed by dimerization via the Fc domains of the heterodimeric bispecific molecules). Interestingly, such tetrameric molecules were not detected in the conditioned media of recombinant expression systems using functional assays, *e.g.*, testing the conditioned media for immunospecific binding to target antigens. Instead, only a dimeric molecule, comprising monomers consisting of a VL, VH and Fc domain, were detected in such functional assays. To test whether stability of the theoretical tetrameric structure was at issue, polypeptides comprising the Fc domain were engineered to further comprise a hinge region while the polypeptides comprising the 'lighter' chain were engineered to further comprise the 6 C-terminal amino acids of the constant domain of the human kappa light chain. When such reengineered 'heavier' and 'lighter' chains were co-expressed in the recombinant expression systems, functional assays detected diabody molecules that were able to immunospecifically bind both of the target antigens and anti-Fc antibodies.

[00394] Materials and Methods

[00395] Construction and Design of Polypeptide Molecules: Nucleic acid expression vectors were designed to produce modified versions of constructs 1 and 2 presented in Example 6.1. Construct 5 (SEQ ID NO: 14) and 6 (SEQ ID NO:15), were created by engineering construct 1 and 2, respectively to further comprise an Fc domain. Construct 7 (SEQ ID NO: 16) was created by engineering construct 1 was to further comprise the sequence FNRGEC (SEQ ID NO: 23) at its C-terminus. Construct 8 (SEQ ID NO:18) was created by engineering construct 2 to further comprise a hinge region and Fc domain (comprising V215A mutation). Schematic representation of constructs 5-8 is shown in FIG. 9.

[00396] PCR and Expression Vector Construction: All PCR and PCR product purification protocols were as described in Example 6.1. Plasmids pMGX0669 and pMGX0667 served as templates for the coding sequences of constructs 1 and 2, respectively. The coding sequences for the of HuIgG Fc domain and/or hinge domain were SEQ ID NO:5 or SEQ ID NO:1 and SEQ ID NO:5, respectively. The coding sequences of

the template DNAs were amplified using forward and reverse primers such that the PCR products would contain overlapping sequences, allowing overlapping PCR to generate the coding sequences of the desired products.

[00397] The coding sequence of construct 1 was amplified from pMGX0669 using forward and reverse primers SEQ ID NO:57 and SEQ ID NO:64, respectively. The coding sequence of construct 2 was amplified from pMGX0667 using forward and reverse primers SEQ ID NO:57 and SEQ ID NO:65, respectively. HuIgG hinge-Fc was amplified using forward and reverse primers SEQ ID NO:67 and SEQ ID NO:68, respectively. Construct 7 (SEQ ID NO:16) was amplified from pMGX0669 using forward and reverse primers SEQ ID NO:57 and SEQ ID NO:69.

[00398] *Overlapping PCR:* Initial PCR products were combined as described below, amplified and purified as described in example 6.1.

[00399] The nucleic acid sequence encoding construct 5, SEQ ID NO:14 (shown schematically in FIG. 9), was amplified by combining the PCR products of the amplifications of construct 1 and HuIgG Fc, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:66, respectively. The nucleic acid sequence encoding construct 6, SEQ ID NO:15 (shown schematically in FIG. 9), was amplified by combining the PCR products of the amplifications of construct 2 and HuIgG Fc, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:66, respectively. The nucleic acid sequence encoding construct 8, SEQ ID NO:18 (shown schematically in FIG. 9), was amplified by combining the PCR products of the amplifications of construct 2 and HuIgG hinge-Fc, and forward and reverse primers SEQ ID NO:57 and SEQ ID NO:68, respectively.

[00400] Final products were cloned into pCIneo mammalian expression vector (Promega, Inc.) as previously described. The plasmid encoding constructs were designated as identified in Table 14:

Table 14. PLASMID CONSTRUCTS

Encoding Construct	Plasmid Designation	Insert
5	pMGX0676	hu2B6VL-hu3G8VH-huFc
6	pMGX0674	hu3G8VL-hu2B6VH-huFc
7	pMGX0677	Hu2B6VL-hu3G8VH-FNRGEC
8	pMGX0678	Hu3G8VL-hu2B6VH-hu hinge-Fc (A215V)

[00401] *Polypeptide/diabody Expression:* Four separate cotransfections into in HEK-293 cells using Lipofectamine 2000, as described in section 6.1, were performed: pMGX0669 and pMGX0674, encoding constructs 1 and 6, respectively;; pMGX0667 and

pMGX0676, encoding constructs 2 and 5, respectively; and pMGX0677 and pMGX0678, encoding constructs 7 and 8, respectively.

[00402] Co-transfection of these plasmids was designed to lead to the expression of a bispecific diabody (CBD) of tetravalency with IgG-like structure, immunospecific for both FcγRIIB and FcγRIIA. An additional cotransfection was also performed: pMGX0674 and pMGX0676, encoding constructs 6 and 5, respectively. Following three days in culture, conditioned media was harvested. The amount of secreted product in the conditioned media was quantitated by anti IgG Fc ELISA using purified Fc as a standard. The concentrations of product in the samples was then normalized based on the quantitation, and the normalized samples used for the remaining assays.

[00403] ELISA: The binding of diabody molecules secreted into the medium was assayed by sandwich ELISA as described, *supra*. Unless indicated, CD32B was used to coat the plate, *i.e.*, as the target protein, and HRP- conjugated CD16 was used as the probe.

[00404] Results

[00405] An ELISA assay was used to test the normalized samples from the recombinant expression systems comprising constructs 1 and 6 (pMGX669-pMGX674), constructs 2 and 5 (pMGX667-pMGX676) and constructs 5 and 6 (pMGX674-pMGX676) for expression of diabody molecules capable of simultaneous binding to CD32B and CD16A (FIG. 10). The ELISA data indicated that co-transfection with constructs 1 and 6 or co-transfection with constructs 2 and 5 failed to produce a product that could bind either or both antigens (FIG. 10, □ and ▲, respectively). However, co-transfection of constructs 5 and 6 lead to secretion of a product capable of binding to both CD32B and CD16 antigens. The latter product was a dimer of constructs 5 and 6, containing one binding site for each antigen with a structure schematically depicted in FIG. 11.

[00406] In order to drive formation of an IgG like heterotetrameric structure, the coding sequence for six additional amino acids was appended to the C-terminal of construct 1, generating construct 7 (SEQ ID NO:16 and shown schematically in FIG. 9). The six additional amino acids, FNRGEC (SEQ ID NO:23), were derived from the C-terminal end of the the Kappa light chain and normally interact with the upper hinge domain of the heavy chain in an IgG molecule. A hinge domain was then engineered into construct 6, generating construct 8 (SEQ ID NO:18 and FIG. 9). Construct 8 additionally comprised an amino-acid mutation in the upper hinge region, A215V. Expression plasmids encoding construct 7 and construct 8, pMGX677 and pMGX678, respectively, were then cotransfected into HEK-293 cells and expressed as described.

[00407] Diabody molecules produced from the recombinant expression system comprising constructs 7 and 8 (pMGX0677 + pMGX0678), were compared in an ELISA assay for binding to CD32B and CD16A to diabody molecules produced from expression systems comprising constructs 1 and 6 (pMGX669 + pMGX674), constructs 2 and 8 (pMGX669 + pMGX678), and constructs 6 and 7 (pMGX677 + pMGX674) (FIG. 12).

[00408] As before, the molecule produced by the expression system comprising constructs 1 and 6 (pMGX669 + pMGX674) proved unable to bind both CD32A and CD16A (FIG. 10 and FIG. 12). In contrast, the product from the co-expression of either constructs 7 and 6 (pMGX0677 + pMGX0674) or from the co-expression of constructs 7 and 8 (pMGX0677-pMGX0678) were able to bind both CD32B and CD16 (FIG. 12). It is noted that construct 7 is analogous to construct 1, with the exception that construct 7 comprises the C-terminal sequence FNRGEC (SEC ID NO:23); and that construct 8 is analogous to construct 6, except that construct 8 comprises a hinge domain and the mutation A215V. The data indicate that the addition of the 6 extra amino-acids from the C-terminus of the C-kappa light chain (FNRGEC; SEQ ID NO:23) to the non-Fc bearing, 'lighter,' chain helped stabilize the formation of the tetrameric IgG-like diabody molecules, regardless of whether the corresponding heavier chain comprised a hinge domain (*i.e.*, pMGX0677 + pMGX0674 and pMGX0677-pMGX0678, FIG. 12). The addition of the hinge domain to the Fc bearing 'heavier' polypeptide, without the addition of the FNRGEC (SEQ ID NO:23) C-terminal sequence to the corresponding 'lighter' chain, was apparently unable to effect similar stabilization (*i.e.*, lack of binding by product of co-transfection of constructs 2 and 8 (pMGX669 + pMGX678)). The structure of the tetrameric diabody molecule is schematically represented in FIG. 13.

6.3 EFFECT OF DOMAIN ORDER AND ADDITIONAL DISULFIDE BONDS ON FORMATION OF TETRAMERIC IgG-LIKE DIABODY

[00409] The effect of additional stabilization between the 'lighter' and 'heavier' polypeptide chains of the tetrameric IgG-like diabody molecule was investigated by substitution of selected residues on the polypeptide chains with cysteines. The additional cysteine residues provide for additional disulfide bonds between the 'heavier' and 'lighter' chains. Additionally, domain order on binding activity was investigated by moving the Fc domain or the hinge-Fc domain from the C-terminal end of the polypeptide chain to the N-terminus. Although the binding activity of the molecule comprising the additional disulfide bonds was not altered relative to earlier constructed diabody molecules with such bonds, transferring the Fc or hinge-Fc domain to the N-terminus of the 'heavier' polypeptide chain

comprising the diabody surprisingly improved binding affinity and/or avidity of the bispecific molecule to one or both of its target antigens.

[00410] Materials and Methods

[00411] Construction and Design of Polypeptide Molecules: Nucleic acid expression vectors were designed to produce modified versions of constructs 5, 6 and 8 presented in Example 6.2. Construct 9 (SEQ ID NO:19) and construct 10 (SEQ ID NO:20) (both shown schematically in FIG. 13) were analogous to constructs 8 and 6, with the exception that Fc domain or hinge-Fc domain, respectively, was shifted from the C-terminus of the polypeptide to the N-terminus. Additionally all Fc domains used were wild-type IgG1 Fc domains. Construct 11, SEQ ID NO:21, (shown schematically in FIG. 14) was analogous to construct 2 from Example 6.1 except that the C-terminus was designed to further comprise the sequence FNRGEC (SEQ ID NO:23). Construct 12, SEQ ID NO:22 (shown schematically in FIG. 14) was analogous to construct 5 from Example 6.2 except that the Fc domain further comprised a hinge region. Also, for constructs 11 and 12, the 2B6 VL domain and 2B6 VH domain comprised a single amino acid modification (G105C and G44C, respectively) such that a glycine in each domain was replaced by cysteine.

[00412] PCR and Expression Vector Construction: All PCR and PCR product purification protocols were as described in Example 6.1 and 6.2

[00413] Overlapping PCR: Final products were constructed, amplified and purified using methods described in example 6.1 and example 6.2.

[00414] Final products were cloned into pCIneo mammalian expression vector (Promega, Inc.) as previously described. The plasmid encoding constructs were designated as identified in Table 15:

Table 15. PLASMID CONSTRUCTS

Encoding Construct	Plasmid Designation	Insert
9	pMGX0719	Huhinge/Fc -hu3G8VL-hu2B6VH
10	pMGX0718	HuFc -hu2B6VL-hu3G8VH
11	pMGX0716	Hu2B6VL(G/C)-hu3G8VH-huhingeFC
12	pMGX0717	Hu3G8VL-hu2B6VH (G/C)-FNRGEC

[00415] Polypeptide/diabody Expression: Three separate cotransfections in to in HEK-293 cells using Lipofectamine 2000, as described in section 6.1, were performed: pMGX0669 and pMGX0719, encoding constructs 1 and 9, respectively; pMGX0669 and pMGX0718, encoding constructs 1 and 10, respectively; and pMGX0617 and pMGX0717, encoding constructs 11 and 12, respectively. Co-transfection of these plasmids was

designed to lead to the expression of a bispecific diabody (CBD) of tetravalency with IgG-like structure, immunospecific for both FcγRIIB and FcγRIIA. Following three days in culture, conditioned media was harvested. The amount of secreted product in the conditioned media was quantitated by anti IgG Fc ELISA using purified Fc as a standard. The concentrations of product in the samples was then normalized based on the quantitation, and the normalized samples used for the remaining assays.

[00416] ELISA: The binding of diabody molecules secreted into the medium was assayed by sandwich ELISA as described, *supra*. Unless indicated, CD32B was used to coat the plate, *i.e.*, as the target protein, and HRP- conjugated CD16 was used as the probe.

[00417] Western Blot: Approximately 15 ml of conditioned medium from the three above-described cotransfections were analyzed by SDS-PAGE under non-reducing conditions. One gel was stained with Simply Blue Safestain (Invitrogen) and an identical gel was transferred to PVDF membrane (Invitrogen) using standard transfer methods. After transfer, the membrane was blocked with 5% dry skim milk in 1X PBS. The membrane was then incubated in 10 ml of 1:8,000 diluted HRP conjugated Goat anti human IgG1 H+L in 2% dry skim milk 1XPBS/0.1% Tween 20 at room temperature for 1 hr with gentle agitation. Following a wash with 1X PBS/0.3% Tween 20, 2X 5 min each, then 20 min at room temperature, the membrane was developed with ECL Western blotting detection system (Amersham Biosciences) according to the manufacturer's instructions. The film was developed in X-ray processor.

[00418] Results

[00419] Conditioned media from the recombinant expression systems comprising constructs 1 and 9; constructs 1 and 10; and constructs 11 and 12 were analyzed by SDS-PAGE (under non reducing conditions) analysis and Western-blotting (using an anti-IgG as the probe). Western blot revealed that the product from the systems comprising constructs 11 and 12 or comprising constructs 9 and 1 predominately formed a single species of molecule of approximately 150 kDa (FIG. 14, lanes 3 and 2, respectively). Both of these products have engineered internal disulfide bonds between the 'lighter' and 'heavier' chains comprising the diabody. In contrast, the molecule without engineered internal disulfide bonds between the 'lighter' and 'heavier' chains, formed of constructs 10 and 1, formed at least two molecular species of molecular weights ~75 and ~100 kDa (FIG. 14, lane 1).

[00420] Despite the results of the Western Blot, each of the three products was found capable of binding both CD32A and CD16 (FIG. 15). Surprisingly, relative to the product comprising a C-terminal hinge-Fc domain (formed of constructs 11 and 12), the product from both systems wherein the Fc (or Fc-hinge) domain was at the amino terminus of the Fc

containing polypeptide chain (*i.e.*, the 'heavier' chain) (constructs 9+1 and constructs 10+1) demonstrated enhanced affinity and/or avidity to one or both of its target peptides (*i.e.* CD32B and/or CD16).

6.4 EFFECT OF INTERNAL/EXTERNAL CLEAVAGE SITE ON PROCESSING OF POLYPROTEIN PRECURSOR AND EXPRESSION OF COVALENT BISPECIFIC DIABODY; DESIGN AND CHARACTERIZATION OF BISPECIFIC DIABODY COMPRISING PORTIONS OF HUMAN IgG LAMBDA CHAIN AND HINGE DOMAIN

[00421] As described herein, the individual polypeptide chains of the diabody or diabody molecule of the invention may be expressed as a single polyprotein precursor molecule. The ability of the recombinant systems described in Examples 6.1-6.3 to properly process and express a functional CBD from such a polyprotein precursor was tested by engineering a nucleic acid to encode, both the first and second polypeptide chains of a CBD separated by an internal cleavage site, in particular, a furin cleavage site. Functional, CBD was isolated from the recombinant system comprising the polyprotein precursor molecule.

[00422] As discussed in Example 6.3, addition of the 6 C-terminal amino acids from the human kappa light chain, FNRGEC (SEQ ID NO:23), was found to stabilize diabody formation -- presumably through enhanced inter-chain interaction between the domains comprising SEQ ID NO:23 and those domains comprising an Fc domain or a hinge-Fc domain. The stabilizing effect of this lambda chain/Fc like interaction was tested in CBD wherein neither polypeptide chain comprised an Fc domain. One polypeptide chain of the diabody was engineered to comprise SEQ ID NO:23 at its C-terminus; the partner polypeptide chain was engineered to comprise the amino acid sequence VEPKSC (SEQ ID NO:79), which was derived from the hinge domain of an IgG. Comparison of this CBD to that comprised of constructs 1 and 2 (from example 6.1) revealed that the CBD comprising the domains derived from hinge domain and lambda chain exhibited slightly greater affinity to one or both of its target epitopes.

[00423] Materials and Methods

- Construction and Design of Polypeptide Molecules: Polyprotein precursor: Nucleic acid expression vectors were designed to produce 2 polyprotein precursor molecules, both represented schematically in FIG. 17. Construct 13 (SEQ ID NO:97) comprised from the N-terminus of the polypeptide chain, the VL domain of 3G8, the VH domain of 2.4G2 (which binds mCD32B), a furin cleavage site, the VL domain of 2.4G2 and the VH domain of 3G8. The nucleotide sequence encoding construct 13 is provided in SEQ

SEQ ID NO:98. Construct 14 (SEQ ID NO:99) (FIG. 17), comprised from the N-terminus of the polypeptide chain, the VL domain of 3G8, the VH domain of 2.4G2 (which binds mCD32B), a furin cleavage site, a FMD (Foot and Mouth Disease Virus Protease C3) site, the VL domain of 2.4G2 and the VH domain of 3G8. The nucleotide sequence encoding construct 14 is provided in SEQ ID NO:100.

[00425] Nucleic acid expression vectors were designed to produce modified versions of constructs 1 and 2 presented in Example 6.1. Construct 15 (SEQ ID NO:101) (FIG.17) was analogous to construct 1 (SEQ ID NO:9), presented in example 6.1, with the exception that the C-terminus of construct 15 comprised the amino acid sequence FNRGEC (SEQ ID NO:23). The nucleic acid sequence encoding construct 15 is provided in SEQ ID NO:102. Construct 16 (SEQ ID NO:103) (FIG. 17) was analogous to construct 2, presented in Example 6.1, with the exception that the C-terminus of construct 16 comprised the amino acid sequence VEPKSC (SEQ ID NO:79). The nucleic acid sequence encoding construct 16 is provided in SEQ ID NO:104.

[00426] PCR and Expression Vector Construction: All PCR and PCR product purification protocols were as described in Example 6.1 and 6.2

[00427] Overlapping PCR: Final products were constructed, amplified and purified using methods described in example 6.1 and example 6.2 with appropriate primers

[00428] Final products were cloned into pCIneo mammalian expression vector (Promega, Inc.) as previously described. The plasmid encoding constructs were designated as identified in Table 16:

Table 16. PLASMID CONSTRUCTS

Encoding Construct	Plasmid Designation	Insert
13	pMGX0750	3G8VL-2.4G2VH-Furin-2.4G2VL-3G8VH
15	pMGX0752	Hu2B6VL-Hu3G8VH-FNRGEC
16	pMGX0753	Hu3G8VL-Hu2B6VH-VEPKSC

[00429] Polypeptide/diabody Expression: One transfection and one cotransfection into HEK-293 cells using Lipofectamine 2000, as described in section 6.1, were performed: single: pMGX0750, encoding construct 13; and cotransfection: pMGX0752 and pMGX0753, encoding constructs 15 and 16, respectively. Following three days in culture, conditioned media was harvested, and secreted product affinity purified as described.

[00430] ELISA: The binding of diabody molecules secreted into the medium was assayed by sandwich ELISA as described, *supra*. Murine CD32B was used to coat the

plate, *i.e.*, as the target protein, and HRP- conjugated CD16A was used as the probe for the product of the co-transfection of constructs 15 and 16. mCD32B was used as the target protein and biotin-conjugated CD16A was used as the probe for the recombinant system comprising construct 13.

[00431] Results

[00432] Conditioned media from the recombinant expression systems comprising constructs 13 was analysed by sandwich ELISA. The ELISA assay tested the binding of the CBD for specificity to either or both of mCD32B and/or CD16 (FIG. 18). CD32B served as the target antigen and CD16A was used as the secondary probe. The positive signal in the ELISA revealed that the heterodimeric h2.4G2-h3G8 CBD produced from the polyprotein precursor had specificity for both antigens.

[00433] Similarly, the purified product generated by cotransfection of the vectors encoding constructs 15 and 16 was tested in an ELISA assay and compared to the product comprised of constructs 1 and 2 (Example 6.1). CD32B served as the target antigen and CD16A was used as the secondary probe. As with the product comprised of constructs 1 and 2, the product of constructs 15 and 16 was found to be capable of simultaneously binding CD32B and CD16A. In fact, the product of constructs 15 and 16 showed slightly enhanced affinity for one or both of the target antigens, *i.e.* CD32B or CD16A. This is perhaps due to increased stability and or fidelity (relative to a wild type VH-VL domain interaction) of the interchain association afforded by the interaction of the lambda chain region, FNRGEC (SEQ ID NO:23) and hinge region VEPKSC (SEQ ID NO:79), which is absent in the product comprised of constructs 1 and 2.

[00434] Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled. Such modifications are intended to fall within the scope of the appended claims.

[00435] All references, patent and non-patent, cited herein are incorporated herein by reference in their entireties and for all purposes to the same extent as if each individual publication or patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

CLAIMS:

1. A diabody molecule comprising a first polypeptide chain and a second polypeptide chain, said chains being covalently bonded to one another, wherein:
 - (a) said first polypeptide chain comprises an amino terminus and a carboxy terminus and from N-terminus to C-terminus:
 - (i) a Domain A comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope,
 - (ii) a Domain B comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope, and
 - (iii) an optionally present Domain C, which if present comprises a hinge region,

wherein said Domain A and said Domain B are covalently linked such that said Domain A and said Domain B do not associate to form an epitope binding site; and
 - (b) said second polypeptide chain comprises an amino terminus and a carboxy terminus and from N-terminus to C-terminus:
 - (i) a Domain D comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2),
 - (ii) a Domain E comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and
 - (iii) a Domain F that is present if Domain C is present, and which comprises a cysteine-containing polypeptide,

wherein said Domain D and said Domain E are covalently linked such that said Domain D and said Domain E do not associate to form an epitope binding site; and
 - (c)
 - (i) said Domain A and said Domain E associate with one another to form a first binding site (VL1)(VH1) that binds said first epitope; and
 - (ii) said Domain B and said Domain D associate with one another to form a second binding site (VL2)(VH2) that binds the second epitope; and
 - (iii) said first and second epitopes are different;

wherein if said optional Domain C is absent, said first polypeptide chain and said second polypeptide chain are covalently linked via a disulfide bond between at least one cysteine residue outside of said Domain A and said Domain B on the first polypeptide chain and at least one cysteine residue outside of said Domain D and said Domain E on the second polypeptide chain, which cysteine residue on the first polypeptide chain is not at the C-terminus of the first polypeptide chain and cysteine residue on the second polypeptide chain is not at the C-terminus of the second polypeptide chain.
2. The diabody molecule of claim 1, wherein said Domain C is present and comprises, from N-terminus to C-terminus, said hinge region, a CH2 domain and a CH3 domain of an Fc region.

3. A diabody molecule comprising a first polypeptide chain and a second polypeptide chain, said chains being covalently bonded to one another wherein:
 - (a) said first polypeptide chain comprises an amino terminus and a carboxy terminus and from N-terminus to C-terminus:
 - (i) a Domain C comprising, from N-terminus to C-terminus, a hinge region, a CH2 domain and a CH3 domain of an Fc region,
 - (ii) a Domain A comprising a binding region of a light chain variable domain of a first immunoglobulin (VL1) specific for a first epitope, and
 - (iii) a Domain B comprising a binding region of a heavy chain variable domain of a second immunoglobulin (VH2) specific for a second epitope,
 wherein said Domain A and said Domain B are covalently linked such that said Domain A and said Domain B do not associate with one another to form an epitope binding site;
 - (b) said second polypeptide chain comprises an amino terminus and a carboxy terminus and:
 - (i) a Domain D comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2),
 - (ii) a Domain E comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and
 - (iii) a Domain F comprising a cysteine-containing polypeptide,
 wherein said Domain D is N-terminal to said Domain E and said Domain D and said Domain E are covalently linked such that said Domain D and said Domain E do not associate with one another to form an epitope binding site; and
 - (c)
 - (i) said Domain A and said Domain E associate with one another to form a first binding site (VL1)(VH1) that binds said first epitope;
 - (ii) said Domain B and said Domain D associate with one another to form a second binding site (VL2)(VH2) that binds said second epitope; and
 - (iii) said first and second epitopes are different.
4. The diabody molecule of claim 1, wherein said Domain C is absent and:
 - (1) said Domain A and said Domain E associate to form a first binding site (VL1)(VH1) that binds said first epitope, which epitope binding site is specific for CD32B; and
 - (2) said Domain B and said Domain D associate to form a second binding site (VL2)(VH2) that binds said second epitope, which epitope binding site is specific for CD16.
5. The diabody molecule of any of claims 1-3, wherein said Domain C is present and the first polypeptide chain and the second polypeptide chain are covalently linked via at least one disulfide bond between at least one cysteine residue outside of said Domain A and said Domain B on the first polypeptide chain and at least one cysteine residue outside of said Domain D and said Domain E on the

- second polypeptide chain.
6. The diabody molecule of any of claims 1-5 wherein said molecule comprises at least one amino acid modification of said Domain A relative to a wild-type Domain A and at least one amino acid modification of said Domain E relative to a wild-type Domain E, which said at least one amino acid modification in each of said Domain A and said Domain E comprises a substitution with cysteine such that said first polypeptide chain and said second polypeptide chain are covalently linked via a disulfide bond between said substituted cysteine residue in said Domain A and said substituted cysteine residue in said Domain E.
 7. The diabody molecule of any of claims 1-6, wherein the first immunoglobulin or second immunoglobulin is a human or humanized immunoglobulin, which human or humanized immunoglobulin is an IgA, IgE, IgD, IgG or IgM.
 8. The diabody molecule of claim 7, wherein the human or humanized immunoglobulin is an IgG, which IgG is selected from the list consisting of IgG1, IgG2, IgG3 and IgG4.
 9. The diabody molecule of any of claims 1-8, wherein the Fc domain is a human or humanized Fc domain.
 10. The diabody molecule of any of claims 1-3 and 5-9, wherein said Domain C is present and wherein at least one epitope binding site is specific for an FcγRI, FcγRII or FcγRIII receptor.
 11. The diabody molecule of claim 10, wherein the Fcγ receptor is a FcγRIII receptor, which FcγRIII receptor is FcγRIIIA (CD16A) receptor or FcγRIIB (CD16B) receptor.
 12. The diabody molecule of claim 11, wherein the FcγRIII receptor is FcγRIIIA (CD16A) receptor.
 13. The diabody molecule of claim 10, wherein the Fcγ receptor is a FcγRII receptor, which FcγRII receptor is FcγRIIA (CD32A) receptor or FcγRIIB (CD32B) receptor.
 14. The diabody molecule of claim 13, wherein the FcγRII receptor is the FcγRIIB (CD32B) receptor.
 15. The diabody molecule of any of claims 1-3 and 5-14, wherein said Domain C is present and wherein at least one epitope binding site is specific for a pathogenic antigen.
 16. The diabody molecule of any of claims 1-3 and 5-14, wherein said Domain C is present and wherein at least one epitope binding site is specific for a toxin or a drug.
 17. The diabody molecule of claim 10, wherein said diabody has an epitope binding site specific for CD32B and an epitope binding site specific for CD16A.

18. The diabody molecule of any of claims 1-3, wherein said Domain C is present and wherein said Domain C comprises a variant Fc region, which variant Fc region comprises at least one amino acid modification relative to the wild type Fc region.
19. The diabody molecule of any of claims 1-3 and 5-18, wherein said Domain C is present and wherein said diabody is a dimer, said dimer comprising
 - (a) a first monomer comprising a set of said first polypeptide chain and said second polypeptide chain, said first polypeptide chain having said Domains A, B and C; and
 - (b) a second monomer comprising a set of said first polypeptide chain and said second polypeptide chain, said first polypeptide chain having said Domains A, B and C;wherein said first and second monomers are covalently linked via at least one disulfide bond between at least one cysteine residue in said Domain C of the first polypeptide chain of each monomer.
20. A nucleic acid molecule comprising a nucleotide sequence encoding the first polypeptide chain of the diabody molecule of any of claims 1-19.
21. A nucleic acid molecule comprising a nucleotide sequence encoding the second polypeptide chain of the diabody molecule of any of claims 1-19.
22. A host cell comprising the nucleic acid molecule of claim 20 or 21.
23. A method of treating a disease or disorder comprising administering to a patient in need thereof the diabody molecule of claim 15.
24. The use of the diabody molecule claim 23, wherein said disease or disorder is cancer.
25. The use of the diabody molecule claim 23, wherein said disease or disorder is an infectious disease.
26. The diabody molecule of claim 3, wherein said second polypeptide chain comprises from N-terminus to C-terminus:
 - (i) said Domain D comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2),
 - (ii) said Domain E comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and
 - (iii) said Domain F comprising said cysteine-containing polypeptide.
27. The diabody molecule of claim 3, wherein said second polypeptide chain comprises from N-terminus to C-terminus:
 - (i) said Domain F comprising said cysteine-containing polypeptide.
 - (ii) said Domain E comprising a binding region of a heavy chain variable domain of the first immunoglobulin (VH1), and
 - (iii) said Domain D comprising a binding region of a light chain variable domain of the second immunoglobulin (VL2).

28. The diabody molecule of any of claims 1-3, substantially as hereinbefore described and with reference to any one of the Examples.

216		226	
+	-----+	-----+	-----+
E	P K S C D - - K T H T C P P - - - - -		IGG1
E	R K C C V - - E - - - - - C P P - - - - -		-IGG2
E	L K T P L G D T T H T C P R C P E P K S C D T P		IGG3
E	S K Y G P - - P - - - - C P S - - - - -		-IGG4
		230	
		-----+	
		- - - - -	C P
		- - - - -	C P
		P P C P R C P E P K S C D T P P P C P R C P	
		- - - - -	C P

FIG. 1A

231
|
+++++
A P E L L G G IgG1
A P P V A G G IgG2
A P E L L G G IgG3
A P E L L G G IgG4

240 260
| |
+++++
P S V F L F P P K P K D T L M I S R T P E V T C V V V D V S H E D P E V K F N W
P S V F L F P P K P K D T L M I S R T P E V T C V V V D V S H E D P E V Q F N W
P S V F L F P P K P K D T L M I S R T P E V T C V V V D V S H E D P E V Q F N W
P S V F L F P P K P K D T L M I S R T P E V T C V V V D V S Q E D P E V Q F N W

280 300
| |
+++++
Y V D G V E V H N A K T K P R E E Q Y N S T Y R V V S V L T V L H Q D W L N G K
Y V D G V E V H N A K T K P R E E Q F N S T F R V V S V L T V V H Q D W L N G K
Y V D G V Q V H N A K T K P R E E Q Q F N S T F R V V S V L T V L H Q N W L D G K
Y V D G V E V H N A K T K P R E E Q F N S T Y R V V S V L T V L H Q D W L N G K

320 340
| |
+++++
E Y K C K V S N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R D E
E Y K C K V S N K Q L P A P I E K T I S K T K G Q P R E P Q V Y T L P P S R E E
E Y K C K V S N K A L P A P I E K T I S K T K G Q P R E P Q V Y T L P P S R E E
E Y K C K V S N K G L P S S I E K T I S K A K G Q P R E P Q V Y T L P P S Q E E

360 380
| |
~~~~~  
L T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V  
M T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P M  
M T K N Q V S L T C L V K G F Y P S D I A V E W E S S G Q P E N N Y N T T P P M  
M T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V

400 420  
| |  
~~~~~  
L D S D G S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T
L D S D G S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T
L D S D G S F F L Y S K L T V D K S R W Q Q G N F F S C S V M H E A L H N R E T
L D S D G S F F L Y S R L T V D K S R W Q E G N V F S C S V M H E A L H N H Y T

440
|
~~~~~  
Q K S L S L S P G K  
Q K S L S L S P G K  
Q K S L S L S P G K  
Q K S L S L S L G K

FIG. 1B




FIG. 2




**FIG. 3**




**FIG. 4A**




FIG. 4B




**FIG. 5**




**FIG. 6A**




**FIG. 6B**




**FIG. 6C**

9/23




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D




FIG. 7E






FIG. 9





FIG. 10



**FIG. 12**




FIG. 13




FIG. 14

M 1 2 3




**FIG. 15A**

3 2 1 M




**FIG. 15B**




FIG. 16




FIG. 17




**FIG. 18**




**FIG. 19**

SEQUENCE LISTING

<110> MacroGenics, Inc.

<120> COVALENT DIABODIES AND USES THEREOF

<130> 11183-058-228

<140>

<141>

<150> 60/671,657

<151> 2005-04-15

<160> 104

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 13

<212> PRT

<213> Homo sapiens

<220>

<223> human IgG1 hinge

<400> 1

|     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Pro | Lys | Ser | Cys | Asp | Lys | Thr | His | Thr | Cys | Pro | Pro |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |

<210> 2

<211> 10

<212> PRT

<213> Homo sapiens

<220>

<223> human IgG2 hinge

<400> 2

|     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Arg | Lys | Cys | Cys | Val | Glu | Cys | Pro | Pro |
| 1   |     |     |     | 5   |     |     |     | 10  |     |

<210> 3

<211> 60

<212> PRT

<213> Homo sapiens

<220>

<223> human IgG3 hinge

<400> 3

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Leu | Lys | Thr | Pro | Leu | Gly | Asp | Thr | Thr | His | Thr | Cys | Pro | Arg | Cys |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |     |
| Pro | Glu | Pro | Lys | Ser | Cys | Asp | Thr | Pro | Pro | Pro | Cys | Pro | Arg | Cys | Pro |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Glu | Pro | Lys | Ser | Cys | Asp | Thr | Pro | Pro | Pro | Cys | Pro | Arg | Cys | Pro | Glu |
|     |     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |

Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg  
50 55 60

<210> 4  
<211> 10  
<212> PRT  
<213> Homo sapiens

<220>  
<223> human IgG4 hinge

<400> 4  
Glu Ser Lys Tyr Gly Pro Pro Cys Pro Ser  
1 5 10

<210> 5  
<211> 217  
<212> PRT  
<213> Homo sapiens

<220>  
<223> human IgG1 Fc domain

<400> 5  
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys  
1 5 10 15  
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val  
20 25 30  
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr  
35 40 45  
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu  
50 55 60  
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His  
65 70 75 80  
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys  
85 90 95  
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln  
100 105 110  
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu  
115 120 125  
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro  
130 135 140  
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn  
145 150 155 160  
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu  
165 170 175  
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val  
180 185 190  
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln  
195 200 205  
Lys Ser Leu Ser Leu Ser Pro Gly Lys  
210 215

<210> 6  
<211> 216  
<212> PRT  
<213> Homo sapiens

<220>

<223> human IgG2 Fc domain

<400> 6

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Pro | Pro | Val | Ala | Gly | Pro | Ser | Val | Phe | Leu | Phe | Pro | Pro | Lys | Pro |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Lys | Asp | Thr | Leu | Met | Ile | Ser | Arg | Thr | Pro | Glu | Val | Thr | Cys | Val | Val |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Val | Asp | Val | Ser | His | Glu | Asp | Pro | Glu | Val | Gln | Phe | Asn | Trp | Tyr | Val |
|     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |
| Asp | Gly | Val | Glu | Val | His | Asn | Ala | Lys | Thr | Lys | Pro | Arg | Glu | Glu | Gln |
|     | 50  |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |
| Phe | Asn | Ser | Thr | Phe | Arg | Val | Val | Ser | Val | Leu | Thr | Val | Val | His | Gln |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     | 80  |     |
| Asp | Trp | Leu | Asn | Gly | Lys | Glu | Tyr | Lys | Cys | Lys | Val | Ser | Asn | Lys | Gly |
|     |     |     | 85  |     |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Leu | Pro | Ala | Pro | Ile | Glu | Lys | Thr | Ile | Ser | Lys | Thr | Lys | Gly | Gln | Pro |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     | 110 |     |     |
| Arg | Glu | Pro | Gln | Val | Tyr | Thr | Leu | Pro | Pro | Ser | Arg | Glu | Glu | Met | Thr |
|     |     | 115 |     |     |     |     | 120 |     |     |     |     | 125 |     |     |     |
| Lys | Asn | Gln | Val | Ser | Leu | Thr | Cys | Leu | Val | Lys | Gly | Phe | Tyr | Pro | Ser |
|     | 130 |     |     |     |     | 135 |     |     |     |     | 140 |     |     |     |     |
| Asp | Ile | Ala | Val | Glu | Trp | Glu | Ser | Asn | Gly | Gln | Pro | Glu | Asn | Asn | Tyr |
| 145 |     |     |     |     | 150 |     |     |     |     | 155 |     |     |     |     | 160 |
| Lys | Thr | Thr | Pro | Pro | Met | Leu | Asp | Ser | Asp | Gly | Ser | Phe | Phe | Leu | Tyr |
|     |     |     |     | 165 |     |     |     |     | 170 |     |     |     |     | 175 |     |
| Ser | Lys | Leu | Thr | Val | Asp | Lys | Ser | Arg | Trp | Gln | Gln | Gly | Asn | Val | Phe |
|     |     |     | 180 |     |     |     |     | 185 |     |     |     |     | 190 |     |     |
| Ser | Cys | Ser | Val | Met | His | Glu | Ala | Leu | His | Asn | His | Tyr | Thr | Gln | Lys |
|     | 195 |     |     |     |     |     | 200 |     |     |     |     | 205 |     |     |     |
| Ser | Leu | Ser | Leu | Ser | Pro | Gly | Lys |     |     |     |     |     |     |     |     |
|     | 210 |     |     |     |     | 215 |     |     |     |     |     |     |     |     |     |

<210> 7

<211> 217

<212> PRT

<213> Homo sapiens

<220>

<223> human IgG3 Fc domain

<400> 7

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Pro | Glu | Leu | Leu | Gly | Gly | Pro | Ser | Val | Phe | Leu | Phe | Pro | Pro | Lys |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Pro | Lys | Asp | Thr | Leu | Met | Ile | Ser | Arg | Thr | Pro | Glu | Val | Thr | Cys | Val |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Val | Val | Asp | Val | Ser | His | Glu | Asp | Pro | Glu | Val | Gln | Phe | Lys | Trp | Tyr |
|     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |
| Val | Asp | Gly | Val | Gln | Val | His | Asn | Ala | Lys | Thr | Lys | Pro | Arg | Glu | Gln |
|     | 50  |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |
| Gln | Phe | Asn | Ser | Thr | Phe | Arg | Val | Val | Ser | Val | Leu | Thr | Val | Leu | His |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     | 80  |     |
| Gln | Asn | Trp | Leu | Asp | Gly | Lys | Glu | Tyr | Lys | Cys | Lys | Val | Ser | Asn | Lys |
|     |     |     | 85  |     |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Ala | Leu | Pro | Ala | Pro | Ile | Glu | Lys | Thr | Ile | Ser | Lys | Thr | Lys | Gly | Gln |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     | 110 |     |     |
| Pro | Arg | Glu | Pro | Gln | Val | Tyr | Thr | Leu | Pro | Pro | Ser | Arg | Glu | Glu | Met |
|     |     | 115 |     |     |     |     | 120 |     |     |     |     | 125 |     |     |     |
| Thr | Lys | Asn | Gln | Val | Ser | Leu | Thr | Cys | Leu | Val | Lys | Gly | Phe | Tyr | Pro |
|     | 130 |     |     |     |     | 135 |     |     |     |     |     | 140 |     |     |     |

```

Ser Asp Ile Ala Val Glu Trp Glu Ser Ser Gly Gln Pro Glu Asn Asn
145          150          155          160
Tyr Asn Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
          165          170          175
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Ile
          180          185          190
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln
          195          200          205
Lys Ser Leu Ser Leu Ser Pro Gly Lys
210          215

```

<210> 8  
 <211> 217  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <223> human IgG4 Fc domain

```

<400> 8
Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
1          5          10          15
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
          20          25          30
Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
          35          40          45
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
          50          55          60
Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
65          70          75          80
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
          85          90          95
Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
          100          105          110
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met
          115          120          125
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
          130          135          140
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
145          150          155          160
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
          165          170          175
Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val
          180          185          190
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
          195          200          205
Lys Ser Leu Ser Leu Ser Leu Gly Lys
210          215

```

<210> 9  
 <211> 237  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody Construct 1

```

<400> 9
Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys

```

```

      1             5             10             15
Glu Lys Val Thr Phe Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn
      20             25             30
Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
      35             40             45
Lys Glu Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly
      50             55             60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala
      65             70             75             80
Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe
      85             90             95
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Gly Gly Gly Ser Gly
      100             105             110
Gly Gly Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys
      115             120             125
Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu
      130             135             140
Ser Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys
      145             150             155             160
Ala Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp Lys Arg Tyr
      165             170             175
Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys
      180             185             190
Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala
      195             200             205
Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly
      210             215             220
Gln Gly Thr Leu Val Thr Val Ser Ser Leu Gly Gly Cys
      225             230             235

```

<210> 10  
 <211> 8  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> linker

<400> 10  
 Gly Gly Gly Ser Gly Gly Gly Gly  
 1 5

<210> 11  
 <211> 244  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody Construct 2

<400> 11  
 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
 1 5 10 15  
 Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp  
 20 25 30  
 Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
 35 40 45  
 Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro Asp

|                                                                 |     |     |
|-----------------------------------------------------------------|-----|-----|
| 50                                                              | 55  | 60  |
| Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser |     |     |
| 65                                                              | 70  | 75  |
| Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn |     | 80  |
|                                                                 | 85  | 90  |
| Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Gly |     | 95  |
|                                                                 | 100 | 105 |
| Gly Gly Ser Gly Gly Gly Gly Gln Val Gln Leu Val Gln Ser Gly Ala |     | 110 |
|                                                                 | 115 | 120 |
| Glu Val Lys Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser |     | 125 |
|                                                                 | 130 | 135 |
| Gly Tyr Thr Phe Thr Asn Tyr Trp Ile His Trp Val Arg Gln Ala Pro |     | 140 |
| 145                                                             | 150 | 155 |
| Gly Gln Gly Leu Glu Trp Ile Gly Val Ile Asp Pro Ser Asp Thr Tyr |     | 160 |
|                                                                 | 165 | 170 |
| Pro Asn Tyr Asn Lys Lys Phe Lys Gly Arg Val Thr Met Thr Val Val |     | 175 |
|                                                                 | 180 | 185 |
| Val Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu Arg Ser Asp |     | 190 |
|                                                                 | 195 | 200 |
| Asp Thr Ala Val Tyr Tyr Cys Ala Arg Asn Gly Asp Ser Asp Tyr Tyr |     | 205 |
| 210                                                             | 215 | 220 |
| Ser Gly Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser |     | 225 |
| 225                                                             | 230 | 235 |
| Leu Gly Gly Cys                                                 |     | 240 |

<210> 12  
 <211> 240  
 <212> PRT

<213> Artificial Sequence

<220>  
 <223> Diabody Construct 3

<400> 12

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys |     |
| 1                                                               | 5   |
| Glu Lys Val Thr Phe Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn |     |
|                                                                 | 20  |
| Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile |     |
|                                                                 | 35  |
| Lys Glu Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly |     |
| 50                                                              | 55  |
| Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala |     |
| 65                                                              | 70  |
| Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe |     |
|                                                                 | 85  |
| Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Gly Gly Gly Ser Gly |     |
|                                                                 | 100 |
| Gly Gly Gly Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys |     |
|                                                                 | 115 |
| Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe |     |
| 130                                                             | 135 |
| Thr Asn Tyr Trp Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu |     |
| 145                                                             | 150 |
| Glu Trp Ile Gly Val Ile Asp Pro Ser Asp Thr Tyr Pro Asn Tyr Asn |     |
|                                                                 | 165 |
| Lys Lys Phe Lys Gly Arg Val Thr Met Thr Val Val Val Ser Thr Ser |     |
|                                                                 | 180 |
|                                                                 | 185 |
|                                                                 | 190 |

```

Thr Ala Tyr Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val
      195                      200                205
Tyr Tyr Cys Ala Arg Asn Gly Asp Ser Asp Tyr Tyr Ser Gly Met Asp
      210                      215                220
Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Leu Gly Gly Cys
225                      230                235                240

```

<210> 13  
 <211> 241  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody Construct 4

```

<400> 13
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1                      5                      10                15
Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp
      20                      25                      30
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
      35                      40                      45
Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro Asp
      50                      55                      60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65                      70                      75                80
Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
      85                      90                      95
Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Gly
      100                     105                     110
Gly Gly Ser Gly Gly Gly Gly Gln Val Thr Leu Arg Glu Ser Gly Pro
      115                     120                     125
Ala Leu Val Lys Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser
130                     135                     140
Gly Phe Ser Leu Ser Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln
145                     150                     155                160
Pro Pro Gly Lys Ala Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp
      165                     170                     175
Asp Lys Arg Tyr Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys
      180                     185                     190
Asp Thr Ser Lys Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro
      195                     200                     205
Val Asp Thr Ala Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe
      210                     215                     220
Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Leu Gly Gly
225                     230                     235                240
Cys

```

<210> 14  
 <211> 470  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody Construct 5

```

<400> 14
Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys

```

|                                                                 |     |     |     |
|-----------------------------------------------------------------|-----|-----|-----|
| 1                                                               | 5   | 10  | 15  |
| Glu Lys Val Thr Phe Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn |     |     |     |
|                                                                 | 20  | 25  | 30  |
| Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile |     |     |     |
|                                                                 | 35  | 40  | 45  |
| Lys Glu Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly |     |     |     |
|                                                                 | 50  | 55  | 60  |
| Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala |     |     |     |
| 65                                                              | 70  | 75  | 80  |
| Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe |     |     |     |
|                                                                 | 85  | 90  | 95  |
| Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Gly Gly Gly Ser Gly |     |     |     |
|                                                                 | 100 | 105 | 110 |
| Gly Gly Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys |     |     |     |
|                                                                 | 115 | 120 | 125 |
| Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu |     |     |     |
| 130                                                             | 135 | 140 |     |
| Ser Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys |     |     |     |
| 145                                                             | 150 | 155 | 160 |
| Ala Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr |     |     |     |
|                                                                 | 165 | 170 | 175 |
| Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys |     |     |     |
|                                                                 | 180 | 185 | 190 |
| Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala |     |     |     |
|                                                                 | 195 | 200 | 205 |
| Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly |     |     |     |
| 210                                                             | 215 | 220 |     |
| Gln Gly Thr Leu Val Thr Val Ser Ser Leu Gly Gly Cys Val Glu Pro |     |     |     |
| 225                                                             | 230 | 235 | 240 |
| Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu |     |     |     |
|                                                                 | 245 | 250 | 255 |
| Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp |     |     |     |
|                                                                 | 260 | 265 | 270 |
| Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp |     |     |     |
| 275                                                             | 280 | 285 |     |
| Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly |     |     |     |
| 290                                                             | 295 | 300 |     |
| Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn |     |     |     |
| 305                                                             | 310 | 315 | 320 |
| Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp |     |     |     |
|                                                                 | 325 | 330 | 335 |
| Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro |     |     |     |
|                                                                 | 340 | 345 | 350 |
| Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu |     |     |     |
|                                                                 | 355 | 360 | 365 |
| Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn |     |     |     |
| 370                                                             | 375 | 380 |     |
| Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile |     |     |     |
| 385                                                             | 390 | 395 | 400 |
| Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr |     |     |     |
|                                                                 | 405 | 410 | 415 |
| Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys |     |     |     |
|                                                                 | 420 | 425 | 430 |
| Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys |     |     |     |
|                                                                 | 435 | 440 | 445 |
| Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu |     |     |     |
| 450                                                             | 455 | 460 |     |
| Ser Leu Ser Pro Gly Lys                                         |     |     |     |
| 465                                                             | 470 |     |     |

<210> 15  
 <211> 477  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody Construct 6

<400> 15  
 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
 1 5 10 15  
 Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp  
 20 25 30  
 Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
 35 40 45  
 Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro Asp  
 50 55 60  
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser  
 65 70 75 80  
 Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn  
 85 90 95  
 Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Gly  
 100 105 110  
 Gly Gly Ser Gly Gly Gly Gly Gln Val Gln Leu Val Gln Ser Gly Ala  
 115 120 125  
 Glu Val Lys Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser  
 130 135 140  
 Gly Tyr Thr Phe Thr Asn Tyr Trp Ile His Trp Val Arg Gln Ala Pro  
 145 150 155 160  
 Gly Gln Gly Leu Glu Trp Ile Gly Val Ile Asp Pro Ser Asp Thr Tyr  
 165 170 175  
 Pro Asn Tyr Asn Lys Lys Phe Lys Gly Arg Val Thr Met Thr Val Val  
 180 185 190  
 Val Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu Arg Ser Asp  
 195 200 205  
 Asp Thr Ala Val Tyr Tyr Cys Ala Arg Asn Gly Asp Ser Asp Tyr Tyr  
 210 215 220  
 Ser Gly Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 225 230 235 240  
 Leu Gly Gly Cys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys  
 245 250 255  
 Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu  
 260 265 270  
 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu  
 275 280 285  
 Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys  
 290 295 300  
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys  
 305 310 315 320  
 Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu  
 325 330 335  
 Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys  
 340 345 350  
 Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys  
 355 360 365  
 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser  
 370 375 380  
 Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys  
 385 390 395 400  
 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln  
 405 410 415

```

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
      420      425      430
Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln
      435      440      445
Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
      450      455      460
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
465      470      475

```

```

<210> 16
<211> 243
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Diabody Construct 7

```

```

<400> 16
Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys
 1      5      10      15
Glu Lys Val Thr Phe Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn
      20      25      30
Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
      35      40      45
Lys Glu Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly
 50      55      60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala
65      70      75      80
Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe
      85      90      95
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Gly Gly Gly Ser Gly
      100     105     110
Gly Gly Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys
      115     120     125
Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu
130     135     140
Ser Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys
145     150     155     160
Ala Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp Lys Arg Tyr
      165     170     175
Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys
      180     185     190
Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala
      195     200     205
Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly
      210     215     220
Gln Gly Thr Leu Val Thr Val Ser Ser Leu Gly Gly Cys Phe Asn Arg
225     230     235     240
Gly Glu Cys

```

```

<210> 17
<211> 10
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Linker plus C-terminus of human IgG-kappa chain

```

&lt;400&gt; 17

Leu Gly Gly Cys Phe Asn Arg Gly Glu Cys  
 1 5 10

&lt;210&gt; 18

&lt;211&gt; 477

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Diabody Construct 8

&lt;400&gt; 18

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
 1 5 10 15  
 Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp  
 20 25 30  
 Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
 35 40 45  
 Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro Asp  
 50 55 60  
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser  
 65 70 75 80  
 Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn  
 85 90 95  
 Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Gly  
 100 105 110  
 Gly Gly Ser Gly Gly Gly Gly Gln Val Gln Leu Val Gln Ser Gly Ala  
 115 120 125  
 Glu Val Lys Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser  
 130 135 140  
 Gly Tyr Thr Phe Thr Asn Tyr Trp Ile His Trp Val Arg Gln Ala Pro  
 145 150 155 160  
 Gly Gln Gly Leu Glu Trp Ile Gly Val Ile Asp Pro Ser Asp Thr Tyr  
 165 170 175  
 Pro Asn Tyr Asn Lys Lys Phe Lys Gly Arg Val Thr Met Thr Val Val  
 180 185 190  
 Val Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu Arg Ser Asp  
 195 200 205  
 Asp Thr Ala Val Tyr Tyr Cys Ala Arg Asn Gly Asp Ser Asp Tyr Tyr  
 210 215 220  
 Ser Gly Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 225 230 235 240  
 Leu Gly Gly Cys Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys  
 245 250 255  
 Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu  
 260 265 270  
 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu  
 275 280 285  
 Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys  
 290 295 300  
 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys  
 305 310 315 320  
 Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu  
 325 330 335  
 Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys  
 340 345 350  
 Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys  
 355 360 365  
 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser

```

      370                      375                      380
Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
385                      390                      395                      400
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
      405                      410                      415
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
      420                      425                      430
Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln
      435                      440                      445
Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
      450                      455                      460
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
465                      470                      475

```

```

<210> 19
<211> 485
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Diabody Construct 9

```

```

<400> 19
Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
1      5      10      15
Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
20      25      30
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
35      40      45
Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
50      55      60
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
65      70      75      80
Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
85      90      95
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
100     105     110
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
115     120     125
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
130     135     140
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
145     150     155     160
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
165     170     175
Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
180     185     190
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
195     200     205
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
210     215     220
Lys Ser Leu Ser Leu Ser Pro Gly Lys Gly Gly Gly Ser Gly Gly Gly
225     230     235     240
Gly Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu
245     250     255
Gly Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe
260     265     270
Asp Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro
275     280     285
Pro Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro

```

|                     |                     |                         |
|---------------------|---------------------|-------------------------|
| 290                 | 295                 | 300                     |
| Asp Arg Phe Ser Gly | Ser Gly Ser Gly Thr | Asp Phe Thr Leu Thr Ile |
| 305                 | 310                 | 315                     |
| Ser Ser Leu Gln Ala | Glu Asp Val Ala Val | Tyr Tyr Cys Gln Gln Ser |
| 325                 | 330                 | 335                     |
| Asn Glu Asp Pro Tyr | Thr Phe Gly Gln Gly | Thr Lys Leu Glu Ile Lys |
| 340                 | 345                 | 350                     |
| Gly Gly Gly Ser Gly | Gly Gly Gly Gln Val | Gln Leu Val Gln Ser Gly |
| 355                 | 360                 | 365                     |
| Ala Glu Val Lys Lys | Pro Gly Ala Ser Val | Lys Val Ser Cys Lys Ala |
| 370                 | 375                 | 380                     |
| Ser Gly Tyr Thr Phe | Thr Asn Tyr Trp Ile | His Trp Val Arg Gln Ala |
| 385                 | 390                 | 395                     |
| Pro Gly Gln Gly Leu | Glu Trp Ile Gly Val | Ile Asp Pro Ser Asp Thr |
| 405                 | 410                 | 415                     |
| Tyr Pro Asn Tyr Asn | Lys Lys Phe Lys Gly | Arg Val Thr Met Thr Val |
| 420                 | 425                 | 430                     |
| Val Val Ser Thr Ser | Thr Ala Tyr Met Glu | Leu Arg Ser Leu Arg Ser |
| 435                 | 440                 | 445                     |
| Asp Asp Thr Ala Val | Tyr Tyr Cys Ala Arg | Asn Gly Asp Ser Asp Tyr |
| 450                 | 455                 | 460                     |
| Tyr Ser Gly Met Asp | Tyr Trp Gly Gln Gly | Thr Thr Val Thr Val Ser |
| 465                 | 470                 | 475                     |
| Ser Leu Gly Gly Cys |                     | 480                     |
| 485                 |                     |                         |

<210> 20  
 <211> 469  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody Construct 10

<400> 20

|                                                                 |  |
|-----------------------------------------------------------------|--|
| Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys |  |
| 1 5 10 15                                                       |  |
| Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val |  |
| 20 25 30                                                        |  |
| Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr |  |
| 35 40 45                                                        |  |
| Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu |  |
| 50 55 60                                                        |  |
| Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His |  |
| 65 70 75 80                                                     |  |
| Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys |  |
| 85 90 95                                                        |  |
| Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln |  |
| 100 105 110                                                     |  |
| Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu |  |
| 115 120 125                                                     |  |
| Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro |  |
| 130 135 140                                                     |  |
| Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn |  |
| 145 150 155 160                                                 |  |
| Tyr Lys Thr Thr Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu     |  |
| 165 170 175                                                     |  |
| Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val |  |
| 180 185 190                                                     |  |
| Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln |  |

[illegible]

```
<210> 21
<211> 466
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Diabody Construct 11
```

```

<400> 21
Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys
 1           5           10           15
Glu Lys Val Thr Phe Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn
          20           25           30
Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
          35           40           45
Lys Glu Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly
 50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala
65           70           75           80
Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe
          85           90           95

Thr Phe Gly Cys Gly Thr Lys Val Glu Ile Lys Gly Gly Gly Ser Gly
          100           105           110

```

Gly Gly Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys  
115 120 125  
Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu  
130 135 140  
Ser Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys  
145 150 155 160  
Ala Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr  
165 170 175  
Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys  
180 185 190  
Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala  
195 200 205  
Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly  
210 215 220  
Gln Gly Thr Leu Val Thr Val Ser Ser Val Glu Pro Lys Ser Cys Asp  
225 230 235 240  
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly  
245 250 255  
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile  
260 265 270  
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu  
275 280 285  
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His  
290 295 300  
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg  
305 310 315 320  
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys  
325 330 335  
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu  
340 345 350  
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr  
355 360 365  
Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu  
370 375 380  
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp  
385 390 395 400  
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val  
405 410 415  
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp  
420 425 430  
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His  
435 440 445  
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro  
450 455 460  
Gly Lys  
465

<210> 22  
<211> 246  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Diabody Construct 12

<400> 22  
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly  
1 5 10 15  
Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp

<220>  
<223> C-terminus of human IgG Kappa chain

<400> 23  
Phe Asn Arg Gly Glu Cys  
1 5

```
<210> 24
<211> 5
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> 2B6 Heavy chain variable region - CDR1
```

```
<400> 24
Asn Tyr Trp Ile His
  1                               5
```

```
<210> 25
<211> 5
<212> PRT
<213> Artificial Sequence
```

<220>  
<223> 3H7 Heavy Chian Variable region - CDR1

<400> 25

Asp Ala Trp Met Asp  
1 5

<210> 26  
<211> 17  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Heavy chain variable region - CDR2

<400> 26  
Val Ile Asp Pro Ser Asp Thr Tyr Pro Asn Tyr Asn Lys Lys Phe Lys  
1 5 10 15  
Gly

<210> 27  
<211> 19  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 3H7 Heavy Chian Variable region - CDR2

<400> 27  
Glu Ile Arg Asn Lys Ala Asn Asn Leu Ala Thr Tyr Tyr Ala Glu Ser  
1 5 10 15  
Val Lys Gly

<210> 28  
<211> 12  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Heavy chain variable region - CDR3

<400> 28  
Asn Gly Asp Ser Asp Tyr Tyr Ser Gly Met Asp Tyr  
1 5 10

<210> 29  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 3H7 Heavy Chian Variable region - CDR3

<400> 29  
Tyr Ser Pro Phe Ala Tyr  
1 5

<210> 30  
<400> 30  
000

<210> 31  
<400> 31  
000

<210> 32  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Light chain variable region - CDR1

<400> 32  
Arg Thr Ser Gln Ser Ile Gly Thr Asn Ile His  
1 5 10

<210> 33  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 3H7 Light Chian Variable region - CDR1

<400> 33  
Arg Ala Ser Gln Glu Ile Ser Gly Tyr Leu Ser  
1 5 10

<210> 34  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Light chain variable region - CDR2

<400> 34  
Asn Val Ser Glu Ser Ile Ser  
1 5

<210> 35  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Light chain variable region - CDR2

<400> 35  
Tyr Val Ser Glu Ser Ile Ser  
1 5

<210> 36  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Light chain variable region - CDR2

<400> 36  
Tyr Ala Ser Glu Ser Ile Ser  
1 5

<210> 37  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 3H7 Light Chian Variable region - CDR2

<400> 37  
Ala Ala Ser Thr Leu Asp Ser  
1 5

<210> 38  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 2B6 Light chain variable region - CDR3

<400> 38  
Gln Gln Ser Asn Thr Trp Pro Phe Thr  
1 5

<210> 39  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> 3H7 Light Chian Variable region - CDR3

<400> 39  
Leu Gln Tyr Val Ser Tyr Pro Tyr Thr  
1 5

<210> 40  
<211> 121  
<212> PRT  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; 2B6 heavy chain variable region

&lt;400&gt; 40

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1           5           10           15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
      20           25           30
Trp Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
      35           40           45
Gly Val Ile Asp Pro Ser Asp Thr Tyr Pro Asn Tyr Asn Lys Lys Phe
      50           55           60
Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
      65           70           75           80
Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
      85           90           95
Ala Arg Asn Gly Asp Ser Asp Tyr Tyr Ser Gly Met Asp Tyr Trp Gly
      100          105          110
Gln Gly Thr Thr Val Thr Val Ser Ser
      115          120

```

&lt;210&gt; 41

&lt;211&gt; 107

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; 2B6 light chain variable region - Hu2B6VL-1

&lt;400&gt; 41

```

Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys
 1           5           10           15
Glu Lys Val Thr Ile Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn
      20           25           30
Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
      35           40           45
Lys Asn Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly
      50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala
      65           70           75           80
Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe
      85           90           95
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
      100          105

```

&lt;210&gt; 42

&lt;211&gt; 107

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Humanized 2B6 light chain variable region - Hu2B6VL-2

&lt;400&gt; 42

```

Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys
 1           5           10           15
Glu Lys Val Thr Ile Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn

```

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ile | His | Trp | Tyr | Gln | Gln | Lys | Pro | Asp | Gln | Ser | Pro | Lys | Leu | Leu | Ile |
|     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |
| Lys | Tyr | Val | Ser | Glu | Ser | Ile | Ser | Gly | Val | Pro | Ser | Arg | Phe | Ser | Gly |
|     |     | 50  |     |     |     | 55  |     |     |     | 60  |     |     |     |     |     |
| Ser | Gly | Ser | Gly | Thr | Asp | Phe | Thr | Leu | Thr | Ile | Asn | Ser | Leu | Glu | Ala |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |
| Glu | Asp | Ala | Ala | Thr | Tyr | Tyr | Cys | Gln | Gln | Ser | Asn | Thr | Trp | Pro | Phe |
|     |     |     |     | 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Thr | Phe | Gly | Gly | Gly | Thr | Lys | Val | Glu | Ile | Lys |     |     |     |     |     |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     |     |     |     |

<210> 43  
 <211> 107  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Humanized 2B6 light chain variable region - Hu2B6VL-3

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Ile | Val | Leu | Thr | Gln | Ser | Pro | Asp | Phe | Gln | Ser | Val | Thr | Pro | Lys |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Glu | Lys | Val | Thr | Ile | Thr | Cys | Arg | Thr | Ser | Gln | Ser | Ile | Gly | Thr | Asn |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Ile | His | Trp | Tyr | Gln | Gln | Lys | Pro | Asp | Gln | Ser | Pro | Lys | Leu | Leu | Ile |
|     |     | 35  |     |     |     | 40  |     |     |     |     |     | 45  |     |     |     |
| Lys | Tyr | Ala | Ser | Glu | Ser | Ile | Ser | Gly | Val | Pro | Ser | Arg | Phe | Ser | Gly |
|     |     | 50  |     |     |     | 55  |     |     |     | 60  |     |     |     |     |     |
| Ser | Gly | Ser | Gly | Thr | Asp | Phe | Thr | Leu | Thr | Ile | Asn | Ser | Leu | Glu | Ala |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |
| Glu | Asp | Ala | Ala | Thr | Tyr | Tyr | Cys | Gln | Gln | Ser | Asn | Thr | Trp | Pro | Phe |
|     |     |     |     | 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Thr | Phe | Gly | Gly | Gly | Thr | Lys | Val | Glu | Ile | Lys |     |     |     |     |     |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     |     |     |     |

<210> 44  
 <211> 107  
 <212> PRT  
 <213> mus sp.

<220>  
 <223> mouse 3H7 Light Chain Variable Region

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asp | Ile | Gln | Met | Thr | Gln | Ser | Pro | Ser | Ser | Leu | Ser | Ala | Ser | Leu | Gly |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Glu | Arg | Val | Ser | Leu | Thr | Cys | Arg | Ala | Ser | Gln | Glu | Ile | Ser | Gly | Tyr |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Leu | Ser | Trp | Leu | Gln | Gln | Lys | Pro | Asp | Gly | Thr | Ile | Arg | Arg | Leu | Ile |
|     |     | 35  |     |     |     | 40  |     |     |     |     |     | 45  |     |     |     |
| Tyr | Ala | Ala | Ser | Thr | Leu | Asp | Ser | Gly | Val | Pro | Lys | Arg | Phe | Ser | Gly |
|     |     | 50  |     |     |     | 55  |     |     |     | 60  |     |     |     |     |     |
| Ser | Trp | Ser | Gly | Ser | Asp | Tyr | Ser | Leu | Thr | Ile | Ser | Ser | Leu | Glu | Ser |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |
| Glu | Asp | Phe | Ala | Asp | Tyr | Tyr | Cys | Leu | Gln | Tyr | Val | Ser | Tyr | Pro | Tyr |
|     |     |     |     | 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Thr | Phe | Gly | Gly | Gly | Thr | Lys | Leu | Glu | Ile | Lys |     |     |     |     |     |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     |     |     |     |

<210> 45  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 45  
 Lys Lys Phe Ser Arg Ser Asp Pro Asn  
 1 5

<210> 46  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 46  
 Gln Lys Phe Ser Arg Leu Asp Pro Asn  
 1 5

<210> 47  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 47  
 Gln Lys Phe Ser Arg Leu Asp Pro Thr  
 1 5

<210> 48  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 48  
 Lys Lys Phe Ser Arg Leu Asp Pro Thr  
 1 5

<210> 49  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 49  
Gln Lys Phe Ser His Leu Asp Pro Thr  
1 5

<210> 50  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 50  
Lys Lys Phe Ser His Leu Asp Pro Thr  
1 5

<210> 51  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 51  
Gln Lys Phe Ser Arg Leu Asp Pro Asn  
1 5

<210> 52  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 52  
Lys Lys Phe Ser Arg Ser Asp Pro Asn  
1 5

<210> 53  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 53  
Gln Lys Phe Ser Arg Leu Asp Pro Thr  
1 5

<210> 54  
<211> 9  
<212> PRT

<213> Artificial Sequence

<220>

<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 54

Gln Lys Phe Ser His Leu Asp Pro Thr  
1 5

<210> 55

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 55

Lys Lys Phe Ser Arg Leu Asp Pro Thr  
1 5

<210> 56

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> IgG binding site of the Fc-gamma-RIIB receptor

<400> 56

Lys Lys Phe Ser His Leu Asp Pro Thr  
1 5

<210> 57

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Lgh321F primer

<400> 57

cgagctagct ctagatgaga tcacagttct ctctac

36

<210> 58

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Lgh318R primer

<400> 58

gcctccgcct ccg gatccgc ctcttttgat ctccaccttg gtccttc

47

<210> 59

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Lgh319F primer

<400> 59  
ggaggcggat ccggaggcgg aggccagggt cagctggtgc agtctgg 47

<210> 60  
<211> 43  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Lgh320R primer

<400> 60  
tttgaattct agcagcctcc cagtgaggag acggtgaccg tgg 43

<210> 61  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Lgh315R primer

<400> 61  
gcctccgcct ccgatccgc ctcccttgat ctcaagcttg gtcccc 46

<210> 62  
<211> 48  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Lgh316F primer

<400> 62  
ggaggcggat ccggaggcgg aggccagggt accctgagag agtctggc 48

<210> 63  
<211> 47  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Lgh317R primer

<400> 63  
tttgaattcc tagcagcctc ccagtgagct cacagtgacc agagtcc 47

<210> 64  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Lgh339R primer

<400> 64

ctcaacgcag cctcccagtg agctcac 27

<210> 65  
 <211> 31  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Lgh340R primer

<400> 65  
 aacgcagcct cccagtgagg agacggtgac c 31

<210> 66  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Lgh366R primer

<400> 66  
 tttgaattct atttaccgag agacagg 27

<210> 67  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Lgh367F primer

<400> 67  
 ctgggaggct gcgcagagcc caaatcttgt gac 33

<210> 68  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Lgh368R primer

<400> 68  
 gtcacaagat ttgggctctg cgcagcctcc cag 33

<210> 69  
 <211> 50  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Lgh369R primer

<400> 69  
 tttgaattct aacactctcc cctgttgaag cagcctccca gtgaggagac 50

<210> 70  
 <211> 118  
 <212> PRT

<213> Artificial Sequence

<220>

<223> humanized anti-CD16A heavy chain variable region(Hu3G8VH-1)

<400> 70

```

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
 1           5           10           15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
      20           25           30
Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
      35           40           45
Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala
      50           55           60
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
      65           70           75           80
Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
      85           90           95
Cys Ala Arg Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr
      100          105          110
Leu Val Thr Val Ser Ser
      115

```

<210> 71

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> humanized anti-CD16A heavy chain variable region(Hu3G8VH-5)

<400> 71

```

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
 1           5           10           15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
      20           25           30
Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
      35           40           45
Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala
      50           55           60
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
      65           70           75           80
Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
      85           90           95
Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr
      100          105          110
Leu Val Thr Val Ser Ser
      115

```

<210> 72

<211> 117

<212> PRT

<213> Artificial Sequence

<220>

<223> humanized anti-CD16A heavy chain variable region(Hu3G8VH-22)

<400> 72

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln

|     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|
| 1   |     | 5   |     | 10  |     | 15  |
| Thr | Leu | Thr | Leu | Thr | Cys | Thr |
|     |     | 20  |     | 25  |     | 30  |
| Gly | Val | Gly | Val | Gly | Trp | Ile |
|     |     | 35  |     | 40  |     | 45  |
| Trp | Leu | Ala | His | Ile | Trp | Trp |
|     |     | 50  |     | 55  |     | 60  |
| Leu | Lys | Ser | Arg | Leu | Thr | Ile |
|     |     | 65  |     | 70  |     | 75  |
| Val | Leu | Thr | Met | Thr | Asn | Met |
|     |     |     | 85  |     | 90  |     |
| Cys | Ala | Arg | Ile | Asn | Pro | Ala |
|     |     | 100 |     | 105 |     | 110 |
| Leu | Val | Thr | Val | Ser |     |     |
|     |     | 115 |     |     |     |     |

<210> 73  
 <211> 111  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> humanized anti-CD16A light chain variable region(Hu3G8VL-1)

<400> 73

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asp | Ile | Val | Met | Thr | Gln | Ser | Pro | Asp | Ser | Leu | Ala | Val | Ser | Leu | Gly |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     | 15  |     |     |
| Glu | Arg | Ala | Thr | Ile | Asn | Cys | Lys | Ala | Ser | Gln | Ser | Val | Asp | Phe | Asp |
|     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |     |
| Gly | Asp | Ser | Phe | Met | Asn | Trp | Tyr | Gln | Gln | Lys | Pro | Gly | Gln | Pro | Pro |
|     |     | 35  |     |     |     | 40  |     |     |     |     |     | 45  |     |     |     |
| Lys | Leu | Leu | Ile | Tyr | Thr | Thr | Ser | Asn | Leu | Glu | Ser | Gly | Val | Pro | Asp |
|     |     | 50  |     |     |     | 55  |     |     |     | 60  |     |     |     |     |     |
| Arg | Phe | Ser | Gly | Ser | Gly | Ser | Gly | Thr | Asp | Phe | Thr | Leu | Thr | Ile | Ser |
|     |     | 65  |     |     | 70  |     |     | 75  |     |     |     |     |     | 80  |     |
| Ser | Leu | Gln | Ala | Glu | Asp | Val | Ala | Val | Tyr | Tyr | Cys | Gln | Gln | Ser | Asn |
|     |     |     | 85  |     |     |     |     | 90  |     |     |     |     |     | 95  |     |
| Glu | Asp | Pro | Tyr | Thr | Phe | Gly | Gln | Gly | Thr | Lys | Leu | Glu | Ile | Lys |     |
|     |     | 100 |     |     |     |     |     | 105 |     |     |     |     | 110 |     |     |

<210> 74  
 <211> 111  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> humanized anti-CD16A light chain variable region(Hu3G8VL-43)

<400> 74

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Asp | Ile | Val | Met | Thr | Gln | Ser | Pro | Asp | Ser | Leu | Ala | Val | Ser | Leu | Gly |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     | 15  |     |     |
| Glu | Arg | Ala | Thr | Ile | Asn | Cys | Lys | Ser | Ser | Gln | Ser | Val | Asp | Phe | Asp |
|     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |     |
| Gly | Asp | Ser | Phe | Met | Asn | Trp | Tyr | Gln | Gln | Lys | Pro | Gly | Gln | Pro | Pro |
|     |     | 35  |     |     |     | 40  |     |     |     |     |     | 45  |     |     |     |
| Lys | Leu | Leu | Ile | Tyr | Thr | Thr | Ser | Ser | Leu | Glu | Ser | Gly | Val | Pro | Asp |
|     |     | 50  |     |     |     | 55  |     |     |     | 60  |     |     |     |     |     |

```

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65              70              75              80
Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
              85              90              95
Ser Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
              100              105              110

```

<210> 75  
 <211> 111  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> humanized anti-CD16A light chain variable region(Hu3G8VL-22)

```

<400> 75
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1              5              10              15
Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Asp Phe Asp
              20              25              30
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
              35              40              45
Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Thr Gly Val Pro Asp
              50              55              60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65              70              75              80
Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
              85              90              95
Ser Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
              100              105              110

```

<210> 76  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> DNA sequence of linker (SEQ ID NO. 10)

<400> 76  
 ggaggcggat ccggaggcgg aggc 24

<210> 77  
 <211> 18  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <223> DNA sequence of C-terminus of human IgG Kappa chain

<400> 77  
 ttcaacaggg gagagtgt 18

<210> 78  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Linker plus C-terminus of human IgG Kappa chain

<400> 78

ctgggaggct gcttcaacag gggagagtgt

30

<210> 79

<211> 6

<212> PRT

<213> Homo sapiens

<220>

<223> Human IgG hinge domain

<400> 79

Val Glu Pro Lys Ser Cys

1

5

<210> 80

<211> 18

<212> DNA

<213> Homo sapiens

<220>

<223> Human IgG hinge domain

<400> 80

gttgagccca aatcttgt

18

<210> 81

<211> 118

<212> PRT

<213> Homo sapiens

<220>

<223> VH domain of 3G8 antibody

<400> 81

Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln  
1 5 10 15

Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Arg Thr Ser  
20 25 30

Gly Met Gly Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu  
35 40 45

Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala  
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn Gln Val  
65 70 75 80

Phe Leu Lys Ile Ala Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr  
85 90 95

Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr  
100 105 110

Leu Val Thr Val Ser Ser  
115

<210> 82

<211> 354

<212> DNA

<213> Homo sapiens

<220>

<223> VH domain of 3G8 antibody

<400> 82

```
caggttaccc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
acttggttctt tctctgggtt ttcactgagg acttctggta tgggtgtagg ctggattcgt 120
cagccttcag ggaagggctc agagtggctg gcacacattt ggtgggatga tgacaagcgc 180
tataatccag ccctgaagag ccgactgaca atctccaagg atacctccag caaccaggta 240
ttcctcaaaa tcgccagtgt ggacactgca gatactgcca catactactg tgctcaaata 300
aaccgccgct ggtttgctta ctggggccaa gggactctgg tcactgtgag ctca 354
```

<210> 83

<211> 354

<212> DNA

<213> Artificial Sequence

<220>

<223> humanized anti-CD16A heavy chain variable region(Hu3G8VH-5)

<400> 83

```
caggttaccc tgagagagtc tggccctgcg ctggtgaagc ccacacagac cctcacactg 60
acttgtagctt tctctgggtt ttcactgagc acttctggta tgggtgtagg ctggattcgt 120
cagcctcccg ggaaggtctc agagtggctg gcacacattt ggtgggatga tgacaagcgc 180
tataatccag ccctgaagag ccgactgaca atctccaagg atacctccaa aaaccaggta 240
gtcctcaciaa tgaccaacat ggaccctgtg gatactgcca catactactg tgctcaaata 300
aaccgccgct ggtttgctta ctggggccaa gggactctgg tcactgtgag ctca 354
```

<210> 84

<211> 111

<212> PRT

<213> Homo sapiens

<220>

<223> VL domain of 3G8 antibody

<400> 84

```
Asp Thr Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1          5          10          15
Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser Val Asp Phe Asp
      20          25          30
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
      35          40          45
Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Ile Pro Ala
      50          55          60
Arg Phe Ser Ala Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
65          70          75          80
Pro Val Glu Glu Glu Asp Thr Ala Thr Tyr Tyr Cys Gln Gln Ser Asn
      85          90          95
Glu Asp Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100          105          110
```

<210> 85

<211> 333

<212> DNA

<213> Homo sapiens

<220>

<223> VL domain of 3G8 antibody

<400> 85  
gacactgtgc tgaccaatc tccagcttct ttggctgtgt ctctagggca gagggccacc 60  
atctcctgca aggccagcca aagtgttgat tttgatggtg atagttttat gaactggtac 120  
caacagaaac caggacagcc acccaaaactc ctcacttata ctacatccaa tctagaatct 180  
gggatcccag ccagggttttag tgccagtggg tctgggacag acttcaccct caacatccat 240  
cctgtggagg aggaggatac tgcaacctat tactgtcagc aaagtaatga ggatccgtac 300  
acgttcggag gggggaccaa gctggaaata aaa 333

<210> 86  
<211> 333  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> humanized anti-CD16A light chain variable region(Hu3G8VL-1)

<400> 86  
gacatcgtga tgaccaatc tccagaactct ttggctgtgt ctctagggga gagggccacc 60  
atcaactgca aggccagcca aagtgttgat tttgatggtg atagttttat gaactggtac 120  
caacagaaac caggacagcc acccaaaactc ctcacttata ctacatccaa tctagaatct 180  
ggggtcccag acagggttttag tgccagtggg tctgggacag acttcaccct caccatcagc 240  
agcctgcagg ctgaggatgt ggcagtttat tactgtcagc aaagtaatga ggatccgtac 300  
acgttcggac aggggaccaa gcttgagatc aaa 333

<210> 87  
<211> 121  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> humanized 2B6 heavy chain variable region

<400> 87

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |    |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|
| Gln | Val | Gln | Leu | Val | Gln | Ser | Gly | Ala | Glu | Val | Lys | Lys | Pro | Gly | Ala | 1   | 5   | 10  | 15 |
| Ser | Val | Lys | Val | Ser | Cys | Lys | Ala | Ser | Gly | Tyr | Thr | Phe | Thr | Asn | Tyr | 20  | 25  | 30  |    |
| Trp | Ile | His | Trp | Val | Arg | Gln | Ala | Pro | Gly | Gln | Gly | Leu | Glu | Trp | Ile | 35  | 40  | 45  |    |
| Gly | Val | Ile | Asp | Pro | Ser | Asp | Thr | Tyr | Pro | Asn | Tyr | Asn | Lys | Lys | Phe | 50  | 55  | 60  |    |
| Lys | Gly | Arg | Val | Thr | Met | Thr | Val | Val | Val | Ser | Thr | Ser | Thr | Ala | Tyr | 65  | 70  | 75  | 80 |
| Met | Glu | Leu | Arg | Ser | Leu | Arg | Ser | Asp | Asp | Thr | Ala | Val | Tyr | Tyr | Cys | 85  | 90  | 95  |    |
| Ala | Arg | Asn | Gly | Asp | Ser | Asp | Tyr | Tyr | Ser | Gly | Met | Asp | Tyr | Trp | Gly | 100 | 105 | 110 |    |
| Gln | Gly | Thr | Thr | Val | Thr | Val | Ser | Ser | 115 | 120 |     |     |     |     |     |     |     |     |    |

<210> 88  
<211> 363  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> humanized 2B6 heavy chain variable region

<400> 88

```

caggttcagc tgggtgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60
tcttgcaagg cttctgggta cacctttacc aactactgga tacactgggt gcgacaggcc 120
cctggacaag ggcttgagtg gattggagtg attgatcctt ctgatactta tccaaattac 180
aataaaaaag tcaagggcag agtcaccatg accgtagtcg tatccacgag cacagcctac 240
atggagctga ggagcctgag atctgacgac acggcctgtg attactgtgc gagaaacggg 300
gattccgatt attactctgg tatggactac tgggggcaag ggaccacggg caccgtctcc 360
tca

```

<210> 89

<211> 111

<212> PRT

<213> Artificial Sequence

<220>

<223> humanized 2B6 light chain variable region

<400> 89

```

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1             5             10             15
Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp
          20             25             30
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
          35             40             45
Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro Asp
          50             55             60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65             70             75             80
Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
          85             90             95
Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
          100             105             110

```

<210> 90

<211> 321

<212> DNA

<213> Artificial Sequence

<220>

<223> humanized 2B6 light chain variable region

<400> 90

```

gaaattgtgc tgactcagtc tccagacttt cagtctgtga ctccaaagga gaaagtcacc 60
ttcacctgca ggaccagtca gagcattggc acaaacatac actggtacca gcagaaacca 120
gatcagtcctc caaagctcct catcaaggag gtttctgagt ctatctctgg agtccatcgc 180
aggttcagtg gcagtggtac tgggacagat ttcacccctca ccatcaatag cctggaagct 240
gaagatgctg caacgtatta ctgtcaacaa agtaatacct ggccgttcac gttcggcgga 300
gggaccaagg tggagatcaa a

```

<210> 91

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> thrombin cleavage recognition site

<400> 91

```

Leu Val Pro Arg Gly Ser
 1             5

```

<210> 92  
 <211> 4  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Factor Xa cleavage recognition site

<220>  
 <221> VARIANT  
 <222> 2  
 <223> Xaa = Glu or Asp

<400> 92  
 Ile Xaa Gly Arg  
 1

<210> 93  
 <211> 5  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> enterokinase cleavage recognition site

<400> 93  
 Asp Asp Asp Asp Lys  
 1 5

<210> 94  
 <211> 4  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> furin cleavage recognition site

<220>  
 <221> VARIANT  
 <222> 2, 3  
 <223> Xaa = Any Amino Acid

<400> 94  
 Arg Xaa Xaa Arg  
 1

<210> 95  
 <211> 4  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> preferred furin cleavage recognition site

<220>  
 <221> VARIANT

<222> 2  
<223> Xaa = Any Amino Acid

<220>  
<221> VARIANT  
<222> 3  
<223> Xaa = Lys or Arg

<400> 95  
Arg Xaa Xaa Arg  
1

<210> 96  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> AcTEV cleavage recognition site

<400> 96  
Glu Asn Leu Tyr Phe Gln Gly  
1 5

<210> 97  
<211> 487  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> covalent diabody polypeptide precursor (Construct 13)

<400> 97  
Asp Thr Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly  
1 5 10 15  
Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser Val Asp Phe Asp  
20 25 30  
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro  
35 40 45  
Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Ile Pro Ala  
50 55 60  
Arg Phe Ser Ala Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His  
65 70 75 80  
Pro Val Glu Glu Glu Asp Thr Ala Thr Tyr Tyr Cys Gln Gln Ser Asn  
85 90 95  
Glu Asp Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly  
100 105 110  
Gly Gly Ser Gly Gly Gly Gly Glu Val Glu Leu Val Glu Ser Gly Gly  
115 120 125  
Gly Leu Val Gln Pro Gly Arg Ser Leu Lys Leu Ser Cys Ala Ala Ser  
130 135 140  
Gly Phe Thr Phe Ser Asp Tyr Tyr Met Ala Trp Val Arg Gln Ala Pro  
145 150 155 160  
Thr Thr Gly Leu Glu Trp Val Ala Ser Ile Ser Tyr Asp Gly Gly Asp  
165 170 175  
Thr His Tyr Arg Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp  
180 185 190  
Asn Ala Lys Ser Ser Leu Tyr Leu Gln Met Asp Ser Leu Arg Ser Glu  
195 200 205

```

Asp Thr Ala Thr Tyr Tyr Cys Ala Thr Glu Thr Thr Gly Ile Pro Thr
  210                215                220
Gly Val Met Asp Ala Trp Gly Gln Gly Val Ser Val Thr Val Ser Ser
  225                230                235                240
Leu Gly Gly Cys Gly Arg Ala Lys Arg Asp Val Gln Met Thr Gln
                245                250                255
Ser Pro Ser Asn Leu Ala Ala Ser Pro Gly Glu Ser Val Ser Ile Asn
                260                265                270
Cys Lys Ala Ser Glu Ser Ile Ser Lys Tyr Leu Ala Trp Tyr Leu Gln
                275                280                285
Lys Pro Gly Lys Ala Asn Lys Leu Leu Met Tyr Asp Gly Ser Thr Leu
                290                295                300
Gln Ser Gly Ile Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp
  305                310                315                320
Phe Thr Leu Thr Ile Arg Ser Leu Glu Pro Glu Asp Phe Gly Leu Tyr
                325                330                335
Tyr Cys Gln Gln His Tyr Glu Tyr Pro Ala Thr Phe Gly Ser Gly Thr
                340                345                350
Lys Leu Glu Ile Lys Gly Gly Gly Ser Gly Gly Gly Gly Gln Val Thr
                355                360                365
Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln Thr Leu Ser
                370                375                380
Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Arg Thr Ser Gly Met Gly
  385                390                395                400
Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu Trp Leu Ala
                405                410                415
His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala Leu Lys Ser
                420                425                430
Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn Gln Val Phe Leu Lys
                435                440                445
Ile Ala Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr Cys Ala Gln
                450                455                460
Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
  465                470                475                480
Val Ser Ser Leu Gly Gly Cys
                485

```

<210> 98  
 <211> 1464  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> covalent diabody polypeptide precursor (DNA sequence of  
 SEQ ID NO. 97)

```

<400> 98
gacactgtgc tgaccaatc tccagcttct ttggctgtgt ctctagggca gagggccacc 60
atctcctgca aggccagcca aagtgtgat ttgatggtg atagttttat gaactggtac 120
caacagaaac caggacagcc acccaaaactc ctcatctata ctacatccaa tctagaatct 180
gggatcccag ccaggtttag tgccagtggg tctgggacag acttcaccct caacatccat 240
cctgtggagg aggaggatac tgcaacctat tactgtcagc aaagtaatga ggatccgtac 300
acgttcggag gggggaccaa gctggaaata aaaggaggcg gatccggagg cggaggcgag 360
gtggagctag tggagtcttg gggaggctta gtgcagcctg gaaggtccct gaaactctcg 420
tgtgcagcct caggattcac tttagtgac tattacatgg cctgggtccg gcagggtcca 480
acgacgggtc tggagtgggt cgcattccatt agttatgatg gtggtgacac tcaactatcg 540
gactccgtga agggccgatt tactatttcc agagataatg caaaaagcag cctatacctg 600
caaatggaca gtctgaggtc tgaggacacg gccacttatt actgtgcaac agagactacg 660
ggaataccta caggtgttat ggatgcctgg ggtcaaggag tttcagtcac tgtctcctca 720
ctgggagggt gcggcgggag agctaagagg gatgtccaga tgaccagtc tccatcta 780

```

```

cttgcctgcct ctccctggaga aagtgtttcc atcaattgca aggcaagtga gagcattagc 840
aagtatttag cctgggtatct acagaaacct gggaaagcaa ataagcttct tatgtacgat 900
gggtcaactt tgcaatctgg aattccatcg aggttcagtg gcagtggatc tggtagacat 960
ttcactctca ccatacagaag cctggagcct gaagattttg gactctatta ctgtcaacag 1020
cattatgaat atccagccac gtccggttct gggaccaagc tggagatcaa aggaggcgga 1080
tccggaggcg gaggccaggt taccctgaaa gagtctggcc ctgggatatt gcagccctcc 1140
cagaccctca gtctgacttg ttctttctct gggttttcac tgaggacttc tggtaggggt 1200
gtaggctgga ttctgcagcc ttcagggaag ggtctagagt ggctggcaca catttggtgg 1260
gatgatgaca agcgctataa tccagccctg aagagccgac tgacaatctc caaggatacc 1320
tccagcaacc aggtattcct caaaatcgcc agtgtggaca ctgcagatac tgccacatac 1380
tactgtgctc aaataaaccc cgctgtggtt gcttactggg gccaaaggac tctggtcact 1440
gtgagctcac tgggaggctg ctac                                     1464

```

<210> 99

<211> 511

<212> PRT

<213> Artificial Sequence

<220>

<223> covalent diabody polyprotein precursor (Construct 14)

<400> 99

```

Asp Thr Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1           5           10           15
Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser Val Asp Phe Asp
          20           25           30
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
          35           40           45
Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Ile Pro Ala
          50           55           60
Arg Phe Ser Ala Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
          65           70           75           80
Pro Val Glu Glu Glu Asp Thr Ala Thr Tyr Tyr Cys Gln Gln Ser Asn
          85           90           95
Glu Asp Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly
          100          105          110
Gly Gly Ser Gly Gly Gly Gly Glu Val Glu Leu Val Glu Ser Gly Gly
          115          120          125
Gly Leu Val Gln Pro Gly Arg Ser Leu Lys Leu Ser Cys Ala Ala Ser
          130          135          140
Gly Phe Thr Phe Ser Asp Tyr Tyr Met Ala Trp Val Arg Gln Ala Pro
          145          150          155          160
Thr Thr Gly Leu Glu Trp Val Ala Ser Ile Ser Tyr Asp Gly Gly Asp
          165          170          175
Thr His Tyr Arg Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
          180          185          190
Asn Ala Lys Ser Ser Leu Tyr Leu Gln Met Asp Ser Leu Arg Ser Glu
          195          200          205
Asp Thr Ala Thr Tyr Tyr Cys Ala Thr Glu Thr Thr Gly Ile Pro Thr
          210          215          220
Gly Val Met Asp Ala Trp Gly Gln Gly Val Ser Val Thr Val Ser Ser
          225          230          235          240
Leu Gly Gly Cys Gly Gly Arg Ala Lys Arg Ala Pro Val Lys Gln Thr
          245          250          255
Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro
          260          265          270
Gly Pro Asp Val Gln Met Thr Gln Ser Pro Ser Asn Leu Ala Ala Ser
          275          280          285
Pro Gly Glu Ser Val Ser Ile Asn Cys Lys Ala Ser Glu Ser Ile Ser
          290          295          300
Lys Tyr Leu Ala Trp Tyr Leu Gln Lys Pro Gly Lys Ala Asn Lys Leu

```

```

305          310          315          320
Leu Met Tyr Asp Gly Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg Phe
          325          330          335
Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Arg Ser Leu
          340          345          350
Glu Pro Glu Asp Phe Gly Leu Tyr Tyr Cys Gln Gln His Tyr Glu Tyr
          355          360          365
Pro Ala Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly
          370          375          380
Ser Gly Gly Gly Gly Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile
385          390          395          400
Leu Gln Pro Ser Gln Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe
          405          410          415
Ser Leu Arg Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln Pro Ser
          420          425          430
Gly Lys Gly Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys
          435          440          445
Arg Tyr Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr
450          455          460
Ser Ser Asn Gln Val Phe Leu Lys Ile Ala Ser Val Asp Thr Ala Asp
465          470          475          480
Thr Ala Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr
          485          490          495
Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Leu Gly Gly Cys
          500          505          510

```

<210> 100

<211> 1536

<212> DNA

<213> Artificial Sequence

<220>

<223> covalent diabody polypeptide precursor (DNA sequence of  
SEQ ID NO. 99)

<400> 100

```

gacactgtgc tgacceaatc tccagcttct ttggctgtgt ctctagggca gagggccacc 60
atctcctgca aggccagcca aagtgttgat ttgatgggtg atagttttat gaactggtag 120
caacagaaac caggacagcc acccaaaactc ctcatctata ctacatcaa tctagaatct 180
gggatcccag ccaggttttag tgccagtggg tctgggacag acttcaccct caacatccat 240
cctgtggagg aggaggatac tgcaacctat tactgtcagc aaagtaatga ggatccgtac 300
acgttcggag gggggaccaa gctggaaata aaaggaggcg gatccggagg cggaggcgag 360
gtggagctag tggagtctgg gggaggctta gtgcagcctg gaaggtccct gaaactctcg 420
tgtgcagcct caggattcac ttccagtgc tttacatgg cctgggtccg gcaggtccca 480
acgacgggtc tggagtgggt cgcattccatt agttatgatg gtggtgacac tcactatcga 540
gactccgtga agggccgatt tactatttcc agagataatg caaaaagcag cctataacctg 600
caaatggaca gtctgaggtc tgaggacacg gccacttatt actgtgcaac agagactacg 660
ggaataccta caggtgttat ggatgcctgg ggtcaaggag ttccagtgc tctctcctca 720
ctgggaggct gcggcgggag agctaagagg gcccctgtga agcagaccct gaacttcgac 780
ctgctgaagc tggccggaga cgtggagagc aaccccgccc ccgatgtcca gatgaccag 840
tctccatcta atcttctgctc ctctcctgga gaaagtgtt ccatcaattg caaggcaagt 900
gagagcatta gcaagtattt agcctggtat ctacagaaac ctgggaaagc aaataagctt 960
cttatgtacg atgggtcaac ttgtcaatct ggaattccat cgagggtcag tggcagtgga 1020
tctggtacag atttcaactc caccatcaga agcctggagc ctgaagattt tggactctat 1080
tactgtcaac agcattatga atatccagcc acgttcggtt ctgggaccaa gctggagatc 1140
aaaggaggcg gatccggagg cggaggccag gttaccctga aagagctgg ccctgggata 1200
ttgcagccct cccagaccct cagctgactg tgttctttct ctgggttttc actgaggact 1260
tctggtatgg gtgtaggctg gattcgtcag ccttcaggga agggctctaga gtggctggca 1320
cacatttggg gggatgatga caagcgtat aatccagccc tgaagagccg actgacaatc 1380
tccaaggata cctccagcaa ccaggtattc ctcaaaatcg ccagtgtgga cactgcagat 1440

```

actgccacat actactgtgc tcaaataaac cccgcctggt ttgcttactg gggccaaggg 1500  
actctgggtca ctgtgagctc actgggagggc tgctag 1536

<210> 101  
<211> 239  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Diabody construct (Construct 15)

<400> 101  
Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys  
1 5 10 15  
Glu Lys Val Thr Phe Thr Cys Arg Thr Ser Gln Ser Ile Gly Thr Asn  
20 25 30  
Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile  
35 40 45  
Lys Glu Val Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly  
50 55 60  
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala  
65 70 75 80  
Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Thr Trp Pro Phe  
85 90 95  
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Gly Gly Gly Ser Gly  
100 105 110  
Gly Gly Gly Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys  
115 120 125  
Pro Thr Gln Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu  
130 135 140  
Ser Thr Ser Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys  
145 150 155 160  
Ala Leu Glu Trp Leu Ala His Ile Trp Trp Asp Asp Lys Arg Tyr  
165 170 175  
Asn Pro Ala Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys  
180 185 190  
Asn Gln Val Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala  
195 200 205  
Thr Tyr Tyr Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly  
210 215 220  
Gln Gly Thr Leu Val Thr Val Ser Ser Phe Asn Arg Gly Glu Cys  
225 230 235

<210> 102  
<211> 720  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Diabody construct (DNA sequence of SEQ ID NO. 101)

<400> 102  
gaaattgtgc tgactcagtc tccagacttt cagtctgtga ctccaaagga gaaagtcacc 60  
ttcacctgca ggaccagtca gagcattggc acaaacatac actggtacca gcagaaacca 120  
gatcagtcct caaagctcct catcaaggag gtttctgagt ctatctctgg agtcccatcg 180  
aggttcagtg gcagtggatc tgggacagat ttcaccctca ccatcaatag cctggaagct 240  
gaagatgctg caacgtatta ctgtcaacaa agtaatacct ggccgttcac gttcggcgga 300  
gggaccaagg tggagatcaa aggaggcgga tccggaggcg gaggccaggt taccctgaga 360  
gagtctggcc ctgcgctggt gaagcccaca cagaccctca cactgacttg taccttctct 420  
gggttttcac tgagcacttc tggtatgggt gtaggctgga ttcgtcagcc tcccgggaag 480

gctctagagt ggctggcaca catttggtgg gatgatgaca agcgctataa tccagccctg 540  
 aagagccgac tgacaatctc caaggatacc tccaaaaacc aggtagtcct cacaatgacc 600  
 aacatggacc ctgtggatac tgccacatac tactgtgctc aaataaacc cgctgggtt 660  
 gcttactggg gccaaaggac tctggtcact gtgagctcat tcaacagggg agagtgttag 720

<210> 103  
 <211> 246  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Diabody construct (Construct 16)

<400> 103  
 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly

|                 |                 |                 |                 |
|-----------------|-----------------|-----------------|-----------------|
| 1               | 5               | 10              | 15              |
| Glu Arg Ala Thr | Ile Asn Cys Lys | Ala Ser Gln Ser | Val Asp Phe Asp |
| 20              | 25              | 30              |                 |
| Gly Asp Ser Phe | Met Asn Trp Tyr | Gln Gln Lys Pro | Gly Gln Pro Pro |
| 35              | 40              | 45              |                 |
| Lys Leu Leu Ile | Tyr Thr Thr Ser | Asn Leu Glu Ser | Gly Val Pro Asp |
| 50              | 55              | 60              |                 |
| Arg Phe Ser Gly | Ser Gly Thr Asp | Phe Thr Leu Thr | Ile Ser         |
| 65              | 70              | 75              | 80              |
| Ser Leu Gln Ala | Glu Asp Val Ala | Val Tyr Tyr Cys | Gln Gln Ser Asn |
| 85              | 90              | 95              |                 |
| Glu Asp Pro Tyr | Thr Phe Gly Gln | Gly Thr Lys Leu | Glu Ile Lys Gly |
| 100             | 105             | 110             |                 |
| Gly Gly Ser Gly | Gly Gly Gly Gln | Val Gln Leu Val | Gln Ser Gly Ala |
| 115             | 120             | 125             |                 |
| Glu Val Lys Lys | Pro Gly Ala Ser | Val Lys Val Ser | Cys Lys Ala Ser |
| 130             | 135             | 140             |                 |
| Gly Tyr Thr Phe | Thr Asn Tyr Trp | Ile His Trp Val | Arg Gln Ala Pro |
| 145             | 150             | 155             | 160             |
| Gly Gln Gly Leu | Glu Trp Ile Gly | Val Ile Asp Pro | Ser Asp Thr Tyr |
| 165             | 170             | 175             |                 |
| Pro Asn Tyr Asn | Lys Lys Phe Lys | Gly Arg Val Thr | Met Thr Val Val |
| 180             | 185             | 190             |                 |
| Val Ser Thr Ser | Thr Ala Tyr Met | Glu Leu Arg Ser | Leu Arg Ser Asp |
| 195             | 200             | 205             |                 |
| Asp Thr Ala Val | Tyr Tyr Cys Ala | Arg Asn Gly Asp | Ser Asp Tyr Tyr |
| 210             | 215             | 220             |                 |
| Ser Gly Met Asp | Tyr Trp Gly Gln | Gly Thr Thr Val | Thr Val Ser Ser |
| 225             | 230             | 235             | 240             |
| Val Glu Pro Lys | Ser Cys         |                 |                 |
| 245             |                 |                 |                 |

<210> 104  
 <211> 741  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Diabody construct (DNA sequence of SEQ ID NO. 103)

<400> 104  
 gacatcgtga tgacccaatc tccagactct ttggctgtgt ctctagggga gagggccacc 60  
 atcaactgca aggccagcca aagtgttgat tttgatgggtg atagttttat gaactggtac 120

|            |             |            |            |            |            |     |
|------------|-------------|------------|------------|------------|------------|-----|
| caacagaaac | caggacagcc  | acccaaactc | ctcatctata | ctacatccaa | tctagaatct | 180 |
| ggggtcccag | acaggttttag | tggcagtggg | tctgggacag | acttcaccct | caccatcagc | 240 |
| agcctgcagg | ctgaggatgt  | ggcagtttat | tactgtcagc | aaagtaatga | ggatccgtac | 300 |
| acgttcggac | aggggaccaa  | gcttgagatc | aaaggaggcg | gatccggagg | cggaggccag | 360 |
| gttcagctgg | tgagctctgg  | agctgaggtg | aagaagcctg | gggcctcagt | gaaggtctcc | 420 |
| tgcaaggctt | ctggttacac  | ctttaccaac | tactggatac | actgggtgcg | acaggcccct | 480 |
| ggacaagggc | ttgagtggat  | tgagtgatt  | gatccttctg | atacttatcc | aaattacaat | 540 |
| aaaaagttca | agggcagagt  | caccatgacc | gtagtcgtat | ccacgagcac | agcctacatg | 600 |
| gagctgagga | gcctgagatc  | tgacgacacg | gccgtgtatt | actgtgcgag | aaacggtgat | 660 |
| tccgattatt | actctggtat  | ggactactgg | gggcaaggga | ccacggtcac | cgtctcctca | 720 |
| gttgagccca | aatcttggtta | g          |            |            |            | 741 |