



(86) **Date de dépôt PCT/PCT Filing Date:** 2008/12/19
(87) **Date publication PCT/PCT Publication Date:** 2009/07/09
(45) **Date de délivrance/Issue Date:** 2019/08/06
(85) **Entrée phase nationale/National Entry:** 2010/06/14
(86) **N° demande PCT/PCT Application No.:** US 2008/013922
(87) **N° publication PCT/PCT Publication No.:** 2009/085221
(30) **Priorité/Priority:** 2007/12/21 (US61/008,680)

(51) **Cl.Int./Int.Cl.** **C12Q 1/689** (2018.01),
C12Q 1/68 (2018.01), **C12Q 1/6813** (2018.01),
C12Q 1/6844 (2018.01), **C07H 21/04** (2006.01)


(72) **Inventeurs/Inventors:**
JAY, CORINNE, FR;
DEIMAN, BIRGIT, NL;
VAN STRIJP, DIANNE, NL;
VAN DE WIEL, PAUL, NL

(73) **Propriétaire/Owner:**
BIOMERIEUX SA, US

(74) **Agent:** SMART & BIGGAR

(54) **Titre : DETECTION DE STAPHYLOCOCCUS AUREUS RESISTANT A LA METHICILLINE**

(54) **Title: DETECTION OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS**



(57) **Abrégé/Abstract:**

The present invention provides improved tests for the detection of methicillin- resistant Staphylococcus aureus. The tests are particularly useful for eliminating false positive results due to the presence of a mixed bacterial population in patient samples.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 July 2009 (09.07.2009)

PCT

(10) International Publication Number
WO 2009/085221 A2

(51) International Patent Classification:
C12Q 1/68 (2006.01)

(74) Agent: MYERS BIGEL SIBLEY & SAJOVEC, P.A.;
P.O. Box 37428, Raleigh, NC 27627 (US).

(21) International Application Number:
PCT/US2008/013922

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
19 December 2008 (19.12.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/008,680 21 December 2007 (21.12.2007) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
BIOMERIEUX SA [US/US]; 100 Rodophe Street,
Durham, NC 27712 (US).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **JAY, Corinne** [FR/FR]; 223 avenue de Grenoble, F-38180 Seyssins (FR). **DEIMAN, Birgit** [NL/NL]; Dille 19, NL-5063 EE Oisterwijk (NL). **VAN STRIJP, Dianne** [NL/NL]; Zoomweg 11, NL-5215 HN 's-Hertogenbosch (NL). **VAN DE WIEL, Paul** [NL/NL]; Aquamarijn 21, NL-5629 HC Eindhoven (NL).

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: DETECTION OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS

Fig. 1



(57) Abstract: The present invention provides improved tests for the detection of methicillin-resistant *Staphylococcus aureus*. The tests are particularly useful for eliminating false positive results due to the presence of a mixed bacterial population in patient samples.

WO 2009/085221 A2

DETECTION OF METHICILLIN-RESISTANT *STAPHYLOCOCCUS AUREUS***FIELD OF THE INVENTION**

[0001] The present invention relates to molecular detection of methicillin-resistant *Staphylococcus aureus* (MRSA). More particularly, the present invention relates to an improved detection of MRSA that reduces false positive results.

5

BACKGROUND OF THE INVENTION

[0002] Methicillin-resistant *Staphylococcus aureus* (MRSA) is a major nosocomial but also community acquired pathogen that can cause serious infections such as surgical wound infections, pneumonia, endocarditis and septicemia. Resistance to methicillin is due to the presence of the *mecA* gene that encodes a modified Penicillin-Binding protein, PBP2a or PBP2', with reduced affinity for B-lactam drugs. The *mecA* gene is carried by a cassette named the SCCmec (Staphylococcal Cassette Chromosome mec; Ito et al., 2001, Antimicrob. Agents Chemother. 45(5):1323-1336; Hiramatsu, et al., 2001, Trends Microbiol. Oct;9(10):486-93), a mobile element that can be incorporated into the chromosome of *S. aureus* and other coagulase negative Staphylococci, mainly *S. epidermidis* and *S. haemolyticus*. SCCmec is characterized by the presence of terminal inverted and direct repeats, a set of site-specific recombinase genes (*ccrA* and *ccrB*), and the *mecA* gene complex (Ito et al., 1999, Antimicrob. Agents Chemother. 43:1449-1458; Katayama et al., 2000, Antimicrob. Agents Chemother. 44:1549-1555). The site of insertion of this *mecA* gene cassette SCCmec into the *Staphylococcus aureus* genome is known and the sequence conserved (Ito et al., 2001, Antimicrob. Agents Chemother. 45:1323-1336). After insertion into the *S. aureus* chromosome, the SCCmec has a left extremity junction region and a right extremity junction region (see FIG. 1), where the SCCmec sequence is contiguous with the *S. aureus* chromosomal sequence. The nucleotide sequence of the regions surrounding the left and right boundaries of SCCmec DNA (i.e. *attL* and *attR*, respectively), as well as those of the regions around the SCCmec DNA integration site (i.e., *attB_{scc}*, the bacterial chromosome attachment site for SCCmec DNA), have previously been analyzed. Sequence analysis of the integration sites revealed that *attB_{scc}* is located at the 3' end of a novel open reading frame (ORF), *orfX*. *orfX* encodes a putative 159-amino acid polypeptide that exhibits sequence homology with some previously identified polypeptides of unknown function (Ito et al.,

30

1999, Antimicrob. Agents Chemother. 43:1449-1458). Organization of the *mecA* region of *SCCmec* has additionally been studied (Oliveira, D.C., et al., 2000, Antimicrob. Agents Chemother. 44(7):1906-1910).

[0003] MRSA can be carried by healthy people without causing any disease but these

5 healthy carriers, when entering the hospital, can contaminate hospitalized patients.

Additionally, a patient can contaminate himself, e.g., if one undergoes surgery, the risk of infection is increased. MRSA healthy carriers constitute a reservoir of MRSA and screening of these carriers must be performed to eradicate the strains by local decontamination. MRSA screening is now recognized as a major tool to reduce the prevalence of MRSA strains in the
10 world. Typically, in an MRSA assay in a patient, a nasal swab is taken from the patient and cultured repeatedly, to determine if an MRSA strain is present. The need to culture could be obviated by an assay for identifying MRSA directly from a nasal swab. Culture identification methods typically require minimally 24 hours, and more typically 72 hours, to obtain results. New chromogenic media (having substrate(s) within the media and, typically, antibiotic (e.g.,
15 cefoxitin) to select methicillin-resistant strains) can potentially restrict this time to result to a 24- 48 hour time period. However, in the case of MRSA infection, results are needed in a matter of hours, since the patient should be isolated until results are obtained. Therefore, a reliable molecular MRSA test which can provide results in a matter of 2-4 hours is highly desirable.

20 **[0004]** Amplification is a well known art, and various methods have been developed, including transcription-based amplification such as transcription-mediated amplification (TMA; U.S. Pat. Nos. 5,766,849 5,399,491; 5,480,784; 5,766,849; and 5,654,142) and nucleic acid sequence-based amplification (NASBA; 5,130,238; 5,409,818; 5,654,142; and 6,312,928), and cycling nucleic acid amplification technologies (thermocycling) such as
25 polymerase chain reaction (PCR; U.S. Pat. Nos. 4,683,195; 4,965,188; 4,683,202) and ligase chain reaction (LCR; U.S. Pat. No. 5,792,607). Known amplification methods also include strand displacement amplification (SDA), self-sustained sequence replication (3SR), Q- β replicase, and cascade rolling circle amplification (CRCA).

[0005] Detection methods utilizing nucleic acids are also well known in the art.

30 Nucleic acids are often labeled for various detection purposes. For example, methods described in U.S. Patent Nos 4,486,539 (Kourlisky); 4,411,955 (Ward); 4,882,269 (Schneider) and 4,213,893 (Carrico), illustrate preparation of labeled detection probes for detecting specific nucleic acid sequences. Probe designs for different detection methods, such as target-capture, HPA, TAQman, molecular beacons and sandwich hybridization have

also been described (e.g., U.S. Pat. No. 4,486,539, and U.S. Pat. No. 4,751,177; 5,210,015; 5,487,972; 5,804,375; 5,994,076). Nucleic acid hybridization techniques and conditions are known to the skilled artisan and have been described for example, in Sambrook et al.

Molecular Cloning A Laboratory Manual, 2nd Ed. Cold Spring Lab. Press, Dec. 1989; U.S.

5 Patent Nos 4,563,419 (Ranki) and 4,851,330 (Kohne) and in Dunn, et al., *Cell* 12, pp. 23-26 (1978) among many other publications. Probe designs for different detection methods are also known, such as target-capture, HPA, TaqMan, molecular beacons and sandwich hybridization (e.g., U.S. Pat. No. 4,486,539, and U.S. Pat. No. 4,751,177; 5,210,015; 5,487,972; 5,804,375; 5,994,076).

10 [0006] Earlier molecular methods developed to detect and identify MRSA based on the detection of the *mecA* gene and *S. aureus*-specific chromosomal sequences have been described. (Saito et al., 1995, J. Clin. Microbiol. 33:2498-2500; Ubukata et al., 1992, J. Clin. Microbiol. 30:1728-1733; Murakami et al., 1991, J. Clin. Microbiol. 29:2240-2244; Hiramatsu et al., 1992, Microbiol. Immunol. 36:445-453). However, positive results for the
15 presence in a sample of both *mecA* gene and *S. aureus* chromosomal sequences cannot guarantee MRSA is present, since, for example, in tests based on the detection of *mecA* and *S. aureus* specific marker, false positives can be observed in the presence of MSSA and methicillin resistant coagulase negative Staphylococcus that possess the *mecA* gene. Furthermore, in tests based on the detection of the cassette junction only, false positives have
20 been observed with methicillin-susceptible *S. aureus* isolates containing a small fragment of the right extremity of the *SCCmec* (see Rupp, J. et al., J. Clin. Microbiol. 44(6): 2317 (2006)). Additionally, Ramakrishnan and Riccelli describe a method for detecting MRSA utilizing oligonucleotide probes having sequences that are complementary to regions near the left junction of the *SCCmec* cassette insertion site, including part of the *SCCmec* cassette
25 sequence and part of the *S. aureus* sequence in the region of insertion (the left extremity junction region) (U.S. patent publication No. US20060057613).

[0007] However, previous attempts to determine MRSA by molecular methods have had difficulties with false positive results. Such results have been postulated to be the result of any of: the presence of a mixed population in swabs, the presence in an MSSA of a
30 residual *SCCmec* right extremity fragment following the deletion of the *mecA* gene and/or non-specific amplification. To date, two concepts for determining resistance to methicillin carried specifically by *S. aureus* have been published:

- the SCC*mec* right extremity junction amplification concept (Hiramatsu et al. WO97/31125; EP 0 887 424; U.S. Pat. No. 6,156,507; and further, Huletsky and Rossbach WO02/099034 (2002); Huletsky et al. *J. Clin. Microbiol.* 42(5): 1875-1884 (2004));

- the immuno-enrichment concept described by François and co-workers (François, P et al. *J. Clin. Microbiol.* 41(1):254-260 (2003); WO02/082086), in which the immuno-enrichment is followed by amplification of three markers (*mecA* gene, *S. aureus*-specific marker, and *S. epidermidis*-specific marker).

[0008] The SCC*mec* right extremity junction concept is based on the amplification of a region covering the right extremity junction region of the SCC*mec* integration site. The principle is the following: the SCC*mec* cassette always integrates the *S. aureus* chromosome upstream of a *S. aureus* specific open reading frame called *orfX*; the PCR assay combines multiple forward primers located on the right part of the cassette, one reverse primer and a beacon probe, both located in the *S. aureus* chromosomal *orfX*, i.e., downstream of the right extremity junction of SCC*mec* with *orfX* ("right extremity junction region" of *orfX*).

15 Hiramatsu et al. describe a test with two forward primers in the right extremity junction region of the cassette to amplify the main SCC*mec* types described at that time (one primer for SCC*mec* types I and II and a second primer for type III). Huletsky et al. set forth that several MRSA strains were not detected if only the two forward primers described by Hiramatsu were used, and they determined new types of cassettes named as MREJ types

20 having sequence variations in the right part of the SCC*mec* cassette. A commercially available (Infectio Diagnostics Inc.) test combines (*refer to* Fig. 1) five forward primers located in the right part of the cassette (one primer was designed for the detection of MREJ types i and ii and the four others for the MREJ types iii, iv, v and vii), one reverse primer located in the *orfX*, and three generic beacons covering the same portion of the *orfX* region

25 and required to identify the *orfX* variants identified. This test is performed in real-time PCR. However, the specificity of this test as reported (Huletsky et al. 2004) shows that 4.6 % of MSSA (26 out of 569 tested) were misidentified. False-positive result has also been reported with another commercial test using a single-locus (right extremity SCC*mec* cassette-*orfX* junction) PCR assay (Rupp, J, et al., *J. Clin. Microbiol.* (44)6: 2317 (2006)).

30 [0009] Thus false positives remain an issue, and there is a strong need for an improved test for MRSA to reduce false positive results obtained with current tests. The challenge for such a test is that, due to the presence of a mixed population in nasal swabs, the following mixtures can be present: one or more of (1) MRSA, (2) methicillin sensitive coagulase-negative Staphylococci (*e.g.*, methicillin sensitive *Staphylococcus epidermidis*

(MSSE)), (3) methicillin sensitive *Staphylococcus aureus* (MSSA), and (4) methicillin resistant coagulase-negative Staphylococci (MR-CNS) (mainly methicillin resistant *Staphylococcus epidermidis* (MRSE)). Furthermore, there has been report of clinical MSSA isolates retaining SCCmec elements without the *mecA* gene. (Donnio, P.-Y., et al., J Clin Microbiol. 2005 August; 43(8): 4191–4193). Because only the presence of a MRSA will lead to decontamination of the carrier, the test must ensure that resistance to methicillin is carried by *S. aureus* and not by *S. epidermidis* (or coagulase (-) Staphylococcus strain). Thus, also, amplification and detection of the *mecA* gene (associated or not with a *S. aureus* specific marker) directly on the mixed population present on the swabs is not appropriate; indeed in both situations (MRSA+MSSE or MSSA+MRSE), both markers (*mecA* and *S. aureus* specific marker) will be detected whereas only the first situation with an MRSA is desired to be specifically detected by the clinician. The present invention addresses primary sources of MRSA false positives and thus provides a much-needed, improved test to detect MRSA that has not been addressed by currently available tests.

15

SUMMARY OF THE INVENTION

[0010] The present invention provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, comprising performing on the sample an amplification and detection reaction utilizing

- a. a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,
- b. a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
- c. a probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

wherein if the sample contains MRSA, hybridization of the probe is detected. In such methods as set forth herein, "a" primer, probe, etc. can mean one or more primer or probe, unless otherwise stated or the context dictates otherwise.

30

[0011] The present invention additionally provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, comprising

- (a) performing on the sample an amplification and detection reaction that detects the presence of a junction of an inserted SCCmec cassette and *Staphylococcus aureus* chromosomal DNA utilizing
- 1) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,
 - 2) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
 - 3) a first probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

- (b) performing on the sample an amplification and detection reaction that detects the presence of the *mecA* gene,

wherein if the sample contains MRSA, the presence of both the target junction and *mecA* in the sample is detected.

[0012] The present invention further provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, comprising

performing on the sample a multiplex amplification reaction wherein the amplification reaction comprises

- a. amplifying and detecting the presence of a junction of an inserted SCCmec cassette and *Staphylococcus aureus* chromosomal DNA utilizing
- 1) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,
 - 2) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
 - 3) a first probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

- b. amplifying and detecting the presence of *mecA* gene,
 wherein if the sample contains MRSA, the presence of both the target junction and *mecA* in
 5 the sample is detected.

[0013] Additionally, the present invention provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA, the method comprising

- a. performing on a sample an amplification reaction which can simultaneously amplify
 10 both (1) a junction of an inserted SCC*mec* cassette and *Staphylococcus aureus* chromosomal DNA and (2) a region of *mecA*, and
 b. detecting, within the products of the amplification, the presence or absence of each of the junction and *mecA*,

wherein if the sample contains MRSA, the presence of both the junction and *mecA* in
 15 the sample is detected.

[0014] The present invention further provides a method of identifying the presence in a sample of a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA which comprises the steps of:

20 bringing, in a single container, the biological sample in contact with

(a) a first oligonucleotide set comprising

(1) a first oligonucleotide having a nucleic acid sequence that specifically hybridizes to an extremity junction region of a SCC*mec* cassette, and

(2) a second oligonucleotide having a nucleic acid sequence that specifically
 25 hybridizes to a *Staphylococcus aureus* chromosomal DNA region flanking said SCC*mec* cassette to form a first reaction product of the biological sample and the first and second oligonucleotides, and

(b) a second oligonucleotide set comprising

(3) a third oligonucleotide having a nucleotide sequence that specifically
 30 hybridizes to a first region of *mecA* nucleic acid and

(4) a fourth oligonucleotide having a nucleotide sequence that specifically hybridizes to a second region of *mecA* nucleic acid to form a second reaction product of the biological sample and the third and fourth oligonucleotides; and

identifying the presence of MRSA by detecting both a first and a second reaction product. The identifying step can, in one embodiment, comprise contacting the first and second reaction products with (1) a first probe capable of specifically hybridizing with the first reaction product and (2) a second probe capable of specifically hybridizing with the second reaction product.

[0015] The present invention additionally provides kits for use in such methods. Specifically, the present invention provides a kit for detection of methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, comprising

- (a) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,
- (b) a second primer capable of specifically hybridizing in chromosomal *Staphylococcus aureus* DNA in a region of the extremity junction, and
- (c) a probe capable of specifically hybridizing fully within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified. Further, the present invention provides a kit for identifying the presence in a sample of a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA which comprises:

a) first amplification and detection oligonucleotide set comprising

- 1) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,
- 2) a second primer capable of specifically hybridizing in chromosomal *Staphylococcus aureus* DNA in a region of the extremity junction, and
- 3) a first probe selected from the group consisting of (1) a first probe capable of specifically hybridizing primarily within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction, and (2) a first probe capable of specifically hybridizing fully within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

77203-162

b) a second amplification and detection oligonucleotide set comprising

4) a third primer capable of specifically hybridizing to a first region of *mecA*,

5) a fourth primer capable of specifically hybridizing to a second region of *mecA*, and

5 6) a second probe capable of specifically hybridizing to a region of *mecA* between the first and second regions of *mecA*,

wherein each of the third primer and the fourth primer is oriented such that, under amplification conditions, the region of *mecA* between the first and second regions of *mecA* is amplified.

10 [0015A] The present invention as claimed relates to:

- a method of detecting in a sample a methicillin-resistant

Staphylococcus aureus (MRSA) having an insertion of an SCC*mec* cassette within

Staphylococcus aureus chromosomal DNA, comprising: performing on the sample an amplification and detection reaction utilizing: a) a first primer capable of specifically

15 hybridizing in an extremity junction region of the SCC*mec* cassette, b) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and c) a probe capable of specifically hybridizing to a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction, wherein each of the first primer and the second primer is oriented such that,
20 under amplification conditions, the junction is amplified, and wherein if the sample contains MRSA, hybridization of the probe is detected;

- a kit for detection of methicillin-resistant *Staphylococcus aureus* (MRSA)

having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA,

comprising: a) a first primer capable of specifically hybridizing in a right extremity junction

25 region of the SCC*mec* cassette, b) a second primer capable of specifically hybridizing in a right extremity junction region of chromosomal *Staphylococcus aureus* DNA, and c) a probe selected from the group consisting of (1) a probe capable of specifically hybridizing primarily within a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction, and (2) a probe capable of specifically hybridizing

30 fully within a region of the SCC*mec* cassette between the region with which the first primer is

77203-162

capable of hybridizing and the junction, wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified;

- a kit for detection of methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, comprising: (a) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette, (b) a second primer capable of specifically hybridizing to chromosomal *Staphylococcus aureus* DNA in the region of the extremity junction, and (c) a probe selected from the group consisting of (1) a first probe capable of specifically hybridizing primarily within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction, and (2) a first probe capable of specifically hybridizing fully within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction, wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified; and
- an oligonucleotide composition comprising: (1) a first oligonucleotide having a nucleic acid sequence that specifically hybridizes to an extremity junction region of a SCCmec cassette; (2) a second oligonucleotide having a nucleic acid sequence that specifically hybridizes to a *Staphylococcus aureus* chromosomal DNA region flanking said SCCmec cassette; and (3) a first probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] **Figure 1** shows generally the region of MRSA chromosome with the inserted SCCmec cassette, indicating the left and right extremity junctions.

- 25 [0017] **Figure 2** demonstrates generally the location of primers and probes in the right extremity junction region of SCCmec/orfX wherein five Primers 2 (“5P2,” arrows) are located in the right part of the cassette, one generic Primer 1 (“P1T7”) is located in *S. aureus* orfX and one generic beacon (“MB”) is located in *S. aureus* orfX (“Approach 1”).

- [0018] **Figure 3** demonstrates generally the location of primers and probes for an inventive method for detection of the right extremity junction of the SCCmec insertion in

77203-162

MRSA: five Primers 2 (“P2,” arrows) located in SCC*mec* right extremity junction region, Primer 1 (“P1,” angled line) located in *orfX*, and five specific probes (lines) located in the right part of the SCC*mec* cassette are used (“Approach 2”).

- [0019] **Figure 4** demonstrates multiplex amplification of *mecA* and the right extremity
 5 junction with the following primers and probes: in *mecA*, Primer 1 (“P1”), Primer 2 (“P2”) and probe (“beacon”); in “Right SCC*mec*-chromosome junction region,” right extremity junction region, Primer 1 (“1P1,” angled line), five Primers 2 (“5P2,” arrows) and five SCC*mec*-specific probes (“5 specific beacons,” lines angled at both ends).

DETAILED DESCRIPTION OF THE INVENTION

- 10 [0020] As discussed herein, the present invention provides that the identification of false positives by the previous molecular methods can be explained in some instances by the presence in MSSA strains of a residual SCC*mec* right extremity fragment following the

deletion of a chromosomal region containing *mecA* or the presence of an SCC which does not contain *mecA*. Additionally, as further disclosed herein, the present invention provides that some portion of the false positives can be due to non specific amplification; indeed, (as shown in FIG. 2) because the reverse primer and the beacons are located in the *orfX* which is common to both MRSA and MSSA, non specific annealing of the forward primer(s) on MSSA chromosome will lead to amplification and detection of MSSA. The present invention addresses both sources of false positives and provides an improved test.

[0021] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of skill in the art to which the disclosed invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are as described.

[0022] As used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise.

[0023] As stated above, the present invention provides solutions to reduce the lack of specificity due to MSSA detection reported with previous detection methods, which solutions can be utilized individually or, ideally, within the same test. Thus, the present invention describes the use of specific beacons (specific for SCC*mec* cassette) rather than generic beacons; this configuration will suppress detection of MSSA amplified due to non-specific amplification. Further, the present invention describes the advantage of combining in multiplex reaction the *mecA* gene detection with the cassette insertion region (*e.g.*, Right SCC*mec*-chromosome junction region) detection, compared to the detection of the cassette insertion region alone; this aspect of the invention can reduce detection of MSSA either due to presence of a residual SCC*mec* right extremity fragment following the deletion of a chromosomal region containing *mecA* or to the presence of an SCC which does not contain *mecA*.

[0024] In a preferred embodiment, the present invention provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA, comprising

performing on the sample an amplification and detection reaction utilizing

- a. a first primer capable of specifically hybridizing in an extremity junction region of the SCC*mec* cassette,

- b. a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
- c. a probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and wherein if the sample contains MRSA, hybridization of the probe is detected.

[0025] The genomic structure of MRSA has been characterized previously. As used in the claims, the “SCCmec cassette” (sometimes referred to as “mecDNA,” e.g., in Hiramatsu U.S. Pat. No. 6,156,507) has the definition as known in the art, i.e., an integrated adventitious DNA existing on a chromosome of MRSA or MR-CNS and including the *mec* gene complex, a set of site-specific recombinase genes (*ccrA* and *ccrB*), and terminal inverted and direct repeats (at both 3’ and 5’ ends). “*mecA* gene” includes all sequences necessary to encode PBP2a or PBP’ (Penicillin Binding Protein) conferring methicillin resistance.

[0026] As known in the art, insertion of the SCCmec cassette into the *S. aureus* chromosome creates two junctions, and two corresponding junction regions, of SCCmec DNA with *S. aureus* chromosomal DNA, wherein the SCCmec sequence is contiguous with the *S. aureus* chromosomal sequence. The junctions, therefore, are located at the left and right extremities of the SCCmec cassette (see FIG. 1). These two regions are named “Right SCCmec-Chromosome Junction” and “Chromosome-Left SCCmec junction” by Ito et al. (Antimicrob. Agents Chemother. May 2001 45(5): 1323-1336, “Structural Comparison of three Types of Staphylococcal Cassette Chromosome *mec* Integrated in the chromosome in Methicillin-Resistant *Staphylococcus aureus*”). At the right extremity junction, the *S. aureus* genomic sequence abutting the SCCmec cassette is the gene *orfX*, which is in some literature referred to as “IntM.” As used in the claims, “extremity junction region” is a region of either SCCmec cassette or *S. aureus* chromosomal nucleic acid within distance of either the right or the left extremity junction, or insertion site, such that a primer can, in a primer extension reaction or a transcription-type (e.g., NASBA or TMA) reaction, be extended across that junction, e.g., within 600 nt, 550nt, 500nt, 450nt, 400nt, 350nt, 300nt, 250nt, 200nt, 150nt, 100nt, or 50nt (in either direction) of the junction. Useful distances may vary depending upon the amplification technology used (e.g., it may be longer distances as new technologies are developed). “Extremity junction region,” therefore, depending upon context used, can refer to a region within the SCCmec DNA or a region within the *S. aureus* chromosomal

DNA; both uses refer to such DNA within distance of the junction such that an appropriately selected primer could, under appropriate, standard extension or amplification conditions, be extended, or transcribed, from it, in the direction of the junction, across the junction. That is, “an extremity junction region of the SCC*mec* cassette” would be a region within the SCC*mec* DNA near its abutment, or integration site, with the *S. aureus* chromosomal DNA; and “an extremity junction region of *orfX*” would be a region within the *orfX* DNA near an abutment with SCC*mec* DNA (an SCC*mec* integration site). Similarly, “an extremity junction region of chromosomal *S. aureus* DNA” would be a region within the chromosomal *S. aureus* DNA near an abutment with SCC*mec* DNA. Alternatively, this region may also be referred to as “chromosomal *S. aureus* DNA in the region of the SCC*mec* extremity junction.” Thus, “right extremity junction region” refers to the region surrounding the junction on the right side of the SCC*mec* cassette, and “left extremity junction region” refers to the region surrounding the junction on the left side of the SCC*mec* cassette (see FIG. 1).

[0027] To further provide a useful method for detecting MRSA, the present invention provides an amplification and detection which detects both the presence of a SCC*mec* insertion junction and *mecA* sequences. More specifically, the present invention additionally provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA, comprising

(a) performing on the sample an amplification and detection reaction that detects the presence of a junction of an inserted SCC*mec* cassette and *Staphylococcus aureus* chromosomal DNA utilizing

- 1) a first primer capable of specifically hybridizing in an extremity junction region of the SCC*mec* cassette,
- 2) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
- 3) a first probe capable of specifically hybridizing to a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

(b) performing on the sample an amplification and detection reaction that detects the presence of *mecA* gene,

wherein if the sample contains MRSA, the presence of both the target junction and *mecA* in the sample is detected.

Such a method is particularly helpful in discriminating MRSA from those, somewhat rare, MSSAs from which *mecA*, but not the complete SCC*mec* cassette, have been deleted.

5 [0028] Advantageously, one can perform a multiplex amplification reaction to detect both the presence of a SCC*mec* insertion junction and *mecA* sequences. Thus, the present invention provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA, comprising

10 performing on the sample a multiplex amplification reaction wherein the amplification reaction comprises

a. amplifying and detecting the presence of a junction of an inserted SCC*mec* cassette and *Staphylococcus aureus* chromosomal DNA utilizing

- 15 1) a first primer capable of specifically hybridizing in an extremity junction region of the SCC*mec* cassette,
- 2) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
- 3) a first probe capable of specifically hybridizing to a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing
- 20 and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

b. amplifying and detecting the presence of *mecA* gene,

25 wherein if the sample contains MRSA, the presence of both the target junction and *mecA* in the sample is detected.

[0029] By “amplify the junction” is meant performing an amplification reaction that produces an amplification product that includes sequences corresponding to nucleic acids both within SCC*mec* abutting the junction and within *S. aureus* chromosomal DNA abutting the same junction. By “amplifying and detecting the presence of *mecA* gene” is meant

30 amplifying and detecting within the sample any portion of a *mecA* gene, for example, the region between primers comprising a nucleic acid sequence set forth in SEQ ID NO: 15 and 16 (which can be detected, for example, utilizing a probe comprising the nucleic acid sequence set forth in SEQ ID NO: 14). Primers and probes can be readily designed for hybridization to the known *mecA* sequence.

[0030] The term “multiplex amplification reaction” means that the specific reagents for amplification of more than one target are contacted together, such that more than one amplification can occur within the same reaction container. Additionally, detection reagents for more than one target can be included. Thus one can conduct a multiplex amplification and detection reaction by placing into contact all of the specific reagents for amplification and detection of more than one target. Thus, in a multiplex reaction, one can amplify multiple target regions in the same reaction. “Simultaneous amplification” wherein individual reactions are allowed to proceed at the same time, but the reactants for more than one amplification reaction are not necessarily all within the same reaction container, or tube, rather, the more than one reactions can be carried out in separate reaction containers, can also be utilized if a multiplex is not desired or feasible. It is understood that, even in a multiplex amplification reaction, each reaction will occur at whatever pace the individual reactions proceed under the provided conditions. Detection can also be “simultaneous,” meaning that, if appropriate probes for each reaction in the reaction container are included, under the appropriate conditions, detection of more than one target can be achieved in either a single reaction container (multiplex) or in more than one reaction container (appropriate probes distributed to the relevant reaction container). Such detection can be performed, if desired, in the same reaction container as the multiplex or simultaneous amplification reaction, and, further, can be performed while amplification reactions continue (*i.e.*, real-time).

[0031] Thus, the present invention provides a method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, the method comprising

a. performing on a sample a multiplex amplification reaction which can amplify both (1) a junction of an inserted SCCmec cassette and *Staphylococcus aureus* chromosomal DNA and (2) a region of *mecA*, and

b. detecting, within the products of the amplification, the presence or absence of each of the junction and *mecA*,

wherein if the sample contains MRSA, the presence of both the junction and *mecA* in the sample is detected.

[0032] Amplification of the junction of an inserted SCCmec cassette and *Staphylococcus aureus* chromosomal DNA can be advantageously achieved by utilizing

- 1) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,

2) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and

3) a first probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of specifically

5 hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified. This probe can be selected to specifically hybridize fully within the SCCmec region. The presence or absence of either the junction or *mecA* can be determined by performing whatever analysis provides detection of
10 the product, e.g., if a labeled probe is used, detection of the hybridized label by the appropriate detection device. Lack of a detectable signal indicates the absence of the target; perception of a detectable signal indicates presence of the target.

[0033] Primers and probes used in a reaction of this invention are capable of specifically hybridizing with a target nucleic acid. Specific hybridization is known in the art,
15 and, typically, specific hybridization is achieved through nucleic acid identity or high similarity of the primer/probe with the target nucleic acid and/or through use of stringent hybridization conditions (e.g., stringent temperature and/or salt conditions). Specific hybridization provides selective hybridization to the target within the reaction.

[0034] A primer "oriented such that, under amplification conditions, the junction is
20 amplified" includes a primer oriented such that, upon hybridization to its specific target nucleic acid, and upon initiation of an amplification reaction including the primer, an amplicon is formed that includes the junction. Such a reaction is designed to amplify across the junction (i.e., to be in sufficiently close proximity of the junction so that a typical amplification reaction would extend across the junction). Thus a primer pair useful for
25 amplifying a junction will typically hybridize to two regions that surround the junction and each primer will be oriented to hybridize in a 5'-3' direction toward the junction. Typically, the primer would be designed to hybridize within 600nt, 500nt, 400nt, 350nt, 300nt, 250nt, 200nt 150nt, 100nt, 50nt, 30nt, 25nt, 20nt, etc. of the junction. A probe for detecting an amplification product is therefore selected to be capable of specifically hybridizing to a
30 region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction. In certain embodiments, such a probe can specifically hybridize fully within or primarily within SCCmec cassette. In one embodiment, in which the probe specifically hybridizes primarily within SCCmec cassette, the region to which the probe hybridizes can additionally include the junction and, therefore, at least one, or two or three or

a few nucleotides of *orfX* that abut the junction. If one, two three or a few nucleotides of the probe hybridize to a region of *orfX* abutting the junction, the probe will, however, be capable of at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% of its nucleotides, preferably contiguous, specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction. Thus, for an amplification of the right extremity junction region, the probe is selected to specifically hybridize to a region of the SCCmec cassette that is 3', or downstream, of the 3' end of the SCCmec cassette primer (when hybridized) and upstream of the junction. Typically, the primer is selected such that amplification product synthesized utilizing it and a second primer (located in an *S. aureus* genomic sequence) will be of approximately 200-350nt in length. While PCR amplification can be designed to generate longer amplicons (e.g., 250, 300, 350, 400, 500, 600, 700, 800, 900, 1000nt), preferred amplicon lengths for either transcription-based (e.g., NASBA or TMA) or PCR-type reactions will be within about 200-300nt (e.g. 150, 200, 250, 300nt) in length. Additionally, for a multiplex amplification reaction, whether transcription-based or PCR-based, an amplicon in the range of 200-300nt or shorter is preferable, to enhance sensitivity of the test.

[0035] As used in the claims, "amplification conditions" are those appropriate for a selected amplification reaction, as are known to those of skill in the art, such as are utilized in various amplification reactions. Such conditions can be optimized for a specific reaction, primers, etc. as also known by the skilled artisan. As is known, such amplification conditions include contact with the required reagents for the amplification, e.g., nucleotides and enzymes, as well as the appropriate selected temperature, salt and pH conditions, among other aspects. Furthermore, as used in the claims, a primer or probe may be a primer or probe set, i.e., multiple primers or probes. Such primer/probe sets can be utilized in a reaction in which more than one type or subtype of MRSA is desired to be amplified and/or detected, and wherein the nucleic acid sequence of the target MRSA region selected for hybridization of the primer and/or probe varies among types and/or subtypes. Individual primers/probes can be designed for each type or subtype, as exemplified herein.

[0036] Specific primers useful for amplifying extremity junction regions can readily be designed, given the teachings herein. Primers for hybridizing to SCCmec right extremity region can include the primers set forth in SEQ ID NOs: 1, 2, 3, 4, 5, and 6, as well as primers comprising these sequences and primers consisting essentially of these sequences. Specific primers for hybridizing to *orfX* can include the primer set forth in SEQ ID NO: 13,

as well as a primer comprising this sequence and a primer consisting essentially of this sequence. Probes for hybridizing to SCC*mec* right extremity region can include the probes set forth in SEQ ID NOs: 7, 8, 9, 10, 11, and 12, as well as probes comprising these sequences and probes consisting essentially of these sequences. Primers for hybridizing to *mecA* can include the primers set forth in SEQ ID NOs: 15 and 16 as well as primers comprising these sequences and primers consisting essentially of these sequences. Either can readily be adapted as a P1-type or a P2 type (for NASBA-type amplifications). Probes for hybridizing to *mecA* can include the probe set forth in SEQ ID NO: 14 as well as probes comprising this sequence and probes consisting essentially of this sequence.

10 [0037] The present invention further provides a method of identifying the presence in a sample of a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA which comprises the steps of:

bringing, in a single container, the biological sample in contact with

- 15 a. a first oligonucleotide set comprising
 - (1) a first oligonucleotide having a nucleic acid sequence that specifically hybridizes to an extremity junction region of a SCC*mec* cassette, and
 - (2) a second oligonucleotide having a nucleic acid sequence that specifically hybridizes to a *Staphylococcus aureus* chromosomal DNA region flanking
 - 20 said SCC*mec* cassette to form a first reaction product of the biological sample and the first and second oligonucleotides, and
 - b. a second oligonucleotide set comprising
 - (3) a third oligonucleotide having a nucleotide sequence that specifically hybridizes to a first region of *mecA* nucleic acid and
 - 25 (4) a fourth oligonucleotide having a nucleotide sequence that specifically hybridizes to a second region of *mecA* nucleic acid to form a second reaction product of the biological sample and the third and fourth oligonucleotides; and
- identifying the presence of MRSA by detecting both a first and a second reaction product. The first and second reaction products can be formed by an amplification reaction.
- 30 Conditions under which the reagents are brought in contact with the sample can be conditions appropriate for a selected amplification reaction. The identifying step can, in one embodiment, comprise contacting the first and second reaction products with (1) a first probe capable of specifically hybridizing with the first reaction product, if present,

and (2) a second probe capable of specifically hybridizing with the second reaction product, if present.

[0038] As used herein, an oligonucleotide "having" a nucleic acid sequence included in a portion of target DNA means the sequence has sufficient identity to the target DNA sequence, or its complement, to specifically and selectively hybridize to that target DNA under stringent hybridization conditions. It includes nucleic acid sequences having full sequence identity to the sequence. In the single container can be included all components of a reaction mixture, tailored to the specific amplification and detection method utilized. Thus, a "reaction mixture" can include all the necessary reactants for performing a reaction, which may include, but not be limited to, buffering agents to maintain pH at a selected level during a reaction, salts, co-factors, scavengers, and the like.

[0039] Generally, amplification reactions producing amplicons (the product of a polynucleotide amplification reaction) are "template-driven" in that base pairing of reactants, either nucleotides or oligonucleotides, have complements in a template polynucleotide that are required for the creation of reaction products. In one aspect, template-driven reactions are primer extensions with a nucleic acid polymerase or oligonucleotide ligations with a nucleic acid ligase. Amplification can include any known or newly designed method of amplification, including those used in published methods (e.g., transcription-based amplification such as transcription-mediated amplification (TMA) and nucleic acid sequence-based amplification NASBA (as exemplified herein), and cycling nucleic acid amplification technologies (thermocycling) such as polymerase chain reaction (PCR), reverse transcriptase PCR (RT-PCR), and ligase chain reaction (LCR), and any method of amplification, e.g., sustained sequence replication (3SR), strand displacement amplification (SDA), branched DNA (bDNA), cycling probe technology (CPT), solid phase amplification (SPA), rolling circle amplification technology (RCA), solid phase RCA, anchored SDA and nuclease dependent signal amplification (NDSA), all of which are known to the skilled artisan. An amplification reaction may be a "real-time" amplification if a detection chemistry is available that permits a reaction product to be measured as the amplification reaction progresses, e.g. real-time PCR or real-time NASBA. Thus this invention includes the use of any nucleic acid amplification method or any other procedure which may be used to increase the sensitivity and/or the rapidity of nucleic acid-based diagnostic tests. The present invention also includes the use of any detection technology including post-amplification detection technologies, any amplification technology combined with detection, any hybridization nucleic acid chips or array technologies, and any amplification chips or combination of amplification and

hybridization chip technologies. Detection and identification by any nucleotide sequencing method is also within the present invention.

[0040] While any suitable amplification methods can be utilized for the present assay, it is noted that amplification reactions that utilize a restriction enzyme, such as DNA-

5 NASBA, can contribute an additional level of specificity to the assay. For example, if one selects a DNA-NASBA amplification reaction having a P1 capable of hybridizing in the *S. aureus orfX* region, the requirement that the correct restriction site be present in the genomic region of the *S. aureus* for amplification to occur greatly reduces or eliminates the likelihood of amplification of a methicillin-resistant coagulase-negative Staphylococcus (*e.g.*,
10 methicillin-resistant *S. epidermidis*) since it should not have the requisite restriction site. This is particularly useful in “approach 2” reactions of this present invention, wherein the probe to detect the junction between bacterial genomic DNA and SCC*mec* DNA is designed to hybridize to SCC*mec* DNA (or primarily to SCC*mec* DNA, with, in one embodiment, one, two, three or a few probe nucleotides hybridizing across the junction to *orfX* DNA), rather
15 than to genomic bacterial DNA. The combination of these features provides a highly specific assay for MRSA. Furthermore, additional specificity can be attained with PCR-type reactions by increasing the stringency of hybridization conditions of the primers. Thus, by increasing the stringency of the selected amplification reaction, by any selected means, one can increase the overall specificity of MRSA assays that include detection reactions that
20 utilize an SCC*mec*-hybridizing probe.

[0041] Furthermore, amplification can be conducted in additional manners to enhance amplification, for example, for transcription-based amplification reactions, (*e.g.*, TMA, NASBA), a blocked “P1” (*i.e.* promoter-bearing) primer can be utilized, if desired. Such blocked primers are typically blocked at their 3’ end from extension of the primer, by a
25 chemical moiety, such as dabsyl, others being known in the art.

[0042] A variety of detection methods can be utilized in this invention. Detection methods utilizing nucleic acid probes are well known in the art. Probes of the present kits and/or for use in the present methods can be labeled by any selected label suitable for the detection method chosen, many of which are known in the art, such as a phosphatase (*e.g.*,
30 alkaline phosphatase), biotin, avidin, a peroxidase (*e.g.*, horseradish peroxidase), digoxigenin, a fluorescent dye (such as Cy3 and Cy5 dyes, fluorescein, FAM, ROX), a chemiluminescent label, a chromophoric label, a radioactive label (*e.g.*, a radioisotope) and a ligand. Probe designs for different detection methods can be utilized, such as target-capture, HPA, TaqMan,

molecular beacons and sandwich hybridization. Hybridization conditions can be selected in accordance with the type of probe and the type of detection reaction selected.

[0043] The present method further provides useful kits for use in such amplification and detection methods. Specifically provided is a kit for detection of methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, the kit comprising a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette, a second primer capable of specifically hybridizing to chromosomal *Staphylococcus aureus* DNA in the region of the extremity junction, and a probe capable of specifically hybridizing fully within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction, wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified. In one preferred embodiment, the extremity junction region is the right extremity junction region. In another embodiment, the extremity junction region is the left extremity junction region. In one embodiment, the first and second primers are provided in a single container. In another embodiment, the first and second primers and the probe are provided in a single container. Thus, a kit of the invention can comprise a first container comprising a first primer capable of specifically hybridizing in a right extremity junction region of the SCCmec cassette and a second primer capable of specifically hybridizing in a right extremity junction region of chromosomal *Staphylococcus aureus* DNA and a second container comprising a probe capable of specifically hybridizing fully within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction. Additionally, a kit of the present invention can comprise a container comprising a first primer capable of specifically hybridizing in a right extremity junction region of the SCCmec cassette, a second primer capable of specifically hybridizing in a right extremity junction region of chromosomal *Staphylococcus aureus* DNA, and a probe capable of specifically hybridizing fully within a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction.

[0044] The present invention further provides a kit for identifying the presence in a sample of a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA which comprises:

a) first amplification and detection oligonucleotide set comprising

1) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,

2) a second primer capable of specifically hybridizing to chromosomal *Staphylococcus aureus* DNA in the region of the extremity junction, and
 3) a first probe selected from the group consisting of (a) a first probe capable of specifically hybridizing primarily within a region of the SCC*mec* cassette between
 5 the region with which the first primer is capable of hybridizing and the junction, and (b) a first probe capable of specifically hybridizing fully within a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under
 10 amplification conditions, the junction is amplified, and

b) a second amplification and detection oligonucleotide set comprising

4) a third primer capable of specifically hybridizing to a first region of *mecA*,

5) a fourth primer capable of specifically hybridizing to a second region of *mecA*,
 and

15 6) a second probe capable of specifically hybridizing to a region of *mecA* between the first and second regions of *mecA*,

wherein each of the third primer and the fourth primer is oriented such that, under amplification conditions, the region of *mecA* between the first and second regions of *mecA* is amplified.

20 [0045] The present invention additionally provides a kit for identifying the presence in a sample of a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA which comprises:

a) first amplification and detection oligonucleotide set comprising

1) a first primer capable of specifically hybridizing in a right extremity junction
 25 region of the SCC*mec* cassette,

2) a second primer capable of specifically hybridizing to chromosomal *Staphylococcus aureus* DNA in the region of the right extremity junction, and

3) a first probe capable of specifically hybridizing primarily, or fully, within a region of the SCC*mec* cassette between the region with which the first primer is
 30 capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and

b) a second amplification and detection oligonucleotide set comprising

4) a third primer capable of specifically hybridizing to a first region of *mecA*,

5) a fourth primer capable of specifically hybridizing to a second region of *mecA*,
and

6) a second probe capable of specifically hybridizing to a region of *mecA* between
the first and second regions of *mecA*,

5 wherein each of the third primer and the fourth primer is oriented such that, under
amplification conditions, the region of *mecA* between the first and second regions of *mecA* is
amplified.

[0046] A probe capable of specifically hybridizing “fully within” a region of the
SCC*mec* cassette between the region with which the first primer is capable of hybridizing and
10 the junction means a probe that hybridizes only within SCC*mec* cassette, and does not
hybridize across the junction, under selected conditions for specific hybridization. A probe
capable of specifically hybridizing “primarily within” SCC*mec* cassette means a probe that
hybridizes within SCC*mec* cassette and can additionally hybridize across the junction and,
therefore, includes at least one, or two or three or a few nucleotides of *orfX* that abut the
15 junction. If one, two three or a few nucleotides of the probe hybridize to a region of *orfX*
abutting the junction, the probe will, however, be capable of at least 75%, at least 80%, at
least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at
least 96%, at least 97%, at least 98%, or at least 99% of its nucleotides, preferably contiguous
nucleotides, specifically hybridizing to a region of the SCC*mec* cassette between the region
20 with which the first primer is capable of hybridizing and the junction. A probe hybridizing
primarily within a region of the SCC*mec* cassette can include a probe hybridizing fully within
a region of the SCC*mec* cassette.

[0047] Probes of this invention, including those included in such kits, can
advantageously be labeled for detection, as known by persons of skill in the art. Labels can
25 appropriately be selected for the specific design and type of amplification reaction to be
performed. Primer and probe reagents can be provided in any of several states, including
dried, lyophilized, pelleted, spray-dried, or in liquid.

[0048] Kits of this invention can include additional elements, such as reagents for a
selected amplification method (e.g., amplification enzyme(s), buffer(s), and/or restriction
30 enzyme(s), among others), control(s), reaction container(s), and the like. In a specific
embodiment, a first, second, third and fourth primers are provided in a single container. In
another embodiment, the first and second amplification and detection oligonucleotide sets are
provided in a single container. Thus, such kits can be useful for performing multiplex
amplifications.

[0049] In one embodiment a kit is provided which comprises a container comprising a first primer capable of specifically hybridizing in an extremity junction region, preferably the right extremity junction region, of the SCC*mec* cassette; a second primer capable of specifically hybridizing in chromosomal *Staphylococcus aureus* DNA in the region of an extremity junction, specifically the right extremity junction region if the first primer hybridizes in the right extremity junction region; a third primer capable of specifically hybridizing to a first region of *mecA*; and a fourth primer capable of specifically hybridizing to a second region of *mecA*. Such a kit can further comprise a container comprising a first probe capable of specifically hybridizing fully within a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction and a second probe capable of specifically hybridizing to a region of *mecA* between the first and second regions of *mecA*. Additionally, in another embodiment of the invention, a kit can be provided which comprises a container comprising a first primer capable of specifically hybridizing in an extremity junction region of the SCC*mec* cassette, preferably the right extremity junction region; a second primer capable of specifically hybridizing in chromosomal *Staphylococcus aureus* DNA in the region of an extremity junction region, specifically the right extremity junction region if the first primer hybridizes in the right extremity junction region; a third primer capable of specifically hybridizing to a first region of *mecA*; a fourth primer capable of specifically hybridizing to a second region of *mecA*; a first probe capable of specifically hybridizing fully within a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction; and a second probe capable of specifically hybridizing to a region of *mecA* between the first and second regions of *mecA*.

[0050] A primer capable of specifically hybridizing in a right extremity junction region of the SCC*mec* cassette can be a nucleic acid comprising a sequence selected from the group consisting of: SEQ ID NO: 1, 2, 3, 4, 5, and 6. Furthermore, such a primer can consist essentially of, or consist of, a sequence selected from the group consisting of: SEQ ID NO: 1, 2, 3, 4, 5, and 6. One or more such primers can be selected; in a preferred embodiment, several primers, each specific for a different type of MRSA, are selected for inclusion in a kit. Such primers are useful in the methods of this invention.

[0051] A primer capable of specifically hybridizing in chromosomal *Staphylococcus aureus* DNA in the region of the right extremity junction can preferably be a primer that is capable of specifically hybridizing to *orfX*. More specifically, in one embodiment, this primer can comprise the nucleic acid set forth as SEQ ID NO: 13. Furthermore, such a primer

can consist essentially of, or consist of, the nucleic acid set forth as SEQ ID NO: 13. Such primers are useful in the methods of this invention.

[0052] A probe capable of specifically hybridizing fully within a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction can, in one embodiment, comprise five or more probes, each capable of hybridizing specifically to a different type of MRSA. In a specific embodiment, such a probe can comprise a sequence selected from the group consisting of SEQ ID NOs: 7, 8, 9, 10, 11, and 12. Furthermore, such a probe can consist essentially of, or consist of, a sequence selected from the group consisting of SEQ ID NOs: 7, 8, 9, 10, 11, and 12. Such probes are useful in the methods of this invention.

[0053] In certain kits of the invention, primers and/or probes for detection of *mecA* gene are included. Such a kit can include a primer set specific for the *mecA* gene, specifically, a first primer capable of specifically hybridizing to a first selected region of *mecA* wherein the first primer is oriented such that, under amplification conditions, the region of *mecA* between the first region of *mecA* and a second selected region of *mecA* is amplified, and a second primer capable of specifically hybridizing to the second selected region of *mecA* wherein the second primer is oriented such that, under amplification conditions with the first primer, the region of *mecA* between the first region of *mecA* and the second region of *mecA* is amplified. In one embodiment, these *mecA* primers can include one or both of SEQ ID NOs: 15 and 16. The kit can further include a probe specific for the region of *mecA* between the two selected *mecA* primers. In a specific embodiment, such a *mecA* probe can comprise the nucleic acid sequence set forth in SEQ ID NO: 14. Furthermore, such a *mecA* probe can consist essentially of the nucleic acid sequence set forth in SEQ ID NO: 14. By providing primer and probe sets for amplifying and detecting both SCC*mec* cassette and *mecA*, within the same kit for use within the same reaction container, a useful multiplex kit for detecting MRSA can be provided.

[0054] It is noted that references to primer and probe sequences that include thymidine can be readily adapted to utilize uridine in substitution for thymidine, where useful for the particular assay. Furthermore, nucleotides may be modified by addition of chemical groups, or substitution of individual residues by analogues (e.g., 2'-O-methoxy versions). Additional such modified nucleotides are known in the art; some examples include hydroxymethyl nucleotides, methylated nucleotides, fluorinated nucleotides, alpha thio phosphate nucleotides, amine-modified nucleotides, methoxy nucleotides, carboxymethyl nucleotides, thio nucleotides, inosine, dihydrouridine, pseudouridine, wybutosine, queuosine,

C7dGTP. Additional modified nucleotides are found in U.S. Pat. Nos. 5,405,950 and 5,633,364 (both, Mock and Lovern). Furthermore, a probe can comprise DNA, RNA, modified DNA or RNA, PNA, other synthetic nucleic acids or nucleic acid substitutes that use nucleotide bases as means of selectively hybridizing to a target.

5 [0055] Throughout this application, a particular oligonucleotide may be exemplified in use as a particular primer type (e.g. NASBA P1-type (linked to a sequence that provides a promoter region when in double-stranded form) or P2-type (used alone or linked to a tag oligonucleotide) primer); however, such use should not limit the use(s) for which the oligonucleotide may be useful. For example, a primer exemplified as a P1 primer may be
10 useful as a P2-type primer. Additionally, primers can be adapted for other amplification methods (e.g., the T7 polymerase promoter region of P1 (for NASBA or TMA) removed for PCR use), as is known by the skilled artisan.

[0056] The nucleic acid sequence of the T7 promoter is well-known to persons skilled in the art, and though a particular sequence is exemplified herein, functional equivalents
15 having slight variations, known in the art or newly designed, may be selected. In a preferred embodiment, the sequence of the T7 promoter is that set forth in SEQ ID NO:17.

[0057] The present method can be utilized on any selected sample, such as a direct patient sample, e.g., nasal or inguinal swab, throat swab, rectal swab, samples from wounds, all particularly suitable for screening, as well as particularly suitable for diagnosis,
20 bronchoalveolar lavage or blood (e.g., septicemia or blood culture). Such samples typically contain a mixed population of organisms. Additionally, if desired, this method can be applied to a sample having only a single bacterial species or strain, e.g., samples utilizing isolation, culture, capture, and/or enrichment of MRSA.

[0058] The present invention is exemplified in the following examples.
25

EXAMPLES

[0059] A study was initiated using NASBA as amplification. Two main approaches were investigated : approach 1 (generic *S. aureus* (*orfX*) probes) and approach 2 (SCC*mec*-
30 specific probes). “Approach 1” is very similar to the configuration used in previous tests; it is based on the use of 5 P2 located in the right part of the cassette, one generic P1 located in *S. aureus orfX* and one generic beacon located in *S. aureus orfX* (FIG. 2). The herein described “approach 2” uses the same five P2 and P1 as approach 1, but five SCC*mec*-specific beacons located in the right part of the cassette are used instead of the generic beacon

(see FIG. 3). Furthermore, is it also described and shown herein that a multiplex reaction utilizing approach 2 with amplification and detection of *mecA* was found to be highly effective at detecting MRSA.

5 **Amplification conditions**

[0060] For all experiments, the following DNA NASBA conditions were used: Chromosomal DNA was used as input material for amplification. For each strain, bacterial suspensions were standardized to 0.5 McFarland (1.5×10^8 CFU/ml) using densitometer (bioMérieux, Marcy l'Etoile, France). DNA was released from bacterial cells using a
10 mechanical lysis technique with glass beads and vortex, and finally serial dilutions were prepared from the lysed suspension.

[0061] Amplification was performed using standard NASBA reagents (40 mM Tris-HCl pH 8.5, 12 mM $MgCl_2$, 90 mM KCl, 15% v/v DMSO, 5 mM DTT, 1 mM each dNTP, 2 mM ATP, 2 mM CTP, 2 mM UTP, 1.5 mM GTP, 0.5 mM ITP ; the concentration of primers
15 (forward primer =P1 and reverse primer= P2) was 0.2 μ M when tested in monoplex and 0.1 μ M when tested in multiplex; the concentration of FAM labeled molecular beacon probe(s) was 0.01 μ M and 0.1 μ M for Cy5 labeled beacon probe.

[0062] Concentration of restriction enzyme (Sau3A-I) was 0.05 Units per μ l in the NASBA reaction. Incubation of the mixture for 15 min at 41°C enabled the restriction
20 enzyme(s) to cut the DNA, this step was followed by 5 min at 95°C for DNA denaturation and restriction enzyme(s) degradation, finally a 3 min step at 41°C was performed before adding NASBA enzymes (0.08 units RNase H, 32 units T7 RNA polymerase, 6.4 units AMV reverse transcriptase and 2.1 μ g BSA). The reaction mixture was mixed by gently vortexing and short centrifugation, and the amplification and real-time detection was started. The
25 reaction mixture was incubated at 41°C in the NucliSens EasyQ Analyzer (NucliSens, BioMérieux) for 90 minutes with fluorescence monitoring every 30 seconds. For FAM detection, the reactions were excited at 485 nm and the emission signal was measured at 518 nm. For ROX, excitation and emission were performed at 578 nm and 604 nm respectively and for Cy5, excitation and emission were performed at 646 nm and 678 nm respectively.

30 [0063] Primers and probes utilized are set forth below. For NASBA reactions, "primer 1" or "P1" is traditionally (as here) used to indicate the primer having at its 5'end a sequence that provides a promoter region (here, for T7 polymerase) when in double-stranded form. The second primer for NASBA, a non-promoter-linked primer, is traditionally, as here, labeled "Primer 2" or "P2."

[0064] The following Table 1 provides sequences for primers and probes utilized in the Examples.

Table 1

Function	SEQ ID NO	Description	Sequence 5' → 3'
P2	1	<i>SCCmec Right extremity junction region MREJ type v</i>	CTCTGCTTTATATTATAAAATTACGGCTG
P2	2	<i>SCCmec Right extremity junction region MREJ type iii</i>	ATTTTCATATATGTAATTCCTCCACATCTC
P2	3	<i>SCCmec Right extremity junction region MREJ type i and ii</i>	AAGACTGCGGAGGCTAA
P2	4	<i>SCCmec Right extremity junction region MREJ type vii</i>	TATTCTTCAAAGATTGAGC
P2	5	<i>SCCmec Right extremity junction region MREJ type iv</i>	CAAATATTATCTCGTAATTTAC
P2	6	<i>SCCmec Right extremity junction region SCCmec type V</i>	TCTAATTTATTTAACATAAAATCAATCCT
Beacon-FAM	7	<i>SCCmec Right extremity junction region MREJ type i and ii</i>	FAM-CGGCGCGT CAAAAATCATGAACCTC ATTACTTATGCGCCG-DabSyl
Beacon-FAM	8	<i>SCCmec Right extremity junction region MREJ type iii</i>	FAM-CGAGCGC AAATTATACACAACCTAA TTTTTAGTGCGCTCG-DabSyl
Beacon-FAM	9	<i>SCCmec Right extremity junction region MREJ type vii</i>	FAM-CGGAGCTAATTTAATAATTTTCTCAT ATTTTTTAGCTCCG-Dabsyl
Beacon-FAM	10	<i>SCCmec Right extremity junction region MREJ type iv</i>	FAM-CGTAACG GATAAAAAACCGCATCATTTGACGTTACG - Dabsyl

Beacon-FAM	11	<i>SCCmec Right extremity junction region MREJ type v</i>	FAM-CGTAGCG GCTGAAATAACCGCATCATTTA CGCTACG-Dabsyl
Beacon-FAM	12	<i>SCCmec Right extremity junction region SCCmec type V</i>	FAM- CGTCACGTAAAATATATTATACAC AATCCGTTTCGTGACG -Dabsyl
P1	13	<i>OrfX Right extremity junction region</i>	aattctaatacgactcactatagggagag TCAAACGGCCTGCACAAGGA
Beacon-Cy5	14	<i>mecA gene</i>	Cy5 -CGTACGGGATCATAGCGTCATTATT CGTACG-Dabsyl
P1	15	<i>mecA P1</i>	aattctaatacgactcactatagggagag GTATTGGCCAATTCCACATTGTTTC
P2	16	<i>mecA P2</i>	CATTGATCGCAACGTTCA
T7 polymerase promoter sequence	17		AATTCTAATACGACTCACTATAGGG
Beacon-FAM	18	Generic- orfX (SCCmec)	FAM - CGTACGGTAGTTACTGCGTTGTAAGACGTAC G -Dabsyl

EXAMPLE 1: NASBA results obtained with approach 1 (generic beacon NASBA) on 21 MSSA strains

- 5 [0065] “Approach 1” is based on the use of 5 P2 located in the right part of the cassette, one generic P1 located in *S. aureus orfX* and one generic beacon located in *S. aureus orfX* (FIG. 2). A DNA NASBA was performed using five *SCCmec* cassette-specific Primers 2 (P2) (SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5), one generic P1 (SEQ ID NO: 13) and one generic FAM-labeled beacon (SEQ ID NO: 18)
- 10 NASBA curves were obtained for MRSA (6 strains belonging to MREJ types i, ii, iii, iv, v and vii) and MSSA. Lysate input corresponds to 10⁵ CFU per NASBA. Max signal ratio (ratio between final signal and initial background) are given in Table 2. This experiment

shows positive max signal ratio for 5 out of 21 MSSA strains (24%) with the generic beacon NASBA (approach 1).

Table 2: max signal ratio obtained for MSSA with generic beacon NASBA (positive signals are in bold italics)

5

strain	Max signal ratio
MSSA 10	1.01
MSSA 11	<i>1.69</i>
MSSA 12	1.02
MSSA 13	1.07
MSSA 14	1.08
MSSA 15	1.00
MSSA 16	<i>2.06</i>
MSSA 17	<i>2.38</i>
MSSA 18	<i>1.52</i>
MSSA 19	1.00
MSSA 23	1.02
MSSA 24	1.00
MSSA 25	1.00
MSSA 26	1.00
MSSA 27	<i>1.38</i>
MSSA 28	1.00
MSSA 29	1.00
MSSA 30	1.00
MSSA 31	1.05
MSSA 32	1.00
MSSA 1034	1.00

EXAMPLE 2: NASBA results obtained with approach 2 (specific beacons NASBA) on 6 MRSA strains and 18 MSSA strains.

10 [0066] To overcome non specific detection of MSSA, “approach 2” was defined and developed. Specifically, beacons located in the right part of the SCCmec cassette instead of in the *orfX* were defined. “Approach 2” uses five specific Primers 2 (P2) and one generic P1, but five specific beacons located in the right part of the cassette are used instead of a generic beacon (see FIG. 3).

15 [0067] A DNA NASBA was performed using five SCCmec cassette-specific primers 2 (P2) (SEQ ID NOs: 1-5), one generic (*orfX*) primer 1 (P1) (SEQ ID NO: 13) and five SCCmec cassette-specific FAM-labeled beacons (SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11). Six MRSA and 18 MSSA strains were tested using a lysate as target corresponding to 10⁵ CFU per NASBA. NASBA curves were
20 obtained on MRSA and MSSA. Max signal ratio obtained are given in Table 3.

[0068] This experiment shows that only one out of 18 MSSA was detected with specific beacons approach (approach 2); this detected MSSA strain is expected to possess the right part of the cassette without the *mecA* gene. This experiment shows that the use of specific beacons (approach 2) significantly reduces the percentage of MSSA that are non specifically detected with the generic beacon approach (approach 1).

Table 3: max signal ratio obtained for MSSA and MRSA with specific beacons NASBA (positive signals are in bold italics)

Strain	<i>mecA</i> PCR	Max signal ratio
neg control	Not applicable	1.0
MSSA 10	neg	1.0
MSSA 11	neg	1.0
MSSA 12	neg	1.0
MSSA 13	neg	1.0
MSSA 14	neg	1.0
MSSA 15	neg	1.0
MSSA 16	neg	1.7
MSSA 17	neg	1.0
MSSA 18	neg	1.0
MSSA 19	neg	1.0
MSSA 23	neg	1.0
MSSA 24	neg	1.0
MSSA 25	neg	1.0
MSSA 26	neg	1.0
MSSA 27	neg	1.0
MSSA 28	neg	1.0
MSSA 29	neg	1.0
MSSA 30	neg	1.0
MRSA 3	pos	1.6
MRSA 7	pos	1.6
MRSA 8	pos	1.7
MRSA 9	pos	2.2
MRSA 10	pos	2.1
MRSA 11	pos	1.9

EXAMPLE 3: Multiplex amplification and detection of *mecA* gene and cassette insertion region

[0069] Using approach 2, some false positives are eliminated, thus providing an improved method for detecting MRSA. However, MSSA strains possessing the cassette without the *mecA* gene can be detected, and a further improvement was investigated. As shown in this example, simultaneous amplification and detection of both the insertion

cassette region, using approach 2, and the *mecA* gene (see FIG. 4) can help reduce detection of such strains (false MRSA positive).

[0070] This example shows the feasibility of a multiplex NASBA for detection of both the *mecA* gene and the cassette junction region (approach 2, which has SCC*mec* cassette-specific beacons) in the same tube. This NASBA makes use of 5 SCC*mec* cassette-specific P2 (SEQ ID NOs: 1-5), 1 P1 (SEQ ID NO: 13) and 5 FAM-labeled SCC*mec* cassette-beacons (SEQ ID NOs: 7-11) for the cassette junction region and 1 P1 (SEQ ID NO: 15), 1 P2 (SEQ ID NO: 16) and 1 ROX-labeled beacon (SEQ ID NO: 14) for *mecA*. A NASBA reaction targeting only the SCC*mec* right extremity junction region, amplified in one tube (five SCC*mec* cassette-specific P2 (SEQ ID NOs: 1-5), five specific SCC*mec* cassette-specific FAM beacons (SEQ ID NOs: 7-11) and one *orfX* P1 (SEQ ID NO: 13)), is also performed for comparison.

[0071] The following table (Table 4) gives max signal ratios obtained for (1) SCC*mec* right extremity junction NASBA (FAM signal) when only the SCC*mec* right extremity junction region is amplified in one tube (five P2, five specific FAM beacons and one P1 ("SCC*mec* junction only") and (2) SCC*mec* right extremity junction NASBA (FAM signal) and *mecA* NASBA (ROX signal) when both NASBA were performed in the same tube ("SCC*mec* junction and *mecA* NASBA in one tube") (max signal ratio are considered positive when > 1.2).

Table 4

	type i			type ii			type iii		
	SCC <i>mec</i> junction only	SCC <i>mec</i> junction and <i>mecA</i> NASBA in one tube		SCC <i>mec</i> junction only	SCC <i>mec</i> junction and <i>mecA</i> NASBA in one tube		SCC <i>mec</i> junction only	SCC <i>mec</i> junction and <i>mecA</i> NASBA in one tube	
CFU/NASBA	FAM	FAM	ROX	FAM	FAM	ROX	FAM	FAM	ROX
1000	1.90	1.84	4.75	2.02	2.07	4.86	1.50	1.39	4.73
100	1.85	1.84	4.75	2.03	1.97	4.74	1.48	1.39	4.53
10	1.87	1.84	4.72	1.88	1.88	4.63	1.49	1.36	4.60
5	1.85	1.85	4.80	1.01	1.69	4.58	1.01	1.40	4.41
	type iv			type v			type vii		
	SCC <i>mec</i> junction only	SCC <i>mec</i> junction and <i>mecA</i> NASBA in one tube		SCC <i>mec</i> junction only	SCC <i>mec</i> junction and <i>mecA</i> NASBA in one tube		SCC <i>mec</i> junction only	SCC <i>mec</i> junction and <i>mecA</i> NASBA in one tube	
CFU/NASBA	FAM	FAM	ROX	FAM	FAM	ROX	FAM	FAM	ROX

1000	<i>1.62</i>	<i>1.64</i>	<i>4.85</i>	<i>1.14</i>	<i>1.39</i>	<i>4.81</i>	<i>2.41</i>	<i>2.43</i>	<i>4.83</i>
100	<i>1.57</i>	<i>1.48</i>	<i>4.58</i>	<i>1.42</i>	<i>1.33</i>	<i>4.69</i>	<i>2.44</i>	<i>2.43</i>	<i>4.81</i>
10	<i>1.61</i>	<i>1.50</i>	<i>4.62</i>	<i>1.43</i>	1.00	<i>4.03</i>	<i>2.40</i>	1.00	<i>4.59</i>
5	<i>1.61</i>	<i>1.44</i>	<i>4.15</i>	<i>1.43</i>	1.01	1.00	<i>2.40</i>	1.01	1.01

Numbers in *bold italics* in Table 4 indicate positive signals.

[0072] These results show that when SCCmec right extremity junction and *mecA* NASBA are performed in the same tube, the limit of detection is as low as 5 CFU/NASBA for both SCCmec right extremity junction region and *mecA* gene with all strain types except types v and vii. Thus, in the case of an MSSA possessing the insertion cassette region without the *mecA* gene, the present test will properly provide a "MRSA negative" result (SCCmec junction (+) plus *mecA* (-).

EXAMPLE 4: Multiplex amplification and detection of *mecA* gene and cassette insertion region

[0073] A multiplex (same reaction tube) amplification and detection of MRSA was performed, utilizing approach 2 (SCCmec cassette-specific beacons) along with *mecA* detection (same conditions as described in Example 3), utilizing six SCCmec cassette-specific P2 (SEQ ID NOs: 1-6), six specific SCCmec cassette-specific FAM beacons (SEQ ID NOs: 7-12) and one *orfX* P1 (SEQ ID NO: 13). Positive detection of the MRSA strains was obtained.

[0075] It is understood that the disclosed invention is not limited to the particular methodology, protocols, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

[0076] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 77203-162 Seq 08-JUL-10 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110>	bioMerieux SA Jay, Corrine van Strijp, Dianne van de Wiel, Paul Deiman, Birgit	
<120>	DETECTION OF METHICILLIN-RESISTANT STAPHYLOCOCCUS AUREUS	
<130>	9250-174WO	
<150>	US 61/008,680	
<151>	2007-12-21	
<160>	18	
<170>	PatentIn version 3.3	
<210>	1	
<211>	29	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	NABSA primer	
<400>	1	
	ctctgcttta tattataaaa ttacggctg	29
<210>	2	
<211>	29	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	NABSA primer	
<400>	2	
	atttcatata tgtaattcct ccacatctc	29
<210>	3	
<211>	17	
<212>	DNA	
<213>	Artificial Sequence	

<220>
 <223> NABSA primer

<400> 3
 aagactgcgg aggctaa 17

<210> 4
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NABSA primer

<400> 4
 tattcttcaa agatttgagc 20

<210> 5
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NABSA primer

<400> 5
 caaatattat ctcgtaattt ac 22

<210> 6
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NABSA primer

<400> 6
 tctaatttat ttaacataaa atcaatcct 29

<210> 7
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (40)..(40)
 <223> 5' Dabsyl modification

<400> 7
 cggcgcgtca aaaatcatga acctcattac ttatgcgccg 40

<210> 8
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (40)..(40)
 <223> 5' Dabsyl modification

<400> 8
 cgagcgcaaa ttatacacia cctaattttt agtgcgctcg

40

<210> 9
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (40)..(40)
 <223> 5' Dabsyl modification

<400> 9
 cggagctaatt ttaataattt tctcatattt tttagctccg

40

<210> 10
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (36)..(36)
 <223> 5' Dabsyl modification

<400> 10
 cgtaacggat aaaaaaccgc atcatttgac gttacg

36

<210> 11
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (36)..(36)
 <223> 5' Dabsyl modification

<400> 11
 cgtagcggct gaaataaccg catcatttac gctacg

36

<210> 12
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (40)..(40)
 <223> 5' Dabsyl modification

<400> 12
 cgtcacgtaa aatatattat acacaatccg ttctgtgacg

40

<210> 13
 <211> 49
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NABSA primer

<400> 13
 aattctaata cgactcacta tagggagagt caaacggcct gcacaagga

49

<210> 14
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' Cy5 modification

<220>
 <221> misc_feature
 <222> (31)..(31)
 <223> 5' Dabsyl modification

<400> 14
 cgtacgggat catagcgtca ttattcgtac g 31

<210> 15
 <211> 54
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NABSA promoter

<400> 15
 aattctaata cgactcacta tagggagagg tattggccaa ttccacattg tttc 54

<210> 16
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NABSA primer

<400> 16
 cattgatcgc aacgttca 18

<210> 17
 <211> 25
 <212> DNA
 <213> Bacteriophage T7

<400> 17
 aattctaata cgactcacta taggg 25

<210> 18
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide probe sequence

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> 3' FAM modification

<220>
 <221> misc_feature
 <222> (32)..(32)
 <223> 5' Dabsyl modification

<400> 18

cgtacggtag ttactgcggt gtaagacgta cg

32

77203-162

CLAIMS:

1. A method of detecting in a sample a methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCCmec cassette within *Staphylococcus aureus* chromosomal DNA, comprising:
 - 5 performing on the sample an amplification and detection reaction utilizing:
 - a) a first primer capable of specifically hybridizing in an extremity junction region of the SCCmec cassette,
 - b) a second primer capable of specifically hybridizing in an extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
 - 10 c) a probe capable of specifically hybridizing to a region of the SCCmec cassette between the region with which the first primer is capable of hybridizing and the junction,
 - wherein each of the first primer and the second primer is oriented such that, under amplification conditions, the junction is amplified, and
 - 15 wherein if the sample contains MRSA, hybridization of the probe is detected.
2. The method of claim 1, wherein the probe comprises more than one probe, each capable of hybridizing specifically to a different type of MRSA.
3. The method of claim 1, wherein the probe comprises five or more probes, each capable of hybridizing specifically to a different type of MRSA.
- 20 4. The method of claim 1, wherein the amplification reaction is selected from the group consisting of transcription-based amplification and PCR.
5. The method of claim 4, wherein the transcription-based amplification is NASBA.

77203-162

6. The method of claim 5, wherein the NASBA is DNA NASBA.
7. The method of claim 1, wherein the first primer comprises more than one primer capable of hybridizing in an extremity junction region of the SCC mec cassette.
8. The method of claim 1, wherein the extremity junction region is the right
5 extremity junction region.
9. The method of claim 8, wherein the second primer comprises the nucleic acid set forth as SEQ ID NO: 13.
10. The method of claim 8, wherein the probe comprises a sequence selected from the group consisting of SEQ ID NO: 7, 8, 9, 10, 11, and 12.
- 10 11. The method of claim 8, wherein the probe comprises at least five nucleic acids, each comprising one of SEQ ID NO: 7, 8, 9, 10, 11, and 12.
12. The method of claim 8, wherein the first primer comprises a nucleic acid comprising a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, and 6.
13. The method of claim 8, wherein the first primer comprises at least five nucleic
15 acids, each comprising one of SEQ ID NO: 1, 2, 3, 4, 5, and 6.
14. A kit for detection of methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC mec cassette within *Staphylococcus aureus* chromosomal DNA, comprising:
 - a) a first primer capable of specifically hybridizing in a right extremity junction
20 region of the SCC mec cassette,
 - b) a second primer capable of specifically hybridizing in a right extremity junction region of chromosomal *Staphylococcus aureus* DNA, and
 - c) a probe selected from the group consisting of (1) a probe capable of specifically hybridizing primarily within a region of the SCC mec cassette between the region

77203-162

with which the first primer is capable of hybridizing and the junction, and (2) a probe capable of specifically hybridizing fully within a region of the SCC*mec* cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that,
5 under amplification conditions, the junction is amplified.

15. The kit of claim 14, wherein the first primer is a nucleic acid comprising a sequence selected from the group consisting of SEQ ID NO: 1, 2, 3, 4, 5, and 6.

16. The kit of claim 14, wherein the second primer is capable of specifically hybridizing to *orfX*.

10 17. The kit of claim 14, wherein the second primer comprises the nucleic acid set forth as SEQ ID NO: 13.

18. The kit of claim 14, wherein the probe comprises more than one probe, each capable of hybridizing specifically to a different type of MRSA.

15 19. The kit of claim 14, wherein the probe is a nucleic acid comprising a sequence selected from the group consisting of SEQ ID NO: 7, 8, 9, 10, 11, and 12.

20. A kit for detection of methicillin-resistant *Staphylococcus aureus* (MRSA) having an insertion of an SCC*mec* cassette within *Staphylococcus aureus* chromosomal DNA, comprising:

20 (a) a first primer capable of specifically hybridizing in an extremity junction region of the SCC*mec* cassette,

(b) a second primer capable of specifically hybridizing to chromosomal *Staphylococcus aureus* DNA in the region of the extremity junction, and

(c) a probe selected from the group consisting of (1) a first probe capable of specifically hybridizing primarily within a region of the SCC*mec* cassette between the region

77203-162

with which the first primer is capable of hybridizing and the junction, and (2) a first probe capable of specifically hybridizing fully within a region of the SCC mec cassette between the region with which the first primer is capable of hybridizing and the junction,

wherein each of the first primer and the second primer is oriented such that,
 5 under amplification conditions, the junction is amplified.


21. An oligonucleotide composition comprising:

(1) a first oligonucleotide having a nucleic acid sequence that specifically hybridizes to an extremity junction region of a SCC mec cassette;

(2) a second oligonucleotide having a nucleic acid sequence that specifically
 10 hybridizes to a *Staphylococcus aureus* chromosomal DNA region flanking said SCC mec cassette; and

(3) a first probe capable of specifically hybridizing to a region of the SCC mec cassette between the region with which the first primer is capable of hybridizing and the junction.

Fig. 1



2/4




Figure 2 : NASBA approach 1

Fig 2

Fig 3




Figure 3 NASBA approach 2

Fig. 4.

