

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2017/089366 A1

(43) International Publication Date
1 June 2017 (01.06.2017)

- (51) **International Patent Classification:**
CI2N 9/54 (2006.01) *CUD 3/386* (2006.01)
- (21) **International Application Number:**
PCT/EP20 16/078492
- (22) **International Filing Date:**
23 November 2016 (23.11.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
15 195967.3 24 November 2015 (24.11.2015) EP
- (71) **Applicant:** NOVOZYMES A/S [DK/DK]; Krogshoejvej 36, 2880 Bagsvaerd (DK).
- (72) **Inventors:** BENIE, Astrid; Krogshoejvej 36, 2880 Bagsvaerd (DK). GJERMANSEN, Morten; Krogshoejvej 36, 2880 Bagsvaerd (DK). OESTERGAARD, Peter, Rahbek; Krogshoejvej 36, 2880 Bagsvaerd (DK).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2017/089366 A1

(54) **Title:** POLYPEPTIDES HAVING PROTEASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME

(57) **Abstract:** The present invention relates to isolated polypeptides having protease activity, and polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

POLYPEPTIDES HAVING PROTEASE ACTIVITY AND POLYNUCLEOTIDES ENCODING SAME

REFERENCE TO A SEQUENCE LISTING

This application contains a Sequence Listing in computer readable form, which is
5 incorporated herein by reference.

BACKGROUND OF THE INVENTION

The present invention relates to isolated polypeptides having protease activity and isolated
nucleic acid sequences encoding the proteases. The invention also relates to nucleic acid
10 constructs, vectors, and host cells, including plant and animal cells, comprising the nucleic acid
sequences, as well as methods for producing and using the proteases, in particular the use of the
proteases in detergents.

DESCRIPTION OF THE RELATED ART

15 The present invention provides polypeptides having protease activity and polynucleotides
encoding the polypeptides.

The detergent industry has for many years implemented different enzymes in detergent
formulations, most commonly used enzymes includes proteases, amylases and lipases each
adapted for removing various types of stains. In addition to the enzymes detergent compositions
20 typically include a complex combination of ingredients. For example, most cleaning products
include surfactant system, bleaching agents or builders. Despite the complexity of current
detergents, there remains a need for developing new detergent compositions comprising new
enzymes and/or enzyme blends.

Traditionally laundering has been done at elevated temperatures and well known detergents
25 have been selected to perform at higher temperatures.

The increased focus on improving the washing processes in order to make them more
environmental friendly has resulted in a global tendency to lowering wash time, pH and temperature
and decreasing the amount of detergent components which may influence the environment
negatively. There is therefore a desire e.g. to launder at lower temperature and therefore a need
30 for detergent proteases having high performance at various conditions such as at low temperature.

A protease from *Bacillus lehensis* (Joshi et al., *Bioresour. Technol.* 2013 131 :76-85,
UNIPROT: I3QII3) is 81.9% identical to the protease of the present invention.

SUMMARY OF THE INVENTION

35 The present invention relates to an isolated polypeptide having protease activity, selected

from the group consisting of:

(a) a polypeptide having at least 78%, at least 79%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84% at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 2;

(b) a polypeptide encoded by a polynucleotide having at least 78%, at least 79%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84% at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1;

(c) a variant of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more (e.g. several) positions; and

(d) a fragment of the polypeptide of (a), (b) or (c) that has protease activity.

The present invention also relates to isolated polynucleotides encoding the polypeptides of the present invention; nucleic acid constructs; recombinant expression vectors; recombinant host cells comprising the polynucleotides; and methods of producing the polypeptides.

The present invention also relates to the use of the proteases of the invention in a cleaning process, and methods of producing detergent compositions.

The present invention also relates to a polynucleotide encoding a signal peptide comprising or consisting of amino acids -111 to -85 of SEQ ID NO: 2, a polynucleotide encoding a propeptide comprising or consisting of amino acids -84 to -1 of SEQ ID NO: 2, or a polynucleotide encoding a signal peptide and a propeptide comprising or consisting of amino acids -111 to -1 of SEQ ID NO: 2, each of which is operably linked to a gene encoding a protein; nucleic acid constructs, expression vectors, and recombinant host cells comprising the polynucleotides; and methods of producing a protein.

OVERVIEW OF SEQUENCE LISTING

SEQ ID NO: 1 is the DNA sequence of S8 protease from *Bacillus* sp-13231 .

SEQ ID NO: 2 is the amino acid sequence encoded by SEQ ID NO: 1.

SEQ ID NO: 3 is a *Bacillus clausii* secretion signal.

SEQ ID: NO: 4 is the mature amino acid sequence of the commercial protease, Savinase.

DEFINITIONS

Polypeptides having protease activity, or proteases, are sometimes also designated

peptidases, proteinases, peptide hydrolases, or proteolytic enzymes. Proteases may be of the exo-type that hydrolyses peptides starting at either end thereof, or of the endo-type that act internally in polypeptide chains (endopeptidases). Endopeptidases show activity on N- and C-terminally blocked peptide substrates that are relevant for the specificity of the protease in question.

5 The term "protease" is defined herein as an enzyme that hydrolyses peptide bonds. This definition of protease also applies to the protease-part of the terms "parent protease" and "protease variant," as used herein. The term "protease" includes any enzyme belonging to the EC 3.4 enzyme group (including each of the thirteen subclasses thereof). The EC number refers to Enzyme Nomenclature 1992 from NC-IUBMB, Academic Press, San Diego, California, including
10 supplements 1-5 published in Eur. J. Bio-chem. 1994, 223, 1-5; Eur. J. Biochem. 1995, 232, 1-6; Eur. J. Biochem. 1996, 237, 1-5; Eur. J. Biochem. 1997, 250, 1-6; and Eur. J. Biochem. 1999, 264, 610-650; respectively. The nomenclature is regularly supplemented and updated; see e.g. the World Wide Web (WWW) at <http://www.chem.qmw.ac.uk/iubmb/enzyme/index.html>.

The proteases of the invention are:

- 15 (a) proteases belonging to the EC 3.4.21 . enzyme group; and/or
 (b) Serine proteases of the peptidase family S8;

as described in Biochem.J. 290:205-218 (1993) and in MEROPS protease database, release, 9.4 (www.merops.ac.uk). The database is described in Rawlings, N.D., Barrett, A.J. & Bateman, A. (2010) MEROPS: the peptidase database. Nucleic Acids Res 38, D227-D233.

20 For determining whether a given protease is a Serine protease, and a family S8 protease, reference is made to the above Handbook and the principles indicated therein. Such determination can be carried out for all types of proteases, be it naturally occurring or wild-type proteases; or genetically engineered or synthetic proteases.

The proteases of family S8 contain the catalytic triad in the order Asp, His, Ser. Mutation of
25 any of the amino acids of the catalytic triad will result in change or loss of enzyme activity. The amino acids of the catalytic triad of the S8 protease 1 from *Bacillus* sp-1 3231 (SEQ ID NO: 2) are probably positions Asp32, His62 and Ser215.

Protease activity can be measured using any assay, in which a substrate is employed, that includes peptide bonds relevant for the specificity of the protease in question. Assay-pH and assay-
30 temperature are likewise to be adapted to the protease in question. Examples of assay-pH-values are pH 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12. Examples of assay-temperatures are 15, 20, 25, 30, 35, 37, 40, 45, 50, 55, 60, 65, 70, 80, 90, or 95°C. Examples of protease substrates are casein, such as Azurine-Crosslinked Casein (AZCL-casein), or suc-AAPF-pNA, Examples of suitable protease assays are described in the experimental part.

35 The term "protease activity" means a proteolytic activity (EC 3.4.21 .) that catalyzes the

hydrolysis of amide bond or a protein by hydrolysis of the peptide bond that link amino acids together in a polypeptide chain. Several assays for determining protease activity are available in the art. For purposes of the present invention, protease activity may be determined using Protazyme AK tablet (cross-linked and dyed casein; from Megazyme) as described in the Examples of the present application. The polypeptides of the present invention have at least 20%, e.g., at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 100% of the protease activity of the mature polypeptide of SEQ ID NO: 2.

The term "allelic variant" means any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

The term "catalytic domain" means the region of an enzyme containing the catalytic machinery of the enzyme.

The term "cDNA" means a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic or prokaryotic cell. cDNA lacks intron sequences that may be present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps, including splicing, before appearing as mature spliced mRNA.

The term "cleaning compositions" and "cleaning formulations," refer to compositions that find use in the removal of undesired compounds from items to be cleaned, such as fabric, carpets, dishware including glassware, contact lenses, hard surfaces such as tiles, zincs, floors, and table surfaces, hair (shampoos), skin (soaps and creams), teeth (mouthwashes, toothpastes), etc. The terms encompasses any materials/compounds selected for the particular type of cleaning composition desired and the form of the product (e.g., liquid, gel, granule, or spray compositions), as long as the composition is compatible with the protease according to the invention and other enzyme(s) used in the composition. The specific selection of cleaning composition materials is readily made by considering the surface, item or fabric to be cleaned, and the desired form of the composition for the cleaning conditions during use. These terms further refer to any composition that is suited for cleaning, bleaching, disinfecting, and/or sterilizing any object and/or surface. It is intended that the terms include, but are not limited to detergent composition (e.g., liquid and/or solid laundry detergents and fine fabric detergents; hard surface cleaning formulations, such as for glass, wood, ceramic and metal counter tops and windows; carpet cleaners; oven cleaners; fabric fresheners; fabric softeners; and textile and laundry pre-spotters, as well as dish detergents).

The term "coding sequence" means a polynucleotide, which directly specifies the amino

acid sequence of a polypeptide. The boundaries of the coding sequence are generally determined by an open reading frame, which begins with a start codon such as ATG, GTG, or TTG and ends with a stop codon such as TAA, TAG, or TGA. The coding sequence may be a genomic DNA, cDNA, synthetic DNA, or a combination thereof.

5 The term "control sequences" means nucleic acid sequences necessary for expression of a polynucleotide encoding a mature polypeptide of the present invention. Each control sequence may be native (*i.e.*, from the same gene) or foreign (*i.e.*, from a different gene) to the polynucleotide encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal
10 peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the polynucleotide encoding a polypeptide.

 The term "detergent composition", includes unless otherwise indicated, granular or powder-
15 form all-purpose or heavy-duty washing agents, especially cleaning detergents; liquid, gel or paste-form all-purpose washing agents, especially the so-called heavy-duty liquid (HDL) types; liquid fine-fabric detergents; hand dishwashing agents or light duty dishwashing agents, especially those of the high-foaming type; machine dishwashing agents, including the various tablet, granular, liquid and rinse-aid types for household and institutional use; liquid cleaning and disinfecting agents,
20 including antibacterial hand-wash types, cleaning bars, mouthwashes, denture cleaners, car or carpet shampoos, bathroom cleaners; hair shampoos and hair-rinses; shower gels, foam baths; metal cleaners; as well as cleaning auxiliaries such as bleach additives and "stain-stick" or pre-treat types.

 The terms "detergent composition" and "detergent formulation" are used in reference to
25 mixtures which are intended for use in a wash medium for the cleaning of soiled objects. In some embodiments, the term is used in reference to laundering fabrics and/or garments (e.g., "laundry detergents"). In alternative embodiments, the term refers to other detergents, such as those used to clean dishes, cutlery, etc. (e.g., "dishwashing detergents"). It is not intended that the present invention be limited to any particular detergent formulation or composition. It is intended that in
30 addition to the protease according to the invention, the term encompasses detergents that contains, e.g., surfactants, builders, chelators or chelating agents, bleach system or bleach components, polymers, fabric conditioners, foam boosters, suds suppressors, dyes, perfume, tannish inhibitors, optical brighteners, bactericides, fungicides, soil suspending agents, anti corrosion agents, enzyme inhibitors or stabilizers, enzyme activators, transferase(s), hydrolytic enzymes, oxido reductases,
35 bluing agents and fluorescent dyes, antioxidants, polymers and solubilizers.

The term "dish washing composition" refers to all forms of compositions for cleaning hard surfaces. The present invention is not restricted to any particular type of dish wash composition or any particular detergent.

5 The term "enzyme detergency benefit" or "detergency" is defined herein as the advantageous effect an enzyme may add to a detergent compared to the same detergent without the enzyme. Important detergency benefits which can be provided by enzymes are stain removal with no or very little visible soils after washing and or cleaning, prevention or reduction of redeposition of soils released in the washing process an effect that also is termed anti-redeposition, restoring fully or partly the whiteness of textiles, which originally were white but after repeated use
10 and wash have obtained a greyish or yellowish appearance an effect that also is termed whitening. Textile care benefits, which are not directly related to catalytic stain removal or prevention of redeposition of soils, are also important for enzyme detergency benefits. Examples of such textile care benefits are prevention or reduction of dye transfer from one fabric to another fabric or another part of the same fabric an effect that is also termed dye transfer inhibition or anti-backstaining,
15 removal of protruding or broken fibers from a fabric surface to decrease pilling tendencies or remove already existing pills or fuzz an effect that also is termed anti-pilling, improvement of the fabric-softness, colour clarification of the fabric and removal of particulate soils which are trapped in the fibers of the fabric or garment. Enzymatic bleaching is a further enzyme detergency benefit where the catalytic activity generally is used to catalyze the formation of bleaching component such
20 as hydrogen peroxide or other peroxides.

The term "expression" includes any step involved in the production of a polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

25 The term "expression vector" means a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide and is operably linked to control sequences that provide for its expression.

The term "fabric" encompasses any textile material. Thus, it is intended that the term encompass garments, as well as fabrics, yarns, fibers, non-woven materials, natural materials, synthetic materials, and any other textile material.

30 The term "fragment" means a polypeptide having one or more (e.g., several) amino acids absent from the amino and/or carboxyl terminus of a mature polypeptide or domain; wherein the fragment has protease activity.

The term "Hard surface cleaning" is defined herein as cleaning of hard surfaces wherein hard surfaces may include floors, tables, walls, roofs etc. as well as surfaces of hard objects such
35 as cars (car wash) and dishes (dish wash). Dish washing includes but are not limited to cleaning of

plates, cups, glasses, bowls, and cutlery such as spoons, knives, forks, serving utensils, ceramics, plastics, metals, china, glass and acrylics.

The term "host cell" means any cell type that is susceptible to transformation, transfection, transduction, or the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication.

The term "improved wash performance" is defined herein as an enzyme such as a protease (also a blend of enzymes, not necessarily only variants but also backbones, and in combination with certain cleaning composition etc.) displaying an alteration of the wash performance of e.g. said protease relative to the wash performance of the another protease or parent protease e.g. by increased stain removal. The term "wash performance" includes wash performance in laundry but also e.g. in dish wash.

The term "isolated" means a substance in a form or environment that does not occur in nature. Non-limiting examples of isolated substances include (1) any non-naturally occurring substance, (2) any substance including, but not limited to, any enzyme, variant, nucleic acid, protein, peptide or cofactor, that is at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature; (3) any substance modified by the hand of man relative to that substance found in nature; or (4) any substance modified by increasing the amount of the substance relative to other components with which it is naturally associated (e.g., multiple copies of a gene encoding the substance; use of a stronger promoter than the promoter naturally associated with the gene encoding the substance). An isolated substance may be present in a fermentation broth sample.

The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. In one aspect, the mature polypeptide is amino acids 1 to 269 of SEQ ID NO: 2 based on amino acid sequencing using Edman degradation chemistry. It is known in the art that a host cell may produce a mixture of two or more different mature polypeptides (*i.e.*, with a different C-terminal and/or N-terminal amino acid) expressed by the same polynucleotide.

The term "mature polypeptide coding sequence" means a polynucleotide that encodes a mature polypeptide having protease activity. In one aspect, the mature polypeptide coding sequence is nucleotides 334-1 140 of SEQ ID NO: 1.

The term "nucleic acid construct" means a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic, which comprises one or more control sequences.

The term "operably linked" means a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of a polynucleotide such that the control sequence directs expression of the coding sequence.

5 The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity".

For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *Trends Genet.* 10 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

(Identical Residues x 100)/(Length of Alignment - Total Number of Gaps in Alignment)

15 For purposes of the present invention, the sequence identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *supra*), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap 20 extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

(Identical Deoxyribonucleotides x 100)/(Length of Alignment - Total Number of Gaps in Alignment)

Stringency conditions: The term "very low stringency conditions" means for probes of at 25 least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 25% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 45°C.

The term "low stringency conditions" means for probes of at least 100 nucleotides in length, 30 prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 25% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 50°C.

The term "medium stringency conditions" means for probes of at least 100 nucleotides in 35 length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml

sheared and denatured salmon sperm DNA, and 35% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 55°C.

5 The term "medium-high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 35% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 60°C.

10 The term "high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 50% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 65°C.

15 The term "very high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 50% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 70°C.

20 The term "subsequence" means a polynucleotide having one or more (e.g., several) nucleotides absent from the 5' and/or 3' end of a mature polypeptide coding sequence; wherein the subsequence encodes a fragment having protease activity.

25 The term "substantially pure polynucleotide" means a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered polypeptide production systems. Thus, a substantially pure polynucleotide contains at most 10%, at most 8%, at most 6%, at most 5%, at most 4%, at most 3%, at most 2%, at most 1%, and at most 0.5% by weight of other polynucleotide material with which it is natively or recombinantly associated. A substantially pure polynucleotide may, however, include naturally occurring 5' and 3' untranslated regions, such as promoters and terminators. Preferably, the polynucleotide is at least 90% pure, e.g., at least 92% pure, at least 94% pure, at least 95% pure, 30 at least 96% pure, at least 97% pure, at least 98% pure, at least 99% pure, and at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form.

35 The term "substantially pure polypeptide" means a preparation that contains at most 10%, at most 8%, at most 6%, at most 5%, at most 4%, at most 3%, at most 2%, at most 1%, and at most 0.5% by weight of other polypeptide material with which it is natively or recombinantly associated. Preferably, the polypeptide is at least 92% pure, e.g., at least 94% pure, at least 95% pure, at least

96% pure, at least 97% pure, at least 98% pure, at least 99%, at least 99.5% pure, and 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides of the present invention are preferably in a substantially pure form. This can be accomplished, for example, by preparing the polypeptide by well known recombinant methods or by classical purification methods.

The term "textile" means any textile material including yarns, yarn intermediates, fibers, non-woven materials, natural materials, synthetic materials, and any other textile material, fabrics made of these materials and products made from fabrics (e.g., garments and other articles). The textile or fabric may be in the form of knits, wovens, denims, non-wovens, felts, yarns, and towelling. The textile may be cellulose based such as natural cellulose, including cotton, flax/linen, jute, ramie, sisal or coir or manmade cellulose (e.g. originating from wood pulp) including viscose/rayon, ramie, cellulose acetate fibers (tricell), lyocell or blends thereof. The textile or fabric may also be non-cellulose based such as natural polyamides including wool, camel, cashmere, mohair, rabbit and silk or synthetic polymer such as nylon, aramid, polyester, acrylic, polypropylene and spandex/elastane, or blends thereof as well as blend of cellulose based and non-cellulose based fibers. Examples of blends are blends of cotton and/or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g. polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers (e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fibers, lyocell). Fabric may be conventional washable laundry, for example stained household laundry. When the term fabric or garment is used it is intended to include the broader term textiles as well.

The term "textile care benefits", which are not directly related to catalytic stain removal or prevention of redeposition of soils, are also important for enzyme detergency benefits. Examples of such textile care benefits are prevention or reduction of dye transfer from one textile to another textile or another part of the same textile an effect that is also termed dye transfer inhibition or anti-backstaining, removal of protruding or broken fibers from a textile surface to decrease pilling tendencies or remove already existing pills or fuzz an effect that also is termed anti-pilling, improvement of the textile-softness, colour clarification of the textile and removal of particulate soils which are trapped in the fibers of the textile. Enzymatic bleaching is a further enzyme detergency benefit where the catalytic activity generally is used to catalyze the formation of bleaching component such as hydrogen peroxide or other peroxides or other bleaching species.

The term "variant" means a polypeptide having protease activity comprising an alteration, *i.e.*, a substitution, insertion, and/or deletion, at one or more (e.g., several) positions. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding one or more

(e.g. several) amino acids, e.g. 1-5 amino acids adjacent to and immediately following the amino acid occupying a position.

The term "wash performance" is used as an enzyme's ability to remove stains present on the object to be cleaned during e.g. wash or hard surface cleaning.

5 The term "whiteness" is defined herein as a broad term with different meanings in different regions and for different customers. Loss of whiteness can e.g. be due to greying, yellowing, or removal of optical brighteners/hueing agents. Greying and yellowing can be due to soil redeposition, body soils, colouring from e.g. iron and copper ions or dye transfer. Whiteness might include one or several issues from the list below: Colorant or dye effects; Incomplete stain removal
10 (e.g. body soils, sebum ect.); Re-deposition (greying, yellowing or other discolorations of the object) (removed soils re-associates with other part of textile, soiled or unsoiled); Chemical changes in textile during application; and Clarification or brightening of colours.

DETAILED DESCRIPTION OF THE INVENTION

15 In an embodiment, the present invention relates to an isolated polypeptide having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 78%, at least 79%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84% at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%,
20 which have protease activity. In one aspect, the polypeptide differ by no more than 20 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 or 19 from the mature polypeptide of SEQ ID NO: 2.

A polypeptide of the present invention preferably comprises or consists of the amino acid sequence of SEQ ID NO: 2 or an allelic variant thereof; or is a fragment thereof having protease
25 activity. In another aspect, the polypeptide comprises or consists of the mature polypeptide of SEQ ID NO: 2. In another aspect, the polypeptide comprises or consists of amino acids 1 to 269 of SEQ ID NO: 2.

In another embodiment, the present invention relates to an isolated polypeptide having protease activity encoded by a polynucleotide that hybridizes under very low stringency conditions,
30 low stringency conditions, medium stringency conditions, medium-high stringency conditions, high stringency conditions, or very high stringency conditions with the mature polypeptide coding sequence of SEQ ID NO: 1, or the full-length complement thereof (Sambrook *et al.*, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York).

The polynucleotide of SEQ ID NO: 1 or a subsequence thereof, as well as the polypeptide
35 of SEQ ID NO: 2 or a fragment thereof may be used to design nucleic acid probes to identify and

clone DNA encoding polypeptides having protease activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic DNA or cDNA of a cell of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes
5 can be considerably shorter than the entire sequence, but should be at least 15, e.g., at least 25, at least 35, or at least 70 nucleotides in length. Preferably, the nucleic acid probe is at least 100 nucleotides in length, e.g., at least 200 nucleotides, at least 300 nucleotides, at least 400 nucleotides, at least 500 nucleotides, at least 600 nucleotides, at least 700 nucleotides, at least 800 nucleotides, or at least 900 nucleotides in length. Both DNA and RNA probes can be used. The
10 probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

A genomic DNA or cDNA library prepared from such other strains may be screened for DNA that hybridizes with the probes described above and encodes a polypeptide having protease activity. Genomic or other DNA from such other strains may be separated by agarose or
15 polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that hybridizes with SEQ ID NO: 1 or a subsequence thereof, the carrier material is used in a Southern blot.

For purposes of the present invention, hybridization indicates that the polynucleotide
20 hybridizes to a labeled nucleic acid probe corresponding to (i) SEQ ID NO: 1; (ii) the mature polypeptide coding sequence of SEQ ID NO: 1; (iii) the full-length complement thereof; or (iv) a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using, for example, X-ray film or any other detection means known in the art.

25 In one aspect, the nucleic acid probe is nucleotides 101 to 1405, nucleotides 188 to 1222, nucleotides 665 to 1222, or nucleotides 800 to 1200 of SEQ ID NO: 1. In another aspect, the nucleic acid probe is a polynucleotide that encodes the polypeptide of SEQ ID NO: 2; the mature polypeptide thereof; or a fragment thereof. In another aspect, the nucleic acid probe is SEQ ID NO:
1.

30 In another embodiment, the present invention relates to an isolated polypeptide having protease activity encoded by a polynucleotide having a sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1 of at least 78%, at least 79%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84% at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at
35 least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%.

In another embodiment, the present invention relates to variants of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more (e.g., several) positions. In an embodiment, the number of amino acid substitutions, deletions and/or insertions introduced into the mature polypeptide of SEQ ID NO: 2 is not more than 20, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19. In another embodiment, the number of positions comprising a substitution, deletion, and/or insertion in the mature polypeptide of SEQ ID NO: 2 is between 1 and 20, such as 1-15, 1-10 or 1-5 positions. In an embodiment, the number of positions comprising a substitution, deletion, and/or insertion in the mature polypeptide of SEQ ID NO: 2 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10. In another embodiment, the number of substitutions, deletions, and/or insertions in the mature polypeptide of SEQ ID NO: 2 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10. In a further embodiment, the number of substitutions in the mature polypeptide of SEQ ID NO: 2 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10. In a further embodiment, the number of conservative substitutions in the mature polypeptide of SEQ ID NO: 2 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10. The amino acid changes may be of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of 1-30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the groups of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. Common substitutions are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, LeuA/al, Ala/Glu, and Asp/Gly.

Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

Essential amino acids in a polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and

Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant variant molecules are tested for protease activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton *et al.*, 1996, *J. Biol. Chem.* 271 : 4699-4708. The active site of the enzyme or other
5 biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Lett.* 309: 59-64. The identity of essential amino acids can also be
10 inferred from an alignment with a related polypeptide.

Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241 : 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156;
15 WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman *et al.*, 1991, *Biochemistry* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire *et al.*, 1986, *Gene* 46: 145; Ner *et al.*, 1988, *DNA* 7: 127).

Mutagenesis/shuffling methods can be combined with high-throughput, automated
20 screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness *et al.*, 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide.

25 The polypeptide may be a hybrid polypeptide in which a region of one polypeptide is fused at the N-terminus or the C-terminus of a region of another polypeptide.

The polypeptide may be a fusion polypeptide or cleavable fusion polypeptide in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide of the present invention. A fusion polypeptide is produced by fusing a polynucleotide encoding another
30 polypeptide to a polynucleotide of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fusion polypeptide is under control of the same promoter(s) and terminator. Fusion polypeptides may also be constructed using intein technology in which fusion polypeptides are created post-translationally (Cooper *et al.*, 1993,
35 *EMBO J.* 12: 2575-2583; Dawson *et al.*, 1994, *Science* 266: 776-779).

A fusion polypeptide can further comprise a cleavage site between the two polypeptides. Upon secretion of the fusion protein, the site is cleaved releasing the two polypeptides. Examples of cleavage sites include, but are not limited to, the sites disclosed in Martin *et al.*, 2003, *J. Ind. Microbiol. Biotechnol.* 3: 568-576; Svetina *et al.*, 2000, *J. Biotechnol.* 76: 245-251 ; Rasmussen-
5 Wilson *et al.*, 1997, *Appl. Environ. Microbiol.* 63: 3488-3493; Ward *et al.*, 1995, *Biotechnology* 13: 498-503; and Contreras *et al.*, 1991, *Biotechnology* 9: 378-381 ; Eaton *et al.*, 1986, *Biochemistry* 25: 505-512; Collins-Racie *et al.*, 1995, *Biotechnology* 13: 982-987; Carter *et al.*, 1989, *Proteins: Structure, Function, and Genetics* 6: 240-248; and Stevens, 2003, *Drug Discovery World* 4: 35-48.

10 **Embodiments**

In certain embodiments of the invention, the protease of the invention exhibits beneficial thermal properties such as thermostability, steam stability, pH properties, such as acid stability, etc.

Acidity/alkalinity properties

15 In certain embodiments of the invention, the protease of the invention exhibits beneficial properties in respect of pH, such as acid stability etc. Stability of the protease at a low pH is beneficial since the protease can have activity in the intestine after passing through the stomach. Stability of the protease at a high pH is beneficial for cleaning and washing since detergent compositions often have an alkaline pH. The proteases of the invention exhibit surprising pH
20 properties, especially pH stability properties which makes them interesting candidates for use in detergents.

Wash performance

In certain embodiments of the invention, the protease of the invention exhibits beneficial
25 wash performance. Thus a detergent with the S8 protease 1 from *Bacillus* sp-13231 is more effective at removing stains compared to detergent without any protease.

Sources of polypeptides having protease activity

A polypeptide having protease activity of the present invention may be obtained from
30 microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a polynucleotide is produced by the source or by a strain in which the polynucleotide from the source has been inserted. In one aspect, the polypeptide obtained from a given source is secreted extracellularly.

35 The polypeptide may be a bacterial polypeptide.

In one aspect, the polypeptide is a protease from a bacterium of the class *Actinobacteria*, such as from the order *Actinomycetales*, or from the suborder *Micromonosporineae*, or from the family *Micromonosporineae*, or from the genera *Genus*. In another aspect, the polypeptide is a *Bacillus sp-13231* polypeptide.

5 Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL). The strain used is publically available under the accession number ATCC 31203 or DSM 4391 1.

10 The polypeptide may be identified and obtained from other sources including microorganisms isolated from nature (e.g., soil, composts, water, etc.) or DNA samples obtained directly from natural materials (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms and DNA directly from natural habitats are well known in the art. A polynucleotide encoding the polypeptide may then be obtained by similarly screening a
15 genomic DNA or cDNA library of another microorganism or mixed DNA sample. Once a polynucleotide encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are known to those of ordinary skill in the art (see, e.g., Sambrook *et al.*, 1989, *supra*).

20 **Polynucleotides**

The present invention also relates to isolated polynucleotides encoding a polypeptide of the present invention, as described herein.

The techniques used to isolate or clone a polynucleotide are known in the art and include isolation from genomic DNA or cDNA, or a combination thereof. The cloning of the polynucleotides
25 from genomic DNA can be effected, e.g., by using the well-known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, e.g., Innis *et al.*, 1990, *PCR: A Guide to Methods and Application*, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligation activated transcription (LAT) and polynucleotide-based amplification (NASBA) may be used. The
30 polynucleotides may be cloned from a strain of *Dactylosporangium*, or a related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the polynucleotide.

Modification of a polynucleotide encoding a polypeptide of the present invention may be necessary for synthesizing polypeptides substantially similar to the polypeptide. The term
35 "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide.

These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, e.g., variants that differ in specific activity, thermostability, pH optimum, or the like. The variants may be constructed on the basis of the polynucleotide presented as the mature polypeptide coding sequence of SEQ ID NO: 1, e.g., a subsequence thereof, and/or by introduction
5 of nucleotide substitutions that do not result in a change in the amino acid sequence of the polypeptide, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford *et al.*, 1991, *Protein Expression and Purification* 2: 95-107.

10

Nucleic acid constructs

The present invention also relates to nucleic acid constructs comprising a polynucleotide of the present invention operably linked to one or more control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

15

A polynucleotide may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides utilizing recombinant DNA methods are well known in the art.

20

The control sequence may be a promoter, a polynucleotide that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any polynucleotide that shows transcriptional activity in the host cell including variant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

25

Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a bacterial host cell are the promoters obtained from the *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus subtilis* *xylA* and *xylB* genes,
30 *Bacillus thuringiensis cryIIIA* gene (Agaisse and Lereclus, 1994, *Molecular Microbiology* 13: 97-107), *E. coli lac* operon, *E. coli trc* promoter (Egon *et al.*, 1988, *Gene* 69: 301-315), *Streptomyces coelicolor* agarase gene (*dagA*), and prokaryotic beta-lactamase gene (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. USA* 75: 3727-3731), as well as the *tac* promoter (DeBoer *et al.*, 1983, *Proc. Natl. Acad. Sci. USA* 80: 21-25). Further promoters are described in "Useful proteins from
35 recombinant bacteria" in Gilbert *et al.*, 1980, *Scientific American* 242: 74-94; and in Sambrook *et*

a., 1989, *supra*. Examples of tandem promoters are disclosed in WO 99/43835.

Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus nidulans* acetamidase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Aspergillus oryzae* TAKA amylase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Rhizomucor miehei* lipase, *Rhizomucor miehei* aspartic proteinase, *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a modified promoter from an *Aspergillus* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus* triose phosphate isomerase gene; non-limiting examples include modified promoters from an *Aspergillus niger* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus nidulans* or *Aspergillus oryzae* triose phosphate isomerase gene); and variant, truncated, and hybrid promoters thereof.

In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos *et al.*, 1992, *Yeast* 8: 423-488.

The control sequence may also be a transcription terminator, which is recognized by a host cell to terminate transcription. The terminator is operably linked to the 3'-terminus of the polynucleotide encoding the polypeptide. Any terminator that is functional in the host cell may be used in the present invention.

Preferred terminators for bacterial host cells are obtained from the genes for *Bacillus clausii* alkaline protease (*aprH*), *Bacillus licheniformis* alpha-amylase (*amyL*), and *Escherichia coli* ribosomal RNA (*rrnB*).

Preferred terminators for filamentous fungal host cells are obtained from the genes for

Aspergillus nidulans anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*.

The control sequence may also be an mRNA stabilizer region downstream of a promoter and upstream of the coding sequence of a gene which increases expression of the gene.

Examples of suitable mRNA stabilizer regions are obtained from a *Bacillus thuringiensis cryIIIA* gene (WO 94/25612) and a *Bacillus subtilis* SP82 gene (Hue *et al.*, 1995, *Journal of Bacteriology* 177: 3465-3471).

The control sequence may also be a leader, a nontranslated region of an mRNA that is important for translation by the host cell. The leader is operably linked to the 5'-terminus of the polynucleotide encoding the polypeptide. Any leader that is functional in the host cell may be used.

Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

The control sequence may also be a polyadenylation sequence; a sequence operably linked to the 3'-terminus of the polynucleotide and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell may be used.

Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, *Mol. Cellular Biol.* 15: 5983-5990.

The control sequence may also be a signal peptide coding region that encodes a signal peptide linked to the N-terminus of a polypeptide and directs the polypeptide into the cell's secretory pathway. The 5'-end of the coding sequence of the polynucleotide may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the polypeptide. Alternatively, the 5'-end of the coding sequence

may contain a signal peptide coding sequence that is foreign to the coding sequence. A foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, a foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell may be used.

Effective signal peptide coding sequences for bacterial host cells are the signal peptide coding sequences obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus stearothermophilus* neutral proteases {*nprT*, *nprS*, *nprM*}, and *Bacillus subtilis* *prsA*. Further signal peptides are described by Simonen and Palva, 1993, *Microbiological Reviews* 57: 109-137.

Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Aspergillus oryzae* TAKA amylase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, *Humicola lanuginosa* lipase, and *Rhizomucor miehei* aspartic proteinase.

Useful signal peptides for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding sequences are described by Romanos *et al*, 1992, *supra*.

The control sequence may also be a propeptide coding sequence that encodes a propeptide positioned at the N-terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to an active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding sequence may be obtained from the genes for *Bacillus subtilis* alkaline protease {*aprE*}, *Bacillus subtilis* neutral protease {*nprT*}, *Myceliophthora thermophila* laccase (WO 95/33836), *Rhizomucor miehei* aspartic proteinase, and *Saccharomyces cerevisiae* alpha-factor.

Where both signal peptide and propeptide sequences are present, the propeptide sequence is positioned next to the N-terminus of a polypeptide and the signal peptide sequence is positioned next to the N-terminus of the propeptide sequence.

It may also be desirable to add regulatory sequences that regulate expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems

include the *lac*, *tac*, and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the *Aspergillus niger* glucoamylase promoter, *Aspergillus oryzae* TAKA alpha-amylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the polynucleotide encoding the polypeptide would be operably linked with the regulatory sequence.

10 **Expression vectors**

The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleotide and control sequences may be joined together to produce a recombinant expression vector that may include one or more convenient restriction sites to allow for insertion or substitution of the polynucleotide encoding the polypeptide at such sites. Alternatively, the polynucleotide may be expressed by inserting the polynucleotide or a nucleic acid construct comprising the polynucleotide into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about expression of the polynucleotide. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vector may be a linear or closed circular plasmid.

The vector may be an autonomously replicating vector, *i.e.*, a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

The vector preferably contains one or more selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to

auxotrophs, and the like.

Examples of bacterial selectable markers are *Bacillus licheniformis* or *Bacillus subtilis* *dal* genes, or markers that confer antibiotic resistance such as ampicillin, chloramphenicol, kanamycin, neomycin, spectinomycin, or tetracycline resistance. Suitable markers for yeast host cells include, but are not limited to, ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are *Aspergillus nidulans* or *Aspergillus oryzae* *amdS* and *pyrG* genes and a *Streptomyces hygroscopicus* *bar* gene.

The vector preferably contains an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or non-homologous recombination. Alternatively, the vector may contain additional polynucleotides for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, 400 to 10,000 base pairs, and 800 to 10,000 base pairs, which have a high degree of sequence identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding polynucleotides. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" means a polynucleotide that enables a plasmid or vector to replicate *in vivo*.

Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB1, pE194, pTA1060, and pAM β I permitting replication in *Bacillus*.

Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and

CEN6.

Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANSI (Gems *et al.*, 1991, *Gene* 98: 61-67; Cullen *et al.*, 1987, *Nucleic Acids Res.* 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the
5 gene can be accomplished according to the methods disclosed in WO 00/24883.

More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of a polypeptide. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells
10 containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, e.g.,
15 Sambrook *et al.*, 1989, *supra*).

Host cells

The present invention also relates to recombinant host cells, comprising a polynucleotide of the present invention operably linked to one or more control sequences that direct the production of
20 a polypeptide of the present invention. A construct or vector comprising a polynucleotide is introduced into a host cell so that the construct or vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene
25 encoding the polypeptide and its source.

The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, e.g., a prokaryote or a eukaryote.

The prokaryotic host cell may be any Gram-positive or Gram-negative bacterium. Gram-positive bacteria include, but are not limited to, *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*,
30 *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, and *Streptomyces*. Gram-negative bacteria include, but are not limited to, *Campylobacter*, *E. coli*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, and *Ureaplasma*.

The bacterial host cell may be any *Bacillus* cell including, but not limited to, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus*
35 *coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus*

megaterium, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis* cells.

The bacterial host cell may also be any *Streptococcus* cell including, but not limited to, *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, and *Streptococcus equi* subsp. *Zooepidemicus* cells.

The bacterial host cell may also be any *Streptomyces* cell including, but not limited to, *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, and *Streptomyces lividans* cells.

The introduction of DNA into a *Bacillus* cell may be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Mol. Gen. Genet.* 168: 111-115), competent cell transformation (see, e.g., Young and Spizizen, 1961, *J. Bacteriol.* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *J. Mol. Biol.* 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, *J. Bacteriol.* 169: 5271-5278). The introduction of DNA into an *E. coli* cell may be effected by protoplast transformation (see, e.g., Hanahan, 1983, *J. Mol. Biol.* 166: 557-580) or electroporation (see, e.g., Dower *et al.*, 1988, *Nucleic Acids Res.* 16: 6127-6145). The introduction of DNA into a *Streptomyces* cell may be effected by protoplast transformation, electroporation (see, e.g., Gong *et al.*, 2004, *Folia Microbiol. (Praha)* 49: 399-405), conjugation (see, e.g., Mazodier *et al.*, 1989, *J. Bacteriol.* 171: 3583-3585), or transduction (see, e.g., Burke *et al.*, 2001, *Proc. Natl. Acad. Sci. USA* 98: 6289-6294). The introduction of DNA into a *Pseudomonas* cell may be effected by electroporation (see, e.g., Choi *et al.*, 2006, *J. Microbiol. Methods* 64: 391-397) or conjugation (see, e.g., Pinedo and Smets, 2005, *Appl. Environ. Microbiol.* 71: 51-57). The introduction of DNA into a *Streptococcus* cell may be effected by natural competence (see, e.g., Perry and Kuramitsu, 1981, *Infect. Immun.* 32: 1295-1297), protoplast transformation (see, e.g., Catt and Jollick, 1991, *Microbios* 68: 189-207), electroporation (see, e.g., Buckley *et al.*, 1999, *Appl. Environ. Microbiol.* 65: 3800-3804), or conjugation (see, e.g., Clewell, 1981, *Microbiol. Rev.* 45: 409-436). However, any method known in the art for introducing DNA into a host cell can be used.

The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

The host cell may be a fungal cell. "Fungi" as used herein includes the phyla *Ascomycota*, *Basidiomycota*, *Chytridiomycota*, and *Zygomycota* as well as the *Oomycota* and all mitosporic fungi (as defined by Hawksworth *et al.*, *In, Ainsworth and Bisby's Dictionary of The Fungi*, 8th edition, 1995, CAB International, University Press, Cambridge, UK).

The fungal host cell may be a yeast cell. "Yeast" as used herein includes ascosporegenous yeast (Endomycetales), basidiosporegenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this

invention, yeast shall be defined as described in *Biology and Activities of Yeast* (Skinner, Passmore, and Davenport, editors, *Soc. App. Bacteriol. Symposium Series No. 9*, 1980).

The yeast host cell may be a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell, such as a *Kluyveromyces lactis*, *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*,
5 *Saccharomyces kluyveri*, *Saccharomyces norbensis*, *Saccharomyces oviformis*, or *Yarrowia lipolytica* cell.

The fungal host cell may be a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision *Eumycota* and *Oomycota* (as defined by Hawksworth *et al.*,
10 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

The filamentous fungal host cell may be an *Acremonium*, *Aspergillus*, *Aureobasidium*,
15 *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

For example, the filamentous fungal host cell may be an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*,
20 *Aspergillus oryzae*, *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, *Ceriporiopsis subvermispora*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*,
25 *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochromum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*,
30 *Trichoderma reesei*, or *Trichoderma viride* cell.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238023, Yelton *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81: 1470-1474, and Christensen *et al.*, 1988, *Bio/Technology* 6: 1419-1422. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, *In* Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito *et al.*, 1983, *J. Bacteriol.* 153: 163; and Hinnen *et al.*, 1978, *Proc. Natl. Acad. Sci. USA* 75: 1920.

Methods of Production

The present invention also relates to methods of producing a polypeptide of the present invention, comprising (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. In a preferred aspect, the cell is a *Bacillus* cell. In a more preferred aspect, the cell is a *Bacillus sp-13231* cell.

The present invention also relates to methods of producing a polypeptide of the present invention, comprising (a) cultivating a recombinant host cell of the present invention under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

The host cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, or small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

The polypeptide may be detected using methods known in the art that are specific for the polypeptides. These detection methods include, but are not limited to, use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide.

The polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, collection, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

5 The polypeptide may be purified by a variety of procedures known in the art including, but not limited to, chromatography (**e.g.**, ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (**e.g.**, preparative isoelectric focusing), differential solubility (**e.g.**, ammonium sulfate precipitation), SDS-PAGE, or extraction (see, **e.g.**, *Protein Purification*, Janson and Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially
10 pure polypeptides.

In an alternative aspect, the polypeptide is not recovered, but rather a host cell of the present invention expressing the polypeptide is used as a source of the polypeptide.

Plants

15 The present invention also relates to isolated plants, **e.g.**, a transgenic plant, plant part, or plant cell, comprising a polynucleotide of the present invention so as to express and produce a polypeptide or domain in recoverable quantities. The polypeptide or domain may be recovered from the plant or plant part.

20 The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as *Festuca*, *Lolium*, temperate grass, such as *Agrostis*, and cereals, **e.g.**, wheat, oats, rye, barley, rice, sorghum, and maize (corn).

25 Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

30 Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, **e.g.**, epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilization of the invention are also considered plant parts, **e.g.**, embryos, endosperms, aleurone and seed coats.

Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

35 The transgenic plant or plant cell expressing the polypeptide or domain may be constructed

in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more expression constructs encoding the polypeptide or domain into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

5 The present invention also relates to methods of producing a polypeptide or domain of the present invention comprising (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide or domain under conditions conducive for production of the polypeptide or domain; and (b) recovering the polypeptide or domain.

10 **Signal peptide and propeptide**

The present invention also relates to an isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids -111 to -85 of SEQ ID NO: 2. The present invention also relates to an isolated polynucleotide encoding a propeptide comprising or consisting of amino acids -84 to -1 of SEQ ID NO: 2. The present invention also relates to an isolated polynucleotide
15 encoding a signal peptide and a propeptide comprising or consisting of amino acids -111 to -1 of SEQ ID NO: 2. The polynucleotides may further comprise a gene encoding a protein, which is operably linked to the signal peptide and/or propeptide. The protein is preferably foreign to the signal peptide and/or propeptide. In one aspect, the polynucleotide encoding the signal peptide is nucleotides 1 to 81 of SEQ ID NO: 1. In another aspect, the polynucleotide encoding the propeptide
20 is nucleotides 82 to 333 of SEQ ID NO: 1. In another aspect, the polynucleotide encoding the signal peptide and the propeptide is nucleotides 1 to 333 of SEQ ID NO: 1.

The present invention also relates to nucleic acid constructs, expression vectors and recombinant host cells comprising such polynucleotides.

25 **Enzyme compositions**

The present invention also relates to compositions comprising a polypeptide of the present invention.

Fermentation broth formulations or cell compositions

30 A composition may be a fermentation broth formulation or a cell composition comprising a polypeptide of the present invention. The fermentation broth product further comprises additional ingredients used in the fermentation process, such as, for example, cells (including, the host cells containing the gene encoding the polypeptide of the present invention which are used to produce the polypeptide of interest), cell debris, biomass, fermentation media and/or fermentation products.
35 In some embodiments, the composition is a cell-killed whole broth containing organic acid(s), killed

cells and/or cell debris, and culture medium.

The term "fermentation broth" as used herein refers to a preparation produced by cellular fermentation that undergoes no or minimal recovery and/or purification. For example, fermentation broths are produced when microbial cultures are grown to saturation, incubated under carbon-limiting conditions to allow protein synthesis (e.g., expression of enzymes by host cells) and secretion into cell culture medium. The fermentation broth can contain unfractionated or fractionated contents of the fermentation materials derived at the end of the fermentation. Typically, the fermentation broth is unfractionated and comprises the spent culture medium and cell debris present after the microbial cells (e.g., filamentous fungal cells) are removed, e.g., by centrifugation.

5
10 In some embodiments, the fermentation broth contains spent cell culture medium, extracellular enzymes, and viable and/or nonviable microbial cells.

In an embodiment, the fermentation broth formulation and cell compositions comprise a first organic acid component comprising at least one 1-5 carbon organic acid and/or a salt thereof and a second organic acid component comprising at least one 6 or more carbon organic acid and/or a salt thereof. In a specific embodiment, the first organic acid component is acetic acid, formic acid, propionic acid, a salt thereof, or a mixture of two or more of the foregoing and the second organic acid component is benzoic acid, cyclohexanecarboxylic acid, 4-methylvaleric acid, phenylacetic acid, a salt thereof, or a mixture of two or more of the foregoing.

15
20 In one aspect, the composition contains an organic acid(s), and optionally further contains killed cells and/or cell debris. In one embodiment, the killed cells and/or cell debris are removed from a cell-killed whole broth to provide a composition that is free of these components.

The fermentation broth formulations or cell compositions may further comprise a preservative and/or anti-microbial (e.g., bacteriostatic) agent, including, but not limited to, sorbitol, sodium chloride, potassium sorbate, and others known in the art.

25
30 The cell-killed whole broth or composition may contain the unfractionated contents of the fermentation materials derived at the end of the fermentation. Typically, the cell-killed whole broth or composition contains the spent culture medium and cell debris present after the microbial cells (e.g., filamentous fungal cells) are grown to saturation, incubated under carbon-limiting conditions to allow protein synthesis. In some embodiments, the cell-killed whole broth or composition contains the spent cell culture medium, extracellular enzymes, and killed filamentous fungal cells. In some embodiments, the microbial cells present in the cell-killed whole broth or composition can be permeabilized and/or lysed using methods known in the art.

A whole broth or cell composition as described herein is typically a liquid, but may contain insoluble components, such as killed cells, cell debris, culture media components, and/or insoluble enzyme(s). In some embodiments, insoluble components may be removed to provide a clarified

liquid composition.

Detergent compositions

In one embodiment, the invention is directed to compositions comprising a polypeptide of the present invention in combination with one or more additional cleaning composition components. Thus
5 one embodiment, the present invention relates to a detergent composition comprising an isolated polypeptide having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 60%, at least 61% at least 62% at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68% at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79% at least 80% at least 81% at least
10 82% at least 83% at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89% at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which have protease activity.

The choice of additional components is within the skill of the artisan and includes conventional ingredients, including the exemplary non-limiting components set forth below. The choice of
15 components may include, for fabric care, the consideration of the type of fabric to be cleaned, the type and/or degree of soiling, the temperature at which cleaning is to take place, and the formulation of the detergent product. Although components mentioned below are categorized by general header according to a particular functionality, this is not to be construed as a limitation, as a component may comprise additional functionalities as will be appreciated by the skilled artisan.

In one embodiment of the present invention, the a polypeptide of the present invention may
20 be added to a detergent composition in an amount corresponding to 0.001-200 mg of protein, such as 0.005-100 mg of protein, preferably 0.01-50 mg of protein, more preferably 0.05-20 mg of protein, even more preferably 0.1-10 mg of protein per liter of wash liquor.

The enzyme(s) of the detergent composition of the invention may be stabilized using
25 conventional stabilizing agents, *e.g.*, a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, *e.g.*, an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, and the composition may be formulated as described in, for example, WO92/19709 and WO92/19708 or the polypeptides according to the invention may be stabilized using peptide aldehydes or ketones such as described in WO
30 2005/1 05826 and WO 2009/1 18375.

Surfactants

The detergent composition may comprise one or more surfactants, which may be anionic
35 and/or cationic and/or non-ionic and/or semi-polar and/or zwitterionic, or a mixture thereof. In a particular embodiment, the detergent composition includes a mixture of one or more nonionic

surfactants and one or more anionic surfactants. The surfactant(s) is typically present at a level of from about 0.1% to 60% by weight, such as about 1% to about 40%, or about 3% to about 20%, or about 3% to about 10%. The surfactant(s) is chosen based on the desired cleaning application, and includes any conventional surfactant(s) known in the art. Any surfactant known in the art for use in detergents
5 may be utilized.

When included therein the detergent will usually contain from about 1% to about 40% by weight, such as from about 5% to about 30%, including from about 5% to about 15%, or from about 20% to about 25% of an anionic surfactant. Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched
10 alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester
15 sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodecenylolefin succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or soap, and combinations thereof.

When included therein the detergent will usually contain from about 1% to about 40% by
20 weight of a cationic surfactant. Non-limiting examples of cationic surfactants include alkyldimethylethanolamine quat (ADMEAQ), cetyltrimethylammonium bromide (CTAB), dimethyldistearylammonium chloride (DSDMAC), alkylbenzyltrimethylammonium, alkyl quaternary ammonium compounds, alkoxyated quaternary ammonium (AQA), and combinations thereof.

When included therein the detergent will usually contain from about 0.2% to about 40% by
25 weight of a non-ionic surfactant, for example from about 0.5% to about 30%, in particular from about 1% to about 20%, from about 3% to about 10%, such as from about 3% to about 5%, or from about 8% to about 12%. Non-limiting examples of non-ionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxyated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE),
30 nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxyated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxy alkyl fatty acid amides, or N-acyl N-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamide, FAGA), as well as products available under the trade names SPAN and TWEEN, and
35 combinations thereof.

When included therein the detergent will usually contain from about 1% to about 40% by weight of a semipolar surfactant. Non-limiting examples of semipolar surfactants include amine oxides (AO) such as alkyldimethylamineoxide, *N*-(coco alkyl)-*N,N*-dimethylamine oxide and *N*-(tallow-alkyl)-*N,N*-bis(2-hydroxyethyl)amine oxide, fatty acid alkanolamides and ethoxylated fatty acid alkanolamides, and combinations thereof.

When included therein the detergent will usually contain from about 1% to about 40% by weight of a zwitterionic surfactant. Non-limiting examples of zwitterionic surfactants include betaine, alkyldimethylbetaine, and sulfobetaine, and combinations thereof.

10 **Hydrotropes**

A hydrotrope is a compound that solubilises hydrophobic compounds in aqueous solutions (or oppositely, polar substances in a non-polar environment). Typically, hydrotropes have both hydrophilic and a hydrophobic character (so-called amphiphilic properties as known from surfactants); however the molecular structure of hydrotropes generally do not favor spontaneous self-aggregation, see e.g. review by Hodgdon and Kaler (2007), Current Opinion in Colloid & Interface Science 12: 121-128. Hydrotropes do not display a critical concentration above which self-aggregation occurs as found for surfactants and lipids forming micellar, lamellar or other well defined meso-phases. Instead, many hydrotropes show a continuous-type aggregation process where the sizes of aggregates grow as concentration increases. However, many hydrotropes alter the phase behavior, stability, and colloidal properties of systems containing substances of polar and non-polar character, including mixtures of water, oil, surfactants, and polymers. Hydrotropes are classically used across industries from pharma, personal care, food, to technical applications. Use of hydrotropes in detergent compositions allow for example more concentrated formulations of surfactants (as in the process of compacting liquid detergents by removing water) without inducing undesired phenomena such as phase separation or high viscosity.

The detergent may contain 0-5% by weight, such as about 0.5 to about 5%, or about 3% to about 5%, of a hydrotrope. Any hydrotrope known in the art for use in detergents may be utilized. Non-limiting examples of hydrotropes include sodium benzene sulfonate, sodium p-toluene sulfonate (STS), sodium xylene sulfonate (SXS), sodium cumene sulfonate (SCS), sodium cymene sulfonate, amine oxides, alcohols and polyglycoethers, sodium hydroxynaphthoate, sodium hydroxynaphthalene sulfonate, sodium ethylhexyl sulfate, and combinations thereof.

Builders and co-builders

The detergent composition may contain about 0-65% by weight, such as about 5% to about 50% of a detergent builder or co-builder, or a mixture thereof. In a dish wash detergent, the level of

builder is typically 40-65%, particularly 50-65%. The builder and/or co-builder may particularly be a chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in laundry detergents may be utilized. Non-limiting examples of builders include zeolites, diphosphates (pyrophosphates), triphosphates such as sodium triphosphate (STP or STPP), carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), ethanolamines such as 2-aminoethan-1-ol (MEA), diethanolamine (DEA, also known as iminodiethanol), triethanolamine (TEA, also known as 2,2',2"-nitrilotriethanol), and carboxymethyl inulin (CMI), and combinations thereof.

The detergent composition may also contain 0-65% by weight, such as about 5% to about 40%, of a detergent co-builder, or a mixture thereof. The detergent composition may include include a co-builder alone, or in combination with a builder, for example a zeolite builder. Non-limiting examples of co-builders include homopolymers of polyacrylates or copolymers thereof, such as poly(acrylic acid) (PAA) or copoly(acrylic acid/maleic acid) (PAA/PMA). Further non-limiting examples include citrate, chelators such as aminocarboxylates, aminopolycarboxylates and phosphonates, and alkyl- or alkenylsuccinic acid. Additional specific examples include 2,2',2"-nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), iminodisuccinic acid (IDS), ethylenediamine-*N,N'*-disuccinic acid (EDDS), methylglycinediacetic acid (MGDA), glutamic acid-*N,N'*-diacetic acid (GLDA), ethylenediaminetetra(methylenephosphonic acid) (HEDP), ethylenediaminetetra(methylenephosphonic acid) (EDTMPA), diethylenetriaminepenta(methylenephosphonic acid) (DTPMPA), *N*-(2-hydroxyethyl)iminodiacetic acid (EDG), aspartic acid-*N*-monoacetic acid (ASMA), aspartic acid-*N,N'*-diacetic acid (ASDA), aspartic acid-*N*-monopropionic acid (ASMP), iminodisuccinic acid (IDA), *N*-(2-sulfomethyl)aspartic acid (SMAS), *N*-(2-sulfoethyl)aspartic acid (SEAS), *N*-(2-sulfomethyl)glutamic acid (SMGL), *N*-(2-sulfoethyl)glutamic acid (SEGL), *N*-methyliminodiacetic acid (MIDA), α -alanine-*N,N'*-diacetic acid (α -ALDA), serine-*N,N'*-diacetic acid (SEDA), isoserine-*N,N'*-diacetic acid (ISDA), phenylalanine-*N,N'*-diacetic acid (PHDA), anthranilic acid-*N,N'*-diacetic acid (ANDA), sulfanilic acid-*N,N'*-diacetic acid (SLDA), taurine-*N,N'*-diacetic acid (TUDA) and sulfomethyl-*N,N'*-diacetic acid (SMDA), *N*-(2-hydroxyethyl)-ethylidenediamine-*N,N',N'*-triacetate (HEDTA), diethanolglycine (DEG), diethylenetriamine penta(methylenephosphonic acid) (DTPMP), aminotris(methylenephosphonic acid) (ATMP), and combinations and salts thereof. Further exemplary builders and/or co-builders are described in, e.g., WO 09/1 02854, US 5977053.

Bleaching systems

The detergent may contain 0-10% by weight, such as about 1% to about 5%, of a bleaching system. Any bleaching system known in the art for use in laundry detergents may be utilized. Suitable bleaching system components include bleaching catalysts, photobleaches, bleach activators,

sources of hydrogen peroxide such as sodium percarbonate and sodium perborates, preformed peracids and mixtures thereof. Suitable preformed peracids include, but are not limited to, peroxy-carboxylic acids and salts, percarbonic acids and salts, perimidic acids and salts, peroxy-monosulfuric acids and salts, for example, Oxone (R), and mixtures thereof. Non-limiting


5 examples of bleaching systems include peroxide-based bleaching systems, which may comprise, for example, an inorganic salt, including alkali metal salts such as sodium salts of perborate (usually mono- or tetra-hydrate), percarbonate, persulfate, perphosphate, persilicate salts, in combination with a peracid-forming bleach activator. The term bleach activator is meant herein as a compound which reacts with peroxygen bleach like hydrogen peroxide to form a peracid. The peracid thus formed

10 constitutes the activated bleach. Suitable bleach activators to be used herein include those belonging to the class of esters amides, imides or anhydrides, Suitable examples are tetracetylene diamine (TAED), sodium 4-[(3,5,5-trimethylhexanoyl)oxy]benzene sulfonate (ISONOBS), diperoxy dodecanoic acid, 4-(dodecanoyloxy)benzenesulfonate (LOBS), 4-(decanoyloxy)benzenesulfonate, 4-(decanoyloxy)benzoate (DOBS), 4-(nonanoyloxy)benzenesulfonate (NOBS), and/or those

15 disclosed in W098/17767. A particular family of bleach activators of interest was disclosed in EP624154 and particularly preferred in that family is acetyl triethyl citrate (ATC). ATC or a short chain triglyceride like Triacin has the advantage that it is environmental friendly as it eventually degrades into citric acid and alcohol. Furthermore acetyl triethyl citrate and triacetin has a good hydrolytical stability in the product upon storage and it is an efficient bleach activator. Finally ATC provides a good building

20 capacity to the laundry additive. Alternatively, the bleaching system may comprise peroxyacids of, for example, the amide, imide, or sulfone type. The bleaching system may also comprise peracids such as 6-(phthalimido)peroxyhexanoic acid (PAP). The bleaching system may also include a bleach catalyst. In some embodiments the bleach component may be an organic catalyst selected from the group consisting of organic catalysts having the following formulae:

25



30

(iii) and mixtures thereof; wherein each R¹ is independently a branched alkyl group containing from 9 to 24 carbons or linear alkyl group containing from 11 to 24 carbons, preferably each R¹ is independently a branched alkyl group containing from 9 to 18 carbons or linear alkyl group

35 containing from 11 to 18 carbons, more preferably each R¹ is independently selected from the

group consisting of 2-propylheptyl, 2-butyloctyl, 2-pentylonyl, 2-hexyldecyl, n- dodecyl, n- tetradecyl, n-hexadecyl, n-octadecyl, iso-nonyl, iso-decyl, iso- tridecyl and iso-pentadecyl. Other exemplary bleaching systems are described, e.g., in WO2007/087258, WO2007/087244, WO2007/087259, WO2007/087242. Suitable photobleaches may for example be sulfonated zinc
5 phthalocyanine

Polymers

The detergent may contain 0-10% by weight, such as 0.5-5%, 2-5%, 0.5-2% or 0.2-1% of a polymer. Any polymer known in the art for use in detergents may be utilized. The polymer may
10 function as a co-builder as mentioned above, or may provide antiredeposition, fiber protection, soil release, dye transfer inhibition, grease cleaning and/or anti-foaming properties. Some polymers may have more than one of the above-mentioned properties and/or more than one of the below-mentioned motifs. Exemplary polymers include (carboxymethyl)cellulose (CMC), polyvinyl alcohol (PVA), poly(vinylpyrrolidone) (PVP), poly(ethyleneglycol) or poly(ethylene oxide) (PEG), ethoxylated
15 poly(ethyleneimine), carboxymethyl inulin (CMI), and polycarboxylates such as PAA, PAA/PMA, poly-aspartic acid, and lauryl methacrylate/acrylic acid copolymers, hydrophobically modified CMC (HM-CMC) and silicones, copolymers of terephthalic acid and oligomeric glycols, copolymers of poly(ethylene terephthalate) and poly(oxyethene terephthalate) (PET-POET), PVP, poly(vinylimidazole) (PVI), poly(vinylpyridine-*N*-oxide) (PVPO or PVPNO) and polyvinylpyrrolidone-
20 vinylimidazole (PVPVI). Further exemplary polymers include sulfonated polycarboxylates, polyethylene oxide and polypropylene oxide (PEO-PPO) and diquaternium ethoxy sulfate. Other exemplary polymers are disclosed in, e.g., WO 2006/130575. Salts of the above-mentioned polymers are also contemplated.

25 Fabric hueing agents

The detergent compositions of the present invention may also include fabric hueing agents such as dyes or pigments which when formulated in detergent compositions can deposit onto a fabric when said fabric is contacted with a wash liquor comprising said detergent compositions thus altering the tint of said fabric through absorption/reflection of visible light. Fluorescent whitening
30 agents emit at least some visible light. In contrast, fabric hueing agents alter the tint of a surface as they absorb at least a portion of the visible light spectrum. Suitable fabric hueing agents include dyes and dye-clay conjugates, and may also include pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Colour Index (C.I.) classifications of
35 Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic Violet and

Basic Red, or mixtures thereof, for example as described in WO2005/03274, WO2005/03275, WO2005/03276 and EP1 876226 (hereby incorporated by reference). The detergent composition preferably comprises from about 0.00003 wt% to about 0.2 wt%, from about 0.00008 wt% to about 0.05 wt%, or even from about 0.0001 wt% to about 0.04 wt% fabric hueing agent. The composition
5 may comprise from 0.0001 wt% to 0.2 wt% fabric hueing agent, this may be especially preferred when the composition is in the form of a unit dose pouch. Suitable hueing agents are also disclosed in, e.g., WO 2007/087257, WO2007/087243.

Additional enzymes

10 The detergent additive as well as the detergent composition may comprise one or more enzymes such as a protease, lipase, cutinase, an amylase, carbohydrase, cellulase, pectinase, mannanase, arabinase, galactanase, xylanase, oxidase, e.g., a laccase, and/or peroxidase.

In general, the properties of the selected enzyme(s) should be compatible with the selected detergent, (i.e., pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.),
15 and the enzyme(s) should be present in effective amounts.

Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered variants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum*
20 disclosed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

Especially suitable cellulases are the alkaline or neutral cellulases having color care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/1 1262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO
25 95/24471, WO 98/12307 and PCT/DK98/00299.

Commercially available cellulases include Celluzyme™, and Carezyme™ (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation).

Proteases: Suitable proteases include those of bacterial, fungal, plant, viral or animal origin e.g. vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. It may be an alkaline protease, such as a serine protease or a metalloprotease. A serine protease may for example be of the S1 family, such as trypsin, or the S8 family such as subtilisin. A metalloproteases protease may for example be a thermolysin from e.g. family M4 or other metalloprotease such as those from M5, M7 or M8 families.
30

The term "subtilases" refers to a sub-group of serine protease according to Siezen et al., Protein Engng. 4 (1991) 719-737 and Siezen et al. Protein Science 6 (1997) 501-523. Serine proteases are a subgroup of proteases characterized by having a serine in the active site, which forms a covalent adduct with the substrate. The subtilases may be divided into 6 sub-divisions, i.e. the Subtilisin family, the Thermitase family, the Proteinase K family, the Lantibiotic peptidase family, the Kexin family and the Pyrolysin family.

Examples of subtilases are those derived from *Bacillus* such as *Bacillus lentus*, *B. alkalophilus*, *B. subtilis*, *B. amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii* described in; US7262042 and WO9/021867, and *subtilisin lentus*, *subtilisin Novo*, *subtilisin Carlsberg*, *Bacillus licheniformis*, *subtilisin BPN'*, *subtilisin 309*, *subtilisin 147* and *subtilisin 168* described in WO89/06279 and protease PD138 described in (WO93/18140). Other useful proteases may be those described in W092/175177, WO01/016285, WO02/026024 and WO02/016547. Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO89/06270, W094/25583 and WO05/040372, and the chymotrypsin proteases derived from *Cellumonas* described in WO05/052161 and WO05/052146.

A further preferred protease is the alkaline protease from *Bacillus lentus* DSM 5483, as described for example in W095/23221, and variants thereof which are described in WO92/21760, W095/23221, EP1921 147 and EP1921 148.

Examples of metalloproteases are the neutral metalloprotease as described in WO07/044993 (Genencor Int.) such as those derived from *Bacillus amyloliquefaciens*.

Examples of useful proteases are the variants described in: W092/19729, WO96/034946, WO98/201 15, WO98/201 16, WO99/01 1768, WO01/44452, WO03/006602, WO04/03186, WO04/041979, WO07/006305, W01 1/036263, W01 1/036264, especially the variants with substitutions in one or more of the following positions: 3, 4, 9, 15, 24, 27, 42, 55, 59, 60, 66, 74, 85, 96, 97, 98, 99, 100, 101, 102, 104, 116, 118, 121, 126, 127, 128, 154, 156, 157, 158, 161, 164, 176, 179, 182, 185, 188, 189, 193, 198, 199, 200, 203, 206, 211, 212, 216, 218, 226, 229, 230, 239, 246, 255, 256, 268 and 269 wherein the positions correspond to the positions of the *Bacillus Lentus* protease shown in SEQ ID NO 1 of WO 2016/001449. More preferred the subtilase variants may comprise the mutations: S3T, V4I, S9R, S9E, A15T, S24G, S24R, K27R, N42R, S55P, G59E, G59D, N60D, N60E, V66A, N74D, N85S, N85R, G96S, G96A, S97G, S97D, S97A, S97SD, S99E, S99D, S99G, S99M, S99N, S99R, S99H, S101A, V102I, V102Y, V102N, S104A, G116V, G116R, H118D, H118N, N120S, S126L, P127Q, S128A, S154D, A156E, G157D, G157P, S158E, Y161A, R164S, Q176E, N179E, S182E, Q185N, A188P, G189E, V193M, N198D, V199I, Y203W, S206G, L21 1Q, L21 1D, N212D, N212S, M216S, A226V, K229L, Q230H, Q239R, N246K, N255W, N255D, N255E, L256E, L256D T268A, R269H. The protease variants are preferably variants of the *Bacillus lentus* protease (Savinase®) shown in SEQ ID NO 1 of WO 2016/001449, the *Bacillus amylolichenifaciens* protease (BPN') shown in SEQ ID NO 2 of WO20 16/00 1449. The protease

variants preferably have at least 80 % sequence identity to SEQ ID NO 1 or SEQ ID NO 2 of WO 2016/001449.

A protease variant comprising a substitution at one or more positions corresponding to positions 171, 173, 175, 179, or 180 of SEQ ID NO: 1 of WO2004/067737, wherein said protease variant has a sequence identity of at least 75% but less than 100% to SEQ ID NO: 1 of
5 WO2004/067737.

Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Duralase™, Durazym™, Relase®, Relase® Ultra, Savinase®, Savinase® Ultra, Primase®, Polarzyme®, Kannase®, Liquanase®, Liquanase® Ultra, Ovozyme®, Coronase®,
10 Coronase® Ultra, Blaze®, Blaze Evity® 100T, Blaze Evity® 125T, Blaze Evity® 150T, Neutrase®, Everlase® and Esperase® (Novozymes A/S), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, , , Purafect Ox®, Purafect OxP®, Puramax®, FN2®, FN3®, FN4®, Excellase®, Excellenz P1000™, Excellenz P1250™, Eraser®, Preferenz P100™, Purafect Prime®, Preferenz P110™, Effectenz P1000™, Purafect®™, Effectenz P1050™, Purafect Ox®™, Effectenz
15 P2000™, Purafast®, Properase®, Opticlean® and Optimase® (Danisco/DuPont), Axapem™ (Gist-Brocades N.V.), BLAP (sequence shown in Figure 29 of US5352604) and variants hereof (Henkel AG) and KAP (*Bacillus alkalophilus* subtilisin) from Kao.

Lipases and Cutinases: Suitable lipases and cutinases include those of bacterial or fungal origin. Chemically modified or protein engineered variant enzymes are included. Examples include
20 lipase from *Thermomyces*, e.g. from *T. lanuginosus* (previously named *Humicola lanuginosa*) as described in EP258068 and EP305216, cutinase from *Humicola*, e.g. *H. insolens* (WO96/13580), lipase from strains of *Pseudomonas* (some of these now renamed to *Burkholderia*), e.g. *P. alcaligenes* or *P. pseudoalcaligenes* (EP218272), *P. cepacia* (EP331376), *P. sp.* strain SD705 (WO95/06720 & WO96/27002), *P. wisconsinensis* (WO96/12012), GDSL-type *Streptomyces*
25 lipases (W0 10/065455), cutinase from *Magnaporthe grisea* (W0 10/1 07560), cutinase from *Pseudomonas mendocina* (US5,389,536), lipase from *Thermobifida fusca* (W0 11/084412), *Geobacillus stearothermophilus* lipase (W01 1/084417), lipase from *Bacillus subtilis* (W0 11/084599), and lipase from *Streptomyces griseus* (W01 1/150157) and *S. pristinaespiralis* (W012/137147).

Other examples are lipase variants such as those described in EP407225, WO92/05249, WO94/01541, W094/25578, W095/14783, WO95/30744, W095/35381, W095/22615, WO96/00292, WO97/04079, WO97/07202, WO00/34450, WO00/60063, WO01/92502, WO07/87508 and WO09/1 09500.

Preferred commercial lipase products include Lipolase™, Lipex™; Lipolex™ and
35 Lipoclean™ (Novozymes A/S), Lumafast (originally from Genencor) and Lipomax (originally from Gist-Brocades).

Still other examples are lipases sometimes referred to as acyl transferases or

perhydrolases, e.g. acyltransferases with homology to *Candida antarctica* lipase A (WO10/1 11143), acyltransferase from *Mycobacterium smegmatis* (WO05/56782), perhydrolases from the CE 7 family (WO09/67279), and variants of the *M. smegmatis* perhydrolase in particular the S54V variant used in the commercial product Gentle Power Bleach from Huntsman Textile Effects Pte Ltd

5 (W01 0/1 00028).

Amylases: Suitable amylases which can be used together with the protease of the invention may be an alpha-amylase or a glucoamylase and may be of bacterial or fungal origin. Chemically modified or protein engineered variants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *Bacillus licheniformis*, described in

10 more detail in GB 1,296,839.

Suitable amylases include amylases having SEQ ID NO: 2 in WO 95/10603 or variants having 90% sequence identity to SEQ ID NO: 3 thereof. Preferred variants are described in WO 94/02597, WO 94/18314, WO 97/43424 and SEQ ID NO: 4 of WO 99/019467, such as variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154,

15 156, 178, 179, 181, 188, 190, 197, 201, 202, 207, 208, 209, 211, 243, 264, 304, 305, 391, 408, and 444.

Different suitable amylases include amylases having SEQ ID NO: 6 in WO 02/010355 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a deletion in positions 181 and 182 and a substitution in position 193.

20 Other amylases which are suitable are hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 of WO 2006/066594 or variants having 90% sequence identity thereof. Preferred variants of this hybrid alpha-amylase are those having a substitution, a deletion or an insertion in one of more of the

25 following positions: G48, T49, G107, H156, A181, N190, M197, I201, A209 and Q264. Most preferred variants of the hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of SEQ ID NO: 4 are those having the substitutions:

M197T;

30 H156Y+A181T+N190F+A209V+Q264S; or

G48A+T49I+G107A+H156Y+A181T+N190F+I201 F+A209V+Q264S.

Further amylases which are suitable are amylases having SEQ ID NO: 6 in WO 99/019467 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a substitution, a deletion or an insertion in one or more of the following

35 positions: R181, G182, H183, G184, N195, I206, E212, E216 and K269. Particularly preferred

amylases are those having deletion in positions R181 and G182, or positions H183 and G184.

Additional amylases which can be used are those having SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 2 or SEQ ID NO: 7 of WO 96/023873 or variants thereof having 90% sequence identity to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7. Preferred variants of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7 are those having a substitution, a deletion or an insertion in one or more of the following positions: 140, 181, 182, 183, 184, 195, 206, 212, 243, 260, 269, 304 and 476, using SEQ ID 2 of WO 96/023873 for numbering. More preferred variants are those having a deletion in two positions selected from 181, 182, 183 and 184, such as 181 and 182, 182 and 183, or positions 183 and 184. Most preferred amylase variants of SEQ ID NO: 1, SEQ ID NO: 2 or SEQ ID NO: 7 are those having a deletion in positions 183 and 184 and a substitution in one or more of positions 140, 195, 206, 243, 260, 304 and 476.

Other amylases which can be used are amylases having SEQ ID NO: 2 of WO 08/153815, SEQ ID NO: 10 in WO 01/66712 or variants thereof having 90% sequence identity to SEQ ID NO: 2 of WO 08/153815 or 90% sequence identity to SEQ ID NO: 10 in WO 01/66712. Preferred variants of SEQ ID NO: 10 in WO 01/66712 are those having a substitution, a deletion or an insertion in one or more of the following positions: 176, 177, 178, 179, 190, 201, 207, 211 and 264.

Further suitable amylases are amylases having SEQ ID NO: 2 of WO 09/061380 or variants having 90% sequence identity to SEQ ID NO: 2 thereof. Preferred variants of SEQ ID NO: 2 are those having a truncation of the C-terminus and/or a substitution, a deletion or an insertion in one or more of the following positions: Q87, Q98, S125, N128, T131, T165, K178, R180, S181, T182, G183, M201, F202, N225, S243, N272, N282, Y305, R309, D319, Q320, Q359, K444 and G475. More preferred variants of SEQ ID NO: 2 are those having the substitution in one or more of the following positions: Q87E,R, Q98R, S125A, N128C, T131 I, T165I, K178L, T182G, M201 L, F202Y, N225E,R, N272E,R, S243Q,A,E,D, Y305R, R309A, Q320R, Q359E, K444E and G475K and/or deletion in position R180 and/or S181 or of T182 and/or G183. Most preferred amylase variants of SEQ ID NO: 2 are those having the substitutions:

N128C+K178L+T182G+Y305R+G475K;

N128C+K178L+T182G+F202Y+Y305R+D319T+G475K;

S125A+N128C+K178L+T182G+Y305R+G475K; or

S125A+N128C+T131 I+T165I+K178L+T182G+Y305R+G475K wherein the variants are C-terminally truncated and optionally further comprises a substitution at position 243 and/or a deletion at position 180 and/or position 181.

Other suitable amylases are the alpha-amylase having SEQ ID NO: 12 in WO01/66712 or a variant having at least 90% sequence identity to SEQ ID NO: 12. Preferred amylase variants are those having a substitution, a deletion or an insertion in one or more of the following positions of

SEQ ID NO: 12 in WO01/66712: R28, R118, N174; R181, G182, D183, G184, G186, W189, N195, M202, Y298, N299, K302, S303, N306, R310, N314; R320, H324, E345, Y396, R400, W439, R444, N445, K446, Q449, R458, N471, N484. Particular preferred amylases include variants having a deletion of D183 and G184 and having the substitutions R118K, N195F, R320K and R458K, and a
5 variant additionally having substitutions in one or more position selected from the group: M9, G149, G182, G186, M202, T257, Y295, N299, M323, E345 and A339, most preferred a variant that additionally has substitutions in all these positions.

Other examples are amylase variants such as those described in WO2011/098531, WO2013/001078 and WO2013/001087.

10 Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™, Stainzyme™, Stainzyme Plus™, Natalase™, Liquozyme X and BAN™ (from Novozymes A/S), and Rapidase™, Purastar™/Effectenz™, Powerase and Preferenz S100 (from Genencor International Inc./DuPont).

Peroxidases/Oxidases: Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered variants are included. Examples of useful
15 peroxidases include peroxidases from *Coprinus*, e.g., from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

Commercially available peroxidases include Guardzyme™ (Novozymes A/S).

The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these
20 enzymes. A detergent additive of the invention, i.e., a separate additive or a combined additive, can be formulated, for example, as a granulate, liquid, slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452
25 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-
30 forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

35 **Adjunct materials**

Any detergent components known in the art for use in laundry detergents may also be utilized. Other optional detergent components include anti-corrosion agents, anti-shrink agents, anti-soil redeposition agents, anti-wrinkling agents, bactericides, binders, corrosion inhibitors, disintegrants/disintegration agents, dyes, enzyme stabilizers (including boric acid, borates, CMC, and/or polyols such as propylene glycol), fabric conditioners including clays, fillers/processing aids, fluorescent whitening agents/optical brighteners, foam boosters, foam (suds) regulators, perfumes, soil-suspending agents, softeners, suds suppressors, tarnish inhibitors, and wicking agents, either alone or in combination. Any ingredient known in the art for use in laundry detergents may be utilized. The choice of such ingredients is well within the skill of the artisan.

Dispersants - The detergent compositions of the present invention can also contain dispersants. In particular powdered detergents may comprise dispersants. Suitable water-soluble organic materials include the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Suitable dispersants are for example described in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc.

Dye Transfer Inhibiting Agents - The detergent compositions of the present invention may also include one or more dye transfer inhibiting agents. Suitable polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof. When present in a subject composition, the dye transfer inhibiting agents may be present at levels from about 0.0001 % to about 10%, from about 0.01% to about 5% or even from about 0.1% to about 3% by weight of the composition.

Fluorescent whitening agent - The detergent compositions of the present invention will preferably also contain additional components that may tint articles being cleaned, such as fluorescent whitening agent or optical brighteners. Where present the brightener is preferably at a level of about 0.01% to about 0.5%.. Any fluorescent whitening agent suitable for use in a laundry detergent composition may be used in the composition of the present invention. The most commonly used fluorescent whitening agents are those belonging to the classes of diaminostilbene-sulfonic acid derivatives, diarylpyrazoline derivatives and bisphenyl-distyryl derivatives. Examples of the diaminostilbene-sulfonic acid derivative type of fluorescent whitening agents include the sodium salts of: 4,4'-bis-(2-diethanolamino-4-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate; 4,4'-bis-(2,4-dianilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate; 4,4'-

bis-(2-anilino-4-(N-methyl-N-2-hydroxy-ethylamino)-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(4-phenyl-2,1,3-triazol-2-yl)stilbene-2,2'-disulfonate; 4,4'-bis-(2-anilino-4(1-methyl-2-hydroxy-ethylamino)-s-triazin-6-ylamino) stilbene-2,2'-disulfonate and sodium 5-(2'-naphtho[1,2-c][1,2,3]triazol-2-yl)-2-[(E)-2-phenylvinyl]benzenesulfonate. Preferred fluorescent whitening agents are Tinopal DMS and Tinopal CBS available from Ciba-Geigy AG, Basel, Switzerland. Tinopal DMS is the disodium salt of 4,4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino) stilbene disulfonate. Tinopal CBS is the disodium salt of 2,2'-bis-(phenyl-styryl)-disulfonate. Also preferred are fluorescent whitening agents is the commercially available Parawhite KX, supplied by Paramount Minerals and Chemicals, Mumbai, India. Other fluorescers suitable for use in the invention include the 1-3-diaryl pyrazolines and the 7-alkylaminocoumarins. Suitable fluorescent brightener levels include lower levels of from about 0.01, from 0.05, from about 0.1 or even from about 0.2 wt % to upper levels of 0.5 or even 0.75 wt%.

Soil release polymers - The detergent compositions of the present invention may also include one or more soil release polymers which aid the removal of soils from fabrics such as cotton and polyester based fabrics, in particular the removal of hydrophobic soils from polyester based fabrics. The soil release polymers may for example be nonionic or anionic terephthalate based polymers, polyvinyl caprolactam and related copolymers, vinyl graft copolymers, polyester polyamides see for example Chapter 7 in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc. Another type of soil release polymers are amphiphilic alkoxyated grease cleaning polymers comprising a core structure and a plurality of alkoxyate groups attached to that core structure. The core structure may comprise a polyalkylenimine structure or a polyalkanolamine structure as described in detail in WO 2009/087523 (hereby incorporated by reference). Furthermore random graft co-polymers are suitable soil release polymers Suitable graft co-polymers are described in more detail in WO 2007/138054, WO 2006/108856 and WO 2006/1 13314 (hereby incorporated by reference). Other soil release polymers are substituted polysaccharide structures especially substituted cellulosic structures such as modified cellulose derivatives such as those described in EP 1867808 or WO 2003/040279 (both are hereby incorporated by reference). Suitable cellulosic polymers include cellulose, cellulose ethers, cellulose esters, cellulose amides and mixtures thereof. Suitable cellulosic polymers include anionically modified cellulose, nonionically modified cellulose, cationically modified cellulose, zwitterionically modified cellulose, and mixtures thereof. Suitable cellulosic polymers include methyl cellulose, carboxy methyl cellulose, ethyl cellulose, hydroxyl ethyl cellulose, hydroxyl propyl methyl cellulose, ester carboxy methyl cellulose, and mixtures thereof.

35

Anti-redeposition agents - The detergent compositions of the present invention may also include one or more anti-redeposition agents such as carboxymethylcellulose (CMC), polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), polyoxyethylene and/or polyethyleneglycol (PEG), homopolymers of acrylic acid, copolymers of acrylic acid and maleic acid, and ethoxylated polyethyleneimines. The cellulose based polymers described under soil release polymers above may also function as anti-redeposition agents.

Other suitable adjunct materials include, but are not limited to, anti-shrink agents, anti-wrinkling agents, bactericides, binders, carriers, dyes, enzyme stabilizers, fabric softeners, fillers, foam regulators, hydrotropes, perfumes, pigments, sod suppressors, solvents, structurants for liquid detergents and/or structure elasticizing agents.

Formulation of detergent products

The detergent composition of the invention may be in any convenient form, e.g., a bar, a homogenous tablet, a tablet having two or more layers, a regular or compact powder, a granule, a paste, a gel, or a regular, compact or concentrated liquid.

Detergent formulation forms: Layers (same or different phases), Pouches, versus forms for Machine dosing unit.

Pouches can be configured as single or multicompartments. It can be of any form, shape and material which is suitable for hold the composition, e.g. without allowing the release of the composition to release of the composition from the pouch prior to water contact. The pouch is made from water soluble film which encloses an inner volume. Said inner volume can be divided into compartments of the pouch. Preferred films are polymeric materials preferably polymers which are formed into a film or sheet. Preferred polymers, copolymers or derivatives thereof are selected polyacrylates, and water soluble acrylate copolymers, methyl cellulose, carboxy methyl cellulose, sodium dextrin, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, malto dextrin, poly methacrylates, most preferably polyvinyl alcohol copolymers and, hydroxypropyl methyl cellulose (HPMC). Preferably the level of polymer in the film for example PVA is at least about 60%. Preferred average molecular weight will typically be about 20,000 to about 150,000. Films can also be of blended compositions comprising hydrolytically degradable and water soluble polymer blends such as polylactide and polyvinyl alcohol (known under the Trade reference M8630 as sold by MonoSol LLC, Indiana, USA) plus plasticisers like glycerol, ethylene glycerol, propylene glycol, sorbitol and mixtures thereof. The pouches can comprise a solid laundry cleaning composition or part components and/or a liquid cleaning composition or part components separated by the water soluble film. The compartment for liquid components can be different in composition than compartments containing solids. Ref:

(US2009/001 1970 A1)

Detergent ingredients can be separated physically from each other by compartments in water dissolvable pouches or in different layers of tablets. Thereby negative storage interaction between components can be avoided. Different dissolution profiles of each of the compartments can also give rise to delayed dissolution of selected components in the wash solution.

Definition/characteristics of the forms:

A liquid or gel detergent, which is not unit dosed, may be aqueous, typically containing at least 20% by weight and up to 95% water, such as up to about 70% water, up to about 65% water, up to about 55% water, up to about 45% water, up to about 35% water. Other types of liquids, including without limitation, alkanols, amines, diols, ethers and polyols may be included in an aqueous liquid or gel. An aqueous liquid or gel detergent may contain from 0-30% organic solvent.

A liquid or gel detergent may be non-aqueous.

15 Laundry soap bars

The polypeptides of the invention may be added to laundry soap bars and used for hand washing laundry, fabrics and/or textiles. The term laundry soap bar includes laundry bars, soap bars, combo bars, syndet bars and detergent bars. The types of bar usually differ in the type of surfactant they contain, and the term laundry soap bar includes those containing soaps from fatty acids and/or synthetic soaps. The laundry soap bar has a physical form which is solid and not a liquid, gel or a powder at room temperature. The term solid is defined as a physical form which does not significantly change over time, i.e. if a solid object (e.g. laundry soap bar) is placed inside a container, the solid object does not change to fill the container it is placed in. The bar is a solid typically in bar form but can be in other solid shapes such as round or oval.

The laundry soap bar may contain one or more additional enzymes, protease inhibitors such as peptide aldehydes (or hydrosulfite adduct or hemiacetal adduct), boric acid, borate, borax and/or phenylboronic acid derivatives such as 4-formylphenylboronic acid, one or more soaps or synthetic surfactants, polyols such as glycerine, pH controlling compounds such as fatty acids, citric acid, acetic acid and/or formic acid, and/or a salt of a monovalent cation and an organic anion wherein the monovalent cation may be for example Na^+ , K^+ or NH_4^+ and the organic anion may be for example formate, acetate, citrate or lactate such that the salt of a monovalent cation and an organic anion may be, for example, sodium formate.

The laundry soap bar may also contain complexing agents like EDTA and HEDP, perfumes and/or different type of fillers, surfactants e.g. anionic synthetic surfactants, builders, polymeric soil release agents, detergent chelators, stabilizing agents, fillers, dyes, colorants, dye transfer

inhibitors, alkoxyated polycarbonates, suds suppressers, structurants, binders, leaching agents, bleaching activators, clay soil removal agents, anti-redeposition agents, polymeric dispersing agents, brighteners, fabric softeners, perfumes and/or other compounds known in the art.

5 The laundry soap bar may be processed in conventional laundry soap bar making equipment such as but not limited to: mixers, plodders, e.g a two stage vacuum plodder, extruders, cutters, logo-stampers, cooling tunnels and wrappers. The invention is not limited to preparing the laundry soap bars by any single method. The premix of the invention may be added to the soap at different stages of the process. For example, the premix containing a soap, an enzyme, optionally one or more additional enzymes, a protease inhibitor, and a salt of a monovalent cation and an
10 organic anion may be prepared and and the mixture is then plodded. The enzyme and optional additional enzymes may be added at the same time as the protease inhibitor for example in liquid form. Besides the mixing step and the plodding step, the process may further comprise the steps of milling, extruding, cutting, stamping, cooling and/or wrapping.

15 **Uses**

The present invention is also directed to methods for using the protease according to the invention or compositions thereof in laundry of textilea and fabrics, such as house hold laundry washing and industrial laundry washing.

The invention is also directed to methods for using the compositions thereof in hard surface
20 cleaning such as Automated Dish Washing (ADW), car wash and cleaning of Industrial surfaces. Thus one embodiment, the present invention relates to the use in cleaning such as laundry or desh wash of a protease according to the invention or a detergent composition comprising a protease according to the present invention having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 60%, at least 61% at least 62% at least 63%, at least 64%, at least 65%, at least
25 66%, at least 67%, at least 68% at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79% at least 80% at least 81% at least 82% at least 83% at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89% at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. Thus in one
30 embodiment, the present invention relates to the use of an isolated polypeptide having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 60%, at least 61% at least 62% at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68% at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79% at least 80% at least 81% at least 82% at least 83% at least
35 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89% at least 90%, at least

91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% which have protease activity in a detergent, a cleaning process and/or laundry process.

Use in detergents. The polypeptides of the present invention may be added to and thus
5 become a component of a detergent composition.

The detergent composition of the present invention may be formulated, for example, as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated
10 for hand or machine dishwashing operations.

In a specific aspect, the present invention provides a detergent additive comprising a polypeptide of the present invention as described herein.

Use of proteases of the invention in detergents

15 The soils and stains that are important for detergent formulators are composed of many different substances, and a range of different enzymes, all with different substrate specificities have been developed for use in detergents both in relation to laundry and hard surface cleaning, such as dishwashing. These enzymes are considered to provide an enzyme detergency benefit, since they specifically improve stain removal in the cleaning process they are applied in as compared to the
20 same process without enzymes. Stain removing enzymes that are known in the art include enzymes such as carbohydrases, amylases, proteases, lipases, cellulases, hemicellulases, xylanases, cutinases, and pectinase.

In one aspect, the present invention concerns the use of an isolated polypeptide having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 60%, at least 61% at least
25 62% at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68% at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79% at least 80% at least 81% at least 82% at least 83% at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89% at least 90%, at
30 least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% in detergent compositions and cleaning processes, such as laundry and hard surface cleaning. Thus, in one aspect, the present invention demonstrates the detergency effect of a variety of exemplary proteases of the invention on various stains and under various conditions. In a particular aspect of the invention the detergent composition and the use in cleaning process concerns the use of a protease of the invention together with at least one of the
35 above mentioned stain removal enzymes.

In a preferred aspect of the present invention, the protease of the invention useful according to the invention may be combined with at least two enzymes. These additional enzymes are described in details in the section "other enzymes" more preferred at least three, four or five enzymes. Preferably, the enzymes have different substrate specificity, e.g., carbolytic activity, proteolytic activity, amylolytic activity, lipolytic activity, hemicellulytic activity or pectolytic activity. The enzyme combination may for example be a protease of the invention with another stain removing enzyme, e.g., a protease of the inveniton and an amylase, a protease of the invention and a cellulase, a protease of the invention and a hemicellulase, a protease of the invention and a lipase, a protease of the invention and a cutinase, a protease of the invention and a pectinase or a protease of the invention and an anti-redeposition enzyme. More preferably, the protease of the invention is combined with at least two other stain removing enzymes, e.g., a protease of the invention, a lipase and an amylase; or a protease of the invention, an amylase and a pectinase; or a protease of the invention, an amylase and a cutinase; or a protease of the invention, an amylase and a cellulase; or a protease of the invention, an amylase and a hemicellulase; or a protease of the invention, a lipase and a pectinase; or a protease of the invention, a lipase and a cutinase; or a protease of the invention, a lipase and a cellulase; or a protease of the invention, a lipase and a hemicellulase. Even more preferably, a protease of the invention may be combined with at least three other stain removing enzymes, e.g., a protease of the invention, an amylase, a lipase and a pectinase; or a protease of the invention, an amylase, a lipase and a cutinase; or a protease of the invention, an amylase, a lipase and a cellulase; or a protease of the invention, an amylase, a lipase and a hemicellulase. A protease of the invention may be combined with any of the enzymes selected from the non-exhaustive list comprising: carbohydrases, such as an amylase, a hemicellulase, a pectinase, a cellulase, a xanthanase or a pullulanase, a peptidase, other proteases or a lipase.

In another embodiment of the present invention, a protease of the invention may be combined with one or more metalloproteases, such as a M4 Metalloprotease, including Neutrase™ or Thermolysin. Such combinations may further comprise combinations of the other detergent enzymes as outlined above.

The cleaning process or the textile care process may for example be a laundry process, a dishwashing process or cleaning of hard surfaces such as bathroom tiles, floors, table tops, drains, sinks and washbasins. Laundry processes can for example be household laundering, but it may also be industrial laundering. Furthermore, the invention relates to a process for laundering of fabrics and/or garments where the process comprises treating fabrics with a washing solution containing a detergent composition, and at least one protease of the invention. The cleaning process or a textile care process can for example be carried out in a machine washing process or

in a manual washing process. The washing solution can for example be an aqueous washing solution containing a detergent composition.

The fabrics and/or garments subjected to a washing, cleaning or textile care process of the present invention may be conventional washable laundry, for example household laundry. Preferably, the major part of the laundry is garments and fabrics, including knits, woven, denims, non-woven, felts, yarns, and towelling. The fabrics may be cellulose based such as natural cellulosics, including cotton, flax, linen, jute, ramie, sisal or coir or manmade cellulosics (e.g., originating from wood pulp) including viscose/rayon, ramie, cellulose acetate fibers (tricell), lyocell or blends thereof. The fabrics may also be non-cellulose based such as natural polyamides including wool, camel, cashmere, mohair, rabbit and silk or synthetic polymer such as nylon, aramid, polyester, acrylic, polypropylen and spandex/elastane, or blends thereof as well as blend of cellulose based and non-cellulose based fibers. Examples of blends are blends of cotton and/or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g., polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers (e.g., rayon/viscose, ramie, flax, linen, jute, cellulose acetate fibers, lyocell).

The last few years there has been an increasing interest in replacing components in detergents, which is derived from petrochemicals with renewable biological components such as enzymes and polypeptides without compromising the wash performance. When the components of detergent compositions change new enzyme activities or new enzymes having alternative and/or improved properties compared to the common used detergent enzymes such as proteases, lipases and amylases is needed to achieve a similar or improved wash performance when compared to the traditional detergent compositions.

The invention further concerns the use of proteases of the invention a proteinaceous stain removing processes. The proteinaceous stains may be stains such as food stains, e.g., baby food, sebum, cocoa, egg, blood, milk, ink, grass, or a combination hereof.

Typical detergent compositions includes various components in addition to the enzymes, these components have different effects, some components like the surfactants lower the surface tension in the detergent, which allows the stain being cleaned to be lifted and dispersed and then washed away, other components like bleach systems removes discolor often by oxidation and many bleaches also have strong bactericidal properties, and are used for disinfecting and sterilizing. Yet other components like builder and chelator softens, e.g., the wash water by removing the metal ions from the liquid.

In a particular embodiment, the invention concerns the use of a composition comprising a protease of the invention, wherein said enzyme composition further comprises at least one or more

of the following a surfactant, a builder, a chelator or chelating agent, bleach system or bleach component in laundry or dish wash.

In a preferred embodiment of the invention, the amount of a surfactant, a builder, a chelator or chelating agent, bleach system and/or bleach component are reduced compared to amount of
5 surfactant, builder, chelator or chelating agent, bleach system and/or bleach component used without the added protease of the invention. Preferably the at least one component which is a surfactant, a builder, a chelator or chelating agent, bleach system and/or bleach component is present in an amount that is 1% less, such as 2% less, such as 3% less, such as 4% less, such as 5% less, such as 6% less, such as 7% less, such as 8% less, such as 9% less, such as 10% less,
10 such as 15% less, such as 20% less, such as 25% less, such as 30% less, such as 35% less, such as 40% less, such as 45% less, such as 50% less than the amount of the component in the system without the addition of protease of the invention, such as a conventional amount of such component. In one aspect, the protease of the invention is used in detergent compositions wherein said composition is free of at least one component which is a surfactant, a builder, a chelator or
15 chelating agent, bleach system or bleach component and/or polymer.

Washing method

The detergent compositions comprising a protease of the present invention are ideally suited for use in laundry applications. Accordingly, the present invention includes a method for
20 laundering a fabric. The method comprises the steps of contacting a fabric to be laundered with a cleaning laundry solution comprising the detergent composition according to the invention. The fabric may comprise any fabric capable of being laundered in normal consumer use conditions. The solution preferably has a pH of from about 5.5 to about 9. The compositions may be employed at concentrations of from about 100 ppm, preferably 500 ppm to about 15,000 ppm in solution. The
25 water temperatures typically range from about 5°C to about 90°C, including about 10°C, about 15°C, about 20°C, about 25°C, about 30°C, about 35°C, about 40°C, about 45°C, about 50°C, about 55°C, about 60°C, about 65°C, about 70°C, about 75°C, about 80°C, about 85°C and about 90°C. The water to fabric ratio is typically from about 1:1 to about 30:1 .

In particular embodiments, the washing method is conducted at a pH of from about 5.0 to
30 about 11.5, or in alternative embodiments, even from about 6 to about 10.5, such as about 5 to about 11, about 5 to about 10, about 5 to about 9, about 5 to about 8, about 5 to about 7, about 5.5 to about 11, about 5.5 to about 10, about 5.5 to about 9, about 5.5 to about 8, about 5.5. to about 7, about 6 to about 11, about 6 to about 10, about 6 to about 9, about 6 to to about 8, about 6 to about 7, about 6.5 to about 11, about 6.5 to about 10, about 6.5 to about 9, about 6.5 to about 8, about
35 6.5 to about 7, about 7 to about 11, about 7 to about 10, about 7 to about 9, or about 7 to about 8,

preferably about 5.5 to about 9, and more preferably about 6 to about 8.

In particular embodiments, the washing method is conducted at a degree of hardness of from about 0°dH to about 30°dH, such as about 1°dH, about 2°dH, about 3°dH, about 4°dH, about 5°dH, about 6°dH, about 7°dH, about 8°dH, about 9°dH, about 10°dH, about 11°dH, about 12°dH, about 13°dH, about 14°dH, about 15°dH, about 16°dH, about 17°dH, about 18°dH, about 19°dH, about 20°dH, about 21°dH, about 22°dH, about 23°dH, about 24°dH, about 25°dH, about 26°dH, about 27°dH, about 28°dH, about 29°dH, about 30°dH. Under typical European wash conditions, the degree of hardness is about 15°dH, under typical US wash conditions about 6°dH, and under typical Asian wash conditions, about 3°dH.

The present invention relates to a method of cleaning a fabric, a dishware or hard surface with a detergent composition comprising a protease of the invention.

A preferred embodiment concerns a method of cleaning, said method comprising the steps of: contacting an object with a cleaning composition comprising a protease of the invention under conditions suitable for cleaning said object. In a preferred embodiment the cleaning composition is a detergent composition and the process is a laundry or a dish wash process.

Still another embodiment relates to a method for removing stains from fabric which comprises contacting said a fabric with a composition comprising a protease of the invention under conditions suitable for cleaning said object.

In a preferred embodiment, the compositions for use in the methods above further comprises at least one additional enzyme as set forth in the "other enzymes" section above, such as an enzyme selected from the group consisting of carbohydrases, peptidases, proteases, lipases, cellulase, xylanases or cutinases or a combination hereof. In yet another preferred embodiment the compositions comprises a reduced amount of at least one or more of the following components a surfactant, a builder, a chelator or chelating agent, bleach system or bleach component or a polymer.

Also contemplated are compositions and methods of treating fabrics (e.g., to desize a textile) using one or more of the protease of the invention. The protease can be used in any fabric-treating method which is well known in the art (see, e.g., U.S. Patent No. 6,077,316). For example, in one aspect, the feel and appearance of a fabric is improved by a method comprising contacting the fabric with a protease in a solution. In one aspect, the fabric is treated with the solution under pressure.

In one embodiment, the protease is applied during or after the weaving of textiles, or during the desizing stage, or one or more additional fabric processing steps. During the weaving of textiles, the threads are exposed to considerable mechanical strain. Prior to weaving on mechanical looms, warp yarns are often coated with sizing starch or starch derivatives in order to increase their

tensile strength and to prevent breaking. The protease can be applied to remove these sizing protein or protein derivatives. After the textiles have been woven, a fabric can proceed to a desizing stage. This can be followed by one or more additional fabric processing steps. Desizing is the act of removing size from textiles. After weaving, the size coating should be removed before further processing the fabric in order to ensure a homogeneous and wash- proof result. Also provided is a method of desizing comprising enzymatic hydrolysis of the size by the action of an enzyme.

Low temperature uses

One embodiment of the invention concerns a method of doing laundry, dish wash or industrial cleaning comprising contacting a surface to be cleaned with a protease of the invention, and wherein said laundry, dish wash, industrial or institutional cleaning is performed at a temperature of about 40°C or below. One embodiment of the invention relates to the use of a protease of the invention in laundry, dish wash or a cleaning process wherein the temperature in laundry, dish wash, industrial cleaning is about 40°C or below.

In another embodiment, the invention concerns the use of a protease of the invention in a protein removing process, wherein the temperature in the protein removing process is about 40°C or below.

Thus, one aspect of the invention concerns the use of a polypeptide having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 60%, at least 61% at least 62% at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68% at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79% at least 80% at least 81% at least 82% at least 83% at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89% at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which have protease activity in laundry, dish wash or a cleaning process wherein the temperature in laundry, dish wash, industrial cleaning is about 40°C or below .

The present invention also relates to the use in laundry, dish wash or industrial cleaning process of a protease of the invention having at least one improved property compared to a subtilisin 309 protease with (SEQ ID NO 4) and wherein the temperature in laundry, dish wash or cleaning process is performed at a temperature of about 40°C or below.

In each of the above-identified methods and uses, the wash temperature is about 40°C or below, such as about 39°C or below, such as about 38°C or below, such as about 37°C or below, such as about 36°C or below, such as about 35°C or below, such as about 34°C or below, such as about 33°C or below, such as about 32°C or below, such as about 31°C or below, such as about 30°C or below, such as about 29°C or below, such as about 28°C or below, such as about 27°C or

below, such as about 26°C or below, such as about 25°C or below, such as about 24°C or below, such as about 23°C or below, such as about 22°C or below, such as about 21°C or below, such as about 20°C or below, such as about 19°C or below, such as about 18°C or below, such as about 17°C or below, such as about 16°C or below, such as about 15°C or below, such as about 14°C or below, such as about 13°C or below, such as about 12°C or below, such as about 11°C or below, such as about 10°C or below, such as about 9°C or below, such as about 8°C or below, such as about 7°C or below, such as about 6°C or below, such as about 5°C or below, such as about 4°C or below, such as about 3°C or below, such as about 2°C or below, such as about 1°C or below.

In another preferred embodiment, the wash temperature is in the range of about 5-40°C, such as about 5-30°C, about 5-20°C, about 5-10°C, about 10-40°C, about 10-30°C, about 10-20°C, about 15-40°C, about 15-30°C, about 15-20°C, about 20-40°C, about 20-30°C, about 25-40°C, about 25-30°C, or about 30-40°C. In a particular preferred embodiment the wash temperature is about 30°C.

In particular embodiments, the low temperature washing method is conducted at a pH of from about 5.0 to about 11.5, or in alternative embodiments, even from about 6 to about 10.5, such as about 5 to about 11, about 5 to about 10, about 5 to about 9, about 5 to about 8, about 5 to about 7, about 5.5 to about 11, about 5.5 to about 10, about 5.5 to about 9, about 5.5 to about 8, about 5.5 to about 7, about 6 to about 11, about 6 to about 10, about 6 to about 9, about 6 to about 8, about 6 to about 7, about 6.5 to about 11, about 6.5 to about 10, about 6.5 to about 9, about 6.5 to about 8, about 6.5 to about 7, about 7 to about 11, about 7 to about 10, about 7 to about 9, or about 7 to about 8, preferably about 5.5 to about 9, and more preferably about 6 to about 9.

In particular embodiments, the low temperature washing method is conducted at a degree of hardness of from about 0°dH to about 30°dH, such as about 1°dH, about 2°dH, about 3°dH, about 4°dH, about 5°dH, about 6°dH, about 7°dH, about 8°dH, about 9°dH, about 10°dH, about 11°dH, about 12°dH, about 13°dH, about 14°dH, about 15°dH, about 16°dH, about 17°dH, about 18°dH, about 19°dH, about 20°dH, about 21°dH, about 22°dH, about 23°dH, about 24°dH, about 25°dH, about 26°dH, about 27°dH, about 28°dH, about 29°dH, about 30°dH. Under typical European wash conditions, the degree of hardness is about 15°dH, under typical US wash conditions about 6°dH, and under typical Asian wash conditions, about 3°dH.

Use in removing egg stains

Another particular embodiment of the invention concerns removal of egg stains. These types of stain are often very difficult to remove completely. Egg stains are particularly problematic in hard surface cleaning such as dish wash where the stains often remain on the plates and cutlery after washing. The metalloproteases of the invention are particularly suitable for removing egg

stains.

Thus, the invention further concerns methods for removing egg stains from textiles, fabrics and/or hard surfaces like dishes and cutlery in particular from fabrics and textiles. A preferred aspect of the invention concerns a method of removing egg stains from textiles and/or fabrics
5 comprising contacting a surface in need of removal of an egg stain with a metalloprotease of the invention. In one embodiment, the invention comprises a method of removing egg stains from textiles and/or fabrics comprising contacting a surface in need of removal of an egg stain with a detergent composition comprising a metalloprotease of the invention. The invention also concerns a method of removing egg stains comprising adding a metalloprotease of the invention to a laundry
10 and/or washing process wherein said textiles and/or fabric comprises various egg stains.

One embodiment of the present invention relates to a method for removal of egg stains from a hard surface or from laundry, the method comprising contacting the egg stain-containing hard surface or the egg stain-containing laundry with a cleaning or detergent composition, preferably a laundry or dish wash composition, containing a metalloprotease of the invention.

15 Another embodiment relates a method for removing egg stains from fabric or textile which comprises contacting the fabric or textile with a cleaning or detergent composition, preferably a laundry or dish wash composition, comprising a metalloprotease of the invention.

A still further embodiment relates to a method for removing egg stains from fabric or textile which comprises contacting said a fabric or textile with a composition comprising a metalloprotease
20 of the invention, wherein said composition further comprises at least one additional enzyme as set forth in the "other enzymes" section above, such as an enzyme selected from the group consisting of a carbohydrase, a peptidase, a protease, a lipase, a cellulase, a xylanase, a cutinase or a combination hereof.

In particular embodiments, the egg removing method is conducted at a pH of from about 5.0
25 to about 11.5, or in alternative embodiments, even from about 6 to about 10.5, such as about 5 to about 11, about 5 to about 10, about 5 to about 9, about 5 to about 8, about 5 to about 7, about 5.5 to about 11, about 5.5 to about 10, about 5.5 to about 9, about 5.5 to about 8, about 5.5 to about 7, about 6 to about 11, about 6 to about 10, about 6 to about 9, about 6 to to about 8, about 6 to about 7, about 6.5 to about 11, about 6.5 to about 10, about 6.5 to about 9, about 6.5 to about 8, about
30 6.5 to about 7, about 7 to about 11, about 7 to about 10, about 7 to about 9, or about 7 to about 8, about 5.5 to about 9, preferably about 6 to about 8.5, and more preferably about 6 to about 8.

In particular embodiments, the egg removing method is conducted at a degree of hardness of from about 0°dH to about 30°dH, such as about 1°dH, about 2°dH, about 3°dH, about 4°dH, about 5°dH, about 6°dH, about 7°dH, about 8°dH, about 9°dH, about 10°dH, about 11°dH, about
35 12°dH, about 13°dH, about 14°dH, about 15°dH, about 16°dH, about 17°dH, about 18°dH, about

19°dH, about 20°dH, about 21°dH, about 22°dH, about 23°dH, about 24°dH, about 25°dH, about 26°dH, about 27°dH, about 28°dH, about 29°dH, about 30°dH. Under typical European wash conditions, the degree of hardness is about 15°dH, under typical US wash conditions about 6°dH, and under typical Asian wash conditions, about 3°dH.

5

EXAMPLES

Materials and methods

Composition of model detergents

10 Wash performance of proteases was investigated in liquid detergents and powder detergents, which had the following compositions:

Table 1: Composition of the model detergents used in the wash experiments (amounts are stated in wt%)	
Laundry liquid model detergent LF	Natrium hydroxide 99 %: 2,95 % Sulfonic acid: 11,52 % Soy fatty acid: 5,5 % Propylenglycol: 5,05 % C13-alkoholethoxylat, 8 EO: 9,45 % Phosphonat, Dequest 2066: 1,00 % Triethanolamin: 2,0 % Coco fatty acid: 4,5 % Natrium citrate, dihydrate: 1,0 % EtOH: 4,63 % Opacifier: 0,12 % Water to 100%
Laundry liquid model detergent J	LAS 5.15% AS 5.00% AEOS 14.18% Coco fatty acid 1.00% AEO 5.00% MEA 0.30% MPG 3.00 EtOH 1.50% DTPA (Na salt) 0.25%

	<p>Sodium citrate 4.00%</p> <p>Sodium formate 1.00%</p> <p>Sodium hydroxide 0.66%</p> <p>Water 58.95%</p>
Laundry liquid model detergent B	<p>Water 30.63%</p> <p>Sodium hydroxide 2.95%</p> <p>Dodecylbenzensulfonic acid 11.52%</p> <p>Fatty acids (Soya) 5.50%</p> <p>Propane-1,2-diol (MPG) 5.05%</p> <p>Water 17.38%</p> <p>C13-alcohol ethoxylate, 10.50%</p> <p>Diethylenetriaminepentakis (methylenephosphonic acid) (DTMPA) 3.08%</p> <p>Triethanolamine (TEA) 2.22%</p> <p>Fatty acids (Coco) 4.50%</p> <p>Sodium citrate monohydrate 1.00%</p> <p>Ethanol 4.63%</p> <p>Syntran 5909 (opacifier) 0.30%</p> <p>Perfume 0.35%</p>
Laundry powder model detergent A	<p>Sodium citrate dihydrate 32.3%</p> <p>Sodium-LAS 24.2%</p> <p>Sodium lauryl sulfate 32.2%</p> <p>Neodol 25-7 (alcohol ethoxylate) 6.4%</p> <p>Sodium sulfate 4.9%</p>
Laundry powder model detergent TB	<p>LAS, sodium salt 10.0%</p> <p>AS, sodium salt 1.8%</p> <p>Soap, sodium salt 2.0%</p> <p>AEO 3.0%</p> <p>Soda ash 14.9%</p> <p>Hydrous sodium silicate 2.5%</p> <p>Zeolite A 15.0%</p> <p>HEDP-Na4 0.13%</p> <p>Sodium citrate 2.0%</p> <p>PCA, copoly(acrylic acid/maleic acid), sodium salt 1.5%</p>

	SRP 0.5%
	Sodium sulfate 13.4%
	Sodium percarbonate 20.0%
	TAED 3.0%
	Foam regulator 1.0%

Table 2: Composition of the model detergents used in the wash experiments (amounts in wt%)

Component name	Model X / kg	Model Y / kg
Na-LAS granules (Marlon ARL)	8,25	4,95
Zeolite 4A (Silkem)	7,5	7,5
Sodium disilicate (Britesil H 265 LC)	6	6
Sodium carbonate (dense)	10	10
Sokalan CP 5 Gran.	0,5	0,5
Sodium sulfate - granulate	17,75	21,05
Total	50	50

2% AEO was added to both Det X and Det Y

	Liquid detergents				Powder detergents			
	Det A	Det B	Det J	Det LF	Det T	Det TB	Det X	DET Y
Dosage (g/L)	3.3	3.3	0.78	8	5.23	5.23	1.75	1.75

5

Test materials

Wash performance of proteases was investigated on the following test materials:

C-05 (Blood/milk/ink on cotton)

10 C-10 (Oil/milk/pigment on cotton)

CS-37 (Full egg with pigment non-aged on cotton)

EMPA 117 (Blood/milk/ink on cotton, extra heat treated)

EMPA 164 (Grass on cotton)

PC-03 (Chocolate-milk/ink on cotton/polyester)

15 PC-05 (Blood/milk/ink on cotton/polyester)

Test materials are obtained from Center for Testmaterials BV, P.O. Box 120, 3133 KT Vlaardingen, the Netherlands and EMPA Testmaterials AG, Movenstrasse 12, CH-9015 St. Gallen, Switzerland.

20 **Wash assays**

Automatic Mechanical Stress Assay (AMSA) for laundry

In order to assess the wash performance in laundry washing experiments were performed, using the Automatic Mechanical Stress Assay (AMSA). With the AMSA, the wash performance of a large quantity of small volume enzyme-detergent solutions can be examined. The AMSA plate has a number of slots for test solutions and a lid firmly squeezing the laundry sample, the textile to be washed against all the slot openings. During the washing time, the plate, test solutions, textile and lid were vigorously shaken to bring the test solution in contact with the textile and apply mechanical stress in a regular, periodic oscillating manner. For further description see WO02/42740 especially the paragraph "Special method embodiments" at page 23-24.

The wash performance was measured as the brightness of the colour of the textile washed. Brightness can also be expressed as the intensity of the light reflected from the sample when illuminated with white light. When the sample is stained the intensity of the reflected light is lower, than that of a clean sample. Therefore the intensity of the reflected light can be used to measure wash performance.

Colour measurements were made with a professional flatbed scanner (Kodak iQsmart, Kodak, Midtager 29, DK-2605 Brandby, Denmark), which is used to capture an image of the washed textile.

To extract a value for the light intensity from the scanned images, 24-bit pixel values from the image were converted into values for red, green and blue (RGB). The intensity value (Int) was calculated by adding the RGB values together as vectors and then taking the length of the resulting vector:

$$Int = \sqrt{r^2 + g^2 + b^2}$$

Terg-O-tometer (TOM) wash assay for laundry

The Terg-O-tometer (TOM) is a medium scale model wash system that can be applied to test 16 different wash conditions simultaneously. A TOM is basically a large temperature controlled water bath with up to 16 open metal beakers submerged into it. Each beaker constitutes one small top loader style washing machine and during an experiment, each of them will contain a solution of a specific detergent/enzyme system and the soiled and unsoiled fabrics where on which performance is tested. Mechanical stress is achieved by a rotating stirring arm, which stirs the liquid within each beaker with 120rpm.

In a TOM experiment, factors such as the ballast to soil ratio and the fabric to wash liquor ratio can be varied. Therefore, the TOM provides the link between small scale experiments, such

as AMSA, and the more time consuming full scale wash experiments in full scale washing machines.

After wash swatches are rinsed for 5 min under tap water and then dried overnight on a tray (12h-20h) at room temperature (around 20-25C) and 40% humidity after wash. The next day
5 swatches are measured at 460 nm with a Macbeth Color-Eye 7000 spectrophotometer.

Protease assays

1) Suc-AAPF-pNA assay:

pNA substrate : Suc-AAPF-pNA (Bachem L-1400).

10 Temperature : Room temperature (25°C)

Assay buffers : 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS,
1mM CaCl₂, 150mM KCl, 0.01% Triton X-100 adjusted to pH-values 2.0,
3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, 10.0, and 11.0 with HCl or NaOH.

15 20 μ I protease (diluted in 0.01% Triton X-100) was mixed with 100 μ I assay buffer. The assay was started by adding 100 μ I pNA substrate (50mg dissolved in 1.0ml DMSO and further diluted 45x with 0.01% Triton X-100). The increase in OD₄₀₅ was monitored as a measure of the protease activity.

20 2) Protazyme AK assay:

Substrate : Protazyme AK tablet (cross-linked and dyed casein; from Megazyme)

Temperature : controlled (assay temperature).

Assay buffers : 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS,
1mM CaCl₂, 150mM KCl, 0.01% Triton X-100, pH 7.0.

25

A Protazyme AK tablet was suspended in 2.0ml 0.01% Triton X-100 by gentle stirring. 500 μ I of this suspension and 500 μ I assay buffer were dispensed in an Eppendorf tube and placed on ice. 20 μ I protease sample (diluted in 0.01% Triton X-100) was added. The assay was initiated by transferring the Eppendorf tube to an Eppendorf thermomixer, which was set to the assay
30 temperature. The tube was incubated for 15 minutes on the Eppendorf thermomixer at its highest shaking rate (1400 rpm.). The incubation was stopped by transferring the tube back to the ice bath. Then the tube was centrifuged in an ice cold centrifuge for a few minutes and 200 μ I supernatant was transferred to a microtiter plate. OD₆₅₀ was read as a measure of protease activity. A buffer blind was included in the assay (instead of enzyme).

3) Suc-AAPX-pNA assay:

pNA substrates : Suc-AAPA-pNA (Bachem L-1 775)
 Suc-AAPR-pNA (Bachem L-1 720)
 5 Suc-AAPD-pNA (Bachem L-1 835)
 Suc-AAPI-pNA (Bachem L-1 790)
 Suc-AAPM-pNA (Bachem L-1 395)
 Suc-AAPV-pNA (Bachem L-1 770)
 Suc-AAPL-pNA (Bachem L-1 390)
 10 Suc-AAPE-pNA (Bachem L-1 710)
 Suc-AAPK-pNA (Bachem L-1 725)
 Suc-AAPF-pNA (Bachem L-1400)
 Temperature : Room temperature (25°C)
 Assay buffer : 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS,
 15 1mM CaCl₂, 150mM KCl, 0.01% Triton X-100, pH 9.0.

20μl protease (diluted in 0.01% Triton X-100) was mixed with 100μl assay buffer. The assay was started by adding 100μl pNA substrate (50mg dissolved in 1.0ml DMSO and further diluted 45x with 0.01% Triton X-100). The increase in OD₄₀₅ was monitored as a measure of the protease activity.

Soybean-Maize Meal Assay (SMM Assay)**Protease activity on soybean-maize meal at pH 3, 4, 5, 6, and 7**

An end-point assay using soybean-maize meal as substrate was used for obtaining the activity profile of the proteases at pH 3-7.

Assay buffers: 100 mM succinic acid, 100 mM HEPES, 100 mM CHES, 100 mM CAPS, 1 mM CaCl₂, 150 mM KCl, 0.01% Triton X-100 adjusted using HCl or NaOH to give pH-values 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, 10.0 and 11.0 when mixing 10 ml assay buffer with 1 g soybean-maize meal (30:70 ratio).

2 mL soybean-maize meal slurry is mixed for 30 min before protease addition and incubation for 3 hours at 40°C (500 rpm). Protease is added via 100 μl 100 mM sodium acetate (NaAc) buffer (9.565 g/l NaAc, 1.75 g/l acetic acid, 5 mM CaCl₂, 0.01% BSA, 0.01% Tween20, pH 6.0). Supernatants are collected after centrifugation (10 min, 4000 rpm, 0°C) and protein activity is determined using a colorimetric assay based on the o-Phthaldialdehyde (OPA) method essentially according to Nielsen et al. (Nielsen, PM, Petersen, D, Dampmann, C. Improved method for

determining food protein degree of hydrolysis. J Food Sci, 2001 , 66: 642-646). This assay detects free a-amino groups and hence protease activity can be measured as an increase in absorbance. First 500 μ I of each supernatant is filtered through a 100 kDa Microcon filter by centrifugation (60 min, 11,000 rpm, 5°C). The samples are diluted 10x in deionized water and 25 μ I of each sample is loaded into a 96 well microtiter plate (5 replicates). Finally 200 μ I OPA reagent is dispensed into all wells and the plate is shaken (10 sec, 750 rpm) and absorbance measured at 340 nm. The level of protease activity is calculated as the difference between absorbance in the enzyme treated sample and the blank sample and expressed as 'Abs x dilution factor'.

10 EXAMPLES

Example 1. Cloning and expression of a S8 protease from *Bacillus* sp-13231

The protease with the nucleotide sequence given in SEQ ID: 1 was derived from a bacterial strain isolated from a soil sample collected in USA. Chromosomal DNA from a pure culture was purified and subjected to full genome sequencing using Illumina technology. The assembled genome sequence and subsequent analysis of the 16S ribosomal subunit gene sequences against the ARB-SILVA database (The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Quast et al. Nucl. Acids Res. 41 (2013): D590-D596.) confirmed the identity of the strain as new species within the genus *Bacillus* which was named *Bacillus* sp-13231 .

20 The gene encoding the protease was amplified by PCR and fused with regulatory elements and homology regions for recombination into the *B. subtilis* genome.

The linear integration construct was a SOE-PCR fusion product (*Horton, R.M., Hunt, H.D., Ho, S.N., Pullen, J.K. and Pease, L.R. (1989) Engineering hybrid genes without the use of restriction enzymes, gene splicing by overlap extension Gene 77: 61-68*) made by fusion of the gene between two *Bacillus subtilis* chromosomal regions along with strong promoters and a chloramphenicol resistance marker. The SOE PCR method is also described in patent application WO 2003095658.

The gene was expressed under the control of a triple promoter system (as described in WO 99/43835), consisting of the promoters from *Bacillus licheniformis* alpha-amylase gene (amyL), *Bacillus amyloliquefaciens* alpha-amylase gene (amyQ), and the *Bacillus thuringiensis* cryIIIA promoter including stabilizing sequence.

The gene was expressed with a *Bacillus clausii* secretion signal (encoding the following amino acid sequence: MKKPLGKIVASTALLISVAFSSSIASA) replacing the native secretion signal.

The SOE-PCR product was transformed into *Bacillus subtilis* and integrated in the

chromosome by homologous recombination into the pectate lyase locus. Subsequently a recombinant *Bacillus subtilis* clone containing the integrated expression construct was grown in liquid culture. The culture broth was centrifuged (20000 x g, 20 min) and the supernatant was carefully decanted from the precipitate and used for purification of the enzyme.

5 The N-terminal AQSVPWG was determined by Edman degradation resulting in a mature polypeptide of 269 amino acids.

Example 2. Determination of Molecular Weight:

Intact molecular weight analyses of the purified enzyme was performed using a MAXIS 11
 10 elec-trospray mass spectrometer (Bruker Daltonik GmbH, Bremen, DE). The samples were diluted to 0,1 mg/ml in MQ water. The diluted sample was applied to an Aeris Widepore C4 column (Phenom-enex). The sample was washed and eluted from the column running an acetonitrile linear gradient and introduced to the electrospray source with a flow of 300 ml/min by an Ultimate 3000
 15 LC system (Dionex). Data analysis is performed with DataAnalysis version 4.2 (Bruker Daltonik GmbH, Bremen, DE). The molecular weight of the sample was calculated by Maximum Entropy deconvolution of the raw data in the range 10.000 to 40.000 Da. The dominant peak observed at molecular weight of 28873.08 Da corresponds to within 0.3 Da of the calculated molecular weight of the mature protease in SEQ ID: 2.

20 Example 3: Wash performance in laundry - Terg-O-tometer (TOM)

In order to assess the wash performance in laundry, washing experiments are performed using the Terg-O-tometer (TOM) assay. The TOM experiments are conducted under the experimental conditions specified below:

Test solution volume	1000 ml
pH	As is
Wash time	20 minutes (Det X), 30 minutes (Det T & A) and 12 minutes (Det J)
Temperature	40°C and 20°C
Water hardness	9°dH - Ca ²⁺ /Mg ²⁺ 7HCO ₃ ⁻ 2:1 :4.5 (Det X) 15°dH - Ca ²⁺ /Mg ²⁺ 7HCO ₃ ⁻ 4:1 :7.5 (Det T)
Protease dosage	10 - 30 nM and 60 nM
Test materials	EMPA 117, C-05, PC-03, CS-37, EMPA 164 and C-10 (2 swatches with 5x5 cm each)
Ballast	Cotton swatches (5x5 cm) from WFK80A and WFK10A in a ratio of

	50/50 are added on top of the swatches to give a total weight of 30g
--	--

The protease shown was tested in TOM wash as described above. The test was performed in a liquid and a powder detergent on several different stains at both 20°C and 40°C and at different protease dosages. Wash performance was compared with a well-known protease for detergent, Savinase (SEQ ID NO 4), a protease derived from *Bacillus clausii*. The results are shown in table 4 and 5.

Table 4: Delta remission of *Bacillus* sp-13231 and Savinase relative to the liquid detergent without protease.

Stain	<i>Bacillus</i> sp-13231			Savinase		
	Det J, 30 nM	Det A, 60 nM		Det J, 30 nM	Det A, 60 nM	
	30°C	20°C	40°C	30°C	20°C	40°C
EMPA 117	4.8	11.9	25.8	4.7	11.6	23.5
C-05	4.2	7.3	14.0	2.7	8.8	15.3
PC-03	3.0	9.5	22.3	3.3	8.4	18.8
CS-37	6.5	6.5	10.5	5.4	9.6	11.0
EMPA164	2.0	6.3	8.6	0.8	4.6	8.5
C-10	3.8	10.3	20.8	3.8	8.3	21.1

The results show that the *Bacillus* sp-13231 S8 protease has good wash performance in liquid detergents on all investigated stains at different temperatures. Especially in Det J and to a certain extent also at 20°C the *Bacillus* sp-13231 S8 protease shows a superior wash performance to the known laundry protease Savinase.

Table 5: Delta intensities of *Bacillus* sp-13231 and Savinase relative to the powder detergents without protease.

Stain	<i>Bacillus</i> sp-13231	Savinase
-------	--------------------------	----------

	Det X, 20°C		Det T, 60 nM		Det X, 20°C		Det T, 60 nM	
	10 nM	30 nM	20°C	40°C	10 nM	30 nM	20°C	40°C
EMPA 117	12.1	1.6	25.0	37.3	7.1	4.8	34.6	37.0
C-05	13.6	5.2	17.3	21.9	10.6	6.4	21.3	26.4
PC-03	9.6	2.0	12.3	28.5	5.6	2.6	22.4	27.4
CS-37	8.7	2.7	6.6	9.9	4.6	9.3	9.8	13.0
EMPA164	2.1	7.3	7.7	13.7	1.0	2.3	9.6	12.7
C-10			15.3	24.6			20.4	24.2

The results show that the *Bacillus sp-13231* S8 protease has good wash performance on all investigated stains at both 20°C and 40°C. Especially in Detergent X at low dosage the *Bacillus sp-13231* S8 protease shows a distinct superior wash performance to the known laundry protease Savinase.

10

15

CLAIMS

1. An isolated polypeptide having protease activity, selected from the group consisting of:
- 5 (a) a polypeptide having at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 2; and
- (b) a polypeptide encoded by a polynucleotide having at least 85%, at least 86%, at
- 10 least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1.
2. The polypeptide of claim 1 comprising or consisting of SEQ ID NO: 2 or the mature
- 15 polypeptide of SEQ ID NO: 2.
3. The polypeptide of claim 2, wherein the mature polypeptide is amino acids 1 to 269 of SEQ ID NO: 2.
- 20 4. The polypeptide of any of claims 1 to 3, which is a variant of the mature polypeptide of SEQ ID NO: 2 comprising a substitution, deletion, and/or insertion at one or more positions.
5. A composition comprising the polypeptide of any of claims 1 to 4.
- 25 6. The composition of claim 5 being a detergent composition such as a composition for laundry or automatic dish washing.
7. The composition of any of claims 5 to 7 further comprising one of more additional enzymes selected among, proteases, amylases, lipases, cutinases, cellulases, endoglucanases,
- 30 xyloglucanases, pectinases, pectin lyases, xanthanases, peroxidaes, haloperoxygenases, catalases, mannanases, or any mixture thereof.
8. The composition of any of claims 5 to 7 comprising one or more components selected from the group consisting of surfactants, builders, chelators or chelating agents, bleach system or bleach
- 35 components, polymers, fabric conditioners, foam boosters, suds suppressors, dyes, perfume,

tannish inhibitors, optical brighteners, bactericides, fungicides, soil suspending agents, anti corrosion agents, enzyme inhibitors or stabilizers, enzyme activators, bluing agents and fluorescent dyes, antioxidants, polymers and solubilizers.

5 9. An isolated polynucleotide encoding the polypeptide of any of claims 1 to 4.

10. A nucleic acid construct or expression vector comprising the polynucleotide of claim 9 operably linked to one or more control sequences that direct the production of the polypeptide in an expression host.

10

11. A recombinant host cell comprising the polynucleotide of claim 9 operably linked to one or more control sequences that direct the production of the polypeptide.

12. A method of producing the polypeptide of any of claims 1 to 4 comprising:

- 15 (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and
(b) recovering the polypeptide.

13. A method of producing a polypeptide having protease activity comprising:

- 20 (a) cultivating the host cell of claim 11 under conditions conducive for production of the polypeptide; and
(b) recovering the polypeptide.

14. A method for the treatment of proteins, comprising the step of adding at least one protease
25 of any one of claims 1 to 4 to at least one protein or protein source.

15. Use of at least one protease of any one of claims 1 to 4 in detergents or a composition according to any of claims 5 to 8 in a cleaning process.

30

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2016/078492

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N9/54 C11D3/386
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
C12N C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal , Sequence Search , BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE REfseq [Onl i ne] embl ; 18 October 2015 (2015-10-18) , "Bacill us sp. JCM 19035" , XP002766325 , Database accessi on no. WP_054794820 the whol e document -----	1,4,9-11
X	EP 1 160 327 A2 (GENENCOR INT [US]) 5 December 2001 (2001-12-05) cl aims ; f i g u r e 4 ; e x a m p l e s -----	1-15
X	W0 2015/089441 A1 (DANISCO US INC [US]) 18 June 2015 (2015-06-18) paragraph [00129] ; cl aims ; e x a m p l e 7 ; sequence 59 -----	1,4-15

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 25 January 2017	Date of mailing of the international search report 10/02/2017
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Le Cornec, Nadi ne
--	---

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2016/078492
--

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 1160327	A2	05-12-2001	AT 250669 T 15-10 -2003
			AU 629814 B2 15- 10- 1992
			AU 3050189 A 06- 09- 1989
			BG 60566 BI 28-08 - 1995
			BR 8905580 A 14-08 - 1990
			CA 1340366 C 02- 02- 1999
			CN 1036602 A 25-10 - 1989
			DE 68912359 DI 03- 03- 1994
			DE 68912359 T2 09-06 - 1994
			DE 68929484 DI 30- 10- 2003
			DE 68929484 T2 22-07 -2004
			DK 501289 A 07- 12- 1989
			EP 0328229 AI 16- 08- 1989
			EP 0571049 AI 24-11 - 1993
			EP 1160327 A2 05-12 -2001
			ES 2061929 T3 16-12 - 1994
			GR 3021934 T3 31- 03- 1997
			IE 63248 BI 05-04 - 1995
			IE 940164 L 11-08 - 1989
			JP 2849141 B2 20-01 - 1999
			JP H02503986 A 22-11 - 1990
			LT 3963 B 27-05 - 1996
			LV 10652 B 20-08 - 1995
			PT 89702 A 04- 10- 1989
			RU 2069695 CI 27-11 - 1996
			US 5336611 A 09-08 - 1994
			WO 8907642 AI 24-08 - 1989
WO 2015089441	AI	18-06-2015	EP 3080262 AI 19-10-2016
			US 2016312204 AI 27-10-2016
			WO 2015089441 AI 18-06-2015