



- (51) **International Patent Classification:**
C07K 1/14 (2006.01)
- (21) **International Application Number:**
PCT/SG2014/000232
- (22) **International Filing Date:**
28 May 2014 (28.05.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (71) **Applicant:** AGENCY FOR SCIENCE, TECHNOLOGY
AND RESEARCH [SG/SG]; 1 Fusionopolis Way, #20-10
Connexis, Singapore 138632 (SG).
- (72) **Inventor:** GAGNON, Peter Stanley; 20 Biopolis Way,
Centros #06-01, Singapore 138668 (SG).
- (74) **Agent:** SPRUSON & FERGUSON (ASIA) PTE LTD;
P.O. Box 1531, Robinson Road Post Office, Singapore
903031 (SG).
- (81) **Designated States** (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

- (84) **Designated States** (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

Published:

— *with international search report (Art. 21(3))*

(54) **Title:** VIRUS REDUCTION METHOD

(57) **Abstract:** Methods are provided for enhancing reduction of virus and viral DNA levels in protein preparations by contacting protein preparations with a surface bearing a primary amine, secondary amine, or both.



WO 2015/183180 A1

VIRUS REDUCTION METHOD

BACKGROUND

[0001] Embodiments disclosed herein relate to methods for reduction of viruses and viral polynucleotides in preparations containing a desired protein to be purified.

[0002] Some virus species are known to be inactivated by means of exposure to extreme conditions such as highly acidic pH. These include most species of lipid-enveloped viruses. Some virus species are known to be resistant to such inactivation. These include many non-lipid enveloped virus species, also referred to as protein coat viruses.

[0003] Anion exchange chromatography is commonly used for the purification of proteins, including removal of contaminating virus species. Anion exchangers used for such applications are frequently quaternary amine ion exchangers. These applications are performed at neutral to slightly alkaline pH because they generally support the strongest virus binding under those conditions (C. Curtis et al, Biotechnol. Bioeng. 84 (2003) 179-185), and such conditions are required to remove contaminating host cell proteins in parallel with removing viruses (P. Gagnon, J. Chromatogr. A 1221 (2012) 57-70. Primary amine-based anion exchangers have also been described for removal of virus and host cell protein at neutral to alkaline operating pH (W. Riordan et al., Biotechnol. Prog. 25 (2009) 1695-1702).

SUMMARY

[0004] In some embodiments, disclosed herein are methods for reducing virus and viral DNA levels in an acidic preparation of a desired protein, the method comprising contacting the acidic preparation of the desired protein with a surface bearing a primary amine, secondary amine, or both.

DETAILED DESCRIPTION

[0005] Methods are provided for enhanced removal of virus and viral DNA from a preparation containing a desired protein, during or following low pH treatment, by contacting a low pH preparation with a solid surface bearing primary and/or secondary amines. As used herein, "low pH" treatment refers to a sufficiently low pH as part of a virus inactivation treatment, *i.e.*, a sufficiently low pH to inactivate at least one virus. Thus, a top end of the pH range will typically not be greater than about 4.0. Thus, in some

5 embodiments, a low pH may range, for example, from about 1.0 to about 4.0, in some
embodiments, or from about 2.0 to about 4.0, or about from 2.0 to about 3.8. Those skilled in the art
will appreciate that the pH may be selected to conform to regulatory standards. In a typical case, the
upper end regulatory limit pH may be about 3.8. In some embodiments, there are provided methods
for reducing virus and viral DNA levels in an acidic preparation of a desired protein, the method
10 comprising contacting the acidic preparation of the desired protein with a surface bearing a primary
amine, secondary amine, or both.

[0006] In some embodiments, an operating pH during the contacting step is in a range selected
from the group consisting of (a) from about 2.0 to about 4.0, (b) from about 2.25 to about 3.75, (c)
from about 2.5 to about 3.5, (d) from about 2.75 to about 3.25, (e) from about 3.5 to about 3.75, and
15 (f) from about 3.25 to about 3.5 and intermediate ranges or intermediate individual values in
between.


[0007] In some embodiments, a salt concentration during the contacting step is range selected
from the group consisting of (a) from about 0.0, or a non-zero value, up to about 0.5 M, (b) from
about 0.05 to about 0.3 M, and (c) from about 0.1 to about 0.2 M. The salt concentration may be any
20 intermediate ranges or intermediate individual values in between these recited ranges.

[0008] In some embodiments, the duration of the contacting step is in a range selected from the
group consisting of (a) from about 15 to about 120 minutes, (b) from about 30 to about 60 minutes,
(c) a non-zero amount less than about 30 minutes, (d) a non-zero amount less than about 15
minutes, (e) a non-zero amount less than about 10 minutes, (f) a non-zero amount less than about 5
25 minutes, (g) a non-zero amount less than about 2 minutes, (h) a non-zero amount less than about 1
minute, and (i) a time required to pass the preparation through a device housing the primary amine-
bearing surface.

[0009] In some embodiments, the surface comprises one or more moieties selected from the
group consisting of ethylenediamine, tris(2-aminoethyl)amine, diethylaminetriamine,
30 triethylenetetraamine, tetraethylenepentamine, polyalkylamine. In some embodiments, the surface
may comprise any primary amine, secondary amine, or combinations thereof. The surface may be
derived from or take the form of any straight chain or branched primary monoamine fragment
attached at the surface of a substrate. Exemplary primary amine moieties include, without
limitation, fragments based on methylamine, ethylamine, propylamine, isopropylamine, butylamine,
35 sec-butylamine, iso-butylamine, any isomeric configuration of hexylamine, any isomeric

5 configuration of heptylamine, octylamines, any isomeric configuration of nonylamine, any isomeric configuration of decylamine, or mixtures or combinations thereof.

[0010] In some embodiments, diamine fragments may be employed. Suitable diamines include, without limitation, diamines of the general formula



10 wherein R is linear or branched alkyl or alkenyl groups having about 1 and about 20 carbon atoms, cycloalkenyl groups having about 1 and about 20 carbon atoms, alkylcycloalkenyl groups having about 1 and about 20 carbon atoms, alkarenyl group having between about 1 and about 20 carbon atoms, aralkenyl group having about 1 and about 20 carbon atoms, or the like or mixtures or combinations thereof. The R group can also include atoms other than carbon and hydrogen such as
15 oxygen, nitrogen, fluorine and/or chlorine.

[0011] Suitable surface amine moieties may be based on diamines and triamines including, without limitation, alkyl diamines, cycloalkyl diamines, alkacycloalkyl diamines, aralkyl diamines, aryl diamines, alkaryl diamines, and the like and analogs thereof and where one or more of the carbon atoms can be replaced with nitrogen atoms, oxygen atoms, or mixtures thereof where the
20 oxygen atoms form carboxy, hydroxy and or ether moieties and the nitrogen atoms form tertiary amine and/or amide moieties and/or one or more hydrogen atoms can be replaced with fluorine atoms, chlorine atoms or mixture thereof and including between 2 and about 20 carbon atoms, including, about 3 to about 15 carbon atoms and including, about 4 to about 10 carbon atoms. Exemplary alkyl diamines include, without limitation, 1,2-diaminoethane (1,2-ethylene diamine),
25 1,2-diaminopropane, 1,3-diaminopropane, 1,2-diaminobutane, 1,3-diaminobutane, 1,4-diaminobutane, 1,2-diaminopentane, 1,3-diaminopentane, 1,4-diaminopentane, 1,5-diaminopentane, and similar higher diaminoalkanes, hexamethylenediamine, aminomethylcyclopentylamine, 1,2-cyclopentanediamine, 1,6-hexanediamine, 1,2-diaminobenzene, lysine (or other diamine amino acids), 1,2-diaminobenzene, 1,4-diamine benzene, 1,2-diphenyl-1,2-ethane diamine, phenylene
30 diamine, 2-hydroxypropylene diamine, hydantoin, N,N-Bis(dihydroxyethyl)ethylenediamine, hexahydrotriazine, aminoethylpiperazine (AEP) or the like, or mixtures or combinations thereof. Examples of other aliphatic diamines and triamines that can be satisfactorily used in making the subject compositions include 1,4-diaminocyclohexane and bis-hexamethylenetriamine. In all of the
aforementioned embodiments, presentation of the amine on the surface of a substrate may be
35 achieved through connectivity to any atom of any of the fragments, including the nitrogen itself. In

5 some embodiments, a secondary amine may also be employed. Such secondary amines may comprise a further straight chain or branched C₁ to C₂₀ fragment.

[0012] In some embodiments, the surface is in the form of one selected from the group consisting of a particle, a membrane, a monolith, a hydrogel-coated skeletal framework, and combinations thereof.

10 [0013] In some embodiments, the contacting step is performed by an operation selected from the group consisting of (a) dispersing primary and/or secondary amine-bearing particles in the preparation, (b) passing the preparation through a device in fluid contact with the surface, (c) recirculating the preparation through a device in providing fluid contact with the surface, and (d) combinations thereof.

15 [0014] It has been unexpectedly discovered that virus binding to primary and/or secondary amines on a solid surface breaks the accepted rules of anion exchange chromatography. Binding strength, as measured by the concentration of salt required to elute bound virus or viral polynucleotides from the primary amine-bearing surface, increases as a function of decreasing pH, in some cases by more than a factor of 2 compared to binding on the same surface at pH 8, by
20 more than a factor of 5 compared to binding on quaternary exchangers at pH 8, and by nearly a factor of 10 compared to binding on quaternary exchangers at pH 6. Binding becomes progressively stronger at lower pH values so that at pH 3, up to 5 M guanidine is required to remove DNA from the primary and/or secondary amine-bearing surface. At the same time, experimental data show that retention of peptides and proteins with a mass lower than about 150
25 kDa is not enhanced compared to pH 8, thereby creating a window in which the desired protein in most preparations does not bind to the primary and/or secondary-amine-bearing surface under the conditions where retention of virus and viral DNA is enhanced. Experimental data show that this translates to a more than 700-fold reduction of non-lipid enveloped viruses such as Hepatitis A virus and Minute virus of Mice, which are not significantly inactivated by treatment with low pH.
30 This is unexpected because retention of solutes such as viruses, DNA, and proteins on quaternary anion exchangers diminishes with decreasing pH, as a result of the solutes becoming more electropositive (and/or less electronegative). This is generally understood to render anion exchange chromatography un-useful for virus removal at acidic pH, and especially at strongly acidic pH values, such as the conditions customarily used for virus inactivation. Without ascribing
35 to any particular theory, the contrary behavior of viruses and viral DNA on primary and/or

5 secondary-amine bearing surfaces may be mediated by hydrogen bonding. This is unexpected because hydrogen bonding is known to involve an electrostatic component that tends to suggest its contribution should parallel ion exchange behavior. Data from competition experiments with selective hydrogen donor-acceptors such as sorbitol nevertheless indicates hydrogen bonding. The behavior is further unexpected because it is distinct and contrary in multiple respects to results
10 known in the art, indicating that retention increases substantially at pH 6 compared to pH 8 for proteins ranging in size from 18 to 78 kDa; indicating that there is no size correlation of increased binding with decreasing pH; and indicating that retention becomes strongest at about pH 6, weakens at lower pH, and is lost at pH 4-5 (US Patent 6,090,288). Supporting the methods disclosed herein, experimental data indicate that there is substantially no enhancement of retention
15 for a protein of about 66 kDa, while there is a strong correlation between size and enhancement, and retention continues to become stronger at pH values below 6 for solutes with a size approaching about 1 MDa and greater, particularly at pH values less than 4, all of which are necessary enabling factors.

[0015] In some embodiments, the operating pH may be less than 6, such as pH 5 or less, or 4 or
20 less, or 3.9 or less, or 3.8 or less, or 3.7 or less, or 3.6 or less, or 3.5 or less, or 3.4 or less, or 3.3 or less, or 3.3 or less, or 3.2 or less, or 3.1 or less, or 3.0 or less, down to the lowest pH that can be tolerated by the desired protein without it sustaining permanent damage that may affect its clinical efficacy or safety. In some embodiments, the tolerable pH may be 2.5 or less, or 2.0 or less, or about 1.0. The lowest tolerable pH will be understood to be unique to each desired product, and
25 not possible to express on a broad generic basis. In some embodiments, the maximum pH will be no greater than about 4 since inactivation of even lipid-enveloped virus species requires a pH below 4. In some embodiments, the pH will be between 3.25 and 3.75.

[0016] In some embodiments, the preparation may be devoid of salts. In some embodiments, the preparation may contain one or more species of salts, wherein the concentration of such salts is a
30 non-zero amount up to 0.01 M, up to 0.05 M, up to 0.1 M, up to 0.2 M, up to 0.3 M, up to 0.4 M, up to 0.5 M, or higher, potentially up to 1.0 M. In some such embodiments, the species of salt may be sodium chloride (NaCl). In some embodiments, the presence of salt may have a negligible effect on virus removal by the primary amine-bearing surface, while in some embodiments the presence of salt may decrease the degree of virus removal achieved by the primary amine-bearing
35 surface. In some embodiments, the presence of salts may either enhance virus inactivation, or

- 5 improve stability of the desired protein, or decrease solubility of the desired proteins. Such effects can be determined for each species of desired protein. In some embodiments, increasing salt concentration within the indicated ranges may limit the degree to which virus removal is enhanced by the primary amine-bearing surface over the degree of inactivation achieved solely by treatment at low pH, but the degree of removal will remain significant.
- 10 **[0017]** In some embodiments, the preparation may contain elements intended to enhance virus inactivation, such as ethacridine, methylene blue, daunomycin, doxorubicin, chlorhexidine, alexidine, benzalkonium chloride, tris(2-aminoethyl)amine, tri(N-butyl)phosphate, a surfactant, a chelating agent, or combinations of such elements.
- 15 **[0018]** In some embodiments, the primary-amine-bearing surface may additionally include other species of electropositive chemical moieties, such as secondary amine groups. In other embodiments, the primary-amine-bearing surface may additionally include other species of electropositive chemical moieties, such as tertiary or quaternary amine groups. In some embodiments, the primary-amine bearing surface may additionally include more than one species of electropositive chemical moieties. In some embodiments, the primary-amine-bearing surface
- 20 may additionally include a combination of electropositive chemical moieties consisting of secondary and tertiary amines; secondary and quaternary amines; tertiary and quaternary amines; secondary, tertiary, and quaternary amines, in any relative proportions.
- 25 **[0019]** In some embodiments, the chemical species that contains the primary amine, may be tris(2-aminoethyl)amine (TREN). In other embodiments, the chemical species that contains the primary amine may be ethylenediamine. In other embodiments the chemical species that contains the primary amine may be polyallylamine or another polymeric primary amine.
- 30 **[0020]** In some embodiments, the primary-amine-bearing surface may include chemical elements that permit the surface as a whole to engage in electrostatic interactions, hydrogen bonds, hydrophobic interactions, pi-pi bonding, and metal coordination interactions.
- [0021]** In some embodiments, the primary-amine-bearing surface may include elements with a charge opposite to the primary amines. In some embodiments, the primary-amine-bearing surface may include elements that are electroneutral, either by virtue of being uncharged or by virtue of containing balanced proportions of negative and positive charges, or both.
- 35 **[0022]** In some embodiments where a plurality of surfaces are present, the respective surfaces may bear the same primary amine-containing species. In some embodiments where a plurality of

5 surfaces are present, the respective surfaces may bear different primary amine-containing species. In some embodiments where a plurality of surfaces are present, some of the respective surfaces may bear species other than primary amines.

[0023] In some embodiments, the primary-amine-bearing solid surface may be in the physical form of a particle, a membrane, a monolith, a hydrogel on a macro-reticulate skeleton. In some
10 such embodiments, more than a single physical unit may be present, such as a plurality of particles, or a plurality of membranes, or a plurality of other physical forms, or combinations of different physical forms. In some such embodiments, a plurality of particles may be packed in a column.

[0024] In some embodiments, a plurality of primary-amine surfaces may be suspended within the
15 preparation containing the desired protein. In some such embodiments, the suspended surfaces may be removed after treatment by sedimentation, or filtration, or flotation, or entrapment in a magnetic field, or entrapment on an oppositely charged surface, or any other expedient means. In some embodiments, the preparation containing the desired protein may be flowed through a device containing one or more primary-amine-bearing fluid-contact surfaces after low pH
20 treatment for virus inactivation has been nominally completed. In some embodiments, the desired protein preparation may be recycled continuously through a device containing one or more primary-amine fluid contact surfaces for a period of time up to the full duration of the low pH treatment. In some embodiments, the desired protein preparation may be contacted with one or more primary amine binding surfaces by any combination of the foregoing modes of contact.

[0025] In one embodiment that illustrates the key elements of the disclosed method, a preparation
25 containing a partially purified desired protein such as an IgG monoclonal antibody is titrated to a pH of 3.5 and held at that pH for 60 minutes. A monolith with a surface modified to bear ethylenediamine is equilibrated to nominally the same pH and salt conditions as the antibody preparation. The preparation is then flowed through the monolith. The monolith may optionally be rinsed with clean buffer at pH 3.5 to
30 maximize recovery of the desired protein. The monolith may then optionally be discarded or regenerated.

[0026] In another embodiment that illustrates the key elements of the disclosed method, a preparation containing a partially purified antibody is titrated to a pH of 3.5. A monolith with a surface modified to bear ethylenediamine is equilibrated to nominally the same pH and salt conditions as the antibody preparation. The preparation is then recirculated through the monolith until the desired protein has been
35 exposed to pH 3.5 for 60 minutes, or some other specified time interval.

5 [0027] In another embodiment that illustrates the key elements of the disclosed method, a preparation containing a partially purified antibody is titrated to a pH of 3.5. A membrane adsorber with a surface modified to bear polyallylamine is equilibrated to nominally the same pH and salt conditions as the antibody preparation. The preparation is then recirculated through the membrane for a period of 60 minutes. In a closely related embodiment, the antibody preparation is incubated at pH 3.5 for 60 minutes
10 then flowed through the membrane adsorber in a single pass.

[0028] In another embodiment that illustrates the key elements of the disclosed method, a preparation containing a partially purified antibody is titrated to a pH of 3.5. Particles with surfaces modified to bear ethylenediamine, or polyallylamine, or other primary-amine-inclusive ligand are equilibrated to nominally the same pH and salt conditions as the antibody preparation. The particles are maintained in a suspended
15 state by stirring for a period of 60 minutes, following which they are removed by sedimentation or filtration or any other expedient method. The particles may optionally be rinsed with clean buffer at pH 3.5 to maximize recovery of the desired protein. In a closely related embodiment, the antibody preparation is incubated at pH 3.5 for 60 minutes, then passed through a column packed with particles bearing ethylenediamine, polyallylamine, tris(2-aminoethyl)amine, or other primary-amine-inclusive ligand.

20 [0029] Terms below are defined so that the embodiments herein may be understood more readily. Additional definitions are set forth throughout the detailed description.

[0030] “**Amines**” are defined as organic compounds derived from ammonia by replacement of one or more hydrogen atoms with organic radicals, often with a hydrocarbon such as a methyl or ethyl group. On a “**primary amine**,” one hydrogen atom is replaced by an organic radical. On a
25 “**secondary amine**,” two hydrogens are replaced by organic radicals. On a “**tertiary amine**,” three hydrogens are replaced by organic radicals. On a “**quaternary amine**,” three hydrogens are replaced by organic radicals and a fourth organic radical replaces an electron pair that exists on less substituted amines. Primary-amine-containing compounds may also contain other types of amines, or non-amine species.

30 [0031] “**Hydrogen bonds**” are defined as weak bonds between two molecules resulting in part from an electrostatic attraction between a proton in one molecule and an electronegative atom in another, and in part from covalent interactions.

[0032] “**Virus**” or “**virion**” refers to an ultramicroscopic (roughly 20 to 300 nm in diameter), metabolically inert, infectious agent that replicates only within the cells of living hosts, mainly
35 bacteria, plants, and animals: composed of an RNA or DNA core, a protein coat, and, in more

5 complex types, a surrounding envelope containing one or two layers of lipids. Examples include but are not limited to a dsDNA virus, a ssDNA virus, a dsRNA virus, a (+)ssRNA virus, a (-)ssRNA virus, a ssRNA RT virus and a dsDNA-RT virus; an adenovirus, a herpesvirus, a poxvirus, a parvovirus, a reovirus, a norovirus, a picornavirus, a togavirus, an orthomyxovirus, a rhabdovirus, a retrovirus, a hepadanvirus, a papillomavirus, a *Human Immunodeficiency Virus* (HIV), an influenza virus, dengue virus, Japanese encephalitis virus, West Nile virus, and
10 bacteriophages. The term virus is understood to include virus particles for use as vectors for gene therapy, for use as vaccines, and as replacements for antibiotics. It is also understood to include so-called pseudovirions, which may be described as virus particles that have been recombinantly modified to conserve their ability to generate protective immunity while eliminating their ability
15 to cause infection, and may also described as xenotropic viral species where distinct viral species spontaneously transfer genes from one to the other within a host that is infected with both.

[0033] “**Virus inactivation**” refers to a process that damages the virus in a way that suspends its ability to reproduce. Heat is commonly used for this purpose, as well exposure to chemical agents or extreme chemical environments such as highly acidic or highly alkaline conditions, or
20 exposure to organic solvents, among others.

[0034] “**Virus removal**” refers to a process that removes virus from a preparation. Such removal may result from physical methods, such as filtration, where the porosity of a filtration device prevents the passage of virus, or size exclusion chromatography, where the size of a virus causes it to be fractionated from smaller proteins. Removal may also result from the chemical
25 association of a virus with a complementary surface, such as a chromatography surface, such as in ion exchange, affinity, hydrophobic interaction, or hydroxyapatite chromatography, or other chromatography methods.

[0035] “**Virus reduction**” refers to the total reduction of virus load from a preparation by inactivation and removal.

30 [0036] “**Viral polynucleotides**” refers to the genetic material of a virus, in some case DNA, in some cases RNA.

[0037] In certain embodiments, the disclosed method can be applied without variation from the general examples described above to enhance virus reduction. The pH of the inactivating solution should be selected to achieve the highest degree of virus inactivation possible without
35 consideration of the solid material bearing a primary amine. Within the effective range of pH

5 values known to inactivate virus without permanently damaging the product of interest, which may be from a value lower than pH 4, to a value of about pH 3, the solid material may be expected to substantially enhance net viral reduction. Time of exposure may also be selected independent of the solid material bearing primary amine. Temperature of exposure may also be selected independent of the solid material bearing primary amine. Addition of antiviral
10 components may be considered, however, it will be prudent to check certain variables to assess their individual and collective impact. Nonionic, zwitterionic, or positively charged additives can be expected to be tolerated without significant interference to the ability of the solid material bearing primary amines to retain virus or viral DNA. Electronegative additives may have a tendency to interfere with virus binding to the solid material bearing primary amines. Conductive
15 additives such as salts may limit the ability of the solid material bearing primary amines to bind virus, though generally, salt concentrations that produce roughly physiological conductivity or less will be well tolerated, and in some embodiments, salt concentrations twice or thrice physiological conductivity will still permit the solid material bearing primary amines to deliver a worthy enhancement to overall virus reduction. Simple experiments can be conducted at various
20 levels of additives to determine the degree to which they may reduce the ability of the solid material bearing primary amines to reduce virus. The technique known as Design of Experiments (DoE) is widely known in the art for minimizing the number of tests required to obtain a statistically valid body of data to identify the optimum conditions for a given set of variables. As a general matter, the conditions that support the most effective virus inactivation independent of
25 the solid material bearing primary amines, will support the greatest overall reduction of virus in combination with the solid material bearing primary amines.

[0038] In some embodiments, the disclosed methods will enhance removal of DNA and/or contaminating proteins in parallel with enhancing virus reduction. Experimental data indicate that this occurs through the same chemical mechanism as virus reduction; unexpectedly
30 increasing with decreasing pH. In some such embodiments, for example where the desired product does bind to the primary-amine-bearing surface, it may be advantageous to employ the lowest practical salt concentration to favor the highest reduction of virus. In other embodiments, for example where the desired protein has an attraction to the primary amine-bearing surface, it may be necessary to increase the salt concentration for the dual purpose of preventing loss of
35 desired product, and to prevent bound desired protein from interfering with virus-binding

5 capacity. In some such embodiments, the increase in salt concentration should generally not be greater than necessary to prevent desired protein loss and/or desired protein interference with virus binding. Since each species of desired protein presents different properties, it is understood that experimentation may be required to determine if salts are required, and to determine their minimum effective concentrations. It is within the ability of persons skilled in the art to
10 determine the effective concentration through simple experiments.

[0039] In some embodiments, it will be beneficial to evaluate primary amine bearing surfaces and devices of various form and composition to determine which provides the most effective contact and highest capacity for virus reduction. As a general matter, membranes offer less flow resistance than monoliths but monoliths offer higher virus binding capacity. Porous particles
15 suspended in the desired protein preparation generally suffer from poor capacity and retarded virus binding kinetics due to the large molecular distances among viruses and charged particles. Columns packed with porous particles offer more rapid kinetics than suspended particles, though they are still slow in comparison to membranes and monoliths and still suffer from low capacity, but porous particles may achieve worthy enhancement of virus reduction nonetheless.

20 EXAMPLES

[0040] **Example 1.** DNA binding on an ethylenediamine-bearing monolith (CIM EDA, BIA Separations) as a function of pH. Genomic DNA from salmon sperm was used as a surrogate for viral DNA. An EDA monolith was equilibrated with buffers ranging from pH 8.0 to 3.5 in separate experiments. DNA suspended in 50 mM Tris, pH 8 was injected onto the monolith. After
25 a wash step to displace unbound sample components, the DNA was eluted in a linear gradient of either NaCl or guanidine. In both cases, results indicate a disproportionate increase in strength of retention beginning at pH 4. Experimental results are provided in Table 1:

Table 1

	pH	buffering species	M NaCl elution	M Guanidine elution
30	8.0	50 mM Tris	1.65	1.84
	7.0	50 mM Hepes	2.00	2.15
	6.0	50 mM MES	2.82	2.79
	5.0	50 mM acetate	3.35	3.06
	4.0	50 mM acetate	4.85	5.39
35	3.0	50 mM acetate	4.85	5.41

5 [0041] **Example 2.** Enhancement of DNA retention on a primary amine-bearing surface versus weakened retention on a quaternary amine-bearing surface. Genomic DNA from salmon sperm was used as a surrogate for viral DNA. Retention of DNA was evaluated at pH 8, 7, and 6 on a primary amine-bearing monolith (CIM EDA, BIA Separations) versus a quaternary amine-bearing monolith (CIM QA, BIA Separations). DNA eluted from QA at 0.6 M NaCl at pH 8.0, 10 weakening to 0.55 M NaCl at pH 7.0, and further weakening to 0.47 M NaCl at pH 6.0. DNA eluted from EDA at 0.83 M NaCl at pH 8.0, 1.11 M NaCl, at pH 7.0, and 1.8 M NaCl at pH 6.0. These results highlight the fact that retention on the primary amine-bearing surface is opposite to the trend, and of dramatically greater magnitude than observed with the quaternary amine-bearing surface.

15 [0042] **Example 3.** Enhancement of virus retention on a primary amine-bearing surface versus weakened retention on a quaternary amine-bearing surface. Bacteriophage M13 was used as a test virus. Retention of DNA was evaluated at pH 8, 7, and 6 on a primary amine-bearing monolith (CIM EDA, BIA Separations) versus a quaternary amine-bearing monolith (CIM QA, BIA Separations). DNA eluted from QA at 0.7 M NaCl at pH 8.0, weakening to 0.61 M NaCl at 20 pH 7.0, and further weakening to 0.53 M NaCl at pH 6.0. DNA eluted from EDA at 1.02 M NaCl at pH 8.0, 1.47 M NaCl, at pH 7.0, and 2.5 M NaCl at pH 6.0. These results highlight the fact that retention on the primary amine-bearing surface is opposite to the trend, and dramatically greater than observed with the quaternary amine-bearing surface.

[0043] **Example 4.** Enhanced reduction of minute virus of mice (MVM) with a primary amine-bearing monolith. A viral cell culture containing MVM was exposed to pH 3.5 for 60 minutes. 25 MVM is known not to be inactivated by these conditions. Solids were removed by filtration and the filtrate was passed through an EDA monolith equilibrated to pH 3.5. MVM content was reduced by a $\log_{10}$ factor of 2.85. This represents about a 700-fold enhancement of virus reduction, and particularly highlights the ability of the disclosed method to enhance removal of 30 nonenveloped virus species, particularly including retroviral species such as MVM.

[0044] **Example 5.** Enhanced reduction of Hepatitis A virus (HAV) with a primary amine-bearing monolith. A viral cell culture containing HAV was exposed to pH 3.5 for 60 minutes. HAV is known not to be inactivated by these conditions. Solids were removed by filtration and the filtrate was passed through an EDA monolith equilibrated to pH 3.5. MVM content was 35 reduced by a $\log_{10}$ factor of 2.85. This represents about a 700-fold enhancement of virus reduction

5 and further highlights the ability of the disclosed method to enhance removal of nonenveloped virus species.

[0045] It will be apparent to the person of skill in the art, that the disclosed methods may also be used for the purification of viruses, especially non-lipid enveloped viruses where the low pH of binding may provide the secondary benefit of inactivating lipid-enveloped viruses in
10 conjunction with purification. For the same reasons, it will be equally apparent that the disclosed methods may also be used for isolation of DNA from preparations that may be contaminated by lipid-enveloped viruses. In both cases, the ability of the disclosed methods to disproportionately enhance the retention of target products according to their size will also favor the elimination of smaller contaminants such as proteins.

15 [0046] All references cited herein are incorporated by reference in their entirety and for all purposes to the same extent as if each individual publication or patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes. To the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede
20 and/or take precedence over any such contradictory material.

[0047] All numbers expressing quantities of ingredients, chromatography conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are approximations that may vary depending upon the
25 desired performance sought to be obtained by the present embodiments.

[0048] Many modifications and variations of the embodiments herein can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only and are not meant to be limiting in any way. It is intended that the specification and examples be considered as exemplary
30 only, with the true scope and spirit of the embodiments being indicated by the following claims.

5 CLAIMS

What is claimed is:

1. A method for reducing virus and viral DNA levels in an acidic preparation of a desired
10 protein, the method comprising contacting the acidic preparation of the desired protein with a surface bearing a primary amine, secondary amine, or both.
2. The method of claim 1, wherein an operating pH during the contacting step is in a range selected from the group consisting of (a) from about 2.0 to about 4.0, (b) from about 2.25 to about 3.75, (c) from about 2.5 to about 3.5, (d) from about 2.75 to about 3.25, (e) from about
15 3.5 to about 3.75, and (f) from about 3.25 to about 3.5.
3. The method of claim 1, wherein a salt concentration during the contacting step is range selected from the group consisting of (a) from about 0.0 to about 0.5 M, (b) from about 0.05 to about 0.3 M, (c) from about 0.1 to about 0.2M.
4. The method of claim 1, wherein the duration of the contacting step is in a range selected
20 from the group consisting of (a) from about 15 to about 120 minutes, (b) from about 30 to about 60 minutes, (c) a non-zero amount less than about 30 minutes, (d) a non-zero amount less than about 15 minutes, (e) a non-zero amount less than about 10 minutes, (f) a non-zero amount less than about 5 minutes, (g) a non-zero amount less than about 2 minutes, (h) a non-zero amount less than about 1 minute, and (i) a time required to pass the preparation through a
25 device housing the primary amine-bearing surface.
5. The method of claim 1, wherein the surface comprises one or more moieties selected from the group consisting of ethylenediamine, tris(2-aminoethyl)amine, diethylaminetriamine, triethylenetetraamine, tetraethylenepentamine, and a polyallylamine.
6. The method of claim 1, wherein the surface is in the form of one selected from the
30 group consisting of a particle, a membrane, a monolith, a hydrogel-coated skeletal framework, and combinations thereof.
7. The method of claim 1, wherein the contacting step is performed by an operation selected from the group consisting of (a) dispersing primary or secondary amine-bearing particles in the preparation, (b) passing the preparation through a device in fluid contact with
35 the primary amine-bearing surface, (c) recirculating the preparation through a device in providing fluid contact with the primary amine-bearing surface, and combinations thereof.

INTERNATIONAL SEARCH REPORT

 International application No.
PCT/SG2014/000232

A. CLASSIFICATION OF SUBJECT MATTER

C07K 1/14 (2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Medline, WPI, EPODOC, CAPLUS, Biosis. Keywords: virus, viral, viral DNA, virus DNA, reduce, decrease, limit, remove, lower, chromatograph, purify, separate, isolate, acid, low pH, protein, peptide, polypeptide, antibody, immunoglobulin, TREN, tris(2-aminoethyl)amine, diethylenetriamine, triethylenetetramine, tetraethylenepentamine, amine, particle, membrane, monolith hydrogel-coated skeletal framework, Gagnon P and related terms.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	Documents are listed in the continuation of Box C	



Further documents are listed in the continuation of Box C



See patent family annex

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 26 August 2014	Date of mailing of the international search report 26 August 2014
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA Email address: pct@ipaaustralia.gov.au	Authorised officer Ishanee Mookerjee AUSTRALIAN PATENT OFFICE (ISO 9001 Quality Certified Service) Telephone No. 0262256167

INTERNATIONAL SEARCH REPORT		International application No.
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		PCT/SG2014/000232
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/062841 A1 (BIO-RAD LABORATORIES INC.) 02 May 2013 [0021], [0025], [0032]-[0034], [0035], [0040], [0041]	1-4, 6, 7
X	WO 2013/180655 A1 (AGENCY FOR SCIENCE, TECHNOLOGY AND RESEARCH) 05 December 2013 [0004], [0007], [0009], [0024], [0033], [0034]	1, 2, 4, 5, 7

Form PCT/ISA/210 (fifth sheet) (July 2009)

INTERNATIONAL SEARCH REPORT Information on patent family members		International application No. PCT/SG2014/000232	
This Annex lists known patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.			
Patent Document/s Cited in Search Report		Patent Family Member/s	
Publication Number	Publication Date	Publication Number	Publication Date
WO 2013/062841 A1	02 May 2013	CN 103906762 A	02 Jul 2014
		US 2013109807 A1	02 May 2013
WO 2013/180655 A1	05 December 2013	None	
End of Annex			
Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001. Form PCT/ISA/210 (Family Annex)(July 2009)			