



(51) International Patent Classification:

A61K 47/68 (2017.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/EP2023/057565

(22) International Filing Date:

23 March 2023 (23.03.2023)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

22163928.9 23 March 2022 (23.03.2022) EP

(71) Applicant: SYNAFFIX B.V. [NL/NL]; Kloosterstraat 9, 5349 AB Oss (NL).

(72) Inventors: LELIEVELDT, Lianne; Biezenstraat 180, 6541 ZZ Nijmegen (NL). VAN GEEL, Remon; Margrietstraat 17, 5396 CM Lithoijen (NL). VAN BERKEL, Sander Sebastiaan; Kloosterstraat 9, 5349 AB Oss (NL). VAN DELFT, Floris Louis; Van Goorstraat 81, 6512 EC Nijmegen (NL).

(74) Agent: NEDERLANDSCH OCTROOIBUREAU; P.O.Box 29720, 2502 LS The Hague (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MU, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, CV, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SC, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT,

LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTIBODY-CONJUGATES FOR TARGETING OF TUMOURS EXPRESSING CARCINOEMBRYONIC ANTIGEN

(57) Abstract: The present invention concerns antibody-conjugates which are especially suitable for the targeting of CEA-expressing cells, in particular tumour cells. The antibody-conjugates according to the invention have structure (1): $AB-[(L^6)_b-\{Z-L-D\}_x]_y$ (1) Herein, AB is an antibody capable of targeting CEA-expressing tumours; L is a linker that links Z to D; Z is a connecting group; L^6 is $-GlcNAc(Fuc)_w-(G)_j-S-(L^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 - 10, S is a sugar or a sugar derivative, GlcNAc is N- acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-N(H)C(O)CH_2-$, $-N(H)C(O)CF_2-$ or $-CH_2-$; D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolbenzodiazepine dimers (PBDs), and analogues or prodrugs thereof; b is 0 or 1; x is 1 or 2; and y is 1, 2, 3 or 4. The invention further concerns a method for preparing the antibody-conjugates of structure (1) and application of the antibody-conjugates of structure (1).



Antibody-conjugates for targeting of tumours expressing carcinoembryonic antigen

Field of the invention

[0001] The present invention is in the field of bioconjugation. More specifically, the present invention relates to antibody-drug conjugates for targeted treatment of patients with cancer, in particular carcinoembryonic antigen (CEA)-expressing tumours.

Background

[0002] A promising approach for targeted treatment of tumours entails the conjugation of a multitude (2 to 8) of highly toxic payloads to a monoclonal antibody, thereby generating an antibody-drug conjugate (ADCs). ADCs are well known in the art, as for example described by Chari et al., *Angew. Chem. Int. Ed.* **2014**, 53, 3796 and Beck et al., *Nat. Rev. Drug Discov.* **2017**, 16, 315–37. Mechanistically, the antibody is designed to bind with high specificity to tumour-associated receptor that is overexpressed versus healthy tissue. The ADC is thought to internalize into the tumour cell after binding to the receptor, then to release the toxic payload upon degradation of the antibody and/or the linker in the lysosome.

[0003] Current ADCs are commonly prepared by various conjugation technologies (summarized in Figure 1), mostly based on conjugation to cysteine side chains with maleimides or to lysine side chains with activated esters. To generate an ADC based on native cysteines naturally engaged in disulfide bonds, the thiol in the side-chain can be liberated by subjection of the antibody to a suitable reducing agent such as TCEP or DTT, followed by treatment with a maleimide-functionalized linker-drug. The resulting ADC will typically consist of a mixture of positional isomers, in case only the total of eight free thiols are not comprehensively alkylated. Alternatively, to generate a site-specific ADC, an antibody can be generated by mutating one or more amino acids at defined positions in the sequence to cysteines, the side-chain of which can be selectively liberated for alkylation by a reduction-oxidation sequence. Commonly employed cysteines for site-specific conjugation are LC-41C (light chain 41C), HC-41C (heavy chain 41C), LC-80C, HC-118C, HC-265C, HC-140C, LC-149C, LC-124C, LC-180C, HC-190C, HC-160C, LC-183C, HC-290C, LC-205C, HC-220C, HC-239C, HC-442C. An additional cysteine can also be inserted into the sequence, for example HC-i239C. Besides maleimides as alkylating agents, reaction of cysteine side chain with haloacetamide or vinylbenzene derivatives has also been reported. Besides reaction to natural amino acid side chains, specific unnatural (non-canonical) amino acids can also be engineered into the amino acid sequence of an antibody, thereby providing a unique handle for chemical conjugation, such as ketone, acetylene, azide, cyclic alkyne or cyclic alkene, for reaction with oxime, azide, alkyne or tetrazine, respectively. However, a disadvantage of the latter approach is that the natural sequence of the antibody must be re-engineered, which besides being time-consuming and costly, may lead to instability issues.

[0004] Conjugation through the glycan by an oxidation–ligation sequence is known in the art and has for example been described by Hamann et al. (*Bioconjugate Chem.* **2002**, 13, 47–58). Chemoenzymatic conjugation through the glycan is known in the art and has been described for

the use of sialyltransferase by Boons et al, *Angew. Chem. Int. Ed.* **2014**, 53, 7179, and for the use of a mutant galactosyltransferase by Zhu et al, *mAbs* **2014**, 6, 1 and Cook et al, *Bioconjugate Chem.* **2016**, 27, 1789.

[0005] Chemoenzymatic conjugation through the glycan including first trimming of the glycan is known in the art and has been described by van Geel et al, *Bioconjugate Chem.* **2015**, 26, 2233 and is schematically depicted in Figure 2. In short, the monoclonal antibody is treated with an endoglycosidase to trim the glycan down to the core GlcNAc (attached directly to Asn-297), followed by transfer of an azido-modified sugar under the action of a glycosyltransferase. Various structures of UDP-azidosugars are depicted in Figure 3. One particularly suitable combination involves that transfer of GalNAz **2b** (2-azidoacetyl-*N*-galactosamine) under the action of a mutant galactosyltransferase GalT(Y289L) as disclosed in WO 2007/095506, EP 2911699 B1 and van Geel et al. An alternative powerful combination entails 6-azidoGalNAc **2d** with native GalNAc-transferase, as has been disclosed in PCT/EP2016/059194. Another useful combination entails GlcNAz with α -1,3-mannosyl-glycoprotein-2- β -*N*-acetylglucosaminyltransferase (MGAT1) as disclosed in WO2021/248048.

[0006] Various cyclooctynes for application in metal-free click chemistry are known in the art (Figure 4). In particular, various cyclooctynes such as DIBO (**I**), DBCO/DIBAC (**J**), s-DIBO (**K**), BCN (**L**) and TMTHSI (**T**) are regularly applied for conjugation to azide.

[0007] A payload for an ADCs is typically a highly cytotoxic molecule, with IC₅₀-value in low nanomolar or picomolar range, in particular low to medium molecular weight compounds (e.g. about 200 to about 2500 Da). Examples of suitable cytotoxin classes for ADCs include anthracyclines, camptothecins, taxanes, tubulysins, enediynes, inhibitory peptides, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, hemiasterlins, BCL-XL inhibitors, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs) and analogues or prodrugs thereof. A representative set of cytotoxic molecules, and/or synthetic derivatives or prodrugs thereof, with suitable attachment point for conjugation to a monoclonal antibody, is depicted in Figure 5.

[0008] Specific examples of anthracyclins suitable of application in ADCs include (but are not limited to) doxorubicin, daunorubicin, nemorubicin and PNU-159,682.

[0009] Specific examples of camptothecins suitable for application in ADCs include (but are not limited to) SN-38, exatecan, exatecan-S, topotecan, silatecan, cositecan, lurtotecan, gimatecan, belotecan, rubitecan, AMDCPT, G-AMDCPT and other synthetic camptothecins the structures of which are depicted in Figure 6. Various novel camptothecins have been disclosed in EP0296597, WO2019/236954, WO2020/200880, WO2020/219287, CN113816969, CN113710277 and US20180200273.

[0010] Specific examples of enediynes suitable for application in ADCs include (but are not limited to) calicheamicin, esperamicins, shishijimicins and namenamicins and other enediynes as summarized by Galm et al., *Chem. Rev.* **2005**, 105, 739-758.

[0011] Specific examples of auristatins suitable for application in ADCs include (but are not limited to) MMAD, MMAE, MMAF and PF-06380101 and other auristatins as summarized by Maderna et

al., *Mol. Pharmaceutics* **2015**, 12, 1798-1812.

[0012] The 7 genes that belong to the CEACAM subgroup show distinct expression patterns on different cell types but share structural homology, as they are composed, in their extracellular domains (ECDs), of Ig variable and constant-like domains, as reported by Kuespert et al. *Curr. Opin. Cell. Biol.* **2006**, 18, 565–571, and Beauchemin et al., *Cancer Metastasis Rev.* **2013**, DOI 10.1007/s10555-013-9444-6, see Figure 8.

[0013] The gene carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5), also known as CD66e, is the founding member of the CEACAM family, and was first identified by Gold and Freedman in 1965. CEACAM5 codes for the protein known as carcinoembryonic antigen (CEA), see Hefta et al. *Proc. Nat. Acad. Sci.* **1988**, 85, 4648-4652. CEA is a glycoprotein involved in cell adhesion and is highly expressed in particular on the surface of colorectal, gastric, lung and uterine tumour cells. The CEACAM5 cDNA encodes a protein that exhibits one variable (V)-like domain, identified as the N domain, followed by three repeating units comprising, in total, six constant C2-like domains (termed A1, B1, A2, B2, A3, and B3, see Figure 8. The Ig domains are preceded by a 34-amino acid signal peptide. In addition, the CEA cDNA structure predicted 12 cysteine residues forming the core of the Ig folds and 28 N-linked glycosylation sites in line with the high carbohydrate content of the purified protein.

[0014] A reference sequence of full length human CEACAM5, including signal peptide (positions 1-34) and pro-peptide (positions 686-702), is available from the GenBank database under accession number AAA51967.1. Domain organisation of human CEACAM5 is as follows:

Human CEACAM5 domains	Positions
Domain N	35-142
Domain A1	143-237
Domain B1	238-320
Domain A2	321-415
Domain B2	416-498
Domain A3	499-593
Domain B3	594-685

Accordingly, the A3-B3 domain of human CEACAM5 consists of amino acids at positions 499-685.

[0015] CEACAM5 has limited expression in normal adult tissues, but is overexpressed in carcinomas of the gastrointestinal tract, the genitourinary and respiratory systems, and breast cancer, as reported by Hammarstrom, *Semin. Cancer Biol.* **1999**, 9, 67–81. Thus, CEACAM5 (CEA) may constitute a therapeutic target suitable for tumour specific targeting approaches, such as immunoconjugates or antibody-drug conjugates (ADCs).

[0016] Circulating CEA in serum is barely detectable in healthy adults (<5 ng/mL), but is shed from tumours, either directly or via the lymphatics, and is then detected in the serum. Because of this property, the level of serum CEA has been used as a clinical marker for diagnosis of cancers and screening for recurrence of cancers, particularly colorectal cancer (Goldenberg et al., *Int. J. Biol. Mark.* **1992**, 7, 183-188; Chau et al. *J. Clin. Oncol.* **2004**, 22, 1420-1429; Flamini et al. *Clin. Cancer Res.* **2006**, 12, 6985-6988.

[0017] The high tumour expression levels of CEA relative to that in normal tissue makes this glycoprotein an attractive receptor for targeted treatment strategies with monoclonal antibodies. Therefore, multiple monoclonal antibodies have been raised, and humanized in various cases, against CEA for research purposes, as diagnostic tools and for therapeutic purposes, see for example various reviews summarizing anti-CEA antibodies, e.g. Nap et al., *Cancer Res.* **1992**, 52, 2329-2339; Sheahan et al., *Am. J. Clin. Path.* **1990**, 94, 157-164. Specific reports on anti-CEA antibodies: US9,695,250 (A5/A240VL and D8/A240VL); Richman et al., *Int. J. Cancer* **1987**, 39, 317-32, Stewart et al., *Cancer Immunol. Immunother.* **1999**, 47, 299-306, Ashraf et al., *Br. J. Canc.* **2009**, 101, 1758-1768, WO199506067 and EP0721470 (PR1A3), Bacac et al. *Clin. Canc. Res.* **2016**, 22, 3286-3297, US8,642,742 and US9,068,008 (antibodies based on VH CH1A1A/CH7A or CH1A1B/CH7A and VL pAC18/2F1); WO2021053587 (AB17, AB72 and AB73); US9,068,008; Sakurai et al., *J. Surg. Oncol.*, **1989**, 42, 39-46 (antibodies NCC-CO-413, 308, 432, 411); Harwood et al., *Br. J. Cancer*, 1982, 54, 72; Ledermann, *Br. J. Cancer*, **1988**, 58, 654; Pedley et al., *Br. J. Cancer*, **1993**, 68, 69-73; Boxer et al., *Br. J. Cancer*, **1992**, 65, 825-831 and US8,394,926 (antibody A5B7); Baek et al. *Canc. Lett.* **2022**, 525, 97-107 (antibody 1G9); Hansen et al., *Cancer* **1993**, 71, 347, US5,874,540 and humanized version thereof as described in US7,803,372, US8,778,342, WO2004032857A2, and WO2015069430A2 (hMN-14 or labeltuzumab); US7,776,330 (antibodies M5A and M5B derived from murine T84.66). Chester et al. have isolated a single chain anti-CEA antibody (MFE-23) from a phage display library to be used in radioimmunoassay and radioimmunotherapy (US5,876,691), and the antibody was subsequently humanized (to SM3E) as described in US7,232,888 and US20050147614). Shinmi et al. *Canc. Med.* **2017**, 6, 798-808 and Iwano et al. *Drug Metab. Dispos.* **2019**, 47, 1240-1246 have isolated and used for targeted chemotherapy anti-CEA antibodies 12-140-1 and 15-1-32. Anti-CEA antibody CEA6 and other variants CEA1-CEA5 and CEA7, (derived from VH sequences T06D10, HBA11, HBB11, HBB6 and VL sequences T06D4, T06D8, T06D12) has been obtained from human phage display libraries (WO1997020932A1 and US5,872,215). Anti-CEA antibody 769-cea-4 was generated by immunization and humanized as described in US9,617,345 and US10,457,739 (tusamitamab or SAR408377). Micromet's MT111 antibody (also known as MEDI-565 antibody of MedImmune) is a bispecific antibody based on A5B7 for binding to human CEACAM5 and human CD3, as reported by Peng et al., *PLoS ONE* **2012**, 7, e3641 and WO2007071426.

[0018] Another CEACAM family member CEACAM6 (also called CD66c) codes for the protein known as NCA-90. CEACAM6 is a non-specific cross-reacting glycoprotein antigen that shares some antigenic determinants with CEACAM5, as described by Kuroki et al., *Biochem. Biophys. Res. Comm.* **1992**, 182, 501-506. CEACAM6 also is expressed on granulocytes and epithelia from various organs and has a broader expression zone in proliferating cells of hyperplastic colonic polyps and adenomas, compared with normal mucosa, as well as by many human cancers. Relatively high serum levels of NCA-90 are found in patients with lung, pancreatic, breast, colorectal, and hepatocellular carcinomas. The amount of CEACAM6 does not correlate with the amount of CEACAM5 expressed, as described by Kuroki et al., *Anticancer. Res.* **1999**, 19, 5599-5606. In addition to structural homology, CEACAMs also share sequence similarity, with 85.7%

sequence identity in the ECDs of CEACAM5 and CEACAM6, in particular.

[0019] Therapeutic antibodies against CEACAM6 are known in the art. Some are not selective for human CEACAM6 (e.g. Immunomedics MN-3, Neo201/h16C3 from Neogenix; both also bound to human CEACAM5). Single domain antibody 2A3 and its fusion mutants (WO2012040824 and Niu et al., *J. Contr. Rel.* **2012**, *10*, 18-24) have not been characterized with respect to selectivity and cross-reactivity for monkey CEACAM5. Mouse antibody 9A6 (Genovac/Aldevron) is the only antibody described to be capable of regulating the immunosuppressive activity of CEACAM6 (Witzens-Harig et al. *Blood* **2013**, *121*, 4493-4503). Potent anti-CECAM6 antibodies for cancer immunotherapy including TPP-3310 were disclosed in WO2016150899.

[0020] ADCs targeting CEACAM5 are known in the art. For example, Govindan et al. *Clin. Canc. Res.* **2009**, *15*, 6052-6061 have reported on the preclinical and clinical evaluation of an ADC known as labetuzumab-govetican obtained by modification of hMN-14 with CL2A linker-drug with SN-38 payload. Also, Shinmi et al. *Canc. Med.* **2017**, *6*, 798-808 and Iwano et al. *Drug Metab. Dispos.* **2019**, *47*, 1240-1246 have preclinically evaluated an anti-CEA ADC conjugated with MMAE payload based on antibodies 12-140-1 and 15-1-32. Thirdly, Decary et al. *Clin. Canc. Res.* **2020**, *26*, 6589-6599 have reported on derivatization and clinical evaluation of the anti-CEA antibody SAR408377 derivatized with DM4 payload, an ADC known as SAR408701 or tusamitamab-ravtansine.

[0021] ADCs against CEACAM6 are known in the art, such as the maytansinoid anti-CEACAM6 antibody reported by Genentech Strickland et al, *J. Pathol.* **2009**, *218*, 380, which in a non-human primate study was shown to induce CEACAM6-dependent haematopoietic toxicity. This toxicity, attributed to accumulation of the antibody drug conjugate in bone marrow and depletion of granulocytes and their cell precursors, was considered by the authors as a major safety concern.

Summary of the invention

[0022] The inventors have developed antibody-conjugates that are highly suitable for targeting CEA-expressing cells, in particular tumours. Thus, the antibody-conjugates according to the invention are highly suitable for treating CEA-positive cancer, especially colorectal cancer, gastric cancer, lung cancer, uterine cancer or pancreatic cancer.

[0023] In a first aspect, the present invention concerns an antibody-conjugate. Related thereto, in a second aspect, the invention concerns a process for preparing the antibody-conjugate according to the invention. In a third aspect, the invention concerns a method for targeting CEA-expressing cells. Related thereto are the first medical use of the antibody-conjugate according to the invention, as well as the second medical use for the treatment of cancer. In a last aspect, the invention concerns the use of a mode of conjugation for increasing the therapeutic index of an antibody-conjugate in the treatment of CEA-expressing tumours.

Detailed description

Definitions

[0024] The verb "to comprise", and its conjugations, as used in this description and in the claims

is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded.

[0025] In addition, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there is one and only one of the elements. The indefinite article "a" or "an" thus usually means "at least one".

[0026] A linker is herein defined as a moiety that connects (covalently links) two or more elements of a compound. A linker may comprise one or more spacer moieties. A spacer-moiety is herein defined as a moiety that spaces (i.e. provides distance between) and covalently links together two (or more) parts of a linker. The linker may be part of e.g. a linker-construct, a linker-conjugate, a linker-payload (e.g. linker-drug) or an antibody-conjugate, as defined below.

[0027] A "hydrophilic group" or "polar linker" is herein defined as any molecular structure containing one or more polar functional groups that imparts improved polarity, and therefore improved aqueous solubility, to the molecule it is attached to. Preferred hydrophilic groups are selected from a carboxylic acid group, an alcohol group, an ether group, a polyethylene glycol group, an amino group, an ammonium group, a sulfonate group, a phosphate group, an acyl sulfamide group or a carbamoyl sulfamide group. In addition to higher solubility other effects of the hydrophilic group include improved click conjugation efficiency, and, once incorporated into an antibody-drug conjugate: less aggregation, improved pharmacokinetics resulting in higher efficacy and in vivo tolerability.

[0028] The term "salt thereof" means a compound formed when an acidic proton, typically a proton of an acid, is replaced by a cation, such as a metal cation or an organic cation and the like. Where applicable, the salt is a pharmaceutically acceptable salt, although this is not required for salts that are not intended for administration to a patient. For example, in a salt of a compound the compound may be protonated by an inorganic or organic acid to form a cation, with the conjugate base of the inorganic or organic acid as the anionic component of the salt. The term "pharmaceutically accepted" salt means a salt that is acceptable for administration to a patient, such as a mammal (salts with counterions having acceptable mammalian safety for a given dosage regime). Such salts may be derived from pharmaceutically acceptable inorganic or organic bases and from pharmaceutically acceptable inorganic or organic acids. "Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound, which salts are derived from a variety of organic and inorganic counter ions known in the art and include, for example, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, etc., and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, formate, tartrate, besylate, mesylate, acetate, maleate, oxalate, etc.

[0029] The term "enediyne" or "enediyne antibiotic" or "enediyne-containing cytotoxin" refers to any cytotoxin characterized by the presence of a 3-ene-1,5-diyne structural feature as part of a cyclic molecule as known in the art and include neocarzinostatin (NCS), C-1027, kedarcidin (KED), maduropeptin (MDP), N1999A2, the sporolides (SPO), the cyanosporasides (CYA and CYN), and the fijiolides, calicheamicins (CAL), the esperamicins (ESP), dynemicin (DYN), namenamycin,

shishijimicin, and unciamycin (UCM).

[0030] The term “alkylaminosugar” as used herein means a tetrahydropyranyl moiety connected to an alcohol function via its 2-position, thereby forming an acetal function, and further substituted by (at least) one N-alkylamino group in position 3, 4 or 5. “N-alkylamino group” in this context refers to an amino group having one methyl, ethyl or 2-propyl group.

[0031] The term “click probe” refers to a functional moiety that is capable of undergoing a click reaction, i.e. two compatible click probes mutually undergo a click reaction such that they are covalently linked in the product. Compatible probes for click reactions are known in the art, and preferably include (cyclic) alkynes and azides. In the context of the present invention, click probe Q in the compound according to the invention is capable of reacting with click probe F on the (modified) protein, such that upon the occurrence of a click reaction, a conjugate is formed wherein the protein is conjugated to the compound according to the invention. Herein, F and Q are compatible click probes.

[0032] An “acylsulfamide moiety” is herein defined as a sulfamide moiety ($\text{H}_2\text{NSO}_2\text{NH}_2$) that is N-acylated or N-carbamoylated on one end of the molecule and N-alkylated (mono or bis) at the other end of the molecule. In the context of the present invention, especially in the examples, this group is also referred to as “HS”.

[0033] A “domain” may be any region of a protein, generally defined on the basis of sequence homologies and often related to a specific structural or functional entity. CEACAM family members are known to be composed of Ig-like domains. The term domain is used in this document to designate either individual Ig-like domains, such as “N-domain” or for groups of consecutive domains, such as “A3-B3 domain”.

[0034] A “coding sequence” or a sequence “encoding” an expression product, such as a RNA, polypeptide, protein, or enzyme, is a nucleotide sequence that, when expressed, results in the production of that RNA, polypeptide, protein, or enzyme, i.e., the nucleotide sequence encodes an amino acid sequence for that polypeptide, protein or enzyme. A coding sequence for a protein may include a start codon (usually ATG) and a stop codon.

[0035] The term “gene” means a DNA sequence that codes for, or corresponds to, a particular sequence of amino acids which comprises all or part of one or more proteins or enzymes, and may or may not include regulatory DNA sequences, such as promoter sequences, which determine for example the conditions under which the gene is expressed. Some genes, which are not structural genes, may be transcribed from DNA to RNA, but are not translated into an amino acid sequence. Other genes may function as regulators of structural genes or as regulators of DNA transcription. In particular, the term gene may be intended for the genomic sequence encoding a protein, i.e. a sequence comprising regulator, promoter, intron and exon sequences.

[0036] The term “glycoprotein” is herein used in its normal scientific meaning and refers to a protein comprising one or more monosaccharide or oligosaccharide chains (“glycans”) covalently bonded to the protein. A glycan may be attached to a hydroxyl group on the protein (O-linked-glycan), e.g. to the hydroxyl group of serine, threonine, tyrosine, hydroxylysine or hydroxyproline, or to an amide function on the protein (N-glycoprotein), e.g. asparagine or arginine, or to a carbon on the protein

(C-glycoprotein), e.g. tryptophan. A glycoprotein may comprise more than one glycan, may comprise a combination of one or more monosaccharide and one or more oligosaccharide glycans, and may comprise a combination of N-linked, O-linked and C-linked glycans. It is estimated that more than 50% of all proteins have some form of glycosylation and therefore qualify as glycoprotein.

5 Examples of glycoproteins include PSMA (prostate-specific membrane antigen), CAL (candida antartica lipase), gp41, gp120, EPO (erythropoietin), antifreeze protein and antibodies.

[0037] The term “glycan” is herein used in its normal scientific meaning and refers to a monosaccharide or oligosaccharide chain that is linked to a protein. The term glycan thus refers to the carbohydrate-part of a glycoprotein. The glycan is attached to a protein via the C-1 carbon of one sugar, which may be without further substitution (monosaccharide) or may be further substituted at one or more of its hydroxyl groups (oligosaccharide). A naturally occurring glycan typically comprises 1 to about 10 saccharide moieties. However, when a longer saccharide chain is linked to a protein, said saccharide chain is herein also considered a glycan. A glycan of a glycoprotein may be a monosaccharide. Typically, a monosaccharide glycan of a glycoprotein consists of a single N-acetylglucosamine (GlcNAc), glucose (Glc), mannose (Man) or fucose (Fuc) covalently attached to the protein. A glycan may also be an oligosaccharide. An oligosaccharide chain of a glycoprotein may be linear or branched. In an oligosaccharide, the sugar that is directly attached to the protein is called the core sugar. In an oligosaccharide, a sugar that is not directly attached to the protein and is attached to at least two other sugars is called an internal sugar. In an oligosaccharide, a sugar that is not directly attached to the protein but to a single other sugar, *i.e.* carrying no further sugar substituents at one or more of its other hydroxyl groups, is called the terminal sugar. For the avoidance of doubt, there may exist multiple terminal sugars in an oligosaccharide of a glycoprotein, but only one core sugar. A glycan may be an O-linked glycan, an N-linked glycan or a C-linked glycan. In an O-linked glycan a monosaccharide or oligosaccharide glycan is bonded to an O-atom in an amino acid of the protein, typically via a hydroxyl group of serine (Ser) or threonine (Thr). In an N-linked glycan a monosaccharide or oligosaccharide glycan is bonded to the protein via an N-atom in an amino acid of the protein, typically via an amide nitrogen in the side chain of asparagine (Asn) or arginine (Arg). In a C-linked glycan a monosaccharide or oligosaccharide glycan is bonded to a C-atom in an amino acid of the protein, typically to a C-atom of tryptophan (Trp).

[0038] The term “antibody” (AB) is herein used in its normal scientific meaning. An antibody is a protein generated by the immune system that is capable of recognizing and binding to a specific antigen. An antibody is an example of a glycoprotein. The term antibody herein is used in its broadest sense and specifically includes monoclonal antibodies, polyclonal antibodies, dimers, multimers, multispecific antibodies (e.g. bispecific antibodies), antibody fragments, and double and single chain antibodies. The term “antibody” is herein also meant to include human antibodies, humanized antibodies, chimeric antibodies and antibodies specifically binding cancer antigen. The term “antibody” is meant to include whole antibodies, but also fragments of an antibody, for example an antibody Fab fragment, F(ab')₂, Fv fragment or Fc fragment from a cleaved antibody, a scFv-Fc fragment, a minibody, a diabody or a scFv. Furthermore, the term includes genetically engineered

antibodies and derivatives of an antibody. Antibodies, fragments of antibodies and genetically engineered antibodies may be obtained by methods that are known in the art.

[0039] An antibody may be a natural or conventional antibody in which two heavy chains are linked to each other by disulfide bonds and each heavy chain is linked to a light chain by a disulfide bond.

5 There are two types of light chain, lambda (l) and kappa (k). The light chain includes two domains or regions, a variable domain (VL) and a constant domain (CL). The heavy chain includes four domains, a variable domain (VH) and three constant domains (CH1, CH2 and CH3, collectively referred to as CH). The variable regions of both light (VL) and heavy (VH) chains determine binding recognition and specificity to the antigen. The constant region domains of the light (CL) and heavy
10 (CH) chains confer important biological properties, such as antibody chain association, secretion, trans-placental mobility, complement binding, and binding to Fc receptors (FcR). The Fv fragment is the N-terminal part of the Fab fragment of an immunoglobulin and consists of the variable portions of one light chain and one heavy chain. The immunoglobulin can be of any type (e.g. IgG, IgE, IgM, IgD, and IgA), class (e.g. IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass, or allotype (e.g.
15 human G1m1, G1m2, Gm3, non-G1m1 [that, is any allotype other than G1m1], G1m17, G2m23, G3m21, G3m28, G3m1.1, G3m5, G3m13, G3m14, G3m10, G3m15, G3m16, G3m6, G3m24, G3m26, G3m27, A2m1, A2m2, Km1, Km2 and Km3) of immunoglobulin molecule. Preferred allotypes for administration include a non-G1m1 allotype (nG1m1), such as G1m17.1, G1m3, G1m3.1, G1m3.2 or G1m3.1.2. More preferably, the allotype is selected from the group consisting
20 of the G1m17.1 or G1m3 allotype. The antibody may be engineered in the Fc-domain to enhance or nihilate binding to Fc-gamma receptors, as summarized by Saunders et al. *Front. Immunol.* **2019**, 10, doi: 10.3389/fimmu.2019.01296 and Ward et al., *Mol. Immunol.* **2015**, 67, 131–141. For example, the combination of Leu234Ala and Leu235Ala (commonly called LALA mutations) eliminate FcγRIIa binding. Elimination of binding to Fc-gamma receptors can also be achieved by
25 mutation of the N297 amino acid to any other amino acid except asparagine, by mutation of the T299 amino acid to any other amino acid except threonine or serine, or by enzymatic deglycosylation or trimming of the fully glycosylated antibody with for example PNGase F or an endoglycosidase. The immunoglobulins can be derived from any species, including human, murine, or rabbit origin. Each chain contains distinct sequence domains.

30 **[0040]** A percentage of “sequence identity” may be determined by comparing the two sequences, optimally aligned over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. A sequence “at least 85% identical to a reference sequence” is a
35 sequence having, on its entire length, 85%, or more, for instance 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity with the entire length of the reference sequence.

[0041] The term “CDR” refers to complementarity-determining region: the specificity of the antibody resides in the structural complementarity between the antibody combining site and the antigenic determinant. Antibody combining sites are made up of residues that are primarily from the
40 hypervariable or complementarity determining regions (CDRs). Occasionally, residues from non-

hypervariable or framework regions (FR) influence the overall domain structure and hence the combining site. Complementarity Determining Regions or CDRs therefore refer to amino acid sequences which together define the binding affinity and specificity of the natural Fv region of a native immunoglobulin binding site. The light and heavy chains of an immunoglobulin each have three CDRs, designated CDR1-L, CDR2-L, CDR3-L and CDR1-H, CDR2-H, CDR3-H, respectively. A conventional antibody antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. "CDR"

[0042] The term "monoclonal antibody" or "mAb" as used herein refers to an antibody molecule of a single amino acid sequence, which is directed against a specific antigen, and is not to be construed as requiring production of the antibody by any particular method. A monoclonal antibody may be produced by a single clone of B cells or hybridoma, but may also be recombinant, i.e. produced by protein engineering.

[0043] The term "chimeric antibody" refers to an engineered antibody which, in its broadest sense, contains one or more regions from one antibody and one or more regions from one or more other antibodies. In an embodiment, a chimeric antibody comprises a VH domain and a VL domain of an antibody derived from a non-human animal, in association with a CH domain and a CL domain of another antibody, in an embodiment, a human antibody. As the non-human animal, any animal such as mouse, rat, hamster, rabbit or the like can be used. A chimeric antibody may also denote a multispecific antibody having specificity for at least two different antigens.

[0044] The term "humanised antibody" refers to an antibody which is wholly or partially of non-human origin and which has been modified to replace certain amino acids, for instance in the framework regions of the VH and VL domains, in order to avoid or minimize an immune response in humans. The constant domains of a humanized antibody are most of the time human CH and CL domains. "Fragments" of (conventional) antibodies comprise a portion of an intact antibody, in particular the antigen binding region or variable region of the intact antibody. Examples of antibody fragments include Fv, Fab, F(ab')₂, Fab', dsFv, (dsFv)₂, scFv, sc(Fv)₂, diabodies, bispecific and multispecific antibodies formed from antibody fragments. A fragment of a conventional antibody may also be a single domain antibody, such as a heavy chain antibody or VHH.

[0045] As used herein "CEA" or "CEACAM5" designates the "carcino-embryonic antigen-related cell adhesion molecule 5", also known as "CD66e" (Cluster of Differentiation 66e). CEA and CEACAM5 are used exchangeably herein.

Description of figures

[0046] Figure 1 shows a representative set of reactive groups (F) that when present in a biomolecule lead to connecting group Z (**1a-1h**) upon reaction with reactive group Q. Functional groups F can be naturally present or may be artificially introduced (engineered) into a biomolecule at any position of choice.

[0047] Figure 2 schematically displays how an antibody conjugate can be obtained from any monoclonal antibody in a two-stage process. In the first stage, an azido-modified UPD-Gal or UDP-GalNAc may be attached to the monoclonal antibody in a one pot process involving (a) trimming of

the glycan by an endoglycosidase (to the core GlcNAc) and (b) attachment of the azido-sugar under the action of a glycosyltransferase (a galactosyltransferase or a mutant thereof or a GalNAc-transferase), thereby generating a β -glycosidic 1-4 linkage between the azido-modified GalNAc and GlcNAc. In the second stage, the azido-modified antibody is reacted with an appropriately functionalized cyclooctyne, thereby generating the antibody conjugate.

[0048] Figure 3 shows several structures of derivatives of UDP sugars of galactosamine, which may be modified with e.g. a 3-mercaptopropionyl group (**2a**), an azidoacetyl group (**2b**), or an azidodifluoroacetyl group (**2c**) at the 2-position, or with an azido group at the 6-position of N-acetyl galactosamine (**2d**).

[0049] Figure 4 shows cyclooctynes (**A-T**) suitable for metal-free click chemistry, which are preferred options for reactive moiety Q.

[0050] Figure 5 shows a set of exemplary toxic payloads for conjugation to various CEA-targeting monoclonal antibodies according to the invention. Point of attachment of a linker (to an amino group present in the payload) is indicated with an arrow. Preferred conjugates according to the present invention contain the payloads, including point of attachment, as shown in figure 5.

[0051] Figure 6 shows the structures of various camptothecins, which are preferred payloads in the context of the present invention.

[0052] Figure 7 depicts the structures of the BCN-linker drugs containing a Val-Cit-PABC or Val-Ala-PABC cleavable linker used to prepare ADCs via click chemistry to the azidosugar-remodeled antibodies, with the following payloads (**3** = exatecan, **4** = MMAE, **5a** = calicheamicin γ_1^I , **5b** = glycine-calicheamicin γ_1^I , **6** = 6-aminohexanoyl-maytansinoid).

[0053] Figure 8 shows the structures of twelve members of the human CEA family, which belong to the CEACAM subgroup. CEACAM-encoded proteins generally have one variable (V)-like Ig domain, identified as the N domain (except CEACAM16 with two N domains), but they differ in the number of constant C2-like Ig domains, identified as A or B, as well as the membrane anchorage.

[0054] Figure 9a shows binding of tusamitamab and labetuzumab to hCEACAM5 (hCD66e), cCEACAM5 (cCD66e) and hCEACAM6 (hCD66c) corrected for the background. Both mAbs show similar affinity for hCEACAM5, do not show any binding to hCEACAM6 and only tusamitamab shows affinity for cCEACAM5. Figure 9b shows binding of tusamitamab and labetuzumab-based ADCs to hCEACAM5. All ADCs still show binding to hCEACAM5, comparable to the corresponding mAb.

[0055] Figure 10 shows the cytotoxicity data on CEACAM5 expressing MKN-45 cells. All ADCs, except negative control B12-3, induce cytotoxicity with increasing concentrations.

[0056] Figure 11a shows the in vivo efficacy data of the colorectal IGR-034P PDX model. First effects of tumour regression is observed for both tusamitamab-3 and labetuzumab-3 after 7 days. Figure 11b shows the body weight plot. Fig. 11c and 11d show the same plots for the full experiment (35 days).

[0057] Figure 12a shows the in vivo efficacy data of the colorectal IGR-002P PDX model. Tumor regression is observed for both labetuzumab-3 and tusamitamab-3 at 10 mg/kg. No effect was observed for tusamitamab-DM4, nor for B12-3 the negative control in this model. Figure 12b shows

-12-

body weight loss for the three groups in which no tumor growth inhibition was observed. This is due to the cachectic nature of the PDX model. Body weight was very constant for the groups that responded well to the treatment.

[0058] Figure 13 shows the in vivo efficacy data obtained with labeltuzumab-3 for several PDX models. Fig. 13(A) – (C) show the plots corresponding to colorectal IC-003P PDX model. Clear tumor regression is observed for this model. Fig. 13(D) – (F) show the plots corresponding to colorectal LRB-010P PDX model. Tumor growth inhibition was observed in the treated group, with a greatly enhanced overall survival (at least 1.5x compared to vehicle). Fig. 13(G) – (I) show the plots corresponding to lung NIC-014 PDX model. Tumor regression is observed for this fast growing model with at least a twofold increased overall survival. Fig. 13(J) – (L) show the plots corresponding to pancreatic SA-083 PDX model. Tumor growth inhibition is also observed for this indication. Tumor growth inhibition is confirmed in another pancreatic PDX model. Fig. 13(M) – (O) show the plots corresponding to pancreatic IM-PAN-07 PDX model with also high tumor growth inhibition.

[0059] Figure 14 shows the in vivo tolerability data obtained with labeltuzumab-3. The body weight change (%) over time is depicted, which was more or less equal for all doses tested. Even up to a dose of 140 mg/kg, no toxicity was observed. Hence the MTD is above 140 mg/kg.

The invention

[0060] In a first aspect, the invention concerns antibody-conjugates of general structure (1)



wherein:


- AB is an antibody capable of targeting CEA-expressing tumours;
- L is a linker that links Z to D;
- Z is a connecting group;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolbenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- b is 0 or 1;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

[0061] Also contemplated within the present invention are salts, preferably pharmaceutically acceptable salts, of the antibody-conjugate according to structure (1).

[0062] In a second aspect, the invention concerns a process for preparing the antibody-conjugate

-13-

according to the invention, comprising reacting the compound according to general structure (2) with an antibody (3). The compound according to general structure (2) comprises a reactive moiety Q and the antibody a reactive moiety F which is capable of reacting with Q in a conjugation reaction, wherein Q and F react to form connecting group Z. In this reaction, a conjugate according to general structure (1) is formed. The process according to this aspect this concerns the following bioconjugation reaction:




[0063] Here below, the antibody-conjugate according to structure (1) is first defined. The structural features of the antibody-conjugate according to structure (1) also apply to the compound according to structure (2) and the antibody according to structure (3), as those are unchanged in the conjugation reaction except for reactive moieties F and Q, which are transformed into connecting group Z upon reaction of the compound according to structure (2) with an antibody according to structure (3).

[0064] In a third aspect, the invention concerns the application antibody-conjugate according to structure (1), for targeting CEA-expressing cells. Related thereto, the invention concerns the first medical use and second medical use of the antibody-conjugate according to structure (1).

[0065] As will be understood by the skilled person, the definition of the chemical moieties, as well as their preferred embodiments, apply to all aspects of the invention.

Antibody-conjugate of general structure (1)

[0066] In a first aspect, the invention concerns antibody-conjugates of general structure (1):



wherein:

- AB is an antibody capable of targeting CEA-expressing tumours;
- b is 0 or 1;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- Z is a connecting group;
- L is a linker that links Z to D;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

Antibody AB

[0067] The antibody-conjugate according to the invention contains an antibody that is capable of targeting CEA-expressing cells, in particular tumour cells. Carcinoembryonic antigen (CEA) is a known target for cancer treatment, and is encoded by the CEACAM5 gene. The term “expressing” is used as common in the art, and refers to overexpression of the target with respect to the expression in healthy tissue. Antibodies capable of targeting CEA-expressing tumours may also be referred to as anti-CEA antibodies, CEA-targeting antibodies or CEA-binding antibodies. Anti-CEA antibodies selectively bind to CEA-expressing cells.

[0068] Anti-CEA antibodies are known in the art, and any suitable one can be used in the context of the present invention. Preferred antibodies are selected from the list consisting of labetuzumab, A5/A240VL, D8/A240VL, PR1A3, CH1A1A/pAC18, CH1A1B/pAC18, AB173, AB14AB72, AB15AB73, NCC-CO-413, NCC-CO-308, NCC-CO-432, NCC-CO-411, A5B7, 1G9, CEA6, T84.66M5A, M5B, MFE-23, SM3E, 12-140-1, 15-1-32, tusamitamab, MT111, and humanized or functional analogues thereof. More preferably, the antibody is PR1A3, SM3E, huMAb2-4, tusamitamab or labetuzumab, even more preferably the antibody is tusamitamab or labetuzumab, most preferably the antibody is labetuzumab.

[0069] Herein, the Fc regions of these antibodies may have one or more mutations, such as 0 – 10 mutations or 0 – 5 mutations. Especially preferred are mutations that change binding to the FcRn receptor to modulate half-life of the antibody. For example, inclusion of mutations Met-to-Tyr, Ser-to-Thr and Thr-to-Glu in the region of amino acids 254 – 260 of the heavy chain, often called YTE, in the IgG1 Fc results in a ~11-fold higher binding of the antibody to human FcRn, thereby increasing the circulation half-life with a factor ~3.5. For example, the YTE mutations for tusamitamab are Met255Tyr, Ser257Thr and Thr259Glu, and for labetuzumab, the YTE mutations are Met254Tyr, Ser256Thr and Thr258Glu. Therefore, in one embodiment, the antibody has an apparent human FcRn binding affinity $K_{D,app}$ of below 2.5×10^{-6} M, preferably in the range of $0.05 - 0.99 \times 10^{-6}$ M, more preferably in the range of $0.1 - 0.49 \times 10^{-6}$ M, most preferably in the range of $0.2 - 0.4 \times 10^{-6}$ M. The apparent binding affinity $K_{D,app}$ may be determined according to Mackness et al. *MABS*, **2019**, 11(7), 1276-1288. In a preferred embodiment, the antibody AB is the YTE mutant of the preferred antibodies defined here above or below.

[0070] Labetuzumab is also known as hMN-14 and may also be defined as containing a light chain sequence according to SEQ ID No. 38 and a heavy chain sequence according to SEQ ID No. 36 or 37, preferably SEQ ID No. 36, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0071] Tusamitamab is also known as huMAb2-3 or as SAR408377 and may also be defined as containing a light chain sequence according to SEQ ID No. 43 and a heavy chain sequence according to SEQ ID No. 41 or 42, preferably SEQ ID No. 41, wherein the sequence identity is at

least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0072] M5B may also be defined as containing a light chain sequence according to SEQ ID No. 8 and a heavy chain sequence according to SEQ ID No. 7, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0073] D8 may also be defined as containing a light chain sequence according to SEQ ID No. 15 and a heavy chain sequence according to SEQ ID No. 14, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0074] SM3E may also be defined as containing a light chain sequence according to SEQ ID No. 19 and a heavy chain sequence according to SEQ ID No. 18, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0075] PR1A3 may also be defined as containing a light chain sequence according to SEQ ID No. 23 and a heavy chain sequence according to SEQ ID No. 22, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0076] huMAb2-4 may also be defined as containing a light chain sequence according to SEQ ID No. 47 and a heavy chain sequence according to SEQ ID No. 46, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. In a particularly preferred embodiment, the antibody according to this embodiment is combined with a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably with a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0077] Alternatively, the antibody is defined by its V_L and V_H domains, which together form the variable domain that binds to the antigen. Thus, in a preferred embodiment, the antibody contains a V_L domain selected from the group consisting of SEQ ID No. 2, 6, 10, 17, 21, 28, 31, 35, 40, 45, 49, 50 and 51, and a V_H domain selected from the group consisting of SEQ ID No. 1, 3, 5, 9, 11, 13, 16, 20, 24, 26, 30, 34, 39, 44 and 48, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a further preferred embodiment, the antibody contains a V_L domain selected from the group consisting of SEQ ID No. 17, 21, 35 and 40, and a V_H domain selected from the group consisting of SEQ ID No. 16, 20, 34 and 39, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a more preferred embodiment, the antibody contains a V_L domain selected from the group consisting of SEQ ID No. 35 and 40, and a V_H domain selected from the group consisting of SEQ ID No. 34 and 39, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a most preferred embodiment, the antibody contains a V_L domain of SEQ ID No. 35, and a V_H domain of SEQ ID No. 34, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %.

[0078] The aforementioned sequence identities refer to the complete sequence of the V_L and V_H domains. While the entire sequences of these domains allow for some variation in the sequence without jeopardizing the binding to CEA, it is preferred that the complementarity-determining regions (CDRs) have a higher sequence identity, to ensure that the binding to CEA is not significantly jeopardized. The location of the CDRs is given in the tables below. Thus, it is preferred that the antibody contains a V_L domain selected from the group consisting of SEQ ID No. 2, 6, 10, 17, 21, 28, 31, 35, 40, 45, 49, 50 and 51, and a V_H domain selected from the group consisting of SEQ ID No. 1, 3, 5, 9, 11, 13, 16, 20, 24, 26, 30, 34, 39, 44 and 48, preferably a V_L domain selected from the group consisting of SEQ ID No. 17, 21, 35 and 40, and a V_H domain selected from the group consisting of SEQ ID No. 16, 20, 34 and 39, more preferably a V_L domain selected from the group consisting of SEQ ID No. 35 and 40, and a V_H domain selected from the group consisting of SEQ ID No. 34 and 39, most preferably a V_L domain of SEQ ID No. 35, and a V_H domain of SEQ ID No. 34, wherein the sequence identity of the CDR is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %. The skilled person understands that the V_L and V_H domains identified above can be combined with a suitable constant domain to form a complete antibody.

[0079] Preferred V_L domains

SEQ ID No.	CDR (amino acid numbers within the sequence)
2	24-34 50-56 89-97

-17-

6	24-38 54-60 93-101
10	23-36 52-62 97-106
17	26-30 49-51 90-95
21	25-34 50-52 89-98
28	24-34 49-56 89-98
31	24-34 49-56 89-98
35	24-34 50-56 89-96
40	26-31 49-51 88-96
45	26-31 49-51 88-96
49	24-34 50-56 89-99
50	24-34 50-56 89-99
51	24-34 50-56 89-99

[0080] Preferred V_H domains

SEQ ID No.	CDR (amino acid numbers within the sequence)
1	31-35 50-66 99-112
3	31-35 50-66 99-110
5	31-36 50-65 99-110
9	31-35 50-68 101-110
11	31-35 50-62 99-108
13	31-35 50-66 99-108
16	26-32 52-57 100-108
20	20-35 50-66 97-110
24	31-35 50-66 99-110
26	31-35 50-66 99-110
30	31-35 50-65 99-110
34	31-35 50-66 99-108
39	26-33 51-58 97-109
44	26-33 51-58 97-109
48	26-35 50-68 99-114

[0081] In an especially preferred embodiment, the antibody contains a V_L domain of SEQ ID No. 2 and a V_H domain of SEQ ID No. 1, or a V_L domain of SEQ ID No. 6 and a V_H domain of SEQ ID No. 3 or 5, or a V_L domain of SEQ ID No. 10 and a V_H domain of SEQ ID No. 9, or a V_L domain of SEQ ID No. 17 and a V_H domain of SEQ ID No. 16, or a V_L domain of SEQ ID No. 21 and a V_H domain of SEQ ID No. 20, or a V_L domain of SEQ ID No. 31 and a V_H domain of SEQ ID No. 30, or a V_L domain of SEQ ID No. 35 and a V_H domain of SEQ ID No. 34, or a V_L domain of SEQ ID No. 40 and a V_H domain of SEQ ID No. 39, or a V_L domain of SEQ ID No. 45 and a V_H domain of SEQ ID No. 44, or a V_L domain of SEQ ID No. 49, 50 or 51 and a V_H domain of SEQ ID No. 48. Herein,

the sequence identities as defined above for the complete sequence as well as for the CDR apply.

[0082] Alternatively, the antibody is defined by its light and heavy chains, which together form the antibody. Thus, in a preferred embodiment, the antibody contains a light chain selected from the group consisting of SEQ ID No. 8, 15, 19, 23, 29, 33, 38, 43 and 47, and a heavy chain selected from the group consisting of SEQ ID No. 4, 7, 12, 14, 18, 22, 25, 27, 32, 36, 37, 41, 42 and 46, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a further preferred embodiment, the antibody contains a light chain selected from the group consisting of SEQ ID No. 19, 23, 38 and 43, and a heavy chain selected from the group consisting of SEQ ID No. 18, 22, 36, 37, 41 and 42, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a more preferred embodiment, the antibody contains a light chain selected from the group consisting of SEQ ID No. 38 and 43, and a heavy chain selected from the group consisting of SEQ ID No. 36 and 41, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %. In a most preferred embodiment, the antibody contains a light chain selected from the group consisting of SEQ ID No. 38, and a heavy chain selected from the group consisting of SEQ ID No. 36, wherein the sequence identity is at least 70 %, preferably at least 75 % or at least 80 %, more preferably at least 85 % or at least 90 % or at least 95 %, most preferably at least 99 % or even 100 %.

[0083] The aforementioned sequence identities refer to the complete sequence of the light and heavy chains. While the entire sequences of these chains allow for some variation in the sequence without jeopardizing the binding to CEA, it is preferred that the CDRs have a higher sequence identity, to ensure that the binding to CEA is not significantly jeopardized. The location of the CDRs is given in the tables below. Thus, it is preferred that the antibody contains a light chain selected from the group consisting of SEQ ID No. 8, 15, 19, 23, 29, 33, 38, 43 and 47, and a heavy chain selected from the group consisting of SEQ ID No. 4, 7, 12, 14, 18, 22, 25, 27, 32, 36, 37, 41, 42 and 46, wherein the sequence identity of the CDR is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.

[0084] Preferred light chains

SEQ ID No.	CDR (amino acid numbers within the sequence)
8	24-38 54-60 93-101
15	23-36 52-62 97-106
19	26-30 49-51 90-95
23	25-34 50-52 89-98
29	24-34 49-56 89-98
33	24-34 49-56 89-98
38	24-34 50-56 89-96

43	26-31 49-51 88-96
47	26-31 49-51 88-96

[0085] Preferred heavy chains

SEQ ID No.	CDR (amino acid numbers within the sequence)
4	31-35 50-66 99-110
7	31-36 50-65 99-110
12	31-35 50-62 99-108
14	31-35 50-66 99-108
18	26-32 52-57 100-108
22	20-35 50-66 97-110
25	31-35 50-66 99-110
27	31-35 50-66 99-110
32	31-35 50-65 99-110
36	31-35 50-66 99-108
37	31-35 50-66 99-108
41	26-33 51-58 97-109
42	26-33 51-58 97-109
46	26-33 51-58 97-109

[0086] In an especially preferred embodiment, the antibody contains a light chain of SEQ ID No. 8 and a heavy chain of SEQ ID No. 7, or a light chain of SEQ ID No. 15 and a heavy chain of SEQ ID No. 14, or a light chain of SEQ ID No. 19 and a heavy chain of SEQ ID No. 18, or a light chain of SEQ ID No. 23 and a heavy chain of SEQ ID No. 22, or a light chain of SEQ ID No. 33 and a heavy chain of SEQ ID No. 32, or a light chain of SEQ ID No. 38 and a heavy chain of SEQ ID No. 36 or 37, preferably 36, or a light chain of SEQ ID No. 43 and a heavy chain of SEQ ID No. 41 or 42, preferably 41, or a light chain of SEQ ID No. 47 and a heavy chain of SEQ ID No. 46. Herein, the sequence identities as defined above for the complete sequence as well as for the CDR apply.

Linker L⁶

[0087] In case reactive group F is directly connected to the antibody, or even part of the antibody structure, linker L⁶ that connects AB to F (for antibodies of structure (3)) or AB to Z (for conjugates of structure (1) is absent and b = 0. This is for example the case for cysteine conjugation and lysine conjugation. Alternatively, reactive group F may also be introduced onto the antibody using a linker L⁶ that connects AB to F (for antibodies of structure (3)) or AB to Z (for conjugates of structure (1), in which case L⁶ is present and b = 1. In case L⁶ is present, reactive group F is typically introduced at the glycan of the antibody. This is for example the case for conjugation via an artificially introduced reactive group F, such as for example using transglutaminase or by enzymatic glycan modification (e.g. glycosyltransferase or α -1,3-mannosyl-glycoprotein-2- β -N-acetylglucosaminyl-

transferase). For example, a modified sugar residue $S(F)_x$ may be introduced at the glycan, extending the glycan with one monosaccharide residue S, which introduces x reactive groups F on the glycan of an antibody. In a most preferred embodiment, conjugation occurs via the glycan of the antibody and $b = 1$. The site of conjugation is preferably at the heavy chain of the antibody.

5 **[0088]** If present, L^6 is a linker that links AB to F or Z, and is represented by $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-S-(L^7)_{w'}$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$. Typically, L^6 is at least partly formed by the glycan of the antibody. All recombinant antibodies, generated in mammalian host systems, contain the
10 conserved N-glycosylation site at the asparagine residue at or close to position 297 of the heavy chain, which is modified by a glycan of the complex type. This naturally occurring glycosylation site of antibodies is preferably used, but other glycosylation sites, including artificially introduced ones, may also be used for the connection of linker L^6 . Thus, in a preferred embodiment, L^6 is connected to an amino acid of the antibody which is located at a position in the range of 250 – 350 of the heavy
15 chain, preferably in the range of 280 – 310 of the heavy chain, more preferably in the range of 295 – 300 of the heavy chain, most preferably at position 297 of the heavy chain.

[0089] The $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-$ of L^6 is the glycan of the antibody, or part thereof. The $-\text{GlcNAc}(\text{Fuc})_b-(\text{G})_j-$ of the glycan thus typically originates from the original antibody, wherein GlcNAc is an N-acetylglucosamine moiety and Fuc is a fucose moiety. Fuc is typically bound to
20 GlcNAc via an α -1,6-glycosidic bond. Normally, antibodies may ($w = 1$) or may not be fucosylated ($w = 0$). In the context of the present invention, the presence of a fucosyl moiety is irrelevant, and similar effects are obtained with fucosylated ($w = 1$) and non-fucosylated ($w = 0$) antibody conjugates. The GlcNAc residue may also be referred to as the core-GlcNAc residue and is the monosaccharide that is directly attached to the peptide part of the antibody.

25 **[0090]** S may be directly connected to the core-GlcNAc(Fuc)_w moiety, i.e. $j = 0$, meaning that the remainder of the glycan is removed from the core-GlcNAc(Fuc)_w moiety before S is attached. Such trimming of glycans is well-known in the art and can be achieved by the action of an endoglycosidase. Alternatively, there are one or more monosaccharide residues present in between the core-GlcNAc(Fuc)_w moiety and S, i.e. j is an integer in the range of 1 – 10, preferably $j = 2 - 5$.
30 In a preferred embodiment, $(\text{G})_j$ is an oligosaccharide fraction comprising j monosaccharide residues G, wherein j is an integer in the range of 2 – 5. $(\text{G})_j$ is connected to the GlcNAc moiety of $\text{GlcNAc}(\text{Fuc})_w$, typically via a β -1,4 bond. In a preferred embodiment, j is 3, 4 or 5. Although any monosaccharide that may be present in a glycan may be employed as G, each G is preferably individually selected from the group consisting of galactose, glucose, N-acetylgalactosamine, N-acetylglucosamine, mannose and N-acetylneuraminic acid. More preferred options for G are galactose, N-acetylglucosamine, mannose. The inventors found that antibody-conjugates having j
35 below 4 show no or hardly any binding to the Fc-gamma receptor, while antibody conjugates having j in the range of 4 – 10 do bind to the Fc-gamma receptor. Thus, by selecting a certain value for j, the desired extent of binding to the Fc-gamma receptor can be obtained. It is thus preferred that $j =$

0, 3, 4, 5, 6, 7, 8, 9 or 10, more preferably $j = 0, 3, 4$ or 5, most preferably the antibody is trimmed and $j = 0$.

[0091] S is a sugar or sugar derivative. The term "sugar derivative" is herein used to indicate a derivative of a monosaccharide sugar, i.e. a monosaccharide sugar comprising substituents and/or functional groups. Suitable examples for S include glucose (Glc), galactose (Gal), mannose (Man), fucose (Fuc), amino sugars and sugar acids, e.g. glucosamine (GlcNH₂), galactosamine (GalNH₂), N-acetylglucosamine (GlcNAc), N-acetylgalactosamine (GalNAc), sialic acid (Sia) which is also referred to as N-acetylneuraminic acid (NeuNAc), and N-acetylmuramic acid (MurNAc), glucuronic acid (GlcA) and iduronic acid (IdoA). Preferably, S is selected from Glc, Gal, GlcNAc and GalNAc.

10 In an especially preferred embodiment, S is GalNAc.

[0092] x is an integer that denotes the number of connecting groups Z (for conjugate (1)) or reactive groups F (for antibody (3)) that are attached to sugar (derivative) S. Thus, the antibody according to the invention contains a moiety S comprising x reactive moieties F. Each of these reactive moieties F are reacted with a reactive moiety Q of the compound according to general structure (2), such that x connecting groups Z are formed and x compounds according to general structure (2) are attached to a single occurrence of S. x is 1 or 2, preferably $x = 1$.

15

[0093] Connecting group Z (for conjugate (1)) or reactive group F (for antibody (3)) may be attached directly to S, or there may be a linker L⁷ present in between S and Z or F. L⁷ is a linker that links S with Z. L⁷ may be present ($w' = 1$ or 2) or absent ($w' = 0$). Typically, each moiety Z may be connected to S via a linker L⁷, thus in one embodiment $w' = 0$ of x. Preferably, L⁷ is absent and each connecting moiety Z is directly attached to S. If present, L⁷ may be selected from -N(H)C(O)CH₂-, -N(H)C(O)CF₂- or -CH₂-. In a preferred embodiment, $x = 1$ and $w' = 0$ or 1, most preferably $x = 1$ and $w' = 0$.

20

[0094] y is an integer that denotes the number of sugar(s) (derivative(s)) S, each having x reactive groups F or connected to x connecting groups Z, that are connected to the antibody. y is 1, 2, 3 or 4, preferably $y = 2$ or 4, most preferably $y = 2$. Thus, the antibody contains y moieties S, each of which comprises x reactive moieties F. Each of these reactive moieties F are reacted with reactive moiety Q of the compound according to general structure (2), such that $x + y$ connecting groups Z are formed and $x + y$ compounds according to general structure (2) are attached to a single antibody. Each compound according to general structure (2) may contain multiple payloads, e.g. by virtue of branching nitrogen atom N* in L. It is preferred that each compound according to general structure (2) contains 1 or 2 occurrences of D, most preferably 2 occurrences of D. In an especially preferred embodiment, linker L¹ contains a branching nitrogen atom N* to which a second occurrence of D is connected.

25

30

[0095] The amount of payload (D) molecules attached to a single antibody is known in the art as the DAR (drug-antibody ratio). In the context of the present invention, it is preferred that DAR is an integer in the range 1 – 8, more preferably 2 or 4, most preferably DAR = 4. Alternatively worded, the DAR is preferably an integer in the range $(x + y)$ to $[(x + y) \times 2]$, most preferably DAR = $[(x + y) \times 2]$. With preferred values for x of 1 and y of 2, the DAR is preferably 4. It will be appreciated that these are theoretical DAR values, and in practice the DAR may slightly deviate from this value, by

35

40

virtue of incomplete conjugation. Typically, the conjugates are obtained as a stochastic mixture of antibody-drug conjugates, with DAR values varying between individual conjugates, and depending on the conjugation technique used the DAR may have a broad distribution (e.g. DAR = 0 – 10) or a narrow distribution (e.g. DAR = 3-4). In case of such mixture, DAR often refers to the average DAR of the mixture. This is well-known in the art of bioconjugation. However, in case the conjugation occurs via the glycan (i.e. $b = 1$ and L^6 is present), the antibody-conjugates according to the invention have a close-to-theoretical DAR. For example, when the theoretical DAR is 4, DAR values above 3.6 or even above 3.8 are readily obtained, indicating that most antibodies in the reaction mixture have reacted completely and have a DAR of 4.

Connecting group Z

[0096] Z is a connecting group, which covalently connects both parts of the conjugate according to the invention. The term “connecting group” herein refers to the structural element, resulting from the reaction between Q and F, connecting one part of the conjugate with another part of the same conjugate. As will be understood by the person skilled in the art, the nature of a connecting group depends on the type of reaction with which the connection between the parts of said compound is obtained. As an example, when the carboxyl group of $R-C(O)-OH$ is reacted with the amino group of H_2N-R' to form $R-C(O)-N(H)-R'$, R is connected to R' via connecting group Z, and Z may be represented by the group $-C(O)-N(H)-$. Since connecting group Z originates from the reaction between Q and F, it can take any form.

[0097] Since more than one reactive moiety F can be present or introduced in an antibody, the antibody-conjugate according to the present invention may contain per biomolecule more than one payload D, such as 1 – 8 payloads D, preferably 1, 2, 3 or 4 payloads D, more preferably 2 or 4 payloads D. The number of payloads is typically an even integer, in view of the symmetric nature of antibodies. In other words, when one side of the antibody is functionalized with F, the symmetrical counterpart will also be functionalized. Alternatively, in case naturally occurring thiol groups of the cysteine residues of a protein are used as F, the value of m can be anything and may vary between individual conjugates.

[0098] In a compound according to structure (1), connecting group Z connects D via linker L to AB, optionally via L^6 . Numerous reactions are known in the art for the attachment of a reactive group Q to a reactive group F. Consequently, a wide variety of connecting groups Z may be present in the conjugate according to the invention. In one embodiment, the reactive group Q is selected from the options described above, preferably as depicted in Figure 1, and complementary reactive groups F and the thus obtained connecting groups Z are known to a person skilled in the art. Several examples of suitable combinations of F and Q, and of connecting group Z^3 that will be present in a bioconjugate when a linker-conjugate comprising Q is conjugated to a biomolecule comprising a complementary reactive group F, are shown in Figure 1.

[0099] For example, when F comprises or is a thiol group, complementary groups Q include N-maleimidyl groups and alkenyl groups, and the corresponding connecting groups Z are as shown

in Figure 1. When F comprises or is a thiol group, complementary groups Q also include allenamide groups.

[0100] For example, when F comprises or is an amino group, complementary groups Q include ketone groups and activated ester groups, and the corresponding connecting groups Z are as shown in Figure 1.

[0101] For example, when F comprises or is a ketone group, complementary groups Q include (O-alkyl)hydroxylamino groups and hydrazine groups, and the corresponding connecting groups Z are as shown in Figure 1.

[0102] For example, when F comprises or is an alkynyl group, complementary groups Q include azido groups, and the corresponding connecting group Z is as shown in Figure 1.

[0103] For example, when F comprises or is an azido group, complementary groups Q include alkynyl groups, and the corresponding connecting group Z is as shown in Figure 1.

[0104] For example, when F comprises or is a cyclopropenyl group, a trans-cyclooctene group or a cyclooctyne group, complementary groups Q include tetrazinyl groups, and the corresponding connecting group Z is as shown in Figure 1. In these particular cases, Z is only an intermediate structure and will expel N₂, thereby generating a dihydropyridazine (from the reaction with alkene) or pyridazine (from the reaction with alkyne).


[0105] Additional suitable combinations of F and Q, and the nature of resulting connecting group Z3 are known to a person skilled in the art, and are e.g. described in G.T. Hermanson, "Bioconjugate Techniques", Elsevier, 3rd Ed. 2013 (ISBN:978-0-12-382239-0), in particular in Chapter 3, pages 229 - 258, incorporated by reference. A list of complementary reactive groups suitable for bioconjugation processes is disclosed in Table 3.1, pages 230 - 232 of Chapter 3 of G.T. Hermanson, "Bioconjugate Techniques", Elsevier, 3rd Ed. 2013 (ISBN:978-0-12-382239-0), and the content of this Table is expressly incorporated by reference herein.

[0106] In a preferred embodiment, connecting group Z is obtained by a cycloaddition or a nucleophilic reaction, preferably wherein the cycloaddition is a [4+2] cycloaddition or a 1,3-dipolar cycloaddition or the nucleophilic reaction is a Michael addition or a nucleophilic substitution. Such a cycloaddition or nucleophilic reaction occurs via a reactive group F, connected to S, and reactive group Q, connected to D via L. Conjugation reactions via cycloadditions or nucleophilic reactions are known to the skilled person, and the skilled person is capable of selecting appropriate reaction partners F and Q, and will understand the nature of the resulting connecting group Z.

[0107] In a first preferred embodiment, Z is formed by a cycloaddition. Preferred cycloadditions are a (4+2)-cycloaddition (e.g. a Diels-Alder reaction) or a (3+2)-cycloaddition (e.g. a 1,3-dipolar cycloaddition). Preferably, the conjugation is the Diels-Alder reaction or the 1,3-dipolar cycloaddition. The preferred Diels-Alder reaction is the inverse-electron demand Diels-Alder cycloaddition. In another preferred embodiment, the 1,3-dipolar cycloaddition is used, more preferably the alkyne-azide cycloaddition, and most preferably wherein Q is or comprises an alkyne group and F is an azido group. Cycloadditions, such as Diels-Alder reactions and 1,3-dipolar cycloadditions are known in the art, and the skilled person knows how to perform them.

-24-

[0108] Preferably, Z contains a moiety selected from the group consisting of a triazole, a cyclohexene, a cyclohexadiene, a [2.2.2]-bicyclooctadiene, a [2.2.2]-bicyclooctene, an isoxazoline, an isoxazolidine, a pyrazoline, a piperazine, a thioether, an amide or an imide group. Triazole moieties are especially preferred to be present in Z. In one embodiment, Z comprises a (hetero)cycloalkene moiety, i.e. formed from Q comprising a (hetero)cycloalkyne moiety. In an alternative embodiment, Z comprises a (hetero)cycloalkane moiety, i.e. formed from Q comprising a (hetero)cycloalkene moiety. In a preferred embodiment, Z has the structure (Z1):



(Z1)

10

[0109] Herein, the bond depicted as --- is a single bond or a double bond. Furthermore:


- ring Z is obtained by a cycloaddition, preferably ring Z is selected from (Za) – (Zj) defined below, wherein the carbon atoms labelled with ** correspond to the two carbon atoms of the bond depicted as --- of (Z1) to which ring Z is fused;
- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- Y^2 is $C(R^{31})_2$, O, S, $S^{(+)}R^{31}$, $S(O)R^{31}$, $S(O)=NR^{31}$ or NR^{31} , wherein $S^{(+)}$ is a cationic sulphur atom counterbalanced by $B^{(-)}$, wherein $B^{(-)}$ is an anion, and wherein each R^{31} individually is R^{15} or a connection with D, connected via L;
- u is 0, 1, 2, 3, 4 or 5;
- u' is 0, 1, 2, 3, 4 or 5, wherein $u + u' = 0, 1, 2, 3, 4, 5, 6, 7$ or 8;
- v = an integer in the range 8 – 16;
- Ring A is formed by the cycloaddition, and is preferably selected from (Za) – (Zj).

30

[0110] In case the bond depicted as --- is a double bond, it is preferred that $u + u' = 4, 5, 6, 7$ or 8. Preferably, the wavy bond labelled with * is connected to S and the wavy bond labelled with ** is connected to L.


-25-

[0111] It is especially preferred that Z comprises a (hetero)cycloalkene moiety, i.e. the bond depicted as --- is a double bond. In a preferred embodiment, Z is selected from the structures (Z2) – (Z20), depicted here below:




[0112] Herein, the connection to L is depicted with the wavy bond. B⁽⁻⁾ is an anion, preferably a pharmaceutically acceptable anion. Ring Z is formed by the cycloaddition reaction, and preferably is a triazole, a cyclohexene, a cyclohexadiene, a [2.2.2]-bicyclooctadiene, a [2.2.2]-bicyclooctene, an isoxazoline, an isoxazolidine, a pyrazoline or a piperazine. Most preferably, ring Z is a triazole ring. Ring Z may have the structure selected from (Za) – (Zm) depicted below, wherein the carbon atoms labelled with ** correspond to the two carbon atoms of the (hetero)cycloalkane ring of (Z2) – (Z20) to which ring Z is fused. Preferred rings Z are selected from (Za) – (Zj), more preferably from (Za), (Zd) and (Zh), most preferably ring Z has structure (Za). Since the connecting group Z is formed by reaction with a (hetero)cycloalkyne in the context of the present embodiment, the bound depicted above as --- is a double bond.


-26-



[0113] In a further preferred embodiment, Z is selected from the structures (Z21) – (Z38) and (Z38a), depicted here below:



-27-



[0114] Herein, the connection to L is depicted with the wavy bond. In structure (Z38), $B^{(-)}$ is an anion, preferably a pharmaceutically acceptable anion. Ring Z is selected from structures (Za) - (Zm), preferably from structures (Za) - (Zj), as defined above.

[0115] In a preferred embodiment, Z comprises a (hetero)cyclooctene moiety or a (hetero)cycloheptene moiety, preferably according to structure (Z8), (Z26), (Z27), (Z28) or (Z37) or (Z38a), more preferably according to structure (Z8), (Z26), (Z27), (Z28) or (Z37), which are optionally substituted. Each of these preferred options for Z are further defined here below.


[0116] Thus, in a preferred embodiment, Z comprises a heterocycloheptene moiety according to structure (Z37), which is optionally substituted. Preferably, the heterocycloheptene moiety according to structure (Z37) is not substituted.

[0117] In a preferred embodiment, Z comprises a (hetero)cyclooctene moiety according to structure (Z8), more preferably according to (Z29), which is optionally substituted. Preferably, the cyclooctene moiety according to structure (Z8) or (Z29) is not substituted. In the context of the present embodiment, Z preferably comprises a (hetero)cyclooctene moiety according to structure (Z39) as shown below, wherein V is $(CH_2)_l$ and l is an integer in the range of 0 to 10, preferably in the range of 0 to 6. More preferably, l is 0, 1, 2, 3 or 4, more preferably l is 0, 1 or 2 and most preferably l is 0 or 1. In the context of group (Z39), l is most preferably 1. Most preferably, Z is according to structure (Z42), defined further below.


[0118] In an alternative preferred embodiment, Z comprises a (hetero)cyclooctene moiety according to structure (Z26), (Z27) or (Z28), which are optionally substituted. In the context of the present embodiment, Z preferably comprises a (hetero)cyclooctene moiety according to structure (Z40) or (Z41) as shown below, wherein Y^1 is O or NR^{11} , wherein R^{11} is independently selected from the group consisting of hydrogen, a linear or branched $C_1 - C_{12}$ alkyl group or a $C_4 - C_{12}$ (hetero)aryl group. The aromatic rings in (Z40) are optionally O-sulfonylated at one or more positions, whereas the rings of (Z41) may be halogenated at one or more positions. Preferably, the (hetero)cyclooctene moiety according to structure (Z40) or (Z41) is not further substituted. Most preferably, Z is according to structure (Z43), defined further below.

-28-

[0119] In an alternative preferred embodiment, Z comprises a heterocycloheptenyl group and is according to structure (Z37).



[0120] In an especially preferred embodiment, Z comprises a cyclooctenyl group and is according to structure (Z42):



Herein:


- the bond labelled with * is connected to S and the wavy bond labelled with ** is connected to L;
- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_5 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- R^{18} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- R^{19} is selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups, the alkyl groups optionally being interrupted by one of more hetero-atoms selected from the group consisting of O, N and S, wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are independently optionally substituted, or R^{19} is a second occurrence of Z (or Q) or D connected via a spacer moiety; and
- l is an integer in the range 0 to 10.

[0121] In a preferred embodiment of the group according to structure (Z42), R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $C_1 - C_6$ alkyl groups, $C_5 - C_6$

-29-

(hetero)aryl groups, wherein R^{16} is hydrogen or $C_1 - C_6$ alkyl, more preferably R^{15} is independently selected from the group consisting of hydrogen and $C_1 - C_6$ alkyl, most preferably all R^{15} are H. In a preferred embodiment of the group according to structure (Z42), R^{18} is independently selected from the group consisting of hydrogen, $C_1 - C_6$ alkyl groups, most preferably both R^{18} are H. In a preferred embodiment of the group according to structure (Z42), R^{19} is H. In a preferred embodiment of the group according to structure (Z42), l is 0 or 1, more preferably l is 1.

[0122] In an especially preferred embodiment, Z comprises a (hetero)cyclooctynyl group and is according to structure (Z43):





Herein:

- the bond labelled with * is connected to S and the wavy bond labelled with ** is connected to L;
- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_5 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- Y is N or CR^{15} .













[0123] In a preferred embodiment of the group according to structure (Z43), R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_6$ alkyl groups, $C_5 - C_6$ (hetero)aryl groups, wherein R^{16} is hydrogen or $C_1 - C_6$ alkyl, more preferably R^{15} is independently selected from the group consisting of hydrogen and $-S(O)_3^{(-)}$. In a preferred embodiment of the group according to structure (Z43), Y is N or CH, more preferably Y = N.

[0124] In an especially preferred embodiment, Z comprises a heterocycloheptynyl group and is according to structure (Z37) or (Z38a), preferably according to structure (Z37), wherein ring Z is a triazole.


-30-




[0125] In an alternative preferred embodiment, Z comprises a (hetero)cycloalkane moiety, i.e. the bond depicted as --- is a single bond. The (hetero)cycloalkane group may also be referred to as a heterocycloalkanyl group or a cycloalkanyl group, preferably a cycloalkanyl group, wherein the (hetero)cycloalkanyl group is optionally substituted. Preferably, the (hetero)cycloalkanyl group is a (hetero)cyclopropanyl group, a (hetero)cyclobutanyl group, a norbornane group, a norbornene group, a (hetero)cycloheptanyl group, a (hetero)cyclooctanyl group, a (hetero)cyclononyl group or a (hetero)cyclodecanyl group, which may all optionally be substituted. Especially preferred are (hetero)cyclopropanyl groups, (hetero)cycloheptanyl group or (hetero)cyclooctanyl groups, wherein the (hetero)cyclopropanyl group, the *trans*-(hetero)cycloheptanyl group or the (hetero)cyclooctanyl group is optionally substituted. Preferably, Z comprises a cyclopropanyl moiety according to structure (Z44), a heterocyclobutane moiety according to structure (Z45), a norbornane or norbornene group according to structure (Z46), a (hetero)cycloheptanyl moiety according to structure (Z47) or a (hetero)cyclooctanyl moiety according to structure (Z48). Herein, Y^3 is selected from $\text{C}(\text{R}^{23})_2$, NR^{23} or O, wherein each R^{23} is individually hydrogen, $\text{C}_1 - \text{C}_8$ alkyl or is connected to L, optionally via a spacer, and the bond labelled --- is a single or double bond. In a further preferred embodiment, the cyclopropanyl group is according to structure (Z49). In another preferred embodiment, the (hetero)cycloheptane group is according to structure (Z50) or (Z51). In another preferred embodiment, the (hetero)cyclooctane group is according to structure (Z52), (Z53), (Z54), (Z55) or (Z56).




[0126] Herein, the R group(s) on Si in (Z50) and (Z51) are typically alkyl or aryl, preferably C₁-C₆ alkyl. Ring Z is typically selected from structures (Zn) – (Zu), wherein the carbon atoms labelled with ** correspond to the two carbon atoms of the (hetero)cycloalkane ring of (Z44) – (Z56) to which ring Z is fused, and the carbon a carbon labelled with * is directly connected to the peptide chain of the antibody. Preferred rings Z are selected from (Zo) – (Zr). Since the connecting group Z is formed by reaction with a (hetero)cycloalkene in the context of the present embodiment, the bond depicted above as --- is a single bond.



[0127] In a second preferred embodiment, Z is formed by a nucleophilic reaction, preferably by a nucleophilic substitution or a Michael addition, preferably by a Michael addition. A preferred Michael reaction is the thiol-maleimide ligation, most preferably wherein Q is maleimide and F is a thiol group. Preferably, the thiol is present in the sidechain of a cysteine residue. In a preferred embodiment, connection group Z comprises a succinimidyl ring or its ring-opened succinic acid amide derivative. Preferred options for connection group Z comprise a moiety selected from (Z57) – (Z71) depicted here below.



-32-



[0128] Herein, the wavy bond(s) labelled with an * is connected to the antibody Ab, optionally via a linker, and the wavy bond without label to the payload, optionally via a linker. In addition, R^{29} is C_{1-12} alkyl, preferably C_{1-4} alkyl, most preferably ethyl, and X^1 is O or S, preferably $X^1 = O$. The nitrogen atom labelled with ** in (Z67)-(Z71) corresponds to the nitrogen atom of the side chain of a lysine residue of the antibody. The carbon atoms of the phenyl group of (Z69) and (Z70) are optionally substituted, preferably optionally fluorinated.

[0129] In a preferred embodiment, connection group Z comprise a moiety selected from (Z1) – (Z71).

10 Linker L

[0130] Linker L connects payload D with connecting group Z (in the conjugate according to structure (1)) or connects payload D with reactive group Q (in the compound according to structure (2)). Linkers are known in the art and may be cleavable or non-cleavable. Linker L preferably contains a self-immolative group or cleavable linker, comprising a peptide spacer and a *para*-aminobenzylloxycarbonyl (PABC) moiety or derivative thereof.

[0131] In a preferred embodiment, linker L as the structure $-(L^1)_n-(L^2)_o-(L^3)_p-(L^4)_q-$, wherein $(L^4)_q$ is connect to payload D and $(L^1)_n$ is connected to Z or Q. Herein L^1 , L^2 , L^3 and L^4 are linkers or linking units and each of n, o, p and q are individually 0 or 1, wherein $n + o + p + q$ is at least 1. In a preferred embodiment, at least linkers L^1 and L^2 are present (i.e. $n = 1$; $o = 1$; $p = 0$ or 1 ; $q = 0$ or 1), more preferably linkers L^1 , L^2 and L^3 are present and L^4 is either present or not (i.e. $n = 1$; $o = 1$; $p = 1$; $q = 0$ or 1). In one embodiment, linkers L^1 , L^2 , L^3 and L^4 are present (i.e. $n = 1$; $o = 1$; $p = 1$; $q = 1$). In one embodiment, linkers L^1 , L^2 and L^3 are present and L^4 is not (i.e. $n = 1$; $o = 1$; $p = 1$; $q = 0$).


[0132] A linker, especially linker L^1 , may contain one or more branch-points for attachment of multiple payloads to a single connecting group. In a preferred embodiment, the linker of the conjugate according to the invention contains a branching moiety. A “branching moiety” in the context of the present invention refers to a moiety that is embedded in a linker connecting three moieties. In other words, the branching moiety comprises at least three bonds to other moieties, typically one bond connecting to Z or Q, one bond to the payload D and one bond to a second payload D. The branching moiety, if present, is preferably embedded in linker L^1 , more preferably

part of Sp^2 or as the nitrogen atom of NR^{13} . Any moiety that contains at least three bonds to other moieties is suitable as branching moiety in the context of the present invention. In a preferred embodiment, the branching moiety is selected from a carbon atom, a nitrogen atom, a phosphorus atom, a (hetero)aromatic ring, a (hetero)cycle or a polycyclic moiety. Most preferably, the branching moiety is a nitrogen atom.

Linker L^1

[0133] Linker L^1 is either absent ($n = 0$) or present ($n = 1$). Preferably, linker L^1 is present and $n = 1$. L^1 may for example be selected from the group consisting of linear or branched C_1 - C_{200} alkylene groups, C_2 - C_{200} alkenylene groups, C_2 - C_{200} alkynylene groups, C_3 - C_{200} cycloalkylene groups, C_5 - C_{200} cycloalkenylene groups, C_8 - C_{200} cycloalkynylene groups, C_7 - C_{200} alkylarylene groups, C_7 - C_{200} arylalkylene groups, C_8 - C_{200} arylalkenylene groups, C_9 - C_{200} arylalkynylene groups. Optionally the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups may be substituted, and optionally said groups may be interrupted by one or more heteroatoms, preferably 1 to 100 heteroatoms, said heteroatoms preferably being selected from the group consisting of O, $S(O)_{y'}$ and NR^{21} , wherein y' is 0, 1 or 2, preferably $y' = 2$, and R^{21} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups.

[0134] In a preferred embodiment, linker L^1 contains a polar group. Such a polar group may be selected from (poly)ethylene glycoldiamines (e.g. 1,8-diamino-3,6-dioxaoctane or equivalents comprising longer ethylene glycol chains), (poly)ethylene glycol or (poly)ethylene oxide chains, (poly)propylene glycol or (poly)propylene oxide chains and 1, z' -diaminoalkanes (wherein z' is the number of carbon atoms in the alkane, preferably $z' = 1 - 10$), $-(O)_a-C(O)-NH-S(O)_2-NR^{13}-$ (as further defined below, see structure (23)), $-C(S(O)_3^{(-)})-$, $-C(C(O)_2^{(-)})-$, $-S(O)_2-$, $-P(O)_2^{(-)}$, $-O(CH_2CH_2O)_t-$, $-NR^{30}(CH_2CH_2NR^{30})_t-$, and the following two structures:




[0135] The polar group may also contain an amino acid, preferably selected from Arg, Glu, Asp, Ser and Thr. Herein, a and R^{13} are further defined below for structure (23). t is an integer in the range of integer in the range of 0 - 15, preferably 1 - 10, more preferably 2 - 5, most preferably $t = 2$ or 4. Each R^{30} is individually H, C_{1-12} alkyl, C_{1-12} aryl, C_{1-12} alkaryl or C_{1-12} aralkyl. Linker L^1 may contain more than one such polar group, such as at least two polar groups. The polar group may also be present in a branch of linker L^1 , which branches off a branching moiety as defined elsewhere. Preferable, a nitrogen or carbon atom is used as branching moiety. It is especially preferred to have a $-O(CH_2CH_2O)_t-$ polar group present in a branch.

[0136] In a preferred embodiment, Linker L^1 is or comprises a sulfamide group, preferably a

-34-

sulfamide group according to structure (23):




(23)

[0137] The wavy lines represent the connection to the remainder of the compound, typically to Q and L², L³, L⁴ or D, preferably to Q and L². Preferably, the (O)_aC(O) moiety is connected to Q and the NR¹³ moiety to L², L³, L⁴ or D, preferably to L².

[0138] In structure (23), a = 0 or 1, preferably a = 1, and R¹³ is selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups and C₃ – C₂₄ (hetero)arylalkyl groups, the C₁ – C₂₄ alkyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups and C₃ – C₂₄ (hetero)arylalkyl groups optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR¹⁴ wherein R¹⁴ is independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups, or R¹³ is D connected to N via a spacer moiety, preferably Sp² as defined below, in one embodiment D is connected to N via –(B)_e–(A)_f–(B)_g–C(O)–.

[0139] In a preferred embodiment, R¹³ is hydrogen or a C₁ – C₂₀ alkyl group, more preferably R¹³ is hydrogen or a C₁ – C₁₆ alkyl group, even more preferably R¹³ is hydrogen or a C₁ – C₁₀ alkyl group, wherein the alkyl group is optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR¹⁴, preferably O, wherein R¹⁴ is independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups. In a preferred embodiment, R¹³ is hydrogen. In another preferred embodiment, R¹³ is a C₁ – C₂₀ alkyl group, more preferably a C₁ – C₁₆ alkyl group, even more preferably a C₁ – C₁₀ alkyl group, wherein the alkyl group is optionally interrupted by one or more O-atoms, and wherein the alkyl group is optionally substituted with an -OH group, preferably a terminal -OH group. In this embodiment it is further preferred that R¹³ is a (poly)ethylene glycol chain comprising a terminal -OH group. In another preferred embodiment, R¹³ is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl and t-butyl, more preferably from the group consisting of hydrogen, methyl, ethyl, n-propyl and i-propyl, and even more preferably from the group consisting of hydrogen, methyl and ethyl. Yet even more preferably, R¹³ is hydrogen or methyl, and most preferably R¹³ is hydrogen.

[0140] In a preferred embodiment, L¹ is according to structure (24):



(24)

[0141] Herein, a and R¹³ are as defined above, Sp¹ and Sp² are independently spacer moieties and b and c are independently 0 or 1. Preferably, b = 0 or 1 and c = 1, more preferably b = 0 and c = 1. In one embodiment, spacers Sp¹ and Sp² are independently selected from the group consisting of linear or branched C₁–C₂₀₀ alkylene groups, C₂–C₂₀₀ alkenylene groups, C₂–C₂₀₀ alkynylene groups, C₃–C₂₀₀ cycloalkylene groups, C₅–C₂₀₀ cycloalkenylene groups, C₈–C₂₀₀ cycloalkynylene

groups, C₇-C₂₀₀ alkylarylene groups, C₇-C₂₀₀ arylalkylene groups, C₈-C₂₀₀ arylalkenylene groups and C₉-C₂₀₀ arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted. When the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups are interrupted by one or more heteroatoms as defined above, it is preferred that said groups are interrupted by one or more O-atoms, and/or by one or more S-S groups.

[0142] More preferably, spacer moieties Sp¹ and Sp², if present, are independently selected from the group consisting of linear or branched C₁-C₁₀₀ alkylene groups, C₂-C₁₀₀ alkenylene groups, C₂-C₁₀₀ alkynylene groups, C₃-C₁₀₀ cycloalkylene groups, C₅-C₁₀₀ cycloalkenylene groups, C₈-C₁₀₀ cycloalkynylene groups, C₇-C₁₀₀ alkylarylene groups, C₇-C₁₀₀ arylalkylene groups, C₈-C₁₀₀ arylalkenylene groups and C₉-C₁₀₀ arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted.

[0143] Even more preferably, spacer moieties Sp¹ and Sp², if present, are independently selected from the group consisting of linear or branched C₁-C₅₀ alkylene groups, C₂-C₅₀ alkenylene groups, C₂-C₅₀ alkynylene groups, C₃-C₅₀ cycloalkylene groups, C₅-C₅₀ cycloalkenylene groups, C₈-C₅₀ cycloalkynylene groups, C₇-C₅₀ alkylarylene groups, C₇-C₅₀ arylalkylene groups, C₈-C₅₀ arylalkenylene groups and C₉-C₅₀ arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted.

[0144] Yet even more preferably, spacer moieties Sp¹ and Sp², if present, are independently selected from the group consisting of linear or branched C₁-C₂₀ alkylene groups, C₂-C₂₀ alkenylene groups, C₂-C₂₀ alkynylene groups, C₃-C₂₀ cycloalkylene groups, C₅-C₂₀ cycloalkenylene groups, C₈-

C₂₀ cycloalkynylene groups, C₇-C₂₀ alkylarylene groups, C₇-C₂₀ arylalkylene groups, C₈-C₂₀ arylalkenylene groups and C₉-C₂₀ arylalkynylene groups, the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted.

[0145] In these preferred embodiments it is further preferred that the alkylene groups, alkenylene groups, alkynylene groups, cycloalkylene groups, cycloalkenylene groups, cycloalkynylene groups, alkylarylene groups, arylalkylene groups, arylalkenylene groups and arylalkynylene groups are unsubstituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, preferably O, wherein R¹⁶ is independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups, preferably hydrogen or methyl.

[0146] Most preferably, spacer moieties Sp¹ and Sp², if present, are independently selected from the group consisting of linear or branched C₁-C₂₀ alkylene groups, the alkylene groups being optionally substituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, wherein R¹⁶ is independently selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₂ - C₂₄ alkenyl groups, C₂ - C₂₄ alkynyl groups and C₃ - C₂₄ cycloalkyl groups, the alkyl groups, alkenyl groups, alkynyl groups and cycloalkyl groups being optionally substituted. In this embodiment, it is further preferred that the alkylene groups are unsubstituted and optionally interrupted by one or more heteroatoms selected from the group of O, S and NR¹⁶, preferably O and/or S-S, wherein R¹⁶ is independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups, preferably hydrogen or methyl.

[0147] Preferred spacer moieties Sp¹ and Sp² thus include -(CH₂)_r-, -(CH₂CH₂)_r-, -(CH₂CH₂O)_r-, -(OCH₂CH₂)_r-, -(CH₂CH₂O)_rCH₂CH₂-, -CH₂CH₂(OCH₂CH₂)_r-, -(CH₂CH₂CH₂O)_r-, -(OCH₂CH₂CH₂)_r-, -(CH₂CH₂CH₂O)_rCH₂CH₂CH₂- and -CH₂CH₂CH₂(OCH₂CH₂CH₂)_r-, wherein r is an integer in the range of 1 to 50, preferably in the range of 1 to 40, more preferably in the range of 1 to 30, even more preferably in the range of 1 to 20 and yet even more preferably in the range of 1 to 15. More preferably n is 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10, more preferably 1, 2, 3, 4, 5, 6, 7 or 8, even more preferably 1, 2, 3, 4, 5 or 6, yet even more preferably 1, 2, 3 or 4.

[0148] Alternatively, preferred linkers L¹ may be represented by -(W)_k-(A)_d-(B)_e-(A)_f-(C(O))_g-, wherein:

- d = 0 or 1, preferably d = 1;
- e = an integer in the range 0 – 10, preferably e = 0, 1, 2, 3, 4, 5 or 6, preferably an integer in the range 1 – 10, most preferably e = 1, 2, 3 or 4;
- f = 0 or 1, preferably f = 0;
- wherein d + e + f is at least 1, preferably in the range 1 – 5; and preferably wherein d + f is at least 1, preferably d + f = 1.

-37-

- $g = 0$ or 1 , preferably $g = 1$;
- $k = 0$ or 1 , preferably $k = 1$;
- A is a sulfamide group according to structure (23);
- B is a $-\text{CH}_2-\text{CH}_2-\text{O}-$ or a $-\text{O}-\text{CH}_2-\text{CH}_2-$ moiety, or $(\text{B})_e$ is a $-(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$ or a $-(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-$ moiety, wherein $e1$ is defined the same way as e ;
- W is $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{OC}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})\text{O}-$, $-\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})-$, $-\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$ or $-(4\text{-Ph})\text{CH}_2\text{NHC}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$, preferably wherein W is $-\text{OC}(\text{O})\text{NH}-$, $-\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$ or $-\text{C}(\text{O})\text{NH}-$, and wherein m is an integer in the range $0 - 10$, preferably $m = 0, 1, 2, 3, 4, 5$ or 6 , most preferably $m = 2$ or 3 ;
- preferably wherein L^1 is connected to Q via $(\text{W})_k$ and to L^2 , L^3 , L^4 or D, preferably to L^2 , via $(\text{C}(\text{O}))_g$, preferably via $\text{C}(\text{O})$.

[0149] In the context of the present embodiment, the wavy lines in structure (23) represent the connection to the adjacent groups such as $(\text{W})_k$, $(\text{B})_e$ and $(\text{C}(\text{O}))_g$. It is preferred that A is according to structure (23), wherein $a = 1$ and $\text{R}^{13} = \text{H}$ or a $\text{C}_1 - \text{C}_{20}$ alkyl group, more preferably $\text{R}^{13} = \text{H}$ or methyl, most preferably $\text{R}^{13} = \text{H}$.

[0150] Preferred linkers L^1 have structure $-(\text{W})_k-(\text{A})_d-(\text{B})_e-(\text{A})_f-(\text{C}(\text{O}))_g-$, wherein:

- (a) $k = 0$; $d = 1$; $g = 1$; $f = 0$; $\text{B} = -\text{CH}_2-\text{CH}_2-\text{O}-$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (b) $k = 1$; $\text{W} = -\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$; $m = 2$; $d = 0$; $(\text{B})_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $f = 0$; $g = 1$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 1$.
- (c) $k = 1$; $\text{W} = -\text{OC}(\text{O})\text{NH}-$; $d = 0$; $\text{B} = -\text{CH}_2-\text{CH}_2-\text{O}-$; $g = 1$; $f = 0$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (d) $k = 1$; $\text{W} = -\text{C}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$; $m = 2$; $d = 0$; $(\text{B})_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $f = 0$; $g = 1$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 4$.
- (e) $k = 1$; $\text{W} = -\text{OC}(\text{O})\text{NH}-$; $d = 0$; $(\text{B})_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $g = 1$; $f = 0$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 4$.
- (f) $k = 1$; $\text{W} = -(4\text{-Ph})\text{CH}_2\text{NHC}(\text{O})(\text{CH}_2)_m\text{C}(\text{O})\text{NH}-$, $m = 3$; $d = 0$; $(\text{B})_e = -(\text{CH}_2-\text{CH}_2-\text{O})_{e1}-\text{CH}_2-\text{CH}_2-$; $g = 1$; $f = 0$; $e1 = 1, 2, 3$ or 4 , preferably $e1 = 4$.
- (g) $k = 0$; $d = 0$; $g = 1$; $f = 0$; $\text{B} = -\text{CH}_2-\text{CH}_2-\text{O}-$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (h) $k = 1$; $\text{W} = -\text{C}(\text{O})\text{NH}-$; $d = 0$; $g = 1$; $f = 0$; $\text{B} = -\text{CH}_2-\text{CH}_2-\text{O}-$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.

[0151] In a preferred embodiment, linker L^1 comprises a branching nitrogen atom, which is located in the backbone between Q or Z and $(\text{L}^2)_o$ and which contains a further moiety D as substituent, which is preferably linked to the branching nitrogen atom via a linker. An example of a branching nitrogen atom is the nitrogen atom NR^{13} in structure (23), wherein R^{13} is connected to a second occurrence of D via a spacer moiety. Alternatively, a branching nitrogen atoms may be located within L^1 according to structure $-(\text{W})_k-(\text{A})_d-(\text{B})_e-(\text{A})_f-(\text{C}(\text{O}))_g-\text{N}^*[-(\text{A})_d-(\text{B})_e-(\text{A})_f-(\text{C}(\text{O}))_g-]_2$, wherein A, B, W, d, e, f, g and k are as defined above and individually selected for each occurrence, and N^* is the branching nitrogen atoms, to which two instances of $-(\text{A})_d-(\text{B})_e-(\text{A})_f-(\text{C}(\text{O}))_g-$ are connected. Herein, both $(\text{C}(\text{O}))_g$ moieties are connected to $-(\text{L}^2)_o-(\text{L}^3)_p-(\text{L}^4)_q-\text{D}$, wherein L^2 , L^3 , L^4 , o, p, q and

D are as defined above and are each selected individually. In a preferred embodiment, each of L^2 , L^3 , L^4 , o, p, q and D are the same for both moieties connected to $(C(O))_g$.

[0152] Preferred linkers L^1 comprising a branching nitrogen atom have structure $-(W)_k-(A)_d-(B)_e-(A)_f-(C(O))_g-N^*[-(A')_d'-(B')_e'-(A')_f'-(C(O))_g']_2$ wherein:

- 5 (i) $k = d = g = e' = 1$; $f = d' = g' = 0$; $W = -C(O)-$; $B = B' = -CH_2-CH_2-O-$; A is according to structure (23) with $a = 0$ and $R^{13} = H$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.
- (j) $k = d = g = e' = g' = 1$; $f = d' = 0$; $W = -C(O)-$; $B = B' = -CH_2-CH_2-O-$; A is according to structure (23) with $a = 0$ and $R^{13} = H$; $e = 1, 2, 3$ or 4 , preferably $e = 2$.

10 Linker L^2

[0153] Linker L^2 is either absent ($o = 0$) or present ($o = 1$). Preferably, linker L^2 is present and $o = 1$. Linker L^2 is a peptide spacer. The peptide spacer is preferably defined by $(NH-CR^{17}-CO)_n$, wherein R^{17} represents an amino acid side chain as known in the art. Herein, the amino acid may be a natural or a synthetic amino acid. Preferably, the amino acid(s) are all in their L-configuration.

15 n is an integer in the range of $1 - 5$, preferably in the range of $2 - 5$. Thus, the peptide spacer preferably contains $1 - 5$ amino acids. Preferably, the peptide is a dipeptide ($n = 2$), tripeptide ($n = 3$) or tetrapeptide ($n = 4$), most preferably the peptide spacer is a dipeptide. Although any peptide spacer may be used, preferably the peptide spacer is selected from Val-Cit, Val-Ala, Val-Lys, Val-Arg, AcLys-Val-Cit, AcLys-Val-Ala, Glu-Val-Ala, Asp-Val-Ala, iGlu-Val-Ala, Glu-Val-Cit, Asp-Val-Cit,


20 iGlu-Val-Cit, Phe-Cit, Phe-Ala, Phe-Lys, Phe-Arg, Ala-Lys, Leu-Cit, Ile-Cit, Trp-Cit, Ala-Ala-Asn, Ala-Asn, Gly-Gly-Phe-Gly and Lys, more preferably Val-Cit, Val-Ala, Glu-Val-Ala, Val-Lys, Phe-Cit, Phe-Ala, Phe-Lys, Ala-Ala-Asn, more preferably Val-Cit, Val-Ala, Ala-Ala-Asn, most preferably Val-Cit or Val-Ala. Herein, AcLys is acetyllysine and iGlu is isoglutamate. In one embodiment, $L^2 = \text{Val-Cit}$. In one embodiment, $L^2 = \text{Val-Ala}$.

25 **[0154]** R^{17} represents the amino acid side chain, preferably selected from the side chains of alanine, cysteine, aspartic acid, glutamic acid, phenylalanine, glycine, histidine, isoleucine, lysine, acetyllysine, leucine, methionine, asparagine, pyrrolysine, proline, glutamine, arginine, serine, threonine, selenocysteine, valine, tryptophan, tyrosine and citrulline. Preferred amino acid side chains are those of Val, Cit, Ala, Lys, Arg, AcLys, Phe, Leu, Ile, Trp, Glu, Asp and Asn, more preferably from the side chains of Val, Cit, Ala, Glu and Lys. Alternatively worded, R^{17} is preferably selected from CH_3 (Ala), $CH_2CH(CH_3)_2$ (Leu), $CH_2CH_2CH_2NHC(O)NH_2$ (Cit), $CH_2CH_2CH_2CH_2NH_2$ (Lys), $CH_2CH_2CH_2NHC(O)CH_3$ (AcLys), $CH_2CH_2CH_2NHC(=NH)NH_2$ (Arg), CH_2Ph (Phe), $CH(CH_3)_2$ (Val), $CH(CH_3)CH_2CH_3$ (Ile), $CH_2C(O)NH_2$ (Asn), $CH_2CH_2C(O)OH$ (Glu), $CH_2C(O)OH$ (Asp) and $CH_2(1H\text{-indol-3-yl})$ (Trp). Especially preferred embodiments of R^{17} are CH_3 (Ala),

30 $CH_2CH_2CH_2NHC(O)NH_2$ (Cit), $CH_2CH_2CH_2CH_2NH_2$ (Lys), $CH_2CH_2C(O)OH$ (Glu) and $CH(CH_3)_2$ (Val). Most preferably, R^{17} is CH_3 (Ala), $CH_2CH_2CH_2NHC(O)NH_2$ (Cit), $CH_2CH_2CH_2CH_2NH_2$ (Lys), or $CH(CH_3)_2$ (Val).

[0155] In an especially preferred embodiment, the peptide spacer may be represented by general structure (L3):

-39-




(L3)

[0156] Herein, R^{17} is as defined above, preferably R^{17} is CH_3 (Val) or $\text{CH}_2\text{CH}_2\text{CH}_2\text{NHC(O)NH}_2$ (Cit). The wavy lines indicate the connection to $(L^1)_n$ and $(L^3)_p$, preferably L^2 according to structure (L3) is connected to $(L^1)_n$ via NH and to $(L^3)_p$ via C(O).

Linker L^3

[0157] Linker L^3 is either absent ($p = 0$) or present ($p = 1$). Preferably, linker L^3 is present and $p = 1$. Linker L^3 is a self-cleavable spacer, also referred to as self-immolative spacer. Preferably, L^3 is para-aminobenzyloxycarbonyl (PABC) derivative, more preferably a PABC derivative according to structure (L4):



(L4)

[0158] Herein, the wavy lines indicate the connection to Q or Z, L^1 or L^2 , and to L^4 or D. Typically, the PABC derivative is connected via NH to Q, Z, L^1 or L^2 , preferably to L^2 , and via O to L^4 or D.

[0159] A is a 5- or 6-membered aromatic or heteroaromatic ring, preferably a 6-membered aromatic or heteroaromatic ring. Suitable 5-membered rings are oxazole, thiazole and furan. Suitable 6-membered rings are phenyl and pyridyl. In a preferred embodiment, A is 1,4-phenyl, 2,5-pyridyl or 3,6-pyridyl. Most preferably, A is 1,4-phenyl.

[0160] R^{21} is selected from H, R^{26} , C(O)OH and C(O)R^{26} , wherein R^{26} is $\text{C}_1 - \text{C}_{24}$ (hetero)alkyl groups, $\text{C}_3 - \text{C}_{10}$ (hetero)cycloalkyl groups, $\text{C}_2 - \text{C}_{10}$ (hetero)aryl groups, $\text{C}_3 - \text{C}_{10}$ alkyl(hetero)aryl groups and $\text{C}_3 - \text{C}_{10}$ (hetero)arylalkyl groups, which are optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR^{28} wherein R^{28} is independently selected from the group consisting of hydrogen and $\text{C}_1 - \text{C}_4$ alkyl groups. Preferably, R^{26} is $\text{C}_3 - \text{C}_{10}$ (hetero)cycloalkyl or polyalkylene glycol. The polyalkylene glycol is preferably a polyethylene glycol or a polypropylene glycol, more preferably $-(\text{CH}_2\text{CH}_2\text{O})_s\text{H}$ or $-(\text{CH}_2\text{CH}_2\text{CH}_2\text{O})_s\text{H}$. The polyalkylene glycol is most preferably a polyethylene glycol, preferably $-(\text{CH}_2\text{CH}_2\text{O})_s\text{H}$, wherein s is an integer in the range 1 – 10, preferably 1 – 5, most preferably $s = 1, 2, 3$ or 4. More preferably, R^{21} is H or C(O)R^{26} , wherein $R^{26} = 4\text{-methyl-piperazine}$ or morpholine. Most preferably, R^{21} is H.

Linker L^4

[0161] Linker L^4 is either absent ($q = 0$) or present ($q = 1$). Preferably, linker L^4 is present and $q = 1$. Linker L^4 is selected from:

- an aminoalkanoic acid spacer according to the structure $-\text{NR}^{22}-(\text{C}_z\text{-alkylene})-\text{C(O)-}$, wherein z is an integer in the range 1 – 20 and R^{22} is H or $\text{C}_1 - \text{C}_4$ alkyl;

-40-

- an ethyleneglycol spacer according to the structure $\text{--NR}^{22}\text{--}(\text{CH}_2\text{--CH}_2\text{--O})_{e6}\text{--}(\text{CH}_2)_{e7}\text{--C(O)--}$, wherein $e6$ is an integer in the range 1 – 10, $e7$ is an integer in the range 1 – 3 and R^{22} is H or $\text{C}_1\text{--C}_4$ alkyl; and
- an diamine spacer according to the structure $\text{--NR}^{22}\text{--}(\text{C}_z\text{-alkylene})\text{--NR}^{22}\text{--}(\text{C(O)})_h\text{--}$, wherein h is 0 or 1, z is an integer in the range 1 – 20 and R^{22} is H or $\text{C}_1\text{--C}_4$ alkyl.

[0162] Linker L^4 may be an aminoalkanoic acid spacer, i.e. $\text{--NR}^{22}\text{--}(\text{C}_z\text{-alkylene})\text{--C(O)--}$, wherein z is an integer in the range 1 to 20, preferably 1 – 10, most preferably 1 – 6. Herein, the aminoalkanoic acid spacer is typically connected to L^3 via the nitrogen atom and to D via the carbonyl moiety. Preferred linkers L^4 are selected from 6-aminohexanoic acid (Ahx, $z = 5$), β -alanine ($z = 2$) and glycine (Gly, $z = 1$), even more preferably 6-aminohexanoic acid or glycine. In one embodiment, $\text{L}^4 = 6\text{-aminohexanoic acid}$. In one embodiment, $\text{L}^4 = \text{glycine}$. Herein, R^{22} is H or $\text{C}_1\text{--C}_4$ alkyl, preferably R^{22} is H or methyl, most preferably R^{22} is H.

[0163] Alternatively, linker L^4 may be an ethyleneglycol spacer according to the structure $\text{--NR}^{22}\text{--}(\text{CH}_2\text{--CH}_2\text{--O})_{e6}\text{--}(\text{CH}_2)_{e7}\text{--C(O)--}$, wherein $e6$ is an integer in the range 1 – 10, preferably $e6$ is in the range 2 – 6, and $e7$ is an integer in the range 1 – 3, preferably $e7$ is 2. Herein, R^{22} is H or $\text{C}_1\text{--C}_4$ alkyl, preferably R^{22} is H or methyl, most preferably R^{22} is H.

[0164] Alternatively, linker L^4 may be a diamine spacer according to the structure $\text{--NR}^{22}\text{--}(\text{C}_z\text{-alkylene})\text{--NR}^{22}\text{--}(\text{C(O)})_h\text{--}$, wherein h is 0 or 1, z is an integer in the range 1 – 20, preferably an integer in the range 2 – 6, even more preferably $z = 2$ or 5, most preferably $z = 2$. R^{22} is H or $\text{C}_1\text{--C}_4$ alkyl. Herein, R^{22} is H or $\text{C}_1\text{--C}_4$ alkyl, preferably R^{22} is H or methyl, most preferably R^{22} is methyl. Herein, h is preferably 1, in which case linker L^4 is especially suited for conjugation via a phenolic hydroxyl group present on payload D.

Payload D

[0165] D represents the target molecule D that is or is to be connected to the antibody, which is also referred to in the art as the payload. D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemeasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof. Alternatively, D is defined as a pharmaceutically active substance, such as an anti-cancer agent, preferably a cytotoxin. Preferably, D is selected from the group consisting of anthracyclines, camptothecins, maytansinoids, enediynes, amanitins, auristatins and pyrrolobenzodiazepine dimers, more preferably D is selected from the group consisting of enediynes, auristatins and camptothecins. In one embodiment, D is an enediyne. In one embodiment, D is an auristatin. In one embodiment, D is a camptothecin. Most preferably, D is a camptothecin.

[0166] In a preferred embodiment, the enediyne is selected from the group consisting of calicheamicins, esperamicins, shishijimicins and namenamicins, more preferably calicheamicin. In another preferred embodiment, the auristatin is selected from the group consisting of MMAD, MMAE, MMAF or PF-06380101, more preferably MMAE or PF-06380101. In another preferred embodiment, the camptothecin is selected from the compounds depicted in Figure 6, preferably

from the group consisting of topotecan, silatecan, cositecan, exatecan, exatecan-S, DXd, SN-38, lurtotecan, gimatecan, belotecan, rubitecan, AMDCPT and G-AMDCPT, more preferably exatecan or DXd, most preferably exatecan.

[0167] In an especially preferred embodiment, D is selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably from the group consisting of calicheamicin, MMAE, and exatecan, most preferably D is exatecan.


[0168] The compound according to general structure (2) may comprise more than one moiety D. When more than one cytotoxin D is present the cytotoxins D may be the same or different, typically they are the same. In a preferred embodiment, the compound according to general structure (2) contains 1 or 2 occurrences of D, most preferably 2 occurrence of D. Typically, the second occurrence of D is present within linker L, which may contain a branching moiety, typically a branching nitrogen atom, that is connected to the second occurrence of D. Preferably, both occurrences of D are connected to the branching moiety via the same linker. Likewise, the antibody-conjugate according to structure (1) may contain more than one moiety D per connecting group Z.

15

Preferred antibody-conjugates

[0169] Preferred antibody-conjugates according to the first aspect are selected from the group consisting of compounds (I) – (III), more preferably (II) or (III), most preferably (II). More preferred antibody-conjugates are selected from (X) – (XIII). In one especially preferred embodiment, the antibody-conjugates is selected from (Xa), (Xib), (XIlg), (XIIf) and (XIIIe). In an even more preferred embodiment, the antibody-conjugates is selected from (XI) and (XIII), more preferably (Xib) or (XIIIe), more preferably the antibody-conjugates is according to (XIII), most preferably according to (XIIIe). The structures of these antibody-conjugates are defined here below.

[0170] Antibody-conjugate (I) has the following structure:



wherein:

- AB, L⁶, Z, D, x and y are as defined above;
- L¹ is a linker represented by -(A)_d-(B)_e-(A)-(C(O))_g-, as defined above;
- L² is Val-Cit or Val-Ala;
- L³ is the PABC derivative according to structure (L4);
- L⁴ is -N-(C_z-alkylene)-C(O)- or -NR²²-(C_z-alkylene)-NR²²-, wherein z and R²² are as defined above;
- q = 0 or 1.

[0171] In the context of antibody-conjugate (I), it is preferred that for L¹, d = 1 (A according to structure (23)), it is preferred that a = 1 and R¹³ = H), e = 2, f = 0 and g = 1. In the context of antibody-conjugate (I), it is preferred that L² = Val-Cit. In the context of antibody-conjugate (I), it is preferred that for L³, R²¹ = H. In the context of antibody-conjugate (I), it is preferred that in case q = 1, then z = 1 or 5.


40

-42-

[0172] In a particularly preferred embodiment, the antibody-conjugate according to structure (I) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

5

[0173] Antibody-conjugate (II) has the following structure:




wherein:

- 10 - AB, L^6 , Z, D, x and y are as defined above;
- L^1 is a linker represented by $-(A)-(B)_e-(C(O))-$, as defined above;
- L^2 is Val-Cit or Val-Ala;
- L^3 is the PABC derivative according to structure (L4), wherein $R^{21} = H$.

15 **[0174]** In the context of antibody-conjugate (II), it is preferred that for L^1 , $e = 2$, A according to structure (23), it is preferred that $a = 1$ and $R^{13} = H$. In the context of antibody-conjugate (II), it is preferred that $L^2 = \text{Val-Cit}$.

20 **[0175]** In a particularly preferred embodiment, the antibody-conjugate according to structure (II) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

[0176] Antibody-conjugate (III) has the following structure:



25 wherein:


- AB, L^6 , Z, D, x and y are as defined above;
- L^1 is a linker represented by $-(A)-(B)_e-(C(O))-$, as defined above;
- L^2 is Val-Cit or Val-Ala;
- L^3 is the PABC derivative according to structure (L4), wherein $R^{21} = H$;
- 30 - L^4 is $-\text{NR}^{22}-(\text{C}_z\text{-alkylene})-\text{NR}^{22}-$, wherein R^{22} is as defined above and z is an integer in the range 1 to 6.

[0177] In the context of antibody-conjugate (III), it is preferred that for L^1 , $e = 2$, and with $a = 1$ and $R^{13} = H$. In the context of antibody-conjugate (III), it is preferred that $L^2 = \text{Val-Cit}$. In the context of antibody-conjugate (III), it is preferred that $z = 2$.

35 **[0178]** In a particularly preferred embodiment, the antibody-conjugate according to structure (III) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

40 **[0179]** Antibody-conjugate (X) has a linker-payload moiety according to the following structure:

-43-



(X)

wherein:


- the wavy line indicates the connection to Z;

5 - L^2 , o and D are as defined above.

[0180] L^2 may be present or absent, preferably L^2 is present and o = 1. For preferred antibody-conjugate (Xa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (Xb), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. The antibody-conjugate (X) preferably has structure (Xa).

10 **[0181]** In a particularly preferred embodiment, the antibody-conjugate according to structure (X) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

15 **[0182]** Antibody-conjugate (XI) has a linker-payload moiety according to the following structure:



(XI)

wherein:

- the wavy line indicates the connection to Z;


20 - L^2 , o and D are as defined above.

[0183] L^2 may be present or absent, preferably L^2 is present and o = 1. For preferred antibody-conjugate (XIa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (XIb), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. The antibody-conjugate (XI) preferably has structure (XIb).

25 **[0184]** In a particularly preferred embodiment, the antibody-conjugate according to structure (XI) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with calicheamicin as payload D.

30 **[0185]** Antibody-conjugate (XII) has a linker-payload moiety according to the following structure:

-44-



(XII)

wherein:

- the wavy line indicates the connection to Z;

5 - L^2 , L^4 , o , q and D are as defined above.

[0186] L^2 may be present or absent, preferably L^2 is present and $o = 1$. For preferred antibody-conjugate (XIIa), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (XIIb), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. Preferably, $R^{17} = CH_2CH_2CH_2NHC(O)NH_2$.

10 **[0187]** L^4 may be present or absent. For preferred antibody-conjugate (XIIc), $q = 0$ and L^4 is absent. For preferred antibody-conjugate (XIId), $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.

15 **[0188]** For preferred antibody-conjugate (XIIe), L^2 is according to structure (L3), R^{17} is CH_3 , $q = 0$ and L^4 is absent. For preferred antibody-conjugate (XIIf), L^2 is according to structure (L3), R^{17} is CH_3 , $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 10, preferably wherein $z = 2$, and R^{22} is H or $C_1 - C_4$ alkyl.

20 **[0189]** For preferred antibody-conjugate (XIIf), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, $q = 0$ and L^4 is absent. For preferred antibody-conjugate (XIIh), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.


25 **[0190]** In the context of antibody-conjugate (XII), it is preferred that z is an integer in the range of 1 – 10, more preferably $z = 2 - 6$, most preferably $z = 2$. In the context of antibody-conjugate (XII), it is further preferred that R^{22} is H or CH_3 .

[0191] In the context of antibody-conjugate (XII), structures (XIIg) and (XIIh) are most preferred.

30 **[0192]** In a particularly preferred embodiment, the antibody-conjugate according to structure (XII) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

-45-

[0193] Antibody-conjugate (**XIII**) has a linker-payload moiety according to the following structure:



(XIII)

wherein:

- the wavy line indicates the connection to Z;
- L^2 , L^4 , o, q and D are as defined above.

[0194] L^2 may be present or absent, preferably L^2 is present and $o = 1$. For preferred antibody-conjugate (**XIIIa**), L^2 is according to structure (L3) and R^{17} is CH_3 . For preferred antibody-conjugate (**XIIIb**), L^2 is according to structure (L3) and R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$. Preferably, $R^{17} = CH_3$.

- 10 **[0195]** L^4 may be present or absent. For preferred antibody-conjugate (**XIIIc**), $q = 0$ and L^4 is absent. For preferred antibody-conjugate (**XIIId**), $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.

- 15 **[0196]** For preferred antibody-conjugate (**XIIIe**), L^2 is according to structure (L3), R^{17} is CH_3 , $q = 0$ and L^4 is absent. For preferred antibody-conjugate (**XIIIff**), L^2 is according to structure (L3), R^{17} is CH_3 , $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 10, preferably wherein z = 2, and R^{22} is H or $C_1 - C_4$ alkyl.

- 20 **[0197]** For preferred antibody-conjugate (**XIIIg**), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, $q = 0$ and L^4 is absent. For preferred antibody-conjugate (**XIIIh**), L^2 is according to structure (L3), R^{17} is $CH_2CH_2CH_2NHC(O)NH_2$, $q = 1$ and L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-$, wherein z is an integer in the range 1 – 20, and R^{22} is H or $C_1 - C_4$ alkyl.

- 25 **[0198]** In the context of antibody-conjugate (**XIII**), it is preferred that z is an integer in the range of 1 – 10, more preferably z = 2 – 6, most preferably z = 2. In the context of antibody-conjugate (**XIII**), it is further preferred that R^{22} is H or CH_3 .

[0199] In the context of antibody-conjugate (**XIII**), structure (**XIIIe**) is most preferred.

- 30 **[0200]** In a particularly preferred embodiment, the antibody-conjugate according to structure (**XIII**) contains a payload D selected from the group consisting of calicheamicin, MMAE, PF-06380101, exatecan and DXd, more preferably a payload D selected from the group consisting of calicheamicin, MMAE, and exatecan, most preferably with exatecan as payload D.

-46-

[0201] In one particularly preferred embodiment, the antibody-conjugate according to the invention is according to structure **(XIb)** as defined above, and the antibody is labetuzumab as defined above or tusamitamab as defined above and the payload is calicheamicin.

5 **[0202]** In one particularly preferred embodiment, the antibody-conjugate according to the invention is according to structure **(XIIIe)** as defined above, and the antibody is labetuzumab as defined above or tusamitamab as defined above and the payload is exatecan.

[0203] In one particularly preferred embodiment, the antibody-conjugate according to the invention is according to structure **(XIIIe)** as defined above, and the antibody is labetuzumab as defined above and the payload is exatecan.

10 **[0204]** It is further preferred that these preferred antibody-conjugates are conjugated through the glycan, i.e. $b = 1$, more preferably a trimmed glycan, i.e. $j = 0$. Herein, it is further preferred that $S = \text{GalNAc}$ and $w' = 0$. Herein, it is further preferred that connecting group Z is formed by an azide-alkyne cycloaddition, preferably connecting group $Z = (Z39)$, wherein ring $Z = (Za)$ and $V = \text{CH}_2$. Herein, it is further preferred that $x = 1$. Herein, it is further preferred that $y = 2$, more preferably that

15 $x = 1$ and $y = 2$.

[0205] In a most preferred embodiment, the antibody-conjugate according to the invention is according to structure **(XIIIe)** as defined above, and the antibody is labetuzumab as defined above and the payload is exatecan, wherein $b = 1$, $e = 0$, $S = \text{GalNAc}$, $w' = 0$, connecting group $Z = (Z39)$, wherein ring $Z = (Za)$ and $V = \text{CH}_2$, $x = 1$ and $y = 2$.

20

Compound according to general structure (2)

[0206] The compound has general structure (2):



25 wherein:

- Q is a reactive moiety;
- L is a linker that links Z to D;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins,
- 30 KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolbenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;

[0207] The compound of general structure (2) may also be referred to as a "linker-drug construct", for containing linker L and payload D of the final conjugate. Compounds according to general formula (2) can be prepared by the skilled person using standard organic synthesis techniques, and

35 as exemplified in the examples. Linker L and payload D are defined above in the context of the conjugate according to structure (1).

Reactive moiety Q

[0208] The compound according to general structure (2) comprises a reactive moiety Q. In the

40 context of the present invention, the term "reactive moiety" may refer to a chemical moiety that

comprises a reactive group, but also to a reactive group itself. For example, a cyclooctynyl group is a reactive group comprising a reactive group, namely a C-C triple bond. Similarly, an *N*-maleimidyl group is a reactive group, comprising a C-C double bond as a reactive group. However, a reactive group, for example an azido reactive group, a thiol reactive group or an alkynyl reactive group, may
5 herein also be referred to as a reactive moiety.

[0209] Q serves as chemical handle for the connection to $S(F)_x$. In other words, Q is reactive towards and complementary to F. Herein, a reactive group is denoted as “complementary” to a reactive group when said reactive group reacts with said reactive group selectively, optionally in the presence of other functional groups. Complementary reactive and functional groups are known to
10 a person skilled in the art, and are described in more detail below. As such, the compound according to general structure (2) is conveniently used in a conjugation reaction, wherein a chemical reaction between Q and F takes place, thereby forming an antibody-conjugate comprising a covalent connection between payload D and the antibody.


[0210] The exact nature of Q, and F, depends on the type of conjugation reaction that is employed.
15 The skilled person will be able to select the appropriate combination of Q and F. Preferably, Q, and thus also F, is reactive in a cycloaddition or a nucleophilic reaction. Thus, Q preferably comprises a click probe, a thiol, a thiol-reactive moiety, an amine or an amine-reactive moiety, more preferably Q is a click probe, a thiol-reactive moiety or an amine-reactive moiety, most preferably Q is a click probe. The click probe is reactive in a cycloaddition (click reaction) and is preferably selected from
20 an azide, a tetrazine, a triazine, a nitron, a nitrile oxide, a nitrile imine, a diazo compound, an *ortho*-quinone, a dioxothiophene, a sydnone, an alkene moiety and an alkyne moiety. Preferably, the click probe comprises or is an alkene moiety or an alkyne moiety, more preferably wherein the alkene is a (hetero)cycloalkene and/or the alkyne is a terminal alkyne or a (hetero)cycloalkyne. Typical thiol-reactive moieties are selected from maleimide moiety, a haloacetamide moiety, an allenamide
25 moiety, a phosphoramidite moiety, a cyanoethynyl moiety, a vinylsulfone, a vinylpyridine moiety or a methylsulfonylphenyloxadiazole moiety. Most preferably, the thiol-reactive moiety comprises or is a maleimide moiety. Typical amine-reactive moieties are selected from N-hydroxysuccinimidyl esters and other activated esters, *p*-nitrophenyl carbonates and other activated carbonates, isocyanates, isothiocyanates, haloacetamides and benzyl halides. In a preferred embodiment, Q is
30 selected from an alkene moiety, an alkyne moiety, a thiol-reactive moiety or an amine-reactive moiety, more preferably an alkene moiety or an alkyne moiety, even more preferably an alkyne moiety. Herein, the alkene is preferably a (hetero)cycloalkene and the alkyne is preferably a terminal alkyne or a (hetero)cycloalkyne. Most preferably, Q is a cyclic (hetero)alkyne moiety. Each of these moieties are further defined here below.

35
[0211] Thus, in an especially preferred embodiment, Q comprises a cyclic (hetero)alkyne moiety. The alkynyl group may also be referred to as a (hetero)cycloalkynyl group, i.e. a heterocycloalkynyl group or a cycloalkynyl group, wherein the (hetero)cycloalkynyl group is optionally substituted. Preferably, the (hetero)cycloalkynyl group is a (hetero)cycloheptynyl group, a (hetero)cyclooctynyl
40 group, a (hetero)cyclononyl group or a (hetero)cyclodecynyl group. Herein, the

-48-

(hetero)cycloalkynes may optionally be substituted. Preferably, the (hetero)cycloalkynyl group is an optionally substituted (hetero)cycloheptynyl group or an optionally substituted (hetero)cyclooctynyl group. Most preferably, the (hetero)cycloalkynyl group is a (hetero)cyclooctynyl group, wherein the (hetero)cyclooctynyl group is optionally substituted.

- 5 **[0212]** In an especially preferred embodiment, Q comprises a (hetero)cycloalkynyl or (hetero)cycloalkenyl group and is according to structure (Q1):




Herein:

- 10 - the bond depicted as $\equiv$ is a double bond or a triple bond;
- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- 15
- 20 - Y^2 is $C(R^{31})_2$, O, S, $S^{(+)}R^{31}$, $S(O)R^{31}$, $S(O)=NR^{31}$ or NR^{31} , wherein $S^{(+)}$ is a cationic sulphur atom counterbalanced by $B^{(-)}$, wherein $B^{(-)}$ is an anion, and wherein each R^{31} individually is R^{15} or a connection with D, connected via L;
- u is 0, 1, 2, 3, 4 or 5;
- u' is 0, 1, 2, 3, 4 or 5, wherein $u + u' = 0, 1, 2, 3, 4, 5, 6, 7$ or 8;
- 25 - v = an integer in the range 0 – 16;

[0213] Typically, $v = (u + u') \times 2$ (when the connection to L, depicted by the wavy bond, is via Y^2) or $[(u + u') \times 2] - 1$ (when the connection to L, depicted by the wavy bond, is via one of the carbon atoms).

- [0214]** In a preferred embodiment of structure (Q1), reactive group Q comprises a (hetero)cycloalkynyl group and is according to structure (Q1a):
- 30




Herein,

-49-

- R^{15} and Y^2 are as defined above
- u is 0, 1, 2, 3, 4 or 5;
- u' is 0, 1, 2, 3, 4 or 5, wherein $u + u' = 4, 5, 6, 7$ or 8 ;
- v = an integer in the range $8 - 16$.

5 **[0215]** In a preferred embodiment, $u + u' = 4, 5$ or 6 , more preferably $u + u' = 5$. In a preferred embodiment, $v = 8, 9$ or 10 , more preferably $v = 9$ or 10 , most preferably $v = 10$.

[0216] In a preferred embodiment, Q is a (hetero)cycloalkynyl group selected from the group consisting of (Q2) – (Q20) and (Q20a) depicted here below.




10 **[0217]** Herein, the connection to L , depicted with the wavy bond, may be to any available carbon or nitrogen atom of Q . The nitrogen atom of (Q10), (Q13), (Q14) and (Q15) may bear the connection to L , or may contain a hydrogen atom or be optionally functionalized. $B^{(-)}$ is an anion, which is preferably selected from $^{(-)}OTf$, $Cl^{(-)}$, $Br^{(-)}$ or $I^{(-)}$, most preferably $B^{(-)}$ is $^{(-)}OTf$. In the conjugation reaction, $B^{(-)}$ does not need to be a pharmaceutically acceptable anion, since $B^{(-)}$ will exchange

15 with the anions present in the reaction mixture anyway. In case (Q19) is used for Q , the negatively

charged counter-ion is preferably pharmaceutically acceptable upon isolation of the conjugate according to the invention, such that the conjugate is readily useable as medicament.

[0218] In a further preferred embodiment, Q is a (hetero)cycloalkynyl group selected from the group consisting of (Q21) – (Q38) and (Q38a) depicted here below.



- 5 **[0219]** In structure (Q38), $B^{(-)}$ is an anion, which is preferably selected from $(-)\text{OTf}$, $\text{Cl}^{(-)}$, $\text{Br}^{(-)}$ or $\text{I}^{(-)}$, most preferably $B^{(-)}$ is $(-)\text{OTf}$.

- [0220]** In a preferred embodiment, Q comprises a (hetero) cyclooctyne moiety or a (hetero)cycloheptyne moiety, preferably according to structure (Q8), (Q26), (Q27), (Q28), (Q37) or (Q38a), more preferably according to structure (Q8), (Q26), (Q27), (Q28) or (Q37), which are
10 optionally substituted. Each of these preferred options for Q are further defined here below.

[0221] Thus, in a preferred embodiment, Q comprises a heterocycloheptyne moiety according to structure (Q37), also referred to as a TMTHSI, which is optionally substituted. Preferably, the heterocycloheptyne moiety according to structure (Q37) is not substituted.


- [0222]** In an alternative preferred embodiment, Q comprises a cyclooctyne moiety according to
15 structure (Q8), more preferably according to (Q29), also referred to as a bicyclo[6.1.0]non-4-yn-9-yl] group (BCN group), which is optionally substituted. Preferably, the cyclooctyne moiety according to structure (Q8) or (Q29) is not substituted. In the context of the present embodiment, Q preferably is a (hetero)cyclooctyne moiety according to structure (Q39) as shown below, wherein V is $(\text{CH}_2)_l$ and l is an integer in the range of 0 to 10, preferably in the range of 0 to 6. More preferably, l is 0,
20 1, 2, 3 or 4, more preferably l is 0, 1 or 2 and most preferably l is 0 or 1. In the context of group

-51-


(Q39), I is most preferably 1. Most preferably, Q is according to structure (Q42), defined further below.

[0223] In an alternative preferred embodiment, Q comprises a (hetero)cyclooctyne moiety according to structure (Q26), (Q27) or (Q28), also referred to as a DIBO, DIBAC, DBCO or ADIBO group, which are optionally substituted. In the context of the present embodiment, Q preferably is a (hetero)cyclooctyne moiety according to structure (Q40) or (Q41) as shown below, wherein Y¹ is O or NR¹¹, wherein R¹¹ is independently selected from the group consisting of hydrogen, a linear or branched C₁ - C₁₂ alkyl group or a C₄ - C₁₂ (hetero)aryl group. The aromatic rings in (Q40) are optionally O-sulfonylated at one or more positions, whereas the rings of (Q41) may be halogenated at one or more positions. Preferably, the (hetero)cyclooctyne moiety according to structure (Q40) or (Q41) is not further substituted. Most preferably, Q is according to structure (Q43), defined further below.

[0224] In an alternative preferred embodiment, Q comprises a heterocycloheptynyl group and is according to structure (Q37).



[0225] In an especially preferred embodiment, Q comprises a cyclooctynyl group and is according to structure (Q42):



Herein:

- R¹⁵ is independently selected from the group consisting of hydrogen, halogen, -OR¹⁶, -NO₂, -CN, -S(O)₂R¹⁶, -S(O)₃⁽⁻⁾, C₁ - C₂₄ alkyl groups, C₅ - C₂₄ (hetero)aryl groups, C₇ - C₂₄ alkyl(hetero)aryl groups and C₇ - C₂₄ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R¹⁵ may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R¹⁶ is independently selected from the group consisting of hydrogen, halogen, C₁ - C₂₄ alkyl groups, C₆ - C₂₄ (hetero)aryl groups, C₇ - C₂₄ alkyl(hetero)aryl groups and C₇ - C₂₄ (hetero)arylalkyl groups;
- R¹⁸ is independently selected from the group consisting of hydrogen, halogen, C₁ - C₂₄ alkyl groups, C₆ - C₂₄ (hetero)aryl groups, C₇ - C₂₄ alkyl(hetero)aryl groups and C₇ - C₂₄ (hetero)arylalkyl groups;


-52-

(hetero)arylalkyl groups;

- R^{19} is selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups, the alkyl groups optionally being interrupted by one or more hetero-atoms selected from the group consisting of O, N and S, wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are independently optionally substituted, or R^{19} is a second occurrence of Q or D connected via a spacer moiety; and
- I is an integer in the range 0 to 10.

[0226] In a preferred embodiment of the reactive group according to structure (Q42), R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $C_1 - C_6$ alkyl groups, $C_5 - C_6$ (hetero)aryl groups, wherein R^{16} is hydrogen or $C_1 - C_6$ alkyl, more preferably R^{15} is independently selected from the group consisting of hydrogen and $C_1 - C_6$ alkyl, most preferably all R^{15} are H. In a preferred embodiment of the reactive group according to structure (Q42), R^{18} is independently selected from the group consisting of hydrogen, $C_1 - C_6$ alkyl groups, most preferably both R^{18} are H. In a preferred embodiment of the reactive group according to structure (Q42), R^{19} is H. In a preferred embodiment of the reactive group according to structure (Q42), I is 0 or 1, more preferably I is 1.

[0227] In an especially preferred embodiment, Q comprises a (hetero)cyclooctynyl group and is according to structure (Q43):





Herein:

- R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-NO_2$, $-CN$, $-S(O)_2R^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_{24}$ alkyl groups, $C_5 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups and wherein the alkyl groups, (hetero)aryl groups, alkyl(hetero)aryl groups and (hetero)arylalkyl groups are optionally substituted, wherein two substituents R^{15} may be linked together to form an optionally substituted annulated cycloalkyl or an optionally substituted annulated (hetero)arene substituent, and wherein R^{16} is independently selected from the group consisting of hydrogen, halogen, $C_1 - C_{24}$ alkyl groups, $C_6 - C_{24}$ (hetero)aryl groups, $C_7 - C_{24}$ alkyl(hetero)aryl groups and $C_7 - C_{24}$ (hetero)arylalkyl groups;
- Y is N or CR^{15} .

[0228] In a preferred embodiment of the reactive group according to structure (Q43), R^{15} is independently selected from the group consisting of hydrogen, halogen, $-OR^{16}$, $-S(O)_3^{(-)}$, $C_1 - C_6$ alkyl groups, $C_5 - C_6$ (hetero)aryl groups, wherein R^{16} is hydrogen or $C_1 - C_6$ alkyl, more preferably R^{15} is independently selected from the group consisting of hydrogen and $-S(O)_3^{(-)}$. In a preferred

embodiment of the reactive group according to structure (Q43), Y is N or CH, more preferably Y = N.







[0229] In an especially preferred embodiment, Q comprises a heterocycloheptynyl group and is according to structure (Q37) or (Q38a), preferably according to structure (Q37)




5

[0230] In an alternative preferred embodiment, Q comprises a cyclic alkene moiety. The alkenyl group Q may also be referred to as a (hetero)cycloalkenyl group, i.e. a heterocycloalkenyl group or a cycloalkenyl group, preferably a cycloalkenyl group, wherein the (hetero)cycloalkenyl group is optionally substituted. Preferably, the (hetero)cycloalkenyl group is a (hetero)cyclopropenyl group, a (hetero)cyclobutenyl group, a norbornene group, a norbornadiene group, a *trans*-(hetero)cycloheptenyl group, a *trans*-(hetero)cyclooctenyl group, a *trans*-(hetero)cyclononenyl group or a *trans*-(hetero)cyclodecenyl group, which may all optionally be substituted. Especially preferred are (hetero)cyclopropenyl groups, *trans*-(hetero)cycloheptenyl group or *trans*-(hetero)cyclooctenyl groups, wherein the (hetero)cyclopropenyl group, the *trans*-(hetero)cycloheptenyl group or the *trans*-(hetero)cyclooctenyl group is optionally substituted. Preferably, Q comprises a cyclopropenyl moiety according to structure (Q44), a heterocyclobutene moiety according to structure (Q45), a norbornene or norbornadiene group according to structure (Q46), a *trans*-(hetero)cycloheptenyl moiety according to structure (Q47) or a *trans*-(hetero)cyclooctenyl moiety according to structure (Q48). Herein, Y³ is selected from C(R²³)₂, NR²³ or O, wherein each R²³ is individually hydrogen, C₁ – C₆ alkyl or is connected to L, optionally via a spacer, and the bond labelled --- is a single or double bond. In a further preferred embodiment, the cyclopropenyl group is according to structure (Q49). In another preferred embodiment, the *trans*-(hetero)cycloheptene group is according to structure (Q50) or (Q51). In another preferred embodiment, the *trans*-(hetero)cyclooctene group is according to structure (Q52), (Q53), (Q54), (Q55) or (Q56).


25




-54-



(Q52)




(Q53)



(Q54)



(Q55)




(Q56)


[0231] Herein, the R group(s) on Si in (Q50) and (Q51) are typically alkyl or aryl, preferably C₁-C₆ alkyl.

[0232] In an alternative preferred embodiment, Q is a thiol-reactive probe. In this embodiment, Q is a reactive group compatible with cysteine conjugation. Such probes are known in the art and may be selected from the group consisting of a maleimide moiety, a haloacetamide moiety, an allenamide moiety, a phosphonamidite moiety, a cyanoethynyl moiety, a vinylsulfone, a vinylpyridine moiety or a methylsulfonylphenyloxadiazole moiety. Most preferably, Q comprises or is a maleimide moiety. Reagents may be monoalkylation type or may be a cross-linker for reaction with two cysteine side-chains.


[0233] In a further preferred embodiment, probe Q is selected from the group consisting of (Q57) – (Q71) depicted here below.




(Q57)




(Q58)




(Q59)




(Q60)




(Q61)




(Q62)



(Q63)




(Q64)




(Q65)




(Q66)




(Q67)




(Q68)



(Q69)



(Q70)



(Q71)


wherein:

- X⁶ is H, halogen, PhS, MeS, preferably a halogen, such as Cl, Br, I;
- X⁷ is halogen, PhS, MeS, preferably a halogen, such as Cl, Br, I;
- R²⁴ is H or C₁₋₁₂ alkyl, preferably H or C₁₋₆ alkyl;

-55-

- R^{25} is H, C_{1-12} alkyl, C_{1-12} aryl, C_{1-12} alkaryl or C_{1-12} aralkyl, preferably H or *para*-methylphenyl;
- wherein the aromatic ring of (Q61) and (Q63) may optionally be a heteroaromatic ring, such as a phenyl or pyridine ring.

- 5 **[0234]** In a preferred embodiment of thiol-reactive probe (Q57), the probe Q is selected from the group consisting of (Q72) – (Q74) depicted here below.



wherein:


- R^{27} is C_{1-12} alkyl, C_{1-12} aryl, C_{1-12} alkaryl or C_{1-12} aralkyl;
- t is an integer in the range of 0 – 15, preferably 1 – 10.

10

[0235] In an alternative preferred embodiment, Q is an amine-reactive probe. In this embodiment, Q is a reactive group compatible with lysine conjugation. Such probes are known in the art and may be selected from the group consisting of *N*-hydroxysuccinimidyl groups, isocyanate groups, isothiocyanate groups and benzoyl halide groups. Most preferably, Q comprises or is an *N*-hydroxysuccinimidyl esters or a *p*-nitrophenyl carbonate moiety.

15

[0236] In a further preferred embodiment, probe Q is selected from the group consisting of (Q75) – (Q79) depicted here below.




Herein, X^2 is halogen, preferably F.

[0237] In a preferred embodiment, Q is selected from the group consisting of (Q1) – (Q79).

20

Antibody according to general structure (3)

[0238] The antibody has general structure (3):



25 wherein:

- AB is an antibody capable of targeting CEA-expressing tumours;
- b is 0 or 1;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;

30

- F is a reactive moiety;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

[0239] The antibody of general structure (3) may also be referred to as a “(modified) antibody”, for being an antibody containing reactive groups F, wherein the reactive groups F are naturally present or the antibody is modified to incorporate the reactive groups F. The (modified) antibody according to general formula (2) can be prepared by the skilled person using standard organic and/or enzymatic synthesis techniques, and as exemplified in the examples. Antibody AB, linker L⁶, b, x and y are defined above in the context of the conjugate according to structure (1).

Reactive moiety F


[0240] F is reactive towards Q in the conjugation reaction defined below, preferably wherein the conjugation reaction is a cycloaddition or a nucleophilic reaction. As the skilled person will understand, the options for F are the same as those for Q, provided that F and Q are reactive towards each other. Thus, F preferably comprises a click probe, a thiol, a thiol-reactive moiety, an amine or an amine-reactive moiety, more preferably F is a click probe, a thiol or an amine, most preferably F is a click probe. The click probe is reactive in a cycloaddition (click reaction) and is preferably selected from an azide, a tetrazine, a triazine, a nitron, a nitrile oxide, a nitrile imine, a diazo compound, an *ortho*-quinone, a dioxothiophene, a sydnone, an alkene moiety and an alkyne moiety. Preferably, the click probe comprises or is an azide, a tetrazine, a triazine, a nitron, a nitrile oxide, a nitrile imine, a diazo compound, an *ortho*-quinone, a dioxothiophene or a sydnone, most preferably an azide. Typical thiol-reactive moieties are selected from maleimide moiety, a haloacetamide moiety, an allenamide moiety, a phosphonamidite moiety, a cyanoethynyl moiety, an *ortho*-quinone moiety, a vinylsulfone, a vinylpyridine moiety or a methylsulfonylphenyloxadiazole moiety. Most preferably, the thiol-reactive moiety comprises or is a maleimide moiety. Typical amine-reactive moieties are selected from N-hydroxysuccinimidyl esters, isocyanates, isothiocyanates and benzyl halides. In a preferred embodiment, F is a click probe or a thiol, more preferably F is an azide or a thiol, most preferably F is an azide.

[0241] More than one reactive group F may be present in the antibody. The reactive group F in the antibody may be naturally present or may be placed in the antibody by a specific technique, for example a (bio)chemical or a genetic technique. The reactive group that is placed in the antibody is prepared by chemical synthesis, for example an azide or a terminal alkyne. Methods of preparing modified antibodies are known in the art, e.g. from WO 2014/065661, WO 2016/170186 and WO 2016/053107, which are incorporated herein by reference. From the same documents, the conjugation reaction between the modified antibody and a linker-drug construct is known to the skilled person.

[0242] Preferably, F is a click probe reactive towards a (hetero)cycloalkene and/or a (hetero)cycloalkyne, and is typically selected from the group consisting of azide, tetrazine, triazine, nitron, nitrile oxide, nitrile imine, diazo compound, *ortho*-quinone, dioxothiophene and sydnone.

Preferred structures for the reactive group are structures (F1) – (F10) depicted here below.

-57-



[0243] Herein, the wavy bond represents the connection to the payload. For (F3), (F4), (F8) and (F9), the payload can be connected to any one of the wavy bonds. The other wavy bond may then be connected to an R group selected from hydrogen, $\text{C}_1 - \text{C}_{24}$ alkyl groups, $\text{C}_2 - \text{C}_{24}$ acyl groups, $\text{C}_3 - \text{C}_{24}$ cycloalkyl groups, $\text{C}_2 - \text{C}_{24}$ (hetero)aryl groups, $\text{C}_3 - \text{C}_{24}$ alkyl(hetero)aryl groups, $\text{C}_3 - \text{C}_{24}$ (hetero)arylalkyl groups and $\text{C}_1 - \text{C}_{24}$ sulfonyl groups, each of which (except hydrogen) may optionally be substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR³² wherein R³² is independently selected from the group consisting of hydrogen and $\text{C}_1 - \text{C}_4$ alkyl groups. The skilled person understands which R groups may be applied for each of the groups F. For example, the R group connected to the nitrogen atom of (F3) may be selected from alkyl and aryl, and the R group connected to the carbon atom of (F3) may be selected from hydrogen, alkyl, aryl, acyl and sulfonyl. Preferably, the reactive moiety F is selected from azides or tetrazines. Most preferably, the reactive moiety F is an azide.

[0244] In a second preferred embodiment, F is a thiol or precursor thereof. Thiol or precursor thereof F is used in the conjugation reaction to connect the linker-drug construct to the (modified) antibody. F is reactive towards thiol-reactive probe Q in a thiol ligation. The thiol preferably the thiol of the side chain of a cysteine amino acid, which are naturally present within the antibody AB, in which case linker L⁶ is not present ($b = 0$), although it may also be synthetically introduced, optionally via a linker L⁶. Thiol precursors in the context of bioconjugation are known in the art, and include disulfides, which may be naturally occurring disulfide bridges present in the antibody or synthetically introduced disulfides, which are reduced as known in the art. Preferably, F is a thiol group of a cysteine side chain.

[0245] In a third preferred embodiment, F is an amine or precursor thereof, preferably an amine. Amine or precursor thereof F is used in the conjugation reaction to connect the linker-drug construct to the (modified) antibody. F is reactive towards amine-reactive probe Q in nucleophilic substitution. The amine is typically a primary amine, preferably the amine of the side chain of a lysine amino acid, which are naturally present within the antibody AB, in which case linker L⁶ is not present ($b = 0$), although it may also be synthetically introduced, optionally via a linker L⁶. Preferably, F is a primary amine group of a lysine side chain.

30

Process for synthesising the antibody-conjugate according to general structure (1)

[0246] In a further aspect, the present invention relates to a process for the preparation of the antibody-conjugate according to the invention, the process comprising the step of reacting Q of the

compound according to the invention with a reactive group F of an antibody. The compound according to general structure (2), and preferred embodiments thereof, are described in more detail above. The present process occurs under conditions such that Q is reacted with F to covalently link the antibody AB (3) to the payload D. In the process according to the invention, Q reacts with F, forming a covalent connection between the antibody and the compound according to the invention. Complementary reactive groups Q and reactive groups F are known to the skilled person and are described in more detail below.

[0247] Any conjugation technique known in the art can be employed to prepare the multifunctional antibody constructs according to the invention. Suitable conjugation techniques include thiol ligation, lysine ligation, cycloadditions (e.g. copper-catalysed click reaction, strain-promoted azide-alkyne cycloaddition, strain-promoted quinone-alkyne cycloaddition). Preferred conjugation techniques used in the context of the present invention include nucleophilic reactions and cycloadditions, preferably wherein the cycloaddition is a [4+2] cycloaddition or a [3+2] cycloaddition and the nucleophilic reaction is a Michael addition or a nucleophilic substitution. Suitable conjugation techniques are for example disclosed in G.T. Hermanson, "Bioconjugate Techniques", Elsevier, 3rd Ed. 2013 (ISBN:978-0-12-382239-0), WO 2014/065661, van Geel et al., *Bioconj. Chem.* **2015**, 26, 2233–2242, PCT/EP2021/050594, PCT/EP2021/050598 and NL 2026947.

[0248] Thus, in a preferred embodiment of the conjugation process according to the invention, conjugation is accomplished via a nucleophilic reaction, such as a nucleophilic substitution or a Michael reaction. A preferred nucleophilic reaction is the acylation of a primary amino group with an activated ester. A preferred Michael reaction is the maleimide-thiol reaction, which is widely employed in bioconjugation.

[0249] Thus, in a preferred embodiment of the conjugation process according to the invention, conjugation is accomplished via a cycloaddition. Preferred cycloadditions are a (4+2)-cycloaddition (e.g. a Diels-Alder reaction) or a (3+2)-cycloaddition (e.g. a 1,3-dipolar cycloaddition). Preferably, the conjugation reaction is the Diels-Alder reaction or the 1,3-dipolar cycloaddition. The preferred Diels-Alder reaction is the inverse-electron demand Diels-Alder cycloaddition. In another preferred embodiment, the 1,3-dipolar cycloaddition is used, more preferably the alkyne-azide cycloaddition, and most preferably wherein Q is or comprises an alkyne group and F is an azido group. Cycloadditions, such as Diels-Alder reactions and 1,3-dipolar cycloadditions are known in the art, and the skilled person knows how to perform them.


[0250] The process according to the present aspect preferably concerns a click reaction, more preferably a 1,3-dipolar cycloaddition, most preferably an alkyne/azide cycloaddition. Most preferably, Q is or comprises an alkyne group and F is an azido group. Click reactions, such as 1,3-dipolar cycloadditions, are known in the art, and the skilled person knows how to perform them.

[0251] Thus, the process for preparing the antibody-conjugate according to the invention according to this aspect comprises reacting the modified antibody of structure (3) with a compound according to structure (2), to obtain the antibody-conjugate of structure (1).

[0252] In a preferred embodiment, the process for preparing the antibody-conjugate according to

the invention comprises:

- (i) contacting an antibody comprising y core *N*-acetylglucosamine (GlcNAc) moieties, wherein $y = 1, 2, 3$ or 4 , with a compound of the formula $S(F)_xP$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety, to obtain a modified antibody according to Formula (26):



wherein

- AB is an antibody capable of targeting CEA-expressing tumours;
- Fuc is fucose;
- w is 0 or 1 ; and

- (ii) reacting the modified antibody with a compound according to structure (2):



wherein:

- Q is a reactive moiety;
- L is a linker that links Z to D ;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolbenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;

to obtain the antibody-conjugate according to structure (1).

Step (i)

[0253] In step (i), an antibody comprising $1, 2, 3$ or 4 core *N*-acetylglucosamine moieties is contacted with a compound of the formula $S(F)_xP$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety. Herein, the antibody is typically an antibody that has been trimmed to a core-GlcNAc residue as described further below. Step (i) affords a modified antibody according to Formula (26).

[0254] The starting material, i.e. the antibody comprising a core-GlcNAc substituent, is known in the art and can be prepared by methods known by the skilled person. In one embodiment, the process according to the invention further comprises the deglycosylation of an antibody glycan having a core *N*-acetylglucosamine, in the presence of an endoglycosidase, in order to obtain an antibody comprising a core *N*-acetylglucosamine substituent, wherein said core *N*-acetylglucosamine and said core *N*-acetylglucosamine substituent are optionally fucosylated. Depending on the nature of the glycan, a suitable endoglycosidase may be selected. The

endoglycosidase is preferably selected from the group consisting of EndoS, EndoA, EndoE, EfEndo18A, EndoF, EndoM, EndoD, EndoH, EndoT and EndoSH and/or a combination thereof, the selection of which depends on the nature of the glycan. EndoSH is described in PCT/EP2017/052792, see Examples 1 – 3, and SEQ. ID No: 1, which is incorporated by reference
5 herein.

[0255] Structural features S and x are defined above for the antibody-conjugate according to the invention, which equally applies to the present aspect. Compounds of the formula $S(F)_xP$, wherein a nucleoside monophosphate or a nucleoside diphosphate P is linked to a sugar derivative $S(F)_x$, are known in the art. For example Wang *et al.*, *Chem. Eur. J.* **2010**, *16*, 13343–13345, Piller *et al.*,
10 *ACS Chem. Biol.* **2012**, *7*, 753, Piller *et al.*, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5459–5462 and WO 2009/102820, all incorporated by reference herein, disclose a number of compounds $S(F)_xP$ and their syntheses. In a preferred embodiment nucleoside mono- or diphosphate P in $S(F)_xP$ is selected from the group consisting of uridine diphosphate (UDP), guanosine diphosphate (GDP), thymidine diphosphate (TDP), cytidine diphosphate (CDP) and cytidine monophosphate (CMP),
15 more preferably P is selected from the group consisting of uridine diphosphate (UDP), guanosine diphosphate (GDP) and cytidine diphosphate (CDP), most preferably P = UDP. Preferably, $S(F)_xP$ is selected from the group consisting of GalNAz-UDP, F₂-GalNAz-UDP (*N*-(azidodifluoro)acetyl-galactosamine), 6-AzGal-UDP, 6-AzGalNAc-UDP (6-azido-6-deoxy-N-acetylgalactosamine-UDP), 4-AzGalNAz-UDP, 6-AzGalNAz-UDP, GlcNAz-UDP, 6-AzGlc-UDP, 6-AzGlcNAz-UDP and 2-(but-
20 3-ynic acid amido)-2-deoxy-galactose-UDP. Most preferably, $S(F)_xP$ is GalNAz-UDP or 6-AzGalNAc-UDP.

[0256] Suitable catalyst that are capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety are known in the art. A suitable catalyst is a catalyst wherefore the specific sugar derivative nucleotide $S(F)_xP$ in that specific process is a substrate. More specifically, the catalyst catalyses
25 the formation of a $\beta(1,4)$ -glycosidic bond. Preferably, the catalyst is selected from the group of galactosyltransferases and N-acetylgalactosaminyltransferases, more preferably from the group of $\beta(1,4)$ -N-acetylgalactosaminyltransferases (GalNAcT) and $\beta(1,4)$ -galactosyltransferases (GalT), most preferably from the group of $\beta(1,4)$ -N-acetylgalactosaminyltransferases having a mutant catalytic domain. Suitable catalysts and mutants thereof are disclosed in WO 2014/065661, WO
30 2016/022027 and WO 2016/170186, all incorporated herein by reference. In one embodiment, the catalyst is a wild-type galactosyltransferase or N-acetylgalactosaminyltransferase, preferably an N-acetylgalactosaminyltransferase. In an alternative embodiment, the catalyst is a mutant galactosyltransferase or N-acetylgalactosaminyltransferases, preferably a mutant N-acetylgalactosaminyltransferase. Mutant enzymes described in WO 2016/022027 and WO
35 2016/170186 are especially preferred. These galactosyltransferase (mutant) enzyme catalysts are able to recognize internal sugars and sugar derivatives as an acceptor. Thus, sugar derivative $S(F)_x$ is linked to the core-GlcNAc substituent in step (i), irrespective of whether said GlcNAc is fucosylated or not.

[0257] Step (i) is preferably performed in a suitable buffer solution, such as for example phosphate,
40 buffered saline (e.g. phosphate-buffered saline, tris-buffered saline), citrate, HEPES, tris and

glycine. Suitable buffers are known in the art. Preferably, the buffer solution is phosphate-buffered saline (PBS) or tris buffer. Step (i) is preferably performed at a temperature in the range of about 4 to about 50 °C, more preferably in the range of about 10 to about 45 °C, even more preferably in the range of about 20 to about 40 °C, and most preferably in the range of about 30 to about 37 °C.

- 5 Step (i) is preferably performed a pH in the range of about 5 to about 9, preferably in the range of about 5.5 to about 8.5, more preferably in the range of about 6 to about 8. Most preferably, step (i) is performed at a pH in the range of about 7 to about 8.

Step (ii)

- 10 **[0258]** In step (ii), the modified antibody is reacted with a compound according to general structure (2), comprising a reactive group Q capable of reacting with reactive group F and a payload D, to obtain the antibody-conjugate according to structure (1), containing connecting group Z resulting from the reaction between Q and F. Such reaction occurs under condition such that reactive group Q is reacted with the reactive group F of the biomolecule to covalently link the antibody to the
15 compound according to general structure (2). Step (ii) may also be referred to as the conjugation reaction.

- [0259]** In a preferred embodiment, in step (ii) an azide on an azide-modified antibody reacts with an alkynyl group, preferably a terminal alkynyl group, or a (hetero)cycloalkynyl group of the compound according to general structure (2), via a cycloaddition reaction. This cycloaddition
20 reaction of a molecule comprising an azide with a molecule comprising a terminal alkynyl group or a (hetero)cycloalkynyl group is one of the reactions that is known in the art as “click chemistry”. In the case of a linker-conjugate comprising a terminal alkynyl group, said cycloaddition reaction needs to be performed in the presence of a suitable catalyst, preferably a Cu(I) catalyst. However, in a preferred embodiment, the linker-conjugate comprises a (hetero)cycloalkynyl group, more
25 preferably a strained (hetero)cycloalkynyl group. When the (hetero)cycloalkynyl is a strained (hetero)cycloalkynyl group, the presence of a catalyst is not required, and said reaction may even occur spontaneously by a reaction called strain-promoted azide-alkyne cycloaddition (SPAAC). This is one of the reactions known in the art as “metal-free click chemistry”.

30 Application

- [0260]** The invention further concerns a method for the treatment of a subject in need thereof, comprising the administration of the antibody-conjugate according to the invention as defined above. The subject in need thereof is typically a cancer patient. The use of antibody-conjugates, such as antibody-drug conjugates, is well-known in the field of cancer treatment, and the antibody-
35 conjugates according to the invention are especially suited in this respect. The method as described is typically suited for the treatment of cancer. In the method according to this aspect, the antibody-conjugate is typically administered in a therapeutically effective dose. The present aspect of the invention can also be worded as an antibody-conjugate according to the invention for use in the treatment of a subject in need thereof, preferably for the treatment of cancer. In other words, this
40 aspect concerns the use of an antibody-conjugate according to the invention for the preparation of

a medicament or pharmaceutical composition for use in the treatment of a subject in need thereof, preferably for use in the treatment of cancer. In the present context, treatment of cancer is envisioned to encompass treating, imaging, diagnosing, preventing the proliferation of, containing and reducing tumours.

5 **[0261]** This aspect of the present invention may also be worded as a method for targeting CEA-expressing cells, in particular CEA-expressing tumour cells, comprising contacting the antibody-conjugate according to the invention with cells that may possibly be CEA-expressing. The method according to this aspect is thus suitable to determine whether the cells are CEA-expressing. These CEA-expressing cells may be present in a subject, in which case the method comprises
10 administering to a subject in need thereof the antibody-conjugate according to the invention. In a preferred embodiment, the cells that may possibly be CEA-expressing are CEA-expressing cells. The targeting of CEA-expressing cells preferably includes one or more of treating, imaging, diagnosing, preventing the proliferation of, containing and reducing CEA-expressing cells, in particular CEA-expressing tumour cells. The method according to this embodiment may be medical
15 or non-medical. Non-medical methods according to the present aspect may be directed to *in vitro* or *ex vivo* targeting CEA-expressing cells, wherein the cells that may possibly be CEA-expressing are present in a sample, e.g. taken from a patient. Such a non-medical method is typically used for the diagnosis of cancer, in particular CEA-positive cancer.

[0262] The CEA-expressing cells preferably express CEACAM1, CEACAM3, CEACAM4,
20 CEACAM5, CEACAM6, CEACAM7, CEACAM8, CEACAM16, CEACAM18, CEACAM19, CEACAM20, CEACAM21, most preferably they express CEACAM5. In the context of the present invention, the subject may suffer from a disorder selected from colorectal cancer, gastric cancer, lung cancer, uterine cancer or pancreatic cancer. Thus, the treatment of a subject in need thereof preferably refers to the treatment of colorectal cancer, gastric cancer, lung cancer or uterine cancer.

25 **[0263]** The inventors have surprisingly found that the antibody-conjugates according to the invention are superior to conventional CEA-targeting antibody-conjugates in terms of safety and/or efficacy, such that the therapeutic index of the antibody-conjugate according to the invention is increased with respect to conventional CEA-targeting antibody-conjugates.

30 Mode of conjugation


[0264] In the context of the present invention, the “mode of conjugation” refers to the process that is used to conjugate a payload D to an antibody AB, as well as to the structural features of the resulting antibody-conjugate, in particular of the linker that connects the payload to the antibody, that are a direct consequence of the process of conjugation. Thus, in one embodiment, the mode
35 of conjugation refers to a process for conjugation a payload to an antibody. In an alternative embodiment, the mode of conjugation refers to structural features of the linker and/or to the attachment point of the linker to the antibody that are a direct consequence of the process for conjugation a payload to an antibody.

[0265] In a further aspect, the invention concerns the use of a mode of conjugation for increasing
40 the therapeutic index of an antibody-conjugate in the treatment of CEA-expressing tumours,

-63-

wherein the mode of conjugation is being used to connect antibody AB with payload D via a linker L. The mode of conjugation mode of conjugation comprises:

- (i) contacting an antibody comprising y core *N*-acetylglucosamine (GlcNAc) moieties, wherein $y = 1, 2, 3$ or 4 , with a compound of the formula $S(F)_xP$ in the presence of a catalyst, wherein $S(F)_x$ is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q , x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the $S(F)_x$ moiety to the core-GlcNAc moiety, to obtain a modified antibody according to Formula (26):



(26)

wherein

- AB is an antibody capable of targeting CEA-expressing tumours;
- Fuc is fucose;
- w is 0 or 1 ; and

- (ii) reacting the modified antibody with a compound according to structure (2):



(2)

wherein:

- Q is a reactive moiety;
- L is a linker that links Z to D ;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemicasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolbenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;

to obtain the antibody-conjugate according to structure (1)

[0266] Preferably, increasing the therapeutic index of an antibody-conjugate is selected from:

- (a) increasing the therapeutic efficacy of the antibody-conjugate; and/or
- (b) increasing the tolerability of the antibody-conjugate.

[0267] Increase in therapeutic efficacy of the antibody-conjugates according to the invention may take the form of a reduction in tumour size and/or a prolonged period of regression, when compared to conventional CEA-targeting ADC. Increase in tolerability of the antibody-conjugates according to the invention may take the form of a reduction in signs of toxicity, compared to administration of a CEA-targeting ADC made with a conventional technology. The reduction in signs may also be referred to as a reduction in symptoms or side-effects of cancer treatment, and may involve one or more clinical signs such as reduced reduction in body weight, reduced reduction in mobility, reduced reduction in food intake and/or one or more toxicity parameters, such as improved blood chemistry, hematology, and/or histopathology.

Examples

[0268] General procedure for transient expression and purification of monoclonal antibodies:

-64-

Various IgGs (tusamitamab, labetuzumab or B12) were transiently expressed in CHO K1 cells by Evitria (Zurich, Switzerland) at 2L, 250 mL and 5 L scale respectively. The supernatant was purified using a HiTrap MabSelect sure column. The supernatant was loaded onto the column followed by washing with at least 10 column volumes of 25 mM Tris pH 7.5, 150 mM NaCl (TBS). Retained protein was eluted with 0.1 M AcOH pH 2.7. The eluted product was immediately neutralized with 2.5 M Tris-HCl pH 8.8 and dialyzed against 20 mM histidine, 150 mM NaCl, pH 7.5. Next the IgG was concentrated (>20 mg/mL) using a Vivaspin Turbo 15 ultrafiltration unit (Sartorius). The sequences of the IgGs is given here below:

[0269] tusamitamab (I) light chain (SEQ ID No. 43):

DIQMTQSPASLSASVDRVTITCRASENIFSYLAWYQQKPGKSPKLLVYNTRTLAEGVPSRFSGSG
SGTDFSLTISSLQPEDFATYYCQHHYGTPFTFGSGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASV
VCLLNNFYPRKAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSTLTLSKADYEKHKVYACEV
THQGLSSPVTKSFNRGEC

[0270] tusamitamab (I) heavy chain (SEQ ID No. 41):

EVQLQESGPGLVKPGGSLSLSCAASGFVSSYDMSWVRQTPERGLEWVAYISSGGGITYAPSTV
KGRFTVSRDNAKNTLYLQMNSLTSEDVAVYYCAAHYFGSSGPFAYWGQGLTVTVSSASTKGPS
VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMIS
RTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK
EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSQSVSMHEALHNHYTQKSLSLSPG

[0271] labetuzumab (II) light chain (SEQ ID No. 38):

DIQLTQSPSSLSASVGDRTITCKASQDVGTSVAWYQQKPGKAPKLLIYWTSTRHTGVPSRFSG
SGSGTDFITFTISLQPEDATYYCQYSLYRSFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTAS
VVCLLNNFYPRKAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSTLTLSKADYEKHKVYACE
VTHQGLSSPVTKSFNRGEC

[0272] labetuzumab (II) heavy chain (SEQ ID No. 36):

EVQLVESGGGVVQPGRSLRLSCSASGFDFTTYWMSWVRQAPGKGLEWIGEIHDPDSSTINYAPSL
KDRFTISRDNANKNTFLQMDSLRLPEDTGVYFCASLYFGFPWFAYWGQGTPTVTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
SSSLGTQTYICNVNHKPSNTKVDKRVKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISR
TPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEEMTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSQSVSMHEALHNHYTQKSLSLSPG
K

[0273] B12 (III) light chain (SEQ ID No. 52):

EIVLTQSPGTLSPGERATFSCRSSHISRRVAWYQHKGQAPRLVIHGVSNNRASGISDRFSG
SGSGTDFTLTITRVEPEDFALYYCQVYGASSYTFGQGTCLERKRTVAAPSVFIFPPSDEQLKSGT
ASVVCLLNNFYPRKAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSTLTLSKADYEKHKVYA
CEVTHQGLSSPVTKSFNRGEC

-65-

[0274] B12 (III) heavy chain (SEQ ID No. 53):

QVQLVQSGAEVKKPGASVKVSCQASGYRFSNFIHWRQAPGQRFQFEWMGWINPYNGNKEFSA
 KFQDRVTFTADTSANTAYMELRSLRSADTAVYYCARVGPYSWDDSPQDNYYMDVWGKGTIVIV
 SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
 5 YSLSSVWTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPPELLGGPSVFLFP
 PKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV
 LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFY
 PSDIAVEWESNGQPENNYKTTTPVLDSGSSFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHY
 TQKSLSLSPGK

10 **[0275]** General procedure for analytical RP-UPLC (DTT treated samples): Prior to RP-UPLC analysis, IgG (10 μ L, 1 mg/mL in PBS pH 7.4) was added to 12.5 mM DTT, 100 mM TrisHCl pH 8.0 (40 μ L) and incubated for 15 minutes at 37 °C. The reaction was quenched by adding 49% acetonitrile, 49% water, 2% formic acid (50 μ L). RP-UPLC analysis was performed on a Waters Acquity UPLC-SQD. The sample (5 μ L) was injected with 0.4 mL/min onto Bioresolve RP
 15 mAb 2.1*150 mm 2.7 μ m (Waters) with a column temperature of 70 °C. A linear gradient was applied in 9 minutes from 30 to 54% acetonitrile in 0.1% TFA and water. Absorbance of eluted peaks was measured at 215 nm followed by automated integration (MassLynx, Waters) to determine reaction conversion.

[0276] General procedure for mass spectral analysis of (modified) monoclonal antibodies: Prior to
 20 mass spectral analysis, IgG was treated with IdeS, which allows analysis of the Fc/2 fragment. For analysis of both light and heavy chain, a solution of 20 μ g (modified) IgG was incubated for 5 minutes at 37 °C with 100 mM DTT in a total volume of 4 μ L. If present, azide-functionalities are reduced to amines under these conditions. For analysis of the Fc/2 fragment, a solution of 20 μ g (modified) IgG was incubated for 1 hour at 37 °C with IdeS/Fabricator™ (1.25 U/ μ L) in phosphate-
 25 buffered saline (PBS) pH 6.6 in a total volume of 10 μ L. Samples were diluted to 80 μ L followed by analysis electrospray ionization time-of-flight (ESI-TOF) on a JEOL AccuTOF. Deconvoluted spectra were obtained using Magtran software.

[0277] General procedure for enzymatic remodeling of IgG to mAb-(6-N₃-GalNAc)₂: IgG (15 mg/mL) was incubated with 1% w/w EndoSH (as described in PCT/EP2017/052792, see Examples
 30 1 – 3, and SEQ. ID No: 1, which is incorporated by reference herein), 3% w/w His-TnGalNAcT (as described in PCT/EP2016/059194, see Examples 3 and 4, and SEQ. ID No: 49, which is incorporated by reference herein), 0.01% AP (Roche) and UDP 6-N₃-GalNAc (compound **2d** in Figure 3, 25 eq compared to IgG) in 6 mM MnCl₂ and TBS for 16 hours at 30 °C. Next, the functionalized IgG was purified using a HiTrap MabSelect sure 5 mL column. After loading of the
 35 reaction mixture the column was washed with TBS+0.2% triton and TBS. The IgG was eluted with 0.1 M AcOH pH 2.7 and neutralized with 2.5 M Tris-HCl pH 8.8. After three times dialysis to 20 mM histidine, 150 mM NaCl pH 7.5, the IgG was concentrated to 15-20 mg/mL using a Vivaspin Turbo 15 ultrafiltration unit (Sartorius).

Preparation of azide-functionalized antibodies: Examples 1 – 3:

Example 1: Preparation of tusamitamab-(6-N₃-GalNAc)₂ (I-N₃)

[0278] According to the general procedure for enzymatic remodeling, tusamitamab was converted to tusamitamab-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24329 Da, approximately 95% of total Fc/2), corresponding to the expected product.

Example 2: Preparation of labetuzumab-(6-N₃-GalNAc)₂ (II-N₃)

[0279] According to the general procedure for enzymatic remodeling, labetuzumab was converted to labetuzumab-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24360 Da, approximately 90% of total Fc/2), corresponding to the expected product.

Example 3: Preparation of B12-(6-N₃-GalNAc)₂ (III-N₃)

[0280] According to the general procedure for enzymatic remodeling, B12 was converted to B12-(6-N₃-GalNAc)₂. Mass spectral analysis of a sample after IdeS treatment showed one major Fc/2 product (observed mass 24330 Da, approximately 90% of total Fc/2), corresponding to the expected product.

Examples 4-8: Synthesis of linker conjugates 3, 4, 5b and 9


Example 4: Preparation of compound 11

[0281] Compound **10** (163 mg, 240 μmol) was added to a mixture of exatecan mesylate (125 mg, 235 μmol) and DIPEA (61 mg, 82 μL, 0.47 mmol) in dry DMF (0.9 mL). After 20 h, the reaction mixture was diluted to 9 mL DCM and purified by gradient column chromatography (0 → 40% MeOH/DCM) to afford **11** (155 mg, 159 μmol, 68%). LCMS (ESI+) calculated for C₅₅H₅₄FN₆O₁₀⁺ (M+H)⁺ 977.39, found 977.72. In addition to **11**, free base of exatecan (82.4 mg, 189 μmol, 20%) was recovered. LCMS (ESI+) calculated for C₂₄H₂₃FN₃O₄⁺ (M+H)⁺ 436.46, found 436.54.

Example 5: Preparation of compound 3

[0282] The synthesis of BCN-HS-(va-PABC-Ex)₂ (**3**) is also described in PCT/EP2021/075401 (example 4), incorporated herein. To a solution of compound **11** (155 mg, 159 μmol) in DMF (1.6 mL) were added Et₃N (73 mg, 101 μL, 0.72 mmol) and a solution of compound **12** (65 mg, 72 μmol) in DMF (1.4 mL). The reaction mixture was stirred for 18 h, diluted with DCM (20 mL) and purified by gradient column chromatography (0 → 40% MeOH/DCM) to afford **3** as a pale-yellow solid (94 mg, 44 μmol, 28%). LCMS (ESI+) calculated for C₁₀₂H₁₁₈F₂N₁₆O₂₉S₂²⁺ (M/2+H)⁺ 1066.88, found 1067.12.


-67-



-68-

Example 6: Synthesis of linker conjugate 5b


[0283] Compound BCN-HS-vc-PABC-CM (**5b**) was prepared according to the procedures described in WO2019110725, incorporated herein. A solution of compound **16** (172 μ L, 21.5 mg, 38.1 μ mol) was added to a solution of compound **15** (190 μ L, 82.6 mg, 38.1 μ mol, 1.0 equiv.) followed by addition of Et₃N (53 μ L, 38.6 mg, 381 μ mol, 10.0 equiv.). The reaction was left to stand for 14.5 hours at room temperature. The reaction mixture was diluted with DCM to a volume of 6 mL and then purified by automated silicagel flash chromatography (0% --> 20% MeOH in DCM) affording unpure product (72.2 mg, 31.0 μ mol) as an off-white film. The unpure product was further purified by prepHPLC (30% --> 90% CH₃CN / H₂O +1% CH₃COOH, Column Xbridge prep C18 5 μ m OBD, 30 x 100mm, runtime 16 minutes), followed by a 2nd purification by automated silicagel flash chromatography (0% --> 20% MeOH in DCM) affording compound **5b** (45.2 mg, 19.4 μ mol, yield 47%) as a colorless film. LCMS (ESI⁺) calculated for C₁₀₀H₁₄₄IN₁₁O₃₆S₄²⁺ (M+2H⁺)/2 1165.39, found 1165.71.

*Example 7. Synthesis of linker conjugate 6*

[0284] The synthesis of BCN-HS-PEG2-HS-(vc-PABA-Ahx-May)₂ (**6**) is also described in PCT/NL2015/050697 (example 55), incorporated herein. A solution of **18** (3.9 mg, 4.3 μ mol) and Et₃N (3.0 μ L, 2.2 mg, 21.5 μ mol) in DMF (1 mL) was added to a solution of H-Val-Cit-PABA-Ahx-maytansin **17** (10 mg, 8.6 μ mol) in DMF (100 μ L). The mixture was allowed to react overnight and concentrated. The residue was purified via reversed phase (C18) HPLC chromatography (30 $\rightarrow$ 90% MeCN, 1% AcOH) in H₂O (1% AcOH) to give product **6** (3.9 mg, 1.32 μ mol, 31%). LRMS (ESI⁺) m/z calcd for C₁₃₆H₁₉₆Cl₂N₂₂O₄₃S₂ (M+2H⁺)/2=1480.13; found 1480.35. As a side-product, the mono-substituted Ahx-maytansin derivative of **6** was isolated (not depicted). LRMS (ESI⁺) calcd for C₈₅H₁₁₉ClN₁₄O₃₁S₂²⁺ m/z 965.36 found 965.54.

Example 8. Preparation of compound 9

- [0285]** A solution of BCN-HS-PEG₂-b-(Glu(OFm)-OH)₂ (**8**, 12.1 mg, 10 μ mol, 1.0 eq) dissolved in anhydrous DMF (180 μ L) was added to a solution of NH₂-Val-Ala-PABC-exatecan (**5b**, Fmoc-deprotected **5**, 19 mg, 25 μ mol, 2.5 eq) in anhydrous DCM (180 μ L), DIPEA (11 μ L, 63 μ mol, 6.2 eq) and HATU (8.9 mg, 23 μ mol, 2.3 eq). After stirring for 2 h at room temperature, the reaction mixture was further diluted with DCM (800 μ L) and purified by flash column chromatography over silicagel (0% $\rightarrow$ 20% MeOH in DCM) to give the product as a clear oil (difficult to determine yield due to presence of DMF). LCMS (ESI⁺) calculated for C₁₄₀H₁₅₀F₂N₁₇O₃₃S⁺ (M/2+H⁺) 1334.01, found 1334.79.
- [0286]** This compound was dissolved in DMF (300 μ L) and triethylamine (21 μ L, 150 μ mol, 15 eq) was added. After 17 h at room temperature, the reaction mixture was diluted with DCM (700 μ L) and purified by flash column chromatography over silicagel (0% $\rightarrow$ 45% MeOH in DCM) to give compound **9** in 44% yield as a yellow solid (10.2 mg, 4.4 μ L). LCMS (ESI⁺) calculated for C₁₁₂H₁₃₀F₂N₁₇O₃₃S⁺ (M/2+H⁺) 1156.2, found 1156.74.



Examples 9 – 15: Conjugation of linker-payloads to (modified) monoclonal antibodies

*Example 9: Conjugation of labetuzumab(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **labetuzumab-3***

[0287] To a solution of labetuzumab(6-N₃-GalNAc)₂ (844 µL, 18.0 mg, 23.71 mg/ml in TBS pH 7.5) was added sodium deoxycholate (200 µL, 110 mM) and BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (53 µL, 10 mM solution in DMF) and propylene glycol (547 µL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 16/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 4 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated for 3h. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 µm, Corning). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and concentrated using a Vivaspin Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius). 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26492 Da, approximately 90% of total Fc/2), corresponding to the conjugate **labetuzumab-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.90.

20

*Example 10: Conjugation of tusamitamab(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **tusamitamab-3***

[0288] To a solution of tusamitamab(6-N₃-GalNAc)₂ (1762 µL, 37.0 mg, 21.06 mg/ml in TBS pH 7.5) was added sodium deoxycholate (247 µL, 110 mM) and BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (74 µL, 10 mM solution in DMF) and propylene glycol (421 µL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 16/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 19 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated for 3h. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 µm, Corning). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and concentrated using a Vivaspin Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius). 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26461 Da, approximately 90% of total Fc/2), corresponding to the conjugate **tusamitamab-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.89.

40

*Example 11: Conjugation of tusamitamab(6-N₃-GalNAc)₂ with BCN-HS-vc-PABC-calicheamicin **5b** to obtain conjugate **tusamitamab-5b***

[0289] To a solution of tusamitamab(6-N₃-GalNAc)₂ (1424 µL, 30.0 mg, 21.06 mg/ml in TBS pH 7.5) was added sodium deoxycholate (200 µL, 110 mM) and BCN-HS-vc-PABC-calicheamicin **5b**

-71-

(15 μ L, 40 mM solution in DMF) and DMF (185 μ L). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 16/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and concentrated using a Vivaspın Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius). 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26715 Da, approximately 90% of total Fc/2), corresponding to the conjugate **tusamitamab-5b**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 1.94.

10

*Example 12: Conjugation of tusamitamab(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(vc-PABA-Ahx-May)₂ **6** to obtain conjugate **tusamitamab-6***

[0290] To a solution of tusamitamab(6-N₃-GalNAc)₂ (1424 μ L, 30.0 mg, 21.06 mg/ml in TBS pH 7.5) was added BCN-HS-PEG2-HS-(vc-PABA-Ahx-May)₂ **6** (80 μ L, 10 mM solution in DMF) and DMF (420 μ L). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 16/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). Subsequently the solution was buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with 20 mM histidine, 6% sucrose buffer pH 6.0, and concentrated using a Vivaspın Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius). 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 27289 Da, approximately 90% of total Fc/2), corresponding to the conjugate **tusamitamab-6**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.85.

*Example 13: Conjugation of B12(6-N₃-GalNAc)₂ with BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** to obtain conjugate **B12-3***

[0291] To a solution of B12(6-N₃-GalNAc)₂ (6.33 mL, 150.0 mg, 23.71 mg/ml in TBS pH 7.5) was added BCN-HS-PEG2-HS-(va-PAB-Ex)₂ **3** (495 μ L, 10 mM solution in DMF) and propylene glycol (7.0 mL). The reaction was incubated overnight at rt. Next the conjugate was purified on a HiLoad 26/600 Superdex200 PG column (Cytiva) on an AKTA Pure (Cytiva). To remove the excess of free payload, 150 mg of active charcoal (Carbon RHC, Filtrox AG) was added and rotated overnight. The charcoal was removed by centrifugation and subsequently filtered over a PES syringe filter (pore 0.20 μ m, Corning). Subsequently the solution was dialysed to 20 mM histidine, 6% sucrose buffer pH 6.0 for two hours at rt and overnight at 4°C. The solution was concentrated using a Vivaspın Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius) and 0.04% Tween-20 was added before filter sterilization. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26462 Da, approximately 90% of total Fc/2), corresponding to the conjugate **B12-3**. RP-UPLC analysis of the sample under reducing conditions showed an average DAR of 3.75.

40

-72-

Example 14: Production of tusamitamab-DM4

[0292] Tusamitamab (17.5 mg, 2.11 mg/ml in PBS) was charged with DMA to give 5% v/v and 7 molar equivalents SPDB-DM4 (20 mM in DMA) and incubated at rt overnight. The conjugate was purified by preparative SEC (HiLoad 26/600 Superdex pg) into 20 mM histidine / 80 mM NaCl / 6% sucrose. Concentration and buffer exchange into 20 mM histidine / 6% sucrose / pH 6 was performed via discontinuous diafiltration in Vivaspins. 0.04% Tween-20 was added before filter sterilization. The average DAR was measured and was 3.4.

10 *Example 15: Conjugation of labetuzumab(6-N₃-GalNAc)₂ with BCN-HS-PEG2-(eva-PAB-Ex)₂ **9** to obtain conjugate **labetuzumab-9***

[0293] To a solution of labetuzumab(6-N₃-GalNAc)₂ (26 µL, 0.5 mg, 19.35 mg/ml in TBS pH 7.5) was added sodium deoxycholate (5 µL, 110 mM) and BCN-HS-PEG2-(eva-PAB-Ex)₂ **9** (1 µL, 10 mM solution in DMF) and propylene glycol (9 µL). The reaction was incubated overnight at rt. Mass spectral analysis of the sample after IdeS treatment showed one major Fc/2 product (observed mass 26674 Da, approximately 95% of total Fc/2), corresponding to the conjugate **labetuzumab-9**. RP-UPLC analysis of the conjugate under reducing conditions showed an average DAR of 3.88.

Examples 16 – 19: In vitro studies

20 *Example 16. hCEACAM5, cCEACAM5 and hCEACAM6 binding assay to mAbs and ADCs using ELISA*

[0294] Nickel NTA plates (Pierce™ Nickel coated plated, ThermoScientific™) were washed three times prior to use. Human CEACAM5 (CD66e protein, His Tag, Sino Biological), human CEACAM6 (CD66c protein, His Tag, Sino Biological) and cynomolgus CEACAM5 (CD66e protein, His Tag, Sino Biological) were dissolved at a concentration of 0.05 µg/mL in 0.1% BSA in PBS (PBA). 100 µL was added to each well and incubated while shaking for 1 hour at room temperature. After removal, the plate was washed 3x with 0.05% Tween-20 in PBS (washing buffer). ADCs were diluted in 0.1% PBA to a final concentration of 8 µg/mL and 100 µL was added to each well (in triplo). ADCs were incubated for 1h at room temperature. Prior to the addition of 100 µL 1:100 dilution of secondary antibody (Goat anti-human IgG, HRP conjugate, Invitrogen) the plate was washed 3x with washing buffer. The plate was incubated again for 1h at room temperature and subsequently washed 3x with washing buffer and 3x with PBS. Finally, 100 µL TMB ELISA substrate (1 Step™ Turbo TMB ELISA substrate, ThermoScientific™) was added and incubated for 1 minute. The absorbance of the colorimetric signal was measured with Infinite® M1000 (Tecan) at 605 nm. Data was plotted corrected for the background (see Figure 9a and 9b).

Example 17: In vitro cytotoxicity

[0295] MKN-45 (CEACAM+, DSMZ ACC-409) cells were diluted in 80% RPMI 1640 + 20% FBS. The cells were dispensed in a 384-well plate, at a density of 200 cells per well in 45 µl medium. The margins of the plate were filled with phosphate-buffered saline. Plated cells were incubated in a

humidified atmosphere of 5 % CO₂ at 37 °C. After 24 hours, 5 µl of compound dilution was added and plates were further incubated. At t=end, 24 µl of ATPlite 1Step™ (PerkinElmer) solution was added to each well, and subsequently shaken for 2 minutes. After 5 minutes of incubation in the dark, the luminescence was recorded on an Envision multimode reader (PerkinElmer).

- 5 **[0296]** Controls: t = 0 signal. On a parallel plate, 45 µl cells were dispensed and incubated in a humidified atmosphere of 5 % CO₂ at 37 °C. After 24 hours 5 µl DMSO-containing HEPES buffer and 24 µl ATPlite 1Step™ solution were mixed, and luminescence measured after 5 minutes incubation (= *luminescence*_{t=0}). Cell growth control. The cellular doubling times of all cell lines are calculated from the t = 0 hours and t = end growth signals of the untreated cells. If the doubling time is out of specification (0.5 – 2.0 times deviating from historic average) the assay is invalidated. Maximum signals. For each cell line, the maximum luminescence was recorded after incubation until t= end without compound in the presence of 0.4% DMSO (= *luminescence*_{untreated,t=end}).
- 10 **[0297]** Data analysis: IC₅₀s were calculated by non-linear regression using IDBS XLfit 5. The percentage growth after incubation until t= end (%-growth) was calculated as follows: 100% x (*luminescence*_{t=end} / *luminescence*_{untreated,t=end}). This was fitted to the ¹⁰log compound concentration (conc) by a 4-parameter logistics curve : %-growth = bottom + (top – bottom) / (1+ 10^{(logIC₅₀ –conc)*hill}), where hill is the Hill-coefficient, and bottom and top the asymptotic minimum and maximum cell growth that the compound allows in that assay. Survival plots show data normalized to cell viability percentage by setting wells without cells at 0% viability and wells with untreated cells at 100% viability (see Figure 10). IC₅₀ values for ADCs are shown in the table below:
- 20

[0298] Table 1. IC₅₀ values for several ADCs and mAbs for MKN-45 cells

	MKN-45, IC ₅₀ (nM)	
labetuzumab-3	6.5	
tusamitamab-DM4	2.6	25
tusamitamab-3	3.4	
tusamitamab-5b	1.9	
tusamitamab-6	0.88	
B12-3	>100	

30

Example 18. Human and mouse serum stability

- [0299]** Stability of ADCs in mice and human plasma was tested. Prior to the assay, the plasma was depleted from all IgG using CaptivA® Protein A agarose (1 mL agarose/mL serum). ADCs were added to the depleted human/mouse serum to a final concentration of 0.1 mg/mL followed by incubation at 37°C. At each time point, 0.5 mL was snap frozen and stored at -80°C until further analysis. To isolate the ADCs after incubation, 20 µl CaptivA® Protein A agarose resin was added to the samples and incubated for 1 hour at room temperature. The resin was washed three times with PBS and subsequently 0.1 M Glycine-HCl pH 2.7 (0.4 mL) was added to elute the ADCs. After elution, the samples were immediately neutralized with 1.0 M Tris pH 8.0 (0.1 mL). The samples were spin-filtered against PBS for three times using Amicon Ultra spin-filter 0.5 mL MWCO 10 kDa
- 35
- 40

(Merck Millipore) and the volume was reduced to 40 μ L, yielding a final ADC concentration of approximately 1 mg/mL. Samples were analyzed on RP-UPLC (DTT reduced) at specific days to determine the DAR and the results are shown in the table below. At $t = 14$ days, the difference with $t = 0$ days is given in percentage.

5

[0300] Table 2. Stability in human and mouse plasma

Plasma	Conjugate	DAR $t=0$	DAR $t=3$	DAR $t=7$	DAR $t=14$
Human	Tusamitamab-DM4	3.40	3.30	3.10	3.00 (88%)
Human	labetuzumab-3	3.90	3.86	3.79	3.70 (95%)
Mouse	Tusamitamab-DM4	3.40	3.35	3.13	3.21 (94%)
Mouse	labetuzumab-3	3.84	3.78	3.78	3.69 (96%)

Example 19. Thermostability in physiological and enhance stress conditions

[0301] The stability of ADCs was tested at elevated temperatures either at physiological conditions (PBS, pH 7.4, 37°C) or enhanced stress conditions (citrate buffered saline, CBS, pH 5.0, 40°C). The ADCs were buffer exchanged using a HiTrap 26-10 desalting column (Cytiva), rinsed with 0.1M NaOH and equilibrated with either PBS or CBS on an AKTA Pure (Cytiva). The solution was concentrated using a Vivaspın Turbo 4 10 kDa MWCO ultrafiltration unit (Sartorius) to a concentration > 1 mg/mL. The concentration of the ADCs was measured, they were diluted to 1 mg/mL and the first measurement, $t=0$, was taken for SE-HPLC analysis as described above. The samples were placed at either 37°C or 40°C and samples were taken at several timepoints t (in days) and aggregation levels were determined, see tables below.

15

[0302] Table 3. Physiological conditions, monomer levels (in percentage):

	$t=0$	$t=1$	$t=2$	$t=5$	$t=7$	$t=14$	$t=21$	$t=28$
Tusamitamab-DM4	99,0%	99,0%	99,0%	98,9%	98,7%	96,8%	97,0%	95,0%
labetuzumab-3	98,5%	98,7%	98,5%	98,9%	98,0%	99,0%	98,6%	98,4%

20

[0303] Table 4. Enhanced stress conditions, monomer levels (in percentage):

	$t=0$	$t=1$	$t=2$	$t=5$	$t=7$	$t=14$	$t=21$	$t=28$
Tusamitamab-DM4	99,0%	85,3%	89,3%	83,6%	80,3%	75,5%	70,5%	62,5%
labetuzumab-3	98,8%	98,5%	99,0%	98,4%	98,6%	08,7%	98,4%	97,7%

Examples 20 – 23: In vivo studies

Example 20: In vivo efficacy study

[0304] Fragments of the patient-derived tumors, frozen in DMSO/SVF/RPMI 1640 medium (5/10/85) and stored in liquid nitrogen, were thawed at 37°C for 5 min. Then they were rinsed twice in RPMI 1640 medium before subcutaneous implantation in immunodeficient mice and then serially transplanted in immunodeficient mice. The CR-IGR-034P, a colorectal PDX cell line was amplified in fifteen healthy female CB17 SCID (CB17/lcr-Prkdcscid/lcrIcoCrI) mice, 6-weeks old at reception

30

(obtained from Charles River), by implantation subcutaneously into the right flank of each animal. When tumor volumes reached 500-1500 mm³, tumors were surgically excised for the engraftment.

[0305] CB17 SCID mice (female) of 6–7 weeks old (obtained from Charles River) received a subcutaneous injection in the right flank region with tumor fragments amplified as described above.

5 When tumors reached an average size of 100-200 mm³, randomization was performed into 10 groups of 8 mice and treatment began. Homogeneity between groups was tested by an analysis of variance (ANOVA).

[0306] The test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. Treatment was initiated on the same day of randomization.

10 Dosing was conducted in a Laminar Flow Cabinet.

[0307] After tumor cells inoculation, the animals were checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects and are recorded in the Vivo

15 Manager database (Biosystemes, France). Tumor volumes are measured every 3-4 days in two dimensions using an caliper, and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be conducted in a Laminar Flow Cabinet. The endpoint of the experiment

20 is a tumor volume of 2,000 mm³, body weight loss over 20% or 28 days, whichever comes first.

[0308] Table 5. Overview of test items and dose levels included in in vivo study

Test item	Dose levels (mg/kg)	Group (N=8)	Administration	Dosing Schedule
vehicle	–	1	IV	Single dose
tusamitamab-DM4	5.7	2	IV	Single dose
labetuzumab-3	5	3	IV	Single dose
	10	4	IV	Single dose
tusamitamab-3	5	5	IV	Single dose
	10	6	IV	Single dose
tusamitamab-5b	3	8	IV	Single dose
	5	9	IV	Single dose
tusamitamab-6	5	7	IV	Single dose
B12-3	10	10	IV	Single dose

Data for efficacy study with the test items above is depicted in Figures 11a – 11d.

25 *Example 21: In vivo efficacy study*

[0309] Fragments of the patient-derived tumors, frozen in DMSO/SVF/RPMI 1640 medium (5/10/85) and stored in liquid nitrogen, were thawed at 37°C for 5 min. Then they were rinsed twice in RPMI 1640 medium before subcutaneous implantation in immunodeficient mice and then serially

transplanted in immunodeficient mice. The CR-IGR-002P, a colorectal PDX cell line was amplified in fifteen healthy female CB17 SCID (CB17/lcr-Prkdcscid/lcrIcoCrI) mice, 6-weeks old at reception (obtained from Charles River), by implantation subcutaneously into the right flank of each animal. When tumor volumes reached 500-1500 mm³, tumors were surgically excised for the engraftment.

5 **[0310]** CB17 SCID mice (female) of 6–7 weeks old (obtained from Charles River) received a subcutaneous injection in the right flank region with tumor fragments amplified as described above. When tumors reached an average size of 100-200 mm³, randomization was performed into 5 groups of 6 mice and treatment began. Homogeneity between groups was tested by an analysis of variance (ANOVA).

10 **[0311]** The test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. Treatment was initiated on the same day of randomization. Dosing was conducted in a Laminar Flow Cabinet.

[0312] After tumor cells inoculation, the animals were checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects and are recorded in the Vivo Manager database (Biosystemes, France). Tumor volumes are measured every 3-4 days in two dimensions using an caliper, and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be conducted in a Laminar Flow Cabinet. The endpoint of the experiment is a tumor volume of 2,000 mm³, body weight loss over 20% or 29 days, whichever comes first.

[0313] Table 6. Overview of test items and dose levels included in in vivo study

Test item	Dose levels (mg/kg)	Group (N=8)	Administration	Dosing Schedule
vehicle	–	1	IV	Single dose
tusamitamab-DM4	5	2	IV	Single dose
labetuzumab-3	10	3	IV	Single dose
tusamitamab-3	10	4	IV	Single dose
B12-3	10	5	IV	Single dose

25 Data for efficacy study with the test items above is depicted in Figure 12A and 12B.

Example 22: In vivo efficacy study of several PDX models, a single mouse trial

[0314] Fragments of the patient-derived tumors, frozen in DMSO/SVF/RPMI 1640 medium (5/10/85) and stored in liquid nitrogen, were thawed at 37°C for 5 min. Then they were rinsed twice in RPMI 1640 medium before subcutaneous implantation in immunodeficient mice and then serially transplanted in immunodeficient mice. The PDX cell lines, see table below, were amplified in 12 healthy female CB17 SCID (CB17/lcr-Prkdcscid/lcrIcoCrI) mice, 6-weeks old at reception (obtained from Charles River) and 9 healthy NSG (NOD.Cg.PrkdcSCID Il2rgtmWjl/SzJ, obtained from

Charles River), by implantation subcutaneously into the right flank of each animal. When tumor volumes reached 500-1500 mm³, tumors were surgically excised for the engraftment.

[0315] CB17 SCID mice (female) of 6–7 weeks old (obtained from Charles River) received a subcutaneous injection in the right flank region with tumor fragments amplified as described above.

5 When tumors reached an average size of 100-200 mm³, randomization was performed and treatment began. Homogeneity between groups was tested by an analysis of variance (ANOVA).

[0316] The test article administration was performed via intravenous injection through tail vein, and the dosing volume was 10 mL/kg. Dose was either vehicle or 10 mg/kg labetuzumab-3. Treatment was initiated on the same day of randomization. Dosing was conducted in a Laminar Flow Cabinet.

10 **[0317]** After tumor cells inoculation, the animals were checked daily for morbidity and mortality. At the time of routine monitoring, the animals are checked for any adverse effects of tumor growth and treatments on normal behavior such as mobility, visual estimation of food and water consumption, body weight gain/loss, eye/hair matting and any other abnormal effects and are recorded in the Vivo Manager database (Biosystemes, France). Tumor volumes are measured every 3-4 days in two
15 dimensions using an caliper, and the volume data are expressed in mm³ using the formula: $V = (L \times W \times W)/2$, where V is tumor volume, L is tumor length (the longest tumor dimension) and W is tumor width (the longest tumor dimension perpendicular to L). Dosing as well as tumor and body weight measurements will be conducted in a Laminar Flow Cabinet. The endpoint of the experiment is a tumor volume of 2,000 mm³, body weight loss over 20% or 1.5 times survival over the vehicle,
20 whichever comes first.

[0318] Tumor growth inhibition is calculated with the following equation:

$$\frac{T}{C} \% = 100 \times \frac{\text{Median tumor volume of treated group at } Dx}{\text{Median tumor volume of vehicle treated group at } Dx}$$

Where DX: Day of tumor volume measurement. The optimal value is the minimal T/C% ratio reflecting the maximal tumor growth inhibition achieved. Tumor growth inhibition is classified as
25 follows:

<20% : high inhibition
21 - 40% : moderately high inhibition
41 – 80% : moderate inhibition
>81% : low inhibition

30

[0319] Table 7. Overview and characteristics of the PDX models

PDX model	Animals for amplification	Indication	CEACAM expression level (H-score)	Intensity of staining (IHC)	Irinotecan sensitive*
CR-IC-003P	SCID mice	Colorectal	ND	2+	no
CR-LRB-010P	SCID mice	Colorectal	ND	1+	yes
LUN-NIC-014	SCID mice	Lung	120	2+	unknown
IM-PAN-007	NSG mice	Pancreatic	270	3+	yes
SA-PAN-083	NSG mice	Pancreatic	70	1+	yes

* Yes means >40% irinotecan response

Figure 13A – 13O show the efficacy plots, body weight plots and Kaplan-Meier plots per PDX model. Table 8 below shows the TGI% per model.

5 Table 8. TGI% per model

PDX model	TGI%	Classification
CR-IC-003P	8.3 %	High inhibition
CR-LRB-010P	29.4 %	Moderately high inhibition
LUN-NIC-014	11.9 %	High inhibition
IM-PAN-007	11.8 %	High inhibition
SA-PAN-083	11.2 %	High inhibition

Example 23: In vivo tolerability study

15 **[0320]** Sprague Dawley rats (female) of 6-7 weeks old acclimatized for a week before being enrolled in the study.

[0321] Randomization: In the tolerability study totally 15 rats were enrolled and randomly allocated to 5 study groups with 3 rats per group. Randomization was performed based on “Matched distribution” method (StudyDirector™ software, version 3.1.399.19). The date of randomization was denoted as day 0.


20 **[0322]** Test Article Administration: The treatment was initiated on the same day of randomization (day 0) per study design. Chosen dose levels were 80, 100, 120 and 140 mg/kg, single dose. All animals were dosed via slow intravenous injection (10 mL/kg).

25 **[0323]** Observation and Data Collection After randomization: The animals were checked daily for morbidity and mortality. During routine monitoring, the animals were checked for any effects of treatments on behavior such as mobility, food and water consumption, body weight gain/loss, eye/hair matting and any other abnormalities. Mortality and observed clinical signs were recorded for individual animals in detail. Dosing as well as body weight measurements were conducted in a Laminar Flow Cabinet. The body weights were measured daily by using StudyDirector™ software (version 3.1.399.19).

30 **[0324]** Body weight of the rats over time with labetuzumab-3 as test item at several dose levels are depicted in Figure 14.

Claims

1. An antibody-conjugate according to general structure (1):



(1)

wherein:

- AB is an antibody capable of targeting CEA-expressing tumours;
- L is a linker that links Z to D;
- Z is a connecting group;
- L^6 is $-\text{GlcNAc}(\text{Fuc})_w-(\text{G})_j-\text{S}-(\text{L}^7)_w-$, wherein G is a monosaccharide, j is an integer in the range of 0 – 10, S is a sugar or a sugar derivative, GlcNAc is N-acetylglucosamine and Fuc is fucose, w is 0 or 1, w' is 0, 1 or 2 and L^7 is $-\text{N}(\text{H})\text{C}(\text{O})\text{CH}_2-$, $-\text{N}(\text{H})\text{C}(\text{O})\text{CF}_2-$ or $-\text{CH}_2-$;
- D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- b is 0 or 1;
- x is 1 or 2; and
- y is 1, 2, 3 or 4.

20

2. The antibody-conjugate according to claim 1, wherein D is selected from the group consisting of anthracyclines, camptothecins, maytansinoids, enediynes, auristatins and pyrrolobenzodiazepine dimers, preferably wherein:

- D is an enediyne, preferably selected from the group consisting of calicheamicins, esperamicins, shishijimicins and namenamicins, more preferably calicheamicin; or
- D is an auristatin, preferably MMAD, MMAE, MMAF or PF-06380101, more preferably MMAE or PF-06380101; or
- D is a camptothecin, preferably selected from the group consisting of topotecan, silatecan, cositecan, exatecan, exatecan-S, DXd, SN-38, lurtotecan, gimatecan, belotecan, rubitecan, AMDCPT and G-AMDCPT, more preferably exatecan or DXd, most preferably exatecan, more preferably D is calicheamicin or exatecan, more preferably D is exatecan.

25

30


3. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is selected from labetuzumab, A5/A240VL, D8/A240VL, PR1A3, CH1A1A/pAC18, CH1A1B/pAC18, AB17, AB72, AB73, NCC-CO-413, NCC-CO-308, NCC-CO-432, NCC-CO-411, A5B7, 1G9, CEA6, M5A, M5B, SM3E, 12-140-1, 15-1-32, huMAb2-4, tusamitamab, and humanized or functional analogues thereof, preferably is PR1A3, SM3E, huMAb2-4, tusamitamab or labetuzumab, more preferably wherein the antibody is tusamitamab or labetuzumab, most preferably wherein the antibody is labetuzumab, optionally wherein the Fc region of the antibody may have one or more mutations.

40

4. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody contains a V_L domain having at least 70 % sequence identity with a sequence selected from the group consisting of SEQ ID No. 2, 6, 10, 17, 21, 28, 31, 35, 40, 45, 49, 50 and 51, and a V_H domain having at least 70 % sequence identity with a sequence selected from the group consisting of SEQ ID No. 1, 3, 5, 9, 11, 13, 16, 20, 24, 26, 30, 34, 39, 44 and 48, preferably wherein the antibody has a complementarity-determining region (CDR) having at least 90 % sequence identity.
5. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is labetuzumab, and/or contains a light chain sequence according to SEQ ID No. 38 and a heavy chain sequence according to SEQ ID No. 36 or 37, preferably SEQ ID No. 36, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
6. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is tusamitamab, and/or contains a light chain sequence according to SEQ ID No. 43 and a heavy chain sequence according to SEQ ID No. 41 or 42, preferably SEQ ID No. 41, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
7. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is SM3E, and/or contains a light chain sequence according to SEQ ID No. 19 and a heavy chain sequence according to SEQ ID No. 18, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
8. The antibody-conjugate according to any one of claims 1 – 2, wherein the antibody is PR1A3, and/or contains a light chain sequence according to SEQ ID No. 23 and a heavy chain sequence according to SEQ ID No. 22, wherein the sequence identity is at least 90 %, preferably at least 95 %, more preferably at least 99 %, most preferably 100 %.
9. The antibody-conjugate according to any one of claims 1 – 8, wherein Z comprises the reaction product of a nucleophilic reaction or a cycloaddition reaction, preferably a 1,3-dipolar cycloaddition reaction.
10. The antibody-conjugate according to any one of claims 1 – 9, wherein linker L has the structure
- $$-(L^1)_n-(L^2)_o-(L^3)_p-(L^4)_q-$$
- wherein:
- L¹, L², L³ and L⁴ are each individually linkers that together link Z to D;
 - n, o, p and q are each individually 0 or 1, provided that n + o + p + q = 1, 2, 3 or 4, preferably wherein n = o = p = 1;


-81-

and wherein:

(a) linker L¹ is represented by:

5 wherein:

- d and d' are individually 0 or 1;
- e and e' are individually an integer in the range 1 – 10;
- f and f' are individually 0, or 1;
- g and g' are individually an integer in the range 0 – 10;
- k = 0 or 1 with the proviso that if k = 1 then d = 0;
- A is a sulfamide group according to structure (23)



(23)

15 wherein a = 0 or 1, and R¹³ is selected from the group consisting of hydrogen, C₁ – C₂₄ alkyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups and C₃ – C₂₄ (hetero)arylalkyl groups, the C₁ – C₂₄ alkyl groups, C₃ – C₂₄ cycloalkyl groups, C₂ – C₂₄ (hetero)aryl groups, C₃ – C₂₄ alkyl(hetero)aryl groups and C₃ – C₂₄ (hetero)arylalkyl groups optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR¹⁴ wherein R¹⁴ is

20 independently selected from the group consisting of hydrogen and C₁ – C₄ alkyl groups, or R¹³ is D connected to N via a spacer moiety, preferably wherein the spacer moiety is

25 $-(B)_g-(C(O))_g-(L^2)_o-(L^3)_p-(L^4)_q-$;

- W is $-OC(O)-$, $-C(O)O-$, $-C(O)NH-$, $-NHC(O)-$, $-OC(O)NH-$, $-NHC(O)O-$, $-C(O)(CH_2)_mC(O)-$, $-C(O)(CH_2)_mC(O)NH-$ or $-(4-Ph)CH_2NHC(O)(CH_2)_mC(O)NH-$, wherein m is an integer in the range 0 – 10;


- B is a $-CH_2-CH_2-O-$ or a $-O-CH_2-CH_2-$ moiety, or (B)_e is a $-(CH_2-CH_2-O)_{e1}-CH_2-CH_2-$ moiety, wherein e₁ is an integer in the range 1 – 10;

- N* is a branching nitrogen atom, to which two instances of $-(A)_d-(B)_e-(A)_f-(C(O))_g-$ are connected and both (C(O))_g moieties are connected to $-(L^2)_o-(L^3)_p-(L^4)_q-D$, wherein L², L³, L⁴, o, p, q and D are each selected individually;

30

and/or

(b) linker L² is a peptide spacer, preferably comprising 1 – 5 amino acids, more preferably a dipeptide, tripeptide or tetrapeptide spacer, most preferably wherein L² is represented by general structure (L3):



(L3)


35

-82-

wherein $R^{17} = CH_3$ or $CH_2CH_2CH_2NHC(O)NH_2$;

and/or

- (c) linker L^3 is a self-immolative spacer, preferably a para-aminobenzyloxycarbonyl (PABC) derivative according to structure (L4).



(L4)


wherein R^{21} is H, R^{26} or $C(O)R^{26}$, wherein R^{26} is $C_1 - C_{24}$ (hetero)alkyl groups, $C_3 - C_{10}$ (hetero)cycloalkyl groups, $C_2 - C_{10}$ (hetero)aryl groups, $C_3 - C_{10}$ alkyl(hetero)aryl groups and $C_3 - C_{10}$ (hetero)arylalkyl groups, which are optionally substituted and optionally interrupted by one or more heteroatoms selected from O, S and NR^{28} wherein R^{28} is independently selected from the group consisting of hydrogen and $C_1 - C_4$ alkyl groups, preferably wherein R^{21} is H or $C(O)R^{26}$, wherein $R^{26} = 4$ -methyl-piperazine or morpholine, most preferably wherein R^{21} is H;

and/or

- (d) linker L^4 is an aminoalkanoic acid spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-C(O)-$, wherein x is an integer in the range 1 – 20 and R^{22} is H or $C_1 - C_4$ alkyl; or linker L^4 is an ethyleneglycol spacer according to the structure $-NR^{22}-(CH_2-CH_2-O)_{e6}-(CH_2)_{e7}-C(O)-$, wherein $e6$ is an integer in the range 1 – 10, $e7$ is an integer in the range 1 – 3 and R^{22} is H or $C_1 - C_4$ alkyl; or linker L^4 is a diamine spacer according to the structure $-NR^{22}-(C_z\text{-alkylene})-NR^{22}-(C(O))_h-$, wherein h is 0 or 1, x is an integer in the range 1 – 10 and R^{22} is H or $C_1 - C_4$ alkyl.


11. The antibody-conjugate according to any one of claims 1 – 10, wherein $b = 1$ and conjugation is via the glycan of the antibody, preferably wherein $j = 0$.

12. The antibody-conjugate according to any one of claims 1 – 11, wherein Z–L–D has a structure selected from the group consisting of (X) – (XII):




(X)


-83-



(XI)



(XII)




(XIII)

5

wherein the wavy line indicates the connection to Z, and L², L⁴, o and q are as defined in claim 10,


preferably wherein Z-L-D has a structure selected from the group consisting of (Xa), (Xib), (XIIg), (XIIle) and (XIIh):

10




(Xa)


-84-



(XIb)




(XIlg)



(XIIIe)


-85-



(XIIh)

wherein the wavy line indicates the connection to Z, and R²² is as defined in claim 10.

13. The antibody-conjugate according to claim 12, which is according to structure (XII) or (XIII), and wherein o = 1 and the antibody is labetuzumab as defined in claim 5, preferably wherein q = 0.
14. The antibody-conjugate according to claim 13, which is according to structure (XII) or (XIII), wherein q = 0, o = 1, L² is a peptide spacer containing 1 – 5 amino acids, the antibody is labetuzumab as defined in claim 5 and the payload is exatecan.
15. The antibody-conjugate according to claim 1, which is according to structure (XIIIe) as defined in claim 12, and wherein the antibody is labetuzumab as defined in claim 5 and the payload is exatecan.
16. A process for preparing the antibody-conjugate according to any one of claims 1 – 15, comprising:
 - (i) contacting an antibody comprising y core N-acetylglucosamine (GlcNAc) moieties, wherein y = 1, 2, 3 or 4, with a compound of the formula S(F)_xP in the presence of a catalyst, wherein S(F)_x is a sugar derivative comprising x reactive groups F capable of reacting with a reactive group Q, x is 1 or 2 and P is a nucleoside mono- or diphosphate, and wherein the catalyst is capable of transferring the S(F)_x moiety to the core-GlcNAc moiety, to obtain a modified antibody according to Formula (26):



(26)

wherein

- AB is an antibody capable of targeting CEA-expressing tumours;
- Fuc is fucose;

-86-

- w is 0 or 1; and

(ii) reacting the modified antibody with a compound according to structure (2):




(2)

5 wherein:


- Q is a reactive moiety;
 - L is a linker that links Z to D;
 - D is selected from the group consisting of anthracyclines, camptothecins, tubulysins, enediynes, amanitins, duocarmycins, maytansinoids, auristatins, eribulins, BCL-XL inhibitors, hemiasterlins, KSP inhibitors, TLR agonists, indolinobenzodiazepine dimers or pyrrolobenzodiazepine dimers (PBDs), and analogues or prodrugs thereof;
- 10 to obtain the antibody-conjugate according to structure (1).

17. The process according to claim 16, wherein the reaction of step (ii) is a nucleophilic reaction or a cycloaddition, preferably a 1,3-dipolar cycloaddition, and preferably wherein Q comprises or is an alkyne moiety and F comprises or is an azide moiety.

18. The process according to claim 16 or 17, wherein Q is a click probe comprising a (hetero)cycloalkyne moiety or a (hetero)cycloalkene moiety, preferably wherein the click probe Q is selected from the group consisting of (Q21) – (Q38) and (Q44) – (Q56):



-87-



wherein

- B is an anion; and
- the R group(s) on Si in (Q50) and (Q51) are alkyl or aryl;

5 and wherein F is a click probe selected from the group consisting of azide, tetrazine, triazine, nitron, nitrile oxide, nitrile imine, diazo compound, *ortho*-quinone, dioxothiophene and sydnone, preferably F is an azide moiety.

- 10 19. Method for targeting CEA-expressing cells, contacting the antibody-conjugate according to any one of claims 1 – 15 with cells that may possibly be CEA-expressing, preferably wherein the cells are CEA-expressing tumour cells.
- 15 20. The method according to claim 19, wherein the targeting CEA-expressing cells includes one or more of treating, imaging, diagnosing, preventing the proliferation of, containing and reducing CEA-expressing cells, in particular CEA-expressing tumour cells.
21. The method according to claim 19 or 20, wherein the subject suffers from colorectal cancer, gastric cancer, lung cancer, uterine cancer or pancreatic cancer.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 11(C)




Fig. 11(D)




Fig. 12(A)




Fig. 12(B)







Fig. 13(A)*Fig. 13(B)**Fig. 13(C)*

Fig. 13(D)




Fig. 13(E)




Fig. 13(F)







Fig. 13(G)*Fig. 13(H)**Fig. 13(I)*




Fig. 13(J)*Fig. 13(K)**Fig. 13(L)*

Fig. 13(M)




Fig. 13(N)




Fig. 13(O)





Fig. 14

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2023/057565

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☒ forming part of the international application as filed.
 - b. ☐ furnished subsequent to the international filing date for the purposes of international search (Rule 13*ter*.1(a)).
☐ accompanied by a statement to the effect that the sequence listing does not go beyond the disclosure in the international application as filed.
2. ☐ With regard to any nucleotide and/or amino acid sequence disclosed in the international application, this report has been established to the extent that a meaningful search could be carried out without a WIPO Standard ST.26 compliant sequence listing.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2023/057565

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K47/68 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2017/137458 A1 (SYNAFFIX BV [NL]) 17 August 2017 (2017-08-17)	1, 2, 9-12, 16-21
Y	page 23, lines 1-17 page 26, line 24 - page 27, line 10 page 32, lines 3-12 examples 11-29; compounds 63, 66, 67, 100, 108, 111 ----- -/--	5-8



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 June 2023

Date of mailing of the international search report

07/07/2023

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bliem, Barbara

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2023/057565

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GAZZAH A. ET AL: "Safety, pharmacokinetics, and antitumor activity of the anti-CEACAM5-DM4 antibody-drug conjugate tusamitamab ravtansine (SAR408701) in patients with advanced solid tumors: first-in-human dose-escalation study", ANNALS OF ONCOLOGY, vol. 33, no. 4, 19 January 2022 (2022-01-19), pages 416-425, XP055950615, NL ISSN: 0923-7534, DOI: 10.1016/j.annonc.2021.12.012	1-4, 6, 19-21
Y	abstract page 423, column 2, paragraph 4 -----	6
Y	DOTAN EFRAT ET AL: "Phase I/II Trial of Labetuzumab Govitecan (Anti-CEACAM5/ SN-38 Antibody-Drug Conjugate) in Patients With Refractory or Relapsing Metastatic Colorectal Cancer", JOURNAL OF CLINICAL ONCOLOGY, vol. 35, no. 29, 17 August 2017 (2017-08-17), XP055950623, abstract -----	5
Y	ASHRAF S Q ET AL: "Humanised IgG1 antibody variants targeting membrane-bound carcinoembryonic antigen by antibody-dependent cellular cytotoxicity and phagocytosis", BRITISH JOURNAL OF CANCER, NATURE PUBLISHING GROUP UK, LONDON, vol. 101, no. 10, 17 November 2009 (2009-11-17), pages 1758-1768, XP002609361, ISSN: 0007-0920, DOI: 10.1038/SJ.BJC.6605355 page 1759, column 2, paragraph 2-3 -----	7, 8
A	WO 2017/137456 A1 (SYNAFFIX BV [NL]) 17 August 2017 (2017-08-17) ----- -/--	1-19

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2023/057565

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	REMON VAN GEEL ET AL: "Chemoenzymatic Conjugation of Toxic Payloads to the Globally Conserved N-Glycan of Native mAbs Provides Homogeneous and Highly Efficacious Antibody&ndash;Drug Conjugates", BIOCONJUGATE CHEMISTRY, vol. 26, no. 11, 10 June 2015 (2015-06-10), pages 2233-2242, XP055249394, US ISSN: 1043-1802, DOI: 10.1021/acs.bioconjchem.5b00224 -----	1-19
A	JORGE M. M. VERKADE ET AL: "Chemoenzymatic Conjugation of Toxic Payloads to the Globally Conserved N-Glycan of Native mAbs Provides Homogeneous and Highly Efficacious Antibody&ndash;Drug Conjugates", ANTIBODIES, vol. 7, no. 1, 20 February 2018 (2018-02-20), page 12, XP055475317, DOI: 10.3390/antib7010012 -----	1-19

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2023/057565

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 2017137458 A1		17-08-2017	CN	108883191 A		23-11-2018
			EP	3413916 A1		19-12-2018
			WO	2017137458 A1		17-08-2017

WO 2017137456 A1		17-08-2017	CN	109152844 A		04-01-2019
			EP	3413920 A1		19-12-2018
			JP	2019507741 A		22-03-2019
			JP	2022058417 A		12-04-2022
			US	2019262467 A1		29-08-2019
			WO	2017137456 A1		17-08-2017
