



- (51) International Patent Classification:  
*C07D 401/12* (2006.01)
- (21) International Application Number:  
PCT/US2015/066741
- (22) International Filing Date:  
18 December 2015 (18.12.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
62/096,077 23 December 2014 (23.12.2014) US
- (71) Applicants: **CERECOR, INC.** [US/US]; 400 E. Pratt Street, Suite 606, Baltimore, MD 21202 (US). **MERCK SHARP DOHME & CORP.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).
- (72) Inventors: **MAZHARI, Reza**; 551 Allegheny Avenue, Towson, MD 21204 (US). **MEZAACHE, Djelila**; 42135 Pepperbush Place, Stone Ridge, VA 20105 (US). **PATERSON, Blake, M.**; 1000 Fells Street, Baltimore, MD 21231 (US). **VORNOV, James**; 3519 Old Court Road, Pikesville, MD 21208 (US). **GARNER, Rachel, M.**; 8113 Belona Ave., Baltimore, MD 21204 (US). **NELSON, Todd**; 8 Orchard Drive, Cream Ridge, NJ 08514 (US).
- (74) Agents: **ZHU, Jin** et al.; Fox Rothschild LLP, 997 Lenox Drive, Building 3, Lawrenceville, NJ 08648-2311 (US).
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report (Art. 21(3))

[Continued on next page]

(54) Title: COMPOUNDS, COMPOSITIONS AND METHODS





FIG. 1A



(57) Abstract: The disclosed subject matter provides certain polymorphic forms of Compound (I) as well as pharmaceutical compositions comprising Compound (I) or such polymorphic forms, and methods of using or making such compounds and pharmaceutical compositions. It has now been discovered that Compound (I) can exist in multiple crystalline forms (polymorphs). One particular crystalline form, Form II, has been found to be more thermodynamically stable and, thus, likely more suitable for bulk preparation and handling than other polymorphic forms. Efficient and economic methods have been developed to prepare Compound (I) and Form II in high purity on a large scale. In animal studies, Form II has demonstrated safety and efficacy in treating depressive disorders and, when micronized, improved absorption compared to non-micronized Form II.

- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*


## COMPOUNDS, COMPOSITIONS AND METHODS

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims benefit under 35 U.S.C. §119(e) of U.S. Provisional Application Serial No. 62/096,077, filed December 23, 2014, which is hereby incorporated  
5 by reference.

### BACKGROUND

U.S. Pat. No. 7,592,360 (issued September 22, 2009) discloses Compound (I):




(I)

10 or 4-methylbenzyl (3S, 4R)-3-fluoro-4-[(pyrimidin-2-ylamino) methyl] piperidine-1-  
carboxylate (referred to in the '360 patent as (3S, 4R)-4-methylbenzyl 3-fluoro-4-  
[(pyrimidin-2-ylamino)methyl]piperidine-1-carboxylate, and elsewhere as MK-0657 or  
CERC-301). A potent selective antagonist of N-methyl-D-aspartate receptor subunit 2B  
(NMDA-GluN2B or NR2B), Compound (I) was initially developed for treatment of  
15 Parkinson's disease (Addy *et al.*, *J. Clin. Psychopharmacol.*, 49:856-864 (2009)). A pilot  
study of Compound (I) in patients with treatment-resistant major depressive disorder  
(TRMDD) showed antidepressant effects in the 17-item Hamilton Depression Rating  
Scale (HAM-D17) and Beck Depression Inventory (BDI) (Ibrahim *et al.*, *J. Clin.*  
*Psychopharmacol.*, 32(4):551-557 (2012)). Due to the study's small sample size, no  
20 definitive conclusions as to Compound (I)'s potential efficacy or safety profile could be  
drawn from the study's preliminary data.

### SUMMARY OF THE DISCLOSURE

The disclosed subject matter provides certain polymorphic forms of Compound (I)



(I)

as well as pharmaceutical compositions comprising Compound (I) or such polymorphic forms, and methods of using or making such compounds and pharmaceutical  
5 compositions.

It has now been discovered that Compound (I) can exist in multiple crystalline forms (polymorphs). One particular crystalline form, Form II, has been found to be more thermodynamically stable and, thus, likely more suitable for bulk preparation and handling than other polymorphic forms. Efficient and economic methods have been developed to  
10 prepare Compound (I) and Form II in high purity on a large scale. In animal studies, Form II has demonstrated safety and efficacy in treating depressive disorders and, when micronized, improved absorption compared to non-micronized Form II.

One aspect of the disclosure provides a compound that is a substantially pure crystalline Form II of Compound (I) exhibiting at least one of:

- 15 (i) an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, comprising peaks of 2-theta angles of about 5.9 and 8.8 degrees;
- (ii) an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, substantially as shown in FIG. 1A;
- (iii) an ultraviolet absorbance spectrum, obtained using methanol as diluent,  
20 substantially as shown in FIG. 2;
- (iv) an infrared spectrum substantially as shown in FIG. 3;
- (v) a proton nuclear magnetic resonance spectrum at about 600 MHz in CD<sub>3</sub>CN substantially as shown in FIG. 4;
- (vi) a <sup>13</sup>C nuclear magnetic resonance spectrum at about 150 MHz in CD<sub>3</sub>CN  
25 substantially as shown in FIG. 5;
- (vii) a thermogravimetric analysis curve substantially as shown in FIG. 6; and
- (viii) a differential scanning calorimetry thermogram substantially as shown in FIG. 7.

Also provided herein is a crystalline Form I of Compound (I) as described in FIG 1B.

Another aspect of the disclosure provides a pharmaceutical composition of the disclosure comprises particles of Compound (I) with an X90 particle size of about 10  $\mu\text{m}$  or less. Suitable particles of Compound (I) include for example microparticles and nanoparticles.

Another aspect of the disclosure provides a method of treating a condition responsive to an NR2B antagonist. The method includes administering to a patient in need thereof an effective amount of a compound or pharmaceutical composition of the present invention.

Another aspect of the disclosure provides a method of treating suicidal ideation, comprising administering Form I or Form II of Compound (I) to a patient who has, is suspected of having, or has been diagnosed with having suicidal ideation.


Another aspect of the disclosure provides a method of targeting N-methyl-D-aspartate (NMDA) receptor subunit 2B (GluN2B) expressed on a cell comprising administering to a patient in need thereof an effective amount of a compound or pharmaceutical composition of the present invention.

Another aspect of the disclosure provides a method of reducing absorption rate of Compound (I) comprising administering to a patient in need thereof an effective amount of the Compound (I) with food, wherein the compound is administered either substantially concurrently with, or up to about 2 hours after, or up to about 30 minutes before administration of food.

Another aspect of the disclosure provides a method of preparing Compound (I), analogs and associated intermediates.

Another aspect of the disclosure provides a method of preparing Compound (I) comprising:


- (i) reacting Compound (8)



(8)

with triflic anhydride to yield a triflate;

(ii) reacting the triflate with ammonia to yield Compound (9)



(9)

(iii) reacting the Compound (9) with 2-chloropyrimidine to yield Compound (I).

Another aspect of the disclosure provides a method of preparing Form II of

Compound (I). By suspending purified Form I of Compound (I) in water for a sufficient period of time or heating solid Form I, conversion to Form II can be readily achieved.

#### BRIEF DESCRIPTION OF FIGURES

**FIG. 1A** shows an X-ray powder diffraction pattern of crystalline Form II of Compound (I).

**FIG. 1B** shows X-ray powder diffraction patterns of crystalline Form I and crystalline Form II of Compound (I).

**FIG. 2** shows an ultraviolet absorbance spectrum of crystalline Form II of Compound (I).

**FIG. 3** shows an infrared spectrum of crystalline Form II of Compound (I).

**FIG. 4** shows proton nuclear magnetic resonance spectrum of crystalline Form II of Compound (I).

**FIG. 5** shows a <sup>13</sup>C nuclear magnetic resonance spectrum of crystalline Form II of Compound (I).

**FIG. 6** shows a thermogravimetric analysis curve of crystalline Form II of Compound (I).

**FIG. 7** shows a differential scanning calorimetry thermogram of crystalline Form II of Compound (I).

**FIG. 8** shows the dose response curve for antidepressant effects of crystalline Form II of Compound (I) relative to its locomotor effect in rats.

**FIG. 9** shows the dose response curve for hypertensive effects of crystalline Form II of Compound (I) in rats.

**FIG. 10A** shows the pharmacokinetic profile (plasma concentrations) of crystalline Form II of Compound (I) in healthy humans in fasted state.

**FIG. 10B** shows the difference in pharmacokinetics of crystalline Form II of Compound (I) in fed versus fasted state in healthy humans.

**FIG. 11** shows effects of a single oral dose of Compound (I) on systolic blood pressure.

**FIG. 12** shows effects of a single oral dose of Compound (I) on systolic blood pressure, with or without prazosin.

**FIG. 13** shows the effect of Compound (I) and ketamine on behaviors in the rat forced swim test 24 hours after administration (\* $p < 0.05$  compare to vehicle group ~ $p > 0.05$ ).

## **DETAILED DESCRIPTION**

### **Definitions**

Unless clearly indicated otherwise, the following terms as used herein have the meanings indicated below.

“Micronized” refers to particles with a diameter within the micron range. Methods of producing micronized particles include friction-based techniques, such as milling (*e.g.*, jet-milling) and grinding, and supercritical fluid-based techniques, such as the Rapid Expansion of Supercritical Solutions (RESS), Supercritical Anti-Solvent (SAS), and Particles from Gas Saturated Solutions (PGSS) methods. In some embodiments, the Compound (I) particles are micronized by jet-milling.

“Particle size” refers to the particle dimension of the active pharmaceutical ingredient (API), such as Compound (I) or Form (II) of Compound (I), as ascertained by laser diffraction particle size analysis, performed for example using an analyzer such as Malvern, Sympatec, Microtac or Horibe.

“X90” refers to the particle size corresponding to 90% of the cumulative undersize distribution by volume.

“X50” refers to the particle size corresponding to 50% of the cumulative undersize distribution by volume.

“X10” refers to the particle size corresponding to 10% of the cumulative undersize distribution by volume.

5 “Mean particle size” or “mean PS” includes D43 particle size.

“D43” refers to the particle size calculated according to the mean diameter over volume or DeBroukere mean.

“Pharmaceutically acceptable excipient” refers to any substance, not itself a therapeutic agent, used as a carrier, diluent, adjuvant, binder, and/or vehicle for delivery of a therapeutic agent to a patient, or added to a pharmaceutical composition to improve its handling or storage properties or to permit or facilitate formation of a compound or pharmaceutical composition into a unit dosage form for administration. Pharmaceutically acceptable excipients are known in the pharmaceutical arts and are disclosed, for example, in *Remington: The Science and Practice of Pharmacy*, 21<sup>st</sup> Ed. (Lippincott Williams & Wilkins, Baltimore, MD, 2005) and *Handbook of Pharmaceutical Excipients*, American Pharmaceutical Association, Washington, D.C., (*e.g.*, 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> Eds., 1986, 1994 and 2000, respectively). As will be known to those in the art, pharmaceutically acceptable excipients can provide a variety of functions and can be described as wetting agents, buffering agents, suspending agents, lubricating agents, emulsifiers, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, and sweeteners. Examples of pharmaceutically acceptable excipients include without limitation: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, cellulose acetate, hydroxypropylmethylcellulose, and hydroxypropylcellulose; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer’s solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters,

polycarbonates and/or polyanhydrides; and (22) other non-toxic compatible substances employed in pharmaceutical formulations.

“Unit dosage form” refers to a physically discrete unit suitable as a unitary dosage for a human or an animal. Each unit dosage form can contain a predetermined amount of a therapeutic agent calculated to produce a desired effect.

“Patient” refers to an animal, such as a mammal, including but not limited to, a human. Hence, the methods disclosed herein can be useful in human therapy and veterinary applications. In one embodiment, the patient is a mammal. In another embodiment, the patient is a human.

“Effective amount” or “therapeutically effective amount” refers to an amount of a compound or pharmaceutical composition which, based on its parameters of efficacy and potential for toxicity and the knowledge of one skilled in the art, produces a desired effect, such as treating or preventing a condition. An effective amount can be administered in for example one, two, three, four or more doses per day or per week.

“Treat” or “treating” refers to attain or attaining a beneficial or desired result, such as a clinical result. In some embodiments, the beneficial or desired result is any one or more of the following: inhibiting or suppressing the onset or development of a condition, reducing the severity of the condition, reducing the number or severity of symptoms associated with the condition, increasing the quality of life of a patient suffering from the condition, decreasing the dose of another medication required to treat the condition, enhancing the effect of another medication a patient is taking for the condition, and prolonging the survival of a patient having the condition. For example, in the case of major depressive disorder (MDD), treating may involve a clinically significant decline in at least one measurable marker or symptom of MDD. Measurable markers include electro-encephalogram (EEG) slow wave activity (SWA) and brain-derived neurotrophic factor (BDNF). The severity of MDD may be assessed, for example, using the Montgomery-Asberg Depression Rating Scale (MADRS), Hamilton Rating Scale for Depression (HAM-D or HRSD, such as HAM-D17, Beck Depression Inventory (BDI), VAS-depression, Hamilton Anxiety Rating Scale (HAM-A), Brief Psychiatric Rating Scale-positive symptoms (BPRS), the Clinician Administered Dissociative States Scale (CADSS), Young Mania Rating Scale (YMRS), Snaith Hamilton Pleasure Scale-Modified Scoring System (SHAPS-M), Wechsler Depression Rating Scale, Raskin Depression

Rating Scale, Inventory of Depressive Symptomatology (IDS), the Quick Inventory of Depressive Symptomatology (QIDS), or any other scale known in the art for rating MDD.

“Prevent” or “preventing” refers to reduce or reducing the probability of that a patient develops a condition which the patient does not have but is at risk of developing.

5 “At risk” denotes that a patient has one or more risk factors, which are measurable parameters that correlate with the development of a condition and are known in the art. A patient having one or more of risk factors has a higher probability of developing the condition than a patient without such risk factors.

“Adjunct” refers to the use of a compound or pharmaceutical composition in  
10 conjunction with at least one additional treatment. As an adjunct, the compound or pharmaceutical composition may improve the efficacy of the at least one additional treatment, such as by achieving a faster response to the at least one additional treatment, reducing the severity of a condition, reducing the number or severity of symptoms associated with a condition, or decreasing the dose of the at least one additional treatment.  
15 The adjunct may be administered with the at least one additional treatment together in a single composition or separately in individual compositions, at substantially the same time or at different times.

A condition that is “responsive to an NR2B antagonist” includes any condition in which administration of an N-methyl-D-aspartate receptor subunit 2B (NR2B) antagonist  
20 treats or prevents the condition, as those terms are defined herein. Hence, a condition whose symptoms are diminished upon administration of an NMDA NR2B antagonist is a condition responsive to an NR2B antagonist. Examples of a condition that is responsive to an NR2B antagonist include Parkinson's disease, neuropathic pain (such as postherpetic neuralgia, nerve injury, “dynias”, vulvodynia, phantom limb pain, root avulsions, painful  
25 diabetic neuropathy, painful traumatic mononeuropathy, painful polyneuropathy, central pain syndromes, and postsurgical pain syndromes, postmastectomy syndrome, postthoracotomy syndrome, stump pain), bone and joint pain (such as osteoarthritis), repetitive motion pain, dental pain, cancer pain, myofascial pain (muscular injury, fibromyalgia), perioperative pain (general surgery, gynecological), chronic pain,  
30 dysmenorrhea, as well as pain associated with angina, and inflammatory pain of varied origins (*e.g.* osteoarthritis, rheumatoid arthritis, rheumatic disease, teno-synovitis and gout), headache, migraine and cluster headache, schizophrenia, stroke, traumatic brain

injury, Alzheimer's disease, cerebral ischemia, amyotrophic lateral sclerosis, Huntington's disease, sensorineural hearing loss, tinnitus, glaucoma, neurological damage caused by epileptic seizures or by neurotoxin poisoning or by impairment of glucose and/or oxygen to the brain, vision loss caused by neurodegeneration of the visual pathway, Restless Leg  
5 Syndrome, multi-system atrophy, non-vascular headache, chronic (or chronic persistent), subchronic or acute cough, primary hyperalgesia, secondary hyperalgesia, primary allodynia, secondary allodynia, or other pain caused by central sensitization, dyskinesias (such as the side effects accompanying normal doses of L-Dopa), depressive disorders (such as, major depressive disorder (MDD) and treatment-resistant MDD), trauma- and  
10 stressor-related disorders (such as acute stress disorder and posttraumatic stress disorder (PTSD)), bipolar disorders with depressive features, anxiety disorders, and obsessive-compulsive and related disorders. *See Diagnostic and Statistical Manual of Mental Disorders, 5<sup>th</sup> Ed.* (American Psychiatric Association, Arlington, VA, 2013) for additional examples of depressive disorders, bipolar disorders, anxiety disorders, obsessive-  
15 compulsive and related disorders, and other conditions.

“Anti-hypertensive” refers to any agent that lowers blood pressure. Examples of an anti-hypertensive include calcium channel blockers, angiotensin converting enzyme inhibitors (ACE inhibitors), angiotensin II receptor antagonists (A-II antagonists), diuretics, beta-adrenergic receptor blockers ( $\beta$ -blockers), vasodilators and alpha-  
20 adrenergic receptor blockers ( $\alpha$ -blockers).

“Administered intermittently” refers to administration at irregular time periods throughout, for example, a 24 hour period, or a 7 day period, or on an as needed basis.

“Reacting” refers to combining or mixing two or more agents under appropriate conditions to produce the indicated or desired product. The indicated or desired product  
25 may not necessarily result directly from reacting the agents; reacting the agents may yield one or more intermediates that ultimately lead to the formation of the indicated or desired product.

“Recrystallization” refers to a purification process whereby a solid compound is dissolved in an appropriate solvent and recrystallized to provide a solid of higher purity.  
30 Types of recrystallization include single solvent, multi-solvent, hot filtration and seeding.

“Slurrying” refers to a purification process of suspending crystals of a compound in an appropriate solvent, stirring the suspension, and isolating the crystals.

"Such as" has the same meaning as "such as but not limited to." Similarly, "include" has the same meaning as "include but not limited to", while "including" has the same meaning as "including but not limited to".

"Between about X and about Y" refers to values between about X and about Y,  
5 including about X and about Y.

Other than in the working examples, or where otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth as used in the specification and claims are to be understood as being modified by the term "about". Accordingly, unless indicated to the contrary, such numbers are approximations that may  
10 vary depending upon the desired properties sought to be obtained by the disclosed subject matter. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should be construed in light of the number of significant digits and ordinary rounding techniques.


While the numerical ranges and parameters setting forth the broad scope of the  
15 disclosed subject matter are approximations, the numerical values set forth in the working examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

### **Compounds**

20 It has been discovered that Compound (I) can exist in multiple crystalline or polymorphic forms. The polymorphic forms of Compound (I) identified to date are: Form I, Form II and Form III. Form I is composed of acicular particles (rods) while Form II has plate-like morphology. Form III has been detected only in polymorphic mixtures. Neither is Compound (I)'s polymorphism nor any of its polymorphic forms disclosed in U.S. Pat.  
25 No. 7,592,360.

Form II, the thermodynamically most stable of the three polymorphs, is a white crystalline powder. This form is an anhydrate with a melting point of 123-124°C. Forms I and III have been shown to convert to Form II both thermally in the solid state as well as in aqueous suspensions.

30 One aspect of the disclosure provides a compound that is crystalline Form II of Compound (I)



(I)

exhibiting at least one of:

- 5 (i) an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, comprising peaks of 2-theta angles of about 5.9 and 8.8 degrees;
- (ii) an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, substantially as shown in FIG. 1A;
- (iii) an ultraviolet absorbance spectrum, obtained using methanol as diluent, substantially as shown in FIG. 2;
- 10 (iv) an infrared spectrum substantially as shown in FIG. 3;
- (v) a proton nuclear magnetic resonance spectrum at about 600 MHz in CD<sub>3</sub>CN substantially as shown in FIG. 4;
- (vi) a <sup>13</sup>C nuclear magnetic resonance spectrum at about 150 MHz in CD<sub>3</sub>CN substantially as shown in FIG. 5;
- 15 (vii) a thermogravimetric analysis curve substantially as shown in FIG. 6; and
- (viii) a differential scanning calorimetry thermogram substantially as shown in FIG. 7.

In one embodiment, the crystalline Form II of Compound (I) is substantially pure. "Substantially pure" refers to crystalline Form II of Compound (I) in isolated form that is  
20 at least about 90% by weight pure or free of impurities, including other polymorphic forms. In one embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 95% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 96% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about  
25 97% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 98% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 99% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I)

is at least about 99.5% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 99.6% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 99.7% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 99.8% by weight pure. In another embodiment, the substantially pure crystalline Form II of Compound (I) is at least about 99.9% by weight pure. Percent purity may be assessed by any method known in the art, such as gas chromatography (GC), column chromatography (CC), liquid chromatography (LC), high-pressure liquid chromatography (HPLC), thin layer chromatography (TLC), mass spectrometry (MS) and/or high-resolution mass spectrometry (HRMS).

In another embodiment, the compound exhibits an X-ray powder diffraction pattern comprising peaks of 2-theta angles of about 5.9 and 8.8 degrees that correspond, respectively, to d-spacing at about 14.9 and 10.0 Angstroms (Å).

In another embodiment, the compound exhibits an X-ray powder diffraction pattern substantially as shown in FIG. 1A. In another embodiment, the compound further exhibits at least one of: an ultraviolet absorbance spectrum substantially as shown in FIG. 2, an infrared spectrum substantially as shown in FIG. 3, a proton nuclear magnetic resonance spectrum substantially as shown in FIG. 4, a <sup>13</sup>C nuclear magnetic resonance spectrum substantially as shown in FIG. 5, a thermogravimetric analysis curve substantially as shown in FIG. 6, and a differential scanning calorimetry thermogram substantially as shown in FIG. 7.

In another embodiment, the compound exhibits an ultraviolet absorbance spectrum substantially as shown in FIG. 2. In another embodiment, the ultraviolet absorbance spectrum comprises an absorbance maximum at about  $236 \pm 2$  nm.

In another embodiment, the compound exhibits an infrared spectrum substantially as shown in FIG. 3.

In another embodiment, the compound exhibits a proton nuclear magnetic resonance spectrum substantially as shown in FIG. 4. In another embodiment, the proton nuclear magnetic resonance spectrum comprises peaks substantially as set out in Table 1.

In another embodiment, the compound exhibits a <sup>13</sup>C nuclear magnetic resonance spectrum comprising peaks substantially as shown in FIG. 5. In another embodiment, the

<sup>13</sup>C nuclear magnetic resonance spectrum comprises peaks substantially as set out in Table 2.

In another embodiment, the compound exhibits a thermogravimetric analysis curve substantially as shown in FIG. 6. In another embodiment, the thermogravimetric analysis curve corresponds to a weight loss of about 0.16% up to about 250° C.

In another embodiment, the compound exhibits a differential scanning calorimetry thermogram substantially as shown in FIG. 7. In another embodiment, the differential scanning calorimetry thermogram comprises an endothermic peak at a temperature of about 124° C.

Also provided herein is a crystalline Form I of Compound (I) as described in FIG. 1B. The purity of Form I can be at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95%.

### **Pharmaceutical Compositions**

Another aspect of the disclosure provides a pharmaceutical composition comprising an effective amount of crystalline Form II of Compound (I) as in any of the embodiments described above. In one embodiment, the crystalline Form II of Compound (I) is substantially pure as in any of the embodiments described above. Also provided herein is a pharmaceutical composition comprising an effective amount of crystalline Form I of Compound (I) as described herein.

Another aspect of the disclosure provides a pharmaceutical composition comprising an effective amount of Compound (I) in particulate form with an X90 particle size of about 10 µm or less. In one embodiment, the X90 particle size is about 8 µm or less. In another embodiment, the X90 particle size is about 6 µm or less. In another embodiment, the X90 particle size is about 5 µm or less. In another embodiment, the X90 particle size is between about 1 µm and about 10 µm. In another embodiment, the X90 particle size is between about 2 µm and about 8 µm. In another embodiment, the X90 particle size is between about 3 µm and about 6 µm. In another embodiment, the X90 particle size is between about 4 µm and about 5 µm. In another embodiment, the X90 particle size is about 4.5 µm. Other exemplary embodiments of the X90 particle size include about 4 µm, about 3.5 µm, about 3 µm, about 2.5 µm, about 2 µm, about 1.5 µm, about 1 µm, about 0.9 µm, about 0.8 µm, about 0.7 µm, about 0.6 µm, about 0.5 µm, about 0.4 µm, about 0.3 µm, about 0.2 µm, and about 0.1 µm. In another embodiment, the

Compound (I) is crystalline Form II. In another embodiment, the Compound (I) is substantially pure crystalline Form II. In another embodiment, the Compound (I) is crystalline Form I.

Another aspect of the disclosure provides a pharmaceutical composition  
5 comprising an effective amount of Compound (I) in particulate form with an X50 particle size of about 5  $\mu\text{m}$  or less. In one embodiment, the X50 particle size is about 4  $\mu\text{m}$  or less. In another embodiment, the X50 particle size is about 3  $\mu\text{m}$  or less. In another embodiment, the X50 particle size is about 3  $\mu\text{m}$  or less. In another embodiment, the X50 particle size is about 2  $\mu\text{m}$  or less. In another embodiment, the X50 particle size is  
10 between about 1  $\mu\text{m}$  and about 5  $\mu\text{m}$ . In another embodiment, the X50 particle size is between about 1  $\mu\text{m}$  and about 4  $\mu\text{m}$ . In another embodiment, the X50 particle size is between about 1  $\mu\text{m}$  and about 3  $\mu\text{m}$ . In another embodiment, the X50 particle size is between about 1  $\mu\text{m}$  and about 2  $\mu\text{m}$ . In another embodiment, the X50 particle size is about 2  $\mu\text{m}$ . In another embodiment, the X50 particle size is about 1.9  $\mu\text{m}$ .

Another aspect of the disclosure provides a pharmaceutical composition  
15 comprising an effective amount of Compound (I) in particulate form with an X10 particle size of about 2  $\mu\text{m}$  or less. In one embodiment, the X10 particle size is about 1  $\mu\text{m}$  or less. In another embodiment, the X10 particle size is between about 0.1  $\mu\text{m}$  and about 1  $\mu\text{m}$ . In another embodiment, the X10 particle size is between about 0.1  $\mu\text{m}$  and about 0.9  $\mu\text{m}$ . In  
20 another embodiment, the X10 particle size is between about 0.5  $\mu\text{m}$  and about 0.9  $\mu\text{m}$ . In another embodiment, the X10 particle size is between about 0.7  $\mu\text{m}$  and about 0.8  $\mu\text{m}$ . In another embodiment, the X10 particle size is about 0.8  $\mu\text{m}$ . In another embodiment, the X50 particle size is about 7.9  $\mu\text{m}$ .

In some embodiments of the pharmaceutical compositions disclosed herein, the  
25 pharmaceutical composition further comprises a pharmaceutically acceptable excipient. Examples of pharmaceutically acceptable excipients include those described above, such as carriers, surface active agents, thickening or emulsifying agents, solid binders, dispersion or suspension aids, solubilizers, colorants, flavoring agents, coatings, disintegrating agents, lubricants, sweeteners, preservatives, isotonic agents, and any  
30 combination thereof. The selection and use of pharmaceutically acceptable excipients is taught, *e.g.*, in *Remington: The Science and Practice of Pharmacy*, 21<sup>st</sup> Ed. (Lippincott Williams & Wilkins, Baltimore, MD, 2005).

In some embodiments of the pharmaceutical compositions disclosed herein, the pharmaceutical composition further comprises at least one additional active agent. Examples of such active agent include: (1) non-steroidal anti-inflammatory agents; (2) COX-2 inhibitors; (3) bradykinin B1 receptor antagonists; (4) sodium channel blockers and antagonists; (5) nitric oxide synthase (NOS) inhibitors; (6) glycine site antagonists; (7) potassium channel openers; (8) AMPA/kainate receptor antagonists; (9) calcium channel antagonists; (10) GABA-A receptor modulators (*e.g.*, a GABA-A receptor agonist); (11) matrix metalloprotease (MMP) inhibitors; (12) thrombolytic agents; (13) opioids such as morphine; (14) neutrophil inhibitory factor (NIF); (15) L-Dopa; (16) carbidopa; (17) levodopa/carbidopa; (18) dopamine agonists such as bromocriptine, pergolide, pramipexole, ropinirole; (19) anticholinergics; (20) amantadine; (21) carbidopa; (22) catechol O-methyltransferase (COMT) inhibitors such as entacapone and tolcapone; (23) Monoamine oxidase B (MAO-B) inhibitors; (24) opiate agonists or antagonists; (25) 5HT receptor agonists or antagonists; (26) NMDA receptor agonists or antagonists; (27) NK1 antagonists; (28) selective serotonin reuptake inhibitors (SSRI) and selective serotonin and norepinephrine reuptake inhibitors (SSNRI); (29) tricyclic antidepressant drugs; (30) norepinephrine modulators; (31) lithium; (32) valproate; (33) D-serine; (34) neurontin (gabapentin); (35) antitussives; (36) antihistamines (*e.g.*, first generation antihistamines); (37) decongestants; (38) expectorants; (39) mucolytics; (40) antipyretics; and (41) analgesics.

In some embodiments of the pharmaceutical compositions disclosed herein, the pharmaceutical composition further contains an anti-hypertensive agent. In some embodiments, the anti-hypertensive agent is an  $\alpha_1$ -adrenoreceptor antagonist or an  $\alpha_2$ -adrenoreceptor antagonist. Non-limiting examples of  $\alpha_1$ -adrenoreceptor antagonists include doxazosin, prazosin, terazosin, indoramin, metabolites thereof, and analogs thereof. Non-limiting examples of  $\alpha_2$ -adrenoreceptor antagonists include clonidine, guanabenz, guanoxabenz, metabolite thereof, and analogs thereof.

The pharmaceutical compositions can be prepared as any appropriate unit dosage form. For example, the pharmaceutical compositions can be formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, as drenches, tablets (such as those targeted for buccal, sublingual and systemic absorption, including over-encapsulation tablets), capsules (such as dry filled,

hard gelatin, soft gelatin or over-encapsulation capsules), caplets, boluses, powders, sachets, granules, pastes, mouth sprays, troches, lozenges, pellets, syrups, suspensions, elixirs, liquids, liposomes, emulsions and microemulsions; or (2) parenteral administration by, for example, subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension. Additionally, the pharmaceutical compositions can be formulated for immediate, sustained, extended, delayed or controlled release.

In one embodiment, the pharmaceutical composition is formulated for oral administration. In another embodiment, the pharmaceutical composition is in tablet or capsule form. In another embodiment, the pharmaceutical composition is in tablet form. In another embodiment, the pharmaceutical composition is in capsule form. In another embodiment, the tablet or capsule is formulated for immediate release. In another embodiment, the tablet or capsule is formulated for sustained, extended, delayed or controlled release.

Tablets can be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing in a suitable machine Compound (I) in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets can be optionally coated or scored and can be formulated so as to provide sustained, extended, delayed or controlled release of Compound (I). Methods of formulating such sustained, extended, delayed or controlled release compositions are known in the art and disclosed in issued U.S. patents, including but not limited to U.S. Patent Nos. 4,369,174, 4,842,866, and the references cited therein. Coatings can be used for delivery of compounds to the intestine (*see, e.g.*, U.S. Patent Nos. 6,638,534, 5,217,720, 6,569,457, and the references cited therein). In addition to tablets, other dosage forms, such as capsules, granulations and gel-caps, can be formulated to provide sustained, extended, delayed or controlled release of Compound (I).

In another embodiment, the pharmaceutical composition is formulated for parenteral administration. Examples of a pharmaceutical composition suitable for parenteral administration include aqueous sterile injection solutions and non-aqueous sterile injection solutions, each containing, for example, anti-oxidants, buffers,

bacteriostats and/or solutes that render the formulation isotonic with the blood of the intended recipient; and aqueous sterile suspensions and non-aqueous sterile suspensions, each containing, for example, suspending agents and/or thickening agents. The formulations can be presented in unit-dose or multi-dose containers, for example, sealed ampules or vials, and can be stored in a freeze dried (lyophilized) condition requiring only the addition of a sterile liquid carrier, such as water, immediately prior to use. In one embodiment, the pharmaceutical composition is formulated for intravenous administration.

In another embodiment, the effective amount of the Compound (I) or crystalline Form (I) or (II) of Compound I is between about 1 mg and about 200 mg. In another embodiment, the effective amount is between about 4 mg and about 16 mg. In another embodiment, the effective amount is between about 4 mg and about 12 mg. In another embodiment, the effective amount is between about 4 mg and about 8 mg. In another embodiment, the effective amount is between about 8 mg and about 20 mg. In another embodiment, the effective amount is between about 12 mg and about 20 mg. In another embodiment, the effective amount is between about 16 mg and about 20 mg. In another embodiment, the effective amount is between about 8 mg and about 20 mg. In another embodiment, the effective amount is between about 8 mg and about 16 mg. In another embodiment, the effective amount is between about 8 mg and about 12 mg. In another embodiment, the effective amount is between about 12 mg and about 20 mg. In another embodiment, the effective amount is between about 12 mg and about 16 mg. In another embodiment, the effective amount is about 4 mg. In another embodiment, the effective amount is about 8 mg. In another embodiment, the effective amount is about 12 mg. In another embodiment, the effective amount is about 16 mg. In another embodiment, the effective amount is about 20 mg.

In connection with the embodiments described herein, there is also provided a kit containing the above disclosed Compound (I) and a secondary agent. In an exemplary embodiment, the kit contains Compound (I) and one or more secondary agent selected from a selective serotonin reuptake inhibitor (SSRI), a selective serotonin and norepinephrine reuptake inhibitor (SSNRI), a tricyclic antidepressant drug, a norepinephrine modulator, an antitussive, an antihistamine, a decongestant, an expectorants, a mucolytics, a antipyretics, an analgesics, and an anti-hypertensive agent.

In some embodiments, two or more ingredients (Compound (I) and secondary agents) may be administered simultaneously.

In some embodiments, the kit may include an anti-hypertensive agent as a secondary. In some embodiments, the anti-hypertensive agent is an  $\alpha_1$ -adrenoreceptor blocker. Non-limiting examples of  $\alpha_1$ -adrenoreceptor antagonists include doxazosin, prazosin, terazosin, indoramin, metabolites thereof, pharmaceutically acceptable salts thereof, and analogs thereof.

In some embodiments, the agents can be administered sequentially. The specific administration mode and dosing schedule can be determined by one of ordinary skill in the art (e.g. a practicing physician) without undue experimentation.

### Methods of Use

Another aspect of the disclosure provides a method of treating or preventing a condition responsive to an NR2B antagonist, comprising administering to a patient in need thereof an effective amount of crystalline Form I or II of Compound (I) or a pharmaceutical composition comprising crystalline Form I or II of Compound (I). In one embodiment, the method is of treating a condition responsive to an NR2B antagonist. In another embodiment, the crystalline Form II is substantially pure as in any of the embodiments described above.

In another embodiment, the condition responsive to an NR2B antagonist is a depressive disorder. In another embodiment, the condition is major depressive disorder (MDD). In another embodiment, the condition is treatment-resistant major depressive disorder (TRMDD). In another embodiment, the condition is bipolar disorder with depressive feature. In another embodiment, the condition is anxiety disorder. In another embodiment, the condition is posttraumatic stress disorder (PTSD). In another embodiment, the condition is a depressive disorder, MDD or TRMDD with suicidal ideation.

In another embodiment, the compound or pharmaceutical composition is administered with food. In another embodiment, the compound or pharmaceutical composition is administered without food.

Also provided is a method of targeting N-methyl-D-aspartate (NMDA) receptor subunit 2B (GluN2B) expressed on a cell by administering to a subject in need thereof a therapeutically effective amount of Form I or Form II of compound I, allowing sufficient

amount of time for the compound to bind to GluN2B. As explained above, GluN2B plays a key role in various diseases including for example schizophrenia, stroke, traumatic brain injury, Alzheimer's disease, cerebral ischemia, amyotrophic lateral sclerosis, Huntington's disease, sensorineural hearing loss, tinnitus, glaucoma, neurological damage caused by epileptic seizures or by neurotoxin poisoning or by impairment of glucose and/or oxygen to the brain, vision loss caused by neurodegeneration of the visual pathway, Restless Leg Syndrome, multi-system atrophy, non-vascular headache, chronic (or chronic persistent), subchronic or acute cough, primary hyperalgesia, secondary hyperalgesia, primary allodynia, secondary allodynia, or other pain caused by central sensitization, dyskinesias (such as the side effects accompanying normal doses of L-Dopa), depressive disorders (such as, major depressive disorder (MDD) and treatment-resistant MDD), trauma- and stressor-related disorders (such as acute stress disorder and posttraumatic stress disorder (PTSD)), bipolar disorders with depressive features, anxiety disorders, and obsessive-compulsive and related disorders. By antagonizing GluN2B with the compound I (Form I or II), the present invention provides a novel approach in treating GluN2B-associated diseases.

Another aspect of the disclosure provides a method of treating or preventing a condition responsive to an NR2B antagonist, comprising administering to a patient in need thereof an effective amount of Compound (I) in particulate form (Form I or Form II) with an X90, X50 or X10 particle size as in any of the embodiments described above. In one embodiment, the method is of treating a condition responsive to an NR2B antagonist. In another embodiment, the Compound (I) is crystalline Form II as in any of the embodiments described above. In another embodiment, the crystalline Form II is substantially pure as in any of the embodiments described above. In another embodiment, the condition responsive to an NR2B antagonist is a depressive disorder. In another embodiment, the condition is major depressive disorder (MDD). In another embodiment, the condition is treatment-resistant major depressive disorder (TRMDD). In another embodiment, the condition is a depressive disorder, MDD or TRMDD with suicidal ideation. In one embodiment, the compound or pharmaceutical composition is administered with food. In another embodiment, the compound or pharmaceutical composition is administered without food.

Another aspect of the disclosure provides a method of treating or preventing suicidal ideation, comprising administering an effective amount of Compound (I) (Form I or Form II) to a patient who has, is suspected of having, or has been diagnosed with having suicidal ideation. In one embodiment, the method is of treating suicidal ideation.

5 In another embodiment, the Compound (I) is in particulate form with an X90, X50 or X10 particle size as in any of the embodiments described above. In another embodiment, the Compound (I) is crystalline Form II as in any of the embodiments described above. In another embodiment, the crystalline Form II is substantially pure as in any of the embodiments described above. In another embodiment, the Compound (I) is crystalline  
10 Form I as in any of the embodiments described above.

In another embodiment, the patient has been diagnosed with having suicidal ideation within about 4 weeks prior to administration of the compound. In another embodiment, the patient has been further diagnosed with having a depressive disorder. In another embodiment, the patient has been further diagnosed with having major depressive  
15 disorder (MDD). In another embodiment, the patient has been further diagnosed with having treatment-resistant major depressive disorder (TRMDD). In one embodiment, the compound or pharmaceutical composition is administered with food. In another embodiment, the compound or pharmaceutical composition is administered without food.

Another aspect of the disclosure provides a method of reducing absorption rate of  
20 Compound (I) comprising administering to a patient in need thereof an effective amount of the Compound (I) (Form I or Form II) with food, wherein the Compound (I) is administered either substantially concurrently with, or up to about 2 hours after, or up to about 30 minutes before administration of food. In one embodiment, the Compound (I) is in particulate form with an X90, X50 or X10 particle size as in any of the embodiments  
25 described above. In another embodiment, the Compound (I) is crystalline Form II as in any of the embodiments described above. In another embodiment, the crystalline Form II is substantially pure as in any of the embodiments described above.

In another embodiment, the method results in a lower  $C_{max}$  or a higher  $T_{max}$  compared to that of a method comprising administering the Compound (I) without food.  
30 In another embodiment, the Compound (I) is administered either substantially concurrently with, or up to about 90 minutes after, or up to about 15 minutes before administration of food. In another embodiment, the Compound (I) is administered either

substantially concurrently with, or up to about 60 minutes after, or up to about 10 minutes before administration of food. In another embodiment, the Compound (I) is administered substantially concurrently with administration of food.

In any of the methods provided herein, administration of the compound or pharmaceutical composition may be via any accepted mode known in the art, such as orally or parenterally. The term "parenterally" includes without limitation subcutaneously, intravenously, intramuscularly, intraperitoneally, intrathecally, intraventricularly, intrasternally, intracranially, by intraosseous injection and by infusion techniques. In one embodiment, the compound or pharmaceutical composition is administered orally. In another embodiment, the compound or pharmaceutical composition is administered parenterally. In another embodiment, the compound or pharmaceutical composition is administered intravenously.

In any of the methods provided herein, the method may further comprise monitoring the patient's blood pressure; and if hypertension is detected, administering an anti-hypertensive to the patient.

In any of the methods provided herein, the compound or pharmaceutical composition may be used in combination with at least one additional active agent as disclosed above. The compound or pharmaceutical composition may be administered with the at least one additional active agent either together in a single composition or separately in individual compositions at substantially the same time or at different times (*e.g.*, sequentially). Examples of such active agent include: (1) non-steroidal anti-inflammatory agents; (2) COX-2 inhibitors; (3) bradykinin B1 receptor antagonists; (4) sodium channel blockers and antagonists; (5) nitric oxide synthase (NOS) inhibitors; (6) glycine site antagonists; (7) potassium channel openers; (8) AMPA/kainate receptor antagonists; (9) calcium channel antagonists; (10) GABA-A receptor modulators (*e.g.*, a GABA-A receptor agonist); (11) matrix metalloprotease (MMP) inhibitors; (12) thrombolytic agents; (13) opioids such as morphine; (14) neutrophil inhibitory factor (NIF); (15) L-Dopa; (16) carbidopa; (17) levodopa/carbidopa; (18) dopamine agonists such as bromocriptine, pergolide, pramipexole, ropinirole; (19) anticholinergics; (20) amantadine; (21) carbidopa; (22) catechol O-methyltransferase (COMT) inhibitors such as entacapone and tolcapone; (23) Monoamine oxidase B (MAO-B) inhibitors; (24) opiate agonists or antagonists; (25) 5HT receptor agonists or antagonists; (26) NMDA receptor agonists or antagonists; (27)

NK1 antagonists; (28) selective serotonin reuptake inhibitors (SSRI) and selective serotonin and norepinephrine reuptake inhibitors (SSNRI); (29) tricyclic antidepressant drugs, (30) norepinephrine modulators; (31) lithium; (32) valproate; (33) D-serine; and (34) neurontin (gabapentin); (35) antitussives; (36) antihistamines (*e.g.*, first generation antihistamines); (37) decongestants; (38) expectorants; (39) mucolytics; (40) antipyretics; and (41) analgesics. In some embodiments, the compound or pharmaceutical composition is administered with at least one additional active agent selected from antitussives, first generation antihistamines, decongestants, expectorants, mucolytics, antipyretics, analgesics.

10 In some embodiments, the compound or pharmaceutical composition is administered as an adjunct to a selective serotonin reuptake inhibitor (SSRI) or selective serotonin and norepinephrine reuptake inhibitor (SSNRI). Examples of SSRI and SSNRI include binedaline, m-chloropiperazine, citalopram, duloxetine, etoperidone, femoxetine, fluoxetine, fluvoxamine, indalpine, indeloxazine, milnacipran, nefazodone, oxaflazone, paroxetine, prolintane, ritanserin, sertraline, tandospirone, venlafaxine and zimeldine. In 15 another embodiment, the method improves the efficacy of the SSRI or SSNRI in the treatment or prevention of a condition responsive to an NR2B antagonist. In another embodiment, the method reduces the time it takes for a patient to respond to treatment with an SSRI or SSNRI. In another embodiment, the method reduces the dose of SSRI or SSNRI that a patient would otherwise need. In another embodiment, the method reduces 20 the severity of a condition or the number or severity of symptoms associated with a condition. In another embodiment, the condition is a depressive disorder. In another embodiment, the condition is major depressive disorder (MDD). In another embodiment, the condition is treatment-resistant major depressive disorder (TRMDD). In another embodiment, the condition is a depressive disorder, MDD or TRMDD with suicidal 25 ideation. The adjunct may be administered with the at least one additional treatment either together in a single composition or separately in individual compositions at substantially the same time or at different times.

In any of the methods provided herein, Compound (I) (Form I or Form II) may be 30 administered in conjunction with an anti-hypertensive agent. The use of Compound (I) may be associated with an increase in blood pressure that is likely due to increased  $\alpha_1$ -adrenergic tone in peripheral arterioles mediated by circulating norepinephrine. The

increase on blood pressure is mostly transient as it relates to arteriolar vasoconstriction, in contrast to other frequent causes of hypertension such as cardiac stimulation, arterial wall hypertrophy, salt retention and fluid overload.

Accordingly, in some embodiments, the anti-hypertensive agent is an  $\alpha_1$ -  
5  $\alpha_1$ -adrenoreceptor, that blocks peripheral  $\alpha_1$ -adrenergic receptors. Non-limiting examples of  $\alpha_1$ -adrenoreceptor antagonists include Doxazosin, Prazosin, Terazosin, indoramin, metabolites thereof, pharmaceutically acceptable salts thereof, and analogs thereof. These agents are highly selective for  $\alpha_1$ -adrenoreceptor subtypes ( $\alpha_1A$ ,  $\alpha_1B$ ,  $\alpha_1D$ ). When given in large doses, they do not inhibit the  $\alpha_2$ -adrenoreceptors ( $\alpha_2A$ ,  $\alpha_2B$ ,  
10  $\alpha_2C$ ), the  $\beta$ -adrenoreceptors ( $\beta_1$ ,  $\beta_2$ ,  $\beta_3$ ), or other receptors such as acetylcholine (muscarinic), dopamine, and 5-hydroxytryptamine (5-HT, serotonin) receptors. In some embodiments, the anti-hypertensive agent is an  $\alpha_2$ -adrenoreceptor antagonist, that reduces central adrenergic outflow. Non-limiting examples of  $\alpha_2$ -adrenoreceptor antagonists include clonidine, guanabenz, guanoxabenz, metabolite  
15 thereof, and analogs thereof. Compound (I) and the anti-hypertensive agent may be administered simultaneously or separately. These agents stimulate  $\alpha_2$ -adrenoreceptors and inhibit brainstem vasomotor center-mediated norepinephrine release.

Other suitable antihypertensives include calcium channel blockers, diuretics and angiotensin converting enzyme (ACE) inhibitors. Non-limiting examples of calcium  
20 channel blockers include diltiazem, amlodipine, verapamil, diltiazem, nifedipine, amlodipine, felodipine, isradipine, nicardipine, cinnarizine, nisoldipine, and pharmaceutically acceptable salts thereof. Non-limiting examples of diuretics include thiazide diuretics, potassium-sparing diuretics, loop diuretics, and pharmaceutically acceptable salts thereof. Non-limiting examples of angiotensin converting enzyme (ACE)  
25 inhibitors include alacepril, benazepril, benazeprilat, captopril, ceronapril, cilazapril, delapril, enalapril, enalaprilat, fosinopril, imidapril, lisinopril, moexipril, moveltopril, perindopril, quinapril, quinaprilat, ramipril, ramiprilat, spirapril, temocapril,trandolapril, zofenopril, and pharmaceutically acceptable salts thereof.

The antihypertensives described herein may be used in any suitable forms.  
30 Examples include doxazosin mesylate, prazosin hydrochloride, and terazosin hydrochloride.

Because the transient increase in blood pressure relates to the increased  $\alpha$ 1-adrenergic tone in peripheral arterioles mediated by circulating norepinephrine, blocking the  $\alpha$ 1-adrenoceptor or  $\alpha$ 2-adrenoceptor is able effectively to mitigate the blood pressure effect. In some embodiments, the active ingredient of the anti-hypertensive agent consists essentially of an  $\alpha$ 1-adrenoceptor antagonist, an  $\alpha$ 2-adrenoceptor antagonist, or both. In some embodiments, the active ingredient of the anti-hypertensive agent consists essentially of an  $\alpha$ 1-adrenoceptor antagonist. The above described  $\alpha$ 1-adrenoceptor antagonist or  $\alpha$ 2-adrenoceptor antagonist can be administered before, during, or after the administration of Compound (I). In exemplary embodiments, the anti-hypertensive agent is administered 1, 2, 5, 10, 15, 20, 30, 60, 100 minutes prior to the administration of Compound (I). In other exemplary embodiments, the anti-hypertensive agent is administered 1, 2, 5, 10, 15, 20, 30, 60, 100 minutes after the administration of Compound (I). In some embodiments, the antihypertensive agent is administered at about the same time or sequentially with the administration of Compound (I). In some embodiments, the  $\alpha$ 1-adrenoceptor antagonist or  $\alpha$ 2-adrenoceptor antagonist is administered only in the above described timeframe to control the transient blood pressure effect from Compound (I). In some embodiments, the  $\alpha$ 1-adrenoceptor antagonist or  $\alpha$ 2-adrenoceptor antagonist is administered within 5, 10, 20, 30, or 60 minutes of the administration of Compound (I).

The effective amount or unit dosage of the hypertensive agent may range from about of between 0.001 to 100 mg/kg. of body weight. In some exemplary embodiments, the effective amount will be about 0.5 mg to 2500 mg, in yet some other embodiments about 5 mg to 50 mg.

In any of the methods provided herein, the compound or pharmaceutical composition may be administered intermittently. In some embodiments, the compound, secondary agent, combination of agents, or pharmaceutical composition may be independently administered once, twice, three times, four times or more per day. In some embodiments, the compound, secondary agent, combination of agents, or pharmaceutical composition may be independently administered once every 1, 2, 3, 4, 5, 6, 7, 10, 14, 15, 10 or 30 days.

In any of the methods provided herein, the effective amount or unit dosage of Compound (I) may be between about 1 mg and about 2000 mg daily, weekly or intermittently. For daily administration, the effective amount may be divided into two,

three or more doses for multiple administrations a day. In one embodiment, the effective amount is between about 1 mg/day and about 60 mg/day. In another embodiment, the effective amount is between about 4 mg/day and about 20 mg/day. In another embodiment, the effective amount is between about 4 mg/day and about 8 mg/day. In another embodiment, the effective amount is about 4 mg/day. In another embodiment, the effective amount is about 8 mg/day. In another embodiment, the effective amount is about 16 mg/day. In another embodiment, the effective amount is about 20 mg/day. In another embodiment, the effective amount is between about 4 mg and about 60 mg intermittently. In another embodiment, the effective amount is between about 4 mg and about 60 mg intermittently throughout a 24 hour period. In another embodiment, the effective amount is between about 4 mg/day and about 20 mg/day intermittently throughout a 24 hour period. In another embodiment, the effective amount is between about 4 mg/day and about 8 mg/day intermittently throughout a 24 hour period.

Compound (I) may be administered intermittently in any of the methods of the present invention described herein. As exemplified in FIG. 13, Compound (I) demonstrated marketed effects in immobility, climbing and swimming 24 hours after administration. Accordingly, Compound (I) can be administered intermittently to achieve sustained therapeutic effects. In some embodiments, the therapeutically effective amount or unit dosage of Compound (I) is between about 4 mg and about 2000 mg intermittently throughout 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 day period. Examples of intermittent administration include the administration of a therapeutically effect amount or a unit dosage of about 1, 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 mg of Compound (I) once every 1, 2, 3, 4, 5, 7, 10, 14, 15, 20, or 30 day period. In some embodiments, Compound (I) exists substantially in Form II. In further examples, such effective amount or unit dosage is administered once every day, once every 7 days, once every two weeks, once every 3 weeks, or once every month. In another embodiment, the effective amount is between about 8 mg and about 200 mg intermittently throughout a 7 day period. In another embodiment, the effective amount is between about 12 mg and about 60 mg intermittently throughout a 7 day period. In another embodiment, the effective amount is between about 16 mg and about 60 mg intermittently throughout a 7

day period. In another embodiment, the effective amount is between about 20 mg and about 60 mg intermittently throughout a 7 day period.

There may be an interval of between about 1 and 30 days between days of administration of Compound (I). Depending on the disease condition, treatment response  
5 and phase of treatment, the length of the interval may vary accordingly. For example, each interval may independently be 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 day long. In an exemplary embodiment, an initial administration of the compound on the first day may be followed by a 2-day interval before a second administration on the third day, which is then followed by a 6-day interval before a third administration.

10 In some embodiments of the methods disclosed herein, the compound or pharmaceutical composition is administered at a weight base dose. Therefore, in some embodiments, the effective amount is about 0.01 to about 1 mg/kg daily or intermittently.

The dose level can be adjusted for intravenous administration. In such case, the compound or pharmaceutical composition can be administered in an amount of between  
15 about 0.01  $\mu\text{g}/\text{kg}/\text{min}$  to about 100  $\mu\text{g}/\text{kg}/\text{min}$ .

In some embodiments, the administration regimen comprises a first initiation phase and a second maintenance phase. The dose in the first phase serves to reach a desirable therapeutic level whereas the dose of the second phase provides a sustained therapeutic effect. The daily dose of the first initiation phase may be more, less than or the same as  
20 that of the maintenance phase. The effective amount of Compound (I) for each phase may be, for example, between about 4 mg and about 60 mg daily or intermittently.

Independently during each phase, the compound or pharmaceutical composition can be administered 1, 2, 3, 4 or more times on the day of administration. The length of the initiation phase may be any suitable period of time from a day to a week or a month,  
25 including for example 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 15 days. The maintenance phase may last from days to years or as long as needed. In exemplary embodiments, the intermittent dosing schedule for initiation phase and maintenance phase independently includes administration of the compound every two days, every three days, every four days, every five days, every six days, every week, every 10 days, every two weeks, or every month.  
30 The effective amount for the initiation phase and the maintenance phase is independently selected from the range or value describe above for daily or intermittent administration.

With each of the above phases, additional adjustment in the dose and dosing schedule can be made. The adjustment will depend on the specific circumstances (for example, the presence or absence of a predisposition to the disease or condition being treated, the severity or expected severity of the disease, or the age or general health of the patient), even doses outside the aforementioned ranges may be in order. For example, the initiation phase may start with a higher daily dose than the dose for rest of the administration regimen. The administration may also include a toleration phase, where the compound is administered to a subject at a dose and for a period sufficient to allow the subject / patient to tolerate the dose without showing any adverse effects. The dose can then be increased at selected intervals of time until a therapeutic dose is achieved. The particular dose given the specific circumstances can be determined by a physician or other health-care professional of ordinary skill.

These compositions of the present invention may contain immediate release, sustained or extended release, delayed release components, or combinations thereof. The preparation of pharmaceutical or pharmacological compositions are known to those of skill in the art in light of the present disclosure. General techniques for formulation and administration are found in "Remington: The Science and Practice of Pharmacy, Twentieth Edition," Lippincott Williams & Wilkins, Philadelphia, Pa. Tablets, capsules, pills, powders, granules, dragées, gels, slurries, ointments, solutions suppositories, injections, inhalants and aerosols are examples of such formulations. For instance, , extended or modified release oral formulation can be prepared using methods known in the art. An extended release form of the pharmaceutical composition may be a matrix tablet or capsule composition.

Another aspect of the disclosure provides a method of increasing bioavailability of Compound (I), comprising administering to a patient in need thereof an effective amount of Compound (I), or a pharmaceutical composition comprising Compound (I), wherein the Compound (I) is in particulate form with an X90, X50 or X10 particle size as in any of the embodiments described above. In one embodiment, the Compound (I) is crystalline Form II as in any of the embodiments described above. In another embodiment, the crystalline Form II is substantially pure as in any of the embodiments described above. In another embodiment, the Compound (I) is crystalline Form I as in any of the embodiments described above.


In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having an X90 particle size of about 10  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering  
5 Compound (I) having an X90 particle size of about 11  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having an X90 particle size of about 12  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having an X90  
10 particle size of about 13  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having an X90 particle size of about 14  $\mu\text{m}$  or higher.

In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having a D43  
15 particle size of about 10  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having a D43 particle size of about 11  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having an D43 particle size of about 12  
20  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having a D43 particle size of about 13  $\mu\text{m}$  or higher. In another embodiment, the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) having a D43 particle size of about 14  $\mu\text{m}$  or higher.

25 In one embodiment, the compound or pharmaceutical composition is administered with food. In another embodiment, the compound or pharmaceutical composition is administered without food.

### **Methods of Manufacture**


Another aspect of the invention provides a method for the conversion from  
30 compound A to compound B under transition metal catalyzed hydrogenation conditions:



Wherein R' is

- 1). OH;
- 2). NH<sub>2</sub>; or
- 5     3). NHHet, wherein Het is a 5 or 6 membered heteroaromatic ring containing 1 or 2 nitrogen ring atoms, thiazolyl, or thiadiazolyl, the NH is linked to a carbon ortho to a nitrogen on the Het ring, and Het is optionally substituted with 1 or 2 substituents, each substituent independently is C1-4alkyl, fluoro, chloro, bromo, or iodo.

Examples of compound A include:



10

Besides Bn (Benzyl) and Bz (Benzoyl), other protecting groups may also be used to prepare analogs of compound A.

Various transition metals, including for example palladium and ruthenium, can be used as the catalyst. The catalyst can be prepared from transition precursor complexes and suitable ligands. For example, ruthenium precursor complexes include [(COD)]RhCl]<sub>2</sub> and similar complexes employing cationic OTf<sup>-</sup> and BF<sub>4</sub><sup>-</sup> salts. Suitable ligands include, for example, Josiphos and Walphos:

15




In some embodiments, compound A may exist in an acid form such as HCl salt, toluenesulfonic acid (TSA) salt, and methanesulfonic acid (MSA) salt. Various solvents,

20


including co-solvents of two or more solvents may be used in the hydrogenation reaction. The exact condition, such as the ratio of the catalyst to the substrate and the amount of the ligand, can be determined by one of ordinary skill in the art without undue experimentation.

- 5 Novel methods have been developed to prepare Compound (I) and crystalline Form I or II of Compound (I) in high purity on a large scale. The methods provided herein differ from the methods disclosed in U.S. Pat. No. 7,592,360 and PCT International Publication No. WO 2006/069287 in many respects, including the type and amount of reactants, intermediates, reaction conditions, and inclusion or omission of specific steps.
- 10 For example, neither of the methods disclosed in the '360 patent and the '287 PCT publication includes steps for converting Form I of Compound (I) into Form II. Furthermore, whereas the methods provided herein use intermediate Compound (8)



(8)


- 15 the method disclosed in the '360 patent uses the intermediate (±)-((cis)-3-fluoro-1-{[(4-methylbenzyl)oxy]carbonyl}piperidin-4-yl)acetic acid



Additionally, the methods provided herein use different reactants, amounts of reactants and reaction conditions than those recited in the '287 PCT publication.

- 20 One aspect of the disclosure provides a method of preparing Compound (I) comprising:


- (i) reacting Compound (8)



(8)

with triflic anhydride to yield a triflate;

(ii) reacting the triflate with ammonia to yield Compound (9)



(9)

5


; and

(iii) reacting the Compound (9) with 2-chloropyrimidine to yield Form I of Compound (I).

In one embodiment, the method further comprises seeding Form I of Compound (II) with Form II of Compound (I).


10

In another embodiment, the method further comprises reacting Compound (6)



(6)

with carbonyldiimidazole and Compound (7)




(7)

15

to yield Compound (8).


In another embodiment, the method further comprises debenzylating Compound (5)



(5)

with hydrogen over palladium to yield Compound (6).

In another embodiment, the method further comprises reducing Compound (4)



(4)

with chloro(1,5-cyclooctadiene)rhodium (I) dimer under hydrogen atmosphere to yield Compound (5).

In another embodiment, the method further comprises purifying the Compound (I).

10 In another embodiment, the purifying comprises slurring or recrystallization. In another embodiment, the purifying comprises slurring followed by recrystallization. In some embodiments, Form I of Compound (I) is purified by recrystallization in a suitable solvent such as ethyl acetate or a co-solvent including ethyl acetate and heptane.

15 Form I of Compound (I) can be converted to Form II in different ways. In some embodiments, Form I of Compound (I) in solid state is heated to a temperature above room temperature for a desirable period of time. Exemplary temperatures for promoting such conversion includes about 50 °C, 60 °C, 70 °C, 80 °C, 90 °C, 100 °C, 105 °C, 110 °C, 115 °C, 120 °C, 125 °C, 130 °C, 135 °C, 140 °C, and 150 °C. In some embodiments, the temperature ranges from about 90-135 °C, all subranges included. In some more  
20 exemplary embodiments, the temperature is about 100 - 105 °C, 100-110 °C, or 100 - 115 °C. The heating can be maintained for a period, for example from about 1 minute to 30 hours, all subranges included. Exemplary length of heating include about 10 minutes, 20

minutes, 30 minutes, 40 minutes, 50 minutes, 1 hour, 2 hours, 3 hours, 5 hours, 10 hours and 24 hours.

Form II of Compound (I) may also be obtained by the slurring of Form I in a suitable solvent or co-solvents. Stirring or heating may be applied. In an exemplary embodiment, Form I of Compound (I) can be suspended in water at room temperature for over 24 hours. If necessary, the time can be further extended to allow the complete conversion of Form I to Form II.

Subsequent to the conversion of Form I of Compound (I), Form II may be isolated using routine steps. In some embodiments, the isolation includes filtration and / or washing with a suitable solvent.

Large scale production of Form II can be achieved by seeding Form I with Form II. Seeding is an important step to induce the growth of crystal material from a small crystal of the target form. A system of a single solvent or multiple solvents may be used for seeding and crystal growth. In some embodiments, the system contains the co-solvent of methanol and acetic acid.


## EXAMPLES


The following examples are presented for illustrative purposes and should not serve to limit the scope of the disclosed subject matter.

### EXAMPLE 1: Preparation of Compound (I) and Crystalline Form II

20

#### Scheme I





Abbreviations used herein denote the following:

5 ACN = acetonitrile

AcOH = acetic acid

BnBr = benzyl bromide

CDI = 1,1'-carbonyldiimidazole

Darco G = activated charcoal

5 DCM = dichloromethane

DIPA = diisopropylamine

DMF = dimethylformamide

EtOAc = ethyl acetate

HCl = hydrogen chloride

10 iProAc = isopropyl acetate

MeOH = methanol

NaBH<sub>4</sub> = sodium borohydride

NaCl = sodium chloride

NaOH = sodium hydroxide

15 NH<sub>3</sub> = ammonia

Pd/C = palladium on carbon

[RhCODCl]<sub>2</sub> = chloro(1,5-cyclooctadiene)rhodium (I) dimer (also known as 1,5-cyclooctadienerhodium (I) chloride dimer, bis(1,5-cyclooctadiene)dirhodium (I) dichloride, di-μ-chlorobis[(1,2,5,6-η)-1,5-cyclooctadiene]dirhodium, rhodium (I) chloride 1,5-cyclooctadiene complex dimer, and [Rh(COD)Cl]<sub>2</sub>)

20

RT = room temperature

THF = tetrahydrofuran

### **Step 1**

Compound 1 was treated with LDA and then reacted with DMF to produce an *in situ* aldehyde intermediate, which was reduced with NaBH<sub>4</sub> to yield Compound 2.

25

Purification of compound 2 was completed by recrystallization.

### **Step 2**

Compound 2 was reacted with benzyl bromide to yield Compound 3.

### **Step 3**

Compound 3 was reduced with NaBH<sub>4</sub> to produce Compound 4 as the HCl salt.

30

Purification of Compound 4 was completed by recrystallization.

### **Step 4**

An asymmetric reduction of compound 4 using Walphos under hydrogen atmosphere produced Compound 5.

#### **Step 5**

Compound 5 was then deprotected (debenzylated) with a further reduction step  
5 using hydrogen over palladium to yield Compound 6.

#### **Step 6**

Compound 6 was coupled with 4-methylbenzylalcohol (compound 7) and carbonyldiimidazole to produce Compound 8.

#### **Step 7**

10 Compound 8 was then converted to an *in-situ* triflate, which was further reacted with ammonia to yield Compound 9.

#### **Step 8**

Compound 9 was reacted with chloropyrimidine to produce crude Form I of Compound (I) or 4-methylbenzyl (3S, 4R)-3-fluoro-4-[(pyrimidin-2-ylamino) methyl]  
15 piperidine-1-carboxylate. Form I of Compound (I) was seeded with Form II of Compound (I). Purification by slurring in the presence of Darco G60 followed by recrystallization produced purified Form II, that was co-milled to the desired particle size distribution.

#### **EXAMPLE 2: X-Ray Powder Diffraction Pattern**

Crystalline Form II of Compound (I) was characterized using X-ray powder  
20 diffraction (XRPD)(FIG. 1A). The XRPD pattern corresponds to an anhydrous form denoted as Form II, which is highly crystalline. In addition to Form II, Compound (I) can exist as another anhydrous polymorph, denoted as Form I. The XRPD patterns of these phases are shown in FIG. 1B. A slurry experiment was performed at room temperature by adding approximately equal proportions of the two anhydrous phases to methyl-*tert*-butyl  
25 ether. The XRPD of the filtered solids from the slurry showed phase conversion to Form II. XRPD analysis of solids recovered from heating Form I to 115°C showed phase conversion to Form II. Based on these experiments, Form II is the thermodynamically stable form between room temperature and 121°C.

#### **EXAMPLE 3: Ultraviolet Absorbance Spectrum**


30 The ultraviolet (UV) absorbance spectrum of crystalline Form II of Compound (I) shown in FIG. 2 was obtained using a Cary 300 Bio UV-Vis spectrophotometer and methanol as diluent. The spectrum is characterized by maxima at  $236 \pm 2$  nm.

**EXAMPLE 4: Infrared Transmittance Spectrum**

The infrared (IR) transmittance spectrum of crystalline Form II of Compound (I), shown in FIG. 3, was obtained with the ATR accessory using a Nicolet Nexus 670 FTIR spectrometer.

5 **EXAMPLE 5: Proton Nuclear Magnetic Resonance Spectrum**

The proton nuclear magnetic resonance ( $^1\text{H}$  NMR) spectrum of crystalline Form II of Compound (I), shown in FIG. 4, was obtained using a Bruker DRX-600 nuclear magnetic resonance (NMR) spectrometer operating at a frequency of 600.13 MHz. The sample concentration was approximately 3.1% (w/v) in  $\text{CD}_3\text{CN}$ . The reference compound was  $\text{CHD}_2\text{CN}$  (1.94 ppm). The spectrum was obtained at  $0^\circ\text{C}$  to sharpen signals that were broad at ambient temperature. Signal assignments following the numbered structural formula of Compound (I) below are provided in Table 1.



(I)

15

**Table 1.  $^1\text{H}$  NMR Signal Assignments**

$\delta_{\text{H}}$ (ppm)	Multiplicity*	Assignment <sup>†, #, §</sup>	$\delta_{\text{H}}$ (ppm)	Multiplicity*	Assignment <sup>†, #, §</sup>
8.24	d, J~4	C <sub>3</sub> H, C <sub>5</sub> H	4.34	o m	C <sub>2</sub> H
7.24	o m	C <sub>2</sub> 'H, C <sub>6</sub> 'H	4.15	o m	C <sub>6</sub> H
7.17	o m	C <sub>3</sub> 'H, C <sub>5</sub> 'H	3.33	o m	C <sub>4a</sub> H <sub>2</sub>
6.54	t, J=4.8	C <sub>4</sub> H	2.99–	o m	C <sub>2</sub> H, C <sub>6</sub> H
6.11	br t, J~6	NH	2.73	s	C <sub>4a</sub> 'H <sub>3</sub>
5.03	o m	C <sub>1a</sub> 'H	2.31	o m	C <sub>4</sub> H
4.78	br d, J <sub>HF</sub> -48.0	C <sub>3</sub> H (rotamer)	2.00	o m	C <sub>5</sub> H
4.73	br d, J <sub>HF</sub> -47.9	C <sub>3</sub> H (rotamer)	1.54	o m	C <sub>5</sub> H
			1.45		

\*Multiplicity: s = singlet; d = doublet; br = broad; t = triplet; m = multiplet; o = overlapped; J = coupling constant in Hertz.

<sup>†</sup>In solution, L-001067743-005K exists as a 1:1 mixture of carbamate rotamers.


Assignments are grouped except where noted by: (rotamer).

<sup>#</sup>The signal at 2.26 ppm is due to H<sub>2</sub>O.

<sup>\$</sup>A multiplet at 3.36 ppm is assigned to a low-level unknown.

### EXAMPLE 6: <sup>13</sup>C Nuclear Magnetic Resonance Spectrum

The <sup>13</sup>C nuclear magnetic resonance (<sup>13</sup>C NMR) spectrum of crystalline Form II of Compound (I) shown in FIG. 5 was obtained using a Bruker DRX-600 NMR spectrometer operating at a frequency of 150.90 MHz. The sample concentration was approximately 3.1% (w/v) in CD<sub>3</sub>CN. The reference compound was CD<sub>3</sub>CN (1.39 ppm for the CD<sub>3</sub> group; the –CN group was observed at 118.45 ppm). The spectrum was obtained at 0°C to sharpen signals that were broad at ambient temperature. Signal assignments following the numbered structural formula of Compound (I) below are provided in Table 2.



(I)

Table 2. <sup>13</sup>C NMR Signal Assignments

$\delta_C$ (ppm)	Assignment*	$\delta_C$ (ppm)	Assignment*
163.56	C <sub>1'</sub>	87.69, d, J <sub>CF</sub> = 174.1	C <sub>3</sub> (rotamer)
158.9 (broad)	C <sub>3'</sub> , C <sub>5'</sub>	67.46, 67.36	C <sub>1a''</sub>
156.29, 156.25	C <sub>1a</sub>	48.40, d, J <sub>CF</sub> = 20.7	C <sub>2</sub> (rotamer)
138.75, 138.71	C <sub>4''</sub>	48.16, d, J <sub>CF</sub> = 20.7	C <sub>2</sub> (rotamer)
135.13	C <sub>1''</sub>	44.06, 43.90	C <sub>6</sub>
129.96	C <sub>3''</sub> , C <sub>5''</sub>	43.09, d, J <sub>CF</sub> = 2.7	C <sub>4a</sub>
128.89, 128.78	C <sub>2''</sub> , C <sub>6''</sub>	39.22, d, J <sub>CF</sub> = 20.3	C <sub>4</sub>
111.34	C <sub>4'</sub>	24.58, 24.38	C <sub>5</sub>
87.88, d, J <sub>CF</sub> = 174.3	C <sub>3</sub> (rotamer)	21.15	C <sub>4'</sub>

**EXAMPLE 7: Thermogravimetric Analysis Curve**

The thermogravimetric (TG) analysis curve for Form II of Compound (I) was obtained under nitrogen flow at a heating rate of 10°C per minute. A weight loss of 0.04% was observed up to 150°C (FIG. 6). Further weight loss above 150°C was attributed to decomposition.

**EXAMPLE 8: Differential Scanning Calorimetry**

The differential scanning calorimetry (DSC) curve for Form II of Compound (I) was obtained under nitrogen flow at a heating rate of 10°C per minute in a crimped pan (FIG. 7). The DSC curve displayed a melting endotherm with extrapolated onset temperature of 121°C, a peak temperature of 123°C, and a heat of fusion of 112.7 J/g.

**EXAMPLE 9: Particle Size Evaluation**

Samples of micronized crystalline Form II of Compound (I) were evaluated for particle size and development of particle size analysis method.

Particle size information was obtained using a Sympatec HELOS/KF particle size analyzer equipped with RODOS/M dispersion module, VIBRI vibratory feeder and an R2 lens (upper limit = 87.5 microns). 100 ± 10 mg sample of micronized Form II of Compound (I) was transferred to the VIBRI and spread evenly to sure a consistent feed rate. Analysis was conducted using a feed rate of 25% and a feed pressure of 3.0 bar. Trigger conditions were as follows: timebase - 100 ms; start - c.opt ≥ 1.0%; valid - always; stop - 5 s, c.opt ≤ 1.0% or 10 s real time. The data was analyzed using Sympatec's WINDOX 5 software. The results are provided in Table 3.

**Table 3. X10, X50 and X90 Particle Size of Micronized Form II**

Replicate	Analysis 1			Analysis 2		
	X10 (µm)	X50 (µm)	X90 (µm)	X10 (µm)	X50 (µm)	X90 (µm)
1	0.78	1.90	4.41	0.79	1.93	4.45
2	0.79	1.93	4.43	0.80	1.95	4.63
3	0.79	1.92	4.46	0.79	1.92	4.42
4	0.78	1.98	4.75	0.79	1.93	4.43
5	0.79	1.94	4.62	0.79	1.93	4.60
6	0.79	1.95	4.48	0.79	1.91	4.42
Average	0.79	1.94	4.53	0.79	1.93	4.49

Replicate	Analysis 1			Analysis 2		
	X10 (µm)	X50 (µm)	X90 (µm)	X10 (µm)	X50 (µm)	X90 (µm)
Standard Deviation	0.01	0.03	0.13	0.00	0.01	0.10
% Relative Standard Deviation	1.27	1.55	2.87	0.00	0.52	2.23

Based on the development studies, the method parameters considered suitable for routine particle size analysis of micronized Compound (I) samples are set forth below:

Calculation Mode:	HRLD	Trigger Conditions:	
Sample Mass:	100 ± 25 mg	Timebase:	100ms
Feeder:	VIBRI	Start:	c.opt ≥ 1.0%
Dispenser:	RODOS/M	Valid:	always
Feed Pressure:	3.0 bar	Stop:	5 s, c.opt ≤ 1.0% or 10 s real time
Feed Rate:	25%	Lens:	R2 (upper limit 87.5 microns)

#### EXAMPLE 10: Effect of Particle Size on Drug Absorption

To test the particle size (PS) effect on absorption, monkey studies were conducted at a 90-mg/kg dose to compare the unmilled and jet-milled Compound (I) as suspensions in 0.5% methocel. Male Rhesus monkeys were administered unmilled and jet-milled crystalline Form II of Compound (I) as suspensions in 0.5% methocel at a dose of 90 mg/kg. The monkeys were fed approximately one hour prior to dosing. As shown in Table 4, the jet-milled Form II with a mean PS of 4.5 µm increased drug exposure by 1.43 fold vs. the unmilled Form II with a mean PS of 14 µm.

**Table 4. Particle Size Effect in Fed Male Rhesus Monkeys\***

Mean parameters	Particle size** (µm)	Surface area (m <sup>2</sup> /g)	AUC <sub>0-24</sub> (µM*hr)	C <sub>max</sub> (µM)	T <sub>max</sub> (hr)
Jet-milled Form II	4.5	2.3	119.3 ± 20.9	9.70 ± 3.00	6.67 ± 1.20
Unmilled Form II	14	1.5	83.7 ± 14.9	6.53 ± 2.76	3.33 ± 1.15

\* The monkeys were fed 10 biscuits approximately one hour prior to dosing.

\*\* PS was measured using bulk drug powder.

**EXAMPLE 11: Affinity for NMDA-GluN2B Receptors**

Radioligand binding assays were performed at room temperature (or 37°C) in 96-well microtiter plates with a final assay volume of 1.0 mL in 20 mM HEPES buffer (pH 7.4) containing 150 mM NaCl. Crystalline Form II of Compound (I) was prepared at 10 mM in dimethylsulfoxide (DMSO) and serially diluted with DMSO to yield 20 µL of each of 10 solutions differing by 3-fold in concentration (Mosser et al., 2003). Nonspecific binding was assessed using *N*-(3-chlorobenzyl)-4-iodobenzimidamide (final concentration, 10 µM), and total binding was measured by addition of DMSO (final concentration, 2%). L(tk-) cell membranes expressing human GluN1a/GluN2B receptors (final concentration, 40 pM) and [<sup>3</sup>H]-[(*E*)-*N*1-(2-methoxybenzyl)-cinnamamide] (final concentration, 1 nM) were added to all wells of the microtiter plate. After 3 h (24 h or 48 h) of incubation at room temperature (or 37°C), samples were filtered through Packard GF/B filters (presoaked in 0.05% polyethylenimine [PEI; Sigma P-3143]) and washed 10 times with 1 mL of cold 20 mM HEPES buffer per wash. After vacuum drying of the filter plates, 50 µL of Packard Microscint-20 was added and the bound radioactivity determined in a Packard TopCount.

Analogous binding experiments, to those described above for the cloned human receptor expressed in L(tk-) cells, were also performed using whole brain homogenate (rat), frontal cortex homogenate (dog and rhesus monkey), and temporal cortex homogenate (human).

Form II potently inhibited radioligand ([<sup>3</sup>H]Compound-2) binding to human NMDA- GluN1a/GluN2B receptors expressed in L(tk-) cells as well as brain tissue homogenates from all tested species (rat, dog, rhesus monkey, human). The binding affinity of Form II determined using human temporal cortex homogenate yielded  $K_i$  values of 3.1 nM (0.0031 µM) and 8.1 nM (0.0081 µM) at room temperature and 37°C, respectively. Findings in other species were consistent with the human data (Table 5).

**Table 5. Binding Affinities to Expressed Human GluN1a/GluN2B Receptors in L(tk-) Membranes and Brain Tissue Homogenates**

Species	$K_i$ (nM)	$K_i$ (nM)	$K_i$ (nM)
	4°C	Room Temp.	37°C
Human GluN1a/GluN2B in L(tk-)	1.8	4.9	31
Human Temporal Cortex	—	3.1	8.1

Species	K <sub>i</sub> (nM) 4°C	K <sub>i</sub> (nM) Room Temp.	K <sub>i</sub> (nM) 37°C
<b>Rhesus</b> Frontal Cortex	—	3.1	14
<b>Dog</b> Frontal Cortex	—	3.2	13
<b>Rat</b> Brain Homogenate	—	3.3	14

Values are geometric means.

#### EXAMPLE 12: Functional Activity and Selectivity for NMDA Receptors

The inhibition of calcium influx into L(tk-) cells expressing either GluN1a/GluN2B or GluN1a/GluN2A human receptors was measured to determine the IC<sub>50</sub> of Compound (I) inhibition of NMDA receptor functions.

GluN1a/GluN2B (or GluN1a/GluN2A) receptor-transfected L(tk-) cells were plated in 96-well format at  $3 \times 10^4$  cells per well and grown for 1 day in normal growth medium (Dulbecco's modified Eagle medium with Na pyruvate, 4500 mg glucose, penicillin/streptomycin, glutamine, 10% fetal calf serum, and 0.5 mg/mL geneticin).

GluN1a/GluN2B (GluN1a/GluN2A) expression in these cells was induced by the addition of 10 nM dexamethasone in the presence of 500  $\mu$ M ketamine for 16-24 h. Crystalline Form II of Compound (I) was prepared in DMSO and serially diluted with DMSO to yield 10 solutions differing by 3-fold in concentration. A 96-well drug plate was prepared by diluting the DMSO solution 250-fold into assay buffer (Mg<sup>2+</sup>-free Hanks Balanced Salt Solution containing 20 mM HEPES, 2 mM CaCl<sub>2</sub>, 0.1% bovine serum albumin, and 250  $\mu$ M probenecid). After induction, the cells were washed twice (Labsystem cell washer; 3-fold dilutions leaving 100  $\mu$ L) with assay buffer and loaded with the calcium fluorescence indicator fluo-3 AM (4  $\mu$ M) in assay buffer containing Pluronic F-127 and 100  $\mu$ M ketamine at 37°C for 1 h. The cells were then washed 8 times with assay buffer leaving 100  $\mu$ L of buffer in each well. Fluorescence intensity was immediately measured in a FLIPR (Fluorometric Imaging Plate Reader; Molecular Devices, Sunnyvale, CA) using an excitation of 488 nm and emission at 530 nm. Five seconds after starting the recording of fluorescence intensity, 50  $\mu$ L of agonist solution (40  $\mu$ M glutamate/glycine; final concentration, 10  $\mu$ M) was added, and after 1 min, when the fluorescence signal was stable, 50  $\mu$ L of Form II and control solutions from the drug plate were added and the fluorescence intensity recorded for another 30 min.

Form II inhibited calcium influx into agonist-stimulated NMDA-GluN1a/GluN2B L(tk-) cells with an  $IC_{50}$  of 3.6 nM (0.0036  $\mu$ M) but had no effect on calcium influx into agonist-stimulated NR1a/NR2A cells at concentrations up to 30,000 nM (30  $\mu$ M). The results demonstrate that Form II is a potent, highly selective antagonist of NMDA  
 5 receptors containing the GluN2B subunit.

**EXAMPLE 13: Single-Dose Pharmacokinetics in Rats and Monkeys (Oral and IV)**

Male Sprague-Dawley rats (N = 8; N = 4 oral and N = 4 intravenous [IV]) weighing approximately 250–280 g and male rhesus monkeys (N = 4, crossover) weighing 6.0–8.8 kg were used for the PK studies. In rats, the IV dose of crystalline Form II of  
 10 Compound (I) at 2 mg/kg (0.4 mL/kg) was administered as a bolus and the oral dose was administered by gavage at 15 mg/kg (5 mL/kg). In monkeys, the IV dose of Form II at 2 mg/kg (0.1 mL/kg) was administered as a slow bolus and the oral dose was administered via nasogastric tube at 15 mg/kg (5 mL/kg). Blood samples were serially collected pre-dose, and at 0.083 (IV only), 0.25, 0.50, 1, 2, 4, 6, 8, 10, 24, 48, 72, and 96 h post-dose.  
 15 The concentrations of Form II in rat and monkey plasma were determined by LC/MS/MS analysis in the positive ion mode using a heated nebulizer interface. The lower limit of quantitation was 2.5 ng/mL (approximately 0.00697  $\mu$ M).

PK parameters are summarized in Table 6. After oral administration to both rats and monkeys, Form II was rapidly absorbed ( $T_{max}$  <1 h for rats and approximately 2 h for  
 20 monkeys) and exhibited good bioavailability in both species (60% in rats and 50% in monkeys). Volume of distribution ( $V_{dss}$ ) was estimated after IV administration and exceeded total body water in rats (mean  $V_{dss}$  = 3.1 L/kg) and was moderate in monkeys (mean  $V_{dss}$  = 2.9 L/kg). High total plasma clearance ( $CL_p$ ) was exhibited after IV administration in both species (mean  $CL_p$  = 28.4 mL/min/kg in rats and 15.1 mL/min/kg  
 25 in monkeys). Half-life ( $t_{1/2}$ ), which was calculated after the IV dose, was longer in monkeys (mean  $t_{1/2}$  = 4.2 h) compared to rats (mean  $t_{1/2}$  = 1.7 h).

**Table 6. Pharmacokinetics in Rats and Monkeys during Oral (PO) and Intravenous (IV) Administration at Doses of 2 and 15 mg/kg**

Parameter <sup>a</sup>	Rat (N = 4)	Monkey (N = 4)
IV Dose (mg/kg)	2	2
$AUC_{(0-\infty)}$ ( $\mu$ M·h)	3.31 (0.41)	6.67 (2.12)

CL <sub>p</sub> (mL/min/kg)	28.4 (3.4)	15.1 (4.8)
Vd <sub>ss</sub> (L/kg)	3.1 (0.3)	2.9 (2.5)
t <sub>1/2</sub> (h)	1.7 (0.4)	4.2 (3.7)
PO Dose (mg/kg)	15	15
AUC <sub>(0-∞)</sub> (μM·h)	14.9 (2.1)	25.1 (11.9)
C <sub>max</sub> (μM)	5.9 (1.1)	3.6 (1.3)
T <sub>max</sub> (h)	0.44 (0.1)	2.1 (1.4)
Bioavailability (%)	60 <sup>b</sup>	50 (12) <sup>c</sup>

<sup>a</sup> Data are presented as mean (S.D.)

<sup>b</sup> Based on AUC<sub>0-∞</sub> (dose normalized) values after IV and PO dosing (non-crossover study design).

5 <sup>c</sup> Based on AUC<sub>0-∞</sub> (dose normalized) values after IV and PO dosing (crossover study design).

AUC, area under the curve; CL<sub>p</sub>, total plasma clearance; C<sub>max</sub>, maximum plasma concentration of drug; T<sub>max</sub>, time to C<sub>max</sub>; t<sub>1/2</sub>, half-life; Vd<sub>ss</sub>, volume of distribution at steady state.

#### 10 **EXAMPLE 14: Acute Depression Model – Forced Swim Test**

Young, adult, male Sprague-Dawley rats were randomly assigned across the treatment groups and were administered vehicle (0.5% MC/0.02% SLS), the reference compound desipramine (20 mg/kg; a tricyclic antidepressant; Sigma, Lot# 078K1326) dissolved in sterile water, or crystalline Form II of Compound (I) (0.1, 0.3, 1, 3, 10, and 30 mg/kg) suspended in 0.5% MC/0.02% SLS, twice on Day 1 (after habituation; ~24 h prior to test, and prior to dark cycle) and once on Day 2 (30 min pre-test for desipramine and 45 min pre-test for Form II and vehicle).

Each Forced Swim chamber was constructed of clear acrylic (height, 40 cm; diameter, 20.3 cm). Rats were subjected to a pre-dose swim test of one 15-min session in cylinders containing water at 23°C ± 1°C, followed approximately 24 h later by the experimental 5-min session. The water level was 16 cm deep during habituation and 30 cm deep during the test. Immobility, climbing, and swimming behaviors were recorded every 5 s for a total of 60 counts per subject. When an animal was unable to maintain a

posture with its nose above water, it was immediately removed from the water and eliminated from the study. Blood was collected at the completion of swim test procedures and plasma was analyzed for Form II concentrations.

Form II (1, 3, 10, and 30 mg/kg) significantly decreased immobility frequency ( $P < 0.001$ ) and significantly increased swimming behavior ( $P < 0.01$  for 1, 3, and 30 mg/kg;  $P < 0.05$  for 10 mg/kg) compared to the vehicle control (FIG. 8), but did not affect climbing behavior except at the dose of 3 mg/kg ( $P < 0.05$ ). Desipramine (20 mg/kg) significantly decreased immobility ( $P < 0.001$ ) and significantly increased climbing behavior ( $P < 0.01$ ) compared to the vehicle control, with no change in swimming behavior. Form II plasma levels were approximately 15, 120, 390, 1420, 4700, and 14110 nM at the time of sampling, corresponding to approximately 5, 29, 56, 83, 94, and 98% RO, respectively, in rats. The ED<sub>50</sub> for increase in frequency of swimming and decrease in immobility were ~0.3 and 0.7 mg/kg, respectively, corresponding to RO of ~30% and 50%.

#### **EXAMPLE 15: Acute Depression Model – Locomotor Assay**

To confirm that the effect of Form II in the forced swim test was not due to a general increase in activity levels, rats were subjected to a locomotor assay following oral Form II administration. Adult male Sprague-Dawley rats (N = 42) were randomly assigned across the treatment groups (vehicle or Compound (I) at 0.1, 0.3, 1, 3, 10, and 30 mg/kg; N = 6/group). Locomotor activity was assessed during the light cycle in photocell-monitored cages (Hamilton Kinder, San Diego, CA). Each cage consisted of a standard plastic rat cage (24 × 45.5 cm) surrounded by a stainless steel frame. Infrared photocell beams were located across the long axis of the frame to measure the ambulatory distance traveled. A second set of beams was placed above the floor and was used to measure rearing activity. Photocell beam interruptions were recorded by a computer system. Filter tops were placed on top of the test enclosures during testing. Rats were administered either vehicle or test compound via oral gavage twice on Day 1 (approximately 24 h before the test and prior to dark cycle) and once on Day 2 (45 min prior to placing in the locomotor cages for a 60-min test). Locomotor activity was captured in 5-min bins.

Form II (1, 3, 10, and 30 mg/kg) significantly increased distance traveled ( $P < 0.01$  for 1 and 3 mg/kg;  $P < 0.001$  for 10 and 30 mg/kg) compared to vehicle control during the first 5 min of testing (timing correlates with time of forced swim test). Form II (1, 3, 10, and 30 mg/kg) significantly increased total distance traveled ( $P < 0.01$  for 1 mg/kg;  $P <$

0.001 for 3, 10, and 30 mg/kg) compared to vehicle control summed over the 60-min test. The ED<sub>50</sub> for increase in locomotor activity was ~2mg/kg, translating to an RO of ~75%, which is higher than the ED<sub>50</sub> for increase in frequency of swimming and decrease in immobility. No locomotor effects were observed for the 0.1 and 0.3 mg/kg dose groups  
5 (FIG. 8).

#### **EXAMPLE 16: Blood Pressure Profile in Chronically Instrumented Rats**

Six (n=6) rats were used to study the effect of crystalline Form II of Compound (I) on systemic blood pressure and heart rate. A single oral gavage dose was administered (volume: 5 mL/kg). Prior to dosing, a 24 hours baseline was recorded. The effect of  
10 crystalline Form II of Compound (I) was recorded for 24 hours. A Latin square was performed on using 3 doses of crystalline Form II of Compound (I) (0.3, 1, and 3 mg/kg) and corresponding vehicle (n=6 per group). Following the completion of the Latin square, two additional dose concentrations were evaluated (0.6 and 10 mg/kg, n=3 per group).

Three (3) rats were used to study the effect of MK-801 (dizocilpine). On day 1,  
15 baseline readings were recorded for 24 hours prior to dosing. On day 2, the rats were administered a single intravenous (tail vein) bolus of 0.9% saline (volume ~0.2 mL) and monitored continuously (via telemetry) for at least 24 hours. On day 3, the rats were administered a single intravenous (tail vein) bolus of MK-801 (200 µg/kg; volume ~0.2 mL) and monitored continuously (via telemetry) for at least 24 hours.

20 In order to investigate a potential mechanism of crystalline Form II of Compound (I), crystalline Form II of Compound (I) was administered in combination with different pharmacological inhibitors. Atenolol (β1 blocker, 1 mg/kg, IV bolus) and prazosin (α1 adrenergic receptor antagonist, 200 µg/kg, IV bolus) were administered 30 minutes prior to crystalline Form II of Compound (I) (1 mg/kg). All IV doses were administered in  
25 manually restrained rats via tail vena-puncture. It should be noted that vehicle controls were also administered.

One hour following a single oral gavage dose, crystalline Form II of Compound (I) elicited a dose-dependent increase in mean arterial pressure (MAP) ( $4.2 \pm 0.9$ ,  $6.8 \pm 1.2$ ,  $15.7 \pm 3.2$ ,  $17.1 \pm 2.3$ , and  $19.1 \pm 2.4$  mmHg for 0.3, 0.6, 1, 3, and 10 mg/kg respectively).  
30 Heart rate (HR) also increased over the dose ranges, however, the dose-dependency of these effects was not evident, as markedly larger but similar effects were noted at 3 and 10 mg/kg (*i.e.*,  $15 \pm 10$  and  $10 \pm 15$ , and  $34 \pm 13$  bpm at 0.3, 0.6, and 1 mg/kg vs.  $66 \pm 9$ , and

71 ± 25 bpm at 3 and 10 mg/kg) (FIG. 9). Thus, crystalline Form II of Compound (I) (when given orally as a single-dose) elicited hemodynamic responses consistent (albeit markedly to a lower extent) with those seen for MK-801, a well-established NMDA antagonist. The latter finding was unexpected for crystalline Form II of Compound (I).

5           Rat activity (recorded via telemetry) followed a similar trend as the heart rate response. That is, the 3 and 10 mg/kg doses had much larger effects than the lower doses (0.012 ± 0.025, 0.003 ± 0.033, and 0.049 ± 0.037 at 0.3, 0.6, and 1 mg/kg versus 0.174 ± 0.048 and 0.232 ± 0.057 at 3 and 10 mg/kg). Interestingly, the hemodynamic changes during the first 3.5 hours post administration were linearly correlated with the level of  
10 activity (as estimated from telemetry recordings); in particular, activity was an excellent predictor of HR ( $R^2 = 0.67$ ), and therefore, the excitatory effects of crystalline Form II of Compound (I) may explain the observed increases in HR (particularly at the higher dose-levels) in rats.

To investigate a potential mechanism of the crystalline Form II of Compound (I)-  
15 medicated hemodynamic changes, crystalline Form II of Compound (I) was administered in combination with either atenolol ( $\beta_1$  AR blocker, 1 mg/kg) or prazosin ( $\alpha_1$  AR antagonist, 200  $\mu$ g/kg). As anticipated, atenolol minimally affected MAP ( $-3.3 \pm 3.1$  mmHg) but markedly reduced HR ( $-87 \pm 11$  bpm), while prazosin reduced MAP ( $18 \pm 2$  mmHg) but minimally reduced HR ( $-9 \pm 24$  bpm). In these settings (*i.e.*,  $\beta_1$  AR or  $\alpha_1$  AR  
20 blockade), crystalline Form II of Compound (I) (at 1 mg/kg PO) was administered and the hemodynamic responses were studied 1 hour post dosing. In  $\beta_1$  AR blocked animals, crystalline Form II of Compound (I) (1 mg/kg) had negligible effects in MAP ( $5.8 \pm 4.8$  mmHg) and HR ( $-7 \pm 17$  bpm) when compared to pre-blockade values (*i.e.*, baseline); however, HR did increase  $53 \pm 17$  bpm post-dosing. Crystalline Form II of Compound (I)  
25 (1 mg/kg), in the setting of  $\alpha_1$  AR blockade, increased HR ( $118 \pm 20$  bpm) but lacked MAP changes ( $-3.8 \pm 2.0$  bpm) when compared to pre-blockade values (*i.e.*, baseline); however, MAP did increase ( $11.3 \pm 2.0$  mmHg) with dosing.

**Table 7. Peak systemic hemodynamics changes 1 hour post single oral gavage dose of Crystalline Form II of Compound (I); represented as vehicle subtracted change from time-controlled baseline**

<i>Parameters</i>	<i>Crystalline Form II of Compound (I)</i>				
	<i>0.3 mg/kg</i>	<i>0.6 mg/kg*</i>	<i>1.0 mg/kg</i>	<i>3.0 mg/kg</i>	<i>10 mg/kg*</i>
<i>SAP (mmHg)</i>	4.3 ± 1.2	6.3 ± 1.6	16.8 ± 3.7	19.2 ± 2.4	20.7 ± 2.4
<i>DAP (mmHg)</i>	4.1 ± 0.9	7.8 ± 1.1	15.2 ± 3.0	15.0 ± 2.0	17.3 ± 1.9
<i>MAP (mmHg)</i>	4.2 ± 0.9	6.8 ± 1.2	15.7 ± 3.2	17.1 ± 2.3	19.1 ± 2.4
<i>PP (mmHg)</i>	0.1 ± 1.0	-1.5 ± 1.4	1.4 ± 1.1	4.0 ± 0.9	3.1 ± 1.3
<i>HR (bpm)</i>	15 ± 10	10 ± 15	34 ± 13	66 ± 9	71 ± 25
<i>Activity (arbitrary unit)</i>	0.012 ±	0.003 ±	0.049 ±	0.174 ±	0.232 ±
	0.025	0.033	0.037	0.048	0.057

Values are mean ± standard error of the mean (n = 6), \* n = 3.

#### 5 **EXAMPLE 17: Human Pharmacokinetic Study**

Twenty-four healthy, young male subjects were assigned to 1 of 3 sequential treatment panels (A, B, and C). For each panel of 8 subjects, 2 subjects received placebo and 6 subjects were administered single ascending oral doses of crystalline Form II of Compound (I) with a minimum 7-day washout between each dose: Panel A (0.1, 0.2, 0.5, 1, and 2 mg); Panel B (2, 4, 8, and 15 mg, and 4 mg with food); and Panel C (15 and 20 mg). Blood samples were collected pre-dose and 0.5, 1, 1.5, 2, 3, 4, 6, 9, 12, 18, 24, 30, 48, and 72 h post dose. Plasma samples were analyzed for Form II concentrations using reversed phase high performance liquid chromatography with tandem mass spectrometric detection (Merck Research Laboratories). The analytical range was 0.5 to 500 nM (0.180 to 180 ng/mL).

Form II was rapidly absorbed (FIG. 10A) with mean  $T_{max}$  within 1 h post-dose across all fasted dose levels and terminal elimination half-life ranging from approximately 12 to 17 h over the 4- to 20-mg dose range.  $C_{max}$  and AUC behaved in a dose-proportional manner over the dose range studied. The 4-mg dose was administered in the fasted and fed states. Dosing in the fed state led to an approximate 1-h delay in  $T_{max}$  and an approximate 56% decrease in  $C_{max}$ , but the overall extent of absorption (AUC) was not affected (FIG. 10B, Table 8).

**Table 8. Plasma Pharmacokinetic Parameters in Healthy Male Volunteers**

<b>Parameter (unit)</b>	<b>2 mg (N = 12)</b>	<b>4 mg (N = 6)</b>	<b>4 mg (fed) (N = 6)</b>	<b>8 mg (N = 6)</b>	<b>15 mg (N = 12)</b>	<b>20 mg (N = 6)</b>
$C_{\max}$ (ng/mL)	51.47 (16.96)	140.59 (22.27)	80.00 (18.46)	211.93 (42.70)	385.21 (99.15)	590.22 (87.09)
$T_{\max}$ (h)	1.08 (0.42)	0.67 (0.26)	2.00 (1.30)	0.92 (0.20)	1.04 (0.33)	1.00 (0.32)
$AUC_{0-t}$ (h*ng/mL)	397.64 (83.40)	1063.42 (196.97)	1080.32 (252.37)	2151.14 (442.25)	3521.08 (702.88)	5593.55 (874.71)
$AUC_{0-\infty}$ (h*ng/mL)	493.25 (101.99)	1110.27 (200.68)	1137.98 (261.23)	2255.09 (473.43)	4396.45 (966.26)	5909.82 (1173.62)
$t_{1/2}$ (h)	16.14 (3.12)	16.92 (1.88)	17.31 (2.20)	16.96 (2.10)	16.79 (4.67)	15.90 (5.91)

Data are presented as mean (S.D.)

AUC, area under the curve;  $C_{\max}$ , maximum plasma concentration of drug;  $T_{\max}$ , time to  $C_{\max}$ ;  $t_{1/2}$ , half-life.

## 5 EXAMPLE 18: Safety Pharmacology Studies

The potential effects of Form II of Compound (I) on the functions of the nervous system, cardiovascular system, and respiratory system were evaluated in a battery of safety pharmacology studies that included an in vitro human ether-a-go-go-related gene (hERG) assay and in vivo studies in rats, dogs, and monkeys. CNS effects were tested in  
10 rats at single doses at 15, 75, and 225 mg/kg.

Effects of Form II on the cardiovascular system were tested in several studies. Potential QTc effects were tested in vitro and in vivo. Form II inhibited hERG current in vitro with an  $IC_{50}$  of approximately 13  $\mu$ M (4660 ng/mL). In anesthetized dogs dosed by continuous IV infusion for 90 min at escalating rates, Form II did not affect heart rhythm  
15 or any ECG parameter, including QTc interval duration, even though the plasma concentration of Form II at the end of infusion was approximately 74  $\mu$ M (25450 ng/mL). In conscious, chair-restrained monkeys given single oral doses at 15, 75, and 150 mg/kg and monitored for 6 h post-dose, Form II did not affect heart rhythm or ECG parameters at  
20 any dose level, even though the peak plasma concentration was approximately 15  $\mu$ M (5375 ng/mL) at the highest dose level.

Hemodynamic effects were tested in dogs and monkeys. Form II did not affect heart rate, arterial blood pressure, or femoral blood flow in anesthetized dogs dosed by continuous IV infusion for 90 min at escalating rates. However, Form II did produce dose-independent increases in systolic (12–15 mm Hg), diastolic (7–10 mm Hg), and mean arterial pressures (9–12 mm Hg) in conscious, chair-restrained monkeys given single oral doses at 15, 75, or 150 mg/kg and monitored for 6 h post-dose. At these dose levels,  $C_{\max}$  was approximately 3, 9, and 15  $\mu\text{M}$ , (1075, 3225, and 5375 ng/mL), respectively.

Respiratory effects were investigated in rats and dogs. Single oral doses of Form II at 15, 75, and 225 mg/kg produced respiratory stimulation in male rats, reflected in increased respiratory rate, tidal volume, and minute ventilation, and decreased PenH (an index of airway resistance). Respiratory function returned to baseline within 5 to 6 h at 15 mg/kg but remained altered at the end of the 6-h monitoring period at  $\geq 75$  mg/kg. In anesthetized dogs dosed at 3 mg/kg by 5-min IV infusion and monitored at intervals for 60 min, Form II did not affect intrapulmonary pressure, peak expiratory flow, airway resistance, lung compliance, tidal volume, respiration rate, or blood pH,  $\text{pCO}_2$ , or  $\text{pO}_2$ . Mean Form II plasma concentration was approximately 30  $\mu\text{M}$  (10600 ng/mL) at the end of infusion and approximately 4  $\mu\text{M}$  (1250 ng/mL) at 60 min post-infusion.

#### **EXAMPLE 19: Neurotoxicity in Rats**

Four groups of 24 Sprague-Dawley rats (12/sex) were given single doses of vehicle (0.5% methylcellulose [MC] and 0.02% sodium lauryl sulfate [SLS] in deionized water) or crystalline Form II of Compound (I) at 10, 30 or 100 mg/kg by oral gavage at a dose volume of 10 mL/kg. An additional group of 12 male rats was given single doses of (5S,10R)-(+)-5-methyl-10,11-dihydro-5H-dibenzo-[a,d]cyclohepten-5,10-imine maleate; NBF, neutral buffered formalin (“MK-801”, a non-competitive antagonist of the NMDA receptor; positive control) at 10 mg/kg by subcutaneous injection at a dose volume of 2 mL/kg. Six rats per sex in each group were terminated and necropsied at 4 to 6 h post-dose, and the remaining rats in each group were terminated and necropsied 3 days post-dose (on Day 4). In-life observations and measurements included body weight and clinical observations. At termination, rats were anesthetized and perfusion fixed. At necropsy, the brain was collected for histopathological evaluation.

Animals in Form II and MK-801 assessment groups were terminated at the scheduled necropsy intervals (4–6 h post-dose or Day 4). All animals were anesthetized

with an isoflurane/oxygen mixture and perfused via the left cardiac ventricle with heparinized 0.001% sodium nitrite in saline. The saline wash was followed by perfusion of 10% neutral buffered formalin (NBF). Brains were harvested, weighed, and stored in 10% NBF.

5           The brains were sectioned into 2 mm coronal sections to produce multiple sections in 3 blocks for each animal. The following brain regions were stained: neocortex, paleocortex, basal nuclei, limbic system, thalamus/hypothalamus, midbrain regions, cerebellum, pons region, and medulla oblongata. All brain sections from all animals sacrificed 4 to 6 h after dosing and all animals sacrificed 3 days after dosing were  
10 embedded in paraffin, sectioned at 5  $\mu$ m, stained with hematoxylin and eosin and examined microscopically. For rats sacrificed on Day 4 (3 days after dosing), serial sections from Blocks 1 and 2 were stained with Fluoro-Jade B (a stain increasing the sensitivity of evaluating the brain for neuronal degeneration) and glial fibrillary acidic protein (a stain for astrocyte reactions) and examined microscopically. Three additional  
15 groups of rats (4 males and 3 females per group) were orally dosed in the same manner with Form II, and 24-h serial blood samples were obtained and analyzed for Compound (I) plasma concentrations and evaluated for systemic exposure.

          There were no Form II-associated morphologic effects at any of the dose levels or time points. In rats, single doses of Form II (10, 30, and 100 mg/kg) did not produce  
20 vacuolation or necrosis in all examined regions of the brain. At these doses, mean  $C_{\max}$  was approximately 4, 14, and 26  $\mu$ M (1433, 5018, and 9319 ng/mL), respectively. By contrast, all of the MK-801 (10 mg/kg)-dosed animals had vacuolation and necrosis in cingulate gyrus neurons, consistent with previous reports (Fix *et al.*, *Brain Res*, 696:194-204 (1995)). At the 4–6 h time point, the animals treated with MK-801 (6 males; Group 5)  
25 all had numerous vacuolated neurons in cortical layers 3 and 4 in the cingulate gyrus region of the cerebral cortex. The appearance of the MK-801-treated males was completely consistent with previously published descriptions. Affected neurons were characterized by numerous, tightly packed, somewhat distinct, vacuoles filling the cytoplasm. At the 4–6 h time point, none of the vehicle controls and none of the Form II-  
30 treated animals had any evidence of cytoplasmic vacuoles in any of the neurons in the cingulate gyrus.

On Day 4, all the animals treated with MK-801 (6 males; Group 5) had necrotic neurons in cortical layers 3 and 4 in the cingulate gyrus region of the cerebral cortex. The appearance of the MK-801-treated males was completely consistent with previously published descriptions (Fix et al., 1996). Using the Fluoro-Jade B stain, necrotic neurons were easily visualized in all of the Day 4 MK-801-treated animals. In the MK-801-treated animals, sections stained (immunohistochemically) for glial fibrillary acidic protein showed a very slight increase of staining in the region of the cingulate gyrus. On Day 4, none of the vehicle controls and none of the Form II-treated animals had any evidence of necrotic neurons in the cingulate gyrus.

#### 10 **EXAMPLE 20: Pharmacodynamic Effects in Healthy Subjects**

A randomized, double-blind, placebo-controlled, parallel-dose group in-patient study is conducted to investigate the pharmacodynamic (PD) effects of crystalline Form II of Compound (I) in healthy subjects.

The study is designed as a three-part, sequential study. The sequential nature of the study ensures that a range of doses are investigated first in a young, healthy population (Part 1) before proceeding to higher doses in the healthy population and the comparator populations (intermediate age and elderly) (Parts 2 and 3). For the purposes of this study, “young” refers to a subject who is 18 to 45 years of age, “intermediate age” refers to a subject who is 46 to 64 years of age, and “elderly” refers to a subject who is 65 years of age or older.

#### **Part 1:**

After a screening period of up to 21 days (Days -23 to -3), 16 healthy, young male and female subjects are randomly assigned to a dose group as shown in Table 9.

**Table 9. Part 1 Parallel Dose Groups (Dosing Days 1-7)**

Age	Group*	Study Drug	Dosing Regimen
Young	1 (N=6)	Crystalline Form II of Compound (I) 8 mg	Two 4-mg tablets + one placebo tablet once-daily with food
Young	2 (N=6)	Crystalline Form II of Compound (I) 12 mg	Three 4-mg tablets once-daily with food
Young	3 (N=4)	Placebo	Three placebo tablets once-daily with food

\* Randomization is stratified such that three male subjects and three female subjects are randomized to each dose group.

**Note:** On Day-1 (time-matched baseline day), all subjects receive placebo in the fed state.

5            Each parallel dose group is balanced for gender (approximately equal number of males and females). The randomized subjects are domiciled from Day -1 through Day 8. On Day -1, the subjects receive placebo tablets in the fed state and undergo ambulatory blood pressure monitoring (ABPM) and safety assessments to establish time-matched baselines. On Days 1-7, the subjects receive the study drug in the fed state and undergo  
10 safety, pharmacokinetic (PK) and pharmacodynamics (PD) assessments. On Days 8-11, follow-up PK and safety assessments are performed.

             Safety assessments include Clinician-Administered Dissociative States Scale (CADSS), Columbia-Suicide Severity Rating Scale (C-SSRS), concomitant medication, adverse event (AE) and vital sign assessments. In PK assessments, blood samples are  
15 taken and analyzed to determine PK parameters such as  $C_{max}$ ,  $T_{max}$ ,  $AUC_{0-24}$ , and  $t_{1/2}$ . PD assessments include ABPM, plasma brain-derived neurotrophic factor (BDNF) measurements, and Profile of Mood States (POMS) assessments. Since BDNF levels have been shown to correlate with the severity of major depressive disorder (MDD) and change in response to antidepressant treatment, BDNF levels are measured in this study as a  
20 biomarker of drug activity.

**Part 1 Safety Review:**

             After completion of Part 1, safety data from Day -1 to Day 8 is reviewed. A decision is made to conduct Part 2 or an alternative dosing regimen (*e.g.*, by changing any combination of dose level, dosing interval, and fed state).

25 **Part 2:**

             After a screening period of up to 21 days (Days -23 to -3), 16 subjects (8 healthy, young male and female subjects, and 8 healthy, intermediate age male and female subjects) are randomly assigned to a dose group as shown in Table 10.

**Table 10. Part 2 Parallel Dose Groups (Dosing Days 1-7)**

Age	Group*	Study Drug†	Dosing Regimen**
Young	4 (N=6)	Crystalline Form II of Compound (I) 16 mg	Four 4-mg tablets once-daily with food
Young	5 (N=2)	Placebo	Four placebo tablets once-daily with food
Intermediate	6 (N=6)	Crystalline Form II of Compound (I) 12 mg	Three 4-mg tablets once-daily with food
Intermediate	7 (N=2)	Placebo	Three placebo tablets once-daily with food

\* Randomization is stratified such that an equal number of males and females are randomized to each dose group.

\*\* The dosing regimen may be changed based on the outcome of the Part 1 safety review.

5 The regimen may be altered by any combination of changing the dose level, dosing interval, or fed state; or repeating a dose level.

† The planned dose level may be changed based on the outcome of Part 1 safety review.

The maximum dose in young subjects is 16 mg and the maximum dose in intermediate age subjects is 12 mg.

10 **Note:** On Day -1 (time-matched baseline day), all subjects receive placebo in the fed state.

As in Part 1, the randomized subjects undergo placebo administration in the fed state and time-matched baseline ABPM and safety assessments on Day -1, followed by once-daily dosing of the study drug and pharmacokinetic (PK) and pharmacodynamics (PD) assessments on Days 8-11, and follow-up PK and safety assessments on Days 8-11. Each parallel dose group is balanced for gender (approximately equal number of males and females).

### **Part 2 Safety Review:**

20 After completion of Part 2, safety data from Day -1 to Day 8 is reviewed. A decision is made to conduct Part 3 or an alternative dosing regimen.

### **Part 3:**

After a screening period of up to 21 days (Days -23 to -3), 16 subjects (8 healthy, young male and female subjects, and 8 healthy, elderly male and female subjects) are randomly assigned to a dose group as shown in Table 11.

**Table 11. Part 3 Parallel Dose Groups (Dosing Days 1-7)**

Age	Group*	Study Drug†	Dosing Regimen**
Young	8 (N=6)	Crystalline Form II of Compound (I) 20 mg	Five 4-mg tablets once-daily with food
Young	9 (N=2)	Placebo	Five placebo tablets once-daily with food
Elderly	10 (N=6)	Crystalline Form II of Compound (I) 12 mg	Three 4-mg tablets once-daily with food
Elderly	11 (N=2)	Placebo	Three placebo tablets once-daily with food

\* Randomization is stratified such that an equal number of males and females are randomized to each dose group.

\*\* The dosing regimen may be changed based on the outcome of the Part 2 safety review.

- 5 The regimen may be altered by any combination of changing the dose level, dosing interval, or fed state; or repeating a dose level.

† The planned dose level may be changed based on the outcome of Part 2 safety review.

The maximum dose in young subjects is 20 mg and the maximum dose in elderly subjects is 16 mg.

- 10 **Note:** On Day -1 (time-matched baseline day), all subjects receive placebo in the fed state.

As in Parts 1 and 2, the randomized subjects undergo placebo administration in the fed state and time-matched baseline ABPM and safety assessments on Day -1, followed by once-daily dosing of the study drug and pharmacokinetic (PK) and pharmacodynamics (PD) assessments on Days 8-11, and follow-up PK and safety assessments on Days 8-11.

- 15 Each parallel dose group is balanced for gender (approximately equal number of males and females).

After completion of Part 3, safety data from Day -1 to Day 8 is reviewed. A decision is made as to whether or not to add additional dosing cohorts to investigate alternate dosing regimens or to repeat a regimen.

- 20 All data is subsequently analyzed for PD effects, PK profiles, safety and tolerability, as well as age and gender effects on PK, blood pressure and other safety parameters such as AE.

Following repeated daily doses of Crystalline Form II of Compound (I), steady-state predose plasma Crystalline Form II of Compound (I) concentrations were achieved

by study day 5. Crystalline Form II of Compound (I) was orally bioavailable with median Tmax values ranged from 2.00 to 3.00 hours on Day 1 and from 1.50 to 3.00 hours on Day 7 of dosing. On Days 1 and 7, Cmax and AUC values increased in an approximately dose-proportional manner over the dose range studied of Crystalline Form II of Compound (I) 8 to 20 mg. Mean plasma Crystalline Form II of Compound (I) concentrations on Day 7  
5 were approximately 20% to 40% higher from 2 to 24 hours postdose in the intermediate age and elderly subgroups, respectively, compared to the young age subgroup, and were approximately 20% higher from 2 to 24 hours postdose in female subjects compared to male subjects.

10 On Day 7, average  $t_{1/2}$  ranged from approximately 17 to 25 hours, average apparent oral clearance values ranged from approximately 3 to 4 L/h, and average apparent volume of distribution ranged from approximately 83 to 116 L. Modest accumulation was observed with seven days of daily Crystalline Form II of Compound (I) dosing as the observed accumulation index was approximately 1.6 to 1.9, on average, and steady-state  
15 accumulation index was 1.1, on average. Average Day 7 to Day 1 Cmax ratios were 1.3 to 1.4.

Clinically modest differences in PK parameters were observed in intermediate age and elderly subjects compared to young subjects, and in female subjects compared to male subjects. On Day 7, average Cmax values were approximately 30% higher in the  
20 intermediate age and elderly subgroups compared to the young age group. On Day 1, average AUC values were approximately 25% to 37% higher in the intermediate age group compared to the young age group and average AUC values were approximately 6% to 29% higher in female subjects compared to male subjects. On Day 7, average AUC values were approximately 48% to 74% higher in the elderly age group compared to the  
25 young age group and average AUC values were approximately 19% to 38% higher in female subjects compared to male subjects. On Day 7, apparent oral clearance values were approximately 18% and 30%, on average, lower in the intermediate age and elderly groups, respectively, compared to the young age group and apparent oral clearance values were approximately 15%, on average, lower in female subjects compared to male subjects.  
30 On Day 7, average  $t_{1/2}$  ranged from approximately 20 to 28 hours, and was approximately 27% and 38% higher in the intermediate age and elderly age groups, respectively,

compared to the young age group. On Day 7, average  $t_{1/2}$  ranged from approximately 21 to 24 hours, and was approximately 16% higher in female subjects compared to male subjects.

**Table 12. Summary of Plasma for Crystalline Form II of Compound (I)  
Pharmacokinetic Parameters by Dose and Day for Young Subjects**

5

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose - Young Subjects By Dose			
		8 mg	12 mg	16 mg	20 mg
$C_{max}$ (ng/mL)	Day 1				
N		6	6	6	6
Mean (SD)		179.33 (18.18)	239.00 (33.36)	305.83 (31.59)	383.50 (57.48)
Min, Max		164, 214	209, 294	264, 346	314, 476
CV%		10.14	13.96	10.33	14.99
$C_{max}$ (ng/mL)	Day 7				
N		4	6	6	6
Mean (SD)		224.00 (30.01)	273.50 (44.00)	412.75 (65.06)	534.33 (97.69)
Min, Max		191, 260	219, 325	331, 483	365, 628
CV%		13.40	16.09	15.76	18.28
$t_{max}$ (h)	Day 1				
N		6	6	6	6
Median		2.00	1.50	3.00	2.25
Min, Max		2.00, 3.03	1.00, 3.00	1.50, 3.00	1.00, 3.00
Mean (SD)		2.17 (0.42)	1.77 (0.70)	2.58 (0.66)	2.17 (0.93)
CV%		19.42	39.52	25.73	42.97
$t_{max}$ (h)	Day 7				
N		4	6	4	6
Median		3.00	2.50	1.75	1.50
Min, Max		2.00, 3.00	1.50, 3.00	1.50, 2.00	1.50, 2.00
Mean (SD)		2.75 (0.50)	2.42 (0.66)	1.75 (0.29)	1.67 (0.26)
CV%		18.18	27.50	16.50	15.49
$AUC_{0-24}$ (h*ng/mL)	Day 1				
N		6	6	6	6
Mean (SD)		1503.0 (134.7)	1857.8 (51.9)	2663.3 (393.9)	3521.5 (706.6)
Min, Max		1322.6, 1713.8	1777.4, 1906.3	2197.0, 3133.1	2507.4, 4265.5
CV%		8.96	2.80	14.79	20.07
$AUC_{0-24}$ (h*ng/mL)	Day 7				
N		4	6	4	6
Mean (SD)		2397.7 (262.1)	3062.5 (512.2)	4301.2 (1411.2)	6039.3 (1530.7)
Min, Max		2107.3, 2688.1	2614.4, 3972.0	2894.4, 6260.5	4087.2, 7880.0
CV%		10.93	16.72	32.81	25.35
$AUC_{last}$ (h*ng/mL)	Day 1				
N		6	6	6	6
Mean (SD)		1488.7 (134.3)	1835.9 (51.68)	2631.8 (392.1)	3476.8 (692.5)

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose - Young Subjects By Dose			
		8 mg	12 mg	16 mg	20 mg
Min, Max		1308.2, 1699.7	1751.6, 1880.6	2171.1, 3101.1	2481.9, 4206.0
CV%		9.02	2.82	14.90	19.92

$AUC_{last}$ (h*ng/mL)	Day 7				
N		4	6	4	6
Mean (SD)		3770.6 (1355.2)	5606.7 (1836.0)	7450.6 (4389.7)	10132 (3267.1)
Min, Max		2534.3, 5650.9	3680.6, 8885.6	3657.8, 13614.4	6494.6, 14653.4
CV%		35.94	32.75	58.92	32.25
$AUC_{inf}$ (h*ng/mL)	Day 7				
N		4	6	4	6
Mean (SD)		4109.8 (1398.1)	5933.1 (2133.5)	8023.5 (4985.1)	10468 (3506.7)
Min, Max		3187.2, 6145.9	3824.4, 9805.7	3692.1, 14922.9	6630.9, 15396.7
CV%		34.02	35.96	62.13	33.50
$AUC_{%extrap}$ (%)	Day 7				
N		4	6	4	6
Mean (SD)		8.61 (8.27)	4.77 (2.72)	5.38 (5.13)	2.88 (1.30)
Min, Max		2.84, 20.48	1.38, 9.38	0.93, 10.77	1.62, 4.83
CV%		96.11	57.15	95.36	45.18
$CL/F$ (L/h)	Day 1				
N		6	6	6	6
Mean (SD)		5.36 (0.48)	6.46 (0.18)	6.12 (0.90)	5.90 (1.31)
Min, Max		4.67, 6.05	6.29, 6.75	5.11, 7.28	4.69, 7.98
CV%		8.91	2.84	14.67	22.17
$CL/F$ (L/h)	Day 7				
N		4	6	4	6
Mean (SD)		3.37 (0.37)	4.00 (0.60)	4.01 (1.21)	3.51 (0.97)
Min, Max		2.98, 3.80	3.02, 4.59	2.56, 5.53	2.54, 4.89
CV%		10.99	14.98	30.30	27.69
$V_z/F$ (L)	Day 1				
N		6	6	6	6
Mean (SD)		101.03 (43.82)	196.87 (93.73)	175.02 (104.02)	147.76 (56.52)
Min, Max		60.75, 172.75	112.49, 376.03	64.13, 288.86	94.54, 256.58
CV%		43.37	47.61	59.43	38.25
$V_{ss}/F$ (L)	Day 7				
N		4	6	4	6
Mean (SD)		82.60 (25.30)	115.44 (14.55)	108.19 (51.29)	93.18 (19.02)
Min, Max		49.78, 108.30	99.20, 133.36	64.42, 182.39	76.51, 123.31
CV%		30.62	12.60	47.41	20.41

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose - Young Subjects By Dose			
		8 mg	12 mg	16 mg	20 mg
$t_{1/2}$ (h)	Day 7				
N		4	6	4	6
Mean (SD)		17.22 (6.20)	20.49 (4.55)	20.00 (10.22)	18.75 (2.22)
Min, Max		10.93, 25.23	15.18, 27.02	11.35, 31.46	16.31, 21.56
CV%		35.98	22.23	51.10	11.84
Dose-Normalized $C_{max}$ (ng/mL/mg)	Day 1				
N		6	6	6	6
Mean (SD)		22.42 (2.27)	19.92 (2.78)	19.11 (1.97)	19.18 (2.87)
Min, Max		20.50, 26.75	17.42, 24.50	16.50, 21.63	15.70, 23.80
CV%		10.14	13.96	10.33	14.99
Dose-Normalized $C_{max}$ (ng/mL/mg)	Day 7				
N		4	6	4	6
Mean (SD)		28.00 (3.75)	22.79 (3.67)	25.80 (4.07)	26.72 (4.88)
Min, Max		23.88, 32.50	18.25, 27.08	20.69, 30.19	18.25, 31.40
CV%		13.40	16.09	15.76	18.28
Dose-Normalized $AUC_{inf}$ (h*ng/mL/mg)	Day 7				
N		4	6	4	6
Mean (SD)		513.73 (174.76)	494.43 (177.79)	501.47 (311.57)	523.40 (175.33)
Min, Max		398.40, 768.23	318.70, 817.14	230.76, 932.68	331.55, 769.83
CV%		34.02	35.96	62.13	33.50
Observed Accumulation Index					
N		4	6	4	6
Mean (SD)		1.58 (0.19)	1.65 (0.31)	1.70 (0.54)	1.71 (0.22)
Min, Max		1.36, 1.77	1.42, 2.23	1.26, 2.41	1.41, 2.00
CV%		12.15	18.56	31.83	12.77

**Table 13. Summary of Plasma for Crystalline Form II of Compound (I)  
Pharmacokinetic Parameters by Dose and Day for Young, Intermediate Age and  
Elderly Subjects at 12 mg Dose**

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose		
		Young	Intermediate Age	Elderly
		12 mg	12 mg	12 mg
$C_{max}$ (ng/mL)	Day 1	6	6	6
N		239.00 (33.36)	257.00 (50.66)	228.83 (30.73)
Mean (SD)		209, 294	190, 345	174, 260
Min, Max		13.96	19.71	13.43
CV%				
$C_{max}$ (ng/mL)	Day 7	6	6	6
N		273.50 (44.00)	360.33 (64.86)	349.33 (47.56)
Mean (SD)		219, 325	247, 432	279, 401
Min, Max		16.09	18.00	13.61
CV%				
$t_{max}$ (h)	Day 1	6	6	6
N		1.50	2.50	1.75
Median		1.00, 3.00	1.00, 4.00	1.50, 3.00
Min, Max		1.77 (0.70)	2.50 (1.05)	1.93 (0.59)
Mean (SD)		39.52	41.95	30.46
CV%				
$t_{max}$ (h)	Day 7	6	6	6
N		2.50	1.50	2.00
Median		1.50, 3.00	1.00, 3.00	1.50, 3.03
Min, Max		2.42 (0.66)	1.83 (0.98)	2.26 (0.62)
Mean (SD)		27.50	53.63	27.52
CV%				
$AUC_{0-24}$ (h*ng/mL)	Day 1	6	6	6
N		1857.8 (51.9)	2001.2 (400.0)	2127.2 (438.9)
Mean (SD)		1777.4, 1906.3	1412.4, 2523.8	1503.8, 2612.2
Min, Max		2.80	19.99	20.63
CV%				
$AUC_{0-24}$ (h*ng/mL)	Day 7	6	6	6
N		3062.5 (512.2)	3841.1 (827.6)	4516.9 (1022.1)
Mean (SD)		2614.4, 3972.0	2570.6, 4903.3	2934.5, 5958.0
Min, Max		16.72	21.55	22.63
CV%				

(continued)

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose		
		Young	Intermediate Age	Elderly
		12 mg	12 mg	12 mg
$AUC_{last}$ (h*ng/mL)	Day 1			
N		6	6	6
Mean (SD)		1835.9 (51.68)	1975.8 (395.6)	2099.4 (431.6)
Min, Max		1751.6, 1880.6	1392.7, 2490.7	1488.2, 2580.7
CV%		2.82	20.02	20.56
$AUC_{last}$ (h*ng/mL)	Day 7			
N		6	6	6
Mean (SD)		5606.7 (1836.0)	7466.8 (1967.8)	9143.7 (2972.6)
Min, Max		3680.6, 8885.6	4807.4, 10204.7	5123.6, 13850.9
CV%		32.75	26.35	32.51
$AUC_{mf}$ (h*ng/mL)	Day 7			
N		6	6	6
Mean (SD)		5933.1 (2133.5)	8104.1 (2270.0)	10304.0 (3904.9)
Min, Max		3824.4, 9805.7	5116.2, 11320.1	5412.3, 16659.1
CV%		35.96	28.01	37.90
$AUC_{%extrap}$ (%)	Day 7			
N		6	6	6
Mean (SD)		4.77 (2.72)	7.51 (1.85)	9.70 (5.93)
Min, Max		1.38, 9.38	5.08, 9.85	2.92, 16.86
CV%		57.15	24.61	61.20
$CL/F$ (L/h)	Day 1			
N		6	6	6
Mean (SD)		6.46 (0.18)	6.22 (1.34)	5.87 (1.32)
Min, Max		6.29, 6.75	4.75, 8.50	4.59, 7.98
CV%		2.84	21.62	22.51
$CL/F$ (L/h)	Day 7			
N		6	6	6
Mean (SD)		4.00 (0.60)	3.27 (0.80)	2.79 (0.72)
Min, Max		3.02, 4.59	2.45, 4.67	2.01, 4.09
CV%		14.98	24.62	25.75
$V_z/F$ (L)	Day 1			
N		6	6	6
Mean (SD)		196.87 (93.73)	201.33 (70.22)	164.01 (47.57)
Min, Max		112.49, 376.03	147.36, 341.22	114.66, 244.64
CV%		47.61	34.88	29.00
$V_{ss}/F$ (L)	Day 7			
N		6	6	6

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose		
		Young	Intermediate Age	Elderly
		12 mg	12 mg	12 mg
Mean (SD)		115.44 (14.55)	121.40 (23.97)	110.72 (31.49)
Min, Max		99.20, 133.36	96.43, 159.29	77.31, 160.37
CV%		12.60	19.74	28.45
$t_{1/2}$ (h)	Day 7			
N		6	6	6
Mean (SD)		20.49 (4.55)	26.09 (2.78)	28.28 (8.11)
Min, Max		15.18, 27.02	22.18, 29.85	18.93, 37.88
CV%		22.23	10.67	28.66
Dose-Normalized $C_{max}$ (ng/mL/mg)	Day 1			
N		6	6	6
Mean (SD)		19.92 (2.78)	21.42 (4.22)	29.11 (3.96)
Min, Max		17.42, 24.50	15.83, 28.75	23.25, 33.42
CV%		13.96	19.71	13.61
Dose-Normalized $C_{max}$ (ng/mL/mg)	Day 7			
N		6	6	6
Mean (SD)		22.79 (3.67)	30.03 (5.40)	29.11 (3.96)
Min, Max		18.25, 27.08	20.58, 36.00	23.25, 33.42
CV%		16.09	18.00	13.61
Dose-Normalized $AUC_{inf}$ (h*ng/mL/mg)	Day 7			
N		6	6	6
Mean (SD)		494.43 (177.79)	675.34 (189.17)	858.67 (325.40)
Min, Max		318.70, 817.14	426.35, 943.34	451.02, 1388.25
CV%		35.96	28.01	37.90
Observed Accumulation Index				
N		6	6	6
Mean (SD)		1.65 (0.31)	1.92 (0.25)	2.12 (0.22)
Min, Max		1.42, 2.23	1.73, 2.41	1.85, 2.34
CV%		18.56	12.91	10.34

(continued)

Steady State Accumulation Index				
N		6	6	6
Mean (SD)		1.01 (0.08)	1.13 (0.16)	1.30 (0.11)
Min, Max		0.92, 1.14	0.93, 1.41	1.13, 1.46
CV%		7.51	14.05	8.74
Day 7/Day 1		6	6	6

PK Parameter Statistic	Study Day	CRYSTALLINE FORM II OF COMPOUND (I) Dose		
		Young	Intermediate Age	Elderly
		12 mg	12 mg	12 mg
$C_{max}$ Ratio		1.15 (0.18)	1.41 (0.17)	1.53 (0.13)
N		0.95, 1.42	1.25, 1.68	1.34, 1.71
Mean (SD)		15.95	11.74	8.26
Min, Max				
CV%				

**Table 14. Summary of Plasma for Crystalline Form II of Compound (I)  
Pharmacokinetic Parameters by Gender and Day for All Crystalline Form II Of  
Compound (I) Doses Pooled**

PK Parameter Statistic	Study Day	Gender	
		Males	Females
$C_{max}$ (ng/mL)	Day 1		
N		18	18
Mean (SD)		261.56 (84.47)	269.61 (66.40)
Min, Max		168, 476	164, 375
CV%		32.30	24.63
$C_{max}$ (ng/mL)	Day 7		
N		16	16
Mean (SD)		355.31 (126.67)	372.94 (107.85)
Min, Max		210, 628	191, 586
CV%		35.65	28.92
$t_{max}$ (h)	Day 1		
N		18	18
Median		2.00	2.00
Min, Max		1.00, 3.03	1.00, 4.00
Mean (SD)		2.12 (0.72)	2.26 (0.81)
CV%		34.06	35.78
$t_{max}$ (h)	Day 7		
N		16	16
Median		1.75	2.00
Min, Max		1.00, 3.00	1.00, 3.03
Mean (SD)		1.88 (0.65)	2.31 (0.68)
CV%		34.43	29.48
$AUC_{0-24}$ (h*ng/mL)	Day 1		
N		18	18
Mean (SD)		2214.7 (788.1)	2343.3 (761.6)
Min, Max		1412.4, 4265.5	1322.6, 4099.3
CV%		35.58	32.50

PK Parameter Statistic	Study Day	Gender	
		Males	Females
$AUC_{0-24}$ (h*ng/mL)	Day 7		
N		16	16
Mean (SD)		3749.3 (1388.3)	4472.9 (1570.0)
Min, Max		2543.3, 7364.8	2107.3, 7880.0
CV%		37.03	35.10

(continued)

$AUC_{last}$ (h*ng/mL)	Day 1		
N		18	18
Mean (SD)		2190.3 (778.5)	2312.5 (749.4)
Min, Max		1392.7, 4206.0	1308.2, 4054.0
CV%		35.54	32.41
$AUC_{last}$ (h*ng/mL)	Day 7		
N		16	16
Mean (SD)		6353.6 (2692.0)	8582.7 (3497.9)
Min, Max		2534.3, 12849.0	3103.4, 14653.4
CV%		42.37	40.76
$AUC_{inf}$ (h*ng/mL)	Day 7		
N		16	16
Mean (SD)		6767.1 (2890.3)	9319.7 (4031.5)
Min, Max		3187.2, 13362.2	3201.4, 16659.1
CV%		42.71	43.26
$AUC_{\%extrap}$ (%)	Day 7		
N		16	16
Mean (SD)		6.06 (5.38)	6.76 (4.26)
Min, Max		0.93, 20.48	1.38, 16.86
CV%		88.81	62.97
$CL/F$ (L/h)	Day 1		
N		18	18
Mean (SD)		6.19 (1.15)	5.79 (0.84)
Min, Max		4.67, 8.50	4.59, 7.28
CV%		18.58	14.53
$CL/F$ (L/h)	Day 7		
N		16	16
Mean (SD)		3.74 (0.83)	3.19 (0.84)
Min, Max		2.72, 5.53	2.01, 4.89
CV%		22.09	26.17
$V_z/F$ (L)	Day 1		
N		18	18
Mean (SD)		147.17 (70.09)	181.51 (79.11)
		60.75, 341.22	83.53, 376.03

PK Parameter Statistic	Study Day	Gender	
		Males	Females
Min, Max		47.62	43.59
CV%			

(continued)

$V_{ss}/F$ (L)	Day 7		
N		16	16
Mean (SD)		107.79 (32.05)	105.19 (26.24)
Min, Max		49.78, 160.37	76.51, 182.39
CV%		29.73	24.95
$t_{1/2}$ (h)	Day 7		
N		16	16
Mean (SD)		20.57 (6.87)	23.84 (6.58)
Min, Max		10.93, 37.88	14.12, 37.50
CV%		33.42	27.61
Dose-Normalized $C_{max}$ (ng/mL/mg)	Day 1		
N		18	18
Mean (SD)		19.78 (3.09)	20.59 (2.86)
Min, Max		14.50, 26.75	16.50, 28.75
CV%		15.62	13.89
Dose-Normalized $C_{max}$ (ng/mL/mg)	Day 7		
N		16	16
Mean (SD)		26.21 (3.94)	27.98 (5.43)
Min, Max		19.67, 32.50	18.25, 36.00
CV%		15.04	19.42
Dose-Normalized $AUC_{inf}$ (h*ng/mL/mg)	Day 7		
N		16	16
Mean (SD)		512.15 (188.98)	698.59 (282.19)
Min, Max		230.76, 937.07	331.55, 1388.25
CV%		36.90	40.39
Observed Accumulation Index			
N		16	16
Mean (SD)		1.72 (0.28)	1.88 (0.37)
Min, Max		1.26, 2.34	1.36, 2.41
CV%		16.08	19.60

PK Parameter Statistic	Study Day	Gender	
		Males	Females
Steady State Accumulation Index			
N		16	16
Mean (SD)		1.15 (0.16)	1.10 (0.15)
Min, Max		0.93, 1.46	0.80, 1.41
CV%		13.91	13.88
Day 7/Day 1 C <sub>max</sub> Ratio			
N		16	16
Mean (SD)		1.36 (0.22)	1.38 (0.27)
Min, Max		1.06, 2.00	0.95, 1.83
CV%		16.44	19.28

Effects of Crystalline Form II of Compound (I) on arterial blood pressure are as follows. Briefly, the average 24 hour and daytime systolic blood pressure (SBP) and diastolic blood pressure (DBP) increased with all Crystalline Form II of Compound (I) doses compared to placebo, with the maximal increases in BP being observed by Day 4, except for the Crystalline Form II of Compound (I) 20 mg dose which increased further from Day 4 to Day 7. The effects of Crystalline Form II of Compound (I) on average nighttime SBP and DBP were smaller and less consistently observed. The effects of Crystalline Form II of Compound (I) on blood pressure appeared similar regardless of age subgroup. The effect of Crystalline Form II of Compound (I) 20 mg on SBP appeared slightly more pronounced in females compared to males on Day 7, however, overall, the BP effects of Crystalline Form II of Compound (I) appeared similar in males and females.

There was a high degree of variability in the plasma BDNF measurements in this study. However, there appeared to be a trend for an increase on average 7-day BDNF values compared to placebo at the higher two doses of 16 and 20 mg.

#### **EXAMPLE 21: Antidepressant Effect in Subjects with MDD and Suicidal Ideation**

A randomized, double-blind, placebo-controlled, sequential parallel study is conducted to evaluate the antidepressant effect of crystalline Form II of Compound (I) after 7 days of treatment.

Following a screening period of up to 14 days, approximately 135 subjects (male and female, 18 to 70 years of age with MDD and experiencing a severe depressive episode with recent active suicidal ideation despite current, stable treatment with an SSRI or SNRI) are randomly assigned to one of three treatment sequences with unequal  
5 distribution as shown in Table 15.

**Table 15. Treatment Sequences**

<b>Treatment Sequence</b>	<b>Period 1 (7 days)</b>	<b>Period 2 (28 days)</b>
1 (N = 27)	Crystalline Form II of Compound (I) 12 mg on Day 0 followed by 8 mg on Days 1-6	Crystalline Form II of Compound (I) 8 mg on Days 7-27 followed by placebo on Days 28-34
2 (N = 54)	Placebo on Days 0-6	Crystalline Form II of Compound (I) (8 mg) on Days 7-34
3 (N = 54)	Placebo on Days 0-6	Placebo on Days 7-34

The randomized subjects participate in two treatment periods (Period 1 and Period 2) and a follow-up safety assessment. In Period 1, the subjects receive their first dose of study drug and undergo safety, PK and efficacy assessments on Day 0. On Day 4 ( $\pm 1$   
10 day), the subjects undergo pre-dose safety, PK and efficacy assessments, followed by dosing procedures and post-dose safety and efficacy assessments. On Day 7 ( $\pm 1$  day), the subjects undergo safety, PK and efficacy assessments, thereby completing Period 1. Upon completion of Period 1 study assessments, the subjects receive their first dose of Period 2 study drug on that same day (Day 7  $\pm 1$  day) and undergo post-dose safety and efficacy  
15 assessments. On Days 11, 14, 21 and 28 ( $\pm 1$  day), the subjects undergo pre-dose safety, PK and efficacy assessments, followed by dosing procedures and post-dose safety and efficacy assessments. On Day 35 ( $\pm 1$  day), the subjects undergo safety, PK and efficacy assessments, thereby concluding Period 2. On Day 49 ( $\pm 2$  days), the subjects undergo follow-up safety assessments.

20 Efficacy is assessed based on the following:

- MDD: HAM-D17, Clinical Global Impression-Severity (CGI-S), and Clinical Global Impression-Improvement (CGI-I);
- suicidality: Beck Scale for Suicide Ideation (BSSI);
- biomarker: plasma BDNF; and
- 25 • genotype: BDNF polymorphism.

Safety is assessed based on the following:

- vital signs (semi-recumbent blood pressure, heart rate and oral body temperature);
- blood pressure measurements performed in triplicate (within 5 minutes) at each time point and the average determined;
- 12-lead electrocardiogram (ECG);
- clinical laboratory testing (hematology, clinical chemistry and urinalysis)
- physical examinations
- adverse event (AE) assessments
- Columbia-Suicide Severity Rating Scale (C-SSRS, “Baseline/Screening” version and “Since Last Visit” version)
- Brief Psychiatric Rating Scale (BPRS)
- concomitant medication assessments

Safety, PK and efficacy data (*e.g.*, changes from baseline in HAM-D17, duration of HAM-D17 response, time to meet HAM-D17 responder criterion ( $\geq 50\%$  decrease from baseline), BSSI scores and changes from baseline, BDNF levels and changes from baseline, CGI-S and CGI-I scores, blood pressure and changes from baseline, and trough concentrations of crystalline Form II of Compound (I)) are assessed to evaluate the antidepressant effect of crystalline Form II of Compound (I), as well as its safety and tolerability as an adjunctive treatment in subjects with MDD, its effect as adjunctive therapy on suicidal ideation, its effects on specific depressive symptoms, its effects on plasma BDNF, the need for a loading dose, the relationship between onset of antidepressant effect and plasma BDNF in subjects with MDD, the relationship between baseline symptoms and rate and magnitude of response in subjects with MDD, and differences of antidepressant effect in subjects with BDNF Val66Val vs. Val66Met polymorphism.

**EXAMPLE 22: Antidepressant Effect in Subjects with MDD**

A randomized, double-blind, placebo-controlled, sequential parallel study is conducted to evaluate the antidepressant effect of a single dose of crystalline Form II of Compound (I) after a single dose of treatment.

Following a screening period of up to 14 days, approximately 60 subjects (male and female, 18 to 65 years of age, diagnosed with MDD without psychotic features, and undergoing stable treatment with an SSRI or SNRI) are randomly assigned to one of three treatment sequences with unequal distribution as shown in Table 16.

5

**Table 16. Treatment Sequences**

<b>Treatment Sequence</b>	<b>Period 1 Study Drug</b>	<b>Period 2 Study Drug</b>
1 (N = 10)	Crystalline Form II of Compound (I) (20 mg) on Day 0	Crystalline Form II of Compound (I) (20 mg) on Day 3
2 (N = 25)	Placebo on Day 0	Crystalline Form II of Compound (I) (20 mg) on Day 3
3 (N = 25)	Placebo on Day 0	Placebo on Day 3

The randomized subjects participate in two treatment periods (Period 1 and Period 2) and follow-up safety and efficacy assessments. On Day 0 of Period 1, the subjects undergo baseline assessments, receive their Period 1 study drug, and undergo post-dose safety, PK and efficacy assessments. On Day 1 of Period 1, the subjects undergo  
 10 additional safety, PK and efficacy assessments, after which the subjects are re-randomized into Period 2 treatment sequences. On Day 3 of Period 2, the subjects undergo baseline assessments, receive their Period 2 study drug, and undergo post-dose safety, PK and efficacy assessments. On Day 4 of Period 2, the subjects undergo additional safety, PK and efficacy assessments. Following Period 2, the subjects undergo follow-up safety and  
 15 efficacy assessments on Day 6, Day 10 ( $\pm 1$  day), and Day 17 ( $\pm 1$  day).

Safety, PK and efficacy data are assessed to evaluate the antidepressant effect of a single dose of crystalline Form II of Compound (I), as well as its safety and tolerability as an adjunctive treatment in subjects with MDD, its effects on specific depressive  
 20 symptoms, the relationship between antidepressant effect and plasma BDNF in subjects with MDD, the relationship between baseline symptoms and rate and magnitude of response in subjects with MDD, and differences of antidepressant effect in subjects with BDNF Val66Val vs. Val66Met polymorphism.

**EXAMPLE 23: Study of the Hemodynamic Effects of Compound (I)**

To determine its systemic hemodynamic effects, Compound (I) was administered  
 25 as a single oral gavage dose to six (n=6) chronically telemetered rats (implantation at least 7 days prior to the dosing day) at doses of 0.3, 0.6, 1, 3, and 10 mg / kg and systemic

blood pressure and heart rate values were recorded (FIG. 11). The hemodynamic effects were compared to Dizocilpine (MK-801) at a dose of 0.2 mg / kg given intravenously (in 0.9% saline). In each animal, a single oral gavage dose of Compound (I) or vehicle (0.5% MC/0.02% SLS) was administered (volume: 5 mL/ kg). A 24-hour recording was performed prior to dosing (vehicle alone) and after each oral dose. In another set of studies, Compound (I) (1 mg / kg) was administered in combination with the  $\alpha$ 1-adrenergic receptor antagonist prazosin ( $\alpha$ 1-adrenergic receptor antagonist, 200  $\mu$ g / kg, IV bolus) to elucidate the underlying mechanism of hypertension. Data were analyzed and compared to baseline, with correction for 24-h predose vehicle control.

The studies in conscious telemetered rats demonstrate that Compound (I), when given orally, increased arterial blood pressure transiently, and in a dose-dependent manner between 0.3 – 1 mg / kg, and this effect plateaued at 1 - 10 mg / kg. Interestingly, the ED50 for blood pressure effects was similar to ED50 for the forced swim test. The magnitude of change in hemodynamics with Compound (I) was significantly less than that of MK-801, a broad NMDA receptor antagonist. In addition, Compound (I) modestly increased heart rate at 0.3 and 0.6 mg/kg doses, and increased HR dose proportionally between 1 and 10 mg/kg. Further, a strong correlation between locomotor activity level and change in heart rate was observed ( $R^2 = 0.67$ ). The changes in heart rate may be partially explained by the central nervous system excitatory effects of Compound (I). Similar dose-dependent movement effects (dose proportional between 1 and 10 mg/kg) were also observed in a study of the locomotor effects of Compound (I) in rats.

As shown in FIG. 12, the effects of a single oral dose of Compound (I) on systolic blood pressure, with or without prazosin, were also investigated. This study demonstrated that  $\alpha$ 1-adrenergic blockade provided protection from Compound (I) mediated increases in blood pressure. Alpha-adrenoceptors predominate in the innervation of the vascular smooth muscle causing vasoconstriction and increases in afterload. The sympathetic nervous system's adrenergic neurotransmitter, norepinephrine, produces its vascular effects by binding to these adrenoceptors in the vasculature. The administration of agents that blocked peripheral  $\alpha$ 1-adrenergic receptors, such as the prazosin quinazoline compounds, effectively blocked and can be used to treat the blood pressure effects of Compound (I). Further, in human studies of Compound (I) administered orally at doses up to 20 mg, the primary hemodynamic observation has been a transient increase in systolic

and diastolic blood pressure, with a maximum and minimum that tends to correlate with TMAX and redistribution time. Heart rates have not been observed to increase to any significant degree, either clinically or statistically. Blood pressure increases in the absence of heart rate increases suggests a vasoconstrictive effect of Compound (I) ( $\alpha_1$ -adrenergic activity) with limited effects on cardiac contractility or inotropy (little to no  $\beta_1$ -adrenergic effect). Increases in tone of the  $\alpha_1$ -adrenergic system in animals and humans, induced either directly with  $\alpha_1$ -agonists, or indirectly with drugs that increase norepinephrine activity, can be easily and safely counteracted by agents that block  $\alpha_1$ -adrenergic receptors in the vasculature, or  $\alpha_1$ -adrenergic blockers.

10 **EXAMPLE 24: Effect of Compound (I) and Ketamine on Behaviors in the Rat**  
**Forced Swim Test 24 Hours after Administration**

This study aimed to evaluate the duration of the antidepressant-like effects of Form II of Compound (I) using the forced swim test in male, Sprague-Dawley rats. Clinically effective antidepressant compounds reduce immobility time and increase swimming in rodents subjected to this test. Rats received Compound (I) (10 mg/kg, p.o.) or ketamine (10 mg/kg, i.p.) and 24 hours later were subjected to testing. As shown in FIG. 13, both ketamine and Form II of Compound (I) reduced immobility and increased swim time significantly in this behavioral procedure ( $p < 0.05$ ). The data represent mean  $\pm$  SEM. Similar to ketamine, these data are consistent with a long lasting antidepressant-like effect of Compound (I).

20 **EXAMPLE 25: Preparation of Tablets**

**Step 1: Preparation for Roller Compaction**

*Dispensing and de-lumping:*

1. The following components were dispensed into individual containers:
  - 25 a. Microcrystalline cellulose (Avicel® PH 102)
  - b. Lactose monohydrate (310)
  - c. Croscarmellose sodium (Ac-Di-Sol®)
  - d. Magnesium stearate (Hyqual®; vegetable source)
  - e. Crystalline Form II of Compound (I)
- 30 2. All ingredients except magnesium stearate were de-lumped through a 20 mesh hand screen. Magnesium stearate was de-lumped through a 60 mesh hand screen.

*Blending and Pre-Roller Compaction Lubrication:*

1. The ingredients were loaded in the v-blender in the following order:
  - a. Approximately half of microcrystalline cellulose
  - b. Crystalline Form II of Compound (I)
  - 5 c. Remaining half of microcrystalline cellulose
  - d. Materials were blended for 8.5 min at 30 rpm.
2. Croscarmellose sodium was then added, followed by lactose monohydrate and blended for 17 min at 30 rpm.
3. Magnesium stearate was added to the v-blender and blended for an  
10 additional 4.5 min at 30 rpm.
4. The blend was discharged in a low-density polyethylene (LDPE) bags.

**Step 2: Preparation of Roller Compaction Granules***Roller Compaction:*

1. The blend was manually loaded into the hopper. Feeder screw and roller  
15 compactor parameters were adjusted to yield a ribbon of the desired density. The parameters were recorded at a preset time interval.
2. The ribbons were collected in a double LDPE bags.
3. Ribbon Reconciliation. The ribbon thickness, weight, and length to yield ribbon density were evaluated.

20 *Milling*

The collected ribbons were milled at  $800 \pm 300$  rpm through a Comil fitted with a round hole (diameter, 0.039") and a round impeller.

*Post Roller Compaction Lubrication*

25 The granules were loaded in the v-blender followed by de-lumped magnesium stearate and blended for a 4.5 min at 30 rpm.

**Step 3: Preparation of Finished Tablets***Tableting*

The blend was manually charged into the hopper. The die fill amount and compression parameters (press speed, fill depth, pre-compression thickness setting,  
30 compression thickness setting, and force feeder speed) were adjusted to yield a tablet with the target weight and hardness. All finished tablets were collected in a double LDPE bags.


The composition of the tablets is provided in Table 17.

Table 17. Composition of Tablets

Component	Function	Strength (label claim)	
		4 mg	
		Qty per tablet (mg)	% w/w
Crystalline Form II of Compound (I)	Active Ingredient	4.0	5.0
Microcrystalline Cellulose (Avicel® PH 102)	Diluent	36.8	46.0
Lactose Monohydrate 310	Diluent	36.8	46.0
Croscarmellose Sodium (Ac-Di-Sol®)	Disintegrant	1.6	2.0
Magnesium Stearate (Hyqual®; vegetable source)	Lubricant	0.8	1.0
Total Weight per Tablet (mg)		80	100

## SPECIFIC EMBODIMENTS

1. A compound which is substantially pure crystalline Form II of Compound (I)



(I)

exhibiting at least one of:


- (i) an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, comprising peaks of 2-theta angle of about 5.9 and 8.8 degrees;
- (ii) an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, substantially as shown in FIG. 1A;
- (iii) an ultraviolet absorbance spectrum, obtained using methanol as diluent, substantially as shown in FIG. 2;
- (iv) an infrared spectrum substantially as shown in FIG. 3;

- (v) a proton nuclear magnetic resonance spectrum at about 600 MHz in CD<sub>3</sub>CN substantially as shown in FIG. 4;
- (vi) a <sup>13</sup>C nuclear magnetic resonance spectrum at about 150 MHz in CD<sub>3</sub>CN substantially as shown in FIG. 5;
- 5 (vii) a thermogravimetric analysis curve substantially as shown in FIG. 6; and
- (viii) a differential scanning calorimetry thermogram substantially as shown in FIG. 7.
2. The compound of embodiment 1, wherein the compound exhibits an X-ray powder diffraction pattern comprising peaks of 2-theta angles of about 5.9 and 8.8 degrees which correspond, respectively, to d-spacing at about 14.9 and 10.0 Angstroms (Å).
- 10 3. The compound of embodiment 1 or 2, wherein the compound exhibits an X-ray powder diffraction pattern substantially as shown in FIG. 1A.
- 15 4. The compound of any one of embodiments 2 and 3, wherein the compound further exhibits at least one of: an ultraviolet absorbance spectrum substantially as shown in FIG. 2, an infrared spectrum substantially as shown in FIG. 3, a proton nuclear magnetic resonance spectrum substantially as shown in FIG. 4, a <sup>13</sup>C nuclear magnetic resonance spectrum substantially as shown in FIG. 5, a thermogravimetric analysis curve substantially as shown in FIG. 6, and a differential scanning calorimetry thermogram substantially as shown in FIG. 7.
- 20 5. The compound of any one of embodiments 1-4, wherein the compound exhibits an ultraviolet absorbance spectrum substantially as shown in FIG. 2.
- 25 6. The compound of embodiment 5, wherein the ultraviolet absorbance spectrum comprises an absorbance maximum at about 236 ± 2 nm.
7. The compound of any one of embodiments 1-4, wherein the compound exhibits an infrared spectrum substantially as shown in FIG. 3.
- 30 8. The compound of any one of embodiments 1-4, wherein the compound exhibits a proton nuclear magnetic resonance spectrum substantially as shown in FIG. 4.

9. The compound of embodiment 8, wherein the proton nuclear magnetic resonance spectrum comprises peaks substantially as set out in Table 1.
10. The compound of any one of embodiments 1-4, wherein the compound exhibits a  $^{13}\text{C}$  nuclear magnetic resonance spectrum comprises peaks substantially as shown in FIG. 5.
- 5 11. The compound of embodiment 10, wherein the  $^{13}\text{C}$  nuclear magnetic resonance spectrum comprises peaks substantially as set out in Table 2.
12. The compound of any one of embodiments 1-4, wherein the compound exhibits a thermogravimetric analysis curve substantially as shown in FIG.
- 10 6.
13. The compound of embodiment 12, wherein the thermogravimetric analysis curve corresponds to a weight loss of about 0.16% up to about 250° C.
14. The compound of any one of embodiments 1-4, wherein the compound exhibits a differential scanning calorimetry thermogram substantially as
- 15 shown in FIG. 7.
15. The compound of embodiment 14, wherein the differential scanning calorimetry thermogram comprises an endothermic peak at a temperature of about 124° C.
16. A pharmaceutical composition comprising an effective amount of the
- 20 compound of any one of embodiments 1-15.
17. The pharmaceutical composition of embodiment 16, further comprising a pharmaceutically acceptable excipient.
18. A pharmaceutical composition of any one of embodiments 16 and 17, wherein the Compound (I) is in particulate form with an X90 particle size
- 25 of about 10  $\mu\text{m}$  or less.
19. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is about 8  $\mu\text{m}$  or less.
20. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is about 6  $\mu\text{m}$  or less.
- 30 21. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is about 5  $\mu\text{m}$  or less.

22. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is between about 1  $\mu\text{m}$  and about 10  $\mu\text{m}$ .
23. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is between about 2  $\mu\text{m}$  and about 8  $\mu\text{m}$ .
- 5 24. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is between about 3  $\mu\text{m}$  and about 6  $\mu\text{m}$ .
25. The pharmaceutical composition of embodiment 18, wherein the X90 particle size is between about 4  $\mu\text{m}$  and about 5  $\mu\text{m}$ .
26. The pharmaceutical composition of embodiment 18, wherein the X90  
10 particle size is about 4.5  $\mu\text{m}$ .
27. The pharmaceutical composition of any one of embodiments 16-26, wherein the pharmaceutical composition is formulated for oral administration.
28. The pharmaceutical composition of embodiment 27, wherein the  
15 pharmaceutical composition is in tablet or capsule form.
29. A method of treating a condition responsive to an NR2B antagonist, comprising administering to a patient in need thereof an effective amount of a compound of any one of embodiments 1-15 or a pharmaceutical composition of any one of embodiments 16-28.
- 20 30. The method of embodiment 29, wherein the condition is a depressive disorder.
31. The method of embodiment 29, wherein the condition is major depressive disorder.
32. The method of embodiment 29, wherein the condition is treatment-resistant  
25 major depressive disorder.
33. The method of any one of embodiments 29-32, wherein the compound is administered as an adjunct to a serotonin reuptake inhibitor or a serotonin and norepinephrine reuptake inhibitor.
34. The method of any one of embodiments 29-33, wherein the compound is  
30 administered intermittently.
35. The method of any one of embodiments 29-34, wherein the effective amount is between about 4 mg and about 60 mg daily or intermittently.

36. The method of any one of embodiments 29-35 where the compound is administered without food.
37. The method of any one of embodiments 29-35 wherein the compound is administered with food.
- 5 38. A pharmaceutical composition comprising particles of Compound (I)




(I)

with an X90 particle size of about 10  $\mu\text{m}$  or less.

- 10 39. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is about 8  $\mu\text{m}$  or less.
40. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is about 6  $\mu\text{m}$  or less.
41. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is about 5  $\mu\text{m}$  or less.
- 15 42. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is between about 1  $\mu\text{m}$  and about 10  $\mu\text{m}$ .
43. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is between about 2  $\mu\text{m}$  and about 8  $\mu\text{m}$ .
44. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is between about 3  $\mu\text{m}$  and about 6  $\mu\text{m}$ .
- 20 45. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is between about 4  $\mu\text{m}$  and about 5  $\mu\text{m}$ .
46. The pharmaceutical composition of embodiment 38, wherein the X90 particle size is about 4.5  $\mu\text{m}$ .
- 25 47. The pharmaceutical composition of any one of embodiments 38-46, further comprising a pharmaceutically acceptable excipient.

48. The pharmaceutical composition of any one of embodiments 38-47, wherein the pharmaceutical composition is formulated for oral administration.
49. The pharmaceutical composition of any one of embodiments 38-47, wherein the pharmaceutical composition is in the form of a tablet or capsule.
50. The pharmaceutical composition of any one of embodiments 38-49, wherein the Compound (I) is in crystalline Form II as characterized in any one of embodiments 1-15.
51. A method of increasing bioavailability of Compound (I)




(I)

comprising administering to a patient in need thereof an effective amount of a pharmaceutical composition of any one of embodiments 38-50.

52. The method of embodiment 51, wherein the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) with an X90 particle size of about 10  $\mu\text{m}$  or higher.
53. The method of embodiment 51, wherein the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) with an X90 particle size of about 11  $\mu\text{m}$  or higher.
54. The method of embodiment 51, wherein the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) with an X90 particle size of about 12  $\mu\text{m}$  or higher.
55. The method of embodiment 51, wherein the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) with an X90 particle size of about 13  $\mu\text{m}$  or higher.


56. The method of embodiment 51, wherein the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) with an X90 particle size of about 14  $\mu\text{m}$  or higher.
57. A method of treating a condition responsive to an NR2B antagonist, comprising administering to a patient in need thereof an effective amount of a pharmaceutical composition of any one of embodiments 38-50.
58. The method of embodiment 57, wherein the condition is a depressive disorder.
59. The method of embodiment 57, wherein the condition is major depressive disorder.
60. The method of embodiment 57, wherein the condition is treatment-resistant major depressive disorder.
61. The method of any one of embodiments 57-60, wherein the compound is administered as an adjunct to a serotonin reuptake inhibitor or a serotonin and norepinephrine reuptake inhibitor.
62. A method of treating suicidal ideation, comprising administering an effective amount of Compound (I)



(I)

- 20 to a patient who has, is suspected of having, or has been diagnosed with having suicidal ideation.
63. The method of embodiment 62, wherein the patient has been diagnosed with having suicidal ideation within about 4 weeks prior to administration of the compound.
64. The method of any one of embodiments 62 and 63, wherein the patient has been further diagnosed with having a depressive disorder.
65. The method of any one of embodiments 62 and 63, wherein the patient has been further diagnosed with having major depressive disorder.

66. The method of any one of embodiments 62 and 63, wherein the patient has been further diagnosed with having treatment-resistant major depressive disorder.
67. The method of any one of embodiments 62-66, wherein the Compound (I) is in crystalline Form II as characterized in any one of embodiments 1-15.
68. The method of any one of embodiments 62-67, wherein the Compound (I) is in a pharmaceutical composition as characterized in any one of embodiments 38-50.
69. The method of any one of embodiments 51-68, wherein the Compound (I) or pharmaceutical composition is administered without food.
70. The method of any one of embodiments 51-68, wherein the Compound (I) or pharmaceutical composition is administered with food.
71. A method of reducing absorption rate of Compound (I)



(I)

comprising administering to a patient in need thereof an effective amount of the Compound (I) with food, wherein the compound is administered either substantially concurrently with, or up to about 2 hours after, or up to about 30 minutes before administration of food.

72. The method of embodiment 71, wherein the method results in a lower  $C_{\max}$  or a higher  $T_{\max}$  compared to that of a method comprising administering the Compound (I) without food.
73. The method of any one of embodiments 71 and 72, wherein the Compound (I) is administered either substantially concurrently with, or up to about 90 minutes after, or up to about 15 minutes before administration of food.
74. The method of any one of embodiments 71 and 72, wherein the Compound (I) is administered either substantially concurrently with, or up to about 60 minutes after, or up to about 10 minutes before administration of food.

75. The method of any one of embodiments 71 and 72, wherein the Compound (I) is administered substantially concurrently with administration of food.

76. The method of any one of embodiments 71-75, wherein the compound is administered orally.


5 77. The method of any one of embodiments 71-76, wherein the compound is in crystalline Form II as characterized in any one of embodiments 1-15.

78. The method of any one of embodiments 71-77, wherein the compound is in a pharmaceutical composition as characterized in any one of embodiments 38-50.

10 79. The method of any one of embodiments 51-78, wherein the effective amount is between about 4 mg and about 60 mg daily or intermittently.

80. The method of any one of embodiments 29-37 and 51-79, further comprising monitoring the patient's blood pressure; and if hypertension is detected, administering an anti-hypertensive to the patient.


15 81. A method of preparing Compound (I)



(I)

comprising:

(i) reacting Compound (8)




(8)

with triflic anhydride to yield a triflate;


(ii) reacting the triflate with ammonia to yield Compound (9)

20




(iii) reacting the Compound (9) with 2-chloropyrimidine to yield Form I of Compound (I).

- 5      82. The method of embodiment 81, further comprising seeding Form I of Compound (I) with Form II of Compound (I).
83. The method of embodiment 81 or 82, further comprising reacting Compound (6)




- 10      with carbonyldiimidazole and Compound (7)




to yield Compound (8).

- 15      84. The method of embodiment 82, further comprising debenzylating Compound (5)



with hydrogen over palladium to yield Compound (6).

85. The method of embodiment 84, further comprising reducing Compound (4)



(4)

- 5 with chloro(1,5-cyclooctadiene)rhodium (I) dimer under hydrogen atmosphere to yield Compound (5).

86. The method of any one of embodiments 81-85, further comprising purifying the Compound (I).

- 10 87. The method of embodiment 86, wherein the purifying comprises slurring or recrystallization.

88. The method of embodiment 85, wherein the purifying comprises slurring followed by recrystallization.


15 It will be apparent to those in the art that specific embodiments of the disclosed subject matter may be directed to one or more of the above- and below-indicated embodiments in any combination.

20 While the invention has been disclosed in some detail by way of illustration and example for purposes of clarity of understanding, it is apparent to those in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. Therefore, the description and examples should not be construed as limiting the scope of the invention.

All references, publications, patents, and patent applications disclosed herein are hereby incorporated by reference in their entirety.

## WHAT IS CLAIMED IS:

1. A compound which is substantially pure crystalline Form II of Compound (I)




(I)

5 exhibiting at least one of:

- a. an X-ray powder diffraction pattern, obtained using copper K $\alpha$  radiation, comprising a peak of 2-theta angle of about 5.9 degree;
- b. an ultraviolet absorbance spectrum, obtained using methanol as diluent, substantially as shown in FIG. 2;
- c. an infrared spectrum substantially as shown in FIG. 3;
- d. a thermogravimetric analysis curve substantially as shown in FIG. 6; and
- e. a differential scanning calorimetry thermogram substantially as shown in FIG. 7.

2. The compound of claim 1, wherein the X-ray powder diffraction pattern comprises peaks of 2-theta angles of about 5.9 and 8.8 degrees.
3. The compound of any one of claims 1-2, wherein the X-ray powder diffraction pattern is substantially as shown in FIG. 1A.
4. The compound of any one of claims 1-3, wherein the thermogravimetric analysis curve of the compound is substantially as shown in FIG. 6.
5. The compound of any one of claims 1-3, wherein the thermogravimetric analysis curve of the compound corresponds to a weight loss of about 0.16% up to about 250° C.
6. The compound of any one of claims 1-5, wherein the differential scanning calorimetry thermogram of the compound comprises an endothermic peak at a temperature of about 124° C.

7. The compound of any one of claims 1-5, wherein the differential scanning calorimetry thermogram of the compound is substantially as shown in FIG. 7.
8. The compound of any one of claims 1-7, which has a purity of at least above 95%.
9. A pharmaceutical composition comprising an effective amount of the compound of any one of claims 1-8.
10. The pharmaceutical composition of claim 9, comprising an active ingredient consisting essentially of the compound of any one of claims 1-8.
11. The pharmaceutical composition of claim 9, wherein the compound is in particulate form with an X90 particle size of about 10  $\mu\text{m}$  or less.
12. The pharmaceutical composition of claim 9, wherein the X90 particle size is between about 3  $\mu\text{m}$  and about 6  $\mu\text{m}$ .
13. The pharmaceutical composition of claim 9, wherein the X90 particle size is about 4.5  $\mu\text{m}$ .
14. The pharmaceutical composition of claim 9, wherein the pharmaceutical composition is in tablet or capsule form.
15. A pharmaceutical composition comprising particles of Compound (I)




(I)

- with an X90 particle size of about 10  $\mu\text{m}$  or less.
16. The pharmaceutical composition of claim 15, wherein the X90 particle size is between about 3  $\mu\text{m}$  and about 6  $\mu\text{m}$ .
17. The pharmaceutical composition of claim 15, wherein the X90 particle size is about 4.5  $\mu\text{m}$ .
18. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition is in tablet or capsule form.

19. The pharmaceutical composition of any one of claims 9 - 18, further comprising an adjunct selected from (1) non-steroidal anti-inflammatory agents; (2) COX-2 inhibitors; (3) bradykinin B1 receptor antagonists; (4) sodium channel blockers and antagonists; (5) nitric oxide synthase (NOS) inhibitors; (6) glycine site antagonists; (7) potassium channel openers; (8) AMPA/kainate receptor antagonists; (9) calcium channel antagonists; (10) GABA-A receptor modulators (*e.g.*, a GABA-A receptor agonist); (11) matrix metalloprotease (MMP) inhibitors; (12) thrombolytic agents; (13) opioids such as morphine; (14) neutrophil inhibitory factor (NIF); (15) L-Dopa; (16) carbidopa; (17) levodopa/carbidopa; (18) dopamine agonists such as bromocriptine, pergolide, pramipexole, ropinirole; (19) anticholinergics; (20) amantadine; (21) carbidopa; (22) catechol O-methyltransferase (COMT) inhibitors such as entacapone and tolcapone; (23) Monoamine oxidase B (MAO-B) inhibitors; (24) opiate agonists or antagonists; (25) 5HT receptor agonists or antagonists; (26) NMDA receptor agonists or antagonists; (27) NK1 antagonists; (28) selective serotonin reuptake inhibitors (SSRI) and selective serotonin and norepinephrine reuptake inhibitors (SSNRI); (29) tricyclic antidepressant drugs, (30) norepinephrine modulators; (31) lithium; (32) valproate; (33) D-serine; (34) neurontin (gabapentin); (35) antitussives; (36) antihistamines (*e.g.*, first generation antihistamines); (37) decongestants; (38) expectorants; (39) mucolytics; (40) antipyretics; and (41) analgesics.
20. A method of treating a condition responsive to an NR2B antagonist, comprising administering to a patient in need thereof an effective amount of a compound of any one of claims 1-8 or a pharmaceutical composition of any one of claims 9-19.
21. The method of claim 20, wherein the condition is a depressive disorder.
22. The method of claim 20, wherein the condition is major depressive disorder.
23. The method of claim 20, wherein the condition is treatment-resistant major depressive disorder.

24. The method of any one of claims 20-23, wherein the compound is administered as an adjunct to a serotonin reuptake inhibitor or a serotonin and norepinephrine reuptake inhibitor.
25. The method of any one of claims 20-24, wherein the compound is administered intermittently.
26. The method of any one of claims 20-25, wherein the effective amount is between about 4 mg and about 60 mg daily or intermittently.
27. The method of any one of claims 20-26 where the compound is administered without food.
28. The method of any one of claims 20-26 wherein the compound is administered with food.
29. A method of treating suicidal ideation, comprising administering an effective amount of Compound (I)



(I)

to a patient who has, is suspected of having, or has been diagnosed with having suicidal ideation.

30. The method of claim 29, wherein the patient has been diagnosed with having suicidal ideation within about 4 weeks prior to administration of the compound.
31. The method of any one of claims 29-30, wherein the patient has been further diagnosed with having a depressive disorder.
32. The method of any one of claims 29-30, wherein the patient has been further diagnosed with having major depressive disorder.
33. The method of any one of claims 29-30, wherein the patient has been further diagnosed with having treatment-resistant major depressive disorder.

34. The method of any one of claims 29-33, wherein Compound (I) is in crystalline Form II as characterized in any one of claims 1-8.
35. The method of any one of claims 29-33, wherein Compound (I) is in a pharmaceutical composition as characterized in any one of claims 9-19.
- 5 36. The method of any one of claims 29-35, wherein the Compound (I) or pharmaceutical composition is administered without food.
37. The method of any one of claims 29-35, wherein the Compound (I) or pharmaceutical composition is administered with food.
38. The method of any one of claims 20 and 29, wherein the effective amount is between about 4 mg and about 60 mg daily or intermittently.
- 10 39. The method of any one of claims 20 and 29, wherein the effective amount is between about 4 mg and about 20 mg daily or intermittently.
40. A method of targeting N-methyl-D-aspartate (NMDA) receptor subunit 2B (GluN2B) expressed on a cell comprising administering to a patient in need thereof an effective amount of a compound of any one of claims 1-8 or a pharmaceutical composition of any one of claims 9-19.
- 15 41. The method of claim 40, wherein the patient is suffering from a condition selected from Parkinson's disease, neuropathic pain, bone and joint pain, repetitive motion pain, dental pain, cancer pain, myofascial pain, perioperative pain, chronic pain, dysmenorrhea, as well as pain associated with angina, and inflammatory pain of varied origins, headache, migraine and cluster headache, schizophrenia, stroke, traumatic brain injury, Alzheimer's disease, cerebral ischemia, amyotrophic lateral sclerosis, Huntington's disease, sensorineural hearing loss, tinnitus, glaucoma, neurological damage caused by epileptic seizures or by neurotoxin poisoning or by impairment of glucose and/or oxygen to the brain, vision loss caused by neurodegeneration of the visual pathway, Restless Leg Syndrome, multi-system atrophy, non-vascular headache, chronic, subchronic or acute cough, primary hyperalgesia, secondary hyperalgesia, primary allodynia, secondary allodynia, or other pain caused by central sensitization, dyskinesias, depressive disorders, trauma- and stressor-related
- 20
- 25
- 30


disorders, bipolar disorders with depressive features, anxiety disorders, and obsessive-compulsive and related disorders.

42. The method of claim 40, wherein the patient is suffering from major depressive disorder.

5 43. The method of claim 40, wherein the patient is suffering from treatment-resistant major depressive disorder.

44. The method of any one of claims 40-43, wherein the compound is administered as an adjunct to a serotonin reuptake inhibitor or a serotonin and norepinephrine reuptake inhibitor.

10 45. A method of increasing bioavailability of Compound (I)




(I)

comprising administering to a patient in need thereof an effective amount of a pharmaceutical composition of any one of claims 15-17.

15 46. The method of claim 45, wherein the method results in a higher area under the curve compared to that of a method comprising administering Compound (I) with an X90 particle size of more than about 10  $\mu\text{m}$ .

47. A method of reducing absorption rate of Compound (I)




(I)

20 comprising administering to a patient in need thereof an effective amount of the Compound (I) with food, wherein the compound is administered either substantially concurrently with, or up to about 2 hours after, or up to about 30 minutes before administration of food.

48. The method of claim 47, wherein the method results in a lower  $C_{\max}$  or a higher  $T_{\max}$  compared to that of a method comprising administering the Compound (I) without food.
49. The method of claim 47, wherein the Compound (I) is administered either  
5 substantially concurrently with, or up to about 90 minutes after, or up to about 15 minutes before administration of food.
50. The method of claim 47, wherein the Compound (I) is administered either substantially concurrently with, or up to about 60 minutes after, or up to about 10 minutes before administration of food.
- 10 51. The method of claim 47, wherein the Compound (I) is administered substantially concurrently with administration of food.
52. The method of any one of claims 47-51, wherein the compound is administered orally.
53. The method of any one of claims 47-51, wherein the compound is in  
15 crystalline Form II as characterized in any one of claims 1-8.
54. The method of any one of claims 47-51, wherein the compound is in a pharmaceutical composition as characterized in any one of claims 9-19.
55. The method of any one of claims 47-54, wherein the effective amount is between about 4 mg and about 60 mg daily or intermittently.
- 20 56. The method of any one of claims 20-55, further comprising administering an anti-hypertensive agent to the patient.
57. The method of claim 56, wherein the anti-hypertensive agent is selected from an  $\alpha_1$ -adrenoreceptor antagonist, an  $\alpha_2$ -adrenoreceptor antagonist, a calcium channel blocker, diuretics, a angiotensin converting enzyme (ACE)  
25 inhibitor, and a combination thereof.
58. The method of any one of claims 56-57, wherein the anti-hypertensive agent is administered within 30 minutes of administering Compound (I).
59. The method of any one of claims 56-58, wherein the anti-hypertensive is selected from doxazosin, prazosin, terazosin, indoramin, clonidine,  
30 guanabenz, and guanoxabenz.


60. The method of any one of claims 56-59, wherein the active ingredient of said anti-hypertensive agent consists essentially of an  $\alpha_1$ -adrenoreceptor antagonist.
61. The method of any one of claims 57-60, wherein the  $\alpha_1$ -adrenoreceptor antagonist is prazosin.
62. A method of preparing Compound (I)



(I)

comprising:


- a. reacting Compound (8)



(8)

with triflic anhydride to yield a triflate;

- b. reacting the triflate with ammonia to yield Compound (9)



(9)

- c. reacting the Compound (9) with 2-chloropyrimidine to yield crude Compound (I);

- d. subjecting said crude Compound (I) to purification steps, said steps comprising dissolving the crude Compound (I) in ethyl acetate at a temperature of above 35 °C, replacing the solvent of ethyl acetate partially or completely with heptane; and
- 5 e. collecting purified Form I of Compound (I).
63. The method of claim 62, further comprising heating heptane to a temperature above 50 °C and cooling heptane to room temperature.
64. A method of preparing Form II of Compound (I) comprising suspending purified Form I of Compound (I) in water for a sufficient period of time.
- 10 65. A method of preparing Form II of compound (I) comprising heating Form I of Compound (I) in solid state to a temperature above 50 °C for a sufficient period of time.
66. The method of claim 65, wherein the temperature is above about 100 °C.




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3

Solvent: CD<sub>3</sub>CN




FIG. 4

Solvent: CD<sub>3</sub>CN




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10A




FIG. 10B




FIG. 11




FIG. 12




FIG. 13

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/066741

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(8) - C07D 401/12 (2016.01) CPC - C07D 401/12 (2016.02) According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC(8) - A01N 43/40; A61K 31/506; A61P 25/16; C07D 211/74, 211/82, 401/06, 401/12 (2016.01) CPC - A01N 43/40; A61K 31/506; C07D 211/74, 211/82, 401/06, 401/12 (2016.02)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched USPC - 546/2, 236, 339; 556/12, 14; IPC(8) - A01N 43/40; A61K 31/506; A61P 25/16; C07D 211/74, 211/82, 401/06, 401/12; CPC - A01N 43/40; A61K 31/506; C07D 211/74, 211/82, 401/06, 401/12 (keyword delimited)		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Orbit, Google Patents, Google, PubChem Search terms used: 4-methylbenzyl (3S, 4R)-3-fluoro-4-[(pyrimidin-2-ylamino) methyl] piperidine-1-carboxylate, MK-0657, CERC-301, crystal, crystalline		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 7,592,360 B2 (LIVERTON et al) 22 September 2009 (22.09.2009) entire document	15-18
Y	US 2012/0289465 A1 (LEE et al) 15 November 2012 (15.11.2012) entire document	16-18
A	EP 1 648 882 B1 (MERCK & CO INC) 06 August 2008 (06.08.2008) entire document	1-3, 15-18
A	US 2008/0086006 A1 (NELSON) 10 April 2008 (10.04.2008) entire document	1-3, 15-18
A	WO 2013/156614 A1 (UCB PHARMA SA) 24 October 2013 (24.10.2013) entire document	1-3, 15-18
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed	
Date of the actual completion of the international search		Date of mailing of the international search report
29 March 2016		17 MAY 2016
Name and mailing address of the ISA/ Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, VA 22313-1450 Facsimile No. 571-273-8300		Authorized officer Blaine R. Copenheaver PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/066741

**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2.  Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3.  Claims Nos.: 4-14, 19-28, 34-44, 53-61  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

See Extra Sheet

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-3, 15-18

- Remark on Protest**
- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/066741

Continued from Box No. III Observations where unity of invention is lacking

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees need to be paid.

Group I: Claims 1-3 and 15-18 are drawn to a crystalline Form II of Compound (I) and compositions thereof.

Group II: Claims 29-33 are drawn to a method of treating suicidal ideation.

Group III: Claims 45-52 are drawn to methods of increasing bioavailability and reducing absorption rate.

Group IV: Claims 62-66 are drawn to methods of preparing Compound (I).

The inventions listed in Groups I, II, III, and IV do not relate to a single general inventive concept under PCT Rule 13.1, because under PCT Rule 13.2 they lack the same or corresponding special technical features for the following reasons:

The special technical features of Group I, a crystalline Form II of Compound (I) and compositions thereof, are not present in Groups II, III, and IV; the special technical features of Group II, a method of treating suicidal ideation, are not present in Groups I, III, and IV; the special technical features of Group III, methods of increasing bioavailability and reducing absorption rate, are not present in Groups I, II, and IV; and the special technical features of Group IV, methods of preparing Compound (I) are not present in groups I, II, and III.

The Groups I, II, III, and IV share the technical features of a Form II of Compound (I). However, these shared technical features do not represent a contribution over the prior art.

Specifically US 2008/0086006 A1 to Nelson teaches a Form II of Compound (I) (Para. [0116], See shown structure ).

The inventions listed in Groups I, II, III, and IV therefore lack unity under Rule 13 because they do not share a same or corresponding special technical feature.