

(19) **DANMARK**

(10) **DK/EP 2579901 T3**



Patent- og
Varemærkestyrelsen

(12) **Oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **A 61 K 39/02 (2006.01)** **C 07 K 14/205 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2019-11-04**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-08-07**
- (86) Europæisk ansøgning nr.: **11793174.1**
- (86) Europæisk indleveringsdag: **2011-06-09**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-04-17**
- (86) International ansøgning nr.: **US2011039832**
- (87) Internationalt publikationsnr.: **WO2011156619**
- (30) Prioritet: **2010-06-09 US 353039 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **The Board of Trustees of The University of Arkansas, 2404 North University Avenue, Little Rock, AR 72207, USA**
- (72) Opfinder: **PUMFORD, Neil, R., 3196 Ozark Acres Drive, Bentonville, AR 72712, USA**
LAYTON, Sherryl, 6111 Bainbridge Drive, Rogers, AR 72758, USA
KWON, Young, Min, 15537 Low Gap Road, West Fork, AR 72774, USA
HARGIS, Billy, 3811 Monte Vallo Manor, Fayetteville, AR 72764, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **VACCINE OG FREMGANGSMÅDER TIL REDUKTION AF CAMPYLOBACTERINFEKTION**
- (56) Fremdragne publikationer:
WO-A2-2006/046017
US-A1- 2007 128 183
US-A1- 2007 249 553
US-A1- 2010 112 002
WYSZYNSKA ET AL.: "Oral immunization of chickens with avirulent *Salmonella* vaccine strain carrying *C. jejuni* 72Dz/92 *cjaA* gene elicits specific humoral immune response associated with protection against challenge with wild-type *Campylobacter*.", *VACCINE.*, vol. 22, 2004, pages 1379-1389, XP002713521,
BUCKLEY A M ET AL.: "Evaluation of live-attenuated *Salmonella* vaccines expressing *Campylobacter* antigens for control of *C. jejuni* in poultry", *VACCINE, ELSEVIER LTD, GB*, vol. 28, no. 4, 22 January 2010 (2010-01-22), pages 1094-1105, XP026855279, ISSN: 0264-410X [retrieved on 2009-10-22]
DATABASE UniProt [Online] 1 November 1996 (1996-11-01), "SubName: Full=Precursor of peptidoglycane associated lipoprotein; Flags: Precursor;", XP002713522, retrieved from EBI accession no. UNIPROT:Q46123 Database accession no. Q46123
DATABASE UniProt [Online] 3 April 2007 (2007-04-03), "SubName: Full=CjaA protein;", XP002713523, retrieved from EBI accession no. UNIPROT:A3ZI25 Database accession no. A3ZI25

Fortsættes ...

LAYTON ET AL.: "Evaluation of Salmonella-Vectored Campylobacter Peptide Epitopes for Reduction of Campylobacter jejuni in Broiler Chickens.", CLIN VACCINE IMMUNOL., vol. 18, no. 3, 2011, pages 449-454, XP002713524,

SKERRA: 'Use of the tetracycline promoter for the tightly regulated production of a murine antibody fragment in Escherichia coli.' GENE, [Online] vol. 151, 1994, pages 131 - 135 Retrieved from the Internet:

<URL:<http://www.ncbi.nlm.nih.gov/pubmed/7828861>> [retrieved on 2011-11-01]

WYSZYNSKA ET AL.: 'Oral immunization of chickens with avirulent Salmonella vaccine strain carrying C. jejuni 72Dz/92 cjaA gene elicits specific humoral immune response associated with protection against challenge with wild-type Campylobacter.' VACCINE. vol. 22, 2004, pages 1379 - 1389

DATABASE UNIPROT 03 April 2007 'CjaA protein' Database accession no. A3ZI25

LAYTON ET AL.: 'Evaluation of Salmonella-Vectored Campylobacter Peptide Epitopes for Reduction of Campylobacter jejuni in Broiler Chickens.' CLIN VACCINE IMMUNOL. vol. 18, no. 3, 2011, pages 449 - 454

DESCRIPTION

SEQUENCE LISTING

[0001] A Sequence Listing accompanies this application. The Sequence Listing was filed with the application as a text file on June 9, 2011.

BACKGROUND

[0002] The repertoire of safe and cost-effective vaccines for generation of mucosal immunity against a variety of agents is limited. The leading bacterial cause of human gastrointestinal disease worldwide is *Campylobacter*. Bacterial gastroenteritis continues to pose a significant threat to the general public in the United States and abroad for the foreseeable future. Infections with *Campylobacter jejuni* occur more frequently than the more publicized infections from *Salmonella* species or *Escherichia coli* O157:H7. The actual burden of illness of *Campylobacter* gastroenteritis nationwide is 500-850 infections/100,000 persons per year.

[0003] Not only is *Campylobacter* the leading cause of bacterial gastroenteritis, but *C. jejuni* has been associated with the neuropathological disease Guillain-Barré Syndrome (GBS). This life-threatening disease may be an immune response to ganglioside-like structures on certain *C. jejuni* strains leading to an autoimmune response against nerve cells. Although GBS is the most important chronic sequelae, *Campylobacter* infection is also associated with a reactive arthritis, which may progress to Reiter's syndrome.

[0004] Vaccination against *Campylobacter* has had limited success using killed whole-cell or protein based vaccines. In addition, there are concerns regarding the development of Guillain-Barré syndrome or other sequelae from killed whole-cell vaccination. A successful vaccine would need to be cost-effective, safe, orally effective, and be produced in large quantities in a very short time-period. At the present time there is no such vaccine.

[0005] Wszyńska, Agnieszka, et al. (2004) *Vaccine* 22.11: 1379-1389, describes a study examining the general efficacy of avirulent *Salmonella* vaccine strains expressing *Campylobacter* antigen as a bivalent chicken vaccine prototype. The authors report that chickens orally immunized with *Salmonella* expressing *Campylobacter cjaA* gene developed serum IgG and mucosal IgA antibody responses against *Campylobacter* membrane proteins and *Salmonella* OMPs, as measured by an ELISA test. The authors further report that protection experiments showed that chicken immunization with avirulent *Salmonella* carrying *Campylobacter cjaA* gene greatly reduced the ability of heterologous wild type *C. jejuni* strain to colonize the bird cecum.

[0006] Buckley, Anthony M., et al. (2010) *Vaccine* 28.4: 1094-1105, describes a study in which

chickens were used to evaluate a *Salmonella enterica* serovar Typhimurium Δ aroA vaccine expressing the *C. jejuni* amino acid binding protein CjaA as a plasmid-borne fusion to the C-terminus of fragment C of tetanus toxin. The authors report that their data support the potential of CjaA- and Peb1A-based vaccines for control of *C. jejuni* in poultry.

SUMMARY

[0007] Vectors and methods for enhancing resistance to *Campylobacter* infection or enhancing the immune response to *Campylobacter* are provided herein.

[0008] In one aspect, the invention provides a vaccine vector the expression of an antigenic polypeptide consisting of the sequence of SEQ ID NO: 7 (cjaD; cj0113; GVSITVEGNCDEWGTDEYNQA), wherein the vector comprises a first polynucleotide sequence encoding the antigenic polypeptide, said first polynucleotide sequence not natively associated with the vector, wherein the vector is a bacterium. The only polypeptide sequence derived from a *C. jejuni* cjaD (cj0113) that is encoded by the polynucleotides of the bacterium is the antigenic polypeptide consisting of the sequence of SEQ ID NO: 7.

[0009] The bacterial vector also includes an immunostimulatory polypeptide not natively associated with the vector. The vaccine vector is capable of eliciting an immune response from a vaccinated subject that includes an IgA antibody response against *Campylobacter*. The response may be protective against challenge with *Campylobacter*.

[0010] In still another aspect, pharmaceutical compositions comprising the vectors provided herein in a pharmaceutically acceptable carrier are disclosed.

[0011] In yet another aspect, vectors for use in methods of inducing a therapeutic immune response directed to *Campylobacter* in a subject are provided. The methods include administering an effective amount of the vectors provided herein to a subject. In one embodiment, the therapeutic immune response includes an IgA antibody response and the response may be protective.

[0012] In still a further aspect, vectors for use in methods of enhancing resistance to *Campylobacter* infection are provided herein. The methods include administering an effective amount of the vectors disclosed herein to the subject such that the subject is resistant to infection after subsequent exposure to *Campylobacter*. In one embodiment the enhanced immune response includes an IgA antibody response and the response may be protective.

BRIEF DESCRIPTION OF THE FIGURES

[0013]

Figure 1 is a graph showing the standard curve for the quantitative PCR showing the number of cultivable *Campylobacter jejuni* on the x-axis as determined by conventional culturing and determination of colony forming units (CFU). The cycle number at which the fluorescence crossed the threshold in the quantitative PCR is shown on the y axis.

Figure 2 is a graph showing the log cfu/g ileum content in chicks vaccinated with saline, or Salmonella vectors expressing *Campylobacter* peptides cj0420 (SEQ ID NO: 9), cj0113 (SEQ ID NO: 7) or cj0982 (SEQ ID NO: 8) (10^8 cfu/chick), 11 days after challenge with *C. jejuni* (5×10^7 cfu/ml). The quantitative PCR was performed on total DNA extracted by conventional methods from mucosal linings of the ileum. The results are presented as mean \pm SEM (n=10). Groups with different lower case letters are significantly different ($p < 0.05$).

Figure 3 is a graph showing the log cfu/g ileum content in chicks vaccinated with saline, the Salmonella vector without an antigenic polypeptide insert or the Salmonella vector with the cj0113 insert (SEQ ID NO: 7) (10^8 cfu/chick), 11 days after challenge with *C. jejuni* (1×10^7 cfu/ml). The quantitative PCR was performed on total DNA extracted by conventional methods from mucosal linings of the ileum. The results are presented as mean \pm SEM (n=10) and the * indicates a significant difference ($P < 0.05$).

Figure 4 is a graph showing the relative levels of anti-*Campylobacter* IgG (S/P ratio) as measured by ELISA at days 21 and 32 after administration of the indicated vectors (10^8 cfu/chick). Groups with different lower case letters are significantly different ($P < 0.05$).

Figure 5 is a graph showing the relative levels of anti-*Campylobacter* sIgA (S/P ratio) in the mucosa of the ileum on day 32 post-vaccination with the indicated vectors (10^8 cfu/chick). Groups with different lower case letters are significantly different ($P < 0.05$).

Figure 6 is a graph showing the serum IgG levels (S/P ratio) at days 21 and 31 post vaccination and sIgA levels (S/P ratio) in the mucosa of the ileum at day 32 post vaccination with saline, the Salmonella vector without an insert or the vector with the antigenic polypeptide, cj0113 (SEQ ID NO: 7) (10^8 cfu/chick). A * indicates a significant difference from controls ($P < 0.05$).

Figure 7 is a graph showing the *C. jejuni* specific serum IgG antibody levels 10 days post vaccination by oral gavage with either *Bacillus subtilis* backbone strain (BSBB) or cj0113 (SEQ ID NO: 7) *Bacillus subtilis* vectored vaccine candidate at 10^8 cfu/chick. Data are presented as mean \pm SEM with the * indicating a significant difference ($P < 0.05$) from both controls.

Figure 8 is a graph showing the *C. jejuni* specific secretory IgA antibody levels 10 days post vaccination by oral gavage with either *Bacillus subtilis* backbone strain (BSBB) or cj0113 (SEQ ID NO: 7) *Bacillus subtilis* vectored vaccine candidate at 10^8 cfu/chick. The mucosa was collected in the ileum region. Data are presented as mean \pm SEM with the * indicating a significant difference ($P < 0.05$) from both controls.

Figure 9 is a graph showing the \log_{10} CFU of *C. jejuni* per gram of ileum content as

enumerated by quantitative PCR. Birds vaccinated with either *Bacillus subtilis* backbone strain (BSBB) or *Bacillus subtilis* vector expressing cj0113 (SEQ ID NO: 7), were challenged with *C. jejuni* at 1×10^8 cfu/chick then enumerated 10 days after by PCR. qPCR was performed on total DNA extracted by conventional methods from mucosal linings of the ileum. The results are presented as mean \log_{10} cfu/gram of ileum content \pm SEM (n=10) and the * indicates a significant difference ($P < 0.05$) from control.

Figure 10 is a graph showing the \log_{10} CFU of *C. jejuni* per gram of turkey ileum content as enumerated by quantitative PCR. Turkeys vaccinated with either backbone strain or cj0113 (SEQ ID NO: 7) *Salmonella* vectored vaccine, were challenged with *C. coli* at 1×10^8 cfu/chick then enumerated 12 days after by PCR. qPCR was performed on total DNA extracted by conventional methods from mucosal linings of the ileum. The results are presented as mean \log_{10} cfu/gram of ileum content \pm SEM (n=10) and the * indicates a significant difference ($P < 0.05$) from control.

DETAILED DESCRIPTION

[0014] Vaccine vectors that elicit mucosal, humoral, and cell-mediated immune responses against multiple serovars of *Campylobacter* offer a promising approach to limit *Campylobacter* gastroenteritis. This project utilizes a novel approach in the development of vaccines by inserting polynucleotide sequences encoding non-native linear epitopes (antigenic polypeptides). The antigenic polypeptides may be used in combination with an immunostimulatory polypeptide such as CD154 (CD40L) or HMGB1 (high mobility group box 1) in the vaccine vector. The antigenic polypeptide and the immunostimulatory polypeptide are not polypeptides found natively associated with the vector. The epitope or antigenic polypeptide and the immunostimulatory polypeptide may be expressed on the surface of recombinant vectors. The vectors are bacterial. The vectors may be live, live and attenuated, or killed prior to administration. Substantial preliminary data, such as that shown in the Examples, demonstrates that *Salmonella* or *Bacillus* constructs expressing a foreign epitope are able to rapidly induce high titer epitope-specific antibodies *in vivo*. Furthermore, co-expression of surface CD154 or HMGB1 effectively enhanced the antibody response against the foreign epitope.

[0015] Recombinant DNA technologies enable relatively easy manipulation of many bacterial species. Some bacteria are mildly or non-pathogenic, but are capable of generating a robust immune response. These bacteria make attractive vaccine vectors for eliciting an immune response to a heterologous, non-native, or foreign antigen. Bacterial vaccine vectors may mimic the natural infection and produce robust and long lasting immunity. Vaccine vectors are often relatively inexpensive to produce and administer. In addition, such vectors can often carry more than one antigen and may provide protection against multiple infectious agents.

[0016] Polynucleotides encoding polypeptide antigens from any number of pathogenic organisms may be inserted into the vaccine vector and expressed to generate antigenic polypeptides. An antigenic polypeptide is a polypeptide that is capable of being specifically recognized by the adaptive immune system. An antigenic polypeptide includes any polypeptide that is immunogenic. The antigenic polypeptides include, but are not limited to, antigens that are pathogen-related, allergen-related, tumor-related or disease-related. Pathogens include viral, parasitic, fungal and bacterial pathogens as well as protein pathogens such as the prions.

[0017] The antigenic polypeptides may be full-length proteins or portions thereof. It is well established that immune system recognition of many proteins is based on a relatively small number of amino acids, often referred to as the epitope. Epitopes may be only 8-10 amino acids. Thus, the antigenic polypeptides described herein may be full-length proteins, 8 amino acid long epitopes or any portion between these extremes. In fact the antigenic polypeptide may include more than one epitope from a single pathogen or protein. Suitably the antigenic polypeptide is a polypeptide that is not natively associated with the vector. Not natively associated includes antigenic polypeptides that may also occur natively in the vector, but that are being expressed recombinantly as an epitope, are being expressed in combination with a different polypeptide as a fusion protein to allow for differential display and differential enhancement of the immune response as compared to the natively expressed polypeptide.

[0018] Multiple copies of the same epitope or multiple epitopes from different proteins may be included in the vaccine vector. It is envisioned that several epitopes or antigens from the same or different pathogens or diseases may be administered in combination in a single vaccine vector to generate an enhanced immune response against multiple antigens. Recombinant vaccine vectors may encode antigens from multiple pathogenic microorganisms, viruses or tumor associated antigens. Administration of vaccine vectors capable of expressing multiple antigens has the advantage of inducing immunity against two or more diseases at the same time.

[0019] The polynucleotides may be inserted into the chromosome of the vaccine vector or encoded on plasmids or other extrachromosomal DNA. Polynucleotides encoding epitopes may be expressed independently (i.e., operably linked to a promoter functional in the vector) or may be inserted into a vaccine vector polynucleotide (i.e., a native polynucleotide or a non-native polynucleotide) that is expressed in the vector. Suitably, the vaccine vector polynucleotide encodes a polypeptide expressed on the surface of the vaccine vector such as a transmembrane protein. The polynucleotide encoding the antigenic polypeptide may be inserted into the vaccine vector polynucleotide sequence in frame to allow expression of the antigenic polypeptide on the surface of the vector. For example, the polynucleotide encoding the antigenic polypeptide may be inserted in frame into a bacterial polynucleotide in a region encoding an external loop region of a transmembrane protein such that the vector polynucleotide sequence remains in frame. See the Examples below in which the antigenic polypeptides are inserted into an external loop of the *lamB* gene of the *Salmonella enteritidis* vector.

[0020] Alternatively, the polynucleotide encoding the antigenic polypeptide may be inserted into a secreted polypeptide. Those of skill in the art will appreciate that the polynucleotide encoding the antigenic polypeptide could be inserted in a wide variety of vaccine vector polynucleotides to provide expression and presentation of the antigenic polypeptide to the immune cells of a subject treated with the vaccine vector. In the Examples, several *Campylobacter* polynucleotides were inserted into the *lamB* coding sequence of *Salmonella enteritidis*. The resulting recombinant bacteria express the inserted antigenic polypeptides on the surface of the bacteria. The polynucleotides may be inserted in *CotB* of *Bacillus subtilis* such that the recombinant bacteria expressed the inserted antigenic polypeptides in spores or into *slp* for surface expression in vegetative bacteria.

[0021] Described herein are polynucleotides encoding full length *Campylobacter* proteins including *cjaD* (SEQ ID NO: 1), *cjaA* (SEQ ID NO: 2) and ACE393 (SEQ ID NO: 3) or an antigenic polypeptide of these proteins. In the Examples, antigenic polypeptides derived from the full-length proteins were used as follows: SEQ ID NO: 7 (a *cjaD* polypeptide called *cj0113*); SEQ ID NO: 8 (a *cjaA* polypeptide called *cj0982*); and SEQ ID NO: 9 (an ACE 393 polypeptide called *cj0420*). The polynucleotides used in the Examples are provided as SEQ ID NOs: 4-6, respectively. The polynucleotides used in the Examples had the antigenic polypeptides of SEQ ID NOs 7-9 separated by serine linkers and linked to CD154 amino acids 140-149 (three amino acids before, after and in between the antigenic polypeptide and the immunostimulatory polypeptide).

[0022] Suitably, the portion of the antigenic polypeptide inserted into the vaccine vector is immunogenic or antigenic. An immunogenic fragment is a peptide or polypeptide capable of eliciting a cellular or humoral immune response. Suitably, an antigenic polypeptide may be 20 or more amino acids, 15 or more amino acids, 10 or more amino acids or 8 or more amino acids of the full-length sequence. Suitably the immune response generated against the target pathogen is a protective immune response. A protective immune response is a response capable of blocking or reducing morbidity or mortality caused by subsequent infection with the target pathogen, namely *Campylobacter*.

[0023] One of skill in the art will appreciate that any of these polynucleotide sequences may be used in combination with any other antigenic polypeptide including polypeptides from other heterologous pathogens or organisms and may also be used in conjunction with polynucleotides encoding immunostimulatory polypeptides such as a polypeptide of CD154 or HMGB1 such as is described in International Application Nos.

[0024] WO2008036675 and WO/2011/091255.

[0025] Polynucleotides encoding immunostimulatory polypeptides that are homologous to proteins of the subject and capable of stimulating the immune system to respond to the foreign epitope may also be inserted into a vector. As described in more detail below, the vector may include a CD154 polypeptide that is capable of binding CD40 in the subject and stimulating the subject to respond to the vector and its associated foreign antigenic polypeptide. In addition, a

vector may include a HMGB1 polypeptide or a functional fragment thereof. As described above with regard to antigenic polypeptides, polynucleotides encoding these polypeptides may be inserted into the chromosome of the vector or maintained extrachromosomally. One of skill in the art will appreciate that these polynucleotides can be inserted in a variety of vector polynucleotides for expression in different parts of the vector or for secretion of the polypeptides.

[0026] The polynucleotide encoding an immunostimulatory polypeptide capable of enhancing the immune response to a non-native antigenic polypeptide may also encode the antigenic polypeptide. The polynucleotide encoding an immunostimulatory polypeptide may be linked to the polynucleotide encoding the antigenic polypeptide, such that in the vaccine vector the immunostimulatory polypeptide and the foreign antigenic polypeptide are present on the same polynucleotide. For example, the antigenic polypeptide and the immunostimulatory polypeptide may be portions of a fusion protein. In the Examples, a polynucleotide encoding a polypeptide of CD154 that is capable of binding to CD40 also encodes an antigenic polypeptide from *cjaD*, *cjaA* or ACE 393 of *Campylobacter*. See SEQ ID NOS: 10-12 in the attached sequence listing for some examples of potential polypeptide sequences and SEQ ID NOS: 4-6 for polynucleotide sequences which encode for optional serine linkers between the antigenic polypeptide, the immunostimulatory polypeptide and the host polypeptide.

[0027] In the Examples, the polynucleotide encoding the *Campylobacter* antigenic polypeptides and the polynucleotide encoding the immunostimulatory polypeptide are both inserted in the outer loop of the transmembrane *lamB* gene. Those of skill in the art will appreciate that vector polynucleotides encoding other transmembrane proteins may also be used. In addition, the antigenic polynucleotides may be extrachromosomal or secreted by the vector. In the Examples, the polynucleotide encoding the *Campylobacter* cj0113 antigen (SEQ ID NO: 7) and the immunostimulatory peptide HMGB1 (SEQ ID NO:20) were expressed from a plasmid carried by a *Bacillus* vector and expressed on the cell surface.

[0028] Suitably, the CD154 polypeptide is fewer than 50 amino acids long, more suitably fewer than 40, fewer than 30 or fewer than 20 amino acids in length. The polypeptide may be between 10 and 15 amino acids, between 10 and 20 amino acids or between 10 and 25 amino acids in length. The CD154 sequence and CD40 binding region are not highly conserved among various species. The CD154 sequences of chicken and human are provided in SEQ ID NO: 13 and SEQ ID NO: 14, respectively.

[0029] The CD40 binding regions of CD154 have been determined for a number of species, including human, chicken, duck, mouse and cattle and are shown in SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO:18, and SEQ ID NO:19, respectively. Although there is variability in the sequences in the CD40 binding region between species, cross-species binding of CD154 to CD40 has been reported. For example, the human CD154 polypeptide was able to enhance the immune response in chickens. Therefore, one may practice the invention using species specific CD154 polypeptides or a heterologous CD154 polypeptide.

[0030] The HMGB1 (High Mobility Group Box-1) protein was first identified as a DNA-binding protein critical for DNA structure and stability. It is a ubiquitously expressed nuclear protein that binds DNA with no sequence specificity. The protein is highly conserved and found in plants to mammals. The zebrafish, chicken and human HMGB1 amino acid sequences are provided in SEQ ID NO: 28, SEQ ID NO: 20 and SEQ ID NO: 27, respectively. The sequence throughout mammals is highly conserved with 98% amino acid identity and the amino acid changes are conservative. Thus an HMGB1 protein from one species can likely substitute for that from another species functionally. The full-length HMGB1 protein or a portion thereof may be used as the HMGB1 polypeptide in the vaccine vectors described herein. HMGB1 has two DNA binding regions termed A box as shown in SEQ ID NO: 21 and 22 and B box as shown in SEQ ID NO: 23 and 24. See Andersson and Tracey, *Annu. Rev. Immunol.* 2011, 29:139-162.

[0031] HMGB1 is a mediator of inflammation and serves as a signal of nuclear damage, such as from necrotic cells. HMGB1 can also be actively secreted by cells of the monocyte/macrophage lineage in a process requiring acetylation of the protein, translocation across the nucleus and secretion. Extracellular HMGB1 acts as a potent mediator of inflammation by signaling via the Receptor for Advanced Glycated End-products (RAGE) and via members of the Toll-like Receptor family (TLR), in particular TLR4. The RAGE binding activity has been identified and requires the polypeptide of SEQ ID NO: 25. TLR4 binding requires the cysteine at position 106 of SEQ ID NO: 20, which is found in the B box region of HMGB1.

[0032] The inflammatory activities of HMGB1 do not require the full-length protein and functional fragments have been identified. The B box has been shown to be sufficient to mediate the pro-inflammatory effects of HMGB1 and thus SEQ ID NO: 23 and 24 are HMGB1 polypeptides or functional fragments thereof within the context of the present invention. In addition, the RAGE binding site and the pro-inflammatory cytokine activity have been mapped to SEQ ID NO: 25 and SEQ ID NO: 26, respectively. Thus, these polypeptides are functional fragments of HMGB1 polypeptides in the context of the present invention.

[0033] Those of skill in the art are capable of identifying HMGB1 polypeptides and fragments thereof capable of stimulating pro-inflammatory cytokine activity, using methods such as those in International Publication No. WO02 092004. Suitably, the HMGB1 polypeptide includes the RAGE binding domain at amino acids 150-183 of SEQ ID NO:20 (SEQ ID NO: 25 or a homolog thereof) and the pro-inflammatory cytokine activity domain between amino acids 89-109 of SEQ ID NO: 20 (SEQ ID NO: 26 or a homolog thereof). In particular, HMGB1 polypeptides and functional fragments or homologs thereof include polypeptides identical to, or at least 99% identical, at least 98% identical, at least 95% identical, at least 90% identical, at least 85% identical, or at least 80% identical to the HMGB1 polypeptides of SEQ ID NOs: 20-28.

[0034] One of skill in the art will appreciate that the HMGB1 polypeptide could be used to enhance the immune response to more than one antigenic polypeptide present in a vector. The polypeptide from HMGB1 stimulates an immune response at least in part by activating dendritic cells and macrophages and thus stimulating production of IL-1, IL-6, IFN- γ and TNF-

α. Suitably, HMGB1 may be expressed on the surface of the vector.

[0035] At least a portion of the antigenic polypeptide and at least a portion of the HMGB1 polypeptide or another immunostimulatory polypeptide may be present on the surface of the vaccine vector. Present on the surface of the vaccine vector includes polypeptides that are comprised within a transmembrane protein, interacting with, covalently or chemically cross-linked to a transmembrane protein, a membrane lipid or membrane anchored carbohydrate. A polypeptide can be comprised within a transmembrane protein by having the amino acids comprising the polypeptide linked via a peptide bond to the N-terminus, C-terminus or anywhere within the transmembrane protein (i.e. inserted between two amino acids of the transmembrane protein or in place of one or more amino acids of the transmembrane protein (i.e. deletion-insertion). Suitably, the polypeptides may be inserted into an external loop of a transmembrane protein. Suitable transmembrane proteins are *cotB* and *lamB*, but those of skill in the art will appreciate many suitable transmembrane proteins are available.

[0036] Alternatively, the polypeptides may be covalently or chemically linked to proteins, lipids or carbohydrates in the membrane, through methods available to persons of skill in the art. For example, di-sulfide bonds or biotin - avidin cross-linking could be used to present the antigenic and HMGB1 polypeptides on the surface of a vaccine vector. Suitably, the antigenic polypeptide and the HMGB1 polypeptide are part of a fusion protein. The two polypeptides may be directly linked via a peptide bond or may be separated by a linker or a section of a third protein into which they are inserted.

[0037] In the Examples, some of the vectors have the *Campylobacter* antigenic polypeptides (cj0113, cj0420 and cj0982) and the immunostimulatory polypeptide (CD154 amino acids 140-149 or HMGB1 or a functional fragment thereof) encoded on the same polynucleotide (*lamB*) such that the sequences are in frame with each other and with the *Salmonella* polynucleotide in which they were inserted. In some aspects linkers may be added between the polynucleotide sequences encoding the antigenic polypeptide and the immunostimulatory polypeptide such that in the expressed polypeptide several amino acids separate the two polypeptides. The linker may be 3 nucleotides encoding a single amino acid, or may be much longer, e.g. 30 nucleotides encoding 10 or more amino acids. In the Examples a 9 nucleotide linker was used and encoded for three serine residues. Those of skill in the art will readily envision many other types of linkers that could be used.

[0038] In addition, the polynucleotides may be present in a single copy or in multiple copies. For example, three copies of the antigenic polypeptide and three copies of the immunostimulatory polypeptide may be found in the same external loop of a transmembrane protein or expressed within several different vector proteins. In alternative aspects the immunostimulatory polypeptide and the antigenic polypeptide may be encoded by distinct polynucleotides.

[0039] Potential vaccine vectors for use in the methods include, but are not limited to, *Bacillus*,

Salmonella (*Salmonella enteritidis*), Shigella, Escherichia (*E. coli*), Yersinia, Bordetella, Lactococcus, Lactobacillus, Streptococcus, Vibrio (*Vibrio cholerae*), and Listeria.

[0040] Suitably, the vaccine vector is a GRAS organism. The vaccine vector may be inactivated or killed such that it is not capable of replicating. Methods of inactivating or killing bacterial vaccine vectors are known to those of skill in the art and include, but are not limited to methods such as formalin inactivation, antibiotic- based inactivation, heat treatment and ethanol treatment. In some aspects the vaccine vector may be a liposome based vector.

[0041] Compositions comprising the vector and a pharmaceutically acceptable carrier are also provided. A pharmaceutically acceptable carrier is any carrier suitable for *in vivo* administration. The pharmaceutically acceptable carrier may include water, buffered solutions, glucose solutions or bacterial culture fluids. Additional components of the compositions may suitably include excipients such as stabilizers, preservatives, diluents, emulsifiers and lubricants. Examples of pharmaceutically acceptable carriers or diluents include stabilizers such as carbohydrates (e.g., sorbitol, mannitol, starch, sucrose, glucose, dextran), proteins such as albumin or casein, protein-containing agents such as bovine serum or skimmed milk and buffers (e.g., phosphate buffer). Especially when such stabilizers are added to the compositions, the composition is suitable for freeze-drying or spray-drying.

[0042] Methods of enhancing immune responses to *Campylobacter* in a subject by administering the vectors described herein are also provided. The vector may contain a HMGB1 polypeptide or a CD154 polypeptide capable of stimulating the immune response to the vector and the antigenic polypeptides described above. The vector is administered to a subject in an amount effective to enhance the immune response of the subject to the non-native antigenic polypeptides. Suitably the immune response to challenge with *Campylobacter* is enhanced.

[0043] Enhancing an immune response includes, but is not limited to enhancing antibody responses. Suitably the IgA response is enhanced, more suitably the secretory IgA response is enhanced after administration of the vector as compared to a control. The control may be the same subject prior to administration of the vector, a comparable subject administered a vector alone or a vector expressing an irrelevant or a non-*Campylobacter* antigenic polypeptide. The antibody response, suitably the IgA response, may be increased as much as two fold, three fold, four fold, five fold or more as compared to the response of a control subject. The enhanced immune response may also result in a reduction of the ability of *Campylobacter* to grow or replicate and colonize the subject after administration of the vectors described herein. Such a reduction may be tested by challenging a subject administered the vector with a *Campylobacter* infection and monitoring the ability of the bacteria to colonize and replicate, i.e. infect, the subject as compared to a control subject. The growth of *Campylobacter* in the subject may be reduced by 1 log, 2 logs, 3 logs, 4 logs, 5 logs or even more. The growth of *Campylobacter* in a subject administered the vector may be below the level of detection.

[0044] In addition, methods of enhancing resistance to *Campylobacter* infection are disclosed.

Briefly, the methods comprise administering to a subject the vectors described above comprising *Campylobacter* antigenic polypeptides in an amount effective to elicit an immune response. Enhancing resistance to *Campylobacter* infection includes but is not limited to reducing the incidence of *Campylobacter* infections, limiting the spread of *Campylobacter* infections from one host to another, reducing *Campylobacter* replication in the subject, invasion or spread within a single host, reduced morbidity associated with *Campylobacter* infections, and reduced duration of a *Campylobacter* infection.

[0045] Administration of the vector may prevent the subject from contracting *Campylobacter* or from exhibiting any outward signs of disease, such as gastroenteritis or GBS. Increased resistance to *Campylobacter* may also include increased antibody production, suitably IgA production. The antibody response, suitably the IgA response, may be increased as much as two fold, three fold, four fold, five fold or more as compared to the response of a control subject. The enhanced immune response may also result in a reduction of the ability of *Campylobacter* to grow or replicate and colonize the subject after administration of the vectors described herein. Such a reduction may be tested by challenging a subject administered the vector with a *Campylobacter* infection and monitoring the ability of the bacteria to colonize and replicate, i.e. infect, the subject as compared to a control subject. The growth of *Campylobacter* in the subject may be reduced by 1 log, 2 logs, 3 logs, 4 logs, 5 logs or even more. The growth of *Campylobacter* in a subject administered the vector may be below the level of detection.

[0046] The antigenic polypeptides for use in all the methods described herein are from *cjaD* as discussed above. The insertion of the antigenic polypeptides into the vector may be accomplished in a variety of ways known to those of skill in the art, including but not limited to the scarless site-directed mutation system described in International Patent Publication No. WO2008/036675. The vector may be a bacterium engineered to express *Campylobacter* antigenic polypeptides in conjunction with polynucleotides capable of enhancing the immune response as discussed above. In particular, a polypeptide of CD154 or HMGB1 may be expressed by the vector to enhance the immune response of the subject to the antigenic polypeptides. The vectors used in these methods may be attenuated or killed prior to administration or use in the methods.

[0047] The useful dosage to be administered will vary depending on the age, weight and species of the subject, the mode and route of administration and the type of pathogen against which an immune response is sought. The composition may be administered in any dose of vector sufficient to evoke an immune response. For bacterial vectors, it is envisioned that doses ranging from 10^3 to 10^{10} bacteria, from 10^4 to 10^9 bacteria, or from 10^5 to 10^7 bacteria are suitable. The composition may be administered only once or may be administered two or more times to increase the immune response. For example, the composition may be administered two or more times separated by one week, two weeks, or by three or more weeks. The bacteria vectors are suitably viable prior to administration, but in some aspects the bacteria vectors may be killed prior to administration. In some aspects the bacteria vectors may be able to replicate in the subject, while in other embodiments the bacteria vectors may

be attenuated and/or may not be capable of replicating in the subject.

[0048] For administration to animals or humans, the compositions may be administered by a variety of means including, but not limited to, intranasally, mucosally, by spraying, intradermally, parenterally, subcutaneously, orally, by aerosol or intramuscularly. Eye-drop administration or addition to drinking water or food are additionally suitable means of administration. For chickens, the compositions may be administered *in ovo*.

[0049] With regard to the methods, a subject includes, but is not limited to, a vertebrate, suitably a mammal, suitably a human, or birds, suitably poultry such as chickens or turkeys. Other animal models of infection may also be used. Enhancing an immune response includes, but is not limited to, inducing a therapeutic or prophylactic effect that is mediated by the immune system of the subject. For example, an immune response is enhanced if the subject is protected from subsequent infection with *Campylobacter*. Specifically, enhancing an immune response may include enhanced production of antibodies, such as demonstrated in Figures 4-8, enhanced class switching of antibody heavy chains, maturation of antigen presenting cells, stimulation of helper T cells, stimulation of cytolytic T cells or induction of T and B cell memory. In the Examples, an increase in the amount of secretory IgA was seen after administration of the vector and was correlated with protection from subsequent *Campylobacter* infection.

[0050] It is envisioned that several epitopes or antigens from the same or different pathogens may be administered in combination in a single vector to generate an enhanced immune response against multiple antigens and their associated pathogens. Recombinant vaccine vectors may encode antigens from multiple pathogenic microorganisms, viruses or tumor associated antigens. Administration of vaccine vectors capable of expressing multiple antigens has the advantage of inducing immunity against two or more diseases at the same time.

[0051] Heterologous polynucleotides encoding antigens can be inserted in the vaccine vector genome at any non-essential site or alternatively may be carried on a plasmid using methods well known in the art. One suitable site for insertion of polynucleotides is within external portions of transmembrane proteins or coupled to sequences which target the heterologous polynucleotide for secretory pathways. One example of a suitable transmembrane protein for insertion of polynucleotides is the *lamB* gene of *Salmonella*. Heterologous polynucleotides include, but are not limited to, polynucleotides encoding antigens selected from pathogenic microorganisms or viruses other than the vaccine vector, i.e., non-native polynucleotides encoding non-native polypeptides.

[0052] The following examples are meant only to be illustrative.

Examples

Attenuation of Salmonella Vaccine Candidate Strains

[0053] *Salmonella enteritidis* phage type 13A (*S. enteritidis*) was attenuated by introducing defined, irreversible deletion mutations in the *aroA* and/or *htrA* gene of the *S. enteritidis* genome as previously described (available as ATCC Deposit Nos: PTA-7871, PTA-7872 and PTA-7873). Briefly, the target gene sequence in the bacterial genome of *S. enteritidis* was replaced with the kanamycin-resistant (Km^R) gene sequence. This was performed using 3S-PCR and electroporation of the 3S-PCR products into electrocompetent *Salmonella* cells containing the pKD46 plasmid. The resulting cell mixture was plated on LB agar plates supplemented with Km to select for positive clones containing a Km^R gene. The Km^R gene was inserted into the genomic region containing the genes of interest (*aroA* or *htrA*) by flanking the Km^R gene with sequences homologous to the genes of interest. Once Km^R mutants were obtained, the deletion mutations were confirmed by PCR and DNA sequencing. All Km^R genes were removed before epitope insertion was started.

Construction of Recombinant Vaccine Candidates

[0054] Three potential candidate antigenic polypeptides were selected: Omp18/cjaD (cj0113), cjaA (cj0982) and ACE393 (cj0420). The polypeptides selected were as follows: cj0113 (GVSTVEGNCDEWGTDEYNQAWMTTSYAPTS; SEQ ID NO: 10), cj0982c (KDIVLDAEIGGVAKGKDGKEKWMTTTSYAPTS; SEQ ID NO: 11), and cj0420 (KVALGVAVPKDSNITSVEDLKDKTLLLLNKGTTADAWMTTSYAPTS; SEQ ID NO: 12), all inserts additionally contain a sequence for amino acids 140-149 of CD 154.

[0055] Recombinant *S. enteritidis* strains containing stable integrated copies of cj0113-CD154 (cj0113), cj0420-CD154 (cj0420) or cj0982c-CD154 (cj0982) were constructed using the method of Cox et al. Scarless and site-directed mutagenesis in *Salmonella enteritidis* chromosome. BMC Biotechnol 2007;7:59. Briefly, an I-SceI enzyme site along with a Km^R gene was introduced into Loop 9 of the *lamB* gene by design of a PCR product which had the I-SceI enzyme site and Km^R gene flanked by approximately 200-300 base pairs of DNA on each side, homologous to the up and downstream regions of Loop 9. Primers used are shown in Table I below. The PCR product was electroporated into electrocompetent attenuated *Salmonella* cells containing the pKD46 plasmid and the resulting cell mixture plated on LB agar plates supplemented with Km to select for positive clones now containing a Km^R gene. After the Sce-I/ Km mutation was made in Loop 9, this region was replaced by a codon-optimized foreign epitope DNA sequence (Burns DM, Beacham IR. Rare codons in *E. coli* and *S. typhimurium* signal sequences. FEBS Lett 1985;189(2):318-24.). This second 3S-PCR reaction produced the foreign epitope insert flanked by Loop 9 up and downstream regions, and the resulting PCR product was electroporated into electrocompetent SE13A containing the Sce-I/ Km mutation described above. Plasmid pBC-I-SceI was also electroporated into the cells along with the insert as the plasmid produces the I-SceI enzyme which recognizes and cleaves a sequence creating a gap at the I-SceI enzyme site in the Loop 9 region of the LamB gene

where the foreign epitope sequences inserted into the SE13A genome. The plasmid also carries with it a chloramphenicol (Cm) resistant gene (Cm^R) as the inserts that will replace the Km^R gene the mutations must have a new selection marker to counter-select against the previous I-SceI/Km mutation. After electroporation, cells were plated on LB agar plates containing 25 µg/mL Cm for the selection of positive mutants.

Table I: PCR Primers

Primer	Amplified Region	Primer Sequence (SEQ ID NO)
lam-up-f	Loop 9 up	5'TGTACAAGTGGACGCCAATC 3' (SEQ ID NO: 29)
lam-up-r	Loop 9 up	5'GTTATCGCCGTCTTTGATATAGCC3' (SEQ ID NO: 30)
lam-dn-f	Loop 9 dn	5'ATTTCCCGTTATGCCGCAGC3' (SEQ ID NO: 31)
lam-dn-r	Loop 9 dn	5'GTTAAACAGAGGGCGACGAG 3'(SEQ ID NO: 32)
Km-f	I-SceI/Km ^r gene	5'GCTATATCAAAGACGGCGATAAC TAACTATAACGGTCCCTAAGG'TAGCGA ATTTCGGGGATCCGTCGA 3'(SEQ ID NO: 33)
Km-r	I-SceI/Km ^r gene	5'GCTGCGGCATAACGGGAAA TGTAGGCTGGAGCTGCTTCG 3' (SEQ ID NO: 34)
Kan4f	Inside Km ^r gene	5'CAAAGCGCTCTGAAGTTCC 3' (SEQ ID NO: 35)
Kan4r	Inside Km ^r gene	5'GCGTGAGGGGATCTTGAAGT 3' (SEQ ID NO: 36)
lam 3f	Outer regions of loop 9	5'GCCATCTCGCTTGGTGATAA 3' (SEQ ID NO: 37)
lam 3r	Outer regions of loop 9	5'CGCTGGTATTTTGC GG TACA 3' (SEQ ID NO: 38)
Cj0113f	Insert with loop 9 up	5'TTCATCGGTACCCCAITCATCACAGTACCTTCAACGGTGATGCTAACACCGGAGGAGGAGT TATCGCCGTCTTTGATATAGCC3' (SEQ ID NO: 39)
Cj0113r	Insert with loop 9 down	5'ATGAATGGGGTACCGATGAATATAACCAGGCGTCTCTCTGGATGACCACCTCCTATGCC CCGACCTCCTCTCTCCATTTCCTGTTATGCCGAGC3' (SEQ ID NO: 40)
Cj0420f	Insert with loop 9 up	5'ATCTTTACCTTTTCGCAACACCACCGATTTCGCAATCCAGAACGATATCTTTGGAGGAGGAGT TATCGCCGTCTTTGATATAGCC3' (SEQ ID NO: 41)
Cj0420r	Insert with loop 9 down	5'GTGTTGCGAAAAGGTAAGATGGTAAAGAAAAATCCCTCTCTGGATGACCACCTCCTATGC GCCGACCTCCTCTCTCCATTTCCTGTTATGCCGAGC3' (SEQ ID NO: 42)
Cj0982c-	Insert with	

Primer	Amplified Region	Primer Sequence (SEQ ID NO)
f	loop 9 up	5'GGTITTTATCTTTCAGATCTTCAACGCTGGTGATGTTGCTATCTTTCGGAACCGCAACACCCA GCCCAACTTTGGAGGAGGAGTTATCGCCGTCCTTGATATAGCC3' (SEQ ID NO: 43)
Cj0982c- r	Insert with loop 9 down	5'AAGATCTGAAAGATAAAACCCCTGCTGCTGAACAAAGGTACCACCGCGGATGCGTCTCCTC CTGGATGACCACCTCCTATGCGCCGACCTCCTCCTCCATTCCCGTTATGCCGCAGC3' (SEQ ID NO: 44)

[0056] Once positive mutation/inserts were suspected, PCR and DNA sequencing were performed to confirm that the insertion sequences are present and correct.

Challenge with *Campylobacter jejuni*

[0057] Three wild-type isolates of *C. jejuni* from broiler chickens were individually grown to log-phase growth, combined, serially diluted and spread plated for conventional culture enumeration as previously described (Cole et al. Effect of aeration and storage temperature on *Campylobacter* concentrations in poultry semen. *Poult Sci* 2004;83:1734-8.). These were diluted to approximately 10^7 to 10^8 cfu/ml for challenge by oral gavage using spectrophotometric density and comparison to a previously-generated standard curve. Empirically determined cfu administered are reported for each of experiment involving challenge (see below).

Vaccination Study 1

[0058] In the first immunization study, 210 day-of-hatch broiler chicks were obtained from a local commercial hatchery and randomly assigned to one of four treatment groups: saline only (Negative control), or one of three vaccine candidate groups: cj0113, cj0420 or cj0982; n=50/pen. Each treatment group was housed in an individual floor pen on fresh pine litter and provided water and feed *ad libitum*. On day-of-hatch, all chicks in each treatment group were inoculated, via oral gavage, with 0.25mL of a solution containing approximately 10^8 cfu/mL of the appropriate treatment. On day 21 post-hatch, all birds in each treatment group were challenged with *C. jejuni*, via oral gavage, with 0.25mL of a solution containing 1×10^7 cfu/ml. On days 3, 11, 21 (prior to booster inoculation) and 32 post-hatch, 10-15 birds from each treatment group were humanely killed and their liver, spleen and cecal tonsils aseptically removed for the determination of organ invasion, colonization and clearance of the *Salmonella* vaccine vector strains. Also, on days 21 and 32 post-hatch, ileum sections were removed and processed for use in qRT-PCR and on day 32 a separate ileum sample was removed and diluted 1:5 in saline and was used to test for secretory immunoglobulin A (sIgA). In addition, blood samples were collected from 10 birds per treatment group and the serum was used for

determining antibody response on days 21 and 32 post-vaccination.

Vaccination Study 2

[0059] In experiment 2, 110 day-of-hatch broiler chicks were obtained from a local commercial hatchery and randomly assigned to one of two treatment groups: saline only (vehicle control) or *Salmonella* vaccine candidate, cj0113, (n=55/pen). Each treatment group was housed in an individual floor pen on fresh pine litter and provided water and feed *ad libitum*. On day-of-hatch, all chicks in each treatment group were inoculated via oral gavage with 0.25mL of a solution containing approximately 10^8 cfu/mL of the appropriate treatment. On day 21 post-hatch, all birds in each treatment group were challenged with *C. jejuni*, via oral gavage, with 0.25 mL of a solution containing 1×10^7 cfu/ml. On days 3, 11, 21 (prior to booster inoculation) and 32 post-hatch, 10-15 birds from each treatment group were humanely killed and their liver, spleen and cecal tonsils aseptically removed for the determination of organ invasion, colonization and clearance of the *Salmonella* vaccine vector strains. Also, on days 21 and 32 post-hatch ileum sections were removed and processed for use in qRT-PCR. In addition, blood samples were collected from 10 birds per treatment group and the serum was used for determining antibody response on days 21 and 32 post-hatch.

Vaccination Study 3

[0060] A third experiment was similar to vaccination experiment 2 (described above) except with the addition of a third group of *S. enteritidis* 13A *aroA/htrA* without the *Campylobacter* epitope (SE13A) as a control for the oral vaccination of the vector itself. All sample collections were the same as vaccination study 2 except on day 32 post-hatch an additional section of ileum was used to harvest the mucosal layer for sIgA as in experiment 1.

Measurement of Campylobacter Antibody Response

[0061] Serum collected from birds in both immunization studies was used in an ELISA to determine relative antibody responses. Briefly, individual wells of a 96-well plate were coated with *C. jejuni*. Antigen adhesion was allowed to proceed overnight at 4°C, the plates were then washed and blocked with Superblock (Pierce) for 1 hour at room temperature. Plates were then incubated for 2 hours with a 1:50 dilution of the previously collected sera. The plates were rinsed again followed by incubation with a Peroxidase-labeled anti-chicken IgG secondary antibody (Jackson Immunolaboratories) for an additional hour. After subsequent rinsing, the plates were developed using a peroxidase substrate kit (BD OptEIA, Fisher Scientific) and absorbances were read on a spectrophotometer at 450nm. Each plate contained a positive control and negative control where a pooled sample from vaccinated chicks and pre-immune chicken serum, respectively, replaced the serum from the treatment groups. The absorbance

obtained for the positive control, negative control and experimental samples were used to calculate Sample to Positive control ratios (S/P ratios) using the following calculation: (sample mean - negative control mean) / (positive control mean - negative control mean) (Brown et al. Detection of antibodies to *Mycoplasma gallisepticum* in egg yolk versus serum samples. J Clin Microbiol 1991;29(12):2901-3 and Davies et al. Evaluation of the use of pooled serum, pooled muscle tissue fluid (meat juice) and pooled faeces for monitoring pig herds for *Salmonella*. J Appl Microbiol 2003;95(5):1016-25.). The ELISA method used for detection of sIgA was similar to the above described assay for serum immunoglobulin except we used goat anti-chicken IgA conjugated with horseradish peroxidase (GenTex) in place of the anti-chicken IgG antibody conjugate.

DNA Isolation and Quantitative PCR for *C. jejuni*

[0062] Total DNA extraction from ileal samples was achieved using the QIAmp DNA Stool Mini Kit (Qiagen). The manufacturer's included protocol was modified slightly in the following ways: ileal contents were removed to include the mucosal layer and diluted 1:5(w/v) with ice cold PBS + .05% Tween 20; one ml of the slurry was added to 1ml of the included ASL Buffer in a 2.0ml microcentrifuge tube, vortexed and heated to 70°C for 5 minutes. Subsequently, the manufactures recommendations were followed to the last step when the DNA was eluted into a final volume of 50ul.

[0063] Quantitative determination of *C. jejuni* was accomplished using a previously published method with slight modifications (Skanseng et al. Comparison of chicken gut colonisation by the pathogens *Campylobacter jejuni* and *Clostridium perfringens* by real-time quantitative PCR. Mol Cell Probes 2006;20(5):269-79). The assay was optimized for use on the MX3005P (Agilent Technology) and Brilliant II QPCR master mix (Agilent Technologies) all other mixture components, primers, probe and cycling conditions remained as published.

[0064] A standard curve (Figure 1) was prepared using a pure culture of *C. jejuni* serially diluted 10-fold and added to a constant background of ileal content; total DNA isolation was done as previously described.

Statistical Analysis

[0065] Data were analyzed using Student's two-tailed *t-test* assuming unequal variances to compare the difference between groups and controls using JMP™ statistic software. A value of $P < 0.05$ was considered significant.

Results

[0066] An excellent correlation of quantification of *C. jejuni* using conventional microbiological enumeration techniques versus the qPCR was found (Fig. 1) with a greater than 99% correlation between the two methods. In experiment 1, we observed significant levels of colonization by the three candidate vectored vaccines within the cecal tonsils by day 3 post-vaccination; as well as significant invasion of the internal organs by the cj0113 expressing vector at the same time point (Table II). However, by day 11 post-vaccination, there was a decline in the amount of colonization of all three vectors and by day 21 post-vaccination, the vectors had been completely cleared from the cecal tonsils as well as the internal organs (Table II). We observed the same trend in our follow up vaccination study (experiment 2), using vector-expressed cj0113 as our vaccine candidate, as shown by the data presented in Table II. Table II.

Percentage of colonization, invasion and clearance of liver, spleen or cecal tonsils by <i>Salmonella</i> following vaccination with one of three <i>Salmonella</i> vectored vaccine candidates or saline gavage.								
	Liver/Spleen				Cecal Tonsils			
	Day 3	Day 11	Day 21	Day 32	Day 3	Day 11	Day 21	Day 32
Experiment 1								
Saline	0	0	0	0	0	0	0	0
cj0420	0	0	0	0	60	0	0	0
cj0113	50	0	0	0	100	40	0	0
cj0982	0	0	0	0	70	20	0	0
Experiment 2								
Saline	0	0	0	0	0	0	0	0
cj0113	0	20	0	0	50	40	0	0

In experiments 1 and 2 incidence of the attenuated recombinant *Salmonella* vaccine vector is represented as the percentage of positive liver, spleen, or cecal tonsils out of 10 birds. Chicks were orally gavaged with approximately 10^8 cfu of the appropriate treatment on day-of-hatch and. On days 3, 11, 21 and 32 post-hatch, 10 birds from each treatment group were euthanatized, and the livers, spleens, and ceca tonsils were collected for the determination (+/-) of the attenuated recombinant *Salmonella* vaccine vectors. The liver and spleen of each bird was pooled and assayed as one sample.

[0067] Chickens were challenged with *C. jejuni* on day 21 post vaccination. Ileal mucosal samples were obtained on days 21 and 32 post vaccination (days 0 and 11 post challenge) and used for DNA sample preparation to enumerate *C. jejuni* within the gut as described above. Vaccination with vector candidates cj0420 and cj0982 caused an approximate 1 log and 2 log reduction ($P < 0.05$), respectively, in the level of *C. jejuni* present in the ileal samples. Using the cj0113 vaccine candidate, there was a marked 4.8 log reduction ($P < 0.05$) of *C. jejuni* in the ileum compared to the control birds (Fig. 2).

[0068] In experiment 2, a repeat of the primary immunization study was done with only the vaccine candidate expressing cj0113. In this study, qPCR data revealed an approximate 5 log reduction of *C. jejuni* in cj0113 SE-vectored vaccine administered to birds when compared to the birds receiving saline only (Table III). Additionally, in experiment 3 vaccination with the cj0113 vector caused an approximate 4 log reduction, to below detectable levels, of *C. jejuni* as compared with the saline or *Salmonella* parent strain which contained no epitope insert (Fig. 3).

Table III.

Enumeration of <i>Campylobacter jejuni</i> by quantitative PCR in chicks 11 days following <i>Campylobacter</i> challenge in Experiment 2 (n=10).			
	Mean <i>C. jejuni</i> Log10 cfu/gm ileum	SD ^a	SE ^b
Saline	5.00	0.98	0.44
cj0113	0.00	0.00	0.00

In experiment 2 *Campylobacter jejuni* quantification was determined in chicks receiving saline or the *Salmonella* vectored vaccine candidate, cj0113 at 10^8 cfu/chick by quantitative PCR 11 days after receiving a *C. jejuni* challenge dose of approximately 10^7 cfu/ml. qPCR was performed on total DNA extracted by conventional methods from mucosal linings of the ileum. The results are presented as mean log10 cfu/gram of ileum content with standard deviation^a and standard error^b (n=10).

[0069] Serum samples collected in each experiment on Days 21 and 32 post- vaccination were used to determine *C. jejuni*- specific IgG antibodies. In the first experiment all three vaccine candidates (cj0420, cj0113, cj0982) caused significantly higher antibody levels at both time points when compared to the group which received only saline (Fig. 4). Also in the first experiment, the group vaccinated with cj0113 showed significantly higher antibody titers when compared to cj0420 and cj0982 (Fig. 4). An ELISA was also used to determine mucosal sIgA antibody levels specific for *Campylobacter*. These data indicate that the vaccine vector cj0113 caused a significant increase in the levels of sIgA when compared to the saline group and the two groups receiving either cj0420 or cj0982 (Fig. 5). The results from the second and third study in which only cj0113 was used as a vaccine candidate showed results similar to experiment 1 with vaccinated birds having significantly higher levels of antigen-specific IgG and sIgA antibodies to *C. jejuni* when compared to the birds receiving only saline (Data for Experiment 3 are shown in Fig. 6). Also, in the third experiment the antibody levels for the backbone strain (SE13) were similar to saline controls (Fig. 6).

Bacillus vectored vaccination study

Production of Heterologous Proteins for Vegetative Cell Expression

[0070] Plasmid pHT10 purchased from MoBioTec/Boca Scientific, Boca Raton, FL (Nguyen et al., 2007) was transformed at the multiple cloning site by addition of a *Bacillus subtilis* codon optimized insertion sequence for cj0113 and HMGB1 (SEQ ID NO: 4 and 20, respectively). DNA sequencing was done to confirm correct sequence insertion. The newly modified plasmid was then transformed into *Bacillus*. Briefly, *Bacillus* cultures were grown overnight at 37°C in HS media (Spizizen's medium supplemented with 0.5% glucose, 50µg/ml DL-tryptophan, 50µg/ml uracil, 0.02% casein hydrolysate, 0.1% yeast extract, 8µg/ml arginine, 0.4µg/ml histidine, 1mM MgSO₄). Inoculate 20ml LS medium (Spizizen's medium supplemented with 0.5% glucose, 5µg/ml DL-tryptophan, 5µg/ml uracil, 0.01% casein hydrolysate, 0.1% yeast extract, 1mM MgSO₄, 2.5mM MgCl₂, 0.5mM CaCl₂) with 1 ml overnight culture and incubate with shaking for 3-4 hours at 30 °C. Withdraw 1 ml of LS culture and add 10µl of 0.1M EGTA and incubate at room temperature for 5 minutes. Add 1-2 µg plasmid DNA, shake for 2 hours at 37°C, and plate on LB plates with selective antibiotics. These transformed *Bacillus* spp. now produce heterologous epitope sequences from *Campylobacter* (cj0113) and HMGB1 when induced with 1mM IPTG.

Vaccination study

[0071] In the vaccination challenge, 100 day-of-hatch broiler chicks were obtained from a local commercial hatchery and randomly assigned to one of four treatment groups: saline only, *Bacillus* vector alone (BSBB) or 10⁶ or 10⁸ *Bacillus* vaccine candidate, cj0113, (n=25/pen). Each treatment group was housed in an individual floor pen on fresh pine litter and provided water and feed *ad libitum*. On day-of-hatch, all chicks in each treatment group were inoculated via oral gavage with 0.25mL of a solution containing approximately 10⁶ cfu/mL of the backbone vector strain or the Cj0113 *Bacillus* vector. On day 10 birds received a booster vaccination of the same treatment they received on day of hatch. On day 21 post-hatch, all birds in each treatment group were challenged with *C. jejuni*, via oral gavage, with 0.25 mL of a solution containing 1x10⁷ cfu/ml prepared as described above. On days 3, 11, 21 (prior to booster inoculation) and 32 post-hatch, 10-15 birds from each treatment group were humanely killed and their liver, spleen and cecal tonsils aseptically removed for the determination of organ invasion, colonization and clearance of the vaccine vector strains. Also, on days 21 and 36 post-hatch ileum sections were removed and processed for use in qRT-PCR. In addition, blood samples were collected from 15 birds per treatment group and the serum was used for determining antibody response on days 21 and 36 post-hatch.

Results

[0072] Serum samples collected on Day 36 post- vaccination were used to determine *C. jejuni*-specific IgG antibodies. Vaccination with the bacillus vector expressing the cj0113 polypeptide caused significantly higher antibody levels when compared to the group which received only

saline or empty vector (Fig. 7). An ELISA was also used to determine mucosal sIgA antibody levels specific for *Campylobacter* at day 36 post vaccination. These data indicate that the vaccine vector expressing cj0113 caused a significant increase in the levels of sIgA when compared to the empty vector (Fig. 8).

[0073] Chickens were challenged with *C. jejuni* on day 24 post vaccination. Ileal mucosal samples were obtained on days 24 and 36 post vaccination (days 0 and 11 post challenge) and used for DNA sample preparation to enumerate *C. jejuni* within the gut as described above. Vaccination with the Bacillus cj0113 vector caused an approximate 3 log reduction ($P < 0.05$) in the level of *C. jejuni* present in the ileal samples (Fig. 9).

Vaccination study in Turkey poults

[0074] Since the *Salmonella*-vectored Cj0113 vaccine has been effective in reducing *Campylobacter* recovery after challenge in chickens we hypothesized that the vaccine may also work in poults. Therefore, to further evaluate the use of this epitope delivery system, an experiment was designed to test the effectiveness of the vaccine against *C. coli* challenge in turkey poults.

[0075] The Δ SE-cj0113 vaccine was constructed as described above. *Salmonella enteritidis* phage type 13A (SE13A) was used as the backbone strain for this vaccine candidate. This isolate was double attenuated by irreversible gene deletions in the *aroA* and *htrA* genes as previously described. Recombinant strains containing these deletions were then modified further to incorporate the cj0113 insert and an immunostimulatory molecule, CD-154 (Δ SE-cj0113). These sequences were integrated as previously described.

[0076] In this experiment, 70 poults were obtained from a local hatchery. They were randomly assigned to one of two treatment groups and tagged. Thirty poults were orally gavaged with 10^8 cfu/poult Δ SE-cj0113 and the remaining poults were sham treated with saline. On day 21, poults were challenged with 1.5×10^8 cfu/poult of *C. coli* by oral gavage. Liver, spleen and cecal tonsils were aseptically removed on day 3 (N=10), day 21 (N=10) and day 35 (N=10) for detection of vector recovery by enrichment in tetrathionate broth and plating on Brilliant Green agar. On days 21 and 35, ingesta (N=10) and tissue (N=5) samples were collected for further analysis. Ingesta from the ileum of cultured poults were analyzed using qPCR for enumeration of *C. coli*. Tissue samples from the same region were collected and total RNA was extracted. Interferon-gamma and TNF-alpha like vaccine responses were evaluated.

Results

[0077] As previously shown in chickens, the poults developed a significant immune response following vaccination with *Salmonella*-vectored Cj0113 vaccine. After challenge, there was a

about a five-log reduction in *Campylobacter coli* in the ileum as compared to vector only controls (Fig. 10). Vaccination with the *Salmonella*-vectored Cj0113 vaccine does appear to work in turkeys similar to the chickens.

SEQUENCE LISTING

[0078]

<110> THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ARKANSAS

<120> VACCINE AND METHODS TO REDUCE CAMPYLOBACTER INFECTION

<130> 5658-00102

<150> US 61/353,039

<151> 2010-06-09

<160> 44

<170> PatentIn version 3.5

<210> 1

<211> 165

<212> PRT

<213> *Campylobacter jejuni* cjaD

<400> 1

Met Lys Lys Ile Leu Phe Thr Ser Ile Ala Ala Leu Ala Val Val Ile
1 5 10 15

Ser Gly Cys Ser Thr Lys Ser Thr Ser Val Ser Gly Asp Ser Ser Val
20 25 30

Asp Ser Asn Arg Gly Ser Gly Gly Ser Asp Gly Trp Asp Ile Asp Ser
35 40 45

Lys Ile Ser Gln Leu Asn Asp Thr Leu Asn Lys Val Tyr Phe Asp Phe
50 55 60

Asp Lys Phe Asn Ile Arg Pro Asp Met Gln Asn Val Val Ser Thr Asn
65 70 75 80

Ala Asn Ile Phe Asn Thr Glu Val Ser Gly Val Ser Ile Thr Val Glu
85 90 95

Gly Asn Cys Asp Glu Trp Gly Thr Asp Glu Tyr Asn Gln Ala Leu Gly
100 105 110

Leu Lys Arg Ala Lys Ala Val Lys Glu Ala Leu Ile Ala Lys Gly Val
115 120 125

Asn Ala Asp Arg Ile Ala Val Lys Ser Tyr Gly Glu Thr Asn Pro Val
 130 135 140

Cys Thr Glu Lys Thr Lys Ala Cys Asp Ala Gln Asn Arg Arg Ala Glu
 145 150 155 160

Phe Lys Leu Ser Arg
 165

<210> 2

<211> 279

<212> PRT

<213> Campylobacter jejuni cjaA

<400> 2

Met Lys Lys Ile Leu Leu Ser Val Leu Thr Ala Phe Val Ala Val Val
 1 5 10 15

Leu Ala Ala Cys Gly Gly Asn Ser Asp Ser Lys Thr Leu Asn Ser Leu
 20 25 30

Asp Lys Ile Lys Gln Asn Gly Val Val Arg Ile Gly Val Phe Gly Asp
 35 40 45

Lys Pro Pro Phe Gly Tyr Val Asp Glu Lys Gly Asn Asn Gln Gly Tyr
 50 55 60

Asp Ile Ala Leu Ala Lys Arg Ile Ala Lys Glu Leu Phe Gly Asp Glu
 65 70 75 80

Asn Lys Val Gln Phe Val Leu Val Glu Ala Ala Asn Arg Val Glu Phe
 85 90 95

Leu Lys Ser Asn Lys Val Asp Ile Ile Leu Ala Asn Phe Thr Gln Thr
 100 105 110

Pro Gln Arg Ala Glu Gln Val Asp Phe Cys Ser Pro Tyr Met Lys Val
 115 120 125

Ala Leu Gly Val Ala Val Pro Lys Asp Ser Asn Ile Thr Ser Val Glu
 130 135 140

Asp Leu Lys Asp Lys Thr Leu Leu Leu Asn Lys Gly Thr Thr Ala Asp
 145 150 155 160

Ala Tyr Phe Thr Gln Asn Tyr Pro Asn Ile Lys Thr Leu Lys Tyr Asp
 165 170 175

Gln Asn Thr Glu Thr Phe Ala Ala Leu Met Asp Lys Arg Gly Asp Ala
 180 185 190

Leu Ser His Asp Asn Thr Leu Leu Phe Ala Trp Val Lys Asp His Pro
 195 200 205

Asp Phe Lys Met Gly Ile Lys Glu Leu Gly Asn Lys Asp Val Ile Ala
210 215 220

Pro Ala Val Lys Lys Gly Asp Lys Glu Leu Lys Glu Phe Ile Asp Asn

225 230 235 240

Leu Ile Ile Lys Leu Gly Gln Glu Gln Phe Phe His Lys Ala Tyr Asp
245 250 255

Glu Thr Leu Lys Ala His Phe Gly Asp Asp Val Lys Ala Asp Asp Val
260 265 270

Val Ile Glu Gly Gly Lys Ile
275

<210> 3

<211> 190

<212> PRT

<213> Campylobacter jejuni ACE 393

<400> 3

Met Lys Lys Val Leu Leu Ser Ser Leu Val Ala Val Ser Leu Leu Ser
1 5 10 15

Thr Gly Leu Phe Ala Lys Glu Tyr Thr Leu Asp Lys Ala His Thr Asp
20 25 30

Val Gly Phe Lys Ile Lys His Leu Gln Ile Ser Asn Val Lys Gly Asn
35 40 45

Phe Lys Asp Tyr Ser Ala Val Ile Asp Phe Asp Pro Ala Ser Ala Glu
50 55 60

Phe Lys Lys Leu Asp Val Thr Ile Lys Ile Ala Ser Val Asn Thr Glu
65 70 75 80

Asn Gln Thr Arg Asp Asn His Leu Gln Gln Asp Asp Phe Phe Lys Ala
85 90 95

Lys Lys Tyr Pro Asp Met Thr Phe Thr Met Lys Lys Tyr Glu Lys Ile
100 105 110

Asp Asn Glu Lys Gly Lys Met Thr Gly Thr Leu Thr Ile Ala Gly Val
115 120 125

Ser Lys Asp Ile Val Leu Asp Ala Glu Ile Gly Gly Val Ala Lys Gly
130 135 140

Lys Asp Gly Lys Glu Lys Ile Gly Phe Ser Leu Asn Gly Lys Ile Lys
145 150 155 160

Arg Ser Asp Phe Lys Phe Ala Thr Ser Thr Ser Thr Ile Thr Leu Ser
165 170 175

Asp Asp Ile Asn Leu Asn Ile Glu Val Glu Ala Asn Glu Lys
 180 185 190

<210> 4

<211> 120

<212> DNA

<213> Campylobacter jejuni cjaD

<400> 4

tctctctccg gtgtagcat caccgttgaa ggtaactgtg atgaatgggg taccgatgaa 60

tataaccagg cgtcctcctc ctggatgacc acctcctatg cgccgacctc ctctcctcc 120

<210> 5

<211> 162

<212> DNA

<213> Campylobacter jejuni cjaA

<400> 5

tctctctcca aagttgogct ggggtgtgog gttccgaaag atagcaacat caccagcgtt 60

gaagatctga aagataaaac cctgctgctg aacaaaggta ccaccgogga tgogtctctc 120

tctggatga ccacctcta tgcgogacc tctcctctcc cc 162

<210> 6

<211> 120

<212> DNA

<213> Campylobacter jejuni ACE 393

<400> 6

tctctctcca aagatatcgt tctggatgog gaaatcogtg gtgttgogaa aggtaaagat 60

ggtaaagaaa aatcctcctc ctggatgacc acctcctatg cgccgacctc ctctcctcc 120

<210> 7

<211> 21

<212> PRT

<213> Campylobacter jejuni Cj0113

<400> 7

Gly Val Ser Ile Thr Val Glu Gly Asn Cys Asp Glu Trp Gly Thr Asp
 1 5 10 15

Glu Tyr Asn Gln Ala
 20

<210> 8

<211> 21

<212> PRT

<213> Campylobacter jejuni Cj0982

<400> 8

Lys Asp Ile Val Leu Asp Ala Glu Ile Gly Gly Val Ala Lys Gly Lys

<213> Artificial Sequence

<220>

<223> Synthetic peptide: Cj0420-CD154

<400> 12

Lys Val Ala Leu Gly Val Ala Val Pro Lys Asp Ser Asn Ile Thr Ser
1 5 10 15

Val Glu Asp Leu Lys Asp Lys Thr Leu Leu Leu Asn Lys Gly Thr Thr
20 25 30

Ala Asp Ala Trp Met Thr Thr Ser Tyr Ala Pro Thr Ser
35 40 45

<210> 13

<211> 272

<212> PRT

<213> Gallus gallus

<220>

<221> misc_feature

<223> CD154 chicken

<400> 13

Met Asn Glu Ala Tyr Ser Pro Ala Ala Pro Arg Pro Met Gly Ser Thr
1 5 10 15

Ser Pro Ser Thr Met Lys Met Phe Met Cys Phe Leu Ser Val Phe Met
20 25 30

Val Val Gln Thr Ile Gly Thr Val Leu Phe Cys Leu Tyr Leu His Met
35 40 45

Lys Met Asp Lys Met Glu Glu Val Leu Ser Leu Asn Glu Asp Tyr Ile
50 55 60

Phe Leu Arg Lys Val Gln Lys Cys Gln Thr Gly Glu Asp Gln Lys Ser
65 70 75 80

Thr Leu Leu Asp Cys Glu Lys Val Leu Lys Gly Phe Gln Asp Leu Gln
85 90 95

Cys Lys Asp Arg Thr Ala Ser Glu Glu Leu Pro Lys Phe Glu Met His
100 105 110

Arg Gly His Glu His Pro His Leu Lys Ser Arg Asn Glu Thr Ser Val
115 120 125

Ala Glu Glu Lys Arg Gln Pro Ile Ala Thr His Leu Ala Gly Val Lys
130 135 140

Ser Asn Thr Thr Val Arg Val Leu Lys Trp Met Thr Thr Ser Tyr Ala
145 150 155 160

Pro Thr Ser Ser Leu Ile Ser Tyr His Glu Gly Lys Leu Lys Val Glu
 165 170 175

Lys Ala Gly Leu Tyr Tyr Ile Tyr Ser Gln Val Ser Phe Cys Thr Lys
 180 185 190

Ala Ala Ala Ser Ala Pro Phe Thr Leu Tyr Ile Tyr Leu Tyr Leu Pro
 195 200 205

Met Glu Glu Asp Arg Leu Leu Met Lys Gly Leu Asp Thr His Ser Thr
 210 215 220

Ser Thr Ala Leu Cys Glu Leu Gln Ser Ile Arg Glu Gly Gly Val Phe
 225 230 235 240

Glu Leu Arg Gln Gly Asp Met Val Phe Val Asn Val Thr Asp Ser Thr
 245 250 255

Ala Val Asn Val Asn Pro Gly Asn Thr Tyr Phe Gly Met Phe Lys Leu
 260 265 270

- <210> 14
- <211> 261
- <212> PRT
- <213> Homo sapiens
- <220>
- <221> misc_feature
- <223> CD154 human

<400> 14

Met Ile Glu Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly
 1 5 10 15

Leu Pro Ile Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu
 20 25 30

Ile Thr Gln Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg
 35 40 45

Arg Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val
 50 55 60

Phe Met Lys Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser
 65 70 75 80

Leu Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys
 85 90 95

Asp Ile Met Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu
 100 105 110

Met Gln Lys Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser
 115 120 125

Glu Ala Ser Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly
 130 135 140

Tyr Tyr Thr Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln
 145 150 155 160

Leu Thr Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr
 165 170 175

Phe Cys Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser
 180 185 190

Leu Cys Leu Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala
 195 200 205

Ala Asn Thr His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His
 210 215 220

Leu Gly Gly Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn
 225 230 235 240

Val Thr Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe
 245 250 255

Gly Leu Leu Lys Leu
 260

- <210> 15
- <211> 11
- <212> PRT
- <213> Homo sapiens
- <220>
- <221> misc_feature
- <223> CD154 partial human

<400> 15
 Trp Ala Glu Lys Gly Tyr Tyr Thr Met Ser Cys
 1 5 10

- <210> 16
- <211> 11
- <212> PRT
- <213> Gallus gallus
- <220>
- <221> misc_feature
- <223> Chicken CD154 peptide

<400> 16
 Trp Met Phe Phe Ser Thr Ala Asp Phe Ser Ser

Trp Met Thr Thr Ser Tyr Ala Phe Thr Ser Ser
 1 5 10

<210> 17

<211> 10

<212> PRT

<213> Anas sp.

<220>

<221> misc_feature

<223> Duck CD154 peptide

<400> 17

Trp Asn Lys Thr Ser Tyr Ala Pro Met Asn
 1 5 10

<210> 18

<211> 10

<212> PRT

<213> Mus sp.

<220>

<221> misc_feature

<223> Mouse CD154 peptide

<400> 18

Trp Ala Lys Lys Gly Tyr Tyr Thr Met Lys
 1 5 10

<210> 19

<211> 10

<212> PRT

<213> Bos taurus

<220>

<221> misc_feature

<223> Cow CD154 peptide

<400> 19

Trp Ala Pro Lys Gly Tyr Tyr Thr Leu Ser
 1 5 10

<210> 20

<211> 190

<212> PRT

<213> Gallus gallus

<220>

<221> misc_feature

<223> Chicken HMGB1 amino acid

<400> 20

Met Gly Lys Gly Asp Pro Lys Lys Pro Arg Gly Lys Met Ser Ser Tyr
 1 5 10 15

Ala Phe Phe Val Gln Thr Cys Arg Glu Glu His Lys Lys Lys His Pro
 20 25 30

Asp Ala Ser Val Asn Phe Ser Glu Phe Ser Lys Lys Cys Ser Glu Arg
 35 40 45

Trp Lys Thr Met Ser Ser Lys Glu Lys Gly Lys Phe Glu Asp Met Ala
 50 55 60

Lys Ala Asp Lys Leu Arg Tyr Glu Lys Glu Met Lys Asn Tyr Val Pro
 65 70 75 80

Pro Lys Gly Glu Thr Lys Lys Lys Phe Lys Asp Pro Asn Ala Pro Lys
 85 90 95

Arg Pro Pro Ser Ala Phe Phe Leu Phe Cys Ser Glu Phe Arg Pro Lys
 100 105 110

Ile Lys Gly Glu His Pro Gly Leu Ser Ile Gly Asp Val Ala Lys Lys
 115 120 125

Leu Gly Glu Met Trp Asn Asn Thr Ala Ala Asp Asp Lys Gln Pro Tyr
 130 135 140

Glu Lys Lys Ala Ala Lys Leu Lys Glu Lys Tyr Glu Lys Asp Ile Ala
 145 150 155 160

Ala Tyr Arg Ala Lys Gly Lys Val Asp Ala Gly Lys Lys Val Val Ala
 165 170 175

Lys Ala Glu Lys Ser Lys Lys Lys Lys Glu Glu Glu Glu Asp
 180 185 190

<210> 21

<211> 85

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide: HMGB1 box a1

<400> 21

Met Gly Lys Gly Asp Pro Lys Lys Pro Arg Gly Lys Met Ser Ser Tyr
 1 5 10 15

Ala Phe Phe Val Gln Thr Cys Arg Glu Glu His Lys Lys Lys His Pro
 20 25 30

Asp Ala Ser Val Asn Phe Ser Glu Phe Ser Lys Lys Cys Ser Glu Arg
 35 40 45

Trp Lys Thr Met Ser Ser Lys Glu Lys Gly Lys Phe Glu Asp Met Ala
 50 55 60

Lys Ala Asp Lys Leu Arg Tyr Glu Lys Glu Met Lys Asn Tyr Val Pro
 65 70 75 80

Pro Lys Gly Glu Thr
 85

<210> 22

<211> 54

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide: HMGB1 box a2

<400> 22

Pro Asp Ala Ser Val Asn Phe Ser Glu Phe Ser Lys Lys Cys Ser Glu
 1 5 10 15

Arg Trp Lys Thr Met Ser Ser Lys Glu Lys Gly Lys Phe Glu Asp Met
 20 25 30

Ala Lys Ala Asp Lys Leu Arg Tyr Glu Lys Glu Met Lys Asn Tyr Val
 35 40 45

Pro Pro Lys Gly Glu Thr
 50

<210> 23

<211> 73

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide: HMGB1 box b1

<400> 23

Lys Asp Pro Asn Ala Pro Lys Arg Pro Pro Ser Ala Phe Phe Leu Phe
 1 5 10 15

Cys Ser Glu Phe Arg Pro Lys Ile Lys Gly Glu His Pro Gly Leu Ser
 20 25 30

Ile Gly Asp Val Ala Lys Lys Leu Gly Glu Met Trp Asn Asn Thr Ala
 35 40 45

Ala Asp Asp Lys Gln Pro Tyr Glu Lys Lys Ala Ala Lys Leu Lys Glu
 50 55 60

Lys Tyr Glu Lys Asp Ile Ala Ala Tyr
 65 70

<210> 24

<211> 69

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide: HMGB1 box b2

<400> 24

Asn Ala Pro Lys Arg Pro Pro Ser Ala Phe Phe Leu Phe Cys Ser Glu
 1 5 10 15

Phe Arg Pro Lys Ile Lys Gly Glu His Pro Gly Leu Ser Ile Gly Asp
 20 25 30

Val Ala Lys Lys Leu Gly Glu Met Trp Asn Asn Thr Ala Ala Asp Asp
 35 40 45

Lys Gln Pro Tyr Glu Lys Lys Ala Ala Lys Leu Lys Glu Lys Tyr Glu
 50 55 60

Lys Asp Ile Ala Ala
 65

<210> 25

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide: HMGB1 RAGE Binding domain

<400> 25

Lys Asp Pro Asn Ala Pro Lys Arg Pro Pro Ser Ala Phe Phe Leu Phe
 1 5 10 15

Cys Ser Glu Phe Arg
 20

<210> 26

<211> 33

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide: HMGB1 proinflammatory cytokine activity

<400> 26

Leu Lys Glu Lys Tyr Glu Lys Asp Ile Ala Ala Tyr Arg Ala Lys Gly
 1 5 10 15

Lys Val Asp Ala Gly Lys Lys Val Val Ala Lys Ala Glu Lys Ser Lys
 20 25 30

Lys

<210> 27

<211> 215

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> HMGB1

<400> 27

Met Gly Lys Gly Asp Pro Lys Lys Pro Arg Gly Lys Met Ser Ser Tyr
 1 5 10 15

Ala Phe Phe Val Gln Thr Cys Arg Glu Glu His Lys Lys Lys His Pro
 20 25 30

Asp Ala Ser Val Asn Phe Ser Glu Phe Ser Lys Lys Cys Ser Glu Arg
 35 40 45

Trp Lys Thr Met Ser Ala Lys Glu Lys Gly Lys Phe Glu Asp Met Ala
 50 55 60

Lys Ala Asp Lys Ala Arg Tyr Glu Arg Glu Met Lys Thr Tyr Ile Pro
 65 70 75 80

Pro Lys Gly Glu Thr Lys Lys Lys Phe Lys Asp Pro Asn Ala Pro Lys
 85 90 95

Arg Pro Pro Ser Ala Phe Phe Leu Phe Cys Ser Glu Tyr Arg Pro Lys
 100 105 110

Ile Lys Gly Glu His Pro Gly Leu Ser Ile Gly Asp Val Ala Lys Lys
 115 120 125

Leu Gly Glu Met Trp Asn Asn Thr Ala Ala Asp Asp Lys Gln Pro Tyr
 130 135 140

Glu Lys Lys Ala Ala Lys Leu Lys Glu Lys Tyr Glu Lys Asp Ile Ala
 145 150 155 160

Ala Tyr Arg Ala Lys Gly Lys Pro Asp Ala Ala Lys Lys Gly Val Val
 165 170 175

Lys Ala Glu Lys Ser Lys Lys Lys Lys Glu Glu Glu Glu Asp Glu Glu
 180 185 190

Asp Glu Glu Asp Glu Glu Glu Glu Glu Asp Glu Glu Asp Glu Asp Glu
 195 200 205

Glu Glu Asp Asp Asp Asp Glu
 210 215

<210> 28

<211> 205

<212> PRT

<213> Danio rerio

<220>

<221> misc_feature

<223> Zebra fish HMGB1

<400> 28

Met Gly Lys Asp Pro Thr Lys Pro Arg Gly Lys Met Ser Ser Tyr Ala
 1 5 10 15

Tyr Phe Val Gln Thr Cys Arg Glu Glu His Lys Lys Lys His Pro Glu
 20 25 30

Ala Thr Val Asn Phe Ser Glu Phe Ser Lys Lys Cys Ser Glu Arg Trp
 35 40 45

Lys Thr Met Ser Ala Lys Glu Lys Gly Lys Phe Glu Asp Met Ala Lys
 50 55 60

Leu Asp Lys Ala Arg Tyr Glu Arg Glu Met Lys Asn Tyr Ile Pro Pro
 65 70 75 80

Lys Gly Glu Lys Lys Lys Arg Phe Lys Asp Pro Asn Ala Pro Lys Arg
 85 90 95

Pro Pro Ser Ala Phe Phe Ile Phe Cys Ser Glu Phe Arg Pro Lys Val
 100 105 110

Lys Glu Glu Thr Pro Gly Leu Ser Ile Gly Asp Val Ala Lys Arg Leu
 115 120 125

Gly Glu Met Trp Asn Lys Ile Ser Ser Glu Glu Lys Gln Pro Tyr Glu
 130 135 140

Lys Lys Ala Ala Lys Leu Lys Glu Lys Tyr Glu Lys Asp Ile Ala Ala
 145 150 155 160

Tyr Arg Ser Lys Gly Lys Val Gly Gly Gly Ala Ala Lys Ala Pro Ser
 165 170 175

Lys Pro Asp Lys Ala Asn Asp Glu Asp Glu Asp Asp Asp Glu Glu Glu
 180 185 190

Asp Glu Asp Asp Asp Asp Glu Glu Glu Glu Asp Asp Glu
 195 200 205

<210> 29

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: lam-up-f; Loop 9 up

<400> 29

tgtacaagtg gacgccaatc 20

<210> 30

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: lam-up-r; Loop 9 up

<400> 30

gttatcgccg tctttgatat agcc 24

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: lam-dn-f; Loop 9 dn

<400> 31

atttcccggt atgccgcagc 20

<210> 32

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: lam-dn-r; Loop 9 dn

<400> 32

gttaaacaga gggcgacgag 20

<210> 33

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Km-f; I-Scel/Kmr gene

<400> 33

.....

gctatatcaa agacggcgat aactaactat aacggtccta aggtagcgaa tttccgggga 60
tccgtcga 68

<210> 34
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic primer: Km-r; I-SceI/Kmr gene

<400> 34
gctgcggcat aacgggaaat gtaggctgga gctgctcg 39

<210> 35
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic primer: Kan4f; Inside Kmr gene

<400> 35
caaaagcgct ctgaagttcc 20

<210> 36
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic primer: Kan4r; Inside Kmr gene

<400> 36
gcgtgagggg atctgaagt 20

<210> 37
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic primer: lam 3f; Outer regions of loop 9

<400> 37
gccatctcgc ttggtgataa 20

<210> 38
<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: lam 3r; Outer regions of loop 9

<400> 38

cgctggtatt ttgcggtaca 20

<210> 39

<211> 84

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Cj0113f; Insert with loop 9 up

<400> 39

ttcatcggta cccattcat cacagttacc ttcaacggtg atgctaacac cggaggagga 60

gttatcgccg tctttgatat agcc 84

<210> 40

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Cj0113r; Insert with loop 9 down

<400> 40

atgaatgggg taccgatgaa tataaccagg cgtcctcctc ctggatgacc acctcctatg 60

cgccgacctc ctctcctccc atttcccgtt atgccgcagc 100

<210> 41

<211> 84

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Cj0420f; Insert with loop 9 up

<400> 41

atctttacct ttcgcaacac caccgatttc cgcattccaga acgatatctt tggaggagga 60

gttatcgccg tctttgatat agcc 84

<210> 42

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Cj0420r; Insert with loop 9 down

<400> 42

```
gtggtgcgaa aggtaaagat ggtaaagaaa aatcctcctc ctggatgacc acctcctatg      60
cgccgacctc ctctcctcc atttcccgtt atgccgcagc                               100
```

<210> 43

<211> 105

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Cj0982c-f; Insert with loop 9 up

<400> 43

```
ggttttatct ttcagatctt caacgctggt gatggtgcta tctttcggaa ccgcaacacc      60
cagcgcaact ttggaggagg agttatcgcc gtctttgata tagcc                               105
```

<210> 44

<211> 121

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic primer: Cj0982c-r; Insert with loop 9 down

<400> 44

```
aagatctgaa agataaaacc ctgctgctga acaaaggtac caccgcggat gcgtcctcct      60
cctggatgac cacctcctat gcgcgcacct cctcctcctc catttcccgt tatgccgcag      120
c                                                                              121
```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2008036675A](#) [0024] [0046]
- [WO2011091255A](#) [0024]
- [WO02092004A](#) [0033]
- [US61353039B](#) [0078]

Non-patent literature cited in the description

- **WYSZYŃSKA, AGNIESZKA et al.** Vaccine, 2004, vol. 22.11, 1379-1389 [0005]
- **BUCKLEY, ANTHONY M. et al.** Vaccine, 2010, vol. 28.4, 1094-1105 [0006]
- **ANDERSSON TRACEY** Annu. Rev. Immunol., 2011, vol. 29, 139-162 [0030]
- **COX et al.** Scarless and site-directed mutagenesis in *Salmonella enteritidis* chromosome BMC Biotechnol, 2007, vol. 7, 59- [0055]
- **BURNS DM BEACHAM IR** Rare codons in *E. coli* and *S. typhimurium* signal sequences FEBS Lett, 1985, vol. 189, 2318-24 [0055]
- **COLE et al.** Effect of aeration and storage temperature on *Campylobacter* concentrations in poultry semen Poultry Sci, 2004, vol. 83, 1734-8 [0057]
- **BROWN et al.** Detection of antibodies to *Mycoplasma gallisepticum* in egg yolk versus serum samples J Clin Microbiol, 1991, vol. 29, 122901-3 [0061]
- **DAVIES et al.** Evaluation of the use of pooled serum, pooled muscle tissue fluid (meat juice) and pooled faeces for monitoring pig herds for *Salmonella* J Appl Microbiol, 2003, vol. 95, 51016-25 [0061]
- **SKANSENG et al.** Comparison of chicken gut colonisation by the pathogens *Campylobacter jejuni* and *Clostridium perfringens* by real-time quantitative PCR Mol Cell Probes, 2006, vol. 20, 5269-7 [0063]
- **NGUYEN et al.** MoBioTec/Boca Scientific, 2007, [0070]

P a t e n t k r a v

- 5 **1.** Bakterie, der er egnet til anvendelse i vaccination af fjerkræ, og ekspressionen af et antigen polypeptid, bestående af sekvensen med SEQ ID NO: 7, hvor bakterien omfatter en første polynukleotidsekvens, der koder for det antigen polypeptid, hvor den første polynukleotidsekvens, der ikke er nativt associeret med bakterien og yderligere omfattende en anden polynukleotidsekvens, der koder for et immunstimulerende polypeptid, der ikke er nativt associeret med bakterien, hvor den eneste polypeptidsekvens, som er afledt af C. jejuni cjaD (cj0113), der er kodet af bakteriens polynukleotider er det antigen polypeptid bestående af sekvensen med SEQ ID NO: 7.
- 10
- 2.** Bakterie ifølge krav 1, hvor bakterien er fra en art udvalgt blandt Salmonella, Escherichia, Bacillus eller Lactobacillus.
- 15
- 3.** Bakterie ifølge et af kravene 1-2, hvor bakterien omfatter mere end en kopi af den første polynukleotidsekvens, mere end en kopi af den anden polynukleotidsekvens eller mere end en kopi af den første og anden polynukleotidsekvens.
- 20
- 4.** Bakterie ifølge et af kravene 1-3, hvor den første polynukleotidsekvens er forbundet i ramme med den anden polynukleotidsekvens, og eventuelt er den første polynukleotidsekvens og den anden polynukleotidsekvens integreret i bakteriens genom.
- 25
- 5.** Bakterie ifølge et af kravene 1-4, hvor det antigen polypeptid og det immunstimulerende polypeptid er udtrykt på vektorens overflade.
- 6.** Bakterie ifølge et af kravene 1-5, hvor den første polynukleotidsekvens og den anden polynukleotidsekvens er sat inden i en tredje polynukleotidsekvens, der koder for en ekstern del af et transmembranprotein.
- 30
- 7.** Bakterie ifølge et af kravene 1-6, hvor det immunstimulerende polypeptid er et CD154-polypeptid, der er i stand til at binde CD40, eller et HMGB1-polypeptid, hvor CD154-polypeptidet har mindre end 50 aminosyrer og omfatter aminosyrerne 140-149 af SEQ ID NO:13 eller er udvalgt blandt SEQ ID NO: 15-
- 35

19, og hvor HMGB1-polypeptidet omfatter et polypeptid udvalgt blandt mindst en blandt SEQ ID NO: 20-28.

5 **8.** Bakterie ifølge et af kravene 1-7, hvor det første polypeptid koder for polypeptidet som angivet i SEQ ID NO: 10.

10 **9.** Bakterie ifølge et af kravene 1-8, hvor bakterien er i stand til at styrke resistensen hos et individ over for en efterfølgende infektion med *Campylobacter* spp. efter indgivelse af bakterien til individet i en virksom mængde.

10. Farmaceutisk sammensætning omfattende bakterien ifølge et af kravene 1-9 og en farmaceutisk acceptabel bærer.

15 **11.** Bakterie ifølge et af kravene 1-9 til anvendelse ved en fremgangsmåde med at inducere en terapeutisk immunreaktion hos et individ, hvor bakterien skal indgives til individet i en mængde, er er virksom til at styrke individets immunreaktion over for det antigene polypeptid.

20 **12.** Bakterie ifølge et af kravene 1-9 til anvendelse ved en fremgangsmåde til at styrke resistens over for *Campylobacter*-infektion hos et individ, hvor bakterien skal indgives til individet i en mængde, der er virksom til at styrke resistensen over for *Campylobacter*-infektion efter efterfølgende udsættelse for *Campylobacter* spp.

25 **13.** Bakterie til anvendelse ifølge krav 11 eller 12, hvor bakterien er svækket eller dræbt inden indgivelse.

30 **14.** Bakterie til anvendelse ifølge krav 11 eller 13, hvor den terapeutiske immunreaktion indbefatter styrket antistofproduktion, og hvor den styrkede antistofproduktion omfatter øget opløselig IgA.

15. Bakterie til anvendelse ifølge krav 12 eller 13, hvor øget resistens er **kendetegnet ved, at** efterfølgende infektion med *Campylobacter* spp. resulterer i nedsat *Campylobacter*-vækst sammenlignet med en kontrol.

35

DRAWINGS




Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.




Figure 8.




Figure 9.




Figure 10.