



(51) International Patent Classification:

A61K 31/47 (2006.01) A61P 35/00 (2006.01)

A61P 1/16 (2006.01)

(21) International Application Number:

PCT/JP2018/018810

(22) International Filing Date:

15 May 2018 (15.05.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/506,900 16 May 2017 (16.05.2017) US

(71) Applicant: EISAI R&D MANAGEMENT CO., LTD.

[JP/JP]; 4-6-10 Koishikawa, Bunkyo-ku, Tokyo, 1128088 (JP).

(72) Inventor: TAMAI Toshiyuki; C/O Eisai Co., Ltd., 4-6-10

Koishikawa, Bunkyo-ku, Tokyo, 1128088 (JP).

(74) Agent: HASEGAWA Yoshiaki et al.; SOEI PATENT AND

LAW FIRM, Marunouchi MY PLAZA (Meiji Yasuda Life Bldg.) 9th fl., 1-1, Marunouchi 2-chome, Chiyoda-ku, Tokyo, 1000005 (JP).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: TREATMENT OF HEPATOCELLULAR CARCINOMA

(57) Abstract: This disclosure provides methods for treating a hepatocellular carcinoma (e.g., unresectable HCC) with lenvatinib or a pharmaceutically acceptable salt thereof. Also encompassed by the disclosure are dosage regimens described herein of lenvatinib or a pharmaceutically acceptable salt thereof for use in treating hepatocellular carcinoma (e.g., unresectable hepatocellular carcinoma) according to any of the methods described herein. Particularly useful dosages and dose modifications upon the occurrence of an adverse event or events are also disclosed.



Description

Title of Invention: TREATMENT OF HEPATOCELLULAR CARCINOMA

Technical Field

[0001] The present application relates generally to methods of treating hepatocellular carcinoma.

Background Art

[0002] Hepatocellular carcinoma (HCC) is the second leading cause of cancer death worldwide and is responsible for nearly 745,000 deaths each year. It usually occurs in a background of chronic liver disease, particularly in cirrhosis, which limits the feasibility of surgical resection. Sorafenib, an oral multikinase inhibitor, extends overall survival when used as a first-line treatment for HCC, demonstrating a median improvement of 2.8 months compared with placebo (10.7 months vs. 7.9 months; hazard ratio [HR]: 0.69; $P < 0.001$) despite a low response rate of 2% (Llovet, *N Engl. J Med.*, 359:378-390, 2008).

[0003] Drug development in HCC in the past 10 years is marked by four failed phase 3 trials (of sunitinib, brivanib, linifanib, and erlotinib plus sorafenib) that did not demonstrate noninferiority (Cheng, *J. Clin. Oncol.*, 31:4067-4075, 2013; Johnson, *J. Clin. Oncol.*, 31:36517-3524, 2013; Cainap, *J. Clin. Oncol.*, 33:172-179, 2015) or superiority (Zhu, *J. Clin. Oncol.*, 33(6):559-66, 2015) to sorafenib in overall survival. Therefore, unresectable HCC represents a highly unmet medical need.

Summary of Invention

[0004] This disclosure relates, in part, to methods of treating a subject with a HCC (e.g., advanced HCC, unresectable HCC (uHCC), or advanced uHCC) with lenvatinib or a pharmaceutically acceptable salt thereof. In some embodiments, lenvatinib or a pharmaceutically acceptable salt thereof is administered as a first-line single agent to patients with unresectable HCC. In some embodiments, the dosage of lenvatinib or a pharmaceutically acceptable salt thereof is modified upon the occurrence of one or more adverse events in the treated subject.

[0005] In a first aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen of lenvatinib or a pharmaceutically acceptable salt thereof that is: (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In certain embodiments, lenvatinib or a pharmaceutically acceptable salt thereof is administered orally. In certain embodiments, lenvatinib or a

pharmaceutically acceptable salt thereof is administered once daily. In certain embodiments, lenvatinib or a pharmaceutically acceptable salt thereof is administered orally, once daily.

[0006] In a second aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen of lenvatinib or a pharmaceutically acceptable salt thereof that is 8 mg/day. In certain embodiments, lenvatinib or a pharmaceutically acceptable salt thereof is administered orally. In certain embodiments, lenvatinib or a pharmaceutically acceptable salt thereof is administered once daily. In certain embodiments, lenvatinib or a pharmaceutically acceptable salt thereof is administered orally, once daily.

[0007] As used throughout this disclosure, a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a specified dose means that lenvatinib or a pharmaceutically acceptable salt thereof is present in the dosage regimen at the specified dose. Although such a dosage regimen can contain additional components, lenvatinib or a pharmaceutically acceptable salt thereof is present only at the specific dose listed. The dose of lenvatinib or a pharmaceutically acceptable salt thereof (e.g., 12 mg, 8 mg, or 4 mg) as used throughout refers to the dose of the free form of lenvatinib.

[0008] In a third aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In carrying out this method, the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity until the first Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a second Grade 3 nonhematologic toxicity during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second Grade 3 nonhematologic toxicity

until the second Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a third Grade 3 nonhematologic toxicity during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third Grade 3 nonhematologic toxicity until the third Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[0009] In a fourth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In carrying out this method, the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity until the first Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In certain embodiments, the human subject develops an occurrence of a second Grade 3 nonhematologic toxicity during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second Grade 3 nonhematologic toxicity until the second Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In certain embodiments, the human subject develops an occurrence of a third Grade 3 nonhematologic toxicity during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third Grade 3 nonhematologic toxicity.

[0010] In a fifth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unre-

resectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In carrying out this method, the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity until the first persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity until the third persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[0011] In a sixth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In

carrying out this method, the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose 4 mg/day. In certain embodiments, the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In certain embodiments, the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity.

[0012] In a seventh aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In carrying out this method, the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject

is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

- [0013] In an eighth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In carrying out this method, the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising

lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In certain embodiments, the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In certain embodiments, the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality.

- [0014] In some embodiments of the third to eighth aspects, following or during treatment with the second dosage regimen, the human subject does not develop an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality. In such embodiments, the method further comprises continuing administration of the second dosage regimen to the human subject (i.e., not lowering the dose being given in the second dosage regimen).
- [0015] In some embodiments of the third to eighth aspects, following or during treatment with the third dosage regimen, the human subject does not develop an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life threatening Grade 4 laboratory abnormality. In such embodiments, the method further comprises continuing administration of the third dosage regimen to the human subject (i.e., not lowering the dose being given in the third dosage regimen).
- [0016] In certain embodiments, the human subject develops an occurrence of a Grade 4 non-hematologic toxicity excluding a non-life-threatening Grade 4 laboratory abnormality during treatment with the above dosage regimens. In such embodiments, the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding non-life-threatening Grade 4 laboratory abnormality.
- [0017] In a ninth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unre-

sectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In carrying out this method, the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria until the first Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria until the second Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria until the third Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria during treatment with the fourth dosage regimen. In such embodiments, the method further comprises terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria until the fourth Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fifth dosage regimen comprising lenvatinib or a pharmaceutically ac-

ceptable salt thereof at a dose of 4 mg every other day.

[0018] In a tenth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma that comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In carrying out this method, the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria until the first Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In certain embodiments, the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria until the second Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In certain embodiments, the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria until the third Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In certain embodiments, the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria during treatment with the fourth dosage regimen. In such embodiments, the method further comprises terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria.

[0019] In some embodiments of the ninth or tenth aspects, following or during treatment with the second dosage regimen, the human subject does not develop an occurrence of a second Grade 3 hematologic toxicity or proteinuria. In such embodiments, the method further comprises continuing administration of the second dosage regimen to the human subject (i.e., not lowering the dose being given in the second dosage regimen).

- [0020] In some embodiments of the ninth or tenth aspects, following or during treatment with the third dosage regimen, the human subject does not develop an occurrence of a third Grade 3 hematologic toxicity or proteinuria. In such embodiments, the method further comprises continuing administration of the third dosage regimen to the human subject (i.e., not lowering the dose being given in the third dosage regimen).
- [0021] In some embodiments of the ninth or tenth aspects, following or during treatment with the fourth dosage regimen, the human subject does not develop an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria. In such embodiments, the method further comprises continuing administration of the fourth dosage regimen to the human subject (i.e., not lowering the dose being given in the fourth dosage regimen).
- [0022] In an eleventh aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In carrying out this method, the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg. In certain embodiments, the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity until the third Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal

to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[0023] In a twelfth aspect, the disclosure provides a method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In carrying out this method, the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen. Thereupon, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In certain embodiments, the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen. In such embodiments, the method further comprises terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In certain embodiments, the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen. In such embodiments, the method further comprises terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity.

[0024] In a thirteenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is

resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a first persistent and intolerable Grade 2 non-hematologic toxicity during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 nonhematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 non-hematologic toxicity excluding nonlife-threatening laboratory abnormality.

[0025] In a fourteenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has

an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. If the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. If the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. If the human subject develops an occurrence of a first persistent and intolerable Grade 2 non-hematologic toxicity during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 nonhematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. If the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the first Grade 3 non-hematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. If the human subject develops an occurrence of a Grade 4 non-hematologic toxicity excluding nonlife-threatening laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 non-

hematologic toxicity excluding nonlife-threatening laboratory abnormality.

[0026] In some embodiments of the eleventh to fourteenth aspects, following or during treatment with the second dosage regimen, the human subject does not develop an occurrence of a second Grade 4 hematologic toxicity. In such embodiments, the method further comprises continuing administration of the second dosage regimen to the human subject (i.e., not lowering the dose being given in the second dosage regimen).

[0027] In some embodiments of the eleventh to fourteenth aspects, following or during treatment with the third dosage regimen, the human subject does not develop an occurrence of a third Grade 4 hematologic toxicity. In such embodiments, the method further comprises continuing administration of the third dosage regimen to the human subject (i.e., not lowering the dose being given in the third dosage regimen).

[0028] In a fifteenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen. In yet other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/

day, and wherein the human subject developed an occurrence of a Grade 3 non-hematologic toxicity during treatment with the prior dosage regimen.

[0029] In a sixteenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 non-hematologic toxicity during treatment with the prior dosage regimen.

[0030] In a seventeenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or

more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen. In yet other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen.

[0031] In an eighteenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen.

[0032] In a nineteenth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight

of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen. In certain instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen.

[0033] In a twentieth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising

lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen.

[0034] In a twenty first aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen. In certain instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen. In other instances, the method comprises ad-

administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen.

[0035] In a twenty second aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen. In yet other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen.

[0036] In a twenty third aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4

mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen. In other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen. In yet other instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen.

[0037] In a twenty fourth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen. In some instances, the method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen

comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen.

[0038] In a twenty fifth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. The human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen. In some instances, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen. The method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen. The method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth

dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In instances wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0039] In a twenty sixth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. The human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen. In some instances, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen. The method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject

develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen. The method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality. In instances wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0040] In a twenty seventh aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In instances where the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In instances where the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the

dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0041] In a twenty eighth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In instances where the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In instances where the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory ab-

normality.

[0042] In a twenty ninth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. The human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen. In some instances, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen. The method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen. The method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day. In instances wherein the human subject develops an

occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0043] In a thirtieth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. The human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen. In some instances, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen. The method further comprises terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen. The method further comprises terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality. In instances wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4

laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0044] In a thirty first aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. In instances where the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In instances where the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0045] In a thirty second aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment

classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In instances where the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In instances where the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality. In the above aspects, hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[0046] In a thirty third aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a Grade 3 hypertension during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the Grade 3 hypertension until the Grade 3 hypertension is controlled at less than or equal to Grade 2, and administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg.

[0047] In a thirty fourth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment

classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. If the human subject develops an occurrence of a Grade 3 hypertension during treatment with the first dosage regimen, then the method further comprises terminating administration of the first dosage regimen after the occurrence of the Grade 3 hypertension until the Grade 3 hypertension is controlled at less than or equal to Grade 2, and administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day.

[0048] In a thirty fifth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a 2 g or greater proteinuria in 24 hours during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the 2 g or greater proteinuria in 24 hours until the proteinuria is less than or equal to 2 g of proteinuria in 24 hours and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg. In the above aspect, the human subject develops an occurrence of a nephrotic syndrome during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the nephrotic syndrome.

[0049] In a thirty sixth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. If the human subject develops an occurrence of a 2 g or greater proteinuria in 24 hours during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the 2 g or greater proteinuria in 24 hours until the proteinuria is less than or equal to 2 g of proteinuria in 24 hours and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day. In the above aspect, the human subject develops an occurrence of a nephrotic syndrome during treatment with the first dosage regimen, and

the method further comprises terminating administration of the dosage regimen after the occurrence of the nephrotic syndrome.

[0050] In a thirty seventh aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg. If the human subject develops an occurrence of a greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation until the QT/QTc interval prolongation improves to less than or equal to 480 ms or baseline and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg.

[0051] In a thirty eighth aspect, the disclosure features a method of treating unresectable hepatocellular carcinoma. The method comprises administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. If the human subject develops an occurrence of a greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation during treatment with the first dosage regimen, then the method further comprises terminating administration of the dosage regimen after the occurrence of the greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation until the QT/QTc interval prolongation improves to less than or equal to 480 ms or baseline and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day if the body weight of the human subject is less than 60 kg.

[0052] In some embodiments of the above aspects, median overall survival is 13.6 months.

[0053] In some embodiments of the above aspects, median overall survival with 95% confidence interval is between 12.1 and 14.9 months.

[0054] In some embodiments of the above aspects, hazard ratio of overall survival compared with sorafenib at a dosage of 400 mg twice daily is 0.92.

- [0055] In some embodiments of the above aspects, hazard ratio of overall survival compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 0.79 and 1.06.
- [0056] In some embodiments of the above aspects, overall survival is shown in Fig. 1 (Kaplan-Meier Plot of Overall Survival).
- [0057] In some embodiments of the above aspects, median progression-free survival is 7.4 months.
- [0058] In some embodiments of the above aspects, wherein median progression-free survival with 95% confidence interval is between 6.9 to 8.8 months.
- [0059] In some embodiments of the above aspects, hazard ratio of progression-free survival compared with sorafenib at a dosage of 400 mg twice daily is 0.66.
- [0060] In some embodiments of the above aspects, hazard ratio of progression-free survival compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 0.57 and 0.77.
- [0061] In some embodiments of the above aspects, progression-free survival is shown in Fig. 2 (Kaplan-Meier Plot of Progression-Free Survival).
- [0062] In some embodiments of the above aspects, median time to progression is 8.9 months.
- [0063] In some embodiments of the above aspects, median time to progression with 95% confidence interval is between 7.4 to 9.2 months.
- [0064] In some embodiments of the above aspects, hazard ratio of time to progression compared with sorafenib at a dosage of 400 mg twice daily is 0.63.
- [0065] In some embodiments of the above aspects, hazard ratio of time to progression compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 0.53 and 0.73.
- [0066] In some embodiments of the above aspects, time to progression is shown in Fig. 6 (Kaplan-Meier Plot of Time to Progression).
- [0067] In some embodiments of the above aspects, the objective response rate is 24.1%.
- [0068] In some embodiments of the above aspects, the odds ratio of objective response rate compared with sorafenib at a dosage of 400 mg twice daily is 3.13.
- [0069] In some embodiments of the above aspects, the odds ratio of objective response rate compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 2.15 to 4.56.
- [0070] In some embodiments of the above aspects, the method comprises achieving the results shown in Table 2 or Table 10 (Efficacy Results in HCC).
- [0071] In some embodiments of the above aspects, the method comprises achieving the results shown in Fig. 15 (Quality of Life).
- [0072] In some embodiments of the above aspects, the human subject consists essentially of

the subject with mild hepatic impairment classified in Child-Pugh class A under Child-Pugh Classification.

[0073] In some embodiments of the above aspects, the human subject is categorized to stage B or stage C based on Barcelona Clinic Liver Cancer (BCLC) staging system.

[0074] In some embodiments of the above aspects, medical management of each of the first, second, and third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicities or non-life-threatening Grade 4 laboratory abnormality is initiated prior to terminating administration of the dosage regimen administered at the time of onset of the Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality.

[0075] In some embodiments of the above aspects, medical management of each of the first, second, third, and fourth Grade 3 hematologic toxicities or proteinuria is initiated prior to terminating administration of the dosage regimen administered at the time of onset of the Grade 3 hematologic toxicities or proteinuria.

[0076] In some embodiments of the above aspects, medical management of each of the first, second, and third Grade 4 hematologic toxicities is initiated prior to terminating administration of the dosage regimen administered at the time of onset of the Grade 4 hematologic toxicities.

[0077] In some embodiments of the above aspects, the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is the same as the second and/or third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality.

[0078] In some embodiments of the above aspects, the first Grade 3 hematologic toxicity or proteinuria is the same as the second and/or third Grade 3 hematologic toxicity or proteinuria.

[0079] In some embodiments of the above aspects, the first Grade 4 hematologic toxicity is the same as the second and/or third Grade 4 hematologic toxicity.

[0080] In some embodiments of any of the above aspects, the Grade 3 nonhematologic toxicity is selected from the group consisting of Grade 3 hypertension, Grade 3 diarrhea, Grade 3 decreased appetite, Grade 3 fatigue, Grade 3 arthralgia, Grade 3 myalgia, Grade 3 decreased weight, Grade 3 dysphonia, Grade 3 nausea, Grade 3 abdominal pain, Grade 3 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 3 vomiting, Grade 3 constipation, Grade 3 rash, and Grade 3 palmar-plantar erythrodysesthesia.

[0081] In some embodiments of any of the above aspects, the Grade 2 or Grade 3 non-hematologic toxicity is selected from the group consisting of Grade 3 hypertension, Grade 2 hypertension, Grade 3 diarrhea, Grade 2 diarrhea, Grade 3 decreased appetite,

Grade 2 decreased appetite, Grade 3 fatigue, Grade 2 fatigue, Grade 3 arthralgia, Grade 2 arthralgia, Grade 3 myalgia, Grade 2 myalgia, Grade 3 decreased weight, Grade 2 decreased weight, Grade 2 alopecia, Grade 3 dysphonia, Grade 2 dysphonia, Grade 3 nausea, Grade 2 nausea, Grade 3 abdominal pain, Grade 2 abdominal pain, Grade 3 QT/QTc interval prolongation, Grade 2 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 2 hypothyroidism, Grade 3 vomiting, Grade 2 vomiting, Grade 3 constipation, Grade 2 constipation, Grade 3 rash, Grade 2 rash, Grade 3 palmar-plantar erythrodysesthesia, and Grade 2 palmar-plantar erythrodysesthesia.

[0082] In some embodiments of any of the above aspects, the Grade 4 laboratory abnormality is selected from the group consisting of Grade 4 increase in aspartate aminotransferase, Grade 4 increase in alanine aminotransferase, Grade 4 increase in alkaline phosphatase, Grade 4 hypokalemia, Grade 4 hyponatremia, Grade 4 hypoglycemia, Grade 4 increase in blood bilirubin, and Grade 4 increase in gamma glutamyl transferase.

[0083] In some embodiments of any of the above aspects, the Grade 3 hematologic toxicity or proteinuria is selected from the group consisting of Grade 3 proteinuria, Grade 3 thrombopenia (thrombocytopenia), Grade 3 anemia, Grade 3 decrease in white blood cell count, Grade 3 neutropenia, and Grade 3 lymphocytopenia.

[0084] In some embodiments of any of the above aspects, the Grade 4 hematologic toxicity is selected from the group consisting of Grade 4 thrombopenia (thrombocytopenia), Grade 4 anemia, Grade 4 decrease in white blood cell count, Grade 4 neutropenia, and Grade 4 lymphocytopenia.

[0085] In some embodiments of the above aspects, lenvatinib or the pharmaceutically acceptable salt thereof is formulated as a capsule.

[0086] In some embodiments of the above aspects, lenvatinib or the pharmaceutically acceptable salt thereof is administered to the human subject orally.

[0087] In some embodiments of the above aspects, lenvatinib or a pharmaceutically acceptable salt thereof is lenvatinib mesylate.

[0088] Also encompassed by the disclosure is a dosage regimen described herein of lenvatinib or a pharmaceutically acceptable salt thereof for use in treating hepatocellular carcinoma (e.g., unresectable hepatocellular carcinoma) according to any of the methods described herein.

[0089] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the exemplary methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by

reference in their entirety. In case of conflict, the present application, including definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0090] Other features and advantages of the invention will be apparent from the following detailed description and from the claims.

Brief Description of Drawings

[0091] [fig.1]Fig. 1 depicts Kaplan-Meier estimates of overall survival by treatment group. CI denotes confidence interval, and HR hazard ratio.

[fig.2]Fig. 2 shows progression-free survival by modified Response Evaluation Criteria in Solid Tumors (mRECIST). CI denotes confidence interval, and HR hazard ratio.

[fig.3]Fig. 3 provides a subgroup analyses of overall survival indicating associated hazard ratio and 95% confidence interval. AFP denotes alpha-fetoprotein, BCLC Barcelona Clinic Liver Cancer, CI confidence interval, and HR hazard ratio.

[fig.4]Fig. 4 provides subgroup analyses of progression-free survival indicating the associated hazard ratio and 95% confidence interval. AFP denotes alpha-fetoprotein, BCLC Barcelona Clinic Liver Cancer, CI confidence interval, and HR hazard ratio.

[fig.5]Fig. 5 provides a schematic representation of the enrollment, randomization, and treatment of the 954 patients in the phase 3 trial.

[fig.6]Fig. 6 is a graph providing a Kaplan-Meier Estimate of time to progression. CI denotes confidence interval, and HR hazard ratio.

[fig.7]Fig. 7 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.8]Fig. 8 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.9]Fig. 9 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.10]Fig. 10 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.11]Fig. 11 depicts waterfall graphs of percentage change in summed diameter of target lesions. The percentage change in lesion size is shown from baseline to nadir.

[fig.12]Fig. 12 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.13]Fig. 13 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.14]Fig. 14 provides Forest plots indicating hazard ratios for time to progression in the subgroup analyses. AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; HR, hazard ratio.

[fig.15]Fig. 15 provides Forest plots of hazard ratio of time to clinically meaningful worsening of QLQC30 questionnaire scores (top) and HCC18 questionnaire scores (bottom) comparing lenvatinib to sorafenib. * Nominal p-value; HR, hazard ratio; LCL, lower control limit; UCL, upper control limit.

Description of Embodiments

[0092] This disclosure provides methods of treating a human subject that has a hepatocellular carcinoma (e.g., advanced HCC, unresectable HCC, or unresectable advanced HCC). The method comprises administering to the subject a starting dose of lenvatinib or a pharmaceutically acceptable salt thereof. The starting dose is determined based on the body weight of the subject: 12 mg/day if the body weight of the human subject is equal to or more than 60 kg, and 8 mg/day if the body weight of the human subject is less than 60 kg. However, if the human subject has moderate hepatic impairment classified in Child-Pugh class B under the Child-Pugh Classification, then the starting dose can be 8 mg/day regardless of the body weight of the subject. If the subject develops one or more adverse events as a result of the treatment with lenvatinib or a pharmaceutically acceptable salt thereof, the disclosure provides modifications of the treatment regimen as well as adjusted dosing regimens (reduced doses of lenvatinib or a pharmaceutically acceptable salt thereof). These dose modifications enable the subject to continue treatment with lenvatinib or a pharmaceutically acceptable salt thereof. If the subject does not develop an adverse reaction as a result of administration of a particular dose of lenvatinib or a pharmaceutically acceptable salt thereof, the subject can be maintained on the same dosage regimen.

[0093] Hepatocellular Carcinoma

Primary liver cancer is the sixth most common cancer worldwide with approximately 782,000 new cases worldwide in 2012. A total of 83% of the world's cases occur in less-developed regions and more than 50% occur in China alone; however, the incidence of liver cancer has been rising in a number of low-rate areas such as Australia, the United Kingdom, and the United States. In all populations, males have higher rates of liver cancer than do females. Liver cancer is the third most common cause of cancer death worldwide, accounting for an estimated 746,000 cancer deaths annually.

- [0094] Hepatocellular carcinoma (HCC) accounts for 85% to 90% of primary cancer of the liver and occurs predominantly in patients with underlying chronic liver disease, in particular cirrhosis. Major causes of cirrhosis include hepatitis B virus (HBV), hepatitis C virus (HCV), and alcoholic liver disease. Hepatitis B is the most frequent underlying cause of HCC, with an estimated 300 million people with chronic infection worldwide. Chronic HBV carriers have a 5- to 15-fold increased risk of developing HCC compared with the general population. Chronic HCV infection is also a major risk factor for HCC. The risk of HCC was 17-fold higher in HCV-infected patients compared with HCV-negative controls.
- [0095] To reduce mortality rates, it is important not only to prevent onset and promote early detection of HCC in patients in whom the hepatitis virus cannot be eradicated, but also to develop effective therapeutic interventions for patients who have already developed HCC.
- [0096] For HCC, at least eight different staging systems are available. These include the Barcelona Clinic Liver Cancer (BCLC) system, the Okuda stage system, the TNM staging system, the JIS score, the CLIP score, the CUPI score, the French classification, and the ER system (Pons et al., HPB, 7(1):35-41, 2005). The BCLC system categorizes HCC based on characteristics of the tumor, liver function, performance status, and cancer-related symptoms. BCLC stage groupings include:
- Very early stage. The tumor is smaller than 2 cm. There is no increased pressure in the portal vein. Bilirubin levels are normal. Surgery is usually recommended.
- Early stage. The tumor is smaller than 5 cm. Liver function varies. There may be no increased pressure in the portal vein, increased portal vein pressure and normal bilirubin levels, or increased portal vein pressure and increased bilirubin levels. People with early-stage disease may be candidates for a liver transplant, surgery, or radiofrequency ablation (RFA).
- Intermediate stage. The tumor may be large or there may be multiple tumors. Doctors usually recommend regional therapies, such as transarterial chemoembolization.
- Advanced stage. The tumor has invaded the portal vein or spread to other parts of the body, such as the lungs and bones. Targeted therapy is generally recommended.
- [0097] Evidence-based guidelines for the diagnosis and treatment of HCC have been established and adopted (El-Serag, et al., Ann. Intern. Med., 139(10):817-23, 2003; Llovet, et al., Lancet, 362(9399):1907-17, 2003). Recent technological advances for patients with early-stage HCC include surgical resection, localized treatments such as radiofrequency ablation (RFA), percutaneous ethanol injection (PEI), cryotherapy and transarterial chemoembolization (TACE). Treatment options for HCC are determined by stage of the disease. For very early stage HCC, surgical resection is the treatment of choice in patients without cirrhosis. Among patients who have underlying cirrhosis, or-

thotopic liver transplantation is the treatment option associated with the lowest risk of tumor recurrence. For patients with early-stage HCC who are not eligible for surgical resection or transplantation, RFA is likely the best alternative treatment, although PEI and cryoablation are also routinely performed. Other locoregional treatment options for HCC include transarterial TACE, which has been shown to improve survival among patients with preserved liver function and unresectable HCC whose disease is either too large or too multifocal for percutaneous ablation techniques, as well as radioembolization, which has been used as palliative treatment in intermediate-stage HCC. However, recurrence due to residual tumor cells is a clinical characteristic of HCC and ultimately leads to an advanced stage where surgery, RFA, and TACE are no longer appropriate.

[0098] This disclosure provides methods of treating unresectable HCC using lenvatinib or a pharmaceutically acceptable salt thereof.

[0099] Lenvatinib


A number of kinase inhibitors have been developed as antitumor agents. For example, a group of compounds having inhibitory activity against receptor tyrosine kinases, such as vascular endothelial growth factor receptor (VEGFR), are known to inhibit angiogenesis and are regarded as a new class of antitumor agents. Lenvatinib is a multi-target receptor tyrosine kinase inhibitor that inhibits the kinase activities of VEGFR1 (FLT1), VEGFR2 (KDR), and VEGFR3 (FLT4). Lenvatinib inhibits other receptor tyrosine kinases that have been implicated in pathogenic angiogenesis, tumor growth, and cancer progression in addition to their normal cellular functions, including fibroblast growth factor (FGF) receptors FGFR1, FGFR2, FGFR3, and FGFR4; re-arranged during transfection receptor (RET), KIT, and platelet-derived growth factor receptor alpha (PDGFR α). Lenvatinib also exhibits antiproliferative activity in hepatocellular carcinoma cell lines dependent on activated FGFR signaling with a concurrent inhibition of FGF-receptor substrate 2 α (FRS2 α) phosphorylation.

[0100] The term “lenvatinib” refers to

4-(3-chloro-4(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide. This compound is disclosed in Example 368 (see, column 270) of U.S. Patent No. 7,253,286. U.S. Patent No. 7,253,286 is incorporated by reference in its entirety herein. The term “pharmaceutically acceptable salt” is not particularly restricted as to the type of salt. Examples of such salts include, but are not limited to, inorganic acid addition salt such as hydrochloric acid salt, sulfuric acid salt, carbonic acid salt, bicarbonate salt, hydrobromic acid salt, and hydriodic acid salt; organic carboxylic acid addition salt such as acetic acid salt, maleic acid salt, lactic acid salt, tartaric acid salt, and trifluoroacetic acid salt; organic sulfonic acid addition salt such as methane-sulfonic acid salt, hydroxymethanesulfonic acid salt, hydroxyethanesulfonic acid salt,

benzenesulfonic acid salt, toluenesulfonic acid salt, and taurine salt; amine addition salt such as trimethylamine salt, triethylamine salt, pyridine salt, procaine salt, picoline salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, N-methylglucamine salt, diethanolamine salt, triethanolamine salt, tris(hydroxymethylamino)methane salt, and phenethylbenzylamine salt; and amino acid addition salt such as arginine salt, lysine salt, serine salt, glycine salt, aspartic acid salt, and glutamic acid salt. In one embodiment, the pharmaceutically acceptable salt is a methanesulfonic acid salt ("mesylate"). The methanesulfonic acid salt form (i.e., the mesylate) of lenvatinib is disclosed in US Patent 7,612,208, which is incorporated by reference herein in its entirety. The chemical name of lenvatinib mesylate is 4-[3-chloro-4-(N'-cyclopropylureido)phenoxy]-7-methoxyquinoline-6-carboxamide methanesulfonate and its chemical structure is provided below:

[Chem.1]



[0101] Lenvatinib mesylate is also referred to as LENVIMA^(registered trademark).

[0102] Lenvatinib mesylate is a white to pale reddish yellow powder. It is slightly soluble in water and practically insoluble in ethanol (dehydrated). The dissociation constant (pKa value) of lenvatinib mesylate is 5.05 at 25°C. The partition coefficient (log P value) is 3.30.

[0103] Administration

As shown in the Examples, which describe, inter alia, the results of an open-label phase 3 human clinical trial in subjects with unresectable HCC, lenvatinib was shown to be noninferior to sorafenib in overall survival (median 13.6 months with lenvatinib vs. 12.3 months with sorafenib; hazard ratio [HR]: 0.92; 95% confidence interval [CI], 0.79 to 1.06). Lenvatinib prolonged progression-free survival (median 7.3 vs. 3.6 months; HR: 0.64; 95% CI, 0.55 to 0.75; P<0.001) versus sorafenib. Objective response rate was 41% with lenvatinib versus 12% with sorafenib (P<0.001). In sum, lenvatinib improved progression-free survival, time to progression, and objective response rate versus sorafenib. Thus, lenvatinib can be used for the treatment of HCC.

[0104] Lenvatinib or a pharmaceutically acceptable salt thereof may be administered orally to a human subject in need thereof (e.g., a human subject having advanced HCC,

uHCC, or advanced uHCC) by any means that the health care provider deems useful.

[0105] For oral administration, the lenvatinib compound can be in the form of, e.g., a tablet, capsule, suspension, or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are capsules, tablets, powders, granules or a suspension, with conventional additives such as lactose, mannitol, corn starch or potato starch; with binders such as crystalline cellulose, cellulose derivatives, acacia, corn starch or gelatins; with disintegrators such as corn starch, potato starch or sodium carboxymethyl-cellulose; and with lubricants such as talc or magnesium stearate. The active ingredient(s) may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable pharmaceutically acceptable carrier.

[0106] In one embodiment, lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) is administered to the human subject as a capsule. The capsule can contain, lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) equivalent to 1 mg, 2 mg, 3 mg, 4 mg, 5 mg, 6 mg, 7 mg, 8 mg, 9 mg, 10 mg, 11 mg, or 12 mg of lenvatinib. In certain instances, the capsule contains lenvatinib mesylate equivalent to 4 mg lenvatinib. In certain instances, the capsule contains lenvatinib mesylate equivalent to 8 mg lenvatinib. In certain instances, the capsule contains lenvatinib mesylate equivalent to 12 mg lenvatinib. In some embodiments, these capsules also contain one or more of the following inactive ingredients: calcium carbonate, mannitol, microcrystalline cellulose, hydroxypropylcellulose, hydroxypropyl cellulose (type H), and talc. In some embodiments, the shell of these capsules is a hypromellose shell and can contain one or more of: titanium dioxide, ferric oxide yellow, and ferric oxide red. The printing ink used on the capsule may contain one or more of: shellac, black iron oxide, potassium hydroxide, and propylene glycol.

[0107] In certain embodiments, lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) is administered to the human subject at a dose of 12 mg once daily. This dose can be administered, e.g., as three 4 mg capsules orally once daily. In other embodiments, lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) is administered to the human subject at a dose of 8 mg once daily. This dose can be administered, e.g., as two 4 mg capsules orally once daily. In some embodiments, lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) is administered to the human subject at a dose of 4 mg once daily. This dose can be administered, e.g., as one 4 mg capsule orally once daily. In some embodiments, lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) is administered to the human subject at a dose of 4 mg every other day. This

dose can be administered, e.g., as one 4 mg capsule orally once every other day.

[0108] It is recommended that the subject take lenvatinib or a pharmaceutically acceptable salt thereof one time each day at about the same time, with or without food.

[0109] If the patient is unable to swallow the lenvatinib capsules whole, the patient may use a cup to measure about one tablespoon of water or apple juice into a glass and place the drug capsules into the liquid without breaking or crushing them. The capsules should be left in the liquid for at least 10 minutes and the contents then stirred for at least 3 minutes. The patient can then drink this mixture. After drinking, the patient should rinse the glass with a small amount of additional water or apple juice and swallow the liquid.

[0110] In certain embodiments, lenvatinib or the pharmaceutically acceptable salt thereof is administered to a subject that has a HCC (e.g., advanced HCC or unresectable HCC) once daily for at least 7 weeks, at least 14 weeks, at least 28 weeks, at least 56 weeks, at least 84 weeks, at least 112 weeks, at least 140 weeks, at least 168 weeks, or at least 196 weeks.

[0111] Methods of Treatment to Control, Reduce, or Prevent Adverse Events

A major problem in treating a subject with a new therapy is the development of a treatment-emergent adverse event(s) (TEAE). A treatment-emergent adverse event is as any adverse event not present in the subject prior to the initiation of the treatment, or any adverse event already present that worsens in either intensity or frequency following exposure to the treatment. In certain embodiments, the adverse event is a persistent and intolerable adverse event.

[0112] The National Cancer Institute Common Terminology Criteria for Adverse Events v4.0 (CTCAE, published: May 28, 2009; v4.03: June 14, 2010) (incorporated by reference herein in its entirety) is a descriptive terminology that can be utilized for adverse event reporting. The CTCAE provides a grading (severity) scale for each adverse event term. An Adverse Event (AE) is any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease temporally associated with the use of a medical treatment or procedure that may or may not be considered related to the medical treatment or procedure. An AE is a term that is a unique representation of a specific event used for medical documentation and scientific analyses. An AE can be graded. The CTCAE grade refers to the severity of the AE. The CTCAE displays Grades 1 through 5 with unique clinical descriptions of severity for each AE based on this guideline:

Grade 1: Mild; asymptomatic or mild symptoms; clinical or diagnostic observations only; intervention not indicated.

Grade 2: Moderate; minimal, local or noninvasive intervention indicated; limiting age-appropriate instrumental Activities of Daily Living (ADL). ["Instrumental ADL"]

refer to preparing meals, shopping for groceries or clothes, using the telephone, managing money, etc.]

Grade 3: Severe or medically significant but not immediately life-threatening; hospitalization or prolongation of hospitalization indicated; disabling; limiting self-care ADL. ["Self-care ADL" refers to bathing, dressing and undressing, feeding self, using the toilet, taking medications, and not bedridden.]

Grade 4: Life-threatening consequences; urgent intervention indicated.

Grade 5: Death related to AE.

[0113] Not all Grades are appropriate for all AEs. Therefore, some AEs are listed in the CTCAE with fewer than five options for Grade selection.

[0114] Therapy with lenvatinib or a pharmaceutically acceptable salt thereof (e.g., lenvatinib mesylate) can lead to treatment-emergent adverse events (see Examples). In certain embodiments, the adverse event associated with therapy with lenvatinib or a pharmaceutically acceptable salt thereof is a persistent and intolerable AE. In certain instances, the persistent and intolerable AE is a Grade 2 AE. In other instances, the persistent and intolerable AE is a Grade 3 AE. In certain embodiments, the adverse event associated with the therapy using lenvatinib or a pharmaceutically acceptable salt thereof is a Grade 4 AE. In yet other instances, the persistent and intolerable AE is a Grade 4 laboratory abnormality. In certain cases, the Grade 2 or Grade 3 AE is a non-hematological toxicity. In other cases, the Grade 3 or Grade 4 AE is a hematological toxicity. In yet other cases, the Grade 3 AE is proteinuria. The most common adverse reactions observed in lenvatinib-treated HCC subjects were, in order of decreasing frequency, hypertension, fatigue, diarrhea, decreased appetite, arthralgia/myalgia, and decreased weight.

[0115] Hypertension is a disorder characterized by a pathological increase in blood pressure; a repeatedly elevation in the blood pressure exceeding 140 over 90 mm Hg, and is graded as follows:

Grade 1: Prehypertension (systolic BP 120 - 139 mm Hg or diastolic BP 80 - 89 mm Hg)

Grade 2: Stage 1 hypertension (systolic BP 140 - 159 mm Hg or diastolic BP 90 - 99 mm Hg); medical intervention indicated; recurrent or persistent (≥ 24 hrs); symptomatic increase by >20 mm Hg (diastolic) or to $>140/90$ mm Hg if previously WNL; monotherapy indicated Pediatric: recurrent or persistent (≥ 24 hrs) BP $>ULN$; monotherapy indicated

Grade 3: Stage 2 hypertension (systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg); medical intervention indicated; more than one drug or more intensive therapy than previously used indicated Pediatric: Same as adult

Grade 4: Life-threatening consequences (e.g., malignant hypertension, transient or

permanent neurologic deficit, hypertensive crisis); urgent intervention indicated

Pediatric: Same as adult

Grade 5: Death

- [0116] Fatigue is a disorder characterized by a state of generalized weakness with a pronounced inability to summon sufficient energy to accomplish daily activities and is graded as follows:

Grade 1: Fatigue relieved by rest.

Grade 2: Fatigue not relieved by rest or limiting instrumental activities of daily living (ADL). Instrumental ADL refer to preparing meals, shopping for groceries or clothes, using the telephone, managing money, etc.

Grade 3: Fatigue not relieved by rest, limiting self-care ADL. Self-care ADL refer to bathing, dressing and undressing, feeding self, using the toilet, taking medications, and not bedridden.

Grade 4: Grade is not available.

Grade 5: Grade is not available.

- [0117] Diarrhea is a disorder characterized by frequent and watery bowel movements and is graded as follows:

Grade 1: Increase of <4 stools per day over baseline; mild increase in ostomy output compared to baseline

Grade 2: Increase of 4 - 6 stools per day over baseline; moderate increase in ostomy output compared to baseline

Grade 3: Increase of ≥ 7 stools per day over baseline; incontinence; hospitalization indicated; severe increase in ostomy output compared to baseline; limiting self-care ADL

Grade 4: Life-threatening consequences; urgent intervention indicated

Grade 5: death

- [0118] Decreased appetite (anorexia) is a disorder characterized by a loss of appetite, and is graded as follows:

Grade 1: Loss of appetite without alteration in eating habits

Grade 2: Oral intake altered without significant weight loss or malnutrition; oral nutritional supplements indicated

Grade 3: Associated with significant weight loss or malnutrition (e.g., inadequate oral caloric and/or fluid intake); tube feeding or TPN indicated

Grade 4: Life-threatening consequences; urgent intervention indicated

Grade 5: Death

- [0119] Arthralgia is a disorder characterized by a sensation of marked discomfort in a joint, and is graded as follows:

Grade 1: Mild pain

Grade 2: Moderate pain; limiting instrumental ADL

Grade 3: Severe pain; limiting self care ADL

Grade 4: Not available

Grade 5: Not available

[0120] Myalgia is a disorder characterized by marked discomfort sensation originating from a muscle or group of muscles, and is graded as follows:

Grade 1: Mild pain

Grade 2: Moderate pain; limiting instrumental ADL

Grade 3: Severe pain; limiting self care ADL

Grade 4: Not available

Grade 5: Not available

[0121] Decreased weight (weight loss) is a finding characterized by a decrease in overall body weight; for pediatrics, less than the baseline growth curve, and is graded as follows:

Grade 1: Weight loss 5 to <10% from baseline; intervention not indicated

Grade 2: 10 - <20% from baseline; nutritional support indicated

Grade 3: $\geq 20\%$ from baseline; tube feeding or TPN indicated

Grade 4: Not available

Grade 5: Not available

[0122] The most common serious adverse reactions ($\geq 2\%$) in LENVIMA-treated patients were hepatic encephalopathy (4%), hepatic failure (3%), ascites (3%), and decreased appetite (2%).

[0123] Hepatic encephalopathy comprises hepatobiliary disorders, and is graded as follows:

Grade 1: Asymptomatic or mild symptoms; clinical or diagnostic observations only; intervention not indicated

Grade 2: Moderate; minimal, local or noninvasive intervention indicated; limiting age appropriate instrumental ADL

Grade 3: Severe or medically significant but not immediately life-threatening; hospitalization or prolongation of existing hospitalization indicated; disabling; limiting self care ADL

Grade 4: Life-threatening consequences; urgent intervention indicated.

Grade 5: Death

[0124] Hepatic failure is a disorder characterized by the inability of the liver to metabolize chemicals in the body. Laboratory test results reveal abnormal plasma levels of ammonia, bilirubin, lactic dehydrogenase, and alkaline phosphatase. Hepatic failure is graded as follows:

Grade 1: Not available

Grade 2: Not available

Grade 3: Asterixis or mild encephalopathy or limiting self-care ADL

Grade 4: Moderate to severe encephalopathy or coma or life-threatening consequences

Grade 5: Death

[0125] Ascites is disorder characterized by accumulation of serous or hemorrhagic fluid in the peritoneal cavity and is graded as follows:

Grade 1: Asymptomatic or clinical or diagnostic observations only or intervention not indicated

Grade 2: Symptomatic or medical intervention indicated

Grade 3: Severe symptoms or invasive intervention indicated

Grade 4: Life-threatening consequences or urgent operative intervention indicated

Grade 5: Death

[0126] Adverse reactions led to dose reduction or interruption in about 62% of patients receiving LENVIMA. The most common adverse reactions (>5%) resulting in dose reduction or interruption of LENVIMA were fatigue (9%), decreased appetite (8%), diarrhea (8%), proteinuria (7%), hypertension (6%), and palmar-plantar erythrodysesthesia syndrome (5%).

[0127] Proteinuria is disorder characterized by laboratory test results that indicate the presence of excessive protein in the urine. It is predominantly albumin, but also globulin, and is graded as follows:

Grade 1: 1+ proteinuria; urinary protein <1.0 g/24 hrs

Grade 2: Adults: 2+ proteinuria; urinary protein 1.0 - 3.4 g/24 hrs; Pediatric: urine P/C (Protein/Creatinine) ratio 0.5 -1.9

Grade 3: Adults: urinary protein ≥3.5 g/24 hrs; Pediatric: urine P/C >1.9

Grade 4: Not Available

Grade 5: Not Available

[0128] Palmar-plantar erythrodysesthesia syndrome is a disorder characterized by redness, marked discomfort, swelling, and tingling in the palms of the hands or the soles of the feet, and is graded as follows:

Grade 1: Minimal skin changes or dermatitis (e.g., erythema, edema, or hyperkeratosis)

without pain

Grade 2: Skin changes (e.g., peeling, blisters, bleeding, edema, or hyperkeratosis) with pain; limiting instrumental ADL

Grade 3: Severe skin changes (e.g., peeling, blisters, bleeding, edema, or hyperkeratosis) with pain; limiting self-care ADL

Grade 4: Not Available

Grade 5: Not Available

[0129] Treatment discontinuation due to adverse reactions occurred in 20% of patients in the

LENVIMA-treated group. The most common adverse reactions ($\geq 1\%$) resulting in discontinuation of LENVIMA were fatigue (1%), hepatic encephalopathy (1%), hyperbilirubinemia (1%), and hepatic failure (1%).

[0130] Hyperbilirubinemia (“Blood bilirubin increased”) is a finding based on laboratory test results that indicate an abnormally high level of bilirubin in the blood. Excess bilirubin is associated with jaundice.

Grade 1: > 1.0 ULN - $1.5 \times$ ULN (upper limit of the normal range (ULN)).

Grade 2: >1.5 ULN - $3.0 \times$ ULN

Grade 3: >3.0 ULN - $10.0 \times$ ULN

Grade 4: $>10.0 \times$ ULN

Grade 5: Not Available

[0131] This disclosure provides dose modifications for therapy comprising lenvatinib or a pharmaceutically acceptable salt thereof upon the occurrence of a treatment-emergent adverse event(s) during the course of treatment. In certain embodiments, a subject who has a baseline body weight of 60 kg or greater and who has a HCC is administered a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day. In other embodiments, a subject who has a baseline body weight of less than 60 kg and who has a HCC is administered a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day. In some embodiments, if a subject who has HCC and has moderate hepatic impairment classified in Child-Pugh class B under the Child-Pugh Classification, then the subject is administered a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose 8 mg/day regardless of the body weight of the subject.

[0132] Exemplary recommendations for dose interruption, reduction, and discontinuation of LENVIMA for some adverse reactions are listed in Table I below.

[0133]

[Table 1]

Table I. Dose Modifications for Exemplary Adverse Reactions

Adverse Reaction	Severity^a	Dose Modifications for LENVIMA
Hypertension	Grade 3	• Withhold for Grade 3 that persists despite optimal antihypertensive therapy. • Resume at reduced dose when hypertension is controlled at less than or equal to Grade 2.
	Grade 4	• Permanently discontinue for life-threatening hypertension.
Cardiac Dysfunction	Grade 3	• Withhold until improves to Grade 0 or 1 or baseline. • Resume at a reduced dose or discontinue depending on the severity and persistence of adverse reaction.
	Grade 4	• Permanently discontinue.
Arterial Thromboembolic Event	Any Grade	• Permanently discontinue.
Hepatotoxicity	Grade 3 or 4	• Withhold until improves to Grade 0 to 1 or baseline. • Either resume at a reduced dose or discontinue depending on severity and persistence of hepatotoxicity. • Permanently discontinue for hepatic failure.
Renal Failure or Impairment	Grade 3 or 4	• Withhold until improves to Grade 0 to 1 or baseline. • Resume at a reduced dose or discontinue depending on severity and persistence of renal impairment.

[Table 2]

Table I. (cont.) Dose Modifications for Exemplary Adverse Reactions

Adverse Reaction	Severity ^a	Dose Modifications for LENVIMA
Proteinuria	2 g or greater proteinuria in 24 hours	- Withhold until less than or equal to 2 grams of proteinuria per 24 hours. - Resume at a reduced dose. - Discontinue for nephrotic syndrome.
Gastrointestinal Perforation	Any Grade	Permanently discontinue.
Fistula Formation	Grade 3 or 4	Permanently discontinue.
QT Prolongation	Greater than 500 ms or greater than 60 ms increase from baseline	- Withhold until improves to less than or equal to 480 ms or baseline. - Resume at a reduced dose.
Reversible Posterior Leukoencephalopathy Syndrome	Any Grade	- Withhold until fully resolved. - Resume at a reduced dose or discontinue depending on severity and persistence of neurologic symptoms.
Other Adverse	Persistent or intolerable Grade 2 or 3 adverse reaction	- Withhold until improves to Grade 0 or 1 or baseline - Resume at reduced dose.
	Grade 4 laboratory abnormality	- Withhold until improves to Grade 0 or 1 or baseline - Resume at reduced dose.
	Grade 4 adverse reaction	Permanently discontinue.

^a National Cancer Institute Common Terminology Criteria for Adverse Events, version 4.0.**[0134] Non-Hematologic Toxicity**

In some cases, the subject may develop a Grade 1 or tolerable Grade 2 adverse reaction (e.g., nonhematological toxicity) after being administered the first dosage regimen. In such instances, treatment of the subject can continue without any changes to the first dosage regimen. Following or during treatment period with the first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day or 8 mg/day, if the human subject does not develop an intolerable Grade 2 or Grade 3 adverse reaction or non-life-threatening Grade 4 laboratory abnormality (e.g., nonhematological toxicity), the dosage regimen can be maintained without any changes to the first dosage regimen.

[0135] In some embodiments, the subject develops an intolerable Grade 2 or Grade 3 adverse reaction or non-life-threatening Grade 4 laboratory abnormality (e.g., non-hematologic toxicity) during the period of treatment with the first dosage regimen that is related to lenvatinib toxicity. In certain instances, the subject develops a Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, or 20 weeks after the administration of the first dosage regimen. In one embodiment, the subject develops an intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality within 12 weeks after the administration of the first dosage regimen. In another embodiment, the subject develops an intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality within 16 weeks after the administration of the first dosage regimen. In certain instances, the intolerable Grade 2 or Grade 3 non-hematologic toxicity is Grade 3 hypertension, Grade 2 hypertension, Grade 3 fatigue, Grade 2 fatigue, Grade 3 palmar-plantar erythrodysesthesia, Grade 2 palmar-plantar erythrodysesthesia, Grade 3 diarrhea, Grade 2 diarrhea, Grade 3 decreased appetite, Grade 2 decreased appetite, Grade 3 fatigue, Grade 2 fatigue, Grade 3 arthralgia, Grade 2 arthralgia, Grade 3 myalgia, Grade 2 myalgia, Grade 3 decreased weight, Grade 2 decreased weight, Grade 2 alopecia, Grade 3 dysphonia, Grade 2 dysphonia, Grade 3 nausea, Grade 2 nausea, Grade 3 abdominal pain, Grade 2 abdominal pain, Grade 3 QT/QTc interval prolongation (Electrocardiogram QT corrected interval prolonged), Grade 2 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 2 hypothyroidism, Grade 3 vomiting, Grade 2 vomiting, Grade 3 constipation, Grade 2 constipation, Grade 3 rash, and Grade 2 rash. In specific cases, the persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity is Grade 3 hypertension, Grade 2 hypertension, Grade 3 fatigue, Grade 2 fatigue, Grade 3 diarrhea, Grade 2 diarrhea, Grade 3 decreased appetite, Grade 2 decreased appetite, Grade 3 arthralgia, Grade 2 arthralgia, Grade 3 myalgia, Grade 2 myalgia, Grade 3 decreased weight, or Grade 2 decreased weight. In certain instances, Grade 2 toxicities can be determined to be tolerable or intolerable by both the subject and healthcare provider. In certain instances, the Grade 4 laboratory abnormality is Grade 4 increase in aspartate aminotransferase, Grade 4 increase in alanine aminotransferase, Grade 4 increase in alkaline phosphatase, Grade 4 hypokalemia, Grade 4 hyponatremia, Grade 4 hypoglycemia, Grade 4 increase in blood bilirubin, or Grade 4 increase in gamma glutamyl transferase. In the above embodiments, the healthcare provider can determine whether the Grade 4 laboratory abnormality is life-threatening or not.

- [0136] If the subject develops an intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality, after being administered the first dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the first dosage regimen and administer to the subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg/day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the second dosage regimen is administered after interruption of the first dosage regimen and after the nonhematologic toxicity observed after the first dosage regimen is resolved to Grade 0-1 or baseline. In some instances, the first dosage regimen is terminated only after commencement of medical management of the intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality. In specific embodiments, the intolerable Grade 2 or Grade 3 adverse reaction is hypertension, fatigue, diarrhea, arthralgia/myalgia, decreased appetite, or decreased weight. If the Grade 3 non-hematologic toxicity is hypertension, in one embodiment, the subject is provided anti-hypertensive therapy and treatment with lenvatinib or a pharmaceutically acceptable salt thereof is resumed at a lower dose (e.g., 8 or 4 mg/day) when hypertension is controlled at less than or equal to Grade 1; however, therapy with lenvatinib or a pharmaceutically acceptable salt thereof is discontinued for life-threatening hypertension.
- [0137] In some cases, even after administration of the second dosage regimen, a subject may develop an adverse reaction (e.g., nonhematologic toxicity). In certain instances, the subject develops an intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the second dosage regimen. The nonhematologic toxicity after the second dosage regimen may be the same as, or different from, the nonhematologic toxicity after the first dosage regimen. The nonhematologic toxicity after the second dosage regimen may be an intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality. In some instances, the nonhematologic toxicity is hypertension. In some instances, the non-hematologic toxicity is fatigue. In some instances, the nonhematologic toxicity is arthralgia. In some instances, the nonhematologic toxicity is myalgia. In some instances, the nonhematologic toxicity is diarrhea. In some instances, the non-hematologic toxicity is decreased appetite. In some instances, the nonhematologic toxicity is decreased weight.

[0138] If the subject develops an intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality, after being administered the second dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the second dosage regimen and administer to the subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg every other day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the third dosage regimen is administered after interruption of the second dosage regimen and after the non-hematologic toxicity observed after the second dosage regimen is resolved to Grade 0-1 or baseline. In some instances, the second dosage regimen is terminated only after commencement of medical management of the intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality. In specific embodiments, the intolerable Grade 2 or Grade 3 nonhematologic toxicity is hypertension, fatigue, diarrhea, decreased appetite, arthralgia/myalgia, or decreased weight. If the Grade 3 nonhematologic toxicity is hypertension, in one embodiment, the subject is provided antihypertensive therapy and treatment with lenvatinib or a pharmaceutically acceptable salt thereof is resumed at a lower dose (e.g., 4 mg every other day) when hypertension is controlled at less than or equal to Grade 1; however, therapy with lenvatinib or a pharmaceutically acceptable salt thereof is discontinued for life-threatening hypertension.

[0139] In some cases, even after administration of the third dosage regimen, a subject may develop an adverse reaction (e.g., nonhematologic toxicity). In certain instances, the subject develops an intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the third dosage regimen. The nonhematologic toxicity after the third dosage regimen may be the same as, or different from, the non-hematologic toxicity after the second dosage regimen. The nonhematologic toxicity after the third dosage regimen may be an intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality. In some instances, the nonhematologic toxicity is hypertension. In some instances, the non-hematologic toxicity is fatigue. In some instances, the nonhematologic toxicity is arthralgia. In some instances, the nonhematologic toxicity is myalgia. In some instances, the nonhematologic toxicity is diarrhea. In some instances, the non-hematologic toxicity is decreased appetite. In some instances, the nonhematologic

toxicity is decreased weight.

- [0140] If the subject develops an intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality, after being administered the third dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg every other day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the third dosage regimen and administer to the subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day (if the patient has a baseline body weight of greater than or equal to 60 kg). If the patient has a baseline body weight of less than 60 kg, in certain instances, lenvatinib therapy may be discontinued. In certain instances, the fourth dosage regimen is administered after interruption of the third dosage regimen and after the nonhematologic toxicity observed after the third dosage regimen is resolved to Grade 0-1 or baseline. In some instances, the third dosage regimen is terminated only after commencement of medical management of the intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality. In specific embodiments, the persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is hypertension, fatigue, diarrhea, decreased appetite, arthralgia/myalgia, or decreased weight. If the Grade 3 nonhematologic toxicity is hypertension, in one embodiment, the subject is provided antihypertensive therapy and treatment with lenvatinib or a pharmaceutically acceptable salt thereof is resumed at a lower dose when hypertension is controlled at less than or equal to Grade 1; however, therapy with lenvatinib or a pharmaceutically acceptable salt thereof is discontinued for life-threatening hypertension.
- [0141] In some embodiments, if the subject develops a Grade 4 nonhematologic toxicity excluding non-life-threatening Grade 4 laboratory abnormality after being administered the first, second, third, or fourth dosage regimen, the healthcare provider can terminate administration of the dosage regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding non-life-threatening Grade 4 laboratory abnormality.
- [0142] In some embodiments, a Grade 3 laboratory abnormality that is not clinically relevant based on a judgment of healthcare provider is excluded from a Grade 3 non-hematologic toxicity.
- [0143] In the above embodiments, a Grade 3 proteinuria is excluded from a Grade 3 non-hematologic toxicity to those dosage regimens.
- [0144] Grades 1 to 3 Hematologic Toxicity and/or Grades 1 to 3 Proteinuria
In some embodiments, the subject may develop a Grade 1 or Grade 2 hematologic toxicity or Grade 1 or Grade 2 proteinuria during the period of treatment with the first

dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg)) that is related to lenvatinib toxicity. In such instances, the subject can continue with treatment with the first dosage regimen. However, in some cases, the subject may develop a Grade 3 adverse reaction (e.g., hematologic toxicity or proteinuria) during the period of treatment with the first dosage regimen that is related to lenvatinib toxicity. In certain instances, the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, or 20 weeks after the administration of the first dosage regimen. In one embodiment, the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria within 12 weeks after the administration of the first dosage regimen. In another embodiment, the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria within 16 weeks after the administration of the first dosage regimen. In certain instances, the Grade 3 hematologic toxicity is Grade 3 thrombopenia (thrombocytopenia; platelet count decreased), Grade 3 anemia (hemoglobin decreased), Grade 3 decrease in white blood cell count (leukocyte count decreased; white blood cell decreased), Grade 3 neutropenia (neutrophil count decreased), or Grade 3 lymphocytopenia (lymphocyte count decreased).

[0145] If the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria, after being administered the first dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the first dosage regimen and administer to the subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the second dosage regimen is administered after interruption of the first dosage regimen and after the nonhematologic toxicity observed after the first dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the first dosage regimen is terminated only after commencement of medical management of the Grade 3 hematologic toxicity. In specific embodiments, the Grade 3 hematologic toxicity is Grade 3 thrombopenia (thrombocytopenia), Grade 3 anemia, Grade 3 decrease in white blood cell count, Grade 3 neutropenia, or Grade 3 lymphocytopenia.

[0146] In some cases, even after administration of the second dosage regimen, a subject may develop a Grade 3 hematologic toxicity or Grade 3 proteinuria. In certain instances, the

subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the second dosage regimen. The Grade 3 hematologic toxicity or Grade 3 proteinuria after the second dosage regimen may be the same as, or different from, the Grade 3 hematologic toxicity or Grade 3 proteinuria after the first dosage regimen. The hematologic toxicity or proteinuria after the second dosage regimen may be a Grade 3 hematologic toxicity or Grade 3 proteinuria.

[0147] If the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria, after being administered the second dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the second dosage regimen and administer to the subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg/day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the third dosage regimen is administered after interruption of the second dosage regimen and after the Grade 3 hematologic toxicity or Grade 3 proteinuria observed after the second dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the second dosage regimen is terminated only after commencement of medical management of the Grade 3 hematologic toxicity or Grade 3 proteinuria.

[0148] In some cases, even after administration of the third dosage regimen, a subject may develop a Grade 3 hematologic toxicity or Grade 3 proteinuria. In certain instances, the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the third dosage regimen. The Grade 3 hematologic toxicity or Grade 3 proteinuria after the third dosage regimen may be the same as, or different from, the Grade 3 hematologic toxicity or Grade 3 proteinuria after the second dosage regimen. The Grade 3 hematologic toxicity or Grade 3 proteinuria after the third dosage regimen may be a Grade 3 hematologic toxicity or Grade 3 proteinuria.

[0149] If the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria, after being administered the third dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the third dosage regimen and administer to the subject a fourth dosage regimen comprising lenvatinib or a pharma-

ceutically acceptable salt thereof at a dose of 4 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg every other day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the fourth dosage regimen is administered after interruption of the third dosage regimen and after the Grade 3 hematologic toxicity or Grade 3 proteinuria observed after the third dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the third dosage regimen is terminated only after commencement of medical management of the Grade 3 hematologic toxicity or Grade 3 proteinuria.

[0150] In some cases, even after administration of the fourth dosage regimen, a subject may develop a Grade 3 hematologic toxicity or Grade 3 proteinuria. In certain instances, the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the third dosage regimen. The Grade 3 hematologic toxicity or Grade 3 proteinuria after the third dosage regimen may be the same as, or different from, the Grade 3 hematologic toxicity or Grade 3 proteinuria after the second dosage regimen. The Grade 3 hematologic toxicity or Grade 3 proteinuria after the third dosage regimen may be a Grade 3 hematologic toxicity or Grade 3 proteinuria.

[0151] If the subject develops a Grade 3 hematologic toxicity or Grade 3 proteinuria, after being administered the fourth dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg every other day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the fourth dosage regimen and administer to the subject a fifth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day (if the patient has a baseline body weight of greater than or equal to 60 kg). In some cases, if the patient has a baseline body weight of less than 60 kg, lenvatinib treatment may be discontinued. In certain instances, the fifth dosage regimen is administered after interruption of the fourth dosage regimen and after the Grade 3 hematologic toxicity or Grade 3 proteinuria observed after the third dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the fourth dosage regimen is terminated only after commencement of medical management of the Grade 3 hematologic toxicity or Grade 3 proteinuria.

[0152] In some embodiments, a Grade 3 laboratory abnormality that is not clinically relevant based on a judgment of healthcare provider is excluded from a Grade 3 hematologic toxicity.

[0153] Grade 4 Hematologic Toxicity

In some embodiments, the subject may develop a Grade 4 hematologic toxicity

during the period of treatment with the first dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg)) that is related to lenvatinib toxicity. In certain instances, the subject develops the Grade 4 hematologic toxicity within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, or 20 weeks after the administration of the first dosage regimen. In one embodiment, the subject develops Grade 4 hematologic toxicity within 12 weeks after the administration of the first dosage regimen. In another embodiment, the subject develops the Grade 4 hematologic toxicity within 16 weeks after the administration of the first dosage regimen. In certain instances, the Grade 4 hematologic toxicity is Grade 4 thrombopenia (thrombocytopenia; platelet count decreased), Grade 4 anemia, Grade 4 decrease in white blood cell count, Grade 4 neutropenia, or Grade 4 lymphocytopenia.

[0154] If the subject develops the Grade 4 hematologic toxicity after being administered the first dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 12 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 8 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the first dosage regimen and administer to the subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg/day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the second dosage regimen is administered after interruption of the first dosage regimen and after the Grade 4 hematologic toxicity observed after the first dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the first dosage regimen is terminated only after commencement of medical management of the Grade 4 hematologic toxicity. In specific embodiments, the Grade 4 hematologic toxicity is Grade 4 thrombopenia (thrombocytopenia), Grade 4 anemia, Grade 4 decrease in white blood cell count, Grade 4 neutropenia, or Grade 4 lymphocytopenia.

[0155] In some cases, even after administration of the second dosage regimen, a subject may develop a Grade 4 hematologic toxicity. In certain instances, the subject develops a Grade 4 hematologic toxicity within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the second dosage regimen. The Grade 4 hematologic toxicity after the second dosage regimen may be the same as, or different from, the Grade 4 hematologic toxicity after the first dosage regimen. The hematologic toxicity after the second

dosage regimen may be a persistent and intolerable Grade 4 hematologic toxicity.

[0156] If the subject develops a Grade 4 hematologic toxicity after being administered the second dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg/day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the second dosage regimen and administer to the subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg every other day (if the patient has a baseline body weight of less than 60 kg). In certain instances, the third dosage regimen is administered after interruption of the second dosage regimen and after the Grade 4 hematologic toxicity observed after the second dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the second dosage regimen is terminated only after commencement of medical management of the Grade 4 hematologic toxicity.

[0157] In some cases, even after administration of the third dosage regimen, a subject may develop a Grade 4 hematologic toxicity. In certain instances, the subject develops a Grade 4 hematologic toxicity within 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, or 12 weeks after the administration of the third dosage regimen. The Grade 4 hematologic toxicity after the third dosage regimen may be the same as, or different from, the Grade 4 hematologic toxicity after the second dosage regimen. The Grade 4 hematologic toxicity after the third dosage regimen may be a persistent and intolerable Grade 4 hematologic toxicity.

[0158] If the subject develops a Grade 4 hematologic toxicity after being administered the third dosage regimen (i.e., lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day (if the patient has a baseline body weight of greater than or equal to 60 kg) or 4 mg every other day (if the patient has a baseline body weight of less than 60 kg)), the healthcare provider can terminate the third dosage regimen and administer to the subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day (if the patient has a baseline body weight of greater than or equal to 60 kg). If the patient has a baseline body weight of less than 60 kg, lenvatinib therapy can, in some embodiments, be discontinued. In certain instances, the fourth dosage regimen is administered after interruption of the third dosage regimen and after the Grade 4 hematologic toxicity observed after the third dosage regimen is resolved to Grade 0-2 or baseline. In some instances, the third dosage regimen is terminated only after commencement of medical management of the Grade 4 hematologic toxicity.

[0159] The dose modifications discussed above can be helpful in permitting a subject who develops an adverse reaction to the lenvatinib therapy (e.g., a nonhematologic toxicity,

a hematologic toxicity, proteinuria, or a laboratory abnormality) to continue with and benefit from the lenvatinib therapy.

[0160] Table II below lists exemplary dose reductions for LENVIMA due to adverse reactions after administration of a first dosage regimen discussed above.

[0161] [Table 3]

Table II. Exemplary Dose Modifications for LENVIMA

Indication	First Dosage Reduction To	Second Dosage Reduction To	Third Dosage Reduction To
HCC (<i>e.g.</i> , unresectable HCC)			
Actual weight 60 kg or greater	8 mg once daily	4 mg once daily	4 mg every other day
Actual weight less than 60 kg	4 mg once daily	4 mg every other day	Discontinue

[0162] The following examples are provided to better illustrate the claimed invention and are not to be interpreted as limiting the scope of the invention. To the extent that specific materials are mentioned, it is merely for purposes of illustration and is not intended to limit the invention. One skilled in the art can develop equivalent means or reactants without the exercise of inventive capacity and without departing from the scope of the invention.

[0163] Examples

Example 1: Lenvatinib for Unresectable Hepatocellular Carcinoma

This Example describes the open-label phase 3 noninferiority study using lenvatinib in unresectable HCC. In this study 478 patients were randomized to lenvatinib (body weight ≥ 60 kg: 12 mg/day; < 60 kg: 8 mg/day) and 476 to sorafenib 400 mg twice daily. The primary endpoint was overall survival. Secondary endpoints included progression-free survival, time to progression, objective response rate, safety, and quality of life.

[0164] Materials & Methods

Study Eligibility

Patients who were eligible for enrollment had unresectable HCC with diagnosis confirmed histologically or cytologically or with diagnosis confirmed clinically in accordance with the American Association for the Study of Liver Diseases criteria. Included patients also had ≥ 1 measureable target lesion, based on modified Response Evaluation Criteria in Solid Tumors (Lencioni 2010); Barcelona Clinic Liver Cancer stage B or C categorization (Bruix Hepatology 2011); Child-Pugh class A; and Eastern Cooperative Oncology Group performance status ≤ 1 . All eligible patients had controlled blood pressure ($\leq 150/90$ mm Hg), and adequate organ function. Patients with $\geq 50\%$ liver occupation, clear bile duct invasion, or portal vein invasion at the

main portal vein were excluded. Patients also were excluded if that had received prior systemic therapy for HCC.

[0165] Study Oversight

The study was approved by all relevant institutional review boards and was conducted in accordance with the Declaration of Helsinki and local laws. The trial was registered before the start of patient enrollment. All patients provided written informed consent before undergoing any study-specific procedures. The study was overseen by an independent data monitoring committee.

[0166] Study Design

This multicenter phase 3 randomized open-label noninferiority study was conducted throughout the Asia-Pacific, European, and North American regions. Patients were recruited from March 1, 2013 through July 30, 2016. Randomization was stratified according to region (Asia-Pacific or Western regions), macroscopic portal vein invasion and/or extrahepatic spread (yes or no), Eastern Cooperative Oncology Group performance status (0 or 1), and body weight (<60 kg or ≥60 kg). Within stratification factors, patients were randomly assigned (1:1) to receive oral lenvatinib at a dosage of 12 mg per day (for body weight ≥60 kg) or 8 mg per day (for body weight <60 kg) or sorafenib at a dosage of 400 mg twice daily in 28-day cycles. Dosage interruptions and reductions for lenvatinib-related toxicities (to 8 and 4 mg per day, or 4 mg every other day) were permitted. Modifications to sorafenib dosage were implemented according to prescribing information in each region.

[0167] Endpoints and Assessments

The primary endpoint was overall survival. Secondary endpoints included progression-free survival, time to progression, objective response rate, and quality of life as measurements including use of EORTC QLQ-C30 (Cocks, J, Clin. Oncol., 29:89-96, 2011, Giesinger, J., Clin. Epidemiol., 69:79-88, 2016) and HCC-specific EORTC QLQ-HCC18 (Chie, Hepatology, 55(4):1122-9, 2012) health questionnaires.

[0168] Tumors were evaluated in accordance with mRECIST (Lencioni R., Semin Liver Dis., 30(1):52-60, 2010); RECIST 1.1 was applied for nonhepatic lesions (Eisenhauer, Eur J Cancer, 45(2):228-47, 2009). The liver was examined with computed tomography or magnetic resonance imaging was performed using a triphasic scanning technique. Assessments were performed every 8 weeks until disease progression. Quality of life questionnaires were administered at baseline, on day 1 of each subsequent treatment cycle, and at the off-treatment visit.

[0169] Safety assessments included recording of vital signs, hematologic and biochemical laboratory testing, urinalysis, and electrocardiography. Adverse events were graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events version 4.0 (CTCAE v4.0, NCI 2013).

[0170] Statistical Analysis

The primary endpoint of overall survival was first tested for noninferiority then for superiority. The required number of events for the primary analysis was 700 deaths, based on the full analysis set. The HR and its 95% confidence interval (CI) were estimated from a Cox proportional hazard model with treatment group as a factor and with the analysis stratified according to the same factors applied for randomization. The noninferiority margin was set at 1.08 based on previous phase 3 trials of sorafenib (Llovet NEJM 2008, Cheng Lancet Oncol 2009).

[0171] A fixed-sequence procedure was followed to control the overall type I error rate of analyses for both the primary and secondary efficacy endpoints at $\alpha=0.05$ (2-sided). After noninferiority was declared, secondary efficacy endpoints were tested. Differences in progression-free survival and time to progression were evaluated using a stratified log-rank test with randomization stratification factors, with the associated HR and its 95% CI. A difference in the objective response rate was evaluated using the Cochran-Mantel-Haenszel chi-square test with randomized stratification factors as strata, with associated odds ratio and its 95% CI. To assess futility, 2 interim analyses (at 30% and 70% of the target number of events) were performed using Bayesian predictive probability in a noninferiority design by the independent data monitoring committee.

[0172] Patients

A total of 954 patients from 20 countries were randomly assigned to receive lenvatinib (478 patients) or sorafenib (476 patients) (Figure 5). The efficacy analysis followed the intent to treat principle. Only patients who received treatment (lenvatinib, 476 patients; sorafenib, 475 patients) were included in the safety analysis. Patient characteristics at baseline were well balanced between treatment groups, with the exception of baseline hepatitis C and alpha-fetoprotein level (Table 1).

[0173]

[Table 4]

Table 1. *Demographic and Disease Characteristics at Baseline.*

	Lenvatinib (N = 478)	Sorafenib (N = 476)	Total (N = 954)
Age - yr			
Mean	61.3	61.2	61.3
Standard Deviation	11.7	12.0	11.8
Age group (yr) - no. (%)			
<65	270 (56.5)	283 (59.5)	553 (58.0)
≥65 to <75	150 (31.4)	126 (26.5)	276 (28.9)
≥75	58 (12.1)	67 (14.1)	125 (13.1)
Sex - no. (%)			
Male	405 (84.7)	401 (84.2)	806 (84.5)
Female	73 (15.3)	75 (15.8)	148 (15.5)
Region - no. (%)			
Western	157 (32.8)	157 (33.0)	314 (32.9)
Asia-Pacific	321 (67.2)	319 (67.0)	640 (67.1)
Race - no. (%)			
White	135 (28.2)	141 (29.6)	276 (28.9)
Asian	334 (69.9)	326 (68.5)	660 (69.2)
Body weight (kg) - no. (%)			
<60	153 (32.0)	146 (30.7)	299 (31.3)
≥60	325 (68.0)	330 (69.3)	655 (68.7)
ECOG performance status - no. (%)			
0	304 (63.6)	301 (63.2)	605 (63.4)
1	174 (36.4)	175 (36.8)	349 (36.6)
Child-Pugh class - no. (%)			
CP-A	475 (99.4)	471 (98.9)	946 (99.2)
CP-B	3 (0.6)	5 (1.1)	8 (0.8)
Macroscopic portal vein invasion - no. (%)			
Yes	109 (22.8)	90 (18.9)	199 (20.9)
No	369 (77.2)	386 (81.1)	755 (79.1)
Extrahepatic spread - no. (%)			
Yes	291 (60.9)	295 (62.0)	586 (61.4)
No	187 (39.1)	181 (38.0)	368 (38.6)
Macroscopic portal vein invasion, extrahepatic spread, or both - no. (%)			
Yes	329 (68.8)	336 (70.6)	665 (69.7)
No	149 (31.2)	140 (29.4)	289 (30.3)
Underlying cirrhosis based on medical history - no. (%)			
Yes	243 (50.8)	231 (48.5)	474 (49.7)
No	235 (49.2)	245 (51.5)	480 (50.3)

[Table 5]

Table 1. (cont.) *Demographic and Disease Characteristics at Baseline.*

	Lenvatinib (N = 478)	Sorafenib (N = 476)	Total (N = 954)
Barcelona Clinic Liver Cancer stage - no. (%)			
B (intermediate stage)	104 (21.8)	92 (19.3)	196 (20.5)
C (advanced stage)	374 (78.2)	384 (80.7)	758 (79.5)
Involved disease sites - no. (%)			
Liver	441 (92.3)	430 (90.3)	871 (91.3)
Lung	163 (34.1)	144 (30.3)	307 (32.2)
Involved disease sites per patient - no. (%)			
1	207 (43.3)	207 (43.5)	414 (43.4)
2	167 (34.9)	183 (38.4)	350 (36.7)
≥3	103 (21.5)	86 (18.1)	189 (19.8)
Etiology of chronic liver disease - no. (%)			
Hepatitis B	251 (52.5)	228 (47.9)	479 (50.2)
Hepatitis C	91 (19.0)	126 (26.5)	217 (22.7)
Alcohol	36 (7.5)	21 (4.4)	57 (6.0)
Other	38 (7.9)	32 (6.7)	70 (7.3)
Unknown	62 (13.0)	69 (14.5)	131 (13.7)
Baseline alpha-fetoprotein level - ng/mL			
No. of patients	471	463	934
Mean	17507.7	16678.5	17096.5
Standard deviation	105137.4	94789.5	100088.8
Median	133.1	71.2	89.0
Range	0–1567470	0–1446396	0–1567470
Baseline alpha-fetoprotein level group (ng/mL) - no. (%)			
<200	255 (53.3)	286 (60.1)	541 (56.7)
≥200	222 (46.4)	187 (39.3)	409 (42.9)
Missing	1 (0.2)	3 (0.6)	4 (0.4)
Concomitant systemic antiviral therapy for Hepatitis B or C - no. (%)	163 (34.1)	149 (31.3)	312 (32.7)
Prior therapy - no. (%)			
Prior anticancer procedures	327 (68.4)	344 (72.3)	671 (70.3)
Radiotherapy	49 (10.3)	60 (12.6)	109 (11.4)

ECOG, Eastern Cooperative Oncology Group.

At the time of data cutoff (November 13, 2016) the median duration of follow-up was 27.7 months in the lenvatinib group and 27.2 months in the sorafenib group.

[0174] Efficacy

Lenvatinib demonstrated noninferiority in overall survival compared with sorafenib. The median overall survival was 13.6 months (95% CI, 12.1 to 14.9) with lenvatinib compared with 12.3 months (95% CI, 10.4 to 13.9) with sorafenib (HR: 0.92; 95% CI, 0.79 to 1.06) (Figure 1). Median overall survival for the lenvatinib treatment arm was numerically longer in all the prespecified strata, with the exception of the patients from Western regions subgroup; and was the same in patients who had neither macroscopic portal vein invasion nor extrahepatic spread (Figure 3). Patients with baseline alpha-fetoprotein <200 ng/mL had longer overall survival than those with alpha-fetoprotein

≥ 200 ng/mL (Figure 3). The HR was <1 favoring lenvatinib in both alpha-fetoprotein subgroups, but there were more patients with baseline alpha-fetoprotein level <200 ng/mL in the sorafenib arm (60.1%) compared with the lenvatinib arm (53.3%, Table 1).

[0175] Lenvatinib demonstrated a statistically significant improvement compared to sorafenib in all secondary efficacy endpoints as determined by tumor assessment based on mRECIST. Median progression-free survival for lenvatinib was 7.4 months (95% CI, 6.9 to 8.8 months) compared with 3.7 months (95% CI, 3.6 to 4.6 months) with sorafenib (HR: 0.66; 95% CI, 0.57 to 0.77; $P < 0.001$) (Figure 2). In addition, progression-free survival in each of the prespecified subgroups was longer with lenvatinib compared with sorafenib (Figure 4). The median time to progression was 8.9 months (95% CI, 7.4 to 9.2 months) for patients in the lenvatinib group compared with 3.7 months (95% CI, 3.6 to 5.4 months) for patients in the sorafenib group (HR: 0.63; 95% CI, 0.53 to 0.73; $P < 0.001$) (Table 2 and Figure 6).

[0176] [Table 6]

Table 2. Efficacy Measures.

Outcome	Lenvatinib (N = 478)	Sorafenib (N = 476)	Hazard Ratio (95% CI)
Median (95% CI) overall survival - mo	13.6 (12.1–14.9)	12.3 (10.4–13.9)	0.92 (0.79–1.06)
Median (95% CI) progression-free survival - mo	7.4 (6.9–8.8)	3.7 (3.6–4.6)	0.66 (0.57–0.77) $P < 0.001$
Median (95% CI) time to progression - mo	8.9 (7.4–9.2)	3.7 (3.6–5.4)	0.63 (0.53–0.73) $P < 0.001$
Objective response rate* - no. (%)	115 (24.1)	44 (9.2)	3.13†
95% CI	20.2–27.9	6.6–11.8	(2.15–4.56)
Complete response	6 (1.3)	2 (0.4)	$P < 0.001$
Partial response	109 (22.8)	42 (8.8)	
Stable disease	246 (51.5)	244 (51.3)	
Durable stable disease lasting ≥ 23 wk	167 (34.9)	139 (29.2)	
Progressive disease	71 (14.9)	147 (30.9)	
Unknown/not evaluable	46 (9.6)	41 (8.6)	
Disease control rate‡ - no. (%)			
95% CI	361 (75.5) 71.7–79.4	288 (60.5) 56.1–64.9	

*The objective response rate was determined according to mRECIST (modified Response Evaluation Criteria in Solid Tumors). †Odds ratio. ‡The disease control rate is computed as follows: complete response + partial response + stable disease. CI, confidence interval.

[0177] Median time to progression with lenvatinib treatment was also numerically longer in

each of the prespecified strata compared with sorafenib (Figures 7-10). Lenvatinib showed an objective response rate of 24.1% versus 9.2% for sorafenib (odds ratio, 3.13; 95% CI, 2.15 to 4.56; $P < 0.001$) (Table 2 and Figure 11). The odds ratio favored lenvatinib in all prespecified subgroups, with the exception of the alcohol etiology subgroup (Figures 12-14). Analysis for overall survival with stratification factors and other subgroups supports the robustness of the noninferiority result (Table 3).

[0178] [Table 7]

Table 3. Overall Survival with Stratification Factors Adjusted by Baseline Characteristics.

Baseline Characteristics	Hazard Ratio for (Lenvatinib/Sorafenib) (95% CI)
Overall	0.916 (0.789–1.064)
Age (<65, ≥65 to <75, ≥75 yr)	0.919 (0.791–1.067)
Sex (male, female)	0.916 (0.789–1.064)
Region (Asia-Pacific, Western)	0.915 (0.789–1.062)
Macroscopic portal vein invasion (yes, no)	0.910 (0.784–1.057)
Extrahepatic spread (yes, no)	0.915 (0.788–1.062)
Macroscopic portal vein invasion, extrahepatic spread or both (yes, no)	0.908 (0.783–1.054)
ECOG-PS (0, ≥1)	0.923 (0.795–1.071)
Body weight (<60 kg, ≥60 kg)	0.923 (0.796–1.071)
AFP at baseline (<200 ng/mL, ≥200 ng/mL)	0.856 (0.736–0.995)
Antiviral therapy for HCB or HCV (yes, no)	0.912 (0.785–1.059)
No. of disease sites at baseline (1, 2, ≥3)	0.878 (0.755–1.020)
Etiology (HBV, HCV, alcohol)	0.855 (0.721–1.013)
Underlying cirrhosis (yes, no)	0.916 (0.789–1.063)
BCLC staging (stage B, stage C)	0.918 (0.791–1.067)
Prior procedure (yes, no)	0.902 (0.777–1.048)

AFP, alpha-fetoprotein; BCLC, Barcelona Clinic Liver Cancer; ECOG-PS, Eastern Cooperative Oncology Group performance status; HBV, hepatitis B virus; HCV, hepatitis C virus.

[0179] Of note, 32.6% patients in the lenvatinib arm and 38.7% in the sorafenib arm received a post-study anticancer medication (including investigational therapy). Of these, 25.3% of patients in the lenvatinib arm and 11.8 % in the sorafenib arm, respectively, received sorafenib during survival follow-up. In the Western region, 26.1% of patients in the lenvatinib arm received any anticancer medication during survival follow-up versus 38.9% in the sorafenib arm (Table 4).

[0180]

[Table 8]

Table 4. *Post-study Anticancer Therapy During Survival Follow-up.*

	Lenvatinib			Sorafenib		
	Asia-Pacific Subgroup (N = 321)	Western Subgroup (N = 157)	Total (N = 478)	Asia-Pacific Subgroup (N = 319)	Western Subgroup (N = 157)	Total (N = 478)
Received any anticancer medication (not given for any procedure) during survival follow-up - no. (%)	115 (35.8)	41 (26.1)	156 (32.6)	123 (38.6)	61 (38.9)	184 (38.7)
Underwent any anticancer procedure during survival follow-up - no. (%)	111 (34.6)	11 (7.0)	122 (25.5)	112 (35.1)	18 (11.5)	130 (27.3)

In the lenvatinib arm, 7.0% of patients in the Western region had any anticancer procedure during follow-up compared with 11.0% of patients in the sorafenib arm in this region.

[0181] Safety and Side-effect Profile

Median duration of study treatment for patients in the lenvatinib group was longer than for patients in the sorafenib group (5.7 vs. 3.7 months). Treatment-emergent adverse events occurred in 98.7% of patients who received lenvatinib and 99.4% of patients who received sorafenib. Adjusted by patient-years, the adverse event rate was 18.9 in the lenvatinib group and 19.7 in the sorafenib group. Treatment-emergent adverse events of grade 3 or higher occurred in 75.0% of patients who received lenvatinib and 66.5% of patients who received sorafenib (adverse event rate: 3.2 vs. 3.3). The most common treatment-emergent adverse events among patients who received lenvatinib (at a dosage of 8 mg or 12 mg per day) were hypertension (8 mg per day, 43.0%; 12 mg per day, 41.8%), diarrhea (35.1%; 40.3%), decreased appetite (33.1%; 34.5%), and decreased weight (28.5%; 32.0%). In the sorafenib arm, the most common treatment-emergent adverse events were palmar-plantar erythrodysesthesia (52.4%), diarrhea (46.3%), hypertension (30.3%), and decreased appetite (26.7%) (Table 5).

[0182]

[Table 9]

Table 5. Adverse Events.

	Lenvatinib (N = 476)		Sorafenib (N = 475)	
Total treatment-related treatment-emergent adverse events—no. (%)	447 (93.9)		452 (95.42)	
Treatment-related treatment-emergent adverse events of grade ≥ 3 —no. (%)	270 (56.7)		231 (48.6)	
Serious treatment-emergent adverse events — no. (%)	205 (43.1)		144 (30.3)	
Treatment emergent adverse events occurring in $\geq 15\%$ of patients in either treatment group	Any grade	Grade ≥ 3	Any grade	Grade ≥ 3
<i>Number of patients (percent)</i>				
Palmar-plantar erythrodysesthesia	128 (26.9)	14 (2.9)	249 (52.4)	54 (11.4)
Diarrhea	184 (38.7)	20 (4.2)	220 (46.3)	20 (4.2)
Hypertension	201 (42.2)	111 (23.3)	144 (30.3)	68 (14.3)
Decreased appetite	162 (34.0)	22 (4.6)	127 (26.7)	6 (1.3)
Decreased weight	147 (30.9)	36 (7.6)	106 (22.3)	14 (2.9)
Fatigue	141 (29.6)	18 (3.8)	119 (25.1)	17 (3.6)
Alopecia	14 (2.9)	0 (0)	119 (25.1)	0 (0)
Proteinuria	117 (24.6)	27 (5.7)	54 (11.4)	8 (1.7)
Dysphonia	113 (23.7)	1 (0.2)	57 (12.0)	0 (0)
Nausea	93 (19.5)	4 (0.8)	68 (14.3)	4 (0.8)
Abdominal pain	81 (17.0)	8 (1.7)	87 (18.3)	13 (2.7)
Decreased platelet count	87 (18.3)	26 (5.5)	58 (12.2)	16 (3.4)
Elevated aspartate aminotransferase	65 (13.7)	24 (5.0)	80 (16.8)	38 (8.0)
Hypothyroidism	78 (16.4)	0 (0)	8 (1.7)	0 (0)
Vomiting	77 (16.2)	6 (1.3)	36 (7.6)	5 (1.1)
Constipation	76 (16.0)	3 (0.6)	52 (10.9)	0 (0)
Rash	46 (9.7)	0 (0)	76 (16.0)	2 (0.4)

[0183] Fatal adverse events occurred throughout treatment and appeared to occur at similar rates in both arms. Fatal adverse events determined by the investigator to be related to lenvatinib treatment occurred in 11 patients (2.3%) and included hepatic failure (3 patients), cerebral hemorrhage (3 patients), and respiratory failure (2 patients). In the sorafenib group, treatment-related fatal adverse events occurred in 4 patients (0.8%) and included tumor hemorrhage, ischemic stroke, respiratory failure, and sudden death

(1 event per patient).

[0184] Lenvatinib dose reduction, drug interruption, and discontinuation due to adverse events occurred in 184 (38.7%), 248 (52.1%), and 94 (19.7%) patients, respectively. In the sorafenib arm, dose reduction, drug interruption, and discontinuation due to adverse events occurred in 185 (38.9%), 193 (40.6%), and 69 (14.5%) patients, respectively. The mean lenvatinib dose intensity was 7.0 mg (87.7%) in the 8 mg/day group and 10.5 mg (87.5%) in the 12 mg/day group. The mean sorafenib dose intensity was 663.8 mg (83.0%).

[0185] Quality of Life

Baseline scores on the EORTC QLQ-C30 and EORTC QLQ-HCC18 health questionnaires were similar in the lenvatinib and sorafenib treatment groups (Figure 15). Following treatment, scores declined in both groups. The analysis of time to clinically meaningful deterioration showed that the role functioning, pain, and diarrhea from QLQ-C30 and nutrition and body image from QLQ-HCC18 deterioration was observed earlier in patients treated with sorafenib than with lenvatinib (nominal $P < 0.05$).

[0186] Patients who received lenvatinib experienced fewer instances of palmar-plantar erythrodysesthesia, diarrhea, and alopecia and more instances of hypertension, proteinuria, dysphonia, and hypothyroidism than did patients who received sorafenib. Although quality of life scores declined in both groups after treatment, a clinically meaningful delay in deterioration for multiple domains was observed with lenvatinib compared with sorafenib.

[0187] The median duration of lenvatinib treatment was 1.5 times longer than that of sorafenib, which may have contributed to the higher incidence of adverse events. When adjusted for treatment duration, almost all episodes were comparable for the lenvatinib and sorafenib arms. The dosages of lenvatinib for HCC (8 or 12 mg per day based on body weight) are lower than the lenvatinib dosage for radioiodine-refractory differentiated thyroid cancer (24 mg per day). In the phase 1 study of lenvatinib in HCC, patients with HCC who received 12 mg of lenvatinib per day and patients with solid tumors who received 25 mg of lenvatinib per days had similar lenvatinib plasma concentration at 24 hours, possibly because lenvatinib is metabolized in the liver. In the current phase 3 study, similar clinical activities and safety profiles were observed for both the 8 mg and 12 mg per day lenvatinib starting dosages.

[0188] In conclusion, the results of this study demonstrated noninferiority of lenvatinib in overall survival to sorafenib and statistically significant and clinically meaningful improvement in progression-free survival, time to progression, and objective response rate. The safety profiles of lenvatinib and sorafenib in this study appear consistent with the known safety profiles of these agents in HCC and no new safety signals were identified. Based on these results, lenvatinib may be a potential treatment option in

advanced HCC.

[0189] Example 2: Dose Interruption and Dose Reduction for Lenvatinib Toxicity

Lenvatinib capsules were taken orally, once daily (QD) in continuous 28-day cycles. The dose of lenvatinib was based on the subject's baseline body weight (BW) as shown below.

[0190] [Table 10]

Drug Name	Baseline BW	Strength	Oral Dose Form	Number Dispensed and Frequency
Lenvatinib	≥60 kg	12 mg	capsule	3 × 4-mg capsules, once daily at the same time each day
	<60 kg	8 mg	capsule	2 × 4-mg capsules, once daily at the same time each day

[0191] Lenvatinib toxicity was managed by treatment interruption, dose reduction, and/or treatment discontinuation. Dose adjustment for management of lenvatinib toxicity (with the exception of hypertension) was done in accordance with Table 6.

[0192]

[Table 11]

Table 6: Lenvatinib Dose Reductions and Interruption Instructions

Dose reductions occurred in succession based on the previous dose level (12, 8, and 4 mg/day, and 4 mg every other day [QOD]). Any dose reduction below 4 mg QOD was discussed with the sponsor. Once the dose was reduced, it was not increased at a later date.		
Nonhematologic Toxicities		
Treatment-Related Toxicity^{a,b}	Management	Dose Adjustment
Grade 1 or Tolerable Grade 2	Continue treatment ^c	No change
Intolerable Grade 2^c or Grade 3^{d,e}		
First occurrence	Interrupted until resolved to Grade 0-1 or baseline	One-level reduction
Second occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-1 or baseline	One-level reduction
Third occurrence ^f (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-1 or baseline	One-level reduction
Fourth occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-1 or baseline	Discuss with sponsor
Grade 4:^g Discontinue Lenvatinib		
Hematologic Toxicities and Proteinuria		
Treatment-Related Toxicity^a	Management	Dose Adjustment
Grade 1 or Grade 2^e	Continue treatment ^c	No change
Grade 3^e		
First occurrence	Interrupted until resolved to Grade 0-2 or Baseline	No change
Second occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	One-level reduction
Third occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	One-level reduction
Fourth occurrence ^f (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	One-level reduction
Fifth occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	Discuss with sponsor

[Table 12]

Table 6 (cont.): Lenvatinib Dose Reductions and Interruption Instructions

Grade 4		
First occurrence	Interrupted until resolved to Grade 0-2 or Baseline	One-level reduction
Second occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	One-level reduction
Third occurrence ^f (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	One-level reduction
Fourth occurrence (same toxicity or new toxicity)	Interrupted until resolved to Grade 0-2 or Baseline	Discuss with sponsor

Note: Grading according to CTCAE v4.0.

ALP = alkaline phosphatase; ALT = alanine aminotransferase; AST = aspartate aminotransferase; CTCAE v4.0 = Common Terminology Criteria for Adverse Events Version 4.0; ULN = upper limit of normal; γ -GTP = international normalized ratio.

a: An interruption of lenvatinib for more than 28 days (due to treatment-related toxicities) required sponsor's approval before treatment was resumed. During treatment interruption, AE assessment was repeated at least every 7 days (until administration was restarted).

b: Initiated optimal medical management for nausea, vomiting, diarrhea, and/or hypothyroidism prior to any lenvatinib treatment, interruption, or dose reduction.

c: Grade 2 toxicities were determined to be tolerable or intolerable by both the subject and investigator. If the Grade 2 toxicity was determined to be intolerable, the dose of study drug was reduced with or without dose interruption. Interruption for Grade 3 toxicity was mandatory.

d: Obese subjects with weight loss did not need to return to baseline or Grade 1 weight loss to restart lenvatinib. If there was no weight loss for at least 1 week, subjects were restarted at the lower dose, and normal body mass index was used for further dose reductions.

e: Not applicable to abnormal clinical laboratory values that were not clinically relevant based on the judgment of the investigator (e.g., ALT, AST, γ -GTP values $<10 \times$ ULN, and sodium).

f: Not applicable for subjects who started at 8 mg QD.

g: Excluding laboratory abnormalities judged to be non-life-threatening, which were managed as Grade 3.

[0193] Management of Hypertension

Hypertension is a recognized side effect of treatment with drugs inhibiting vascular endothelial growth factor (VEGF) signaling. Subjects enrolled in the trial had BP $\leq 150/90$ mm Hg at the time of study entry and, if they were known to be hypertensive, had been on a stable dose of antihypertensive therapy for at least 1 week before Cycle 1/Day 1. Early detection and effective management of hypertension were important to minimize the need for lenvatinib dose interruptions and reductions.

[0194] Antihypertensive agents were started as soon as elevated BP (systolic BP ≥ 140 mm Hg or diastolic BP ≥ 90 mm Hg) was confirmed on 2 assessments a minimum of 1-hour apart. One BP assessment was defined as the mean value of 3 measurements at least 5 minutes apart. The choice of antihypertensive treatment was individualized to the subject's clinical circumstances and followed standard medical practice. For previously normotensive subjects, appropriate antihypertensive therapy was started when systolic BP ≥ 140 mm Hg or diastolic BP ≥ 90 mm Hg was first observed on 2 assessments a minimum of 1-hour apart. For those subjects who were already on antihypertensive medication, treatment modification may have been necessary if hypertension persisted. For subjects with hypertension and proteinuria, appropriate therapy, e.g., angiotensin-converting enzyme inhibitor or angiotensin-II receptor antagonist was preferred.

[0195] Lenvatinib was withheld in any instance where a subject was at imminent risk to develop a hypertensive crisis or had significant risk factors for severe complications of uncontrolled hypertension (e.g., BP $\geq 160/100$ mm Hg, significant risk factors for cardiac disease, intracerebral hemorrhage, or other significant co-morbidities). Once the subject was on the same antihypertensive medications for at least 48 hours and the BP was controlled, lenvatinib was resumed as described below.

[0196] During the Treatment Period, both in the Randomization Phase and in the Extension Phase, subjects with systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg had their BP monitored every 2 weeks (on Day 15 or more frequently as clinical indicated) until systolic BP was ≤ 150 mm Hg and diastolic BP was ≤ 95 mm Hg for 3 consecutive months. If a repeat event of systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg occurred, the subject resumed the Day 15 evaluation until systolic BP was ≤ 150 mm Hg and diastolic BP was ≤ 95 mm Hg for 3 consecutive months.

[0197] The following guidelines were followed for the management of systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg confirmed on repeat measurements after 1 hour:

Lenvatinib was continued and antihypertensive therapy was instituted for subjects not already receiving antihypertensive medication

For those subjects already on antihypertensive medication, dose or medication choice was modified as per the investigator.

If systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg persisted despite maximal antihypertensive therapy, then lenvatinib administration was interrupted and restarted at a dose of 8 mg QD (one dose level reduction as specified in Table 6) only when systolic BP ≤ 150 mm Hg and diastolic BP ≤ 95 mm Hg and the subject was on a stable dose of antihypertensive medication for at least 48 hours.

If systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg recurred on the 8-mg QD dose despite optimal management of hypertension with antihypertensive med-

ications (either by dose increase or the addition of a different class of antihypertensive), then lenvatinib administration was interrupted and restarted at a dose of 4-mg QD (1 more dose level reduction as specified in Table 6) only when systolic BP ≤ 150 mm Hg and diastolic BP ≤ 95 mm Hg and the subject was on a stable dose of antihypertensive medication for at least 48 hours.

If systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg recurred on the 4-mg QD dose despite optimal management of hypertension with antihypertensive medications (either by dose increase or the addition of a different class of antihypertensive), then lenvatinib administration was interrupted and restarted at a dose of 4-mg QOD (1 more dose level reduction as specified in Table 6) only when systolic BP ≤ 150 mm Hg and diastolic BP ≤ 95 mm Hg and the subject was on a stable dose of antihypertensive medication for at least 48 hours.

If systolic BP ≥ 160 mm Hg or diastolic BP ≥ 100 mm Hg recurred on the 4-mg QOD dose despite optimal management of hypertension with antihypertensive medications (either by dose increase or the addition of a different class of antihypertensive), then lenvatinib administration was interrupted and restart of study medication was discussed with the sponsor.

[0198] The following guidelines were followed for the management of Grade 4 hypertension (life-threatening consequences):

Appropriate medical management was instituted.

Study drug was discontinued.

[0199] Management of Proteinuria

Regular assessment for proteinuria was conducted. Guidelines for assessment and management of proteinuria were as follows:

Initial episode of proteinuria: If proteinuria $\geq 2+$ was detected on urine dipstick testing, study drug was continued and a 24-hour urine collection for total protein was obtained as soon as possible within 72 hours to verify the grade of proteinuria. Grading according to CTCAE v4.0 was based on the 24-hour urine collection for total protein result. Additionally, a spot protein-creatinine ratio test was performed on the 24-hour urine sample as soon as possible within 72 hours. Management of lenvatinib administration was based on the grade of proteinuria according to 7.

During the Treatment Period, both in the Randomization Phase and the Extension Phase, urine dipstick testing for subjects with proteinuria $\geq 2+$ was performed every 2 weeks (on Day 15 or more frequently as clinically indicated) until the results were 1+ or negative for 3 consecutive months. Any subsequent increases in the level of proteinuria $\geq 2+$ on urine dipstick testing were confirmed with a 24-hour urine collection and graded according to the dose reduction and interruption instructions provided in 7. A spot protein-creatinine ratio test was performed on the 24 hour urine sample as soon

as possible. If a new event of proteinuria $\geq 2+$ occurred, the subject resumed the Day 15 urine dipstick testing for evaluation of proteinuria until results were 1+ or negative for 3 consecutive months.

[0200] A 24-hour urine collection for protein quantitation was required in the following situations:

The first (initial) occurrence of 2+, 3+, or 4+ proteinuria on urine dipstick while on study drug

A subsequent apparent increase in severity of urine dipstick proteinuria (from the prior measurement which was $\geq 2+$) occurring on the same lenvatinib dose level

When there was a lenvatinib dose reduction and on follow-up, the urine protein dipstick result is 2+, 3+, or 4+ (at the new dose level)

[0201] The 24-hour urine collection was not required in the following situations:

Persistence of the same severity of proteinuria by urine dipstick at the same study dose level when a 24-hour urine collection has already been collected at that dose level)

Subsequent occurrences of 2+, 3+, or 4+ proteinuria by urine dipstick when the subject was off study drug

[0202] Management of Hepatotoxicity

Regular monitoring of liver function tests (e.g., alanine transaminase [ALT], aspartate transaminase [AST], bilirubin levels) was conducted as clinically indicated. If signs occurred indicating a decrease in liver function by 1 grade or more from Baseline, the instructions in Table 6 were followed. Appropriate supportive care was provided together with close monitoring. If hepatic failure occurred the study drug was discontinued.

[0203] Management of Thromboembolic Events

Subjects were advised to pay attention to the symptoms suggestive of venous thromboembolic events, which included acute onset of dyspnea, chest pain, cough, hemoptysis, tachypnea, tachycardia, cyanosis, and deep vein thrombosis signs including lower extremity swelling, redness, and warmth to touch or tenderness. If any of these signs or symptoms appeared, subjects were instructed to report such signs and symptoms promptly to the treating physician. If a thromboembolic event was confirmed, instructions contained in Table 6 were followed. If a subject experienced life-threatening (Grade 4) thromboembolic reactions, including pulmonary embolism, the study drug was discontinued.

[0204] Management of Posterior Reversible Encephalopathy Syndrome

In clinical studies with lenvatinib, events of posterior reversible encephalopathy syndrome (PRES), a neurological disorder that can present with headache, seizure, lethargy, confusion, altered mental function, blindness, and other visual or neurological

disturbances, were reported in less than 1% of lenvatinib-treated subjects. Mild to severe hypertension also could be present. A magnetic resonance imaging (MRI) was necessary to confirm the diagnosis of PRES. In subjects with signs or symptoms of PRES, appropriate measures were to be taken to control blood pressure, and instructions in Table 6 were followed.

[0205] Example 3: Adverse Events that Required Dose Reduction or Interruption of Study
 Drug

A summary of TEAEs occurring in $\geq 10\%$ of subjects in the lenvatinib or sorafenib treatment arm, in decreasing order of frequency in the total lenvatinib arm, is presented in Table 7.

[0206]

[Table 13]

Table 7: Treatment-emergent Adverse Events Occurring in at Least 10% of Subjects in Either Treatment Arm, by Preferred Term

	Lenvatinib			Sorafenib (N = 475) n (%)
	8 mg ^a (N = 151) n (%)	12 mg ^a (N = 325) n (%)	Total (N = 476) n (%)	
Preferred Term				
<i>Subjects with Any TEAEs</i>	151 (100.0)	319 (98.2)	470 (98.7)	472 (99.4)
Hypertension	65 (43.0)	136 (41.8)	201 (42.2)	144 (30.3)
Diarrhea	53 (35.1)	131 (40.3)	184 (38.7)	220 (46.3)
Decreased appetite	50 (33.1)	112 (34.5)	162 (34.0)	127 (26.7)
Weight decreased	43 (28.5)	104 (32.0)	147 (30.9)	106 (22.3)
Fatigue	42 (27.8)	99 (30.5)	141 (29.6)	119 (25.1)
Palmar-plantar erythrodysesthesia syndrome	35 (23.2)	93 (28.6)	128 (26.9)	249 (52.4)
Proteinuria	37 (24.5)	80 (24.6)	117 (24.6)	54 (11.4)
Dysphonia	28 (18.5)	85 (26.2)	113 (23.7)	57 (12.0)
Nausea	24 (15.9)	69 (21.2)	93 (19.5)	68 (14.3)
Platelet count decreased	26 (17.2)	61 (18.8)	87 (18.3)	58 (12.2)
Abdominal pain	19 (12.6)	62 (19.1)	81 (17.0)	87 (18.3)
Hypothyroidism	25 (16.6)	53 (16.3)	78 (16.4)	8 (1.7)
Vomiting	22 (14.6)	55 (16.9)	77 (16.2)	36 (7.6)
Constipation	19 (12.6)	57 (17.5)	76 (16.0)	52 (10.9)
Blood bilirubin increased	23 (15.2)	48 (14.8)	71 (14.9)	63 (13.3)
Pyrexia	24 (15.9)	45 (13.8)	69 (14.5)	63 (13.3)
Ascites	21 (13.9)	47 (14.5)	68 (14.3)	44 (9.3)
Edema peripheral	23 (15.2)	43 (13.2)	66 (13.9)	33 (6.9)
Aspartate aminotransferase increased	21 (13.9)	44 (13.5)	65 (13.7)	80 (16.8)
Abdominal pain upper	21 (13.9)	37 (11.4)	58 (12.2)	40 (8.4)
Asthenia	14 (9.3)	40 (12.3)	54 (11.3)	48 (10.1)
Alanine aminotransferase increased	17 (11.3)	36 (11.1)	53 (11.1)	52 (10.9)
Back pain	11 (7.3)	39 (12.0)	50 (10.5)	31 (6.5)
Rash	18 (11.9)	28 (8.6)	46 (9.7)	76 (16.0)
Stomatitis	11 (7.3)	34 (10.5)	45 (9.5)	56 (11.8)
Alopecia	5 (3.3)	9 (2.8)	14 (2.9)	119 (25.1)

[Table 14]

[Data cutoff date: Nov 13, 2016; Percentages are based on the total number of subjects within the relevant treatment group in the Safety Analysis Set. Display is in decreasing order of frequency of TEAEs in the lenvatinib total group. Subjects with 2 or more TEAEs in the same preferred term were counted only once. Adverse Event terms were coded using MedDRA version 19.1. MedDRA = Medical Dictionary for Regulatory Activities; TEAE = treatment-emergent adverse event. ^a: 8 mg and 12 mg were the lenvatinib starting doses based on the subjects' body weight (<60 kg, ≥60 kg) at Baseline.]

[0207] A TEAE occurred in all but 6 and 3 subjects in the lenvatinib and sorafenib arms, respectively. The most frequently reported TEAEs (>30% of subjects) were hypertension, diarrhea, decreased appetite, and weight decreased for lenvatinib and palmar-plantar erythrodysesthesia (PPE) syndrome, diarrhea, and hypertension for sorafenib. Adverse events that occurred in $\geq 10\%$ more subjects in the lenvatinib arm than in the sorafenib arm were: hypertension (42.2% vs 30.3%), proteinuria (24.6% vs 11.4%), dysphonia (23.7% vs 12.0%), and hypothyroidism (16.4% vs 1.7%). These AEs are consistent with the known safety profile of lenvatinib in other cancer indications. Other AEs with a subject incidence of $< 10\%$ but that were reported in a higher proportion of lenvatinib-treated subjects compared with sorafenib (≥ 10 episodes in 1 treatment arm and a difference between treatments of ≥ 0.1 episodes per SY) when adjusted by treatment duration included: proteinuria (0.5 vs 0.31 episodes per SY), dysphonia (0.4 vs 0.28), hypothyroidism (0.24 vs 0.03), WBC count decreased (0.25 vs 0.15), neutrophil count decreased (0.21 vs 0.06), and hepatic encephalopathy (0.17 vs 0.04).

[0208] An algorithm of dose interruption followed by dose reduction was used for the management of lenvatinib toxicity (see, Table 6). Unadjusted for treatment duration, adverse events (AEs) leading to dose reduction or interruption were reported in 61.8% of subjects in the lenvatinib arm and 55.6% of subjects in the sorafenib arm (Table 8).

[0209]

[Table 15]

Table 8: Treatment-emergent Adverse Events Leading to Dose Reduction/Interruption of Study Drug in at Least 1% of Subjects in Either Treatment Arm, by System Organ Class and Preferred Term – Safety Analysis Set

System Organ Class Preferred Term	Lenvatinib			Sorafenib
	8 mg ^a (N = 151) n (%)	12 mg ^a (N = 325) n (%)	Total (N = 476) n (%)	(N = 475) n (%)
Subjects with any TEAEs leading to Dose Reduction or Interruption	81 (53.6)	213 (65.5)	294 (61.8)	264 (55.6)
Blood and lymphatic system disorders	5 (3.3)	11 (3.4)	16 (3.4)	13 (2.7)
Neutropenia	0 (0.0)	6 (1.8)	6 (1.3)	0 (0.0)
Thrombocytopenia	2 (1.3)	4 (1.2)	6 (1.3)	4 (0.8)
Anemia	2 (1.3)	1 (0.3)	3 (0.6)	8 (1.7)
Gastrointestinal disorders	22 (14.6)	65 (20.0)	87 (18.3)	72 (15.2)
Diarrhea	8 (5.3)	28 (8.6)	36 (7.6)	35 (7.4)
Nausea	2 (1.3)	14 (4.3)	16 (3.4)	9 (1.9)
Vomiting	1 (0.7)	11 (3.4)	12 (2.5)	6 (1.3)
Ascites	3 (2.0)	7 (2.2)	10 (2.1)	5 (1.1)
Abdominal pain	2 (1.3)	4 (1.2)	6 (1.3)	8 (1.7)
Stomatitis	2 (1.3)	4 (1.2)	6 (1.3)	2 (0.4)
Abdominal pain upper	2 (1.3)	3 (0.9)	5 (1.1)	2 (0.4)
Esophageal varices hemorrhage	0 (0.0)	5 (1.5)	5 (1.1)	1 (0.2)
General disorders and administration site conditions	17 (11.3)	50 (15.4)	67 (14.1)	35 (7.4)
Fatigue	5 (3.3)	22 (6.8)	27 (5.7)	17 (3.6)
Asthenia	3 (2.0)	13 (4.0)	16 (3.4)	9 (1.9)
Pyrexia	4 (2.6)	8 (2.5)	12 (2.5)	5 (1.1)
Edema peripheral	2 (1.3)	4 (1.2)	6 (1.3)	3 (0.6)
Malaise	2 (1.3)	3 (0.9)	5 (1.1)	0 (0.0)
Hepatobiliary disorders	2 (1.3)	25 (7.7)	27 (5.7)	19 (4.0)
Hepatic function abnormal	0 (0.0)	2 (0.6)	2 (0.4)	9 (1.9)

[Table 16]

Table 8 (cont.): Treatment-emergent Adverse Events Leading to Dose Reduction/Interruption of Study Drug in at Least 1% of Subjects in Either Treatment Arm, by System Organ Class and Preferred Term – Safety Analysis Set

System Organ Class Preferred Term	Lenvatinib			Sorafenib (N = 475) n (%)
	8 mg ^a (N = 151) n (%)	12 mg ^a (N = 325) n (%)	Total (N = 476) n (%)	
Investigations	24 (15.9)	57 (17.5)	81 (17.0)	52 (10.9)
Platelet count decreased	9 (6.0)	13 (4.0)	22 (4.6)	12 (2.5)
Blood bilirubin increased	8 (5.3)	12 (3.7)	20 (4.2)	14 (2.9)
Weight decreased	3 (2.0)	17 (5.2)	20 (4.2)	4 (0.8)
Neutrophil count decreased	4 (2.6)	7 (2.2)	11 (2.3)	5 (1.1)
Aspartate aminotransferase increased	4 (2.6)	5 (1.5)	9 (1.9)	18 (3.8)
Gamma-glutamyltransferase increased	1 (0.7)	6 (1.8)	7 (1.5)	4 (0.8)
Alanine aminotransferase increased	2 (1.3)	4 (1.2)	6 (1.3)	9 (1.9)
White blood cell count decreased	3 (2.0)	3 (0.9)	6 (1.3)	4 (0.8)
Metabolism and nutrition disorders	21 (13.9)	31 (9.5)	52 (10.9)	19 (4.0)
Decreased appetite	15 (9.9)	21 (6.5)	36 (7.6)	15 (3.2)
Dehydration	3 (2.0)	3 (0.9)	6 (1.3)	0 (0.0)
Hyponatremia	2 (1.3)	3 (0.9)	5 (1.1)	0 (0.0)
Musculoskeletal and connective tissue disorders	4 (2.6)	14 (4.3)	18 (3.8)	9 (1.9)
Myalgia	1 (0.7)	4 (1.2)	5 (1.1)	0 (0.0)
Nervous system disorders	7 (4.6)	28 (8.6)	35 (7.4)	14 (2.9)
Hepatic encephalopathy	4 (2.6)	16 (4.9)	20 (4.2)	3 (0.6)
Renal and urinary disorders	5 (3.3)	35 (10.8)	40 (8.4)	9 (1.9)
Proteinuria	4 (2.6)	29 (8.9)	33 (6.9)	7 (1.5)
Respiratory, thoracic and mediastinal disorders	4 (2.6)	12 (3.7)	16 (3.4)	12 (2.5)
Dyspnea	2 (1.3)	6 (1.8)	8 (1.7)	4 (0.8)
Skin and subcutaneous tissue disorders	5 (3.3)	23 (7.1)	28 (5.9)	111 (23.4)
Palmar-plantar erythrodysesthesia syndrome	4 (2.6)	21 (6.5)	25 (5.3)	88 (18.5)
Rash	0 (0.0)	2 (0.6)	2 (0.4)	10 (2.1)
Rash maculo-papular	0 (0.0)	0 (0.0)	0 (0.0)	5 (1.1)

[Table 17]

Table 8 (cont.): Treatment-emergent Adverse Events Leading to Dose Reduction/Interruption of Study Drug in at Least 1% of Subjects in Either Treatment Arm, by System Organ Class and Preferred Term – Safety Analysis Set

	Lenvatinib			Sorafenib (N = 475)
	8 mg ^a (N = 151)	12 mg ^a (N = 325)	Total (N = 476)	
System Organ Class				
Preferred Term	n (%)	n (%)	n (%)	n (%)
Vascular disorders	3 (2.0)	30 (9.2)	33 (6.9)	19 (4.0)
Hypertension	3 (2.0)	26 (8.0)	29 (6.1)	18 (3.8)

[The data cutoff date for this table was Nov 13, 2016. Percentages are based on the total number of subjects within the relevant treatment group in the Safety Analysis Set. Subjects with 2 or more TEAEs reported in the same system organ class or preferred term were only counted once. Adverse Events terms were coded using MedDRA version 19.1. MedDRA = Medical Dictionary for Regulatory Activities; TEAE = treatment-emergent adverse event. ^a: 8 mg and 12 mg were the lenvatinib starting doses based on the subjects' body weight (<60 kg, ≥60 kg) at baseline.]

- [0210] Adverse events leading to dose reduction or interruption of lenvatinib were most frequently (≥10% of subjects) coded to the SOC of Gastrointestinal disorders (18.3%), Investigations (17.0%), General disorders and Administration Site Conditions (14.1%), and Metabolism and Nutrition disorders (10.9%).
- [0211] Adverse events that led to dose reduction or interruption in 5% or more of subjects in the lenvatinib or sorafenib arm, respectively, were decreased appetite (7.6% vs 3.2%), diarrhea (7.6% vs 7.4%), proteinuria (6.9% vs 1.5%), hypertension (6.1% vs 3.8%), fatigue (5.7% vs 3.6%), and PPE syndrome (5.3% vs 18.5%). Except for PPE syndrome, which occurred more frequently with sorafenib, these treatment-emergent adverse events (TEAEs) led to dose modification either at similar rates in the 2 treatment arms, or at a higher frequency in the lenvatinib arm.
- [0212] Other TEAEs that led to dose modification in a higher percentage of subjects in the lenvatinib arm (total incidence >2% to <5% but [approximately] >2 times the rate) than in the sorafenib arm were hepatic encephalopathy (4.2% vs 0.6%), weight decreased (4.2% vs 0.8%), pyrexia (2.5% vs 1.1%), vomiting (2.5% vs 1.3%), neutrophil count decreased (2.3% vs 1.1%), and ascites (2.1% vs 1.1%). The only TEAE that led to dose modification in a higher percentage of subjects in the sorafenib arm (total incidence >2% to <5% but at least double the rate) than in the lenvatinib arm was AST increased (3.8% vs 1.9%).
- [0213] As of the data cutoff date for the primary analysis, the majority of subjects had ended treatment; treatment was ongoing for 27 (5.6%) lenvatinib and 25 (5.3%) sorafenib subjects. The most frequent reason for discontinuation of treatment was disease pro-

gression. Fewer subjects ended treatment due to disease progression in the lenvatinib arm (311; 65.1%) than in the sorafenib arm (347; 72.9%). Adverse events were cited by the investigators as the reason for ending treatment in 13.2% (n=63) of lenvatinib and 9.0% (n=43) of sorafenib subjects (see, Table 9).

[0214] [Table 18]

Table 9: Subject Disposition and Reasons for Discontinuation of Treatment during Randomization Phase

	Lenvatinib (N=478) n (%)	Sorafenib (N=476) n (%)	Total (N=954) n (%)
Randomized	478 (100.0)	476 (100.0)	954 (100.0)
Not treated	2 (0.4)	1 (0.2)	3 (0.3)
Treated	476 (99.6)	475 (99.8)	951 (99.7)
Treatment ongoing at data cutoff date	27 (5.6)	25 (5.3)	52 (5.5)
Discontinued Treatment^b	451 (94.4)	451 (94.7)	902 (94.5)
Primary Reason for Discontinuation of Treatment			
Disease Progression	311 (65.1)	347 (72.9)	658 (69.0)
Adverse event	63 (13.2)	43 (9.0)	106 (11.1)
Subject choice	28 (5.9)	15 (3.2)	43 (4.5)
Lost to follow-up	3 (0.6)	1 (0.2)	4 (0.4)
Withdrawal of consent	9 (1.9)	5 (1.1)	14 (1.5)
Other ^a	37 (7.7)	40 (8.4)	77 (8.1)
Discontinued Treatment but in Survival Follow-up	82 (17.2)	82 (17.2)	164 (17.2)

Data cutoff date: Nov 13, 2016.

Percentages are based on the total number of subjects within the relevant treatment group in the Full Analysis Set.

a: These reasons were collected as investigator comments under the "Other" category in the case report form.

b: Includes the 3 subjects who were not treated after randomization.

[0215] The percentage of the subjects who discontinued lenvatinib treatment due to adverse events was 13.2% (n=62). This low percentage of discontinuation could not have been achieved without the use of dose modifications (see Table 6).

[0216] Example 4: Efficacy Results in Hepatocellular Carcinoma Clinical Study

The efficacy of LENVIMA was evaluated in a randomized, open-label, multicenter, international study (REFLECT; NCT0761266) conducted in patients with previously untreated unresectable hepatocellular carcinoma (HCC). The study enrolled adults with Child-Pugh A and Barcelona Clinic Liver Cancer (BCLC) Stage C or B HCC who were ineligible for local liver-directed therapy; had an ECOG PS of 0 or 1; had

received no prior systemic therapy for HCC; and had at least one measurable target lesion according to modified RECIST for HCC. Efficacy results are summarized in Table 10.

[0217] [Table 19]

Table 10: Efficacy Results in Hepatocellular Carcinoma in REFLECT		
	LENVIMA N= 478	Sorafenib N=476
Overall Survival		
Number of deaths (%)	351 (73)	350 (74)
Median OS in months (95% CI)	13.6 (12.1, 14.9)	12.3 (10.4, 13.9)
Hazard Ratio (95% CI) ^a	0.92 (0.79,1.06)	
Progression-Free Survival ^b (mRECIST)		
Number of Events (%)	311 (65)	323 (68)
Median PFS in months (95% CI)	7.3 (5.6, 7.5)	3.6 (3.6, 3.7)
Hazard Ratio (95% CI) ^a	0.64 (0.55, 0.75)	
P-value	<0.001	
Objective Response Rate ^b (mRECIST)		
Objective response rate	41%	12%
Complete responses, n (%)	10 (2.1)	4 (0.8)
Partial responses, n (%)	184 (38.5)	55 (11.6)
95% CI	(36%, 45%)	(9%, 15%)
Odds Ratio (95% CI)	5.01 (3.59, 7.01)	
P-value	<0.001	
Progression-Free Survival ^b (RECIST 1.1)		
Number of Events (%)	307 (64)	320 (67)
Median PFS in months (95% CI)	7.3 (5.6, 7.5)	3.6 (3.6, 3.9)
Hazard Ratio (95% CI) ^a	0.65 (0.56, 0.77)	
Objective Response Rate ^b (RECIST 1.1)		
Objective response rate	19%	7%
Complete responses, n (%)	2 (0.4)	1 (0.2)
Partial responses, n (%)	88 (18.4)	30 (6.3)
95% CI	(15%, 22%)	(4%, 9%)
Odds Ratio (95% CI)	3.3 (2.2, 5.1)	

CI = confidence interval; ECOG PS = Eastern Cooperative Oncology Group Performance Status; HR = hazard ratio; OS = overall survival.

Non-inferiority margin for hazard ratio (lenvatinib vs sorafenib) is 1.08.

Per independent review.

[0218] Example 5: Drug Interaction Studies: Effect of Lenvatinib on Other Drugs

Clinical Studies with Substrates of CYP3A4 or CYP2C8: There is no projected significant drug-drug interaction risk between lenvatinib and midazolam (a CYP3A4 substrate) or repaglinide (a CYP2C8 substrate).

- [0219] In Vitro Studies with Substrates of CYP or UDP-glucuronosyltransferase (UGT): Lenvatinib inhibits CYP2C8, CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2D6, and CYP3A. Lenvatinib does not inhibit CYP2A6 and CYP2E1. Lenvatinib induces CYP3A, but it does not induce CYP1A1, CYP1A2, CYP2B6, and CYP2C9.
- [0220] Lenvatinib inhibits UGT1A1, UGT1A4, and UGT1A9 in vitro, but likely only inhibits UGT1A1 in vivo in the gastrointestinal tract based on the expression of the enzyme in tissues. Lenvatinib does not inhibit UGT1A6, UGT2B7 or aldehyde oxidase. Lenvatinib does not induce UGT1A1, UGT1A4, UGT1A6, UGT1A9, or UGT2B7.
- [0221] In Vitro Studies with Substrates of Transporters: Lenvatinib does not have the potential to inhibit MATE1, MATE2-K, OCT1, OCT2, OAT1, OAT3, BSEP, OATP1B1, or OATP1B3 in vivo.
- [0222] CROSS-REFERENCE TO RELATED APPLICATIONS
- This application claims the benefit of priority of U.S. Provisional Appl. No. 62/506,900, filed May 16, 2017, the contents of which are incorporated by reference herein in their entirety.
- [0223] OTHER EMBODIMENTS
- While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

Claims

- [Claim 1] A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a dosage regimen of lenvatinib or a pharmaceutically acceptable salt thereof that is: (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg.
- [Claim 2] A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a dosage regimen of lenvatinib or a pharmaceutically acceptable salt thereof that is 8 mg/day.
- [Claim 3] A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises:
- (a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity until the first Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 3 nonhematologic toxicity during treatment with the second dosage regimen;
 - (b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 nonhematologic toxicity until the second Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage

regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 3 non-hematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 nonhematologic toxicity until the third Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[Claim 4]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity until the first Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second Grade 3 nonhematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 nonhematologic toxicity until the second Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third Grade 3 nonhematologic toxicity

during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 nonhematologic toxicity.

[Claim 5]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the oc-

currence of the third persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity until the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[Claim 6]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 non-

hematologic toxicity.

[Claim 7]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic

toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[Claim 8]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3

nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality.

[Claim 9]

The method of any one of claims 2 to 8, wherein the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding non-life-threatening Grade 4 laboratory abnormality during treatment with the first, second, third or fourth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding the non-life-threatening Grade 4 laboratory abnormality.

[Claim 10]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria until the first Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 3 hematologic

toxicity or proteinuria during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria until the second Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria during treatment with the third dosage regimen;

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria until the third Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria during treatment with the fourth dosage regimen; and

(d) terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria until the fourth Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fifth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[Claim 11]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria during treatment with the first

dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria until the first Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria until the second Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria during treatment with the third dosage regimen;

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria until the third Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria during treatment with the fourth dosage regimen; and

(d) terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria.

[Claim 12]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprises lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

- (a) terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen;
- (b) terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen; and
- (c) terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity until the third Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[Claim 13]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprises lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

- (a) terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4

hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity.

[Claim 14]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day;

(II) wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria until the first Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body

weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria until the second Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria during treatment with the third dosage regimen;

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria until the third Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria during treatment with the fourth dosage regimen; and

(d) terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria until the fourth Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fifth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day; or

(III) wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and admin-

istering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity until the third Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject an fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day.

[Claim 15]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory ab-

normality until the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality;

(II) wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria until the first Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria

until the second Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria during treatment with the third dosage regimen;

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria until the third Grade 3 hematologic toxicity or proteinuria is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria during treatment with the fourth dosage regimen; and

(d) terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria; or (III) wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity.

- [Claim 16] The method of any one of claims 10 to 15, wherein the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding non-life-threatening Grade 4 laboratory abnormality during treatment with the first, second, third, fourth or fifth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding the non-life-threatening Grade 4 laboratory abnormality.
- [Claim 17] A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,
- (I) wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:
- (a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the second dosage regimen;
- (b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the second Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt

thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the third dosage regimen;

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the third Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the fourth dosage regimen; and

(d) terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the fourth Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fifth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day;

(II) wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is

less than 60 kg, wherein the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity until the third Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day;

(III) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 nonhematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 non-

hematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 non-hematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 nonhematologic toxicity is resolved to Grade 0-1 or baseline, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day;

(IV) wherein the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the first Grade 3 non-hematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the second Grade 3

nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the third Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day; or

(V) wherein the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality during treatment with the first, second, third, fourth or fifth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality.

[Claim 18]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

(I) wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria

excluding non-clinically relevant laboratory abnormality until the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a second Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the second Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a third Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the third dosage regimen;

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the third Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a fourth Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the fourth dosage regimen; and

(d) terminating administration of the fourth dosage regimen after the occurrence of the fourth Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality;

(II) wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the oc-

currence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second Grade 4 hematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 4 hematologic toxicity until the second Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third Grade 4 hematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 4 hematologic toxicity;

(III) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 nonhematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 nonhematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other

day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 nonhematologic toxicity, and with or without interruption of the first dosage regimen;

(IV) wherein the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the second Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality; or

(V) wherein the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, and the method further comprises terminating admin-

istration of the dosage regimen after the occurrence of the Grade 4 non-hematologic toxicity excluding nonlife-threatening laboratory abnormality.

[Claim 19]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

(I) wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality until the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg;

(II) wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg;

(III) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 non-

hematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 non-hematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg; (IV) wherein the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg; or (V) wherein the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality.

[Claim 20]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, (I) wherein the human subject develops an occurrence of a first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory ab-

normality until the first Grade 3 hematologic toxicity or proteinuria excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day;

(II) wherein the human subject develops an occurrence of a first Grade 4 hematologic toxicity during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 4 hematologic toxicity until the first Grade 4 hematologic toxicity is resolved to Grade 0-2 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day;

(III) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 nonhematologic toxicity during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 non-hematologic toxicity, and with or without interruption of the first dosage regimen until the first persistent and intolerable Grade 2 non-hematologic toxicity is resolved to Grade 0-1 or baseline, administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day;

(IV) wherein the human subject develops an occurrence of a first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality until the first Grade 3 nonhematologic toxicity excluding non-clinically relevant laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day; or

(V) wherein the human subject develops an occurrence of a Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage

regimen after the occurrence of the Grade 4 nonhematologic toxicity excluding nonlife-threatening laboratory abnormality.

[Claim 21] The method of any one of claims 3 to 9 and 14 to 18, wherein medical management of each of the first, second, and third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicities or non-life-threatening Grade 4 laboratory abnormality is initiated prior to terminating administration of the dosage regimen administered at the time of onset of the Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality.

[Claim 22] The method of any one of claims 10, 11 and 14 to 18, wherein medical management of each of the first, second, third, and fourth Grade 3 hematologic toxicities or proteinuria is initiated prior to terminating administration of the dosage regimen administered at the time of onset of the Grade 3 hematologic toxicities or proteinuria.

[Claim 23] The method of any one of claims 12 to 18, wherein medical management of each of the first, second, and third persistent and intolerable Grade 4 hematologic toxicities is initiated prior to terminating administration of the dosage regimen administered at the time of onset of the Grade 4 hematologic toxicities.

[Claim 24] The method of any one of claims 3 to 9 and 14 to 18, wherein the first persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality is the same as the second and/or third persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality.

[Claim 25] The method of any one of claims 10, 11 and 14 to 18, wherein the first Grade 3 hematologic toxicity or proteinuria is the same as the second and/or third Grade 3 hematologic toxicity or proteinuria.

[Claim 26] The method of any one of claims 12 to 18, wherein the first Grade 4 hematologic toxicity is the same as the second and/or third Grade 4 hematologic toxicity.

[Claim 27] The method of any one of claims 3 to 9 and 14 to 20, wherein the Grade 3 nonhematologic toxicity is selected from the group consisting of Grade 3 hypertension, Grade 3 diarrhea, Grade 3 arthralgia, Grade 3 myalgia, Grade 3 decreased appetite, Grade 3 fatigue, Grade 3 decreased weight, Grade 3 dysphonia, Grade 3 nausea, Grade 3 abdominal pain, Grade 3 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 3 vomiting, Grade 3 constipation, Grade 3 rash,

and Grade 3 palmar-plantar erythrodysesthesia.

[Claim 28]

The method of any one of claims 5 to 9 and 14 to 20, wherein the Grade 2 or Grade 3 nonhematologic toxicity is selected from the group consisting of Grade 3 hypertension, Grade 2 hypertension, Grade 3 diarrhea, Grade 2 diarrhea, Grade 3 decreased appetite, Grade 2 decreased appetite, Grade 3 arthralgia, Grade 2 arthralgia, Grade 3 myalgia, Grade 2 myalgia, Grade 3 fatigue, Grade 2 fatigue, Grade 3 decreased weight, Grade 2 decreased weight, Grade 2 alopecia, Grade 3 dysphonia, Grade 2 dysphonia, Grade 3 nausea, Grade 2 nausea, Grade 3 abdominal pain, Grade 2 abdominal pain, Grade 3 QT/QTc interval prolongation, Grade 2 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 2 hypothyroidism, Grade 3 vomiting, Grade 2 vomiting, Grade 3 constipation, Grade 2 constipation, Grade 3 rash, Grade 2 rash, Grade 3 palmar-plantar erythrodysesthesia, and Grade 2 palmar-plantar erythrodysesthesia.

[Claim 29]

The method of any one of claims 7 to 9 and 14 to 20, wherein the Grade 4 laboratory abnormality is selected from the group consisting of Grade 4 increase in aspartate aminotransferase, Grade 4 increase in alanine aminotransferase, Grade 4 increase in alkaline phosphatase, Grade 4 hypokalemia, Grade 4 hyponatremia, Grade 4 hypoglycemia, Grade 4 increase in blood bilirubin, and Grade 4 increase in gamma glutamyl transferase.

[Claim 30]

The method of any one of claims 10, 11, and 14 to 20, wherein the Grade 3 hematologic toxicity or proteinuria is selected from the group consisting of Grade 3 proteinuria, Grade 3 thrombopenia (thrombocytopenia), Grade 3 anemia, Grade 3 decrease in white blood cell count, Grade 3 neutropenia, and Grade 3 lymphocytopenia.

[Claim 31]

The method of any one of claims 12 to 20, wherein the Grade 4 hematologic toxicity is selected from the group consisting of Grade 4 thrombopenia (thrombocytopenia), Grade 4 anemia, Grade 4 decrease in white blood cell count, Grade 4 neutropenia, and Grade 4 lymphocytopenia.

[Claim 32]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma:
a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the

body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen;

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen; or

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen.

[Claim 33]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification: a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen; or

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen.

[Claim 34]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma:

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity during treatment with the prior dosage regimen; a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity during treatment with the prior dosage regimen; or a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject

developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen.

[Claim 35]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification: a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen; or a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity during treatment with the prior dosage regimen.

[Claim 36]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma: a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen; a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the

human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen; or a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen.

[Claim 37]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification: a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 nonhematologic toxicity or non-life-threatening Grade 4 laboratory abnormality during treatment with the prior dosage regimen; or a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a persistent and intolerable Grade 2 or Grade 3 non-hematologic toxicity or non-life-threatening Grade 4 laboratory ab-

[Claim 38]

normality during treatment with the prior dosage regimen.

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma:

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen;

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen;

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen; or

a dosage regimen comprising lenvatinib or a pharmaceutically ac-

ceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen.

[Claim 39]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification: a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen;

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen; or

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 3 hematologic toxicity or proteinuria during treatment with the prior dosage regimen.

[Claim 40]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma:

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the

body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen;

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen; or

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the body weight of the human subject is equal to or more than 60 kg, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen.

[Claim 41]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification: a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen; or

a dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject was previously treated with a prior dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, and wherein the human subject developed an occurrence of a Grade 4 hematologic toxicity during treatment with the prior dosage regimen.

[Claim 42] The method of claim 32 or 33, wherein the Grade 3 nonhematologic toxicity is selected from the group consisting of Grade 3 hypertension, Grade 3 diarrhea, Grade 3 arthralgia, Grade 3 myalgia, Grade 3 decreased appetite, Grade 3 fatigue, Grade 3 decreased weight, Grade 3 dysphonia, Grade 3 nausea, Grade 3 abdominal pain, Grade 3 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 3 vomiting, Grade 3 constipation, Grade 3 rash, and Grade 3 palmar-plantar erythrodysesthesia.

[Claim 43] The method of any one of claims 34 to 37, wherein the Grade 2 or Grade 3 nonhematologic toxicity is selected from the group consisting of Grade 3 hypertension, Grade 2 hypertension, Grade 3 diarrhea, Grade 2 diarrhea, Grade 3 decreased appetite, Grade 2 decreased appetite, Grade 3 arthralgia, Grade 2 arthralgia, Grade 3 myalgia, Grade 2 myalgia, Grade 3 fatigue, Grade 2 fatigue, Grade 3 decreased weight, Grade 2 decreased weight, Grade 2 alopecia, Grade 3 dysphonia, Grade 2 dysphonia, Grade 3 nausea, Grade 2 nausea, Grade 3 abdominal pain, Grade 2 abdominal pain, Grade 3 QT/QTc interval prolongation, Grade 2 QT/QTc interval prolongation, Grade 3 hypothyroidism, Grade 2 hypothyroidism, Grade 3 vomiting, Grade 2 vomiting, Grade 3 constipation, Grade 2 constipation, Grade 3 rash, Grade 2 rash, Grade 3 palmar-plantar erythrodysesthesia, and Grade 2 palmar-plantar erythrodysesthesia.

[Claim 44] The method of any one of claims 36 or 37, wherein the Grade 4 laboratory abnormality is selected from the group consisting of Grade 4 increase in aspartate aminotransferase, Grade 4 increase in alanine aminotransferase, Grade 4 increase in alkaline phosphatase, Grade 4 hypokalemia, Grade 4 hyponatremia, Grade 4 hypoglycemia, Grade 4 increase in blood bilirubin, and Grade 4 increase in gamma glutamyl transferase.

[Claim 45] The method of claim 38 or 39, wherein the Grade 3 hematologic toxicity or proteinuria is selected from the group consisting of Grade 3

proteinuria, Grade 3 thrombopenia (thrombocytopenia), Grade 3 anemia, Grade 3 decrease in white blood cell count, Grade 3 neutropenia, and Grade 3 lymphocytopenia.

[Claim 46]

The method of claim 40 or 41, wherein the Grade 4 hematologic toxicity is selected from the group consisting of Grade 4 thrombopenia (thrombocytopenia), Grade 4 anemia, Grade 4 decrease in white blood cell count, Grade 4 neutropenia, and Grade 4 lymphocytopenia.

[Claim 47]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i)

4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day; or

(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 48]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality; or

(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third or fourth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity,

2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 49]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg; or
(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or im-

pairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 50]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day; or

(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that Grade 3 hypertension, Grade 4 hypertension, Grade 3 cardiac dysfunction, Grade 4 cardiac dysfunction, any grade arterial thromboembolic event, Grade 3 hepatotoxicity, Grade 4 hepatotoxicity, 2 g or greater proteinuria in 24 hours, Grade 3 renal failure or impairment, Grade 4 renal failure or impairment, any Grade gastrointestinal perforation, Grade 3 fistula, Grade 4 fistula, a greater than 500 ms QT/QTc interval prolongation, a greater than 60 ms increase from baseline QT/QTc interval prolongation, and any Grade reversible

posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3 or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 51]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 4 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg every other day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3

adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and, if the body weight of the human subject is equal to or more than 60 kg, administering to the human subject a fourth dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day; or

(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third, or fourth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 52]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises:

(a) terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and ad-

ministering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day, wherein the human subject develops an occurrence of a second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the second dosage regimen;

(b) terminating administration of the second dosage regimen after the occurrence of the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the second persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a third dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg every other day, wherein the human subject develops an occurrence of a third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the third dosage regimen; and

(c) terminating administration of the third dosage regimen after the occurrence of the third persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality; or

(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first, second, third or fourth dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 53]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is

less than 60 kg,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg; or

(II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality;

provided that hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 54]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day,

(I) wherein the human subject develops an occurrence of a first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4

laboratory abnormality until the first persistent and intolerable Grade 2 or Grade 3 adverse reaction or Grade 4 laboratory abnormality is resolved to Grade 0-1 or baseline, and administering to the human subject a second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day; or (II) wherein the human subject develops an occurrence of a Grade 4 adverse reaction excluding Grade 4 laboratory abnormality during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the Grade 4 adverse reaction excluding Grade 4 laboratory abnormality; provided that hypertension, cardiac dysfunction, arterial thromboembolic event, hepatotoxicity, proteinuria, renal failure or impairment, gastrointestinal perforation, fistula, QT/QTc interval prolongation, and reversible posterior leukoencephalopathy syndrome are excluded from the persistent and intolerable Grade 2, Grade 3, or Grade 4 adverse reaction or Grade 4 laboratory abnormality.

[Claim 55]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg, wherein the human subject develops an occurrence of a Grade 3 hypertension during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the Grade 3 hypertension until the Grade 3 hypertension is controlled at less than or equal to Grade 2, and administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg.

[Claim 56]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first

dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a Grade 3 hypertension during treatment with the first dosage regimen, and the method further comprises terminating administration of the first dosage regimen after the occurrence of the Grade 3 hypertension until the Grade 3 hypertension is controlled at less than or equal to Grade 2, and administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day.

[Claim 57]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a 2 g or greater proteinuria in 24 hours during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the 2 g or greater proteinuria in 24 hours until the proteinuria is less than or equal to 2 g of proteinuria in 24 hours and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg,

provided that the human subject develops an occurrence of a nephrotic syndrome during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the nephrotic syndrome.

[Claim 58]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a 2 g or greater

proteinuria in 24 hours during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the 2 g or greater proteinuria in 24 hours until the proteinuria is less than or equal to 2 g of proteinuria in 24 hours and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day,

provided that the human subject develops an occurrence of a nephrotic syndrome during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the nephrotic syndrome.

[Claim 59]

A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 12 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 8 mg/day if the body weight of the human subject is less than 60 kg,

wherein the human subject develops an occurrence of a greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation until the QT/QTc interval prolongation improves to less than or equal to 480 ms or baseline and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of (i) 8 mg/day if the body weight of the human subject is equal to or more than 60 kg or (ii) 4 mg/day if the body weight of the human subject is less than 60 kg.

[Claim 60]


A method of treating unresectable hepatocellular carcinoma, the method comprising administering to a human subject that has an unresectable hepatocellular carcinoma and moderate hepatic impairment classified in Child-Pugh class B under Child-Pugh Classification a first dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 8 mg/day, wherein the human subject develops an occurrence of a greater than

500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation during treatment with the first dosage regimen, and the method further comprises terminating administration of the dosage regimen after the occurrence of the greater than 500 ms QT/QTc interval prolongation or a greater than 60 ms increase from baseline QT/QTc interval prolongation until the QT/QTc interval prolongation improves to less than or equal to 480 ms or baseline and, administering to the human subject the second dosage regimen comprising lenvatinib or a pharmaceutically acceptable salt thereof at a dose of 4 mg/day.


- [Claim 61] The method of any one of claims 1 to 60, wherein median overall survival is 13.6 months.
- [Claim 62] The method of any one of claims 1 to 60, wherein median overall survival with 95% confidence interval is between 12.1 and 14.9 months.
- [Claim 63] The method of any one of claims 1 to 60, wherein hazard ratio of overall survival compared with sorafenib at a dosage of 400 mg twice daily is 0.92.
- [Claim 64] The method of any one of claims 1 to 60, wherein hazard ratio of overall survival compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 0.79 and 1.06.
- [Claim 65] The method of any one of claims 1 to 60, wherein overall survival is 12 to 15 months.
- [Claim 66] The method of any one of claims 1 to 60, wherein median progression-free survival is 7.3 months.
- [Claim 67] The method of any one of claims 1 to 60, wherein median progression-free survival with 95% confidence interval is between 5.6 to 7.5 months.
- [Claim 68] The method of any one of claims 1 to 60, wherein hazard ratio of progression-free survival compared with sorafenib at a dosage of 400 mg twice daily is 0.64.
- [Claim 69] The method of any one of claims 1 to 60, wherein hazard ratio of progression-free survival compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 0.55 and 0.75.
- [Claim 70] The method of any one of claims 1 to 60, wherein progression-free survival is 5 to 8 months.
- [Claim 71] The method of any one of claims 1 to 60, wherein median time to progression is 7.3 months.

- [Claim 72] The method of any one of claims 1 to 60, wherein median time to progression with 95% confidence interval is between 5.6 to 7.5 months.
- [Claim 73] The method of any one of claims 1 to 60, wherein hazard ratio of time to progression compared with sorafenib at a dosage of 400 mg twice daily is 0.65.
- [Claim 74] The method of any one of claims 1 to 60, wherein hazard ratio of time to progression compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 0.56 and 0.77.
- [Claim 75] The method of any one of claims 1 to 60, wherein time to progression is shown in Fig. 6 (Kaplan-Meier Plot of Time to Progression).
- [Claim 76] The method of any one of claims 1 to 60, wherein objective response rate is 19%.
- [Claim 77] The method of any one of claims 1 to 60, wherein odds ratio of objective response rate compared with sorafenib at a dosage of 400 mg twice daily is 3.13.
- [Claim 78] The method of any one of claims 1 to 60, wherein odds ratio of objective response rate compared with sorafenib at a dosage of 400 mg twice daily with 95% confidence interval is between 2.15 to 4.56.
- [Claim 79] The method of any one of claims 1 to 60, wherein the method comprises achieving the results shown in Table 10 (Efficacy Results in HCC).
- [Claim 80] The method of any one of claims 1 to 60, wherein the method comprises achieving the results shown in Fig. 15 (Quality of Life).
- [Claim 81] The method of any one of claims 1 to 80, wherein the human subject consists essentially of the subject with mild hepatic impairment classified in Child-Pugh class A under Child-Pugh Classification.
- [Claim 82] The method of any one of claims 1 to 81, wherein the human subject is categorized to stage B or stage C based on Barcelona Clinic Liver Cancer (BCLC) staging system.
- [Claim 83] The method of any one of claims 1 to 82, wherein lenvatinib or the pharmaceutically acceptable salt thereof is formulated as a capsule.
- [Claim 84] The method of any one of claims 1 to 83, wherein lenvatinib or the pharmaceutically acceptable salt thereof is administered to the human subject orally.
- [Claim 85] The method of any one of claims 1 to 84, wherein lenvatinib or a pharmaceutically acceptable salt thereof is lenvatinib mesylate.

[Fig. 1]




[Fig. 2]




[Fig. 3]


Overall Survival




[Fig. 4]

Progression-free Survival

[Fig. 5]




[Fig. 6]




[Fig. 7]

Protocol: E7080-G000-304


Figure 14.2.4.3
Forest Plot of Hazard Ratio for Lenvatinib vs Sorafenib in Time to Progression with Stratification Factors in IxRS
Full Analysis Set




[Fig. 8]




[Fig. 9]



[Fig. 10]




[Fig. 11]



[Fig. 12]

Protocol: E7080-G000-304


Figure 14.2.4.3
Forest Plot of Hazard Ratio for Lenvatinib vs Sorafenib in Time to Progression with Stratification Factors in IvRS
Full Analysis Set




[Fig. 13]

Protocol: E7080-G000-304



Figure 14.2.4.3
Forest Plot of Hazard Ratio for Lenvatinib vs Sorafenib in Time to Progression with Stratification Factors in hRS
Full Analysis Set



[Fig. 14]



[Fig. 15]



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2018/018810

A. CLASSIFICATION OF SUBJECT MATTER		
Int.Cl. A61K31/47(2006.01)i, A61P1/16(2006.01)i, A61P35/00(2006.01)i		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Int.Cl. A61K31/47, A61P1/16, A61P35/00		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Published examined utility model applications of Japan 1922-1996 Published unexamined utility model applications of Japan 1971-2018 Registered utility model specifications of Japan 1996-2018 Published registered utility model applications of Japan 1994-2018		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CAplus/REGISTRY/MEDLINE/EMBASE/BIOSIS (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	IKEDA, Kenji et al., Phase 2 study of lenvatinib in patients with advanced hepatocellular carcinoma, J. Gastroenterol., 2016.10.04, Vol.52, pp.512-519, [online], [retrieved on 2018.07.27], Retrieved from the Internet:<DOI:10.1007/s00535-016-1263-4>, the whole document, especially, abstract, p.513	1, 3, 5, 7, 9-10, 12, 14, 16, 17, 19, 21-32, 34, 36, 38, 40, 42, 47, 49, 51, 53, 55, 57, 59, 61, 85
Y	'study design and treatment', table 1, p.517 right-hand column lines 4-13, table S1	2, 4, 6, 8-9, 11, 13, 15-16, 18, 20-31, 33, 35, 37, 39, 41-46, 48, 50, 52, 54, 56, 58, 60-80, 82-85
☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
30.07.2018		07.08.2018
Name and mailing address of the ISA/JP		Authorized officer
Japan Patent Office		IMAMURA, Akiko
3-4-3, Kasumigaseki, Chiyoda-ku, Tokyo 100-8915, Japan		4C 3847
		Telephone No. +81-3-3581-1101 Ext. 3452

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2018/018810

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	IKEDA, Masafumi et al., Safety and pharmacokinetics of lenvatinib in patients with advanced hepatocellular carcinoma, Clin. Cancer Res., 2015.10.23, [online], [retrieved on 2018.07.27], Retrieved from the Internet:<DOI:10.1158/1078-0432.CCR-15-1354>, the whole document, especially, abstract, table s1-2, p.12 lines 5-6	2, 82-85 2, 4, 6, 8-9, 11 , 13, 15-16, 18 , 20-31, 33, 35 , 37, 39, 41-46 , 48, 50, 52, 54 , 56, 58, 60-80 , 82-85
P, X	KUDO, Masatoshi et al., Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial, Lancet, 2018.02.09, Vol.391, pp.1163-1173, [online], [retrieved on 2018.07.27], Retrieved from the Internet:<DOI:10.1016/s0140-6736(18)30207-1>, the whole document	1, 3, 5, 7, 9-10 , 12, 14, 16, 17 , 19, 21-32, 34 , 36, 38, 40, 42 , 47, 49, 51, 53 , 55, 57, 59, 61 -85
P, X	Pharmaceuticals Interview Form `LENVIMA`, Ver.5, PMDA, 2018.03, the whole document	1-85
A	Pharmaceuticals Interview Form `LENVIMA`, Ver.1, PMDA, 2015.03, the whole document	1-85
A	MIZUSHIMA, Toru, Drug repositioning, BIO INDUSTRY, Vol.31, No.11, 2014.11.12, pp.4-10, the whole document	1-85