



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

(51) Int.Cl.: **A 61 K 9/66 (2006.01)** **A 61 K 9/00 (2006.01)** **A 61 K 31/395 (2006.01)**
A 61 K 31/498 (2006.01) **A 61 K 31/7048 (2006.01)** **A 61 P 31/04 (2006.01)**

(45) Oversættelsen bekendtgjort den: **2019-01-21**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2018-12-05**

(86) Europæisk ansøgning nr.: **09708397.6**

(86) Europæisk indleveringsdag: **2009-02-05**

(87) Den europæiske ansøgnings publiceringsdag: **2010-11-10**

(86) International ansøgning nr.: **AU2009000129**

(87) Internationalt publikationsnr.: **WO2009097651**

(30) Prioritet: **2008-02-08 US 65144**

(84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR**

(73) Patenthaver: **Red Hill Biopharma Ltd., 22 Givati Street, 52232 Ramat-Gan, Israel**

(72) Opfinder: **BORODY, Thomas Julius, Level 1, 229 Great North Road, Five Dock, New South Wales 2046, Australien**
GOSSELIN, Patrick, 200, Armand-Frappier, Laval, Quebec H7V 4A6, Canada

(74) Fuldmægtig i Danmark: **Inspicos P/S, Kogle Alle 2, 2970 Hørsholm, Danmark**

(54) Benævnelse: **Methods and compositions for treating inflammatory bowel disease**

(56) Fremdragne publikationer:
WO-A1-98/43667
BOSQUEE L ET AL: "Cervical lymphadenitis caused by a fastidious mycobacterium closely related to Mycobacterium genavense in an apparently immunocompetent woman: Diagnosis by culture-free microbiological methods", JOURNAL OF CLINICAL MICROBIOLOGY, vol. 33, no. 10, 1995, pages 2670-2674, XP002624769, ISSN: 0095-1137
BORODY ET AL: "Anti-mycobacterial therapy in Crohn's disease heals mucosa with longitudinal scars", DIGESTIVE AND LIVER DISEASE, W.B. SAUNDERS, vol. 39, no. 5, 13 April 2007 (2007-04-13) , pages 438-444, XP022045863, ISSN: 1590-8658, DOI: DOI:10.1016/J.DLD.2007.01.008
BORODY, T. J. ET AL.: 'Treatment of severe Crohn's disease using antimycobacterial triple therapy - apprbaching a cure?' DIGESTIVE AND LIVER DISEASE vol. 34, no. 1, January 2002, pages 29 - 38
SELBY, W. ET AL.: 'Two-tear combination antibiotic therapy with clarithromycin, rifabutin, and clofazimine for Crohn's disease.' GASTROENTEROLOGY vol. 132, 2007, pages 2313 - 2319
YAJKO, D.M. ET AL.: 'In vitro activities of rifabutin, azithromycin, ciprofloxacin, clarithromycin, clofazimine, ethambutol, and amikacin in combinations of two, three and four drugs against mycobacterium avium.'
ANTIMICROBIAL AGENTS AND CHEMOTHERAPY vol. 40, no. 3, March 1996, pages 743 - 749
DUBE, M.P. ET AL.: 'Successful short-term suppression of clarithromycin-resistant Mycobacterium avium

complex bacteremia in AIDS.' CLINICAL INFECTIOUS DISEASES vol. 28, January 1999, pages 136 - 138

DESCRIPTION

Background

[0001] Inflammatory bowel disease (IBD) is a disorder of unknown etiology characterized typically by diarrhea, cramping, abdominal pains, weight loss and rectal bleeding, tiredness, anemia, fistulae, perforations, obstruction of the bowel and frequent need for surgical intervention. It encompasses a number of disorders including Crohn's disease, ulcerative colitis, indeterminate colitis, microscopic colitis and collagenous colitis. Such disorders may at times begin clinically with a more benign or milder presentation, resembling Irritable Bowel Syndrome (IBS) which can subsequently progress to increasing inflammation accompanying the IBS and may ultimately develop full-blown IBD. The precise causes of IBD and IBS remain unknown. However, there has been a rapidly growing evidence base that *Mycobacterium avium* subspecies *paratuberculosis* (MAP), and perhaps its various strains and sub-strains, are involved in a variety of different diseases and may play an infective role in a significant proportion of patients with Crohn's disease and may co-exist in other inflammatory bowel disorders listed above.

[0002] Accordingly, there was a need for an effective treatment of MAP-infected IBD, and in particular Crohn's disease. U.S. Patent 6,277,836 to Borody ("Borody") describes treatment of IBD using rifabutin in combination with the macrolide clarithromycin and clofazimine. These were prescribed to be ingested simultaneously but as separate tablets and capsules. It was found that taking the capsules and tablets simultaneously caused unwanted interactions of the medications including a marked elevation in the serum of rifabutin at the expense of suppressing the clarithromycin, whose serum concentrations found later in pharmacokinetic studies, came close to suboptimal even at the recommended oral drug doses, threatening resistance development (Hafrier, R., et al., *Antimicrobial Agents and Chemotherapy*, 1998, 42, 631-639). Conversely, this produced a situation where some patients had blood drug levels which were bordering on adverse effect ranges, e.g., possibly close to causing leucopenia or uveitis. In addition, clofazimine levels with separate drugs were slow to reach equilibrium.

[0003] Borody et al. *Digestive and Liver Disease*, April 2007, 39, 438-444, and Borody et al. *Digest. Liver dis.* 2002;34:29-38 describe anti-mycobacterial therapy in Crohn's disease.

[0004] Considering the above described unwanted interactions and the undesired results of Borody, an improved formulation was desired to address these shortcomings.

Summary of the Invention

[0005] In one aspect, the present disclosure provides a pharmaceutical composition comprising rifabutin, clarithromycin, clofazimine, and a pharmaceutically acceptable carrier,

wherein the amount of clofazimine is 10-15% w/w relative to the amount of clarithromycin and 20-25% w/w relative to the amount of rifabutin.

[0006] The present disclosure provides a method of increasing the reduced metabolism of rifabutin caused by clarithromycin in a patient, comprising co-administering clofazimine with rifabutin and clarithromycin to the patient, wherein the amount of clofazimine is 6-18% w/w relative to the amount of clarithromycin.

[0007] Also provided is a method of reducing the increased metabolism of clarithromycin caused by rifabutin in a patient, comprising co-administering clofazimine with rifabutin and clarithromycin to the patient, wherein the amount of clofazimine is 6-18% w/w relative to the amount of clarithromycin.

[0008] The present disclosure provides a method of treating a patient suffering from or susceptible to a *Mycobacterium paratuberculosis* infection, comprising co-administering to the patient in a single dosage form rifabutin, clarithromycin, and clofazimine in a $9 \pm 0.5:19 \pm 0.5:2 \pm 0.5$ w/w/w ratio.

[0009] Also contemplated is a method of inhibiting occurrence of a *Mycobacterium paratuberculosis* infection in a patient, comprising simultaneously co-administering to the patient in need thereof (i) 225 mg \pm 2 mg rifabutin, (ii) 475 mg \pm 2 mg clarithromycin, and (iii) 50 mg \pm 1 mg clofazimine once each day.

Brief Description of the Drawings

[0010] The following figures depict illustrative aspects of present compositions and methods and not intended to be limiting in any way.

Figures 1a-b depict mean clarithromycin plasma concentration-time profiles in linear and semi-logarithmic plots, respectively.

Figures 2a-b depict mean 14-hydroxyclarithromycin plasma concentration-time profiles in linear and semi-logarithmic plots, respectively.

Figures 3a-b depict mean rifabutin plasma concentration-time profiles in linear and semi-logarithmic plots, respectively.

Figures 4a-b depict mean 25-O-desacetylrifabutin plasma concentration-time profiles in linear and semi-logarithmic plots, respectively.

Figures 5a-b depict mean clofazimine plasma concentration-time profiles in linear and semi-logarithmic plots, respectively.

Detailed Description

1. Compositions

[0011] The present description provides compositions comprising rifabutin, clarithromycin, and clofazimine and methods of using same. The rifabutin, clarithromycin, and clofazimine of the compositions are provided in ratios that yield improved pharmacokinetic properties. The present compositions reduce potentially deleterious elevations of rifabutin serum concentration that resulted from administration of earlier known formulations. In contrast to earlier known formulations, the present compositions further provide the advantage of maintaining patient blood drug levels well below adverse effect ranges, e.g., below ranges implicating leucopenia or uveitis. Moreover, the subject compositions also maintain higher levels of serum concentrations of clarithromycin as compared to earlier formulations, thereby inhibiting resistance development. Furthermore, the present compositions allow reaching minimum effective patient clofazimine serum levels faster than with previous formulations.

[0012] Present compositions according to the invention comprise rifabutin, clarithromycin, clofazimine, and a pharmaceutically acceptable carrier, wherein the amount of clofazimine is 10-15% w/w relative to the amount of clarithromycin (such as, 9-12%, or 0-1 1% w/w) and 20-25% w/w relative to the amount of rifabutin (such as, 20-23%, or 21-23%).

[0013] The present compositions may comprise rifabutin, clarithromycin, and clofazimine in about a 9:19:2 w/w/w ratio, wherein each of the variables are free to vary ± 2 , 1, 0.5, or 0.25 (e.g., 9 ± 0.5 :19 ± 5 :2 ± 0.5). For example in some instances, the present compositions comprise 90mg rifabutin (± 30 , 20, 10, 5, 2, or 1mg), 190mg clarithromycin (± 60 , 40, 20, 10, 5, 2, or 1mg), and 20mg clofazimine (± 10 , 7, 5, 2, or 1 mg). The present compositions comprise 45 mg rifabutin (± 15 , 10, 7, 5, 2, or 1mg), 95mg clarithromycin (± 30 , 20, 10, 5, 2, or 1mg), and 10mg clofazimine (± 6 , 5, 2, or 1mg). The pharmaceutical compositions according to the invention are solid oral dosage forms.

[0014] The present compositions further comprise an absorption enhancer that may improve bioavailability of one or more of the active ingredients. The amount of absorption enhancer is between 300-700% w/w relative to the amount of clofazimine including 400-600% or 450-550% or 475-525%. The absorption enhancer is polyethylene glycol (PEG), having an average molecular weight of between 5000-12000 or 7000-9000 or 7500-8500, for example PEG 8000.

[0015] The present compositions may further include one or more additional excipients, such as MCC-Tabulose type 200, Mg Stearate, SLS-Emal 10Pwd HD, a polysorbate (such as, polysorbate 80), or a combination thereof, including all of these. In some instances, the present compositions include both polyethylene glycol and a polysorbate, such as polysorbate 80, wherein the amount of polysorbate is 30-120% w/w relative to the amount of clofazimine

(such as 50-100%, 50-85%, or 60-75%). Additional excipients contemplated for use with the present compositions are described further below.

[0016] The present compositions may further include one or more ionic or non-ionic surfactants. In particular, the present compositions may comprise sodium lauryl sulfate. The present compositions are provided in a single dosage form, for example a tablet, capsule, caplet or lozenge, etc. Additional contemplated dosage forms are described further below.

2 Methods of Treatment

[0017] The present compositions are useful for treating a patient suffering from or susceptible to a *Mycobacterium paratuberculosis* (MAP) infection. In some instances, such treatments include the treatment of inflammatory bowel disease (IBD), such as Crohn's disease, ulcerative colitis, indeterminate colitis, microscopic colitis and collagenous colitis, in addition to sarcoidosis. In preferred embodiments, the present methods are useful for the treatment of Crohn's disease or colitis.

[0018] Hence, the present methods include a method of treating a patient suffering from or susceptible to a *Mycobacterium paratuberculosis* infection, comprising co-administering to the patient in a single dosage form rifabutin, clarithromycin, and clofazimine according to claim 1.

[0019] In some instances, the rifabutin, clarithromycin, and clofazimine are co-administered once each day for a first period of treatment (for example, 1-3 weeks, including 1 week, 2 weeks or three weeks).

[0020] For the compositions employed in the present methods, the rifabutin, the clarithromycin, and the clofazimine are co-formulated into a single dosage form.

[0021] Any of the above-mentioned compositions are contemplated for use with the present methods. For example, in some instances the present methods contemplate use of compositions comprising an absorption enhancer that may improve bioavailability of one or more of the active ingredients. The amount of absorption enhancer is between 300-700% w/w relative to the amount of clofazimine (such as, 400-600%, 450-550%, or 475-525%). The absorption enhancer is polyethylene glycol, for having an average molecular weight of between 5000-12000, 7000-9000, or 7500-8500, for example PEG 8000.

5. Dosage Forms

[0022] The present compositions may be available in the form of a tablet containing at least one of rifabutin, clarithromycin, and clofazimine in a powdered form. In some instances two or all of rifabutin, clarithromycin, and clofazimine are in a powdered form. Alternatively, present

compositions may be in the form of a tablet capsule containing at least one of rifabutin, clarithromycin, and clofazimine in a microencapsulated form. In some instances, two or all of rifabutin, clarithromycin, and clofazimine are in a microencapsulated form. As another possibility, present compositions may be in the form of a tablet capsule containing at least one of rifabutin, clarithromycin, and clofazimine in a powdered form, and the remaining agents present in a microencapsulated form. As a further possibility, present compositions may be in the form of a tablet capsule containing one or more of rifabutin, clarithromycin, and clofazimine present in a microgranulated form. In additional possibilities, present compositions may be in the form of a tablet containing one or more of rifabutin, clarithromycin, and clofazimine within a capsule, a capsule containing one or more of rifabutin, clarithromycin, and clofazimine within a tablet, a capsule containing one or more of rifabutin, clarithromycin, and clofazimine within an outer capsule containing the other agents, or any combination of the above.

[0023] In a further embodiment, the present compositions comprise an inner capsule containing rifabutin, within an outer capsule containing clarithromycin and clofazimine, wherein clarithromycin and clofazimine may be present in powdered, microencapsulated, or microgranulated forms.

[0024] The present methods may be carried out by administration of one or more tablets/capsules containing rifabutin, clarithromycin, and clofazimine as described above R-rifabutin, clarithromycin, and clofazimine are administered simultaneously in one dose.

[0025] The present compositions may be prepared by means known in the art for the preparation of pharmaceutical compositions including blending, grinding, homogenizing, suspending, dissolving, emulsifying, dispersing, and, where appropriate, mixing of rifabutin, clarithromycin, and clofazimine together with selected excipients, diluents, carriers and adjuvants.

[0026] For oral administration, the present compositions may be in the form of tablets, lozenges, pills, troches, capsules, elixirs, powders, including lyophilized powders, and granules. Slow-release, or delayed-release, forms may also be prepared, for example in the form of coated particles, multi-layer tablets or microgranules.

[0027] Solid forms for oral administration may contain pharmaceutically acceptable binders, sweeteners, disintegrating agents, diluents, flavorings, coating agents, preservatives, lubricants, and/or time delay agents. Suitable binders include gum acacia, gelatin, corn starch, gum tragacanth, sodium alginate, carboxymethylcellulose or polyethylene glycol (PEG). Suitable sweeteners include sucrose, lactose, glucose, aspartame or saccharine. Suitable disintegrating agents include corn starch, methylcellulose, polyvinylpyrrolidone, xanthan gum, bentonite, alginic acid or agar. Suitable diluents include lactose, sorbitol, mannitol, dextrose, kaolin, cellulose, calcium carbonate, calcium silicate or dicalcium phosphate. Suitable flavoring agents include peppermint oil, oil of wintergreen, cherry, orange, or raspberry flavoring. Suitable coating agents include polymers or copolymers of acrylic acid and/or methacrylic acid and/or their esters, waxes, fatty alcohols, zein, shellac or gluten. Suitable preservatives include

sodium benzoate, vitamin E, alpha-tocopherol, ascorbic acid, methyl paraben, propyl paraben or sodium bisulphite. Suitable lubricants include magnesium stearate, stearic acid, sodium oleate, sodium chloride or talc. Suitable time delay agents include glyceryl monostearate or glyceryl distearate.

[0028] Variations, modifications, and other implementations of what is described herein will occur to those of ordinary skill.

Exemplification

[0029] With aspects of the present compositions and methods now being generally described, these will be more readily understood by reference to the following examples.

1. Bioavailability Study

[0030] The objectives of this study were to 1) determine and compare the rate and extent of absorption and 2) to assess the safety and tolerability of 2 test formulations of a combination product of clarithromycin, rifabutin and clofazimine (herein after "triple combination").

[0031] This study followed a randomized, open-label, single-dose, 1-way 2-arm parallel design in 24 normal, healthy, non-smoking male and female subjects. All subjects completed the study, and their data were used for pharmacokinetic and statistical analyses.

[0032] Subjects were admitted to the clinic the day before dosing, and remained until the 24.00 hour post-dose blood draw, at which time they were allowed to leave the clinic and after which they were required to return for subsequent blood draws. Following a high fat meal, subjects received 2 triple combination capsules (dry formulation) or 2 triple combination capsules (PEG formulation) on Day 1 of the study period.

[0033] During the study, 19 blood samples were collected from each subject, for pharmacokinetic and statistical analyses. Over the course of the entire study, approximately 236.5 mL of blood was collected from male subjects and approximately 241.5 mL of blood was collected from female subjects. These volumes include all required samples, as described further below.

[0034] The randomization scheme was computer-generated and subjects were assigned a 15 treatment sequence before Period I dosing.

[0035] This was an open-label study; however, the bioanalytical group was blinded to the randomization scheme. This scheme was made available for statistical and reporting purposes only after the completion of the bioanalytical portion of the study.

[0036] Water was provided *ad libitum* until 1.0 hour pre-dose and after 1.0 hour post-dose. With the exception of the whole milk provided to all subjects during the high fat content meal, the only fluid intake allowed during this time was 240 mL of ambient temperature dosing water.

[0037] Following an overnight fast of at least 10 hours, subjects began a high fat content meal 30 minutes prior to drug administration. Subjects consumed this meal in 30 minutes or less; however, the study drug was administered 30 minutes after the start of the meal. The FDA standard high-fat content breakfast consisted of the following: 2 eggs fried in butter, 2 strips of bacon, 2 slices of toast with butter, 4 ounces of hash brown potatoes and 8 fluid ounces (~240 mL) of whole milk.

[0038] No food was allowed for at least 4 hours post-dose. At 4.5, 9.5, and 13.5 hours post-dose, standardized meals and beverages were provided to the subjects. All meals and beverages were free of alcohol, grapefruit products, xanthines and caffeine and were identical for both study treatments.

Treatments Administered

[0039] Following an overnight fast of at least 10 hours, and 30 minutes after the start of a high fat content meal, subjects received one of the following treatments at 0.00 hour on Day 1 of the study period according to a randomization scheme:

Treatment A: 2 triple combination capsules (dry formulation) with 240 mL of ambient temperature water (Treatment Dose = 190 mg of clarithromycin, 90 mg of rifabutin and 20 mg of clofazimine).

Treatment B: 2 triple combination capsules (PEG formulation) with 240 mL of ambient temperature water (Treatment Dose = 190 mg of clarithromycin, 90 mg of rifabutin and 20 mg of clofazimine).

[0040] The medications were administered orally. The drugs were given 1 or more capsules at a time. All capsules were ingested within 1 minute.

Blood Processing

[0041] Approximately 236.5 mL of blood was collected from male subjects and 241.5 mL of blood was collected from female subjects over the study period, as detailed in:

Table 1:

Procedure	Volume Taken per Subject	
	Male Subjects	Female Subjects
Pre-Study	19.5 mL	19.5 mL,
Serum p-CG Tests	N/AP	5 mL
Interim Lab Tests (Biochemistry and Haematology)	13.5 ml	13.5 mL
PK Blood Samples	190 mL	190 mL
End-of-Study Examination	13.5 mL	13.5 mL
Total Blood Volume	236.5 mL	241.5 mL,

[0042] During the study period, 19 blood samples (1 x 4 mL and 1 x 6 mL tube for each sampling time point) were collected from each subject by direct venipuncture or by Vasofix® intravenous catheter using pre-cooled, labelled blood collection tubes containing potassium ethylenediaminetetraacetic acid (EDTA) as the anticoagulant. Blood samples were collected at 0.00 (pre-dose), 0.25, 0.50, 0.75, 1.00, 1.50, 2.00, 2.50, 3.00, 4.00, 6.00, 8.00, 10.00, 12.00, 16.00, 24.00, 48.00, 60.00, and 96.00 hours post-dose. The clock times of all blood draws for plasma concentration analyses were recorded.

[0043] The 6 mL tubes were used to measure clarithromycin and 14-hydroxyclarithromycin. The 4 mL tubes were used to measure rifabutin, 25-O-desacetylrifabutin and clofazimine.

[0044] The blood samples were stored in an ice bath before centrifugation and were centrifuged as soon as possible under refrigerated conditions (at 4°C) at 3500 rpm for 7 minutes. The collected plasma from each blood collection tube was aliquotted into pre-cooled labelled polypropylene tubes. A minimum of 1.5 mL of plasma was transferred from the 6 mL tubes into the first polypropylene tube, and all remaining plasma, if any, was transferred into a second polypropylene tube. The samples were kept in an ice bath, and flash frozen in an upright position, then stored at -70 ± 10°C until assayed.

[0045] In another procedure, the blood samples were stored in an ice bath before centrifugation and were centrifuged as soon as possible under refrigerated conditions (at 4°C) at 3500 rpm for 7 minutes. The collected plasma from each blood collection tube was aliquotted into pre-cooled labelled polypropylene tubes. A minimum of 0.8 mL of plasma was transferred from the 4 mL tubes into the first polypropylene tube, and all remaining plasma, if any, was transferred into a second polypropylene tube. The samples were kept in an ice bath, and flash frozen in an upright position, then stored at -70 ± 10°C until assayed.

[0046] Upon completion of the clinical portion of the study, all samples were analysed for clarithromycin and 14-hydroxyclarithromycin in the plasma samples or for rifabutin, 25-O-desacetylrifabutin and clofazimine in the plasma samples.

Measurements

[0047] The direct measurements of this study were the plasma concentrations of clarithromycin and 4-hydroxyclarithromycin performed, and rifabutin, 25-0-desacetylrifabutin, and clofazimine performed.

[0048] The pharmacokinetic parameters were derived from the plasma clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-0-desacetylrifabutin, and clofazimine concentrations.

Bioanalyses

[0049] Clarithromycin and 14-hydroxyclarithromycin - Information about these analytes was obtained using routine methods known in the art.

[0050] Rifabutin, 25-0-Desacetylrifabutin, and Clofazimine - Rifabutin, 25-0-desacetylrifabutin, clofazimine, and the internal standard, diltiazem, were extracted by solid phase extraction into an organic medium from 0.20 mL of human plasma. An aliquot of this extract was injected into a High Performance Liquid Chromatography system and detected using a mass spectrometer. The analytes were separated by reverse phase chromatography. Evaluation of the assay was carried out by the construction of an eight (8) point calibration curve (excluding zero concentration) covering the range of 9.996 ng/mL to 1279.470 ng/mL for rifabutin, 2.499 ng/mL to 319.917 ng/mL for 25-0-desacetylrifabutin, and 4.997 ng/mL to 639.586 ng/L, for clofazimine in human plasma.

[0051] The slope and intercept of the calibration curves were determined through weighted linear regression analysis ($I/\text{conc.}^2$). Two calibration curves and duplicate QC samples (at 3 concentration levels) were analysed along with each batch of the study samples. Peak area ratios were used to determine the concentration of the standards, quality control samples, and the unknown study samples from the calibration curves.

Pharmacokinetic Analysis

[0052] The following pharmacokinetic parameters for clarithromycin, rifabutin and clofazimine and the metabolites 14-hydroxyclarithromycin and 25-0-desacetylrifabutin were calculated by standard non-compartmental methods: AUC_{0-t} , $AUC_{0-\infty}$, $AUC_{0-t} / AUC_{0-\infty}$, C_{\max} , T_{\max} , $t_{1/2}$, K_{cl} , and M/P ratio.

[0053] Using General Linear Model (GLM) procedures in Statistical Analysis System (SAS), analysis of variance (ANOVA) was performed on In-transformed AUC_{0-t} , $AUC_{0-\infty}$, $AUC_{0-t} / AUC_{0-\infty}$

$AUC_{0-\text{inf}}$, and C_{max} and on untransformed $t_{1/2}$, K_{el} , and M/P ratio at the significance level of 0.05. The intra-subject coefficient of variation (CV) was calculated using the Mean Square Error (MSE) from the ANOVA table. The ratio of geometric means and the 90% geometric confidence interval (90% C.I.) were calculated based on the difference in the Least Squares Means of the In-transformed AUC_{0-t} , $AUC_{0-\text{inf}}$, $AUC_{0-t} / AUC_{0-\text{inf}}$, and C_{max} between the dry and PEG formulations. T_{max} was analysed using nonparametric methods.

[0054] The pharmacokinetic parameters for clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-O-desacetylrifabutin, and clofazimine derived for both treatments were:

Primary parameters:

[0055]

- AUC_{0-t} = area under the concentration-time curve from time zero to time of last measurable concentration, calculated using the linear trapezoidal rule
- $AUC_{0-\text{inf}}$ = area under the concentration-time curve from time zero to infinity
- C_{max} = maximum plasma concentration after dosing

Secondary parameters:

[0056]

- $AUC_{0-t} / AUC_{0-\text{inf}}$ = Ratio of AUC_{0-t} to $AUC_{0-\text{inf}}$
- T_{max} = time to reach peak plasma concentration
- K_{el} = first order terminal elimination rate constant
- $t_{1/2}$ = terminal half-life
- M/P ratio = Metabolite / Parent ratio for $AUC_{0-\text{inf}}$ -(the conversion to molar units occurs prior to the computation of the ratio).

[0057] The arithmetic mean, standard deviation (SD) and CV were calculated for plasma clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-O-desacetylrifabutin, and clofazimine concentrations for each sampling time and formulation, and for the PK parameters AUC_{0-t} , $AUC_{0-\text{inf}}$, $AUC_{0-t} / AUC_{0-\text{inf}}$, C_{max} , T_{max} , $t_{1/2}$, K_{el} , and M/P ratio.

[0058] ANOVAs (with the following factors: treatment, period, sequence, subject within sequence) were performed on the In-transformed data for AUC_{0-t} , $AUC_{0-\text{inf}}$, $AUC_{0-t} / AUC_{0-\text{inf}}$,

C_{max} . ANOVAs were also performed on the untransformed data to compare the $t_{1/2}$, K_{el} , and M/P ratio. All ANOVAs were performed with the SAS GLM Procedure. T_{max} was analysed using nonparametric methods. The equality of treatment effect in both arms was evaluated using Wilcoxon rank-sum tests. For all analyses, effects were considered statistically significant if the p-value associated with 'F' was less than or equal to 0.050.

[0059] Based on the ANOVA results and the pair-wise comparisons of the ln-transformed AUC_{0-t} , AUC_{0-inf} , AUC_{0-t}/AUC_{0-inf} , C_{max} data, the intra-subject CV, the relative ratios of the geometric means (calculated according to the formula: $e^{[dry-PEG]} \times 100\%$), and the 90% geometric C.I. were determined.

[0060] Twenty-four subjects participated in this study, and samples from the 24 completing subjects (12 per arm) were assayed for drug concentration.

[0061] The principal statistical software used was SAS®, version 8.2. All analyses were performed on the platform of the SAS® suite of statistical programs, using coded procedures that have been written and verified by the staff in the Pharmacokinetics and Statistics Department of BCR.

[0062] Twenty-four subjects (12 males and 12 females) with a mean age of 31 years (range = 18 to 45 years) were enrolled in and completed the study. The completing subjects' mean height was 170 cm (range = 149 to 187 cm) and their mean weight was 71 kg (range = 48 to 104 kg). The subjects' mean BMI was 24.7 kg/m² (range = 18.6 to 29.7 kg/m²). The completing subjects consisted of 9 Caucasians, 5 Asians, 5 Blacks and 4 Hispanics and 1 Mulatto.

Bioanalytical Results

[0063] Clarithromycin and 14-Hydroxyclarithromycin - Information about these analytes was obtained using routine methods known in the art.

[0064] Rifabutin, 25-O-Desacetylrifabutin, and Clofazimine - The plasma samples were analysed for rifabutin, 25-O-desacetylrifabutin, and clofazimine.

[0065] Accuracy and precision of this method were evaluated both within run (intra-assay - Table 2) and between runs (inter-assay - Table 3) by the analysis of the lowest limit of quantification (LLOQ) and Quality Control samples at 3 different concentrations (QC HIGH, QC MED and QC LOW) in human plasma prepared in the range of the calibration/standard curve. The accuracy and precision determined, at each concentration level, were reported as percent relative error (%RE) and percent coefficient of variation (%CV), respectively.

Intra-Batch (TABLE 2)

	Parent	Metabolite	Parent
Analyte name	rifabutin	25-0-desacetylrifabutin	clofazimine
QC Intraday precision range (%)	4.2 to 8.7	5.3 to 10.0	2.3 to 7.7
QC Intraday accuracy range (%)	-3.9 to 1.1	-7.5 to 9.3	0.6 to 7.0

Inter-Batch (TABLE 3)

	Parent	Metabolite	Parent
Analyte name	rifabutin	25-0-desacetylrifabutin	clofazimine
QC Interday precision range (%)	5.1 to 7.6	4.9 to 10.4	5.1 to 7.2
QC Interday accuracy range (%)	-0.2 to 3.0	-1.4 to 3.9	0.4 to 5.2

[0066] Long Term Stability in Matrix - Long term stability of rifabutin, 25-0-desacetylrifabutin, and clofazimine in human plasma can be determined by comparing the concentration of freshly prepared (not frozen) QC samples (QC LOW and QC HIGH) with aged QC samples of the same concentration.

Pharmacokinetic Profiles

[0067] Mean clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-0-desacetylrifabutin, and clofazimine plasma concentration-time profiles (linear and semi-logarithmic plots) are presented in Figures 1, 2, 3, 4, and 5, respectively.

[0068] The mean pharmacokinetic parameters for clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-0-desacetylrifabutin, and clofazimine are summarized below in Tables 4, 5, 6, 7, and 8, respectively.

PHARMACOKINETIC PARAMETERS FOR CLARITHROMYCIN (TABLE 4):

Pharmacokinetic Parameters	Geometric Mean (%CV)	
	Arithmetic Mean \pm SD	
	Triple Combination Capsules	
	Triple Combination Capsules	Triple Combination Capsules
	(dry formulation)	(PEG formulation)
	(A) (n=12)	(B) (n=12)
AUC _{0-t} (nghr/mL.)	2388.54 (66.38)	2972.86 (54.97)
	3123.93 \pm 2073.75	3450.66 \pm 1896.84
AUC _{0-inf}	2462.93 (65.56)	3049.00 (54.50)

Pharmacokinetic Parameters	Geometric Mean (%CV)	
	Arithmetic Mean ± SD	
	Triple Combination Capsules	Triple Combination Capsules
	(dry formulation)	(PEG formulation)
(A) (n=12)	(B) (n=12)	
(ng hr/mL)	3194.76 ± 2094.48	3520.31 ± 1918.52
AUC _{0-t} / AUC _{0-inf}	96.98 (1.76)	97.50 (1.96)
(%)	96.99 ± 1.70	97.52 ± 1.91
C _{max} (ng/mL)	364.30 (55.76)	485.54 (46.14)
	450.69 ± 251.30	549.18 ± 253.41
T _{max} (hr) *	4.00 (1.00 - 6.00)	4.00 (2.00 - 6.00)
T _{1/2} (hr)	3.13 ± 0.72	3.17 ± 0.52
K _{el} (hr ⁻¹)	2.35E-01 ± 6.74E-02	2.24E-01 ± 3.57E-02

* median (min - max)

PHARMACOKINETIC PARAMETERS FOR 14-HYDROXYCLARITHROMYCIN (TABLE 5):

Pharmacokinetic Parameters	Geometric Mean (%CV)	
	Arithmetic Mean ± SD	
	Triple Combination Capsules (A)	Triple Combination Capsules
	(dry formulation)	(PEG formulation)
(n=12)	(B) (n=12)	
AUC _{0-t} (ng hr/mL.)	2671.07 (49.71)	2868.16 (34.51)
	3015.37 ± 1499.06	3119.41 ± 1076.45
AUC _{0-inf} (ng hr/mL)	2704.76 (49.93)	2904.20 (33.94)
	3055.45 ± 1525.66	3145.08 ± 1067.45
AUC _{0-t} / AUC _{0-inf} (%)	98.75 (1.05)	98.76 (1.61)
	98.76 ± 1.03	98.77 ± 1.59
C _{max} (ng/mL)	261.17 (48.85)	296.82 (34.49)
	292.43 ± 142.86	324.26 ± 111.82
T _{max} (hr) *	4.00 (2.00 - 8.03)	4.00 (2.50 - 6.00)
T _{1/2} (hr)	7.52 ± 2.44	6.66 ± 1.82
K _{el} (hr ⁻¹)	1.04E-01 ± 4.21E-02	1.12E-01 ± 3.14E-02
M/P Ratio (hr)	1.20 ± 0.63	0.99 ± 0.37

* median (min-max)

PHARMACOKINETIC PARAMETERS FOR RIFABUTIN (TABLE 6):

Pharmacokinetic Parameters	Geometric Mean (%CV) Arithmetic Mean \pm SD	
	Triple Combination Capsules	Triple Combination Capsules
	(dry formulation)	(PEG formulation)
	(A) (n=12)	(B) (n=12)
AUC _{0-t} (ng hr/mL.)	1461.26 (57.71)	1897.71 (36.42)
	1633.40 \pm 942.60	2023.01 \pm 736.79
AUC _{0-inf} (ng hr/mL)	1499.70 (31.80)	2047.97 (39.63)
	1577.68 \pm 501.76†	2200.58 \pm 872.14‡
AUC _{0-t} / AUC _{0-inf} (%)	83.65 (6.71)	81.83 (8.94)
	83.84 \pm 5.63	82.14 \pm 7.34
C _{max} (ng/mL)	142.75 (39.94)	160.87 (26.66)
	151.41 \pm 60.47	166.39 \pm 44.37
T _{max} (hr) *	6.00 (2.50 - 6.12)	6.00 (4.00 - 8.00)
T _{1/2} (hr)	10.80 \pm 5.68†	14.43 \pm 6.83‡
K _{el} (hr ⁻¹)	8.44E-02 \pm 4.77E-02†	6.07E-02 \pm 3.30E-02‡

*median (min - max) † n=9 ‡ n=8

PHARMACOKINETIC PARAMETERS FOR 25-O-DESACETYL RIFABUTIN (TABLE 7):

Pharmacokinetic Parameters	Geometric Mean (%CV) Arithmetic Mean \pm SD	
	Triple Combination Capsules	Triple Combination Capsules
	(dry formulation)	(PEG formulation)
	(A) (n=12)	(B) (n=12)
AUC _{0-t} (ng hr/mL.)	1461.26 (57.71)	1897.71 (36.42)
	1633.40 \pm 942.60	2023.01 \pm 736.79
AUC _{0-inf} (ng hr/mL)	1499.70 (31.80)	2047.97 (39.63)
	1577.68 \pm 501.76†	2200.58 \pm 872.14‡
AUC _{0-t} / AUC _{0-inf} (%)	83.65 (6.71)	81.83 (8.94)
	83.84 \pm 5.63	82.14 \pm 7.34
C _{max} (ng/mL)	142.75 (39.94)	160.87 (26.66)
	151.41 \pm 60.47	166.39 \pm 44.37
T _{max} (hr) *	6.00 (2.50 - 6.12)	6.00 (4.00 - 8.00)
T _{1/2} (hr)	10.80 \pm 5.68†	14.43 \pm 6.83‡
K _{el} (hr ⁻¹)	8.44E-02 \pm 4.77E-02†	6.07E-02 \pm 3.30E-02‡

median (min-max)† n=9 ≠ n=8

PHARMACOKINETIC PARAMETERS FOR CLOFAZIMINE (TABLE 8):

Pharmacokinetic Parameters	Geometric Mean (%CV) Arithmetic Mean ± SD	
	Triple Combination	Triple Combination
	Capsules (dry formulation)	Capsules (PEG formulation)
	(A) (n=12)	(B) (n=12)
AUC _{0-t} (ng hr/mL.)	696.93 (55.71) 829.07 ± 461.86	680.75 (47.37) 769.72 ± 364.62
AUC _{0-inf} (ng hr/mL)	1242.28 (37.19) 1304.12 ± 484.95†	1030.78 (41.69) 1088.58 ± 453.78‡
AUC _{0-t} / AUC _{0-inf} (%)	76.08 (17.55) 77.55 ± 13.61	64.28 (27.44) 68.26 ± 18.73
C _{max} (ng/mL)	33.01 (54.75) 38.32 ± 20.98	27.82 (38.14) 29.82 ± 11.37
T _{max} (hr) *	8.00 (2.50 - 24.00)	8.00 (4.00 - 12.00)
T _{1/2} (hr)	23.25 ± 4.49†	21.28 1 8.27‡
K _{el} (hr ⁻¹)	3.07E-02 ± 5.75E-03†	3.55E-02 ± 1.14E-02‡

* median (min-max) † n=6 ≠ n=3

[0069] The relative bioavailability analysis results for AUC_{0-t}, AUC_{0-inf}, AUC_{0-t} / AUC_{0-inf}, C_{max}, and for clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-O-desacetylrifabutin, and clofazimine are summarized above in Tables 9, 10, 11, 12, and 13, respectively.

RELATIVE BIOAVAILABILITY ASSESSMENTS FOR CLARITHROMYCIN (TABLE 9):

Parameter	90% C.I.	Ratio of Means
AUC _{0-t}	47.58% to 135.69%	80.34%
AUC _{0-inf}	48.33% to 135.03%	80.78%
C _{max}	46.57% to 120.88%	75.03%

RELATIVE BIOAVAILABILITY ASSESSMENTS FOR 14-HYDROXYCLARITHROMYCIN (TABLE 10):

Parameter	90% C.I.	Ratio of Means
AUC _{0-t}	65.11% to 133.20%	93.13%
AUC _{0-inf}	65.43% to 132.57%	93.13%
C _{max}	61.72% to 125.43%	87.99%

RELATIVE BIOAVAILABILITY ASSESSMENTS FOR RIFABUTIN (TABLE 11):

Parameter	90% C.I.	Ratio of Means
AUC _{0-t}	57.12% to 103.80%	77.00%
AUC _{0-inf}	53.02% to 101.13%	73.23%
C _{max}	71.44% to 110.23%	88.74%

RELATIVE BIOAVAILABILITY ASSESSMENTS FOR 25-0-DESACETYL RIFABUTIN (TABLE 12):

Parameter	90% C.I.	Ratio of Means
AUC _{0-t}	48.10% to 113.68%	73.95%
AUC _{0-inf}	56.74% to 130.54%	86.06%
C _{max}	55.19% to 97.71%	73.43%

RELATIVE BIOAVAILABILITY ASSESSMENTS FOR CLOFAZIMINE (TABLE 13):

Parameter	90% C.I.	Ratio of Means
AUC _{0-t}	66.12% to 158.52%	102.38%
AUC _{0-inf}	75.50% to 192.39%	120.52%
C _{max}	83.26% to 169.02%	118.63%

Pharmacokinetic Conclusions

[0070] Based on data from 12 completing subjects per arm, the pharmacokinetics of clarithromycin, 14-hydroxyclarithromycin, rifabutin, 25-0-desacetyl rifabutin, and clofazimine data were assessed from the following treatments:

Treatment A: Triple Combination Capsules (dry formulation)

Treatment B: Triple Combination Capsules (PEG formulation)

Pharmacokinetic Analysis of Clarithromycin:

[0071] The peak and systemic exposures of clarithromycin were ~25% and ~20% lower after the single oral dose of triple combination capsules (dry formulation) when compared to triple combination capsules (PEG formulation). Also, the statistical results indicated that the 90% confidence intervals of the geometric mean ratios (dry/PEG) for AUC_{0-t}, AUC_{0-inf}, and C_{max} were 47.58% to 135.69%, 48.33% to 135.03%, and 46.57% to 120.88%, respectively.

[0072] A single dose of 250 mg of clarithromycin resulted in a C_{max} of 780 ± 250 ng/mL. The results obtained from dry and PEG formulations of triple combination capsules were

approximately proportional to literature findings. Chu *et al.* (1993) reported that the rise of clarithromycin peak concentrations occur non-linearly to dose which might explain any slight disproportionality between the literature values and those obtained from triple combination capsules dry and PEG formulations.

[0073] There was however, no significant difference in the rate of exposure (T_{max}) of clarithromycin between the 2 formulations, indicating a similar rate of absorption between the dry and PEG formulations (Median T_{max} 4.00 hours). Similarly, the elimination half-life was also found to be similar between the dry and PEG formulations (p value>0.05).

Pharmacokinetic Analysis of 14-Hydroxyclarithromycin:

[0074] The peak and systemic exposures of the metabolite, 14-hydroxyclarithromycin were ~12% and ~7% lower after the single oral dose of triple combination capsules (dry formulation) when compared to triple combination capsules (PEG formulation). Also, the statistical results indicated that the 90% confidence intervals of the geometric mean (dry/PEG) for AUC_{0-t} AUC_{0-inf} , and C_{max} were 65.11% to 133.20%, 65.43% to 132.57%, and 61.72% to 125.43%, respectively.

[0075] Similar to the parent compound, there was no significant difference in the rate of exposure (T_{max}) of 14-hydroxyclarithromycin between the 2 formulations, indicating a similar rate of absorption between the dry and PEG formulations (Median T_{max} 4.00 hours). Similarly, the elimination half-life was also found to be similar between the dry and PEG formulations (p values>0.05).

Pharmacokinetic Analysis of Rifabutin:

[0076] The peak and systemic exposures of rifabutin were ~11% and ~23% lower after the single oral dose of triple combination capsules (dry formulation) when compared to triple combination capsules (PEG formulation). Also, the statistical results indicated that the 90% confidence intervals of the geometric mean ratios (dry/PEG) for AUC_{0-t} AUC_{0-inf} , and C_{max} were 57.12% to 103.80%, 53.02% to 101.13%, and 71.44% to 110.23%, respectively.

[0077] Gatti *et al.* (1998) conducted a comparative study of rifabutin absorption and disposition in HIV-infected patients with or without wasting syndrome. They found that the C_{max} (peak concentration) was 340 ± 140 ng/ml, in 10 HIV patients without wasting syndrome after a single 300 mg dose of rifabutin administered under fasting conditions.

[0078] There was however, no significant difference in the rate of exposure (T_{max}) of rifabutin between the 2 formulations, indicating a similar rate of absorption between the dry and PEG

formulations (Median T_{max} 6.00 hours). Similarly, the elimination half-life was also found to be similar between the dry and PEG formulations (p values>0.05).

Pharmacokinetic Analysis of 25-O-Desacetylrifabutin:

[0079] The peak and systemic exposures of the metabolite 25-O-desacetylrifabutin were ~26% lower after the single oral dose of triple combination capsules (dry formulation) when compared to triple combination capsules (PEG formulation). Also, the statistical results indicated that the 90% confidence intervals of the geometric mean ratios (dry/PEG) for AUC_{0-t} and C_{max} were 55.19% and 97.71%, respectively. Approximately 65% of subjects were excluded from the statistical analysis of AUC_{0-inf} , K_{el} , and $t_{1/2}$ due to the AUC_{0-inf} extrapolation being more than 20%. Hence, the pharmacokinetic discussion was not based on the outcome of AUC_{0-inf} .

[0080] Similar to the parent compound, there was no significant difference in the rate of exposure (T_{max}) of 25-O-desacetylrifabutin between the 2 formulations, indicating a similar rate of absorption between the dry and the PEG formulation (Median dry and PEG T_{max} 6.01 hours and 7.04 hours, respectively). Similarly, the elimination half-life was also found to be similar between the dry and PEG formulations (p values>0.05).

Pharmacokinetic Analysis of Clofazimine:

[0081] The peak exposure of clofazimine was ~19% (C_{max}) higher after the single oral dose of triple combination capsules (dry formulation) when compared to triple combination capsules (PEG formulation). However, the total systemic exposure (AUC_{0-t}) was found to be similar between the 2 formulations, with a geometric mean ratio of 102%. The statistical results indicated that the 90% confidence intervals of the geometric mean ratios (dry/PEG) for AUC_{0-t} and C_{max} were 66.12% to 158.52%, and 83.26% to 169.02%, respectively. Approximately 50% (Treatment A) and 75% (Treatment B) of subjects were excluded from the statistical analysis of AUC_{0-inf} , K_{el} and $t_{1/2}$ due to the AUC_{0-inf} extrapolation being more than 20%. Hence, the pharmacokinetic discussion was not based on the outcome of AUC_{0-inf} .

[0082] Nix *et al.* (2004) reported proportional values after administration of a 200 mg dose. The C_{max} was found to be 227 ng/mL. These values are proportional to the values obtained from administration of triple combination capsules in the current study.

[0083] There was however, no significant difference in the rate of exposure (T_{max}) of clofazimine between the 2 formulations, indicating a similar rate of absorption between the dry and PEG formulations (Median T_{max} 8.00 hours). Similarly, the elimination half-life was also

found to be similar between the dry and PEG formulations (p values>0.05).

Conclusion:

[0084] The relative bioavailability of clarithromycin, rifabutin, clofazimine, and their metabolites were assessed by measuring and comparing the peak and total systemic exposures from the 2 treatments (using AUC_{0-t} , $AUC_{0-\infty}$, and C_{max}).

[0085] The dry/PEG geometric mean ratios of the total systemic exposures (AUCs) for clarithromycin, rifabutin and their metabolites were lower by ~7%-26%. Similarly, the dry/PEG geometric mean ratios of the peak systemic exposures (C_{max}) for clarithromycin, rifabutin, and their metabolites were found to be ~11%-26% lower when compared to the PEG formulation. However, the total systemic exposures for clofazimine (AUC_{0-t}) were similar between the dry formulation and the PEG formulation. The peak exposure of the dry formulation was ~19% (C_{max}) higher than that of the PEG formulation. There was no significant difference in the time to peak concentration for any of the analytes from either the dry or the PEG formulation treatment group.

[0086] Overall, triple combination (dry and PEG formulations) were well tolerated as a single-dose of about 190 mg of clarithromycin, about 90 mg of rifabutin, and about 20 mg of clofazimine, and no significant safety issues emerged.

2 . C_{max} Comparisons with Literature Values

[0087] Clofazimine - C_{max} fed = 227 ng/ml after 200 mg dose (Nix, et al., 2004).

[0088] Triple combination (dry form) 38.32 ± 20.98 ng/mL. Triple combination (PEG form). 29.82 ± 11.37 ng/mL (Bioavailability study with 20 mg). 20 mg dose gives C_{max} of 23 ng/mL.

[0089] Rifabutin - C_{max} (peak concentration) was 340 ± 140 ng/mL in 10 HIV patients without wasting syndrome after a single 300-mg dose of rifabutin administered fasting (Comparative study of rifabutin absorption and disposition in HIV-infected patients with or without wasting syndrome. Gatti G, Di Biagio A, De Pascalis C, Guerra M, Bassetti M, Bassetti D. Int Conf AIDS. 1998; 12: 554 (abstract no. 32171)).

[0090] Triple combination (dry form) 151.41 ± 60.47 ng/mL. Triple combination (PEG Form) 166.39 ± 44.37 ng/mL (Bioavailability study with 90 mg). 90 mg dose gives C_{max} of 102 ng/mL.

[0091] Clarithromycin - 500 mg (four 125-mg capsules, Abbott Laboratories) every 12 hours for 5 doses. C_{max} 2410 ± 670 mg/L and 660 ± 210 ng/mL for metabolite. Single dose of 250

mg resulted in C_{max} of 780 ± 250 ng/mL.

[0092] Triple combination (dry form) 450.69 ± 251.30 , Triple combination (PEG Form) 549.18 ± 253.41 (Bioavailability study with 190 mg). 190 mg dose gives C_{max} of 593 ng/mL.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US6277836B **[0002]**

Non-patent literature cited in the description

- **HAFRIER, R. et al.** *Antimicrobial Agents and Chemotherapy*, 1998, vol. 42, 631-639 **[0002]**
- **BORODY et al.** *Digestive and Liver Disease*, 2007, vol. 39, 438-444 **[0003]**
- **BORODY et al.** *Digest. Liver dis.*, 2002, vol. 34, 29-38 **[0003]**
- **GATTI GDI BIAGIO ADE PASCALIS CGUERRA MBASSETTI MBASSETTI D.** *Int Conf AIDS.*, 1998, vol. 12, 554- **[0089]**

FREMGANGSMÅDER OG SAMMENSÆTNINGER TIL BEHANDLING AF
INFLAMMATORISK TARMSYGDOM

PATENTKRAV

5

1. Farmaceutisk sammensætning, der omfatter

rifabutin;

10

clarithromycin;

clofazimin;

polyethylenglycol;

15

og en farmaceutisk acceptabel bærer,

hvor den farmaceutiske sammensætning er en fast oral doseringsform,

20

hvor polyethylenglycolen,

(i) har en gennemsnitlig molekylvægt på mellem 5000 og 12000 dalton, og

(ii) udgør mellem 300 % og 700 % (vægt/vægt) i forhold til mængden af

25 clofazimin

hvor mængden af clofazimin udgør 10-15 % (vægt/vægt) i forhold til
mængden af clarithromycin og 20-25 % (vægt/vægt) i forhold til mængden af rifabutin.

30 2. Sammensætning ifølge krav 1, hvor rifabutin, clarithromycin og clofazimin
forekommer i et forhold på $9 \pm 0,5:19 \pm 0,5:2 \pm 0,5$ (vægt/vægt/vægt).

3. Sammensætning ifølge krav 1, hvor polyethylenglycolen har en
gennemsnitlig molekylvægt på 7000-9000 dalton.

4. Sammensætning ifølge krav 1, der endvidere omfatter mikrokristallinsk cellulose (MCC), Mg-stearat, natriumlaurylsulfat (SLS), polysorbat 80 eller en kombination deraf.

DRAWINGS




Figure 1a




Figure 1b




Figure 2a




Figure 2b




Figure 3a




Figure 3b




Figure 4a




Figure 4b




Figure 5a




Figure 5b