



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/70	A1	(11) International Publication Number: WO 00/12099 (43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/20139 (22) International Filing Date: 2 September 1999 (02.09.99) (30) Priority Data: 09/145,820 2 September 1998 (02.09.98) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th floor, Oakland, CA 94607-5200 (US). (72) Inventors: SONG, Hongjun; Department of Biology, University of California, San Diego, 3110 Pacific Hall, 9500 Gilman Drive, La Jolla, CA 92093-0357 (US). POO, Mu-ming; Department of Biology, University of California, San Diego, 3110 Pacific Hall, 9500 Gilman Drive, La Jolla, CA 92093-0357 (US). MING, Guo-li; Department of Biology, University of California, San Diego, 3110 Pacific Hall, 9500 Gilman Drive, La Jolla, CA 92093-0357 (US). TESSIER-LAVIGNE, Marc; Department of Anatomy, University of California, San Francisco, 513 Parnassus Avenue, Rm. S-1470, San Francisco, CA 94143-0452 (US). HE, Zhigang; Department of Anatomy, University of California, San Francisco, 513 Parnassus Avenue, Rm. S-1470, San Francisco, CA 94143-0452 (US).	(74) Agent: OSMAN, Richard, Aron; Science & Technology Law Group, 75 Denise Drive, Hillsborough, CA 94010 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: PROMOTERS OF NEURAL REGENERATION		
(57) Abstract <p>The invention provides methods and compositions for promoting neural cell growth and/or regeneration. The general methods involve contacting with an activator of a cyclic nucleotide dependent protein kinase a neural cell subject to growth repulsion mediated by a neural cell growth repulsion factor. The activator may comprise a direct or an indirect activator of the protein kinase; the repulsion factor typically comprises one or more natural, endogenous proteins mediating localized repulsion or inhibition of neural cell growth; and the target cells are generally vertebrate neurons, typically injured mammalian neurons. The subject compositions include mixtures comprising a neural cell, an activator of a cyclic nucleotide dependent protein kinase and a neural cell growth repulsion factor.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Promoters of Neural Regeneration

Inventors: Hong-jun Song, Mu-Ming Poo, Guo-li Ming, Marc Tessier-Lavigne, Zhigang He

5 The research carried out in the subject application was supported in part by NIH grant NS22764. The government may have rights in any patent issuing on this application.

INTRODUCTION

Field of the Invention

10 The field of this invention is nerve cell growth regulation.

Background

 The failure of the adult central nervous system (CNS) to regenerate after injury is a major clinical problem, affecting some 200,000 people in the United States alone. Despite
15 intensive research, an effective approach in promoting significant regeneration of CNS nerve fibers remains lacking. The inability of CNS to regenerate is partly due to inhibitory factors associated with myelin, a cellular structure surrounding the nerve fibers.

 There are currently few effective methods that can promote significant nerve regeneration of severed or damaged nerve fibers. A number of endogenous molecules are
20 known to modulate neural cell growth (1). These factors may exert either attractive or repulsive action on the extension of axonal growth cones (1,2). Experiments in mammals have shown that blocking of some of the inhibitory factors by antibodies could promote regeneration of severed axons in the spinal cord and lead to functional recovery of limb movements.

25 Recent reports have shown that attractive responses to netrins, mediated by the DCC/UNC-40 family of proteins, can be converted to repulsion by coexpression of proteins of the UNC-5 family (3). In addition, attractive effects of brain-derived neurotrophic factor (BDNF) and netrin-1 on *Xenopus* spinal neurites in culture can be converted to repulsion by inhibition of protein kinase A activity (4, 5). We disclose here the opposite phenomenon: that
30 by specific pharmacological manipulations, the action of neural cell inhibitory (or repulsive) factors can be reversed. Such pharmacological treatments change the inhibitory nature of those inhibitory factors into supportive factors and thus promote nerve regeneration.

SUMMARY OF THE INVENTION

We have found that the action of many protein factors that either inhibit or promote nerve growth are mediated through two general mechanisms. Factors such as neurotrophins including brain-derived neurotrophic factor (BDNF) and neurotrophin 3 (NT-3) as well as certain netrins can promote nerve growth in vitro. Nerve fibers grow towards these factors when presented as a localized source. On the other hand, factors such as Semaphorin III, myelin-associated glycoprotein (MAG) and purified myelin, cause collapse of the nerve terminal and block nerve growth. When these factors present as a localized source, nerve fibers grow away from these factors. We have further found that the inhibitory or repulsive effects of the latter factors can be reversed by pharmacological treatments that activate cyclic nucleotide dependent protein kinases, e.g. by increasing the level of cyclic nucleotides, cAMP and cGMP. In addition, we found the action of protein factors that promote nerve growth, e.g. BDNF and NT-3, can also be enhanced by these cyclic nucleotides. Thus by pharmacological interventions that elevate the cyclic nucleotide levels, we are able to help nerve regeneration associated with injuries of the nervous system.

Accordingly, the invention provides methods and compositions for promoting neural cell growth and/or regeneration. The general methods involve contacting with an activator of a cyclic nucleotide dependent protein kinase a neural cell subject to growth repulsion mediated by a neural cell growth repulsion factor. The activator may comprise a direct or indirect activator of the protein kinase, including cyclic nucleotide analogs, activators of a cyclic nucleotide cyclase, NO inducers, inhibitors of a cyclic nucleotide phosphodiesterase, etc. The repulsion factor typically comprises one or more natural, endogenous proteins mediating localized repulsion or inhibition of neural cell growth. Examples include neural cell guidance proteins such as semaphorins, CNS myelin fractions or components thereof such as MAG, etc. The target cells are generally vertebrate neurons, typically injured mammalian neurons in situ. The subject compositions include mixtures comprising a neural cell, an activator of a cyclic nucleotide dependent protein kinase and a neural cell growth repulsion factor.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1A, 1B. (A) The dependence of turning responses on the Sema III concentration in the pipette. Turning angles (mean \pm SEM) ($n=8$) in normal medium (filled symbols) and in medium

containing 8-br-cGMP (100 μ M, open symbols), and for heat-inactivated Sema III (squares). The total number of neurons examined is shown in parenthesis. '*': Significantly different from the data set at zero concentration ($p < 0.05$, Kruskal-Wallis test). (B) Correlation of turning angles induced by a Sema III gradient in normal medium (first turn) and in medium containing 8-br-cGMP (second turn) for the same neuron. The line represents best linear fit of the data ($r = -0.88$, $p = 0.02$).

Fig. 2A-2D. Growth cone turning in a gradient of Sema III. (A) Effects of manipulating cGMP-dependent activities. Angular positions of all growth cones at the end of the one-hour exposure to a Sema III gradient (50 μ g/ml in the pipette) are shown in a cumulative distribution plot for the following conditions: normal medium, medium containing 8-br-cGMP (100 μ M), Sp-cGMPS (10 μ M), Protoporphyrin-9 (PP-9, 10 μ M), S-Nitroso-N-acetylpenicillamine (SNAP, 300 μ M), and Rp-cGMPS (10 μ M). Isolated symbols along the abscissa are median values for corresponding data shown above. (B) Effects of manipulating cAMP-dependent activities. Shown are distributions of turning angles for the following conditions: normal culture medium, medium containing 8-br-cGMP (100 μ M), Rp-cAMPS (20 μ M), Sp-cAMPS (20 μ M), both 8-br-cGMP (100 μ M) and Rp-cAMPS (20 μ M), or both 8-br-cGMP (100 μ M) and Sp-cAMPS (20 μ M). (C) Distribution of turning angles in the absence or presence of α -28 (20 μ g/ml) in normal medium and in medium containing 8-br-cGMP (100 μ M). '*': significantly different ($p < 0.01$, Kolmogorov-Smirnov test). (D) Effects of reducing $[Ca^{2+}]_o$ on turning responses in a Sema III gradient. Due to increased growth rate, the turning in 1 μ M $[Ca^{2+}]_o$ was assayed 30 min after the onset of the gradient. Distribution of turning angles in normal (1 mM) or low (1 μ M) Ca^{2+} medium in the absence or presence of 8-br-cGMP (100 μ M). '‡': not significantly different ($p > 0.2$, Kolmogorov-Smirnov test).

Fig. 3. Percentage of intact growth cones from the explants treated with cyclic nucleotides and/or Sema III. Error bars refer to SEM. '*': Significantly different from the set without pretreatment with 8-br-cGMP and Sp-cAMPS ($P < 0.001$, t-test)

Fig. 4A-4D. (A) Effects of manipulating cyclic nucleotide levels on turning induced by a rMAG gradient (150 μ g/ml in the pipette) in normal medium, in medium containing 8-br-

cGMP (100 μM) or Sp-cAMPS (20 μM), and by heat-inactivated rMAG in normal medium. Results from the latter were significantly different from all three other groups ($p < 0.01$, Kolmogorov-Smirnov test). (B) Effects of reducing $[\text{Ca}^{2+}]_o$ on turning responses induced by rMAG. Distribution of turning angles in normal (1 mM) or low (1 μM) Ca^{2+} medium in the absence or presence of Sp-cAMPS (20 μM). **: Significantly different ($p < 0.05$, Kolmogorov-Smirnov test). (C) Turning responses induced by a gradient of netrin-1 (5 $\mu\text{g}/\text{ml}$ in the pipette) (5), in normal medium and in medium containing Rp-cGMPS (10 μM), 8-br-cGMP (100 μM), Rp-cAMPS (20 μM), or both Rp-cAMPS (20 μM) and 8-br-cGMP (100 μM). Also shown is the control distribution of turning angles observed when the pipette contained only culture medium (no gradient), which is significantly different from all other data sets ($p < 0.05$, Kolmogorov-Smirnov test). (D) Turning induced by a NT-3 gradient (50 $\mu\text{g}/\text{ml}$ in the pipette) in normal medium and in medium containing Rp-cAMPS (20 μM) or Rp-cGMPS (10 μM), or in medium containing 1 μM Ca^{2+} . All data sets were significantly different from the no gradient control ($p < 0.05$, Kolmogorov-Smirnov test).

15

Fig 5. Growth cone turning induced by purified myelin. A gradient of purified myelin (25 $\mu\text{g}/\text{ml}$ in the pipette) was applied in normal culture medium (open circles), in medium containing Sp-cAMPS (20 μM , solid circles), or 8-br-cGMP (100 μM , solid squares). Isolated symbols along the abscissa are median values for corresponding data shown above.

20

Fig 6A, 6B. Effects of purified myelin-associated glycoprotein (MAG) on growth cones of *Xenopus* spinal neurons. A gradient of purified MAG (150 $\mu\text{g}/\text{ml}$ in the pipette) was applied in normal culture medium (open circles), in medium containing Sp-cAMPS (20 μM , solid circles), or in medium containing 8-br-cGMP (100 μM , solid squares). (A) Shown are distributions of turning angles under different conditions. Isolated symbols along the abscissa are median values for corresponding data shown above. (B) Shown are distributions of net neurite extension in the gradient under different conditions.

25

Fig 7A, 7B. Effects of reducing external Ca^{2+} . (A) Purified MAG-induced turning depends on $[\text{Ca}^{2+}]_o$. Shown are distributions of turning angles in normal medium in the absence (solid circles) or presence of Sp-cAMPS (20 μM , solid squares), and in medium containing 1 μM $[\text{Ca}^{2+}]_o$ in the absence (open circles) or presence of Sp-cAMPS (20 μM , open squares),

30

respectively. Isolated symbols along the abscissa are median values for corresponding data shown above. (B) Purified myelin-induced turning depends on $[Ca^{2+}]_o$. Shown are distributions of turning angles in normal medium in the absence (solid circles) or presence of Sp-cAMPS (20 μ M, solid squares), and in medium containing 1 μ M $[Ca^{2+}]_o$ in the absence (open circles) or presence of Sp-cAMPS (20 μ M, open squares), respectively.

DETAILED DESCRIPTION OF THE INVENTION

The general methods involve contacting with an activator of a cyclic nucleotide dependent protein kinase a neural cell subject to growth repulsion mediated by a neural cell growth repulsion factor. Preferred activators enhance the activity of at least one of PKA or PKG. A wide variety of direct and indirect activators of cyclic nucleotide dependent protein kinases are known in the art, or readily identified in assays such as immuno, kinase and cell based assays. Indirect activators are agents which increase the activity of the protein kinase without directly interacting with the kinase, and include any agent which increases the functional activity of the corresponding cyclic nucleotide (e.g. by increasing its synthesis, increasing its availability, decreasing its degradation, etc.). Exemplary activators include cyclic nucleotide analog agonists, activators of cyclic nucleotide cyclases, NO inducers, inhibitors of cyclic nucleotide phosphodiesterases, drugs such as KT5720, etc. Additional activators are readily made by screening candidate agents for activation of the targeted protein kinase, inhibition of a targeted phosphodiesterase (e.g. cAMP or cGMP phosphodiesterase), activation of a targeted cyclase (e.g. guanylate or adenylate cyclase), etc. in conventional in vitro or cell based assays.

The repulsion factor typically comprises one or more natural, endogenous agents mediating localized repulsion or inhibition of the targeted neural cell growth, which repulsion or inhibition is reversible by increasing the activity of a cyclic nucleotide dependent protein kinase in the cell. Such factors are generally present at the site of neuronal cells in situ, particularly at the cite of CNS axons, and provide an endogenous inhibition to nerve cell growth and/or regeneration. A wide variety of such factors are known or are readily identified in cell based assays, such as described herein. Exemplary agents capable of acting as repulsion factors include neural cell guidance proteins such as some semaphorins, netrins, CNS myelin fractions or components thereof such as MAG, etc.

The target cells are generally vertebrate neurons, typically injured mammalian neurons

in situ. A wide variety of methods may be used to effect the contacting of the cell with the activator. For example, for CNS administration, a variety of techniques are available for promoting transfer of therapeutic agents across the blood brain barrier including disruption by surgery or injection, drugs which transiently open adhesion contact between CNS vasculature endothelial cells, and compounds which facilitate translocation through such cells. The compositions may also be amenable to direct injection or infusion, intraocular administration, or within/on implants e.g. fibers such as collagen fibers, in osmotic pumps, grafts comprising appropriately transformed cells, etc.

In a preferred embodiment, the activator is delivered locally and its distribution is restricted. For example, a particular method of administration involves coating, embedding or derivatizing fibers, such as collagen fibers, protein polymers, etc. with therapeutic agents, see also Otto et al. (1989) *J Neuroscience Research* 22, 83-91 and Otto and Unsicker (1990) *J Neuroscience* 10, 1912-1921. Another particular embodiment is adapted from treatment of spinal cord injuries, e.g. Schulz MK, et al., *Exp Neurol.* 1998 Feb; 149(2): 390-397; Guest JD, et al., *J Neurosci Res.* 1997 Dec 1; 50(5): 888-905; Schwab ME, et al., *Spinal Cord.* 1997 Jul; 35(7): 469-473; Tatagiba M, et al., *Neurosurgery.* 1997 Mar; 40(3): 541-546. For example, the subject compositions improve corticospinal tract (CST) regeneration following thoracic spinal cord injury by promoting CST regeneration into human Schwann cell grafts in the methods of Guest et al. (supra). For these data, the human grafts are placed to span a midthoracic spinal cord transection in the adult nude rat, a xenograft tolerant strain. Activators (see Table 1) incorporated into a fibrin glue are placed in the same region. Anterograde tracing from the motor cortex using the dextran amine tracers, Fluororuby (FR) and biotinylated dextran amine (BDA), are performed. Thirty-five days after grafting, the CST response is evaluated qualitatively by looking for regenerated CST fibers in or beyond grafts and quantitatively by constructing camera lucida composites to determine the sprouting index (SI), the position of the maximum termination density (MTD) rostral to the GFAP-defined host/graft interface, and the longitudinal spread (LS) of bulbous end terminals. The latter two measures provide information about axonal die-back. In control animals (graft only), the CST do not enter the SC graft and undergo axonal die-back. As shown in Table 1, the activators dramatically reduce axonal die-back and cause sprouting.

Table I. In Vivo Neuronal Regeneration with Exemplary Activator Formulations

<u>Activator</u>	<u>Formulation</u>	<u>Reduced</u> <u>Die-Back</u>	<u>Promote</u> <u>Sprouting</u>	
	1. Forsklin	5 uM	++++	++++
5	2. 7 β -Deacetyl-7 β -[γ -(morpholino)butyryl]-forsklin	5 uM	++++	++++
	3. 6 β -[β' -(Piperidino)-propionyl]-forsklin	5 uM	++++	++++
	4. 3-Isobutyl-1-methylxanthine (IBMX)	25-100 uM	++++	++++
10	5. Rolipram	2 uM	++++	++++
	6. 8-bromo-cAMP	100 uM	++++	++++
	7. 8-chloro-cAMP	100 uM	++++	++++
	8. 8-(4-chlorophenylthio)-cAMP	100 uM	++++	++++
	9. Dibutyryl-cAMP	100 uM	++++	++++
15	10. Dioctanoyl-cAMP	100 uM	++++	++++
	11. Sp-cAMPS	20 uM	++++	++++
	12. Sp-8-bromo-cAMPS	20 uM	++++	++++
	13. 8-br-cGMP	100 uM	++++	++++
	14. 8-(4-chlorophenylthio)-cGMP	100 uM	++++	++++
20	15. Dibutyryl-cGMP	100 uM	++++	++++
	16. Glyco-SNAP-1	300 uM	++++	++++
	17. Glyco-SNAP-2	300 uM	++++	++++
	18. S-Nitroso-N-acetylpenicillamine	300 uM	++++	++++
	19. NOC-18	100 uM	++++	++++
25	20. NOR-3	100 uM	++++	++++
	21. Protoporphyrin-9	10 uM	++++	++++

In another demonstration of in vivo therapeutic activity, the subject activators are incorporated in the implantable devices described in US Pat No. 5,656,605 and tested for the promotion of in vivo regeneration of peripheral nerves. Prior to surgery, 18 mm surgical-grade silicon rubber tubes (I.D. 1.5 mm) are prepared with or without guiding filaments (four 10-0 monofilament nylon) and filled with test compositions comprising the activators of

Table 1. Experimental groups consist of: 1. Guiding tubes plus Biomatrix 1™ (Biomedical Technologies, Inc., Stoughton, Mass) ; 2. Guiding tubes plus Biomatrix plus filaments; 3-23. Guiding tubes plus Biomatrix 1™ plus activators 1-21 of Table 1 (supra).

5 The sciatic nerves of rats are sharply transected at mid-thigh and guide tubes containing the test substances with and without guiding filaments sutured over distances of approximately 2 mm to the end of the nerves. In each experiment, the other end of the guide tube is left open. This model simulates a severe nerve injury in which no contact with the distal end of the nerve is present.

10 After four weeks, the distance of regeneration of axons within the guide tube is tested in the surviving animals using a functional pinch test. In this test, the guide tube is pinched with fine forceps to mechanically stimulate sensory axons. Testing is initiated at the distal end of the guide tube and advanced proximally until muscular contractions are noted in the lightly anesthetized animal. The distance from the proximal nerve transection point is the parameter measured. For histological analysis, the guide tube containing the regenerated
15 nerve is preserved with a fixative. Cross sections are prepared at a point approximately 7 mm from the transection site. The diameter of the regenerated nerve and the number of myelinated axons observable at this point are used as parameters for comparison.

20 Measurements of the distance of nerve regeneration document the therapeutic effect of groups 3-23. Similarly, plots of the diameter of the regenerated nerve measured at a distance of 7 mm into the guide tube as a function of the presence or absence of one or more activators of the device demonstrate a similar therapeutic effect of all 21 activators tested. No detectable nerve growth is measured at the point sampled in the guide tube with the matrix-forming material alone. The presence of guiding filaments plus the matrix-forming material (no activator) induces only very minimal regeneration at the 7 mm measurement point,
25 whereas dramatic results, as assessed by the diameter of the regenerating nerve, are produced by the device which consisted of the guide tube, guiding filaments and activator compositions. Finally, treatments using guide tubes comprising either a matrix-forming material alone, or a matrix-forming material in the presence of guiding filaments, result in no measured growth of myelinated axons. In contrast, treatments using a device comprising
30 guide tubes, guiding filaments, and matrix containing activator compositions consistently result in axon regeneration, with the measured number of axons being increased markedly by the presence of guiding filaments.

The amount of activator administered depends on the activator, formulation, route of administration, etc. and is generally empirically determined. For example, with cyclic nucleotide activators delivered locally in a solid matrix or semi-solid phase, the administered dose is typically in the range of about 2 mg up to about 2,000 mg, although variations will necessarily occur depending on the target, the host, and the route of administration, etc..

In one embodiment, the invention provides the subject activators combined with a pharmaceutically acceptable excipient suitable for contacting target neuronal cells in situ, such as CNS administration, including as sterile saline or other medium, gelatin, an oil, etc. to form pharmaceutically acceptable compositions. The compositions and/or compounds may be administered alone or in combination with any convenient carrier, solid or semi-solid matrix, diluent, etc. and such administration may be provided in single or multiple dosages. Useful carriers and matrices include solid, semi-solid or liquid media including water and non-toxic organic solvents. In another embodiment, the invention provides the subject compounds in the form of a pro-drug, which can be metabolically converted to the subject compound by the recipient host. A wide variety of pro-drug formulations are known in the art. The compositions may be provided in any convenient form including tablets, capsules, fibers, guides, osmotic pumps, etc. (see, e.g. US Pat Nos. 5,656,605; 5,660,849 and 5,735,863 for delivery systems particularly suited for CNS administration). As such the compositions, in pharmaceutically acceptable dosage units or in bulk, may be incorporated into a wide variety of containers and materials. For example, dosage units may be included in a variety of containers including microcapsules, pumps, fibers, etc.

The compositions may be advantageously combined and/or used in combination with other therapeutic or prophylactic agents, different from the subject compounds. In many instances, administration in conjunction with the subject compositions enhances the efficacy of such agents. For example, the compounds may be advantageously used in conjunction with other neurogenic agents, neurotrophic factors, growth factors, anti-inflammatories, antibiotics etc.; and mixtures thereof, see e.g. *Goodman & Gilman's The Pharmacological Basis of Therapeutics*, 9th Ed., 1996, McGraw-Hill, esp. Chabner et al., *Antineoplastic Agents* at pp.1233.

The subject compositions include ex vivo mixtures comprising a neural cell, an activator of a cyclic nucleotide dependent protein kinase and a neural cell growth repulsion factor. Such mixtures may be used in in vitro screens for identifying suitable activators,

optimizing formulations, delivery concentrations, etc., etc.

The following experimental section and examples are offered by way of illustration and not by way of limitation.

5

EXAMPLES

Collapsin-1/Semaphorin III/D (Sema III), a diffusible member of the semaphorin family can repel or cause collapse of growth cones in culture (6). Defects in Sema III knock-out mice suggest that Sema III creates exclusion zones for axons or drives axonal fasciculation through surround repulsion (7). We analyzed the effect of a microscopic gradient of Sema III on growth cones of the cultured *Xenopus* spinal neuron (8). Sema III-containing saline was applied in pulses from a micropipette positioned 100 μm from the center of the growth cone and at a 45° angle with respect to the original direction of neurite extension. Most growth cones grew away from the pipette. The repulsive response was dose-dependent with a minimal response occurring at an effective concentration of about 10 ng/ml at the growth cone (8). Heat-inactivated Sema III was ineffective (Fig. 1A). The repulsive turning was initiated by active protrusion of filopodia in the direction away from the pipette, with no obvious growth cone collapse during the turning process (9). The rate of neurite extension was unaffected by the presence of the Sema III gradient.

When 8-br-cGMP (10) or Sp-cGMPS (11), membrane permeable agonists of endogenous cGMP signaling pathways, was present in the culture medium (e.g. a gradient of Sema III -50 $\mu\text{g}/\text{ml}$ in the pipette- applied in normal culture medium or in medium containing 100 μM 8-br-cGMP), nearly all growth cones turned toward rather than away from the pipette in the same Sema III gradient (Fig. 2A). Protoporphyrin-9 (PP-9), a guanylate cyclase activator (12), has similar effect (Fig. 2A). Application of a nitric oxide (NO) donor S-Nitroso-N-acetylpenicillamine (SNAP), which activates soluble guanylate cyclase by releasing NO (13), abolished the repulsive turning response without causing a significant attractive response. On the other hand, bath application of Rp-cGMPS (11), a cGMP antagonist and a specific inhibitor of protein kinase G, did not affect the growth cone response. Thus cGMP regulates the direction of growth cone turning induced by Sema III. NO and cGMP can regulate the establishment of the central connections of developing retinal axons and stimulate the synapse formation of developing and regenerating olfactory neurons (14). In contrast to the effect of cGMP analogues, we found that cAMP analogues had no

30

significant effect on the repulsion induced by Sema III gradients (Fig. 2B). However, a cAMP antagonist Rp-cAMPS, but not agonist Sp-cAMPS (15), blocked the conversion of the turning response in the presence of 8-br-cGMP (Fig. 2B), indicating interaction between cAMP- and cGMP-dependent pathways in these neurons.

5 The opposite turning responses are not due to behaviors of different types of neurons in these *Xenopus* cultures (16), and the turning response can be converted in the same neuron. After a repulsive turning response was first elicited by a Sema III gradient, the same neuron was tested by the same gradient in the presence of 8-br-cGMP (100 μ M). We found that the growth cone exhibited an attractive response. Repulsive response was restored after 8-br-cGMP was washed away. Thus the response of the growth cone to a Sema III gradient does not desensitize with time and can be switched between repulsion and attraction in a cGMP-dependent manner. The extent of repulsion versus attraction for each individual neuron appears to be correlated (Fig. 1B), indicating that cGMP affects only the directionality of the response, not the extent to which the growth cone turns.

10 Neuropilins are receptors for several members of the semaphorin family (17-19), and antibodies against the extracellular domain of neuropilin-1 blocks the effects of Sema III *in vitro* (17, 18). A function-blocking antiserum to rat neuropilin-1 (α -28) cross-reacts with the *Xenopus* protein (20). We found that bath application of the α -28 antiserum abolished both the repulsion induced by Sema III under normal conditions and the attraction towards Sema III in the presence of 8-br-cGMP (Fig. 2C). Thus neuropilin-1 function is required for both repulsion and attraction of these growth cones induced by Sema III.

15 Cyclic nucleotides also change the responsiveness of developing rat dorsal root ganglion (DRG) axons to Sema III. When added to cultures of DRG explants (21), 8-br-cGMP, but not Sp-cAMPS, inhibited the collapsing activity of Sema III in a dose-dependent manner (Fig. 3), while 8-br-cGMP or Sp-cAMPS alone had no detectable effect on growth cones in these cultures. In these experiments higher concentrations of 8-br-cGMP (0.5 - 5 mM) were needed to inhibit the collapsing activity of Sema III, suggesting that the turning response of *Xenopus* spinal neurons might be more sensitive to modulation by cGMP than is the collapse of growth cones of rat DRG neurons (9).

20 To examine whether the responses to other repulsive factors can be converted, we studied myelin-associated glycoprotein (MAG), a component of myelin and an inhibitor of axonal regeneration (22). A soluble proteolytic fragment of MAG consisting of its

extracellular domain is released in abundance from myelin *in vivo*, and can potently inhibit axon regeneration (23). We found that a gradient of recombinant protein consisting of the extracellular domain of MAG (rMAG) (24) repelled growth cones of *Xenopus* spinal neurons (Fig. 4A). However, the repulsion by rMAG was not affected by the addition of 8-br-cGMP.

5 On the other hand, when the Sp-cAMPS was added to the medium, the growth cone responses were converted to attraction in the same rMAG gradient (Fig. 4A). Thus the turning response induced by rMAG can be modulated by cAMP-dependent activities. This is reminiscent of the turning responses induced by gradients of BDNF and netrin-1 (4, 5), although the latter factors are normally attractive and are converted to repulsive by inhibition
10 of cAMP-dependent activities.

Cytosolic Ca^{2+} regulates growth cone motility (25). An increase in cytosolic Ca^{2+} levels correlate with growth cone collapse induced by some myelin-associated proteins, but not other factors (26). To examine the involvement of Ca^{2+} in Sema III and rMAG-induced turning, we reduced the extracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_o$) from the normal level of 1
15 mM to 1 μM (4). This resulted in a 2-3 fold increase in the rate of neurite extension, but no change in the turning responses in a Sema III gradient (Fig. 2D). In contrast, the repulsion and attraction induced by gradients of rMAG were both abolished by the reduction of $[\text{Ca}^{2+}]_o$ (Fig. 4B). Thus growth cone turning induced by rMAG, but not Sema III, requires normal $[\text{Ca}^{2+}]_o$.

20 The above studies on growth cone turning induced by Sema III and MAG point to the existence of two distinct pathways involving cAMP and cGMP, with different Ca^{2+} -dependences. Growth cone turning induced by BDNF, acetylcholine, and netrin-1, but not NT-3, depends on both $[\text{Ca}^{2+}]_o$ and cAMP (4, 5). Here, we found that inhibition or activation of cGMP-dependent pathways by Rp-cGMPS or 8-br-cGMP, respectively, did not affect the
25 attractive turning towards netrin-1 ($p > 0.1$, Kolmogorov-Smirnov test) (Fig. 4C). Moreover, repulsive turning induced by the same netrin-1 gradient in the presence of Rp-cAMPS was also not affected by 8-br-cGMP. On the other hand, the attractive response in a NT-3 gradient was converted to repulsive response by inhibiting cGMP-dependence pathways with Rp-cGMPS, while depletion of $[\text{Ca}^{2+}]_o$ or addition of Rp-cAMPS had no effect on the
30 attractive turning (Fig. 4D). Thus, the dependence of turning behavior on $[\text{Ca}^{2+}]_o$ for the four factors examined here (rMAG, netrin-1, Sema III and NT-3) correlates with a dependence on cAMP, not cGMP.

In addition to recombinant Collapsin I/Semaphorin III (Sema III) and myelin-associated glycoprotein (rMAG, consisting of the extracellular domain of MAG), we also tested purified myelin-associated glycoprotein (MAG) and myelin (which have many components) and obtained results similar to those with rMAG (see, Figs. 5-7). For these studies, myelin was made from bovine brain corpus callosum. The tissue was first homogenized. The homogenate then was filtered through a cheese cloth and overlaid on 0.85M sucrose and centrifuged for 25 min at 75,000 g. The material at the interface was collected and resuspended in ice-water and centrifuged for 15 min at 10,000 g. This step was repeated once more, and the purified myelin was collected and washed twice with ice-water. To obtain the soluble form of myelin protein, the myelin fractions were solubilized in 60 mM octylglucoside at 10 mg/ml at 4°C for 1 hr. The supernatants were collected and dialyzed against PBS and F12 medium. Native MAG was purified from myelin after extraction in 1% octylglucoside and separation by DEAE-Sepharose column as described in McKerracher et al., *Neuron* 13, 805-811, 1994 and Li et al., *Journal of Neuroscience Research*, 46, 404-414, 1996.

Studies of the turning response of *Xenopus* neuronal growth cones induced by a number of diffusible factors (4, 5), including those examined here, have implicated cAMP and cGMP in setting the neuronal response to different guidance cues. The guidance cues examined can all be either attractive or repulsive, depending on the status of cytosolic cyclic nucleotides. Manipulations to increase the level of cyclic nucleotide activity favor attraction and manipulations to decrease the level of cyclic nucleotide activity favor repulsion. Since cyclic nucleotides are known to serve as second messengers for a large number of cell surface receptors (27), the response of a growth cone to a particular guidance cue may thus depend critically on other coincident signals received by the neuron. The susceptibility to conversion between attraction and repulsion may enable a growing axon to respond differentially to the same guidance cue at different points along its journey to its final target (28). The reversal of the action of repulsive factors by elevated cyclic nucleotides provides means for promoting nerve regeneration in the central nervous system (CNS), since effective regeneration in the CNS is blocked by inhibitory factors (22, 29), and modulation of cyclic nucleotide levels helps relieve this inhibition and therefore help stimulate regeneration.

Parentetical references and notes:

1. R. Keynes and G.M.W. Cook, *Cell* 83, 161 (1995); M. Tessier-Lavigne and C.S.

- Goodman, *Science* **274**, 1123 (1996).
2. M. Tessier-Lavigne, M. Placzek, A.G. Lumsden, J. Doss, T.M. Jessell, *Nature* **336**, 775 (1988); N. Ishii, W.G. Wadsworth, B.D. Stern, J.G. Culotti, E.M. Hedgecock, *Neuron* **9**, 873 (1992); T. Serafini et al., *Cell* **78**, 409 (1994); T.E. Kennedy, T. Serafini, J.R. de la Torre, M. Tessier-Lavigne, *ibid.* p. 425; S.A. Colamarino and M. Tessier-Lavigne, *ibid.* **81**, 621 (1995).
- 5 3. M. Hamelin, Y. Zhou, M.-W. Su, I.M. Scott, J.G. Culotti, *Nature* **364**, 327 (1993).
4. H-j. Song, G-l. Ming, M-m. Poo, *ibid.* **388**, 275 (1997).
5. G-l. Ming et al., *Neuron* **19**, 1225 (1997).
6. Y. Luo, D. Raible, J.A. Raper, *Cell* **75**, 217 (1993); A.L. Kolodkin, D.J. Matthes, C.S. Goodman, *ibid.* p. 1389; E.K. Messersmith et al., *ibid.* **14**, 949 (1995); A.W. Puschel, R.H. Adams, H. Betz, *Mole. Cell. Neurosci.* **7**, 419 (1996).
- 10 7. O. Behar, J.A. Golden, H. Mashimo, F.J. Schoen, M.C. Fishman, *Nature* **383**, 525 (1996); M. Taniguchi et al., *ibid.* **19**, 519 (1997).
8. Dissociated *Xenopus* spinal neurons were prepared as described (4,5). Isolated spinal neurons were used for experiments at room temperature (22°-24°C) 14-22 hour after plating. Semaphorin III was purified from conditioned medium of stable 293-EBNA cell lines secreting Semaphorin III-AP as described (17). To inactivate Semaphorin III activity, the supernatant containing Semaphorin III-AP was heated at 85°C for 45 minutes [J. Fan and J.A. Raper, *Neuron* **14**, 263 (1995)]. Microscopic gradients of diffusible factors were produced as described (4,5). The average concentration of factors at the growth cone was about 10³-fold lower than that in the pipette [A.M. Lohof, M. Quilian, Y. Dan and M-m. Poo, *J. Neurosci.* **12**, 1253 (1992)], and a concentration gradient of 5-10 % was created across a growth cone 10 µm wide, 100 µm away from the tip of the ejecting pipette. The original direction of neurite growth was defined by the last 10-µm segment of the neurite. The turning angle was defined by the angle between the original direction of neurite extension and a straight line connecting the positions of the growth cone at the onset and the end of the one hour period. Only growth cones with a net extension > 5 µm over the one-hour period were scored.
- 15 20 25
9. No obvious collapse of growth cones was observed even when a higher concentration (up to 100 µg/ml of Semaphorin III in the pipette) was applied to cultured *Xenopus* spinal neurons. We have not examined whether a uniform concentration of Semaphorin III causes collapse, as it does for chick and rat growth cones.
- 30 10. M.A. Schwarzschild and R.E. Zigmond, *J. Neurochem.* **56**, 400 (1991). Pharmacological

agents were added to the culture medium at least 30 minutes before the gradient was applied and were present during the experiments. The final concentrations of pharmacological agents in the medium were listed.

11. E. Butt, M. van Bemmelen, L. Fischer, U. Walter, B. Jastorff, *FEBS Lett.* **263**, 47 (1990).

5 12. M.S. Wolin, K.S. Wood, L.J. Ignarro, *J. Biol. Chem.* **257**, 13312 (1982).

13. E. Southam and J. Garthwaite, *Neurosci. Lett.* **130**, 107 (1991).

14. H.H. Wu, C.V. Williams, S.C. McLoon, *Science* **26**, 1593 (1994); A.J. Roskams, D.S. Bredt, T.M. Dawson, G.V. Ronnett, *Neuron* **13**, 289 (1994); S.M. Gibbs and J.W. Truman, *Neuron* **20**, 83 (1998).

10 15. J.D. Rothermel and L.H. Parker Botelho, *J. Biochem.* **251**, 757 (1988).

16. These *Xenopus* cultures contain spinal neurons sensitive to various neurotransmitters [J.L. Bixby and N.C. Spitzer, *J. Physiol.* **353**, 143 (1984)] exhibiting a wide-range of capability in secreting acetylcholine [J. Evers, M. Laser, Y.A. Sun, Z.P. Xie, M-m. Poo, *J. Neurosci.* **9**, 1523 (1989)]. The variability of turning responses observed within a population of these neurons (see Fig. 2) could be attributed to either the heterogeneity of neuronal types or variability in the cytoplasmic level of cyclic nucleotides or other signal transduction components.

15

17. Z. He and M. Tessier-Lavigne, *Cell* **90**, 739 (1997).

18. A.L. Kolodkin et al., *ibid.* p. 753.

20

19. H. Chen, A. Chedotal, Z. He, C.S. Goodman and M. Tessier-Lavigne, *Neuron* **19**, 547 (1997).

20. The anti-neuropilin-1 antibody (α -28) is directed against the ectodomain of rat neuropilin-1, purified on a protein-A agarose column as described (17), and does not cross-react with neuropilin-2 (19). Using α -28, a putative *Xenopus* neuropilin was detected by Western blot. Interaction of α -28 with putative *Xenopus* neuropilin in cultured *Xenopus* spinal neurons was confirmed by immunostaining. For antibody blocking experiments, α -28 (20 μ g/ml) were added 30 minutes before the onset of the gradient. During the turning assay, the concentration of α -28 was 5 μ g/ml.

25

21. DRG explants derived from E14 rat embryos were cultured with 25 ng/ml NGF on plates pre-coated with poly-D-lysine and laminin for 20 hours before experiments as described (17). Pharmacological agents were added an hour before the collapse assay. The collapse assay was performed on sensory axons from these explants using Sema III-AP-containing medium

30

essentially as described (17). For visualization, growth cones were stained with rhodamine-phalloidin, then washed and mounted.

22. G. Mukhopadhyay, P. Doherty, F.S. Walsh, P.R. Crocker, M.T. Filbin, *Neuron* **13**, 757 (1994); L. McKerracher et al., *ibid.* p. 805; M. Schafer, M. Fruttiger, D. Montag, M.

5 Schachner, R. Martini, *ibid.* **16**, 1107 (1996).

23. S. Sato, R.H. Quarles, R.O. Brady, W.W. Tourtellotte, *Ann. Neurol.* **15**, 264 (1984); S. Tang et al., *Mol. Cell Neurosci.* **9**, 333 (1997).

24. To generate rMAG, SF9 cells were infected with baculovirus expressing the extracellular domain of MAG and conditioned medium was collected 5-6 days after infection and purified
10 as described [M. Li et al., *J. Neurosci. Res.* **46**, 404 (1996)]. For experiments using inactivated rMAG, rMAG was heated at 80°C for 35 min.

25. S.B. Kater, M.P. Mattson, C. Cohan, J.A. Conner, *Trends. Neurosci.* **11**, 317 (1988).

26. J.K. Ivins, J.A. Raper, R.N. Pittman, *J. Neurosci.* **11**, 1597 (1991); C.E. Bandtlow, M.F. Schmidt, T.D. Hassinger, M.E. Schwab, S.B. Kater, *Science* **259**, 80 (1993); J. Loschinger et
15 al., *J. Neurobiol.* **33**, 825 (1997).

27. J. R. Cooper, F.E. Bloom, R.H. Roth, *The Biochemical Basis of Neuropharmacology* (7th ed, Oxford University Press, New York, 1996); R. Laufer and J. Changeux, *EMBO J.* **6**, 901 (1987); J.H. Fong and D.E. Ingber, *Biochem. Biophys. Res. Commun.* **221**, 19 (1996).

28. Attraction of commissural axons towards the floor plate in the developing spinal cord
20 might be switched off (or even converted to a repulsive response) after the contact with the floor plate cells, allowing further axonal growth past the floor plate [see R. Shirasaki, R. Katsumata, F. Murakami, *Science* **279**, 105 (1998)].

29. M.E. Schwab and D. Bartholdi, *Physiol. Rev.* **76**, 319 (1996).

WHAT IS CLAIMED IS:

1. A method for promoting neural cell growth, the method comprising the step of contacting with a composition comprising an effective amount of an activator of a cyclic nucleotide dependent protein kinase a neural cell subject to growth repulsion mediated by a neural cell growth repulsion factor, whereby the growth of the cell is promoted.
5
2. The method of claim 1, wherein the activator comprises an active component selected from a cyclic nucleotide analog, an activator of a cyclic nucleotide cyclase, an NO inducer and an inhibitor of a cyclic nucleotide phosphodiesterase.
10
3. The method of claim 1, wherein the activator comprises an active component selected from an adenylate cyclase activator selected from Forskolin, 7β -Deaceyl- 7β -[γ -(morpholino)butyryl]-forskolin and 6β -[β' -(Piperidino)-propionyl]-forskolin; a phosphodiesterase inhibitor selected from 3-Isobutyl-1-methylxanthine (IBMX) and Rolipram; a PKA activator selected from 8-bromo-cAMP, 8-chloro-cAMP, 8-(4-chlorophenylthio)-cAMP, Dibutyryl-cAMP, Dioctanoyl-cAMP, Sp-cAMPS and Sp-8-bromo-cAMPS; a PKG activator selected from 8-br-cGMP, 8-(4-chlorophenylthio)-cGMP and Dibutyryl-cGMP; an NO donor selected from Glyco-SNAP-1, Glyco-SNAP-2, S-Nitroso-N-acetylpenicillamine, NOC-18 and NOR-3; and a guanylate cyclase activator selected from
15
20 Protoporphyrin-9.
4. The method of claim 1, wherein the repulsion factor comprises an active component selected from a semaphorin, a netrin, a MAG and a CNS myelin fraction.
- 25 5. The method of claim 1, wherein the protein kinase is protein kinase A or G.
6. The method of claim 1, wherein the cell is a mammalian neuron in situ.
7. A method of promoting the growth, repair or regeneration of a neuronal axon subject to growth inhibition by an endogenous neural cell growth repulsion factor, comprising
30 locally administering to a patient in need thereof at the axon a therapeutically effective amount of an activator of a cyclic nucleotide dependent protein kinase.

8. The method of claim 7, wherein the activator comprises an active component selected from a cyclic nucleotide analog, an activator of a cyclic nucleotide cyclase, an NO inducer and an inhibitor of a cyclic nucleotide phosphodiesterase.

5 9. The method of claim 7, wherein the activator comprises an active component selected from an adenylate cyclase activator selected from Forskolin, 7 β -Deaceyl-7 β -[γ -
(morpholino)butyryl]-forsklin and 6 β -[β' -(Piperidino)- propionyl]-forsklin; a
phosphodiesterase inhibitor selected from 3-Isobutyl-1-methylxanthine (IBMX) and
10 Rolipram; a PKA activator selected from 8-bromo-cAMP, 8-chloro-cAMP, 8-(4-
chlorophenylthio)-cAMP, Dibutyryl-cAMP, Dioctanoyl-cAMP, Sp-cAMPS and Sp-8-bromo-
cAMPS; a PKG activator selected from 8-br-cGMP, 8-(4-chlorophenylthio)-cGMP and
Dibutyryl-cGMP; an NO donor selected from Glyco-SNAP-1, Glyco-SNAP-2, S-Nitroso-N-
acetylpenicillamine, NOC-18 and NOR-3; and a guanylate cyclase activator selected from
15 Protoporphyrin

15

10. The method of claim 7, wherein the repulsion factor comprises an active component selected from a semaphorin, a netrin, a MAG and a CNS myelin fraction.

11. The method of claim 7, wherein the protein kinase is protein kinase A or G.

20

12. A mixture comprising a neural cell, an activator of a cyclic nucleotide dependent protein kinase and a neural cell growth repulsion factor.

13. The mixture of claim 12, wherein the activator comprises an active component
25 selected from a cyclic nucleotide analog, an activator of a cyclic nucleotide cyclase, an NO
inducer and an inhibitor of a cyclic nucleotide phosphodiesterase.

14. The mixture of claim 12, wherein the activator comprises an active component
selected from an adenylate cyclase activator selected from Forskolin, 7 β -Deaceyl-7 β -[γ -
30 (morpholino)butyryl]-forsklin and 6 β -[β' -(Piperidino)- propionyl]-forsklin; a
phosphodiesterase inhibitor selected from 3-Isobutyl-1-methylxanthine (IBMX) and
Rolipram; a PKA activator selected from 8-bromo-cAMP, 8-chloro-cAMP, 8-(4-

chlorophenylthio)-cAMP, Dibutyryl-cAMP, Dioctanoyl-cAMP, Sp-cAMPS and Sp-8-bromo-cAMPS; a PKG activator selected from 8-br-cGMP, 8-(4-chlorophenylthio)-cGMP and Dibutyryl-cGMP; an NO donor selected from Glyco-SNAP-1, Glyco-SNAP-2, S-Nitroso-N-acetylpenicillamine, NOC-18 and NOR-3; and a guanylate cyclase activator selected from
5 Protoporphyrin

15. The mixture of claim 12, wherein the repulsion factor comprises an active component selected from a semaphorin, a netrin, a MAG and a CNS myelin fraction.

10 16. The mixture of claim 12, wherein the protein kinase is protein kinase A or G.

17. The mixture of claim 12, wherein the cell is a mammalian neuron.

FIG. 1A





FIG. 1B






FIG. 2A

FIG. 2B

FIG. 2C

FIG. 2D

FIG. 3




FIG. 4B




FIG. 4A




FIG. 4D

Turning angle (degree)

FIG. 4C

Turning angle (degree)

Growth cones with turning angle $\leq X$ (%)

FIG. 5




FIG. 6A







FIG. 6B





INTERNATIONAL SEARCH REPORT

Int. application No.
PCT/US99/20139

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A61K 31/70 US CL :514/42, 47, 48; 435/375 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/42, 47, 48; 435/375 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched none Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST 1.2, Registry, Medline, HCAPLUS		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	RYDEL, R.E. et al. cAMP Analogs Promote Survival and Neurite Outgrowth in Cultures of Rat Sympathetic and Sensory Neurons Independently of Nerve Growth Factor. Proceedings of the National Academy of Sciences of the United States of America, February 1988, Vol. 85, pages 1257-1261, see the abstract.	1-17
X	GENAIN, C.P. et al. Prevention of Autoimmune Demyelination in Non-Human Primates by a cAMP-Specific Phosphodiesterase Inhibitor. Proceedings of the National Academy of Sciences of the United States of America. April 1995, Vol. 92, pages 3601-3605, see the abstract.	1-17
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* "A" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* * & * later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 26 NOVEMBER 1999	Date of mailing of the international search report 23 DEC 1999	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer  KATHLEEN KAHLER FONDA Telephone No. (703) 308-1235	