



(86) Date de dépôt PCT/PCT Filing Date: 2016/01/07
(87) Date publication PCT/PCT Publication Date: 2016/07/14
(45) Date de délivrance/Issue Date: 2021/07/20
(85) Entrée phase nationale/National Entry: 2017/07/05
(86) N° demande PCT/PCT Application No.: US 2016/012496
(87) N° publication PCT/PCT Publication No.: 2016/112195
(30) Priorités/Priorities: 2015/01/09 (US62/101,969);
2015/04/20 (US62/150,236)

(51) Cl.Int./Int.Cl. C12N 15/63(2006.01),
A61K 39/235(2006.01), C12N 15/861(2006.01)

(72) Inventeurs/Inventors:
JONES, FRANK R., US;
GABITZSCH, ELIZABETH, US;
LATCHMAN, YVETTE, US;
RICE, ADRIAN, US

(73) Propriétaire/Owner:
ETUBICS CORPORATION, US

(74) Agent: SMART & BIGGAR LLP

(54) Titre : METHODES ET COMPOSITIONS D'IMMUNOTHERAPIE COMBINEE
(54) Title: METHODS AND COMPOSITIONS FOR COMBINATION IMMUNOTHERAPY

(57) Abrégé/Abstract:

Methods for generating immune responses using adenovirus vectors that allow multiple vaccinations with the same adenovirus vector and vaccinations in individuals with preexisting immunity to adenovirus are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 July 2016 (14.07.2016)

(10) International Publication Number
WO 2016/112195 A1

(51) International Patent Classification:
C12N 15/63 (2006.01) *A61K 39/235* (2006.01)
C12N 15/861 (2006.01)

(74) Agents: **MCANDREW, Christopher W.** et al.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, California 94304 (US).

(21) International Application Number:
PCT/US2016/012496

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
7 January 2016 (07.01.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/101,969 9 January 2015 (09.01.2015) US
62/150,236 20 April 2015 (20.04.2015) US

(71) Applicant: **ETUBICS CORPORATION** [US/US]; 410 West Harrison Street, Suite 100, Seattle, Washington 98119 (US).

(72) Inventors; and
(71) Applicants : **JONES, Frank R.** [US/US]; 410 West Harrison Street, Suite 100, Seattle, Washington 98119 (US). **GABITZSCH, Elizabeth**; 410 West Harrison Street, Suite 100, Seattle, Washington 98119 (US). **LATCHMAN, Yvette**; 410 West Harrison Street, Suite 100, Seattle, Washington 98119 (US). **RICE, Adrian**; 410 West Harrison Street, Suite 100, Seattle, Washington 98119 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2016/112195 A1

(54) Title: METHODS AND COMPOSITIONS FOR COMBINATION IMMUNOTHERAPY

(57) Abstract: Methods for generating immune responses using adenovirus vectors that allow multiple vaccinations with the same adenovirus vector and vaccinations in individuals with preexisting immunity to adenovirus are provided.

METHODS AND COMPOSITIONS FOR COMBINATION IMMUNOTHERAPY

CROSS-REFERENCE

[0001] This application claims priority to U.S. Provisional Application No. 62/101,969, filed January 9, 2015, and U.S. Provisional Application No. 62/150,236, filed April 20, 2015.

RELATED APPLICATIONS

[0002] This application is related to PCT App. No. PCT/US2013/032688 filed on, March 15, 2013.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0003] This invention was made with United States government support under Contract No. HHSN 261200900059C, awarded by the National Cancer Institute (NCI); and Contract No. HHSN 261201100097C, awarded by the NCI; Grant No. 1R43CA134063, awarded by the NCI; Grant No. 2R44CA134063 awarded by the NCI; Grant No. 1R43CA186357 awarded by the NCI; Grant No. 1R43DE021973 awarded by the National Institute of Dental and Craniofacial Research (NIDCR); and Grant No. 2R44DE021973 awarded by the NIDCR. The United States government may have certain rights in the invention.

BACKGROUND OF THE INVENTION

[0004] Vaccines help the body fight disease by training the immune system to recognize and destroy harmful substances and diseased cells. Vaccines can be largely grouped into two types, preventive and treatment vaccines. Prevention vaccines are given to healthy people to prevent the development of specific diseases, while treatment vaccines, also referred to as immunotherapies, are given to a person who has been diagnosed with disease to help stop the disease from growing and spreading or as a preventive. Viral vaccines are currently being developed to help fight infectious diseases and cancers. These viral vaccines work by inducing expression of a small fraction of genes associated with a disease within the host's cells, which in turn, enhance the host's immune system to identify and destroy diseased cells. As such, clinical response of a viral vaccine can depend on the ability of vaccine to obtain a high level immunogenicity and have sustained long-term expression. Immune checkpoints, such as immune inhibitory pathways, can play a critical in modulating the duration and amplitude of physiological immune responses underlying immunogenicity. By combining the administration of a vaccine with drugs that inhibit the immune checkpoint immune inhibitory pathways one may be able to enhance the efficacy and effectiveness of a vaccine in a patient.

[0005] Cancer immunotherapy achieved by delivering tumor-associated antigens (TAA) may have survival benefits; however, limitations to these strategies exist and more immunologically potent vaccines are needed. A variety of vaccination strategies including co-administration of adjuvants and immune stimulating cytokines have been employed to address the low immunogenicity of self-tumor

antigens. Alternatives include the use of recombinant viral vectors that inherently provide innate pro-inflammatory signals and express the antigen of interest. Adenovirus serotype-5 (Ad5)-based immunotherapeutics have been repeatedly used in humans to induce robust T-cell-mediated immune (CMI) responses while remaining safe. Although, Ad5 vectors have been manufactured in large quantities and are stable for storage and delivery for outpatient administration, a major obstacle to the use of first generation (E1-deleted) Ad5-based vectors is the high frequency of pre-existing anti-Ad5 neutralizing antibodies (NAb), which may be present in a host from prior wild type adenovirus infection or induction of Ad5 NAb by repeated injections with Ad5-based vaccines, each resulting in inadequate immune stimulation against the target TAA.

[0006] A major problem with adenovirus (Ad) vectors has been their inability to sustain long-term transgene expression due largely to host immune responses that eliminates the Ad vector and virally-transduced cells in immune-competent subjects. Use of first generation Ad vector vaccines is severely limited by preexisting or induced immunity of vaccines to Ad. Most humans have Ad5 NAb, and two-thirds have lympho-proliferative responses against Ad. An Ad vector vaccine carrying an HIV-1 envelope gene was incapable of re-immunizing a primed immune response using DNA without adjuvant. Single immunizations of non-human primates were unable to generate transgene-specific antibodies to HIV proteins, or alter overall T-cell responses.

[0007] Numerous mechanisms by which preexisting immunity interferes with Ad vector vaccines exist, but one major concern is the presence of NAb followed by CMI elimination of Ad infected antigen harboring cells. Both of these responses can be directed to several Ad proteins. Although increasing vaccine doses may increase induction of desired CMI responses in Ad-immune animals, often unacceptable adverse effects result in animals and humans. Using first generation Ad5 vectors, a heterologous prime-boost regimen using naked (non-vectored) DNA as the priming vaccination, followed by an Ad5 vector immunization may result in a subsequent immune response against Ad5 such that one cannot administer a further re-immunization (boost) with the same (or a different) Ad vector vaccine that utilizes the same viral backbone. Thus, using this approach with current first generation Ad5 vectors can abrogate further use of Ad5 vector immunizations.

[0008] First generation (E1-deleted) Ad vector vaccines express Ad late genes, albeit at a decreased level and over a longer time period than wild-type Ad virus, and vaccine antigens are presented to the immune system simultaneously with highly immunogenic Ad capsid proteins. Due to antigenic competition, the immune responses generated are less likely directed to desired vaccine epitopes and more likely directed to the Ad-derived antigens. One of the major problems facing Ad5 based vectors is the high propensity of pre-existing immunity to Ads in the human population, and how this may preclude the use of conventional, Ad5 [E1-] deleted (first generation Ads) in most human populations, for any additional vaccine application. A major obstacle to the use of first generation (E1-deleted) Ad5-based vectors is the

high frequency of pre-existing anti-adenovirus type 5 neutralizing antibodies. These antibodies can be present in a potential vaccine due to either prior wild type adenovirus infection and/or induction of adenovirus neutralizing antibodies by repeated injections with Ad5-based vaccines, each resulting in inadequate immune stimulation against the target TAA.

[0009] Thus, there remains more effective cancer vaccine vector candidates are needed. Cancer targeting Ad vaccine vectors that allow multiple vaccinations and vaccinations in individuals with preexisting immunity to Ad are needed. While cancer immunotherapy achieved by delivering tumor-associated antigens (TAA) provides survival benefits, limitations to these strategies exist and more immunologically potent vaccines are needed.

SUMMARY OF THE INVENTION

[0010] To overcome these challenges, the present invention provides combination multi-targeted vaccines, immunotherapies and methods for enhanced therapeutic response to complex diseases such as infectious diseases and cancers. The present disclosure relates compositions, methods and kits for generating an immune response in an individual to fight infectious diseases and cancer. The present disclosure provides compositions, methods and kits for generating an immune response against a target antigen or cells expressing or presenting a target antigen or a target antigen signature comprising at least one target antigen.

[0011] It has been discovered that Ad5 [E1-, E2b-] vectors are not only are safer than, but appear to be superior to Ad5 [E1-] vectors in regard to induction of antigen specific immune responses, making them much better suitable as a platform to deliver MUC1, T and/or CEA vaccines that can result in a clinical response. In other cases, immune induction may take months. Ad5 [E1-, E2b-] vectors not only are safer than, but appear to be superior to Ad5 [E1-] vectors in regard to induction of antigen specific immune responses, making them much better suitable as a platform to deliver MUC1c, T and/or CEA vaccines that can result in a clinical response.

[0012] Various embodiments of the invention, by taking advantage of the new Ad5 [E1-, E2b-] vector system in delivering a long sought-after need for a develop a therapeutic vaccine against MUC1, T and/or CEA, overcome barriers found with other Ad5 systems and permit the immunization of people who have previously been exposed to Ad5. In other embodiments of the invention, by taking advantage of the new Ad5 [E1-, E2b-] vector system in delivering a long sought-after need for a develop a therapeutic vaccine against MUC1c, T and/or CEA, overcome barriers found with other Ad5 systems and permit the immunization of people who have previously been exposed to Ad5. In other embodiments of the invention, by taking advantage of the new Ad5 [E1-, E2b-] vector system in delivering a long sought-after need for a develop a therapeutic vaccine against MUC1n, T or CEA, overcome barriers found with other Ad5 systems and permit the immunization of people who have previously been exposed to Ad5.

[0013] To address the low immunogenicity of self-tumor antigens, a variety of advanced, multi-component vaccination strategies including co-administration of adjuvants and immune stimulating cytokines are provided. The invention relates to recombinant viral vectors that provide innate pro-inflammatory signals, while simultaneously engineered to express the antigen of interest. Of particular interest are adenovirus serotype-5 (Ad5)-based immunotherapeutics that have been repeatedly used in humans to induce robust T-cell-mediated immune (CMI) responses, all while maintaining an extensive safety profile.

[0014] In one aspect, a composition is provided comprising a recombinant replication defective viral vector comprising a sequence encoding a MUC1-C antigen, wherein the sequence encoding the MUC1-C antigen has at least 80% sequence identity to SEQ ID NO:5 or SEQ ID NO:6. In some embodiments, the MUC1-C antigen comprises a sequence with at least 80% sequence identity to SEQ ID NO:9.

[0015] In one aspect, a composition is provided comprising a recombinant replication defective viral vector comprising a sequence encoding a Brachyury antigen, wherein sequence encoding the Brachyury antigen has at least 80% sequence identity to SEQ ID NO:7 or SEQ ID NO:8. In some embodiments, an immune response against the antigen or cells expressing the antigen is induced in a human administered the viral vector.

[0016] In one aspect, a composition is provided comprising a recombinant replication defective viral vector comprising a sequence encoding at least two antigens selected from the group consisting of a MUC1-C antigen, a Brachyury antigen, and a CEA antigen. In some embodiments, an immune response against the at least two antigens or cells expressing the at least two antigens is induced in a human administered the viral vector.

[0017] In some embodiments, the immune response comprises generation of an antibody to the antigen. In some embodiments, the immune response comprises cell mediated immunity (CMI). In some embodiments, the sequence encoding the MUC1-C antigen has at least 80% sequence identity to SEQ ID NO:5 or SEQ ID NO:6. In some embodiments, the sequence encoding the Brachyury antigen has at least 80% sequence identity to SEQ ID NO:7 or SEQ ID NO:8. In some embodiments, the sequence encoding the CEA antigen has at least 80% sequence identity to SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:3. In some embodiments, the antigen comprises a modification of 25, 15, 10, 5, or less amino acids. In some embodiments, the antigen comprises a modification in 1 amino acid. In some embodiments, the recombinant viral vector is selected from the group consisting of: retrovirus, lentivirus, cytomegalovirus, Sendai virus, HPV virus, and adenovirus. In some embodiments, the recombinant viral vector comprises a replication defective adenovirus vector. In some embodiments, the recombinant viral vector comprises a replication defective adenovirus 5 vector. In some embodiments, the replication defective adenovirus vector comprises a deletion in an E2b gene region. In some embodiments, the replication defective adenovirus vector comprises a deletion in an E1 gene region. In some embodiments, the replication

defective adenovirus vector comprises a deletion in an E3 gene region. In some embodiments, the replication defective adenovirus vector comprises a deletion in an E4 gene region. In some embodiments, the recombinant viral vector effects overexpression of the antigen in transfected cells. In some embodiments, the recombinant viral induces a specific immune response against cells expressing the antigen in a human that is at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or 25 fold over basal. In some embodiments, the human has an inverse Ad5 neutralizing antibody titer of greater than 50, 75, 100, 125, 150, 160, 175, or 200. In some embodiments, the human has an inverse Ad5 neutralizing antibody titer of greater than 250, 500, 750, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, or 4767. In some embodiments, the immune response is measured as antigen specific antibody response.

[0018] In some embodiments, the immune response is measured as antigen specific cell-mediated immunity (CMI). In some embodiments, the immune response is measured as antigen specific IFN- γ secretion. In some embodiments, the immune response is measured as antigen specific IL-2 secretion. In some embodiments, the immune response against the antigen is measured by ELISpot assay. In some embodiments, the antigen specific CMI is greater than 25, 50, 75, 100, 150, 200, 250, or 300 IFN- γ spot forming cells (SFC) per 10^6 peripheral blood mononuclear cells (PBMC). In some embodiments, the immune response is measured by T-cell lysis of CAP-1 pulsed antigen-presenting cells, allogeneic antigen expressing cells from a tumor cell line or from an autologous tumor. In some embodiments, the composition further comprises an immunogenic component. In some embodiments, the immunogenic component comprises a cytokine selected from the group of IFN- γ , TNF α , IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13. In some embodiments, the immunogenic component is selected from the group consisting of IL-7, a nucleic acid encoding IL-7, a protein with substantial identity to IL-7, and a nucleic acid encoding a protein with substantial identity to IL-7. In some embodiments, the CEA antigen comprises a modification that comprises a substitution of aspartate at a position corresponding to position 610 in SEQ ID NO:3. In some embodiments, the composition further comprises a molecular composition comprising an immune pathway checkpoint modulator. In some embodiments, the immune pathway checkpoint modulator activates or potentiates an immune response. In some embodiments, the immune pathway checkpoint modulator inhibits an immune response inhibitor. In some embodiments, the immune pathway checkpoint inhibits an immune response. In some embodiments, the immune pathway checkpoint modulator targets an endogenous immune pathway checkpoint protein or fragment thereof selected from the group consisting of: PD1, PDL1, PDL2, CD28, CD80, CD86, CTLA4, B7RP1, ICOS, B7RPI, B7-H3, B7-H4, BTLA, HVEM, KIR, TCR, LAG3, CD137, CD137L, OX40, OX40L, CD27, CD70, CD40, CD40L, TIM3, GAL9, ADORA, CD276, VTCN1, IDO1, KIR3DL1, HAVCR2, VISTA, and CD244. In some embodiments, the immune pathway checkpoint modulator targets a PD1 protein. In some embodiments, the molecular composition comprises

siRNAs, antisense, small molecules, mimic, a recombinant form of a ligand, a recombinant form of a receptor, antibodies, or a combination thereof.

[0019] In one aspect, a method of selecting a human for administration of the compositions is provided comprising: determining a HLA subtype of the human; and administering the composition to the human, if the HLA subtype is determined to be one of a preselected subgroup of HLA subtypes. In some embodiments, the preselected subgroup of HLA subtypes comprises one or more of HLA-A2, HLA-A3, and HLA-A24.

[0020] In one aspect, a method of treating a human for cancer or an infectious disease is provided comprising administering the recombinant viral vector to the human.

[0021] In one aspect, a method of generating an immune response in a human to MUC1-C, Brachyury, CEA, or any combination thereof is provided comprising administering to the human the composition. In some embodiments, the administering step is repeated at least once. In some embodiments, the administering step is repeated after about 2, 3, 4, 5, or 6 weeks following a previous administering step. In some embodiments, the administering step is repeated after about 2, 3, 4, 5, or 6 months following a previous administering step. In some embodiments, the administering step is repeated twice.

[0022] In one aspect, a method of treatment is provided comprising: selecting a first phase of treatment and a second phase of treatment; during the first phase, administering to a human a total of 3 times, in about 3 week intervals, a first composition comprising a first replication defective adenovirus vector encoding a MUC1-C antigen; and during the second phase, administering to the human a total of 3 times, in about 3 month intervals, a second composition comprising a second replication defective adenovirus vector encoding an antigen that induces an immune response in a human against cells expressing the MUC1-C antigen.

[0023] In one aspect, a method of treatment is provided comprising: selecting a first phase and a second phase of treatment; during the first phase, administering to a human a total of 3 times, in about 3 week intervals, a first composition comprising a first replication defective adenovirus vector encoding a Brachyury antigen; and during the second phase, administering to the human a total of 3 times, in about 3 month intervals, a second composition comprising a second replication defective adenovirus vector encoding an antigen that induces an immune response in a human against cells expressing the Brachyury antigen.

[0024] In one aspect, a method of treatment is provided comprising: selecting a first phase of treatment and a second phase of treatment; during the first phase, administering to a human a total of 3 times, in about 3 week intervals, a first composition comprising a first replication defective adenovirus vector encoding at least two antigens selected from the group consisting of a MUC1-C antigen, a Brachyury antigen, and a CEA antigen; and during the second phase, administering to the human a total of 3 times, in about 3 month intervals, a second composition comprising a second replication defective adenovirus

vector encoding an antigen that induces an immune response in a human against cells expressing the at least two antigens. In some embodiments, the second phase starts about 3 months after the end of the first phase.

[0025] In one aspect, a method of treatment is provided comprising: selecting a first phase of treatment and a second phase of treatment; during the first phase, administering to a human, a total of n times, a first composition comprising a first replication defective adenovirus vector encoding a Brachyury antigen; during the second phase, administering the human, a total of m times, a second composition comprising a second replication defective adenovirus vector encoding an antigen that induces an immune response in a human against cells expressing the Brachyury antigen.

[0026] In one aspect, a method of treatment is provided comprising: selecting a first phase of treatment and a second phase of treatment; during the first phase, administering to a human, a total of n times, a first composition comprising a first replication defective adenovirus vector encoding a MUC1-C antigen; during the second phase, administering the human, a total of m times, a second composition comprising a second replication defective adenovirus vector encoding an antigen that induces an immune response in a human against cells expressing the MUC1-C antigen.

[0027] In one aspect, a method of treatment is provided comprising: selecting a first phase of treatment and a second phase of treatment; during the first phase, administering to a human, a total of n times, a first composition comprising a first replication defective adenovirus vector encoding at least two antigens selected from the group consisting of a MUC1-C antigen, a Brachyury antigen, and a CEA antigen; during the second phase, administering the human, a total of m times, a second composition comprising a second replication defective adenovirus vector encoding the at least two antigens that induces an immune response in a human against cells expressing the at least two antigens. In some embodiments, n is greater than 1. In some embodiments, n is 3. In some embodiments, m is greater than 1. In some embodiments, m is 3. In some embodiments, the first phase is at least 2, 3, 4, 5, 6, 7, or 8 weeks. In some embodiments, the second phase is at least 2, 3, 4, 5, 6, 7, or 8 months. In some embodiments, the second phase starts 3-16 weeks after first phase ends. In some embodiments, in the first phase two administrations of the replication defective adenovirus are at least 18 days apart. In some embodiments, in the first phase two administrations of the replication defective adenovirus are about 21 days apart. In some embodiments, in the first phase two administrations of the replication defective adenovirus are at most 24 days apart. In some embodiments, in the second phase two administrations of the replication defective adenovirus are at least 10 weeks apart. In some embodiments, in the second phase two administrations of the replication defective adenovirus are about 13 weeks apart. In some embodiments, in the second phase two administrations of the replication defective adenovirus are at most 16 weeks apart. In some embodiments, the method further comprises administering a molecular composition comprising an immune pathway checkpoint modulator.

[0028] In one aspect, a method of treatment is provided comprising: selecting a first phase of treatment and a second phase of treatment; during the first phase, administering to a human, a total of n times, a first composition comprising a first replication defective adenovirus vector encoding an antigen that induces an immune response in a human against cells expressing a MUC1-C, Brachyury, or CEA antigen; and during the second phase, administering the human, a total of m times, a second composition comprising a second replication defective adenovirus vector encoding an antigen that is capable of inducing an immune response directed towards cells expressing MUC1-C, Brachyury, or CEA antigen in a human; wherein a molecular composition comprising and an immune pathway checkpoint modulator is administered during the first phase, the second phase, or both.

[0029] In one aspect, a method of treating a subject in need thereof is provided, comprising administering to the subject: (a) a recombinant replication deficient adenovirus vector encoding (i) a MUC1-C antigen, (ii) a Brachyury antigen, or (iii) at least two antigens selected from the group consisting of a MUC1-C antigen, a Brachyury antigen, and a CEA antigen; and (b) a molecular composition comprising an immune pathway checkpoint modulator; thereby generating an immune response in the subject. In some embodiments, (a) and (b) are administered in series. In some embodiments, (a) and (b) are administered at the same time. In some embodiments, (a) and (b) are administered a month apart.

[0030] In some embodiments, the immune pathway checkpoint modulator activates or potentiates an immune response. In some embodiments, the immune pathway checkpoint modulator inhibits an immune response inhibitor. In some embodiments, the immune pathway checkpoint modulator inhibits an immune response. In some embodiments, the immune pathway checkpoint modulator targets an endogenous immune pathway checkpoint protein or fragment thereof selected from the group consisting of: PD1, PDL1, PDL2, CD28, CD80, CD86, CTLA4, B7RP1, ICOS, B7RPI, B7-H3, B7-H4, BTLA, HVEM, KIR, TCR, LAG3, CD137, CD137L, OX40, OX40L, CD27, CD70, CD40, CD40L, TIM3, GAL9, ADORA, CD276, VTCN1, IDO1, KIR3DL1, HAVCR2, VISTA, and CD244. In some embodiments, the immune pathway checkpoint modulator targets a PD1 protein. In some embodiments, the molecular composition comprises siRNAs, antisense, small molecules, mimic, a recombinant form of a ligand, a recombinant form of a receptor, antibodies, or a combination thereof.

[0031] In some embodiments, the immune response is increased at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or 25 fold. In some embodiments, the first replication defective adenovirus vector and the second replication defective adenovirus vector are the same. In some embodiments, the first replication defective adenovirus vector comprises a replication defective adenovirus 5 vector. In some embodiments, the second replication defective adenovirus vector comprises a replication defective adenovirus 5 vector. In some embodiments, the human expresses a human leukocyte antigen of serotype HLA-A2, HLA-A3, or HLA-A24. In some embodiments, the first replication defective adenovirus vector comprises a deletion in

an E2b gene region. In some embodiments, the first replication defective adenovirus vector further comprises a deletion in an E1 gene region. In some embodiments, the first replication defective adenovirus vector further comprises a deletion in an E3 gene region. In some embodiments, the first replication defective adenovirus vector further comprises a deletion in an E4 gene region. In some embodiments, the second replication defective adenovirus vector comprises a deletion in an E2b gene region. In some embodiments, the second replication defective adenovirus vector further comprises a deletion in an E1 gene region. In some embodiments, the second replication defective adenovirus vector further comprises a deletion in an E3 gene region. In some embodiments, the second replication defective adenovirus vector further comprises a deletion in an E4 gene region. In some embodiments, the first composition, the second composition, or both, comprises at least 1.0×10^{11} , 2.0×10^{11} , 3.0×10^{11} , 3.5×10^{11} , 4.0×10^{11} , 4.5×10^{11} , 4.8×10^{11} , 4.9×10^{11} , 4.95×10^{11} , or 4.99×10^{11} virus particles comprising the recombinant nucleic acid vector. In some embodiments, the first composition, the second composition, or both, comprises at most 7.0×10^{11} , 6.5×10^{11} , 6.0×10^{11} , 5.5×10^{11} , 5.2×10^{11} , 5.1×10^{11} , 5.05×10^{11} , or 5.01×10^{11} virus particles. In some embodiments, the first composition, the second composition, or both, comprises 1.0 - 7.0×10^{11} or 1.0 - 5.5×10^{11} virus particles. In some embodiments, the first composition, the second composition, or both, comprises 4.5 - 5.5×10^{11} virus particles. In some embodiments, the first composition, the second composition, or both, comprises 4.8 - 5.2×10^{11} virus particles. In some embodiments, the first composition, the second composition, or both, comprises 4.9 - 5.1×10^{11} virus particles. In some embodiments, the first composition, the second composition, or both, comprises 4.95 - 5.05×10^{11} virus particles. In some embodiments, the first composition, the second composition, or both, comprises 4.99 - 5.01×10^{11} virus particles. In some embodiments, the first phase is at least 2, 3, 4, 5, 6, 7, or 8 weeks. In some embodiments, the second phase is at least 2, 3, 4, 5, 6, 7, or 8 months. In some embodiments, the immune response is measured as antigen specific antibody response. In some embodiments, the immune response is measured as antigen specific cell-mediated immunity (CMI). In some embodiments, the immune response is measured as antigen specific IFN- γ secretion. In some embodiments, the immune response is measured as antigen specific IL-2 secretion. In some embodiments, the immune response against the antigen is measured by ELISpot assay. In some embodiments, the antigen specific CMI is greater than 25, 50, 75, 100, 150, 200, 250, or 300 IFN- γ spot forming cells (SFC) per 10^6 peripheral blood mononuclear cells (PBMC). In some embodiments, the immune response is measured by T-cell lysis of CAP-1 pulsed antigen-presenting cells, allogeneic antigen expressing cells from a tumor cell line or from an autologous tumor. In some embodiments, a first or a second replication defective adenovirus infects dendritic cells in the human and wherein the infected dendritic cells present the antigen, thereby inducing the immune response. In some embodiments, the administering steps comprise subcutaneous (sc) administration. In some embodiments, the human carries an inverse Ad5 neutralizing antibody titer that is of greater than 50, 75, 100, 125, 150, 160, 175, 200, 225, 250, 275, or 300 prior to the

administering step. In some embodiments, the human has an inverse Ad5 neutralizing antibody titer of greater than 250, 500, 750, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, or 4767. In some embodiments, the human is not concurrently being treated by any one of steroids, corticosteroids, and immunosuppressive agents. In some embodiments, the human does not have an autoimmune disease. In some embodiments, the human does not have inflammatory bowel disease, systemic lupus erythematosus, ankylosing spondylitis, scleroderma, multiple sclerosis, viral hepatitis, or HIV. In some embodiments, the human has or may have in the future an infectious disease. In some embodiments, the human has autoimmune related thyroid disease or vitiligo. In some embodiments, the human has or may have in the future a proliferative disease cancer. In some embodiments, the human has colorectal adenocarcinoma, metastatic colorectal cancer, advanced MUC1-C, Brachyury, or CEA expressing colorectal cancer, breast cancer, lung cancer, bladder cancer, or pancreas cancer. In some embodiments, the human has at least 1, 2, or 3 sites of metastatic disease. In some embodiments, the human comprises cells overexpressing MUC1-C, Brachyury, or CEA. In some embodiments, the cells overexpressing MUC1-C, Brachyury, or CEA, overexpress the MUC1-C, Brachyury, or CEA by at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 times over a baseline MUC1-C, Brachyury, or CEA expression in a non-cancer cell. In some embodiments, the cells overexpressing MUC1-C, Brachyury, or CEA comprise cancer cells. In some embodiments, the subject has a diagnosed disease predisposition. In some embodiments, the subject has a stable disease. In some embodiments, the subject has a genetic predisposition for a disease. In some embodiments, the disease is a cancer. In some embodiments, the cancer is selected from the group consisting of prostate cancer, colon cancer, breast cancer, or gastric cancer. In some embodiments, the cancer is prostate cancer. In some embodiments, the subject has a Gleason score of 6 or less. In some embodiments, the subject has a Gleason score greater than 6. In some embodiments, the first or the second replication defective adenovirus vector comprises a sequence with at least 80% sequence identity to SEQ ID NO:3. In some embodiments, the first or the second replication defective adenovirus vector comprises a region with at least 80% sequence identity to a region in SEQ ID NO:3 selected from 26048-26177, 26063-26141, 1-103, 54-103, 32214-32315, and 32214-32262. In some embodiments, the first or the second replication defective adenovirus vector comprises a region with at least 80% sequence identity to a region in SEQ ID NO:3 between positions 1057 and 3165. In some embodiments, the first or second replication defective adenovirus vector comprises a sequence encoding a MUC1-C, Brachyury, or CEA antigen; wherein the MUC1-C antigen is encoded by a sequence with at least 80% sequence identity to SEQ ID NO:5 or SEQ ID NO:6; wherein the Brachyury antigen is encoded by a sequence with at least 80% sequence identity to SEQ ID NO:7 or SEQ ID NO:8; wherein the CEA antigen is encoded by a sequence with at least 80% sequence identity to SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:3.

[0032] In one aspect a kit for inducing an immune response in a human is provided comprising: a composition comprising a therapeutic solution of a volume in the range of 0.8-1.2 mL, the therapeutic

solution comprising at least 1.0×10^{11} virus particles; wherein the virus particles comprise a recombinant replication defective adenovirus vector; a composition comprising of a therapeutic solution of a molecular composition comprising an immune pathway checkpoint modulator and; instructions.

[0033] In some embodiments, the therapeutic solution comprises $1.0-5.5 \times 10^{11}$ virus particles. In some embodiments, adenovirus vector is capable of effecting overexpression of the modified MUC1-C, Brachyury, or CEA in transfected cells. In some embodiments, therapeutic solution comprises a first, second and third replication defective adenovirus vector each comprising an antigen selected from the group consisting of a MUC1-C antigen, a Brachyury antigen, a CEA antigen, and combinations thereof. In some embodiments, the adenovirus vector comprises a nucleic acid sequence encoding an antigen that induces a specific immune response against MUC1-C, Brachyury, or CEA expressing cells in a human. In some embodiments, the immune pathway checkpoint modulator targets an endogenous immune pathway checkpoint protein or fragment thereof selected from the group consisting of: PD1, PDL1, PDL2, CD28, CD80, CD86, CTLA4, B7RP1, ICOS, B7RPI, B7-H3, B7-H4, BTLA, HVEM, KIR, TCR, LAG3, CD137, CD137L, OX40, OX40L, CD27, CD70, CD40, CD40L, TIM3, GAL9, ADORA, CD276, VTCN1, IDO1, KIR3DL1, HAVCR2, VISTA, and CD244. In some embodiments, the molecular composition comprises siRNAs, antisense, small molecules, mimic, a recombinant form of a ligand, a recombinant form of a receptor, antibodies, or a combination thereof. In some embodiments, the instructions are for the treatment of a proliferative disease or cancer. In some embodiments, the instructions are for the treatment of an infectious disease. In some embodiments, the adenovirus vector comprises a replication defective adenovirus 5 vector. In some embodiments, the therapeutic solution comprises at least 1.0×10^{11} , 2.0×10^{11} , 3.0×10^{11} , 3.5×10^{11} , 4.0×10^{11} , 4.5×10^{11} , 4.8×10^{11} , 4.9×10^{11} , 4.95×10^{11} , or 4.99×10^{11} virus particles comprising the recombinant nucleic acid vector. In some embodiments, the therapeutic solution comprises at most 7.0×10^{11} , 6.5×10^{11} , 6.0×10^{11} , 5.5×10^{11} , 5.2×10^{11} , 5.1×10^{11} , 5.05×10^{11} , or 5.01×10^{11} virus particles. In some embodiments, the therapeutic solution comprises $1.0-7.0 \times 10^{11}$ or $1.0-5.5 \times 10^{11}$ virus particles. In some embodiments, the therapeutic solution comprises $4.5-5.5 \times 10^{11}$ virus particles. In some embodiments, the therapeutic solution comprises $4.8-5.2 \times 10^{11}$ virus particles. In some embodiments, the therapeutic solution comprises $4.9-5.1 \times 10^{11}$ virus particles. In some embodiments, the therapeutic solution comprises $4.95-5.05 \times 10^{11}$ virus particles. In some embodiments, the kit further comprises an immunogenic component. In some embodiments, the immunogenic component comprises a cytokine selected from the group of IFN- γ , TNF α , IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13. In some embodiments, the immunogenic component is selected from the group consisting of IL-7, a nucleic acid encoding IL-7, a protein with substantial identity to IL-7, and a nucleic acid encoding a protein with substantial identity to IL-7.

[0034] Various embodiments of the claimed invention relate to a composition comprising: (a) a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a MUC1 antigen; (b) a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a Brachyury antigen; and (c) a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a CEA antigen, wherein the MUC1 antigen is a MUC1-C antigen.

[0034A] Various embodiments of the claimed invention also relate to a composition for treating a subject in need thereof, wherein the composition comprises: a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a MUC1 antigen, a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a Brachyury antigen; and a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a CEA antigen, characterized in that the composition is for administration to the subject in need thereof; and induces an immune response against the MUC1 antigen, the Brachyury antigen, or the CEA antigen or cells expressing the MUC1 antigen, the Brachyury antigen, or the CEA antigen, wherein the subject in need thereof has cancer, wherein the MUC1 antigen is a MUC1-C antigen.

BRIEF DESCRIPTION OF THE DRAWINGS

[0035] **FIG. 1** exemplifies a bar graph showing antibody levels from mice immunized with Ad5-null (empty vector). Mice were immunized three times with Ad5-null viral particles (VPs) at 14 day intervals. Anti-Ad antibody (neutralizing antibody) levels increased after each immunization.

[0036] **FIG. 2** exemplifies a bar graph showing neutralizing antibody (NAb) levels from mice immunized with Ad5-null. Mice were immunized three times with Ad5-null VPs at 14 day intervals. Neutralizing antibody levels increased after each immunization. Optical density readings indicate the presence of viable target cells.

[0037] **FIG. 3** exemplifies a bar graph showing the induction of NAb in C57Bl/6 mice after injections with Ad5-null vector platform. Increasing levels of NAb were induced in mice after repeated injections with Ad particles.

[0038] **FIG. 4A** exemplifies a bar graph showing INF- γ levels secreted from splenocytes from Ad5-immune mice immunized with Ad5 [E1-]-CEA or Ad5 [E1-, E2b-]-CEA. A significantly elevated response is shown in the Ad5 [E1-, E2b-]-CEA immunized group.

[0039] **FIG. 4B** exemplifies a bar graph showing IL-2 secreted from splenocytes from Ad5-immune mice immunized with Ad5 [E1-]-CEA or Ad5 [E1-, E2b-]-CEA. A significantly elevated response is shown in the Ad5 [E1-, E2b-]-CEA immunized group.

[0040] **FIG. 5** exemplifies a bar graph showing serum aspartate aminotransferase (AST) levels in control mice and mice vaccinated with 1×10^{10} Ad5 [E1-]-CEA VPs or Ad5 [E1-, E2b-]-CEA VPs.

[0041] **FIG. 6** exemplifies a line graph showing tumor volume over time in Ad5-immune C57Bl/6 mice injected with MC38 CEA-expressing tumor cells and subsequently treated (Vac) with Ad5

[E1-, E2b-]-CEA vaccine. Tumor size is shown to be significantly reduced by days 19-21 compared to untreated tumor-bearing mice.

[0042] **FIG. 7** exemplifies a graph showing tumor weight from 7 treated and 7 untreated Ad5-immune MC38 tumor-bearing mice. A significant ($p=0.0124$) reduction in tumor weight in mice treated with Ad5 [E1-, E2b-]-CEA is shown.

[0043] **FIG. 8** is an immunoblot using a mouse monoclonal antibody against Gag and exemplifies Gag production by A549 cells infected with Ad5 [E1-, E2b-]-Gag. A549 whole cell lysate was infected with Ad5 [E1-, E2b-]-gag or Ad5-null at a multiplicity of infection (MOI) of 200 for 44 h. The upper band (55kDa) comprises the gag precursor. The lower band (41kDa) comprises the p17/p24 gag complex.

[0044] **FIG. 9** is a graph exemplifying the effect of multiple immunizations on inducing a greater cell-mediated immunity (CMI) response. Ad5-Naïve BALB/C mice (n=5/group) were immunized once or three times at 14 day intervals with 1×10^{10} Ad5 [E1-]-null VPs, Ad5 [E1-, E2b-]-null VPs, Ad5 [E1-]-Gag VPs, Ad5 [E1-, E2b-]-Gag VPs, or buffer alone. IFN- γ secretion from splenocytes was assessed by ELISpot analysis 14 days after the final immunization. Positive control splenocytes were exposed to Concanavalin A (Con A).

[0045] **FIG. 10A** exemplifies results of an ELISpot INF- γ analysis of peripheral blood mononuclear cells (PBMCs) from Cynomolgus Macaques (n=3) pre-immunized against wild type Ad5. Elevated levels of INF- γ induction ($P < 0.05$) are shown. Positive control splenocytes were exposed to Con A.

[0046] **FIG. 10B** exemplifies the results of an ELISpot IL-2 analysis from **FIG 10A**.

[0047] **FIG. 11** exemplifies a graph depicting the number of spot-forming cells (SFCs) from mice vaccinated with recombinant Ad5 CEA expression vectors secreting IFN- γ . The reduction in SFCs from mice vaccinated with Ad5 [E1-, E2b-]-CEA compared to the reduction in SFCs from Ad5 [E1-]-CEA vaccinated mice is shown.

[0048] **FIG. 12A** exemplifies a Kaplan-Meier survival plot of 7 patients in cohorts 1 and 2 treated with Ad5 [E1-, E2b-]-CEA(6D).

[0049] **FIG. 12B** exemplifies a Kaplan-Meier survival plot of 21 patients in cohort 3 and Ph II treated with Ad5 [E1-, E2b-]-CEA(6D).

[0050] **FIG. 12C** exemplifies a Kaplan-Meier survival plot of 6 patients in cohort 5 treated with Ad5 [E1-, E2b-]-CEA(6D).

[0051] **FIG. 12D** exemplifies a Kaplan-Meier survival plot of all 34 patients treated with Ad5 [E1-, E2b-]-CEA(6D).

[0052] **FIG. 13A** exemplifies a Kaplan-Meier survival plot of 28 cancer patients treated three times with Ad5 [E1-, E2b-]-CEA(6D).

[0053] **FIG. 13B** exemplifies a Kaplan-Meier survival plot of 27 colorectal cancer patients treated with Ad5 [E1-, E2b-]-CEA(6D).

[0054] **FIG. 14** exemplifies CEA-directed CMI responses in treated patients. CMI (IFN- γ secretion) was assessed at baseline (Pre) and after administrations of Ad5 [E1-, E2b-]-CEA(6D) (Post). The highest CMI responses (regardless of time point) observed in the patients after treatment showed a dose response. The highest CMI levels occurred in patients that received the highest dose of 5×10^{11} VP (Cohort 5). The CMI responses were significantly elevated for Cohort 3/Phase II ($p=0.0002$; *Mann-Whitney test*) and Cohort 5 ($p=0.0317$; *Mann-Whitney test*) as compared to their baseline (Pre) values. Response specificity was shown by the lack of reactivity with the irrelevant antigens

β -galactosidase and HIV-gag. Positive control PBMCs were exposed to Con A.

[0055] **FIG. 15** exemplifies CEA directed CMI responses in treated patients. CMI (IFN- γ secretion) was assessed at baseline (week 0) and 3 weeks after the last immunotherapy (week 9) for patients in all 4-dose cohorts. A dose response is shown and the highest CMI level occurred in patients that received the highest dose. The CMI response with the highest dose was significantly elevated ($P<0.02$; *Mann-Whitney test*). Response specificity was shown by the lack of reactivity with the irrelevant antigens β -galactosidase and HIV-gag. Positive control PBMCs were exposed to Con A.

[0056] **FIG. 16A** exemplifies Ad5 immune responses in patients receiving immunizations with Ad5 [E1-, E2b-]-CEA(6D) vaccine. Ad5 NAb titers to Ad5 were determined in patients at baseline (week 0) and 3 weeks (week 9) after the third immunization. The number of IFN- γ secreting PBMCs from patients specific for Ad5 was determined by ELISpot. The Ad5 NAb titers were significantly elevated at week 9 ($p<0.0001$; *Mann-Whitney test*).

[0057] **FIG. 16B** exemplifies Ad5 immune responses in patients receiving immunizations with Ad5 [E1-, E2b-]-CEA(6D) vaccine. CMI responses were determined in patients at baseline (week 0) and 3 weeks (week 9) after the third immunization. The number of IFN- γ secreting PBMCs from patients specific for Ad5 was determined by ELISpot. The Ad5 CMI responses were significantly elevated at week 9 ($P < 0.01$; *Mann-Whitney test*).

[0058] **FIG. 17A** exemplifies CEA-specific immunity in patients receiving immunizations with Ad5 [E1-, E2b-]-CEA(6D) vaccine and comparisons with Ad5 immunity. The mean CEA specific immune responses in patients ($n=19$) who received 1×10^{11} VP of Ad5 [E1-, E2b-]-CEA(6D) as measured by IFN- γ secretion of PBMC in patients with none to low pre-existing Ad5 immunity ($NAb<200$) is shown compared to the CEA specific immune response of patients with high pre-existing Ad5 immunity ($NAb\geq 200$) prior to the initiation of treatment with Ad5

[E1-, E2b-]-CEA(6D). There was no significant difference between the two groups at any time point tested ($p>0.4$, *Mann-Whitney test*)

[0059] **FIG. 17B** exemplifies a plot of the correlation between pre-existing Ad5 NAb activity and the highest levels of induced CEA CMI responses. The r^2 value (0.0155) indicates there is no correlation between pre-existing Ad5 NAb activity and CEA CMI ELISpot responses.

[0060] **FIG. 17C** exemplifies a plot of the correlation between vector induced Ad5 NAb activity and CEA CMI responses. The r^2 value (0.0069) indicates there is no correlation between vector-induced Ad5 NAb activity and CEA CMI ELISpot responses.

[0061] **FIG. 18** exemplifies a Kaplan-Meier survival plot demonstrating the effect of Ad5 [E1-, E2b-]-CEA(6D) immunotherapy in 32 metastatic colorectal cancer patients (mCRC) patients treated with Ad5 [E1-, E2b-]-CEA(6D).

[0062] **FIG. 19** exemplifies CMI responses (IFN- γ secretion) at baseline (Pre-immunized) and after administrations of Ad5 [E1-, E2b-]-CEA(6D) (Post- immunized) in mCRC patients. The highest CMI

responses (regardless of time point) observed in the patients after treatment revealed a dose response. The highest CMI levels occurred in patients receiving the highest dose (Cohort 5) and were significantly elevated ($p<0.02$; *Mann-Whitney test*). Response specificity is shown by lack of reactivity with irrelevant antigens β -galactosidase and HIV-gag. For positive controls, PBMCs were exposed to Con A.

[0063] **FIG. 20** exemplifies CMI responses (ELISpot IFN- γ SFCs) in immunized mCRC patients assessed at weeks 0, 3, 6, and 9. CMI responses increased during immunizations.

[0064] **FIG. 21** exemplifies cytotoxic T-cell (CTL) mediated cytotoxicity responses (ELISpot granzyme B secreting SFCs) assessed for pre-immunized (week 0) and post-immunized (weeks 6-9) treatments. Responses increased after immunizations ($P < 0.05$, *Wilcoxon test*).

[0065] **FIG. 22** exemplifies CMI responses in follow-up PBMC samples from 5 immunized mCRC patients as assessed by ELISpot IFN- γ SFCs. CMI responses peaked at week 9 and decreased by week 26 after treatment was stopped.

[0066] **FIG. 23** exemplifies CMI responses in mice immunized against CEA, MUC1, and Brachyury as assessed by ELISpot assays for IFN- γ secretion from splenocytes (IFN- γ SFCs). IFN- γ SFCs were detected in multi-targeted immunized mice (CEA(6D), mMUC1-C, and Brachyury) but not control mice injected with Ad5-Null (empty vector). Response specificity of the ELISpot assay was confirmed by lack of reactivity to irrelevant SIV-nef or SIV-vif peptide antigens. A positive control included cells exposed to Con A.

[0067] **FIG. 24A** exemplifies a bar graph of results from flow cytometry on polyfunctional CD8 α $+$ splenocyte cells expressing IFN- γ and TNF- α in mice immunized with CEA(6D), mMUC1-C, and Brachyury, but not in controls injected with Ad5-null. Specificity of the responses was confirmed by lack of reactivity to media alone or irrelevant SIV-nef or SIV-vif peptides.

[0068] **FIG. 24B** exemplifies a bar graph of results from flow cytometry on polyfunctional CD4 $+$ splenocyte cells expressing IFN- γ and TNF- α in mice immunized with CEA(6D), mMUC1-C, and Brachyury, but not in controls injected with Ad5-null. Specificity of the responses was confirmed by lack of reactivity to media alone or irrelevant SIV-nef or SIV-vif peptides.

[0069] **FIG. 25A** exemplifies a bar graph showing statistical significance ($p<0.0001$) of CEA antibody-dependent cellular cytotoxicity (ADCC) responses induced in mice immunized with CEA(6D), mMUC1-C, and Brachyury, but not in control mice.

[0070] **FIG. 25B** exemplifies a bar graph showing statistical significance ($p<0.0001$) of CEA complement-dependent cellular cytotoxicity (CDCC) responses induced in mice immunized with CEA(6D), mMUC1-C, and Brachyury, but not in control mice.

[0071] **FIG. 26A** exemplifies results from C57Bl/6 mice ($n=7$ /group) subcutaneously inoculated in the left flank with MC38 tumor cells expressing MUC1 tumor cells and administered 1×10^{10}

Ad5-null VPs or 1×10^{10} Ad5 [E1-, E2b-]-mMUC1-C VPs in the right flank on days 0, 7, 14. Mice treated with Ad5 [E1-, E2b-]-mMUC1-C had significantly ($p < 0.05$) smaller tumors on days 15 and 18 compared to controls and significantly longer survival. Experiments were terminated on day 36.

[0072] **FIG. 26B** exemplifies that the mice treated with Ad5 [E1-, E2b-]-mMUC1-C as described for Fig. 26A had significantly ($p < 0.05$) longer survival on days 15 and 18 compared to controls.

[0073] **FIG. 27** exemplifies results from C57Bl/6 mice subcutaneously inoculated in the left flank with MC38 tumor cells expressing Brachyury and administered 1×10^{10} Ad5-null VPs ($n=4$) or 1×10^{10} Ad5 [E1-, E2b-]-Brachyury VPs ($n=5$) in the right flank on days 5, 11, and 17. Tumors were smaller in treated mice on days 15, 19, and 22.

[0074] **FIG. 28** exemplifies the effects of HPV immunotherapy in C57Bl/6 mice ($n=7$ /group) implanted with HPV-E6/E7 expressing TC-1 tumor cells (day 0) and treated by immunotherapy on days 10, 17, 24 with 1×10^{10} Ad5-null VPs plus 100 μ g control IgG (intraperitoneal), 1×10^{10} Ad5-null VPs plus 100 μ g anti-PD1, 1×10^{10} Ad5 [E1-, E2b-]-HPV-E6/E7 VPs plus 100 μ g mouse IgG, or 1×10^{10} Ad5 [E1-, E2b-]-HPV-E6/E7 VPs plus 100 μ g anti-PD1. Immunotherapy with or without anti-PD1 resulted in significant inhibition of tumor growth by day 23 ($p < 0.05$). All control mice were terminated by day 23 due to tumor mass.

[0075] **FIG. 29** exemplifies combination multi-targeted HPV immunotherapy effects on tumor growth and regression in 2 mice treated by immunotherapy with anti-PD1 injections in **FIG. 28**. Tumor growth peaked on days 20 and 27, respectively, and then regressed thereafter.

[0076] **FIG. 30A** exemplifies expression of Brachyury protein in human dendritic cells (DCs) infected with Ad5 [E1-, E2b-]-Brachyury. SW620 tumor cells were used as positive control. Actin was used as a loading control. Expression of Brachyury was robust in DCs infected with Ad5 [E1-, E2b-]-Brachyury.

[0077] **FIG. 30B** exemplifies expression of MUC1 protein in human DCs infected with Ad5 [E1-, E2b-]-MUC1. SW620 tumor cells were used as positive control. Actin was used as a loading control. MUC1 expression was observed in human DCs infected with Ad5 [E1-, E2b-]-MUC1 vector as compared to DCs infected with Ad5 [E1-, E2b-]-null.

[0078] **FIG. 31A** exemplifies analyses of IFN- γ secreted from splenocytes from C57Bl/6 mice ($n = 5$ /group) vaccinated three times at 2-week intervals with 1×10^{10} Ad5 [E1-, E2b-]-Brachyury VPs, 1×10^{10} Ad5 [E1-, E2b-]-CEA VPs, 1×10^{10} Ad5 [E1-, E2b-]-MUC1 VPs, or Tri-Ad5 (1:1:1 mixture of 1×10^{10} Ad5 [E1-, E2b-]-Brachyury VPs, 1×10^{10} Ad5 [E1-, E2b-]-CEA VPs, and 1×10^{10} Ad5 [E1-, E2b-]-MUC1 VPs). Controls received 3×10^{10} Ad5-null VPs. Splenocytes were collected 14 days after the final vaccination and assessed for IFN- γ secretion by ELISpot assay. Positive control splenocytes were exposed to Con A. Significant differences ($p < 0.05$) between columns are reported in p-values. Not significant = ns.

[0079] **FIG. 31B** exemplifies analyses of IL-2 secreted from splenocytes from the vaccinated mice described in **FIG. 31A**. Splenocytes were assessed for IL-2 secretion by ELISpot assay.

[0080] **FIG. 32A** exemplifies a graph of analyses of CD8⁺ and multifunctional cellular populations following vaccination of C57Bl/6 mice (n=5/group) vaccinated three times at 2-week intervals with 1x10¹⁰ Ad5 [E1-, E2b-]-Brachyury VPs, 1x10¹⁰ Ad5 [E1-, E2b-]-CEA VPs, 1x10¹⁰ Ad5 [E1-, E2b-]-MUC1 VPs or Tri-Ad5 (1:1:1 mixture of 1x10¹⁰ Ad5 [E1-, E2b-]-Brachyury VPs, 1x10¹⁰ Ad5 [E1-, E2b-]-CEA VPs, and 1x10¹⁰ Ad5 [E1-, E2b-]-MUC1 VPs). Controls received 3x10¹⁰ Ad5 [E1-, E2b-]-null VPs. Splenocytes were collected 14 days after the final vaccination and were assessed by FACS for CD8⁺ cells secreting IFN- γ and TNF- α . Positive control splenocytes were exposed to Con A.

[0081] **FIG. 32B** exemplifies a graph of FACS analyses of CD4⁺ cells and multifunctional cellular populations from the vaccinated mice described in **FIG. 32A**.

[0082] **FIG. 32C** exemplifies a graph of FACS analyses of CD8 α^+ cells secreting IFN- γ and TNF- α from the vaccinated mice described in **FIG. 32A**.

[0083] **FIG. 32D** exemplifies a graph of FACS analyses of CD4⁺ cells secreting IFN- γ and TNF- α from the vaccinated mice described in **FIG. 32A**.

[0084] **FIG. 33A** exemplifies an ELISA analysis of CEA IgG levels in mice vaccinated three times with 1x10¹⁰ Ad5 [E1-, E2b-]-CEA VPs, Tri-Ad5 (1:1:1 mixture of 1x10¹⁰ Ad5 [E1-, E2b-]-Brachyury VPs, 1x10¹⁰ Ad5 [E1-, E2b-]-CEA VPs, and 1x10¹⁰ Ad5 [E1-, E2b-]-MUC1 VPs), or 3x10¹⁰ Ad5 [E1-, E2b-]-null VPs.

[0085] **FIG. 33B** exemplifies a CDC assay against MC38-CEA2 cells using the mice described in **FIG. 33A**.

[0086] **FIG. 34** exemplifies effects of immunotherapy on tumor volume in C57Bl/6 mice (n=7/group) inoculated subcutaneously in the left flank with 1x10⁶ MC-38-MUC1 cells and 1x10¹⁰ Ad5 [E1-, E2b-]-MUC1 VPs or Tri-Ad5 (1:1:1 mixture of 1x10¹⁰ Ad5 [E1-, E2b-]-CEA VPs, 1x10¹⁰ Ad5 [E1-, E2b-]-MUC1 VPs, and 1x10¹⁰ Ad5 [E1-, E2b-]-Brachyury VPs (3x10¹⁰ VP total)). Control mice received 3x10¹⁰ Ad5 [E1-, E2b-]-null VPs. (*) indicates days when Ad5 [E1-, E2b-]-MUC1 treated mice had significantly smaller (p<0.05) tumors than control mice and (^) indicates days when Tri-Ad5-treated mice had significantly smaller (p<0.05) tumors than control mice. No significant difference (p>0.1) between Ad5 [E1-, E2b-]-MUC1 and Tri-Ad5-treated mice was seen.

[0087] **FIG. 35** exemplifies an immunoblot showing expression of CEA in A549 cells infected with Ad5 [E1-, E2b-]-CEA. (A) Negative Control. (B) Protein molecular weight marker. (C) Negative. (D) CEA Reference Material (30 ng). (E) Ad5 [E1-, E2b-]-CEA lysate (20 μ L). (F) Ad5 [E1-, E2b-]-CEA lysate (20 μ L). (G) Negative A549 cells. Recombinant CEA was used as a positive control and uninfected A549 cells served as a negative control.

[0088] **FIG. 36** exemplifies CMI dose responses as measured by ELISpot of splenocytes from C57BL/6 mice (n=5/group) immunized three times at 14-day intervals with doses of 1×10^8 , 1×10^9 or 1×10^{10} Ad5 [E1-, E2b-]-E6/E7 VPs and assessed 14 days after the final immunization. The greatest induction of CMI was achieved with the 1×10^{10} VP dose. Positive control splenocytes were exposed to Con A.

[0089] **FIG. 37A** exemplifies activation of CD8- α^+ /IFN- γ^+ splenocytes after immunization of C57BL/6 mice (n=5/group) immunized three times at two week intervals with 1×10^{10} VP Ad5 [E1-, E2b-]-E6/E7 VPs. Controls received 1×10^{10} Ad5 [E1-, E2b-]-null VPs. Splenocytes collected 14 days after the final immunization were assessed by flow cytometry for. For positive controls, splenocytes were exposed to PMA/ionomycin.

[0090] **FIG. 37B** exemplifies activation of CD8- α^+ /IFN- γ^+ /TNF- α^+ splenocytes after immunization of mice as described in **FIG. 37A**.

[0091] **FIG. 38A** exemplifies changes in tumor size from immunotherapy of C57BL/6 mice (n=5/group) implanted on day 0 with 2×10^5 non-palpable HPV-E6/E7 TC-1 tumor cells and administered 1×10^{10} Ad5 [E1-, E2b-]-null VPs or 1×10^{10} Ad5 [E1-, E2b-]-E6/E7 VPs on days 1, 8 and 15. Tumor size was determined and volumes calculated according to the formula $V = (a^2 \times b)/2$. Analysis of significance was performed between experimental and vector control groups using unpaired t-tests and significance is denoted by * (p<0.05) and ** (p<0.01).

[0092] **FIG. 38B** exemplifies a survival curve of the mice as described in **FIG. 38A** that was plotted and compared using the Mantel-Cox test. Significance is denoted by ** (p<0.01).

[0093] **FIG. 39A** exemplifies changes in tumor size from immunotherapy of C57BL/6 mice (n=4/group) implanted on day 0 with 2×10^5 small palpable HPV-E6/E7 TC-1 tumor cells and administered 1×10^{10} Ad5 [E1-, E2b-]-null VPs or 1×10^{10} Ad5 [E1-, E2b-]-E6/E7 VPs on days 6, 13 and 20. Tumor size was determined and volumes calculated according to the formula $V = (a^2 \times b)/2$. Analysis of significance was performed between experimental and vector control groups using unpaired t-tests and significance is denoted by ** (p<0.01).

[0094] **FIG. 39B** exemplifies a survival curve of the mice as described in **FIG. 39A** that was plotted and compared using the Mantel-Cox test. Significance is denoted by ** (p<0.01).

[0095] **FIG. 40A** exemplifies changes in tumor size from immunotherapy of C57BL/6 mice (n=4/group) implanted on day 0 with 2×10^5 large established HPV-E6/E7 TC-1 tumor cells and administered 1×10^{10} Ad5 [E1-, E2b-]-null VPs or 1×10^{10} Ad5 [E1-, E2b-]-E6/E7 VPs on days 13, 20 and 27. Tumor size was determined and volumes calculated according to the formula $V = (a^2 \times b)/2$. Analysis of significance was performed between experimental and vector control groups using unpaired t-tests and significance is denoted by ** (p<0.01).

[0096] **FIG. 40B** exemplifies a survival curve of the mice as described in **FIG. 40A** that was plotted and compared using the Mantel-Cox test. Significance is denoted by ** (p<0.01).

[0097] **FIG. 41A** exemplifies changes in tumor size from C57BL/6 mice (n=7/group) inoculated on day 0 with 2×10^5 TC-1 tumor cells and administered treatments on days 10, 17, and 24 with 1×10^{10} Ad5 [E1-, E2b-]-null VPs plus 100 μ g of isotype control rat IgG. Tumor size was determined and volumes calculated according to the formula $V = (a^2 \times b)/2$. Tumor growth kinetics represents individual mice in each group.

[0098] **FIG. 41B** exemplifies changes in tumor size from C57BL/6 mice (n=7/group) inoculated on day 0 with 2×10^5 TC-1 tumor cells and administered treatments on days 10, 17, and 24 with 1×10^{10} Ad5 [E1-, E2b-]-null VPs plus 100 μ g anti-PD1 antibody.

[0099] **FIG. 41C** exemplifies changes in tumor size from C57BL/6 mice (n=7/group) inoculated on day 0 with 2×10^5 TC-1 tumor cells and administered treatments on days 10, 17, and 24 with 1×10^{10} Ad5 [E1-, E2b-]-E6/E7 VPs plus 100 μ g isotype control rat IgG.

[0100] **FIG. 41D** exemplifies changes in tumor size from C57BL/6 mice (n=7/group) inoculated on day 0 with 2×10^5 TC-1 tumor cells and administered treatments on days 10, 17, and 24 with 1×10^{10} Ad5 [E1-, E2b-]-E6/E7 VPs plus 100 μ g anti-PD1.

[00101] **FIG. 42** exemplifies a survival curve for C57BL/6 mice (n=7/group) treated as those in **FIGs. 41A-D**. The experiment was terminated on day 52 following tumor implantation. Mice treated with Ad5 [E1-, E2b-]-E6/E7 and control antibody exhibited significantly ($p < 0.008$) longer survival compared to both groups of control mice (Ad5 [E1-, E2b-]-null and control antibody or Ad5 [E1-, E2b-]-null and anti-PD1 antibody). 2 of 7 (29%) Ad5 [E1-, E2b-]-E6/E7 and control antibody treated mice remained alive at day 52. Mice treated with Ad5 [E1-, E2b-]-E6/E7 plus anti PD1 antibody exhibited significantly ($p < 0.0006$) longer survival as compared to both groups of controls. 4 of 7 (57%) Ad5 [E1-, E2b-]-E6/E7 plus anti-PD1 antibody treated mice remained alive at day 52.

[00102] **FIG. 43A** exemplifies that Ad5 [E1-, E2b-]-E6/E7 promotes the recruitment of CD8+ tumor-infiltrating lymphocytes (TILs) into TC-1 tumors. C57BL/6 mice (n=5/group) were implanted with 2×10^5 TC-1 tumor cells. Twelve days after implantation mice began treatment with Ad5 [E1-, E2b-]-null empty vector plus control IgG, Ad5 [E1-, E2b-]-null plus anti-PD1, Ad5 [E1-, E2b-]-E6/E7 plus control IgG, or Ad5 [E1-, E2b-]-E6/E7 plus anti-P-1. Vaccine was administered subcutaneously weekly and anti-PD1 antibodies were administered via intraperitoneal injection every 3-4 days and tumors were analyzed on day 27. Ad5 [E1-, E2b-]-E6/E7 treatment significantly decreases the ratio of Treg/CD8+ TILs. Analysis of significance was performed using unpaired t-tests and significance is denoted by ns ($p > 0.05$), * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$), or **** ($p < 0.0001$).

[00103] **FIG. 43B** exemplifies that the reduction in the ratio of Treg/CD8+ TILs of **FIG. 43A** reduction is not driven by a reduction in the number of Tregs.

[00104] **FIG. 43C** exemplifies that the reduction in the ratio of Treg/CD8+ TILs of **FIG. 43A** is driven through an increase in the number of CD8+ TILs.

[00105] **FIG.44A** exemplifies that Ad5 [E1-, E2b-]-E6/E7 plus anti-PD1 antibody combination therapy promotes a pro-inflammatory tumor microenvironment. C57BL/6 mice (n=5/group) were tumor implanted, treated, and tumors were analyzed as in **FIGs. 43A-C**. The frequency of PD1⁺ CD4⁺ and CD8⁺ TILs is increased in tumors from mice treated with Ad5 [E1-, E2b-]-E6/E7. Tumors from mice treated with a combination of Ad5 [E1-, E2b-]-E6/E7 and anti-PD1 have a significantly lower frequency of PD1⁺ CD4⁺ and CD8⁺ TILs (A), LAG-3⁺ CD8⁺ TILs (B), and (C). Analysis of significance was performed using unpaired t-tests and significance is denoted by ns (p>0.05), * (p<0.05), ** (p<0.01), or *** (p<0.001).

[00106] **FIG.44B** exemplifies that tumors from mice treated with a combination of Ad5 [E1-, E2b-]-E6/E7 and anti-PD1 as in **FIG. 44A** have a significantly lower frequency of LAG-3⁺ CD8⁺ TILs bringing these levels more in line with tumors from control mice.

[00107] **FIG.44C** exemplifies that tumors from mice treated with a combination of Ad5 [E1-, E2b-]-E6/E7 and anti-PD1 as in **FIG. 44A** have a significantly reduced expression level of PDL1.

DETAILED DESCRIPTION OF THE INVENTION

[00108] The following passages describe different aspects of the invention in greater detail. Each aspect of the invention may be combined with any other aspect or aspects of the invention unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous.

[00109] Unless otherwise indicated, any embodiment can be combined with any other embodiment. A variety of aspects of this invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range as if explicitly written out. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range. When ranges are present, the ranges include the range endpoints.

[00110] Compared to first generation adenovirus vectors, certain embodiments of the Second Generation E2b deleted adenovirus vectors of the present invention contain additional deletions in the DNA polymerase gene (pol) and deletions of the pre-terminal protein (pTP). E2b deleted vectors have up to a 13 kb gene-carrying capacity as compared to the 5 to 6 kb capacity of First Generation adenovirus vectors, easily providing space for nucleic acid sequences encoding any of a variety of target antigens. The E2b deleted adenovirus vectors also have reduced adverse reactions as compared to first generation adenovirus vectors.

[00111] The innate immune response to wild type Ad can be complex, and it appears that Ad proteins expressed from adenovirus vectors play an important role. Specifically, the deletions of pre-terminal protein and DNA polymerase in the E2b deleted vectors appear to reduce inflammation during the first 24 to 72 h following injection, whereas First Generation adenovirus vectors stimulate inflammation during this period. In addition, it has been reported that the additional replication block created by E2b deletion also leads to a 10,000 fold reduction in expression of Ad late genes, well beyond that afforded by E1, E3 deletions alone. The decreased levels of Ad proteins produced by E2b deleted adenovirus vectors effectively reduce the potential for competitive, undesired, immune responses to Ad antigens, responses that prevent repeated use of the platform in Ad immunized or exposed individuals. The reduced induction of inflammatory response by second generation E2b deleted vectors results in increased potential for the vectors to express desired vaccine antigens during the infection of antigen presenting cells (i.e., dendritic cells), decreasing the potential for antigenic competition, resulting in greater immunization of the vaccine to the desired antigen relative to identical attempts with First Generation adenovirus vectors. E2b deleted adenovirus vectors provide an improved Ad-based vaccine candidate that is safer, more effective, and more versatile than previously described vaccine candidates using First Generation adenovirus vectors. Thus, first generation, E1-deleted Adenovirus subtype 5 (Ad5)-based vectors, although promising platforms for use as cancer vaccines, are impeded in activity by naturally occurring or induced Ad-specific neutralizing antibodies. Without being bound by theory, Ad5-based vectors with deletions of the E1 and the E2b regions (Ad5 [E1-, E2b-]), the latter encoding the DNA polymerase and the pre-terminal protein, for example by virtue of diminished late phase viral protein expression, may avoid immunological clearance and induce more potent immune responses against the encoded tumor antigen transgene in Ad-immune hosts.

[00112] The present invention relates to methods and compositions (e.g., viral vectors) for generating immune responses against target antigens, in particular, those associated or related to infectious disease or proliferative cell disease such as cancer. The present invention relates to methods and compositions for generating immune responses in an individual against target antigens, in particular, those related to cell proliferation diseases such as cancer. In various aspects of the invention, compositions and methods described herein relate to generating an immune response in an individual against cells expressing and/or presenting a target antigen or a target antigen signature comprising at least one target antigen. The present invention provides compositions and methods for immunotherapy against human papilloma virus (HPV) using a viral gene delivery platform to immunize against HPV genes E6 and E7 combined with PD1 checkpoint blockade. These compositions and methods utilize an Ad5 [E1-, E2b-]-E6/E7 vaccine combined with an immune pathway checkpoint modulator.

[00113] The compositions and methods can be used to generate an immune response against a target antigen expressed and/or presented by a cell. For example, the compositions and methods can be used to

generate immune responses against a carcinoembryonic antigen (CEA), such as CEA expressed or presented by a cell. For example, the compositions and methods can be used to generate an immune response against CEA(6D) expressed or presented by a cell. For example, the compositions and methods can be used to generate an immune response against Mucin 1 (MUC1) expressed and/or presented by a cell. For example, the compositions and methods can be used to generate an immune response against MUC1c expressed and/or presented by a cell. For example, the compositions and methods can be used to generate an immune response against Brachyury (T protein (T)) expressed and/or presented by a cell.

[00114] The compositions and methods can be used to generate an immune response against multiple target antigens expressed and/or presented by a cell. For example, the compositions and methods can be used to generate an immune response against MUC1c, T, or any combination thereof. For example, the compositions and methods can be used to generate an immune response against T and CEA. For example, the compositions and methods can be used to generate an immune response against MUC1c and CEA. For example, the compositions and methods can be used to generate an immune response against MUC1 and T. For example, the compositions and methods can be used to generate an immune response against MUC1c, T, and CEA.

[00115] A modified form of CEA, MUC1c, or T can be used in a vaccine directed to raising an immune response against CEA, MUC1c, or T, or cells expressing and/or presenting CEA, MUC1c, or T. In particular, the present invention provides an improved Ad-based vaccine such that multiple vaccinations against one or more antigenic target entity can be achieved. In some embodiments, the improved Ad-based vaccine comprises a replication defective adenovirus carrying a target antigen, a fragment, a variant or a variant fragment thereof, such as Ad5 [E1-,E2b-]-CEA(6D). Variants or fragments of target antigens, such as CEA, MUC1c, or T, can be selected based on a variety of factors, including immunogenic potential. A mutant CEA, CEA(6D) can utilized for its increased capability to raise an immune response relative to the CEA(WT). Importantly, vaccination can be performed in the presence of preexisting immunity to the Ad or administered to subjects previously immunized multiple times with the Ad vector of the present invention or other Ad vectors. The Ad vectors can be administered to subjects multiple times to induce an immune response against an antigen of interest, such as CEA, MUC1c, or T, including but not limited to, the production of antibodies and CMI responses against one or more target antigens.

[00116] The following passages describe different aspects of the invention in greater detail. Each aspect of the invention may be combined with any other aspect or aspects of the invention unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous. As used herein, unless otherwise indicated, the article "a" means one or more unless explicitly otherwise provided for. As used herein, unless otherwise indicated, terms such as "contain," "containing," "include," "including,"

and the like mean “comprising.” As used herein, unless otherwise indicated, the term “or” can be conjunctive or disjunctive. As used herein, unless otherwise indicated, any embodiment can be combined with any other embodiment.

[00117] An “adenovirus” (Ad) refers to non-enveloped DNA viruses from the family Adenoviridae. These viruses can be found in, but are not limited to, human, avian, bovine, porcine and canine species. The present invention contemplates the use of any Ad from any of the four genera of the family Adenoviridae (e.g., Aviadenovirus, Mastadenovirus, Atadenovirus and Siadenovirus) as the basis of an E2b deleted virus vector, or vector containing other deletions as described herein. In addition, several serotypes are found in each species. Ad also pertains to genetic derivatives of any of these viral serotypes, including but not limited to, genetic mutations, deletions or transpositions.

[00118] A “helper adenovirus” or “helper virus” refers to an Ad that can supply viral functions that a particular host cell cannot (the host may provide Ad gene products such as E1 proteins). This virus is used to supply, in trans, functions (e.g., proteins) that are lacking in a second virus, or helper dependent virus (e.g., a gutted or gutless virus, or a virus deleted for a particular region such as E2b or other region as described herein); the first replication-incompetent virus is said to “help” the second, helper dependent virus thereby permitting the production of the second viral genome in a cell.

[00119] An “adenovirus 5 null (Ad5-null)” refers to a non-replicating Ad that does not contain any heterologous nucleic acid sequences for expression.

[00120] A “first generation adenovirus” refers to an Ad that has the early region 1 (E1) deleted. In additional cases, the early region 3 (E3) may also be deleted.

[00121] “Gutted” or “gutless” refers to an Ad vector that has been deleted of all viral coding regions.

[00122] “Transfection” refers to the introduction of foreign nucleic acid into eukaryotic cells.

Exemplary means of transfection include calcium phosphate-DNA co-precipitation, DEAE-dextran-mediated transfection, polybrene-mediated transfection, electroporation, microinjection, liposome fusion, lipofection, protoplast fusion, retroviral infection, and biolistics.

[00123] “Stable transfection” or “stably transfected” refers to the introduction and integration of foreign nucleic acid, DNA or RNA, into the genome of the transfected cell. The term “stable transfectant” refers to a cell which has stably integrated foreign DNA into the genomic DNA.

[00124] A “reporter gene” indicates a nucleotide sequence that encodes a reporter molecule (e.g., an enzyme). A “reporter molecule” is detectable in any of a variety of detection systems, including, but not limited to, enzyme-based detection assays (e.g., ELISA, histochemical assays), fluorescent, radioactive, and luminescent systems. The *E. coli* β -galactosidase gene, green fluorescent protein (GFP), the human placental alkaline phosphatase gene, the chloramphenicol acetyltransferase (CAT) gene; and other reporter genes may be employed.

[00125] A “heterologous sequence” refers to a nucleotide sequence that is ligated to, or is manipulated to become ligated to, a nucleic acid sequence to which it is not ligated in nature, or to which it is ligated at a different location in nature. Heterologous nucleic acid may include a naturally occurring nucleotide sequence or some modification relative to the naturally occurring sequence.

[00126] A “transgene” refers to any gene coding region, either natural or heterologous nucleic acid sequences or fused homologous or heterologous nucleic acid sequences, introduced into cells or a genome of subject. Transgenes may be carried on any viral vector used to introduce transgenes to the cells of the subject.

[00127] A “second generation adenovirus” refers to an Ad that has all or parts of the E1, E2, E3, and, in certain embodiments, E4 DNA gene sequences deleted (removed) from the virus.

[00128] A “subject” refers to any animal, including, but not limited to, humans, non-human primates (e.g., rhesus or other types of macaques), mice, pigs, horses, donkeys, cows, sheep, rats and fowls.

[00129] An “immunogenic fragment” refers to a fragment of a polypeptide that is specifically recognized (i.e., specifically bound) by a B-cell and/or T-cell surface antigen receptor resulting in a generation of an immune response specifically against a fragment.

[00130] A “target antigen” or “target protein” refers to a molecule, such as a protein, against which an immune response is to be directed.

[00131] “E2b deleted” refers to a DNA sequence mutated in such a way so as to prevent expression and/or function of at least one E2b gene product. Thus, in certain embodiments, “E2b deleted” is used in relation to a specific DNA sequence that is deleted (removed) from an Ad genome. E2b deleted or “containing a deletion within an E2b region” refers to a deletion of at least one base pair within an E2b region of an Ad genome. Thus, in certain embodiments, more than one base pair is deleted and in further embodiments, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 base pairs are deleted. In another embodiment, a deletion is of more than 150, 160, 170, 180, 190, 200, 250, or 300 base pairs within an E2b region of an Ad genome. An E2b deletion may be a deletion that prevents expression and/or function of at least one E2b gene product and therefore, encompasses deletions within exons of encoding portions of E2b-specific proteins as well as deletions within promoter and leader sequences. In certain embodiments, an E2b deletion is a deletion that prevents expression and/or function of one or both a DNA polymerase and a preterminal protein of an E2b region. In a further embodiment, “E2b deleted” refers to one or more point mutations in a DNA sequence of this region of an Ad genome such that one or more encoded proteins is non-functional. Such mutations include residues that are replaced with a different residue leading to a change in an amino acid sequence that result in a nonfunctional protein.

[00132] “E1-deleted” refers to a DNA sequence that is mutated in such a way so as to prevent expression and/or function of at least one E1 gene product. Thus, in certain embodiments, “E1 deleted” is

used in relation to a specific DNA sequence that is deleted (removed) from the Ad genome. E1 deleted or “containing a deletion within the E1 region” refers to a deletion of at least one base pair within the E1 region of the Ad genome. Thus, in certain embodiments, more than one base pair is deleted and in further embodiments, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 base pairs are deleted. In another embodiment, the deletion is of more than 150, 160, 170, 180, 190, 200, 250, or 300 base pairs within the E1 region of the Ad genome. An E1 deletion may be a deletion that prevents expression and/or function of at least one E1 gene product and therefore, encompasses deletions within exons of encoding portions of E1-specific proteins as well as deletions within promoter and leader sequences. In certain embodiments, an E1 deletion is a deletion that prevents expression and/or function of one or both of a trans-acting transcriptional regulatory factor of the E1 region. In a further embodiment, “E1 deleted” refers to one or more point mutations in the DNA sequence of this region of an Ad genome such that one or more encoded proteins is non-functional. Such mutations include residues that are replaced with a different residue leading to a change in the amino acid sequence that result in a nonfunctional protein.

[00133] “Generating an immune response” or “inducing an immune response” refers to a statistically significant change, e.g., increase or decrease, in the number of one or more immune cells (T-cells, B-cells, antigen-presenting cells, dendritic cells, neutrophils, and the like) or in the activity of one or more of these immune cells (CTL activity, HTL activity, cytokine secretion, change in profile of cytokine secretion, etc.).

[00134] The terms “nucleic acid” and “polynucleotide” are used essentially interchangeably herein. Polynucleotides of the invention may be single-stranded (coding or antisense) or double-stranded, and may be DNA (e.g. genomic, cDNA, or synthetic) or RNA molecules. RNA molecules may include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials. An isolated polynucleotide, as used herein, means that a polynucleotide is substantially away from other coding sequences. For example, an isolated DNA molecule as used herein does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. This refers to the DNA molecule as originally isolated, and does not exclude genes or coding regions later added to the segment recombinantly in the laboratory.

[00135] As will be understood by those skilled in the art, the polynucleotides of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express target antigens as described herein, fragments of

antigens, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

[00136] Typically, polynucleotide variants will contain one or more substitutions, additions, deletions and/or insertions, preferably such that the immunogenicity of the epitope of the polypeptide encoded by the variant polynucleotide or such that the immunogenicity of the heterologous target protein is not substantially diminished relative to a polypeptide encoded by the native polynucleotide sequence. In some cases, the one or more substitutions, additions, deletions and/or insertions may result in an increased immunogenicity of the epitope of the polypeptide encoded by the variant polynucleotide. As described elsewhere herein, the polynucleotide variants can encode a variant of the target antigen, or a fragment (e.g., an epitope) thereof wherein the propensity of the variant polypeptide or fragment (e.g., epitope) thereof to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished relative to the native polypeptide. The polynucleotide variants can encode a variant of the target antigen, or a fragment thereof wherein the propensity of the variant polypeptide or fragment thereof to react with antigen-specific antisera and/or T-cell lines or clones is substantially increased relative to the native polypeptide.

[00137] The term “variants” should also be understood to encompass homologous genes of xenogenic origin. In particular embodiments, variants or fragments of target antigens are modified such that they have one or more reduced biological activities. For example, an oncogenic protein target antigen may be modified to reduce or eliminate the oncogenic activity of the protein, or a viral protein may be modified to reduce or eliminate one or more activities of the viral protein. An example of a modified CEA protein is a CEA having a N610D mutation, resulting in a variant protein with increased immunogenicity.

[00138] When comparing polynucleotide sequences, two sequences are “identical” if the sequence of nucleotides in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A “comparison window” as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software using default parameters. Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman, *Add. APL. Math* 2:482 (1981), by the identity alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity methods of Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA), or by inspection. One example of algorithms that are suitable for determining percent sequence identity and

sequence similarity are the BLAST and BLAST 2.0 algorithms. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program uses as defaults a word length (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

[00139] The “percentage of sequence identity” can be determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence and multiplying the results by 100 to yield the percentage of sequence identity.

[00140] It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a particular antigen of interest, or fragment thereof, as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

COMPOSITIONS

Viral Vectors for Immunotherapies and Vaccines

[00141] Recombinant viral vectors can be used to express protein coding genes or antigens (e.g., TAAs and/or IDAAs). The advantages of recombinant viral vector based vaccines and immunotherapy include high efficiency gene transduction, highly specific delivery of genes to target cells, induction of robust immune responses, and increased cellular immunity. The present disclosure provides for recombinant adenovirus vectors comprising deletions or insertions of crucial regions of the viral genome. The viral vectors of provided by the present disclosure can comprise heterologous nucleic acid sequences that encode one or more target antigens of interest, or variants, fragments or fusions thereof, against which it is desired to generate an immune response.

[00142] Suitable viral vectors that can be used with the methods and compositions of the present disclosure include but are not limited to retroviruses, lentiviruses, provirus, Vaccinia virus, adenoviruses, adeno-associated viruses, self-complementary adeno-associated virus, Cytomegalovirus, or Sendai virus. In some embodiments, the viral vector can be replication-competent. In some embodiments, the viral vector can be replication-defective. For replication-defective viral vectors, the viruses' genome can have the coding regions necessary for additional rounds of replication and packaging replaced with other genes, or deleted. These viruses are capable of infecting their target cells and delivering their viral payload, but then fail to continue the typical lytic pathway that leads to cell lysis and death. Depending on the viral vector, the typical maximum length of an allowable DNA or cDNA insert in a replication-defective viral vector is can be about 8-10 kilobases (kB).

[00143] Retroviruses have been used to express antigens, such as an enveloped, single-stranded RNA virus that contains reverse transcriptase. Retrovirus vectors can be replication-defective. Retrovirus vectors can be of murine or avian origin. Retrovirus vectors can be from Moloney murine leukemia virus (MoMLV). Retrovirus vectors can be used that require genome integration for gene expression. Retrovirus vectors can be used to provide long-term gene expression. For example, retrovirus vectors can have a genome size of approximately 7–11 kb and the vector can harbor 7–8 kb long foreign DNA inserts. Retrovirus vectors can be used to display low immunogenicity and most patients do not show pre-existing immunity to retroviral vectors. Retrovirus vectors can be used to infect dividing cells. Retrovirus vectors can be used to not infect non-dividing cells.

[00144] Lentivirus vectors have been used to express antigens. Lentiviruses constitute a subclass of retroviruses. Lentivirus vectors can be used to infect non-dividing cells. Lentivirus vectors can be used to infect dividing cells. Lentivirus vectors can be used to infect both non-dividing and dividing cells. Lentiviruses generally exhibit broader tropism than retroviruses. Several proteins such as tat and rev regulate the replication of lentiviruses. These regulatory proteins are typically absent in retroviruses. HIV is an exemplary lentivirus that can be engineered into a transgene delivery vector. The advantages of

lentivirus vectors are similar to those of retroviral vectors. Although lentiviruses can potentially trigger tumorigenesis, the risk is lower than that of retroviral vectors, as the integration sites of lentiviruses are away from the sites harboring cellular promoters. HIV-based vectors can be generated, for example, by deleting the HIV viral envelope and some of the regulatory genes not required during vector production. Instead of parental envelope, several chimeric or modified envelope vectors are generated because it determines the cell and tissue specificity.

[00145] Cytomegalovirus (CMV) vectors have been used to express antigens and is a member of the herpesviruses. Species-specific CMVs can be used (e.g., human CMV (HCMV), e.g., human herpesvirus type 5. HCMV contains a 235-kb double-stranded linear DNA genome surrounded by a capsid. The envelope contains glycoproteins gB and gH, which bind to cellular receptors.

[00146] Sendai virus (SeV) vectors have been used to express antigens. SeV is an enveloped, single-stranded RNA virus of the family Paramyxovirus. The SeV genome encodes six protein and two envelope glycoproteins, HN and F proteins, that mediate cell entry and determine its tropism. SeV vectors that lack F protein can be used as a replication-defective virus to improve the safety of the vector. SeV vector produced in a packaging cell can be used to express the F protein. An F gene-deleted and transgene-inserted genome can be transfected into a packaging cell. SeV contains RNA dependent RNA polymerase and viral genome localizes to the cytoplasm. This ensures that fast gene expression occurs soon after infection and the genotoxic advantage of SeV. SeV vectors can be used to exhibit highly efficient gene transfer. SeV vectors can be used to transduce both dividing and non-dividing cells. SeV vectors can be used to transduce non-dividing cells. SeV vectors can be used to transduce dividing cells. SeV vectors can be used, for example, to efficiently transduce human airway epithelial cells. SeV vectors can be, for example, administered by a mucosal (e.g., oral and nasal) route. Intranasal administration can be used to potentially reduce the influence of a pre-existing immunity to SeV, as compared to intramuscular administration. Compared to other viral vectors, its transgene capacity (3.4 kb) is low. SeV is highly homologous to the human parainfluenza type 1 (hPIV-1) virus; thus, a pre-existing immunity against hPIV-1 can work against the use of SeV.

[00147] Human papillomavirus (HPV) vectors can be used to express antigens. For example, by modifying oncogenes in the genome, such as by deletion or insertion of crucial regions of the HPV viral genome, a recombinant vector can be engineered to increase predictability of infection and reduce unwanted side effects. An exemplary HPV vector is a fusion vector with an adenovirus vector. An exemplary HPV vector is Ad5 [E1-, E2b-]-HPV-E6/E7 viral vector comprising a modified non-oncogenic and fused HPV-E6/E7.

Adenovirus Vectors

[00148] In general, adenoviruses are attractive for clinical because they can have a broad tropism, they can infect a variety of dividing and non-dividing cell types and they can be used systemically as well as

through more selective mucosal surfaces in a mammalian body. In addition, their relative thermostability further facilitates their clinical use. Adenoviruses are a family of DNA viruses characterized by an icosahedral, non-enveloped capsid containing a linear double-stranded genome. Generally, adenoviruses are found as non-enveloped viruses comprising double-stranded DNA genome approximated ~30–35 kilobases in size. Of the human Ads, none are associated with any neoplastic disease, and only cause relatively mild, self-limiting illness in immunocompetent individuals. The first genes expressed by the virus are the E1 genes, which act to initiate high-level gene expression from the other Ad5 gene promoters present in the wild type genome. Viral DNA replication and assembly of progeny virions occur within the nucleus of infected cells, and the entire life cycle takes about 36 hr with an output of approximately 10^4 virions per cell. The wild type Ad5 genome is approximately 36 kb, and encodes genes that are divided into early and late viral functions, depending on whether they are expressed before or after DNA replication. The early/late delineation is nearly absolute, since it has been demonstrated that super-infection of cells previously infected with an Ad5 results in lack of late gene expression from the super-infecting virus until after it has replicated its own genome. Without bound by theory, this is likely due to a replication dependent *cis*-activation of the Ad5 major late promoter (MLP), preventing late gene expression (primarily the Ad5 capsid proteins) until replicated genomes are present to be encapsulated. The composition and methods of the invention take advantage of feature in the development of advanced generation Ad vectors/vaccines. The linear genome of the adenovirus is generally flanked by two origins for DNA replication (ITRs) and has eight units for RNA polymerase II-mediated transcription. The genome carries five early units E1A, E1B, E2, E3, E4, and E5, two units that are expressed with a delay after initiation of viral replication (IX and IVa2), and one late unit (L) that is subdivided into L1–L5. Some adenoviruses can further encode one or two species of RNA called virus-associated (VA) RNA.

[00149] Adenoviruses that induce innate and adaptive immune responses in human patient are provided. By deletion or insertion of crucial regions of the viral genome, recombinant vectors are provided that have been engineered to increase their predictability and reduce unwanted side effects. In some aspects, the invention provides for an adenovirus vector comprising the genome deletion or insertion selected from the group consisting of: E1A, E1B, E2, E3, E4, E5, IX, IVa2, L1, L2, L3, L4, and L5, and any combination thereof.

[00150] The present disclosure provides for recombinant adenovirus vectors comprising an altered capsid. Generally, the capsid of an adenovirus is primarily comprises 20 triangular facets of an icosahedron each icosahedron contains 12 copies of hexon trimers. In addition there are also other several additional minor capsid proteins, IIIa, VI, VIII, and IX.

[00151] The present disclosure provides for recombinant adenovirus vectors comprising one or more altered fiber proteins. In general the fiber proteins, which also form trimers, are inserted at the 12 vertices into the pentameric penton bases. The fiber can comprise of a thin N-terminal tail, a shaft, and a knob

domain. The shaft can comprise a variable numbers of β -strand repeats. The knob can comprise one or more loops A, B, C, D, E, F, G, H, I, J. The fiber knob loops can bind to cellular receptors. The present disclosure provides for adenovirus vectors to be used in vaccine systems for the treatment of cancers and infectious diseases.

[00152] Suitable adenoviruses that can be used with the present methods and compositions of the disclosure include but are not limited to species-specific adenovirus including human subgroups A, B1, B2, C, D, E and F or their crucial genomic regions as provided herein, which subgroups can further classified into immunologically distinct serotypes. Further, suitable adenoviruses that can be used with the present methods and compositions of the disclosure include, but are not limited to, species-specific adenovirus or their crucial genomic regions identified from primates, bovines, fowls, reptiles, or frogs.

[00153] Some adenoviruses serotypes preferentially target distinct organs. Serotypes such as AdHu1, AdHu2, and AdHu5 (subgenus C), generally effect the upper respiratory, while subgenera A and F effect gastrointestinal organs. The present disclosure provides for recombinant adenovirus vectors to be used in preferentially target distinct organs for the treatment of organ-specific cancers or organ-specific infectious diseases. In some applications the recombinant adenovirus vector is altered to reduce tropism to a specific organ in a mammal. In some applications the recombinant adenovirus vector is altered to increase tropism to a specific organ in a mammal.

[00154] The tropism of an adenovirus can be determined by their ability to attach to host cell receptors. In some instances the process of host cell attachment can involve the initial binding of the distal knob domain of the fiber to a host cell surface molecule followed by binding of the RGD motif within the penton base with α V integrins. The present disclosure provides for recombinant adenovirus vectors with altered tropism such that they can be genetic engineered to infect specific cell types of a host. The present disclosure provides for recombinant adenovirus vectors with altered tropism for the treatment of cell-specific cancers or cell-specific infectious diseases. The present disclosure provides for recombinant adenovirus vectors with altered fiber knob from one or more adenoviruses of subgroups A, B, C, D, or F, or a combination thereof or the insertion of RGD sequences. In some applications the recombinant adenovirus vectors comprising an altered fiber knob results in a vector with reduced tropism for one or more particular cell types. In some applications the recombinant adenovirus vectors comprising an altered fiber knob results in a vector with enhanced tropism for one or more particular cell types. In some applications the recombinant adenovirus vectors comprising an altered fiber knob results in a vector with reduced product-specific B or T-cell responses. In some applications the recombinant adenovirus vectors comprising an altered fiber knob results in a vector with enhanced product-specific B or T-cell responses.

[00155] The present disclosure provides for recombinant adenovirus vectors that are coated with other molecules to circumvent the effects of virus-neutralizing antibodies or improve transduction in to a host cell. The present disclosure provides for recombinant adenovirus vectors that are coated with an adaptor

molecule that aids in the attachment of the vector to a host cell receptor. By way of example an adenovirus vector can be coated with adaptor molecule that connects coxsackie Ad receptor (CAR) with CD40L resulting in increased transduction of dendritic cells, thereby enhancing immune responses in a subject. Other adenovirus vectors similarly engineered for enhancing the attachment to other target cell types are also included by the present disclosure.

Ad5 vectors

[00156] Studies in humans and animals have demonstrated that pre-existing immunity against Ad5 can be an inhibitory factor to commercial use of Ad-based vaccines. The preponderance of humans have antibody against Ad5, the most widely used subtype for human vaccines, with two-thirds of humans studied having lympho-proliferative responses against Ad5. This pre-existing immunity can inhibit immunization or re-immunization using typical Ad5 vaccines and may preclude the immunization of a vaccine against a second antigen, using an Ad5 vector, at a later time. Overcoming the problem of pre-existing anti-vector immunity has been a subject of intense investigation. Investigations using alternative human (non-Ad5 based) Ad5 subtypes or even non-human forms of Ad5 have been examined. Even if these approaches succeed in an initial immunization, subsequent vaccinations may be problematic due to immune responses to the novel Ad5 subtype. To avoid the Ad5 immunization barrier, and improve upon the limited efficacy of first generation Ad5 [E1-] vectors to induce optimal immune responses, various embodiments of the invention relate to a next generation Ad5 vector based vaccine platform.

[00157] First generation, or E1-deleted adenovirus vectors Ad5 [E1-] are constructed such that a transgene replaces only the E1 region of genes. Typically, about 90% of the wild-type Ad5 genome is retained in the vector. Ad5 [E1-] vectors have a decreased ability to replicate and cannot produce infectious virus after infection of cells that do not express the Ad5 E1 genes. The recombinant Ad5 [E1-] vectors are propagated in human cells (e.g., 293 cells) allowing for Ad5 [E1-] vector replication and packaging. Ad5 [E1-] vectors have a number of positive attributes; one of the most important is their relative ease for scale up and cGMP production. Currently, well over 220 human clinical trials utilize Ad5 [E1-] vectors, with more than two thousand subjects given the virus sc, im, or iv. Additionally, Ad5 vectors do not integrate; their genomes remain episomal. Generally, for vectors that do not integrate into the host genome, the risk for insertional mutagenesis and/or germ-line transmission is extremely low if at all. Conventional Ad5 [E1-] vectors have a carrying capacity that approaches 7 kb.

[00158] Ad5-based vectors with deletions of the E1 and the E2b regions (Ad5 [E1-, E2b-]), the latter encoding the DNA polymerase and the pre-terminal protein, by virtue of diminished late phase viral protein expression, provide an opportunity to avoid immunological clearance and induce more potent immune responses against the encoded tumor antigen transgene in Ad-immune hosts. The new Ad5 platform has additional deletions in the E2b region, removing the DNA polymerase and the preterminal protein genes. The Ad5 [E1-, E2b-] platform has an expanded cloning capacity that is sufficient to allow

inclusion of many possible genes. Ad5 [E1-, E2b-] vectors have up to about 12 kb gene-carrying capacity as compared to the 7 kb capacity of Ad5 [E1-] vectors, providing space for multiple genes if needed. In some embodiments, an insert of more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 kb is introduced into an Ad5 vector, such as the Ad5 [E1-, E2b-] vector. Deletion of the E2b region confers advantageous immune properties on the Ad5 vectors of the invention, often eliciting potent immune responses to target transgene antigens while minimizing the immune responses to Ad viral proteins.

[00159] In various embodiments, Ad5 [E1-, E2b-] vectors of the invention induce a potent CMI, as well as antibodies against the vector expressed vaccine antigens even in the presence of Ad immunity. Ad5 [E1-, E2b-] vectors also have reduced adverse reactions as compared to Ad5 [E1-] vectors, in particular the appearance of hepatotoxicity and tissue damage. A key aspect of these Ad5 vectors is that expression of Ad late genes is greatly reduced. For example, production of the capsid fiber proteins could be detected *in vivo* for Ad5 [E1-] vectors, while fiber expression was ablated from Ad5 [E1-, E2b-] vector vaccines. The innate immune response to wild type Ad is complex. Proteins deleted from the Ad5 [E1-, E2b-] vectors generally play an important role. Specifically, Ad5 [E1-, E2b-] vectors with deletions of preterminal protein or DNA polymerase display reduced inflammation during the first 24 to 72 h following injection compared to Ad5 [E1-] vectors. In various embodiments, the lack of Ad5 gene expression renders infected cells invisible to anti-Ad activity and permits infected cells to express the transgene for extended periods of time, which develops immunity to the target.

[00160] Various embodiments of the invention contemplate increasing the capability for the Ad5 [E1-, E2b-] vectors to transduce dendritic cells, improving antigen specific immune responses in the vaccine by taking advantage of the reduced inflammatory response against Ad5 [E1-, E2b-] vector viral proteins and the resulting evasion of pre-existing Ad immunity.

Replication defective Ad5 vector

[00161] Attempts to overcome anti-Ad immunity have included use of alternative Ad serotypes and/or alterations in the Ad5 viral capsid protein each with limited success and the potential for significantly altering biodistribution of the resultant vaccines. Therefore, a completely novel approach was attempted by further reducing the expression of viral proteins from the E1 deleted Ad5 vectors, proteins known to be targets of pre-existing Ad immunity. Specifically, a novel recombinant Ad5 platform has been described with deletions in the early 1 (E1) gene region and additional deletions in the early 2b (E2b) gene region (Ad5 [E1-, E2b-]). Deletion of the E2b region (that encodes DNA polymerase and the pre-terminal protein) results in decreased viral DNA replication and late phase viral protein expression. This vector platform can be used to induce CMI responses in animal models of cancer and infectious disease and more importantly, this recombinant Ad5 gene delivery platform overcomes the barrier of Ad5 immunity and can be used in the setting of pre-existing and/or vector-induced Ad immunity thus enabling multiple homologous administrations of the vaccine. In particular embodiments, the present invention

relates to a replication defective adenovirus vector of serotype 5 comprising a sequence encoding an immunogenic polypeptide. The immunogenic polypeptide may be a mutant, natural variant, or a fragment thereof.

[00162] In some embodiments, the replication defective adenovirus vector comprises a modified sequence encoding a polypeptide with at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to a wild-type immunogenic polypeptide or a fragment thereof. In some embodiments, the replication defective adenovirus vector comprises a modified sequence encoding a subunit of a wild-type polypeptide. The compositions and methods of the invention, in some embodiments, relate to an adenovirus-derived vector comprising at least 60% sequence identity to SEQ. ID. NO:3.

[00163] In some embodiments, an adenovirus-derived vector, optionally relating to a replication defective adenovirus, comprises a sequence with at least 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.8%, 99.9% identity to SEQ. ID. NO:3 or a sequence generated from SEQ. ID. NO:3 by alternative codon replacements. In various embodiments, the adenovirus-derived vectors described herein have a deletion in the E2b region, and optionally, in the E1 region, the deletion conferring a variety of advantages to the use of the vectors in immunotherapy as described herein.

[00164] Certain regions within the adenovirus genome serve essential functions and may need to be substantially conserved when constructing the replication defective adenovirus vectors of the invention. These regions are further described in Lauer et al., J. Gen. Virol., 85, 2615-25 (2004), Leza et al., J. Virol., p. 3003-13 (1988), and Miralles et al., J. Bio Chem., Vol. 264, No. 18, p. 10763-72 (1983). Recombinant nucleic acid vectors comprising a sequence with identity values of at least 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.8%, 99.9% to a portion of SEQ. ID. NO:3, such as a portion comprising at least about 100, 250, 500, 1000 or more bases of SEQ. ID. NO:3 are within the bounds of the invention.

[00165] The present invention contemplates the use of E2b deleted adenovirus vectors, such as those described in U.S. Pat. Nos. 6,063,622; 6,451,596; 6,057,158; 6,083,750; and 8,298,549.

The vectors with deletions in the E2b regions in many cases cripple viral protein expression and/or decrease the frequency of generating replication competent Ad (RCA). Propagation of these E2b deleted adenovirus vectors can be done utilizing cell lines that express the deleted E2b gene products. Such packaging cell lines are provided herein; e.g., E.C7 (formally called C-7), derived from the HEK-2p3 cell line.

[00166] Further, the E2b gene products, DNA polymerase and preterminal protein, can be constitutively expressed in E.C7, or similar cells along with the E1 gene products. Transfer of gene segments from the Ad genome to the production cell line has immediate benefits: (1) increased carrying capacity; and, (2) a decreased potential of RCA generation, typically requiring two or more independent recombination events to generate RCA. The E1, Ad DNA polymerase and/or preterminal protein expressing cell lines

used in the present invention can enable the propagation of adenovirus vectors with a carrying capacity approaching 13 kb, without the need for a contaminating helper virus. In addition, when genes critical to the viral life cycle are deleted (e.g., the E2b genes), a further crippling of Ad to replicate or express other viral gene proteins occurs. This can decrease immune recognition of infected cells, and extend durations of foreign transgene expression.

[00167] E1, DNA polymerase, and preterminal protein deleted vectors are typically unable to express the respective proteins from the E1 and E2b regions. Further, they may show a lack of expression of most of the viral structural proteins. For example, the major late promoter (MLP) of Ad is responsible for transcription of the late structural proteins L1 through L5. Though the MLP is minimally active prior to Ad genome replication, the highly toxic Ad late genes are primarily transcribed and translated from the mLP only after viral genome replication has occurred. This cis-dependent activation of late gene transcription is a feature of DNA viruses in general, such as in the growth of polyoma and SV-40. The DNA polymerase and preterminal proteins are important for Ad replication (unlike the E4 or protein IX proteins). Their deletion can be extremely detrimental to adenovirus vector late gene expression, and the toxic effects of that expression in cells such as APCs.

[00168] The adenovirus vectors can include a deletion in the E2b region of the Ad genome and, optionally, the E1 region. In some cases, such vectors do not have any other regions of the Ad genome deleted. The adenovirus vectors can include a deletion in the E2b region of the Ad genome and deletions in the E1 and E3 regions. In some cases, such vectors have no other regions deleted. The adenovirus vectors can include a deletion in the E2b region of the Ad genome and deletions in the E1, E3 and partial or complete removal of the E4 regions. In some cases, such vectors have no other deletions. The adenovirus vectors can include a deletion in the E2b region of the Ad genome and deletions in the E1 and/or E4 regions. In some cases, such vectors contain no other deletions. The adenovirus vectors can include a deletion in the E2a, E2b and/or E4 regions of the Ad genome. In some cases, such vectors have no other deletions. The adenovirus vectors can have the E1 and/or DNA polymerase functions of the E2b region deleted. In some cases, such vectors have no other deletions. The adenovirus vectors can have the E1 and/or the preterminal protein functions of the E2b region deleted. In some cases, such vectors have no other deletions. The adenovirus vectors can have the E1, DNA polymerase and/or the preterminal protein functions deleted. In some cases, such vectors have no other deletions. The adenovirus vectors can have at least a portion of the E2b region and/or the E1 region. In some cases, such vectors are not gutted adenovirus vectors. In this regard, the vectors may be deleted for both the DNA polymerase and the preterminal protein functions of the E2b region. The adenovirus vectors can have a deletion in the E1, E2b and/or 100K regions of the adenovirus genome. The adenovirus vectors can comprise vectors having the E1, E2b and/or protease functions deleted. In some cases, such vectors have no other deletions. The adenovirus vectors can have the E1 and/or the E2b regions deleted, while the fiber genes have been

modified by mutation or other alterations (for example to alter Ad tropism). Removal of genes from the E3 or E4 regions may be added to any of the adenovirus vectors mentioned. In certain embodiments, the adenovirus vector may be a gutted adenovirus vector.

[00169] Other regions of the Ad genome can be deleted. A “deletion” in a particular region of the Ad genome refers to a specific DNA sequence that is mutated or removed in such a way so as to prevent expression and/or function of at least one gene product encoded by that region (e.g., E2b functions of DNA polymerase or preterminal protein function). Deletions encompass deletions within exons encoding portions of proteins as well as deletions within promoter and leader sequences. A deletion within a particular region refers to a deletion of at least one base pair within that region of the Ad genome. More than one base pair can be deleted. For example, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 base pairs can be deleted from a particular region. The deletion can be more than 150, 160, 170, 180, 190, 200, 250, or 300 base pairs within a particular region of the Ad genome. These deletions can prevent expression and/or function of the gene product encoded by the region. For example, a particular region of the Ad genome can include one or more point mutations such that one or more encoded proteins is non-functional. Such mutations include residues that are replaced with a different residue leading to a change in the amino acid sequence that result in a nonfunctional protein. Exemplary deletions or mutations in the Ad genome include one or more of E1a, E1b, E2a, E2b, E3, E4, L1, L2, L3, L4, L5, TP, POL, IV, and VA regions. Deleted adenovirus vectors can be made, for example, using recombinant techniques.

[00170] Ad vectors for use in the present invention can be successfully grown to high titers using an appropriate packaging cell line that constitutively expresses E2b gene products and products of any of the necessary genes that may have been deleted. HEK-293-derived cells that not only constitutively express the E1 and DNA polymerase proteins, but also the Ad-preterminal protein, can be used. E.C7 cells can be used, for example, to grow high titer stocks of the adenovirus vectors.

[00171] To delete critical genes from self-propagating adenovirus vectors, proteins encoded by the targeted genes can first be coexpressed in HEK-293 cells, or similar, along with E1 proteins. For example, those proteins which are non-toxic when coexpressed constitutively (or toxic proteins inducibly-expressed) can be selectively utilized. Coexpression in HEK-293 cells of the E1 and E4 genes is possible (for example utilizing inducible, not constitutive, promoters). The E1 and protein IX genes, a virion structural protein, can be coexpressed. Further coexpression of the E1, E4, and protein IX genes is also possible. E1 and 100K genes can be expressed in trans-complementing cell lines, as can E1 and protease genes.

[00172] Cell lines coexpressing E1 and E2b gene products for use in growing high titers of E2b deleted Ad particles can be used. Useful cell lines constitutively express the approximately 140 kDa Ad-DNA polymerase and/or the approximately 90 kDa preterminal protein. Cell lines that have high-level,

constitutive coexpression of the E1, DNA polymerase, and preterminal proteins, without toxicity (e.g. E.C7), are desirable for use in propagating Ad for use in multiple vaccinations. These cell lines permit the propagation of adenovirus vectors deleted for the E1, DNA polymerase, and preterminal proteins.

[00173] The recombinant Ad of the present invention can be propagated using, for example, tissue culture plates containing E.C7 cells infected with Ad vector virus stocks at an appropriate MOI (e.g., 5) and incubated at 37 °C for 40-96 h. The infected cells can be harvested, resuspended in 10 mM Tris-Cl (pH 8.0), and sonicated, and the virus can be purified by two rounds of cesium chloride density centrifugation. The virus containing band can be desalted over a column, sucrose or glycerol can be added, and aliquots can be stored at -80 °C. Virus can be placed in a solution designed to enhance its stability, such as A195. The titer of the stock can be measured (e.g., by measurement of the optical density at 260 nm of an aliquot of the virus after lysis). Plasmid DNA, either linear or circular, encompassing the entire recombinant E2b deleted adenovirus vector can be transfected into E.C7, or similar cells, and incubated at 37 °C until evidence of viral production is present (e.g. cytopathic effect). Conditioned media from cells can be used to infect more cells to expand the amount of virus produced before purification. Purification can be accomplished, for example, by two rounds of cesium chloride density centrifugation or selective filtration. Virus may be purified by chromatography using commercially available products or custom chromatographic columns.

[00174] The compositions of the present invention can comprise enough virus to ensure that cells to be infected are confronted with a certain number of viruses. Thus, in various embodiments, the present invention provides a stock of recombinant Ad, such as an RCA-free stock of recombinant Ad. Viral stocks can vary considerably in titer, depending largely on viral genotype and the protocol and cell lines used to prepare them. Viral stocks can have a titer of at least about 10⁶, 10⁷, or 10⁸ pfu/mL, or higher, such as at least about 10⁹, 10¹⁰, 10¹¹, or 10¹² pfu/mL. Depending on the nature of the recombinant virus and the packaging cell line, a viral stock of the present invention can have a titer of even about 10¹³ particles/ml or higher.

[00175] A replication defective adenovirus vector (e.g., SEQ. ID. NO.:3) can comprise a sequence encoding a target antigen, a fragment thereof, or a variant thereof, at a suitable position. In some embodiments, a replication defective adenovirus vector (e.g., SEQ. ID. NO.:3) can comprise a sequence encoding a target antigen described herein, or a fragment, a variant, or a variant fragment thereof, at a position replacing the nucleic acid sequence encoding a CEA or a variant CEA (e.g., SEQ. ID. NO.:1). In some embodiments, a replication defective adenovirus vector (e.g., SEQ. ID. NO.:3) can comprise a sequence encoding a target antigen described herein, or a fragment, a variant, or a variant fragment thereof, at a position replacing the nucleic acid sequence encoding a CEA or a variant CEA (e.g., SEQ. ID. NO.:2). In some embodiments, a replication defective adenovirus vector (e.g., SEQ. ID. NO.:3) can comprise a sequence encoding a target antigen described herein, or a fragment, a variant, or a variant

fragment thereof, at a position replacing the nucleic acid sequence encoding a MUC1 or a variant MUC1 (e.g., SEQ. ID. NO.:5, SEQ. ID. NO.:6 or SEQ. ID. NO.:9). In some embodiments, a replication defective adenovirus vector (e.g., SEQ. ID. NO.:3) can comprise a sequence encoding a target antigen described herein, or a fragment, a variant, or a variant fragment thereof, at a position replacing the nucleic acid sequence encoding a T or a variant T (e.g., SEQ. ID. NO.:7 or SEQ. ID. NO.:8).

Polynucleotides and Variants Encoding Antigen Targets

[00176] The present disclosure further provides nucleic acid sequences, also referred to herein as polynucleotides that encode one or more target antigens of interest, or fragments or variants thereof. As such, the present invention provides polynucleotides that encode target antigens from any source as described further herein, vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors. In order to express a desired target antigen polypeptide, nucleotide sequences encoding the polypeptide, or functional equivalents, can be inserted into an appropriate Ad vector (e.g., using recombinant techniques). The appropriate adenovirus vector may contain the necessary elements for the transcription and translation of the inserted coding sequence and any desired linkers. Methods which are well known to those skilled in the art may be used to construct these adenovirus vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

[00177] Polynucleotides may comprise a native sequence (i.e., an endogenous sequence that encodes a target antigen polypeptide/protein/epitope of the invention or a portion thereof) or may comprise a sequence that encodes a variant, fragment, or derivative of such a sequence. Polynucleotide sequences can encode target antigen proteins. In some embodiments, polynucleotides represent a novel gene sequence optimized for expression in specific cell types that may substantially vary from the native nucleotide sequence or variant but encode a similar protein antigen.

[00178] In other related embodiments, polynucleotide variants have substantial identity to native sequences encoding proteins (e.g., target antigens of interest), for example those comprising at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a native polynucleotide sequence encoding the polypeptides (e.g., BLAST analysis using standard parameters). These values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Polynucleotides can encode a protein comprising for example at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a protein sequence encoded by a native polynucleotide sequence.

[00179] Polynucleotides can comprise at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 11, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, or 1000 or more contiguous nucleotides encoding a polypeptide (e.g., target protein antigens), and all intermediate lengths there between. “Intermediate lengths”, in this context, refers to any length between the quoted values, such as 16, 17, 18, 19, etc.; 21, 22, 23, etc.; 30, 31, 32, etc.; 50, 51, 52, 53, etc.; 100, 101, 102, 103, etc.; 150, 151, 152, 153, etc.; including all integers through 200-500; 500-1,000, and the like. A polynucleotide sequence may be extended at one or both ends by additional nucleotides not found in the native sequence encoding a polypeptide, such as an epitope or heterologous target protein. This additional sequence may consist of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides or more, at either end of the disclosed sequence or at both ends of the disclosed sequence.

[00180] The polynucleotides, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, expression control sequences, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. Illustrative polynucleotide segments with total lengths of about 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

[00181] A mutagenesis approach, such as site-specific mutagenesis, can be employed to prepare target antigen sequences. Specific modifications in a polypeptide sequence can be made through mutagenesis of the underlying polynucleotides that encode them. Site-specific mutagenesis can be used to make mutants through the use of oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. For example, a primer comprising about 14 to about 25 nucleotides or so in length can be employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered. Mutations may be made in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

[00182] Mutagenesis of polynucleotide sequences can be used to alter one or more properties of the encoded polypeptide, such as the immunogenicity of an epitope comprised in a polypeptide or the oncogenicity of a target antigen. Assays to test the immunogenicity of a polypeptide include, but are not limited to, T-cell cytotoxicity assays (CTL/chromium release assays), T-cell proliferation assays,

intracellular cytokine staining, ELISA, ELISpot, etc. Other ways to obtain sequence variants of peptides and the DNA sequences encoding them can be employed. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

[00183] Polynucleotide segments or fragments encoding the polypeptides of the present invention may be readily prepared by, for example, directly synthesizing the fragment by chemical means. Fragments may be obtained by application of nucleic acid reproduction technology, such as PCR, by introducing selected sequences into recombinant vectors for recombinant production.

[00184] A variety of vector/host systems may be utilized to contain and produce polynucleotide sequences. Exemplary systems include microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA vectors; yeast transformed with yeast vectors; insect cell systems infected with virus vectors (e.g., baculovirus); plant cell systems transformed with virus vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

[00185] Control elements or regulatory sequences present in an Ad vector may include those non-translated regions of the vector-enhancers, promoters, and 5' and 3' untranslated regions. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, sequences encoding a polypeptide of interest may be ligated into an Ad transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells. In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

[00186] Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest (e.g., ATG initiation codon and adjacent sequences). Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used. Specific termination sequences, either for transcription or translation, may also be incorporated in order to achieve efficient translation of the sequence encoding the polypeptide of choice.

[00187] A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products (e.g., target antigens), can be used (e.g., using polyclonal or monoclonal antibodies specific for the product). Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing

monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed.

[00188] The Ad vectors can comprise a product that can be detected or selected for, such as a reporter gene whose product can be detected, such as by fluorescence, enzyme activity on a chromogenic or fluorescent substrate, and the like, or selected for by growth conditions. Exemplary reporter genes include green fluorescent protein (GFP), β -galactosidase, chloramphenicol acetyltransferase (CAT), luciferase, neomycin phosphotransferase, secreted alkaline phosphatase (SEAP), and human growth hormone (HGH). Exemplary selectable markers include drug resistances, such as neomycin (G418), hygromycin, and the like.

[00189] The Ad vectors can also comprise a promoter or expression control sequence. The choice of the promoter will depend in part upon the targeted cell type and the degree or type of control desired.

Promoters that are suitable within the context of the present invention include, without limitation, constitutive, inducible, tissue specific, cell type specific, temporal specific, or event-specific. Examples of constitutive or nonspecific promoters include the SV40 early promoter, the SV40 late promoter, CMV early gene promoter, bovine papilloma virus promoter, and adenovirus promoter. In addition to viral promoters, cellular promoters are also amenable within the context of this invention. In particular, cellular promoters for the so-called housekeeping genes are useful (e.g., β -actin). Viral promoters are generally stronger promoters than cellular promoters. Inducible promoters may also be used. These promoters include MMTV LTR, inducible by dexamethasone, metallothionein, inducible by heavy metals, and promoters with cAMP response elements, inducible by cAMP, heat shock promoter. By using an inducible promoter, the nucleic acid may be delivered to a cell and will remain quiescent until the addition of the inducer. This allows further control on the timing of production of the protein of interest. Event-type specific promoters (e.g., HIV LTR) can be used, which are active or upregulated only upon the occurrence of an event, such as tumorigenicity or viral infection, for example. The HIV LTR promoter is inactive unless the tat gene product is present, which occurs upon viral infection. Some event-type promoters are also tissue-specific. Preferred event-type specific promoters include promoters activated upon viral infection.

[00190] Examples of promoters include promoters for α -fetoprotein, α -actin, myo D, carcinoembryonic antigen, VEGF-receptor; FGF receptor; TEK or tie 2; tie; urokinase receptor; E- and P-selectins; VCAM-1; endoglin; endosialin; α V- β 3 integrin; endothelin-1; ICAM-3; E9 antigen; von Willebrand factor; CD44; CD40; vascular-endothelial cadherin; notch 4, high molecular weight melanoma-associated antigen; prostate specific antigen-1, probasin, FGF receptor, VEGF receptor, erb B2; erb B3; erb B4; MUC-1; HSP-27; int-1; int-2, CEA, HBEGF receptor; EGF receptor; tyrosinase, MAGE, IL-2 receptor; prostatic acid phosphatase, probasin, prostate specific membrane antigen, α -crystallin, PDGF receptor, integrin receptor, α -actin, SM1 and SM2 myosin heavy chains, calponin-h1, SM22 α -angiotensin

receptor, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, immunoglobulin heavy chain, immunoglobulin light chain, and CD4.

[00191] Repressor sequences, negative regulators, or tissue-specific silencers may be inserted to reduce non-specific expression of the polynucleotide. Multiple repressor elements may be inserted in the promoter region. Repression of transcription is independent of the orientation of repressor elements or distance from the promoter. One type of repressor sequence is an insulator sequence. Such sequences inhibit transcription and can silence background transcription. Negative regulatory elements can be located in the promoter regions of a number of different genes. The repressor element can function as a repressor of transcription in the absence of factors, such as steroids, as does the NSE in the promoter region of the ovalbumin gene. These negative regulatory elements can bind specific protein complexes from oviduct, none of which are sensitive to steroids. Three different elements are located in the promoter of the ovalbumin gene. Oligonucleotides corresponding to portions of these elements can repress viral transcription of the TK reporter. One of the silencer elements shares sequence identity with silencers in other genes (TCTCTCCNA (SEQ ID NO:11)).

[00192] Elements that increase the expression of the desired target antigen can be incorporated into the nucleic acid sequence of the Ad vectors described herein. Exemplary elements include internal ribosome binding sites (IRESs). IRESs can increase translation efficiency. As well, other sequences may enhance expression. For some genes, sequences especially at the 5' end may inhibit transcription and/or translation. These sequences are usually palindromes that can form hairpin structures. In some cases, such sequences in the nucleic acid to be delivered are deleted. Expression levels of the transcript or translated product can be assayed to confirm or ascertain which sequences affect expression. Transcript levels may be assayed by any known method, including Northern blot hybridization, RNase probe protection and the like. Protein levels may be assayed by any known method, including ELISA.

Antigen-Specific Immunotherapies and Vaccines

[00193] The present disclosure provides for single antigen or combination antigen immunization against MUC1, MUC1c, MUC1n, T, or CEA utilizing such vectors and other vectors as provided herein. The present disclosure provides for therapeutic vaccines against MUC1, MUC1c, MUC1n, T, or CEA. The present disclosure provides for prophylactic vaccines against MUC1, MUC1c, MUC1n, T, or CEA. Further, in various embodiments, the composition and methods provide herein can lead to clinical responses, such as altered disease progression or life expectancy.

[00194] Ad5 [E1-] vectors encoding a variety of antigens can be used to efficiently transduce 95% of *ex vivo* exposed DC's to high titers of the vector. Importantly, the inventors have discovered increasing levels of foreign gene expression in the DC with increasing multiplicities of infection (MOI) with the vector. DCs infected with Ad5 [E1-] vectors can encode a variety of antigens (including the tumor antigens MART-1, MAGE-A4, DF3/MUC1, p53, hugp100 melanoma antigen, polyoma virus middle -T

antigen) that have the propensity to induce antigen specific CTL responses, have an enhanced antigen presentation capacity, and/or have an improved ability to initiate T-cell proliferation in mixed lymphocyte reactions. Immunization of animals with dendritic cells (DCs) previously transduced by Ad5 vectors encoding tumor specific antigens can be used to induce significant levels of protection for the animals when challenged with tumor cells expressing the respective antigen. Interestingly, intra-tumoral injection of Ads encoding IL-7 is less effective than injection of DCs transduced with IL-7 encoding Ad5 vectors at inducing antitumor immunity. *Ex vivo* transduction of DCs by Ad5 vectors is contemplated by the present disclosure. *Ex vivo* DC transduction strategies can be used to induce recipient host tolerance. For example, Ad5 mediated delivery of the CTLA4Ig into DCs can block interactions of the DCs CD80 with CD28 molecules present on T-cells.

[00195] Ad5 vector capsid interactions with DCs may trigger several beneficial responses, which may be enhancing the propensity of DCs to present antigens encoded by Ad5 vectors. For example, immature DCs, though specialized in antigen uptake, are relatively inefficient effectors of T-cell activation. DC maturation coincides with the enhanced ability of DCs to drive T-cell immunity. In some instances, the compositions and methods of the invention take advantage of an Ad5 infection resulting in direct induction of DC maturation. Ad vector infection of immature bone marrow derived DCs from mice may upregulate cell surface markers normally associated with DC maturation (MHC I and II, CD40, CD80, CD86, and ICAM-1) as well as down-regulation of CD11c, an integrin down regulated upon myeloid DC maturation. In some instances, Ad vector infection triggers IL-12 production by DCs, a marker of DC maturation. Without being bound by theory, these events may possibly be due to Ad5 triggered activation of NF- κ B pathways. Mature DCs can be efficiently transduced by Ad vectors, and do not lose their functional potential to stimulate the proliferation of naive T-cells at lower MOI, as demonstrated by mature CD83 $+$ human DC (derived from peripheral blood monocytes). However, mature DCs may also be less infectable than immature ones. Modification of capsid proteins can be used as a strategy to optimize infection of DC by Ad vectors, as well as enhancing functional maturation, for example using the CD40L receptor as a viral vector receptor, rather than using the normal CAR receptor infection mechanisms.

[00196] In various embodiments the compositions and methods of the invention comprising an Ad5 [E1-, E2b-] vector(s) MUC1, T and CEA vaccine effect of increased overall survival (OS) within the bounds of technical safety. In various embodiments the compositions and methods of the invention comprising an Ad5 [E1-, E2b-] vector(s) MUC1c, T and CEA vaccine effect of increased overall survival (OS) within the bounds of technical safety. In various embodiments the compositions and methods of the invention comprising an Ad5 [E1-, E2b-] vector(s) MUC1n, T and CEA vaccine effect of increased overall survival (OS) within the bounds of technical safety.

[00197] In some embodiments, the antigen targets are associated with benign tumors. In some embodiments, the antigens targeted are associated with pre-cancerous tumors.

[00198] In some embodiments, the antigens targeted are associated with carcinomas, in situ carcinomas, metastatic tumors, neuroblastoma, sarcomas, myosarcoma, leiomyosarcoma, retinoblastoma, hepatoma, rhabdomyosarcoma, plasmacytomas, adenomas, gliomas, thymomas, or osteosarcoma. In some embodiments, the antigens targeted are associated with a specific type of cancer such as neurologic cancers, brain cancer, thyroid cancer, head and neck cancer, melanoma, leukemia, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), and chronic lymphocytic leukemia (CLL), non-Hodgkin's lymphoma, multiple myeloma, Hodgkin's disease, breast cancer, bladder cancer, prostate cancer, colorectal cancer, colon cancer, kidney cancer, renal cell carcinoma, pancreatic cancer, esophageal cancer, lung cancer, mesothelioma, ovarian cancer, cervical cancer, endometrial cancer, uterine cancer, germ cell tumors, testicular cancer, gastric cancer, or other cancers, or any clinical (e.g. TNM, Histopathological, Staging or Grading systems or a combination thereof) or molecular subtype thereof. In some embodiments, the antigens targeted are associated with a specific clinical or molecular subtype of cancer. By way of example, breast cancer can be divided into at least four molecular subtypes including Luminal A, Luminal B, Triple negative/basal-like, and HER2 type. By way of example, Prostate cancer can be subdivided TNM, Gleason score, or molecular expression of the PSA protein.

[00199] As noted above, the adenovirus vectors of the present invention comprise nucleic acid sequences that encode one or more target proteins or antigens of interest. In this regard, the vectors may contain nucleic acid encoding 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more different target antigens of interest. The target antigens may be a full length protein or may be a fragment (e.g., an epitope) thereof. The adenovirus vectors may contain nucleic acid sequences encoding multiple fragments or epitopes from one target protein of interest or may contain one or more fragments or epitopes from numerous different target proteins of interest. A target antigen may comprise any substance against which it is desirable to generate an immune response but generally, the target antigen is a protein. A target antigen may comprise a full length protein, a subunit of a protein, an isoform of a protein, or a fragment thereof that induces an immune response (i.e., an immunogenic fragment). A target antigen or fragment thereof may be modified, e.g., to reduce one or more biological activities of the target antigen or to enhance its immunogenicity. The target antigen or target protein can be MUC1, MUC1c, MUC1n, T, CEA, or any combination thereof.

[00200] In certain embodiments, immunogenic fragments bind to an MHC class I or class II molecule. An immunogenic fragment may "bind to" an MHC class I or class II molecule if such binding is detectable using any assay known in the art. For example, the ability of a polypeptide to bind to MHC class I may be evaluated indirectly by monitoring the ability to promote incorporation of ¹²⁵I labeled β -2-

microglobulin (β -2m) into MHC class I/ β 2m/peptide heterotrimeric complexes. Alternatively, functional peptide competition assays that are known in the art may be employed. Immunogenic fragments of polypeptides may generally be identified using well known techniques. Representative techniques for identifying immunogenic fragments include screening polypeptides for the ability to react with antigen-specific antisera and/or T-cell lines or clones. An immunogenic fragment of a particular target polypeptide is a fragment that reacts with such antisera and/or T-cells at a level that is not substantially less than the reactivity of the full length target polypeptide (e.g., in an ELISA and/or T-cell reactivity assay). In other words, an immunogenic fragment may react within such assays at a level that is similar to or greater than the reactivity of the full length polypeptide. Such screens may be performed using methods known in the art.

[00201] In some embodiments, the viral vectors of the present invention comprise heterologous nucleic acid sequences that encode one or more proteins, variants thereof, fusions thereof, or fragments thereof, that can modulate the immune response. In some embodiments, the viral vector of the present invention encodes one or more antibodies against specific antigens, such as anthrax protective antigen, permitting passive immunotherapy. In some embodiments, the viral vectors of the present invention comprise heterologous nucleic acid sequences encoding one or more proteins having therapeutic effect (e.g., anti-viral, anti-bacterial, anti-parasitic, or anti-tumor function). In some embodiments the Second Generation E2b deleted adenovirus vectors comprise a heterologous nucleic acid sequence. In some embodiments, the heterologous nucleic acid sequence is CEA, MUC1, MUC1c, MUC1n, T, a variant, a portion, or any combination thereof.

[00202] Target antigens include, but are not limited to, antigens derived from a variety of tumor proteins. In some embodiments, parts or variants of tumor proteins are employed as target antigens. In some embodiments, parts or variants of tumor proteins being employed as target antigens have a modified, for example, increased ability to effect and immune response against the tumor protein or cells containing the same. A vaccine of the present invention can vaccinate against an antigen. A vaccine can also target an epitope. An antigen can be a tumor cell antigen. An epitope can be a tumor cell epitope. Such a tumor cell epitope may be derived from a wide variety of tumor antigens, such as antigens from tumors resulting from mutations, shared tumor specific antigens, differentiation antigens, and antigens overexpressed in tumors. Tumor-associated antigens (TAAs) may be antigens not normally expressed by the host; they can be mutated, truncated, misfolded, or otherwise abnormal manifestations of molecules normally expressed by the host; they can be identical to molecules normally expressed but expressed at abnormally high levels; or they can be expressed in a context or environment that is abnormal. Tumor-associated antigens may be, for example, proteins or protein fragments, complex carbohydrates, gangliosides, haptens, nucleic acids, other biological molecules or any combinations thereof.

[00203] Illustrative tumor proteins useful in the present invention include, but are not limited to any one or more of, WT1, HPV E6, HPV E7, p53, MAGE-A1, MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A6, MAGE-A10, MAGE-A12, BAGE, DAM-6, DAM-10, GAGE-1, GAGE-2, GAGE-8, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE-7B, NA88-A, NY-ESO-1, MART-1, MC1R, Gp100, PSA, PSM, Tyrosinase, TRP-1, TRP-2, ART-4, CAMEL, CEA, Cyp-B, Her2/neu, BRCA1, Brachyury, Brachyury (TIVS7-2, polymorphism), Brachyury (IVS7 T/C polymorphism), T Brachyury, T, hTERT, hTRT, iCE, MUC1, MUC1 (VNTR polymorphism), MUC1c, MUC1n, MUC2, PRAME, P15, RU1, RU2, SART-1, SART-3, WT1, AFP, β -catenin/m, Caspase-8/m, CEA, CDK-4/m, ELF2M, GnT-V, G250, HSP70-2M, HST-2, KIAA0205, MUM-1, MUM-2, MUM-3, Myosin/m, RAGE, SART-2, TRP-2/INT2, 707-AP, Annexin II, CDC27/m, TPI/mber-abl, ETV6/AML, LDLR/FUT, Pml/RAR α , and TEL/AML1. In some embodiments, the viral vector comprises a target antigen sequence encoding a modified polypeptide selected from WT1, HPV E6, HPV E7, p53, MAGE-A1, MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A6, MAGE-A10, MAGE-A12, BAGE, DAM-6, DAM-10, GAGE-1, GAGE-2, GAGE-8, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE-7B, NA88-A, NY-ESO-1, MART-1, MC1R, Gp100, PSA, PSM, Tyrosinase, TRP-1, TRP-2, ART-4, CAMEL, CEA, Cyp-B, Her2/neu, BRCA1, Brachyury, Brachyury (TIVS7-2, polymorphism), Brachyury (IVS7 T/C polymorphism), T Brachyury, T, hTERT, hTRT, iCE, MUC1, MUC1 (VNTR polymorphism), MUC1c, MUC1n, MUC2, PRAME, P15, RU1, RU2, SART-1, SART-3, WT1, AFP, β -catenin/m, Caspase-8/m, CEA, CDK-4/m, ELF2M, GnT-V, G250, HSP70-2M, HST-2, KIAA0205, MUM-1, MUM-2, MUM-3, Myosin/m, RAGE, SART-2, TRP-2/INT2, 707-AP, Annexin II, CDC27/m, TPI/mber-abl, ETV6/AML, LDLR/FUT, Pml/RAR α , and TEL/AML1 wherein the polypeptide or a fragment thereof has at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to the described sequence.

[00204] Illustrative tumor proteins useful in the present invention include, but are not limited to any one or more of CEA, human epidermal growth factor receptor 2 (*HER2/neu*), human epidermal growth factor receptor 3 (HER3), Human papillomavirus (HPV), MUC1, Prostate-specific antigen (PSA), alpha-actinin-4, ARTC1, CAR-ABL fusion protein (b3a2), B-RAF, CASP-5, CASP-8, beta-catenin, Cdc27, CDK4, CDKN2A, COA-1, dek-can fusion protein, EFTUD2, Elongation factor 2, ETV6-AML1 fusion protein, FLT3-ITD, FN1, GPNMB, LDLR-fucosyltransferase fusion protein, HLA-A2d, HLA-A1 ld, hsp70-2, KIAAO205, MART2, ME1, MUM-1f, MUM-2, MUM-3, neo-PAP, Myosin class I, NFYC, OGT, OS-9, p53, pml-RARalpha fusion protein, PRDX5, PTPRK, K-ras, N-ras, RBAF600, SIRT2, SNRPD1, SYT-SSX1- or -SSX2 fusion protein, TGF-betaRII, triosephosphate isomerase, BAGE-1, GnTVf, HERV-K-MEL, KK-LC-1, KM-HN-1, LAGE-1, MAGE-A9, MAGE-C2, mucink, NA-88, NY-ESO-1/LAGE-2, SAGE, Sp17, SSX-2, SSX-4, TAG-1, TAG-2, TRAG-3, TRP2-INT2g, XAGE-1b, gp100/Pmel17, Kallikrein 4, gammaglobin-A, Melan-A/MART-1, NY-BR-1, OA1, PSA, RAB38/NY-MEL-1, TRP-1/gp75, TRP-2, tyrosinase, adipophilin, AIM-2, ALDH1A1, BCLX (L), BCMA, BING-4,

CPSF, cyclin D1, DKK1, ENAH (hMena), EP-CAM, EphA3, EZH2, FGF5, G250/MN/CAIX, HER-2/neu, IL13Ralpha2, intestinal carboxyl esterase, alpha fetoprotein, M-CSFT, MCSP, mdm-2, MMP-2, MUC1, p53, PBF, PRAME, PSMA, RAGE-1, RGS5, RNF43, RU2AS, secernin 1, SOX10, STEAP1, survivin, Telomerase, and/or VEGF.

[00205] Tumor-associated antigens may be antigens from infectious agents associated with human malignancies. Examples of infectious agents associated with human malignancies include Epstein-Barr virus, Helicobacter pylori, Hepatitis B virus, Hepatitis C virus, Human herpesvirus-8, Human immunodeficiency virus, Human papillomavirus, Human T-cell leukemia virus, liver flukes, and Schistosoma haematobium.

CEA Antigen Targets

[00206] CEA represents an attractive target antigen for immunotherapy since it is over-expressed in nearly all colorectal cancers and pancreatic cancers, and is also expressed by some lung and breast cancers, and uncommon tumors such as medullary thyroid cancer, but is not expressed in other cells of the body except for low-level expression in gastrointestinal epithelium. CEA contains epitopes that may be recognized in an MHC restricted fashion by T-cells.

[00207] The inventors have discovered that multiple homologous immunizations with Ad5 [E1-, E2b-]-CEA(6D), encoding the tumor antigen CEA, induced CEA-specific cell-mediated immune (CMI) responses with antitumor activity in mice despite the presence of pre-existing or induced Ad5-neutralizing antibody. In the present phase I/II study, cohorts of patients with advanced colorectal cancer were immunized with escalating doses of Ad5 [E1-, E2b-]-CEA(6D). CEA-specific CMI responses were observed despite the presence of pre-existing Ad5 immunity in a majority (61.3%) of patients. Importantly, there was minimal toxicity, and overall patient survival (48% at 12 months) was similar regardless of pre-existing Ad5 neutralizing antibody titers. The results demonstrate that, in cancer patients, the novel Ad5 [E1-, E2b-] gene delivery platform generates significant CMI responses to the tumor antigen CEA in the setting of both naturally acquired and immunization-induced Ad5-specific immunity.

[00208] CEA antigen specific CMI can be, for example, greater than 10, 20, 30, 40, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 5000, 10000, or more IFN- γ spot forming cells (SFC) per 10^6 peripheral blood mononuclear cells (PBMC). In some embodiments, the immune response is raised in a human subject with a preexisting inverse Ad5 neutralizing antibody titer of greater than 50, 100, 150, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 6000, 7000, 8000, 9000, 1000, 12000, 15000 or higher. The immune response may comprise a cell-mediated immunity and/or a humoral immunity as described herein. The immune response may be measured by one or more of intracellular cytokine staining (ICS), ELISpot, proliferation assays, cytotoxic T-cell assays including chromium release or equivalent assays, and gene expression analysis using any number

of polymerase chain reaction (PCR) or RT-PCR based assays, as described herein and to the extent they are available to a person skilled in the art, as well as any other suitable assays known in the art for measuring immune response.

[00209] In some embodiments, the replication defective adenovirus vector comprises a modified sequence encoding a subunit with at least 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to a wild-type subunit of the polypeptide.

[00210] The immunogenic polypeptide may be a mutant CEA or a fragment thereof. In some embodiments, the immunogenic polypeptide comprises a mutant CEA with an Asn->Asp substitution at position 610. In some embodiments, the replication defective adenovirus vector comprises a sequence encoding a polypeptide with at least 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to the immunogenic polypeptide. In some embodiments, the sequence encoding the immunogenic polypeptide comprises the sequence of SEQ. ID. NO:1.

[00211] In some embodiments, the sequence encoding the immunogenic polypeptide comprises a sequence with at least 70% 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to SEQ. ID. NO:1 or a sequence generated from SEQ. ID. NO:1 by alternative codon replacements. In some embodiments, the immunogenic polypeptide encoded by the adenovirus vectors comprise up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, or more point mutations, such as single amino acid substitutions or deletions, as compared to a wild-type human CEA sequence.

[00212] In some embodiments, the immunogenic polypeptide comprises a sequence from SEQ. ID. NO.:2 or a modified version, e.g. comprising up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, or more point mutations, such as single amino acid substitutions or deletions, of SEQ. ID. NO.:2.

[00213] Members of the CEA gene family are subdivided into three subgroups based on sequence similarity, developmental expression patterns and their biological functions: the CEA-related Cell Adhesion Molecule (CEACAM) subgroup containing twelve genes (*CEACAM1*, *CEACAM3-CEACAM8*, *CEACAM16* and *CEACAM18-CEACAM21*), the Pregnancy Specific Glycoprotein (PSG) subgroup containing eleven closely related genes (*PSGI-PSGII*) and a subgroup of eleven pseudogenes (*CEACAMP1-CEACAMP11*). Most members of the CEACAM subgroup have similar structures consist of an extracellular Ig-like domains composed of a single N-terminal V-set domain, with structural homology to the immunoglobulin variable domains, followed by varying numbers of C2-set domains of A or B subtypes, a transmembrane domain and a cytoplasmic domain. There are two members of CEACAM subgroup (*CEACAM16* and *CEACAM20*) that show a few exceptions in the organization of their structures. *CEACAM16* contains two Ig-like V-type domains at its N and C termini and *CEACAM20* contains a truncated Ig-like V-type 1 domain. The CEACAM molecules can be anchored to the cell

surface via their transmembrane domains (*CEACAM5* thought *CEACAM8*) or directly linked to glycoprophosphatidylinositol (GPI) lipid moiety (*CEACAM5*, *CEACAM18* thought *CEACAM21*).

[00214] CEA family members are expressed in different cell types and have a wide range of biological functions. CEACAMs are found prominently on most epithelial cells and are present on different leucocytes. In humans, *CEACAM1*, the ancestor member of CEA family, is expressed on the apical side of epithelial and endothelial cells as well as on lymphoid and myeloid cells. *CEACAM1* mediates cell-cell adhesion through hemophilic (*CEACAM1* to *CEACAM1*) as well as heterothallic (e.g., *CEACAM1* to *CEACAM5*) interactions. In addition, *CEACAM1* is involved in many other biological processes, such as angiogenesis, cell migration, and immune functions. *CEACAM3* and *CEACAM4* expression is largely restricted to granulocytes, and they are able to convey uptake and destruction of several bacterial pathogens including *Neisseria*, *Moraxella*, and *Haemophilus* species.

[00215] Thus, in various embodiments, compositions and methods of the invention relate to raising an immune response against a CEA, selected from the group consisting of *CEACAM1*, *CEACAM3*, *CEACAM4*, *CEACAM5*, *CEACAM6*, *CEACAM7*, *CEACAM8*, *CEACAM16*, *CEACAM18*, *CEACAM19*, *CEACAM20*, *CEACAM21*, *PSG1*, *PSG2*, *PSG3*, *PSG4*, *PSG5*, *PSG6*, *PSG7*, *PSG8*, *PSG9*, and *PSG11*. An immune response may be raised against cells, e.g. cancer cells, expressing or overexpressing one or more of the CEAs, using the methods and compositions of the invention. In some embodiments, the overexpression of the one or more CEAs in such cancer cells is over 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 fold or more compared to non-cancer cells.

Mucin Family Antigen Targets

[00216] The human mucin family (MUC1 to MUC21) includes secreted and transmembrane mucins that play a role in forming protective mucous barriers on epithelial surfaces in the body. These proteins function in to protecting the epithelia lining the respiratory, gastrointestinal tracts, and lining ducts in important organs such as, for example the mammary gland, liver, stomach, pancreas, and kidneys.

[00217] MUC1 (CD227) is a TAA that is over-expressed on a majority of human carcinomas and several hematologic malignancies. MUC1 (GenBank: X80761.1, NCBI: NM_001204285.1) and activates many important cellular pathways known to be involved in human disease. MUC1 is a heterodimeric protein formed by two subunits that is commonly overexpressed in several human cancers. MUC1 undergoes autoproteolysis to generate two subunits MUC1n and MUC1c that, in turn, form a stable noncovalent heterodimer.

[00218] The MUC1 C-terminal subunit (MUC1c) can comprise a 58 aa extracellular domain (ED), a 28 aa transmembrane domain (TM) and a 72 aa cytoplasmic domain (CD). The MUC1c also can contain a “CQC” motif that can allow for dimerization of MUC1 and it can also impart oncogenic function to a cell. In some cases MUC1 can in part oncogenic function through inducing cellular signaling via MUC1c. MUC1c can interact with EGFR, ErbB2 and other receptor tyrosine kinases and contributing to

the activation of the PI3K→AKT and MEK→ERK cellular pathways. In the nucleus, MUC1c activates the Wnt/β-catenin, STAT and NF-κB RelA cellular pathways. In some cases MUC1 can impart oncogenic function through inducing cellular signaling via MUC1n. The MUC1 N-terminal subunit (MUC1n) can comprise variable numbers of 20 amino acid tandem repeats that can be glycosylated. MUC1 is normally expressed at the surface of glandular epithelial cells and is over-expressed and aberrantly glycosylated in carcinomas. MUC1 is a TAA that can be utilized as a target for tumor immunotherapy. Several clinical trials have been and are being performed to evaluate the use of MUC1 in immunotherapeutic vaccines. Importantly, these trials indicate that immunotherapy with MUC1 targeting is safe and may provide survival benefit.

[00219] However, clinical trials have also shown that MUC1 is a relatively poor immunogen. To overcome this, the inventors have identified a T lymphocyte immune enhancer peptide sequence in the C terminus region of the MUC1 oncoprotein (MUC1-C or MUC1c). Compared with the native peptide sequence, the agonist in their modified MUC1-C (a) bound HLA-A2 at lower peptide concentrations, (b) demonstrated a higher avidity for HLA-A2, (c) when used with antigen-presenting cells, induced the production of more IFN-γ by T-cells than with the use of the native peptide, and (d) was capable of more efficiently generating MUC1-specific human T-cell lines from cancer patients. Importantly, T-cell lines generated using the agonist epitope were more efficient than those generated with the native epitope for the lysis of targets pulsed with the native epitope and in the lysis of HLA-A2 human tumor cells expressing MUC1. Additionally, the inventors have identified additional CD8+ cytotoxic T lymphocyte immune enhancer agonist epitopes of MUC1-C.

[00220] The present disclosure provides a potent MUC1-C modified for immune enhancer capability (mMUC1-C or MUC1-C or MUC1c). The present disclosure provides a potent MUC1-C modified for immune enhancer capability incorporated it into a recombinant Ad5 [E1-, E2b-] platform to produce a new and more potent immunotherapeutic vaccine. For example, the immunotherapeutic vaccine can be Ad5 [E1-, E2b-]-mMUC1-C for treating MUC1 expressing cancers or infectious diseases.

[00221] Post-translational modifications play an important role in controlling protein function in the body and in human disease. For example, in addition to proteolytic cleavage discussed above, MUC1 can have several post-translational modifications such as glycosylation, sialylation, palmitoylation, or a combination thereof at specific amino acid residues. Provided herein are immunotherapies targeting glycosylation, sialylation, phosphorylation, or palmitoylation modifications of MUC1.

[00222] MUC1 can be highly glycosylated (N- and O-linked carbohydrates and sialic acid at varying degrees on serine and threonine residues within each tandem repeat, ranging from mono- to penta-glycosylation). Differentially O-glycosylated in breast carcinomas with 3,4-linked GlcNAc. N-glycosylation consists of high-mannose, acidic complex-type and hybrid glycans in the secreted form

MUC1/SEC, and neutral complex-type in the transmembrane form, MUC1/TM.4. The present disclosure provides for immunotherapies targeting differentially O-glycosylated forms of MUC1.

[00223] Further, MUC1 can be sialylated. Membrane-shed glycoproteins from kidney and breast cancer cells have preferentially sialylated core 1 structures, while secreted forms from the same tissues display mainly core 2 structures. The O-glycosylated content is overlapping in both these tissues with terminal fucose and galactose, 2- and 3-linked galactose, 3- and 3,6-linked GalNAc-ol and 4-linked GlcNAc predominating. The present disclosure provides for immunotherapies targeting various sialylation forms of MUC1. Dual palmitoylation on cysteine residues in the CQC motif is required for recycling from endosomes back to the plasma membrane. The present disclosure provides for immunotherapies targeting various palmitoylation forms of MUC1.

[00224] Phosphorylation can affect MUC1's ability to induces specific cell signaling responses that are important for human health. The present disclosure provides for immunotherapies targeting various phosphorylated forms of MUC1. For example, MUC1 can be phosphorylated on tyrosine and serine residues in the C-terminal domain. Phosphorylation on tyrosines in the C-terminal domain can increase nuclear location of MUC1 and β -catenin. Phosphorylation by PKC delta can induce binding of MUC1 to β -catenin/CTNNB1 and decrease formation of β -catenin/E-cadherin complexes. Src-mediated phosphorylation of MUC1 can inhibits interaction with GSK3B. Src- and EGFR-mediated phosphorylation of MUC1 on Tyr-1229 can increase binding to β -catenin/CTNNB1. GSK3B-mediated phosphorylation of MUC1 on Ser-1227 can decrease this interaction but restores the formation of the β -catenin/E-cadherin complex. PDGFR-mediated phosphorylation of MUC1 can increase nuclear colocalization of MUC1CT and CTNNB1. The present disclosure provides for immunotherapies targeting different phosphorylated forms of MUC1, MUC1c and MUC1n known to regulate its cell signaling abilities.

[00225] The disclosure provides for immunotherapies that modulate MUC1c cytoplasmic domain and its functions in the cell. The disclosure provides for immunotherapies that comprise modulating a CQC motif in MUC1c. The disclosure provides for immunotherapies that comprise modulating the extracellular domain (ED), the transmembrane domain (TM), the cytoplasmic domain (CD) of MUC1c, or a combination thereof. The disclosure provides for immunotherapies that comprise modulating MUC1c's ability to induce cellular signaling through EGFR, ErbB2 or other receptor tyrosine kinases. The disclosure provides for immunotherapies that comprise modulating MUC1c's ability to induce PI3K \rightarrow AKT, MEK \rightarrow ERK, Wnt/ β -catenin, STAT, NF- κ B RelA cellular pathways, or combination thereof. In some embodiments, the MUC1c immunotherapy can further comprise CEA.

[00226] The disclosure also provides for immunotherapies that modulate MUC1n and its cellular functions. The disclosure also provides for immunotherapies comprising tandem repeats of MUC1n, the

glycosylation sites on the tandem repeats of MUC1n, or a combination thereof. In some embodiments, the MUC1n immunotherapy further comprises CEA.

[00227] The disclosure also provides vaccines comprising MUC1n, MUC1c, CEA, or a combination thereof. The disclosure provides vaccines comprising MUC1c and CEA. The disclosure also provides vaccines targeting MUC1n and CEA. In some embodiments, the antigen combination is contained in one vector as provided herein. In some embodiments, the antigen combination is contained in a separate vector as provided herein.

[00228] The present invention relates to a replication defective adenovirus vector of serotype 5 comprising a sequence encoding an immunogenic polypeptide. The immunogenic polypeptide may be an isoform of MUC1 or a subunit or a fragment thereof. In some embodiments, the replication defective adenovirus vector comprises a sequence encoding a polypeptide with at least 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to the immunogenic polypeptide. In some embodiments, the sequence encoding the immunogenic polypeptide comprises the sequence of SEQ. ID. NO.:5. In some embodiments, the sequence encoding the immunogenic polypeptide comprises the following sequence identified by SEQ. ID. NO.:6. In some embodiments, the sequence encoding the immunogenic polypeptide comprises the following sequence identified by SEQ. ID. NO.:9. In some embodiments, the sequence encoding the immunogenic polypeptide comprises a sequence with at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to SEQ. ID. NO:5, SEQ. ID. NO:6, SEQ. ID. NO.:9 or a sequence generated from SEQ. ID. NO.:5, SEQ. ID. NO:6, SEQ. ID. NO.:9 by alternative codon replacements. In some embodiments, the immunogenic polypeptide encoded by the adenovirus vectors described herein comprising up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, or more point mutations, such as single amino acid substitutions or deletions, as compared to a wild-type human MUC1 sequence.

Brachyury Antigen Targets

[00229] The disclosure also provides for immunotherapies that comprise one or more antigens to Brachyury. Brachyury (also known as the “T” protein in humans) is a member of the T-box family of transcription factors that play key roles during early development, mostly in the formation and differentiation of normal mesoderm and is characterized by a highly conserved DNA-binding domain designated as T-domain. The epithelial to mesenchymal transition (EMT) is a key step during the progression of primary tumors into a metastatic state in which Brachyury plays a crucial role. The expression of Brachyury in human carcinoma cells induces changes characteristic of EMT, including up-regulation of mesenchymal markers, down-regulation of epithelial markers, and an increase in cell migration and invasion. Conversely, inhibition of Brachyury resulted in down-regulation of mesenchymal markers and loss of cell migration and invasion and diminished the ability of human tumor cells to form metastases. Brachyury can function to mediate epithelial-mesenchymal transition and promotes invasion.

[00230] The disclosure also provides for immunotherapies that modulate Brachyury effect on epithelial-mesenchymal transition function in cell proliferation diseases, such as cancer. The disclosure also provides for immunotherapies that modulate Brachyury's ability to promote invasion in cell proliferation diseases, such as cancer. The disclosure also provides for immunotherapies that modulate the DNA binding function of T-box domain of Brachyury. In some embodiments, the Brachyury immunotherapy can further comprise one or more antigens to CEA or MUC1, MUC1c or MUC1n.

[00231] Brachyury expression is nearly undetectable in most normal human tissues and is highly restricted to human tumors and often overexpressed making it an attractive target antigen for immunotherapy. In human, Brachyury is encoded by the T gene (GenBank: AJ001699.1, NCBI: NM_003181.3). There are at least two different isoforms produced by alternative splicing found in humans. Each isoform has a number of natural variants.

[00232] Brachyury is immunogenic and Brachyury-specific CD8+ T-cells expanded in vitro can lyse Brachyury expressing tumor cells. These features of Brachyury make it an attractive TAA for immunotherapy. The Brachyury protein is a T-box transcription factor. It can bind to a specific DNA element, a near palindromic sequence "TCACACCT" through a region in its N-terminus, called the T-box to activate gene transcription when bound to such a site.

[00233] The disclosure also provides vaccines comprising Brachyury, CEA, or a combination thereof. In some embodiments, the antigen combination is contained in one vector as provided herein. In some embodiments, the antigen combination is contained in a separate vector as provided herein.

[00234] In particular embodiments, the present invention relates to a replication defective adenovirus vector of serotype 5 comprising a sequence encoding an immunogenic polypeptide. The immunogenic polypeptide may be an isoform of Brachyury or a subunit or a fragment thereof. In some embodiments, the replication defective adenovirus vector comprises a sequence encoding a polypeptide with at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to the immunogenic polypeptide. In some embodiments, the sequence encoding the immunogenic polypeptide comprises the following sequence identified by SEQ. ID. NO.:7. In some embodiments, the replication defective adenovirus vector comprises a sequence encoding a polypeptide with at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to the immunogenic polypeptide. In some embodiments, the sequence encoding the immunogenic polypeptide comprises the sequence of SEQ. ID. NO.:8. In some embodiments, the sequence encoding the immunogenic polypeptide comprises a sequence with at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% identity to SEQ. ID. NO:7, SEQ. ID. NO:8 or a sequence generated from SEQ. ID. NO:7, SEQ. ID. NO:8 by alternative codon replacements. In some embodiments, the immunogenic polypeptide encoded by the adenovirus vectors described herein comprising up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, or more

point mutations, such as single amino acid substitutions or deletions, as compared to a wild-type human Brachyury sequence.

Infectious Disease Associated Antigen Targets

[00235] Target antigens of the present disclosure include, but are not limited to, antigens derived from any of a variety of infectious agents such as parasites, bacteria, virus, prions, and the like. An infectious agent may refer to any living organism capable of infecting a host. Infectious agents include, for example, bacteria, any variety of viruses, such as, single stranded RNA viruses, single stranded DNA viruses, fungi, parasites, and protozoa.

[00236] Examples of infectious disease associated target antigens that can be used with the compositions and the methods of the present disclosure can be derived from the following:

Actinobacillus spp., Actinomyces spp., Adenovirus (types 1, 2, 3, 4, 5 et 7), Adenovirus (types 40 and 41), Aerococcus spp., Aeromonas hydrophila, Ancylostoma duodenale, Angiostrongylus cantonensis, Ascaris lumbricoides, Ascaris spp., Aspergillus spp., Babesia spp, B. microti, Bacillus anthracis, Bacillus cereus, Bacteroides spp., Balantidium coli, Bartonella bacilliformis, Blastomyces dermatitidis, Bluetongue virus, Bordetella bronchiseptica, Bordetella pertussis, Borrelia afzelii, Borrelia burgdorferi, Borrelia garinii, Brachamella catarrhalis, Brucella spp. (B. abortus, B. canis, B. melitensis, B. suis), Brugia spp., Burkholderia, (Pseudomonas) mallei, Burkholderia (Pseudomonas) pseudomallei, California serogroup, Campylobacter fetus subsp. Fetus, Campylobacter jejuni, C. coli, C. fetus subsp. Jejuni, Candida albicans, Capnocytophaga spp., Chikungunya virus, Chlamydia psittaci, Chlamydia trachomatis, Citrobacter spp., Clonorchis sinensis, Clostridium botulinum, Clostridium difficile, Clostridium perfringens, Clostridium tetani, Clostridium spp. (with the exception of those species listed above), Coccidioides immitis, Colorado tick fever virus, Corynebacterium diphtheriae, Coxiella burnetii, Coxsackievirus, Creutzfeldt-Jakob agent, Kuru agent, Crimean-Congo hemorrhagic fever virus, Cryptococcus neoformans, Cryptosporidium parvum, Cytomegalovirus, Cyclospora cayetanensis, Dengue virus (1, 2, 3, 4), Diphtheroids, Eastern (Western) equine encephalitis virus, Ebola virus, Echinococcus granulosus, Echinococcus multilocularis, Echovirus, Edwardsiella tarda, Entamoeba histolytica, Enterobacter spp., Enterovirus 70, Epidermophyton floccosum, Ehrlichia spp., Ehrlichia sennetsu, Microsporum spp., Trichophyton spp., Epstein-Barr virus, Escherichia coli, enterohemorrhagic, Escherichia coli, enteroinvasive, Escherichia coli, enteropathogenic, Escherichia coli, enterotoxigenic, Fasciola hepatica, Francisella tularensis, Fusobacterium spp., Gemella haemolysans, Giardia lamblia, Guanarito virus, Haemophilus ducreyi, Haemophilus influenzae (group b), Hantavirus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Hepatitis D virus, Hepatitis E virus, Herpes simplex virus, Herpesvirus simiae, Histoplasma capsulatum, Human coronavirus, Human immunodeficiency virus, Human papillomavirus, Human rotavirus, Human T-lymphotrophic virus, Influenza virus including H5N1, Junin virus/Machupo virus, Klebsiella spp., Kyasanur Forest disease virus, Lactobacillus spp.,

Lassa virus, Legionella pneumophila, Leishmania major, Leishmania infantum, Leishmania spp., Leptospira interrogans, Listeria monocytogenes, Lymphocytic choriomeningitis virus, Machupo virus, Marburg virus, Measles virus, Micrococcus spp., Moraxella spp., Mycobacterium spp. (other than *M. bovis*, *M. tuberculosis*, *M. avium*, *M. leprae*), Mycobacterium tuberculosis, *M. bovis*, Mycoplasma hominis, *M. orale*, *M. salivarium*, *M. fermentans*, Mycoplasma pneumoniae, *Naegleria fowleri*, *Necator americanus*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Neisseria* spp. (other than *N. gonorrhoeae* and *N. meningitidis*), *Nocardia* spp., Norwalk virus, Omsk hemorrhagic fever virus, *Onchocerca volvulus*, *Opisthorchis* spp., *Parvovirus B19*, *Pasteurella* spp., *Peptococcus* spp., *Peptostreptococcus* spp., *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium* spp., *Plesiomonas shigelloides*, Powassan encephalitis virus, *Proteus* spp., *Pseudomonas* spp. (other than *P. mallei*, *P. pseudomallei*), Rabies virus, Respiratory syncytial virus, Rhinovirus, *Rickettsia akari*, *Rickettsia prowazekii*, *R. Canada*, *Rickettsia rickettsii*, Rift Valley virus, Ross river virus/O'Nyong-Nyong virus, Rubella virus, *Salmonella choleraesuis*, *Salmonella paratyphi*, *Salmonella typhi*, *Salmonella* spp. (with the exception of those species listed above), *Schistosoma* spp., Scrapie agent, *Serratia* spp., *Shigella* spp., *Sindbis* virus, *Sporothrix schenckii*, *St. Louis* encephalitis virus, Murray Valley encephalitis virus, *Staphylococcus aureus*, *Streptobacillus moniliformis*, *Streptococcus agalactiae*, *Streptococcus faecalis*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus salivarius*, *Taenia saginata*, *Taenia solium*, *Toxocara canis*, *T. cati*, *T. cruzi*, *Toxoplasma gondii*, *Treponema pallidum*, *Trichinella* spp., *Trichomonas vaginalis*, *Trichuris trichiura*, *Trypanosoma brucei*, *Trypanosoma cruzi*, *Ureaplasma urealyticum*, *Vaccinia* virus, *Varicella-zoster* virus, eastern equine encephalitis virus (EEEV), severe acute respiratory virus (SARS), Venezuelan equine encephalitis virus (VEEV), Vesicular stomatitis virus, *Vibrio cholerae*, serovar 01, *Vibrio parahaemolyticus*, West Nile virus, *Wuchereria bancrofti*, Yellow fever virus, *Yersinia enterocolitica*, *Yersinia pseudotuberculosis*, and *Yersinia pestis*. Target antigens may include proteins, or variants or fragments thereof, produced by any of the infectious organisms.

[00237] A number of viruses are associated with viral hemorrhagic fever, including filoviruses (e.g., Ebola, Marburg, and Reston), arenaviruses (e.g. Lassa, Junin, and Machupo), and bunyaviruses. In addition, phleboviruses, including, for example, Rift Valley fever virus, have been identified as etiologic agents of viral hemorrhagic fever. Etiological agents of hemorrhagic fever and associated inflammation may also include paramyxoviruses, particularly respiratory syncytial virus. In addition, other viruses causing hemorrhagic fevers in man have been identified as belonging to the following virus groups: togavirus (Chikungunya), flavivirus (dengue, yellow fever, Kyasanur Forest disease, Omsk hemorrhagic fever), nairovirus (Crimian-Congo hemorrhagic fever) and hantavirus (hemorrhagic fever with renal syndrome, nephropathic epidemia). Furthermore, Sin Nombre virus was identified as the etiologic agent of the 1993 outbreak of hantavirus pulmonary syndrome in the American Southwest.

[00238] Target antigens may include viral coat proteins, i.e., influenza neuraminidase and hemagglutinin, HIV gp160 or derivatives thereof, HIV Gag, HIV Nef, HIV Pol, SARS coat proteins, herpes virion proteins, WNV proteins, etc. Target antigens may also include bacterial surface proteins including pneumococcal PsaA, PspA, LytA, surface or virulence associated proteins of bacterial pathogens such as *Nisseria gonorrhoea*, outer membrane proteins or surface proteases.

HPV Antigen Targets

[00239] Target antigens may also include proteins, or variants or fragments thereof, associated with human papillomavirus (HPV), such as oncoproteins E6 and E7. In certain embodiments, the oncoprotein may be modified to produce a non-oncogenic variant or a variant having reduced oncogenicity relative to the wild type protein. For example, the portion of the peptide that is responsible for binding a tumor suppressor protein (e.g., p53 and pRb) may be deleted or modified so that it no longer interacts with the tumor suppressor protein. As another example, an oncoprotein that is a kinase, such as Her2/neu, may be kinase-inactivated, e.g., by point mutation. In some instances, two or more target antigens may be used during immunization. For example, the E6 and E7 antigens can be combined in a fusion protein, or separate vectors containing heterologous nucleotides encoding the modified or unmodified E6 and E7 target antigens are used in combination. For example, an Ad5-E6 vector can be administered with an Ad5-E7 vector. In this example, the Ad5-E6 vector and Ad5-E7 vector may be administered simultaneously or they may be administered sequentially.

[00240] High-risk human papillomavirus (HPV) such as HPV type-16 is associated with the etiology of cervical and more than 90% of HPV-related head and neck squamous cell carcinomas. Preventive vaccines such as human papillomavirus bivalent [Types 16 and 18] vaccine and recombinant human papillomavirus quadrivalent [Types 6, 11, 16, and 18] vaccine can be a primary defense against HPV associated cancers by preventing infection with the virus but reports indicate that they are not effective for active immunotherapy of established disease. The HPV early 6 (E6) and early 7 (E7) genes are expressed at high levels in HPV-induced cancers and are involved in the immortalization of primary human epidermal cells. Thus, these are ideal targets for tumor-specific immunotherapy because unlike many other tumor-associated antigens these viral antigens are “non-self” and thus do not have the potential to induce autoimmunity.

[00241] Disclosed herein is an immunotherapy against human papilloma virus (HPV) using a viral gene delivery platform to immunize against HPV 16 genes E6 and E7 (Ad5 [E1-, E2b-]-E6/E7) combined with programmed death-ligand 1 (PD1) blockade. Also disclosed herein is an immunotherapeutic comprised of a gene delivery vehicle (Ad5 [E1-, E2b-]) carrying modified genes for HPV type-16 E6 and E7. The HPV E6 and E7 genes can be modified to render them non-oncogenic while retaining the antigenicity necessary to produce an immune response against HPV induced tumors. The modified genes can lack the capacity to degrade p53, pRb, and PTPN13. The modified genes can be incorporated into a vaccine (Ad5

[E1-, E2b-]-E6/E7). The Ad5 [E1-, E2b-]-E6/E7 vaccine can retain the ability to induce an HPV-specific cell-mediated immune (CMI) response and can synergize with standard clinical therapy, enhancing immune-mediated clearance of an HPV-E6/E7 expressing tumor.

[00242] A balance between activation and inhibitory signals regulates the interaction between T lymphocytes and tumor cells, wherein T cell responses are initiated through antigen recognition by T-cell receptors (TCRs). In some cases, when combined with chemotherapy/radiation treatment in HPV-E6/E7 expressing tumor bearing mice, immunotherapy treatment with Ad5 [E1-, E2b-]-E6/E7 can result in significant improvement in overall survival as compared to subjects that receive chemotherapy/radiation alone.

Personalized Tumor Associated Antigens

[00243] In certain embodiments tumor associated antigens used with the compositions and methods of the invention may be identified directly from an individual with a proliferative disease or cancer. In certain embodiments, cancers may include benign tumors, metastatic tumors, carcinomas, or sarcomas and the like. In some embodiments, a personalized tumor antigen may comprise MUC1, MUC1c, MUC1n, T, or CEA characterized from a patient and further utilized as the target antigen as a whole, in part or as a variant.

[00244] In this regard, screens can be carried out using a variety of known technologies to identify tumor target antigens from an individual. For example, in one embodiment, a tumor biopsy is taken from a patient, RNA is isolated from the tumor cells and screened using a gene chip (for example, from Affymetrix, Santa Clara, Calif.) and a tumor antigen is identified. Once the tumor target antigen is identified, it may then be cloned, expressed and purified using techniques known in the art.

[00245] This target antigen can then be linked to one or more epitopes or incorporated or linked to cassettes or viral vectors described herein and administered to the patient in order to alter the immune response to the target molecule isolated from the tumor. In this manner, “personalized” immunotherapy and vaccines are contemplated within the context of the invention. Where cancer is genetic, that is inherited, for example the patient has been identified to have a BRAC1 or BRAC2 mutation, the vaccine can be used prophylactically. When the cancer is sporadic the immunotherapy can be used to reduce the size of the tumor, enhance overall and reduce reoccurrence of the cancer in a subject.

Combination Immunotherapies and Vaccines

[00246] The present disclosure provides for a combination immunotherapy and vaccine compositions for the treatment of cancer and infectious diseases. In some aspects, combination immunotherapies and vaccines provided herein can comprise a multi-targeted immunotherapeutic approach against antigens associated with the development of cancer such as tumor associated antigen, (TAA) or antigens known to be involved in a particular infectious disease, such as infectious disease associated antigen (IDAA). In some aspects, combination immunotherapies and vaccines provided herein can comprise a multi-targeted

antigen signature immunotherapeutic approach against antigens associated with the development of cancer or infectious disease. The compositions and methods of the invention, in various embodiments, provide viral based vectors expressing a variant of MUC1, MUC1c, MUC1n, T, and/or CEA for immunization of a disease, as provided herein. These vectors can raise an immune response against MUC1, MUC1c, MUC1n, T, and/or CEA.

[00247] In some aspects, the vector comprises at least one antigen. In some aspects, the vector comprises at least two antigens. In some aspects, the vector comprises at least three antigens. In some aspects, the vector comprises more than three antigens. In some aspects, the vaccine formulation comprises 1:1 ratio of vector to antigen. In some aspects, the vaccine comprises 1:2 ratio of vector to antigen. In some aspects, the vaccine comprises 1:3 ratio of vector to antigen. In some aspects, the vaccine comprises 1:4 ratio of vector to antigen. In some aspects, the vaccine comprises 1:5 ratio of vector to antigen. In some aspects, the vaccine comprises 1:6 ratio of vector to antigen. In some aspects, the vaccine comprises 1:7 ratio of vector to antigen. In some aspects, the vaccine comprises 1:8 ratio of vector to antigen. In some aspects, the vaccine comprises 1:9 ratio of vector to antigen. In some aspects, the vaccine comprises 1:10 ratio of vector to antigen.

[00248] In some aspects, the vaccine is a combination vaccine, wherein the vaccine comprises at least two vectors each containing at least a single antigen. In some aspects the vaccine is a combination vaccine, wherein the vaccine comprises at least three vectors each containing at least a single antigen target. In some aspects the vaccine is a combination vaccine, wherein the vaccine comprises more than three vectors each containing at least a single antigen.

[00249] In some aspects, the vaccine is a combination vaccine, wherein the vaccine comprises at least two vectors, wherein a first vector of the at least two vectors comprises at least a single antigen and wherein a second vector of the at least two vectors comprises at least two antigens. In some aspects, the vaccine is a combination vaccine, wherein the vaccine comprises at least three vectors, wherein a first vector of the at least three vectors comprises at least a single antigen and wherein a second vector of the at least three vectors comprises at least two antigens. In some aspects, the vaccine is a combination vaccine, wherein the vaccine comprises three or more vectors, wherein a first vector of the three or more vectors comprises at least a single antigen and wherein a second vector of the three or more vectors comprises at least two antigens. In some aspects the vaccine is a combination vaccine, wherein the vaccine comprises more than three vectors each containing at least two antigens.

[00250] When a mixture of different antigens are simultaneously administered or expressed from a same or different vector in an individual, they may compete with one another. As a result the formulations comprising different concentration and ratios of expressed antigens in a combination immunotherapy or vaccine must be evaluated and tailored to the individual or group of individuals to ensure that effective and sustained immune responses occur after administration.

[00251] Composition that comprises multiple antigens can be present at various ratios. For example, formulations with more than vector can have various ratios. For example, immunotherapies or vaccines can have two different vectors in a stoichiometry of 1:1, 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:15, 1:20, 1:30, 2:1, 2:3, 2:4, 2:5, 2:6, 2:7, 2:8, 3:1, 3:3, 3:4, 3:5, 3:6, 3:7, 3:8, 3:1, 3:3, 3:4, 3:5, 3:6, 3:7, 3:8, 4:1, 4:3, 4:5, 4:6, 4:7, 4:8, 5:1, 5:3, 5:4, 5:6, 5:7, 5:8, 6:1, 6:3, 6:4, 6:5, 6:7, 6:8, 7:1, 7:3, 7:4, 7:5, 7:6, 7:8, 8:1, 8:3, 8:4, 8:5, 8:6, or 8:7. For example, immunotherapies or vaccines can have three different vectors in a stoichiometry of: 1:1:1, 1:2:1, 1:3:1, 1:4:1, 1:5:1, 1:6:1, 1:7:1, 1:8:1, 2:1:1, 2:3:1, 2:4:1, 2:5:1, 2:6:1, 2:7:1, 2:8:1, 3:1, 3:3:1, 3:4:1, 3:5:1, 3:6:1, 3:7:1, 3:8:1, 3:1:1, 3:3:1, 3:4:1, 3:5:1, 3:6:1, 3:7:1, 3:8:1, 4:1:1, 4:3:1, 4:4:1, 4:5:1, 4:6:1, 4:7:1, 4:8:1, 5:1:1, 5:3:1, 5:4:1, 5:5:1, 5:6:1, 5:7:1, 5:8:1, 6:1:1, 6:3:1, 6:4:1, 6:5:1, 6:6:1, 6:7:1, 6:8:1, 7:1:1, 7:3:1, 7:4:1, 7:5:1, 7:6:1, 7:7:1, 7:8:1, 8:1:1, 8:3:1, 8:4:1, 8:5:1, 8:6:1, 8:7:1, 8:8:1, 1:1:2, 1:2:2, 1:3:2, 1:4:2, 1:5:2, 1:6:2, 1:7:2, 1:8:2, 2:1:2, 2:3:2, 2:4:2, 2:5:2, 2:6:2, 2:7:2, 2:8:2, 3:1:2, 3:3:2, 3:4:2, 3:5:2, 3:6:2, 3:7:2, 3:8:2, 3:1:2, 3:3:2, 3:4:2, 3:5:2, 3:6:2, 3:7:2, 3:8:2, 4:1:2, 4:3:2, 4:4:2, 4:5:2, 4:6:2, 4:7:2, 4:8:2, 5:1:2, 5:3:2, 5:4:2, 5:5:2, 5:6:2, 5:7:2, 5:8:2, 6:1:2, 6:3:2, 6:4:2, 6:5:2, 6:6:2, 6:7:2, 6:8:2, 7:1:2, 7:3:2, 7:4:2, 7:5:2, 7:6:2, 7:7:2, 7:8:2, 8:1:2, 8:3:2, 8:4:2, 8:5:2, 8:6:2, 8:7:2, 8:8:2, 1:1:3, 1:2:3, 1:3:3, 1:4:3, 1:5:3, 1:6:3, 1:7:3, 1:8:3, 2:1:3, 2:3:3, 2:4:3, 2:5:3, 2:6:3, 2:7:3, 2:8:3, 3:1:3, 3:3:3, 3:4:3, 3:5:3, 3:6:3, 3:7:3, 3:8:3, 3:1:3, 3:3:3, 3:4:3, 3:5:3, 3:6:3, 3:7:3, 3:8:3, 4:1:3, 4:3:3, 4:4:3, 4:5:3, 4:6:3, 4:7:3, 4:8:3, 5:1:3, 5:3:3, 5:4:3, 5:5:3, 5:6:3, 5:7:3, 5:8:3, 6:1:3, 6:3:3, 6:4:3, 6:5:3, 6:6:3, 6:7:3, 6:8:3, 7:1:3, 7:3:3, 7:4:3, 7:5:3, 7:6:3, 7:7:3, 7:8:3, 8:1:3, 8:3:3, 8:4:3, 8:5:3, 8:6:3, 8:7:3, 8:8:3, 1:1:4, 1:2:4, 1:3:4, 1:4:4, 1:5:4, 1:6:4, 1:7:4, 1:8:4, 2:1:4, 2:3:4, 2:4:4, 2:5:4, 2:6:4, 2:7:4, 2:8:4, 3:1:4, 3:3:4, 3:4:4, 3:5:4, 3:6:4, 3:7:4, 3:8:4, 4:1:4, 4:3:4, 4:4:4, 4:5:4, 4:6:4, 4:7:4, 4:8:4, 5:1:4, 5:3:4, 5:4:4, 5:5:4, 5:6:4, 5:7:4, 5:8:4, 6:1:4, 6:3:4, 6:4:4, 6:5:4, 6:6:4, 6:7:4, 6:8:4, 7:1:4, 7:3:4, 7:4:4, 7:5:4, 7:6:4, 7:7:4, 7:8:4, 8:1:4, 8:3:4, 8:4:3, 8:5:4, 8:6:4, 8:7:4, 8:8:4, 1:1:5, 1:2:5, 1:3:5, 1:4:5, 1:5:5, 1:6:5, 1:7:5, 1:8:5, 2:1:5, 2:3:5, 2:4:5, 2:5:5, 2:6:5, 2:7:5, 2:8:5, 3:1:5, 3:3:5, 3:4:5, 3:5:5, 3:6:5, 3:7:5, 3:8:5, 3:1:5, 3:3:5, 3:4:5, 3:5:5, 3:6:5, 3:7:5, 3:8:5, 4:1:5, 4:3:5, 4:4:5, 4:5:5, 4:6:5, 4:7:5, 4:8:5, 5:1:5, 5:3:5, 5:4:5, 5:5:5, 5:6:5, 5:7:5, 5:8:5, 6:1:5, 6:3:5, 6:4:5, 6:5:5, 6:6:5, 6:7:5, 6:8:5, 7:1:5, 7:3:5, 7:4:5, 7:5:5, 7:6:5, 7:7:5, 7:8:5, 8:1:5, 8:3:5, 8:4:5, 8:5:5, 8:6:5, 8:7:5, 8:8:5, 1:1:6, 1:2:6, 1:3:6, 1:4:6, 1:5:6, 1:6:6, 1:7:6, 1:8:6, 2:1:6, 2:3:6, 2:4:6, 2:5:6, 2:6:6, 2:7:6, 2:8:6, 3:1:6, 3:3:6, 3:4:6, 3:5:6, 3:6:6, 3:7:6, 3:8:6, 3:1:6, 3:3:6, 3:4:6, 3:5:6, 3:6:6, 3:7:6, 3:8:6, 4:1:6, 4:3:6, 4:4:6, 4:5:6, 4:6:6, 4:7:6, 4:8:6, 5:1:6, 5:3:6, 5:4:6, 5:5:6, 5:6:6, 5:7:6, 5:8:6, 6:1:6, 6:3:6, 6:4:6, 6:5:6, 6:6:6, 6:7:6, 6:8:6, 7:1:6, 7:3:6, 7:4:6, 7:5:6, 7:6:6, 7:7:6, 7:8:6, 8:1:6, 8:3:6, 8:4:6, 8:5:6, 8:6:5, 8:7:6, 8:8:6, 1:1:7, 1:2:7, 1:3:7, 1:4:7, 1:5:7, 1:6:7, 1:7:7, 1:8:7, 2:1:7, 2:3:7, 2:4:7, 2:5:7, 2:6:7, 2:7:7, 2:8:7, 3:1:7, 3:3:7, 3:4:7, 3:5:7, 3:6:7, 3:7:7, 3:8:7, 3:1:7, 3:3:7, 3:4:7, 3:5:7, 3:6:7, 3:7:7, 3:8:7, 4:1:7, 4:3:7, 4:4:7, 4:5:7, 4:6:7, 4:7:7, 4:8:7, 5:1:7, 5:3:7, 5:4:7, 5:5:7, 5:6:7, 5:7:7, 5:8:7, 6:1:7, 6:3:7, 6:4:7, 6:5:7, 6:6:7, 6:7:7, 6:8:7, 7:1:7, 7:3:7, 7:4:7, 7:5:7, 7:6:7, 7:7:7, 7:8:7, 8:1:7, 8:3:7, 8:4:7, 8:5:7, 8:6:5, 8:7:7, or 8:8:7.

[00252] In some aspects the present disclosure provides combination immunotherapies comprising multi-targeted immunotherapeutic directed TAAs. In some aspects the present disclosure provides combination immunotherapies comprising multi-targeted immunotherapeutic directed to IDAAs.

[00253] The present disclosure provides for a combination immunotherapies or vaccines comprising: at least two, at least three, or more than three different target antigens comprising a sequence encoding a modified CEA, MUC1c, and/or Brachyury. For example, a combination immunotherapy or vaccine can comprise at least two, at least three, or more than three different target antigens comprising a sequence encoding a modified CEA, MUC1c, and/or Brachyury, wherein the modified CEA comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, or 99.9% to SEQ ID NO:1 or SEQ ID NO:2; wherein the modified MUC1c comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, or 99.9% to SEQ ID NO:5 or SEQ ID NO:6; and wherein the modified Brachyury comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, 99.5%, 99.9% to SEQ ID NO:7 or SEQ ID NO:8. In some embodiments, the modified CEA comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% or 100% SEQ ID NO:1 or SEQ ID NO:2 and has a Asn->Asp substitution at position 610.

[00254] In some aspects the present disclosure provides combination immunotherapies comprising multi-targeted immunotherapeutic directed to TAAs and molecular compositions comprising an immune pathway checkpoint modulator that targets at least one immune checkpoint protein of the immune inhibitory pathway. In some aspects the present disclosure provides combination immunotherapies comprising multi-targeted immunotherapeutic directed to IDAAs and molecular compositions comprising an immune pathway checkpoint modulator that targets at least one immune checkpoint protein of the immune inhibitory pathway. The present disclosure provides for a combination immunotherapies or vaccines comprising: at least two, at least three, or more than three different target antigens comprising a sequence encoding a modified CEA, MUC1c, and/or Brachyury, and at least one molecular composition comprising an immune pathway checkpoint modulator. For example, a combination immunotherapy or vaccine can comprise at least two, at least three, or more than three different target antigens comprising a sequence encoding a modified CEA, MUC1c, and/or Brachyury, wherein the modified CEA comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, or 99.9% to SEQ ID NO:1 or SEQ ID NO:2; wherein the modified MUC1c comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.5%, or 99.9% to SEQ ID NO:5 or SEQ ID NO:6; wherein the modified Brachyury comprises a sequence with an identity value of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, 99.5%, 99.9% to SEQ ID NO:7 or SEQ ID NO:8, and at least one molecular composition comprising an immune pathway checkpoint modulator. In some embodiments, the modified CEA comprises a sequence with an identity value of at least 70%, 75%, 80%,

85%, 90%, 95%, 98%, 99%, 99.5%, 99.9% or 100% SEQ ID NO:1 or SEQ ID NO:2 and has a Asn->Asp substitution at position 610.

[00255] In some embodiments, the immune pathway checkpoint modulator targets CTLA4. In some embodiments, the immune pathway checkpoint modulator targets PD1. In some embodiments, the immune pathway checkpoint modulator targets PDL1. In some embodiments, the immune pathway checkpoint modulator targets LAG3. In some embodiments, the immune pathway checkpoint modulator targets B7-H3. In some embodiments, the immune pathway checkpoint modulator targets B7-H4. In some embodiments, the immune pathway checkpoint modulator targets TIM3. In some embodiment, the immune pathway checkpoint modulator is a monoclonal or polyclonal antibody directed to PD1, PDL1, PDL2, CD28, CD80, CD86, CTLA4, B7RP1, ICOS, B7RPI, B7-H3, B7-H4, BTLA, HVEM, KIR, TCR, LAG3, CD137, CD137L, OX40, OX40L, CD27, CD70, CD40, CD40L, TIM3 (*i.e.*, HAVcr2), GAL9, and A2aR.

[00256] In some embodiments, at least one of the recombinant nucleic acid vector is a replication defective adenovirus vector that comprises a replication defective adenovirus 5 vector comprising a first identity value. In some embodiments, the replication defective adenovirus vector comprises a deletion in the E2b region. In some embodiments, the replication defective adenovirus vector further comprises a deletion in the E1 region. In some embodiments, the first identity value is at least 90%. In some embodiments, the first identity value is at least 95%. In some embodiments, the first identity value is at least 99%. In some embodiments, the first identity value is 100%. In some embodiments, the first identity value is at least 90%. In some embodiments, the first identity value is at least 95%. In some embodiments, the first identity value is at least 99%. In some embodiments, the first identity value is 100%. In some embodiments, the first identity value is at least 90%. In some embodiments, the first identity value is at least 95%. In some embodiments, the first identity value is at least 99%. In some embodiments, the first identity value is 100%.

[00257] In some embodiments, at least one of the recombinant nucleic acid vector comprises a sequence with at least 80%, 85%, 90%, 95%, 99%, or 100% sequence identity to SEQ ID NO:3. In some embodiments, the recombinant nucleic acid vector comprises a region with at least 80%, 85%, 90%, 95%, 99%, or 100% sequence identity to a region in SEQ ID NO:3, wherein the region is selected from 26048-26177, 26063-26141, 1-103, 54-103, 32214-32315, and 32214-32262. In some embodiments, the recombinant nucleic acid vector further comprises a region encoding a peptide with at least 80%, 85%, 90%, 95%, 99%, or 100% sequence identity to a peptide encoded by a region in SEQ ID NO:3 between positions 1057 and 3165.

Immunological Fusion Partner Antigen Targets

[00258] The viral vectors of the present invention may also include nucleic acid sequences that encode proteins that increase the immunogenicity of the target antigen. In this regard, the protein produced

following immunization with the viral vector containing such a protein may be a fusion protein comprising the target antigen of interest fused to a protein that increases the immunogenicity of the target antigen of interest.

[00259] In one embodiment, such an immunological fusion partner is derived from a *Mycobacterium* sp., such as a *Mycobacterium tuberculosis*-derived Ra12 fragment. Ra12 compositions and methods for their use in enhancing the expression and/or immunogenicity of heterologous polynucleotide/polypeptide sequences are described in U.S. Patent Application 60/158,585 and U.S. Patent No. 7,009,042. Briefly, Ra12 refers to a polynucleotide region that is a subsequence of a *Mycobacterium tuberculosis* MTB32A nucleic acid. MTB32A is a serine protease of 32 kDa encoded by a gene in virulent and avirulent strains of *M. tuberculosis*. The nucleotide sequence and amino acid sequence of MTB32A have been described (see, e.g., U.S. Patent Application 60/158,585; Skeiky et al., *Infection and Immun.* 67:3998-4007 (1999)). C-terminal fragments of the MTB32A coding sequence express at high levels and remain as soluble polypeptides throughout the purification process. Moreover, Ra12 may enhance the immunogenicity of heterologous immunogenic polypeptides with which it is fused. One Ra12 fusion polypeptide comprises a 14 kDa C-terminal fragment corresponding to amino acid residues 192 to 323 of MTB32A. Other Ra12 polynucleotides generally comprise at least about 15, 30, 60, 100, 200, 300, or more nucleotides that encode a portion of a Ra12 polypeptide. Ra12 polynucleotides may comprise a native sequence (i.e., an endogenous sequence that encodes a Ra12 polypeptide or a portion thereof) or may comprise a variant of such a sequence. Ra12 polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions such that the biological activity of the encoded fusion polypeptide is not substantially diminished, relative to a fusion polypeptide comprising a native Ra12 polypeptide. Variants can have at least about 70%, 80%, or 90% identity, or more, to a polynucleotide sequence that encodes a native Ra12 polypeptide or a portion thereof.

[00260] An immunological fusion partner can be derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenzae* B. In some cases, a protein D derivative comprises approximately the first third of the protein (e.g., the first N-terminal 100-110 amino acids). A protein D derivative may be lipidated. Within certain embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes, which may increase the expression level in *E. coli* and may function as an expression enhancer. The lipid tail may ensure optimal presentation of the antigen to antigen presenting cells. Other fusion partners include the non-structural protein from influenza virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

[00261] The immunological fusion partner can be the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus can be employed. Within another embodiment, a repeat portion of LYTA may be incorporated into a fusion polypeptide. A repeat portion can, for example, be found in the C-terminal region starting at residue 178. One particular repeat portion incorporates residues 188-305.

[00262] In some embodiments, the antigen target comprises an immunogenic component comprising a cytokine selected from the group of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13. In some embodiments, the antigen target further comprises one or more immunogenic component comprises a cytokine selected from the group of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13. In some embodiments, the antigen target comprises an immunogenic component comprising a nucleic acid encoding of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13., a protein with substantial identity to of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13, and a nucleic acid encoding a protein with substantial identity to of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13.

[00263] In some embodiments, the antigen target is fused or linked to an immunogenic component comprising a cytokine selected from the group of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13. In some embodiments, the antigen target is co-expressed in a cell with an immunogenic component comprising a cytokine selected from the group of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13.

Immune Pathway Checkpoint Modulators

[00264] In some embodiments, compositions of the present invention are administered with one or more immune checkpoint modulators. A balance between activation and inhibitory signals regulates the interaction between T lymphocytes and disease cells, wherein T-cell responses are initiated through antigen recognition by the T-cell receptor (TCR). The inhibitory pathways and signals are referred to as immune checkpoints. In normal circumstances, immune checkpoints play a critical role in control and prevention of autoimmunity and also protect from tissue damage in response to pathogenic infection.

[00265] The present disclosure provides combination immunotherapies comprising viral vector based vaccines and compositions for modulating immune checkpoint inhibitory pathways for the treatment of cancer and infectious diseases. In some embodiments, modulating is increasing expression or activity of a

gene or protein. In some embodiments, modulating is decreasing expression or activity of a gene or protein. In some embodiments, modulating affects a family of genes or proteins.

[00266] In general, the immune inhibitory pathways are initiated by ligand-receptor interactions. It is now clear that in diseases, the disease can co-opt immune-checkpoint pathways as mechanism for inducing immune resistance in a subject.

[00267] The induction of immune resistance or immune inhibitory pathways in a subject by a given disease can be blocked by molecular compositions such as siRNAs, antisense, small molecules, mimic, a recombinant form of ligand, receptor or protein, or antibodies (which can be an Ig fusion protein) that are known to modulate one or more of the Immune Inhibitory Pathways. For example, preliminary clinical findings with blockers of immune-checkpoint proteins, such as Cytotoxic T-lymphocyte-associated antigen 4 (CTLA4) and programmed cell death protein 1 (PD1) have shown promise for enhancing antitumor immunity.

[00268] Because diseased cells can express multiple inhibitory ligands, and disease-infiltrating lymphocytes express multiple inhibitory receptors, dual or triple blockade of immune checkpoints proteins may enhance anti-disease immunity. Combination immunotherapies as provide herein can comprise one or more molecular compositions of the following immune-checkpoint proteins: PD1, PDL1, PDL2, CD28, CD80, CD86, CTLA4, B7RP1, ICOS, B7RPI, B7-H3 (also known as CD276), B7-H4 (also known as B7-S1, B7x and VCTN1), BTLA (also known as CD272), HVEM, KIR, TCR, LAG3 (also known as CD223), CD137, CD137L, OX40, OX40L, CD27, CD70, CD40, CD40L, TIM3 (also known as HAVcr2), GAL9, and A2aR. In some embodiments, the molecular composition comprises siRNAs. In some embodiments, the molecular composition comprises a small molecule. In some embodiments, the molecular composition comprises a recombinant form of a ligand. In some embodiments, the molecular composition comprises a recombinant form of a receptor. In some embodiments, the molecular composition comprises an antibody. In some embodiments, the combination therapy comprises more than one molecular composition and/or more than one type of molecular composition. As it will be appreciated by those in the art, future discovered proteins of the immune checkpoint inhibitory pathways are also envisioned to be encompassed by the present disclosure.

[00269] In some embodiments, combination immunotherapies comprise molecular compositions for the modulation of CTLA4. In some embodiments, combination immunotherapies comprise molecular compositions for the modulation PD1. In some embodiments, combination immunotherapies comprise molecular compositions for the modulation PDL1. In some embodiments, combination immunotherapies comprise molecular compositions for the modulation LAG3. In some embodiments, combination immunotherapies comprise molecular compositions for the modulation B7-H3. In some embodiments, combination immunotherapies comprise molecular compositions for the modulation B7-H4. In some embodiments, combination immunotherapies comprise molecular compositions for the modulation TIM3.

In some embodiments, modulation is an increase or enhancement of expression. In other embodiments, modulation is the decrease or absence of expression.

[00270] Two exemplary immune checkpoint inhibitors include the cytotoxic T lymphocyte associated antigen-4 (CTLA-4) and the programmed cell death protein-1 (PD1). CTLA-4 can be expressed exclusively on T-cells where it regulates early stages of T-cell activation. CTLA-4 interacts with the co-stimulatory T-cell receptor CD28 which can result in signaling that inhibits T-cell activity. Once TCR antigen recognition occurs, CD28 signaling may enhance TCR signaling, in some cases leading to activated T-cells and CTLA-4 inhibits the signaling activity of CD28. The present disclosure provides immunotherapies as provided herein in combination with anti-CTLA-4 monoclonal antibody for the treatment of proliferative disease and cancer. The present disclosure provides immunotherapies as provided herein in combination with CTLA-4 molecular compositions for the treatment of proliferative disease and cancer.

[00271] Programmed death cell protein ligand-1 (PDL1) is a member of the B7 family and is distributed in various tissues and cell types. PDL1 can interact with PD1 inhibiting T-cell activation and CTL mediated lysis. Significant expression of PDL1 has been demonstrated on various human tumors and PDL1 expression is one of the key mechanisms in which tumors evade host antitumor immune responses. Programmed death-ligand 1 (PDL1) and programmed cell death protein-1 (PD1) interact as immune checkpoints. This interaction can be a major tolerance mechanism which results in the blunting of anti-tumor immune responses and subsequent tumor progression. PD1 is present on activated T cells and PDL1, the primary ligand of PD1, is often expressed on tumor cells and antigen-presenting cells (APC) as well as other cells, including B cells. Significant expression of PDL1 has been demonstrated on various human tumors including HPV-associated head and neck cancers. PDL1 interacts with PD1 on T cells inhibiting T cell activation and cytotoxic T lymphocyte (CTL) mediated lysis. The present disclosure provides immunotherapies as provided herein in combination with anti-PD1 or anti-PDL1 monoclonal antibody for the treatment of proliferative disease and cancer. The present disclosure provides immunotherapies as provided herein in combination with PD1 or anti-PDL1 molecular compositions for the treatment of proliferative disease and cancer. The present disclosure provides immunotherapies as provided herein in combination with anti-CTLA-4 and anti-PD1 monoclonal antibodies for the treatment of proliferative disease and cancer. The present disclosure provides immunotherapies as provided herein in combination with anti-CTLA-4 and PDL1 monoclonal antibodies for the treatment of proliferative disease and cancer. The present disclosure provides immunotherapies as provided herein in combination with anti-CTLA-4, anti-PD1, PDL1, monoclonal antibodies, or a combination thereof, for the treatment of proliferative disease and cancer.

[00272] Provided herein are Ad5 [E1-, E2b-]-E6/E7 immunizations combined with PD1 blockade that can increase an anti-tumor effect. The inventors have characterized a CMI response induced by the Ad5

[E1-, E2b-]-E6/E7 vaccine and show kinetics of an anti-tumor response to evaluate the therapeutic potential of treating small versus large established tumors.

[00273] Immune checkpoint molecules can be expressed by T cells. Immune checkpoint molecules can effectively serve as "brakes" to down-modulate or inhibit an immune response. Immune checkpoint molecules include, but are not limited to Programmed Death 1 (PD1, also known as PDCD1 or CD279, accession number: NM_005018), Cytotoxic T-Lymphocyte Antigen 4 (CTLA-4, also known as CD152, GenBank accession number AF414120.1), LAG3 (also known as CD223, accession number: NM_002286.5), Tim3 (also known as HAVCR2, GenBank accession number: JX049979.1), BTLA (also known as CD272, accession number: NM_181780.3), BY55 (also known as CD160, GenBank accession number: CR541888.1), TIGIT (also known as IVSTM3, accession number: NM_173799), LAIR1 (also known as CD305, GenBank accession number: CR542051.1), SIGLECIO (GeneBank accession number: AY358337.1), 2B4 (also known as CD244, accession number: NM_001166664.1), PPP2CA, PPP2CB, PTPN6, PTPN22, CD96, CRTAM, SIGLEC7, SIGLEC9, TNFRSF10B, TNFRSF10A, CASP8, CASP10, CASP3, CASP6, CASP7, FADD, FAS, TGFBRII, TGFRBRI, SMAD2, SMAD3, SMAD4, SMAD10, SKI, SKIL, TGIF1, ILIORA, IL10RB, HMOX2, IL6R, IL6ST, EIF2AK4, CSK, PAG1, SIT1, FOXP3, PRDM1, BATF, GUCY1A2, GUCY1A3, GUCY1B2, GUCY1B3 which directly inhibit immune cells. For example, PD1 can be combined with an adenoviral vaccine of the present invention to treat a patient in need thereof. **Table 1**, without being exhaustive, shows exemplary immune checkpoint genes that can be inactivated to improve the efficiency of the adenoviral vaccine of the present invention. Immune checkpoints gene can be selected from such genes listed in **Table 1** and others involved in co-inhibitory receptor function, cell death, cytokine signaling, arginine tryptophan starvation, TCR signaling, Induced T-reg repression, transcription factors controlling exhaustion or anergy, and hypoxia mediated tolerance.

Table 1

Gene Symbol	NCBI # (GRCh38.p2)	Start	Stop	Genome location
ADORA2A	135	24423597	24442360	22q11.23
CD276	80381	73684281	73714518	15q23-q24
VTCN1	79679	117143587	117270368	1p13.1
BTLA	151888	112463966	112499702	3q13.2
CTLA4	1493	203867788	203873960	2q33
IDO1	3620	39913809	39928790	8p12-p11
KIR3DL1	3811	54816438	54830778	19q13.4
LAG3	3902	6772483	6778455	12p13.32
PDCD1	5133	241849881	241858908	2q37.3
HAVCR2	84868	157085832	157109237	5q33.3
VISTA	64115	71747556	71773580	10q22.1
CD244	51744	160830158	160862902	1q23.3
CISH	1154	50606454	50611831	3p21.3

[00274] The combination of an adenoviral-based vaccine and an immune pathway checkpoint modulator may result in reduction in cancer recurrences in treated patients, as compared to either agent alone. In yet another embodiment of this invention the combination of an adenoviral-based vaccine and an immune pathway checkpoint modulator may result in reduction in the presence or appearance of metastases or micro metastases in treated patients, as compared to either agent alone. In another embodiment, the combination of a adenoviral-based vaccine and an immune pathway checkpoint modulator may result improved overall survival of treated patients, as compared to either agent alone. In some cases, the combination of an adenoviral vaccine and an immune pathway checkpoint modulator may increase the frequency or intensity of tumor-specific T cell responses in patients compared to either agent alone.

[00275] The present invention also discloses the use of immune checkpoint inhibition to improve performance of an adenoviral vector-based vaccine. The immune checkpoint inhibition may be administered at the time of the vaccine. The immune checkpoint inhibition may also be administered after a vaccine. Immune checkpoint inhibition may occur simultaneously to an adenoviral vaccine administration. Immune checkpoint inhibition may occur 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, or 60 minutes after vaccination. Immune checkpoint inhibition may also occur 1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23, or 24 hours post vaccination. In some cases, immune inhibition may occur 1, 2, 3, 4, 5, 6, or 7 days after vaccination. Immune checkpoint inhibition may occur at any time before or after vaccination.

[00276] In another aspect, the invention pertains to a vaccine comprising an antigen and an immune pathway checkpoint modulator. The invention can pertain to a method for treating a subject having a condition that would benefit from downregulation of an immune checkpoint, PD1 for example, and its natural binding partner(s) on cells of the subject.

[00277] An immune pathway checkpoint modulator may be combined with an adenoviral vaccine comprising any antigen. For example, an antigen can be MUC1c, HER3, Brachyury, HER2NEU, CEA, or PSA. An immune pathway checkpoint modulator may produce a synergistic effect when combined with a vaccine. An immune pathway checkpoint modulator may also produce an additive effect when combined with a vaccine.

Formulations

[00278] The present invention provides pharmaceutical compositions comprising a vaccination regime that can be administered either alone or together with a pharmaceutically acceptable carrier or excipient, by any routes, and such administration can be carried out in both single and multiple dosages. More particularly, the pharmaceutical composition can be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hand candies, powders, sprays, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, such oral

pharmaceutical formulations can be suitably sweetened and/or flavored by means of various agents of the type commonly employed for such purposes. The compositions described throughout can be formulated into a pharmaceutical medicament and be used to treat a human or mammal, in need thereof, diagnosed with a disease, *e.g.*, cancer.

[00279] For administration, viral vector stock can be combined with an appropriate buffer, physiologically acceptable carrier, excipient or the like. In certain embodiments, an appropriate number of virus vector particles (VP) are administered in an appropriate buffer, such as, sterile PBS or saline. In certain embodiment's viral vector compositions disclosed herein are provided in specific formulations for subcutaneously, parenterally, intravenously, intramuscularly, or even intraperitoneally administration. In certain embodiments, formulations in a solution of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, squalene-based emulsion, Squalene-based oil-in-water emulsions, water-in-oil emulsions, oil-in-water emulsions, nonaqueous emulsions, water-in-paraffin oil emulsion, and mixtures thereof and in oils. In other embodiments, viral vectors may be provided in specific formulations for pill form administration by swallowing or by suppository.

[00280] Illustrative pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (see, *e.g.*, U.S. Pat. No. 5,466,468). Fluid forms to the extent that easy syringability exists may be preferred. Forms that are stable under the conditions of manufacture and storage are within the bounds of this invention. In various embodiments, forms are preserved against the contaminating action of microorganisms, such as bacteria, molds and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and/or by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and thimerosal. It may be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[00281] In one embodiment, for parenteral administration in an aqueous solution, the solution can be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art in light of the present disclosure. For example, one

dosage may be dissolved in 1 mL of isotonic NaCl solution and either added to 1000 mL of hypodermoclysis fluid or injected at the proposed site of infusion, (see, e.g., "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage may occur depending on the condition of the subject being treated.

[00282] Carriers of formulation can comprise any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

[00283] In certain embodiments, the viral vectors of the invention may be administered in conjunction with one or more immunostimulants, such as an adjuvant. An immunostimulant refers to essentially any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an antigen. One type of immunostimulant comprises an adjuvant. Many adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, Bortadella pertussis or *Mycobacterium tuberculosis* derived proteins. Certain adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories); Merck Adjuvant 65 (Merck and Company, Inc.) AS-2 (SmithKline Beecham); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF, IFN- γ , TNF α , IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and/or IL-13, and others, like growth factors, may also be used as adjuvants.

[00284] Within certain embodiments of the invention, the adjuvant composition can be one that induces an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (e.g., IFN- γ , TNF α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (e.g., IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient may support an immune response that includes Th1- and/or Th2-type responses. Within certain embodiments, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. Thus, various embodiments of the invention relate to therapies raising an immune response against a target antigen, for example MUC1, MUC1c, MUC1n, T, or CEA, using cytokines, e.g. IFN- γ , TNF α , IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and/or IL-13 supplied concurrently with a replication defective viral vector treatment. In some

embodiments, a cytokine or a nucleic acid encoding a cytokine, is administered together with a replication defective viral described herein. In some embodiments, cytokine administration is performed prior or subsequent to viral vector administration. In some embodiments, a replication defective viral vector capable of raising an immune response against a target antigen, for example MUC1, MUC1c, MUC1n, T, and/or CEA, further comprises a sequence encoding a cytokine.

[00285] Certain illustrative adjuvants for eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, such as 3-de-O-acylated monophosphoryl lipid A, together with an aluminum salt. MPL® adjuvants are commercially available (*see, e.g.*, U.S. Pat. Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. (*see, e.g.*, WO 96/02555, WO 99/33488 and U.S. Pat. Nos. 6,008,200 and 5,856,462). Immunostimulatory DNA sequences can also be used. Another adjuvant for use in the present invention comprises a saponin, such as Quil A, or derivatives thereof, including QS21 and QS7 (Aquila Biopharmaceuticals Inc.), Escin; Digitonin; or Gypsophila or Chenopodium quinoa saponins. Other formulations may include more than one saponin in the adjuvant combinations of the present invention, e.g., combinations of at least two of the following group comprising QS21, QS7, Quil A, β -escin, or digitonin.

[00286] In some embodiments, the compositions may be delivered by intranasal sprays, inhalation, and/or other aerosol delivery vehicles. The delivery of drugs using intranasal microparticle resins and lysophosphatidyl-glycerol compounds can be employed (*see, e.g.*, U.S. Pat. No. 5,725,871). Likewise, illustrative transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix can be employed (*see, e.g.*, U.S. Pat. No. 5,780,045).

[00287] Liposomes, nanocapsules, microparticles, lipid particles, vesicles, and the like, can be used for the introduction of the compositions of the present invention into suitable host cells/organisms. Compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like. Alternatively, compositions of the present invention can be bound, either covalently or non-covalently, to the surface of such carrier vehicles. Liposomes can be used effectively to introduce genes, various drugs, radiotherapeutic agents, enzymes, viruses, transcription factors, allosteric effectors and the like, into a variety of cultured cell lines and animals. Furthermore, the use of liposomes does not appear to be associated with autoimmune responses or unacceptable toxicity after systemic delivery. In some embodiments, liposomes are formed from phospholipids dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (i.e. multilamellar vesicles (MLVs)).

[00288] In some embodiments, the invention provides for pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds

in a stable and reproducible way. To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μm) may be designed using polymers able to be degraded *in vivo*.

[00289] The compositions described throughout can comprise or be administered with a chemotherapeutic agent (e.g., a chemical compound useful in the treatment of cancer). Chemotherapeutic cancer agents that can be used in combination with the disclosed T cell include, but are not limited to, mitotic inhibitors (vinca alkaloids), such as vincristine, vinblastine, vindesine and NavelbineTM (vinorelbine,5'-noranhydroblastine); topoisomerase I inhibitors, such as camptothecin compounds (e.g., CamptosarTM (irinotecan HCL), HycamtinTM (topotecan HCL) and other compounds derived from camptothecin and its analogues); podophyllotoxin derivatives, such as etoposide, teniposide and mitopodazole; alkylating agents such as cisplatin, cyclophosphamide, nitrogen mustard, trimethylene thiophosphoramide, carmustine, busulfan, chlorambucil, belustine, uracil mustard, chlomaphazin, and dacarbazine; antimetabolites such as cytosine arabinoside, fluorouracil, methotrexate, mercaptourine, azathioprine, and procarbazine; antibiotics, such as doxorubicin, bleomycin, dactinomycin, daunorubicin, mithramycin, mitomycin, mytomycin C, and daunomycin; anti-tumor antibodies; dacarbazine; azacytidine; amsacrine; melphalan; ifosfamide; and mitoxantrone.

[00290] Compositions disclosed herein can be administered in combination with other anti-tumor agents, including cytotoxic/antineoplastic agents and anti-angiogenic agents. Cytotoxic/anti-neoplastic agents can be defined as agents who attack and kill cancer cells. Some cytotoxic/anti-neoplastic agents can be alkylating agents, which alkylate the genetic material in tumor cells, *e.g.*, cis-platin, cyclophosphamide, nitrogen mustard, trimethylene thiophosphoramide, carmustine, busulfan, chlorambucil, belustine, uracil mustard, chlomaphazin, and dacabazine. Other cytotoxic/anti-neoplastic agents can be antimetabolites for tumor cells, *e.g.*, cytosine arabinoside, fluorouracil, methotrexate, mercaptourine, azathioprine, and procarbazine. Other cytotoxic/anti-neoplastic agents can be antibiotics, *e.g.*, doxorubicin, bleomycin, dactinomycin, daunorubicin, mithramycin, mitomycin, mytomycin C, and daunomycin. There are numerous liposomal formulations commercially available for these compounds. Still other cytotoxic/anti-neoplastic agents can be mitotic inhibitors (vinca alkaloids). These include vincristine, vinblastine and etoposide. Miscellaneous cytotoxic/anti-neoplastic agents include taxol and its derivatives, L-asparaginase, anti-tumor antibodies, dacarbazine, azacytidine, amsacrine, melphalan, VM-26, ifosfamide, mitoxantrone, and vindesine.

[00291] Anti-angiogenic agents can also be used. Suitable anti-angiogenic agents for use in the disclosed methods and compositions include anti-VEGF antibodies, including humanized and chimeric antibodies, anti-VEGF aptamers and antisense oligonucleotides. Other inhibitors of angiogenesis include angiostatin, endostatin, interferons, interleukin 1 (including α and β) interleukin 12, retinoic acid, and tissue inhibitors of metalloproteinase-1 and -2. (TIMP-1 and -2). Small molecules, including topoisomerases such as razoxane, a topoisomerase II inhibitor with anti-angiogenic activity, can also be used.

[00292] Other anti-cancer agents that can be used in combination with the disclosed invention include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; avastin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflorenthine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulazole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofosine; interleukin II (including recombinant interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1; interferon alfa-n3; interferon beta-I a; interferon gamma-I b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprolol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tegafuran; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprime; thioguanine; thioteplatin; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride. Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene;

adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplastic; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstaurosporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentanthraquinones; cycloplatam; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; daclizimab; decitabine; dehydrodidemnin B; deslorelin; dexamethasone; dexifosfamide; dextrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; manostatin A; marimastat; masoprolac; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone;

meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaumomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentrozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; ralitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrophostins; UBC inhibitors; ubenimex; urogenital sinus-derived

growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer. In one embodiment, the anti-cancer drug is 5-fluorouracil, taxol, or leucovorin.

METHODS

[00293] Compositions and methods of the invention, in various embodiments, take advantage of human cytolytic T-cells (CTLs), such as those that recognize MUC1, T, or CEA epitopes which bind to selected MHC molecules, e.g. HLA- A2, A3, and A24. Individuals expressing MHC molecules of certain serotypes, e.g. HLA- A2, A3, and A24 may be selected for therapy using the methods and compositions of the invention. For example, individuals expressing MHC molecules of certain serotypes, e.g. HLA- A2, A3, and A24, may be selected for a therapy including raising an immune response against MUC1, T, or CEA, using the methods and compositions described herein.

[00294] In various embodiments, these T-cells can be generated by in vitro cultures using antigen-presenting cells pulsed with the epitope of interest to stimulate peripheral blood mononuclear cells. In addition, T-cell lines can also be generated after stimulation with MUC1, T, or CEA latex beads, MUC1, T, or CEA protein-pulsed plastic adherent peripheral blood mononuclear cells, or DCs sensitized with MUC1, T, or CEA RNA. T-cells can also be generated from patients immunized with a vaccine vector encoding MUC1, T, or CEA immunogen. HLA A2-presented peptides from MUC1, T, or CEA can further be found in primary gastrointestinal tumors. In various embodiments, the invention relates to an HLA A2 restricted epitope of MUC1, T, or CEA, CAP-1, a nine amino acid sequence (YLSGANLNL; SEQ. ID. NO.:4), with ability to stimulate CTLs from cancer patients immunized with vaccine-MUC1, T, or CEA. Cap-1(6D) (YLSGADLNL; SEQ. ID. NO.:10) is a peptide analog of CAP-1. Its sequence includes a heteroclitic (nonanchor position) mutation, resulting in an amino acid change from Asn to Asp, enhancing recognition by the T-cell receptor. The Asn to Asp mutation appears to not cause any change in the binding of the peptide to HLA A2. Compared with the non-mutated CAP-1 epitope, Cap-1(6D) can enhance the sensitization of CTLs by 100 to 1,000 times. CTL lines can be elicited from peripheral blood mononuclear cells of healthy volunteers by in vitro sensitization to the Cap-1(6D) peptide, but not significantly to the CAP-1 peptide. These cell lines can lyse human tumor cells expressing endogenous CEA. Thus, polypeptide sequences comprising CAP-1 or CAP-1(6D), nucleic acid sequences encoding such sequences, an adenovirus vectors; for example replication defective adenovirus vectors, comprising such nucleic acid sequences are within the bounds of the invention.

Methods of Treatment

[00295] The adenovirus vectors of the present invention can be used in a number of vaccine settings for generating an immune response against one or more target antigens as described herein. The present invention provides methods of generating an immune response against any target antigen, such as those

described elsewhere herein. The adenovirus vectors are of particular importance because of the unexpected finding that they can be used to generate immune responses in subjects who have preexisting immunity to Ad and can be used in vaccination regimens that include multiple rounds of immunization using the adenovirus vectors, regimens not possible using previous generation adenovirus vectors.

[00296] Generally, generating an immune response comprises an induction of a humoral response and/or a cell-mediated response. It may be desirable to increase an immune response against a target antigen of interest. Generating an immune response may involve a decrease in the activity and/or number of certain cells of the immune system or a decrease in the level and/or activity of certain cytokines or other effector molecules. A variety of methods for detecting alterations in an immune response (e.g., cell numbers, cytokine expression, cell activity) are known in the art and are useful in the context of the instant invention. Illustrative methods useful in this context include intracellular cytokine staining (ICS), ELISpot, proliferation assays, cytotoxic T-cell assays including chromium release or equivalent assays, and gene expression analysis using any number of polymerase chain reaction (PCR) or RT-PCR based assays.

[00297] Generating an immune response can comprise an increase in target antigen-specific CTL activity of between 1.5 and 5 fold in a subject administered the adenovirus vectors of the invention as compared to a control. In another embodiment, generating an immune response comprises an increase in target-specific CTL activity of about 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 11.5, 12, 12.5, 15, 16, 17, 18, 19, 20, or more fold in a subject administered the adenovirus vectors as compared to a control.

[00298] Generating an immune response can comprise an increase in target antigen-specific HTL activity, such as proliferation of helper T-cells, of between 1.5 and 5 fold in a subject administered the adenovirus vectors of the invention that comprise nucleic acid encoding the target antigen as compared to an appropriate control. In another embodiment, generating an immune response comprises an increase in target-specific HTL activity of about 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 11.5, 12, 12.5, 15, 16, 17, 18, 19, 20, or more fold as compared to a control. In this context, HTL activity may comprise an increase as described above, or decrease, in production of a particular cytokine, such as interferon- γ (IFN- γ), interleukin-1 (IL-1), IL-2, IL-3, IL-6, IL-7, IL-12, IL-15, tumor necrosis factor- α (TNF- α), granulocyte macrophage colony-stimulating factor (GM-CSF), granulocyte-colony stimulating factor (G-CSF), or other cytokine. In this regard, generating an immune response may comprise a shift from a Th2 type response to a Th1 type response or in certain embodiments a shift from a Th1 type response to a Th2 type response. In other embodiments, generating an immune response may comprise the stimulation of a predominantly Th1 or a Th2 type response.

[00299] Generating an immune response can comprise an increase in target-specific antibody production of between 1.5 and 5 fold in a subject administered the adenovirus vectors of the present

invention as compared to an appropriate control. In another embodiment, generating an immune response comprises an increase in target-specific antibody production of about 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 11.5, 12, 12.5, 15, 16, 17, 18, 19, 20, or more fold in a subject administered the adenovirus vector as compared to a control.

[00300] Thus the present invention provides methods for generating an immune response against a target antigen of interest comprising administering to the individual an adenovirus vector comprising: a) a replication defective adenovirus vector, wherein the adenovirus vector has a deletion in the E2b region, and b) a nucleic acid encoding the target antigen; and readministering the adenovirus vector at least once to the individual; thereby generating an immune response against the target antigen. In certain embodiments, the present invention provides methods wherein the vector administered is not a gutted vector. In particular embodiments, the target antigen may be a wild-type protein, a fragment, a variant, or a variant fragment thereof. In some embodiments, the target antigen comprises MUC1, MUC1c, MUC1n, T, or CEA, a fragment, a variant, or a variant fragment thereof.

[00301] In a further embodiment, the present invention provides methods for generating an immune response against a target antigen in an individual, wherein the individual has preexisting immunity to Ad, by administering to the individual an adenovirus vector comprising: a) a replication defective adenovirus vector, wherein the adenovirus vector has a deletion in the E2b region, and b) a nucleic acid encoding the target antigen; and readministering the adenovirus vector at least once to the individual; thereby generating an immune response against the target antigen. In particular embodiments, the target antigen may be a wild-type protein, a fragment, a variant, or a variant fragment thereof. In some embodiments, the target antigen comprises MUC1, MUC1c, MUC1n, T, or CEA, a fragment, a variant, or a variant fragment thereof.

[00302] With regard to preexisting immunity to Ad, this can be determined using methods known in the art, such as antibody-based assays to test for the presence of Ad antibodies. Further, in certain embodiments, the methods of the present invention include first determining that an individual has preexisting immunity to Ad then administering the E2b deleted adenovirus vectors of the invention as described herein.

[00303] One embodiment of the invention provides a method of generating an immune response against one or more target antigens in an individual comprising administering to the individual a first adenovirus vector comprising a replication defective adenovirus vector, wherein the adenovirus vector has a deletion in the E2b region, and a nucleic acid encoding at least one target antigen; administering to the individual a second adenovirus vector comprising a replication defective adenovirus vector, wherein the adenovirus vector has a deletion in the E2b region, and a nucleic acid encoding at least one target antigen, wherein the at least one target antigen of the second adenovirus vector is the same or different from the at least one target antigen of the first adenovirus vector. In particular embodiments, the target antigen may be a

wild-type protein, a fragment, a variant, or a variant fragment thereof. In some embodiments, the target antigen comprises MUC1, MUC1c, MUC1n, T, or CEA, a fragment, a variant, or a variant fragment thereof.

[00304] Thus, the present invention contemplates multiple immunizations with the same E2b deleted adenovirus vector or multiple immunizations with different E2b deleted adenovirus vectors. In each case, the adenovirus vectors may comprise nucleic acid sequences that encode one or more target antigens as described elsewhere herein. In certain embodiments, the methods comprise multiple immunizations with an E2b deleted adenovirus encoding one target antigen, and re-administration of the same adenovirus vector multiple times, thereby inducing an immune response against the target antigen. In some embodiments, the target antigen comprises MUC1, MUC1c, MUC1n, T, or CEA, a fragment, a variant, or a variant fragment thereof.

[00305] In a further embodiment, the methods comprise immunization with a first adenovirus vector that encodes one or more target antigens, and then administration with a second adenovirus vector that encodes one or more target antigens that may be the same or different from those antigens encoded by the first adenovirus vector. In this regard, one of the encoded target antigens may be different or all of the encoded antigens may be different, or some may be the same and some may be different. Further, in certain embodiments, the methods include administering the first adenovirus vector multiple times and administering the second adenovirus multiple times. In this regard, the methods comprise administering the first adenovirus vector 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more times and administering the second adenovirus vector 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more times. The order of administration may comprise administering the first adenovirus one or multiple times in a row followed by administering the second adenovirus vector one or multiple times in a row. In certain embodiments, the methods include alternating administration of the first and the second adenovirus vectors as one administration each, two administrations each, three administrations each, and so on. In certain embodiments, the first and the second adenovirus vectors are administered simultaneously. In other embodiments, the first and the second adenovirus vectors are administered sequentially. In some embodiments, the target antigen comprises MUC1, MUC1c, MUC1n, T, or CEA, a fragment, a variant, or a variant fragment thereof.

[00306] As would be readily understood by the skilled artisan, more than two adenovirus vectors may be used in the methods of the present invention. Three, 4, 5, 6, 7, 8, 9, 10 or more different adenovirus vectors may be used in the methods of the invention. In certain embodiments, the methods comprise administering more than one E2b deleted adenovirus vector at a time. In this regard, immune responses against multiple target antigens of interest can be generated by administering multiple different adenovirus vectors simultaneously, each comprising nucleic acid sequences encoding one or more target antigens.

[00307] The adenovirus vectors can be used to generate an immune response against a cancer, such as carcinomas or sarcomas (e.g., solid tumors, lymphomas and leukemia). The adenovirus vectors can be used to generate an immune response against a cancer, such as neurologic cancers, melanoma, non-Hodgkin's lymphoma, Hodgkin's disease, leukemia, plasmocytomas, adenomas, gliomas, thymomas, breast cancer, prostate cancer, colorectal cancer, kidney cancer, renal cell carcinoma, uterine cancer, pancreatic cancer, esophageal cancer, lung cancer, ovarian cancer, cervical cancer, testicular cancer, gastric cancer, multiple myeloma, hepatoma, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), and chronic lymphocytic leukemia (CLL), or other cancers.

[00308] Methods are also provided for treating or ameliorating the symptoms of any of the infectious diseases or cancers as described herein. The methods of treatment comprise administering the adenovirus vectors one or more times to individuals suffering from or at risk from suffering from an infectious disease or cancer as described herein. As such, the present invention provides methods for vaccinating against infectious diseases or cancers in individuals who are at risk of developing such a disease. Individuals at risk may be individuals who may be exposed to an infectious agent at some time or have been previously exposed but do not yet have symptoms of infection or individuals having a genetic predisposition to developing a cancer or being particularly susceptible to an infectious agent. Individuals suffering from an infectious disease or cancer described herein may be determined to express and/or present a target antigen, which may be used to guide the therapies herein. For example, an example can be found to express and/or present a target antigen and an adenovirus vector encoding the target antigen, a variant, a fragment or a variant fragment thereof may be administered subsequently.

[00309] The present invention contemplates the use of adenovirus vectors for the in vivo delivery of nucleic acids encoding a target antigen, or a fragment, a variant, or a variant fragment thereof. Once injected into a subject, the nucleic acid sequence is expressed resulting in an immune response against the antigen encoded by the sequence. The adenovirus vector vaccine can be administered in an "effective amount", that is, an amount of adenovirus vector that is effective in a selected route or routes of administration to elicit an immune response as described elsewhere herein. An effective amount can induce an immune, immunoprotective or other immunotherapeutic response without significant adverse effects generally associated with typical vaccines. Once vaccinated, subjects may be monitored to determine the efficacy of the vaccine treatment. Monitoring the efficacy of vaccination may be performed by any method known to a person of ordinary skill in the art. In some embodiments, blood or fluid samples may be assayed to detect levels of antibodies. In other embodiments, ELISpot assays may be

performed to detect a cell-mediated immune response from circulating blood cells or from lymphoid tissue cells.

[00310] Routes and frequency of administration of the therapeutic compositions described herein, as well as dosage, may vary from individual to individual, and from disease to disease, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (e.g., intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (e.g., by aspiration), in pill form (e.g. swallowing, suppository for vaginal or rectal delivery). In certain embodiments, between 1 and 10 doses may be administered over a 52 week period. In certain embodiments, 6 doses are administered, at intervals of 1 month, and further booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. As such, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more doses may be administered over a 1 year period or over shorter or longer periods, such as over 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 week periods. Doses may be administered at 1, 2, 3, 4, 5, or 6 week intervals or longer intervals.

[00311] A vaccine can be infused over a period of less than about 4 hours, and more preferably, over a period of less than about 3 hours. For example, the first 25-50 mg could be infused within 30 minutes, preferably even 15 min, and the remainder infused over the next 2-3 hrs. More generally, the dosage of an administered vaccine construct may be administered as one dosage every 2 or 3 weeks, repeated for a total of at least 3 dosages. Or, the construct may be administered twice per week for 4-6 weeks. The dosing schedule can optionally be repeated at other intervals and dosage may be given through various parenteral routes, with appropriate adjustment of the dose and schedule. Compositions of the present invention can be administered to a patient in conjunction with (e.g., before, simultaneously, or following) any number of relevant treatment modalities.

[00312] A suitable dose is an amount of an adenovirus vector that, when administered as described above, is capable of promoting a target antigen immune response as described elsewhere herein. In certain embodiments, the immune response is at least 10-50% above the basal (i.e., untreated) level. In certain embodiments, the immune response is at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 110, 125, 150, 200, 250, 300, 400, 500 or more over the basal level. Such response can be monitored by measuring the target antigen(s) antibodies in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing patient tumor or infected cells in vitro, or other methods known in the art for monitoring immune responses. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome of the disease in question in vaccinated patients as compared to non-vaccinated patients. In some embodiments, the improved clinical outcome comprises treating disease, reducing the symptoms of a disease, changing the progression of a disease, or extending life.

[00313] In general, an appropriate dosage and treatment regimen provides the adenovirus vectors in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome for the particular disease being treated in treated patients as compared to non-treated patients. The monitoring data can be evaluated over time. The progression of a disease over time can be altered. Such improvements in clinical outcome would be readily recognized by a treating physician. Increases in preexisting immune responses to a target protein can generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

[00314] While one advantage of the present invention is the capability to administer multiple vaccinations with the same or different adenovirus vectors, particularly in individuals with preexisting immunity to Ad, the adenoviral vaccines of this invention may also be administered as part of a prime and boost regimen. A mixed modality priming and booster inoculation scheme may result in an enhanced immune response. Thus, one aspect of this invention is a method of priming a subject with a plasmid vaccine, such as a plasmid vector comprising a target antigen of interest, by administering the plasmid vaccine at least one time, allowing a predetermined length of time to pass, and then boosting by administering the adenovirus vector. Multiple primings, e.g., 1-4, may be employed, although more may be used. The length of time between priming and boost may typically vary from about four months to a year, but other time frames may be used. In certain embodiments, subjects may be primed 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more times with plasmid vaccines, and then boosted 4 months later with the adenovirus vector.

[00315] Any of the compositions provided herein may be administered to an individual. "Individual" may be used interchangeably with "subject" or "patient." An individual may be a mammal, for example a human or animal such as a non-human primate, a rodent, a rabbit, a rat, a mouse, a horse, a donkey, a goat, a cat, a dog, a cow, a pig, or a sheep. In embodiments, the individual is a human. In embodiments, the individual is a fetus, an embryo, or a child. In some cases, the compositions provided herein are administered to a cell *ex vivo*. In some cases, the compositions provided herein are administered to an individual as a method of treating a disease or disorder. In some embodiments, the individual has a genetic disease. In some cases, the individual is at risk of having the disease, such as any of the diseases described herein. In some embodiments, the individual is at increased risk of having a disease or disorder caused by insufficient amount of a protein or insufficient activity of a protein. If an individual is "at an increased risk" of having a disease or disorder, the method involves preventative or prophylactic treatment. For example, an individual can be at an increased risk of having such a disease or disorder because of family history of the disease. Typically, individuals at an increased risk of having such a

disease or disorder benefit from prophylactic treatment (*e.g.*, by preventing or delaying the onset or progression of the disease or disorder).

[00316] In some cases, a subject does not have a disease. In some cases, the treatment of the present invention is administered before onset of a disease. A subject may have undetected disease. A subject may have a low disease burden. A subject may also have a high disease burden. In certain cases, a subject may be administered a treatment of the present invention according to a grading scale. A grading scale can be a Gleason classification. A Gleason classification reflects how different tumor tissue is from normal prostate tissue. It uses a scale from 1 to 5. A physician gives a cancer a number based on the patterns and growth of the cancer cells. The lower the number, the more normal the cancer cells look and the lower the grade. The higher the number, the less normal the cancer cells look and the higher the grade. In certain cases, a treatment may be administered to a patient with a low Gleason score.

Preferably, a patient with a Gleason score of 3 or below may be administered a treatment of the present invention.

[00317] Various embodiments of the invention relate to compositions and methods for raising an immune response against one or more MUC1, MUC1c, MUC1n, T, or CEA antigens in selected patient populations. Accordingly, methods and compositions of the invention may target patients with a cancer including but not limited to carcinomas or sarcomas such as neurologic cancers, melanoma, non-Hodgkin's lymphoma, Hodgkin's disease, leukemia, plasmacytomas, adenomas, gliomas, thymomas, breast cancer, prostate cancer, colorectal cancer, kidney cancer, renal cell carcinoma, uterine cancer, pancreatic cancer, esophageal cancer, lung cancer, ovarian cancer, cervical cancer, testicular cancer, gastric cancer, multiple myeloma, hepatoma, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), and chronic lymphocytic leukemia (CLL), or other cancers can be targeted for therapy. In some cases, the targeted patient population may be limited to individuals having colorectal adenocarcinoma, metastatic colorectal cancer, advanced MUC1, MUC1c, MUC1n, T, or CEA expressing colorectal cancer, head and neck cancer, liver cancer, breast cancer, lung cancer, bladder cancer, or pancreas cancer. A histologically confirmed diagnosis of a selected cancer, for example colorectal adenocarcinoma, may be used. A particular disease stage or progression may be selected, for example, patients with one or more of a metastatic, recurrent, stage III, or stage IV cancer may be selected for therapy with the methods and compositions of the invention. In some embodiments, patients may be required to have received and, optionally, progressed through other therapies including but not limited to fluoropyrimidine, irinotecan, oxaliplatin, bevacizumab, cetuximab, or panitumumab containing therapies. In some cases, individual's refusal to accept such therapies may allow the patient to be included in a therapy eligible pool with methods and compositions of the invention. In some embodiments, individuals to receive therapy using the methods and compositions of the invention may be required to have an estimated life expectancy of at least, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 18, 21,

or 24 months. The patient pool to receive a therapy using the methods and compositions of the invention may be limited by age. For example, individuals who are older than 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 25, 30, 35, 40, 50, 60, or more years old can be eligible for therapy with methods and compositions of the invention. For another example, individuals who are younger than 75, 70, 65, 60, 55, 50, 40, 35, 30, 25, 20, or fewer years old can be eligible for therapy with methods and compositions of the invention.

[00318] In some embodiments, patients receiving therapy using the methods and compositions of the invention are limited to individuals with adequate hematologic function, for example with one or more of a WBC count of at least 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000 or more per microliter, a hemoglobin level of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or higher g/dL, a platelet count of at least 50,000; 60,000; 70,000; 75,000; 90,000; 100,000; 110,000; 120,000; 130,000; 140,000; 150,000 or more per microliter; with a PT-INR value of less than or equal to 0.8, 1.0, 1.2, 1.3, 1.4, 1.5, 1.6, 1.8, 2.0, 2.5, 3.0, or higher, a PTT value of less than or equal to 1.2, 1.4, 1.5, 1.6, 1.8, 2.0 X ULN or more. In various embodiments, hematologic function indicator limits are chosen differently for individuals in different gender and age groups, for example 0-5, 5-10, 10-15, 15-18, 18-21, 21-30, 30-40, 40-50, 50-60, 60-70, 70-80 or older than 80.

[00319] In some embodiments, patients receiving therapy using the methods and compositions of the invention are limited to individuals with adequate renal and/or hepatic function, for example with one or more of a serum creatinine level of less than or equal to 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2 mg/dL or more, a bilirubin level of .8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2 mg/dL or more, while allowing a higher limit for Gilbert's syndrome, for example, less than or equal to 1.5, 1.6, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, or 2.4 mg/dL, an ALT and AST value of less than or equal to less than or equal to 1.5, 2.0, 2.5, 3.0 x upper limit of normal (ULN) or more. In various embodiments, renal or hepatic function indicator limits are chosen differently for individuals in different gender and age groups, for example 0-5, 5-10, 10-15, 15-18, 18-21, 21-30, 30-40, 40-50, 50-60, 60-70, 70-80 or older than 80.

[00320] In some embodiments, the K-ras mutation status of individuals who are candidates for a therapy using the methods and compositions of the invention can be determined. Individuals with a preselected K-ras mutational status can be included in an eligible patient pool for therapies using the methods and compositions of the invention.

[00321] In various embodiments, patients receiving therapy using the methods and compositions of the invention are limited to individuals without concurrent cytotoxic chemotherapy or radiation therapy, a history of, or current, brain metastases, a history of autoimmune disease, such as but not restricted to, inflammatory bowel disease, systemic lupus erythematosus, ankylosing spondylitis, scleroderma, multiple sclerosis, thyroid disease and vitiligo, serious intercurrent chronic or acute illness, such as

cardiac disease (NYHA class III or IV), or hepatic disease, a medical or psychological impediment to probable compliance with the protocol, concurrent (or within the last 5 years) second malignancy other than non-melanoma skin cancer, cervical carcinoma *in situ*, controlled superficial bladder cancer, or other carcinoma *in situ* that has been treated, an active acute or chronic infection including: a urinary tract infection, HIV (e.g. as determined by ELISA and confirmed by Western Blot), and chronic hepatitis, or concurrent steroid therapy (or other immuno-suppressives, such as azathioprine or cyclosporin A). In some cases, patients with at least 3, 4, 5, 6, 7, 8, 9, or 10 weeks of discontinuation of any steroid therapy (except that used as pre-medication for chemotherapy or contrast-enhanced studies) may be included in a pool of eligible individuals for therapy using the methods and compositions of the invention.

[00322] In some embodiments, patients receiving therapy using the methods and compositions of the invention include individuals with thyroid disease and vitiligo.

[00323] In various embodiments, samples, for example serum or urine samples, from the individuals or candidate individuals for a therapy using the methods and compositions of the invention may be collected. Samples may be collected before, during, and/or after the therapy for example, within 2, 4, 6, 8, 10 weeks prior to the start of the therapy, within 1 week, 10 day, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, or 12 weeks from the start of the therapy, within 2, 4, 6, 8, 10 weeks prior to the start of the therapy, within 1 week, 10 day, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, 9 weeks, or 12 weeks from the start of the therapy, in 1 week, 10 day, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, 9 weeks, or 12 weeks intervals during the therapy, in 1 month, 3 month, 6 month, 1 year, 2 year intervals after the therapy, within 1 month, 3 months, 6 months, 1 year, 2 years, or longer after the therapy, for a duration of 6 months, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 years or longer. The samples may be tested for any of the hematologic, renal, or hepatic function indicators described herein as well as suitable others known in the art, for example a β -HCG for women with childbearing potential. In that regard, hematologic and biochemical tests, including cell blood counts with differential, PT, INR and PTT, tests measuring Na, K, Cl, CO_2 , BUN, creatinine, Ca, total protein, albumin, total bilirubin, alkaline phosphatase, AST, ALT and glucose are within the bounds of the invention. In some embodiments, the presence or the amount of HIV antibody, Hepatitis BsAg, or Hepatitis C antibody are determined in a sample from individuals or candidate individuals for a therapy using the methods and compositions of the invention. Biological markers, such as antibodies to MUC1, MUC1c, MUC1n, T, or CEA or the neutralizing antibodies to Ad5 vector can be tested in a sample, such as serum, from individuals or candidate individuals for a therapy using the methods and compositions of the invention. In some cases, one or more samples, such as a blood sample can be collected and archived from an individuals or candidate individuals for a therapy using the methods and compositions of the invention. Collected samples can be assayed for immunologic evaluation. Individuals or candidate individuals for a therapy using the methods and compositions of the invention can be evaluated in imaging studies, for example using CT scans or MRI of the chest,

abdomen, or pelvis. Imaging studies can be performed before, during, or after therapy using the methods and compositions of the invention, during, and/or after the therapy, for example, within 2, 4, 6, 8, 10 weeks prior to the start of the therapy, within 1 week, 10 day, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, or 12 weeks from the start of the therapy, within 2, 4, 6, 8, 10 weeks prior to the start of the therapy, within 1 week, 10 day, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, 9 weeks, or 12 weeks from the start of the therapy, in 1 week, 10 day, 2 week, 3 week, 4 week, 6 week, 8 week, 9 week, or 12 week intervals during the therapy, in 1 month, 3 month, 6 month, 1 year, 2 year intervals after the therapy, within 1 month, 3 months, 6 months, 1 year, 2 years, or longer after the therapy, for a duration of 6 months, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 years or longer.

Dosages and Administration

[00324] Compositions and methods of the invention contemplate various dosage and administration regimens during therapy. Patients may receive one or more replication defective adenovirus or adenovirus vector, for example Ad5 [E1-, E2B-]-CEA(6D), Ad5 [E1-, E2b-]-MUC1, Ad5 [E1-, E2b-]-MUC1c, Ad5 [E1-, E2b-]-MUC1n, Ad5 [E1-, E2b-]-T that is capable of raising an immune response in an individual against a target antigen described herein. In various embodiments, the replication defective adenovirus is administered at a dose that suitable for effecting such immune response. In some cases, the replication defective adenovirus is administered at a dose that is greater than or equal to 1×10^9 , 2×10^9 , 3×10^9 , 4×10^9 , 5×10^9 , 6×10^9 , 7×10^9 , 8×10^9 , 9×10^9 , 1×10^{10} , 2×10^{10} , 3×10^{10} , 4×10^{10} , 5×10^{10} , 6×10^{10} , 7×10^{10} , 8×10^{10} , 9×10^{10} , 1×10^{11} , 2×10^{11} , 3×10^{11} , 4×10^{11} , 5×10^{11} , 6×10^{11} , 7×10^{11} , 8×10^{11} , 9×10^{11} , 1×10^{12} , 1.5×10^{12} , 2×10^{12} , 3×10^{12} , or more virus particles (VP) per immunization. In some cases, the replication defective adenovirus is administered at a dose that is less than or equal to 1×10^9 , 2×10^9 , 3×10^9 , 4×10^9 , 5×10^9 , 6×10^9 , 7×10^9 , 8×10^9 , 9×10^9 , 1×10^{10} , 2×10^{10} , 3×10^{10} , 4×10^{10} , 5×10^{10} , 6×10^{10} , 7×10^{10} , 8×10^{10} , 9×10^{10} , 1×10^{11} , 2×10^{11} , 3×10^{11} , 4×10^{11} , 5×10^{11} , 6×10^{11} , 7×10^{11} , 8×10^{11} , 9×10^{11} , 1×10^{12} , 1.5×10^{12} , 2×10^{12} , 3×10^{12} , or more virus particles per immunization. In various embodiments, a desired dose described herein is administered in a suitable volume of formulation buffer, for example a volume of about 0.1-10 mL, 0.2-8mL, 0.3-7mL, 0.4-6 mL, 0.5-5 mL, 0.6-4 mL, 0.7-3 mL, 0.8-2 mL, 0.9-1.5 mL, 0.95-1.2 mL, or 1.0-1.1 mL. Those of skill in the art appreciate that the volume may fall within any range bounded by any of these values (e.g., about 0.5 mL to about 1.1 mL).

Administration of virus particles can be through a variety of suitable paths for delivery, for example it can be by injection (e.g., intracutaneously, intramuscularly, intravenously or subcutaneously), intranasally (e.g., by aspiration), in pill form (e.g. swallowing, suppository for vaginal or rectal delivery). In some embodiments, a subcutaneous delivery may be preferred and can offer greater access to dendritic cells.

[00325] Administration of virus particles to an individual may be repeated. Repeated deliveries of virus particles may follow a schedule or alternatively, may be performed on an as needed basis. For example,

an individual's immunity against a target antigen, for example MUC1, T and/or CEA, may be tested and replenished as necessary with additional deliveries. In some embodiments, schedules for delivery include administrations of virus particles at regular intervals. Joint delivery regimens may be designed comprising one or more of a period with a schedule and/or a period of need based administration assessed prior to administration. For example, a therapy regimen may include an administration, such as subcutaneous administration once every three weeks then another immunotherapy treatment every three months until removed from therapy for any reason including death. Another example regimen comprises three administrations every three weeks then another set of three immunotherapy treatments every three months. Another example regimen comprises a first period with a first number of administrations at a first frequency, a second period with a second number of administrations at a second frequency, a third period with a third number of administrations at a third frequency, etc., and optionally one or more periods with undetermined number of administrations on an as needed basis. The number of administrations in each period can be independently selected and can for example be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more. The frequency of the administration in each period can also be independently selected, can for example be about every day, every other day, every third day, twice a week, once a week, once every other week, every three weeks, every month, every six weeks, every other month, every third month, every fourth month, every fifth month, every sixth month, once a year etc. The therapy can take a total period of up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 30, 36 months or more. The scheduled interval between immunizations may be modified so that the interval between immunizations is revised by up to a fifth, a fourth, a third, or half of the interval. For example, for a 3-week interval schedule, an immunization may be repeated between 20 and 28 days (3 weeks -1 day to 3 weeks +7 days). For the first 3 immunizations, if the second and/or third immunization is delayed, the subsequent immunizations may be shifted allowing a minimum amount of buffer between immunizations. For example, for a three week interval schedule, if an immunization is delayed, the subsequent immunization may be scheduled to occur no earlier than 17, 18, 19, or 20 days after the previous immunization.

[00326] Compositions of the invention, such as Ad Ad5 [E1-, E2B-]-CEA(6D), Ad5 [E1-, E2B-]-MUC1, Ad5 [E1-, E2B-]-MUC1c, Ad5 [E1-, E2B-]-MUC1n, Ad5 [E1-, E2B-]-T virus particles, can be provided in various states, for example, at room temperature, on ice, or frozen. Compositions may be provided in a container of a suitable size, for example a vial of 2 mL vial. In one embodiment, 1 2ml vial with 1.0 mL of extractable vaccine contains 5×10^{11} total virus particles/mL. Storage conditions including temperature and humidity may vary. For example, compositions for use in therapy may be stored at room temperature, 4 °C, -20 °C, or lower.

[00327] In various embodiments, general evaluations are performed on the individuals receiving treatment according to the methods and compositions of the invention. One or more of any tests may be

performed as needed or in a scheduled basis, such as on weeks 0, 3, 6 etc. A different set of tests may be performed concurrent with immunization vs. at time points without immunization.

[00328] General evaluations may include one or more of medical history, ECOG Performance Score, Karnofsky performance status, and complete physical examination with weight by the attending physician. Any other treatments, medications, biologics, or blood products that the patient is receiving or has received since the last visit may be recorded. Patients may be followed at the clinic for a suitable period, for example approximately 30 minutes, following receipt of vaccine to monitor for any adverse reactions. Local and systemic reactogenicity after each dose of vaccine will be assessed daily for a selected time, for example for 3 days (on the day of immunization and 2 days thereafter). Diary cards may be used to report symptoms and a ruler may be used to measure local reactogenicity. Immunization injection sites may be assessed. CT scans or MRI of the chest, abdomen, and pelvis may be performed.

[00329] In various embodiments, hematological and biochemical evaluations are performed on the individuals receiving treatment according to the methods and compositions of the invention. One or more of any tests may be performed as needed or in a scheduled basis, such as on weeks 0, 3, 6 etc. A different set of tests may be performed concurrent with immunization vs. at time points without immunization. Hematological and biochemical evaluations may include one or more of blood test for chemistry and hematology, CBC with differential, Na, K, Cl, CO₂, BUN, creatinine, Ca, total protein, albumin, total bilirubin, alkaline phosphatase, AST, ALT, glucose, and ANA

[00330] In various embodiments, biological markers are evaluated on individuals receiving treatment according to the methods and compositions of the invention. One or more of any tests may be performed as needed or in a scheduled basis, such as on weeks 0, 3, 6 etc. A different set of tests may be performed concurrent with immunization vs. at time points without immunization.

[00331] Biological marker evaluations may include one or more of measuring antibodies to MUC1, MUC1c, MUC1n, CEA or the Ad5 vector, from a serum sample of adequate volume, for example about 5ml Biomarkers (e.g., CEA or CA15-3) may be reviewed if determined and available.

[00332] In various embodiments, an immunological assessment is performed on individuals receiving treatment according to the methods and compositions of the invention. One or more of any tests may be performed as needed or in a scheduled basis, such as on weeks 0, 3, 6 etc. A different set of tests may be performed concurrent with immunization vs. at time points without immunization.

[00333] Peripheral blood, for example about 90mL may be drawn prior to each immunization and at a time after at least some of the immunizations, to determine whether there is an effect on the immune response at specific time points during the study and/or after a specific number of immunizations.

Immunological assessment may include one or more of assaying peripheral blood mononuclear cells (PBMC) for T-cell responses to MUC1, MUC1c, MUC1n, T or CEA using ELISpot, proliferation assays,

multi-parameter flow cytometric analysis, and cytotoxicity assays. Serum from each blood draw may be archived and sent and determined.

[00334] In various embodiments, a tumor assessment is performed on individuals receiving treatment according to the methods and compositions of the invention. One or more of any tests may be performed as needed or in a scheduled basis, such as prior to treatment, on weeks 0, 3, 6 etc. A different set of tests may be performed concurrent with immunization vs. at time points without immunization. Tumor assessment may include one or more of CT or MRI scans of chest, abdomen, or pelvis performed prior to treatment, at a time after at least some of the immunizations and at approximately every three months following the completion of a selected number, for example 2, 3, or 4, of first treatments and for example until removal from treatment.

[00335] Immune responses against a target antigen described herein, such as CEA, may be evaluated from a sample, such as a peripheral blood sample of an individual using one or more suitable tests for immune response, such as ELISpot, cytokine flow cytometry, or antibody response. A positive immune response can be determined by measuring a T-cell response. A T-cell response can be considered positive if the mean number of spots adjusted for background in six wells with antigen exceeds the number of spots in six control wells by 10 and the difference between single values of the six wells containing antigen and the six control wells is statistically significant at a level of $p \leq 0.05$ using the Student's t-test. Immunogenicity assays may occur prior to each immunization and at scheduled time points during the period of the treatment. For example, a time point for an immunogenicity assay at around week 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 18, 20, 24, 30, 36, or 48 of a treatment may be scheduled even without a scheduled immunization at this time. In some cases, an individual may be considered evaluable for immune response if they receive at least a minimum number of immunizations, for example 1, 2, 3, 4, 5, 6, 7, 8, 9, or more immunizations.

[00336] In some embodiments, disease progression or clinical response determination is made according to the RECIST 1.1 criteria among patients with measurable/evaluable disease. In some embodiments, therapies using the methods and compositions of the invention affect a Complete Response (CR; disappearance of all target lesions for target lesions or disappearance of all non-target lesions and normalization of tumor marker level for non-target lesions) in an individual receiving the therapy. In some embodiments, therapies using the methods and compositions of the invention affect a Partial Response (PR; at least a 30% decrease in the sum of the LD of target lesions, taking as reference the baseline sum LD for target lesions) in an individual receiving the therapy.

[00337] In some embodiments, therapies using the methods and compositions of the invention affect a Stable Disease (SD; neither sufficient shrinkage to qualify for PR nor sufficient increase to qualify for PD, taking as reference the smallest sum LD since the treatment started for target lesions) in an individual receiving the therapy. In some embodiments, therapies using the methods and compositions of the

invention affect an Incomplete Response/ Stable Disease (SD; persistence of one or more non-target lesion(s) or/and maintenance of tumor marker level above the normal limits for non-target lesions) in an individual receiving the therapy. In some embodiments, therapies using the methods and compositions of the invention affect a Progressive Disease (PD; at least a 20% increase in the sum of the LD of target lesions, taking as reference the smallest sum LD recorded since the treatment started or the appearance of one or more new lesions for target lesions or persistence of one or more non-target lesion(s) or/and maintenance of tumor marker level above the normal limits for non-target lesions) in an individual receiving the therapy.

KITS

[00338] The compositions, immunotherapy or vaccines may be supplied in the form of a kit. The kits of the present disclosure may further comprise instructions regarding the dosage and or administration including treatment regimen information.

[00339] In some embodiments, kits comprise the compositions and methods for providing combination multi-targeted cancer immunotherapy. In some embodiments, kits comprise the compositions and methods for the combination multi-targeted treatment of an infectious disease. In some embodiment's kits may further comprise components useful in administering the kit components and instructions on how to prepare the components. In some embodiments, the kit can further comprise software for conducting monitoring patient before and after treatment with appropriate laboratory tests, or communicating results and patient data with medical staff.

[00340] The components comprising the kit may be in dry or liquid form. If they are in dry form, the kit may include a solution to solubilize the dried material. The kit may also include transfer factor in liquid or dry form. If the transfer factor is in dry form, the kit will include a solution to solubilize the transfer factor. The kit may also include containers for mixing and preparing the components. The kit may also include instrument for assisting with the administration such for example needles, tubing, applicator, inhalant, syringe, pipette, forceps, measured spoon, eye dropper or any such medically approved delivery vehicle. The kits or drug delivery systems of the present invention also will typically include a means for containing compositions of the present disclosure in close confinement for commercial sale and distribution.

EXAMPLES

Peptides and Vectors

[00341] The following HLA-A2 and HLA-A24 binding peptides were used in this and other examples: (a) the HLA-A2 binding CEA agonist peptide CAPI-6D (YLSGADLNL (SEQ ID NO:10)), (b) the HLA-A2 MUC1 agonist peptide P93L (ALWGQDVTSV (SEQ ID NO:12)), (c) the HLA-A24 binding MUC1 agonist peptide C6A (KYHPMSEYAL (SEQ ID NO:13)), and (d) the HLA-A2 binding brachyury agonist peptide (WLLPGTSTV (SEQ ID NO:14)). All peptides were greater than 96% pure.

[00342] Ad5 [E1-, E2b-]-brachyury, Ad5 [E1-, E2b-]-CEA and Ad5 [E1-, E2b-]-MUC1 were constructed and produced. Briefly, the transgenes were sub-cloned into the E1 region of the Ad5 [E1-, E2b-] vector using a homologous recombination-based approach. The replication deficient virus was propagated in the E.C7 packaging cell line, CsCl₂ purified, and titered. Viral infectious titer was determined as plaque-forming units (PFUs) on an E.C7 cell monolayer. The VP concentration was determined by sodium dodecyl sulfate (SDS) disruption and spectrophotometry at 260 nm and 280 nm. The CEA transgene also contained a modified CEA containing the highly immunogenic epitope CAPI-6D.

[00343] The sequence encoding for the human brachyury protein (T, NM_003181.3) was modified by introducing the enhancer T-cell HLA-A2 epitope (WLLPGTSTV) and removal of a 25 amino acid fragment involved in DNA binding. The resulting construct was subsequently subcloned into the Ad5 vector to generate the Ad5 [E1-, E2b-]-brachyury construct.

[00344] The MUC1 molecule consisted of two regions: the N-terminus (MUC1-n), which is the large extracellular domain of MUC1, and the C-terminus (MUC1-c), which has three regions: a small extracellular domain, a single transmembrane domain, and a cytoplasmic tail. The cytoplasmic tail contained sites for interaction with signaling proteins and acts as an oncogene and a driver of cancer motility, invasiveness and metastasis. For construction of the Ad5 [E1-, E2b-]-MUC1, the entire MUC1 transgene, including eight agonist epitopes, was subcloned into the Ad5 vector. The agonist epitopes included in the Ad5 [E1-, E2b-]-MUC1 vector bind to HLA-A2 (epitope P93L in the N-terminus, V1A and V2A in the VNTR region, and C1A, C2A and C3A in the C-terminus), HLA-A3 (epitope C5A), and HLA-A24 (epitope C6A in the C-terminus). The Tri-Ad5 vaccine was produced by combining of 10¹⁰ VP of Ad5 [E1-, E2b-]-Brachyury, Ad5 [E1-, E2b-]-CEA and Ad5 [E1-, E2b-]-MUC1 at a ratio of 1:1:1 (3x10¹⁰ VP total).

EXAMPLE 1: Multiple Injections of Ad5Null Adenovirus Vector Produces Anti-Adenovirus

Antibodies

[00345] This example shows that multiple injections of Ad5-null results in the production of anti-adenovirus antibodies in the injected subjects.

[00346] It was demonstrated that the Ad5-null adenovirus vector that does not contain any heterologous nucleic acid sequences, generated a neutralizing immune response in mice. In one experiment, female Balb/c mice aged 5-7 weeks were immunized with Ad5Null viral particles at 14 day intervals. To determine the presence of anti-adenovirus antibodies, an enzyme linked immunosorbent assay (ELISA) was used. For this ELISA, 10^9 viral particles were coated onto microtiter wells in 100 μ L of 0.05M carbonate/bicarbonate buffer, pH 9.6, and incubated overnight at room temperature. For a standard immunoglobulin G (IgG) reference curve, 200 ng, 100 ng, 50 ng, 25 ng, and 0 ng of purified mouse IgG were coated onto microtiter wells as described above. After incubation, all wells were washed 3 times with 250 μ L of 1% bovine serum albumin (BSA) in phosphate buffered saline (PBS), pH 7.4. After washing, 250 μ L of BSA/PBS was added to all and incubated for 30 minutes at room temperature to block unbound sites. After incubation, all wells were washed 3 times with 250 μ L of BSA/PBS. After washing, 200 μ L of a 1/100 serum dilution in BSA/PBS was added to wells and incubated for 1 hour at room temperature. For a positive control, 200 μ L of a 1/10000 dilution of anti-adenovirus antiserum in BSA/PBS was added to wells. Control wells contained BSA/PBS only. After incubation, all wells were washed 3 times with 250 μ L of BSA/PBS. After washing, 200 μ L of a 1/10000 dilution of peroxidase conjugated γ -chain specific goat anti-mouse IgG (Sigma Chemicals) in BSA/PBS were added to each well and incubated for 1 hour at room temperature. After incubation, all wells were washed 3 times with 250 μ L of BSA/PBS. After washing, 200 μ L of developing reagent (0.5 mg/mL 1,2-phenylene-diamine in 0.2M potassium phosphate buffer, pH 5.0, containing 0.06% hydrogen peroxide) was added to each well and incubated for 30-40 minutes at room temperature. After incubation, the color reaction was stopped by addition of 50 μ L 5N HCl to each well. All wells were then read in a microwell plate reader at 492 nm. After readings were obtained, the optical density readings of unknown samples were correlated with the standard IgG curve to obtain the ngs of IgG bound per well. This was performed using the INSTAT statistical package.

ELISA to detect antibodies against CEA

[00347] ELISA plates were coated with 100 ng of human CEA (Sigma-Aldrich) in 0.05 M carbonate-bicarbonate buffer pH 9.6 and incubated overnight at room temperature. Plates were washed three times with phosphate buffered saline containing 1% Tween-20 (PBS-T) and then blocked with PBS containing 1% BSA for 60 min at room temperature. After an additional three washes, serum diluted 1/50 in PBS-T was added to the wells and the plates were incubated for 1 hour at room temperature. Peroxidase labeled goat anti-mouse immunoglobulin (Ig) G (γ -chain specific) (Sigma-Aldrich) antibody at a 1:5000 dilution was added to the wells after washings and plates were incubated for 1 hour. Plates were washed three times and 1,2-phenylene-diamine substrate solution was added to each well. The reaction was stopped by adding 10% phosphoric acid. Absorbance was measured at 492 nm on a SpectraMax 190 ELISA reader. The nanogram equivalents of IgG bound to CEA per well was obtained by reference to a standard curve

generated using purified mouse IgG and developed at the same time as the CEA ELISA. The results were analyzed and quantitated using SoftMax Pro 6.3 software.

[00348] Significant levels (P<0.001) of anti-adenovirus IgG antibody were detected in mice 2 weeks after a first injection with 10^{10} Ad-5-null (FIG. 1). A significantly higher level (P<0.001) was observed 2 weeks after a second injection with 10^{10} adenovirus. Significantly higher (P<0.001) levels of antibody were continued to be observed 2 weeks after a third injection with 10^{10} Ad5-null. Each value represents the average of triplicate determinations from pooled sera of 5 mice in each group. Multiple injections of Ad5-null resulted in production of anti-adenovirus antibodies in the subjects.

[00349] To determine the presence of neutralizing antibody to Ad, the following assay was utilized. A HEK-293T-cell line was cultured in 200 μ L of culture medium consisting of DMEM containing 10% fetal calf serum (DMEM/FCS) in microwell tissue culture plates at a cell concentration of 2×10^3 cells per well for 24 hours at 37 °C in 5% CO₂. After incubation, 100 μ L of culture medium was removed from triplicate wells and mixed with 20 μ L of DMEM/FCS containing viral particles (VP). After mixing, the 120 μ L mixture was added back to the respective microwells. In another set of triplicate wells, 100 μ L of culture medium was removed and mixed with 20 μ L of heat inactivated (56 °C for 1 h) Ad immune mouse serum previously incubated with VP for one hour at room temperature. After mixing, the 120 μ L mixture was added back to the respective wells. In triplicate cell control wells, 20 μ L of DMEM/FCS was added to control for total culture medium volume. Triplicate medium-only control wells contained 220 μ L of DMEM/FCS. The tissue culture plate was incubated for an additional 3 days at 37 °C in 5% CO₂. After incubation, 40 μ L of PROMEGA cell viability reagent (Owen's reagent) was added to all wells and incubated for 75 minutes at 37 °C in 5% CO₂. In this assay, the Owen's reagent (MTS tetrazolium compound) is bioreduced by viable cells into a colored formazan product that is soluble in tissue culture medium. The quantity of formazan product as measured by absorbance at 490 nm is directly proportional to the number of living cells in culture. After incubation, 150 μ L was removed from each well and transferred to another microwell plate for optical density readings. Optical density readings at 492 nm were subsequently obtained using a microwell plate reader.

[00350] To detect the presence of neutralizing antibodies to Ad, groups of 5 mice each were injected once, twice, or three times with 10^{10} Ad5-null at two week intervals. Two weeks after the final injection of virus, mice were bled, pooled, and assessed for neutralizing antibody as described above using 4×10^7 VP incubated with or without heat inactivated sera. Cells cultured alone served as a control group. Normal mice and mice injected one time with Ad5null did not exhibit significant levels of neutralizing antibody (FIG. 2). Mice injected two times with Ad exhibited significant (P<0.05) levels of neutralizing antibody as compared with cells incubated with virus only. Mice injected three times with Ad5-null also exhibited significant (P<0.01) levels of neutralizing antibody as compared with cells incubated with virus only.

EXAMPLE 2: The Ad5 [E1-]-CEA Vector Vaccine Induces CEA Specific Immune Response Upon Re-Immunization in Ad5 Immune Mice

[00351] This example shows that the Ad5 [E1-, E2b-] vector platform induces CMI responses against the tumor associated antigen (TAA) carcinoembryonic antigen (CEA) in the presence of pre-existing Ad5 immunity in mice.

Characterization of Ad5 CEA Vectors

[00352] Initial studies were performed to confirm CEA gene expression of two Ad5-CEA vector platforms. It was first determined that the CEA antigen could be expressed on cells transfected with the vaccine vector platforms. A549 cells were obtained from ATCC and transfected with Ad5 [E1-]-CEA or Ad5 [E1-, E2b-]-CEA. Western blot analysis revealed that cells transfected with the vector platforms expressed CEA antigen. (**FIG. 35**)

Methods

[00353] A549 cells were inoculated at a MOI of 555 VPs/cell with Ad5 [E1-, E2b-]-CEA. Cells were incubated for 48 hours at 37°C in 5% CO₂. After 48 hours cells were harvested and washed with PBS and freeze/thawed three times. The whole cell lysate was heated for 70 °C for 10 min prior to loading on the gel. Recombinant CEA control was loaded at 30ng/Lane and the prepared lysate at 20 µL/lane. Sample loading buffer was included as an additional negative control and the positive controls were Magic Mark CP Western markers and the recombinant CEA. The gel was transferred to a nitrocellulose membrane and blocked with SuperBlock Blocking solution for 60 min. The membrane was probed with mouse monoclonal anti-CEA primary antibody (1:1000) and a secondary anti-mouse HRP (1:2500) conjugated antibody. The membrane was washed three times then incubated with SuperSignal chemiluminescent reagent and banding was visualized by exposing X-ray film to the membrane followed by development.

Induction of Ad5 Immunity in Mice

[00354] To assess the levels of Ad5 immunity that could be induced, groups of Ad5 naive C57Bl/6 mice were injected subcutaneously with the Ad5 vector platform (VP). Twenty eight to forty two days later, serum samples were collected and assessed for endpoint Ad5 NAb titers. As shown in **FIG. 3**, undetectable Ad5 NAb titers (endpoint Ad5 NAb titer < 1/25) were observed in normal control mice. Ad5 NAb (endpoint titers of 1/25 to 1/50) was detectable after one injection but dramatically increased after three injections of 10¹⁰ Ad5. Therefore, in additional Ad5 immune studies, mice were injected twice with 10¹⁰ Ad5 VP to render the animals Ad5 immune.

Immunization of Ad5 Immune Mice with Ad5 [E1-]-CEA or Ad5 [E1-, E2b-]-CEA.

[00355] These experiments were designed to determine and compare the immunization induction potential of Ad5 [E1-]-CEA and Ad5 [E1-, E2b-]-CEA vaccines in Ad5 immune mice. Groups of female C57Bl/6 mice, 4 to 8 weeks old, were immunized 2 times at 2 week intervals with 10¹⁰ Ad5-null VP. Two weeks following the last Ad5-null immunization, the mice were immunized 3 times at weekly

intervals with 10^{10} VP of Ad5 [E1-]-CEA or Ad5 [E1-, E2b-]-CEA. Two weeks following the last immunization, mice were euthanized and their spleens and sera harvested for analyses.

[00356] CMI responses were assessed by ELISpot assays performed on splenocytes exposed to intact CEA antigen. Splenocytes from Ad5 immune C57Bl/6 mice that were immunized subcutaneously with Ad5 E1-]-CEA or Ad5 [E1-, E2b-]-CEA were harvested and assessed for the number of IFN- γ and IL-2 secreting cells as described above. Significantly elevated numbers of both IFN- γ and IL-2 secreting cells were observed in spleens assayed from mice immunized with Ad5 [E1-, E2b-]-CEA as compared to immunized Ad5 [E1-]-CEA mice (**FIG. 4A and FIG. 4B**). Specificity studies revealed that immunizations with Ad5 CEA vectors induced specific CEA associated CMI responses and not responses against other irrelevant antigens such as the HIV-gag protein or β -galactosidase. These results demonstrate that immunization of Ad5 immune mice with Ad5 [E1-, E2b-]-CEA induce significantly higher CMI responses.

Lack of Adverse Liver Effects in Immunized Mice

[00357] Toxicity studies were performed on serum from Ad5 immune female C57Bl/6 mice immunized with Ad5 [E1-]-CEA, Ad5 [E1-, E2b-]-CEA as described above. Ad5 naive or Ad5 immune mice injected with buffer alone served as controls. Three days after the third immunization, aspartate aminotransferase (AST) levels were assessed on the blood samples to determine liver toxicity due to the treatment. AST levels were not elevated over controls following immunization with either vector (**FIG. 5**). Alanine aminotransferase (ALT) levels were also assessed and similar results were observed.

Ad5 [E1-, E2b-]-CEA Immunotherapy in Ad5 Immune Tumor Bearing Mice

[00358] Based upon the successful immunological results observed above, studies in which MC38 tumors were established in mice and then treated were performed as described below. For these studies a CEA expressing MC38 murine cell line was used. This cell line has been genetically modified to express human CEA and can be implanted into C57Bl/6 mice. After tumor establishment, the mice were treated with the novel Ad5 [E1-, E2b-]-CEA vector platform. To determine if Ad5 immune tumor bearing mice could be treated with the Ad5 [E1-, E2b-]-CEA vector, C57Bl/6 mice were injected two times subcutaneously with 10^{10} Ad5 [E1-]-null VP at 14 day intervals to render the mice Ad5 immune. Two weeks after the last injection, two groups of 7 C57Bl/6 mice were injected subcutaneously with 10^6 CEA expressing MC38 tumor cells. Seven days later, when tumors were palpable, one group of mice was treated by distal subcutaneous injection with 10^{10} VP of Ad5 [E1-, E2b-]-CEA on days 7, 13 and 19. A group of 7 injection buffer only treated C57Bl/6 mice served as untreated controls. All mice were monitored for tumor size over a 21 day period and tumor volumes were determined as previously described.

[00359] The tumor growth by day 19 was significantly reduced in the Ad5 [E1-, E2b-]-CEA treated mice and remained so (**FIG. 6**). At the end of the study (Day 22), the mice were sacrificed and the tumors

were excised and weighed. Tumor measurements were taken and volumes were determined. Statistical analysis was performed using the Bonferroni post-tests analysis with PRISM software. The tumors in the mice treated with Ad5 [E1-, E2b-]-CEA were significantly ($P<0.05$) smaller in weight than the untreated controls (**FIG. 7**).

[00360] At the termination of the study, spleens were collected from mice and the CEA specific CMI response was determined by ELISpot assay. CEA specific IFN- γ secretion response was significantly higher in mice immunized with Ad5 [E1-, E2b-]-CEA than in mice who received MC-38 tumor cells alone. These results indicate that treatment of CEA expressing tumors in Ad5 immunized mice using the Ad5 [E1-, E2b-]-CEA vaccine can significantly decrease tumor growth progression.

EXAMPLE 3: Quantitative ELISA for CEA expression on A549 cells after infection

[00361] This example shows a dose response evaluation using the Ad5 [E1-, E2b-] CEA vector to transduce the human cancerous lung cell line, A-549. The results show that the CEA antigen can be expressed in a dose dependent manner.

Experimental Design

[00362] On day one, of the assay a BD Falcon Tissue Culture 96-well plate was seeded with A549 cells passaged three days prior (lot# 30Jul02, passage p+23), (7.7×10^3 cells/well) and placed into a 37 ± 2 °C incubator with a $5 \pm 2\%$ CO₂ atmosphere overnight.

[00363] The next day, a dilution series of the test article were prepared and replicate wells were inoculated at levels ranging from 1.56×10^3 to 2.5×10^4 viral particles/well. Untreated A549 cells were used to serve as the mock sample. On day four of the assay wells were treated with a 10% Triton X-100 solution for analysis by ELISA to measure CEA concentration. For the ELISA, a microtiter plate was coated overnight with an anti-CEA capture antibody (abcam Carcino embryonic antigen CEA antibody [(NCRC16(AKA161))]). The wells were washed to remove unbound reactants, and the plate was blocked with a Phosphate Buffered Saline (PBS) solution containing 1% Tween 20 to fall within the range of the standard curve. After the blocking period, the wells were washed, and samples, controls, and standards were incubated in assigned triplicate wells. Unbound reactants were removed by washing, and a rabbit polyclonal antibody to a CEA detection antibody was added. After incubation, the wells were washed and incubated with 3,3',5,5'-tetramethylbenzidine (TMB), the peroxidase substrate. The substrate formed a colored product in the presence of the enzyme, reaction was stopped with 1 M phosphoric acid solution, and the absorbance was determined on a calibrated microplate reader. A calibration curve was generated from standards containing known concentrations of CEA, and the curve was used to determine the concentration of CEA in the samples. The quantity of CEA produced per virus particle was calculated from the concentration of CEA measured by ELISA, after adjusting for dilution and multiplicity of infection (MOI). The value determined in a similar manner for culture media alone was subtracted to compensate for background levels present in the media. The sample analysis is shown in **Table 2**.

TABLE 2

Sample ID	A ₄₅₀ - A ₅₄₀ Rep 1	Rep 2	Mean	SD	RSD	Blank Subtr	ELISA Dil'n Fact	CEA (ng/ml)	Avg CEA (ng/mL)	Total CEA (ng/mL)	CEA (ng/vp)
10-002917 Well 1 at 2.5E+04 vp	1.3387	1.2977	1.318	0.029	2.2%	1.244	1000	7,731	7,691	7,621	0.30
	0.7121	0.6789	0.693	0.023	3.4%	0.622	2000	7,797			
10-002917 Well 2 at 2.5E+04 vp	1.1717	1.1329	1.152	0.027	2.4%	1.078	1000	6,563	6,049	5,979	0.48
	0.6222	0.6151	0.619	0.005	0.8%	0.545	2000	6,975			
10-002917 Well 3 at 2.5E+04 vp	1.1659	2.0492	1.608	0.625	38.9%	1.534	1000	10,264	4,286	4,216	0.67
	0.6131	0.5946	0.604	0.013	2.2%	0.530	2000	6,815			
10-002917 Well 1 at 1.25E+04 vp	1.1051	1.0759	1.091	0.021	1.9%	1.017	1000	6,169	2,690	2,620	0.84
	0.5970	0.5716	0.584	0.018	3.1%	0.510	2000	6,602			
10-002917 Well 2 at 1.25E+04 vp	1.0652	1.0376	1.051	0.020	1.9%	0.977	1000	5,919	1,271	1,201	0.77
	0.5726	0.5770	0.575	0.003	0.5%	0.501	2000	6,506			
10-002917 Well 3 at 1.25E+04 vp	0.9731	0.9514	0.962	0.015	1.6%	0.888	1000	5,383	70	0	-
	0.5049	0.4970	0.501	0.006	1.1%	0.427	2000	5,716			
10-002917 Well 1 at 6.25E+03 vp	0.7601	0.7210	0.741	0.028	3.7%	0.667	1000	4,141	70	0	-
	0.4041	0.3881	0.396	0.011	2.9%	0.322	2000	4,566			
10-002917 Well 2 at 6.25E+03 vp	0.7157	0.7068	0.711	0.006	0.9%	0.637	1000	3,979	1,271	1,201	0.77
	0.3893	0.3843	0.387	0.004	0.9%	0.313	2000	4,465			
10-002917 Well 3 at 6.25E+03 vp	0.7360	0.7188	0.727	0.012	1.7%	0.653	1000	4,065	70	0	-
	0.3995	0.3807	0.390	0.013	3.4%	0.316	2000	4,499			
10-002917 Well 1 at 3.13E+03 vp	0.8920	0.8878	0.890	0.003	0.3%	0.816	500	2,483	70	0	-
	0.4573	0.4613	0.459	0.003	0.6%	0.385	1000	2,631			
10-002917 Well 2 at 3.13E+03 vp	0.8615	0.8544	0.858	0.005	0.6%	0.784	500	2,393	1,271	1,201	0.77
	0.4425	0.4406	0.442	0.001	0.3%	0.368	1000	2,538			
10-002917 Well 3 at 3.13E+03 vp	1.0518	1.0464	1.049	0.004	0.4%	0.975	500	2,953	70	0	-
	0.5519	0.5565	0.554	0.003	0.6%	0.480	1000	3,141			
10-002917 Well 1 at 1.56E+03 vp	1.8771	1.8616	1.869	0.011	0.6%	1.795	100	1,351	1,271	1,201	0.77
	1.1963	1.1695	1.183	0.019	1.6%	1.109	200	1,354			
10-002917 Well 2 at 1.56E+03 vp	1.7435	1.7436	1.744	0.000	0.0%	1.670	100	1,179	70	0	-
	1.0960	1.0788	1.087	0.012	1.1%	1.013	200	1,229			
10-002917 Well 3 at 1.56E+03 vp	1.7801	1.8098	1.795	0.021	1.2%	1.721	100	1,245	70	0	-
	1.1041	1.1263	1.115	0.016	1.4%	1.041	200	1,264			
10-002917 Well 1 Mock	1.2509	1.2278	1.239	0.016	1.3%	1.165	10	72	70	0	-
	0.7146	0.6952	0.705	0.014	1.9%	0.631	20	79			
10-002917 Well 2 Mock	1.2290	1.2382	1.234	0.007	0.5%	1.160	10	71	70	0	-
	0.7246	0.7133	0.719	0.008	1.1%	0.645	20	80			
10-002917 Well 3 Mock	0.9769	0.9750	0.976	0.001	0.1%	0.902	10	55	70	0	-
	0.5579	0.5454	0.552	0.009	1.6%	0.478	20	63			

EXAMPLE 4: Schedule, dose, route of immunization safety data

[00364] Initial pre-clinical studies were performed to evaluate and confirm that an Ad5 [E1-, E2b-] vector platform could express the antigen proteins on transfected cells. A-549 cells were transfected with vaccine platforms and analyzed by Western Blot Analysis (**FIG. 8**). Antigen proteins such as HIV-gag, HIV-pol, or HIV-nef were observed to be expressed on cells once they were transfected with the Ad5 [E1-, E2b-] vector platforms.

[00365] A dose response evaluation was performed using the Ad5 [E1-, E2b-] vector platform and demonstrated that 10^{10} virus particles (VP) is a dose that results in a desired CMI response against a transgene product in a murine model. CMI responses were assessed by utilizing an ELISpot assay to detect interferon- γ (IFN- γ) and IL-2 secreting cells (splenocytes) from spleens of mice. Furthermore, in murine and non-human primate (NHP) models, three immunizations using 10^{10} VP separated by two weeks to four weeks, respectively, resulted in the desired CMI responses. In mice, a greater degree of CMI responses were observed after multiple immunizations as compared with one immunization only (**FIG. 9**).

[00366] In a NHP model, the animals were rendered Ad5 immune by injection with wild type Ad5 virus. When Ad5 neutralizing antibody titers reached 1:50 or greater, which confirmed that the animals were immune to Ad5, they were immunized intradermally three times at 30 day intervals with Ad5-[E1-, E2b-]-gag at a dose of 1×10^{10} VP. 32 days after last vaccination and 124 days after the first immunization (wild type Ad5), the NAb titers were equal to or greater than 1:1000. After immunizations, the presence of robust CMI responses was detected, when peripheral blood mononuclear cells (PBMCs) of animals were assessed for IFN- γ and IL-2 secreting cells (**FIG. 10**).

[00367] In addition to the preliminary immunology studies performed in the initial vaccine trial in 3 NHP shown above, toxicity studies were also performed on the same NHP vaccinated with Ad5 [E1-, E2b-]-HIV gag. Animal temperatures and weights were assessed during the study period. The animals gained weight as they grew during the study period. No temperature differences were observed during the study period. Hematology studies were also performed on the vaccinated NHP. There appeared to be a small increase in the white blood cell count 2 weeks after the second vaccination that normalized thereafter.

[00368] Other than fluctuation in values, there appeared to be no other differences in hematology values during the course of the study. Chemistry values were also determined in the NHP during the course of the study. Alkaline phosphatase levels declined slightly during the course of the study but remained in the normal range. Albumin levels declined slightly during the course of the study but remained in the normal. There were no other differences observed in the blood chemistries during the course of the study. The route of immunization in this clinical study is chosen since the preponderance of DCs reside in the dermis.

[00369] A desired level of CMI response was induced using the Ad5 [E1-, E2b-] platform employing CEA and other transgenes. Using an Ad5 [E1-, E2b-]-CEA vector platform, both non-Ad5 immune and Ad5 pre-immunized mice were injected three times with the vaccine. After immunizations, the splenocytes from mice were assessed by ELISpot for IFN- γ secreting cells. Elevated CMI responses were observed after immunizations and the levels of CMI responses were similar in both non-Ad5 immune and Ad5 pre-immunized mice (FIG. 11). These results indicate that robust CMI responses can be induced despite the presence of pre-existing Ad5 immunity. A III clinical study was designed using three immunizations separated by three weeks via a needle subcutaneous delivery method.

Rationale for schedule, dose, route of administration

[00370] A clinical study design flowed from pre-clinical and clinical studies in animals and humans using the Ad5 [E1-, E2b-] vector platform. A dose response evaluation using the Ad5 [E1-, E2b-] vector platform was performed demonstrating that 10^{10} VPs is a dose which results in a desired CMI response against a transgene product in a murine model. Furthermore, in murine and non-human primate (NHP) models three immunizations using 10^{10} VP separated by two to four weeks respectively resulted in the desired CMI. The route of immunization is chosen since a preponderance of dendritic cells (DCs) reside in the dermis. Using this premise, multiple murine and NHP studies were performed using a subcutaneous injection protocol. A desired level of circulating CMI was induced using the Ad5 [E1-, E2b-] platform employing CEA and other transgenes. A phase III clinical study followed using three immunizations separated by three weeks via a needle subcutaneous (SQ) delivery method continuing immunotherapy treatment every three months until removed from study for any reason including death.

Summary

[00371] The cDNA sequence containing the modified CEA with the CAP1(6D) mutation was produced. Clinical grade Ad5 [E1-, E2b-]-CEA(6D) was constructed and manufactured using the E.C7 cell line. A total of 34 patients (32 colorectal cancer patients, one bladder cancer patient, and one lung cancer patient) were entered into the Phase I/II clinical study under IND14325. The majority received all three scheduled immunotherapy treatments with Ad5 [E1-, E2b-]-CEA(6D). Five patients who stopped immunotherapy early did so due to significant disease progression. RECIST 1.0 criteria using CT or MRI scans obtained at baseline and after treatments were completed. Toxicity was assessed according to the National Cancer Institute Common Terminology Criteria for Adverse Events (CTCAE) version 4.0. Peripheral blood CEA levels, hematology, serum chemistries, and anti-nuclear antibody titers were compared between baseline and 9 weeks following the initiation of immunotherapy. Survival was measured from the day of the first immunization until death from any cause.

[00372] A total of 94 treatments were administered to patients. No dose limiting toxicity or serious adverse events (SAE) that resulted in treatment discontinuation at any treatment dose level. There was only one significant change in a blood hematology value. As a group, the basophil count was

significantly lower at week 9, three weeks after treatment ended. However, this value remained in the normal range for basophil counts and, overall, there appeared to be no significant biological effect. With a median follow-up of 7.4 months, all 34 patients as a group (cohorts 1, 2, 3/phase II, and cohort 5) experienced a 12-month survival proportion of 41.4%. Of the 34 patients entered into the study, 28 patients received the three immunotherapy treatments and experienced a 12-month survival proportion of 55%. For the colorectal adenocarcinoma patients, 27 patients received the three immunotherapy treatments and experienced a 12-month survival proportion of 53%. A dose response to increasing levels of Ad5 [E1-, E2b-]-CEA(6D) was observed with the highest cell-mediated immune (CMI) responses occurring in patients that received the highest dose of 5×10^{11} VP of Ad5 [E1-, E2b-]-CEA(6D). When the highest CEA specific CMI responses were compared with pre-existing or vector induced Ad5 NAb activity, there was no correlation between levels of CEA specific CMI and Ad5 neutralizing antibody (NAb) level. These clinical trial data lead us to believe that there is sufficient data to advance to a randomized Phase III trial for the treatment of metastatic colorectal adenocarcinoma with overall survival as the clinical endpoint.

Protocol schema and patient treatment.

[00373] The clinical study was performed under an FDA-approved Investigational New Drug Exemption and registered at ClinicalTrials.gov. Participants were recruited from medical oncology clinics at Duke University Medical Center (NC) and Medical Oncology Associates (WA). Patients provided informed consent approved by the respective Institutional Review Boards (IRB). Eligibility requirements included metastatic cancer expressing CEA and adequate hematologic, renal, and hepatic function. Trial participants were required to have received treatment with standard therapy known to have a possible overall survival benefit or refused such therapy. Exclusion criteria included chemotherapy or radiation within the prior 4 weeks, history of autoimmune disease, viral hepatitis, HIV, or use of immunosuppressive agents. Patients who had been receiving bevacizumab or cetuximab for at least 3 months prior to enrollment were permitted to continue receiving these antibodies. Prior CEA immunotherapy was permitted. The study employed a standard 3 + 3 dose escalation strategy with dose limiting toxicities (DLT) defined as grade 3 or 4 major organ toxicity. The Ad5 [E1-, E2b-]-CEA(6D) doses were delivered to patients as follows: cohort 1: dose of 1×10^9 VP in 0.5 mL subcutaneously (SQ) in the same thigh every 3 weeks for 3 immunizations.; cohort 2: dose of 1×10^{10} VP in 0.5 mL SQ every 3 weeks for 3 treatments; cohort 3: dose of 1×10^{11} in 0.5 mL SQ every 3 weeks for 3 treatments. Following establishment of the dose of 1×10^{11} VP as safe, an additional 12 patients received Ad5 [E1-, E2b-]-CEA(6D) at this dose and schedule (phase II cohort). After completing the phase II cohort, an additional cohort (cohort 5) of six (6) patients received a dose of 5×10^{11} VP in 2.5 mL SQ every 3 weeks for 3 treatments to determine safety of the highest achievable dose. PMBC were collected from patients just prior to the immunizations at weeks 0, 3, 6, and three weeks following the last treatment. The PBMC

were frozen in liquid nitrogen until ELISpot assays were performed. In cohort 5, fresh PBMC were analyzed in preliminary flow cytometry assays for polyfunctional CD8+ T lymphocytes.

Assessment of clinical activity.

[00374] Clinical activity was assessed according to Response Evaluation Criteria in Solid Tumors (RECIST 1.0 criteria) using computed tomography (CT) or magnetic resonance imaging (MRI) scans obtained at baseline and after treatments were completed. Toxicity was assessed according to the National Cancer Institute Common Terminology Criteria for Adverse Events (CTCAE) version 4. Peripheral blood CEA levels, hematology, serum chemistries, and anti-nuclear antibody titers were compared at baseline and 3 weeks following the final treatment. Survival was measured from the day of the first immunization until death from any cause

Analysis of CMI responses by ELISpot Assay.

[00375] An ELISpot assay for IFN- γ secreting lymphocytes was performed. Briefly, isolated PBMCs (2×10^5 cells/well) from individual patient samples were incubated 36–40 h with a CEA peptide pool (15mers with 11aa overlap covering full length CEA with the 6D modification; 0.1 μ g/well) to stimulate IFN- γ producing T-cells. CMI responses to Ad5 were determined after exposure of patient PBMC to Ad5-null (empty vector). Cells stimulated with concanavalin A (Con A) at a concentration of 0.25 μ g/well served as positive controls. Colored spot-forming cells (SFC) were counted using an Immunospot ELISpot plate reader and responses were considered to be positive if 50 SFC were detected/ 10^6 cells after subtraction of the negative control and SFC were ≥ 2 -fold higher than those in the negative control wells

Determination of Ad5 neutralizing antibody (NAb) titers.

[00376] Endpoint Ad5 NAb titers were determined. Briefly, dilutions of heat inactivated test sera in 100 μ L of DMEM containing 10% fetal calf serum were mixed with 4×10^7 VP of Ad5 [E1-]-null and incubated for 60 minutes at room temperature. The samples were added to microwells containing HEK293 cells cultured in DMEM containing 10% heat inactivated calf serum at 2×10^3 cells/well for 24 hours at 37 °C in 5% CO₂. The mixture was incubated for an additional 72 hours at 37 °C in 5% CO₂. An MTS tetrazolium bioreduction assay was used to measure cell killing and endpoint Ad5 NAb titers. Endpoint titers with a value less than 1:25 were assigned a value of 0.

Statistics.

[00377] Statistical analyses comparing immune responses were performed employing the Mann-Whitney test (PRISM, Graph Pad). Survival comparisons were performed employing Kaplan-Meier plots (PRISM, Graph Pad). Ad5 NAb titer and CEA-specific CMI were analyzed as continuous variables. The association of Ad5 NAb titer with change in CEA-specific CMI was tested with the Spearman correlation coefficient. The association of Ad5 NAb titer with survival was tested with the Wald test of the proportional hazards model. All tests used a 2-sided α of 0.05.

Demographics: All Patients.

[00378] Thirty two patients with metastatic colorectal cancer, one with lung cancer and one with bladder cancer, median age 58 (range 38-77), who had failed a median of three prior chemotherapeutic regimens (range: 2->5), had a median performance status of 90% (range 70%-100%), and had a median of three sites of metastatic disease (range 1-5), were enrolled (Table 2). The majority of patients were able to receive all three immunizations. Five patients who stopped immunizations prior to completion of all three treatments did so due to significant disease progression. Patient demographics are shown in

Table 3.**TABLE 3**

Patient ID/ Cohort	Dose (VP)	Dx	Age	Sex	KPS	# prior CTx	Mets (# sites)	# doses	⁺⁺ Status after tx	Survival (Mos)	base-line	CEA Week 9
002/1	10 ⁹	C	67	M	70	>3	4	3	PD	3 (-)	98.8	867.4
003/1	10 ⁹	R	63	M	100	5	2	3	PD	9 (-)	195.1	472.2
004/1	10 ⁹	C	53	F	100	2	3	3	PD	11 (-)	65.4	196.8
005/2	10 ¹⁰	C	60	M	100	3	3	3	SD	12 (+)	2.5	3.7 (7 month follow-up)
007/2	10 ¹⁰	C	52	M	80	2	5	1	PD	1 (-)	120.7	Not Done
008/2	10 ¹⁰	C	42	F	100	3	3	3	PD	12 (+)	3.0	3.1
010/2	10 ¹⁰	C	58	M	90	3	3	3	PD	12 (-)	7.1	5.8
011/3	10 ¹¹	R	50	M	100	5	1	3	PD	12 (+)	21.0	25.9
012/3	10 ¹¹	C	48	M	100	1	2	3	PD	12 (+)	5.8	18.4
013/3	10 ¹¹	R	62	M	100	3	2	2	PD	4 (-)	172.9	Not Done
500/3	10 ¹¹	C	55	M	80	4	3	3	PD	12 (+)	3.2	11.5
015/3	10 ¹¹	C	58	F	80	3	4	3	PD	10 (-)	2.0	2.4
016/3	10 ¹¹	C	53	F	100	3	4	3	PD	6 (-)	6.1	12.7
017/3*	10 ¹¹	R	52	F	90	3	2	3	PD	3 (-)	204.8	Not Done
501/II	10 ¹¹	R	54	M	90	1	1	3	PD	12 (+)	17.1	96.4
502/II	10 ¹¹	C	66	F	80	1	2	2	PD	3 (-)	2549.5	Not Done
503/II	10 ¹¹	Bl	73	M	70	4	5	1	PD	0.25 (-)	Not Done	Not Done
019/II	10 ¹¹	C	69	M	90	1	3	3	PD	12 (+)	264.3	638.0
020/II^	10 ¹¹	C	59	M	100	5	4	3	SD	12 (+)	2.2	2.2
021/II^	10 ¹¹	C	51	F	100	4	3	3	PD	12 (+)	2.0	2.7
506/II	10 ¹¹	C	77	F	80	2	2	3	PD	3 (-)	16.5	38.2
023/II	10 ¹¹	C	51	F	100	3	4	3	PD	4 (-)	32.4	211.4
504/II	10 ¹¹	C	57	M	90	3	3	3	PD	12 (+)	424.7	2073.6
507/II	10 ¹¹	R	58	M	90	2	2	3	PD	12 (+)	<0.5	0.6
508/II	10 ¹¹	L	67	M	100	2	0	3	Unknown	12 (+)	109.2	Not Done
024/II	10 ¹¹	C	67	M	90	2	3	3	PD	12 (+)	7.8	6.4
025/II	10 ¹¹	C	62	F	100	2	4	3	PD	7 (-)	391.2	Not Done
026/II	10 ¹¹	C	53	M	100	3	2	2	PD	4 (-)	4057.5	7859.1 (treatment #2)
030/5	5x10 ¹¹	C	38	M	90	4	3	3	PD	8 (+)	9.2	18.7
031/5	5x10 ¹¹	R	72	F	90	4	2	3	SD	7 (+)	3.9	5.6
032/5@	5x10 ¹¹	R	53	M	90	4	3	3	PD	6 (-)	31.9	75.4
033/5@	5x10 ¹¹	R	48	F	90	>3	2	3	PD	5 (-)	21.3	21.1
034/5	5x10 ¹¹	C	62	M	100	5	4	3	PD	6 (+)	1.9	2.4
035/5	5x10 ¹¹	C	60	F	90	3	5	2	PD	2 (-)	9.5	Not Done

Dx = diagnosis (Bl= bladder cancer; C= colon cancer; L= lung cancer; R= rectal cancer) KPS=Karnofsky

Performance Status *concurrent cetuximab; ^concurrent bevacizumab; @ concurrent panitumumab

++ Represents disease status at 9 weeks post-initiation of immunizations

PD = Progressive Disease; SD = Stable Disease

(+) Alive; (-) Dead at last follow-up; survival rounded off to nearest month

Demographics: Colorectal Adenocarcinoma Patients

[00379] Thirty two patients, median age 57.5 (range 38-77) with metastatic colorectal cancer, who had failed a median of three prior chemotherapeutic regimens (range: 2->5), had a median performance status of 90% (range 70%-100%), and had a median of three sites of metastatic disease (range 1-5), were enrolled (Table 2). The majority was able to receive all three immunizations. Four patients who stopped immunizations early did so due to significant disease progression. The colorectal adenocarcinoma patient demographic compares favorably with previously published studies of patients with chemotherapy-refractory colorectal cancer.

Adverse Effects

[00380] A total of 94 immunization treatments were administered to all patients. There was no dose limiting toxicity and no serious adverse events that resulted in treatment discontinuation at any vaccine dose level. The most common toxicity (Table 3) was a self-limited, injection site reaction. Other reactions that occurred at a low frequency include fever, flu-like symptoms, anorexia, chills, nausea, and headache. These symptoms were also self-limiting and did not require intervention other than symptomatic measures such as acetaminophen.

Summary of Hematology, Chemistry, and ANA values pre and post treatment

[00381] Biological effects of Ad5 [E1-, E2b-]-CEA(6D) injections were monitored by recording blood hematology, chemistry, and anti-nuclear antibody (ANA) values of individual patients in case record forms (CRFs). Of 34 total patients entered into the trial, 28 received all three treatments with Ad5 [E1-, E2b-]-CEA(6D). For the 28 patients which received all three treatments, the blood hematology, chemistry, and ANA values at week 0 (prior to first treatment) were compared with those obtained at week 9 (three weeks after the third treatment). As shown in Table 4 below, there were no significant changes in chemistry or ANA values after treatments with Ad5 [E1-, E2b-]-CEA(6D). There was only one significant change in the blood hematology values. The basophil count was significantly (P=0.0403) lower at week 9 after treatments. However, this value remained in the normal range for basophil counts and overall there were no significant biological effects.

Clinical Outcomes

[00382] CEA levels at baseline and week 9 were assessed in patients. Among those with CEA levels available at baseline and follow-up, three (patients 010, 020, and 024) had no increase in CEA levels at the end of the immunization period while the remaining patients showed increased CEA levels. There

were three patients with stable disease who remained so during the 9 week study period. All other patients experienced some level of progressive disease (Table 2). Of the seven patients in cohorts 1 and 2, there were five deaths and two patients remained alive at 12 months following the initiation of immunization. Of the 21 patients in cohort 3 and phase II, there were 10 deaths and all the remaining 11 patients were alive at 12 months, respectively. Of the six patients in cohort 5, there were three deaths and three patients were alive at 6, 7, and 8 months, respectively.

[00383] Of the 34 patients enrolled into the study, two patients received one treatment, four patients received two immunization treatments, and the remaining 28 patients received all three immunization treatments. All patients were followed for survival and Kaplan-Meier plots and survival proportions performed (PRISM software). Patient deaths were determined by information gathered from the social security death index (SSDI) database and clinical charts.

[00384] The seven patients in cohorts 1 and 2 experienced a 12-month survival proportion of 29% (**Figure 12A**). Of the patients in cohorts 1 and 2, patient 004 survived 11 months and received additional post-immunization treatments with bevacizumab, folfox, and xeloda. Patient 003 survived nine months and received irradiation treatment after Ad5 [E1-, E2b-]-CEA(6D)immunizations. Patient 005, alive at 12+ months, received irradiation treatment and entered another clinical trial after immunizations. Patient 010 survived up to 12 months and entered two clinical trials after immunizations. Patients 002, 007, and 008 received no further treatments after immunizations and survived 3, 1, and 12+ months, respectively.

[00385] The 21 patients in cohort 3 and phase II experienced a 12-month survival proportion of 48% (**FIG. 12B**). Of the patients in cohort 3 and phase II, one patient (017) received concurrent cetuximab during immunizations. Patients 020 and 021 received concurrent bevacizumab during immunizations. Patient 011 surviving over 12 months received radiation treatment after Ad5 [E1-, E2b-]-CEA(6D) immunizations. Patients 012 and 016 survived over 12 months and 6 months, respectively, and received additional chemotherapy treatment after immunizations. Patient 013 survived 4 months and received treatment with nexavar after immunizations. Patient 015 survived 10 months and received follow-on treatment with cetuximab. Patient 019 survived over 12 months and received treatment with bevacizumab and xeliri after protocol immunizations. Patient 020 survived over 12 months and received treatment with bevacizumab after immunizations. Patient 021 survived over 12 months and received follow-on treatment with bevacizumab and xeloda. Patient 500 survived over 12 months and received treatment with cetuximab and xeloda and entered a clinical trial after immunizations. Patient 501 survived over 12 months and received treatment with cetuximab and irinotecan after Ad5 [E1-, E2b-]-CEA(6D) immunizations. Patient 508 has survived over 12 months; however, further data on the characteristics of this patient was unable to be obtained. Patients 017, 023, 024, 025, 026, 502, 504, 506, and 507 received no further treatment after immunizations and survived 3, 4, 12+, 7, 4, 3, 12+, 3, and 12+ months, respectively (+ means still alive at the time of writing).

[00386] The six patients in cohort 5 experienced a 12-month survival proportion of 50% (**FIG. 12C**). Of the patients in this cohort, one patient (030) is currently alive at 8 months and received treatment with pazopanib and threshold 302 chemotherapy after Ad5 [E1-, E2b-]-CEA(6D) immunizations. Patient 031 is currently alive at 7 months and has not received further treatment after immunizations. Patient 032 received concurrent panitumumab, survived 6 months, and received treatment with folfox after immunizations. Patient 033 received concurrent panitumumab, survived 5 months with no additional therapy after immunizations. Patient 034 is currently alive at 6+ months and received radiation and treatment with xeloda after immunizations. Patient 035 received two treatments and survived 2 months.

[00387] With a median survival of 7.4 months, all 34 patients as a group (cohorts 1, 2, 3/phase II, and cohort 5) experienced a 12-month survival proportion of 41% (**FIG. 12D**). Of the 34 patients entered in to the study, 28 patients received the three immunization treatments and experienced a 12-month survival proportion of 55% (**FIG. 13A**) with a median survival of 10.625 months. For the colorectal adenocarcinoma patients, 27 patients received the three immunization treatments and experienced a 12-month survival proportion of 53% (**FIG. 13B**) with a median survival of 10.00 months.

Evaluation of immune parameters in treated metastatic colorectal cancer patients

[00388] A secondary objective was to evaluate CEA specific immune responses following immunization treatments with the product.

[00389] Dendritic cells were generated from the peripheral blood mononuclear cells (PBMCs) of a prostate cancer patient (HLA-A2⁺ and -A24⁺) enrolled in a clinical trial employing a PSA-TRICOM vaccine in combination with ipilimumab; using PBMCs from this patient post-vaccination, individual T-cell lines specific for CEA, MUC1, and brachyury were unable to be established. Briefly, PBMCs were isolated using lymphocyte separation medium gradient, resuspended in AIM-V medium (2x10⁷ cells) and allowed to adhere in a 6-well plate for 2 hours. Adherent cells were cultured for 5 days in AIM-V medium containing 100ng/ml of recombinant human (rh) GM-CSF and 20ng/ml of rhIL-4. The culture medium was replenished every 3 days.

[00390] As determined by ELISA, no antibody activity directed against CEA was observed. CMI responses in colorectal cancer patients treated in cohort 1, cohort 2, cohort 3/Phase II, and cohort 5 were assessed. PBMCs were isolated prior to Ad5 [E1-, E2b-]-CEA(6D) treatment and after all treatments as well as three weeks following the last treatment from patients. CEA specific ELISpot assays were performed on PBMCs to determine the numbers of interferon γ (IFN- γ) secreting lymphocytes after exposure to CEA peptides *in vitro*. The highest CMI responses during immunizations were determined, regardless of time point (weeks 3, 6, or 9) in the patients treated in cohort 1, cohort 2, cohort 3/Phase II, and cohort 5. This analysis revealed a dose response to increasing levels of product. The highest CMI levels occurred in patients that received the highest dose of 5x10¹¹ VP (Cohort 5) (**FIG. 14**).

Determination of Induced CMI Responses to CEA.

[00391] ELISpot analysis was performed on cryopreserved PBMC samples drawn before each immunization and after completion of the final immunization to assess CEA-specific CMI responses. A dose response effect with the highest magnitude CEA-specific CMI responses occurring in patients who received the highest dose of Ad5 [E1-, E2b-]-CEA(6D) was observed(**FIG. 14**). Of the doses received, 0/3 (0%) patients in cohort 1 exhibited positive CEA-directed CMI responses, 1/4 (25%) patients in cohort 2 exhibited positive CEA-directed CMI responses, 10/19 (53%) patients in cohort 3/phase II exhibited positive CEA-directed CMI responses, and 4/6 (67%) patients in cohort 5 exhibited positive CEA-directed CMI responses. The time course of induction of CEA-specific CMI (**Figure 15**) demonstrated that there may be plateau in the magnitude of CEA CMI prior to the last dose. In the largest group of patients who received the same dose (cohort 3 plus phase II), a significant increase over baseline in the average CEA-directed CMI responses at the 6 week evaluation ($P<0.05$, Mann-Whitney test) was observed, averaging 94 SFC/106 PBMC, which increased further by the 9 week evaluation (**FIG. 15**). One patient (patient ID 13) had a highly elevated baseline CEA-specific immune response (1100 SFC) and had elevated CMI at week six (2305 SFC) but did not return for 9-week evaluation and was not included in CEA CMI analysis.

[00392] Ad5 NAb and CMI against Ad5 was also measured and correlated with CEA-specific CMI. Each patient had their serum and PBMC sample tested at baseline (prior to treatment) and at 9 weeks after completion of 3 treatments. Nineteen of 31 colorectal cancer patients (61%) tested in this study had Ad5 neutralizing activity in serum samples prior to the onset of treatment with Ad5 [E1-, E2b-]-CEA(6D). The mean pre-treatment Ad5 NAb titer value obtained among all patients was $1:189 \pm 1:71$ SEM (geometric mean 1:21) and the mean pre-treatment Ad5 NAb titer among seropositive patients was $1:308 \pm 1:108$ (geometric mean 1:146). Analysis of serum samples from patients who received 3 immunizations revealed Ad5 NAb titers that were significantly increased ($P<0.0001$, *Mann-Whitney test*) by week 9 (mean $1:4767 \pm 1:1225$ SEM (geometric mean 1:1541) when compared with their respective baseline values (**FIG. 16**). Analysis of PBMC for CMI responses to Ad5 also revealed a significant increase ($P<0.01$, Mann-Whitney test) in Ad5 directed CMI responses after immunizations with Ad5 [E1-, E2b-]-CEA(6D) (**FIG. 16**).

[00393] Comparison of week 9 CEA-directed CMI responses from patients with low baseline pre-existing Ad5 immunity ($\text{Ad5 NAb} \geq 200$) verses those with high baseline Ad5 immunity ($\text{Ad5 NAb} > 200$) revealed no significant difference in immune responses ($P>0.4$, Mann Whitney test) (**FIG. 17**). Further, when the highest CEA specific CMI responses were compared with pre-existing or vector induced Ad5 NAb activity, there was no correlation between levels of CEA CMI and Ad5 NAb activity (**FIG. 17**). These data indicate that immunizations with Ad5 [E1-, E2b-]-CEA(6D) were not only able to overcome self-tolerance, but were also able to induce CEA-specific immune responses in colorectal cancer patients

despite the presence of pre-existing and/or immunization induced Ad5 immunity. Together these clinical trial data support the advancement to a Phase III clinical trial with overall survival as the primary endpoint.

Adverse Effects, Hematology, Chemistry, and ANA values

[00394] In this Phase I/II trial, the Ad5 [E1-, E2b-]-CEA(6D) was demonstrated to be suitable to be manufactured to scale, as well be easily and repeatedly administered by conventional subcutaneous injection techniques. The most common adverse effects were site of injection reactions and flu-like symptoms consisting of fever, chills, headache, and nausea. There was no impact on blood hematology or serum chemistries and, overall, the treatments were well tolerated. Specifically, no SAE were noted, and no treatments were stopped due to adverse events, indicating that a dose limitation to use of Ad5 [E1-, E2b-]-CEA(6D) in this clinical application had not been met.

[00395] These data suggest that patients with advanced colorectal cancer which are treated with Ad5 [E1-, E2b-]-CEA(6D) do not have serious adverse effects and may experience extension of life even if they have pre-existing immunity to Ad5. The results of this trial were encouraging enough to advance to a large, randomized, single agent trial. The observation that some of the patients experienced an increase of CMI which is dose dependent, indicates that this may play a role in their clinical outcome.

Table 4. Adverse Events

Adverse Events	# Events	Unrelated/Unlikely	Possible	Probably/Definite	*Grade	**% Incidence
Injection Site Reaction	21			21	G1 (19); G2 (2)	22.3
Pain (all types)	17	17			G1 (8); G2 (7); G3 (2)	18.1
Fever	10	4	2	4	G1 (7); G2 (3)	10.6
Flu-like symptoms	10	3	5	2	G1 (9); G2 (1)	10.6
Fatigue	8	6	2		G1 (5); G2 (2); G3 (1)	8.5
Shortness of Breath	6	6			G1 (3); G2 (3)	6.4
Anorexia	5	4	1		G1 (3); G2 (2)	5.3
Chills	5	1	1	3	G1 (5)	5.3
Nausea	5	4	1		G1 (5)	5.3
Constipation	5	5			G1 (3); G2 (2)	5.3
Edema	5	5			G1 (3); G2 (2)	5.3
Vomiting	4	4			G1 (4)	4.3
Hypertension	3	3			G1 (2); G2 (1)	3.2
Anemia	3	3			G1 (1); G2 (1); G3 (1)	3.2
Cough	2	2			G1 (2)	2.1
Depression	2	2			G1 (2)	2.1
Diarrhea	2	2			G1 (2)	2.1
Headache	2	1	1		G1 (2)	2.1
Hypoalbuminemia	2	2			G1 (1); G2 (1)	2.1
Hypokalemia	2	2			G1 (1); G2 (1)	2.1

Adverse Events	# Events	Unrelated/Unlikely	Possible	Probably/Definite	*Grade	**% Incidence
Pleural Effusion	2	2			G2 (1); G3 (1)	2.1
Alkaline Phosphatase Increase	2	2			G1 (1); G3 (1)	2.1
Myalgia	2		2		G1 (2)	2.1
Night Sweats	2	2			G1 (1); G2 (1)	2.1
Sleep	2	2			G1 (2)	2.1
Low Magnesium	2	2			G1 (2)	2.1
Abdominal Bloating	1	1			G1 (1)	1.1
Abdominal Distention	1	1			G3 (1)	1.1
Abdominal Swelling	1	1			G2 (1)	1.1
Abdominal Wound	1	1			G2 (1)	1.1
ALT Increase	1	1			G1 (1)	1.1
AST Increase	1	1			G2 (1)	1.1
Biliary Obstruction	1	1			G3 (1)	1.1
Bowel Obstruction	1	1			G3 (1)	1.1
Cold	1	1			G1 (1)	1.1
Dyspnea	1	1			G3 (1)	1.1
Dysuria	1	1			G1 (1)	1.1
Frequent Urination	1	1			G1 (1)	1.1
GI Disorder	1	1			G3 (1)	1.1
Extra Pyramidal Movements	1	1			G1 (1)	1.1
Insomnia	1	1			G1 (1)	1.1
Herpes Simplex	1	1			G1 (1)	1.1
Hypotension	1	1			G1 (1)	1.1
Loss of Appetite	1	1			G1 (1)	1.1
Low White Blood Cells	1	1			G1 (1)	1.1
Numbness/Sensation in Fingertips	1	1			G1 (1)	1.1
Onset of Menses	1	1			G1 (1)	1.1
Poor Quality Sleep	1	1			G1 (1)	1.1
Presyncope	1	1			G2 (1)	1.1
Pruritis	1	1			G1 (1)	1.1
Rash-Right Lower Eye Lid	1	1			G1 (1)	1.1
Red/Swelling Right Upper Eyelid	1	1			G1 (1)	1.1
Renal Calculi	1	1			G2 (1)	1.1
Runny Nose	1		1		G1 (1)	1.1
Shallow Breathing	1	1			G1 (1)	1.1
Skin Rash	1	1			G1 (1)	1.1
Vaginal Discharge	1	1			G1 (1)	1.1
Concentration	1				G1 (1)	1.1
Weight Loss	1	1			G2 (1)	1.1

Adverse Events	# Events	Unrelated/Unlikely	Possible	Probably/Definite	*Grade	**% Incidence
Arthritis Joint Inflammation	1	1			G1 (1)	1.1
Flushing	1		1		G1 (1)	1.1
Acute Renal Failure	Disease progression				G3 (1)	1.1

* Parenthesis () indicates numbers of events.

**Based on 94treatments

TABLE 5. Hematology, Chemistry, and ANA values

Hematology Test	Week 0 value (Mean \pm SEM)	Week 9 value (Mean \pm SEM)
Hgb (g/dL)	13.09 \pm 0.313	12.48 \pm 0.413
Hct (%)	39.63 \pm 0.875	37.92 \pm 1.140
Plts (x10 ⁹ /L)	225.1 \pm 20.76	247.3 \pm 23.57
WBC (x10 ³ /mm ³)	6.81 \pm 0.532	8.21 \pm 0.741
Neutrophils (%)	64.46 \pm 2.068	67.28 \pm 3.268
Lymphocytes (%)	23.23 \pm 1.874	18.34 \pm 2.071
Monocytes (%)	8.86 \pm 0.462	7.68 \pm 0.569
Eosinophils (%)	3.97 \pm 0.677	3.16 \pm 0.685
Basophils (%)	0.52 \pm 0.056	0.38 \pm 0.048
Chemistry Test	Week 0 value (Mean \pm SEM)	Week 9 value (Mean \pm SEM)
Na (mEq/L)	139.2 \pm 0.424	137.9 \pm 0.718
K (mEq/L)	3.90 \pm 0.085	3.80 \pm 0.073
Cl (mEq/L)	105.0 \pm 0.561	103.3 \pm 1.061
CO ₂ (mEq/L)	27.8 \pm 0.374	27.63 \pm 0.458
BUN (mg/dL)	17.0 \pm 1.136	17.1 \pm 1.611
Creatinine (mg/dL)	0.81 \pm 0.046	0.86 \pm 0.054
Glucose (mg/dL)	121.8 \pm 7.458	123.5 \pm 7.885
Ca (mg/dL)	8.84 \pm 0.075	8.87 \pm 0.073
Total protein (g/dL)	6.95 \pm 0.078	6.67 \pm 0.100
Albumin (g/dL)	3.78 \pm 0.085	3.62 \pm 0.113
AST (U/L)	31.71 \pm 3.846	31.88 \pm 3.506
ALT (U/L)	27.83 \pm 4.228	25.67 \pm 3.414
Alkaline phosphatase (U/L)	107.0 \pm 13.30	124.0 \pm 17.40
Bilirubin (mg/dL)	0.78 \pm 0.071	0.75 \pm 0.079
*ANA Test	Week 0 value (Mean \pm SEM)	Week 9 value (Mean \pm SEM)
Titer	103.3 \pm 51.04	123.3 \pm 50.56

*Values represent inverse of the titer and are from patients with positive values.

Discussion

[00396] Adenoviral vectors have significant potential for use as cancer therapeutic vaccines because of their propensity to induce robust adaptive immune responses specifically against transgene products in general. However, recombinant first generation Ad5 [E1-] vectors used in homologous prime/boost regimens have been greatly limited in their potential efficacy due to the presence of pre-existing Ad5

immunity as well as vector induced immunity. Specifically, Ad5-directed immunity mitigates immune responses to TAA that have been incorporated into earlier generation Ad5 [E1-] based platforms. The Ad5 [E1-, E2b-] platform utilized in the present study was intended to accommodate a homologous prime-boost regimen, by avoiding presentation of antigens that are the targets of pre-existing Ad5 immunity. Since CEA has been identified as one of the priority cancer antigens by the National Cancer Institute, this TAA was investigated as a transgene to be incorporated into the new Ad5 [E1-, E2b-] vector platform for use as a cancer therapeutic vaccine. CEA expression in adults is normally limited to low levels in the gastrointestinal epithelium, whereas, CEA is over-expressed in adenocarcinomas of the colon and rectum and in many breast, lung, and pancreas cancers. The HLA A2 restricted CAP1(6D) modification of CEA was chosen because compared with the wild type CAP1 epitope, CAP1(6D) can enhance the sensitization of CTLs and has been included in recent CEA-based vaccine constructs. Although HLA type was not tested for because a full length CEA was used that is not HLA-restricted, A*0201 is the allele observed most frequently in Caucasians (allele frequency 0.2717) and is common in other populations. However, it is possible to test patients for HLA type and utilize the relationship between HLA type and clinical and/or CMI responses.

[00397] Multiple subcutaneous immunizations employing three administrations of a single dose level (1×10^{10} VP) of this class of Ad5 vaccine expressing the TAA CEA, (Ad5 [E1-, E2b-]-CEA(6D)) were tested in a pre-clinical murine model of CEA expressing cancer. In mice with pre-existing Ad5 immunity, the induction of potent CEA directed CMI responses were demonstrated that resulted in anti-tumor activity and noted that these CMI and anti-tumor responses were significantly greater than those responses induced by a current generation Ad5 [E1-] based vector vaccine. In additional animal models (both cancer and infectious disease targeted) it was demonstrated that multiple subcutaneous immunizations with vaccines based on the new Ad5 [E1-, E2b-] platform induce CMI responses that were superior to those of current generation Ad5 [E1-] based vaccines, can overcome the barrier of Ad5 immunity, and can be utilized in multiple immunization regimens requiring a generation of robust CMI responses. The greatest magnitude of CEA-directed CMI responses occurred in patients receiving the highest dose of the vector. A CEA-directed CMI response was induced in a dose-responsive manner despite the presence of pre-existing and/or vector induced Ad5 immunity. No CEA directed antibody responses were observed either pre- or post-vaccination employing an ELISA technique. A population of polyfunctional CD8+ T-cells (those that secrete more than one cytokine when activated) after immunizations were also observed, a sign of greater functionality of T-cells induced by the vaccine. These data support the use of the Ad5 [E1 , E2b-]-CEA(6D) vector in homologous prime-boost regimens designed to induce and increase CEA-directed CMI responses in patients with advanced colorectal adenocarcinoma, as well as any number of other vaccine amenable diseases or applications.

[00398] As compared to earlier generation Ad5 [E1-] vectors containing deletion in the early 1 (E1) gene region, the Ad5 [E1-, E2b-] vector platform with additional deletions in the early 2b (E2b) gene region exhibits significantly reduced inflammatory responses directed at the vector. This can result in longer transgene expression and a reduction in elimination of transgene expressing cells (e.g., antigen presenting cells) that would otherwise occur due to induced inflammatory responses. Since Ad5 late gene antigen expression is significantly reduced as compared to earlier generation Ad5 platforms, this could enable the Ad5 [E1-, E2b-] platform to evade Ad5 immune mediated neutralizing activity for significantly longer periods of time resulting in greater longevity and amplification of TAA expression. In addition, an E2b gene product, a polymerase, is a target of human cellular memory immune responses to Ad5 infection and its elimination from the vaccine could be furthering its capability in the setting of pre-existing Ad5 immunity. Without being bound by theory, the extended and/or greater expression of TAA by the vector in this milieu could result in a more effective immune response against the target antigen. However, it is also possible that this vector configuration produces better transgene expression, different biodistribution, or different innate/adaptive immune effects that impact the effectiveness of this vector, rather than escape from pre-existing immunity.

[00399] Of interest is the observation that treated patients in this study exhibited favorable survival probability. Overall, all 25 patients treated at least 2 times with Ad5 [E1-, E2b-]-CEA(6D) exhibited a 12-month survival probability of 48% and this was achieved despite the presence of significant levels of pre-existing Ad5 neutralizing antibody titers. Without being bound by theory, by engaging the patient's immune system, active immunotherapeutics, such as Ad5 [E1-, E2b-]-CEA(6D), could induce continuous immunologic anti-tumor responses over a long period of time that could result in a "deceleration" or alteration in specific aspects of the rapid growth rate or spread of the tumor not measured by standard response assessments. Indeed, slower tumor progression in Ad5 immune mice harboring established CEA-expressing tumors following treatment with Ad5 [E1-, E2b-]-CEA(6D) were observed. Moreover, it has been noted that overall survival might be the only true parameter for determination of clinical efficacy of any potential cancer (immune) therapy.

EXAMPLE 5: Clinical study of CEA (6D) immunotherapeutic in metastatic colorectal cancer

[00400] Phase I/II clinical trial in mCRC using Ad5 [E1-, E2b-]-CEA(6D): The objectives of this first in man phase I/II dosing trial were to assess safety, evaluate CEA-specific immune responses in mCRC patients, and obtain data on overall survival. The trial was performed under FDA-approved IND14325 and registered at ClinicalTrials.gov (NCT01147965). The Ad5 [E1-, E2b-]-CEA(6D) doses were administered subcutaneous (sc) every 3 weeks for 3 treatments as follows: Cohort 1 (3 patients) received 10^9 VP each treatment; Cohort 2 (3 patients) received 10^{10} VP; Cohorts 3/4 (21 patients) received 10^{11} VP; and Cohort 5 (5 patients) received 5×10^{11} VP.

Patient Demographics:

[00401] Thirty-two mCRC patients with CEA expressing cancers were enrolled into the study. The 32 mCRC patients enrolled had a median age 57.5 (range 38-77), had failed a median of three prior chemotherapeutic regimens (range: 2-5), had a median performance status of 90% (range 70%-100%), and had a median of three sites of metastatic disease (range 1-5). The majority of patients received all three immunizations. Patients who stopped immunizations early did so due to significant disease progression. The mCRC demographics compared favorably with previously published studies with chemotherapy-refractory mCRC.

Adverse Effects:

[00402] There were no dose-limiting toxicities and no serious adverse events that resulted in treatment discontinuation at any dose level. The most common toxicity was a self-limited, injection site reaction. The blood hematology, chemistry, and anti-nuclear antibody (ANA) values at week 0 (before the first treatment) were compared with those obtained at week 9. There were no biologically relevant changes in chemistry, hematology, or ANA values.

Clinical Outcomes:

[00403] Patients were washed out from anti-cancer treatments for 30 days prior to immunotherapy. Both Kras wild type and mutated patients were enrolled.

[00404] The mCRC patients were followed for long-term survival and Kaplan-Meier survival plots performed (PRISM software). Events were determined from the social security death index database and clinical charts. Median overall survival was 11 months and overall survival was 30% at 18 months and 20% at 29 months (**FIG. 18**).

Immune Responses:

[00405] A secondary objective was to evaluate CEA-specific immune responses during immunotherapy. Again, since the mCRC patients represented the highest number of patients treated, immunogenicity studies were performed on them. As determined by ELISA there was no detectable antibody activity directed against CEA in serum samples.

[00406] CMI responses of mCRC patients treated in all cohorts were assessed. Peripheral blood mononuclear cells (PBMC) were isolated from patients before and after each immunotherapy treatment, as well as three weeks after the last treatment. CEA-specific ELISpot assays were performed on PBMC as previously described to determine the number of IFN- γ -secreting cells (SFC) after exposure to CEA peptides. The highest CMI responses during immunizations, regardless of time point (weeks 3, 6, or 9) were determined. This analysis revealed a dose-response relationship with increasing levels of vaccine (**FIG. 19**); the highest CMI levels occurred in patients who received the highest dose of 5×10^{11} VP (Cohort 5) and the responses were observed despite the presence of pre-existing Ad5 immunity in a

majority (61%) of patients. Analysis of serial PBMC samples over time indicated that CEA directed CMI responses were induced and increased over the course of 3 immunizations (**FIG. 20**).

[00407] Briefly, CMI (IFN- γ secretion) was assessed at baseline (Pre) and after administrations of Ad5 [E1-, E2b-]-CEA(6D) (Post). The highest CMI responses (regardless of time point) observed in the patients after treatment revealed a dose response. The highest CMI levels occurred in patients receiving the highest dose (Cohort 5) and was significantly elevated ($P < 0.02$; Mann-Whitney test). Specificity of the responses was demonstrated by lack of reactivity with irrelevant antigens β -galactosidase and HIV-gag. For positive controls, PBMCs were exposed to concanavalin A. Values = Mean \pm SEM (**FIG. 19**).

[00408] Briefly, CMI (ELISpot IFN- γ SFC) responses in immunized mCRC patients were assessed at weeks 0, 3, 6, and 9. Note the increase in CMI responses during immunizations. Values=Mean \pm SEM (**FIG. 20**).

Additional immune analyses on treated mCRC patients using flow cytometry.

[00409] Flow cytometry testing on PBMC from a patient with high CMI activity (>1000 ELISpot IFN- γ SFC) after exposure to CEA peptides revealed CD8+/IFN- γ + cells (3.5%), CD8+/IFN- γ +/TNF- α + cells (1.0%), CD4+/IFN- γ + cells (0.4%), and CD4+/IFN- γ +/TNF- α + cells (0.2%) demonstrating that polyfunctional cells were present.

[00410] Patient PBMC samples were also tested for HLA-A2 positivity by flow cytometry and 63% of the samples tested were HLA-A2 positive. When the highest CMI response level achieved per patient was assessed in association with the presence of HLA-A2, there was no significant difference observed between HLA-A2+ and HLA-A2- patients (264.6 ± 119.0 IFN- γ SFC versus 165.6 ± 108.1 IFN- γ SFC).

[00411] To assess the induction of cytolytic T lymphocyte (CTL) responses, ELISpot assays for granzyme B activity were performed. Granzyme B is a key mediator of target cell death via the granule-mediated pathway and the release of granzyme B by cytolytic lymphocytes upon effector-target interaction is an indicator of CTLs. The granzyme B ELISpot assay is a superior alternative to the ^{51}Cr -release assay since it is significantly more sensitive and provides an estimation of cytotoxic effector cell frequency.

[00412] Increased granzyme B activity in PBMCs were observed after immunization (**FIG. 16**).

Importantly, in an extended follow-up on PBMC samples from 5 mCRC patients, CMI responses were observed to decrease after immunizations (**FIG. 22**) were stopped indicating that further “booster” immunizations may be required to maintain induced CEA directed CMI activity.

[00413] Briefly, CTL responses (ELISpot granzyme B secreting SFC) were assessed pre (week 0) and post (week 6-9) treatment. Responses increased after immunizations ($P < 0.05$ i). Values = Mean \pm SEM (**FIG. 21**).

[00414] Briefly, PBMC samples from 5 patients as assessed by ELISpot IFN- γ SFC. CMI responses peaked at week 9 and decreased by week 26 after treatment was stopped (**FIG. 22**).

[00415] These data indicate that multiple homologous immunizations with Ad5 [E1-, E2b-]-CEA(6D) induce CEA-specific CMI immune responses in patients despite pre-existing Ad5-neutralizing activity. These data further indicate that importantly, there was minimal toxicity and a favorable overall survival profile was observed. Finally, the results indicate that the novel Ad5 [E1-, E2b-]-CEA(6D) gene delivery/expression platform can overcome tolerance to TAA and generate significant CMI responses to CEA in the setting of naturally acquired Ad5-specific immunity.

EXAMPLE 6: GLP Production of Clinical Grade Multi-targeted Vaccine

[00416] This example shows the production of clinical-grade multi-target vaccine using good laboratory practice (GLP) standards. Previously, the Ad5 [E1-, E2b-]-CEA(6D) product was produced using a 5 L Cell Bioreactor under GLP conditions in accordance with good manufacturing practice standards. This example shows that the Ad5 [E1-, E2b-]-mMUC1-C and the Ad5 [E1-, E2b-]-Brachyury products can be produced in a 5 L Cell Bioreactor using a similar approaches.

[00417] Briefly, vials of the E.C7 manufacturing cell line will be thawed, transferred into a T225 flasks, and initially cultured at 37 °C in 5% CO₂ in DMEM containing 10% FBS/4 mM L-glutamine. After expansion, the E.C7 cells will be expanded using 10-layered CellSTACKS (CS-10) and transitioned to FreeStyle serum-free medium (SFM). The E.C7 cells will be cultured in SFM for 24 hours at 37 °C in 5% CO₂ to a target density of 5x10⁵ cells/mL in the Cell Bioreactor. The E.C7 cells will then be infected with Ad5 [E1-, E2b-]-mMUC1-C or Ad5 [E1-, E2b-]-Brachyury, respectively, and cultured for 48 hours.

[00418] Mid-stream processing will be performed in an identical manner as that used to prepare clinical grade Ad5 [E1-, E2b-]-CEA(6D) product under IND14325. 30 minutes before harvest, Benzonase nuclease will be added to the culture to promote better cell pelleting for concentration. After pelleting by centrifugation, the supernatant will be discarded and the pellets re-suspended in Lysis Buffer containing 1% Polysorbate-20 for 90 minutes at room temperature. The lysate will then be treated with Benzonase and the reaction quenched by addition of 5M NaCl. The slurry will be centrifuged and the pellet discarded. The lysate will be clarified by filtration and subjected to a two-column ion exchange procedure.

[00419] To purify the vaccine products, a two-column anion exchange procedure will be performed. A first column will be packed with Q Sepharose XL resin, sanitized, and equilibrated with loading buffer. The clarified lysate will be loaded onto the column and washed with loading buffer. The vaccine product will be eluted and the main elution peak (eluate) containing Ad5 [E1-, E2b-]-mMUC1-C or Ad5 [E1-, E2b-]-Brachyury carried forward to the next step. A second column will be packed with Source 15Q resin, sanitized, and equilibrated with loading buffer. The eluate from the first anion exchange column will be loaded onto the second column and the vaccine product eluted with a gradient starting at 100% Buffer A (20 mM Tris, 1 mM MgCl₂, pH 8.0) running to 50% Buffer B (20 mM Tris, 1 mM MgCl₂, 2M NaCl, pH 8.0). The elution peak containing Ad5 [E1-, E2b-]-mMUC1-C or Ad5 [E1-, E2b-]-Brachyury

will be collected and stored overnight at 2-8 °C. The peak elution fraction will be processed through a tangential flow filtration (TFF) system for concentration and diafiltration against formulation buffer (20 mM Tris, 25 mM NaCl, 2.5% (v/v) glycerol, pH 8.0). After processing, the final vaccine product will be sterile filtered, dispensed into aliquots, and stored at ≤ -60 °C. A highly purified product approaching 100% purity is typically produced and similar results for these products are predicted.

[00420] The concentration and total number of VP product produced will be determined spectrophotometrically. Product purity is assessed by HPLC. Infectious activity is determined by performing an Ad5 hexon-staining assay for infectious particles using kits.

[00421] Western blots will be performed using lysates from vector transfected A549 cells to verify mMUC1-C or Brachyury expression. Quality control tests will be performed to determine that the final vaccine products are mycoplasma-free, have no microbial bioburden, and exhibit endotoxin levels less than 2.5 endotoxin units (EU) per mL. To confirm immunogenicity, the individual vectors will be tested in mice as described below (Example 7).

EXAMPLE 7: Immunogenicity of multi-targeted CEA, MUC1, T viral vector

[00422] This example shows immunogenicity results using the multi-targeted vaccine directed to CEA, MUC1 and T (Brachyury). Each viral vector product was tested for purity, infectivity, and antigen expression, as described herein and each passed these criteria.

Vaccination and splenocyte preparation

[00423] Female C57BL/6 mice (n=5) were injected s.c. with 10^{10} VP of Ad5 [E1-, E2b-]-brachyury or Ad5 [E1-, E2b-]-CEA or Ad5 [E1-, E2b-]-MUC1 or a combination of 10^{10} VP of all three viruses at a ratio of 1:1:1 (Tri-Ad5). Control mice were injected with 3×10^{10} VP of Ad-null (no transgene insert). Doses were administered in 25 μ l of injection buffer (20mM HEPES with 3% sucrose) and mice were vaccinated three times at 14-day intervals. Fourteen days after the final injection spleens and sera were collected. Sera were frozen at -20 °C. Splenocyte suspensions were generated by gently crushing the spleens through a 70 μ M nylon cell strainer (BD Falcon, San Jose, CA). Red cells were removed by the addition of red cell lysis buffer (Sigma-Aldrich, St. Louis, MO) and the splenocytes were washed twice and resuspended in R10 (RPMI 1640 supplemented with L- glutamine (2 mM), HEPES (20 mM), penicillin 100 U/ml and streptomycin 100 μ g/ml, and 10% fetal bovine serum. Splenocytes were assayed for cytokine production by ELISPOT and flow cytometry.

Immunogenicity studies:

[00424] Previous studies assess induced immunity generated by the multi-targeted vaccine mixture. Immunization with Ad5 [E1-, E2b-] vectors is dose dependent and 1×10^{10} VP per dose was routinely used. Groups (N=5) of C57Bl/6 mice were used.

[00425] This study C57Bl/6 mice were injected subcutaneously 3 times at 2-week intervals with tri-immunization comprising 3×10^{10} viral particles (VP) Ad5 [E1-, E2b-]-null (empty vector controls) or

with 3×10^{10} VP containing a 1:1:1 mixture of Ad5 [E1-, E2b-]-CEA(6D), Ad5 [E1-, E2b-]-mMUC1-C, and Ad5 [E1-, E2b-]-Brachyury.

[00426] Two weeks after the last immunization CMI activity was determined employing ELISpot assays for IFN- γ secreting cells (SFC) after exposure of splenocytes to CEA, MUC1, or Brachyury peptide pools, respectively, as described (8-12,14).

[00427] Significant CMI responses to were detected in immunized mice (**FIG. 23**). Flow cytometry utilizing intracellular cytokine staining was performed on spleen cells after exposure to CEA peptides to assess the quantity of activated CD4+ and CD8+ T-cells.

[00428] Briefly, CMI responses against CEA, MUC1, and Brachyury as assessed by ELISpot assays for IFN- γ secreting splenocytes (SFC) were detected in multi-targeted immunized mice but not control mice (injected with Ad5-Null empty vector). Specificity of the ELISpot assay responses was confirmed by lack of reactivity to irrelevant SIV-nef or SIV-vif peptide antigens. A positive control included cells exposed to concanavalin A (Con A, data not shown). Values = Mean \pm SEM (**FIG. 23**).

ELISpot assay

[00429] Brachyury-, CEA- and MUC1-specific IFN- γ - or IL-2-secreting T cells were determined by ELISpot assay from freshly isolated mouse splenocytes, as described above. Briefly, 2×10^5 splenocytes were stimulated with 0.2 μ g/well of overlapping 15-mer peptides in a single pool derived from brachyury or CEA, or MUC1. Cells were stimulated with Con A at a concentration of 0.0625 μ g/per well as a positive control and overlapping 15-mer complete peptides pools derived from SIV-Nef and SIV-Vif were used as irrelevant peptide controls. The numbers of SFCs were determined using an Immunospot ELISpot plate reader and results were reported as the number of SFCs per 10^6 splenocytes.

[00430] TNF- α and IFN- γ expressing polyfunctional CD4+ and CD8+ cells were detected in immunized but not control splenocytes (**FIG. 24**). Testing was also performed on sera to detect induced CEA antibody using a previously described quantitative ELISA assay with purified CEA protein. Polyfunctional CD8+ (top) and CD4+ (bottom) cells expressing IFN- γ and TNF- α in mice immunized with the multi-targeted vaccine (right) but not in controls injected with Ad5-null (left). Specificity of the responses was confirmed by lack of reactivity to media alone or irrelevant SIV-nef or SIV-vif peptides. Values = Mean \pm SEM (**FIG. 24**).

[00431] Significant antibody responses to CEA were detected in immunized but not control mice (**FIG. 8**). To determine the level of complement dependent cellular cytotoxicity (CDCC), a CDCC test was performed using murine MC38-CEA target cells. Significant CDCC activity was detected in sera of immune, but not control mice (injected with Ad5-null) (**FIG. 25**). Significant ($P < 0.0001$) CEA antibody (left) and CDCC (right) responses were induced in mice immunized with the multi-targeted vaccine but not in control mice. Values = Mean \pm SEM (**FIG. 25**).

Intracellular cytokine stimulation

[00432] Splenocytes were prepared as indicated for above. Stimulation assays were performed using 1×10^6 live splenocytes per well in 96-well U-bottom plates. Pools of overlapping peptides spanning the entire coding sequences of brachyury, CEA and MUC1 were synthesized as 15-mers with 11-amino acid overlaps and lyophilized peptide pools were dissolved in DMSO. Similarly constructed peptide pools corresponding to SIV-Vif and SIV-Nef served as off-target controls. Splenocytes in R10 media (RPMI 1640, 10% fetal bovine serum, and antibiotics) were stimulated by the addition of peptide pools at 2 $\mu\text{g/mL}/\text{peptide}$ for 6h at 37 °C and 5% CO₂, with protein transport inhibitor (GolgiStop, BD) added 2h into the incubation. Stimulated splenocytes were then stained for lymphocyte surface markers CD8 α and CD4, fixed, permeabilized, and then stained for the intracellular accumulation of IFN- γ and TNF α . Antibodies against mouse CD8 α , CD4, IFN- γ , and TNF α were used and staining was performed in the presence of anti-CD16/CD32. Flow cytometry was performed and analyzed in BD Accuri C6 Software.

Complement-dependent cytotoxicity assay (CDC)

[00433] MC38-CEA2 tumor cells were cultured overnight at a density of 2×10^4 cells per well in 96-well tissue culture microplates. Pooled heat inactivated mouse sera were added at a 1:50 dilution and incubated at 37°C for 1 hour. Rabbit serum was then added at a 1:50 dilution as a source of complement and cells were incubated an additional 2.5 hours at 37 °C. Cell culture supernatants were assayed using Promega Cytotox 96 non-radioactive cytotoxicity assay, according to the manufacturer's instructions. Percent lysis of MC38-CEA2 cells was calculated by the formula % lysis = (experimental – target spontaneous) / (target maximum – target spontaneous) x 100%.

Anti-tumor immunotherapy studies:

[00434] Studies were conducted to test the anti-tumor capability of Ad5 [E1-, E2b-]-based tri-vaccines in immunotherapy studies in mice with established CEA, MUC1, or Brachyury expressing tumors, respectively. In this study the anti-tumor activity of the individual components of the Ad5 [E1-, E2b-]-based tri-vaccine was assessed.

[00435] Groups (n=7) of C57Bl/6 mice were injected subcutaneously in the right flank with 5×10^5 CEA, MUC1, and/or Brachyury expressing murine tumor cells. After palpable tumors were detected, mice were treated by 3 subcutaneous injections with 1×10^{10} VP each of Ad5 [E1-, E2b-]-null (no transgene, e.g., empty vector), Ad5 [E1-, E2b-]-CEA(6D), Ad5 [E1-, E2b-]-mMUC1-C, and/or Ad5 [E1-, E2b-]-Brachyury, respectively. Tumor volumes were calculated and tumor growth curves were plotted. 7-10 mice/group are sufficient for statistical evaluation of treatment.

[00436] For *in vivo* tumor treatment studies, female C57BL/6 mice, 8–10 weeks old, were implanted with 10^6 MC38-MUC1 cells s.c. in the left flank. Mice were treated three times at a 7-day interval with 10^{10} VP Adeno-MUC1 or Tri-Ad5. Control mice were injected with 3×10^{10} VP of Adeno-null. Tumor growth was assessed by measuring two opposing dimensions (a, b) and the volume calculated according

to the formula $V=(a \times b)^2/2$ where the shorter dimension was "a". Tumor studies were terminated when tumors reached 1500 mm^3 or became severely ulcerated.

[00437] Significant anti-tumor activity and increased survival in MUC1 expressing tumor-bearing mice treated by immunotherapy (**FIG. 26**). Importantly, flow cytometry was used to determine that the MC38-MUC1 cell line expressed PDL1 and anti-tumor activity was achieved despite its presence. C57Bl/6 mice (n=7/group) were inoculated with MC38-MUC1 expressing tumor cells subcutaneously in the left flank and administered 10^{10} VP of Ad5-Null (empty vector) or 10^{10} VP of Ad5 [E1-, E2b-]-mMUC1-C in the right flank on days 0, 7, 14.

[00438] Mice treated with Ad5 [E1-, E2b-]-mMUC1-C had significantly ($P < 0.05$) smaller tumors on days 15 and 18 as compared to controls (top) and significantly longer survival (bottom). Values = Mean \pm SEM. Experiment was terminated on day 36 (**FIG. 26**).

[00439] Immunotherapy of mice with Ad5 [E1-, E2b-]-Brachyury resulted in smaller tumors (**FIG. 27**). Briefly, C57Bl/6 mice were inoculated with MC38-Brachyury expressing tumor cells subcutaneously in the left flank and administered 10^{10} VP of Ad5-Null (empty vector) (N=4) or 10^{10} VP of Ad5 [E1-, E2b-]-Brachyury (N=5) in the right flank on days 5, 11, and 17. Tumors were smaller in treated mice on days 15, 19, and 22. Values = Mean \pm SEM (**FIG. 27**)

[00440] Larger numbers of mice will be treated to show significant anti-tumor activity and to combine immunotherapy with immune pathway checkpoint modulators, such as anti-checkpoint inhibitor antibodies, to determine if anti-tumor activity is enhanced.

EXAMPLE 8: In vitro validation of human T-cell activation by recombinant viral vectors

Tumor cell culture

[00441] Human colon carcinoma SW620 (HLA-A2 $^+$, HLA-A24 $^+$, brachyury $^+$, MUC1 $^+$, CEA $^+$) and pancreatic carcinoma ASPC-1 (HLA-A1 $^+$, HLA-A26 $^+$, MUC1 $^+$) cell lines were used. Cell cultures were free of mycoplasma and maintained in complete medium (RPMI-1640 supplemented with 10% FBS, 100 U/ml penicillin, 100 μ g/ml streptomycin and 2 mM L-glutamine).

Detection of cytokines

[00442] Supernatants of T cells stimulated for 24 h with DCs infected with adenovirus vectors or peptide-pulsed DCs in IL-2-free medium were evaluated for secretion of IFN- γ using an ELISA kit. The antigen-specific T-cell lines used in this analysis have been reported previously: (a) an HLA-A2 CEA-specific CTL, (b) an HLA-A2 MUC1-specific CTL, (c) an HLA-A24 MUC1-specific CTL, and (d) an HLA-A2 brachyury-specific CTL.

Cytotoxic assay

[00443] In brief, target cells were labeled with 50 μ Ci of ^{111}In oxide at 37 °C for 20 min and used at 3,000 cells/well in 96-well round-bottom culture plates. T cells were added at different ratios and incubated at 37 °C for 16 h. Supernatants were harvested for gamma counting. Determinations were

carried out in triplicate and SDs were calculated. Spontaneous release was determined by incubating target cells with medium alone and complete lysis was determined by incubating with 0.25% Triton X-100. Specific lysis was calculated with the use of the following formula: Lysis (%) = [observed release (CPM) - spontaneous release (CPM)] / [Complete release (CPM)- spontaneous release (CPM)] x 100.

[00444] An *in vitro* study was performed to assess dendritic cells (DC) function and antigen-specific T-cell activation using the Ad5 [E1-, E2b-]-CEA(6D), and Ad5 [E1-, E2b-]-Brachyury vectors. Briefly, human DC were cultured 6-7 days in media with IL-2 and GM-CSF. DC cultured in AIM-V medium were infected with Ad5 [E1-, E2b-]-CEA(6D), or Ad5 [E1-, E2b-]-Brachyury, or Ad5 [E1-, E2b-]-null (empty vector) and incubated for 2 days. Uninfected DCs were used as controls.

[00445] Human DC cells were then analyzed by flow cytometry for expression of CD80 (a marker for DC maturation), CD83 (a marker for DC maturation), CD86 (a DC marker) and DR (a DC marker) as measured by mean fluorescence intensity. As shown in **Table 6**, infection of DC with Ad5 [E1-, E2b-]-based vectors induced activation and maturation of human DCs.

Table 6

Treatment	MOI	% CD80+ cells	% CD83+ cells	% CD86+ cells	% DR+ cells
Non-infected controls	0	20.2%	33.6%	99.5%	94.4%
Ad5 [E1-, E2b-]-null infected	10,000	40.7%	40.9%	99.2%	98.0%
Ad5 [E1-, E2b-]-CEA(6D) infected	10,000	56.8%	48.7%	99.2%	98.3%

[00446] Dendritic cells (2×10^5) in 1 mL of AIM-V medium were infected with adenovirus vectors (Ad5 [E1-, E2b-]-CEA, Ad5 [E1-, E2b-]-MUC1, Ad5 [E1-, E2b-]-brachyury, and Ad5 [E1-, E2b-]-null at indicated multiplicity of infection (MOI of 10,000 or 20,000) for 1 hour in 6-well plates. AIM-V medium (4 mL) was then added to each well and incubated for an additional 2 days. To analyze the efficacy of transgene expression, DCs were harvested and analyzed using flow cytometry and Western blot. For phenotypic analysis, DCs were stained for the expression of CD80, CD83, CD86, CEA, and HLA-DR using BV421-conjugated anti-CD80, PerCP Cy5.5-conjugated anti-CD83, APC-Cy7-conjugated anti-HLA-DR, PE-conjugated anti-CD86, and FITC-conjugated anti-CEA. As shown in **Table 7**, Ad5 vectors induced maturation and activation of human DCs. Indicated adenovirus vectors were used at a concentration of 10,000 or 20,000 multiplicity of infection (MOI) for infection of human dendritic cells (DCs). Expression of CD80, CD83, CD86 and HLA-DR were analyzed by flow cytometry. Results are expressed in % of positive cells (mean fluorescence intensity).

Table 7

Treatment	MOI	CD80	CD83	CD86	HLA-DR
Control	0	20.2 (146)	33.6 (259)	99.5 (4586)	94.4 (1355)
Ad5 [E1-, E2b-]-null	10,000	40.7 (162)	40.9 (253)	99.2 (3794)	98.0 (4489)
Ad5 [E1-, E2b-]-null	20,000	47.4 (169)	46.8 (266)	98.6 (3012)	95.7 (3203)
Ad5 [E1-, E2b-]-CEA	10,000	56.8 (189)	48.7 (262)	99.2 (3877)	98.3 (6553)
Ad5 [E1-, E2b-]-CEA	20,000	54.5 (185)	46.6 (271)	99.1 (3628)	98.0 (6015)
Ad5 [E1-, E2b-]-MUC1	10,000	41.4 (167)	42.4 (251)	98.5 (3178)	96.9 (5227)
Ad5 [E1-, E2b-]-MUC1	20,000	46.7 (172)	44.3 (260)	98.9 (3591)	97.2 (5779)

[00447] To determine if infected human DCs could stimulate human antigen-specific T-cell lines to secrete IFN- γ , infected DCs were incubated with antigen-specific T-cell lines and tested for IFN- γ secreting activity. As shown in **Table 8**, only the CEA or Brachyury antigen-specific HLA-A2+ T-cell lines incubated with respective HLA-A2+ DC infected with Ad5 [E1-, E2b-]-CEA(6D) or Ad5 [E1-, E2b-]-Brachyury were stimulated to secrete IFN- γ . No IFN- γ secretion was detected in the controls. Infection of HLA-A2+ DC with Ad5 [E1-, E2b-]-based vectors can activate antigen-specific T-cells to produce IFN- γ .

Table 8

Treatment	MOI	Antigen-specific T-cell line	
		CEA (HLA-A2+) (pg IFN- γ /10 ⁵ cells/mL)	Brachyury (HLA-A2+) (pg IFN- γ /10 ⁵ cells/mL)
Non-infected controls	0	<15.6	<15.6
No DC	0	<15.6	<15.6
Ad5 [E1-, E2b-]-null infected	10,000	<15.6	<15.6
Ad5 [E1-, E2b-]-CEA(6D) infected	10,000	122.7	Not Done
Ad5 [E1-, E2b-]-Brachyury infected	10,000	Not Done	145.6

[00448] HLA-A2+ DC were infected with a mixture of Ad5 [E1-, E2b-]-CEA(6D) and Ad5 [E1-, E2b-]-Brachyury. Infected DCs were used to generate specific cytotoxic HLA-A2+ T-lymphocytes (CTL) using autologous PBMC. The autologous DCs were used as APC for three in vitro stimulations (IVSs). Autologous B cells pulsed with CEA or Brachyury were used to re-stimulate antigen-specific CTLs after the 3 IVSs. The effector T-cell to target tumor cell ratio was 30:1 and the percent cytolytic activity was determined. As shown in **Table 9**, infection of HLA-A2+ DC with a mixture of Ad5 [E1-, E2b-]-CEA(6D) and Ad5 [E1-, E2b-]-Brachyury can generate antigen-specific cytolytic T-cells. Cytolytic activity was detected in an HLA-A2 dependent manner.

Table 9

Antigen-specific T-cell line	% lysis of SW620 cells (HLA-A2+, CEA, and Brachyury expressing)	% lysis of ASPC-1 cells (HLA-A1+ and CEA expressing)
CEA-specific HLA-A2+ T-cells	42.4%	4.3%
Brachyury-specific HLA-A2+ T-cells	64.4%	8.3%

[00449] To determine if infected human DCs could stimulate human antigen-specific T-cell lines to lyse target tumor cells. Human DCs (6-day culture in IL-4 and granulocyte-macrophage colony-stimulating factor (GM-CSF) 2×10^4 cells/well in 0.5 mL of AIM-V) were infected with indicated adenovirus vectors at 20,000 multiplicity of infection (MOI). After 48 hours, DCs were washed and used for stimulation of human antigen-specific T cells. Results are expressed in pg/ml of IFN- γ per 1×10^5 T cells/ml. **Table 10** demonstrates that infection of human DCs with recombinant adenovirus vectors encoding CEA, MUC1, or brachyury, can activate antigen specific T-cell lines. Numbers in bold indicate a significant enhancement of IFN- γ secretion compared to corresponding wells with uninfected DCs. [-- indicates that the assay was not performed.]

Table 10

DCs infected with	Antigen-specific T-cell lines			
	CEA	MUC1 (HLA-A2)	MUC1 (HLA-A24)	Brachyury
Ad5 [E1-, E2b-]-null	<15.6	<15.6	<15.6	<15.6
Ad5 [E1-, E2b-]-brachyury	<15.6	--	--	351.9
Ad5 [E1-, E2b-]-MUC1	<15.6	335.2	806.4	--
Ad5 [E1-, E2b-]-CEA	350.0	<15.6	<15.6	--
Uninfected DCs	<15.6	<15.6	<15.6	<15.6
T cells only	<15.6	<15.6	<15.6	<15.6

[00450] Human DCs (6-day culture in IL-4 and GM-CSF) from an HLA-A2 and -A24 donor were infected with Tri-Ad5 vector at 2×10^4 /well (24-well plate) in 0.5 mL of AIM-V. Tri-Ad5 vectors were used at 20,000 MOI for 1 hour and then 1.5 mL of AIM-V were added to each well. Infected DCs were incubated for 48 hours and then washed and used for stimulation of human antigen-specific T cells. Results are expressed in pg of IFN- γ per 1×10^5 T cells/ml. Numbers in bold indicate a significant enhancement of IFN- γ secretion compared to corresponding wells with uninfected DCs.

[00451] CEA-, MUC1- and brachyury-specific cytotoxic T lymphocytes (CTLs) were generated. Dendritic cells ($1-2 \times 10^5$ /well in 1 mL of AIM-V) were infected with 20,000 MOI of Tri-Ad5, as described above. Infected DCs were used as APCs for stimulation of autologous nonadherent cells at an

effector-APC ratio of 10:1. Cultures were incubated for 3 days at 37 °C in a humidified atmosphere containing 5% CO₂. The cultures were then supplemented with rhIL-2 for 7 days; IL-2 containing medium was replenished every 3 days. The 10-day stimulation constituted one in vitro stimulation (IVS) cycle. Autologous vector-infected DCs were used as APCs for three IVS. Autologous peptide-pulsed B cells were used to re-stimulate antigen-specific CTLs after three IVS. T-cell lines were maintained in medium containing IL-7 and IL-15 (10 ng/ml).

[00452] Human DCs from a prostate cancer patient (6-day culture in IL-4 and GM-CSF; 2x10⁴ cells/well in 0.5 mL of AIM-V) were infected with Tri-Ad5 at 20,000 MOI. After 48 h, infected DCs were washed and used to generate specific cytotoxic T lymphocytes (CTLs) using autologous peripheral blood mononuclear cells (PBMCs) as effectors. Following 3 cycles of in vitro stimulations, autologous peptides-pulsed B cells were used as antigen-presenting cells. Results are expressed in pg/ml of IFN-γ. [-- indicates that the assay was not performed.] As shown in **Table 11**, infection of human dendritic cells can generate antigen-specific T cells to brachyury, MUC1 and CEA and produce IFN-γ when stimulated with autologous B cells pulsed with the peptides.

Table 11

Antigen-specific T-cell lines	Peptides (10 µg/ml)			
	CEA	MUC1(A2)	MUC1(A24)	Brachyury
T-Brachyury	<15.6	--	--	243
T-MUC1 (A2)	<15.6	174	--	--
T-MUC1 (A24)	<15.6	--	206	--
T-CEA	211	<15.6	--	--

[00453] CEA-, MUC1- and brachyury-specific cytotoxic T lymphocytes (CTLs) were generated. Dendritic cells (1-2x10⁵/well in 1 mL of AIM-V) were infected with 20,000 MOI of Tri-Ad5, as described above. Infected DCs were used as APCs for stimulation of autologous nonadherent cells at an effector-APC ratio of 10:1. Cultures were incubated for 3 days at 37 °C in a humidified atmosphere containing 5% CO₂. The cultures were then supplemented with rhIL-2 for 7 days; IL-2 containing medium was replenished every 3 days. The 10-day stimulation constituted one in vitro stimulation (IVS) cycle. Autologous vector-infected DCs were used as APCs for three IVS. Autologous peptide-pulsed B cells were used to re-stimulate antigen-specific CTLs after three IVS. T-cell lines were maintained in medium containing IL-7 and IL-15 (10 ng/ml).

[00454] Human DCs (6-day culture in IL-4 and GM-CSF) from an HLA-A2 and -A24 donor were infected with Tri-Ad5 vector at 2 x 10⁴/well (24-well plate) in 0.5 mL of AIM-V. Tri-Ad5 vectors were used at 20,000 MOI for 1 hour and then 1.5 mL of AIM-V were added to each well. Infected DCs were

incubated for 48 hours and then washed and used for stimulation of human antigen-specific T cells. Results are expressed in pg of IFN- γ per 1×10^5 T cells/ml. Infection of human dendritic cells with Tri-Ad5 vectors encoding transgenes can activate antigen-specific T cell lines to produce IFN- γ . As shown in **Table 12**, infection of human dendritic cells with Tri-Ad5 can generate antigen-specific T cells to brachyury, MUC1 and CEA and produce IFN- γ when stimulated with autologous B cells pulsed with the corresponding peptides. Numbers in bold indicate a significant enhancement of IFN- γ secretion compared to corresponding wells with uninfected DCs.

Table 12

DCs infected with	Antigen-specific T-cell lines			
	CEA (HLA-A2)	MUC1 (HLA-A2)	MUC1 (HLA-A24)	Brachyury (HLA-A2)
Tri-Ad5	480	236	763	496
Ad5 [E1, E2b]-null	<15.6	<15.6	<15.6	<15.6
Uninfected DCs	<15.6	<15.6	<15.6	<15.6
T cells only	<15.6	<15.6	<15.6	<15.6

[00455] These studies show that Ad5 [E1-, E2b-]-based vectors can induce maturation of human DCs in vitro, infected DCs can stimulate antigen-specific human T-cells, and infected DCs can generate antigen-specific human CTLs. In the studies reported here, multi-TAA targeted immunotherapy (Tri-Ad5), which consisted of a mixture of three Ad5 vectors expressing different TAAs, is as efficient in the activation of human T cells as the use of each of the adenovirus vectors alone, with only minor differences. Nine different *in vivo* parameters were measured via vaccinating mice with each of the Ad5 [E1-, E2b-]-CEA, Ad5 [E1-, E2b-]-MUC1, and Ad5 [E1-, E2b-]-brachyury vectors individually versus vaccination with Tri-Ad5. Of the 21 assays performed, statistical differences observed were (a) an enhanced number of MUC1-specific splenocytes and CD8 $^+$ IFN-producing and multifunctional CD8 $^+$ T cells, and (b) more CEA-specific CD8 $^+$ IFN-producing T cells, in the mice vaccinated with one vector than in the Tri-Ad5 vaccinated mice. On the other hand, the Tri-Ad5 vaccinated mice produced more CEA-specific IL-2-producing cells than the Ad5 [E1-, E2b-]-CEA vaccinated mice. In the other 16/21 assays, however, there were no statistical differences in the results between the use of the individual vector versus the use of the Tri-Ad5 in terms of antigen-specific activation of (a) splenocytes for IFN- γ and IL-2 production, (b) CD8 $^+$ T cells for IFN- γ production, (c) CD4 $^+$ T cells for IFN- γ production, (d) multifunctional CD8 $^+$ T cells for IFN- γ and TNF- α production, (e) multifunctional CD4 $^+$ T cells for IFN- γ and TNF- α production, and (f) production of antigen-specific antibodies. There was also no difference in anti-tumor activity using a single vector (Ad5 [E1-, E2b-]-MUC1) versus the Tri-Ad5 vaccine; while both vaccines did not eliminate the tumor, both vaccines reduced the tumor growth rate in a similar manner. While Tri-Ad5

was not as efficient in T-cell activation in some assays, the potential ability of the Tri-Ad5 platform to overcome the TAA heterogeneity that exists in human solid tumors far outweighs the relatively minor differences in potency of T-cell activation of Tri-Ad5 vs. individual vectors in some assays.

[00456] CEA, MUC1 and brachyury are all human TAAs and are not expressed in murine solid tumors. Moreover, human solid tumors are very heterogeneous with respect to expression of different TAAs. It would be extremely difficult to transfect a murine tumor cell line with all three transgenes to define the effect of vaccination of Tri-Ad5 vs. each vector alone. The targeting of a murine tumor expressing MUC1 was chosen because the single Ad5 [E1-, E2b-]-MUC1 vector was more potent in some of the murine T-cell assays compared to the Tri-Ad5 than the CEA and brachyury vectors compared to Tri-Ad5. Thus this appeared to be the most stringent model to compare the Tri-Ad5 platform to a single vector platform. The studies reported herein were designed to provide the rationale for potential clinical studies as a vaccine immunotherapy, or use in combination with other therapeutics, using this novel adenovirus vaccine delivery platform (Ad5 [E1-, E2b-] targeting a diverse range of TAA transgenes in the Tri-Ad5 regimen.

[00457] While the checkpoint inhibitor antibodies have shown evidence of clinical activity in melanoma and squamous non-small cell lung cancer, clinical benefit in other cancer types has been observed in a minority of patients. For some tumor types, such as colorectal cancer and prostate cancer, the anti-PDL1/PD1 checkpoint inhibitors have shown little clinical activity. One hypothesis that has been put forth for the lack of PDL1/PD1 therapeutic activity in some patients is the lack of T-cell infiltrates in tumors. Consequently, if a vaccine targeting TAAs in the tumor would result in the presence of antigen-specific T cells in the tumor microenvironment, then an immune pathway checkpoint modulator employed in combination or following vaccination would be able to “release the brakes” of the tumor-infiltrating anergized T cells leading to clinical effect.

EXAMPLE 9: Immunotherapy for HPV

[00458] This example shows that the combination therapy as provided herein reduces tumors and a murine model of HPV-associated cancers which express HPV early 6 (E6) and early 7 (E7) oncogenes. Briefly, tumor bearing HPV murine model mice were treated with the Ad5 [E1-, E2b-]-HPV-E6/E7 vaccine comprising a modified non-oncogenic and fused HPV-E6/E7 gene and treated tumor bearing mice. The TC-1 murine tumor cell line used to induce tumors in the mice expressed HPV-E6/E7 as well as PDL1.

[00459] Flow cytometry was used to examine the tumors in HPV tumor bearing mice and calculated the abundance of various tumor infiltrating lymphocytes (TILs). The TILs in the tumors were observed to have the following cell types and abundance levels: myeloid derived suppressor cells (MDSC) ($7.5\% \pm 2.4$ SD), FoxP3 expressing T regulatory (Treg) lymphocytes ($7.1\% \pm 2.5$), PD1 expressing CD8 α + lymphocytes ($25.5\% \pm 16.4$), and LAG-3 expressing CD8 α + lymphocytes ($4.1\% \pm 2.3$). This profile of TILs indicates that a suppressive immune pathway is present in the tumors.

[00460] The addition of immune checkpoint inhibitor antibodies, such as anti-PD1 antibody, to the Ad5 [E1-, E2b-]-HPV-E6/E7 immunotherapy resulted in greater anti-tumor responses (**FIG. 28**). Additionally two mice in this group were observed to have essentially complete tumor regression (**FIG. 29**). Briefly, C57Bl/6 mice (n=7/group) were implanted with HPV-E6/E7 expressing TC-1 tumor cells (day 0) and treated by immunotherapy on days 10, 17, 24 with 10^{10} VP of Ad5-null (empty vector) plus 100 μ g control IgG (intraperitoneal), 10^{10} VP of Ad5-Null (empty vector) plus 100 μ g anti-PD1, 10^{10} VP of Ad5 [E1-, E2b-]-HPV-E6/E7 plus 100 μ g mouse IgG, or 10^{10} VP of Ad5 [E1-, E2b-]-HPV-E6/E7 plus 100 μ g anti-PD1. Immunotherapy with or without anti-PD1 resulted in significant inhibition of tumor growth by day 23 (P<0.05). All control mice were terminated by day 23 due to tumor mass. Values = Mean \pm SEM (**FIG. 28**).

[00461] Tumor growth and regression in 2 mice treated by immunotherapy with anti-PD1 injections were analyzed (**FIG. 29**). Tumor growth peaked on days 20 & 27, respectively, and then regressed thereafter. This study demonstrates that the combination of immune checkpoint inhibitor drugs in combination with use of multi-targeted vaccines can enhance anti-tumor effects.

[00462] Immunotherapy against human papilloma virus (HPV) using a viral gene delivery platform to immunize against HPV 16 genes E6 and E7 (Ad5 [E1-, E2b-]-E6/E7) combined with programmed death-ligand 1 (PD1) blockade was investigated to determine whether it could increase therapeutic effect as compared to the vaccine alone. Ad5 [E1-, E2b-]-E6/E7 as a single agent induced HPV-E6/E7 cell-mediated immunity. Immunotherapy using Ad5 [E1-, E2b-]-E6/E7 resulted in clearance of small tumors and an overall survival benefit in mice with larger established tumors. When immunotherapy was combined with immune checkpoint blockade, an increased level of anti-tumor activity against large tumors was observed. Analysis of the tumor microenvironment in Ad5 [E1-, E2b-]-E6/E7 treated mice revealed elevated CD8⁺ tumor infiltrating lymphocytes (TILs); however, induction of suppressive mechanisms such as programmed death-ligand 1 (PDL1) expression on tumor cells and an increase in PD1⁺ TILs was seen. When Ad5 [E1-, E2b-]-E6/E7 immunotherapy was combined with anti-PD1 antibody, CD8⁺ TILs were observed at the same level but a reduction in tumor PDL1 expression on tumor cells and reduced PD1⁺ TILs providing a mechanism by which combination therapy favors a tumor clearance state and a rationale for pairing antigen-specific vaccines with immune pathway checkpoint modulators in future clinical trials.

[00463] Herein, Ad5 [E1-, E2b-]-E6/E7 immunizations combined with PD1 blockade were examined for an increase an anti-tumor effect. Also, the CMI response induced by the Ad5 [E1-, E2b-]-E6/E7 vaccine was characterized and the kinetics of an anti-tumor response was determined to evaluate the therapeutic potential of treating small versus large established tumors. To investigate a possible mechanism of action, the relationship between the levels of effector T cells and suppressor T cells within

the parenchyma of the tumor was evaluated and lymphocyte populations and expression of co-inhibitory molecules that may play a role in the observed anti-tumor responses were characterized.

Viral Construction

[00464] Ad5 [E1-, E2b-]-E6/E7 was constructed and produced. Briefly, the transgenes were sub-cloned into the Ad5 [E1-, E2b-] vector using a homologous recombination-based approach and the replication deficient virus was propagated in the E.C7 packaging cell line, CsCl₂ purified, and titered. Viral infectious titer was determined as plaque forming units (PFU) on an E.C7 cell monolayer. The viral particle (VP) concentration was determined by sodium dodecyl sulfate (SDS) disruption and spectrophotometry at 260 nm and 280 nm. As a vector control, Ad5 [E1-, E2b-]-null was employed, which is the Ad5 platform backbone with no transgene insert.

Immunization and splenocyte preparation

[00465] Female C57BL/6 mice (n=5/group) were injected subcutaneously (SQ) with varying doses of Ad5 [E1-, E2b-]-E6/E7 or Ad5 [E1-, E2b-]-null. Doses were administered in 25µL injection buffer (20mM HEPES with 3% sucrose) and mice were immunized three times at 14-day intervals. Fourteen days after the final injection, spleens and sera were collected. Serum from mice was frozen at -20°C until evaluation. Suspensions of splenocytes were generated by disrupting the spleen capsule and gently pressing the contents through a 70 µm nylon cell strainer. Red blood cells were lysed by the addition of red cell lysis buffer and after lysis, the splenocytes were washed twice in R10 (RPMI 1640 supplemented with L-glutamine (2mM), HEPES (20mM) (Corning, Corning, NY), penicillin (100 U/ml) and streptomycin (100 µg/mL), and 10% fetal bovine serum. Splenocytes were assayed for cytokine production by ELISpot and flow cytometry.

Enzyme-Linked Immunosorbent Spot (ELISpot) assay

[00466] HPV E6 and E7 specific interferon-γ (IFN-γ) secreting T cells were determined by ELISpot assays using freshly isolated mouse splenocytes prepared as described above. The ELISpot assay was performed. Pools of overlapping peptides spanning the entire coding sequences of HPV E6 and E7 were synthesized as 15-mers with 11-amino acid overlaps (and lyophilized peptide pools were dissolved in DMSO. Splenocytes (2x10⁵ cells) were stimulated with 2 µg/mL peptide of overlapping 15-mer peptides in pools derived from E6 or E7. Cells were stimulated with Concanavalin A (Con A) at a concentration of 0.06 µg/per well as a positive control. Overlapping 15-mer complete peptide pools derived from SIV-Nef (AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH) were used as irrelevant peptide controls. The numbers of Spot Forming Cells (SFC) were determined using an Immunospot ELISpot plate reader (and results reported as the number of SFC per 10⁶ splenocytes.

Intracellular Cytokine Stimulation

[00467] Splenocytes were prepared as described for the ELISpot assay above. Stimulation assays were performed using 10⁶ live splenocytes per well in 96-well U-bottom plates. Splenocytes in R10 media

were stimulated by the addition of HPV E6, HPV E7, or SIV-Nef peptide pools at 2 μ g/mL/peptide for 6 h at 37 °C in 5% CO₂, with protcin transport inhibitor (GolgiStop, BD) added two hours after initiation of incubation. Stimulated splenocytes were then stained for lymphocyte surface markers CD8 α and CD4, fixed with paraformaldehyde, permeabilized, and stained for intracellular accumulation of IFN- γ and TNF- α . Fluorescent-conjugated antibodies against mouse CD8 α (clone 53-6.7), CD4 (clone RM4-5), IFN- γ (clone XMG1.2), and TNF- α (clone MP6-XT22) were purchased from BD and staining was performed in the presence of anti-CD16/CD32 (clone 2.4G2). Flow cytometry was performed using an Accuri C6 Flow Cytometer (BD) and analyzed using BD Accuri C6 Software.

Tumor immunotherapy

[00468] For *in vivo* tumor immunotherapy studies, female C57BL/6 mice, 8–10 weeks old, were implanted with 2 \times 10⁵ TC-1 HPV-E6/E7 expressing tumor cells SQ in the left flank. Mice were treated three times at 7-day intervals with SQ injections of 10¹⁰ VP Ad5 [E1-, E2b-]-E6/E7. Control mice were injected with 10¹⁰ VP Ad5 [E1-, E2b-]-null under the same protocol. In combinational studies, mice were given 100 μ g of rat anti-PD1 (clone RMP1-14) or an isotype rat control antibody (clone 2A3) IP at the same time as immunization. Rat anti-PD1 antibody and rat IgG_{2a} isotype control antibodies were purchased from BioXcell. Tumor size was measured by two opposing dimensions (a, b) and volume was calculated according to the formula $V=(a^2xb)/2$ where a was the shorter dimension. Animals were euthanized when tumors reached 1500mm³ or when tumors became ulcerated.

Analysis of tumor infiltrating cells (TILs) by flow cytometry

[00469] Four groups of 8–10 week old female C57BL/6 mice (n=5/group) were implanted with 2 \times 10⁵ TC-1 tumor cells SQ in the left flank at day 0. Two of these groups were immunized SQ with 10¹⁰ VP Ad5 [E1-, E2b-]-null vector control and the other two groups SQ with 10¹⁰ VP Ad5 [E1-, E2b-]-E6/E7 vaccine. These immunizations were administered twice at 7-day intervals starting on day 12. In addition to immunizations, mice in one Ad5 [E1-, E2b-]-E6/E7 group and one Ad5 [E1-, E2b-]-null group were administered 100 μ g rat anti-PD1 (clone RMP1-14) SQ at days 12 and 16 and 100 μ g hamster anti-PD1 (clone J43) at days 19 and 23 to increase the effective dose of anti-PD1. To control for treatment with these immune pathway checkpoint modulators, mice in the remaining Ad5 [E1-, E2b-]-E6/E7 and Ad5 [E1-, E2b-]-null groups were administered the relevant rat and hamster control IgG antibodies on the same days. Hamster anti-PD1 antibody and isotype control were purchased from BioXcell. At day 27, tumors were measured, excised, and weighed. Tumors were minced and digested with a mixture of collagenase IV (1mg/ml), hyaluronidase (100 μ g/ml), and DNase IV (200U/ml) in Hank's Balanced Salt Solution (HBSS) at room temperature for 30 min and rotating at 80 rpm. Enzymes were purchased from Sigma-Aldrich. After digestion, the tumor suspension was placed through a 70 μ m nylon cell strainer and centrifuged. Red cells were removed by the addition of red cell lysis buffer (Sigma-Aldrich) and after

lysis, the tumor suspensions were washed twice in phosphate buffered saline (PBS) containing 1% (w/v) bovine serum albumin and resuspended in fluorescent activated cell sorting (FACS) buffer (PBS pH 7.2, 1% fetal bovine serum, and 2 mM EDTA) for staining. Fluorescent-conjugated antibodies against CD45 (30-F11), CD4 (RM4-5), and PDL1 (MIH5) were purchased from BD. Fluorescent-conjugated antibodies against CD8 β (H35-17.2), CD25 (PC61.5), FoxP3 (FJK-16s), PD1 (RMP1-30), LAG-3 (C9B7W), and CTLA4 (UC10-4B9) were all purchased from eBioscience. Surface staining was performed for 30 minutes at 4 °C in 100 μ L FACS buffer containing anti-CD16/CD32 (clone 2.4G2). Stained cells were washed in FACS buffer, fixed with paraformaldehyde, and (if needed) permeabilized in permeabilization buffer (eBioscience) before staining with fluorescent-conjugated anti-FoxP3 or anti-CTLA4 for 60 minutes at 4°C in 100 μ L permeabilization buffer containing anti-CD16/CD32 (clone 2.4G2). Cells were washed with permeabilization buffer, washed back into FACS buffer, and a fixed volume of each sample was analyzed by flow cytometry using a BD Accuri C6 flow cytometer. Tumor cells were defined as CD45 $^{+}$ events in a scatter gate that includes small and large cells. CD4 $^{+}$ TILs were defined as CD45 $^{+}$ /CD4 $^{+}$ events in a lymphocyte scatter gate. CD8 $^{+}$ TILs were defined as CD45 $^{+}$ /CD8 β $^{+}$ events in a lymphocyte scatter gate. Regulatory T cells (Tregs) were defined as CD45 $^{+}$ /CD4 $^{+}$ /CD25 $^{+}$ /FoxP3 $^{+}$ events in a lymphocyte scatter gate. Effector CD4 $^{+}$ T cells were defined as CD45 $^{+}$ /CD4 $^{+}$ /CD25 $^{-}$ /FoxP3 $^{-}$ events in a lymphocyte scatter gate. Isotype-matched control antibodies were used to determine positive expression of FoxP3, PDL1, PD1, LAG-3, and CTLA4. Flow cytometry was performed using an Accuri C6 Flow Cytometer (BD) and analyzed in BD Accuri C6 Software.

HPV-E6/E7 specific cell mediated immune responses induced by Ad5 [E1-, E2b-]-E6/E7

[00470] A study was performed to determine the effect of increasing doses of Ad5 [E1-, E2b-]-E6/E7 immunizations on the induction of CMI responses in mice. Groups of C57BL/6 mice (n=5/group) were immunized SQ three times at 14-day intervals with 10⁸, 10⁹, or 10¹⁰ VP Ad5 [E1-, E2b-]-E6/E7. Control mice received 10⁸ VP, 10⁹ VP, or 10¹⁰ VP Ad5 [E1-, E2b-]-null (empty vector controls). Two weeks after the last immunization, splenocyte CMI responses were assessed by ELISpot analysis for IFN- γ secreting cells. A dose effect was observed and the highest CMI response level was obtained by immunizations with 10¹⁰ VP Ad5 [E1-, E2b-]-E6/E7. No responses were detected in control mice injected with Ad5 [E1-, E2b-]-null.

[00471] Intracellular accumulation of IFN- γ and TNF- α in both CD8 α $^{+}$ and CD4 $^{+}$ splenocytes populations was also determined in mice immunized with 10¹⁰ VP Ad5 [E1-, E2b-]-E6/E7. Intracellular cytokine staining (ICS) after stimulation with overlapping peptide pools revealed E6 and E7 antigen-specific IFN- γ accumulation in CD8 α $^{+}$ lymphocytes isolated from all mice immunized with Ad5 [E1-, E2b-]-E6/E7. Peptide-stimulated splenocytes were also stained for the intracellular accumulation of TNF-

α , and a significant population of multifunctional (IFN- γ ⁺/TNF- α ⁺) CD8 α ⁺ splenocytes specific for both E6 and E7 were able to be detected.

Treatment of HPV-E6/E7 expressing tumors

[00472] The anti-tumor effect of immunotherapy treatment in mice bearing HPV-E6/E7 TC-1 tumors was investigated. These tumor cells expressed PDL1 as assessed by flow cytometry analysis. When labeled with PE-conjugated anti-PDL1, the TC-1 cells had a median fluorescent intensity (MFI) of 537 whereas cells labeled with a PE-conjugated isotype control antibody had an MFI of 184, demonstrating the presence of the immune suppressive PDL1 on the surface of the TC-1 cells (data not shown). Two groups of C57BL/6 mice (n=5/group) were inoculated with 2×10^5 TC-1 tumor cells SQ into the right subcostal area on day 0. On days 1, 8, and 14 mice were treated by SQ injections of 10^{10} VP Ad5 [E1-, E2b-]-null (vector control) or 10^{10} VP Ad5 [E1-, E2b-]-E6/E7. All mice were monitored for tumor size and tumor volumes were calculated. Mice immunized with Ad5 [E1-, E2b-]-E6/E7 had significantly smaller tumors than control mice beginning on day 12 (p<0.01) and remained significantly smaller for the remainder of the experiment (p<0.02), including 3 of 5 mice showing complete tumor regression. Tumors in mice from the vector control treated group began reaching the threshold for euthanasia starting on day 26 and all mice in this group were euthanized by day 33, whereas mice in the Ad5 [E1-, E2b-]-E6/E7 treated group were all alive with complete tumor regression of small tumors (<150mm³) at the end of experiment on day 36.

[00473] To determine if immunotherapy with Ad5 [E1-, E2b-]-E6/E7 was effective against larger tumors, TC-1 tumor cells tumors were implanted in two groups of C57BL/6 mice (n=4/group) and then delayed weekly treatment with Ad5 [E1-, E2b-]-E6/E7 for 6 days post tumor implantation, at a time when tumors were small but palpable. Mice beginning treatment on day 6 initially demonstrated tumor growth similar to the control group; however, beginning on day 16, tumor regression was observed. The tumors in mice that began treatment on day 6 were significantly smaller (p<0.05) than the control group beginning on day 20 and 3 of 4 mice had complete regression by day 27. Ad5 [E1-, E2b-]-E6/E7 administration beginning on day 6 also conferred a significant survival benefit (p<0.01).

[00474] Finally, to determine if immunotherapy with Ad5 [E1-, E2b-]-E6/E7 was effective against large established tumors, TC-1 tumor cells were implanted in two groups of C57BL/6 mice (n=4/group) then delayed weekly treatment with Ad5 [E1-, E2b-]-E6/E7 until 13 days post tumor implantation, when tumors were ~100mm³. In this treatment group, initial tumor growth was observed to be similar to the control group but some mice in the control group reached euthanasia criteria on day 23, preventing analysis of significance at further time points. However, tumor volumes in the Ad5 [E1-, E2b-]-E6/E7 treated group were below the euthanasia threshold through day 29, at which point tumors from all mice in the vector control group had exceeded 1500mm³ and were euthanized (Figure 5B). These results indicate that in the TC-1 tumor model the Ad5 [E1-, E2b-]-E6/E7 immunotherapeutic was a potent inhibitor of

tumor growth and lead to significant overall survival benefit, however complete clearance of tumors was only observed when treatment was initiated in smaller tumors. Furthermore, these results demonstrate that, despite the presence of immune suppressing PDL1 on tumor cells, immunotherapeutic treatment with Ad5 [E1-, E2b-]-E6/E7 resulted in significant inhibition of tumor growth.

Combination immunotherapy with immune checkpoint inhibition

[00475] To determine if the therapeutic effect of Ad5 [E1-, E2b-]-E6/E7 could be improved in the setting of large tumors, anti-PD1 antibody was co-administered. Four groups of mice (n=7/group) were implanted with 2×10^5 TC-1 tumor cells on day 0 and beginning on day 10 the mice received weekly administrations of SQ 10^{10} VP Ad5 [E1-, E2b-]-E6/E7 plus IP 100 μ g anti-PD1, 10^{10} VP Ad5 [E1-, E2b-]-null plus 100 μ g anti-PD1, 10^{10} VP Ad5 [E1-, E2b-]-E6/E7 plus 100 μ g rat IgG_{2a} isotype control, or 10^{10} VP Ad5 [E1-, E2b-]-null plus 100 μ g rat IgG_{2a} isotype control. Tumor size was monitored over time and mice were euthanized when tumor size exceeded 1500 mm³ or when tumor ulceration was present. Control mice that received Ad5 [E1-, E2b-]-null plus 100 μ g rat IgG_{2a} isotype control (Figure 6A) and mice treated with Ad5 [E1-, E2b-]-null plus 100 μ g anti-PD1 (Figure 6B) exhibited a similar tumor growth pattern (Figure 6B). No significant survival benefit was observed between these two groups. Mice that received Ad5 [E1-, E2b-]-E6/E7 plus rat IgG_{2a} isotype control had a delayed tumor growth pattern as compared to the controls and 2 of the mice had tumor regressions to near baseline level at day 52 post tumor implantation (Figure 6C). Four of the 7 mice that received Ad5 [E1-, E2b-]-E6/E7 and anti-PD1 had tumor regression starting at day 25, and two of these resulted in tumor clearance through the end of experiment at day 53 (Figure 6D).

[00476] Mice treated with Ad5 [E1-, E2b-]-E6/E7 plus rat IgG_{2a} isotype control (Figure 7) also experienced a survival benefit with 28.6% of the animals surviving at termination of the study whereas 100% of the control mice (Ad5 [E1-, E2b-]-null plus rat IgG_{2a} isotype control) and the Ad5 [E1-, E2b-]-null plus anti-PD1 treated mice had to be terminated by day 28 and 32, respectively (Figure 7). Mice treated with both Ad5 [E1-, E2b-]-E6/E7 and anti-PD1 antibody had the greatest treatment benefit (Figure 7), demonstrating delayed tumor growth and a significant improvement ($P \leq 0.0006$) in survival as compared to the controls.

[00477] Mouse anti-rat IgG antibody responses were induced by the second injection (endpoint antibody titer 1:200 by ELISA, data not shown) with rat anti-PD1 antibody and these responses were dramatically increased by the third injection (endpoint antibody titer 1:4000 to 1:8000 by ELISA, data not shown). This anti-rat antibody response may explain why no anti-tumor activity was observed after injections with anti-PD1 antibody alone. Also, it is likely that the first and possibly the second injections of anti-PD1 antibody combined with Ad5 [E1-, E2b-]-E6/E7 immunotherapy were effective but the third injection with anti-PD1 was effectively neutralized by the induced mouse anti-rat IgG response.

Tumor microenvironment following combination immunotherapy

[00478] To analyze cell populations that contributed to delayed tumor growth and survival in Ad5 [E1-, E2b-]-E6/E7 treated mice, tumor infiltrating lymphocytes (TILs) were by flow cytometry. Four groups of mice were implanted with 2×10^5 TC-1 cells and began treatment 10 days later with two weekly immunizations of Ad5 [E1-, E2b-]-E6/E7 plus PD1 antibody. On day 27 whole tumors were collected and processed as described in the materials and methods. The number of infiltrating CD8⁺ T cells per mg of tumor was significantly increased in the Ad5 [E1-, E2b-]-E6/E7 treated groups as compared to the groups that received Ad5 [E1-, E2b-]-null (Figure 8C). Anti-PD1 antibody treatment had little or no effect on the number of infiltrating CD8⁺ T cells (Figure 8C). There was no difference between any of the four groups, in terms of the number of infiltrating Tregs (CD4⁺CD25⁺Foxp3⁺) per mg of tumor (Figure 8B). However, the increase in CD8⁺ T cells led to a decrease in the Treg:CD8⁺ T cell ratio in the tumor microenvironment when the mice were treated with the Ad5 [E1-, E2b-]-E6/E7 vaccine or Ad5 [E1-, E2b-]-E6/E7 vaccine plus anti-PD1 antibody treatment (Figure 8A).

[00479] To further study the synergistic/additive effect of anti-PD1 antibody to Ad5 [E1-, E2b-]-E6/E7 immunotherapy, the expression of PD1, LAG-3, and CTLA-4 was examined on TILs. The expression of these co-inhibitory molecules on T cells within the tumor microenvironment has been shown to down regulate activation of antigen-specific T cells. Immunizations with Ad5 [E1-, E2b-]-E6/E7 plus control antibody treatment significantly increased the fraction of PD1⁺ and LAG-3⁺ CD8⁺ TILs, whereas, expression of these co-inhibitory molecules on CD4⁺ TILs was unaffected by this treatment. The percentage of CD4⁺ and CD8⁺ TILs expressing CTLA-4 was not significantly affected by vaccine treatment (data not shown). Combining anti-PD1 antibody injections with Ad5 [E1-, E2b-]-E6/E7 vaccine treatment resulted in a significant reduction in the fraction of PD1⁺ CD8⁺ and CD4⁺ TILs, as compared with those found in tumors from mice treated with Ad5 [E1-, E2b-]-E6/E7 plus control antibody ($p=0.0083$ for CD8⁺ TILs and $p=0.0016$ for CD4⁺ TILs). Furthermore the fraction of PD1⁺ CD8⁺ TILs was decreased to the level of expression observed in the Ad5 [E1-, E2b-]-null treated control groups, and the fraction of PD1⁺ CD4⁺ TILs was significantly reduced to below that observed in the control groups ($p=0.0016$, Figure 9A). In addition, the percentage of LAG-3⁺ CD8⁺ TILs was also observed to decrease when the Ad5 [E1-, E2b-]-E6/E7 immunization was combined with the anti-PD1 checkpoint inhibitor ($p=0.0363$, Figure 9B). Since it has previously been shown that vaccine treatment can enhance PDL1 expression on tumor cells *ex vivo*, the expression of PDL1 was examined on tumor cells. There was an augmentation in the median fluorescence intensity of PDL1 on tumor cells after vaccine treatment. However, PDL1 expression was reduced in mice treated with the combination of Ad5 [E1-, E2b-]-E6/E7 and anti-PD1 antibody, although this level was still significantly expressed above that observed in Ad5 [E1-, E2b-]-null treated control mice.

[00480] In summary, the data demonstrate that Ad5 [E1-, E2b-]-E6/E7 can induce HPV-E6/E7 directed CMI responses in a dose dependent manner, which results in upregulation of PDL1 on tumor cells. Multiple homologous immunizations in tumor bearing mice with the highest dose of vaccine resulted in significant anti-tumor activity and increased survival, particularly in mice bearing small tumors. Importantly, a greater degree of anti-tumor activity was achieved when immunotherapy with Ad5 [E1-, E2b-]-E6/E7 was combined with anti-PD1 in mice with large tumors. Overall, immunizations with the Ad5 [E1-, E2b-]-E6/E7 vaccine combined with anti-PD1 antibody results in an increase in CD8⁺ and CD4⁺ effector populations that have a less exhaustive/anergic phenotype and therefore favor the balance to a more pro-inflammatory state in the tumor microenvironment. The observation that the combined treatment was associated with reductions in large tumor mass indicates that immunotherapy with Ad5 [E1-, E2b-]-E6/E7 combined with anti-PD1 antibody might increase clinical effectiveness during the immunotherapy of patients with HPV-associated head and neck or cervical cancers. Furthermore, the data suggests that clinical trials with the Ad5 [E1-, E2b-]-E6/E7 vaccine should be combined with an immune pathway checkpoint modulator and remains a high priority.

[00481] The various embodiments described above can be combined to provide further embodiments. Aspects of the embodiments can be modified, if necessary to employ concepts of the various patents, application and publications to provide yet further embodiments. These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

SEQUENCE LISTING

[00482] This description contains a sequence listing in electronic form in ASCII text format. A copy of the sequence listing is available from the Canadian Intellectual Property Office.

CLAIMS

WHAT IS CLAIMED IS:

1. A composition comprising:
 - (a) a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a MUC1 antigen;
 - (b) a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a Brachyury antigen; and
 - (c) a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a CEA antigen,
wherein the MUC1 antigen is a MUC1-C antigen.
2. The composition of claim 1, wherein an amino acid sequence of the MUC1-C antigen has at least 90% sequence identity to SEQ ID NO: 9.
3. The composition of claim 1 or 2, wherein the nucleotide sequence encoding the Brachyury antigen comprises SEQ ID NO: 8.
4. The composition of claim 1, 2 or 3, wherein the nucleotide sequence encoding the CEA antigen comprises SEQ ID NO: 1.
5. The composition of any one of claims 1 to 4, wherein the viral vector comprising the nucleotide sequence encoding the MUC1-C antigen, the viral vector comprising the nucleotide sequence encoding the Brachyury antigen, and the viral vector comprising the nucleotide sequence encoding the CEA antigen comprise a deletion in an E3 gene region, an E4 gene region, or any combination thereof.

6. The composition of any one of claims 1 to 5, further comprising an immunogenic component, wherein the immunogenic component comprises a cytokine selected from the group of IFN- γ , TNF α IL-2, IL-8, IL-12, IL-18, IL-7, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, and IL-13.

7. The composition of any one of claims 1 to 6, wherein a viral vector comprises a sequence encoding at least two antigens selected from the group consisting of the MUC1-C antigen, the Brachyury antigen, and the CEA antigen.

8. The composition of any one of claims 1 to 7, wherein an amino acid sequence of the CEA antigen comprises SEQ ID NO: 10.

9. A composition for treating a subject in need thereof, wherein the composition comprises:

a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a MUC1 antigen, a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a Brachyury antigen; and a recombinant, replication defective adenovirus 5 viral vector comprising an E2b gene region deletion, an E1 gene region deletion, and a nucleotide sequence encoding a CEA antigen, characterized in that the composition is for administration to the subject in need thereof; and

induces an immune response against the MUC1 antigen, the Brachyury antigen, or the CEA antigen or cells expressing the MUC1 antigen, the Brachyury antigen, or the CEA antigen,

wherein the subject in need thereof has cancer,

wherein the MUC1 antigen is a MUC1-C antigen.

10. The composition of claim 9, wherein the composition comprises 5×10^{11} virus particles (VPs) of the viral vector comprising the nucleotide sequence encoding the MUC1-C antigen, at

least 5×10^{11} virus particles (VPs) of the viral vector comprising the nucleotide sequence encoding the Brachyury antigen, and at least 5×10^{11} virus particles (VPs) of the viral vector comprising the nucleotide sequence encoding the CEA antigen.

11. The composition of claim 9 or 10, wherein the immune response is at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or 25 fold over basal.

12. The composition of claim 9, 10, or 11, wherein the immune response is measured as antigen specific antibody response, as antigen specific cell-mediated immunity (CMI), as antigen specific IFN- γ secretion, as antigen specific IL-2 secretion, by ELISpot assay, or any combination thereof.

13. The composition of any one of claims 9 to 12, wherein the compound is for subcutaneous administration.

14. The composition of any one of claims 9 to 13, wherein the subject has colorectal adenocarcinoma; metastatic colorectal cancer; advanced MUC1-C, Brachyury, or CEA expressing colorectal cancer; breast cancer; lung cancer; bladder cancer; or pancreas cancer.

15. The composition of any one of claims 9 to 14, wherein the subject in need thereof is a human.

16. The composition of any one of claims 9 to 15, wherein an amino acid sequence of the MUC1-C antigen has at least 90% sequence identity to SEQ ID NO: 9, the nucleotide sequence encoding the Brachyury comprises SEQ ID NO: 8, the nucleotide sequence encoding the CEA antigen comprises SEQ ID NO: 1, an amino acid sequence of the CEA antigen comprises SEQ ID NO: 10, or any combination thereof.

1/38




FIG. 1





FIG. 2

2/38



3/38




FIG. 4A




FIG. 4B

4/38




FIG. 5




FIG. 6

5/38




FIG. 7




FIG. 8

6/38




FIG. 9

7/38




FIG. 10A

FIG. 10B

8/38




FIG. 11

9/38




FIG. 12B




FIG. 12D




FIG. 12A




FIG. 12C

10/38




FIG. 13B




FIG. 13A

11/38




FIG. 14

12/38




FIG. 15

13/38




FIG. 16B




FIG. 16A

14/38




FIG. 17A

FIG. 17B




FIG. 17C




FIG. 18




FIG. 19

16/38




FIG. 21




FIG. 20

17/38




FIG. 23




FIG. 22

18/38




FIG. 24A




FIG. 24B

19/38




FIG. 25A

FIG. 25B

20/38




FIG. 26A




FIG. 26B




FIG. 27




FIG. 28

22/38




FIG. 29

23/38




FIG. 30A



FIG. 30B

24/38




FIG. 31A

FIG. 31B

25/38




FIG. 32B

FIG. 32A

26/38




FIG. 32D

FIG. 32C

27/38




FIG. 33A




FIG. 33B




FIG. 34




FIG. 35

29/38




FIG. 36

30/38




FIG. 37A

FIG. 37B




FIG. 38A




FIG. 38B

32/38




FIG. 39A




FIG. 39B

33/38




FIG. 40A




FIG. 40B

34/38




FIG. 41A




FIG. 41B

35/38




FIG. 41C




FIG. 41D

36/38




FIG. 42

37/38




FIG. 43B





FIG. 43A




FIG. 43C

38/38




FIG. 44B




FIG. 44A




FIG. 44C