

(12) STANDARD PATENT APPLICATION (11) Application No. AU 2022217319 A9
(19) AUSTRALIAN PATENT OFFICE

(54) Title
TRICYCLIC COMPOUNDS AND USES THEREOF

(51) International Patent Classification(s)
C07D 471/04 (2006.01) **A61P 35/00** (2006.01)
A61K 31/444 (2006.01) **C07D 401/14** (2006.01)
A61K 31/496 (2006.01) **C07D 487/04** (2006.01)

(21) Application No: **2022217319** (22) Date of Filing: **2022.01.28**

(87) WIPO No: **WO22/166844**

(30) Priority Data

(31) Number	(32) Date	(33) Country
202110166021.0	2021.02.05	CN
202210069346.1	2022.01.21	CN

(43) Publication Date: **2022.08.11**

(48) Corrigenda Journal Date: **2024.10.17**

(71) Applicant(s)
HUTCHMED LIMITED

(72) Inventor(s)
DAI, Guangxiu;XIAO, Kun;DENG, Wei

(74) Agent / Attorney
WRAYS PTY LTD, L7 863 Hay St, Perth, WA, 6000, AU



(51) International Patent Classification:

C07D 471/04 (2006.01) A61K 31/444 (2006.01)

C07D 487/04 (2006.01) A61K 31/496 (2006.01)

C07D 401/14 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/CN2022/074753

(22) International Filing Date:

28 January 2022 (28.01.2022)

Published:

— with international search report (Art. 21(3))

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

202110166021.0 05 February 2021 (05.02.2021) CN

202210069346.1 21 January 2022 (21.01.2022) CN

(71) Applicant: **HUTCHMED LIMITED** [CN/CN]; Building 4, 720 Cailun Road, Pilot Free Trade Zone, Shanghai 201203 (CN).


(72) Inventors: **DAI, Guangxiu**; Building 4, 720 Cailun Road, Pilot Free Trade Zone, Shanghai 201203 (CN). **XIAO, Kun**; Building 4, 720 Cailun Road, Pilot Free Trade Zone, Shanghai 201203 (CN). **DENG, Wei**; Building 4, 720 Cailun Road, Pilot Free Trade Zone, Shanghai 201203 (CN).

(74) Agent: **ZHONGZI LAW OFFICE**; 7F, New Era Building, 26 Pinganli Xidajie, Xicheng District, Beijing 100034 (CN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

(54) Title: TRICYCLIC COMPOUNDS AND USES THEREOF



(I)

(57) Abstract: Tricyclic compounds of formula (I), pharmaceutical compositions comprising same, methods for preparing same, and uses thereof, wherein each variable is as defined in the description.



TRICYCLIC COMPOUNDS AND USES THEREOF

Technical Field

5 The present invention relates to tricyclic compounds, pharmaceutical compositions comprising same, methods for preparing same, and uses thereof.

Background Art

10 SHP2 (Src homology-2 domain-containing protein tyrosine phosphatase-2) is a non-receptor protein tyrosine phosphohydrolase encoded by a PTPN11 gene. SHP2 contains two SH2 domains (N-terminal SH2 domain and C-terminal SH2 domain), a catalytic domain and a C-terminal sequence containing two tyrosine phosphorylation sites. The non-receptor protein tyrosine phosphohydrolase subfamily contains two members, SHP1 and SHP2. The SHP1 and SHP2 proteins have 61% similarity in amino acid sequences, and have up to 75% amino acid identity in the catalytic domains (PTP). SHP1 is mainly expressed in hematopoietic cells and some epithelial cells, and is mainly involved in the negative regulation of intracellular signaling. SHP2 is widely expressed in various organs of the human body, and plays an important physiological role in the growth and development and homeostasis of the human body. After being stimulated by growth factors or hormones, SHP2 participates in the signal regulation and transduction of many signaling pathways, including RAS-ERK, JAK-STAT, PI3K-AKT, and PD1-PD-L1, thereby promoting many biological functions, such as cell proliferation, differentiation and migration.

25 Mutation or overexpression of SHP2 would lead to hereditary developmental diseases and tumors. It is reported that in the hereditary developmental diseases, 90% of the LEOPARD Syndrome cases and 50% of the Noonan Syndrome cases have been found to have gain-of-function (GOF) mutations in the PTPN11 gene. Mutations in the PTPN11 gene have also been reported in hematologic malignancies, such as myelodysplastic syndrome (10%), B-cell acute lymphocytic leukemia (7%), and juvenile acute myelogenous leukemia (5%). Mutations in SHP2 are rarely found in solid tumors, but overexpression/activation of SHP2 is closely related to the occurrence of a variety of tumors. SHP2 expression in invasive ductal breast cancer has increased by 70%;


overexpression of the SHP2 binding protein, GAB2 has also been detected in 10%-15% of breast cancers; and overexpression of SHP2 in melanoma is often closely related to poor prognosis.

SHP2 is closely related to tumors, making it becoming an attractive anti-tumor drug target. Currently, several selective SHP2 inhibitors such as TNO155 (Novartis), RMC-4630 (Revolution Medicines/Sanofi), JAB-3068 (Jacobio), and RLY-1971 (Relay Therapeutics) have entered clinical studies; however, no SHP2 inhibitor has been approved for marketing.

Therefore, SHP2 inhibitors stand for the development of an attractive therapy for related diseases, especially cancer, Noonan Syndrome and LEOPARD Syndrome.

Summary of the Invention

The present invention provides a compound of formula (I):



or a pharmaceutically acceptable salt thereof, or a solvate, a racemic mixture, an enantiomer, a diastereomer or a tautomer thereof, wherein

ring A is a benzene ring or a pyridine ring;

Z is CH₂, O, S or NH;

R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄, -SR₅ and -SR₆, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-CN, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and -(C₁₋₆ alkyl)-CN; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, phenyl, 4-8 membered heterocyclyl and 5-12

membered heteroaryl; R_6 is chosen from $-\text{CO}(\text{C}_{1-6} \text{ alkyl})$, $-\text{CO}(\text{C}_{3-8} \text{ cycloalkyl})$, $-\text{CO}(4-8 \text{ membered heterocyclyl})$, $-\text{CONH}_2$, $-\text{CONH}(\text{C}_{1-6} \text{ alkyl})$, $-\text{CONH}(\text{C}_{3-8} \text{ cycloalkyl})$, $-\text{CONH}(4-8 \text{ membered heterocyclyl})$, $-\text{CON}(\text{C}_{1-6} \text{ alkyl})_2$, $-(\text{C}_{1-6} \text{ alkyl})-\text{O}-(\text{C}_{1-6} \text{ alkyl})$, $-(\text{C}_{1-6} \text{ alkyl})-\text{OH}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{NH}(\text{C}_{1-6} \text{ alkyl})$, $-(\text{C}_{1-6} \text{ alkyl})-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$ and $-(\text{C}_{1-6} \text{ alkyl})-\text{NHCO}(\text{C}_{1-6} \text{ alkyl})$, wherein the $\text{C}_{1-6} \text{ alkyl}$ of R_6 is optionally substituted with one or more groups independently chosen from: halogen, $-\text{CN}$, $-\text{OH}$ and $-\text{O}(\text{C}_{1-6} \text{ alkyl})$; and the above-mentioned $\text{C}_{3-8} \text{ cycloalkyl}$, phenyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: halogen, $-\text{CN}$, $-\text{CONH}_2$, $-\text{OH}$, oxo, $-\text{NH}_2$, $\text{C}_{1-6} \text{ alkyl}$, $\text{C}_{2-6} \text{ alkenyl}$, $\text{C}_{2-6} \text{ alkynyl}$, $\text{C}_{1-6} \text{ haloalkyl}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{OH}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{O}-(\text{C}_{1-6} \text{ alkyl})$, $-(\text{C}_{1-6} \text{ alkyl})-\text{CN}$, $-\text{O}(\text{C}_{1-6} \text{ alkyl})$, $-\text{O}(\text{C}_{1-6} \text{ haloalkyl})$, $-\text{S}(\text{C}_{1-6} \text{ alkyl})$, $-\text{NH}(\text{C}_{1-6} \text{ alkyl})$ and $-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$;

R_1' is independently chosen from halogen, $-\text{CN}$, $-\text{CONH}_2$, $-\text{OH}$, $-\text{NH}_2$, $\text{C}_{1-6} \text{ alkyl}$, $\text{C}_{1-6} \text{ haloalkyl}$, $\text{C}_{3-8} \text{ cycloalkyl}$, 4-8 membered heterocyclyl, $-(\text{C}_{1-6} \text{ alkyl})-\text{OH}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{O}-(\text{C}_{1-6} \text{ alkyl})$, $-(\text{C}_{1-6} \text{ alkyl})-\text{CN}$, $-\text{O}(\text{C}_{1-6} \text{ alkyl})$, $-\text{O}(\text{C}_{1-6} \text{ haloalkyl})$, $-\text{O}(\text{C}_{3-8} \text{ cycloalkyl})$, $-\text{O}(4-8 \text{ membered heterocyclyl})$, $-\text{S}(\text{C}_{1-6} \text{ alkyl})$, $-\text{NH}(\text{C}_{1-6} \text{ alkyl})$ and $-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$, wherein the $\text{C}_{3-8} \text{ cycloalkyl}$ and 4-8 membered heterocyclyl are each optionally substituted with one or more halogen;

n is 0, 1, 2 or 3;

R_2 is chosen from $-\text{NH}_2$, $-\text{NH}(\text{C}_{1-6} \text{ alkyl})$, $-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$, oxo and $-\text{OH}$;

Cy_1 is 5-12 membered heterocyclyl or 5-12 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-\text{CN}$, $-\text{CONH}_2$, $-\text{OH}$, oxo, $-\text{NH}_2$, $\text{C}_{1-6} \text{ alkyl}$, $\text{C}_{2-6} \text{ alkenyl}$, $\text{C}_{2-6} \text{ alkynyl}$, $\text{C}_{1-6} \text{ haloalkyl}$, $\text{C}_{3-8} \text{ cycloalkyl}$, 4-8 membered heterocyclyl, $-(\text{C}_{1-6} \text{ alkyl})-\text{OH}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{O}-(\text{C}_{1-6} \text{ alkyl})$, $-(\text{C}_{1-6} \text{ alkyl})-\text{CN}$, $-\text{O}(\text{C}_{1-6} \text{ alkyl})$, $-\text{O}(\text{C}_{1-6} \text{ haloalkyl})$, $-\text{O}(\text{C}_{3-8} \text{ cycloalkyl})$, $-\text{O}(4-8 \text{ membered heterocyclyl})$, $-\text{S}(\text{C}_{1-6} \text{ alkyl})$, $-\text{NH}(\text{C}_{1-6} \text{ alkyl})$, $-\text{N}(\text{C}_{1-6} \text{ alkyl})_2$, $-\text{NHCO}(\text{C}_{1-6} \text{ alkyl})$, $-\text{CONH}(\text{C}_{1-6} \text{ alkyl})$ and $-\text{CON}(\text{C}_{1-6} \text{ alkyl})_2$, wherein the $\text{C}_{3-8} \text{ cycloalkyl}$ and 4-8 membered heterocyclyl are each optionally substituted with one or more halogen;

Cy_2 is phenyl or 5-14 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-\text{CN}$, $-\text{CONH}_2$, $-\text{OH}$, oxo, $\text{C}_{1-6} \text{ alkyl}$, $\text{C}_{2-6} \text{ alkenyl}$, $\text{C}_{2-6} \text{ alkynyl}$, $\text{C}_{1-6} \text{ haloalkyl}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{OH}$, $-(\text{C}_{1-6} \text{ alkyl})-\text{O}-(\text{C}_{1-6} \text{ alkyl})$, $-(\text{C}_{1-6} \text{ alkyl})-\text{CN}$, $-\text{O}(\text{C}_{1-6} \text{ alkyl})$, $-\text{O}(\text{C}_{1-6} \text{ haloalkyl})$, $-\text{O}(\text{C}_{3-8} \text{ cycloalkyl})$, $-\text{O}(4-8 \text{ membered heterocyclyl})$, $-\text{S}(\text{C}_{1-6} \text{ alkyl})$, $-\text{NR}_7\text{R}_8$, $-\text{NHCO}(\text{C}_{1-6} \text{ alkyl})$, $-\text{CONH}(\text{C}_{1-6} \text{ alkyl})$

and -CON(C₁₋₆ alkyl)₂, wherein R₇ and R₈ are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-CN, C₃₋₈ cycloalkyl, phenyl, 3-8 membered heterocyclyl and 5-6 membered heteroaryl; and L is absent, or L is S, O, NH, C₁₋₆ alkyl, C₂₋₆ alkenyl or C₂₋₆ alkynyl.

5 The above-mentioned compounds and the active compounds (including compounds of general formula and specific compounds) disclosed in the context of the present invention, and pharmaceutically acceptable salts thereof, or solvates, racemic mixtures, enantiomers, diastereomers or tautomers thereof are collectively referred to herein as “compounds of the present invention”.

10 The present invention also provides a pharmaceutical composition, comprising the compounds of the present invention, and optionally comprising a pharmaceutically acceptable excipient.

The present invention also provides a method of *in vivo* or *in vitro* inhibiting the activity of SHP2, comprising contacting SHP2 with an effective amount of the
15 compounds of the present invention.

The present invention also provides a method of treating or preventing a disease mediated by SHP2 or at least in part by SHP2, comprising administering to the subject in need thereof an effective amount of the compounds of the present invention.

The present invention also provides a method of treating or preventing cancer,
20 Noonan Syndrome or LEOPARD Syndrome, comprising administering to the subject in need thereof an effective amount of the compounds of the present invention.

The present invention also provides the use of the compounds of the present invention in the treatment or prevention of a disease mediated by SHP2 or at least in part by SHP2.

25 The present invention also provides the use of the compounds of the present invention in the treatment or prevention of cancer, Noonan Syndrome or LEOPARD Syndrome.

The present invention also provides the use of the compounds of the present invention in the manufacture of a medicament for treating or preventing a disease
30 mediated by SHP2 or at least in part by SHP2.

The present invention also provides the use of the compounds of the present invention in the manufacture of a medicament for treating or preventing cancer, Noonan Syndrome or LEOPARD Syndrome.

5 The present invention also provides the compounds of the present invention for *in vivo* or *in vitro* inhibiting the activity of SHP2.

The present invention also provides the compounds of the present invention for use as a medicament.

10 The present invention also provides the compounds of the present invention for use as a medicament for treating or preventing a disease mediated by SHP2 or at least in part by SHP2, especially for treating or preventing cancer, Noonan Syndrome or LEOPARD Syndrome.

15 The present invention also provides a pharmaceutical combination, comprising the compounds of the present invention and at least one additional therapeutic agent, wherein the additional therapeutic agent is preferably chosen from: an anti-neoplastic active agent, an anti-inflammatory agent or an immunomodulator, wherein the anti-neoplastic active agent includes a chemotherapeutic agent, an immune checkpoint inhibitor or agonist, and a targeted therapeutic agent.

20 The present invention also provides a kit for treating or preventing a disease mediated by SHP2 or at least in part by SHP2. The kit can comprise the pharmaceutical composition of the present invention and instructions for use, and the pharmaceutical composition comprises the compounds of the present invention.

Detailed Description of the Invention

Definitions

25 As used in the present application, the following words, phrases and symbols are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise.

30 A dash (“-”) that is not between two letters or symbols is used to indicate a point of attachment for a substituent. For example, -OR₆ refers to the attachment of R₆ to the rest of the molecule through an oxygen atom.

The term “alkyl” as used herein refers to a straight or branched saturated hydrocarbon radical containing 1-18 carbon atoms (C₁₋₁₈), preferably 1-10 carbon atoms

(C₁₋₁₀), more preferably 1-6 carbon atoms (C₁₋₆), and further more preferably 1-4 carbon atoms (C₁₋₄) or 1-3 carbon atoms (C₁₋₃). When the term “alkyl” is prefixed with “C_{a-b}”, it means the number of carbon atoms in the alkyl, where a is the minimum number of carbons in the alkyl and b is the maximum number of carbons in the alkyl. For example, “C₁₋₆ alkyl” refers to an alkyl containing 1-6 carbon atoms. “C₁₋₃ alkyl” refers to an alkyl containing 1-3 carbon atoms. Examples of C₁₋₆ alkyl include, but are not limited to, methyl, ethyl, propyl (e.g. n-propyl, i-propyl), butyl (e.g. n-butyl, i-butyl, s-butyl and t-butyl), pentyl (e.g. n-pentyl, i-pentyl, neo-pentyl), hexyl, and the like. When used as a linker (e.g., in the definition of L) or between two dashes (“-”) (e.g., -(C₁₋₆ alkyl)-OH), the alkyl refers to an alkylene.

The term “alkenyl” as used herein refers to a straight or branched unsaturated hydrocarbon radical containing one or more, for example 1, 2, or 3 carbon-carbon double bonds (C=C) and 2-18 carbon atoms (C₂₋₁₈), preferably 2-10 carbon atoms (C₂₋₁₀), more preferably 2-6 carbon atoms (C₂₋₆), and further more preferably 2-4 carbon atoms (C₂₋₄). When the term “alkenyl” is prefixed with “C_{a-b}”, it means the number of carbon atoms in the alkenyl, where a is the minimum number of carbons in the alkenyl and b is the maximum number of carbons in the alkenyl. For example, “C₂₋₆ alkenyl” refers to an alkenyl containing 2-6 carbon atoms. “C₂₋₄ alkenyl” refers to an alkenyl containing 2-4 carbon atoms. Examples of C₂₋₆ alkenyl include, but are not limited to, vinyl, propenyl (e.g. 2-propenyl), and butenyl (e.g. 2-butenyl), and the like. The point of attachment for the alkenyl can be on or not on the double bonds. When used as a linker (e.g., in the definition of L), the alkenyl refers to an alkenylene.

The term “alkynyl” as used herein refers to a straight or branched unsaturated hydrocarbon radical containing one or more, for example 1, 2, or 3, carbon-carbon triple bonds (C≡C) and 2-18 carbon atoms (C₂₋₁₈), preferably 2-10 carbon atoms (C₂₋₁₀), more preferably 2-6 carbon atoms (C₂₋₆), and further more preferably 2-4 carbon atoms (C₂₋₄). When the term “alkynyl” is prefixed with “C_{a-b}”, it means the number of carbon atoms in the alkynyl, where a is the minimum number of carbons in the alkynyl and b is the maximum number of carbons in the alkynyl. For example, “C₂₋₆ alkynyl” refers to an alkynyl containing 2-6 carbon atoms. “C₂₋₄ alkynyl” refers to an alkynyl containing 2-4 carbon atoms. Examples of C₂₋₆ alkynyl include, but are not limited to, ethynyl, propynyl (e.g. 2-propynyl), and butynyl (e.g. 2-butynyl), and the like. The point of attachment for

the alkynyl can be on or not on the triple bonds. When used as a linker (e.g., in the definition of L), the alkynyl refers to an alkynylene.

The term “halogen” or “halo” as used herein means fluoro, chloro, bromo, and iodo, preferably fluoro, chloro and bromo, more preferably fluoro and chloro.

5 The term “haloalkyl” as used herein refers to an alkyl radical, as defined herein, in which one or more, for example 1, 2, 3, 4, or 5, or all hydrogen atoms are replaced with halogen atoms, and when more than one hydrogen atoms are replaced with halogen atoms, the halogen atoms may be the same or different from each other. In one embodiment, the term “haloalkyl” as used herein refers to an alkyl radical, as defined
10 herein, in which two or more, such as 2, 3, 4, or 5, or all hydrogen atoms are replaced with halogen atoms, wherein the halogen atoms are identical to each other. In another embodiment, the term “haloalkyl” as used herein refers to an alkyl radical, as defined herein, in which two or more hydrogen atoms, such as 2, 3, 4, or 5, or all hydrogen atoms are replaced with halogen atoms, wherein the halogen atoms are different from each other.
15 When the term “haloalkyl” is prefixed with “C_{a-b}”, it means the number of carbon atoms in the haloalkyl, where a is the minimum number of carbons in the haloalkyl and b is the maximum number of carbons in the haloalkyl. For example, “C₁₋₆ haloalkyl” refers to a haloalkyl as defined herein containing 1-6 carbon atoms. “C₁₋₄ haloalkyl” refers to a haloalkyl as defined herein containing 1-4 carbon atoms. Examples of C₁₋₆ haloalkyl
20 include, but are not limited to -CF₃, -CHF₂, -CH₂F, -CH₂CF₃, -CH(CF₃)₂, and the like.

 The term “cycloalkyl” as used herein refers to saturated or partially unsaturated cyclic hydrocarbon radical having 3-12 ring carbon atoms (C₃₋₁₂), such as 3-8 ring carbon atoms (C₃₋₈), 5-7 ring carbon atoms (C₅₋₇), 4-7 ring carbon atoms (C₄₋₇) or 3-6 ring carbon atoms (C₃₋₆), which may have one or more rings, such as 1, 2, or 3 rings, preferably 1 or 2
25 rings. When the term “cycloalkyl” is prefixed with “C_{a-b}”, it means the number of carbon atoms in the cycloalkyl, where a is the minimum number of carbons in the cycloalkyl and b is the maximum number of carbons in the cycloalkyl. For example, “C₃₋₈ cycloalkyl” or “3-8 membered cycloalkyl” refers to a cycloalkyl containing 3-8 ring carbon atoms; “C₃₋₆ cycloalkyl” or “3-6 membered cycloalkyl” refers to a cycloalkyl containing 3-6 ring
30 carbon atoms. The cycloalkyl may include a fused or bridged ring, or a spirocyclic ring. The rings of the cycloalkyl may be saturated or have one or more, for example, one or two double bonds (i.e. partially unsaturated), but not fully conjugated, and not an aryl as

defined herein. Examples of C₃₋₈ cycloalkyl include, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, spiro[2.2]pentyl, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclopentadienyl, cyclohexenyl, etc.

The term “heterocyclyl” or “heterocycle” as used herein can be used

- 5 interchangeably and each refers to saturated or partially unsaturated cyclic radicals having 3-12 ring atoms, such as 5-12 ring atoms (5-12 membered heterocyclyl), 3-8 ring atoms (3-8 membered heterocyclyl), 4-8 ring atoms (4-8 membered heterocyclyl), 4-6 ring atoms (4-6 membered heterocyclyl) or 4-5 ring atoms (4-5 membered heterocyclyl), and containing one or more, for example 1, 2 or 3, preferably 1 or 2 heteroatoms
- 10 independently chosen from N, O and S in the rings, with the remaining ring atoms being carbon; it may have one or more rings, for example 1, 2 or 3, preferably 1 or 2 rings. The heterocyclyl also includes those wherein the N or S heteroatom are optionally oxidized to various oxidation states. The point of attachment of heterocyclyl can be on the N heteroatom or carbon. For example, “4-8 membered heterocyclyl” represents a
- 15 heterocyclyl having 4-8 (4, 5, 6, 7 or 8) ring atoms comprising at least one, such as 1, 2 or 3, preferably 1 or 2 heteroatoms independently chosen from N, O and S; “4-6 membered heterocyclyl” represents a heterocyclyl having 4-6 (4, 5 or 6) ring atoms comprising at least one, preferably 1 or 2 heteroatoms independently chosen from N, O and S (preferably N and O), which is preferably a monocyclic ring; and “4-5 membered
- 20 heterocyclyl” represents a heterocyclyl having 4-5 ring atoms comprising at least one, preferably 1 or 2 heteroatoms independently chosen from N, O and S (preferably N and O), which is a monocyclic ring. The heterocyclyl also includes a fused or bridged ring, or a spirocyclic ring. The rings of the heterocyclyl may be saturated or have one or more, for example, one or two double bonds (i.e. partially unsaturated), but not fully conjugated,
- 25 and not a heteroaryl as defined herein. Examples of heterocyclyl include, but are not limited to: 3-8 membered heterocyclyl, 4-8 membered heterocyclyl, 4-6 membered heterocyclyl and 4-5 membered heterocyclyl, such as oxetanyl, azetidiny, pyrrolidyl, tetrahydrofuranyl, dioxolanyl, tetrahydropyranyl, morpholinyl, thiomorpholinyl, piperidyl, piperazinyl, tetrahydropyridyl, dihydropyrimidyl, dihydropyrazinyl,
- 30 pyrazolidinyl and oxaspiro[3.3]heptyl, preferably oxetanyl (such as oxetan-3-yl), azetidiny, tetrahydropyranyl, morpholinyl (such as morpholino), piperazinyl (such as

piperazin-1-yl), tetrahydropyridyl (such as 1,2,3,6-tetrahydropyridyl), dihydropyrimidyl (such as 1,6-dihydropyrimidyl).

The term “aryl” or “aromatic ring” as used herein can be used interchangeably and each refers to carbocyclic hydrocarbon radical of 6 to 14 carbon atoms consisting of one ring or more fused rings, wherein at least one ring is an aromatic ring. Examples of aryl include, but are not limited to phenyl, naphthalenyl, 1,2,3,4-tetrahydronaphthalenyl, phenanthryl, indenyl, indanyl, azulenyl, preferably phenyl and naphthalenyl.

The term “heteroaryl” or “heteroaromatic ring” as used herein can be used interchangeably and each refers to: mono-, bi-, or tri- ring system having 5-15 ring atoms, preferably 5-14 ring atoms, more preferably 5-12 ring atoms, further preferably 5-10 ring atoms, and most preferably 5-6 or 8-10 ring atoms, wherein at least one ring is 5- or 6-membered aromatic ring containing one or more, for example 1 to 4, heteroatoms independently chosen from N, O, and S, wherein S and N may be optionally oxidized to various oxidation states. When the total number of S and O atoms in the heteroaryl group exceeds 1, said S and O heteroatoms are not adjacent to one another. Preferably, the heteroaryl is 5-12 membered heteroaryl. For example, the heteroaryl includes:

a 5-6 membered monocyclic heteroaryl, i.e., a monocyclic ring aromatic hydrocarbyl having 5 or 6 ring atoms, wherein the ring atoms include one or more, such as 1, 2 or 3 heteroatoms independently chosen from N, O and S (preferably N), and the remaining ring atoms are carbon atoms; and the heteroaryl is preferably triazolyl, pyridyl, pyrazinyl, pyrimidyl, pyrazolyl, imidazolyl, isoxazolyl, triazinyl, oxazolyl, thiadiazolyl, and pyridazinyl, more preferably pyridyl (such as pyridin-4-yl, pyridin-3-yl), pyrazinyl, pyrimidyl, and triazinyl (such as 1,2,4-triazinyl), and

a 8-10 membered bicyclic heteroaryl, i.e., a bicycle aromatic hydrocarbyl having 8, 9 or 10 ring atoms, wherein the ring atoms include one or more, such as 1, 2, 3 or 4, preferably 1, 2 or 3 heteroatoms independently chosen from N, O and S (preferably N), and the remaining ring atoms are carbon atoms, wherein at least one ring is an aromatic ring; which is preferably imidazo[1,2-c]pyrimidyl, 1*H*-pyrrolo[2,3-b]pyridyl, indazolyl, imidazo[1,2-a]pyrazinyl, imidazo[1,5-a]pyrazinyl, pyrrolo[1,2-a]pyrazinyl, pyrazolo[1,5-a]pyrazinyl, [1,2,4]triazolo[1,5-a]pyrazinyl, [1,2,4]triazolo[4,3-c]pyrimidyl, [1,2,4]triazolo[1,5-c]pyrimidyl, and 1,2,3,4-tetrahydro-1,5-naphthyridinyl.

Examples of heteroaryl include, but are not limited to: 5-6 membered monocyclic heteroaryl, such as pyridyl, N-oxide pyridyl, pyrazinyl, pyrimidyl, triazinyl (such as 1,2,4-triazinyl, 1,3,5-triazinyl), pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, oxadiazolyl (such as 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl and 1,3,4-oxadiazolyl), thiazolyl, isothiazolyl, thiadiazolyl, tetrazolyl, triazolyl, thienyl, furanyl, pyranyl, pyrrolyl, and pyridazinyl; and a 8-10 membered bicyclic heteroaryl, such as benzoxazolyl, benzoisoxazolyl, benzothienyl, benzothiazolyl, benzoisothiazolyl, imidazopyrimidyl (such as imidazo[1,2-c]pyrimidyl), imidazopyrazinyl (such as imidazo[1,2-a]pyrazinyl and imidazo[1,5-a]pyrazinyl), imidazopyridyl (such as imidazo[1,2-a]pyridyl), imidazopyridazinyl (such as imidazo[1,2-b]pyridazinyl), pyrrolopyrazinyl (such as pyrrolo[1,2-a]pyrazinyl), pyrrolopyridyl (such as 1*H*-pyrrolo[2,3-b]pyridyl), pyrrolopyrimidyl (such as pyrrolo[3,4-d]pyrimidyl), pyrazolopyrazinyl (such as pyrazolo[1,5-a]pyrazinyl), pyrazolopyridyl (such as 1*H*-pyrazolo[3,4-b]pyridyl), pyrazolopyrimidyl (such as pyrazolo[1,5-a]pyrimidyl), triazolopyrimidyl (such as [1,2,4]triazolo[4,3-c]pyrimidyl and [1,2,4]triazolo[1,5-c]pyrimidyl), triazolopyrazinyl (such as [1,2,4]triazolo[1,5-a]pyrazinyl), triazolopyridyl (such as [1,2,4]triazolo[4,3-a]pyridyl and [1,2,4]triazolo[1,5-a]pyridyl), tetrazolopyridyl (such as tetrazolo[1,5-a]pyridyl), benzofuranyl, indolyl, indazolyl, purinyl, quinoliny, isoquinoliny, 6,7-dihydro-4*H*-pyrazolo[5,1-c][1,4]oxazinyl, and 1,2,3,4-tetrahydro-1,5-naphthyridinyl.

The term “-OH” as used herein refers to hydroxyl radical.

The term “-CN” as used herein refers to cyano radical.

The term “oxo” as used herein refers to =O.


The term “optional” or “optionally” as used herein means that the subsequently described event or circumstance may or may not occur, and the description includes instances wherein the event or circumstance occur and instances in which it does not occur. For example, “optionally substituted with one or more” includes unsubstituted and substituted with 1, 2, 3 or more substituents as described. It will be understood by those skilled in the art, with respect to any group containing one or more substituents, that such groups are not intended to introduce any substitution or substitution patterns that are sterically impractical, chemically incorrect, synthetically non-feasible and/or inherently unstable.

The term “substituted” or “substituted with...”, as used herein, means that one or more (such as, 1, 2, 3 or 4) hydrogens on the designated atom or group are replaced with one or more (such as 1, 2, 3 or 4) substituents, preferably the substituents chosen from the indicated group of substituents or radicals, provided that the designated atom’s normal
5 valence is not exceeded. The said substituents may be the same or different from each other. The term “substituted with one or more groups chosen from” or “substituted with one or more” as used herein means that one or more hydrogens on the designated atom or group are independently replaced with one or more radicals from the indicated group of substituents or radicals, wherein the said radicals may be the same or different from each
10 other. Preferably, “substituted with one or more groups chosen from” or “substituted with one or more” means that the designated atom or group is substituted with 1, 2, 3, or 4 radicals independently chosen from the indicated group of substituents or radicals, wherein the said radicals may be the same or different from each other. In some embodiments, when a substituent is oxo (i.e., =O), then 2 hydrogens on a single atom are
15 replaced by the oxo. An optional substituent can be any radicals, provided that combinations of substituents and/or variables result in a chemically correct and stable compound. A chemically correct and stable compound is meant to imply a compound that is sufficiently robust to survive sufficient isolation from a reaction mixture to be able to identify the chemical structure of the compound. Preferably, substituents are those
20 exemplified in the compounds of the examples of the present application.

Unless otherwise specified, substituents are named into the core structure. For example, it is to be understood that when (cycloalkyl)alkyl is listed as a possible substituent, the point of attachment of this substituent to the core structure is in the alkyl portion.

25 When a structural formula herein contains an asterisk “*”, it means that the chiral center (or chiral axis) at the “*” mark in the compound is a single configuration of (R) configuration or (S) configuration; wherein the content of the single-configuration compound marked with “*” is at least 90% (e.g., 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9%, 100%, or any value between these listed values). For
30 example, some compounds of the present invention are axially chiral compounds, such as the following compound of formula (a), and its structural formula contains an asterisk “*”,

which means that the compound is a compound of formula (b) or compound of formula (c) in a single configuration.



It will be appreciated by the person of ordinary skill in the art ("POSITA") that some of the compounds of formula (I) may contain one or more chiral centers (or chiral axes) and therefore exist in two or more stereoisomeric forms. The racemates of these isomers, the individual isomers and mixtures enriched in one enantiomer, as well as diastereomers when there are two chiral centers (or chiral axes), and mixtures partially enriched with specific diastereomers are within the scope of the present invention. It will be further appreciated by the POSITA that the present invention includes all the individual stereoisomers (e.g. enantiomers, diastereomers), racemic mixtures or partially resolved mixtures of the compounds of formula (I) and, where appropriate, the individual tautomeric forms thereof.

The term "axial chirality" as used herein is a special case of chirality. The molecule has a chiral axis, with multiple groups arranged around the axis, and their arrangement makes the molecule unable to superimpose with its mirror image. Axial chirality is most commonly found in asymmetric biaromatic ring (e.g. biphenyl) compounds with limited rotation, such as 1,1'-bi-(2-naphthol).

The term "stereoisomers" as used herein refers to compounds that have the same chemical constitution but differ in the arrangement of atoms or groups in space. Stereoisomers include enantiomers, diastereomers and the like.

The terms "enantiomers" and "enantiomeric forms" as used herein can be used interchangeably and refer to two stereoisomers of a compound that are non-superimposable mirror images of each other.

The terms "diastereomers" and "diastereomeric forms" as used herein can be used interchangeably and refer to stereoisomers that have two or more chiral centers (or chiral axes) and whose molecules are not mirror images of each other. Diastereomers have different physical properties, such as melting points, boiling points, spectral properties, or

biological activities. A mixture of diastereomers can be separated by high-resolution analytical methods such as electrophoresis and chromatography such as HPLC.

In some embodiments, the present invention provides compounds of various stereoisomeric purities, that is, enantiomeric or diastereomeric purity expressed in different “ee” or “de” values. In some embodiments, the compound of formula (I) described herein has an enantiomeric purity of at least 60% ee (e.g., 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9%, 100% ee, or any value between these listed values). In some embodiments, the compound of formula (I) described herein has an enantiomeric purity of greater than 99.9% ee. In some embodiments, the compound of formula (I) described herein has a diastereomeric purity of at least 60% de (e.g., 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9%, 100% de, or any value between these listed values). In some embodiments, the compound of formula (I) described herein has a diastereomeric purity of greater than 99.9% de.

The term “enantiomeric excess” or “ee” refers to the amount of one enantiomer relative to the other. For a mixture of R and S enantiomers, the percentage of enantiomeric excess is defined as $|R - S| * 100$, where R and S are the mole or weight fractions of the respective enantiomers in the mixture, $R + S = 1$. If the optical rotation of a chiral substance is known, the percentage of enantiomeric excess is defined as $([a]_{\text{obs}}/[a]_{\text{max}}) * 100$, wherein $[a]_{\text{obs}}$ is the optical rotation of the enantiomeric mixture, and $[a]_{\text{max}}$ is the optical rotation of the pure enantiomer.

The term “diastereomeric excess” or “de” refers to the amount of one diastereomer relative to the other, and is defined by analogy based on the enantiomeric excess. Therefore, for a mixture of diastereomers D1 and D2, the percentage of diastereomeric excess is defined as $|D1 - D2| * 100$, wherein D1 and D2 are the mole or weight fractions of the respective diastereomers in the mixture, $D1 + D2 = 1$.

The diastereomeric excess and enantiomeric excess can be measured by a number of analytical techniques (including nuclear magnetic resonance spectroscopy, chiral column chromatography and/or optical polarimetry) according to conventional protocols well known to a person skilled in the art.

The racemates can be used as such or can be resolved into their individual isomers. The resolution can afford stereochemically pure compounds or mixtures enriched in one

or more isomers. Methods for separation of isomers are well known (cf. Allinger N. L. and Eliel E. L. in *“Topics in Stereochemistry”*, Vol. 6, Wiley Interscience, 1971) and

include physical methods such as chromatography using a chiral adsorbent. Individual isomers can be prepared in chiral form from chiral precursors. Alternatively, individual isomers can be separated chemically from a mixture by: forming diastereomeric salts with a chiral acid (such as the individual enantiomers of 10-camphorsulfonic acid, camphoric acid, alpha-bromocamphoric acid, tartaric acid, diacetyltartaric acid, malic acid, pyrrolidone-5-carboxylic acid, and the like), fractionally crystallizing the salts, and then freeing one or both of the resolved bases, optionally repeating the process, so as to obtain either or both substantially free of the other; i.e., in a form having an optical purity of > 95%. Alternatively, the racemates can be covalently linked to a chiral compound (auxiliary) to produce diastereomers which can be separated by chromatography or by fractional crystallization after which time the chiral auxiliary is chemically removed to afford the pure enantiomers.

The term “tautomer” as used herein refers to constitutional isomers of compounds generated by rapid movement of an atom in two positions in a molecule. Tautomers readily interconvert into each other, e.g., enol form and ketone form are typical tautomers.

A “pharmaceutically acceptable salt” is intended to mean a salt of a free acid or base of a compound of Formula (I) that is non-toxic, biologically tolerable, or otherwise biologically suitable for administration to the subject to be treated or prevented. For example, an acid addition salt includes such as a salt derived from an inorganic acid and an organic acid. For examples, see, generally, S. M. Berge, et al., “Pharmaceutical Salts”, J. Pharm. Sci., 1977, 66:1-19, and Handbook of Pharmaceutical Salts, Properties, Selection, and Use, Stahl and Wermuth, Eds., Wiley-VCH and VHCA, Zurich, 2002.

In addition, if a compound described herein is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid addition salt. Conversely, if the product is a free base, an acid addition salt, particularly a pharmaceutically acceptable acid addition salt, may be produced by dissolving the free base in a suitable solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds. The POSITA will recognize various synthetic methodologies that may be used without undue experimentation to prepare non-toxic pharmaceutically acceptable acid addition salts or base addition salts.

The term “solvates” means solvent addition forms that contain either stoichiometric or non-stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the solid state, thus forming a solvate. If the solvent is water, the solvate formed is a hydrate, when the solvent is alcohol, the solvate
5 formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water, or less than one molecule of water, with one molecule of the substances in which the water retains its molecular state as H₂O, such combination being able to form one or more hydrates, for example, hemihydrate, monohydrate, and dihydrate.

10 As used herein, the terms “group(s)” and “radical(s)” are synonymous and are intended to indicate functional groups or fragments of molecules attachable to other fragments of molecules.

The term “active ingredient” is used to indicate a chemical substance which has biological activity. In some embodiments, an “active ingredient” is a chemical substance
15 having pharmaceutical utility.

The term “pharmaceutical combination” as used herein means a product obtained by mixing or combining two or more active ingredients, including fixed and non-fixed combinations of active ingredients, such as a kit, and a pharmaceutical composition. The term “fixed combination” means that two or more active ingredients (such as compounds
20 of the present invention and additional therapeutic agents) are administered simultaneously to a patient in the form of a single entity or dose. The term “non-fixed combination” means that two or more active ingredients (such as compounds of the present invention and additional therapeutic agents) are administered simultaneously, in parallel or successively to a patient in separate entities, wherein the administration
25 provides the patient with a therapeutically effective level of the compound.

The terms “treating” or “treatment” or “prevention” of a disease or disorder, in the context of achieving therapeutic benefit, refer to administering one or more pharmaceutical substances, especially compounds of the present invention to a subject that has the disease or disorder, or has a symptom of a disease or disorder, or has a
30 predisposition toward a disease or disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease or disorder, the symptoms of the disease or disorder, or the predisposition toward the disease or disorder.

In some embodiments, the disease or disorder is cancer, such as solid tumors or hematologic malignancies, including leukemia, lymphoma and myeloma. In another embodiment, the disease or disorder is Noonan Syndrome or LEOPARD Syndrome.

The terms “treating”, “contacting” and “reacting,” in the context of a chemical reaction, mean adding or mixing two or more reagents under appropriate conditions to produce the indicated and/or the desired product. It should be appreciated that the reaction which produces the indicated and/or the desired product may not necessarily result directly from the combination of two reagents which were initially added, i.e., there may be one or more intermediates which are produced in the mixture which ultimately lead to the formation of the indicated and/or the desired product.

The term “effective amount” as used herein refers to an amount or dose of a SHP2 inhibitor sufficient to generally bring about a therapeutic benefit in patients in need of treatment or prevention for a disease or disorder mediated by SHP2 or at least in part by SHP2. Effective amounts or doses of the active ingredient of the present disclosure may be ascertained by methods such as modeling, dose escalation studies or clinical trials, and by taking into consideration factors, e.g., the mode or route of administration or drug delivery, the pharmacokinetics of the agent, the severity and course of the disease or disorder, the subject’s previous or ongoing therapy, the subject’s health status and response to drugs, and the judgment of the attending physician.

An exemplary dose is in the range of from about 0.0001 to about 200 mg of active agent per kg of subject’s body weight per day, such as from about 0.001 to 100 mg/kg/day, or about 0.01 to 35 mg/kg/day, or about 0.1 to 10 mg/kg daily in single or divided dosage units (e.g., BID, TID, QID). For a 70-kg human, an illustrative range for a suitable dosage amount is from about 0.05 to about 7 g/day, or about 0.2 to about 5 g/day. Once improvement of the patient’s disease or disorder has occurred, the dose may be adjusted for maintenance treatment. For example, the dosage or the frequency of administration, or both, may be reduced as a function of the symptoms, to a level at which the desired therapeutic effect is maintained. Of course, if symptoms have been alleviated to an appropriate level, treatment may cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of symptoms.

The term “inhibition” or “inhibiting” indicates a decrease in the baseline activity of a biological activity or process. The term “inhibition of SHP2 activity” is a practical

pharmaceutical activity for purposes of this disclosure and refers to a decrease in the activity of SHP2 as a direct or indirect response to the presence of the compound of the present invention, relative to the activity of SHP2 in the absence of the compound of the present invention. The decrease in activity may be due to the direct interaction of the compound of the present invention with SHP2, or due to the interaction of the compound of the present invention, with one or more other factors that in turn affect the SHP2 activity. For example, the presence of the compound of the present invention may decrease the SHP2 activity by directly binding to the SHP2, by causing (directly or indirectly) another factor to decrease the SHP2 activity, or by (directly or indirectly) decreasing the amount of SHP2 present in the cell or organism.

The term “subject” or “patient” as used herein means mammals and non-mammals. Mammals means any member of the mammalia class including, but not limited to, humans; non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, and swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice, and guinea pigs; and the like. Examples of non-mammals include, but are not limited to, birds, and the like. The term “subject” or “patient” does not denote a particular age or sex. In some embodiments, the subject or patient is a human.


In general, the term “about” is used herein to modify a numerical value above or below the stated value by a variance of 20%.

Technical and scientific terms used herein and not specifically defined have the meaning commonly understood by the POSITA to which the present disclosure pertains.

All numerical ranges herein shall be interpreted as disclosing each numerical value and subset of numerical values within the range, regardless of whether they are specifically otherwise disclosed. For example, when referring to any range of values, it should be regarded as referring to every value within the range of values, for example, every integer within the range of values. For example, C₁₋₆ as used herein represents the inclusion of 1, 2, 3, 4, 5 or 6 C. The invention relates to all values falling within the ranges, all smaller ranges and the upper or lower limits of the numerical range.

Detailed Description of Embodiments

Embodiment 1. A compound of formula (I):



or a pharmaceutically acceptable salt thereof, or a solvate, a racemic mixture, an enantiomer, a diastereomer or a tautomer thereof, wherein

ring A is a benzene ring or a pyridine ring;

5 Z is CH₂, O, S or NH;

R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄, -SR₅ and -SR₆, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-CN, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and -(C₁₋₆ alkyl)-CN; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, phenyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl; R₆ is chosen from -CO(C₁₋₆ alkyl), -CO(C₃₋₈ cycloalkyl), -CO(4-8 membered heterocyclyl), -CONH₂, -CONH(C₁₋₆ alkyl), -CONH(C₃₋₈ cycloalkyl), -CONH(4-8 membered heterocyclyl), -CON(C₁₋₆ alkyl)₂, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-NH(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-N(C₁₋₆ alkyl)₂ and -(C₁₋₆ alkyl)-NHCO(C₁₋₆ alkyl), wherein the C₁₋₆ alkyl of R₆ is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH and -O(C₁₋₆ alkyl); and the above-mentioned C₃₋₈ cycloalkyl, phenyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: halogen, -CN, -CONH₂, -OH, oxo, -NH₂, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-CN, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -S(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl) and -N(C₁₋₆ alkyl)₂;

R₁' is independently chosen from halogen, -CN, -CONH₂, -OH, -NH₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-

O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-CN, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl) and -N(C₁₋₆ alkyl)₂, wherein the C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more halogen;


5 n is 0, 1, 2 or 3;

R₂ is chosen from -NH₂, -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, oxo and -OH;

Cy₁ is 5-12 membered heterocyclyl or 5-12 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -CONH₂, -OH, oxo, -NH₂, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -
 10 (C₁₋₆ alkyl)-CN, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCO(C₁₋₆ alkyl), -CONH(C₁₋₆ alkyl) and -CON(C₁₋₆ alkyl)₂, wherein the C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more halogen;

15 Cy₂ is phenyl or 5-14 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -CONH₂, -OH, oxo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-CN, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -NR₇R₈, -NHCO(C₁₋₆ alkyl), -CONH(C₁₋₆ alkyl)
 20 and -CON(C₁₋₆ alkyl)₂, wherein R₇ and R₈ are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-CN, C₃₋₈ cycloalkyl, phenyl, 3-8 membered heterocyclyl and 5-6 membered heteroaryl; and
 L is absent, or L is S, O, NH, C₁₋₆ alkyl, C₂₋₆ alkenyl or C₂₋₆ alkynyl.

Embodiment 2. The compound or the pharmaceutically acceptable salt
 25 thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1, wherein the compound is a compound of formula (IA):



Embodiment 3. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1 or 2, wherein Z is CH₂ or O; and preferably, Z is CH₂.

5 **Embodiment 4.** The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of embodiments 1-3, wherein R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄ and -SR₅, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-
10 8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, C₃₋₈ cycloalkyl
15 and 4-8 membered heterocyclyl; R₃ is independently chosen from hydrogen and C₁₋₆ alkyl; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl; and the above-mentioned C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: halogen, -CN, -CONH₂, -OH, oxo,
20 -NH₂, C₁₋₆ alkyl and -O(C₁₋₆ alkyl).

Embodiment 5. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 4, wherein R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -
25 OH, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl, wherein the C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆
30 alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -CONH(C₁₋₆ alkyl), -CONH(C₁₋₆ alkyl)-O-

(C₁₋₆ alkyl), -CON(C₁₋₆ alkyl)₂, -CON(C₁₋₆ alkyl)(C₁₋₆ alkyl-O-C₁₋₆ alkyl), -CONH(C₃₋₈ cycloalkyl), -COOH, -COO(C₁₋₆ alkyl), -CO(C₁₋₆ alkyl), -CO(4-8 membered heterocyclyl) and -CO(4-8 membered heterocyclyl)-O-(C₁₋₆ alkyl); more preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂, -NHCONH₂, -CONH(C₁₋₃ alkyl), -CONH(CH₂CH₂)-O-(CH₃), -CON(CH₃)₂, -CON(CH₃)(CH₂CH₂-O-CH₃), -CONH(cyclopropyl), -COOH, -COO(CH₃), -CO(CH₃), -CO(azetidiny) or -CO(azetidiny)-O-(CH₃); and most preferably, R₁ is ethynyl, or ethynyl substituted with -CONH(C₁₋₃ alkyl).

Embodiment 6. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 4, wherein R₁ is chosen from -NR₃R₄ and -SR₅, wherein the R₃ is independently chosen from hydrogen and C₁₋₆ alkyl; and R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl, wherein the C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl.

Embodiment 7. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of embodiments 1-6, wherein R₁' is chosen from halogen, -CN, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0.


Embodiment 8. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of embodiments 1-7, wherein R₂ is chosen from -NH₂ and oxo; and preferably, R₂ is -NH₂.

Embodiment 9. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of embodiments 1-8, wherein Cy₁ is 5-12 membered heterocyclyl or 5-12 membered heteroaryl, preferably 5-10 membered heterocyclyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heterocyclyl or 5-9 membered heteroaryl, which is optionally substituted with one or

more groups independently chosen from: oxo, -NH₂, -CN, halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl and -(C₁₋₆ alkyl)-OH.


Embodiment 10. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 9, wherein Cy₁ is chosen from 1,6-dihydropyrimidyl, pyrazinyl, pyrimidyl, 1,2,4-triazinyl, imidazopyrimidyl, triazolopyrimidyl, imidazopyrazinyl, pyrrolopyrazinyl, pyrazolopyrazinyl and triazolopyrazinyl, each of which is optionally substituted with one or more groups independently chosen from: oxo, -NH₂, -CN, halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl and -(C₁₋₆ alkyl)-OH.

Embodiment 11. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 10, wherein Cy₁ is chosen from




, each of


which is optionally substituted with one or more groups independently chosen from: -NH₂, -CN, halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl and -(C₁₋₆ alkyl)-OH; preferably, Cy₁ is





each of which is optionally substituted with one or more groups independently chosen from: -NH₂, -CN, halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl and -(C₁₋₆ alkyl)-OH;




each of which is optionally substituted with one or more groups independently chosen from: -NH₂, C₁₋₆ alkyl and -(C₁₋₆ alkyl)-OH;

further preferably, Cy₁ is , which is optionally substituted with one or


more groups independently chosen from: -NH₂ and C₁₋₆ alkyl; or Cy₁ is , which is optionally substituted with one or more groups independently chosen from: C₁₋₆

alkyl; or Cy₁ is , which is optionally substituted with one or more groups

5 independently chosen from: -NH₂, C₁₋₆ alkyl and -(C₁₋₆ alkyl)-OH; or Cy₁ is , which is optionally substituted with one or more groups independently chosen from: -NH₂ and C₁₋₆ alkyl.

Embodiment 12. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the
10 tautomer thereof according to any one of embodiments 1-11, wherein Cy₂ is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably phenyl, 5-6 membered heteroaryl or 9-10 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇
15 and R₈ are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-CN and C₃₋₈ cycloalkyl.

Embodiment 13. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1, wherein the compound is a compound of
20 formula (II):



wherein

Z is CH₂ or O; and preferably, Z is CH₂;

R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄ and -SR₅, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently
 10 chosen from hydrogen and C₁₋₆ alkyl; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl; and the above-mentioned C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is
 15 optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -CONH(C₁₋₆ alkyl) and -CON(C₁₋₆ alkyl)₂; further preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂, -CONH(CH₃) or -CON(CH₃)₂; and more preferably, R₁ is ethynyl;

20 R₁' is chosen from halogen, -CN, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0;


R₂ is -NH₂;

R₉ and R₁₀ are each independently chosen from hydrogen, -NH₂, halogen, C₁₋₆ alkyl and C₁₋₆ haloalkyl; and preferably, R₉ and R₁₀ are each independently chosen from
 25 hydrogen and C₁₋₆ alkyl;

Cy₂ is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably phenyl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently
 30 chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and

L is absent.

Embodiment 14. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1, wherein the compound is a compound of formula (III):



wherein

Z is CH₂;

R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄ and -SR₅, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently chosen from hydrogen and C₁₋₆ alkyl; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl; and the above-mentioned C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -CONH(C₁₋₆ alkyl) and -CON(C₁₋₆ alkyl)₂; further preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂ or -CONH(CH₃); and more preferably, R₁ is ethynyl;

R₁' is chosen from halogen, -CN, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0;

R_2 is chosen from $-NH_2$ and oxo; and preferably, R_2 is $-NH_2$;

R_{11} , R_{12} and R_{13} are each independently chosen from hydrogen, $-NH_2$, $-CN$, C_{1-6} alkyl and C_{1-6} haloalkyl; preferably, R_{11} , R_{12} and R_{13} are each independently chosen from hydrogen and C_{1-6} alkyl; and more preferably, R_{11} , R_{12} and R_{13} are all hydrogen;


5 Cy_2 is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6}$ alkyl), $-S(C_{1-6}$ alkyl) and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, $-(C_{1-6}$ alkyl)- OH , $-(C_{1-6}$ alkyl)- $O-(C_{1-6}$ alkyl)

10 and C_{3-8} cycloalkyl; and

L is absent, or L is S ; and preferably, L is S .

Embodiment 15. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1, wherein the compound is a compound of

15 formula (IV):



wherein

Z is CH_2 or O ; and preferably, Z is CH_2 ;

R_1 is C_{2-6} alkynyl, wherein the C_{2-6} alkynyl is optionally substituted with one or

20 more groups independently chosen from: halogen, $-CN$, $-OH$, $-NH_2$, C_{3-8} cycloalkyl, 4-8 membered heterocyclyl, $-O(C_{1-6}$ alkyl), $-O(C_{1-6}$ haloalkyl), $-O(C_{3-8}$ cycloalkyl), $-O(4-8$ membered heterocyclyl), $-S(C_{1-6}$ alkyl), $-S(C_{3-8}$ cycloalkyl), $-S(4-8$ membered heterocyclyl), $-NH(C_{1-6}$ alkyl), $-N(C_{1-6}$ alkyl) $_2$, $-NH-CN$, $-NHCONH_2$, $-NHCO(C_{1-6}$ alkyl), $-CONR_aR_b$, $-COOR_c$ and $-COR_d$, wherein R_a , R_b , R_c and R_d are each independently

25 chosen from hydrogen, C_{1-6} alkyl, $-(C_{1-6}$ alkyl)- $O-(C_{1-6}$ alkyl), $-(C_{1-6}$ alkyl)- OH , C_{3-8} cycloalkyl and 4-8 membered heterocyclyl; the above-mentioned C_{3-8} cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups

independently chosen from: C_{1-6} alkyl and $-O(C_{1-6} \text{ alkyl})$; preferably, R_1 is C_{2-6} alkynyl, wherein the C_{2-6} alkynyl is optionally substituted with one or more groups independently chosen from: $-OH$, $-O(C_{1-6} \text{ alkyl})$, $-NH(C_{1-6} \text{ alkyl})$, $-N(C_{1-6} \text{ alkyl})_2$, $-CONH(C_{1-6} \text{ alkyl})$ and $-CON(C_{1-6} \text{ alkyl})_2$; further preferably, R_1 is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with $-OH$, $-OCH_3$, $-NH(CH_3)$ or $-N(CH_3)_2$; and more preferably, R_1 is ethynyl;

R_1' is chosen from halogen, $-O(C_{1-6} \text{ alkyl})$ and $-S(C_{1-6} \text{ alkyl})$, and n is 0 or 1; preferably, R_1' is halogen, and n is 0 or 1; and more preferably, n is 0;

R_2 is $-NH_2$;


R_{14} is chosen from hydrogen, $-NH_2$ and C_{1-6} alkyl; preferably, R_{14} is hydrogen or $-NH_2$; and more preferably, R_{14} is hydrogen;

R_{14}' is C_{1-6} alkyl;

Cy_2 is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heteroaryl or 9-10 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6} \text{ alkyl})$, $-S(C_{1-6} \text{ alkyl})$ and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, $-(C_{1-6} \text{ alkyl})-OH$ and $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$; and

L is absent, or L is S ; and preferably, L is S .

Embodiment 16. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1, wherein the compound is a compound of formula (V):



wherein

Z is CH_2 or O ; and preferably, Z is CH_2 ;

R_1 is C_{2-6} alkynyl, wherein the C_{2-6} alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C_{3-8} cycloalkyl, 4-8 membered heterocyclyl, -O(C_{1-6} alkyl), -O(C_{1-6} haloalkyl), -O(C_{3-8} cycloalkyl), -O(4-8 membered heterocyclyl), -S(C_{1-6} alkyl), -S(C_{3-8} cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C_{1-6} alkyl), -N(C_{1-6} alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C_{1-6} alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, -(C_{1-6} alkyl)-O-(C_{1-6} alkyl), -(C_{1-6} alkyl)-OH, -(C_{1-6} alkyl)-CN, C_{3-8} cycloalkyl and 4-8 membered heterocyclyl; and the above-mentioned C_{3-8} cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C_{1-6} alkyl and -O(C_{1-6} alkyl); preferably, R_1 is C_{2-6} alkynyl, wherein the C_{2-6} alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -O(C_{1-6} alkyl), -NH(C_{1-6} alkyl), -N(C_{1-6} alkyl)₂, -NHCONH₂, -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C_{1-6} alkyl, -(C_{1-6} alkyl)-O-(C_{1-6} alkyl), C_{3-8} cycloalkyl and 4-8 membered heterocyclyl, wherein the C_{3-8} cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C_{1-6} alkyl and -O(C_{1-6} alkyl); more preferably, R_1 is C_{2-6} alkynyl, wherein the C_{2-6} alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C_{1-6} alkyl), -NH(C_{1-6} alkyl), -N(C_{1-6} alkyl)₂, -NHCONH₂, -CONH(C_{1-6} alkyl), -CONH(C_{1-6} alkyl)-O-(C_{1-6} alkyl), -CON(C_{1-6} alkyl)₂, -CON(C_{1-6} alkyl)(C_{1-6} alkyl-O- C_{1-6} alkyl), -CONH(C_{3-8} cycloalkyl), -COOH, -COO(C_{1-6} alkyl), -CO(C_{1-6} alkyl), -CO(4-8 membered heterocyclyl) and -CO(4-8 membered heterocyclyl)-O-(C_{1-6} alkyl); further preferably, R_1 is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂, -NHCONH₂, -CONH(C_{1-3} alkyl), -CONH(CH₂CH₂)-O-(CH₃), -CON(CH₃)₂, -CON(CH₃)(CH₂CH₂-O-CH₃), -CONH(cyclopropyl), -COOH, -COO(CH₃), -CO(CH₃), -CO(azetidiny) or -CO(azetidiny)-O-(CH₃); and most preferably, R_1 is ethynyl substituted with -CONH(CH₃), -CONH(CH₂CH₃) or -CONH(CH₂CH₂)-O-(CH₃);

R_1' is chosen from halogen, -O(C_{1-6} alkyl) and -S(C_{1-6} alkyl), and n is 0 or 1;

preferably, R_1' is halogen, and n is 0 or 1; and more preferably, n is 0;


R_2 is -NH₂;

R_{15} and R_{15}' are each independently chosen from hydrogen, $-NH_2$, $-CN$, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl and $-(C_{1-6} \text{ alkyl})-OH$; preferably, R_{15} and R_{15}' are each independently chosen from hydrogen, $-NH_2$, C_{1-6} alkyl and $-(C_{1-6} \text{ alkyl})-OH$; and more preferably, both R_{15} and R_{15}' are hydrogen;


- 5 Cy_2 is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heteroaryl or 9-10 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6} \text{ alkyl})$, $-S(C_{1-6} \text{ alkyl})$ and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, $-(C_{1-6} \text{ alkyl})-$
 10 OH , $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$ and C_{3-8} cycloalkyl; and
 L is absent, or L is S; and preferably, L is S.

Embodiment 17. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of embodiments 1-16, wherein Cy_2 is phenyl,
 15 pyridyl, pyrimidyl, indazolyl, pyrrolopyridyl or 1,2,3,4-tetrahydro-1,5-naphthyridinyl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6} \text{ alkyl})$, $-S(C_{1-6} \text{ alkyl})$ and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, $-(C_{1-6} \text{ alkyl})-$
 OH , $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$ and C_{3-8} cycloalkyl.


20 **Embodiment 18.** The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 17, wherein




each of which is optionally substituted with one or more groups independently
 25 chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6} \text{ alkyl})$, $-S(C_{1-6} \text{ alkyl})$ and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, $-(C_{1-6} \text{ alkyl})-$
 OH , $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$ and C_{3-8} cycloalkyl;
 preferably, Cy_2 is chosen from





each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and

5 more preferably, Cy₂ is chosen from , each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl.











10 **Embodiment 19.** The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 18, wherein











Cy₂ is , which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl);
or













Cy₂ is , which is optionally substituted with one or more groups independently chosen from: halogen, C₁₋₆ alkyl, -O(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and preferably, both R₇ and R₈ are hydrogen;
or















Cy₂ is , which is optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH and -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl);
25 and preferably, both R₇ and R₈ are hydrogen.

Embodiment 20. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to embodiment 1, which is chosen from:

No.	Structural formula	No.	Structural formula
1		2	
3		4	
5		6	
7		8	
9		10	













11 and 28	 +	12 and 64	 +
13		14	
15		16	
17		18	
19		21	











20 and 24		22	
		23	
25		26	
27		29	
30		31	
32		33	

34		35	
36		37	
38		39	
40		41	
42		43	
44		45	
46		47	

48 and 49	 +	50 and 51	
52		53	
54		55	
56 and 57	 +	58 and 59	 +
60		61	













62		63	
65 and 66		67	
		68	
69		70	
71		72	
73		74	
75		76	

77		78	
79		80	
81		82	
83		84	
85		86	
87		90	















88 and 89		91	
93		94	
95		96	
97		98	
99		100	














101		104	
102 and 103		106	
107		112	
108 and 109		110 and 111	
113 and 114		115	
		116	

117		120	
118 and 119		121	
		122	
124 and 125		123	
		126	
127		128	
129		130	












131		132	
133		134	
135		136	
137		138	
139		140	
141		142	















143		144	
145		146	
147		148	
149		150	
151		152	
153		154	
155		156	

157		158	
159		160	
161 and 162	 + 	163 and 164	 + 
165		166	
167		168	
169		170	

171		172	
173		174	
175		176	
177		178	
179		180	
181 and 182		183	
		184	
















185		186	
187		188	
189		194	
190 and 191		192 and 193	
195		196	
197		198	

















199		200 and 201	
202			
203		204	
205		206	
207 and 208		209 and 210	
211		212	

213		214	
215		216	
217 and 218		220 and 221	
219		222	
223		224	
225		226	
227		228	

229		230	
231		232	
233		234	
235		236	
237		238	
239		240	
241		242	
243		244	

245		246	
247		248	
249		250	
251		252	
253		254	
255		256	
257		258	
259		260	

261		262	
263		264	
265		266	
267 and 268		269	
		270	
271		272	
273		274	
275		276	

277		278	
279		280	
281		282	
283		284	
285		286	
287		288	
289		290	
291		292	

293		294	
295		296	
297		298	
299		300	
301		302	
303		304	
305			

Embodiment 21. A pharmaceutical composition, comprising the compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20, and optionally comprising a pharmaceutically acceptable excipient.

Embodiment 22. A method of *in vivo* or *in vitro* inhibiting the activity of SHP2, comprising contacting SHP2 with an effective amount of the compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20.

Embodiment 23. Use of the compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20 in the manufacture of a medicament for treating or preventing a disease mediated by SHP2 or at least in part by SHP2, and preferably for treating or preventing cancer, Noonan Syndrome and LEOPARD Syndrome, wherein the cancer is preferably a solid tumor or hematologic malignancy, including leukemia, lymphoma and myeloma; and the cancer is more preferably chosen from breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma, acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia, chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, and myeloma (such as multiple myeloma).

Embodiment 24. A method of treating or preventing a disease in a subject, comprising administering to the subject in need thereof an effective amount of the compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20, wherein the disease is a disease mediated by SHP2 or at least in part by SHP2, and the disease is preferably cancer, Noonan Syndrome and LEOPARD Syndrome, wherein the cancer is preferably a solid tumor or hematologic malignancy, including leukemia, lymphoma and myeloma; and the cancer is more preferably chosen from breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma, acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia,

chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, and myeloma (such as multiple myeloma).

Embodiment 25. The compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20, for use as a medicament.

Embodiment 26. The compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20, for use in treating or preventing a disease mediated by SHP2 or at least in part by SHP2, and preferably for use in treating or preventing cancer, Noonan Syndrome and LEOPARD Syndrome, wherein the cancer is preferably a solid tumor or hematologic malignancy, including leukemia, lymphoma and myeloma; and the cancer is more preferably chosen from breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma, acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia, chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, and myeloma (such as multiple myeloma).


Embodiment 27. A pharmaceutical combination, comprising the compound and/or the pharmaceutically acceptable salt thereof according to any one of embodiments 1-20, and at least one additional therapeutic agent, wherein the additional therapeutic agent is preferably chosen from: an anti-neoplastic active agent, an anti-inflammatory agent or an immunomodulator, wherein the anti-neoplastic active agent includes a chemotherapeutic agent, an immune checkpoint inhibitor or agonist, and a targeted therapeutic agent.

The various embodiments of the present invention (including the following examples) and the features of the various embodiments should be interpreted as being arbitrarily combined with each other, and the various solutions obtained from these mutual combinations are all included in the scope of the present invention, just like the solutions obtained from the mutual combinations specifically and individually set forth herein, unless clearly stated otherwise in the context.

General synthetic methods

The compound of formula (I) and/or a pharmaceutically acceptable salt thereof described herein can be synthesized using commercially available starting materials, by methods known in the art, or methods disclosed in the present patent application. The synthetic routes shown in Scheme 1 to Scheme 4 illustrate the general synthetic methods of the compounds of the present invention.

Scheme 1:




Ring A, Z, R_1 , R_1' , n, L and Cy_2 thereof are defined as for formula (I); Pg is an amino protecting group; and R_9 and R_{10} are each independently chosen from hydrogen, - NH_2 , halogen, C_{1-6} alkyl and C_{1-6} haloalkyl.

As shown in Scheme 1, a compound of formula II-A is reacted with acetonitrile under the catalysis of hydrogen chloride to obtain a compound of formula II-B. The compound of formula II-B is reacted with a corresponding malonate under alkaline conditions (sodium ethoxide/ethanol) to obtain a compound of formula II-C. The compound of formula II-C is reacted with 2,4,6-triisopropyl benzenesulfonyl chloride to obtain a compound of formula II-D. The compound of formula II-D is reacted with a compound of formula II-E under alkaline conditions (Et_3N or DIEA) to obtain a

compound of formula II-F. The compound of formula II-F is deprotected with an acid to obtain a compound of formula II-G.

Scheme 2:



- 5 Ring A, Z, R_1 , R_1' , n, L and Cy_2 thereof are defined as for formula (I); Pg is an amino protecting group; and R_{11} , R_{12} and R_{13} are each independently chosen from hydrogen, $-NH_2$, $-CN$, C_{1-6} alkyl and C_{1-6} haloalkyl.


As shown in Scheme 2, a compound of formula III-A and a compound of formula III-B are subjected to a nucleophilic substitution reaction under alkaline conditions (Et_3N or DIEA) to obtain a compound of formula III-C. The compound of formula III-C and a sulfur-containing sodium salt or boronic ester are subjected to a coupling reaction under the catalysis of palladium to obtain a compound of formula III-D. Palladium-catalyzed coupling reaction is carried out under suitable conditions. The base used can be chosen from Cs_2CO_3 , K_2CO_3 , DIEA, etc., and the catalyst used can be chosen from $Pd_2(dba)_3$, $Pd(PPh_3)_4$, $Pd(dppf)Cl_2 \cdot CH_2Cl_2$, etc. The compound of formula III-D is deprotected with an acid to obtain a compound of formula III-E.

10

15

Scheme 3:


57



Ring A, Z, R_1 , R_1' , n, L and Cy_2 thereof are defined as for formula (I); Pg is an amino protecting group; R_{14} is chosen from hydrogen, $-NH_2$ and C_{1-6} alkyl; and R_{14}' is C_{1-6} alkyl.

- 5 As shown in Scheme 3, a compound of formula IV-A is reacted with a compound of formula IV-B under BOP and DBU conditions to obtain a compound of formula IV-C. The compound of formula IV-C is reacted with NIS to obtain a compound of formula IV-D. The compound of formula IV-D and a sulfur-containing sodium salt or boronic ester are subjected to a coupling reaction under the catalysis of palladium to obtain a
- 10 compound of formula IV-E. Palladium-catalyzed coupling reaction is carried out under suitable conditions. The base used can be chosen from Cs_2CO_3 , K_2CO_3 , DIEA, etc., and the catalyst used can be chosen from $Pd_2(dba)_3$, $Pd(PPh_3)_4$, $Pd(dppf)Cl_2 \cdot CH_2Cl_2$, etc. The compound of formula IV-E is deprotected with an acid to obtain a compound of formula IV-F.

15 **Scheme 4:**



Ring A, R_1 , R_1' , n, L and Cy_2 thereof are defined as for formula (I); Pg is an amino protecting group; and R_{15} is chosen from hydrogen, $-NH_2$, $-CN$, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl and $-(C_{1-6} \text{ alkyl})-OH$.

- 5 As shown in Scheme 4, a compound of formula V-A is reacted with a compound of formula V-B under alkaline conditions (Et_3N or DIEA) to obtain a compound of formula V-C. The compound of formula IV-C is reacted with DIBAL-H to obtain a compound of formula V-D. The compound of formula V-D and a sulfur-containing sodium salt or boronic ester are subjected to a coupling reaction under the catalysis of palladium and
- 10 deprotected with an acid to obtain a compound of formula V-E. Palladium-catalyzed coupling reaction is carried out under suitable conditions. The base used can be chosen from CS_2CO_3 , K_2CO_3 , DIEA, etc., and the catalyst used can be chosen from $Pd_2(dba)_3$, $Pd(PPh_3)_4$, $Pd(dppf)Cl_2 \cdot CH_2Cl_2$, etc.

The substituents of the compounds thus obtained can be further modified to provide

15 other desired compounds. Synthetic chemistry transformations are described, for example, in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995) and subsequent editions thereof.

- 20 Before use, the compound(s) of the present invention can be purified by column chromatography, high performance liquid chromatography, crystallization or other suitable methods.

Pharmaceutical Compositions and Utility

The compound of the present invention (e.g., a compound of any of the examples as described herein) is used, alone or in combination with one or more additional therapeutic agents, to formulate pharmaceutical compositions. A pharmaceutical composition comprises: (a) an effective amount of the compounds of the present invention; (b) a pharmaceutically acceptable excipient (e.g., one or more pharmaceutically acceptable carriers); and optionally (c) at least one additional therapeutic agent.

A pharmaceutically acceptable excipient refers to an excipient that is compatible with active ingredients of the composition (and in some embodiments, capable of stabilizing the active ingredients) and not deleterious to the subject to be treated. For example, solubilizing agents, such as cyclodextrins (which form specific, more soluble complexes with the compounds of the present invention), can be utilized as pharmaceutical excipients for delivery of the active ingredients. Examples of other excipients include colloidal silicon dioxide, magnesium stearate, cellulose, sodium lauryl sulfate, and pigments such as D&C Yellow # 10. Suitable pharmaceutically acceptable excipients are disclosed in Remington's Pharmaceutical Sciences, A. Osol, a standard reference text in the art.

A pharmaceutical composition comprising a compound of the present invention can be administered in various known manners, such as orally, topically, rectally, parenterally, by inhalation spray, or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques.

A pharmaceutical composition described herein can be prepared in the form of tablet, capsule, sachet, dragee, powder, granule, lozenge, powder for reconstitution, liquid preparation, or suppository. In some embodiments, a pharmaceutical composition comprising a compound of the present invention is formulated for intravenous infusion, topical administration, or oral administration.

An oral composition can be any orally acceptable dosage form including, but not limited to, tablets, capsules, emulsions, and aqueous suspensions, dispersions and

solutions. Commonly used carriers for tablets include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added to tablets. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions or emulsions are administered orally, the active ingredient
5 can be suspended or dissolved in an oily phase combined with emulsifying or suspending agents. If desired, certain sweetening, flavoring, or coloring agents can be added.

In some embodiments, the compound of the present invention can be present in an amount of 1, 5, 10, 15, 20, 25, 50, 75, 80, 85, 90, 95, 100, 125, 150, 200, 250, 300, 400 and 500 mg in a tablet. In some embodiments, the compound of the present invention can
10 be present in an amount of 1, 5, 10, 15, 20, 25, 50, 75, 80, 85, 90, 95, 100, 125, 150, 200, 250, 300, 400 and 500 mg in a capsule.

A sterile injectable composition (e.g., aqueous or oleaginous suspension) can be formulated according to techniques known in the art using suitable dispersing or wetting agents (for example, Tween 80) and suspending agents. The sterile injectable
15 composition can also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the pharmaceutically acceptable vehicles and solvents that can be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium (e.g.,
20 synthetic mono- or di-glycerides). Fatty acids, such as oleic acid and its glyceride derivatives, and natural pharmaceutically acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions, can be used as sterile injectable medium. These oil solutions or suspensions can also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents.

25 An inhalation composition can be prepared according to techniques well known in the art of pharmaceutical formulation and can be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

30 A topical composition can be formulated in form of oil, cream, lotion, ointment, and the like. Suitable carriers for the composition include vegetable or mineral oils, white petrolatum (white soft paraffin), branched chain fats or oils, animal fats and high

molecular weight alcohols (greater than C12). In some embodiments, the pharmaceutically acceptable carrier is one in which the active ingredient is soluble. Emulsifiers, stabilizers, humectants and antioxidants may also be included as well as agents imparting color or fragrance, if desired. Additionally, transdermal penetration enhancers may be employed in those topical formulations. Examples of such enhancers can be found in U.S. Patent Nos. 3,989,816 and 4,444,762.

Creams may be formulated from a mixture of mineral oil, self-emulsifying beeswax and water in which mixture the active ingredient, dissolved in a small amount of an oil, such as almond oil, is admixed. An example of such a cream is one which includes, by weight, about 40 parts water, about 20 parts beeswax, about 40 parts mineral oil and about 1 part almond oil. Ointments may be formulated by mixing a solution of the active ingredient in a vegetable oil, such as almond oil, with warm soft paraffin and allowing the mixture to cool. An example of such an ointment is one which includes about 30% by weight almond oil and about 70% by weight white soft paraffin.

Suitable *in vitro* assays can be used to evaluate the effect of the compounds of the present invention in inhibiting the activity of SHP2. The compounds of the present invention can further be examined for additional effects in preventing or treating cancer by *in vivo* assays. For example, the compound of the present invention can be administered to an animal (e.g., a mouse model) having cancer and its therapeutic effects can be accessed. If the pre-clinical results are successful, the dosage range and administration route for animals, such as humans, can be projected.

The compound of the present invention can be shown to have sufficient pre-clinical practical utility to merit clinical trials hoped to demonstrate a beneficial therapeutic or prophylactic effect, for example, in subjects with cancer.

As used herein, the term “cancer” refers to a cellular disorder characterized by uncontrolled or dysregulated cell proliferation, decreased cellular differentiation, inappropriate ability to invade surrounding tissue, and/or ability to establish new growth at ectopic sites. The term “cancer” includes, but is not limited to, solid tumors and hematologic malignancies, such as leukemia, lymphoma or myeloma. The term “cancer” encompasses diseases of skin, tissues, organs, bone, cartilage, blood, and vessels. The term “cancer” further encompasses primary cancer, and metastatic cancer, recurrent cancer and refractory cancer.

Non-limiting examples of solid tumors include pancreatic cancer; bladder cancer; colorectal cancer; colon cancer; breast cancer, including metastatic breast cancer; prostate cancer, including androgen-dependent and androgen-independent prostate cancer; testicular cancer; renal cancer, including, e.g., metastatic renal cell carcinoma; urothelial carcinoma; liver cancer; hepatocellular cancer; lung cancer, including, e.g., non-small cell lung cancer (NSCLC), bronchioloalveolar carcinoma (BAC), and adenocarcinoma of the lung; ovarian cancer, including, e.g., progressive epithelial or primary peritoneal cancer; cervical cancer; endometrial cancer; gastric cancer; esophageal cancer; head and neck cancer, including, e.g., squamous cell carcinoma of the head and neck; skin cancer, including, e.g., melanoma and basal carcinoma; neuroendocrine cancer, including metastatic neuroendocrine tumors; brain tumors, including, e.g., glioma, anaplastic oligodendroglioma, adult glioblastoma multiforme, and adult anaplastic astrocytoma; bone cancer; sarcoma, including, e.g., Kaposi's sarcoma; adrenal carcinoma; mesothelioma; mesothelial carcinoma; choriocarcinoma; muscle carcinoma; connective tissue carcinoma; and thyroid carcinoma.

Non-limiting examples of hematologic malignancies include acute myelogenous leukemia (AML); juvenile acute myelogenous leukemia; chronic myelogenous leukemia (CML), including accelerated phase CML and CML blastic phase (CML-BP); acute lymphocytic leukemia (ALL); B-cell acute lymphocytic leukemia (B-ALL); chronic lymphocytic leukemia (CLL), including high risk CLL; human acute monocytic leukemia (M(5)); hairy cell leukemia; lymphocytic leukemia; chronic lymphoid leukemia; myelogenous leukemia; acute lymphoblastic leukemia; small lymphotic lymphoma (SLL); lymphoblastic lymphoma; Hodgkin's lymphoma; non-Hodgkin's lymphoma (NHL); mantle cell lymphoma (MCL); B-cell lymphoma; T-cell lymphoma; diffuse large B-cell lymphoma (DLBCL); large B-cell lymphoma (LBCL); follicular lymphoma; marginal zone lymphoma; Burkitt's lymphoma; non-Burkitt's highly degree B cell malignant lymphoma; extranodal marginal-zone B-cell lymphoma; multiple myeloma (MM); Waldenstrom macroglobulinemia; myelodysplastic syndrome (MDS), including refractory anemia (RA), refractory anemia with ring sideroblasts (RARS), refractory anemia with excess of blasts (RAEB) and refractory anemia with excess blasts in transformation (RAEB-T); and myeloproliferative syndrome.

In some embodiments, solid tumor is breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma.

In some embodiments, hematologic malignancy is acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia, chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, myeloma (such as multiple myeloma).

The compound of the present invention can be used to achieve a beneficial therapeutic or prophylactic effect, for example, in subjects with cancer.

In addition, the compounds of the present invention (e.g., a compound of any of the examples as described herein) can be administered in combination with additional therapeutic agents for the treatment of diseases or disorders described herein, such as cancer. The additional therapeutic agents may be administered separately with the compound of the present invention or included with such an ingredient in a pharmaceutical composition according to the disclosure, such as a fixed-dose combination drug product. In some embodiments, additional therapeutic agents are those that are known or discovered to be effective in the treatment of diseases mediated by SHP2 or at least in part by SHP2, such as another SHP2 inhibitor or a compound active against another target associated with the particular disease. The combination may serve to increase efficacy (e.g., by including in the combination a compound potentiating the potency or effectiveness of the compound of the present invention), decrease one or more side effects, or decrease the required dose of the compound of the present invention.

In some embodiments, the compounds of the present invention (e.g., a compound of any of the examples as described herein) can be administered in combination with additional therapeutic agents, such as anti-neoplastic active agents, anti-inflammatory agents, or immunomodulators, wherein the anti-neoplastic active agents include chemotherapeutic agents, immune checkpoint inhibitors or agonists, and targeted

therapeutic agents. The term “anti-neoplastic active agent” as used herein refers to any agent that is administered to a subject suffering from cancer for the purposes of treating the cancer, such as a chemotherapeutic agent, an immune checkpoint inhibitor or agonist, and a targeted therapeutic agent.

5 Non-limiting examples of chemotherapeutic agents include topoisomerase I inhibitors (e.g., irinotecan, topotecan, camptothecin and analogs or metabolites thereof, and doxorubicin); topoisomerase II inhibitors (e.g., etoposide, teniposide, mitoxantrone, idarubicin, and daunorubicin); alkylating agents (e.g., melphalan, chlorambucil, busulfan, thiotepa, ifosfamide, carmustine, lomustine, semustine, streptozocin, decarbazine,
10 methotrexate, mitomycin C, and cyclophosphamide); DNA intercalators (e.g., cisplatin, oxaliplatin, and carboplatin); free radical generators such as bleomycin; nucleoside mimetics (e.g., 5-fluorouracil, capecitabine, gemcitabine, fludarabine, cytarabine, azacitidine, mercaptopurine, thioguanine, pentostatin, and hydroxyurea); paclitaxel, docetaxel, and related analogs; vincristine, vinblastin, and related analogs; thalidomide
15 and related analogs (e.g., CC-5013 and CC-4047).

Non-limiting examples of immune checkpoint inhibitors or agonists include PD-1 inhibitors, for example, anti-PD-1 antibodies, such as pembrolizumab, nivolumab, and PDR001 (spartalizumab); PD-L1 inhibitors, for example, anti-PD-L1 antibodies, such as atezolizumab, durvalumab, and avelumab; CTLA-4 inhibitors, such as anti-CTLA-4
20 antibodies, for example ipilimumab; and BTLA inhibitors, LAG-3 inhibitors, TIM3 inhibitors, TIGIT inhibitors, VISTA inhibitors, OX-40 agonists, and the like.

Targeted therapeutic agents include various small molecule or macromolecular targeted therapeutic agents, and non-limiting examples thereof include: protein tyrosine kinase inhibitors (such as imatinib mesylate and gefitinib); proteasome inhibitors (such as
25 bortezomib); NF- κ B inhibitors, including I κ B kinase inhibitors; KRAS G12C inhibitors; ERK inhibitors; CDK4/6 inhibitors; PI3K δ inhibitors; SYK inhibitors; Bcl2 inhibitors; IDO inhibitors; A2AR inhibitors; BRAF inhibitors (such as dabrafenib); MEK inhibitors (such as trametinib); mTOR inhibitors (such as rapamycin); anti-CD40 antibodies (such as APX005M, RO7009789); antibodies that bind to proteins overexpressed in cancer to
30 down-regulate cell replication, such as anti-CD20 antibodies (such as rituximab, ibritumomab tiuxetan, and tositumomab), anti-Her2 monoclonal antibodies (such as trastuzumab), anti-EGFR antibodies (such as cetuximab) and anti-VEGF antibodies (such

as bevacizumab); anti-angiogenic drugs, such as lenalidomide; and other protein or enzyme inhibitors, these proteins or enzymes are known to be upregulated, overexpressed or activated in cancers, and the inhibition of which can down-regulate cell replication.

5

EXAMPLES

The examples below are intended to be purely exemplary and should not be considered to be limiting in any way. Efforts have been made to ensure the accuracy with respect to numbers used (for example, amounts, temperature, etc.), but those skilled in the art should understand that some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric. All MS data were determined by Agilent 6120 or Agilent 1100. All NMR data were generated using a Varian 400 MR machine. All reagents and materials, except synthesized intermediates, used in the present invention are commercially available. All compound names except the reagents are generated by Chemdraw 16.0.

If there is any atom with empty valence(s) in any one of the structures disclosed herein, the empty balance(s) is (are) the hydrogen atom(s) which is (are) omitted for convenience purpose.

In the present application, in the case of inconsistency of the name and structure of a compound, when the two of which are both given for the compound, it is subject to the structure of the compound, unless the context shows that the structure of the compound is incorrect and the name is correct.

List of abbreviations used in the following examples:

AcOH	Acetic acid
(Boc) ₂ O	Di-tert-butyl dicarbonate
BOP	Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate
CD ₃ OD	Deuterated methanol
CDCl ₃	Deuterated chloroform
<i>m</i> -CPBA	<i>m</i> -chloroperoxybenzoic acid
DBU	1,8-diazabicyclo-undec-7-ene
DCM	Dichloromethane


DIBAL-H	Diisobutylaluminium hydride
DIEA	<i>N,N</i> -diisopropylethylamine
DMAP	4-dimethylaminopyridine
DMA	<i>N,N</i> -dimethylacetamide
DMF	<i>N,N</i> -dimethylformamide
DMSO-d ₆	Deuterated dimethyl sulfoxide
EA/EtOAc	Ethyl acetate
Et ₃ N	Triethylamine
EtOH	Ethanol
g	Gram
HMDSLi	Lithium hexamethyldisilazide
HATU	2-(7-azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
KOAc	Potassium acetate
L	Liter
LDA	Lithium diisopropylamide
M	Mole/liter
MeCN	Acetonitrile
MeI	Iodomethane
MeOH	Methanol
MeSO ₃ H	Methanesulfonic acid
mg	Milligram
mL	Milliliter
mmol	Millimole
mol	Mole
NaOMe	Sodium methoxide
NBS	N-bromosuccinimide
NCS	<i>N</i> -chlorosuccinimide
NFSI	<i>N</i> -fluorobisbenzenesulfonamide
NIS	N-iodosuccinimide
Ti(OEt) ₄	Ethyl titanate
TMSCN	Trimethylsilyl cyanide

$\text{Pd}_2(\text{dba})_3$	Tris(dibenzylidene acetone)dipalladium
$\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$	[1,1'-bis(diphenylphosphino) ferrocene]palladium dichloride dichloromethane complex
$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$	Bis(triphenylphosphine)palladium chloride
PE	Petroleum ether
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
Xant-phos	4,5-bis(diphenylphosphino)-9,9-dimethylxanthene

Example 1 Synthesis of Compounds

Intermediate I-A1

Sodium 2-amino-3-chloropyridine-4-thiolate



Step 1: Methyl 3-((2-amino-3-chloropyridin-4-yl)thio)propanoate

Under nitrogen, 3-chloro-4-iodopyridin-2-amine (10.0 g, 39.3 mmol), methyl 3-mercaptopropanoate (5.20 g, 42.8 mmol), palladium acetate (0.44 g, 1.97 mmol), Xant-phos (2.27 g, 3.93 mmol) and DIEA (10.2 g, 78.6 mmol) were placed in 1,4-dioxane (160 mL). The reaction solution was refluxed and stirred for 2 hours, cooled to room temperature and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product (9.70 g, yield 100%). $[\text{M}+\text{H}]^+$ 201.1

Step 2: Sodium 2-amino-3-chloropyridine-4-thiolate

Under nitrogen, to a solution of methyl 3-((2-amino-3-chloropyridin-4-yl)thio)propanoate (9.70 g, 39.3 mmol) in tetrahydrofuran was added 2 M sodium ethoxide/ethanol solution (20 mL, 40 mmol). The reaction was stirred at room temperature for 2 hours and concentrated in vacuum under reduced pressure. To the resulting residue was added dichloromethane and the mixture was stirred. A solid was precipitated and the mixture was filtered. The filter cake was collected to give the target product (7.18 g, yield 100%). $[\text{M}+2\text{H}-\text{Na}]^+$ 161.0


The intermediates in the table below were prepared by following the steps for preparing intermediate I-A1 from corresponding starting materials and reagents:

Intermediates	Structural formula	LC-MS [M+2H-Na] ⁺	Intermediates	Structural formula	LC-MS [M+2H-Na] ⁺
I-A3		219.0	I-A5		289.0
I-A6		128.1	I-A7		215.0
I-A10		127.2	I-A20		180.1

Intermediate I-A2

5

3-((2-amino-3-chloropyridin-4-yl)thio)-6-chloropyrazin-2-amine



Step 1: Methyl 3-((3-amino-5-chloropyrazin-2-yl)thio)propanoate

Under nitrogen, 3-bromo-6-chloropyrazin-2-amine (500 mg, 2.4 mmol), methyl 3-mercaptopropanoate (317 mg, 2.6 mmol), Pd₂(dba)₃ (110 mg, 0.12 mmol), Xant-phos (138 mg, 0.24 mmol) and DIEA (620 mg, 4.8 mmol) were placed in 1,4-dioxane (20 mL). The reaction solution was refluxed and stirred for 16 hours, cooled to room temperature and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product (460 mg, yield 78%). [M+H]⁺ 248.0

Step 2: Sodium 3-((3-amino-5-chloropyrazin-2-yl)thio)propanoate

Under nitrogen, to a solution of methyl 3-((3-amino-5-chloropyrazin-2-yl)thio)propanoate (460 mg, 1.87 mmol) in tetrahydrofuran was added 2 M sodium ethoxide/ethanol solution. The reaction was stirred at room temperature for 2 hours and concentrated in vacuum under reduced pressure. To the resulting residue was added dichloromethane and the mixture was stirred. A solid was precipitated and the mixture

was filtered. The filter cake was collected to give the target product (400 mg, yield 118%). $[M+2H-Na]^+$ 162.0


Step 3: 3-((2-amino-3-chloropyridin-4-yl)thio)-6-chloropyrazin-2-amine

Under nitrogen, sodium 3-amino-5-chloropyrazine-2-thiolate (300 mg, 1.63 mmol),
 5 3-chloro-4-iodopyridin-2-amine (414 mg, 1.63 mmol), $Pd_2(dba)_3$ (75 mg, 0.08 mmol),
 Xant-phos (93 mg, 0.16 mmol) and DIEA (0.55 mL, 3.26 mmol) were placed in 1,4-
 dioxane (20 mL). The reaction solution was refluxed and stirred for 4 hours and cooled to
 room temperature. The reaction solution was concentrated in vacuum under reduced
 pressure, and the resulting residue was purified with silica gel column chromatography
 10 (water/methanol) to give the target product (300 mg, yield 65%). $[M+H]^+$ 288.0

The intermediates in the table below were prepared by following the steps for
 preparing intermediate I-A2 from corresponding starting materials and reagents:

Interme diate	Structural formula	LC-MS $[M+H]^+$	Interme diate	Structural formula	LC-MS $[M+H]^+$
I-A9		254.0	I-A11		302.0
I-A12		317.0	I-A13		301.0
I-A14		269.0	I-A15		273.0
I-A16		258.0	I-A17		287.0
I-A18		254.0	I-A19		307.0
I-A21		293.0	I-A22		288.0
I-A23		274.0	I-A24		444.0


I-A25		$[M+Na]^+$ 428.9	I-A26		257.0
I-A27		240.1	I-A28		272.0
I-A29		257.0	I-A30		292.0
I-A31		298.1	I-A32		291.1
I-A33		286.0	I-A34		272.1
I-A35		287.0	I-A36		443.0
I-A37		254.0	I-A38		288.0
I-A39		255.0	I-A40		273.0

Intermediate I-A4**6-amino-2-hydroxy-5-iodo-3-methylpyrimidin-4(3H)-one**

- 5 Under nitrogen, 6-amino-2-hydroxy-3-methylpyrimidin-4(3H)-one (1.41 g, 10 mmol) and NIS (2.47 g, 11 mmol) were placed in tetrahydrofuran (20 mL) and stirred at room temperature for 16 hours. The mixture was filtered, and the filter cake was collected to give the target product (2.40 g, yield 90%). $[M+H]^+$ 267.9

Intermediate I-A8

Tert-butyl (8-bromo-5-(methylsulfinyl)imidazo[1,2-c]pyrimidin-7-yl)(tert-butoxycarbonyl)carbamate



Step 1: 5-bromo-2-(methylthio)pyrimidine-4,6-diamine

At 0°C, to a solution of 2-(methylthio)pyrimidine-4,6-diamine (5 g, 32 mmol) in *N,N*-dimethylformamide (50 mL) was added NBS (6.2 g, 35 mmol). The reaction was stirred at room temperature for 12 hours, and the reaction solution was poured into water (200 mL), filtered and dried to give the target product as a yellow solid (5.7 g, yield 76%). $[M+H]^+$ 234.9, 236.9

Step 2: 8-bromo-5-(methylthio)imidazo[1,2-c]pyrimidin-7-amine

To a solution of 5-bromo-2-(methylthio)pyrimidine-4,6-diamine (5.7 g, 24.2 mmol) in *N,N*-dimethylformamide (70 mL) was added 40% 2-chloroacetaldehyde aqueous solution (7.1 g, 36 mmol). The reaction was stirred at 80°C for 2 hours, and the reaction solution was poured into water (500 mL), adjusted with solid sodium hydroxide to a pH value of 10 and extracted with ethyl acetate. The organic phases were collected and combined, and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product (3.0 g, yield 48%). $[M+H]^+$ 258.9, 260.9

Step 3: Tert-butyl (8-bromo-5-(methylthio)imidazo[1,2-c]pyrimidin-7-yl)(tert-butoxycarbonyl)carbamate

To a solution of 8-bromo-5-(methylthio)imidazo[1,2-c]pyrimidin-7-amine (3.0 g, 11.6 mmol) in tetrahydrofuran (40 mL) were added $(Boc)_2O$ (7.6 g, 24.8 mmol) and DMAP (283 mg, 2.3 mmol). The reaction was stirred at room temperature for 12 hours, and the reaction solution was poured into water (200 mL) and extracted with ethyl acetate. The organic phases were collected and combined, and concentrated in vacuum under


reduced pressure, and the resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product (3.6 g, yield 68%). $[M+H]^+$ 459.2, 461.2. ^1H NMR (400 MHz, CDCl_3): δ 7.71 (d, $J = 1.4$ Hz, 1H), 7.57 (d, $J = 1.4$ Hz, 1H), 2.72 (s, 3H), 1.42 (s, 18H).

Step 4: Tert-butyl (8-bromo-5-(methylsulfinyl)imidazo[1,2-c]pyrimidin-7-yl)(tert-butoxycarbonyl)carbamate

At 0°C , to a solution of tert-butyl (8-bromo-5-(methylthio)imidazo[1,2-c]pyrimidin-7-yl)(tert-butoxycarbonyl)carbamate (400 mg, 0.87 mmol) in dichloromethane (10 mL) was added m-chloroperoxybenzoic acid (530 mg, 2.6 mmol). The reaction was stirred at 0°C for 2 hours, and a saturated sodium hydrogen sulfite aqueous solution (2 mL) was added thereto. The reaction solution was adjusted with a saturated sodium bicarbonate aqueous solution to a pH value of 8 and extracted with dichloromethane. The organic phases were combined, dried over anhydrous sodium sulfate, and concentrated to give a crude target product (500 mg, yield 120%), which was used in the reaction of the next step directly. $[M+H]^+$ 475.0

Intermediates I-B1 and I-B2

Enantiomers of 1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate



Step 1: N-(2,3-dichlorophenyl)acetimidamide

2,3-dichloroaniline (13.0 g, 80.0 mmol) was placed in 1 M hydrogen chloride/acetonitrile (160 mL, 160 mmol). The reaction was stirred at 120°C for 16 hours and concentrated in vacuum under reduced pressure to give the target product (19.5 g, yield 120%), which was used in the reaction of the next step directly.

Step 2: 1-(2,3-dichlorophenyl)-2-methylpyrimidine-4,6(1H,5H)-dione

N-(2,3-dichlorophenyl)acetimidamide (19.5 g, 80 mmol) and diethyl malonate (25.6 g, 160 mmol) were placed in ethanol (80 mL), and to the solution was added 2 M sodium ethoxide/ethanol solution (120 mL, 240 mmol). The reaction was stirred at 120°C for 16 hours and concentrated in vacuum under reduced pressure. The resulting residue was dissolved in water (100 mL), and the solution was adjusted with 6 M hydrochloric acid to a pH value of 2, and a solid was precipitated. The mixture was filtered, and the filter cake was collected and dried under reduced pressure to give the target product (10.5 g, yield 49%). $[M+H]^+$ 271.0



Step 3: 1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate

To a solution of 3-(2,3-dichlorophenyl)-6-hydroxy-2-methylpyrimidin-4(3H)-one (10.5 g, 38.7 mmol) and 2,4,6-triisopropylbenzenesulfonic acid (17.7 g, 58.5 mmol) in dichloromethane (50 mL) were added DMAP (240 mg, 1.9 mmol) and triethylamine (9.9 g, 97.4 mmol). The reaction was stirred at room temperature for 2 hours and concentrated in vacuum under reduced pressure. The resulting residue was purified with silica gel column chromatography (ethyl acetate/petroleum ether=1/3) to give the target product (18.0 g, yield 87%). $[M+H]^+$ 537.2

Step 4: Enantiomers of 1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate

1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate (19.0 g) was resolved by chiral HPLC to obtain a pair of enantiomers. Chiral HPLC resolution conditions: column: IG-H (0.46 cm I.D. × 15 cm L); mobile phase: carbon dioxide/ethanol = 60 : 40; flow rate: 2.5 mL/minute; detector: UV 254 nm. First eluent (intermediate I-B2, 8.66 g, RT = 0.808 minutes), ee% = 100%. Second eluent (intermediate I-B1, 9.66 g, RT = 1.236 minutes), ee% = 99.94%.

The intermediates in the table below were prepared by following the steps 1-3 for preparing intermediates I-B1 and I-B2 from corresponding starting materials and reagents:

Intermediates	Structural formula	LC-MS $[M+H]^+$	Intermediates	Structural formula	LC-MS $[M+H]^+$
I-B3		505.2	I-B5		521.2

I-B6		503.1	I-B7		469.2
I-B8		503.2	I-B9		517.2
I-B10		537.1	I-B11		528.2
I-B12		505.2	I-B14		509.2
I-B15		505.2	I-B16		503.2
I-B17		499.2	I-B18		533.2
I-B19		521.1	I-B20		547.2
I-B23		515.2	I-B25		505.2
I-B27		521.2	I-B28		547.2

I-B29		521.1	I-B30		555.2
I-B31		537.2	I-B34		533.1
I-B37		571.2	I-B38		537.2

The intermediates in the table below were prepared by following the steps 1-3 for preparing intermediates I-B1 and I-B2 from corresponding substituted aniline, diethyl 2-methylmalonate or diethyl 2-fluoromalonate, and reagents:



Intermediates	Structural formula	LC-MS [M+H] ⁺	Intermediates	Structural formula	LC-MS [M+H] ⁺
I-B4		519.2	I-B13		551.1
I-B24		535.2	I-B35		561.2
I-B42		555.1			

5

The intermediates in the table below were prepared from intermediate I-B13 by chiral resolution by following the step 4 for preparing intermediates I-B1 and I-B2:


Chiral HPLC resolution conditions: column: IG-H (0.46 cm I.D. × 15 cm L); mobile phase: carbon dioxide/ethanol = 80 : 20; flow rate: 2.5 mL/minute; detector: UV 254 nm.

First eluent (intermediate I-B21, RT = 2.784 minutes), ee% = 100%. Second eluent (intermediate I-B22, RT = 3.119 minutes), ee% = 99.92%.

Intermediates	Structural formula	LC-MS [M+H] ⁺
I-B21		551.1
I-B22		551.1

Intermediate I-B26

5 1-(2-chloropyridin-4-yl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate



Step 1: *N*-(2-chloropyridin-4-yl)acetimidamide

2-chloro-4-iodopyridine (3.78 g, 15.8 mmol), acetimidamide hydrochloride (1.92 g, 20.3 mmol), cuprous iodide (301 mg, 1.58 mmol), cesium carbonate (13.2 g, 40.6 mmol) and *N,N*-dimethylformamide (22 mL) were placed in a sealed tube. The reaction was stirred at 90°C for 12 hours and cooled to room temperature, and acetonitrile (200 mL) was added thereto. The mixture was filtered, and the filtrate was concentrated in vacuum under reduced pressure to give the target product (1.75 g, yield 70%), which was used in the reaction of the next step directly. [M+H]⁺ 170.0

Step 2: 3-(2-chloropyridin-4-yl)-6-hydroxy-2-methylpyrimidin-4(3*H*)-one

N-(2-chloropyridin-4-yl)acetimidamide (1.70 g, 10.0 mmol), bis(2,4,6-trichlorophenyl)malonate (5.55 g, 12 mmol) and tetrahydrofuran (20 mL) were placed in

a sealed tube. The reaction was stirred at 90°C for 16 hours and concentrated in vacuum under reduced pressure. The resulting residue was purified with silica gel column chromatography (water/methanol) to give the target product (1.19 g, yield 50%). $[M+H]^+$ 238.0


5 **Step 3: 1-(2-chloropyridin-4-yl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate**

The target product was prepared by following the step 3 for preparing intermediates I-B1 and I-B2 from corresponding starting materials and reagents. $[M+H]^+$ 504.2

The intermediates in the table below were prepared by following the steps for

10 preparing intermediate I-B26 from corresponding starting materials and reagents:

Intermediates	Structural formula	LC-MS $[M+H]^+$	Intermediates	Structural formula	LC-MS $[M+H]^+$
I-B32		539.2	I-B33		535.2
I-B36		528.2	I-B39		591.1
I-B40		515.2	I-B41		551.2
I-B43		546.2	I-B44		562.1
I-B45		542.1	I-B46		567.1
I-B47		551.2			

Intermediate I-B48**2-amino-1-(2,3-dichlorophenyl)-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate****Step 1: 1-(2,3-dichlorophenyl)thiourea**

The solution of 2,3-dichloroaniline (6.48 g, 40 mmol) and benzoyl isothiocyanate (9.79 g, 60 mmol) in acetone (20 mL) was stirred at reflux for 30 minutes. The reaction solution was poured into ice-water, filtered. The filter cake was washed with cold acetone, collected, added with 1 M NaOH aqueous solution (50 mL) and stirred at 80°C for 3 hours. After cooling to room temperature, the mixture was extracted with DCM. The organic layer was collected and concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with PE/EA) to give target product as white solid (8.2 g, 93% yield).

Step 2: 3-(2,3-dichlorophenyl)-6-hydroxy-2-mercaptopyrimidin-4(3H)-one

The solution of 1-(2,3-dichlorophenyl)thiourea (1.1 g, 5.0 mmol), diethyl malonate (1.6 g, 10.0 mmol), 18-crown-6 (661 mg, 2.5 mmol) and 2 M NaOMe/MeOH (5.0 mL, 10.0 mmol) in 1,4-dioxane (15 mL) was stirred at 70°C for 3 hours. The reaction solution was purified by silica gel column chromatography (eluting with DCM/MeOH) to give target product as yellow solid (1.8 g, 99% yield). $[M+H]^+$ 289.0

Step 3: 3-(2,3-dichlorophenyl)-6-hydroxy-2-(methylthio)pyrimidin-4(3H)-one

At 0°C, to the solution of 3-(2,3-dichlorophenyl)-6-hydroxy-2-mercaptopyrimidin-4(3*H*)-one (1.45 g, 5.0 mmol) in THF (10 mL) was added MeI (1.42 g, 10.0 mmol) and stirred at room temperature for 3 hours. The reaction solution was concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with DCM/MeOH) to give target product as yellow oil (1.18 g, 78% yield).

Step 4: 1-(2,3-dichlorophenyl)-2-(methylthio)-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate

The solution of 3-(2,3-dichlorophenyl)-6-hydroxy-2-(methylthio)pyrimidin-4(3*H*)-one (1.18 g, 3.9 mmol), bis(2,4,6-trichlorophenyl) malonate (1.77 g, 5.9 mmol), DMAP (24 mg, 0.20 mmol) and Et₃N (790 mg, 7.8 mmol) in DCM (25 mL) was stirred at room temperature for 2 hours. The reaction solution was concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with PE/EA) to give target product as white solid (500 mg, 22% yield). [M+H]⁺ 568.4


Step 5: 1-(2,3-dichlorophenyl)-2-(methylsulfonyl)-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate

At 0°C, to the solution of 1-(2,3-dichlorophenyl)-2-(methylthio)-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate (500 mg, 0.88 mmol) in DCM (30 mL) was added the solution of *m*-CPBA (757 mg, 4.4 mmol) in DCM dropwise, and stirred at room temperature for 4 hours. The reaction solution was concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with PE/EA) to give target product (420 mg, 80% yield).

Step 6: 2-amino-1-(2,3-dichlorophenyl)-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate

1-(2,3-dichlorophenyl)-2-(methylsulfonyl)-6-oxo-1,6-dihydropyrimidin-4-yl 2,4,6-triisopropylbenzenesulfonate (420 mg, 0.70 mmol) was dissolved in 0.5 M NH₃/THF solution (7.0 mL, 3.5 mmol) and stirred at room temperature for 6 hours. The reaction solution was concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with PE/EA) to give target product as yellow oil (120 mg, 48% yield). [M+H]⁺ 538.2.

(R)-2-methyl-N-((S)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)propane-2-sulfinamide



Step 1: 1-(tert-butyl) 4-ethyl 4-(4-bromobenzyl)piperidine-1,4-dicarboxylate

At -78°C, under nitrogen, to a solution of 1-tert-butyl-4-ethylpiperidine-1,4-dicarboxylate (20.6 g, 80 mmol) in anhydrous tetrahydrofuran (100 mL) was added dropwise 2 M LDA/tetrahydrofuran solution (52 mL, 104 mmol). The reaction was stirred at -78°C for 2 hours, and a solution of 1-bromo-4-(bromomethyl)benzene (19.4 g, 80 mmol) in anhydrous tetrahydrofuran was added dropwise thereto. The reaction was stirred at -78°C for 3 hours and warmed to room temperature, and the reaction solution was poured into water and extracted with ethyl acetate. The organic phase was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated in vacuum under reduced pressure to give the target product as a white solid, which was used in the reaction of the next step directly.

Step 2: 4-(4-bromobenzyl)-1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid

1-(tert-butyl) 4-ethyl 4-(4-bromobenzyl)piperidine-1,4-dicarboxylate (80 mmol) and lithium hydroxide (18.7 g, 780 mmol) were placed in ethanol (200 mL) and water (100 mL). The reaction was stirred at 90°C for 16 hours and concentrated in vacuum under reduced pressure. The residue was dissolved in water and washed with ethyl acetate/petroleum ether (volume ratio 1 : 1). The aqueous phase was collected, adjusted with 6 M hydrochloric acid to a pH value of 3, and extracted with ethyl acetate. The organic phase was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated in vacuum under reduced pressure to give the target product as a white solid

(34.0 g, two-step yield 107%), which was used in the reaction of the next step directly.
[M+H-56]⁺ 342.2

Step 3: Tert-butyl 6-bromo-1-oxo-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate

5 4-(4-bromobenzyl)-1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid (34.0 g, 85 mmol) was placed in polyphosphoric acid (200 mL). The reaction was stirred at 120°C for 16 hours, cooled to room temperature, and dissolved by adding water, and the mixture was adjusted with sodium hydroxide to a pH value of 9. To the solution was added (Boc)₂O, and the mixture was stirred at room temperature for 16 hours. The reaction
10 solution was extracted with ethyl acetate, and the organic phase was collected and concentrated in vacuum under reduced pressure. The resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product as a yellow solid (23.4 g, yield 72%). [M+H-56]⁺ 324.0

Step 4: Tert-butyl (R,E)-6-bromo-1-((tert-butylsulfinyl)imino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate

15 Under nitrogen, tert-butyl 6-bromo-1-oxo-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (23.4 g, 62 mmol) and (R)-2-methylpropane-2-sulfinamide (28.7 g, 237 mmol) were placed in Ti(OEt)₄ (200 mL). The reaction was stirred at 80°C for 16 hours, and the reaction solution was poured into water/ethyl acetate and filtered. The organic
20 phase was washed with saturated brine, dried over anhydrous sodium sulfate, and concentrated in vacuum under reduced pressure to give the target product, which was used in the reaction of the next step directly.

Step 5: Tert-butyl (S)-6-bromo-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate

25 Under nitrogen, tert-butyl (R,E)-6-bromo-1-((tert-butylsulfinyl)imino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (62 mmol) was placed in anhydrous tetrahydrofuran (200 mL). Sodium borohydride (9.07 g, 240 mmol) was added in batches to the mixture at -78°C, and the resulting solution was stirred at this temperature for 30 minutes, warmed to room temperature, poured into water, and extracted with ethyl
30 acetate. The organic phase was collected and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography

(petroleum ether/ethyl acetate) to give the target product as a yellow solid (14.5 g, yield 48%). $[M+H]^+$ 485.2



Step 6: Tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-6-(((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate

Under nitrogen, tert-butyl (S)-6-bromo-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (6.0 g, 12.4 mmol), ethynyltrimethylsilane (20 ml), $Pd(PPh_3)_2Cl_2$ (0.87 g, 1.2 mmol), cuprous iodide (0.24 g, 1.2 mmol), triethylamine (40 mL) and *N,N*-dimethylformamide (40 mL) were placed in a sealed tube. The reaction was stirred at 90°C for 16 hours and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product.

Step 7: (R)-2-methyl-N-((S)-5-(((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)propane-2-sulfinamide

To a solution of tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-6-(((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (12.4 mmol) in dichloromethane (60 mL) was added dropwise methanesulfonic acid (3.6 g, 37.1 mmol). The reaction was stirred at room temperature for 30 minutes, and the reaction solution was adjusted with aqueous ammonia to a pH value of 8 under ice bath cooling. The organic phase was collected and concentrated in vacuum under reduced pressure to give the target product as a brown solid (3.9 g, two-step yield 78%). $[M+H]^+$ 403.2

The intermediates in the table below were prepared by following the steps for preparing intermediate I-C3 from corresponding starting materials and reagents:


Intermediates	Structural formula	LC-MS $[M+H]^+$	Intermediates	Structural formula	LC-MS $[M+H]^+$
I-C4		385.2	I-C6		403.2

I-C7		375.2	I-C10		388.4
I-C12		403.2	I-C13		375.2
I-C14		397.2	I-C15		389.2
I-C16		374.2	I-C17		345.2
I-C18		393.2	I-C19		421.2
I-C26		421.2	I-C30		403.2
I-C31		389.0	I-C34		285.2

Intermediate I-C20

3-((*S*)-1-(((*R*)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-methylpropiolamide

84



Step 1: tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-6-(3-(methylamino)-3-oxoprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate






Under N₂, tert-butyl (S)-6-bromo-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (4.85 g, 10.0 mmol), Pd(PPh₃)₂Cl₂ (701 mg, 0.1 mmol), CuI (380 mg, 0.2 mmol), KOAc (2.94 g, 30.0 mmol) and DMSO (100 mL) were placed in three-necked flask. At 90 °C, to the stirred mixture was added dropwise the solution of N-methylpropiolamide (2.49 g, 30.0 mmol) in DMSO (50 mL) during 2 hours, and then stirred for additional 3 hours. The reaction solution was poured into water and filtered. The solid was collected and purified by silica gel column chromatography (eluting with DCM/EA) to give target product.

Step 2: 3-((S)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-methylpropiolamide

In an ice-bath, to the solution of tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-6-(3-(methylamino)-3-oxoprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate in DCM (15 mL) was added MeSO₃H (3.84 g, 40.0 mmol) dropwise. The reaction solution was stirred at room temperature for 30 minutes, adjusted pH to 8 with aqueous ammonia in an ice-bath. The organic layer was collected, concentrated in vacuum under reduced pressure to give target product as yellow solid (2.2 g, 57% yield of two steps). [M+H]⁺ 388.2


The intermediates in the table below were prepared by following the steps for preparing intermediate I-C20 from corresponding starting materials and reagents:

Intermediates	Structural formula	LC-MS [M+H] ⁺	Intermediates	Structural formula	LC-MS [M+H] ⁺
I-C21		402.2	I-C23		402.2

I-C24		406.2	I-C25		414.2
I-C27		416.2	I-C33		432.2
I-C36		414.2			

Intermediate I-C35

3-((*S*)-1-(((*R*)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-(2-methoxyethyl)-*N*-methylpropiolamide



Step 1: 3-((*S*)-1'-(tert-butoxycarbonyl)-1-(((*R*)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)propionic acid

To the solution of tert-butyl (*S*)-1-(((*R*)-tert-butylsulfinyl)amino)-6-(3-methoxy-3-oxoprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (1.0 g, 2.05 mmol; prepared by following the steps 1-6 for preparing intermediate I-C3) in EtOH/water (10 mL/2 mL) was added LiOH (245 mg, 10.23 mmol), stirred at 85°C for 1 hour and concentrated in vacuum under reduced pressure. The residue was dissolved in

water, extracted with EA. The aqueous layer was collected, adjusted pH to 6 with AcOH, extracted with DCM. The organic layer was collected, dried over anhydrous Na₂SO₄, and concentrated in vacuum under reduced pressure to give target product as white solid (920 mg, 95% yield). [M+H]⁺ 475.2.

5 **Step 2: tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-6-(3-((2-methoxyethyl)(methyl)amino)-3-oxoprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate**

To the solution of 3-((S)-1'-(tert-butoxycarbonyl)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)propionic acid (477 mg, 1.0 mmol), 2-methoxy-*N*-methylethan-1-amine (178 mg, 2.0 mmol) and HATU (760 mg, 2.0 mmol) in
10 DMF (5 mL) was added Et₃N (202 mg, 2.0 mmol) dropwise. The reaction mixture was stirred at room temperature for 1 hour, poured into water and filtered. The solid was collected to give target product.


15 **Step 3: 3-((S)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-(2-methoxyethyl)-*N*-methylpropiolamide**

The target product was prepared by following the step 7 for preparing intermediate I-C3 from corresponding starting materials and reagents (380 mg, 85% yield of two steps). [M+H]⁺ 446.2

20

Intermediate I-C1



(R)-*N*-((S)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide



To a solution of tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-5-
25 ((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (1.08 g, 2.2 mmol) in dichloromethane (10 mL) was added dropwise trifluoroacetic acid (5 mL). The reaction was stirred at room temperature for 5 minutes and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column

chromatography (water/methanol) to give the target product (0.25 g, yield 35%). $[M+H]^+$ 331.2


The intermediates in the table below were prepared by following the steps for preparing intermediate I-C1 from corresponding starting materials and reagents:

Intermediate	Structural formula	LC-MS $[M+H]^+$	Intermediate	Structural formula	LC-MS $[M+H]^+$
I-C2		331.2	I-C22		227.2

5

Intermediate I-C5

(R)-2-methyl-N-((R)-6-((trimethylsilyl)ethynyl)-3H-spiro[benzofuran-2,4'-piperidin]-3-yl)propane-2-sulfinamide



10 Step 1: 2-(4-bromo-2-fluorophenyl)-2-((trimethylsilyl)oxy)acetonitrile

Under nitrogen, to a solution of 4-bromo-2-fluorobenzaldehyde (6.84 g, 33.7 mmol) and DMAP (50 mg) in acetonitrile (50 mL) was added dropwise TMSCN (3.78 g, 38.0 mmol). The reaction was stirred at room temperature for 4 hours and concentrated in vacuum under reduced pressure, and the residue was used in the reaction of the next step directly.

15 Step 2: Tert-butyl 4-(4-bromo-2-fluorobenzoyl)-4-hydroxypiperidine-1-carboxylate

At -78°C, under nitrogen, 2-(4-bromo-2-fluorophenyl)-2-((trimethylsilyl)oxy)acetonitrile was dissolved in anhydrous tetrahydrofuran (150 mL),

and 1 M LiHMDS/tetrahydrofuran solution (37.1 mL, 37.1 mmol) was added dropwise thereto. The reaction was stirred at -78°C for 1.5 hours, and a solution of tert-butyl 4-oxopiperidine-1-carboxylate (7.39 g, 37.1 mmol) in anhydrous tetrahydrofuran (30 mL) was added dropwise thereto. The reaction was stirred at -78°C for 3 hours, and 1 M hydrochloric acid (200 mL) was added dropwise thereto. The reaction solution was warmed to room temperature, the organic phase was collected, and the aqueous phase was extracted with ethyl acetate. The organic phases were combined, washed with saline, dried over anhydrous sodium sulfate and concentrated in vacuum under reduced pressure, and the residue was used in the reaction of the next step directly.





Step 3: Tert-butyl 6-bromo-3-oxo-3H-spiro[benzofuran-2,4'-piperidine]-1'-carboxylate

Tert-butyl 4-(4-bromo-2-fluorobenzoyl)-4-hydroxypiperidine-1-carboxylate (13.56 g, 33.7 mmol) was dissolved in 1,4-dioxane, and potassium t-butoxide (4.16 g, 37.1 mmol) was added thereto. The reaction was stirred at 85°C for 3 hours and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (petroleum ether/ethyl acetate) to give the target product (6.6 g, yield 51.2%). $[M+H]^+$ 326.0

Step 4: (R)-2-methyl-N-((R)-6-((trimethylsilyl)ethynyl)-3H-spiro[benzofuran-2,4'-piperidin]-3-yl)propane-2-sulfonamide


The target product was prepared by following the steps 4-7 for preparing intermediate I-C3 from corresponding starting materials and reagents. $[M+H]^+$ 405.2

The intermediates in the table below were prepared by following the steps for preparing intermediate I-C5 from corresponding starting materials and reagents:

Intermediate	Structural formula	LC-MS $[M+H]^+$	Intermediate	Structural formula	LC-MS $[M+H]^+$
I-C9		387.2	I-C28		404.2
I-C29		390.2	I-C32		405.2

Intermediates I-C8 and I-C8'

(R)-N-((S)-5-(3-hydroxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide and (R)-N-((S)-5-(3-(tert-butoxy)prop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide



Step 1: Tert-butyl (S)-6-(3-((tert-butyldimethylsilyl)oxy)prop-1-yn-1-yl)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1-carboxylate

The target product was prepared by following the step 6 for preparing intermediate I-C3 from corresponding starting materials and reagents. $[M+H]^+$ 575.2

Step 2: (R)-N-((S)-5-(3-hydroxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide and (R)-N-((S)-5-(3-(tert-butoxy)prop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide

The mixture of target products I-C8 and I-C8' was prepared by following the step 7 for preparing intermediate I-C3 from corresponding starting materials and reagents. $[M+H]^+$ 361.2, 417.2


Step 3: (R)-N-((S)-5-(3-hydroxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide

The target product was prepared by following the steps for preparing intermediate I-C1 from corresponding starting materials and reagents. $[M+H]^+$ 361.2

Intermediate I-C11

(R)-2-methyl-N-((S)-5-((1-methyl-1H-pyrazol-4-yl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)propane-2-sulfinamide

90



Step 1: Tert-butyl (S)-1-(((R)-tert-butylsulfinyl)amino)-6-((1-methyl-1H-pyrazol-4-yl)amino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate

Under nitrogen, tert-butyl (S)-6-bromo-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidine]-1'-carboxylate (485 mg, 1.0 mmol), 1-methyl-1H-pyrazol-4-amine (135 mg, 1.4 mmol), Xant-phos (35 mg, 0.06 mmol), Pd₂(dba)₃ (27 mg, 0.03 mmol) and CsCO₃ (650 mg, 2.0 mmol) were placed in 1,4-dioxane (15 mL). The reaction was stirred at 110°C for 16 hours and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (water/methanol) to give the target product. [M+H]⁺ 502.3


Step 2: (R)-2-methyl-N-((S)-5-((1-methyl-1H-pyrazol-4-yl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)propane-2-sulfinamide

The target product was prepared by following the steps for preparing intermediate I-C1 from corresponding starting materials and reagents. [M+H]⁺ 402.2

15

Compound 1

(S)-1'-(8-((2-amino-3-chloropyridin-4-yl)thio)imidazo[1,2-c]pyrimidin-5-yl)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1-amine



Step 1: (R)-N-((S)-5-ethynyl-1'-(8-iodoimidazo[1,2-c]pyrimidin-5-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide

20

Intermediate I-C1 (149 mg, 0.45 mmol), 5-chloro-8-iodoimidazo[1,2-c]pyrimidine (126 mg, 0.45 mmol) and diisopropylethylamine (116 mg, 0.90 mmol) were placed in *N,N*-dimethylformamide (2 mL). The reaction was stirred at room temperature for 30 minutes. The reaction solution was purified with silica gel column chromatography (water/methanol) to give the target product (132 mg, yield 51%). $[M+H]^+$ 574.1

Step 2: (R)-*N*-((S)-1'-(8-((2-amino-3-chloropyridin-4-yl)thio)imidazo[1,2-c]pyrimidin-5-yl)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide





Under nitrogen, (R)-*N*-((S)-5-ethynyl-1'-(8-iodoimidazo[1,2-c]pyrimidin-5-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide (132 mg, 0.23 mmol), intermediate I-A1 (42 mg, 0.23 mmol), Xant-phos (13.3 mg, 0.023 mmol), $Pd_2(dba)_3$ (10.5 mg, 0.012 mmol) and diisopropylethylamine (59 mg, 0.46 mmol) were placed in 1,4-dioxane (4 mL). The reaction was stirred at 100°C for 3 hours and concentrated in vacuum under reduced pressure, and the resulting residue was purified with silica gel column chromatography (water/methanol) to give the target product (101 mg, yield 72%). $[M+H]^+$ 606.2





Step 3: (S)-1'-(8-((2-amino-3-chloropyridin-4-yl)thio)imidazo[1,2-c]pyrimidin-5-yl)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1-amine





(R)-*N*-((S)-1'-(8-((2-amino-3-chloropyridin-4-yl)thio)imidazo[1,2-c]pyrimidin-5-yl)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide (101 mg, 0.17 mmol) was dissolved in 2 M hydrogen chloride/methanol solution. The reaction was stirred at room temperature for 3 minutes. Under ice bath cooling, the reaction solution was diluted with dichloromethane (15 mL) and adjusted with aqueous ammonia to a pH value of 8. The organic phase was collected and concentrated in vacuum under reduced pressure, and the resulting residue was purified with thin layer chromatography (dichloromethane/methanol = 12/1) to give the target product (76 mg, two-step yield 91%). $[M+H]^+$ 502.1. ^1H NMR (400 MHz, CD_3OD): δ 8.08-7.97 (m, 1H), 7.85-7.79 (m, 1H), 7.57-7.51 (m, 1H), 7.51-7.45 (m, 1H), 7.39-7.28 (m, 3H), 5.93-5.82 (m, 1H), 4.06-3.96 (m, 3H), 3.43-3.33 (m, 3H), 3.20-3.13 (m, 1H), 2.86-2.74 (m, 1H), 2.14-2.05 (m, 1H), 2.04-1.95 (m, 1H), 1.76-1.65 (m, 1H), 1.53-1.42 (m, 1H).





The compounds in the table below were prepared by following the steps for preparing compound 1 from corresponding intermediates and reagents:





Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Intermediates
3		560.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.06-7.99 (m, 1H), 7.85-7.80 (m, 1H), 7.59-7.51 (m, 2H), 7.38-7.30 (m, 3H), 5.88-5.79 (m, 1H), 4.06-3.96 (m, 3H), 3.60-3.52 (m, 4H), 3.43-3.33 (m, 6H), 3.21-3.14 (m, 1H), 2.87-2.75 (m, 1H), 2.15-2.05 (m, 1H), 2.04-1.94 (m, 1H), 1.75-1.64 (m, 1H), 1.54-1.43 (m, 1H).	I-A3 I-C1
7		546.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.04-8.01 (m, 1H), 7.84-7.81 (m, 1H), 7.57-7.53 (m, 2H), 7.38-7.31 (m, 3H), 5.86-5.82 (m, 1H), 4.07-3.96 (m, 3H), 3.72-3.67 (m, 2H), 3.55-3.50 (m, 2H), 3.45-3.34 (m, 3H), 3.21-3.14 (m, 1H), 2.86-2.77 (m, 1H), 2.15-2.04 (m, 1H), 2.04-1.95 (m, 1H), 1.73-1.67 (m, 1H), 1.52-1.45 (m, 1H).	I-A5 I-C1
8		502.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.00 (s, 1H), 7.81-7.77 (m, 1H), 7.54-7.50 (m, 1H), 7.48 (s, 1H), 7.46-7.43 (m, 1H), 7.31-7.27 (m, 1H), 7.21-7.16 (m, 1H), 5.87-5.83 (m, 1H), 4.01-3.94 (m, 3H), 3.41-3.39 (m, 1H), 3.37-3.34 (m, 1H), 3.31-3.27 (m, 1H), 3.20-3.12 (m, 1H), 2.82-2.75 (m, 1H), 2.12-2.02 (m, 1H), 2.02-1.92 (m, 1H), 1.72-1.62 (m, 1H), 1.50-1.40 (m, 1H).	I-A1 I-C2
9		469.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.03-7.97 (m, 1H), 7.86-7.82 (m, 1H), 7.81-7.78 (m, 1H), 7.57-7.51 (m, 1H), 7.37-7.29 (m, 3H), 6.25-6.18 (m, 1H), 4.02-3.94 (m, 3H), 3.42-3.34 (m, 3H), 3.20-3.13 (m, 1H), 2.85-2.76 (m, 1H), 2.13-2.05 (m, 1H), 2.03-1.95 (m, 1H), 1.74-1.64 (m, 1H), 1.54-1.43 (m, 1H).	I-A6 I-C1





10		556.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.03-7.99 (m, 1H), 7.83-7.79 (m, 1H), 7.56-7.50 (m, 2H), 7.37-7.29 (m, 3H), 5.86-5.77 (m, 1H), 4.46-4.37 (m, 1H), 4.04-3.96 (m, 3H), 3.44-3.34 (m, 3H), 3.19-3.14 (m, 1H), 2.85-2.75 (m, 1H), 2.41-2.31 (m, 2H), 2.13-2.05 (m, 1H), 2.03-1.94 (m, 3H), 1.79-1.66 (m, 3H), 1.52-1.43 (m, 1H).	I-A7 I-C1
27		469.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.00 (s, 1H), 7.83 (d, J = 5.5 Hz, 1H), 7.79 (d, J = 1.6 Hz, 1H), 7.54 (d, J = 1.6 Hz, 1H), 7.49 (s, 1H), 7.32-7.27 (m, 1H), 7.24-7.17 (m, 1H), 6.21 (d, J = 5.5 Hz, 1H), 4.03-3.93 (m, 3H), 3.40 (s, 1H), 3.37-3.33 (m, 2H), 3.23-3.12 (m, 1H), 2.86-2.75 (m, 1H), 2.14-2.04 (m, 1H), 2.04-1.92 (m, 1H), 1.75-1.63 (m, 1H), 1.52-1.40 (m, 1H).	I-A6 I-C3
40		546.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.02 (s, 1H), 7.84-7.78 (m, 1H), 7.56-7.50 (m, 1H), 7.50-7.44 (m, 2H), 7.31-7.25 (m, 1H), 7.23-7.17 (m, 1H), 5.91-5.82 (m, 1H), 4.29 (s, 2H), 4.07-3.93 (m, 3H), 3.42-3.32 (m, 5H), 3.21-3.14 (m, 1H), 2.86-2.75 (m, 1H), 2.13-1.95 (m, 2H), 1.73-1.64 (m, 1H), 1.50-1.42 (m, 1H).	I-A1 I-C7
71		559.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.03 (s, 1H), 7.84-7.80 (m, 1H), 7.56-7.51 (m, 1H), 7.50-7.44 (m, 2H), 7.31-7.25 (m, 1H), 7.23-7.17 (m, 1H), 5.89-5.85 (m, 1H), 4.05-3.96 (m, 3H), 3.49-3.45 (m, 2H), 3.42-3.33 (m, 2H), 3.23-3.14 (m, 1H), 2.85-2.77 (m, 1H), 2.36 (s, 6H), 2.14-2.06 (m, 1H), 2.03-1.95 (m, 1H), 1.74-1.64 (m, 1H), 1.52-1.42 (m, 1H).	I-A1 I-C10





72		558.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.06 (s, 1H), 7.86 (s, 1H), 7.79-7.69 (m, 2H), 7.60-7.53 (m, 2H), 7.51-7.47 (m, 1H), 5.94-5.85 (m, 1H), 4.19-4.11 (m, 2H), 3.53-3.48 (m, 2H), 3.46-3.38 (m, 2H), 3.34-3.30 (m, 2H), 2.38 (s, 6H), 2.23-2.15 (m, 2H), 1.70-1.62 (m, 2H).	I-A1 I-C10
75		573.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74 (d, J = 1.6 Hz, 1H), 7.51-7.41 (m, 3H), 7.28 (dd, J = 7.7, 1.2 Hz, 1H), 7.20 (d, J = 7.8 Hz, 1H), 5.79 (d, J = 5.6 Hz, 1H), 4.05-3.97 (m, 3H), 3.47 (s, 2H), 3.42-3.33 (m, 2H), 3.18 (d, J = 16.2 Hz, 1H), 2.81 (d, J = 16.1 Hz, 1H), 2.53 (s, 3H), 2.36 (s, 6H), 2.15-2.04 (m, 1H), 2.03-1.93 (m, 1H), 1.69 (d, J = 13.0 Hz, 1H), 1.47 (d, J = 13.3 Hz, 1H).	I-A1 I-C10
76		532.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.03 (s, 1H), 7.83 (d, J = 1.6 Hz, 1H), 7.55-7.42 (m, 3H), 7.27 (dd, J = 7.7, 1.3 Hz, 1H), 7.20 (d, J = 7.7 Hz, 1H), 5.87 (d, J = 5.6 Hz, 1H), 4.37 (s, 2H), 4.04-3.97 (m, 3H), 3.43-3.35 (m, 2H), 3.18 (d, J = 16.1 Hz, 1H), 2.81 (d, J = 16.0 Hz, 1H), 2.16-1.95 (m, 2H), 1.69 (d, J = 12.2 Hz, 1H), 1.47 (d, J = 12.9 Hz, 1H).	I-A1 I-C8
97		546.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.06-8.00 (m, 1H), 7.85-7.81 (m, 1H), 7.56-7.51 (m, 1H), 7.50-7.46 (m, 1H), 7.42-7.37 (m, 1H), 7.32-7.26 (m, 1H), 7.24-7.19 (m, 1H), 5.90-5.85 (m, 1H), 4.35 (s, 2H), 4.06-3.97 (m, 3H), 3.43 (s, 3H), 3.42-3.35 (m, 2H), 3.27-3.22 (m, 1H), 2.93-2.85 (m, 1H), 2.12-1.98 (m, 2H), 1.74-1.67 (m, 1H), 1.56-1.49 (m, 1H).	I-A1 I-C13

98		535.3	¹ H NMR (400 MHz, CD ₃ OD): δ 8.01 (s, 1H), 7.85-7.80 (m, 2H), 7.55 (d, J = 1.3 Hz, 1H), 7.44 (s, 1H), 7.27-7.14 (m, 2H), 6.24-6.17 (m, 1H), 4.03-3.94 (m, 3H), 3.54 (t, J = 6.7 Hz, 2H), 3.42-3.32 (m, 2H), 3.31 (s, 3H), 3.15 (d, J = 15.8 Hz, 1H), 3.07 (d, J = 6.7 Hz, 2H), 2.78 (d, J = 15.8 Hz, 1H), 2.12-1.93 (m, 2H), 1.74-1.65 (m, 1H), 1.52-1.44 (m, 1H).	I-A6 I-C14
100		513.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.01 (s, 1H), 7.85-7.80 (m, 2H), 7.56-7.54 (m, 1H), 7.51-7.44 (m, 1H), 7.33-7.28 (m, 1H), 7.26-7.20 (m, 1H), 6.23-6.19 (m, 1H), 4.30 (s, 2H), 4.04-3.95 (m, 3H), 3.43-3.32 (m, 5H), 3.22-3.14 (m, 1H), 2.90-2.80 (m, 1H), 2.12-1.97 (m, 2H), 1.73-1.66 (m, 1H), 1.54-1.46 (m, 1H).	I-A6 I-C7
117		527.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.87-7.80 (m, 1H), 7.72 (s, 1H), 7.50-7.41 (m, 2H), 7.31-7.25 (m, 1H), 7.23-7.18 (m, 1H), 6.20-6.15 (m, 1H), 4.34-4.25 (m, 2H), 4.02-3.92 (m, 3H), 3.43-3.31 (m, 5H), 3.21-3.15 (m, 1H), 2.84-2.76 (m, 1H), 2.55 (s, 3H), 2.13-2.05 (m, 1H), 2.02-1.94 (m, 1H), 1.74-1.62 (m, 1H), 1.50-1.42 (m, 1H).	I-A6 I-C7
128		545.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.04-8.02 (m, 1H), 7.83-7.82 (m, 1H), 7.60-7.58 (m, 1H), 7.55-7.53 (m, 1H), 7.49-7.47 (m, 1H), 7.45-7.42 (m, 1H), 7.32-7.28 (m, 1H), 5.88-5.86 (m, 1H), 4.05-3.98 (m, 3H), 3.44-3.35 (m, 2H), 3.26-3.20 (m, 1H), 2.91-2.82 (m, 1H), 2.15-2.07 (m, 1H), 2.04-1.97 (m, 1H), 1.74-1.67 (m, 1H), 1.51-1.43 (m, 1H).	I-A1 I-C16

132		512.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.25-8.17 (m, 1H), 8.06 (s, 1H), 7.87-7.80 (m, 1H), 7.58-7.52 (m, 1H), 7.47 (s, 1H), 7.31-7.25 (m, 1H), 7.23-7.17 (m, 1H), 6.83-6.76 (m, 1H), 4.30 (s, 2H), 4.09-3.91 (m, 3H), 3.44-3.33 (m, 5H), 3.22-3.15 (m, 1H), 2.85-2.79 (m, 1H), 2.52 (s, 3H), 2.13-2.05 (m, 1H), 2.03-1.96 (m, 1H), 1.75-1.65 (m, 1H), 1.53-1.42 (m, 1H).	I-A10 I-C7
147		547.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.37-8.26 (m, 1H), 8.21-8.11 (m, 1H), 7.53-7.49 (m, 1H), 7.45 (s, 1H), 7.31-7.27 (m, 1H), 7.23-7.20 (m, 1H), 5.94-5.90 (m, 1H), 5.20-5.09 (m, 2H), 4.30 (s, 2H), 3.95 (s, 1H), 3.67-3.57 (m, 2H), 3.45-3.36 (m, 3H), 3.25-3.20 (m, 1H), 2.86-2.80 (m, 1H), 2.07-1.99 (m, 1H), 1.97-1.88 (m, 1H), 1.72-1.65 (m, 1H), 1.48-1.42 (m, 1H).	I-A1 I-C7
148		547.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.35-8.27 (m, 1H), 8.19-8.13 (m, 1H), 7.53-7.48 (m, 1H), 7.45 (s, 1H), 7.31-7.25 (m, 1H), 7.24-7.18 (m, 1H), 5.97-5.88 (m, 1H), 5.22-5.06 (m, 2H), 4.30 (s, 2H), 3.95 (s, 1H), 3.67-3.55 (m, 2H), 3.45-3.37 (m, 3H), 3.24-3.18 (m, 1H), 2.86-2.78 (m, 1H), 2.06-1.98 (m, 1H), 1.96-1.88 (m, 1H), 1.73-1.64 (m, 1H), 1.47-1.40 (m, 1H).	I-A1 I-C7
156		502.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.92-7.88 (m, 1H), 7.76-7.73 (m, 1H), 7.5-7.47 (m, 1H), 7.47-7.44 (m, 1H), 7.34-7.30 (m, 1H), 7.24-7.20 (m, 1H), 7.13-7.10 (m, 1H), 5.71-5.66 (m, 1H), 4.58-4.50 (m, 2H), 3.96 (s, 1H), 3.56-3.45 (m, 2H), 3.41 (s, 1H), 3.24-3.19 (m, 1H), 2.87-2.80 (m, 1H), 2.03-1.94 (m, 1H), 1.92-1.83 (m, 1H), 1.71-1.64 (m, 1H), 1.48-1.41 (m, 1H).	I-A1 I-C3

171		531.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.02 (s, 1H), 7.88-7.78 (m, 2H), 7.59-7.52 (m, 1H), 7.32 (s, 1H), 7.07-6.99 (m, 1H), 6.25-6.19 (m, 1H), 4.31 (s, 2H), 4.06-3.95 (m, 3H), 3.47-3.32 (m, 5H), 3.26-3.17 (m, 1H), 2.87-2.76 (m, 1H), 2.17-1.96 (m, 2H), 1.78-1.67 (m, 1H), 1.55-1.43 (m, 1H).	I-A6 I-C18
172		559.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.05-8.03 (m, 1H), 7.85-7.83 (m, 1H), 7.61-7.57 (m, 1H), 7.56-7.54 (m, 1H), 7.51-7.48 (m, 1H), 7.45-7.42 (m, 1H), 7.32-7.29 (m, 1H), 5.90-5.87 (m, 1H), 4.06-3.99 (m, 3H), 3.42-3.36 (m, 2H), 3.26-3.21 (m, 1H), 2.89-2.84 (m, 1H), 2.81 (s, 3H), 2.16-2.08 (m, 1H), 2.05-1.98 (m, 1H), 1.75-1.68 (m, 1H), 1.51-1.45 (m, 1H).	I-A1 I-C20
175		530.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.24-8.20 (m, 1H), 8.08 (s, 1H), 7.88-7.82 (m, 1H), 7.58-7.54 (m, 1H), 7.33 (s, 1H), 7.08-7.01 (m, 1H), 6.84-6.78 (m, 1H), 4.31 (s, 2H), 4.10-3.98 (m, 3H), 3.47-3.37 (m, 5H), 3.26-3.22 (m, 1H), 2.89-2.79 (m, 1H), 2.54 (s, 3H), 2.16-2.07 (m, 1H), 2.03-2.00 (m, 1H), 1.79-1.69 (m, 1H), 1.55-1.47 (m, 1H).	I-A10 I-C18
176		520.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.91 (d, J = 2.5 Hz, 1H), 7.75 (s, 1H), 7.46 (d, J = 5.5 Hz, 1H), 7.33 (s, 1H), 7.11 (d, J = 2.5 Hz, 1H), 7.06 (d, J = 9.2 Hz, 1H), 5.69 (d, J = 5.6 Hz, 1H), 4.60-4.47 (m, 2H), 3.99 (s, 1H), 3.61-3.43 (m, 3H), 3.27-3.21 (m, 1H), 2.85-2.81 (m, 1H), 2.04-1.83 (m, 2H), 1.72-1.68 (m, 1H), 1.48-1.44 (m, 1H).	I-A1 I-C19





177		559.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.93-7.89 (m, 1H), 7.78-7.74 (m, 1H), 7.59-7.55 (m, 1H), 7.49-7.46 (m, 1H), 7.45-7.40 (m, 1H), 7.33-7.28 (m, 1H), 7.14-7.11 (m, 1H), 5.72-5.67 (m, 1H), 4.59-4.51 (m, 2H), 4.01-3.96 (m, 1H), 3.58-3.45 (m, 2H), 3.27-3.23 (m, 1H), 2.90-2.83 (m, 1H), 2.80 (s, 3H), 2.05-1.96 (m, 1H), 1.93-1.85 (m, 1H), 1.74-1.65 (m, 1H), 1.49-1.40 (m, 1H).	I-A1 I-C20
178		468.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.26-8.22 (m, 1H), 7.92-7.88 (m, 1H), 7.78-7.76 (m, 1H), 7.51-7.47 (m, 1H), 7.34-7.31 (m, 1H), 7.24-7.21 (m, 1H), 7.13-7.12 (m, 1H), 6.70-6.67 (m, 1H), 4.60-4.50 (m, 2H), 3.97 (s, 1H), 3.58-3.48 (m, 2H), 3.41 (s, 1H), 3.25-3.20 (m, 1H), 2.88-2.81 (m, 1H), 2.53 (s, 3H), 2.04-1.96 (m, 1H), 1.94-1.84 (m, 1H), 1.73-1.64 (m, 1H), 1.50-1.43 (m, 1H).	I-A10 I-C3
179		503.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.32 (s, 1H), 7.93 (s, 1H), 7.54-7.44 (m, 2H), 7.32 (dd, J = 7.7, 1.1 Hz, 1H), 7.23 (d, J = 7.8 Hz, 1H), 5.79 (d, J = 5.5 Hz, 1H), 5.51-5.22 (m, 1H), 3.94 (s, 1H), 3.67-3.51 (m, 2H), 3.41 (s, 1H), 3.24 (d, J = 16.1 Hz, 1H), 2.85 (d, J = 16.1 Hz, 1H), 2.04-1.79 (m, 2H), 1.67 (d, J = 13.6 Hz, 1H), 1.44 (d, J = 13.7 Hz, 1H).	I-A1 I-C3
180		516.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.18 (d, J = 0.7 Hz, 1H), 7.91 (d, J = 0.7 Hz, 1H), 7.56 (d, J = 5.5 Hz, 1H), 7.49 (s, 1H), 7.32 (dd, J = 7.7, 1.1 Hz, 1H), 7.22 (d, J = 7.7 Hz, 1H), 5.82 (d, J = 5.5 Hz, 1H), 4.70-4.56 (m, 2H), 3.96 (s, 1H), 3.60-3.47 (m, 2H), 3.41 (s, 1H), 3.26-3.17 (m, 1H), 2.84 (d, J = 16.0 Hz, 1H), 2.42 (s, 3H), 2.01-1.81 (m, 2H), 1.68 (d, J = 12.0 Hz, 1H), 1.45 (d, J = 13.3 Hz, 1H).	I-A1 I-C3





196		573.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.91 (d, J = 2.4 Hz, 1H), 7.75 (s, 1H), 7.61 (s, 1H), 7.52-7.43 (m, 2H), 7.33-7.31 (m, 1H), 7.13-7.11 (m, 1H), 5.69 (d, J = 5.5 Hz, 1H), 4.61-4.46 (m, 2H), 4.00 (s, 1H), 3.53-3.49 (m, 2H), 3.32 (s, 3H), 3.26-3.22 (m, 1H), 3.00 (s, 3H), 2.89-2.85 (m, 1H), 2.03-1.83 (m, 2H), 1.71-1.67 (m, 1H), 1.46-1.42 (m, 1H).	I-A1 I-C21
234		478.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.36 (s, 1H), 7.66 (d, J = 5.5 Hz, 1H), 7.36 (s, 1H), 7.22-7.12 (m, 2H), 6.04 (d, J = 5.5 Hz, 1H), 4.73-4.60 (m, 2H), 3.92 (s, 1H), 3.46-3.35 (m, 2H), 3.18 (d, J = 16.0 Hz, 1H), 2.79 (d, J = 15.9 Hz, 1H), 2.00 (s, 3H), 1.92-1.72 (m, 2H), 1.63 (d, J = 12.9 Hz, 1H), 1.41 (d, J = 13.4 Hz, 1H).	I-A1 I-C17
242		539.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.46-8.42 (m, 1H), 8.29-8.22 (m, 2H), 7.65-7.61 (m, 1H), 7.57-7.53 (m, 1H), 7.48-7.40 (m, 2H), 7.30-7.26 (m, 1H), 4.37-4.27 (m, 2H), 3.95 (s, 1H), 3.35-3.31 (m, 1H), 3.28-3.17 (m, 2H), 2.85-2.76 (m, 4H), 1.93-1.84 (m, 1H), 1.81-1.72 (m, 1H), 1.64-1.59 (m, 1H), 1.41-1.36 (m, 1H).	I-A20 I-C20
278		559.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.44 (s, 1H), 7.69-7.62 (m, 2H), 7.59-7.53 (m, 1H), 7.40 (d, J = 7.9 Hz, 1H), 6.12 (d, J = 5.5 Hz, 1H), 4.59-4.43 (m, 2H), 4.36 (s, 1H), 3.57-3.46 (m, 2H), 3.31-3.30 (m, 1H), 3.27-3.19 (m, 2H), 3.12 (d, J = 16.8 Hz, 1H), 1.99-1.86 (m, 2H), 1.74-1.63 (m, 2H), 1.15 (t, J = 7.3 Hz, 3H).	I-A1 I-C23






The compounds in the table below were prepared by following the steps 1 and 3 for preparing compound 1 from corresponding intermediates and reagents:





Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Intermediates
2		478.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.65-7.55 (m, 2H), 7.42-7.28 (m, 3H), 5.92 (d, J = 5.5 Hz, 1H), 4.33-4.22 (m, 2H), 3.95 (s, 1H), 3.42 (s, 1H), 3.24-3.09 (m, 3H), 2.82-2.76 (m, 1H), 1.85-1.67 (m, 2H), 1.59-1.55 (m, 1H), 1.40-1.33 (m, 1H).	I-A2 I-C1
86		522.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.71-7.58 (m, 2H), 7.48 (s, 1H), 7.32 (d, J = 7.6 Hz, 1H), 7.24 (d, J = 7.6 Hz, 1H), 5.99-5.92 (m, 1H), 4.45-4.20 (m, 4H), 3.95 (s, 1H), 3.45 (s, 3H), 3.31-3.15 (m, 3H), 2.87-2.76 (m, 1H), 1.95-1.70 (m, 2H), 1.61 (d, J = 13.2 Hz, 1H), 1.39 (d, J = 13.2 Hz, 1H).	I-A2 I-C7
95		480.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.66-7.59 (m, 2H), 7.52 (s, 1H), 7.37-7.33 (m, 1H), 7.27-7.22 (m, 1H), 5.98-5.93 (m, 1H), 4.39-4.29 (m, 2H), 3.95 (s, 1H), 3.33-3.13 (m, 4H), 2.87-2.78 (m, 1H), 1.95-1.86 (m, 1H), 1.82-1.72 (m, 1H), 1.66-1.59 (m, 1H), 1.43-1.38 (m, 1H).	I-A2 I-C3
112		489.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.88 (d, J = 5.2 Hz, 1H), 7.53-7.51 (m, 1H), 7.44 (s, 1H), 7.35-7.10 (m, 2H), 6.11 (d, J = 5.0 Hz, 1H), 4.35-4.20 (m, 4H), 3.91 (s, 1H), 3.39 (s, 3H), 3.22-3.04 (m, 1H).	I-A9 I-C7





				3H), 2.83-2.70 (m, 1H), 1.90-1.63 (m, 2H), 1.58-1.54 (m, 1H), 1.36-1.32 (m, 1H).	
126		445.2		¹ H NMR (400 MHz, CD ₃ OD): δ 7.97-7.89 (m, 1H), 7.63-7.47 (m, 2H), 7.35 (d, J = 7.8 Hz, 1H), 7.24 (d, J = 7.8 Hz, 1H), 6.22-6.12 (m, 1H), 4.38-4.24 (m, 2H), 3.95 (s, 1H), 3.44 (s, 1H), 3.29-3.16 (m, 3H), 2.87-2.76 (m, 1H), 1.93-1.84 (m, 1H), 1.81-1.73 (m, 1H), 1.65-1.59 (m, 1H), 1.44-1.39 (m, 1H).	I-A9 I-C3
129		521.1		¹ H NMR (400 MHz, CD ₃ OD): δ 7.61-7.55 (m, 3H), 7.45-7.41 (m, 1H), 7.30-7.26 (m, 1H), 5.93-5.89 (m, 1H), 4.34-4.23 (m, 2H), 3.94 (s, 1H), 3.26-3.15 (m, 3H), 2.85-2.77 (m, 1H), 1.91-1.82 (m, 1H), 1.78-1.69 (m, 1H), 1.61-1.55 (m, 1H), 1.37-1.32 (m, 1H).	I-A2 I-C16
131		492.2		¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.51 (m, 2H), 7.33 (s, 1H), 7.22-7.07 (m, 2H), 5.89 (d, J = 5.6 Hz, 1H), 4.31-4.19 (m, 2H), 3.86 (s, 1H), 3.23-3.06 (m, 3H), 2.72 (d, J = 16.0 Hz, 1H), 1.97 (s, 3H), 1.87-1.77 (m, 1H), 1.75-1.63 (m, 1H), 1.53 (d, J = 13.4 Hz, 1H), 1.33 (d, J = 13.4 Hz, 1H).	I-A2 I-C17
134		459.2		¹ H NMR (400 MHz, CD ₃ OD): δ 7.93-7.83 (m, 1H), 7.54 (s, 1H), 7.35 (s, 1H), 7.22-7.09 (m, 2H), 6.16-6.07 (m, 1H), 4.37-4.12 (m, 2H), 3.88 (s, 1H), 3.25-3.09 (m, 3H), 2.83-2.67 (m, 1H), 1.99 (s, 3H), 1.87-1.78 (m, 1H), 1.76-1.67 (m, 1H), 1.59-1.52 (m, 1H), 1.38-1.32 (m, 1H).	I-A9 I-C17





155		534.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃ =2/1): δ 8.28-8.22 (m, 2H), 7.58-7.56 (m, 1H), 7.52 (s, 1H), 7.42-7.36 (m, 1H), 7.27-7.21 (m, 1H), 5.95-5.89 (m, 1H), 4.34-4.26 (m, 2H), 3.95 (s, 1H), 3.28-3.23 (m, 4H), 3.20-3.14 (m, 1H), 2.84-2.77 (m, 1H), 1.91-1.82 (m, 1H), 1.79-1.71 (m, 1H), 1.65-1.59 (m, 1H), 1.43-1.36 (m, 1H), 1.19-1.11 (m, 3H).	I-A15 I-C23
187		549.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.56 (m, 2H), 7.44-7.40 (m, 1H), 7.30-7.26 (m, 1H), 5.89-5.87 (m, 1H), 3.97 (s, 1H), 3.72-3.64 (m, 2H), 3.19-3.14 (m, 1H), 3.13-3.06 (m, 2H), 2.82-2.76 (m, 4H), 2.37 (s, 3H), 2.03-1.95 (m, 1H), 1.92-1.83 (m, 1H), 1.63-1.58 (m, 1H), 1.41-1.35 (m, 1H).	I-A11 I-C20
188		564.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.78-7.75 (m, 1H), 7.58-7.55 (m, 1H), 7.44-7.40 (m, 1H), 7.30-7.26 (m, 1H), 6.20-6.17 (m, 1H), 3.98-3.96 (m, 4H), 3.73-3.66 (m, 2H), 3.20-3.04 (m, 3H), 2.83-2.76 (m, 4H), 2.38 (s, 3H), 2.04-1.97 (m, 1H), 1.94-1.83 (m, 1H), 1.63-1.58 (m, 1H), 1.40-1.35 (m, 1H).	I-A12 I-C20
189		548.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.04-8.01 (m, 1H), 7.59-7.55 (m, 1H), 7.44-7.40 (m, 1H), 7.30-7.26 (m, 1H), 6.51-6.48 (m, 1H), 3.97 (s, 1H), 3.75-3.67 (m, 2H), 3.20-3.15 (m, 1H), 3.15-3.06 (m, 2H), 2.84-2.76 (m, 4H), 2.59 (s, 3H), 2.38 (s, 3H), 2.04-1.97 (m, 1H), 1.94-1.84 (m, 1H), 1.64-1.58 (m, 1H), 1.41-1.35 (m, 1H).	I-A13 I-C20

203		563.3	¹ H NMR (400 MHz, CD ₃ OD): δ 7.61-7.58 (m, 1H), 7.58-7.55 (m, 1H), 7.45-7.40 (m, 1H), 7.30-7.25 (m, 1H), 5.88-5.84 (m, 1H), 3.96 (s, 1H), 3.71-3.62 (m, 2H), 3.19-3.13 (m, 1H), 3.12-3.04 (m, 2H), 3.00 (s, 3H), 2.81-2.75 (m, 1H), 2.36 (s, 3H), 2.02-1.94 (m, 1H), 1.91-1.82 (m, 1H), 1.63-1.56 (m, 1H), 1.39-1.32 (m, 1H).	I-A11 I-C21
204		578.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.77-7.73 (m, 1H), 7.61-7.58 (m, 1H), 7.46-7.41 (m, 1H), 7.31-7.26 (m, 1H), 6.19-6.15 (m, 1H), 3.98 (s, 1H), 3.94 (s, 3H), 3.73-3.62 (m, 2H), 3.19-3.14 (m, 1H), 3.14-3.05 (m, 2H), 3.00 (s, 3H), 2.84-2.77 (m, 1H), 2.36 (s, 3H), 2.03-1.93 (m, 1H), 1.93-1.82 (m, 1H), 1.63-1.56 (m, 1H), 1.41-1.34 (m, 1H).	I-A12 I-C21
205		478.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.94-7.91 (m, 1H), 7.62-7.60 (m, 1H), 7.47-7.45 (m, 1H), 7.31-7.27 (m, 1H), 7.21-7.16 (m, 1H), 6.03-6.00 (m, 1H), 4.58-4.51 (m, 2H), 3.89 (s, 1H), 3.38 (s, 1H), 3.23-3.11 (m, 3H), 2.80-2.73 (m, 1H), 1.84-1.75 (m, 1H), 1.71-1.63 (m, 1H), 1.55-1.49 (m, 1H), 1.34-1.29 (m, 1H).	I-A38 I-C3
213		445.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.93-7.91 (m, 1H), 7.91-7.88 (m, 1H), 7.48-7.44 (m, 1H), 7.31-7.27 (m, 1H), 7.20-7.17 (m, 1H), 6.23-6.20 (m, 1H), 4.56-4.49 (m, 2H), 3.89 (s, 1H), 3.37 (s, 1H), 3.21-3.11 (m, 3H), 2.79-2.73 (m, 1H), 1.82-1.74 (m, 1H), 1.71-1.63 (m, 1H), 1.54-1.49 (m, 1H), 1.33-1.31 (m, 1H).	I-A39 I-C3





215		492.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.94-7.91 (s, 1H), 7.62-7.59 (m, 1H), 7.37-7.33 (m, 1H), 7.20-7.12 (m, 2H), 6.03-6.00 (m, 1H), 4.57-4.50 (m, 2H), 3.90 (s, 1H), 3.24-3.15 (m, 2H), 3.14-3.09 (m, 1H), 2.80-2.72 (m, 1H), 1.98 (s, 3H), 1.81-1.73 (m, 1H), 1.71-1.63 (m, 1H), 1.54-1.48 (m, 1H), 1.35-1.30 (m, 1H).	I-A38 I-C17
216		516.3	¹ H NMR (400 MHz, CD ₃ OD): δ 7.90-7.85 (m, 1H), 7.57-7.54 (m, 1H), 7.42-7.38 (m, 1H), 7.28-7.24 (m, 1H), 6.13-6.09 (m, 1H), 3.98-3.94 (m, 1H), 3.69-3.59 (m, 2H), 3.17-3.04 (m, 3H), 2.80-2.75 (m, 4H), 2.35 (s, 3H), 2.01-1.93 (m, 1H), 1.90-1.82 (m, 1H), 1.61-1.56 (m, 1H), 1.39-1.34 (m, 1H).	I-A14 I-C20
219		563.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.59-7.54 (m, 2H), 7.43-7.39 (m, 1H), 7.28-7.24 (m, 1H), 5.88-5.84 (m, 1H), 3.96 (s, 1H), 3.71-3.62 (m, 2H), 3.28-3.24 (m, 2H), 3.17-3.05 (m, 3H), 2.83-2.75 (m, 1H), 2.36 (s, 3H), 2.02-1.93 (m, 1H), 1.91-1.83 (m, 1H), 1.62-1.56 (m, 1H), 1.40-1.34 (m, 1H), 1.17-1.12 (m, 3H).	I-A11 I-C23
225		575.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.59-7.51 (m, 2H), 7.41-7.37 (m, 1H), 7.27-7.22 (m, 1H), 5.88-5.83 (m, 1H), 3.94 (s, 1H), 3.72-3.61 (m, 2H), 3.18-3.03 (m, 3H), 2.81-2.71 (m, 2H), 2.35 (s, 3H), 2.01-1.92 (m, 1H), 1.90-1.82 (m, 1H), 1.61-1.55 (m, 1H), 1.38-1.32 (m, 1H), 0.82-0.70 (m, 2H), 0.60-0.49 (m, 2H).	I-A11 I-C25
226		549.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃ =2/1): δ 7.57-7.55 (m, 1H), 7.53 (s, 1H), 7.51 (s, 1H), 7.41-7.36 (m, 1H), 7.25-7.20 (m, 1H), 5.97-5.90 (m, 1H), 4.26-4.19 (m, 2H), 3.93 (s, 1H), 3.28-3.23 (m, 2H), 3.22-3.11 (m, 3H), 2.82-2.75 (m, 1H), 1.87-1.78 (m, 1H),	I-A2 I-C23






				1.76-1.67 (m, 1H), 1.61-1.55 (m, 1H), 1.40-1.33 (m, 1H), 1.17-1.11 (m, 3H).	
227		520.1		¹ H NMR (400 MHz, CD ₃ OD): δ 8.32-8.30 (m, 1H), 8.27-8.23 (m, 1H), 7.59-8.57 (m, 1H), 7.54 (s, 1H), 7.43-7.38 (m, 1H), 7.29-7.25 (m, 1H), 5.93-5.91 (m, 1H), 4.37-4.30 (m, 2H), 3.95 (s, 1H), 3.28-3.17 (m, 3H), 2.85-2.78 (m, 4H), 1.92-1.84 (m, 1H), 1.80-1.72 (m, 1H), 1.64-1.58 (m, 1H), 1.40-1.34 (m, 1H).	I-A15 I-C20
229		535.2		¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.52 (m, 3H), 7.41-7.39 (m, 1H), 7.27-7.25 (m, 1H), 5.91 (d, J = 5.5 Hz, 1H), 4.26 (br, 2H), 3.93 (s, 1H), 3.35-3.33 (m, 1H), 3.24-3.13 (m, 3H), 2.78 (s, 3H), 1.90-1.70 (m, 2H), 1.59-1.55 (m, 1H), 1.36-1.32 (m, 1H).	I-A2 I-C20
231		505.2		¹ H NMR (400 MHz, CD ₃ OD): δ 8.31-8.29 (m, 1H), 8.27-8.25 (m, 1H), 8.17-8.13 (m, 1H), 7.57-7.55 (m, 1H), 7.44-7.40 (m, 1H), 7.37-7.34 (m, 1H), 7.30-7.26 (m, 1H), 7.25-7.22 (m, 1H), 4.36-4.29 (m, 2H), 3.96 (s, 1H), 3.29-3.24 (m, 2H), 3.24-3.18 (m, 1H), 2.85-2.79 (m, 4H), 1.93-1.85 (m, 1H), 1.81-1.73 (m, 1H), 1.65-1.59 (m, 1H), 1.41-1.35 (m, 1H).	I-A16 I-C20
232		534.2		¹ H NMR (400 MHz, CD ₃ OD): δ 8.26-8.23 (m, 1H), 7.62-7.60 (m, 1H), 7.58-7.55 (m, 1H), 7.44-7.39 (m, 1H), 7.30-7.26 (m, 1H), 6.00-5.98 (m, 1H), 3.98 (s, 1H), 3.75-3.67 (m, 2H), 3.21-3.11 (m, 3H), 2.83-2.77 (m, 4H), 2.55 (s, 3H), 2.05-1.97 (m, 1H), 1.95-1.86 (m, 1H), 1.67-1.60 (m, 1H), 1.44-1.37 (m, 1H).	I-A17 I-C20





238		534.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.13-8.08 (m, 1H), 7.59 (s, 1H), 7.56 (s, 1H), 7.44-7.39 (m, 1H), 7.31-7.25 (m, 1H), 7.24-7.19 (m, 1H), 7.15-7.09 (m, 1H), 4.33-4.24 (m, 2H), 3.95 (s, 1H), 3.34 (s, 1H), 3.27-3.16 (m, 4H), 2.87-2.76 (m, 1H), 1.93-1.81 (m, 1H), 1.80-1.69 (m, 1H), 1.62-1.56 (m, 1H), 1.38-1.34 (m, 1H), 1.21-1.12 (m, 3H).	I-A40 I-C23
241		492.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.58 (d, J = 5.5 Hz, 1H), 7.48 (s, 1H), 7.30 (d, J = 7.7 Hz, 1H), 7.19 (d, J = 7.7 Hz, 1H), 5.88 (d, J = 5.5 Hz, 1H), 3.94 (s, 1H), 3.68-3.64 (m, 2H), 3.39 (s, 1H), 3.15-3.02 (m, 3H), 2.77-2.73 (m, 1H), 2.37 (s, 3H), 2.01-1.82 (m, 2H), 1.60-1.56 (m, 1H), 1.40-1.36 (m, 1H).	I-A11 I-C3
246		540.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.68 (dd, J = 4.7, 1.4 Hz, 1H), 8.42 (s, 1H), 8.19 (d, J = 7.4 Hz, 1H), 7.67 (dd, J = 8.0, 4.7 Hz, 1H), 7.55 (s, 1H), 7.41 (dd, J = 7.7, 1.3 Hz, 1H), 7.27 (d, J = 7.8 Hz, 1H), 4.29-4.16 (m, 2H), 3.94 (s, 1H), 3.30-3.16 (m, 3H), 2.84-2.77 (m, 4H), 1.98-1.72 (m, 2H), 1.61 (d, J = 13.5 Hz, 1H), 1.36 (d, J = 13.6 Hz, 1H).	I-A21 I-C20
247		549.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74 (s, 1H), 7.64-7.55 (m, 2H), 7.47-7.41 (m, 1H), 7.34-7.27 (m, 1H), 5.89 (s, 1H), 4.37-4.24 (m, 2H), 3.97 (s, 1H), 3.30-3.12 (m, 5H), 2.91-2.77 (m, 1H), 1.96-1.81 (m, 1H), 1.81-1.69 (m, 1H), 1.64-1.57 (m, 1H), 1.40-1.34 (m, 1H), 1.20-1.14 (m, 3H).	I-A22 I-C23






248		520.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃): δ 8.11-8.05 (m, 1H), 7.55 (s, 1H), 7.50 (s, 1H), 7.42-7.37 (m, 1H), 7.25-7.21 (m, 1H), 7.17-7.10 (m, 2H), 4.26-4.19 (m, 2H), 3.95 (s, 1H), 3.24-3.10 (m, 3H), 2.89-2.74 (m, 4H), 1.87-1.78 (m, 1H), 1.75-1.67 (m, 1H), 1.63-1.57 (m, 1H), 1.41-1.34 (m, 1H).	I-A40 I-C20
249		535.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.54 (s, 1H), 7.73 (d, J = 5.5 Hz, 1H), 7.57 (s, 1H), 7.43 (d, J = 7.7 Hz, 1H), 7.29 (d, J = 7.8 Hz, 1H), 6.50 (d, J = 5.5 Hz, 1H), 4.39-4.28 (m, 2H), 3.97 (s, 1H), 3.41-3.32 (m, 3H), 3.28-3.19 (m, 2H), 2.84 (d, J = 16.2 Hz, 1H), 1.98-1.75 (m, 2H), 1.66 (d, J = 13.7 Hz, 1H), 1.40 (d, J = 13.9 Hz, 1H), 1.17 (t, J = 7.3 Hz, 3H).	I-A23 I-C23
250		521.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.53 (s, 1H), 7.73 (d, J = 5.5 Hz, 1H), 7.56 (s, 1H), 7.45-7.39 (m, 1H), 7.29 (d, J = 7.8 Hz, 1H), 6.49 (d, J = 5.5 Hz, 1H), 4.38-4.28 (m, 2H), 3.96 (s, 1H), 3.41-3.32 (m, 2H), 3.23 (d, J = 16.3 Hz, 1H), 2.87-2.79 (m, 4H), 1.97-1.73 (m, 2H), 1.65 (d, J = 13.2 Hz, 1H), 1.40 (d, J = 13.5 Hz, 1H).	I-A23 I-C20
252		463.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.32 (d, J = 1.3 Hz, 1H), 8.27 (d, J = 1.3 Hz, 1H), 7.59 (d, J = 5.5 Hz, 1H), 7.48 (s, 1H), 7.32 (dd, J = 7.7, 1.3 Hz, 1H), 7.21 (d, J = 7.7 Hz, 1H), 5.97-5.90 (m, 1H), 4.4-4.29 (m, 2H), 3.94 (s, 1H), 3.41 (s, 1H), 3.36-3.30 (m, 2H), 3.20-3.16 (m, 1H), 2.82-2.78 (m, 1H), 1.92-1.70 (m, 2H), 1.64-1.58 (m, 1H), 1.44-1.37 (m, 1H).	I-A15 I-C3





253		569.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.62-7.59 (m, 1H), 7.58-7.55 (m, 1H), 7.45-7.41 (m, 1H), 7.32-7.27 (m, 1H), 5.79 (s, 1H), 4.36-4.24 (m, 2H), 3.96 (s, 1H), 3.28-3.18 (m, 3H), 2.88-2.77 (m, 4H), 1.93-1.83 (m, 1H), 1.79-1.69 (m, 1H), 1.63-1.57 (m, 1H), 1.38-1.33 (m, 1H).	I-A24 I-C20
254		554.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.40-8.34 (m, 1H), 8.33-8.26 (m, 1H), 7.61-7.54 (m, 1H), 7.47-7.41 (m, 1H), 7.34-7.27 (m, 1H), 5.79 (s, 1H), 4.43-4.33 (m, 2H), 3.97 (s, 1H), 3.38-3.20 (m, 3H), 2.90-2.76 (m, 4H), 1.97-1.86 (m, 1H), 1.82-1.73 (m, 1H), 1.69-1.61 (m, 1H), 1.44-1.36 (m, 1H).	I-A25 I-C20
255		583.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.61-7.59 (m, 1H), 7.59-7.55 (m, 1H), 7.46-7.41 (m, 1H), 7.32-7.27 (m, 1H), 5.80 (s, 1H), 4.34-4.26 (m, 2H), 3.95 (s, 1H), 3.30-3.16 (m, 5H), 2.86-2.79 (m, 1H), 1.92-1.84 (m, 1H), 1.79-1.70 (m, 1H), 1.63-1.57 (m, 1H), 1.38-1.32 (m, 1H), 1.20-1.14 (m, 3H).	I-A24 I-C23
256		568.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.37-8.35 (m, 1H), 8.31-8.26 (m, 1H), 7.59-7.55 (m, 1H), 7.46-7.41 (m, 1H), 7.32-7.27 (m, 1H), 5.78 (s, 1H), 4.41-4.33 (m, 2H), 3.96 (s, 1H), 3.37-3.20 (m, 5H), 2.88-2.80 (m, 1H), 1.96-1.86 (m, 1H), 1.81-1.72 (m, 1H), 1.68-1.61 (m, 1H), 1.42-1.36 (m, 1H), 1.20-1.13 (m, 3H).	I-A25 I-C23
257		518.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.28 (dd, J = 9.4, 1.2 Hz, 2H), 7.57 (s, 1H), 7.50 (d, J = 5.6 Hz, 1H), 7.43 (d, J = 7.7 Hz, 1H), 7.29 (d, J = 7.7 Hz, 1H), 6.06 (t, J = 5.3 Hz, 1H), 4.38-4.30 (m, 2H), 3.96 (s, 1H), 3.35-3.31 (m, 2H), 3.28-3.19 (m, 3H), 2.83 (d, J	I-A26 I-C23





				δ 16.3 Hz, 1H), 1.95-1.73 (m, 2H), 1.63 (d, J = 12.7 Hz, 1H), 1.38 (d, J = 13.2 Hz, 1H), 1.17 (t, J = 7.3 Hz, 3H).	
259		550.2		^1H NMR (400 MHz, CD_3OD): δ 8.31 (s, 1H), 7.65 (d, J = 5.3 Hz, 2H), 7.51-7.41 (m, 1H), 6.91-6.83 (m, 1H), 6.03 (d, J = 5.3 Hz, 1H), 4.20 (s, 1H), 3.91-3.72 (m, 2H), 3.47-3.39 (m, 2H), 3.33-3.26 (m, 2H), 2.60 (s, 3H), 2.22-2.11 (m, 1H), 2.09-1.98 (m, 2H), 1.94-1.83 (m, 1H), 1.24-1.15 (m, 3H).	I-A17 I-C28
260		501.2		^1H NMR (400 MHz, CD_3OD): δ 8.31-8.26 (m, 2H), 7.92-7.89 (m, 1H), 7.58-7.55 (m, 1H), 7.45-7.41 (m, 1H), 7.31-7.27 (m, 1H), 6.21-6.18 (m, 1H), 4.38-4.31 (m, 2H), 3.98-3.95 (m, 1H), 3.30-3.18 (m, 5H), 2.86-2.81 (m, 1H), 1.94-1.86 (m, 1H), 1.81-1.73 (m, 1H), 1.66-1.60 (m, 1H), 1.42-1.36 (m, 1H), 1.18-1.14 (m, 3H).	I-A27 I-C23
261		515.2		^1H NMR (400 MHz, CD_3OD): δ 8.30-8.26 (m, 1H), 7.93-7.90 (m, 1H), 7.59-7.56 (m, 1H), 7.44-7.41 (m, 1H), 7.30-7.27 (m, 1H), 6.26-6.24 (m, 1H), 3.99-3.97 (m, 1H), 3.73-3.66 (m, 2H), 3.29-3.11 (m, 5H), 2.83-2.78 (m, 1H), 2.55 (s, 3H), 2.06-1.99 (m, 1H), 1.95-1.87 (m, 1H), 1.67-1.61 (m, 1H), 1.44-1.38 (m, 1H), 1.18-1.14 (m, 3H).	I-A18 I-C23
262		487.2		^1H NMR (400 MHz, CD_3OD): δ 8.31-8.26 (m, 2H), 7.92-7.88 (m, 1H), 7.58-7.54 (m, 1H), 7.45-7.40 (m, 1H), 7.31-7.27 (m, 1H), 6.22-6.18 (m, 1H), 4.38-4.31 (m, 2H), 3.98-3.94 (m, 1H), 3.30-3.18 (m, 3H), 2.86-2.79 (m, 4H), 1.94-1.86 (m, 1H), 1.81-1.73 (m, 1H), 1.66-1.60 (m, 1H), 1.42-1.36 (m, 1H).	I-A27 I-C20



264		519.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.56 (s, 2H), 7.47 (d, J = 5.6 Hz, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.28 (d, J = 7.8 Hz, 1H), 5.96 (t, J = 5.3 Hz, 1H), 4.35-4.23 (m, 2H), 3.94 (s, 1H), 3.26-3.16 (m, 3H), 2.84-2.78 (m, 4H), 1.91-1.69 (m, 2H), 1.59 (d, J = 13.3 Hz, 1H), 1.34 (d, J = 13.5 Hz, 1H).	I-A28 I-C20
265		504.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.27 (dd, J = 9.5, 1.3 Hz, 2H), 7.56 (s, 1H), 7.49 (d, J = 5.5 Hz, 1H), 7.42 (d, J = 7.6 Hz, 1H), 7.28 (d, J = 7.7 Hz, 1H), 6.05 (t, J = 5.3 Hz, 1H), 4.38-4.28 (m, 2H), 3.95 (s, 1H), 3.30-3.18 (m, 3H), 2.86-2.76 (m, 4H), 1.95-1.71 (m, 2H), 1.62 (d, J = 12.9 Hz, 1H), 1.38 (d, J = 13.2 Hz, 1H).	I-A29 I-C20
266		536.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.20 (s, 1H), 7.60-7.49 (m, 2H), 7.39-7.33 (m, 1H), 6.80-6.75 (m, 1H), 5.95-5.90 (m, 1H), 4.11 (s, 1H), 3.79-3.61 (m, 2H), 3.35-3.31 (m, 1H), 3.26-3.24 (m, 1H), 2.74 (s, 3H), 2.50 (s, 3H), 2.12-2.02 (m, 1H), 1.98-1.88 (m, 2H), 1.83-1.75 (m, 1H).	I-A17 I-C29
269		552.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.19-8.13 (m, 1H), 8.12-8.06 (m, 1H), 7.78-7.71 (m, 1H), 7.54 (s, 1H), 7.43-7.38 (m, 1H), 7.29-7.23 (m, 1H), 4.29-4.19 (m, 2H), 3.93 (s, 1H), 3.28-3.15 (m, 5H), 2.84-2.75 (m, 1H), 1.90-1.81 (m, 1H), 1.77-1.69 (m, 1H), 1.62-1.55 (m, 1H), 1.37-1.32 (m, 1H), 1.20-1.09 (m, 3H).	I-A30 I-C23
270		506.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.43-8.25 (m, 2H), 7.66-7.57 (m, 2H), 7.51-7.44 (m, 1H), 7.37-7.29 (m, 1H), 6.01-5.93 (m, 1H), 4.46-4.32 (m, 2H), 4.01 (s, 1H), 3.40-3.38 (m, 1H), 3.33-3.22 (m, 2H), 2.93-2.82 (m, 1H), 1.99-1.89 (m, 1H), 1.87-1.74 (m, 1H).	I-A15 I-C16

271		520.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.24 (s, 1H), 7.63-7.55 (m, 2H), 7.46-7.36 (m, 1H), 7.30-7.25 (m, 1H), 5.99-5.94 (m, 1H), 3.97 (s, 1H), 3.77-3.64 (m, 2H), 3.21-3.10 (m, 3H), 2.84-2.76 (m, 1H), 2.54 (s, 3H), 2.07-1.82 (m, 2H), 1.63 (d, J = 13.2 Hz, 1H), 1.40 (d, J = 13.2 Hz, 1H).	I-A17 I-C16
272		559.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.53 (s, 1H), 7.65 (d, J = 5.5 Hz, 1H), 7.55 (s, 1H), 7.42 (d, J = 7.6 Hz, 1H), 7.28 (d, J = 7.7 Hz, 1H), 6.02 (d, J = 5.5 Hz, 1H), 4.42-4.31 (m, 2H), 3.96 (s, 1H), 3.40-3.32 (m, 2H), 3.28-3.18 (m, 3H), 2.83 (d, J = 16.7 Hz, 1H), 1.96-1.86 (m, 1H), 1.81-1.72 (m, 1H), 1.65 (d, J = 13.9 Hz, 1H), 1.39 (d, J = 14.5 Hz, 1H), 1.15 (t, J = 7.3 Hz, 3H).	I-A31 I-C23
273		552.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.33 (d, J = 1.3 Hz, 1H), 8.26 (d, J = 1.2 Hz, 1H), 7.55 (s, 1H), 7.44-7.38 (m, 1H), 7.27 (d, J = 7.8 Hz, 1H), 5.43 (d, J = 1.4 Hz, 1H), 4.42-4.29 (m, 2H), 3.94 (s, 1H), 3.37-3.30 (m, 2H), 3.28-3.18 (m, 3H), 2.82 (d, J = 16.4 Hz, 1H), 1.95-1.85 (m, 1H), 1.81-1.71 (m, 1H), 1.62 (d, J = 11.7 Hz, 1H), 1.37 (d, J = 11.7 Hz, 1H), 1.15 (t, J = 7.3 Hz, 3H).	I-A32 I-C23
274		519.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.57 (s, 1H), 7.48-7.38 (m, 2H), 7.26 (d, J = 7.6 Hz, 1H), 5.93 (t, J = 5.3 Hz, 1H), 3.95 (s, 1H), 3.72-3.61 (m, 2H), 3.18-3.03 (m, 3H), 2.77 (d, J = 16.5 Hz, 1H), 2.36 (s, 3H), 2.02-1.82 (m, 2H), 1.59 (d, J = 12.2 Hz, 1H), 1.36 (d, J = 11.8 Hz, 1H).	I-A33 I-C16

275		505.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.59-7.53 (m, 2H), 7.48-7.39 (m, 2H), 7.27 (d, J = 7.8 Hz, 1H), 5.95 (t, J = 5.3 Hz, 1H), 4.32-4.21 (m, 2H), 3.94 (s, 1H), 3.26-3.13 (m, 3H), 2.80 (d, J = 16.3 Hz, 1H), 1.93-1.67 (m, 2H), 1.57 (d, J = 12.9 Hz, 1H), 1.34 (d, J = 13.5 Hz, 1H).	I-A34 I-C16
276		535.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.59-7.55 (m, 2H), 7.43-7.40 (m, 1H), 7.28-7.25 (m, 1H), 5.88-5.85 (m, 1H), 3.95 (s, 1H), 3.71-3.63 (m, 2H), 3.18-3.04 (m, 3H), 2.81-2.74 (m, 1H), 2.36 (s, 3H), 2.02-1.94 (m, 1H), 1.91-1.83 (m, 1H), 1.62-1.56 (m, 1H), 1.39-1.33 (m, 1H).	I-A11 I-C16
277		520.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃ =3/1): δ 8.11-8.08 (m, 1H), 7.62-7.55 (m, 2H), 7.47-7.43 (m, 1H), 7.30-7.27 (m, 1H), 5.84-5.81 (m, 1H), 4.40-4.33 (m, 2H), 3.99 (s, 1H), 3.32-3.18 (m, 3H), 2.87-2.82 (m, 1H), 2.46 (s, 3H), 1.94-1.86 (m, 1H), 1.82-1.74 (m, 1H), 1.69-1.62 (m, 1H), 1.46-1.40 (m, 1H).	I-A35 I-C16
280		554.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.14-8.12 (m, 1H), 7.58-7.53 (m, 1H), 7.44-7.40 (m, 1H), 7.30-7.26 (m, 1H), 5.65 (s, 1H), 4.40-4.32 (m, 2H), 3.94 (s, 1H), 3.28-3.18 (m, 3H), 2.85-2.78 (m, 1H), 2.44 (s, 3H), 1.92-1.84 (m, 1H), 1.78-1.70 (m, 1H), 1.65-1.59 (m, 1H), 1.39-1.33 (m, 1H).	I-A36 I-C16
281		487.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.08 (s, 1H), 7.92-7.84 (m, 1H), 7.56 (s, 1H), 7.46-7.39 (m, 1H), 7.32-7.24 (m, 1H), 6.17-6.09 (m, 1H), 4.41-4.28 (m, 2H), 3.95 (s, 1H), 3.28-3.15 (m, 3H), 2.88-2.76 (m, 1H), 2.44 (s, 3H), 1.94-1.82 (m, 1H), 1.81-1.69 (m, 1H), 1.66-	I-A37 I-C16


290		548.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.27-8.23 (m, 1H), 7.63-7.56 (m, 2H), 7.47-7.39 (m, 1H), 7.32-7.26 (m, 1H), 6.00-5.96 (m, 1H), 3.98 (s, 1H), 3.76-3.67 (m, 2H), 3.29-3.25 (m, 2H), 3.21-3.11 (m, 3H), 2.85-2.76 (m, 1H), 2.55 (s, 3H), 2.06-1.97 (m, 1H), 1.96-1.85 (m, 1H), 1.68-1.59 (m, 1H), 1.45-1.36 (m, 1H), 1.26-1.14 (m, 3H).	I-A17 I-C23
292		522.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃ =2/1): δ 8.35-8.33 (m, 1H), 8.33-8.29 (m, 1H), 7.65-7.57 (m, 2H), 7.48-7.42 (m, 1H), 6.88-6.82 (m, 1H), 5.99-5.95 (m, 1H), 4.52-4.46 (m, 1H), 4.42-4.35 (m, 1H), 4.15 (s, 1H), 3.56-3.48 (m, 2H), 2.83 (s, 3H), 2.04-1.97 (m, 2H), 1.94-1.83 (m, 2H).	I-A15 I-C29
293		536.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃): δ 8.35-8.24 (m, 2H), 7.62-7.54 (m, 2H), 7.43 (dd, J = 8.3, 1.6 Hz, 1H), 6.82 (d, J = 8.3 Hz, 1H), 5.94 (d, J = 5.6 Hz, 1H), 4.50-4.42 (m, 1H), 4.40-4.32 (m, 1H), 4.13 (s, 1H), 3.53-3.44 (m, 2H), 3.29-3.24 (m, 2H), 2.02-1.94 (m, 2H), 1.90-1.81 (m, 2H), 1.15 (t, J = 7.3 Hz, 3H).	I-A15 I-C28
296		546.6	¹ H NMR (400 MHz, CD ₃ OD): δ 8.33-8.31 (m, 1H), 8.28-8.26 (m, 1H), 7.61-7.53 (m, 2H), 7.43-7.41 (m, 1H), 7.29-7.27 (m, 1H), 5.93 (d, J = 5.6 Hz, 1H), 4.42-4.27 (m, 2H), 3.96 (s, 1H), 3.30-3.15 (m, 3H), 2.89-2.68 (m, 2H), 1.93-1.85 (m, 1H), 1.81-1.75 (m, 1H), 1.66-1.60 (m, 1H), 1.42-1.36 (m, 1H), 0.80-0.74 (m, 2H), 0.59-0.53 (m, 2H).	I-A15 I-C25

297		564.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.38-8.26 (m, 2H), 7.69-7.57 (m, 2H), 7.49-7.43 (m, 1H), 7.35-7.27 (m, 1H), 5.99-5.90 (m, 1H), 4.43-4.32 (m, 2H), 3.99 (s, 1H), 3.55-3.51 (m, 2H), 3.49-3.45 (m, 2H), 3.42-3.36 (m, 4H), 3.32-3.21 (m, 2H), 2.91-2.81 (m, 1H), 1.97-1.87 (m, 1H), 1.85-1.75 (m, 1H), 1.66 (d, J = 13.3 Hz, 1H), 1.41 (d, J = 13.3 Hz, 1H).	I-A15 I-C33
298		548.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.32-8.24 (m, 2H), 7.57 (s, 1H), 7.49 (d, J = 5.5 Hz, 1H), 7.44 (d, J = 9.2 Hz, 1H), 7.29 (d, J = 7.7 Hz, 1H), 6.05 (t, J = 5.3 Hz, 1H), 4.39-4.29 (m, 2H), 3.96 (s, 1H), 3.51-3.47 (m, 2H), 3.46-3.41 (m, 2H), 3.36 (s, 3H), 3.29-3.26 (m, 2H), 3.22 (d, J = 16.4 Hz, 1H), 2.83 (d, J = 16.3 Hz, 1H), 1.93-1.71 (m, 2H), 1.63 (d, J = 13.6 Hz, 1H), 1.38 (d, J = 13.5 Hz, 1H).	I-A26 I-C33
300		521.2	¹ H NMR (400 MHz, CDCl ₃): δ 8.35-8.09 (m, 2H), 7.81-7.63 (m, 1H), 7.55 (s, 1H), 7.50-7.38 (m, 1H), 7.24-7.12 (m, 1H), 6.17-6.01 (m, 1H), 4.93-4.76 (m, 2H), 4.30-4.17 (m, 2H), 4.04-3.95 (m, 1H), 3.83 (s, 3H), 3.34-3.21 (m, 2H), 3.16-3.05 (m, 1H), 2.83-2.65 (m, 1H), 1.94-1.78 (m, 2H), 1.68-1.61 (m, 1H), 1.42-1.30 (m, 3H).	I-A15 I-C34
301		578.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.26 (dd, J = 16.9, 1.0 Hz, 2H), 7.64-7.52 (m, 2H), 7.42 (t, J = 6.2 Hz, 1H), 7.27 (d, J = 7.6 Hz, 1H), 5.98 (dd, J = 5.5, 1.0 Hz, 1H), 4.36-4.22 (m, 2H), 3.96 (s, 1H), 3.86 (t, J = 5.4 Hz, 1H), 3.66-3.60 (m, 2H), 3.59-3.56 (m, 1H), 3.38-3.31 (m, 5H), 3.29-3.26 (m, 2H), 3.19 (d, J = 16.2 Hz, 1H), 3.02-3.01 (m, 1H), 2.81 (d, J = 16.3 Hz, 1H), 1.93-1.75 (m, 2H), 1.62 (d, J = 12.9 Hz, 1H), 1.39 (d, J = 12.9 Hz, 1H).	I-A15 I-C35

302		534.2	¹H NMR (400 MHz, CD₃OD/CDCl₃=1/1): δ 8.23-8.19 (m, 2H), 7.58 (d, J = 5.6 Hz, 1H), 7.52 (s, 1H), 7.40 (d, J = 7.7 Hz, 1H), 7.23 (d, J = 7.7 Hz, 1H), 5.98 (d, J = 5.6 Hz, 1H), 4.25-4.21 (m, 2H), 3.97 (s, 1H), 3.35-3.30 (m, 1H), 3.28-3.23 (m, 4H), 3.16-3.10 (m, 1H), 2.99 (s, 3H), 2.83-2.77 (m, 1H), 1.89-1.71 (m, 2H), 1.65-1.59 (m, 1H), 1.46-1.40 (m, 1H).	I-A15 I-C21
303		546.2	¹H NMR (400 MHz, CD₃OD/CDCl₃=1/1): δ 8.29-8.22 (m, 2H), 7.64-7.61 (m, 1H), 7.55-7.52 (m, 1H), 7.44-7.41 (m, 1H), 7.29-7.25 (m, 1H), 6.04-6.00 (m, 1H), 4.38-4.30 (m, 4H), 4.14-4.09 (m, 2H), 4.01-3.98 (m, 1H), 3.37-3.28 (m, 2H), 3.21-3.13 (m, 1H), 2.88-2.79 (m, 1H), 2.45-2.34 (m, 2H), 1.94-1.78 (m, 2H), 1.70-1.64 (m, 1H), 1.49-1.43 (m, 1H).	I-A15 I-C36

Compound 4

(S)-6-amino-2-(1-amino-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-methyl-5-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)pyrimidin-4(3*H*)-one



4

Step 1: (R)-*N*-((S)-1'-(4-amino-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide

6-amino-2-hydroxy-3-methylpyrimidin-4(3*H*)-one (170 mg, 1.20 mmol), BOP (1.06 g, 2.40 mmol) and DBU (366 mg, 2.40 mmol) were placed in anhydrous acetonitrile (10 mL), and the reaction was stirred at room temperature for 30 minutes. Intermediate I-C6 (242 mg, 0.60 mmol) was added thereto, and the reaction was stirred at room temperature for 16 hours. The reaction solution was purified with silica gel column chromatography (water/methanol) to give the target product (95 mg, yield 15%). $[M+H]^+$ 526.3

Step 2: (R)-*N*-((S)-1'-(4-amino-5-iodo-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide

Under nitrogen, (R)-*N*-((S)-1'-(4-amino-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide (95 mg, 0.18 mmol) and NIS (45 mg, 0.20 mmol) were placed in *N,N*-dimethylformamide (2 mL). The reaction was stirred at room temperature for 1 hour, and the reaction solution was purified with silica gel column chromatography (water/methanol) to give the target product (70 mg, yield 60%). $[M+H]^+$ 652.3





Step 3: (R)-N-((S)-1'-(4-amino-1-methyl-5-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide

Under nitrogen, (R)-N-((S)-1'-(4-amino-5-iodo-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide (70 mg, 0.11 mmol), 1-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrolo[2,3-*b*]pyridine (36 mg, 0.14 mmol), Pd(PPh₃)₄ (12 mg, 0.01 mmol) and potassium carbonate (45 mg, 0.32 mmol) were placed in 1,4-dioxane (3 mL) and water (0.3 mL). The reaction was reacted at 100°C for 1 hour, and the reaction solution was purified with silica gel column chromatography (water/methanol) to give the target product (10 mg, yield 14%). [M+H]⁺ 656.3

Step 4: (S)-6-amino-2-(1-amino-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-methyl-5-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)pyrimidin-4(3*H*)-one


(R)-N-((S)-1'-(4-amino-1-methyl-5-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1-yl)-2-methylpropane-2-sulfinamide (10 mg, 0.015 mmol) was dissolved in 2 M hydrogen chloride/methanol solution. The reaction was stirred at room temperature for 3 minutes. Under ice bath cooling, the reaction solution was diluted with dichloromethane (15 mL) and adjusted with aqueous ammonia to a pH value of 8. The organic phase was collected and concentrated in vacuum under reduced pressure. The resulting residue and potassium carbonate (10 mg, 0.072 mmol) were placed in methanol (0.5 mL). The mixture was stirred at room temperature for 10 minutes and purified with thin layer chromatography (dichloromethane/methanol = 15/1) to give the target product (5.5 mg, yield 75%). [M+H]⁺ 480.2. ¹H NMR (400 MHz, CD₃OD): δ 8.26-8.20 (m, 1H), 7.82-7.75 (m, 1H), 7.37-7.30 (m, 4H), 7.11-7.07 (m, 1H), 3.99-3.96 (m, 1H), 3.90 (s, 3H), 3.54-3.49 (m, 2H), 3.47 (s, 3H), 3.41-3.39 (m, 1H), 3.16-3.07 (m, 3H), 2.78-2.72 (m, 1H), 2.05-1.98 (m, 1H), 1.96-1.86 (m, 1H), 1.65-1.59 (m, 1H), 1.44-1.37 (m, 1H).

The compounds in the table below were prepared by following the steps for preparing compound 4 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Intermediates
5		508.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.56 (m, 1H), 7.36-7.29 (m, 3H), 6.17-6.11 (m, 1H), 3.97-3.94 (m, 1H), 3.67-3.59 (m, 2H), 3.42 (s, 3H), 3.40-3.39 (m, 1H), 3.22-3.07 (m, 3H), 2.78-2.71 (m, 1H), 2.02-1.95 (m, 1H), 1.92-1.84 (m, 1H), 1.62-1.58 (m, 1H), 1.41-1.36 (m, 1H).	I-A1 I-A4 I-C1
6		474.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.33-8.30 (m, 1H), 8.29-8.25 (m, 1H), 8.16-8.12 (m, 1H), 8.11-8.07 (m, 1H), 7.80-7.76 (m, 1H), 7.66-7.60 (m, 1H), 7.38-7.30 (m, 3H), 7.23-7.18 (m, 1H), 4.02-3.99 (m, 1H), 3.96 (s, 3H), 3.82-3.75 (m, 2H), 3.42-3.39 (m, 1H), 3.27-3.20 (m, 2H), 3.17-3.12 (m, 1H), 2.82-2.75 (m, 1H), 2.16-2.08 (m, 1H), 2.05-1.98 (m, 1H), 1.73-1.66 (m, 1H), 1.52-1.45 (m, 1H).	I-C1
25		489.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.29-8.25 (m, 1H), 7.81-7.77 (m, 1H), 7.54 (s, 1H), 7.43 (d, J = 1.5 Hz, 1H), 7.38-7.30 (m, 3H), 7.20 (d, J = 1.3 Hz, 1H), 7.13-7.09 (m, 1H), 4.01 (s, 1H), 3.94 (s, 3H), 3.85-3.77 (m, 2H), 3.41 (s, 1H), 3.26-3.11 (m, 3H), 2.81-2.77 (m, 1H), 2.09-1.96 (m, 2H), 1.68-1.64 (m, 1H), 1.48-1.44 (m, 1H).	I-A8 I-C6
120		518.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.39-8.30 (m, 2H), 8.19 (s, 1H), 8.15 (s, 1H), 7.85 (d, J = 1.5 Hz, 1H), 7.69 (d, J = 1.5 Hz, 1H), 7.52 (s, 1H), 7.36-7.31 (m, 1H), 7.29-7.22 (m, 2H), 4.35 (s, 2H), 4.04 (s, 1H), 4.01 (s, 3H), 3.90-3.79 (m, 2H), 3.46 (s, 3H), 3.32-3.19 (m, 3H), 2.89-2.82 (m, 1H), 2.23-2.03 (m, 2H), 1.75 (d, J = 12.8 Hz, 1H), 1.53 (d, J = 12.8 Hz, 1H).	I-C7

Compound 299

(S)-5-((2-amino-3-chloropyridin-4-yl)thio)-2-(1-amino-6-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-methylpyrimidin-4(3H)-one



299

- 5 **Step 1: (R)-N-((S)-1'-(5-iodo-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide**

Intermediate I-C3 (0.88 g, 2.19 mmol), 2-chloro-5-iodo-3-methylpyrimidin-4(3H)-one (0.59 g, 2.19 mmol) and DIEA (0.57 g, 4.38 mmol) were placed in DMA (5 mL) and
 10 stirred at 90°C for 3 hours. The reaction solution was purified by silica gel column chromatography (eluting with water/MeOH) to give target product (0.82 g, 59% yield).
 [M+H]⁺ 637.2

- 15 **Step 2: (R)-N-((S)-1'-(5-((2-amino-3-chloropyridin-4-yl)thio)-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide**

Under N₂, (R)-N-((S)-1'-(5-iodo-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide (637 mg, 1.0 mmol), intermediate I-A1 (337 mg, 1.85 mmol), 4,7-dimethoxy-1,10-phenanthroline (156 mg, 0.65 mmol), CuI (43 mg, 0.23 mmol) and
 20 anhydrous K₃PO₄ (420 mg, 1.98 mmol) were placed in 1,4-dioxane (10 mL). The reaction solution was stirred at 100°C for 3 hours, and concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with water/MeOH) to give target product (506 mg, 76% yield). [M+H]⁺ 669.2

Step 3: (*S*)-5-((2-amino-3-chloropyridin-4-yl)thio)-2-(1-amino-6-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-methylpyrimidin-4(3*H*)-one


(*R*)-*N*-((*S*)-1'-(5-((2-amino-3-chloropyridin-4-yl)thio)-1-methyl-6-oxo-1,6-dihydropyrimidin-2-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide (506 mg, 0.76 mmol) was dissolved in 2 M HCl/MeOH solution (6 mL). The reaction solution was stirred at room temperature for 5 minutes, diluted with DCM (15 mL) and adjusted pH to 8 with aqueous ammonia in an ice-bath. The organic layer was collected and concentrated in vacuum under reduced pressure. The residue was dissolved in MeOH (5 mL), added K₂CO₃ powder (522 mg, 3.78 mmol) and stirred at room temperature for 15 minutes. The reaction solution was purified by thin layer chromatography (eluting with DCM/MeOH = 12/1) to give target product (96 mg, 26% yield). [M+H]⁺ 493.2. ¹H NMR (400 MHz, CD₃OD): δ 8.16-8.11 (m, 1H), 7.62-7.56 (m, 1H), 7.51-7.48 (m, 1H), 7.35-7.30 (m, 1H), 7.24-7.19 (m, 1H), 6.17-6.10 (m, 1H), 3.99 (s, 1H), 3.77-3.69 (m, 2H), 3.53 (s, 3H), 3.42 (s, 1H), 3.29-3.19 (m, 2H), 3.18-3.12 (m, 1H), 2.85-2.76 (m, 1H), 2.05-1.96 (m, 1H), 1.94-1.86 (m, 1H), 1.68-1.60 (m, 1H), 1.46-1.38 (m, 1H).

The compounds in the table below were prepared by following the steps for preparing compound 299 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Intermediates
288		507.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.14-8.09 (m, 1H), 7.62-7.54 (m, 1H), 7.40-7.32 (m, 1H), 7.21-7.11 (m, 2H), 6.17-6.08 (m, 1H), 3.93 (s, 1H), 3.75-3.67 (m, 2H), 3.51 (s, 3H), 3.27-3.18 (m, 2H), 3.14-3.06 (m, 1H), 2.80-2.69 (m, 1H), 2.02-1.95 (m, 4H), 1.92-1.83 (m, 1H), 1.65-1.57 (m, 1H), 1.43-1.36 (m, 1H).	I-A1 I-C17
291		537.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.16-8.11 (m, 1H), 7.61-7.57 (m, 1H), 7.49-7.45 (m, 1H), 7.33-7.28 (m, 1H), 7.24-7.20 (m, 1H), 6.16-6.11 (m, 1H), 4.31 (s, 2H), 3.99 (s, 1H), 3.77-3.69 (m, 2H), 3.53 (s, 3H), 3.42 (s, 3H), 3.28-3.20 (m, 2H), 3.17-3.12 (m, 1H), 2.85-2.76 (m, 1H), 2.05-1.97 (m, 1H), 1.95-1.85 (m, 1H), 1.66-1.60 (m, 1H), 1.46-1.39 (m, 1H).	I-A1 I-C7
294		507.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.12 (s, 1H), 7.60 (d, J = 5.6 Hz, 1H), 7.49 (s, 1H), 7.31 (d, J = 7.7 Hz, 1H), 7.20 (d, J = 7.7 Hz, 1H), 6.10 (d, J = 5.6 Hz, 1H), 4.11 (q, J = 6.9 Hz, 2H), 3.95 (s, 1H), 3.66-3.58 (m, 2H), 3.41 (s, 1H), 3.29-3.19 (m, 2H), 3.13 (d, J = 16.1 Hz, 1H), 2.76 (d, J = 16.1 Hz, 1H), 2.06-1.86 (m, 2H), 1.63 (d, J = 13.5 Hz, 1H), 1.40 (d, J = 13.2 Hz, 1H), 1.34 (t, J = 6.9 Hz, 3H).	I-A1 I-C3
295		495.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.14 (s, 1H), 7.61-7.57 (m, 1H), 7.53-7.49 (m, 1H), 7.35-7.28 (m, 1H), 6.80-6.74 (m, 1H), 6.17-6.12 (m, 1H), 4.14 (s, 1H), 3.87-3.80 (m, 1H), 3.77-3.71 (m, 1H), 3.54 (s, 3H), 3.50-3.41 (m, 2H), 3.34 (s, 1H), 2.17-2.09 (m, 1H), 2.04-1.94 (m, 2H), 1.88-1.80 (m, 1H).	I-A1 I-C32

Compound 12

(S)-6-(1-amino-6-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-(2,3-dichlorophenyl)-2-methylpyrimidin-4(3H)-one



5 **Step 1: (R)-N-((S)-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide**

Intermediate I-B1 (2.0 g, 3.73 mmol), intermediate I-C3 (1.5 g, 3.73 mmol) and DIEA (1.9 g, 14.9 mmol) were placed in *N,N*-dimethylformamide (5 mL). The reaction was stirred at 100°C for 3 hours, and the reaction solution was purified with silica gel column chromatography (water/methanol) to give the target product.

Step 2: (R)-N-((S)-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide




15 (R)-N-((S)-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-5-((trimethylsilyl)ethynyl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide was dissolved in 2 M hydrogen chloride/methanol solution (5 mL, 10 mmol). The reaction was stirred at room temperature for 10 minutes, and under ice bath cooling, aqueous ammonia (3 mL) and water (30 mL) were added thereto. The mixture was extracted with dichloromethane. The organic phases were collected and combined, and concentrated in vacuum under reduced pressure to give a crude target product. $[M+H]^+$ 551.2





Step 3: (S)-6-(1-amino-6-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-(2,3-dichlorophenyl)-2-methylpyrimidin-4(3H)-one





The crude (R)-N-((S)-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-5-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfonamide and potassium carbonate (2.1 g, 14.9 mmol) were placed in methanol (15 mL), and the mixture was stirred at room temperature for 30 minutes. Water was added thereto, and the mixture was extracted with dichloromethane. The organic phases were collected and combined, and concentrated in vacuum under reduced pressure, and the residue was purified with silica gel column chromatography (dichloromethane/methanol) to give the target product (490 mg, three-step yield 27%).





[M+H]⁺ 479.2. ¹H NMR (400 MHz, CD₃OD): δ 7.70 (dd, J = 8.2, 1.5 Hz, 1H), 7.52-7.45 (m, 2H), 7.38 (dd, J = 7.9, 1.5 Hz, 1H), 7.33-7.27 (m, 1H), 7.19 (d, J = 7.7 Hz, 1H), 5.48 (s, 1H), 4.51-4.05 (m, 2H), 3.92 (s, 1H), 3.39 (s, 1H), 3.26- 3.12 (m, 3H), 2.77 (d, J = 16.1 Hz, 1H), 2.06 (s, 3H), 1.89-1.67 (m, 2H), 1.63-1.54(m, 1H), 1.39-1.31 (m, 1H).





The compounds in the table below were prepared by following the steps for preparing compound 12 from corresponding intermediates and reagents:





Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Intermediates
11		481.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.71 (dd, J = 8.1, 1.5 Hz, 1H), 7.48 (t, J = 8.0 Hz, 1H), 7.41-7.31 (m, 2H), 7.02 (dd, J = 7.6, 1.3 Hz, 1H), 6.87 (s, 1H), 5.54 (s, 1H), 4.57-4.18 (m, 2H), 4.11 (s, 1H), 3.47-3.34 (m, 3H), 2.07 (s, 3H), 1.96-1.89 (m, 2H), 1.84-1.76 (m, 2H).	I-B1 I-C5
28		481.1	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃): δ 7.62-7.55 (m, 1H), 7.40-7.34 (m, 1H), 7.29-7.24 (m, 1H), 7.23-7.18 (m, 1H), 7.05-6.99 (m, 1H), 6.90-6.86 (m, 1H), 5.46 (s, 1H), 4.37-4.11 (m, 2H), 4.06 (s, 1H), 3.43-3.33 (m, 2H), 3.13 (s, 1H), 2.05 (s, 3H), 1.90-1.78 (m, 3H), 1.76-1.67 (m, 1H).	I-B2 I-C5
13		449.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.47-7.39 (m, 1H), 7.36-7.30 (m, 2H), 7.14-7.09 (m, 1H), 7.04-6.99 (m, 1H), 6.88-6.84 (m, 1H), 5.54-5.50 (m, 1H), 4.44-4.21 (m, 2H), 4.12-4.07 (m, 1H), 3.43-3.34 (m, 3H), 2.13 (s, 3H), 1.94-1.88 (m, 2H), 1.82-1.75 (m, 2H).	I-B3 I-C5






14		463.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.48-7.40 (m, 1H), 7.38-7.29 (m, 2H), 7.14-7.08 (m, 1H), 7.04-6.97 (m, 1H), 6.88-6.82 (m, 1H), 4.12-4.08 (m, 1H), 3.99-3.91 (m, 1H), 3.91-3.83 (m, 1H), 3.45-3.36 (m, 3H), 2.12 (s, 3H), 2.07-2.01 (m, 1H), 2.00 (s, 3H), 1.93-1.84 (m, 2H), 1.83-1.74 (m, 1H).	I-B4 I-C5
15		447.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.50-7.37 (m, 2H), 7.36-7.27 (m, 2H), 7.22-7.16 (m, 1H), 7.13-7.08 (m, 1H), 5.50-5.42 (m, 1H), 4.35-4.08 (m, 2H), 3.94-3.88 (m, 1H), 3.38 (s, 1H), 3.25-3.11 (m, 3H), 2.81-2.72 (m, 1H), 2.11 (s, 3H), 1.85-1.76 (m, 1H), 1.75-1.66 (m, 1H), 1.59-1.52 (m, 1H), 1.38-1.31 (m, 1H).	I-B3 I-C3
16		463.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.67-7.65 (m, 1H), 7.49 (s, 1H), 7.40-7.28 (m, 3H), 7.25-7.18 (m, 1H), 5.50 (s, 1H), 4.50-4.10 (m, 2H), 3.94 (s, 1H), 3.42-3.34 (m, 1H), 3.30-3.12 (m, 3H), 2.83-2.75 (m, 1H), 2.13 (s, 3H), 1.85-1.63 (m, 2H), 1.63-1.57 (m, 1H), 1.41-1.33 (m, 1H).	I-B5 I-C3
17		445.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.54-7.48 (m, 2H), 7.47-7.45 (m, 1H), 7.37-7.34 (m, 1H), 7.32-7.27 (m, 1H), 7.23-7.16 (m, 2H), 5.47 (s, 1H), 4.33-4.15 (m, 2H), 3.90 (s, 1H), 3.40 (s, 1H), 3.26-3.10 (m, 3H), 2.82-2.69 (m, 1H), 2.09 (s, 3H), 1.86-1.75 (m, 1H), 1.74-1.63 (m, 1H), 1.60-1.49 (m, 1H), 1.36-1.30 (m, 1H).	I-B6 I-C3





18		447.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.55-7.49 (m, 2H), 7.39-7.36 (m, 1H), 7.36-7.31 (m, 1H), 7.25-7.18 (m, 1H), 7.05-6.99 (m, 1H), 6.86 (s, 1H), 5.52 (s, 1H), 4.47-4.17 (m, 2H), 4.09 (s, 1H), 3.46-3.34 (m, 3H), 2.11 (s, 3H), 1.95-1.87 (m, 2H), 1.82-1.75 (m, 2H).	I-B6 I-C5
19		413.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.62-7.51 (m, 3H), 7.49 (d, J = 7.5 Hz, 1H), 7.33-7.26 (m, 2H), 7.15 (d, J = 7.5 Hz, 1H), 7.03 (s, 1H), 5.61 (s, 1H), 4.55-4.34 (m, 3H), 3.58 (s, 1H), 3.52-3.39 (m, 2H), 2.14 (s, 3H), 2.09-1.99 (m, 2H), 1.93-1.79 (m, 2H).	I-B7 I-C5
20		479.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.1, 1.5 Hz, 1H), 7.48 (t, J = 8.0 Hz, 1H), 7.40-7.26 (m, 4H), 5.48 (s, 1H), 4.48-4.10 (m, 2H), 3.93 (s, 1H), 3.39 (s, 1H), 3.26-3.11 (m, 3H), 2.76 (d, J = 15.7 Hz, 1H), 2.06 (s, 3H), 1.90-1.67 (m, 2H), 1.63-1.51 (m, 1H), 1.40-1.32 (m, 1H).	I-B1 I-C6
24		479.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.78-7.71 (m, 1H), 7.55-7.49 (m, 1H), 7.44-7.40 (m, 1H), 7.39-7.32 (m, 3H), 5.52 (s, 1H), 4.31 (s, 2H), 3.97 (s, 1H), 3.37 (s, 1H), 3.31-3.22 (m, 2H), 3.18 (d, J = 15.8 Hz, 1H), 2.80 (d, J = 15.8 Hz, 1H), 2.10 (s, 3H), 1.92-1.82 (m, 1H), 1.76 (m, 1H), 1.66-1.58 (m, 1H), 1.44-1.36 (m, 1H).	I-B2 I-C6





21		447.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.47-7.39 (m, 1H), 7.37-7.26 (m, 4H), 7.14-7.08 (m, 1H), 5.49-5.45 (m, 1H), 4.39-4.08 (m, 2H), 3.94-3.90 (m, 1H), 3.40 (s, 1H), 3.27-3.17 (m, 2H), 3.16-3.10 (m, 1H), 2.80-2.71 (m, 1H), 1.85-1.76 (m, 1H), 1.74-1.66 (m, 1H), 1.59-1.53 (m, 1H), 1.37-1.32 (m, 1H).	I-B3 I-C6
22		445.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.56-7.51 (m, 2H), 7.35-7.28 (m, 3H), 7.27-7.23 (m, 2H), 5.49-5.46 (m, 1H), 4.40-4.10 (m, 2H), 3.94-3.90 (m, 1H), 3.39 (s, 1H), 3.27-3.17 (m, 2H), 3.16-3.10 (m, 1H), 2.79-2.72 (m, 1H), 2.09 (s, 3H), 1.86-1.76 (m, 1H), 1.75-1.65 (m, 1H), 1.59-1.53 (m, 1H), 1.37-1.32 (m, 1H).	I-B8 I-C6
23		411.2	¹ H NMR (400 MHz, DMSO-d ₆): δ 7.57-7.40 (m, 3H), 7.36-7.19 (m, 5H), 5.37 (s, 1H), 4.24-4.00 (m, 2H), 3.89-3.75 (m, 1H), 3.20-3.18 (m, 2H), 3.13-3.01 (m, 3H), 2.68-2.56 (m, 2H), 1.99 (s, 3H), 1.79-1.69 (m, 2H), 1.63-1.42 (m, 2H).	I-B7 I-C6
26		445.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.55-7.48 (m, 2H), 7.38-7.27 (m, 4H), 7.24-7.16 (m, 1H), 5.47 (s, 1H), 4.46-4.00 (m, 2H), 3.91 (s, 1H), 3.40 (s, 1H), 3.26-3.15 (m, 2H), 3.15-3.08 (m, 1H), 2.79-2.69 (m, 1H), 2.09 (s, 3H), 1.85-1.75 (m, 1H), 1.75-1.63 (m, 1H), 1.61-1.49 (m, 1H), 1.37-1.29 (m, 1H).	I-B6 I-C6





29		459.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.45 (m, 1H), 7.33-7.28 (m, 1H), 7.25-7.18 (m, 2H), 7.07-7.01 (m, 1H), 6.83-6.77 (m, 1H), 5.55-5.47 (s, 1H), 4.25 (br, 2H), 3.93-3.91 (m, 1H), 3.87 (s, 3H), 3.41-3.36 (m, 1H), 3.26-3.10 (m, 3H), 2.83-2.73 (m, 1H), 2.12 (s, 3H), 1.85-1.76 (m, 1H), 1.76-1.65 (m, 1H), 1.60-1.53 (m, 1H), 1.38-1.32 (m, 1H).	I-B9 I-C3
30		523.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.68 (m, 1H), 7.50-7.43 (m, 2H), 7.40-7.35 (m, 1H), 7.30-7.26 (m, 1H), 7.22-7.18 (m, 1H), 5.50-5.46 (m, 1H), 4.43-4.12 (m, 4H), 3.95-3.91 (m, 1H), 3.40 (s, 3H), 3.27-3.10 (m, 3H), 2.84-2.74 (m, 1H), 2.06 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.68 (m, 1H), 1.61-1.55 (m, 1H), 1.40-1.34 (m, 1H).	I-B1 I-C7
31		565.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.2, 1.5 Hz, 1H), 7.48 (t, J = 8.0 Hz, 1H), 7.42 (s, 1H), 7.38 (dd, J = 7.9, 1.5 Hz, 1H), 7.26 (d, J = 7.7 Hz, 1H), 7.18 (d, J = 7.8 Hz, 1H), 5.48 (s, 1H), 4.43-4.09 (m, 4H), 3.91 (s, 1H), 3.26-3.11 (m, 3H), 2.77 (d, J = 16.0 Hz, 1H), 2.06 (s, 3H), 1.89-1.79 (m, 1H), 1.76-1.66 (m, 1H), 1.66-1.53 (m, 1H), 1.39-1.32 (m, 1H), 1.26 (s, 9H).	I-B1 I-C8'
32		509.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.1, 1.4 Hz, 1H), 7.53-7.34 (m, 3H), 7.26 (d, J = 6.8 Hz, 1H), 7.18 (d, J = 7.8 Hz, 1H), 5.48 (s, 1H), 4.34-4.09 (m, 4H), 3.91 (s, 1H), 3.25-3.11 (m, 3H), 2.81-2.71 (m, 1H), 2.06 (s, 3H), 1.88-1.78 (m, 1H), 1.76-1.67 (m, 1H), 1.62-1.53 (m, 1H), 1.40-1.31 (m, 1H).	I-B1 I-C8





33		479.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.66-7.57 (m, 2H), 7.55-7.43 (m, 2H), 7.33-7.27 (m, 1H), 7.23-7.17 (m, 1H), 5.48 (s, 1H), 4.35-4.25 (m, 2H), 3.93 (s, 1H), 3.39 (s, 1H), 3.28-3.12 (m, 3H), 2.82-2.74 (m, 1H), 2.05 (s, 3H), 1.91-1.67 (m, 2H), 1.61-1.57 (m, 1H), 1.38-1.34 (m, 1H).	I-B10 I-C3
34		470.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.00-7.94 (m, 1H), 7.77-7.72 (m, 1H), 7.69-7.62 (m, 1H), 7.47 (s, 1H), 7.33-7.27 (m, 1H), 7.22-7.16 (m, 1H), 5.48 (s, 1H), 4.45-4.08 (m, 2H), 3.91 (s, 1H), 3.39 (s, 1H), 3.26-3.10 (m, 3H), 2.85-2.70 (m, 1H), 2.06 (s, 3H), 1.87-1.68 (m, 2H), 1.60-1.53 (m, 1H), 1.39-1.33 (m, 1H).	I-B11 I-C3
35		447.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.63-7.53 (m, 1H), 7.46 (s, 1H), 7.32-7.27 (m, 1H), 7.24-7.16 (m, 3H), 5.47 (s, 1H), 4.51-4.00 (m, 2H), 3.91 (s, 1H), 3.39 (s, 1H), 3.27-3.10 (m, 3H), 2.81-2.72 (m, 1H), 2.14 (s, 3H), 1.88-1.79 (m, 1H), 1.75-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.36-1.31 (m, 1H).	I-B12 I-C3
36		537.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.67 (m, 1H), 7.52-7.41 (m, 2H), 7.39-7.33 (m, 1H), 7.30-7.24 (m, 1H), 7.23-7.16 (m, 1H), 4.33-4.26 (m, 2H), 3.96-3.86 (m, 3H), 3.40 (s, 3H), 3.25-3.19 (m, 2H), 3.16-3.11 (m, 1H), 2.82-2.72 (m, 1H), 2.05 (s, 3H), 2.00 (s, 3H), 1.95-1.88 (m, 1H), 1.85-1.77 (m, 1H), 1.60-1.54 (m, 1H), 1.39-1.34 (m, 1H).	I-B13 I-C7

39		451.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.84 (s, 1H), 7.71-7.65 (m, 1H), 7.54-7.46 (m, 2H), 7.33-7.27 (m, 1H), 7.23-7.18 (m, 1H), 7.11-7.05 (m, 1H), 5.54 (s, 1H), 4.39-4.17 (m, 2H), 3.94 (s, 1H), 3.39 (s, 1H), 3.28-3.13 (m, 3H), 2.84-2.74 (m, 1H), 2.06 (s, 3H), 1.90-1.81 (m, 1H), 1.79-1.70 (m, 1H), 1.61-1.56 (m, 1H), 1.40-1.34 (m, 1H).	I-B14 I-C3
41		447.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.47-7.35 (m, 2H), 7.31-7.26 (m, 1H), 7.25-7.11 (m, 3H), 5.48-5.44 (m, 1H), 4.46-4.00 (m, 2H), 3.91 (s, 1H), 3.33 (s, 1H), 3.28-3.09 (m, 3H), 2.82-2.68 (m, 1H), 2.20-1.98 (m, 3H), 1.90-1.79 (m, 1H), 1.75-1.55 (m, 2H), 1.38-1.31 (m, 1H).	I-B15 I-C3
42		445.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.65-7.60 (m, 1H), 7.52-7.46 (m, 3H), 7.40-7.37 (m, 1H), 7.32-7.28 (m, 1H), 7.22-7.18 (m, 1H), 5.49-5.47 (m, 1H), 4.42-4.12 (m, 2H), 3.94-3.91 (m, 1H), 3.39 (s, 1H), 3.26-3.12 (m, 3H), 2.82-2.74 (m, 1H), 2.04 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.68 (m, 1H), 1.61-1.54 (m, 1H), 1.39-1.33 (m, 1H).	I-B16 I-C3
43		441.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.39 (m, 2H), 7.33-7.27 (m, 1H), 7.22-7.16 (m, 1H), 7.07-7.01 (m, 1H), 6.85-6.77 (m, 2H), 5.51-5.45 (m, 1H), 4.38-4.10 (m, 2H), 3.94-3.89 (m, 1H), 3.39 (s, 1H), 3.24-3.11 (m, 3H), 2.81-2.73 (m, 1H), 2.10 (s, 3H), 1.85-1.77 (m, 1H), 1.75-1.66 (m, 1H), 1.59-1.52 (m, 1H), 1.38-1.31 (m, 1H).	I-B17 I-C3
44		475.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.51-7.45 (m, 2H), 7.32-7.28 (m, 1H), 7.21-7.18 (m, 1H), 7.09-7.05 (m, 1H), 7.00-6.98 (m, 1H), 5.48-5.47 (m, 1H), 4.40-4.12 (m, 2H), 3.93-3.91 (m, 1H), 3.39 (s, 1H), 3.25-3.12 (m, 3H), 2.81-2.74 (m, 1H), 2.08 (s, 3H), 1.87-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.38-1.33 (m, 1H).	I-B18 I-C3





45		463.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.54-7.48 (m, 1H), 7.48-7.43 (m, 1H), 7.43-7.38 (m, 1H), 7.31-7.23 (m, 2H), 7.21-7.16 (m, 1H), 5.49-5.45 (m, 1H), 4.35-4.12 (m, 2H), 3.93-3.88 (m, 1H), 3.39 (s, 1H), 3.25-3.11 (m, 3H), 2.81-2.72 (m, 1H), 2.11 (s, 3H), 1.85-1.76 (m, 1H), 1.74-1.66 (m, 1H), 1.59-1.52 (m, 1H), 1.38-1.31 (m, 1H).	I-B19 I-C3
46		489.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.44 (m, 1H), 7.34-7.27 (m, 2H), 7.20-7.15 (m, 2H), 7.14-7.10 (m, 1H), 5.48-5.45 (m, 1H), 4.32-4.12 (m, 4H), 3.92-3.89 (m, 1H), 3.39 (s, 1H), 3.25-3.11 (m, 3H), 2.80-2.73 (m, 1H), 2.10 (s, 3H), 1.85-1.76 (m, 1H), 1.74-1.65 (m, 1H), 1.58-1.52 (m, 1H), 1.45 (t, J = 7.0 Hz, 3H), 1.36-1.30 (m, 1H).	I-B20 I-C3
48		493.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.67 (m, 1H), 7.51-7.45 (m, 2H), 7.39-7.33 (m, 1H), 7.32-7.27 (m, 1H), 7.21-7.16 (m, 1H), 3.96-3.85 (m, 3H), 3.39 (s, 1H), 3.28-3.17 (m, 2H), 3.17-3.09 (m, 1H), 2.80-2.71 (m, 1H), 2.05 (s, 3H), 2.00 (s, 3H), 1.96-1.87 (m, 1H), 1.85-1.75 (m, 1H), 1.60-1.52 (m, 1H), 1.38-1.32 (m, 1H).	I-B21 I-C3
49		493.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.67 (m, 1H), 7.52-7.44 (m, 2H), 7.39-7.34 (m, 1H), 7.32-7.27 (m, 1H), 7.21-7.17 (m, 1H), 3.99-3.85 (m, 3H), 3.39 (s, 1H), 3.28-3.19 (m, 2H), 3.18-3.10 (m, 1H), 2.82-2.72 (m, 1H), 2.05 (s, 3H), 2.00 (s, 3H), 1.96-1.86 (m, 1H), 1.86-1.76 (m, 1H), 1.60-1.54 (m, 1H), 1.39-1.34 (m, 1H).	I-B22 I-C3





50		550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.68 (m, 1H), 7.51-7.46 (m, 1H), 7.44 (s, 1H), 7.40-7.33 (m, 1H), 7.29-7.24 (m, 1H), 7.21-7.15 (m, 1H), 3.97-3.85 (m, 3H), 3.49-3.41 (m, 2H), 3.28-3.18 (m, 2H), 3.17-3.09 (m, 1H), 2.82-2.72 (m, 1H), 2.36 (s, 6H), 2.05 (s, 3H), 2.00 (s, 3H), 1.97-1.86 (m, 1H), 1.86-1.75 (m, 1H), 1.62-1.52 (m, 1H), 1.39-1.33 (m, 1H).	I-B21 I-C10
51		550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.72-7.68 (m, 1H), 7.52-7.45 (m, 1H), 7.44 (s, 1H), 7.39-7.34 (m, 1H), 7.29-7.23 (m, 1H), 7.21-7.15 (m, 1H), 4.00-3.80 (m, 3H), 3.49-3.42 (m, 2H), 3.28-3.18 (m, 2H), 3.17-3.08 (m, 1H), 2.81-2.70 (m, 1H), 2.36 (s, 6H), 2.05 (s, 3H), 2.00 (s, 3H), 1.96-1.86 (m, 1H), 1.86-1.75 (m, 1H), 1.61-1.51 (m, 1H), 1.40-1.32 (m, 1H).	I-B22 I-C10
52		537.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.67 (m, 1H), 7.50-7.45 (m, 1H), 7.44 (s, 1H), 7.40-7.34 (m, 1H), 7.29-7.24 (m, 1H), 7.21-7.16 (m, 1H), 5.47 (s, 1H), 4.51-3.95 (m, 2H), 3.91 (s, 1H), 3.48-3.43 (m, 2H), 3.28-3.17 (m, 2H), 3.17-3.10 (m, 1H), 2.83-2.67 (m, 1H), 2.35 (s, 6H), 2.05 (s, 3H), 1.88-1.77 (m, 1H), 1.76-1.65 (m, 1H), 1.61-1.52 (m, 1H), 1.40-1.30 (m, 1H).	I-B1 I-C10
53		457.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.50-7.28 (m, 4H), 7.22-7.10 (m, 2H), 7.01-6.97 (m, 1H), 5.48 (s, 1H), 4.34-4.14 (m, 2H), 3.91 (s, 1H), 3.39 (s, 1H), 3.25-3.09 (m, 3H), 2.79-2.75 (m, 1H), 2.49 (s, 3H), 2.09 (s, 3H), 1.86-1.64 (m, 2H), 1.57-1.53 (m, 1H), 1.36-1.32 (m, 1H).	I-B23 I-C3





54		477.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.52-7.38 (m, 3H), 7.32-7.24 (m, 2H), 7.20-7.18 (m, 1H), 3.97-3.83 (m, 3H), 3.39 (s, 1H), 3.26-3.08 (m, 3H), 2.77-2.75 (m, 1H), 2.08-1.72 (m, 8H), 1.58-1.54 (m, 1H), 1.37-1.33 (m, 1H).	I-B24 I-C3
55		447.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.46 (s, 1H), 7.35-6.95 (m, 5H), 5.46 (s, 1H), 4.34-4.14 (m, 2H), 3.90 (s, 1H), 3.39 (s, 1H), 3.27-3.09 (m, 3H), 2.78-2.74 (m, 1H), 2.13 (s, 3H), 1.86-1.63 (m, 2H), 1.57-1.53 (m, 1H), 1.36-1.32 (m, 1H).	I-B25 I-C3
60		446.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.57-8.52 (m, 1H), 7.57-7.53 (m, 1H), 7.49-7.44 (m, 1H), 7.39-7.35 (m, 1H), 7.32-7.27 (m, 1H), 7.21-7.16 (m, 1H), 5.46 (s, 1H), 4.24 (br, 2H), 3.91 (s, 1H), 3.39 (s, 1H), 3.26-3.11 (m, 3H), 2.80-2.73 (m, 1H), 2.13 (s, 3H), 1.85-1.76 (m, 1H), 1.74-1.66 (m, 1H), 1.60-1.53 (m, 1H), 1.38-1.31 (m, 1H).	I-B26 I-C3
61		463.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.45 (m, 3H), 7.37-7.29 (m, 2H), 7.26-7.22 (m, 1H), 5.53 (s, 1H), 4.31 (s, 2H), 3.97 (s, 1H), 3.44 (s, 1H), 3.31-3.17 (m, 3H), 2.86-2.78 (m, 1H), 2.15-2.06 (m, 3H), 1.93-1.74 (m, 2H), 1.66-1.60 (m, 1H), 1.44-1.38 (m, 1H).	I-B27 I-C3





62		489.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.52 (s, 1H), 7.49-7.43 (m, 1H), 7.35 (d, J = 7.0 Hz, 1H), 7.28-7.20 (m, 2H), 6.99 (d, J = 7.0 Hz, 1H), 5.53 (s, 1H), 4.41-4.18 (m, 4H), 3.96 (s, 1H), 3.38 (s, 1H), 3.31-3.17 (m, 3H), 2.87-2.79 (m, 1H), 2.10 (s, 3H), 1.93-1.72 (m, 2H), 1.66-1.59 (m, 1H), 1.53-1.45 (m, 3H), 1.44-1.37 (m, 1H).	I-B28 I-C3
63		550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.1, 1.4 Hz, 1H), 7.52 (s, 1H), 7.47 (t, J = 8.0 Hz, 1H), 7.37 (dd, J = 7.9, 1.4 Hz, 1H), 7.32 (s, 1H), 6.98 (d, J = 8.1 Hz, 1H), 6.86 (d, J = 2.0 Hz, 1H), 6.65 (dd, J = 8.1, 2.3 Hz, 1H), 5.47 (s, 1H), 4.49-4.06 (m, 2H), 3.84-3.81 (m, 4H), 3.27-3.16 (m, 2H), 3.01 (d, J = 15.2 Hz, 1H), 2.66 (d, J = 15.1 Hz, 1H), 2.05 (s, 3H), 1.84-1.65 (m, 2H), 1.60-1.52 (m, 1H), 1.44-1.36 (m, 1H).	I-B1 I-C11
64		479.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.2, 1.5 Hz, 1H), 7.52-7.45 (m, 2H), 7.38 (dd, J = 7.9, 1.5 Hz, 1H), 7.33-7.27 (m, 1H), 7.19 (d, J = 7.7 Hz, 1H), 5.48 (s, 1H), 4.51-4.05 (m, 2H), 3.92 (s, 1H), 3.39 (s, 1H), 3.26-3.12 (m, 3H), 2.77 (d, J = 16.1 Hz, 1H), 2.06 (s, 3H), 1.89-1.67 (m, 2H), 1.63-1.54 (m, 1H), 1.39-1.31 (m, 1H).	I-B2 I-C3
68		463.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.22 (m, 6H), 5.48 (s, 1H), 4.40-4.10 (m, 3H), 3.44 (s, 1H), 3.28-3.11 (m, 3H), 2.94-2.90 (m, 1H), 2.09 (s, 3H), 1.87-1.42 (m, 4H).	I-B29 I-C3





69		497.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.54-7.37 (m, 3H), 7.33-7.16 (m, 2H), 5.48 (s, 1H), 4.40-4.05 (m, 2H), 3.92 (s, 1H), 3.41 (s, 1H), 3.33-3.13 (m, 3H), 2.81-2.75 (m, 1H), 2.07 (s, 3H), 1.90-1.65 (m, 2H), 1.61-1.52 (m, 1H), 1.41-1.32 (m, 1H).	I-B30 I-C3
70		554.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.50-7.41 (m, 3H), 7.32-7.17 (m, 2H), 5.48 (s, 1H), 4.50-4.07 (m, 2H), 3.95 (s, 1H), 3.47 (s, 2H), 3.27-3.09 (m, 3H), 2.84-2.80 (m, 1H), 2.36 (s, 6H), 2.07 (s, 3H), 1.88-1.63 (m, 2H), 1.60-1.56 (m, 1H), 1.40-1.36 (m, 1H).	I-B30 I-C10
73		479.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.65-7.60 (m, 1H), 7.57-7.51 (m, 2H), 7.47 (s, 1H), 7.33-7.28 (m, 1H), 7.22-7.17 (m, 1H), 5.48 (s, 1H), 4.55-3.99 (m, 2H), 3.92 (s, 1H), 3.40 (s, 1H), 3.28-3.11 (m, 3H), 2.85-2.70 (m, 1H), 2.08 (s, 3H), 1.88-1.79 (m, 1H), 1.75-1.66 (m, 1H), 1.61-1.54 (m, 1H), 1.39-1.32 (m, 1H).	I-B31 I-C3
74		520.3	¹ H NMR (400 MHz, CD ₃ OD): δ 7.60-7.53 (m, 1H), 7.52-7.44 (m, 2H), 7.35-7.28 (m, 2H), 7.27-7.19 (m, 1H), 5.53 (s, 1H), 4.32 (s, 2H), 3.98 (s, 1H), 3.51 (s, 2H), 3.31-3.13 (m, 3H), 2.87-2.79 (m, 1H), 2.41 (s, 6H), 2.10 (s, 3H), 1.92-1.83 (m, 1H), 1.82-1.72 (m, 1H), 1.68-1.58 (m, 1H), 1.45-1.37 (m, 1H).	I-B27 I-C10





77		481.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.52-7.42 (m, 2H), 7.35-7.28 (m, 2H), 7.24-7.17 (m, 1H), 5.48 (s, 1H), 4.28 (br, 2H), 3.94 (s, 1H), 3.40 (s, 1H), 3.27-3.12 (m, 3H), 2.85-2.74 (m, 1H), 2.08 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.62-1.55 (m, 1H), 1.40-1.33 (m, 1H).	I-B32 I-C3
78		477.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.50-7.42 (m, 1H), 7.34-7.27 (m, 1H), 7.22-7.12 (m, 2H), 7.10-7.04 (m, 1H), 5.47 (s, 1H), 4.27 (br, 2H), 4.01 (s, 3H), 3.92 (s, 1H), 3.40 (s, 1H), 3.26-3.12 (m, 3H), 2.82-2.74 (m, 1H), 2.12 (s, 3H), 1.88-1.78 (m, 1H), 1.76-1.66 (m, 1H), 1.61-1.54 (m, 1H), 1.38-1.32 (m, 1H).	I-B33 I-C3
80		479.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.67 (m, 1H), 7.52-7.46 (m, 1H), 7.42-7.36 (m, 2H), 7.33-7.28 (m, 1H), 7.23-7.17 (m, 1H), 5.49 (s, 1H), 4.29 (br, 2H), 3.97 (s, 1H), 3.67 (s, 1H), 3.36-3.19 (m, 3H), 2.90-2.80 (m, 1H), 2.06 (s, 3H), 1.87-1.70 (m, 2H), 1.62-1.56 (m, 1H), 1.43-1.36 (m, 1H).	I-B1 I-C12
82		521.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.48-7.40 (m, 1H), 7.30-7.25 (m, 1H), 7.22-7.12 (m, 2H), 7.11-7.04 (m, 1H), 5.47 (s, 1H), 4.42-4.15 (m, 4H), 4.01 (s, 3H), 3.92 (s, 1H), 3.41 (s, 3H), 3.26-3.12 (m, 3H), 2.83-2.73 (m, 1H), 2.12 (s, 3H), 1.88-1.78 (m, 1H), 1.76-1.66 (m, 1H), 1.61-1.53 (m, 1H), 1.38-1.32 (m, 1H).	I-B33 I-C7





83		534.3	¹ H NMR (400 MHz, CD ₃ OD): δ 7.46-7.41 (m, 1H), 7.29-7.25 (m, 1H), 7.20-7.12 (m, 2H), 7.10-7.04 (m, 1H), 5.47 (s, 1H), 4.27 (br, 2H), 4.01 (s, 3H), 3.92 (s, 1H), 3.46 (s, 2H), 3.26-3.12 (m, 3H), 2.82-2.73 (m, 1H), 2.36 (s, 6H), 2.12 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.38-1.32 (m, 1H).	I-B33 I-C10
84		536.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.65-7.60 (m, 1H), 7.57-7.51 (m, 2H), 7.44 (s, 1H), 7.30-7.24 (m, 1H), 7.22-7.16 (m, 1H), 5.47 (s, 1H), 4.55-4.02 (m, 2H), 3.92 (s, 1H), 3.46 (s, 2H), 3.27-3.12 (m, 3H), 2.82-2.71 (m, 1H), 2.36 (s, 6H), 2.08 (s, 3H), 1.87-1.78 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.55 (m, 1H), 1.39-1.33 (m, 1H).	I-B31 I-C10
85		509.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.65-7.59 (m, 1H), 7.57-7.51 (m, 2H), 7.43 (s, 1H), 7.29-7.23 (m, 1H), 7.21-7.17 (m, 1H), 5.48 (s, 1H), 4.51-4.03 (m, 4H), 3.92 (s, 1H), 3.27-3.18 (m, 2H), 3.17-3.12 (m, 1H), 2.81-2.74 (m, 1H), 2.08 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.39-1.33 (m, 1H).	I-B31 I-C8
90		475.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.50-7.40 (m, 2H), 7.32-7.17 (m, 3H), 6.96 (d, J = 7.8 Hz, 1H), 5.48 (s, 1H), 4.42-4.15 (m, 2H), 3.99-3.90 (m, 4H), 3.40 (s, 1H), 3.28-3.13 (m, 3H), 2.81-2.72 (m, 1H), 2.04 (s, 3H), 1.90-1.64 (m, 2H), 1.59-1.55 (m, 1H), 1.37-1.33 (m, 1H).	I-B34 I-C3





92		503.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.48 (s, 1H), 7.30 (d, J = 7.7 Hz, 1H), 7.25-7.14 (m, 3H), 7.03-6.99 (m, 1H), 4.09 (q, J = 6.9 Hz, 2H), 3.96-3.80 (m, 3H), 3.40 (s, 1H), 3.26-3.12 (m, 3H), 2.79-2.75 (m, 1H), 2.06-1.81 (m, 8H), 1.58-1.54 (m, 1H), 1.42-1.32 (m, 4H).	I-B35 I-C3
93		514.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.14-8.05 (m, 1H), 7.91-7.83 (m, 1H), 7.67-7.60 (m, 1H), 7.44 (s, 1H), 7.31-7.25 (m, 1H), 7.23-7.17 (m, 1H), 5.48 (s, 1H), 4.56-3.97 (m, 4H), 3.92 (s, 1H), 3.41 (s, 3H), 3.28-3.10 (m, 3H), 2.86-2.70 (m, 1H), 2.06 (s, 3H), 1.89-1.79 (m, 1H), 1.78-1.66 (m, 1H), 1.62-1.55 (m, 1H), 1.39-1.33 (m, 1H).	I-B36 I-C7
94		470.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.15-8.08 (m, 1H), 7.92-7.85 (m, 1H), 7.69-7.59 (m, 2H), 7.51-7.43 (m, 1H), 7.38-7.30 (m, 1H), 5.51 (s, 1H), 4.63-3.99 (m, 3H), 3.53 (s, 1H), 3.24-3.04 (m, 2H), 2.98 (s, 1H), 2.85 (s, 1H), 2.07 (s, 3H), 1.90-1.71 (m, 2H), 1.71-1.52 (m, 2H).	I-B36 I-C3
96		523.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.67 (m, 1H), 7.51-7.46 (m, 1H), 7.41-7.34 (m, 2H), 7.31-7.25 (m, 1H), 7.24-7.17 (m, 1H), 5.49 (s, 1H), 4.49-4.06 (m, 4H), 3.97 (s, 1H), 3.43 (s, 3H), 3.28-3.16 (m, 3H), 2.88-2.80 (m, 1H), 2.06 (s, 3H), 1.86-1.71 (m, 2H), 1.61-1.55 (m, 1H), 1.43-1.36 (m, 1H).	I-B1 I-C13





99		545.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.78-7.65 (m, 1H), 7.48 (t, J = 8.1 Hz, 1H), 7.43-7.35 (m, 2H), 7.22 (d, J = 7.9 Hz, 1H), 7.16 (d, J = 7.8 Hz, 1H), 5.48 (s, 1H), 4.48-4.05 (m, 2H), 3.91 (s, 1H), 3.61-3.50 (m, 2H); 3.31 (s, 3H), 3.26-3.17 (m, 2H), 3.15-3.03 (m, 3H), 2.74 (d, J = 15.8 Hz, 1H), 2.06 (s, 3H), 1.84 (t, J = 10.4 Hz, 1H), 1.72 (t, J = 10.5 Hz, 1H), 1.58 (d, J = 13.3 Hz, 1H), 1.37 (d, J = 13.5 Hz, 1H).	I-B1 I-C14
101		543.4	¹ H NMR (400 MHz, CD ₃ OD): 7.41 (s, 1H), 7.25-7.20 (m, 1H), 7.20-7.11 (m, 2H), 7.12-7.02 (m, 1H), 5.47 (s, 1H), 4.52-4.06 (m, 2H), 4.01 (s, 3H), 3.91 (s, 1H), 3.58-3.49 (m, 2H), 3.31 (s, 3H), 3.27-3.17 (m, 2H), 3.14-3.03 (m, 3H), 2.77-2.71 (m, 1H), 2.12 (s, 3H), 1.89-1.77 (m, 1H), 1.77-1.64 (m, 1H), 1.62-1.52 (m, 1H), 1.41-1.32 (m, 1H).	I-B33 I-C14
104		519.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.48-7.40 (m, 2H), 7.30-7.18 (m, 3H), 6.98-6.94 (m, 1H), 5.48 (s, 1H), 4.40-4.20 (m, 4H), 3.98-3.90 (m, 4H), 3.41 (s, 3H), 3.27-3.09 (m, 3H), 2.82-2.72 (m, 1H), 2.05 (s, 3H), 1.90-1.64 (m, 2H), 1.60-1.56 (m, 1H), 1.38-1.34 (m, 1H).	I-B34 I-C7
106		513.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.04-7.96 (m, 1H), 7.80-7.69 (m, 2H), 7.52 (s, 1H), 7.35 (d, J = 7.6 Hz, 1H), 7.24 (d, J = 7.6 Hz, 1H), 5.54 (s, 1H), 4.33 (s, 2H), 3.97 (s, 1H), 3.44 (s, 1H), 3.32-3.16 (m, 3H), 2.87-2.79 (m, 1H), 2.10 (s, 3H), 1.95-1.71 (m, 2H), 1.63 (d, J = 13.6 Hz, 1H), 1.41 (d, J = 13.6 Hz, 1H).	I-B37 I-C3





107		479.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.99-7.91 (m, 1H), 7.91-7.81 (m, 1H), 7.80-7.72 (m, 1H), 7.55-7.45 (m, 2H), 7.35 (d, J = 7.5 Hz, 1H), 7.24 (d, J = 7.5 Hz, 1H), 5.51 (s, 1H), 4.32 (s, 2H), 3.97 (s, 1H), 3.34 (s, 1H), 3.31-3.11 (m, 3H), 2.87-2.75 (m, 1H), 2.08 (s, 3H), 1.95-1.71 (m, 2H), 1.63 (d, J = 12.9 Hz, 1H), 1.40 (d, J = 12.9 Hz, 1H).	I-B38 I-C3
115		537.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.66 (m, 1H), 7.52-7.42 (m, 2H), 7.40-7.35 (m, 1H), 7.34-7.25 (m, 1H), 7.21-7.12 (m, 1H), 5.48 (s, 1H), 4.49-4.70 (m, 4H), 3.91 (s, 1H), 3.71-3.53 (m, 2H), 3.26-3.10 (m, 3H), 2.82-2.71 (m, 1H), 2.06 (s, 3H), 1.87-1.79 (m, 1H), 1.75-1.66 (m, 1H), 1.60-1.53 (m, 1H), 1.38-1.32 (m, 1H), 1.26-1.16 (m, 3H).	I-B1 I-C15
121		533.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.77-7.70 (m, 1H), 7.51-7.41 (m, 3H), 7.31 (d, J = 7.5 Hz, 1H), 7.20 (d, J = 7.8 Hz, 1H), 5.67 (s, 1H), 4.76-4.04 (m, 2H), 3.94 (s, 1H), 3.40 (s, 1H), 3.37-3.31 (m, 2H), 3.18 (d, J = 16.1 Hz, 1H), 2.80 (d, J = 16.2 Hz, 1H), 1.95-1.83 (m, 1H), 1.81-1.69 (mz, 1H), 1.67-1.58 (m, 1H), 1.44-1.34 (m, 1H).	I-B39 I-C3
127		522.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.69 (m, 1H), 7.58-7.55 (m, 1H), 7.51-7.46 (m, 1H), 7.45-7.41 (m, 1H), 7.40-7.37 (m, 1H), 7.30-7.26 (m, 1H), 5.49 (s, 1H), 4.28 (br, 2H), 3.95 (s, 1H), 3.29-3.16 (m, 3H), 2.85-2.78 (m, 1H), 2.07 (s, 3H), 1.90-1.81 (m, 1H), 1.78-1.68 (m, 1H), 1.63-1.56 (m, 1H), 1.38-1.32 (m, 1H).	I-B1 I-C16





130		577.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.76-7.71 (m, 1H), 7.49-7.44 (m, 3H), 7.30-7.19 (m, 2H), 5.67 (s, 1H), 4.51-4.02 (m, 4H), 3.94 (s, 1H), 3.41 (s, 3H), 3.37-3.30 (m, 2H), 3.18 (d, J = 16.1 Hz, 1H), 2.79 (d, J = 16.1 Hz, 1H), 1.95-1.82 (m, 1H), 1.81-1.69 (m, 1H), 1.63 (d, J = 13.7 Hz, 1H), 1.39 (d, J = 13.5 Hz, 1H).	I-B39 I-C7
133		457.2	¹ H NMR (400 MHz, CD ₃ OD) δ 7.52-7.41 (m, 3H), 7.35-7.27 (m, 2H), 7.22-7.15 (m, 2H), 5.48 (s, 1H), 4.39-4.11 (m, 2H), 3.91 (s, 1H), 3.40 (s, 1H), 3.26-3.10 (m, 3H), 2.80-2.70 (m, 1H), 2.45 (s, 3H), 2.03 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.53 (m, 1H), 1.38-1.32 (m, 1H).	I-B40 I-C3
135		497.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.71 (d, J = 8.4 Hz, 1H), 7.49 (t, J = 7.7 Hz, 1H), 7.39 (d, J = 7.3 Hz, 1H), 7.32 (s, 1H), 7.04 (d, J = 8.9 Hz, 1H), 5.49 (s, 1H), 4.53-4.07 (m, 2H), 3.95 (s, 1H), 3.51 (s, 1H), 3.27-3.16 (m, 3H), 2.78 (d, J = 16.0 Hz, 1H), 2.07 (s, 3H), 1.91-1.70 (m, 2H), 1.61 (d, J = 13.0 Hz, 1H), 1.37 (d, J = 12.8 Hz, 1H).	I-B1 I-C19
157		493.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.51-7.44 (m, 1H), 7.36-7.28 (m, 2H), 7.22-7.14 (m, 2H), 5.48 (s, 1H), 4.28 (br, 2H), 4.00 (d, J = 1.1 Hz, 3H), 3.94 (s, 1H), 3.40 (s, 1H), 3.26-3.14 (m, 3H), 2.84-.74 (m, 1H), 2.06 (s, 3H), 1.89-1.79 (m, 1H), 1.77-1.67 (m, 1H), 1.62-1.55 (m, 1H), 1.40-1.34 (m, 1H).	I-B41 I-C2





158		537.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.41 (m, 1H), 7.35-7.27 (m, 2H), 7.23-7.14 (m, 2H), 5.53-5.44 (m, 1H), 4.40-4.15 (m, 4H), 4.00 (s, 3H), 3.95 (s, 1H), 3.41 (s, 3H), 3.26-3.13 (m, 3H), 2.85-2.76 (m, 1H), 2.06 (s, 3H), 1.88-1.79 (m, 1H), 1.77-1.68 (m, 1H), 1.62-1.55 (m, 1H), 1.41-1.34 (m, 1H).	I-B41 I-C7
160		555.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.69 (m, 1H), 7.51-7.47 (m, 1H), 7.39-7.35 (m, 1H), 7.29 (s, 1H), 7.04-7.00 (m, 1H), 4.30 (s, 2H), 4.00-3.86 (m, 3H), 3.41 (s, 3H), 3.27-3.12 (m, 3H), 2.78-2.74 (m, 1H), 2.08-1.82 (m, 8H), 1.60-1.58 (m, 1H), 1.38-1.36 (m, 1H).	I-B22 I-C18
165		497.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.75 (dd, J = 8.1, 1.2 Hz, 1H), 7.54-7.41 (m, 3H), 7.31 (d, J = 7.9 Hz, 1H), 7.20 (d, J = 7.7 Hz, 1H), 4.50-4.41 (m, 2H), 3.92 (s, 1H), 3.43-3.35 (m, 3H), 3.18 (d, J = 16.0 Hz, 1H), 2.79 (d, J = 16.2 Hz, 1H), 2.04 (s, 3H), 1.96-1.73 (m, 2H), 1.60 (d, J = 13.5 Hz, 1H), 1.36 (d, J = 13.4 Hz, 1H).	I-B42 I-C3
166		541.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.71 (dd, J = 8.2, 1.4 Hz, 1H), 7.49 (t, J = 8.0 Hz, 1H), 7.39 (dd, J = 7.9, 1.4 Hz, 1H), 7.29 (s, 1H), 7.02 (d, J = 9.3 Hz, 1H), 5.49 (s, 1H), 4.51-4.14 (m, 4H), 3.95 (s, 1H), 3.41 (s, 3H), 3.27-3.15 (m, 3H), 2.78 (d, J = 16.4 Hz, 1H), 2.07 (s, 3H), 1.90-1.70 (m, 2H), 1.61 (d, J = 13.0 Hz, 1H), 1.37 (d, J = 12.8 Hz, 1H).	I-B1 I-C18


167		515.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.75 (dd, J = 8.1, 1.5 Hz, 1H), 7.51 (t, J = 8.0 Hz, 1H), 7.43 (dd, J = 8.0, 1.5 Hz, 1H), 7.32 (s, 1H), 7.04 (d, J = 9.3 Hz, 1H), 4.51-4.39 (m, 2H), 3.95 (s, 1H), 3.51 (s, 1H), 3.44-3.33 (dd, m, 2H), 3.22 (d, J = 16.3 Hz, 1H), 2.79 (d, J = 16.2 Hz, 1H), 2.04 (s, 3H), 1.95-1.77 (m, 2H), 1.62 (d, J = 13.4 Hz, 1H), 1.37 (d, J = 13.1 Hz, 1H).	I-B42 I-C19
168		511.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.75-7.68 (m, 1H), 7.52-7.46 (m, 1H), 7.41-7.35 (m, 1H), 7.32 (s, 1H), 7.08-7.00 (m, 1H), 4.01-3.86 (m, 3H), 3.52 (s, 1H), 3.28-3.15 (m, 3H), 2.82-2.70 (m, 1H), 2.06 (s, 3H), 2.01 (s, 3H), 1.97-1.89 (m, 1H), 1.89-1.80 (m, 1H), 1.63-1.56 (m, 1H), 1.40-1.34 (m, 1H).	I-B22 I-C19
169		541.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74 (dd, J = 8.1, 1.4 Hz, 1H), 7.51 (t, J = 8.0 Hz, 1H), 7.46-7.41 (m, 2H), 7.28 (d, J = 8.8 Hz, 1H), 7.20 (d, J = 7.7 Hz, 1H), 4.50-4.39 (m, 2H), 4.30 (s, 2H), 3.92 (s, 1H), 3.41 (s, 3H), 3.39-3.32 (m, 2H), 3.18 (d, J = 16.1 Hz, 1H), 2.79 (d, J = 16.1 Hz, 1H), 2.04 (s, 3H), 1.96-1.75 (m, 2H), 1.59 (d, J = 13.9 Hz, 1H), 1.36 (d, J = 12.5 Hz, 1H).	I-B42 I-C7
170		511.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74 (dd, J = 8.1, 1.5 Hz, 1H), 7.51 (t, J = 8.0 Hz, 1H), 7.43 (dd, J = 7.9, 1.5 Hz, 1H), 7.36 (s, 1H), 7.21-7.11 (m, 2H), 4.49-4.39 (m, 2H), 3.90 (s, 1H), 3.42-3.34 (m, 2H), 3.15 (d, J = 15.9 Hz, 1H), 2.77 (d, J = 15.8 Hz, 1H), 2.04 (s, 3H), 1.99 (s, 3H), 1.93-1.75 (m, 2H), 1.58 (d, J = 14.1 Hz, 1H), 1.37 (d, J = 13.0 Hz, 1H).	I-B42 I-C17

173		536.2	¹ H NMR (400 MHz, DMSO-d ₆): δ 8.66 (s, 1H), 7.79 (s, 1H), 7.59-7.43 (m, 3H), 7.40-7.32 (m, 1H), 7.30-7.23 (m, 1H), 5.40 (s, 1H), 4.16 (s, 1H), 3.84 (s, 1H), 3.17-3.02 (m, 4H), 2.69-2.63 (m, 3H), 1.98 (s, 1H), 1.91-1.69 (m, 3H), 1.63-1.48 (m, 2H), 1.23 (s, 3H), 1.08-0.96 (m, 1H).	I-B1 I-C20
174		550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.70 (m, 1H), 7.57 (s, 1H), 7.52-7.47 (m, 1H), 7.43-7.40 (m, 1H), 7.39-7.36 (m, 1H), 7.30-7.27 (m, 1H), 4.00-3.88 (m, 3H), 3.27-3.16 (m, 3H), 2.84-2.75 (m, 4H), 2.07 (s, 3H), 2.01 (s, 3H), 1.99-1.89 (m, 1H), 1.87-1.79 (m, 1H), 1.62-1.56 (m, 1H), 1.38-1.34 (m, 1H).	I-B22 I-C20
183		500.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.85-7.79 (m, 1H), 7.48 (s, 1H), 7.40-7.34 (m, 1H), 7.33-7.28 (m, 1H), 7.23-7.18 (m, 1H), 5.49 (s, 1H), 4.55-4.14 (m, 2H), 4.11 (s, 3H), 3.92 (s, 1H), 3.42 (s, 1H), 3.29-3.12 (m, 3H), 2.84-2.72 (m, 1H), 2.08 (s, 3H), 1.88-1.78 (m, 1H), 1.77-1.66 (m, 1H), 1.63-1.53 (m, 1H), 1.40-1.32 (m, 1H).	I-B43 I-C3 by-product
184		504.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.01-7.95 (m, 1H), 7.66-7.61 (m, 1H), 7.48 (s, 1H), 7.34-7.29 (m, 1H), 7.24-7.17 (m, 1H), 5.49 (s, 1H), 4.53-4.06 (m, 2H), 3.93 (s, 1H), 3.41 (s, 1H), 3.29-3.11 (m, 3H), 2.84-2.73 (m, 1H), 2.09 (s, 3H), 1.90-1.79 (m, 1H), 1.78-1.67 (m, 1H), 1.64-1.55 (m, 1H), 1.40-1.34 (m, 1H).	I-B44 I-C3

185		470.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.48 (s, 1H), 7.35-7.27 (m, 1H), 7.24-7.18 (m, 1H), 7.17-7.11 (m, 1H), 6.22-6.12 (m, 1H), 5.46 (s, 1H), 4.43-4.06 (m, 2H), 3.93 (s, 1H), 3.40 (s, 1H), 3.28-3.11 (m, 3H), 2.84-2.72 (m, 1H), 2.16 (s, 3H), 1.88-1.78 (m, 1H), 1.76-1.66 (m, 1H), 1.62-1.51 (m, 1H), 1.40-1.33 (m, 1H).	I-B45 I-C3 by-product
186		514.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.45 (s, 1H), 7.31-7.26 (m, 1H), 7.24-7.18 (m, 1H), 7.17-7.11 (m, 1H), 6.21-6.12 (m, 1H), 5.46 (s, 1H), 4.44-4.07 (m, 4H), 3.93 (s, 1H), 3.42 (s, 3H), 3.27-3.12 (m, 3H), 2.83-2.74 (m, 1H), 2.16 (s, 3H), 1.88-1.78 (m, 1H), 1.77-1.66 (m, 1H), 1.61-1.53 (m, 1H), 1.39-1.33 (m, 1H).	I-B45 I-C7 by-product
194		550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.71 (m, 1H), 7.60 (s, 1H), 7.50-7.45 (m, 2H), 7.40-7.38 (m, 1H), 7.32-7.30 (m, 1H), 5.49 (s, 1H), 4.28 (br, 2H), 3.97 (s, 1H), 3.33 (s, 3H), 3.26-3.20 (m, 3H), 3.01 (s, 3H), 2.84-2.82 (m, 1H), 2.07 (s, 3H), 1.91-1.68 (m, 2H), 1.65-1.55 (m, 1H), 1.41-1.31 (m, 1H).	I-B1 I-C21
195		564.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.70 (m, 1H), 7.60 (s, 1H), 7.53-7.43 (m, 2H), 7.40-7.36 (m, 1H), 7.32-7.28 (m, 1H), 4.01-3.86 (m, 3H), 3.33 (s, 3H), 3.27-3.18 (m, 3H), 3.02 (s, 3H), 2.85-2.81 (m, 1H), 2.10-1.85 (m, 8H), 1.61-1.57 (m, 1H), 1.38-1.34 (m, 1H).	I-B22 I-C21

197		528.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.64-7.61 (m, 1H), 7.45 (s, 1H), 7.30-7.27 (m, 1H), 7.24-7.20 (m, 2H), 5.48 (s, 1H), 4.43-4.19 (m, 4H), 4.18-4.17 (m, 3H), 3.93 (s, 1H), 3.43-3.40 (m, 3H), 3.28-3.20 (m, 2H), 3.19-3.14 (m, 1H), 2.81-2.76 (m, 1H), 2.23-2.06 (m, 3H), 1.88-1.80 (m, 1H), 1.76-1.69 (m, 1H), 1.61-1.55 (m, 1H), 1.39-1.34 (m, 1H).	I-B45 I-C7
198		488.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.96-7.89 (m, 1H), 7.55-7.45 (m, 2H), 7.34-7.29 (m, 1H), 7.24-7.18 (m, 1H), 5.49 (s, 1H), 4.54-3.97 (m, 2H), 3.93 (s, 1H), 3.41 (s, 1H), 3.30-3.11 (m, 3H), 2.85-2.71 (m, 1H), 2.10 (s, 3H), 1.89-1.80 (m, 1H), 1.78-1.68 (m, 1H), 1.63-1.55 (m, 1H), 1.42-1.33 (m, 1H).	I-B43 I-C22
202		538.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.56-7.53 (m, 1H), 7.49-7.42 (m, 1H), 7.42-7.38 (m, 1H), 7.33-7.28 (m, 1H), 7.28-7.25 (m, 1H), 5.47 (s, 1H), 4.27 (br, 2H), 3.94 (s, 1H), 3.26-3.15 (m, 3H), 2.84-2.76 (m, 4H), 2.07 (s, 3H), 1.88-1.79 (m, 1H), 1.77-1.67 (m, 1H), 1.61-1.55 (m, 1H), 1.37-1.31 (m, 1H).	I-B32 I-C20
206		489.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.45-7.39 (m, 2H), 7.28-7.16 (m, 3H), 6.95 (d, J = 7.8 Hz, 1H), 5.47 (s, 1H), 4.26 (br, 2H), 4.04 (s, 1H), 3.94 (s, 3H), 3.27-3.10 (m, 3H), 2.89-2.85 (m, 1H), 2.06-1.98 (m, 6H), 1.83-1.69 (m, 2H), 1.61-1.41 (m, 2H).	I-B34 I-C17

211		509.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.45 (m, 1H), 7.33-7.29 (m, 2H), 7.22-7.18 (m, 2H), 5.47 (s, 1H), 4.27 (br, 2H), 3.97 (s, 3H), 3.93 (s, 1H), 3.39 (s, 1H), 3.27-3.12 (m, 3H), 2.83-2.75 (m, 1H), 2.05 (s, 3H), 1.87-1.78 (m, 1H), 1.76-1.68 (m, 1H), 1.60-1.55 (m, 1H), 1.39-1.34 (m, 1H).	I-B46 I-C3
212		493.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.45 (s, 1H), 7.32-7.28 (m, 1H), 7.24-7.15 (m, 3H), 5.47 (s, 1H), 4.26 (br, 2H), 3.95 (s, 3H), 3.93 (s, 1H), 3.39 (s, 1H), 3.25-3.13 (m, 3H), 2.82-2.75 (m, 1H), 2.06 (s, 3H), 1.87-1.78 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.39-1.34 (m, 1H).	I-B47 I-C3
214		495.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.42 (m, 1H), 7.37-7.33 (m, 1H), 7.33-7.28 (m, 1H), 7.20-7.11 (m, 2H), 5.47 (s, 1H), 4.26 (br, 2H), 3.89 (s, 1H), 3.27-3.17 (m, 2H), 3.15-3.09 (m, 1H), 2.79-2.72 (m, 1H), 2.07 (s, 3H), 1.98 (s, 3H), 1.86-1.77 (m, 1H), 1.76-1.65 (m, 1H), 1.60-1.53 (m, 1H), 1.40-1.33 (m, 1H).	I-B32 I-C17
228		497.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.72-7.68 (m, 1H), 7.50-7.43 (m, 2H), 7.39-7.35 (m, 1H), 7.00 (d, J = 9.3 Hz, 1H), 5.47 (s, 1H), 4.25 (br, 2H), 3.88 (s, 1H), 3.66 (s, 1H), 3.25-3.10 (m, 3H), 2.79-2.77 (m, 1H), 2.05 (s, 3H), 1.85-1.67 (m, 2H), 1.58-1.54 (m, 1H), 1.38-1.34 (m, 1H).	I-B1 I-C26


245		480.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.75-7.68 (m, 1H), 7.55-7.45 (m, 2H), 7.38-7.31 (m, 2H), 7.27-7.21 (m, 1H), 5.20 (s, 1H), 4.28 (s, 2H), 3.96 (s, 1H), 3.44 (s, 1H), 3.28-3.15 (m, 3H), 2.86-2.77 (m, 1H), 1.91-1.69 (m, 2H), 1.62-1.54 (m, 1H), 1.41-1.36 (m, 1H).	I-B48 I-C3
-----	---	-------	--	---------------

The optically pure diastereomers in the table are subjected to chiral HPLC under the following conditions (flow rate: 15 mL/minute; detector: UV 254 nm):

Compounds	Column	Mobile phase	RT/minute	de%
20	IG (2 × 25 cm)	Acetonitrile/ethanol = 10/90	26.084	100%
24			17.095	100%
11	IG (2 × 25 cm)	Acetonitrile/ethanol = 10/90	16.291	100%
28			14.413	100%
12	IG (2 × 25 cm)	Acetonitrile/ethanol = 10/90	16.514	100%
64			11.431	100%
48	IG (2 × 25 cm)	Acetonitrile/ethanol = 10/90	5.640	100%
49			6.740	100%
50	IG (2 × 25 cm)	Acetonitrile/ethanol = 10/90	8.240	100%
51			10.836	100%

Compound 81

(S)-6-(1-amino-6-(oxetan-3-ylthio)-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-(2,3-dichlorophenyl)-2-methylpyrimidin-4(3H)-one



81

Step 1: (R)-N-((S)-5-bromo-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfonamide

The target product was prepared by following the step 1 for preparing compound 12 from corresponding starting materials and reagents. $[M+H]^+$ 639.1

Step 2: Ethyl 3-(((S)-1-(((R)-tert-butylsulfonyl)amino)-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)thio)propanoate

The target product was prepared by following the step 1 for preparing intermediate I-A1 from corresponding starting materials and reagents. $[M+H]^+$ 691.1

Step 3: Ethyl (S)-3-((1-amino-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)thio)propanoate

The target product was prepared by following the step 2 for preparing compound 12 from corresponding starting materials and reagents. $[M+H]^+$ 587.2




Step 4: Sodium (S)-1-amino-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-dihydropyrimidin-4-yl)-1,3-dihydrospiro[indene-2,4'-piperidine]-6-thiolate




5 The target product was prepared by following the step 2 for preparing intermediate I-A1 from corresponding starting materials and reagents.

Step 5: (S)-6-(1-amino-6-(oxetan-3-ylthio)-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-(2,3-dichlorophenyl)-2-methylpyrimidin-4(3H)-one

Sodium (S)-1-amino-1'-(1-(2,3-dichlorophenyl)-2-methyl-6-oxo-1,6-
10 dihydropyrimidin-4-yl)-1,3-dihydrospiro[indene-2,4'-piperidine]-6-thiolate was dissolved in *N,N*-dimethylformamide (3 mL), and 3-bromooxetane (93 mg, 0.68 mmol) was added thereto. The reaction was stirred at room temperature for 30 minutes, and water (10 mL) was added thereto. The mixture was extracted with dichloromethane. The organic phases were collected and combined, and concentrated in vacuum under reduced pressure, and
15 the residue was purified with silica gel column chromatography (dichloromethane/methanol) to give the target product (13 mg, yield 7.0%). $[M+H]^+$ 543.2. ^1H NMR (400 MHz, CD_3OD): δ 7.71 (dd, $J = 8.1, 1.4$ Hz, 1H), 7.48 (t, $J = 8.0$ Hz, 1H), 7.38 (dd, $J = 7.9, 1.4$ Hz, 1H), 7.29 (s, 1H), 7.21-7.10 (m, 2H), 5.48 (s, 1H), 5.04 (t, $J = 6.8$ Hz, 2H), 4.58-4.47 (m, 3H), 4.44-4.10 (m, 2H), 3.91 (s, 1H), 3.25-3.20 (m, 2H),
20 3.13 (d, $J = 15.9$ Hz, 1H), 2.75 (d, $J = 15.9$ Hz, 1H), 2.06 (s, 3H), 1.90-1.79 (m, 1H), 1.77-1.67 (m, 1H), 1.62-1.54 (m, 1H), 1.39-1.32 (m, 1H).


The compounds in the table below were prepared by following the steps for preparing compound 81 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Interme diates
37		523.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.1, 1.5 Hz, 1H), 7.48 (t, J = 8.0 Hz, 1H), 7.42-7.36 (m, 2H), 7.25 (d, J = 7.6 Hz, 1H), 7.18 (d, J = 7.7 Hz, 1H), 5.48 (s, 1H), 4.65 (q, J = 6.5 Hz, 1H), 4.44-4.10 (s, 2H), 3.91 (s, 1H), 3.25-3.24 (m, 1H), 3.23-3.21 (m, 1H), 3.17-3.12 (m, 1H), 2.80-2.74 (m, 1H), 2.06 (s, 3H), 1.86-1.68 (m, 2H), 1.61-1.53 (m, 1H), 1.45 (d, J = 6.6 Hz, 3H), 1.39-1.33 (m, 1H).	I-B1 I-C4
38		523.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.70 (dd, J = 8.1, 1.5 Hz, 1H), 7.48 (t, J = 8.0 Hz, 1H), 7.42-7.36 (m, 2H), 7.24 (dd, J = 7.7, 1H), 7.18 (d, J = 7.8 Hz, 1H), 5.46 (d, J = 9.9 Hz, 1H), 4.65 (q, J = 6.6 Hz, 1H), 4.41-4.10 (m, 2H), 3.90 (s, 1H), 3.26-3.17 (m, 2H), 3.14 (d, J = 16.1 Hz, 1H), 2.76 (d, J = 16.0 Hz, 1H), 2.05 (s, 3H), 1.86-1.68 (m, 2H), 1.60-1.54 (m, 1H), 1.45 (dd, J = 6.6, 3H), 1.38-1.32 (m, 1H).	I-B1 I-C4
47		481.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.71 (dd, J = 8.1, 1.5 Hz, 1H), 7.53-7.43 (m, 2H), 7.39 (dd, J = 7.9, 1.5 Hz, 1H), 7.31 (dd, J = 8.3, 1.8 Hz, 1H), 6.76 (d, J = 8.3 Hz, 1H), 5.54 (s, 1H), 4.58-4.19 (m, 2H), 4.10 (s, 1H), 3.48-3.35 (m, 2H), 3.30 (s, 1H), 1.97-1.87 (m, 2H), 1.84-1.77 (m, 2H).	I-B1 I-C9

67		522.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.73-7.71 (m, 1H), 7.63 (s, 1H), 7.51-7.45 (m, 2H), 7.40-7.36 (m, 2H), 5.51 (s, 1H), 4.44-4.30 (m, 3H), 4.07 (s, 2H), 3.27-3.08 (m, 4H), 2.75 (s, 3H), 2.07 (s, 3H), 1.82-1.56 (m, 4H).	I-B1 I-C4
79		493.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.74-7.68 (m, 1H), 7.51-7.45 (m, 1H), 7.41-7.37 (m, 1H), 7.36 (s, 1H), 7.21-7.16 (m, 1H), 7.16-7.12 (m, 1H), 5.48 (s, 1H), 4.52-4.01 (m, 2H), 3.90 (s, 1H), 3.28-3.17 (m, 2H), 3.16-3.10 (m, 1H), 2.81-2.72 (m, 1H), 2.06 (s, 3H), 1.99 (s, 3H), 1.86-1.78 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.39-1.34 (m, 1H).	I-B1 I-C4
91		545.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.03 (s, 1H), 7.83 (d, J = 1.6 Hz, 1H), 7.54 (d, J = 1.6 Hz, 1H), 7.51-7.44 (m, 2H), 7.31-7.19 (m, 2H), 5.87 (d, J = 5.6 Hz, 1H), 4.06-3.95 (m, 3H), 3.61 (s, 2H), 3.45-3.31 (m, 3H), 3.22-3.15 (m, 1H), 2.49 (s, 3H), 2.14-1.96 (m, 2H), 1.72-1.68 (m, 1H), 1.52-1.48 (m, 1H).	I-A1 I-C4

Compound 87

(S)-(3-(1-amino-6-(3-methoxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-6-((2-aminopyrimidin-4-yl)thio)pyrazin-2-yl)methanol



- 5 **Step 1: Methyl 6-bromo-3-((S)-1-(((R)-tert-butylsulfinyl)amino)-6-(3-methoxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)pyrazine-2-carboxylate**

The target product was prepared by following the step 1 for preparing compound 1 from corresponding starting materials and reagents. $[M+H]^+$ 589.1





- 10 **Step 2: (R)-N-((S)-1'-(5-bromo-3-(hydroxymethyl)pyrazin-2-yl)-5-(3-methoxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide**

- Under nitrogen, at -78°C , to a solution of methyl 6-bromo-3-((S)-1-(((R)-tert-butylsulfinyl)amino)-6-(3-methoxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)pyrazine-2-carboxylate (380 mg, 0.64 mmol) in anhydrous THF (15 mL) was added dropwise 1.0 M DIBAL-H solution (3.2 mL, 3.2 mmol). The reaction mixture was stirred at -78°C for 2 hours, warmed to room temperature and stirred for 20 minutes. $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ and water (20 mL) were added thereto, and the mixture was filtered. The aqueous layer was extracted with dichloromethane. The organic phases were collected and combined, washed with saturated brine, dried over anhydrous Na_2SO_4 , and concentrated in vacuum under reduced pressure. The residue was purified with silica gel column chromatography (water/methanol) to give the target product (186 mg, yield 52%). $[M+H]^+$ 561.2

Step 3: (S)-(3-(1-amino-6-(3-methoxyprop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-6-((2-aminopyrimidin-4-yl)thio)pyrazin-2-yl)methanol


The target product was prepared by following the steps 2-3 for preparing compound 1 from corresponding starting materials and reagents. $[M+H]^+$ 504.2. ^1H NMR (400 MHz, CD_3OD): δ 8.31 (s, 1H), 7.90 (d, $J = 5.5$ Hz, 1H), 7.45 (s, 1H), 7.27 (d, $J = 7.7$ Hz, 1H), 7.19 (d, $J = 7.8$ Hz, 1H), 6.28 (d, $J = 5.5$, 1H), 4.66 (s, 2H), 4.30 (s, 2H), 3.93 (s, 1H), 3.84- 3.75 (m, 2H), 3.41 (s, 3H), 3.26-3.17 (m, 2H), 3.13 (d, $J = 16.0$ Hz, 1H), 2.76 (d, $J = 16.1$ Hz, 1H), 2.03-1.83 (m, 2H), 1.61 (d, $J = 12.2$ Hz, 1H), 1.39 (d, $J = 12.6$ Hz, 1H).

The compounds in the table below were prepared by following the steps for preparing compound 87 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Intermediates
116		460.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.31 (s, 1H), 7.90 (d, J = 5.5 Hz, 1H), 7.48 (s, 1H), 7.30 (d, J = 7.7 Hz, 1H), 7.19 (d, J = 7.7 Hz, 1H), 6.28 (d, J = 5.5 Hz, 1H), 4.66 (s, 2H), 3.93 (s, 1H), 3.85-3.76 (m, 2H), 3.39 (s, 1H), 3.26-3.18 (m, 2H), 3.14 (d, J = 16.1 Hz, 1H), 2.76 (d, J = 16.1 Hz, 1H), 2.03-1.94 (m, 1H), 1.93-1.82 (m, 1H), 1.61 (d, J = 12.6 Hz, 1H), 1.39 (d, J = 13.4 Hz, 1H).	I-A6 I-C3
122		503.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.33-8.29 (m, 1H), 7.91-7.89 (m, 1H), 7.59-7.55 (m, 1H), 7.44-7.40 (m, 1H), 7.29-7.26 (m, 1H), 6.29-6.27 (m, 1H), 4.67 (s, 2H), 3.96 (s, 1H), 3.81 (br, 2H), 3.25-3.15 (m, 3H), 2.84-2.75 (m, 1H), 2.03-1.98 (m, 1H), 1.94-1.84 (m, 1H), 1.65-1.59 (m, 1H), 1.41-1.35 (m, 1H).	I-A6 I-C16
123		536.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.28-8.27 (m, 1H), 7.61-7.59 (m, 1H), 7.58-7.56 (m, 1H), 7.44-7.40 (m, 1H), 7.29-7.26 (m, 1H), 6.05-6.02 (m, 1H), 4.67 (s, 2H), 3.97 (s, 1H), 3.84 (br, 2H), 3.26-3.15 (m, 3H), 2.83-2.77 (m, 1H), 2.04-1.97 (m, 1H), 1.92-1.84 (m, 1H), 1.65-1.59 (m, 1H), 1.41-1.35 (m, 1H).	I-A1 I-C16
199		564.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.62-7.54 (m, 2H), 7.41 (d, J = 7.7 Hz, 1H), 7.28 (d, J = 7.8 Hz, 1H), 5.88 (d, J = 5.6 Hz, 1H), 4.64 (s, 2H), 3.96 (s, 1H), 3.93-2.84 (m, 2H), 3.28-3.15 (m, 3H), 2.84-2.77 (m, 4H), 2.48 (s, 3H), 2.04-1.84 (m, 2H), 1.61 (d, J = 12.3 Hz, 1H), 1.38 (d, J = 12.9 Hz, 1H).	I-A1 I-C20

Compound 159

(*S*)-1-(3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)prop-2-yn-1-yl)urea



159

5 Step 1: *N,N*-bis(4-methoxybenzyl)prop-2-yn-1-amine

To the solution of propargylamine (550 mg, 10 mmol) in MeCN (30 mL) was added *p*-methoxybenzyl chloride (3.4 g, 22 mmol) and K₂CO₃ (4.1 g, 30 mmol). The reaction solution was heated to 60 °C and stirred for 16 hours. After cooling to room temperature, added water (100 mL), extracted with EA (100 mL x 2). The organic layer was collected, combined, and concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with PE/EA) to give target product (2.1 g, 71% yield). [M+H]⁺ 296.1

Step 2: *N*-(4-methoxybenzyl)-*N*-(prop-2-yn-1-yl)cyanamide

N,N-bis(4-methoxybenzyl)prop-2-yn-1-amine (2.1 g, 7.1 mmol), BrCN (1.5 g, 14.2 mmol) and K₂CO₃ (2.2 g, 16.3 mmol) were placed in 1,4-dioxane (70 mL), stirred at room temperature for 20 hours, added water (50 mL), extracted with EA (50 mL x 2). The organic layer was collected, combined, and concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with PE/EA) to give target product (750 mg, 50% yield). [M+Na]⁺ 223.2

Step 3: (*R*)-*N*-((*S*)-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-5-(3-(*N*-(4-methoxybenzyl)cyanamido)prop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide

Under N₂, (*R*)-*N*-((*S*)-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-5-bromo-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide (424 mg, 0.67 mmol; prepared by following the step 1 for preparing compound 1 from intermediates I-A2 and I-C4), *N*-(4-methoxybenzyl)-*N*-(prop-2-yn-1-yl)cyanamide (200 mg, 1 mmol), Pd(PPh₃)₂Cl₂ (94 mg, 0.13 mmol), CuI (25 mg, 0.13 mmol) and DIEA (1 mL) were placed in DMF (3 mL). The reaction solution was heated to 90 °C and stirred for 20 hours. The mixture was concentrated in vacuum under reduced pressure. The residue was purified by silica gel column chromatography (eluting with water/MeOH) to give target product (70 mg, 14% yield). [M+H]⁺ 756.3

Step 4: (*S*)-*N*-(3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)prop-2-yn-1-yl)-*N*-(4-methoxybenzyl)cyanamide

(*R*)-*N*-((*S*)-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-5-(3-(*N*-(4-methoxybenzyl)cyanamido)prop-1-yn-1-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-3-yl)-2-methylpropane-2-sulfinamide (70 mg, 0.09 mmol) was dissolved in 2 M HCl/MeOH solution (1 mL) and stirred at room temperature for 10 minutes. The mixture was adjusted to pH = 9 with aqueous ammonia, added water (10 mL), extracted with DCM (10 mL x 2). The organic layer was collected, combined, and concentrated in vacuum under reduced pressure to give target product (50 mg, 85% yield). [M+H]⁺ 652.2

Step 5: (*S*)-1-(3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)prop-2-yn-1-yl)urea

To (*S*)-*N*-(3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)prop-2-yn-1-yl)-*N*-(4-methoxybenzyl)cyanamide (50 mg, 0.07 mmol) was added TFA (2 mL) and stirred at 50 °C for 1 hour. After cooling to room temperature, the mixture was concentrated in vacuum under reduced pressure, added saturated NaHCO₃ aqueous solution (10 mL), extracted with DCM (10 mL x 2). The organic layer was collected, combined, and concentrated in vacuum under reduced pressure. The residue was purified by thin layer


chromatography (eluting with DCM/MeOH) to give target product (10 mg, 25% yield).

$[M+H]^+$ 550.2. ^1H NMR (400 MHz, CD_3OD): δ 7.59-7.56 (m, 2H), 7.42 (s, 1H), 7.30-7.25 (m, 1H), 7.22-7.18 (m, 1H), 5.94-5.86 (m, 1H), 4.33-4.24 (m, 2H), 4.09 (s, 2H), 3.96 (s, 1H), 3.25-3.13 (m, 3H), 2.84-2.80 (m, 1H), 1.88-1.67 (m, 2H), 1.58-1.56 (m, 1H),

5 1.40-1.38 (m, 1H).

Compound 222

(S)-3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)-3-chloropyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-methylpropiolamide



Step 1: 3-((S)-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)-3-chloropyrazin-2-yl)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-methylpropiolamide

To the solution of 3-((S)-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-methylpropiolamide (96 mg, 0.15 mmol; prepared by following the step 1 for preparing compound 1 from intermediates I-A2 and I-C20) in DMF (3 mL) was added NCS (40 mg, 0.30 mmol) and stirred at room temperature for 30 minutes. The reaction solution was purified by silica gel column chromatography (eluting with water/MeOH) to give target product.

15

20




Step 2: (S)-3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)-3-chloropyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-methylpropiolamide

The target product was prepared by following the step 3 for preparing compound 1 from corresponding starting materials and reagents (26 mg, 31% yield of two steps).

25


[M+H]⁺ 569.2. ¹H NMR (400 MHz, CD₃OD): δ 7.60 (d, J = 5.5 Hz, 1H), 7.55 (s, 1H), 7.40 (d, J = 7.7 Hz, 1H), 7.26 (d, J = 7.8 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 4.08-4.01 (m, 2H), 3.94 (s, 1H), 3.22-3.12 (m, 3H), 2.81-2.74 (m, 4H), 1.99-1.82 (m, 2H), 1.58 (d, J = 13.4 Hz, 1H), 1.36 (d, J = 13.2 Hz, 1H).

- 5 The compounds in the table below were prepared by following the steps for preparing compound 222 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Interme diates
223		587.2	¹ H NMR (400 MHz, CD ₃ OD) δ 7.62-7.51 (m, 2H), 7.07 (d, J = 9.3 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 4.06-4.02 (m, 2H), 3.92 (s, 1H), 3.21-3.11 (m, 3H), 2.84-2.80 (m, 1H), 2.78 (s, 3H), 1.96-1.81 (m, 2H), 1.59-1.55 (m, 1H), 1.40-1.36 (m, 1H).	I-A2 I-C24
224		583.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.61 (d, J = 5.5 Hz, 1H), 7.55 (s, 1H), 7.40 (d, J = 7.9 Hz, 1H), 7.26 (d, J = 7.7 Hz, 1H), 5.95 (d, J = 5.5 Hz, 1H), 4.08-4.02 (m, 2H), 3.94 (s, 1H), 3.26-3.10 (m, 5H), 2.78 (d, J = 16.0 Hz, 1H), 2.03-1.82 (m, 2H), 1.58 (d, J = 15.0 Hz, 1H), 1.36 (d, J = 15.3 Hz, 1H), 1.15 (t, J = 7.3 Hz, 3H).	I-A2 I-C23
230		595.2	¹ H NMR (400 MHz, CD ₃ OD/CDCl ₃): δ 7.60-7.58 (m, 1H), 7.53 (s, 1H), 7.41-7.35 (m, 1H), 7.27-7.22 (m, 1H), 5.96-5.92 (m, 1H), 4.07-4.00 (m, 2H), 3.93 (s, 1H), 3.21-3.11 (m, 3H), 2.81-2.70 (m, 2H), 1.98-1.82 (m, 2H), 1.59-1.53 (m, 1H), 1.37-1.32 (m, 1H), 0.78-0.72 (m, 2H), 0.60-0.52 (m, 2H).	I-A2 I-C25

Compound 239


(*S*)-3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)-3-fluoropyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-methylpropiolamide



At 0°C, to the solution of (*S*)-3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-methylpropiolamide (compound 229; 165 mg, 0.31 mmol) in DMF (2 mL) and MeCN (3 mL) was added NFSI (97 mg, 0.31 mmol) and stirred at room temperature for 3 hours. The reaction solution was purified by silica gel column chromatography (eluting with water/MeOH) and thin layer chromatography (eluting with DCM/MeOH) to give target product (28 mg, 16% yield). $[M+H]^+$ 553.2. ^1H NMR (400 MHz, CD_3OD): δ 7.63-7.57 (m, 1H), 7.54 (s, 1H), 7.42-7.38 (m, 1H), 7.26 (d, $J = 7.7$ Hz, 1H), 5.99 (d, $J = 5.5$ Hz, 1H), 4.30-4.26 (m, 2H), 3.94 (s, 1H), 3.34-3.32 (m, 1H), 3.26-3.01 (m, 2H), 2.82-2.78 (m, 4H), 1.95-1.72 (m, 2H), 1.59-1.55 (m, 1H), 1.35-1.31 (m, 1H).

Compound 240

(*S*)-3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)-3-bromopyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-methylpropiolamide




To the solution of (*S*)-3-(1-amino-1'-(6-amino-5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-*N*-methylpropiolamide (compound 229; 60 mg, 0.093 mmol) in DMF (1.5 mL) was added NBS (33 mg, 0.188 mmol) and stirred at room temperature for 1 hour. The reaction solution was purified by silica gel column chromatography (eluting with water/MeOH) and thin layer chromatography (eluting with DCM/MeOH) to give target product (15 mg,

26% yield). $[M+H]^+$ 613.1. ^1H NMR (400 MHz, CD_3OD): δ 7.62 (d, $J = 5.5$ Hz, 1H), 7.57 (s, 1H), 7.41 (d, $J = 7.7$ Hz, 1H), 7.27 (d, $J = 7.8$ Hz, 1H), 5.96 (d, $J = 5.6$ Hz, 1H), 4.09-3.91 (m, 3H), 3.24-3.09 (m, 3H), 2.87-2.73 (m, 4H), 2.04-1.80 (m, 2H), 1.62-1.58 (m, 1H), 1.41-1.37 (m, 1H).

5

Compound 243

(S)-3-(1-amino-1'-(5-(3,4-dihydro-1,5-naphthyridin-1(2H)-yl)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-ethylpropiolamide



10 **Step 1: 3-((S)-1'-((R)-tert-butylsulfinyl)amino)-1'-(5-(3,4-dihydro-1,5-naphthyridin-1(2H)-yl)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-ethylpropiolamide**

Under N_2 , 3-((S)-1'-(5-bromopyrazin-2-yl)-1-((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-ethylpropiolamide (80 mg, 0.14 mmol; 15 prepared by following the step 1 for preparing compound 1 from 2,5-dibromopyrazine and intermediate I-C23), 1,2,3,4-tetrahydro-1,5-naphthyridine (37 mg, 0.28 mmol), $\text{Pd}_2(\text{dba})_3$ (9 mg, 0.01 mmol), Xant-phos (12 mg, 0.02 mmol) and Cs_2CO_3 (91 mg, 0.28 mmol) were placed in 1,4-dioxane (10 mL). The reaction mixture was heated to 100°C and stirred for 16 hours. After concentration in vacuum under reduced pressure, the 20 residue was purified by silica gel column chromatography (eluting with water/MeOH, 0.05% formic acid) to give target product.

Step 2: (S)-3-(1-amino-1'-(5-(3,4-dihydro-1,5-naphthyridin-1(2H)-yl)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)-N-ethylpropiolamide


The target product was prepared by following the step 3 for preparing compound 1 from corresponding starting materials and reagents (25 mg, 35% yield). $[M+H]^+$ 508.3. ^1H NMR (400 MHz, CD_3OD): δ 8.06 (d, $J = 2.4$ Hz, 2H), 7.83 (dd, $J = 4.7, 1.3$ Hz, 1H), 7.56 (s, 1H), 7.42 (d, $J = 7.6$ Hz, 1H), 7.28 (d, $J = 7.8$ Hz, 1H), 7.08 (dd, $J = 8.4, 1.2$ Hz, 1H), 7.03-6.95(m, 1H), 4.25-4.12 (m, 2H), 3.95 (s, 1H), 3.72-3.65 (m, 2H), 3.29-3.14 (m, 5H), 2.99 (t, $J = 6.5$ Hz, 2H), 2.81 (d, $J = 16.3$ Hz, 1H), 2.18-2.09 (m, 2H), 1.96-1.75 (m, 2H), 1.60 (d, $J = 12.5$ Hz, 1H), 1.37 (d, $J = 12.7$ Hz, 1H), 1.16 (t, $J = 7.3$ Hz, 3H).

The compounds in the table below were prepared by following the steps for preparing compound 243 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Interme diates
244		509.3	¹ H NMR (400 MHz, CD ₃ OD): δ 8.37 (s, 1H), 7.94 (d, J = 4.6 Hz, 1H), 7.55 (s, 1H), 7.41 (d, J = 7.5 Hz, 1H), 7.28 (t, J = 8.8 Hz, 2H), 7.10-7.02 (m, 1H), 4.58-4.46 (m, 2H), 3.93 (s, 1H), 3.85-3.77 (m, 2H), 3.28-3.18 (m, 5H), 2.99 (t, J = 6.6 Hz, 2H), 2.81 (d, J = 16.4 Hz, 1H), 2.19-2.09 (m, 2H), 1.93-1.70 (m, 2H), 1.58 (d, J = 12.1 Hz, 1H), 1.34 (d, J = 12.1 Hz, 1H), 1.15 (t, J = 7.3 Hz, 3H).	I-C23
251		494.3	¹ H NMR (400 MHz, CD ₃ OD): δ 8.11-8.06 (m, 2H), 7.88-7.85 (m, 1H), 7.60 (s, 1H), 7.47-7.42 (m, 1H), 7.33-7.28 (m, 1H), 7.14-7.09 (m, 1H), 7.05-6.99 (m, 1H), 4.27-4.14 (m, 2H), 3.98 (s, 1H), 3.76-3.67 (m, 2H), 3.29-3.16 (m, 3H), 3.05-2.98 (m, 2H), 2.88-2.79 (m, 4H), 2.21-2.12 (m, 2H), 1.99-1.87 (m, 1H), 1.86-1.75 (m, 1H), 1.67-1.58 (m, 1H), 1.43-1.37 (m, 1H).	I-C20
258		522.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.00 (s, 1H), 7.96-7.89 (m, 1H), 7.57 (s, 1H), 7.49-7.37 (m, 2H), 7.30-7.23 (m, 1H), 7.09-7.02 (m, 1H), 3.97 (s, 1H), 3.81-3.74 (m, 2H), 3.37-3.32 (m, 2H), 3.30-3.25 (m, 2H), 3.18-3.11 (m, 1H), 3.06-2.94 (m, 4H), 2.82-2.72 (m, 1H), 2.47 (s, 3H), 2.14-2.07 (m, 2H), 2.06-1.99 (m, 1H), 1.96-1.86 (m, 1H), 1.66-1.58 (m, 1H), 1.44-1.36 (m, 1H), 1.25-1.11 (m, 3H).	I-C23

Compound 279

(S)-3-(1-amino-1'-(5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)propionic acid




5 Step 1: 3-((S)-1'-(5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)propionic acid

To the solution of ethyl 3-((S)-1'-(5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1-(((R)-tert-butylsulfinyl)amino)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)propionate (320 mg, 0.5 mmol; prepared by following the step 1 for preparing
 10 compound 1 from intermediates I-A15 and I-C30) in EtOH/water was added LiOH (120 mg, 5.0 mmol). The reaction solution was stirred at 70°C for 1 hour, adjusted pH to 5 with MeSO₃H. After concentration in vacuum under reduced pressure, the residue was purified by silica gel column chromatography (eluting with DCM/MeOH) to give target product (80 mg, 26% yield).

15 Step 2: (S)-3-(1-amino-1'-(5-((2-amino-3-chloropyridin-4-yl)thio)pyrazin-2-yl)-1,3-dihydrospiro[indene-2,4'-piperidin]-6-yl)propionic acid


The target product was prepared by following the step 3 for preparing compound 1 from corresponding starting materials and reagents (30 mg, 45% yield). [M+H]⁺ 507.2. ¹H NMR (400 MHz, CD₃OD): δ 8.45-8.30 (m, 2H), 7.73-7.54 (m, 3H), 7.49-7.38 (m, 1H),
 20 6.00-5.91 (m, 1H), 4.55-4.46 (m, 2H), 4.41-4.34 (m, 1H), 3.46-3.38 (m, 2H), 3.27-3.21 (m, 2H), 1.91-1.67 (m, 4H).

The compounds in the table below were prepared by following the steps for preparing compound 279 from corresponding intermediates and reagents:

Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	Interme diates
283		536.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.66 (s, 1H), 7.62 (d, J = 5.6 Hz, 1H), 7.57-7.54 (m, 1H), 7.41-7.38 (m, 1H), 5.90 (d, J = 5.6 Hz, 1H), 4.41 (s, 1H), 3.83-3.68 (m, 2H), 3.25-3.10 (m, 4H), 2.42 (s, 3H), 2.04-1.91 (m, 2H), 1.74-1.63 (m, 2H).	I-A11 I-C31

Compounds 56 and 57

Diastereomers of (S)-6-(1-amino-6-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-(2-chlorophenyl)-2-methylpyrimidin-4(3H)-one



5

(S)-6-(1-amino-6-ethynyl-1,3-dihydrospiro[indene-2,4'-piperidin]-1'-yl)-3-(2-chlorophenyl)-2-methylpyrimidin-4(3H)-one (compound 42) was resolved by chiral HPLC to obtain a pair of diastereomers. Chiral HPLC conditions: column: IC (2 × 25 cm); mobile phase: acetonitrile/ethanol = 10 : 90; flow rate: 15 mL/minute; detector: UV 254

10 nm.





First eluent (compound 56, RT = 6.942 minutes), de% = 100%, [M+H]⁺ 445.2. ¹H NMR (400 MHz, CD₃OD): δ 7.65-7.59 (m, 1H), 7.53-7.46 (m, 3H), 7.41-7.36 (m, 1H), 7.33-7.28 (m, 1H), 7.22-7.17 (m, 1H), 5.48 (s, 1H), 4.27 (br, 2H), 3.92 (s, 1H), 3.39 (s, 1H), 3.27-3.13 (m, 3H), 2.82-2.74 (m, 1H), 2.04 (s, 3H), 1.88-1.79 (m, 1H), 1.77-1.67 (m, 1H), 1.61-1.54 (m, 1H), 1.39-1.32 (m, 1H).





15





Second eluent (compound 57, RT = 9.352 minutes), de% = 100%, [M+H]⁺ 445.2. ¹H NMR (400 MHz, CD₃OD): δ 7.66-7.60 (m, 1H), 7.54-7.45 (m, 3H), 7.41-7.36 (m, 1H), 7.33-7.27 (m, 1H), 7.22-7.17 (m, 1H), 5.48 (s, 1H), 4.27 (br, 2H), 3.93 (s, 1H), 3.39 (s, 1H), 3.27-3.12 (m, 3H), 2.83-2.74 (m, 1H), 2.04 (s, 3H), 1.88-1.78 (m, 1H), 1.77-1.68 (m, 1H), 1.61-1.54 (m, 1H), 1.40-1.33 (m, 1H).



20





The compounds in the table below were prepared by following the chiral resolution conditions of compounds 56 and 57:







Compounds	Structural formula	LC-MS [M+H] ⁺	¹ H NMR	
58		475.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.52-7.45 (m, 2H), 7.33-7.27 (m, 1H), 7.21-7.19 (m, 1H), 7.10-7.04 (m, 1H), 7.01-6.97 (m, 1H), 5.48 (s, 1H), 4.26 (br, 2H), 3.92 (s, 1H), 3.82 (s, 3H), 3.39 (s, 1H), 3.26-3.11 (m, 3H), 2.81-2.74 (m, 1H), 2.08 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.39-1.33 (m, 1H).	Obtained by resolution of compound 44
59		475.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.52-7.45 (m, 2H), 7.32-7.28 (m, 1H), 7.21-7.17 (m, 1H), 7.09-7.05 (m, 1H), 7.01-6.97 (m, 1H), 5.48 (s, 1H), 4.26 (br, 2H), 3.92 (s, 1H), 3.82 (s, 3H), 3.39 (s, 1H), 3.26-3.12 (m, 3H), 2.81-2.74 (m, 1H), 2.08 (s, 3H), 1.87-1.78 (m, 1H), 1.76-1.68 (m, 1H), 1.61-1.54 (m, 1H), 1.39-1.33 (m, 1H).	
65		463.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.66-7.63 (m, 1H), 7.49 (s, 1H), 7.40-7.26 (m, 3H), 7.24-7.20 (m, 1H), 5.48 (s, 1H), 4.30-4.26 (m, 2H), 4.00 (s, 1H), 3.41 (s, 1H), 3.26-3.13 (m, 3H), 2.86-2.82 (m, 1H), 2.12 (s, 3H), 1.89-1.65 (m, 2H), 1.60-1.56 (m, 1H), 1.42-1.36 (m, 1H).	Obtained by resolution of compound 16
66		463.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.68-7.62 (m, 1H), 7.55 (s, 1H), 7.44-7.40 (m, 1H), 7.37-7.26 (m, 3H), 5.50 (s, 1H), 4.40-4.16 (m, 3H), 3.48 (s, 1H), 3.27-2.96 (m, 4H), 2.12 (s, 3H), 1.80-1.74 (m, 2H), 1.61-1.53 (m, 2H).	





88		477.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.44 (m, 1H), 7.33-7.27 (m, 1H), 7.21-7.12 (m, 2H), 7.10-7.04 (m, 1H), 5.47 (s, 1H), 4.26 (br, 2H), 4.01 (s, 3H), 3.92 (s, 1H), 3.39 (s, 1H), 3.26-3.13 (m, 3H), 2.81-2.75 (m, 1H), 2.12 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.60-1.54 (m, 1H), 1.38-1.32 (m, 1H).	Obtained by resolution of compound 78
89		477.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.51-7.44 (m, 1H), 7.34-7.27 (m, 1H), 7.22-7.12 (m, 2H), 7.10-7.03 (m, 1H), 5.48 (s, 1H), 4.27 (br, 2H), 4.01 (s, 3H), 3.94 (s, 1H), 3.40 (s, 1H), 3.24-3.13 (m, 3H), 2.84-2.75 (m, 1H), 2.12 (s, 3H), 1.88-1.78 (m, 1H), 1.76-1.66 (m, 1H), 1.61-1.53 (m, 1H), 1.39-1.32 (m, 1H).	
102		477.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.53-7.36 (m, 3H), 7.34-7.23 (m, 2H), 7.20-7.18 (m, 1H), 3.95-3.85 (m, 3H), 3.40 (s, 1H), 3.26-3.07 (m, 3H), 2.79-2.75 (m, 1H), 2.08-1.74 (m, 8H), 1.59-1.55 (m, 1H), 1.38-1.34 (m, 1H).	Obtained by resolution of compound 54
103		477.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.62-7.39 (m, 3H), 7.36-7.24 (m, 2H), 7.22-7.18 (m, 1H), 3.98-3.81 (m, 3H), 3.41 (s, 1H), 3.27-3.08 (m, 3H), 2.79-2.75 (m, 1H), 2.09-1.74 (m, 8H), 1.58-1.54 (m, 1H), 1.38-1.34 (m, 1H).	







108		497.2	¹ H NMR (400 MHz, CD ₃ OD) δ 7.53-7.38 (m, 3H), 7.31 (d, J = 7.4 Hz, 1H), 7.20 (d, J = 7.5 Hz, 1H), 5.48 (s, 1H), 4.27 (br, 2H), 3.94 (s, 1H), 3.40 (s, 1H), 3.30-3.10 (m, 3H), 2.81-2.77 (m, 1H), 2.07 (s, 3H), 1.90-1.67 (m, 2H), 1.60-1.56 (m, 1H), 1.39-1.35 (m, 1H).	Obtained by resolution of compound 69
109		497.2	¹ H NMR (400 MHz, CD ₃ OD) δ 7.55-7.41 (m, 3H), 7.32 (d, J = 7.5 Hz, 1H), 7.21 (d, J = 7.5 Hz, 1H), 5.48 (s, 1H), 4.28 (br, 2H), 3.95 (s, 1H), 3.41 (s, 1H), 3.27-3.09 (m, 3H), 2.87-2.76 (m, 1H), 2.07 (s, 3H), 1.90-1.63 (m, 2H), 1.60-1.56 (m, 1H), 1.40-1.36 (m, 1H).	
110		475.2	¹ H NMR (400 MHz, CD ₃ OD) δ 7.55-7.20 (m, 5H), 6.98-6.94 (m, 1H), 5.48 (s, 1H), 4.27 (br, 2H), 4.02-3.94 (m, 4H), 3.42 (s, 1H), 3.26-3.12 (m, 3H), 2.87-2.83 (m, 1H), 2.05 (s, 3H), 1.90-1.63 (m, 2H), 1.60-1.56 (m, 1H), 1.43-1.39 (m, 1H).	Obtained by resolution of compound 90
111		475.2	¹ H NMR (400 MHz, CD ₃ OD) δ 7.52-7.18 (m, 5H), 6.98-6.94 (m, 1H), 5.48 (s, 1H), 4.20 (br, 2H), 3.99-3.93 (m, 4H), 3.42 (s, 1H), 3.26-3.16 (m, 3H), 2.84-2.80 (m, 1H), 2.05 (s, 3H), 1.90-1.63 (m, 2H), 1.60-1.56 (m, 1H), 1.41-1.37 (m, 1H).	

113		481.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.51-7.42 (m, 2H), 7.34-7.28 (m, 2H), 7.22-7.17 (m, 1H), 5.48 (s, 1H), 4.28 (s, 2H), 3.92 (s, 1H), 3.40 (s, 1H), 3.27-3.12 (m, 3H), 2.83-2.74 (m, 1H), 2.08 (s, 3H), 1.90-1.80 (m, 1H), 1.77-1.66 (m, 1H), 1.62-1.54 (m, 1H), 1.40-1.33 (m, 1H).	Obtained by resolution of compound 77
114		481.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.53-7.39 (m, 2H), 7.34-7.27 (m, 2H), 7.22-7.17 (m, 1H), 5.48 (s, 1H), 4.27 (br, 2H), 3.92 (s, 1H), 3.40 (s, 1H), 3.26-3.12 (m, 3H), 2.83-2.74 (m, 1H), 2.08 (s, 3H), 1.88-1.78 (m, 1H), 1.76-1.67 (m, 1H), 1.61-1.54 (m, 1H), 1.39-1.33 (m, 1H).	
118		513.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.08-7.93 (m, 1H), 7.83-7.64 (m, 2H), 7.52 (s, 1H), 7.36 (d, J = 7.6 Hz, 1H), 7.25 (d, J = 7.6 Hz, 1H), 5.54 (s, 1H), 4.35 (s, 2H), 3.98 (s, 1H), 3.45 (s, 1H), 3.32-3.17 (m, 3H), 2.88-2.75 (m, 1H), 2.10 (s, 3H), 1.95-1.71 (m, 2H), 1.68-1.58 (m, 1H), 1.46-1.37 (m, 1H).	Obtained by resolution of compound 106
119		513.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.05-7.98 (m, 1H), 7.79-7.69 (m, 2H), 7.53 (s, 1H), 7.36 (d, J = 7.8 Hz, 1H), 7.25 (d, J = 7.8 Hz, 1H), 5.55 (s, 1H), 4.34 (s, 2H), 3.99 (s, 1H), 3.46 (s, 1H), 3.32-3.16 (m, 3H), 2.89-2.78 (m, 1H), 2.10 (s, 3H), 1.94-1.72 (m, 2H), 1.68-1.60 (m, 1H), 1.47-1.39 (m, 1H).	

124		541.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.53-7.42 (m, 3H), 7.32-7.18 (m, 2H), 5.48 (s, 1H), 4.50-4.10 (m, 4H), 3.93 (s, 1H), 3.41 (s, 3H), 3.26-3.13 (m, 3H), 2.81-2.75 (m, 1H), 2.07 (s, 3H), 1.90-1.65 (m, 2H), 1.60-1.56 (m, 1H), 1.38-1.34 (m, 1H).	Prepared from I-B30 and I-C7 by following the steps for preparing compound 12, and obtained by further resolution
125		541.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.51-7.42 (m, 3H), 7.31-7.18 (m, 2H), 5.48 (s, 1H), 4.40-4.20 (m, 4H), 3.93 (s, 1H), 3.41 (s, 3H), 3.26-3.09 (m, 3H), 2.81-2.77 (m, 1H), 2.07 (s, 3H), 1.86-1.65 (m, 2H), 1.60-1.56 (m, 1H), 1.39-1.35 (m, 1H).	
161		493.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.51-7.45 (m, 1H), 7.36-7.29 (m, 2H), 7.22-7.14 (m, 2H), 5.48 (s, 1H), 4.28 (br, 2H), 4.00 (s, 3H), 3.94 (s, 1H), 3.41 (s, 1H), 3.26-3.13 (m, 3H), 2.83-2.75 (m, 1H), 2.06 (s, 3H), 1.88-1.79 (m, 1H), 1.77-1.70 (m, 1H), 1.61-1.55 (m, 1H), 1.40-1.34 (m, 1H).	Obtained by resolution of compound 157
162		493.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.50-7.45 (m, 1H), 7.35-7.29 (m, 2H), 7.22-7.14 (m, 2H), 5.48 (s, 1H), 4.28 (br, 2H), 4.00 (s, 3H), 3.92 (s, 1H), 3.40 (s, 1H), 3.25-3.14 (m, 3H), 2.82-2.75 (m, 1H), 2.06 (s, 3H), 1.89-1.79 (m, 1H), 1.77-1.68 (m, 1H), 1.61-1.55 (m, 1H), 1.39-1.33 (m, 1H).	

163	 	537.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.48-7.41 (m, 1H), 7.35-7.26 (m, 2H), 7.22-7.14 (m, 2H), 5.48 (s, 1H), 4.30 (br, 4H), 4.03-3.97 (m, 3H), 3.92 (s, 1H), 3.41 (s, 3H), 3.27-3.13 (m, 3H), 2.83-2.74 (m, 1H), 2.06 (s, 3H), 1.88-1.79 (m, 1H), 1.76-1.67 (m, 1H), 1.61-1.55 (m, 1H), 1.39-1.33 (m, 1H).	Obtained by resolution of compound 158
164		537.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.41 (m, 1H), 7.35-7.26 (m, 2H), 7.22-7.14 (m, 2H), 5.48 (s, 1H), 4.30 (br, 4H), 4.03-3.97 (m, 3H), 3.93 (s, 1H), 3.41 (s, 3H), 3.26-3.13 (m, 3H), 2.82-2.76 (m, 1H), 2.06 (s, 3H), 1.87-1.79 (m, 1H), 1.76-1.68 (m, 1H), 1.61-1.55 (m, 1H), 1.40-1.33 (m, 1H).	
181	 	532.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.69 (s, 1H), 7.60-7.58 (m, 1H), 7.48-7.44 (m, 2H), 7.27-7.25 (m, 1H), 6.98-6.96 (m, 1H), 5.52 (s, 1H), 4.48-4.20 (m, 3H), 3.96 (s, 3H), 3.30-3.14 (m, 4H), 2.81 (s, 3H), 2.07 (s, 3H), 1.90-1.54 (m, 4H).	Prepared from I-B34 and I-C20 by following the steps for preparing compound 12, and obtained by further resolution
182		532.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.64 (s, 1H), 7.57-7.34 (m, 3H), 7.26-7.24 (m, 1H), 6.98-6.96 (m, 1H), 5.51 (s, 1H), 4.48-4.20 (m, 3H), 3.96 (s, 3H), 3.29-2.99 (m, 4H), 2.80 (s, 3H), 2.06 (s, 3H), 1.83-1.50 (m, 4H).	

190		500.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.86 (d, J = 8.3 Hz, 1H), 7.52 (s, 1H), 7.41 (d, J = 8.3 Hz, 1H), 7.35 (d, J = 7.6 Hz, 1H), 7.25 (d, J = 7.6 Hz, 1H), 5.53 (s, 1H), 4.34 (s, 2H), 4.16 (s, 3H), 3.98 (s, 1H), 3.45 (s, 1H), 3.32-3.17 (m, 3H), 2.87-2.78 (m, 1H), 2.12 (s, 3H), 1.93-1.72 (m, 2H), 1.63 (d, J = 12.1 Hz, 1H), 1.41 (d, J = 12.1 Hz, 1H).	Obtained by resolution of compound 183
191		500.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.86 (d, J = 8.4 Hz, 1H), 7.52 (s, 1H), 7.41 (d, J = 8.4 Hz, 1H), 7.36 (d, J = 7.7 Hz, 1H), 7.25 (d, J = 7.7 Hz, 1H), 5.53 (s, 1H), 4.34 (s, 2H), 4.16 (s, 3H), 3.98 (s, 1H), 3.45 (s, 1H), 3.33-3.14 (m, 3H), 2.87-2.78 (m, 1H), 2.12 (s, 3H), 1.93-1.71 (m, 2H), 1.63 (d, J = 13.6 Hz, 1H), 1.42 (d, J = 13.6 Hz, 1H).	
192		504.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.01 (d, J = 8.2 Hz, 1H), 7.67 (d, J = 8.2 Hz, 1H), 7.52 (s, 1H), 7.35 (d, J = 7.7 Hz, 1H), 7.24 (d, J = 7.7 Hz, 1H), 5.53 (s, 1H), 4.33 (s, 2H), 3.97 (s, 1H), 3.44 (s, 1H), 3.32-3.11 (m, 3H), 2.87-2.79 (m, 1H), 2.13 (s, 3H), 1.94-1.72 (m, 2H), 1.63 (d, J = 12.8 Hz, 1H), 1.41 (d, J = 12.8 Hz, 1H).	Obtained by resolution of compound 184
193		504.1	¹ H NMR (400 MHz, CD ₃ OD): δ 8.01 (d, J = 8.3 Hz, 1H), 7.67 (d, J = 8.3 Hz, 1H), 7.52 (s, 1H), 7.35 (d, J = 7.8 Hz, 1H), 7.25 (d, J = 7.8 Hz, 1H), 5.53 (s, 1H), 4.32 (s, 2H), 3.98 (s, 1H), 3.45 (s, 1H), 3.32-3.16 (m, 3H), 2.87-2.78 (m, 1H), 2.13 (s, 3H), 1.94-1.70 (m, 2H), 1.63 (d, J = 13.3 Hz, 1H), 1.41 (d, J = 13.3 Hz, 1H).	

200	 	550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.56 (s, 1H), 7.42 (d, J = 7.7 Hz, 1H), 7.36-7.26 (m, 2H), 7.20-7.14 (m, 1H), 5.49 (s, 1H), 4.45-4.10 (m, 2H), 4.01 (d, J = 1.2 Hz, 3H), 3.95 (s, 1H), 3.28-3.15 (m, 3H), 2.86-2.73 (m, 4H), 2.07 (s, 3H), 1.90-1.69 (m, 2H), 1.59 (d, J = 13.3 Hz, 1H), 1.35 (d, J = 12.3 Hz, 1H).	Prepared from I-B41 and I-C20 by following the steps for preparing compound 12, and obtained by further resolution
201		550.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.56 (s, 1H), 7.42 (d, J = 7.7 Hz, 1H), 7.36-7.26 (m, 2H), 7.20-7.14 (m, 1H), 5.49 (s, 1H), 4.45-4.10 (m, 2H), 4.01 (d, J = 1.2 Hz, 3H), 3.95 (s, 1H), 3.28-3.15 (m, 3H), 2.86-2.73 (m, 4H), 2.07 (s, 3H), 1.90-1.69 (m, 2H), 1.59 (d, J = 13.3 Hz, 1H), 1.35 (d, J = 12.3 Hz, 1H).	
207	 	507.2	¹ H NMR (400 MHz, CDCl ₃): δ 7.33 (s, 1H), 7.23-7.08 (m, 3H), 6.99-6.95 (m, 1H), 5.44 (s, 1H), 4.15 (br, 2H), 4.02 (s, 3H), 3.92 (s, 1H), 3.17-3.10 (m, 2H), 3.06-3.02 (m, 1H), 2.67-2.63 (m, 1H), 2.04-2.02 (m, 6H), 1.87-1.67 (m, 2H), 1.32-1.26 (m, 2H).	Prepared from I-B41 and I-C17 by following the steps for preparing compound 12, and obtained by further resolution
208		507.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.38-7.27 (m, 2H), 7.20-7.10 (m, 3H), 5.46 (s, 1H), 4.21 (br, 2H), 3.99 (s, 3H), 3.88 (s, 1H), 3.24-3.05 (m, 3H), 2.76-2.72 (m, 1H), 2.09-1.93 (m, 6H), 1.84-1.64 (m, 2H), 1.58-1.54 (m, 1H), 1.37-1.33 (m, 1H).	

209		538.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.56-7.53 (s, 1H), 7.49-7.39 (m, 2H), 7.33-7.25 (m, 2H), 5.47 (s, 1H), 4.27 (br, 2H), 3.95 (s, 1H), 3.27-3.15 (m, 3H), 2.85-2.74 (m, 4H), 2.07 (s, 3H), 1.89-1.79 (m, 1H), 1.77-1.67 (m, 1H), 1.62-1.55 (m, 1H), 1.38-1.31 (m, 1H).	Obtained by resolution of compound 202
210		538.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.57-7.53 (m, 1H), 7.50-7.39 (m, 2H), 7.33-7.26 (m, 2H), 5.47 (s, 1H), 4.25 (br, 2H), 3.96 (s, 1H), 3.27-3.15 (m, 3H), 2.86-2.76 (m, 4H), 2.07 (s, 3H), 1.87-1.79 (m, 1H), 1.77-1.68 (m, 1H), 1.61-1.55 (m, 1H), 1.39-1.33 (m, 1H).	
217		495.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.42 (m, 1H), 7.37-7.33 (m, 1H), 7.32-7.27 (m, 1H), 7.21-7.17 (m, 1H), 7.15-7.11 (m, 1H), 5.47 (s, 1H), 4.26 (br, 2H), 3.90 (s, 1H), 3.26-3.17 (m, 2H), 3.15-3.09 (m, 1H), 2.79-2.72 (m, 1H), 2.07 (s, 3H), 1.98 (s, 3H), 1.85-1.77 (m, 1H), 1.75-1.67 (m, 1H), 1.59-1.53 (m, 1H), 1.40-1.34 (m, 1H).	Obtained by resolution of compound 214
218		495.1	¹ H NMR (400 MHz, CD ₃ OD): δ 7.49-7.42 (m, 1H), 7.37-7.33 (m, 1H), 7.32-7.27 (m, 1H), 7.22-7.17 (m, 1H), 7.17-7.11 (m, 1H), 5.47 (s, 1H), 4.26 (br, 2H), 3.91 (s, 1H), 3.27-3.18 (m, 2H), 3.15-3.09 (m, 1H), 2.81-2.72 (m, 1H), 2.07 (s, 3H), 1.98 (s, 3H), 1.85-1.77 (m, 1H), 1.75-1.67 (m, 1H), 1.60-1.53 (m, 1H), 1.41-1.33 (m, 1H).	




220		511.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.59-7.33 (m, 3H), 7.20-7.16 (m, 2H), 5.47 (s, 1H), 4.25 (br, 2H), 3.95 (s, 1H), 3.27-3.05 (m, 3H), 2.82-2.78 (m, 1H), 2.18-1.92 (m, 6H), 1.82-1.68 (m, 2H), 1.60-1.52 (m, 1H), 1.44-1.36 (m, 1H).	Prepared from I-B30 and I-C17 by following the steps for preparing compound 12, and obtained by further resolution
221		511.2	¹ H NMR (400 MHz, CD ₃ OD): δ 7.54-7.28 (m, 3H), 7.18-7.12 (m, 2H), 5.47 (s, 1H), 4.24 (br, 2H), 3.89 (s, 1H), 3.26-3.06 (m, 3H), 2.76-2.72 (m, 1H), 2.20-1.90 (m, 6H), 1.82-1.64 (m, 2H), 1.58-1.52 (m, 1H), 1.38-1.32 (m, 1H).	
267		470.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.00-7.94 (m, 1H), 7.77-7.73 (m, 1H), 7.69-7.64 (m, 1H), 7.47 (s, 1H), 7.34-7.27 (m, 1H), 7.22-7.18 (m, 1H), 5.49 (s, 1H), 4.50-3.98 (m, 2H), 3.92 (s, 1H), 3.39 (s, 1H), 3.28-3.20 (m, 2H), 3.19-3.11 (m, 1H), 2.84-2.72 (m, 1H), 2.06 (s, 3H), 1.89-1.78 (m, 1H), 1.78-1.67 (m, 1H), 1.61-1.56 (m, 1H), 1.39-1.34 (m, 1H).	Obtained by resolution of compound 34
268		470.2	¹ H NMR (400 MHz, CD ₃ OD): δ 8.01-7.94 (m, 1H), 7.78-7.72 (m, 1H), 7.70-7.62 (m, 1H), 7.47 (s, 1H), 7.33-7.27 (m, 1H), 7.23-7.16 (m, 1H), 5.49 (s, 1H), 4.47-4.02 (m, 2H), 3.92 (s, 1H), 3.39 (s, 1H), 3.28-3.19 (m, 2H), 3.19-3.12 (m, 1H), 2.83-2.71 (m, 1H), 2.06 (s, 3H), 1.89-1.78 (m, 1H), 1.77-1.67 (m, 1H), 1.61-1.55 (m, 1H), 1.40-1.33 (m, 1H).	








The diastereomers in the table are subjected to chiral HPLC under the following conditions (flow rate: 15 mL/minute; detector: UV 254 nm):








Compounds	Column	Mobile phase	RT/minute	de%
58	IC (2 × 25 cm)	Acetonitrile/ethanol = 10:90	7.180	100%
59			9.684	100%
65	IC (2 × 25 cm)	Acetonitrile/ethanol = 10:90	6.427	100%
66			7.045	100%
88	IC (2 × 25 cm)	Acetonitrile/ethanol = 10:90	6.136	93.39%
89			6.832	93.92%
102	ODH (2 × 25 cm)	n-Heptane/ethanol = 50:50	10.838	100%
103			12.127	99.30%
108	IC (2 × 25 cm)	Acetonitrile/ethanol = 10:90	6.311	96.97%
109			7.702	97.01%
110	ODH (2 × 25 cm)	Acetonitrile/ethanol = 10:90	14.376	100%
111			23.609	100%
113	IC (2 × 25 cm)	Acetonitrile/ethanol = 10:90	5.515	100%
114			6.581	100%
118	IC (2 × 25 cm)	Acetonitrile /ethanol = 40:60	4.193	97.78%
119			4.660	97.17%
124	IC (2 × 25 cm)	Acetonitrile/ethanol = 10:90	8.555	100%
125			10.658	96.90%
161	IC (2×25 cm)	Acetonitrile/ethanol = 10:90	6.693	99.01%
162			8.071	95.18%
163	IC (2×25 cm)	Acetonitrile/ethanol = 10:90	9.344	100%
164			11.191	93.96%
181	ODH (2×25 cm)	Acetonitrile/ethanol = 50:50	5.810	100%
182			7.066	100%
190	IA (2×25 cm)	Acetonitrile/ethanol = 10:90	12.754	100%
191			19.343	100%
192	ADH (2×25 cm)	Acetonitrile/ethanol = 10:90	5.340	100%
193			14.222	100%
200	IA (2×25 cm)	Acetonitrile/ethanol = 10:90	5.599	100%
201			10.573	100%
207	IC (2×25 cm)	Ethanol	18.586	99.815
208			25.797	99.57%



209	IC (2×25 cm)	Ethanol	10.679	100%
210			14.798	100%
217	IC (2×25 cm)	Ethanol	10.544	100%
218			15.425	100%
220	IA (2×25 cm)	Acetonitrile/ethanol = 10:90	6.682	100%
221			13.507	100%
267	ODH (2×25 cm)	Acetonitrile/ethanol = 10:90	7.894	100%
268			14.016	100%

The compounds in the table below were prepared by following the steps for preparing the above compounds from corresponding intermediates and reagents:

Compounds	Structural formula	Molecular weight
136		509.43
137		504.42
138		509.43

139		529.48
140		544.50
141		460.96
142		484.99
143		477.97
144		552.11
145		567.13

146		563.05
149		509.65
150		524.66
151		543.10
152		558.12
153		494.64
154		509.65

304		576.12
305		505.04

Example 2 Assay of Full-length SHP2 Phosphatase Activity

1. Reagents and materials

- Human full-length SHP2 recombinant protein: BPS Bioscience, Cat# 79018;
- SHP2 substrate DiFMUP (1 mM): BPS Bioscience, Cat# 79769;
- SHP2 activating peptide (100 μ M): BPS Bioscience, Cat# 79319-2;
- DTT: Merck, Cat# DTT-RO;
- 384-well plate: Corning, Cat# 3575;
- 96-well plate: Thermo Fisher Scientific, Cat# 249952;
- Instrument: EnVision 2104, PerkinElmer.

2. Preparation of reaction solutions

The test compound was dissolved in DMSO and diluted with DMSO to 100.0 μ M, and the compound was further 3-fold diluted with DMSO to: 100.00, 33.33, 11.11, 3.70, 1.23, 0.41, 0.14 and 0.05 μ M. Then 4 μ L of the compound at different dilution concentrations was added to 96 μ L of an enzymatic reaction buffer to prepare a 4X test compound, wherein DMSO was at the concentration of 4% (DMSO was at the final concentration of 1%).

Preparation of 1X enzymatic reaction buffer: the 5X reaction buffer (250 mM HEPES, 500 mM NaCl, 2.5 mM EDTA, 0.005% Brij-35 and 0.01% BSA, pH 7.2) was diluted 5-fold with deionized water, and then DTT was added thereto so that the 1X enzymatic reaction buffer contained 5 mM DTT.

Preparation of 4X mixed solution of SHP2 enzyme/activating peptide: the SHP2 enzyme (75.5 nM) and activating peptide (100 μ M) were diluted with the enzymatic reaction buffer to prepare a 4X mixed solution of SHP2 enzyme/activating

peptide (0.12 nM SHP2 and 2 μ M activating peptide), so that the SHP2 enzyme and activating peptide were at the final concentrations of 0.03 nM and 0.5 μ M in the enzymatic reaction system, respectively.

- Preparation of 2X DiFMUP substrate: 1 mM DiFMUP was diluted 100-fold with the enzymatic reaction buffer to prepare a 2X substrate (10 μ M), so that the substrate DiFMUP was at the final concentration of 5 μ M in the enzymatic reaction system.

3. Experimental steps

- To the corresponding wells of the 384-well plate was added 2.5 μ L of 4X test compound or 2.5 μ L of 4% DMSO solution, and centrifugation was carried out at 1000 rpm for 30 seconds.

- To the test compound well and positive control well was added 2.5 μ L of 4X mixed solution of SHP2 enzyme/activating peptide while to the negative control well was added 2.5 μ L of 1X enzymatic reaction buffer; centrifugation was carried out at 1000 rpm for 30 seconds and incubation was carried out at room temperature for 30 minutes.

- To each well was added 5 μ L of 2X DiFMUP substrate to initiate the enzymatic reaction, then the plate was transiently oscillated and centrifuged at 1000 rpm for 30 seconds, and the 384-well plate was sealed with a sealing membrane and incubated at 25°C on a low-speed shaker (100 rpm) in the dark for 60 minutes.

- The fluorescence value (relative fluorescence units, RFU) of each well was detected on EnVision2104 (excitation: 355 nm, emission: 460 nm).

4. Data analysis

$$\text{Inhibition rate (\%)} = 100 - \frac{\text{RFU of compound well} - \text{RFU of negative control well}}{\text{RFU of positive control well} - \text{RFU of negative control well}} \times 100$$

wherein

- the RFU of compound well refers to the fluorescence reading at 460 nm of the well containing the test compounds;

- the RFU of negative control well refers to the fluorescence reading at 460 nm of the background well containing 1% DMSO and the enzymatic reaction buffer;

• the RFU of positive control well refers to the fluorescence reading at 460 nm of the well containing 1% DMSO and the mixed solution of SHP2 enzyme/activating peptide.

IC₅₀ values were calculated following formula 205: $y = A + ((B-A)/(1 +$

5 $((C/X)^D)))$, by using the XL-Fit 5.0 software.

5. Test results

Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)
1	0.002	23	0.757	45	0.011	67	0.001
2	0.005	24	0.412	46	0.096	68	0.130
3	0.009	25	0.012	47	0.011	69	0.005
4	0.004	26	0.158	48	0.063	70	0.002
5	0.002	27	0.001	49	0.003	71	0.001
6	0.420	28	> 1	50	0.015	72	0.978
7	0.010	29	0.025	51	0.001	73	0.253
8	0.003	30	0.002	52	< 0.0005	74	0.004
9	0.004	31	0.005	53	0.018	75	0.002
10	0.035	32	0.001	54	0.003	76	0.002
11	0.097	33	0.310	55	0.026	77	0.005
12	0.001	34	0.004	56	0.002	78	0.003
13	> 1	35	0.017	57	0.028	79	0.001
14	> 1	36	0.006	58	0.013	80	0.004
15	0.008	37	0.044	59	0.248	81	0.002
16	0.004	38	0.046	60	0.013	82	0.002
17	0.018	39	0.021	61	0.003	83	0.001
18	> 1	40	0.002	62	0.005	84	0.067
19	> 1	41	0.021	63	0.004	85	0.050
20	0.012	42	0.006	64	0.184	86	0.002
21	0.541	43	0.006	65	0.002	87	0.001
22	0.348	44	0.060	66	0.024	88	0.0008
89	0.015	101	0.012	114	0.053	126	0.002
90	0.0011	102	0.002	115	0.002	127	0.0006
91	0.0007	103	0.017	116	0.0005	128	0.0009
92	0.011	104	0.004	117	0.002	129	0.0006

93	0.016	106	0.009	118	0.003	130	> 1.0
94	0.007	107	0.739	119	0.178	131	0.003
95	0.003	108	0.001	120	0.045	132	0.003
96	0.005	109	0.118	121	> 1.0	133	0.029
97	0.002	110	0.249	122	0.0006	134	0.0014
98	0.003	111	0.002	123	0.0006	135	0.005
99	0.002	112	0.002	124	0.0013	147	0.003
100	0.002	113	0.002	125	0.096	148	0.004
155	0.001	193	0.001	231	<0.0005	269	0.0009
156	0.006	194	0.001	232	0.002	270	0.0004
157	0.002	195	0.001	233	0.0011	271	0.001
158	0.003	196	0.003	234	0.0028	272	0.017
159	0.001	197	0.003	235	0.0017	273	0.001
160	0.013	198	0.016	236	0.0013	274	0.001
161	0.001	199	0.001	237	0.0039	275	0.0005
162	0.034	200	0.025	238	0.0013	276	0.002
163	0.001	201	0.001	239	0.003	277	0.001
164	0.042	202	0.001	240	0.0015	278	0.005
165	0.006	203	0.002	241	0.0012	279	0.0004
166	0.006	204	0.006	242	0.0007	280	0.011
167	0.025	205	0.012	243	0.0007	281	0.010
168	0.010	206	0.003	244	0.011	282	0.001
169	0.006	207	0.002	245	<0.0005	283	0.001
170	0.009	208	0.058	246	0.133	284	0.002
171	0.006	209	0.001	247	0.001	285	0.085
172	0.001	210	0.052	248	<0.0005	286	0.002
173	0.001	211	0.017	249	0.001	287	0.012
174	0.006	212	0.023	250	0.001	288	0.002
175	0.014	213	0.068	251	0.001	289	0.002
176	0.038	214	0.004	252	0.001	290	0.001
177	0.002	215	0.025	253	0.001	291	0.001
178	0.002	216	0.001	254	0.003	292	0.0009
179	0.004	217	0.001	255	0.003	293	0.001
180	0.013	218	0.104	256	0.004	294	0.001
181	0.044	219	0.003	257	<0.0005	295	0.002

182	0.001	220	0.317	258	0.010	296	0.0003
183	0.002	221	0.001	259	0.004	297	0.0003
184	0.003	222	0.004	260	0.001	298	0.0003
185	0.014	223	0.004	261	0.0009	299	0.001
186	0.042	224	0.009	262	0.0003	300	0.001
187	0.001	225	0.002	263	0.009	301	0.0004
188	0.005	226	0.001	264	0.0006	302	0.0004
189	0.001	227	0.001	265	0.0004	303	0.0005
190	0.253	228	0.203	266	0.005		
191	0.001	229	0.001	267	>0.1		
192	>1.0	230	0.010	268	0.001		

Example 3 Assay of intracellular pERK1/2 (Thr202/Tyr204) phosphorylation

1. Reagents and materials

- pERK1/2 (Thr202/Tyr204) HTRF kit: Cisbio, Cat# 64ERKPEH;
- Cell line: Miapaca2, ATCC, CRL-1420;
- OptiPlate™-384-well plate: PerkinElmer, Cat# 6007299;
- 96-well plate: Corning, Cat# 353072;
- Instrument: EnVision2104, PerkinElmer.

2. Preparation of reaction solutions

The test compound was dissolved in DMSO and diluted with DMSO to 600.0 μ M, and the compound was further 3-fold diluted with DMSO to: 600.0, 200.0, 66.7, 22.2, 7.4, 2.5, 0.82 and 0.27 μ M. Then 10 μ L of the compound at different dilution concentrations was added to 190 μ L of a DMEM medium to prepare a 10X test compound, wherein DMSO was at the concentration of 5% (DMSO was at the final concentration of 0.5%).

1X cell lysis buffer: 4X cell lysis stock solution (provided by the kit) was diluted 4-fold with deionized water, and then 1% 100X blocking stock solution (provided by the kit) was added thereto.

pERK1/2 detection solution (prepared just before use): the pERK1/2 d2 antibody (provided by the kit) and pERK1/2 Cryptate antibody (provided by the kit) were diluted with the detection solution (provided by the kit) at a ratio of 1 : 1 : 38.

3. Experimental steps

- Miapaca2 cells were inoculated into a 96-well plate at a density of 10000 cells/well at 90 μ L/well and cultured in a cell incubator at 5% CO₂ and 37°C overnight.

- 10 μ L of 10X test compound was added to the 90 μ L cell culture 96-well plate; to the cell positive control well was added 10 μ L of 5% DMSO culture solution; and the plate was cultured in a cell incubator at 5% CO₂ and 37°C for 2 hours.

- The medium in the 96-well plate was removed; 50 μ L of 1X cell lysis buffer was added to each well; the plate was placed in a microplate shaker; and lysis under shaking was carried out at 900 rpm at room temperature for 1 hour.

- 16 μ L of the lysis buffer in the 96-well plate was taken and transferred to a 384-well plate, and centrifugation was carried out at 1000 rpm for 30 seconds; then 4 μ L of pERK1/2 detection solution was added to each well and centrifugation was carried out at 1000 rpm for 30 seconds; and the 384-well plate was sealed with a sealing membrane and incubated at 25°C on a low-speed shaker (100 rpm) in the dark for 2 hours.

- The fluorescence value (relative fluorescence units, RFU) of each well was detected on EnVision2104 (emission 1: 665 nm, emission 2: 615 nm).

4. Data analysis

$$\text{Fluorescence ratio} = \frac{\text{RFU}_{665 \text{ nm}}}{\text{RFU}_{615 \text{ nm}}}$$

$$\text{Inhibition rate (\%)} = 100 - \frac{\text{Fluorescence ratio of compound well} - \text{fluorescence ratio of negative control well}}{\text{Fluorescence ratio of positive control well} - \text{fluorescence ratio of negative control well}} \times 100$$

wherein

- the fluorescence ratio of compound well refers to the fluorescence ratio of the well containing the test compounds;
- the fluorescence ratio of negative control well refers to the fluorescence ratio of the background containing the cell lysis buffer and no Miapaca2 cells;
- the fluorescence ratio of positive control well refers to the fluorescence ratio of the well containing 0.5% DMSO and Miapaca2 cells.

IC₅₀ values were calculated following formula 205: $y = A + ((B-A)/(1 + ((C/X)^D)))$, by using the XL-Fit 5.0 software.

5. Test results

Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)
1	0.018	32	0.006	55	0.388	78	0.043
2	0.023	34	0.244	56	0.085	79	0.060
3	0.081	35	0.274	57	0.526	80	0.125
4	0.160	36	0.033	58	0.260	81	0.089
5	0.023	37	0.599	60	> 1.0	82	0.043
6	> 3	38	0.343	61	0.078	83	0.057
7	0.028	39	0.294	62	0.105	84	> 1
8	0.008	40	0.006	63	0.177	85	0.785
9	0.038	41	0.115	64	0.891	86	0.006
10	0.220	42	0.133	65	0.060	87	0.002
11	1.148	43	0.117	66	0.271	88	0.026
12	0.007	44	0.415	67	0.039	89	0.153
15	0.087	45	0.261	68	0.510	90	0.053
16	0.034	46	0.648	69	0.066	91	0.017
17	0.203	47	0.115	70	0.141	92	0.117
20	0.052	48	0.231	71	0.022	93	0.301
25	0.151	49	0.018	72	> 1.0	94	0.155
26	> 3	50	0.318	73	> 1.0	95	0.002
27	0.011	51	0.021	74	0.223	96	0.055
29	0.489	52	0.037	75	0.046	97	0.011
30	0.008	53	0.364	76	0.004	98	0.020
31	0.037	54	0.036	77	0.056	99	0.040
100	0.012	108	0.007	115	0.026	122	0.007
101	0.096	109	0.259	116	0.001	123	0.004
102	0.012	110	0.959	117	0.024	124	0.015
103	0.080	111	0.015	118	0.040	125	0.175
104	0.040	112	0.006	119	0.698	126	0.042
106	0.084	113	0.014	120	0.385	127	0.013
107	> 1.0	114	0.284	121	> 1.0	128	0.015
129	0.003	131	0.004	132	0.009	133	0.147
134	0.005	135	0.056	147	0.010	148	0.013
155	0.002	193	0.024	231	0.007	269	0.007

156	0.010	194	0.018	232	0.008	270	0.002
157	0.025	195	0.011	233	0.014	271	0.004
158	0.031	196	0.008	234	0.153	272	0.252
159	0.005	197	0.029	235	0.009	273	0.005
160	0.056	198	0.191	236	0.003	274	0.006
161	0.013	199	0.005	237	0.004	275	0.002
162	0.155	200	0.512	238	0.004	276	0.004
163	0.014	201	0.006	239	0.015	277	0.011
164	0.154	202	0.040	240	0.018	278	0.034
165	0.039	203	0.007	241	0.010	279	0.183
166	0.038	204	0.054	242	0.013	280	0.035
167	0.124	205	0.029	243	0.019	281	0.167
168	0.034	206	0.069	244	0.139	282	0.003
169	0.125	207	0.009	245	0.012	283	0.737
170	0.143	208	0.187	246	0.688	284	0.005
171	0.038	209	0.019	247	0.003	285	0.962
172	0.008	210	0.527	248	0.002	286	0.010
173	0.018	211	0.161	249	0.031	287	0.140
174	0.033	212	0.223	250	0.072	288	0.004
175	0.122	213	0.447	251	0.024	289	0.004
176	0.041	214	0.100	252	0.003	290	0.004
177	0.007	215	0.167	253	0.010	291	0.009
178	0.048	216	0.045	254	0.011	292	0.006
179	0.112	217	0.049	255	0.008	293	0.005
180	0.251	218	0.944	256	0.012	294	0.018
181	0.447	219	0.007	257	0.001	295	0.022
182	0.023	220	>1	258	0.188	296	0.002
183	0.050	221	0.015	259	0.023	297	0.001
184	0.067	222	0.006	260	0.005	298	0.001
185	>1.0	223	0.023	261	0.007	299	0.002
186	>1.0	224	0.020	262	0.007	300	0.013
187	0.003	225	0.006	263	0.204	301	0.005
188	0.014	226	0.002	264	0.002	302	0.005
189	0.007	227	0.002	265	0.0009	303	0.009
190	0.672	228	>1	266	0.026		

191	0.021	229	0.002	267	>1		
192	>1.0	230	0.031	268	0.063		

Example 4 Assay of cell proliferation on Miapaca2 3D spheroids

1. Reagents and materials

- CellTiter-Glo ® 3D Cell Viability Assay Kit: Promega, Cat# G9683;
- CellCarrier Spheroid ULA 96-well plate: Corning, Cat# 4515;
- Instrument: Envision, Perkinelmer;
- Cell line: Miapaca2, ATCC, Cat# CRL-1420.

2. Preparation of reaction solutions

- The test compound was dissolved in DMSO and diluted with DMSO to 3000.0 μ M, and the compound was further 3-fold diluted with DMSO to: 1000.0, 333.3, 111.1, 37.0, 12.3, 4.1, and 1.4 μ M. Then 2 μ L of the compound at different dilution concentrations was added to 198 μ L of a 1640 medium to prepare a 10X test compound, wherein DMSO was at the concentration of 1% (DMSO was at the final concentration of 0.1%).

3. Experimental methods

- Day0: Digest cells and count cell numbers. Miapaca2 cells were inoculated into a spheroid ULA 96-well plate at a density of 300 cells/well at 100 μ L/well and cultured in a cell incubator at 5% CO₂ and 37°C.
- Day2: The cell spheroids formed at day2. Add 10 μ L of 10X test compounds to the 96-well plate. To the cell positive control well was added 10 μ L of 1% DMSO 1640 medium. Incubate the cell spheroids in a cell incubator at 5% CO₂ and 37°C for additional 5 days.
- Cell viability assay: Add 50 μ L of CellTiter-Glo reagent to each well, place the plate in a microplate shaker, and lysis under shaking at 900 rpm at room temperature for 5 minutes. Then incubate the plate at room temperature in the dark for 30 minutes.
- The luminescence of each well was detected on Envision2104.

4. Data analysis

- $Cell\ Survival\% = (Lum(d7\ treatment) - Lum(d2\ cell)) / (Lum(d7\ cell) - Lum(d2\ cell)) \times 100$

Wherein:

- *Lum(d7 treatment)* refers to the luminescence of the test compound treated cells on day7;
- *Lum(d2 cell)* refers to the luminescence of the 0.1% DMSO treated Miapaca2 cells on day2;
- *Lum(d7 cell)* refers to the luminescence of the 0.1% DMSO treated Miapaca2 cells on day7.

IC₅₀ values were calculated following formula 205: $y=A+((B-A)/(1+((C/X)^D)))$, by using the XL-Fit 5.0 software.


5. Test results

Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)	Compound no.	IC ₅₀ (μM)
135	0.095	191	0.058	237	0.0024	271	0.002
147	0.048	193	0.090	238	0.001	273	0.002
148	0.047	194	0.036	239	0.0021	274	0.003
155	0.0009	195	0.030	240	0.005	275	0.002
156	0.031	196	0.021	241	0.0038	276	0.003
157	0.028	197	0.038	242	0.002	277	0.005
158	0.032	199	0.012	243	0.015	278	0.016
159	0.006	201	0.018	244	0.041	279	0.042
160	0.041	202	0.046	245	0.017	280	0.035
161	0.008	203	0.016	247	0.001	281	0.024
163	0.020	204	0.121	248	0.0004	282	0.002
165	0.071	205	0.068	249	0.019	283	0.243
166	0.110	206	0.078	250	0.014	284	0.004
168	0.077	207	0.035	251	0.029	286	0.009
170	0.162	209	0.038	252	0.001	288	0.002
171	0.039	216	0.028	253	0.002	289	0.002
172	0.009	217	0.062	254	0.010	290	0.0003
173	0.024	219	0.006	255	0.003	291	0.0003
174	0.053	221	0.063	256	0.006	292	0.001
175	0.100	222	0.015	257	0.0003	293	0.0009
176	0.097	225	0.002	259	0.005	294	0.009
177	0.004	226	0.0004	260	0.002	295	0.004

178	0.041	227	0.0004	261	0.002	296	0.0007
179	0.065	229	0.0005	262	0.001	297	0.0003
182	0.018	230	0.0074	264	0.0004	298	0.0003
183	0.068	231	0.0008	265	0.0004	299	0.001
184	0.096	232	0.0009	266	0.010	300	0.016
187	0.003	233	0.0032	268	0.017	301	0.001
188	0.047	235	0.0009	269	0.003	302	0.001
189	0.015	236	0.001	270	0.0006		

CLAIMS

1. A compound of formula (I):



or a pharmaceutically acceptable salt thereof, or a solvate, a racemic mixture, an enantiomer, a diastereomer or a tautomer thereof, wherein

ring A is a benzene ring or a pyridine ring;

Z is CH₂, O, S or NH;

- 10 R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄, -SR₅ and -SR₆, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -
- 15 NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-CN, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and -(C₁₋₆ alkyl)-CN; R₄ and R₅ are each
- 20 independently chosen from C₃₋₈ cycloalkyl, phenyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl; R₆ is chosen from -CO(C₁₋₆ alkyl), -CO(C₃₋₈ cycloalkyl), -CO(4-8 membered heterocyclyl), -CONH₂, -CONH(C₁₋₆ alkyl), -CONH(C₃₋₈ cycloalkyl), -CONH(4-8 membered heterocyclyl), -CON(C₁₋₆ alkyl)₂, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-NH(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-N(C₁₋₆ alkyl)₂ and -(C₁₋₆ alkyl)-
- 25 NHCO(C₁₋₆ alkyl), wherein the C₁₋₆ alkyl of R₆ is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH and -O(C₁₋₆ alkyl); and the above-mentioned C₃₋₈ cycloalkyl, phenyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: halogen, -CN, -CONH₂, -OH, oxo, -NH₂, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆

haloalkyl, $-(C_{1-6} \text{ alkyl})-OH$, $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$, $-(C_{1-6} \text{ alkyl})-CN$, $-O(C_{1-6} \text{ alkyl})$, $-O(C_{1-6} \text{ haloalkyl})$, $-S(C_{1-6} \text{ alkyl})$, $-NH(C_{1-6} \text{ alkyl})$ and $-N(C_{1-6} \text{ alkyl})_2$;

R_1' is independently chosen from halogen, $-CN$, $-CONH_2$, $-OH$, $-NH_2$, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-8} cycloalkyl, 4-8 membered heterocyclyl, $-(C_{1-6} \text{ alkyl})-OH$, $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$, $-(C_{1-6} \text{ alkyl})-CN$, $-O(C_{1-6} \text{ alkyl})$, $-O(C_{1-6} \text{ haloalkyl})$, $-O(C_{3-8} \text{ cycloalkyl})$, $-O(4-8 \text{ membered heterocyclyl})$, $-S(C_{1-6} \text{ alkyl})$, $-NH(C_{1-6} \text{ alkyl})$ and $-N(C_{1-6} \text{ alkyl})_2$, wherein the C_{3-8} cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more halogen;

n is 0, 1, 2 or 3;


R_2 is chosen from $-NH_2$, $-NH(C_{1-6} \text{ alkyl})$, $-N(C_{1-6} \text{ alkyl})_2$, oxo and $-OH$;

Cy_1 is 5-12 membered heterocyclyl or 5-12 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-CONH_2$, $-OH$, oxo, $-NH_2$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{3-8} cycloalkyl, 4-8 membered heterocyclyl, $-(C_{1-6} \text{ alkyl})-OH$, $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$, $-(C_{1-6} \text{ alkyl})-CN$, $-O(C_{1-6} \text{ alkyl})$, $-O(C_{1-6} \text{ haloalkyl})$, $-O(C_{3-8} \text{ cycloalkyl})$, $-O(4-8 \text{ membered heterocyclyl})$, $-S(C_{1-6} \text{ alkyl})$, $-NH(C_{1-6} \text{ alkyl})$, $-N(C_{1-6} \text{ alkyl})_2$, $-NHCO(C_{1-6} \text{ alkyl})$, $-CONH(C_{1-6} \text{ alkyl})$ and $-CON(C_{1-6} \text{ alkyl})_2$, wherein the C_{3-8} cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more halogen;

Cy_2 is phenyl or 5-14 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-CONH_2$, $-OH$, oxo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, $-(C_{1-6} \text{ alkyl})-OH$, $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$, $-(C_{1-6} \text{ alkyl})-CN$, $-O(C_{1-6} \text{ alkyl})$, $-O(C_{1-6} \text{ haloalkyl})$, $-O(C_{3-8} \text{ cycloalkyl})$, $-O(4-8 \text{ membered heterocyclyl})$, $-S(C_{1-6} \text{ alkyl})$, $-NR_7R_8$, $-NHCO(C_{1-6} \text{ alkyl})$, $-CONH(C_{1-6} \text{ alkyl})$ and $-CON(C_{1-6} \text{ alkyl})_2$, wherein R_7 and R_8 are each independently chosen from hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, $-(C_{1-6} \text{ alkyl})-OH$, $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$, $-(C_{1-6} \text{ alkyl})-CN$, C_{3-8} cycloalkyl, phenyl, 3-8 membered heterocyclyl and 5-6 membered heteroaryl; and L is absent, or L is S, O, NH, C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl.

2. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1, wherein the compound is a compound of formula (IA):

199



3. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1 or 2, wherein Z is CH₂ or O; and preferably, Z is CH₂.

4. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of claims 1-3, wherein R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄ and -SR₅,

wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently chosen from hydrogen and C₁₋₆ alkyl; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-12 membered heteroaryl; and the above-mentioned C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-12

membered heteroaryl are each optionally substituted with one or more groups independently chosen from: halogen, -CN, -CONH₂, -OH, oxo, -NH₂, C₁₋₆ alkyl and -O(C₁₋₆ alkyl).

5. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 4, wherein R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆

alkyl), C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl, wherein the C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently
5 chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -CONH(C₁₋₆ alkyl), -CONH(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -CON(C₁₋₆ alkyl)₂, -CON(C₁₋₆ alkyl)(C₁₋₆ alkyl-O-C₁₋₆ alkyl), -CONH(C₃₋₈ cycloalkyl), -COOH, -COO(C₁₋₆ alkyl), -CO(C₁₋₆ alkyl), -CO(4-8 membered heterocyclyl) and -CO(4-8 membered heterocyclyl)-O-(C₁₋₆ alkyl); more preferably, R₁ is ethynyl, propynyl or butynyl, each of which is
10 unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂, -NHCONH₂, -CONH(C₁₋₃ alkyl), -CONH(CH₂CH₂)-O-(CH₃), -CON(CH₃)₂, -CON(CH₃)(CH₂CH₂-O-CH₃), -CONH(cyclopropyl), -COOH, -COO(CH₃), -CO(CH₃), -CO(azetidiny) or -CO(azetidiny)-O-(CH₃); and most preferably, R₁ is ethynyl, or ethynyl substituted with -CONH(C₁₋₃ alkyl).

15
6. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 4, wherein R₁ is chosen from -NR₃R₄ and -SR₅, wherein the R₃ is independently chosen from hydrogen and C₁₋₆ alkyl; and R₄ and R₅ are each independently chosen from
20 C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl, wherein the C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl.

7. The compound or the pharmaceutically acceptable salt thereof, or the solvate,
25 the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of claims 1-6, wherein R₁' is chosen from halogen, -CN, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0.


30 8. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according

to any one of claims 1-7, wherein R_2 is chosen from $-NH_2$ and oxo; and preferably, R_2 is $-NH_2$.

9. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of claims 1-8, wherein Cy_1 is 5-12 membered heterocyclyl or 5-12 membered heteroaryl, preferably 5-10 membered heterocyclyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heterocyclyl or 5-9 membered heteroaryl, which is optionally substituted with one or more groups independently chosen from: oxo, $-NH_2$, $-CN$, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl and $-(C_{1-6} \text{ alkyl})-OH$.

10. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 9, wherein Cy_1 is chosen from 1,6-dihydropyrimidyl, pyrazinyl, pyrimidyl, 1,2,4-triazinyl, imidazopyrimidyl, triazolopyrimidyl, imidazopyrazinyl, pyrrolopyrazinyl, pyrazolopyrazinyl and triazolopyrazinyl, each of which is optionally substituted with one or more groups independently chosen from: oxo, $-NH_2$, $-CN$, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl and $-(C_{1-6} \text{ alkyl})-OH$.


11. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 10, wherein Cy_1 is chosen from



, each of


which is optionally substituted with one or more groups independently chosen from: -

NH_2 , $-CN$, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl and $-(C_{1-6} \text{ alkyl})-OH$; preferably, Cy_1 is





each of which is optionally substituted with one or more groups independently chosen from: $-NH_2$, $-CN$, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl and $-(C_{1-6} \text{ alkyl})-OH$;


more preferably, Cy_1 is chosen from




5 each of which is optionally substituted with one or more groups independently chosen from: $-NH_2$, C_{1-6} alkyl and $-(C_{1-6} \text{ alkyl})-OH$;

further preferably, Cy_1 is , which is optionally substituted with one or

more groups independently chosen from: $-NH_2$ and C_{1-6} alkyl; or Cy_1 is , which is optionally substituted with one or more groups independently chosen from: C_{1-6}


10 alkyl; or Cy_1 is , which is optionally substituted with one or more groups

independently chosen from: $-NH_2$, C_{1-6} alkyl and $-(C_{1-6} \text{ alkyl})-OH$; or Cy_1 is , which is optionally substituted with one or more groups independently chosen from: $-NH_2$ and C_{1-6} alkyl.

15 **12.** The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of claims 1-11, wherein Cy_2 is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably phenyl, 5-6 membered heteroaryl or 9-10 membered heteroaryl, each of which is optionally

20 substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6} \text{ alkyl})$, $-S(C_{1-6} \text{ alkyl})$ and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, $-(C_{1-6} \text{ alkyl})-OH$, $-(C_{1-6} \text{ alkyl})-O-(C_{1-6} \text{ alkyl})$, $-(C_{1-6} \text{ alkyl})-CN$ and C_{3-8} cycloalkyl.

13. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1, wherein the compound is a compound of formula (II):



5 wherein

Z is CH₂ or O; and preferably, Z is CH₂;

R₁ is chosen from C₂₋₆ alkynyl, -NR₃R₄ and -SR₅, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; R₃ is independently
 15 chosen from hydrogen and C₁₋₆ alkyl; R₄ and R₅ are each independently chosen from C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl; and the above-mentioned C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is
 20 optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -CONH(C₁₋₆ alkyl) and -CON(C₁₋₆ alkyl)₂; further preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂, -CONH(CH₃) or -CON(CH₃)₂; and more preferably, R₁ is ethynyl;

25 R₁' is chosen from halogen, -CN, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0;


R₂ is -NH₂;

R_9 and R_{10} are each independently chosen from hydrogen, $-NH_2$, halogen, C_{1-6} alkyl and C_{1-6} haloalkyl; and preferably, R_9 and R_{10} are each independently chosen from hydrogen and C_{1-6} alkyl;

Cy_2 is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably phenyl, each of which is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6}$ alkyl), $-S(C_{1-6}$ alkyl) and $-NR_7R_8$, wherein R_7 and R_8 are each independently chosen from hydrogen, $-(C_{1-6}$ alkyl)- OH , $-(C_{1-6}$ alkyl)- $O-(C_{1-6}$ alkyl) and C_{3-8} cycloalkyl; and

L is absent.

14. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1, wherein the compound is a compound of formula (III):



wherein

Z is CH_2 ;

R_1 is chosen from C_{2-6} alkynyl, $-NR_3R_4$ and $-SR_5$, wherein the C_{2-6} alkynyl is optionally substituted with one or more groups independently chosen from: halogen, $-CN$, $-OH$, $-NH_2$, C_{3-8} cycloalkyl, 4-8 membered heterocyclyl, $-O(C_{1-6}$ alkyl), $-O(C_{1-6}$ haloalkyl), $-O(C_{3-8}$ cycloalkyl), $-O(4-8$ membered heterocyclyl), $-S(C_{1-6}$ alkyl), $-S(C_{3-8}$ cycloalkyl), $-S(4-8$ membered heterocyclyl), $-NH(C_{1-6}$ alkyl), $-N(C_{1-6}$ alkyl) $_2$, $-NH-CN$, $-NHCONH_2$, $-NHCO(C_{1-6}$ alkyl), $-CONR_aR_b$, $-COOR_c$ and $-COR_d$, wherein R_a , R_b , R_c and R_d are each independently chosen from hydrogen, C_{1-6} alkyl, $-(C_{1-6}$ alkyl)- $O-(C_{1-6}$ alkyl), $-(C_{1-6}$ alkyl)- OH , C_{3-8} cycloalkyl and 4-8 membered heterocyclyl; R_3 is independently chosen from hydrogen and C_{1-6} alkyl; R_4 and R_5 are each independently chosen from C_{3-8} cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl; and the above-

mentioned C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl and 5-6 membered heteroaryl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -CONH(C₁₋₆ alkyl) and -CON(C₁₋₆ alkyl)₂; further preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂ or -CONH(CH₃); and more preferably, R₁ is ethynyl;

R₁' is chosen from halogen, -CN, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1;

preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0;


R₂ is chosen from -NH₂ and oxo; and preferably, R₂ is -NH₂;

R₁₁, R₁₂ and R₁₃ are each independently chosen from hydrogen, -NH₂, -CN, C₁₋₆ alkyl and C₁₋₆ haloalkyl; preferably, R₁₁, R₁₂ and R₁₃ are each independently chosen from hydrogen and C₁₋₆ alkyl; and more preferably, R₁₁, R₁₂ and R₁₃ are all hydrogen;

Cy₂ is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and

L is absent, or L is S; and preferably, L is S.

15. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1, wherein the compound is a compound of formula (IV):



wherein

Z is CH₂ or O; and preferably, Z is CH₂;

R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; the above-mentioned C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -CONH(C₁₋₆ alkyl) and -CON(C₁₋₆ alkyl)₂; further preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -OCH₃, -NH(CH₃) or -N(CH₃)₂; and more preferably, R₁ is ethynyl;

R₁' is chosen from halogen, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0;

R₂ is -NH₂;


R₁₄ is chosen from hydrogen, -NH₂ and C₁₋₆ alkyl; preferably, R₁₄ is hydrogen or -NH₂; and more preferably, R₁₄ is hydrogen;

R₁₄' is C₁₋₆ alkyl;

Cy₂ is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heteroaryl or 9-10 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH and -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl); and

L is absent, or L is S; and preferably, L is S.

16. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1, wherein the compound is a compound of formula (V):



5 wherein

Z is CH₂ or O; and preferably, Z is CH₂;

R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, -NH₂, C₃₋₈ cycloalkyl, 4-8 membered heterocyclyl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O(C₃₋₈ cycloalkyl), -O(4-8 membered heterocyclyl), -S(C₁₋₆ alkyl), -S(C₃₋₈ cycloalkyl), -S(4-8 membered heterocyclyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NH-CN, -NHCONH₂, -NHCO(C₁₋₆ alkyl), -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-CN, C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl; and the above-mentioned C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -CONR_aR_b, -COOR_c and -COR_d, wherein R_a, R_b, R_c and R_d are each independently chosen from hydrogen, C₁₋₆ alkyl, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl, wherein the C₃₋₈ cycloalkyl and 4-8 membered heterocyclyl are each optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -O(C₁₋₆ alkyl); more preferably, R₁ is C₂₋₆ alkynyl, wherein the C₂₋₆ alkynyl is optionally substituted with one or more groups independently chosen from: -OH, -CONH₂, -O(C₁₋₆ alkyl), -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -NHCONH₂, -CONH(C₁₋₆ alkyl), -CONH(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl), -CON(C₁₋₆ alkyl)₂, -CON(C₁₋₆ alkyl)(C₁₋₆ alkyl-O-C₁₋₆ alkyl), -CONH(C₃₋₈ cycloalkyl), -COOH, -

COO(C₁₋₆ alkyl), -CO(C₁₋₆ alkyl), -CO(4-8 membered heterocyclyl) and -CO(4-8 membered heterocyclyl)-O-(C₁₋₆ alkyl); further preferably, R₁ is ethynyl, propynyl or butynyl, each of which is unsubstituted or substituted with -OH, -CONH₂, -OCH₃, -NH(CH₃), -N(CH₃)₂, -NHCONH₂, -CONH(C₁₋₃ alkyl), -CONH(CH₂CH₂)-O-(CH₃), -CON(CH₃)₂, -CON(CH₃)(CH₂CH₂-O-CH₃), -CONH(cyclopropyl), -COOH, -COO(CH₃), -CO(CH₃), -CO(azetidiny) or -CO(azetidiny)-O-(CH₃); and most preferably, R₁ is ethynyl substituted with -CONH(CH₃), -CONH(CH₂CH₃) or -CONH(CH₂CH₂)-O-(CH₃);

R₁' is chosen from halogen, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl), and n is 0 or 1; preferably, R₁' is halogen, and n is 0 or 1; and more preferably, n is 0;

10 R₂ is -NH₂;


R₁₅ and R₁₅' are each independently chosen from hydrogen, -NH₂, -CN, halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl and -(C₁₋₆ alkyl)-OH; preferably, R₁₅ and R₁₅' are each independently chosen from hydrogen, -NH₂, C₁₋₆ alkyl and -(C₁₋₆ alkyl)-OH; and more preferably, both R₁₅ and R₁₅' are hydrogen;

15 Cy₂ is phenyl or 5-14 membered heteroaryl, preferably phenyl or 5-10 membered heteroaryl, and more preferably 5-6 membered heteroaryl or 9-10 membered heteroaryl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and

L is absent, or L is S; and preferably, L is S.


17. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to any one of claims 1-16, wherein Cy₂ is phenyl, pyridyl, pyrimidyl, indazolyl, pyrrolopyridyl or 1,2,3,4-tetrahydro-1,5-naphthyridinyl, each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl.

18. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 17, wherein




each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl;


preferably, Cy₂ is chosen from



each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and


more preferably, Cy₂ is chosen from , each of which is optionally substituted with one or more groups independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl), -S(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl.

19. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 18, wherein




independently chosen from: halogen, -CN, -OH, C₁₋₆ alkyl, C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl) and -S(C₁₋₆ alkyl);

or



, which is optionally substituted with one or more groups independently chosen from: halogen, C₁₋₆ alkyl, -O(C₁₋₆ alkyl) and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH, -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl) and C₃₋₈ cycloalkyl; and preferably, both R₇ and R₈ are hydrogen;







5 or

























, which is optionally substituted with one or more groups independently chosen from: C₁₋₆ alkyl and -NR₇R₈, wherein R₇ and R₈ are each independently chosen from hydrogen, -(C₁₋₆ alkyl)-OH and -(C₁₋₆ alkyl)-O-(C₁₋₆ alkyl); and preferably, both R₇ and R₈ are hydrogen.

10













20. The compound or the pharmaceutically acceptable salt thereof, or the solvate, the racemic mixture, the enantiomer, the diastereomer or the tautomer thereof according to claim 1, which is chosen from:













No.	Structural formula	No.	Structural formula
1		2	
3		4	
5		6	

7		8	
9		10	
11 and 28		12 and 64	
13		14	
15		16	












17		18	
19		21	
20 and 24		22	
		23	
25		26	
27		29	











30		31	
32		33	
34		35	
36		37	
38		39	
40		41	
42		43	











44		45	
46		47	
48 and 49	 + 	50 and 51	 + 
52		53	
54		55	












56 and 57		58 and 59	
60		61	
62		63	
65 and 66		67	
		68	
69		70	

71		72	
73		74	
75		76	
77		78	
79		80	
81		82	
83		84	













85		86	
87		90	
88 and 89		91	
		92	
93		94	
95		96	















97		98	
99		100	
101		104	
102 and 103		106	
107		112	

108 and 109		110 and 111	
113 and 114		115	
117		116	
118 and 119		120	
		121	
		122	












124 and 125		123	
127		126	
129		128	
131		130	
133		132	
135		136	

137		138	
139		140	
141		142	
143		144	
145		146	
147		148	
149		150	

151		152	
153		154	
155		156	
157		158	
159		160	
161 and 162		163 and 164	

















165		166	
167		168	
169		170	
171		172	
173		174	
175		176	
177		178	

179		180	
181 and 182		183	
		184	
185		186	
187		188	
189		194	

















190 and 191		192 and 193	
195		196	
197		198	
199		200 and 201	
202			
203		204	

205		206	
207 and 208		209 and 210	
211		212	
213		214	
215		216	
217 and 218		220 and 221	








219		222	
223		224	
225		226	
227		228	
229		230	
231		232	
233		234	
235		236	

237		238	
239		240	
241		242	
243		244	
245		246	
247		248	
249		250	
251		252	

253		254	
255		256	
257		258	
259		260	
261		262	
263		264	
265		266	

267 and 268		269	
		270	
271		272	
273		274	
275		276	
277		278	
279		280	
281		282	

283		284	
285		286	
287		288	
289		290	
291		292	
293		294	
295		296	
297		298	

299		300	
301		302	
303		304	
305			

21. A pharmaceutical composition, comprising the compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20, and
 5 optionally comprising a pharmaceutically acceptable excipient.

22. A method of *in vivo* or *in vitro* inhibiting the activity of SHP2, comprising contacting SHP2 with an effective amount of the compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20.

10

23. Use of the compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20 in the manufacture of a medicament for treating or preventing a disease mediated by SHP2 or at least in part by SHP2, and preferably for treating or preventing cancer, Noonan Syndrome and LEOPARD Syndrome, wherein the
 15 cancer is preferably a solid tumor or hematologic malignancy, including leukemia, lymphoma and myeloma; and the cancer is more preferably chosen from breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal

cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma, acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia, chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, and myeloma (such as multiple myeloma).

24. A method of treating or preventing a disease in a subject, comprising administering to the subject in need thereof an effective amount of the compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20, wherein the disease is a disease mediated by SHP2 or at least in part by SHP2, and the disease is preferably cancer, Noonan Syndrome and LEOPARD Syndrome, wherein the cancer is preferably a solid tumor or hematologic malignancy, including leukemia, lymphoma and myeloma; and the cancer is more preferably chosen from breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma, acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia, chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, and myeloma (such as multiple myeloma).

25. The compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20, for use as a medicament.

26. The compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20, for use in treating or preventing a disease mediated by SHP2 or at least in part by SHP2, and preferably for use in treating or preventing cancer, Noonan Syndrome and LEOPARD Syndrome, wherein the cancer is preferably a solid tumor or hematologic malignancy, including leukemia, lymphoma and myeloma; and the cancer is more preferably chosen from breast cancer, melanoma, glioblastoma, esophageal cancer, gastric cancer, colon cancer, colorectal cancer, pancreatic cancer, lung cancer, head and neck cancer (such as squamous cell carcinoma of the head and neck), liver cancer, renal cancer, ovarian cancer, cervical cancer, prostate cancer, endometrial cancer, thyroid carcinoma, sarcoma, adrenal carcinoma, acute myelogenous leukemia (AML), juvenile acute myelogenous leukemia, chronic myelogenous leukemia (CML), acute lymphocytic leukemia (ALL), B-cell acute lymphocytic leukemia (B-ALL), acute lymphoblastic leukemia, chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), large B-cell lymphoma (LBCL), B-cell lymphoma, T-cell lymphoma, mantle cell lymphoma, follicular lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, myelodysplastic syndrome, and myeloma (such as multiple myeloma).

27. A pharmaceutical combination, comprising the compound and/or the pharmaceutically acceptable salt thereof according to any one of claims 1-20, and at least one additional therapeutic agent, wherein the additional therapeutic agent is preferably chosen from: an anti-neoplastic active agent, an anti-inflammatory agent or an immunomodulator, wherein the anti-neoplastic active agent includes a chemotherapeutic agent, an immune checkpoint inhibitor or agonist, and a targeted therapeutic agent.