

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2013217933 B2**

(54) Title
Methods of treating fibrosis

(51) International Patent Classification(s)
A61K 31/00 (2006.01) **A61K 31/357** (2006.01)
A61K 31/196 (2006.01) **A61P 1/04** (2006.01)
A61K 31/343 (2006.01)

(21) Application No: **2013217933** (22) Date of Filing: **2013.02.08**

(87) WIPO No: **WO13/117744**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/644,544	2012.05.09	US
12425027.5	2012.02.09	EP

(43) Publication Date: **2013.08.15**

(44) Accepted Journal Date: **2017.11.30**

(71) Applicant(s)
Nogra Pharma Limited

(72) Inventor(s)
Baroni, Sergio;Bellinvia, Salvatore;Viti, Francesca

(74) Agent / Attorney
Shelston IP Pty Ltd., Level 21, 60 Margaret Street, Sydney, NSW, 2000, AU

(56) Related Art
JOHNSON, L. A. et al. "Intestinal fibrosis is reduced by early elimination of inflammation in a mouse model of IBD" Inflamm. Bowel Dis. (2012) vol. 18, no. 3, pages 460-471
WO 2008/104557 A1
WO 2007/010514 A2
WO 2010/091892 A2
YU, J. et al. International Journal of Biochemistry and Cell Biology (2010) vol. 42, no. 6, pages 948-957



(51) International Patent Classification:

A61K 31/196 (2006.01) *A61K 31/357* (2006.01)
A61K 31/343 (2006.01) *A61P 1/04* (2006.01)

(21) International Application Number:

PCT/EP2013/052617

(22) International Filing Date:

8 February 2013 (08.02.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12425027.5 9 February 2012 (09.02.2012) EP
61/644,544 9 May 2012 (09.05.2012) US

(71) Applicant: **NOGRA PHARMA LIMITED** [IE/IE]; 33,
Sir John Rogerson's Quay, 2 Dublin (IE).

(72) Inventors; and

(71) Applicants : **BARONI, Sergio** [IT/IT]; Via Piazzola 3, I-24030 Villa D'Adda (BG) (IT). **BELLINIA, Salvatore** [IT/IT]; Via Brusafiera 12, I-33170 Pordenone (IT). **VITI, Francesca** [IT/IT]; Via Benedetto Croce, 44, I-20099 Sesto San Giovanni (IT).

(74) Agents: **LANE, Cathal Michael** et al.; **TOMKINS & CO**,
5 Dartmouth Road, 6 Dublin (IE).(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with information concerning authorization of rectification of an obvious mistake under Rule 91.3 (b) (Rule 48.2(i))

(88) Date of publication of the international search report:

21 November 2013


(15) Information about Correction:

Previous Correction:

see Notice of 3 October 2013

(54) Title: METHODS OF TREATING FIBROSIS

FIG. 1



(57) Abstract: The present disclosure is directed in part to methods of treating fibrosis, e.g., hepatic fibrosis and/or intestinal fibrosis, comprising administering to a patient in need thereof an effective amount of a disclosed compound.

METHODS OF TREATING FIBROSIS

[0001] This application claims priority to EP12425027.5, filed February 9, 2012, and U.S.S.N. 61/644,544, filed May 9, 2012, each of which is incorporated by reference in its entirety.

SEQUENCE LISTING

[0001.1] The instant application contains a Sequence Listing which has been submitted in electronically in text format and is hereby incorporated by reference in its entirety. Said text copy, created on May 21, 2013, is named PS966PCT_SL.txt and is 3,284 bytes in size.

BACKGROUND

[0002] Fibrosis is the formation of excess fibrous connective tissue in an organ or tissue in a reparative or reactive process, e.g., healing, usually because of injury or long-term inflammation. Fibrosis causes the affected tissues to harden and/or swell and reduces the flow of fluids through these tissues. As result, tissues with fibrosis may not be able to function properly.

[0003] For example, hepatic fibrosis can be described as the wound-healing response to chronic hepatic injury as a result of, for example, alcohol and/or drug abuse, viral and parasitic infection (e.g., hepatitis such as hepatitis B or C), nonalcoholic steatohepatitis (NASH), iron and copper overload, and autoimmune diseases. All chronic liver diseases can lead to liver fibrosis with the principle causes chronic viral hepatitis B and alcoholic liver disease. The hepatic endocannabinoid system mediates both pro- and anti-fibrogenic effects by activating distinct signaling pathways that differentially effect proliferation and death of fibrogenic cell types. Over time this process can result in cirrhosis of the liver in which the architectural organization of the functional units of the liver becomes so disrupted that blood flow through the liver and liver function become disrupted.

[0004] Liver fibrosis represents a common and difficult challenge of worldwide importance. At present, the only curative treatment for end stage cirrhosis is transplantation, but even in the developed world, the number of donor organs available and the clinical condition of the potential recipient limit the applicability of this technique. The development of fibrosis, and particularly cirrhosis, is associated with a significant morbidity and mortality. Thus, there is a considerable imperative to develop antifibrotic strategies that are applicable to liver fibrosis.


[0005] Renal fibrosis causes significant morbidity and mortality as the primary acquired lesion leading to the need for dialysis or kidney transplantation. Fibrosis can occur in either the filtering or reabsorptive component of the nephron, the functional unit of the kidney. Fibrosis can also occur in the heart, e.g., cardiac fibrosis can occur as a thickening of a heart valve.

[0006] Intestinal fibrosis is a common complication of inflammatory bowel disease (IBD) that can become symptomatic and may require surgical intervention if stricture formation ensues. Most of the traditional and novel mechanisms underlying intestinal fibrosis are associated with chronic inflammation.

[0007] Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

SUMMARY

[0007.1] According to a first aspect, the present invention provides a method of treating fibrosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound, wherein the compound is




2-methoxy-3-(4'-aminophenyl) propionic acid (compound 34) represented by the formula: or a pharmaceutically acceptable salt or a stereoisomer thereof, wherein the fibrosis is selected from the group consisting of hepatic fibrosis, intestinal fibrosis, and idiopathic pulmonary fibrosis.

[0007.2] According to a second aspect, the present invention provides a method of the invention, wherein the compound is (S)-2-methoxy-3-(4'-aminophenyl) propionic acid, or a pharmaceutically acceptable salt thereof.

[0007.3] According to a third aspect, the present invention provides a method of the invention, wherein the compound is (R)-2-methoxy-3-(4'-aminophenyl) propionic acid, or a pharmaceutically acceptable salt thereof

[0007.4] According to a fourth aspect, the present invention provides a method of treating fibrosis, comprising administering to a patient in need thereof, a pharmaceutical preparation comprising a therapeutically effective amount of a compound, wherein the compound is 2-methoxy-3-(4'-aminophenyl) propionic acid or a pharmaceutically acceptable salt or a stereoisomer thereof; and one or more pharmaceutically acceptable carriers, wherein the fibrosis is selected from the group consisting of hepatic fibrosis, intestinal fibrosis, and idiopathic pulmonary fibrosis.

[0007.5] According to a fifth aspect, the present invention provides a use of a compound, wherein the compound is 2-methoxy-3-(4'-aminophenyl) propionic acid (compound 34) represented by the formula:




or a pharmaceutically acceptable salt or a stereoisomer thereof, in the manufacture of a medicament for treating fibrosis, wherein the fibrosis is selected from the group consisting of hepatic fibrosis, intestinal fibrosis, and idiopathic pulmonary fibrosis

[0007.6] Unless the context clearly requires otherwise, throughout the description and the claims, the words “comprise”, “comprising”, and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of “including, but not limited to”.

[0007.7] Provided herein, for example, are methods for treating fibrosis comprising administering compounds disclosed herein to a patient. Compound for use in such methods are also provided. For example, a method of preventing or treating hepatic fibrosis and/or intestinal fibrosis in a subject or patient in need thereof or suffering from same is provided, comprising administering a therapeutically and/or pharmaceutically acceptable

amount of a compound represented by:



R₁₀ is selected from the group consisting of H and C₁-C₆alkyl;

R₁₁ is selected from the group consisting of H, C₁-C₆alkyl, and -C(O)-C₁-C₆alkyl;


R₅ is C₁-C₆alkyl; and pharmaceutically acceptable salts thereof.

[0008] Suitably, in some embodiments, R₅ can be H.

[0009] Suitably, the C₁-C₆alkyl functionalities of R₁₀, R₁₁ and/or R₅ may be independently selected from the group consisting of: methyl, ethyl, propyl, pentyl and hexyl. The alkyl functionality may be linear or branched. Methyl and ethyl alkyl functionalities are particularly preferred with methyl being the most preferred alkyl functionality.

[0010] Suitably, the nitrogen may be attached to the ring at the 2', 3', 4' or 5' positions. Preferably, the nitrogen is attached at the 3' or 4' position. The 4' position is most preferred.

[0011] Exemplary contemplated compounds may include formulas A' and A'', wherein R₁₀, R₁₁ and R₅ are provided above and herein:



[0012] Suitably, in some embodiments, R₅ can be H.

[0013] Suitably, the C₁-C₆alkyl functionalities of R₁₀, R₁₁ and/or R₅ may be independently selected from the group consisting of: methyl, ethyl, propyl, pentyl and hexyl. The alkyl functionality may be linear or branched. Methyl and ethyl alkyl functionalities are particularly preferred with methyl being the most preferred alkyl functionality. In certain embodiments, R₅ is methyl, ethyl, propyl or is e.g., methyl. In certain embodiments, R₅ is methyl, ethyl, propyl or is e.g., methyl.

[0014] Also provided herein is a method of preventing or treating hepatic or intestinal fibrosis, comprising administering to a patient in need thereof, a pharmaceutical preparation comprising a PPAR_γ agent such as a compound disclosed herein. Contemplated herein are methods of preventing or treating renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, idiopathic pulmonary fibrosis, myelofibrosis, retroperitoneal fibrosis, and/or nephrogenic systemic fibrosis, comprising administering to a patient in need thereof, a pharmaceutical preparation comprising a PPAR_γ agent such as a compound disclosed herein. For example, methods disclosed herein may include methods wherein the patient is human. Accordingly, the compounds described herein may be used in the treatment and/or prevention of these diseases/conditions.

[0015] These and other aspects and advantages of the disclosure will become apparent upon consideration of the following figures, detailed description, and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] Figure 1 depicts the body weight evolution of mice upon administration of CCl₄, JWH-133 and compound 34.

[0017] Figure 2 (A) depicts an explanation of metavir scores and (B) depicts the effect of CCl₄, JWH-133+ CCl₄ and compound 34+ CCl₄ on the liver and specifies a metavir score for each group.

- [0018] Figure 3 (A) depicts the metavir score and (B) adapted metavir score of livers from mice subjected to CCl₄, JWH-133+ CCl₄ and compound 34+ CCl₄.
- [0019] Figure 4 depicts the effect of CCl₄, JWH-133+ CCl₄ and compound 34+ CCl₄ on α -SMA and collagen-1.
- [0020] Figure 5 depicts the intestinal fibrosis study schedule.
- [0021] Figure 6 depicts the effect of compound 34 on the colons of mice having intestinal fibrosis.
- [0022] Figure 7 depicts the anti-inflammatory effect of compound 34 on the colons of mice having intestinal fibrosis.
- [0023] Figure 8 depicts the effect of compound 34 on the colons of mice having intestinal fibrosis.
- [0024] Figure 9 depicts the effect of compound 34 on the colons of mice having intestinal fibrosis using an immunohistochemical analysis of the main markers of fibrosis (Collagen I-III, CTGF, SMAD 2/3, PDGF, α -SMA, and TGF- β 1).
- [0025] Figure 10 depicts the effect of compound 34 on the colons of mice having intestinal fibrosis using a Western Blot analysis.
- [0026] Figure 11 shows a body weight curve in mice treated with DSS and DSS+GED.
- [0027] Figure 12 shows the level of fibrosis in the colon of mice treated with DSS and DSS+GED compared to control mice receiving regular water.
- [0028] Figure 13 depicts microscopic findings of colons from DSS and DSS + GED mice chronic colitis. Colon tissues were fixed in 4% PFA and transversal sections (4 μ m) were stained with May-Grunwald-Giemsa(H&E) and Masson's trichrome. DSS mice showed severe inflammatory degree and fibrosis both in submucosa and serosa compared to controls, while GED treatment determined the restoring of normal intestinal wall architecture. Data are expressed as mean \pm SEM; * = p < 0.05; ** = p < 0.01 and *** = p < 0.005 vs H₂O.
- [0029] Figure 14 shows immunoistochemistry and immunoblotting of the main markers of fibrosis. The microphotograph shows a significant reduction of Collagen I-III (a), α -SMA (b) in DSS mice treated with GED, compared to the mice with DSS-induced chronic colitis. The protein levels of these markers were measured also by Western blotting confirming the expression profile induced by GED. Representative blots are shown. * = p < 0.05, ** = p < 0.01. N = 6 mice for the control group, 8 mice for DSS group, 8 mice for DSS + GED group.
- [0030] Figure 15 depicts a microphotograph showing a significant reduction TGF- β 1 (a), SMAD 2/3(b), CTGF (c) in the DSS mice treated with GED, compared to the mice with DSS-induced chronic colitis.
- [0031] Figure 16 shows the effect of GED administration on IL-13 protein levels in chronic DSS-treated mice and on its downstream mediators, TGF- β 1 and CTGF. IL-13, TGF- β 1 and CTGF protein levels were determined by Western blot analysis. Data are presented as percentage of mean controls values. Representative blots are shown. ** = p < 0.01. N = 6 mice for the control group, 8 mice for DSS group, 8 mice for DSS + GED group.

DETAILED DESCRIPTION

[0032] The disclosure is based, in part, upon the discovery that certain compounds disclosed herein have the ability to prevent or treat fibrosis, for example, in the liver or in the colon. In one aspect, the disclosure is directed to methods of preventing or treating fibrosis, e.g. hepatic fibrosis and/or intestinal fibrosis, in patients in need thereof. The disclosed methods comprise administering a compound disclosed herein.

[0033] Before further description, certain terms employed in the specification, examples and appended claims are collected here. These definitions should be read in light of the remainder of the disclosure and understood as by

a person of skill in the art. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art.

[0034] A “patient,” “subject” or “host” to be treated by the subject method may mean either a human or non-human animal, e.g. a small mammal such as a mouse or rat, and including horse, cow, dog, cat, etc.

[0035] The term “therapeutic agent” is art-recognized and refers to any chemical moiety that is a biologically, physiologically, or pharmacologically active substance that acts locally and/or systemically in a subject. Examples of therapeutic agents, also referred to as “drugs”, are described in well-known literature references such as the Merck Index, the Physicians Desk Reference, and The Pharmacological Basis of Therapeutics, and they include, without limitation, medicaments; vitamins; mineral supplements; substances used for the treatment, prevention, diagnosis, cure or mitigation of a disease or illness; substances which affect the structure or function of the body; or pro-drugs, which become biologically active or more active after they have been placed in a physiological environment.

[0036] The term “therapeutic effect” is art-recognized and refers to a local and/or systemic effect in animals, particularly mammals, and more particularly humans caused by a pharmacologically active substance. The term thus means any substance intended for use in the diagnosis, cure, mitigation, treatment or prevention of disease or in the enhancement of desirable physical or mental development and/or conditions in an animal or human. The phrase “therapeutically-effective amount” means that amount of such a substance that produces some desired local or systemic effect at a reasonable benefit/risk ratio applicable to any treatment. The therapeutically effective amount of such substance will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. For example, certain compositions of the present invention may be administered in a sufficient amount to produce a at a reasonable benefit/risk ratio applicable to such treatment.

[0037] The term “treating” is art-recognized and refers to curing as well as ameliorating at least one symptom of any condition or disease.

[0038] The term “alkyl” is art-recognized, and includes saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In certain embodiments, a straight chain or branched chain alkyl has about 30 or fewer carbon atoms in its backbone (e.g., C₁-C₃₀ for straight chain, C₃-C₃₀ for branched chain), and alternatively, about 20 or fewer, e.g. from 1 to 6 carbons. Likewise, cycloalkyls have from about 3 to about 10 carbon atoms in their ring structure, and alternatively about 5, 6 or 7 carbons in the ring structure. The term “alkyl” is also defined to include halosubstituted alkyls.

[0039] Moreover, the term “alkyl” (or “lower alkyl”) includes “substituted alkyls”, which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents may include, for example, a hydroxyl, a carbonyl (such as a carboxyl, an alkoxycarbonyl, a formyl, or an acyl), a thiocarbonyl (such as a thioester, a thioacetate, or a thioformate), an alkoxyl, a phosphoryl, a phosphonate, a phosphinate, an amino, an amido, an amidine, an imine, a cyano, a nitro, an azido, a sulfhydryl, an alkylthio, a sulfate, a sulfonate, a sulfamoyl, a sulfonamido, a sulfonyl, a heterocyclyl, an aralkyl, or an aromatic or heteroaromatic moiety. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain may themselves be substituted, if appropriate. For instance, the substituents of a substituted alkyl may include substituted and unsubstituted forms of amino, azido, imino, amido, phosphoryl (including phosphonate and phosphinate), sulfonyl (including sulfate, sulfonamido, sulfamoyl and sulfonate), and silyl groups, as well as

ethers, alkylthios, carbonyls (including ketones, aldehydes, carboxylates, and esters), -CN and the like. Exemplary substituted alkyls are described below. Cycloalkyls may be further substituted with alkyls, alkenyls, alkoxys, alkylthios, aminoalkyls, carbonyl-substituted alkyls, -CN, and the like.

[0040] The terms ortho, meta and para are art-recognized and refer to 1,2-, 1,3- and 1,4-disubstituted benzenes, respectively. For example, the names 1,2-dimethylbenzene and ortho-dimethylbenzene are synonymous.

[0041] The definition of each expression, e.g. alkyl, m, n, and the like, when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

[0042] Certain compounds contained in compositions of the present invention may exist in particular geometric or stereoisomeric forms. In addition, compounds of the present invention may also be optically active. The present invention contemplates all such compounds, including cis- and trans-isomers, *R*- and *S*-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

[0043] It will be understood that “substitution” or “substituted with” includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, or other reaction.

[0044] The term “substituted” is also contemplated to include all permissible substituents of organic compounds. In a broad aspect, the permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. Illustrative substituents include, for example, those described herein above. The permissible substituents may be one or more and the same or different for appropriate organic compounds. For purposes of this invention, the heteroatoms such as nitrogen may have hydrogen substituents and/or any permissible substituents of organic compounds described herein which satisfy the valences of the heteroatoms. This invention is not intended to be limited in any manner by the permissible substituents of organic compounds.

[0045] For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 67th Ed., 1986-87, inside cover. Also for purposes of this invention, the term “hydrocarbon” is contemplated to include all permissible compounds having at least one hydrogen and one carbon atom. In a broad aspect, the permissible hydrocarbons include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic organic compounds that may be substituted or unsubstituted.


[0046] The term “pharmaceutically-acceptable salts” is art-recognized and refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds, including, for example, those contained in compositions of the present invention.

[0047] The term “pharmaceutically acceptable carrier” is art-recognized and refers to a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting any subject composition or component thereof from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the subject composition and its components and not injurious to the patient. Some examples of materials which may serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as

sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

Compounds

[0048] Compounds contemplated for use in one or more of the disclosed methods include compounds represented by formula I, or a pharmaceutically acceptable salt, enantiomer or stereoisomer thereof:



wherein:

R^1 and R^2 are each independently selected from the group consisting of H and C_{1-6} alkyl; or R^1 and R^2 may be taken together with the nitrogen atom to which they are attached to form an aromatic or aliphatic ring with 5 or 6 atoms;

Y and Z are independently selected from the group consisting of H, OH, COOH, $-OR^3$, $-CH(OR_3)COOH$; and

R^3 is selected from the group consisting of H, phenyl, benzyl, vinyl, allyl, C_{1-6} alkyl or C_{1-6} alkyl substituted by one or more halogens; or a pharmaceutically acceptable salt thereof.


[0049] Suitably, the C_1 - C_6 alkyl functionalities may be independently selected from the group consisting of: methyl, ethyl, propyl, pentyl and hexyl. The alkyl functionality may be linear or branched. Methyl and ethyl alkyl functionalities are particularly preferred with methyl being the most preferred alkyl functionality.

[0050] In an embodiment, Y may be H or COOH. For example, Y may be H and Z may be $CH(OR^3)COOH$, or Y may be COOH and Z may be $-OR^3$. In some embodiments, R^3 may be methyl, ethyl, n-propyl, or isopropyl.

[0051] In other embodiments, the NR^1R^2 moiety of Formula I may be in the 4' position or may be in the 3' position. In certain embodiments, R^1 and R^2 are H.

[0052] Suitably, in some embodiments, R_3 can be H.

[0053] Exemplary compounds also include those represented by formulas IIa or IIb or a pharmaceutically acceptable salt, enantiomer or stereoisomer of:



wherein:

R^1 and R^2 are independently selected from the group consisting of H and C_{1-6} alkyl; or R^1 and R^2 may be taken together with the nitrogen atom to which they are attached to form an aromatic or aliphatic ring with 5 or 6 atoms;

R^6 is selected from the group consisting of $-NR^9OH$, OH, and $-OR^9$;

R^9 is C_{1-6} alkyl;

R^4 is selected from the group consisting of H, halo, phenyl, benzyl, vinyl, allyl, C_{1-6} alkyl and C_{1-6} alkyl substituted by one or more halogens;

R^5 and R^7 are independently selected from the group consisting of hydrogen or halo, or


R^4 and R^5 , or R^4 and R^6 together, form a fused heterocyclic ring with 5 or 6 atoms, optionally substituted with halo or C_{1-6} alkyl; and

A is a fused heterocyclic ring; or a pharmaceutically acceptable salt thereof.

Suitably, the C_1 - C_6 alkyl functionalities may be independently selected from the group consisting of: methyl, ethyl, propyl, pentyl and hexyl. The alkyl functionality may be linear or branched. Methyl and ethyl alkyl functionalities are particularly preferred with methyl being the most preferred alkyl functionality.

[0054] In certain embodiments, the NR^1R^2 moiety of formula IIa may be in the 4' position or may be in the 3' position. In certain embodiments, R^1 and R^2 are H. R^9 in some embodiments, may be methyl, ethyl, n-propyl, or isopropyl.

[0055] In some embodiments a contemplated compound can be represented by




wherein p is 1 or 2;

R^6 is selected from the group consisting of OH or $-OR^9$, wherein R^9 is defined above; and

R^{10} is selected independently for each occurrence from the group consisting of H, halo, or C_{1-6} alkyl, e.g., methyl or ethyl.

[0056] The disclosure also provides, at least in part, compounds represented by the formulas depicted below, for use in the disclosed methods.

[0057] For example, contemplated herein for use in disclosed methods are compounds of formula III:



III

wherein X is C_1 - C_3 alkylene, optionally substituted with one, two or three substituents selected from halogen or hydroxyl;

R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

R_2 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl;

R_3 is independently selected, for each occurrence from the group consisting of hydrogen, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, cyano, C_3 - C_6 cycloalkyl, halogen, hydroxyl, and nitro;

R_4 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl;

R_5 is C_1 - C_6 alkyl;


or pharmaceutically acceptable salts or N-oxides thereof.

Suitably, in some embodiments, R_5 can be H.


Suitably, the C_1 - C_6 alkyl functionalities may be independently selected from the group consisting of: methyl, ethyl, propyl, pentyl and hexyl. The alkyl functionality may be linear or branched. Methyl and ethyl alkyl functionalities are particularly preferred with methyl being the most preferred alkyl functionality.

[0058] In one embodiment, R_1 of formula III can be C_1 - C_6 alkyl, such as methyl. In one embodiment, R_2 can be hydrogen. In another embodiment, R_3 can be selected from the group consisting of hydrogen, C_1 - C_6 alkyl, halogen, and hydroxyl. In a further embodiment, R_3 can be hydrogen. In one embodiment, R_4 and R_5 can each be C_1 - C_6 alkyl. In another embodiment, R_4 may be hydrogen and R_5 may be methyl. In one embodiment, X may be $(CH_2)_n$, wherein n is 1 or 2, such as 1.

[0059] In another embodiment, the moiety $-NR_2-COR_1$ of formula III can be in the meta or the para position relative to X as shown in formulas IIIa and IIIb.



[0060] The disclosure also provides compounds represented by formula IV, as depicted below, which may be used in the disclosed methods, e.g.:



wherein R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;


R_2 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl;

R_3 is independently selected, for each occurrence from the group consisting of hydrogen, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, cyano, C_3 - C_6 cycloalkyl, halogen, hydroxyl, and nitro;

R_5 is hydrogen or C_1 - C_6 alkyl;

or pharmaceutically acceptable salts or N-oxides thereof.

[0061] Compounds of Formula V are also contemplated as shown below:



wherein R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

R_3 is independently selected, for each occurrence from the group consisting of hydrogen, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, cyano, C_3 - C_6 cycloalkyl, halogen, hydroxyl, and nitro;

R_4 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl;


R_5 is hydrogen or C_1 - C_6 alkyl; and

A is a fused five or six membered heterocycle;

or pharmaceutically acceptable salts or N-oxides thereof.

[0062] In one embodiment, R_1 can be C_1 - C_6 alkyl, such as methyl. In another embodiment, R_1 and R_3 can each be C_1 - C_6 alkyl, such as methyl. In one embodiment, R_2 can be hydrogen.

[0063] In some embodiments, a compound can be represented by




wherein p is 1 or 2;

R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

R_4 and R_8 are each independently selected from the group consisting of hydrogen and C_1 - C_6 alkyl;

or pharmaceutically acceptable salts or N-oxides thereof.

[0064] For example, in one embodiment, contemplated compounds for use in disclosed methods include compounds of formula A:




R_{10} is selected from the group consisting of H and C_1 - C_6 alkyl (e.g. methyl);

R_{11} is selected from the group consisting of H, C_1 - C_6 alkyl, and $-C(O)$ - C_1 - C_6 alkyl; (e.g., R_{11} may be H, methyl, $-C(O)$ -methyl, or $-C(O)$ -ethyl).


R_5 is C_1 - C_6 alkyl (e.g., R_5 may be methyl, ethyl, or propyl);

or pharmaceutically acceptable salts or N-oxides thereof.

[0065] For example, contemplated herein for use in disclosed methods are formulas A' and A'':




[0066] Exemplary compounds contemplated herein for use in disclosed methods include:



or a pharmaceutically acceptable salt thereof.

[0067] In some embodiments, contemplated compounds include: 4-amino-N-hydroxy-2-methoxybenzamide (compound 13); 6-methoxy quinoline-5-carboxylic acid (compound 36); 6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid (compound 37); and 5-diisopropylaminosalicylic acid (compound 38).

[0068] Other exemplary compounds include those represented by:




[0069] Compounds contemplated herein include racemic mixtures, and enantiomers of compounds, for example: (±)-2-hydroxy-3-(3'-aminophenyl) propionic acid (compound 20); (±)-2-methoxy-2-(4'-aminophenyl) acetic acid (compound 23); (±)-2-ethoxy-2-(3'-aminophenyl) acetic acid (compound 32); (±)-2-ethoxy-2-(4'-aminophenyl) acetic acid (compound 33); (±)-2-methoxy-3-(4'-aminophenyl) propionic acid (compound 34) "±34" (racemic form); (±)-2-ethoxy-3-(4'-aminophenyl) propionic acid (compound 39); (±)-2-ethoxy-3-(3'-aminophenyl) propionic acid (compound 40).


[0070] For example, the compounds used in the methods disclosed herein can be enantiomers of the following racemic mixtures: (R,S)-2-hydroxy-2-(3-aminophenyl)acetic acid (compound 10); (R,S)-2-hydroxy-2-(4-aminophenyl)acetic acid (compound 11); (R,S)-2-hydroxy-3-(4'-aminophenyl)propionic acid (compound 21); (R,S)-2-methoxy-2-(3'-aminophenyl)acetic acid (compound 22); (R,S)-2-methoxy-3-(3'-aminophenyl)propionic acid (compound 35); (R,S)-2-methoxy-3-(4-aminophenyl)propionic acid (compound 34), as well as enantiomers, e.g.: (+) 2-S-methoxy-3-(4-aminophenyl)propionic acid; (-) 2-R-methoxy-3-(4-aminophenyl)propionic acid. In some embodiments, (R,S)-2-hydroxy-3-(4'-aminophenyl)propionic acid (compound 21) or (±)-2-hydroxy-3-(3'-aminophenyl) propionic acid (compound 20) are preferred.

[0071] Other racemic type mixtures of compounds contemplated include: e.g. (±)-2-hydroxy-2-(3'-aminophenyl)acetic acid (compound 10); (±)-2-hydroxy-2-(4'-aminophenyl)acetic acid (compound 11); (±)-2-hydroxy-3-(4'-aminophenyl)propionic acid (compound 21) and (±)-2-methoxy-2-(3'-aminophenyl)acetic acid (compound 22).

[0072] Further compounds contemplated for use in the disclosed methods: 5-aminosalicylo-hydroxamic acid (compound 5); 3-dimethylaminosalicylic acid (compound 6); 2-methoxy-4-aminobenzoic acid (compound 7); 2-methoxy-5-aminobenzoic acid (compound 8); 5-methylaminosalicylic acid (compound 9); 4-methylaminosalicylic acid (compound 12); 4-acetylaminosalicylic acid (compound 16); 2-ethoxy-4-aminobenzoic acid (compound 18); 2-ethoxy-5-aminobenzoic acid (compound 19); 4-dimethylaminosalicylic acid (compound 24); 2-ethoxy-4-aminobenzoylhydroxamic acid (compound 25); 6-hydroxyquinoline-5-carboxylic acid (compound 27); 2-(2-propyl)oxy-4-aminobenzoic acid (compound 30); 4-(1-piperazinyl)salicylic acid (compound 41); (R,S) 5-oxaquinoline-6-carboxylic acid (compound 15); 6-methoxy quinoline-5-carboxylic acid (compound 36); 6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid (compound 37); 5-diisopropylaminosalicylic acid (compound 38); and 4-diisopropylaminosalicylic acid (compound 42).

[0073] Contemplated compounds, and pharmaceutical compositions, comprising at least one compound, may be selected from the group consisting of: N-acetyl-(R)-(-)-3-(4-aminophenyl)-2-methoxypropionic acid, N-acetyl-(S)-(-)-3-(4-aminophenyl)-2-methoxypropionic acid; and racemic N-acetyl-(S)-(-)-3-(4-aminophenyl)-2-methoxypropionic acid, or a compound selected from:





or pharmaceutically acceptable salts or N-oxides thereof.

[0074] Other contemplated compounds include compound 20, compound 21, 4-acetamido-N-hydroxy-2-methoxybenzamide; 1-acetyl-6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid, 5-acetamido-2hydroxybenzoic acid (e.g., acetylated 5-aminosalicylic acid) or pharmaceutically acceptable salts or N-oxides thereof.

[0075] Also contemplated herein for use in the disclosed methods are compositions that include a compound represented by the depicted formulas and e.g., a pharmaceutically acceptable carrier.

[0076] Methods for making contemplated compounds may be found for example in WO2007/010516 and WO2007/010514, each hereby incorporated by reference in their entirety.

Methods

[0077] Methods of preventing or treating fibrosis, such as hepatic fibrosis and/or intestinal fibrosis, form part of this disclosure. Such methods may comprise administering to a patient in need thereof or a patient at risk, a pharmaceutical preparation comprising a PPARy agent such as a compound disclosed herein, e.g., Formulas I, IIa, or IIb, e.g., compounds 17, 29, 34 or 39. For example, a method of preventing or treating hepatic fibrosis is provided comprising administering to a patient in need thereof a compound disclosed herein. Alternatively, a method of preventing or treating intestinal fibrosis is provided comprising administering to a patient in need thereof a compound disclosed herein.

[0078] Patients treated using an above method may or may not have detectable fibrosis. In some embodiments, the patient has at least about a 5%, 10%, 20%, 30%, 40% or even 50% or more reduction in the amount of fibrosis present in the patient after administering a disclosed compound, e.g., compounds 17, 29, 34 or 39, after e.g. 1 day, 2 days, 1 week, 1 month or 6 months or more. Administering such a compound may be on, e.g., at least a daily basis. The compound may be administered orally. The delay of clinical manifestation of fibrosis in a patient as a consequence of administering a compound disclosed here may be at least e.g., 6 months, 1 year, 18 months or even 2 years or more as compared to a patient who is not administered a compound such as one disclosed herein.

[0079] A patient in need may have hepatic fibrosis that has developed into cirrhosis. A patient at risk of hepatic fibrosis may include those patients with hepatitis B, hepatitis C or nonalcoholic steatohepatitis (NASH). NASH is included in the spectrum of nonalcoholic fatty liver diseases, including steatosis and cirrhosis. NASH is a component of the metabolic syndrome, which is characterized by obesity, type 2 diabetes mellitus, and dyslipidemia, and can eventually lead to hepatocellular carcinoma.

[0080] Methods of treating disorders associated with hepatic fibrosis are also provided, such as the treatment of at least one of: certain storage diseases and inborn errors of metabolism, such as, alpha 1-antitrypsin deficiency, copper storage diseases (e.g., Wilson's disease), fructosemia, galactosemia, glycogen storage diseases (e.g., Types III, IV, VI, IX and X), iron-overload syndromes (e.g., hemochromatosis), lipid abnormalities (e.g. Gaucher's disease), peroxisomal disorders (e.g., Zellweger syndrome), and tyrosinemia; bacterial infections (e.g., brucellosis); parasitic infections (e.g., echinococcosis); NASH; viral infections (e.g., hepatitis B or hepatitis C, including chronic hepatitis B or C); Budd-Chiari syndrome; heart failure; hepatic veno-occlusive disease; and portal vein thrombosis. Methods of treating congenital hepatic fibrosis are also contemplated. The composition may be administered orally.

[0081] Abuse of drugs and/or alcohol has been implicated in cases of hepatic fibrosis. Contemplated herein are methods of treating hepatic fibrosis in a patient with a history of drug and/or alcohol abuse. For example, a patient with a history of abusing at least one of the following: alcohol, amiodarone, chlorpromazine, isoniazid, methotrexate, methyl dopa, oxyphenisatin and, tolbutamide.

[0082] A patient at risk of intestinal fibrosis may include those patients with ulcerative colitis, inflammatory bowel disease, or Crohn's disease. A patient at risk may also include those patients with an early age at diagnosis of Crohn's or colitis, extensive and/or severe of colonic disease, patients with the presence of primary sclerosing cholangitis, and/or patient's having a family history of cancer.

[0083] Methods of treating disorders associated with intestinal fibrosis are also provided, such as the treatment of at least one of: ulcerative colitis, inflammatory bowel disease, or Crohn's disease.

[0084] Contemplated herein are methods of preventing or treating renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, idiopathic pulmonary fibrosis, myelofibrosis, retroperitoneal fibrosis, or nephrogenic systemic fibrosis, comprising administering to a patient in need thereof, a pharmaceutical preparation comprising a PPAR γ agent such as a compound disclosed herein.

[0085] The compounds of the invention can be used alone or in combination with each other where by at least two compounds of the invention are used together in a single composition or as part of a treatment regimen. The compounds of the invention may also be used in combination with other drugs for treating drug and/or alcohol abuse, renal fibrosis, cardiac fibrosis, endomyocardial fibrosis, idiopathic pulmonary fibrosis, myelofibrosis, retroperitoneal fibrosis, or nephrogenic systemic fibrosis, drug and/or alcohol abuse.

[0086] Generally, a therapeutically effective amount of active component will be in the range of from about 0.1 mg/kg to about 100 mg/kg, optionally from about 1 mg/kg to about 100 mg/kg, optionally from about 1 mg/kg to 10 mg/kg. The amount administered will depend on variables such as the type and extent of disease or indication to be treated, the overall health status of the particular patient, the relative biological efficacy of the binding protein delivered, the formulation of the binding protein, the presence and types of excipients in the formulation, and the route of administration. The initial dosage administered may be increased beyond the upper level in order to rapidly achieve the desired blood-level or tissue level, or the initial dosage may be smaller than the optimum and the daily dosage may be progressively increased during the course of treatment depending on the particular situation. Human dosage can be optimized, e.g., in a conventional Phase I dose escalation study designed to run from 0.5 mg/kg to 20

mg/kg. Dosing frequency can vary, depending on factors such as route of administration, dosage amount and the disease condition being treated. Exemplary dosing frequencies are once per day, once per week and once every two weeks.

[0087] Contemplated formulations or compositions comprise a disclosed compound and typically include a pharmaceutically acceptable carrier.

[0088] Compositions contemplated herein may be administered by various means, depending on their intended use, as is well known in the art. For example, if compositions of the present invention are to be administered orally, they may be formulated as tablets, capsules, granules, powders or syrups. Alternatively, formulations of the present invention may be administered parenterally as injections (intravenous, intramuscular or subcutaneous), drop infusion preparations or suppositories. For application by the ophthalmic mucous membrane route, compositions of the present invention may be formulated as eyedrops or eye ointments. These formulations may be prepared by conventional means, and, if desired, the compositions may be mixed with any conventional additive, such as an excipient, a binder, a disintegrating agent, a lubricant, a corrigent, a solubilizing agent, a suspension aid, an emulsifying agent or a coating agent.

[0089] In formulations of the subject invention, wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants may be present in the formulated agents.

[0090] Subject compositions may be suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal, aerosol and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of composition that may be combined with a carrier material to produce a single dose vary depending upon the subject being treated, and the particular mode of administration.

[0091] Methods of preparing these formulations include the step of bringing into association compositions of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association agents with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

[0092] Formulations suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia), each containing a predetermined amount of a subject composition thereof as an active ingredient. Compositions of the present invention may also be administered as a bolus, electuary, or paste.

[0093] In solid dosage forms for oral administration (capsules, tablets, pills, film-coated tablets, sugar-coated tablets, powders, granules and the like), the subject composition is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, acetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate,

magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

[0094] Formulations and compositions may include micronized crystals of the disclosed compounds. Micronization may be performed on crystals of the compounds alone, or on a mixture of crystals and a part or whole of pharmaceutical excipients or carriers. Mean particle size of micronized crystals of a disclosed compound may be for example about 5 to about 200 microns, or about 10 to about 110 microns.

[0095] A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the subject composition moistened with an inert liquid diluent. Tablets, and other solid dosage forms, such as film coated tablets or sugar coated tablets, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art.

[0096] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the subject composition, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, cyclodextrins and mixtures thereof.

[0097] Suspensions, in addition to the subject composition, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

[0098] Formulations for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing a subject composition with one or more suitable non-irritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the body cavity and release the active agent. Formulations which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

[0099] Dosage forms for transdermal or topical administration of a subject composition include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active component may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

[00100] The ointments, pastes, creams and gels may contain, in addition to a subject composition, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

[00101] Powders and sprays may contain, in addition to a subject composition, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays

may additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

[00102] Compositions and compounds of the present invention may alternatively be administered by aerosol. This is accomplished by preparing an aqueous aerosol, liposomal preparation or solid particles containing the compound. A non-aqueous (e.g., fluorocarbon propellant) suspension could be used. Sonic nebulizers may be used because they minimize exposing the agent to shear, which may result in degradation of the compounds contained in the subject compositions.

[00103] Ordinarily, an aqueous aerosol is made by formulating an aqueous solution or suspension of a subject composition together with conventional pharmaceutically acceptable carriers and stabilizers. The carriers and stabilizers vary with the requirements of the particular subject composition, but typically include non-ionic surfactants (Tweens, Pluronics, or polyethylene glycol), innocuous proteins like serum albumin, sorbitan esters, oleic acid, lecithin, amino acids such as glycine, buffers, salts, sugars or sugar alcohols. Aerosols generally are prepared from isotonic solutions.

[00104] Pharmaceutical compositions of this invention suitable for parenteral administration comprise a subject composition in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or non-aqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

[00105] Examples of suitable aqueous and non-aqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate and cyclodextrins. Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. The efficacy of treatment with the subject compositions may be determined in a number of fashions known to those of skill in the art.

[00106] Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where processes are described as having, including, or comprising specific process steps, the processes also consist essentially of, or consist of, the recited processing steps. Except where indicated otherwise, the order of steps or order for performing certain actions are immaterial so long as the invention remains operable. Moreover, unless otherwise noted, two or more steps or actions may be conducted simultaneously.

EXAMPLES

[00107] The disclosure is further illustrated by the following examples. The following examples are provided for illustration purposes only, and are not to be construed as limiting the scope or content of the disclosure in any way.

Example 1: Liver Fibrosis Study

[00108] Liver fibrosis was induced in Male C57bl6 mice (Charles River, l'Arbresle, France) by injection of carbon tetrachloride (CCl₄). The C57bl6 mice weighed 22-25 g and were maintained in laboratory conditions for 1 week before experiment. Animals were housed 5 per cage with food and water available *ad libitum*.

[00109] The study consisted of 5 test groups (see Table 1): control gavage (CG), control intraperitoneal (CIP), carbon tetrachloride (CCl₄), compound 34 and JWH-133 agent (see Julien *et al.*, (2005) *Gastroenterology* 128: 742-755 (for JWH-133, a CB2-specific agonist and anti-fibrogenic agent)).

TABLE 1

Test Group	Treatment	Dose Schedule
control oral gavage (5 mice)	Carboxymethylcellulose solution (oral gavage vehicle) and olive oil (CCl ₄ vehicle)	Daily for >5 weeks
control intraperitoneal (5 mice)	DMSO/Tween (JWH-133 vehicle) and olive oil (CCl ₄ vehicle)	Daily for >5 weeks
carbon tetrachloride (10 mice)	CCl ₄ in olive oil via intraperitoneal administration (200 uL)	3 times a week (every two days) for >5 weeks
compound 34 (10 mice)	30 mM solution of compound 34 in 0.5% CMC administered via oral gavage (200 uL) and CCl ₄ in olive oil via intraperitoneal administration (200 uL)	34: Daily for >5 weeks CCl ₄ : 3 times a week for >5 weeks
JWH-133 agent (10 mice)	3 mg/kg of JWH-133 in DMSO/Tween via intraperitoneal administration (100 uL) and CCl ₄ in olive oil via intraperitoneal administration (200 uL)	JWH: Daily for >5 weeks CCl ₄ : 3 times a week for >5 weeks

[00110] The effectiveness of compound 34 (specific ligand of PPAR γ) as compared to JWH-133 (anti-fibrogenic agent) was evaluated. Compound 34 was administered orally in combination with CCl₄ injections for ca. 5 weeks. Compound 34 was dosed daily at 200 uL of a 30 mM solution. JWH-133 was administered daily via intraperitoneal injections at 3 mg/kg (dose volume = 100 uL). The model of liver fibrosis is induced by chronic exposure to increased dosages of CCl₄ for 5 weeks. The first week, mice received an intraperitoneal injection of CCl₄, resuspended in olive oil, at the dosage of 80 μ L/kg of mice body weight twice a week. At week 2, 3 injections of CCl₄ at the dosage of 160 μ L/kg were performed every two days. At week 3, 3 injections of CCl₄ at the dosage of 240 μ L/kg were performed every two days. At week 4, 3 injections of CCl₄ at the dosage of 320 μ L/kg were performed every two days. Mice were euthanized, at week 5, 3 days after the last CCl₄ administration.

[00111] Body weights were recorded for the duration of the experiment. As seen in FIG. 1, there was no detected difference in the body weight evolution between the different groups of mice.

[00112] Livers were excised and aliquots were snap frozen in liquid nitrogen and kept at -80°C until analyzed. A portion of each liver part was fixed in 10% formalin for histology.

Example 2: Histology Analysis of Liver Fibrosis Study

[00113] The formalin-fixed liver tissue was processed and 5 μ m thick paraffin sections were stained with hematoxylin and eosin (H&E) and Picrosirius red staining (specific coloration for collagen in tissue sections) to evaluate the degree of fibrosis in the liver. Each section was analyzed blindly by 3 different experimenters for the quantification of the degree of fibrosis for each lobe of the liver (5) using the Metavir score (see FIG 2A for Metavir Score chart). The photographs seen in FIG. 2B were obtained with a 20x objective. The control group had a F0 score and the CCl₄ group had a F3 score, indicating fibrosis of the liver. Compound 34 (GED) showed a greater reduction in fibrosis than the JWH-133 group with respective scores of F1 for Compound 34 and F2 for JWH-133.

[00114] As expected, no fibrosis was observed in the two groups of control mice (Control gavage (CG) and Control IP (CIP)). In the group of fibrotic mice (CCl₄ group) receiving only vehicle, a mean score 2.5 \pm 0.5 was observed corresponding to moderate to severe fibrosis according to the Metavir score (see FIG. 3A). For fibrotic

mice receiving JWH-133 or compound 34 (GED), a score of 2 ± 0 and 2.1 ± 0.2 was observed, respectively. A decrease in the degree of fibrosis from severe to moderate fibrosis was observed with similar efficacy for both JWH-133 and GED.

[00115] These results were confirmed using an adapted Metavir score (FIG. 3B): each liver section was analyzed blindly by 2 investigators for the quantification at low magnification (X5) of > 30 portal tracts classified from F0 to F4 according to the METAVIR score (Score (of 1 section) = Σ (PTF1 + PTF2 + PTF3 + PTF4)). Using this score it was demonstrated that JWH and GED decrease respectively by 14.2% and 12.5% the fibrotic score compared to untreated animals and that the therapeutic effect of the 2 tested molecules seems similar.

Example 3: Gene Expression Analysis of Liver Fibrosis Study

[00116] The effect of administration of compound 34 (GED) and the specific CB2 agonist (JWH-133) was evaluated on the level of gene expression of cytokines (IL-1 β , TNF- α) and genes involved in the mechanism of fibrosis by quantitative Real Time PCR.

[00117] Total RNA was isolated from livers using Rneasy kit (Macherey Nagel, Hoerd, France) according to the manufacturer's instructions. RNA quantification was performed using spectrophotometry. After treatment at 37°C for 30 min with 20-50 units of RNase-free DNase I (Roche Diagnostics Corporation, Indianapolis, IN, USA), oligo-dT primers (Roche Diagnostics Corporation, Indianapolis, USA) were used to synthesize single-stranded cDNA. mRNAs were quantified using SYBR green Master Mix (Applera, Courtaboeuf, France) with murine specific oligonucleotides (see Table 2) in a GeneAmp Abiprism 7000 (Applera, Courtaboeuf, France). In each assay, calibrated and no-template controls were included. Each sample was run in triplicate. SYBR green dye intensity was analyzed using the Abiprism 7000 SDS software (Applera, Courtaboeuf, France). All results were normalized to the unaffected housekeeping gene β -actin.

[00118]

TABLE 2

Murine Genes	Primer Sequences (5' \rightarrow 3')
β -actin	F: 5'-gggTCagAAggATTCCTATg-3' (SEQ ID NO: 1) R: 5'-ggTCTCAAACATgATCTggg-3' (SEQ ID NO: 2)
TNF- α	F: 5'-TgggAgTAgACAaggTACAACCC-3' (SEQ ID NO: 3) R: 5' CATCTTTCTCAAATTCgAgTgACAA-3' (SEQ ID NO: 4)
IL-1 β	F: 5'-gATCCACACTCTCCAgCTgCA-3' (SEQ ID NO: 5) R: 5'-CAACCAACAAGTgATATTCTCCATg-3' (SEQ ID NO: 6)
Collagen-1	F: 5'-GAG TAC TGG ATC GAC CCT AAC CAA -3' (SEQ ID NO: 7) R: 5'-ACA CAG GTC TGA CCT GTC TCC AT -3' (SEQ ID NO: 8)
α -SMA	F: 5'-CCT GAC GGG CAG GTG ATC -3' (SEQ ID NO: 9) R: 5'-ATG AAA GAT GGC TGG AAG AGA GTC T -3' (SEQ ID NO: 10)

[00119] There was no significant modification of cytokine gene expression in the different groups of mice. With regards to the genes involved in the mechanism of fibrosis, there was an observed increase of expression of SMA and collagen-1 in animals with fibrosis (CCl₄ group) compared to control animals (CG and CIP groups). As seen in FIG. 4 and Table 3, there was a decrease in the level of α -SMA and collagen-1 in the GED and JWH-133

groups of mice as compared to the CCl₄ group (fibrosis control). For the level of collagen-1, the decrease was significant for the group of mice treated by compound 34 compared to the CCl₄ mice with respectively, 42.81±14.81, p=0.03 vs. 75.15±5.23. This biological data along with the histological improvement observed in mice treated with compound 34 suggests that compound 34 may have therapeutic anti-fibrotic properties.

[00120]

TABLE 3

Group	Collagen-1	α -SMA
CCl ₄	138.12±34.68	65.51±22.24
JWH-133 + CCl ₄	87.59±16.97	30.07±4.7
Compound 34 + CCl ₄	75.15±5.23, p=0.03	42.81±14.81

Example 4: Liver Blood Enzyme Analysis of Liver Fibrosis Study

[00121] The effect of oral administration of compound 34 (GED) and the specific CB2 agonist (JWH-133) was evaluated on different liver biochemical parameters in the serum of mice: hepatic enzymes (alanine aminotransferase (AST) and aspartate aminotransferase (ALT)), Gamma GT and the Alkaline phosphatase.

[00122] As expected, 4 days after the last administration of CCl₄, no enhancement of AST, ALT and Alk P was observed in any groups of mice. Compound 34 was not associated with any enhancement of blood liver enzymes suggesting an absence of liver toxicity.

[00123] As evaluated by histology and liver markers, JWH-133 and compound 34 have similar anti-fibrotic effects in mice with liver fibrosis induced by repeated injections of CCl₄. As expected, JWH decreases hepatic fibrotic lesions by 14% and normalizes the liver concentration of collagen-1 mRNA. Compound 34 has a similar effect without liver toxicity. The data suggests that compound 34 may be the first compound with both anti-inflammatory and anti-fibrotic properties.

Example 5: Intestinal Fibrosis Study

[00124] This example describes an experiment to evaluate the effect of compound 34 on intestinal fibrosis. Chronic colitis was induced in C57bl6 mice by 3 cycles of 2.5% DSS (40,000–50,000 MW, TdB consultancy AB, Sweden) dissolved in sterile drinking water for 5 days followed by 7 days of regular drinking water.

[00125] The study consisted of 3 test groups (see Table 4): control group, DSS control group, GED (compound 34) group.

TABLE 4

Test Group	Treatment	Dose Schedule
control (10 mice)	H ₂ O	Daily for 36 days
DSS control (25 mice)	2.5% DSS in drinking H ₂ O	5 days on, 7 days off for 36 days
compound 34 (25 mice)	30 mM solution of compound 34 in 0.5% CMC administered via oral gavage (200 μ L) and 2.5% DSS in drinking H ₂ O	cmpd. 34: daily from day 13-36 DSS: 5 days on, 7 days off for 36 days

[00126] Compound 34 was used at its optimal dosage (30 mM) and administered by oral gavage daily after the second cycle of DSS and until euthanasia. Mice receiving DSS were checked twice a week for development of colitis by monitoring body weight. Euthanasia of mice was performed one week after the third cycle of DSS (see FIG. 5). Evaluation of the size and weight of the colon was evaluated *post mortem*.

Example 6: Analysis of Intestinal Fibrosis Study

A. Colon ratio weight/size

[00127] Intestinal fibrosis is characterized by a shortening and thickening of the colon and adhesences. The measure of the ratio weight/size of the colon is an indicator of the level of inflammation and fibrosis. As seen in FIG. 6A, a significant increase of 124% in the ratio of colon weight/size was observed in the DSS mice receiving the vehicle compared to control mice receiving only regular water, with respectively 44.98 ± 6.31 vs 20.11 ± 3.91 , $p < 0.05$. These results are illustrated by an important shortening of the colon in DSS mice (FIG 6B). Compound 34 causes a 34% reduction in the profibrotic effect of DSS leading to a significant decrease ratio weight/size of the colon of 38.13 ± 7.82 vs 44.98 ± 6.31 ($p < 0.05$) compared to DSS mice receiving the vehicle (FIG. 6A). Accordingly, it reduces the morphological signs of fibrosis such as the shortening and the thickening of the colon and it decreases the collagen deposit in the colon (FIG. 6B).

B. Inflammation level estimated by histology

[00128] Rings of the transverse part of the colon were sampled (*post mortem*) and fixed in 4% formaldehyde and embedded in paraffin for histological analysis. Sections (4 μ m) were stained with May-Grunwald-Giemsa and a multiparametric histological scoring (0 to 18) was performed blindly by two investigators. This staining allowed the quantification of inflammation. Histological grades, reflecting the inflammation level, were assigned based on the intensity of cellular infiltrate in the mucosa, its extension in sub-mucosa layers and the presence of epithelial lesions (see Table 5).

TABLE 5

Segment	Grade	Description
Severity	0	None
	1	Slight
	2	Moderate
	3	Severe
Extent	0	None
	1	Mucosa
	2	Mucosa and submucosa
	3	Transmural
Regeneration	4	No Tissue Repair
	3	Surface epithelium not intact
	2	Regeneration with crypt depletion
	1	Almost complete regeneration
	0	Complete regeneration or normal tissue
Crypt damage	0	None
	1	Basal 1/3 damaged
	2	Basal 2/3 damaged

	3	Only surface epithelium intact
	4	Entire crypt and epithelium lost
Percentage involvement	1	1-25%
	2	26-50%
	3	51-75%
	4	76-100%

[00129] The inflammatory status was evaluated at the histological level for each colon of mice using a classical MGG staining. As expected, significant and strong inflammation was observed in the group of mice receiving the DSS compared to control mice receiving regular water, with respectively 3.77 ± 0.80 vs. 0.50 ± 0.33 , $p < 0.05$ (see FIG 7). This result confirms the induction of inflammation by the 3 cycles of DSS. A significant decrease in the inflammation level was observed in DSS mice receiving compound 34 as compared to DSS mice receiving only the vehicle with respectively, 2.18 ± 0.48 vs. 3.77 ± 0.80 , $p < 0.05$.

C. Level of Fibrosis

C1. Evaluation by Picrosirius Red staining

[00130] Samples of colon obtained were promptly fixed with 10% buffered formalin in PBS (pH 7.4) for 3 h, dehydrated in graded ethanol and embedded in low-temperature-fusion paraffin. Formalin-fixed colonic tissue was processed and 5 μ m thick paraffin sections were stained with Picrosirius red staining for histological analysis. Pictures were obtained with a x20 objective. Each section was analyzed blindly by 3 different experimenters for the quantification of the degree of fibrosis using a score from 0 to 4.

[00131] A significant increase of the level of fibrosis was observed in the colon of DSS mice compared to control mice with respectively 2.27 ± 0.12 vs 1.12 ± 0.07 (see FIG. 8). This result confirms that chronic inflammation induced by 3 cycles of DSS induces intestinal fibrosis. It also confirms that administration of compound 34 to DSS mice decreases the level of fibrosis compared to DSS mice receiving only the vehicle (1.625 ± 0.15 vs. 2.27 ± 0.12 , $p < 0.05$).

C2. Evaluation by Trichrome Masson's staining

[00132] A trichrome masson's and collagen staining were used in this evaluation of the level of colonic fibrosis. Samples of colon obtained were promptly fixed with 10% buffered formalin in PBS (pH 7.4) for 3 h, dehydrated in graded ethanol and embedded in low-temperature-fusion paraffin. Serial 3 μ m sections were incubated for 40 min in methanol and 3% hydrogen peroxide solution and then rinsed in PBS. Intestinal fibrosis was scored as absent, mild, or severe, depending on the density and extent of trichrome-positive connective tissue staining and disruption of tissue architecture.

C3. Evaluation of collagen levels

[00133] For the immunohistochemical analysis, tissue specimens from the colon were fixed in fresh 4% paraformaldehyde (PFA)/ PBS solution for 3h at room temperature, dehydrated in a graded ethanol series, and embedded in low-temperature-fusion paraffin. 3- μ m-thick sections were incubated in methanol for 40 min and then in 3% hydrogen peroxide for 5 min. Samples were incubated overnight with specific antibodies against collagen types I-III (Abcam), connective tissue growth factor (CTGF) (Abcam), Platelet Derived Growth factor (PDGF), SMAD2/3, the main markers of fibrosis. The samples were washed for 5 min with PBS and incubated with streptavidin-biotin-peroxidase conjugated secondary antibody (Dako LSAB Corporation, cod K0675, Dako-

Cytomation, Milano). After a wash in PBS for 10 min the sections were subjected to incubation with 3,3'-diaminobenzidine-tetrahydrochloride for 1-3 min.

[00134] The specificity of immune reaction was performed omitting the primary antibodies. Finally the samples were stained with hematoxylin of Mayer and observed under the Olympus BX51 Light Microscope (Olympus, Optical Co. Ltd., Tokyo, Japan).

[00135] In the immunohistochemical analysis (x 20) of collagens I-VII in colon from DSS and control mice, the staining of collagens I-VII in large intestine of DSS mice was localized mainly within connective tissue of submucosa and muscularis propria, where strong staining of collagen was observed (see FIG. 9). As shown in FIG. 9, administration of compound 34 is associated with a decrease in the collagen staining indicating that it has improved the fibrosis and reduces the development of intestinal fibrosis.

[00136] In addition to anti-inflammatory effects, administration of compound 34 decreases rapidly molecular events leading to fibrosis and prevents the intestinal fibrotic lesions induced by DSS in mice.

[00137] For the Western blot analysis, 0.5 cm of frozen colonic samples were homogenized in RIPA Buffer containing 50 mM Tris HCl pH 7.6, 150 mM NaCl, 1.5 mM MgCl₂, 5mM EDTA, 1% Triton-X and 10% Glycerol, supplemented with 100 mM Sodium Fluoride (NaF), 2 mM Sodium Orthovanadate (Na₃VO₄), 10 mM Sodium Pyrophosphate (NaPPi), 1 mM phenylmethanesulphonyl fluoride (PMSF) and a classical protease-inhibitor cocktail containing 10 µg/mL Leupeptin and Aprotinin.

[00138] For each sample, 30 µg of protein was separated by PAGE and electroblotted. Nitrocellulose membranes (100% pure) were incubated (according with a specific protocol for each antibody) with primary antibodies directed against CTGF, Collagen-I and GAPDH (purchased from Abcam, Cambridge, UK; 1:1000 for 2 h at RT), diluted in 5% non-fat milk in 0.1% TBS-t. Membranes were subsequently washed in TBS-t 0.1% and incubated with secondary horseradish peroxidase-conjugated antibodies (anti-Rabbit and anti-Mouse; Sigma Aldrich; 1:20000 for 1 h at RT) diluted in 5% non-fat milk in 0.1% TBS-t. Finally, immunodetection was performed with SuperSignal West Pico chemiluminescent substrate (Thermo Scientific Pierce, Erembodegem) according to manufacturer's protocol. Membranes were exposed to autoradiography film (Fuji Photo Film Co., Dusseldorf, Germany). Optical density of target bands was determined using a computer-assisted densitometer and the ImageJ public domain software (W. S., Rasband, ImageJ, U. S. National Institutes of Health, Bethesda, MD; <http://rsb.info.nih.gov/ij/>, 1997–2011). Tissue levels were expressed as units of Optical Density (OD) per quantity of total proteins, normalized with internal control GAPDH and the results were expressed as percentage of control groups.

[00139] Using the IHC methods and specific antibodies, a reduction of the main markers of fibrosis (Collagen I-III, CTGF, SMAD 2/3, PDGF, α-SMA, and TGF-β1) was observed in the colon of DSS mice having received the GED compared to colitic mice receiving the vehicle (see FIG. 9A-9E).

[00140] These results were confirmed using Western blot method and the quantification of different fibrosis markers was performed on the whole colon samples (see FIG. 10). As seen in Table 6, a significant decrease of the expression of collagen I, CTGF and IL-13 was observed in the colitic mice treated by GED compared to colitic mice receiving only the vehicle (Collagen I 90.75±19.91 vs. 284.4±63.86, p=0.012, for CTG 55.75±17.53 vs. 115.1±16.88, p=0.029 and IL-13 88.88±14.01 vs. 189.5±28.71, p=0.007).

TABLE 6

	DSS	DSS +GED	p
α-SMA	125.4±19.45	97.88±15.66	0.282

Collagen I	284.4±63.86	90.75±19.91	0.012
CTGF	115.1±16.88	55.75±17.53	0.029
IL-13	189.5±28.71	88.88±14.01	0.007
TGF-β1	153.6±26.25	130.5±28.02	0.5

Example 7: Intestinal Fibrosis Study

[00141] This example describes an experiment to evaluate the effect of compound 34 (GED) in treating intestinal fibrosis.

MATERIALS AND METHODS***In vitro* experiments**

[00142] A total of 60 wild type C57BL/6 mice, purchased from Janvier (Le Genest-St-Isle, France) were included in the study. All mice were maintained in a specific pathogen-free facility of the Pasteur Institute in Lille (France). Animal experiments were performed according to the governmental guidelines N° 68/609/CEE.

Induction of chronic colitis

[00143] Chronic colitis and fibrosis were induced in mice by oral administration of 2.5% (w/v) DSS in drinking water, for three cycles (5 day DSS, 7 days of water). Animals were monitored daily for food and fluid intake and were weighed at the beginning of the study and thereafter regularly every three days.

Experimental design

[00144] The mice were randomly divided into two groups: i. DSS and ii. DSS + GED. Each group consisted of 25 mice and was compared with 10 control animals, which received only water. GED (30mg/kg/mouse) was dissolved in a solution containing 0.5% Carboxymethylcellulose sodium salt (CMC; MW: 90,000 Da; Sigma Aldrich) and 1% Tween 80 and was administrated daily by oral gavage (100µl/mouse). GED was administered at the beginning of second cycle.

Assessment of colitis course

[00145] Animals were observed daily for fluid intake, weight changes, and examined for signs of colitis including weight loss, diarrhoea, rectal bleeding and prolapse as well as signs of systemic inflammation such as piloerection, lethargy, and periorbital exudates.

Samples recovery and preparation

[00146] Finally, seven days after the last DSS cycle administration, the animals of each group were sacrificed by cervical dislocation under deep CO₂ anaesthesia and undergone to laparotomy. The colon were visualized and rapidly excised. The length of the whole colon- rectum and the weight of the distal 8 cm of colon-rectum were measured.

Assessment of macroscopic and microscopic colonic lesions

[00147] The macroscopic colonic lesions including dilation, thickness and stenosis were scored by an independent observer, unaware of the mice treatment. Tissue specimens from the colon were dissected and immediately frozen or fixed in fresh 4% formaldehyde (FA)/ PBS solution for 3h at room temperature followed by the standard procedure for paraffin embedding. Paraffin embedded tissues transversally sectioned at 3 µm were subject to Hematoxylin/Eosin staining, to assess the inflammatory degree, and Masson trichrome to better assess connective tissue and fibrosis. The stained sections were then blindly observed under an Olympus BX51 Light Microscope (Olympus, Optical Co. Ltd., Tokyo, Japan) by two pathologists which have performed the histological score evaluating: i. presence of ulcers (0=absent, 1= little ulcers, 2= big ulcers); ii. the inflammatory degree (0=absent 1= mild, 2= moderate and 3=severe); iii. depth of lesions (0=absent, 1= lesions extending in

the submucosa, 2= lesions in the muscularispropria and 3= lesions in the serosa); iv. degree of fibrosis (0=absent 1= mild, 2= moderate and 3=severe). The degree of intestinal inflammation was assessed according to the density and extent of chronic inflammatory infiltrate, loss of goblet cells, and bowel wall thickening. A total microscopic score will be obtained as sum of all scores (maximum score possible =10).

[00148] Intestinal fibrosis was scored as absent, mild, or severe, depending on the density and extent of trichrome-positive connective tissue staining and disruption of tissue architecture.

Immunohistochemistry

[00149] Tissue specimens from the colon, dissected and fixed in fresh 4% formaldehyde (FA)/ PBS solution for 3 hours at room temperature, were dehydrated in a graded ethanol series, and embedded in low-temperature-fusion paraffin. 3- μ m-thick sections were incubated in a solution of methanol and 3% hydrogen peroxide for 45 min.

[00150] Samples were incubated overnight with specific antibodies against α -smooth muscle actin (α -SMA, Abcam), collagen types I-III (Abcam), TGF- β 1 (Abcam), connective tissue growth factor (CTGF) (Abcam), pSmad3 and Smad3 (Cell Signaling), Smad7 (Imgenex) PPAR γ (Cell Signaling). The samples were washed 5 min with PBS and incubated with streptavidin-biotin-peroxidase conjugated secondary antibody (Dako LSAB Corporation, cod K0675, Dako-Cytomation, Milano). After one wash in PBS for 10 min the sections are subject to incubation with 3,3-diaminobenzidine-tetrahydrochloride for 1-3 min. The counterstaining of nuclei was obtained by Hematoxilin Eosin. The specificity of immune reaction was effected keeping off the primary antibodies. Finally the samples were stained with hematoxylin of Mayer and observed under the Olympus BX51 Light Microscope (Olympus, Optical Co. Ltd., Tokyo, Japan).

Western blot analysis

[00151] 0.5 cm of frozen colonic samples were cut and mechanically homogenized in RIPA Buffer containing 50 mMTrisHCl pH 7.6, 150 mMNaCl, 1.5 mM MgCl₂, 5mM EDTA, 1% Triton-X and 10% Glycerol, supplemented with 100 mM Sodium Fluoride (NaF), 2 mM Sodium Orthovanadate (Na₃VO₄), 10 mM Sodium Pyrophosphate (NaPPi), 1 mMphenylmethanesulphone (PMSF) and a classical protease-inhibitor cocktail containing 10 μ g/mL Leupeptin and Aprotinin.

[00152] 30 μ g of protein for each sample were separated by PAGE and electroblotted. 100% pure Nitrocellulose membranes were incubated (according with a specific protocol for each antibody) with primary antibodies directed against TGF- β 1, CTGF, α -SMA, Collagen-I and GAPDH (purchased from Abcam, Cambridge, UK; 1:1000 for 2 h at RT), and IL-13 (purchased from Antibodies online; 1:1000 for 2h at RT) diluted in 5% non fat-milk in 0,1% TBS-t. Membranes were subsequently washed in TBS-t 0.1% and incubated with secondary horseradish peroxidase-conjugated antibodies (anti-Rabbit and anti-Mouse; Sigma Aldrich; 1:20000 for 1 h at RT) diluted in 5% non fat-milk in 0,1% TBS-t. Finally, immunodetection was performed with SuperSignal West Pico chemoluminescent substrate (Thermo Scientific Pierce, Erembodegem) according to manufacturer's protocol. Membranes were exposed to autoradiography film (Fuji Photo Film Co., Dusseldorf, Germany). Optical density of target bands was determined using a computer-assisted densitometer and the ImageJ public domain software (W. S., Rasband, ImageJ, U. S. National Institutes of Health, Bethesda, MD; <http://rsb.info.nih.gov/ij/>, 1997–2011). Tissue levels were expressed as units of Optical Density (OD) per quantity of total proteins, normalizing with internal control GAPDH and the results were expressed as percentage of control groups.

In vivo experiments

Cell cultures

[00153] Human colon cancer cell line HT-29 (ATCC HTB-38) and colonic intestinal fibroblasts (ATCC human CDD-18 co) were grown in Dulbecco's modified Eagle's medium (DMEM) and Alpha-Modified Eagle's medium, respectively, 100 U/ml penicillin and 100 µg/ml streptomycin supplemented with 10% fetal bovine serum (FBS). Cell cultures were maintained in a humidified atmosphere of 95% air and 5% CO₂ at 37°C.

PPAR γ knock down cells generation

[00154] PPAR γ knock down HT29 were obtained using the pSUPER.retro system (OligoEngine). Forward and reverse target sequences corresponding to nucleotides 105-123 of the human PPAR γ mRNA (5'-GCCCTTCACTACTGTTGAC-3' (SEQ ID NO: 11)) were cloned into the BglII/XhoI restriction sites of the pSUPERretro vector (pRS) giving the ShPPAR construct. A negative control pRS plasmid containing the sequence 5'-ACGCTGAGTACTTCGAAAT-3' (SEQ ID NO: 12) targeted against the luciferase gene was also generated (ShLuc construct). Both constructions were transfected in HT-29 and Caco-2 cells using Nucleofector technology from AmaxaBiosystems, according to the manufacturer's protocol. Stably transfected clones were selected 24h post-transfection with complete culture medium supplemented with puromycin (5µg/ml). The silencing of PPAR γ expression was checked by quantitative RT-PCR and western-blot analysis. Once established, ShPPAR and ShLuc cell lines were maintained in complete medium supplemented with 2.5µg/ml puromycin.

Experimental design

[00155] Fibrotic phenotype was induced by stimulating hCCD-18 and HT29 with TGF- β 1ng/mL and 10 ng/mL for 4 days, respectively, dissolved in serum-free medium. 1mM for GED (195.22 g/mol) was administrated during the differentiation period. In addition, the potential dependence on PPAR γ activation has been investigated by using GW9662, a specific antagonist of this receptor. 10⁻⁵ M GW9662 has been administrated 4h before to start the TGF β -induced differentiation.

Quantitative RT-PCR

[00156] Total RNA was extracted with a Nucleospin RNA kit (Macherey-Nagel, Hoerd, France). After RNase inactivation, the total RNA was cleaned of trace genomic DNA via a DNase treatment and eluted in RNase-free, DEPC-free water. The purity of the RNA was evaluated by UV spectroscopy on a Nanodrop system from 220 to 350 nm and profiling on an Agilent 2100 bioanalyzer. One µg of total RNA was used to perform a Quantitative RT-PCR by using LightCyclerFastStart DNA Master SYBR Green I from Roche Diagnostics (Indianapolis, IN) according to the manufacturer's protocol.

RESULTS

In vivo experiments

Clinical and macroscopic findings on chronic colitis in mice

[00157] Chronic oral administration of DSS induced loss of body weight in all chronic mice starting from the last day of first DSS cycle (day 5). At day 10 the mice receiving DSS showed a lower body weight compared to the control mice with respectively 23.88 ± 0.31 vs. 25.27 ± 0.35 p<0,05 (Fig. 11)

[00158] Moderate to severe diarrhea were observed in 70% of DSS treated mice and 50% of GED mice on the seventh day of DSS oral administration. In the same period systemic manifestations, such as piloerection, periorbital exudates and/or lethargy, were observed in 40% of DSS mice and 30% of GED mice. At no stage frank rectal bleeding or rectal prolapse were observed. The mice were sacrificed on day 15 and day 30, respectively, and the colon was subjected to macroscopic evaluation by measuring of colon weight (referred to the last 8 cm of the distal large bowel) and colon length and by observing the presence of adhesions, strictures, dilatation, thickness that was

scored as 0 if absent; 1 if mild or moderate; and 2 if severe. The total macroscopic score was calculated as sum of all these score.

The effects of GED administration on course and macroscopic findings of DSS-induced chronic colitis in mice are summarized in Table 7, indicating that GED ameliorates macroscopic lesions in and chronic DSS-induced colitis and reduce the morphological signs of fibrosis.

[00159] All the macroscopic features observed in DSS treatments showed, in both models, significant variation that indicated an effective ability of DSS to induce a severe tissue damage. All macroscopic lesions were ameliorated by the GED treatment determining significant differences between DSS and DSS plus GED total macroscopic scores. Colon weight and colon weight/body weight ratio showed no significant variation following the GED treatment, while colon length appeared significantly restored. Direct observation of colons showed, indeed, an evident shortening and dilatation in majority of DSS-treated samples and a remission of these features on GED treated mice.

TABLE 7. Effect of GED Administration on Course and Macroscopic Findings of Chronic colitis

Parameters	CHRONIC		
	H ₂ O	DSS	DSS + GED
Duration of DSS treatment	30 days	30 days	15 days
Mortality (n. mice)	0	2	0
Body weight	28,95 ± 0,331	25,93 ± 0,223**	25,132 ± 0,458**
Colon weight	0,171 ± 0,005	0,283 ± 0,006***	0,269 ± 0,007***
Colon weight/body weight ratio	0,29 ± 0,0051	0,26 ± 0,007**	0,27 ± 0,012**/¥¥
Colon length	8,52 ± 0,093	6,304 ± 0,09**	7,092 ± 0,16**/
Dilation	np	1,35 ± 0,165**	1,12 ± 0,185**/¥¥
Thickness	np	1 ± 0,154**	0,48 ± 0,108**/¥
Stenosis	np	0,364 ± 0,151**	0,32 ± 0,114**/¥
Adhesions	np	0,348 ± 0,135**	0,12 ± 0,1¥¥
Total macroscopic score	np	3,112 ± 0,31 **	2,02 ± 0,26** /¥¥

Data are expressed as mean ± SEM; np = not present; * = p < 0.05 vs H₂O; ** = p < 0.01 vs H₂O; *** = p < 0.005 vs H₂O; ¥ = p < 0.05 vs DSS; ¥¥ = p < 0.01 vs DSS; ¥¥¥ = p < 0.005 vs DSS

[00160] Thus, the ratio weight/size of the colon was used as further indicator of the level of inflammation and fibrosis. A significant increase of 124% in the ratio was observed in the DSS-receiving mice compared to control mice receiving only regular water, with respectively 44.98 ± 6.31 vs. 20.11 ± 3.91 (p < 0.05). GED led to a significant decrease of ratio weight/size of the colon compared to DSS mice (38.13 ± 7.82 vs. 44.98 ± 6.31 (p < 0.05)) causing a 26% reduction respect to the profibrotic effect of DSS. Accordingly, GED is able to reduce the morphological signs of fibrosis such as shortening and thickening of the colon.

Microscopic findings on chronic colitis

[00161] Several histological parameters were scored on the bases of intensity of cellular infiltrate in the mucosa, inflammation extension in sub-mucosa layers and the presence of epithelial lesions and collagen deposition. A significant and strong inflammation was observed in the group of mice receiving the DSS compared to control mice receiving regular water. The histological features in both models showed mucosal ulceration and degeneration, decrease of goblet cells, wide inflammatory cellular infiltration, and submucosal edema. All these findings appeared more emphasized in DSS-induced chronic colitis and a wide thickening of the colon wall was evident. In addition, the chronic model showed an evident regeneration of crypts and a possible spontaneous recovering of inflammatory degree in the long-term treatments (Fig. 12).

[00162] The level of fibrosis was observed by Masson's trichrome staining, specific for collagen deposition. A significant increase of the level of fibrosis was observed not only in the colon of DSS mice compared to control mice but also between the two treatments (Fig. 12). This evidence, together with the observation of a lower inflammatory degree, seems consistent with the postulate that fibrosis begins from an inflammatory response but, subsequently, may progress independently from it.

[00163] Total microscopic scores were assigned and demonstrated that high inflammatory degree and levels of fibrosis, induced by 1 and 3 cycles of DSS, decreased significantly following the daily oral administration of GED (3.36 ± 0.55 for GED vs. 6.45 ± 0.82 for DSS, $p < 0.01$) (Fig.12).

Evaluation of tissue levels of main fibrosis markers

[00164] The tissue levels of main markers of fibrosis in the chronic model of DSS-induced colitis were evaluated by immunohistochemical analysis and confirmed by immunoblotting. Thus, a marked reduction of Collagen I-III (Fig. 13a) and α -SMA (Fig. 13b), the specific markers of fibroblasts differentiation, in colons of mice subjected to the oral administration of GED treatment, compared to mice receiving only DSS, was observed. The expression levels of Collagen I-III and α -SMA were measured on 6 control mice, 8 DSS mice and 8 DSS + GED mice. Both immunochemistry and western blotting showed that all of main fibrosis markers were significantly increased in the group of mice receiving the DSS compared to control mice receiving regular water, and that daily administration of GED is associated with their reduced expression. The expression levels of each protein were expressed as percentage of the mean of control group and each marker of interest was normalized with a housekeeping protein like GAPDH. A marked increase of collagen I-III expression (184%) in DSS mice compared with the control group, as well as moderated upregulation of 25% for α -SMA, was observed. The oral administration of GED in mice with DSS-induced chronic colitis caused a reduction in expression levels of these proteins, more significantly marked for Collagen I-III ($90.75 \% \pm 19.9 \%$ for GED vs. $284.4 \% \pm 63.8 \%$ for DSS, $p < 0.005$).

GED controls the TGF β /Smad pathway and its specific activator, IL13.

[00165] The ability of GED to control the TGF β /Smad pathway at different levels was observed by immunohistochemical analysis and demonstrated a marked reduction of different members involved in this pathway, such as TGF- β 1, Smad3, CTGF (data not shown).

[00166] The benefit of GED in the prevention or treatment of intestinal fibrosis relating to variations in IL-13 expression levels was observed Fig. 14a). IL-13 is a specific activator of the TGF β /Smad pathway and this aspect was confirmed by consequent decrease in TGF- β 1 (Fig. 14b) and CTGF (Fig. 14c) expression levels. The increased expression of IL-13 in DSS mice was significantly restored by oral administration of GED with $88.88 \% \pm 14.01 \%$ for GED vs. $189.5 \% \pm 28.71\%$ for DSS, $p < 0.01$. The consequent reduction of expression levels of TGF- β 1 and CTGF were confirmed by immunoblotting analysis, with marked significance for CTGF ($55.57 \% \pm 17.53\%$ for

GED vs. 115.1% \pm 16.88% for DSS, $p < 0.05$). The expression levels were expressed as percentage of optical density in comparison to control group and normalized with GAPDH.

INCORPORATION BY REFERENCE


[00167] The entire disclosure of each of the patent documents and scientific articles referred to herein is incorporated by reference for all purposes.

EQUIVALENTS

[00168] The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting on the invention described herein. Scope of the invention is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to be embraced therein.

We claim:


1. A method of treating fibrosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound, wherein the compound is 2-methoxy-3-(4'-aminophenyl) propionic acid (compound 34)



represented by the formula: COCC(c1ccc(N)cc1)C(=O)O or a pharmaceutically acceptable salt or a stereoisomer thereof, wherein the fibrosis is selected from the group consisting of hepatic fibrosis, intestinal fibrosis, and idiopathic pulmonary fibrosis.

2. The method of claim 1, wherein the compound is (S)-2-methoxy-3-(4'-aminophenyl) propionic acid, or a pharmaceutically acceptable salt thereof.
3. The method of claim 1, wherein the compound is (R)-2-methoxy-3-(4'-aminophenyl) propionic acid, or a pharmaceutically acceptable salt thereof.
4. The method of any one of claims 1 to 3, wherein the fibrosis is hepatic fibrosis.
5. The method of any one of claims 1 to 3, wherein the fibrosis is intestinal fibrosis.
6. The method of any one of claims 1 to 3, wherein the fibrosis is idiopathic pulmonary fibrosis.
7. A method of treating fibrosis, comprising administering to a patient in need thereof, a pharmaceutical preparation comprising a therapeutically effective amount of a compound, wherein the compound is 2-methoxy-3-(4'-aminophenyl) propionic acid or a pharmaceutically acceptable salt or a stereoisomer thereof; and one or more pharmaceutically acceptable carriers, wherein the fibrosis is selected from the group consisting of hepatic fibrosis, intestinal fibrosis, and idiopathic pulmonary fibrosis.
8. The method of claim 7, wherein the compound is (S)-2-methoxy-3-(4'-aminophenyl) propionic acid or a pharmaceutically acceptable salt thereof.
9. The method of claim 7, wherein the compound is (R)-2-methoxy-3-(4'-aminophenyl) propionic acid or a pharmaceutically acceptable salt thereof.
10. The method of any one of claims 1 to 9, wherein the compound, or a pharmaceutically acceptable salt or a stereoisomer thereof, or the pharmaceutical preparation is administered orally.
11. The method of any one of claims 1 to 10, wherein the patient is human.
12. The method of any one of claims 7 to 11, wherein the fibrosis is hepatic fibrosis.
13. The method of any one of claims 7 to 11, wherein the fibrosis is intestinal fibrosis.
14. The method of any one of claims 7 to 11, wherein the fibrosis is idiopathic pulmonary fibrosis.
15. The method of any one of claims 1 to 14, wherein the patient is currently suffering or has suffered from hepatitis B or hepatitis C.
16. The method of any one of claims 1 to 15, wherein the patient has cirrhosis.
17. The method of any one of claims 1 to 16, wherein the patient is also suffering from Crohn's disease, inflammatory bowel disease or ulcerative colitis.

18. Use of a compound, wherein the compound is 2-methoxy-3-(4'-aminophenyl) propionic acid (compound 34)



represented by the formula: or a pharmaceutically acceptable salt or a stereoisomer thereof, in the manufacture of a medicament for treating fibrosis, wherein the fibrosis is selected from the group consisting of hepatic fibrosis, intestinal fibrosis, and idiopathic pulmonary fibrosis.

19. The use according to claim 18, wherein the compound is (S)-2-methoxy-3-(4'-aminophenyl) propionic acid, or a pharmaceutically acceptable salt thereof.

20. The use according to claim 18, wherein the compound is (R)-2-methoxy-3-(4'-aminophenyl) propionic acid, or a pharmaceutically acceptable salt thereof.

FIG. 1




FIG. 2A

Metavir score (Activity A0→A3 & Fibrosis F0→F4)		
	Activity (Necrosis & inflammation)	Fibrosis
0	A0	F0
low	A1	F1
moderate	A2	F2
severe	A3	F3
Cirrhosis		F4




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6A




FIG. 6B




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 9D




FIG. 9E




FIG. 10




Fig. 11




Fig. 12




Figure 13




Figure 14




Figure. 15




Figure. 16

