

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 August 2008 (28.08.2008)

PCT

(10) International Publication Number
WO 2008/103392 A2

(51) International Patent Classification:

A61K 39/00 (2006.01) *A61K 39/395* (2006.01)
A61K 38/00 (2006.01) *A61P 37/00* (2006.01)
C12N 5/06 (2006.01)

(21) International Application Number:

PCT/US2008/002256

(22) International Filing Date:

21 February 2008 (21.02.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/890,964 21 February 2007 (21.02.2007) US

(71) Applicants and

(72) Inventors: **DONDA, Alena** [CH/CH]; Les Dailles, CH-1673 Auboranges (CH). **MACH, Jean-pierre** [CH/CH]; 300 Route de Lausanne, CH-1293 Bellevue (CH). **STIRNEMANN, Kathrin** [CH/CH]; Puentstrasse 22, CH-8047 Zurich (CH). **ZAUDERER, Maurice** [US/US]; 44 Woodland Road, Pittsford, NY 14534 (US).

(74) Agents: **HAANES, Elizabeth, J.** et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— *without international search report and to be republished upon receipt of that report*

(54) Title: MODULATION OF NKT CELL ACTIVITY WITH ANTIGEN-LOADED CD1D MOLECULES

Figure 1

TCR downmodulation upon iNKT activation



(57) **Abstract:** The invention is directed to methods of modulating an immune response in an animal, comprising administering a composition comprising one or more soluble CD1d complexes, in particular non-specific soluble CD1d complexes. Soluble CD1d complexes comprise a soluble CD1d polypeptide, a β 2-microglobulin polypeptide, and a ceramide-like glycolipid antigen bound to the CD1d antigen binding groove, and in certain embodiments, an immunogen. The administration of compositions of the present invention affects the activity of CD1d-restricted NKT cells, and in particular, allows for multiple administrations without causing CD1d-restricted NKT cell anergy.

MODULATION OF NKT CELL ACTIVITY WITH ANTIGEN-LOADED CD1d MOLECULES

Background Of The Invention

Field of the Invention

[0001] The invention generally relates to the field of immunology.

Background Art

[0002] The natural immune system strikes a complex balance between highly aggressive, protective immune responses to foreign pathogens and the need to maintain tolerance to normal tissues. In recent years there has been increasing recognition that interactions among many different cell types contribute to maintaining this balance. Such interactions can, for example, result in polarized responses with either production of pro-inflammatory cytokines (*e.g.*, interferon- γ) by TH1 type T cells or production of interleukin-4 (IL-4) by TH2 type T cells that suppress TH1 activity. In a number of different animal models, T cell polarization to TH1 has been shown to favor protective immunity to tumors or infectious pathogens whereas T cell polarization to TH2 can be a critical factor in preventing development of cell-mediated autoimmune disease. The conditions that determine whether immune stimulation will result in aggressive cell-mediated immunity or in down regulation of such responses are highly localized in the sense that each tissue is comprised of a distinctive set of antigen presenting cells (APC) and lymphocyte lineages that interact to favor different immune responses. For example, under optimal conditions, the dendritic cells (DC) localized in a normal tissue may represent predominantly a lineage and stage of maturation that favors tolerogenic interactions and serves as a barrier to cell-mediated autoimmunity whereas a tumor or site of infection will attract mature myeloid dendritic cells that stimulate potent cell-mediated immune responses.

[0003] CD1d-restricted NKT cells are a unique class of non-conventional T cells that appear to play an important role in defining the outcome of immune stimulation in the

- 2 -

local environment. They share with the larger class of NKT cells the expression of markers of both the T cell and natural killer (NK) cell lineages. As such, NKT cells are considered as part of innate immunity like NK cells and in humans their frequency in normal individuals can be as high as 2.0% of total T lymphocytes (Gumperz *et al.*, 2002. J Exp Med 195:625; Lee *et al.*, 2002. J Exp Med 195:637).

[0004] CD1d-restricted NKT cells are distinguished from other NKT cells by their specificity for lipid and glycolipid antigens presented by the monomorphic MHC class Ib molecule, CD1d (Kawano *et al.*, Science 278 (1997), pp. 1626–1629). CD1d is a non-MHC encoded molecule that associates with β 2-microglobulin and is structurally related to classical MHC class I molecules. CD1d has a hydrophobic antigen-binding pocket that is specialized for binding the hydrocarbon chains of lipid tails or hydrophobic peptides (Zeng *et al.*, Science 277 (1997), pp. 339–345). CD1d is known to bind a marine sponge derived α -glycosylated sphingolipid, α -galactosylceramide (α -GalCer), and related molecules such as ceramide-like glycolipid antigens with α -linked galactose or glucose but not mannose (Kawano *et al.*, Science 278 (1997), pp. 1626–1629; and Zeng *et al.*, Science 277 (1997), pp. 339–345). As discussed below, the ability to activate CD1d-restricted NKT cells by stimulation with α -GalCer or related molecules bound to CD1d of antigen presenting cells has greatly facilitated functional analysis of this non-conventional T cell subset. In the absence of inflammation, CD1d-restricted NKT cells have been shown to localize preferentially in certain tissues like thymus, liver and bone marrow (Wilson *et al.*, 2002. Trends Mol Med 8:225) and antitumor activity of NKT cells has been mainly investigated in mouse liver metastasis.

[0005] NKT cells have an unusual ability of secreting both TH1 and TH2 cytokines and potent cytotoxic as well as regulatory functions have been documented in inflammation, autoimmunity and tumor immunity (Bendelac *et al.*, (1995) Science 268:863; Chen and Paul. 1997. J Immunol 159:2240; and Exley *et al.*, 1997. J Exp Med 186:109).

[0006] Among the CD1d-restricted NKT cells is a subset, referred to herein as "iNKT cells," that express a highly conserved $\alpha\beta$ T cell receptor (TCR). In man this invariant TCR is comprised of V α 24J α 15 in association with V β 11 whereas in mice the

- 3 -

receptor comprises the highly homologous V α 14J α 18 and V β 8.2. Other CD1d-restricted NKT cells express more variable TCR. Both TCR invariant and TCR variant classes of CD1d-restricted T cells can be detected by binding of CD1d-tetramers loaded with α -GalCer (Benlagha *et al.*, J Exp Med 191 (2000), pp. 1895–1903; Matsuda *et al.*, J Exp Med 192 (2000), pp. 741–754; and Karadimitris *et al.*, Proc Natl Acad Sci USA 98 (2001), pp. 3294–3298). CD1d-restricted NKT cells, as defined in this application (CD1d-restricted NKT), include cells that express either invariant or variant TCR and that bind or are activated by CD1d loaded with either α -GalCer or with related ceramide-like glycolipid antigens. CD1d-restricted NKT cells, as defined in this application (CD1d-NKT), include cells that express either invariant or variant TCR and that bind or are activated by CD1d loaded with either α -GalCer or with related sphingolipids that have α -linked galactose or glucose including molecules such as OCH, which differs from α -GalCer by having a shortened long-chain sphingosine base (C5 vs. C14) and acyl chain (C24 vs. C26) (Miyamoto *et al.*, Nature 2001 413:531-4).

[0007] CD1d-restricted NKT have been shown to have direct cytotoxic activity against targets that express CD1d. It is likely, however, that the effect of CD1d-restricted NKT on immune responses is amplified through recruitment of other lymphocytes either by direct interaction or, perhaps even more importantly, by indirect recruitment through interaction with DC. CD1d-restricted NKT have the unique ability to secrete large quantities of IL-4 and IFN- γ early in an immune response. Secretion of IFN- γ induces activation of DC which produce interleukin-12 (IL-12). IL-12 stimulates further IFN- γ secretion by NKT cells and also leads to activation of NK cells which secrete more IFN- γ .

[0008] Since CD1d-restricted NKT are able to rapidly secrete large amounts of both IL-4 and IFN- γ , the polarization of immune responses will depend on whether the effect of pro-inflammatory IFN- γ or anti-inflammatory IL-4 cytokines predominate. This has been reported to be, in part, a function of the relative frequency of different subsets of CD1d-restricted NKT. These subsets include (i) an invariant CD1d-restricted NKT population that is negative for both CD4 and CD8 and that gives rise

- 4 -

to predominantly a TH1 type response including secretion of pro-inflammatory IFN- γ and TNF- α and (ii) a separate population of CD1d-restricted NKT that is CD4⁺ and that gives rise to both a TH1 type and TH2 type response including secretion of the anti-inflammatory Th2-type cytokines IL-4, IL-5, IL-10 and IL-13 (Lee *et al.*, J Exp Med 2002; 195 :637-41; and Gumperz *et al.*, J Exp Med 2002; 195 :625-36). In addition, NKT cell activity is differentially modulated by depending on the particular ceramide-like glycolipid bound to CD1d (see, *e.g.*, US Patent Application Publication No. 2006/0052316). Local factors that influence activation of CD1d-restricted NKT subsets include the cytokine environment and, importantly, the DC that are recruited to that environment.

[0009] A number of indirect mechanisms contribute to the protective effect of CD1d-restricted NKT cells. Activation of NKT cells by administration of α -GalCer *in vivo* results in concomitant activation of NK cells (Eberl and MacDonald, Eur. J. Immunol. 30 (2000), pp. 985–992; and Carnaud *et al.*, J. Immunol. 163 (1999), pp. 4647–4650). In mice deficient in NKT cells, α -GalCer is unable to induce cytotoxic activity by NK cells. NKT cells also enhance the induction of classical MHC class I restricted cytotoxic T cells (Nishimura *et al.*, Int Immunol 2000; 12 :987-94; and Stober *et al.*, J Immunol 2003; 170:2540-8).

[0010] The availability of a defined antigen, *e.g.*, α -GalCer and related antigens, that can be employed to specifically activate CD1d-restricted NKT cells has made it possible to examine the role of these non-conventional T cells in a variety of immune responses.

[0011] Indeed, α -GalCer has significant promise as a therapeutic agent or adjuvant. For example, α -GalCer administration has a dramatic effect on a number of different microbial infections, including protective effects in murine malaria, fungal and hepatitis B virus infections (Kakimi *et al.*, J Exp Med 192 (2000), pp. 921–930; Gonzalez-Aseguinolaza *et al.*, Proc Natl Acad Sci USA 97 (2000), pp. 8461–8466; and Kawakami *et al.*, Infect Immun 69 (2001), pp. 213–220). Dramatic effects of administration of α -GalCer have also been observed in animal models of tumor immunity. For example, stimulation with α -GalCer suppresses lung and liver

- 5 -

metastases in an NKT dependent manner (Smyth *et al.*, 2002. Blood 99:1259). In addition, α -GalCer has been shown to have a protective effect against certain autoimmune diseases, including type 1 diabetes and experimental autoimmune encephalomyelitis (EAE, a well-known model system for multiple sclerosis) (Hong S, *et al.* Nat. Med. 2001;7:1052–1056 and Miyamoto K. *et al.* Nature. 2001;413:531–534).

[0012] However, NKT cells, upon restimulation with α -GalCer, become unresponsive, *e.g.*, reduced in their capacity to proliferate, produce cytokines, transactivate other cell types, and prevent tumor metastasis. Parekh, VV, *et al.* J. Clin. Invest. 115:2572-2583 (2005). Accordingly, there remains a need in the art for methods of stimulating NKT cells multiple times without causing the NKT cells to become nonresponsive.

Summary Of The Invention

[0013] In one embodiment, the present invention is directed to a method of modulating an immune response in an animal, comprising administering to an animal in need of immune modulation a composition comprising: (a) non-specific CD1d complex which comprises: (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen; (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with the CD1d polypeptide; and (iii) a ceramide-like glycolipid antigen bound to the CD1d polypeptide; and (b) a carrier; wherein the non-specific CD1d complex is administered in an amount sufficient to affect the activity of NKT cells in the animal.

[0014] In another embodiment, the present invention is directed to a method of treating a disease in an animal, comprising administering to an animal with said disease a composition comprising: (a) non-specific CD1d complex which comprises: (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen; (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with the CD1d polypeptide; and (iii) a ceramide-like glycolipid antigen bound to the CD1d

- 6 -

polypeptide; and (b) a carrier; wherein said composition is administered in an amount sufficient to alter the progression of the disease.

[0015] In another embodiment the present invention is directed to a method of inhibiting an anergic effect of a ceramide-like glycolipid antigen on NKT cell activity, comprising: stimulating NKT cells with the ceramide-like glycolipid antigen as part of a CD1d complex, where the complex comprises: (a) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen; (b) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with the CD1d polypeptide; and (c) the ceramide-like glycolipid antigen bound to the CD1d polypeptide; and restimulating said NKT cells one or more times with said complex; wherein said NKT cells are activated in response to said stimulation, and wherein said NKT cells are reactivated in response to said restimulation by said complex.

[0016] In yet another embodiment, the present invention is directed to a method of modulating an immune response to an immunogen in an animal, comprising administering to an animal in need thereof a composition comprising: (a) an immunogen; (b) a CD1d complex, said complex comprising: (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen; (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with the CD1d polypeptide; and (iii) a ceramide-like glycolipid antigen bound to the CD1d polypeptide; and (c) a carrier; wherein the CD1d complex is administered in an amount sufficient to modulate the immune response against the immunogen relative to administration of the immunogen in the absence of the CD1d complex.

[0017] In yet another embodiment the present invention is directed to a method of treating a disease in an animal, comprising administering to an animal in need thereof a composition comprising: (a) an immunogen; (b) a CD1d complex which comprises: (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen; (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with the CD1d polypeptide; and (iii) a ceramide-like glycolipid antigen bound to the CD1d

- 7 -

polypeptide; and (c) a carrier; wherein an immune response against the immunogen is effective in treating the disease, and wherein the CD1d complex is administered in an amount sufficient to modulate the immune response against the immunogen relative to administration of the immunogen in the absence of the CD1d complex.

[0018] In yet another embodiment the present invention is directed to a method of preventing a disease in an animal, comprising administering to an animal in need thereof a composition comprising: (a) an immunogen; (b) a CD1d complex which comprises: (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen; (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with the CD1d polypeptide; and (iii) a ceramide-like glycolipid antigen bound to the CD1d polypeptide; and (c) a carrier; wherein an immune response against the immunogen is effective in treating the disease, and wherein the CD1d complex is administered in an amount sufficient to modulate the immune response against the immunogen relative to administration of the immunogen in the absence of the CD1d complex.

[0019] The present invention is further directed to a composition comprising any CD1d complex described herein.

Brief Description of the Figures

[0020] Figure 1: *In vivo* biological activity of α GalCer-loaded sCD1d and CD1d-anti-HER2 fusion protein shown by the transient disappearance of liver iNKT cells 20 hours after i.p. injection with PBS (control), α GalCer 5 μ g, or α GalCer/sCD1d 20 μ g or α GalCer/CD1d-anti-HER2 fusion 40 μ g (loaded or unloaded). Frequency of iNKT cells was measured by flow cytometry using CD1d-Tetramer-Extravidin-PE and anti CD3 FITC.

[0021] Figure 2: a Sustained IFN γ production by liver and spleen iNKT cells after several injections of CD1d/anti-HER2 fusion. Liver and spleen lymphocytes were isolated 20 minutes after the sixth injection of either PBS (control, white bar), α GalCer 0.4 μ g (grey bar), or α GalCer/CD1d-anti-HER2 fusion protein 40 μ g (black bar) and cultured for 1 hour in presence of Golgi Plug reagent. NKT cells were then

- 8 -

stained with anti NK1.1-PE and anti CD3-FITC antibodies, fixed and stained for intracellular IFN γ with anti IFN γ -APC. Graph shows percentage of IFN γ producing NKT cells (gated on NK1.1+ CD3+ cells). **b** Sustained IFN γ production by liver NKT cells after several injections i.v. and *in vitro* rechallenge with α GalCer or α GalCer loaded recombinant CD1d molecules. Liver lymphocytes were isolated after 5 injections i.v. of either PBS (Control), α GalCer, α GalCer/CD1d-anti-HER2 fusion protein (Fusion) or α GalCer/sCD1d (sCD1d) and stimulated *in vitro* for 6 hours in presence of Golgi Plug as indicated. (PBS, white; α GalCer, light grey; α GalCer/CD1d-anti-HER2 fusion, black; α GalCer/sCD1d, dark grey) Cells were stained for FACS analysis as described in **a**.

[0022] Figure 3: iNKT expansion in blood during systemic treatment with α GalCer loaded recombinant CD1d molecules. Mice were bled after the third injection of either PBS (control) α GalCer (0.4 μ g), α GalCer/CD1d-anti-Her2 fusion protein (Fusion, 40 μ g), or α GalCer/sCD1d (sCD1d, 20 μ g). NKT cells were stained in PBMCs using the CD1d-Tetramer-PE and anti CD3 FITC antibody. **a** representative dot blot of one mouse from each group. **b** graph representing several mice per group expressed as CD1d tetramer positive and CD3+ cells as percentage of total PBMC.

[0023] Figure 4: Sustained iNKT activation *in vivo* due to repeated injections of α GalCer/sCD1d prevents formation of lung metastases. Mice were treated 5 times with PBS (Control, white), α GalCer (0.4 μ g, light grey), or α GalCer/sCD1d (sCD1d, 20 μ g, black) and then grafted with 700,000 B16 wild type cells plus co-injection of the respective treatment. 2 naive groups were included that had no pretreatment and got only tumor cell graft + co-injection of α GalCer (striped grey) or α GalCer/sCD1d (sCD1d, dark striped grey) respectively. Lung metastases formation was analysed 2 weeks after graft with the ImageJ k-means clustering program and results are expressed as percent of black metastatic surface over total lung surface. P values for α GalCer/sCD1d pretreated and naive+ α GalCer groups compared to control $*P < 0.04$, and compared to α GalCer pretreated group $*P < 0.02$.

[0024] Figure 5: Precoating experiment. B16-HER2 (**a**) and B16 wild type (B16 wt) cells (**b**) were precoated for 1 hour with 0.4 μ g/ml α GalCer, 40 μ g/ml α GalCer/CD1d-

- 9 -

Her2 Fusion protein (Fusion), 10µg/ml Herceptin, or 20µg/ml αGalCer/sCD1d (sCD1d) and, with or without a previous wash, cells were injected i.v. Lung metastases were analyzed 3 weeks after graft with the ImageJ k-means clustering program and results are expressed as percentage of black metastatic surface over total lung surface. * $P < 0.005$ compared to PBS control.

[0025] Figure 6: *In vivo* anti tumor activity - Systemic Treatment. **a** Mice were grafted i.v. with 700.000 B16-HER2 cells and i.v. treatment was started 48 hours later. Mice were injected i.v. 5 times every 2 to 3 days with PBS (control), αGalCer (0.4µg), αGalCer/sCD1d (sCD1d, 25µg), or αGalCer/CD1d-anti-HER2 fusion (Fusion, 40µg). Lung metastases were analysed after 3 weeks by the ImageJ k-means clustering program and expressed as percent metastatic surface over total lung surface. Treatment with αGalCer/CD1d-anti-HER2 fusion protein significantly inhibited the metastases formation. * $P < 0.005$ versus control, * $P < 0.06$ versus αGalCer **b** Treatment was started 6 days after graft. Mice were injected i.v. with 700,000 B16-HER2 cells and treatment with PBS (Control), αGalCer, or αGalCer/CD1d-anti-HER2 fusion protein (Fusion) was started 6 days after graft. Mice were treated 3x i.v. every 2-3 days and lung metastases were analyzed after 3 weeks as described above. Only treatment with the CD1d-anti HER2 fusion protein significantly reduced metastatic growth. * $P < 0.01$ versus control.

[0026] Figure 7: Transactivation of NK, DCs and T cells by αGalCer/sCD1d or αGalCer/CD1d-anti-HER2 fusion protein activated iNKT cells. **a** Increase of liver NK cell numbers 20 hours after i.p. treatment with either PBS (control, white bar), 5µg αGalCer (light grey bar), or 20µg αGalCer-loaded sCD1d or 40µg αGalCer/CD1d-anti-HER2 fusion protein (αGalCer-sCD1d, dark grey and αGalCer-Fusion, black bars, respectively). Cells were stained with anti NK1.1-PE and anti CD3-FITC and analysed by flow cytometry. NK cells are reported as NK1.1 positive/CD3 negative population. **b c** Induction of DC maturation and T cell proliferation by αGalCer/sCD1d activated NKT cells *in vivo*. Splenocytes were isolated after five treatments i.v. with either PBS (control), αGalCer, or αGalCer/sCD1d and cultured for 4 days with GM-CSF and then 3 more days with either PBS (control, white bars),

- 10 -

α GalCer (grey bars), or α GalCer/sCD1d (sCD1d, black bars). Cells were then stained with anti CD11c-FITC and biotinylated anti CD40 and Streptavidin-PE to detect double positive mature DCs (c), or with anti NK1.1-PE and anti CD3-FITC to detect CD3 single positive T cells (d) by flow cytometry.

[0027] Figure 8: α GalCer loaded sCD1d acts as a strong adjuvant for the expansion of antigen-specific T cells upon active immunization. **a** kinetics of expansion of H-2K^b/OVA specific CTLs. Mice were primed with 200 μ g ovalbumin either as such (i.v.) or with Montanide adjuvant (3:7 ratio, s.c.) or with 1 μ g α GalCer or 20 μ g α GalCer/sCD1d (i.v.). Mice were bled every 5-7 days and PBMC were stained with H-2K^b/OVA tetramer+anti-CD8. Results are expressed as tetramer+ and CD8+ NKT cells as a percentage of total CD8+ T cells. Mice were boosted with OVA peptide (20 μ g) on day 22 employing the same adjuvants and route of injection as for primary immunization. **b** Ten days later, mice were sacrificed and splenocytes were cultured another five days either with no stimuli or with the same stimuli and adjuvants as for the *in vivo* boost. Frequency of OVA specific T cells was measured as in **a**. **c** The presence of mature DC in the same cultures as in **b** was analyzed by surface staining with anti CD11c and anti CD40 antibodies.

Detailed Description Of The Invention

[0028] The present invention provides compositions and methods which are useful for modulating, *i.e.*, either eliciting, inhibiting, or stimulating, an immune response. The compounds comprise one or more CD1d complexes comprising a ceramide-like glycolipid antigen bound to a soluble CD1d polypeptide fragment associated with beta-2 microglobulin. In certain embodiments, the soluble CD1d complexes of the present invention are non-specific, *i.e.*, they are not targeted to any particular tissue, cell, or cell surface marker. Soluble CD1d complexes for use in the methods of the present invention modulate an immune response by affecting the activity of CD1d-restricted natural killer T ("NKT") cells. Soluble CD1d complexes as described herein are useful for stimulating desirable immune responses, for example, immune responses against infectious agents or cancer; or for inhibiting undesirable immune

- 11 -

responses, such as allergic responses, allograft rejections, and autoimmune diseases. In certain embodiments, soluble CD1d complexes of the present invention are administered with an immunogen and function as an adjuvant by, for example, increasing or modulating the immune response to the immunogen.

Definitions

[0029] It is to be noted that the term "a" or "an" entity refers to one or more of that entity; for example, "a vector" is understood to represent one or more vectors. As such, the terms "a" (or "an"), "one or more," and "at least one" can be used interchangeably herein.

[0030] As used herein, the term "polypeptide" is intended to encompass a singular "polypeptide" as well as plural "polypeptides," and refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). The term "polypeptide" refers to any chain or chains of two or more amino acids, and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides, "protein," "amino acid chain," or any other term used to refer to a chain or chains of two or more amino acids, are included within the definition of "polypeptide," and the term "polypeptide" may be used instead of, or interchangeably with any of these terms. The term "polypeptide" is also intended to refer to the products of post-expression modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, or modification by non-naturally occurring amino acids. A polypeptide may be derived from a natural biological source or produced by recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It may be generated in any manner, including by chemical synthesis.

[0031] A polypeptide of the invention may be of a size of about 3 or more, 5 or more, 10 or more, 20 or more, 25 or more, 50 or more, 75 or more, 100 or more, 200 or more, 500 or more, 1,000 or more, or 2,000 or more amino acids. Polypeptides may have a defined three-dimensional structure, although they do not necessarily have such structure. Polypeptides with a defined three-dimensional structure are referred to

- 12 -

as folded, and polypeptides which do not possess a defined three-dimensional structure, but rather can adopt a large number of different conformations, and are referred to as unfolded. As used herein, the term glycoprotein refers to a protein coupled to at least one carbohydrate moiety that is attached to the protein via an oxygen-containing or a nitrogen-containing side chain of an amino acid residue, *e.g.*, a serine residue or an asparagine residue.

[0032] By an "isolated" polypeptide or a fragment, variant, or derivative thereof is intended a polypeptide that is not in its natural milieu. No particular level of purification is required. For example, an isolated polypeptide can be removed from its native or natural environment. Recombinantly produced polypeptides and proteins expressed in host cells are considered isolated for purposes of the invention, as are native or recombinant polypeptides which have been separated, fractionated, or partially or substantially purified by any suitable technique.

[0033] Also included as polypeptides of the present invention are fragments, derivatives, analogs, or variants of the foregoing polypeptides, and any combination thereof. The terms "fragment," "variant," "derivative" and "analog" when referring to polypeptides of the present invention include any polypeptides that retain at least some of the biological, antigenic, or immunogenic properties of the corresponding native polypeptide. Fragments of polypeptides of the present invention include proteolytic fragments, as well as deletion fragments, in addition to other specific fragments discussed elsewhere herein. Variants of polypeptides of the present invention include fragments as described above, and also polypeptides with altered amino acid sequences due to amino acid substitutions, deletions, or insertions. Variants may occur naturally or be non-naturally occurring. Non-naturally occurring variants may be produced using art-known mutagenesis techniques. Variant polypeptides may comprise conservative or non-conservative amino acid substitutions, deletions or additions. Derivatives of polypeptides of the present invention, are polypeptides which have been altered so as to exhibit additional features not found on the native polypeptide. Examples include fusion proteins. Variant polypeptides may also be referred to herein as "polypeptide analogs." As used herein a "derivative" of a polypeptide refers to a subject polypeptide having one

- 13 -

or more residues chemically derivatized by reaction of a functional side group. Also included as "derivatives" are those peptides which contain one or more naturally occurring amino acid derivatives of the twenty standard amino acids. For example, 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine.

[0034] The term "polynucleotide" is intended to encompass a singular nucleic acid as well as plural nucleic acids, and refers to an isolated nucleic acid molecule or construct, *e.g.*, messenger RNA (mRNA), virally-derived RNA, or plasmid DNA (pDNA). A polynucleotide may comprise a conventional phosphodiester bond or a non-conventional bond (*e.g.*, an amide bond, such as found in peptide nucleic acids (PNA)). The term "nucleic acid" refers to any one or more nucleic acid segments, *e.g.*, DNA or RNA fragments, present in a polynucleotide. By "isolated" nucleic acid or polynucleotide is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, a recombinant polynucleotide encoding a therapeutic polypeptide contained in a vector is considered isolated for the purposes of the present invention. Further examples of an isolated polynucleotide include recombinant polynucleotides maintained in heterologous host cells or purified (partially or substantially) polynucleotides in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the present invention, as well as positive and negative strand forms, and double-stranded forms, of pestivirus vectors disclosed herein.

[0035] Isolated polynucleotides or nucleic acids according to the present invention further include such molecules produced synthetically. In addition, a polynucleotide or a nucleic acid may be or may include a regulatory element such as a promoter, ribosome binding site, or a transcription terminator.

[0036] As used herein, a "coding region" is a portion of nucleic acid which consists of codons translated into amino acids. Although a "stop codon" (TAG, TGA, or TAA) is not translated into an amino acid, it may be considered to be part of a coding region, if present, but any flanking sequences, for example promoters, ribosome binding sites, transcriptional terminators, introns, 5' and 3' non-translated regions, and the like, are

- 14 -

not part of a coding region. Two or more coding regions of the present invention can be present in a single polynucleotide construct, *e.g.*, on a single vector, or in separate polynucleotide constructs, *e.g.*, on separate (different) vectors. Furthermore, any vector may contain a single coding region, or may comprise two or more coding regions, *e.g.*, a vector of the present invention may encode one or more polyproteins, which are post- or co-translationally separated into the final proteins via proteolytic cleavage. In addition, a vector, polynucleotide, or nucleic acid of the invention may encode heterologous coding regions, either fused or unfused to a first or second nucleic acid encoding of the invention, or variant or derivative thereof. Heterologous coding regions include without limitation specialized elements or motifs, such as a secretory signal peptide or a heterologous functional domain.

[0037] In certain embodiments, the polynucleotide or nucleic acid is DNA. In the case of DNA, a polynucleotide comprising a nucleic acid, which encodes a polypeptide normally may include a promoter and/or other transcription or translation control elements operably associated with one or more coding regions. An operable association is when a coding region for a gene product, *e.g.*, a polypeptide, is associated with one or more regulatory sequences in such a way as to place expression of the gene product under the influence or control of the regulatory sequence(s). Two DNA fragments (such as a polypeptide coding region and a promoter associated therewith) are "operably associated" if induction of promoter function results in the transcription of mRNA encoding the desired gene product and if the nature of the linkage between the two DNA fragments does not interfere with the ability of the expression regulatory sequences to direct the expression of the gene product or interfere with the ability of the DNA template to be transcribed. Thus, a promoter region would be operably associated with a nucleic acid encoding a polypeptide if the promoter was capable of effecting transcription of that nucleic acid. The promoter may be a cell-specific promoter that directs substantial transcription of the DNA only in predetermined cells. Other transcription control elements, besides a promoter, for example enhancers, operators, repressors, and transcription termination signals, can be operably associated with the polynucleotide to direct cell-specific transcription. Suitable promoters and other transcription control regions are disclosed herein.

[0038] A variety of transcription control regions are known to those skilled in the art. These include, without limitation, transcription control regions, which function in vertebrate cells, such as, but not limited to, promoter and enhancer segments from cytomegaloviruses (*e.g.*, the immediate early promoter, in conjunction with intron-A), simian virus 40 (*e.g.*, the early promoter), and retroviruses (such as, *e.g.*, Rous sarcoma virus). Other transcription control regions include those derived from vertebrate genes such as actin, heat shock protein, bovine growth hormone and rabbit β -globin, as well as other sequences capable of controlling gene expression in eukaryotic cells. Additional suitable transcription control regions include tissue-specific promoters and enhancers as well as lymphokine-inducible promoters (*e.g.*, promoters inducible by interferons or interleukins).

[0039] Similarly, a variety of translation control elements are known to those of ordinary skill in the art. These include, but are not limited to ribosome binding sites, translation initiation and termination codons, and elements derived from viral systems (particularly an internal ribosome entry site, or IRES, also referred to as a CITE sequence).

[0040] In other embodiments, a polynucleotide of the present invention is RNA, for example, in the form of messenger RNA (mRNA). RNA of the present invention may be single stranded or double stranded.

[0041] Polynucleotide and nucleic acid coding regions of the present invention may be associated with additional coding regions which encode secretory or signal peptides, which direct the secretion of a polypeptide encoded by a polynucleotide of the present invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal peptide or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Those of ordinary skill in the art are aware that polypeptides secreted by vertebrate cells generally have a signal peptide fused to the N-terminus of the polypeptide, which is cleaved from the complete or "full length" polypeptide to produce a secreted or "mature" form of the polypeptide. In certain embodiments, the native signal peptide, *e.g.*, an immunoglobulin heavy chain or light chain signal peptide is used, or a functional derivative of that sequence that retains the

- 16 -

ability to direct the secretion of the polypeptide that is operably associated with it. Alternatively, a heterologous mammalian signal peptide, or a functional derivative thereof, may be used. For example, the wild-type leader sequence may be substituted with the leader sequence of human tissue plasminogen activator (TPA) or mouse β -glucuronidase.

[0042] The term "construct" refers to an engineered vector.

[0043] The term "artificial" refers to a synthetic, or non-host cell derived composition, *e.g.*, a chemically-synthesized oligonucleotide.

[0044] As discussed in more detail below, a functional antigen-loaded soluble fragment of a CD1d polypeptide, including both CD1d and β -2 microglobulin subunits, is referred to herein as a "soluble CD1d complex." The antigen to be loaded onto the CD1d polypeptide is a glycolipid, typically a ceramide-like glycolipid, *e.g.*, an alpha-galactosylceramide, *e.g.*, α -GalCer. "Ceramide-like glycolipids," as referred to herein include glycolipids with α -linked galactose or glucose. Examples of glycolipid antigens which bind to CD1d are found, *e.g.*, in Porcelli, U.S. Patent Appl. Publ. No. 2006/0052316, Tsuji, U.S. Patent Appl. Publ. No. 2006/0211856, Jiang, U.S. Patent Appl. Publ. No. 2006/0116331, Hirokazu *et al.*, U.S. Patent Appl. Publ. No. 2006/0074235, Tsuji *et al.*, U.S. Patent Appl. Publ. No. 2005/0192248, Tsuji, U.S. Patent Application No. 2004/0127429, and Tsuji *et al.*, U.S. Patent Application No. 2003/0157135, all of which are incorporated by reference herein in their entireties.

[0045] The term "non-specific soluble CD1d complex" refers to a soluble CD1d complex which has not been engineered to be targeted to any specific organ, tissue, cell, or cell-surface molecule. A "non-specific soluble CD1d complex" is, however, capable of interacting with NKT cells, in a way similar to that in which a cell-surface-expressed CD1d molecule, when loaded with antigen, would interact. In contrast to a "non-specific soluble CD1d complex" is a "targeted CD1d complex," which is fused or conjugated to an antibody or other binding molecule, thus targeting the complex to a specific organ, tissue, cell, or cell-surface marker. Targeted CD1d complexes exert their effect on NKT cells locally, *e.g.*, in the vicinity of a tumor. See, *e.g.*, Bruno *et al.* U.S. Patent Appl. Publ. No. 2006/0269540, incorporated herein by reference in its entirety.

[0046] Antibodies are constructed of one, or several, units, each of which consists of two heavy (H) polypeptide chains and two light (L) polypeptide chains. The H and L chains are made up of a series of domains. The L chains, of which there are two major types (κ and λ), consists of two domains. The H chains are of several types, including μ , δ , and γ (of which there are several subclasses), α and ϵ . In humans, there are eight genetically and structurally identified antibody classes and subclasses as defined by heavy chain isotypes: IgM, IgD, IgG3, IgG1, IgG2, IgG4, IgE, and IgA. Further, for example, "IgG" means an antibody of the G class, and that, "IgG1" refers to an IgG molecules of subclass 1 of the G class. IgG1 antibodies, like all antibodies of the IgG class, are comprised of 4 domains, one of which is variable and the other 3 are constant. An Fab antibody fragment is comprised of an intact light chain and a truncated heavy chain that each comprise two domains, one variable and one constant.

[0047] As used herein, the term "antibody" (Ab) or "monoclonal antibody" (MAb) is meant to include intact molecules as well as antibody portions (such as, for example, Fab and F(ab')₂ portions and Fv fragments) which are capable of specifically binding to a cell surface marker. Such portions are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab portions) or pepsin (to produce F(ab')₂ portions). Especially preferred in the compounds of the invention are Fab portions. Alternatively, antigen-binding portions can be produced through the application of recombinant DNA technology.

[0048] In addition, the immunoglobulin may be a single chain antibody ("SCA"). These may consist of single chain Fv fragments ("scFv") in which the variable light ("V[L]") and variable heavy ("V[H]") domains are linked by a peptide bridge or by disulfide bonds. Also, the immunoglobulin may consist of single V[H]domains (dAbs) which possess antigen-binding activity. See, *e.g.*, G. Winter and C. Milstein, *Nature* 349:295 (1991); R. Glockshuber *et al.*, *Biochemistry* 29:1362 (1990); and, E. S. Ward *et al.*, *Nature* 341:544 (1989).

[0049] Also preferred for use in the present invention are chimeric monoclonal antibodies, preferably those chimeric antibodies having specificity toward a tumor associated surface membrane antigen, a surface membrane antigen of a tissue or organ affected by autoimmune disease, or an antigen of a pathogen infected cell. As used in

- 18 -

this example, the term "chimeric antibody" refers to a monoclonal antibody comprising a variable region, *i.e.* binding region, from one source or species and at least a portion of a constant region derived from a different source or species, usually prepared by recombinant DNA techniques.

[0050] Encompassed by the term "chimeric antibody" is the concept of "humanized antibody", that is those antibodies in which the framework or "complementarity" determining regions ("CDR") have been modified to comprise the CDR of an immunoglobulin of different specificity as compared to that of the parent immunoglobulin. In certain embodiments, a murine CDR is grafted into the framework region of a human antibody to prepare the "humanized antibody". See, *e.g.*, L. Riechmann *et al.*, Nature 332:323 (1988); M. S. Neuberger *et al.*, Nature 314:268 (1985).

[0051] In other embodiments, fully human antibodies or fragments thereof are used in the compositions and methods of the invention, preferably those fully human antibodies having specificity toward a tumor associated surface membrane antigen, a surface membrane antigen of a tissue or organ affected by autoimmune disease, or an antigen of a pathogen infected cell. Methods have been described for selection of fully human antibodies in human immunoglobulin transgenic mice, from libraries of human immunoglobulin genes constructed in phage and expressed in bacteria or constructed in a mammalian viral expression vector for expression in mammalian cells, and from human hybridoma cells. A method for selection of fully human antibodies from libraries of human immunoglobulin genes constructed in vaccinia virus is described in Zauderer, M. *et al.* WO 01/72995, published 4 October 2001, the disclosure of which is incorporated by reference herein.

[0052] In certain embodiments, targeted CD1d complexes of the present invention comprise, instead of, or in addition to an antibody, a specific binding molecule, *e.g.*, a receptor or ligand that has a matching or counterpart ligand or receptor expressed on a cell surface of a target cell. In these embodiments, the targeted CD1d complex comprises a ligand or receptor specific for a cell surface marker. Examples include: CD4 coupled to CD1d for interaction with HIV infected cells; chemokine or

- 19 -

chemokine receptor coupled to CD1d for interaction with DC subset; or heregulins coupled to CD1d for interaction with ErbB2 positive tumor cells.

- [0053] In one embodiment, the antibody is specific for a cell surface marker of a tumor cell. In another embodiment, the antibody is specific for a cell surface marker of a CD1d-restricted NKT cell. In another embodiment, the antibody is specific for a cell surface marker of a target tissue of autoimmune disease or inflammatory response. In another embodiment, the antibody is specific for an infectious agent or a cell surface marker of an infected cell or tissue.
- [0054] In another embodiment, the antibody is specific for a cell surface marker of a professional antigen presenting cell, *e.g.*, a dendritic cell.
- [0055] The term "antigen" and the related term "antigenic" as used herein refers to a substance that binds specifically to an antibody or to a T-cell receptor.
- [0056] The term "immunogen" and the related term "immunogenic" as used herein refers to the ability to induce an immune response, including an antibody and/or a cellular immune response in an animal, preferably a mammal. It is quite likely that an immunogen will also be antigenic, but an "antigen," because of its size or conformation, may not necessarily be an "immunogen." An "immunogenic composition" induces an immune response in a subject, *e.g.*, antibodies that specifically recognize one or more antigens, contained within that "immunogenic composition."
- [0057] The term "immune response" is meant to include any activity of cells of the immune system in response to an antigen or immunogen. Such activities include, but are not limited to production of antibodies, cytotoxicity, lymphocyte proliferation, release of cytokines, inflammation, phagocytosis, antigen presentation, and the like. An immune response which is highly specific to a given antigen or immunogen, *e.g.*, production of specific antibodies or production of specific T lymphocytes is referred to herein as an "adaptive immune response." An immune response which is not specific to a given antigen, *e.g.*, release of cytokines by NK and NKT cells, is referred to herein as an "innate immune response." Examples of immune responses include an antibody response or a cellular, *e.g.*, cytotoxic T-cell, response.

- [0058] The terms "protective immune response" or "therapeutic immune response" refer to an immune response to an immunogen which in some way prevents or at least partially arrests disease symptoms, side effects or progression. By "protective" is meant that the immune response is induced in a subject animal which has not contracted a disease, where the immune response alleviates, reduces, moderates or, in some cases fully prevents disease symptoms if the animal later contracts or is susceptible to that disease. By "therapeutic" is meant that the immune response is induced in a subject animal which has the disease, where the immune response alleviates, reduces, moderates, or in some cases fully eliminates disease symptoms.
- [0059] The term "modulating an immune response" is meant to refer to any way in which a given immune response is increased, decreased, or changed by a composition or treatment relative to the immune response without that composition or treatment. For example, use of an adjuvant to increase an immune response to an antigen is considered modulation of that immune response. Decrease in an immune response, *e.g.*, prevention of autoimmunity, is also a modulation. In addition, changing an immune response, *e.g.*, from a TH2 response to a TH1 response, is a modulation of an immune response.
- [0060] The term "anergy" refers to a specific kind of immune modulation, in which certain cells of the immune system are rendered non-responsive to antigen stimulus. An example would be the ability of free α -GalCer, upon multiple administrations to an animal, to render the NKT cells of that animal non-responsive to stimulus, *e.g.*, unable to proliferate or produce cytokines.
- [0061] The term "adjuvant" refers to any material having the ability to (1) alter or increase the immune response to a particular antigen or (2) increase or aid an effect of a pharmacological agent. In certain embodiments, a soluble CD1d complex of the present invention, *e.g.*, a non-specific soluble CD1d complex, functions as an adjuvant upon administration with an immunogen. Other suitable adjuvants include, but are not limited to, cytokines and growth factors; bacterial components (*e.g.*, endotoxins, in particular superantigens, exotoxins and cell wall components); aluminum-based salts; calcium-based salts; silica; polynucleotides; toxoids; serum

proteins, viruses and virally-derived materials, poisons, venoms, imidazoquiniline compounds, poloxamers, and cationic lipids.

[0062] A great variety of materials have been shown to have adjuvant activity through a variety of mechanisms. Any compound which may increase the expression, antigenicity or immunogenicity of the polypeptide is a potential adjuvant. Potential adjuvants include, but are not limited to: inert carriers, such as alum, bentonite, latex, and acrylic particles; pluronic block polymers, such as TiterMax® (block copolymer CRL-8941, squalene (a metabolizable oil) and a microparticulate silica stabilizer), depot formers, such as Freund's adjuvant, surface active materials, such as saponin, lysolecithin, retinal, Quil A, liposomes, and pluronic polymer formulations; macrophage stimulators, such as bacterial lipopolysaccharide; alternate pathway complement activators, such as insulin, zymosan, endotoxin, and levamisole; and non-ionic surfactants, such as poloxamers, poly(oxyethylene)-poly(oxypropylene) tri-block copolymers.

[0063] In certain embodiments, the adjuvant is a cytokine. A composition of the present invention can comprise one or more cytokines, chemokines, or compounds that induce the production of cytokines and chemokines. Examples include, but are not limited to granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), colony stimulating factor (CSF), erythropoietin (EPO), interleukin 2 (IL-2), interleukin-3 (IL-3), interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 6 (IL-6), interleukin 7 (IL-7), interleukin 8 (IL-8), interleukin 10 (IL-10), interleukin 12 (IL-12), interleukin 15 (IL-15), interleukin 18 (IL-18), interferon alpha (IFN α), interferon beta (IFN β), interferon gamma (IFN γ), interferon omega (IFN ω), interferon tau (IFN τ), interferon gamma inducing factor I (IGIF), transforming growth factor beta (TGF- β), RANTES (regulated upon activation, normal T-cell expressed and presumably secreted), macrophage inflammatory proteins (*e.g.*, MIP-1 alpha and MIP-1 beta), *Leishmania* elongation initiating factor (LEIF), and Flt-3 ligand.

[0064] The term "vertebrate" is intended to encompass a singular "vertebrate" as well as plural "vertebrates" and comprises mammals and birds, as well as fish, reptiles, and amphibians.

- 22 -

[0065] The term "mammal" is intended to encompass a singular "mammal" and plural "mammals," and includes, but is not limited to humans; primates such as apes, monkeys (*e.g.*, owl, squirrel, cebus, rhesus, African green, patas, cynomolgus, and cercopithecus), orangutans, baboons, gibbons, and chimpanzees; canids such as dogs and wolves; felids such as cats, lions, and tigers; equines such as horses, donkeys, and zebras, food animals such as cows, pigs, and sheep; ungulates such as deer and giraffes; ursids such as bears; and others such as rabbits, mice, ferrets, seals, whales. In particular, the mammal can be a human subject, a food animal or a companion animal.

[0066] The term "bird" is intended to encompass a singular "bird" and plural "birds," and includes, but is not limited to feral water birds such as ducks, geese, terns, shearwaters, and gulls; as well as domestic avian species such as turkeys, chickens, quail, pheasants, geese, and ducks. The term "bird" also encompasses passerine birds such as starlings and budgerigars.

Soluble CD1d Complexes

[0067] As mentioned above, soluble CD1d complexes of the present invention can be used both to prevent a disease, and also to therapeutically treat a disease. In individuals already suffering from a disease, the present invention is used to further stimulate or modulate the immune system of the animal, thus reducing or eliminating the symptoms associated with that disease or disorder. As defined herein, "treatment" refers to the use of one or more compositions of the present invention to prevent, cure, retard, or reduce the severity of given disease symptoms in an animal, and/or result in no worsening of the disease over a specified period of time in an animal which has already contracted the disease and is thus in need of therapy. The term "prevention" refers to the use of one or more compositions of the present invention to generate immunity in an animal which has not yet contracted a disease, thereby preventing or reducing disease symptoms if the vertebrate is later disposed to develop that disease. The methods of the present invention therefore may be referred to as therapeutic methods or preventative or prophylactic methods. It is not required that any composition of the present invention provide total immunity to a disease agent or

- 23 -

totally cure or eliminate all disease symptoms. As used herein, an "animal in need of therapeutic and/or preventative immunity" refers to an individual for whom it is desirable to treat, *i.e.*, to prevent, cure, retard, or reduce the severity of certain disease symptoms, and/or result in no worsening of disease over a specified period of time.

[0068] The present invention provides methods of modulating an immune response comprising administering to an animal a composition which comprises an antigen-loaded soluble CD1d molecule, which interact with, and thereby affects, the activity of CD1d-restricted NKT cells. The monomorphic CD1d molecule is suitable for activation of a broad spectrum of CD1d-restricted NKT in an entire species. Soluble CD1d molecules, which include both CD1d and β -2 microglobulin subunits, are loaded with a ceramide-like glycolipid antigen, for example, α -GalCer, to produce a soluble CD1d complex. Soluble CD1d complexes of the present invention, *e.g.*, non-specific soluble CD1d complexes, can be the primary or only active ingredient in a composition of the present invention, for example for treatment of cancer. In other embodiments, soluble CD1d complexes of the present invention may be used as an adjuvant in combination with a specific immunogen, thereby, stimulating, increasing, modulating, or otherwise altering an immune response to that immunogen relative to administration of the immunogen without the soluble CD1d complex. The present invention further encompasses pharmaceutical compositions which comprise an immunogen and a soluble CD1d complex adjuvant.

[0069] Moreover, soluble CD1d complexes of the present invention may be used as a diagnostic or therapeutic agent not only for cancer and infectious diseases but also for a large class of autoimmune and inflammatory diseases that result from a failure to down modulate cell-mediated immune responses.

[0070] Soluble CD1d complexes for use in the methods of the present invention comprise a soluble fragment of a CD1d polypeptide sufficient to bind β 2-microglobulin as well as a ceramide-like glycolipid antigen, a β 2-microglobulin polypeptide, and a ceramide-like glycolipid antigen, *e.g.*, α -GalCer. The ceramide-like glycolipid antigen is bound in the antigen binding groove of the CD1d molecule.

[0071] As taught by WO 9964597, published 16 December 1999 and incorporated herein by reference, it is possible to introduce mutations into β 2-microglobulin that

- 24 -

increase affinity for the class I heavy chain so as to facilitate assembly and increase stability of the CD1d complex in the fusion protein. In certain embodiments, a soluble CD1d polypeptide is linked to β 2-microglobulin as a fusion protein. In certain embodiments, the β 2-microglobulin polypeptide is linked via its C-terminus to the N-terminus of the soluble CD1d polypeptide. Such fusion constructs can be made using conventional recombinant nucleic acid techniques. The fusion may be direct or may contain spacers. A short linker amino acid sequence may be inserted between the CD1d polypeptide and the β 2-microglobulin polypeptide. If a linker sequence is included, this sequence will preferably contain at least 3 and not more than 30 amino acids. More preferably, the linker is about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, or 25 amino acids long. Generally, the linker consists of short glycine/serine spacers, but any known amino acid may be used. Examples of linkers known to those skilled in the art include (Gly₄Ser)₃ (SEQ ID NO:3) and (Gly₄Ser)₂Gly₃AlaSer (SEQ ID NO:4).

[0072] Alternatively, the CD1d and β 2-microglobulin polypeptides may be chemically linked. A number of reagents capable of cross-linking proteins are known in the art, illustrative entities include: azidobenzoyl hydrazide, N-[4-(p-azidosalicylamino)butyl]-3'-[2'-pyridyldithio] propionamide), bis-sulfosuccinimidyl suberate, dimethyladipimide, disuccinimidyltartrate, N- γ -maleimidobutyryloxysuccinimide ester, N-hydroxy sulfosuccinimidyl-4-azidobenzoate, N-succinimidyl [4-azidophenyl]-1,3'-dithiopropionate, N-succinimidyl [4-iodoacetyl]aminobenzoate, glutaraldehyde, formaldehyde and succinimidyl 4-[N-maleimidomethyl] cyclohexane-1-carboxylate.

[0073] In certain embodiments, multiple CD1d complexes are linked together through a multivalent compound. The CD1d complexes may be linked to the multivalent compound through any site. In a preferred embodiment soluble CD1d polypeptides are linked to the multivalent compound through the CD1d carboxyl terminus. These compounds typically comprise 2 or more CD1d complexes. The compounds may comprise 2, 3, 4, 5, 6, 7, 8, 9 or 10 CD1d complexes.

[0074] Examples of multivalent compounds are chicken avidin or streptavidin (Shin, S.U. *et al.*, *J. Immunology* 158: 4797-4804 (1997)) to which biotinylated CD1d

- 25 -

complexes are bound (Altman, J. *et al*, *Science* 274:94-96 (1996); Boniface, J.J. *et al*., *Immunity* 9:459-66 (1998)); or a leucine zipper system.

[0075] Alternatively, CD1d and β 2-microglobulin polypeptides can be genetically modified by including sequences encoding amino acid residues with chemically reactive side chains such as Cys or His. Such amino acids with chemically reactive side chains may be positioned in a variety of positions on the CD1d and β 2-microglobulin polypeptides, preferably distal to the site(s) where β 2-microglobulin and CD1d interact. Suitable side chains can be used to chemically link two or more assembled CD1d complexes to a suitable dendrimer particle. Dendrimers are synthetic chemical polymers that can have any one of a number of different functional groups on their surface (D. Tomalia, *Aldrichimica Acta* 26:91:101 (1993)). Exemplary dendrimers for use in accordance with the present invention include *e.g.* E9 starburst polyamine dendrimer and E9 combburst polyamine dendrimer, which can link cysteine residues. The CD1d and/or β 2-microglobulin polypeptides are modified to introduce a cysteine residue at the carboxyl terminus. Following synthesis in eukaryotic cells, a complete cysteine modified CD1d complex is assembled *in vitro*. Cysteine modified CD1d and/or β 2-microglobulin polypeptides will react with the maleimide groups on the various peptide backbones with either two, three, or four modified lysine residues for formation of CD1d dimers, trimers, and tetramers.

[0076] Cochran, J.R. *et al*., *Immunity* 12:241-50 (2000) describe the use of chemically synthesized peptide-based cross-linking reagents in which two or more thiol-reactive maleimide groups are linked to lysine side chains in a flexible peptide of 8 to 19 residues containing glycine, serine, and glutamic acid in addition to the modified lysine residues. Isolated CD1d and/or β 2-microglobulin polypeptides are modified to introduce a cysteine residue at the carboxyl terminus. Cysteine modified CD1d and/or β 2-microglobulin polypeptides react with the maleimide groups on the various peptide backbones with either two, three, or four modified lysine residues for formation of dimers, trimers, and tetramers.

[0077] Another means of assembling polymeric CD1d complexes is to exploit the observation that defined amino acid substitutions in the GCN4 leucine zipper dimerization domain results in formation of highly stable trimeric and tetrameric

- 26 -

structures of the synthetic peptide (Harbury, P.B. *et al.*, *Science* 262:1401-7 (1993)). For example, multivalent CD1d complexes are constructed by attaching a modified GCN4-zipper to the carboxyl terminus of soluble CD1d or β 2-microglobulin polypeptides. Tetravalent CD1d complexes could be assembled from a mixture of CD1d complexes each separately fused to a modified GCN4-zipper motif.

[0078] The attachment site(s) on a soluble CD1d complex for binding to a multivalent compound may be naturally occurring, or may be introduced through genetic engineering. The site will be a specific binding pair member or one that is modified to provide a specific binding pair member, where the complementary pair has a multiplicity of specific binding sites. Binding to the complementary binding member can be a chemical reaction, epitope-receptor binding or hapten-receptor binding where a hapten is linked to the subunit chain.

[0079] In a preferred embodiment, the CD1d and/or β 2 microglobulin contain an amino acid sequence which is a recognition site for a modifying enzyme. Modifying enzymes include BirA, various glycosylases, farnesyl protein transferase, and protein kinases. The group introduced by the modifying enzyme, *e.g.* biotin, sugar, phosphate, farnesyl, etc. provides a complementary binding pair member, or a unique site for further modification, such as chemical cross-linking, biotinylation, etc. that will provide a complementary binding pair member.

[0080] For example, the CD1d molecule may be engineered to contain a site for biotinylation, for example a BirA-dependent site. The multivalent compound can be avidin or can be linked to avidin either directly or indirectly.

[0081] Both the soluble CD1d and β 2-microglobulin polypeptides useful in the present invention may be autologous to any mammalian or avian species, for example, primates (*esp.* humans), rodents, rabbits, equines, bovines, canines, felines, etc. β 2-microglobulin is typically not inflammatory *in vivo*. However, it is preferable to employ β 2-microglobulin derived from the same species as is to be treated so as to reduce the risk of a xenogeneic immune response.

Soluble CD1d Polypeptides

- [0082] In certain embodiments, the non-specific CD1d complex comprises soluble CD1d polypeptides and polypeptide fragments, which associates with β_2 -microglobulin and binds antigen, *e.g.*, ceramide-like glycolipid. The CD1d molecule is a member of the family of major histocompatibility complex (MH+HC) antigen-like glycoproteins which associate with β_2 -microglobulin and are expressed at the surface of cortical thymocytes, B cells, dendritic cells, Langerhans cells in the skin, and gastrointestinal epithelial cells. CD1d is mainly expressed on dendritic cells or epithelial cells of the gastrointestinal tract. The CD1 family members are involved in the presentation of glycolipids as antigens. In particular, CD1d regulates cytokine tone through activation of a distinct subset of T-lymphocytes, namely NK1 T cells which secrete IL-4 and INF- γ . All of the CD1 glycoproteins have been cloned and analyzed. For a detailed discussion of CD1 glycoproteins, and in particular CD1d, *see, e.g.*, Balk *et al.*, *Proc. Natl. Acad. Sci. USA* 86:252-256 (1989); Kojo *et al.*, *Biochem. Biophys. Res. Comm.* 276:107-111 (2000); Kojo *et al.*, *J. Rheumatology* 30:2524-2528 (2003); Kang and Cresswell, *Nature Immunology* 5:175-181 (2004); Im *et al.*, *J. Biol. Chem.* 279:299-310 (2004); Dutronc and Porcelli, *Tissue Antigens* 60:337-353 (2002) which are incorporated by reference herein in their entirety.

Domains of CD1d

- [0083] Full-length CD1d consists of a signal sequence, an extracellular domain, a transmembrane domain and a cytoplasmic domain. The full-length CD1d polypeptide is 335 amino acids in length.
- [0084] The following polypeptide sequence was reported as the human CD1d sequence and has the accession number NP_001757 in Genbank.

- [0085] Full-Length Human CD1d (SEQ ID NO:1):

MGCLLFLLW ALLQAWGSAE VPQRLFPLRC LQISSFANSS WTRTDGLAWL
 GELQTHSWSN DSDTVRSLKP WSQGTFSDDQ WETLQHIFRV YRSSFTRDVK
 EFAKMLRLSY PLELQVSAGC EVHPGNASNN FFHVAFQGKD ILSFQGTSWE
 PTQEAPLWVN LAIQVLNQDK WTRETVQWLL NGTCPQFVSG LLESGKSELK
 KQVKPKAWLS RGPSPGPGRLLLVCHVSGFY PKPVWVKWMR

- 28 -

GEQEQQGTQP GDILPNADET WYLRATLDVV AGEAAGLSCR VKHSSLEGQD
IVLYWGGSYTSMGLIALAVL ACLLFLLIVG FTSRFKRQTS YQGVV

[0086] A variant of human CD1d includes, but is not limited to, a polypeptide with the following mutation: T64S.

[0087] The sequence of mouse CD1d can be found on Genbank with the following accession number: NP_031665. The sequence of rat CD1d can be found on Genbank with the following accession number: NP_058775. The sequence of sheep CD1d can be found on Genbank with the following accession numbers: O62848 and Q29422. The sequence of chimpanzee CD1d can be found on Genbank with the following accession number: NP_001065272. The sequence of rabbit CD1d can be found on Genbank with the following accession number: P23043. All of the above Genbank accession numbers are incorporated herein by reference.

[0088] The accession number was reported as the mouse CD1d: NP_031665 in Genbank.

[0089] The extracellular domain of CD1d consists of three domains: the $\alpha 1$ domain, the $\alpha 2$ domain, and the $\alpha 3$ domain. The $\alpha 1$ and $\alpha 2$ domains comprise the antigen binding sites. The $\alpha 3$ domain includes a β_2 -microglobulin association site.

[0090] The CD1d domain designations used herein are defined as follows:

Table 1. CD1d domains

Domain	CD1d (human)
Signal Seq.	1-19
Extracellular	20-301
$\alpha 1$ domain	20-108
$\alpha 2$ domain	109-201
$\alpha 3$ domain	202-295
Transmembrane	302-322
Cytoplasmic	323-335

- [0091] As one of skill in the art will appreciate, the beginning and ending residues of the domains listed above may vary depending upon the computer modeling program used or the method used for determining the domain.
- [0092] Some embodiments of the invention provide a CD1d complex, *e.g.*, a non-specific CD1d complex, which comprises a soluble CD1d polypeptide or polypeptide fragment. Specifically, soluble CD1d polypeptides of the present invention include fragments, variants, or derivative thereof of a soluble CD1d polypeptide. Table 1 above describes the various domains of the CD1d polypeptide. Soluble CD1d polypeptides of the invention generally comprise a portion or all of the extracellular domain of the polypeptides, including the $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains. Soluble CD1d polypeptides generally lack some or all of the transmembrane domain and cytoplasmic domain. As one of skill in the art would appreciate, the entire extracellular domain of CD1d may comprise additional or fewer amino acids on either the C-terminal or N-terminal end of the extracellular domain polypeptide.
- [0093] Soluble human CD1d polypeptides for use in the methods of the present invention include, but are not limited to, a soluble CD1d polypeptide comprising, consisting essentially of, or consisting of an amino acid sequence identical to a reference amino acid sequence, except for up to twenty amino acid substitutions, wherein said reference amino acid sequence is selected from the group consisting of amino acids a to 295 of SEQ ID NO:1, amino acids 21 to b of SEQ ID NO:1, and a to b of SEQ ID NO:1, wherein a is any integer from 1 to 100, and b is any integer from 201 to 301, and wherein said soluble CD1d polypeptide associates with β_2 -microglobulin and binds a ceramide-like glycolipid antigen. In one embodiment, the soluble CD1d polypeptide comprises amino acids 21 to 295 of SEQ ID NO:1. In another embodiment, the soluble CD1d polypeptide comprises amino acids 20-295, 20-296, 20-297, 20-298, 20-299, 20-300 and 20 to 301 of SEQ ID NO:1.
- [0094] By "a reference amino acid sequence" is meant the specified sequence without the introduction of any amino acid substitutions. As one of ordinary skill in the art would understand, if there are no substitutions, the "isolated polypeptide" of the invention comprises an amino acid sequence which is identical to the reference amino acid sequence.

- 30 -

- [0095] Soluble CD1d polypeptides described herein may have various alterations such as substitutions, insertions or deletions. Exemplary amino acids that can be substituted in the polypeptide include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).
- [0096] Corresponding fragments of soluble CD1d polypeptides at least 70%, 75%, 80%, 85%, 90%, or 95% identical to the polypeptides and reference polypeptides described herein are also contemplated.
- [0097] As known in the art, "sequence identity" between two polypeptides is determined by comparing the amino acid sequence of one polypeptide to the sequence of a second polypeptide. When discussed herein, whether any particular polypeptide is at least about 70%, 75%, 80%, 85%, 90% or 95% identical to another polypeptide can be determined using methods and computer programs/software known in the art such as, but not limited to, the BESTFIT program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). BESTFIT uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981), to find the best segment of homology between two sequences. When using BESTFIT or any other sequence alignment program to determine whether a particular sequence is, for example, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference polypeptide sequence and that gaps in homology of up to 5% of the total number of amino acids in the reference sequence are allowed.
- [0098] In methods of the present invention, a soluble CD1d polypeptide or polypeptide fragment of the invention may be administered directly as a preformed polypeptide. In certain embodiments, however, the soluble CD1d polypeptide or

- 31 -

fragment thereof is associated with β_2 -microglobulin, and is bound to a ceramide-like glycolipid antigen.

β_2 -microglobulin polypeptides

[0099] In certain embodiments, a CD1d complex of the invention comprises a β_2 -microglobulin polypeptide, which associates with a soluble CD1d polypeptide or polypeptide fragment. β_2 -microglobulin is present on the surface of all nucleated cells as the small extracellular subunit of the major histocompatibility complex (MHC) class I molecule and actively participates in the immune response. For a detailed discussion of β_2 -microglobulin, *see, e.g.*, Peterson *et al.*, *Adv. Cancer Res.* 24:115-163 (1977); Sege *et al.*, *Biochemistry* 20:4523-4530 (1981); which are incorporated by reference herein in their entirety.

β_2 -microglobulin domains

[0100] Full-length β_2 -microglobulin is a secreted protein which comprises a signal sequence and Ig-like domain. The full-length CD1d polypeptide is 119 amino acids in length.

[0101] The following polypeptide sequence was reported as the human β_2 -microglobulin sequence and has the accession number NP_004039 in Genbank.

[0102] Full-Length Human β_2 -microglobulin (SEQ ID NO:2):

MSRSVALAVL ALLSLSGLEA IQRTPKIQVY SRHPAENGKS
NFLNCYVSGF HPSDIEVDLL KNGERIEKVE HSDLSFSKDW
SFYLLYYTEF TPTEKDEYAC RVNHVTLSP KIVKWDRDM

[0103] Variants of human β_2 -microglobulin include, but are not limited to, polypeptides with one or more of the following mutations: A20G, P52Q, S55V, and Y86YS.

[0104] The sequence of mouse β_2 -microglobulin can be found on Genbank with the following accession number: NP_033865. The sequence of pig β_2 -microglobulin can be found on Genbank with the following accession number: NP_999143. The sequence of rat β_2 -microglobulin can be found on Genbank with the following accession number: NP_036644. The sequence of chimpanzee β_2 -microglobulin can

- 32 -

be found on Genbank with the following accession number: NP_001009066. The sequence of rabbit β_2 -microglobulin can be found on Genbank with the following accession number: P01885. The sequence of sheep β_2 -microglobulin can be found on Genbank with the following accession number: NP_001009284.

[0105] The β_2 -microglobulin domain designations used herein are defined as in Table 2:

Table 2. β_2 -microglobulin domains

Domain	β_2 -microglobulin (human)
Signal Seq.	1-20
β_2 -microglobulin	21-119
Ig domain	25-113 or 22-116

[0106] As one of skill in the art will appreciate, the beginning and ending residues of the domains listed above may vary depending upon the computer modeling program used or the method used for determining the domain.

[0107] Some embodiments of the invention provide a CD1d complex, *e.g.*, a non-specific CD1d complex, which comprises a β_2 -microglobulin polypeptide or polypeptide fragment. β_2 -microglobulin polypeptides of the present invention include fragments, variants, or derivative thereof of a β_2 -microglobulin polypeptide. Table 2 above describes the various domains of the β_2 -microglobulin polypeptide. β_2 -microglobulin polypeptides of the invention generally comprise a portion or all of the secreted portion of the polypeptides.

[0108] Human β_2 -microglobulin polypeptides for use in the methods of the present invention include, but are not limited to, a β_2 -microglobulin polypeptide comprising, consisting essentially of, or consisting of an amino acid sequence identical to a reference amino acid sequence, except for up to twenty amino acid substitutions, wherein said reference amino acid sequence is selected from the group consisting of amino acids a to 119 of SEQ ID NO:2, amino acids 21 to b of SEQ ID NO: 2, and a to b of SEQ ID NO:2, wherein a is any integer from 15 to 25, and b is any integer from 100 to 119, wherein said β_2 -microglobulin polypeptide associates with CD1d and

- 33 -

supports binding of ceramide-like glycolipid antigens. In one embodiment, the β_2 -microglobulin polypeptide comprises amino acids 21 to 113 of SEQ ID NO:2. In one embodiment, the β_2 -microglobulin polypeptide comprises amino acids 21 to 119 of SEQ ID NO:2.

[0109] By "a reference amino acid sequence" is meant the specified sequence without the introduction of any amino acid substitutions. As one of ordinary skill in the art would understand, if there are no substitutions, the "isolated polypeptide" of the invention comprises an amino acid sequence which is identical to the reference amino acid sequence.

[0110] β_2 -microglobulin polypeptides described herein may have various alterations such as substitutions, insertions or deletions. Exemplary amino acids that can be substituted in the polypeptide include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

[0111] Corresponding fragments of β_2 -microglobulin polypeptides at least 70%, 75%, 80%, 85%, 90%, or 95% identical to the polypeptides and reference polypeptides described herein are also contemplated.

[0112] In methods of the present invention, a β_2 -microglobulin polypeptide or polypeptide fragment of the invention is typically administered directly as a preformed polypeptide. In certain embodiments, the β_2 -microglobulin polypeptide or fragment thereof is associated with a soluble CD1d polypeptide.

[0113] A soluble CD1d polypeptide may contain some or all of the amino acids from the transmembrane domain, provided that the polypeptide is still capable of remaining soluble in an aqueous, *e.g.*, a physiological solution. Preferably, not more than about 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1, and preferably none of the amino acids of the transmembrane domain will be included.

- 34 -

- [0114] Additionally, fragments of β_2 -microglobulin are useful in the present invention. To be useful in the present invention, the fragment of β_2 -microglobulin would have to retain the ability to associate with the CD1d molecule. Preferably, not more than about 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1, and preferably none of the amino acids of β_2 -microglobulin will be deleted.
- [0115] One may wish to introduce a small number of amino acids at the polypeptide termini of either the soluble CD1d polypeptide or the β_2 -microglobulin polypeptide, usually not more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1. The deletion or insertion of amino acids will usually be as a result of the needs of the construction, providing for convenient restriction sites, addition of processing signals, ease of manipulation, improvement in levels of expression, or the like. In addition, one may wish to substitute one or more amino acids with a different amino acid for similar reasons, usually not substituting more than about 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acids in any one domain.
- [0116] The soluble CD1d polypeptide and β_2 -microglobulin polypeptide may be separately produced and allowed to associate to form a stable heteroduplex complex, or both of the subunits may be expressed in a single cell.
- [0117] Soluble CD1d polypeptides and β_2 -microglobulin polypeptides for use in the methods and compositions of the present invention may be isolated from a multiplicity of cells, *e.g.*, transformed cell lines JY, BM92, WIN, MOC, and MG, and CHO using a variety of techniques known to those skilled in the art.
- [0118] Additionally, the amino acid sequences of CD1d and β_2 -microglobulin from a variety of species are known, and the polynucleotides encoding these polypeptides have been cloned, therefore, the polypeptides can be made using recombinant methods. The coding regions for the CD1d and β_2 microglobulin chains or their fusion products are inserted into expression vectors, expressed separately in an appropriate host, such as *E. coli*, yeast, insect cells, mammalian cells or other suitable cells, and the recombinant proteins obtained are recombined in the presence of a ceramide like glycolipid antigen (*e.g.* α -GalCer).

Fusion Proteins, Modified Proteins and Conjugated Polypeptides

- [0119] Some embodiments of the invention involve the use of a soluble CD1d polypeptide and/or β_2 -microglobulin polypeptide fused to a heterologous polypeptide moiety to form a fusion protein. Such fusion proteins can be used to accomplish various objectives, *e.g.*, increased serum half-life, improved bioavailability, *in vivo* targeting to a specific organ or tissue type, improved recombinant expression efficiency, improved host cell secretion, ease of purification, and higher avidity. Depending on the objective(s) to be achieved, the heterologous moiety can be inert or biologically active. Also, it can be chosen to be stably fused to the soluble CD1d polypeptide or β_2 -microglobulin polypeptide of the invention or to be cleavable, *in vitro* or *in vivo*. Heterologous moieties to accomplish these other objectives are known in the art.
- [0120] As an alternative to expression of a fusion protein, a chosen heterologous moiety can be preformed and chemically conjugated to the soluble CD1d polypeptide or β_2 -microglobulin polypeptide of the invention. In most cases, a chosen heterologous moiety will function similarly, whether fused or conjugated to the soluble CD1d polypeptide or β_2 -microglobulin polypeptide. Therefore, in the following discussion of heterologous amino acid sequences, unless otherwise noted, it is to be understood that the heterologous sequence can be joined to the soluble CD1d polypeptide or β_2 -microglobulin polypeptide in the form of a fusion protein or as a chemical conjugate.
- [0121] Soluble CD1d polypeptides or β_2 -microglobulin polypeptides for use in the treatment methods disclosed herein include derivatives that are modified, *i.e.*, by the covalent attachment of any type of molecule such that covalent attachment does not prevent the soluble CD1d polypeptide and β_2 -microglobulin polypeptide from associating and binding antigen to form a CD1d complex, *e.g.*, a non-specific CD1d complex. For example, but not by way of limitation, the soluble CD1d polypeptides and/or β_2 -microglobulin polypeptides of the present invention may be modified *e.g.*, by glycosylation, acetylation, pegylation, phosphylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be

- 36 -

carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[0122] Soluble CD1d polypeptides or β_2 -microglobulin polypeptides for use in the treatment methods disclosed herein can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, *i.e.*, peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. Soluble CD1d polypeptides and/or β_2 -microglobulin polypeptides may be modified by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in the soluble CD1d polypeptides or β_2 -microglobulin polypeptide including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini, or on moieties such as carbohydrates. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given soluble CD1d polypeptides or β_2 -microglobulin polypeptide. Also, a given soluble CD1d polypeptides or β_2 -microglobulin polypeptide may contain many types of modifications. Soluble CD1d polypeptides or β_2 -microglobulin polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic soluble CD1d polypeptides or β_2 -microglobulin polypeptides may result from posttranslational natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA

- 37 -

mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, *Proteins - Structure And Molecular Properties*, T. E. Creighton, W. H. Freeman and Company, New York 2nd Ed., (1993); *Posttranslational Covalent Modification Of Proteins*, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter *et al.*, *Meth Enzymol* 182:626-646 (1990); Rattan *et al.*, *Ann NY Acad Sci* 663:48-62 (1992)).

[0123] A heterologous polypeptide to which the soluble CD1d polypeptides or β_2 -microglobulin polypeptide is fused may be useful therapeutically or useful to target the soluble CD1d polypeptides and/or β_2 -microglobulin polypeptide. Soluble CD1d fusion polypeptides and/or β_2 -microglobulin fusion proteins can be used to accomplish various objectives, *e.g.*, increased serum half-life, improved bioavailability, *in vivo* targeting to a specific organ or tissue type, improved recombinant expression efficiency, improved host cell secretion, ease of purification, and higher avidity. Depending on the objective(s) to be achieved, the heterologous moiety can be inert or biologically active. Also, it can be chosen to be stably fused to the soluble CD1d polypeptides or β_2 -microglobulin polypeptide or to be cleavable, *in vitro* or *in vivo*. Heterologous moieties to accomplish these various objectives are known in the art.

[0124] Various heterologous amino acid sequences, *i.e.*, polypeptide moieties or "carriers," for increasing the *in vivo* stability, *i.e.*, serum half-life, of therapeutic polypeptides are known. Examples include serum albumins such as, *e.g.*, bovine serum albumin (BSA) or human serum albumin (HSA).

[0125] Due to its long half-life, wide *in vivo* distribution, and lack of enzymatic or immunological function, essentially full-length human serum albumin (HSA), or an HSA fragment, is commonly used as a heterologous moiety. Through application of methods and materials such as those taught in Yeh *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:1904-08 (1992) and Syed *et al.*, *Blood* 89:3243-52 (1997), HSA can be used to form a fusion protein or polypeptide conjugate that displays pharmacological activity of a CD1d complex, *e.g.*, a non-specific CD1d complex of the invention, while displaying significantly increased *in vivo* stability, *e.g.*, 10-fold to 100-fold higher. The C-terminus of the HSA can be fused to the N-terminus of the soluble CD1d

- 38 -

polypeptides or β_2 -microglobulin polypeptide moiety. Since HSA is a naturally secreted protein, the HSA signal sequence can be exploited to obtain secretion of the fusion protein into the cell culture medium when the fusion protein is produced in a eukaryotic, *e.g.*, mammalian, expression system.

[0126] Some embodiments of the invention employ a soluble CD1d polypeptide or β_2 -microglobulin polypeptide moiety fused to a hinge and Fc region, *i.e.*, the C-terminal portion of an immunoglobulin heavy chain constant region.

[0127] Potential advantages of a soluble CD1d polypeptide-Fc fusion or a β_2 -microglobulin polypeptide-Fc fusion include solubility, *in vivo* stability, and multivalency, *e.g.*, dimerization. The Fc region used can be an IgA, IgD, or IgG Fc region (hinge-CH2-CH3). Alternatively, it can be an IgE or IgM Fc region (hinge-CH2-CH3-CH4). An IgG Fc region is generally used, *e.g.*, an IgG1 Fc region or IgG4 Fc region. Materials and methods for constructing and expressing DNA encoding Fc fusions are known in the art and can be applied to obtain fusions without undue experimentation. Some embodiments of the invention employ a fusion protein such as those described in Capon *et al.*, U.S. Patent Nos. 5,428,130 and 5,565,335.

[0128] Fully intact, wild-type Fc regions display effector functions that normally are unnecessary and undesired in an Fc fusion protein used in the methods of the present invention. Therefore, certain binding sites typically are deleted from the Fc region during the construction of the secretion cassette. For example, since coexpression with the light chain is unnecessary, the binding site for the heavy chain binding protein, Bip (Hendershot *et al.*, *Immunol. Today* 8:111-14 (1987)), is deleted from the CH2 domain of the Fc region of IgE, such that this site does not interfere with the efficient secretion of the immunofusion. Transmembrane domain sequences, such as those present in IgM, also are generally deleted.

[0129] The IgG1 Fc region is most often used. Alternatively, the Fc region of the other subclasses of immunoglobulin gamma (gamma-2, gamma-3 and gamma-4) can be used in the secretion cassette. The IgG1 Fc region of immunoglobulin gamma-1 is generally used in the secretion cassette and includes at least part of the hinge region, the CH2 region, and the CH3 region. In some embodiments, the Fc region of immunoglobulin gamma-1 is a CH2-deleted-Fc, which includes part of the hinge

- 39 -

region and the CH3 region, but not the CH2 region. A CH2-deleted-Fc has been described by Gillies *et al.*, *Hum. Antibod. Hybridomas* 1:47 (1990). In some embodiments, the Fc region of one of IgA, IgD, IgE, or IgM, is used.

[0130] Soluble CD1d polypeptide-Fc fusion proteins or β_2 -microglobulin-polypeptide-moiety-Fc fusion proteins can be constructed in several different configurations. In one configuration the C-terminus of the soluble CD1d or β_2 -microglobulin polypeptide moiety is fused directly to the N-terminus of the Fc hinge moiety. In a slightly different configuration, a short polypeptide, *e.g.*, 2-10 amino acids, is incorporated into the fusion between the N-terminus of the soluble CD1d or β_2 -microglobulin polypeptide moiety and the C-terminus of the Fc moiety. In the alternative configuration, the short polypeptide is incorporated into the fusion between the C-terminus of the soluble CD1d or β_2 -microglobulin polypeptide moiety and the N-terminus of the Fc moiety. Such a linker provides conformational flexibility, which may improve biological activity in some circumstances. If a sufficient portion of the hinge region is retained in the Fc moiety, the soluble CD1d polypeptide-Fc fusion or β_2 -microglobulin-polypeptide-moiety-Fc fusion will dimerize, thus forming a divalent molecule. A homogeneous population of monomeric Fc fusions will yield monospecific, bivalent dimers. A mixture of two monomeric Fc fusions each having a different specificity will yield bispecific, bivalent dimers.

[0131] Soluble CD1d or β_2 -microglobulin polypeptides of the invention can be fused to a polypeptide tag. The term "polypeptide tag," as used herein, is intended to mean any sequence of amino acids that can be attached to, connected to, or linked to a soluble CD1d or β_2 -microglobulin polypeptide and that can be used to identify, purify, concentrate or isolate the soluble CD1d or β_2 -microglobulin polypeptide. The attachment of the polypeptide tag to the soluble CD1d or β_2 -microglobulin polypeptide may occur, *e.g.*, by constructing a nucleic acid molecule that comprises: (a) a nucleic acid sequence that encodes the polypeptide tag, and (b) a nucleic acid sequence that encodes a soluble CD1d or β_2 -microglobulin polypeptide. Exemplary polypeptide tags include, *e.g.*, amino acid sequences that are capable of being post-translationally modified, *e.g.*, amino acid sequences that are biotinylated. Other

- 40 -

exemplary polypeptide tags include, *e.g.*, amino acid sequences that are capable of being recognized and/or bound by an antibody (or fragment thereof) or other specific binding reagent. Polypeptide tags that are capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent include, *e.g.*, those that are known in the art as "epitope tags." An epitope tag may be a natural or an artificial epitope tag. Natural and artificial epitope tags are known in the art, including, *e.g.*, artificial epitopes such as FLAG, Strep, or poly-histidine peptides. FLAG peptides include the sequence Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (SEQ ID NO:5) or Asp-Tyr-Lys-Asp-Glu-Asp-Asp-Lys (SEQ ID NO:6) (Einhauer, A. and Jungbauer, A., *J. Biochem. Biophys. Methods* 49:1-3:455-465 (2001)). The Strep epitope has the sequence Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO:7). The VSV-G epitope can also be used and has the sequence Tyr-Thr-Asp-Ile-Glu-Met-Asn-Arg-Leu-Gly-Lys (SEQ ID NO:8). Another artificial epitope is a poly-His sequence having six histidine residues (His-His-His-His-His-His (SEQ ID NO:9)). Naturally-occurring epitopes include the influenza virus hemagglutinin (HA) sequence Tyr-Pro-Tyr-Asp-Val-Pro-Asp-Tyr-Ala-Ile-Glu-Gly-Arg (SEQ ID NO:10) recognized by the monoclonal antibody 12CA5 (Murray *et al.*, *Anal. Biochem.* 229:170-179 (1995)) and the eleven amino acid sequence from human c-myc (Myc) recognized by the monoclonal antibody 9E10 (Glu-Gln-Lys-Leu-Leu-Ser-Glu-Glu-Asp-Leu-Asn (SEQ ID NO:11) (Manstein *et al.*, *Gene* 162:129-134 (1995)). Another useful epitope is the tripeptide Glu-Glu-Phe which is recognized by the monoclonal antibody YL 1/2. (Stammers *et al.* *FEBS Lett.* 283:298-302(1991)).

- [0132] In certain embodiments, the soluble CD1d or β_2 -microglobulin polypeptide and the polypeptide tag may be connected via a linking amino acid sequence. As used herein, a "linking amino acid sequence" may be an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases. Amino acid sequences that can be recognized and/or cleaved by one or more proteases are known in the art. Exemplary amino acid sequences include, but are not limited to, those that are recognized by the following proteases: factor VIIa, factor IXa, factor Xa, APC, t-PA, u-PA, trypsin, chymotrypsin, enterokinase, pepsin, cathepsin B,H,L,S,D, cathepsin G, renin, angiotensin converting enzyme, matrix metalloproteases

(collagenases, stromelysins, gelatinases), macrophage elastase, Cir, and Cis. The amino acid sequences that are recognized by the aforementioned proteases are known in the art. Exemplary sequences recognized by certain proteases can be found, *e.g.*, in U.S. Patent No. 5,811,252.

[0133] Polypeptide tags can facilitate purification using commercially available chromatography media.

[0134] By fusing a soluble CD1d or β_2 -microglobulin polypeptide moiety at the amino and carboxy termini of a suitable fusion partner, bivalent or tetravalent forms of a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention can be obtained. For example, a soluble CD1d or β_2 -microglobulin polypeptide moiety can be fused to the amino and carboxy termini of an Ig moiety to produce a bivalent monomeric polypeptide containing two soluble CD1d or β_2 -microglobulin polypeptide moieties. Upon dimerization of two of these monomers, by virtue of the Ig moiety, a tetravalent form of a soluble CD1d or β_2 -microglobulin polypeptide is obtained. Such multivalent forms can be used to achieve increased binding affinity for the target. Multivalent forms of a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention also can be obtained by placing soluble CD1d or β_2 -microglobulin polypeptide moieties in tandem to form concatamers, which can be employed alone or fused to a fusion partner such as Ig or HSA.

Conjugated Polymers (other than polypeptides)

[0135] Some embodiments of the invention involve a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention wherein one or more polymers are conjugated (covalently linked) to the soluble CD1d or β_2 -microglobulin polypeptide. Examples of polymers suitable for such conjugation include polypeptides (discussed above), sugar polymers and polyalkylene glycol chains. Typically, but not necessarily, a polymer is conjugated to the soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention for the purpose of improving one or more of the following: solubility, stability, or bioavailability.

- [0136] The class of polymer generally used for conjugation to a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention is a polyalkylene glycol. Polyethylene glycol (PEG) is most frequently used. PEG moieties, *e.g.*, 1, 2, 3, 4 or 5 PEG polymers, can be conjugated to each soluble CD1d or β_2 -microglobulin polypeptide to increase serum half life, as compared to the soluble CD1d or β_2 -microglobulin polypeptide alone. PEG moieties are non-antigenic and essentially biologically inert. PEG moieties used in the practice of the invention may be branched or unbranched.
- [0137] The number of PEG moieties attached to the soluble CD1d or β_2 -microglobulin polypeptide and the molecular weight of the individual PEG chains can vary. In general, the higher the molecular weight of the polymer, the fewer polymer chains attached to the polypeptide. Usually, the total polymer mass attached to a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment is from 20 kDa to 40 kDa. Thus, if one polymer chain is attached, the molecular weight of the chain is generally 20-40 kDa. If two chains are attached, the molecular weight of each chain is generally 10-20 kDa. If three chains are attached, the molecular weight is generally 7-14 kDa.
- [0138] The polymer, *e.g.*, PEG, can be linked to the soluble CD1d or β_2 -microglobulin polypeptide through any suitable, exposed reactive group on the polypeptide. The exposed reactive group(s) can be, *e.g.*, an N-terminal amino group or the epsilon amino group of an internal lysine residue, or both. An activated polymer can react and covalently link at any free amino group on the soluble CD1d or β_2 -microglobulin polypeptide. Free carboxylic groups, suitably activated carbonyl groups, hydroxyl, guanidyl, imidazole, oxidized carbohydrate moieties and mercapto groups of the soluble CD1d or β_2 -microglobulin polypeptide (if available) also can be used as reactive groups for polymer attachment.
- [0139] In a conjugation reaction, from about 1.0 to about 10 moles of activated polymer per mole of polypeptide, depending on polypeptide concentration, is typically employed. Usually, the ratio chosen represents a balance between maximizing the reaction while minimizing side reactions (often non-specific) that can impair the desired pharmacological activity of the soluble CD1d or β_2 -microglobulin

- 43 -

polypeptide moiety. Preferably, at least 50% of the biological activity (as demonstrated, *e.g.*, in any of the assays described herein or known in the art) of the soluble CD1d or β_2 -microglobulin polypeptide is retained, and most preferably nearly 100% is retained.

[0140] The polymer can be conjugated to the soluble CD1d or β_2 -microglobulin polypeptide using conventional chemistry. For example, a polyalkylene glycol moiety can be coupled to a lysine epsilon amino group of the soluble CD1d or β_2 -microglobulin polypeptide. Linkage to the lysine side chain can be performed with an N-hydroxysuccinimide (NHS) active ester such as PEG succinimidyl succinate (SS-PEG) and succinimidyl propionate (SPA-PEG). Suitable polyalkylene glycol moieties include, *e.g.*, carboxymethyl-NHS and norleucine-NHS, SC. These reagents are commercially available. Additional amine-reactive PEG linkers can be substituted for the succinimidyl moiety. These include, *e.g.*, isothiocyanates, nitrophenylcarbonates (PNP), epoxides, benzotriazole carbonates, SC-PEG, tresylate, aldehyde, epoxide, carbonylimidazole and PNP carbonate. Conditions are usually optimized to maximize the selectivity and extent of reaction. Such optimization of reaction conditions is within ordinary skill in the art.

[0141] PEGylation can be carried out by any of the PEGylation reactions known in the art. *See, e.g., Focus on Growth Factors*, 3: 4-10, 1992 and European patent applications EP 0 154 316 and EP 0 401 384. PEGylation may be carried out using an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer).

[0142] PEGylation by acylation generally involves reacting an active ester derivative of polyethylene glycol. Any reactive PEG molecule can be employed in the PEGylation. PEG esterified to N-hydroxysuccinimide (NHS) is a frequently used activated PEG ester. As used herein, "acylation" includes without limitation the following types of linkages between the therapeutic protein and a water-soluble polymer such as PEG: amide, carbamate, urethane, and the like. *See, e.g., Bioconjugate Chem.* 5: 133-140, 1994. Reaction parameters are generally selected to avoid temperature, solvent, and pH conditions that would damage or inactivate the soluble CD1d or β_2 -microglobulin polypeptide.

- 44 -

- [0143] Generally, the connecting linkage is an amide and typically at least 95% of the resulting product is mono-, di- or tri-PEGylated. However, some species with higher degrees of PEGylation may be formed in amounts depending on the specific reaction conditions used. Optionally, purified PEGylated species are separated from the mixture, particularly unreacted species, by conventional purification methods, including, *e.g.*, dialysis, salting-out, ultrafiltration, ion-exchange chromatography, gel filtration chromatography, hydrophobic exchange chromatography, and electrophoresis.
- [0144] PEGylation by alkylation generally involves reacting a terminal aldehyde derivative of PEG with a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention in the presence of a reducing agent. In addition, one can manipulate the reaction conditions to favor PEGylation substantially only at the N-terminal amino group of the soluble CD1d or β_2 -microglobulin polypeptide, *i.e.* a mono-PEGylated protein. In either case of mono-PEGylation or poly-PEGylation, the PEG groups are typically attached to the protein via a -CH₂-NH- group. With particular reference to the -CH₂- group, this type of linkage is known as an "alkyl" linkage.
- [0145] Derivatization via reductive alkylation to produce an N-terminally targeted mono-PEGylated product exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization. The reaction is performed at a pH that allows one to take advantage of the pK_a differences between the epsilon-amino groups of the lysine residues and that of the N-terminal amino group of the protein. By such selective derivatization, attachment of a water-soluble polymer that contains a reactive group, such as an aldehyde, to a protein is controlled: the conjugation with the polymer takes place predominantly at the N-terminus of the protein and no significant modification of other reactive groups, such as the lysine side chain amino groups, occurs.
- [0146] The polymer molecules used in both the acylation and alkylation approaches are selected from among water-soluble polymers. The polymer selected is typically modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the degree of polymerization may be controlled as

provided for in the present methods. An exemplary reactive PEG aldehyde is polyethylene glycol propionaldehyde, which is water stable, or mono C1-C10 alkoxy or aryloxy derivatives thereof (*see, e.g.*, Harris *et al.*, U.S. Pat. No. 5,252,714). The polymer may be branched or unbranched. For the acylation reactions, the polymer(s) selected typically have a single reactive ester group. For reductive alkylation, the polymer(s) selected typically have a single reactive aldehyde group. Generally, the water-soluble polymer will not be selected from naturally occurring glycosyl residues, because these are usually made more conveniently by mammalian recombinant expression systems.

[0147] Methods for preparing a PEGylated soluble CD1d or β_2 -microglobulin polypeptide of the invention generally include the steps of (a) reacting a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the molecule becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined case-by-case based on known parameters and the desired result. For example, a larger the ratio of PEG to protein, generally leads to a greater the percentage of poly-PEGylated product.

[0148] Reductive alkylation to produce a substantially homogeneous population of mono-polymer/soluble CD1d or β_2 -microglobulin polypeptide generally includes the steps of: (a) reacting a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention with a reactive PEG molecule under reductive alkylation conditions, at a pH suitable to permit selective modification of the N-terminal amino group of soluble CD1d or β_2 -microglobulin; and (b) obtaining the reaction product(s).

[0149] For a substantially homogeneous population of mono-polymer/soluble CD1d or β_2 -microglobulin polypeptide, the reductive alkylation reaction conditions are those that permit the selective attachment of the water-soluble polymer moiety to the N-terminus of a soluble CD1d or β_2 -microglobulin polypeptide or polypeptide fragment of the invention. Such reaction conditions generally provide for pKa differences between the lysine side chain amino groups and the N-terminal amino group. For

- 46 -

purposes of the present invention, the pH is generally in the range of 3-9, typically 3-6.

[0150] In some embodiments, the polyalkylene glycol moiety is coupled to a cysteine group of the soluble CD1d or β_2 -microglobulin polypeptide. Coupling can be effected using, *e.g.*, a maleimide group, a vinylsulfone group, a haloacetate group, or a thiol group.

[0151] Optionally, the soluble CD1d or β_2 -microglobulin polypeptide is conjugated to the polyethylene-glycol moiety through a labile bond. The labile bond can be cleaved in, *e.g.*, biochemical hydrolysis, proteolysis, or sulfhydryl cleavage. For example, the bond can be cleaved under *in vivo* (physiological) conditions.

[0152] The reactions may take place by any suitable method used for reacting biologically active materials with inert polymers, generally at about pH 5-8, *e.g.*, pH 5, 6, 7, or 8, if the reactive groups are on the alpha amino group at the N-terminus. Generally the process involves preparing an activated polymer and thereafter reacting the protein with the activated polymer to produce the soluble protein suitable for formulation.

Vectors

[0153] Vectors comprising nucleic acids encoding soluble CD1d or β_2 -microglobulin polypeptides may be used to produce CD1d complexes, *e.g.*, non-specific CD1d complexes for use in the methods of the invention. The choice of vector and expression control sequences to which such nucleic acids are operably linked depends on the functional properties desired, *e.g.*, protein expression, and the host cell to be transformed.

[0154] Expression control elements useful for regulating the expression of an operably linked coding sequence are known in the art. Examples include, but are not limited to, inducible promoters, constitutive promoters, secretion signals, and other regulatory elements. When an inducible promoter is used, it can be controlled, *e.g.*, by a change in nutrient status, or a change in temperature, in the host cell medium.

[0155] The vector can include a prokaryotic replicon, *i.e.*, a DNA sequence having the ability to direct autonomous replication and maintenance of the recombinant DNA

- 47 -

molecule extra-chromosomally in a bacterial host cell. Such replicons are well known in the art. In addition, vectors that include a prokaryotic replicon may also include a gene whose expression confers a detectable marker such as a drug resistance. Examples of bacterial drug-resistance genes are those that confer resistance to ampicillin or tetracycline.

[0156] Vectors that include a prokaryotic replicon can also include a prokaryotic or bacteriophage promoter for directing expression of the coding gene sequences in a bacterial host cell. Promoter sequences compatible with bacterial hosts are typically provided in plasmid vectors containing convenient restriction sites for insertion of a DNA segment to be expressed. Examples of such plasmid vectors are pUC8, pUC9, pBR322 and pBR329 (BioRad® Laboratories), pPL and pKK223 (Pharmacia). Any suitable prokaryotic host can be used to express a recombinant DNA molecule encoding a protein used in the methods of the invention.

[0157] For the purposes of this invention, numerous expression vector systems may be employed. For example, one class of vector utilizes DNA elements which are derived from animal viruses such as bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (RSV, MMTV or MOMLV) or SV40 virus. Others involve the use of polycistronic systems with internal ribosome binding sites. Additionally, cells which have integrated the DNA into their chromosomes may be selected by introducing one or more markers which allow selection of transfected host cells. The marker may provide for prototrophy to an auxotrophic host, biocide resistance (*e.g.*, antibiotics) or resistance to heavy metals such as copper. The selectable marker gene can either be directly linked to the DNA sequences to be expressed, or introduced into the same cell by cotransformation. The neomycin phosphotransferase (neo) gene is an example of a selectable marker gene (Southern *et al.*, *J. Mol. Anal. Genet.* 1:327-341 (1982)). Additional elements may also be needed for optimal synthesis of mRNA. These elements may include signal sequences, splice signals, as well as transcriptional promoters, enhancers, and termination signals.

[0158] Of course, any expression vector which is capable of eliciting expression in eukaryotic cells may be used in the present invention. Examples of suitable vectors

- 48 -

include, but are not limited to plasmids pcDNA3, pHCMV/Zeo, pCR3.1, pEF1/His, pIND/GS, pRc/HCMV2, pSV40/Zeo2, pTRACER-HCMV, pUB6/V5-His, pVAX1, and pZeoSV2 (available from Invitrogen, San Diego, CA), and plasmid pCI (available from Promega, Madison, WI). Additional eukaryotic cell expression vectors are known in the art and are commercially available. Typically, such vectors contain convenient restriction sites for insertion of the desired DNA segment. Exemplary vectors include pSVL and pKSV-10 (Pharmacia), pBPV-1, pml2d (International Biotechnologies), pTDT1 (ATCC 31255), retroviral expression vector pMIG and pLL3.7, adenovirus shuttle vector pDC315, and AAV vectors. Other exemplary vector systems are disclosed *e.g.*, in U.S. Patent 6,413,777.

[0159] In general, screening large numbers of transformed cells for those which express suitably high levels of a soluble CD1d or β_2 -microglobulin polypeptide is routine experimentation which can be carried out, for example, by robotic systems.

[0160] Frequently used regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and enhancers derived from retroviral LTRs, cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (*e.g.*, the adenovirus major late promoter (AdmlP)), polyoma and strong mammalian promoters such as native immunoglobulin and actin promoters. For further description of viral regulatory elements, and sequences thereof, *see e.g.*, Stinski, U.S. Pat. No. 5,168,062; Bell, U.S. Pat. No. 4,510,245; and Schaffner, U.S. Pat. No. 4,968,615.

[0161] The recombinant expression vectors may carry sequences that regulate replication of the vector in host cells (*e.g.*, origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (*see, e.g.*, Axel, U.S. Pat. Nos. 4,399,216; 4,634,665 and 5,179,017). For example, typically the selectable marker gene confers resistance to a drug, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Frequently used selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr- host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

- [0162] Vectors encoding soluble CD1d or β_2 -microglobulin polypeptides can be used for transformation of a suitable host cell. Transformation can be by any suitable method. Methods for introduction of exogenous DNA into mammalian cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be introduced into mammalian cells by viral vectors.
- [0163] Transformation of host cells can be accomplished by conventional methods suited to the vector and host cell employed. For transformation of prokaryotic host cells, electroporation and salt treatment methods can be employed (Cohen *et al.*, *Proc. Natl. Acad. Sci. USA* 69:2110-14 (1972)). For transformation of vertebrate cells, electroporation, cationic lipid or salt treatment methods can be employed. *See, e.g.*, Graham *et al.*, *Virology* 52:456-467 (1973); Wigler *et al.*, *Proc. Natl. Acad. Sci. USA* 76:1373-76 (1979).
- [0164] The host cell line used for protein expression may be of mammalian origin; those skilled in the art are credited with ability to determine particular host cell lines which are best suited for the desired gene product to be expressed therein. Exemplary host cell lines include, but are not limited to NSO, SP2 cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (*e.g.*, Hep G2), A549 cells DG44 and DUXB11 (Chinese Hamster Ovary lines, DHFR minus), HELA (human cervical carcinoma), CVI (monkey kidney line), COS (a derivative of CVI with SV40 T antigen), R1610 (Chinese hamster fibroblast) BALBC/3T3 (mouse fibroblast), HAK (hamster kidney line), SP2/O (mouse myeloma), P3x63-Ag3.653 (mouse myeloma), BFA-1c1BPT (bovine endothelial cells), RAJI (human lymphocyte) and 293 (human kidney). Host cell lines are typically available from commercial services, the American Tissue Culture Collection or from published literature.
- [0165] Expression of polypeptides from production cell lines can be enhanced using known techniques. For example, the glutamine synthetase (GS) system is commonly used for enhancing expression under certain conditions. *See, e.g.*, European Patent





- 50 -





Nos. 0 216 846, 0 256 055, and 0 323 997 and European Patent Application No. 89303964.4.

Ceramide-like Glycolipid Antigens






- [0166] Ceramide like glycolipid antigens useful within the present invention include any which are capable of modulating an immune response in an animal when presented in conjunction with a CD1d molecule. The antigens may be derived from foreign antigens or from autoantigens. Further, the antigens may be synthetic. Suitable antigens are disclosed, *e.g.*, in Porcelli, U.S. Patent Appl. Publ. No. 2006/0052316, Tsuji, U.S. Patent Appl. Publ. No. 2006/0211856, Jiang, U.S. Patent Appl. Publ. No. 2006/0116331, Hirokazu *et al.*, U.S. Patent Appl. Publ. No. 2006/0074235, Tsuji *et al.*, U.S. Patent Appl. Publ. No. 2005/0192248, Tsuji, U.S. Patent Application No. 2004/0127429, and Tsuji *et al.*, U.S. Patent Application No. 2003/0157135, which are incorporated herein by reference. In certain embodiments, the ceramide-like glycolipid antigen is α -GalCer.
- [0167] Other Ceramide-like glycolipid antigens for use in the present invention include, but are not limited to, the antigens in Table 3.

TABLE 3




Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
DB04-1 (KRN7000)		α -D-Gal	C26:0	C18 aminotriol	858.32		+++	+++	KRN7000. Strong agonist, mixed IL-4 and IFN γ response both <i>in vivo</i> and <i>in vitro</i> .
DB01-1		α -D-Gal	C24:0	C18 aminotriol	830.27		++	++	Similar to KRN7000, but slightly less potent. Strong agonist, mixed IL-4 and IFN γ response both <i>in vivo</i> and <i>in vitro</i> .
DB02-1		α -D-Glu	C24:0	C18 aminotriol	830.27		+	+	Moderate to weak iNKT agonist <i>in vitro</i> and <i>in vivo</i> . Gives reasonably strong early IL-4 response, with reduced late IFN γ like DB03-4 or OCH.
DB03-4		α -D-Gal	C20; 11,14 <i>cis</i> dienoic	C18 aminotriol	770.13		+++	+++	"Type 2 Cytokine bias": Strong inducer of iNKT cell IL-4 response, with blunted IFN γ and NK cell transactivation.






Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
DB03-5		α -D-Gal	C20; 5,8,11,14 <i>cis</i> tetraenoic (arachidona te)	C18 aminotriol	766.10		++	++	"Type 2 Cytokine bias": Strong inducer of iNKT cell IL-4 response, with blunted IFN γ and NK cell transactivation.
DB03-8	PI-11	α -L-Fuc	C24:0	C18 aminotriol	814.27		+	+/-	Weak iNKT agonist activity <i>in vitro</i> . No serum cytokine response <i>in vivo</i> , but exacerbates SLE in NZB/W F1 mice. Possible antagonist/partial agonist.
DB04-9	PI-14	β -D-Man	C20, 11,14 <i>cis</i> dienoic	C18 aminotriol	770.13		++	++	Similar to DB03-4. Most active among the β -Man analogues tested to date (7/30/05)
DB05-9	PI-19	α -D-Gal	C18:2 (10t, 12c, conj)	C18 aminotriol	742.08		++	ND	A potent Th2-biased agonist <i>in vitro</i> . Not tested yet <i>in vivo</i> (12/22/06)
DB05-10	PI-20	α -D-Gal	C18:3 (9c,11t, 13c, conj)	C18 aminotriol	740.06		++	ND	Similar to DB05-9, possibly slightly more potent

- 53 -




Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
DB05-11	PI-21	α -D-Gal	C18:2 (9c, 11c, conj)	C18 aminotriol	742.08		++	ND	A potent Th2-biased analogue similar to DB05-9, possibly more active in proliferation assay
DB05-12	PI-22	α -D-Gal	C20:2 (11c, 13t, conj)	C18 aminotriol	770.13		++	ND	Similar to DB05-11, but possibly less potent and less Th2-bias
DB05-14	PI-24	α -D-Gal	C18:3 (8t, 10t, 12c, conj)	C18 aminotriol	740.06		+++	ND	Very active in prolif, strong IL-4 secretion bias
DB05-15	PI-26	α -D-Gal	C18:2 (9c, 11t, conj)	C18 aminotriol	742.08		++	ND	Similar to DB05-14
DB05-16	PI-27	α -D-Gal	C18:2 (9t, 11t, conj)	C18 aminotriol	742.08		++	ND	Similar to DB05-14
DB05-17	PI-29	α -D-Gal	C18:3 (9c, 11t, 13t, conj)	C18 aminotriol	740.06		++	ND	Weak in splenocyte prolif, but moderately strong in cytokine secretion assays with moderate IL-4 predominance


- 54 -

Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
DB06-14		α -L-Fuc	C26:0	C18 aminotriol	842.32		+	ND	Weakly active in splenocyte stimulation <i>in vitro</i> . Minimal or no prolifer, but moderate IL-4 secretion at high concentration with weak IFNg (needs to be compared directly to DB03-8)
DB06-15		α -D-Glu	C20:2 (cis 11,14)	C18 aminotriol	770.13		+	ND	Moderately active in splenocyte prolifer; Weak IL-4 and even weaker IFNg secretion
AH04-1	RMN 3-84	α -D-Gal	C24:0	C9 aminotriol	704.03		++	++	Identical to OCH
AH04-2		α -D-Gal	C24:0	C9 aminodiol	688.03		++	ND	Similar activity to OCH in B6 splenocyte assay. About a half log more potent than OCH in hybridoma stimulation assay. No info on recognition by human iNKT cells.

Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
YTC03-15		α -D-Gal	C18:0	C18 aminotriol	746.11		+	ND	EXTREMELY active when presented to mouse iNKT hybridoma by human CD1d+ HeLa cells
YTC03-17		α -D-Gal	Biphenylacetate	C18 aminotriol	705.96		++	-	Very strong agonist in some <i>in vitro</i> studies, but no activity detectable <i>in vivo</i> (cytokine stimulation).
YTC03-24		α -D-Gal	dimethoxyphenylacetate	C18 aminotriol	657.83		+	-	Possible enhanced IL-4 relative to IFN γ <i>in vitro</i> with splenocytes from NZB/W F1 mice
YTC03-30		α -D-Gal	fluorephenylacetate	C18 aminotriol	615.77		+	-	Possible enhanced IL-4 relative to IFN γ <i>in vitro</i> with splenocytes from NZB/W F1 mice. EXTREMELY active when presented to mouse iNKT hybridoma by human CD1d+ HeLa cells
YTC03-33		α -D-Gal	methoxyphenylacetate	C18 aminotriol	627.81		+	ND	

- 56 -

Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
YTC03-34		α -D-Gal	C10:0	C18 aminotriol	633.90		+	+	Moderate activity with mouse iNKT hybridoma. Possible TH2 skewing of cytokine response? Active <i>in vivo</i> with good IL-4 production assoc. with weak IFN γ and weak IL-12 p70 (very interesting compound!)
SKRN7000		α -D-Gal	C26:0	C18 aminotriol	874.39		-	ND	S-glycoside of KRN7000 from D. Bundle. No agonist activity in iNKT hybridoma assay. However, seemed to reduce autoreactivity of iNKT cell hybridoma, so might be worth evaluating as an antagonist.
RF03-1		α -D-Gal	C26:0	C19 aminodiol	856.35		+/-	ND	C-glycoside of KRN7000. Reported to be strong agonist, with enhanced IFN γ production <i>in vivo</i> . May have slower kinetics of activation.

Compound Name		CHO group	N-linked group	sphingoid base	MW	Structure	Activity in vitro	Activ-ity in vivo	Comments
Bronx	UK/other								
OCH	(AH04-1)	α -D-Gal	C24:0	C9 aminotriol	704.03		++	++	Famous "TH2-skewing" analogue. Weaker agonist than DB03-4 in our hands, and skewing of cytokine response not very impressive in most assays. Also, seems NOT to be recognized by human iNKT cells.

- [0168] Ceramide-like glycolipid antigens are bound or associated with soluble CD1d polypeptides by standard methods known to those of ordinary skill in the art. For example, a preparation of purified CD1d protein (including both the CD1d and β_2 -microglobulin subunits) is mixed with a 2.5 molar excess of α -GalCer, insuring that every CD1d protein is bound with antigen. Excess α -GalCer is then removed, *e.g.*, by chromatographic methods, *e.g.*, size exclusion FPLC.
- [0169] In certain embodiments, compositions for use in the methods of the present invention further comprise another component, *e.g.*, a polypeptide with immunological activity. Preferably, the protein with immunological activity is a costimulatory molecule, such as a Saponin, a toll-like receptor ("TLR"), B7.1 or B7.2. "B7" is used herein to generically refer to either B7.1 or B7.2. In one embodiment, a costimulatory molecule, *e.g.*, the extracellular domain of B7-1 (CD80) or B7-2 (CD86) that interacts with CD28 on T- and NK-cells, is administered as an amino terminal fusion to β_2 -microglobulin incorporated into the structure of a soluble CD1d complex for use in the present invention. See, *e.g.*, WO 9964597, published 16 Dec 1999. Alternatively, a costimulatory molecule is administered as an amino-terminal fusion to the CD1d heavy chain. In certain embodiments, incorporation of a costimulatory molecule, *e.g.*, a B7 signaling molecule in the compositions of the invention allows more effective and prolonged activation of CD1d-restricted NKT cells by the soluble CD1d complex.
- [0170] In other embodiments, the compositions for use in the methods of the present invention further comprise adjuvant components, *e.g.*, Toll-like receptor (TLR) agonists. Examples of TLR agonist adjuvants which may be effective, include, but are not limited to: N-acetylmuramyl-L-alanine-D-isoglutamine (MDP), lipopolysaccharides (LPS), genetically modified and/or degraded LPS, alum, glucan, colony stimulating factors (*e.g.*, EPO, GM-CSF, G-CSF, M-CSF, PEGylated G-CSF, SCF, IL-3, IL6, PIXY 321), interferons (*e.g.*, γ -interferon, α -interferon), interleukins (*e.g.*, IL-2, IL-7, IL-12, IL-15, IL-18), saponins (*e.g.*, QS21), monophosphoryl lipid A (MPL), 3 De-O-acylated monophosphoryl lipid A (3D-MPL), unmethylated CpG sequences, 1-methyl tryptophan, arginase inhibitors, cyclophosphamide, antibodies that block immunosuppressive functions (*e.g.*, anti-CTLA4 antibodies), lipids (such as

palmitic acid residues), tripalmitoyl-S-glycerylcystein lyseryl-serine (P₃ CSS), and Freund's adjuvant.

[0171] Alternatively or additionally, compositions of the present invention may further comprise a lymphokine or cytokine that modulates immune cell activation such as interleukins IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-10, IL-12, IL-15, IL-18; granulocyte-macrophage colony stimulating factor (GM-CSF); transforming growth factor (TGF, *e.g.*, TGF α and TGF β); α interferons (*e.g.* IFN α); β interferons (*e.g.* IFN β); γ interferons (*e.g.* IFN γ) or lymphocyte function-associated protein, such as LFA-1 or LFA-3; or an intercellular adhesion molecule, such as ICAM-1 or ICAM-2.

[0172] Compositions of the present invention may contain a homogenous or heterogeneous population of antigens and/or costimulatory molecules. That is, each soluble CD1d polypeptide in the composition may be linked to the same ceramide-like glycolipid antigen or soluble CD1d polypeptides may be linked to different antigens. Likewise, various soluble CD1d complexes may be associated with the same costimulatory molecules or different costimulatory molecules.

[0173] The soluble CD1d complexes of the present invention, or compositions comprising same may be labeled, so as to be directly detectable, or will be used in conjunction with secondary labeled immunoreagents which will specifically bind the compound for example, for detection or diagnostic purposes. Labels of interest may include dyes, enzymes, chemilumescers, particles, radioisotopes, or other directly or indirectly detectable agent. Alternatively, a second stage label may be used, *e.g.* labeled antibody directed to one of the constituents of the compound of the invention.

[0174] Examples of suitable enzyme labels include malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast-alcohol dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, and acetylcholine esterase.

[0175] Examples of suitable radioisotopic labels include ³H, ¹¹¹In, ¹²⁵I, ¹³¹I, ³²P, ³⁵S, ¹⁴C, ⁵¹Cr, ⁵⁷To, ⁵⁸Co, ⁵⁹Fe, ⁷⁵Se, ¹⁵²Eu, ⁹⁰Y, ⁶⁷Cu, ²¹⁷Pb, ²¹¹At, ²¹²Pb, ⁴⁷Sc,

- 60 -

¹⁰⁹Pd, etc. Examples of suitable non-radioactive isotopic labels include ¹⁵⁷Gd, ⁵⁵Mn, ¹⁶²Dy, ⁵²Tr, and ⁵⁶Fe.

[0176] Examples of suitable fluorescent labels include an ¹⁵²Eu label, a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin label, a phycocyanin label, an allophycocyanin label, an o-phthaldehyde label, and a fluorescamine label.

[0177] Examples of suitable toxin labels include diphtheria toxin, ricin, and cholera toxin.

[0178] Examples of chemiluminescent labels include a luminal label, an isoluminal label, an aromatic acridinium ester label, an imidazole label, an acridinium salt label, an oxalate ester label, a luciferin label, a luciferase label, and an aequorin label.

[0179] Examples of nuclear magnetic resonance contrasting agents include heavy metal nuclei such as Gd, Mn, and Fe.

[0180] Typical techniques for binding the above-described labels to polypeptides are provided by Kennedy *et al.*, Clin. Chim. Acta 70:1-31 (1976), and Schurs *et al.*, Clin. Chim. Acta 81:1-40 (1977). Coupling techniques mentioned in the latter are the glutaraldehyde method, the periodate method, the dimaleimide method, the m-maleimidobenzyl-N-hydroxy-succinimide ester method, all of which methods are incorporated by reference herein.

Immunogenic And Therapeutic Molecules

[0181] An "immunogenic polypeptide" is meant to encompass any antigenic or immunogenic polypeptides including poly-aminoacid materials having epitopes or combinations of epitopes. As used herein, an immunogenic polypeptide is a polypeptide which, when introduced into a vertebrate, reacts with the immune system molecules of the vertebrate, *i.e.*, is antigenic, and/or induces an immune response in the vertebrate, *i.e.*, is immunogenic. It is quite likely that an immunogenic polypeptide will also be antigenic, but an antigenic polypeptide, because of its size or conformation, may not necessarily be immunogenic. Examples of antigenic and immunogenic polypeptides include, but are not limited to, polypeptides from infectious agents such as bacteria, viruses, parasites, or fungi, allergens such as those

- 61 -

from pet dander, plants, dust, and other environmental sources, as well as certain self polypeptides, for example, tumor-associated antigens.

[0182] Antigenic and immunogenic polypeptides of the present invention can be used to prevent or treat, *i.e.*, cure, ameliorate, lessen the severity of, or prevent or reduce contagion of viral, bacterial, fungal, and parasitic infectious diseases, as well as to treat allergies.

[0183] In addition, antigenic and immunogenic polypeptides of the present invention can be used to prevent or treat, *i.e.*, cure, ameliorate, or lessen the severity of cancer including, but not limited to, cancers of oral cavity and pharynx (*i.e.*, tongue, mouth, pharynx), digestive system (*i.e.*, esophagus, stomach, small intestine, colon, rectum, anus, anal canal, anorectum, liver, gallbladder, pancreas), respiratory system (*i.e.*, larynx, lung), bones, joints, soft tissues (including heart), skin, melanoma, breast, reproductive organs (*i.e.*, cervix, endometrium, ovary, vulva, vagina, prostate, testis, penis), urinary system (*i.e.*, urinary bladder, kidney, ureter, and other urinary organs), eye, brain, endocrine system (*i.e.*, thyroid and other endocrine), lymphoma (*i.e.*, hodgkin's disease, non-hodgkin's lymphoma), multiple myeloma, leukemia (*i.e.*, acute lymphocytic leukemia, chronic lymphocytic leukemia, acute myeloid leukemia, chronic myeloid leukemia).

[0184] Examples of viral antigenic and immunogenic polypeptides include, but are not limited to, adenovirus polypeptides, alphavirus polypeptides, calicivirus polypeptides, *e.g.*, a calicivirus capsid antigen, coronavirus polypeptides, distemper virus polypeptides, Ebola virus polypeptides, enterovirus polypeptides, flavivirus polypeptides, hepatitis virus (AE) polypeptides, *e.g.*, a hepatitis B core or surface antigen, herpesvirus polypeptides, *e.g.*, a herpes simplex virus or varicella zoster virus glycoprotein, immunodeficiency virus polypeptides, *e.g.*, the human immunodeficiency virus envelope or protease, infectious peritonitis virus polypeptides, influenza virus polypeptides, *e.g.*, an influenza A hemagglutinin, neuraminidase, or nucleoprotein, leukemia virus polypeptides, Marburg virus polypeptides, orthomyxovirus polypeptides, papilloma virus polypeptides, parainfluenza virus polypeptides, *e.g.*, the hemagglutinin/neuraminidase, paramyxovirus polypeptides, parvovirus polypeptides, pestivirus polypeptides,

picorna virus polypeptides, *e.g.*, a poliovirus capsid polypeptide, pox virus polypeptides, *e.g.*, a vaccinia virus polypeptide, rabies virus polypeptides, *e.g.*, a rabies virus glycoprotein G, reovirus polypeptides, retrovirus polypeptides, and rotavirus polypeptides.

[0185] Examples of bacterial antigenic and immunogenic polypeptides include, but are not limited to, *Actinomyces* polypeptides, *Bacillus* polypeptides, *e.g.*, immunogenic polypeptides from *Bacillus anthracis*, *Bacteroides* polypeptides, *Bordetella* polypeptides, *Bartonella* polypeptides, *Borrelia* polypeptides, *e.g.*, *B. burgdorferi* OspA, *Brucella* polypeptides, *Campylobacter* polypeptides, *Capnocytophaga* polypeptides, *Chlamydia* polypeptides, *Clostridium* polypeptides, *Corynebacterium* polypeptides, *Coxiella* polypeptides, *Dermatophilus* polypeptides, *Enterococcus* polypeptides, *Ehrlichia* polypeptides, *Escherichia* polypeptides, *Francisella* polypeptides, *Fusobacterium* polypeptides, *Haemobartonella* polypeptides, *Haemophilus* polypeptides, *e.g.*, *H. influenzae* type b outer membrane protein, *Helicobacter* polypeptides, *Klebsiella* polypeptides, L form bacteria polypeptides, *Leptospira* polypeptides, *Listeria* polypeptides, *Mycobacteria* polypeptides, *Mycoplasma* polypeptides, *Neisseria* polypeptides, *Neorickettsia* polypeptides, *Nocardia* polypeptides, *Pasteurella* polypeptides, *Peptococcus* polypeptides, *Peptostreptococcus* polypeptides, *Pneumococcus* polypeptides, *Proteus* polypeptides, *Pseudomonas* polypeptides, *Rickettsia* polypeptides, *Rochalimaea* polypeptides, *Salmonella* polypeptides, *Shigella* polypeptides, *Staphylococcus* polypeptides, *Streptococcus* polypeptides, *e.g.*, *S. pyogenes* M proteins, *Treponema* polypeptides, and *Yersinia* polypeptides, *e.g.*, *Y. pestis* F1 and V antigens.

[0186] Examples of tumor-associated antigenic and immunogenic polypeptides include, but are not limited to, tumor-specific immunoglobulin variable regions, GM2, Tn, sTn, Thompson-Friedenreich antigen (TF), Globo H, Le(y), MUC1, MUC2, MUC3, MUC4, MUC5AC, MUC5B, MUC7, carcinoembryonic antigens, beta chain of human chorionic gonadotropin (hCG beta), C35, HER2/neu, CD20, PSMA, EGFRvIII, KSA, PSA, PSCA, GP100, MAGE 1, MAGE 2, TRP 1, TRP 2, tyrosinase, MART-1, PAP, CEA, BAGE, MAGE, RAGE, and related proteins.

- 63 -

[0187] Compositions of the present invention may further comprise other therapeutic agents. The therapeutic agent or agents may be linked to or otherwise associated with the soluble the CD1d complex. Examples of therapeutic agents include, but are not limited to, antimetabolites, alkylating agents, anthracyclines, antibiotics, and anti-mitotic agents. Antimetabolites include methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine. Alkylating agents include mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin. Anthracyclines include daunorubicin (formerly daunomycin) and doxorubicin (also referred to herein as adriamycin). Additional examples include mitozantrone and bisantrene. Antibiotics include dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC). Antimytotic agents include vincristine and vinblastine (which are commonly referred to as vinca alkaloids). Other cytotoxic agents include procarbazine, hydroxyurea, asparaginase, corticosteroids, mytotane (O,P'-(DDD)), interferons. Further examples of cytotoxic agents include, but are not limited to, ricin, doxorubicin, taxol, cytochalasin B, gramicidin D, ethidium bromide, etoposide, tenoposide, colchicin, dihydroxy anthracin dione, 1-dehydrotestosterone, and glucocorticoid.

[0188] Analogs and homologs of such therapeutic and cytotoxic agents are encompassed by the present invention. For example, the chemotherapeutic agent aminopterin has a correlative improved analog namely methotrexate. Further, the improved analog of doxorubicin is an Fe-chelate. Also, the improved analog for 1-methylnitrosourea is lomustine. Further, the improved analog of vinblastine is vincristine. Also, the improved analog of mechlorethamine is cyclophosphamide.

NKT Activity Assays

[0189] The ability of a composition of the present invention to modulate an immune response can be readily determined by an *in vitro* assay. NKT cells for use in the assays include transformed NKT cell lines, or NKT cells which are isolated from a mammal, *e.g.*, from a human or from a rodent such as a mouse. NKT cells can be

- 64 -

isolated from a mammal by sorting cells that bind CD1d:α-GalCer tetramers. See, for example, Benlagha *et al.*, J Exp Med 191 (2000), pp. 1895–1903; Matsuda *et al.*, J Exp Med 192 (2000), pp. 741–754; and Karadimitris *et al.*, Proc Natl Acad Sci USA 98 (2001), pp. 3294–3298. A suitable assay to determine if a compound of the present invention is capable of modulating the activity of NKT cells is conducted by coculturing NKT cells and antigen presenting cells, adding the particular compound of interest to the culture medium that targets either the antigen presenting cells or the NKT cells directly, and measuring IL-4 or IFN-γ production. A significant increase or decrease in IL-4 or IFN-γ production over the same co-culture of cells in the absence of the compound of the invention or, preferably, in the presence of a compound of the invention with a non-targeting antibody indicates stimulation or inhibition of NKT cells.

[0190] The NKT cells employed in the assays are incubated under conditions suitable for proliferation. For example, an NKT cell hybridoma is suitably incubated at about 37°C and 5% CO₂ in complete culture medium (RPMI 1640 supplemented with 10% FBS, penicillin/streptomycin, L-glutamine and 5x10⁻⁵ M 2-mercaptoethanol). Serial dilutions of the compound can be added to the NKT cell culture medium. Suitable concentrations of the compound added to the NKT cells typically will be in the range of from 10⁻¹² to 10⁻⁶ M. Use of antigen dose and APC numbers giving slightly submaximal NKT cell activation is preferred to detect stimulation or inhibition of NKT cell responses by the compounds of the invention.

[0191] Alternatively, rather than measurement of an expressed protein such as IL-4 or IFN-γ, modulation of NKT cell activation can be determined by changes in antigen-dependent T cell proliferation as measured by radiolabelling techniques as are recognized in the art. For example, a labeled (*e.g.*, tritiated) nucleotide may be introduced to an assay culture medium. Incorporation of such a tagged nucleotide into DNA serves as a measure of T cell proliferation. This assay is not suitable for NKT cells that do not require antigen presentation for growth, *e.g.*, NKT cell hybridomas. A difference in the level of T cell proliferation following contact with the compound of the invention indicates the compound modulates activity of the T cells. For example, a decrease in NKT cell proliferation indicates the compound can suppress an immune

- 65 -

response. An increase in NKT cell proliferation indicates the compound can stimulate an immune response.

[0192] Additionally, the ^{51}Cr release assay, described below, can be used to determine cytotoxic activity.

[0193] These *in vitro* assays can be employed to select and identify soluble CD1d complexes and compositions comprising same that are capable of modulating an immune response. Assays described above, *e.g.*, measurement of IL-4 or IFN- γ production or NKT cell proliferation, are employed to determine if contact with the compound modulates T cell activation.

[0194] *In vivo* assays also may be suitably employed to determine the ability of a composition of the invention to modulate the activity of NKT cells. For example, a composition of interest can be assayed for its ability to stimulate NKT cell activation or inhibit tumor growth. For example, a composition of the invention can be administered to a mammal such as a mouse, before or after challenge with a tumorigenic dose of transformed cells and the presence or size of growing tumors may be monitored.

[0195] Compositions of the present invention further comprise a suitable carrier. Such compositions comprise a therapeutically effective amount of the soluble CD1d complex and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The formulation should suit the mode of administration.

Methods Of Treatment

[0196] The present invention also includes a method of modulating, *i.e.*, either stimulating or inhibiting an immune response, comprising administering to an animal an effective amount of a composition comprising a soluble CD1d complex loaded with a ceramide-like glycolipid antigen as described herein.

[0197] The present invention further provides a method of treating a disease in an animal, comprising administering to an animal with that disease, or prone to contract that disease, a composition comprising a soluble CD1d complex loaded with a ceramide-like glycolipid antigen as described herein.

- 66 -

- [0198] According to these methods, a composition of the present invention is administered in an amount sufficient to alter the progression of said disease.
- [0199] Certain embodiments of the present invention include a method of reducing or eliminating the anergic response of NKT cells to multiple administrations of ceramide-like glycolipid antigens administered by themselves, which are therefore presented to NKT cells in the context of cell-bound CD1d. It has been shown that multiple administrations of α -GalCer, administered by itself, causes NKT cells to become non-responsive for an extended period of time. The present invention, in which glycolipids such as α -GalCer are administered as part of a soluble CD1d complex, protects NKT cells from anergy in response to antigen, and allows for a prolonged response upon multiple administrations. Accordingly, NKT cells are activated in response to stimulation with soluble CD1d complexes loaded with a ceramide-like glycolipid antigen of the present invention and furthermore, NKT cells can be reactivated in response to restimulation by soluble CD1d complexes loaded with a ceramide-like glycolipid antigen of the present invention.
- [0200] In certain embodiments, soluble CD1d complexes for use in the methods described herein are "non-specific" soluble CD1d complexes, *i.e.*, they are not targeted to a specific organ, tissue, cell, or cell surface marker, rather they are administered systemically.
- [0201] According to the methods of the present invention, a composition comprising a soluble CD1d complex is administered to modulate an immune response in an animal, *e.g.*, a vertebrate, *e.g.*, a mammal, *e.g.*, a human. In certain embodiments, the methods of the present invention result in the enhancement of an immune response, *e.g.*, to an immunogen delivered before, after, or concurrently with a soluble CD1d complex. Administration of soluble CD1d complexes of the invention, *e.g.*, with an immunogen, may typically result in the release of cytokines from immune cells, *e.g.*, NKT cells or NK cells. Cytokines released in response to administration of compositions of the invention may be those associated with a TH1-type immune response, *e.g.*, interferon gamma and TNF-alpha. Alternatively, or in addition, administration of compositions of the present invention may result in the release of cytokines associated with a TH2-type immune response, *e.g.*, IL-4, IL-5, IL-10, or IL-

- 67 -

13. Alternatively, or in addition, administration of compositions of the present invention may result in the release of other cytokines, *e.g.*, IL-2, IL-1 β , IL-12, IL-17, IL-23, TNF- β /LT, MCP-2, oncostatin-M, and RANTES. Methods to modulate the type of cytokines released include varying the ceramide-like glycolipid antigen of the soluble CD1d complex. Choosing and testing various ceramide-like glycolipid antigens for their effect on cytokine release from NKT or other immune cells may be performed using *in vitro* assays described elsewhere herein and in Porcelli, U.S. Patent Appl. Publ. No. 2006/0052316, as well as by additional methods well-known by those of ordinary skill in the art. Administration of soluble CD1d complexes of the present invention and compositions comprising same may further modulate an immune response by inducing proliferation of NKT cells, and also by inducing recruitment and or activation of other immune cells including, but not limited to NK cells, CTLs, other T lymphocytes, *e.g.*, CD8⁺ or CD4⁺ T lymphocytes, dendritic cells, B lymphocytes, and others.

[0202] In certain embodiments, administration of soluble CD1d complexes of the present invention and compositions comprising same results in the suppression or inhibition of an undesired immune response, *e.g.*, inflammation or autoimmunity.

[0203] In certain embodiments, administration of soluble CD1d complexes of the present invention and compositions comprising same affects one or more NKT cell activities such as, but not limited to cell proliferation, the production of one or more cytokines, or recruitment and/or activation of non-NKT immune system cells including, but not limited to NK cells, CTLs, other T lymphocytes, *e.g.*, CD8⁺ or CD4⁺ T lymphocytes, dendritic cells, B lymphocytes, and others.

[0204] Certain embodiments of the present invention involve use of soluble CD1d complexes of the invention as adjuvants, *i.e.*, to modulate an immune response to a specific immunogen. Accordingly, the present invention provides a method of modulating an immune response to an immunogen in an animal, where the method comprises administering to an animal in need of such modulation a composition comprising an immunogen, a soluble CD1d complex loaded with a ceramide-like glycolipid antigen as described elsewhere herein, and a suitable carrier. According to this embodiment, the soluble CD1d complex is administered in an amount sufficient

- 68 -

to modulate the immune response against the immunogen relative to administration of the immunogen without the soluble CD1d complex. A soluble CD1d complex for use as an adjuvant as described herein may in certain embodiments be a non-specific soluble CD1d complex. In other embodiments, a soluble CD1d complex for use as an adjuvant may be targeted to a particular organ, tissue, cell or cell surface marker as described, *e.g.*, in Bruno *et al.* U.S. Patent Appl. Publ. No. 2006/0269540.

[0205] In certain embodiments, soluble CD1d complexes of the present invention and compositions comprising same are administered with an immunogen as a therapeutic vaccine, *e.g.*, to an animal already suffering from a disease such as cancer. According to these methods, the immune response elicited by the immunogen/adjuvant composition is effective in treating, *e.g.*, affecting the outcome of the disease by reducing symptoms or lessening the severity of the disease, and the non-specific CD1d complex is administered in an amount sufficient to modulate the immune response against the immunogen relative to administration of the immunogen in the absence of the non-specific soluble CD1d complex. Alternatively, soluble CD1d complexes of the present invention and compositions comprising same are administered with an immunogen as a prophylactic vaccine, *i.e.*, to prevent, or reduce symptoms to a disease, such as an infectious disease that might be contracted by that animal in the future. According to these methods, the immune response elicited by the immunogen/adjuvant composition is effective in preventing, *e.g.*, affecting the outcome of the disease by reducing symptoms or lessening the severity of the disease, and the non-specific CD1d complex is administered in an amount sufficient to modulate the immune response against the immunogen relative to administration of the immunogen in the absence of the non-specific soluble CD1d complex.

[0206] The present invention also provides immunogen/adjuvant compositions for use in the methods described herein. Such compositions comprise an immunogen and a soluble CD1d complex as described elsewhere herein. Immunogen/adjuvant compositions of the present invention typically include non-specific soluble CD1d complex, but may, in certain embodiments, include targeted soluble CD1d complexes.

[0207] The methods and compositions as described herein are useful for raising an immune response and treating hyperproliferative disorders. Examples of

hyperproliferative disorders that can be treated by the compounds of the invention include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

[0208] Similarly, other hyperproliferative disorders can also be treated by the compounds of the invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenström's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

[0209] The methods and compositions as described herein are also useful for raising an immune response against infectious agents. Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated by the compounds of the invention. Examples of viruses, include, but are not limited to the following DNA and RNA viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Flaviviridae, Hepadnaviridae (hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (*e.g.*, Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (*e.g.*, Influenza), Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (*e.g.*, Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (*e.g.*, Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, encephalitis, eye infections (*e.g.*, conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), meningitis, opportunistic infections (*e.g.*, AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, measles, mumps, parainfluenza, rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (*e.g.*, Kaposi's, warts), and viremia.

- 70 -

[0210] Similarly, bacterial or fungal agents that can cause disease or symptoms can be treated or prevented by the methods and compositions of the invention. These include, but are not limited to, the following Gram-Negative and Gram-positive bacterial families and fungi: Actinomycetales (*e.g.*, *Corynebacterium*, *Mycobacterium*, *Nocardia*), Aspergillosis, Bacillaceae (*e.g.*, *Anthrax*, *Clostridium*), Bacteroidaceae, Blastomycosis, *Bordetella*, *Borrelia*, Brucellosis, Candidiasis, *Campylobacter*, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, Enterobacteriaceae (*Klebsiella*, *Salmonella*, *Serratia*, *Yersinia*), *Erysipelothrix*, *Helicobacter*, Legionellosis, Leptospirosis, *Listeria*, Mycoplasmatales, Neisseriaceae (*e.g.*, *Acinetobacter*, *Gonorrhea*, *Menigococcal*), Pasteurellaceae Infections (*e.g.*, *Actinobacillus*, *Haemophilus*, *Pasteurella*), *Pseudomonas*, Rickettsiaceae, Chlamydiaceae, Syphilis, and Staphylococcal. These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (*e.g.*, AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, *Gonorrhea*, meningitis, Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (*e.g.*, cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections.

[0211] Moreover, the methods and compositions of the present invention may be used to treat or prevent diseases caused by parasitic agents. These include, but are not limited to that can be treated by the compounds of the invention include, but are not limited to, the following families: amebiasis, babesiosis, coccidiosis, cryptosporidiosis, *dientamoebiasis*, dourine, ectoparasitic, giardiasis, helminthiasis, leishmaniasis, theileriasis, toxoplasmosis, trypanosomiasis, and trichomonas.

[0212] Additionally, the methods and compositions of the present invention may be used to treat or prevent autoimmune diseases. An autoimmune disease is characterized by the attack by the immune system on the tissues of the victim. In

- 71 -

autoimmune diseases, the recognition of tissues as "self" apparently does not occur, and the tissue of the afflicted subject is treated as an invader--*i.e.*, the immune system sets about destroying this presumed foreign target. The compounds of the present invention are therefor useful for treating autoimmune diseases by desensitizing the immune system to these self antigens by, for example, immune deviation away from a destructive HG1 type response.

[0213] Examples of autoimmune diseases which may be treated using the compounds of the present invention include, but are not limited to Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, multiple sclerosis, myasthenia gravis, neuritis, ophthalmia, bullous pemphigoid, pemphigus, polyendocrinopathies, purpura, Reiter's Disease, Stiff-Man Syndrome, autoimmune thyroiditis, systemic lupus erythematosus, autoimmune pulmonary inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, autoimmune inflammatory eye disease, autoimmune hemolysis, psoriasis, juvenile diabetes, primary idiopathic myxedema, autoimmune asthma, scleroderma, chronic hepatitis, hypogonadism, pernicious anemia, vitiligo, alopecia areata, Coeliac disease, autoimmune enteropathy syndrome, idiopathic thrombocytic purpura, acquired splenic atrophy, idiopathic diabetes insipidus, infertility due to antispermatozoan antibodies, sudden hearing loss, sensorineural hearing loss, polymyositis, autoimmune demyelinating diseases, transverse myelitis, ataxic sclerosis, progressive systemic sclerosis, dermatomyositis, polyarteritis nodosa, idiopathic facial paralysis, cryoglobulinemia, inflammatory bowel diseases, Hashimoto's disease, adrenalitis, hypoparathyroidism, and ulcerative colitis.

[0214] Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by the methods and compositions of the invention. In one embodiment, the invention provides for effective delivery of signals that inhibit or skew cytokine production by NKT cells resulting in reduced immune responses or immune deviation. For example, the methods and compositions of the invention can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

- [0215] The methods and compositions of the invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of the compounds of the invention that inhibit or result in immune deviation of an immune response may be an effective therapy in preventing organ rejection or GVHD.
- [0216] Certain compositions of the invention described elsewhere herein which can inhibit an immune response upon administration are also useful for treating and/or preventing atherosclerosis; otitis; regional enteritis; adult respiratory distress syndrome; local manifestations of drug reactions, such as dermatitis, etc.; inflammation-associated or allergic reaction patterns of the skin; atopic dermatitis and infantile eczema; contact dermatitis; psoriasis; lichen planus; allergic enteropathies; allergic rhinitis; bronchial asthma; hypersensitivity or destructive responses to infectious agents; poststreptococcal diseases, *e.g.* cardiac manifestations of rheumatic fever, and the like.
- [0217] According to the disclosed methods, compositions for use in the methods of the present invention can be administered, for example, by intramuscular (*i.m.*), subcutaneous (*s.c.*), or intrapulmonary routes. Other suitable routes of administration include, but are not limited to intratracheal, transdermal, intraocular, intranasal, inhalation, intracavity, intravenous (*i.v.*), intraductal (*e.g.*, into the pancreas) and intraparenchymal (*i.e.*, into any tissue) administration. Transdermal delivery includes, but not limited to intradermal (*e.g.*, into the dermis or epidermis), transdermal (*e.g.*, percutaneous) and transmucosal administration (*i.e.*, into or through skin or mucosal tissue). Intracavity administration includes, but not limited to administration into oral, vaginal, rectal, nasal, peritoneal, or intestinal cavities as well as, intrathecal (*i.e.*, into spinal canal), intraventricular (*i.e.*, into the brain ventricles or the heart ventricles), inraatrial (*i.e.*, into the heart atrium) and sub arachnoid (*i.e.*, into the sub arachnoid spaces of the brain) administration.

Pharmaceutical Compositions

- [0218] Soluble CD1d complexes of the present invention may be administered in pharmaceutical compositions either with or without an immunogen, in combination with one or more pharmaceutically acceptable excipients. It will be understood that, when administered to a human patient, the total daily usage of the pharmaceutical compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the type and degree of the response to be achieved; the specific composition of another agent, if any, employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the composition; the duration of the treatment; drugs (such as a chemotherapeutic agent) used in combination or coincidental with the specific composition; and like factors well known in the medical arts. Suitable formulations, known in the art, can be found in Remington's Pharmaceutical Sciences (latest edition), Mack Publishing Company, Easton, PA.
- [0219] A pharmaceutical composition to be used in a given therapeutic treatment will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the compounds alone), the site of delivery of the compound, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" of the compounds of the invention for purposes herein is thus determined by such considerations.
- [0220] Pharmaceutical compositions of the invention may be administered orally, intravenously, rectally, parenterally, intracisternally, intradermally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, creams, drops or transdermal patch), buccally, or as an oral or nasal spray. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

- 74 -

[0221] The pharmaceutical compositions are administered in an amount which is effective for treating and/or prophylaxis of the specific indication. In most cases, the dosage is from about 1 $\mu\text{g/kg}$ to about 30 mg/kg body weight daily, taking into account the routes of administration, symptoms, etc. However, the dosage can be as low as 0.001 $\mu\text{g/kg}$.

[0222] As a general proposition, the total pharmaceutically effective amount of the compositions administered parenterally per dose will be in the range of about 1 $\mu\text{g/kg/day}$ to 100 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. If given continuously, the composition is typically administered at a dose rate of about 1 $\mu\text{g/kg/hour}$ to about 5 mg/kg/hour , either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution or bottle solution may also be employed.

[0223] The compositions of the invention may also be suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, *e.g.*, films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (U. Sidman *et al.*, Biopolymers 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (R. Langer *et al.*, J. Biomed. Mater. Res. 15:167-277 (1981), and R. Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (R. Langer *et al.*, *Id.*) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped compositions of the present invention. Liposomes are prepared by methods known per se: DE 3,218,121; Epstein, *et al.*, Proc. Natl. Acad. Sci. USA 82:3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal therapy.

- [0224] For parenteral administration, in one embodiment, a composition of the invention is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, *i.e.*, one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compositions that are known to be deleterious to polypeptides.
- [0225] Generally, the formulations are prepared by contacting a soluble CD1d complex and optionally an immunogen of the invention uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes. Suitable formulations, known in the art, can be found in Remington's Pharmaceutical Sciences (latest edition), Mack Publishing Company, Easton, PA.
- [0226] The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, *e.g.*, polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.
- [0227] The compositions are typically formulated in such vehicles at a non-limiting concentration of about 0.01 µg/ml to 100 mg/ml, for example about 0.01 µg/ml to 10

- 76 -

mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of salts.

[0228] Compositions to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (*e.g.*, 0.2 micron membranes). Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0229] Compositions of the invention ordinarily will be stored in unit or multi-dose containers, for example, sealed ampules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized composition using bacteriostatic Water-for-Injection.

[0230] Dosaging may also be arranged in a patient specific manner to provide a predetermined concentration of activity in the blood, as determined by an RIA technique, for instance. Thus patient dosaging may be adjusted to achieve regular on-going trough blood levels, as measured by RIA, on the order of from 50 to 1000 ng/ml, preferably 150 to 500 ng/ml.

[0231] Compositions of the invention are useful for administration to any animal, preferably a mammal (such as apes, cows, horses, pigs, boars, sheep, rodents, goats, dogs, cats, chickens, monkeys, rabbits, ferrets, whales, and dolphins), and more preferably a human.

[0232] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such containers can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the compositions of the present invention may be employed in conjunction with other therapeutic compositions.

- 77 -

[0233] Other therapeutic compositions useful for administration along with a composition of the present invention include cytotoxic drugs, particularly those which are used for cancer therapy. Such drugs include, in general, alkylating agents, anti-proliferative agents, tubulin binding agents and the like. Preferred classes of cytotoxic agents include, for example, the anthracycline family of drugs, the vinca drugs, the mitomycins, the bleomycins, the cytotoxic nucleosides, the pteridine family of drugs, diynenes, and the podophyllotoxins. Particularly useful members of those classes include, for example, adriamycin, carminomycin, daunorubicin, aminopterin, methotrexate, methopterin, dichloromethotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, cytosine arabinoside, podophyllotoxin, or podophyllotoxin derivatives such as etoposide or etoposide phosphate, melphalan, vinblastine, vincristine, leurosine, vindesine, leurosine and the like. As noted previously, one skilled in the art may make chemical modifications to the desired compound in order to make reactions of that compound more convenient for purposes of preparing conjugates of the invention.

[0234] The compositions of the invention can be used to treat tumor-bearing animals, including humans, to generate an immune response against tumor cells. The generation of an adequate and appropriate immune response leads to tumor regression *in vivo*. Such "vaccines" can be used either alone or in combination with other therapeutic regimens, including but not limited to chemotherapy, radiation therapy, surgery, bone marrow transplantation, etc. for the treatment of tumors. For example, surgical or radiation techniques could be used to debulk the tumor mass, after which, the vaccine formulations of the invention can be administered to ensure the regression and prevent the progression of remaining tumor masses or micrometastases in the body. Alternatively, administration of the "vaccine" can precede such surgical, radiation or chemotherapeutic treatment.

[0235] Alternatively, the compositions of the invention can be used to immunize or "vaccinate" tumor-free subjects to prevent tumor formation. With the advent of genetic testing, it is now possible to predict a subject's predisposition for certain cancers. Such subjects, therefore, may be immunized using a compound comprising one or more antigenic ligands derived from tumors.





- [0236] Suitable preparations of such vaccines include injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in liquid prior to injection, may also be prepared. The preparation may also be emulsified, or the polypeptides encapsulated in liposomes. The active immunogenic ingredients are often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine preparation may also include minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine.
- [0237] Compositions of the present invention which comprise a soluble CD1d complex and an immunogen may further comprise additional adjuvants. Examples of adjuvants which may be effective, include, but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine, N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine, GM-CSF, QS-21 (investigational drug, Progenics Pharmaceuticals, Inc.), DETOX (investigational drug, Ribi Pharmaceuticals), BCG, and CpG rich oligonucleotides.
- [0238] Compositions of the present invention which comprise a soluble CD1d complex and an immunogen may further comprise additional adjuvants which are also Toll-like receptor (TLR) agonists. Examples of TLR agonist adjuvants which may be effective, include, but are not limited to: N-acetylmuramyl-L-alanine-D-isoglutamine (MDP), lipopolysaccharides (LPS), genetically modified and/or degraded LPS, alum, glucan, colony stimulating factors (e.g., EPO, GM-CSF, G-CSF, M-CSF, PEGylated G-CSF, SCF, IL-3, IL-6, PIXY 321), interferons (e.g., γ -interferon, α -interferon), interleukins (e.g., IL-2, IL-7, IL-12, IL-15, IL-18), saponins (e.g., QS21), monophosphoryl lipid A (MPL), 3 De-O-acylated monophosphoryl lipid A (3D-MPL), unmethylated CpG sequences, 1-methyl tryptophan, arginase inhibitors, cyclophosphamide, antibodies that block immunosuppressive functions (e.g., anti-CTLA4 antibodies), lipids (such as palmitic acid residues), tripalmitoyl-S-glycerylcystein lyseryl-serine (P₃ CSS), and Freund's adjuvant. Other adjuvant

- 79 -

examples include compounds such as isatoribin and its derivatives (Anadys Pharmaceuticals) or imidazoquinolinamines, such as imiquimod and resiquimod (Dockrell & Kinghorn, *J. Antimicrob. Chemother.*, 48:751-755 (2001) and Hemmi *et al.*, *Nat. Immunol.*, 3:196-200 (2002), guanine ribonucleosides, such as C8-substituted or N7, C-8-disubstituted guanine ribonucleosides (Lee *et al.*, *Proc. Natl. Acad. Sci. USA*, 100:6646-6651 (2003) and the compounds that are disclosed in Pat. Pub. Nos. JP-2005-089,334; WO99/32122; WO98/01448 WO05/092893; and WO05/092892, and TLR-7 agonist SM360320 (9-benzyl-8-hydroxy-2-(2-methoxy-ethoxy)adenine) disclosed in Lee *et al.*, *Proc Natl Acad Sci USA*, 103(6):1828-1833 (2006).

[0239] In addition to isatoribin, other TLR agonist adjuvants include 9-benzyl-8-hydroxy-2-(2-methoxyethoxy)adenine (SM360320), Actilon.TM. (Coley Pharmaceutical Group, Inc.), and the following compounds by Sumitomo Pharmaceutical Co, Ltd.:

- 80 -



[0240] Other adjuvants which may be used in conjunction with the composition of the present invention are disclosed in PCT Pub. No. WO 2005/000348, U.S. Pat. Pub. No. 2007/0292418, and U.S. Pat. Pub. No. 2007/0287664.

[0241] The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc.

[0242] In an alternate embodiment, compositions of the present invention may be used in adoptive immunotherapeutic methods for the activation of NKT lymphocytes that are histocompatible with the patient. (for methods of adoptive immunotherapy, see, *e.g.*, Rosenberg, U.S. Patent No. 4,690,915, issued September 1, 1987; Zarling, *et al.*, U.S. Patent No. 5,081,029, issued January 14, 1992). Such NKT lymphocytes may be isolated from the patient or a histocompatible donor. The NKT lymphocytes are activated *in vitro* by exposure to a composition of the invention. Activated NKT lymphocytes are expanded and inoculated into the patient in order to transfer NKT cell immunity directed against the particular antigenic peptide or peptides.

[0243] The compositions of the present invention may further comprise other compounds which modulate an immune response, for example, cytokines. The term "cytokine" refers to polypeptides, including, but not limited to, interleukins (*e.g.*, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, and IL-18), α interferons (*e.g.*, IFN- α), β interferon (IFN- β), γ interferons (*e.g.*, IFN- γ), , colony stimulating factors (CSFs, *e.g.*, CSF-1, CSF-2, and CSF-3), granulocyte- macrophage colony stimulating factor (GMCSF), transforming growth factor (TGF, *e.g.*, TGF α and TGF β), and insulin-like growth factors (IGFs, *e.g.*, IGF-I and IGF-II).

[0244] In certain embodiments, therapeutic compositions useful in systemic administration, include soluble CD1d complexes of the present invention complexed to a delivery vehicle. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site, for example, ligands for targeting the vehicle to a tissue of interest. Targeting vehicles for other tissues and organs are well known to skilled artisans. In other embodiments, soluble CD1d complexes of the present invention are non-specific, *i.e.*, they are not targeted to any particular tissue, organ, cell, or cell surface marker.

[0245] Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling *et al.*, Proc. Natl. Acad. Sci. USA 189:11277-11281, 1992, which is incorporated herein by reference). Oral

- 82 -

delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (*e.g.*, DMSO) that is capable of passing into the skin.

[0246] Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian.

[0247] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, *Molecular Cloning A Laboratory Manual*, 2nd Ed., Sambrook *et al.*, ed., Cold Spring Harbor Laboratory Press: (1989); *Molecular Cloning: A Laboratory Manual*, Sambrook *et al.*, ed., Cold Springs Harbor Laboratory, New York (1992), *DNA Cloning*, D. N. Glover ed., Volumes I and II (1985); *Oligonucleotide Synthesis*, M. J. Gait ed., (1984); Mullis *et al.* U.S. Pat. No: 4,683,195; *Nucleic Acid Hybridization*, B. D. Hames & S. J. Higgins eds. (1984); *Transcription And Translation*, B. D. Hames & S. J. Higgins eds. (1984); *Culture Of Animal Cells*, R. I. Freshney, Alan R. Liss, Inc., (1987); *Immobilized Cells And Enzymes*, IRL Press, (1986); B. Perbal, *A Practical Guide To Molecular Cloning* (1984); the treatise, *Methods In Enzymology*, Academic Press, Inc., N.Y.; *Gene Transfer Vectors For Mammalian Cells*, J. H. Miller and M. P. Calos eds., Cold Spring Harbor Laboratory (1987); *Methods In Enzymology*, Vols. 154 and 155 (Wu *et al.* eds.); *Immunochemical Methods In Cell And Molecular Biology*, Mayer and Walker, eds., Academic Press, London (1987); *Handbook Of*

Experimental Immunology, Volumes I-IV, D. M. Weir and C. C. Blackwell, eds., (1986); Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); and in Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, Maryland (1989).

[0248] General principles of antibody engineering are set forth in Antibody Engineering, 2nd edition, C.A.K. Borrebaeck, Ed., Oxford Univ. Press (1995). General principles of protein engineering are set forth in Protein Engineering, A Practical Approach, Rickwood, D., *et al.*, Eds., IRL Press at Oxford Univ. Press, Oxford, Eng. (1995). General principles of antibodies and antibody-hapten binding are set forth in: Nisonoff, A., Molecular Immunology, 2nd ed., Sinauer Associates, Sunderland, MA (1984); and Steward, M.W., Antibodies, Their Structure and Function, Chapman and Hall, New York, NY (1984). Additionally, standard methods in immunology known in the art and not specifically described are generally followed as in Current Protocols in Immunology, John Wiley & Sons, New York; Stites *et al.* (eds), Basic and Clinical -Immunology (8th ed.), Appleton & Lange, Norwalk, CT (1994) and Mishell and Shiigi (eds), Selected Methods in Cellular Immunology, W.H. Freeman and Co., New York (1980).

[0249] Standard reference works setting forth general principles of immunology include Current Protocols in Immunology, John Wiley & Sons, New York; Klein, J., Immunology: The Science of Self-Nonself Discrimination, John Wiley & Sons, New York (1982); Kennett, R., *et al.*, eds., Monoclonal Antibodies, Hybridoma: A New Dimension in Biological Analyses, Plenum Press, New York (1980); Campbell, A., "Monoclonal Antibody Technology" in Burden, R., *et al.*, eds., Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 13, Elsevier, Amsterdam (1984), Kubly Immunology 4th ed. Ed. Richard A. Goldsby, Thomas J. Kindt and Barbara A. Osborne, H. Freeman & Co. (2000); Roitt, I., Brostoff, J. and Male D., Immunology 6th ed. London: Mosby (2001); Abbas A., Abul, A. and Lichtman, A., Cellular and Molecular Immunology Ed. 5, Elsevier Health Sciences Division (2005); Kontermann and Dubel, Antibody Engineering, Springer Verlag (2001); Sambrook and Russell, Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press (2001); Lewin, Genes VIII, Prentice Hall (2003); Harlow and Lane, Antibodies: A

- 84 -

Laboratory Manual, Cold Spring Harbor Press (1988); Dieffenbach and Dveksler, PCR Primer Cold Spring Harbor Press (2003).

[0250] All of the references cited above, as well as all references cited herein, are incorporated herein by reference in their entireties.

Examples

Materials and Methods

[0251] **Construction of soluble β 2m-CD1d ("CD1d") and β 2m-CD1d-4D5 scFv fusion ("CD1d fusion") proteins.** Mouse beta2-microglobulin (β 2m), soluble CD1, and the murine anti-HER2 antibody single chain 4D5 scFv were cloned by PCR. Total RNA was extracted from the CD1d transfected mouse RMA.S cell line using the RNeasy Mini Kit (QIAGEN). For the anti-HER2 antibody part, the plasmid pIG6-4D5 containing the scFv fragment derived from the mouse anti-HER2 antibody 4D5 was used as template (Worn, A., and Pluckthun, A. *FEBS Lett*: 427, 357-361 (1998)). Briefly, the entire mouse β 2m was amplified with a Hind III site at the N-terminus for subsequent cloning in the PEAK 8 expression vector (EdgeBioSystems, MD, USA) and an Nhe I site at its C-terminus for its ligation to the N-terminal sequence of the α 1 domain of CD1d with the insertion of a sequence encoding a flexible glycine/serine-rich peptide linker (GGGGSGGSGSGGG (SEQ ID NO:12)). The primers used for this PCR were 5'-TTAAGCTTATGGCTCGCTCGGTGA (SEQ ID NO:13) and 5'-AAGATATCGCTAGCTCCACCTCCAGAACCGGATCCACCTGATCCACCTC CACCCATGTCTCGATCCCAGTAGA (SEQ ID NO:14). The C-terminus of the soluble CD1d fragment was either directly fused to a 6xHis tag via a small linker (SSGSGG (SEQ ID NO:15)) (for soluble β 2m-CD1d) or ligated to the N-terminus of the 4D5 scFv fragment via the same flexible linker as above (GGGGSGGSGSGGG (SEQ ID NO:12)) and a Nar I restriction site (for the CD1d-4D5 fusion). The primers for this PCR were 5'-TTCTCGAGGCTAGCCAGCAAAA GAATTACACCTTC (sense, SEQ ID NO:16) and 5'-TTGAATTCGGCGCCTCCACCTCCAGAACCGGATCCACCTGATCCACCTCCA

- 85 -

CCGCCCACGGGTGCTTGCCTGGCAT (reverse, SEQ ID NO:17). The DNA fragment for the anti-HER2 4D5 scFv was fused at the C-terminus to the small glycine-/serine-rich linker (SSGSGG (SEQ ID NO:18) followed by the 6xHis tag, a stop codon and a Not I site for subcloning. The PCR primers were 5'-TTCTCGAGGGCGCCGACTACAAAGATATCGTTAT (sense, SEQ ID NO:19) and 5'-AAGCGGCCGCTTAATGGTGGTGATGATGATGTCCTCCAGAACCAGAAGAA ACGGTAACGGTGGTA (reverse, SEQ ID NO:20). PCR was performed using Pwo Polymerase (Roche) and amplified DNA fragments were cloned into pCR®-Blunt vector (Invitrogen) and sequenced to ensure no mutation was introduced. Using the described restriction sites, a two or three-part ligation reaction was performed to join the β 2m and linker to the soluble CD1d and, in the case of the fusion, to the 4D5 scFv DNA part with concomitant subcloning into the pEAK8 expression vector (EdgeBiosystems).

[0252] **Recombinant protein production by transient transfection.** The human cell line HEK293EBNA was adapted to serum-free suspension growth in Excell™-293 medium (JRH Biosciences, Lexington, KS) in 1-liter glass bottles placed on an orbital shaker (Kühner AG, Switzerland). For large scale transfection in suspension cultures, cells were seeded in serum-free RPMI medium (with 25 mM Hepes, Cambrex Biosciences, Verviers, Belgium) at a density of 2×10^6 cells/ml, and transfected using linear 25-kD polyethyleneimine, as described in Baldi, L., *et al. Biotechnol Prog* 21:148-153 (2005). The addition to the DNA/PEI mix of 1% (corresponding to 25 ng/ml) pEGFP-N1 plasmid DNA (Clontech, Palo Alto, CA) allowed direct visual estimation of transfection efficiency under a fluorescence light microscope (Zeiss Axiovert). Four hours post-transfection, the culture was diluted by adding one volume of Pro-293s medium (Cambrex). After 6 days, the culture was centrifuged, and the supernatant was saved for protein purification.

[0253] **Affinity purification of recombinant proteins and α GalCer loading on CD1d.** The His-tagged soluble CD1d and the CD1d-4D5 fusion proteins were purified from the HEK293 supernatants using Ni-NTA resin batch-wise (Ni-NTA Superflow, QIAGEN) and bound proteins were eluted with 0.25 M Imidazole. Purity was analyzed on a 10% SDS-PAGE. Depending the batch, the yield could reach up to

- 86 -

10mg/L supernatant of pure soluble or anti-HER2 CD1d fusion proteins. After purification, CD1d was loaded overnight at RT with a three-fold molar excess of α GalCer and the unbound glycolipid was removed by FPLC (Superdex™ 200 or Sephacryl S100, Pharmacia Biotech). Binding to the HER2 tumor antigen and proper folding of the CD1d protein were assessed on the B16-HER2 and SKBR3 target cell lines by flow cytometry using either an anti-His or anti-CD1d mAbs (BD Biosciences).

[0254] Generation of CD1d tetramer. For this purpose, the soluble β 2m-CD1d molecule was modified at the C-terminus by the addition of a cysteine after the stretch of histidine residues. After a mild reduction with 0.5 mM β -mercaptoethanol for 30 min at 30C and purification on a PD10 desalting column, the pure recombinant protein was biotinylated on the cystein residue by the chemical coupling of a biotin-maleimide linker (EZ-linked BM, Pierce) in 2M excess overnight at RT. The excess of linker was removed by gel filtration on a Superdex S200 column (Pharmacia Biotech). The biotinylated CD1d was loaded with α GalCer as described above and was tetramerized on Extravidin-PE (Sigma) and resulting complex was used at 5 μ g/ml for NKT cell staining.

[0255] Mice, cell lines and antibodies. Female mice C57BL/6, 6-8 weeks old, were purchased from Harlan (Zeist, Holland). The B16-F10 melanoma cell line (ATCC/CRL-6475) was stably transfected with the human HER2 antigen (Cesson, V., *et al. Clin Cancer Res* 12:7422-7430 (2006)). Transfected cells were selected with 1.2 mg/ml G418 and grafted intravenously (i.v.) into naïve mice. High HER2 expressing clones were established from lung metastases and expanded in DMEM supplied with 10% FCS, antibiotics and 1.2 mg/ml G418. Expression of HER2 was monitored by flow cytometry (BD FACScan) with 10 μ g/ml of the humanized anti-HER2 monoclonal antibody Herceptin (Trastuzumab, Hoffmann-La-Roche) and goat anti-human IgG-FITC (Sigma). All other antibodies, unless specified, were from Becton Dickinson (BD Biosciences). The software used was Cell Quest (BD Biosciences).

[0256] Isolation of liver, lung and spleen lymphocytes. Mouse livers and lungs were homogenized with 100 μ m strainer and lymphocytes were isolated using a

- 87 -

Percoll density gradient (Amersham Biosciences). After two washings, cells were either re-challenged or directly analyzed by fluorescence cytometry. Spleens were homogenized with a 70 μ m strainer, debris were eliminated by spontaneous sedimentation and splenocytes were recovered by centrifugation. They were then depleted from B cells by incubation with anti-CD19 MACS microbeads and elution on LD columns according to manufacturer recommendation (Myltenyi Biotec).

[0257] **Serum cytokines.** Serum TNF α was measured by the Quantikine immunassay kit from R&D Systems Inc. (Minneapolis, USA).

[0258] **Quantification of lung metastasis.** Metastatic nodule surface area was measured on 2560x1920 pixels photographs of the whole organs, taken on an Zeiss Stemi SV11 dissection microscope (Carl Zeiss Jena Germany) equipped with a ProgRes-C10plus Color Camera (Jenoptik, Jena, Germany). Each image was analyzed with the ImageJ program (rsb.info.nih.gov/ij/) using a k-means clustering algorithm (plugin available on ij-plugins.sourceforge.net). Images were segmented into 9 segments (cluster tolerance 1x10⁻⁴, randomization seed: 48) and the appropriate segments were selected based on the color of the metastasis reaching the surface of the organ. The area of the sum of selected color segments was expressed as percentage of the area occupied by the lungs. Both sides of the lung were analyzed. This method takes into account both the number and the size of the tumor nodules. Since nodules are often heterogenous the percentage of metastatic lung surface area is a more sensitive measure of tumor growth than just counting tumor nodules.

Example 1

Sustained activation of iNKT cells with repeated injections of recombinant α GalCer-loaded CD1d molecules

[0259] This Example demonstrates that a soluble monomeric form of CD1d loaded with α GalCer can fully activate mouse iNKT cells *in vitro* and *in vivo* and that this activation is due to the complex per se and not to the *in vivo* release of α GalCer. In particular, this Example shows that iNKT cells remain responsive following repeated injections of α GalCer/sCD1d and α GalCer/CD1d-anti HER2 (α GalCer/sCD1d linked

- 88 -

to a scFv fragment of anti-Her2 antibody). This is in sharp contrast to stimulation with α GalCer alone, which was previously shown to induce unresponsiveness after a single injection. Parameters of iNKT activation include invTCR (the invariant T cell receptor of NKT cells) downmodulation, production of IFN γ , increased iNKT frequency during systemic treatment and resistance to tumor development after pretreatment.

[0260] **InvTCR downmodulation:** free α GalCer (5 μ g), sCD1d (20 μ g) and CD1d-anti HER2 fusion (40 μ g), loaded or not with α GalCer, were injected intraperitoneally into 6 groups of 5 mice. Mice were sacrificed 20 hours later. Liver lymphocytes were prepared as described above and stained with α GalCer/CD1d tetramer + anti-CD3 for analysis by FACS. **Figure 1** shows that inv TCR downmodulation occurred following a single injection of either free α GalCer or recombinant CD1d molecules loaded with α GalCer. The following experiments demonstrate that this result is not due to the *in vivo* release of α GalCer but is due to direct stimulation by the complex per se since the outcome of the two modes of stimulation differ.

[0261] **Sustained production of IFN γ after repeated i.v. injections:** cytokine production was tested by IntraCellular Cytokine Staining (ICCS). Liver and spleen lymphocytes (2x10⁶/ml) from treated mice (5 to 6 i.v. injections every three days) were activated "*in vitro*" or "*ex vivo*" by either 200ng/ml α GalCer or by α GalCer/CD1d-anti HER2 fusion (10 μ g/ml) bound to plastic-coated anti-His antibody. Golgi Plug (BD) was added after 1 hour to block secretion and after a total of 6 hours incubation, activated lymphocytes were stained with different antibody combinations to gate on NKT cells. Anti-CD3-FITC was generally tested with α GalCer-CD1d tetramer-PE or, in case of NKT TCR downmodulation, anti-CD3-FITC was used with NK1.1-PerCP. After fixation and permeabilization with Cytofix/Cytoperm (BD), intracellular IFN γ was detected with an APC-labeled anti-IFN γ monoclonal antibody. Cells were analyzed by flow cytometry on a FACS Calibur (Cell Quest Software; BD). For *ex vivo* measurement following an i.v. injection, the procedure was similar except that the six hour incubation was omitted. **Figure 2** summarizes the results. For "*ex vivo*" measurements, liver and spleen cells

were analyzed (**Fig. 2a**), whereas for “*in vitro*” analysis, data show only liver cells (**Fig. 2b**). Results demonstrate that in the two settings, the α GalCer/CD1d-anti HER2 fusion and the soluble α GalCer/sCD1d stimulate iNKT to produce IFN γ following repeated *in vivo* stimulation with the same material, whereas free α GalCer is only active in naïve (PBS control) mice and has no more effect in α GalCer-treated mice. No significant amount of IL-4 was measured in any of the mice (data not shown), suggesting that under these conditions, activated iNKT cells develop a pro-inflammatory cytokine bias.

[0262] **Increased frequency of iNKT cells during systemic treatment:** normally iNKT cells are not detectable in mouse peripheral blood as shown on **Figure 3**. In contrast, they became clearly detectable upon systemic treatment with recombinant α GalCer/CD1d molecules, whereas there was no change with α GalCer alone. **Figure 3a** gives representative dot plots of FACS staining with CD1d-tetramer+anti-CD3 on PBMC 3 days after the third injection with either PBS, free α GalCer, α GalCer/CD1d-anti HER2, or α GalCer/sCD1d (five mice per group, i.v., injection every 3 days). Percentages of iNKT are given in **Figure 3b**.

[0263] **Pretreatment generates tumor protection:** the sustained activation of iNKT cells by recombinant α GalCer/CD1d molecules was further demonstrated by the following experiment. Pretreatment of mice with five sequential i.v. injections of α GalCer/sCD1d (5x25 μ g every 3 days) rendered them resistant to B16 melanoma cells (700,000 cells) co-injected with α GalCer/sCD1d (25 μ g) (**Fig. 4**). As previously reported by Parekh *et al.* Parekh, V.V. *et al. J Clin Invest* 115:2572-2583 (2005), mice pretreated with free α GalCer (5x0.4 μ g) were not able to block lung metastasis when α GalCer was co-injected with the tumor cells. This is most likely due to the anergic state of iNKT cells induced by free α GalCer. In contrast, in naïve mice, opposite results were obtained. Co-injection of α GalCer alone completely blocked tumor development, while co-injection of α GalCer/sCD1d had no effect. The key finding is that multiple prior injections of α GalCer loaded recombinant CD1d molecules but not of free α GalCer confers resistance to tumor development. This supports a sustained

- 90 -

activation of iNKT cells exhibiting cytotoxic activity against tumor cells directly and/or indirectly through activation of other cells such as NK cells.

Example 2

The α GalCer/CD1d-anti HER2 fusion protein has anti tumor activity when targeted to HER2-expressing tumors

[0264] This Example demonstrates that the sustained activation of iNKT cells can be redirected to the tumor site by fusing the CD1d to an anti-tumor antibody fragment.

[0265] **Precoating experiments:** as a first approach to test the anti-tumor activity of the α GalCer/CD1d fused or not fused to the anti HER2 scFv, B16 melanoma cells, wild type or stably transfected with the human HER2 antigen, were pre-incubated with either (i) α GalCer alone; (ii) α GalCer/CD1d-anti HER2 fusion; (iii) α GalCer/sCD1d; or (iv) intact anti HER2 mAb (Herceptin), before being injected i.v. into naïve mice (**Fig. 5a,b**). As already reported Kawano, T. *et al. Science* 278:1626-1629 (1997), co-injection of α GalCer with tumor cells completely inhibited tumor development, whether the tumor cells expressed or did not express the HER2 antigen. This affect may be caused by a transient association of the α GalCer with the tumor cell surface, or by transient uptake of the free α GalCer. In contrast, the α GalCer/CD1d-anti HER2 fusion inhibited tumor metastases only when HER2 was expressed on the tumor cells. This effect was maintained even after several washings of the tumor cells, indicating that the anti tumor effect was due to specifically bound anti-HER2 fusion protein. Intact anti-HER2 mAb was unable to inhibit lung metastases of B16-HER2 tumor cells, supporting that the antitumor effect of the bound fusion protein was NKT cell mediated. Soluble α GalCer/sCD1d was also not able to block tumor growth of wild-type B16 melanoma cells confirming that recombinant CD1d molecules need to be bound to the tumor cell to block metastasis development. In these precoating settings, the anti tumor activity of the CD1d-anti HER2 was not superior to the already optimal effect of free α GalCer co-injected with tumor cells. However, the dependence on HER2 binding suggests that the CD1d-anti HER2 fusion can be efficiently targeted to HER2-expressing cancer cells and may

- 91 -

redirect iNKT cells to the tumor site. In addition, these results demonstrate that the HER2-dependent anti tumor activity is due to the α GalCer/CD1d-anti HER2 per se and not to *in vivo* release of α GalCer.

[0266] **Systemic treatment:** In view of the success obtained in precoating experiments, systemic treatments were started at different time points after the injection of B16-HER2 melanoma cells. **Figure 6a** illustrates results obtained when treatment was initiated 48 hours after injecting B16-HER2 tumor cells. The mean of tumor metastasis in each group of five mice is expressed as the percentage of lung surface invaded by melanin-loaded nodules quantified as described in Materials and Methods. Free α GalCer administered in a series of 5 injections starting two days after tumor graft had no significant anti tumor effect. In contrast, α GalCer/CD1d-anti HER2 fusion protein administered on the same schedule had a potent anti metastatic effect with an average of 7% of the lungs invaded by melanin as compared with 35% in untreated animals ($p < 0.005$). Even when treatment was started six days after injection of the tumor cells, the α GalCer/CD1d-anti HER2 fusion protein still had a significant anti tumor effect with 60% less metastasis than in untreated or free α GalCer-treated mice ($p < 0.01$) (**Fig. 6b**). Interestingly, treatment with α GalCer/sCD1d had anti tumor activity which was very variable from mouse to mouse as shown by the large standard deviation (**Fig. 6a**). This in spite of the fact that the experiment of NKT TCR down-modulation did demonstrate that soluble α GalCer/sCD1d was functional to the same extent as the CD1d-anti HER2 fusion (**Fig. 1**).

[0267] Altogether, the efficient anti tumor activity obtained with the α GalCer/CD1d-anti HER2 fusion protein indicates that activated iNKT cells can be efficiently re-directed to the tumor site. Most importantly, these experiments again demonstrate that NKT cells can be repeatedly stimulated with α GalCer administered in association with CD1d, in this case, linked to anti-HER2 scFv, and also untargeted, as illustrated in Example 1.

Example 3

iNKT activated by recombinant CD1d molecules retain their capacity to transactivate NK cells, DC and T lymphocytes

- [0268] In absence of a specific antigen: Several reports have described the capacity of iNKT cells to transactivate NK, DC and T cells upon activation with free α GalCer or α GalCer-pulsed dendritic cells (Nieda, M., *et al. Blood* 103:383-389 (2004); Hermans, I.F., *et al. J Immunol* 171:5140-5147 (2003); Smyth, M.J., *et al. J Exp Med* 201:1973-1985 (2005)). This Example demonstrates that these modulating properties of iNKT were retained upon their activation with recombinant α GalCer/sCD1d complexes. Regarding NK cells, their frequency indeed increased after a single injection of free α GalCer or of α GalCer/sCD1d with or without anti HER2, as shown by the increase of the NK1.1⁺ CD3⁻ cell population (**Fig. 7a**). Markers for DC maturation were also analyzed after 5 days *in vitro* culture of splenocytes isolated at the end of a systemic treatment, as described in Example 1. The percentage of CD11c⁺ CD40⁺ double positive cells was increased after *in vitro* stimulation with α GalCer/sCD1d indicating that activated iNKT cells promote DC maturation (**Fig. 7b**). This positive effect of α GalCer/sCD1d on DC *in vitro* was seen in splenocytes from α GalCer or α GalCer/sCD1d-treated mice, whereas α GalCer alone had no effect in any of the mice. The conventional T cell population of CD3⁺ cells negative for CD1d tetramer and NK1.1, was analyzed in the same splenocytes cultures and it was significantly increased after *in vitro* stimulation by recombinant α GalCer/sCD1d complex while α GalCer alone had no significant effect (**Fig. 7c**). From 7% of total spleen cells without stimulation, percentage of CD3⁺ cells increased to 17, 27 and 30%, respectively in naïve, α GalCer and α GalCer/sCD1d treated mice. In contrast, α GalCer alone had no significant effect in any of the mice (**Fig. 7c**).
- [0269] **Together with active OVA immunization**: An adjuvant effect has previously been attributed to free α GalCer and its potential use in vaccination has been proposed (Silk, J.D., *et al. J Clin Invest* 114:1800-1811 (2004)). The present data suggest that α GalCer/sCD1d would have superior adjuvant properties compared to free α GalCer.

- 93 -

In this context, the adjuvant effect on the expansion of antigen-specific T cells in the model of C57 BL/6 mice immunized with ovalbumin was investigated (Hermans, I.F., *et al. J Immunol* 171:5140-5147 (2003)). After priming, specific H-2K^b/OVA₂₅₇₋₂₆₄ tetramer CD8 double positive T cells could be detected only in mice that received 200 µg ovalbumin together with αGalCer/sCD1d, whereas no H-2K^b/OVA₂₅₇₋₂₆₄ tetramer CD8 double positive T cells could be detected in mice primed with the same amount of antigen together with montanide or free αGalCer as adjuvant, (**Fig. 8a**). Furthermore, after boosting these same mice with OVA peptide in the same adjuvants *in vivo* and then 5 days *in vitro* culture without further stimulation, the frequency of K^b/OVA specific CTLs in the spleens reached up to 45% of total CD8 in mice that received a boost of OVA peptide with αGalCer/sCD1d, whereas the frequency was 12% in mice boosted with peptide and αGalCer alone (**Fig. 8b**). A further increase could be obtained by *in vitro* re-stimulation with the same respective stimuli and the frequency of specific T cells reached close to 70% in both αGalCer and αGalCer/sCD1d-treated mice (**Fig. 8b**). In the same cultures, there was also an increased frequency of mature DC following restimulation with αGalCer/sCD1d but not free αGalCer as shown by CD11c+CD40+ FACS staining (**Fig. 8c**)

[0270] The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.

- 94 -

WHAT IS CLAIMED IS:

1. A method of modulating an immune response in an animal, comprising administering to an animal in need of immune modulation a composition comprising:
 - (a) non-specific CD1d complex, said complex comprising:
 - (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;
 - (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and
 - (iii) a ceramide-like glycolipid antigen bound to said CD1d polypeptide; and
 - (b) a carrier;wherein said non-specific CD1d complex is administered in an amount sufficient to affect the activity of NKT cells in said animal.
2. A method of treating a disease in an animal, comprising administering to an animal with said disease a composition comprising:
 - (a) non-specific CD1d complex, said complex comprising:
 - (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;
 - (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and
 - (iii) a ceramide-like glycolipid antigen bound to said CD1d polypeptide; and

- 95 -

- (b) a carrier;

wherein said composition is administered in an amount sufficient to alter the progression of said disease.

3. A method of inhibiting an anergic effect of a ceramide-like glycolipid antigen on NKT cell activity, comprising:

stimulating NKT cells with said ceramide-like glycolipid antigen as part of a CD1d complex, said complex comprising:

- (a) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;

- (b) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and

- (c) said ceramide-like glycolipid antigen bound to said CD1d polypeptide;

and

restimulating said NKT cells one or more times with said complex;

wherein said NKT cells are activated in response to said stimulation, and wherein said NKT cells are reactivated in response to said restimulation by said complex.

4. A method of modulating an immune response to an immunogen in an animal, comprising administering to an animal in need thereof a composition comprising:

- (a) an immunogen;

- (b) a CD1d complex, said complex comprising:

- (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;

- 96 -

(ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and

(iii) a ceramide-like glycolipid antigen bound to said CD1d polypeptide; and

(c) a carrier;

wherein said CD1d complex is administered in an amount sufficient to modulate the immune response against said immunogen relative to administration of said immunogen in the absence of said CD1d complex.

5. A method of treating a disease in an animal, comprising administering to an animal in need thereof a composition comprising:

(a) an immunogen;

(b) a CD1d complex, said complex comprising:

(i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;

(ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and

(iii) a ceramide-like glycolipid antigen bound to said CD1d polypeptide; and

(c) a carrier;

wherein an immune response against said immunogen is effective in treating said disease, and wherein said CD1d complex is administered in an amount sufficient to modulate the immune response against said immunogen relative to administration of said immunogen in the absence of said CD1d complex.

- 97 -

6. A method of preventing a disease in an animal, comprising administering to an animal in need thereof a composition comprising:

- (a) an immunogen;
- (b) a CD1d complex, said complex comprising:
 - (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;
 - (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and
 - (iii) a ceramide-like glycolipid antigen bound to said CD1d polypeptide; and
- (c) a carrier;

wherein an immune response against said immunogen is effective in treating said disease, and wherein said CD1d complex is administered in an amount sufficient to modulate the immune response against said immunogen relative to administration of said immunogen in the absence of said CD1d complex.

7. The method of any one of claims 1 to 6, wherein said composition additionally comprises an adjuvant.
8. The method of claim 7, wherein said adjuvant is a Toll-like receptor (TLR) agonist selected from the group consisting of: monophosphoryl lipid A (MPL), CpG, and BCG.
9. The method of claim 8, wherein said adjuvant is MPL.

- 98 -

10. The method of any one of claims 1 to 9, wherein said soluble CD1d polypeptide comprises an amino acid sequence at least 90% identical to amino acids 21 to 295 of SEQ ID NO:1.
11. The method of claim 10, wherein said soluble CD1d polypeptide comprises an amino acid sequence at least 95% identical to amino acids 21 to 295 of SEQ ID NO:1.
12. The method of claim 11, wherein said soluble CD1d polypeptide comprises amino acids 21 to 295 of SEQ ID NO:1.
13. The method of claim 12, wherein said soluble CD1d polypeptide comprises amino acids 1 to 295 of SEQ ID NO:1.
14. The method of any one of claims 1 to 9, wherein said soluble CD1d polypeptide comprises amino acids 20 to 295 of SEQ ID NO:1, except for at least one but less than 10 conservative amino acid substitutions.
15. The method of any one of claims 1 to 14, wherein said soluble CD1d polypeptide further comprises a heterologous polypeptide.
16. The method of any one of claims 1 to 15, wherein said β 2-microglobulin polypeptide comprises an amino acid sequence at least 90% identical to amino acids 21 to 113 of SEQ ID NO:2.
17. The method of claim 16, wherein said β 2-microglobulin polypeptide comprises amino acids 21 to 113 of SEQ ID NO:2.
18. The method of claim 17, wherein said β 2-microglobulin polypeptide comprises amino acids 1 to 119 of SEQ ID NO:2.


- 99 -

19. The method of any one of claims 1 to 15, wherein said β 2-microglobulin polypeptide comprises amino acids 21 to 113 of SEQ ID NO:2, except for at least one but less than 5 conservative amino acid substitutions.
20. The method of any one of claims 1 to 19, wherein said β 2-microglobulin polypeptide further comprises a heterologous polypeptide.
21. The method of any one of claims 1 to 20, wherein said β 2-microglobulin polypeptide is associated with said soluble CD1d polypeptide via normal disulfide linkages.
22. The method of any one of claims 1 to 21, wherein said β 2-microglobulin polypeptide is attached to said soluble CD1d polypeptide as a fusion polypeptide.
23. The method of claim 22, wherein said β 2-microglobulin polypeptide is situated at the N-terminus of said soluble CD1d polypeptide.
24. The method of any one of claims 22 or 23, wherein said β 2-microglobulin polypeptide and said soluble CD1d polypeptide are separated by a peptide linker.
25. The method of any one of claims 1 to 24, wherein two or more soluble CD1d complexes are associated together as a multivalent soluble CD1d complex.
26. The method of any one of claim 1 to 25, wherein said ceramide-like glycolipid antigen comprises compound selected from the group consisting of an α -galactosylceramide and α -glycosylceramide.
27. The method of claim 26, wherein said ceramide-like glycolipid antigen comprises an α -galactosylceramide.

- 100 -

28. The method of claim 27, wherein said α -galactosylceramide comprises the formula:

Formula I



wherein


R1 is a linear or branched C_1 - C_{27} alkene with at least one $C=C$ bond but not $-(CH_2)_7CH=CH(CH_2)_7CH_3$, or R1 is $C(OH)-R3$ wherein R3 is a linear or branched C_1 - C_{26} alkene with at least one $C=C$ bond; and

R2 is one of the following (a)-(e):

- (a) $-(CH_2)(CH_2)_xCH_3$,
- (b) $-CH(OH)(CH_2)_xCH_3$,
- (c) $-CH(OH)(CH_2)_xCH(CH_3)_2$,
- (d) $-CH=CH(CH_2)_xCH_3$,
- (e) $-CH(OH)(CH_2)_xCH(CH_3)CH_2CH_3$,

wherein X is an integer ranging from 5-17.

29. The method of claim 27, wherein said ceramide-like glycolipid antigen comprises (2S, 3S, 4R)-1-O-(α -galactopyranosyl)-N-hexacosanoyl-2-amino-1,3,4-octadecanetriol (KRN7000, (2S,3S,4R)-1-(α -D-galactopyranosyloxy)-2-hexacosanoylamino-3,4-octa-decanediol), having the structure:



- 101 -

30. The method of any one of claims 1 to 28, wherein said ceramide-like glycolipid antigen is modified to have a shortened long-chain sphingosine base (C5 vs.C14) and acyl chain (C24 vs.C26).
31. The method of claim 30, wherein said modified ceramide-like glycolipid antigen is an -OCH analog with a long-chain sphingosine base shortened from C14 to C5 and acyl chain from C26 to C24.
32. The method of any one of claims 1 to 31, wherein said composition further comprises a costimulatory molecule.
33. The method of claim 32, wherein said costimulatory molecule is selected from the group consisting of: Toll Like Receptors, and B71.
34. The method of any one of claims 1 to 33, wherein said composition further comprises an agent selected from the group consisting of: a chemotherapeutic agent, an antibody, and an antigen.
35. The method of any one of claims 1 and 4 to 34, wherein said immune response is elicited.
36. The method of any one of claims 1 and 4 to 34, wherein said immune response is enhanced.
37. The method of any one of claims 1 and 4 to 36, wherein said immune response comprises release of a cytokine.
38. The method of claim 37, wherein said cytokine is selected from the group consisting of IFN-gamma, TNF-alpha, and both cytokines.

- 102 -

39. The method of claim 37, wherein said cytokine is selected from the group consisting of IL-4, IL-10, IL-5, IL-13, and a combination of said cytokines.
40. The method of any one of claims 1 and 4 to 39, wherein said immune response comprises recruitment of cells selected from the group consisting of NK cells, CTLs, B lymphocytes, dendritic cells, macrophages, neutrophils, and a combination of two or more of said cells.
41. The method of any one of claims 1 and 4 to 34, where in said immune response is inhibited.
42. The method of claim 41, wherein said immune response is inflammation.
43. The method of claim 41, wherein said immune response is autoimmunity.
44. The method of any one of claims 1, 3, and 10-34, wherein said NKT cell activity is selected from the group consisting of cell proliferation, the production of a cytokine, recruitment or activation of a non-NKT immune system cell, or a combination of two or more of said activities.
45. The method of any one of claims 1 to 44, wherein said animal is a vertebrate.
46. The method of claim 45, wherein said vertebrate is a mammal.
47. The method of claim 46, wherein said mammal is a human.
48. The method of any one of claims 1 to 47, comprising administering 1 μ g to about 30 mg of said non-specific CD1d complex to said animal.

- 103 -

49. The method of claim 3, wherein activation of said NKT cells comprises an effect selected from the group consisting of NKT cell proliferation, the production of a cytokine, recruitment or activation of a non-NKT immune system cell, or a combination of two or more of said effects.
50. The method of claim 49, wherein activation of said NKT cells comprises NKT cell proliferation.
51. The method of claim 49, wherein activation of said NKT cells comprises the production of a cytokine.
52. The method of claim 51, wherein said cytokine is selected from the group consisting of interferon-gamma, TNF-alpha, IL-4, IL-2, IL-1 β , IL-5, IL-10, IL-12, IL-13, IL-17, IL-23, TNF- β /LT, MCP-2, oncostatin-M, RANTES, and a combination of two or more of said cytokines.
53. The method of claim 51, wherein activation of said NKT cells comprises recruitment or activation of a non-NKT immune system cell.
54. The method of claim 49, wherein said non-NKT immune system cell is selected from the group consisting of an NK cell, a CD4⁺ T cell, a CD8⁺ T cell, a B cell, a dendritic cell, and a combination of two or more of said cells.
55. The method of claim 3, wherein reactivation of said NKT cells comprises an effect selected from the group consisting of NKT cell proliferation, the production of a cytokine, recruitment or activation of a non-NKT immune system cell, or a combination of two or more of said effects.
56. The method of claim 55, wherein reactivation of said NKT cells comprises NKT cell proliferation.

- 104 -

57. The method of claim 55, wherein reactivation of said NKT cells comprises the production of a cytokine.
58. The method of claim 57, wherein said cytokine is selected from the group consisting of interferon-gamma, TNF-alpha, IL-4, IL-2 IFN gamma, IL-4, IL-5, IL-10, and IL-13, IL-1 β , IL-12, IL-17, IL-23, TNF- β /LT, MCP-2, oncostatin-M, RANTES, and a combination of two or more of said cytokines.
59. The method of claim 55, wherein reactivation of said NKT cells comprises recruitment or activation of a non-NKT immune system cell.
60. The method of claim 59, wherein said non-NKT immune system cell is selected from the group consisting of an NK cell, a CD4+ T cell, a CD8+ T cell, a dendritic cell, a macrophage, a neutrophil, and a combination of two or more of said cells.
61. The method of any one of claims 3 and 6 to 60, wherein said CD1d complex is non-specific.
62. The method of any one of claims 3 and 6 to 60, wherein said CD1d complex further comprises a targeting antibody or fragment thereof associated with said soluble CD1d polypeptide, wherein said antibody or fragment thereof is specific for a cell surface marker.
63. The method of claim 62, wherein said antibody or fragment thereof is an scFv fragment.
64. The method of claim 62, wherein said antibody or fragment thereof is a F(ab) fragment.

- 105 -

65. The method of claim 59, wherein said antibody or fragment thereof is a F(ab')₂ fragment.
66. The method of claim 62, wherein said antibody or fragment thereof is a full-length antibody.
67. The method of claim 62, wherein said cell surface marker is a cell surface marker of tumor cells.
68. The method of claim 62, wherein said cell surface marker is a cell surface marker of dendritic cells.
69. The method of claim 62, wherein said cell surface marker is a cell surface marker of a target of autoimmune or inflammatory disease.
70. The method of claim 62, wherein said cell surface marker is a cell surface marker of an infected cell or tissue.
71. The method of any one of claims 4 to 69, wherein said immunogen is selected from the group consisting of a bacterial polypeptide, a fungal polypeptide, a parasite polypeptide, an allergenic polypeptide, a tumor specific polypeptide, fragments, variants, derivatives of any of said polypeptides, membrane lipids, glycolipids, glycosylceramides, and a combination of two or more of said immunogens.
72. The method of any one of claims 2, 5, and 6 to 70, wherein said disease is associated with an infectious agent.
73. The method of any one of claims 2, 5, and 6 to 70, wherein said disease is associated with abnormal cell growth.

- 106 -

74. The method of any one of claims 2, 5, and 6 to 70, wherein said disease is an autoimmune disease.
75. The method of any one of claims 2, 5, and 6 to 70, wherein said disease is an inflammatory disease.
76. The method of any one of claims 2, 5, and 10 to 74 wherein said treating comprises, curing, altering disease progression, prolonging survival, alleviating symptoms, or inhibiting symptoms associated with said disease.
77. The method of any one of claims 1 to 76, wherein said administration is by a route selected from the group consisting of intramuscularly, intravenously, intratracheally, intranasally, transdermally, interdermally, subcutaneously, intraocularly, vaginally, rectally, intraperitoneally, intraintestinally, by inhalation, or by a combination of two or more of said routes.
78. An composition comprising:
 - (a) an immunogen;
 - (b) a CD1d complex, said complex comprising:
 - (i) an isolated soluble CD1d polypeptide sufficient to associate with β 2-microglobulin and bind a ceramide-like glycolipid antigen;
 - (ii) an isolated polypeptide comprising β 2-microglobulin or a fragment thereof associated with said CD1d polypeptide; and
 - (iii) a ceramide-like glycolipid antigen bound to said CD1d polypeptide; and
 - (c) a carrier.

- 107 -

79. The composition of claim 78, wherein the composition additionally comprises an adjuvant.
80. The composition of claim 79, wherein the adjuvant is a TLR agonist selected from the group consisting of: MPL, CpG, and BCG.
81. The composition of claim 80, wherein the adjuvant is MPL.


Figure 1**TCR downmodulation upon iNKT activation**



Figure 2**Sustained production of IFN γ upon repeated stimulation of iNKT
with α GalCer-loaded CD1d molecules****a*****Ex vivo* analysis****b*****In vitro* re-challenge**

Figure 3

Higher frequencies of iNKT cells in the blood upon systemic treatment with recombinant CD1d molecules





Figure 4**Pretreatment of mice with aGalCer-loaded sCD1d confers protection against tumor development**

Figure 5

α GalCer/CD1d-anti HER2 fusion protein blocks tumor development only when bound to HER2-expressing cancer cells




Figure 6

Systemic treatment with the α GalCer/CD1d-anti HER2 fusion protein inhibits lung metastasis development induced by B16-HER2 cells



a**b**

Figure 7
 α GalCer/CD1d molecules promote innate and adaptive immune responses induced by activated iNKT cells




Figure 8
 α GalCer/sCD1d protein acts as adjuvant in the development of an antigen-specific immune response

