



HU000027098T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 027 098**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 09 718467**
(22) A bejelentés napja: **2009. 03. 05.**
(96) Az európai bejelentés bejelentési száma:
EP 20090718467
(97) Az európai bejelentés közzétételi adatai:
EP 2254870 A2 **2009. 09. 11.**
(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2254870 B1 **2016. 02. 24.**
(51) Int. Cl.: **C07D 231/56** (2006.01)
A61K 314/15 (2006.01)
A61P 13/00 (2006.01)
A61P 25/00 (2006.01)
A61P 29/00 (2006.01)
(86) A nemzetközi (PCT) bejelentési szám:
PCT/EP 09/052590
(87) A nemzetközi közzétételi szám:
WO 09109618

(30) Elsőbbségi adatok: 08425139 2008. 03. 07. EP	(73) Jogosult(ak): Aziende Chimiche Riunite Angelini Francesco A.C.R.A.F. S.p.A., 00181 Roma (IT)
(72) Feltaláló(k): GUGLIELMOTTI, Angelo, I-00149 Roma (IT) FURLOTTI, Guido, I-00151 Roma (IT) MANGANO, Giorgina, I-00139 Roma (IT) CAZZOLLA, Nicola, I-00041 Albano Laziale (Roma) (IT) GAROFALO, Barbara, I-00179 Roma (IT)	(74) Képviselő: SBGK Szabadalmi Ügyvivői Iroda, Budapest

- (54) **1-Benzil-3-hidroximetilindazol-származékok és azok alkalmazása MCP-1, CX3CR1 expresszió alapú betegségek kezelésében**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 254 870 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
24.02.2016 Bulletin 2016/08

(51) Int Cl.:
C07D 231/56 (2006.01) **A61K 31/415** (2006.01)
A61P 13/00 (2006.01) **A61P 25/00** (2006.01)
A61P 29/00 (2006.01)

(21) Application number: **09718467.5**

(86) International application number:
PCT/EP2009/052590

(22) Date of filing: **05.03.2009**

(87) International publication number:
WO 2009/109618 (11.09.2009 Gazette 2009/37)

(54) **1-BENZYL-3-HYDROXYMETHYLINDAZOLE DERIVATIVES AND USE THEREOF IN THE TREATMENT OF DISEASES BASED ON THE EXPRESSION OF MCP-1, CX3CR1**

1-BENZYL-3-HYDROXYMETHYLINDAZOL DERIVATE UND DEREN VERWENDUNG ZUR BEHANDLUNG VON ERKRANKUNGEN IN VERBINDUNG MIT DER EXPRESSION VON MCP-1, CX3CR1

DÉRIVÉS DE 1-BENZYL-3-HYDROXYMETHYLINDAZOLE ET LEUR UTILISATION POUR TRAITER LES TROUBLES LIÉS À L'EXPRESSION DU MCP-1, CX3CR1

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR

(30) Priority: **07.03.2008 EP 08425139**

(43) Date of publication of application:
01.12.2010 Bulletin 2010/48

(73) Proprietor: **Aziende Chimiche Riunite Angelini
Francesco A.C.R.A.F. S.p.A.
00181 Roma (IT)**

(72) Inventors:
• **GUGLIELMOTTI, Angelo**
I-00149 Roma (IT)
• **FURLOTTI, Guido**
I-00151 Roma (IT)
• **MANGANO, Giorgina**
I-00139 Roma (IT)
• **CAZZOLLA, Nicola**
I-00041 Albano Laziale (Roma) (IT)
• **GAROFALO, Barbara**
I-00179 Roma (IT)

(74) Representative: **Allaix, Roberto
Marchi & Partners S.r.l.
Via Vittor Pisani, 13
20124 Milano (IT)**

(56) References cited:
EP-A- 0 382 276 EP-A- 0 510 748
WO-A-99/04770

- **Silvina Gazzaniga ET AL: "Targeting Tumor-Associated Macrophages and Inhibition of MCP-1 Reduce Angiogenesis and Tumor Growth in a Human Melanoma Xenograft", Journal of Investigative Dermatology, vol. 127, no. 8, 26 April 2007 (2007-04-26), pages 2031-2041, XP055124836, ISSN: 0022-202X, DOI: 10.1038/sj.jid.5700827**
- **C Zoja ET AL: "Bindarit retards renal disease and prolongs survival in murine lupus autoimmune disease", Kidney International, 1 March 1998 (1998-03-01), pages 726-734, XP055124837, UNITED STATES DOI: 10.1046/j.1523-1755.1998.00804.x Retrieved from the Internet: URL: <http://www.ncbi.nlm.nih.gov/pubmed/9507220> [retrieved on 2014-06-24]**
- **A. Guglielmotti ET AL: "Amelioration of rat adjuvant arthritis by therapeutic treatment with bindarit, an inhibitor of MCP-1 and TNF-[alpha] production", Inflammation Research, vol. 51, no. 5, 1 May 2002 (2002-05-01), pages 252-258, XP055124839, ISSN: 1023-3830, DOI: 10.1007/PL00000301**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 254 870 B1

Description

[0001] The present invention relates to 1-benzyl-3-hydroxymethylindazole derivatives, to a pharmaceutical composition comprising them, and to their use in the treatment of diseases based on the expression of MCP-1, CX3CR1 and p40.

[0002] In particular, the present invention relates to novel 1-benzyl-3-hydroxymethylindazole derivatives according to formula (I) below, and to a pharmaceutical composition comprising them together with a pharmaceutically acceptable vehicle. In addition, the present invention relates to the use of 1-benzyl-3-hydroxymethylindazole derivatives for preparing a pharmaceutical composition that is active in the treatment of diseases based on the expression of MCP-1, CX3CR1 and p40, and to their use in a method for treating or preventing diseases based on the expression of MCP-1, CX3CR1 and p40.

BACKGROUND OF THE ART

[0003] As is known, MCP-1 (Monocyte Chemotactic Protein-1) is a protein belonging to the β subfamily of chemokines. MCP-1 has powerful chemotactic action on monocytes and exerts its action also on T lymphocytes, mastocytes and basophils (Rollins B.J., Chemokines, Blood 1997; 90: 909-928; M. Baggiolini, Chemokines and leukocyte traffic, Nature 1998; 392: 565-568).

[0004] Other chemokines belonging to the β subfamily are, for example, MCP-2 (Monocyte Chemotactic Protein-2), MCP-3, MCP-4, MIP-1 α and MIP-1 β , RANTES.

[0005] The β subfamily differs from the α subfamily in that, in the structure, the first two cysteines are adjacent for the β subfamily, whereas they are separated by an intervening amino acid for the α subfamily.

[0006] MCP-1 is produced by various types of cells (leukocytes, platelets, fibroblasts, endothelial cells and smooth muscle cells).

[0007] Among all the known chemokines, MCP-1 shows the highest specificity for monocytes and macrophages, for which it constitutes not only a chemotactic factor but also an activation stimulus, consequently inducing processes for producing numerous inflammatory factors (superoxides, arachidonic acid and derivatives, cytokines/chemokines) and amplifying the phagocytic activity.

[0008] The secretion of chemokines in general, and of MCP-1 in particular, is typically induced by various pro-inflammatory factors, for instance interleukin-1 (IL-1), interleukin-2 (IL-2), TNF α (Tumour Necrosis Factor α), interferon- γ and bacterial lipopolysaccharide (LPS).

[0009] Prevention of the inflammatory response by blocking the chemokine/chemokine receptor system represents one of the main targets of pharmacological intervention (Gerard C. and Rollins B.J., Chemokines and disease. Nature Immunol. 2001; 2:108-115).

[0010] There is much evidence to suggest that MCP-1 plays a key role during inflammatory processes and has been indicated as a new and validated target in various pathologies.

[0011] Evidence of a considerable physiopathological contribution of MCP-1 has been obtained in the case of patients with articular and renal inflammatory diseases (rheumatoid arthritis, lupus nephritis, diabetic nephropathy and rejection following transplant).

[0012] However, more recently, MCP-1 has been indicated among the factors involved in inflammatory pathologies of the CNS (multiple sclerosis, Alzheimer's disease, HIV-associated dementia) and other pathologies and conditions, with and without an obvious inflammatory component, including atopic dermatitis, colitis, interstitial lung pathologies, restenosis, atherosclerosis, complications following a surgical intervention (for instance angioplasty, arterectomy, transplant, organ and/or tissue replacement, prosthesis implant), cancer (adenomas, carcinomas and metastases) and even metabolic diseases such as insulin resistance and obesity.

[0013] In addition, despite the fact that the chemokine system is involved in controlling and overcoming viral infections, recent studies have demonstrated that the response of certain chemokines, and in particular of MCP-1, may have a harmful role in the case of host-pathogen interactions. In particular, MCP-1 has been indicated among the chemokines that contribute towards organ and tissue damage in pathologies mediated by alpha viruses characterized by monocyte/macrophage infiltration in the joints and muscles (Mahalingam S. et al. Chemokines and viruses: friend or foes? Trends in Microbiology 2003; 11: 383-391; Rulli N. et al. Ross River Virus: molecular and cellular aspects of disease pathogenesis. 2005; 107: 329-342).

[0014] Monocytes are the main precursors of macrophages and dendritic cells, and play a critical role as mediators of inflammatory processes. CX3CR1, with its ligand CX3CL1 (fractalkine), represents a key factor in regulating the migration and adhesiveness of monocytes. CX3CR1 is expressed in monocytes, whereas CX3CL1 is a transmembrane chemokine in endothelial cells. Genetic studies in man and in animal models have demonstrated an important role in the physiopathology of inflammatory diseases of CX3CR1 and CX3CL1. There is in fact much evidence to suggest a key contribution of CX3CR1 and of its ligand in the pathogenesis and progression of articular, renal, gastrointestinal and vascular inflammatory diseases (e.g. rheumatoid arthritis, lupus nephritis, diabetic nephropathy, Crohn's disease, ulcer-

ative colitis, restenosis and atherosclerosis).

[0015] The expression of CX3CR1 is over-regulated in T cells, which are believed to accumulate in the synovium of patients suffering from rheumatoid arthritis. In addition, the expression of CX3CL1 is over-regulated in endothelial cells and fibroblasts present in the synovium of these patients. Consequently, the CX3CR1/CX3CL1 system plays an important role in controlling the type of cell and the mode of infiltration of the synovium and contributes towards the pathogenesis of rheumatoid arthritis (Nanki T. et al., "Migration of CX3CR1-positive T cells producing type 1 cytokines and cytotoxic molecules into the synovium of patients with rheumatoid arthritis", *Arthritis & Rheumatism* (2002), vol. 46, No. 11, pp. 2878-2883).

[0016] In patients suffering from renal damage, the majority of the inflammatory leukocytes that infiltrate the kidneys express CX3CR1, and in particular it is expressed on two of the main cell types involved in the most common inflammatory renal pathologies and in kidney transplant rejection, T cells and monocytes (Segeer S. et al., "Expression of the fractalkine receptor (CX3CR1) in human kidney diseases", *Kidney International* (2002) 62, pp. 488-495).

[0017] Participation of the CX3CR1/CX3CL1 system has been suggested also in inflammatory bowel diseases (IBD). In point of fact, in the case of patients suffering from IBD (e.g. Crohn's disease, ulcerative colitis), a significant increase in the production of CX3CL1 by the intestinal capillary system and a significant increase in CX3CR1-positive cells have been demonstrated, both at the circulatory level and in the mucosa (Sans M. et al., "Enhanced recruitment of CX3CR1 + T cells by mucosal endothelial cell-derived fractalkine in inflammatory bowel diseases", *Gastroenterology* 2007, vol. 132, No. 1, pp. 139-153).

[0018] Even more interesting is the demonstration of the key role played by the CX3CR1/CX3CL1 system in vascular damage and in particular under pathological conditions, for instance atherosclerosis and restenosis. CX3CR1 is indicated as a critical factor in the process of infiltration and accumulation of monocytes in the vascular wall, and CX3CR1 polymorphism in man is associated with a reduced prevalence of atherosclerosis, coronary disorders and restenosis (Liu P. et al., "Cross-talk among Smad, MAPK and integrin signalling pathways enhances adventitial fibroblast functions activated by transforming growth factor-1 and inhibited by Gax" *Arterioscler. Thromb. Vasc. Biol.* 2008; McDermott D.H. et al., "Chemokine receptor mutant CX3CR1-M280 has impaired adhesive function and correlates with protection from cardiovascular diseases in humans", *J. Clin. Invest.* 2003; Niessner A. et al., *Thrombosis and Haemostasis* 2005).

[0019] IL-12 and IL-23 are members of a small family of proinflammatory heterodimeric cytokines. Both cytokines share a common subunit, p40, which is covalently bonded either to the p35 subunit to produce the mature form of IL-12, or to the p19 subunit to produce the mature form of IL-23. The receptor for IL-12 is constituted by the subunits IL-12R1 and IL-12R β 2, while the receptor for IL-23 is constituted by the subunits IL-12R β 1 and IL-23R.

[0020] IL-12 and IL-23 are mainly expressed by activated dendritic cells and by phagocytes. The receptors for the two cytokines are expressed on the T and NK cells, and NK T cells, but low levels of complexes of the receptor for IL-23 are also present in monocytes, macrophages and dendritic cells.

[0021] Despite these similarities, there is much evidence to suggest that IL-12 and IL-23 control different immunological circuits. In point of fact, whereas IL-12 controls the development of Th1 cells, which are capable of producing gamma-interferon (IFN- γ), and increases the cytotoxic, antimicrobial and antitumoral response, IL-23 regulates a circuit that leads to the generation of CD4⁺ cells, which are capable of producing IL-17. The induction of IL-23-dependent processes leads to the mobilization of various types of inflammatory cell, for instance T_H-17, and it has been demonstrated as being crucial for the pathogenesis of numerous inflammatory pathologies mediated by immunological responses.

[0022] Typical examples of pathologies associated with the expression of p40 are chronic inflammatory diseases of the articular apparatus (e.g. rheumatoid arthritis), of the dermatological apparatus (e.g. psoriasis) and of the gastrointestinal apparatus (e.g. Crohn's disease). However, IL-23 also exerts a role in promoting tumour incidence and growth. In point of fact, IL-23 regulates a series of circuits in the tumoral microenvironment, stimulating angiogenesis and the production of inflammation mediators.

[0023] Psoriasis is a chronic inflammatory skin disease that affects 3% of the world's population (Koo J. *Dermatol. Clin.* 1996; 14:485-96; Schon M.P. et al., *N. Engl. J. Med.* 2005; 352: 1899-912). A type-1 aberrant immune response has been correlated with the pathogenesis of psoriasis, and the cytokines that induce this response, such as IL-12 and IL-23, may represent suitable therapeutic objects. The expression of IL-12 and IL-23, which share the subunit p40, is significantly increased in psoriasis plaques, and preclinical studies have demonstrated a role of these cytokines in the pathogenesis of psoriasis. More recently, the treatment of anti-IL-12 and IL-23 monoclonal antibodies of patients suffering from psoriasis proved to be effective in improving the signs of progression and seriousness of the disease and has subsequently reinforced the role of IL-12 and IL-23 in the physiopathology of psoriasis.

[0024] Crohn's disease is a chronic inflammatory pathology of the digestive apparatus and may affect any region thereof - from the mouth to the anus. It typically afflicts the terminal tract of the ileum and well-defined areas of the large intestine. It is often associated with systemic autoimmune disorders, such as mouth ulcers and rheumatic arthritis. Crohn's disease affects over 500 000 people in Europe and 600 000 people in the United States.

[0025] Crohn's disease is a pathology associated with a Th1 cell-mediated excessive activity of cytokines. IL-12 is a key cytokine in the initiation of the inflammatory response mediated by Th1 cells. Crohn's disease is characterized by

increased production of IL-12 by cells presenting the antigen in intestinal tissue, and of gamma-interferon (IFN- γ) and TNF α by lymphocytes and intestinal macrophages. These cytokines induce and support the inflammatory process and thickening of the intestinal wall, which are characteristic signs of the pathology. Preclinical and clinical evidence has demonstrated that inhibition of IL-12 is effective in controlling the inflammatory response in models of intestinal inflammation and/or in patients suffering from Crohn's disease.

[0026] The relationship between cancer and inflammation is now an established fact. Many forms of tumours originate from sites of inflammation, and inflammation mediators are often produced in tumours.

[0027] IL-23 has been identified as a cytokine associated with cancer and, in particular, the expression of IL-23 is significantly high in samples of human carcinomas when compared with normal adjacent tissues. In addition, the absence of a significant expression of IL-23 in the normal adjacent tissues suggests an over-regulation of IL-23 in tumours, reinforcing its role in tumour genesis.

[0028] European patent EP-B-0 382 276 describes a number of 1-benzyl-3-hydroxymethylindazole derivatives endowed with analgesic activity. In turn, European patent EP-B-0 510 748 describes, on the other hand, the use of these derivatives for preparing a pharmaceutical composition that is active in the treatment of autoimmune diseases. Finally, European patent EP-B-1 005 332 describes the use of these derivatives for preparing a pharmaceutical composition that is active in treating diseases derived from the production of MCP-1. 2-Methyl-2-[[1-(phenylmethyl)-1H-indazol-3-yl]methoxy]propanoic acid is thought to be capable of inhibiting, in a dose-dependent manner, the production of MCP-1 and TNF- α induced in vitro in monocytes from LPS and *Candida albicans*, whereas the same compound showed no effects in the production of cytokines IL-1 and IL-6, and of chemokines IL-8, MIP-1 α , and RANTES (Sironi M. et al., "A small synthetic molecule capable of preferentially inhibiting the production of the CC chemokine monocyte chemoattractant protein-1", European Cytokine Network. Vol. 10, No. 3, 437-41, September 1999).

[0029] European patent application EP-A-1 185 528 relates to the use of triazine derivatives for inhibiting the production of IL-12. European patent application EP-A-1 188 438 and EP-A-1 199 074 relate to the use of inhibitors of the enzyme PDE4, for instance Rolipram, Ariflo and diazepam-indole derivatives, in the treatment and prevention of diseases associated with excessive production of IL-12. European patent application EP-A-1 369 119 relates to the use of hyaluronane with a molecular weight of between 600 000 and 3 000 000 daltons for controlling and inhibiting the expression of IL-12. European patent application EP-A-1 458 687 relates to the use of pyrimidine derivatives for treating diseases related to an overproduction of IL-12. European patent application EP-A-1 819 341 relates to the use of nitrogenous heterocyclic compounds, for instance pyridine, pyrimidine and triazine derivatives, for inhibiting the production of IL-12 (or of other cytokines, such as IL-23 and IL-27 which stimulate the production of IL-12). European patent application EP-A-1 827 447 relates to the use of pyrimidine derivatives for treating diseases related to an overproduction of IL-12, IL-23 and IL-27.

[0030] European patent applications EP-A-1 869 055, EP-A-1 869 056 and EP-A-1 675 862 describe 1,3-thiazolo-4,5-pyrimidine derivatives that are capable of acting as CX3CR1 receptor antagonists.

[0031] Despite the activity developed thus far, there is still felt to be a need for novel pharmaceutical compositions and compounds that are effective in the treatment of diseases based on the expression of MCP-1, CX3CR1 and p40.


[0032] The Applicant has found, surprisingly, novel 1-benzyl-3-hydroxymethylindazole derivatives with pharmacological activity.

[0033] The Applicant has found, surprisingly, that the novel 1-benzyl-3-hydroxymethylindazole derivatives according to formula (I) of the present invention are capable of reducing the production of the chemokine MCP-1.

[0034] More surprisingly, the Applicant has found that the novel 1-benzyl-3-hydroxymethylindazole derivatives according to formula (I) of the present invention are capable of reducing the expression of the chemokine MCP-1.

[0035] Even more surprisingly, the Applicant has found that the 1-benzyl-3-hydroxymethylindazole derivatives according to formula (I) of the present invention are capable of reducing the expression of the subunit p40 involved in the production of the cytokines IL-12 and IL-23, and the expression of the receptor CX3CR1.

[0036] Thus, in a first aspect, the present invention consists of a compound of formula (I)



in which:

A may be $-X_1-$ or $-X_1-OC(R_9)(R_{10})-$, in which

X_1 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms, and R_8 and R_{10} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

Y may be $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$, in which

R_{11} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, or R_{11} together with R_{12} forms a 4- to 7-membered heterocycle,

R_{12} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms, or R_{12} together with R_{11} , forms a 4- to 7-membered heterocycle,

R_{13} and R_{15} , which may be identical or different, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R_{14} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

X_2 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms,

X_3 may be OH, NH_2 , $NHOH$ or $NHNH_2$,

R_1 and R_2 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R_3 , R_4 and R_8 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro and trifluoromethyl; with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

R_5 may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, nitro and trifluoromethyl, or R_5 together with one from between R_6 and R_7 forms a ring having 5 or 6 carbon atoms; with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

R_6 and R_7 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, or together form a group $C=O$, or one from between R_6 and R_7 , together with R_5 , forms a ring having 5 or 6 carbon atoms.


[0037] In a second aspect, the present invention relates to a pharmaceutical composition comprising the novel 1-benzyl-3-hydroxymethylindazole derivatives according to formula (I) or a pharmaceutically acceptable salt thereof together with at least one pharmaceutically acceptable vehicle.

[0038] The over-regulation and/or the increase of the expression of the above mentioned MCP-1, CX3CR1, and p40,

the latest resulting consequently in IL-12 and/or IL-23 expression/production, which results in a development of a pathology and/or a disease is often referred in the art with the term "overexpression". For the purpose of the present invention, the term expression is intended to include overexpression as known in the art.

[0039] Surprisingly, the Applicant has found that the novel 1-benzyl-3-hydroxymethylindazole derivatives may be used for the preparation of a pharmaceutical composition that is active in the case of diseases based on the expression of the chemokine MCP-1, of the subunit p40, and consequently of the cytokines IL-12 and IL-23, and of the receptor CX3CR1.

[0040] Thus, in a further aspect, the present invention relates to the use of a compound of formula (I)



in which:

A may be $-X_1-$ or $-X_1-OC(R_9)(R_{10})-$, in which

X_1 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms, and

R_9 and R_{10} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

Y may be $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$, in which

R_{11} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, or R_{11} together with R_{12} forms a 4- to 7-membered heterocycle,

R_{12} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms, or R_{12} together with R_{11} , forms a 4- to 7-membered heterocycle,

R_{13} and R_{15} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R_{14} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

X_2 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms, X_3 may be OH, NH_2 , $NHOH$ or $NHNH_2$,

R_1 and R_2 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R_3 , R_4 and R_8 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro and trifluoromethyl; with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

R_5 may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, nitro and trifluoromethyl, or R_5 together with one from among R_6 and R_7 forms a ring having 5 or 6 carbon atoms; with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms, and

R₆ and R₇, which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, or together form a group C=O, or one from among R₆ and R₇, together with R₅, forms a ring having 5 or 6 carbon atoms,

for preparing a pharmaceutical composition for the treatment of diseases based on the expression of MCP-1, CX3CR1 and p40.

[0041] In addition, in a further aspect, the present invention relates to a method for treating or preventing diseases based on the expression of MCP-1, CX3CR1 and p40, characterized by the administration to a person in need thereof an effective amount of the compound of formula (I) previously described.

DETAILED DESCRIPTION OF THE INVENTION

[0042] Preferably, in formula (I) previously described, residue A is represented by the group X₁ or X₁-OC(R₉)(R₁₀), where X₁ is an alkyl group having from 1 to 3 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 3 carbon atoms or one or more alkoxy groups having 1 or 2 carbon atoms, and R₉ and R₁₀, which may be identical or different each other, are hydrogen, an alkyl group having from 1 to 3 carbon atoms or an alkoxy group having 1 or 2 carbon atoms.

[0043] More preferably, residue A is represented by the group X₁ or X₁-OC(R₉)(R₁₀), where X₁ is the group CH₂, the group CH₂CH₂ or the group C(CH₃)₂, and R₉ and R₁₀, which may be identical or different each other, are hydrogen or a CH₃ group.

[0044] Advantageously, residue A is chosen from the group comprising a group CH₂, a group CH₂CH₂, a group C(CH₃)₂, a group CH₂CH₂OCH₂, a group CH₂CH₂OC(CH₃)₂ and a group CH₂CH₂CH₂OC(CH₃)₂.

[0045] Preferably, in formula (I) described previously, residue Y is represented by the group N(R₁₁)(R₁₂), N(R₁₃)N(R₁₄)(R₁₅), N(R₁₃)-X₂-N(R₁₄)(R₁₅) or N(R₁₃)-X₂-CO-X₃.

[0046] More preferably, in formula (I) described previously, residue Y is represented by the group N(R₁₁)(R₁₂), N(R₁₃)N(R₁₄)(R₁₅) or N(R₁₃)-X₂-N(R₁₄)(R₁₅).

[0047] Advantageously, R₁₁ is represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, or R₁₁ together with R₁₂ forms a 5- or 6-membered heterocycle.

[0048] More preferably, R₁₁ is represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, or R₁₁ together with R₁₂ forms a 5- or 6-membered heterocycle.

[0049] Advantageously, R₁₂ is represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 3 carbon atoms, or R₁₂ together with R₁₁ forms a 5- or 6-membered heterocycle.

[0050] More preferably, R₁₂ is represented by a hydrogen atom, an alkyl group having 1 or 2 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 3 carbon atoms, or R₁₂ together with R₁₁ forms a 5- or 6-membered heterocycle.

[0051] Preferably, R₁₃ and R₁₅, which may be identical or different each other, are represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms or an alkoxy group having 1 or 2 carbon atoms.

[0052] More preferably, R₁₃ and R₁₅, which may be identical or different each other, are represented by a hydrogen atom, an alkyl group having 1 or 2 carbon atoms, or an alkoxy group having 1 or 2 carbon atoms.

[0053] Preferably, R₁₄ is represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, COR', COOR', CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 3 carbon atoms.

[0054] More preferably, R₁₄ is represented by a hydrogen atom, an alkyl group having 1 or 2 carbon atoms, or CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 3 carbon atoms.

[0055] Advantageously, X₂ is represented by an alkyl group having from 1 to 3 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 3 carbon atoms or one or more alkoxy groups having 1 or 2 carbon atoms.

[0056] Advantageously, X₃ is represented by the groups OH, NH₂, NHOH or NNNH₂.

[0057] Preferably, R₁ and R₂, which may be identical or different each other, are represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms or an alkoxy group having 1 or 2 carbon atoms.

[0058] Preferably, R₃, R₄ and R₈, which may be identical or different each other, are represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, a Br, Cl or F atom, an OH group, a nitro group, a trifluoromethyl group or a group N(R')(R'') or N(R')COR''; -CN, -CONR'R'', -SO₂NR'R'', -SO₂R', with R' and R'', which may be identical or different each other, represented by a hydrogen atom and an alkyl group having from 1 to 3 carbon atoms.

[0059] Advantageously, R₅ is represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, a halogen atom, an OH group, or R₅, together with one from among R₆ and R₇, forms a ring having 5 or 6 carbon atoms.

[0060] Preferably, R₆ and R₇, which may be identical or different each other, are represented by a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, or together form a group C=O, or one from among R₆ and R₇, together with R₅, forms a ring having 5 or 6 carbon atoms.

[0061] In the case of certain substituents, the compound of formula (I) according to the present invention may be an asymmetric carbon atom and may then be in the form of stereoisomers and enantiomers.

[0062] Depending on the nature of the substituents, the compound of formula (I) may form addition salts with physiologically acceptable organic or mineral acids or bases.

[0063] Typical examples of suitable physiologically acceptable mineral acids are hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid and nitric acid.

[0064] Typical examples of suitable physiologically acceptable organic acids are acetic acid, ascorbic acid, benzoic acid, citric acid, fumaric acid, lactic acid, maleic acid, methanesulfonic acid, oxalic acid, para-toluenesulfonic acid, benzenesulfonic acid, succinic acid, tannic acid and tartaric acid.

[0065] Typical examples of suitable physiologically acceptable mineral bases are hydroxides, carbonates and hydrogen carbonates of ammonium, calcium, magnesium, sodium and potassium, for instance ammonium hydroxide, calcium hydroxide, magnesium carbonate, sodium hydrogen carbonate and potassium hydrogen carbonate.

[0066] Typical examples of suitable physiologically acceptable organic bases are: arginine, betaine, caffeine, choline, N,N-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, N-methylglucamine, glucamine, glucosamine, histidine, N-(2-hydroxyethyl)piperidine, N-(2-hydroxyethyl)pyrrolidine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, theobromine, triethylamine, trimethylamine, tripropylamine and tromethamine.

[0067] Depending on the nature of the substituents, the compound of formula (I) may form esters with physiologically acceptable organic acids or bases.

[0068] The compounds of the present invention also include the prodrugs, stereoisomers, enantiomers and pharmaceutically acceptable salts or esters of the compounds represented by formula (I) described in the claims. The prodrug of a compound of formula (I) is a substance in substantially inactive form, which, when administered to a living being, is metabolized into a compound of formula (I).

[0069] The terms "pharmaceutically acceptable" and "physiologically acceptable" are intended to define, without any particular limitation, any material suitable for preparing a pharmaceutical composition to be administered to a living being.

[0070] The compounds according to formula (I) of the present invention may be used for the preparation of a pharmaceutical composition that is active in the treatment of diseases (or pathologies) based on the expression of the chemokine MCP-1, the cytokine p40, the subunit p40 (involved in the production of cytokines IL-12 and IL-23) and the receptor CX3CR1.

[0071] Preferably, the pathologies associated with the expression of MCP-1 and CX3CR1 are articular diseases, renal diseases, cardiovascular diseases, metabolic syndrome, obesity, diabetes, insulin resistance and cancer.

[0072] In particular, the pathologies associated with the expression of MCP-1 are rheumatoid arthritis, arthritis induced by viral infections, psoriatic arthritis, arthrosis, lupus nephritis, diabetic nephropathy, glomerulonephritis, polycystic kidney disease, interstitial lung disease, fibrosis, multiple sclerosis, Alzheimer's disease, HIV-associated dementia, atopic dermatitis, psoriasis, vasculitis, restenosis, atherosclerosis, myocardial infarction, angina, acute coronary diseases, adenomas, carcinomas and metastases, metabolic diseases, transplants, organ replacements, tissue replacements and prosthesis implants.

[0073] In particular, the pathologies associated with the expression of CX3CR1 are rheumatoid arthritis, lupus nephritis, diabetic nephropathy, Crohn's disease, ulcerative colitis, coronary disorders, restenosis, atherosclerosis, myocardial infarction, angina.

[0074] Preferably, the pathologies associated with the expression of p40, and thus of IL-12 and IL-23, are autoimmune diseases, such as chronic degenerative inflammatory diseases, and cancer.

[0075] In particular, the pathologies associated with the expression of p40 are rheumatoid arthritis, psoriasis, glomerulonephritis, diabetes, lupus erythematosus, diabetes, Crohn's disease, and tumours such as, for example, colon carcinomas, breast carcinomas, lung carcinomas and prostate carcinomas, and skin and CNS neoplasias.

[0076] Preferably, the pharmaceutical compositions of the present invention are prepared in suitable dosage forms comprising an effective dose of at least one compound of formula (I), or a pharmaceutically acceptable salt, or ester thereof, and at least one pharmaceutically acceptable vehicle.

[0077] Examples of pharmaceutically acceptable vehicles known in the prior art are, for example, glidants, binders, disintegrants, fillers, diluents, flavourings, colorants, fluidizers, lubricants, preserving agents, humectants, absorbents and sweeteners.

[0078] Useful examples of pharmaceutically acceptable excipients are sugars, such as lactose, glucose or sucrose,

starches, such as corn starch and potato starch, cellulose and derivatives thereof, for instance sodium carboxymethyl-cellulose, ethylcellulose and cellulose acetate, gum tragacanth, malt, gelatin, talc, cocoa butter, waxes, oils, such as groundnut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil, glycols such as propylene glycol, polyols such as glycerol, sorbitol, mannitol and polyethylene glycol, esters such as ethyl oleate and ethyl laurate, agar-agar, and the like.

[0079] Examples of suitable dosage forms are tablets, capsules, coated tablets, granules, solutions and syrups for oral administration; medicated plasters, solutions, pastes, creams and ointments for transdermal administration; suppositories for rectal administration and sterile solutions for injection or aerosol administration.

[0080] Other suitable dosage forms are sustained-release forms and liposome-based forms, for either the oral or injection route.

[0081] The dosage forms may also contain other conventional ingredients such as: preserving agents, stabilizers, surfactants, buffers, osmotic pressure regulators, emulsifiers, sweeteners, colorants, flavourings and the like.

[0082] When required for particular therapies, the pharmaceutical composition of the present invention may contain other pharmacologically active ingredients whose simultaneous administration is useful.

[0083] The amount of compound of formula (I) or of pharmaceutically acceptable salt, or ester thereof in the pharmaceutical composition of the present invention may vary within a wide range as a function of known factors, for instance the type of pathology to be treated, the severity of the disease, the body weight of the patient, the dosage form, the chosen route of administration, the number of daily administrations and the efficacy of the chosen compound of formula (I). However, the optimum amount may be determined simply and routinely by a person skilled in the art.

[0084] Typically, the amount of compound of formula (I) or of pharmaceutically acceptable salt, or ester thereof in the pharmaceutical composition of the present invention will be such that it ensures a level of administration of between 0.001 and 100 mg/kg/day. Preferably, the level of administration is between 0.05 and 50 mg/kg/day and even more preferably between 0.1 and 10 mg/kg/day.

[0085] The dosage forms of the pharmaceutical composition of the present invention may be prepared according to techniques that are well known to pharmaceutical chemists, including mixing, granulation, compression, dissolution, sterilization and the like.


[0086] The activity of the compounds of the present invention on MCP-1 and CX3CR1 was demonstrated *in vitro* in human monocytes via techniques of gene expression analysis with "real-time" RT-PCR and by protein production analysis via an immunoenzymatic test. As is known to those skilled in the art, the abovementioned experimental models are considered useful for checking the activity of the compounds with regard to the expression and production of MCP-1 and the expression of CX3CR1. Consequently, the abovementioned models may be considered as predictive of the activity in man for the treatment of pathologies characterized by the expression and production of MCP-1, by the expression of CX3CR1 and by inflammatory conditions with the presence of infiltrates rich in monocytes and macrophages.

[0087] The activity of the compounds of the present invention on p40 was demonstrated *in vitro* in human monocytes via gene expression analysis techniques via "real-time" RT-PCR. As is known to those skilled in the art, the abovementioned experimental model is useful for checking the activity of compounds with regard to the expression of p40 and may be considered as predictive of the activity in man for the treatment of pathologies characterized by the expression of p40.

[0088] The preparation of the compounds of general formula (I) may be performed according to one of the following procedures.

Method A:

[0089]




[0090] In method A, the compounds of general formula (III) are reacted with the compounds of formula (IV). The substituents R_1 to R_8 , A and Y have the meanings given previously for the compounds of formula (I) and Q indicates a leaving group chosen from the group comprising halogen, CH_3SO_3^- and $p\text{-CH}_3\text{PhSO}_3^-$.

[0091] Method A is performed according to conventional techniques. For example, the alcohols of formula (III) are reacted, respectively, with the derivatives of formula (IV) in which Q is a leaving group preferably chosen from the group comprising a chlorine atom, a bromine atom and a methanesulfonyl group. The reaction is performed in the presence of a suitable base and in a suitable solvent. The bases that may preferably be used are NaH, butyllithium and lithium diisopropylamide, whereas the solvents that are suitable for this type of reaction are preferably polar aprotic solvents such as tetrahydrofuran, diethyl ether or 1,4-dioxane. The reaction temperature is preferably between room temperature and the reflux temperature of the solvent used. Reactions of this type may last from a few hours to a few days.

Method B:

[0092]




[0093] In method B, the compounds of general formula (V) are reacted with the compounds of formula (VI). The substituents R_1 to R_8 , A and Y have the meanings given previously for the compounds of formula (I) and Q indicates a leaving group chosen from the group comprising halogen, CH_3SO_3^- and $p\text{-CH}_3\text{PhSO}_3^-$.

[0094] Method B is performed according to conventional techniques. For example, the alcohols of formula (VI) are reacted, respectively, with the derivatives of formula (V) in which Q is a leaving group preferably chosen from the group comprising a chlorine atom, a bromine atom and a methanesulfonyl group. The reaction is performed in the presence of a suitable base and in a suitable solvent. The bases that may preferably be used are NaH, butyllithium and lithium diisopropylamide, whereas the solvents that are suitable for this type of reaction are preferably polar aprotic solvents such as tetrahydrofuran, diethyl ether or 1,4-dioxane. The reaction temperature is preferably between room temperature and the reflux temperature of the solvent used. Reactions of this type may last from a few hours to a few days.

Method C:

[0095]



[0096] In method C, the products of general formula (IX) are reacted with the products of general formula (X). The substituents R_1 to R_8 , A and Y have the meanings given previously for the compounds of formula (I) and T may be

hydrogen or an alkyl group.

[0097] Method C is performed using conventional techniques.

[0098] When T is hydrogen, the method involves the use of a suitable agent for activating the carboxylic acid, such as thionyl chloride, and an aprotic solvent. Preferably, the solvents used are toluene, tetrahydrofuran and dichloromethane. The reaction may be performed in the presence of a base, which may be either organic, such as triethylamine or diisopropylethylamine, or mineral, such as sodium methoxide. Preferably, the reaction is performed at temperatures of between 0°C and the reflux temperature of the solvent used. The reaction time generally ranges from 1 to 24 hours.

[0099] When T is an alkyl group, the method involves the use of a suitable mineral base, such as sodium methoxide or NaOH and an excess of amine of general formula (X). The reaction may generally be performed using a suitable aprotic solvent, preferably toluene or dioxane or the same reactive amine (X) if its physical state makes this possible. The reaction is preferably performed at temperatures above room temperature, for times that range from a few hours to several days.

[0100] The examples that follow are intended to illustrate the present invention without, however, limiting it in any way.

Preparative examples

[0101] The compounds of formula (I) listed in Table A below were prepared using the preparation methods described previously.

TABLE A

No.	A	Y	Groups R							
			1	2	3	4	5	6	7	8
9	C(CH ₃) ₂	NH ₂	H	H	H	H	H	H	H	H
10	"	NHOH	"	"	"	"	"	"	"	"
11	"	NHNH ₂	"	"	"	"	"	"	"	"
12	"	NHNHCONH ₂	"	"	"	"	"	"	"	"
13	"	NH(CH ₂) ₂ N(CH ₃) ₂	"	"	"	"	"	"	"	"
14*	"	NH-(2-pyridine)	"	"	"	"	"	"	"	"
15*	"	NH-[1-(4-CH ₃)-piperazine]	"	"	"	"	"	"	"	"
26	"	NHCH ₂ COOH	"	"	"	"	"	"	"	"
27*	"	NH-NH-Ph	"	"	"	"	"	"	"	"
28	"	N-morpholine	"	"	"	"	"	"	"	"
29	"	NH ₂	"	"	"	"	"	"	"	5-OCH ₃
30	"	"	"	"	p-Cl	"	"	"	"	H
31	"	"	"	"	p-Cl	m-Cl	"	"	"	"
32	"	NHOH	"	"	H	H	"	"	"	5-CN
33	"	"	"	"	p-OCH ₃	"	"	"	"	5-CN
34	"	"	"	"	H	"	"	CO		H
35	"	"	"	"	"	"	(CH ₂) ₃		H	H
36	CH ₂	"	"	"	p-OCH ₃	"	H	H	"	"
37	"	NH ₂	"	"	H	"	"	"	"	5-CONH ₂
38	C(CH ₃) ₂ CH ₂ OC(CH ₃) ₂	"	"	"	"	"	"	"	"	H
39	C(CH ₃) ₂ CH ₂ OCH ₂	"	"	"	"	"	"	"	"	"
40	"	NHOH	"	"	"	"	"	"	"	"
41	"	"	"	"	p-OCH ₃	"	"	"	"	"
42	CH ₂ CH ₂ OC(CH ₃) ₃	"	"	"	H	"	"	"	"	"

(continued)

No.	A	Y	Groups R							
			1	2	3	4	5	6	7	8
43	CH ₂ CH ₂ CH ₂ OC(CH ₃) ₂	"	"	"	"	"	"	"	"	"
44	CH ₂ CH ₂ CH ₂ OCH ₂	"	"	"	"	"	"	"	"	"
45	"	NH ₂	"	"	"	"	"	"	"	"
46	CH ₂ CH ₂ OCH ₂	"	"	"	"	"	"	"	"	"
47	"	NHOH	"	"	"	"	"	"	"	"
48	"	NHNH ₂	"	"	"	"	"	"	"	"
* reference compound not part of the invention										

[0102] The details of the preparation of most compounds of Table A are given hereinbelow. The remaining compounds were prepared with similar techniques using suitable starting products and reagents.

Preparation of compound 9

2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanamide

9a) methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoate

[0103] A suspension of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (20 g; 0.062 mol) [prepared as described in patent application EP 382 276] in methanol (300 ml) was treated at 0°C with gaseous HCl for 4 hours. The mixture was then poured into water (500 ml) and the product was extracted with diethyl ether (3×250 ml). The combined organic phases were washed with 5% sodium bicarbonate solution (2×100 ml) and then with water (50 ml). The solvent was evaporated off under reduced pressure and the crude residue was purified by crystallization from hexane.

[0104] 18.5 g of methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoate were obtained.

m.p. = 66°-67°C

¹H-NMR (CDCl₃, δ ppm): 1.56 (s, 6 H), 3.77 (s, 3 H), 4.87 (s, 2 H), 5.54 (s, 2 H), 7.0-7.4 (m, 8 H), 8.03 (d, J = 7.61 Hz, 1 H).

9b) 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanamide

[0105] A solution of methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (15 g; 0.044 mol) in methanol (250 ml) stirred at room temperature was treated with gaseous NH₃ for 48 hours. The solvent was then evaporated off under reduced pressure and the crude residue was purified by crystallization from ethyl acetate.

[0106] 7.3 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanamide were obtained.

m.p. = 111 °-112°C

¹H-NMR (CDCl₃, δ ppm): 1.58 (s, 6 H), 4.88 (s, 2 H), 5.57 (s, 2 H), 5.64 (bs, 1 H), 6.94 (bs, 1 H), 7.08-7.43 (m, 8 H), 7.77 (d, J = 8.04 Hz, 1 H).

Preparation of compound 10

2-[(1-benzyl-1H-indazol-3-yl)methoxy]-N-hydroxy-2-methylpropanamide

10a) methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoate

[0107] A suspension of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (20 g; 0.062 mol) in methanol (300 ml) was treated at 0°C with gaseous HCl for 4 hours. The mixture was then poured into water (500 ml) and the product was extracted with diethyl ether (3×250 ml). The combined organic phases were washed with 5% sodium bicarbonate solution (2×100 ml) and then with water (50 ml). The solvent was evaporated off under reduced pressure and the crude residue was purified by crystallization from hexane.

[0108] 18.5 g of methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid were thus obtained.

m.p. = 66°-67°C

¹H-NMR (CDCl₃, δ ppm): 1.56 (s, 6 H), 3.77 (s, 3 H), 4.87 (s, 2 H), 5.54 (s, 2 H), 7.0-7.4 (m, 8 H), 8.03 (d, J = 7.61 Hz, 1 H).

10b) 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-N-hydroxy-2-methylpropanamide

[0109] A solution of KOH (9.7 g, 0.173 mol) in methanol (25 ml) was added at room temperature to a solution of hydroxylamine hydrochloride (8.0 g, 0.115 mol) in methanol (45 ml). The resulting mixture was stirred at 0°C for 30 minutes and then filtered, and the solution was added to a solution of methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (19 g, 0.057 mol) in methanol (30 ml) and CHCl₃ (15 ml). The mixture was stirred at room temperature for 72 hours. The reaction was stopped by concentrating the solvent under reduced pressure and washing the residue taken up in water (100 ml) with diethyl ether (3×50 ml). The aqueous phase was acidified with 2N HCl and the product was extracted with diethyl ether (3×100 ml). The organic phase was then concentrated under reduced pressure and the resulting crude residue was purified by double crystallization from a 1/1 hexane/ethyl acetate mixture.

[0110] 6 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-N-hydroxy-2-methylpropanamide were obtained.

m.p. = 115°-116°C

¹H-NMR (CDCl₃, δ ppm): 1.58 (s, 6 H), 4.87 (s, 2 H), 5.57 (s, 2 H), 7.12-7.41 (m, 8 H), 8.10 (bs, 1 H), 7.72 (d, J = 8.09 Hz, 1 H), 10.05 (bs, 1 H).

Preparation of compound 11

2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanohydrazide

[0111] To a solution of 1 M hydrazine hydrate (100 ml, 0.100 mol) stirred at 80°C was added portionwise methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoate (20 g, 0.060 mol). Once the addition was complete, the mixture was stirred for 30 minutes at the same temperature and then heated at 120°C for 2 hours. The reaction was stopped by diluting the mixture with water (600 ml) and extracting the product with diethyl ether (4×200 ml). The combined organic phases were then extracted with 2N HCl (4×200 ml). The acidic phase was then brought to basic pH with 10N NaOH and again extracted with diethyl ether (4×200 ml). The combined organic phases were concentrated under reduced pressure and the crude residue was purified by crystallization from a 1/1 hexane/ethyl acetate mixture.

[0112] 16 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanohydrazide were obtained.

m.p. = 92°-93°C

¹H-NMR (DMSO-d₆, δ ppm): 1.41 (s, 6 H), 4.30 (s, 2 H), 4.73 (s, 2 H), 5.62 (s, 2 H), 7.15 (ddd, J = 8.00; 6.98; 0.80 Hz, 1 H), 7.20-7.34 (m, 5 H), 7.38 (ddd, J = 8.37; 6.98; 1.17 Hz, 1 H), 7.66 (d, J = 8.48 Hz, 1 H), 7.90 (dt, J = 8.11; 0.91 Hz, 1 H), 8.84 (s, 1 H).

Preparation of compound 12

2-{2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoyl}hydrazine-carboxamide

[0113] To a solution of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanohydrazide (7.0 g, 0.021 mol) in water (100 ml) and 3N HCl (15 ml) stirred at room temperature was slowly added a solution of KOCN (1.8 g, 0.022 mol) in water (30 ml). The mixture was stirred at this same temperature for 30 minutes and the reaction was then stopped by filtering off the solid thus formed. The isolated solid was crystallized from 95° ethanol.

[0114] 6.0 g of 2-{2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoyl}-hydrazinecarboxamide were obtained.

m.p. = 174°-175°C

¹H-NMR (DMSO-d₆, δ ppm): 1.43 (s, 6 H), 4.80 (s, 2 H), 5.62 (s, 2 H), 5.90 (s, 2 H), 7.14 (t, J = 7.45 Hz, 1 H), 7.20-7.43 (m, 6 H), 7.66 (d, J = 8.48 Hz, 1 H), 7.77 (s, 1 H), 7.96 (d, J = 8.04 Hz, 1 H), 9.37 (s, 1 H).

Preparation of compound 13

2-[(1-benzyl-1H-indazol-3-yl)methoxy]-N-[2-(dimethylamino)ethyl]-2-methylpropanamide hydrochloride

[0115] To a solution of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid, prepared as described in EP 382 276 (19.5 g, 0.060 mol), in toluene (150 ml) stirred at room temperature was added triethylamine (8.4 ml, 0.060 mol). The mixture was then cooled to 0°C and a solution of ethyl chloroformate (7.0 ml, 0.073 mol) in toluene (30 ml) was added. Once the addition was complete, the mixture was stirred at this same temperature for 30 minutes, and N,N-dimethylethane-1,2-diamine (7.0 ml, 0.064 mol) dissolved in toluene (30 ml) was then added. Once the addition was complete, the mixture was stirred at room temperature for 24 hours. The mixture was then diluted with diethyl ether (500 ml) and extracted with 1 N HCl (4×50 ml). The combined acidic aqueous phases were washed with diethyl ether (3×50 ml) and then brought to basic pH with 10N NaOH and extracted again with diethyl ether (4×150 ml). The organic phase was then washed with water (2×50 ml) and with saturated NaCl solution (2×50 ml). The solution was dried with anhydrous

Na₂SO₄ and the solvent was then evaporated off under reduced pressure.

[0116] The residue obtained was dissolved in ethyl acetate (150 ml) and treated at room temperature with a solution of HCl in ethanol (about 5N). The solid thus formed was filtered off and purified by crystallization from a 9/1 ethyl acetate/ethanol mixture.

[0117] 12.1 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-N-[2-(dimethylamino)-ethyl]-2-methylpropanamide hydrochloride were obtained.

m.p. = 136°-137°C

¹H-NMR (CDCl₃, δ ppm): 1.54 (s, 6 H), 2.75 (d, J = 4.76 Hz, 6 H), 3.01-3.13 (m, 2 H), 3.69 (q, J = 6.46 Hz, 2 H), 4.87 (s, 2 H), 5.61 (s, 2 H), 7.13-7.41 (m, 8 H), 7.69 (t, J = 5.85 Hz, 1 H), 7.80 (d, J = 8.05 Hz, 1 H), 12.68 (bs, 1 H).

Preparation of compound 14 (reference compound)

2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methyl-N-pyrid-2-ylpropanamide

[0118] 8 g (0.024 mol) of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methyl-propanoic acid were treated at room temperature with 30% sodium methoxide (4.33 ml, 0.024 mol) for 5 minutes. The solvent was then evaporated off under reduced pressure and the residue obtained was suspended in toluene (160 ml), followed by addition thereto, at room temperature, of 2-aminopyridine (6.8 g, 0.072 mol). To the mixture, stirred at room temperature, was slowly added a solution of thionyl chloride (2.1 ml, 0.029 mol) in toluene (40 ml). Once the addition was complete, the mixture was stirred for 24 hours. The solid produced was then filtered off and the solvent was evaporated off under reduced pressure. The crude residue obtained was purified by crystallization from isopropanol.

[0119] 3 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methyl-N-pyrid-2-ylpropanamide were obtained.

m.p. = 121°-122°C

¹H-NMR (CDCl₃, δ ppm): 1.65 (s, 6 H), 4.97 (s, 2 H), 5.60 (s, 2 H), 6.98-7.04 (m, 1 H), 7.15 (ddd, J = 8.04; 5.77; 2.12 Hz, 1 H), 7.20-7.37 (m, 7 H), 7.64-7.73 (m, 1 H), 7.84 (d, J = 8.18 Hz, 1 H), 8.22-8.27 (m, 2 H), 9.32 (bs, 1 H).

Preparation of compound 15 (reference compound)

1-benzyl-3-[[1,1-dimethyl-2-(4-methylpiperazin-1-yl)-2-oxoethoxy]methyl]-1H-indazole

[0120] 50 ml of 1-methylpiperazine (0.452 mol) were treated at room temperature with a methanolic solution of 30% sodium methoxide (9 ml, 0.045 mol) in methanol (5 ml). To the solution, stirred at room temperature, was added methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (15 g, 0.045 mol). The mixture was stirred at reflux for 18 hours. The reaction was stopped by pouring the mixture into water (500 ml) and extracting the product with diethyl ether (3×150 ml). The combined organic phases were extracted with 3N HCl (3×100 ml). The acidic phase was then brought to basic pH with 10N NaOH and extracted with diethyl ether (3×150 ml). The combined organic phases were concentrated under reduced pressure. The residue obtained was purified by crystallization from a 1/3 hexane/ethyl acetate mixture.

[0121] 6 g of 1-benzyl-3-[[1,1-dimethyl-2-(4-methylpiperazin-1-yl)-2-oxoethoxy]methyl]-1H-indazole were thus obtained.

m.p. = 97°-98°C

¹H-NMR (CDCl₃, δ ppm): 1.60 (s, 6 H), 2.21 (s, 3 H), 2.24-2.47 (m, 4 H), 3.47-4.16 (m, 4 H), 4.82 (s, 2 H), 5.56 (s, 2 H), 7.09-7.38 (m, 8 H), 7.76 (d, J = 8.18 Hz, 1 H).

Preparation of compound 26

[0122] N-{2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoyl}glycine 4.17 g of glycine ethyl ester (0.040 mol) were treated at room temperature with a solution of 30% sodium methoxide in methanol (0.46 ml) and the solvent was then evaporated off under reduced pressure. The residue obtained was added to a solution of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (10 g, 0.04 mol) in dichloromethane (DCM) (100 ml) while stirring at room temperature. The mixture was then cooled to 0°C, and a solution of N,N'-dicyclohexylcarbodiimide (6.73 g, 0.033 mol) in DCM (25 ml) was added thereto. The mixture was stirred at room temperature for 24 hours.

[0123] The reaction was stopped by filtering off the solid thus formed and concentrating the solvent under reduced pressure. The crude residue was purified via flash chromatography on silica gel, using a 1/1 hexane/ethyl acetate mixture as eluent.

[0124] The product obtained was suspended in water (100 ml) and NaOH (1.28 g, 0.032 mol) was added thereto. The mixture was stirred at 50°C for 16 hours, and then cooled and acidified with concentrated HCl. The solid thus obtained was purified by crystallization from ethanol.

[0125] 5.2 g of N-{2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoyl}-glycine were obtained.

m.p. = 157°-158°C

¹H-NMR (CDCl₃, δ ppm): 1.55 (s, 6 H), 3.95 (d, J = 5.74 Hz, 2 H), 4.89 (s, 2 H), 5.52 (s, 2 H), 7.0-7.4 (m, 8 H), 7.6 (bt, 1 H), 7.77 (d, J = 7.82 Hz, 1 H), 10.38 (bs, 1 H).

Preparation of compound 27 (reference compound)

2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methyl-N'-phenylpropanohydrazide

[0126] 24 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid (0.074 mol) were treated at room temperature with a solution of 30% sodium methoxide in methanol (13 ml, 0.074 mol) for 10 minutes, the solvent was then evaporated off under reduced pressure and the residue was suspended in toluene (240 ml), and phenylhydrazine (29.1 ml, 0.296 mol) was added thereto. To the mixture, kept stirring at this same temperature, was slowly added a solution of thionyl chloride (6.3 ml, 0.088 mol) in toluene (50 ml). Once the addition was complete, the mixture was stirred for 24 hours.

[0127] The solid formed was then filtered off and the solvent was evaporated off under reduced pressure. The crude residue was washed with hexane (3×100 ml) and with a 10/1 hexane/ethyl acetate mixture (30 ml).

[0128] The crude residue obtained was purified by successive crystallizations from isopropanol.

[0129] 11 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methyl-N'-phenylpropanohydrazide were obtained.

m.p. = 124°-125°C

¹H-NMR (CDCl₃, δ ppm): 1.61 (s, 2 H), 4.94 (s, 2 H), 5.53 (s, 2 H), 6.10 (bd, J = 4.70 Hz, 1 H), 6.7-7.0 (m, 3 H), 7.1-7.4 (m, 13 H), 7.78 (dt, J₁, J₂ = 7.78; 1.19 Hz, 1 H), 8.99 (bd, J = 4.70 Hz, 1 H).

Preparation of compound 28

1-benzyl-3-[(1,1-dimethyl-2-morpholin-4-yl-2-oxoethoxy)methyl]-1H-indazole

[0130] 72 g (0.222 mol) of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid were treated at room temperature with a methanolic solution of 30% sodium methoxide (39 ml; 0.222 mol) for 10 minutes, the solvent was then evaporated off under reduced pressure and the residue obtained was suspended in anhydrous toluene (750 ml). To the suspension, stirred at room temperature, was added morpholine (77.6 ml; 0.888 mol) followed by slow addition of a solution of thionyl chloride (19.3 ml; 0.266 mol) in toluene (150 ml). The mixture was stirred for 24 hours and the reaction was then stopped by filtering off the solid thus formed. The solution was concentrated under reduced pressure and the crude residue obtained was purified by crystallization from isopropanol.

[0131] 14 g of 1-benzyl-3-[(1,1-dimethyl-2-morpholin-4-yl-2-oxoethoxy)methyl]-1H-indazole were thus obtained.

m.p. = 135°-137°C

¹H-NMR (DMSO-d₆, δ ppm): 1.47 (s, 6 H), 3.1-4.0 (2bs, 8 H), 4.73 (s, 2 H), 5.83 (s, 2 H), 7.0-7.9 (m, 9 H).

Preparation of compound 29

2-[(1-benzyl-5-methoxy-1H-indazol-3-yl)methoxy]-2-methylpropanamide

29a) benzyl 1-benzyl-5-methoxy-1H-indazole-3-carboxylate

[0132] A suspension of 5-methoxy-1H-indazole-3-carboxylic acid (21.5 g; 0.11 mol) and 60% NaH (10.5 g; 0.44 mol) in N,N-dimethylformamide (DMF) (200 ml) was stirred at 70°C for 1 hour. Benzyl chloride (32.9 g; 0.26 mol) was then added slowly to the suspension and the mixture was stirred at 70°C for 4 hours. The reaction was stopped by cooling the mixture to room temperature and pouring the mixture into water and ice. The product was extracted with ethyl acetate (3×250 ml). The combined organic phases were concentrated under reduced pressure. The crude residue thus obtained was purified by successive crystallizations from 95° ethanol, to give 18 g of benzyl 1-benzyl-5-methoxy-1H-indazole-3-carboxylate with a melting point of 107-109°C.

¹H-NMR (CDCl₃, δ ppm): 3.78 (s, 3 H), 5.51 (s, 2 H), 6.9-7.6 (m, 13 H).

29b) (1-benzyl-5-methoxy-1H-indazol-3-yl)methanol

[0133] To a solution of benzyl 1-benzyl-5-methoxy-1H-indazole-3-carboxylate (17.7 g; 0.05 mol), diethyl ether (100 ml) and tetrahydrofuran (THF) (170 ml) stirred at room temperature was slowly added LiAlH₄ (3.8 g; 0.1 mol). Once the addition was complete, the suspension was stirred at reflux for 24 hours. The reaction was stopped by destroying the

excess LiAlH₄ via addition of water (40 ml) and 5N NaOH (10 ml). The organic phase was separated out and the solvent was evaporated off under reduced pressure. The crude residue obtained was purified by crystallization from 95° ethanol to give 14 g of (1-benzyl-5-methoxy-1H-indazol-3-yl)methanol with a melting point of 97-98°C.

¹H-NMR (CDCl₃, δ ppm): 3.3 (bs, 1 H), 3.80 (s, 3 H), 4.92 (s, 2 H), 5.47 (s, 2 H), 6.9-7.5 (m, 8 H).

29c) 1-benzyl-3-(chloromethyl)-5-methoxy-1H-indazole

[0134] To a solution of (1-benzyl-5-methoxy-1H-indazol-3-yl)methanol (18 g; 0.07 mol) in chloroform (200 ml) stirred at room temperature was slowly added thionyl chloride (15.8 g; 0.13 mol). Once the addition was complete, the solution was refluxed for 24 hours. The reaction was then stopped by cooling the mixture to room temperature and evaporating off the solvent under reduced pressure. The residue was then taken up several times in toluene and concentrated under reduced pressure. The crude residue obtained was purified by crystallization from hexane to give 9.5 g of 1-benzyl-3-(chloromethyl)-5-methoxy-1H-indazole with a melting point of 78-80°C.

¹H-NMR (CDCl₃, δ ppm): 3.85 (s, 3 H), 4.97 (s, 2 H), 5.51 (s, 2 H), 6.9-7.4 (m, 8 H).

29d) 2-[(1-benzyl-5-methoxy-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid

[0135] To a solution containing 1-benzyl-3-(chloromethyl)-5-methoxy-1H-indazole (2.95 g; 0.01 mol) and ethyl 3-hydroxy-3-methylbutanoate (1.98 g; 0.015 mol) in DMF (30 ml) stirred at room temperature was slowly added 60% NaH (0.36 g; 0.015 mol). The mixture was then heated at 40°C for 24 hours. The reaction was stopped by cooling the suspension to room temperature and adding water (200 ml). The solvent was evaporated off under reduced pressure and the residue was treated at reflux with NaOH (0.84 g; 0.021 mol) in water (6 ml) and 95° ethanol (6 ml) for 6 hours. The mixture was then cooled to room temperature and diluted with water (50 ml). The alkaline phase was washed with diethyl ether (2×20 ml) and then acidified with concentrated HCl and extracted with diethyl ether (3×30 ml).

[0136] The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by crystallization from 10/1 hexane/ethyl acetate mixture to give 0.8 g of 2-[(1-benzyl-5-methoxy-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid with a melting point of 128-130°C.

¹H-NMR (DMSO-d₆, δ ppm): 1.44 (s, 6 H), 3.77 (s, 3 H), 4.69 (s, 2 H), 5.55 (s, 2 H), 7.02 (dd, J = 9.15; 2.38 Hz, 1 H), 7.17-7.33 (m, 5 H), 7.41 (d, J = 2.38 Hz, 1 H), 7.55 (d, J = 9.15 Hz, 1 H), 12.79 (s, 1 H).

29e) 2-[(1-benzyl-5-methoxy-1H-indazol-3-yl)methoxy]-2-methylpropanamide

[0137] The product was prepared via the procedures described in the preparation of compound 9, using 2-[(1-benzyl-5-methoxy-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid as starting reagent.

Preparation of compound 30

2-[[1-(4-chlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanamide

30a) [1-(4-chlorobenzyl)-1H-indazol-3-yl]methanol

[0138] To a suspension of 60% NaH (2.7 g; 0.07 mol) in toluene (200 ml) was added 1-benzyl-3-hydroxymethylindazole (10 g; 0.07 mol). The mixture was brought to the boiling point and left stirring at reflux for 1 hour. 4-chlorobenzyl chloride (14.4 g; 0.09 mol) was then added. The mixture was then stirred at reflux for 4 hours. The reaction was stopped by cooling the mixture to room temperature and adding water (50 ml). The organic phase was separated out and washed, respectively, with 2N HCl (50 ml) and water (5×50 ml). The solvent was evaporated off under reduced pressure. The crude residue thus obtained was purified by flash chromatography on silica gel, using as eluent a 3/1 hexane/ethyl acetate mixture. The product obtained was crystallized from a 5/1 hexane/ethyl acetate mixture to give 4.4 g of [1-(4-chlorobenzyl)-1H-indazol-3-yl]methanol.

m.p. = 102°-104°C

¹H-NMR (CDCl₃, δ ppm): 3.5 (bs, 1 H), 5.01 (s, 2 H), 5.37 (s, 2 H), 6.8-7.5 (m, 7 H), 7.81 (d, J = 7.82 Hz, 1 H).

30b) 2-[[1-(4-chlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid

[0139] To a suspension of NaOH (15.6 g; 0.39 mol) in acetone (50 ml) was added [1-(4-chlorobenzyl)-1H-indazol-3-yl]methanol (9.1 g; 0.03 mol). To the mixture was slowly added a solution of chloroform (7.2 ml; 0.09 mol) and acetone (7.2 ml; 0.1 mol). The addition caused refluxing of the mixture of solvents. Once the addition was complete, the mixture was refluxed for 1 hour. The reaction was stopped by cooling the mixture to room temperature and evaporating off the

solvent under reduced pressure. The resulting crude residue was taken up in toluene (100 ml) and water (50 ml). The aqueous phase was separated from the organic phase and then washed with toluene (2×50 ml). The combined organic phases were extracted with water (3×50 ml). The combined aqueous phases were washed with hexane (2×30 ml) and then acidified with 2N HCl and stirred at room temperature. The solid thus obtained was filtered off and crystallized first from a 5/1 water/acetic acid mixture, and then from toluene, to give 4.0 g of 2-[[1-(4-chlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid.

m.p. = 186°-188°C.

¹H-NMR (CDCl₃, δ ppm): 1.61 (s, 6 H), 4.91 (s, 2 H), 5.54 (s, 2 H), 7.0-7.5 (m, 7 H), 8.07 (s, 1 H), 10.3 (bs, 1 H).

30c) 2-[[1-(4-chlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanamide

[0140] The product was prepared via the procedures described in the preparation of compound 9, using 2-[[1-(4-chlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid as starting reagent.

Preparation of compound 31

2-[[1-(3,4-dichlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanamide

31a) [1-(3,4-dichlorobenzyl)-1H-indazol-3-yl]methanol

[0141] The product was obtained via the method described in Example 30a), using 3,4-chlorobenzyl chloride as reagent. The product obtained was purified by crystallization from a 1/1 hexane/ethyl acetate mixture.

m.p. = 118°-120°C

¹H-NMR (CDCl₃, δ ppm): 3.1-3.3 (m, 1 H), 4.9-5.2 (m, 2 H), 5.38 (s, 2 H), 6.89 (dd, J₁, J₂ = 8.27; 2.05 Hz, 1 H), 7.1-7.5 (m, 5 H), 7.82 (dt, J₁, J₂ = 8.01; 0.93, 1 H).

31b) 2-[[1-(3,4-dichlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid

[0142] The product was obtained via the method described in Example 30b), using [1-(3,4-dichlorobenzyl)-1H-indazol-3-yl]methanol as starting reagent. The product obtained was purified by crystallization from toluene.

m.p. = 174°-176°C

¹H-NMR (DMSO-d₆, δ ppm): 1.44 (s, 6 H), 4.76 (s, 2 H), 5.64 (s, 2 H), 7.12-7.22 (m, 2 H), 7.41 (t, J = 7.68 Hz, 1 H), 7.54 (d, J = 2.01 Hz, 1 H), 7.58 (d, J = 8.42 Hz, 1 H), 7.72 (d, J = 8.42 Hz, 1 H), 7.95 (d, J = 8.05 Hz, 1 H), 12.81 (bs, 1 H).

31c) 2-[[1-(3,4-dichlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanamide

[0143] The product was prepared via the procedures described in the preparation of compound 9, using 2-[[1-(3,4-dichlorobenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid as starting reagent.

Preparation of compound 34

2-[[1-(benzoyl-1H-indazol-3-yl)ethoxy]-N-hydroxy-2-methylpropanamide

34a) (1-tritylindazol-3-yl)methanol

[0144] To a solution containing isobutyl 1H-indazole-3-carboxylate (280 g; 1.28 mol) in chloroform (2 L) stirred at room temperature were added triethylamine (300 ml; 2.16 mol) and triphenylchloromethane (400 g; 1.4 mol). The solution was stirred at room temperature for 4 days, and water (500 ml) was then added. The organic phase was separated out and concentrated under reduced pressure. The crude residue obtained was used without further purification in the following step.

[0145] To a solution of the crude isobutyl 1-tritylindazole-3-carboxylate (180 g; 0.39 mol) in THF (1 L) stirred at room temperature was added slowly a suspension of LiAlH₄ (18 g; 0.48 mol) in THF (100 ml). Once the addition was complete, the mixture was stirred at room temperature for 30 minutes, and the reaction was then stopped by cooling the mixture to 0°C and successively adding water (40 ml), 2N NaOH (40 ml) and water (60 ml). The solid thus formed was filtered off and the solution was concentrated under reduced pressure. The crude residue obtained was purified by crystallization from a 1/1 hexane/ethyl acetate mixture. 120 g of (1-tritylindazol-3-yl)methanol were obtained.

m.p. = 192°-193°C

¹H-NMR (CDCl₃, δ ppm): 2.51 (t, J = 6.98 Hz, 1 H), 4.90 (d, J = 6.98 Hz, 2 H), 6.2-6.5 (m, 1 H), 6.9-7.4 (m, 17 H), 7.6-7.8

(m, 1 H).

34b) 2-(1-tritylindazol-3-ylmethoxy)-2-methylpropanoic acid

[0146] To a suspension of (1-tritylindazol-3-yl)methanol (78 g; 0.20 mol), acetone (260 ml) and water (0.5 ml) stirred at room temperature were added NaOH (76 g; 1.9 mol) and, slowly, a 1/1 chloroform/acetone mixture (100 ml). The reaction is exothermic and the rate of addition was adjusted so as to keep the reaction temperature close to the boiling point. 30 minutes after the end of addition, the reaction was stopped by cooling the mixture to room temperature and evaporating off the solvent under reduced pressure. The residue was taken up in water (500 ml) and washed with diethyl ether (3×100 ml). The aqueous phase was then acidified with concentrated HCl and the product was extracted with toluene (3×250 ml). The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by crystallization from a 3/7 hexane/ethyl acetate mixture to give 22 g of 2-(1-tritylindazol-3-ylmethoxy)-2-methylpropanoic acid.

m.p. = 179°-180°C.

¹H-NMR (CDCl₃, δ ppm): 1.53 (s, 6 H), 4.88 (s, 2 H), 6.3-6.5 (m, 1 H), 6.9-7.5 (m, 17 H), 7.8-8.0 (m, 1 H), 9.3 (bs, 1 H).

34c) 2-(1H-indazol-3-ylmethoxy)-2-methylpropanoic acid

[0147] To a solution of 2-(1-tritylindazol-3-ylmethoxy)-2-methylpropanoic acid (83 g; 0.174 mol) in dichloromethane (DCM) (900 ml) stirred at room temperature was added para-toluenesulfonic acid (PTSA) (50 g; 0.29 mol). The mixture was stirred for 30 minutes at room temperature and then poured into 5N NaOH (400 ml). The organic phase was separated out and washed with water (300 ml). The combined aqueous phases were acidified with concentrated HCl and then extracted with ethyl acetate (5×300 ml). The combined organic phases were evaporated under reduced pressure and the resulting crude residue was purified by crystallization from a 1/1 hexane/ethyl acetate mixture.

[0148] 42 g of 2-(1H-indazol-3-ylmethoxy)-2-methylpropanoic acid were obtained.

m.p. = 135°-137°C.

¹H-NMR (DMSO-d₆, δ ppm): 1.46 (s, 6 H), 4.77 (s, 2 H), 7.0-7.6 (m, 3 H), 7.94 (d, J = 7.88 Hz, 1 H).

34d) 2-[(1-benzoyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid

[0149] To a solution of 2-(1H-indazol-3-ylmethoxy)-2-methylpropanoic acid (6 g; 0.026 mol) in acetone (50 ml) stirred at room temperature was added K₂CO₃ (6.8 g; 0.049 mol) and then, slowly, a solution of benzoyl chloride (5 ml; 0.043 mol) in acetone (30 ml). The mixture was stirred at room temperature for 24 hours and then poured into water (1 L). The solution was then brought to basic pH with 5N NaOH and washed with diethyl ether (3×150 ml). The alkaline phase was then acidified with concentrated HCl and extracted with diethyl ether (3×300 ml). The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by crystallization from a 1/1 hexane/ethyl acetate mixture. 2 g of 2-[(1-benzoyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid were obtained.

m.p. = 132°-135°C.

¹H-NMR (CDCl₃, δ ppm): 1.61 (s, 6 H), 4.93 (s, 2 H), 7.41 (t, J = 7.60 Hz, 1 H), 7.46-7.66 (m, 4 H), 8.02-8.09 (m, 3 H), 8.68 (bs, 1 H), 8.53 (d, J = 8.42 Hz, 1 H).

34e) 2-[(1-benzoyl-1H-indazol-3-yl)methoxy]-N-hydroxy-2-methylpropanamide

[0150] The product was prepared via the procedures described in the preparation of compound 10, using 2-[(1-benzoyl-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid as starting reagent.

[0151] 0.34 g of 2-[(1-benzoyl-1H-indazol-3-yl)methoxy]-N-hydroxy-2-methylpropanamide were thus obtained.

Preparation of compound 35 2-[(1-(1,2,3,4-tetrahydronaphth-1-yl)-1H-indazol-3-yl)methoxy]-N-hydroxy-2-methylpropanamide

35a) 2-methyl-2-[(1-(1,2,3,4-tetrahydronaphth-1-yl)-1H-indazol-3-yl)methoxy]-propanoic acid

[0152] To a solution of 2-methyl-2-[(1H-indazol-3-yl)methoxy]propanoic acid (26 g; 0.093 mol) in DMF (200 ml) was added 60% NaH (10 g; 0.25 mol) and the mixture was stirred for 10 minutes at 60°C. 1-chloro-1,2,3,4-tetrahydronaphthalene (0.217 mol) was then added to the mixture and the whole was stirred at 60°C for 18 hours. The reaction was stopped by pouring the mixture into water (1 L), followed by acidifying with 5 N HCl and extracting the product with diethyl ether (3×250 ml). The residue obtained was dissolved in 95° ethanol (100 ml) and treated at reflux with 1N NaOH (200 ml) for 2 hours. The solution was then cooled to room temperature and washed with diethyl ether (3×200 ml). The

alkaline phase was then acidified with concentrated HCl and extracted with ethyl acetate (3×300 ml). The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by crystallization from ethyl acetate. 31 g of 2-methyl-2-[[1-(1,2,3,4-tetrahydronaphth-1-yl)-1H-indazol-3-yl]methoxy]-propanoic acid were thus obtained.

m.p. = 132°-134°C

¹H-NMR (CDCl₃, δ ppm): 1.60 (d, J = 2.15 Hz, 6 H), 1.84-2.02 (m, 1H), 2.03-2.18 (m, 1 H), 2.23-2.46 (m, 2 H), 2.90 (dt, J = 16.50; 4.85 Hz, 1 H), 2.98-3.12 (m, 1 H), 4.95 (s, 2 H), 5.92 (dd, J = 8.83; 6.52 Hz, 1 H), 6.71 (d, J = 7.76 Hz, 1 H), 6.93-7.05 (m, 2 H), 7.08-7.32 (m, 4 H), 7.85 (d, J = 8.09 Hz, 1 H).

35b) 2-[[1-(1,2,3,4-tetrahydronaphth-1-yl)-1H-indazol-3-yl]methoxy]-N-hydroxy-2-methylpropanamide.

[0153] The product was obtained using the same procedure described in the preparation of compound 10, using 2-methyl-2-[[1-(1,2,3,4-tetrahydronaphth-1-yl)-1H-indazol-3-yl]methoxy]-propanoic acid as starting material.

Preparation of compound 36

{[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy}-N-hydroxyacetamide

36a) [1-(4-methoxybenzyl)-1H-indazol-3-yl]methanol

[0154] To a suspension of 60% NaH (2.7 g; 0.07 mol) in toluene (200 ml) was added 1-benzyl-3-hydroxymethylindazole (10 g; 0.07 mol). The mixture was brought to the boiling point and left stirring at reflux for 1 hour. 4-Methoxybenzyl chloride (14 g; 0.09 mol) was then added. The mixture was then stirred at reflux for 4 hours. The reaction was stopped by cooling the mixture to room temperature and adding water (50 ml). The organic phase was separated out and washed, respectively, with 2N HCl (50 ml) and water (5×50 ml). The solvent was evaporated off under reduced pressure. The crude residue thus obtained was purified by flash chromatography on silica gel, using as eluent a 3/2 hexane/ethyl acetate mixture. The product obtained was crystallized from a 5/1 hexane/ethyl acetate mixture to give 5.1 g of [1-(4-methoxybenzyl)-1H-indazol-3-yl]methanol with a melting point of 95-97°C.

¹H-NMR (CDCl₃, δ ppm): 3.43 (t, J = 6.9 Hz, 1 H), 3.67 (s, 3 H), 4.98 (d, J = 6.9 Hz, 2 H), 5.36 (s, 2 H), 6.5-6.8 (m, 2 H), 6.9-7.4 (m, 7 H), 7.80 (d, J = 7.86 Hz, 1 H).

36b) {[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy}acetic acid

[0155] A suspension containing [1-(4-methoxybenzyl)-1H-indazol-3-yl]methanol (6 g; 0.022 mol), bromoacetic acid (4 g; 0.03 mol) and 50% NaH (3 g; 0.066 mol) in THF (170 ml) was stirred at reflux for 72 hours. The reaction was then stopped by diluting with a suspension with water and ice (300 ml) and washing the aqueous phase with diethyl ether (3×150 ml). The aqueous phase was acidified with concentrated HCl. The solid thus formed was filtered off and purified by crystallization from isopropanol. 4.5 g of {[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy}acetic acid were thus obtained.

36c) {[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy}-N-hydroxyacetamide

[0156] The product was obtained with the method described for the preparation of compound 10, using {[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy}acetic acid as starting material.

Preparation of compound 39

{2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy}acetamide

39a) 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropan-1-ol

[0157] To a suspension of LiAlH₄ (4.48 g; 0.118 mol) in diethyl ether (100 ml) stirred at room temperature was slowly added a solution of methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropanoate, prepared according to the method described in EP 0 382 276, (20 g; 0.06 mol) in diethyl ether (200 ml) and THF (50 ml). Once the addition was complete, the mixture was stirred at room temperature for 30 minutes and the reaction was then completed by adding 10 N NaOH (20 ml) and water (40 ml). The solvent was evaporated off under reduced pressure and the oily residue was purified by distillation at 0.01 mmHg at 190°C. The solid product thus obtained was crystallized from isopropanol.

[0158] 11 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropan-1-ol were thus obtained.

m.p. = 52°-53°C

¹H-NMR (CDCl₃, δ ppm): 1.34 (s, 6 H), 2.50 (bs, 1 H), 3.51 (s, 2 H), 4.87 (s, 2 H), 5.55 (s, 2 H), 7.14 (ddd, J = 8.04; 6.21; 1.68 Hz, 1 H), 7.17-7.38 (m, 7 H), 7.78 (dt, J = 8.08; 1.00 Hz, 1H).

39b) {2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy}acetic acid

[0159] To a solution of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropan-1-ol (11.0g, 0.04 mol) in dry tetrahydrofuran (THF) (100 ml) stirred at room temperature was added 60% sodium hydride (1.6 g, 0.04 mol). The mixture was heated at the boiling point for 2 hours and then cooled to room temperature, and a solution of ethyl bromoacetate (7.4 g, 0.044 mol) in THF (7 ml) was slowly added thereto. Once the addition was complete, the mixture was refluxed for a further 2 hours. The reaction was stopped by cooling to room temperature and evaporating off the solvent under reduced pressure. The residue was taken up in 2N NaOH (100 ml) and the product was extracted with diethyl ether (3×150 ml). The combined organic phases were concentrated under reduced pressure.

[0160] The crude residue was taken up in a solution of NaOH (1.9 g, 0.045 mol) in a 1/1 water/ethanol mixture (160 ml). The mixture was then stirred at reflux for 2 hours. The reaction was stopped by concentrating the solvent under reduced pressure, and the residue was taken up in water (100 ml) and washed with diethyl ether (3×50 ml). The alkaline phase was then acidified with concentrated HCl and the solid formed was filtered off. The product was purified by double crystallization from a 1/2 hexane/ethyl acetate mixture. 4.6 g of {2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy}acetic acid were thus obtained.

39c) {2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy} acetamide

[0161] The product was prepared via the procedures described in the preparation of compound 9, using {2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy}acetic acid as starting reagent.

Preparation of compound 40

{2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy}-N-hydroxyacetamide

[0162] The product was prepared via the procedures described in the preparation of compound 10, using {2-[(1-benzyl-1H-indazol-3-yl)methoxy]-2-methylpropoxy}acetic acid as starting reagent.

Preparation of compound 41

{2-[(1-(4-methoxybenzyl)-1H-indazol-3-yl)methoxy]-2-methylpropoxy}-N-hydroxy-acetamide

41a) [1-(4-methoxybenzyl)-1H-indazol-3-yl]methanol

[0163] To a suspension of 60% NaH (2.7 g; 0.07 mol) in toluene (200 ml) was added 1-benzyl-3-hydroxymethylindazole (10 g; 0.07 mol). The mixture was brought to the boiling point and left stirring at reflux for 1 hour. 4-Methoxybenzyl chloride (14 g; 0.09 mol) was then added. The mixture was then stirred at reflux for 4 hours. The reaction was stopped by cooling the mixture to room temperature and adding water (50 ml). The organic phase was separated out and washed, respectively, with 2N HCl (50 ml) and water (5×50 ml). The solvent was evaporated off under reduced pressure. The crude residue thus obtained was purified by flash chromatography on silica gel, using as eluent a 3/2 hexane/ethyl acetate mixture. The product obtained was crystallized from a 5/1 hexane/ethyl acetate mixture to give 5.1 g of [1-(4-methoxybenzyl)-1H-indazol-3-yl]methanol with a melting point of 95-97°C.

¹H-NMR (CDCl₃, δ ppm): 3.43 (t, J = 6.9 Hz, 1 H), 3.67 (s, 3 H), 4.98 (d, J = 6.9 Hz, 2 H), 5.36 (s, 2 H), 6.5-6.8 (m, 2 H), 6.9-7.4 (m, 7 H), 7.80 (d, J = 7.86 Hz, 1 H).

41b) 2-[(1-(4-methoxybenzyl)-1H-indazol-3-yl)methoxy]-2-methylpropanoic acid

[0164] To a suspension of NaOH (15.6 g; 0.39 mol) in acetone (50 ml) was added [1-(4-methoxybenzyl)-1H-indazol-3-yl]methanol (8.7 g; 0.03 mol). To the mixture was slowly added a solution of chloroform (7.2 ml; 0.09 mol) and acetone (7.2 ml; 0.1 mol). The addition caused refluxing of the mixture of solvents. Once the addition was complete, the mixture was refluxed for 1 hour. The reaction was stopped by cooling the mixture to room temperature and evaporating off the solvent under reduced pressure. The resulting crude residue was taken up in toluene (100 ml) and water (50 ml). The aqueous phase was separated from the organic phase and then washed with toluene (2×50 ml). The combined organic phases were extracted with water (3×50 ml). The combined aqueous phases were washed with hexane (2×30 ml) and

then acidified with 2N HCl and stirred at room temperature. The solid thus obtained was filtered off and crystallized first from a 5/1 water/acetic acid mixture, and then from toluene, to give 4.8 g of 2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid with a melting point of 169-171°C.

¹H-NMR (DMSO-d₆, δ ppm): 1.44 (s, 6 H), 3.69 (s, 3 H), 4.74 (s, 2 H), 5.52 (s, 2 H), 6.82-6.90 (m, 2 H), 7.13 (t, J = 7.50 Hz, 1 H), 7.18-7.26 (m, 2 H), 7.36 (t, J = 7.23 Hz, 1 H), 7.66 (d, J = 8.42 Hz, 1 H), 7.92 (dd, J = 8.14; 1.01 Hz, 1 H), 12.76 (s, 1 H).

41c) 2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropan-1-ol

[0165] The product was obtained with the method described for the preparation of compound 39a, using 2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropanoic acid as starting material.

41 d) (2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropoxy)acetic acid

[0166] The product was obtained with the method described for the preparation of compound 39b, using 2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropan-1-ol as starting material.

41e) (2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropoxy)-N-hydroxy-acetamide

[0167] The product was obtained with the method described for the preparation of compound 10, using (2-[[1-(4-methoxybenzyl)-1H-indazol-3-yl]methoxy]-2-methylpropoxy)acetic acid as starting material.

Preparation of compound 42

2-2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethoxy]-N-hydroxy-2-methylpropanamide

42a) 2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethanol

[0168] To a solution of NaOH (2.8 g; 0.07 mol) in ethylene glycol (150 ml) stirred at room temperature was added 1-benzyl-3-chloromethylindazole, prepared as described in EP 0 382 276 (17.6 g; 0.07 mol). The solution was heated at 130°C for 4 hours and then cooled to room temperature and the solvent was evaporated off under reduced pressure. The residue was taken up in water (100 ml) and the product was extracted with ethyl acetate (3×100 ml). The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by crystallization from an approximately 1/1 hexane/ethyl acetate mixture.

[0169] 13.8 g of 2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethanol were thus obtained.

m.p. = 67°-69°C

¹H-NMR (CDCl₃, δ ppm): 2.15 (bs, 1 H), 3.61-3.82 (m, 4 H), 4.97 (s, 2 H), 5.57 (s, 2 H), 7.11-7.38 (m, 8 H), 7.81 (dt, J = 8.15; 0.97 Hz, 1 H).

42b) 2-2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethoxy]-2-methylpropanoic acid

[0170] To a mixture of 2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethanol (35 g, 0.124 mol) and NaOH (63 g, 1.57 mol) in acetone (180 ml) and water (1 ml), stirred at room temperature, was slowly added a 1/1 chloroform/acetone mixture (80 ml). During the addition, the temperature rose to the reflux point. Once the addition was complete, the solvent was evaporated off under reduced pressure and the residue was taken up in water (100 ml) and washed with diethyl ether (3×50 ml). The aqueous phase was acidified with glacial acetic acid and then extracted with diethyl ether (3×150 ml). The combined organic phases were concentrated under reduced pressure. The crude residue obtained was purified by crystallization from a 1/1 hexane/ethyl acetate mixture.

[0171] 12.4 g of 2-2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethoxy]-2-methylpropanoic acid were obtained.

m.p. = 94°-95°C

¹H-NMR (CDCl₃, δ ppm): 1.43 (s, 6 H), 3.56-3.63 (m, 2 H), 3.65-3.71 (m, 2 H), 5.03 (s, 2 H), 5.58 (s, 2 H), 7.13-7.39 (m, 8 H), 7.83 (dt, J = 8.05; 0.82 Hz, 1 H), 9.60 (bs, 1 H).

42c) 2-2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethoxy]-N-hydroxy-2-methylpropanamide

[0172] The product was obtained with the method described for the preparation of compound 34e), using 2-2-[[1-(benzyl-1H-indazol-3-yl)methoxy]ethoxy]-2-methylpropanoic acid as starting material.

Preparation of compound 43

2-{3-[(1-benzyl-1H-indazol-3-yl)methoxy]propoxy}-N-hydroxy-2-methylpropanamide

5 43a) 3-[(1-benzyl-1H-indazol-3-yl)methoxy]propan-1-ol

[0173] To a solution of NaOH (2.8 g; 0.07 mol) in 1,3-propanediol (150 ml) stirred at room temperature was added 1-benzyl-3-chloromethylindazole, prepared as described in EP 0 382 276 (17.6 g; 0.07 mol). The solution was heated at 130°C for 4 hours and then cooled to room temperature, and the solvent was evaporated off under reduced pressure. The residue was taken up in water (100 ml) and the product was extracted with ethyl acetate (3×100 ml). The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by flash chromatography on silica gel, using an approximately 1/1 hexane/ethyl acetate mixture as eluent.

[0174] 10.5 g of 3-[(1-benzyl-1H-indazol-3-yl)methoxy]propan-1-ol were thus obtained.

¹H-NMR (CDCl₃, δ ppm): 1.85 (q, J = 5.83 Hz, 2 H), 2.75 (bs, 1 H), 3.71 (t, J = 7.74 Hz, 4 H), 4.91 (s, 2 H), 5.55 (s, 2 H), 7.0-7.4 (m, 8 H), 7.80 (d, J = 7.77 Hz, 1H).

43b) 2-{3-[(1-benzyl-1H-indazol-3-yl)methoxy]propoxy}-2-methyl-propanoic acid

[0175] The product was obtained via the method described in example 42b), using as starting material 3-[(1-benzyl-1H-indazol-3-yl)methoxy]propan-1-ol. The product was purified by double crystallization from a 7/3 hexane/ethyl acetate mixture.

m.p. = 57°-59°C

¹H-NMR (DMSO-d₆, δ ppm): 1.25 (s, 6 H), 1.72 (quint., J = 6.40 Hz, 2 H), 3.37 (t, J = 6.40 Hz, 2 H), 3.53 (t, J = 6.40 Hz, 2 H), 4.77 (s, 2 H), 5.62 (s, 2 H), 7.14 (ddd, J = 8.00; 7.00; 0.73 Hz, 1 H), 7.19-7.33 (m, 5 H), 7.38 (ddd, J = 8.37; 7.00; 0.91 Hz, 1 H), 7.66 (d, J = 8.42 Hz, 1 H), 7.79 (dt, J = 8.05; 0.91 Hz, 1 H), 12.46 (s, 1 H).

43c) 2-{3-[(1-benzyl-1H-indazol-3-yl)methoxy]propoxy}-N-hydroxy-2-methylpropanamide

[0176] The product was obtained with the method described for the preparation of compound 10, using 2-{3-[(1-benzyl-1H-indazol-3-yl)methoxy]propoxy}-2-methylpropanoic acid as starting material.

Preparation of compound 46

{2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}acetamide

46a) 2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethanol

[0177] To a solution of NaOH (2.8 g; 0.07 mol) in ethylene glycol (150 ml) stirred at room temperature was added 1-benzyl-3-chloromethylindazole, prepared as described in EP 0 382 276 (17.6 g; 0.07 mol). The solution was heated at 130°C for 4 hours and then cooled to room temperature and the solvent was evaporated off under reduced pressure. The residue was taken up in water (100 ml) and the product was extracted with ethyl acetate (3×100 ml). The combined organic phases were concentrated under reduced pressure and the crude residue obtained was purified by crystallization from an approximately 1/1 hexane/ethyl acetate mixture.

[0178] 13.8 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethanol were thus obtained.

m.p. = 67°-69°C

¹H-NMR (CDCl₃, δ ppm): 2.15 (bs, 1 H), 3.61-3.82 (m, 4 H), 4.97 (s, 2 H), 5.57 (s, 2 H), 7.11-7.38 (m, 8 H), 7.81 (dt, J = 8.15; 0.97 Hz, 1 H).

46b) {2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}acetic acid

[0179] To a solution of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethanol (11.28 g, 0.04 mol) in dry tetrahydrofuran (THF) (100 ml) stirred at room temperature was added 60% sodium hydride (1.6 g, 0.04 mol). The mixture was heated at the boiling point for 2 hours and then cooled to room temperature, and a solution of ethyl bromoacetate (7.4 g, 0.044 mol) in THF (7 ml) was slowly added thereto. Once the addition was complete, the mixture was refluxed for a further 2 hours. The reaction was stopped by cooling to room temperature and evaporating off the solvent under reduced pressure. The residue was taken up in 2N NaOH (100 ml) and the product was extracted with diethyl ether (3×150 ml). The combined organic phases were concentrated under reduced pressure.

[0180] The crude residue was taken up in a solution of NaOH (1.9 g, 0.045 mol) in a 1/1 water/ethanol mixture (160

ml). The mixture was then stirred at reflux for 2 hours. The reaction was stopped by concentrating the solvent under reduced pressure, and the residue was taken up in water (100 ml) and washed with diethyl ether (3×50 ml). The alkaline phase was then acidified with concentrated HCl and the solid formed was filtered off. The product was purified by double crystallization from a 1/3 hexane/ethyl acetate mixture.

[0181] 5.8 g of {2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}acetic acid were obtained.

m.p. = 101°-103°C

¹H-NMR (CDCl₃, δ ppm): 3.66-3.79 (m, 4H), 4.13 (s, 2 H), 5.01 (s, 2 H), 5.58 (s, 2 H), 7.66 (bs, 1 H), 7.14-7.40 (m, 8 H), 7.84 (dt, J = 8.11; 0.99 Hz, 1 H).

46c) {2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}acetamide

[0182] The product was obtained with the method described for the preparation of compound 9, using {2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy} acetic acid as starting material.

Preparation of compound 47

{2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}-N-hydroxy-acetamide

[0183] The product was obtained with the method described for the preparation of compound 10 using {2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}acetic acid as starting material.

Preparation of compound 48

{2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy}-acetohydrazide

[0184] A suspension of {2-[(1-benzyl-1H-indazol-3-yl)methoxy]ethoxy} acetic acid (19 g; 0.06 mol) in methanol (300 ml) was treated at 0°C with gaseous HCl for 4 hours. The mixture was then poured into water (500 ml) and the product was extracted with diethyl ether (3×250 ml). The combined organic phases were washed with 5% sodium bicarbonate solution (2×100 ml) and then with water (50 ml). The solvent was evaporated off under reduced pressure and the crude residue was purified by crystallization from hexane. 16 g of methyl 2-[(1-benzoyl-1H-indazol-3-yl)methoxy]-acetate was thus obtained.

[0185] To a solution of 1 M hydrazine hydrate (100 ml, 0.100 mol) stirred at 80°C was added portion wise methyl 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-acetate (19 g, 0.06 mol). Once the addition was complete, the mixture was stirred for 30 minutes at the same temperature and then heated at 120°C for 2 hours. The reaction was stopped by diluting the mixture with water (600 ml) and extracting the product with diethyl ether (4×200 ml). The combined organic phases were then extracted with 2N HCl (4×200 ml). The acidic phase was then brought to basic pH with 10N NaOH and again extracted with diethyl ether (4×200 ml). The combined organic phases were concentrated under reduced pressure and the crude residue was purified by crystallization from a 1/1 hexane/ethyl acetate mixture.

[0186] 15 g of 2-[(1-benzyl-1H-indazol-3-yl)methoxy]-acetohydrazide were thus obtained.

Example 1

Analysis of the gene expression of MCP-1 in a human monocyte line

[0187] The capacity of the compounds to inhibit the expression of MCP-1 by lipopolysaccharide (LPS)-stimulated MonoMac6 cells was evaluated. The cells were placed in 96-well plates at a concentration of 50 000 cells/well. The compounds were tested at the maximum soluble concentration given in Table 1 (in the range 30-300 μM) and incubated for 1 hour. The cells were then stimulated with LPS (100 ng/ml) for 4 hours.

[0188] The total RNA was extracted from the cell pellet using the RNeasy mini kit (Qiagen), reverse-transcribed with the TaqMan Reverse transcription reagents synthesis kit (Applied Biosystems) and the cDNA obtained was used for the real-time PCR reaction. The amplification was obtained in 96-well plates using the ABI Prism 7000 sequence detection system (Applied Biosystems), by applying the following temperature profile: 50°C for 2 minutes, 95°C for 10 minutes and 45 cycles at 95°C for 15 seconds and 60°C for 1 minute. For the amplification, a set of primers and probe specific for human MCP-1 was used (Applied Biosystems, RefSeq NM_002982.3). A set of primers and probe for β-actin was used in separate wells as an internal control of the samples for the purposes of normalization. Once the reaction had taken place, the fluorescence data were analysed using the ABI Prism 7000 SDS software, by calculating the threshold cycle (Ct) for each sample and subsequently performing a relative quantification via the ΔΔCt method.

[0189] The results obtained, expressed as a percentage of inhibition, are collated in Table 1 below.

TABLE 1

No.	% inhibition	[μ M]
9	99	300
10	61	150
11	68	150
12	20	300
13	72	75
14 *	49	30
15 *	40	75
* reference compound not part of the invention		

[0190] As shown by the results obtained and given in Table 1, the compounds were capable of significantly inhibiting the LPS-induced expression of MCP-1 in a human monocyte line, and showed a reduction in the levels of specific mRNA between 20% and 99%.

Example 2

Measurement of the production of MCP-1 in a human monocyte line

[0191] The capacity of the compounds to inhibit the expression of the protein MCP-1 by lipopolysaccharide (LPS)-stimulated MonoMac6 cells was evaluated. The cells were placed in 96-well plates at a concentration of 50 000 cells/well. The compounds were tested at the maximum soluble concentration given in Table 2 (in the range 30-300 μ M) and incubated for 1 hour. The cells were then stimulated with LPS (100 ng/ml) for 20 hours.

[0192] The amount of MCP-1 produced was measured in the supernatants, suitably diluted with buffer, by means of an immunoenzymatic test (ELISA) using a commercial kit (ELISA MCP-1/JE, R&D Systems).

[0193] The results obtained, expressed as a percentage of inhibition, are collated in Table 2 below.

TABLE 2

No.	% inhibition	[μ M]
9	98	300
10	91	150
11	79	150
12	48	300
13	74	75
14*	70	30
15*	59	75
27*	82	30
* reference compound not part of the invention		

[0194] As shown by the results obtained and given in Table 2, the compounds were capable of significantly inhibiting the LPS-induced expression of MCP-1 in a human monocyte line, and showed a reduction in the levels of produced protein between 48% and 98%.

Example 3Analysis of the gene expression of CX3CR1 in a human monocyte line

[0195] The capacity of the compounds to inhibit the expression of CX3CR1 by lipopolysaccharide (LPS)-stimulated MonoMac6 cells was evaluated. The cells were placed in 96-well plates at a concentration of 50 000 cells/well. The compounds were tested at the maximum soluble concentration given in Table 3 (in the range 30-300 μ M) and incubated for 1 hour. The cells were then stimulated with LPS (100 ng/ml) for 20 hours.

[0196] The total RNA was extracted from the cell pellet using the RNeasy mini kit (Qiagen), reverse-transcribed with the TaqMan Reverse transcription reagents synthesis kit (Applied Biosystems) and the cDNA obtained was used for the real-time PCR reaction. The amplification was obtained in 96-well plates using the ABI Prism 7000 sequence detection system (Applied Biosystems), by applying the following temperature profile: 50°C for 2 minutes, 95°C for 10 minutes and 45 cycles at 95°C for 15 seconds and 60°C for 1 minute. For the amplification, a set of primers and probe specific for human CX3CR1 was used (Applied Biosystems, RefSeq NM_001337.3). A set of primers and probe for β -actin was used in separate wells as an internal control of the samples for the purposes of normalization. Once the reaction had taken place, the fluorescence data were analysed using the ABI Prism 7000 SDS software, by calculating the threshold cycle (Ct) for each sample and subsequently performing a relative quantification via the $\Delta\Delta$ Ct method.

[0197] The results obtained, expressed as a percentage of inhibition, are collated in Table 3 below.

TABLE 3

No.	% inhibition	[μ M]
9	99	300
10	95	150
11	90	150
12	83	300
13	100	75
14 *	44	30
15 *	89	75
* reference compound not part of the invention		

[0198] As shown by the results obtained and given in Table 3, the compounds were capable of significantly inhibiting the LPS-induced expression of CX3CR1 in a human monocyte line, and showed a reduction in the levels of specific mRNA between 44% and 100%.

Example 4Analysis of the gene expression of p40 in a human monocyte line

[0199] The capacity of the compounds to inhibit the expression of p40 by lipopolysaccharide (LPS)-stimulated MonoMac6 cells was evaluated. The cells were placed in 96-well plates at a concentration of 50 000 cells/well. The compounds were tested at the maximum soluble concentration given in Table 4 (in the range 30-300 μ M) and incubated for 1 hour. The cells were then stimulated with LPS (100 ng/ml) for 4 hours.

[0200] The total RNA was extracted from the cell pellet using the RNeasy mini kit (Qiagen), reverse-transcribed with the TaqMan Reverse transcription reagents synthesis kit (Applied Biosystems) and the cDNA obtained was used for the real-time PCR reaction. The amplification was obtained in 96-well plates using the ABI Prism 7000 sequence detection system (Applied Biosystems), by applying the following temperature profile: 50°C for 2 minutes, 95°C for 10 minutes and 45 cycles at 95°C for 15 seconds and 60°C for 1 minute. For the amplification, a set of primers and probe specific for human p40 was used (Applied Biosystems, RefSeq NM_002187.2). A set of primers and probe for β -actin was used in separate wells as an internal control of the samples for the purposes of normalization. Once the reaction had taken place, the fluorescence data were analysed using the ABI Prism 7000 SDS software, by calculating the threshold cycle (Ct) for each sample and subsequently performing a relative quantification via the $\Delta\Delta$ Ct method.

[0201] The results obtained, expressed as a percentage of inhibition, are collated in Table 4 below.


TABLE 4

No.	% inhibition	[μ M]
9	96	300
10	54	150
11	44	150
13	59	75
14 *	27	30
15 *	33	75
* reference compound not part of the invention		

[0202] As shown by the results obtained and given in Table 4, the compounds were capable of significantly inhibiting the LPS-induced expression of p40 in a human monocyte line, and showed a reduction in the levels of specific mRNA between 33% and 96%.

Claims

1. Compound of formula (I)



in which:

A may be $-X_1-$ or $-X_1-OC(R_9)(R_{10})-$, in which

X_1 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms, and R_9 and R_{10} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

Y may be $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$, in which

R_{11} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, or R_{11} together with R_{12} forms a 4- to 7-membered heterocycle,

R_{12} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms, or R_{12} together with R_{11} , forms a 4- to 7-membered heterocycle,

R_{13} and R_{15} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R₁₄ may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, COR', COOR', CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

X₂ may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms,

X₃ may be OH, NH₂, NHOH or NHNH₂,

R₁ and R₂, which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,


R₃, R₄ and R₈, which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, -OH, -N(R')(R''), -N(R')COR'', -CN, -CONR'R'', -SO₂NR'R'', -SO₂R', nitro and trifluoromethyl; with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

R₅ may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, -OH, -N(R')(R''), -N(R')COR'', nitro and trifluoromethyl, or R₅ together with one from among R₆ and R₇ forms a ring having 5 or 6 carbon atoms; with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

R₆ and R₇, which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, or together form a group C=O, or one from among R₆ and R₇, together with R₅, forms a ring having 5 or 6 carbon atoms.

2. Compound according to Claim 1, **characterized in that** X₁ is an alkyl group having from 1 to 3 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 3 carbon atoms or one or more alkoxy groups having 1 or 2 carbon atoms.
3. Compound according to Claim 1, **characterized in that** X₁ is chosen from the group comprising a group CH₂, a group CH₂CH₂ or a group C(CH₃)₂, and R₉ and R₁₀, which may be identical or different each other, are hydrogen or a CH₃ group.
4. Compound according to Claim 1, **characterized in that** residue A is chosen from the group comprising a group CH₂, a group CH₂CH₂, a group C(CH₃)₂, a group CH₂CH₂OCH₂, a group CH₂CH₂OC(CH₃)₂ and a group CH₂CH₂CH₂OC(CH₃)₂.
5. Compound according to Claim 1, **characterized in that** R₁₁ is a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, or R₁₁ together with R₁₂ forms a 5- or 6-membered heterocycle.
6. Compound according to Claim 1, **characterized in that** R₁₂ is a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 3 carbon atoms, or R₁₂ together with R₁₁ forms a 5- or 6-membered heterocycle.
7. Compound according to Claim 1, **characterized in that** R₁₃ and R₁₅, which may be identical or different each other, are a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms or an alkoxy group having 1 or 2 carbon atoms.
8. Compound according to Claim 1, **characterized in that** R₁₄ is a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, COR', COOR', CON(R')(R'') with R' and R'', which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 3 carbon atoms.
9. Compound according to Claim 1, **characterized in that** X₂ is an alkyl group having from 1 to 3 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 3 carbon atoms or one or more alkoxy groups having 1 or 2 carbon atoms.
10. Compound according to Claim 1, **characterized in that** X₃ is chosen from the group comprising OH, NH₂, NHOH and NHNH₂.
11. Compound according to Claim 1, **characterized in that** R₁ and R₂, which may be identical or different each other, are a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms or an alkoxy group having 1 or 2 carbon atoms.

12. Compound according to Claim 1, **characterized in that** R_3 , R_4 and R_8 , which may be identical or different each other, are chosen from the group comprising a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, a Br, Cl or F atom, an OH group, a nitro group, a trifluoromethyl group or a group $N(R')(R'')$ or $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, with R' and R'' , which may be identical or different each other, represented by a hydrogen atom and an alkyl group having from 1 to 3 carbon atoms.
13. Compound according to Claim 1, **characterized in that** R_5 is chosen from the group comprising a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having 1 or 2 carbon atoms, a halogen atom, an OH group, or R_5 , together with one from among R_6 and R_7 , forms a ring having 5 or 6 carbon atoms.
14. Compound according to Claim 1, **characterized in that** R_6 and R_7 , which may be identical or different each other, are chosen from the group comprising a hydrogen atom, an alkyl group having from 1 to 3 carbon atoms, or together form a group $C=O$, or one from among R_6 and R_7 , together with R_5 , forms a ring having 5 or 6 carbon atoms.
15. Pharmaceutical composition comprising the compound of formula (I) according to any one of the preceding claims, or a pharmaceutically acceptable salt, or ester thereof, and at least one pharmaceutically acceptable vehicle.
16. Pharmaceutical composition according to Claim 15, **characterized in that** the said pharmaceutically acceptable salt is an addition salt with physiologically acceptable organic or mineral acids or bases.
17. Pharmaceutical composition according to Claim 16, **characterized in that** the said physiologically acceptable acids are chosen from the group comprising hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid, acetic acid, ascorbic acid, benzoic acid, citric acid, fumaric acid, lactic acid, maleic acid, methanesulfonic acid, oxalic acid, para-toluenesulfonic acid, benzenesulfonic acid, succinic acid, tannic acid and tartaric acid.
18. Pharmaceutical composition according to Claim 16, **characterized in that** the said physiologically acceptable bases are chosen from the group comprising ammonium hydroxide, calcium hydroxide, magnesium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, arginine, betaine, caffeine, choline, N,N-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, N-methylglucamine, glucamine, glucosamine, histidine, N-(2-hydroxyethyl)piperidine, N-(2-hydroxyethyl)pyrrolidine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, theobromine, triethylamine, trimethylamine, tripropylamine and tromethamine.
19. Pharmaceutical composition according to Claim 15, **characterized in that** the said pharmaceutically acceptable ester is formed with physiologically acceptable organic acids or alcohols.
20. Pharmaceutical composition according to any one of Claims 15 to 19, **characterized in that** the said composition contains a stereoisomer or an enantiomer of the compound of formula (I), or a pharmaceutically acceptable salt, or ester thereof, or a mixture thereof.
21. Pharmaceutical composition according to any one of Claims 15 to 20, **characterized in that** the said pharmaceutically acceptable vehicle is chosen from the group comprising glidants, binders, disintegrants, fillers, diluents, flavourings, colorants, fluidizers, lubricants, preserving agents, humectants, absorbents and sweeteners.
22. Use of a compound of formula (I)



in which:

A may be $-X_1-$ or $-X_1-OC(R_9)(R_{10})-$, in which

X_1 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms, and R_9 and R_{10} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

Y may be $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$, in which

R_{11} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, or R_{11} together with R_{12} forms a 4- to 7-membered heterocycle,

R_{12} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms, or R_{12} together with R_{11} , forms a 4- to 7-membered heterocycle,

R_{13} and R_{15} , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R_{14} may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, COR' , $COOR'$, $CON(R')(R'')$ with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,

X_2 may be an alkyl group having from 1 to 5 carbon atoms, optionally substituted with one or more alkyl groups having from 1 to 5 carbon atoms or one or more alkoxy groups having from 1 to 3 carbon atoms,

X_3 may be OH, NH_2 , $NHOH$ or $NHNH_2$,

R_1 and R_2 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms or an alkoxy group having from 1 to 3 carbon atoms,

R_3 , R_4 and R_8 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro and trifluoromethyl; with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms, R_5 may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a halogen atom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, nitro and trifluoromethyl, or R_5 together with one from among R_6 and R_7 forms a ring having 5 or 6 carbon atoms; with R' and R'' , which may be identical or different each other, represented by hydrogen and an alkyl group having from 1 to 5 carbon atoms,


R_6 and R_7 , which may be identical or different each other, may be hydrogen, an alkyl group having from 1 to 5 carbon atoms, or together form a group $C=O$, or one from among R_6 and R_7 , together with R_5 , forms a ring having 5 or 6 carbon atoms,

for the preparation of a pharmaceutical composition for treating diseases based on the expression of MCP-1, CX3CR1 and p40, wherein said diseases based on the expression of MCP-1 and CX3CR1 are chosen from the group comprising articular diseases, renal diseases, cardiovascular diseases, metabolic syndrome, obesity, diabetes, insulin resistance and cancer, and said diseases based on the expression of p40 are chosen from the group comprising autoimmune diseases, chronic degenerative inflammatory diseases, and cancer.

23. Use according to Claim 22, **characterized in that** the said diseases based on the expression of MCP-1 are chosen from the group comprising rheumatoid arthritis, arthritis induced by viral infections, psoriatic arthritis, arthrosis, lupus nephritis, diabetic nephropathy, glomerulonephritis, polycystic kidney disease, interstitial lung disease, fibrosis, multiple sclerosis, Alzheimer's disease, HIV-associated dementia, atopic dermatitis, psoriasis, vasculitis, restenosis, atherosclerosis, myocardial infarction, angina, acute coronary diseases, adenomas, carcinomas and metastases, and metabolic diseases.
24. Use according to Claim 22, **characterized in that** the said diseases based on the expression of CX3CR1 are chosen from the group comprising rheumatoid arthritis, lupus nephritis, diabetic nephropathy, Crohn's disease, ulcerative colitis, coronary disorders, restenosis, atherosclerosis, myocardial infarction, and angina.
25. Use according to Claim 22, **characterized in that** the said diseases based on the expression of p40 are chosen from the group comprising rheumatoid arthritis, psoriasis, glomerulonephritis, diabetes, lupus erythematosus, diabetes, Crohn's disease, and tumours.

Patentansprüche

1. Verbindung gemäß Formel (I)



worin

A für -X₁- oder -X₁-OC(R₉)(R₁₀)- stehen kann, wobei

X₁ für eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen kann, die gegebenenfalls mit einer oder mehreren Alkylgruppen mit 1 bis 5 Kohlenstoffatomen oder mit einer oder mehreren Alkoxygruppen mit 1 bis 3 Kohlenstoffatomen substituiert sein kann.

und

R₉ und R₁₀, die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen kann,

Y für $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$ stehen kann, wobei R_{11} für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen kann oder R_{11} zusammen mit R_{12} einen 4- bis 7-gliedrigen Heterocyclus bildet,

R₁₂ für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, COR', COOR', CON(R')(R'') stehen kann, wobei R' und R'', die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen, oder R₁₂ zusammen mit R₁₁ einen 4- bis 7-gliedrigen Heterocyclus bildet.

R₁₃ und R₁₅, die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen können,

R₁₄ für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, COR', COOR', CON(R')(R'') stehen kann, wobei R' und R'', die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen.

X₂ für eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen kann, die gegebenenfalls mit einer oder mehreren Alkylgruppen mit 1 bis 5 Kohlenstoffatomen oder mit einer oder mehreren Alkoxygruppen mit 1 bis 3 Kohlenstoffatomen substituiert sein kann.

X_3 für OH , NH_2 , $NHOH$ oder $NHNH_2$ stehen kann,

R₁ und R₂, die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen kann,

R₃, R₄ und R₈, die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoff-

atomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, ein Halogenatom, -OH, -N(R')(R''), -N(R')COR'', -CN, -CONR'R'', -SO₂NR'R'', -SO₂R', Nitro und Trifluormethyl stehen können; wobei R' und R'', die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen,

R₅ für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, ein Halogenatom, -OH, -N(R')(R''), -N(R')COR'', Nitro und Trifluormethyl stehen kann, oder R₅ zusammen mit einem der Reste R₆ oder R₇ einen Ring mit 5 oder 6 Kohlenstoffatomen bildet; wobei R' und R'', die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen,

R₆ oder R₇, die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen können, oder zusammen eine C=O-Gruppe bilden, oder einer der Reste R₆ oder R₇ zusammen mit R₅ einen Ring mit 5 oder 6 Kohlenstoffatomen bildet.

2. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** X₁ eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen ist, die gegebenenfalls mit einer oder mehreren Alkylgruppen mit 1 bis 3 Kohlenstoffatomen oder einer oder mehreren Alkoxygruppen mit 1 oder 2 Kohlenstoffatomen substituiert ist.

3. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** X₁ unter einer CH₂-Gruppe, CH₂CH₂-Gruppe oder einer C(CH₃)₂-Gruppe ausgewählt ist, und R₉ und R₁₀, die gleich oder verschieden sein können, für Wasserstoff oder eine CH₃-Gruppe stehen.

4. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** der Rest A unter einer CH₂-Gruppe, einer CH₂CH₂-Gruppe, einer C(CH₃)₂-Gruppe, einer CH₂CH₂OCH₂-Gruppe, einer CH₂CH₂OC(CH₃)₂-Gruppe und einer CH₂CH₂CH₂OC(CH₃)₂-Gruppe ausgewählt ist.

5. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₁₁ für ein Wasserstoffatom, eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen, eine Alkoxygruppe mit 1 oder 2 Kohlenstoffatomen steht, oder R₁₁ zusammen mit R₁₂ einen 5- oder 6-gliedrigen Heterocyclus bildet.

6. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₁₂ für ein Wasserstoffatom, eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen, eine Alkoxygruppe mit 1 oder 2 Kohlenstoffatomen, CON(R')(R'') steht, wobei R' und R'', die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen stehen oder R₁₂ zusammen mit R₁₁ einen 5- oder 6-gliedrigen Heterocyclus bildet.

7. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₁₃ und R₁₅, die gleich oder verschieden sein können, für ein Wasserstoffatom, eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 oder 2 Kohlenstoffatomen stehen.

8. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₁₄ für ein Wasserstoffatom, eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen, COR', COOR', CON(R')(R'') steht, wobei R' und R'', die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen stehen.

9. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** X₂ für eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen steht, die gegebenenfalls mit einer oder mehreren Alkylgruppen mit 1 bis 3 Kohlenstoffatomen oder einer oder mehreren Alkoxygruppen mit 1 oder 2 Kohlenstoffatomen substituiert ist.

10. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** X₃ unter OH, NH₂, NHOH und NHNH₂ ausgewählt ist.

11. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₁ und R₂, die gleich oder verschieden sein können, für ein Wasserstoffatom, eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen oder einer Alkoxygruppe mit 1 oder 2 Kohlenstoffatomen stehen.

12. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₃, R₄ und R₈, die gleich oder verschieden sein können, unter einem Wasserstoffatom, einer Alkylgruppe mit 1 bis 3 Kohlenstoffatomen, einer Alkoxygruppe mit 1 oder 2 Kohlenstoffatomen, einem Br-, Cl- oder F-Atom, einer OH-Gruppe, einer Nitrogruppe, einer Trifluormethyl-Gruppe oder einer Gruppe N(R')(R'') oder -N(R')COR'', -CN, -CONR'R'', -SO₂NR'R'', -SO₂R' ausgewählt sind, wobei R' und R'', die gleich oder verschieden sein können, für ein Wasserstoffatom und eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen stehen.

13. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R₅ unter einem Wasserstoffatom, einer Alkylgruppe

mit 1 bis 3 Kohlenstoffatomen, einer Alkoxygruppe mit 1 oder 2 Kohlenstoffatomen, einem Halogenatom, einer OH-Gruppe ausgewählt ist, oder R_5 zusammen mit einem der Reste R_6 oder R_7 einen Ring mit 5 oder 6 Kohlenstoffatomen bildet.

14. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, dass** R_6 und R_7 , die gleich oder verschieden sein können, unter einem Wasserstoffatom, einer Alkylgruppe mit 1 bis 3 Kohlenstoffatomen ausgewählt sind, oder zusammen eine Gruppe $C=O$ bilden, oder einer der Reste R_6 oder R_7 zusammen mit R_5 einen Ring mit 5 oder 6 Kohlenstoffatomen bildet.

15. Pharmazeutische Zusammensetzung umfassend die Verbindung nach Formel (I) nach einem der vorhergehenden Ansprüche oder ein pharmazeutisch annehmbares Salz oder einen Ester davon und mindestens ein pharmazeutisch annehmbares Vehikel.

16. Pharmazeutische Zusammensetzung nach Anspruch 15, **dadurch gekennzeichnet, dass** das pharmazeutische annehmbare Salz ein Additionssalz mit physiologisch annehmbaren organischen oder mineralischen Säuren oder Basen ist.

17. Pharmazeutische Zusammensetzung nach Anspruch 16, **dadurch gekennzeichnet, dass** die physiologisch annehmbaren Säuren unter Chlorwasserstoffsäure, Bromwasserstoffsäure, Schwefelsäure, Phosphorsäure, Salpetersäure, Essigsäure, Ascorbinsäure, Benzoesäure, Zitronensäure, Fumarsäure, Milchsäure, Maleinsäure, Methansulfonsäure, Oxalsäure, Paratoluolsulfonsäure, Benzolsulfonsäure, Bernsteinsäure, Gerbsäure und Weinsäure ausgewählt sind.


18. Pharmazeutische Zusammensetzung nach Anspruch 16, **dadurch gekennzeichnet, dass** die physiologisch annehmbaren Basen unter Ammoniumhydroxid, Calciumhydroxid, Magnesiumcarbonat, Natriumhydrogencarbonat, Kaliumhydrogencarbonat, Arginin, Betain, Koffein, Cholin, N,N-Dibenzylethyldiamin, Diethylamin, 2-Diethylaminoethanol, 2-Dimethylaminoethanol, Ethanolamin, Ethylendiamin, N-Ethylmorpholin, N-Ethylpiperidin, N-Methylglucamin, Glucamin, Glucosamin, Histidin, N-(2-Hydroxyethyl)piperidin, N-(2-Hydroxyethyl)pyrrolidin, Isopropylamin, Lysin, Methylglucamin, Morpholin, Piperazin, Piperidin, Theobromin, Triethylamin, Trimethylamin, Tripropylamin und Tromethamin ausgewählt sind.

19. Pharmazeutische Zusammensetzung nach Anspruch 15, **dadurch gekennzeichnet, dass** der pharmazeutisch annehmbare Ester mit physiologisch annehmbaren organischen Säuren oder Alkoholen gebildet wird.

20. Pharmazeutische Zusammensetzung nach einem der Ansprüche 15 bis 19, **dadurch gekennzeichnet, dass** die Zusammensetzung ein Stereoisomer oder ein Enantiomer der Verbindung nach Formel (I) enthält oder ein pharmazeutisch annehmbares Salz oder einen Ester davon oder eine Mischung davon.

21. Pharmazeutische Zusammensetzung nach einem der Ansprüche 15 bis 20, **dadurch gekennzeichnet, dass** das pharmazeutisch annehmbare Vehikel unter Gleitmitteln, Bindemitteln, Sprengmitteln, Füllstoffen, Verdünnungsmitteln, Geschmacksstoffen, Farbstoffen, Verflüssigern, Schmiermitteln, Konservierungsmitteln, Feuchthaltemitteln, Absorptionsmitteln und Süßungsmitteln ausgewählt ist.

22. Verwendung einer Verbindung der Formel (I)



worin

A für $-X_1-$ oder $-X_1-OC(R_9)(R_{10})-$ stehen kann, wobei

X_1 für eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen kann, die gegebenenfalls mit einer oder mehreren Alkylgruppen mit 1 bis 5 Kohlenstoffatomen oder mit einer oder mehreren Alkoxygruppen mit 1 bis 3 Kohlenstoffatomen substituiert sein kann,
und

R_9 und R_{10} , die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen kann,

Y für $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$ stehen kann, wobei R_{11} für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen kann oder R_{11} zusammen mit R_{12} einen 4- bis 7-gliedrigen Heterocyclus bildet,

R_{12} für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, COR' , $COOR'$, $CON(R')(R'')$ stehen kann, wobei R' und R'' , die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen, oder R_{12} zusammen mit R_{11} einen 4- bis 7-gliedrigen Heterocyclus bildet,

R_{13} und R_{15} , die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen können,

R_{14} für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, COR' , $COOR'$, $CON(R')(R'')$ stehen kann, wobei R' und R'' , die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen,

X_2 für eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen kann, die gegebenenfalls mit einer oder mehreren Alkylgruppen mit 1 bis 5 Kohlenstoffatomen oder mit einer oder mehreren Alkoxygruppen mit 1 bis 3 Kohlenstoffatomen substituiert sein kann,

X_3 für OH , NH_2 , $NHOH$ oder $NHNH_2$ stehen kann,

R_1 und R_2 , die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen oder eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen stehen kann,

R_3 , R_4 und R_8 , die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, ein Halogenatom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, $-CN$, $-CONR''R''$, $-SO_2NR''R''$, $-SO_2R'$, Nitro und Trifluormethyl stehen können; wobei R' und R'' , die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen,

R_5 für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, ein Halogenatom, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, Nitro und Trifluormethyl stehen kann, oder R_5 zusammen mit einem der Reste R_6 oder R_7 einen Ring mit 5 oder 6 Kohlenstoffatomen bildet; wobei R' und R'' , die gleich oder verschieden sein können, für Wasserstoff und eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen,

R_6 oder R_7 , die gleich oder verschieden sein können, für Wasserstoff, eine Alkylgruppe mit 1 bis 5 Kohlenstoffatomen stehen können, oder zusammen eine $C=O$ -Gruppe bilden, oder einer der Reste R_6 oder R_7 zusammen mit R_5 einen Ring mit 5 oder 6 Kohlenstoffatomen bildet,

zur Herstellung einer pharmazeutischen Zusammensetzung zur Behandlung von Krankheiten, die auf der Expression von MCP-1, CX3CR1 und p40 basieren, wobei die Krankheiten, die auf der Expression von MCP-1 und CX3CR1 basieren, unter Gelenkserkrankungen, Nierenerkrankungen, kardiovaskulären Erkrankungen, metabolischem Syndrom, Fettleibigkeit, Diabetes, Insulinresistenz und Krebs ausgewählt sind, und die auf der Expression von p40 basierenden Erkrankungen unter Autoimmunerkrankungen, chronisch degenerativen Entzündungskrankheiten und Krebs ausgewählt sind.


23. Verwendung nach Anspruch 22, **dadurch gekennzeichnet, dass** die Krankheiten, die auf der Expression von MCP-1 basieren, unter rheumatoider Arthritis, Arthritis, die durch virale Infektionen induziert ist, Psoriasis Arthritis, Arthrose, Lupusnephritis, diabetischer Nephropathie, Glomerulonephritis, polyzystischer Nierenerkrankung, interstitieller Lungenerkrankung, Fibrose, Multipler Sklerose, Alzheimerscher Krankheit, HIVassoziierter Demenz, atopischer Dermatitis, Psoriasis, Vaskulitis, Restenose, Arteriosklerose, Herzinfarkt, Angina, akuter Herzgefäßerkrankung, Adenomen, Karzinomen und Metastasen und metabolischen Krankheiten ausgewählt sind.

24. Verwendung nach Anspruch 22, **dadurch gekennzeichnet, dass** die Erkrankungen, die auf der Expression von CX3CR1 basieren, unter rheumatoider Arthritis, Lupusnephritis, diabetischer Nephropathie, Morbus Crohn, Colitis ulcerosa, koronaren Erkrankungen, Restenose, Arteriosklerose, Herzinfarkt und Angina ausgewählt sind.

25. Verwendung nach Anspruch 22, **dadurch gekennzeichnet, dass** die Erkrankungen, die auf der Expression von p40 basieren, unter rheumatoider Arthritis, Psoriasis, Glomerulonephritis, Diabetes, Lupus erythematoses, Diabetes, Morbus Crohn und Tumoren ausgewählt sind.

Revendications

1. Composé de formule (I)



dans laquelle :

A peut être $-X_1-$ ou $-X_1-OC(R_9)(R_{10})-$, formules dans lesquelles

X_1 peut représenter un groupe alkyle comportant de 1 à 5 atomes de carbone, facultativement substitué par un ou plusieurs groupes alkyles comportant de 1 à 5 atomes de carbone ou un ou plusieurs groupes alcoxy comportant de 1 à 3 atomes de carbone, et

R_9 et R_{10} , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone ou un groupe alcoxy comportant de 1 à 3 atomes de carbone,

Y peut être $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$, formules dans lesquelles

R_{11} peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, ou R_{11} conjointement avec R_{12} forme un hétérocycle de 4 à 7 chaînons,

R_{12} peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, COR' , $COOR'$, $CON(R')(R'')$ avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone, ou R_{12} conjointement avec R_{11} , forme un hétérocycle de 4 à 7 chaînons,

R_{13} et R_{15} , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone ou un groupe alcoxy comportant de 1 à 3 atomes de carbone,

R_{14} peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, COR' , $COOR'$, $CON(R')(R'')$ avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone,

X_2 peut représenter un groupe alkyle comportant de 1 à 5 atomes de carbone, facultativement substitué par un ou plusieurs groupes alkyles comportant de 1 à 5 atomes de carbone ou un ou plusieurs groupes alcoxy comportant de 1 à 3 atomes de carbone,

X_3 peut être un groupe OH, NH_2 , $NHOH$ ou $NHNH_2$,

R_1 et R_2 , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone ou un groupe alcoxy comportant de 1 à 3 atomes de carbone, R_3 , R_4 et R_8 , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, un atome d'halogène, un groupe $-OH$, $-N(R')(R'')$, $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro et trifluorométhyle ; avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone,

R_5 peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, un atome d'halogène, un groupe $-OH$, $-N(R')(R'')$, $-N(R')COR''$, nitro

et trifluorométhyle, ou R_5 conjointement avec l'un parmi R_6 et R_7 forme un noyau comportant 5 ou 6 atomes de carbone ; avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome

d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone,

R_6 et R_7 , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, ou forment conjointement un groupe $C=O$, ou l'un parmi R_6 et R_7 , conjointement avec R_5 , forme un noyau comportant 5 ou 6 atomes de carbone.

2. Composé selon la revendication 1, **caractérisé en ce que** X_1 représente un groupe alkyle comportant de 1 à 3 atomes de carbone, facultativement substitué par un ou plusieurs groupes alkyles comportant de 1 à 3 atomes de carbone ou un ou plusieurs groupes alcoxy comportant 1 ou 2 atomes de carbone.

3. Composé selon la revendication 1, **caractérisé en ce que** X_1 est choisi dans le groupe comprenant un groupe CH_2 , un groupe CH_2CH_2 ou un groupe $C(CH_3)_2$, et R_9 et R_{10} , qui peuvent être identiques ou différents l'un de l'autre, sont un atome d'hydrogène ou un groupe CH_3 .

4. Composé selon la revendication 1, **caractérisé en ce que** le résidu A est choisi dans le groupe comprenant un groupe CH_2 , un groupe CH_2CH_2 , un groupe $C(CH_3)_2$, un groupe $CH_2CH_2OCH_2$, un groupe $CH_2CH_2OC(CH_3)_2$ et un groupe $CH_2CH_2CH_2OC(CH_3)_2$.

5. Composé selon la revendication 1, **caractérisé en ce que** R_{11} est un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone, un groupe alcoxy comportant 1 ou 2 atomes de carbone, ou R_{11} conjointement avec R_{12} forme un hétérocycle de 5 ou 6 chaînons.

6. Composé selon la revendication 1, **caractérisé en ce que** R_{12} est un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone, un groupe alcoxy comportant 1 ou 2 atomes de carbone, $CON(R')(R'')$ avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 3 atomes de carbone, ou R_{12} conjointement avec R_{11} forme un hétérocycle de 5 ou 6 chaînons.

7. Composé selon la revendication 1, **caractérisé en ce que** R_{13} et R_{15} , qui peuvent être identiques ou différents l'un de l'autre, sont un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone ou un groupe alcoxy comportant 1 ou 2 atomes de carbone.

8. Composé selon la revendication 1, **caractérisé en ce que** R_{14} est un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone, COR' , $COOR'$, $CON(R')(R'')$ avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 3 atomes de carbone.

9. Composé selon la revendication 1, **caractérisé en ce que** X_2 représente un groupe alkyle comportant de 1 à 3 atomes de carbone, facultativement substitué par un ou plusieurs groupes alkyles comportant de 1 à 3 atomes de carbone ou un ou plusieurs groupes alcoxy comportant 1 ou 2 atomes de carbone.

10. Composé selon la revendication 1, **caractérisé en ce que** X_3 est choisi dans le groupe comprenant OH , NH_2 , $NHOH$ et $NHNH_2$.

11. Composé selon la revendication 1, **caractérisé en ce que** R_1 et R_2 , qui peuvent être identiques ou différents l'un de l'autre, sont un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone ou un groupe alcoxy comportant 1 ou 2 atomes de carbone.

12. Composé selon la revendication 1, **caractérisé en ce que** R_3 , R_4 et R_8 , qui peuvent être identiques ou différents l'un de l'autre, sont choisis dans le groupe comprenant un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone, un groupe alcoxy comportant 1 ou 2 atomes de carbone, un atome de Br , Cl ou F , un groupe OH , un groupe nitro, un groupe trifluorométhyle ou un groupe $N(R')(R'')$ ou $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 3 atomes de carbone.

13. Composé selon la revendication 1, **caractérisé en ce que** R_5 est choisi dans le groupe comprenant un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone, un groupe alcoxy comportant 1 ou 2 atomes

de carbone, un atome d'halogène, un groupe OH, ou R₅, conjointement avec l'un parmi R₆ et R₇, forme un noyau comportant 5 ou 6 atomes de carbone.

5 14. Composé selon la revendication 1, **caractérisé en ce que** R₆ et R₇, qui peuvent être identiques ou différents l'un de l'autre, sont choisis dans le groupe comprenant un atome d'hydrogène, un groupe alkyle comportant de 1 à 3 atomes de carbone, ou forment conjointement un groupe C=O, ou l'un parmi R₆ et R₇, conjointement avec R₅, forme un noyau comportant 5 ou 6 atomes de carbone.

10 15. Composition pharmaceutique comprenant le composé de formule (I) selon l'une quelconque des revendications précédentes, ou un sel ou un ester pharmaceutiquement acceptable de celui-ci, et au moins un véhicule pharmaceutiquement acceptable.

15 16. Composition pharmaceutique selon la revendication 15, **caractérisée en ce que** ledit sel pharmaceutiquement acceptable est un sel d'addition avec des acides ou des bases organiques ou minéraux physiologiquement acceptables.

20 17. Composition pharmaceutique selon la revendication 16, **caractérisée en ce que** lesdits acides physiologiquement acceptables sont choisis dans le groupe comprenant l'acide chlorhydrique, l'acide bromhydrique, l'acide sulfurique, l'acide phosphorique, l'acide nitrique, l'acide acétique, l'acide ascorbique, l'acide benzoïque, l'acide citrique, l'acide fumarique, l'acide lactique, l'acide maléique, l'acide méthanesulfonique, l'acide oxalique, l'acide para-toluènesulfonique, l'acide benzènesulfonique, l'acide succinique, l'acide tannique et l'acide tartrique.


25 18. Composition pharmaceutique selon la revendication 16, **caractérisée en ce que** lesdites bases physiologiquement acceptable sont choisies dans le groupe comprenant l'hydroxyde d'ammonium, l'hydroxyde de calcium, le carbonate de magnésium, l'hydrogénocarbonate de sodium, l'hydrogénocarbonate de potassium, l'arginine, la bétaïne, la caféine, la choline, la N,N-dibenzyléthylènediamine, la diéthylamine, le 2-diéthylaminoéthanol, le 2-diméthylaminoéthanol, l'éthanolamine, l'éthylènediamine, la N-éthylmorpholine, la N-éthylpipéridine, la N-méthylglucamine, la glucamine, la glucosamine, l'histidine, la N-(2-hydroxyéthyl)pipéridine, la N-(2-hydroxyéthyl)pyrrolidine, l'isopropylamine, la lysine, la méthylglucamine, la morpholine, la pipérazine, la pipéridine, la théobromine, la triéthylamine, la triméthylamine, la tripropylamine et la trométhamine.

35 19. Composition pharmaceutique selon la revendication 15, **caractérisée en ce que** ledit ester pharmaceutiquement acceptable est formé avec des acides organiques ou des alcools physiologiquement acceptables.

40 20. Composition pharmaceutique selon l'une quelconque des revendications 15 à 19, **caractérisée en ce que** ladite composition contient un stéréoisomère ou un énantiomère du composé de formule (I), ou un sel ou un ester pharmaceutiquement acceptable de celui-ci, ou un mélange de ceux-ci.

45 21. Composition pharmaceutique selon l'une quelconque des revendications 15 à 20, **caractérisée en ce que** ledit véhicule pharmaceutiquement acceptable est choisi dans le groupe comprenant les glissants, les liants, les agents de désintégration, les charges, les diluants, les arômes, les colorants, les fluidifiants, les lubrifiants, les agents de conservation, les humectants, les absorbants et les édulcorants.

50 22. Utilisation d'un composé de formule (I)



dans laquelle :

A peut être $-X_1-$ ou $-X_1-OC(R_9)(R_{10})-$, formules dans lesquelles

X_1 peut représenter un groupe alkyle comportant de 1 à 5 atomes de carbone, facultativement substitué par un ou plusieurs groupes alkyles comportant de 1 à 5 atomes de carbone ou un ou plusieurs groupes alcoxy comportant de 1 à 3 atomes de carbone, et

R_9 et R_{10} , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone ou un groupe alcoxy comportant de 1 à 3 atomes de carbone,

Y peut être $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$, formules dans lesquelles

R_{11} peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, ou R_{11} conjointement avec R_{12} forme un hétérocycle de 4 à 7 chaînons,

R_{12} peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, COR' , $COOR'$, $CON(R')(R'')$ avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone, ou R_{12} conjointement avec R_{11} , forme un hétérocycle de 4 à 7 chaînons,

R_{13} et R_{15} , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone ou un groupe alcoxy comportant de 1 à 3 atomes de carbone,

R_{14} peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, COR' , $COOR'$, $CON(R')(R'')$ avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone,

X_2 peut représenter un groupe alkyle comportant de 1 à 5 atomes de carbone, facultativement substitué par un ou plusieurs groupes alkyles comportant de 1 à 5 atomes de carbone ou un ou plusieurs groupes alcoxy comportant de 1 à 3 atomes de carbone,

X_3 peut être OH, NH_2 , $NHOH$ ou $NHNH_2$,

R_1 et R_2 , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone ou un groupe alcoxy comportant de 1 à 3 atomes de carbone, R_3 , R_4 et R_8 , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, un atome d'halogène, un groupe -OH, $-N(R')(R'')$, $-N(R')COR''$, -CN, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro et trifluorométhyle ;

avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone,

R_5 peut être un atome d'hydrogène, un groupe alkyle comportant de 1 à 5 atomes de carbone, un groupe alcoxy comportant de 1 à 3 atomes de carbone, un atome d'halogène, un groupe -OH, $-N(R')(R'')$, $-N(R')COR''$, nitro et trifluorométhyle, ou R_5 conjointement avec l'un parmi R_6 et R_7 forme un noyau comportant 5 ou 6 atomes de carbone ; avec R' et R'' , qui peuvent être identiques ou différents l'un de l'autre, représentés par un atome d'hydrogène et un groupe alkyle comportant de 1 à 5 atomes de carbone,

R_8 et R_7 , qui peuvent être identiques ou différents l'un de l'autre, peuvent être un atome d'hydrogène, un groupe

alkyle comportant de 1 à 5 atomes de carbone, ou forment conjointement un groupe C=O, ou l'un parmi R₆ et R₇, conjointement avec R₅, forme un noyau comportant 5 ou 6 atomes de carbone, pour la préparation d'une composition pharmaceutique pour le traitement de maladies basées sur l'expression de MCP-1, CX3CR1 et p40, dans laquelle lesdites maladies basées sur l'expression de MCP-1 et CX3CR1 sont choisies dans le groupe comprenant les maladies articulaires, les maladies rénales, les maladies cardiovasculaires, le syndrome métabolique, l'obésité, le diabète, l'insulinorésistance et le cancer, et lesdites maladies basées sur l'expression de p40 sont choisies dans le groupe comprenant les maladies auto-immunes, les maladies inflammatoires dégénératives chroniques et le cancer.

23. Utilisation selon la revendication 22, **caractérisée en ce que** lesdites maladies basées sur l'expression de MCP-1 sont choisies dans le groupe comprenant la polyarthrite rhumatoïde, l'arthrite induite par des infections virales, le rhumatisme psoriasique, l'arthrose, la néphropathie lupique, la néphropathie diabétique, la glomérulonéphrite, la maladie kystique des reins, la maladie pulmonaire interstitielle, la fibrose, la sclérose en plaques, la maladie d'Alzheimer, la démence associée au VIH, la dermatite atopique, le psoriasis, la vasculite, la resténose, l'athérosclérose, l'infarctus du myocarde, l'angine de poitrine, les maladies coronaires aiguës, les adénomes, les carcinomes et les métastases et les maladies métaboliques.
24. Utilisation selon la revendication 22, **caractérisée en ce que** lesdites maladies basées sur l'expression de CX3CR1 sont choisies dans le groupe comprenant la polyarthrite rhumatoïde, la néphropathie lupique, la néphropathie diabétique, la maladie de Crohn, la recto-colite hémorragique, les troubles coronariens, la resténose, l'athérosclérose, l'infarctus du myocarde et l'angine de poitrine.
25. Utilisation selon la revendication 22, **caractérisée en ce que** lesdites maladies basées sur l'expression de p40 sont choisies dans le groupe comprenant la polyarthrite rhumatoïde, le psoriasis, la glomérulonéphrite, le diabète, le lupus érythémateux, le diabète, la maladie de Crohn et les tumeurs.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0382276 B [0028]
- EP 0510748 B [0028]
- EP 1005332 B [0028]
- EP 1185528 A [0029]
- EP 1188438 A [0029]
- EP 1199074 A [0029]
- EP 1369119 A [0029]
- EP 1458687 A [0029]
- EP 1819341 A [0029]
- EP 1827447 A [0029]
- EP 1869055 A [0030]
- EP 1869056 A [0030]
- EP 1675862 A [0030]
- EP 382276 A [0103] [0115]
- EP 0382276 A [0157] [0168] [0173] [0177]


Non-patent literature cited in the description

- **ROLLINS B.J.** *Chemokines, Blood*, 1997, vol. 90, 909-928 [0003]
- **M. BAGGIOLINI.** Chemokines and leukocyte traffic. *Nature*, 1998, vol. 392, 565-568 [0003]
- **GERARD C. ; ROLLINS B.J.** Chemokines and disease. *Nature Immunol.*, 2001, vol. 2, 108-115 [0009]
- **MAHALINGAM S. et al.** Chemokines and viruses: friend or foes. *Trends in Microbiology*, 2003, vol. 11, 383-391 [0013]
- **RULLI N. et al.** *Ross River Virus: molecular and cellular aspects of disease pathogenesis*, 2005, vol. 107, 329-342 [0013]
- **NANKI T. et al.** Migration of CX3CR1-positive T cells producing type 1 cytokines and cytotoxic molecules into the synovium of patients with rheumatoid arthritis. *Arthritis & Rheumatism*, 2002, vol. 46 (11), 2878-2883 [0015]
- **SEGERER S. et al.** Expression of the fractalkine receptor (CX3CR1) in human kidney diseases. *Kidney International*, 2002, vol. 62, 488-495 [0016]
- **SANS M. et al.** Enhanced recruitment of CX3CR1 + T cells by mucosal endothelial cell-derived fractalkine in inflammatory bowel diseases. *Gastroenterology*, 2007, vol. 132 (1), 139-153 [0017]
- **LIU P. et al.** Cross-talk among Smad, MAPK and integrin signalling pathways enhances adventitial fibroblast functions activated by transforming growth factor-1 and inhibited by Gax. *Arterioscler. Thromb. Vasc. Biol.*, 2008 [0018]
- **MCDERMOTT D.H. et al.** Chemokine receptor mutant CX3CR1-M280 has impaired adhesive function and correlates with protection from cardiovascular diseases in humans. *J. Clin. Invest.*, 2003 [0018]
- **NIESSNER A. et al.** *Thrombosis and Haemostasis*, 2005 [0018]
- **KOO. J.** *Dermatol. Clin.*, 1996, vol. 14, 485-96 [0023]
- **SCHON M.P. et al.** *N. Engl. J. Med.*, 2005, vol. 352, 1899-912 [0023]
- **SIRONI M. et al.** A small synthetic molecule capable of preferentially inhibiting the production of the CC chemokine monocyte chemoattractant protein-1. *European Cytokine Network.*, September 1999, vol. 10 (3), 437-41 [0028]

1-Benzil-3-hidroxi-metilindazol-származékok és azok alkalmazása MCP-1, CX3CR1
expresszió alapú betegségek kezelésében

SZABADALMI IGÉNYPONTOK

1. Az (I) általános képletű vegyület



ahol:

A jelentése $-X_1-$ vagy $-X_1-OC(R_9)(R_{10})-$ általános képletű csoport, ahol

X_1 jelentése 1-5 szénatomos alkilcsoport, mely adott esetben egy vagy több, 1-5 szénatomos alkilcsoporttal vagy egy vagy több, 1-3 szénatomos alkoxycsoporttal lehet szubsztituálva, és

R_9 and R_{10} jelentése azonos vagy különböző, éspedig hidrogénatom, 1-5 szénatomos alkilcsoport vagy 1-3 szénatomos alkoxycsoport,

Y jelentése $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$ általános képletű csoport, ahol

R_{11} hidrogénatomot, 1-5 szénatomos alkilcsoportot, 1-3 szénatomos alkoxycsoportot képvisel, vagy R_{11} és R_{12} együtt egy 4-7-tagú heterociklusos csoportot jelent,

R_{12} hidrogénatom, 1-5 szénatomos alkilcsoport, 1-3 szénatomos alkoxycsoport, COR' , $COOR'$, $CON(R')(R'')$ általános képletű csoport lehet, ahol R' és R'' , melyek jelentése azonos vagy különböző, hidrogénatomot és 1-5 szénatomos alkilcsoportot képviselnek, vagy R_{12} és R_{11} együtt 4-7-tagú heterociklusos csoportot képez,

R_{13} és R_{15} , melyek lehetnek azonosak vagy különbözőek, hidrogénatomot, 1-5 szénatomos alkilcsoportot vagy 1-3 szénatomos alkoxycsoportot jelentenek,

R_{14} jelentése hidrogénatom, 1-5 szénatomos alkilcsoport, COR' , $COOR'$, $CON(R')(R'')$ általános képletű csoport, ahol

R' and R'' , melyek lehetnek azonosak vagy különbözőek, hidrogénatomot és 1-5 szénatomos alkilcsoportot jelenthetnek,

X_2 jelentése 1-5 szénatomos alkilcsoport, mely adott esetben egy vagy több, 1-5 szénatomos alkilcsoporttal vagy egy vagy több, 1-3 szénatomos alkoxicsoporthal lehet helyettesítve,

X_3 jelentése OH , NH_2 , $NHOH$ vagy $NHNH_2$ csoport,

R_1 and R_2 , melyek lehetnek azonos vagy különböző jelentésűek, hidrogénatomot, 1-5 szénatomos alkilcsoportot vagy 1-3 szénatomos alkoxicsoporthat képviselnek,

R_3 , R_4 and R_8 , melyek lehetnek azonos vagy különböző jelentésűek, hidrogénatomot, 1-5 szénatomos alkilcsoportot, 1-3 szénatomos alkoxicsoporthat, halogénatomot, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, $-CN$, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro- és trifluorometilcsoportot jelenthetnek; az általános képletű csoportokban R' és R'' jelentése lehet azonos vagy különböző, és pedig hidrogénatom és 1-5 szénatomos alkilcsoport,

R_5 jelentése hidrogénatom, 1-5 szénatomos alkilcsoport, 1-3 szénatomos alkoxicsoporthat, halogénatomot, $-OH$, $-N(R')(R'')$, $-N(R')COR''$, nitro- és trifluorometilcsoportot, vagy R_5 az R_6 és R_7 csoportok egyikével együtt 5-6 szénatomos gyűrűt képez; ahol az általános képletű csoportokban R' és R'' , melyek jelentése lehet azonos vagy különböző, hidrogénatomot és 1-5 szénatomos alkilcsoportot képviselhet,

R_6 és R_7 , melyek lehetnek azonos vagy különböző jelentésűek, hidrogénatomot, 1-5 szénatomos alkilcsoportot képviselhetnek, vagy együtt egy $C=O$ -csoportot képviselhetnek, vagy R_6 és R_7 egyike és R_5 együttesen, egy 5-6 szénatomos gyűrűt képez.

2. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy X_1 jelentése 1-3 szénatomos alkilcsoport, mely adott esetben egy vagy több, 1-3 szénatomos alkilcsoporttal, vagy egy vagy több, 1 vagy 2 szénatomos alkoxicsoporthal lehet helyettesítve.

3. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy X_1 a CH_2 -csoport, CH_2CH_2 -csoport vagy $C(CH_3)_2$ -csoport közül van kiválasztva, és R_9 és R_{10} , melyek lehetnek azonosak vagy különbözőek, hidrogénatomot vagy CH_3 -csoportot jelentenek.

4. Az 1. igényponi szerinti vegyület, azzal jellemezve, hogy az A csoport a CH_2 -csoport, CH_2CH_2 -csoport, $C(CH_3)_2$ -csoport, $CH_2CH_2OCH_2$ -csoport, $CH_2CH_2OC(CH_3)_2$ -csoport, és a $CH_2CH_2CH_2OC(CH_3)_2$ -csoport közül van kiválasztva.

5. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_{11} jelentése hidrogénatom, 1-3 szénatomos alkilcsoport, 1 vagy 2 szénatomos alkoxics csoport, vagy R_{11} és R_{12} együtt egy 5- vagy 6-tagú heterociklusos csoportot képez.

6. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_{12} jelentése hidrogénatom, 1-3 szénatomos alkilcsoport, 1 vagy 2 szénatomos alkoxics csoport, $\text{CON}(R')(R'')$ általános képletű csoport, ahol R' és R'' , melyek jelentése lehet azonos vagy különböző, hidrogénatomot, és 1-3 szénatomos alkilcsoportot jelenthetnek, vagy R_{12} és R_{11} együtt egy 5- vagy 6-tagú heterociklusos csoportot képvisel.

7. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_{13} és R_{15} , melyek jelentése lehet azonos vagy különböző, hidrogénatomot, 1-3 szénatomos alkilcsoportot, vagy 1 vagy 2 szénatomos alkoxics csoportot képviselnek.

8. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_{14} jelentése hidrogénatom, 1-3 szénatomos alkilcsoport, COR' , COOR' , $\text{CON}(R')(R'')$ általános képletű csoport, ahol az általános képletekben R' és R'' , melyek azonosak vagy különbözőek lehetnek, hidrogénatomot és 1-3 szénatomos alkilcsoportot képviselhetnek.

9. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy X_2 jelentése 1-3 szénatomos alkilcsoport, mely adott esetben egy vagy több, 1-3 szénatomos alkilcsoporttal vagy egy vagy több, 1 vagy 2 szénatomos alkoxics csoporttal lehet helyettesítve.

10. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy X_3 az OH, NH_2 , NHOH és NHNH_2 csoportok közül van kiválasztva.

11. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_1 és R_2 , melyek jelentése lehet azonos vagy különböző, hidrogénatomot, 1-3 szénatomos alkilcsoportot, vagy 1 vagy 2 szénatomos alkoxics csoportot jelentenek.

12. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_3 , R_4 és R_8 , melyek jelentése azonos vagy különböző, a hidrogénatom, 1-3 szénatomos alkil-, 1 vagy 2 szénatomos alkoxics csoport, Br-, Cl- vagy F-atom, OH-csoport, nitrocsoport, trifluorometilcsoport, vagy $\text{N}(R')(R'')$ vagy $-\text{N}(R')\text{COR}''$, $-\text{CN}$, $-\text{CONR}'R''$, $-\text{SO}_2\text{NR}'R''$, $-\text{SO}_2R'$ általános képletű csoport, ahol R' és R'' jelentése azonos vagy különböző, és hidrogénatom vagy 1-3 szénatomos alkilcsoport lehet, csoportból van kiválasztva.

13. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_5 a hidrogénatom, 1-3 szénatomos alkilcsoport, 1 vagy 2 szénatomos alkoxics csoport, halogénatom, OH-csoport közül van kiválasztva, vagy R_5 az R_6 és R_7 -csoport valamelyikével egy 5 vagy 6 szénatomos gyűrűt képez.

14. Az 1. igénypont szerinti vegyület, azzal jellemezve, hogy R_6 és R_7 , melyek jelentése azonos vagy különböző, a hidrogénatom, 1-3 szénatomos alkilcsoport közül van kiválasztva, vagy együttesen $C=O$ -csoportot képeznek, vagy R_6 és R_7 -csoport közül az egyik, és R_5 együttesen egy 5 vagy 6 szénatomos gyűrűt képez.

15. Gyógyszerkészítmény, mely az előző igénypontok bármelyike szerinti (I) általános képletű vegyületet, vagy gyógyászatilag elfogadható sóját vagy annak észterét, és legalább egy gyógyászatilag elfogadható vivőanyagot tartalmaz.

16. A 15. igénypont szerinti gyógyszerkészítmény, azzal jellemezve, hogy az említett gyógyászatilag elfogadható só fiziológiailag elfogadható szerves vagy ásványi savakkal vagy bázisokkal képezett addíciós só.

17. A 16. igénypont szerinti gyógyszerkészítmény, azzal jellemezve, hogy az említett fiziológiailag elfogadható savak a sósav, hidrogénbromid, kénsav, foszforsav, salétromsav, ecetsav, aszkorbinsav, benzoészav, citromsav, fumársav, tejsav, maleinsav, metánszulfonsav, oxálsav, para-toluolszulfonsav, benzolszulfonsav, borostyánkősav, csersav és borkősav csoportból vannak kiválasztva.

18. A 16. igénypont szerinti gyógyszerkészítmény, azzal jellemezve, hogy az említett fiziológiailag elfogadható bázis az ammóniumhidroxid, kalciumhidroxid, magnéziumkarbonát, nátrium-hidrogén-karbonát, kálium-hidrogén-karbonát, arginin, betain, koffein, kolin, N,N-dibenziletiléndiamin, dietilamin, 2-dietilaminoetanol, 2-dimetilaminoetanol, etanolamin, etilénediamin, N-etilmorfolin, N-etilpiperidin, N-metilglükamin, glükamin, glükózamin, hisztidin, N-(2-hidroxietyl)-piperidin, N-(2-hidroxietyl)-pirrolidin, izopropilamin, lizin, metilglükamin, mofolin, piperazin, piperidin, teobromin, trietilamin, trimetilamin, tripropilamin és trometamin közül van kiválasztva.


19. A 15. igénypont szerinti gyógyszerkészítmény, azzal jellemezve, hogy az említett gyógyászatilag elfogadható észter gyógyászatilag elfogadható szerves savakkal vagy alkoholokkal képezett észter.

20. A 15-19. igénypontok bármelyike szerinti gyógyszerkészítmény, azzal jellemezve, hogy az említett készítmény az (I) általános képletű vegyület sztereoizomerét vagy egy enantiomerjét vagy gyógyászatilag elfogadható sóját vagy észterét, vagy ezek keverékét tartalmazza.

21. A 15-20. igénypontok bármelyike szerinti gyógyszerkészítmény, azzal jellemezve, hogy az említett gyógyászatilag elfogadható vivőanyag a glidánsokat, kötőanyagokat, szét- és elősegítő anyagokat, töltőanyagokat, hígítókat, ízesítőanyagokat, színezőanyagokat,

fluidizálószerkeket, kenőanyagokat, konzerválószerkeket, nedvesítőszerkeket, abszorbenseket, és édesítőszerkeket tartalmazó csoportból vannak kiválasztva.

22. Az (I) általános képletű vegyület alkalmazása



ahol

A jelentése $-X_1-$ vagy $-X_1-OC(R_9)(R_{10})-$ általános képletű csoport, ahol

X_1 jelentése 1-5 szénatomos alkilcsoport, mely adott esetben egy vagy több, 1-5 szénatomos alkilcsoporttal vagy egy vagy több, 1-3 szénatomos alkoxycsoporttal lehet helyettesítve, és

R_9 és R_{10} , melyek lehetnek azonosak vagy különbözőek, hidrogénatomot, 1-5 szénatomos alkilcsoportot vagy 1-3 szénatomos alkoxycsoport jelentenek,

Y jelentése $N(R_{11})(R_{12})$, $N(R_{13})O(R_{14})$, $N(R_{13})N(R_{14})(R_{15})$, $N(R_{13})-X_2-N(R_{14})(R_{15})$, $N(R_{13})-X_2-CO-X_3$ általános képletű csoport, ahol

R_{11} hidrogénatomot, 1-5 szénatomos alkilcsoportot, 1-3 szénatomos alkoxycsoportot képviselhet, vagy R_{11} és R_{12} együtt 4-7-tagú heterociklusos csoportot képez,

R_{12} jelentése hidrogénatom, 1-5 szénatomos alkilcsoport, 1-3 szénatomos alkoxycsoport, COR' , $COOR'$, $CON(R')(R'')$ általános képletű csoport, ahol R' és R'' , melyek lehetnek azonosak vagy különbözőek, hidrogénatomot és 1-5 szénatomos alkilcsoportot képviselhetnek, vagy R_{12} és R_{11} együttesen egy 4-7-tagú heterociklusos csoportot képezhet,

R_{13} és R_{15} , melyek azonos vagy különböző jelentésűek, hidrogénatomot, 1-5 szénatomos alkilcsoportot, vagy 1-3 szénatomos alkoxycsoportot képviselnek,

R_{14} jelentése hidrogénatom, 1-5 szénatomos alkilcsoport, COR' , $COOR'$, $CON(R')(R'')$ általános képletű csoport, ahol

R' and R'' jelentése azonos vagy különböző, éspedig hidrogénatom és 1-5 szénatomos alkilcsoport,

X_2 jelentése 1-5 szénatomos alkilcsoport, mely adott esetben egy vagy több, 1-5 szénatomos alkilcsoporttal vagy egy vagy több, 1-3 szénatomos alkoxicssoporttal lehet helyettesítve,

X_3 jelentése OH, NH_2 , NHOH vagy $NHNH_2$ csoport,

R_1 és R_2 , melyek azonosak vagy különbözőek, hidrogénatomot, 1-5 szénatomos alkilcsoportot, vagy 1-3 szénatomos alkoxicssoportot képviselnek,

R_3 , R_4 és R_5 , melyek azonos vagy különböző jelentésűek, hidrogénatomot, 1-5 szénatomos alkilcsoportot, 1-3 szénatomos alkoxicssoportot, halogénatomot, -OH, $-N(R')(R'')$, $-N(R')COR''$, -CN, $-CONR'R''$, $-SO_2NR'R''$, $-SO_2R'$, nitro- és trifluormetilcsoportot jelentenek; R' és R'' , melyek jelentése azonos vagy különböző, hidrogénatomot és 1-5 szénatomos alkilcsoportot képviselnek,

R_6 hidrogénatom, 1-5 szénatomos alkilcsoport, 1-3 szénatomos alkoxicssoport, halogénatom, -OH, $-N(R')(R'')$, $-N(R')COR''$, nitro- és trifluormetilcsoportot képvisel, vagy R_5 és R_6 vagy R_7 egyike együttesen 5 vagy 6 szénatomos gyűrűt képvisel; ahol R' és R'' , melyek jelentése azonos vagy különböző, hidrogénatom vagy 1-5 szénatomos alkilcsoport lehet,

R_6 és R_7 , melyek azonosak vagy különbözőek, hidrogénatomot, 1-5 szénatomos alkilcsoportot jelentenek, vagy együttesen C=O-csoportot képeznek, vagy R_6 és R_7 egyike, és R_5 együttesen, egy 5- vagy 6-szénatomos gyűrűt képez,

olyan gyógyszerkészítmények előállítására,

melyek MCP-1, CX_3CR_1 és p40 expresszió alapú betegségek kezelésére szolgálnak, ahol az említett, MCP-1 és CX_3CR_1 expresszió alapú betegségek az ízületi betegségek, vesebetegségek, kardiovaszkuláris betegségek, metabolikus szindróma, elhízás, diabétesz, inzulinrezisztencia, és rák csoportból vannak kiválasztva, és az említett, p40 expresszió alapú betegségek az autoimmun betegségeket, krónikus degeneratív gyulladásos betegségeket, és rákot tartalmazó csoportból vannak kiválasztva.

23. A 22. igénypont szerinti alkalmazás, azzal jellemezve, hogy az említett, MCP-1 expresszió alapú betegségek a rheumatoid arthritis, vírusfertőzés által indukált arthritis, psoriasis arthritis, arthrosis, lupus nephritis, diabeteses nephropathia, glomerulonephritis, policisztás vesebetegség, interstitialis tüdőbetegség, fibrózis, sclerosis multiplex, Alzheimer-kór, HIV-vel társult demencia, atópiás dermatitis, psoriasis, vasculitis, restenosis, atherosclerosis, adenómák, karcinómák és metasztázisok, és metabolikus betegségek csoportból vannak kiválasztva.

24. A 22. igénypont szerinti alkalmazás, azzal jellemezve, hogy az említett, CX3CR1 expresszió alapú betegségek a rheumatoid arthritis, lupus nephritis, diabeteses nephropathia, Chron-betegség, fekélyes colitis, koronáriás betegségek, restenosis, atherosclerosis, miokardiális infarktus, és angina csoportból vannak kiválasztva.

25. A 22. igénypont szerinti alkalmazás, azzal jellemezve, hogy az említett, p40 expresszió alapú betegségek a rheumatoid arthritis, psoriasis, glomerulonephritis, diabetes, lupus erythematosus, diabetes, Chron-betegség és tumorok közül vannak kiválasztva.

A meghatalmazotti

~~Derzsi Katalin~~
~~szociális ügyvivő~~
SZOK Szociális Ügyvivő
H-1002 Budapest, Andrássy út 133
Telefon: 461-1000 Fax: 461-1091
Email: derzsi@szok.hu