

[19] Patents Registry
The Hong Kong Special Administrative Region
香港特別行政區
專利註冊處

[11] 1245801 B
EP 3230321 B1

[12] **STANDARD PATENT (R) SPECIFICATION**
轉錄標準專利說明書

[21] Application no. 申請編號 18104137.6	[51] Int. Cl. C07K 16/28 (2006.01) A61K 39/00 (2006.01) C07K 14/005 (2006.01) C07K 14/725 (2006.01) A61K 48/00 (2006.01) C12N 15/86 (2006.01) C12N 15/62 (2006.01) C12N 5/0783 (2010.01)
[22] Date of filing 提交日期 26.03.2018	

[54] BCMA CHIMERIC ANTIGEN RECEPTORS
BCMA 嵌合抗原受體

[30] Priority 優先權 12.12.2014 US US201462091419P 03.08.2015 US US201562200505P	[73] Proprietor 專利所有人 Bluebird Bio, Inc. 60 Binney Street Cambridge, MA 02142 UNITED STATES OF AMERICA
[43] Date of publication of application 申請發表日期 31.08.2018	[72] Inventor 發明人 MORGAN, Richard FRIEDMAN, Kevin
[45] Date of publication of grant of patent 批予專利的發表日期 15.05.2020	[74] Agent and / or address for service 代理人及/或送達地址 ROBIN BRIDGE & JOHN LIU 6th Floor, Emperor Commercial Centre, 39 Des Voeux Road Central, HONG KONG
[86] International application no. 國際申請編號 PCT/US2015/064269	
[87] International publication no. and date 國際申請發表編號及日期 WO2016/094304 16.06.2016	
EP Application no. & date 歐洲專利申請編號及日期 EP 15868392.0 07.12.2015	
EP Publication no. & date 歐洲專利申請發表編號及日期 EP 3230321 18.10.2017	
Date of grant in designated patent office 指定專利當局批予專利日期 28.08.2019	



(11) **EP 3 230 321 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
28.08.2019 Bulletin 2019/35

(21) Application number: **15868392.0**

(22) Date of filing: **07.12.2015**

(51) Int Cl.:
C07K 16/28 ^(2006.01) **A61K 39/00** ^(2006.01)
C07K 14/005 ^(2006.01) **C07K 14/725** ^(2006.01)
A61K 48/00 ^(2006.01) **C12N 15/86** ^(2006.01)
C12N 15/62 ^(2006.01) **C12N 5/0783** ^(2010.01)

(86) International application number:
PCT/US2015/064269

(87) International publication number:
WO 2016/094304 (16.06.2016 Gazette 2016/24)

(54) **BCMA CHIMERIC ANTIGEN RECEPTORS**
CHIMÄRE BCMA-ANTIGENREZEPTOREN
RÉCEPTEURS DE L'ANTIGÈNE CHIMÉRIQUE BCMA

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **12.12.2014 US 201462091419 P**
03.08.2015 US 201562200505 P

(43) Date of publication of application:
18.10.2017 Bulletin 2017/42

(73) Proprietor: **Bluebird Bio, Inc.**
Cambridge, MA 02142 (US)

(72) Inventors:
• **MORGAN, Richard**
Center Harbor, New Hampshire 03226-1254 (US)
• **FRIEDMAN, Kevin**
Medford, Massachusetts 02155 (US)

(74) Representative: **Cooley (UK) LLP**
Dashwood
69 Old Broad Street
London EC2M 1QS (GB)

(56) References cited:
WO-A1-2013/154760

- **WANG XIULI ET AL: "CS-1 Re-Directed Central Memory T Cell Therapy for Multiple Myeloma", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 124, no. 21, 9 December 2014 (2014-12-09), XP009189090, ISSN: 0006-4971**
- **R. Et Al. Uchibori, "400. CD269 (BCMA)-specific CAR-expressing T cells dramatically eradicate myeloma cells from bone marrow of an orthotopic multiple myeloma mouse model.", Molecular Therapy, vol. 24, abstract 400., 1 May 2016 (2016-05-01), XP055482537, DOI: [https://www.cell.com/molecular-therapy-family/molecular-therapy/fulltext/S1525-0016\(16\)33209-9](https://www.cell.com/molecular-therapy-family/molecular-therapy/fulltext/S1525-0016(16)33209-9) Retrieved from the Internet: URL:[https://www.cell.com/molecular-therapy-family/molecular-therapy/fulltext/S1525-0016\(16\)33209-9](https://www.cell.com/molecular-therapy-family/molecular-therapy/fulltext/S1525-0016(16)33209-9) [retrieved on 2018-06-08]**
- **CARPENTER ET AL.: 'B- cell maturation antigen is a promising target for adoptive T- cell therapy of multiple myeloma.' CLIN CANCER RES vol. 19, no. 8, pages 2048 - 2060, XP002727959**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**BACKGROUND****Technical Field**

[0001] Described herein are improved compositions and methods for treating B cell related conditions. More particularly, improved chimeric antigen receptors (CARs) comprising murine anti-BCMA antibodies or antigen binding fragments thereof, immune effector cells genetically modified to express these CARs, and use of these compositions to effectively treat B cell related conditions are disclosed.

Description of the Related Art

[0002] Several significant diseases involve B lymphocytes, *i.e.*, B cells. Abnormal B cell physiology can also lead to development of autoimmune diseases including, but not limited to systemic lupus erythematosus (SLE). Malignant transformation of B cells leads to cancers including, but not limited to lymphomas, *e.g.*, multiple myeloma and non-Hodgkins' lymphoma.

[0003] The large majority of patients having B cell malignancies, including non-Hodgkin's lymphoma (NHL) and multiple myeloma (MM), are significant contributors to cancer mortality. The response of B cell malignancies to various forms of treatment is mixed. Traditional methods of treating B cell malignancies, including chemotherapy and radiotherapy, have limited utility due to toxic side effects. Immunotherapy with anti-CD19, anti-CD20, anti-CD22, anti-CD23, anti-CD52, anti-CD80, and anti-HLA-DR therapeutic antibodies have provided limited success, due in part to poor pharmacokinetic profiles, rapid elimination of antibodies by serum proteases and filtration at the glomerulus, and limited penetration into the tumor site and expression levels of the target antigen on cancer cells. Attempts to use genetically modified cells expressing chimeric antigen receptors (CARs) have also met with limited success. In addition, the therapeutic efficacy of a given antigen binding domain used in a CAR is unpredictable: if the antigen binding domain binds too strongly, the CAR T cells induce massive cytokine release resulting in a potentially fatal immune reaction deemed a "cytokine storm," and if the antigen binding domain binds too weakly, the CAR T cells do not display sufficient therapeutic efficacy in clearing cancer cells.

[0004] WO 2013/154760 A1, Carpenter, R.O. et. al., Clin Cancer Res, vol. 19, no. 8, 2048-2060 (2013) and Wang, X, et. al., Blood, vol. 124, no. 21, abstract 1114 (2014) all disclose polynucleotides encoding anti-BCMA-CARs.

BRIEF SUMMARY

[0005] In a first aspect of the invention there is provided a polynucleotide encoding a CAR, wherein the polynucleotide sequence is set forth in SEQ ID NO: 10.

[0006] Preferred embodiments of the invention in any of its various aspects are as described below or as defined in the sub claims.

BRIEF DESCRIPTION OF SEVERAL VIEWS OF THE DRAWINGS**[0007]**

Figure 1 shows a schematic of murine B cell maturation antigen (muBCMA) CAR constructs.

Figure 2a shows the amount of IFN γ released from anti-BCMA02 CAR T cells, anti-BCMA10 CART cells, and CAR19 Δ T cells after the cells were co-cultured for 24 hours with K562 cells expressing BCMA.

Figure 2b shows the amount of IFN γ released from anti-BCMA02 CAR T cells, anti-BCMA10 CART cells, and CAR19 Δ T cells after the cells were co-cultured for 24 hours with K562 cells that lack BCMA expression compared to K562 cells expressing BCMA.

Figure 3 shows the amount of inflammatory cytokines in growth media from untransduced control T cells, anti-BCMA02 CAR T cells, anti-BCMAIO CART cells, and CAR19 Δ T cells, stimulated 10 days prior to the assay.

Figure 4 shows the amount of inflammatory cytokines produced by anti-BCMA02 CAR T cells, anti-BCMAIO CART cells, and CAR19 Δ T cells in the absence of antigen stimulation.

Figure 5 shows the expression of phenotypic markers of activation at the end of anti-BCMA CAR T cell manufacturing. HLA-DR and CD25 expression was measured in anti-BCMA02 CAR T cells, anti-BCMA10 CAR T cells, and CAR19Δ T cells.

Figure 6 shows the levels of activated caspase-3, a necessary step in apoptosis and important for AICD in anti-BCMA10 CAR T cells and anti-BCMA02 CAR T cells in the absence of antigen stimulation.

Figure 7 shows the amount of inflammatory cytokine release in anti-BCMA02 and anti-BCMA10 CAR T cells in media containing fetal bovine serum (FBS), human AB serum (HABS), or 100ng/ml soluble BCMA.

Figures 8A shows the tumor volume in NOD *scid* gamma (NSG) mice with ~100mm³ experimental sub-cutaneous human multiple myeloma (RPMI-8226) tumors. Mice were treated with vehicle, 10⁷ anti-BCMA02 CAR T cells, 10⁷ anti-BCMA10 CAR T cells, or Bortezomib (velcade).

Figures 8B shows the tumor volume in NOD *scid* gamma (NSG) mice with ~100mm³ experimental sub-cutaneous human multiple myeloma (RPMI-8226) tumors. Mice were treated with vehicle, 10⁷ anti-BCMA02 CAR T cells, 10⁷ anti-BCMA10 CAR T cells, or Bortezomib (velcade).

Figure 9 shows the level of BCMA expression on lymphoma and leukemia cell lines (circles) and the activity of anti-BCMA CAR T cells to each cell line (IFN γ release, boxes). BCMA-negative (BCMA-) tumor cell lines: myelogenous leukemia (K562), acute lymphoblastic leukemia (NALM-6 and NALM-16); Mantle cell lymphoma (REC-1); or Hodgkin's lymphoma (HDLM-2) showed little or no IFN γ release. BCMA-positive (BCMA+) tumor cell lines: B cell chronic lymphoblastic leukemia (MEC-1), Mantle cell lymphoma (JeKo-1), Hodgkin's lymphoma (RPMI-6666), Burkitt's lymphoma (Daudi cells and Ramos cells), and multiple myeloma (RPMI-8226) showed substantial IFN γ release.

Figure 10A shows the *in vivo* activity of vehicle, anti-CD19Δ CAR T cells, anti-CD 19 CAR T cells, and anti-BCMA CAR T cells to BCMA expressing Burkitt's lymphoma cells (Daudi cells) in an NSG mouse model when CAR T cells are administered to the mice at 8 days post tumor induction.

Figure 10B shows the *in vivo* activity of vehicle, anti-CD19Δ CAR T cells, anti-CD19 CAR T cells, and anti-BCMA CAR T cells to BCMA expressing Burkitt's lymphoma cells (Daudi cells) in an NSG mouse model when CAR T cells are administered to the mice at 18 days post tumor induction.

Figure 11 shows potent *in vitro* activity of anti-BCMA CAR T cells achieved with a 50 percent reduction anti-BCMA CAR expression. (A) T cell populations were transduced with between 4x10⁸ and 5x10⁷ transducing units of a lentivirus encoding an anti-BCMA CAR molecule (MOI of 5 to 40). The resulting T cell populations were normalized to contain 26 $\pm$ 4% anti-BCMA CAR-positive T cells. (B) MFI of the normalized anti-BCMA CAR T cells ranged from 885 to 1875 as assayed by flow cytometry. (C) K562 cells and K562-BCMA cells were co-cultured with normalized anti-BCMA CAR T cells at a 20:1 or 10:1 effector (E; T cell) to target (T; 1:1 mix of K562 and K562 BCMA cells) ratio showed comparable cytolytic activity.

Figure 12 shows the reliability of the manufacturing process for anti-BCMA CAR T cells. (A) anti-BCMA CAR T cell products manufactured from PBMCs of 11 individual donors show comparable levels of expansion compared to a matched culture of untransduced donor T cells. (B) anti-BCMA CAR T cell products manufactured from the 11 donors showed comparable lentiviral transduction efficiency (VCN). (C) The frequency of anti-BCMA CAR positive T cells was measured by flow cytometry and BCMA expression was found to be comparable across all donors. (D) anti-BCMA CAR T cell products manufactured from the 11 donors showed therapeutically relevant levels of IFN γ release when exposed to BCMA-expressing K562 cells.

Figure 13 shows Venn diagrams for co-expression of CD127, CD197 and CD38 in CD62L positive anti-BCMA02 T cells that have been cultured in the presence of IL-2 or IL-2 and ZSTK474 for ten days. ZSTK474-treated anti-BCMA02 CAR T cells showed an increase in the percentage of cells co-expressing CD127, CD197 and CD38 compared to anti-BCMA CAR T cells cultured with IL-2 alone.

Figure 14 shows an increased percentage of CD8 expressing anti-BCMA02 CAR T cells in cultures treated IL-2 and ZSTK474 (n=7) compared to cultures treated with IL-2 alone. CD8 expression was determined using a fluorescently-labeled anti-CD8 antibody and flow cytometry.

Figure 15 shows the amount of IFN- γ released by anti-BCMA02 CAR T cells from 14 donors after culture with IL-2 alone or with IL-2 and ZSTK474. At the end of the culture period, an equivalent number of anti-BCMA02 CAR T cells were re-cultured for 24 hours in media alone. The amount of IFN- γ released in 24 hours was quantified by ELISA. Culture in ZSTK474 did not significantly increase anti-BCMA02 CAR T cell tonic cytokine release compared to anti-BCMA02 CAR T cells cultured with IL-2 alone.

Figure 16 shows anti-tumor activity of anti-BCMA02 CAR T cells treated with IL-2, or IL-2 and ZSTK474, or a truncated signaling deficient anti-BCMA02 (tBCMA02) CAR T cell treated with IL-2 and ZSTK474 in an aggressive Daudi tumor model. Complete tumor regression was observed in 50% of mice administered the anti-BCMA02 CAR T cells treated with IL-2 and ZSTK474.

Figure 17 shows anti-tumor activity of anti-BCMA02 CAR T cells treated with IL-2, or IL-2 and ZSTK474 in a multiple myeloma tumor (RPMI-8226) model. Animals treated with IL-2- or IL-2 and ZSTK474-cultured anti-BCMA02 CAR T cells completely prevented tumor outgrowth.

BRIEF DESCRIPTION OF THE SEQUENCE IDENTIFIERS

[0008]

SEQ ID NOs: 1-3 set forth amino acid sequences of exemplary light chain CDR sequences for BCMA CARs contemplated herein.

SEQ ID NOs: 4-6 set forth amino acid sequences of exemplary heavy chain CDR sequences for BCMA CARs contemplated herein.

SEQ ID NO: 7 sets forth an amino acid sequence of an exemplary light chain sequences for BCMA CARs contemplated herein.

SEQ ID NO: 8 sets forth an amino acid sequence of an exemplary heavy chain sequences for BCMA CARs contemplated herein.

SEQ ID NO: 9 sets forth an amino acid sequence of an exemplary BCMA CAR contemplated herein.

SEQ ID NO: 10 set forth a polynucleotide sequence that encode an exemplary BCMA CAR contemplated herein.

SEQ ID NO: 11 sets forth the amino acid sequence of human BCMA.

SEQ ID NO: 12-22 set for the amino acid sequence of various linkers.

SEQ ID NOs: 23-35 set for the amino acid sequence of protease cleavage sites and self-cleaving polypeptide cleavage sites.

SEQ ID NO: 36 sets for the polynucleotide sequence of a vector encoding a BCMA CAR.

DETAILED DESCRIPTION

A. OVERVIEW

[0009] Described herein are improved compositions and methods for treating B cell related conditions. As used herein, the term "B cell related conditions" relates to conditions involving inappropriate B cell activity and B cell malignancies.

[0010] In particular, the disclosure relates to improved adoptive cell therapy of B cell related conditions using genetically modified immune effector cells. Genetic approaches offer a potential means to enhance immune recognition and elimination of cancer cells. One promising strategy is to genetically engineer immune effector cells to express chimeric antigen receptors (CAR) that redirect cytotoxicity toward cancer cells. However, existing adoptive cell immunotherapies for treating B cell disorders present a serious risk of compromising humoral immunity because the cells target antigens expressed on all of, or the majority of, B cells. Accordingly, such therapies are not clinically desirable and thus, a need in the art remains for more efficient therapies for B cell related conditions that spare humoral immunity.

[0011] The improved compositions and methods of adoptive cell therapy disclosed herein, provide genetically modified

immune effector cells that can readily be expanded, exhibit long-term persistence *in vivo*, and reduce impairment of humoral immunity by targeting B cells expression B cell maturation antigen (BCMA, also known as CD269 or tumor necrosis factor receptor superfamily, member 17; TNFRSF17).

[0012] BCMA is a member of the tumor necrosis factor receptor superfamily (see, e.g., Thompson et al., J. Exp. Medicine, 192(1): 129-135, 2000, and Mackay et al., Annu. Rev. Immunol, 21: 231-264, 2003. BCMA binds B-cell activating factor (BAFF) and a proliferation inducing ligand (APRIL) (see, e.g., Mackay et al., 2003 and Kalled et al., Immunological Reviews, 204: 43-54, 2005). Among nonmalignant cells, BCMA has been reported to be expressed mostly in plasma cells and subsets of mature B-cells (see, e.g., Laabi et al., EMBO J., 77(1): 3897-3904, 1992; Laabi et al., Nucleic Acids Res., 22(7): 1147-1154, 1994; Kalled et al., 2005; O'Connor et al., J. Exp. Medicine, 199(1): 91-97, 2004; and Ng et al., J. Immunol., 73(2): 807-817, 2004. Mice deficient in BCMA are healthy and have normal numbers of B cells, but the survival of long-lived plasma cells is impaired (see, e.g., O'Connor et al., 2004; Xu et al., Mol. Cell. Biol, 21(12): 4067-4074, 2001; and Schiemann et al., Science, 293(5537): 2 111-21 14, 2001). BCMA RNA has been detected universally in multiple myeloma cells and in other lymphomas, and BCMA protein has been detected on the surface of plasma cells from multiple myeloma patients by several investigators (see, e.g., Novak et al., Blood, 103(2): 689-694, 2004; Neri et al., Clinical Cancer Research, 73(19): 5903-5909, 2007; Bellucci et al., Blood, 105(10): 3945-3950, 2005; and Moreaux et al., Blood, 703(8): 3148-3157, 2004.

[0013] CARs comprising murine anti-BCMA antibody sequences may be highly efficacious compared to BCMA CARs comprising particular human antibody sequences; may undergo robust *in vivo* expansion; and recognize human B cells expressing BMCA; may show cytotoxic activity against the BCMA expressing B cells; and may show signs of inducing a cytokine storm, a potentially fatal condition where the cytokines released by activated T cells create a sudden inflammatory response in the system that spurs a noninfectious fever.

[0014] In particular, a CAR comprising a murine anti-BCMA antibody or antigen binding fragment, a transmembrane domain, and one or more intracellular signaling domains is provided.

[0015] In particular, an immune effector cell is genetically modified to express a CAR contemplated herein is provided. T cells expressing a CAR are referred to herein as CAR T cells or CAR modified T cells.

[0016] The genetically modified immune effector cells contemplated herein, may be administered to a patient with a B cell related condition, e.g., an autoimmune disease associated with B cells or a B cell malignancy.

[0017] The practice of the present disclosure will employ, unless indicated specifically to the contrary, conventional methods of chemistry, biochemistry, organic chemistry, molecular biology, microbiology, recombinant DNA techniques, genetics, immunology, and cell biology that are within the skill of the art, many of which are described below for the purpose of illustration. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al., Molecular Cloning: A Laboratory Manual (3rd Edition, 2001); Sambrook, et al., Molecular Cloning: A Laboratory Manual (2nd Edition, 1989); Maniatis et al., Molecular Cloning: A Laboratory Manual (1982); Ausubel et al., Current Protocols in Molecular Biology (John Wiley and Sons, updated July 2008); Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology, Greene Pub. Associates and Wiley-Interscience; Glover, DNA Cloning: A Practical Approach, vol. I & II (IRL Press, Oxford, 1985); Anand, Techniques for the Analysis of Complex Genomes, (Academic Press, New York, 1992); Transcription and Translation (B. Hames & S. Higgins, Eds., 1984); Perbal, A Practical Guide to Molecular Cloning (1984); Harlow and Lane, Antibodies, (ColdSpring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1998) Current Protocols in Immunology Q. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach and W. Strober, eds., 1991); *Annual Review of Immunology*; as well as monographs in journals such as *Advances in Immunology*.

B. DEFINITIONS

[0018] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, preferred embodiments of compositions, methods and materials are described herein. For the purposes of the present disclosure, the following terms are defined below.

[0019] The articles "a," "an," and "the" are used herein to refer to one or to more than one (*i.e.*, to at least one, or to one or more) of the grammatical object of the article. By way of example, "an element" means one element or one or more elements.

[0020] The use of the alternative (e.g., "or") should be understood to mean either one, both, or any combination thereof of the alternatives.

[0021] The term "and/or" should be understood to mean either one, or both of the alternatives.

[0022] As used herein, the term "about" or "approximately" refers to a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2% or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length. The term "about" or "approximately" may refer a range of quantity, level, value, number, frequency, percentage,

dimension, size, amount, weight or length $\pm 15\%$, $\pm 10\%$, $\pm 9\%$, $\pm 8\%$, $\pm 7\%$, $\pm 6\%$, $\pm 5\%$, $\pm 4\%$, $\pm 3\%$, $\pm 2\%$, or $\pm 1\%$ about a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

[0023] Throughout this specification, unless the context requires otherwise, the words "comprise", "comprises" and "comprising" will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements. By "consisting of" is meant including, and limited to, whatever follows the phrase "consisting of." Thus, the phrase "consisting of" indicates that the listed elements are required or mandatory, and that no other elements may be present. By "consisting essentially of" is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the phrase "consisting essentially of" indicates that the listed elements are required or mandatory, but that no other elements are present that materially affect the activity or action of the listed elements.

[0024] Reference throughout this specification to "one embodiment," "an embodiment," "a particular embodiment," "a related embodiment," "a certain embodiment," "an additional embodiment," or "a further embodiment" or combinations thereof means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present invention. Thus, the appearances of the foregoing phrases in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments. It is also understood that the positive recitation of a feature in one embodiment, serves as a basis for excluding the feature in a particular embodiment.

C. CHIMERIC ANTIGEN RECEPTORS

[0025] Genetically engineered receptors that redirect cytotoxicity of immune effector cells toward B cells are provided. These genetically engineered receptors referred to herein as chimeric antigen receptors (CARs). CARs are molecules that combine antibody-based specificity for a desired antigen (e.g., BCMA) with a T cell receptor-activating intracellular domain to generate a chimeric protein that exhibits a specific anti-BCMA cellular immune activity. As used herein, the term, "chimeric," describes being composed of parts of different proteins or DNAs from different origins.

[0026] CARs contemplated herein, comprise an extracellular domain (also referred to as a binding domain or antigen-specific binding domain) that binds to BCMA, a transmembrane domain, and an intracellular signaling domain. Engagement of the anti-BCMA antigen binding domain of the CAR with BCMA on the surface of a target cell results in clustering of the CAR and delivers an activation stimulus to the CAR-containing cell. The main characteristic of CARs are their ability to redirect immune effector cell specificity, thereby triggering proliferation, cytokine production, phagocytosis or production of molecules that can mediate cell death of the target antigen expressing cell in a major histocompatibility (MHC) independent manner, exploiting the cell specific targeting abilities of monoclonal antibodies, soluble ligands or cell specific co-receptors.

[0027] A CAR may comprise an extracellular binding domain that comprises a murine anti-BCMA-specific binding domain; a transmembrane domain; one or more intracellular co-stimulatory signaling domains; and a primary signaling domain.

[0028] A CAR may comprise an extracellular binding domain that comprises a murine anti-BCMA antibody or antigen binding fragment thereof; one or more hinge domains or spacer domains; a transmembrane domain including; one or more intracellular co-stimulatory signaling domains; and a primary signaling domain.

1. BINDING DOMAIN

[0029] CARs contemplated herein may comprise an extracellular binding domain that comprises a murine anti-BCMA antibody or antigen binding fragment thereof that specifically binds to a human BCMA polypeptide expressed on a B cell. As used herein, the terms, "binding domain," "extracellular domain," "extracellular binding domain," "antigen-specific binding domain," and "extracellular antigen specific binding domain," are used interchangeably and provide a CAR with the ability to specifically bind to the target antigen of interest, e.g., BCMA. The binding domain may be derived either from a natural, synthetic, semi-synthetic, or recombinant source.

[0030] The terms "specific binding affinity" or "specifically binds" or "specifically bound" or "specific binding" or "specifically targets" as used herein, describe binding of an anti-BCMA antibody or antigen binding fragment thereof (or a CAR comprising the same) to BCMA at greater binding affinity than background binding. A binding domain (or a CAR comprising a binding domain or a fusion protein containing a binding domain) "specifically binds" to a BCMA if it binds to or associates with BCMA with an affinity or K_a (i.e., an equilibrium association constant of a particular binding interaction with units of $1/M$) of, for example, greater than or equal to about

$10^5 M^{-1}$. A binding domain may (or a fusion protein thereof) bind to a target with a K_a greater than or equal to about $10^6 M^{-1}$, $10^7 M^{-1}$, $10^8 M^{-1}$, $10^9 M^{-1}$, $10^{10} M^{-1}$, $10^{11} M^{-1}$, $10^{12} M^{-1}$, or $10^{13} M^{-1}$. "High affinity" binding domains (or single chain

fusion proteins thereof) refers to those binding domains with a K_d of at least 10^7 M⁻¹, at least 10^8 M⁻¹, at least 10^9 M⁻¹, at least 10^{10} M⁻¹, at least 10^{11} M⁻¹, at least 10^{12} M⁻¹, at least 10^{13} M⁻¹, or greater.

[0031] Alternatively, affinity may be defined as an equilibrium dissociation constant (K_d) of a particular binding interaction with units of M (e.g., 10^{-5} M to 10^{-13} M, or less). Affinities of binding domain polypeptides and CAR proteins according to the present disclosure can be readily determined using conventional techniques, e.g., by competitive ELISA (enzyme-linked immunosorbent assay), or by binding association, or displacement assays using labeled ligands, or using a surface-plasmon resonance device such as the Biacore T100, which is available from Biacore, Inc., Piscataway, NJ, or optical biosensor technology such as the EPIC system or EnSpire that are available from Corning and Perkin Elmer respectively (see also, e.g., Scatchard et al. (1949) Ann. N.Y. Acad. Sci. 51:660; and U.S. Patent Nos. 5,283,173; 5,468,614, or the equivalent).

[0032] The affinity of specific binding may be about 2 times greater than background binding, about 5 times greater than background binding, about 10 times greater than background binding, about 20 times greater than background binding, about 50 times greater than background binding, about 100 times greater than background binding, or about 1000 times greater than background binding or more.

[0033] The extracellular binding domain of a CAR may comprise an antibody or antigen binding fragment thereof. An "antibody" refers to a binding agent that is a polypeptide comprising at least a light chain or heavy chain immunoglobulin variable region which specifically recognizes and binds an epitope of an antigen, such as a peptide, lipid, polysaccharide, or nucleic acid containing an antigenic determinant, such as those recognized by an immune cell.

[0034] An "antigen (Ag)" refers to a compound, composition, or substance that can stimulate the production of antibodies or a T cell response in an animal, including compositions (such as one that includes a cancer-specific protein) that are injected or absorbed into an animal. An antigen reacts with the products of specific humoral or cellular immunity, including those induced by heterologous antigens, such as the disclosed antigens. The target antigen may be an epitope of a BCMA polypeptide.

[0035] An "epitope" or "antigenic determinant" refers to the region of an antigen to which a binding agent binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5, about 9, or about 8-10 amino acids in a unique spatial conformation.

[0036] Antibodies include antigen binding fragments thereof, such as Camel Ig, Ig NAR, Fab fragments, Fab' fragments, F(ab)'₂ fragments, F(ab)'₃ fragments, Fv, single chain Fv proteins ("scFv"), bis-scFv, (scFv)₂, minibodies, diabodies, triabodies, tetrabodies, disulfide stabilized Fv proteins ("dsFv"), and single-domain antibody (sdAb, Nanobody) and portions of full length antibodies responsible for antigen binding. The term also includes genetically engineered forms such as chimeric antibodies (for example, humanized murine antibodies), heteroconjugate antibodies (such as, bispecific antibodies) and antigen binding fragments thereof. See also, Pierce Catalog and Handbook, 1994-1995 (Pierce Chemical Co., Rockford, IL); Kuby, J., Immunology, 3rd Ed., W. H. Freeman & Co., New York, 1997.

[0037] As would be understood by the skilled person and as described elsewhere herein, a complete antibody comprises two heavy chains and two light chains. Each heavy chain consists of a variable region and a first, second, and third constant region, while each light chain consists of a variable region and a constant region. Mammalian heavy chains are classified as α , δ , ϵ , γ , and μ . Mammalian light chains are classified as λ or κ . Immunoglobulins comprising the α , δ , ϵ , γ , and μ heavy chains are classified as immunoglobulin (Ig)A, IgD, IgE, IgG, and IgM. The complete antibody forms a "Y" shape. The stem of the Y consists of the second and third constant regions (and for IgE and IgM, the fourth constant region) of two heavy chains bound together and disulfide bonds (inter-chain) are formed in the hinge. Heavy chains γ , α and δ have a constant region composed of three tandem (in a line) Ig domains, and a hinge region for added flexibility; heavy chains μ and ϵ have a constant region composed of four immunoglobulin domains. The second and third constant regions are referred to as "CH2 domain" and "CH3 domain", respectively. Each arm of the Y includes the variable region and first constant region of a single heavy chain bound to the variable and constant regions of a single light chain. The variable regions of the light and heavy chains are responsible for antigen binding.

[0038] Light and heavy chain variable regions contain a "framework" region interrupted by three hypervariable regions, also called "complementarity-determining regions" or "CDRs." The CDRs can be defined or identified by conventional methods, such as by sequence according to Kabat *et al* (Wu, TT and Kabat, E. A., J Exp Med. 132(2):211-50, (1970); Borden, P. and Kabat E. A., PNAS, 84: 2440-2443 (1987); (see, Kabat et al., Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services, 1991), or by structure according to Chothia *et al* (Chothia, C. and Lesk, A.M., J Mol. Biol., 196(4): 901-917 (1987), Chothia, C. et al, Nature, 342: 877 - 883 (1989)).

[0039] The sequences of the framework regions of different light or heavy chains are relatively conserved within a species, such as humans. The framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains, serves to position and align the CDRs in three-dimensional space. The CDRs are primarily responsible for binding to an epitope of an antigen. The CDRs of each chain are typically referred to as CDR1, CDR2, and CDR3, numbered sequentially starting from the N-terminus, and are also typically identified by the chain in which

the particular CDR is located. Thus, the CDRs located in the variable domain of the heavy chain of the antibody are referred to as CDRH1, CDRH2, and CDRH3, whereas the CDRs located in the variable domain of the light chain of the antibody are referred to as CDRL1, CDRL2, and CDRL3. Antibodies with different specificities (*i.e.*, different combining sites for different antigens) have different CDRs. Although it is the CDRs that vary from antibody to antibody, only a limited number of amino acid positions within the CDRs are directly involved in antigen binding. These positions within the CDRs are called specificity determining residues (SDRs). Illustrative examples of light chain CDRs that are suitable for constructing humanized BCMA CARs contemplated herein include, but are not limited to the CDR sequences set forth in SEQ ID NOs: 1-3. Illustrative examples of heavy chain CDRs that are suitable for constructing humanized BCMA CARs contemplated herein include, but are not limited to the CDR sequences set forth in SEQ ID NOs: 4-6.

[0040] References to "V_H" or "VH" refer to the variable region of an immunoglobulin heavy chain, including that of an antibody, Fv, scFv, dsFv, Fab, or other antibody fragment as disclosed herein. References to "V_L" or "VL" refer to the variable region of an immunoglobulin light chain, including that of an antibody, Fv, scFv, dsFv, Fab, or other antibody fragment as disclosed herein.

[0041] A "monoclonal antibody" is an antibody produced by a single clone of B lymphocytes or by a cell into which the light and heavy chain genes of a single antibody have been transfected. Monoclonal antibodies are produced by methods known to those of skill in the art, for instance by making hybrid antibody-forming cells from a fusion of myeloma cells with immune spleen cells. Monoclonal antibodies include humanized monoclonal antibodies.

[0042] A "chimeric antibody" has framework residues from one species, such as human, and CDRs (which generally confer antigen binding) from another species, such as a mouse. A CAR contemplated herein may comprise antigen-specific binding domain that is a chimeric antibody or antigen binding fragment thereof.

[0043] A "humanized" antibody is an immunoglobulin including a human framework region and one or more CDRs from a non-human (for example a mouse, rat, or synthetic) immunoglobulin. The non-human immunoglobulin providing the CDRs is termed a "donor," and the human immunoglobulin providing the framework is termed an "acceptor."

[0044] A murine anti-BCMA antibody or antigen binding fragment thereof, may include but is not limited to a Camel Ig (a camelid antibody (VHH)), Ig NAR, Fab fragments, Fab' fragments, F(ab')₂ fragments, F(ab')₃ fragments, Fv, single chain Fv antibody ("scFv"), bis-scFv, (scFv)₂, minibody, diabody, triabody, tetrabody, disulfide stabilized Fv protein ("dsFv"), and single-domain antibody (sdAb, Nanobody).

[0045] "Camel Ig" or "camelid VHH" as used herein refers to the smallest known antigen-binding unit of a heavy chain antibody (Koch-Nolte, et al, FASEB J., 21: 3490-3498 (2007)). A "heavy chain antibody" or a "camelid antibody" refers to an antibody that contains two VH domains and no light chains (Riechmann L. et al, J. Immunol. Methods 231:25-38 (1999); WO94/04678; WO94/25591; U.S. Patent No. 6,005,079).

[0046] "IgNAR" of "immunoglobulin new antigen receptor" refers to class of antibodies from the shark immune repertoire that consist of homodimers of one variable new antigen receptor (VNAR) domain and five constant new antigen receptor (CNAR) domains. IgNARs represent some of the smallest known immunoglobulin-based protein scaffolds and are highly stable and possess efficient binding characteristics. The inherent stability can be attributed to both (i) the underlying Ig scaffold, which presents a considerable number of charged and hydrophilic surface exposed residues compared to the conventional antibody VH and VL domains found in murine antibodies; and (ii) stabilizing structural features in the complementary determining region (CDR) loops including inter-loop disulphide bridges, and patterns of intra-loop hydrogen bonds.

[0047] Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

[0048] "Fv" is the minimum antibody fragment which contains a complete antigen-binding site. In one embodiment, a two-chain Fv species consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv (scFv) species, one heavy- and one light-chain variable domain can be covalently linked by a flexible peptide linker such that the light and heavy chains can associate in a "dimeric" structure analogous to that in a two-chain Fv species. It is in this configuration that the three hypervariable regions (HVRs) of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six HVRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three HVRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0049] The Fab fragment contains the heavy- and light-chain variable domains and also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in

which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0050] The term "diabodies" refers to antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) in the same polypeptide chain (VH-VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies may be bivalent or bispecific. Diabodies are described more fully in, for example, EP 404,097; WO 1993/01161; Hudson et al., *Nat. Med.* 9:129-134 (2003); and Hollinger et al., *PNAS USA* 90: 6444-6448 (1993). Triabodies and tetrabodies are also described in Hudson et al., *Nat. Med.* 9:129-134 (2003).

[0051] "Single domain antibody" or "sdAb" or "nanobody" refers to an antibody fragment that consists of the variable region of an antibody heavy chain (VH domain) or the variable region of an antibody light chain (VL domain) (Holt, L., et al, *Trends in Biotechnology*, 21(11): 484-490).

[0052] "Single-chain Fv" or "scFv" antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain and in either orientation (e.g., VL-VH or VH-VL). Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv, see, e.g., Pluckthün, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., (Springer-Verlag, New York, 1994), pp. 269-315.

[0053] A CAR contemplated herein may comprise antigen-specific binding domain that is a murine scFv. Single chain antibodies may be cloned from the V region genes of a hybridoma specific for a desired target. The production of such hybridomas has become routine. A technique which can be used for cloning the variable region heavy chain (VH) and variable region light chain (VL) has been described, for example, in Orlandi et al., *PNAS*, 1989; 86: 3833-3837.

[0054] The antigen-specific binding domain may be a murine scFv that binds a human BCMA polypeptide. Illustrative examples of variable heavy

chains that are suitable for constructing BCMA CARs contemplated herein include, but are not limited to the amino acid sequences set forth in SEQ ID NO: 8. Illustrative examples of variable light chains that are suitable for constructing BCMA CARs contemplated herein include, but are not limited to the amino acid sequences set forth in SEQ ID NO: 7.

[0055] BCMA-specific binding domains provided herein also comprise one, two, three, four, five, or six CDRs. Such CDRs may be nonhuman CDRs or altered nonhuman CDRs selected from CDRL1, CDRL2 and CDRL3 of the light chain and CDRH1, CDRH2 and CDRH3 of the heavy chain. In certain embodiments, a BCMA-specific binding domain comprises (a) a light chain variable region that comprises a light chain CDRL1, a light chain CDRL2, and a light chain CDRL3, and (b) a heavy chain variable region that comprises a heavy chain CDRH1, a heavy chain CDRH2, and a heavy chain CDRH3.

2. LINKERS

[0056] The CARs contemplated herein may comprise linker residues between the various domains, e.g., , added for appropriate spacing and conformation of the molecule. The linker may be a variable region linking sequence. A "variable region linking sequence," is an amino acid sequence that connects the V_H and V_L domains and provides a spacer function compatible with interaction of the two sub-binding domains so that the resulting polypeptide retains a specific binding affinity to the same target molecule as an antibody that comprises the same light and heavy chain variable regions. CARs contemplated herein, may comprise one, two, three, four, or five or more linkers. The length of a linker may be about 1 to about 25 amino acids, about 5 to about 20 amino acids, or about 10 to about 20 amino acids, or any intervening length of amino acids. In some embodiments, the linker is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more amino acids long.

[0057] Illustrative examples of linkers include glycine polymers $(G)_n$; glycine-serine polymers $(G_{1-5}S_{1-5})_n$, where n is an integer of at least one, two, three, four, or five; glycine-alanine polymers; alanine-serine polymers; and other flexible linkers known in the art. Glycine and glycine-serine polymers are relatively unstructured, and therefore may be able to serve as a neutral tether between domains of fusion proteins such as the CARs described herein. Glycine accesses significantly more phi-psi space than even alanine, and is much less restricted than residues with longer side chains (see Scheraga, *Rev. Computational Chem.* 11173-142 (1992)). The ordinarily skilled artisan will recognize that design of a CAR in particular embodiments can include linkers that are all or partially flexible, such that the linker can include a flexible linker as well as one or more portions that confer less flexible structure to provide for a desired CAR structure.

[0058] Other exemplary linkers include, but are not limited to the following amino acid sequences: GGG; DGGGS (SEQ ID NO: 12); TGEKP (SEQ ID NO: 13) (see, e.g., Liu et al., *PNAS* 5525-5530 (1997)); GGRR (SEQ ID NO: 14) (Pomerantz *et al.* 1995, supra); $(GGGGS)_n$ wherein $n = 1, 2, 3, 4$ or 5 (SEQ ID NO: 15) (Kim et al., *PNAS* 93, 1156-1160 (1996.); EGKSSGSGSESKVD (SEQ ID NO: 16) (Chaudhary et al., 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87:1066-1070); KESGSVSSEQLAQFRSLD (SEQ ID NO: 17) (Bird et al., 1988, *Science* 242:423-426), GGRRGGGS (SEQ ID NO: 18); LRQRDGERP (SEQ ID NO: 19); LRQKDGGSERP (SEQ ID NO: 20); $LRQKd(GGGGS)_2$ ERP (SEQ ID NO: 21). Alternatively, flexible linkers can be rationally designed using a computer program capable of modeling both DNA-binding sites and the peptides themselves (Desjarlais & Berg, *PNAS* 90:2256-2260 (1993), *PNAS* 91:11099-11103 (1994) or

by phage display methods. In one embodiment, the linker comprises the following amino acid sequence: GSTSGSG-KPGSGEGSTKG (SEQ ID NO: 22) (Cooper et al., Blood, 101(4): 1637-1644 (2003)).

3. SPACER DOMAIN

[0059] The binding domain of the CAR may be followed by one or more "spacer domains," which refers to the region that moves the antigen binding domain away from the effector cell surface to enable proper cell/cell contact, antigen binding and activation (Patel et al., Gene Therapy, 1999; 6: 412-419). The hinge domain may be derived either from a natural, synthetic, semi-synthetic, or recombinant source. A spacer domain may be a portion of an immunoglobulin, including, but not limited to, one or more heavy chain constant regions, e.g., CH2 and CH3. The spacer domain can include the amino acid sequence of a naturally occurring immunoglobulin hinge region or an altered immunoglobulin hinge region.

[0060] The spacer domain may comprise the CH2 and CH3 domains of IgG1 or IgG4.

4. HINGE DOMAIN

[0061] The binding domain of the CAR is generally followed by one or more "hinge domains," which plays a role in positioning the antigen binding domain away from the effector cell surface to enable proper cell/cell contact, antigen binding and activation. A CAR generally comprises one or more hinge domains between the binding domain and the transmembrane domain (TM). The hinge domain may be derived either from a natural, synthetic, semi-synthetic, or recombinant source. The hinge domain can include the amino acid sequence of a naturally occurring immunoglobulin hinge region or an altered immunoglobulin hinge region.

[0062] An "altered hinge region" refers to (a) a naturally occurring hinge region with up to 30% amino acid changes (e.g., up to 25%, 20%, 15%, 10%, or 5% amino acid substitutions or deletions), (b) a portion of a naturally occurring hinge region that is at least 10 amino acids (e.g., at least 12, 13, 14 or 15 amino acids) in length with up to 30% amino acid changes (e.g., up to 25%, 20%, 15%, 10%, or 5% amino acid substitutions or deletions), or (c) a portion of a naturally occurring hinge region that comprises the core hinge region (which may be 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15, or at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acids in length). One or more cysteine residues in a naturally occurring immunoglobulin hinge region may be substituted by one or more other amino acid residues (e.g., one or more serine residues). An altered immunoglobulin hinge region may alternatively or additionally have a proline residue of a wild type immunoglobulin hinge region substituted by another amino acid residue (e.g., a serine residue).

[0063] Other illustrative hinge domains suitable for use in the CARs described herein include the hinge region derived from the extracellular regions of type 1 membrane proteins such as CD8 α , CD4, CD28 and CD7, which may be wild-type hinge regions from these molecules or may be altered. The hinge domain may comprise a CD8 α hinge region.

5. TRANSMEMBRANE (TM) DOMAIN

[0064] The "transmembrane domain" is the portion of the CAR that fuses the extracellular binding portion and intracellular signaling domain and anchors the CAR to the plasma membrane of the immune effector cell. The TM domain may be derived either from a natural, synthetic, semi-synthetic, or recombinant source. The TM domain may be derived from (i.e., comprise at least the transmembrane region(s) of) the alpha, beta or zeta chain of the T-cell receptor, CD3 ϵ , CD3 ζ , CD4, CD5, CD8 α , CD9, CD 16, CD22, CD27, CD28, CD33, CD37, CD45, CD64, CD80, CD86, CD 134, CD137, CD152, CD 154, and PD1. The TM domain may be synthetic and predominantly comprise hydrophobic residues such as leucine and valine.

[0065] The CARs contemplated herein may comprise a TM domain derived from CD8 α . A CAR contemplated herein may comprise a TM domain derived from CD8 α and a short oligo- or polypeptide linker, preferably between 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids in length that links the TM domain and the intracellular signaling domain of the CAR. A glycine-serine based linker provides a particularly suitable linker.

6. INTRACELLULAR SIGNALING DOMAIN

[0066] CARs contemplated herein may comprise an intracellular signaling domain. An "intracellular signaling domain," refers to the part of a CAR that participates in transducing the message of effective BCMA CAR binding to a human BCMA polypeptide into the interior of the immune effector cell to elicit effector cell function, e.g., activation, cytokine production, proliferation and cytotoxic activity, including the release of cytotoxic factors to the CAR-bound target cell, or other cellular responses elicited with antigen binding to the extracellular CAR domain.

[0067] The term "effector function" refers to a specialized function of an immune effector cell. Effector function of the T cell, for example, may be cytolytic activity or help or activity including the secretion of a cytokine. Thus, the term

"intracellular signaling domain" refers to the portion of a protein which transduces the effector function signal and that directs the cell to perform a specialized function. While usually the entire intracellular signaling domain can be employed, in many cases it is not necessary to use the entire domain. To the extent that a truncated portion of an intracellular signaling domain is used, such truncated portion may be used in place of the entire domain as long as it transduces the effector function signal. The term intracellular signaling domain is meant to include any truncated portion of the intracellular signaling domain sufficient to transducing effector function signal.

[0068] It is known that signals generated through the TCR alone are insufficient for full activation of the T cell and that a secondary or co-stimulatory signal is also required. Thus, T cell activation can be said to be mediated by two distinct classes of intracellular signaling domains: primary signaling domains that initiate antigen-dependent primary activation through the TCR (e.g., a TCR/CD3 complex) and co-stimulatory signaling domains that act in an antigen-independent manner to provide a secondary or co-stimulatory signal. A CAR contemplated herein may comprise an intracellular signaling domain that comprises one or more "co-stimulatory signaling domain" and a "primary signaling domain."

[0069] Primary signaling domains regulate primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary signaling domains that act in a stimulatory manner may contain signaling motifs which are known as immunoreceptor tyrosine-based activation motifs or ITAMs.

[0070] Illustrative examples of ITAM containing primary signaling domains that are of particular use include those derived from TCR ζ , FcR γ , FcR β , CD3 γ , CD3 δ , CD3 ϵ , CD3 ζ , CD22, CD79a, CD79b, and CD66d. In particular, a CAR may comprise a CD3 ζ primary signaling domain and one or more co-stimulatory signaling domains. The intracellular primary signaling and co-stimulatory signaling domains may be linked in any order in tandem to the carboxyl terminus of the transmembrane domain.

[0071] CARs contemplated herein comprise one or more co-stimulatory signaling domains to enhance the efficacy and expansion of T cells expressing CAR receptors. As used herein, the term, "co-stimulatory signaling domain," or "co-stimulatory domain", refers to an intracellular signaling domain of a co-stimulatory molecule. Co-stimulatory molecules are cell surface molecules other than antigen receptors or Fc receptors that provide a second signal required for efficient activation and function of T lymphocytes upon binding to antigen. Illustrative examples of such co-stimulatory molecules include CARD11, CD2, CD7, CD27, CD28, CD30, CD40, CD54 (ICAM), CD83, CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD152 (CTLA4), CD223 (LAG3), CD270 (HVEM), CD273 (PD-L2), CD274 (PD-L1), CD278 (ICOS), DAP10, LAT, NKD2C SLP76, TRIM, and ZAP70. A CAR may comprise one or more co-stimulatory signaling domains selected from the group consisting of CD28, CD137, and CD134, and a CD3 ζ primary signaling domain.

[0072] A CAR may comprise CD28 and CD137 co-stimulatory signaling domains and a CD3 ζ primary signaling domain.

[0073] A CAR may comprise CD28 and CD134 co-stimulatory signaling domains and a CD3 ζ primary signaling domain.

[0074] A CAR may comprise CD137 and CD134 co-stimulatory signaling domains and a CD3 ζ primary signaling domain.

[0075] A CARs contemplated herein may comprise a murine anti-BCMA antibody or antigen binding fragment thereof that specifically binds to a BCMA polypeptide expressed on B cells.

[0076] A CAR may comprises a murine anti-BCMA scFv that binds a BCMA polypeptide; a transmembrane domain derived from a polypeptide selected from the group consisting of: alpha, beta or zeta chain of the T-cell receptor, CD3 ϵ , CD3 ζ , CD4, CD5, CD8 α , CD9, CD 16, CD22, CD27, CD28, CD33, CD37, CD45, CD64, CD80, CD86, CD 134, CD137, CD152, CD 154, and PD1; and one or more intracellular co-stimulatory signaling domains from a co-stimulatory molecule selected from the group consisting of: CARD11, CD2, CD7, CD27, CD28, CD30, CD40, CD54 (ICAM), CD83, CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD152 (CTLA4), CD223 (LAG3), CD270 (HVEM), CD273 (PD-L2), CD274 (PD-L1), CD278 (ICOS), DAP10, LAT, NKD2C SLP76, TRIM, and ZAP70; and a primary signaling domain from TCR ζ , FcR γ , FcR β , CD3 γ , CD3 δ , CD3 ϵ , CD3 ζ , CD22, CD79a, CD79b, and CD66d.

[0077] A CAR may comprise a murine anti-BCMA scFv that binds a BCMA polypeptide; a hinge domain selected from the group consisting of: IgG1 hinge/CH2/CH3, IgG4 hinge/CH2/CH3, and a CD8 α hinge; a transmembrane domain derived from a polypeptide selected from the group consisting of: alpha, beta or zeta chain of the T-cell receptor, CD3 ϵ , CD3 ζ , CD4, CD5, CD8 α , CD9, CD 16, CD22, CD27, CD28, CD33, CD37, CD45, CD64, CD80, CD86, CD 134, CD137, CD152, CD 154, and PD1; and one or more intracellular co-stimulatory signaling domains from a co-stimulatory molecule selected from the group consisting of: CARD11, CD2, CD7, CD27, CD28, CD30, CD40, CD54 (ICAM), CD83, CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD152 (CTLA4), CD223 (LAG3), CD270 (HVEM), CD273 (PD-L2), CD274 (PD-L1), CD278 (ICOS), DAP10, LAT, NKD2C SLP76, TRIM, and ZAP70; and a primary signaling domain from TCR ζ , FcR γ , FcR β , CD3 γ , CD3 δ , CD3 ϵ , CD3 ζ , CD22, CD79a, CD79b, and CD66d.

[0078] A CAR may comprise a murine anti-BCMA scFv that binds a BCMA polypeptide; a hinge domain selected from the group consisting of: IgG1 hinge/CH2/CH3, IgG4 hinge/CH2/CH3, and a CD8 α hinge; a transmembrane domain derived from a polypeptide selected from the group consisting of: alpha, beta or zeta chain of the T-cell receptor, CD3 ϵ , CD3 ζ , CD4, CD5, CD8 α , CD9, CD 16, CD22, CD27, CD28, CD33, CD37, CD45, CD64, CD80, CD86, CD 134, CD137, CD152, CD 154, and PD1; a short oligo- or polypeptide linker, preferably between 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids in length that links the TM domain to the intracellular signaling domain of the CAR; and one or more intracellular

co-stimulatory signaling domains from a co-stimulatory molecule selected from the group consisting of: CARD11, CD2, CD7, CD27, CD28, CD30, CD40, CD54 (ICAM), CD83, CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD152 (CTLA4), CD223 (LAG3), CD270 (HVEM), CD273 (PD-L2), CD274 (PD-L1), CD278 (ICOS), DAP10, LAT, NKD2C SLP76, TRIM, and ZAP70; and a primary signaling domain from TCR ζ , FcR γ , FcR β , CD3 γ , CD3 δ , CD3 ϵ , CD3 ζ , CD22, CD79a, CD79b, and CD66d.

[0079] A CAR may comprise a murine anti-BCMA scFv that binds a BCMA polypeptide; a hinge domain comprising an IgG1 hinge/CH2/CH3 polypeptide and a CD8 α polypeptide; a CD8 α transmembrane domain comprising a polypeptide linker of about 3 to about 10 amino acids; a CD137 intracellular co-stimulatory signaling domain; and a CD3 ζ primary signaling domain.

[0080] A CAR may comprise a murine anti-BCMA scFv that binds a BCMA polypeptide; a hinge domain comprising a CD8 α polypeptide; a CD8 α transmembrane domain comprising a polypeptide linker of about 3 to about 10 amino acids; a CD 134 intracellular co-stimulatory signaling domain; and a CD3 ζ primary signaling domain.

[0081] A CAR may comprise a murine anti-BCMA scFv that binds a BCMA polypeptide; a hinge domain comprising a CD8 α polypeptide; a CD8 α transmembrane domain comprising a polypeptide linker of about 3 to about 10 amino acids; a CD28 intracellular co-stimulatory signaling domain; and a CD3 ζ primary signaling domain.

[0082] Moreover, the design of the CARs contemplated herein enable improved expansion, long-term persistence, and tolerable cytotoxic properties in T cells expressing the CARs compared to non-modified T cells or T cells modified to express other CARs.

D. POLYPEPTIDES

[0083] The present disclosure contemplates, in part, CAR polypeptides and fragments thereof, cells and compositions comprising the same, and vectors that express polypeptides. A polypeptide comprising one or more CARs as set forth in SEQ ID NO: 9 may be provided.

[0084] "Polypeptide," "polypeptide fragment," "peptide" and "protein" are used interchangeably, unless specified to the contrary, and according to conventional meaning, *i.e.*, as a sequence of amino acids. Polypeptides are not limited to a specific length, *e.g.*, they may comprise a full length protein sequence or a fragment of a full length protein, and may include post-translational modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. The CAR polypeptides contemplated herein may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. Illustrative examples of suitable signal sequences useful in CARs disclosed herein include, but are not limited to the IgG1 heavy chain signal sequence and the CD8 α signal sequence. Polypeptides can be prepared using any of a variety of well-known recombinant and/or synthetic techniques. Polypeptides contemplated herein specifically encompass the CARs of the present disclosure, or sequences that have deletions from, additions to, and/or substitutions of one or more amino acid of a CAR as disclosed herein.

[0085] An "isolated peptide" or an "isolated polypeptide" and the like, as used herein, refer to *in vitro* isolation and/or purification of a peptide or polypeptide molecule from a cellular environment, and from association with other components of the cell, *i.e.*, it is not significantly associated with *in vivo* substances. Similarly, an "isolated cell" refers to a cell that has been obtained from an *in vivo* tissue or organ and is substantially free of extracellular matrix.

[0086] Polypeptides include "polypeptide variants." Polypeptide variants may differ from a naturally occurring polypeptide in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the above polypeptide sequences. For example, it may be desirable to improve the binding affinity and/or other biological properties of the CARs by introducing one or more substitutions, deletions, additions and/or insertions into a binding domain, hinge, TM domain, co-stimulatory signaling domain or primary signaling domain of a CAR polypeptide. Preferably, polypeptides include polypeptides having at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% amino acid identity thereto.

[0087] Polypeptides include "polypeptide fragments." Polypeptide fragments refer to a polypeptide, which can be monomeric or multimeric, that has an amino-terminal deletion, a carboxyl-terminal deletion, and/or an internal deletion or substitution of a naturally-occurring or recombinantly-produced polypeptide. A polypeptide fragment can comprise an amino acid chain at least 5 to about 500 amino acids long. It will be appreciated that, fragments may be at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 150, 200, 250, 300, 350, 400, or 450 amino acids long. Particularly useful polypeptide fragments include functional domains, including antigen-binding domains or fragments of antibodies. In the case of a murine anti-BCMA antibody, useful fragments include, but are not limited to: a CDR region, a CDR3 region of the heavy or light chain; a variable region of a heavy or light chain; a portion of an antibody chain or variable region including two CDRs; and the like.

[0088] The polypeptide may also be fused in-frame or conjugated to a linker or other sequence for ease of synthesis,

purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support.

[0089] As noted above, polypeptides may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of a reference polypeptide can be prepared by mutations in the DNA. Methods

for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985, Proc. Natl. Acad. Sci. USA. 82: 488-492), Kunkel et al., (1987, Methods in Enzymol, 154: 367-382), U.S. Pat. No. 4,873,192, Watson, J. D. et al., (Molecular Biology of the Gene, Fourth Edition, Benjamin/Cummings, Menlo Park, Calif., 1987) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff et al., (1978) Atlas of Protein Sequence and Structure (Natl. Biomed. Res. Found., Washington, D.C.).

[0090] A variant may contain conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. Modifications may be made in the structure of the polynucleotides and polypeptides and still obtain a functional molecule that encodes a variant or derivative polypeptide with desirable characteristics. When it is desired to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, variant polypeptide, one skilled in the art, for example, can change one or more of the codons of the encoding DNA sequence, e.g., according to Table 1.

TABLE 1- Amino Acid Codons

Amino Acids	One letter code	Three letter code	Codons						
Alanine	A	Ala	GCA	GCC	GCG	GCU			
Cysteine	C	Cys	UGC	UGU					
Aspartic acid	D	Asp	GAC	GAU					
Glutamic acid	E	Glu	GAA	GAG					
Phenylalanine	F	Phe	UUC	UUU					
Glycine	G	Gly	GGA	GGC	GGG	GGU			
Histidine	H	His	CAC	CAU					
Isoleucine	I	Iso	AUA	AUC	AUU				
Lysine	K	Lys	AAA	AAG					
Leucine	L	Leu	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	M	Met	AUG						
Asparagine	N	Asn	AAC	AAU					
Proline	P	Pro	CCA	CCC	CCG	CCU			
Glutamine	Q	Gln	CAA	CAG					
Arginine	R	Arg	AGA	AGG	CGA	CGC	CGG	CGU	
Serine	S	Ser	AGC	AGU	UCA	UCC	UCG	UCU	
Threonine	T	Thr	ACA	ACC	ACG	ACU			
Valine	V	Val	GUA	GUC	GUG	GUU			
Tryptophan	W	Trp	UGG						
Tyrosine	Y	Tyr	UAC	UAU					

[0091] Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological activity can be found using computer programs well known in the art, such as DNASTAR™ software. Preferably, amino acid changes in the protein variants disclosed herein are conservative amino acid changes, i.e., substitutions of similarly charged or uncharged amino acids. A conservative amino acid change involves substitution of one of a family of amino acids which are related in their side chains. Naturally occurring amino acids are generally divided into four families: acidic (aspartate, glutamate), basic (lysine, arginine, histidine), non-polar (alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), and uncharged polar (glycine, asparagine, glutamine, cysteine, serine,

threonine, tyrosine) amino acids. Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids. In a peptide or protein, suitable conservative substitutions of amino acids are known to those of skill in this art and generally can be made without altering a biological activity of a resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Benjamin/Cummings Pub. Co., p.224). Exemplary conservative substitutions are described in U.S. Provisional Patent Application No. 61/241,647.

[0092] In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982). Each amino acid has been assigned

a hydropathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cysteine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

[0093] It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.*, still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity.

[0094] As detailed in U.S. Patent No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 $\pm$ 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

[0095] As outlined above, amino acid substitutions may be based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like.

[0096] Polypeptide variants further include glycosylated forms, aggregative conjugates with other molecules, and covalent conjugates with unrelated chemical moieties (e.g., pegylated molecules). Covalent variants can be prepared by linking functionalities to groups which are found in the amino acid chain or at the N- or C-terminal residue, as is known in the art. Variants also include allelic variants, species variants, and muteins. Truncations or deletions of regions which do not affect functional activity of the proteins are also variants.

[0097] Where expression of two or more polypeptides is desired, the polynucleotide sequences encoding them can be separated by an IRES sequence as discussed elsewhere herein. Two or more polypeptides can be expressed as a fusion protein that comprises one or more self-cleaving polypeptide sequences.

[0098] Polypeptides include fusion polypeptides. Fusion polypeptides and polynucleotides encoding fusion polypeptides may be provided, e.g., CARs. Fusion polypeptides and fusion proteins refer to a polypeptide having at least two, three, four, five, six, seven, eight, nine, or ten or more polypeptide segments. Fusion polypeptides are typically linked C-terminus to N-terminus, although they can also be linked C-terminus to C-terminus, N-terminus to N-terminus, or N-terminus to C-terminus. The polypeptides of the fusion protein can be in any order or a specified order. Fusion polypeptides or fusion proteins can also include conservatively modified variants, polymorphic variants, alleles, mutants, subsequences, and interspecies homologs, so long as the desired transcriptional activity of the fusion polypeptide is preserved. Fusion polypeptides may be produced by chemical synthetic methods or by chemical linkage between the two moieties or may generally be prepared using other standard techniques. Ligated DNA sequences comprising the fusion polypeptide are operably linked to suitable transcriptional or translational control elements as discussed elsewhere herein.

[0099] A fusion partner may comprise a sequence that assists in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments or to facilitate transport of the fusion protein through the cell membrane.

[0100] Fusion polypeptides may further comprise a polypeptide cleavage signal between each of the polypeptide domains described herein. In addition, polypeptide site can be put into any linker peptide sequence. Exemplary polypeptide cleavage signals include polypeptide cleavage recognition sites such as protease cleavage sites, nuclease cleavage sites (e.g., rare restriction enzyme recognition sites, self-cleaving ribozyme recognition sites), and self-cleaving viral oligopeptides (see deFelipe and Ryan, 2004. *Traffic*, 5(8); 616-26).

[0101] Suitable protease cleavage sites and self-cleaving peptides are known to the skilled person (see, e.g., in Ryan et al., 1997. *J. Gener. Virol.* 78, 699-722; Scymczak et al. (2004) *Nature Biotech.* 5, 589-594). Exemplary protease

cleavage sites include, but are not limited to the cleavage sites of potyvirus NIa proteases (e.g., tobacco etch virus protease), potyvirus HC proteases, potyvirus P1 (P35) proteases, byovirus NIa proteases, byovirus RNA-2-encoded proteases, aphthovirus L proteases, enterovirus 2A proteases, rhinovirus 2A proteases, picorna 3C proteases, comovirus 24K proteases, nepovirus 24K proteases, RTSV (rice tungro spherical virus) 3C-like protease, PYVF (parsnip yellow fleck virus) 3C-like protease, heparin, thrombin, factor Xa and enterokinase. Due to its high cleavage stringency, TEV (tobacco etch virus) protease cleavage sites are preferred in one embodiment, e.g., EXXYXQ(G/S) (SEQ ID NO: 23), for example, ENLYFQG (SEQ ID NO: 24) and ENLYFQS (SEQ ID NO: 25), wherein X represents any amino acid (cleavage by TEV occurs between Q and G or Q and S).

[0102] Self-cleaving peptides may include those polypeptide sequences obtained from potyvirus and cardiovirus 2A peptides, FMDV (foot-and-mouth disease virus), equine rhinitis A virus, Thossea asigna virus and porcine teschovirus.

[0103] The self-cleaving polypeptide site may comprise a 2A or 2A-like site, sequence or domain (Donnelly et al., 2001. J. Gen. Virol. 82:1027-1041).

TABLE 2: Exemplary 2A sites include the following sequences:

SEQ ID NO: 26	LLNFDLLKLAGDVESNPGP
SEQ ID NO: 27	TLNFDLLKLAGDVESNPGP
SEQ ID NO: 28	LLKLAGDVESNPGP
SEQ ID NO: 29	NFDLLKLAGDVESNPGP
SEQ ID NO: 30	QLLNFDLLKLAGDVESNPGP
SEQ ID NO: 31	APVKQTLNFDLLKLAGDVESNPGP
SEQ ID NO: 32	VTELLYRMKRAETCYCPRLAIHPTEARHKQKIVAPVKQT
SEQ ID NO: 33	LNFDLLKLAGDVESNPGP
SEQ ID NO: 34	LLAIHPTEARHKQKIVAPVKQTLNFDLLKLAGDVESNPGP
SEQ ID NO: 35	EARHKQKIVAPVKQTLNFDLLKLAGDVESNPGP

[0104] A polypeptide contemplated herein may comprise a CAR polypeptide.

E. POLYNUCLEOTIDES

[0105] A polynucleotide encoding one or more CAR polypeptides may be provided, e.g., SEQ ID NO: 10. As used herein, the terms "polynucleotide" or "nucleic acid" refers to messenger RNA (mRNA), RNA, genomic RNA (gRNA), plus strand RNA (RNA(+)), minus strand RNA (RNA(-)), genomic DNA (gDNA), complementary DNA (cDNA) or recombinant DNA. Polynucleotides include single and double stranded polynucleotides. Preferably, polynucleotides include polynucleotides or variants having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to any of the reference sequences described herein (see, e.g., Sequence Listing), typically where the variant maintains at least one biological activity of the reference sequence. Contemplated are polynucleotides comprising expression vectors, viral vectors, and transfer plasmids, and compositions, and cells comprising the same.

[0106] Polynucleotides may be provided that encode at least about 5, 10, 25, 50, 100, 150, 200, 250, 300, 350, 400, 500, 1000, 1250, 1500, 1750, or 2000 or more contiguous amino acid residues of a polypeptide, as well as all intermediate lengths. It will be readily understood that "intermediate lengths," in this context, means any length between the quoted values, such as 6, 7, 8, 9, etc., 101, 102, 103, etc.; 151, 152, 153, etc.; 201, 202, 203, etc.

[0107] As used herein, the terms "polynucleotide variant" and "variant" and the like refer to polynucleotides displaying substantial sequence identity with a reference polynucleotide sequence or polynucleotides that hybridize with a reference sequence under stringent conditions that are defined hereinafter. These terms include polynucleotides in which one or more nucleotides have been added or deleted, or replaced with different nucleotides compared to a reference polynucleotide. In this regard, it is well understood in the art that certain alterations inclusive of mutations, additions, deletions and substitutions can be made to a reference polynucleotide

whereby the altered polynucleotide retains the biological function or activity of the reference polynucleotide.

[0108] The recitations "sequence identity" or, for example, comprising a "sequence 50% identical to," as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a "percentage of sequence identity" may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical

nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. Included are nucleotides and polypeptides having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to any of the reference sequences described herein, typically where the polypeptide variant maintains at least one biological activity of the reference polypeptide.

[0109] Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include "reference sequence," "comparison window," "sequence identity," "percentage of sequence identity," and "substantial identity". A "reference sequence" is at least 12 but frequently 15 to 18 and often at least 25 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (*i.e.*, only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window" refers to a conceptual segment of at least 6 contiguous positions, usually about 50 to about 100, more usually about 100 to about 150 in which a sequence is compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. The comparison window may comprise additions or deletions (*i.e.*, gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, WI, USA) or by inspection and the best alignment (*i.e.*, resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul et al., 1997, Nucl. Acids Res. 25:3389. A detailed discussion of sequence analysis can be found in Unit 19.3 of Ausubel et al., Current Protocols in Molecular Biology, John Wiley & Sons Inc, 1994-1998, Chapter 15.

[0110] As used herein, "isolated polynucleotide" refers to a polynucleotide that has been purified from the sequences which flank it in a naturally-occurring state, e.g., a DNA fragment that has been removed from the sequences that are normally adjacent to the fragment. An "isolated polynucleotide" also refers to a complementary DNA (cDNA), a recombinant DNA, or other polynucleotide that does not exist in nature and that has been made by the hand of man.

[0111] Terms that describe the orientation of polynucleotides include: 5' (normally the end of the polynucleotide having a free phosphate group) and 3' (normally the end of the polynucleotide having a free hydroxyl (OH) group). Polynucleotide sequences can be annotated in the 5' to 3' orientation or the 3' to 5' orientation. For DNA and mRNA, the 5' to 3' strand is designated the "sense," "plus," or "coding" strand because its sequence is identical to the sequence of the premessenger (premRNA) [except for uracil (U) in RNA, instead of thymine (T) in DNA]. For DNA and mRNA, the complementary 3' to 5' strand which is the strand transcribed by the RNA polymerase is designated as "template," "antisense," "minus," or "non-coding" strand. As used herein, the term "reverse orientation" refers to a 5' to 3' sequence written in the 3' to 5' orientation or a 3' to 5' sequence written in the 5' to 3' orientation.

[0112] The terms "complementary" and "complementarity" refer to polynucleotides (*i.e.*, a sequence of nucleotides) related by the base-pairing rules. For example, the complementary strand of the DNA sequence 5' A G T C A T G 3' is 3' T C A G T A C 5'. The latter sequence is often written as the reverse complement with the 5' end on the left and the 3' end on the right, 5' C A T G A C T 3'. A sequence that is equal to its reverse complement is said to be a palindromic sequence. Complementarity can be "partial," in which only some of the nucleic acids' bases are matched according to the base pairing rules. Or, there can be "complete" or "total" complementarity between the nucleic acids.

[0113] Moreover, it will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide, or fragment of variant thereof, as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated, for example polynucleotides that are optimized for human and/or primate codon selection. Further, alleles of the genes comprising the polynucleotide sequences provided herein may also be used. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides.

[0114] The term "nucleic acid cassette" as used herein refers to genetic sequences within a vector which can express a RNA, and subsequently a protein. The nucleic acid cassette contains the gene of interest, e.g., a CAR. The nucleic acid cassette is positionally and sequentially oriented within the vector such that the nucleic acid in the cassette can be transcribed into RNA, and when necessary, translated into a protein or a polypeptide, undergo appropriate post-translational modifications required for activity in the transformed cell, and be translocated to the appropriate compartment for biological activity by targeting to appropriate intracellular compartments or secretion into extracellular compartments.

Preferably, the cassette has its 3' and 5' ends adapted for ready insertion into a vector, *e.g.*, it has restriction endonuclease sites at each end. Preferably, the nucleic acid cassette contains the sequence of a chimeric antigen receptor used to treat a B cell malignancy. The cassette can be removed and inserted into a plasmid or viral vector as a single unit.

[0115] In particular, polynucleotides include at least one polynucleotide-of-interest. As used herein, the term "polynucleotide-of-interest" refers to a polynucleotide encoding a polypeptide (*i.e.*, a polypeptide-of-interest), inserted into an expression vector that is desired to be expressed. A vector may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 polynucleotides-of-interest. The

polynucleotide-of-interest may encode a polypeptide that provides a therapeutic effect in the treatment or prevention of a disease or disorder. Polynucleotides-of-interest, and polypeptides encoded therefrom, include both polynucleotides that encode wild-type polypeptides, as well as functional variants and fragments thereof. In particular embodiments, a functional variant has at least 80%, at least 90%, at least 95%, or at least 99% identity to a corresponding wild-type reference polynucleotide or polypeptide sequence. A functional variant or fragment may have at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% of a biological activity of a corresponding wild-type polypeptide.

[0116] The polynucleotide-of-interest may not encode a polypeptide but serves as a template to transcribe miRNA, siRNA, or shRNA, ribozyme, or other inhibitory RNA. A polynucleotide may comprise a polynucleotide-of-interest encoding a CAR and one or more additional polynucleotides-of-interest including but not limited to an inhibitory nucleic acid sequence including, but not limited to: an siRNA, an miRNA, an shRNA, and a ribozyme.

[0117] As used herein, the terms "siRNA" or "short interfering RNA" refer to a short polynucleotide sequence that mediates a process of sequence-specific post-transcriptional gene silencing, translational inhibition, transcriptional inhibition, or epigenetic RNAi in animals (Zamore et al., 2000, *Cell*, 101, 25-33; Fire et al., 1998, *Nature*, 391, 806; Hamilton et al., 1999, *Science*, 286, 950-951; Lin et al., 1999, *Nature*, 402, 128-129; Sharp, 1999, *Genes & Dev.*, 13, 139-141; and Strauss, 1999, *Science*, 286, 886). An siRNA may comprise a first strand and a second strand that have the same number of nucleosides; however, the first and second strands are offset such that the two terminal nucleosides on the first and second strands are not paired with a residue on the complementary strand. In certain instances, the two nucleosides that are not paired are thymidine residues. The siRNA should include a region of sufficient homology to the target gene, and be of sufficient length in terms of nucleotides, such that the siRNA, or a fragment thereof, can mediate down regulation of the target gene. Thus, an siRNA includes a region which is at least partially complementary to the target RNA. It is not necessary that there be perfect complementarity between the siRNA and the target, but the correspondence must be sufficient to enable the siRNA, or a cleavage product thereof, to direct sequence

specific silencing, such as by RNAi cleavage of the target RNA. Complementarity, or degree of homology with the target strand, is most critical in the antisense strand. While perfect complementarity, particularly in the antisense strand, is often desired, some embodiments include one or more, but preferably 10, 8, 6, 5, 4, 3, 2, or fewer mismatches with respect to the target RNA. The mismatches are most tolerated in the terminal regions, and if present are preferably in a terminal region or regions, *e.g.*, within 6, 5, 4, or 3 nucleotides of the 5' and/or 3' terminus. The sense strand need only be sufficiently complementary with the antisense strand to maintain the overall double-strand character of the molecule.

[0118] In addition, an siRNA may be modified or include nucleoside analogs. Single stranded regions of an siRNA may be modified or include nucleoside analogs, *e.g.*, the unpaired region or regions of a hairpin structure, *e.g.*, a region which links two complementary regions, can have modifications or nucleoside analogs. Modification to stabilize one or more 3'- or 5'-terminus of an siRNA, *e.g.*, against exonucleases, or to favor the antisense siRNA agent to enter into RISC are also useful. Modifications can include C3 (or C6, C7, C12) amino linkers, thiol linkers, carboxyl linkers, non-nucleotidic spacers (C3, C6, C9, C12, abasic, triethylene glycol, hexaethylene glycol), special biotin or fluorescein reagents that come as phosphoramidites and that have another DMT-protected hydroxyl group, allowing multiple couplings during RNA synthesis. Each strand of an siRNA can be equal to or less than 30, 25, 24, 23, 22, 21, or 20 nucleotides in length. The strand is preferably at least 19 nucleotides in length. For example, each strand can be between 21 and 25 nucleotides in length. Preferred siRNAs have a duplex region of 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotide pairs, and one or more overhangs of 2-3 nucleotides, preferably one or two 3' overhangs, of 2-3 nucleotides.

[0119] As used herein, the terms "miRNA" or "microRNA" refer to small non-coding RNAs of 20-22 nucleotides, typically excised from ~70 nucleotide fold-back RNA precursor structures known as pre-miRNAs. miRNAs negatively regulate their targets in one of two ways depending on the degree of complementarity between the miRNA and the target. First, miRNAs that bind with perfect or nearly perfect complementarity to protein-coding mRNA sequences induce the RNA-mediated interference (RNAi) pathway. miRNAs that exert their regulatory effects by binding to imperfect complementary sites within the 3' untranslated regions (UTRs) of their mRNA targets,

repress target-gene expression post-transcriptionally, apparently at the level of translation, through a RISC complex that is similar to, or possibly identical with, the one that is used for the RNAi pathway. Consistent with translational control, miRNAs that use this mechanism reduce the protein levels of their target genes, but the mRNA levels of these genes are only minimally affected. miRNAs encompass both naturally occurring miRNAs as well as artificially designed miRNAs that can specifically target any mRNA sequence. For example, the skilled artisan can design short hairpin RNA

constructs expressed as human miRNA (e.g., miR-30 or miR-21) primary transcripts. This design adds a Drosha processing site to the hairpin construct and has been shown to greatly increase knockdown efficiency (Pusch *et al.*, 2004). The hairpin stem consists of 22-nt of dsRNA (e.g., antisense has perfect complementarity to desired target) and a 15-19-nt loop from a human miR. Adding the miR loop and miR30 flanking sequences on either or both sides of the hairpin results in greater than 10-fold increase in Drosha and Dicer processing of the expressed hairpins when compared with conventional shRNA designs without microRNA. Increased Drosha and Dicer processing translates into greater siRNA/miRNA production and greater potency for expressed hairpins.

[0120] As used herein, the terms "shRNA" or "short hairpin RNA" refer to double-stranded structure that is formed by a single self-complementary RNA strand. shRNA constructs containing a nucleotide sequence identical to a portion, of either coding or non-coding sequence, of the target gene are preferred for inhibition. RNA sequences with insertions, deletions, and single point mutations relative to the target sequence have also been found to be effective for inhibition. Greater than 90% sequence identity, or even 100% sequence identity, between the inhibitory RNA and the portion of the target gene is preferred. The length of the duplex-forming portion of an shRNA may be at least 20, 21 or 22 nucleotides in length, e.g., corresponding in size to RNA products produced by Dicer-dependent cleavage. The shRNA construct may be at least 25, 50, 100, 200, 300 or 400 bases in length. The shRNA construct may be 400-800 bases in length. shRNA constructs are highly tolerant of variation in loop sequence and loop size.

[0121] As used herein, the term "ribozyme" refers to a catalytically active RNA molecule capable of site-specific cleavage of target mRNA. Several subtypes have been described, e.g., hammerhead and hairpin ribozymes. Ribozyme catalytic activity and stability can be improved by substituting deoxyribonucleotides for ribonucleotides at noncatalytic bases. While ribozymes that cleave mRNA at site-specific recognition sequences can be used to destroy particular mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA has the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art.

[0122] A preferred method of delivery of a polynucleotide-of-interest that comprises an siRNA, an miRNA, an shRNA, or a ribozyme comprises one or more regulatory sequences, such as, for example, a strong constitutive pol III, e.g., human U6 snRNA promoter, the mouse U6 snRNA promoter, the human and mouse HI RNA promoter and the human tRNA-val promoter, or a strong constitutive pol II promoter, as described elsewhere herein.

[0123] The polynucleotides regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters and/or enhancers, untranslated regions (UTRs), signal sequences, Kozak sequences, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, internal ribosomal entry sites (IRES), recombinase recognition sites (e.g., LoxP, FRT, and Att sites), termination codons, transcriptional termination signals, and polynucleotides encoding self-cleaving polypeptides, epitope tags, as disclosed elsewhere herein or as known in the art, such that their overall length may vary considerably. It is therefore contemplated that a polynucleotide fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

[0124] Polynucleotides can be prepared, manipulated and/or expressed using any of a variety of well-established techniques known and available in the art. In order to express a desired polypeptide, a nucleotide sequence encoding the polypeptide, can be inserted into appropriate vector. Examples of vectors are plasmid, autonomously replicating sequences, and transposable elements. Additional exemplary vectors include, without limitation, plasmids, phagemids, cosmids, artificial chromosomes such as yeast artificial chromosome (YAC), bacterial artificial chromosome (BAC), or P1-derived artificial chromosome (PAC), bacteriophages such as lambda phage or M13 phage, and animal viruses. Examples of categories of animal viruses useful as vectors include, without limitation, retrovirus (including lentivirus), adenovirus, adeno-associated virus, herpesvirus (e.g., herpes simplex virus), poxvirus, baculovirus, papillomavirus, and papovavirus (e.g., SV40). Examples of expression vectors are pCneo vectors (Promega) for expression in mammalian cells; pLenti4/V5-DEST™, pLenti6/V5-DEST™, and pLenti6.2N5-GW/lacZ (Invitrogen) for lentivirus-mediated gene transfer and expression in mammalian cells. The coding sequences of the chimeric proteins disclosed herein can be ligated into such expression vectors for the expression of the chimeric protein in mammalian cells.

[0125] A vector encoding a CAR contemplated herein may comprise the polynucleotide sequence set forth in SEQ ID NO: 36.

[0126] The vector may be an episomal vector or a vector that is maintained extrachromosomally. As used herein, the term "episomal" refers to a vector that is able to replicate without integration into host's chromosomal DNA and without gradual loss from a dividing host cell also meaning that said vector replicates extrachromosomally or episomally. The vector is engineered to harbor the sequence coding for the origin of DNA replication or "ori" from a lymphotropic herpes virus or a gamma herpesvirus, an adenovirus, SV40, a bovine papilloma virus, or a yeast, specifically a replication origin of a lymphotropic herpes virus or a gamma herpesvirus corresponding to oriP of EBV. In a particular aspect, the lymphotropic herpes virus may be Epstein Barr virus (EBV), Kaposi's sarcoma herpes virus (KSHV), Herpes virus

saimiri (HS), or Marek's disease virus (MDV). Epstein Barr virus (EBV) and Kaposi's sarcoma herpes virus (KSHV) are also examples of a gamma herpesvirus. Typically, the host cell comprises the viral replication transactivator protein that activates the replication.

[0127] The "control elements" or "regulatory sequences" present in an expression vector are those non-translated regions of the vector-origin of replication, selection cassettes, promoters, enhancers, translation initiation signals (Shine Dalgarno sequence or Kozak sequence) introns, a polyadenylation sequence, 5' and 3' untranslated regions-which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including ubiquitous promoters and inducible promoters may be used.

[0128] A vector for use in practicing the present disclosure including, but not limited to expression vectors and viral vectors, will include exogenous, endogenous, or heterologous control sequences such as promoters and/or enhancers. An "endogenous" control sequence is one which is naturally linked with a given gene in the genome. An "exogenous" control sequence is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological techniques) such that transcription of that gene is directed by the linked enhancer/promoter. A heterologous control sequence is an exogenous sequence that is from a different species than the cell being genetically manipulated.

[0129] The term "promoter" as used herein refers to a recognition site of a polynucleotide (DNA or RNA) to which an RNA polymerase binds. An RNA polymerase initiates and transcribes polynucleotides operably linked to the promoter. Promoters operative in mammalian cells may comprise an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated and/or another sequence found 70 to 80 bases upstream from the start of transcription, a CNCAAT region where N may be any nucleotide.

[0130] The term "enhancer" refers to a segment of DNA which contains sequences capable of providing enhanced transcription and in some instances can function independent of their orientation relative to another control sequence. An enhancer can function cooperatively or additively with promoters and/or other enhancer elements. The term "promoter/enhancer" refers to a segment of DNA which contains sequences capable of providing both promoter and enhancer functions.

[0131] The term "operably linked", refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. The term may refer to a functional linkage between a nucleic acid expression control sequence (such as a promoter, and/or enhancer) and a second polynucleotide sequence, *e.g.*, a polynucleotide-of-interest, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

[0132] As used herein, the term "constitutive expression control sequence" refers to a promoter, enhancer, or promoter/enhancer that continually or continuously allows for transcription of an operably linked sequence. A constitutive expression control sequence may be a "ubiquitous" promoter, enhancer, or promoter/enhancer that allows expression in a wide variety of cell and tissue types or a "cell specific," "cell type specific," "cell lineage specific," or "tissue specific" promoter, enhancer, or promoter/enhancer that allows expression in a restricted variety of cell and tissue types, respectively.

[0133] Illustrative ubiquitous expression control sequences suitable for use include, but are not limited to, a cytomegalovirus (CMV) immediate early promoter, a viral simian virus 40 (SV40) (*e.g.*, early or late), a Moloney murine leukemia virus (MoMLV) LTR promoter, a Rous sarcoma virus (RSV) LTR, a herpes simplex virus (HSV) (thymidine kinase) promoter, H5, P7.5, and P11 promoters from vaccinia virus, an elongation factor 1-alpha (EF1a) promoter, early growth response 1 (EGR1), ferritin H (FerH), ferritin L (FerL), Glyceraldehyde 3-phosphate dehydrogenase (GAPDH), eukaryotic translation initiation factor 4A1 (EIF4A1), heat shock 70kDa protein 5 (HSPA5), heat shock protein 90kDa beta, member 1 (HSP90B1), heat shock protein 70kDa (HSP70), β -kinesin (β -KIN), the human ROSA 26 locus (Irions et al., Nature Biotechnology 25, 1477 - 1482 (2007)), a Ubiquitin C promoter (UBC), a phosphoglycerate kinase-1 (PGK) promoter, a cytomegalovirus enhancer/chicken β -actin (CAG) promoter, a β -actin promoter and a myeloproliferative sarcoma virus enhancer, negative control region deleted, dl587rev primer-binding site substituted (MND) promoter (Challita et al., J Virol. 69(2):748-55 (1995)).

[0134] A vector may comprise a MND promoter.

[0135] A vector may comprise an EF1a promoter comprising the first intron of the human EF1a gene.

[0136] A vector may comprise an EF1a promoter that lacks the first intron of the human EF1a gene.

[0137] It may be desirable to express a polynucleotide comprising a CAR from a T cell specific promoter.

[0138] As used herein, "conditional expression" may refer to any type of conditional expression including, but not limited to, inducible expression; repressible expression; expression in cells or tissues having a particular physiological, biological, or disease state, *etc.* This definition is not intended to exclude cell type or tissue specific expression. Provided is conditional expression of a polynucleotide-of-interest, *e.g.*, expression is controlled by subjecting a cell, tissue, organism, *etc.*, to a treatment or condition that causes the polynucleotide to be expressed or that causes an increase or decrease in expression of the polynucleotide encoded by the polynucleotide-of-interest.

[0139] Illustrative examples of inducible promoters/systems include, but are not limited to, steroid-inducible promoters such as promoters for genes encoding glucocorticoid or estrogen receptors (inducible by treatment with the corresponding hormone), metallothionein promoter (inducible by treatment with various heavy metals), MX-1 promoter (inducible by interferon), the "GeneSwitch" mifepristone-regulatable system (Sirin et al., 2003, *Gene*, 323:67), the cumate inducible gene switch (WO 2002/088346), tetracycline-dependent regulatory systems, etc.

[0140] Conditional expression can also be achieved by using a site specific DNA recombinase. The vector may comprise at least one (typically two) site(s) for recombination mediated by a site specific recombinase. As used herein, the terms "recombinase" or "site specific recombinase" include excisive or integrative proteins, enzymes, co-factors or associated proteins that are involved in recombination reactions involving one or more recombination sites (e.g., two, three, four, five, seven, ten, twelve, fifteen, twenty, thirty, fifty, etc.), which may be wild-type proteins (see Landy, *Current Opinion in Biotechnology* 3:699-707 (1993)), or mutants, derivatives (e.g., fusion proteins containing the recombination protein sequences or fragments thereof), fragments, and variants thereof. Illustrative examples of recombinases suitable for use include, but are not limited to: Cre, Int, IHF, Xis, Fli, Fis, Hin, Gin, ϕ C31, Cin, Tn3 resolvase, TndX, XerC, XerD, TnpX, Hjc, Gin, SpCCE1, and ParA.

[0141] The vectors may comprise one or more recombination sites for any of a wide variety of site specific recombinases. It is to be understood that the target site for a site specific recombinase is in addition to any site(s) required for integration of a vector, e.g., a retroviral vector or lentiviral vector. As used herein, the terms "recombination sequence," "recombination site," or "site specific recombination site" refer to a particular nucleic acid sequence to which a recombinase recognizes and binds.

[0142] For example, one recombination site for Cre recombinase is loxP which is a 34 base pair sequence comprising two 13 base pair inverted repeats (serving as the recombinase binding sites) flanking an 8 base pair core sequence (see FIG. 1 of Sauer, B., *Current Opinion in Biotechnology* 5:521-527 (1994)). Other exemplary loxP sites include, but are not limited to: lox511 (Hoess et al., 1996; Bethke and Sauer, 1997), lox5171 (Lee and Saito, 1998), lox2272 (Lee and Saito, 1998), m2 (Langer et al., 2002), lox71 (Albert et al., 1995), and lox66 (Albert et al., 1995).

[0143] Suitable recognition sites for the FLP recombinase include, but are not limited to: FRT (McLeod, et al., 1996), F₁, F₂, F₃ (Schlake and Bode, 1994), F₄, F₅ (Schlake and Bode, 1994), FRT(LE) (Senecoff et al., 1988), FRT(RE) (Senecoff et al., 1988).

[0144] Other examples of recognition sequences are the attB, attP, attL, and attR sequences, which are recognized by the recombinase enzyme λ Integrase, e.g., ϕ C31. The ϕ C31 SSR mediates recombination only between the heterotypic sites attB (34 bp in length) and attP (39 bp in length) (Groth et al., 2000). attB and attP, named for the attachment sites for the phage integrase on the bacterial and phage genomes, respectively, both contain imperfect inverted repeats that are likely bound by ϕ C31 homodimers (Groth et al., 2000). The product sites, attL and attR, are effectively inert to further ϕ C31-mediated recombination (Belteki et al., 2003), making the reaction irreversible. For catalyzing insertions, it has been found that attB-bearing DNA inserts into a genomic attP site more readily than an attP site into a genomic attB site (Thyagarajan et al., 2001; Belteki et al., 2003). Thus, typical strategies position by homologous recombination an attP-bearing "docking site" into a defined locus, which is then partnered with an attB-bearing incoming sequence for insertion.

[0145] As used herein, an "internal ribosome entry site" or "IRES" refers to an element that promotes direct internal ribosome entry to the initiation codon, such as ATG, of a cistron (a protein encoding region), thereby leading to the cap-independent translation of the gene. See, e.g., Jackson et al., 1990. *Trends Biochem Sci* 15(12):477-83) and Jackson and Kaminski. 1995. *RNA* 1(10):985-1000. The vectors may include one or more polynucleotides-of-interest that encode one or more polypeptides. To achieve efficient translation of each of the plurality of polypeptides, the polynucleotide sequences may be separated by one or more IRES sequences or polynucleotide sequences encoding self-cleaving polypeptides.

[0146] As used herein, the term "Kozak sequence" refers to a short nucleotide sequence that greatly facilitates the initial binding of mRNA to the small subunit of the ribosome and increases translation. The consensus Kozak sequence is (GCC)RCCATGG, where R is a purine (A or G) (Kozak, 1986. *Cell*. 44(2):283-92, and Kozak, 1987. *Nucleic Acids Res.* 15(20):8125-48). The vectors may comprise polynucleotides that have a consensus Kozak sequence and that encode a desired polypeptide, e.g., a CAR.

[0147] A polynucleotide or cell harboring the polynucleotide may utilize a suicide gene, including an inducible suicide gene to reduce the risk of direct toxicity and/or uncontrolled proliferation. The suicide gene may not be immunogenic to the host harboring the polynucleotide or cell. A certain example of a suicide gene that may be used is caspase-9 or caspase-8 or cytosine deaminase. Caspase-9 can be activated using a specific chemical inducer of dimerization (CID).

[0148] The vectors may comprise gene segments that cause the immune effector cells, e.g., T cells, to be susceptible to negative selection in vivo. By "negative selection" is meant that the infused cell can be eliminated as a result of a change in the in vivo condition of the individual. The negative selectable phenotype may result from the insertion of a gene that confers sensitivity to an administered agent, for example, a compound. Negative selectable genes are known in the art, and include, inter alia the following: the Herpes simplex virus type I thymidine kinase (HSV-I TK) gene (Wigler

et al., Cell 11:223, 1977) which confers ganciclovir sensitivity; the cellular hypoxanthine phosphoribosyltransferase (HPRT) gene, the cellular adenine phosphoribosyltransferase (APRT) gene, and bacterial cytosine deaminase, (Mullen et al., Proc. Natl. Acad. Sci. USA. 89:33 (1992)).

[0149] Genetically modified immune effector cells, such as T cells, may comprise a polynucleotide further comprising a positive marker that enables the selection of cells of the negative selectable phenotype *in vitro*. The positive selectable marker may be a gene which, upon being introduced into the host cell expresses a dominant phenotype permitting positive selection of cells carrying the gene. Genes of this type are known in the art, and include, inter alia, hygromycin-B phosphotransferase gene (hph) which confers resistance to hygromycin B, the amino glycoside phosphotransferase gene (neo or aph) from Tn5 which codes for resistance to the antibiotic G418, the dihydrofolate reductase (DHFR) gene, the adenosine deaminase gene (ADA), and the multi-drug resistance (MDR) gene.

[0150] Preferably, the positive selectable marker and the negative selectable element are linked such that loss of the negative selectable element necessarily also is accompanied by loss of the positive selectable marker. Even more preferably, the positive and negative selectable markers are fused so that loss of one obligatorily leads to loss of the other. An example of a fused polynucleotide that yields as an expression product a polypeptide that confers both the desired positive and negative selection features described above is a hygromycin phosphotransferase thymidine kinase fusion gene (HyTK). Expression of this gene yields a polypeptide that confers hygromycin B resistance for positive selection *in vitro*, and ganciclovir sensitivity for negative selection *in vivo*. See Lupton S. D., et al, Mol. and Cell. Biology 11:3374- 3378, 1991. In addition, the polynucleotides encoding the chimeric receptors may be in retroviral vectors containing the fused gene, particularly those that confer hygromycin B resistance for positive selection *in vitro*, and ganciclovir sensitivity for negative selection *in vivo*, for example the HyTK retroviral vector described in Lupton, S. D. et al. (1991), *supra*. See also the publications of PCT US91/08442 and PCT/US94/05601, by S. D. Lupton, describing the use of bifunctional selectable fusion genes derived from fusing a dominant positive selectable markers with negative selectable markers.

[0151] Preferred positive selectable markers are derived from genes selected from the group consisting of hph, nco, and gpt, and preferred negative selectable markers are derived from genes selected from the group consisting of cytosine deaminase, HSV-I TK, VZV TK, HPRT, APRT and gpt. Especially preferred markers are bifunctional selectable fusion genes wherein the positive selectable marker is derived from hph or neo, and the negative selectable marker is derived from cytosine deaminase or a TK gene or selectable marker. Inducible Suicide Genes

F. VIRAL VECTORS

[0152] A cell (*e.g.*, an immune effector cell) may be transduced with a retroviral vector, *e.g.*, a lentiviral vector, encoding a CAR. For example, an immune effector cell is transduced with a vector encoding a CAR that comprises a murine anti-BCMA antibody or antigen binding fragment thereof that binds a BCMA polypeptide, with an intracellular signaling domain of CD3 ζ , CD28, 4-1BB, Ox40, or any combinations thereof. Thus, these transduced cells can elicit a CAR-mediated cytotoxic response.

[0153] Retroviruses are a common tool for gene delivery (Miller, 2000, Nature. 357: 455-460). A retrovirus may be used to deliver a polynucleotide encoding a chimeric antigen receptor (CAR) to a cell. As used herein, the term "retrovirus" refers to an RNA virus that reverse transcribes its genomic RNA into a linear double-stranded DNA copy and subsequently covalently integrates its genomic DNA into a host genome. Once the virus is integrated into the host genome, it is referred to as a "provirus." The provirus serves as a template for RNA polymerase II and directs the expression of RNA molecules which encode the structural proteins and enzymes needed to produce new viral particles.

[0154] Illustrative retroviruses suitable for use, include, but are not limited to: Moloney murine leukemia virus (M-MuLV), Moloney murine sarcoma virus (MoMSV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), gibbon ape leukemia virus (GaLV), feline leukemia virus (FLV), spumavirus, Friend murine leukemia virus, Murine Stem Cell Virus (MSCV) and Rous Sarcoma Virus (RSV) and lentivirus.

[0155] As used herein, the term "lentivirus" refers to a group (or genus) of complex retroviruses. Illustrative lentiviruses include, but are not limited to: HIV (human immunodeficiency virus; including HIV type 1, and HIV type 2); visna-maedi virus (VMV) virus; the caprine arthritis-encephalitis virus (CAEV); equine infectious anemia virus (EIAV); feline immunodeficiency virus (FIV); bovine immune deficiency virus (BIV); and simian immunodeficiency virus (SIV). In one embodiment, HIV based vector backbones (*i.e.*, HIV cis-acting sequence elements) are preferred. A lentivirus may be used to deliver a polynucleotide comprising a CAR to a cell.

[0156] Retroviral vectors and more particularly lentiviral vectors may be used. Accordingly, the term "retrovirus" or "retroviral vector", as used herein is meant to include "lentivirus" and "lentiviral vectors" respectively.

[0157] The term "vector" is used herein to refer to a nucleic acid molecule capable transferring or transporting another nucleic acid molecule. The transferred nucleic acid is generally linked to, *e.g.*, inserted into, the vector nucleic acid molecule. A vector may include sequences that direct autonomous replication in a cell, or may include sequences sufficient to allow integration into host cell DNA. Useful vectors include, for example, plasmids (*e.g.*, DNA plasmids or

RNA plasmids), transposons, cosmids, bacterial artificial chromosomes, and viral vectors. Useful viral vectors include, e.g., replication defective retroviruses and lentiviruses.

[0158] As will be evident to one of skill in the art, the term "viral vector" is widely used to refer either to a nucleic acid molecule (e.g., a transfer plasmid) that includes virus-derived nucleic acid elements that typically facilitate transfer of the nucleic acid molecule or integration into the genome of a cell or to a viral particle that mediates nucleic acid transfer. Viral particles will typically include various viral components and sometimes also host cell components in addition to nucleic acid(s).

[0159] The term viral vector may refer either to a virus or viral particle capable of transferring a nucleic acid into a cell or to the transferred nucleic acid itself. Viral vectors and transfer plasmids contain structural and/or functional genetic elements that are primarily derived from a virus. The term "retroviral vector" refers to a viral vector or plasmid containing structural and functional genetic elements, or portions thereof, that are primarily derived from a retrovirus. The term "lentiviral vector" refers to a viral vector or plasmid containing structural and functional genetic elements, or portions thereof, including LTRs that are primarily derived from a lentivirus. The term "hybrid vector" refers to a vector, LTR or other nucleic acid containing both retroviral, e.g., lentiviral, sequences and non-lentiviral viral sequences. A hybrid vector may refer to a vector or transfer plasmid comprising retroviral e.g., lentiviral, sequences for reverse transcription, replication, integration and/or packaging.

[0160] The terms "lentiviral vector," "lentiviral expression vector" may be used to refer to lentiviral transfer plasmids and/or infectious lentiviral particles. Where reference is made herein to elements such as cloning sites, promoters, regulatory elements, heterologous nucleic acids, etc., it is to be understood that the sequences of these elements are present in RNA form in the lentiviral particles and are present in DNA form in the DNA plasmids.

[0161] At each end of the provirus are structures called "long terminal repeats" or "LTRs." The term "long terminal repeat (LTR)" refers to domains of base pairs located at the ends of retroviral DNAs which, in their natural sequence context, are direct repeats and contain U3, R and U5 regions. LTRs generally provide functions fundamental to the expression of retroviral genes (e.g., promotion, initiation and polyadenylation of gene transcripts) and to viral replication.

The LTR contains

numerous regulatory signals including transcriptional control elements, polyadenylation signals and sequences needed for replication and integration of the viral genome. The viral LTR is divided into three regions called U3, R and U5. The U3 region contains the enhancer and promoter elements. The U5 region is the sequence between the primer binding site and the R region and contains the polyadenylation sequence. The R (repeat) region is flanked by the U3 and U5 regions. The LTR composed of U3, R and U5 regions and appears at both the 5' and 3' ends of the viral genome. Adjacent to the 5' LTR are sequences necessary for reverse transcription of the genome (the tRNA primer binding site) and for efficient packaging of viral RNA into particles (the Psi site).

[0162] As used herein, the term "packaging signal" or "packaging sequence" refers to sequences located within the retroviral genome which are required for insertion of the viral RNA into the viral capsid or particle, see e.g., Clever et al., 1995. J. of Virology, Vol. 69, No. 4; pp. 2101-2109. Several retroviral vectors use the minimal packaging signal (also referred to as the psi [Ψ] sequence) needed for encapsidation of the viral genome. Thus, as used herein, the terms "packaging sequence," "packaging signal," "psi" and the symbol "Ψ," are used in reference to the non-coding sequence required for encapsidation of retroviral RNA strands during viral particle formation.

[0163] Vectors may comprise modified 5' LTR and/or 3' LTRs. Either or both of the LTR may comprise one or more modifications including, but not limited to, one or more deletions, insertions, or substitutions. Modifications of the 3' LTR are often made to improve the safety of lentiviral or retroviral systems by rendering viruses replication-defective. As used herein, the term "replication-defective" refers to virus that is not capable of complete, effective replication such that infective virions are not produced (e.g., replication-defective lentiviral progeny). The term "replication-competent" refers to wild-type virus or mutant virus that is capable of replication, such that viral replication of the virus is capable of producing infective virions (e.g., replication-competent lentiviral progeny).

[0164] "Self-inactivating" (SIN) vectors refers to replication-defective vectors, e.g., retroviral or lentiviral vectors, in which the right (3') LTR enhancer-promoter region, known as the U3 region, has been modified (e.g., by deletion or substitution) to prevent viral transcription beyond the first round of viral replication. This is because the right (3') LTR U3 region is used as a template for the left (5') LTR U3 region during viral replication and, thus, the viral transcript cannot be made without the U3 enhancer-promoter. The 3' LTR may be modified such that the U5 region is replaced, for example, with an ideal poly(A) sequence. It should be noted that modifications to the LTRs such as modifications to the 3' LTR, the 5' LTR, or both 3' and 5' LTRs, are also included.

[0165] An additional safety enhancement is provided by replacing the U3 region of the 5' LTR with a heterologous promoter to drive transcription of the viral genome during production of viral particles. Examples of heterologous promoters which can be used include, for example, viral simian virus 40 (SV40) (e.g., early or late), cytomegalovirus (CMV) (e.g., immediate early), Moloney murine leukemia virus (MoMLV), Rous sarcoma virus (RSV), and herpes simplex virus (HSV) (thymidine kinase) promoters. Typical promoters are able to drive high levels of transcription in a Tat-independent manner. This replacement reduces the possibility of recombination to generate replication-competent virus because

there is no complete U3 sequence in the virus production system. The heterologous promoter may have additional advantages in controlling the manner in which the viral genome is transcribed. For example, the heterologous promoter can be inducible, such that transcription of all or part of the viral genome will occur only when the induction factors are present. Induction factors include, but are not limited to, one or more chemical compounds or the physiological conditions such as temperature or pH, in which the host cells are cultured.

[0166] Viral vectors may comprise a TAR element. The term "TAR" refers to the "trans-activation response" genetic element located in the R region of lentiviral (e.g., HIV) LTRs. This element interacts with the lentiviral trans-activator (tat) genetic element to enhance viral replication. However, this element is not required in embodiments wherein the U3 region of the 5' LTR is replaced by a heterologous promoter.

[0167] The "R region" refers to the region within retroviral LTRs beginning at the start of the capping group (*i.e.*, the start of transcription) and ending immediately prior to the start of the poly A tract. The R region is also defined as being flanked by the U3 and U5 regions. The R region plays a role during reverse transcription in permitting the transfer of nascent DNA from one end of the genome to the other.

[0168] As used herein, the term "FLAP element" refers to a nucleic acid whose sequence includes the central polypurine tract and central termination sequences (cPPT and CTS) of a retrovirus, *e.g.*, HIV-1 or HIV-2. Suitable FLAP elements are described in U.S. Pat. No. 6,682,907 and in Zennou, et al., 2000, Cell, 101:173. During HIV-1 reverse transcription, central initiation of the plus-strand DNA at the central polypurine tract (cPPT) and central termination at the central termination sequence (CTS) lead to the formation of a three-stranded DNA structure: the HIV-1 central DNA flap. While not wishing to be bound by any theory, the DNA flap may act as a cis-active determinant of lentiviral genome nuclear import and/or may increase the titer of the virus. The retroviral or lentiviral vector backbones may comprise one or more FLAP elements upstream or downstream of the heterologous genes of interest in the vectors. For example, a transfer plasmid may include a FLAP element. A vector may comprise a FLAP element isolated from HIV-1.

[0169] Retroviral or lentiviral transfer vectors may comprise one or more export elements. The term "export element" refers to a cis-acting post-transcriptional regulatory element which regulates the transport of an RNA transcript from the nucleus to the cytoplasm of a cell. Examples of RNA export elements include, but are not limited to, the human immunodeficiency virus (HIV) rev response element (RRE) (*see e.g.*, Cullen et al., 1991, J. Virol. 65: 1053; and Cullen et al., 1991, Cell 58: 423), and the hepatitis B virus post-transcriptional regulatory element (HPRE). Generally, the RNA export element is placed within the 3' UTR of a gene, and can be inserted as one or multiple copies.

[0170] Expression of heterologous sequences in viral vectors may be increased by incorporating posttranscriptional regulatory elements, efficient polyadenylation sites, and optionally, transcription termination signals into the vectors. A variety of posttranscriptional regulatory elements can increase expression of a heterologous nucleic acid at the protein, *e.g.*, woodchuck hepatitis virus posttranscriptional regulatory element (WPRE; Zufferey et al., 1999, J. Virol., 73:2886); the posttranscriptional regulatory element present in hepatitis B virus (HPRE) (Huang et al., Mol. Cell. Biol., 5:3864); and the like (Liu et al., 1995, Genes Dev., 9:1766). Vectors may comprise a posttranscriptional regulatory element such as a WPRE or HPRE. Vectors may lack or do not comprise a posttranscriptional regulatory element (PTE) such as a WPRE or HPRE because in some instances these elements increase the risk of cellular transformation and/or do not substantially or significantly increase the amount of mRNA transcript or increase mRNA stability. Therefore, vectors may lack or do not comprise a PTE. In other embodiments, vectors lack or do not comprise a WPRE or HPRE as an added safety measure.

[0171] Elements directing the efficient termination and polyadenylation of the heterologous nucleic acid transcripts increases heterologous gene expression. Transcription termination signals are generally found downstream of the polyadenylation signal. Vectors may comprise a polyadenylation sequence 3' of a polynucleotide encoding a polypeptide to be expressed. The term "polyA site" or "polyA sequence" as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript by RNA polymerase II. Polyadenylation sequences can promote mRNA stability by addition of a polyA tail to the 3' end of the coding sequence and thus, contribute to increased translational efficiency. Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. Illustrative examples of polyA signals that can be used in a vector, includes an ideal polyA sequence (*e.g.*, AATAAA, ATATAA, AGTAAA), a bovine growth hormone polyA sequence (BGHPA), a rabbit β -globin polyA sequence (r β gpA), or another suitable heterologous or endogenous polyA sequence known in the art.

[0172] A retroviral or lentiviral vector may further comprise one or more insulator elements. Insulator elements may contribute to protecting lentivirus-expressed sequences, *e.g.*, therapeutic polypeptides, from integration site effects, which may be mediated by cis-acting elements present in genomic DNA and lead to deregulated expression of transferred sequences (*i.e.*, position effect; *see, e.g.*, Burgess-Beusse et al., 2002, Proc. Natl. Acad. Sci., USA, 99:16433; and Zhan et al., 2001, Hum. Genet., 109:471). Transfer vectors may comprise one or more insulator element the 3' LTR and upon integration of the provirus into the host genome, the provirus comprises the one or more insulators at both the 5' LTR or 3' LTR, by virtue of duplicating the 3' LTR. Suitable insulators for use include, but are not limited to, the chicken β -globin insulator (*see* Chung et al., 1993, Cell 74:505; Chung et al., 1997, PNAS 94:575; and Bell et al., 1999, Cell

98:387,). Examples of insulator elements include, but are not limited to, an insulator from an β -globin locus, such as chicken HS4.

[0173] Most or all of the viral vector backbone sequences may be derived from a lentivirus, e.g., HIV-1. However, it is to be understood that many different sources of retroviral and/or lentiviral sequences can be used, or combined and numerous substitutions and alterations in certain of the lentiviral sequences may be accommodated without impairing the ability of a transfer vector to perform the functions described herein. Moreover, a variety of lentiviral vectors are known in the art, see Naldini *et al.*, (1996a, 1996b, and 1998); Zufferey *et al.*, (1997); Dull *et al.*, 1998, U.S. Pat. Nos. 6,013,516; and 5,994,136, many of which may be adapted to produce a viral vector or transfer plasmid.

[0174] The vectors may comprise a promoter operably linked to a polynucleotide encoding a CAR polypeptide. The vectors may have one or more LTRs, wherein either LTR comprises one or more modifications, such as one or more nucleotide substitutions, additions, or deletions. The vectors may further comprise one of more accessory elements to increase transduction efficiency (e.g., a cPPT/FLAP), viral packaging (e.g., a Psi (Ψ) packaging signal, RRE), and/or other elements that increase therapeutic gene expression (e.g., poly (A) sequences), and may optionally comprise a WPRE or HPRE.

[0175] The transfer vector may comprise a left (5') retroviral LTR; a central polypurine tract/DNA flap (cPPT/FLAP); a retroviral export element; a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; and a right (3') retroviral LTR; and optionally a WPRE or HPRE.

[0176] The transfer vector may comprise a left (5') retroviral LTR; a retroviral export element; a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; a right (3') retroviral LTR; and a poly (A) sequence; and optionally a WPRE or HPRE. Also provided is a lentiviral vector comprising: a left (5') LTR; a cPPT/FLAP; an RRE; a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; a right (3') LTR; and a polyadenylation sequence; and optionally a WPRE or HPRE.

[0177] Also provided is a lentiviral vector comprising: a left (5') HIV-1 LTR; a Psi (Ψ) packaging signal; a cPPT/FLAP; an RRE; a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; a right (3') self-inactivating (SIN) HIV-1 LTR; and a rabbit β -globin polyadenylation sequence; and optionally a WPRE or HPRE.

[0178] Also provided is a vector comprising: at least one LTR; a central polypurine tract/DNA flap (cPPT/FLAP); a retroviral export element; and a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; and optionally a WPRE or HPRE.

[0179] Also provided is a vector comprising at least one LTR; a cPPT/FLAP; an RRE; a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; and a polyadenylation sequence; and optionally a WPRE or HPRE.

[0180] Also provided is at least one SIN HIV-1 LTR; a Psi (Ψ) packaging signal; a cPPT/FLAP; an RRE; a promoter active in a T cell, operably linked to a polynucleotide encoding CAR polypeptide contemplated herein; and a rabbit β -globin polyadenylation sequence; and optionally a WPRE or HPRE.

[0181] The vector may be an integrating viral vector.

[0182] The vector may be an episomal or non-integrating viral vector.

[0183] Vectors contemplated herein, may comprise non-integrating or integration defective retrovirus. An "integration defective" retrovirus or lentivirus may refer to retrovirus or lentivirus having an integrase that lacks the capacity to integrate the viral genome into the genome of the host cells. The integrase protein may be mutated to specifically decrease its integrase activity. Integration-incompetent lentiviral vectors are obtained by modifying the pol gene encoding the integrase protein, resulting in a mutated pol gene encoding an integrative deficient integrase. Such integration-incompetent viral vectors have been described in patent application WO 2006/010834. Illustrative mutations in the HIV-1 pol gene suitable to reduce integrase activity include, but are not limited to: H12N, H12C, H16C, H16V, S81 R, D41A, K42A, H51A, Q53C, D55V, D64E, D64V, E69A, K71A, E85A, E87A, D116N, D116I, D116A, N120G, N120I, N120E, E152G, E152A, D35E, K156E, K156A, E157A, K159E, K159A, K160A, R166A, D167A, E170A, H171A, K173A, K186Q, K186T, K188T, E198A, R199c, R199T, R199A, D202A, K211A, Q214L, Q216L, Q221 L, W235F, W235E, K236S, K236A, K246A, G247W, D253A, R262A, R263A and K264H.

[0184] Illustrative mutations in the HIV-1 pol gene suitable to reduce integrase activity include, but are not limited to: D64E, D64V, E92K, D116N, D116I, D116A, N120G, N120I, N120E, E152G, E152A, D35E, K156E, K156A, E157A, K159E, K159A, W235F, and W235E.

[0185] An integrase may comprise a mutation in one or more of amino acids, D64, D116 or E152. An integrase may comprise a mutation in the amino acids, D64, D116 and E152. A defective HIV-1 integrase may comprise a D64V mutation.

[0186] A "host cell" includes cells electroporated, transfected, infected, or transduced *in vivo*, *ex vivo*, or *in vitro* with a recombinant vector or a polynucleotide. Host cells may include packaging cells, producer cells, and cells infected with viral vectors. Host cells infected with viral vector of the invention are administered to a subject in need of therapy. The term "target cell" may be used interchangeably with host cell and refers to transfected, infected, or transduced cells of a desired cell type. The target cell may be a T cell.

[0187] Large scale viral particle production is often necessary to achieve a reasonable viral titer. Viral particles are produced by transfecting a transfer vector into a packaging cell line that comprises viral structural and/or accessory genes, e.g., gag, pol, env, tat, rev, vif, vpr, vpu, vpx, or nef genes or other retroviral genes.

[0188] As used herein, the term "packaging vector" refers to an expression vector or viral vector that lacks a packaging signal and comprises a polynucleotide encoding one, two, three, four or more viral structural and/or accessory genes. Typically, the packaging vectors are included in a packaging cell, and are introduced into the cell via transfection, transduction or infection. Methods for transfection, transduction or infection are well known by those of skill in the art. A retroviral/lentiviral transfer vector can be introduced into a packaging cell line, via transfection, transduction or infection, to generate a producer cell or cell line. The

packaging vectors can be introduced into human cells or cell lines by standard methods including, e.g., calcium phosphate transfection, lipofection or electroporation. The packaging vectors may be introduced into the cells together with a dominant selectable marker, such as neomycin, hygromycin, puromycin, blastocidin, zeocin, thymidine kinase, DHFR, Gln synthetase or ADA, followed by selection in the presence of the appropriate drug and isolation of clones. A selectable marker gene can be linked physically to genes encoding by the packaging vector, e.g., by IRES or self-cleaving viral peptides.

[0189] Viral envelope proteins (env) determine the range of host cells which can ultimately be infected and transformed by recombinant retroviruses generated from the cell lines. In the case of lentiviruses, such as HIV-1, HIV-2, SIV, FIV and EIV, the env proteins include gp41 and gp120. Preferably, the viral env proteins expressed by packaging cells are encoded on a separate vector from the viral gag and pol genes, as has been previously described.

[0190] Illustrative examples of retroviral-derived env genes which can be employed include, but are not limited to: MLV envelopes, 10A1 envelope, BAEV, FeLV-B, RD114, SSAV, Ebola, Sendai, FPV (Fowl plague virus), and influenza virus envelopes. Similarly, genes encoding envelopes from RNA viruses (e.g., RNA virus families of Picornaviridae, Calciviridae, Astroviridae, Togaviridae, Flaviviridae, Coronaviridae, Paramyxoviridae, Rhabdoviridae, Filoviridae, Orthomyxoviridae, Bunyaviridae, Arenaviridae, Reoviridae, Birnaviridae, Retroviridae) as well as from the DNA viruses (families of Hepadnaviridae, Circoviridae, Parvoviridae, Papovaviridae, Adenoviridae, Herpesviridae, Poxviridae, and Iridoviridae) may be utilized. Representative examples include, FeLV, VEE, HFVW, WDSV, SFV, Rabies, ALV, BIV, BLV, EBV, CAEV, SNV, ChTLV, STLV, MPMV, SMRV, RAV, FuSV, MH2, AEV, AMV, CT10, and EIAV.

[0191] Envelope proteins for pseudotyping a virus include, but are not limited to any of the following virus: Influenza A such as H1N1, H1N2, H3N2 and H5N1 (bird flu), Influenza B, Influenza C virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Hepatitis D virus, Hepatitis E virus, Rotavirus, any virus of the Norwalk virus group, enteric adenoviruses, parvovirus, Dengue fever virus, Monkey pox, Mononegavirales, Lyssavirus such as rabies virus, Lagos bat virus, Mokola virus, Duvenhage virus, European bat virus 1 & 2 and

Australian bat virus, Ephemerovirus, Vesiculovirus, Vesicular Stomatitis Virus (VSV), Herpesviruses such as Herpes simplex virus types 1 and 2, varicella zoster, cytomegalovirus, Epstein-Bar virus (EBV), human herpesviruses (HHV), human herpesvirus type 6 and 8, Human immunodeficiency virus (HIV), papilloma virus, murine gammaherpesvirus, Arenaviruses such as Argentine hemorrhagic fever virus, Bolivian hemorrhagic fever virus, Sabia-associated hemorrhagic fever virus, Venezuelan hemorrhagic fever virus, Lassa fever virus, Machupo virus, Lymphocytic choriomeningitis virus (LCMV), Bunyaviridae such as Crimean-Congo hemorrhagic fever virus, Hantavirus, hemorrhagic fever with renal syndrome causing virus, Rift Valley fever virus, Filoviridae (filovirus) including Ebola hemorrhagic fever and Marburg hemorrhagic fever, Flaviviridae including Kaysanur Forest disease virus, Omsk hemorrhagic fever virus, Tick-borne encephalitis causing virus and Paramyxoviridae such as Hendra virus and Nipah virus, variola major and variola minor (smallpox), alphaviruses such as Venezuelan equine encephalitis virus, eastern equine encephalitis virus, western equine encephalitis virus, SARS-associated coronavirus (SARS-CoV), West Nile virus, any encephalitis causing virus.

[0192] Provided are packaging cells which produce recombinant retrovirus, e.g., lentivirus, pseudotyped with the VSV-G glycoprotein.

[0193] The terms "pseudotype" or "pseudotyping" as used herein, refer to a virus whose viral envelope proteins have been substituted with those of another virus possessing preferable characteristics. For example, HIV can be pseudotyped with vesicular stomatitis virus G-protein (VSV-G) envelope proteins, which allows HIV to infect a wider range of cells because HIV envelope proteins (encoded by the env gene) normally target the virus to CD4⁺ presenting cells. Lentiviral envelope proteins are pseudotyped with VSV-G. Provided are packaging cells which produce recombinant retrovirus, e.g., lentivirus, pseudotyped with the VSV-G envelope glycoprotein.

[0194] As used herein, the term "packaging cell lines" is used in reference to cell lines that do not contain a packaging signal, but do stably or transiently express viral structural proteins and replication enzymes (e.g., gag, pol and env) which are necessary for the correct packaging of viral particles. Any suitable cell line can be employed to prepare packaging cells. Generally, the cells are mammalian cells. The cells used to produce the packaging cell line may be human cells. Suitable cell lines which can be used include, for example, CHO cells, BHK cells, MDCK cells, C3H 10T1/2 cells, FLY cells, Psi-2 cells, BOSC 23 cells, PA317 cells, WEHI cells, COS cells, BSC 1 cells, BSC 40 cells, BMT 10 cells, VERO cells, W138 cells, MRC5 cells, A549 cells, HT1080 cells, 293 cells, 293T cells, B-50 cells, 3T3 cells, NIH3T3 cells,

HepG2 cells, Saos-2 cells, Huh7 cells, HeLa cells, W163 cells, 211 cells, and 211A cells. The packaging cells are preferably 293 cells, 293T cells, or A549 cells. The cells are preferably may be A549 cells.

[0195] As used herein, the term "producer cell line" refers to a cell line which is capable of producing recombinant retroviral particles, comprising a packaging cell line and a transfer vector construct comprising a packaging signal. The production of infectious viral particles and viral stock solutions may be carried out using conventional techniques. Methods of preparing viral stock solutions are known in the art and are illustrated by, e.g., Y. Soneoka et al. (1995) Nucl. Acids Res. 23:628-633, and N. R. Landau et al. (1992) J. Virol. 66:5110-5113. Infectious virus particles may be collected from the packaging cells using conventional techniques. For example, the infectious particles can be collected by cell lysis, or collection of the supernatant of the cell culture, as is known in the art. Optionally, the collected virus particles may be purified if desired. Suitable purification techniques are well known to those skilled in the art.

[0196] The delivery of a gene(s) or other polynucleotide sequence using a retroviral or lentiviral vector by means of viral infection rather than by transfection is referred to as "transduction." Retroviral vectors may be transduced into a cell through infection and provirus integration. A target cell, e.g., a T cell, is "transduced" if it comprises a gene or other polynucleotide sequence delivered to the cell by infection using a viral or retroviral vector. A transduced cell may comprise one or more genes or other polynucleotide sequences delivered by a retroviral or lentiviral vector in its cellular genome. Host cells transduced with viral vector that expresses one or more polypeptides, may be administered to a subject to treat and/or prevent a B cell malignancy. Other methods relating to the use of viral vectors in gene therapy, which may be utilized can be found in, e.g., Kay, M. A. (1997) Chest 111(6 Supp.):138S-142S; Ferry, N. and Heard, J. M. (1998) Hum. Gene Ther. 9:1975-81; Shiratory, Y. et al. (1999) Liver 19:265-74; Oka, K. et al. (2000) Curr. Opin. Lipidol. 11:179-86; Thule, P. M. and Liu, J. M. (2000) Gene Ther. 7:1744-52; Yang, N. S. (1992) Crit. Rev. Biotechnol. 12:335-56; Alt, M. (1995) J. Hepatol. 23:746-58; Brody, S. L. and Crystal, R. G. (1994) Ann. N.Y. Acad. Sci. 716:90-101; Strayer, D. S. (1999) Expert Opin. Investig. Drugs 8:2159-2172; Smith-Arica, J. R. and Bartlett, J. S. (2001) Curr. Cardiol. Rep. 3:43-49; and Lee, H. C. et al. (2000) Nature 408:483-8.

G. GENETICALLY MODIFIED CELLS

[0197] Contemplated are cells genetically modified to express the CARs contemplated herein, for use in the treatment of B cell related conditions. As used herein, the term "genetically engineered" or "genetically modified" refers to the addition of extra genetic material in the form of DNA or RNA into the total genetic material in a cell. The terms, "genetically modified cells," "modified cells," and, "redirected cells," are used interchangeably. As used herein, the term "gene therapy" refers to the introduction of extra genetic material in the form of DNA or RNA into the total genetic material in a cell that restores, corrects, or modifies expression of a gene, or for the purpose of expressing a therapeutic polypeptide, e.g., a CAR.

[0198] The CARs contemplated herein may be introduced and expressed in immune effector cells so as to redirect their specificity to a target antigen of interest, e.g., a BCMA polypeptide. An "immune effector cell," is any cell of the immune system that has one or more effector functions (e.g., cytotoxic cell killing activity, secretion of cytokines, induction of ADCC and/or CDC).

[0199] Immune effector cells can be autologous/autogeneic ("self") or non-autologous ("non-self," e.g., allogeneic, syngeneic or xenogeneic).

[0200] "Autologous," as used herein, refers to cells from the same subject.

[0201] "Allogeneic," as used herein, refers to cells of the same species that differ genetically to the cell in comparison.

[0202] "Syngeneic," as used herein, refers to cells of a different subject that are genetically identical to the cell in comparison.

[0203] "Xenogeneic," as used herein, refers to cells of a different species to the cell in comparison. Preferably, the cells are allogeneic. Illustrative immune effector cells used with the CARs contemplated herein include T lymphocytes. The terms "T cell" or "T lymphocyte" are art-recognized and are intended to include thymocytes, immature T lymphocytes, mature T lymphocytes, resting T lymphocytes, or activated T lymphocytes. A T cell can be a T helper (Th) cell, for example a T helper 1 (Th1) or a T helper 2 (Th2) cell. The T cell can be a helper T cell (HTL; CD4⁺ T cell) CD4⁺ T cell, a cytotoxic T cell (CTL; CD8⁺ T cell), CD4⁺CD8⁺ T cell, CD4-CD8- T cell, or any other subset of T cells. Other illustrative populations of T cells suitable for use include naive T cells and memory T cells.

[0204] As would be understood by the skilled person, other cells may also be used as immune effector cells with the CARs as described herein. In particular, immune effector cells also include NK cells, NKT cells, neutrophils, and macrophages. Immune effector cells also include progenitors of effector cells wherein such progenitor cells can be induced to differentiate into an immune effector cells *in vivo* or *in vitro*. Thus, immune effector cell may include progenitors of immune effectors cells such as hematopoietic stem cells (HSCs) contained within the CD34⁺ population of cells derived from cord blood, bone marrow or mobilized peripheral blood which upon administration in a subject differentiate into mature immune effector cells, or which can be induced *in vitro* to differentiate into mature immune effector cells.

[0205] As used herein, immune effector cells genetically engineered to contain BCMA-specific CAR may be referred

to as, "BCMA-specific redirected immune effector cells."

[0206] The term, "CD34+cell," as used herein refers to a cell expressing the CD34 protein on its cell surface. "CD34," as used herein refers to a cell surface glycoprotein (e.g., sialomucin protein) that often acts as a cell-cell adhesion factor and is involved in T cell entrance into lymph nodes. The CD34+ cell population contains hematopoietic stem cells (HSC), which upon administration to a patient differentiate and contribute to all hematopoietic lineages, including T cells, NK cells, NKT cells, neutrophils and cells of the monocyte/macrophage lineage.

[0207] Also provided are methods for making the immune effector cells which express the CAR contemplated herein. The method may comprise transfecting or transducing immune effector cells isolated from an individual such that the immune effector cells express one or more CAR as described herein. The immune effector cells may be isolated from an individual and genetically modified without further manipulation *in vitro*. Such cells can then be directly re-administered into the individual. The immune effector cells may be first activated and stimulated to proliferate *in vitro* prior to being genetically modified to express a CAR. In this regard, the immune effector cells may be cultured before and/or after being genetically modified (i.e., transduced or transfected to express a CAR contemplated herein).

[0208] Prior to *in vitro* manipulation or genetic modification of the immune effector cells described herein, the source of cells may be obtained from a subject. In particular, the CAR-modified immune effector cells may comprise T cells. T cells can be obtained from a number of sources including, but not limited to, peripheral blood mononuclear cells, bone marrow, lymph nodes tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. T cells may be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled person, such as sedimentation, e.g., FICOLL™ separation. Cells from the circulating blood of an individual may be obtained by apheresis. The apheresis product typically contains lymphocytes, including T cells, monocytes, granulocyte, B cells, other nucleated white blood cells, red blood cells, and platelets. The cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing. The cells can be washed with PBS or with another suitable solution that lacks calcium, magnesium, and most, if not all other, divalent cations. As would be appreciated by those of ordinary skill in the art, a washing step may be accomplished by methods known to those in the art, such as by using a semiautomated flowthrough centrifuge. For example, the Cobe 2991 cell processor, the Baxter CytoMate, or the like. After washing, the cells may be resuspended in a variety of biocompatible buffers or other saline solution with or without buffer. The undesirable components of the apheresis sample may be removed in the cell directly resuspended culture media.

[0209] T cells may be isolated from peripheral blood mononuclear cells (PBMCs) by lysing the red blood cells and depleting the monocytes, for example, by centrifugation through a PERCOLL™ gradient. A specific subpopulation of T cells, expressing one or more of the following markers: CD3, CD28, CD4, CD8, CD45RA, and CD45RO, can be further isolated by positive or negative selection techniques. In one embodiment, a specific subpopulation of T cells, expressing CD3, CD28, CD4, CD8, CD45RA, and CD45RO is further isolated by positive or negative selection techniques. For example, enrichment of a T cell population by negative selection can be accomplished with a combination of antibodies directed to surface markers unique to the negatively selected cells. One method for use herein is cell sorting and/or selection via negative magnetic immunoadherence or flow cytometry that uses a cocktail of monoclonal antibodies directed to cell surface markers present on the cells negatively selected. For example, to enrich for CD4⁺ cells by negative selection, a monoclonal antibody cocktail typically includes antibodies to CD14, CD20, CD11b, CD16, HLA-DR, and CD8. Flow cytometry and cell sorting may also be used to isolate cell populations of interest for use.

[0210] PBMC may be directly genetically modified to express CARs using methods contemplated herein. After isolation of PBMC, T lymphocytes may be further isolated and, both cytotoxic and helper T lymphocytes may be sorted into naive, memory, and effector T cell subpopulations either before or after genetic modification and/or expansion.

[0211] CD8⁺ cells can be obtained by using standard methods. CD8⁺ cells may be further sorted into naive, central memory, and effector cells by identifying cell surface antigens that are associated with each of those types of CD8⁺ cells.

[0212] Naive CD8⁺ T lymphocytes may be characterized by the expression of phenotypic markers of naive T cells including CD62L, CCR7, CD28, CD3, CD 127, and CD45RA.

[0213] Memory T cells may be present in both CD62L⁺ and CD62L⁻ subsets of CD8⁺ peripheral blood lymphocytes. PBMC are sorted into CD62L-CD8⁺ and CD62L⁺CD8⁺ fractions after staining with anti-CD8 and anti-CD62L antibodies. The expression of phenotypic markers of central memory T cells include CD45RO, CD62L, CCR7, CD28, CD3, and CD127 may be negative for granzyme B. Central memory T cells may be CD45RO⁺, CD62L⁺, CD8⁺ T cells.

[0214] Effector T cells may be negative for CD62L, CCR7, CD28, and CD127, and positive for granzyme B and perforin.

[0215] CD4⁺T cells may be further sorted into subpopulations. For example, CD4⁺ T helper cells can be sorted into naive, central memory, and effector cells by identifying cell populations that have cell surface antigens. CD4⁺ lymphocytes can be obtained by standard methods. Naive CD4⁺ T lymphocytes may be CD45RO⁻, CD45RA⁺, CD62L⁺ CD4⁺ T cell. Central memory CD4⁺cells may be CD62L positive and CD45RO positive. Effector CD4⁺ cells may be CD62L and CD45RO negative.

[0216] The immune effector cells, such as T cells, can be genetically modified following isolation using known methods, or the immune effector cells can be activated and expanded (or differentiated in the case of progenitors) *in vitro* prior to

being genetically modified. The immune effector cells, such as T cells, are genetically modified with the chimeric antigen receptors contemplated herein (e.g., transduced with a viral vector comprising a nucleic acid encoding a CAR) and then are activated and expanded *in vitro*. T cells can be activated and expanded before or after genetic modification to express a CAR, using methods as described, for example, in U.S. Patents 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; 6,867,041; and U.S. Patent Application Publication No. 20060121005.

[0217] Generally, the T cells are expanded by contact with a surface having attached thereto an agent that stimulates a CD3 TCR complex associated signal and a ligand that stimulates a co-stimulatory molecule on the surface of the T cells. T cell populations may be stimulated by contact with an anti-CD3 antibody, or antigen-binding fragment thereof, or an anti-CD2 antibody immobilized on a surface, or by contact with a protein kinase C activator (e.g., bryostatin) in conjunction with a calcium ionophore. Costimulation of accessory molecules on the surface of T cells, is also contemplated.

[0218] PBMCs or isolated T cells may be contacted with a stimulatory agent and costimulatory agent, such as anti-CD3 and anti-CD28 antibodies, generally attached to a bead or other surface, in a culture medium with appropriate cytokines, such as IL-2, IL-7, and/or IL-15. To stimulate proliferation of either CD4⁺ T cells or CD8⁺ T cells, an anti-CD3 antibody and an anti-CD28 antibody. Examples of an anti-CD28 antibody include 9.3, B-T3, XR-CD28 (Diacione, Besancon, France) can be used as can other methods commonly known in the art (Berg et al., Transplant Proc. 30(8):3975-3977, 1998; Haanen et al., J. Exp. Med. 190(9): 13191328, 1999; Garland et al., J. Immunol Meth. 227(1-2):53-63, 1999). Anti-CD3 and anti-CD28 antibodies attached to the same bead serve as a "surrogate" antigen presenting cell (APC). The T cells may be activated and stimulated to proliferate with feeder cells and appropriate antibodies and cytokines using methods such as those described in US6040177; US5827642; and WO2012129514.

[0219] Artificial APC (aAPC) made by engineering K562, U937, 721.221, T2, and C1R cells to direct the stable expression and secretion, of a variety of co-stimulatory molecules and cytokines. K32 or U32 aAPCs may be used to direct the display of one or more antibody-based stimulatory molecules on the aAPC cell surface. Expression of various combinations of genes on the aAPC enables the precise determination of human T-cell activation requirements, such that aAPCs can be tailored for the optimal propagation of T-cell subsets with specific growth requirements and distinct functions. The aAPCs support *ex vivo* growth and long-term expansion of functional human CD8 T cells without requiring the addition of exogenous cytokines, in contrast to the use of natural APCs. Populations of T cells can be expanded by aAPCs expressing a variety of costimulatory molecules including, but not limited to, CD137L (4-1BBL), CD134L (OX40L), and/or CD80 or CD86. Finally, the aAPCs provide an efficient platform to expand genetically modified T cells and to maintain CD28 expression on CD8 T cells. aAPCs provided in WO 03/057171 and US2003/0147869 are referred to.

[0220] CD34⁺ cells may be transduced with a nucleic acid construct. The transduced CD34⁺ cells may differentiate into mature immune effector cells *in vivo* following administration into a subject, generally the subject from whom the cells were originally isolated. CD34⁺ cells may be stimulated *in vitro* prior to exposure to or after being genetically modified with a CAR as described herein, with one or more of the following cytokines: Flt-3 ligand (FLT3), stem cell factor (SCF), megakaryocyte growth and differentiation factor (TPO), IL-3 and IL-6 according to the methods described previously (Asheuer et al., 2004; Imren, et al., 2004).

[0221] Also provided are a population of modified immune effector cells for the treatment of cancer, the modified immune effector cells comprising a CAR as disclosed herein. For example, a population of modified immune effector cells are prepared from peripheral blood mononuclear cells (PBMCs) obtained from a patient diagnosed with B cell malignancy described herein (autologous donors). The PBMCs form a heterogeneous population of T lymphocytes that can be CD4⁺, CD8⁺, or CD4⁺ and CD8⁺.

[0222] The PBMCs also can include other cytotoxic lymphocytes such as NK cells or NKT cells. An expression vector carrying the coding sequence of a CAR contemplated herein can be introduced into a population of human donor T cells, NK cells or NKT cells. Successfully transduced T cells that carry the expression vector can be sorted using flow cytometry to isolate CD3 positive T cells and then further propagated to increase the number of these CAR protein expressing T cells in addition to cell activation using anti-CD3 antibodies and or anti-CD28 antibodies and IL-2 or any other methods known in the art as described elsewhere herein. Standard procedures are used for cryopreservation of T cells expressing the CAR protein T cells for storage and/or preparation for use in a human subject. The *in vitro* transduction, culture and/or expansion of T cells may be performed in the absence of non-human animal derived products such as fetal calf serum and fetal bovine serum. Since a heterogeneous population of PBMCs is genetically modified, the resultant transduced cells are a heterogeneous population of modified cells comprising a BCMA targeting CAR as contemplated herein.

[0223] A mixture of, e.g., one, two, three, four, five or more, different expression vectors can be used in genetically modifying a donor population of immune effector cells wherein each vector encodes a different chimeric antigen receptor protein as contemplated herein. The resulting modified immune effector cells forms a mixed population of modified cells, with a proportion of the modified cells expressing more than one different CAR proteins.

[0224] Also provided is a method of storing genetically modified murine, human or humanized CAR protein expressing immune effector cells which target a BCMA protein, comprising cryopreserving the immune effector cells such that the

cells remain viable upon thawing. A fraction of the immune effector cells expressing the CAR proteins can be cryopreserved by methods known in the art to provide a permanent source of such cells for the future treatment of patients afflicted with the B cell related condition. When needed, the cryopreserved transformed immune effector cells can be thawed, grown and expanded for more such cells.

[0225] As used herein, "cryopreserving," refers to the preservation of cells by cooling to sub-zero temperatures, such as (typically) 77 K or -196° C. (the boiling point of liquid nitrogen). Cryoprotective agents are often used at sub-zero temperatures to prevent the cells being preserved from damage due to freezing at low temperatures or warming to room temperature. Cryopreservative agents and optimal cooling rates can protect against cell injury. Cryoprotective agents which can be used include but are not limited to dimethyl sulfoxide (DMSO) (Lovelock and Bishop, Nature, 1959; 183: 1394-1395; Ashwood-Smith, Nature, 1961; 190: 1204-1205), glycerol, polyvinylpyrrolidone (Rinfret, Ann. N.Y. Acad. Sci., 1960; 85: 576), and polyethylene glycol (Sloviter and Ravdin, Nature, 1962; 196: 48). The preferred cooling rate is 1° to 3° C/minute. After at least two hours, the T cells have reached a temperature of -80° C. and can be placed directly into liquid nitrogen (-196° C.) for permanent storage such as in a long-term cryogenic storage vessel.

H. T CELL MANUFACTURING METHODS

[0226] The T cells manufactured by the methods contemplated herein provide improved adoptive immunotherapy compositions. Without wishing to be bound to any particular theory, it is believed that the T cell compositions manufactured by the methods contemplated herein are imbued with superior properties, including increased survival, expansion in the relative absence of differentiation, and persistence *in vivo*. A method of manufacturing T cells may comprise contacting the cells with one or more agents that modulate a PI3K cell signaling pathway. In one embodiment, a method of manufacturing T cells comprises contacting the cells with one or more agents that modulate a PI3K/Akt/mTOR cell signaling pathway. In various embodiments, the T cells may be obtained from any source and contacted with the agent during the activation and/or expansion phases of the manufacturing process. The resulting T cell compositions are enriched in developmentally potent T cells that have the ability to proliferate and express one or more of the following biomarkers: CD62L, CCR7, CD28, CD27, CD122, CD127, CD197, and CD38. Populations of cell comprising T cells, that have been treated with one or more PI3K inhibitors may be enriched for a population of CD8+ T cells co-expressing one or more or, or all of, the following biomarkers: CD62L, CD127, CD197, and CD38.

[0227] Modified T cells comprising maintained levels of proliferation and decreased differentiation may be manufactured. T cells may be manufactured by stimulating T cells to become activated and to proliferate in the presence of one or more stimulatory signals and an agent that is an inhibitor of a PI3K cell signaling pathway.

[0228] The T cells can then be modified to express an anti-BCMA CARs. The T cells may be modified by transducing the T cells with a viral vector comprising an anti-BCMA CAR contemplated herein. The T cells may be modified prior to stimulation and activation in the presence of an inhibitor of a PI3K cell signaling pathway. T cells may be modified after stimulation and activation in the presence of an inhibitor of a PI3K cell signaling pathway. T cells may be modified within 12 hours, 24 hours, 36 hours, or 48 hours of stimulation and activation in the presence of an inhibitor of a PI3K cell signaling pathway.

[0229] After T cells are activated, the cells are cultured to proliferate. T cells may be cultured for at least 1, 2, 3, 4, 5, 6, or 7 days, at least 2 weeks, at least 1, 2, 3, 4, 5, or 6 months or more with 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more rounds of expansion.

[0230] T cell compositions may be manufactured in the presence of one or more inhibitors of the PI3K pathway. The inhibitors may target one or more activities in the pathway or a single activity. Without wishing to be bound to any particular theory, it is contemplated that treatment or contacting T cells with one or more inhibitors of the PI3K pathway during the stimulation, activation, and/or expansion phases of the manufacturing process preferentially increases young T cells, thereby producing superior therapeutic T cell compositions.

[0231] A method for increasing the proliferation of T cells expressing an engineered T cell receptor is provided. Such methods may comprise, for example, harvesting a source of T cells from a subject, stimulating and activating the T cells in the presence of one or more inhibitors of the PI3K pathway, modification of the T cells to express an anti-BCMA CAR, e.g., anti-BCMA02 CAR, and expanding the T cells in culture.

[0232] A method for producing populations of T cells enriched for expression of one or more of the following biomarkers: CD62L, CCR7, CD28, CD27, CD122, CD127, CD197, and CD38. Young T cells may comprise one or more of, or all of the following biological markers: CD62L, CD127, CD197, and CD38.

The young T cells may lack expression of CD57, CD244, CD160, PD-1, CTLA4, TIM3, and LAG3 are provided. As discussed elsewhere herein, the expression levels young T cell biomarkers is relative to the expression levels of such markers in more differentiated T cells or immune effector cell populations.

[0233] Peripheral blood mononuclear cells (PBMCs) may be used as the source of T cells in the T cell manufacturing methods contemplated herein. PBMCs form a heterogeneous population of T lymphocytes that can be CD4+, CD8+, or CD4+ and CD8+ and can include other mononuclear cells such as monocytes, B cells, NK cells and NKT cells. An

expression vector comprising a polynucleotide encoding an engineered TCR or CAR contemplated herein can be introduced into a population of human donor T cells, NK cells or NKT cells. Successfully transduced T cells that carry the expression vector can be sorted using flow cytometry to isolate CD3 positive T cells and then further propagated to increase the number of the modified T cells in addition to cell activation using anti-CD3 antibodies and or anti-CD28 antibodies and IL-2, IL-7, and/or IL-15 or any other methods known in the art as described elsewhere herein.

[0234] Manufacturing methods contemplated herein may further comprise cryopreservation of modified T cells for storage and/or preparation for use in a human subject. T cells are cryopreserved such that the cells remain viable upon thawing. When needed, the cryopreserved transformed immune effector cells can be thawed, grown and expanded for more such cells. As used herein, "cryopreserving," refers to the preservation of cells by cooling to sub-zero temperatures, such as (typically) 77 K or -196° C. (the boiling point of liquid nitrogen). Cryoprotective agents are often used at sub-zero temperatures to prevent the cells being preserved from damage due to freezing at low temperatures or warming to room temperature. Cryopreservative agents and optimal cooling rates can protect against cell injury. Cryoprotective agents which can be used include but are not limited to dimethyl sulfoxide (DMSO) (Lovelock and Bishop, Nature, 1959; 183: 1394-1395; Ashwood-Smith, Nature, 1961; 190: 1204-1205), glycerol, polyvinylpyrrolidone (Rinfret, Ann. N.Y. Acad. Sci., 1960; 85: 576), and polyethylene glycol (Sloviter and Ravdin, Nature, 1962; 196: 48). The preferred cooling rate is 1° to 3° C/minute. After at least two hours, the T cells have reached a temperature of -80° C. and can be placed directly into liquid nitrogen (-196° C.) for permanent storage such as in a long-term cryogenic storage vessel.

1. T CELLS

[0235] Also contemplated is the manufacture of improved CAR T cell compositions. T cells used for CAR T cell production may be autologous/autogeneic ("self") or non-autologous ("non-self," e.g., allogeneic, syngeneic or xenogeneic). The T cells may be obtained from a mammalian subject. The T cells are obtained from a primate subject. The T cells are obtained from a human subject.

[0236] T cells can be obtained from a number of sources including, but not limited to, peripheral blood mononuclear cells, bone marrow, lymph nodes tissue, cord blood, thymus issue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. T cells may be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled person, such as sedimentation, e.g., FICOLL™ separation. Cells from the circulating blood of an individual may be obtained by apheresis. The apheresis product typically contains lymphocytes, including T cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. The cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing. The cells can be washed with PBS or with another suitable solution that lacks calcium, magnesium, and most, if not all other, divalent cations. As would be appreciated by those of ordinary skill in the art, a washing step may be accomplished by methods known to those in the art, such as by using a semiautomated flowthrough centrifuge. For example, the Cobe 2991 cell processor, the Baxter CytoMate, or the like. After washing, the cells may be resuspended in a variety of biocompatible buffers or other saline solution with or without buffer. The undesirable components of the apheresis sample may be removed in the cell directly resuspended culture media.

[0237] A population of cells comprising T cells, e.g., PBMCs, may be used in the manufacturing methods contemplated herein. An isolated or purified population of T cells may be used in the manufacturing methods contemplated herein. Cells can be isolated from peripheral blood mononuclear cells (PBMCs) by lysing the red blood cells and depleting the monocytes, for example, by centrifugation through a PERCOLL™ gradient. After isolation of PBMC, both cytotoxic and helper T lymphocytes can be sorted into naïve, memory, and effector T cell subpopulations either before or after activation, expansion, and/or genetic modification.

[0238] A specific subpopulation of T cells, expressing one or more of the following markers: CD3, CD4, CD8, CD28, CD45RA, CD45RO, CD62, CD 127, and HLA-DR can be further isolated by positive or negative selection techniques. A specific subpopulation of T cells, expressing one or more of the markers selected from the group consisting of i) CD62L, CCR7, CD28, CD27, CD122, CD127, CD197; or ii) CD38 or CD62L, CD127, CD197, and CD38, may be further isolated by positive or negative selection techniques. In various embodiments, the manufactured T cell compositions do not express or may not substantially express one or more of the following markers: CD57, CD244, CD160, PD-1, CTLA4, TIM3, and LAG3.

[0239] Expression of one or more of the markers selected from the group consisting of CD62L, CD127, CD197, and CD38 is increased at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, at least 10 fold, at least 25 fold, or more compared to a population of T cells activated and expanded without a PI3K inhibitor.

[0240] Expression of one or more of the markers selected from the group consisting of CD57, CD244, CD160, PD-1, CTLA4, TIM3, and LAG3 is decreased at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, at least 10 fold, at least 25 fold, or more compared to a population of T cells activated and expanded with a PI3K inhibitor.

[0241] The manufacturing methods contemplated herein may increase the number CAR T cells comprising one or more markers of naïve or developmentally potent T cells. Without wishing to be bound to any particular theory, the present inventors believe that treating a population of cells comprising T cells with one or more PI3K inhibitors results in an increase an expansion of developmentally potent T cells and provides a more robust and efficacious adoptive CAR T cell immunotherapy compared to existing CAR T cell therapies.

[0242] Illustrative examples of markers of naïve or developmentally potent T cells increased in T cells manufactured using the methods contemplated herein include, but are not limited to CD62L, CD127, CD197, and CD38. Naïve T cells may not express or may not substantially express one or more of the following markers: CD57, CD244, CD160, PD-1, BTLA, CD45RA, CTLA4, TIM3, and LAG3.

[0243] With respect to T cells, the T cell populations resulting from the various expansion methodologies contemplated herein may have a variety of specific phenotypic properties, depending on the conditions employed. Expanded T cell populations comprise one or more of the following phenotypic markers: CD62L, CD127, CD197, CD38, and HLA-DR.

[0244] Such phenotypic markers may include enhanced expression of one or more of, or all of CD62L, CD127, CD197, and CD38. CD8⁺ T lymphocytes characterized by the expression of phenotypic markers of naïve T cells including CD62L, CD127, CD197, and CD38 may be expanded.

[0245] T cells characterized by the expression of phenotypic markers of central memory T cells including CD45RO, CD62L, CD127, CD197, and CD38 and negative for granzyme B may be expanded. The central memory T cells may be CD45RO⁺, CD62L⁺, CD8⁺ T cells.

[0246] CD4⁺ T lymphocytes characterized by the expression of phenotypic markers of naïve CD4⁺ cells including CD62L and negative for expression of CD45RA and/or CD45RO may be expanded. CD4⁺ cells characterized by the expression of phenotypic markers of central memory CD4⁺ cells may include CD62L and CD45RO positive. Effector CD4⁺ cells may be CD62L positive and CD45RO negative.

[0247] The T cells may be isolated from an individual and activated and stimulated to proliferate *in vitro* prior to being genetically modified to express an anti-BCMA CAR. In this regard, the T cells may be cultured before and/or after being genetically modified (*i.e.*, transduced or transfected to express an anti-BCMA CAR contemplated herein).

2. ACTIVATION AND EXPANSION

[0248] In order to achieve sufficient therapeutic doses of T cell compositions, T cells are often subject to one or more rounds of stimulation, activation and/or expansion. T cells can be activated and expanded generally using methods as described, for example, in U.S. Patents 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; and 6,867,041. T cells modified to express an anti-BCMA CAR can be activated and expanded before and/or after the T cells are modified. In addition, T cells may be contacted with one or more agents that modulate the PI3K cell signaling pathway before, during, and/or after activation and/or expansion. T cells manufactured by the methods contemplated herein may undergo one, two, three, four, or five or more rounds of activation and expansion, each of which may include one or more agents that modulate the PI3K cell signaling pathway.

[0249] A costimulatory ligand may be presented on an antigen presenting cell (*e.g.*, an aAPC, dendritic cell, B cell, and the like) that specifically binds a cognate costimulatory molecule on a T cell, thereby providing a signal which, in addition to the primary signal provided by, for instance, binding of a TCR/CD3 complex, mediates a desired T cell response. Suitable costimulatory ligands include, but are not limited to, CD7, B7-1 (CD80), B7-2 (CD86), PD-L 1, PD-L2, 4-1BBL, OX40L, inducible costimulatory ligand (ICOS-L), intercellular adhesion molecule (ICAM), CD30L, CD40, CD70, CD83, HLA-G, MICA, MICB, HVEM, lymphotoxin beta receptor, ILT3, ILT4, an agonist or antibody that binds Toll ligand receptor, and a ligand that specifically binds with B7-H3.

[0250] A costimulatory ligand comprises an antibody or antigen binding fragment thereof that specifically binds to a costimulatory molecule present on a T cell, including but not limited to, CD27, CD28, 4-1BB, OX40, CD30, CD40, PD-1, ICOS, lymphocyte function-associated antigen- 1 (LFA-1), CD7, LIGHT, NKG2C, B7-H3, and a ligand that specifically binds with CD83.

[0251] Suitable costimulatory ligands further include target antigens, which may be provided in soluble form or expressed on APCs or aAPCs that bind engineered TCRs or CARs expressed on modified T cells.

[0252] A method for manufacturing T cells contemplated herein may comprise activating a population of cells comprising T cells and expanding the population of T cells. T cell activation can be accomplished by providing a primary stimulation signal through the T cell TCR/CD3 complex or via stimulation of the CD2 surface protein and by providing a secondary costimulation signal through an accessory molecule, *e.g.*, CD28.

[0253] The TCR/CD3 complex may be stimulated by contacting the T cell with a suitable CD3 binding agent, *e.g.*, a CD3 ligand or an anti-CD3 monoclonal antibody. Illustrative examples of CD3 antibodies include, but are not limited to, OKT3, G19-4, BC3, and 64.1.

[0254] A CD2 binding agent may be used to provide a primary stimulation signal to the T cells. Illustrative examples

of CD2 binding agents include, but are not limited to, CD2 ligands and anti-CD2 antibodies, e.g., the T11.3 antibody in combination with the T11.1 or T11.2 antibody (Meuer, S. C. et al. (1984) Cell 36:897-906) and the 9.6 antibody (which recognizes the same epitope as TI 1.1) in combination with the 9-1 antibody (Yang, S. Y. et al. (1986) J. Immunol. 137:1097-1100). Other antibodies which bind to the same epitopes as any of the above described antibodies can also be used. Additional antibodies, or combinations of antibodies, can be prepared and identified by standard techniques as disclosed elsewhere herein.

[0255] In addition to the primary stimulation signal provided through the TCR/CD3 complex, or via CD2, induction of T cell responses requires a second, costimulatory signal. A CD28 binding agent can be used to provide a costimulatory signal. Illustrative examples of CD28 binding agents include but are not limited to: natural CD 28 ligands, e.g., a natural ligand for CD28 (e.g., a member of the B7 family of proteins, such as B7-1(CD80) and B7-2 (CD86); and anti-CD28 monoclonal antibody or fragment thereof capable of crosslinking the CD28 molecule, e.g., monoclonal antibodies 9.3, B-T3, XR-CD28, KOLT-2, 15E8, 248.23.2, and EX5.3D10.

[0256] The molecule providing the primary stimulation signal, for example a molecule which provides stimulation through the TCR/CD3 complex or CD2, and the costimulatory molecule may be coupled to the same surface.

[0257] Binding agents that provide stimulatory and costimulatory signals may be localized on the surface of a cell. This can be accomplished by transfecting or transducing a cell with a nucleic acid encoding the binding agent in a form suitable for its expression on the cell surface or alternatively by coupling a binding agent to the cell surface.

[0258] The molecule providing the primary stimulation signal, for example a molecule which provides stimulation through the TCR/CD3 complex or CD2, and the costimulatory molecule may be displayed on antigen presenting cells.

[0259] In one embodiment, the molecule providing the primary stimulation signal, for example a molecule which provides stimulation through the TCR/CD3 complex or CD2, and the costimulatory molecule may be provided on separate surfaces.

[0260] One of the binding agents that provide stimulatory and costimulatory signals may be soluble (provided in solution) and the other agent(s) is provided on one or more surfaces.

[0261] The binding agents that provide stimulatory and costimulatory signals may both provided in a soluble form (provided in solution).

[0262] The methods for manufacturing T cells contemplated herein may comprise activating T cells with anti-CD3 and anti-CD28 antibodies.

[0263] T cell compositions manufactured by the methods contemplated herein comprise T cells activated and/or expanded in the presence of one or more agents that inhibit a PI3K cell signaling pathway. T cells modified to express an anti-BCMA CAR can be activated and expanded before and/or after the T cells are modified. A population of T cells may be activated, modified to express an anti-BCMA CAR, and then cultured for expansion.

[0264] T cells manufactured by the methods contemplated herein may comprise an increased number of T cells expressing markers indicative of high proliferative potential and the ability to self-renew but that do not express or express substantially undetectable markers of T cell differentiation. These T cells may be repeatedly activated and expanded in a robust fashion and thereby provide an improved therapeutic T cell composition.

[0265] A population of T cells activated and expanded in the presence of one or more agents that inhibit a PI3K cell signaling pathway may be expanded at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, at least 10 fold, at least 25 fold, at least 50 fold, at least 100 fold, at least 250 fold, at least 500 fold, at least 1000 fold, or more compared to a population of T cells activated and expanded without a PI3K inhibitor.

[0266] A population of T cells characterized by the expression of markers young T cells may be activated and expanded in the presence of one or more agents that inhibit a PI3K cell signaling pathway and may be expanded at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, at least 10 fold, at least 25 fold, at least 50 fold, at least 100 fold, at least 250 fold, at least 500 fold, at least 1000 fold, or more compared the population of T cells activated and expanded without a PI3K inhibitor.

[0267] Expanding T cells activated by the methods contemplated herein may further comprise culturing a population of cells comprising T cells for several hours (about 3 hours) to about 7 days to about 28 days or any hourly integer value in between. The T cell composition may be cultured for 14 days. T cells may be cultured for about 21 days. The T cell compositions may be cultured for about 2-3 days. Several cycles of stimulation/activation/expansion may also be desired such that culture time of T cells can be 60 days or more.

[0268] Conditions appropriate for T cell culture may include an appropriate media (e.g., Minimal Essential Media or RPMI Media 1640 or, X-vivo 15, (Lonza)) and one or more factors necessary for proliferation and viability including, but not limited to serum (e.g., fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, IL-21, GM-CSF, IL- 10, IL- 12, IL-15, TGF β , and TNF- α or any other additives suitable for the growth of cells known to the skilled artisan.

[0269] Further illustrative examples of cell culture media include, but are not limited to RPMI 1640, Clicks, AIM-V, DMEM, MEM, a-MEM, F-12, X-Vivo 15, and X-Vivo 20, Optimizer, with added amino acids, sodium pyruvate, and vitamins, either serum-free or supplemented with an appropriate amount of serum (or plasma) or a defined set of

hormones, and/or an amount of cytokine(s) sufficient for the growth and expansion of T cells.

[0270] Illustrative examples of other additives for T cell expansion include, but are not limited to, surfactant, piasmanate, pH buffers such as HEPES, and reducing agents such as N-acetylcysteine and 2-mercaptoethanol

[0271] Antibiotics, e.g., penicillin and streptomycin, are included only in experimental cultures, not in cultures of cells that are to be infused into a subject. The target cells are maintained under conditions necessary to support growth, for example, an appropriate temperature (e.g., 37° C) and atmosphere (e.g., air plus 5% CO₂).

[0272] PBMCs or isolated T cells may be contacted with a stimulatory agent and costimulatory agent, such as anti-CD3 and anti-CD28 antibodies, generally attached to a bead or other surface, in a culture medium with appropriate cytokines, such as IL-2, IL-7, and/or IL-15.

[0273] Artificial APC (aAPC) made by engineering K562, U937, 721.221, T2, and C1R cells to direct the stable expression and secretion, of a variety of costimulatory molecules and cytokines. K32 or U32 aAPCs may be used to direct the display of one or more antibody-based stimulatory molecules on the aAPC cell surface. Populations of T cells can be expanded by aAPCs expressing a variety of costimulatory molecules including, but not limited to, CD137L (4-1BBL), CD134L (OX40L), and/or CD80 or CD86. Finally, the aAPCs provide an efficient platform to expand genetically modified T cells and to maintain CD28 expression on CD8 T cells. aAPCs provided in WO 03/057171 and US2003/0147869.

3. AGENTS

[0274] A method for manufacturing T cells is provided that expands undifferentiated or developmentally potent T cells comprising contacting T cells with an agent that modulates a PI3K pathway in the cells. A method for manufacturing T cells is provided that expands undifferentiated or developmentally potent T cells comprising contacting T cells with an agent that modulates a PI3K/AKT/mTOR pathway in the cells. The cells may be contacted prior to, during, and/or after activation and expansion. The T cell compositions retain sufficient T cell potency such that they may undergo multiple rounds of expansion without a substantial increase in differentiation.

[0275] As used herein, the terms "modulate," "modulator," or "modulatory agent" or comparable term refer to an agent's ability to elicit a change in a cell signaling pathway. A modulator may increase or decrease an amount, activity of a pathway component or increase or decrease a desired effect or output of a cell signaling pathway. The modulator may be an inhibitor. The modulator is an activator.

[0276] An "agent" refers to a compound, small molecule, e.g., small organic molecule, nucleic acid, polypeptide, or a fragment, isoform, variant, analog, or derivative thereof used in the modulation of a PI3K/AKT/mTOR pathway.

[0277] A "small molecule" refers to a composition that has a molecular weight of less than about 5 kD, less than about 4 kD, less than about 3 kD, less than about 2 kD, less than about 1 kD, or less than about .5kD. Small molecules may comprise nucleic acids, peptides, polypeptides, peptidomimetics, peptoids, carbohydrates, lipids, components thereof or other organic or inorganic molecules. Libraries of chemical and/or biological

mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays. Examples of methods for the synthesis of molecular libraries can be found in: (Carell *et al.*, 1994a; Carell *et al.*, 1994b; Cho *et al.*, 1993; DeWitt *et al.*, 1993; Gallop *et al.*, 1994; Zuckermann *et al.*, 1994).

[0278] An "analog" refers to a small organic compound, a nucleotide, a protein, or a polypeptide that possesses similar or identical activity or function(s) as the compound, nucleotide, protein or polypeptide or compound having the desired activity but need not necessarily comprise a sequence or structure that is similar or identical to the sequence or structure of the preferred embodiment.

[0279] A "derivative" refers to either a compound, a protein or polypeptide that comprises an amino acid sequence of a parent protein or polypeptide that has been altered by the introduction of amino acid residue substitutions, deletions or additions, or a nucleic acid or nucleotide that has been modified by either introduction of nucleotide substitutions or deletions, additions or mutations. The derivative nucleic acid, nucleotide, protein or polypeptide possesses a similar or identical function as the parent polypeptide.

[0280] The agent that modulates a PI3K pathway may activate a component of the pathway. An "activator," or "agonist" refers to an agent that promotes, increases, or induces one or more activities of a molecule in a PI3K/AKT/mTOR pathway including, without limitation, a molecule that inhibits one or more activities of a PI3K.

[0281] The agent that modulates a PI3K pathway may inhibit a component of the pathway. An "inhibitor" or "antagonist" refers to an agent that inhibits, decreases, or reduces one or more activities of a molecule in a PI3K pathway including, without limitation, a PI3K. The inhibitor may be a dual molecule inhibitor. The inhibitor may inhibit a class of molecules have the same or substantially similar activities (a pan-inhibitor) or may specifically inhibit a molecule's activity (a selective or specific inhibitor). Inhibition may also be irreversible or reversible.

[0282] The inhibitor may have an IC₅₀ of at least 1nM, at least 2nM, at least 5nM, at least 10nM, at least 50nM, at least 100nM, at least 200nM, at least 500nM, at least 1μM, at least 10μM, at least 50μM, or at least 100μM. IC₅₀ determinations can be accomplished using any conventional techniques known in the art. For example, an IC₅₀ can be determined by measuring the activity of a given enzyme in the presence of a range of concentrations of the inhibitor

under study. The experimentally obtained values of enzyme activity then are plotted against the inhibitor concentrations used. The concentration of the inhibitor that shows 50% enzyme activity (as compared to the activity in the absence of any inhibitor) is taken as the "IC50" value. Analogously, other inhibitory concentrations can be defined through appropriate determinations of activity.

[0283] T cells may be contacted or treated or cultured with one or more modulators of a PI3K pathway at a concentration of at least 1nM, at least 2nM, at least 5nM, at least 10nM, at least 50nM, at least 100nM, at least 200nM, at least 500nM, at least 1 μ M, at least 10 μ M, at least 50 μ M, at least 100 μ M, or at least 1 M.

[0284] T cells may be contacted or treated or cultured with one or more modulators of a PI3K pathway for at least 12 hours, 18 hours, at least 1, 2, 3, 4, 5, 6, or 7 days, at least 2 weeks, at least 1, 2, 3, 4, 5, or 6 months or more with 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more rounds of expansion.

a. PI3K/Akt/mTOR pathway

[0285] The phosphatidylinositol-3 kinase/Akt/mammalian target of rapamycin pathway serves as a conduit to integrate growth factor signaling with cellular proliferation, differentiation, metabolism, and survival. PI3Ks are a family of highly conserved intracellular lipid kinases. Class IA PI3Ks are activated by growth factor receptor tyrosine kinases (RTKs), either directly or through interaction with the insulin receptor substrate family of adaptor molecules. This activity results in the production of phosphatidylinositol-3,4,5-trisphosphate (PIP3) a regulator of the serine/threonine kinase Akt. mTOR acts through the canonical PI3K pathway via 2 distinct complexes, each characterized by different binding partners that confer distinct activities. mTORC1 (mTOR in complex with PRAS40, raptor, and mLST8/GbL) acts as a downstream effector of PI3K/Akt signaling, linking growth factor signals with protein translation, cell growth, proliferation, and survival. mTORC2 (mTOR in complex with rictor, mSIN1, protor, and mLST8) acts as an upstream activator of Akt.

[0286] Upon growth factor receptor-mediated activation of PI3K, Akt is recruited to the membrane through the interaction of its pleckstrin homology domain with PIP3, thus exposing its activation loop and enabling phosphorylation at threonine 308 (Thr308) by the constitutively active phosphoinositide-dependent protein kinase 1 (PDK1). For maximal activation, Akt is also phosphorylated by mTORC2, at serine 473 (Ser473) of its C-terminal hydrophobic motif. DNA-PK and HSP have also been shown to be important in the regulation of Akt activity. Akt activates mTORC1 through inhibitory phosphorylation of

TSC2, which along with TSC1, negatively regulates mTORC1 by inhibiting the Rheb GTPase, a positive regulator of mTORC1. mTORC1 has 2 well-defined substrates, p70S6K (referred to hereafter as S6K1) and 4E-BP1, both of which critically regulate protein synthesis. Thus, mTORC1 is an important downstream effector of PI3K, linking growth factor signaling with protein translation and cellular proliferation.

b. PI3K Inhibitors

[0287] As used herein, the term "PI3K inhibitor" refers to a nucleic acid, peptide, compound, or small organic molecule that binds to and inhibits at least one activity of PI3K. The PI3K proteins can be divided into three classes, class 1 PI3Ks, class 2 PI3Ks, and class 3 PI3Ks. Class 1 PI3Ks exist as heterodimers consisting of one of four p110 catalytic subunits (p110 α , p110 β , p110 δ , and p110 γ) and one of two families of regulatory subunits. A PI3K inhibitor preferably targets the class 1 PI3K inhibitors. A PI3K inhibitor may display selectivity for one or more isoforms of the class 1 PI3K inhibitors (i.e., selectivity for p110 α , p110 β , p110 δ , and p110 γ , or one or more of p110 α , p110 β , p110 δ , and p110 γ). A PI3K inhibitor may not display isoform selectivity and be considered a "pan-PI3K inhibitor." A PI3K inhibitor may compete for binding with ATP to the PI3K catalytic domain.

[0288] A PI3K inhibitor can, for example, target PI3K as well as additional proteins in the PI3K-AKT-mTOR pathway. A PI3K inhibitor may target both mTOR and PI3K and can be referred to as either an mTOR inhibitor or a PI3K inhibitor. A PI3K inhibitor that only targets PI3K can be referred to as a selective PI3K inhibitor. A selective PI3K inhibitor may be understood to refer to an agent that exhibits a 50% inhibitory concentration with respect to PI3K that is at least 10-fold, at least 20-fold, at least 30-fold, at least 50-fold, at least 100-fold, at least 1000-fold, or more, lower than the inhibitor's IC50 with respect to mTOR and/or other proteins in the pathway.

[0289] Exemplary PI3K inhibitors may inhibit PI3K with an IC50 (concentration that inhibits 50% of the activity) of about 200 nM or less, preferably about 100 nM or less, even more preferably about 60 nM or less, about 25 nM, about 10 nM, about 5 nM, about 1 nM, 100 μ M, 50 μ M, 25 μ M, 10 μ M, 1 μ M, or less. A PI3K inhibitor may inhibit PI3K with an IC50 from about 2 nM to about 100 nM, more preferably from about 2 nM to about 50 nM, even more preferably from about 2 nM to about 15 nM.

[0290] Illustrative examples of PI3K inhibitors suitable for use in the T cell manufacturing methods contemplated herein include, but are not limited to, BKM120 (class 1 PI3K inhibitor, Novartis), XL147 (class 1 PI3K inhibitor, Exelixis), (pan-PI3K inhibitor, GlaxoSmithKline), and PX-866 (class 1 PI3K inhibitor; p110 α , p110 β , and p110 γ isoforms, Oncothyreon).

[0291] Other illustrative examples of selective PI3K inhibitors include, but are not limited to BYL719, GSK2636771,

TGX-221, AS25242, CAL-101, ZSTK474, and IPI-145.

[0292] Further illustrative examples of pan-PI3K inhibitors include, but are not limited to BEZ235, LY294002, GSK1059615, TG100713, and GDC-0941.

c. AKT Inhibitors

[0293] As used herein, the term "AKT inhibitor" refers to a nucleic acid, peptide, compound, or small organic molecule that inhibits at least one activity of AKT. AKT inhibitors can be grouped into several classes, including lipid-based inhibitors (e.g., inhibitors that target the pleckstrin homology domain of AKT which prevents AKT from localizing to plasma membranes), ATP-competitive inhibitors, and allosteric inhibitors. AKT inhibitors may act by binding to the AKT catalytic site. Akt inhibitors may act by inhibiting phosphorylation of downstream AKT targets such as mTOR. AKT activity may be inhibited by inhibiting the input signals to activate Akt by inhibiting, for example, DNA-PK activation of AKT, PDK-1 activation of AKT, and/or mTORC2 activation of Akt.

[0294] AKT inhibitors can target all three AKT isoforms, AKT1, AKT2, AKT3 or may be isoform selective and target only one or two of the AKT isoforms. An AKT inhibitor can target AKT as well as additional proteins in the PI3K-AKT-mTOR pathway. An AKT inhibitor that only targets AKT can be referred to as a selective AKT inhibitor. A selective AKT inhibitor can be understood to refer to an agent that exhibits a 50% inhibitory concentration with respect to AKT that is at least 10-fold, at least 20-fold, at least 30-fold, at least 50-fold, at least 100-fold, at least 1000-fold, or more lower than the inhibitor's IC₅₀ with respect to other proteins in the pathway.

[0295] Exemplary AKT inhibitors may inhibit AKT with an IC₅₀ (concentration that inhibits 50% of the activity) of about 200 nM or less, preferably about 100 nM or less, even more preferably about 60 nM or less, about 25 nM, about 10 nM, about 5 nM, about 1 nM, 100 μ M, 50 μ M, 25 μ M, 10 μ M, 1 μ M, or less. An AKT may inhibit AKT with an IC₅₀ from about 2 nM to about 100 nM, more preferably from about 2 nM to about 50 nM, even more preferably from about 2 nM to about 15 nM.

[0296] Illustrative examples of AKT inhibitors for use in combination with auristatin based antibody-drug conjugates include, for example, perifosine (Keryx), MK2206 (Merck), VQD-002 (VioQuest), XL418 (Exelixis), GSK690693, GDC-0068, and PX316 (PROLX Pharmaceuticals).

[0297] An illustrative, non-limiting example of a selective Akt1 inhibitor is A-674563.

[0298] An illustrative, non-limiting example of a selective Akt2 inhibitor is CCT128930.

[0299] The Akt inhibitor DNA-PK activation of Akt, PDK-1 activation of Akt, mTORC2 activation of Akt, or HSP activation of Akt.

[0300] Illustrative examples of DNA-PK inhibitors include, but are not limited to, NU7441, PI-103, NU7026, PIK-75, and PP-121.

d. mTOR Inhibitors

[0301] The terms "mTOR inhibitor" or "agent that inhibits mTOR" refers to a nucleic acid, peptide, compound, or small organic molecule that inhibits at least one activity of an mTOR protein, such as, for example, the serine/threonine protein kinase activity on at least one of its substrates (e.g., p70S6 kinase 1, 4E-BP1, AKT/PKB and eEF2). mTOR inhibitors are able to bind directly to and inhibit mTORC1, mTORC2 or both mTORC1 and mTORC2.

[0302] Inhibition of mTORC1 and/or mTORC2 activity can be determined by a reduction in signal transduction of the PI3K/Akt/mTOR pathway. A wide variety of readouts can be utilized to establish a reduction of the output of such signaling pathway. Some non-limiting exemplary readouts include (1) a decrease in phosphorylation of Akt at residues, including but not limited to S473 and T308; (2) a decrease in activation of Akt as evidenced, for example, by a reduction of phosphorylation of Akt substrates including but not limited to FoxO1/O3a T24/32, GSK3 α/β , S21/9, and TSC2 T1462; (3) a decrease in phosphorylation of signaling molecules downstream of mTOR, including but not limited to ribosomal S6 S240/244, 70S6K T389, and 4EBP1 T37/46; and (4) inhibition of proliferation of cancerous cells. The mTOR inhibitors may be active site inhibitors. These are mTOR inhibitors that bind to the ATP binding site (also referred to as ATP binding pocket) of mTOR and inhibit the catalytic activity of both mTORC1 and mTORC2. One class of active site inhibitors suitable for use in the T cell manufacturing methods contemplated herein are dual specificity inhibitors that target and directly inhibit both PI3K and mTOR. Dual specificity inhibitors bind to both the ATP binding site of mTOR and PI3K. Illustrative examples of such inhibitors include, but are not limited to: imidazoquinazolines, wortmannin, LY294002, PI-103 (Cayman Chemical), SF1126 (Semafore), BGT226 (Novartis), XL765 (Exelixis) and NVP-BEZ235 (Novartis).

[0303] Another class of mTOR active site inhibitors suitable for use in the methods contemplated herein selectively inhibit mTORC1 and mTORC2 activity relative to one or more type I phosphatidylinositol 3-kinases, e.g., PI3 kinase α , β , γ , or δ . These active site inhibitors bind to the active site of mTOR but not PI3K. Illustrative examples of such inhibitors include, but are not limited to: pyrazolopyrimidines, Torin1 (Guertin and Sabatini), PP242 (2-(4-Amino-1-isopropyl-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-1H-indol-5-ol), PP30, Ku-0063794, WAY-600 (Wyeth), WAY-687 (Wyeth), WAY-354

(Wyeth), and AZD8055 (Liu et al., Nature Review, 8, 627-644, 2009). I

[0304] A selective mTOR inhibitor refers to an agent that exhibits a 50% inhibitory concentration (IC₅₀) with respect to mTORC1 and/or mTORC2, that is at least 10-fold, at least 20-fold, at least 50-fold, at least 100-fold, at least 1000-fold, or more, lower than the inhibitor's IC₅₀ with respect to one, two, three, or more type I PI3-kinases or to all of the type I PI3-kinases.

[0305] Another class of mTOR inhibitors contemplated are referred to herein as "rapalogs". As used herein the term "rapalogs" refers to compounds that specifically bind to the mTOR FRB domain (FKBP rapamycin binding domain), are structurally related to rapamycin, and retain the mTOR inhibiting properties. The term rapalogs excludes rapamycin. Rapalogs include esters, ethers, oximes, hydrazones, and hydroxylamines of rapamycin, as well as compounds in which functional groups on the rapamycin core structure have been modified, for example, by reduction or oxidation. Pharmaceutically acceptable salts of such compounds are also considered to be rapamycin derivatives. Illustrative examples of rapalogs suitable for use in the methods contemplated herein include, without limitation, temsirolimus (CC1779), everolimus (RAD001), deforolimus (AP23573), AZD8055 (AstraZeneca), and OSI-027 (OSI). The agent may be the mTOR inhibitor rapamycin (sirolimus).

[0306] Exemplary mTOR inhibitors for use may inhibit either mTORC1, mTORC2 or both mTORC1 and mTORC2 with an IC₅₀ (concentration that inhibits 50% of the activity) of about 200 nM or less, preferably about 100 nM or less, even more preferably about 60 nM or less, about 25 nM, about 10 nM, about 5 nM, about 1 nM, 100 μ M, 50 μ M, 25 μ M, 10 μ M, 1 μ M, or less. A mTOR inhibitor for use inhibits either mTORC1, mTORC2 or both mTORC1 and mTORC2 with an IC₅₀ from about 2 nM to about 100 nM, more preferably from about 2 nM to about 50 nM, even more preferably from about 2 nM to about 15 nM.

[0307] Exemplary mTOR inhibitors may inhibit either PI3K and mTORC1 or mTORC2 or both mTORC1 and mTORC2 and PI3K with an IC₅₀ (concentration that inhibits 50% of the activity) of about 200 nM or less, preferably about 100 nM or less, even more preferably about 60 nM or less, about 25 nM, about 10 nM, about 5 nM, about 1 nM, 100 μ M, 50 μ M, 25 μ M, 10 μ M, 1 μ M, or less. A mTOR inhibitor for use may inhibit PI3K and mTORC1 or mTORC2 or both mTORC1 and mTORC2 and PI3K with an IC₅₀ from about 2 nM to about 100 nM, more preferably from about 2 nM to about 50 nM, even more preferably from about 2 nM to about 15 nM.

[0308] Further illustrative examples of mTOR inhibitors suitable for use contemplated herein include, but are not limited to AZD8055, INK128, rapamycin, PF-04691502, and everolimus.

[0309] mTOR has been shown to demonstrate a robust and specific catalytic activity toward the physiological substrate proteins, p70 S6 ribosomal protein kinase I (p70S6K1) and eIF4E binding protein 1 (4EBP1) as measured by phosphor-specific antibodies in Western blotting.

[0310] The inhibitor of the PI3K/AKT/mTOR pathway may be a s6 kinase inhibitor selected from the group consisting of: BI-D1870, H89, PF-4708671, FMK, and AT7867.

I. COMPOSITIONS AND FORMULATIONS

[0311] The compositions contemplated herein may comprise one or more polypeptides, polynucleotides, vectors comprising same, genetically modified immune effector cells, *etc.*, as contemplated herein. Compositions include, but are not limited to pharmaceutical compositions. A "pharmaceutical composition" refers to a composition formulated in pharmaceutically-acceptable or physiologically-acceptable solutions for administration to a cell or an animal, either alone, or in combination with one or more other modalities of therapy. It will also be understood that, if desired, the compositions may be administered in combination with other agents as well, such as, *e.g.*, cytokines, growth factors, hormones, small molecules, chemotherapeutics, pro-drugs, drugs, antibodies, or other various pharmaceutically-active agents. There is virtually no limit to other components that may also be included in the compositions, provided that the additional agents do not adversely affect the ability of the composition to deliver the intended therapy.

[0312] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0313] As used herein "pharmaceutically acceptable carrier, diluent or excipient" includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, surfactant, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals. Exemplary pharmaceutically acceptable carriers include, but are not limited to, to sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; tragacanth; malt; gelatin; talc; cocoa butter, waxes, animal and vegetable fats, paraffins, silicones, bentonites, silicic acid, zinc oxide; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and

polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and any other compatible substances employed in pharmaceutical formulations.

[0314] In particular embodiments, compositions comprise an amount of CAR-expressing immune effector cells contemplated herein. As used herein,

the term "amount" refers to "an amount effective" or "an effective amount" of a genetically modified therapeutic cell, *e.g.*, T cell, to achieve a beneficial or desired prophylactic or therapeutic result, including clinical results.

[0315] A "prophylactically effective amount" refers to an amount of a genetically modified therapeutic cell effective to achieve the desired prophylactic result. Typically but not necessarily, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount is less than the therapeutically effective amount.

[0316] A "therapeutically effective amount" of a genetically modified therapeutic cell may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the stem and progenitor cells to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the virus or transduced therapeutic cells are outweighed by the therapeutically beneficial effects. The term "therapeutically effective amount" includes an amount that is effective to "treat" a subject (*e.g.*, a patient). When a therapeutic amount is indicated, the precise amount of the compositions to be administered can be determined by a physician with consideration of individual differences in age, weight, tumor size, extent of infection or metastasis, and condition of the patient (subject). It can generally be stated that a pharmaceutical composition comprising the T cells described herein may be administered at a dosage of 10^2 to 10^{10} cells/kg body weight, preferably 10^5 to 10^6 cells/kg body weight, including all integer values within those ranges. The number of cells will depend upon the ultimate use for which the composition is intended as will the type of cells included therein. For uses provided herein, the cells are generally in a volume of a liter or less, can be 500 mL or less, even 250 mL or 100 mL or less. Hence the density of the desired cells is typically greater than 10^6 cells/mL and generally is greater than 10^7 cells/mL, generally 10^8 cells/mL or greater. The clinically relevant number of immune cells can be apportioned into multiple infusions that cumulatively equal or exceed 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , or 10^{12} cells. Particularly since all the infused cells will be redirected to a particular target antigen (*e.g.*, κ or λ light chain), lower numbers of cells, in the range of 10^6 /kilogram (10^6 - 10^{11} per patient) may be administered. CAR expressing cell compositions may be administered multiple times at dosages within these ranges. The cells may be allogeneic, syngeneic,

xenogeneic, or autologous to the patient undergoing therapy. If desired, the treatment may also include administration of mitogens (*e.g.*, PHA) or lymphokines, cytokines, and/or chemokines (*e.g.*, IFN- γ , IL-2, IL-12, TNF- α , IL-18, and TNF- β , GM-CSF, IL-4, IL-13, Flt3-L, RANTES, MIP1 α , etc.) as described herein to enhance induction of the immune response.

[0317] Generally, compositions comprising the cells activated and expanded as described herein may be utilized in the treatment and prevention of diseases that arise in individuals who are immunocompromised. In particular, compositions comprising the CAR-modified T cells contemplated herein are used in the treatment of B cell malignancies. The CAR-modified T cells may be administered either alone, or as a pharmaceutical composition in combination with carriers, diluents, excipients, and/or with other components such as IL-2 or other cytokines or cell populations. Pharmaceutical compositions contemplated herein comprise an amount of genetically modified T cells, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients.

[0318] Pharmaceutical compositions comprising a CAR-expressing immune effector cell population, such as T cells, may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (*e.g.*, aluminum hydroxide); and preservatives. Compositions of the present invention are preferably formulated for parenteral administration, *e.g.*, intravascular (intravenous or intraarterial), intraperitoneal or intramuscular administration.

[0319] The liquid pharmaceutical compositions, whether they be solutions, suspensions or other like form, may include one or more of the following: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. An injectable pharmaceutical composition is preferably sterile.

[0320] Compositions contemplated herein comprise an effective amount of CAR-expressing immune effector cells, alone or in combination with one or more therapeutic agents. Thus, the CAR-expressing immune effector cell compositions may be administered alone or in combination with other known cancer treatments, such as radiation therapy, chemotherapy, transplantation, immunotherapy, hormone therapy, photodynamic therapy, *etc.* The compositions may also be administered in combination with antibiotics. Such therapeutic agents may be accepted in the art as a standard treatment

for a particular disease state as described herein, such as a particular cancer. Exemplary therapeutic agents contemplated include cytokines, growth factors, steroids, NSAIDs, DMARDs, anti-inflammatories, chemotherapeutics, radiotherapeutics, therapeutic antibodies, or other active and ancillary agents.

[0321] Compositions comprising CAR-expressing immune effector cells disclosed herein may be administered in conjunction with any number of chemotherapeutic agents. Illustrative examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN™); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine resins; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2', 2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g. paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, N.J.) and doxetaxel (TAXOTERE®, Rhne-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylomithine (DMFO); retinoic acid derivatives such as Targretin™ (bexarotene), Panretin™ (alitretinoin); ONTAK™ (denileukin diftitox); esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on cancers such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0322] A variety of other therapeutic agents may be used in conjunction with the compositions described herein. The composition comprising CAR-expressing immune effector cells may be administered with an anti-inflammatory agent. Anti-inflammatory agents or drugs include, but are not limited to, steroids and glucocorticoids (including betamethasone, budesonide, dexamethasone, hydrocortisone acetate, hydrocortisone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone), nonsteroidal anti-inflammatory drugs (NSAIDs) including aspirin, ibuprofen, naproxen, methotrexate, sulfasalazine, leflunomide, anti-TNF medications, cyclophosphamide and mycophenolate.

[0323] Other exemplary NSAIDs are chosen from the group consisting of ibuprofen, naproxen, naproxen sodium, Cox-2 inhibitors such as VIOXX® (rofecoxib) and CELEBREX® (celecoxib), and sialylates. Exemplary analgesics are chosen from the group consisting of acetaminophen, oxycodone, tramadol or propoxyphene hydrochloride. Exemplary glucocorticoids are chosen from the group consisting of cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, or prednisone. Exemplary biological response modifiers include molecules directed against cell surface markers (e.g., CD4, CD5, etc.), cytokine inhibitors, such as the TNF antagonists (e.g., etanercept (ENBREL®), adalimumab (HUMIRA®) and infliximab (REMICADE®), chemokine inhibitors and adhesion molecule inhibitors. The biological response modifiers include monoclonal antibodies as well as recombinant forms of molecules. Exemplary DMARDs include azathioprine, cyclophosphamide, cyclosporine, methotrexate, penicillamine, leflunomide, sulfasalazine, hydroxychloroquine, Gold (oral (auranofin) and intramuscular) and minocycline.

[0324] Illustrative examples of therapeutic antibodies suitable for combination with the CAR modified T cells contemplated herein, include but are not limited to, bavixumab, bevacizumab (avastin), bivatuzumab, blinatumomab, conatumumab, daratumumab, duligotumab, dacetuzumab, dalotuzumab, elotuzumab (HuLuc63), gemtuzumab, ibritumomab, indatuximab, inotuzumab, lorvotuzumab, lucatumumab, milatuzumab, moxetumomab, ocaratuzumab, ofatumumab, rituximab, siltuximab, teprotumumab, and ublituximab.

[0325] In certain embodiments, the compositions described herein are administered in conjunction with a cytokine. By "cytokine" as used herein is meant a generic term for proteins released by one cell population that act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormones such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor-alpha and -beta; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF-beta; platelet-growth factor; transforming growth factors (TGFs) such as TGF-alpha and TGF-beta; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon-alpha, beta, and -gamma; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1alpha, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12; IL-15, IL-21, a tumor necrosis factor such as TNF-alpha or TNF-beta; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture, and biologically active equivalents of the native sequence cytokines.

[0326] A composition may comprise CAR T cells contemplated herein that are cultured in the presence of a PI3K inhibitor as disclosed herein and express one or more of the following markers: CD3, CD4, CD8, CD28, CD45RA, CD45RO, CD62, CD127, and HLA-DR can be further isolated by positive or negative selection techniques. A composition may comprise a specific subpopulation of T cells, expressing one or more of the markers selected from the group consisting of CD62L, CCR7, CD28, CD27, CD122, CD127, CD197; and CD38 or CD62L, CD127, CD197, and CD38, is further isolated by positive or negative selection techniques. Compositions may not express or do not substantially express one or more of the following markers: CD57, CD244, CD160, PD-1, CTLA4, TIM3, and LAG3.

[0327] Expression of one or more of the markers selected from the group consisting of CD62L, CD127, CD197, and CD38 is increased at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, at least 10 fold, at least 25 fold, or more compared to a population of T cells activated and expanded without a PI3K inhibitor.

[0328] Expression of one or more of the markers selected from the group consisting of CD57, CD244, CD160, PD-1, CTLA4, TIM3, and LAG3 is decreased at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, at least 10 fold, at least 25 fold, or more compared to a population of T cells activated and expanded with a PI3K inhibitor.

J. THERAPEUTIC METHODS

[0329] The genetically modified immune effector cells contemplated herein provide improved methods of adoptive immunotherapy for use in the treatment of B cell related conditions that include, but are not limited to immunoregulatory conditions and hematological malignancies.

[0330] The specificity of a primary immune effector cell may be redirected to B cells by genetically modifying the primary immune effector cell with a CAR contemplated herein. A viral vector may be used to genetically modify an immune effector cell with a particular polynucleotide encoding a CAR comprising a murine anti-BCMA antigen binding domain that binds a BCMA polypeptide; a hinge domain; a transmembrane (TM) domain, a short oligo- or polypeptide linker, that links the TM domain to the intracellular signaling domain of the CAR; and one or more intracellular co-stimulatory signaling domains; and a primary signaling domain.

[0331] Also disclosed is type of cellular therapy where T cells are genetically modified to express a CAR that targets BCMA expressing B cells. Anti-BCMA CAR T cells may be cultured in the presence of IL-2 and a PI3K inhibitor to increase the therapeutic properties and persistence of the CAR T cells. The CAR T cell are then infused to a recipient in need thereof. The infused cell is able to kill disease causing B cells in the recipient. Unlike antibody therapies, CAR T cells are able to replicate *in vivo* resulting in long-term persistence that can lead to sustained cancer therapy.

[0332] The CAR T cells may undergo robust *in vivo* T cell expansion and can persist for an extended amount of time. The CAR T cells may evolve into specific memory T cells that can be reactivated to inhibit any additional tumor formation or growth.

[0333] Compositions comprising immune effector cells comprising the CARs contemplated herein may be used in the treatment of conditions associated with abnormal B cell activity.

[0334] Illustrative examples of conditions that can be treated, prevented or ameliorated using the immune effector cells comprising the CARs contemplated herein include, but are not limited to: systemic lupus erythematosus, rheumatoid arthritis, myasthenia gravis, autoimmune hemolytic anemia, idiopathic thrombocytopenia purpura, anti-phospholipid syndrome, Chagas' disease, Grave's disease, Wegener's granulomatosis, poly-arteritis nodosa, Sjogren's syndrome,

pemphigus vulgaris, scleroderma, multiple sclerosis, anti-phospholipid syndrome, ANCA associated vasculitis, Good-pasture's disease, Kawasaki disease, and rapidly progressive glomerulonephritis.

[0335] The modified immune effector cells may also have application in plasma cell disorders such as heavy-chain disease, primary or immunocyte-associated amyloidosis, and monoclonal gammopathy of undetermined significance (MGUS).

[0336] As used herein, "B cell malignancy" refers to a type of cancer that forms in B cells (a type of immune system cell) as discussed *infra*.

[0337] Compositions comprising CAR-modified T cells contemplated herein may be used in the treatment of hematologic malignancies, including but not limited to B cell malignancies such as, for example, multiple myeloma (MM) and non-Hodgkin's lymphoma (NHL).

[0338] Multiple myeloma is a B cell malignancy of mature plasma cell morphology characterized by the neoplastic transformation of a single clone of these types of cells. These plasma cells proliferate in BM and may invade adjacent bone and sometimes the blood. Variant forms of multiple myeloma include overt multiple myeloma, smoldering multiple myeloma, plasma cell leukemia, non-secretory myeloma, IgD myeloma, osteosclerotic myeloma, solitary plasmacytoma of bone, and extramedullary plasmacytoma (see, for example, Braunwald, et al. (eds), Harrison's Principles of Internal Medicine, 15th Edition (McGraw-Hill 2001)).

[0339] Non-Hodgkin lymphoma encompasses a large group of cancers of lymphocytes (white blood cells). Non-Hodgkin lymphomas can occur at any age and are often marked by lymph nodes that are larger than normal, fever, and weight loss. There are many different types of non-Hodgkin lymphoma. For example, non-Hodgkin's lymphoma can be divided into aggressive (fast-growing) and indolent (slow-growing) types. Although non-Hodgkin lymphomas can be derived from B cells and T-cells, as used herein, the term "non-Hodgkin lymphoma" and "B cell non-Hodgkin lymphoma" are used interchangeably. B cell non-Hodgkin lymphomas (NHL) include Burkitt lymphoma, chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), diffuse large B cell lymphoma, follicular lymphoma, immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, and mantle cell lymphoma. Lymphomas that occur after bone marrow or stem cell transplantation are usually B cell non-Hodgkin lymphomas.

[0340] Chronic lymphocytic leukemia (CLL) is an indolent (slow-growing) cancer that causes a slow increase in immature white blood cells called B lymphocytes, or B cells.

Cancer cells spread through the blood and bone marrow, and can also affect the lymph nodes or other organs such as the liver and spleen. CLL eventually causes the bone marrow to fail. Sometimes, in later stages of the disease, the disease is called small lymphocytic lymphoma.

[0341] Methods comprising administering a therapeutically effective amount of CAR-expressing immune effector cells contemplated herein or a composition comprising the same, to a patient in need thereof, alone or in combination with one or more therapeutic agents, are provided. The cells may be used in the treatment of patients at risk for developing a condition associated with abnormal B cell activity or a B cell malignancy. Thus, the present disclosure provides methods for the treatment or prevention of a condition associated with abnormal B cell activity or a B cell malignancy comprising administering to a subject in need thereof, a therapeutically effective amount of the CAR-modified cells contemplated herein.

[0342] As used herein, the terms "individual" and "subject" are often used interchangeably and refer to any animal that exhibits a symptom of a disease, disorder, or condition that can be treated with the gene therapy vectors, cell-based therapeutics, and methods disclosed elsewhere herein. A subject may include any animal that exhibits symptoms of a disease, disorder, or condition of the hematopoietic system, e.g., a B cell malignancy, that can be treated with the gene therapy vectors, cell-based therapeutics, and methods disclosed elsewhere herein. Suitable subjects (e.g., patients) include laboratory animals (such as mouse, rat, rabbit, or guinea pig), farm animals, and domestic animals or pets (such as a cat or dog). Non-human primates and, preferably, human patients, are included. Typical subjects include human patients that have a B cell malignancy, have been diagnosed with a B cell malignancy, or are at risk or having a B cell malignancy. As used herein, the term "patient" refers to a subject that has been diagnosed with a particular disease, disorder, or condition that can be treated with the gene therapy vectors, cell-based therapeutics, and methods disclosed elsewhere herein.

[0343] As used herein "treatment" or "treating," includes any beneficial or desirable effect on the symptoms or pathology of a disease or pathological condition, and may include even minimal reductions in one or more measurable markers of the disease or condition being treated. Treatment can involve optionally either the reduction or amelioration of symptoms of the disease or condition, or the delaying of the progression of the disease or condition. "Treatment" does not necessarily indicate complete eradication or cure of the disease or condition, or associated symptoms thereof.

[0344] As used herein, "prevent," and similar words such as "prevented," "preventing" etc., indicate an approach for preventing, inhibiting, or reducing the likelihood of the occurrence or recurrence of, a disease or condition. It also refers to delaying the onset or recurrence of a disease or condition or delaying the occurrence or recurrence of the symptoms of a disease or condition. As used herein, "prevention" and similar words also includes reducing the intensity, effect, symptoms and/or burden of a disease or condition prior to onset or recurrence of the disease or condition.

[0345] By "enhance" or "promote," or "increase" or "expand" refers generally to the ability of a composition contemplated herein, e.g., a genetically modified T cell or vector encoding a CAR, to produce, elicit, or cause a greater physiological response (*i.e.*, downstream effects) compared to the response caused by either vehicle or a control molecule/composition. A measurable physiological response may include an increase in T cell expansion, activation, persistence, and/or an increase in cancer cell killing ability, among others apparent from the understanding in the art and the description herein. An "increased" or "enhanced" amount is typically a "statistically significant" amount, and may include an increase that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1, e.g., 1.5, 1.6, 1.7, 1.8, etc.) the response produced by vehicle or a control composition.

[0346] By "decrease" or "lower," or "lessen," or "reduce," or "abate" refers generally to the ability of composition contemplated herein to produce, elicit, or cause a lesser physiological response (*i.e.*, downstream effects) compared to the response caused by either vehicle or a control molecule/composition. A "decrease" or "reduced" amount is typically a "statistically significant" amount, and may include an decrease that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1, e.g., 1.5, 1.6, 1.7, 1.8, etc.) the response (reference response) produced by vehicle, a control composition, or the response in a particular cell lineage.

[0347] By "maintain," or "preserve," or "maintenance," or "no change," or "no substantial change," or "no substantial decrease" refers generally to the ability of a composition contemplated herein to produce, elicit, or cause a lesser physiological response (*i.e.*, downstream effects) in a cell, as compared to the response caused by either vehicle, a control molecule/composition, or the response in a particular cell lineage. A comparable response is one that is not significantly different or measurable different from the reference response.

[0348] A method of treating a B cell related condition in a subject in need thereof comprises administering an effective amount, e.g., therapeutically effective amount of a composition comprising genetically modified immune effector cells contemplated herein. The quantity and frequency of administration will be determined by such factors as the condition of the patient, and the type and severity of the patient's disease, although appropriate dosages may be determined by clinical trials.

[0349] The amount of T cells in the composition administered to a subject may be at least 0.1×10^5 cells, at least 0.5×10^5 cells, at least 1×10^5 cells, at least 5×10^5 cells, at least 1×10^6 cells, at least 0.5×10^7 cells, at least 1×10^7 cells, at least 0.5×10^8 cells, at least 1×10^8 cells, at least 0.5×10^9 cells, at least 1×10^9 cells, at least 2×10^9 cells, at least 3×10^9 cells, at least 4×10^9 cells, at least 5×10^9 cells, or at least 1×10^{10} cells. In particular embodiments, about 1×10^7 CAR T cells to about 1×10^9 CAR T cells, about 2×10^7 CAR T cells to about 0.9×10^9 CAR T cells, about 3×10^7 CAR T cells to about 0.8×10^9 CAR T cells, about 4×10^7 CAR T cells to about 0.7×10^9 CAR T cells, about 5×10^7 CAR T cells to about 0.6×10^9 CAR T cells, or about 5×10^7 CAR T cells to about 0.5×10^9 CAR T cells are administered to a subject.

[0350] The amount of T cells in the composition administered to a subject may be at least 0.1×10^4 cells/kg of bodyweight, at least 0.5×10^4 cells/kg of bodyweight, at least 1×10^4 cells/kg of bodyweight, at least 5×10^4 cells/kg of bodyweight, at least 1×10^5 cells/kg of bodyweight, at least 0.5×10^6 cells/kg of bodyweight, at least 1×10^6 cells/kg of bodyweight, at least 0.5×10^7 cells/kg of bodyweight, at least 1×10^7 cells/kg of bodyweight, at least 0.5×10^8 cells/kg of bodyweight, at least 1×10^8 cells/kg of bodyweight, at least 2×10^8 cells/kg of bodyweight, at least 3×10^8 cells/kg of bodyweight, at least 4×10^8 cells/kg of bodyweight, at least 5×10^8 cells/kg of bodyweight, or at least 1×10^9 cells/kg of bodyweight. About 1×10^6 CAR T cells/kg of bodyweight to about 1×10^8 CAR T cells/kg of bodyweight, about 2×10^6 CAR T cells/kg of bodyweight to about 0.9×10^8 CAR T cells/kg of bodyweight, about 3×10^6 CAR T cells/kg of bodyweight to about 0.8×10^8 CAR T cells/kg of bodyweight, about 4×10^6 CAR T cells/kg of bodyweight to about 0.7×10^8 CAR T cells/kg of bodyweight, about 5×10^6 CAR T cells/kg of bodyweight to about 0.6×10^8 CAR T cells/kg of bodyweight, or about 5×10^6 CAR T cells/kg of bodyweight to about 0.5×10^8 CAR T cells/kg of bodyweight may be administered to a subject.

[0351] One of ordinary skill in the art would recognize that multiple administrations of the compositions may be required to effect the desired therapy. For example a composition may be administered 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more times over a span of 1 week, 2 weeks, 3 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 1 year, 2 years, 5, years, 10 years, or more.

[0352] It may be desirable to administer activated immune effector cells to a subject and then subsequently redraw blood (or have an apheresis performed), activate immune effector cells therefrom, and reinfuse the patient with these activated and expanded immune effector cells. This process can be carried out multiple times every few weeks. In certain embodiments, immune effector cells can be activated from blood draws of from 10cc to 400cc. Immune effector cells may be activated from blood draws of 20cc, 30cc, 40cc, 50cc, 60cc, 70cc, 80cc, 90cc, 100cc, 150cc, 200cc, 250cc, 300cc, 350cc, or 400cc or more. Not to be bound by theory, using this multiple blood draw/multiple reinfusion protocol may serve to select out certain populations of immune effector cells.

[0353] The administration of the compositions contemplated herein may be carried out in any convenient manner, including by aerosol inhalation, injection, ingestion, transfusion, implantation or transplantation. Preferably, compositions

are administered parenterally. The phrases "parenteral administration" and "administered parenterally" as used herein refers to modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravascular, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intratumoral, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion. In one embodiment, the compositions contemplated herein are administered to a subject by direct injection into a tumor, lymph node, or site of infection.

[0354] A subject in need thereof may be administered an effective amount of a composition to increase a cellular immune response to a B cell related condition in the subject. The immune response may include cellular immune responses mediated by cytotoxic T cells capable of killing infected cells, regulatory T cells, and helper T cell responses. Humoral immune responses, mediated primarily by helper T cells capable of activating B cells thus leading to antibody production, may also be induced. A variety of techniques may be used for analyzing the type of immune responses induced by the compositions, which are well described in the art; e.g., Current Protocols in Immunology, Edited by: John E. Coligan, Ada M. Kruisbeek, David H. Margulies, Ethan M. Shevach, Warren Strober (2001) John Wiley & Sons, NY, N.Y.

[0355] In the case of T cell-mediated killing, CAR-ligand binding initiates CAR signaling to the T cell, resulting in activation of a variety of T cell signaling pathways that induce the T cell to produce or release proteins capable of inducing target cell apoptosis by various mechanisms. These T cell-mediated mechanisms include (but are not limited to) the transfer of intracellular cytotoxic granules from the T cell into the target cell, T cell secretion of pro-inflammatory cytokines that can induce target cell killing directly (or indirectly via recruitment of other killer effector cells), and up regulation of death receptor ligands (e.g. FasL) on the T cell surface that induce target cell apoptosis following binding to their cognate death receptor (e.g. Fas) on the target cell.

[0356] Also provided is a method of treating a subject diagnosed with a B cell related condition comprising removing immune effector cells from a subject diagnosed with a BCMA-expressing B cell related condition, genetically modifying said immune effector cells with a vector comprising a nucleic acid encoding a CAR as contemplated herein, thereby producing a population of modified immune effector cells, and administering the population of modified immune effector cells to the same subject. Preferably, the immune effector cells comprise T cells.

[0357] Also provided are methods for stimulating an immune effector cell mediated immune modulator response to a target cell population in a subject comprising the steps of administering to the subject an immune effector cell population expressing a nucleic acid construct encoding a CAR molecule.

[0358] The methods for administering the cell compositions described herein includes any method which is effective to result in reintroduction of *ex vivo* genetically modified immune effector cells that either directly express a CAR in the subject or on reintroduction of the genetically modified progenitors of immune effector cells that on introduction into a subject differentiate into mature immune effector cells that express the CAR. One method comprises transducing peripheral blood T cells *ex vivo* with a nucleic acid construct and returning the transduced cells into the subject.

EXAMPLES

EXAMPLE 1

CONSTRUCTION OF BCMA CARS

[0359] CARs containing murine anti-BCMA scFv antibodies were designed to contain an MND promoter operably linked to anti-BCMA scFv, a hinge and transmembrane domain from CD8 α and a CD137 co-stimulatory domains followed by the intracellular signaling domain of the CD3 ζ chain. See, e.g., Figure 1. The BCMA CAR shown in Figure 1 comprises a CD8 α signal peptide (SP) sequence for the surface expression on immune effector cells. The polynucleotide sequence of an exemplary BCMA CAR is set forth in SEQ ID NO: 10; an exemplary polypeptide sequences of a BCMA CAR is set forth in SEQ ID NO: 9; and a vector map of an exemplary CAR construct is shown in Figure 1. Table 3 shows the Identity, Genbank Reference, Source Name and Citation for the various nucleotide segments of an BCMA CAR lentiviral vector.

Table 3.

Nucleatides	Identity	GenBank Reference	Source Name	Citation
1-185	pUC19 plasmid backbone	Accession #L09137.2 nt 1 - 185	pUC19	New England Biolabs
185-222	Linker	Not applicable	Synthetic	Not applicable

EP 3 230 321 B1

(continued)

Nucleotides	Identity	GenBank Reference	Source Name	Citation
223-800	CMV	Not Applicable	pHCMV	(1994) PNAS 91: 9564-68
801-1136	R, U5, PBS, and packaging sequences	Accession #M19921.2 nt 454-789	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
1137-1139	Gag start codon (ATG) changed to stop codon (TAG)	Not Applicable	Synthetic	Not applicable
1140-1240	HIV-1 gag sequence	Accession #M19921.2 nt 793-893	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
1241-1243	HIV -1 gag sequence changed to a second stop codon	Not Applicable	Synthetic	Not applicable
1244-1595	HIV-1 gag sequence	Accession #M19921.2 nt 897-1248	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
1596-1992	HIV-1 pol cPPT/CTS	Accession #M19921.2 nt 4745-5125	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
1993-2517	HIV-1, isolate HXB3 env region (RRE)	Accession #M14100.1 nt 1875-2399	PgTAT-CMV	Malim, M. H. Nature (1988) 335:181-183
2518-2693	HIV-1 env sequences S/A	Accession #M19921.2 nt 8290-8470	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
2694-2708	Linker	Not applicable	Synthetic	Not applicable
2709-3096	MND	Not applicable	pccl-c-MNDU3c-x2	Challita et al. (1995) J.Virol. 69: 748-755
3097-3124	Linker	Not applicable	Synthetic	Not applicable
3125-3187	Signal peptide	Accession # NM_001768	CD8a signal peptide	Not applicable
3188-3934	BCMA02 scFv (V _L 1-linker-V _H 0)	Not applicable	Synthetic	Not applicable
3935-4141	CD8a hinge and TM	Accession # NM_001768	CD8a hinge and TM	Milone et al (2009) Mol Ther 17(8): 1453-64
4144-4269	CD137 (4-1BB) signaling domain	Accession # NM_001561	CD137 signaling domain	Milone et al (2009) Mol Ther 17(8): 1453-64
4270-4606	CD3-ζ signaling domain	Accession # NM_000734	CD3-ζ signaling domain	Milone et al (2009) Mol Ther 17(8): 1453-64

(continued)

Nucleotides	Identity	GenBank Reference	Source Name	Citation
4607-4717	HIV-1 ppt and part of 3'U3	Accession #M19921.2 nt 9005-9110	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
4718-4834	HIV-1 part of U3 (399bp deletion) and R	Accession #M19921.2 nt 9511-9627	pNL4-3	Maldarelli, et.al. (1991) J Virol: 65(11): 5732-43
4835-4858	Synthetic polyA	Not applicable	Synthetic	Levitt, N. Genes & Dev (1989) 3:1019-1025
4859-4877	Linker	Not applicable	Synthetic	Not Applicable
4878-7350	pUC19 backbone	Accession #L09137.2 nt 2636-2686	pUC19	New England Biolabs

EXAMPLE 2**EVALUATION OF A MURINE BCMA CAR****Introduction**

[0360] Adoptive transfer of T cells genetically engineered with chimeric antigen receptors (CAR) has emerged as a promising approach to treat cancers. A CAR is an artificial molecule comprised of an antigen reactive single chain variable fragment (scFv) fused to T cell signaling domains via a transmembrane region. In this example, a CAR molecule specific to B cell maturation antigen (BCMA) was evaluated. BCMA is expressed on multiple myeloma, plasmacytoma, and some lymphomas yet normal expression is limited to plasma cells (Avery *et al.*, 2003; Carpenito *et al.*, 2009; Chiu *et al.*, 2007).

[0361] Anti-BCMA02 CAR was constructed using sequences from a mouse anti-BCMA antibody (C11D5.3). Anti-BCMA10 CAR was constructed using modified sequences and is a "humanized" version of anti-BCMA02 CAR. In a series of *in vitro* assays, anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells both exhibited tumor specificity, high CAR expression, and caused potent reactivity to antigen expressing targets. Anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells were shown to have comparable reactivity to BCMA-expressing tumor cell lines. Although both anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells were capable of causing regressions in a mouse tumor model, anti-BCMA10 CAR T cells displayed antigen-independent inflammatory cytokine secretion, and thus, have the potential to cause clinical toxicities associated high cytokine levels.

Results*Tonic inflammatory cytokine release from anti-BCMA10 T cells associated with apoptosis*

[0362] BCMA protein is detectable in the serum of patients with multiple myeloma (Sanchez *et al.*, 2012). Average serum BCMA in multiple myeloma patients was 10ng/mL but peaked at levels up to 100ng/mL. The impact of physiological soluble BCMA levels on the anti-BCMA CAR T cell candidates was evaluated.

[0363] IFN γ release from anti-BCMA02 CAR T cells, anti-BCMA10 CAR T cells, and CAR19 Δ T cells was examined after a 24hour culture with soluble BCMA (Figure 2a). Anti-BCMA02 CAR T cells responded with minimal cytokine release after 24 hour culture with up to 1ug/mL BCMA. In contrast, anti-BCMA10 CAR T cells responded with increasing levels of IFN γ that were proportional to the concentration of soluble BCMA added to the culture. At 100ng/mL BCMA, the maximum levels reported in multiple myeloma patients, anti-BCMA10 CAR T cells secreted 82.1ng/ml IFN γ compared to 28.8ng/ml IFN γ secreted by anti-BCMA02 CAR T cells. IFN γ was even detected in several co-culture experiments with anti-BCMA10 CAR T cells plus control cell lines that lacked BCMA antigen (Figure 2b, K562 co-culture). These

data suggested that anti-BCMA10 CAR T cells had increased sensitivity to stimulation by soluble BCMA and the potential for antigen-independent cytokine responses in T cells.

[0364] The potential of tonic cytokine secretion from anti-BCMA02 CAR T cells, anti-BCMA10 CAR T cells (10 days from culture initiation), and CAR19 Δ T cells was examined. After manufacture of CAR T cells, growth media from anti-BCMA02 CAR T cell, anti-BCMA10 CAR T cell, and CAR19 Δ T cell cultures were analyzed for the presence of inflammatory cytokines. Despite the absence of antigen stimulation, anti-BCMA10 CAR T cell cultures contained greater than 10ng/mL IFN γ compared to less than 1ng/mL of IFN γ in anti-BCMA02 CAR T cell cultures (Figure 3). Anti-BCMA10 CAR T cell cultures also contained significantly ($p < 0.001$) more TNF α . To further quantify the amount of cytokine produced by anti-BCMA10 CAR T cells without antigen stimulation, cytokine release was measured from 5×10^4 CAR T cells during a 24 hour culture. anti-BCMA10 CAR T cells produced significantly higher amounts of inflammatory cytokines MIP1 α , IFN γ , GM-CSF, MIP1 β , IL-8, and TNF α compared to anti-BCMA02 CAR T cells (Figure 4, $p < 0.0001$). MIP1 α and IFN γ concentrations were the highest among all cytokines examined. Anti-BCMA10 CAR T cells produced 4.7ng MIP1 α /5 $\times 10^4$ cells/24 hours, 3.0ng IFN γ /5 $\times 10^4$ cells/24 hours and ~1ng/5 $\times 10^4$ cells/24 hours or less of the other cytokines. No significant differences in the anti-inflammatory cytokines IL-10, IL-2, and IL-4 were detected.

[0365] The expression of phenotypic markers of T cell activation at the end of anti-BCMA10 CAR T cell manufacturing were measured to examine whether tonic inflammatory cytokine secretion was indicative of a hyperactive state in anti-BCMA10 CAR T cells. HLA-DR and CD25 are surface markers that normally exhibit peak expression 12-24 hours after T cell activation and then diminish with time. CAR T cells prepared from three normal donors showed that an average 40 $\pm$ 2% of anti-BCMA02 CAR T cells expressed HLA-DR. The expression of HLA-DR in these cells was comparable to untransduced (43 $\pm$ 2.3%) T cells and CAR19 Δ (32 $\pm$ 2.2%) control T cells. In contrast, 88 $\pm$ 1.2% anti-BCMA10 CAR T cells expressed HLA-DR (Figure 5). Expression of another activation marker CD25 was also higher on anti-BCMA10 CAR T cells compared to anti-BCMA02 CAR T cells (53 $\pm$ 0.9% vs 35 $\pm$ 2.4%). Therefore, anti-BCMA10 CAR T cells exhibited phenotypic characteristics of activated T cells in the absence of added antigens.

[0366] Hyperactivity in T cells is often associated with activation-induced cell death (AICD) by apoptosis. Levels of activated caspase-3 were measured to examine whether hyperactivity of anti-BCMA10 CAR T cells could result in higher apoptotic levels compared to anti-BCMA02 CAR T cells. 48% of anti-BCMA10 CAR T cells from two donors had active caspase-3 compared to 16% of anti-BCMA02 CAR T cells (Figure 6). Thus, in the absence of added BCMA antigen, anti-BCMA10 CAR T cells contain a higher frequency of apoptotic cells associated with increased activation and inflammatory cytokine secretion compared to anti-BCMA02 CAR T cells.

[0367] anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells were evaluated for whether the CAR T cells could selectively respond to low BCMA levels or be cross reactive to an unrelated antigen in the human serum used for T cell growth. T anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells were maintained in media lacking human serum for two days and then switched into media containing fetal bovine serum (FBS), human serum (HABS), or HABS in the presence or absence of 100ng/mL soluble BCMA (Figure 7). IFN γ release was assayed 24 hours later by ELISA. Both anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells responded to soluble BCMA. However, anti-BCMA10 CAR T cells secreted 10-times more IFN γ than anti-BCMA02 CAR T cells. In the absence of BCMA, only anti-BCMA10 CAR T cells released IFN γ regardless of culture in fetal bovine serum (FBS) ($p = 0.0002$) or human AB serum (HABS) ($p = 0.0007$). These data suggested that inflammatory cytokine secretion was intrinsic to anti-BCMA10 CAR T cells.

Inferior anti-tumor function of anti-BCMA10 CAR T cells in mouse model of multiple myeloma

[0368] Hyperactivation and increased apoptosis could negatively impact CAR T cell persistence in patients and ultimately clinical efficacy. The anti-tumor function of anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells was examined in a mouse tumor model. NOD scid gamma (NSG) mice with ~100mm³ experimental sub-cutaneous human multiple myeloma (RPMI-8226) tumors were treated with 10⁷ anti-BCMA02 CAR T cells, 10⁷ anti-BCMA10 CAR T cells, or Bortezomib (velcade). RPMI-8226 growth was monitored with calipers. In two independent experiments (Figures 8a and 8b), Bortezomib controlled tumor growth compared to vehicle control animals. Animals adoptively transferred with anti-BCMA02 CAR T cells exhibited rapid and durable tumor clearance (inset graphs magnify early tumor regressions). Adoptive transfer of anti-BCMA10 CAR T cells also caused tumor regressions but was delayed in both experiments compared to anti-BCMA02 CAR T cells.

Conclusions

[0369] anti-BCMA02 CAR T cells and anti-BCMA10 CAR T cells exhibited comparable antitumor function in in vitro assays, but anti-BCMA10 CAR T cells had characteristics that could negatively impact safety and efficacy in patient treatment. Anti-BCMA10 CAR T cells responded robustly with inflammatory cytokine secretion after exposure to physiological levels of BCMA protein. Cytokine storm or cytokine release syndrome is a known clinical toxicity associated with CAR T cell therapies. Concerns over cytokine release to BCMA were worsened after observation of tonic activity

of anti-BCMA10 CAR T cells. Even in the absence of antigen-stimulation, anti-BCMA10 CAR T cells released high levels of inflammatory cytokines. Persistent cytokine secretion has the potential to cause substantial clinical toxicities as well as negatively impact anti-tumor function. Indeed, we found higher composition of apoptotic cells and inferior anti-tumor function in anti-BCMA10 CAR T cells compared to anti-BCMA02 CAR T cell cultures in a mouse model of multiple myeloma.

References

- [0370] Avery et al., (2003). BAFF selectively enhances the survival of plasmablasts generated from human memory B cells. *J Clin Invest*, 112(2), 286-297.
- [0371] Carpenito et al., (2009). Control of large, established tumor xenografts with genetically retargeted human T cells containing CD28 and CD137 domains. *Proc Natl Acad Sci USA*, 106(9), 3360-3365.
- [0372] Chiu et al., (2007). Hodgkin lymphoma cells express TACI and BCMA receptors and generate survival and proliferation signals in response to BAFF and APRIL. *Blood*, 109(2), 729-739.
- [0373] Sanchez et al. (2012). Serum B-cell maturation antigen is elevated in multiple myeloma and correlates with disease status and survival. *Br J Haematol*, 158(6), 727-738.

EXAMPLE 3

MINIMAL BCMA EXPRESSION ON LYMPHOMAS ACTIVATES ANTI-BCMA CAR T CELLS

[0374] The level of BCMA expression on lymphoma and leukemia cell lines (Daudi and Raji) was measured in order to determine if the expression was sufficient to activate anti-BCMA02 CAR T cells.

[0375] BCMA expression on lymphoma, leukemia, and multiple myeloma cells was quantitated using flow cytometry. In this assay, the relative BCMA expression on the cells was assessed by correlating the fluorescence intensity of BCMA expression to a known number of bound antibodies (antibody binding capacity, ABC). BCMA expression levels in the lymphoma cell lines were compared to BCMA expression levels a multiple myeloma cell line (RPMI-8226) known to activate anti-BCMA02 CAR T cells. 12590 ± 1275 BCMA02 molecules were expressed on the surface of RPMI-8226 cells. By contrast, Daudi cells expressed 1173 ± 234 BCMA02 molecules and JeKo-1 cells (a Mantle cell lymphoma cell line) expressed only 222 ± 138 BCMA02 molecules (Figure 9, circles).

[0376] In another set of experiments the activity of anti-BCMA02 CAR T cells to the minute levels of BCMA observed on lymphoma and leukemia cell lines was tested (Figure 9, boxes). Anti-BCMA02 CAR T cells were generated using standard methods and activity was assessed by IFN γ ELISA after co-culture with BCMA-positive and BCMA-negative tumor cell lines. Reactivity of anti-BCMA02 CAR T cells correlated with the relative amount of BCMA mRNA expression (above a threshold) and/or the density of the BCMA receptor on the surface of various tumor cell lines after co-culture (Figure 9). Little, if any, IFN γ is released upon co-culture of BCMA CAR T cells with BCMA-negative (BCMA-) tumor cell lines: myelogenous leukemia (K562), acute lymphoblastic leukemia (NALM-6 and NALM-16); Mantle cell lymphoma (REC-1); or Hodgkin's lymphoma (HDLM-2). In contrast, substantial amounts of IFN γ was released upon co-culture of BCMA02 CAR T cells with BCMA-positive (BCMA+) tumor cell lines: B cell chronic lymphoblastic leukemia (MEC-1), Mantle cell lymphoma (JeKo-1), Hodgkin's lymphoma (RPMI-6666), Burkitt's lymphoma (Daudi cells and Ramos cells), and multiple myeloma (RPMI-8226).

[0377] The reactivity of anti-BCMA02 CAR T cells to BCMA expressing Burkitt's lymphoma cells (Daudi cells) extended to *in vivo* animal studies. Daudi cells also express CD19. The *in vivo* activity of anti-BCMA02 CAR T cells was compared to the *in vivo* activity of anti-CD 19 CAR T cells. NOD scid gamma (NSG) mice were injected IV with 2×10^6 Daudi cells and allowed to accumulate a large systemic tumor burden before being treated with CAR T cells. CAR T cells were administered at 8 days and 18 days post-tumor induction (Figures 10A and 10B, respectively). The vehicle and negative control (anti-CD19 Δ CAR T cells) failed to prevent tumor growth, as shown by log-phase increases in bioluminescence, resulting in weight loss and death (Figure 10A, leftmost two mouse panels). Anti-CD 19 and anti-BCMA02 CAR T cells prevented tumor growth, resulting in maintenance of body weight and survival. Anti-CD 19 and anti-BCMA02 CAR T cells were equally effective when administered on Day 8 (Figure 10A, rightmost two mouse panels). Anti-BCMA02 CAR T cells were also effective in decreasing tumor burden when administered at 18 days post-tumor induction. Figure 10B, rightmost panel.

EXAMPLE 4

POTENT *IN VITRO* ACTIVITY OF ANTI-BCMA CAR T CELLS

[0378] Potent *in vitro* activity of anti-BCMA02 CAR T cells was achieved with a 50 percent reduction anti-BCMA02

CAR expression. T cell populations were transduced with between 4×10^8 and 5×10^7 transducing units of a lentivirus encoding an anti-BCMA CAR molecule. The resulting T cell populations showed reduced anti-BCMA CAR T cell frequency (assayed as percent positive) and reduced expression of anti-BCMA CAR molecules (assayed as mean fluorescence intensity: MFI).

[0379] The impact of reduced CAR molecule expression on anti-BCMA activity was determined. The frequency of anti-BCMA CAR-positive T cells was normalized with untransduced T cells to contain $26 \pm 4\%$ BCMA-reactive T cells (Figure 11A). MFI of the normalized anti-BCMA CAR T cells ranged from 885 to 1875 (Figure 11B). K562 is a CML cell line that lacks BCMA expression. K562 cells were engineered to express BCMA and were used in an *in vitro* cytolytic assay to assess activity of anti-BCMA CAR T cells with varied BCMA CAR expression (Figure 11C). K562 cells were labeled with cell trace violet while K562 cells stably expressing BCMA (K562-BCMA) were labeled with CFSE. T cells, K562 cells, and K562-BCMA cells were harvested, washed, and resuspended in media lacking exogenous cytokines. Cells were cultured at a 20:1 or 10:1 effector (E; T cell) to target (T; 1:1 mix of K562 and K562 BCMA cells) ratio for 4 h in a 37 °C, 5% CO₂ incubator. Cells were then stained with Live/Dead and analyzed by FACS. Cytotoxicity was determined by the difference in the ratio of K562:K562-BCMA cells normalized to conditions lacking T cells.

EXAMPLE 5

ANTI-BCMA CAR T CELL MANUFACTURING PROCESS

[0380] Unique anti-BCMA CAR T cell products are manufactured for each patient treatment. The reliability of the manufacturing process for anti-BCMA CAR T cell products was evaluated by generating anti-BCMA CAR T cells from 11 individual normal donor PBMC. Anti-BCMA CAR T cell expansion from each donor was comparable to a matched untransduced culture performed in parallel (Figure 12A).

[0381] At the end of the culture period (day 10), T cell transduction efficiency was assessed by quantitating the number of integrated lentiviruses with qPCR and lentiviral-specific primer sets (vector copy number, VCN). Anti-BCMA CAR T cell cultures from the 11 donors showed comparable lentiviral transduction efficiency (Figure 12B). The frequency of anti-BCMA CAR positive T cells was measured by flow cytometry and BCMA expression was found to be comparable across all donors (Figure 12C).

[0382] The activity of each anti-BCMA CAR T cell product was assessed by IFN γ -release after co-culture with K562 cells engineered to express BCMA. All anti-BCMA CAR T cell products exhibited therapeutically relevant levels of IFN γ release when exposed to BCMA-expressing K562 cells (Figure 12D).

EXAMPLE 6

CD62L, CD127, CD197, AND CD38 EXPRESSION ON CAR T CELLS TREATED WITH IL-2 OR IL-2 AND ZSTK474

[0383] CAR T cells cultured with IL-2 and ZSTK474 show increased CD62L expression compared to CAR T cells cultured with IL-2 alone. Expression analysis of 29 additional cell surface markers on anti-BCMA CAR T cells cultured with IL-2 and ZSTK474 was performed using multiparameter mass cytometry (CyTOF) and compared with CAR T cells cultured in IL-2 alone. Three additional markers (CD127, CD197, and CD38) showed increased expression in the IL-2 + ZSTK474 treated CAR T cells compared to CAR T cells treated with IL-2 alone. Thus, co-expression of CD62L, CD127, CD197, and CD38 further stratified ZSTK474-cultured CAR

T cells. After culture in media containing IL-2, 7.44% of anti-BCMA CAR T co-expressed CD127, CD197 and CD38 compared to 24.5% of anti-BCMA CAR T cells cultured with IL-2 and ZSTK474. The Venn diagram in Figure 13 illustrates the co-expression of CD127, CD197 and CD38 in CD62L positive anti-BCMA CAR T cells.

EXAMPLE 7

ZSTK474 TREATMENT INCREASES THE FREQUENCY OF CD8 T CELLS

[0384] CD8 expression was quantified in anti-BCMA CAR T cells treated with IL-2 alone or IL-2 and ZSTK474. CD8 expression was determined using a fluorescently-labeled anti-CD8 antibody and flow cytometry. Anti-BCMA CAR T cells from seven normal donors cultured with IL-2 and ZSTK474 had significantly higher CD8 expression compared to anti-BCMA CAR T cells cultured with IL-2 alone. Figure 14.

EXAMPLE 8

LACK OF ANTIGEN-INDEPENDENT ACTIVITY IN ZSTK474 TREATED ANTI-BCMA CAR T CELLS

[0385] Tonic activity of CAR T cells in the absence of antigen has been associated with reduced biological activity. Tonic activity of anti-BCMA02 CAR T cells was assessed by quantifying interferon- γ (IFN- γ) release in the absence of antigen after culture in the presence of IL-2 and ZSTK474 compared to standard culture conditions with IL-2 alone. Anti-BCMA CAR T cells cultures were prepared using a system directly scalable to large clinical manufacturing processes. Briefly, peripheral blood mononuclear cells (PBMC) were cultured in static flasks in media containing IL-2 (CellGenix) and antibodies specific for CD3 and CD28 (Miltenyi Biotec). 2×10^8 transducing units of lentivirus encoding anti-BCMA CARs were added one day after culture initiation.

[0386] Anti-BCMA02 CAR T cells were maintained in log-phase by adding fresh media containing IL-2 and an optimized dose of ZSTK474 for a total of ten days of culture. At the end of manufacture, an equivalent number of anti-BCMA02 CAR T cells were re-cultured for 24 hours in media alone. The amount of IFN- γ released in 24 hours was quantified by ELISA. In this assay IFN- γ levels below 200pg/mL represent no tonic activity. Figure 15 shows the amount of IFN- γ released by anti-BCMA02 CAR T cells from 14 donors is consistent with lacking tonic activity whether or not the CAR T cells are cultured with ZSTK474.

EXAMPLE 9

ZSTK474 TREATED ANTI-BCMA02 CAR T CELLS SHOW THERAPEUTIC ACTIVITY IN A LYMPHOMA TUMOR MODEL

[0387] Daudi tumors were used to interrogate the anti-tumor activity of anti-BCMA02 CAR T cells cultured with IL-2 or IL-2 and ZSTK474. Daudi cells express a low level of BCMA protein and provide an aggressive and difficult to treat lymphoma tumor model.

[0388] 2×10^6 Daudi tumor cells were labeled with a firefly luciferase gene and injected into NOD scid IL-2 receptor gamma chain knockout mice (NSG) by intravenous injection. After tumors were allowed to form, 1×10^7 CAR T cells were injected in to tumor bearing mice. Mice were injected with i) anti-BCMA02 CAR T cells treated for ten days with IL-2 or IL-2 and ZSTK474; or ii) a truncated signaling deficient anti-BCMA02 (tBCMA02) CAR T cell treated for ten days with IL-2 and ZSTK474. Tumor growth was monitored by bioluminescence using a Xenogen-IVIS Imaging system.

[0389] Complete tumor regression was observed in 50% of mice administered the anti-BCMA02 CAR T cells treated with IL-2 and ZSTK474. Figure 16.

EXAMPLE 10

ZSTK474 TREATED CAR T CELLS SHOW THERAPEUTIC ACTIVITY IN A MOUSE MODEL OF HUMAN MYELOMA

[0390] Animals with 100mm^3 sub cutaneous multiple myeloma tumors (RPMI-8226) were infused with equivalent CAR T cell doses (1×10^6 anti-BCMA02 CAR-positive T cells) or unmodified T cells from a matched T cell donor (untransduced). Anti-BCMA CAR T cells were treated with IL-2 or IL-2 and ZSTK474 as described in Example 8.

[0391] Animals treated with IL-2- or IL-2 and ZSTK474-cultured anti-BCMA02 CAR T cells completely prevented tumor outgrowth. Figure 17. In contrast, animals treated with untransduced or vehicle were unable to control tumor growth. Figure 17.

SEQUENCE LISTING

[0392]

<110> bluebird bio, Inc.
Morgan, Richard
Friedman, Kevin

<120> BCMA CHIMERIC ANTIGEN RECEPTORS

<130> BLBD-043/02WO

<150> US 62/091,419

<151> 2014-12-12

<150> US 62/200,505

<151> 2015-08-03

5

<160> 36

<170> PatentIn version 3.5

10

<210> 1

<211> 15

<212> PRT

<213> Mus musculus

15

<400> 1

Arg	Ala	Ser	Glu	Ser	Val	Thr	Ile	Leu	Gly	Ser	His	Leu	Ile	His
1				5					10					15

20

<210> 2

<211> 7

<212> PRT

<213> mus musculus

25

<400> 2

Leu	Ala	Ser	Asn	Val	Gln	Thr
1				5		

30

<210> 3

<211> 9

<212> PRT

<213> mus musculus

35

<400> 3

Leu	Gln	Ser	Arg	Thr	Ile	Pro	Arg	Thr
1				5				

40

<210> 4

<211> 5

<212> PRT

<213> Mus musculus

45

<400> 4

Asp	Tyr	Ser	Ile	Asn
1				5

50

<210> 5

<211> 17

<212> PRT

<213> Mus musculus

55

<400> 5

EP 3 230 321 B1

Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe Arg
1 5 10 15

5 Gly

<210> 6

<211> 8

<212> PRT

10 <213> Mus musculus

<400> 6

15 Asp Tyr Ser Tyr Ala Met Asp Tyr
1 5

<210> 7

<211> 111

<212> PRT

20 <213> Mus musculus

<400> 7

25 Asp Ile Val Leu Thr Gln Ser Pro Pro Ser Leu Ala Met Ser Leu Gly
1 5 10 15

Lys Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Thr Ile Leu
20 25 30

Gly Ser His Leu Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45

35 Thr Leu Leu Ile Gln Leu Ala Ser Asn Val Gln Thr Gly Val Pro Ala
50 55 60

40 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp
65 70 75 80

Pro Val Glu Glu Asp Asp Val Ala Val Tyr Tyr Cys Leu Gln Ser Arg
85 90 95

45 Thr Ile Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

50 <210> 8

<211> 117

<212> PRT

<213> Mus musculus

55 <400> 8

EP 3 230 321 B1

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15
 5 Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 10 Ser Ile Asn Trp Val Lys Arg Ala Pro Gly Lys Gly Leu Lys Trp Met
 35 40 45
 Gly Trp Ile Asn Thr Glu Thr Arg Glu Pro Ala Tyr Ala Tyr Asp Phe
 50 55 60
 15 Arg Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 20 Leu Gln Ile Asn Asn Leu Lys Tyr Glu Asp Thr Ala Thr Tyr Phe Cys
 85 90 95
 25 Ala Leu Asp Tyr Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser
 100 105 110
 Val Thr Val Ser Ser
 115
 30 <210> 9
 <211> 493
 <212> PRT
 <213> Artificial Sequence
 35 <220>
 <223> anti-BCMA02 CAR
 40 <400> 9
 Met Ala Leu Pro Val Thr Ala Leu Leu Leu Pro Leu Ala Leu Leu Leu
 1 5 10 15
 45 His Ala Ala Arg Pro Asp Ile Val Leu Thr Gln Ser Pro Pro Ser Leu
 20 25 30
 50 Ala Met Ser Leu Gly Lys Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu
 35 40 45
 Ser Val Thr Ile Leu Gly Ser His Leu Ile His Trp Tyr Gln Gln Lys
 50 55 60
 55

EP 3 230 321 B1

	Pro	Gly	Gln	Pro	Pro	Thr	Leu	Leu	Ile	Gln	Leu	Ala	Ser	Asn	Val	Gln	
	65					70					75					80	
5	Thr	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Arg	Thr	Asp	Phe	
					85					90					95		
10	Thr	Leu	Thr	Ile	Asp	Pro	Val	Glu	Glu	Asp	Asp	Val	Ala	Val	Tyr	Tyr	
				100					105					110			
15	Cys	Leu	Gln	Ser	Arg	Thr	Ile	Pro	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	
			115					120					125				
20	Leu	Glu	Ile	Lys	Gly	Ser	Thr	Ser	Gly	Ser	Gly	Lys	Pro	Gly	Ser	Gly	
		130					135					140					
25	Glu	Gly	Ser	Thr	Lys	Gly	Gln	Ile	Gln	Leu	Val	Gln	Ser	Gly	Pro	Glu	
	145					150					155					160	
30	Leu	Lys	Lys	Pro	Gly	Glu	Thr	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	
					165					170					175		
35	Tyr	Thr	Phe	Thr	Asp	Tyr	Ser	Ile	Asn	Trp	Val	Lys	Arg	Ala	Pro	Gly	
				180					185					190			
40	Lys	Gly	Leu	Lys	Trp	Met	Gly	Trp	Ile	Asn	Thr	Glu	Thr	Arg	Glu	Pro	
			195					200					205				
45	Ala	Tyr	Ala	Tyr	Asp	Phe	Arg	Gly	Arg	Phe	Ala	Phe	Ser	Leu	Glu	Thr	
		210					215					220					
50	Ser	Ala	Ser	Thr	Ala	Tyr	Leu	Gln	Ile	Asn	Asn	Leu	Lys	Tyr	Glu	Asp	
	225					230					235					240	
55	Thr	Ala	Thr	Tyr	Phe	Cys	Ala	Leu	Asp	Tyr	Ser	Tyr	Ala	Met	Asp	Tyr	
					245					250					255		
60	Trp	Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Ala	Ala	Ala	Thr	Thr	
				260					265					270			
65	Thr	Pro	Ala	Pro	Arg	Pro	Pro	Thr	Pro	Ala	Pro	Thr	Ile	Ala	Ser	Gln	
			275					280					285				
70	Pro	Leu	Ser	Leu	Arg	Pro	Glu	Ala	Cys	Arg	Pro	Ala	Ala	Gly	Gly	Ala	
		290					295					300					
75	Val	His	Thr	Arg	Gly	Leu	Asp	Phe	Ala	Cys	Asp	Ile	Tyr	Ile	Trp	Ala	
	305					310					315					320	

EP 3 230 321 B1

Pro Leu Ala Gly Thr Cys Gly Val Leu Leu Leu Ser Leu Val Ile Thr
325 330 335

5 Leu Tyr Cys Lys Arg Gly Arg Lys Lys Leu Leu Tyr Ile Phe Lys Gln
340 345 350

10 Pro Phe Met Arg Pro Val Gln Thr Thr Gln Glu Glu Asp Gly Cys Ser
355 360 365

Cys Arg Phe Pro Glu Glu Glu Glu Gly Gly Cys Glu Leu Arg Val Lys
370 375 380

15 Phe Ser Arg Ser Ala Asp Ala Pro Ala Tyr Gln Gln Gly Gln Asn Gln
385 390 395 400

20 Leu Tyr Asn Glu Leu Asn Leu Gly Arg Arg Glu Glu Tyr Asp Val Leu
405 410 415

25 Asp Lys Arg Arg Gly Arg Asp Pro Glu Met Gly Gly Lys Pro Arg Arg
420 425 430

Lys Asn Pro Gln Glu Gly Leu Tyr Asn Glu Leu Gln Lys Asp Lys Met
435 440 445

30 Ala Glu Ala Tyr Ser Glu Ile Gly Met Lys Gly Glu Arg Arg Arg Gly
450 455 460

35 Lys Gly His Asp Gly Leu Tyr Gln Gly Leu Ser Thr Ala Thr Lys Asp
465 470 475 480

40 Thr Tyr Asp Ala Leu His Met Gln Ala Leu Pro Pro Arg
485 490

<210> 10

<211> 1485

<212> DNA

45 <213> Artificial Sequence

<220>

<223> anti-BCMA02 CAR

50 <400> 10

atggcactcc ccgtcaccgc ccttctcttg cccctcgccc tgctgctgca tgctgccagg	60
cccgacattg tgctcactca gtcacctccc agcctggcca tgagcctggg aaaaagggcc	120
accatctcct gtagagccag tgagtccgtc acaatcttgg ggagccatct tattcactgg	180
tatcagcaga agcccgggca gcctccaacc cttcttattc agctcgcgtc aaacgtccag	240

EP 3 230 321 B1

	acgggtgtac ctgccagatt ttctggtagc ggggtcccga ctgattttac actgaccata	300
	gatccagtgg aagaagacga tgtggccgtg tattattgtc tgcagagcag aacgattcct	360
5	cgcacatttg gtgggggtac taagctggag attaaggga gcacgtccgg ctcaggaag	420
	ccgggctccg gcgagggaag cacgaagggg caaattcagc tgggtccagag cggacctgag	480
10	ctgaaaaaac ccggcgagac tgttaagatc agttgtaaag catctggcta taccttcacc	540
	gactacagca taaattgggt gaaacgggccc cctggaaagg gcctcaaag gatgggttg	600
	atcaataaccg aaactagggga gcctgcttat gcatatgact tccgcgggag attcgccttt	660
15	tcactcgaga catctgcctc tactgcttac ctccaaataa acaacctcaa gtatgaagat	720
	acagccactt acttttgcgc cctcgactat agttacgcca tggactactg gggacagga	780
	acctccgtta ccgtcagttc cgcggccgca accacaacac ctgctccaag gccccccaca	840
20	cccgtccaa ctatagccag ccaaccattg agcctcagac ctgaagcttg caggcccga	900
	gcaggaggcg ccgtccatac gcgaggcctg gacttcgcgt gtgatattta tatttgggcc	960
	cctttggccg gaacatgtgg ggtgttgctt ctctcccttg tgatcactct gtattgtaag	1020
25	cgcgggagaa agaagctcct gtacatcttc aagcagcctt ttatgcgacc tgtgcaaacc	1080
	actcaggaag aagatgggtg ttcattgccgc ttccccgagg aggaagaagg aggggtgtgaa	1140
30	ctgaggggtga aattttctag aagcgccgat gctcccgcat atcagcaggg tcagaatcag	1200
	ctctacaatg aattgaatct cggcaggcga gaagagtacg atgttctgga caagagacgg	1260
	ggcagggatc ccgagatggg gggaaagccc cggagaaaaa atcctcagga ggggtgtac	1320
35	aatgagctgc agaaggacaa gatggctgaa gcctatagcg agatcggaat gaaaggcgaa	1380
	agacgcagag gcaaggggca tgacggtctg taccagggtc tctctacagc caccaaggac	1440
40	acttatgatg cgttgcatat gcaagccttg ccaccccgt aatga	1485
	<210> 11	
	<211> 184	
	<212> PRT	
	<213> Homo sapiens	
45	<400> 11	
50		
55		

EP 3 230 321 B1

Asp Gly Gly Gly Ser
1 5

EP 3 230 321 B1

<210> 13
 <211> 5
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Flexible peptide linker
 <400> 13
 10
 Thr Gly Glu Lys Pro
 1 5
 <210> 14
 <211> 4
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Flexible peptide linker
 20
 <400> 14
 25
 Gly Gly Arg Arg
 1
 <210> 15
 <211> 5
 <212> PRT
 <213> Artificial Sequence
 30
 <220>
 <223> Flexible peptide linker
 35
 <400> 15
 40
 Gly Gly Gly Gly Ser
 1 5
 <210> 16
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 45
 <220>
 <223> Flexible peptide linker
 <400> 16
 50
 Glu Gly Lys Ser Ser Gly Ser Gly Ser Glu Ser Lys Val Asp
 1 5 10
 <210> 17
 <211> 18
 <212> PRT
 <213> Artificial Sequence
 55
 <220>

EP 3 230 321 B1

<223> Flexible peptide linker

<400> 17

⁵

Lys	Glu	Ser	Gly	Ser	Val	Ser	Ser	Glu	Gln	Leu	Ala	Gln	Phe	Arg	Ser
1				5					10					15	

Leu Asp

10

<210> 18

 $\langle 211 \rangle$ 8

<212> PRT

<213> Artificial Sequence

15

<220>

<223> Flexible peptide linker

<400> 18

20

Gly Gly Arg Arg Gly Gly Gly Ser
1 5

<210> 19

25

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

30

<223> Flexible peptide linker

<400> 19

Leu Arg Gln Arg Asp Gly Glu Arg Pro
1 5

35

<210> 20

<211> 12

<212> PRT

40

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Flexible peptide linker

45

<400> 20

Leu Arg Gln Lys Asp Gly Gly Gly Ser Glu Arg Pro
1 5 10

50

<210> 21

<211> 16

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Flexible peptide linker

<400> 21

EP 3 230 321 B1

	Leu Arg Gln Lys Asp Gly Gly Gly Ser Gly Gly Gly Ser Glu Arg Pro
	1 5 10 15

5 <210> 22
 <211> 18
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Flexible peptide linker

<400> 22

	Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser Thr
15	1 5 10 15

Lys Gly

20 <210> 23
 <211> 7
 <212> PRT
 <213> Artificial Sequence

25 <220>
 <223> protease cleavage site

<220>
 <221> MISC_FEATURE
 30 <222> (2)..(3)
 <223> Xaa = Any amino acid

<220>
 <221> MISC_FEATURE
 35 <222> (5)..(5)
 <223> Xaa = Any amino acid

<220>
 <221> MISC_FEATURE
 40 <222> (7)..(7)
 <223> Xaa is Gly or Ser

<400> 23

45	Glu Xaa Xaa Tyr Xaa Gln Xaa
	1 5

<210> 24
 <211> 7
 50 <212> PRT
 <213> Artificial Sequence

<220>
 <223> protease cleavage site
 55

<400> 24

EP 3 230 321 B1

Glu Asn Leu Tyr Phe Gln Gly
1 5

5 <210> 25
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
10 <223> protease cleavage site

<400> 25

15 Glu Asn Leu Tyr Phe Gln Ser

1 5

20 <210> 26
<211> 19
<212> PRT
<213> Artificial Sequence

25 <220>
<223> protease cleavage site

<400> 26

30 Leu Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn
1 5 10 15

Pro Gly Pro
35 <210> 27
<211> 19
<212> PRT
<213> Artificial Sequence

40 <220>
<223> protease cleavage site

<400> 27

45 Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn
1 5 10 15

50 Pro Gly Pro

<210> 28
<211> 14
<212> PRT
55 <213> Artificial Sequence

<220>
<223> protease cleavage site

EP 3 230 321 B1

<400> 28

Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro Gly Pro
1 5 10

<210> 29

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 29

Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro Gly
1 5 10 15

Pro

<210> 30

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 30

Gln Leu Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser
1 5 10 15

Asn Pro Gly Pro
20

<210> 31

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<223> protease cleavage site

<400> 31

Ala Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly
1 5 10 15

Asp Val Glu Ser Asn Pro Gly Pro
20

<210> 32

<211> 40

<212> PRT

EP 3 230 321 B1

<213> Artificial Sequence

<220>

<223> protease cleavage site

5

<400> 32

10

Val Thr Glu Leu Leu Tyr Arg Met Lys Arg Ala Glu Thr Tyr Cys Pro
1 5 10 15

Arg Pro Leu Leu Ala Ile His Pro Thr Glu Ala Arg His Lys Gln Lys
20 25 30

15

Ile Val Ala Pro Val Lys Gln Thr

35

40

20

<210> 33

<211> 18

<212> PRT

<213> Artificial Sequence

25

<220>

<223> protease cleavage site

<400> 33

30

Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro
1 5 10 15

35

Gly Pro

<210> 34

<211> 40

<212> PRT

<213> Artificial Sequence

40

<220>

<223> protease cleavage site

45

<400> 34

Leu Leu Ala Ile His Pro Thr Glu Ala Arg His Lys Gln Lys Ile Val
1 5 10 15

50

Ala Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly
20 25 30

55

Asp Val Glu Ser Asn Pro Gly Pro
35 40

<210> 35

EP 3 230 321 B1

<211> 33
 <212> PRT
 <213> Artificial Sequence

5 <220>
 <223> protease cleavage site

<400> 35

10 Glu Ala Arg His Lys Gln Lys Ile Val Ala Pro Val Lys Gln Thr Leu
 1 5 10 15

15 Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp Val Glu Ser Asn Pro Gly
 20 25 30

Pro

20 <210> 36
 <211> 7350
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> anti-BCMA02 CAR vector

<400> 36

30

35

40

45

50

55

EP 3 230 321 B1

	tcgcgcggttt	cggatgatgac	ggtgaaaacc	tctgacacat	gcagctcccg	gagacggtca	60
	cagcttgtct	gtaagcggat	gccgggagca	gacaagcccg	tcagggcgcg	tcagcgggtg	120
5	ttggcgggtg	tcggggctgg	cttaactatg	cggcatcaga	gcagattgta	ctgagagtgc	180
	accatcatat	gccagcctat	ggtgacattg	attattgact	agttattaat	agtaatcaat	240
	tacgggggtca	ttagttcata	gcccataat	ggagttccgc	gttacataac	ttacggtaaa	300
10	tggcccgct	ggctgaccgc	ccaacgaccc	ccgcccattg	acgtcaataa	tgacgtatgt	360
	tcccatagta	acgccaatag	ggactttcca	ttgacgtcaa	tgggtggagt	atttacggta	420
	aactgcccac	ttggcagtac	atcaagtgtg	tcatatgccg	agtacgcccc	ctattgacgt	480
15	caatgacggt	aaatggcccg	cctggcatta	tgcccagtac	atgaccttat	gggactttcc	540
	tacttggcag	tacatctacg	tattagtcac	cgctattacc	atggtgatgc	ggttttggca	600
20	gtacatcaat	gggcgtggat	agcggtttga	ctcacgggga	tttccaagtc	tccaccccat	660
	tgacgtcaat	gggagtttgt	tttggcacca	aatcaacgg	gactttccaa	aatgtcgtaa	720
	caactccgcc	ccattgacgc	aaatgggagg	taggcgtgta	cgggtggagg	tctatataag	780
25	cagagctcgt	ttagtgaacc	gggtctctct	ggtagacca	gatctgagcc	tgggagctct	840
	ctggctaact	agggaaccca	ctgcttaagc	ctcaataaag	cttgccttga	gtgctcaaag	900
	tagtgtgtgc	ccgtctgttg	tgtgactctg	gtaactagag	atccctcaga	cccttttagt	960
30	cagtgtggaa	aatctctagc	agtggcgccc	gaacagggac	ttgaaagcga	aagtaaagcc	1020
	agaggagatc	tctcgacgca	ggactcggct	tgctgaagcg	cgcacggcaa	gaggcgaggg	1080
35	gcggcgactg	gtgagtacgc	caaaaatttt	gactagcgga	ggctagaagg	agagagtagg	1140
	gtgcgagagc	gtcgggtatta	agcgggggag	aattagataa	atgggaaaaa	attcggttaa	1200
	ggccaggggg	aaagaaacaa	tataaactaa	aacatatagt	tagggcaagc	aggagctag	1260
40	aacgattcgc	agttaatcct	ggccttttag	agacatcaga	aggctgtaga	caaatactgg	1320
	gacagctaca	accatccctt	cagacaggat	cagaagaact	tagatcatta	tataatacaa	1380
	tagcagtcct	ctatttgtgt	catcaaagga	tagatgtaaa	agacaccaag	gaagccttag	1440
45	ataagataga	ggaagagcaa	aacaaaagta	agaaaaaggc	acagcaagca	gcagctgaca	1500
	caggaaacaa	cagccaggtc	agccaaaatt	accctatagt	gcagaacctc	caggggcaaa	1560

50

55

EP 3 230 321 B1

	tggtacatca	ggccatatca	cctagaactt	taaattaaga	cagcagtaca	aatggcagta	1620
	ttcatccaca	atthttaaag	aaaagggggg	attggggggg	acagtgcagg	ggaaagaata	1680
5	gtagacataa	tagcaacaga	catacaaact	aaagaattac	aaaaacaaat	tacaaaaatt	1740
	caaaatthtt	gggtthtatta	cagggacagc	agagatccag	tttggaagg	accagcaaag	1800
	ctcctctgga	aaggtgaagg	ggcagtagta	atacaagata	atagtgcacat	aaaagtagtg	1860
10	ccaagaagaa	aagcaaagat	catcagggat	tatggaaaac	agatggcagg	tgatgattgt	1920
	gtggcaagta	gacaggatga	ggattaacac	atggaaaaga	ttagtaaaac	accatagctc	1980
	tagagcgatc	ccgatcttca	gacctggagg	aggagatatg	agggacaatt	ggagaagtga	2040
15	attatataaa	tataaagtag	taaaaattga	accattagga	gtagcaccca	ccaaggcaaa	2100
	gagaagagt	gtgcagagag	aaaaaagagc	agtgggaata	ggagctttgt	tccttgggtt	2160
	cttgggagca	gcagggaagca	ctatggggcg	agcgtcaatg	acgctgcagg	tacaggccag	2220
20	acaattattg	tctggtatag	tgcagcagca	gaacaatttg	ctgagggcta	ttgagggcga	2280
	acagcatctg	ttgcaactca	cagtctgggg	catcaagcag	ctccaggcaa	gaatcctggc	2340
25	tgtggaaaga	tacctaaagg	atcaacagct	cctggggatt	tggggttgct	ctggaaaact	2400
	catttgcacc	actgctgtgc	cttggaatgc	tagttggagt	aataaatctc	tggaacagat	2460
	ttggaatcac	acgacctgga	tggagtggga	cagagaaatt	aacaattaca	caagcttggt	2520
30	aggthtaaga	atagtthttg	ctgtacttht	tatagtgaat	agagttaggc	agggatattc	2580
	accattatcg	tttcagaccc	acctcccaac	cccaggggga	cccagcaggc	ccgaaggaat	2640
	agaagaagaa	ggtggagaga	gagacagaga	cagatccatt	cgattagtga	acggatccat	2700
35	ctcgacggaa	tgaagagccc	cacctgtagg	tttggcaagc	taggatcaag	gttaggaaca	2760
	gagagacagc	agaatatggg	ccaaacagga	tatctgtggt	aagcagttcc	tgccccggct	2820
	cagggccaag	aacagttgga	acagcagaat	atgggccaaa	caggatatct	gtggtaagca	2880
40	gttcctgccc	cggctcaggg	ccaagaacag	atggtcccca	gatgcggtcc	cgccctcagc	2940
	agthttctaga	gaaccatcag	atgthttccag	ggtgccccaa	ggacctgaaa	tgacctgtg	3000
	ccttatttga	actaaccaat	cagttcgctt	ctcgcttctg	ttcgcgcgct	tctgctcccc	3060
45	gagctcaata	aaagagccca	caaccctca	ctcggcgcga	ttcacctgac	gcgtctacgc	3120
	caccatggca	ctccccgtca	ccgcccttct	cttgccctct	gccctgctgc	tgcatgctgc	3180
	cagggccgac	attgtgctca	ctcagtcacc	tcccagcctg	gccatgagcc	tgggaaaaag	3240
50	ggccaccatc	tcctgtagag	ccagtgagtc	cgtcacaaat	ttggggagcc	atcttattca	3300
	ctggtatcag	cagaagcccg	ggcagcctcc	aacccttctt	attcagctcg	cgtcaaacgt	3360
55	ccagacgggt	gtacctgcca	gattthttctg	tagcgggtcc	cgcaactgatt	ttacactgac	3420
	catagatcca	gtggaagaag	acgatgtggc	cgtgtattat	tgtctgcaga	gcagaacgat	3480

EP 3 230 321 B1

	tcctcgaca	tttgggtggg	gtactaagct	ggagattaag	ggaagcacgt	ccggctcagg	3540
	gaagccgggc	tccggcgagg	gaagcacgaa	ggggcaaatt	cagctggtcc	agagcggacc	3600
5	tgagctgaaa	aaacccggcg	agactgttaa	gatcagttgt	aaagcatctg	gctatacctt	3660
	caccgactac	agcataaatt	gggtgaaacg	ggcccctgga	aagggcctca	aatggatggg	3720
	ttggatcaat	accgaaacta	gggagcctgc	ttatgcatat	gacttccgcg	ggagattcgc	3780
10	cttttctactc	gagacatctg	cctctactgc	ttacctcaa	ataaacaacc	tcaagtatga	3840
	agatacagcc	acttactttt	gcgccctcga	ctatagttac	gccatggact	actggggaca	3900
	gggaacctcc	gttaccgtca	gttccgcggc	cgcaaccaca	acacctgctc	caaggccccc	3960
15	cacacccgct	ccaactatag	ccagccaacc	attgagcctc	agacctgaag	cttgacaggcc	4020
	cgcagcagga	ggcgccgtcc	atacgcgagg	cctggacttc	gcgtgtgata	tttatatttg	4080
20	ggccccctttg	gccggaacat	gtgggggtgtt	gcttctctcc	cttgtgatca	ctctgtattg	4140
	taagcgcggg	agaaagaagc	tcctgtacat	cttcaagcag	ccttttatgc	gacctgtgca	4200
	aaccactcag	gaagaagatg	ggtgttcacg	ccgcttcccc	gaggaggaag	aaggaggggtg	4260
25	tgaactgagg	gtgaaatfff	ctagaagcgc	cgatgctccc	gcataatcagc	agggtcagaa	4320
	tcagctctac	aatgaattga	atctcggcag	gcgagaagag	tacgatgttc	tggacaagag	4380
	acggggcgagg	gatccccgaga	tggggggaaa	gccccggaga	aaaaatcctc	aggagggggtt	4440
30	gtacaatgag	ctgcagaagg	acaagatggc	tgaagcctat	agcgagatcg	gaatgaaagg	4500
	cgaaagacgc	agaggcaagg	ggcatgacgg	tctgtaccag	ggtctctcta	cagccaccaa	4560
35	ggacacttat	gatgcgttgc	atatgcaagc	cttgccaccc	cgctaattgac	aggtaccttt	4620
	aagaccaatg	acttacaagg	cagctgtaga	tcttagccac	tttttaaaaag	aaaagggggg	4680
	actggaagg	ctaattcact	cccaaagaag	acaagatctg	ctttttgcct	gtactgggtc	4740
40	tctctgggtta	gaccagatct	gagcctggga	gctctctggc	taactaggga	acccactgct	4800
	taagcctcaa	taaagcttgc	cttgagtgtc	tcaatgtgtg	tgttggtttt	ttgtgtgtcg	4860
	aaattctagc	gattctagct	tggcgtaatc	atggtcacag	ctgtttcctg	tgtgaaattg	4920
45	ttatccgctc	acaattccac	acaacatacg	agccggaagc	ataaagtgtg	aagcctgggg	4980
	tgcctaataga	gtgagctaac	tcacattaat	tgcgttgccg	tactgcccg	ctttccagtc	5040
50	gggaaacctg	tcgtgccagc	tgcattaatg	aatcgcccaa	cgcgcgggga	gaggcggttt	5100
	gcgtattggg	cgctcttccg	cttcctcgct	cactgactcg	ctgcgctcgg	tcgttcggct	5160
	gcggcgagcg	gtatcagctc	actcaaaggc	ggtaatacgg	ttatccacag	aatcagggga	5220
55	taacgcagga	aagaacatgt	gagcaaaagg	ccagcaaaag	gccaggaacc	gtaaaaaggc	5280
	cgcgttgctg	gcgtttttcc	ataggctccg	ccccctgac	gagcatcaca	aaaatcgacg	5340

EP 3 230 321 B1

	ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt ttccccctgg	5400
	aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc tgtccgcctt	5460
5	tctcccttcg ggaagcgtgg cgctttctca tagctcacgc tgtaggatc tcagttcgg	5520
	gtaggtcggt cgctccaagc tgggctgtgt gcacgaaccc cccgttcagc ccgaccgctg	5580
	cgccttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact	5640
10	ggcagcagcc actggtaaca ggattagcag agcgaggat gtaggcgggtg ctacagagtt	5700
	cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttggt tctgcgctct	5760
	gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac	5820
15	cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc	5880
	tcaagaagat cctttgatct tttctacggg gtctgacgct cagtggaacg aaaactcacg	5940
	ttaagggatt ttggtcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta	6000
20	aaaatgaagt tttaaatcaa tctaaagtat atatgagtaa acttggtctg acagttacca	6060
	atgcttaatc agtgaggcac ctatctcagc gatctgtcta tttcgttcat ccatagttgc	6120
25	ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg gccccagtgc	6180
	tgcaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc	6240
	agccggaagg gccgagcgca gaagtgggtc tgcaacttta tccgcctcca tccagtctat	6300
30	taattgttgc cgggaagcta gagtaagtag ttcgccagtt aatagtttgc gcaacgttgt	6360
	tgccattgct acaggcatcg tgggtgtcacg ctcgctggtt ggtatggctt cattcagctc	6420
	cggttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag	6480
35	ctccttcggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat cactcatggt	6540
	tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac	6600
	tgggtgagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg	6660
40	cccggcgtca atacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat	6720
	tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc	6780
	gatgtaaccc actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc	6840
45	tgggtgagca aaaacaggaa ggcaaatgc cgcaaaaaag ggaataaggg cgacacggaa	6900
	atgttgaata ctcatactct tcctttttca atattattga agcatttatc agggttattg	6960
	tctcatgagc ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttccgcg	7020
50	cacatttccc cgaaaagtgc cacctgggac tagctttttg caaaagccta ggcctccaaa	7080
	aaagcctcct cactacttct ggaatagctc agaggccgag gcggcctcgg cctctgcata	7140
55	aataaaaaaa attagtcagc catggggcgg agaatgggcg gaactgggcg gagttagggg	7200
	cgggatgggc ggagttaggg gcgggactat ggttgctgac taattgagat gagcttgcat	7260

gccgacattg attattgact agtcctaag aaaccattct tatcatgaca ttaacctata 7320
 aaaataggcg tatcacgagg ccttttcgtc 7350

5

Claims

- 10 1. A polynucleotide encoding a CAR, wherein the polynucleotide sequence is set forth in SEQ ID NO: 10.
2. A vector comprising the polynucleotide of claim 1.
- 15 3. The vector of claim 2, wherein the vector is an expression vector, an episomal vector, a viral vector, a retroviral vector, or a lentiviral vector.
- 20 4. The vector of claim 2, wherein the lentiviral vector is selected from the group consisting essentially of: human immunodeficiency virus 1 (HIV-1); human immunodeficiency virus 2 (HIV-2), visna-maedi virus (VMV) virus; caprine arthritis-encephalitis virus (CAEV); equine infectious anemia virus (EIAV); feline immunodeficiency virus (FIV); bovine immune deficiency virus (BIV); and simian immunodeficiency virus (SIV).
- 25 5. The vector according to any one of claims 2 to 4, comprising a left (5') retroviral LTR, a Psi (Ψ) packaging signal, a central polypurine tract/DNA flap (cPPT/FLAP), a retroviral export element; a promoter operably linked to the polynucleotide; and a right (3') retroviral LTR.
6. The vector of claim 5, further comprising a heterologous polyadenylation sequence.
7. The vector of claim 5 or claim 6, wherein the promoter of the 5' LTR is replaced with a heterologous promoter.
- 30 8. The vector of claim 7, wherein the heterologous promoter is a cytomegalovirus (CMV) promoter, a Rous Sarcoma Virus (RSV) promoter, or an Simian Virus 40 (SV40) promoter.
9. The vector of any one of claims 5 to 8, wherein the 5' LTR or 3' LTR is a lentivirus LTR.
- 35 10. The vector of any one of claims 5 to 8, wherein the 3' LTR is a self-inactivating (SIN) LTR.
- 40 11. The vector of any one of claims 5 to 10, wherein the polyadenylation sequence is a bovine growth hormone polyadenylation or signal rabbit β -globin polyadenylation sequence.
- 45 12. The vector of any one of claims 5 to 11, wherein the promoter operably linked to the polynucleotide of claim 1 is selected from the group consisting of: a cytomegalovirus immediate early gene promoter (CMV), an elongation factor 1 alpha promoter (EF1- α), a phosphoglycerate kinase-1 promoter (PGK), a ubiquitin-C promoter (UBQ-C), a cytomegalovirus enhancer/chicken beta-actin promoter (CAG), polyoma enhancer/herpes simplex thymidine kinase promoter (MC1), a beta actin promoter (β -ACT), a simian virus 40 promoter (SV40), and a myeloproliferative sarcoma virus enhancer, negative control region deleted, dl587rev primer-binding site substituted (MND) promoter.
13. An *ex vivo* or *in vitro* immune effector cell comprising the vector of any one of claims 2 to 12.
- 50 14. The *ex vivo* or *in vitro* immune effector cell of claim 13, wherein the immune effector cell is selected from the group consisting of: a T lymphocyte and a natural killer (NK) cell.
15. A composition comprising the *ex vivo* or *in vitro* immune effector cell of claim 13 or claim 14 and a physiologically acceptable excipient.

55

Patentansprüche

1. Ein Polynucleotid, das für einen CAR kodiert, wobei die Polynucleotidsequenz in SEQ ID NO: 10 dargelegt ist.

2. Ein Vektor, der das Polynucleotid gemäß Anspruch 1 beinhaltet.
3. Vektor gemäß Anspruch 2, wobei der Vektor ein Expressionsvektor, ein episomaler Vektor, ein viraler Vektor, ein retroviraler Vektor oder ein lentiviraler Vektor ist.
4. Vektor gemäß Anspruch 2, wobei der lentivirale Vektor aus der Gruppe ausgewählt ist, die im Wesentlichen aus den Folgenden besteht: dem Human Immunodeficiency Virus 1 (HIV-1); dem Human Immunodeficiency Virus 2 (HIV-2), dem Visna-Maedi-Virus (VMV); dem Caprine-Arthritis-Encephalitis-Virus (CAEV); dem Equine-Infektiöse-Anämie-Virus (EIAV); dem Feline-Immunodeficiency-Virus (FIV); dem Bovine-Immunodeficiency-Virus (BIV); und dem Simian-Immunodeficiency-Virus (SIV).
5. Vektor gemäß einem der Ansprüche 2 bis 4, der Folgendes beinhaltet: eine linke (5') retrovirale LTR, ein Psi (ψ) Packsignal, eine zentrale Polypurin-Trakt/DNA-Flap (cPPT/FLAP), ein retrovirales Exportelement; einen Promoter, der betriebsfähig mit dem Polynucleotid verknüpft ist; und eine rechte (3') retrovirale LTR.
6. Vektor gemäß Anspruch 5, der ferner eine heterologe Polyadenylierungssequenz beinhaltet.
7. Vektor gemäß Anspruch 5 oder Anspruch 6, wobei der Promoter der 5'-LTR durch einen heterologen Promoter ersetzt ist.
8. Vektor gemäß Anspruch 7, wobei der heterologe Promoter ein Cytomegalovirus(CMV)-Promoter, ein Rous-Sarkom-Virus(RSV)-Promoter oder ein Simian-Virus-40(SV40)-Promoter ist.
9. Vektor gemäß einem der Ansprüche 5 bis 8, wobei die 5'-LTR oder 3'-LTR eine Lentivirus-LTR ist.
10. Vektor gemäß einem der Ansprüche 5 bis 8, wobei die 3'-LTR eine selbstinaktivierende (SIN) LTR ist.
11. Vektor gemäß einem der Ansprüche 5 bis 10, wobei die Polyadenylierungssequenz eine Rinderwachstumshormon-Polyadenylierungs- oder Signal-Kaninchen- β -Globin-Polyadenylierungssequenz ist.
12. Vektor gemäß einem der Ansprüche 5 bis 11, wobei der Promoter, der betriebsfähig mit dem Polynucleotid gemäß Anspruch 1 verknüpft ist, aus der Gruppe ausgewählt ist, die aus den Folgenden besteht: einem Cytomegalovirus-Immediate-Early-Gene-Promoter (CMV), einem Elongation-Factor-1-alpha-Promoter (EF1- α), einem Phosphoglyceratkinase-1-Promoter (PGK), einem Ubiquitin-C-Promoter (UBQ-C), einem Cytomegalovirus-Enhancer/Chicken-Beta-Actin-Promoter (CAG), einem Polyom-Enhancer/Herpes-simplex-Thymidinkinase-Promoter (MC1), einem Beta-Actin-Promoter (β -ACT), einem Simian-Virus-40-Promoter (SV40) und einem Myeloproliferativen Sarkom-Virus-Enhancer-Promoter mit negativer Kontrollregion deletiert, dl587rev-Primer-Bindungsstelle substituiert (MND) .
13. Eine *Ex-vivo*- oder *In-vitro*-Immuneffektorzelle, die den Vektor gemäß einem der Ansprüche 2 bis 12 beinhaltet.
14. *Ex-vivo*- oder *In-vitro*-Immuneffektorzelle gemäß Anspruch 13, wobei die Immuneffektorzelle aus der Gruppe ausgewählt ist, die aus Folgenden besteht: einem T-Lymphozyten und einer natürlichen Killer(NK)-Zelle.
15. Eine Zusammensetzung, die die *Ex-vivo*- oder *In-vitro*-Immuneffektorzelle gemäß Anspruch 13 oder Anspruch 14 und einen physiologisch akzeptablen Hilfsstoff beinhaltet.

Revendications

1. Polynucléotide codant un récepteur antigénique chimérique (CAR), dans lequel la séquence du polynucléotide est présentée dans la SEQ ID N° 10.
2. Vecteur comprenant le polynucléotide selon la revendication 1.
3. Vecteur selon la revendication 2, dans lequel le vecteur est un vecteur d'expression, un vecteur épisomal, un vecteur viral, un vecteur rétroviral ou un vecteur lentiviral.
4. Vecteur selon la revendication 2, dans lequel le vecteur lentiviral est choisi dans le groupe essentiellement constitué

par : le virus de l'immunodéficience humaine 1 (VIH-1) ; le virus de l'immunodéficience humaine 2 (VIH-2) ; le virus visna-maedi (VVM) ; le virus de l'arthrite-encéphalite caprine (VAEC) ; le virus de l'anémie infectieuse équine (VAIE) ; le virus de l'immunodéficience féline (VIF) ; le virus de l'immunodéficience bovine (VIB) ; et le virus de l'immunodéficience simienne (VIS).

- 5
10
15
20
25
30
35
40
45
50
55
5. Vecteur selon l'une quelconque des revendications 2 à 4, comprenant une longue séquence répétée terminale (LTR) rétrovirale gauche (5') ; un signal d'encapsidation Psi (Ψ) ; une séquence central polypurine tract/DNA flap (cPPT/FLAP) ; un élément d'exportation rétroviral ; un promoteur lié fonctionnellement au polynucléotide ; et une LTR rétrovirale droite (3').
6. Vecteur selon la revendication 5, comprenant en outre une séquence de polyadénylation hétérologue.
7. Vecteur selon la revendication 5 ou la revendication 6, dans lequel le promoteur de la LTR 5' est remplacé par un promoteur hétérologue.
8. Vecteur selon la revendication 7, dans lequel le promoteur hétérologue est un promoteur de cytomégalo virus (CMV), un promoteur du virus du sarcome de Rous (VSR) ou un promoteur du virus simien 40 (VS40).
9. Vecteur selon l'une quelconque des revendications 5 à 8, dans lequel la LTR 5' ou la LTR 3' est une LTR de lentivirus.
10. Vecteur selon l'une quelconque des revendications 5 à 8, dans lequel la LTR 3' est une LTR auto-inactivante (SIN) .
11. Vecteur selon l'une quelconque des revendications 5 à 10, dans lequel la séquence de polyadénylation est une séquence de polyadénylation de l'hormone de croissance bovine ou une séquence signal de polyadénylation de la β -globine de lapin.
12. Vecteur selon l'une quelconque des revendications 5 à 11, dans lequel le promoteur lié fonctionnellement au polynucléotide selon la revendication 1 est choisi dans le groupe constitué par : un promoteur du gène précoce immédiat du cytomégalo virus (CMV), un promoteur du facteur d'élongation 1 alpha (EF 1- α), un promoteur de la phosphoglycérate kinase 1 (PGK), un promoteur de l'ubiquitine C (UBQ-C), un activateur de cytomégalo virus/promoteur de bêta-actine de poulet (CAG), un activateur de polyome/promoteur de thymidine kinase du virus herpès simplex (MC1), un promoteur de bêta-actine (β -ACT), un promoteur du virus simien 40 (VS40), et un activateur du virus du sarcome myéloprolifératif/promoteur à région de contrôle négative supprimée, site de liaison à l'amorce d1587rev substitué (MND).
13. Cellule effectrice immunitaire *ex vivo* ou *in vitro* comprenant le vecteur selon l'une quelconque des revendications 2 à 12.
14. Cellule effectrice immunitaire *ex vivo* ou *in vitro* selon la revendication 13, dans laquelle la cellule effectrice immunitaire est choisie dans le groupe constitué par : un lymphocyte T et une cellule tueuse naturelle (NK).
15. Composition comprenant la cellule effectrice immunitaire *ex vivo* ou *in vitro* selon la revendication 13 ou la revendication 14 et un excipient physiologiquement acceptable.






FIG. 1

*FIG. 2A*

*FIG. 2B*




FIG. 3

24 hour cytokine secretion
5e4 CAR T cells in 200ul




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8A




FIG. 8B




FIG. 9




FIG. 10A




FIG. 10B




FIG. 11B





FIG. 11C

*FIG. 12A*




FIG. 12B




FIG. 12C




FIG. 12D




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2013154760 A1, Carpenter, R.O. [0004]
- US 5283173 A [0031]
- US 5468614 A [0031]
- WO 9404678 A [0045]
- WO 9425591 A [0045]
- US 6005079 A [0045]
- EP 404097 A [0050]
- WO 199301161 A [0050]
- US 4873192 A [0089]
- US 61241647 A [0091]
- US 4554101 A [0094]
- WO 2002088346 A [0139]
- US 9108442 W [0150]
- US 9405601 W [0150]
- US 6682907 B [0168]
- US 6013516 A, Dull [0173]
- US 5994136 A [0173]
- WO 2006010834 A [0183]
- US 6352694 B [0216] [0248]
- US 6534055 B [0216] [0248]
- US 6905680 B [0216] [0248]
- US 6692964 B [0216] [0248]
- US 5858358 A [0216] [0248]
- US 6887466 B [0216] [0248]
- US 6905681 B [0216] [0248]
- US 7144575 B [0216] [0248]
- US 7067318 B [0216] [0248]
- US 7172869 B [0216] [0248]
- US 7232566 B [0216] [0248]
- US 7175843 B [0216] [0248]
- US 5883223 A [0216] [0248]
- US 6905874 B [0216] [0248]
- US 6797514 B [0216] [0248]
- US 6867041 B [0216] [0248]
- US 20060121005 A [0216]
- US 6040177 A [0218]
- US 5827642 A [0218]
- WO 2012129514 A [0218]
- WO 03057171 A [0219] [0273]
- US 20030147869 A [0219] [0273]
- US 62091419 B [0392]
- US 62200505 B [0392]

Non-patent literature cited in the description

- *Clin Cancer Res*, 2013, vol. 19 (8), 2048-2060 [0004]
- **WANG, X.** *Blood*, 2014, vol. 124 (21) [0004]
- **THOMPSON et al.** *J. Exp. Medicine*, 2000, vol. 192 (1), 129-135 [0012]
- **MACKAY et al.** *Annu. Rev. Immunol.*, 2003, vol. 21, 231-264 [0012]
- **KALLED et al.** *Immunological Reviews*, 2005, vol. 204, 43-54 [0012]
- **LAABI et al.** *EMBO J.*, 1992, vol. 77 (1), 3897-3904 [0012]
- **LAABI et al.** *Nucleic Acids Res.*, 1994, vol. 22 (7), 1147-1154 [0012]
- **O'CONNOR et al.** *J. Exp. Medicine*, 2004, vol. 199 (1), 91-97 [0012]
- **NG et al.** *J. Immunol.*, 2004, vol. 73 (2), 807-817 [0012]
- **XU et al.** *Mol. Cell. Biol.*, 2001, vol. 21 (12), 4067-4074 [0012]
- **SCHIEMANN et al.** *Science*, 2001, vol. 293 (5537), 2111-2114 [0012]
- **NOVAK et al.** *Blood*, 2004, vol. 103 (2), 689-694 [0012]
- **NERI et al.** *Clinical Cancer Research*, 2007, vol. 73 (19), 5903-5909 [0012]
- **BELLUCCI et al.** *Blood*, 2005, vol. 105 (10), 3945-3950 [0012]
- **MOREAUX et al.** *Blood*, 2004, vol. 703 (8), 3148-3157 [0012]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. 2001 [0017]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. 1989 [0017]
- **MANIATIS et al.** *Molecular Cloning: A Laboratory Manual*. 1982 [0017]
- **AUSUBEL et al.** *Current Protocols in Molecular Biology*. John Wiley and Sons, 2008 [0017]
- *Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology*. Greene Pub. Associates and Wiley-Interscience [0017]
- **GLOVER.** *DNA Cloning: A Practical Approach*. IRL Press, 1985, vol. I, II [0017]
- **ANAND.** *Techniques for the Analysis of Complex Genomes*. Academic Press, 1992 [0017]
- *Transcription and Translation*. 1984 [0017]
- **PERBAL.** *A Practical Guide to Molecular Cloning*. 1984 [0017]

- **HARLOW ; LANE.** Antibodies. Cold Spring Harbor Laboratory Press, 1998 [0017]
- Current Protocols in Immunology. 1991 [0017]
- **SCATCHARD et al.** *Ann. N.Y. Acad. Sci.*, 1949, vol. 51, 660 [0031]
- Pierce Catalog and Handbook. Pierce Chemical Co, 1994 [0036]
- **KUBY. J.**, Immunology. W. H. Freeman & Co, 1997 [0036]
- **WU, TT ; KABAT, E. A.** *J Exp Med*, 1970, vol. 132 (2), 211-50 [0038]
- **BORDEN, P. ; KABAT E. A.** *PNAS*, 1987, vol. 84, 2440-2443 [0038]
- **KABAT et al.** Sequences of Proteins of Immunological Interest. U.S. Department of Health and Human Services, 1991 [0038]
- **CHOITHIA, C. ; LESK, A.M.** *J Mol. Biol.*, 1987, vol. 196 (4), 901-917 [0038]
- **CHOITHIA, C. et al.** *Nature*, 1989, vol. 342, 877-883 [0038]
- **KOCH-NOLTE et al.** *FASEB J.*, 2007, vol. 21, 3490-3498 [0045]
- **RIECHMANN L. et al.** *J. Immunol. Methods*, 1999, vol. 231, 25-38 [0045]
- **HUDSON et al.** *Nat. Med.*, 2003, vol. 9, 129-134 [0050]
- **HOLLINGER et al.** *PNAS USA*, 1993, vol. 90, 6444-6448 [0050]
- **HOLT, L. et al.** *Trends in Biotechnology*, vol. 21 (11), 484-490 [0051]
- **PLUCKTHÜN.** The Pharmacology of Monoclonal Antibodies. Springer-Verlag, 1994, vol. 113, 269-315 [0052]
- **ORLANDI et al.** *PNAS*, 1989, vol. 86, 3833-3837 [0053]
- **SCHERAGA.** *Rev. Computational Chem.*, 1992, 11173-142 [0057]
- **LIU et al.** *PNAS*, 1997, 5525-5530 [0058]
- **KIM et al.** *PNAS*, 1996, vol. 93, 1156-1160 [0058]
- **CHAUDHARY et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1990, vol. 87, 1066-1070 [0058]
- **BIRD et al.** *Science*, 1988, vol. 242, 423-426 [0058]
- **DESJARLAIS ; BERG.** *PNAS*, 1993, vol. 90, 2256-2260 [0058]
- *PNAS*, 1994, vol. 91, 11099-11103 [0058]
- **COOPER et al.** *Blood*, 2003, vol. 101 (4), 1637-1644 [0058]
- **PATEL et al.** *Gene Therapy*, 1999, vol. 6, 412-419 [0059]
- **KUNKEL.** *Proc. Natl. Acad. Sci. USA.*, 1985, vol. 82, 488-492 [0089]
- **KUNKEL et al.** *Methods in Enzymol*, 1987, vol. 154, 367-382 [0089]
- **WATSON, J. D. et al.** Molecular Biology of the Gene. Benjamin/Cummings, 1987 [0089]
- **DAYHOFF et al.** Atlas of Protein Sequence and Structure. Natl. Biomed. Res. Found, 1978 [0089]
- **WATSON et al.** Molecular Biology of the Gene. The Benjamin/Cummings Pub. Co, 1987, 224 [0091]
- **DEFELIPE ; RYAN.** *Traffic*, 2004, vol. 5 (8), 616-26 [0100]
- **RYAN et al.** *J. Gener. Virol.*, 1997, vol. 78, 699-722 [0101]
- **SCYMCZAK et al.** *Nature Biotech.*, 2004, vol. 5, 589-594 [0101]
- **DONNELLY et al.** *J. Gen. Virol.*, 2001, vol. 82, 1027-1041 [0103]
- **ALTSCHUL et al.** *Nucl. Acids Res.*, 1997, vol. 25, 3389 [0109]
- **AUSUBEL et al.** Current Protocols in Molecular Biology. John Wiley & Sons Inc, 1994 [0109]
- **ZAMORE et al.** *Cell*, 2000, vol. 101, 25-33 [0117]
- **FIRE et al.** *Nature*, 1998, vol. 391, 806 [0117]
- **HAMILTON et al.** *Science*, 1999, vol. 286, 950-951 [0117]
- **LIN et al.** *Nature*, 1999, vol. 402, 128-129 [0117]
- **SHARP.** *Genes & Dev.*, 1999, vol. 13, 139-141 [0117]
- **STRAUSS.** *Science*, 1999, vol. 286, 886 [0117]
- **IRIONS et al.** *Nature Biotechnology*, 2007, vol. 25, 1477-1482 [0133]
- **CHALLITA et al.** *J Virol.*, 1995, vol. 69 (2), 748-55 [0133]
- **SIRIN et al.** *Gene*, 2003, vol. 323, 67 [0139]
- **LANDY.** *Current Opinion in Biotechnology*, 1993, vol. 3, 699-707 [0140]
- **SAUER, B.** *Current Opinion in Biotechnology*, 1994, vol. 5, 521-527 [0142]
- **JACKSON et al.** *Trends Biochem Sci*, 1990, vol. 15 (12), 477-83 [0145]
- **JACKSON ; KAMINSKI.** *RNA*, 1995, vol. 1 (10), 985-1000 [0145]
- **KOZAK.** *Cell*, 1986, vol. 44 (2), 283-92 [0146]
- **KOZAK.** *Nucleic Acids Res.*, 1987, vol. 15 (20), 8125-48 [0146]
- **WIGLER et al.** *Cell*, 1977, vol. 11, 223 [0148]
- **MULLEN et al.** *Proc. Natl. Acad. Sci. USA.*, 1992, vol. 89, 33 [0148]
- **LUPTON S. D. et al.** *Mol. and Cell. Biology*, 1991, vol. 1 (1), 3374-3378 [0150]
- **MILLER.** *Nature*, 2000, vol. 357, 455-460 [0153]
- **CLEVER et al.** *J. of Virology*, 1995, vol. 69 (4), 2101-2109 [0162]
- **ZENNOU et al.** *Cell*, 2000, vol. 101, 173 [0168]
- **CULLEN et al.** *J. Virol.*, 1991, vol. 65, 1053 [0169]
- **CULLEN et al.** *Cell*, 1991, vol. 58, 423 [0169]
- **ZUFFEREY et al.** *J. Virol.*, 1999, vol. 73, 2886 [0170]
- **HUANG et al.** *Mol. Cell. Biol.*, vol. 5, 3864 [0170]
- **LIU et al.** *Genes Dev.*, 1995, vol. 9, 1766 [0170]
- **BURGESS-BEUSSE et al.** *Proc. Natl. Acad. Sci., USA*, 2002, vol. 99, 16433 [0172]
- **ZHAN et al.** *Hum. Genet.*, 2001, vol. 109, 471 [0172]
- **CHUNG et al.** *Cell*, 1993, vol. 74, 505 [0172]
- **CHUNG et al.** *PNAS*, 1997, vol. 94, 575 [0172]
- **BELL et al.** *Cell*, 1999, vol. 98, 387 [0172]

- **Y. SONEOKA et al.** *Nucl. Acids Res.*, 1995, vol. 23, 628-633 [0195]
- **N. R. LANDAU et al.** *J. Virol.*, 1992, vol. 66, 5110-5113 [0195]
- **KAY, M. A.** *Chest*, 1997, vol. 111 (6), 138S-142S [0196]
- **FERRY, N. ; HEARD, J. M.** *Hum. Gene Ther.*, 1998, vol. 9, 1975-81 [0196]
- **SHIRATORY, Y. et al.** *Liver*, 1999, vol. 19, 265-74 [0196]
- **OKA, K. et al.** *Curr. Opin. Lipidol.*, 2000, vol. 11, 179-86 [0196]
- **THULE, P. M. ; LIU, J. M.** *Gene Ther.*, 2000, vol. 7, 1744-52 [0196]
- **YANG, N. S.** *Crit. Rev. Biotechnol.*, 1992, vol. 12, 335-56 [0196]
- **ALT, M. J.** *Hepatology*, 1995, vol. 23, 746-58 [0196]
- **BRODY, S. L. ; CRYSTAL, R. G.** *Ann. N.Y. Acad. Sci.*, 1994, vol. 716, 90-101 [0196]
- **STRAYER, D. S.** *Expert Opin. Investig. Drugs*, 1999, vol. 8, 2159-2172 [0196]
- **SMITH-ARICA, J. R. ; BARTLETT, J. S.** *Curr. Cardiol. Rep.*, 2001, vol. 3, 43-49 [0196]
- **LEE, H. C. et al.** *Nature*, 2000, vol. 408, 483-8 [0196]
- **BERG et al.** *Transplant Proc.*, 1998, vol. 30 (8), 3975-3977 [0218]
- **HAANEN et al.** *J. Exp. Med.*, 1999, vol. 190 (9), 1319-1328 [0218]
- **GARLAND et al.** *J. Immunol Meth.*, 1999, vol. 227 (1-2), 53-63 [0218]
- **LOVELOCK ; BISHOP.** *Nature*, 1959, vol. 183, 1394-1395 [0225] [0234]
- **ASHWOOD-SMITH.** *Nature*, 1961, vol. 190, 1204-1205 [0225] [0234]
- **RINFRET.** *Ann. N.Y. Acad. Sci.*, 1960, vol. 85, 576 [0225] [0234]
- **SLOVITER ; RAVDIN.** *Nature*, 1962, vol. 196, 48 [0225] [0234]
- **MEUER, S. C. et al.** *Cell*, 1984, vol. 36, 897-906 [0254]
- **YANG, S. Y. et al.** *J. Immunol.*, 1986, vol. 137, 1097-1100 [0254]
- **LIU et al.** *Nature Review*, 2009, vol. 8, 627-644 [0303]
- **Harrison's Principles of Internal Medicine.** McGraw-Hill, 2001 [0338]
- **Current Protocols in Immunology.** John Wiley & Sons, 2001 [0354]
- **PNAS**, 1994, vol. 91, 9564-68 [0359]
- **MALDARELLI.** *J Virol*, 1991, vol. 65 (11), 5732-43 [0359]
- **MALIM, M. H.** *Nature*, 1988, vol. 335, 181-183 [0359]
- **CHALLITA.** *J. Virol.*, 1995, vol. 69, 748-755 [0359]
- **MILONE.** *Mol Ther*, 2009, vol. 17 (8), 1453-64 [0359]
- **LEVITT, N.** *Genes & Dev*, 1989, vol. 3, 1019-1025 [0359]
- **AVERY et al.** BAFF selectively enhances the survival of plasmablasts generated from human memory B cells. *J Clin Invest*, 2003, vol. 112 (2), 286-297 [0370]
- **CARPENITO et al.** Control of large, established tumor xenografts with genetically retargeted human T cells containing CD28 and CD137 domains. *Proc Natl Acad Sci USA*, 2009, vol. 106 (9), 3360-3365 [0371]
- **CHIU et al.** Hodgkin lymphoma cells express TACI and BCMA receptors and generate survival and proliferation signals in response to BAFF and APRIL. *Blood*, 2007, vol. 109 (2), 729-739 [0372]
- **SANCHEZ et al.** Serum B-cell maturation antigen is elevated in multiple myeloma and correlates with disease status and survival. *Br J Haematol*, 2012, vol. 158 (6), 727-738 [0373]