



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2425174 C 2013/07/30

(11)(21) **2 425 174**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2001/04/04

(87) Date publication PCT/PCT Publication Date: 2002/04/25

(45) Date de délivrance/Issue Date: 2013/07/30

(85) Entrée phase nationale/National Entry: 2003/04/07

(86) N° demande PCT/PCT Application No.: US 2001/010981

(87) N° publication PCT/PCT Publication No.: 2002/032955

(30) Priorité/Priority: 2000/10/18 (US60/241,510)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
A61K 38/17 (2006.01), *A61P 15/00* (2006.01),
A61P 15/18 (2006.01), *C07H 21/00* (2006.01),
C07K 14/47 (2006.01), *C12Q 1/68* (2006.01),
G01N 33/53 (2006.01)

(72) Inventeurs/Inventors:
NELSON, LAWRENCE M., US;
TONG, ZHI-BIN, US

(73) Propriétaire/Owner:
THE GOVERNMENT OF THE UNITED STATES OF
AMERICA, AS REPRESENTED BY THE
SECRETARY, DEPARTMENT OF HEALTH &
HUMAN SERVICES, US

(74) Agent: SMART & BIGGAR

(54) Titre : GENE HUMAIN INDISPENSABLE A LA FERTILITE

(54) Title: HUMAN GENE CRITICAL TO FERTILITY

(57) **Abrégé/Abstract:**

Disclosed herein are novel nucleic acid and protein sequences that are essential to fertility. In particular, human Mater cDNA and protein sequences are provided. Functional MATER is required for female fertility; zygotes that arise from Mater null oocytes do not progress beyond the two-cell stage. Methods are described for using Mater molecules in diagnoses, prognosis, and treatment of fertility and reduced fertility. Also provided are methods for using MATER as a contraceptive agent. The disclosure also describes compounds involved in such methods, and the identification of such compounds.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number
WO 02/32955 A1

(51) International Patent Classification⁷: **C07K 14/47**,
C12N 15/12, C12Q 1/68, G01N 33/53, C12N 15/11, 5/06,
A01K 67/027, C07K 16/18, G01N 33/50, A61K 38/17,
A61P 15/00, 15/18

VA 22015-4295 (US). **TONG, Zhi-Bin** [CN/US]; 7826
Oracle Place, Potomac, MD 20854 (US).

(21) International Application Number: PCT/US01/10981

(74) Agent: **HARDING, Tanya, M.**; Klarquist, Sparkman,
Campbell, Leigh & Winston, L, LP, One World Trade
Center, Suite 1600, 121 SW Salmon Street, Portland, OR
97204 (US).

(22) International Filing Date: 4 April 2001 (04.04.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/241,510 18 October 2000 (18.10.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **THE
GOVERNMENT OF THE UNITED STATES OF
AMERICA, as represented by THE SECRETARY
DEPARTMENT OF HEALTH & HUMAN SER-
VICES, THE NATIONAL INSTITUTES OF HEALTH**
[US/US]; Office of Technology Transfer, Suite 325, 601
Executive Boulevard, Rockville, MD 20852 (US).

Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **NELSON,
Lawrence, M.** [US/US]; 9513 Millgate Place, Burke,

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: HUMAN GENE CRITICAL TO FERTILITY

(57) Abstract: Disclosed herein are novel nucleic acid and protein sequences that are essential to fertility. In particular, human *Mater* cDNA and protein sequences are provided. Functional MATER is required for female fertility; zygotes that arise from *Mater* null oocytes do not progress beyond the two-cell stage. Methods are described for using *Mater* molecules in diagnoses, prognosis, and treatment of fertility and reduced fertility. Also provided are methods for using MATER as a contraceptive agent. The disclosure also describes compounds involved in such methods, and the identification of such compounds.



WO 02/32955 A1

-1-

HUMAN GENE CRITICAL TO FERTILITY**FIELD**

The present disclosure is generally related to fertility, including the mechanisms controlling it, diseases that arise from defects in such mechanisms, and methods of influencing (either inhibiting or enhancing) fertility.

BACKGROUND

Premature ovarian failure (POF) is a term used to describe certain types of infertility in women. As many as 1% of all women in the United States are thought to be afflicted with POF, which manifests as menopausal-type symptoms, including infertility, in women under the age of 40. Many different diseases and conditions can cause POF, including underlying chromosomal defects (e.g., X-chromosome fragility), chemotherapy, or radiation treatment. Autoimmunity is a well-established mechanism of premature ovarian failure (see Yan *et al.*, *J Womens Health Gend Based Med.* 9:275-87, 2000; and Kalantaridou & Nelson, *J Am Med Womens Assoc.* 53:18-20, 1998). In autoimmune infertility, a woman's ovaries are attacked by cells of her own immune system, leading to a condition known as autoimmune oophoritis (inflammation of the ovary). Autoimmune disease can develop in response to a single inciting antigen and then spread to involve other antigenic molecules of the same organ (Kaufman, *Nature* 366:69-72, 1993). Therefore, identifying the autoantigen target in an organ-specific autoimmune disease is essential to understanding its pathogenesis.

An experimental animal (mouse) model has been used to gain insight into the mechanisms of human autoimmune oophoritis. Removal of the thymus (thymectomy) in neonatal mice (about three days old) induces experimental autoimmune oophoritis in certain strains of mice (Taguchi *et al.*, *Clin Exp Immunol.* 42:324-331, 1980). This experimentally induced condition leads to the production of high levels of anti-ooplasm antibodies and sterility, accompanied by follicular degeneration; the progression of the condition appears to closely parallel human autoimmune oophoritis (Kalantaridou & Nelson, *J Am Med Womens Assoc.* 53:18-20, 1998).

Maternal products control the developmental program until embryonic genome activation takes place. Maternal effect genes that are important in early embryonic development have been well documented in *Drosophila* and *Xenopus* (Morisato & Anderson, *Annu. Rev. Genet.* 29:371-399, 1995; Newport & Kirschner, *Cell* 30:687-696, 1982), but their presence has only been inferred in mammals (Gardner, *Hum. Reprod. Update* 2:1-27, 1999). In mice, embryonic transcription is first detected in the late 1-cell zygote stage and is required for development beyond the 2-cell stage (Schultz, *Bioessays* 15:531-538, 1993; Flach *et al.*, *EMBO J.* 1:681-686, 1982; Latham *et al.*, *Mol. Reprod. Dev.* 35:140-150, 1993). The factors governing this transition from the maternal to the embryonic genome are unknown.

A critical transition in development occurs with the switch from dependence on proteins stored in the egg to those that result from activation of the embryonic genome. This shift which

-2-

occurs at the two-cell stage in mice is dependent on maternal factors. Gene transcription and protein translation are active during murine oogenesis and RNAs and proteins accumulate within oocytes. However, germ cells become transcriptionally inactive late in oogenesis and much of the maternal RNA is degraded during meiotic maturation and ovulation of the egg into the oviduct. Thus, few maternal gene products persist past the two-cell embryo stage and none have been demonstrated directly to affect early development (Schultz, *Bioessays* 15, 531-538, 1993; Gardner, *Hum. Reprod. Update* 2, 3-27, 1996).

SUMMARY OF THE DISCLOSURE

Described herein is the human MATER protein, an approximately 135 kDa (predicted estimated molecular weight) cytoplasmic protein expressed in mammalian oocytes that is required for female fertility. Zygotes that arise from a MATER null oocyte do not progress beyond the two-cell stage.

Some embodiments are an isolated human MATER protein predicted to have an estimated molecular weight of about 125 kDa to about 135 kDa, in some embodiments more particularly about 134.2 kDa. For instance, this estimated molecular weight may be obtained by SDS-polyacrylamide gel electrophoresis and Western blotting, for instance using an antibody raised against a C-terminal peptide of murine MATER (e.g., residues 1093 to 1111 of SEQ ID NO: 6). MATER proteins include an amino acid sequence as set forth in SEQ ID NOs: 2 or 4 or 24, or a sequence having at least 65% sequence identity with SEQ ID NOs: 2 or 4 or 24. Certain specific examples of such proteins may contain one or more conservative variants within such sequences. The provided human MATER proteins have MATER protein biological activity, for instance in that they can complement a *Mater* null phenotype.

In certain embodiments, the human MATER protein is an autoantigen associated with autoimmune infertility. The protein may be expressed in oocyte cytoplasm, for instance the oocyte cytoplasm of a mammal such as a human.

One specific embodiment is thus an isolated human MATER protein predicted to have an estimated molecular weight of about 135 kDa, wherein the human MATER protein comprises amino acid sequences as set forth in SEQ ID NO: 2 and/or SEQ ID NO: 4 and/or SEQ ID NO: 24, the protein is an oocyte cytoplasm-specific autoantigen associated with autoimmune infertility, and it can complement a *Mater* null phenotype by permitting progression of an embryo beyond the two-cell stage.

Further embodiments are isolated nucleic acid molecules that encode such a MATER protein. Examples of such nucleic acid molecules include a sequence as set forth in SEQ ID NOs: 1 or 3 or 23, or a sequence having at least 82% sequence identity with SEQ ID NOs: 1 or 3 or 23. Certain examples of such nucleic acid molecules may contain one or more conservative variants within such sequences. Certain examples hybridizes with a nucleic acid probe that includes the sequence shown in SEQ ID NO: 1 or SEQ ID NO: 3 or SEQ ID NO: 23 under wash conditions of 55° C, 0.2 x SSC and 0.1% SDS, or under wash conditions of 50° C, 2 x SSC, 0.1 % SDS.

-3-

Also provided are recombinant nucleic acid molecules that include a promoter sequence operably linked to a MATER nucleic acid molecule and cells transformed with such a recombinant nucleic acid molecule.

Methods of detecting a biological condition associated with an abnormal *Mater* nucleic acid or an abnormal MATER expression or an autoimmune response to MATER in a subject are also provided. Examples of such a biological condition include infertility (such as autoimmune infertility) or reduced fertility, or an increased susceptibility to infertility or reduced fertility. Such methods can involve detecting an abnormal *Mater* nucleic acid or an abnormal *Mater* expression or the autoimmune response to MATER in the subject. In specific examples of such methods, the abnormal *Mater* nucleic acid or abnormal *Mater* expression includes an alteration in cellular level of *Mater* nucleic acid or MATER protein, in comparison to a normal level. The abnormal *Mater* expression may include an increased or decreased expression of *Mater* in a subject.

Specific provided methods involve determining whether the subject has circulating autoantibodies that recognize an epitope of a MATER protein, wherein presence of such autoantibodies indicates the infertility or reduced fertility of the subject, or an increased susceptibility of the subject to infertility or reduced fertility. Other specific methods involve reacting at least one *Mater* molecule contained in a clinical sample from the subject with a reagent that includes a *Mater* specific binding agent (such as an oligonucleotide or a MATER protein specific binding agent (e.g., an antibody or functional fragment thereof), to form a *Mater*:agent complex.

Also provided are methods for detecting a predisposition to infertility or reduced fertility or for presymptomatic screening of an individual for infertility or reduced fertility.

Specific methods of detecting a biological condition provided herein involve *in vitro* amplifying a *Mater* nucleic acid prior to detecting the abnormal *Mater* nucleic acid. Amplification can be performed, for instance, using at least one oligonucleotide primer derived from a MATER protein encoding sequence. Examples of such oligonucleotides may include at least 15, for instance at least 20 or at least 23, contiguous nucleotides from SEQ ID NO: 1 or SEQ ID NO: 3 or SEQ ID NO: 23.

Also provided are oligonucleotide primers used in such methods, recombinant DNA vectors that contain such nucleic acid molecules, and recombinant nucleic acid molecules that include a promoter sequence operably linked (in sense or antisense orientation) to such a nucleic acid molecule. Such vectors and recombinant nucleic acid molecules can be transformed into cells or animals (e.g., non-human animals); such transformed cells and animals are also provided.

In certain examples of the provided methods, the *Mater* molecule is a MATER encoding sequence. In some of such examples, a *Mater*:agent complex is detected by nucleotide hybridization, for instance where the agent is a labeled nucleotide probe. Such probes may include a sequence as shown in SEQ ID NOs: 1 or 3 or 23, fragments of these sequences (for instance, fragments of 23 or more nucleotides). Other probes may contain a sequence that shares at least 70% sequence identity with a sequence as shown in SEQ ID NOs: 1 or 3 or 23. Such nucleotide probes are also provided.

-4-

In other examples of the provided methods, the *Mater* molecule is a MATER protein, which may contain the sequence of SEQ ID NOs: 2 or 4 or 24, a sequence sharing at least 65% sequence identity with SEQ ID NOs: 2 or 4 or 24, or conservative variants thereof. In some of such examples, Mater:agent complexes are detected by Western blot assay or by ELISA. In certain methods
5 provided, the *Mater*-specific binding agent is a MATER-specific antibody or a functional fragment thereof, for instance a polyclonal or monoclonal antibody. In specific examples, the antibody recognizes a peptide that includes the sequence of SEQ ID NOs: 2 or 4 or 24, or an antigenic fragment of one of these peptides.

Also provided herein are kits, including kits for detecting an over- or under-abundance of
10 MATER protein or *Mater* nucleic acid (for instance, in a sample from a subject, such as a mammal), which kits include a MATER protein specific binding agent (such as an antibody or a functional fragment thereof). In certain examples, the agent is capable of specifically binding to an epitope within the amino acid sequence shown in SEQ ID NOs: 2 or 4 or 24, an amino acid sequences that differ from these by one or more conservative amino acid substitutions, an amino acid sequences
15 having at least 65% sequence identity to these sequences, or an antigenic fragment of any of these sequences. Particular examples of such kits further include a means for detecting binding of the MATER protein binding agent to a MATER polypeptide. In certain examples of these kits, the overabundance or underabundance of MATER protein or Mater nucleic acid that is detected results in altered infertility.

20 Other provided kits are for detecting a genetic mutation (*e.g.*, a mutation in a *Mater* sequence) in a sample of nucleic acid. Such kits may include an oligonucleotide capable of specifically hybridizing with a *Mater* nucleic acid (which may be provided in a first container), and a fluorescent labeled nucleic acid probe (for instance, of about 5 to 500 nucleotides) that is fully complementary to the oligonucleotide (which may be provided in a second container).

25 Specific kit embodiments provided herein are for determining whether or not a subject has a biological condition associated with an abnormal *Mater* expression by detecting an underabundance of MATER protein in a sample of tissue and/or body fluids from the subject. Such kits include an antibody specific for MATER protein and instructions for using the kit. Such instructions may indicate steps for performing a method to detect the presence of MATER protein in the sample (for
30 instance, using a method described herein); and analyzing data generated by the method, wherein the instructions indicate that underabundance of MATER protein in the sample indicates that the individual has or is predisposed to the biological condition. Specific examples of such kits further include a detectable antibody that binds to the MATER protein specific antibody.

Other specific kit embodiments include a MATER protein specific antibody (which may be
35 provided in a container), a negative control sample which may be provided in a container), instructions for using the kit. Such instructions may indicating steps for performing a test assay to detect a quantity of MATER protein in a test sample of tissue and/or bodily fluid from the subject (such as a test assay provided herein), performing a negative control assay to detect a quantity of MATER protein in the negative control sample; and comparing data generated by the test assay and

63198-1404

-5-

negative control assay. In specific examples of such kits, the instructions indicate that a quantity of MATER protein in the test sample that is less than the quantity of MATER protein in the negative control sample indicates that the
5 subject has the biological condition. Some of such kits further include a detectable antibody that binds to the antibody specific for MATER protein (which may be provided in a separate container).

Also provided are methods of modifying the level
10 of expression of a MATER protein in a subject, for instance by expressing in the subject a recombinant genetic construct that includes a promoter operably linked (in either sense or antisense orientation) to a nucleic acid molecule that includes at least 23 consecutive nucleotides of
15 SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 23 or a sequence at least 70% identical to SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 23. In particular examples, expression of the nucleic acid molecule changes expression of the MATER protein.

20 A further embodiment provides methods of screening for a compound useful in influencing MATER-mediated fertility in a mammal. Such methods involve determining if a test compound (for instance, when it is applied to a test cell) binds to or interacts with a MATER protein, such as a
25 human MATER protein, or a variant or fragment thereof, and selecting a compound that so binds. In particular examples of such methods, binding of the compound inhibits a MATER protein biological activity. Also encompassed herein are compounds selected by these methods, for instance such
30 compounds for use as contraceptives or fertility enhancers.

The foregoing and other features and advantages of the invention will become more apparent from the following

63198-1404

-5a-

detailed description of several embodiments, which proceeds with reference to the accompanying figures.

According to one aspect of the present invention, there is provided an isolated human MATER protein comprising an amino acid sequence at least 95% identical to full length SEQ ID NO: 24, wherein the human MATER protein complements a *Mater* null phenotype.

According to another aspect of the present invention, there is provided an isolated nucleic acid molecule encoding the isolated human MATER protein as described above, comprising a nucleotide sequence at least 95% identical to full length SEQ ID NO: 23.

According to still another aspect of the present invention, there is provided a recombinant nucleic acid molecule comprising a promoter sequence operably linked to the isolated nucleic acid molecule as described above.

According to yet another aspect of the present invention, there is provided a cell transformed with the recombinant nucleic acid molecule as described above.

According to a further aspect of the present invention, there is provided a method of detecting a *Mater* nucleic acid in a human subject, comprising detecting a labeled nucleotide probe that hybridizes to a *Mater* nucleotide sequence at least 95% identical to full length SEQ ID NO: 23.

According to yet a further aspect of the present invention, there is provided a method for determining whether a sample from a human subject has circulating autoantibodies that recognize an epitope of a MATER protein, comprising reacting a sample from the human subject with a human MATER protein comprising a MATER amino acid sequence at least 95% identical full length to SEQ ID NO: 24 to form a MATER:autoantibody complex, wherein detection of the MATER:autoantibody complex indicates the presence of autoantibodies in the sample.

63198-1404

-5b-

According to still a further aspect of the present invention, there is provided a method for detecting Mater nucleic acid expression in a human subject, comprising: reacting at least one human Mater nucleic acid contained in a clinical sample from the human subject, wherein the human Mater nucleic acid is a MATER
5 encoding nucleic acid comprising a Mater nucleotide sequence at least 95% identical to full length SEQ ID NO: 23 with a Mater oligonucleotide, wherein the Mater oligonucleotide comprises at least 23 contiguous nucleotides having at least 95% sequence identity to the corresponding sequence within SEQ ID NO: 23 to form a *Mater:Mater* oligonucleotide complex; and detecting the *Mater:Mater*
10 oligonucleotide complex, thereby detecting Mater nucleic acid expression in the subject.

According to another aspect of the present invention, there is provided a method for detecting MATER protein expression in a human subject, comprising: reacting at least one human MATER protein contained in a clinical sample from the
15 human subject, wherein the human MATER protein is a MATER protein comprising an amino acid sequence at least 95% identical to full length SEQ ID NO: 24 with a MATER protein specific antibody or fragment thereof that binds specifically to an epitope within an amino acid sequence at least 95% identical to full length SEQ ID NO: 24 to form a MATER:MATER-specific antibody complex; and detecting the
20 MATER:MATER-specific antibody complex, thereby detecting MATER protein expression in the subject.

According to another aspect of the present invention, there is provided an oligonucleotide primer for use in the method as described above.

According to yet another aspect of the present invention, there is
25 provided a recombinant DNA vector comprising the isolated nucleic acid as described above.

63198-1404

-5c-

According to yet another aspect of the present invention, there is provided a recombinant nucleic acid molecule comprising a promoter sequence operably linked to the isolated nucleic acid as described above.

According to another aspect of the present invention, there is provided
5 a cell transformed with the recombinant nucleic acid molecule as described above.

According to yet another aspect of the present invention, there is provided a kit for detecting MATER protein in a subject using the method as described above, comprising a MATER protein specific antibody or fragment thereof together with instructions for detecting a MATER protein.

10 According to yet another aspect of the present invention, there is provided a kit for detecting a MATER nucleic acid in a sample from a human subject, comprising: (a) a first container containing an oligonucleotide comprising at least 23 contiguous nucleotides having at least 95% sequence identity to the corresponding sequence within SEQ ID NO: 23, which specifically hybridizes with a *Mater* nucleic
15 acid at least 95% identical to full length SEQ ID NO: 23; and (b) a second container containing a fluorescent labeled nucleic acid probe that is fully complementary to the oligonucleotide.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows two digitized gels, and a series of digitized micrographs,
20 detailing the developmental expression of murine MATER.

FIG. 1A is a digitized image of a Northern blots showing RNase protection of *Mater* and ZP3 transcripts. RNase protection assays were performed using total RNA from 50 oocytes, eggs or embryos. ³²P-labeled antisense probes and protected fragment lengths for *Mater* and ZP3 were 180/139 nt and 257/205 nt,
25 respectively.

63198-1404

-5d-

FIG. 1B is a digitized image of a Western blot, showing MATER protein (approximately 125 kD) amounts in the indicated murine cell types. MATER protein was assayed in 25 oocytes, eggs or embryos by immunoblotting with monospecific antisera to MATER.

5 *FIG. 1C* is a series of digitized micrographs, showing the cellular localization of MATER protein. Protein localization was determined using fluorescent labeled antibody, by imaging oocytes, eggs and embryos with laser-scanning confocal microscopy alone (upper images) or after superimposition on a Nomarski image (lower images). Scale bar, 50 μ m.

10 *FIG. 2* shows the characterization of targeted *Mater* locus disruptions.

-6-

FIG. 2A is a schematic representation of the normal murine *Mater* allele (upper), the targeting construct (middle) and the null allele (bottom). The 5' and 3' probes used to assess targeting were outside the region of DNA homology.

FIG. 2B is a series of digitized Northern blots, showing that the *Mater* null allele was
5 detected in XbaI digested ES cell DNA as a 3.5 kb fragment with the 5' (left) or as a 4.0 kb fragment with the 3' (middle) and neo (right) probes.

FIG. 2C is a pair of digitized Northern blots, showing that *Mater* transcripts were detected in normal and heterozygous (half the abundance of normal), but not in homozygous null ovaries (*left panel*). ZP3 transcripts (control) were present in all three genotypes (*right panel*).

10 FIG. 3 is a series of digitized micrographs and corresponding bar graphs, showing the *in vivo* development defects of embryos derived from *Mater* null female mice.

After gonadotrophin induced ovulation, female mice were mated with normal males and oviducts from normal (FIG. 3A, 3D, 3G, and 3J) or *Mater* null (FIG. 3B, 3E, 3H, and 3K) females were flushed one (FIG. 3A and 3B), two (FIG. 3D and 3E), three (FIG. 3G and 3H) and four (FIG. 3J
15 and 3K) days later. The unfixed embryos were photographed using Nomarski optics. The arrows point to pronuclei in 1-cell zygotes (FIG. 3A and 3B). Scale bar, 50 μ m.

The bar graphs indicate developmental progress of the average number of embryos derived from *Mater* null (■) and normal (□) females at one (FIG. 3C), two (FIG. 3F), three (FIG. 3I), and four (FIG. 3L) days after mating. Each bar represents the average of 4-5 experiments $\pm$ s.e.m.

20 FIG. 4 shows that *de novo* transcription and translation occurs in murine embryos lacking *Mater*.

FIG. 4A-4F show digitized micrograph images of murine embryos. Newly synthesized RNA was measured by BrUTP incorporation into the nucleus of one- (FIG. 4A and 4B) and 2-cell (FIG. 4C and 4D) embryos derived from *Mater* null (FIG. 4A and 4C) and normal (FIG. 4B and 4D) females
25 using laser-scanning confocal microscopy and a monoclonal antibody to BrUTP.

FIG. 4G is a bar graph, showing the quantity of BrUTP incorporated in 2-cell embryos with (+) and without (-) *Mater* using arbitrary fluorescence units after subtraction of that obtained with uninjected controls (FIG. 4E and 4F). Scale bar, 50 μ m.

FIG. 4H is a pair of digitized northern blots, showing RNase protection assays of 28S ribosomal RNA using total RNA from 50 growing oocytes, eggs and 2-cell embryos isolated from females with (+) and without (-) *Mater*. ^{32} P-labeled antisense probes and protected fragment lengths for 28S-rRNA were 153 nt and 115 nt, respectively.

FIG. 4I is a series of digitized fluorographs, showing *de novo* protein synthesis in 1-cell zygotes (left panel) and 2-cell embryos (middle and right panels) that do (+) or do not (-) contain
35 *Mater*. Each lane contained proteins from 10 embryos that were dissolved in sample buffer directly (left and middle panels) or after partial purification extraction of TRC (65-75 kDa) and p35 (right panels).

-7-

FIG. 5 shows ovarian histology, eggs and the number of eggs produced from *Mater* null females. The number of ovulated eggs and their morphology indicating maturation or degeneration were not remarkably different between normal and *Mater* null mice.

FIG. 5A and 5B are digitized images of ovarian slices from a 4 week old (FIG. 5A) and 8 week old (FIG. 5B) *Mater* null mouse. The 4 week old *Mater* null ovarian histology (FIG. 5A) displays different stages of normal ovarian follicles with oogenesis, which are indistinguishable from normal ovarian histology. In the 8 week old sample (FIG. 5B), there are a number of corpora lutea indicating normal spontaneous ovulation, similar to follicular luteinization in the wild-type mouse ovary.

FIG. 5C is a digitized photograph of ovulated eggs from the *Mater* null mice, produced in response to exogenous PMSG and hCG. Morphology appears similar to that of ovulated eggs from wild-type mice.

FIG. 5D is a bar graphs showing the numbers (mean $\pm$ s.e.m.) of the ovulated eggs from the *Mater* null and wild-type mice.

FIG. 6 is a digitized image comparing the gross morphological appearance of the uterus and ovaries of normal and *Mater* null mice. Ovaries and uteri of *Mater* null female mice (right) were indistinguishable from the ovaries and uteri from the wild-type female mice (left).

FIG. 7 is a series of digitized micrographs, showing oocyte-specific expression of *Mater* transcripts in human by *in situ* hybridization. Both [³⁵S]-labeled antisense and sense probes were synthesized by *in vitro* transcription using the cloned human *Mater* cDNA as a template. The frozen human ovarian sections were hybridized with the radiolabeled sense (FIG. 7A and 7C) and antisense (FIG. 7B and 7D) probes. The slides were stained with hematoxylin and eosin. For each probe, bright-field (FIG. 7A and 7B) and dark-field (FIG. 7C and 7D) images are shown.

FIG. 8 is a pair of digitized micrographs, showing oocyte-specific expression of human MATER protein. Frozen human ovarian sections were incubated with rabbit antisera (1:200) against a C-terminal peptide of mouse MATER protein, and FITC-conjugated goat anti-rabbit IgG antisera were used as the second antibody to detect human MATER protein in the oocyte (FIG. 8A). The corresponding phase contrast image is shown in FIG. 8B.

FIG. 9 is an alignment of the amino acid sequence of human MATER (SEQ ID NO: 24) and murine MATER (SEQ ID NO: 6) proteins. Identical amino acids are identified; similar amino acids are indicated with a plus (+) symbol.

SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. In the accompanying sequence listing:

-8-

SEQ ID NO: 1 shows the nucleic acid sequence of human *Mater* cDNA fragment 1.

SEQ ID NO: 2 shows the amino acid sequence of the human MATER peptide encoded by fragment 1 (SEQ ID NO: 1).

SEQ ID NO: 3 shows the nucleic acid sequence of human *Mater* cDNA fragment 2.

5 SEQ ID NO: 4 shows the amino acid sequence of the human MATER peptide encoded by fragment 2 (SEQ ID No: 3).

SEQ ID NO: 5 shows the nucleic acid sequence of murine *Mater* cDNA (GenBank Accession number NM_011860.1 and AF074018; individual exons are also listed in AF143559-AF143573).

10 SEQ ID NO: 6 shows the amino acid sequence of the murine MATER protein (GenBank Accession number NP_035990).

SEQ ID NOs: 7-22 shows several synthetic oligonucleotides useful as probes and/or primers.

SEQ ID NO: 23 shows the nucleic acid sequence of the human *Mater* cDNA, and the amino acid sequence so encoded.

15 SEQ ID NO: 24 shows the amino acid sequence of the human MATER protein.

DETAILED DESCRIPTION OF THE DISCLOSURE

I. Abbreviations

ES: embryonic stem

20 ***Mater***: Maternal Antigen That Embryos Require

OP1: Ooplasm-specific Protein 1

POF: Premature Ovarian Failure

TRC: Transcription Related Complex

ZP3: *zona pellucida* glycoprotein 3

25

II. Terms

Unless otherwise noted, technical terms are used according to conventional usage.

Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew *et al.* (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

30

In order to facilitate review of the various embodiments of the invention, the following explanations of specific terms are provided:

35

Abnormal: Deviation from normal characteristics. Normal characteristics can be found in a control, a standard for a population, etc. For instance, where the abnormal condition is an autoimmune disease condition, such as autoimmune infertility, sources of normal characteristics

might include an individual who is not suffering from the autoimmune disorder, a population standard of individuals believed not to be suffering from the disease, etc.

In some examples, abnormal may refer to a condition that is associated with a disease. The term "associated with" includes an increased risk of developing the disease as well as the disease
5 itself. For instance, a certain abnormality (such as an abnormality in an *Mater* nucleic acid or *MATER* protein expression) can be described as being associated with the biological conditions of altered (*e.g.*, reduced) fertility and tendency to develop autoimmune infertility.

An abnormal nucleic acid, such as an abnormal *Mater* nucleic acid, is one that is different in some manner to a normal (wildtype) nucleic acid. Such abnormality includes but is not necessarily
10 limited to: (1) a mutation in the nucleic acid (such as a point mutation (*e.g.*, a single nucleotide polymorphism) or short deletion or duplication of a few to several nucleotides); (2) a mutation in the control sequence(s) associated with that nucleic acid such that replication or expression of the nucleic acid is altered (such as the functional inactivation of a promoter); (3) a decrease in the amount or copy number of the nucleic acid in a cell or other biological sample (such as a deletion of the nucleic acid,
15 either through selective gene loss or by the loss of a larger section of a chromosome or under expression of the mRNA); and (4) an increase in the amount or copy number of the nucleic acid in a cell or sample (such as a genomic amplification of part or all of the nucleic acid or the overexpression of an mRNA), each compared to a control or standard. It will be understood that these types of abnormalities can co-exist in the same nucleic acid or in the same cell or sample; for instance, a
20 genomic-amplified nucleic acid sequence may also contain one or more point mutations. In addition, it is understood that an abnormality in a nucleic acid may be associated with, and in fact may cause, an abnormality in expression of the corresponding protein.

Abnormal protein expression, such as abnormal *MATER* protein expression, refers to expression of a protein that is in some manner different to expression of the protein in a normal
25 (wildtype) situation. This includes but is not necessarily limited to: (1) a mutation in the protein such that one or more of the amino acid residues is different; (2) a short deletion or addition of one or a few amino acid residues to the sequence of the protein; (3) a longer deletion or addition of amino acid residues, such that an entire protein domain or sub-domain is removed or added; (4) expression of an increased amount of the protein, compared to a control or standard amount; (5) expression of an
30 decreased amount of the protein, compared to a control or standard amount; (6) alteration of the subcellular localization or targeting of the protein; (7) alteration of the temporally regulated expression of the protein (such that the protein is expressed when it normally would not be, or alternatively is not expressed when it normally would be); and (8) alteration of the localized (*e.g.*, organ or tissue specific) expression of the protein (such that the protein is not expressed where it
35 would normally be expressed or is expressed where it normally would not be expressed), each compared to a control or standard.

Controls or standards appropriate for comparison to a sample, for the determination of abnormality, include samples believed to be normal as well as laboratory values, even though possibly arbitrarily set, keeping in mind that such values may vary from laboratory to laboratory.

-10-

Laboratory standards and values may be set based on a known or determined population value and may be supplied in the format of a graph or table that permits easy comparison of measured, experimentally determined values.

Antisense, Sense, and Antigene: Double-stranded DNA (dsDNA) has two strands, a 5' -> 3' strand, referred to as the plus strand, and a 3' -> 5' strand (the reverse complement), referred to as the minus strand. Because RNA polymerase adds nucleic acids in a 5' -> 3' direction, the minus strand of the DNA serves as the template for the RNA during transcription. Thus, the RNA formed will have a sequence complementary to the minus strand and identical to the plus strand (except that U is substituted for T).

Antisense molecules are molecules that are specifically hybridizable or specifically complementary to either RNA or the plus strand of DNA. Sense molecules are molecules that are specifically hybridizable or specifically complementary to the minus strand of DNA. Antigene molecules are either antisense or sense molecules directed to a dsDNA target.

Binding or stable binding: An oligonucleotide binds or stably binds to a target nucleic acid if a sufficient amount of the oligonucleotide forms base pairs or is hybridized to its target nucleic acid, to permit detection of that binding. Binding can be detected by either physical or functional properties of the target:oligonucleotide complex. Binding between a target and an oligonucleotide can be detected by any procedure known to one skilled in the art, including both functional and physical binding assays. Binding may be detected functionally by determining whether binding has an observable effect upon a biosynthetic process such as expression of a gene, DNA replication, transcription, translation, and the like.

Physical methods of detecting the binding of complementary strands of DNA or RNA are well known in the art, and include such methods as DNase I or chemical footprinting, gel shift and affinity cleavage assays, Northern blotting, dot blotting and light absorption detection procedures. For example, one method that is widely used, because it is so simple and reliable, involves observing a change in light absorption of a solution containing an oligonucleotide (or an analog) and a target nucleic acid at 220 to 300 nm as the temperature is slowly increased. If the oligonucleotide or analog has bound to its target, there is a sudden increase in absorption at a characteristic temperature as the oligonucleotide (or analog) and target disassociate from each other, or melt.

The binding between an oligomer and its target nucleic acid is frequently characterized by the temperature (T_m) at which 50% of the target sequence remains hybridized to a perfectly matched probe or complementary strand. A higher (T_m) means a stronger or more stable complex relative to a complex with a lower (T_m).

cDNA (complementary DNA): A piece of DNA lacking internal, non-coding segments (introns) and transcriptional regulatory sequences. cDNA may also contain untranslated regions (UTRs) that are responsible for translational control in the corresponding RNA molecule. cDNA is usually synthesized in the laboratory by reverse transcription from messenger RNA extracted from cells.

-11-

DNA (deoxyribonucleic acid): DNA is a long chain polymer which comprises the genetic material of most living organisms (some viruses have genes comprising ribonucleic acid (RNA)).

The repeating units in DNA polymers are four different nucleotides, each of which comprises one of the four bases, adenine (A), guanine (G), cytosine (C), and thymine (T) bound to a deoxyribose sugar to which a phosphate group is attached. Triplets of nucleotides (referred to as codons) code for each amino acid in a polypeptide, or for a stop signal. The term codon is also used for the corresponding (and complementary) sequences of three nucleotides in the mRNA into which the DNA sequence is transcribed.

Unless otherwise specified, any reference to a DNA molecule is intended to include the reverse complement of that DNA molecule. Except where single-strandedness is required by the text herein, DNA molecules, though written to depict only a single strand, encompass both strands of a double-stranded DNA molecule. Thus, a reference to the nucleic acid molecule that encodes a specific protein, or a fragment thereof, encompasses both the sense strand and its reverse complement. Thus, for instance, it is appropriate to generate probes or primers from the reverse complement sequence of the disclosed nucleic acid molecules.

Deletion: The removal of a sequence of DNA, the regions on either side of the removed sequence being joined together.

Gene amplification or genomic amplification: An increase in the copy number of a gene or a fragment or region of a gene or associated 5' or 3' region, as compared to the copy number in normal tissue. An example of a genomic amplification is an increase in the copy number of an oncogene. A "gene deletion" is a deletion of one or more nucleic acids normally present in a gene sequence and, in extreme examples, can include deletions of entire genes or even portions of chromosomes.

Hybridization: Oligonucleotides and their analogs hybridize by hydrogen bonding, which includes Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary bases. Generally, nucleic acid consists of nitrogenous bases that are either pyrimidines (cytosine (C), uracil (U), and thymine (T)) or purines (adenine (A) and guanine (G)). These nitrogenous bases form hydrogen bonds between a pyrimidine and a purine, and the bonding of the pyrimidine to the purine is referred to as "base pairing." More specifically, A will hydrogen bond to T or U, and G will bond to C. "Complementary" refers to the base pairing that occurs between two distinct nucleic acid sequences or two distinct regions of the same nucleic acid sequence.

"Specifically hybridizable" and "specifically complementary" are terms that indicate a sufficient degree of complementarity such that stable and specific binding occurs between the oligonucleotide (or its analog) and the DNA or RNA target. The oligonucleotide or oligonucleotide analog need not be 100% complementary to its target sequence to be specifically hybridizable. An oligonucleotide or analog is specifically hybridizable when binding of the oligonucleotide or analog to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA, and there is a sufficient degree of complementarity to avoid non-specific binding of the oligonucleotide or analog to non-target sequences under conditions where specific binding is desired,

63198-1404

-12-

for example under physiological conditions in the case of *in vivo* assays or systems. Such binding is referred to as specific hybridization.

Hybridization conditions resulting in particular degrees of stringency will vary depending upon the nature of the hybridization method of choice and the composition and length of the hybridizing nucleic acid sequences. Generally, the temperature of hybridization and the ionic strength (especially the Na^+ concentration) of the hybridization buffer will determine the stringency of hybridization, though waste times also influence stringency. Calculations regarding hybridization conditions required for attaining particular degrees of stringency are discussed by Sambrook *et al.* (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, chapters 9 and 11.

For present purposes, "stringent conditions" encompass conditions under which hybridization will only occur if there is less than 25% mismatch between the hybridization molecule and the target sequence. "Stringent conditions" may be broken down into particular levels of stringency for more precise definition. Thus, as used herein, "moderate stringency" conditions are those under which molecules with more than 25% sequence mismatch will not hybridize; conditions of "medium stringency" are those under which molecules with more than 15% mismatch will not hybridize, and conditions of "high stringency" are those under which sequences with more than 10% mismatch will not hybridize. Conditions of "very high stringency" are those under which sequences with more than 6% mismatch will not hybridize.

***In vitro* amplification:** Techniques that increases the number of copies of a nucleic acid molecule in a sample or specimen. An example of amplification is the polymerase chain reaction, in which a biological sample collected from a subject is contacted with a pair of oligonucleotide primers, under conditions that allow for the hybridization of the primers to nucleic acid template in the sample. The primers are extended under suitable conditions, dissociated from the template, and then re-annealed, extended, and dissociated to amplify the number of copies of the nucleic acid. The product of *in vitro* amplification may be characterized by electrophoresis, restriction endonuclease cleavage patterns, oligonucleotide hybridization or ligation, and/or nucleic acid sequencing, using standard techniques. Other examples of *in vitro* amplification techniques include strand displacement amplification (see U.S. Patent No. 5,744,311); transcription-free isothermal amplification (see U.S. Patent No. 6,033,881); repair chain reaction amplification (see WO 90/01069); ligase chain reaction amplification (see EP-A-320 308); gap filling ligase chain reaction amplification (see U.S. Patent No. 5,427,930); coupled ligase detection and PCR (see U.S. Patent No. 6,027,889); and NASBA™ RNA transcription-free amplification (see U.S. Patent No. 6,025,134).

Injectable composition: A pharmaceutically acceptable fluid composition including at least one active ingredient. The active ingredient is usually dissolved or suspended in a physiologically acceptable carrier, and the composition can additionally include minor amounts of one or more non-toxic auxiliary substances, such as emulsifying agents, preservatives, and pH buffering agents and the like. Such injectable compositions that are useful for use with the provided nucleotides and proteins are conventional; appropriate formulations are well known in the art.

-13-

Isolated: An "isolated" biological component (such as a nucleic acid molecule, protein or organelle) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, *i.e.*, other chromosomal and extra-chromosomal DNA and RNA, proteins and organelles. Nucleic acids and proteins that have been

5 "isolated" include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

MATER protein: A cytosol-localized protein (SEQ ID NO: 24) of approximately 125 to approximately 135 kDa (estimated molecular weight based on gel mobility) that is essential for

10 female fertility. The acronym stands for Maternal Antigen That Embryos Require. *Mater* is a single-copy gene found on human Chromosome 19. Zygotes that arise from *Mater* null (*Mater*^{-/-}) oocytes do not progress beyond the two-cell stage. Thus, *Mater* represents a novel maternal effect gene that is required for embryonic survival and early development in mammals.

Proteins can be identified as MATER proteins by comparing their activity and other physical

15 characteristics to a prototypical MATER protein, such as the human or murine MATER protein. MATER protein biological activity can be described in terms of the ability of a protein to complement (substantially replace the lost function in) a *Mater* null mutant. The ability of a protein to complement a *Mater* mutant may be readily determined by introducing the gene encoding the protein into a *Mater* mutant animal system (such as the *Mater* null mice described herein) using standard

20 methods. If the encoded protein has MATER protein biological activity, this will be manifested as a proportion of the transgenic progeny animals having a relatively wild-type phenotype for those characteristics linked to the *Mater* mutant (*e.g.*, infertility of *Mater* null females due to failure of *Mater* null oocytes to progress beyond the two-cell stage).

Other MATER protein physical characteristics that can be examined when evaluating a

25 hypothetical MATER protein include the molecular weight of the protein (approximately 125-135 kDa, see Example 1, though this value may vary somewhat from species to species), the subcellular localization of the protein (human MATER is expressed in oocytes (Example 4)), and is predicted to be specifically expressed in the cytoplasm and excluded from the nucleus, as occurs in mice (see Example 1), and the temporal and spatial regulation of the mRNA (*Mater* transcript is produced in the

30 maturing oocyte, see Examples 1 and 3). Antibodies that recognize one MATER protein (*e.g.*, a murine MATER) may recognize a MATER protein from another species (*e.g.*, human MATER) (see Example 4); thus, hypothetical MATER proteins can be further examined and identified based on recognition by anti-MATER antibodies produced against MATER proteins from other species. The identity of the human MATER protein can therefore be confirmed for instance by immunological

35 identification using an antibody raised against the murine MATER protein, or an epitope of that protein.

Nucleotide: "Nucleotide" includes, but is not limited to, a monomer that includes a base linked to a sugar, such as a pyrimidine, purine or synthetic analogs thereof, or a base linked to an

-14-

amino acid, as in a peptide nucleic acid (PNA). A nucleotide is one monomer in a polynucleotide. A nucleotide sequence refers to the sequence of bases in a polynucleotide.

Oligonucleotide: An oligonucleotide is a plurality of joined nucleotides joined by native phosphodiester bonds, between about 6 and about 300 nucleotides in length. An oligonucleotide analog refers to moieties that function similarly to oligonucleotides but have non-naturally occurring portions. For example, oligonucleotide analogs can contain non-naturally occurring portions, such as altered sugar moieties or inter-sugar linkages, such as a phosphorothioate oligodeoxynucleotide. Functional analogs of naturally occurring polynucleotides can bind to RNA or DNA, and include peptide nucleic acid (PNA) molecules.

Particular oligonucleotides and oligonucleotide analogs can include linear sequences up to about 200 nucleotides in length, for example a sequence (such as DNA or RNA) that is at least 6 bases, for example at least 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100 or even 200 bases long, or from about 6 to about 50 bases, for example about 10-25 bases, such as 12, 15, 20, or 23 bases.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

Open reading frame: A series of nucleotide triplets (codons) coding for amino acids without any internal termination codons. These sequences are usually translatable into a peptide.

Ortholog: Two nucleic acid or amino acid sequences are orthologs of each other if they share a common ancestral sequence and diverged when a species carrying that ancestral sequence split into two species. Orthologous sequences are also homologous sequences.

Parenteral: Administered outside of the intestine, *e.g.*, not via the alimentary tract. Generally, parenteral formulations are those that will be administered through any possible mode except ingestion. This term especially refers to injections, whether administered intravenously, intrathecally, intramuscularly, intraperitoneally, or subcutaneously, and various surface applications including intranasal, intradermal, and topical application, for instance.

Peptide Nucleic Acid (PNA): An oligonucleotide analog with a backbone comprised of monomers coupled by amide (peptide) bonds, such as amino acid monomers joined by peptide bonds.

Pharmaceutically acceptable carriers: The pharmaceutically acceptable carriers useful with the compositions provided herein are conventional. Martin, *Remington's Pharmaceutical Sciences*, published by Mack Publishing Co., Easton, PA, 19th Edition, 1995, describes compositions and formulations suitable for pharmaceutical delivery of the nucleotides and proteins herein disclosed.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced

salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (*e.g.*, powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain minor
5 amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Polymorphism: Variant in a sequence of a gene. Polymorphisms can be those variations (nucleotide sequence differences) that, while having a different nucleotide sequence, produce functionally equivalent gene products, such as those variations generally found between individuals,
10 different ethnic groups, geographic locations. The term polymorphism also encompasses variations that produce gene products with altered function, *i.e.*, variants in the gene sequence that lead to gene products that are not functionally equivalent. This term also encompasses variations that produce no gene product, an inactive gene product, or increased gene product. The term polymorphism may be used interchangeably with allele or mutation, unless context clearly dictates otherwise.

15 Polymorphisms can be referred to, for instance, by the nucleotide position at which the variation exists, by the change in amino acid sequence caused by the nucleotide variation, or by a change in some other characteristic of the nucleic acid molecule that is linked to the variation (*e.g.*, an alteration of a secondary structure such as a stem-loop, or an alteration of the binding affinity of the nucleic acid for associated molecules, such as polymerases, RNases, and so forth).

20 **Probes and primers:** Nucleic acid probes and primers can be readily prepared based on the nucleic acid molecules provided as indicators of disease or disease progression. It is also appropriate to generate probes and primers based on fragments or portions of these nucleic acid molecules. Also appropriate are probes and primers specific for the reverse complement of these sequences, as well as probes and primers to 5' or 3' regions.

25 A probe comprises an isolated nucleic acid attached to a detectable label or other reporter molecule. Typical labels include radioactive isotopes, enzyme substrates, co-factors, ligands, chemiluminescent or fluorescent agents, haptens, and enzymes. Methods for labeling and guidance in the choice of labels appropriate for various purposes are discussed, *e.g.*, in Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989) and Ausubel *et al.* (In *Current*
30 *Protocols in Molecular Biology*, John Wiley & Sons, New York, 1998).

Primers are short nucleic acid molecules, for instance DNA oligonucleotides 10 nucleotides or more in length. Longer DNA oligonucleotides may be about 15, 20, 25, 30 or 50 nucleotides or more in length. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then the primer
35 extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, *e.g.*, by the polymerase chain reaction (PCR) or other *in vitro* nucleic-acid amplification methods known in the art.

Methods for preparing and using nucleic acid probes and primers are described, for example, in Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989), Ausubel

63198-1404

-16-

et al. (ed.) (In *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1998), and Innis *et al.* (*PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc., San Diego, CA, 1990). Amplification primer pairs (for instance, for use with polymerase chain reaction amplification) can be derived from a known sequence such as the *Mater* sequences described herein, for example, by using computer programs intended for that purpose such as Primer™ (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, MA).

One of ordinary skill in the art will appreciate that the specificity of a particular probe or primer increases with its length. Thus, for example, a primer comprising 30 consecutive nucleotides of a *MATER* protein encoding nucleotide will anneal to a target sequence, such as another homolog of the designated *MATER* protein, with a higher specificity than a corresponding primer of only 15 nucleotides. Thus, in order to obtain greater specificity, probes and primers can be selected that comprise at least 20, 23, 25, 30, 35, 40, 45, 50 or more consecutive nucleotides of a *MATER* protein-encoding nucleotide sequences.

Also provided are isolated nucleic acid molecules that comprise specified lengths of the disclosed *Mater* nucleotide sequences. Such molecules may comprise at least 10, 15, 20, 23, 25, 30, 35, 40, 45 or 50 or more consecutive nucleotides of these sequences or more, and may be obtained from any region of the disclosed sequences (*e.g.*, a *Mater* nucleic acid may be apportioned into halves or quarters based on sequence length, and isolated nucleic acid molecules may be derived from the first or second halves of the molecules, or any of the four quarters, etc.). A *Mater* cDNA or other encoding sequence also can be divided into smaller regions, *e.g.* about eighths, sixteenths, twentieths, fiftieths and so forth, with similar effect.

Another mode of division is to select the 5' (upstream) and/or 3' (downstream) region associated with a *Mater* gene.

Nucleic acid molecules may be selected that comprise at least 10, 15, 20, 25, 30, 35, 40, 50 or 100 or more consecutive nucleotides of any of these or other portions of a human *Mater* nucleic acid molecule, such as those disclosed herein, and associated flanking regions. Thus, representative nucleic acid molecules might comprise at least 10 consecutive nucleotides of the human cDNA fragments shown in SEQ ID NOs: 1 and 2, or the full human *MATER* cDNA shown in SEQ ID NO: 24.

Protein: A biological molecule expressed by a gene or recombinant or synthetic coding sequence and comprised of amino acids.

Purified: The term "purified" does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified protein preparation is one in which the protein referred to is more pure than the protein in its natural environment within a cell or within a production reaction chamber (as appropriate).

Recombinant: A recombinant nucleic acid is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination can be accomplished by chemical synthesis or,

more commonly, by the artificial manipulation of isolated segments of nucleic acids, *e.g.*, by genetic engineering techniques.

Sequence identity: The similarity between two nucleic acid sequences, or two amino acid sequences, is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are. Homologs or orthologs of human MATER protein, and the corresponding cDNA or gene sequence, will possess a relatively high degree of sequence identity when aligned using standard methods. This homology will be more significant when the orthologous proteins or genes or cDNAs are derived from species that are more closely related (*e.g.*, human and chimpanzee sequences), compared to species more distantly related (*e.g.*, human and *C. elegans* sequences).

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith & Waterman *Adv. Appl. Math.* 2: 482, 1981; Needleman & Wunsch *J. Mol. Biol.* 48: 443, 1970; Pearson & Lipman *Proc. Natl. Acad. Sci. USA* 85: 2444, 1988; Higgins & Sharp *Gene*, 73: 237-244, 1988; Higgins & Sharp *CABIOS* 5: 151-153, 1989; Corpet *et al. Nuc. Acids Res.* 16, 10881-90, 1988; Huang *et al. Computer Appls. in the Biosciences* 8, 155-65, 1992; and Pearson *et al. Meth. Mol. Bio.* 24, 307-31, 1994. Altschul *et al. (J. Mol. Biol.* 215:403-410, 1990), presents a detailed consideration of sequence alignment methods and homology calculations.

The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul *et al. J. Mol. Biol.* 215:403-410, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, MD) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. By way of example, for comparisons of amino acid sequences of greater than about 30 amino acids, the Blast 2 sequences function is employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per residue gap cost of 1). When aligning short peptides (fewer than around 30 amino acids), the alignment should be performed using the Blast 2 sequences function, employing the PAM30 matrix set to default parameters (open gap 9, extension gap 1 penalties).

An alternative indication that two nucleic acid molecules are closely related is that the two molecules hybridize to each other under stringent conditions. Stringent conditions are sequence-dependent and are different under different environmental parameters. Generally, stringent conditions are selected to be about 5° C to 20° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence remains hybridized to a perfectly matched probe or complementary strand. Conditions for nucleic acid hybridization and calculation of stringencies can be found in Sambrook *et al. (In Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989) and Tijssen (*Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes* Part I, Chapter 2, Elsevier, New York, 1993). Nucleic acid molecules that hybridize under stringent conditions to a human MATER protein-encoding sequence

-18-

will typically hybridize to a probe based on either an entire human MATER protein-encoding sequence or selected portions of the encoding sequence under wash conditions of 2x SSC at 50° C.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences, due to the degeneracy of the genetic code. It is understood that changes in nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid molecules that all encode substantially the same protein.

Specific binding agent: An agent that binds substantially only to a defined target. Thus a protein-specific binding agent binds substantially only the specified protein. By way of example, as used herein, the term "MATER-protein specific binding agent" includes anti-MATER protein antibodies (and functional fragments thereof) and other agents (such as soluble receptors) that bind substantially only to the MATER protein.

Anti-MATER protein antibodies may be produced using standard procedures described in a number of texts, including Harlow and Lane (*Antibodies, A Laboratory Manual*, CSHL, New York, 1988). The determination that a particular agent binds substantially only to the specified protein may readily be made by using or adapting routine procedures. One suitable *in vitro* assay makes use of the Western blotting procedure (described in many standard texts, including Harlow and Lane (*Antibodies, A Laboratory Manual*, CSHL, New York, 1988)). Western blotting may be used to determine that a given protein binding agent, such as an anti- MATER protein monoclonal antibody, binds substantially only to the MATER protein.

Shorter fragments of antibodies can also serve as specific binding agents. For instance, Fabs, Fvs, and single-chain Fvs (SCFvs) that bind to a specified protein would be specific binding agents. These antibody fragments are defined as follows: (1) Fab, the fragment which contains a monovalent antigen-binding fragment of an antibody molecule produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain; (2) Fab', the fragment of an antibody molecule obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule; (3) (Fab')₂, the fragment of the antibody obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; (4) F(ab')₂, a dimer of two Fab' fragments held together by two disulfide bonds; (5) Fv, a genetically engineered fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and (6) single chain antibody ("SCA"), a genetically engineered molecule containing the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule. Methods of making these fragments are routine.

Subject: Living multi-cellular vertebrate organisms, a category that includes both human and non-human mammals.

Target sequence: "Target sequence" is a portion of ssDNA, dsDNA or RNA that, upon hybridization to a therapeutically effective oligonucleotide or oligonucleotide analog, results in the inhibition of expression. For example, hybridization of therapeutically effectively oligonucleotide to

63198-1404

-19-

a *Mater* target sequence results in inhibition of MATER expression. Either an antisense or a sense molecule can be used to target a portion of dsDNA, since both will interfere with the expression of that portion of the dsDNA. The antisense molecule can bind to the plus strand, and the sense molecule can bind to the minus strand. Thus, target sequences can be ssDNA, dsDNA, and RNA.

5 **Transformed:** A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

10 **Vector:** A nucleic acid molecule as introduced into a host cell, thereby producing a transformed host cell. A vector may include nucleic acid sequences that permit it to replicate in a host cell, such as an origin of replication. A vector may also include one or more selectable marker genes and other genetic elements known in the art.

15 Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. In case of conflict, the present specification, including

20 explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

25 III. *Human MATER Protein and Nucleic Acid Sequences*

Embodiments provide MATER proteins and *Mater* nucleic acid molecules, including cDNA sequences. The prototypical *Mater* sequences are the murine and human sequences, and the use of these sequences to produce transgenic animals having increased or decreased levels of MATER protein is provided, as are diagnostic methods to detect defects or alterations in *Mater* expression or

30 MATER protein production.

The full-length cDNA for human *Mater* (SEQ ID NO: 23) is determined to be 3900 base pairs long (which is somewhat longer the mouse *Mater* cDNA, GenBank Accession # NM_011860.1, SEQ ID NO: 5) and its ORF encodes a protein of 1200 amino acids, having a predicted molecular weight of approximately 135.2 kDa and a predicted pI of about 6.08. The human *Mater* cDNA

35 comprises the sequences shown in SEQ ID NO: 1 and SEQ ID NO: 3, which are thought to be aligned approximately with the seventh and eleventh through fifteenth exons of the mouse *Mater* sequence, respectively. The human *Mater* mRNA is expressed in oocytes, as shown by *in situ* hybridization experiments (Example 3).

-20-

Human MATER protein (SEQ ID NO: 24) has been specifically localized by immunofluorescence to human oocytes (Example 4). The ATG initiation codon of the *Mater* cDNA lies within the context of the ANNATG motif associated with vertebrate initiator codon (Kozak, 1991, *J. Biol. Chem.* 266:19867). The human MATER protein comprises the deduced protein sequences (SEQ ID NOs: 2 and 4) corresponding to the herein described human *Mater* cDNA sequences (SEQ ID NOs: 1 and 3, respectively). These protein fragments show low but significant sequence homology with the OP1/MATER protein from mouse (GenBank # NP_035990, SEQ ID NO: 6). In particular, SEQ ID NO: 2 is 54% identical to residues 257-638 of murine MATER (SEQ ID NO: 6), while SEQ ID NO: 4 is 64% identical to residues 854-1111 of murine MATER. The overall human MATER protein sequence (SEQ ID NO: 24) shares approximately 50% similarity with the murine MATER protein sequence (SEQ ID NO: 6); the relevant sequence alignment is shown in FIG. 9. The depicted comparison between the human and murine MATER proteins was conducted using the BLAST 2 Sequences Program with parameter conditions of Expect 10 and Filter closed (NIH-NCBI).

Certain regions of the human *Mater* cDNA have been identified through the sequencing of human HTGSs, though no function or identity had been previously assigned to those sequences. See, for instance, GenBank accession numbers: AC024580 (GI = 7705010, published May 4, 2000); AC012107 (GI = 6088020, published October 20, 1999, and updated as GI = 7329252, published March 28, 2000); and AC023887 (GI = 7631054, published April 21, 2000). These fragmentary and overlapping human sequences are up to 207 nucleotides in length and are scattered throughout the murine *Mater* cDNA sequence. The human ESTs individually share up to 89% identity with the murine *Mater* cDNA over short, discontinuous regions. Oligonucleotides according to the current disclosure may be chosen to avoid these ESTs. Overall, the first 75 nucleotides of human *Mater* (1-75) and the corresponding deduced amino acids (1-25) were determined by comparison to published human genomic DNA sequences, while the remainder of the *Mater* sequence was determined by direct cloning and sequencing of human ovarian cDNAs.

Mater is a single-copy maternal effect gene, the protein product of which is required for early embryonic survival. Although eggs lacking MATER protein can be fertilized, morphologic signs of deterioration are observed as early as the 2-cell stage beyond which the mutant embryos do not progress. The human MATER protein with 1200 amino acids is predicted to have similar molecular domains to those seen in the murine MATER protein (1111 amino acids). As described by Tong *et al.* (*Mamm. Genome* 11:281-287, 2000), the mouse MATER protein contains a five-fold hydrophilic repeat (26-27 amino acid) near its N-terminus, a short leucine-zipper and a fourteen-fold leucine-rich repeat (28-29 amino acid) near its carboxyl terminus (based on a comparison to the murine MATER protein, as described in Tong *et al.*, *Mamm. Genome* 11, 281-287, 2000). The hydrophilic repeat has low homology with a cytoskeletal protein (neurofilament), raising the possibility that this region mediates interactions that anchor MATER in the cytoplasm. The presence of the leucine-rich domain, as well as a short leucine zipper, both motifs known to mediate protein-protein interactions (Kajava, *J. Mol. Biol.* 277, 519-527, 1998; Buchanan & Gay, *Prog. Biophys. Mol.*

-21-

Biol. 65, 1-44, 1996), suggests that MATER may affect embryonic progression through intermediate factor(s), one or more of which binds directly to MATER in the cytoplasm.

Using antisera against a mouse MATER peptide (residues 1093 through 1111 of murine MATER, SEQ ID NO: 6), the inventors have demonstrated that human MATER protein is present in oocytes in human ovary sections (Example 4). Likewise, using mouse-derived nucleotide sense and antisense probes for *in situ* hybridization experiments, it has been demonstrated that human oocytes express *Mater* mRNA (Example 3).

The inventors have further characterized murine MATER as to its function in mammalian embryogenesis. This protein is a "maternal" protein, generated in the oocyte prior to fertilization, and therefore encoded for by a maternal gene. MATER is unusual in that it persists (as measured by protein level) into the late blastocyst stage of embryonic development. Functional MATER is required for female fertility; zygotes that arise from *Mater* null oocytes do not progress beyond the two-cell stage; this is true regardless of what *Mater* genotype male produced the sperm. The protein is cytoplasmic, with definite exclusion from the nucleus.

The invention is illustrated by the following non-limiting Examples.

EXAMPLE 1

Characterization of the Murine MATER protein

This example provides several methods for examining MATER proteins and nucleic acids in a mammalian system; the murine system is used.

Methods:

Isolation of oocytes, eggs and embryos. All experiments were conducted in compliance with the guidelines of the Animal Care and Use Committee under an approved animal study protocol.

Resting, growing and fully grown oocytes were dissected from 1, 10 and 21 day old (d/o) mouse ovaries, respectively, and eggs were isolated from gonadotrophin stimulated female mice (Tong *et al.*, *J. Biol. Chem.* 270, 849-853, 1995; Rankin *et al.*, *Development* 125, 2415-2424, 1998). Embryos were obtained by mating gonadotrophin stimulated females with males and 1-cell zygotes, 2-cell embryos, morulae and blastocysts were flushed from the oviducts at 1, 2, 3 and 4 days, respectively, counting the morning after hCG administration as day 1. Embryos were either incubated in M16 media (37 °C, 5% CO₂), fixed in 1% paraformaldehyde for subsequent confocal microscopy, or frozen at -80 °C for RNA and protein analyses. Ovaries were fixed (10% formalin) and embedded in paraffin prior to sectioning and staining with hematoxylin and eosin (American Histolabs, Gaithersburg, Maryland).

Detection of transcripts and proteins. *Mater*, ZP3, β -actin, cyclophilin and 28S-rRNA transcripts were detected by RNase protection assays as previously described (Tong & Nelson, *Endocrinology* 140, 3720-3726, 1999; Tong *et al.*, *J. Biol. Chem.* 270, 849-853, 1995) using an RPA II kit (Ambion, Austin, TX).

63198-1404

-22-

Rabbits were immunized with a KLH conjugated MATER peptide (amino acids 1093-1111) to obtain a monospecific antisera. After incubating immunoblots with MATER antisera (1:1000, two hours, 20 °C), antibody binding was detected by ECL using a HRP-conjugated goat anti-rabbit antibody (Amersham Pharmacia Biotech, Piscataway, NJ). Fixed oocytes, eggs or embryos were
 5 incubated with MATER antisera (1:8000, 4 °C, 16 hours) and imaged by laser-scanning confocal microscopy (LSM 5; Zeiss, Thornwood, NY) using Cy⁵-conjugated goat anti-rabbit IgG (1:200).

Protein synthesis and immunoblotting. Embryos were incubated in 100 µl of M16 medium containing L-³⁵S-methionine (0.5 Ci/ml, Amersham) at 37 °C for 4 hours. At the end of radiolabeling, 1-cell and 2-cell embryos were harvested at 26 hours and 48 hours post-hCG/mating, respectively.
 10 After washing with 1% BSA in 10 mM Tris-HCl (pH 7.4), 1mM EDTA and 140 mM NaCl, ten embryos were dissolved directly in 10 µl of sample buffer (1) directly or (2) after treatment with 2% Triton[™] X-100/0.3 M KCl to extract the TRC and p35 (Conover *et al.*, *Dev. Biol.* 144, 392-404, 1991). After SDS-PAGE and fluorography as previously reported (Tong *et al.*, *J. Biol. Chem.* 270, 849-853, 1995), radioactive incorporation was determined by a phosphoimager and ImageQuant[™] software
 15 (Molecular Dynamics, Amersham Pharmacia Biotech, Piscataway, NJ).

Production of Mater null mice. Constructs use to produce *Mater* null mice are shown in FIG. 2. To construct a targeting vector in pPNT (Tybulewicz *et al.*, *Cell* 65, 1153-1163, 1991), a 1.5 kbp SacI-EcoRI DNA fragment containing the first two exons of murine *Mater* (Tong *et al.*, *Mamm. Genome* 11, 281-287, 2000) was inserted between the PGK-Neo and PGK-TK cassettes and a 2 kbp
 20 EcoRI-BamHI fragment containing exons 3 and 4 was inserted upstream of the PCK-Neo. After linearization and electroporation into embryonic stem (ES) cells (Redmond *et al.*, *Nat. Genet.* 20, 344-351, 1998), the presence of the mutant allele in chemically selected clones was detected as 3.5 kbp and 4.0 kbp XbaI fragments with 5' and 3' (or Neo) probes, respectively. Both the 5' and 3' probes detected the normal allele as a 9.4 kbp XbaI fragment. C57BL/6 blastocysts were injected
 25 with 8-12 ES cells derived from five independently selected clones. Two cell lines that gave rise to coat-color chimeric animals were mated with C57BL/6 females and transmitted the *Mater* mutation through the germ line. The absence of *Mater* transcripts in homozygous null females was confirmed by northern blot analysis as described previously (Tong & Nelson, *Endocrinology* 140, 3720-3726, 1999).

30 *Reproductive performance.* Vaginal smears were obtained daily to examine four or more estrus cycles of 6-10 week-old female mice (Rugh, *The Mouse: Its Reproduction and Development* 210 Oxford University Press, Oxford, 1991). Mating behavior was evaluated by the presence of a vaginal plug on the morning after mating with fertile males.

Zygotic gene transcription assays. A 5-10 µl aliquot of 100 mM BrUTP, 140 mM KCl and
 35 2mM Pipes, pH 7.4 was microinjected into the cytoplasm of embryos isolated 28 hours (1-cell embryos) and 48 hours (2-cell embryos) following hCG and mating, essentially as previously described (Bouniol *et al.*, *Exp. Cell Res.* 218, 57-62, 1995). After incubation in M16 medium in 5% CO₂ for one hour at 37 °C, the injected embryos were fixed in 1% paraformaldehyde.

-23-

BrUTP incorporation into RNA was assayed with a mouse anti-BrdU monoclonal antibody (Sigma, 1:1000, 16 hours, 4 °C). After washing in PBS containing 3% BSA, a FITC-conjugated goat anti-mouse IgG antibody was used to image the embryos by confocal microscopy; fluorescence was recorded in arbitrary units. This assay primarily reflects *de novo* RNA polymerase II activity, because poor penetration of the monoclonal antibody into nucleoli prevents measurement of BrUTP incorporation into ribosomal RNA (Wansink, *et al.*, *J. Cell Biol.* 122, 283-293, 1993).

Results

Mater transcript and protein expression

Mater mRNA was first detected as oocytes entered into their growth phase (30-50 µm) (FIG. 1A). *Mater* transcripts were most abundant in fully-grown oocytes (75-85 µm) and, like many oocyte transcripts (Bachvarova, in *Developmental Biology: Oogenesis* 1, 453-524, Plenum Press, New York, 1985; Epifano *et al.*, *Development* 121, 1947-1956, 1995), were degraded during meiotic maturation and subsequent ovulation. *Mater* mRNA was not detected in the early embryo (FIG. 1A), although its presence in EST databases derived from early embryos suggest that some transcripts may have escaped destruction.

The initial accumulation of MATER protein during oogenesis was similar to that observed for *Mater* transcripts (compare FIG. 1A with FIG. 1B). However, unlike the virtual disappearance of the transcripts, the MATER protein persisted during preimplantation development until the late blastocyst stage (FIG. 1B). Using confocal microscopy, MATER protein was located in the cytoplasm of growing oocytes and noticeably excluded from the nucleus. MATER was more concentrated in the cortical region of oocytes and early embryos (one- and two-cell stages), and this peripheral pattern persisted later in preimplantation development with a higher concentration of MATER in the outer trophectodermal cells compared to the inner cell mass of blastocysts (FIG. 1C).

Mater null animals

To determine the function of MATER protein, *Mater* null mouse lines were generated, in which homozygous females did not express either *Mater* transcripts or protein. *Mater* null mice were born in the expected Mendelian ratios with equal sex distribution. No phenotypic abnormalities were observed from birth through adulthood.

Ovaries and uteri from *Mater* null mice appeared morphologically normal (FIG. 6). *Mater* null ovaries had a normal complement of primordial follicles and all stages of follicular development were normally represented; *corpora lutea*, indicating past spontaneous ovulation, were also present (FIG. 5).

Ova lacking *Mater* were fertilized normally *in vivo* (FIG. 1). The number and morphology of zygotes and 2-cell embryos from the *Mater* null females were similar to those from normal females (FIG. 5). However, by 3 or 4 days after mating the embryos from *Mater* null females still remained at the 2-cell stage or had begun to degenerate (compare FIG. 3D and 3G with 3E and 3H). Thus, fertilization is normal in *Mater* null females and the resulting zygotes can progress through the first

cleavage, but subsequent development is arrested at the 2-cell stage. This indicates that an arrest of early embryogenesis accounts for the infertility in *Mater* null females.

Two mouse lines with a *Mater* null mutation (T54 and T85) were generated (FIG. 2A and 2B) and neither *Mater* transcript (FIG. 2C) nor protein (FIG. 2D) were detected in homozygous null females.

Reproductive phenotype of *Mater* null animals

Matings of heterozygous *Mater* null parents produced average sized litters (8.5 ± 1.9) with normal Mendelian and equal sex distribution of the null allele in progeny. Homozygous null animals appeared normal and males were as fertile as normal littermates. However, homozygous null females produced no litters, even after five months of mating (Table 1) which implicated *Mater* as a maternal effect gene.

Table 1: Fertility and Mendelian allele ratio of mice with *Mater* mutation

Mating pairs		Litter sizes	Mendelian ratios		
$\sigma \times \phi$	n*	mean $\pm$ s.e.m.	+/+	+/-	-/-
<i>Mater</i> ^{+/+} x <i>Mater</i> ^{+/+}	11	7.5 ± 1.8	1.00	0	0
<i>Mater</i> ^{+/+} x <i>Mater</i> ^{+/-}	10	7.4 ± 1.9	n.d. [†]	n.d.	n.d.
<i>Mater</i> ^{+/+} x <i>Mater</i> ^{-/-}	9	0	n.a. [‡]	n.a.	n.a.
<i>Mater</i> ^{-/-} x <i>Mater</i> ^{+/+}	8	7.6 ± 2.4	0	1.00	0
<i>Mater</i> ^{-/-} x <i>Mater</i> ^{+/-}	18	8.4 ± 1.9	0	0.47	0.53
<i>Mater</i> ^{+/-} x <i>Mater</i> ^{+/-}	15	8.5 ± 1.9	0.21	0.57	0.22

* n, the number of mating pairs, [†] n.d., not done, [‡] n.a., not applicable.

Sexually mature *Mater* null females exhibited regular 5.67 ± 0.22 days estrus cycles, similar to those observed in normal females (5.40 ± 0.23 days). Mating occurred normally, as demonstrated by the appearance of vaginal plugs in about 70% of either *Mater* null or normal females, after exposure to fertile males. The *Mater* null ovary had a normal complement of primordial follicles, with all stages of follicular development as well as *corpora lutea* indicating past spontaneous ovulations (FIG. 5A and 5B). In addition, the *Mater* null females ovulated normally after stimulation with exogenous gonadotropins. However, homozygous null females produced no litters even after five months of mating with normal males, while the homozygous null males and heterozygous females had normal fertility.

The mean number ($\pm$ s.e.m.) of ovulated eggs recovered from null (27.6 ± 4.5 , n=14) was not statistically different ($p > 0.05$) from that obtained from normal (27.7 ± 4.3 , n=12) females (FIG. 5D). Morphologically, the two groups of eggs were indistinguishable from one another (FIG. 5C).

-25-

Fertilization and embryo development in *Mater* null animals

To evaluate *in vivo* fertilization and early development, embryos were isolated from normal and *Mater* null females at 1, 2, 3 and 4 days after mating with normal males. Fertilization occurred in eggs lacking MATER, as demonstrated by the presence of two pronuclei (arrows, FIG. 3A and 3B).

5 The number and morphology of 1-cell zygotes and 2-cell embryos obtained *in vivo* from the *Mater* null females were similar to those observed from normal females, although the 2-cell mutant embryos appeared less healthy, displaying cytoplasmic granulations (FIG. 3A-3F). Embryos obtained from the *Mater* null female mice 3 or 4 days after mating still remained at the 2-cell stage (or had begun to degenerate), while embryos from normal mice had progressed to the morula or blastocyst stages,
10 respectively

An equal percentage (~70%) of fertilized 1-cell zygotes derived from *Mater* null and normal females progressed *in vitro* to the 2-cell stage. However, after four days in culture, embryos lacking MATER remained arrested at the 2-cell stage (most were degenerating) (FIG. 3K); embryos from the normal females developed to the blastocyst stage in the same time (FIG. 3J). Thus, although
15 fertilization appears normal and zygotes without MATER can progress through the first cleavage, subsequent embryonic development is arrested at the 2-cell stage.

Characterization of transcription in *Mater* null animals

The early arrest phenotype seen in *Mater* null mice is reminiscent of the 2-cell block that
20 occurs after exposure of mouse embryos to α -amanitin (Schultz, *Bioessays* 15:531-538, 1993; Flach *et al.*, *EMBO J.* 1, 681-686, 1982), a mushroom toxin that binds to RNA polymerase II and prevents elongation of transcription beyond dinucleotides (Vaisius & Wieland, *Biochemistry* 21, 3097-3101, 1982). Normal embryonic transcription is first detected late in the 1-cell zygote at a level that is ~20% that of the 2-cell embryo (Aoki *et al.*, *Dev. Biol.* 181, 296-307, 1997), but these nascent
25 transcripts are poorly, if at all, translated into protein (Nothias *et al.*, *EMBO J.* 15, 5715-5725, 1996).

To assess *de novo* RNA polymerase II activity of the early *Mater* null embryo, bromo-UTP (BrUTP) incorporation was assayed with a monoclonal antibody specific to BrUTP using laser-scanning confocal microscopy, essentially as described previously (Bouniol *et al.*, *Exp. Cell Res.* 218, 57-62, 1995) (FIG. 4A-4G). Morphologically there was a dramatic difference in the amount of
30 BrUTP detected in the nuclei of normal embryos 28 hours (FIG. 4B) and 48 hours (FIG. 4D) after mating compared to those lacking MATER (FIG. 4A and 4C). BrUTP incorporation in embryos without MATER was marginally greater than that of uninjected controls (FIG. 4E and 4F) and *de novo* transcription was less than 5% that observed in normal 2-cell embryos (FIG. 4G).

Characterization of TRC in *Mater* null animals

During meiotic maturation and ovulation, more than 50% of maternal RNA is lost in normal mice (Bachvarova, in *Developmental Biology: Oogenesis* 1, 453-524, Plenum Press, New York, 1985). This degradative process also occurs in females lacking MATER with loss of ribosomal RNA

-26-

(FIG. 4H) and at least some mRNAs (β -actin, ZP3, cyclophilin, and GAPDH). Normally, the subsequent activation of the embryonic genome at the 2-cell stage is preceded by the transient transcription and translation of a subset of gene products, some of which form the distinctive Transcription Related Complex (TRC) (Bolton *et al.*, *J. Embryol. Exp. Morphol.* 79:139-163, 1984; Nothias *et al.*, *EMBO J.* 15:5715-5725, 1996; Conover *et al.*, *Dev. Biol.* 144:392-404, 1991). This complex, first observed in the early 2-cell embryo, is not formed in the presence of α -amanitin.

To assay for the TRC, 1- and 2-cell embryos derived from *Mater* null females mated with normal males were incubated with ^{35} S-methionine. No global differences in the amount of ^{35} S-methionine incorporation or in protein profiles were noted in comparing normal 1-cell zygotes and those lacking MATER (FIG. 4I, left panel). Although TRC complexes were detected in embryos without MATER at the 2-cell stage (FIG. 4I, middle and right panels), they appeared somewhat less abundant (~60% of normal), which could reflect abnormal protein synthesis in the embryos lacking MATER. This indicates that MATER is not absolutely critical for initiation of all transcription-translation machinery in early embryos.

Although this striking decrease in transcription could reflect generalized morbidity of the embryo, a similar decrease in 1-cell zygotes, which appear quite healthy, as well as absence of an equally dramatic decrease in de novo protein synthesis (FIG. 4I) suggest that the depressed levels of transcription are due either directly, or indirectly to the absence of MATER.

Proposed function(s) of mammalian MATER

The homology of the leucine-rich repeat domain of mammalian MATER protein with porcine ribonuclease inhibitor implies that MATER may act as an inhibitor of cytoplasmic RNase. If this is true, RNA in embryos lacking MATER might be subject to extensive degradation, perhaps an accentuation of the normal turnover of RNA that occurs during the maternal-to-zygotic transition in the early embryo. The degradation of RNAs, including those that encode proteins required for transcription, could account for the phenotype observed in *Mater* null embryos. This seems unlikely; however, because in the absence of *Mater* mutant embryos incorporate ^{35} S-methionine into proteins and no major difference in protein profiles are observed in comparison with normal embryos. These results indicate that sufficient mRNA, tRNA and ribosomal RNA are present in the mutant embryos to support protein biosynthesis, although it does not preclude the degradation of a targeted subset of RNA required for early embryonic survival.

Transcription-dependent protein synthesis occurs in two phases in the 2-cell embryo, an early minor activation (2-4 hours after the first mitotic division) that is characterized by appearance of the TRC proteins, and a later major activation that occurs in G₂ (8-10 hours after the first mitotic division). The later activation initiates developmental programs required for progression beyond the 2-cell stage (Schultz, *Bioessays* 15, 531-538, 1993). The observed decrease in embryonic transcription in embryos lacking MATER could result from deterioration of transcription machinery, decreased access to genomic DNA or abnormalities in the chromatin template within the nucleus.

-27-

However, the presence of the TRC proteins in *Mater* null animals indicates that MATER is not critical for the early phase of transcription-dependent protein synthesis in 2-cell embryos. It further suggests that the transcription machinery, and its ability to access nuclear DNA, is largely intact in *Mater* null embryos (as it appears to be in *Mater* null oocytes, which are also transcriptionally active).

5 The TRC proteins resulting from the early burst of transcription are associated with the nucleus (Schultz, *Bioessays* 15, 531-538, 1993). These proteins could relieve the late transcriptional repression required for progression beyond the 2-cell stage via mechanisms involving MATER. Additionally, it has been reported that mouse oocytes and 1-cell embryos lack coactivator(s) required for enhancer-dependent transcription in 2-cell embryos (Lawinger *et al.*, *J. Biol. Chem.* 274, 8002-10 8011, 1999). The nature of these coactivators is unknown; they may arise by modification of pre-existing maternal proteins or from early transcription-dependent protein synthesis in the 2-cell embryo. Such process may involve cytoplasmic MATER. Alternatively, MATER might be involved in one or more interactions that are required for cell cycle progression beyond the 2-cell stage. Identification and characterization of proteins that interact with MATER will provide insights into the 15 role of this maternal protein in promoting embryonic survival and early development.

Mater is the first maternal effect gene demonstrated to play a critical role in early mammalian development and the observed sterile phenotype raises the possibility that the absence of a similar molecule could be a cause of human infertility.

20

EXAMPLE 2

Identification of the Human *Mater* cDNA

The human *Mater* cDNA was identified based on its homology to the mouse *Mater* cDNA sequence by searching the collection of human High Throughput Genomic (HTG) Sequences, maintained by NCBI and the National Library of Medicine. Using the isolated and discontinuous 25 nucleotide fragments identified in this search, the inventors devised oligonucleotide primers pairs (for instance, SEQ ID NOs: 7 and 8, 19 and 20, and 21 and 22) that were used to amplify *Mater* sequence from purified human ovarian DNA or mRNA/cDNA, and to isolate a partial cDNA clone from a human cDNA library, using standard techniques.

The sequence of two long fragments of the human *Mater* cDNA is shown in SEQ ID NO: 1 30 and 3. Together, these two long cDNA fragments constitute 2213 nucleotides of the complete human *Mater* cDNA.

The deduced human MATER protein portions encoded cDNA fragment 1 (SEQ ID NO: 1) and fragment 2 (SEQ ID NO: 3) are shown in SEQ ID NOs: 2 and 4, respectively.

Using these sequences, the remainder of the human *Mater* cDNA has been identified; the 35 sequence of the full length cDNA is shown in SEQ ID NO: 23. Overall, the first 75 nucleotides of human *Mater* (1-75) and the corresponding deduced amino acids (1-25) were determined by comparison to published human genomic DNA sequences, while the remainder of the *Mater* sequence was determined by direct cloning and sequencing of human ovarian cDNAs. The complete sequence of the human MATER protein is shown in SEQ ID NO: 24.

-28-

EXAMPLE 3:**Localization of Human *Mater* transcript**

This example provides one method for detecting expression of the human *Mater* transcript.

5 To produce *in situ* hybridization probes, human genomic DNA was amplified using the following primers:

5'-primer: 5'- TTTCACATGAACATCCTTCTCC-3' (SEQ ID NO: 7);

3'-primer: 5'-AGTGCTGGAGGCAGAAGGAAG-3' (SEQ ID NO: 8).

10

The resultant amplified nucleic acid molecule product size was 496 bp, and is predicted to fall within exon 7 of the human *Mater* gene, based on comparison to the murine *Mater* exon-intron map. This fragment was subcloned into pBluscript vector as a DNA template. Both ³⁵S-UTP-labeled sense and antisense sequences were synthesized by *in vitro* transcription using T3 and T7 RNA polymerases.

15

In situ hybridization was carried out essentially as described previously (Tong & Nelson, *Endocrinology* 140:3720-3726, 1999). Briefly, the probes were prepared by alkaline hydrolysis and hybridized with frozen human ovarian sections at 60 °C for 24 hours. After dipping in Kodak NTB-2 emulsion, the slides were exposed to film for 2-3 days and the autoradiographs developed.

20

The frozen human ovarian sections were hybridized with the radio-labeled sense (FIG. 7A and 7C) and antisense (B and D) probes. The slides were stained with hematoxylin and eosin. For each probe, bright-field (A and B) and dark-field (C and D) images are displayed. *Mater* transcript was localized to oocytes.

EXAMPLE 4:

25

Localization of Human MATER Protein

This example provides one method for examining the localization of human MATER protein.

30

In situ localization was carried out essentially as described for murine samples (see, Tong and Nelson, *Endocrinology* 140:3720-3726, 1999). Briefly, rabbit polyclonal antibody to a murine MATER C-terminal peptide (residues 1093 through 1111 of SEQ ID NO: 6) was prepared as described in Example 9C. Frozen human ovarian sections were incubated with this antiserum (1:200), and FITC-conjugated goat anti-rabbit IgG antiserum was used as the secondary antibody to detect human MATER protein in the oocyte (FIG. 8A). FIG. 8B shows the phase contrast images corresponding to the samples in FIG. 8A.

35

EXAMPLE 5:**Methods of Making Human *Mater* cDNA**

The original means by which the *Mater* cDNA was identified and obtained is described above. With the provision of the sequence of the large portions of the MATER protein (SEQ ID

-29-

NOs: 2 and 4) and encoding nucleic acid molecules (SEQ ID NOs: 1 and 3), nucleotide amplification (such as polymerase chain reaction (PCR)) now may be utilized in a simple method for producing the *Mater* cDNA.

Total RNA is extracted from human cells by any one of a variety of methods well known to those of ordinary skill in the art. Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989) and Ausubel *et al.* (In *Current Protocols in Molecular Biology*, Greene Publ. Assoc. and Wiley-Intersciences, 1992) provide descriptions of methods for RNA isolation. Because *MATER* is expressed in oocytes, human cell lines derived from oocytes or ovaries can be used as a source of such RNA. The extracted RNA is then used as a template for performing reverse transcription-polymerase chain reaction (RT-PCR) amplification of cDNA. Methods and conditions for RT-PCR are described in Kawasaki *et al.*, (In *PCR Protocols, A Guide to Methods and Applications*, Innis *et al.* (eds.), 21-27, Academic Press, Inc., San Diego, California, 1990).

The selection of amplification primers will be made according to the portion(s) of the cDNA that is to be amplified. Primers may be chosen to amplify a segment of a cDNA or the entire cDNA molecule. Variations in amplification conditions may be required to accommodate primers and amplicons of differing lengths and composition; such considerations are well known in the art and are discussed for instance in Innis *et al.* (*PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc., San Diego, CA, 1990). By way of example, the majority of the coding portion of the human *MATER* cDNA molecule (approximately 2.8 kb) may be amplified using the following combination of primers:

5'-primer (part of predicted exon 7):

5'-CAGGAATTGGGAAATCGGCTCTCTAG-3' (SEQ ID NO.: 9);

3'-primer (end of predicted exon 15):

5'-CCAAATGCTTTGTGTTTATTTAATTCC-3' (SEQ ID NO.: 10).

One of ordinary skill can use the full length human *Mater* cDNA sequence provided herein (SEQ ID NO: 23) to devise primers that could be used to amplify the entire cDNA using known techniques.

The following set of four primers can be used to amplify ~950 bp at the 5'-end of human *Mater* cDNA:

The 5'-primers are the universal adapter primers (ADP) for 5'-RACE-PCR as follows (OriGene Technologies, Inc, Rockville, MD):

ADP1: 5'-CGGAATTCGTCACCTCAGCG-3' (SEQ ID NO.: 11)

ADP2: 5'-AGCGCGTGAATCAGATCG-3' (SEQ ID NO.: 12)

The 3'-primers are gene-specific primers (GSP) for human *MATER* cDNA within exon 7, as follows:

GSP1: 5'-ATTCCCTGGTAGAGTCCACCTTGC-3' (SEQ ID NO.: 13)

GSP2: 5'-ACAGCACGATCCTTCTGGCTAGAG-3' (SEQ ID NO.: 14).

-30-

GSP1 is used as a primer for reverse transcription of human ovarian RNA to synthesize cDNA. The reverse transcribed cDNAs are then adapted at the 5'-end by ADP1 primer, then amplified for the first round of PCR using ADP1 and GSP1 primers. The second round of PCR amplification is followed using PCR products from the first round as template, and ADP2 and GSP2 as primers.

5

Fragment 1 of the human *Mater* cDNA (SEQ ID NO: 1) may be amplified using the following combination of primers:

5'-primer: 5'-CAAGCTCCGGTGACGGAGATCAT-3' (SEQ ID NO.: 15);

3'-primer: 5'-AGCTGGAGGCAGAAGGAAGATG-3' (SEQ ID NO.: 16).

10

The 3'-terminal region of the human *Mater* cDNA molecule (also referred to herein as human *Mater* cDNA fragment 2, SEQ ID NO: 3) may be amplified using the following combination of primers:

5'-primer: 5'-TCTGGCCTCAGCCCTCGTCAGCTTGAC-3' (SEQ ID NO.: 17);

15

3'-primer: 5'-CCAAATGCTTTGTGTTTATTTAATTCC-3' (SEQ ID NO.: 18).

These primers are illustrative only; one skilled in the art will appreciate that many different primers may be derived from the provided cDNA sequence in order to amplify particular the indicated regions *Mater* cDNA, as well as to complete the sequence of the human *Mater* cDNA.

20

Re-sequencing of PCR products obtained by these amplification procedures is recommended; this will facilitate confirmation of the amplified sequence and will also provide information on natural variation on this sequence in different populations or species.

Oligonucleotides derived from the provided *Mater* sequences provided may be used in such sequencing methods.

25

Orthologs of human *Mater* can be cloned in a similar manner, where the starting material consists of cells taken from a non-human species. Orthologs will generally share at least 80% sequence homology with the disclosed human *Mater* cDNA. Where the non-human species is more closely related to humans, the sequence homology will in general be greater. Closely related orthologous *Mater* molecules may share at least 82%, at least 85%, at least 90%, at least 91%, at least 93%, at least 95%, or at least 98% sequence homology with the disclosed human sequences.

30

Oligonucleotides derived from the human *Mater* cDNA sequence (SEQ ID NO: 23), or fragments thereof (such as SEQ ID NOs: 1 and 3), are encompassed within the scope of the present disclosure. Preferably, such oligonucleotide primers will comprise a sequence of at least 23 consecutive nucleotides of the *Mater* nucleic acid sequence. To enhance amplification specificity, oligonucleotide primers comprising at least 25, 30, 35, 40, 45 or 50 consecutive nucleotides of these sequences may also be used. These primers for instance may be obtained from any region of the disclosed sequences. By way of example, the human *Mater* cDNA, ORF and gene sequences may be apportioned into about halves or quarters based on sequence length, and the isolated nucleic acid molecules (*e.g.*, oligonucleotides) may be derived from the first or second halves of the molecules, or

35

-31-

any of the four quarters. The murine *Mater* cDNA, shown in SEQ ID NO: 5, can be used to illustrate this. The human *Mater* cDNA is 3447 nucleotides in length and so may be hypothetically divided into about halves (nucleotides 1-1723 and 1724-3447) or about quarters (nucleotides 1-862, 863-1723, 1724-2586 and 2587-3447). The human cDNA can be likewise apportioned, or can be described in terms of the fragments presented herein (e.g., fragment 1, corresponding to around about nucleotide 1700 through around about 1950 of murine *Mater*; and fragment 2 of the human *Mater* cDNA, corresponding to the 3'-terminal region of murine *Mater* cDNA, around about nucleotide 2500 through the polyA tail of the murine *Mater* cDNA).

Nucleic acid molecules may be selected that comprise at least 15, 20, 23, 25, 30, 35, 40, 50 or 100 consecutive nucleotides of any of these or other portions of the human *Mater* cDNA. Thus, representative nucleic acid molecules might comprise at least 15 consecutive nucleotides of the human *Mater* cDNA (SEQ ID NO: 23), or fragment 1 or fragment 2 of the disclosed human *Mater* coding sequence (SEQ ID NOs: 1 and 3, respectively).

EXAMPLE 6

Cloning of the *Mater* Genomic Sequence (or Gene)

The *Mater* cDNA sequence and fragments described above does not contain introns, upstream transcriptional promoter or regulatory regions or downstream transcriptional regulatory regions of the *Mater* gene. It is possible that some mutations in the *Mater* gene that may lead to defects in embryo development, infertility, or reduced fertility are not included in the cDNA but rather are located in other regions of the *Mater* gene. Mutations located outside of the open reading frame that encodes the MATER protein are not likely to affect the functional activity of the protein, but rather are likely to result in altered levels of the protein in the cell. For example, mutations in the promoter region of the *Mater* gene may prevent transcription of the gene and therefore lead to the complete absence of the MATER protein in the cell.

Additionally, mutations within intron sequences in the genomic gene may also prevent expression of the MATER protein. Following transcription of a gene containing introns, the intron sequences are removed from the RNA molecule in a process termed splicing prior to translation of the RNA molecule which results in production of the encoded protein. When the RNA molecule is spliced to remove the introns, the cellular enzymes that perform the splicing function recognize sequences around the intron/exon border and in this manner recognize the appropriate splice sites. If there is a mutation within the sequence of the intron close to the junction of the intron with an exon, the enzymes may not recognize the junction and may fail to remove the intron. If this occurs, the encoded protein will likely be defective. Thus, mutations inside the intron sequences within the *Mater* gene (termed "splice site mutations") may also lead to defects in embryo development. However, knowledge of the exon structure and intronic splice site sequences of the *Mater* gene is required to define the molecular basis of these abnormalities. The provision herein of the *Mater* cDNA sequence enables the cloning of the entire *Mater* gene (including the promoter and other regulatory regions and the intron sequences) and the determination of its nucleotide sequence. With

-32-

this information in hand, diagnosis of a genetic predisposition to fertility defects based on DNA analysis will comprehend all possible mutagenic events at the *Mater* locus.

The *Mater* gene may be isolated by one or more routine procedures, including direct sequencing of one or more BAC or PAC clones that contain the *Mater* sequence.

5 With the sequences of human *Mater* cDNA and gene in hand, primers derived from these sequences may be used in diagnostic tests (described below) to determine the presence of mutations in any part of the genomic *Mater* gene of a patient. Such primers will be oligonucleotides comprising a fragment of sequence from the *Mater* gene (intron sequence, exon sequence or a sequence spanning an intron-exon boundary) and may include at least 10 consecutive nucleotides of the *Mater* cDNA or
10 gene. It will be appreciated that greater specificity may be achieved by using primers of greater lengths. Thus, in order to obtain enhanced specificity, the primers used may comprise 15, 17, 20, 23, 25, 30, 40 or even 50 consecutive nucleotides of the *Mater* cDNA or gene. Furthermore, with the provision of the *Mater* intron sequence information the analysis of a large and as yet untapped source of patient material for mutations will now be possible using methods such as chemical cleavage of
15 mismatches (Cotton *et al.*, *Proc. Natl. Acad. Sci. USA* 85:4397-4401, 1985; Montandon *et al.*, *Nucleic Acids Res.* 9:3347-3358, 1989) and single-strand conformational polymorphism analysis (Orita *et al.*, *Genomics* 5:874-879, 1989).

Additional experiments may be performed to identify and characterize regulatory elements flanking the *Mater* gene. These regulatory elements may be characterized by standard techniques
20 including deletion analyses wherein successive nucleotides of a putative regulatory region are removed and the effect of the deletions are studied by either transient or long-term expression analyses experiments. The identification and characterization of regulatory elements flanking the genomic *Mater* gene may be made by functional experimentation (deletion analyses, etc.) in mammalian cells by either transient or long-term expression analyses.

25 It will be apparent to one skilled in the art that either the genomic clone or the cDNA or sequences derived from these clones may be utilized in applications, including but not limited to, studies of the expression of the *Mater* gene, studies of the function of the MATER protein, the generation of antibodies to the MATER protein diagnosis and therapy of MATER deleted or mutated patients to prevent or treat the defects in embryo development. Descriptions of applications
30 describing the use of *Mater* cDNA, or fragments thereof, are therefore intended to comprehend the use of the genomic *Mater* gene.

It will also be apparent to one skilled in the art that homologs of this gene may now be cloned from other species, such as the rat or a monkey, by standard cloning methods. Such homologs will be useful in the production of animal models of onset and development of autoimmune infertility,
35 and early embryogenesis. In general, such orthologous *Mater* molecules will share at least 70% sequence identity with the human *Mater* nucleic acid disclosed herein; more closely related orthologous sequences will share at least 75%, at least 80%, at least 90%, at least 95%, or at least 98% sequence identity with this sequence.

EXAMPLE 7***Mater* Nucleic Acid and Amino Acid Sequence Variants**

With the provision of human *MATER* protein fragments and corresponding nucleic acid sequences herein, the creation of variants of these sequences is now enabled.

Variant *MATER* proteins include proteins that differ in amino acid sequence from the human *MATER* sequences disclosed but that share at least 65% amino acid sequence homology with the provided human *MATER* protein. Other variants will share at least 70%, at least 75%, at least 80%, at least 90%, at least 95%, or at least 98% amino acid sequence homology. Manipulation of the nucleotide sequence of *Mater* using standard procedures, including for instance site-directed mutagenesis or PCR, can be used to produce such variants. The simplest modifications involve the substitution of one or more amino acids for amino acids having similar biochemical properties. These so-called conservative substitutions are likely to have minimal impact on the activity of the resultant protein. Table 2 shows amino acids that may be substituted for an original amino acid in a protein, and which are regarded as conservative substitutions.

Table 2

	Original Residue	Conservative Substitutions
	Ala	ser
	Arg	lys
20	Asn	gln; his
	Asp	glu
	Cys	ser
	Gln	asn
	Glu	asp
25	Gly	pro
	His	asn; gln
	Ile	leu; val
	Leu	ile; val
	Lys	arg; gln; glu
30	Met	leu; ile
	Phe	met; leu; tyr
	Ser	thr
	Thr	ser
	Trp	tyr
35	Tyr	trp; phe
	Val	ile; leu

More substantial changes in enzymatic function or other protein features may be obtained by selecting amino acid substitutions that are less conservative than those listed in Table 2. Such changes include changing residues that differ more significantly in their effect on maintaining polypeptide backbone structure (*e.g.*, sheet or helical conformation) near the substitution, charge or hydrophobicity of the molecule at the target site, or bulk of a specific side chain. The following substitutions are generally expected to produce the greatest changes in protein properties: (a) a hydrophilic residue (*e.g.*, seryl or threonyl) is substituted for (or by) a hydrophobic residue (*e.g.*, leucyl, isoleucyl, phenylalanyl, valyl or alanyl); (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain (*e.g.*, lysyl, arginyl, or histadyl) is

-34-

substituted for (or by) an electronegative residue (*e.g.*, glutamyl or aspartyl); or (d) a residue having a bulky side chain (*e.g.*, phenylalanine) is substituted for (or by) one lacking a side chain (*e.g.*, glycine).

Variant MATER encoding sequences may be produced by standard DNA mutagenesis techniques, for example, M13 primer mutagenesis. Details of these techniques are provided in

5 Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989), Ch. 15. By the use of such techniques, variants may be created that differ in minor ways from the human MATER sequences disclosed. DNA molecules and nucleotide sequences that are derivatives of those specifically disclosed herein, and which differ from those disclosed by the deletion, addition, or

10 substitution of nucleotides while still encoding a protein that has at least 82% sequence identity with the human MATER encoding sequence disclosed (SEQ ID NOs: 1 and 3), are comprehended by this disclosure. Also comprehended are more closely related nucleic acid molecules that share at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% nucleotide sequence homology with the disclosed *Mater* sequences. In their most simple form, such variants may differ from the disclosed

15 sequences by alteration of the coding region to fit the codon usage bias of the particular organism into which the molecule is to be introduced. The full length human *Mater* cDNA (SEQ ID NO: 23) is also encompassed in the disclosure, and as a molecule that comprises both fragment 1 (SEQ ID NO: 1) and fragment 2 (SEQ ID NO: 3), and encoding the human MATER protein with described physical characteristics and biological properties.

Alternatively, the coding region may be altered by taking advantage of the degeneracy of the

20 genetic code to alter the coding sequence such that, while the nucleotide sequence is substantially altered, it nevertheless encodes a protein having an amino acid sequence substantially similar to the disclosed human MATER protein sequences. For example, because of the degeneracy of the genetic code, four nucleotide codon triplets – (GCT, GCG, GCC and GCA) - code for alanine. The coding sequence of any specific alanine residue within the human MATER protein, therefore, could be

25 changed to any of these alternative codons without affecting the amino acid composition or characteristics of the encoded protein. Based upon the degeneracy of the genetic code, variant DNA molecules may be derived from the cDNA and gene sequences disclosed herein using standard DNA mutagenesis techniques as described above, or by synthesis of DNA sequences. Thus, this disclosure also encompasses nucleic acid sequences that encode a MATER protein, but which vary from the

30 disclosed nucleic acid sequences by virtue of the degeneracy of the genetic code.

Variants of the MATER protein may also be defined in terms of their sequence identity with the prototype human MATER protein. As described above, human MATER proteins share at least 65%, at least 70%, at least 75%, at least 80%, at least 90%, at least 95%, or at least 98% amino acid sequence identity with the human MATER protein (SEQ ID NO: 24) or fragments disclosed herein

35 (such as SEQ ID NOs: 2 and 4). Nucleic acid sequences that encode such proteins/fragments readily may be determined simply by applying the genetic code to the amino acid sequence of an MATER protein or fragment, and such nucleic acid molecules may readily be produced by assembling oligonucleotides corresponding to portions of the sequence.

-35-

Nucleic acid molecules that are derived from the human *Mater* cDNA nucleic acid sequences disclosed include molecules that hybridize under stringent conditions to the disclosed prototypical *Mater* nucleic acid molecules, or fragments thereof. Stringent conditions are hybridization at 65° C in 6 x SSC, 5 x Denhardt's solution, 0.5% SDS and 100 µg/ml sheared salmon testes DNA, followed by 5 15-30 minute sequential washes in 2 x SSC, 0.5% SDS, followed by 1 x SSC, 0.5% SDS and finally 0.2 x SSC, 0.5% SDS, at 65° C.

Low stringency hybridization conditions (to detect less closely related homologs) are performed as described above but at 50°C (both hybridization and wash conditions); however, depending on the strength of the detected signal, the wash steps may be terminated after the first 2 x 10 SSC wash.

Human *Mater* nucleic acid encoding molecules (including the cDNA shown in SEQ ID NO: 23 and fragments shown in SEQ ID NOs: 1 and 3, and nucleic acids comprising these sequences), and orthologs and homologs of these sequences, may be incorporated into transformation or expression vectors.

15

EXAMPLE 8

Expression of MATER Protein

With the provision of human *Mater* cDNA sequence fragments, and methods for determining and cloning the full length human *Mater* cDNA, the expression and purification of the 20 MATER protein by standard laboratory techniques is now enabled. Purified human MATER protein may be used for functional analyses, antibody production, diagnostics, and patient therapy. Furthermore, the DNA sequence of the *Mater* cDNA can be manipulated in studies to understand the expression of the gene and the function of its product. Mutant forms of the human MATER may be isolated based upon information contained herein, and may be studied in order to detect alteration in 25 expression patterns in terms of relative quantities, cellular localization, tissue specificity and functional properties of the encoded mutant MATER protein. Partial or full-length cDNA sequences, which encode for the subject protein, may be ligated into bacterial expression vectors. Methods for expressing large amounts of protein from a cloned gene introduced into *Escherichia coli* (*E. coli*) may be utilized for the purification, localization and functional analysis of proteins. For example, fusion 30 proteins consisting of amino terminal peptides encoded by a portion of the *E. coli lacZ* or *trpE* gene linked to MATER proteins may be used to prepare polyclonal and monoclonal antibodies against these proteins. Thereafter, these antibodies may be used to purify proteins by immunoaffinity chromatography, in diagnostic assays to quantitate the levels of protein and to localize proteins in tissues and individual cells by immunofluorescence. Such antibodies may be specific for epitope 35 tags, which can be added to the expression construct for identification an/or purification purposes.

Intact native protein may also be produced in *E. coli* in large amounts for functional studies. Methods and plasmid vectors for producing fusion proteins and intact native proteins in bacteria are described in Sambrook *et al.* (Sambrook *et al.*, In *Molecular Cloning: A Laboratory Manual*, Ch. 17,

-36-

CSHL, New York, 1989). Such fusion proteins may be made in large amounts, are easy to purify, and can be used to elicit antibody response. Native proteins can be produced in bacteria by placing a strong, regulated promoter and an efficient ribosome binding site upstream of the cloned gene. If low levels of protein are produced, additional steps may be taken to increase protein production; if high levels of protein are produced, purification is relatively easy. Suitable methods are presented in Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989) and are well known in the art. Often, proteins expressed at high levels are found in insoluble inclusion bodies. Methods for extracting proteins from these aggregates are described by Sambrook *et al.* (In *Molecular Cloning: A Laboratory Manual*, Ch. 17, CSHL, New York, 1989). Vector systems suitable for the expression of *lacZ* fusion genes include the pUR series of vectors (Ruther and Muller-Hill, *EMBO J.* 2:1791, 1983), pEX1-3 (Stanley and Luzio, *EMBO J.* 3:1429, 1984) and pMR100 (Gray *et al.*, *Proc. Natl. Acad. Sci. USA* 79:6598, 1982). Vectors suitable for the production of intact native proteins include pKC30 (Shimatake and Rosenberg, *Nature* 292:128, 1981), pKK177-3 (Amann and Brosius, *Gene* 40:183, 1985) and pET-3 (Studier and Moffatt, *J. Mol. Biol.* 189:113, 1986). MATer fusion proteins may be isolated from protein gels, lyophilized, ground into a powder and used as an antigen. The DNA sequence can also be transferred from its existing context to other cloning vehicles, such as other plasmids, bacteriophages, cosmids, animal viruses and yeast artificial chromosomes (YACs) (Burke *et al.*, *Science* 236:806-812, 1987). These vectors may then be introduced into a variety of hosts including somatic cells, and simple or complex organisms, such as bacteria, fungi (Timberlake and Marshall, *Science* 244:1313-1317, 1989), invertebrates, plants, and animals (Pursel *et al.*, *Science* 244:1281-1288, 1989), which cells or organisms are rendered transgenic by the introduction of the heterologous *Mater* cDNA.

For expression in mammalian cells, the cDNA sequence may be ligated to heterologous promoters, such as the simian virus (SV) 40 promoter in the pSV2 vector (Mulligan and Berg, *Proc. Natl. Acad. Sci. USA* 78:2072-2076, 1981), and introduced into cells, such as monkey COS-1 cells (Gluzman, *Cell* 23:175-182, 1981), to achieve transient or long-term expression. The stable integration of the chimeric gene construct may be maintained in mammalian cells by biochemical selection, such as neomycin (Southern and Berg, *J. Mol. Appl. Genet.* 1:327-341, 1982) and mycophenolic acid (Mulligan and Berg, *Proc. Natl. Acad. Sci. USA* 78:2072-2076, 1981).

DNA sequences can be manipulated with standard procedures such as restriction enzyme digestion, fill-in with DNA polymerase, deletion by exonuclease, extension by terminal deoxynucleotide transferase, ligation of synthetic or cloned DNA sequences, site-directed sequence-alteration via single-stranded bacteriophage intermediate or with the use of specific oligonucleotides in combination with nucleic acid amplification.

The cDNA sequence (or portions derived from it) or a mini gene (a cDNA with an intron and its own promoter) may be introduced into eukaryotic expression vectors by conventional techniques. These vectors are designed to permit the transcription of the cDNA in eukaryotic cells by providing regulatory sequences that initiate and enhance the transcription of the cDNA and ensure its proper splicing and polyadenylation. Vectors containing the promoter and enhancer regions of the SV40 or

-37-

long terminal repeat (LTR) of the Rous Sarcoma virus and polyadenylation and splicing signal from SV40 are readily available (Mulligan *et al.*, *Proc. Natl. Acad. Sci. USA* 78:1078-2076, 1981; Gorman *et al.*, *Proc. Natl. Acad. Sci. USA* 78:6777-6781, 1982). The level of expression of the cDNA can be manipulated with this type of vector, either by using promoters that have different activities (for
 5 example, the baculovirus pAC373 can express cDNAs at high levels in *S. frugiperda* cells (Summers and Smith, In *Genetically Altered Viruses and the Environment*, Fields *et al.* (Eds.) 22:319-328, CSHL Press, Cold Spring Harbor, New York, 1985) or by using vectors that contain promoters amenable to modulation, for example, the glucocorticoid-responsive promoter from the mouse mammary tumor virus (Lee *et al.*, *Nature* 294:228, 1982). The expression of the cDNA can be
 10 monitored in the recipient cells 24 to 72 hours after introduction (transient expression).

In addition, some vectors contain selectable markers such as the *gpt* (Mulligan and Berg, *Proc. Natl. Acad. Sci. USA* 78:2072-2076, 1981) or *neo* (Southern and Berg, *J. Mol. Appl. Genet.* 1:327-341, 1982) bacterial genes. These selectable markers permit selection of transfected cells that exhibit stable, long-term expression of the vectors (and therefore the cDNA). The vectors can be
 15 maintained in the cells as episomal, freely replicating entities by using regulatory elements of viruses, such as papilloma (Sarver *et al.*, *Mol. Cell Biol.* 1:486-496, 1981) or Epstein-Barr (Sugden *et al.*, *Mol. Cell Biol.* 5:410-413, 1985). Alternatively, one can also produce cell lines that have integrated the vector into genomic DNA. Both of these types of cell lines produce the gene product on a continuous basis. One can also produce cell lines that have amplified the number of copies of
 20 the vector (and therefore of the cDNA as well) to create cell lines that can produce high levels of the gene product (Alt *et al.*, *J. Biol. Chem.* 253:1357-1370, 1978).

The transfer of DNA into eukaryotic, in particular human or other mammalian cells, is now a conventional technique. Recombinant expression vectors can be introduced into the recipient cells as pure DNA (transfection) by, for example, precipitation with calcium phosphate (Graham and vander
 25 Eb, *Virology* 52:466, 1973) or strontium phosphate (Brash *et al.*, *Mol. Cell Biol.* 7:2013, 1987), electroporation (Neumann *et al.*, *EMBO J* 1:841, 1982), lipofection (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413, 1987), DEAE dextran (McCuthan *et al.*, *J. Natl. Cancer Inst.* 41:351, 1968), microinjection (Mueller *et al.*, *Cell* 15:579, 1978), protoplast fusion (Schafner, *Proc. Natl. Acad. Sci. USA* 77:2163-2167, 1980), or pellet guns (Klein *et al.*, *Nature* 327:70, 1987). Alternatively, the
 30 cDNA, or fragments thereof, can be introduced by infection with virus vectors. Systems are developed that use, for example, retroviruses (Bernstein *et al.*, *Gen. Engr'g* 7:235, 1985), adenoviruses (Ahmad *et al.*, *J. Virol.* 57:267, 1986), or Herpes virus (Spaete *et al.*, *Cell* 30:295, 1982). Techniques of use in packaging long transcripts can be found in Kochanek *et al.* (*Proc. Natl. Acad. Sci. USA* 93:5731-5739, 1996) Parks *et al.* (*Proc. Natl. Acad. Sci. USA* 93:13565-13570, 1996)
 35 and Parks and Graham (*J. Virol.* 71:3293-3298, 1997). MATER encoding sequences can also be delivered to target cells *in vitro* via non-infectious systems, for instance liposomes.

These eukaryotic expression systems can be used for studies of MATER encoding nucleic acids and mutant forms of these molecules, the MATER protein and mutant forms of this protein. Such uses include, for example, the identification of regulatory elements located in the 5' region of the

-38-

Mater gene on genomic clones that can be isolated from human genomic DNA libraries using the information contained herein. The eukaryotic expression systems also may be used to study the function of the normal complete protein, specific portions of the protein, or of naturally occurring or artificially produced mutant proteins.

5 Using the above techniques, expression vectors containing the *Mater* gene sequence or cDNA, or fragments or variants or mutants thereof, can be introduced into human cells, mammalian cells from other species or non-mammalian cells, as desired. The choice of cell is determined by the purpose of the treatment. For example, monkey COS cells (Gluzman, *Cell* 23:175-182, 1981) that produce high levels of the SV40 T antigen and permit the replication of vectors containing the SV40
10 origin of replication may be used. Similarly, Chinese hamster ovary (CHO), mouse NIH 3T3 fibroblasts or human fibroblasts or lymphoblasts (as described herein) may be used.

 Embodiments described herein thus encompass recombinant vectors that comprise all or part of a MATER encoding sequence, such as the *Mater* gene or cDNA or variants thereof, for expression in a suitable host. The *Mater* DNA is operatively linked in the vector to an expression control
15 sequence in the recombinant DNA molecule so that the MATER polypeptide can be expressed. The expression control sequence may be selected from the group consisting of sequences that control the expression of genes of prokaryotic or eukaryotic cells and their viruses and combinations thereof. The expression control sequence may be specifically selected from the group consisting of the *lac* system, the *trp* system, the *tac* system, the *trc* system, major operator and promoter regions of phage
20 lambda, the control region of fd coat protein, the early and late promoters of SV40, promoters derived from polyoma, adenovirus, retrovirus, baculovirus and simian virus, the promoter for 3-phosphoglycerate kinase, the promoters of yeast acid phosphatase, the promoter of the yeast alpha-mating factors and combinations thereof.

 The host cell, which may be transfected with a vector, may be selected from the group
25 consisting of *E. coli*, *Pseudomonas*, *Bacillus subtilis*, *Bacillus stearothermophilus* or other bacilli; other bacteria; yeast; fungi; insect; mouse or other animal; or plant hosts; or human tissue cells.

 It is appreciated that for mutant or variant *Mater* DNA sequences, similar systems are employed to express and produce the mutant product.

30

EXAMPLE 9

Production of an Antibody to MATER Protein

 Monoclonal or polyclonal antibodies may be produced to either the normal MATER protein or mutant forms of this protein. Optimally, antibodies raised against the MATER protein would specifically detect the MATER protein. That is, such antibodies would recognize and bind the
35 MATER protein and would not substantially recognize or bind to other proteins found in human cells. Antibodies the human MATER protein may recognize MATER from other species, such as murine MATER, and vice versa.

 The determination that an antibody specifically detects the MATER protein is made by any one of a number of standard immunoassay methods; for instance, the Western blotting technique

-39-

(Sambrook *et al.*, In *Molecular Cloning: A Laboratory Manual*, CSHL, New York, 1989). To determine that a given antibody preparation (such as one produced in a mouse) specifically detects the MATER protein by Western blotting, total cellular protein is extracted from human cells (for example, lymphocytes) and electrophoresed on a sodium dodecyl sulfate-polyacrylamide gel. The proteins are then transferred to a membrane (for example, nitrocellulose or PVDF) by Western blotting, and the antibody preparation is incubated with the membrane. After washing the membrane to remove non-specifically bound antibodies, the presence of specifically bound antibodies is detected by the use of (by way of example) an anti-mouse antibody conjugated to an enzyme such as alkaline phosphatase. Application of an alkaline phosphatase substrate 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium results in the production of a dense blue compound by immunolocalized alkaline phosphatase. Antibodies that specifically detect the MATER protein will, by this technique, be shown to bind to the MATER protein band (which will be localized at a given position on the gel determined by its molecular weight, which is approximately 125 kDa based on gel-mobility estimation for murine MATER. Non-specific binding of the antibody to other proteins may occur and may be detectable as a weak signal on the Western blot. The non-specific nature of this binding will be recognized by one skilled in the art by the weak signal obtained on the Western blot relative to the strong primary signal arising from the specific antibody-MATER protein binding.

Substantially pure MATER protein suitable for use as an immunogen is isolated from the transfected or transformed cells as described above. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon (Millipore, Bedford, Massachusetts) or similar filter device, to the level of a few micrograms per milliliter. Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

A. Monoclonal Antibody Production by Hybridoma Fusion

Monoclonal antibody to epitopes of the MATER protein identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler and Milstein (*Nature* 256:495-497, 1975) or derivative methods thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein over a period of a few weeks. The mouse is then sacrificed, and the antibody-producing cells of the spleen isolated. The spleen cells are fused with mouse myeloma cells using polyethylene glycol, and the excess un-fused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). Successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate, where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall (*Enzymol.* 70(A):419-439, 1980), and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Harlow and Lane (*Antibodies, A Laboratory Manual*, CSHL, New York, 1988).

-40-

B. Polyclonal Antibody Production by Immunization

Polyclonal antiserum containing antibodies to heterogeneous epitopes of a single protein can be prepared by immunizing suitable animals with the expressed protein (Example 8), which optionally can be modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than others and may require the use of carriers and adjuvant, examples of which are known. Also, host animals vary in response to site of inoculations and dose, with either inadequate or excessive doses of antigen resulting in low titer antisera. A series of small doses (ng level) of antigen administered at multiple intradermal sites appear to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis *et al.* (*J. Clin. Endocrinol. Metab.* 33:988-991, 1971).

Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof begins to fall, as determined semi-quantitatively (for example, by double immunodiffusion in agar against known concentrations of the antigen). See, for example, Ouchterlony *et al.* (In *Handbook of Experimental Immunology*, Wier, D. (ed.) chapter 19. Blackwell, 1973). Plateau concentration of antibody is usually in the range of about 0.1 to 0.2 mg/ml of serum (about 12 μ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher (*Manual of Clinical Immunology*, Ch. 42, 1980).

C. Antibodies Raised against Synthetic Peptides

A third approach to raising antibodies against the MATER protein is to use synthetic peptides synthesized on a commercially available peptide synthesizer based upon the predicted amino acid sequence of the MATER protein.

By way of example only, mouse MATER C-terminal peptide (residues 1093 through 1111 of SEQ ID NO: 6) was conjugated with KLH to immunize the female rabbits (two) every two-weeks. Starting from the third immunization, a small amount (~3 ml) of blood was collected from the immunized rabbits to examine the titer of the anti-peptide antibodies using the peptide as antigen and ELISA method. Immunizations continued until the antibodies reached maximal titers, which occurred in about ten immunizations, and then the rabbits were sacrificed to bleed in preparation for the sera. The resultant preparation was used both to characterize murine MATER (Example 1) and human MATER (Example 4).

D. Antibodies Raised by Injection of MATER Encoding Sequence

Antibodies may be raised against the MATER protein by subcutaneous injection of a recombinant DNA vector that expresses the MATER protein into laboratory animals, such as mice. Delivery of the recombinant vector into the animals may be achieved using a hand-held form of the Biolistic system (Sanford *et al.*, *Particulate Sci. Technol.* 5:27-37, 1987), as described by Tang *et al.* (*Nature* 356:152-154, 1992). Expression vectors suitable for this purpose may include those that

-41-

express the *MATER* encoding sequence under the transcriptional control of either the human β -actin promoter or the cytomegalovirus (CMV) promoter.

Antibody preparations prepared according to these protocols are useful in quantitative
5 immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample.

EXAMPLE 10

10 DNA-Based Diagnosis

The *Mater* sequence information presented herein can be used in the area of genetic testing for predisposition to reduced fertility or infertility, such as autoimmune infertility, owing to defects in *Mater*, such as deletion, duplication or mutation. The gene sequence of the *Mater* gene, including
15 intron-exon boundaries is also useful in such diagnostic methods. Individuals carrying mutations in the *Mater* gene (or a portion thereof), or having duplications or heterozygous or homozygous deletions of the *Mater* gene, may be detected at the DNA level with the use of a variety of techniques. For such a diagnostic procedure, a biological sample of the subject, which biological sample contains either DNA or RNA derived from the subject, is assayed for a mutated, duplicated or deleted *Mater* gene. Suitable biological samples include samples containing genomic DNA or RNA obtained from
20 body cells, such as those present in peripheral blood, urine, saliva, tissue biopsy, surgical specimen, amniocentesis samples and autopsy material. The detection in the biological sample of either a mutant *Mater* gene, a mutant *Mater* RNA, or a duplicated or homozygously or heterozygously deleted *Mater* gene, may be performed by a number of methodologies, examples of which are discussed below.

25 One embodiment of such detection techniques for the identification of unknown mutations is the amplification (e.g., polymerase chain reaction amplification) of reverse transcribed RNA (RT-PCR) isolated from a subject, followed by direct DNA sequence determination of the products. The presence of one or more nucleotide differences between the obtained sequence and the prototypical *Mater* cDNA sequence, and especially, differences in the ORF portion of the nucleotide sequence, are
30 taken as indicative of a potential *Mater* gene mutation.

Alternatively, DNA extracted from a biological sample may be used directly for amplification. Direct amplification from genomic DNA would be appropriate for analysis of the entire *Mater* gene including regulatory sequences located upstream and downstream from the open reading frame, or intron/exon borders. Reviews of direct DNA diagnosis have been presented by
35 Caskey (*Science* 236:1223-1228, 1989) and by Landegren *et al.* (*Science* 242:229-237, 1989).

Other mutation scanning techniques appropriate for detecting unknown within amplicons derived from DNA or cDNA could also be performed. These techniques include direct sequencing (without sequencing), single-strand conformational polymorphism analysis (SSCP) (for instance, see Hongyo *et al.*, *Nucleic Acids Res.* 21:3637-3642, 1993), chemical cleavage (including HOT cleavage)

-42-

(Bateman *et al.*, *Am. J. Med. Genet.* 45:233-240, 1993; reviewed in Ellis *et al.*, *Hum. Mutat.* 11:345-353, 1998), denaturing gradient gel electrophoresis (DGGE), ligation amplification mismatch protection (LAMP), and enzymatic mutation scanning (Taylor and Deeble, *Genet. Anal.* 14:181-186, 1999), followed by direct sequencing of amplicons with putative sequence variations.

5 Further studies of *Mater* genes isolated from female subjects displaying infertility, particularly autoimmune infertility subjects, or their relatives, may reveal particular mutations, genomic amplifications, or deletions, which occur at a high frequency within this population of individuals. In such case, rather than sequencing the entire *Mater* gene, DNA diagnostic methods can be designed to specifically detect the most common, or most closely disease-linked, MATER defects.

10 The detection of specific DNA mutations may be achieved by methods such as hybridization using allele specific oligonucleotides (ASOs) (Wallace *et al.*, CSHL Symp. Quant. Biol. 51:257-261, 1986), direct DNA sequencing (Church and Gilbert, *Proc. Natl. Acad. Sci. USA* 81:1991-1995, 1988), the use of restriction enzymes (Flavell *et al.*, *Cell* 15:25-41, 1978; Geever *et al.*, 1981), discrimination on the basis of electrophoretic mobility in gels with denaturing reagent (Myers and Maniatis, Cold Spring Harbor Symp. Quant. Biol. 51:275-284, 1986), RNase protection (Myers *et al.*, *Science* 230:1242-1246, 1985), chemical cleavage (Cotton *et al.*, *Proc. Natl. Acad. Sci. USA* 85:4397-4401, 1985), and the ligase-mediated detection procedure (Landegren *et al.*, *Science* 241:1077-1080, 1988).

Oligonucleotides specific to normal or mutant sequences are chemically synthesized using commercially available machines. These oligonucleotides are then labeled radioactively with isotopes (such as ³²P) or non-radioactively, with tags such as biotin (Ward and Langer, *Proc. Natl. Acad. Sci. USA* 78:6633-6657, 1981), and hybridized to individual DNA samples immobilized on membranes or other solid supports by dot-blot or transfer from gels after electrophoresis. These specific sequences are visualized by methods such as autoradiography or fluorometric (Landegren *et al.*, *Science* 242:229-237, 1989) or colorimetric reactions (Gebeyehu *et al.*, *Nucleic Acids Res.* 15:4513-4534, 1987). Using an ASO specific for a normal allele, the absence of hybridization would indicate a mutation in the particular region of the gene, or deleted *Mater* gene. In contrast, if an ASO specific for a mutant allele hybridizes to a clinical sample, this would indicate the presence of a mutation in the region defined by the ASO.

Sequence differences between normal and mutant forms of the *Mater* gene may also be revealed by the direct DNA sequencing method of Church and Gilbert (*Proc. Natl. Acad. Sci. USA* 81:1991-1995, 1988). Cloned DNA segments may be used as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with nucleic acid amplification, *e.g.*, PCR (Wrichnik *et al.*, *Nucleic Acids Res.* 15:529-542, 1987; Wong *et al.*, *Nature* 330:384-386, 1987; Stoflet *et al.*, *Science* 239:491-494, 1988). In this approach, a sequencing primer that lies within the amplified sequence is used with double-stranded PCR product or single-stranded template generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabeled nucleotides or by automatic sequencing procedures with fluorescent tags.

-43-

Sequence alterations may occasionally generate fortuitous restriction enzyme recognition sites or may eliminate existing restriction sites. Changes in restriction sites are revealed by the use of appropriate enzyme digestion followed by conventional gel-blot hybridization (Southern, *J. Mol. Biol.* 98:503-517, 1975). DNA fragments carrying the restriction site (either normal or mutant) are
5 detected by their reduction in size or increase in corresponding restriction fragment numbers. Genomic DNA samples may also be amplified by PCR prior to treatment with the appropriate restriction enzyme; fragments of different sizes are then visualized under UV light in the presence of ethidium bromide after gel electrophoresis.

Genetic testing based on DNA sequence differences may be achieved by detection of
10 alteration in electrophoretic mobility of DNA fragments in gels, with or without denaturing reagent. Small sequence deletions and insertions can be visualized by high-resolution gel electrophoresis. For example, a PCR product with small deletions is clearly distinguishable from a normal sequence on an 8% non-denaturing polyacrylamide gel (WO 91/10734; Nagamine *et al.*, *Am. J. Hum. Genet.* 45:337-339, 1989). DNA fragments of different sequence compositions may be distinguished on denaturing
15 formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific "partial-melting" temperatures (Myers *et al.*, *Science* 230:1242-1246, 1985). Alternatively, a method of detecting a mutation comprising a single base substitution or other small change could be based on differential primer length in a PCR. For example, an invariant primer could be used in addition to a primer specific for a mutation. The PCR
20 products of the normal and mutant genes can then be differentially detected in acrylamide gels.

In addition to conventional gel-electrophoresis and blot-hybridization methods, DNA fragments may also be visualized by methods where the individual DNA samples are not immobilized on membranes. The probe and target sequences may be both in solution, or the probe sequence may be immobilized (Saiki *et al.*, *Proc. Nat. Acad. Sci. USA* 86:6230-6234, 1989). A variety of detection
25 methods, such as autoradiography involving radioisotopes, direct detection of radioactive decay (in the presence or absence of scintillant), spectrophotometry involving calorogenic reactions and fluorometry involved fluorogenic reactions, may be used to identify specific individual genotypes.

If multiple mutations are encountered frequently in the *Mater* gene, a system capable of detecting such multiple mutations likely will be desirable. For example, a nucleic acid amplification
30 reaction with multiple, specific oligonucleotide primers and hybridization probes may be used to identify all possible mutations at the same time (Chamberlain *et al.*, *Nucl. Acids Res.* 16:1141-1155, 1988). The procedure may involve immobilized sequence-specific oligonucleotide probes (Saiki *et al.*, *Proc. Nat. Acad. Sci. USA* 86:6230-6234, 1989).

35

EXAMPLE 11

Quantitation of MATER Protein

An alternative method of diagnosing *Mater* gene deletion, amplification, or mutation is to quantitate the level of MATER protein in the cells of a subject. This diagnostic tool would be useful for detecting reduced levels of the MATER protein that result from, for example, mutations in the

-44-

promoter regions of the *Mater* gene or mutations within the coding region of the gene that produce truncated, non-functional or unstable polypeptides, as well as from deletions of the entire *Mater* gene. Alternatively, duplications of the *Mater* gene may be detected as an increase in the expression level of this protein. The determination of reduced or increased MATER protein levels would be an
5 alternative or supplemental approach to the direct determination of *Mater* gene deletion, duplication or mutation status by the methods outlined above.

The availability of antibodies specific to the MATER protein will facilitate the quantitation of cellular MATER protein by one of a number of immunoassay methods, which are well known in the art and are presented in Harlow and Lane (*Antibodies, A Laboratory Manual*, CSHL, New York,
10 1988).

For the purposes of quantitating the MATER protein, a biological sample of the subject, which sample includes cellular proteins, is required. Such a biological sample may be obtained from body cells, such as those present in peripheral blood, urine, saliva, tissue biopsy, amniocentesis samples, surgical specimens and autopsy material. In particular, female reproductive cells (*e.g.*, ova)
15 or embryos are appropriate samples. Quantitation of MATER protein is achieved by immunoassay and compared to levels of the protein found in healthy cells (*e.g.*, cells from a female known not to suffer from decreased fertility). A significant (*e.g.*, 10% or greater, for instance, 20%, 25%, 30%, 50% or more) reduction in the amount of MATER protein in the cells of a subject compared to the amount of MATER protein found in normal human cells would be taken as an indication that the
20 subject may have deletions or mutations in the *Mater* gene locus, whereas a significant (*e.g.*, 10% or greater, for instance, 20%, 25%, 30%, 50% or more) increase would indicate that a duplication or enhancing mutation had occurred.

EXAMPLE 12

25 Detection of Serum Antibody against MATER Protein

The MATER family of proteins was first identified as autoantibodies involved in the pathogenesis of certain cases of autoimmune infertility. With the provision herein of human MATER protein sequences and encoding nucleic acids, methods for the detection and diagnosis of such fertility failure are now enabled.

30 Autoantibodies that recognize an epitope of the human MATER protein can be detected in samples from a subject, for instance serum or other fluid, using known immunological techniques. The presence of such autoantibodies (*e.g.*, circulating autoantibodies specific for a MATER epitope) indicates that the subject suffers from MATER-mediated infertility or reduced fertility, or has an increased susceptibility to suffer from one of these conditions.

35 Many techniques are commonly known in the art for the detection and quantification of antigen. Most commonly, the purified antigen will be bound to a substrate, the antibody of the sample will bind via its Fab portion to this antigen, the substrate will then be washed and a second, labeled antibody will then be added which will bind to the Fc portion of the antibody that is the subject of the assay. The second, labeled antibody will be species specific, *i.e.*, if the serum is from a

-45-

human, the second, labeled antibody will be anti-human-IgG antibody. The specimen will then be washed and the amount of the second, labeled antibody that has been bound will be detected and quantified by standard methods.

Examples of methods for the detection of antibodies in biological samples, including methods employing dip strips or other immobilized assay devices, are disclosed for instance in the following patents: U.S. Patents No. 5,965,356 (Herpes simplex virus type specific seroassay); 6,114,179 (Method and test kit for detection of antigens and/or antibodies); 6,077,681 (Diagnosis of motor neuropathy by detection of antibodies); 6,057,097 (Marker for pathologies comprising an auto-immune reaction and/or for inflammatory diseases); and 5,552,285 (Immunoassay methods, compositions and kits for antibodies to oxidized DNA bases).

EXAMPLE 13

Suppression of MATER Expression.

A reduction of MATER protein expression in a transgenic cell may be obtained by introducing into cells an antisense construct based on the *Mater* encoding sequence, including the human *Mater* cDNA or fragments thereof (for instance, the cDNA fragments shown in SEQ ID NO: 1 and 3) or gene sequence or flanking regions thereof. For antisense suppression, a nucleotide sequence from an MATER encoding sequence, *e.g.* all or a portion of the *Mater* cDNA or gene, is arranged in reverse orientation relative to the promoter sequence in the transformation vector. Other aspects of the vector may be chosen as discussed above (Example 8).

The introduced sequence need not be the full length human *Mater* cDNA (SEQ ID NO: 23) or gene, and need not be exactly homologous to the equivalent sequence found in the cell type to be transformed. Thus, portions or fragments of the murine cDNA (SEQ ID NO: 5) could also be used to knock out expression of the human *Mater* gene. Generally, however, where the introduced sequence is of shorter length, a higher degree of homology to the native *Mater* sequence will be needed for effective antisense suppression. The introduced antisense sequence in the vector may be at least 30 nucleotides in length, and improved antisense suppression typically will be observed as the length of the antisense sequence increases. The length of the antisense sequence in the vector advantageously may be greater than 100 nucleotides, and can be up to about the full length of the human *Mater* cDNA or gene. For suppression of the *Mater* gene itself, transcription of an antisense construct results in the production of RNA molecules that are the reverse complement of mRNA molecules transcribed from the endogenous *Mater* gene in the cell.

Although the exact mechanism by which antisense RNA molecules interfere with gene expression has not been elucidated, it is believed that antisense RNA molecules bind to the endogenous mRNA molecules and thereby inhibit translation of the endogenous mRNA.

Suppression of endogenous MATER expression can also be achieved using ribozymes. Ribozymes are synthetic RNA molecules that possess highly specific endoribonuclease activity. The production and use of ribozymes are disclosed in U.S. Patent No. 4,987,071 to Cech and U.S. Patent No. 5,543,508 to Haselhoff. The inclusion of ribozyme sequences within antisense RNAs may be

-46-

used to confer RNA cleaving activity on the antisense RNA, such that endogenous mRNA molecules that bind to the antisense RNA are cleaved, which in turn leads to an enhanced antisense inhibition of endogenous gene expression.

Finally, dominant negative mutant forms of MATER may be used to block endogenous
5 MATER activity.

EXAMPLE 14

MATER Knockout and Overexpression Transgenic Animals

Mutant organisms that under-express or over-express MATER protein are useful for research.
10 Such mutants allow insight into the physiological and/or pathological role of MATER in a healthy and/or pathological organism. These mutants are “genetically engineered,” meaning that information in the form of nucleotides has been transferred into the mutant’s genome at a location, or in a combination, in which it would not normally exist. Nucleotides transferred in this way are said to be “non-native.” For example, a non-*Mater* promoter inserted upstream of a native *Mater* gene would be
15 non-native. An extra copy of a *Mater* gene or other encoding sequence on a plasmid, transformed into a cell, would be non-native, whether that extra copy was *Mater* derived from the same or a different species.

Mutants may be, for example, produced from mammals, such as mice, that either over-express or under-express MATER protein, or that do not express MATER at all. Over-expression mutants are
20 made by increasing the number of MATER-encoding sequences (such as genes) in the organism, or by introducing an MATER-encoding sequence into the organism under the control of a constitutive or inducible or viral promoter such as the mouse mammary tumor virus (MMTV) promoter or the whey acidic protein (WAP) promoter or the metallothionein promoter. Mutants that under-express MATER may be made by using an inducible or repressible promoter, or by deleting the *Mater* gene, or by
25 destroying or limiting the function of the *Mater* gene, for instance by disrupting the gene by transposon insertion.

Antisense genes may be engineered into the organism, under a constitutive or inducible promoter, to decrease or prevent MATER expression, as discussed above in Example 13.

A gene is “functionally deleted” when genetic engineering has been used to negate or reduce
30 gene expression to negligible levels. When a mutant is referred to in this application as having the *Mater* gene altered or functionally deleted, this refers to the *Mater* gene and to any ortholog of this gene. When a mutant is referred to as having “more than the normal copy number” of a gene, this means that it has more than the usual number of genes found in the wild-type organism, *e.g.*, in the diploid mouse or human.

35 A mutant mouse over-expressing MATER may be made by constructing a plasmid having the *Mater* gene driven by a promoter, such as the mouse mammary tumor virus (MMTV) promoter or the whey acidic protein (WAP) promoter. This plasmid may be introduced into mouse oocytes by microinjection. The oocytes are implanted into pseudopregnant females, and the litters are assayed for insertion of the transgene. Multiple strains containing the transgene are then available for study.

-47-

WAP is quite specific for mammary gland expression during lactation, and MMTV is expressed in a variety of tissues including mammary gland, salivary gland and lymphoid tissues. Many other promoters might be used to achieve various patterns of expression, *e.g.*, the metallothionein promoter.

5 An inducible system may be created in which the subject expression construct is driven by a promoter regulated by an agent that can be fed to the mouse, such as tetracycline. Such techniques are well known in the art.

A mutant knockout animal (*e.g.*, mouse) from which the *Mater* gene is deleted or otherwise disabled can be made by removing coding regions of the *Mater* gene from embryonic stem cells. The methods of creating deletion mutations by using a targeting vector have been described (see, for instance, Thomas and Capecch, *Cell* 51:503-512, 1987). One specific example of the production of a *Mater* null mouse is described above, in Example 1.

EXAMPLE 15

15 Gene Therapy

Gene therapy approaches for combating MATER-mediated fertility defects in subjects, or for causing MATER-mediated infertility in subjects, are now made possible.

Retroviruses have been considered the preferred vector for experiments in gene therapy, with a high efficiency of infection and stable integration and expression (Orkin *et al.*, *Prog. Med. Genet.* 7:130-142, 1988). The full-length *Mater* gene or cDNA can be cloned into a retroviral vector and driven from either its endogenous promoter or, for instance, from the retroviral LTR (long terminal repeat). Other viral transfection systems may also be utilized for this type of approach, including adenovirus, adeno-associated virus (AAV) (McLaughlin *et al.*, *J. Virol.* 62:1963-1973, 1988), Vaccinia virus (Moss *et al.*, *Annu. Rev. Immunol.* 5:305-324, 1987), Bovine Papilloma virus (Rasmussen *et al.*, *Methods Enzymol.* 139:642-654, 1987) or members of the herpesvirus group such as Epstein-Barr virus (Margolskee *et al.*, *Mol. Cell. Biol.* 8:2837-2847, 1988).

Recent developments in gene therapy techniques include the use of RNA-DNA hybrid oligonucleotides, as described by Cole-Strauss, *et al.* (*Science* 273:1386-1389, 1996). This technique may allow for site-specific integration of cloned sequences, thereby permitting accurately targeted gene replacement.

In addition to delivery of *Mater* to cells using viral vectors, it is possible to use non-infectious methods of delivery. For instance, lipidic and liposome-mediated gene delivery has recently been used successfully for transfection with various genes (for reviews, see Templeton and Lasic, *Mol. Biotechnol.* 11:175-180, 1999; Lee and Huang, *Crit. Rev. Ther. Drug Carrier Syst.* 14:173-206; and Cooper, *Semin. Oncol.* 23:172-187, 1996). For instance, cationic liposomes have been analyzed for their ability to transfect monocytic leukemia cells, and shown to be a viable alternative to using viral vectors (de Lima *et al.*, *Mol. Membr. Biol.* 16:103-109, 1999). Such cationic liposomes can also be targeted to specific cells through the inclusion of, for instance,

63198-1404

-48-

monoclonal antibodies or other appropriate targeting ligands (Kao *et al.*, *Cancer Gene Ther.* 3:250-256, 1996).

EXAMPLE 16

5

Identification of Therapeutic Compounds

The human MATER molecules disclosed herein can be used to identify (screen for) compounds that are useful in influencing MATER-mediated fertility in a mammal, either by blocking (inhibiting) the activity of MATER (and thereby reducing fertility) or enhancing MATER activity (and thereby increasing fertility).

10

Such screening methods can include determining if a test compound binds directly to or otherwise interacts with a MATER protein, or a variant or fragment of a MATER protein. Proteins that do bind to such a molecule are select for further characterization.

15

In specific embodiments, the compound being tested for activity is applied to a cell for instance a test cell (*e.g.*, a developing oocyte or embryo of a mammal). The activity of the MATER protein in that test cell is then measured, for instance by determining whether the oocyte or embryo progresses beyond the two-cell stage. If application of the test compound alters proportion of embryos that progress beyond two cells, then that compound is selected as a likely candidate for further characterization. In particular examples, a test agent that opposes or inhibits a MATER activity is selected for further study, for example by exposing the agent to a mammalian female reproductive system *in vivo*, to determine if *in vivo* fertility is inhibited. Such identified compounds may be useful as contraceptive agents.

20

Specific examples of compounds likely to be effective at inhibiting MATER activity, and therefore effective as contraceptives, include antibodies specifically directed to epitopes of the MATER protein. The general concept of immunocontraceptives has been described (see, *e.g.*, U.S. Patents No. 5,637,300 and 6,027,727, describing contraceptive antibodies directed to proteins of the *zona pellucida*).

25

Alternatively, compounds that increase the proportion of embryos that progress beyond two cells (for instance, in an animal system known to be defective for fertility) are selected for further study as possible fertility enhancing agents. Similar screens can be used to identify compounds that mimic the activity of MATER protein, for instance in a *Mater* null animal.

30

In addition, it is suggested that MATER protein perform its function within the cytoplasm through interacting with other unknown protein. Physical blockage of such interaction is expected to arrest the MATER function. Candidates for such interactions are being identified based on interactions in a yeast two-hybrid system (Fields and Songs, *Nature* 340:245, 1989). Once molecular domains for the protein interaction are known, molecules can be designed directly based to specifically inhibit or otherwise interfere with MATER protein interactions with these other proteins.

35

Compounds selected using these methods are comprehended by this disclosure.

EXAMPLE 17**Kits**

Kits are provided which contain the necessary reagents for determining *Mater* gene copy number, such as probes or primers specific for the *Mater* gene, as well as written instructions. Kits are also provided to determine abnormal expression of *Mater* mRNA (*i.e.*, containing probes) or MATER protein (*i.e.*, containing a MATER-specific binding agent). Instructions provided in the diagnostic kits can include calibration curves or charts to compare with the determined (*e.g.*, experimentally measured) values.

A. Kits for Detection of *Mater* Genomic Sequences

The nucleotide sequences disclosed herein, and fragments thereof, can be supplied in the form of a kit for use in detection of *Mater* genomic sequences and/or diagnosis of infertility or reduced fertility. In such a kit, an appropriate amount of one or more of the *Mater*-specific oligonucleotide primers is provided in one or more containers. The oligonucleotide primers may be provided suspended in an aqueous solution or as a freeze-dried or lyophilized powder, for instance. The container(s) in which the oligonucleotide(s) are supplied can be any conventional container that is capable of holding the supplied form, for instance, microfuge tubes, ampoules, or bottles. In some applications, pairs of primers may be provided in pre-measured single use amounts in individual, typically disposable, tubes or equivalent containers. With such an arrangement, the sample to be tested for the presence of *Mater* genomic amplification can be added to the individual tubes and *in vitro* amplification carried out directly.

The amount of each oligonucleotide primer supplied in the kit can be any appropriate amount, depending for instance on the market to which the product is directed. For instance, if the kit is adapted for research or clinical use, the amount of each oligonucleotide primer provided would likely be an amount sufficient to prime several *in vitro* amplification reactions. Those of ordinary skill in the art know the amount of oligonucleotide primer that is appropriate for use in a single amplification reaction. General guidelines may for instance be found in Innis *et al.* (*PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc., San Diego, CA, 1990), Sambrook *et al.* (*In Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York, 1989), and Ausubel *et al.* (*In Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1998).

A kit may include more than two primers, in order to facilitate the PCR *in vitro* amplification of *Mater* sequences, for instance the *Mater* gene, specific exon(s) or other portions of the gene, or the 5' or 3' flanking region thereof.

In some embodiments, kits may also include the reagents necessary to carry out PCR *in vitro* amplification reactions, including, for instance, DNA sample preparation reagents, appropriate buffers (*e.g.*, polymerase buffer), salts (*e.g.*, magnesium chloride), and deoxyribonucleotides (dNTPs). Written instructions may also be included.

Kits may in addition include either labeled or unlabeled oligonucleotide probes for use in detection of the *in vitro* amplified *Mater* sequences. The appropriate sequences for such a probe will

-50-

be any sequence that falls between the annealing sites of the two provided oligonucleotide primers, such that the sequence the probe is complementary to is amplified during the *in vitro* amplification reaction.

It may also be advantageous to provided in the kit one or more control sequences for use in the amplification reactions. The design of appropriate positive control sequences is well known to one of ordinary skill in the appropriate art.

B. Kits for Detection of *Mater* mRNA Expression

Kits similar to those disclosed above for the detection of *Mater* genomic sequences can be used to detect *Mater* mRNA expression levels. Such kits may include an appropriate amount of one or more of the oligonucleotide primers for use in reverse transcription amplification reactions, similarly to those provided above, with art-obvious modifications for use with RNA.

In some embodiments, kits for detection of *Mater* mRNA expression levels may also include the reagents necessary to carry out RT-PCR *in vitro* amplification reactions, including, for instance, RNA sample preparation reagents (including *e.g.*, an RNase inhibitor), appropriate buffers (*e.g.*, polymerase buffer), salts (*e.g.*, magnesium chloride), and deoxyribonucleotides (dNTPs). Written instructions may also be included.

Kits in addition may include either labeled or unlabeled oligonucleotide probes for use in detection of the *in vitro* amplified target sequences. The appropriate sequences for such a probe will be any sequence that falls between the annealing sites of the two provided oligonucleotide primers, such that the sequence the probe is complementary to is amplified during the PCR reaction.

It also may be advantageous to provided in the kit one or more control sequences for use in the RT-PCR reactions. The design of appropriate positive control sequences is well known to one of ordinary skill in the appropriate art.

Alternatively, kits may be provided with the necessary reagents to carry out quantitative or semi-quantitative Northern analysis of *Mater* mRNA. Such kits include, for instance, at least one *Mater*-specific oligonucleotide for use as a probe. This oligonucleotide may be labeled in any conventional way, including with a selected radioactive isotope, enzyme substrate, co-factor, ligand, chemiluminescent or fluorescent agent, hapten, or enzyme.

C. Kits For Detection of MATER Protein or Peptide Expression

Kits for the detection of MATER protein expression, for instance MATER at least one target (*e.g.*, MATER) protein specific binding agent (*e.g.*, a polyclonal or monoclonal antibody or antibody fragment) and may include at least one control. The MATER protein specific binding agent and control may be contained in separate containers. The kits may also include means for detecting MATER:agent complexes, for instance the agent may be detectably labeled. If the detectable agent is not labeled, it may be detected by second antibodies or protein A for example which may also be provided in some kits in one or more separate containers. Such techniques are well known.

63198-1404

-51-

Additional components in some kits include instructions for carrying out the assay. Instructions will allow the tester to determine whether MATER expression levels are altered, for instance in comparison to a control sample. Reaction vessels and auxiliary reagents such as chromogens, buffers, enzymes, etc. may also be included in the kits.

5 By way of example only, an effective and convenient immunoassay kit such as an enzyme-linked immunosorbent assay can be constructed to test anti-MATER antibody in human serum, as reported for detection of non-specific anti-ovarian antibodies (Wheatcroft *et al*, *Clin. Exp. Immunol.* 96:122-128, 1994; Wheatcroft *et al*, *Hum. Reprod.* 12:2617-2622, 1997). Expression vectors can be constructed using the human MATER cDNA to produce the recombinant human MATER protein in
10 either bacteria or baculovirus (as described in Example 8). By affinity purification, unlimited amounts of pure recombinant MATER protein can be produced.

 An assay kit could provide the recombinant protein as an antigen and enzyme-conjugated goat anti-human IgG as a second antibody as well as the enzymatic substrates. Such kits can be used to test if the patient sera contain antibodies against human MATER.

15

 This disclosure provides *Mater* nucleic acids and proteins, including the human *Mater* molecules described above. The disclosure further provides methods employing these molecules, including methods to predict and/or diagnose infertility, reduced fertility, or reproductive failure in females, as well as treatments for such infertility and reduced fertility, and contraceptives. It will be
20 apparent that the precise details of the molecules and methods described may be varied or under- or overexpression, are also contemplated. Such kits will include modified without departing from the scope of the claimed invention. The inventors claim all such modifications and variations that fall within the scope of the claims below.

SEQUENCE LISTING

<110> THE GOVERNMENT OF THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH & HUMAN SERVICES, THE NATIONAL INSTITUTES OF HEALTH

Nelson, Lawrence M.

Tong, Zhi-Bin

<120> HUMAN GENE CRITICAL TO FERTILITY

<130> 4239-55874

<150> 60/241,510

<151> 2000-10-18

<160> 24

<170> PatentIn version 3.0

<210> 1

<211> 1157

<212> DNA

<213> homo sapiens

<400> 1

```

caagctccgg tgacggagat catgtcccga ccagaaaggc tgttgttcat cattgacggt      60
ttcgatgacc tgggctctgt cctcaacaat gacacaaagc tctgcaaaga ctgggctgag      120
aagcagcctc cggttcaccct catacgcagt ctgctgagga aggtcctgct ccctgagtcc      180
ttcctgatcg tcaccgtcag agacgtgggc acagagaagc tcaagtcaga ggtcgtgtct      240
ccccgttacc tgtagttag aggaatctcc ggggaacaaa gaatccactt gtccttgag      300
cgcggggattg gtgagcatca gaagacacaa ggggtgcgtg cgatcatgaa caaccgtgag      360
ctgctcgacc agtgccaggt gcccgccgtg ggctctctca tctgcgtggc cctgcagctg      420
caggacgtgg tgggggagag cgtcgccccc ttcaaccaa cgctcacagg cctgcacgcc      480
gcttttgtgt ttcatcagct caccctcga ggcgtgggtc ggcgtgtct caatctggag      540
gaaagagttg tcctgaagcg cttctgccgt atggctgtgg agggagtgtg gaataggag      600
tcagtgtttg acggtgacga cctcatggtt caaggactcg gggagtctga gctccgtgct      660
ctgtttcaca tgaacatcct tctcccagac agccactgtg aggagtacta caccttcttc      720
cacctcagtc tccaggactt ctgtgccgcc ttgtactacy tgtagaggg cctggaaatc      780
gagccagctc tctgccctct gtacgttgag aagacaaaaga ggtccatgga gcttaaacag      840
gcaggcttcc atatccactc gctttggatg aagcgtttct tgtttggcct cgtgagcgaa      900
gacgtaagga ggccactgga ggtcctgctg ggctgtcccc ttcccctggg ggtgaagcag      960
aagcttctgc actgggtctc tctgttgggt cagcagccta atgccaccac cccaggagac     1020
accctggacg ccttccactg tcttttcgag actcaagaca aagagtttgt tcgcttggca     1080
ttaaacagct tccaagaagt gtggcttccg attaaccaga acctggactt gatagcatct     1140
tccttctgcc tccagct                                     1157

```

<210> 2

<211> 385

<212> PRT

<213> homo sapiens

<400> 2

```

Gln Ala Pro Val Thr Glu Ile Met Ser Arg Pro Glu Arg Leu Leu Phe
1           5           10           15
Ile Ile Asp Gly Phe Asp Asp Leu Gly Ser Val Leu Asn Asn Asp Thr
20           25           30

```

Lys Leu Cys Lys Asp Trp Ala Glu Lys Gln Pro Pro Phe Thr Leu Ile
 35 40 45
 Arg Ser Leu Leu Arg Lys Val Leu Leu Pro Glu Ser Phe Leu Ile Val
 50 55 60
 Thr Val Arg Asp Val Gly Thr Glu Lys Leu Lys Ser Glu Val Val Ser
 65 70 75 80
 Pro Arg Tyr Leu Leu Val Arg Gly Ile Ser Gly Glu Gln Arg Ile His
 85 90 95
 Leu Leu Leu Glu Arg Gly Ile Gly Glu His Gln Lys Thr Gln Gly Leu
 100 105 110
 Arg Ala Ile Met Asn Asn Arg Glu Leu Leu Asp Gln Cys Gln Val Pro
 115 120 125
 Ala Val Gly Ser Leu Ile Cys Val Ala Leu Gln Leu Gln Asp Val Val
 130 135 140
 Gly Glu Ser Val Ala Pro Phe Asn Gln Thr Leu Thr Gly Leu His Ala
 145 150 155 160
 Ala Phe Val Phe His Gln Leu Thr Pro Arg Gly Val Val Arg Arg Cys
 165 170 175
 Leu Asn Leu Glu Arg Val Val Leu Lys Arg Phe Cys Arg Met Ala
 180 185 190
 Val Glu Gly Val Trp Asn Arg Lys Ser Val Phe Asp Gly Asp Asp Leu
 195 200 205
 Met Val Gln Gly Leu Gly Glu Ser Glu Leu Arg Ala Leu Phe His Met
 210 215 220
 Asn Ile Leu Leu Pro Asp Ser His Cys Glu Glu Tyr Tyr Thr Phe Phe
 225 230 235 240
 His Leu Ser Leu Gln Asp Phe Cys Ala Ala Leu Tyr Tyr Val Leu Glu
 245 250 255
 Gly Leu Glu Ile Glu Pro Ala Leu Cys Pro Leu Tyr Val Glu Lys Thr
 260 265 270
 Lys Arg Ser Met Glu Leu Lys Gln Ala Gly Phe His Ile His Ser Leu
 275 280 285
 Trp Met Lys Arg Phe Leu Phe Gly Leu Val Ser Glu Asp Val Arg Arg
 290 295 300
 Pro Leu Glu Val Leu Leu Gly Cys Pro Val Pro Leu Gly Val Lys Gln
 305 310 315 320
 Lys Leu Leu His Trp Val Ser Leu Leu Gly Gln Gln Pro Asn Ala Thr
 325 330 335
 Thr Pro Gly Asp Thr Leu Asp Ala Phe His Cys Leu Phe Glu Thr Gln
 340 345 350
 Asp Lys Glu Phe Val Arg Leu Ala Leu Asn Ser Phe Gln Glu Val Trp
 355 360 365
 Leu Pro Ile Asn Gln Asn Leu Asp Leu Ile Ala Ser Ser Phe Cys Leu
 370 375 380
 Gln
 385

<210> 3
 <211> 1075
 <212> DNA
 <213> homo sapiens

<400> 3
 tctggcctca gccctcgtca gcaaccggag cttgacacac ctgtgcctat ccaacaacag 60
 cctggggaac gaaggtgtaa atctactgtg tcgatccatg aggcttcccc actgtagtct 120
 gcagaggctg atgctgaatc agtgccacct ggacacggct ggctgtggtt ctcttgcaact 180
 tgcgcttatg ggtaactcat ggctgacyca cctgagcctt agcatgaacc ctgtggaaga 240
 caatggcgtg aagcttctgt gcgaggtcat gagagaacca tcttgtcatc tccaggacct 300

```

ggagttggta aaatgtcatc tcaccgccgc gtgctgtgag agtctgtcct gtgtgatctc 360
gaggagcaga cacctgaaga gcctggatct cacggacaat gccctgggtg acggtgggtg 420
tgctgcgctg tgcgaggagac tgaagcaaaa gaacagtgtt ctgacgagac tcgggttgaa 480
ggcatgtgga ctgacttctg attgctgtga ggcactctcc ttggcccttt cctgcaaccg 540
gcatctgacc agtctaaacc tgggtgcagaa taacttcagt cccaaaggaa tgatgaagct 600
gtgttcggcc tttgcctgtc ccacgtctaa cttacagata attgggctgt ggaaatggca 660
gtaccctgtg caaataagga agctgctgga ggaagtgcag ctactcaagc cccgagtcgt 720
aattgacggg agttggcatt cttttgatga agatgaccgg tactggtgga aaaactgaag 780
atacggaaac ctgccccact cacacccatc tgatggagga actttaaacg ctgttttctc 840
agagcaagct atgcacctgg gagttccttc tcaaagatgg agaattgatt ctgattctca 900
caaagccctc aatggtagtg attcttctgt gttcactcta cgttgggttac tggatttgaa 960
ggctagagac cttcaagtca taggactcag tatctgtgaa atgtccgtca tatctcagag 1020
catatagagg gaattaaata aacacaaagc atttggaata aaaaaaaaaa aaaaaa 1075

```

```

<210> 4
<211> 258
<212> PRT
<213> homo sapiens

```

```

<400> 4
Leu Ala Ser Ala Leu Val Ser Asn Arg Ser Leu Thr His Leu Cys Leu
1          5          10          15
Ser Asn Asn Ser Leu Gly Asn Glu Gly Val Asn Leu Leu Cys Arg Ser
20          25          30
Met Arg Leu Pro His Cys Ser Leu Gln Arg Leu Met Leu Asn Gln Cys
35          40          45
His Leu Asp Thr Ala Gly Cys Gly Ser Leu Ala Leu Ala Leu Met Gly
50          55          60
Asn Ser Trp Leu Thr His Leu Ser Leu Ser Met Asn Pro Val Glu Asp
65          70          75          80
Asn Gly Val Lys Leu Leu Cys Glu Val Met Arg Glu Pro Ser Cys His
85          90          95
Leu Gln Asp Leu Glu Leu Val Lys Cys His Leu Thr Ala Ala Cys Cys
100         105         110
Glu Ser Leu Ser Cys Val Ile Ser Arg Ser Arg His Leu Lys Ser Leu
115         120         125
Asp Leu Thr Asp Asn Ala Leu Gly Asp Gly Gly Val Ala Ala Leu Cys
130         135         140
Glu Gly Leu Lys Gln Lys Asn Ser Val Leu Thr Arg Leu Gly Leu Lys
145         150         155         160
Ala Cys Gly Leu Thr Ser Asp Cys Cys Glu Ala Leu Ser Leu Ala Leu
165         170         175
Ser Cys Asn Arg His Leu Thr Ser Leu Asn Leu Val Gln Asn Asn Phe
180         185         190
Ser Pro Lys Gly Met Met Lys Leu Cys Ser Ala Phe Ala Cys Pro Thr
195         200         205
Ser Asn Leu Gln Ile Ile Gly Leu Trp Lys Trp Gln Tyr Pro Val Gln
210         215         220
Ile Arg Lys Leu Leu Glu Val Gln Leu Leu Lys Pro Arg Val Val
225         230         235         240
Ile Asp Gly Ser Trp His Ser Phe Asp Glu Asp Asp Arg Tyr Trp Trp
245         250         255
Lys Asn

```

```

<210> 5
<211> 3447

```

<212> DNA

<213> Mus musculus

<400> 5

aaagcacaat	gggtcctcca	gaaaaagaaa	gtaaagcaat	cttgaaagca	cgtggattgg	60
aagaggaaca	gaagtcagaa	agaaaaatga	cttctccaga	aaacgacagt	aaatcaatcc	120
agaaagacca	aggaccagag	caggagcaga	catcagaaaag	cacaatgggt	cctccagaaa	180
aagacagtaa	agcaatcttg	aaagcacgtg	gatttgaaga	ggaacagaag	tcagaaagca	240
caatgtctcc	ttcagaaaat	gtcagtagag	caatcctgaa	agacagtgga	tcagaagaag	300
tggacaggc	gtcagaaaaga	aaaatgactt	ctccagaaaa	cgacagtaaa	tcaatccaga	360
aagaccaagg	accagagcag	gagcagacat	cagaaacctt	acaatctaag	gaagaagatg	420
aagtgcagag	ggcagataaa	gataatggag	gtgacttaca	agactacaag	gcccattgtg	480
ttgctaagtt	cgacacaaat	gtggatctac	actatgacag	cccagagatg	aaattattgt	540
ctgatgcttt	taaaccatac	cagaaaacct	tccagcctca	caccattatc	ctacatggaa	600
gaccaggagt	tgggaagtca	gcttttgcca	gaagtattgt	tcttggtgg	gcacagggta	660
aactcttcca	aaaaatgtcc	tttgtcatct	tcttctctgt	tagagaaata	aagtggacag	720
agaagagcag	tttggcacag	ctgattgcta	aggagtgtcc	agactcctgg	gatctagtga	780
caaagatcat	gtcccaacca	gaaagactct	tgtttgtcat	agatggcttg	gatgatggg	840
actctgtcct	ccaacatgat	gatatgacac	tatccagaga	ctggaaggat	gaacagccca	900
tatacatcct	gatgtacagc	ctcctgagga	atgctctctt	acctcagtcc	tttctcatca	960
ttaccaccag	aaacacaggc	ttagaaaaac	tcaagtcaat	ggttgtgtcc	cccctctata	1020
tactggttga	aggactgtct	gcatcaagga	gatctcagct	ggtcctcgag	aacatctcca	1080
atgagtctga	tagaatacaa	gtcttccatt	ctctgataga	aaatcaccag	ctgtttgacc	1140
aatgccaggc	cccctctgtg	tgtctctgg	tctgtgaggg	tctacagcta	cagaagaaac	1200
tgggaaagag	atgcacccta	ccctgccaga	ctctcaccgg	tttgtatgcc	acgttggtgt	1260
ttcaccagct	caccttgaaa	aggccttccc	agagcgctct	cagtcaggaa	gaacagatta	1320
ctctagtggg	tttgtgcatg	atggcagctg	aaggagtgtg	gaccatgagg	tcggtgttct	1380
atgatgatga	cctgaagaac	tatagcctaa	aggagtctga	gatcttgagg	ctctttcaca	1440
tgaacatcct	tctccagggt	ggccacaaca	gtgagcagtg	ttatgttttc	tcccacctca	1500
gcctgcagga	tttctttgct	gccttatatt	atgtttttaga	agggctggag	gaatggaatc	1560
agcatttttg	cttcattgaa	aaccaaagga	gcatcatgga	ggtgaagaga	actgacgaca	1620
ctcgccctct	cgggatgaag	cgtttcttat	ttggcctcat	gaacaaggat	atcttgaaga	1680
ctctggagggt	tctgtttgaa	tatcccgtag	ttccaaactgt	tgagcagaag	ctccaaact	1740
gggtctctct	gatagctcag	cagggtcaatg	gcaccagccc	aatggacacc	ctggatgcct	1800
tctattgtct	atttgagtct	caggatgaag	agtttgttgg	cggggctctc	aaacgcttcc	1860
aagaagtgtg	gctgtgtgatt	aaccagaaga	tggacttgaa	ggtctcttcc	tactgtctca	1920
agcactgtca	gaacttgaag	gcaatccggg	tggatatcag	agacctctc	tcggtagata	1980
atactctcga	gctgtgccct	gttggttactg	tccaggagac	acaatgtaag	cccctcctca	2040
tggagtgggtg	gggaaacttc	tgctctgtgc	ttggcagcct	ccggaacttg	aaggagctgg	2100
acttggggcga	cagcatcctg	agtcaacggg	ccatgaagat	actgtgcctc	gagctgcgga	2160
atcagtcctg	cagaatacag	aagctgacgt	ttaagagtgc	agaggtagtg	tctggcctga	2220
aacatctctg	gaagctcctt	tttagcaatc	aaaacttaaa	gtacctcaat	ctaggggaaca	2280
ctcccatgaa	ggatgatgac	atgaagttag	cctgcgaagc	gctgaaacat	ccaaagtgtc	2340
ccgtggagac	tctgagggtg	gattcctgtg	agtttaacct	cattgggtat	gagatgatct	2400
ccacgcttct	tatttcaacc	accaggctaa	agtgtctcag	cctggccaaa	aatagagtgg	2460
gagtaaaaag	catgatatcc	cttgggaatg	ccttgagtag	ctcaatgtgt	ctactgcaaa	2520
agttgatact	ggacaactgt	ggcctcacac	ctgccagctg	ccaccttctg	gtctcagccc	2580
ttttcagcaa	ccagaacttg	acacacctgt	gcctgtcaaa	caacagcctg	gggactgaag	2640
gagtgcacaa	gctgtgtcag	ttcctgagga	atccagaatg	tgctctccag	cggctgatac	2700
tgaatcactg	caacattgta	gatgatgctt	atggcttctc	ggcaatgaga	cttgcaaaaca	2760
acacaaaagt	gaccacactg	agcctgacca	tgaacccctg	aggggatggg	gcaatgaagc	2820
tactgtgtga	agctttaaag	gaacctactt	gttaccttca	agaactggaa	ctagtggact	2880
gccaactcac	acagaactgc	tgcgaggacc	tggcctgtat	gatcacaaca	accaagcact	2940
taaaaagttt	ggatcttggg	aacaacgccc	tgggtgacaa	aggagtcata	accctgtgtg	3000
agggactgaa	gcaaagtagc	agctccctga	ggagacttgg	gttgggggca	tgtaagttga	3060
cttccaattg	ctgtgaggca	ttgtcattgg	ccatctcttg	caacctcac	ctgaacagcc	3120
taaacctggg	gaagaatgac	ttcagtacat	cggggatgtt	gaagctgtgc	tctgcgttcc	3180
aatgccctgt	ctctaacctg	gggataattg	gcctgtggaa	gcaggagtac	tatgcccgag	3240
tgagaagaca	gctggaggaa	gttgagtttg	tcaagcccca	cgtggtgatt	gatggtgatt	3300

```

ggatgctag tgatgaagat gaccgaaact ggtggaaaaa ctgaagacat gagccccctc 3360
tccttcacgt cctagcactg cagtatctgt gaaatgtttg tctcaccttg gaggatgtag 3420
caagaatgaa ataaacacag catttag 3447

```

```

<210> 6
<211> 1111
<212> PRT
<213> Mus musculus

```

```

<400> 6
Met Gly Pro Pro Glu Lys Glu Ser Lys Ala Ile Leu Lys Ala Arg Gly
1      5      10      15
Leu Glu Glu Glu Gln Lys Ser Glu Arg Lys Met Thr Ser Pro Glu Asn
20     25     30
Asp Ser Lys Ser Ile Gln Lys Asp Gln Gly Pro Glu Gln Glu Gln Thr
35     40     45
Ser Glu Ser Thr Met Gly Pro Pro Glu Lys Asp Ser Lys Ala Ile Leu
50     55     60
Lys Ala Arg Gly Leu Glu Glu Glu Gln Lys Ser Glu Ser Thr Met Ser
65     70     75     80
Pro Ser Glu Asn Val Ser Arg Ala Ile Leu Lys Asp Ser Gly Ser Glu
85     90     95
Glu Val Glu Gln Ala Ser Glu Arg Lys Met Thr Ser Pro Glu Asn Asp
100    105    110
Ser Lys Ser Ile Gln Lys Asp Gln Gly Pro Glu Gln Glu Gln Thr Ser
115    120    125
Glu Thr Leu Gln Ser Lys Glu Glu Asp Glu Val Thr Glu Ala Asp Lys
130    135    140
Asp Asn Gly Gly Asp Leu Gln Asp Tyr Lys Ala His Val Ile Ala Lys
145    150    155    160
Phe Asp Thr Ser Val Asp Leu His Tyr Asp Ser Pro Glu Met Lys Leu
165    170    175
Leu Ser Asp Ala Phe Lys Pro Tyr Gln Lys Thr Phe Gln Pro His Thr
180    185    190
Ile Ile Leu His Gly Arg Pro Gly Val Gly Lys Ser Ala Leu Ala Arg
195    200    205
Ser Ile Val Leu Gly Trp Ala Gln Gly Lys Leu Phe Gln Lys Met Ser
210    215    220
Phe Val Ile Phe Phe Ser Val Arg Glu Ile Lys Trp Thr Glu Lys Ser
225    230    235    240
Ser Leu Ala Gln Leu Ile Ala Lys Glu Cys Pro Asp Ser Trp Asp Leu
245    250    255
Val Thr Lys Ile Met Ser Gln Pro Glu Arg Leu Leu Phe Val Ile Asp
260    265    270
Gly Leu Asp Asp Met Asp Ser Val Leu Gln His Asp Asp Met Thr Leu
275    280    285
Ser Arg Asp Trp Lys Asp Glu Gln Pro Ile Tyr Ile Leu Met Tyr Ser
290    295    300
Leu Leu Arg Lys Ala Leu Leu Pro Gln Ser Phe Leu Ile Ile Thr Thr
305    310    315    320
Arg Asn Thr Gly Leu Glu Lys Leu Lys Ser Met Val Val Ser Pro Leu
325    330    335
Tyr Ile Leu Val Glu Gly Leu Ser Ala Ser Arg Arg Ser Gln Leu Val
340    345    350
Leu Glu Asn Ile Ser Asn Glu Ser Asp Arg Ile Gln Val Phe His Ser
355    360    365
Leu Ile Glu Asn His Gln Leu Phe Asp Gln Cys Gln Ala Pro Ser Val
370    375    380

```

Cys	Ser	Leu	Val	Cys	Glu	Ala	Leu	Gln	Leu	Gln	Lys	Lys	Leu	Gly	Lys	385	390	395	400
Arg	Cys	Thr	Leu	Pro	Cys	Gln	Thr	Leu	Thr	Gly	Leu	Tyr	Ala	Thr	Leu	405	410		415
Val	Phe	His	Gln	Leu	Thr	Leu	Lys	Arg	Pro	Ser	Gln	Ser	Ala	Leu	Ser	420	425		430
Gln	Glu	Glu	Gln	Ile	Thr	Leu	Val	Gly	Leu	Cys	Met	Met	Ala	Ala	Glu	435	440		445
Gly	Val	Trp	Thr	Met	Arg	Ser	Val	Phe	Tyr	Asp	Asp	Asp	Leu	Lys	Asn	450	455		460
Tyr	Ser	Leu	Lys	Glu	Ser	Glu	Ile	Leu	Ala	Leu	Phe	His	Met	Asn	Ile	465	470		475
Leu	Leu	Gln	Val	Gly	His	Asn	Ser	Glu	Gln	Cys	Tyr	Val	Phe	Ser	His	485	490		495
Leu	Ser	Leu	Gln	Asp	Phe	Phe	Ala	Ala	Leu	Tyr	Tyr	Val	Leu	Glu	Gly	500	505		510
Leu	Glu	Glu	Trp	Asn	Gln	His	Phe	Cys	Phe	Ile	Glu	Asn	Gln	Arg	Ser	515	520		525
Ile	Met	Glu	Val	Lys	Arg	Thr	Asp	Asp	Thr	Arg	Leu	Leu	Gly	Met	Lys	530	535		540
Arg	Phe	Leu	Phe	Gly	Leu	Met	Asn	Lys	Asp	Ile	Leu	Lys	Thr	Leu	Glu	545	550		555
Val	Leu	Phe	Glu	Tyr	Pro	Val	Ile	Pro	Thr	Val	Glu	Gln	Lys	Leu	Gln	565	570		575
His	Trp	Val	Ser	Leu	Ile	Ala	Gln	Gln	Val	Asn	Gly	Thr	Ser	Pro	Met	580	585		590
Asp	Thr	Leu	Asp	Ala	Phe	Tyr	Cys	Leu	Phe	Glu	Ser	Gln	Asp	Glu	Glu	595	600		605
Phe	Val	Gly	Gly	Ala	Leu	Lys	Arg	Phe	Gln	Glu	Val	Trp	Leu	Leu	Ile	610	615		620
Asn	Gln	Lys	Met	Asp	Leu	Lys	Val	Ser	Ser	Tyr	Cys	Leu	Lys	His	Cys	625	630		635
Gln	Asn	Leu	Lys	Ala	Ile	Arg	Val	Asp	Ile	Arg	Asp	Leu	Leu	Ser	Val	645	650		655
Asp	Asn	Thr	Leu	Glu	Leu	Cys	Pro	Val	Val	Thr	Val	Gln	Glu	Thr	Gln	660	665		670
Cys	Lys	Pro	Leu	Leu	Met	Glu	Trp	Trp	Gly	Asn	Phe	Cys	Ser	Val	Leu	675	680		685
Gly	Ser	Leu	Arg	Asn	Leu	Lys	Glu	Leu	Asp	Leu	Gly	Asp	Ser	Ile	Leu	690	695		700
Ser	Gln	Arg	Ala	Met	Lys	Ile	Leu	Cys	Leu	Glu	Leu	Arg	Asn	Gln	Ser	705	710		715
Cys	Arg	Ile	Gln	Lys	Leu	Thr	Phe	Lys	Ser	Ala	Glu	Val	Val	Ser	Gly	725	730		735
Leu	Lys	His	Leu	Trp	Lys	Leu	Leu	Phe	Ser	Asn	Gln	Asn	Leu	Lys	Tyr	740	745		750
Leu	Asn	Leu	Gly	Asn	Thr	Pro	Met	Lys	Asp	Asp	Asp	Met	Lys	Leu	Ala	755	760		765
Cys	Glu	Ala	Leu	Lys	His	Pro	Lys	Cys	Ser	Val	Glu	Thr	Leu	Arg	Leu	770	775		780
Asp	Ser	Cys	Glu	Leu	Thr	Ile	Ile	Gly	Tyr	Glu	Met	Ile	Ser	Thr	Leu	785	790		795
Leu	Ile	Ser	Thr	Thr	Arg	Leu	Lys	Cys	Leu	Ser	Leu	Ala	Lys	Asn	Arg	805	810		815
Val	Gly	Val	Lys	Ser	Met	Ile	Ser	Leu	Gly	Asn	Ala	Leu	Ser	Ser	Ser	820	825		830
Met	Cys	Leu	Leu	Gln	Lys	Leu	Ile	Leu	Asp	Asn	Cys	Gly	Leu	Thr	Pro	835	840		845

7

Ala Ser Cys His Leu Leu Val Ser Ala Leu Phe Ser Asn Gln Asn Leu
 850 855 860
 Thr His Leu Cys Leu Ser Asn Asn Ser Leu Gly Thr Glu Gly Val Gln
 865 870 875 880
 Gln Leu Cys Gln Phe Leu Arg Asn Pro Glu Cys Ala Leu Gln Arg Leu
 885 890 895
 Ile Leu Asn His Cys Asn Ile Val Asp Asp Ala Tyr Gly Phe Leu Ala
 900 905 910
 Met Arg Leu Ala Asn Asn Thr Lys Leu Thr His Leu Ser Leu Thr Met
 915 920 925
 Asn Pro Val Gly Asp Gly Ala Met Lys Leu Leu Cys Glu Ala Leu Lys
 930 935 940
 Glu Pro Thr Cys Tyr Leu Gln Glu Leu Glu Leu Val Asp Cys Gln Leu
 945 950 955 960
 Thr Gln Asn Cys Cys Glu Asp Leu Ala Cys Met Ile Thr Thr Thr Lys
 965 970 975
 His Leu Lys Ser Leu Asp Leu Gly Asn Asn Ala Leu Gly Asp Lys Gly
 980 985 990
 Val Ile Thr Leu Cys Glu Gly Leu Lys Gln Ser Ser Ser Ser Leu Arg
 995 1000 1005
 Arg Leu Gly Leu Gly Ala Cys Lys Leu Thr Ser Asn Cys Cys Glu
 1010 1015 1020
 Ala Leu Ser Leu Ala Ile Ser Cys Asn Pro His Leu Asn Ser Leu
 1025 1030 1035
 Asn Leu Val Lys Asn Asp Phe Ser Thr Ser Gly Met Leu Lys Leu
 1040 1045 1050
 Cys Ser Ala Phe Gln Cys Pro Val Ser Asn Leu Gly Ile Ile Gly
 1055 1060 1065
 Leu Trp Lys Gln Glu Tyr Tyr Ala Arg Val Arg Arg Gln Leu Glu
 1070 1075 1080
 Glu Val Glu Phe Val Lys Pro His Val Val Ile Asp Gly Asp Trp
 1085 1090 1095
 Tyr Ala Ser Asp Glu Asp Arg Asn Trp Trp Lys Asn
 1100 1105 1110

<210> 7
 <211> 22
 <212> DNA
 <213> synthetic oligonucleotide

<400> 7
 tttcacatga acatccttct cc

22

<210> 8
 <211> 21
 <212> DNA
 <213> synthetic oligonucleotide

<400> 8
 agtgctggag gcagaaggaa g

21

<210> 9
 <211> 26
 <212> DNA
 <213> synthetic oligonucleotide

<400> 9
caggaattgg gaaatcggct ctctag 26

<210> 10
<211> 27
<212> DNA
<213> synthetic oligonucleotide

<400> 10
ccaaatgctt tgtgtttatt taattcc 27

<210> 11
<211> 19
<212> DNA
<213> synthetic oligonucleotide

<400> 11
cggaattcgt cactcagcg 19

<210> 12
<211> 18
<212> DNA
<213> synthetic oligonucleotide

<400> 12
agcgcgtgaa tcagatcg 18

<210> 13
<211> 24
<212> DNA
<213> synthetic oligonucleotide

<400> 13
attccctggg agagtccacc ttgc 24

<210> 14
<211> 24
<212> DNA
<213> synthetic oligonucleotide

<400> 14
acagcacgat ccttctggct agag 24

<210> 15
<211> 23
<212> DNA
<213> synthetic oligonucleotide

<400> 15
caagctccgg tgacggagat cat 23

<210> 16
<211> 22

<212> DNA
 <213> synthetic oligonucleotide

 <400> 16
 agctggaggc agaaggaaga tg 22

<210> 17
 <211> 27
 <212> DNA
 <213> synthetic oligonucleotide

 <400> 17
 tctggcctca gccctcgtca gcttgac 27

<210> 18
 <211> 27
 <212> DNA
 <213> synthetic oligonucleotide

 <400> 18
 ccaaattgctt tgtgtttatt taattcc 27

<210> 19
 <211> 21
 <212> DNA
 <213> synthetic oligonucleotide

 <400> 19
 tctggcctca gccctcgtca g 21

<210> 20
 <211> 21
 <212> DNA
 <213> synthetic oligonucleotide

 <400> 20
 actgaagtta ttctgcacca g 21

<210> 21
 <211> 20
 <212> DNA
 <213> synthetic oligonucleotide

 <400> 21
 tgactttctga ttgctgtgag 20

<210> 22
 <211> 20
 <212> DNA
 <213> synthetic oligonucleotide

 <400> 22
 ttccaccagt accggtcatc 20

<210> 23
 <211> 3900
 <212> DNA
 <213> homo sapiens

<220>
 <221> CDS
 <222> (1)..(3600)

<400> 23
 atg aag gtt gca gga gga ctt gaa ctt gga gct gct gct ctg ctc tca 48
 Met Lys Val Ala Gly Gly Leu Glu Leu Gly Ala Ala Ala Leu Leu Ser
 1 5 10 15
 gca tca cca cgt gct ctt gtc act ctt tcc aca ggt cct act tgc tct 96
 Ala Ser Pro Arg Ala Leu Val Thr Leu Ser Thr Gly Pro Thr Cys Ser
 20 25 30
 ata tta cca aag aat cca ctt ttc ccc caa aac ctg agc tct cag cct 144
 Ile Leu Pro Lys Asn Pro Leu Phe Pro Gln Asn Leu Ser Ser Gln Pro
 35 40 45
 tgt atc aag atg gaa gga gac aaa tcg ctc acc ttt tcc agc tac ggg 192
 Cys Ile Lys Met Glu Gly Asp Lys Ser Leu Thr Phe Ser Ser Tyr Gly
 50 55 60
 ctg caa tgg tgt ctc tat gag cta gac aag gaa gaa ttt cag aca ttc 240
 Leu Gln Trp Cys Leu Tyr Glu Leu Asp Lys Glu Glu Phe Gln Thr Phe
 65 70 75 80
 aag gaa tta cta aag aag aaa tct tca gaa tcg acc aca tgc tct att 288
 Lys Glu Leu Leu Lys Lys Lys Ser Ser Glu Ser Thr Thr Cys Ser Ile
 85 90 95
 cca cag ttt gaa atc gag aat gcc aac gtg gaa tgt ctg gca ctc ctc 336
 Pro Gln Phe Glu Ile Glu Asn Ala Asn Val Glu Cys Leu Ala Leu Leu
 100 105 110
 ttg cat gag tat tat gga gca tcg ctg gcc tgg gct acg tcc att agc 384
 Leu His Glu Tyr Tyr Gly Ala Ser Leu Ala Trp Ala Thr Ser Ile Ser
 115 120 125
 atc ttt gaa aac atg aac ctg cga acc ctc tcg gag aag gca cgg gat 432
 Ile Phe Glu Asn Met Asn Leu Arg Thr Leu Ser Glu Lys Ala Arg Asp
 130 135 140
 gac atg aaa aga cat tca cca gaa gat cct gaa gca acg atg act gac 480
 Asp Met Lys Arg His Ser Pro Glu Asp Pro Glu Ala Thr Met Thr Asp
 145 150 155 160
 caa gga cca agc aag gaa aaa gtg cca gga att tca caa gct gtg caa 528
 Gln Gly Pro Ser Lys Glu Lys Val Pro Gly Ile Ser Gln Ala Val Gln
 165 170 175
 caa gat agt gcc aca gct gca gag aca aaa gaa cag gaa att tca caa 576
 Gln Asp Ser Ala Thr Ala Ala Glu Thr Lys Glu Gln Glu Ile Ser Gln
 180 185 190

gct atg gaa caa gaa ggt gcc aca gca gca gag aca gaa gaa caa gaa Ala Met Glu Gln Glu Gly Ala Thr Ala Ala Glu Thr Glu Glu Gln Glu 195 200 205	624
att tca caa gct atg gaa caa gaa ggt gcc aca gca gca gag aca gaa Ile Ser Gln Ala Met Glu Gln Glu Gly Ala Thr Ala Ala Glu Thr Glu 210 215 220	672
gaa caa gga cat gga ggt gac aca tgg gac tac aag agt cac gtg atg Glu Gln Gly His Gly Gly Asp Thr Trp Asp Tyr Lys Ser His Val Met 225 230 235 240	720
acc aaa ttc gct gag gag gag gat gta cgt cgt agt ttt gaa aac act Thr Lys Phe Ala Glu Glu Glu Asp Val Arg Arg Ser Phe Glu Asn Thr 245 250 255	768
gct gct gac tgg ccg gaa atg caa acg ttg gct ggt gct ttt gat tca Ala Ala Asp Trp Pro Glu Met Gln Thr Leu Ala Gly Ala Phe Asp Ser 260 265 270	816
gac cgg tgg ggc ttc cgg cct cgc acg gtg gtt ctg cac gga aag tca Asp Arg Trp Gly Phe Arg Pro Arg Thr Val Val Leu His Gly Lys Ser 275 280 285	864
gga att ggg aaa tcg gct cta gcc aga agg atc gtg ctg tgc tgg gcg Gly Ile Gly Lys Ser Ala Leu Ala Arg Arg Ile Val Leu Cys Trp Ala 290 295 300	912
caa ggt gga ctc tac cag gga atg ttc tcc tac gtc ttc ttc ctc ccc Gln Gly Gly Leu Tyr Gln Gly Met Phe Ser Tyr Val Phe Phe Leu Pro 305 310 315 320	960
gtt aga gag atg cag cgg aag aag gag agc agt gtc aca gag ttc atc Val Arg Glu Met Gln Arg Lys Lys Glu Ser Ser Val Thr Glu Phe Ile 325 330 335	1008
tcc agg gag tgg cca gac tcc cag gct ccg gtg acg gag atc atg tcc Ser Arg Glu Trp Pro Asp Ser Gln Ala Pro Val Thr Glu Ile Met Ser 340 345 350	1056
cga cca gaa agg ctg ttg ttc atc att gac ggt ttc gat gac ctg ggc Arg Pro Glu Arg Leu Leu Phe Ile Ile Asp Gly Phe Asp Asp Leu Gly 355 360 365	1104
tct gtc ctc aac aat gac aca aag ctc tgc aaa gac tgg gct gag aag Ser Val Leu Asn Asn Asp Thr Lys Leu Cys Lys Asp Trp Ala Glu Lys 370 375 380	1152
cag cct ccg ttc acc ctc ata cgc agt ctg ctg agg aag gtc ctg ctc Gln Pro Pro Phe Thr Leu Ile Arg Ser Leu Leu Arg Lys Val Leu Leu 385 390 395 400	1200
cct gag tcc ttc ctg atc gtc acc gtc aga gac gtg ggc aca gag aag Pro Glu Ser Phe Leu Ile Val Thr Val Arg Asp Val Gly Thr Glu Lys 405 410 415	1248
ctc aag tca gag gtc gtg tct ccc cgt tac ctg tta gtt aga gga atc Leu Lys Ser Glu Val Val Ser Pro Arg Tyr Leu Leu Val Arg Gly Ile 420 425 430	1296

tcc ggg gaa caa aga atc cac ttg ctc ctt gag cgc ggg att ggt gag Ser Gly Glu Gln Arg Ile His Leu Leu Leu Glu Arg Gly Ile Gly Glu 435 440 445	1344
cat cag aag aca caa ggg ttg cgt gcg atc atc aac aac cgt gag ctg His Gln Lys Thr Gln Gly Leu Arg Ala Ile Ile Asn Asn Arg Glu Leu 450 455 460	1392
ctc gac cag tgc cag gtg ccc gcc gtg ggc tct ctc atc tgc gtg gcc Leu Asp Gln Cys Gln Val Pro Ala Val Gly Ser Leu Ile Cys Val Ala 465 470 475 480	1440
ctg cag ctg cag gac gtg gtg ggg gag agc gtc gcc ccc ttc aac caa Leu Gln Leu Gln Asp Val Val Gly Glu Ser Val Ala Pro Phe Asn Gln 485 490 495	1488
acg ctc aca ggc ctg cac gcc gct ttt gcg ttt cat cag ctc acc cct Thr Leu Thr Gly Leu His Ala Ala Phe Ala Phe His Gln Leu Thr Pro 500 505 510	1536
cga ggc gtg gtc cgg cgc tgt ctc aat ctg gag gaa aga gtt gtc ctg Arg Gly Val Val Arg Arg Cys Leu Asn Leu Glu Glu Arg Val Val Leu 515 520 525	1584
aag cgc ttc tgc cgt atg gct gtg gag gga gtg tgg aat agg aag tca Lys Arg Phe Cys Arg Met Ala Val Glu Gly Val Trp Asn Arg Lys Ser 530 535 540	1632
gtg ttt gat ggt gac gac ctc atg gtt caa gga ctc ggg gag tct gag Val Phe Asp Gly Asp Asp Leu Met Val Gln Gly Leu Gly Glu Ser Glu 545 550 555 560	1680
ctc cgt gct ctg ttt cac atg aac atc ctt ctc cca gac agc cac tgt Leu Arg Ala Leu Phe His Met Asn Ile Leu Leu Pro Asp Ser His Cys 565 570 575	1728
gag gag tac tac acc ttc ttc cac ctc agt ctc cag gac ttc tgt gcc Glu Glu Tyr Tyr Thr Phe Phe His Leu Ser Leu Gln Asp Phe Cys Ala 580 585 590	1776
gcc ttg tac tac gtg tta gag ggc ctg gaa atc gag cca gct ctc tgc Ala Leu Tyr Tyr Val Leu Glu Gly Leu Glu Ile Glu Pro Ala Leu Cys 595 600 605	1824
cct ctg tac gtt gag aag aca aag agg tcc atg gag ctt aaa cag gca Pro Leu Tyr Val Glu Lys Thr Lys Arg Ser Met Glu Leu Lys Gln Ala 610 615 620	1872
ggc ttc cat atc cac tcg ctt tgg atg aag cgt ttc ttg ttt ggc ctc Gly Phe His Ile His Ser Leu Trp Met Lys Arg Phe Leu Phe Gly Leu 625 630 635 640	1920
gtg agc gaa gac gta agg agg cca ctg gag gtc ctg ctg ggc tgt ccc Val Ser Glu Asp Val Arg Arg Pro Leu Glu Val Leu Leu Gly Cys Pro 645 650 655	1968
gtt ccc ctg ggg gtg aag cag aag ctt ctg cac tgg gtc tct ctg ttg Val Pro Leu Gly Val Lys Gln Lys Leu Leu His Trp Val Ser Leu Leu 660 665 670	2016

ggt cag cag cct aat gcc acc acc cca gga gac acc ctg gac gcc ttc Gly Gln Gln Pro Asn Ala Thr Thr Pro Gly Asp Thr Leu Asp Ala Phe 675 680 685	2064
cac tgt ctt ttc gag act caa gac aaa gag ttt gtt cgc ttg gca tta His Cys Leu Phe Glu Thr Gln Asp Lys Glu Phe Val Arg Leu Ala Leu 690 695 700	2112
aac agc ttc caa gaa gtg tgg ctt ccg att aac cag aac ctg gac ttg Asn Ser Phe Gln Glu Val Trp Leu Pro Ile Asn Gln Asn Leu Asp Leu 705 710 715 720	2160
ata gca tct tcc ttc tgc ctc cag cac tgt ccg tat ttg cgg aaa att Ile Ala Ser Ser Phe Cys Leu Gln His Cys Pro Tyr Leu Arg Lys Ile 725 730 735	2208
cgg gtg gat gtc aaa ggg atc ttc cca aga gat gag tcc gct gag gca Arg Val Asp Val Lys Gly Ile Phe Pro Arg Asp Glu Ser Ala Glu Ala 740 745 750	2256
tgt cct gtg gtc cct cta tgg atg cgg gat aag acc ctc att gag gag Cys Pro Val Val Pro Leu Trp Met Arg Asp Lys Thr Leu Ile Glu Glu 755 760 765	2304
cag tgg gaa gat ttc tgc tcc atg ctt ggc acc cac cca cac ctg cgg Gln Trp Glu Asp Phe Cys Ser Met Leu Gly Thr His Pro His Leu Arg 770 775 780	2352
cag ctg gac ctg ggc agc agc atc ctg aca gag cgg gcc atg aag acc Gln Leu Asp Leu Gly Ser Ser Ile Leu Thr Glu Arg Ala Met Lys Thr 785 790 795 800	2400
ctg tgt gcc aag ctg agg cat ccc acc tgc aag ata cag acc ctg atg Leu Cys Ala Lys Leu Arg His Pro Thr Cys Lys Ile Gln Thr Leu Met 805 810 815	2448
ttt aga aat gca cag att acc cct ggt gtg caa cac ctc tgg aga atc Phe Arg Asn Ala Gln Ile Thr Pro Gly Val Gln His Leu Trp Arg Ile 820 825 830	2496
gtc atg gcc aac cgt aac cta aga tcc ctc aac ttg gga ggc acc cac Val Met Ala Asn Arg Asn Leu Arg Ser Leu Asn Leu Gly Gly Thr His 835 840 845	2544
ctg aag gaa gag gat gta agg atg gcg tgt gaa gcc tta aaa cac cca Leu Lys Glu Glu Asp Val Arg Met Ala Cys Glu Ala Leu Lys His Pro 850 855 860	2592
aaa tgt ttg ttg gag tct ttg agg ctg gat tgc tgt gga ttg acc cat Lys Cys Leu Leu Glu Ser Leu Arg Leu Asp Cys Cys Gly Leu Thr His 865 870 875 880	2640
gcc tgt tac ctg aag atc tcc caa atc ctt acg acc tcc ccc agc ctg Ala Cys Tyr Leu Lys Ile Ser Gln Ile Leu Thr Thr Ser Pro Ser Leu 885 890 895	2688
aaa tct ctg agc ctg gca gga aac aag gtg aca gac cag gga gta acg Lys Ser Leu Ser Leu Ala Gly Asn Lys Val Thr Asp Gln Gly Val Thr 900 905 910	2736

cct ctc agt gat gcc ttg agg gtc tcc cag tgc gcc ctg cag aag ctg Pro Leu Ser Asp Ala Leu Arg Val Ser Gln Cys Ala Leu Gln Lys Leu 915 920 925	2784
ata ctg gag gac tgt ggc atc aca gcc acg ggt tgc cag agt ctg gcc Ile Leu Glu Asp Cys Gly Ile Thr Ala Thr Gly Cys Gln Ser Leu Ala 930 935 940	2832
tca gcc ctc gtc agc aac cgg agc ttg aca cac ctg tgc cta tcc aac Ser Ala Leu Val Ser Asn Arg Ser Leu Thr His Leu Cys Leu Ser Asn 945 950 955 960	2880
aac agc ctg ggg aac gaa ggt gta aat cta ctg tgt cga tcc atg agg Asn Ser Leu Gly Asn Glu Gly Val Asn Leu Leu Cys Arg Ser Met Arg 965 970 975	2928
ctt ccc cac tgt agt ctg cag agg ctg atg ctg aat cag tgc cac ctg Leu Pro His Cys Ser Leu Gln Arg Leu Met Leu Asn Gln Cys His Leu 980 985 990	2976
gac acg gct ggc tgt ggt tct ctt gca ctt gcg ctt atg ggt aac tca Asp Thr Ala Gly Cys Gly Ser Leu Ala Leu Ala Leu Met Gly Asn Ser 995 1000 1005	3024
tgg ctg acg cac ctg agc ctt agc atg aac cct gtg gaa gac aat Trp Leu Thr His Leu Ser Leu Ser Met Asn Pro Val Glu Asp Asn 1010 1015 1020	3069
ggc gtg aag ctt ctg tgc gag gtc atg aga gaa cca tct tgt cat Gly Val Lys Leu Leu Cys Glu Val Met Arg Glu Pro Ser Cys His 1025 1030 1035	3114
ctc cag gac ctg gag ttg gta aag tgt cat ctc acc gcc gcg tgc Leu Gln Asp Leu Glu Leu Val Lys Cys His Leu Thr Ala Ala Cys 1040 1045 1050	3159
tgt gag agt ctg tcc tgt gtg atc tcg agg agc aga cac ctg aag Cys Glu Ser Leu Ser Cys Val Ile Ser Arg Ser Arg His Leu Lys 1055 1060 1065	3204
agc ctg gat ctc acg gac aat gcc ctg ggt gac ggt ggg gtt gct Ser Leu Asp Leu Thr Asp Asn Ala Leu Gly Asp Gly Gly Val Ala 1070 1075 1080	3249
gcg ctg tgc gag gga ctg aag caa aag aac agt gtt ctg acg aga Ala Leu Cys Glu Gly Leu Lys Gln Lys Asn Ser Val Leu Thr Arg 1085 1090 1095	3294
ctc ggg ttg aag gca tgt gga ctg act tct gat tgc tgt gag gca Leu Gly Leu Lys Ala Cys Gly Leu Thr Ser Asp Cys Cys Glu Ala 1100 1105 1110	3339
ctc tcc ttg gcc ctt tcc tgc aac cgg cat ctg acc agt cta aac Leu Ser Leu Ala Leu Ser Cys Asn Arg His Leu Thr Ser Leu Asn 1115 1120 1125	3384
ctg gtg cag aat aac ttc agt ccc aaa gga atg atg aag ctg tgt Leu Val Gln Asn Asn Phe Ser Pro Lys Gly Met Met Lys Leu Cys 1130 1135 1140	3429

tcg gcc	ttt gcc	tgt ccc	acg tct	aac tta	cag ata	att ggg	ctg	3474
Ser Ala	Phe Ala	Cys Pro	Thr Ser	Asn Leu	Gln Ile	Ile Gly	Leu	
1145			1150			1155		

tgg aaa	tgg cag	tac cct	gtg caa	ata agg	aag ctg	ctg gag	gaa	3519
Trp Lys	Trp Gln	Tyr Pro	Val Gln	Ile Arg	Lys Leu	Leu Glu	Glu	
1160			1165			1170		

gtg cag	cta ctc	aag ccc	cga gtc	gta att	gac ggt	agt tgg	cat	3564
Val Gln	Leu Leu	Lys Pro	Arg Val	Val Ile	Asp Gly	Ser Trp	His	
1175			1180			1185		

tct ttt	gat gaa	gat gac	cgg tac	tgg tgg	aaa aac	tgaagatacg	3610
Ser Phe	Asp Glu	Asp Asp	Arg Tyr	Trp Trp	Lys Asn		
1190			1195		1200		

gaaacctgcc	ccactcacac	ccatctgatg	gaggaacttt	aaacgctgtt	ttctcagagc	3670
aagctatgca	cctgggagtt	ccttctcaaa	gatggagaat	gatttctgat	tctcaciaaag	3730
ccctcaatgg	tagtgattct	tctgtgttca	ctctacgttg	gttactggat	ttgaaggcta	3790
gagaccttca	agtcatagga	ctcagtarct	gtgaaatgtc	cgtcataatct	cagagcatat	3850
agaggggaatt	aaataaacac	aaagcatttg	gaaaaaaaaa	aaaaaaaaaa		3900

<210> 24
 <211> 1200
 <212> PRT
 <213> homo sapiens

<400> 24

Met Lys Val Ala Gly Gly Leu Glu Leu Gly Ala Ala Ala Leu Leu Ser	
1 5 10 15	
Ala Ser Pro Arg Ala Leu Val Thr Leu Ser Thr Gly Pro Thr Cys Ser	
20 25 30	
Ile Leu Pro Lys Asn Pro Leu Phe Pro Gln Asn Leu Ser Ser Gln Pro	
35 40 45	
Cys Ile Lys Met Glu Gly Asp Lys Ser Leu Thr Phe Ser Ser Tyr Gly	
50 55 60	
Leu Gln Trp Cys Leu Tyr Glu Leu Asp Lys Glu Glu Phe Gln Thr Phe	
65 70 75 80	
Lys Glu Leu Leu Lys Lys Lys Ser Ser Glu Ser Thr Thr Cys Ser Ile	
85 90 95	
Pro Gln Phe Glu Ile Glu Asn Ala Asn Val Glu Cys Leu Ala Leu Leu	
100 105 110	
Leu His Glu Tyr Tyr Gly Ala Ser Leu Ala Trp Ala Thr Ser Ile Ser	
115 120 125	
Ile Phe Glu Asn Met Asn Leu Arg Thr Leu Ser Glu Lys Ala Arg Asp	
130 135 140	
Asp Met Lys Arg His Ser Pro Glu Asp Pro Glu Ala Thr Met Thr Asp	
145 150 155 160	
Gln Gly Pro Ser Lys Glu Lys Val Pro Gly Ile Ser Gln Ala Val Gln	
165 170 175	
Gln Asp Ser Ala Thr Ala Ala Glu Thr Lys Glu Gln Glu Ile Ser Gln	
180 185 190	
Ala Met Glu Gln Glu Gly Ala Thr Ala Ala Glu Thr Glu Glu Gln Glu	
195 200 205	
Ile Ser Gln Ala Met Glu Gln Glu Gly Ala Thr Ala Ala Glu Thr Glu	
210 215 220	
Glu Gln Gly His Gly Gly Asp Thr Trp Asp Tyr Lys Ser His Val Met	
225 230 235 240	

Thr	Lys	Phe	Ala	Glu	Glu	Glu	Asp	Val	Arg	Arg	Ser	Phe	Glu	Asn	Thr	245	250	255
Ala	Ala	Asp	Trp	Pro	Glu	Met	Gln	Thr	Leu	Ala	Gly	Ala	Phe	Asp	Ser	260	265	270
Asp	Arg	Trp	Gly	Phe	Arg	Pro	Arg	Thr	Val	Val	Leu	His	Gly	Lys	Ser	275	280	285
Gly	Ile	Gly	Lys	Ser	Ala	Leu	Ala	Arg	Arg	Ile	Val	Leu	Cys	Trp	Ala	290	295	300
Gln	Gly	Gly	Leu	Tyr	Gln	Gly	Met	Phe	Ser	Tyr	Val	Phe	Phe	Leu	Pro	305	310	315
Val	Arg	Glu	Met	Gln	Arg	Lys	Lys	Glu	Ser	Ser	Val	Thr	Glu	Phe	Ile	325	330	335
Ser	Arg	Glu	Trp	Pro	Asp	Ser	Gln	Ala	Pro	Val	Thr	Glu	Ile	Met	Ser	340	345	350
Arg	Pro	Glu	Arg	Leu	Leu	Phe	Ile	Ile	Asp	Gly	Phe	Asp	Asp	Leu	Gly	355	360	365
Ser	Val	Leu	Asn	Asn	Asp	Thr	Lys	Leu	Cys	Lys	Asp	Trp	Ala	Glu	Lys	370	375	380
Gln	Pro	Pro	Phe	Thr	Leu	Ile	Arg	Ser	Leu	Leu	Arg	Lys	Val	Leu	Leu	385	390	395
Pro	Glu	Ser	Phe	Leu	Ile	Val	Thr	Val	Arg	Asp	Val	Gly	Thr	Glu	Lys	405	410	415
Leu	Lys	Ser	Glu	Val	Val	Ser	Pro	Arg	Tyr	Leu	Leu	Val	Arg	Gly	Ile	420	425	430
Ser	Gly	Glu	Gln	Arg	Ile	His	Leu	Leu	Leu	Glu	Arg	Gly	Ile	Gly	Glu	435	440	445
His	Gln	Lys	Thr	Gln	Gly	Leu	Arg	Ala	Ile	Ile	Asn	Asn	Arg	Glu	Leu	450	455	460
Leu	Asp	Gln	Cys	Gln	Val	Pro	Ala	Val	Gly	Ser	Leu	Ile	Cys	Val	Ala	465	470	475
Leu	Gln	Leu	Gln	Asp	Val	Val	Gly	Glu	Ser	Val	Ala	Pro	Phe	Asn	Gln	485	490	495
Thr	Leu	Thr	Gly	Leu	His	Ala	Ala	Phe	Ala	Phe	His	Gln	Leu	Thr	Pro	500	505	510
Arg	Gly	Val	Val	Arg	Arg	Cys	Leu	Asn	Leu	Glu	Glu	Arg	Val	Val	Leu	515	520	525
Lys	Arg	Phe	Cys	Arg	Met	Ala	Val	Glu	Gly	Val	Trp	Asn	Arg	Lys	Ser	530	535	540
Val	Phe	Asp	Gly	Asp	Asp	Leu	Met	Val	Gln	Gly	Leu	Gly	Glu	Ser	Glu	545	550	555
Leu	Arg	Ala	Leu	Phe	His	Met	Asn	Ile	Leu	Leu	Pro	Asp	Ser	His	Cys	565	570	575
Glu	Glu	Tyr	Tyr	Thr	Phe	Phe	His	Leu	Ser	Leu	Gln	Asp	Phe	Cys	Ala	580	585	590
Ala	Leu	Tyr	Tyr	Val	Leu	Glu	Gly	Leu	Glu	Ile	Glu	Pro	Ala	Leu	Cys	595	600	605
Pro	Leu	Tyr	Val	Glu	Lys	Thr	Lys	Arg	Ser	Met	Glu	Leu	Lys	Gln	Ala	610	615	620
Gly	Phe	His	Ile	His	Ser	Leu	Trp	Met	Lys	Arg	Phe	Leu	Phe	Gly	Leu	625	630	635
Val	Ser	Glu	Asp	Val	Arg	Arg	Pro	Leu	Glu	Val	Leu	Leu	Gly	Cys	Pro	645	650	655
Val	Pro	Leu	Gly	Val	Lys	Gln	Lys	Leu	Leu	His	Trp	Val	Ser	Leu	Leu	660	665	670
Gly	Gln	Gln	Pro	Asn	Ala	Thr	Thr	Pro	Gly	Asp	Thr	Leu	Asp	Ala	Phe	675	680	685
His	Cys	Leu	Phe	Glu	Thr	Gln	Asp	Lys	Glu	Phe	Val	Arg	Leu	Ala	Leu	690	695	700

Asn Ser Phe Gln Glu Val Trp Leu Pro Ile Asn Gln Asn Leu Asp Leu
 705 710 715 720
 Ile Ala Ser Ser Phe Cys Leu Gln His Cys Pro Tyr Leu Arg Lys Ile
 725 730 735
 Arg Val Asp Val Lys Gly Ile Phe Pro Arg Asp Glu Ser Ala Glu Ala
 740 745 750
 Cys Pro Val Val Pro Leu Trp Met Arg Asp Lys Thr Leu Ile Glu Glu
 755 760 765
 Gln Trp Glu Asp Phe Cys Ser Met Leu Gly Thr His Pro His Leu Arg
 770 775 780
 Gln Leu Asp Leu Gly Ser Ser Ile Leu Thr Glu Arg Ala Met Lys Thr
 785 790 795 800
 Leu Cys Ala Lys Leu Arg His Pro Thr Cys Lys Ile Gln Thr Leu Met
 805 810 815
 Phe Arg Asn Ala Gln Ile Thr Pro Gly Val Gln His Leu Trp Arg Ile
 820 825 830
 Val Met Ala Asn Arg Asn Leu Arg Ser Leu Asn Leu Gly Gly Thr His
 835 840 845
 Leu Lys Glu Glu Asp Val Arg Met Ala Cys Glu Ala Leu Lys His Pro
 850 855 860
 Lys Cys Leu Leu Glu Ser Leu Arg Leu Asp Cys Cys Gly Leu Thr His
 865 870 875 880
 Ala Cys Tyr Leu Lys Ile Ser Gln Ile Leu Thr Thr Ser Pro Ser Leu
 885 890 895
 Lys Ser Leu Ser Leu Ala Gly Asn Lys Val Thr Asp Gln Gly Val Thr
 900 905 910
 Pro Leu Ser Asp Ala Leu Arg Val Ser Gln Cys Ala Leu Gln Lys Leu
 915 920 925
 Ile Leu Glu Asp Cys Gly Ile Thr Ala Thr Gly Cys Gln Ser Leu Ala
 930 935 940
 Ser Ala Leu Val Ser Asn Arg Ser Leu Thr His Leu Cys Leu Ser Asn
 945 950 955 960
 Asn Ser Leu Gly Asn Glu Gly Val Asn Leu Leu Cys Arg Ser Met Arg
 965 970 975
 Leu Pro His Cys Ser Leu Gln Arg Leu Met Leu Asn Gln Cys His Leu
 980 985 990
 Asp Thr Ala Gly Cys Gly Ser Leu Ala Leu Ala Leu Met Gly Asn Ser
 995 1000 1005
 Trp Leu Thr His Leu Ser Leu Ser Met Asn Pro Val Glu Asp Asn
 1010 1015 1020
 Gly Val Lys Leu Leu Cys Glu Val Met Arg Glu Pro Ser Cys His
 1025 1030 1035
 Leu Gln Asp Leu Glu Leu Val Lys Cys His Leu Thr Ala Ala Cys
 1040 1045 1050
 Cys Glu Ser Leu Ser Cys Val Ile Ser Arg Ser Arg His Leu Lys
 1055 1060 1065
 Ser Leu Asp Leu Thr Asp Asn Ala Leu Gly Asp Gly Val Ala
 1070 1075 1080
 Ala Leu Cys Glu Gly Leu Lys Gln Lys Asn Ser Val Leu Thr Arg
 1085 1090 1095
 Leu Gly Leu Lys Ala Cys Gly Leu Thr Ser Asp Cys Cys Glu Ala
 1100 1105 1110
 Leu Ser Leu Ala Leu Ser Cys Asn Arg His Leu Thr Ser Leu Asn
 1115 1120 1125
 Leu Val Gln Asn Asn Phe Ser Pro Lys Gly Met Met Lys Leu Cys
 1130 1135 1140
 Ser Ala Phe Ala Cys Pro Thr Ser Asn Leu Gln Ile Ile Gly Leu
 1145 1150 1155

Trp	Lys	Trp	Gln	Tyr	Pro	Val	Gln	Ile	Arg	Lys	Leu	Leu	Glu	Glu
1160						1165					1170			
Val	Gln	Leu	Leu	Lys	Pro	Arg	Val	Val	Ile	Asp	Gly	Ser	Trp	His
1175						1180					1185			
Ser	Phe	Asp	Glu	Asp	Asp	Arg	Tyr	Trp	Trp	Lys	Asn			
1190						1195					1200			

63198-1404

-52-

CLAIMS:

1. An isolated human MATER protein comprising an amino acid sequence at least 95% identical to full-length SEQ ID NO: 24, wherein the human MATER protein complements a *Mater* null phenotype.
- 5 2. The isolated human MATER protein of claim 1, wherein the protein is expressed in oocyte cytoplasm.
3. The isolated human MATER protein of claim 1 or claim 2, comprising the amino acid sequence as set forth in SEQ ID NO: 24.
4. An isolated nucleic acid molecule encoding the isolated human MATER
10 protein according to claim 1, comprising a nucleotide sequence at least 95% identical to full-length SEQ ID NO: 23.
5. The isolated nucleic acid molecule of claim 4, comprising a nucleotide sequence as set forth in SEQ ID NO: 23.
6. A recombinant nucleic acid molecule comprising a promoter sequence
15 operably linked to the isolated nucleic acid molecule according to claim 4.
7. A cell transformed with the recombinant nucleic acid molecule according to claim 6.
8. A method of detecting a *Mater* nucleic acid in a human subject, comprising detecting a labeled nucleotide probe that hybridizes to a *Mater* nucleotide
20 sequence at least 95% identical to full-length SEQ ID NO: 23.
9. A method for determining whether a sample from a human subject has circulating autoantibodies that recognize an epitope of a MATER protein, comprising:

reacting a sample from the human subject with a human MATER protein comprising a MATER amino acid sequence at least 95% identical to full-
25 length SEQ ID NO: 24 to form a MATER:autoantibody complex, wherein detection of

63198-1404

-53-

the MATER:autoantibody complex indicates the presence of autoantibodies in the sample.

10. A method for detecting Mater nucleic acid expression in a human subject, comprising:

5 reacting at least one human Mater nucleic acid contained in a clinical sample from the human subject, wherein the human Mater nucleic acid is a MATER encoding nucleic acid comprising a Mater nucleotide sequence at least 95% identical to full-length SEQ ID NO: 23 with a Mater oligonucleotide, wherein the Mater oligonucleotide comprises at least 23 contiguous nucleotides having at least 95%
10 sequence identity to the corresponding sequence within SEQ ID NO: 23 to form a *Mater:Mater* oligonucleotide complex; and

detecting the *Mater:Mater* oligonucleotide complex, thereby detecting Mater nucleic acid expression in the subject.

11. A method for detecting MATER protein expression in a human subject,
15 comprising:

reacting at least one human MATER protein contained in a clinical sample from the human subject, wherein the human MATER protein is a MATER protein comprising an amino acid sequence at least 95% identical to full-length SEQ ID NO: 24 with a MATER protein specific antibody or fragment thereof that binds
20 specifically to an epitope within an amino acid sequence at least 95% identical to full-length SEQ ID NO: 24 to form a MATER:MATER-specific antibody complex; and

detecting the MATER:MATER-specific antibody complex, thereby detecting MATER protein expression in the subject.

12. The method of claim 10 or claim 11, wherein the sample comprises a
25 reproductive cell.

63198-1404

-54-

13. The method of claim 8, further comprising *in vitro* amplifying a *Mater* nucleic acid prior to detecting the *Mater* nucleic acid.
14. The method of claim 8, wherein the *Mater* nucleic acid is *in vitro* amplified using at least one oligonucleotide primer comprising at least a portion of a
5 MATER protein encoding sequence at least 95% identical to the corresponding sequence within SEQ ID NO: 23.
15. The method of claim 14, wherein at least one oligonucleotide primer comprises at least 23 contiguous nucleotides having at least 95% sequence identity to the corresponding sequence within SEQ ID NO: 23.
- 10 16. An oligonucleotide primer for use in the method of claim 15.
17. A recombinant DNA vector comprising the isolated nucleic acid according to claim 4 or 5.
18. A recombinant nucleic acid molecule comprising a promoter sequence operably linked to the isolated nucleic acid of claim 4 or 5.
- 15 19. The recombinant nucleic acid molecule according to claim 18, wherein the nucleic acid sequence is in antisense orientation relative to the promoter sequence.
20. A cell transformed with the recombinant nucleic acid molecule according to claim 18 or 19.
- 20 21. The method of claim 10, wherein the *Mater* oligonucleotide is a labeled nucleotide probe comprising at least 23 contiguous nucleotides having at least 95% sequence identity to the corresponding sequence within SEQ ID NO: 23.
22. The method of claim 11, wherein the MATER:MATER-specific antibody complexes are detected by Western blot assay.

63198-1404

-55-

23. The method of claim 11, wherein the MATER:MATER-specific antibody complexes are detected by ELISA.

24. The method of claim 11, wherein the MATER-specific antibody is generated against a C-terminal peptide of a MATER protein.

5 25. A kit for detecting MATER protein in a subject using the method of claim 11, comprising a MATER protein specific antibody or fragment thereof that binds specifically to an epitope within an amino acid sequence at least 95% identical to full-length SEQ ID NO: 24 together with instructions for detecting a MATER protein.

10 26. The kit of claim 25, further comprising a means for detecting binding of the MATER protein specific antibody or fragment thereof to a MATER polypeptide.

27. A kit for detecting a MATER nucleic acid in a sample from a human subject, comprising:

(a) a first container containing an oligonucleotide comprising at least 23
15 contiguous nucleotides having at least 95% sequence identity to the corresponding sequence within SEQ ID NO: 23, which specifically hybridizes with a *Mater* nucleic acid at least 95% identical to full length SEQ ID NO: 23; and

(b) a second container containing a fluorescent labeled nucleic acid probe that is fully complementary to the oligonucleotide.

20 28. The kit of claim 27, wherein the fluorescent labeled nucleic acid probe has a length of between 5 and 500 nucleotides.




29. The isolated nucleic acid molecule according to claim 4, wherein the molecule hybridizes with a nucleic acid probe comprising the complement of the sequence shown in SEQ ID NO: 23 under wash conditions of 55°C,
25 0.2 x SSC and 0.1% SDS.

63198-1404



-56-

30. The isolated nucleic acid molecule according to claim 4, wherein the molecule hybridizes with a nucleic acid probe comprising the complement of the sequence shown in SEQ ID NO: 23 under wash conditions of 50°C, 2 x SSC and 0.1 % SDS.

1/9




2/9






3/9





FIG. 3a - l



4/9




5/9




6/9

FIG. 6



7/9



8/9

FIG. 8a




FIG. 8b




FIG. 9

9/9

Human: 1 MKVAGGLELGAA 12
 13 ALLSASPRALVTLSTGPTCSILPKNLFPPQNLSSQPCIKMEGDKSLTFSSYGLQWCLYEL 72
 73 DKEEFQTFKELKKKSSESTTCSTIPQFEIENANVECLALLHEYYGASLAWATSI SIFEN 132

Mouse: 1 MGPPEKESKAI 11

Human: 133 MNLRTLSEKARDDMKRHSPE-DPEATMTDQGSPSKE-----KVPGISQ 173
 + R L E+ + + K SPE D ++ DQGP +E K G+ +
 Mouse: 12 LKARGLEEEQKSERKMTSPENDSKSIQKQDQGPQEQTSESTMGPPEKDSKAILKARGLEE 71

Human: 174 AVQQDS-----ATAAETKEQEISQAMEQEGATAAETEEQEISQAMEQEGATAAET 223
 + +S A ++ +E+ QA E++ T+ E + + I + E +ET
 Mouse: 72 EQKSESTMSPSENVSRAILKDSGSEEVEQASERK-MTSPENDSKSIQKQDQGPQEQTSET 130

Human: 224 -----EEQGHGGDTWDYKSHVMTKFAEEEDVRRSFENTAADWPQMOTLAGAFDS 272
 ++ +GGD DYK+HV+ KF D+ D PEM+ L+ AF
 Mouse: 131 LQSKEEDEVTEADKNDNGDLQDYKAHVIAKFDTSVDLHY-----DSPMKLLSDAFKP 183

Human: 273 DRWGRFRPTVVLHGKSGIGKSALARRIVLCWAQGGLYQGMFSYVFFLPVREMQRKKESSV 332
 + F+P T++LHG+ G+GKSALAR IVL WAQG L+Q M S+V F VRE++ ++SS+
 Mouse: 184 YQKTFQPHITILLHGRPGVCKSALARISIVLGWAQGKLFQKM-SFVIFSVREIKWTEKSSL 242

Human: 333 TEFISREWPDQAPVTEIMSRPERLLFIIDGFDDLGSVL--NNDTKLCKDWAEEKPPFTLI 391
 + I++E PDS VT+IMS+PERLLF+IDG DD+ SVL ++D L +DW ++QP + L+
 Mouse: 243 AQLIAKECPDSWDLVTKIMSQPERLLFVIDGLDMDSVLQHDDMTLSRDWKDEQPIIYIM 302

Human: 392 RSLLRKVLPPESFLIVTVRDVGTEKLKSEVVSRYLLVRGISGEQRIHLLERIGEHQK 451
 SLLRK LLP+SFLI+T R+ G EKLKS VVSP Y+LV G+S +R L+LE E +
 Mouse: 303 YSLLRKALLPQSFLIITRNTGLEKLKSMVVSPLYLLVEGLSASRRSRLVLENISNESDR 362

Human: 452 TQGLRAIINNRELLDQCQVPVAVGSLICVALQLQDVVGESVAPFNQTLTGLHAAFAFHQLT 511
 Q ++I N +L DQCQ P+V SL+C ALQLQ +G+ QTLTGL+A FHQLT
 Mouse: 363 IQVFHSLIENHQLFDQCQAPSVCSLVCEALQLQKKGKRCCTLPCQTLTGLYATLVFHHQLT 422

Human: 512 PRGVVRRCLNLEERVVLKRFCRMAVEGVWNRKSVFDGDDLMVQCLGESELRALFHMNILL 571
 + + L+ EE++ L C MA RGVW +SVF DDL L ESE+ ALFHMNILL
 Mouse: 423 LKRPSQSALSQEEQITLVGLCMMAEGVWMTMSRVFYDDDLKNYSLKSEILALFHMNILL 482

Human: 572 PDSH-CEEYYTFFHLSLQDFCAALYVLEGLEI-EPALCPLYVEKTKRSMELKQAGFHIH 629
 H E+ Y F HLSLQDF AALYVLEGLE C ++E + ME+K+
 Mouse: 483 QVGHNSEQCYVFSHLSLQDFFAALYVLEGLEEWNQHFC--FIENQRSIMEVKKTD-DTR 539

Human: 630 SLWMKRFLFGLVSEDVRRPLEVLLGCPVPLGVKQKLLHWVSSLGQQPNATTPGDTLDFAH 689
 L MKRFLFGL+++D+ + LEVL PV V+QKL HWVSL+ QQ N T+P DTLDFA+
 Mouse: 540 LLGMRFLFGLMNKDILKTLEVLFEYPVPTVEQKLQHWVSLIAQQVNGTSPMDTLDFAF 599

Human: 690 CLFETQDKEFVRLALNSFQEVVLPINQNLDIASSFCLQHCFYLRKIRVDVKGIFPRDES 749
 CLFE+QD+EFV AL FQEVWL INQ +DL SS+CL+HC L+ IRVD++ + D +
 Mouse: 600 CLFESQDEEFVGGALKRFQEVWLLINQKMDLKVSSYCLKHCQNLKAIKIRVDIRDLSDVNT 659

Human: 750 AEACPVVPLWMRD-KTLLIEQWEDFCSMLGTHPHLRQLDLGSSILTERAMKTLCAKLHP 808
 E CPVV + K L+ E W +FCS+LG+ +L++LDIG SIL++RAMK LC +LR+
 Mouse: 660 LELCPVVTVQETQCKPLMEWGNFCVSLGSLRNLKELDLGDSILSQRAMKILCLELRNQ 719

Human: 809 TCKIQTLMFRNAQITPGVQHLWRIVMANRNLRLSLNLGGTHLKEEDVRMACEALKHPKCLL 868
 +C+IQ L F++A++ G++HLW+++ +N+NL+ LNLG T +K++D+++ACEALKHPKC +
 Mouse: 720 SCRIQKLTFKSAEVVSGCLKHLWKLLFSNQNLKYLNLGNTPMKDDDMKLACEALKHPKCSV 779

Human: 869 ESLRLDCCGLTHACYLKISQILTTSPSLKSLSLAGNKVTDQGVTPLSDALRVSCALQKL 928
 E+LRLD C LT Y IS +L ++ LK LSLA N+V + + L +AL S C LQKL
 Mouse: 780 ETLRLDSCELTIIGYEMISTLLISTRLKCLSLAKNRVGVKSMISLGNALSSSMCLLQKL 839

Human: 929 ILEDGCTITATGCQSLASALVSNRSLTHLCLSNNSLGNLGNVLLCRSMRPLPHCSLQRLMLN 988
 IL++CG+T C L SAL SN++LTHLCLSNNSLG EGV LC+ +R P C+LQRL+LN
 Mouse: 840 ILDNCGLTPASCHLLVSALFSNQNLTHLCLSNNSLGTGEGVQQLCQFLRNPECALQRLILN 899

Human: 989 QCHLDTAGCGLALALMGNSWLTHLSLSMNPVEDNGVKLLCEVMREPSCHLQDLELVKCH 1048
 C++ G LA+ L N+ LTHLSL+MNPV D +KLCE ++EP+C+LQ+LELV C
 Mouse: 900 HCNIVDDAYGFLAMRLANNTKLTHLSLTMPVGDGAMKLLCEALKEPTCYLQLELVDCQ 959

Human: 1049 LTAACCESLSCVISRSRHLKSLDLTDNALGDGGVAALCEGLKQKNSVLTRELGLKACGLTS 1108
 LT CCE L+C+I+ ++HLKSLDL +NALGD GV LCEGLKQ +S L RLGL AC LTS
 Mouse: 960 LTQCCEDLACMITTTKHLKSLDLGNALGDKGVITLCEGLKQSSSLRLRLGLGACKLTS 1019

Human: 1109 DCEALSLALSNCNRHLTSLNLVQNNFSPKGMMLKCSAFACPTSNLQIIGLWKWQYFVQIR 1168
 +CCEALSLA+SCN HL SLNLV+N+FS GM+KLCSAF CP SNL IIGLWK +Y ++R
 Mouse: 1020 NCCEALSLAISCNPHLNSLNLVKNDFSTSGMLKLCSAFQCPVSNLGIIGLWKQYFVARV 1079

Human: 1169 KLEEVQLLKPRVVIDGSWHSFDEDDRYWKN 1200
 + LEEV+ +KP VVIDG W++ DEDDR WKN
 Mouse: 1080 RQLEEVFVKPHVVIDGDWYASDEDDRNWKN 1111