

(19) **DANMARK**

(10) **DK/EP 3342879 T3**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 12 Q 1/6886 (2018.01)**
- (45) Oversættelsen bekendtgjort den: **2019-09-23**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-09-11**
- (86) Europæisk ansøgning nr.: **17211034.8**
- (86) Europæisk indleveringsdag: **2017-12-29**
- (87) Den europæiske ansøgnings publiceringsdag: **2018-07-04**
- (30) Prioritet: **2016-12-30 US 201662440883 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **ONCOLOGY VENTURE A/S, Venlighedsvej 1, 2970 Hørsholm, Danmark**
- (72) Opfinder: **Knudsen, Steen, c/o Medical Prognosis Institute Inc., 9977 N 90th Street Suite 175, Scottsdale, AZ 85258, USA**
- (74) Fuldmægtig i Danmark: **Orsnes Patent ApS, Forskerparken 10, 5230 Odense M, Danmark**
- (54) Benævnelse: **FREMANGSMÅDER TIL FORUDSIGELSE AF LÆGEMIDDELMODTAGELIGHED HOS CANCERPATIENTER**
- (56) Fremdragne publikationer:
EP-A1- 2 123 258
WO-A1-2011/032563
WO-A2-2011/098578
US-A1- 2014 294 730
ARIENTI C ET AL: "Activity of lipoplatin in tumor and in normal cells in vitro", ANTI-CANCER DRUGS 200811 GB, vol. 19, no. 10, November 2008 (2008-11), pages 983-990, XP009504980, ISSN: 0959-4973
N. CASAGRANDE ET AL: "Preclinical Activity of the Liposomal Cisplatin Lipoplatin in Ovarian Cancer", CLINICAL CANCER RESEARCH, vol. 20, no. 21, 17 September 2014 (2014-09-17), pages 5496-5506, XP055469342, US ISSN: 1078-0432, DOI: 10.1158/1078-0432.CCR-14-0713
Oncology Venture ET AL: "Press release issued by Oncology Venture Sweden AB Oncology Venture presents LiPlaCis on AACR in New Orleans", , 4 March 2016 (2016-03-04), XP055469349, Retrieved from the Internet: URL: <http://www.oncologyventure.com/wp-content/uploads/2016/03/Press-release-OV-LiPla-Cis-Poster-Presentation-AACR-2016.pdf> [retrieved on 2018-04-23]
DE JONGE M J A ET AL: "Early cessation of the clinical development of LiPlaCis, a liposomal cisplatin formulation", EUROPEAN JOURNAL OF CANCER, ELSEVIER, AMSTERDAM, NL, vol. 46, no. 16, 1 November 2010 (2010-11-01), pages 3016-3021, XP027431952, ISSN: 0959-8049, DOI: 10.1016/J.EJCA.2010.07.015 [retrieved on 2010-08-26]
TENI BOULIKAS: "Clinical overview on Lipoplatin (TM) : a successful liposomal formulation of cisplatin", EXPERT OPINION ON INVESTIGATIONAL DRUGS, vol. 18, no. 8, 15 July 2009 (2009-07-15), pages 1197-1218, XP055256539, UK ISSN: 1354-3784, DOI: 10.1517/13543780903114168
MAMOT CHRISTOPH ET AL: "Liposome-based approaches to overcome anticancer drug resistance.", DRUG

Fortsættes ...

DESCRIPTION

FIELD OF THE INVENTION

[0001] The invention pertains to methods of using biomarkers in cancer patients to predict responsiveness of the cancer patients to treatment.

BACKGROUND

[0002] DNA microarrays can be used to measure gene expression in tumor samples from patients and to facilitate diagnosis. Gene expression can reveal the presence of cancer in a patient in addition to the type, stage, and origin. Gene expression may even have a role in predicting the efficacy of cancer therapies. In recent decades, the National Cancer Institute (NCI) has tested cancer therapeutics for their effect in limiting the growth of 60 human cancer cell lines. The NCI has also measured gene expression in those 60 cancer cell lines using DNA microarrays. Various studies have explored the relationship between gene expression and therapeutic effect using the NCI datasets.

[0003] During cancer treatment, critical time is often lost due to a trial and error approach to finding an effective therapy. In addition, cancer cells often develop resistance to a previously effective therapy. In such situations, patient outcome would be greatly improved by early detection of such resistance to therapy.

[0004] Thus, there exists a need in the art for methods, devices, and kits that can predict the responsiveness of cancer patients to a medical treatment for cancer.

SUMMARY OF THE INVENTION

[0005] The invention features methods for detecting a level of one or more biomarkers (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 1-4, such as COL5A2 (SEQ ID NO: 73 or 211)) in a patient having cancer, such as a patient having cancer (e.g., a human) that is resistant to one or more cancer therapies other than a secretory phospholipase A₂ (sPLA₂) hydrolysable, cisplatin-containing liposome (e.g., a patient having breast cancer that is resistant to one or more cancer therapies other than the liposome), and for determining responsiveness of a cancer patient (e.g., a patient having breast cancer) to treatment with the liposome. The invention also features methods of treating cancer in a patient (e.g., a patient having breast cancer or a treatment resistant form thereof) that include administering the sPLA₂ hydrolysable, cisplatin-containing liposome to the patient, in which the patient is or has been determined to be responsive to the liposome according to the diagnostic methods described herein. Platinum comprising liposomes have been used previously in the state of the art. Casagrande et al., Clinical Cancer Research, volume 20, no. 21 from September 2014 have been teaching an alternative cisplatin comprising liposome compound that was tested on ovarian cancer. It was found that certain expression markers were differentially expressed in parallel and correlated to treatment success. For example CASPASE-9, -8, -3 were induced / activated, whereas Bcl-2 was downregulated. Bax again was upregulated. Furthermore it was found that EGFR expression was down-regulated, c.f. the RESULTS section. The system is however not based on the enzyme sPLA₂.

[0006] Exemplary types of cancer that can be diagnosed or treated with the methods include, e.g., breast cancer, prostate cancer, ovarian cancer (e.g., ovarian adenocarcinoma or embryonal carcinoma), liver cancer (e.g., HCC or hepatoma), myeloma (e.g., multiple myeloma), colorectal cancer (e.g., colon cancer and rectal cancer), leukemia (e.g., acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, and chronic leukemia), myelodysplastic syndrome, lymphoma (e.g., diffuse large B-cell lymphoma, cutaneous T-cell lymphoma, peripheral T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Waldenstrom's macroglobulinemia, and lymphocytic lymphoma), cervical cancer, esophageal cancer, melanoma, glioma (e.g., oligodendroglioma), pancreatic cancer (e.g., adenocarcinoma, signet ring cell carcinoma, hepatoid carcinoma, colloid carcinoma, islet cell carcinoma, and pancreatic neuroendocrine carcinoma), gastrointestinal stromal tumor, sarcoma (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, leiomyosarcoma, Ewing's sarcoma, and rhabdomyosarcoma), breast

cancer (e.g., medullary carcinoma), ER-positive cancer, bladder cancer, head and neck cancer (e.g., squamous cell carcinoma of the head and neck), lung cancer (e.g., non-small cell lung carcinoma, large cell carcinoma, bronchogenic carcinoma, and papillary adenocarcinoma), metastatic cancer, oral cavity cancer, uterine cancer, testicular cancer (e.g., seminoma and embryonal carcinoma), skin cancer (e.g., squamous cell carcinoma, and basal cell carcinoma), thyroid cancer (e.g., papillary carcinoma and medullary carcinoma), brain cancer (e.g., astrocytoma and craniopharyngioma), stomach cancer, intra-epithelial cancer, bone cancer, biliary tract cancer, eye cancer, larynx cancer, kidney cancer (e.g., renal cell carcinoma and Wilms tumor), gastric cancer, blastoma (e.g., nephroblastoma, medulloblastoma, hemangioblastoma, neuroblastoma, and retinoblastoma), polycythemia vera, chordoma, synovioma, mesothelioma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, cystadenocarcinoma, bile duct carcinoma, choriocarcinoma, epithelial carcinoma, ependymoma, pinealoma, acoustic neuroma, schwannoma, meningioma, pituitary adenoma, nerve sheath tumor, cancer of the small intestine, cancer of the endocrine system, cancer of the penis, cancer of the urethra, cutaneous or intraocular melanoma, a gynecologic tumor, solid tumors of childhood, or neoplasms of the central nervous system. For example, the cancer may be a solid tumor (e.g., breast cancer) or a hematological cancer (e.g., cancer of the blood, such as lymphoma (e.g., acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), or Hodgkin's lymphoma)).

[0007] A first aspect of the invention features a method for detecting a level of a biomarker (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 1 and/or 2, such as COL5A2 (SEQ ID NO: 73 or 211)) in a patient having cancer (e.g., breast cancer), such as a patient having cancer that is resistant to one or more cancer therapies other than a sPLA₂ hydrolysable, cisplatin-containing liposome (e.g., a patient with breast cancer that is resistant to one or more cancer therapies other than the liposome). The method includes (a) contacting a sample (e.g., a tumor sample) from the patient including one or more nucleic acid molecules with a device (e.g., a microarray) including: i) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of sensitivity selected from those listed in Tables 1 and/or 3 or a complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)); and/or ii) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of resistance selected from those listed in Tables 2 and/or 4 or a complement thereof (e.g., SFN (SEQ ID NO: 96 or 324)); and (b) detecting a level of the one or more biomarkers of sensitivity or the complement thereof and/or the one or more biomarkers of resistance or the complement thereof in the sample by detecting hybridization between the one or more single-stranded nucleic acid molecules of the device and the one or more nucleic acid molecules of the sample (e.g., by performing microarray analysis or quantitative reverse transcriptase polymerase chain reaction (qRT-PCR)). The level of the biomarker(s) may be detected by determining the level of a messenger RNA (mRNA) corresponding to the biomarker(s); e.g., an mRNA expressed from COL5A2 (SEQ ID NO: 73 or 211)) or a complementary DNA (cDNA) thereof.

[0008] In the first aspect, the one or more cancer therapies may include surgery, radiation, or a therapeutic agent. In particular, the therapeutic agent may be selected from the group consisting of docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, and rituximab.

[0009] A second aspect of the invention features a method of determining responsiveness of a patient having cancer (e.g., one of the cancers noted above, such as breast cancer) to the liposome. In particular, the patient may have a cancer that is resistant to one or more cancer therapies other than the liposome, such as breast cancer that is resistant to one or more cancer therapies other than the liposome. The method includes a) contacting a sample (e.g., a tumor sample) from the patient including one or more nucleic acid molecules with a device (e.g., a microarray or a device for performing a qRT-PCR reaction) including: i) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of sensitivity selected from those listed in Tables 1 and/or 3 or a complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)); and/or ii) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of resistance selected from those listed in Tables 2 and/or 4 or a

complement thereof (e.g., SFN (SEQ ID NO: 96 or 324)); and b) detecting a level of the one or more biomarkers of sensitivity or the complement thereof and/or the one or more biomarkers of resistance or the complement thereof by detecting hybridization between the single-stranded nucleic acid molecules of the device and the one or more nucleic acid molecules of the sample. The patient is determined to be responsive to the liposome if: i) the level of the biomarkers of sensitivity or the complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)) is substantially similar to the level of the biomarkers of sensitivity or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome; and/or ii) the level of the biomarkers of resistance or the complement thereof (e.g., SFN (SEQ ID NO: 96 or 324)) is substantially dissimilar to the level of the biomarkers of resistance or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to the liposome.

[0010] The method of the second aspect can further include administering the liposome to the patient if: i) the level of the biomarkers of sensitivity or the complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)) is substantially similar to the level of the biomarkers of sensitivity or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome; and/or ii) the level of the biomarkers of resistance or the complement thereof (e.g., SFN (SEQ ID NO: 96 OR 324)) is substantially dissimilar to the level of the biomarkers of resistance or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to the liposome. The method can further include administering one or more cancer therapies other than the liposome to the patient if: i) the level of the biomarkers of sensitivity or the complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)) is substantially dissimilar to the level of the biomarkers of sensitivity or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome; and/or ii) the level of the biomarkers of resistance or the complement thereof (e.g., SFN (SEQ ID NO: 96 OR 324)) is substantially similar to the level of the biomarkers of resistance or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to the liposome.

[0011] In particular, the one or more of the cancer therapies includes surgery, radiation, or a therapeutic agent, such as docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, acliarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, and rituximab.

[0012] A third aspect of the invention features a method that includes contacting a sample from a patient having a cancer that is resistant to one or more cancer therapies (e.g., breast cancer), such as a patient having cancer that is resistant to one or more cancer therapies other than the liposome (e.g., a patient with breast cancer that is resistant to one or more cancer therapies other than the liposome) including one or more nucleic acid molecules, with a device including: i) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of sensitivity selected from the biomarkers of Tables 1 and 3 or a complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)), in which the biomarker of sensitivity is not C1QR1 (SEQ ID NO: 13), SLA (SEQ ID NO: 48), PTPN7 (SEQ ID NO: 77), CENTB1 (SEQ ID NO: 37), IFI16 (SEQ ID NO: 17 or 261), ARHGEF6 (SEQ ID NO: 36 or 294), CD3D (SEQ ID NO: 81), ARHGAP15 (SEQ ID NO: 30), HCLS1 (SEQ ID NO: 16 or 259), CD53 (SEQ ID NO: 282), PTPRCAP (SEQ ID NO: 8), or PTPRC (SEQ ID NO: 10, 18, 25, or 243); and/or ii) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of resistance selected from those listed in Tables 2 and/or 4 or a complement thereof (e.g., SFN (SEQ ID NO: 96 or 324)). The method also includes detecting a level of the one or more biomarkers of sensitivity or the complement thereof and/or the one or more biomarkers of resistance or the complement thereof in the sample by detecting hybridization between the one or more single-stranded nucleic acid molecules of the device and the one or more nucleic acid molecules of the sample (e.g., by performing microarray analysis or quantitative reverse transcriptase polymerase chain reaction (qRT-PCR)). The level of the biomarker(s) may be detected by determining the level of a messenger RNA (mRNA) corresponding to the biomarker(s); e.g., a mRNA expressed from COL5A2 (SEQ ID NO: 73 or 211)) or a complementary DNA (cDNA) thereof.

[0013] The method of the third aspect can further include that the patient is determined to be responsive to the liposome if: i) the level of the biomarkers of sensitivity or the complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)) is substantially similar to the level of the biomarkers of sensitivity or the complement thereof in a cell (e.g., a cancer cell) or

tissue (e.g., a tumor tissue) known to be sensitive to the liposome; and/or ii) the level of the biomarkers of resistance or the complement thereof (e.g., SFN (SEQ ID NO: 96 or 324)) is substantially dissimilar to the level of the biomarkers of resistance or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to the liposome. The method can also include administering the liposome to the patient if: i) the level of the biomarkers of sensitivity or the complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)) is substantially similar to the level of the biomarkers of sensitivity or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome; and/or ii) the level of the biomarkers of resistance or the complement thereof (e.g., SFN (SEQ ID NO: 96 or 324)) is substantially dissimilar to the level of the biomarkers of resistance or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to the liposome.

[0014] In any of the above aspects of the invention, the method can further include detecting a level of PLA2G2A (SEQ ID NO: 380) or a complement thereof in a tumor sample from the patient. In particular, the method can include contacting the tumor sample with one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of PLA2G2A (SEQ ID NO: 380) or a complement thereof and detecting a level of PLA2G2A or the complement thereof in the sample. Optionally, the detecting occurs by performing microarray analysis or qRT-PCR. The method can further include administering the liposome to the patient when a level of PLA2G2A is detected in the sample from the patient. The method can also further include detecting sPLA₂ protein in a tumor sample from the patient, such as by contacting the tumor sample with an anti-sPLA₂ antibody and detecting binding between the sPLA₂ protein and the anti-sPLA₂ antibody. Additionally, the method can include administering the liposome to the patient when the sPLA₂ protein is detected in the sample from the patient.

[0015] The invention also features a method of treating cancer in a patient in need thereof (e.g., one of the cancers noted above, such as breast cancer) that includes administering the liposome to the patient, in which the patient has been determined to be responsive to the liposome according to the method of the first or second aspect of the invention. In particular, the patient may have a cancer that is resistant to one or more cancer therapies other than the liposome (e.g., a patient with breast cancer that is resistant to one or more cancer therapies other than the liposome).

[0016] A fourth aspect of the invention features a method of treating a patient having cancer (e.g., one of the cancers noted above, such as breast cancer). In particular, the patient may have a cancer that is resistant to one or more cancer therapies other than the liposome (e.g., a patient with breast cancer that is resistant to one or more cancer therapies other than the liposome). The method includes a) contacting a sample (e.g., a tumor sample) from the patient including one or more nucleic acid molecules with a device (e.g., a microarray or a device for performing a qRT-PCR reaction) including: i) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of sensitivity selected from those listed in Tables 1 and/or 3 or a complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211)); and/or ii) one or more single-stranded nucleic acid molecules capable of specifically hybridizing with nucleotides of one or more biomarkers of resistance selected from those listed in Tables 2 and/or 4 or a complement thereof (e.g., SFN (SEQ ID NO: 96 OR 324)); b) detecting a level of the one or more biomarkers of sensitivity or the complement thereof and/or the one or more biomarkers of resistance or the complement thereof by detecting hybridization between the one or more single-stranded nucleic acid molecules of the device and the one or more nucleic acid molecules of the sample; and c) administering the liposome to the patient if: i) the level of the biomarkers of sensitivity or the complement thereof is substantially similar to the level of the biomarkers of sensitivity or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome; and/or ii) the level of the biomarkers of resistance or the complement thereof is substantially dissimilar to the level of the biomarkers of resistance or the complement thereof in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to the liposome.

[0017] The method of the fourth aspect of the invention may further include administering one or more additional therapies (e.g., surgery, radiation, or a therapeutic agent) to the patient prior to, concurrently with, or after administration of the liposome. In particular, the therapeutic agent may be selected from the group consisting of docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, cisplatin, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin,

ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, and rituximab. The therapeutic agent can be administered parenterally (e.g. intravenously, intramuscularly, transdermally, intradermally, intra-arterially, intracranially, subcutaneously, intraorbitally, intraventricularly, intraspinally, intraperitoneally, or intranasally), enterally, or topically.

[0018] In the fourth aspect of the invention, the liposome may be administered parenterally (e.g. intravenously, intramuscularly, transdermally, intradermally, intra-arterially, intracranially, subcutaneously, intraorbitally, intraventricularly, intraspinally, intraperitoneally, or intranasally), enterally, or topically. Preferably, the liposome is administered by intravenous injection. The liposome may be administered to the patient at a dose of about 60 mg/m² to about 120 mg/m² of cisplatin (e.g., 60-120 mg/m²). In particular, the liposome is administered to the patient at a dose of about 60 mg/m² of cisplatin weekly, about 75 mg/m² of cisplatin weekly, or about 90 mg/m² of cisplatin weekly. The method may include administering the liposome to the patient in a treatment regimen at least once per one, two, or three weeks or on day 1 and day 8 of a 3 week treatment regimen, in which the treatment regimen may optionally be repeated two to twenty times.

[0019] In the fourth aspect of the invention, the contacting step (a) and the detecting step (b) may occur prior to, concurrent, or after administration of the liposome to the patient. Additionally, the contacting step (a) and the detecting step (b) may occur two or more times, e.g., during treatment with the liposome. For example, the contacting step (a) and the measuring step (b) may occur two or more times to assess the continued sensitivity of the patient to the liposome.

[0020] In any of the above aspects of the invention, the cancer is selected from a solid tumor cancer and a hematological cancer. For example, the cancer is breast cancer, acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), Hodgkin's lymphoma, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, non-small cell lung carcinoma (NSCLC), prostate cancer, ovarian cancer, colon cancer, bladder cancer, and squamous cell carcinoma of the head and neck (SCCHN). In particular, the cancer is breast cancer, such as an estrogen receptor-positive (ERpos) breast cancer and/or a metastatic form of breast cancer.

[0021] In the second or fourth aspect of the invention, the cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome and/or the cell or tissue known to be resistant to the liposome is of the same type as a cell or tissue in the sample from the patient or from which the one or more nucleic acid molecules of the sample are derived. In particular, the cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to the liposome and/or the cell or tissue known to be resistant to the liposome is of the same type of cancer (e.g., breast cancer) as a cell or tissue in the sample from the patient or from which the one or more nucleic acid molecules of the sample are derived.

[0022] In any of the above aspects of the invention, the patient may exhibit cancer relapse (e.g., relapse of breast cancer), such as relapse after a first cancer treatment and prior to treatment with a therapeutic agent other than the liposome. Alternatively, the patient may have not been administered a treatment for cancer. Additionally, the patient may not have been determined to be resistant to the liposome.

[0023] In any of the above aspects of the invention, the device can include at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, at least ten, or more single-stranded nucleic acid molecules capable of specifically hybridizing with the nucleotides of one or more biomarkers of sensitivity selected from the biomarkers of Tables 1 and 3 or a complement thereof (e.g., COL5A2 (SEQ ID NO: 73 or 211); and/or at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, at least ten, or more single-stranded nucleic acid molecules capable of specifically hybridizing with the nucleotides of one or more biomarkers of resistance selected from the biomarkers of Tables 2 and 4 or a complement thereof (e.g., SFN (SEQ ID NO: 96 OR 324)). In particular, one or more of the single-stranded nucleic acid molecules of the device have a length in the range of 10 to 100 nucleotides in length (e.g., a length in the range of 20 to 60 nucleotides). For example, the one or more single-stranded nucleic acid molecules are labeled or immobilized on a solid substrate.

[0024] In any of the above aspects of the invention, the method may include converting the level of the one or more biomarkers of sensitivity or the complement thereof (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 1 and 3, such as COL5A2 (SEQ ID NO: 73 or 211)) and/or the one or more biomarkers of resistance or the complement thereof (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 2 and 4, such as SFN (SEQ ID NO: 96 OR 324)) into a mean score, in which the mean score indicates the responsiveness of the patient to

the liposome. The method can further include subtracting the mean score for one or more of the biomarkers of resistance (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 2 and 4, such as SFN (SEQ ID NO: 96 OR 324)) from the mean score for one or more of the biomarkers of sensitivity (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 1 and 3, such as COL5A2 (SEQ ID NO: 73 or 211) to obtain a difference score, in which the difference score indicates the responsiveness of the patient to the liposome. In particular, the mean score and/or the difference score above a cutoff value indicates that the patient is responsive to the liposome, such as if the cutoff value is about 0.25, about 0.3, about 0.35, about 0.4, about 0.45, about 0.5, or greater.

[0025] In any of the above aspects of the invention, the device can be a microarray, such as a deoxyribonucleic acid (DNA)-based platform. Alternatively, the device can be for performing a qRT-PCR reaction (e.g., the device is used with a system for detecting the amplification product, for example, by fluorescence or by another method). The methods may also utilize both a microarray and a qRT-PCR. Thus, the level of the biomarkers of sensitivity (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 1 and 3, such as COL5A2 (SEQ ID NO: 73 or 211) and/or the biomarkers of resistance (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 2 and 4, such as SFN (SEQ ID NO: 96 OR 324)) can be measured using qRT-PCR. In particular, the level of the one or more biomarkers of sensitivity or the complement thereof (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 1 and 3, such as COL5A2 (SEQ ID NO: 73 or 211)) and/or the one or more biomarkers of resistance or the complement thereof (e.g., one, two, three, four, five, ten, twenty, or all of the biomarkers shown in Tables 2 and 4, such as SFN (SEQ ID NO: 96 OR 324)) are detected by performing microarray analysis or qRT-PCR. Additionally, the nucleic acid molecules of the sample may include mRNA or a cDNA thereof.

[0026] In any of the above aspects of the invention, the biomarker of sensitivity may be selected from one or more of COL5A2 (SEQ ID NO: 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), FNBP1 (SEQ ID NO: 14 or 28), CBFEB (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 or 54), and CAP350 (SEQ ID NO: 20 or 61). The biomarker of resistance may be selected from one or more of S SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), and LRP5 (SEQ ID NO: 112).

[0027] For example, the biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211) and ITGA4 (SEQ ID NO: 1). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), and MSN (SEQ ID NO: 2). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), and FAM46A (SEQ ID NO: 3 OR 280). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), and ITGB2 (SEQ ID NO: 4). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), and DOCK2 (SEQ ID NO: 5 OR 223). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), and EVL (SEQ ID NO: 6). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), and SACS (SEQ ID NO: 7). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), and PTPRCAP (SEQ ID NO: 8). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), and EBI2 (SEQ ID NO: 9). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), and PTPRC (SEQ ID NO: 10, 18, 25, OR 243). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, OR 243), and ANP32E (SEQ ID NO: 11). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, OR 243), ANP32E (SEQ ID NO: 11), and SFPQ (SEQ ID NO: 12, 38 OR 272). The biomarkers of sensitivity may include COL5A2

(SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, OR 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 OR 272), and C1QR1 (SEQ ID NO: 13). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 OR 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 OR 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, OR 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 OR 272), C1QR1 (SEQ ID NO: 13), and FNBP1 (SEQ ID NO: 14 OR 28). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), and CBFB (SEQ ID NO: 15). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO: 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), and SFRS7 (SEQ ID NO: 19 or 54). The biomarkers of sensitivity may include COL5A2 (SEQ ID NO: 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 or 54), and CAP350 (SEQ ID NO: 20 or 61).

[0028] For example, the biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324) and LISCH7 (SEQ ID NO: 97). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), and EPB41L4B (SEQ ID NO: 98). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), and MST1R (SEQ ID NO: 99). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), and ITGB4 (SEQ ID NO: 100). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), and DBNDD2 (SEQ ID NO: 102 OR 365). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), and TACSTD1 (SEQ ID NO: 104). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), and MISP (SEQ ID NO: 105). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), and KRT8 (SEQ ID NO: 106). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), and JUP (SEQ ID NO: 107 OR 400). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 OR 400), and KRT18 (SEQ ID NO: 108 OR 306). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 OR 400), KRT18 (SEQ ID NO: 108 OR 306, and FA2H (SEQ ID NO: 109). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 OR 400), KRT18 (SEQ ID NO: 108 OR 306, FA2H (SEQ ID NO: 109), and MGAT4B (SEQ ID NO: 110). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 OR 400), KRT18 (SEQ ID NO: 108 OR 306, FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), and DSG2 (SEQ ID NO: 111 OR 312). The biomarkers of resistance may include SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 OR 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 OR 400), KRT18 (SEQ ID NO: 108 OR 306, FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 OR 312), and LRP5 (SEQ ID NO: 112).

Definitions

[0029] As used herein, "a" or "an" means "at least one" or "one or more" unless otherwise indicated. In addition, the singular forms "a", "an", and "the" include plural referents unless the context clearly dictates otherwise.

[0030] As used herein, "about" refers to an amount $\pm$ 10% of the recited value.

[0031] By "biomarker" is meant a nucleic acid molecule (e.g., a mRNA or its complement, for example, a cDNA, or the nucleic acid sequence of all or a fragment of a gene) or a protein encoded by the nucleic acid molecule present in, or from, a cell or tissue. The level of the biomarker correlates to the responsiveness (e.g., sensitivity or resistance) of the cell or tissue (and thus, the patient in which the cell or tissue resides or the patient from which the cell or tissue was obtained) to a cancer treatment (e.g., secretory phospholipase A₂ (sPLA₂) hydrolysable liposome). In particular, biomarkers of sensitivity correspond to the nucleic acid molecules or the complements thereof (e.g., a mRNA or its complement or a cDNA thereof) shown in Tables 1 and 3, or the proteins encoded by the nucleic acid molecules, and biomarkers of resistance correspond to the nucleic acid molecules or the complements thereof (e.g., a mRNA or its complement or a cDNA thereof) shown in Tables 2 and 4, or the proteins encoded by the nucleic acid molecules.

[0032] The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals (e.g., humans) that is typically characterized by unregulated cell proliferation. Examples of cancer include, but are not limited to, breast cancer (e.g., medullary carcinoma or ER-positive breast cancer), prostate cancer, ovarian cancer (e.g., ovarian adenocarcinoma or embryonal carcinoma), liver cancer (e.g., hepatocellular carcinoma (HCC) or hepatoma), myeloma (e.g., multiple myeloma), colorectal cancer (e.g., colon cancer and rectal cancer), leukemia (e.g., acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, and chronic leukemia), myelodysplastic syndrome, lymphoma (e.g., diffuse large B-cell lymphoma, cutaneous T-cell lymphoma, peripheral T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Waldenstrom's macroglobulinemia, and lymphocytic lymphoma), cervical cancer, esophageal cancer, melanoma, glioma (e.g., oligodendroglioma), pancreatic cancer (e.g., adenosquamous carcinoma, signet ring cell carcinoma, hepatoid carcinoma, colloid carcinoma, islet cell carcinoma, and pancreatic neuroendocrine carcinoma), gastrointestinal stromal tumor, sarcoma (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, leiomyosarcoma, Ewing's sarcoma, and rhabdomyosarcoma), bladder cancer, head and neck cancer (e.g., squamous cell carcinoma of the head and neck), lung cancer (e.g., non-small cell lung carcinoma, large cell carcinoma, bronchogenic carcinoma, and papillary adenocarcinoma), metastatic cancer, oral cavity cancer, uterine cancer, testicular cancer (e.g., seminoma and embryonal carcinoma), skin cancer (e.g., squamous cell carcinoma and basal cell carcinoma), thyroid cancer (e.g., papillary carcinoma and medullary carcinoma), brain cancer (e.g., astrocytoma and craniopharyngioma), stomach cancer, intra-epithelial cancer, bone cancer, biliary tract cancer, eye cancer, larynx cancer, kidney cancer (e.g., renal cell carcinoma and Wilms tumor), gastric cancer, blastoma (e.g., nephroblastoma, medulloblastoma, hemangioblastoma, neuroblastoma, and retinoblastoma), polycythemia vera, chordoma, synovioma, mesothelioma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, cystadenocarcinoma, bile duct carcinoma, choriocarcinoma, epithelial carcinoma, ependymoma, pinealoma, acoustic neuroma, schwannoma, meningioma, pituitary adenoma, nerve sheath tumor, cancer of the small intestine, cancer of the endocrine system, cancer of the penis, cancer of the urethra, cutaneous or intraocular melanoma, a gynecologic tumor, solid tumors of childhood, and neoplasms of the central nervous system. The term cancer includes solid tumors (e.g., breast cancer) and hematological cancers (e.g., cancer of the blood, such as lymphoma (e.g., diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), and Hodgkin's lymphoma)).

[0033] "Gene" as used herein indicates a coding or noncoding gene whose activity can be determined by measuring RNA (e.g., mRNA) transcribed from the gene, or its complement. Examples include protein coding genes, microRNAs, small nuclear RNAs and other RNAs with catalytic, regulatory or coding properties.

[0034] To "inhibit growth" as used herein means causing a reduction in cell growth (e.g., cancer cell growth, such as the NCI60 cancer cell lines) *in vivo* or *in vitro* by, e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% or more, as evident by a reduction in the proliferation of cells exposed to a treatment (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome described herein), relative to the proliferation of cells in the absence of the treatment. Growth inhibition may be the result of a treatment (e.g., treatment with the sPLA₂ hydrolysable, cisplatin-containing liposome) that induces apoptosis in a cell, induces necrosis in a cell, slows cell cycle progression, disrupts cellular metabolism, induces cell lysis, or induces some other mechanism that reduces the proliferation of cells.

[0035] "Microarray" as used herein means a device employed by any method that quantifies one or more subject oligonucleotides, e.g., RNA, DNA, cDNA, or analogues thereof, at a time. For example, many DNA microarrays, including those made by Affymetrix (e.g., an Affymetrix HG-U133A or HG-U133_Plus_2 array), use several probes for determining the level of a single biomarker. The DNA microarray may contain oligonucleotide probes that may be, e.g., full-length cDNAs complementary to an RNA or cDNA fragments that hybridize to part of an RNA. The DNA microarray may also contain modified versions of DNA or RNA, such as locked nucleic acids or LNA. Exemplary RNAs include mRNA, miRNA, and miRNA precursors.

[0036] "NCI60" as used herein means a panel of 60 cancer cell lines from lung, colon, breast, ovarian, leukemia, renal, melanoma, prostate, and brain cancers including the following cancer cell lines: NSCLC_NCIH23, NSCLC_NCIH522, NSCLC_A549ATCC, NSCLC_EKVX, NSCLC_NCIH226, NSCLC_NCIH332M, NSCLC_H460, NSCLC_HOP62, NSCLC_HOP92, COLON_HT29, COLON_HCC-2998, COLON_HCT116, COLON_SW620, COLON_COLO205, COLON_HCT15, COLON_KM12, BREAST_MCF7, BREAST_MCF7ADPr, BREAST_MDAMB231, BREAST_HS578T, BREAST_MDAMB435, BREAST_MDN, BREAST_BT549, BREAST_T47D, OVAR_OVCAR3, OVAR_OVCAR4, OVAR_OVCAR5, OVAR_OVCAR8, OVAR_IGROV1, OVAR_SKOV3, LEUK_CCRFCM, LEUK_K562, LEUK_MOLT4, LEUK_HL60, LEUK_RPMI8266, LEUK_SR, RENAL_UO31, RENAL_SN12C, RENAL_A498, RENAL_CAK11, RENAL_RXF393, RENAL_7860, RENAL_ACHN, RENAL_TK10, MELAN_LOXIMVI, MELAN_MALME3M, MELAN_SKMEL2, MELAN_SKMEL5, MELAN_SKMEL28, MELAN_M14, MELAN_UACC62, MELAN_UACC257, PROSTATE_PC3, PROSTATE_DU145, CNS_SNB19, CNS_SNB75, CNS_U251, CNS_SF268, CNS_SF295, and CNS_SF539.

[0037] The terms "patient" and "subject," as used interchangeably herein, refer to any animal (e.g., a mammal, such as a human). A patient to be treated or tested for responsiveness to a treatment (e.g., treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome) according to the methods described herein may be one who has been diagnosed with a cancer, such as those described herein, e.g., breast cancer, acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), Hodgkin's lymphoma, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, non-small cell lung carcinoma (NSCLC), prostate cancer, ovarian cancer, colon cancer, bladder cancer, or squamous cell carcinoma of the head and neck (SCCHN). Diagnosis may be performed by any method or technique known in the art, such as x-ray, MRI, or biopsy, and may also be confirmed by a physician. To minimize exposure of a patient to drug treatments that may not be therapeutic, the patient may be determined to be either responsive or non-responsive to a cancer treatment, such as an sPLA₂ hydrolysable, cisplatin-containing liposome, according to the methods described herein prior to treatment.

[0038] As used herein, the term "percent (%) sequence identity" refers to the percentage of nucleic acid residues of a candidate sequence, e.g., a probe or primer of the invention, that are identical to the nucleic acid residues of a reference sequence, e.g., a biomarker sequence of the invention, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity (e.g., gaps can be introduced in one or both of the candidate and reference sequences for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). Alignment for purposes of determining percent sequence identity can be achieved in various ways that are within the skill in the art, for instance, using computer software, such as BLAST, BLAST-2, BLAST-P, BLAST-N, BLAST-X, WU-BLAST-2, ALIGN, ALIGN-2, CLUSTAL, Megalign (DNASTAR). In addition, those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve optimal alignment over the length of the sequences being compared.

[0039] "Resistant" or "resistance" as used herein means that a cell (e.g., a cancer cell), a tissue containing the cell (e.g., a tumor), or a patient containing the cell or tissue having cancer (e.g., a human having cancer) is non-responsive to treatment with an anti-cancer agent (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome). In particular, the treatment reduces the growth of a resistant cell (e.g., the cancer cell) *in vitro* by less than about 40%, 30%, 20%, 10%, 5%, 1%, or less, relative to the growth of a less resistant cell not exposed to the treatment. Resistance to treatment may be determined by a cell proliferation assay, e.g., a cell-based assay, which measures the growth of treated cells as a function of the absorbance of the cells of an incident light beam, such as the NCI60 assays described herein. In this assay, greater absorbance indicates greater cell growth, and thus, resistance to the treatment.

[0040] A cancer patient (e.g., a patient having breast cancer, acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell

lymphoma (PTCL), Hodgkin's lymphoma, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, non-small cell lung carcinoma (NSCLC), prostate cancer, ovarian cancer, colon cancer, bladder cancer, or squamous cell carcinoma of the head and neck (SCCHN)) may also have resistance to a cancer therapy other than an sPLA₂ hydrolysable, cisplatin-containing liposome, such as surgery, radiation, or a therapeutic agent (e.g., docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, or rituximab).

[0041] The terms "responsive" and "responsiveness," as used herein, refer to the likelihood that a cancer treatment (e.g., treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome) has (e.g., induces) a desired effect in a cell (e.g., a cancer cell), a tissue (e.g., a tumor), or a patient having cancer (e.g., a human having cancer). For example, the desired effect can include inhibition of the growth of a cancer cell *in vitro* by more than 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% relative to the growth of a cancer cell not exposed to the treatment. The desired effect can also include reduction in tumor mass by, e.g., about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. Responsiveness to treatment may be determined by a cell proliferation assay, e.g., a cell-based assay, which measures the growth of treated cells as a function of the absorbance of the cells of an incident light beam, such as the NCI60 assays described herein. In this assay, lesser absorbance indicates lesser cell growth, and thus, sensitivity to the treatment. A greater reduction in growth indicates more sensitivity to the treatment. In particular, "responsiveness" is a measure of the sensitivity or resistance of a patient to a treatment for cancer (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome).

[0042] The term "sample," as used herein, refers to any specimen (such as cells, tissue (e.g., a tissue sample obtained by biopsy), blood, serum, plasma, urine, cerebrospinal fluid, or pancreatic fluid) taken from a subject. Preferably, the sample is taken from a portion of the body affected by a cancer (e.g., a biopsy of the cancer tissue, such as breast cancer tissue). Biopsy may involve fine needle aspiration biopsy, core needle biopsy (e.g., stereotactic core needle biopsy, vacuum-assisted core biopsy, or magnetic resonance imaging (MRI) guided biopsy), or surgical biopsy (e.g., incisional biopsy or excisional biopsy). The sample may undergo additional purification and processing, for example, to remove cell debris and other unwanted molecules. Additional processing may further involve producing cDNA molecules corresponding to nucleic acid molecules (e.g., mRNA) in the sample and/or amplification of the nucleic acid molecules, e.g., using PCR, such as RT-PCR. The standard methods of sample purification, such as removal of unwanted molecules, are known in the art.

[0043] The terms "secretory phospholipase A₂ (sPLA₂) hydrolyzable, cisplatin-containing liposome," "sPLA₂ hydrolysable, cisplatin-containing liposome," and "the liposome" as used herein refer to an antitumor agent that is a liposomal formulation of cisplatin. The sPLA₂ hydrolysable, cisplatin-containing liposome (e.g., LiPlaCis®, LiPlasome Pharma) is formulated to release an encapsulated drug (e.g., cisplatin) from the core of a hydrophobic layer into tumor tissue. Since sPLA₂ protein is associated with tumor tissue, sPLA₂ hydrolysable liposomes may be used to preferentially deliver encapsulated drugs (e.g., cisplatin) to the tumor tissue. An sPLA₂ hydrolysable, cisplatin-containing liposome is described in, e.g., U.S. Patent Application Publication No. 2012/0177726 and de Jonge et al. (Eur J Cancer. 46(16):3016-21, 2010).

[0044] "Sensitive" and "sensitivity" as used herein refer to a cell (e.g., a cancer cell), a tissue containing the cell (e.g., a tumor), or a patient containing the cell or tissue having cancer (e.g., a human having cancer) that is responsive to treatment, such as an anti-cancer agent (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome) or radiation treatment. In particular, the treatment inhibits the growth of the cell (e.g., the cancer cell) *in vitro* by about 70%, 80%, 90%, 95%, 99% or 100% relative to the growth of a cell not exposed to the treatment. Sensitivity to treatment may be determined by a cell proliferation assay, e.g., a cell-based assay, which measures the growth of treated cells as a function of the absorbance of the cells of an incident light beam, such as the NCI60 assays described herein. In this assay, lesser absorbance indicates lesser cell growth, and thus, sensitivity to the treatment.

[0045] The term "specific hybridization" as used herein refers to when complementary nucleic acid sequences form a stable

duplex under high stringency conditions, such as high hybridization temperature and low salt in hybridization buffers, which permit only hybridization between nucleic acid sequences that are highly similar. Nucleic acids are referred to as "complementary" that contain nucleotides or nucleotide homologues that can form hydrogen bonds according to Watson-Crick base-pairing rules (e.g., G with C, A with T or A with U) or other hydrogen bonding motifs such as for example diaminopurine with T, 5-methyl C with G, 2-thiothymidine with A, inosine with C, pseudoisocytosine with G, etc. Anti-sense RNA may be complementary to other oligonucleotides, e.g., mRNA.

[0046] "Treatment," "medical treatment," to "treat," and "therapy," as used interchangeably herein, refer to administering or exposing a patient having cancer (e.g., a human) to an anti-cancer agent (e.g., a drug such as an sPLA₂ hydrolysable, cisplatin-containing liposome, a protein, an antibody, a nucleic acid, a chemotherapeutic agent, or a radioactive agent), or to some other form of medical intervention used to treat or prevent a disease, disorder, or condition (e.g., surgery, cryotherapy, radiation therapy, or combinations thereof). In particular, a medical treatment can be or can include administration of an sPLA₂ hydrolysable, cisplatin-containing liposome. For example, the treatment may be of a cancer, such as a solid tumor or a hematological cancer. Examples of cancer include, e.g., breast cancer (e.g., medullary carcinoma or an ER-positive breast cancer), prostate cancer, ovarian cancer (e.g., ovarian adenocarcinoma or embryonal carcinoma), liver cancer (e.g., hepatocellular carcinoma (HCC) or hepatoma), myeloma (e.g., multiple myeloma), colorectal cancer (e.g., colon cancer and rectal cancer), leukemia (e.g., acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, and chronic leukemia), myelodysplastic syndrome, lymphoma (e.g., diffuse large B-cell lymphoma, cutaneous T-cell lymphoma, peripheral T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Waldenstrom's macroglobulinemia, and lymphocytic lymphoma), cervical cancer, esophageal cancer, melanoma, glioma (e.g., oligodendroglioma), pancreatic cancer (e.g., adenosquamous carcinoma, signet ring cell carcinoma, hepatoid carcinoma, colloid carcinoma, islet cell carcinoma, and pancreatic neuroendocrine carcinoma), gastrointestinal stromal tumor, sarcoma (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, leiomyosarcoma, Ewing's sarcoma, and rhabdomyosarcoma), bladder cancer, head and neck cancer (e.g., squamous cell carcinoma of the head and neck), lung cancer (e.g., non-small cell lung carcinoma, large cell carcinoma, bronchogenic carcinoma, and papillary adenocarcinoma), metastatic cancer, oral cavity cancer, uterine cancer, testicular cancer (e.g., seminoma and embryonal carcinoma), skin cancer (e.g., squamous cell carcinoma and basal cell carcinoma), thyroid cancer (e.g., papillary carcinoma and medullary carcinoma), brain cancer (e.g., astrocytoma and craniopharyngioma), stomach cancer, intra-epithelial cancer, bone cancer, biliary tract cancer, eye cancer, larynx cancer, kidney cancer (e.g., renal cell carcinoma and Wilms tumor), gastric cancer, blastoma (e.g., nephroblastoma, medulloblastoma, hemangioblastoma, neuroblastoma, and retinoblastoma), polycythemia vera, chordoma, synovioma, mesothelioma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, cystadenocarcinoma, bile duct carcinoma, choriocarcinoma, epithelial carcinoma, ependymoma, pinealoma, acoustic neuroma, schwannoma, meningioma, pituitary adenoma, nerve sheath tumor, cancer of the small intestine, cancer of the endocrine system, cancer of the penis, cancer of the urethra, cutaneous or intraocular melanoma, a gynecologic tumor, solid tumors of childhood, and neoplasms of the central nervous system. Radiation therapy includes the administration of a radioactive agent to a patient or exposure of a patient to radiation. The radiation may be generated from sources, such as particle accelerators and related medical devices or agents that emit, e.g., X-radiation, gamma radiation, or electron (Beta radiation) beams. A treatment may be or further include surgery, e.g., to remove a tumor from a subject or living organism.

[0047] Other features and advantages of the invention will be apparent from the following Detailed Description, the drawings, and the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0048] Figure 1 is a graph grouping predicted sensitivity to a secretory phospholipase A₂ (sPLA₂) hydrolysable, cisplatin-containing liposome by cancer type. Each gray circle represents the predicted liposome sensitivity of one patient calculated as the difference between the mean of the levels of the biomarkers of sensitivity (e.g., Tables 1 and/or 3) and the mean of the levels of the biomarkers of resistance for the patient (e.g., Tables 2 and/or 4). Patients are grouped according to cancer type. The median predicted sensitivity (black bar) for a cancer type is related to the relative response rate for that cancer type. The predictions are used for relative comparisons to compare cancer types and cannot be used for absolute predictions of response rate for a given cancer type. The predictions are normalized to a scale of 0 to 100 for all 3,522 patients.

DETAILED DESCRIPTION OF THE INVENTION

[0049] I have discovered that the levels of one or more of the biomarkers shown in Tables 1-4 (e.g., as single biomarkers or combinations of biomarkers) may be detected in a patient having cancer and used to predict the responsiveness of the patient to a secretory phospholipase A₂ (sPLA₂) hydrolysable, cisplatin-containing liposome. These patients may already be determined to be resistant to a therapy other than an sPLA₂ hydrolysable, cisplatin-containing liposome, such as docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxin, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, or rituximab.

[0050] A device, such as a microarray, with one or more single-stranded oligonucleotide probes that have substantial identity (e.g., at least 85%, 90%, 95%, 99%, or 100% sequence identity) to a sequence that is complementary or identical to the nucleic acid sequence of one or more biomarkers shown in Tables 1-4 can be used according to the methods described herein to assess the responsiveness of a cancer patient to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. For example, the probes can be used to detect one or more (e.g., two, three, four, five, ten, twenty, or all) of the biomarkers of sensitivity listed in Tables 1 and 3, such as COL5A2 (SEQ ID NO 73 or 211), in a sample (e.g., a tumor sample) from a patient having cancer (e.g., breast cancer). Additionally, the probes can be used to detect one or more (e.g., two, three, four, five, ten, twenty, or all) of the biomarkers of resistance listed in Tables 2 and 4, such as SFN (SEQ ID NO: 96 or 324), in a sample (e.g., a tumor sample) from a patient having cancer (e.g., breast cancer).

[0051] Accordingly, the invention features individual biomarkers (e.g., COL5A2 (SEQ ID NO 73 or 211) or SFN (SEQ ID NO: 96 or 324)) and sets of biomarkers shown in Tables 1-4 that can be used to determine the responsiveness of a cancer patient to an sPLA₂ hydrolysable, cisplatin-containing liposome at various stages of disease progression (e.g., patients diagnosed with cancer or patients after cancer recurrence) and at different times during the treatment process (e.g., prior to administration of any cancer treatment, after administration of one or more cancer treatments other than an sPLA₂ hydrolysable, cisplatin-containing liposome, prior to administration of the liposome, or during administration of the liposome). The individual biomarkers (e.g., COL5A2 (SEQ ID NO 73 or 211) or SFN (SEQ ID NO: 96 or 324)) and sets of biomarkers shown in Tables 1-4 can be used in combination with a level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ protein to determine the responsiveness of a cancer patient to an sPLA₂ hydrolysable, cisplatin-containing liposome. Additionally, the methods can be used to determine responsiveness of a cancer patient to an sPLA₂ hydrolysable, cisplatin-containing liposome that is resistant to one or more cancer therapies other than the liposome, such as docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, AP0010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxin, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, or rituximab.

[0052] In particular, the invention provides methods for determining whether a patient may be responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome by, e.g., detecting the level (e.g., mRNA or a protein produced therefrom) of one or more of the biomarkers shown in Tables 1-4 (e.g., COL5A2 (SEQ ID NO 73 or 211)) in a biological sample (e.g., a tumor biopsy) obtained from the subject using a device (e.g., a microarray or a protein array). The level of one or more of the biomarkers of sensitivity may then be compared to the level of the biomarkers in a cell or tissue known to be sensitive or

resistant to the liposome to determine the patient's responsiveness to the liposome. The patient is determined to be responsive to the liposome if the level of the one or more of the biomarkers of sensitivity (e.g., one or more of COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38, or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDI3 (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), and LAT2 (SEQ ID NO: 220)) is substantially similar to the level of the biomarkers of sensitivity in a cell or tissue known to be sensitive to the liposome (e.g., from a patient sensitive to the liposome). The patient is also determined to be responsive to the liposome if the level of one or more of the biomarkers of resistance (e.g., one or more of SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), and IFI27 (SEQ ID NO: 321)) is substantially dissimilar to the level of the biomarkers of resistance in a cell or tissue known to be resistant to the liposome (e.g., from a patient resistant to the liposome). Additionally, the patient is determined to be responsive to the liposome if a level of PLA2G2A or sPLA₂ is detected in a tumor sample from the patient and the level of the biomarkers of sensitivity (e.g., COL5A2 (SEQ ID NO 73 or 211)) is substantially similar to the level of the biomarkers of sensitivity in a tissue known to be sensitive to the liposome and/or the level of the biomarkers of resistance is substantially dissimilar to the level of the biomarkers of resistance (e.g., SFN (SEQ ID NO: 96 or 324)) in a tissue known to be resistant to the liposome.

[0053] The invention also features methods of treating a patient having cancer, such as a patient having a cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome, by detecting the levels of one or more of the biomarkers shown in Tables 1-4 (e.g., COL5A2 (SEQ ID NO: 73 or 211) in a sample (e.g., a tumor sample) from the patient, and then administering the liposome based on the levels of the biomarkers. In particular, a patient having cancer may be administered the liposome if the level of one or more biomarkers of sensitivity is substantially similar to the level of the biomarkers of sensitivity in a cell or tissue known to be sensitive to the liposome. Additionally, a patient having cancer may be administered the liposome if the level of one or more biomarkers of resistance is substantially dissimilar to the level of the biomarkers of resistance in a cell or tissue known to be resistant to the liposome. Thus, the methods can be used to treat cancer patients predicted to be responsive to the liposome, such as patients having, e.g., breast cancer (e.g., a metastatic form of breast cancer), prostate cancer, ovarian cancer, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, estrogen receptor-positive (ERpos) breast cancer, non-small cell lung carcinoma (NSCLC), colon cancer, bladder cancer, squamous cell carcinoma of the head and neck (SCCHN), acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), and Hodgkin's lymphoma. Alternatively, a patient having cancer may not be administered the liposome if the level of one or more biomarkers of sensitivity is substantially dissimilar to the level of the biomarkers of sensitivity in a cell or tissue known to be sensitive to the liposome. Likewise, a patient having cancer may not be administered the liposome if the level of one or more biomarkers of resistance is substantially similar to the level of the biomarkers of resistance in a cell or tissue known to be resistant to the liposome. Additionally, the patient may not be administered the liposome if a level of PLA2G2A or the protein product thereof (sPLA₂) is not detected in a tumor sample from the patient in combination with the methods.

[0054] Methods are described herein for identifying biomarkers of drug responsiveness, detecting levels of one or more biomarkers of sensitivity and one or more biomarkers of resistance in cancer patients, determining the responsiveness of a cancer patient to an sPLA₂ hydrolysable, cisplatin-containing liposome, and treating cancer patients with the liposome. Also described are devices and kits for use in these methods.

Methods for Identifying Biomarkers of Drug Responsiveness

[0055] The invention features methods for identifying biomarkers (e.g., one or more of the biomarkers of Tables 1-4) for determining the responsiveness of a cancer patient to a cancer treatment, such as an sPLA₂ hydrolysable, cisplatin-containing liposome. Such methods can involve, for example, an algorithm based on growth inhibition values (GI50) of cell lines (e.g., NCI60 cell lines) subjected to treatment with the liposome, followed by measurement of gene expression (e.g., using a microarray (e.g., an Affymetrix HG-U133A or HG-U133_Plus_2 array)).

Methodology of the In Vitro Cancer Growth Inhibition Screen

[0056] The human tumor cell lines of the cancer screening panel may be grown in RPMI 1640 medium containing 5% fetal bovine serum and 2 mM L-glutamine. Cells may be inoculated into 96 well microtiter plates in 100 μ L at plating densities ranging from 5,000 to 40,000 cells/well depending on the doubling time of individual cell lines. After cell inoculation, the microtiter plates may be incubated at 37°C, 5% CO₂, 95% air and 100% relative humidity for 24 hours prior to addition of experimental agent (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome).

[0057] After 24 hours, two plates of each cell line may be fixed *in situ* with TCA, to represent a measurement of the cell population for each cell line at the time of agent addition (Tz). Experimental agents may be solubilized in dimethyl sulfoxide at 400-fold the desired final maximum test concentration and stored frozen prior to use. At the time of agent (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome) addition, an aliquot of frozen concentrate may be thawed and diluted to twice the desired final maximum test concentration with complete medium containing 50 μ g/ml Gentamicin. A total of four additional 10-fold or ½ log serial dilutions are made to provide a total of five concentrations plus control. Aliquots of 100 μ L of these different agent dilutions are added to the appropriate microtiter wells already containing 100 μ L of medium, resulting in the required final agent concentrations.

[0058] Following agent (e.g., a sPLA₂ hydrolysable, cisplatin-containing liposome) addition, the plates may be incubated for an additional 48 h at 37°C, 5% CO₂, 95% air, and 100% relative humidity. For adherent cells, the assay may be terminated by the addition of cold TCA. Cells may be fixed *in situ* by the gentle addition of 50 μ L of cold 50% (w/v) TCA (final concentration, 10% TCA) and incubated for 60 minutes at 4°C. The supernatant may be discarded, and the plates may be washed five times with tap water and air-dried. Sulforhodamine B (SRB) solution (100 μ L) at 0.4% (w/v) in 1% acetic acid may be added to each well, and the plates may be incubated for 10 minutes at room temperature. After staining, unbound dye may be removed by washing five times with 1% acetic acid and the plates may be air-dried. Bound stain may be subsequently solubilized with 10 mM trizma base, and the absorbance may be read on an automated plate reader at a wavelength of 515 nm. For suspension cells, the methodology may be the same, except that the assay may be terminated by fixing settled cells at the bottom of the wells by gently adding 50 μ L of 80% TCA (final concentration, 16 % TCA). Using the seven absorbance measurements [time zero, (Tz), control growth, (C), and test growth in the presence of agent (e.g., the liposome) at the five concentration levels (Ti)], the percentage growth may be calculated at each of the agent concentration levels. Percentage growth inhibition may be calculated as:

$[(Ti - Tz) / (C - Tz)] \times 100$ for concentrations for which $Ti \geq Tz$

$[(Ti - Tz) / Tz] \times 100$ for concentrations for which $Ti < Tz$

[0059] Three dose response parameters may be calculated for each experimental agent (e.g., the liposome). Growth inhibition of 50% (GI50) is calculated from $[(Ti - Tz) / (C - Tz)] \times 100 = 50$, which is the agent (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome) concentration resulting in a 50% reduction in the net protein increase (as measured by SRB staining) in control cells during incubation with the test agent. The agent concentration resulting in total growth inhibition (TGI) is calculated from $Ti = Tz$. The LC50 (concentration of the agent, such as an sPLA₂ hydrolysable, cisplatin-containing liposome, resulting in a 50% reduction in the measured protein at the end of the compound treatment as compared to that at the beginning) indicating a net loss of cells following treatment is calculated from $[(Ti - Tz) / Tz] \times 100 = -50$. Values are calculated for each of these three parameters if the level of activity is reached; however, if the effect is not reached or is exceeded, the value for that parameter is expressed as greater or less than the maximum or minimum concentration tested.

Gene Expression and Growth Inhibition Analysis

[0060] The gene expression measurements of NCI60 cancer cell lines can be obtained from a publicly available database (e.g., the National Cancer Institute and the Massachusetts Institute of Technology). Each dataset can be normalized so that sample expression measured by different chips can be compared. The preferred method of normalization is the logit transformation, which may be performed for each gene y on each chip, as follows:

$$\text{logit}(y) = \log \left[\frac{(y - \text{background})}{(\text{saturation} - y)} \right],$$

where background is calculated as the minimum intensity measured on the chip minus 0.1% of the signal intensity range: $\text{min} - 0.001 * (\text{max} - \text{min})$, and saturation is calculated as the maximum intensity measured on the chip plus 0.1% of the signal intensity range: $\text{max} + 0.001 * (\text{max} - \text{min})$. The resulting logit transformed data may then be z-transformed to mean zero and standard deviation 1.

[0061] Next, gene expression can be correlated to cancer cell growth inhibition. Growth inhibition data (GI50) of the NCI60 cell lines in the presence of a cancer treatment, such as an sPLA₂ hydrolysable, cisplatin-containing liposome, can be obtained from the NCI. The correlation between the logit-transformed expression level of each gene in each cell line and the logarithm of GI50 (the concentration of a given compound that results in a 50% inhibition of growth) can be calculated, e.g., using the Pearson correlation coefficient or the Spearman Rank-Order correlation coefficient. Instead of using GI50s, any other measure of patient sensitivity to a given treatment (e.g., an sPLA₂ hydrolysable, cisplatin-containing liposome) may be correlated to gene expression levels of the patient. Since a plurality of measurements may be available for a single gene, the most accurate determination of correlation coefficient can be, e.g., the median of the correlation coefficients calculated for all probes measuring expression of the same gene.

[0062] For example, the median correlation coefficient of gene expression measured on a probe to growth inhibition or patient sensitivity to an sPLA₂ hydrolysable, cisplatin-containing liposome can be calculated for all genes of interest. Genes that have a median correlation above about 0.25 (e.g., above 0.25, 0.26, 0.27, 0.28, 0.29, 0.30, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.40, or higher (e.g., 0.3 or higher)), can be used as biomarkers of sensitivity for assessing responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than the liposome) to the liposome. Likewise, genes that have a median correlation below about -0.25 (e.g., below -0.25, -0.26, -0.27, -0.28, -0.29, -0.30, -0.31, -0.32, -0.33, -0.34, -0.35, -0.36, -0.37, -0.38, -0.39, -0.40, or lower (e.g., -0.3 or lower)), can be used as biomarkers of resistance for assessing responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than the liposome) to the liposome. Preferably, the correlation coefficient of a biomarker of sensitivity will exceed 0.25, while the correlation coefficient of a biomarker of resistance will be less than -0.25. The result is a list of biomarker genes that correlate to sensitivity or resistance to the liposome, as shown in Tables 1 and 3 and Tables 2 and 4, respectively.

Cancer Types

[0063] The methods, devices, and kits of the invention can be used for prognosing, monitoring, treating, and/or reducing cancer in a subject suffering from, diagnosed with, or susceptible to cancer. Non-limiting examples of cancers that can be prognosed, monitored, treated (e.g., by administering an sPLA₂ hydrolysable, cisplatin-containing liposome), or reduced using the methods include hematological and solid tumors. In particular, cancers include, e.g., breast cancer (e.g., an estrogen receptor-positive (ERpos) breast cancer or a metastatic form of breast cancer), prostate cancer, ovarian cancer (e.g., ovarian adenocarcinoma or embryonal carcinoma), liver cancer (e.g., hepatocellular carcinoma (HCC) or hepatoma), myeloma (e.g., multiple myeloma), colorectal cancer (e.g., colon cancer and rectal cancer), leukemia (e.g., acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, and chronic leukemia), myelodysplastic syndrome, lymphoma (e.g., diffuse large B-cell lymphoma, cutaneous T-cell lymphoma, peripheral T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Waldenstrom's macroglobulinemia, and lymphocytic lymphoma), cervical cancer, esophageal cancer, melanoma, glioma (e.g., oligodendroglioma), pancreatic cancer (e.g., adenosquamous carcinoma, signet ring cell carcinoma, hepatoid carcinoma, colloid carcinoma, islet cell carcinoma, and pancreatic neuroendocrine carcinoma), gastrointestinal stromal tumor, sarcoma (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, leiomyosarcoma, Ewing's sarcoma, and rhabdomyosarcoma), breast cancer (e.g., medullary carcinoma), bladder cancer, head and neck cancer (e.g., squamous cell carcinoma of the head and neck), lung cancer (e.g., non-small cell lung carcinoma, large cell carcinoma, bronchogenic carcinoma, and papillary adenocarcinoma), metastatic cancer, oral cavity cancer, uterine cancer, testicular cancer (e.g., seminoma and embryonal

carcinoma), skin cancer (e.g., squamous cell carcinoma and basal cell carcinoma), thyroid cancer (e.g., papillary carcinoma and medullary carcinoma), brain cancer (e.g., astrocytoma and craniopharyngioma), stomach cancer, intra-epithelial cancer, bone cancer, biliary tract cancer, eye cancer, larynx cancer, kidney cancer (e.g., renal cell carcinoma and Wilms tumor), gastric cancer, blastoma (e.g., nephroblastoma, medulloblastoma, hemangioblastoma, neuroblastoma, and retinoblastoma), polycythemia vera, chordoma, synovioma, mesothelioma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, cystadenocarcinoma, bile duct carcinoma, choriocarcinoma, epithelial carcinoma, ependymoma, pinealoma, acoustic neuroma, schwannoma, meningioma, pituitary adenoma, nerve sheath tumor, cancer of the small intestine, cancer of the endocrine system, cancer of the penis, cancer of the urethra, cutaneous or intraocular melanoma, a gynecologic tumor, solid tumors of childhood, and neoplasms of the central nervous system.

[0064] In particular, the methods are useful for prognosing, monitoring, treating, or preventing, e.g., breast cancer (e.g., an ERpos breast cancer or a metastatic form of breast cancer), prostate cancer, ovarian cancer, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, non-small cell lung carcinoma (NSCLC), colon cancer, bladder cancer, squamous cell carcinoma of the head and neck (SCCHN), acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), and Hodgkin's lymphoma. For example, the cancer can be breast cancer, such as Stage 0, Stage I, Stage II, Stage III, or Stage IV breast cancer. In particular, the cancer may be breast cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome, such as doxorubicin, cyclophosphamide, fluorouracil (5-FU), methotrexate, docetaxel, epirubicin, paclitaxel, carboplatin, trastuzumab, vinorelbine, vinorelbin, radiation, and/or surgery. For instance, the breast cancer is medullary carcinoma. The breast cancer may also be, e.g., a metastatic form of breast cancer.

Methods for Detecting Biomarker Levels in Cancer Patients

[0065] A cancer patient can be assessed for sensitivity or resistance to an sPLA₂ hydrolysable, cisplatin-containing liposome by detecting a level of a biomarker (e.g., one or more of the biomarkers of Tables 1-4) in a biological sample obtained from the cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than the liposome). Additionally, the detection of sPLA₂ and/or a level of PLA2G2A (SEQ ID NO: 380) in a tumor sample from the cancer patient can be used in combination with the level of the biomarker (e.g., one or more of the biomarkers of Tables 1-4) to determine patient sensitivity or resistance to the liposome. The biological sample can include, for example, cells, tissue (e.g., a tissue sample obtained by biopsy), blood, serum, plasma, urine, sputum, cerebrospinal fluid, lymph tissue or fluid, or pancreatic fluid. For example, the biological sample can be fresh frozen or formalin-fixed paraffin embedded (FFPE) tissue obtained from the subject, such as a tumor sample (e.g., a biopsy) from the tissue of interest (e.g., prostate, ovarian, lung, lymph nodes, thymus, spleen, bone marrow, breast, colorectal, pancreatic, cervical, bladder, gastrointestinal, head, or neck tissue).

RNA Extraction and Measurement of Biomarker Levels

[0066] Cell samples or tissue samples may be snap frozen in liquid nitrogen until processing. RNA may be extracted using, e.g., Trizol Reagent from Invitrogen following manufacturer's instructions, and detected directly or converted to cDNA for detection. RNA may be amplified using, e.g., MessageAmp kit from Ambion following manufacturer's instructions. Amplified RNA may be quantified using, e.g., HG-U133A or HG-U133_Plus2 GeneChip from Affymetrix Inc. or a compatible apparatus, e.g., the GCS3000Dx GeneChip® System from Affymetrix Inc., using the manufacturer's instructions. The resulting biomarker level measurements may be further analyzed as described herein. The procedures described can be implemented using, e.g., R software available from R-Project (www.r-project.org) and supplemented with packages available from Bioconductor (www.bioconductor.org).

[0067] The level of one or more of the biomarkers shown in Tables 1-4 (e.g., COL5A2 (SEQ ID NO: 73 or 211)) may be measured in a biological sample (e.g., a tumor sample) obtained from the cancer patient (e.g., a patient having any of the cancer types described herein, such as a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) using, e.g., polymerase chain reaction (PCR), reverse transcriptase PCR (RT-PCR), quantitative real-time PCR (qRT-PCR), an array (e.g., a microarray), a genechip, pyrosequencing,

nanopore sequencing, sequencing by synthesis, sequencing by expansion, single molecule real time technology, sequencing by ligation, microfluidics, infrared fluorescence, next generation sequencing (e.g., RNA-Seq techniques), Northern blots, Western blots, Southern blots, NanoString nCounter technologies (e.g., those described in U.S. Patent Application Nos. US 2011/0201515, US 2011/0229888, and US 2013/0017971, proteomic techniques (e.g., mass spectrometry or protein arrays), and combinations thereof. In combination with the level of these biomarkers, the level of PLA2G2A (SEQ ID NO: 380) or the gene product thereof (sPLA₂) may be detected using, e.g., PCR, RT-PCR, qRT-PCR, an array (e.g., a microarray), a genechip, pyrosequencing, nanopore sequencing, sequencing by synthesis, sequencing by expansion, single molecule real time technology, sequencing by ligation, microfluidics, infrared fluorescence, next generation sequencing (e.g., RNA-Seq techniques), Northern blots, Western blots, Southern blots, NanoString nCounter technologies, an anti-sPLA₂ antibody, proteomic techniques (e.g., mass spectrometry or protein arrays), and combinations thereof.

Devices

[0068] Devices of the invention can be used for detecting a level of one or more biomarkers shown in Tables 1-4 and/or a level of PLA2G2A (SEQ ID NO: 380). The device may include at least one (or one type of) single-stranded nucleic acid (e.g., a probe) having at least 85% sequence identity (e.g., 85%, 90%, 95%, 97%, 98%, 99%, or 100% sequence identity) to a nucleic acid sequence that is complementary or identical to at least 5 (e.g., at least 10, at least 15, at least 20, or more) consecutive nucleotides of one or more biomarkers shown in Tables 1-4 (e.g., COL5A2 (SEQ ID NO 73 or 211) or SFN (SEQ ID NO: 96 or 324)) and/or PLA2G2A (SEQ ID NO: 380), in which the at least one single-stranded nucleic acid is sufficient for the detection of the level of the one or more biomarkers. The device may be used to detect the level of a given biomarker by specific hybridization between the single-stranded nucleic acid and the biomarker (e.g., an mRNA, genomic DNA, or non-coding RNA), a nucleic acid of the biomarker (e.g., an mRNA), or a complementary nucleic acid thereof. The device may be or include a microarray. The device may also include or be used with reagents and materials for next generation sequence (e.g., sequencing by synthesis). The device may also include or be used with NanoString reagents and at least one nCounter cartridge. The device may be or include a protein array, which contains one or more protein binding moieties (e.g., proteins, antibodies, nucleic acids, aptamers, affibodies, lipids, phospholipids, small molecules, labeled variants of any of the above, and any other moieties useful for protein detection as well known in the art) capable of detectably binding to the polypeptide product(s) of one or more biomarkers shown in Tables 1-4 and/or PLA2G2A (SEQ ID NO: 380). The device may also be a cartridge for measuring an amplification product resulting from hybridization between one or more nucleic acid molecules from the patient and at least one single-stranded nucleic acid single-stranded nucleic acid molecules of the device, such as a device for performing qRT-PCR.

Microarrays

[0069] The levels of the biomarkers (e.g., the biomarkers listed in Tables 1-4 (e.g., COL5A2 (SEQ ID NO: 73 or 211)) and/or PLA2G2A (SEQ ID NO: 380)) may be determined using high-throughput expression profiling platforms, such as microarrays. In particular, a microarray for use in the methods for assessing the responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) to the liposome contains or is produced by generating oligonucleotide probes (e.g., DNA, cDNA, or RNA probes) capable of hybridizing to one or more biomarkers of interest (e.g., one or more of the biomarkers of Tables 1-4) and/or PLA2G2A (SEQ ID NO: 380) or the complement sequences thereof. Each probe can have, e.g., at least 10, 15, 20, 25, 30, or more contiguous nucleic acid residues (e.g., at least 15) that are complementary or identical to a nucleic acid sequence of a selected biomarker. The probe nucleic sequence can also have at least 85% (e.g., 90%, 95%, 99%, or 100%) sequence identity to the nucleic acid sequence of the gene coding the biomarker (e.g., COL5A2 (SEQ ID NO 73 or 211)) or the complement sequence thereof. In particular, the probe sequences can be complementary to all or a portion of the nucleic acid sequence of the biomarker(s).

[0070] For example, microarrays of the invention for determining responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome can include probes for one or more (e.g., at least 5, 10, 15, or 20 or more (e.g., all)) biomarkers of sensitivity shown in Tables 1 and 3, such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38, or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO:

15), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), or LAT2 (SEQ ID NO: 220). Microarrays for determining responsiveness to the liposome can also include probes for one or more (e.g., at least 5, 10, 15, or 20 or more (e.g., all)) biomarkers of resistance listed in Tables 2 and 4, such as SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321).

[0071] Microarrays for determining responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome can also include probes for one or more (e.g., at least 5, 10, 15, or 20 or more (e.g., all)) biomarkers of sensitivity and biomarkers of resistance shown in Tables 1-4, such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38, or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFβ (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). Additionally, microarrays of the invention for determining responsiveness to the liposome can include one or more probes for PLA2G2A (SEQ ID NO: 380).

[0072] A microarray probe may be single-stranded or double-stranded. The probe may be labeled (e.g., detectably labeled with a fluorescent molecule, dye molecule, small molecule, epitope tag, barcode sequence, polypeptide, or any other detectable molecule). Probes can be detectably labeled and immobilized on a solid support to form the microarray. Additionally, probes can be detectably labeled and included in, e.g., a tube, such as probes labeled with a fluorescent molecule. For example, probes can be either prefabricated and spotted to the surface or directly synthesized on to the surface (*in situ*) of the microarray. The microarray can also be configured such that the sequence and position of each member (e.g., probe) of the array is known. For example, a selection of biomarkers whose levels correlates with an increased likelihood of responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome can be arrayed on a solid support. Hybridization of a labeled probe with a particular target nucleic acid (e.g., an mRNA corresponding to one or more biomarkers of Tables 1-4 and/or PLA2G2A (SEQ ID NO: 380)) indicates that the sample from which the mRNA was derived expresses that biomarker (e.g., the biomarker of sensitivity or resistance to an sPLA₂ hydrolysable, cisplatin-containing liposome).

PCR-based techniques

[0073] As few as one to thirty (e.g., 5 to 30 or 10 to 30, or at least the first 15 of the biomarkers listed in Tables 1-4) biomarkers, optionally in combination with PLA2G2A (SEQ ID NO: 380), may be used to determine patient responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome using the methods described herein. Tissue or cell samples from a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) can be conveniently assayed for levels using PCR analysis, such as quantitative real-time PCR (qRT-PCR), or quantitative loop-mediated isothermal amplification (q-LAMP). For example, an mRNA corresponding to a biomarker of Tables 1-4 and/or PLA2G2A (SEQ ID NO: 380) can be detected in a biological

sample by (a) producing cDNA from the sample by reverse transcription using at least one primer; (b) amplifying the cDNA so produced using a target polynucleotide as sense and antisense primers to amplify target cDNAs therein; and (c) detecting the presence of the amplified target cDNA using polynucleotide probes. The primers and probes including the target sequences shown in Tables 1-4, such as COL5A2 (SEQ ID NO 73 or 211) and/or SFN (SEQ ID NO: 96 or 324), and/or the sequence of PLA2G2A (SEQ ID NO: 380) may be used to detect the levels of one or more of the indicated biomarkers using PCR. The methods can include one or more steps that allow determination of the levels of target mRNA in a biological sample (e.g., by simultaneously examining the levels of a comparative control mRNA sequence or "housekeeping" gene, such as an actin family member or GAPDH). The primers for these PCR-based assays may be labeled for detection according to methods known in the art.

Sequencing

[0074] The levels of the biomarkers shown in Tables 1-4, such as COL5A2 (SEQ ID NO 73 or 211), SFN (SEQ ID NO: 96 or 324), and/or PLA2G2A (SEQ ID NO: 380) may be determined using sequencing technologies, such as next generation sequencing platforms (e.g., RNA-Seq), as described in Mortazavi et al., Nat. Methods 5: 621-628, 2008. RNA-Seq is a robust technology for monitoring biomarker levels by direct sequencing of the RNA molecules in a sample. This methodology may include fragmentation of RNA to an average length of, e.g., 200 nucleotides, conversion to cDNA by random priming, and synthesis of double-stranded cDNA (e.g., using the PROTOCOLSCRIPT® First Strand cDNA Synthesis Kit from New England Biosciences). The cDNA may then be converted into a molecular library for sequencing by addition of sequence adapters for each library (e.g., from ILLUMINA®/Solexa), and the resulting 50 to 100 nucleotide reads are mapped onto the genome. Exemplary sequencing platforms suitable for use according to the methods include, e.g., pyrosequencing, ILLUMINA® sequencing by synthesis, SOLID® sequencing, ION TORRENT® sequencing, and SMRT® sequencing.

Methods of Determining Patient Responsiveness to an sPLA₂ Hydrolysable, Cisplatin-containing Liposome

[0075] The invention features diagnostic methods for the detection and screening of cancer patients (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) that may be responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome using one or more of the biomarkers shown in Tables 1-4 (e.g., COL5A2 (SEQ ID NO 73 or 211) or SFN (SEQ ID NO: 96 or 324)), optionally in combination with PLA2G2A (SEQ ID NO: 380). The methods of the invention may be used for predicting a patient's responsiveness to the liposome, and optionally, treating the cancer patient throughout the progression of cancer and/or in cases of recurrence (e.g., after a first line treatment, a second line treatment, and/or a third line treatment).

[0076] The invention provides individual biomarkers (e.g., COL5A2 (SEQ ID NO: 73 or 211) and sets of biomarkers (e.g., two or more of the biomarkers listed in Tables 1-4), the levels of which, as detected in a biological sample (e.g., a tumor sample, such as a biopsy) obtained from a cancer patient (e.g., a human with cancer), are indicative of responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The biomarkers were identified using methods similar to those previously described in, e.g., Chen et al. (Mol. Cancer Ther. 11:34-33, 2012), Wang et al. (J. Nat. Cancer Inst. 105: 1284-1291, 2013), and Knudsen et al. (PLoS One, 9: e87415, 2014).

[0077] In particular, an algorithm based on growth inhibition values (GI₅₀) of a cell line (e.g., NCI60 cells) subjected to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome and gene expression is determined (e.g., by microarray analysis, reverse transcriptase polymerase chain reaction (RT-PCR), quantitative real-time PCR (qPCR), or next generation sequencing). After normalization, genes with, e.g., a Pearson correlation coefficient greater than about 0.25 or below about -0.25 can be classified as biomarkers of sensitivity or resistance, respectively. In particular, a correlation coefficient of about 0.25 or greater is a statistically significant cut-off known in the art for establishing whether the levels of the biomarker, e.g., the biomarkers shown in Tables 1-4, correlate with the likelihood of cancer treatment sensitivity, such as sensitivity to the liposome. Thus, a correlation coefficient of about 0.25 or greater or about -0.25 or lower can be used to identify biomarkers, such as the biomarkers of Tables 1-4, for predicting patient responsiveness to treatment with the liposome according to the methods described herein.

Comparison of Biomarker Levels

[0078] One or more biomarkers of sensitivity and/or resistance, identified as described herein, can be used to predict responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome by measuring the level of the biomarkers in a biological sample obtained from the cancer patient, optionally in combination with the level of PLA2G2A (SEQ ID NO: 380) in a tumor sample obtained from the cancer patient. A single biomarker (e.g., any of the biomarkers of Tables 1-4, such as COL5A2 (SEQ ID NO: 73 or 211) or SFN (SEQ ID NO: 96 or 324)) may be used to determine the responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) to the liposome.

[0079] For instance, after determining the level of a biomarker of sensitivity (e.g., any of the biomarkers of Tables 1 and 3, such as COL5A2 (SEQ ID NO: 73 or 211)) in a sample (e.g., a tumor sample) from the cancer patient, the level of the biomarker(s) in the sample may be compared to the level of the biomarker(s) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. If the level of the biomarker(s) in the sample from the cancer patient is substantially similar (e.g., identical to or has the same trend of level) to the level of the biomarker(s) in the cell or tissue known to be sensitive to the liposome, then the cancer patient is predicted to be responsive to treatment with the liposome. Alternatively, if the level of the biomarker(s) in the sample from the cancer patient is substantially dissimilar to the level of the biomarker(s) in the cell or tissue known to be sensitive to the liposome, then the cancer patient is predicted to be non-responsive to treatment with the liposome.

[0080] The level of the biomarker of sensitivity (e.g., any of the biomarkers of Tables 1 and 3, such as COL5A2 (SEQ ID NO: 73 or 211)) in a sample (e.g., a tumor sample) from the cancer patient may also be compared to the level of the biomarker(s) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. If the level of the biomarker(s) in the sample from the cancer patient is substantially similar to the level of the biomarker(s) in the cell or tissue known to be resistant to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be non-responsive to treatment with the liposome. Alternatively, if the level of the biomarker(s) in the sample from the cancer patient is substantially dissimilar to the level of the biomarker(s) in the cell or tissue known to be sensitive to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be responsive to treatment with the liposome.

[0081] For instance, after determining the level of a biomarker of resistance (e.g., any of the biomarkers of Tables 2 and 4, such as SFN (SEQ ID NO: 96 or 324)) in a sample (e.g., a tumor sample) from the cancer patient, the level of the biomarker(s) in the sample may be compared to the level of the biomarker(s) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. If the level of the biomarker(s) in the sample from the cancer patient is substantially dissimilar to the level of the biomarker(s) in the cell or tissue known to be sensitive to the liposome, then the cancer patient is predicted to be responsive to treatment with the liposome. Alternatively, if the level of the biomarker(s) in the sample from the cancer patient is substantially similar (e.g., identical to or has the same trend of level) to the level of the biomarker(s) in the cell or tissue known to be sensitive to the liposome, then the cancer patient is predicted to be non-responsive to treatment with the liposome.

[0082] The level of a biomarker of resistance (e.g., any of the biomarkers of Tables 2 and 4, such as SFN (SEQ ID NO: 96 or 324)) in a sample (e.g., a tumor sample) from the cancer patient may also be compared to the level of the biomarker(s) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. If the level of the biomarker(s) in the sample from the cancer patient is substantially similar to the level of the biomarker(s) in the cell or tissue known to be resistant to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be responsive to treatment with the liposome. Alternatively, if the level of the biomarker(s) in the sample from the cancer patient is substantially dissimilar to the level of the biomarker(s) in the cell or tissue known to be sensitive to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be non-responsive to treatment with the liposome.

[0083] The responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) to an sPLA₂ hydrolysable, cisplatin-containing liposome can also be predicted by comparing the level of a biomarker (e.g., COL5A2 (SEQ ID NO: 73 or 211)) to the level of the biomarker in one or more cells or tissues (e.g., from a cancer patient population) known to be sensitive to treatment with the liposome and one or more cells or tissues (e.g., from a cancer patient population) known to be resistant to treatment with the liposome. In particular, the patient may be determined to be responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the level of the biomarker(s) is more similar to the level of the biomarker(s) in

a cell or tissue known to be sensitive to treatment with the liposome than to a cell or tissue known to be resistant to treatment with the liposome. Alternatively, the patient may be determined to be non-responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the level of the biomarker(s) is more similar to the level of the biomarker(s) in a cell or tissue known to be resistant to treatment with the liposome than to a cell or tissue known to be sensitive to treatment with the liposome.

[0084] Additionally, one or more biomarkers of sensitivity (e.g., one or more of COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38, or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), or LAT2 (SEQ ID NO: 220)) and one or more biomarkers of resistance (e.g., one or more of SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321)) may be used in combination to determine the responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. For example, the predicted responsiveness of a cancer patient may be determined from, e.g., the difference score, which may be defined as the difference between the mean of the level of the one or more biomarkers of sensitivity of Tables 1 and 3 and the mean of the level of the one or more biomarkers of resistance of Tables 2 and 4.

[0085] The difference score of the cancer patient can then be compared to the difference score based on the level of the biomarkers in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. In particular, the patient may be determined to be responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the difference score is substantially similar to the level of the biomarkers in a cell or tissue known to be sensitive to treatment with the liposome. Alternatively, the patient may be determined to be non-responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the difference score is substantially similar to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome. Additionally, the patient may be determined to be responsive to treatment with the liposome if the difference score is substantially similar to the level of the biomarkers in a cell or tissue known to be sensitive to treatment with the liposome than a cell or tissue known to be resistant to treatment with the liposome. Alternatively, the patient may be determined to be non-responsive to treatment with the liposome if the difference score is substantially similar to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome than a cell or tissue known to be sensitive to treatment with the liposome.

[0086] One or more biomarkers of sensitivity and/or resistance, identified as described herein, can be used to predict responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome by measuring the level of the biomarkers in a biological sample obtained from the cancer patient. A single biomarker (e.g., any of the biomarkers of Tables 1-4, such as COL5A2 (SEQ ID NO: 73 or 211)), may be used to determine the responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) to an sPLA₂ hydrolysable, cisplatin-containing liposome. After determining the level of the biomarker(s) in a sample (e.g., a tumor sample) from the cancer patient, the level of the biomarker(s) in the sample may be compared to the level of the biomarker(s) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. If the level of the biomarker(s) in the sample from the cancer patient corresponds to (e.g., is identical to or has the same trend of level as) the level of the biomarker(s) in the cell or tissue known to be sensitive to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be responsive to treatment with the liposome. Alternatively, if the level of the biomarker(s) in the sample from the cancer patient does not correspond to the level of the biomarker(s) in the cell or tissue known to be sensitive to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be non-responsive to treatment with the

liposome.

[0087] The level of the biomarker (e.g., COL5A2 (SEQ ID NO: 73 or 211)) in a sample from the cancer patient may also be compared to the level of the biomarker in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be resistant to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. If the level of the biomarker in the sample from the cancer patient corresponds to the level of the biomarker in the cell or tissue known to be resistant to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be non-responsive to treatment with the liposome. Alternatively, if the level of the biomarker in the sample from the cancer patient does not correspond to the level of the biomarker in the cell or tissue known to be resistant to an sPLA₂ hydrolysable, cisplatin-containing liposome, then the cancer patient is predicted to be responsive to treatment with the liposome.

[0088] The responsiveness of a cancer patient (e.g., a patient having cancer that is resistant to one or more cancer therapies other than the liposome) to an sPLA₂ hydrolysable, cisplatin-containing liposome can also be predicted by comparing the level of a biomarker (e.g., COL5A2 (SEQ ID NO: 73 or 211)) to the level of the biomarker in one or more cells or tissues (e.g., from a cancer patient population) known to be sensitive to treatment with the liposome and one or more cells or tissues (e.g., from a cancer patient population) known to be resistant to treatment with the liposome. In particular, the patient may be responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the level of the biomarker(s) corresponds to the level of the biomarker(s) in a cell or tissue known to be sensitive to treatment with the liposome relative to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome. Alternatively, the patient may be non-responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the level of the biomarker(s) corresponds to the level of the biomarker(s) in a cell or tissue known to be resistant to treatment with the liposome relative to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome.

[0089] For example, the patient may be responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the difference score (e.g., the difference between the mean of the level of the one or more biomarkers of sensitivity of Tables 1 and 3 and the mean of the level of the one or more biomarkers of resistance of Tables 2 and 4) corresponds to the level of the biomarkers in a cell or tissue known to be sensitive to treatment with the liposome. Alternatively, the patient may be non-responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the difference score corresponds to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome. Additionally, the patient may be responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the difference score corresponds to the level of the biomarkers in a cell or tissue known to be sensitive to treatment with the liposome relative to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome. Alternatively, the patient may be non-responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome if the difference score corresponds to the level of the biomarkers in a cell or tissue known to be resistant to treatment with the liposome relative to the level of the biomarkers in a cell or tissue known to be sensitive to treatment with the liposome.

[0090] Preferably, the cell or tissue known to be either sensitive or resistant to an sPLA₂ hydrolysable, cisplatin-containing liposome is of the same cancer type as the cancer patient with an unknown responsiveness to the liposome. For example, the cancer patient and the cell or tissue known to be either sensitive or resistant to an sPLA₂ hydrolysable, cisplatin-containing liposome may both have a cancer type selected from a solid tumor or a hematological cancer, e.g., breast cancer (e.g., a ER-positive breast cancer or medullary carcinoma), prostate cancer, ovarian cancer (e.g., ovarian adenocarcinoma or embryonal carcinoma), liver cancer (e.g., hepatocellular carcinoma (HCC) or hepatoma), myeloma (e.g., multiple myeloma), colorectal cancer (e.g., colon cancer and rectal cancer), leukemia (e.g., acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, and chronic leukemia), myelodysplastic syndrome, lymphoma (e.g., diffuse large B-cell lymphoma, cutaneous T-cell lymphoma, peripheral T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Waldenstrom's macroglobulinemia, and lymphocytic lymphoma), cervical cancer, esophageal cancer, melanoma, glioma (e.g., oligodendroglioma), pancreatic cancer (e.g., adenosquamous carcinoma, signet ring cell carcinoma, hepatoid carcinoma, colloid carcinoma, islet cell carcinoma, and pancreatic neuroendocrine carcinoma), gastrointestinal stromal tumor, sarcoma (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, leiomyosarcoma, Ewing's sarcoma, and rhabdomyosarcoma), bladder cancer, head and neck cancer (e.g., squamous cell carcinoma of the head and neck), lung cancer (e.g., non-small cell lung carcinoma, large cell carcinoma, bronchogenic carcinoma, and papillary adenocarcinoma), metastatic cancer, oral cavity cancer, uterine cancer, testicular cancer (e.g., seminoma and embryonal carcinoma), skin cancer (e.g., squamous cell

carcinoma and basal cell carcinoma), thyroid cancer (e.g., papillary carcinoma and medullary carcinoma), brain cancer (e.g., astrocytoma and craniopharyngioma), stomach cancer, intra-epithelial cancer, bone cancer, biliary tract cancer, eye cancer, larynx cancer, kidney cancer (e.g., renal cell carcinoma and Wilms tumor), gastric cancer, blastoma (e.g., nephroblastoma, medulloblastoma, hemangioblastoma, neuroblastoma, and retinoblastoma), polycythemia vera, chordoma, synovioma, mesothelioma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, cystadenocarcinoma, bile duct carcinoma, choriocarcinoma, epithelial carcinoma, ependymoma, pinealoma, acoustic neuroma, schwannoma, meningioma, pituitary adenoma, nerve sheath tumor, cancer of the small intestine, cancer of the endocrine system, cancer of the penis, cancer of the urethra, cutaneous or intraocular melanoma, a gynecologic tumor, solid tumors of childhood, and neoplasms of the central nervous system.

[0091] In particular, the cancer of the patient and the cell or tissue with known resistance or sensitivity to an sPLA₂ hydrolysable, cisplatin-containing liposome is, e.g., breast cancer (e.g., ERpos breast cancer and/or a metastatic form of breast cancer), acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large B-cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), Hodgkin's lymphoma, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, non-small cell lung carcinoma (NSCLC), prostate cancer, ovarian cancer, colon cancer, bladder cancer, or squamous cell carcinoma of the head and neck (SCCHN).

[0092] Machine learning techniques such as Neural Networks, Support Vector Machines, K Nearest Neighbor, and Nearest Centroids may also be employed to develop models that discriminate patients sensitive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome from those resistant to treatment with the liposome using biomarker levels as model variables which assign each patient a classification as sensitive or resistant to treatment with the liposome. Machine learning techniques used to classify patients using various measurements are described in U.S. Patent No. 5,822,715; U.S. Patent Application Publication Nos. 2003/0073083, 2005/0227266, 2005/0208512, 2005/0123945, 2003/0129629, and 2002/0006613; and in Vapnik V. N. Statistical Learning Theory, John Wiley & Sons, New York, 1998; Hastie et al., 2001, The Elements of Statistical Learning: Data Mining, Inference, and Prediction, Springer, N.Y.; Agresti, 1996, An Introduction to Categorical Data Analysis, John Wiley & Sons, New York; V. Tresp et al., "Neural Network Modeling of Physiological Processes," in Hanson S. J. et al. (Eds.), Computational Learning Theory and Natural Learning Systems 2, MIT Press, 1994.

Biomarkers of Sensitivity and Resistance

[0093] The levels of one or more of the biomarkers of Tables 1-4 can be used to determine cancer patient responsiveness to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. Once determined to be responsive using the methods of the invention, the patient can be treated with the liposome alone or the liposome and one or more of the other therapies described herein.

Table 1. mRNA biomarkers of sensitivity to cisplatin. Dashes mean that the Affymetrix probeset has not been mapped to a specific gene. Affymetrix IDs refer to the array type HG-U133A.

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
ITGA4	213416_at	0.46	CAGGCCTCTCAGATACAAGGGGAAC	1
MSN	200600_at	0.45	ATAGCTGCCTTAAAGTCAGTAAGT	2
FAM46A	221766_s_at	0.41	CACCATGCTGGCTATCCGGGTGTTA	3
ITGB2	202803_s_at	0.39	CTCCACTCTGACTGGCACAGTCTTT	4
DOCK2	213160_at	0.39	GATTCCTGAACTCAAGGTACCAGCA	5
EVL	217838_s_at	0.39	GATCATCGACGCCATCAGGCAGGAG	6
SACS	213262_at	0.38	GTGTGGTTGAACAGGATGCAATCTT	7
PTPRCAP	204960_at	0.37	GCTTCCCAAGATGCCATGGCTGGAC	8
EBI2	205419_at	0.37	GCAGGACTTCCCTTATAAAGCAAAA	9
PTPRC	212587_s_at	0.37	GATTATAACCGTGTTGAACTCTCTG	10
ANP32E	221505_at	0.37	GTTTTCGGTCCTATTTTAATGCTCT	11

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
SFPQ	201586_s_at	0.36	AAAGACCAACAAATCTCAAGCCCTA	12
C1QR1	202878_s_at	0.36	GGTCTGTTCTTGTAGATAATGCCCT	13
FNBP1	213940_s_at	0.36	TGCTGGCCACGGATTTTGACGACGA	14
CBFB	202370_s_at	0.35	GGTGTGTACAGCTCACATGTTTAC	15
HCLS1	202957_at	0.35	GGTTTGCCTCATTGTGCTATTTGCC	16
IFI16	208965_s_at	0.35	ATAAGCATTGATTCTGCATTTCTG	17
PTPRC	212588_at	0.35	GCATTAGTCCAATGTCTTTTAAAG	18
SFRS7	213649_at	0.35	ATCATGCTGAGGCGCCTTGCAAATC	19
CAP350	204373_s_at	0.34	ATGACTGGTATGATAGCTCTTGACA	20
IGLL1	206660_at	0.34	CAATCCAAGCATAACTCAGTGACGC	21
DOCK10	219279_at	0.34	GAATGTGTAGCTCAAATGCAAACCA	22
WASPIP	202664_at	0.33	TTCCCTCCTTATAGTCAAGGACCGT	23
FLI1	204236_at	0.33	TGACCTCGGTACAAAAGCAGTTTT	24
PTPRC	207238_s_at	0.33	GAACAGTTTGTACAGACGTATGCTT	25
IFI16	208966_x_at	0.33	TACAACACTATACATACACACCACC	26
HDGFRP3	209524_at	0.33	TTATGCCAGCTTATATTGTGAGAAC	27
FNBP1	212288_at	0.33	GAGTTGCCTGTTTGTCTCTGGAGAT	28
SEPT6	212414_s_at	0.33	GCTGCAGTGTAGATGGCTCTTGTTT	29
ARHGAP15	218870_at	0.33	ACGTTGTCACCGGAGCACTGAAGAT	30
RASSF2	203185_at	0.32	ATAGCAGCACACATTTTCACGTTTC	31
GMFG	204220_at	0.32	AAGACCGGCAGATGGTGGTGCTGGA	32
SYNCRIP	209025_s_at	0.32	ATTGCGCTCAAGTCCATTGCGCTGT	33
HDGFRP3	209526_s_at	0.32	GCATGAAGTTGCCCTTAACCACTAA	34
ARHGEF6	209539_at	0.32	TAACCATGCTTACACACTAAACTAT	35
TMEM5	204808_s_at	0.31	TGCCCGGTCCGAGTAAACACAGAAT	36
CENTB1	205213_at	0.31	GATGTCAACTGGGTCAATGGGGGCC	37
SFPQ	214016_s_at	0.31	GTTGGCTGATATTGGAGTGCTCATT	38
BCAT1	214452_at	0.31	CCTTTTGTACTTCACTCAGATACTA	39
LCP1	208885_at	0.3	TAAGCATCCTTAGGGTTCTGCCTCT	40
CORO1A	209083_at	0.3	CTCATCTCCCTCAAGGATGGCTACG	41
SLC4A7	209884_s_at	0.3	TGTGAATCATCCTGCCTTTCAAATT	42
RAFTLIN	212646_at	0.3	TACAAACCACATTACTTCTGTCACT	43
CKIP-1	218223_s_at	0.3	GTCCCGGATCCAGGACCTGGTAGCA	44
SNRP70	201221_s_at	0.29	AGTGAAGAGGTCGTCCTCTCCATCT	45
BNIP3	201849_at	0.29	GCTGAAGGCACCTACTCAGTATCTT	46
SLA	203761_at	0.29	TAAGCATTCCGTCCATCTAAGCTCA	47
MFNG	204153_s_at	0.29	TGATGGAGCATAACGGGTCCCAGCC	48
LOC57821	206721_at	0.29	ATGATTCTTAGGGTCTGTGTACTT	49
CBLB	209682_at	0.29	GTTCCATTTCTCTCATTCAACAAGAT	50
QKI	212636_at	0.29	GAGGCCAAGAAATCCATGTTGTTT	51
ZRF1	213097_s_at	0.29	AAAGCTGTGAATCTGTTCCCTGCTG	52
FTL	213187_x_at	0.29	ATGAGCTCCCAGATTGCTCAGAATT	53
SFRS7	214141_x_at	0.29	TCCCCATCAGGAAGTCCCTCGCAGAA	54
VIM	201426_s_at	0.28	TGAGTCCCTGGAACGCCAGATGCGT	55
PWP1	201606_s_at	0.28	TTAGAGCCAGTCTTCACACTCGGAA	56

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
AKAP7	205771_s_at	0.28	AAAACCTCCCGGTATGATGATTGT	57
AF1Q	211071_s_at	0.28	TCAGTGGGCACAGTTCTTCAGCTAC	58
DICER1	213229_at	0.28	ACTAGCTCATTATTTCCATCTTTGG	59
PDE4DIP	213388_at	0.28	AATTATGAGTTTCTATCTGTGTCCA	60
CAP350	213956_at	0.28	GGGAAGTCCACATAGCGTCATTAAT	61
AIF1	215051_x_at	0.28	TTCAGCTACCCTGACTTTCTCAGGA	62
TRAF3	221571_at	0.28	GGCATGATGTCCGGTGATTTCTGTA	63
MBNL1	201152_s_at	0.27	ACTCTTGAGGGTTGATTATGCTGCA	64
FMNL1	204789_at	0.27	GGACCTCATCTCTGAGCTGAAACGG	65
TMEFF1	205122_at	0.27	GTTGGTGTTTAAAGATCTGAAGTGT	66
IL6R	205945_at	0.27	GAAGCACCATAACTTTGTTTAGCCC	67
SIVA	210792_x_at	0.27	ACAGCATGAGGCGGCCGGGAGCTG	68
MCAM	211340_s_at	0.27	GCTATGGTTATATTAGCACCAACT	69
POLR2I	212955_s_at	0.27	GGCCGACAACAGCTGCATCTATGTC	70
T3JAM	213888_s_at	0.27	TGAAAAAGGGTTTCTATTCTCTCTG	71
C1orf24	217967_s_at	0.27	AGTATCAGTCGGTGCAACAGTTGGC	72
COL5A2	221730_at	0.27	TGAAGTTGATCCTGAGACTCTTGAA	73
LAPTM5	201720_s_at	0.26	TACTCAGAGGTGTGACCCTCGCCAG	74
JARID1A	202040_s_at	0.26	GTCGTACTATCTTACTGAGCCACAG	75
CUGBP2	202156_s_at	0.26	AAGGCGTAACGAGTTCATCTTTCTT	76
PTPN7	204852_s_at	0.26	CCTTGATACCAGCTCTCTGTGGAAA	77
LCP2	205269_at	0.26	AAATCACTAAACCTCGTTTTCTCAG	78
RASA4	212706_at	0.26	AGCGTCCTTATCTTTCAGAGCTACA	79
FTL	212788_x_at	0.26	AAACCCAGACGCCATGAAAGCTGC	80
CD3D	213539_at	0.26	GGGAACACTGCTCTCAGACATTACA	81
EIF4A1	214805_at	0.26	CTTTTTCTGGGTCATGCTGCAACA	82
NKTR	215338_s_at	0.26	GATGGGGTGCATGTAGTCTTTGGAC	83
C1orf24	217966_s_at	0.26	GAAGGTGTGATCTGTGGGACTGTCT	84
C2orf33	219137_s_at	0.26	GTACGTTTTTACTCAGTTCATGCGT	85
TMEM22	219569_s_at	0.26	GCTTCTCGTGCTGCACATATTTCT	86
GIMAP6	219777_at	0.26	GTGAACAGACTTGAACTCCAGAGC	87
RAP1B	200833_s_at	0.25	ATCATTTTCAGGCTTCTGCAGCTGT	88
SRRM1	201225_s_at	0.25	GCATGTTGTTTGCCAGGACACTGTG	89
PWP1	201608_s_at	0.25	TTGTGCTTGCTCTTCAGATGGATGG	90
EDG1	204642_at	0.25	TAGCCAGGATCCTTGGTGTCTTAGG	91
CD47	211075_s_at	0.25	GCGGCGTGTATACCAATGCATGGCC	92
CG018	213375_s_at	0.25	GAATAACTTTTGCTGTTGTGCTAA	93
TPK1	221218_s_at	0.25	TGGCCCGCGTGATTGTGGCATTAA	94
COL5A2	221729_at	0.25	CATAACTGTTAGACTTCCCGTTTCT	95

Table 2. mRNA biomarkers of resistance to cisplatin. Dashes mean that the Affymetrix probeset has not been mapped to a specific gene. Affymetrix IDs refer to the array type HG-U133A.

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
SFN	33323_r_at	-0.48	TCAATAAAGTTCCCCTGTGACACTC	96
LISCH7	208190_s_at	-0.47	CTCCCCTATGATGGGCGGCTACTGG	97

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
EPB41L4B	220161_s_at	-0.47	ATCAGTTGATTCTTGTGCCATTTTT	98
MST1R	205455_at	-0.46	TGAGCCAGTGAGGGCAGTCCTGCAA	99
ITGB4	204990_s_at	-0.45	GCATCATCACCATAGAGTCCCAGGA	100
SFN	209260_at	-0.45	TCTTGCTCCAAAGGGCTCCGTGGAG	101
C20orf35	218094_s_at	-0.45	ATACGCCCTTGGCACAGTCGGATGA	102
SFN	33322_i_at	-0.45	GTCTGCTGGGTGTGACCATGTTTCC	103
TACSTD1	201839_s_at	-0.43	GTGCGTGGGACGAAGACATCTTTGA	104
C19orf21	212925_at	-0.42	TGGTCCCCTTCACCTGGGAGAAAAG	105
KRT8	209008_x_at	-0.41	GGGCCAAGCAGGACATGGCGCGGCA	106
JUP	201015_s_at	-0.4	AGCTTCAGACTCAAGTACCCATTCT	107
KRT18	201596_x_at	-0.4	GAGCTGCTGAGACGACGCTCACAGA	108
FA2H	219429_at	-0.39	GAGAAGCAGTTTGACGGACCTTGTG	109
MGAT4B	220189_s_at	-0.38	GGTGATTCTGAGCGAGATCTTCCTG	110
DSG2	217901_at	-0.37	GCAGCCTTGGAAACCTAACCTGCCT	111
LRP5	209468_at	-0.36	CCTGCAGCACCAGCGTGTGTGACAG	112
GJB3	215243_s_at	-0.36	ACTTGGCTCAGTGGAAAGCCCTCTTT	113
TACSTD2	202286_s_at	-0.35	ACATTGCCCGGAAACTCAGTCTATT	114
LAD1	203287_at	-0.35	GCTGTGGATCTGTTTGGCCAGGGTC	115
AGR2	209173_at	-0.35	GTTAGAGCCGATATCACTGGAAGAT	116
HTATIP2	209448_at	-0.35	AGATTTGTCAGCCCTATCTCAAACCT	117
LOC57228	209679_s_at	-0.35	AGGTCTTCCCAGAGGCTGGATACCA	118
BCL2L1	212312_at	-0.35	GTCTTCCCTACCTCAGGCAGGAAGG	119
GPX2	202831_at	-0.34	CTACCCTTATGATGACCCATTTTCC	120
SOX9	202935_s_at	-0.34	AAATGCTCTTATTTTCCAACAGCT	121
TPBG	203476_at	-0.34	GTGTATAGTGTTTTACCCTCTTCTT	122
LGALS4	204272_at	-0.34	TCATCAAGGGCTATGTGCCTCCCAC	123
PHLDA1	217996_at	-0.34	CCCCGCACCAGATCAAGTAGTTTGG	124
PLEK2	218644_at	-0.34	CCCTCCTACCAGATGACACAGACAA	125
TNFRSF21	218856_at	-0.34	TGTATGGTTTTACCTGGACACCGT	126
IER3	201631_s_at	-0.33	AACTCCGTCTGTCTACTGTGTGAGA	127
RAI3	203108_at	-0.33	CCCACTGGCCTGAATCTACACTGGA	128
BENE	209373_at	-0.33	ACATTACATCCGTGGATTCTCCTGC	129
MGC50853	212400_at	-0.33	GGCCCTGGGCCAGGGTGATTGGACT	130
RAI3	212444_at	-0.33	TTTAGCCCTCATGACTGTATTTTCT	131
CLIC3	219529_at	-0.33	ACACGCTGCAGATCGAGGACTTTCT	132
CLDN3	203954_x_at	-0.32	ACCGGCAGCCCTGGAAGGGGCACTT	133
FGFR4	204579_at	-0.32	TACCAGCAGGAGGTTCTGGGCCTCT	134
PPARG	208510_s_at	-0.32	CATCTTTCAGGGCTGCCAGTTTCGC	135
FBP1	209696_at	-0.32	GGGCTACGCCAAGGACTTTGACCCT	136
CPNE3	202119_s_at	-0.31	AATCTAGTCACCTAACCTTGTGGTT	137
AREG	205239_at	-0.31	ATTTCAAATTTCTGCATTACGGA	138
VIL1	205506_at	-0.31	AACACCTGTCCATTGAAGATTTAC	139
GATA6	210002_at	-0.31	GACATTCTTATGCTTCTTTTACAAC	140
TCF7L2	212761_at	-0.31	AATGTTTCCTAACAGTTGTGATGTT	141
PP1201	217730_at	-0.31	GGGTGAAGAGAGACTCGGTGCGGGC	142

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
FLJ20847	219053_s_at	-0.31	CGACCGCCTGTATGTTTGTGTAATT	143
GPR172A	222155_s_at	-0.31	AAGGCCTATCAGCTTCTATCAGCCC	144
ITGA6	201656_at	-0.3	GTCACCTGGTCTGTTTGCAATTTGATA	145
ZNF165	206683_at	-0.3	AGCTCAAACTTGCTAGGCATCAGA	146
FLNB	208613_s_at	-0.3	GCAGCAAAGCTGGCTCCAACATGCT	147
MCCC2	209623_at	-0.3	AAACACTATCTACTTCCTTTGTCAT	148
FLJ20273	218035_s_at	-0.3	GAGGATCATGCCCTTAGCAAGTACT	149
TMEM16A	218804_at	-0.3	AACATCATTTTAGCAAAGGCCAGGA	150
RAB11FIP1	219681_s_at	-0.3	TGTCCTTGTTACATTGAGGTTAAGA	151
SLC3A2	200924_s_at	-0.29	TCCCTACTGCATGGGGACTTCCACG	152
EFNA1	202023_at	-0.29	CCACCTTCACCTCGGAGGGACGGAG	153
SORL1	203509_at	-0.29	TAATTACACGTTACCGTCCAAGCA	154
PLS1	205190_at	-0.29	TTCCCTTTCTACCATTGATTTAAAT	155
GALIG	208949_s_at	-0.29	AGTACTGGTTGAACCTGACCACTTC	156
EHD1	209038_s_at	-0.29	AAATACATAAGCTAGTTTCTGTTCT	157
NR2F2	209120_at	-0.29	GTAACGTGATTGATTCAGTATCTTA	158
SERPINB1	213572_s_at	-0.29	AATACATCCGATGCGTAGATTCTTG	159
PCK2	202847_at	-0.28	AGAATGCTCGGGTGCTAGACTGGAT	160
ARF6	203311_s_at	-0.28	GGACGGACTCTATGAGGGGGCTCACA	161
TGFA	205016_at	-0.28	GGAATGACTCAAATGCCCCAAACCA	162
CST6	206595_at	-0.28	TCCTCTCAGCTCCTAAAGCACAACT	163
PXN	211823_s_at	-0.28	ACATGTTGCGACCCAAGTGTGGCGG	164
SORL1	212560_at	-0.28	TTTCAGATGGAGTACCAGCACCGAA	165
SLC39A4	219215_s_at	-0.28	TGGCACTCGCGGTTGGAGTCAGCGA	166
GCNT3	219508_at	-0.28	GGCCATCTATGGGACTGAACTTTGA	167
S100A11	200660_at	-0.27	GAAGAACTGGACACCAACAGTGAT	168
ITPR3	201189_s_at	-0.27	GCTGTAGCCAGTGCAGACCTCACTG	169
DHCR7	201790_s_at	-0.27	AGGTGTCCAGTACCTAATCACGCTC	170
TCIRG1	204158_s_at	-0.27	TTGCCGTGATGACCGTGGCTATCCT	171
NR2F2	209121_x_at	-0.27	GAATACGTTAGGAGCCAGTACCCCA	172
SLC25A1	210010_s_at	-0.27	GAAGCTGCTCAACAAAGTGTGGAAG	173
SERPINB6	211474_s_at	-0.27	GGAATGTCCCAGACAGACCTGTCTC	174
ARTN	216052_x_at	-0.27	CCTTCATGGACGTCAACAGCACCTG	175
LOC51123	218059_at	-0.27	GGCCCGGATATGGCTCGTGGACAGC	176
S100A14	218677_at	-0.27	AGGAGTCTCCACCAGAGGGAGGCTC	177
FCGRT	218831_s_at	-0.27	GAGCACCCTACTGCTGCATTGTGC	178
RAB20	219622_at	-0.27	ACTCTGACATTCTTGTCTCAAGC	179
SPDEF	220192_x_at	-0.27	CCAGCATTTCCAGAGCAGAGCCTAC	180
PNAS-4	221648_s_at	-0.27	GCGTGTCTTGAGTTCCATGCAAATT	181
PXN	201087_at	-0.26	AATGGTGACAGTCCAAACCACTCCA	182
TPD52L2	201379_s_at	-0.26	GGCCCTGCATGTCAGATGGCGTGGT	183
ALDH3A2	202054_s_at	-0.26	TGATCATAAATTCTCCCCAACTATA	184
ARF6	203312_x_at	-0.26	AAAGTTGCCAAGATGCTCCTTGTTG	185
GPA33	205929_at	-0.26	GTCTCACCCAAGTGCAGTTTACTAT	186

Gene	Affymetrix ID	Correlation	Affymetrix Probe Sequence	SEQ ID NO:
-	208540_x_at	-0.26	GACGGAGTTCCTAAGCTTCATGAAT	187
FLNB	208614_s_at	-0.26	TCAGCCTGGGCAGTCTTACCAAAT	188
TSPAN-1	209114_at	-0.26	TGCTGTGGCTTCACCAACTATACGG	189
CDH17	209847_at	-0.26	CCTTGACTCCTTTGGTATTTCACTG	190
SERPINB1	212268_at	-0.26	ACAGCAGGCATCGCAACTTTCTGCA	191
LCN2	212531_at	-0.26	CAAGAGCTACAATGTCACCTCCGTC	192
KIAA0984	213913_s_at	-0.26	GTTTGTCTCTTGTGTTCTGAAGGA	193
ACSL5	218322_s_at	-0.26	CTCTCTAGTTAGATATCTGACTTGG	194
MUC13	218687_s_at	-0.26	TCCAGCCTCGGGGTGTAGGTTTCTG	195
FAM11B	219253_at	-0.26	ACTCGTCTCACGCCGTGTTTGAGAT	196
SH2D3A	219513_s_at	-0.26	GCCAGAGTTCAAATGTGACTCCACC	197
ANXA2	201590_x_at	-0.25	CAAGCCCCTGTATTTTGCTGATCGG	198
TM4SF3	203824_at	-0.25	AGACCACAGATATCTTCTAGACATA	199
NT5E	203939_at	-0.25	GTCAGTGTAAATCATTCTTAAGCCC	200
TETRA	209215_at	-0.25	AAGGCTGTCAGGGCTTCTGTTTGTT	201
CTBP2	210835_s_at	-0.25	GTAGACACCTGCACGCATAGGATTG	202
SCD	211708_s_at	-0.25	TTGCCACTTCTTGCATATGCTGT	203
DNMBP	212838_at	-0.25	GCCATTCCAGAAGTAGCTTATCCTA	204
TMC5	219580_s_at	-0.25	CCAATACCCACCGTGATGACTTGA	205

Table 3. mRNA biomarkers of sensitivity to an sPLA₂ hydrolysable, cisplatin-containing liposome. Dashes mean that the Affymetrix probeset has not been mapped to a specific gene. Affymetrix IDs refer to the array type HG-U133A.

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
CALD1	212077_at	10861321835689.1	AATTCTCTGTTATCTTTACGAGGTA	206
COL6A2	209156_s_at	8535698909744.43	CACGAGAAGGACTATGACAGCCTGG	207
FERMT2	209210_s_at	5291552917682.63	TGATTTGCCACAATGTCCTTAAGTC	208
BNIP3	201849_at	5145685657339.48	GCTGAAGGCACCTACTCAGTATCTT	209
RAB31	217762_s_at	4734688539598.5	AGACCTGGCACTTCAGTAACTCAGC	210
COL5A2	221730_at	4647176466910.36	GACTCTTGAAGTAATGGCTGATCCT	211
MPO	203948_s_at	4518211644157.6	GGGACTTTGTCAACTGCAGTACACT	212
SRPX	204955_at	4340511505629.07	CCTTCTTTACTCCATCATGGCTGG	213
ARHGDIB	201288_at	4263392661779.67	ATCACTAACAGGTCTTTGACTCAGG	214
TMEM47	209656_s_at	4156685173988.01	GAATTCATGGTATCCTGGTTATTTT	215
CSRP2	207030_s_at	3960151282910.27	AACTACTGTGAAATTCTACCAGCAT	216
DPYSL3	201431_s_at	3876388962016.02	GACACCTGAGCCTGGATTTTCACTC	217
HTRA1	201185_at	3845854940391.73	TCAAACGGCCGAAGTTGCCTCTTTT	218
SLC39A6	202088_at	3547148987590.88	ATACTAGGCCTGTCTGTGGCATTCT	219
LAT2	221581_s_at	3545380986375.43	GGATTTAGGATAAGCTGTCACCCAG	220
ENAH	217820_s_at	3385939870513.75	GGTCAGCAACCTCTTTTGATTTTGT	221
RPS4Y1	201909_at	3384951327956.31	GACAGGTGAACATTCCGCCTGGTC	222
DOCK2	213160_at	3367491706976.35	GATTCCTGAACTCAAGGTACCAGCA	223
COL1A1	202311_s_at	3222960671378.67	TGTTCTTTTTTGTCAAAGTCTATT	224
GMFG	204220_at	3013566458581.29	AGGTGTTTCGAAATCCGCACCACTGA	225
CYR61	201289_at	2999506373414.97	GTGGAGTTGATGACTTTCTGTTTTT	226

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
RHOB	212099_at	2978300392812.93	TGCAGGTCATGCACACAGTTTTGAT	227
COR01A	209083_at	2968352455386.15	GCTCCAGAAGCGCTTGGACAGGCTG	228
ID4	209291_at	2948241975028.96	GGCATAATGGCAAATCCTTCAAGCA	229
RARRES2	209496_at	2907180844659.6	CCCCATAGAGACCCAAAGTTCTGCGG	230
SOX4	201417_at	2862450307972.36	GTAAACCACATCTTTTTTGCACCTT	231
NID1	202007_at	2798544570884.12	CACTTTTTGTATTATCGTTTCATA	232
CALD1	201616_s_at	2776573094080.12	GACGCAGGACGAGCTCAGTTGTAGA	233
SERPINE2	212190_at	2767126943194.04	TGTTGTGCAGTGTGCCTGTCACTAC	234
CTSL1	202087_s_at	2681524741399.96	CACCTACTGACTTTGCATTTTCGTT	235
C3orf14	219288_at	2679480387909.32	GGTGGTTTCTCTTGAGACTCGTTAC	236
DKK3	202196_s_at	2608335983440.84	TTGGCAGTTGCATTAGTAACCTTGA	237
SCRN1	201462_at	2582074623391.62	TCATGTGCACATGCCGTTGCAGCAC	238
MT1M	217546_at	2555792977629.17	CGTTGGAGAACTGCAGCTGCTGTGC	239
PLAU	205479_s_at	2529115320523.6	AGCAGCTGAGGTCTCTTGAGGGAGC	240
NREP	201310_s_at	2514590941976.06	CATTGGCCTGAGTTTCTTGTGCATT	241
HLA-B	208729_x_at	2501423496784.03	GAGCCTACCTGGAGGGCGAGTGCCT	242
PTPRC	212588_at	2494855639496.51	GTTTTCAATTTTGCATGCTCGATTA	243
HDGFRP3	209524_at	2438222715080.89	TTATGTGTACATTATTGTTGCTATT	244
CELF2	202157_s_at	2427790438608.2	CTTCCCGGTCACTGGTAACAATAGC	245
SFRP1	202037_s_at	2413217767593.8	GTACCTGTGGGTTAGCATCAAGTTC	246
HLA-B	211911_x_at	2358346288074.42	CTGAGAGCCTACCTGGAGGGCCTGT	247
LOX	215446_s_at	2354236167712.24	TTGGGCCTTTTATCTGTCTATCCAT	248
CLU	208791_at	2341547177698.15	CAGTGTGACAAGTGCCGGGAGATCT	249
SH3BGRL	201312_s_at	2249866543302.91	AGAATCTTTTCTATGCCTCTATTCC	250
INHBA	210511_s_at	2238550007854.02	GCCATATAGCAGGCACGTCCGGGTC	251
MMP1	204475_at	2203074303300.14	GGCAAGGGATAACTCTTCTAACACA	252
WIPF1	202664_at	2194537285288.12	TTCCCTCCTTATAGTCAAGGACCGT	253
ADAMTS1	222162_s_at	2144423953975.08	AATAACGCAAATGGCTTCCTCTTTC	254
THY1	208850_s_at	2141423198789.74	GGCCTAGCACGGACATGGTCTGTCC	255
UHL1	201387_s_at	2140899985376.98	TGATGGACGAATGCCTTTTCCGGTG	256
MYH10	212372_at	2139390916542.17	GATCCTCTGCAATGTGCTTGAAAAC	257
TYMS	202589_at	2131876162229.91	TCACAAGCTATTCCTCAAATCTGA	258
HCLS1	202957_at	2089924252642.24	TGATGAGCTTTCCTTTGATCCGGAC	259
HLA-B	209140_x_at	2085546519988.6	GAGACAGCTGTCTTGTGAGGGACTG	260
IFI16	208966_x_at	2061722348570.95	TACACACCACCATATATACTAGCTG	261
PRKCB	207957_s_at	2037662863122.06	GTGTAGGTGAATGCAAACCTCATCG	262
BNIP3	201848_s_at	2008580245730.46	TTCCTCTTTAAACACCCGAAGCGCA	263
TUSC3	213423_x_at	1987545095813.27	AACTGTTCTGACTTTATACTATTT	264
WNT5A	205990_s_at	1982235386738.35	GCATAATGATATTCACATCCCCTCA	265
CALD1	201617_x_at	1981280027254.5	TGTTGTTTCTGCACTTTATAATAAA	266
HLA-C	216526_x_at	195599731784.71	AGAGGTGGGGCTGGATGTCTCCATC	267
IL1R1	202948_at	1955342562611.76	AAGTGCAAAGTTATCCCCATCTTC	268
AUTS2	212599_at	1927738178390.84	TACTTACACCCAAACAGATCCTGAA	269
THBS2	203083_at	1912997768879.9	TTGCGTGTGGAGCTGTATTCCCGAG	270

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
CHRD1	209763_at	1895325557387.3	CCCTTTCACTGTTCTCACAGGACAT	271
SFPQ	214016_s_at	1886539698542.15	GTTGGCTGATATTGGAGTGCTCATT	272
CXCL12	209687_at	1857308403453.12	CAGCAGGGTTTCAGGTTCCAATCAG	273
HOXC6	206858_s_at	1831591158444.48	CTGTATTGTGGTCTCTGTATTTAT	274
PLAGL1	209318_x_at	1827870818957.99	ACATCCAAAATGACGGCTGCTATAT	275
RDX	212397_at	1815278384492.07	GTGGACCTACTATTTCATGTTTTGA	276
HNRNPH1	213619_at	1813815711802.08	GCTTAAACTTACGTGCCTTACAGGT	277
KRAS	214352_s_at	1802923545775.42	CATGCAGACTGTTAGCTTTTACCTT	278
IL8	211506_s_at	1788698391848.43	GTCAGTGCATAAAGACATACTCCAA	279
FAM46A	221766_s_at	1787987145165.06	GGAGTCCTATTTGCAGAACCACTTT	280
QKI	212265_at	1787672566876.18	ATAACCAACCTATTGCCTATGAGAA	281
CD53	203416_at	1777870731216.97	CGAATTAGTCTCCAGCCTCTAAATA	282
LAPTM5	201720_s_at	1763708973603.65	TCGGGTCTCTCCATAATTCAGCCCA	283
FOXG1	206018_at	1752375753099.1	ACGATTGCCTTCAGTTTGTGTTGTG	284
MST4	218499_at	1732353014841.79	AATTCTTTTTATTGGTGCCTATATT	285
GAPDH	AFFX- HUMGAPDH/M33197_M_at	1692594771893.01	AAGCTCACTGGCATGGCCTTCCGTG	286
TUBB2B	214023_x_at	1672014039622.35	GAGATATTTCTGAATTACTGTTGTA	287
GAPDH	212581_x_at	1649610188507.54	TTTGACGCTGGGGCTGGCATTGCC	288
CEBPD	203973_s_at	1623762464226.23	GGACAGCAGACTGCCGGTAACGCGC	289
PLAU	211668_s_at	1604895332856.59	GCTCTGAAGTCACCACCAAATGCT	290
CAV1	203065_s_at	1604187716818.41	GGTGCCAATTTCAAGTTCCAAGTTG	291
GAPDH	AFFX- HUMGAPDH/M33197_3_at	1601834913853.31	TAGGGAGCCGCACCTTGTCATGTAC	292
---	213158_at	1597303398144.17	ACGTATATTTACCTGTGACTTGTAT	293
ARHGEF6	209539_at	1586970619512.16	TAAACTGCTGCCCGTAGAGGCCTTT	294
PRKCB	209685_s_at	1580850725622.13	TGGATGTTAGCGGTACTCTTCCACT	295
SRGN	201859_at	1549790579490.15	TTTTCTGGATATCTGTGATTTTC	296
TLE4	204872_at	1549011037374.17	ACTGTGCGTTGTACATAGTTCTAAT	297
LOC100506558 MATN2	202350_s_at	1544181853329.71	GAACACTGGCCATAGGAAATGCTGT	298
BHLHE40	201170_s_at	1537151135133.25	GATCCTTTCTGTAGGCTAATTCCTC	299
SGCE	204688_at	1519398433064.38	AACGCAGCAGAACTTGCCACATCAG	300
---	222288_at	1511518722955.02	GAAGCTTGGCTTTAGTGGTAGAATG	301
PCBP2	204031_s_at	1507948521040.68	AGCCTGGCTCAATATCTAATCAATG	302
TFAP2A	204653_at	1493277682055.65	GAACCTCAAACATTTGGGACCACCT	303
SPON1	209436_at	1472949317341.51	CCACCCTAGTGTCTCATGTTTGTAT	304
COL4A2	211966_at	1468135692764.19	TGGTGATGTCTGCTACTATGCCAGC	305

Table 4. mRNA biomarkers of sensitivity to the liposome. Dashes mean that the Affymetrix probeset has not been mapped to a specific gene. Affymetrix IDs refer to the array type HG-U133A.

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
KRT18	201596_x_at	-22426211704708.5	AAGCTGGAGGCTGAGATCGCCACCT	306
LGALS3	208949_s_at	-11456296973610.8	CACCTTAACCCACGCTTCAATGAGA	307

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
DSP	200606_at	-10269594517738.5	TGGAATGAGTCTCCTTTAGTTTCAG	308
IGFBP4	201508_at	-8435796702432.14	AGAGACATGTACCTTGACCATCGTC	309
SPINT2	210715_s_at	-8294729535462.05	TGGAAATCCTCTAGGAGGCTCCTCC	310
CDH1	201131_s_at	-7786548077136.61	TGTGTGGGTGCTGATAATTGTGTAT	311
DSG2	217901_at	-7061991934030.4	TACTCTCCATCATCTAGAATTGTT	312
RAB25	218186_at	-6195270978776.59	GCACCCTCAGGGTCTTAAGGTCTTC	313
PTPRF	200636_s_at	-6131832886305.69	GTACACAGTCTGTTTTCTATTTGTT	314
SOX9	202936_s_at	-5835576205162.92	TGGGCTGCCTTATATTGTGTGTGTG	315
LYZ	213975_s_at	-5458342909996.32	TAACCCAGACTTAATCTTGAATGAT	316
IER3	201631_s_at	-5365171123958.73	GAGACTTCGGCGGACCATTAGGAAT	317
PERP	217744_s_at	-5097068499548.16	ATGCACGTGAACTTAACACTTTAT	318
SOX9	202935_s_at	-5050052756141.07	AGTTGAACAGTGTGCCCTAGCTTTT	319
ATP1B1	201243_s_at	-4753436553865.35	GATCTTGATTTCAGTCAGGTTAAAA	320
IFI27	202411_at	-4636709898452.9	CCAAAGTGGTCAGGGTGGCCTCTGG	321
PHLDA2	209803_s_at	-4623467982538.76	GGACGAGTCGGACCGAGGCTAGGAC	322
CTTN	201059_at	-4563342040423.69	ATTTGTGGCCACTCACTTTGTAGGA	323
SFN	209260_at	-4455761701170.73	TCTTGCTCCAAAGGGCTCCGTGGAG	324
MALL	209373_at	-4327230558082.54	CTCCTCCATGAGTCTGACATCTCGG	325
S100A11	200660_at	-4322815561525.15	GGTTGAGGAGAGGCTCCAGACCCGC	326
TSPAN13	217979_at	-4261036366041.2	ACAGCAACTTGTCAAACCTAAGCAT	327
AKR1C3	209160_at	-4207721689216.25	ACGCAGAGGACGTCTCTATGCCGGT	328
FAT1	201579_at	-4082641838983.11	GTAGTCATTCACTTCTAGCTGTACA	329
DSTN	201021_s_at	-4020978397283.39	GTAGCTGATGAAGTATGTGCGATTT	330
EFEMP1	201842_s_at	-3992766849062.55	GATGATCTTCTGTGGTGCTTAAGGA	331
TFF3	204623_at	-3853023482644	CTGTGATTGCTGCCAGGCACTGTTT	332
HSPB1	201841_s_at	-3835026328384.26	TTCACGCGGAAATACACGCTGCCCC	333
SDC1	201286_at	-3731984524505.92	TCATCTGCTGGTCCGTGGGACGGTG	334
PLAC8	219014_at	-3720610591317.68	GAAGGAGAGCCATGCGTACTTTCTA	335
TPBG	203476_at	-3655713541808.07	GTGTATAGTGTTTTACCCTCTTCTT	336
LCN2	212531_at	-3340240709988.96	CAGGACTTTTGTTCAGGTTGCCAG	337
CEACAM6	203757_s_at	-3279054777343.26	GTGCAGTTTCTGACACTTGTTGTTG	338
ELF3	210827_s_at	-3241469160886.13	GGGAGCACCGTGATGGAGAGGACAG	339
CLDN3	203953_s_at	-3192796314939.69	AAGGCCAAGATCACCATCGTGGCAG	340
TPD52L1	203786_s_at	-3049121447681.89	TATTCAAATGGCCCCCTCCAGAAAGT	341
VAMP8	202546_at	-2969692217517	AAGCCACATCTGAGCACTTCAAGAC	342
C1orf106	219010_at	-2931724791122.81	GTTCCAAGAACTCTGGTGTCTGACC	343
RBM47	218035_s_at	-2891974033193.95	GAGGATCATGCCCTTAGCAAGTACT	344
C3	217767_at	-2846605120573.62	GGTCTACGCCTATTACAACCTGGAG	345
CAPN2	208683_at	-2829130992700.86	AATCGTTCTCCTTACAATCAAGTTC	346
ERBB3	202454_s_at	-2788407249074.31	GGAAGTGGCTCTTATGTGTGCCCTT	347
SLPI	203021_at	-2755718313124.09	TCTGTCTCCTAAGAAATCTGCCCA	348
SPATS2L	222154_s_at	-2729322838596.83	GAGGCTCAGTTAGCAACCTGTGTTG	349
ERBB2	216836_s_at	-2698032874395.93	AGACTGTCCCTGAAACCTAGTACTG	350
SERPINB1	212268_at	-2694341115802.62	ACTTTCTGCATGTTGATGCCCGAAG	351
CEACAM6	211657_at	-2643169692661.57	GTTCTTGATTGTATTGCCAGGGG	352

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
AKR1B10	206561_s_at	-2617913243059.4	AAAAACCGCAGCCAGGTTCTGATC	353
ID1	208937_s_at	-2607302720347.48	GACATGAACGGCTGTTACTCACGCC	354
PPAP2C	209529_at	-2576535604785.95	TGTTCTTGCGCTGTATGTGCAGGC	355
AQP3	39248_at	-2561344001860.94	CTTCTACAGGCTTTTGGGAAGTAGG	356
PODXL	201578_at	-2559443301040.98	TGGAGGACACAGATGACTCTTTGGT	357
PRR15L	219127_at	-2483388299723.69	GAGTGGGTGGGGAATTTTCTCCTCT	358
EMP2	204975_at	-2470436470609.79	CTGCACCTTCATCAGCGGCATGATG	359
MYO10	201976_s_at	-2463058577194.03	TATAAACCACTCTTCAACAGCTGGC	360
SERPINB1	213572_s_at	-2374385129062.88	AATACATCCGATGCGTAGATTCTTG	361
SDC4	202071_at	-2371552687950.61	TGGCTTAGCCTGGGCAGGTCGTGTC	362
CRABP2	202575_at	-2354608471952.81	GAGCAGGGTCTCTCTAAAGGGGACT	363
HTATIP2	209448_at	-2354028532889.45	GTCTCTGAGTTACAAAAGTGCTAAT	364
DBNDD2 SYS1 SYS1-DBNDD2	218094_s_at	-2352744142308.53	ACCAGTTTTTGGCTTACTCCTGAGA	365
ESRP1	219121_s_at	-2312028194710.22	TTGTCTACACTCAGGCTGCAGTATT	366
HSD17B11	217989_at	-2304068718020.79	TCCTGAGAGATACCTCACATTCCAA	367
GFPT1	202722_s_at	-2272343431090.56	GGTTAGCCTTAGTTTCTCAGACTTG	368
S100A14	218677_at	-2240432231078.46	TGTCCTCATCTCTGCAAAGTTCAGC	369
IGFBP7	201162_at	-2225724813680	TTCCCAAGGACAGGCTTCAGCATCA	370
PTPRF	200637_s_at	-2190473907894.45	CTCCTACGCAGATGCTGTCACTGGC	371
HMGA1	206074_s_at	-2178312788057.87	TGAGCAAGGGGGCCCGAATCGACCA	372
YWHAZ	200641_s_at	-2145016988259.93	AAGCCTGCTCTCTTGCAAAGACAGC	373
SCD	200832_s_at	-2143962895648.8	TAACATAAGGTGCCTCAGTTTTCC	374
SH3YL1	204019_s_at	-2139236372988.65	CATATGGCATCTCTCAACTTTTCTT	375
UCP2	208998_at	-2139031352031.13	GAAAGTTCAGCCAGAATCTTCGTCC	376
F3	204363_at	-2113802654784.93	GGGCAGCTTCCTAATATGCTTTACA	377
AZGP1	209309_at	-2089576575474.55	GCCTGTCTTGAGTAGACTTGGACCC	378
LIMCH1	212327_at	-2089195209441.08	GATCCACCTCATATGTGAGTCCGTC	379
PLA2G2A	203649_s_at	-2069037053701.26	CGCTGCTGTGTCACTCATGACTGTT	380
ITGB5	201125_s_at	-2028321449243.62	GCCTGTTGAAGGTACATCGTTTGCA	381
ABCC3	208161_s_at	-2007168680009.07	TCTCCCGATTCCCAACTGAGTGTTA	382
DDR1 MIR4640	207169_x_at	-2000582844983.07	AGGCAATTTTAATCCCCTGCACTAG	383
GATA3	209604_s_at	-1995114130212.84	GGACAAACTGCCAGTTTTGTTTCCT	384
CYB561	209163_at	-1981172434786.63	GTTCTTCAATCAGCTGGCACACACT	385
C10orf116	203571_s_at	-1962923571527.29	ACCACCCAGGAAACCATCGACAAGA	386
PTPRF	200635_s_at	-1924144465806.05	AAGGACAGAACATTGCCTTCCTCGT	387
DKK1	204602_at	-1893211415469.31	GGATATACAAGTTCTGTGGTTTCAG	388
SERPINB5	204855_at	-1863934443254.52	GTGGTTGGCACTAGACTGGTGGCAG	389
ARHGAP29	203910_at	-1818117319379.63	ATGTACTTGTTCTACCTGGATTGTC	390
GAS6	202177_at	-1817533234900.07	CGCGGCTGCATGACACTGGAGGTCA	391
LAMB3	209270_at	-1817170377879.96	GGTGCCCGGATCCAGAGTGTGAAGA	392
KLF5	209212_s_at	-1814910338390.4	CTCCATCCTATGCTGCTACAATTGC	393
MAOA	212741_at	-1811716715860.48	TGAATGCCAGTCCAGATGTGCCTAG	394
NET1	201830_s_at	-1789348130490.25	TTACATTCATTTAACCTGCCGATTA	395
CYBA	203028_s_at	-1775049034494.02	CACCCAGTGGTACTTTGGTGCCTAC	396

Gene	Affymetrix ID	Covariance	Affymetrix Probe Sequence	SEQ ID NO:
TGM2	201042_at	-1772139742186.19	AGTGCTGGTCACTAACCAACAAGGT	397
ALDH2	201425_at	-1757839520621.92	CTCTCTGAAACGCTTCCTATAACTC	398
HSPA1A HSPA1B	200799_at	-1730673434053.48	TTGTCAGTTCTCAATTTCTGTGTT	399
JUP	201015_s_at	-1729139912998.84	ATTATCGCTTTATGTTTTGGTTAT	400
HSPA1A HSPA1B	200800_s_at	-1722098969341.57	GGGGCTCAAGGGCAAGATCAGCGAG	401
F11R	221664_s_at	-1642391094616.93	GAATAGGTATCTTGAGCTTGGTTCT	402
HBG1 HBG2 LOC10065 3006 LOC10065 3319	204419_x_at	-1595966820539.76	ACACTCGCTTCTGGAACGTCTGAGG	403
KLF4	221841_s_at	-1553919884310.19	AATCTATATTTGTCTCCGATCAAC	404
CA12	214164_x_at	-1551710888005.42	ACAAGGCCCAGGCTGGGGCCAGGGC	405

[0094] The biomarker COL5A2 (SEQ ID NO 73 or 211) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker COL5A2 (SEQ ID NO 73 or 211) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of COL5A2 (SEQ ID NO 73 or 211) in the patient sample may then be compared, e.g., to the level of COL5A2 (SEQ ID NO 73 or 211) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0095] The biomarker COL5A2 (SEQ ID NO 73 or 211) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker COL5A2 (SEQ ID NO 73 or 211). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0096] The biomarker ITGA4 (SEQ ID NO: 1) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker ITGA4 (SEQ ID NO: 1) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of ITGA4 (SEQ ID NO: 1) in the patient sample may then

be compared, e.g., to the level of ITGA4 (SEQ ID NO: 1) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0097] The biomarker ITGA4 (SEQ ID NO: 1) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ITGA4 (SEQ ID NO: 1). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0098] The biomarker MSN (SEQ ID NO: 2) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker MSN (SEQ ID NO: 2) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of MSN (SEQ ID NO: 2) in the patient sample may then be compared, e.g., to the level of MSN (SEQ ID NO: 2) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0099] The biomarker MSN (SEQ ID NO: 2) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor

sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker MSN (SEQ ID NO: 2). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0100] The biomarker FAM46A (SEQ ID NO: 3 or 280) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker FAM46A (SEQ ID NO: 3 or 280) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of FAM46A (SEQ ID NO: 3 or 280) in the patient sample may then be compared, e.g., to the level of FAM46A (SEQ ID NO: 3 or 280) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0101] The biomarker FAM46A (SEQ ID NO: 3 or 280) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker FAM46A (SEQ ID NO: 3 or 280). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0102] The biomarker ITGB2 (SEQ ID NO: 4) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker ITGB2 (SEQ ID NO: 4) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of ITGB2 (SEQ ID NO: 4) in the patient sample may then be compared, e.g., to the level of ITGB2 (SEQ ID NO: 4) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0103] The biomarker ITGB2 (SEQ ID NO: 4) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ITGB2 (SEQ ID NO: 4). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ITGB2 (SEQ ID NO: 4). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0104] The biomarker DOCK2 (SEQ ID NO: 5 or 223) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker DOCK2 (SEQ ID NO: 5 or 223) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of DOCK2 (SEQ ID NO: 5 or 223) in the patient sample may then be compared, e.g., to the level of DOCK2 (SEQ ID NO: 5 or 223) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0105] The biomarker DOCK2 (SEQ ID NO: 5 or 223) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker DOCK2 (SEQ ID NO: 5 or 223). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0106] The biomarker EVL (SEQ ID NO: 6) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker EVL (SEQ ID NO: 6) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or

a device (e.g., a microarray). As is described above, the level of EVL (SEQ ID NO: 6) in the patient sample may then be compared, e.g., to the level of EVL (SEQ ID NO: 6) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0107] The biomarker EVL (SEQ ID NO: 6) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker EVL (SEQ ID NO: 6). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0108] The biomarker SACS (SEQ ID NO: 7) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SACS (SEQ ID NO: 7) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SACS (SEQ ID NO: 7) in the patient sample may then be compared, e.g., to the level of SACS (SEQ ID NO: 7) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0109] The biomarker SACS (SEQ ID NO: 7) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target

sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SACS (SEQ ID NO: 7). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0110] The biomarker PTPRCAP (SEQ ID NO: 8) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to the liposome. The level of the biomarker PTPRCAP (SEQ ID NO: 8) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of PTPRCAP (SEQ ID NO: 8) in the patient sample may then be compared, e.g., to the level of PTPRCAP (SEQ ID NO: 8) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0111] The biomarker PTPRCAP (SEQ ID NO: 8) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker PTPRCAP (SEQ ID NO: 8). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0112] The biomarker EBI2 (SEQ ID NO: 9) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker EBI2 (SEQ ID NO: 9) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of EBI2 (SEQ ID NO: 9) in the patient sample may then be compared, e.g., to the level of EBI2 (SEQ ID NO: 9) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0113] The biomarker EBI2 (SEQ ID NO: 9) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN

(SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISF (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker EBI2 (SEQ ID NO: 9). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0114] The biomarker PTPRC (SEQ ID NO: 10, 18, 25, or 243) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker PTPRC (SEQ ID NO: 10, 18, 25, or 243) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of PTPRC (SEQ ID NO: 10, 18, 25, or 243) in the patient sample may then be compared, e.g., to the level of PTPRC (SEQ ID NO: 10, 18, 25, or 243) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0115] The biomarker PTPRC (SEQ ID NO: 10, 18, 25, or 243) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISF (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker PTPRC (SEQ ID NO: 10, 18, 25, or 243). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0116] The biomarker ANP32E (SEQ ID NO: 11) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the

biomarker ANP32E (SEQ ID NO: 11) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of ANP32E (SEQ ID NO: 11) in the patient sample may then be compared, e.g., to the level of ANP32E (SEQ ID NO: 11) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0117] The biomarker ANP32E (SEQ ID NO: 11) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ANP32E (SEQ ID NO: 11). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0118] The biomarker SFPQ (SEQ ID NO: 12, 38 or 272) may be used to assess a cancer patient's (e.g. a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SFPQ (SEQ ID NO: 12, 38 or 272) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SFPQ (SEQ ID NO: 12, 38 or 272) in the patient sample may then be compared, e.g., to the level of SFPQ (SEQ ID NO: 12, 38 or 272) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0119] The biomarker SFPQ (SEQ ID NO: 12, 38 or 272) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318),

ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SFPQ (SEQ ID NO: 12, 38 or 272). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0120] The biomarker C1QR1 (SEQ ID NO: 13) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker C1QR1 (SEQ ID NO: 13) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of C1QR1 (SEQ ID NO: 13) in the patient sample may then be compared, e.g., to the level of C1QR1 (SEQ ID NO: 13) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0121] The biomarker C1QR1 (SEQ ID NO: 13) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker C1QR1 (SEQ ID NO: 13). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0122] The biomarker FNBP1 (SEQ ID NO: 14 or 28) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker FNBP1 (SEQ ID NO: 14 or 28) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of FNBP1 (SEQ ID NO: 14 or 28) in the patient sample may then be compared, e.g., to the level of FNBP1 (SEQ ID NO: 14 or 28) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0123] The biomarker FNBP1 (SEQ ID NO: 14 or 28) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ

ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker FNBP1 (SEQ ID NO: 14 or 28). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0124] The biomarker SFN (SEQ ID NO: 96 or 324) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SFN (SEQ ID NO: 96 or 324) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SFN (SEQ ID NO: 96 or 324) in the patient sample may then be compared, e.g., to the level of SFN (SEQ ID NO: 96 or 324) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0125] The biomarker SFN (SEQ ID NO: 96 or 324) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SFN (SEQ ID NO: 96 or 324). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0126] The biomarker LISCH7 (SEQ ID NO: 97) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker LISCH7 (SEQ ID NO: 97) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of LISCH7 (SEQ ID NO: 97) in the patient sample may then be compared, e.g., to the level of LISCH7 (SEQ ID NO: 97) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0127] The biomarker LISCH7 (SEQ ID NO: 97) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LISCH7 (SEQ ID NO: 97). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0128] The biomarker EPB41L4B (SEQ ID NO: 98) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker EPB41L4B (SEQ ID NO: 98) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of EPB41L4B (SEQ ID NO: 98) in the patient sample may then be compared, e.g., to the level of EPB41L4B (SEQ ID NO: 98) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0129] The biomarker EPB41L4B (SEQ ID NO: 98) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214),

TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker EPB41L4B (SEQ ID NO: 98). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0130] The biomarker MST1R (SEQ ID NO: 99) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker MST1R (SEQ ID NO: 99) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of MST1R (SEQ ID NO: 99) in the patient sample may then be compared, e.g., to the level of MST1R (SEQ ID NO: 99) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0131] The biomarker MST1R (SEQ ID NO: 99) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker MST1 R (SEQ ID NO: 99). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0132] The biomarker ITGB4 (SEQ ID NO: 100) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker ITGB4 (SEQ ID NO: 100) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of ITGB4 (SEQ ID NO: 100) in the patient sample may then be compared, e.g., to the level of ITGB4 (SEQ ID NO: 100) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0133] The biomarker ITGB4 (SEQ ID NO: 100) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ITGB4 (SEQ ID NO: 100). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0134] The biomarker DBNDD2 (SEQ ID NO: 102 or 365) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker DBNDD2 (SEQ ID NO: 102 or 365) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of DBNDD2 (SEQ ID NO: 102 or 365) in the patient sample may then be compared, e.g., to the level of DBNDD2 (SEQ ID NO: 102 or 365) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome and, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0135] The biomarker DBNDD2 (SEQ ID NO: 102 or 365) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker DBNDD2 (SEQ ID NO: 102 or 365). In particular, the patient is determined to be responsive to the liposome if the

level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0136] The biomarker TACSTD1 (SEQ ID NO: 104) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker TACSTD1 (SEQ ID NO: 104) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of TACSTD1 (SEQ ID NO: 104) in the patient sample may then be compared, e.g., to the level of TACSTD1 (SEQ ID NO: 104) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0137] The biomarker TACSTD1 (SEQ ID NO: 104) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker TACSTD1 (SEQ ID NO: 104). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0138] The biomarker MISP (SEQ ID NO: 105) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker MISP (SEQ ID NO: 105) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of MISP (SEQ ID NO: 105) in the patient sample may then be compared, e.g., to the level of MISP (SEQ ID NO: 105) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0139] The biomarker MISP (SEQ ID NO: 105) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), KRT8 (SEQ

ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker MISP (SEQ ID NO: 105). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0140] The biomarker KRT8 (SEQ ID NO: 106) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker KRT8 (SEQ ID NO: 106) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of KRT8 (SEQ ID NO: 106) in the patient sample may then be compared, e.g., to the level of KRT8 (SEQ ID NO: 106) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0141] The biomarker KRT8 (SEQ ID NO: 106) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker KRT8 (SEQ ID NO: 106). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0142] The biomarker JUP (SEQ ID NO: 107 or 400) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker JUP (SEQ ID NO: 107 or 400) may be assessed using nucleic acid amplification methods (e.g., PCR,

such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of JUP (SEQ ID NO: 107 or 400) in the patient sample may then be compared, e.g., to the level of JUP (SEQ ID NO: 107 or 400) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0143] The biomarker JUP (SEQ ID NO: 107 or 400) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker JUP (SEQ ID NO: 107 or 400). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0144] The biomarker KRT18 (SEQ ID NO: 108 or 306) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker KRT18 (SEQ ID NO: 108 or 306) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of KRT18 (SEQ ID NO: 108 or 306) in the patient sample may then be compared, e.g., to the level of KRT18 (SEQ ID NO: 108 or 306) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0145] The biomarker KRT18 (SEQ ID NO: 108 or 306) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g.,

a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker KRT18 (SEQ ID NO: 108 or 306). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0146] The biomarker FA2H (SEQ ID NO: 109) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker FA2H (SEQ ID NO: 109) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of FA2H (SEQ ID NO: 109) in the patient sample may then be compared, e.g., to the level of FA2H (SEQ ID NO: 109) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0147] The biomarker FA2H (SEQ ID NO: 109) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFβ (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker FA2H (SEQ ID NO: 109). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0148] The biomarker MGAT4B (SEQ ID NO: 110) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker MGAT4B (SEQ ID NO: 110) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of MGAT4B (SEQ ID NO: 110) in the patient sample may then be compared, e.g., to the level of MGAT4B (SEQ ID NO: 110) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0149] The biomarker MGAT4B (SEQ ID NO: 110) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223),

EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA2 may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker MGAT4B (SEQ ID NO: 110). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0150] The biomarker DSG2 (SEQ ID NO: 111 or 312) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker DSG2 (SEQ ID NO: 111 or 312) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of DSG2 (SEQ ID NO: 111 or 312) in the patient sample may then be compared, e.g., to the level of DSG2 (SEQ ID NO: 111 or 312) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0151] The biomarker DSG2 (SEQ ID NO: 111 or 312) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker COL5A2 (SEQ ID NO 73 or 211). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0152] The biomarker LRP5 (SEQ ID NO: 112) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker LRP5 (SEQ ID NO: 112) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of LRP5 (SEQ ID NO: 112) in the patient sample may then be compared, e.g., to the level of LRP5 (SEQ ID NO: 112) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0153] The biomarker LRP5 (SEQ ID NO: 112) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1 R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LRP5 (SEQ ID NO: 112). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0154] The biomarker SFRS7 (SEQ ID NO: 19 or 54) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SFRS7 (SEQ ID NO: 19 or 54) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SFRS7 (SEQ ID NO: 19 or 54) in the patient sample may then be compared, e.g., to the level of SFRS7 (SEQ ID NO: 19 or 54) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0155] The biomarker SFRS7 (SEQ ID NO: 19 or 54) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6

(SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SFRS7 (SEQ ID NO: 19 or 54). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0156] The biomarker CAP350 (SEQ ID NO: 20 or 61) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker CAP350 (SEQ ID NO: 20 or 61) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of CAP350 (SEQ ID NO: 20 or 61) in the patient sample may then be compared, e.g., to the level of CAP350 (SEQ ID NO: 20 or 61) in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0157] The biomarker CAP350 (SEQ ID NO: 20 or 61) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker CAP350 (SEQ ID NO: 20 or 61). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0158] The biomarker CALD1 (SEQ ID NO: 206) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The expression level of the biomarker CALD1 (SEQ ID NO: 206) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the expression level of CALD1 (SEQ ID NO: 206) in the patient sample may then be compared, e.g., to the expression level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0159] The biomarker CALD1 (SEQ ID NO: 206) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker CALD1 (SEQ ID NO: 206). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0160] The biomarker COL6A2 (SEQ ID NO: 207) may be used to assess a cancer patient's (e.g. a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The expression level of the biomarker COL6A2 (SEQ ID NO: 207) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the expression level of COL6A2 (SEQ ID NO: 207) in the patient sample may then be compared, e.g., to the expression level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0161] The biomarker COL6A2 (SEQ ID NO: 207) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker COL6A2 (SEQ ID NO: 207). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement

thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0162] The biomarker FERMT2 (SEQ ID NO: 208) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker FERMT2 (SEQ ID NO: 208) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of FERMT2 (SEQ ID NO: 208) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0163] The biomarker FERMT2 (SEQ ID NO: 208) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker FERMT2 (SEQ ID NO: 208). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0164] The biomarker BNIP3 (SEQ ID NO: 209 or 263) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker BNIP3 (SEQ ID NO: 209 or 263) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of BNIP3 (SEQ ID NO: 209 or 263) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0165] The biomarker BNIP3 (SEQ ID NO: 209 or 263) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H

(SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker BNIP3 (SEQ ID NO: 209 or 263). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0166] The biomarker RAB31 (SEQ ID NO: 210) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker RAB31 (SEQ ID NO: 210) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of RAB31 (SEQ ID NO: 210) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0167] The biomarker RAB31 (SEQ ID NO: 210) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker RAB31 (SEQ ID NO: 210). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0168] The biomarker MPO (SEQ ID NO: 212) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker MPO (SEQ ID NO: 212) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of MPO (SEQ ID NO: 212) in the patient sample may

then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0169] The biomarker MPO (SEQ ID NO: 212) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker MPO (SEQ ID NO: 212). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0170] The biomarker SRPX (SEQ ID NO: 213) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SRPX (SEQ ID NO: 213) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SRPX (SEQ ID NO: 213) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0171] The biomarker SRPX (SEQ ID NO: 213) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on

the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SRPX (SEQ ID NO: 213). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0172] The biomarker ARHGDIB (SEQ ID NO: 214) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker ARHGDIB (SEQ ID NO: 214) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of ARHGDIB (SEQ ID NO: 214) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0173] The biomarker ARHGDIB (SEQ ID NO: 214) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISF (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ARHGDIB (SEQ ID NO: 214). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0174] The biomarker TMEM47 (SEQ ID NO: 215) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker TMEM47 (SEQ ID NO: 215) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of TMEM47 (SEQ ID NO: 215) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0175] The biomarker TMEM47 (SEQ ID NO: 215) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18,

25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker TMEM47 (SEQ ID NO: 215). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0176] The biomarker CSRP2 (SEQ ID NO: 216) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker CSRP2 (SEQ ID NO: 216) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of CSRP2 (SEQ ID NO: 216) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0177] The biomarker CSRP2 (SEQ ID NO: 216) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker CSRP2 (SEQ ID NO: 216). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0178] The biomarker DPYSL3 (SEQ ID NO: 217) may be used to assess a cancer patient's (e.g., a patient having cancer

that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker DPYSL3 (SEQ ID NO: 217) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of DPYSL3 (SEQ ID NO: 217) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0179] The biomarker DPYSL3 (SEQ ID NO: 217) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker DPYSL3 (SEQ ID NO: 217). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0180] The biomarker HTRA1 (SEQ ID NO: 218) may be used to assess a cancer patient's (e.g, a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker HTRA1 (SEQ ID NO: 218) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of HTRA1 (SEQ ID NO: 218) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0181] The biomarker HTRA1 (SEQ ID NO: 218) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ

ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker HTRA1 (SEQ ID NO: 218). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0182] The biomarker SLC39A6 (SEQ ID NO: 219) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SLC39A6 (SEQ ID NO: 219) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SLC39A6 (SEQ ID NO: 219) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0183] The biomarker SLC39A6 (SEQ ID NO: 219) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDI (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SLC39A6 (SEQ ID NO: 219). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0184] The biomarker LAT2 (SEQ ID NO: 220) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker LAT2 (SEQ ID NO: 220) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of LAT2 (SEQ ID NO: 220) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0185] The biomarker LAT2 (SEQ ID NO: 220) may be used as the sole biomarker to predict cancer patient responsiveness

to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LAT2 (SEQ ID NO: 220). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0186] The biomarker LGALS3 (SEQ ID NO: 307) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker LGALS3 (SEQ ID NO: 307) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of LGALS3 (SEQ ID NO: 307) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0187] The biomarker LGALS3 (SEQ ID NO: 307) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LGALS3 (SEQ ID NO: 307). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to

the liposome.

[0188] The biomarker IGFBP4 (SEQ ID NO: 309) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker IGFBP4 (SEQ ID NO: 309) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of IGFBP4 (SEQ ID NO: 309) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0189] The biomarker IGFBP4 (SEQ ID NO: 309) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDI (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker IGFBP4 (SEQ ID NO: 309). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0190] The biomarker SPINT2 (SEQ ID NO: 310) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SPINT2 (SEQ ID NO: 310) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SPINT2 (SEQ ID NO: 310) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0191] The biomarker SPINT2 (SEQ ID NO: 310) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID

NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDI1B (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SPINT2 (SEQ ID NO: 310). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0192] The biomarker CDH1 (SEQ ID NO: 311) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker CDH1 (SEQ ID NO: 311) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of CDH1 (SEQ ID NO: 311) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0193] The biomarker CDH1 (SEQ ID NO: 311) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDI1B (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker CDH1 (SEQ ID NO: 311). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0194] The biomarker RAB25 (SEQ ID NO: 313) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker RAB25 (SEQ ID NO: 313) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of RAB25 (SEQ ID NO: 313) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's

responsiveness to the liposome.

[0195] The biomarker RAB25 (SEQ ID NO: 313) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker RAB25 (SEQ ID NO: 313). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0196] The biomarker TMEM47 (SEQ ID NO: 215) may be used to assess a cancer patient's (e.g. a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker TMEM47 (SEQ ID NO: 215) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of TMEM47 (SEQ ID NO: 215) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0197] The biomarker TMEM47 (SEQ ID NO: 215) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination

with the biomarker TMEM47 (SEQ ID NO: 215). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0198] The biomarker CSRP2 (SEQ ID NO: 216) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker CSRP2 (SEQ ID NO: 216) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of CSRP2 (SEQ ID NO: 216) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0199] The biomarker CSRP2 (SEQ ID NO: 216) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDI (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker CSRP2 (SEQ ID NO: 216). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0200] The biomarker DPYSL3 (SEQ ID NO: 217) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker DPYSL3 (SEQ ID NO: 217) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of DPYSL3 (SEQ ID NO: 217) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0201] The biomarker DPYSL3 (SEQ ID NO: 217) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98),

MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker DPYSL3 (SEQ ID NO: 217). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0202] The biomarker HTRA1 (SEQ ID NO: 218) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker HTRA1 (SEQ ID NO: 218) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of HTRA1 (SEQ ID NO: 218) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0203] The biomarker HTRA1 (SEQ ID NO: 218) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker HTRA1 (SEQ ID NO: 218). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0204] The biomarker SLC39A6 (SEQ ID NO: 219) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the

biomarker SLC39A6 (SEQ ID NO: 219) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SLC39A6 (SEQ ID NO: 219) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0205] The biomarker SLC39A6 (SEQ ID NO: 219) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SLC39A6 (SEQ ID NO: 219). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0206] The biomarker LAT2 (SEQ ID NO: 220) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker LAT2 (SEQ ID NO: 220) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of LAT2 (SEQ ID NO: 220) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0207] The biomarker LAT2 (SEQ ID NO: 220) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1

(SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LAT2 (SEQ ID NO: 220). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0208] The biomarker LGALS3 (SEQ ID NO: 307) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker LGALS3 (SEQ ID NO: 307) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of LGALS3 (SEQ ID NO: 307) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0209] The biomarker LGALS3 (SEQ ID NO: 307) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LGALS3 (SEQ ID NO: 307). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0210] The biomarker DSP (SEQ ID NO: 308) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker DSP (SEQ ID NO: 308) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of DSP (SEQ ID NO: 308) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0211] The biomarker DSP (SEQ ID NO: 308) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN

(SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker DSP (SEQ ID NO: 308). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0212] The biomarker IGFBP4 (SEQ ID NO: 309) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker IGFBP4 (SEQ ID NO: 309) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of IGFBP4 (SEQ ID NO: 309) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0213] The biomarker IGFBP4 (SEQ ID NO: 309) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker IGFBP4 (SEQ ID NO: 309). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0214] The biomarker SPINT2 (SEQ ID NO: 310) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SPINT2 (SEQ ID NO: 310) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SPINT2 (SEQ ID NO: 310) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0215] The biomarker SPINT2 (SEQ ID NO: 310) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SPINT2 (SEQ ID NO: 310). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0216] The biomarker CDH1 (SEQ ID NO: 311) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker CDH1 (SEQ ID NO: 311) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of CDH1 (SEQ ID NO: 311) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0217] The biomarker CDH1 (SEQ ID NO: 311) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213),

RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker CDH1 (SEQ ID NO: 311). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0218] The biomarker PTPRF (SEQ ID NO: 314, 371, or 387) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker PTPRF (SEQ ID NO: 314, 371, or 387) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of PTPRF (SEQ ID NO: 314, 371, or 387) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0219] The biomarker PTPRF (SEQ ID NO: 314, 371, or 387) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker PTPRF (SEQ ID NO: 314, 371, or 387). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0220] The biomarker SOX9 (SEQ ID NO: 121, 315, or 319) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker SOX9 (SEQ ID NO: 121, 315, or 319) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of SOX9 (SEQ ID NO: 121, 315, or 319) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0221] The biomarker SOX9 (SEQ ID NO: 121, 315, or 319) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker SOX9 (SEQ ID NO: 121, 315, or 319). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0222] The biomarker LYZ (SEQ ID NO: 316) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The expression level of the biomarker LYZ (SEQ ID NO: 316) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the expression level of LYZ (SEQ ID NO: 316) in the patient sample may then be compared, e.g., to the expression level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0223] The biomarker LYZ (SEQ ID NO: 316) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker LYZ (SEQ ID NO: 316). In particular, the patient is determined to be responsive to the liposome if the level of the

biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0224] The biomarker IER3 (SEQ ID NO: 127 or 317) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker IER3 (SEQ ID NO: 127 or 317) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of IER3 (SEQ ID NO: 127 or 317) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0225] The biomarker IER3 (SEQ ID NO: 127 or 317) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker IER3 (SEQ ID NO: 127 or 317). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0226] The biomarker PERP (SEQ ID NO: 318) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker PERP (SEQ ID NO: 318) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of PERP (SEQ ID NO: 318) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0227] The biomarker PERP (SEQ ID NO: 318) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CFBF (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H

(SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), ATP1B1 (SEQ ID NO: 320), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker PERP (SEQ ID NO: 318). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0228] The biomarker ATP1B1 (SEQ ID NO: 320) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker ATP1B1 (SEQ ID NO: 320) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of ATP1B1 (SEQ ID NO: 320) in the patient sample may then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0229] The biomarker ATP1B1 (SEQ ID NO: 320) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBFB (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), or IFI27 (SEQ ID NO: 321). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker ATP1B1 (SEQ ID NO: 320). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

[0230] The biomarker IFI27 (SEQ ID NO: 321) may be used to assess a cancer patient's (e.g., a patient having cancer that is resistant to one or more cancer therapies, such as one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome) responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome. The level of the biomarker IFI27 (SEQ ID NO: 321) may be assessed using nucleic acid amplification methods (e.g., PCR, such as qRT-PCR) or a device (e.g., a microarray). As is described above, the level of IFI27 (SEQ ID NO: 321) in the patient sample may

then be compared, e.g., to the level of in a cell (e.g., a cancer cell) or tissue (e.g., a tumor tissue) known to be sensitive or resistant to treatment with the liposome, which can be used as a reference to determine the cancer patient's responsiveness to the liposome.

[0231] The biomarker IFI27 (SEQ ID NO: 321) may be used as the sole biomarker to predict cancer patient responsiveness to treatment with the liposome or in combination with one or more additional biomarkers (e.g., one, two, three, four, five, ten, or all of the biomarkers shown in Tables 1-4), such as COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), RAB25 (SEQ ID NO: 313), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), LAT2 (SEQ ID NO: 220), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), or ATP1B1 (SEQ ID NO: 320). The level of the biomarker(s) may be determined using, e.g., a microarray, PCR, or other techniques described herein, for example, using a nucleic acid probe sequence based on the target sequences shown in Tables 1-4. Optionally, the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ may be detected in a tumor sample from the patient and used to determine the cancer patient's responsiveness to the liposome in combination with the biomarker IFI27 (SEQ ID NO: 321). In particular, the patient is determined to be responsive to the liposome if the level of the biomarker or the complement thereof is substantially similar to the level of the biomarker or the complement thereof in a cell or tissue known to be sensitive to the liposome and/or the level of the biomarker or the complement thereof is substantially dissimilar to the level of the biomarker or the complement thereof in a cell or tissue known to be resistant to the liposome.

Methods of Treatment

[0232] The diagnostic methods of the invention permit the assessment of whether a patient is likely to be responsive to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome, and can thus be used to direct the patient's treatment (e.g., as a first line therapy and/or as a second or third line therapy). A patient to be treated or tested for responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome according to the methods may include, e.g., a patient that has been diagnosed with cancer, a patient that has not received a cancer treatment (e.g., the liposome, an anti-cancer agent other than the liposome, or radiation), a patient that has received a cancer treatment (e.g., an anti-cancer agent other than the liposome or radiation), or a patient during treatment with the liposome. For example, the patient may have a solid tumor or a hematological cancer, such as a cancer type selected from breast cancer (e.g., medullary carcinoma, ER-positive breast cancer, and/or a metastatic form of breast cancer), prostate cancer, ovarian cancer (e.g., ovarian adenocarcinoma or embryonal carcinoma), liver cancer (e.g., hepatocellular carcinoma (HCC) or hepatoma), myeloma (e.g., multiple myeloma), colorectal cancer (e.g., colon cancer and rectal cancer), leukemia (e.g., acute myeloid leukemia, acute lymphoid leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, and chronic leukemia), myelodysplastic syndrome, lymphoma (e.g., diffuse large B-cell lymphoma, cutaneous T-cell lymphoma, peripheral T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, Waldenstrom's macroglobulinemia, and lymphocytic lymphoma), cervical cancer, esophageal cancer, melanoma, glioma (e.g., oligodendroglioma), pancreatic cancer (e.g., adenosquamous carcinoma, signet ring cell carcinoma, hepatoid carcinoma, colloid carcinoma, islet cell carcinoma, and pancreatic neuroendocrine carcinoma), gastrointestinal stromal tumor, sarcoma (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, leiomyosarcoma, Ewing's sarcoma, and rhabdomyosarcoma), bladder cancer, head and neck cancer (e.g., squamous cell carcinoma of the head and neck), lung cancer (e.g., non-small cell lung carcinoma, large cell carcinoma, bronchogenic carcinoma, and papillary adenocarcinoma), metastatic cancer, oral cavity cancer, uterine cancer, testicular cancer (e.g., seminoma and embryonal carcinoma), skin cancer (e.g., squamous cell carcinoma and basal cell carcinoma), thyroid cancer (e.g., papillary carcinoma and medullary carcinoma), brain cancer (e.g., astrocytoma

and craniopharyngioma), stomach cancer, intra-epithelial cancer, bone cancer, biliary tract cancer, eye cancer, larynx cancer, kidney cancer (e.g., renal cell carcinoma and Wilms tumor), gastric cancer, blastoma (e.g., nephroblastoma, medulloblastoma, hemangioblastoma, neuroblastoma, and retinoblastoma), polycythemia vera, chordoma, synovioma, mesothelioma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, cystadenocarcinoma, bile duct carcinoma, choriocarcinoma, epithelial carcinoma, ependymoma, pinealoma, acoustic neuroma, schwannoma, meningioma, pituitary adenoma, nerve sheath tumor, cancer of the small intestine, cancer of the endocrine system, cancer of the penis, cancer of the urethra, cutaneous or intraocular melanoma, a gynecologic tumor, solid tumors of childhood, and neoplasms of the central nervous system. In particular, the cancer of the patient is, e.g., breast cancer (e.g., ERpos breast cancer and/or a metastatic form of breast cancer), acute myelogenous leukemia (AML), acute lympho-blastic leukemia (ALL), chronic lymphocytic leukemia (CLL), myelodysplastic syndrome (MDS), chronic myelogenous leukemia - chronic phase (CMLCP), diffuse large β -cell lymphoma (DLBCL), cutaneous T-cell lymphoma (CTCL), peripheral T-cell lymphoma (PTCL), Hodgkin's lymphoma, hepatocellular carcinoma (HCC), cervical cancer, renal cell carcinoma (RCC), esophageal cancer, melanoma, glioma, pancreatic cancer, gastrointestinal stromal tumors (GIST), sarcoma, non-small cell lung carcinoma (NSCLC), prostate cancer, ovarian cancer, colon cancer, bladder cancer, or squamous cell carcinoma of the head and neck (SCCHN).

[0233] The patient may have a cancer (e.g., breast cancer, such as an ERpos breast cancer and/or a metastatic form of breast cancer) that is resistant to one or more cancer therapies other than an sPLA₂ hydrolysable, cisplatin-containing liposome (e.g., docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, and rituximab), surgery, or radiation. The patient may also have experienced a recurrence following a treatment with a cancer therapy other than the liposome, surgery, or radiation.

[0234] A patient found to be responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome according to the methods of the invention may be preferentially selected for treatment with the liposome. For example, a patient can be identified as responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome by determining the level of one or more biomarkers (e.g., one or more of the biomarkers shown in Tables 1-4, such as COL5A2 (SEQ ID NO: 73 or 211)) in a biological sample (e.g., a tumor sample) obtained from the patient, and subsequently administered the liposome. Optionally, the level of one or more of the biomarkers (e.g., one or more of the biomarkers shown in Tables 1-4, such as COL5A2 (SEQ ID NO: 73 or 211)) can be used in combination with the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ to identify patients responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome. Alternatively, a patient can be identified as less likely to be responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome by determining the level of one or more biomarkers (e.g., one or more of the biomarkers shown in Tables 1-4, such as COL5A2 (SEQ ID NO: 73 or 211)) in a biological sample obtained from the patient. Likewise, a patient may be less likely to be responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome if the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ is not detected in a tumor sample from the patient.

[0235] If the patient exhibits levels of one or more biomarkers indicative of non-responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome, the patient may be treated with or offered a treatment with an agent other than the liposome. Additionally, the patient may be treated with or offered a treatment with an agent other than the liposome if the level of PLA2G2A (SEQ ID NO: 380) or sPLA₂ is not detected in a tumor sample from the patient. In particular, the patient may be treated with, e.g., radiation and/or administration of a therapeutic agent, such as docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA),

cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, or rituximab.

Administration of an sPLA₂ hydrolysable, cisplatin-containing liposome

[0236] Once a patient has been determined to be responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome, according to the methods described herein, the liposome may be administered to the patient, for example, parenterally or enterally. Enteral routes of the liposome administration include oral, buccal, sublabial, sublingual, or by inhalation. Parenteral routes of the liposome administration include intravenous, transdermal, intradermal, intramuscular, intra-arterial, intracranial, subcutaneous, intraorbital, intraventricular, intraspinal, intraperitoneal, or intranasal. The preferred route for administration of the sPLA₂ hydrolysable, cisplatin-containing liposome may be intravenous, such as intravenous infusion or as a bolus injection. For example, the sPLA₂ hydrolysable, cisplatin-containing liposome is administered by intravenous infusion over a time period of, e.g., 15 minutes, 30 minutes, 45 minutes, 60 minutes, 75 minutes, 90 minutes, 105 minutes, 120 minutes, 135 minutes, 150 minutes, 180 minutes, or more. The sPLA₂ hydrolysable, cisplatin-containing liposome may also be administered as, e.g., a bolus injection.

[0237] An sPLA₂ hydrolysable, cisplatin-containing liposome can be administered at, e.g., a dose of about 60 mg/m² to 120 mg/m² of cisplatin, such as about 75 mg/m² of cisplatin. For example, an sPLA₂ hydrolysable, cisplatin-containing liposome may be administered at a dose of about 60 mg/m² of cisplatin, 65 mg/m² of cisplatin, 70 mg/m² of cisplatin, 75 mg/m² of cisplatin, 80 mg/m² of cisplatin, 85 mg/m² of cisplatin, 90 mg/m² of cisplatin, 95 mg/m² of cisplatin, 100 mg/m² of cisplatin, 105 mg/m² of cisplatin, 110 mg/m² of cisplatin, 115 mg/m² of cisplatin, or 120 mg/m² of cisplatin. An sPLA₂ hydrolysable, cisplatin-containing liposome may be administered at a frequency of, e.g., at least once hourly, once daily, twice daily, once weekly, once every two weeks, once every three weeks, once every four weeks, once monthly, once every two months, once every three months, once every six months, or once every year.

[0238] In particular, an sPLA₂ hydrolysable, cisplatin-containing liposome may be administered at a treatment regimen of, e.g., about 75 mg/m² of cisplatin on day 1 and day 8 of a 3 week cycle (1 cycle) for 1 to 3 cycles or more. Additionally, the sPLA₂ hydrolysable, cisplatin-containing liposome may be administered at a treatment regimen of, e.g., about 60 mg/m² of cisplatin on day 1 and day 8 of a 3 week cycle (1 cycle) for 1 to 3 cycles or more. The sPLA₂ hydrolysable, cisplatin-containing liposome may also be administered at a treatment regimen of, e.g., about 90 mg/m² of cisplatin on day 1 and day 8 of a 3 week cycle (1 cycle) for 1 to 3 cycles or more. The treatment regimen may be repeated one to five times, one to ten times, one to fifteen times, one to twenty times, or more. The administration of the sPLA₂ hydrolysable, cisplatin-containing liposome can be repeated at such a frequency for a certain period of time, followed by a period without treatment. Such repeated administrations can occur over a course of therapy lasting a specified length of time (e.g., at least 1 week, 2 weeks, 3 weeks, 1 month, 2 months, 3 months, 6 months, 8 months, 10 months, 12 months, 18 months, 24 months, 36 months, 48 months, or 60 months).

[0239] An sPLA₂ hydrolysable, cisplatin-containing liposome can be administered in a pharmaceutical composition that includes one or more pharmaceutically acceptable carriers, excipients, or diluents. Examples of suitable carriers, excipients, or diluents of the liposome include, but are not limited to, sodium chloride solution, saline, sterile water, polyalkylene glycols, oils of vegetable origin, hydrogenated naphthalenes, suitable buffer, 1,3-butanediol, and/or Ringer's solution. In particular, the sPLA₂ hydrolysable, cisplatin-containing liposome is administered in a pharmaceutical composition that includes about 0.9% sodium chloride. Other exemplary carriers, excipients, or diluents are described in the Handbook of Pharmaceutical Excipients, 6th Edition, Rowe et al., Eds., Pharmaceutical Press (2009).

sPLA₂ Hydrolysable Liposomes

[0240] An sPLA₂ hydrolysable, cisplatin-containing liposome for use in the methods may include anionic lipids, neutral phospholipids, and polymer-conjugated lipids. For example, the liposome includes about 40% to about 75% (mol/mol) of a neutral phospholipid, about 20% to about 45% (mol/mol) of an anionic lipid, and about 3% to about 6% (mol/mol) of a

polymer-conjugated lipid. In particular, the liposome includes distearoyl phosphatidyl glycerol (DSPG) in an amount of about 25% (mol/mol), distearoyl phosphatidyl choline (DSPC) in an amount of about 70% (mol/mol), and [poly(ethylene glycol)]-distearoyl phosphatidyl ethanolamine (DSPE-PEG) in an amount of about 5% (mol/mol).

[0241] The sPLA₂ hydrolysable liposome may include, e.g., about 20% to about 45% (mol/mol) of an anionic lipid, such as about 20% to about 40% (mol/mol) anionic lipid, about 20% to about 30% (mol/mol) anionic lipid, about 25% to about 40% (mol/mol) anionic lipid, about 25% to about 35% (mol/mol) anionic lipid, or about 25% to about 30% (mol/mol) anionic lipid. The anionic lipid may include one or more phospholipids, such as phosphatidyl inositol (PI), phosphatidyl serine (PS), bisphosphatidyl glycerol (DPG), phosphatidic acid (PA), phosphatidyl alcohol (PEOH), or phosphatidyl glycerol (PG). In particular, the anionic phospholipid is PG including stearoyl chains, such as DSPG in an amount of about 25% (mol/mol) of the liposomal composition.

[0242] The sPLA₂ hydrolysable, cisplatin-containing liposome may also include one or more neutral phospholipids, such as phosphatidyl choline (PC) or phosphatidylethanolamine (PE), in an amount of about 40% to about 75% (mol/mol) of the liposomal composition. Preferably, the neutral phospholipid is PC, such as PC in an amount of about 50% to about 70% (mol/mol). In particular, the neutral lipid also includes stearoyl chains, such as the liposome includes DSPC in an amount of about 70% (mol/mol).

[0243] Additionally, the liposome may include one or more polymer-conjugated lipid, such as PEG [poly(ethylene glycol)], PAcM [poly(N-acryloylmorpholine)], PVP [poly(vinylpyrrolidone)], PLA [poly(lactide)], PG [poly(glycolide)], POZO [poly(2-methyl-2-oxazoline)], PVA [poly(vinyl alcohol)], HPMC (hydroxypropylmethylcellulose), PEO [poly(ethylene oxide)], chitosan [poly(D-glucosamine)], PAA [poly(aminoacid)], polyHEMA [Poly(2-hydroxyethylmethacrylate)], and co-polymers thereof. Preferably, the liposome includes a polymer-conjugated lipid in an amount of at least 2%, such as an amount of polymer-conjugated lipid of at least 5% and no more than 15% (mol/mol) or at least 3% and no more than 6% (mol/mol). For example, the liposome includes DSPE-PEG in an amount of about 5% (mol/mol).

Kits

[0244] Kits of the invention can be used for determining the responsiveness of a cancer patient (e.g., a patient having a solid tumor cancer, such as breast cancer, or a hematological cancer, such as lymphoma (e.g., cutaneous T-cell lymphoma (CTCL)) to an sPLA₂ hydrolysable, cisplatin-containing liposome. Kits of the invention can include reagents and/or materials for, e.g., collecting and/or purifying nucleic acids from biological samples (such as those obtained from a patient to be treated with a target drug(s) of the invention), reagents for amplifying such nucleic acids to produce an amplified sample, and/or at least one device of the invention. Reagents for amplifying nucleic acids may include, e.g., PCR reagents, including but not limited to DNA polymerase, RNA polymerase, PCR buffer, magnesium chloride solutions, nucleic acid primers (e.g., primers designed to target particular biomarkers of responsiveness to a target drug(s) of interest), and/or any other PCR reagents as are well known in the art. In particular, kits useful in the method may include one or more of the following: a kit for RNA extraction from tumors (e.g., Trizol for mRNA, mirVana miRNA isolation kit from Ambion Inc), a kit for RNA labeling (e.g., MessageAmp from Ambion Inc., FlashTag from Genisphere Inc), a microarray for measuring biomarker levels (e.g., HG-U133A, HG-U133_Plus2 or miRNA-1.0 from Affymetrix Inc), a microarray hybridization station and scanner (e.g., GeneChip System 3000Dx from Affymetrix Inc), and/or software for analyzing the levels of biomarkers as described in herein (e.g., implemented in R from R-Project or S-Plus from Insightful Corp.).

[0245] For example, a kit of the invention can include one or more probes capable of detecting one or more biomarkers of Tables 1-4 (e.g., the kit may include probes for one or more of the biomarkers of Tables 1-4) and, optionally, one or more probes for detecting PLA2G2A (SEQ ID NO: 380). Such probes can, for example, include nucleic acids capable of specifically hybridizing to the biomarker(s) based on nucleic acid sequence complementarity. In particular, a probe has at least 85% sequence identity (e.g., 85%, 90%, 95%, 97%, 98%, 99%, or 100% sequence identity) to a nucleic acid sequence that is complementary or identical to at least 5 (e.g., at least 15 or more) consecutive nucleotides of one or more biomarkers. The probes can be attached to a solid surface, such as a microarray. The kit may include NanoString capture probes, NanoString reporter probes, and/or one or more nCounter cartridges. The kit may include reagents for next generation sequencing, including but not limited to poly(T) oligonucleotides, dye terminators, sequencing adapters, adapter ligation reagents, reverse transcriptase, primers (e.g., random primers), DNA-cleaving enzymes, polymerases, and/or any combination thereof. The kit may also be one that includes a protein array and/or reagents for detection of the polypeptide product(s) of one or more biomarkers of Tables 1-4 (e.g., antibodies or antigen-binding fragments that specifically bind to

the one or more biomarkers), and optionally, an anti-sPLA₂ antibody for detecting the sPLA₂ protein.

[0246] The following examples are intended to illustrate, rather than limit, the invention.

EXAMPLES

Example 1. Identification of biomarkers of sensitivity and resistance to cisplatin using Affymetrix HG-U133A arrays.

[0247] DNA chip measurements of the 60 cancer cell lines of the NCI60 data set were performed using Affymetrix HG-U133A arrays and logit normalized. For each array, the logit transformation was performed followed by a Z-transformation to mean zero and SD 1, and correlated to growth inhibition ($\log(\text{GI50})$). Growth inhibition data of cisplatin against the same cell lines were downloaded from the National Cancer Institute. Each gene's expression in each cell line was correlated to the growth of those cell lines ($\log(\text{GI50})$) in the presence of cisplatin. The Pearson correlation coefficient was then determined to identify genes positively and negatively correlated to sensitivity to cisplatin. Tables 1 and 2 show the top positively correlated genes (the biomarkers of sensitivity) and negatively correlated genes (the biomarkers of resistance) using the Affymetrix HG-U133A arrays. These biomarkers of sensitivity and resistance shown in Tables 1 and 2, respectively, may be used individually or in combination to identify patients that are responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome.

Example 2. Identification of biomarkers of sensitivity and resistance to an sPLA₂ hydrolysable, cisplatin-containing liposome using Affymetrix HG-U133A arrays.

[0248] DNA chip measurements of the 60 cancer cell lines of the NCI60 data set were also performed using HG-U133_Plus_2 arrays and logit normalized. For each array, the logit transformation was performed followed by a Z-transformation to mean zero and SD 1, and correlated to growth inhibition ($\log(\text{GI50})$). Growth inhibition data of a sPLA₂ hydrolysable, cisplatin-containing liposome against the same cell lines were downloaded from the National Cancer Institute. Each gene's expression in each cell line was correlated to the growth of those cell lines ($\log(\text{GI50})$) in the presence of the liposome. The covariance (Pearson correlation coefficient multiplied by standard deviation) was then determined to identify genes positively and negatively correlated to sensitivity to the liposome. Tables 3 and 4 show the top positively correlated genes (the biomarkers of sensitivity) and negatively correlated genes (the biomarkers of resistance) using the Affymetrix HG-U133A arrays, respectively. These biomarkers of sensitivity and resistance shown in Tables 3 and 4, respectively, may be used individually or in combination to identify patients that are responsive to an sPLA₂ hydrolysable, cisplatin-containing liposome.

Example 3. Predicting responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome in various cancer patient populations.

[0249] An mRNA-based predictor of responsiveness to an sPLA₂ hydrolysable, cisplatin-containing liposome developed according to the methods of the invention was applied to 3,522 patients having a variety of cancers. Each patient had a pre-treatment measurement of gene expression with an Affymetrix array. The predicted liposome sensitivity of each patient was calculated as the difference between the mean of the levels of the biomarkers of sensitivity (Table 1) and the mean of the levels of the biomarkers of resistance (Table 2) for the patient. When the patients were grouped by cancer types, and cancer types predicted to be more responsive to the liposome were identified (Figure 1). Of 27 different cancer types, patients with hematological cancer types were predicted to be responsive to the liposome treatment than patients with solid tumor cancers.

[0250] The median of the boxplots shown in Figure 1 is a cutoff that may be used to separate patients predicted to be responsive to treatment with the sPLA₂ hydrolysable, cisplatin-containing liposome from patients predicted to be non-responsive to the liposome treatment for a given cancer type. Values above the median indicate patients predicted to be responsive to the liposome, while values below the median indicate patients predicted to be non-responsive to the

liposome. For a test sample from an individual patient, it is useful to compare the test sample to the reference population for the same cancer type. If the test sample is above the median for the reference population of the same cancer type, then the patient is predicted to be responsive to the liposome treatment. If the test sample is below the median for the reference population of the same cancer type, then the patient is predicted to be non-responsive to the liposome treatment. This method for predicting patient responsiveness can also be used when the reference cancer population consists of only two patients: a patient responsive to the liposome treatment and a patient non-responsive to the liposome treatment.

Example 4. Determining the presence of secreted phospholipase A₂ (sPLA₂).

[0251] The presence of secreted sPLA₂ in the tumor tissue of a patient having cancer can be used in combination with the expression of the biomarkers listed in Tables 1-4 to determine the responsiveness of a patient to an sPLA₂ hydrolysable, cisplatin-containing liposome. Expression of the enzyme sPLA₂ is required for degradation of the liposomes that deliver cisplatin to the tumor tissue of the patient and can be measured using standard immunocytochemistry techniques with a monoclonal antibody against sPLA₂-IIA, e.g. Clone SCACC353 from Cayman Chemical. Any staining in this assay indicates the presence of sPLA₂ and suggests susceptibility to the liposome. Alternatively, the expression of sPLA₂-IIA can be detected using a microarray including one or more probes for PLA2G2A (SEQ ID NO: 380) or qRT-PCR to determine a level of PLA2G2A (SEQ ID NO: 380) in a tumor sample from the patient. While there is a negative covariance between PLA2G2A (SEQ ID NO: 380) and the liposome response in cancer cell lines *in vitro*, there is a positive correlation between detection of a level of PLA2G2A in a tumor tissue sample (Mirtti et al. APMIS 117: 151-161, 2009).

Example 5. Predicting responsiveness of breast cancer patients to the liposome.

[0252] The diagnostic methods of the present invention can be used to predict the responsiveness of a breast cancer patient to treatment with an sPLA₂ hydrolysable, cisplatin-containing liposome. In particular, the breast cancer patient may be one that has not previously received any cancer treatment or one that has received a cancer treatment other than the liposome. Moreover, the patient may be one diagnosed with breast cancer and/or one with recurrence of breast cancer. A biological sample (e.g., a breast cancer tissue sample) may be obtained from the patient through methods well known in the art. The sample may be frozen and/or prepared, e.g., by formalin fixation and paraffin embedding. In particular, mRNA can be isolated from the sample and a gene expression profile can be determined, e.g., using a microarray platform, such as the Affymetrix HG-U133A or HG-U133_Plus_2 array, for one or more of the biomarkers shown in Tables 1-4. One or more of the biomarkers shown in Tables 1-4 can also be measured, e.g., by sequencing or PCR-based techniques, such as those described herein.

[0253] For example, the level of one or more biomarkers of sensitivity to the liposome can be determined in the sample from the patient, such as one or more of COL5A2 (SEQ ID NO 73 or 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 or 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 or 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25, or 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38, or 272), C1QR1 (SEQ ID NO: 13), FNBP1 (SEQ ID NO: 14 or 28), CBF3 (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 or 54), CAP350 (SEQ ID NO: 20 or 61), CALD1 (SEQ ID NO: 206), COL6A2 (SEQ ID NO: 207), FERMT2 (SEQ ID NO: 208), BNIP3 (SEQ ID NO: 209 or 263), RAB31 (SEQ ID NO: 210), MPO (SEQ ID NO: 212), SRPX (SEQ ID NO: 213), ARHGDIB (SEQ ID NO: 214), TMEM47 (SEQ ID NO: 215), CSRP2 (SEQ ID NO: 216), DPYSL3 (SEQ ID NO: 217), HTRA1 (SEQ ID NO: 218), SLC39A6 (SEQ ID NO: 219), and LAT2 (SEQ ID NO: 220). In particular, the biomarker is COL5A2 (SEQ ID NO 73 or 211). The level of one or more biomarkers of resistance to the liposome can also be determined in the sample from the patient, such as one or more of SFN (SEQ ID NO: 96 or 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 or 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 or 400), KRT18 (SEQ ID NO: 108 or 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 or 312), LRP5 (SEQ ID NO: 112), LGALS3 (SEQ ID NO: 307), DSP (SEQ ID NO: 308), IGFBP4 (SEQ ID NO: 309), SPINT2 (SEQ ID NO: 310), CDH1 (SEQ ID NO: 311), RAB25 (SEQ ID NO: 313), PTPRF (SEQ ID NO: 314, 371, or 387), SOX9 (SEQ ID NO: 121, 315, or 319), LYZ (SEQ ID NO: 316), IER3 (SEQ ID NO: 127 or 317), PERP (SEQ ID NO: 318), ATP1B1 (SEQ ID NO: 320), and IFI27 (SEQ ID NO: 321). In particular, the biomarker is SFN (SEQ ID NO: 96 or 324). The breast cancer patient may be responsive to the liposome if the level of one or more of the biomarkers of sensitivity is substantially similar to the level of the biomarkers of sensitivity in a cell or tissue known to be sensitive to the liposome. The breast cancer patient may also be responsive to the liposome if the level of one or more of the biomarkers of resistance is substantially dissimilar to the level

of the biomarkers of resistance in a cell or tissue known to be resistant to the liposome.

[0254] Additionally, the presence of secreted phospholipase A2 (sPLA₂-IIA) in the tumor tissue of a patient having breast cancer can be used in combination with one or more of the biomarkers of Tables 1-4 to determine the responsiveness of the breast cancer patient to the sPLA₂ hydrolysable, cisplatin-containing liposome. The presence of sPLA₂ can be measured using standard immunocytochemistry techniques with a monoclonal antibody against sPLA₂-IIA, such as Clone SCACC353 from Cayman Chemical. Any staining of the tumor tissue using an anti-sPLA₂ antibody indicates the presence of sPLA₂ and suggests that the breast cancer patient may be responsive to treatment with the liposome. Alternatively, the expression of sPLA₂-IIA can be detected using a microarray including one or more probes that are capable of specifically hybridizing with PLA2G2A (SEQ ID NO: 380) or using qRT-PCR to determine the level of PLA2G2A (SEQ ID NO: 380).

[0255] If the breast cancer patient is predicted to be responsive to treatment with the sPLA₂ hydrolysable, cisplatin-containing liposome, then the patient can be administered the liposome, such as intravenous administration of the liposome at about 60 to about 120 mg/m² on day 1 and day 8 of a three week regimen (e.g., 75 mg/m² of the liposome). Conversely, if the patient is predicted to be non-responsive to treatment with the liposome, then the patient can be treated with one or more therapies other than the liposome, such as surgery, radiation, or a non-liposome therapeutic agent (e.g., docetaxel, cabazitaxel, mitoxantrone, estramustine, prednisone, carboplatin, bevacizumab, paclitaxel, gemcitabine, doxorubicin, topotecan, etoposide, tamoxifen, letrozole, sorafenib, fluorouracil, capecitabine, oxaliplatin, interferon-alpha, 5-fluorouracil (5-FU), a histone deacetylase (HDAC) inhibitor, ipilimumab, bortezomib, carfilzomib, thalidomide, lenalidomide, pomalidomide, dexamethasone, cyclophosphamide, vincristine, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustine, depsipeptide, erlotinib, busulfan, epirubicin, arsenic trioxide, bendamustine, fulvestrant, teniposide, adriamycin, decitabine, estramustine, azaguanine, aclarubicin, mitomycin, paclitaxel, taxotere, APO010, ara-c, methylprednisolone, methotrexate, methyl-gag, belinostat, idarubicin, IL4-PR38, valproic acid, all-trans retinoic acid (ATRA), cytoxan, suberoylanilide hydroxamic acid, leukeran, fludarabine, vinblastine, dacarbazine, hydroxyurea, tegafur, daunorubicin, mechlorethamine, streptozocin, carmustine, mercaptopurine, dactinomycin, tretinoin, ifosfamide, floxuridine, thioguanine, PSC 833, herceptin, celecoxib, iressa, anastrozole, or rituximab).

Example 6. Clinical validation of identified biomarkers of resistance and sensitivity.

[0256] The biomarkers of sensitivity and resistance identified in Example 1 and shown in Tables 1 and 2 have been validated in five clinical trials. Table 5 below summarizes the results of these studies. In each trial, the overall response rate (ORR) or overall survival (OS) is substantially higher in the top 10% of patients identified as responsive to treatment (e.g., the sPLA₂ hydrolysable, cisplatin-containing liposome) using the biomarkers of sensitivity and resistance than in the bottom 10% of patients identified as responsive to treatment using the biomarkers of sensitivity and resistance. Trials 1-3 are retrospective analyses of published data including Affymetrix assays of pre-treatment biopsies from patients treated with platinum, cisplatin, or cisplatin and vinorelbine, respectively, having ovarian cancer, breast cancer, or non-small cell lung carcinoma (NSCLC), respectively. In Trial 3, the biomarkers of sensitivity and resistance were used to evaluate responsiveness to cisplatin without an additional predictor of responsiveness to vinorelbine. In Trial 4, the biomarkers of sensitivity and resistance were used to analyze archived tumor samples of NSCLC patients. Trial 5 is a prospective trial of tumor samples from breast cancer patients screened with the biomarkers of sensitivity and resistance of Tables 1 and 2, respectively, prior to treatment with the sPLA₂ hydrolysable, cisplatin-containing liposome. Only the breast cancer patients in the top 67% of patients predicted to be responsive to liposome treatment and positive for sPLA₂ presence in a tumor sample from the patient, as determined using an immunohistochemistry assay, were treated with the liposome (trial ongoing until N=15). All studies were conducted with the Affymetrix HG-U133A or HG-U133Plus2 array.

Table 5. Summary of clinical trials conducted using biomarkers of sensitivity and resistance (Tables 1 and 2, respectively) to treatment, such as the sPLA₂ hydrolysable, cisplatin-containing liposome. The overall response rate (ORR) and overall survival (OS) are shown for patients predicted to be in the top 10% or bottom 10% of responsiveness to the treatment for each respective trial.

Trial No.	1	2	3	4	5
Cancer	Ovarian	Breast	NSCLC	NSCLC	Breast, etc.
Drug	Platinum	Cisplatin	Cisplatin and Vinorelbine	Cisplatin and Vinorelbine	sPLA ₂ hydrolysable, cisplatin- containing liposome
Level of Evidence	Retrospective	Retrospective	Retrospective	Prospective archival	Prospective screening

Setting	adjuvant	neoadj	adjuvant	adjuvant	metastat
Assay	Fresh	Fresh	Fresh	FFPE	FFPE
N	28	24	71	95	11 (so far)
Top 10%	ORR 100%	ORR 100%	3Y OS 100%	3Y OS 100%	ORR 100%
Bottom 10%	ORR 67%	ORR 50%	3Y OS 71%	3Y OS 55%	0%

[0257] As shown in Table 5, the ORR and OS in the top 10% of patients predicted to be responsive to treatment was 100% and the bottom 10% of patients predicted to be responsive to treatment was substantially lower, ranging from 50% to 71%, in all trials including patients with ovarian cancer, breast cancer, and NSCLC. In trial number 5, also with a cutoff of 33% (top 67%) there is a 67% probability of response and a median of 18 weeks to progression of disease, whereas patients below this cutoff have a 0% probability of response and a median of 3 weeks to progression. These five clinical trials indicate that the biomarkers of sensitivity and resistance of Tables 1-4 can be used singly or in combination with detection of sPLA₂ in patient tumor samples to identify cancer patients that are likely to be responsive to treatment with, e.g., a sPLA₂ hydrolysable, cisplatin-containing liposome.

SEQUENCE LISTING

[0258]

<110> Oncology Venture ApS

<120> METHODS FOR PREDICTING DRUG RESPONSIVENESS IN CANCER PATIENTS

<130> 51167-004001

<160> 405

<170> PatentIn version 3.5

<210> 1

<211> 25

<212> DNA

<213> Homo sapiens

<400> 1

caggcctctc agatacaagg ggaac 25

<210> 2

<211> 25

<212> DNA

<213> Homo sapiens

<400> 2

atagctgcct taaagtcagt aactt 25

<210> 3

<211> 25

<212> DNA

<213> Homo sapiens

<400> 3

caccatgctg gctatccggg tgтта 25

<210> 4

<211> 25

<212> DNA

<213> Homo sapiens

<400> 4

ctccactctg actggcacag tcttt 25

<210> 5
<211> 25
<212> DNA
<213> Homo sapiens

<400> 5
gattcctgaa ctcaaggtac cagca 25

<210> 6
<211> 25
<212> DNA
<213> Homo sapiens

<400> 6
gatcatcgac gccatcaggc aggag 25

<210> 7
<211> 25
<212> DNA
<213> Homo sapiens

<400> 7
gtgtggttga acaggatgca atctt 25

<210> 8
<211> 25
<212> DNA
<213> Homo sapiens

<400> 8
gcttccaag atgccatggc tggac 25

<210> 9
<211> 25
<212> DNA
<213> Homo sapiens

<400> 9
gcaggacttc cttataaag caaaa 25

<210> 10
<211> 25
<212> DNA
<213> Homo sapiens

<400> 10
gattataacc gtgttgaact ctctg 25

<210> 11
<211> 25
<212> DNA
<213> Homo sapiens

<400> 11
gttttcggtc ctattttaat gctct 25

<210> 12
<211> 25
<212> DNA
<213> Homo sapiens

<400> 12

aaagaccaac aaatctcaag cccta 25

<210> 13

<211> 25

<212> DNA

<213> Homo sapiens

<400> 13

ggtctgttct ttagataat gccct 25

<210> 14

<211> 25

<212> DNA

<213> Homo sapiens

<400> 14

tgctggccac ggatttgac gacga 25

<210> 15

<211> 25

<212> DNA

<213> Homo sapiens

<400> 15

ggtgtgtgac agctcacatg ttac 25

<210> 16

<211> 25

<212> DNA

<213> Homo sapiens

<400> 16

ggttgccctc attgtgctat ttgcc 25

<210> 17

<211> 25

<212> DNA

<213> Homo sapiens

<400> 17

ataagcattg attcctgcat ttctg 25

<210> 18

<211> 25

<212> DNA

<213> Homo sapiens

<400> 18

gcatttagtc caatgtcttt ttaag 25

<210> 19

<211> 25

<212> DNA

<213> Homo sapiens

<400> 19

atcatgctga ggcgcttgc aaatc 25

<210> 20

<211> 25

<212> DNA

<213> Homo sapiens

<400> 20
atgactggta tgatagctct tgaca 25

<210> 21
<211> 25
<212> DNA
<213> Homo sapiens

<400> 21
caatccaagc ataactcagt gacgc 25

<210> 22
<211> 25
<212> DNA
<213> Homo sapiens

<400> 22
gaatgtgtag ctcaaatgca aacca 25

<210> 23
<211> 25
<212> DNA
<213> Homo sapiens

<400> 23
ttccctcctt atagtcaagg accgt 25

<210> 24
<211> 25
<212> DNA
<213> Homo sapiens

<400> 24
tgacctcggc cacaaaagca gtttt 25

<210> 25
<211> 25
<212> DNA
<213> Homo sapiens

<400> 25
gaacagtttg tacagacgta tgctt 25

<210> 26
<211> 25
<212> DNA
<213> Homo sapiens

<400> 26
tacaacacta tacatacaca ccacc 25

<210> 27
<211> 25
<212> DNA
<213> Homo sapiens

<400> 27
ttatgccagc ttatattgtg agaac 25

<210> 28
<211> 25
<212> DNA

<213> Homo sapiens

<400> 28
gagttgcctg ttgtctctg gagat 25

<210> 29
<211> 25
<212> DNA
<213> Homo sapiens

<400> 29
gctgcagtgt agatggctct tgttt 25

<210> 30
<211> 25
<212> DNA
<213> Homo sapiens

<400> 30
acgttgctac cggagcactg aagat 25

<210> 31
<211> 25
<212> DNA
<213> Homo sapiens

<400> 31
atagcagcac acatttcac gtttc 25

<210> 32
<211> 25
<212> DNA
<213> Homo sapiens

<400> 32
aagaccggca gatgtggtg ctgga 25

<210> 33
<211> 25
<212> DNA
<213> Homo sapiens

<400> 33
atttggtca agtccattg gctgt 25

<210> 34
<211> 25
<212> DNA
<213> Homo sapiens

<400> 34
gcatgaagtt gcccttaacc actaa 25

<210> 35
<211> 25
<212> DNA
<213> Homo sapiens

<400> 35
taaccatgct tacacactaa actat 25

<210> 36
<211> 25

<212> DNA
<213> Homo sapiens

<400> 36
tgcccggtcg gagtaaacac agaat 25

<210> 37
<211> 25
<212> DNA
<213> Homo sapiens

<400> 37
gatgtcaact gggatcaatgg gggcc 25

<210> 38
<211> 25
<212> DNA
<213> Homo sapiens

<400> 38
gttggtgat attggagtgc tcatt 25

<210> 39
<211> 25
<212> DNA
<213> Homo sapiens

<400> 39
cctttgtac ttcactcaga tacta 25

<210> 40
<211> 25
<212> DNA
<213> Homo sapiens

<400> 40
taagcatcct tagggttctg cctct 25

<210> 41
<211> 25
<212> DNA
<213> Homo sapiens

<400> 41
ctcatctccc tcaaggatgg ctacg 25

<210> 42
<211> 25
<212> DNA
<213> Homo sapiens

<400> 42
tgtgaatcat cctgccttc aaatt 25

<210> 43
<211> 25
<212> DNA
<213> Homo sapiens

<400> 43
tacaaccac attactctg tcaact 25

<210> 44

<211> 25
<212> DNA
<213> Homo sapiens

<400> 44
gtcccgatc caggacctgg tagca 25

<210> 45
<211> 25
<212> DNA
<213> Homo sapiens

<400> 45
agtgaagagg tcgtctctc catct 25

<210> 46
<211> 25
<212> DNA
<213> Homo sapiens

<400> 46
gctgaaggca cctactcagt atctt 25

<210> 47
<211> 25
<212> DNA
<213> Homo sapiens

<400> 47
taagcattcc gtccatctaa gctca 25

<210> 48
<211> 25
<212> DNA
<213> Homo sapiens

<400> 48
tgatggagca taacgggtcc cagcc 25

<210> 49
<211> 25
<212> DNA
<213> Homo sapiens

<400> 49
atgatttctt agggctgtg tactt 25

<210> 50
<211> 25
<212> DNA
<213> Homo sapiens

<400> 50
gttcatttc ttcattcac aagat 25

<210> 51
<211> 25
<212> DNA
<213> Homo sapiens

<400> 51
gaggccaaga aattccatgt tgttt 25

<210> 52
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 52
 aaagctgtga atctgttccc tgctg 25

 <210> 53
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 53
 atgagctccc agattcgtca gaatt 25

 <210> 54
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 54
 tcccatcag gaagtctctg cagaa 25

 <210> 55
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 55
 tgagtcctg gaacgccaga tgcgt 25

 <210> 56
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 56
 ttagagccag tctcacact cggaa 25

 <210> 57
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 57
 aaaacttccc cggatatgatg attgt 25

 <210> 58
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 58
 tcagtgggca cagttcttca gctac 25

 <210> 59
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 59

actagctcat tattccatc ttgg 25

<210> 60
<211> 25
<212> DNA
<213> Homo sapiens

<400> 60
aattatgagt ttctatctgt gtcca 25

<210> 61
<211> 25
<212> DNA
<213> Homo sapiens

<400> 61
gggaagtcca catagcgta ttaat 25

<210> 62
<211> 25
<212> DNA
<213> Homo sapiens

<400> 62
ttcagctacc ctgactttct cagga 25

<210> 63
<211> 25
<212> DNA
<213> Homo sapiens

<400> 63
ggcatgatgt ccggtgattt ctgta 25

<210> 64
<211> 25
<212> DNA
<213> Homo sapiens

<400> 64
actcttgagg gttgattatg ctgca 25

<210> 65
<211> 25
<212> DNA
<213> Homo sapiens

<400> 65
ggacctcatc tctgagctga aacgg 25

<210> 66
<211> 25
<212> DNA
<213> Homo sapiens

<400> 66
gttgggtgtt aaagatctga agtgt 25

<210> 67
<211> 25
<212> DNA
<213> Homo sapiens

<400> 67
gaagcaccat aactttgttt agccc 25

<210> 68
<211> 25
<212> DNA
<213> Homo sapiens

<400> 68
acagcatgag gcggccgggg agctg 25

<210> 69
<211> 25
<212> DNA
<213> Homo sapiens

<400> 69
gctatggta tattagcacc aaact 25

<210> 70
<211> 25
<212> DNA
<213> Homo sapiens

<400> 70
ggccgacaac agctgcatct atgtc 25

<210> 71
<211> 25
<212> DNA
<213> Homo sapiens

<400> 71
tgaaaagggt ttctattct ctctg 25

<210> 72
<211> 25
<212> DNA
<213> Homo sapiens

<400> 72
agtatcagtc ggtgcaacag ttggc 25

<210> 73
<211> 25
<212> DNA
<213> Homo sapiens

<400> 73
tgaagtgat cctgagactc ttgaa 25

<210> 74
<211> 25
<212> DNA
<213> Homo sapiens

<400> 74
tactcagagg tgtgaccctc gccag 25

<210> 75
<211> 25
<212> DNA

<213> Homo sapiens

<400> 75
gtcgtactat ctactgagc cacag 25

<210> 76
<211> 25
<212> DNA
<213> Homo sapiens

<400> 76
aaggcgtaac gagtcatct ttctt 25

<210> 77
<211> 25
<212> DNA
<213> Homo sapiens

<400> 77
ccttgatacc agctctctgt ggaaa 25

<210> 78
<211> 25
<212> DNA
<213> Homo sapiens

<400> 78
aaatcactaa acctcgttt ctcag 25

<210> 79
<211> 25
<212> DNA
<213> Homo sapiens

<400> 79
agcgtcctta tcttcagag ctaca 25

<210> 80
<211> 25
<212> DNA
<213> Homo sapiens

<400> 80
aaaccccaga cgccatgaaa gctgc 25

<210> 81
<211> 25
<212> DNA
<213> Homo sapiens

<400> 81
gggaacactg ctctcagaca ttaca 25

<210> 82
<211> 25
<212> DNA
<213> Homo sapiens

<400> 82
cttttctg ggtcatgctg caaca 25

<210> 83
<211> 25

<212> DNA
 <213> Homo sapiens

 <400> 83
 gatggggtgc atgtagtctt tggac 25

 <210> 84
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 84
 gaaggtgtga tctgtgggac tgtct 25

 <210> 85
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 85
 gtacgttttt actcagttca tgcgt 25

 <210> 86
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 86
 gcttctcgtg ctgcacatat ttctt 25

 <210> 87
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 87
 gtgaacagac ttgaaactcc agagc 25

 <210> 88
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 88
 atcattttca ggcttctgca gctgt 25

 <210> 89
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 89
 gcatgttgtt tgccaggaca ctgtg 25

 <210> 90
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 90
 ttgtgcttgc tctcagatg gatgg 25

 <210> 91

<211> 25
 <212> DNA
 <213> Homo sapiens

<400> 91
 tagccaggat cctgggtgc ctagg 25

<210> 92
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 92
 gcggcgtgta taccaatgca tggcc 25

<210> 93
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 93
 gaataacttt tggctgtgt gctaa 25

<210> 94
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 94
 tggcccgct gattgtggca tttaa 25

<210> 95
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 95
 cataactgtt agacttcccg ttct 25

<210> 96
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 96
 tcaataaagt tcccctgtga cactc 25

<210> 97
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 97
 ctcccctatg atgggaggct actgg 25

<210> 98
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 98
 atcagttgat tctgtgcc tttt 25

<210> 99
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 99
 tgagccagtg agggcagtcc tgcaa 25

 <210> 100
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 100
 gcatcatcac catagagtcc cagga 25

 <210> 101
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 101
 tcttgctcca aagggtccg tggag 25

 <210> 102
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 102
 atacgccctt ggcacagtcg gatga 25

 <210> 103
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 103
 gtctgctggg tgtgaccatg ttcc 25

 <210> 104
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 104
 gtgcgtggga cgaagacatc ttga 25

 <210> 105
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 105
 tggccccctt cacctgggag aaaag 25

 <210> 106
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 106

gggccaaagca ggacatggcg cggca 25

<210> 107

<211> 25

<212> DNA

<213> Homo sapiens

<400> 107

agcttcagac tcaagtaccc attct 25

<210> 108

<211> 25

<212> DNA

<213> Homo sapiens

<400> 108

gagctgtcta gacgacgctc acaga 25

<210> 109

<211> 25

<212> DNA

<213> Homo sapiens

<400> 109

gagaagcagt ttgacggacc ttgtg 25

<210> 110

<211> 25

<212> DNA

<213> Homo sapiens

<400> 110

ggtgattctg agcgagatct tcttg 25

<210> 111

<211> 25

<212> DNA

<213> Homo sapiens

<400> 111

gcagccttgg aaacctaacg tgcct 25

<210> 112

<211> 25

<212> DNA

<213> Homo sapiens

<400> 112

cctgcagcac cgacgtgtgt gacag 25

<210> 113

<211> 25

<212> DNA

<213> Homo sapiens

<400> 113

acttggtca gtggaagccc tcttt 25

<210> 114

<211> 25

<212> DNA

<213> Homo sapiens

<400> 114
acattgcccg gaaactcagt ctatt 25

<210> 115
<211> 25
<212> DNA
<213> Homo sapiens

<400> 115
gctgtggatc tgtttggcca gggtc 25

<210> 116
<211> 25
<212> DNA
<213> Homo sapiens

<400> 116
gttagagccg atatcactgg aagat 25

<210> 117
<211> 25
<212> DNA
<213> Homo sapiens

<400> 117
agatttgta gccctatctc aaact 25

<210> 118
<211> 25
<212> DNA
<213> Homo sapiens

<400> 118
aggtctccc agaggctgga tacca 25

<210> 119
<211> 25
<212> DNA
<213> Homo sapiens

<400> 119
gtcttccta cctcaggcag gaagg 25

<210> 120
<211> 25
<212> DNA
<213> Homo sapiens

<400> 120
ctacccttat gatgacccat tticc 25

<210> 121
<211> 25
<212> DNA
<213> Homo sapiens

<400> 121
aaatgctctt attttccaa cagct 25

<210> 122
<211> 25
<212> DNA

<213> Homo sapiens

<400> 122
gtgtatagtg tttaccctc ttctt 25

<210> 123
<211> 25
<212> DNA
<213> Homo sapiens

<400> 123
tcatcaaggg ctatgtgcct cccac 25

<210> 124
<211> 25
<212> DNA
<213> Homo sapiens

<400> 124
ccccgcacca gatcaagtag ttgg 25

<210> 125
<211> 25
<212> DNA
<213> Homo sapiens

<400> 125
ccctcctacc agatgacaca gacaa 25

<210> 126
<211> 25
<212> DNA
<213> Homo sapiens

<400> 126
tgtatggtt tcacctggac accgt 25

<210> 127
<211> 25
<212> DNA
<213> Homo sapiens

<400> 127
aactccgtct gtctactgtg tgaga 25

<210> 128
<211> 25
<212> DNA
<213> Homo sapiens

<400> 128
cccactggcc tgaatctaca ctgga 25

<210> 129
<211> 25
<212> DNA
<213> Homo sapiens

<400> 129
acattacatc cgtggattct cctgc 25

<210> 130
<211> 25

<212> DNA
<213> Homo sapiens

<400> 130
ggccctgggc caggtgatt ggact 25

<210> 131
<211> 25
<212> DNA
<213> Homo sapiens

<400> 131
tttagccctc atgactgtat ttct 25

<210> 132
<211> 25
<212> DNA
<213> Homo sapiens

<400> 132
acacgctgca gatcgaggac ttct 25

<210> 133
<211> 25
<212> DNA
<213> Homo sapiens

<400> 133
accggcagcc ctggaagggg cactt 25

<210> 134
<211> 25
<212> DNA
<213> Homo sapiens

<400> 134
taccagcagg aggttctggg cctct 25

<210> 135
<211> 25
<212> DNA
<213> Homo sapiens

<400> 135
catctttcag ggctgccagt ttcgc 25

<210> 136
<211> 25
<212> DNA
<213> Homo sapiens

<400> 136
gggctacgcc aaggactttg accct 25

<210> 137
<211> 25
<212> DNA
<213> Homo sapiens

<400> 137
aatctagtca cctaaccttg tgggt 25

<210> 138

<211> 25
<212> DNA
<213> Homo sapiens

<400> 138
atttcaaaat ttctgcattc acgga 25

<210> 139
<211> 25
<212> DNA
<213> Homo sapiens

<400> 139
aacacctgtc cattgaagat ttcac 25

<210> 140
<211> 25
<212> DNA
<213> Homo sapiens

<400> 140
gacattctta tgcttctttt acaac 25

<210> 141
<211> 25
<212> DNA
<213> Homo sapiens

<400> 141
aatgtttcct aacagttgtg atgtt 25

<210> 142
<211> 25
<212> DNA
<213> Homo sapiens

<400> 142
gggtgaagag agactcgggtg cgggc 25

<210> 143
<211> 25
<212> DNA
<213> Homo sapiens

<400> 143
cgaccgcctg tatgtttgtg taatt 25

<210> 144
<211> 25
<212> DNA
<213> Homo sapiens

<400> 144
aaggcctatc agcttctatc agccc 25

<210> 145
<211> 25
<212> DNA
<213> Homo sapiens

<400> 145
gtcactgggc tgtttgcatt tgata 25

<210> 146
<211> 25
<212> DNA
<213> Homo sapiens

<400> 146
agctcaaaac ttgctaggca tcaga 25

<210> 147
<211> 25
<212> DNA
<213> Homo sapiens

<400> 147
gcagcaaagc tggctccaac atgct 25

<210> 148
<211> 25
<212> DNA
<213> Homo sapiens

<400> 148
aaacactatc tacttccttt gtcac 25

<210> 149
<211> 25
<212> DNA
<213> Homo sapiens

<400> 149
gaggatcatg cccttagcaa gtact 25

<210> 150
<211> 25
<212> DNA
<213> Homo sapiens

<400> 150
aacatcatct tagcaaaggc cagga 25

<210> 151
<211> 25
<212> DNA
<213> Homo sapiens

<400> 151
tgtccttggt acattgaggt taaga 25

<210> 152
<211> 25
<212> DNA
<213> Homo sapiens

<400> 152
tcctactgc atggggactt ccacg 25

<210> 153
<211> 25
<212> DNA
<213> Homo sapiens

<400> 153

ccaccttcac ctggaggga cggag 25

<210> 154

<211> 25

<212> DNA

<213> Homo sapiens

<400> 154

taattacacg ttcaccgtcc aagca 25

<210> 155

<211> 25

<212> DNA

<213> Homo sapiens

<400> 155

ttccctttct accattgatt taaat 25

<210> 156

<211> 25

<212> DNA

<213> Homo sapiens

<400> 156

agtactggtt gaacctgacc acttc 25

<210> 157

<211> 25

<212> DNA

<213> Homo sapiens

<400> 157

aaatacataa gctagtttct gttct 25

<210> 158

<211> 25

<212> DNA

<213> Homo sapiens

<400> 158

gtaacgtgat tgattcagta tctta 25

<210> 159

<211> 25

<212> DNA

<213> Homo sapiens

<400> 159

aatacatccg atgcgtagat tcttg 25

<210> 160

<211> 25

<212> DNA

<213> Homo sapiens

<400> 160

agaatgctcg ggtgctagac tggat 25

<210> 161

<211> 25

<212> DNA

<213> Homo sapiens

<400> 161
ggacggactc tatgaggggc tcaca 25

<210> 162
<211> 25
<212> DNA
<213> Homo sapiens

<400> 162
ggaatgactc aaatgcccaa aacca 25

<210> 163
<211> 25
<212> DNA
<213> Homo sapiens

<400> 163
tcctctcagc tcctaaagca caact 25

<210> 164
<211> 25
<212> DNA
<213> Homo sapiens

<400> 164
acatgttcgc acccaagtgt ggcgg 25

<210> 165
<211> 25
<212> DNA
<213> Homo sapiens

<400> 165
ttcagatgg agtaccagca ccgaa 25

<210> 166
<211> 25
<212> DNA
<213> Homo sapiens

<400> 166
tggcactcgc ggttgagtc agcga 25

<210> 167
<211> 25
<212> DNA
<213> Homo sapiens

<400> 167
ggccatctat gggactgaac ttga 25

<210> 168
<211> 25
<212> DNA
<213> Homo sapiens

<400> 168
gaagaaactg gacaccaaca gtgat 25

<210> 169
<211> 25
<212> DNA

<213> Homo sapiens

<400> 169
gctgtagcca gtgcagacct cactg 25

<210> 170
<211> 25
<212> DNA
<213> Homo sapiens

<400> 170
aggtgtccag tacctaata cgctc 25

<210> 171
<211> 25
<212> DNA
<213> Homo sapiens

<400> 171
ttgccgtgat gaccgtggct atcct 25

<210> 172
<211> 25
<212> DNA
<213> Homo sapiens

<400> 172
gaatacgta ggagccagta cccca 25

<210> 173
<211> 25
<212> DNA
<213> Homo sapiens

<400> 173
gaagctgctc aacaaagtgt ggaag 25

<210> 174
<211> 25
<212> DNA
<213> Homo sapiens

<400> 174
ggaatgtccc agacagacct gtctc 25

<210> 175
<211> 25
<212> DNA
<213> Homo sapiens

<400> 175
cctcatgga cgtcaacagc acctg 25

<210> 176
<211> 25
<212> DNA
<213> Homo sapiens

<400> 176
ggcccgata tggctcgtgg acagc 25

<210> 177
<211> 25

<212> DNA
<213> Homo sapiens

<400> 177
aggagtctcc accagagga ggctc 25

<210> 178
<211> 25
<212> DNA
<213> Homo sapiens

<400> 178
gagcaccact actgctgcat tgtgc 25

<210> 179
<211> 25
<212> DNA
<213> Homo sapiens

<400> 179
actctgacat ttctgttct caagc 25

<210> 180
<211> 25
<212> DNA
<213> Homo sapiens

<400> 180
ccagcatttc cagagcagag cctac 25

<210> 181
<211> 25
<212> DNA
<213> Homo sapiens

<400> 181
gcggtgtcttg agttccatgc aaatt 25

<210> 182
<211> 25
<212> DNA
<213> Homo sapiens

<400> 182
aatggtgaca gtccaaacca ctcca 25

<210> 183
<211> 25
<212> DNA
<213> Homo sapiens

<400> 183
ggccctgcat gtcagatggc gtggt 25

<210> 184
<211> 25
<212> DNA
<213> Homo sapiens

<400> 184
tgatcataaa ttctcccaa ctata 25

<210> 185

<211> 25
 <212> DNA
 <213> Homo sapiens

<400> 185
 aaagttgccca agatgctcct tgttg 25

<210> 186
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 186
 gtctcaccca actgcagttt actat 25

<210> 187
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 187
 gacggagttc ctaagcttca tgaat 25

<210> 188
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 188
 tcagcctggg cagtcttacc aaaat 25

<210> 189
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 189
 tgctgtggct tcaccaacta tacgg 25

<210> 190
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 190
 ccttgactcc ttggtattt cactg 25

<210> 191
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 191
 acagcaggca tcgcaacttt ctgca 25

<210> 192
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 192
 caagagctac aatgtcacct ccgtc 25

<210> 193
<211> 25
<212> DNA
<213> Homo sapiens

<400> 193
gtttgtctct tgggttctg aagga 25

<210> 194
<211> 25
<212> DNA
<213> Homo sapiens

<400> 194
ctctctagtt agatatctga ctgg 25

<210> 195
<211> 25
<212> DNA
<213> Homo sapiens

<400> 195
tccagcctcg ggggttaggt ttctg 25

<210> 196
<211> 25
<212> DNA
<213> Homo sapiens

<400> 196
actcgtctca cgccgtgttt gagat 25

<210> 197
<211> 25
<212> DNA
<213> Homo sapiens

<400> 197
gccagagttc aaatgtgact ccacc 25

<210> 198
<211> 25
<212> DNA
<213> Homo sapiens

<400> 198
caagcccctg tatattgctg atcgg 25

<210> 199
<211> 25
<212> DNA
<213> Homo sapiens

<400> 199
agaccacaga tatctctag acata 25

<210> 200
<211> 25
<212> DNA
<213> Homo sapiens

<400> 200

gtcactgtaa atcattctta agccc 25

<210> 201

<211> 25

<212> DNA

<213> Homo sapiens

<400> 201

aaggctgtca gggcttctgt ttgtt 25

<210> 202

<211> 25

<212> DNA

<213> Homo sapiens

<400> 202

gtagacacct gcacgcatag gattg 25

<210> 203

<211> 25

<212> DNA

<213> Homo sapiens

<400> 203

ttgccacttt ctgcgatat gctgt 25

<210> 204

<211> 25

<212> DNA

<213> Homo sapiens

<400> 204

gccattccag aagtagctta tccta 25

<210> 205

<211> 25

<212> DNA

<213> Homo sapiens

<400> 205

ccaatacccc accgtgatga ctga 25

<210> 206

<211> 25

<212> DNA

<213> Homo sapiens

<400> 206

aattctctgt tatctttacg aggta 25

<210> 207

<211> 25

<212> DNA

<213> Homo sapiens

<400> 207

cacgagaagg actatgacag cctgg 25

<210> 208

<211> 25

<212> DNA

<213> Homo sapiens

<400> 208
tgatttgcca caatgtcctt aactc 25

<210> 209
<211> 25
<212> DNA
<213> Homo sapiens

<400> 209
gctgaaggca cctactcagt atctt 25

<210> 210
<211> 25
<212> DNA
<213> Homo sapiens

<400> 210
agacctggca cttcagtaac tcagc 25

<210> 211
<211> 25
<212> DNA
<213> Homo sapiens

<400> 211
gactcttgaa gtaatggctg atcct 25

<210> 212
<211> 25
<212> DNA
<213> Homo sapiens

<400> 212
gggactttgt caactgcagt acact 25

<210> 213
<211> 25
<212> DNA
<213> Homo sapiens

<400> 213
cctttctta ctccatcatg gctgg 25

<210> 214
<211> 25
<212> DNA
<213> Homo sapiens

<400> 214
atcactaaca ggtctttgac tcagg 25

<210> 215
<211> 25
<212> DNA
<213> Homo sapiens

<400> 215
gaattcatgg tatcctggtt atttt 25

<210> 216
<211> 25
<212> DNA

<213> Homo sapiens
 <400> 216
 aactactgtg aaattctacc agcat 25
 <210> 217
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 217
 gacacctgag cctggathtt cactc 25
 <210> 218
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 218
 tcaaacggcc gaagttgcct ctttt 25
 <210> 219
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 219
 atactaggcc tgtctgtggc attct 25
 <210> 220
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 220
 ggatttagga taagctgtca cccag 25
 <210> 221
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 221
 ggtcagcaac ctctttgat ttgt 25
 <210> 222
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 222
 gacaggtgaa cattccgcc tggtc 25
 <210> 223
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 223
 gattcctgaa ctcaaggtac cagca 25
 <210> 224
 <211> 25

<212> DNA
<213> Homo sapiens

<400> 224
tgttcctttt tgttcaaagt ctatt 25

<210> 225
<211> 25
<212> DNA
<213> Homo sapiens

<400> 225
aggtgttcga aatccgcacc actga 25

<210> 226
<211> 25
<212> DNA
<213> Homo sapiens

<400> 226
gtggagttga tgactttctg ttttc 25

<210> 227
<211> 25
<212> DNA
<213> Homo sapiens

<400> 227
tgcaggtcat gcacacagtt ttgat 25

<210> 228
<211> 25
<212> DNA
<213> Homo sapiens

<400> 228
gctccagaag cgcttggaca ggctg 25

<210> 229
<211> 25
<212> DNA
<213> Homo sapiens

<400> 229
ggcataatgg caaatccttc aagca 25

<210> 230
<211> 25
<212> DNA
<213> Homo sapiens

<400> 230
cccatagag acccaagttc tgcgg 25

<210> 231
<211> 25
<212> DNA
<213> Homo sapiens

<400> 231
gtaaaccaca tctttttgc acttt 25

<210> 232

<211> 25
 <212> DNA
 <213> Homo sapiens

<400> 232
 cacttttgt attatcgtt tcata 25

<210> 233
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 233
 gacgcaggac gagctcagtt gtaga 25

<210> 234
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 234
 tgttgtcag tgtgcctgc actac 25

<210> 235
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 235
 cacttactga cttgcattt tcgtt 25

<210> 236
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 236
 ggtggtttct cttgagactc gttac 25

<210> 237
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 237
 ttggcagtg cattagtaac ttgga 25

<210> 238
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 238
 tcatgtgcac atgccgttgc agcac 25

<210> 239
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 239
 cgttgagaa ctgcagctgc tgtgc 25

<210> 240
<211> 25
<212> DNA
<213> Homo sapiens

<400> 240
agcagctgag gtctcttgag ggagc 25

<210> 241
<211> 25
<212> DNA
<213> Homo sapiens

<400> 241
cattggcctg agtttctgt gcatt 25

<210> 242
<211> 25
<212> DNA
<213> Homo sapiens

<400> 242
gagcctacct ggagggcgag tgcgt 25

<210> 243
<211> 25
<212> DNA
<213> Homo sapiens

<400> 243
gtttcaatt ttgcatgctc gatta 25

<210> 244
<211> 25
<212> DNA
<213> Homo sapiens

<400> 244
ttatgtgtac attattgttg ctatt 25

<210> 245
<211> 25
<212> DNA
<213> Homo sapiens

<400> 245
cttcccggtc actggaaca atagc 25

<210> 246
<211> 25
<212> DNA
<213> Homo sapiens

<400> 246
gtacctgtgg gtagcatca agttc 25

<210> 247
<211> 25
<212> DNA
<213> Homo sapiens

<400> 247

ctgagagcct acctggaggg cctgt 25

<210> 248

<211> 25

<212> DNA

<213> Homo sapiens

<400> 248

ttgggccttt tatctgtcta tccat 25

<210> 249

<211> 25

<212> DNA

<213> Homo sapiens

<400> 249

cagtgtgaca agtgccggga gatct 25

<210> 250

<211> 25

<212> DNA

<213> Homo sapiens

<400> 250

agaatctttt ctatgcctct attcc 25

<210> 251

<211> 25

<212> DNA

<213> Homo sapiens

<400> 251

gccatatagc aggcacgtcc gggtc 25

<210> 252

<211> 25

<212> DNA

<213> Homo sapiens

<400> 252

ggcaagggat aactcttcta acaca 25

<210> 253

<211> 25

<212> DNA

<213> Homo sapiens

<400> 253

ttccctcctt atagtcaagg accgt 25

<210> 254

<211> 25

<212> DNA

<213> Homo sapiens

<400> 254

aataacgcaa atggcttctt ctttc 25

<210> 255

<211> 25

<212> DNA

<213> Homo sapiens

<400> 255
ggcctagcac ggacatggtc tgtcc 25

<210> 256
<211> 25
<212> DNA
<213> Homo sapiens

<400> 256
tgatggacga atgccttttc cgggtg 25

<210> 257
<211> 25
<212> DNA
<213> Homo sapiens

<400> 257
gatcctctgc aatgtgcttg aaaac 25

<210> 258
<211> 25
<212> DNA
<213> Homo sapiens

<400> 258
tcacaagcta ttcctcaaa tctga 25

<210> 259
<211> 25
<212> DNA
<213> Homo sapiens

<400> 259
tgatgagctt tccttgatc cggac 25

<210> 260
<211> 25
<212> DNA
<213> Homo sapiens

<400> 260
gagacagctg tctgtgagg gactg 25

<210> 261
<211> 25
<212> DNA
<213> Homo sapiens

<400> 261
tacacaccac catatatact agctg 25

<210> 262
<211> 25
<212> DNA
<213> Homo sapiens

<400> 262
gtgtaggatga atgcaaactc catcg 25

<210> 263
<211> 25
<212> DNA

<213> Homo sapiens

<400> 263
ttcctcttta aacacccgaa gcgca 25

<210> 264
<211> 25
<212> DNA
<213> Homo sapiens

<400> 264
aactgttcct gactttatac tattt 25

<210> 265
<211> 25
<212> DNA
<213> Homo sapiens

<400> 265
gcataatgat attcacatcc cctca 25

<210> 266
<211> 25
<212> DNA
<213> Homo sapiens

<400> 266
tgttgtttct gcacttata ataaa 25

<210> 267
<211> 25
<212> DNA
<213> Homo sapiens

<400> 267
agaggtgggg ctggatgtct ccatc 25

<210> 268
<211> 25
<212> DNA
<213> Homo sapiens

<400> 268
aagtgcaaag ttattcccca tcttc 25

<210> 269
<211> 25
<212> DNA
<213> Homo sapiens

<400> 269
tacttacacc caaacagatc ctgaa 25

<210> 270
<211> 25
<212> DNA
<213> Homo sapiens

<400> 270
ttgcgtgtgg agctgtattc ccgag 25

<210> 271
<211> 25

<212> DNA
<213> Homo sapiens

<400> 271
ccctttcact gttctcacag gacat 25

<210> 272
<211> 25
<212> DNA
<213> Homo sapiens

<400> 272
gttggctgat attggagtgc tcatt 25

<210> 273
<211> 25
<212> DNA
<213> Homo sapiens

<400> 273
cagcagggtt tcaggtcca atcag 25

<210> 274
<211> 25
<212> DNA
<213> Homo sapiens

<400> 274
ctgtatttgt ggtctctgta ttat 25

<210> 275
<211> 25
<212> DNA
<213> Homo sapiens

<400> 275
acatccaaaa tgacggctgc tatat 25

<210> 276
<211> 25
<212> DNA
<213> Homo sapiens

<400> 276
gtggacccta ctattcatgt ttga 25

<210> 277
<211> 25
<212> DNA
<213> Homo sapiens

<400> 277
gcttaaactt acgtgcctta caggt 25

<210> 278
<211> 25
<212> DNA
<213> Homo sapiens

<400> 278
catgcagact gttagctttt acctt 25

<210> 279

<211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 279
 gtcagtgcac aaagacatac tccaa 25

 <210> 280
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 280
 ggagtcctat ttgcagaacc acttt 25

 <210> 281
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 281
 ataaccaacc tattgcctat gagaa 25

 <210> 282
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 282
 cgaattagtc tccagcctct aaata 25

 <210> 283
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 283
 tcgggtctct ccataattca gccca 25

 <210> 284
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 284
 acgattgcct tcagttgtg ttgtg 25

 <210> 285
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 285
 aattctttt attggtgcct atatt 25

 <210> 286
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 286
 aagctcactg gcatggcctt ccgtg 25

<210> 287
<211> 25
<212> DNA
<213> Homo sapiens

<400> 287
gagatatttc tgaattactg ttgta 25

<210> 288
<211> 25
<212> DNA
<213> Homo sapiens

<400> 288
tttgacgctg gggctggcat tgccc 25

<210> 289
<211> 25
<212> DNA
<213> Homo sapiens

<400> 289
ggacagcaga ctgccggtaa cgcgc 25

<210> 290
<211> 25
<212> DNA
<213> Homo sapiens

<400> 290
gctctgaagt caccacaaa atgct 25

<210> 291
<211> 25
<212> DNA
<213> Homo sapiens

<400> 291
ggtgccaatt tcaagttcca agttg 25

<210> 292
<211> 25
<212> DNA
<213> Homo sapiens

<400> 292
tagggagccg cacctgtca tgtac 25

<210> 293
<211> 25
<212> DNA
<213> Homo sapiens

<400> 293
acgtatatat acctgtgact tgtat 25

<210> 294
<211> 25
<212> DNA
<213> Homo sapiens

<400> 294

taaactgctg cccgtagagg ccttt 25

<210> 295

<211> 25

<212> DNA

<213> Homo sapiens

<400> 295

tggaagttag cggtagcttt ccact 25

<210> 296

<211> 25

<212> DNA

<213> Homo sapiens

<400> 296

tttctctgga tatctgtgta ttttc 25

<210> 297

<211> 25

<212> DNA

<213> Homo sapiens

<400> 297

actgtgcgtt gtacatagtt ctaat 25

<210> 298

<211> 25

<212> DNA

<213> Homo sapiens

<400> 298

gaacactggc cataggaaat gctgt 25

<210> 299

<211> 25

<212> DNA

<213> Homo sapiens

<400> 299

gataccttct gtaggctaata tcttc 25

<210> 300

<211> 25

<212> DNA

<213> Homo sapiens

<400> 300

aacgcagcag aactgccac atcag 25

<210> 301

<211> 25

<212> DNA

<213> Homo sapiens

<400> 301

gaagcttggc ttagtggtga gaatg 25

<210> 302

<211> 25

<212> DNA

<213> Homo sapiens

<400> 302
agcctggctc aatatctaata caatg 25

<210> 303
<211> 25
<212> DNA
<213> Homo sapiens

<400> 303
gaactcaaaa catttgggac cacct 25

<210> 304
<211> 25
<212> DNA
<213> Homo sapiens

<400> 304
ccaccctagt gtctcatgtt tgtat 25

<210> 305
<211> 25
<212> DNA
<213> Homo sapiens

<400> 305
tggatgatgc tgctactatg ccagc 25

<210> 306
<211> 25
<212> DNA
<213> Homo sapiens

<400> 306
aagctggagg ctgagatcgc cacct 25

<210> 307
<211> 25
<212> DNA
<213> Homo sapiens

<400> 307
cactttaacc cagcctcaa tgaga 25

<210> 308
<211> 25
<212> DNA
<213> Homo sapiens

<400> 308
tggaatgagt ctctttagt ttcag 25

<210> 309
<211> 25
<212> DNA
<213> Homo sapiens

<400> 309
agagacatgt accttgacca tcgtc 25

<210> 310
<211> 25
<212> DNA

<213> Homo sapiens

<400> 310
tggaatcct ctaggaggct cctcc 25

<210> 311
<211> 25
<212> DNA
<213> Homo sapiens

<400> 311
tgtgtgggtg ctgataattg tgtat 25

<210> 312
<211> 25
<212> DNA
<213> Homo sapiens

<400> 312
tactctcca tcatctagaa ttgtt 25

<210> 313
<211> 25
<212> DNA
<213> Homo sapiens

<400> 313
gcaccctcag ggtcttaagg tcttc 25

<210> 314
<211> 25
<212> DNA
<213> Homo sapiens

<400> 314
gtacacagtc tgtttctat ttgtt 25

<210> 315
<211> 25
<212> DNA
<213> Homo sapiens

<400> 315
tgggctgcct tatatttgtg gtgtg 25

<210> 316
<211> 25
<212> DNA
<213> Homo sapiens

<400> 316
taaccagac ttaatctga atgat 25

<210> 317
<211> 25
<212> DNA
<213> Homo sapiens

<400> 317
gagacttcgg cggaccatta ggaat 25

<210> 318
<211> 25

<212> DNA
 <213> Homo sapiens

 <400> 318
 atgcacgtga aacttaacac ttat 25

 <210> 319
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 319
 agttgaacag tgtgccctag ctttt 25

 <210> 320
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 320
 gatcttgtat tcagtcaggt taaaa 25

 <210> 321
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 321
 ccaaagtggc cagggcggcc tctgg 25

 <210> 322
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 322
 ggacgagtcg gaccgaggct aggac 25

 <210> 323
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 323
 atttgtggcc actcacttg tagga 25

 <210> 324
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 324
 tctgtcca aagggctccg tggag 25

 <210> 325
 <211> 25
 <212> DNA
 <213> Homo sapiens

 <400> 325
 ctctccatg agtctgacat ctggg 25

 <210> 326

<211> 25
 <212> DNA
 <213> Homo sapiens

<400> 326
 gggtgaggag aggctccaga cccgc 25

<210> 327
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 327
 acagcaactt gtcaaaccta agcat 25

<210> 328
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 328
 acgcagagga cgtctctatg ccggt 25

<210> 329
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 329
 gtagtcattc atttctagct gtaca 25

<210> 330
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 330
 gtagctgatg aagtatgtcg cattt 25

<210> 331
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 331
 gatgatcttc tgtggtgctt aagga 25

<210> 332
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 332
 ctgtgattgc tgccaggcac tgttc 25

<210> 333
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 333
 ttcacgcgga aatacacgct gcccc 25

<210> 334
<211> 25
<212> DNA
<213> Homo sapiens

<400> 334
tcactgtctg gtccgtggga cgggtg 25

<210> 335
<211> 25
<212> DNA
<213> Homo sapiens

<400> 335
gaaggagagc catgctact ttcta 25

<210> 336
<211> 25
<212> DNA
<213> Homo sapiens

<400> 336
gtgtatagtg tttaccctc ttctt 25

<210> 337
<211> 25
<212> DNA
<213> Homo sapiens

<400> 337
caggactttt gttccaggtt gccag 25

<210> 338
<211> 25
<212> DNA
<213> Homo sapiens

<400> 338
gtgcagttc tgacactgt tgttg 25

<210> 339
<211> 25
<212> DNA
<213> Homo sapiens

<400> 339
gggagcaccg tgatggagag gacag 25

<210> 340
<211> 25
<212> DNA
<213> Homo sapiens

<400> 340
aaggccaaga tcaccatcgt ggcag 25

<210> 341
<211> 25
<212> DNA
<213> Homo sapiens

<400> 341

tattcaaatg gccctccag aaagt	25
<210> 342	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 342	
aagccacatc tgagcactc aagac	25
<210> 343	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 343	
gttccaagaa ctctggtgtc tgacc	25
<210> 344	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 344	
gaggatcatg cccttagcaa gtact	25
<210> 345	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 345	
ggtctacgcc tattacaacc tggag	25
<210> 346	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 346	
aatcgttctc ctacaatca agttc	25
<210> 347	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 347	
ggaactaggc tcttatgtgt gcctt	25
<210> 348	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 348	
tctgtctcc taagaaatct gccca	25
<210> 349	
<211> 25	
<212> DNA	
<213> Homo sapiens	

<400> 349
gaggctcagt tagcaacctg tgttg 25

<210> 350
<211> 25
<212> DNA
<213> Homo sapiens

<400> 350
agactgtccc tgaaacctag tactg 25

<210> 351
<211> 25
<212> DNA
<213> Homo sapiens

<400> 351
actttctgca tgtgatgcc cgaag 25

<210> 352
<211> 25
<212> DNA
<213> Homo sapiens

<400> 352
gttcttgat tgtatgccc agggg 25

<210> 353
<211> 25
<212> DNA
<213> Homo sapiens

<400> 353
aaaaaccgca gccacgggtc tgatc 25

<210> 354
<211> 25
<212> DNA
<213> Homo sapiens

<400> 354
gacatgaacg gctgttactc acgcc 25

<210> 355
<211> 25
<212> DNA
<213> Homo sapiens

<400> 355
tgttctggc gctgtatgtg caggc 25

<210> 356
<211> 25
<212> DNA
<213> Homo sapiens

<400> 356
cttctacagg ctttgggaa gtagg 25

<210> 357
<211> 25
<212> DNA

<213> Homo sapiens
 <400> 357
 tggaggacac agatgactct ttggt 25
 <210> 358
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 358
 gagtgggtgg ggaattttct cctct 25
 <210> 359
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 359
 ctgcaccttc atcagcggca tgatg 25
 <210> 360
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 360
 tataaaccac tctcaacag ctggc 25
 <210> 361
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 361
 aatacatccg atgcgtagat tcttg 25
 <210> 362
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 362
 tggcttagcc tgggcaggtc gtgtc 25
 <210> 363
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 363
 gagcagggtc tctctaaagg ggact 25
 <210> 364
 <211> 25
 <212> DNA
 <213> Homo sapiens
 <400> 364
 gtctctgagt tacaaaagtg ctaat 25
 <210> 365
 <211> 25

<212> DNA
<213> Homo sapiens

<400> 365
accagttttt ggcttactcc tgaga 25

<210> 366
<211> 25
<212> DNA
<213> Homo sapiens

<400> 366
ttgtctacac tcaggctgca gtatt 25

<210> 367
<211> 25
<212> DNA
<213> Homo sapiens

<400> 367
tctgagaga tacctcacat tccaa 25

<210> 368
<211> 25
<212> DNA
<213> Homo sapiens

<400> 368
ggtagcctt agtttctcag acttg 25

<210> 369
<211> 25
<212> DNA
<213> Homo sapiens

<400> 369
tgtctcatc tctgcaaagt tcagc 25

<210> 370
<211> 25
<212> DNA
<213> Homo sapiens

<400> 370
ttccaagga caggcttcag catca 25

<210> 371
<211> 25
<212> DNA
<213> Homo sapiens

<400> 371
ctoctacgca gatgctgtca ctggc 25

<210> 372
<211> 25
<212> DNA
<213> Homo sapiens

<400> 372
tgagcaaggg ggcccgaatc gacca 25

<210> 373

<211> 25
<212> DNA
<213> Homo sapiens

<400> 373
aagcctgctc tcttgcaaag acagc 25

<210> 374
<211> 25
<212> DNA
<213> Homo sapiens

<400> 374
taactataag gtgcctcagt ttcc 25

<210> 375
<211> 25
<212> DNA
<213> Homo sapiens

<400> 375
catatggcat ctctcaactt ttctt 25

<210> 376
<211> 25
<212> DNA
<213> Homo sapiens

<400> 376
gaaagttcag ccagaatctt cgtcc 25

<210> 377
<211> 25
<212> DNA
<213> Homo sapiens

<400> 377
gggcagcttc ctaatatgct ttaca 25

<210> 378
<211> 25
<212> DNA
<213> Homo sapiens

<400> 378
gcctgtcttg agtagacttg gaccc 25

<210> 379
<211> 25
<212> DNA
<213> Homo sapiens

<400> 379
gatccacctc atatgtgagt ccgtc 25

<210> 380
<211> 25
<212> DNA
<213> Homo sapiens

<400> 380
cgctgctgtg tcaactcatga ctgtt 25

<210> 381
<211> 25
<212> DNA
<213> Homo sapiens

<400> 381
gcctgttgaa ggtacatcgt ttgca 25

<210> 382
<211> 25
<212> DNA
<213> Homo sapiens

<400> 382
tctcccgatt cccaactgag tgta 25

<210> 383
<211> 25
<212> DNA
<213> Homo sapiens

<400> 383
aggcaatttt aatcccctgc actag 25

<210> 384
<211> 25
<212> DNA
<213> Homo sapiens

<400> 384
ggacaaactg ccagtttgt ttct 25

<210> 385
<211> 25
<212> DNA
<213> Homo sapiens

<400> 385
gttcttcaat cagctggcac acact 25

<210> 386
<211> 25
<212> DNA
<213> Homo sapiens

<400> 386
accaccagg aaaccatga caaga 25

<210> 387
<211> 25
<212> DNA
<213> Homo sapiens

<400> 387
aaggacagaa cattgccttc ctgt 25

<210> 388
<211> 25
<212> DNA
<213> Homo sapiens

<400> 388

ggatatacaa gttctgtggt ttcag 25
<210> 389
<211> 25
<212> DNA
<213> Homo sapiens

<400> 389
gtggttgga ctagactggt ggcag 25

<210> 390
<211> 25
<212> DNA
<213> Homo sapiens

<400> 390
atgtactgt tctacctga ttgtc 25

<210> 391
<211> 25
<212> DNA
<213> Homo sapiens

<400> 391
cgcggtgca tgacctga ggtca 25

<210> 392
<211> 25
<212> DNA
<213> Homo sapiens

<400> 392
ggtgcccgga tccagatgt gaaga 25

<210> 393
<211> 25
<212> DNA
<213> Homo sapiens

<400> 393
ctccatccta tgctgtaca attgc 25

<210> 394
<211> 25
<212> DNA
<213> Homo sapiens

<400> 394
tgaatgccag tccagatgtg cctag 25

<210> 395
<211> 25
<212> DNA
<213> Homo sapiens

<400> 395
ttacattcat ttaacctgcc gatta 25

<210> 396
<211> 25
<212> DNA
<213> Homo sapiens

<400> 396
caccacgtgg tacttggtg cctac 25

<210> 397
<211> 25
<212> DNA
<213> Homo sapiens

<400> 397
agtgctggc actaaccaac aaggt 25

<210> 398
<211> 25
<212> DNA
<213> Homo sapiens

<400> 398
ctctctgaaa cgcttcctat aactc 25

<210> 399
<211> 25
<212> DNA
<213> Homo sapiens

<400> 399
ttgtcagttc tcaatttcct gtgtt 25

<210> 400
<211> 25
<212> DNA
<213> Homo sapiens

<400> 400
attatcgctt tatgttttg gttat 25

<210> 401
<211> 25
<212> DNA
<213> Homo sapiens

<400> 401
ggggctcaag ggcaagatca gcgag 25

<210> 402
<211> 25
<212> DNA
<213> Homo sapiens

<400> 402
gaataggtat cttgagcttg gttct 25

<210> 403
<211> 25
<212> DNA
<213> Homo sapiens

<400> 403
acactcgctt ctggaacgtc tgagg 25

<210> 404
<211> 25
<212> DNA

<213> Homo sapiens

<400> 404

aatctatatt tgtctccga tcaac 25

<210> 405

<211> 25

<212> DNA

<213> Homo sapiens

<400> 405

acaaggccca ggctggggcc agggc 25

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US20120177726A \[0043\]](#)
- [US20110201515A \[0067\]](#)
- [US20110229888A \[0067\]](#)
- [US20130017971A \[0067\]](#)
- [US5822715A \[0092\]](#)
- [US20030073083A \[0092\]](#)
- [US20050227266A \[0092\]](#)
- [US20050206512A \[0092\]](#)
- [US20050123945A \[0092\]](#)
- [US20030129629A \[0092\]](#)
- [US20020006613A \[0092\]](#)

Non-patent literature cited in the description

- **CASAGRANDE et al.** Clinical Cancer Research, 2014, vol. 20, 21 [0005]
- **DE JONGE et al.** Eur J Cancer., 2010, vol. 46, 163016-21 [0043]
- **MORTAZAVI et al.** Nat. Methods, 2008, vol. 5, 621-628 [0074]
- **CHEN et al.** Mol. Cancer Ther., 2012, vol. 11, 34-33 [0076]
- **WANG et al.** J. Nat. Cancer Inst., 2013, vol. 105, 1284-1291 [0076]
- **KNUDSEN et al.** PLoS One, 2014, vol. 9, e87415- [0076]
- **VAPNIK V N.** Statistical Learning Theory John Wiley & Sons 19980000 [0092]
- The Elements of Statistical Learning: Data Mining **HASTIE et al.** Inference, and Prediction Springer 20010000 [0092]
- **AGRESTI** An Introduction to Categorical Data Analysis John Wiley & Sons 19960000 [0092]
- Neural Network Modeling of Physiological Processes **V. TRESP et al.** Computational Learning Theory and Natural Learning Systems 2 MIT Press 19940000 [0092]
- Handbook of Pharmaceutical Excipients Pharmaceutical Press 20090000 [0239]
- **MIRTTI et al.** APMIS, 2009, vol. 117, 151-161 [0251]

PATENTKRAV

1. Fremgangsmåde in-vitro til test af ekspresion i et cancervæv fra en patient af én eller flere sensitivitets- eller resistens biomarkører for et sekretorisk phospholipase A₂- (sPLA₂) hydrolyserbart, cisplatin-holdigt liposom, der omfatter bestemmelse af et niveau af PLA2G2A eller et komplement deraf i en tumorprøve fra patienten og:

(a) bestemmelse af et ekspressionsniveau i vævsprøven af én eller flere sensitivitsbiomarkører, der er udvalgt fra Tabel 1 eller et komplement deraf, hvor sensitivitet refererer til en celled eller et vævs modtagelighed for behandling med det sPLA₂-hydrolyserbare, cisplatin-holdige liposom; eller

(b) bestemmelse af et ekspressionsniveau i vævsprøven af én eller flere resistensbiomarkører, der er udvalgt fra Tabel 2 eller et komplement deraf, hvor resistens henviser til en celled eller et vævs manglende modtagelighed for behandling med det sPLA₂-hydrolyserbare, cisplatin-holdige liposom.

2. Fremgangsmåde ifølge krav 1, hvor:

(a) resultaterne fra trin (a) bestemmes til i det væsentlige at svare til ekspressionsniveauet for den ene eller flere sensitivitsbiomarkører i en referencecelle eller et referencevæv, der er kendt for at have sensitivitet, hvor sensitiviteten er baseret på GI50-data fra NCI 60-cellelinjer; eller

(b) resultaterne fra trin (a) bestemmes til i det væsentlige at svare til ekspressionsniveauet for den ene eller flere resistensbiomarkører i en referencecelle eller et referencevæv, der er kendt for at have resistens, hvor resistensen er baseret på GI50-data fra NCI 60-cellelinjer.

3. Fremgangsmåde ifølge krav 1 eller 2, hvor prøven er en tumorprøve.

4. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 3, hvor:

(a) ekspressionsniveauet for den ene eller flere sensitivitsbiomarkører i prøven bestemmes ved indsamling af nukleinsyremolekyler fra prøven og påvisning af ekspressionsniveauet for den ene eller flere biomarkører ved anvendelse af en anordning, hvortil én eller flere enkeltstrengede nukleinsyremolekyler er bundet, hvor

nukleinsyremolekylet specifik hybridiserer til den ene eller flere sensitivitsbiomarkører, der er udvalgt fra Tabel 1; eller

(b) ekspressionsniveauet for den ene eller flere resistensbiomarkører i prøven bestemmes ved indsamling af nukleinsyremolekyler fra prøven og påvisning af ekspressionsniveauet for den ene eller flere biomarkører ved anvendelse af en anordning, hvortil én eller flere enkeltstrengede nukleinsyremolekyler er bundet, hvor nukleinsyremolekylet specifik hybridiserer til den ene eller flere resistensbiomarkører, der er udvalgt fra Tabel 2.

5. Fremgangsmåde ifølge krav 4, hvor:

- (a) anordningen er et mikroarray eller er beregnet til udførelse af en kvantitativ omvendt transkriptasepolymerasekædereaktion (qRT-PCR); og
- (b) anordningens ene eller flere enkeltstrengede nukleinsyremolekyler har en længde i intervallet fra 10 til 100 nukleotider i længden; og
- (c) det ene eller flere enkeltstrengede nukleinsyremolekyler er mærkede eller immobiliserede på et fast substrat.

6. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 5, hvor det ene eller flere af det enkeltstrengede nukleinsyremolekyler har en længde i intervallet fra 20 til 60 nukleotider.

7. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 6, hvor:

- (a) canceren hos patienten bestemmes til at være modtagelig for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom, hvis niveauet af den ene eller flere sensitivitsbiomarkører eller komplementet deraf i prøven i det væsentlige svarer til niveauet af den ene eller flere sensitivitsbiomarkører eller komplementet deraf i den referencecelle eller det referencevæv, der er kendt for at være sensitiv(t) over for det sPLA₂-hydrolyserbare liposom; eller
- (b) canceren hos patienten bestemmes til at være modtagelig for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom, hvis niveauet af den ene eller flere resistensbiomarkører eller komplementet deraf i prøven i det væsentlige svarer til niveauet af den ene eller flere resistensbiomarkører eller komplementet deraf i den referencecelle

eller det referencevæv, der er kendt for at være sensitiv(t) over for det sPLA₂-hydrolyserbare liposom; eller

(c) canceren hos patienten bestemmes til at være ikke-modtagelig for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom, hvis niveauet af den ene eller flere sensitivitsbiomarkører eller komplementet deraf i prøven i det væsentlige adskiller sig fra niveauet af den ene eller flere sensitivitsbiomarkører eller komplementet deraf i den referencecelle eller det referencevæv, der er kendt for at være sensitiv(t) over for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom; eller

(d) canceren hos patienten bestemmes til at være ikke-modtagelig for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom, hvis niveauet af den ene eller flere resistensbiomarkører eller komplementet deraf i prøven i det væsentlige svarer til niveauet af den ene eller flere resistensbiomarkører eller komplementet deraf i den referencecelle eller det referencevæv, der er kendt for at være resistent over for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom.

8. Fremgangsmåde ifølge krav 2, hvor den referencecelle eller det referencevæv, der er kendt for at være sensitiv(t) over for liposomet, eller cellen eller vævet, der er kendt for at være resistent over for liposomet, er af samme cancertype som en celle eller et væv i den tumorprøve, hvorfra den ene eller flere nukleinsyremolekyler er afledt.

9. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 8, hvor den ene eller flere sensitivitsbiomarkører ikke er C1QR1 (SEQ ID NO: 13), SLA (SEQ ID NO: 48), PTPN7 (SEQ ID NO: 77), CENTB1 (SEQ ID NO: 37), IFI16 (SEQ ID NO: 17 eller 261), ARHGEF6 (SEQ ID NO: 36 eller 294), CD3D (SEQ ID NO: 81), ARHGAP15 (SEQ ID NO: 30), HCLS1 (SEQ ID NO: 16 eller 259), CD53 (SEQ ID NO: 282), PTPRCAP (SEQ ID NO: 8) eller PTPRC (SEQ ID NO: 10, 18, 25 eller 243).

10. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 9, hvor canceren er udvalgt fra:

(a) gruppen, der består af en solid tumorcancer og en hæmatologisk cancer; eller

(b) gruppen, der består af brystcancer, akut myeloid leukæmi (AML), akut lymfoblastisk leukæmi (ALL), kronisk lymfocytisk leukæmi (CLL), myelodysplastisk syndrom (MDS),

kronisk myeloid leukæmi – kronisk fase (CMLCP), diffust stort B-cellelymfom (DLBCL), kutant T-cellelymfom (CTCL), periferisk T-cellelymfom (PTCL), Hodgkins lymfom, hepatocellulært karcinom (HCC), cervixcancer, renalcellekarcinom (RCC), øsofagus cancer, melanom, gliom, pankreascancer, gastrointestinale stromatumorer (GIST), sarkom, ikke-småcellet lungekarcinom (NSCLC), prostatacancer, ovariecancer, coloncancer, blærecancer og pladeepitelkarcinom i hoved og hals (SCCHN); eller

(c) gruppen, der består af ubehandlet cancer og recidiverende cancer efter én eller flere behandlinger.

11. Fremgangsmåde ifølge krav 10, hvor brystcanceren er østrogenreceptor-positiv (ER-pos) eller er en metastatisk form af brystcancer.

12. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 10, hvor den ene eller flere sensitivetsbiomarkører er

(a) udvalgt fra én eller flere af COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 eller 272), FNBP1 (SEQ ID NO: 14 eller 28), CBFB (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 eller 54) og CAP350 (SEQ ID NO: 20 eller 61); eller

(b) COL5A2 (SEQ ID NO: 73 eller 211) og ITGA4 (SEQ ID NO: 1); eller

(c) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1) og MSN (SEQ ID NO: 2); eller

(d) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2) og FAM46A (SEQ ID NO: 3 eller 280); eller

(e) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4) og DOCK2 (SEQ ID NO: 5 eller 223); eller

(f) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223) og EVL (SEQ ID NO: 6); eller

- (g) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6) og SACS (SEQ ID NO: 7); eller
- (h) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7) og PTPRCAP (SEQ ID NO: 8); eller
- (i) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8) og EBI2 (SEQ ID NO: 9); eller
- (j) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9) og PTPRC (SEQ ID NO: 10, 18, 25 eller 243); eller
- (k) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25 eller 243) og ANP32E (SEQ ID NO: 11); eller
- (l) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25 eller 243), ANP32E (SEQ ID NO: 11) og SFPQ (SEQ ID NO: 12, 38 eller 272); eller
- (m) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25 eller 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 eller 272) og C1QR1 (SEQ ID NO: 13); eller
- (n) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25 eller 243), ANP32E (SEQ ID NO: 11), SFPQ

(SEQ ID NO: 12, 38 eller 272), C1QR1 (SEQ ID NO: 13) og FNBP1 (SEQ ID NO: 14 eller 28); eller

(o) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), PTPRCAP (SEQ ID NO: 8), EBI2 (SEQ ID NO: 9), PTPRC (SEQ ID NO: 10, 18, 25 eller 243), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 eller 272), C1QR1 (SEQ ID NO: 13), FNBP I (SEQ ID NO: 14 eller 28) og CBFB (SEQ ID NO: 15); eller

(p) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 eller 272), FNBP1 (SEQ ID NO: 14 eller 28), CBFB (SEQ ID NO: 15) og SFRS7 (SEQ ID NO: 19 eller 54); eller

(q) COL5A2 (SEQ ID NO: 73 eller 211), ITGA4 (SEQ ID NO: 1), MSN (SEQ ID NO: 2), FAM46A (SEQ ID NO: 3 eller 280), ITGB2 (SEQ ID NO: 4), DOCK2 (SEQ ID NO: 5 eller 223), EVL (SEQ ID NO: 6), SACS (SEQ ID NO: 7), EBI2 (SEQ ID NO: 9), ANP32E (SEQ ID NO: 11), SFPQ (SEQ ID NO: 12, 38 eller 272), FNBP1 (SEQ ID NO: 14 eller 28), CBFB (SEQ ID NO: 15), SFRS7 (SEQ ID NO: 19 eller 54) og CAP350 (SEQ ID NO: 20 eller 61).

13. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 11, hvor den ene eller flere resistensbiomarkører er

(a) udvalgt fra én eller flere af SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 eller 400), KRT18 (SEQ ID NO: 108 eller 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 eller 312), LRP5 (SEQ ID NO: 112), KRT18 (SEQ ID NO: 108 eller 306), DSG2 (SEQ ID NO: 111 eller 312), SOX9 (SEQ ID NO: 121, 315 eller 319) og IER3 (SEQ ID NO: 127 eller 317); eller

(b) SFN (SEQ ID NO: 96 eller 324) og LISCH7 (SEQ ID NO: 97); eller

(c) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97) og EPB41L4B (SEQ ID NO: 98); eller

- (d) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98) og MST1R (SEQ ID NO: 99); eller
- (e) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99) og ITGB4 (SEQ ID NO: 100); eller
- (f) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100) og DBNDD2 (SEQ ID NO: 102 eller 365); eller
- (g) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365) og TACSTDI (SEQ ID NO: 104); eller
- (h) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104) og MISP (SEQ ID NO: 105); eller
- (i) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTDI (SEQ ID NO: 104), MISP (SEQ ID NO: 105) og KRT8 (SEQ ID NO: 106); eller
- (j) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106) og JUP (SEQ ID NO: 107 eller 400); eller
- (k) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 eller 400) og KRT18 (SEQ ID NO: 108 eller 306); eller
- (l) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 eller 400), KRT18 (SEQ ID NO: 108 eller 306) og FA2H (SEQ ID NO: 109); eller
- (m) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102

eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 eller 400), KRT18 (SEQ ID NO: 108 eller 306), FA2H (SEQ ID NO: 109) og MGAT4B (SEQ ID NO: 110); eller

(n) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 eller 400), KRT18 (SEQ ID NO: 108 eller 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110) og DSG2 (SEQ ID NO: 111 eller 312); eller

(o) SFN (SEQ ID NO: 96 eller 324), LISCH7 (SEQ ID NO: 97), EPB41L4B (SEQ ID NO: 98), MST1R (SEQ ID NO: 99), ITGB4 (SEQ ID NO: 100), DBNDD2 (SEQ ID NO: 102 eller 365), TACSTD1 (SEQ ID NO: 104), MISP (SEQ ID NO: 105), KRT8 (SEQ ID NO: 106), JUP (SEQ ID NO: 107 eller 400), KRT18 (SEQ ID NO: 108 eller 306), FA2H (SEQ ID NO: 109), MGAT4B (SEQ ID NO: 110), DSG2 (SEQ ID NO: 111 eller 312) og LRP5 (SEQ ID NO: 112); eller

(p) KRT18 (SEQ ID NO: 108 eller 306); eller

(q) KRT18 (SEQ ID NO: 108 eller 306) og DSG2 (SEQ ID NO: 111 eller 312); eller

(r) KRT18 (SEQ ID NO: 108 eller 306), DSG2 (SEQ ID NO: 111 eller 312) og SOX9 (SEQ ID NO: 121, 315 eller 319); eller

(s) KRT18 (SEQ ID NO: 108 eller 306), DSG2 (SEQ ID NO: 111 eller 312), SOX9 (SEQ ID NO: 121, 315 eller 319) og IER3 (SEQ ID NO: 127 eller 317).

14. Fremgangsmåde ifølge et hvilket som helst af krav 1 til 12, hvor fremgangsmåden endvidere omfatter:

(a) omdannelse af niveauet af den ene eller flere sensitivitsbiomarkører eller komplementet deraf til en middelscore, hvor middelscoren indikerer patientens modtagelighed for det sPLA₂-hydrolyserbare, cisplatin-holdige liposom; eller

(b) omdannelse af niveauet af den ene eller flere resistensbiomarkører eller komplementet deraf til en middelscore, hvor middelscoren indikerer patientens modtagelighed for de sPLA₂-hydrolyserbare, cisplatin-holdige liposomer.

15. Fremgangsmåde ifølge krav 13, hvor fremgangsmåden endvidere omfatter fratrækning af middelscoren for den ene eller flere af resistensbiomarkørerne fra

middelscoren for den ene eller flere af sensitivitsbiomarkørerne for at opnå en forskelsscore, hvor forskelsscoren indikerer patientens modtagelighed for det sPLA2-hydrolyserbare, cisplatin-holdige liposom.

16. Fremgangsmåde ifølge krav 14, hvor fremgangsmåden endvidere omfatter sammenligning af forskelsscoren med en afskæringsværdi, hvor forskelsscoren over en afskæringsværdi indikerer, at patienten er modtagelig for det sPLA2-hydrolyserbare, cisplatin-holdige liposom.

17. Fremgangsmåde ifølge krav 16, hvor afskæringsværdien er ca. 0,25, ca. 0,35, ca. 0,4, ca. 0,45, ca. 0,5 eller større.

18. Sammensætning, der omfatter et sekretorisk phospholipase A2- (sPLA2) hydrolyserbart cisplatin-holdigt liposom til anvendelse i behandling af en patient med cancer, der er diagnosticeret ifølge fremgangsmåden ifølge et hvilket som helst af krav 1 til 17.

19. Sammensætning ifølge krav 18, hvor sammensætningen er formuleret til administration til patienten:

- (a) som en enkelt behandling til cancerpatienten eller
- (b) som en behandling til cancerpatienten i kombination med én eller flere andre supplerende cancerterapier end det sPLA2-hydrolyserbare liposom, hvor de supplerende cancerterapier omfatter kirurgi, stråling eller et terapeutisk middel, og hvor eventuelt det terapeutiske middel er udvalgt fra gruppen bestående af docetaxel, cabazitaxel, mitoxantron, estramustin, prednison, carboplatin, bevacizumab, paclitaxel, gemcitabin, doxorubicin, topotecan, etoposid, tamoxifen, letrozol, sorafenib, fluoruracil, capecitabin, oxaliplatin, interferon-alpha, 5-fluoruracil (5-FU), en histondeacetylase- (HDAC) hæmmer, ipilimumab, bortezomib, carfilzomib, thalidomid, lenalidomid, pomalidomid, dexamethason, cyclophosphamid, vincristin, melphalan, tegafur, irinotecan, cetuximab, leucovorin, SN-38, everolimus, temsirolimus, bleomycin, lomustin, depsipeptid, erlotinib, busulfan, epirubicin, arsentrioxid, bendamustin, fulvestrant, teniposid, adriamycin, decitabin, estramustin, azaguanin, aclarubicin, mitomycin, paclitaxel, taxotere, APO010,

ara-c, methylprednisolon, methotrexat, methyl-gag, belinostat, idarubicin, IL4-PR38, valproesyre, hel-trans-retinsyre (ATRA), cytoxan, suberoylanilid-hydroxamsyre, leukeran, fludarabin, vinblastin, dacarbazin, hydroxyurea, tegafur, daunorubicin, mechlorethamin, streptozocin, carmustin, mercaptopurin, dactinomycin, tretinoin, ifosfamid, floxuridin, thioguanin, PSC 833, herceptin, celecoxib, iressa, anastrozol og rituximab; eller

(c) via en intravenøs, intramuskulær, transdermal, intradermal, intraarteriel, intrakraniel, subkutan, intraorbital, intraventrikulær, intraspinal, intraperitoneal eller intranasal vej; eller

(d) ved en dosis på ca. 60 mg/m² cisplatin, ca. 75 mg/m² cisplatin eller ca. 90 mg/m² cisplatin; eller

(e) ved en frekvens på mindst én, to eller tre uger eller på dag 1 og dag 8 af en 3 ugers behandling; eller

(f) i en behandling, der er gentagne doser to til tyve gange.

DRAWINGS




FIG. 1