



(86) Date de dépôt PCT/PCT Filing Date: 2017/10/13
(87) Date publication PCT/PCT Publication Date: 2018/05/24
(45) Date de délivrance/Issue Date: 2021/06/15
(85) Entrée phase nationale/National Entry: 2019/04/16
(86) N° demande PCT/PCT Application No.: DK 2017/050342
(87) N° publication PCT/PCT Publication No.: 2018/091050
(30) Priorités/Priorities: 2016/11/18 (US15/356,175);
2016/11/18 (DK PCT/DK2016/050377)

(51) Cl.Int./Int.Cl. *A61K 9/00* (2006.01),
A61K 31/00 (2006.01), *A61K 47/26* (2006.01),
A61K 9/20 (2006.01)
(72) Inventeur/Inventor:
WITTORFF, HELLE, DK
(73) Propriétaire/Owner:
FERTIN PHARMA A/S, DE
(74) Agent: SMART & BIGGAR LLP

(54) Titre : COMPRIME COMPRENANT UN LIANT SEPARÉ ET DE L'ERYTHRITOL COMPRESSIBLE DE FAÇON NON DIRECTE

(54) Title: TABLET COMPRISING SEPARATE BINDER AND NON-DIRECT COMPRESSION ERYTHRITOL

(57) **Abrégé/Abstract:**

The invention relates to an oral delivery vehicle tablet, the delivery vehicle tablet being formed by compression of a plurality of particles, wherein the oral delivery vehicle tablet comprises sugar alcohol(s) in an amount of 40 to 99.9 % by weight of the delivery vehicle tablet, wherein the oral delivery vehicle tablet comprises non-direct compressible (non-DC) erythritol particles in an amount of 15-50% by weight of the delivery vehicle tablet, wherein the oral delivery vehicle tablet comprises one or more binders other than binders granulated to the sugar alcohol(s) in an amount of 0.1 to 6% by weight of the tablet, wherein the delivery vehicle tablet comprises a plurality of further sugar alcohol particles in an amount of at least 10% by weight of the delivery vehicle tablet and wherein said further sugar alcohols particles have a composition which is different from said non-DC erythritol.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

24 May 2018 (24.05.2018)



(10) International Publication Number

WO 2018/091050 A1

(51) International Patent Classification:

A61K 9/00 (2006.01)

A61K 47/26 (2006.01)

A61K 9/20 (2006.01)

A61K 31/00 (2006.01)

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

(21) International Application Number:

PCT/DK2017/050342

Published:

— with international search report (Art. 21(3))

(22) International Filing Date:

13 October 2017 (13.10.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/DK2016/050377

18 November 2016 (18.11.2016) DK

15/356,175

18 November 2016 (18.11.2016) US

(71) Applicant: FERTIN PHARMA A/S [DK/DK]; Dandyvej

19, 7100 Vejle (DK).

(72) Inventor: WITTORFF, Helle; Johannebjergparken 25,

7120 Vejle Øst (DK).

(74) Agent: PATENTGRUPPEN A/S; Aaboulevarden 31, 4th

floor, DK-8000 Aarhus C (DK).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: TABLET COMPRISING SEPARATE BINDER AND ERYTHRITOL

(57) Abstract: The invention relates to an oral delivery vehicle tablet, the delivery vehicle tablet being formed by compression of a plurality of particles, wherein the oral delivery vehicle tablet comprises sugar alcohol(s) in an amount of 40 to 99.9 % by weight of the delivery vehicle tablet, wherein the oral delivery vehicle tablet comprises non-direct compressible (non-DC) erythritol particles in an amount of 15-50% by weight of the delivery vehicle tablet, wherein the oral delivery vehicle tablet comprises one or more binders other than binders granulated to the sugar alcohol(s) in an amount of 0.1 to 6% by weight of the tablet, wherein the delivery vehicle tablet comprises a plurality of further sugar alcohol particles in an amount of at least 10% by weight of the delivery vehicle tablet and wherein said further sugar alcohols particles have a composition which is different from said non-DC erythritol.



WO 2018/091050 A1

TABLET COMPRISING SEPARATE BINDER AND NON-DIRECT COMPRESSION ERYTHRITOL**Field of the invention**

The invention relates to an oral delivery tablet.

5

Background of the invention

In the prior art it is well-known to provide oral delivery vehicles made by compression and where a substantial part of the delivery vehicle is comprised of a sugar alcohol. It is also well-known that such delivery vehicles may be very attractive for delivery of active ingredients orally. Such substances may include pharmaceuticals, nutrients, flavors, vitamins, smoke inhibitors, dietary supplements, etc.

10

A challenge related to such delivery vehicles is that the user of such vehicles, e.g. tablets, is more and more focused on convenience and attractive taste. Such requirements do very often conflict with what is technically possible. This is in particular relevant to delivery vehicles in the form of tablets of compressed particles.

15

EP application 0 922 464 A1 discloses a quickly disintegrable compression-molded material such as tablets produced through conventional tableting processes. The tablets comprise granulated erythritol and exhibits rapid disintegration and dissolution when placed in the oral cavity or water.

20

EP 0 922 464 is thus a representative example of one of many references using granulated polyols, such as erythritol for compressed tablets.

25

It is an object of the invention to provide an oral delivery vehicle suitable for delivery of such active ingredients in oral cavity of a user, and it is an object to obtain a delivery vehicle which may deliver such active ingredients in the oral cavity of a user in such a way that the user finds the delivery process attractive.

Summary of the invention

The invention relates to an oral delivery vehicle tablet, the delivery vehicle tablet being formed by compression of a plurality of particles,

5

wherein the oral delivery vehicle tablet comprises sugar alcohol(s) in an amount of 40 to 99.9% by weight of the delivery vehicle tablet,

wherein the oral delivery vehicle tablet comprises non-direct compressible (non-DC) erythritol particles in an amount of 15- 50% by weight of the delivery vehicle tablet,

10

wherein the oral delivery vehicle tablet comprises one or more binders other than binders granulated to the sugar alcohol(s) in an amount of 0.1 to 6% by weight of the tablet,

15

wherein the oral delivery vehicle tablet comprises a plurality of further sugar alcohol particles in an amount of at least 10% by weight of the delivery vehicle tablet and wherein said further sugar alcohols particles have a composition which is different from said non-DC erythritol.

20

Binder is one of an important excipient to be added in tablet formulation. Binders are the substances that promotes cohesiveness.

In the present context, the non-DC erythritol particles are understood and defined with reference to their trade grade.

25

Other more physical understandings, not conflicting with the above definition, as that non-DC erythritol particles are particles which has not been preprocessed by granulation with other sugar alcohols or binders for the purpose of obtaining so-called direct compressible particles (DC). Such particles of erythritol may typically consist of erythritol or at least comprise very high quantities of erythritol.

30

The present invention benefits from a synergy between the non-DC erythritol and the further sugar alcohol particles. The further sugar alcohols may be e.g. sorbitol which is direct compressible by nature or it may be other sugar alcohols which have been preprocessed, e.g. by granulation with a suitable binder. At the same time, the non-DC erythritol particles serves as a means for salivation which is both attractive to the user and also serves for the purpose of dissolving the further sugar alcohol particles when the tablet is chewed as fast as possible.

The use of high content of non-DC erythritol particles was not expected to work by the inventors of the present invention since the general expectation in the art of tablet pressing is that only DC grade erythritol would be possible to use in a tablet.

In the present context it is noted that the use of one or more binders other than binders which may be in the tablet and be granulated to the non-erythritol sugar alcohol(s) provides an attractive synergy with the non-DC erythritol with respect to sensorics while still obtaining a mechanically stable tablet.

The one or more binders other than binders which may be in the tablet and be granulated to the non-erythritol sugar alcohol(s) may be in the form of particles and they may therefore obtain the desired effect as separate added binder instead of requiring granulation.

In the present context, it is noted that the use of one or more separate added binders suitable for direct compression other than binders suitable for (pre)granulations which may be in the tablet and be (pre)granulated to non-erythritol sugar alcohols provides an attractive effect/combination with the non-DC erythritol with respect to sensorics while still obtaining a mechanically stable tablet.

A non-DC grade of erythritol is typically understood as a reference to commercial understanding and labeling of products and where non-DC grade is designating a product which is not considered suitable for compression. DC is within the art referring to a product enabling the manufacture of tablets of the specified or referred particles by direct compression. A non-DC composition of erythritol is typically understood as

a composition comprising or consisting of particles of pure erythritol, i.e. particles which have not been subject to e.g. granulation with sugar alcohols for the purpose of obtaining particles which as commercially regarded suitable for direct compression.

The binder suitable for direct compression may be present in a tablet according to the invention as one single binder or a combination of binders. A typical content of erythritol in the individual particles of a commercially available erythritol is higher than 95%, such as 98-about 100% by weight of the individual particles.

10 In an embodiment of the invention the binder grade is suitable for direct compression.

When applying a binder according to the provisions of the invention, it is possible to obtain a very advantageous watering effect as these binders suitable for direct compression may then be added to the formulation without being pre-granulated to the non-DC erythritol but still contributing to the stability of the tablet without changing the sensoric and restricting the watering effect of the non-DC erythritol.

20 In an embodiment of the invention the non-DC erythritol particles comprises about 96-100% erythritol, such as 98-100% erythritol by weight, such as 99.5% erythritol by weight of the particles.

In an embodiment of the invention the non-DC erythritol particles are non-granulated.

Suitable binders include Gum Arabic, Methyl Cellulose, Liquid glucose, Tragacanth, Ethyl Cellulose, Gelatin, Hydroxy Propyl Methyl Cellulose (HPMC), Starches, Hydroxy Propyl Cellulose (HPC), Pregelatinized Starch, Sodium Carboxy Methyl Cellulose (NaCMC), Alginic Acid, Polyvinyl Pyrrolidone (PVP), Maltodextrine (MD); Cellulose, Polyethylene Glycol (PEG), Polyvinyl Alcohols, Polymethacrylates, Copovidone or Microcrystalline Cellulose (MCC), alone or in combination.

30

In an embodiment of the invention the one or more binders comprises one or more cellulose binders.

In an embodiment of the invention the one or more binders comprises microcrystalline cellulose (MCC), hydroxypropyl cellulose (HPC) or hydroxypropylmethyl cellulose (HPMC) or any combination thereof.

5

In an embodiment of the invention the oral tablet comprises hydroxypropyl cellulose (HPC) binder in the amount of 0.1 to 6% by weight of the tablet, such as 0.1 to 5%, such as 0.1 to 4%, such as 0.1 to 3%, such as 0.1 to 2% by weight of the tablet.

10 HPC may be applied as a particular attractive binder as the binder, when compared with erythritol exhibits an advantageous sensoric experience when compared to other well-known binders, such as maltodextrin. In particular, the user of HPC lower than 4 % by weight of the tablet is advantageous, such as 0.1 to 3%, such as 0.1 to 2% by weight of the tablet.

15

In an embodiment of the invention the non-DC erythritol particles are particles that are not granulated and the one or more binders are present as separate components in the tablet.

20 In an embodiment of the invention the non-DC erythritol particles are particles consisting of erythritol and not being pregranulated together with the one or more binders that are present in the tablet as separate components.

25 According to the invention the non-DC erythritol particles are particles substantially consisting of erythritol. Thus, the single particles of non-DC erythritol available on the market may include about 96-100% erythritol, such as 98-100% by weight, such as 99.5% by weight.

30 In an embodiment of the invention the non-DC erythritol particles are non-granulated.

Suitable sugar alcohols typically constitute from about 40 to about 100% by weight of the tablet, such as about 80 to about 99% by weight of the tablet.

When including gum base in the formulation sugar alcohols typically constitute from about 5 to about 95% by weight of the tablet, more typically about 20 to about 80% by weight such as 30 to 70% or 30 to 60% by weight of the tablet.

5

High intensity artificial sweetening agents can also be used alone or in combination with the above sweeteners. Preferred high intensity sweeteners include, but are not limited to sucralose, aspartame, salts of acesulfame, alitame, saccharin and its salts, cyclamic acid and its salts, glycyrrhizin, dihydrochalcones, thaumatin, monellin, stevioside (natural intensity sweetener) and the like, alone or in combination. In order to provide longer lasting sweetness and flavor perception, it may be desirable to encapsulate or otherwise control the release of at least a portion of the artificial sweeteners. Techniques such as wet granulation, wax granulation, spray drying, spray chilling, fluid bed coating, conservation, encapsulation in yeast cells and fiber extrusion may be used to achieve desired release characteristics. Encapsulation of sweetening agents can also be provided using another tablet component such as a resinous compound.

Usage level of the artificial sweetener will vary considerably and will depend on factors such as potency of the sweetener, rate of release, desired sweetness of the product, level and type of flavor used and cost considerations. Thus, the active level of artificial sweetener may vary from about 0.001 to about 8% by weight (preferably from about 0.02 to about 8% by weight). When carriers used for encapsulation are included, the usage level of the encapsulated sweetener will be proportionately higher. Combinations of sugar and/or non-sugar sweeteners may be used in the chewing gum formulation.

A tablet according to the invention may, if desired, include one or more fillers/texturisers including as examples, magnesium and calcium carbonate, sodium sulphate, ground limestone, silicate compounds such as magnesium and aluminum silicate, kaolin and clay, aluminum oxide, silicium oxide, talc, titanium oxide, mono-,

di- and tri-calcium phosphates, cellulose polymers, such as wood, and combinations thereof.

In an embodiment of the invention the non-DC erythritol particles are non-DC grade
5 particles, where non-DC stands for particles which are not marketed as directly compressible in a tableting process.

The use of non-DC erythritol when compared to conventionally applied direct compressible erythritol (DC) has shown remarkable effects to the users perception of
10 the delivery vehicle when chewed. This may partly be due to the somewhat larger size of non-DC erythritol, when compared to DC-erythritol, but is may also be a result of a high content of erythritol in the individual particles applied for compression. DC-erythritol, which for obvious reasons are marketed and applied for compression purposes, does not result in such improved salivation effect and taste.

15 It should be noted that the terminology non-DC is easily understood within the field of technology. Suppliers of sugar alcohol provides clear guidance to the user as for the ability for use in relation to compression of tablets. A non-DC particle in this connection is referred to as a particle which is not expressly recommended by the
20 supplier for compression. Example of a non-DC grade of erythritol is Zeros (TM) erythritol 16952F supplied by Cargill whereas an example of a direct compressible (DC) grade of erythritol include ZerosTM DC 16966 also supplied by Cargill.

In an embodiment of the invention the non-DC erythritol particles are defined as non-
25 DC grade with reference to the Compressibility Index according to European Pharmacopeia 6.0 and where non-DC erythritol particles are having a compressibility index which is greater than 21%.

In an embodiment of the invention the non-DC erythritol particles are defined as non-
30 DC with reference to the Compressibility Index according to European Pharmacopeia 6.0 and where non-DC erythritol particles are having a compressibility index which is greater than 21% and less than 37%.

In an embodiment of the invention the further sugar alcohol particles are directly compressible (DC).

- 5 The terminology directly compressible is well-known within the art of tableting, i.e. in technical field of compression of particles in a gathered compressed tablet. Directly compressible is routinely referred to as DC by many manufactures of such particles.

In an embodiment of the invention the further sugar alcohol particles are defined as
10 DC with reference to the Compressibility Index according to European Pharmacopeia 6.0 and where said further sugar alcohol particles are having a compressibility index which is less than 21%, such as less than 15%, such as less than 10%.

In an embodiment of the invention the further sugar alcohol particles includes particles
15 comprising sugar alcohols selected from the group of sorbitol, erythritol, xylitol, lactitol, maltitol, mannitol or isomalt or any combination thereof.

An example of such DC grade of sugar alcohol includes ISOMALT DC 101 provided
20 by BENEIO-Palatinit GmbH.

A further example of a DC grade sugar alcohol includes Zerose™ DC 16966 supplied
by Cargill. It is here noted that the understanding of direct compressible is simply designated by DC. Unless otherwise stated, this terminology will be applied
25 throughout the present application.

In an embodiment of the invention friability of the delivery vehicle tablet is less than
2%, such as less than 1.5%, wherein friability is measured according to European Pharmacopoeia 9.1, test method 2.9.7. by using a pharmaceutical friability-tester PTF
30 10E from Pharma Test.

According to an embodiment of the invention is has been showed that a desired
friability may in fact be obtained even with a stable tablet and an attractive mouthfeel

and taste. The desired mouthfeel is still obtained in spite of the fact that binders would be expected compromising the desired mouthfeel and taste in a compressed tablet and in spite of the fact that the use of the non-DC erythritol particles should compromise the friability when used in such a high amount.

5

In an embodiment of the invention friability of the delivery vehicle tablet is greater than 0.2%, wherein friability is measured according to European Pharmacopoeia 9.1, test method 2.9.7. by using a pharmaceutical friability-tester PTF 10E from Pharma Test.

10

In an embodiment of the invention said non-DC erythritol particles and said further sugar alcohol particles are of different composition,

wherein said non-DC erythritol particles have an average particle size which is larger
15 than the average particle size of said further sugar alcohol particles,

wherein the average particle size is determined according to European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving.

20

The composition of minimum two different types of sugar alcohol particles, the first sugar alcohol particles and the second sugar alcohol particles, may be different by having different chemical nature, different hygroscopicity, different solubility, different particle size, different flowability and/or different morphology may provide
25 the user with a unique sensory experience

In an embodiment of the invention the particle size of at least 80% of the non-DC erythritol particles is greater than 200 micron, such as greater than 225 micron, such as greater than 250 micron and

30

wherein the particle size is determined according to European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving.

- 5 According to this embodiment, it has been realized that the size of the non-DC erythritol particles features an improved salivation and mouthfeel combined with a feasible breaking force of the tablet

10 In an embodiment of the invention the particle size of less than 20% of the non-DC erythritol particles is smaller than 250 micron, such as less than 15% of the non-DC erythritol particles is smaller than 250 micron, such as less than 10% of the non-DC erythritol particles is smaller than 250 micron and

15 wherein the particle size is determined according to European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving.

In an embodiment of the invention the tablet comprises more than 20% by weight of compressed non-DC erythritol particles

20

wherein the resistance to crunching of the tablet is greater than 60N, such as greater than 70N, such as greater than 80N, such as greater than 90N, such as greater than 100 N, such as greater than 110, such as greater than 130N such as greater than 150N,

25 wherein the resistance to crunching of the tablet is less than 300N, such as less than 250N, such as less than 200N,

wherein the resistance to crunching is determined according to European Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

30

In an embodiment of the invention the delivery vehicle tablet comprises at least one module. A module in the context of the invention is referring to a group of particles

which has been compressed into a volume which is comparable to the size of the tablet. In the case of a tablet which is compressed as one module, the complete tablet is regarded as a module. A tablet comprising two or more module will thus have module sizes which each are comparable to the volume of the complete tablet. Comparable in
5 the present context means that the modules are not understood as small particles and a module should at least be greater than 1/20 of the complete tablet volume, preferably greater than 1/10 of the complete tablet volume.

In an embodiment of the invention the delivery vehicle tablet comprises at least one
10 module,

wherein the module comprises more than 20% by weight of compressed non-DC erythritol particles

15 wherein the resistance to crunching of the module is greater than 60N, such as greater than 70N, such as greater than 80N, such as greater than 90N such as greater than 100 N,

wherein the resistance to crunching is determined according to European
20 Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

The module may typically be gathered from a plurality of compressed particles and have a weight which is greater than 0.2 gram and less than 10.
25

In an embodiment of the invention a module is defined as a plurality of particles being compressed together to form a gathered module of particles.

A one-module tablet, i.e. a tablet where the complete tablet is understood as a one
30 single module, may be advantageous as the tablet from a manufacturing perspective may be easier to manufacture as the manufacturing process would not need to take into consideration the that module should be cohered with other modules.

In an embodiment of the invention the delivery vehicle tablet comprises a plurality of delivery vehicle tablet modules.

- 5 In an embodiment of the invention the plurality of modules are slice-like layers.

In an embodiment of the invention the delivery vehicle tablet comprises at least two modules,

- 10 wherein the delivery vehicle tablet comprises more than 10% by weight of compressed non-DC erythritol particles

wherein the resistance to crunching of a first module comprising of compressed non-DC erythritol particles is less than 150N,

15

wherein the resistance to crunching of a second module is more than 100N and more than the resistance to crunching of the first module,

- 20 wherein the second module comprises less compressed non-DC erythritol particles with respect to weight than the first module,

wherein the resistance to crunching of the delivery vehicle tablet is higher than the resistance to crunching of the second module when the second module is separated from the tablet,

25

wherein the resistance to crunching is determined according to European Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

- 30 According to a further embodiment of the invention, the delivery vehicle may be in the form of a multi-module delivery vehicle. This multi-module may have the benefit that the overall hardness of the compressed delivery vehicle may be improved by the

application of a module which by itself has a higher hardness than the module comprising the substantial amount of non-DC erythritol. Non-DC erythritol may have the effect that the module in which large amount of the non-Dc erythritol features a relatively low hardness. This may under certain circumstances be not desirable due to the fact that such low hardness may reflect a relatively low cohesiveness of the vehicle, thereby resulting in an increased friability of the vehicle. Under some circumstances this relatively low hardness may on the other hand reflect that the non-DC is present in substantial amounts and the low hardness may also make the non-DC easily and fast available in particular for the purposes of the second action. When applying a further module having high hardness such as a module formed by substantial amounts of compressed sugar alcohols such as sorbitol, xylitol, isomalt, etc, this support module may both delivery hardness and when broken into pieces during the initial chew benefit from the salivation effect obtained through the non-DC erythritol of the supported module. The third action, i.e. dissolving of sugar alcohols in not only obtained in the non-DC erythritol module but it is also obtained in relation to the sugar alcohols of the support module featuring less salivation effect.

Preferably, the tablet should consist of less than 5 modules, such as less than 4 modules.

In an embodiment of the invention, the tablet may be formed of two modules.

In an embodiment of the invention the oral delivery vehicle tablet comprises 0-60 percent by weight of gum base, such as 10-60 percent by weight of gum base.

An advantage of the above embodiment may be that due to the relatively fast disintegration, the chewing together of the gum base by mastication in the oral cavity of obtained relatively fast.

In an embodiment of the invention the oral delivery vehicle tablet is free of gum base.

30

In an embodiment of the invention the oral delivery vehicle tablet comprises a pharmaceutically active ingredient.

According to an embodiment of the invention, the obtained taste and mouthfeel facilitates an attractive masking of otherwise not to well-tasting chemical substances. One such substance is nicotine, which when delivered through the mouth may cause a sensation referred to as burning. The present invention is advantageous for taste-
5 masking purposes. The fast salivation and significant cooling may partly taste-mask even this badly-tasting substance. Such taste-masking may of course be obtained in relation to several other chemical substances.

10 In an embodiment of the invention the pharmaceutically active ingredient comprises nicotine.

In an embodiment of the invention the oral delivery vehicle tablet comprises an enhancer.

15

In an embodiment of the invention the enhancer comprises pH controlling agent, such as buffering agent.

In an embodiment of the invention the enhancer comprises a mucoadhesive agent.

20

According to a further embodiment of the invention binders in separate particles may be used without compromising the advantageous taste and mouthfeel according to the invention. This is highly surprising as it appears that the presence of binders or other
25 compression aiding substances appears to counteracts the desired effect, i.e. improved taste and mouthfeel.

In an embodiment of the invention the delivery vehicle tablet comprises flavor in an amount of 1-10% by weight of the delivery vehicle, such as 1-6% by weight of the
30 delivery vehicle, such as 2-6% by weight of the delivery vehicle.

In embodiments of the present invention, the tablet comprises one or more flavoring agents selected from the group consisting of essential oils, essences, extracts, powders, acids, coconut, coffee, chocolate, vanilla, grape fruit, orange, lime, menthol, liquorice, caramel aroma, honey aroma, peanut, walnut, cashew, hazelnut, almonds, pineapple, strawberry, raspberry, apple, pear, peach, apricot, blackberry, cherry, pineapple, plum essence, clove oil, bay oil, anise, thyme, cedar leaf oil, nutmeg, cinnamon, peppermint, wintergreen, spearmint, eucalyptus, mint, or any combination thereof.

In an embodiment of the invention the flavor is a powder flavor.

In an embodiment of the invention the flavor further includes a salivation agent.

The salivation agent may e.g. include Optaflow® supplied by Symrise AG. Such agent further encourages saliva production, thereby providing improved mouthfeel and supporting base and flavor performance.

In an embodiment of the invention the weight of non-DC erythritol particles contained in the oral delivery vehicle tablet is greater than 0.3 gram, such as greater than 0.4 gram, such as greater than 0.5 gram, such as greater than 0.6 gram, such as greater than 0.7 gram, such as greater than 0.8 gram, such as greater than 0.9 gram, such as greater than 1.0 gram.

According to a further embodiment of the invention, the amount of non-DC erythritol particles is relatively high. It is in particular high when considering that the erythritol in conventional sense is not regarded attractive for compression, but the mouthfeel and salivation perceived by the user is there improved significantly, when compared to low amounts or the same amounts of DC erythritol.

In an embodiment of the invention the weight of said non-DC erythritol particles contained in the oral delivery vehicle tablet is less than 3.0 gram, such as less than 2.0 gram, such as less than greater than 1.5 gram.

In an embodiment of the invention wherein the oral delivery vehicle tablet has a weight of between 0.5 and 4.0 grams.

5 In an embodiment of the invention wherein the weight ratio between said non-DC erythritol particles and said further sugar alcohol particles in the delivery vehicle tablet is greater than 0.3, such as greater than 0.4, such as greater than 0.5.

10 The weight ratio between non-DC erythritol particles and further sugar alcohol particles has proven significant according to an embodiment of the invention in the sense that a relatively high amount of non-DC erythritol particles must be present in order to obtain the mouthfeel and taste obtained through the invention. However, this taste and mouthfeel also resides in the so-called further sugar alcohol particles. An example of such sugar alcohol particle is xylitol, which, together with the non-DC erythritol particles may provide a mouthfeel which is unique and very attractive to test
15 panels.

In an embodiment of the invention the weight ratio between said non-DC erythritol particles and said further sugar alcohol particles in the delivery vehicle tablet is greater than 0.3, such as greater than 0.4, such as greater than 0.5 and wherein said further
20 sugar alcohol particles are DC sugar alcohol particles.

The weight ratio between non-DC erythritol particles and further sugar alcohol particles has proven significant as mentioned above in relation to the direct sensation and mouthfeel experienced by the user, but is has moreover addressed the challenge in
25 relation to mouthfeel when DC sugar alcohol particles crumbles during the initial chew. The mechanical stability of the tablet is much desired when the tablet is in its non-chewed form, but a fast disintegration and dissolving is desirable when the tablet is chew due to the fact that user of the tablet dislike a sandy mouthfeel induced through small hard-pressed crumbles of DC sugar alcohol. The use of a very high amount of
30 non-DC erythritol particles will facilitate a perceived fast dissolving and disintegration of the tablet after the initial chew.

In an embodiment of the invention the weight ratio between said non-DC erythritol particles and said further sugar alcohol particles in the delivery vehicle tablet is smaller than 0.7, such as smaller than 0.6, such as smaller than 0.55.

- 5 In an embodiment of the invention the weight ratio between said non-DC erythritol particles and said further sugar alcohol particles in the delivery vehicle tablet is smaller than 0.7, such as smaller than 0.6, such as smaller than 0.55 and wherein said further sugar alcohol particles are DC sugar alcohol particles.
- 10 The weight ratio between non-DC erythritol particles and further sugar alcohol particles is important for the purpose of obtaining an advantageous taste and mouthfeel. By having an upper limit of this ratio the chewer will moreover also experience a desirable crunch sensation when starting chewing the tablet, the crunch being obtained through the use of substantial amounts of DC sugar alcohol particles
- 15 and the non-DC erythritol particles.

According to an advantageous embodiment of the invention, the tablet may comprise filler.

- 20 In embodiments of the present invention, the tablet comprises filler in an amount of 0.1 to 50% by weight of the tablet, wherein the filler is hydrophobic and wherein at least 90% of the filler is contained in the tablet throughout the chewing of a user during a chewing period of at least 10 minutes.
- 25 In some embodiments of the invention, a buffer is added, the buffer being selected from the group consisting of a tris buffers, amino acid buffers, carbonate, including monocarbonate, bicarbonate or sesquicarbonate, glycerinate, phosphate, glycerophosphate, acetate, glyconate or citrate of an alkali metal, such as potassium and sodium, e.g. trisodium and tripotassium citrate, or ammonium, and mixtures
- 30 thereof.

When buffer is used, a preferred buffer is sodium bicarbonate and/or sodium carbonate. In some embodiments buffer is not part of the chewing gum formulation. In some other embodiments, buffer is part of the chewing gum formulation.

- 5 In some embodiments of the invention, the amount of buffer is 0.5 to 10% by weight of the tablet.

In some embodiments of the invention the buffer is selected from the group consisting of a carbonate, including mon carbonate, bicarbonate or sesquicarbonate, glycerinate,
10 phosphate, glycerophosphate, acetate, glyconate or citrate of an alkali metal, such as potassium and sodium, e.g. trisodium and tripotassium citrate, or ammonium, tris buffer, amino acids, and mixtures thereof.

In an embodiment of the invention the oral delivery vehicle tablet comprises
15 said non-DC erythritol particles in an amount of 30 to 60% by weight of the tablet, said further sugar alcohol particles in an amount of 55 to 20 % by weight of the tablet, wherein said further sugar alcohol particles comprises xylitol, isomalt or sorbitol alone or in any combination thereof, and wherein said further sugar alcohol particles are
20 directly compressible (DC),

filler in an amount of 5 to 15% by weight of the tablet, and wherein the tablet comprises powdered flavor in an amount of 2 to 6% by weight of the tablet.

- 25 In an embodiment of the invention said further sugar alcohol comprises xylitol in an amount of more than 85% by weight of the further sugar alcohols.

In an embodiment of the invention the oral delivery tablet comprises two separate modules,
30 wherein said modules are layers,

wherein a first layer comprises said non-DC erythritol particles in an amount of 30 to 60% by weight of the modules and wherein said non-DC erythritol particles has a size which is larger than 250 micron,

- 5 wherein the particle size is determined according to the European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving,

- wherein a second layer comprises DC sugar alcohol particles in an amount of 80-
10 100%,

wherein the first layer has a resistance to crunching which is less than the resistance to crunching of the second layer and

- 15 wherein the resistance to crunching is determined according to European Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

- Moreover, the invention relates to a method of manufacturing an oral delivery vehicle.
20

- From the above it may be understood that it is possible to obtain a delivery vehicle, which, when applied as a chewing tablet may perform a fast disintegration in three different actions. The three actions are in principle not starting at the same time, and it should be understood that overlapping may occur due to the logic interaction between
25 these actions.

- The first action in this embodiment of the invention is where the user deposits the delivery vehicle in the mouth and chews the delivery vehicle. This chewing infers a mechanical crunching of the delivery vehicle. This has the consequence that the
30 vehicle gradually, but still very fast, disintegrates into smaller distinct pieces of particles.

The second action, which is significant for obtaining the effect of this embodiment of the invention is that the non-DC erythritol has an attractive effect in the mouth of a user. It provides a cooling effect and a very attractive mouthfeel and taste for the user.

5 Just as important is that the applied non-DC erythritol results in a significant salivation effect in the mouth of a user. This supplements the perceived mouthfeel of the user as this salivation effect is surprisingly convenient.

The third action is obtained as a result of the overall salivation effect. The salivation
10 effect has the further benefit that smaller pieces of compressed particles of fast dissolvable sugar alcohols are dissolved faster due to the high amount of saliva. A fast dissolving of distinct particles in the mouth of the user is highly attractive to the user. Distinct particles in the mouth of a user in relation to chewing of a delivery vehicle is not perceived well by a user. It may, if the mouth feeling lasts too long result in a
15 sandy/grain mouth feel.

Generally according to the terminology applied in the application a delivery vehicle tablet may be understood a oral tablet.

20 According to an advantageous embodiment of the invention, the delivery vehicle may surprisingly be compressed as a one-module delivery vehicle, where the non-DC erythritol is mixed with other sugar alcohol(s) and optional other relevant substances such as pharmaceuticals, flavors, binders etc. It has thus been shown that a delivery vehicle featuring an acceptable hardness and at the same time the desired taste and
25 mouthfeel when chewed may be obtained.

In an embodiment of the invention, the delivery vehicle tablet may moreover comprise flavors, tableting aids, anti-caking agents, emulsifiers, antioxidants, enhancers, absorption enhancers, buffers, high intensity sweeteners, softeners, colors, or any
30 combination thereof.

In an embodiment of the invention the binder grade is suitable for direct compression.

It is advantageous that the applied binder is a grade which is suitable for direct compression when using it with non-DC erythritol.

5 In an embodiment of the invention the tablet comprises one or more binders in the amount of 0.1 to 3% by weight of the tablet, such as 0.1 to 2%, such as 0.1 to 1.5%, such as 0.1 to 1%, such as 0.2 to 0.5% by weight of the tablet. Even very low amounts of these binder, which are not pre-granulated to erythritol prior to compression provides a tablet having great mechanical stability and offers advantageous exploitation of the erythritol with respect to sensorics.

Detailed description

In the following raw materials will refer to the mixed particles to be compressed into a tablet according to embodiments of the invention unless otherwise stated.

- 5 The following description outlines explanations of how the inventive oral delivery vehicle tablet may be produced and further details of what may be added to the inventive composition.

Typically, the process of manufacture of the inventive oral delivery vehicle tablet may
10 be performed in a single tablet press, such as a rotary tablet press. But it may be a benefit under some circumstances to apply a separate tablet press.

Preferably, the upper punch is convex which gives the upper face of the pressed tablet a concave form.

15

It should of course be noted that the shape of the punches may vary depending of the desired tablet shape.

In some embodiments of the invention, pressing of the tablets are performed at a force
20 of 20 to 50 kN.

Important raw materials of the inventive tablet are non-DC erythritol particles and one or more further sugar alcohol particles. The further sugar alcohols particles are preferable sugar alcohols other than the non-DC erythritol particles.

- 25 In the present context, the non-DC erythritol particles are referred to as particles which has not been preprocessed by granulation with other sugar alcohols or binders for the purpose of obtaining so-called direct compressible particles.

The further sugar alcohol particles may typically refer to sugar alcohols known within
30 the art as being direct compressible (DC).

According to a further embodiment of the invention, the applied non-DC erythritol particles is best characterized as being not-directly compressible (non-DC). The use of non-DC erythritol when compared to conventionally applied direct compressible erythritol (DC) has shown remarkable effects to the users perception of the delivery vehicle when chewed. This may partly be due to the somewhat larger size of non-DC erythritol, when compared to DC-erythritol, but it may also be a result of a high content of erythritol in the individual particles applied for compression. DC-erythritol, which for obvious reasons are marketed and applied for compression purposes, does not result in such improved salivation effect and mouthfeel.

10

It should be noted that the terminology non-DC is easily understood within the field of technology. Suppliers of sugar alcohol provides clear guidance to the user as for the ability for use in relation to compression of tablets. A non-DC particle in this connection is referred to as a particle which is not expressly recommended by the supplier for compression. Example of a non-DC grade of erythritol is Zerose (TM) erythritol 16952F supplied by Cargill whereas an example of a direct compressible (DC) grade of erythritol include Zerose™ DC 16966 also supplied by Cargill.

15

Direct compressible sugar alcohols may include sorbitol which is DC by nature, DC grades of Xylitol, DC grades of Erythritol, DC grades of Mannitol, DC grades of maltitol, DC grades of Lactitol, Isomalt or other suitable DC grades of sugar alcohols.

20

The present invention benefits from a synergy between the non-DC erythritol particles and the further sugar alcohol particles. The further sugar alcohols may be e.g. sorbitol which is direct compressible by nature or it may be other sugar alcohols which has been preprocessed, e.g. by granulation with a suitable binder, to obtain particles which when compressed may encapsulate the non-DC erythritol particles into a mechanically stable tablet. At the same time the DC erythritol particles serves as a means for salivation which is both attractive to the user and also serves for the purpose of dissolving the further sugar alcohol particles when the tablet is chewed as fast as possible.

25

30

According to embodiments of the invention, encapsulated flavors or active ingredients may be added to the final blend of raw materials prior to compression.

5 Different methods of encapsulating flavors or active ingredients, which may both refer to flavors or active ingredients mixed into the raw materials to be compressed into the chewing gum may e.g. include spray drying, spray cooling, film coating, coascervation, Double emulsion method (Extrusion technology) or prilling.

10 Materials to be used for the above-mentioned encapsulation methods may e.g. include Gelatine, Wheat protein, Soya protein, Sodium caseinate, Caseine, Gum arabic, Mod. starch, Hydrolyzed starches (maltodextrines), Alginates, Pectin, Carregeenan, Xanthan gum, Locus bean gum, Chitosan, Bees wax, Candelilla wax, Carnauba wax, Hydrogenated vegetable oils, Zein and/or Sucrose.

15 Preferably, these ingredients should be added subsequent to any significant heating or mixing. In other words, the active ingredients should preferably be added immediately prior to the compression of the final tablet.

20 If applying the present invention in relation to chewing gum, the adding of active ingredients may be cautiously blended with pre-mixed gum base granulates and further ingredients, immediately prior to the final compression of the tablet.

In one embodiment the tablet according to the invention comprises a pharmaceutically, cosmetically or biologically active substance. Examples of such active substances, a
25 comprehensive list of which is found e.g. in WO 00/25598, include drugs, dietary supplements, antiseptic agents, pH adjusting agents, anti-smoking agents and substances for the care or treatment of the oral cavity and the teeth such as hydrogen peroxide and compounds capable of releasing urea during chewing. Examples of useful active substances in the form of
30 antiseptics include salts and derivatives of guanidine and biguanidine (for instance chlorhexidine diacetate) and the following types of substances with limited water-

solubility: quaternary ammonium compounds (e.g. ceraminate, chloroxylenol, crystal violet, chloramine), aldehydes (e.g. paraformaldehyde), derivatives of dequaline, polynoxyline, phenols (e.g. thymol, p-chlorophenol, cresol), hexachlorophene, salicylic anilide compounds, triclosan, halogenes (iodine, iodophores, chloramine, dichlorocyanuric acid salts), alcohols (3,4 dichlorobenzyl alcohol, benzyl alcohol, phenoxyethanol, phenylethanol), cf. also Martindale, The Extra Pharmacopoeia, 28th edition, pages 547-578; metal salts, complexes and compounds with limited water-solubility, such as aluminum salts, (for instance aluminum potassium sulphate $AlK(SO_4)_2 \cdot 12H_2O$) and salts, complexes and compounds of boron, barium, strontium, iron, calcium, zinc, (zinc acetate, zinc chloride, zinc gluconate), copper (copper chloride, copper sulphate), lead, silver, magnesium, sodium, potassium, lithium, molybdenum, vanadium should be included; other compositions for the care of mouth and teeth: for instance; salts, complexes and compounds containing fluorine (such as sodium fluoride, sodium monofluorophosphate, aminofluorides, stannous fluoride), phosphates, carbonates and selenium. Further active substances can be found in J. Dent. Res. Vol. 28 No. 2, pages 160-171, 1949.

Examples of active substances in the form of agents adjusting the pH in the oral cavity include: acids, such as adipic acid, succinic acid, fumaric acid, or salts thereof or salts of citric acid, tartaric acid, malic acid, acetic acid, lactic acid, phosphoric acid and glutaric acid and acceptable bases, such as carbonates, hydrogen carbonates, phosphates, sulphates or oxides of sodium, potassium, ammonium, magnesium or calcium, especially magnesium and calcium.

Active ingredients may comprise the below mentioned compounds or derivatives thereof but are not limited thereto: Acetaminophen, Acetylsalicylic acid, Buprenorphine, Bromhexin, Celcoxib, Codeine, Diphenhydramin, Diclofenac, Etoricoxib, Ibuprofen, Indometacin, Ketoprofen, Lumiracoxib, Morphine, Naproxen, Oxycodon, Parecoxib, Piroxicam, Pseudoefedrin, Rofecoxib, Tenoxicam, Tramadol, Valdecoxib, Calciumcarbonat, Magaldrate, Disulfiram, Bupropion, Nicotine, Azithromycin, Clarithromycin, Clotrimazole, Erythromycin, Tetracycline, Granisetron, Ondansetron, Prometazin, Tropisetron, Brompheniramine, Ceterizin, leco-Ceterizin, Chlorcyclizine,

Chlorpheniramin, Chlorpheniramin, Difenhidramine, Doxylamine, Fenofenadin, Guaifenesin, Loratidin, des-Loratidin, Phenyltoloxamine, Promethazin, Pyridamine, Terfenadin, Troxerutin, Methyldopa, Methylphenidate, Benzalcon. Chloride, Benzeth. Chloride, Cetylpyrid. Chloride, Chlorhexidine, Ecabet-sodium, Haloperidol, 5 Allopurinol, Colchicine, Theophylline, Propanolol, Prednisolone, Prednisone, Fluoride, Urea, Actot, Glibenclamide, Glipizide, Metformin, Miglitol, Repaglinide, Rosiglitazone, Apomorfin, Cialis, Sildenafil, Vardenafil, Diphenoxylate, Simethicone, Cimetidine, Famotidine, Ranitidine, Ratinidine, cetirizin, Loratadine, Aspirin, Benzocaine, Dextrometorphan, Phenylpropanolamine, Pseudoephedrine, Cisapride, 10 Domperidone, Metoclopramide, Acyclovir, Dioctylsulfosucc., Phenolphthalein, Almotriptan, Eletriptan, Ergotamine, Migea, Naratriptan, Rizatriptan, Sumatriptan, Zolmitriptan, Aluminum salts, Calcium salts, Ferro salts, Ag-salts, Zinc-salts, Amphotericin B, Chlorhexidine, Miconazole, Triamcinolonacetoneid, Melatonine, Phenobarbitol, Caffeine, Benzodiazepiner, Hydroxyzine, Meprobamate, 15 Phenothiazine, Buclizine, Brometazine, Cinnarizine, Cyclizine, Difenhidramine, Dimenhydrinate, Buflomedil, Amphetamine, Caffeine, Ephedrine, Orlistat, Phenylephedrine, Phenylpropanolamin, Pseudoephedrine, Sibutramin, Ketoconazole, Nitroglycerin, Nystatin, Progesterone, Testosterone, Vitamin B12, Vitamin C, Vitamin A, Vitamin D, Vitamin E, Pilocarpin, Aluminumaminoacetat, Cimetidine, 20 Esomeprazole, Famotidine, Lansoprazole, Magnesiumoxide, Nizatide and or Ratinidine.

The invention is suitable for increased or accelerated release of active agents selected among the group of dietary supplements, oral and dental compositions, antiseptic 25 agents, pH adjusting agents, anti-smoking agents, sweeteners, flavorings, aroma agents or drugs. Some of those will be described below.

The active agents to be used in connection with the present invention may be any substance desired to be released from the tablet. The active agents, for which a 30 controlled and/or accelerated rate of release is desired, are primarily substances with a limited water-solubility, typically below 10 g/100 ml inclusive of substances which are totally water-insoluble. Examples are medicines, dietary supplements, oral

compositions, anti-smoking agents, highly potent sweeteners, pH adjusting agents, flavorings etc.

Other active ingredients are, for instance, paracetamol, benzocaine, cinnarizine,
5 menthol, carvone, caffeine, chlorhexidine-di-acetate, cyclizine hydrochloride, 1,8-cineol, nandrolone, miconazole, mystatine, sodium fluoride, nicotine, cetylpyridinium chloride, other quaternary ammonium compounds, vitamin E, vitamin A, vitamin D, glibenclamide or derivatives thereof, progesterone, acetylsalicylic acid, dimenhydrinate, cyclizine, metronidazole, sodium hydrogen carbonate, the active
10 components from ginkgo, the active components from propolis, the active components from ginseng, methadone, oil of peppermint, salicylamide, hydrocortisone or astemizole.

Examples of active agents in the form of dietary supplements are for instance salts and
15 compounds having the nutritive effect of vitamin B2 (riboflavin), B12, folic acid, folic acid, niacine, biotine, poorly soluble glycerophosphates, amino acids, the vitamins A, D, E and K, minerals in the form of salts, complexes and compounds containing calcium, phosphorus, magnesium, iron, zinc, copper, iodine, manganese, chromium, selenium, molybdenum, potassium, sodium or cobalt.

20

Furthermore, reference is made to lists of nutritionists accepted by the authorities in different countries such as for instance US code of Federal Regulations, Title 21, Section 182.5013.182.5997 and 182.8013-182.8997.

25 Examples of active agents in the form of compounds for the care or treatment of the oral cavity and the teeth are for instance bound hydrogen peroxide and compounds capable of releasing urea during chewing.

Examples of active agents in the form of antiseptics are for instance salts and
30 compounds of guanidine and biguanidine (for instance chlorhexidine diacetate) and the following types of substances with limited water-solubility: quaternary ammonium compounds (for instance ceramium, chloroxylonol, crystal violet, chloramine),

aldehydes (for instance paraformaldehyde), compounds of dequaline, polynoxyline, phenols (for instance thymol, para chlorophenol, cresol) hexachlorophene, salicylic anilide compounds, triclosan, halogenes (iodine, iodophores, chloroamine, dichlorocyanuric acid salts), alcohols (3,4 dichlorobenzyl alcohol, benzyl alcohol, phenoxyethanol, phenylethanol), cf. furthermore Martindale, The Extra Pharmacopoeia, 28th edition, pages 547-578; metal salts, complexes and compounds with limited water-solubility, such as aluminum salts, (for instance aluminum potassium sulphate $AlK(SO_4)_2 \cdot 12H_2O$) and furthermore salts, complexes and compounds of boron, barium, strontium, iron, calcium, zinc, (zinc acetate, zinc chloride, zinc gluconate), copper (copper chloride, copper sulfate), lead, silver, magnesium, sodium, potassium, lithium, molybdenum, vanadium should be included; other compositions for the care of mouth and teeth: for instance; salts, complexes and compounds containing fluorine (such as sodium fluoride, sodiummonofluorophosphate, amino fluorides, stannous fluoride), phosphates, carbonates and selenium.

Cf. furthermore J. Dent. Res. Vol. 28 No. 2, pages 160-171, 1949, wherein a wide range of tested compounds is mentioned.

Examples of active agents in the form of agents adjusting the pH in the oral cavity include for instance: acceptable acids, such as adipic acid, succinic acid, fumaric acid, or salts thereof or salts of citric acid, tartaric acid, malic acid, acetic acid, lactic acid, phosphoric acid and glutaric acid and acceptable bases, such as carbonates, hydrogen carbonates, phosphates, sulfates or oxides of sodium, potassium, ammonium, magnesium or calcium, especially magnesium and calcium.

Examples of active agents in the form of anti-smoking agents include for instance: nicotine, tobacco powder or silver salts, for instance silver acetate, silver carbonate and silver nitrate.

In a further embodiment, the sucrose fatty acid esters may also be utilized for increased release of sweeteners including for instance the so-called highly potent sweeteners, such as for instance saccharin, cyclamate, aspartame, thaumatin, dihydrocalcones,

stevioside, glycyrrhizin or salts or compounds thereof. For increased released of sweetener, the sucrose fatty acids preferable have a content of palmitate of at least 40% such as at least 50%.

- 5 Further examples of active agents are medicines of any type.

Examples of active agents in the form of medicines include caffeine, salicylic acid, salicyl amide and related substances (acetylsalicylic acid, choline salicylate, magnesium salicylate, sodium salicylate), paracetamol, salts of pentazocine
 10 (pentazocine hydrochloride and pentazocinelactate), buprenorphine hydrochloride, codeine hydrochloride and codeine phosphate, morphine and morphine salts (hydrochloride, sulfate, tartrate), methadone hydrochloride, ketobemidone and salts of ketobemidone (hydrochloride), beta-blockers, (propranolol), calcium antagonists, verapamil hydrochloride, nifedipine as well as suitable substances and salts thereof
 15 mentioned in Pharm. Int., Nov.85, pages 267-271, Barney H. Hunter and Robert L. Talbert, nitroglycerine, erythrityl tetranitrate, strychnine and salts thereof, lidocaine, tetracaine hydrochloride, etorphine hydrochloride, atropine, insulin, enzymes (for instance papain, trypsin, amyloglucosidase, glucoseoxidase, streptokinase, streptodornase, dextranase, alpha amylase), polypeptides (oxytocin, gonadorelin,
 20 (LH.RH), desmopressin acetate (DDAVP), isoxsuprine hydrochloride, ergotamine compounds, chloroquine (phosphate, sulfate), isosorbide, demoxytocin, heparin.
 Other active ingredients include beta-lupeol, Letigen®, Sildenafil citrate and derivatives thereof.

25 Dental products include Carbamide, CPP Caseine Phospho Peptide; Chlorhexidine, Chlorhexidine di acetate, Chlorhexidine Chloride, Chlorhexidine di gluconate, Hexetidine, Strontium chloride, Potassium Chloride, Sodium bicarbonate, Sodium carbonate, Fluor containing ingredients, Fluorides, Sodium fluoride, Aluminum fluoride.

30

Ammonium fluoride, Calcium fluoride, Stannous fluoride, Other fluor containing ingredients Ammonium fluorosilicate, Potassium fluorosilicate, Sodium fluorosilicate,

Ammonium monofluorophosphate, Calcium monofluorophosphate, Potassium monofluorophosphate, Sodium monofluorophosphate, Octadecentyl Ammonium fluoride, Stearyl Trihydroxyethyl Propylenediamine Dihydrofluoride, Vitamins include A, B1, B2, B6, B12, Folinic acid, Folic acid, niacin, Pantothenic acid, biotine, C, D, E, K. Minerals include Calcium, phosphor, magnesium, iron, Zinc, Cupper, Iod, Mangan, Crom, Selene, Molybden. Other active ingredients include: Q10®, enzymes. Natural drugs including Ginkgo Biloba, ginger, and fish oil.

The invention also relates to use of migraine drugs such as Serotonin antagonists: Sumatriptan, Zolmitriptan, Naratriptan, Rizatriptan, Eletriptan; nausea drugs such as Cyclizin, Cinnarizin, Dimenhydramin, Difenhydrinat; hay fever drugs such as Cetrizin, Loratidin, pain relief drugs such as Buprenorfin, Tramadol, oral disease drugs such as Miconazol, Amphotericin B, Triamcinolonaceton; and the drugs Cisaprid, Domperidon, Metoclopramid. In a preferred embodiment the invention relates to the release of Nicotine and its salts.

In an embodiment of the invention, the powdered tablet materials beside the already described sugar alcohols selected from the group consisting of bulk sweeteners, flavors, dry-binders, tableting aids, anti-caking agents, emulsifiers, antioxidants, enhancers, absorption enhancers, buffers, high intensity sweeteners, softeners, colors, or any combination thereof.

Suitable sugar alcohols typically constitute from about 40 to about 100% by weight of the tablet, such as about 80 to about 99% by weight of the tablet.

When including gum base in the formulation sugar alcohols typically constitute from about 5 to about 95% by weight of the tablet, more typically about 20 to about 80% by weight such as 30 to 70% or 30 to 60% by weight of the tablet.

High intensity artificial sweetening agents can also be used alone or in combination with the above sweeteners. Preferred high intensity sweeteners include, but are not limited to sucralose, aspartame, salts of acesulfame, alitame, saccharin and its salts, cyclamic acid and its salts, glycyrrhizin, dihydrochalcones, thaumatin, monellin,

stevioside (natural intensity sweetener) and the like, alone or in combination. In order to provide longer lasting sweetness and flavor perception, it may be desirable to encapsulate or otherwise control the release of at least a portion of the artificial sweeteners. Techniques such as wet granulation, wax granulation, spray drying, spray
5 chilling, fluid bed coating, conservation, encapsulation in yeast cells and fiber extrusion may be used to achieve desired release characteristics. Encapsulation of sweetening agents can also be provided using another tablet component such as a resinous compound.

10 Usage level of the artificial sweetener will vary considerably and will depend on factors such as potency of the sweetener, rate of release, desired sweetness of the product, level and type of flavor used and cost considerations. Thus, the active level of artificial sweetener may vary from about 0.001 to about 8% by weight (preferably from about 0.02 to about 8% by weight). When carriers used for encapsulation are
15 included, the usage level of the encapsulated sweetener will be proportionately higher. Combinations of sugar and/or non-sugar sweeteners may be used in the oral delivery vehicle formulation.

A tablet according to the invention may, if desired, include one or more
20 fillers/texturisers including as examples, magnesium and calcium carbonate, sodium sulphate, ground limestone, silicate compounds such as magnesium and aluminum silicate, kaolin and clay, aluminum oxide, silicium oxide, talc, titanium oxide, mono-, di- and tri-calcium phosphates, cellulose polymers and combinations thereof.

Examples**Examples 1-7. Preparation of oral delivery vehicle tablet.**

Rw material (wt%)	Ex1	Ex2	Ex3	Ex4	Ex5	Ex6	Ex7
Non DC Erythritol	27.5	27.5	27.5	27.5	27.5	27,5	27.5
Erythritol DC	-	-	-	-	-	-	-
Xylitol DC	39	39	39	39	39	39	39
Isomalt DC	20.14	17.14	14.14	15.14	10.14	10.14	0.14
CaCO ₃ DC	7	7	7	7	7	7	7
Flavor	5	5	5	5	5	5	5
Acesulfame- K	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Sucralose	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Color	0.16	0.16	0.16	0.16	0.16	0.16	0.16
Magnesium Stearate	1	1	1	1	1	1	1
HPC	-	3	6	-	-	-	-
CMC	-	-	-	5	10	-	-
MD	-	-	-	-	-	10	20

Table 1 – Oral delivery vehicle compositions for 1.6 gram tablets. Amounts are given
5 in wt-% of the tablet.

Rw material (wt%)	Ex8	Ex 9	Ex10	Ex11	Ex12	Ex13
Non-DC	27.5	27.5	27.5	27.5	-	27.5
Erythritol						
Erythritol DC	-	-	-	-	27.5	-
Xylitol DC	39	39	39	39	39	39
Isomalt DC	17.14	15.14	10.14	0.14	20.14	20.14
CaCO3 DC	7	7	7	7	7	7
Flavor	5	5	5	5	5	5
Acesulfame- K	0.05	0.05	0.05	0.05	0.05	0.05
Sucralose	0.15	0.15	0.15	0.15	0.15	0.15
Color	0.16	0.16	0.16	0.16	0.16	0.16
Magnesium Stearate	1	1	1	1	1	1
HPMC	3	5	-	-	-	-
MCC	-	-	10	20	-	-

Table 2 – Oral delivery vehicle compositions for 1.6 gram tablets. Amounts are given in wt-% of the tablet.

Rw material (wt%)	Ex14	Ex15	Ex16	Ex17	Ex18	Ex19	Ex20
Non-DC Erythritol	27.5	27.5	27.5	27.5	27.5	27.5	27.5
Xylitol DC	39	39	39	39	39	39	39.5
Isomalt DC	19.64	19.14	18.14	17.14	19.14	17.14	15.14
CaCO ₃ DC	7	7	7	7	7	7	7
Flavor	5	5	5	5	5	5	5
Acesulfame- K	0.05	0.05	0.05	0.05	0.05	0.05	0.05 10
Sucralose	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Color	0.16	0.16	0.16	0.16	0.16	0.16	0.16
Magnesium Stearate	1	1	1	1	1	1	1
HPC	0.5	1	2	3	-	-	- 15
MCC	-	-	-	-	1	3	5

Table 3 – Oral delivery vehicle compositions for 1.6 gram tablets. Amounts are given
20 in wt-% of the tablet.

The materials described above are generally well-known to the person skilled within
the art. As for the various tested binders, these are listed below.

- 25 HPC: Hydroxy propyl cellulose. KlucelTM Nutra D from Ashland
CMC: Carboxymethyl cellulose. WalocelTM CRT 30 PA from Dow
MD: Maltodextrin. C DryTM MD 01904 from Cargill
HPMC: Hydroxypropylmethylcellulose. MethocelTM K4M from Dow
MCC: Microcrystalline cellulose. EnduranceTM MCC VE-090e from FMC

Process flow

The compositions indicated in the above tables 1-3 are processed into six corresponding tablets, as examples 1-20.

- 5 For each of the examples 1-20 the raw materials are sieved with a 1600 micron sieve and then weighed into the proper amount according to the exemplified compositions of table 1.

- 10 The weighed amounts are then added to a Turbula mixer in a stainless steel container and blended at 50 rpm for 4 minutes and then adding magnesium stearate and blending one additional minute.

- The mixtures are then tableted by means of a Piccola RIVA DC-SC-041-2. A Fette 3090i may also applied.

- 15 The resulting tablets according to Examples 1-20 are then obtained by tableting with a suitable pressure force at about 28-30kN as main compression force.

Ex	Resistance to crunch	Friability (%)	Processability	Sensorics	Suitable Fast dissolving Chewable tablet	WE 1-5 1 low 5 high
1 Ref	153	1.55	Good	Good	Nice crunchy fast dissolving tablet Fresh taste	5
2 3 % HPC	167	0.75	Good	Acc	Nice crunchy chew Almost as fast dissolving as reference Slightly fatty mouth feel	4
3 6 % HPC	170	0.90	Good	Poor	Slower dissolving Big residual particles in mouth for a long time Fatty mouth feel	3
4 5 % CMC	134	15,3	Acc	Poor	Sticky unpleasant feeling	1
5 10 % CMC	167	3,2	Acc	Poor	Unacceptable sticky feeling Does not dissolve or provide watering effect	1
6 10 % MD	150	4.0	Acc	Poor	A more crumble break Off taste	4

					Erythritol crunchiness Leaves a fatty mouthfeel on tough and teeth	
7 20 % MD	130	13.3	Acc	Poor	A very crumble break Off taste Less crunchy Fatty mouthfeel	3
8 3 % HPMC	160	0.87	Good	Acc	Closer to standard, but slightly more fatty mouthfeel Flavor fresh mint	4
9 5 % HPMC	163	1,43	Good	Poor	Different texture, more fatty texture feeling Flavor fresh mint	3
10 10 % MCC	167	1.2	Good	Poor	Initial chew, but does not disintegrate in the right way. More fatty mouthfeel and less taste. Sticks to teeth	3
11 20 % MCC	184	1.0	Good	Poor	Unacceptable sticky Does not dissolve quickly Fatty mouth feel No fresh taste	2

12 ERC DC	210	0.8	Poor	Poor	Too hard for a chewable tablet, does not dissolve the right way.	3
-----------------	-----	-----	------	------	---	---

Table 4 – Evaluation of Examples 1-12. WE = Watering effect, Sensorics= Total sensory experience - Good/Acceptable(Acc)/Poor

Ex	Resistance to crunch	Friability (%)	Processability	Sensorics	Suitable Fast dissolving Chewable tablet	WE 1-5 1 low 5 high
Ex 13 Ref	144	1.42	Good	Good	Nice crunchy fast dissolving tablet Fresh taste	5
Ex 14 0.5 % HPC	154	1.12	Good	Good	Nice crunchy fast dissolving tablet Fresh taste Similar to standard	5
Ex 15 1 % HPC	162	0.87	Good	Good	Nice crunchy fast dissolving tablet Fresh taste Similar to standard	5
Ex 16 2 % HPC	143	0.78	Good	Acc	A bit slower dissolved Fresh and strong mint Erythritol crunchiness Very close to standard	5
Ex 17 3 % HPC	172	0.72	Good	Acc	Fresh and strong mint Erythritol crunchiness A bit slower dissolved Slightly fatty mouthfeel	5
Ex 18 1 % MCC	154	1.06	Good	Acc	A bit more crunchy Fresh mint Dissolve slightly different	5

Ex 19 3 % MCC	166	0.93	Good	Acc	A bit more chruncy Fresh mint Dissolve slower	5
Ex 20 5 % MCC	157	0.80	Good	Poor	Harder and more flaky crunch, slower dissolved A bit sticky Fresh mint	4

Table 5 – Evaluation of Examples 13-20. WE = Watering effect, Sensorics= Total sensory experience - Good/Acceptable(Acc)/Poor

The examples in the above examples 1-20 of Table 4 and 5 a number of relevant parameters: Resistance to crunch given in N, Friability were measured according to European Pharmacopoeia 9.1, test method 2.9.7. by using a pharmaceutical friability-tester PTF 10E from Pharma Test, Processability were evaluated as good, acceptable
5 or poor with respect to suitability for a process flow during manufacture of the tablets, total sensory experience were subjectively evaluated by a test panel as Good, Acceptable (Acc) or Poor, Suitability as fast dissolving chewable tablet was evaluated with a qualitative feedback from a test panel and the Watering effect was evaluated at a scale from 1-5 by a test panel, where 1 designates the lowest watering effect and 5
10 designates the highest watering effect.

The evaluation of the examples shows binder added to the tablets has a variable degree of effect with respect to friability. It is in particular noted that the maltodextrin (MD) of examples 6 and 7 has a very high friability of 4 and 13, 3% respectively even when
15 used in high amounts. It is clear that such a binder is therefore not able to contain or stabilize the non-DC erythritol in the final tablet. It is moreover noted that the binder has a negative effect of the processability and also has too much off taste. It is nevertheless noted that the desired watering effect is relatively high.

It is also noted that the HPMC of examples 8, i.e. a tablet with an amount of 3% by weight of HPMC actually provides a tablet where the friability is very low, the
20 watering effect is high, the sensory evaluation close to standard, i.e. with no binder, and the processability is acceptable.

It is also noted that HPC is even more attractive considering both sensorics, processability and friability compared to other tested binders.

25

The evaluation of examples 14-20 also shows that an acceptable friability, lower as around 1.2 may be obtained with different cellulose binders at the same time as an acceptable sensory evaluation and a really attractive watering effect.

CLAIMS:

1. Oral delivery tablet, the delivery tablet being formed by compression of a plurality of particles,

wherein the oral delivery tablet comprises sugar alcohol(s) in an amount of 40 to 99.9 % by weight of the delivery tablet,

wherein the oral delivery tablet comprises non-direct compressible (non-DC) erythritol particles in an amount of 15-50% by weight of the delivery tablet,

wherein the oral delivery tablet comprises one or more binders other than binders granulated to the sugar alcohol(s) in an amount of 0.1 to 6% by weight of the delivery tablet,

10 wherein the oral delivery tablet comprises a plurality of further sugar alcohol particles in an amount of at least 10% by weight of the delivery tablet,

wherein said plurality of further sugar alcohol particles have a composition which is different from said non-DC erythritol particles, and

wherein said non-DC erythritol particles are particles consisting of erythritol.
- 15 2. Oral delivery tablet according to claim 1, wherein said non-DC erythritol particles comprise about 96 to about 100% erythritol by weight of said non-DC erythritol particles.
3. Oral delivery tablet according to claim 1 or 2, wherein said non-DC erythritol particles are not-granulated.
4. Oral delivery tablet according to any one of claims 1-3, wherein the one or more binders
20 comprise a cellulose binder.
5. Oral delivery tablet according to any one of claims 1-4, wherein the one or more binders are selected from the group consisting of microcrystalline cellulose (MCC), hydroxypropyl cellulose (HPC), hydroxypropylmethyl cellulose (HPMC), and any combination thereof.

6. Oral delivery tablet according to any one of claims 1-5, wherein the oral tablet comprises hydroxypropyl cellulose (HPC) binder in the amount of 0.1 to 6% by weight of the tablet.

7. Oral delivery tablet according to any one of claims 1-6, wherein said non-DC erythritol particles are particles that are not granulated and the one or more binders are present as separate
5 components in the tablet.

8. Oral delivery tablet according to any one of claims 1-6, wherein said non-DC erythritol particles are particles consisting of erythritol and not being granulated together with the one or more binders that are present in the tablet as separate components.

9. Oral delivery tablet according to any one of claims 1-8, wherein said non-DC erythritol
10 particles are defined as non-DC grade with reference to the Compressibility Index according to European Pharmacopeia 6.0, and where said non-DC erythritol particles are having a compressibility index which is greater than 21%.

10. Oral delivery tablet according to any one of claims 1-9, wherein said non-DC erythritol particles are defined as non-DC with reference to the Compressibility Index according to
15 European Pharmacopeia 6.0, and said non-DC erythritol particles are having a compressibility index which is greater than 21% and less than 37%.

11. Oral delivery tablet according to any one of claims 1-10, wherein said plurality of further sugar alcohol particles are directly compressible (DC).

12. Oral delivery tablet according to any one of claims 1-11, wherein said plurality of further
20 sugar alcohol particles are defined as DC with reference to the Compressibility Index according to European Pharmacopeia 6.0, and where said plurality of further sugar alcohol particles are having a compressibility index which is less than 21%.

13. Oral delivery tablet according to any one of claims 1-12, wherein said plurality of further sugar alcohol particles include particles comprising sugar alcohols selected from the group of
25 sorbitol, erythritol, xylitol, lactitol, maltitol, mannitol, isomalt and any combination thereof.

14. Oral delivery tablet according to any one of claims 1-13, wherein friability of the delivery tablet is less than 2%, and wherein friability is measured according to European Pharmacopoeia 9.1, test method 2.9.7. by using a pharmaceutical friability-tester PTF 10E from Pharma Test.

5 15. Oral delivery tablet according to any one of claims 1-14, wherein friability of the delivery tablet is greater than 0.2%, and wherein friability is measured according to European Pharmacopoeia 9.1, test method 2.9.7. by using a pharmaceutical friability-tester PTF 10E from Pharma Test.

10 16. Oral delivery tablet according to any one of claims 1-15, wherein said non-DC erythritol particles and said plurality of further sugar alcohol particles are of different composition, wherein said non-DC erythritol particles have an average particle size which is larger than the average particle size of said plurality of further sugar alcohol particles, and wherein the average particle size is determined according to the European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving.

15 17. Oral delivery tablet according to any one of claims 1-16, wherein the particle size of at least 80% of said non-DC erythritol particles is greater than 200 micron, and wherein the particle size is determined according to European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving.

20 18. Oral delivery tablet according to any one of claims 1-17, wherein the particle size of less than 20% of said non-DC erythritol particles is smaller than 250 micron, and wherein the particle size is determined according to European Pharmacopoeia 9.1 when using test method 2.9.38 particle size distribution estimation by analytical sieving.

25 19. Oral delivery tablet according to any one of claims 1-18, wherein the tablet comprises more than 20% by weight of said non-DC erythritol particles, wherein the resistance to crunching of the tablet is greater than 60N, wherein the resistance to crunching of the tablet is less than 300N, and wherein the resistance to crunching is determined according to European Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

20. Oral delivery tablet according to any one of claims 1-18, wherein the delivery tablet comprises at least one module, wherein the at least one module comprises more than 20% by weight of a plurality of compressed non-DC erythritol particles, wherein the resistance to crunching of the at least one module is greater than 60N, and wherein the resistance to crunching is determined according to the European Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

21. Oral delivery tablet according to claim 20, wherein the at least one module is defined as a plurality of particles being compressed together to form a module.

22. Oral delivery tablet according to claim 21, wherein the delivery tablet comprises a plurality of modules.

23. Oral delivery tablet according to claim 22, where said plurality of modules are layers.

24. Oral delivery tablet according to any one of claims 1-18, wherein the delivery tablet comprises at least two modules, wherein the delivery tablet comprises more than 10% by weight of a plurality of compressed non-DC erythritol particles, wherein the resistance to crunching of a first of the two modules comprising the plurality of compressed non-DC erythritol particles is less than 200N, wherein the resistance to crunching of a second of the two modules is more than 100N and more than the resistance to crunching of the first module, wherein the second module comprises less of the plurality of compressed non-DC erythritol particles with respect to weight than the first module, wherein the resistance to crunching of the delivery tablet is higher than the resistance to crunching of the second module when the second module is separated from the tablet, and wherein the resistance to crunching is determined according to European Pharmacopoeia 9.1, test method 2.9.8. by using a pharmaceutical resistance to crunching tester model Pharma Test type PTB 311.

25. Oral delivery tablet according to any one of claims 1-24, wherein the oral delivery tablet comprises 0-60 percent by weight of a gum base.

26. Oral delivery tablet according to any one of claims 1-24, wherein the oral delivery tablet does not comprise a gum base.

27. Oral delivery tablet according to any one of claims 1-26, wherein said oral delivery tablet comprises an active pharmaceutical ingredient.

28. Oral delivery tablet according to claim 27, wherein the active pharmaceutical ingredient comprises nicotine.

5 29. Oral delivery tablet according to any one of claims 1-28, wherein the oral delivery tablet comprises an enhancer.

30. Oral delivery tablet according to claim 29, wherein the enhancer comprises pH controlling agent.

10 31. Oral delivery tablet according to any one of claims 29-30, wherein the enhancer comprises a mucoadhesive agent.

32. Oral delivery tablet according to any one of claims 1-31, wherein the tablet comprises components for the care of mouth and teeth.

33. Oral delivery tablet according to any one of claims 1-32, wherein the delivery tablet comprises a flavor in an amount of 1-10% by weight of the delivery tablet.

15 34. Oral delivery tablet according to claim 33, wherein the flavor further includes a salivation agent.

35. Oral delivery tablet according to any one of claims 1-34, wherein the weight of said non-DC erythritol particles contained in the oral delivery tablet is greater than 0.3 gram.

20 36. Oral delivery tablet according to any one of claims 1-35, wherein the weight of said non-DC erythritol particles contained in the oral delivery tablet is less than 3.0 gram.

37. Oral delivery tablet according to any one of claims 1-36, wherein the oral delivery tablet has a weight of between 0.5 and 4.0 grams.

38. Oral delivery tablet according to any one of claims 1-37, wherein the weight ratio between said non-DC erythritol particles and said plurality of further sugar alcohol particles in the delivery tablet is greater than 0.3.

39. Oral delivery tablet according to any one of claims 1-38, wherein the weight ratio between
5 said non-DC erythritol particles and said plurality of further sugar alcohol particles in the delivery tablet is greater than 0.3, and wherein said plurality of further sugar alcohol particles are DC sugar alcohol particles.

40. Oral delivery tablet according to any one of claims 1-39, wherein the weight ratio between
10 said non-DC erythritol particles and said plurality of further sugar alcohol particles in the delivery tablet is smaller than 0.7.

41. Oral delivery tablet according to any one of claims 1-40, wherein the weight ratio between said non-DC erythritol particles and said plurality of further sugar alcohol particles in the delivery tablet is smaller than 0.7 and wherein said plurality of further sugar alcohol particles are DC sugar alcohol particles.

15 42. Oral delivery tablet according to any one of claims 1-41, wherein the oral delivery tablet comprises said non-DC erythritol particles in an amount of 30 to 60% by weight of the delivery tablet, said plurality of further sugar alcohol particles in an amount of 20 to 55% by weight of the delivery tablet, wherein said plurality of further sugar alcohol particles are directly compressible (DC) and comprise xylitol, isomalt or sorbitol alone or in any combination, a filler
20 in an amount of 5 to 15% by weight of the delivery tablet, and a powdered flavor in an amount of 2 to 6% by weight of the delivery tablet.

43. Oral delivery tablet according to claim 42, wherein said plurality of further sugar alcohol particles comprise xylitol in an amount of more than 85% by weight of the said plurality of further sugar alcohols.

25 44. Oral delivery tablet according to any one of claims 1-43, wherein the binder grade is suitable for direct compression.

45. Oral delivery tablet according to any one of claims 1-44, wherein the tablet comprises the one or more binders in an amount of 0.1 to 3% by weight of the tablet.